

Phenolic Compounds in Different Olive Varieties

Marco Esti,* Luciano Cinquanta, and Ennio La Notte

Dipartimento di Scienze e Tecnologie Agro-Alimentari, Ambientali e Microbiologiche, Università degli Studi del Molise, Via F. De Sanctis, 86100 Campobasso, Italy

Phenolic compounds in different olive varieties were determined by HPLC analysis over 2 years. Demethyloleuropein was found in only two (Coratina and Leccino) of the eight varieties studied, so it could be used as varietal marker. Elenolic acid glucoside and hydroxytyrosol can be considered indicators of maturation for olives. In fact, as the olives ripen, their tenor increases whereas oleuropein decreases.

Keywords: Phenolic compounds; olive fruit; oleuropein; demethyloleuropein; 3,4-dihydroxyphenylethanol; elenolic acid glucoside

INTRODUCTION

Phenolic compounds in olive fruits are important factors to consider in order to evaluate the quality of virgin olive oil since they are partly responsible for its autoxidation stability (Vazquez et al., 1975; Gutierrez et al., 1977; Montedoro, 1972; Perrin, 1992) and organoleptic characteristics (Vazquez, 1978). Moreover, they have pharmacological properties (Maestro Duràn et al., 1994), are natural antioxidants (Sheabar and Neeman, 1988; Chimi et al., 1988, 1991; Le Tutour and Guedon, 1992), and inhibit Gram-positive microorganisms involved in the fermentation of olive fruit (Fleming et al., 1973; Brenes et al., 1992, 1995). Oleuropein, the main component that produces bitterness in olives, is a heterosidic ester of elenolic acid and 3,4-dihydroxyphenylethanol (Panizzi et al., 1960). As the fruit ripens, oleuropein progressively decreases (Amiot et al., 1986a,b, 1990) and phenolic compounds, such as demethyloleuropein and 3,4-dihydroxyphenylethanol accumulate (Brenes et al., 1995), as does the nonphenolic compound elenolic acid glucoside, which only corresponds to the secoiridoid part of oleuropein (Amiot et al., 1989).

Other natural phenols that have been identified in olives are the flavonol glycosides, such as quercetin 3-rutinoside (rutin) and luteolin 7-glucoside. They are found in varying amounts, depending on the level of ripeness, and the variety (Ragazzi et al., 1973; Vazquez et al., 1974; Vlahov, 1992).

Since the phenolic content of virgin olive oil is influenced by the variety, location, degree of ripeness and the type of oil extraction procedure used (Solinas et al., 1978; Montedoro and Garofolo, 1984; Amiot et al., 1986; Solinas, 1987; Servili et al., 1996), the aim of this study was to assess the composition and evolution (during the harvest-time) of the phenolic compounds in olives, so as to identify when the olives are at their best maturation stage for harvesting in order to guarantee an optimum phenolic content in the relative oils that are currently under study. In addition, differences in the phenolic compound composition of the varieties typical to the area are being studied.

MATERIALS AND METHODS

Plant Material. In November 1995, approximately 5 kg of olives (*Olea europea* L.) from the Gentile (Larino), Gentile (Colletorto), Coratina, and Leccino varieties were hand-picked in the Molise region at different stages of ripeness.

In November 1996, a further 5 kg were picked in 1 week at peak harvest-time. The varieties consisted of Gentile (Larino), Gentile (Colletorto), Gentile (Santacroce), Coratina, Peranzana, Rosciola, Saligna, and Leccino.

Sampling was limited to the period when olives, in the area considered, are harvested and processed.

Analysis. A random selection of 100 g of pulp was immediately frozen in nitrogen and ground. The phenolic compounds were extracted from this powder and purified according to Amiot et al. (1986). Detection and quantification was carried out by HPLC in a Waters 600 apparatus (Milford, MA 01757) with a photodiode array detector (Waters 991) at 20 °C. The 25 cm × 4 mm i.d. column used was filled with Supelcosil ABZ + Plus, 5 μm (Supelco, Inc. Bellefonte, PA), and the flow rate was 1.3 mL/min. The volume of injection was 20 μL. The mobile phase consisted of orthophosphoric acid in water (pH = 2.5) and acetonitrile, with a gradient from 15% to 40% of acetonitrile and the total running time was 20 min.

Oleuropein, quercetin 3-rutinoside, and luteolin 7-glucoside were quantified by using the external standard method at 280 nm for the first compound and at 340 nm for the others. 3,4-Dihydroxyphenylethanol and demethyloleuropein were detected at 280 nm and elenolic acid glucoside was detected at 240 nm and were expressed with the extinction coefficient of oleuropein. 3,4-Dihydroxyphenylethanol, demethyloleuropein and elenolic acid glucoside detections were based on the retention time and the spectrum in a 230–350 nm range.

The 1995 olive fruit samples were also subjected to water and fat analysis.

Determination of water content: sample (100–200 g) was weighed and then dried overnight at 50 °C in a vacuum oven (≤100 mmHg). The sample was cooled for 30 min in a dessiccator and reweighed.

Determination of fat content: a dried sample (2–5 g) was submitted to Soxhlet extraction, using petroleum ether as solvent. After solvent evaporation, the flask containing fat was dried at 100 °C, cooled in a dessiccator, and reweighed.

All the measurements were repeated on triplicate samples; the data reported are the mean of them.

Color. A colorimeter (chroma meter type CR-200b, Minolta, Japan) was used to assess the fruits' color and the Hunter colorimetric system was applied (L^* , lightness; a^* , redness; b^* , yellowness) (Hunter, 1979). The measurement was made on 20 fruit samples.

* Corresponding author (telephone 0039-874-94442; fax 0039-874-98903).

Table 1. Composition and Physical Parameters of Different Varieties of Olive Fruit at Harvest Time in 1995^a

| varieties | harvest-time | fat, % d.b. | water, % | maturation index | puncture force (kg) | <i>L</i> * | <i>a</i> * | <i>b</i> * |
|----------------------|--------------|-------------|----------|------------------|---------------------|------------|-------------|-------------|
| Gentile (Larino) | 11/01/95 | 44.7 | 49.4 | 2.2 | 0.47 ± 0.04 | 44.0 ± 3.2 | -3.1 ± 0.2 | 12.2 ± 0.9 |
| Gentile (Larino) | 11/15/95 | 44.2 | 54.8 | 2.6 | 0.49 ± 0.05 | 42.7 ± 5.2 | -6.5 ± 0.4 | 17.3 ± 1.5 |
| Gentile (Larino) | 11/30/95 | 46.3 | 50.6 | 3.9 | 0.40 ± 0.05 | 38.1 ± 3.8 | -1.8 ± 0.3 | 9.8 ± 0.8 |
| Gentile (Colletorto) | 11/01/95 | 36.9 | 43.1 | 4.2 | 0.37 ± 0.03 | 32.7 ± 4.3 | 1.0 ± 0.1 | -0.1 ± 0.02 |
| Gentile (Colletorto) | 11/15/95 | 42.6 | 53.5 | 4.9 | 0.39 ± 0.05 | 29.3 ± 4.0 | -0.5 ± 0.04 | 2.7 ± 0.16 |
| Gentile (Colletorto) | 11/30/95 | 43.5 | 50.1 | 4.8 | 0.50 ± 0.06 | 30.7 ± 2.9 | -3.8 ± 0.4 | 2.3 ± 0.28 |
| Coratina | 11/01/95 | 38.2 | 50.9 | 0.4 | 0.46 ± 0.07 | 52.0 ± 4.8 | -14.4 ± 0.9 | 24.7 ± 2.6 |
| Coratina | 11/15/95 | 37.0 | 54.0 | 0.8 | 0.60 ± 0.05 | 53.6 ± 3.5 | -19.5 ± 1.2 | 27.6 ± 2.2 |
| Coratina | 11/30/95 | 38.5 | 52.2 | 0.7 | 0.43 ± 0.04 | 48.8 ± 4.2 | -17.6 ± 1.5 | 26.9 ± 2.5 |
| Leccino | 11/01/95 | 37.9 | 47.2 | 4.8 | 0.48 ± 0.08 | 28.4 ± 3.4 | -2.1 ± 0.1 | 1.9 ± 0.12 |
| Leccino | 11/15/95 | 37.5 | 43.4 | 4.5 | 0.48 ± 0.05 | 31.0 ± 3.7 | -1.8 ± 0.1 | 3.6 ± 0.29 |

^a Means ±SD of 20 determinations.

Table 2. Phenolic Compounds (mg/g of Pulp) in Different Varieties of Olive Fruit during Harvest Time in 1995^a

| varieties | harvest-time | 3,4-dihydroxyphenylethanol | elenolic acid glucoside | demethyloleuropein | quercetin 3-rutinoside | luteolin 7-glucoside | oleuropein |
|----------------------|--------------|----------------------------|-------------------------|--------------------|------------------------|----------------------|-------------|
| Gentile (Larino) | 11/01/95 | 0.15 ± 0.03 | 0.32 ± 0.07 | | 0.21 ± 0.03 | 0.30 ± 0.02 | 1.12 ± 0.15 |
| Gentile (Larino) | 11/15/95 | 0.48 ± 0.05 | 1.21 ± 0.19 | | 0.29 ± 0.02 | 0.30 ± 0.03 | 1.45 ± 0.19 |
| Gentile (Larino) | 11/30/95 | 0.62 ± 0.08 | 1.08 ± 0.31 | | 0.40 ± 0.05 | 0.41 ± 0.05 | 0.87 ± 0.09 |
| Gentile (Colletorto) | 11/01/95 | 0.40 ± 0.07 | 0.41 ± 0.05 | | 0.36 ± 0.04 | 0.15 ± 0.01 | 2.08 ± 0.29 |
| Gentile (Colletorto) | 11/15/95 | 0.97 ± 0.12 | 0.95 ± 0.01 | | 0.40 ± 0.05 | 0.23 ± 0.03 | 1.91 ± 0.21 |
| Gentile (Colletorto) | 11/30/95 | 1.10 ± 0.15 | 1.10 ± 0.19 | | 0.54 ± 0.06 | 0.42 ± 0.05 | 1.50 ± 0.19 |
| Coratina | 11/01/95 | 0.30 ± 0.06 | 0.38 ± 0.09 | 1.32 ± 0.17 | 0.14 ± 0.02 | 0.16 ± 0.01 | 1.44 ± 0.21 |
| Coratina | 11/15/95 | 0.47 ± 0.07 | 0.32 ± 0.06 | 1.11 ± 0.12 | 0.20 ± 0.01 | 0.23 ± 0.02 | 1.62 ± 0.18 |
| Coratina | 11/30/95 | 0.52 ± 0.06 | 0.37 ± 0.04 | 0.88 ± 0.07 | 0.29 ± 0.02 | 0.24 ± 0.02 | 1.21 ± 0.16 |
| Leccino | 11/01/95 | 0.37 ± 0.05 | 0.78 ± 0.09 | 0.41 ± 0.05 | 0.37 ± 0.04 | 0.39 ± 0.05 | 0.97 ± 0.11 |
| Leccino | 11/15/95 | 0.60 ± 0.08 | 1.41 ± 0.17 | 1.12 ± 0.15 | 0.41 ± 0.03 | 0.41 ± 0.03 | 0.85 ± 0.09 |

^a Means ±SD of three determinations.

Maturation Index. The maturation index was determined according to the method proposed by the National Institute of Agronomical Research of Spain, San Jaén Station (Solinas et al., 1987). Briefly the empirical procedure consists in distributing a randomly taken sample of 100 olives in 8 groups according to the skin colors: bright green (group $N=0$), green-yellowish (group $N=1$), green with reddish spots (group $N=2$), reddish-brown (group $N=3$), black with white flesh (group $N=4$), black with < 50% purple flesh (group $N=5$), black with ≥ 50% purple flesh (group $N=6$), black with 100% purple flesh (group $N=7$). The index is given by $\Sigma(N_i n_i)/100$ where N_i is the group number and n_i is the olive number in that group. Maturation index values range from 0 ($N=0$ bright green olives, $n=100$) to 7 ($N=100$ purple flesh olives, $n=100$).

Firmness. The firmness was determined on 20 fruits by resistance of flesh to penetration, using a puncture test (fruit pressure tester, FT 011, Effegi, Italy). The puncture force was measured with an aluminium probe (\varnothing_{16} -in. diameter) and the result expressed in kilograms.

Statistical Analysis. The correlations among the chemico-physical parameters of the olive fruits were calculated by using the StatView SE program (Abacus Concepts, Inc., Berkeley, CA, 1984).

Chemicals and Reference Compounds. The reagents (Carlo Erba, Milan, Italy) were analytical or HPLC grade, as required. Oleuropein, luteolin-7-glucoside, and rutin were purchased from Extrasynthèse (Z. I. Lyon-Nord, Genay, France).

RESULTS AND DISCUSSION

The composition and physical parameters of the olive fruits harvested in 1995 are given in Table 1. Water content was affected by seasonal pluviometric behavior. The highest amount was recorded on Nov 15th after 3 days of rain. Differences in fat content, measured on a dry basis, depended on the variety and harvest-time. The correlation coefficients between the chemico-physical parameters of the olive fruits showed that the maturation index (m.i.) was highly correlated to the colorimetric values: m.i. vs. L^* ($r = -0.982$, $P \leq 0.0001$), m.i. vs. a^* ($r = 0.891$, $P \leq 0.0001$), m.i. vs. b^* ($r = -0.963$, $P \leq 0.0001$). Moreover, high correlations between the colo-

rimetric variables were observed: L vs. a^* ($r = -0.873$, $P \leq 0.0001$); L vs. b^* ($r = 0.965$, $P \leq 0.0001$); a^* vs. b^* ($r = -0.938$, $P \leq 0.0001$). The color values of the 1995 varieties, as they ripened, showed the highest values of lightness and yellowness and lowest values of redness were seen in Coratina olives, since the black maturation phase did not occur in the harvest-time considered. Unlike other fruit, differences in firmness, which depended on the time of harvest and the variety (Bourne, 1979), were not found to be correlated to either the colorimetric values or maturation index. Figure 1 shows the HPLC chromatogram of phenolic extracts from drupes of Coratina at 240 nm with the related UV spectra in the range 230–340 nm. 3,4-Dihydroxyphenylethanol ($t_R = 4.09$ min) changed during harvest-time in the four varieties studied in 1995 (Table 2), increasing fourfold in Gentile (Larino), from 0.15 to 0.62 mg/g pulp; about three times in Gentile (Colletorto), from 0.40 to 1.10 mg/g of pulp; and nearly twice in Coratina and Leccino. In the interval considered (1 month) oleuropein ($t_R = 14.69$ min) decreased slightly in the four varieties, while elenolic acid glucoside increased. The data confirm that elenolic acid glucoside and 3,4-dihydroxyphenylethanol are indicators of the maturation of olives, since they increase as the fruit ripens, whereas the amount of oleuropein decreases. This could be correlated to the increased activity of the hydrolytic enzymes (Amiot et al., 1986). In particular glycosidases catalyse the hydrolysis of the oleuropein, with the production of oleuropein aglycon and the dialdehydic form of elenolic acid linked to 3,4-dihydroxyphenylethanol (Lo Scalzo et al., 1993; Montedoro et al., 1993; Limiroli et al., 1995). However, due to its hydrophilic nature, 3,4-dihydroxyphenylethanol, the most important antioxidant in virgin olive oil (Chimi et al., 1988), goes into both the water and the oil during the extraction process in amounts that are related to the distribution coefficient and processing conditions. Therefore, in freshly extracted oil, low concentrations of 3,4-dihy-




Figure 1. Ultraviolet spectra of phenolic compounds in the Coratina variety. Peak number (240 nm): (1) 3,4-dihydroxyphenylethanol; (2) elenolic acid glucoside; (3) demethyloleuropein; (4) quercetin 3-rutinoside; (5) luteolin 7-glucoside; (6) oleuropein.

Table 3. Phenolic Compounds (mg/g of Pulp) in Different Varieties of Olive Fruits in 1996^a

| varieties | harvest-time | 3,4-dihydroxyphenylethanol | elenolic acid glucoside | demethyloleuropein | quercetin 3-rutinoside | luteolin 7-glucoside | oleuropein |
|----------------------|--------------|----------------------------|-------------------------|--------------------|------------------------|----------------------|-------------|
| Leccino | 11/14/96 | 0.54 ± 0.07 | 0.69 ± 0.08 | 1.54 ± 0.18 | 0.41 ± 0.05 | 0.31 ± 0.02 | 0.96 ± 0.07 |
| Gentile (Larino) | 11/12/96 | 0.26 ± 0.03 | 0.99 ± 0.10 | | 0.28 ± 0.02 | 0.23 ± 0.03 | 2.37 ± 0.28 |
| Gentile (S. Croce) | 11/14/96 | 0.23 ± 0.02 | 0.69 ± 0.08 | | 0.19 ± 0.02 | 0.17 ± 0.02 | 0.26 ± 0.03 |
| Gentile (Colletorto) | 11/15/96 | | 0.87 ± 0.06 | | 0.66 ± 0.07 | 0.46 ± 0.06 | 2.06 ± 0.19 |
| Peranzana | 11/17/96 | 0.25 ± 0.03 | 1.07 ± 0.15 | | 0.44 ± 0.03 | 0.44 ± 0.03 | 3.45 ± 0.36 |
| Rosciola | 11/15/96 | 0.48 ± 0.07 | 0.46 ± 0.06 | | 0.37 ± 0.05 | 0.60 ± 0.05 | 0.80 ± 0.09 |
| Saligna | 11/14/96 | 0.05 ± 0.01 | 0.57 ± 0.07 | | 0.25 ± 0.02 | 0.29 ± 0.04 | 0.83 ± 0.07 |
| Coratina | 11/17/96 | 0.31 ± 0.02 | 0.68 ± 0.05 | 2.45 ± 0.26 | 0.32 ± 0.04 | 0.25 ± 0.03 | 0.31 ± 0.05 |

^a Means ±SD of three determinations.

droxyphenylethanol are generally found (Esti et al., 1996), compared with the high concentration of complex phenols containing 3,4-dihydroxyphenylethanol due to their less hydrophilic nature. Moreover, since oleuropein decreases and 3,4-dihydroxyphenylethanol increases, delaying harvest-time, in the period considered, could be less effective in maintaining the oleuropein derivatives in olive oil. The trend for both quercetin 3-rutinoside and luteolin 7-glucoside content was similar, increasing slightly in the later-picked fruit. The differences found between Coratina and Leccino (an earlier-ripening variety) in demethyloleuropein behavior during ripening (Table 2) were probably due to the varieties being at different stages of maturation on the day of analysis. As previously reported (Amiot et al., 1989), demethyloleuropein can be used as a varietal marker, since it was found in Coratina and Leccino only. This finding was confirmed in the second year of the experiment, when eight varieties were compared (Table

3). Demethyloleuropein content was 1.5 and 2.4 mg/g of olive pulp in Leccino and Coratina respectively, while no trace was found in the other varieties. All the varieties, except for Gentile (Colletorto), contained 3,4-dihydroxyphenylethanol and ranged from 0.05 mg/g in Saligna to 0.53 mg/g in Leccino. The highest oleuropein (3.5 mg/g) content was found in Peranzana, which is a later-ripening variety.

The elenolic acid glucoside ranged from 0.45 mg/g in Rosciola to 1.1 mg/g in Peranzana. There was a slight difference in luteolin 7-glucoside and quercetin 3-rutinoside content between the varieties.

CONCLUSION

Demethyloleuropein was found in only two (Coratina and Leccino) of the eight varieties studied over the 2 years, so it could be used as varietal marker. Elenolic acid glucoside and 3,4-dihydroxyphenylethanol can be

considered indicators of maturation for olives. In fact, as the olives ripen, their tenor increases, whereas oleuropein decreases. Moreover, since oleuropein decreases and 3,4-dihydroxyphenylethanol increases and due to their different distribution coefficients in both the water and the oil during the extraction process, delaying the harvest-time in the period considered could be less effective to maintaining the oleuropein derivatives in olive oil. Lastly, a highly significant correlation was found between the maturation index and the Hunter color of the olive fruit.

LITERATURE CITED

- Amiot, M. J.; Fleuriet, A.; Macheix, J. J. Caractérisation des produits de dégradation de l'oleuropein (Technological de-bittering process of olives: characterization of fruits before and during alkaline treatment). *Groupe Phenolic* **1986**, 364–369.
- Amiot, M. J.; Fleuriet, A.; Macheix, J. J. Importance and evolution of phenolic compounds in olive during growth and maturation. *J. Agric. Food Chem.* **1986**, *34*, 823–826.
- Amiot, M. J.; Fleuriet, A.; Macheix, J. J. Accumulation of oleuropein derivatives during olive maturation. *Phytochemistry* **1989**, *28*, 67–69.
- Amiot, M. J.; Tacchini, M.; Fleuriet, A.; Macheix, J. J. Le processus technologique de désamérisation des olives: caractérisation des fruits avant et pendant le traitement alcalin. *Sci. Aliment.* **1990**, *10*, 619–631.
- Bourne, M. C. Theory and application of the puncture test in food texture measurement. In *Food texture and Rheology*; Sherman, P., Ed.; Academic Press: New York; 1979.
- Brenes, M.; Garcia, P.; Duran, M. C.; Garrido, A. Concentration of phenolic compounds change in storage brines of ripe olives. *J. Food Sci.* **1992**, *58*, 347–350.
- Brenes, M.; Rejano, L.; Garcia, P.; Sánchez, A. H.; Garrido, A. Biochemical changes in phenolic compounds during Spanish-style green olive processing. *J. Agric. Food Chem.* **1995**, *43*, 2702–2706.
- Chimi, H.; Sadiq, A.; Le Tutour, B.; Rahmani, M. Contribution à l'étude comparative des pouvoirs antioxydants dans l'huile d'olive du tyrosol, de l'hydroxytyrosol, de l'acide caféique, de l'oleuropeine et du BHT. *Rev. Fr. Corps Gras* **1988**, *35*, 339–344.
- Chimi, H.; Cilliard, P.; Rahmani, M. Peroxil and hydroxyl radical scavenging activity of some natural phenolic extracts. *J. Am. Oil Chem. Soc.* **1991**, *68*, 307–312.
- Esti, M.; Cinquanta, L.; Carrone, A.; Trivisonno, M. C.; La Notte, E.; Gambacorta, G. Composti antiossidanti e parametri qualitativi di oli vergini di oliva prodotti in Molise (Antioxydative compounds and quality parameters in virgin olive oils produced in Molise). *Riv. Ital. Sostanze Grasse* **1996**, *73*, 147–150.
- Fleming, H. P.; Walter, W. M., Jr.; Etchells, J. L. Antimicrobial properties of oleuropein and products of its hydrolysis from green olives. *Appl. Microbiol.* **1973**, *26*, 777–782.
- Gutierrez Gonzales-Quijano, R.; Janer del Valle, C.; Janer del Valle, M. L.; Gutierrez Rosales, F.; Vasquez Roncero, A. Relationship between polyphenol content and the quality and stability of virgin olive oil. *Grasas Aceites* **1977**, *28*, 101–106.
- Hunter, R. S. Scales for the measurement of colour difference. In *The measurement of Appearance*; Wiley: New York, 1979.
- Le Tutour, B.; Guedon, D. Antioxydative activities of *Olea europea* leaves and related phenolic compounds. *Phytochemistry* **1992**, *31*, 1173–1178.
- Limiroli, R.; Consonni, R.; Ottolina, G.; Marsilio, V.; Bianchi, G.; Zetta, L. ¹H and ¹³C NMR Characterization of new Oleuropein Aglycones. *J. Chem. Soc., Perkin Trans. 1* **1995**, *1*, 1519–1523.
- Lo Scalzo, R.; Scarpati, M. L. A new Secoiridoid from Olive Wastewaters. *J. Nat. Prod.* **1993**, *56*, 4, 621–623.
- Maestro-Durán, R.; León Cabello, R.; Ruiz Gutierrez, V. Phenolic compounds from olive (*Olea europea*). *Grasas Aceites* **1994**, *45*, 265–269.
- Montedoro, G. Phenolic substances present in virgin olive oil. Note I: Identification of phenolic acids and their antioxidant power. *Sci. Tecnol. Aliment.* **1972**, 177–186.
- Montedoro, G. F.; Garofolo, L. The qualitative characteristics of virgin olive oils. The influence of variables such as variety, environment, preservation, extraction, conditioning of the finished product. *Riv. Ital. Sostanze Grasse* **1984**, *61*, 157–168.
- Montedoro, G. F.; Servili, M.; Baldioli, M.; Selvaggini, R.; Miniati, E.; Macchioni, A. Simple and Hydrolyzable Compounds in Virgin Olive Oil. 3. Spectroscopic Characterizations of the Secoiridoid Derivatives. *J. Agric. Food Chem.* **1993**, *41*, 2228–2234.
- Panizzi, L.; Scarpati, M. L.; Oriente, E. G. Structure of oleuropein, bitter glycoside with hypotensive action of olive oil. Note II. *Gazz. Chim. Ital.* **1960**, *90*, 1449–1485.
- Perrin, J. L. Les composés mineurs et les antioxygènes naturels de l'olive et de son huile. *Rev. Fr. Corps Gras* **1992**, *39*, 25–32.
- Ragazzi, E.; Veronese, G.; Guiotto, A. Demethyloleuropein, a new glycoside isolated from ripe olives. *Ann. Chim.* **1973**, *63*, 13–20.
- Servili, M.; Baldioli, M.; Begliomini, A.; Montedoro G. Effetto dei complessi colloidali e dell'attività enzimatica endogena delle olive sulla composizione chimica degli oli vergini di oliva. In *Ricerche e innovazioni nell'industria alimentare*; Porretto, S., Ed.; Pinerolo, Italy, 1996; Vol. II, pp 851–860.
- Sheabar, F. Z.; Neeman, I. Separation and concentration of natural antioxidants from the rape of olives. *J. Am. Oil Chem. Soc.* **1988**, *65*, 990–993.
- Solinas, M.; Di Giovacchino, L.; Mascolo, A. The polyphenols of olives and olive oil. Note III: influence of temperature and kneading time on the oil polyphenol content. *Riv. Ital. Sostanze Grasse* **1978**, *55*, 19–23.
- Solinas, M. HRGC analysis of phenolic components in virgin olive oils in relation to the ripening and the variety of olives. *Riv. Ital. Sostanze Grasse* **1987**, *64*, 255–262.
- Vazquez Roncero, A.; Graciani Constante, E.; Maestro Duran, R. Phenolic compounds in olive fruits. I. Polyphenols in pulp. *Grasas Aceites* **1974**, *25*, 269–279.
- Vazquez Roncero, A.; Janer del Valle, C.; Janer del Valle, M. L. Polyphenols content and stability of olive oils. *Grasas Aceites* **1975**, *26*, 14–18.
- Vazquez Roncero, A. Les polyphénols de l'huile d'olive et leur influence sur les caractéristiques de l'huile. *Rev. Fr. Corps Gras* **1978**, *25*, 21–26.
- Vlahov, G. Flavonoids in three olive (*Olea europea*) fruit varieties during maturation. *J. Sci. Food Agric.* **1992**, *58*, 157–159.

Received for review May 8, 1997. Revised manuscript received September 29, 1997. Accepted October 21, 1997.® This work was financially supported by University Consorzio of Molise and Campobasso Province.

JF970391+

® Abstract published in *Advance ACS Abstracts*, December 15, 1997.