

CONTENTS

Introduction

Sources of Variability in Biological Monitoring

Dermal Absorption

Pharmacokinetic Models Used in Setting BEIs®

Documentation: Adopted BEIs®

Acetone

Acetylcholinesterase inhibiting pesticides

Aniline – 2004 Adoption

Arsenic, Elemental and soluble inorganic compounds

Benzene

1,3-Butadiene – 2006 Adoption

Cadmium and inorganic compounds

Carbon disulfide

Carbon monoxide

Chlorobenzene

Chromium (VI) – 2004 Adoption

Cobalt

Cyclohexanol – 2004 Adoption

Cyclohexanone – 2004 Adoption

Dichloromethane – 2005 Adoption

N,N-Dimethylacetamide

N,N-Dimethylformamide

2-Ethoxyethanol [EGEE] and
2-Ethoxyethyl acetate [EGEEA]

Ethyl benzene

Fluorides

Furfural

n-Hexane

n-Hexane – 2003 Adoption

Lead

Mercury

Methanol

Methanol – 2005 Update

Methemoglobin inducers – 2006 Update

2-Methoxyethanol [EGME] and

2-Methoxyethyl acetate [EGMEA]

Methyl n-butyl ketone – 2003 Adoption

Methyl chloroform

4,4'-Methylene bis(2-chloroaniline)

Methyl ethyl ketone

Methyl isobutyl ketone

Nitrobenzene

Parathion

Pentachlorophenol

Phenol – 2006 Update

Polycyclic Aromatic Hydrocarbons
[PAHs] – 2005 Adoption

2-Propanol – 2006 Adoption

Styrene

Styrene – 2003 Adoption

Tetrachloroethylene

Tetrahydrofuran

Toluene

Trichloroethylene

Vanadium pentoxide

Xylenes