

CONTENTS

Preface	xiii
List of Contributors	xv
1 DENSITY, THERMAL PROPERTIES, AND THE GLASS TRANSITION TEMPERATURE OF GLASSES	1
<i>Steve Feller</i>	
Part I: Introduction to Physical Properties and Their Uses	1
Part II: Density	2
1.1 Density: Experimental Background and Theory	2
1.1.1 Overview	2
1.1.2 Experimental Methods and Theory	3
1.1.3 Instrumentation Used for Determining Density	7
1.1.4 Analysis of Data, Extraction of Useful Information, and Other Ways to Express Density	8
1.1.5 Case Studies from Some Glass Systems	13
1.1.6 Conclusion to Density Measurements	19
Part III: Thermal Effects with a Focus on the Glass Transition Temperature	20
1.2 OVERVIEW	20
1.3 EXPERIMENTAL METHODS AND THEORY	20
1.3.2 Differential Thermal Analysis	22
1.4 INSTRUMENTATION USED FOR DETERMINING T_g AND RELATED THERMAL EVENTS	23
1.4.1 DSCs	23
1.4.2 Differential Thermal Analysis	23
1.5 ANALYSIS OF DATA AND EXTRACTION OF USEFUL INFORMATION	25
1.6 CASE STUDIES FROM GLASS SYSTEMS	26
1.6.1 The Glass Transition Temperatures of Barium Borosilicate Glasses [18]	26
1.6.2 Stability Parameters in Lithium Borate Glasses [18]	27
1.7 CONCLUSION TO THERMAL PROPERTIES	30
ACKNOWLEDGMENTS	30
REFERENCES	31

2	INFRARED SPECTROSCOPY OF GLASSES	32
	<i>E.I. Kamitsos</i>	
2.1	INTRODUCTION	32
2.2	BACKGROUND AND THEORY	34
2.2.1	Refractive Index and Dielectric Function	34
2.2.2	Reflectance Spectroscopy of Bulk Materials	36
2.2.3	Infrared Spectra of Thin Films	42
2.3	INSTRUMENTATION	44
2.4	ANALYSIS OF INFRARED DATA	48
2.4.1	Bulk Glasses	48
2.4.2	Thin Films of Amorphous Materials	52
2.5	CASE STUDIES	54
2.5.1	Bulk Glasses	54
2.5.2	Glass Thin Films	63
2.6	CONCLUSIONS	68
	ACKNOWLEDGMENTS	69
	REFERENCES	69
3	RAMAN SPECTROSCOPY OF GLASSES	74
	<i>Rui M. Almeida and Luís F. Santos</i>	
3.1	INTRODUCTION	74
3.2	BACKGROUND	76
3.2.1	Theory	76
3.2.2	Selection Rules	78
3.2.3	Depolarization of Raman Lines	79
3.3	INSTRUMENTATION AND DATA ANALYSIS	80
3.3.1	Light Source	81
3.3.2	Sample Compartment	82
3.3.3	Spectrometer	82
3.3.4	Detector	83
3.3.5	Micro-Raman Spectrometers	84
3.3.6	Resolution	85
3.3.7	Data Analysis	86
3.4	CASE STUDIES	87
3.4.1	Structural Effects of Alkali Incorporation in Silicate Glasses	87
3.4.2	Phase Separation Mechanisms in Transition Metal Phosphate Glasses	92

3.4.3	Raman Study of Niobium Germanosilicate Glasses And Glass-Ceramics	96
3.4.4	Raman Spectroscopy of Chalcogenide Glasses	99
3.5	CONCLUSIONS	103
	ACKNOWLEDGMENTS	103
	REFERENCES	103
4	BRILLOUIN LIGHT SCATTERING	107
	<i>John Kieffer</i>	
4.1	INTRODUCTION	107
4.2	BACKGROUND AND THEORY	110
4.3	INSTRUMENTATION	117
4.4	DATA ANALYSIS AND INFORMATION CONTENT	126
4.5	EXAMPLES OF CASE STUDIES	133
4.5.1	Room-Temperature Glass	133
4.5.2	Temperature Dependence, Glass Transition, and Visco-Elasticity	137
4.5.3	Spatially Confined Systems (e.g., Thin Films)	146
4.5.4	Systems Under Pressure	149
4.5.5	Mechanically Fragile Systems, Soft Matter, and Gels	151
4.6	SUMMARY	154
	REFERENCES	155
5	NEUTRON DIFFRACTION TECHNIQUES FOR STRUCTURAL STUDIES OF GLASSES	158
	<i>Alex C. Hannon</i>	
5.1	INTRODUCTION	158
5.2	INSTRUMENTATION	159
5.2.1	The Neutron	159
5.2.2	The Interactions between a Neutron and a Sample	160
5.2.3	Neutron Sources	161
5.2.4	Neutron Diffractometers	164
5.3	THEORETICAL ASPECTS OF NEUTRON DIFFRACTION ON GLASSES	169
5.3.1	The Static Approximation	169
5.3.2	Scattering from a Single Nucleus	169
5.3.3	Scattering from an Assembly of Nuclei	170
5.3.4	Isotropic Samples	171
5.3.5	Coherent and Incoherent (Distinct and Self) Scattering	171

5.3.6	Atomic Vibrations	173
5.3.7	Real-space Correlation Functions	180
5.4	THE APPLICATION OF NEUTRON DIFFRACTION TO STUDIES OF GLASS STRUCTURE	186
5.4.1	Experimental Corrections	186
5.4.2	Resolution	190
5.4.3	Peak Fitting and Integration	194
5.4.4	Normalization of Data	198
5.4.5	Scattering at low Q	200
5.4.6	Sample-Related Difficulties	203
5.4.7	Partial Correlation Functions	209
5.4.8	Interpretation of Results	218
5.4.9	Modeling	226
5.4.10	The PDF Method	229
	ACKNOWLEDGMENTS	231
	REFERENCES	231
	FURTHER READING	240
6	X-RAY DIFFRACTION FROM GLASS	241
	<i>Christopher J. Benmore</i>	
6.1	INTRODUCTION	241
6.2	BACKGROUND/THEORY	244
6.3	ANALYSIS OF DATA, EXTRACTION OF USEFUL INFORMATION	249
6.4	INSTRUMENTATION	255
6.5	CASE STUDIES	258
6.5.1	SiO ₂ and Oxide Glasses	258
6.5.2	Chalcogenide Glasses	263
6.5.3	Amorphous Materials, Gels, Foams and Fibers	264
6.6	CONCLUSIONS	264
	ACKNOWLEDGMENTS	266
	REFERENCES	267
7	XAFS SPECTROSCOPY AND GLASS STRUCTURE	271
	<i>Giuseppe Dalba and Francesco Rocca</i>	
7.1	INTRODUCTION	271
7.2	THE ORIGINS OF X-RAY ABSORPTION SPECTRA	272
7.3	XAFS INSTRUMENTATION	274

7.4	THE PHYSICAL MECHANISM OF XAFS	278
7.5	EXAFS	279
7.5.1	EXAFS Formula for Glasses	282
7.6	XAFS DATA ANALYSIS	284
7.6.1	Corrections for Instrumental Errors	284
7.6.2	Pre-edge Background Subtraction	284
7.6.3	Post-edge Background Subtraction	285
7.6.4	Normalization	286
7.6.5	Conversion to k -Space, Choice of Threshold Energy E_0 and Weighting	286
7.6.6	Transformation from k -Space to R -Space	286
7.6.7	Fourier Filtering: Reverse Transformation: from R -Space to k -Space	287
7.6.8	Log Amplitude Ratio and Phases Difference Method	288
7.6.9	Fitting Procedure	288
7.7	EXAFS ACCURACY AND LIMITATIONS	289
7.8	XANES	290
7.9	XAFS SPECTROSCOPY APPLIED TO GLASS STRUCTURE: SOME EXAMPLES	291
7.9.1	Silicate Glasses	292
7.9.2	Silica Glass	294
7.9.3	Silica at High Temperature	294
7.9.4	Silica and Germania Glasses under High Pressure	297
7.9.5	Nanoparticles Embedded in Glasses	300
7.9.6	Study of Ionic Conductivity in Superionic Conducting Glasses Doped with AgI	307
7.10	SUMMARY AND CONCLUSIONS	309
	REFERENCES	310

8 NUCLEAR MAGNETIC RESONANCE SPECTROSCOPY OF GLASSES 315

Scott Kroeker

8.1	INTRODUCTION	315
8.2	THEORETICAL BACKGROUND	316
8.2.1	Zeeman Effect	316
8.2.2	Magnetic Shielding	318
8.2.3	Quadrupolar Interaction	319
8.2.4	Dipolar Interactions	320
8.2.5	High Resolution Methodologies	320

8.3	INSTRUMENTATION	323
8.3.1	Magnet	323
8.3.2	Probe	325
8.3.3	Radiofrequency Components	326
8.3.4	Computer Control	326
8.3.5	Measurement Uncertainty	327
8.4	DATA ANALYSIS AND STRUCTURAL INTERPRETATION	329
8.4.1	Chemical Shift Assignments	329
8.4.2	Information from Quadrupolar Effects	330
8.4.3	Low- γ Nuclei	332
8.4.4	Paramagnetic Effects	333
8.5	CASE STUDIES	333
8.5.1	Borophosphate Glasses	333
8.5.2	Aluminosilicate Glasses	336
8.5.3	Borosilicate Glasses	337
8.5.4	Modifier Cations in Alkali Borate Glasses	340
8.6	CONCLUSIONS	341
	ACKNOWLEDGMENTS	342
	REFERENCES	342

9 ADVANCED DIPOLAR SOLID STATE NMR SPECTROSCOPY OF GLASSES 345

Hellmut Eckert

9.1	INTRODUCTION	345
9.2	THEORETICAL ASPECTS	347
9.2.1	Direct Magnetic Dipole–Dipole Coupling	348
9.2.2	Indirect Magnetic Dipole–Dipole Coupling	349
9.3	HETERONUCLEAR EXPERIMENTS	349
9.3.1	Spin Echo Double Resonance	349
9.3.2	Rotational Echo Double Resonance	350
9.3.3	Rotational Echo Adiabatic Passage Double Resonance	353
9.3.4	Cross-polarization	354
9.3.5	Connectivity Studies Based on the Detection of Indirect Spin–Spin Interactions	358
9.3.6	Instrumental Considerations and Caveats.	358
9.4	HOMONUCLEAR EXPERIMENTS	360
9.4.1	Static Spin Echo Decay Spectroscopy	360
9.4.2	Homonuclear Dipolar Recoupling Experiments	362
9.4.3	Instrumental Considerations and Caveats	369

9.5	CASE STUDIES	370
9.5.1	Spatial Distributions of Mobile Ions in Alkali Silicate and Borate Glasses	370
9.5.2	Connectivity Distribution in 70 SiO ₂ -30 [(Al ₂ O ₃) _x (P ₂ O ₅) _{1-x}] Glasses	374
9.5.3	Speciations and Connectivity Distributions in Borophosphate and Thioborophosphate Glasses	380
	ACKNOWLEDGMENTS	386
	REFERENCES	386
10	ATOM PROBE TOMOGRAPHY OF GLASSES	391
	<i>Daniel Schreiber and Joseph V. Ryan</i>	
10.1	INTRODUCTION	391
10.2	BACKGROUND AND THEORY	392
10.3	INSTRUMENTATION	395
10.3.1	APT Specimen Preparation	399
10.3.2	Experimental Procedure and Parameters	401
10.3.3	Data Reconstruction	403
10.4	ANALYSIS METHODS	409
10.4.1	Estimating Error	412
10.5	CASE STUDIES	417
10.5.1	Composition	418
10.5.2	Interfaces	420
10.5.3	Conclusions	424
	ACKNOWLEDGMENTS	425
	REFERENCES	425
	Index	431

INDEX

Note: Page number followed by f and t indicates figure and table respectively.

- Ab initio* molecular dynamics, 255, 291
- Absolute shielding, 318–319
- Absorbance, 36
- Absorption coefficient (μ), 273, 275, 277–279, 284–286
- Accelerator-based neutron sources, 162–164, 163f
- Acetone, 5
- Adiabatic modulus, 154
- Aerodynamic levitation technique, 260
- AgI (silver iodide), 63, 307, 308f, 309f
- Aging of gels, 151–152
- Airy function, 118
- Alkali borates, 10, 13t, 16–19, 59
- Alkali incorporation in silicate glasses, structural effects of, 87–92, 88f, 90f, 91f
- Alkali metal halide salts, as matrix materials, 32–33
- Alkaline borates, 8, 9f, 10, 12f, 55–59, 57f, 58f, 61–63, 62f, 68
- Alkali silicates, 14f, 87, 91, 97, 139, 292–294, 370–371, 374
- Aluminosilicate glasses, 333, 336–337, 358, 369, 394
- American Society for Testing and Materials (ASTM) methods, for density determination, 7
- Anisotropy, 319
- Anomalous dispersion technique, 216
- Anomalous X-ray scattering (AXS) technique, 245, 254, 265
- Anti-Stokes Raman scattering, 77, 84
- APT. *See* Atom probe tomography (APT)
- Archimedes' principle, and density determination, 3
- sink-float method, 4–5, 7
- wet/dry weighing, 3–4, 7
- Argon laser, for Raman spectroscopy, 81–82
- As-cast glass, 93
- Asymmetry parameter (η), 319, 331
- Atomic-force microscopy (AFM), 392
- Atomic vibrations, effect of, on diffraction, 176–180, 179f
- Atom probe tomography (APT), 391–392
- advancements in, 391–392
- analysis methods, 409–417, 411f
- background and theory, 392–395, 393f, 394f
- case studies, 417–418
- composition, 418–420, 418f, 420f, 421t
- interfaces, 420–424, 422f, 423f
- data reconstruction, 403–404
- mass-spectra analysis, 404–406, 405f
- tomographic reconstruction, 406–409, 407f
- error estimation
- common errors, 416–417
- compositional error, 412–414
- spatial resolution, 414–416, 414f, 415f
- experimental procedure and parameters, 401–403, 404t
- field-ion microscopy (FIM) and, 392–393
- instrumentation, 395–398, 396f, 397f
- limitations of, 392
- local electrode atom probe (LEAP), 392, 395, 396f
- nearest-neighbor analysis, 412
- proximity histogram method, 410–412, 411f
- roughness of buried interfaces, determination of, 412

- Atom probe tomography (APT) (*Continued*)
 specimen preparation, 399–401, 400f,
 401f
 unique feature of, 392
 voltage-pulsed, 391
- Attenuation coefficient, 128
- Auger electron yield, 274
- AX₂ glasses, 75
- Background subtraction. *See also* XAFS
 (X-ray absorption fine structure)
 spectroscopy
 post-edge, 285, 285f
 pre-edge, 284, 285f
- Backscattering, 124, 124f
- Ball and spoke models of glass structure, 75
- Ball-and-stick model, 227, 254
- Barium borosilicate glasses, glass transition
 temperature of, 26–27, 26f–28f
- Bas protocol, 406
- Beamstop alignment, 258
- Beer–Lambert law, 170, 273
- Berberman effect, 32–33
- Bhatia–Thornton formalism, 248–249, 259
- Boltzmann distribution, 318
- Bond-valence (BV), concept of, 220
- Bora (B₂O₃), 8, 9, 10f, 13t, 16, 16f, 17f, 19,
 54, 55, 63, 138–145, 138f, 150f, 151,
 176, 177f, 178f, 181f, 182, 224, 229,
 382
- Borate glasses, infrared spectroscopy and,
 54
 borate network structure, 54–59, 56f–58f
 metal ion-site interactions, 59–63, 60f–62f
- Boron anomaly, 54
- Boron, as shielding material in neutron
 diffractometers, 189
- Borophosphate glasses
 NMR spectroscopy and, 333–336, 334f,
 335f
 speciations and connectivity distributions
 in, 380–386, 381f–385f
- Borosilicate glasses
 barium, 26–27, 26f–28f
 calcium, 27, 27f
 glass transition temperature, 26–27,
 26f–28f
 lithium, 27, 28f
 NMR spectroscopy, 337–339, 338f, 339f
- Boroxol groups, 224
- Bose–Einstein thermal phonon population
 factor, 86–87
- Boson peak, 83, 86, 89, 98–99
- Bragg–Brentano reflection geometry,
 255
 and synchrotron high energy X-ray,
 255–256, 256f
- Bragg's law, 165
- Bridging angle, 75
- Bridging oxygen (BO), 10, 87, 92, 221, 294,
 322, 325, 331, 334, 336, 338, 338f,
 347, 371, 381
- Brillouin light scattering, 108–110
 background and theory, 110–117
 dielectric permittivity/dielectric constant,
 112
 dielectric susceptibility of scattering
 medium, 111
 dynamic structure factor, 115
 intensity of scattered light, 111, 114
 isothermal compressibility of scattering
 medium, 113
 magnitude of electric field, 111
 Pockels coefficient, 114
 scattering volume, 112
 strength of induced polarization density,
 111
 total dipole moment, 111
 wavelengths and propagation direction of
 phonon, 114–115, 115f
- Brillouin spectra, information from,
 108–109
- case studies, 133
 mechanically fragile systems, 151–154,
 152f, 153f
 room-temperature glass, 133–136, 134f,
 134t, 135f
 systems under pressure, 149–151, 149f,
 150f
 temperature dependence, glass transition
 and visco-elasticity, 137–146, 137f,
 142f, 145f
 thin films, 146–149, 148f
- challenge with, 117–118
- data analysis and information, 126–132,
 127f, 131f
- instrumentation for, 117–126, 119f, 121f,
 122f, 124f, 125f

- loss modulus from, 109
- scattering geometries for bulk samples, 124, 124f
- Brillouin peaks, 109
- BrukerVertex 80v vacuum spectrometer, 45–46, 46f
- Bulk modulus, 108, 113, 132, 134t, 137, 139, 153
- Bulk viscosity, 129
- Carr-Purcell-Meiboom-Gill (CPMG) pulse, 332
- CCDs. *See* Charge coupled devices (CCDs)
- Central transition, 320, 323f, 328, 331, 347, 361
- Cesium borate glass system, density of, 17, 17f
- Chalcogenide glasses, 81, 158, 260, 350, 383
 - covalently bonded, 242
 - Raman spectroscopy of, 99–103, 100f–102f
 - X-ray diffraction and, 263–264, 264f
- Charge coupled devices (CCDs), 83–85, 123
- Chemical shielding anisotropy, 319
- Chemical shift, 319, 329–330
- Classical dispersion theory, 38
- Coefficient of finesse, 118, 126
- Coherent anti-Stokes Raman spectroscopy (CARS), 74
- Coherent scattering, 171–173, 174f, 175f
- Coherent scattering length, 172
- Complex modulus, 132
- Compositional error, 412–414
- Concentration profile, 410
- Contrast matching, 203
- Convolution model, 41
- Coordination number, 194
 - for glass, 194–198
- Coordination polyhedra, for glasses, 221–224, 222t, 223f
- Coordination shell, 221, 280–282, 284, 286–290, 292, 294, 299, 304, 306, 310
- Copper molybdo-phosphates, 50–52, 50f, 51t
- Correlation function, in real space, 180–186, 181f, 183f
- Corrosion interface, 417, 420–423
- Cristobalite, 139, 295–297
- Critical cooling rates, 29
- Cross-polarization, 354–358, 355f, 357f
- Cross-section (neutron scattering), 162, 164–166, 169–170. *See also* Neutron diffraction (ND)
- Cryogen-free magnets, 324
- Crystal contamination, neutron diffraction and, 207–209, 208f
- Crystallite models, 218, 219f, 220f
- Cubic spline procedure, 285
- Cumulants method, 283
- Curl-of-curl identity, 113
- Dark currents, 250
- de Broglie wavelength, 161
- Debye equation, 171, 176–177, 180
- Debye–Waller factor, 177, 180, 194, 280
- Deconvolution, 50–51, 126, 127, 327, 331, 333, 334f, 358, 381f, 384f, 406, 409, 419, 424
- Density, 1–2, 19
 - of alkali borates glasses, 16–19, 16f–19f
 - data analysis and useful information, 8
 - extraction of atomic level volumes directly from density, 10, 12f, 13, 13t
 - molar volume, 8, 9f
 - packing fraction, 9–10, 11t, 12f
 - volume per mole glass former, 8–9, 10f
 - defined, 2
 - determination of
 - direct determination of mass and volume, 3
 - general considerations in, 7–8
 - gradient density column, 7
 - instrumentation for, 7
 - pycnometry, 5–6, 5f, 6f
 - sink-float method, 4–5
 - wet/dry weighing, 3–4
 - of lithium silicate glasses, 13–16, 14f, 15f, 16t
 - overview, 2
- Density functional theory (DFT), 255
- Depolarization, of Raman lines, 79–80
- Depolarization ratio (DR), 79–80
- Detector efficiency, 408
- Detector memory effects, 258
- 2D flat plate area detectors, 250
- Diborate, 17, 17f, 19, 62, 63, 66, 67
- Dielectric constant, 33, 35, 112, 114
- Dielectric filter, 82
- Dielectric function, 34–36

- Dielectric permittivity, 112
- Differential pair distribution function, 248
- Differential scanning calorimetry (DSC),
20–22, 21f
modulated, 22
and practical considerations, 22
TA[®] modulated DSC, 23, 24f
thermogram from, 21f
- Differential thermal analysis (DTA), 22–23,
23f, 24f, 25
- Diffraction grating, 85
- Diffractometer, 159, 162, 164–169. *See also*
Neutron diffraction (ND)
- Diiodomethane, 5
- Diode lasers, for Raman systems, 82
- Dipolar interactions, 320
- Dipolar recoupling techniques, 348
- Dipolar solid state NMR spectroscopy,
advanced, 345–346
case studies
borophosphate and thioborophosphate
glasses, 380–386, 381f–385f
connectivity distribution in 70 SiO₂–30
[(Al₂O₃)_x(P₂O₅)_{1-x}] glasses, 374, 376f,
377–380, 378f, 379f
spatial distributions of mobile ions in
alkali silicate and borate glasses,
370–374, 370f, 372f, 373f, 375f
heteronuclear experiments
cross-polarization, 354–358, 355f,
357f
indirect spin–spin interactions,
connectivity studies on, 358, 359f
instrumental considerations and caveats,
358–360
rotational echo adiabatic passage double
resonance, 353–354, 355f
rotational echo double resonance,
350–353, 352f, 354f
spin echo double resonance, 349–350,
351f
homonuclear experiments
dipolar recoupling experiments, 362–369
indirect spin–spin interactions, recovery
of, 367–369, 367f, 368f
instrumental considerations and caveats,
369–370
second moment measurements, 364–367,
365f, 366f
static spin echo decay spectroscopy,
360–362, 360f
two-dimensional correlation methods,
362–364, 363f, 364f
for medium-range glass structure, 345–346
spatial correlations in network glasses, 346f
theory related to, 347–348
direct magnetic dipole–dipole coupling,
348–349
indirect spin–spin coupling, 349
total spin Hamiltonian, 347
- Dipole-dipole interactions, 321, 346–350,
354, 356, 360–362, 369, 371, 374, 385
- Dipole moment, 76, 79, 111, 160, 316
- Direct dipolar coupling, 320
- Dispersion analysis, 38–42, 39f, 40f
- Dispersive spectrometer, 82–83
- Distinct scattering, 172
- DL_POLY package, 228
- Double-null isotopic substitution technique,
209, 214
- Double-quantum (DQ) coherences, 362–367
- DQ-DRENAR (double-quantum-based
dipolar recoupling effects nuclear
alignment reduction), 365, 365f, 366f
- DR. *See* Depolarization ratio (DR)
- DSC. *See* Differential scanning calorimetry
(DSC)
- DTA. *See* Differential thermal analysis (DTA)
- Dynamic structure factor, 115
- Dynamic viscosity, 128
- Dy₇Ni₃, 202f, 203, 210f, 211, 212f, 213, 214
- EBS. *See* Electrostatic bond strength (EBS)
- Edge filters, 84
- Eight minus *n*-rule, 263
- Elastic scattering, 108
- Elastic storage modulus, 128
- Elasto-optic coefficients, 110, 114
- Electric susceptibility tensor, 78
- Electrostatic bond strength (EBS), 219
- Empirical potential structure refinement
(EPSR), 228, 229, 254–255
- Energy discriminating solid state detectors,
250
- Epoxy cure, 152–153, 153f
- Equation of continuity, 115
- EXAFS. *See* Extended X-ray absorption fine
structure (EXAFS)

- Excitation wavelength, 81
- Extended range ordering, 249
- Extended X-ray absorption fine structure (EXAFS), 254, 273, 279–282, 280f, 281f. *See also* XAFS (X-ray absorption fine structure) spectroscopy
- Faber–Ziman formalism, 247, 248
- Faber–Ziman partial structure factors, 179
- Fabry–Perot cavity, 118–120, 119f, 122, 122f
- Fabry–Perot interferometer, 118–122, 119f, 121f, 122f
- Fast-ion-conducting glasses, 307
- FDMNES (Finite Difference Method Near-Edge Simulation) software, 296
- Fermi's Golden rule, 278
- Fictive temperature, 25, 33, 241
- Field desorption, 393
- Field-ion microscopy (FIM), 391. *See also* Atom probe tomography (APT)
- Filon's quadrature, 185
- Finesse, 120
- First difference, isotopic substitution, 211
- First sharp diffraction peak (FSDP), 224, 226, 251, 252, 258, 259, 263, 264f
- Fitting procedure, 288–289
- Focused ion beam (FIB), 391, 395, 399–401
- Focus software, for curve-fitting, 42
- Form factor, 160, 171, 186, 211, 215, 245–246, 245f, 246f, 252, 260, 265, 266f
- Four-coordinated boron (N_4 ; f_2), 16, 16f, 26–27, 26f, 57, 58f, 335, 335f
- Fourier artifacts, 253
- Fourier noise, 248
- Fourier's second law, 116
- Fourier-transform infrared (FT-IR) spectrometer, 44–48. *See also* Infrared (IR) spectroscopy
- Fourier-transform (FT) Raman spectrometer, 83
- Free-induction decay (FID), 325, 326f, 327
- Fresnel formulae, 37
- FSDP. *See* First sharp diffraction peak (FSDP)
- FT-IR spectrometer. *See* Fourier-transform infrared (FT-IR) spectrometer
- Full width at half maximum (FWHM), 82
- Gas ionization chambers, 275
- Gas lasers, for Raman spectroscopy, 81–82
- Gas pycnometry, 5–6
- Gaussian broadening, 41
- Gaussian dispersion analysis, 41–42
- Gels, 151–152, 264
- Generalized hydrodynamic approach, 115
- General Materials diffractometer (GEM), 168–169, 168f
- Germania (GeO_2), 297–299, 298f–300f
- Germanosilicate glasses, Raman study of, 96–99, 97f–99f
- $GeSe_2$, 214, 246f, 247f, 251f, 259, 263, 265f
- GeS_2 – Sb_2S_3 , 100–101, 100f, 101f
- Glass ceramics, 96–99, 304, 367, 383
- Glass-in-glass phase separation phenomena, 92. *See also* Raman spectroscopy nucleation phenomenon, 92 spinodal decomposition, 92
- Glass stability parameters, 28t
- Glass transition temperature, 2, 54, 55, 62, 62f, 68, 139, 140f, 141, 147, 151, 201, 248
- of barium borosilicate glasses, 26–27, 26f–28f
- determination of, 20 (*see also* Thermal properties)
- Glitches, 284
- Goldschmidt radii, 219
- Gradient density column method, for density determination, 7
- Group theory, 76
- Hamiltonian, 346–349, 354, 361, 362, 364, 365, 370
- Hartmann–Hahn matching condition, 356, 358
- Heteronuclear correlation experiments, 354–358, 355f, 357f
- Hooke's law, 133
- Hydrogen contamination, neutron diffraction and, 206–207, 207f
- Image compression factor, 408
- INADEQUATE experiment, 368f, 369
- Incoherent scattering, 171–173, 174f, 175f
- Indirect spin–spin coupling, 320
- indirect spin–spin coupling tensor, 349

- Inelastic neutron scattering, 74
- Inelastic scattering, 108
- Infrared (IR) spectroscopy, 32–34
 - applications in glasses, 32
 - assessment of glass films by, 32–33
 - bulk materials, reflectance spectroscopy of, 36–37
 - dispersion analysis, 38–42, 39f, 40f
 - Kramers–Krönig transformation, 37–38
- case studies
 - bulk glasses, 54–63
 - glass thin films, 63–67
- data analysis
 - bulk glasses, 48–52, 49f, 50f, 51t
 - thin films of amorphous materials, 52–54, 52f, 53f
- instrumentation for, 44–48
- refractive index and dielectric function, 34–36
- short-range order structure and, 32
- thin films, infrared spectra of, 42–44
- Institut Laue-Langevin (ILL), 162
- Instrumental errors, 284
- Interference fringes, 42–43
- Interferograms, 45–47, 47f
- Intermediate range order (IRO), 17, 20, 76, 89, 100, 158, 226, 228, 229, 243, 258, 263, 264, 271, 299
- Inverse methods, 185
- Ion-conducting channels, 371
- Ion coordination, 11t
- Ionization, 275, 275f, 392–396, 393f, 394f, 399, 402, 403, 406, 407, 413, 416, 419
- Isoconcentration surfaces, 410
- Isomorphic substitution, 254
- Isomorphous substitution, 215
- Isothermal modulus, 154
- Isotopic incoherence, 171
- Isotopic substitution, 209–214, 210f, 212f
- Jahn–Teller effect, 197, 223
- J-resolved spectroscopy, 367–369, 367f
- Kinematic viscosity, 116
- Klein–Nishina formula, 250
- Kramers–Krönig transformation, 37–38, 48, 145
- Krogh-Moe/Norman technique, 251
- Kronecker delta, 194
- k-space, 281, 286–287
- Lambert's law, 36
- Lamb modes, 146–147
- Landau–Placzek ratio, 116
- Larmor frequency, 318, 325
- Laser monochromator, 82
- Lasers, for Raman spectroscopy, 81–82
- Lever rule, 14, 15f
- Lift out methods, 399
- Light scattering, 107. *See also* Brillouin light scattering
 - thermal phonons and, 108
- Light sources, for Raman spectroscopy, 81–82
- Linear elasticity, constants of, 134t
- Lithium borate glasses
 - density of, 16, 16f
 - stability parameters in, 27–30, 28t, 29f, 30t
- Lithium borates, 10f, 16, 16f, 25, 25f, 27–30, 30t, 371, 372f
- Lithium diborate ($\text{Li}_2\text{O}-2\text{B}_2\text{O}_3$), 63–66, 64f–66f
- Lithium silicate glasses
 - density of, 13–16
 - lever rule, 14, 15f
 - Q^i structural units, 14, 14f
 - sink-float method, 13, 14f
 - volume per mole silica, 15, 15f
 - volumes of silicate units, 16, 16t
- Lithium silicates, 10f, 13–16, 14f, 15f, 350, 371, 372f, 373f
- Local electrode atom probe (LEAP), 392, 395, 396f
- Local magnification effect, 414–415, 415f
- Longitudinal modulus, 133
- Longitudinal optical (LO), 33, 34, 40, 41
- Longitudinal viscosity, 129
- Lorch function, 190, 192
- LTCP glass, phase transformations in, 92–96, 93f–95f
- Lyddane–Sachs–Teller relationship, 41
- Magic-angle spinning (MAS), 315, 320–323, 321f–323f, 347. *See also* Nuclear magnetic resonance (NMR) spectroscopy

- Magnet, 323–325
Magnetic shielding, 318–319, 347
Magnetization, 112, 318, 324, 325, 328f, 354, 356, 358, 362–363, 369
Magnetogyric ratio, 317, 324, 332
Magnification, of point projection, 407–408
MAS. *See* Magic-angle spinning (MAS)
Mass spectrum, 404–406
Maxwell's equations, 112
Maxwell's relation, 34
Mean free path (MFP), of photoelectron, 280
Measurement uncertainties, in
 NMR spectroscopy, 327–329, 328f
Mechanical loss modulus, 128
Medium range structure, 345
Mesostructure, 345
Michelson interferometer, 44. *See also*
 Infrared (IR) spectroscopy
Microchannel plate (MCP) detector, 397
Micro-Raman spectrometers, 80f, 84–85
Mode coupling theory, 115
Modeling techniques, 226–229
Moderator, 162
Modification function, 190, 192
Modified Atomic Form Factor (MAFF), 246
Modifier cations, in alkali borate glasses,
 340–341, 340f, 341f
Modulated differential scanning calorimetry
 (MDSC), 22
Molar volumes, 8
 of alkali and alkaline-earth borate glass
 system, 9f
 of alkali borates, 17, 18f
 stiffness and, 18–19, 19f
Molecular dynamics (MD), 2, 57, 151, 228,
 255, 260, 284, 291, 293, 294, 297, 371
Momentum transfer, 165
Monochromator, 44, 81–84, 86, 87, 166, 168,
 194, 249, 256f, 276, 284
Mott–Bethe formula, 245
Multi-channel detectors, 83–84
Multiple quantum magic angle spinning
 (MQMAS), 321, 331, 337–339
Multiple scattering (MS), 186, 187, 250, 258,
 278, 279f, 288, 290, 294, 299, 304, 310

Nanostructure, 345. *See also* Dipolar solid
 state NMR spectroscopy, advanced
Navier–Stokes equation, 116

NBO. *See* Non-bridging oxygen (NBO)
ND. *See* Neutron diffraction (ND)
Near and Intermediate Range Order
 Diffractometer (NIMROD), 203, 207
Nearest-neighbor analysis, 412
Network-modifying cations, 340–341
Neutron diffraction (ND)
 assembly of nuclei, scattering from,
 170–171
 atomic vibrations
 and diffraction, 176–180, 179f
 effect of inelasticity on self scattering,
 173–176, 176f–178f
 coherent and incoherent scattering,
 171–173, 174f, 175f
 experimental corrections, 186–189, 188f,
 189t
 inelastic scattering processes, 161
 interactions between neutron and sample,
 160–161, 161f
 intermediate range order (IRO) and, 158,
 226, 228, 229
 isotropic samples, 171
 modeling techniques, 226–229
 neutron diffractometer
 general principles, 164–166, 166f
 pulsed source diffractometers, 167–169,
 167f, 168f
 reactor source diffractometers, 166–167,
 167f
 neutrons and, 159, 159t
 normalization of data, 198–200, 199f
 partial correlation functions
 isotopic substitution, 209–214, 210f,
 212f
 other methods, 214–216, 216f, 217f
PDF (pair distribution function) method,
 229–231
peak fitting and integration, 194–198, 196f,
 197t
real-space correlation functions, 180–186,
 181f, 183f
real-space resolution, 190–194, 191f,
 193f
reciprocal-space resolution, 190
results, interpretation of, 218
 bonding and bond lengths, 218–221
 comparison with crystalline structures,
 224, 225f

Neutron diffraction (ND) (*Continued*)

- coordination polyhedra, 221–224, 222t, 223f
- crystallite models, 218, 219f, 220f
- first sharp diffraction peak, 224, 226
- sample-related difficulties, 203
- crystal contamination, 207–209, 208f
- hydrogen contamination, 206–207, 207f
- nuclear resonances, 204–206, 204t, 205f
- scattering at low Q
 - homogeneous samples, 200–201
 - inhomogeneous samples, 201–203, 202f
- short range order (SRO) and, 158, 226, 227

single nucleus, scattering from, 169–170

sources of neutron

- accelerator sources, 162–164, 163f
- reactor sources, 161–162
- static approximation, 169
- for structural studies of glasses, 158
- and x-ray diffraction, 171, 173

Neutron diffractometer, 162. *See also*

Neutron diffraction (ND)

Niobium germanosilicate glasses, Raman study of, 96–99, 97f–99f

NMR spectroscopy, 354. *See* Nuclear magnetic resonance (NMR) spectroscopy

Non-bridging oxygen (NBO)

- alkali incorporation in silicate glasses and, 87

Raman spectroscopy for, 75

Normalization, 198–200, 286

Notch filters, 84

Nuclear fission, 159, 161–162

Nuclear magnetic resonance (NMR)

spectroscopy, 315–316

background and theory

- dipolar interactions, 320
- high resolution methodologies, 320–323, 321f–323f
- magnetic shielding, 318–319
- quadrupolar interaction, 319–320
- Zeeman effect, 316–318, 316f, 317t, 318f

case studies, 333

- aluminosilicate glasses, 336–337, 337f
- borophosphate glasses, 333–336, 334f, 335f

- borosilicate glasses, 337–339, 338f, 339f
- modifier cations in alkali borate glasses, 340–341, 340f, 341f

data analysis and structural interpretation, 329

chemical shift assignments, 329–330

low- γ nuclei, 332

paramagnetic effects, 333

quadrupolar effects, 330–332

instrumentation, 323

computer control, 326–327

magnet, 323–325

measurement uncertainty, 327–329, 328f

probe, 325–326

radiofrequency transmitter and receiver, 326

nuclear properties of selected nuclides, 317t

spin description of NMR experiment, 318f

Nuclear resonances, 204–206, 204t, 205f

Null technique, 209

Nutation frequencies, 326

Nyquist sampling theorem, 289

Packing fraction, 9–10

of alkali and alkaline-earth borates, 12f

ion coordination and size, 11t

Pair distribution function (PDF), 183, 229–231

Paramagnetic effect, 333

Paramagnetic ions, 333

Partial correlation function, 209–218, 210f, 212f, 216f, 217f

PDFfit program, 231

Peak fitting, 86, 90f, 91, 92, 101, 185, 194–198, 248, 250, 282, 283, 330

Percus–Yevick equation, 211

Percus–Yevick model, 254

Phase separation mechanisms, in phosphate glasses, 92–96, 93f–95f

Phonons, 33, 78, 86, 108–109, 114, 117–118, 123, 124, 125f, 126, 133, 136, 146–148

Phosphate glasses, 92–96, 192, 195, 197, 215, 223, 347, 362, 366f, 369

Photoluminescence yield mode (PLY), 275

Photomultiplier tube (PMT), 83

Placzek correction, 175–176

PLAP. *See* Pulsed laser atom probe (PLAP)

Platelet scattering geometry, 125–126, 125f

PMT. *See* Photomultiplier tube (PMT)

- Pockels elasto-optic coefficients, 114
- Poisson ratio, 108, 134t, 143, 144, 153, 153f, 154
- Polarizability, 35, 76, 78–79, 96, 111, 112
- Polymer-cellulose, 148, 148f
- Porod approximation, 203
- Potassium diborate ($K_2O-2B_2O_3$), 48, 49f
- Potassium silicates, 91f, 92, 140f
- Potassium tellurites, 145, 145f
- Powder pattern, 32, 33, 63, 165, 168, 187, 190, 201, 218, 241, 249, 257, 307, 321f, 347, 399
- π pulse, 350, 353, 363f, 367f
- Pressure, 5, 6, 110, 149–151, 149f, 150f, 297–299
- Pressure-induced amorphization, 264
- Probe, NMR, 325–326
- Proximity histogram, 410–412, 411f
- Pulsed laser atom probe (PLAP), 394–395
- Pulsed laser deposition (PLD), 63
- Pulsed source diffractometers, 167–169, 167f, 168f
- Pulse fraction, 396
- Pycnometry, for density determination, 5–6, 5f, 6f
- advantages of, 6
 - disadvantages of, 6
 - helium, use of, 5
 - instrumentation for, 7
- Quadrupolar asymmetry parameter, 331
- Quadrupolar coupling constant, 319, 330
- Quadrupolar interactions, 319–320, 347
- Quadrupolar product, 331
- Quadrupole moment, 319, 331, 332, 347
- Quantachrome[®] manual pycnometer, 6f
- Q units, 10, 14–15, 15f, 16t
- Q-value, 200–203, 205, 224, 245–247, 250, 265
- Radial distribution function, 248
- Radial distribution function (RDF), 282
- of monatomic system, 180
- Radio frequency driven recoupling (RFDR), 362, 363f
- Raman effect, 75, 76. *See also* Raman spectroscopy
- Raman lines, depolarization of, 79–80
- Raman shift, 75
- Raman spectroscopy, 74–76
- anti-Stokes Raman effect, 77
 - Boson peak, 83, 86
 - case studies on, 87
 - alkali incorporation in silicate glasses, structural effects of, 87–92, 88f, 90f, 91f
 - germanosilicate glasses and glass ceramics, 96–99, 97f–99f
 - phase separation mechanisms in phosphate glasses, 92–96, 93f–95f
 - Raman study of chalcogenide glasses, 99–103, 100f–102f
 - data analysis, 86–87
 - depolarization of Raman lines, 79–80
 - elastic Rayleigh scattering, 77–78
 - electric susceptibility tensor, 78
 - instrumentation, 80–81, 80f
 - detector, 83–84
 - light sources, 81–82
 - micro-Raman spectrometers, 84–85
 - resolution, 85–86
 - sample compartment, 82
 - spectrometers, 82–83
 - laser beam for, 76
 - linearly induced polarization, 78
 - Raman effect, 75, 76
 - Raman shift, 77
 - selection rules for, 78–79
 - spontaneous, mechanisms of, 77, 77f
 - theory, 76–78
 - use of, 75
- Random network theory, of glasses, 242
- Zachariasen–Wright network model of glass, 242, 243f
- Range File, 406
- Rare earths, 206, 215, 265, 303–307
- Rayleigh scattering, 77–78
- Reactor neutron sources, 161–162
- Reactor source diffractometers, 166–167, 167f
- Real-space correlation functions, 180–186, 181f, 183f
- Real-space resolution, 190–194, 191f, 193f
- REAPDOR. *See* Rotational echo adiabatic passage double resonance (REAPDOR)
- Reciprocal-space resolution, 190

- REDOR. *See* Rotational echo double resonance (REDOR)
- Reflectance spectroscopy, of bulk materials, 36–42
- Refocused INADEQUATE method, 368f, 369, 380, 384, 385f
- Refractive index, 34–36
- Region of interest (ROI), 83, 327, 399, 401f
- Relaxational modulus, 129
- Renishaw Raman Imaging Microprobe System 3000, 92
- Resolution, neutron scattering, 190–194
- Resolution, of Raman system, 85–86
- Retardation, 45
- Reverse Monte Carlo (RMC) method, 227–229, 254–255
- Root-mean-squared (RMS) roughness, 412
- Rotational echo adiabatic passage double resonance (REAPDOR), 353–354, 355f
- Rotational echo double resonance (REDOR), 350–353, 352f
constant time (CT-REDOR), 353, 354f
- R-space, 281, 286–287
- R7T7 glass, 418
- SANS. *See* Small angle neutron scattering (SANS)
- Satellite transitions, 320, 328
- 90°-scattering, 124, 124f
- Scattering length of nucleus, 160
- Scherrer equation, 218
- Secondary ion mass spectrometry (SIMS), 392
- Second difference, isotopic substitution, 213
- Second moment, 348
- SEDOR. *See* Spin echo double resonance (SEDOR)
- Selection rules, for Raman spectroscopy of glasses, 78–79
- Self scattering, 172
effect of inelasticity on, 173–176, 176f–178f
- Shear modulus, 133
- Short range order (SRO), 13t, 32, 34, 54, 56, 59, 67, 68, 76, 158, 226, 227, 244, 263, 271, 290, 316, 329, 333, 345. *See also* Nuclear magnetic resonance (NMR) spectroscopy
- Silica (SiO_2), 13, 15, 15f, 16, 34t, 47, 87–89, 97, 100, 136, 258–263, 294–297, 398
- Silica aluminophosphates, 366, 374–380
- Silicate glasses, alkali incorporation in, structural effects of, 87–92, 88f, 90f, 91f
- Silver borates, 283, 307, 381
- SIMPSON program, 370
- Single-mode diode lasers, 82
- Single scattering (SS), 278, 279f, 282, 288
- Sink-float method, for density determination, 4
acetone-diiodomethane solution, use of, 5
advantages of, 4–5
disadvantages of, 5
instrumentation for, 7
lithium silicate glasses and, 13, 14f
- Small Angle Neutron Diffractometer for Amorphous and Liquid Samples (SANDALS), 206–207
- Small angle neutron scattering (SANS), 201–203
- Small angle X-ray scattering (SAXS), 265
- Snell's law, 125–126
- Soda lime silicates, 301–303
- Sodium borates, 136, 137f, 140f, 336, 370f, 371, 374, 375f
- Sodium-calcium silicates, 292
- Sodium metasilicate glass, polarized Raman spectra of, 88, 88f
- Sodium phosphate, 366
- Sodium silicates, 87, 293, 370–371, 370f, 374
- Sodium tetrasilicate glass, Raman spectra of, 76
- SON68 glass, 418–420, 418f, 421t, 423, 424
- Soper's GudrunN program, 187
- Spallation, 164
- Spatial resolution
of APT, 414–416, 414f, 415f
of Raman spectrometers, 85–86
- Spectrometers, 82
dispersive, 82–83
Fourier-transform, 83
- Spin echo decay method, 360–262, 360f
- Spin echo double resonance (SEDOR), 349–350, 351f
- Spin incoherence, 171
- Spin-lattice relaxation times, 328
- Spinning sidebands, 321, 322f, 328, 332, 334f

- Splat-quenched glass samples, 54
Splice method, 283
Spontaneous Raman effect, 77, 77f. *See also* Raman spectroscopy
Static approximation, 169, 174
Static spin echo decay spectroscopy, 360–262, 360f
Stiffness, and molar volumes, 18–19, 18f, 19f
Stimulated Raman scattering (SRS), 74
Stokes effect, 75, 77. *See also* Brillouin light scattering; Raman spectroscopy
Structure factor, 114–116, 165, 177, 179, 201, 243–245, 247, 247f, 249, 252–254, 256–260, 263–265
Superionic conducting glasses, ionic conductivity in, doped with AgI, 307, 308f, 309f
Surface-enhanced Raman spectroscopy (SERS), 74

Tandem Fabry–Pérot interferometer, 121–122, 122f
Tandem interferometry, 120–122, 122f
Terbium phospho-aluminate, 229
Terminal oxygens (TOs), 193
Thermal diffusivity, 128
Thermal properties, 20, 30. *See also* Glass transition temperature
 data analysis and useful information, 25, 25f, 26f
 experimental methods and theory
 calorimetry, 20–22, 21f, 24f
 differential thermal analysis, 22–23, 23f, 24f
 instrumentation used, 23–25, 24f
 stability parameters in lithium borate glasses, 27–30, 28t, 29f, 30t
Thin films, 33–34, 36
 of amorphous materials, 52–54, 52f, 53f
 Brillouin light scattering and, 146–149, 148f
 infrared spectra of, 42–44
 infrared spectroscopy and, 63–67
Thioborophosphate glasses, speciations and connectivity distributions in, 380–386, 381f–385f
Time-of-flight mass spectrometry (TOF-MS), 391. *See also* Atom probe tomography (APT)

Time-of-flight neutron diffractometer, 167–168, 167f
Time-of-flight (T-O-F) technique, 164
Tin borate glass, 208, 208f
Tip-enhanced Raman spectroscopy (TERS), 74
Ti₆₂Zr₃₈ alloy, 187
Tomographic reconstruction, 406–409
Total electron yield (TEY), 274, 277–278
Total pair distribution function, 248
Transition energy, 317
Transmission electron microscopy (TEM), 392
Transmittance, 36, 42
Transverse optical (TO), 33, 34, 40–41
Triborate, 17, 17f, 19
Turing's number, 213

Ultralow frequency filters, 84
Uniqueness problem, 227
UO₂, 230
U₄O₉, 225f, 230–231

Vanadium, role in neutron scattering, 187–188
Van Vleck formulae, 348
Vibrational density of states (VDOS), 75
Vibrational spectroscopy, 32, 74. *See also* Infrared (IR) spectroscopy
Virtual image phased array (VIPA) spectrometer, 122–123
Vitreous silica, 46, 47f, 48, 89
Voigt model, 42
Volume per mole glass former, 8–9
 in lithium borate glass system, 10f
 in lithium germanate glass system, 10f
 in lithium silicate glass system, 10f

Wavefunction, 79, 160, 170, 278, 290
Waveguides, 304, 305f, 305t, 306f
Wet/dry weighing method, for density determination, 3
 advantages of, 3–4
 disadvantages of, 4
 instrumentation for, 7
Wide angle neutron scattering (WANS), 203
Wide angle X-ray scattering (WAXS), 265
Wideband uniform rate smooth truncation (WURST) pulse, 327, 332

- XAFS (X-ray absorption fine structure)
 spectroscopy, 272
 data analysis, 284
 conversion to k -space, threshold energy and weighting, 286
 corrections for instrumental errors, 284
 fitting procedures, 288–289
 Fourier filtering, 287, 287f
 log amplitude ratio and phases difference method, 288
 normalization, 286
 post-edge background subtraction, 285, 285f
 pre-edge background subtraction, 284, 285f
 transformation from k -space to R -space, 286–287
- EXAFS, 273, 279–282, 280f, 281f
 accuracy and limitations, 289–290
 cumulants, 283
 effective distribution, 283
 formula for glasses, 282–284
 radial distribution function, 282
 single-shell EXAFS, 282
- for glass structure investigations, 272, 291–292
- Ag nanoparticles in soda-lime silicate glasses, 301–303, 302f–304f
- ionic conductivity in superionic glasses doped with AgI, 307, 308f, 309f
- nanoparticles embedded in glasses, 300–307
- rare-earth impurities dispersed in glassy matrices, 303–307, 305f, 305t, 306f
- silica and germania glasses under high pressure, 297–299, 298f–301f
- silica at high temperature, 294–297, 295f, 296f
- silica glass, 294, 295f
- silicate glasses, 292–294, 293f
- instrumentation, 274–278
 apparatus for transmission measurements, 275, 275f
 detection modes, 275f
 direct measurements, 274
 duration of measurements, 276
 Energy Dispersive-XAFS spectrometers, 276, 276f
 fluorescence detection mode, 276–277, 277f
 indirect measurements, 274–275
 Quick-XAFS apparatuses, 276
 scanning transmission mode, 276
 synchrotron beamline for XAFS measurements, 274, 274f
 total electron yield detection mode (TEY), 277–278
 physical mechanism of, 278
 multiple scattering, 278, 279f
 single scattering, 278, 279f
 theory, 272–274, 273f, 274f
 XANES, 273, 290–291, 291f
 X-ray absorption spectra, origins of, 272–274
- XANES. *See* X-ray absorption near edge structure (XANES)
- X-ray Absorption Fine Structure (XAFS), 213, 230–231
- X-ray absorption near edge structure (XANES), 273, 290–291, 291f, 354.
See also XAFS (X-ray absorption fine structure) spectroscopy
- X-ray absorption spectroscopy (XAS), 271–272. *See also* XAFS (X-ray absorption fine structure) spectroscopy
- X-ray diffraction, 241–244
 background and theory, 244–249, 245f–247f
 average scattering, 246–247
 Bhatia–Thornton formalism, 248–249
 differential pair distribution function, 248
 electron distribution function, 246
 electron form factors, 245
 Modified Atomic Form Factor, 246
 pseudo-atomic function, 246
 pseudo-nuclear function, 247
 radial distribution function, 248
 sharpening function, 246
 total pair distribution function, 248
 X-ray form factors, 245
- case studies
 amorphous materials, gels, foams and fibers, 264
 chalcogenide glasses, 263–264, 264f
 SiO₂ and oxide glasses, 258–263, 259f, 261f, 262f

- data analysis and useful information, 249–255, 251f–253f
 - beam polarization effects, 249
 - bond angle information, 255
 - consistency checks, 251
 - detector corrections, 250
 - first sharp diffraction peak (FSDP), 251–252
 - orientational correlation functions, 255
 - Q*-dependent corrections, 250
 - ring size distribution, 255
 - sample-dependent absorption corrections, 249–250
 - X-ray PDF function, 254
- future perspective, 265–266, 266f
- instrumentation, 255–258, 256f, 257f, 258t
- and neutron diffraction, 244, 254
- X-ray pair distribution function, 243–244, 244f
- X-ray PDF technique, 242–244
- X-ray diffraction (XRD), 271–272
- X-ray excited optical luminescence (XEOL) spectra, 278
- X-ray fluorescence yield (FLY), 274, 276–277, 277f
- X-ray free electron lasers (X-FEL), 266
- X-ray photoemission spectroscopy (XPS), 392
- XTAL (computer program), 224, 230
- Young's modulus, 108, 134t, 153, 154
- Yttria (Y_2O_3), 199–200, 199f
- Zeeman effect, 316–318, 316f, 317t, 318f
- Zero-quantum (ZQ) coherences, 362–364