



Vol. 1216, issue 16, 17 April 2009

## CONTENTS



Abstracted/indexed in: Analytical Abstracts, Biochemical Abstracts, Biological Abstracts, Chemical Abstracts, Chemical Titles, Chromatography Abstracts, Elsevier BIOBASE/Current Awareness in Biological Sciences, Current Contents/Life Sciences, Current Contents/ Physical, Chemical & Earth Sciences, Deep-Sea Research/Part B: Oceanographic Literature Review, EMBASE/Excerpta Medica, Index Medicus, Mass Spectrometry Bulletin, PASCAL-CNRS, Referativnyi Zhurnal, Research Alert and Science Citation Index. Also covered in the abstract and citation database SCOPUS®. Full text available on ScienceDirect®.

### 32ND INTERNATIONAL SYMPOSIUM ON HIGH PERFORMANCE LIQUID PHASE SEPARATIONS AND RELATED TECHNIQUES, BALTIMORE, MD, USA, 10-16 MAY 2008

Foreword	
by G. Guiochon (Knoxville, TN, USA) and S.C. Jacobson (Bloomington, IN, USA)	3421
Ultraviolet thermal lensing detection of amino acids	
by F. Yu (Stanford, CA, USA), A.A. Kachanov (Stanford and Santa Clara, CA, USA), S. Koulikov, A. Wainright (Santa Clara, CA, USA) and R.N. Zare (Stanford, CA, USA)	3423
Effect of analyte properties on the kinetic performance of liquid chromatographic separations	
by A. de Villiers (Matieland, South Africa), F. Lynen and P. Sandra (Ghent, Belgium)	3431
Optimization of two-dimensional gradient liquid chromatography separations	
by P. Česla, T. Hájek and P. Jandera (Pardubice, Czech Republic)	3443
Smart Templates for peak pattern matching with comprehensive two-dimensional liquid chromatography	
by S.E. Reichenbach (Lincoln, NE, USA), P.W. Carr, D.R. Stoll (Minneapolis, MN, USA) and Q. Tao (Lincoln, NE, USA)	3458
Selecting an "orthogonal" column during high-performance liquid chromatographic method development for samples that may contain non-ionized solutes	
by J.W. Dolan and L.R. Snyder (Amity, OR, USA)	3467
Statistical optimization of the silylation reaction of a mercaptosilane with silanol groups on the surface of silica gel	
by R. Dabre (Darmstadt, Germany and Vienna, Austria), A. Schwämmle (Darmstadt, Germany), M. Lämmerhofer and W. Lindner (Vienna, Austria)	3473
Adsorption behaviour of a quinidine carbamate-based chiral stationary phase: Role of the additive	
by R. Arnell (Södertälje, Sweden), P. Forssén, T. Fornstedt (Uppsala, Sweden), R. Sardella, M. Lämmerhofer and W. Lindner (Vienna, Austria)	3480
Effect of microemulsion component purity on the chromatographic figures of merit in chiral microemulsion electrokinetic chromatography	
by A.B. Kojtari and J.P. Foley (Philadelphia, PA, USA)	3488
Evaluation of alternatives to warfarin as probes for Sudlow site I of human serum albumin. Characterization by high-performance affinity chromatography	
by K.S. Joseph, A.C. Moser, S.B.G. Basiaga, J.E. Schiel and D.S. Hage (Lincoln, NE, USA)	3492
Measurement of free hydroxytyrosol in microdialysates from blood and brain of anesthetized rats by liquid chromatography with fluorescence detection	
by Y.-T. Wu, L.-C. Lin and T.-H. Tsai (Taipei, Taiwan)	3501
Determination of steroid sex hormones in wastewater by stir bar sorptive extraction based on poly(vinylpyridine-ethylene dimethacrylate) monolithic material and liquid chromatographic analysis	
by X. Huang, J. Lin, D. Yuan and R. Hu (Xiamen, China)	3508
Using sweeping micellar electrokinetic chromatography to analyze $\Delta^9$ -tetrahydrocannabinol and its major metabolites	
by H.-L. Su, L.-I. Feng, H.-P. Jen and Y.-Z. Hsieh (Hsinchu, Taiwan)	3512
Strategy for the isolation of native dehydrogenases with potential for biosensor development from the organism <i>Hyphomicrobium zavarzinii</i> ZV580	
by F. Hilbrig, V. Jérôme, M. Salzig and R. Freitag (Bayreuth, Germany)	3518
Quantitative determination of fenoterol and fenoterol derivatives in rat plasma using on-line immunoextraction and liquid chromatography/mass spectrometry	
by H.S. Kim, D. Siluk and I.W. Wainer (Baltimore, MD, USA)	3526
Determination of free and protein-bound glutathione in HepG2 cells using capillary electrophoresis with laser-induced fluorescence detection	
by Y. Wang, Y. Xie, M. Bernier and I.W. Wainer (Baltimore, MD, USA)	3533
Study of the human plasma proteome of rheumatoid arthritis	
by X. Zheng, S.-I. Wu, M. Hincapie and W.S. Hancock (Boston, MA, USA)	3538
Screening and monitoring antiretrovirals and antivirals in the serum of acquired immunodeficiency syndrome patients by micellar liquid chromatography	
by M.A. Raviolo (Córdoba, Argentina), I.C. Brea (Vila-real, Spain) and J. Esteve-Romero (Castelló, Spain)	3546
On-line separation of native proteins by two-dimensional liquid chromatography using a single column	
by X. Geng, C. Ke, G. Chen, P. Liu, F. Wang, H. Zhang and X. Sun (Xi'an, China)	3553
Determination of arylamines and aminopyridines in pharmaceutical products using in-situ derivatization and liquid chromatography-mass spectrometry	
by G. Vanhoenacker, E. Dumont, F. David (Kortrijk, Belgium), A. Baker (Loughborough, UK) and P. Sandra (Kortrijk, Belgium)	3563

(Continued overleaf)

Contents (continued)

Thermodynamic investigations on shape selective separation behaviors of poly(4-vinylpyridine)-grafted silica for polycyclic aromatic hydrocarbons in both normal-phase and reversed-phase high-performance liquid chromatography by U.G. Gautam, M.P. Gautam, T. Sawada, M. Takafuji and H. Ihara (Kumamoto, Japan) .....	3571
Methylammonium formate as a mobile phase modifier for totally aqueous reversed-phase liquid chromatography by S. Grossman and N.D. Danielson (Oxford, OH, USA) .....	3578
C <sub>18</sub> , C <sub>8</sub> , and perfluoro reversed phases on diamond for solid-phase extraction by G. Saini, L.A. Wiest, D. Herbert, K.N. Biggs (Provo, UT, USA), A. Dadson, M.A. Vail (Orem, UT, USA) and M.R. Linford (Provo, UT, USA) .....	3587
Using subcritical/supercritical fluid chromatography to separate acidic, basic, and neutral compounds over an ionic liquid-functionalized stationary phase by F.-M. Chou, W.-T. Wang and G.-T. Wei (Min-Hsiung, Taiwan) .....	3594