



ELSEVIER

available at www.sciencedirect.comjournal homepage: www.elsevier.com/locate/aca

Period V.637 Issue 1-2

Contact "Library Services": info@dss.go.th

CONTENTS

(Full texts are incorporated in CJELSEVIER, a file in the Chemical Journals Online database available on STN International; Abstracted, indexed in: Aluminium Abstracts; Anal. Abstr.; Biol. Abstr.; BIOSIS; Chem. Abstr.; Curr. Contents Phys. Chem. Earth Sci.; Engineered Materials Abstracts; Excerpta Medica; Index Med.; Life Sci.; Mass Spectrum. Bull.; Material Business Alerts; Metals Abstracts; Sci. Citation Index). Also covered in the abstract and citation database SCOPUS®. Full text available on ScienceDirect®.

Special Issue:**Papers presented at EuroResidue VI 2008****Preface**

H.F. De Brabander, L.A. Van Ginkel, S. Sterk and A. Bergwerff (Bilthoven, The Netherlands) 1

Review article

Analysis of thyreostats: A history of 35 years
J. Vanden Bussche, H. Noppe, K. Verheyden, K. Wille (Merelbeke, Belgium), G. Pinel, B. Le Bizec (Nantes, France) and
H.F. De Brabander (Merelbeke, Belgium) 2

Chemometrics

Risks of antibiotic residues in milk following intramammary and intramuscular treatments in dairy sheep
A. Pengov and A. Kirbis (Ljubljana, Slovenia) 13

Extraction and Sample Handling

Development and validation of a confirmatory method for the determination of sulphonamides in milk by liquid chromatography with diode array detection
V. Gamba, C. Terzano (Brescia, Italy), L. Fioroni, S. Moretti (Perugia, Italy), G. Dusi (Brescia, Italy) and R. Galarini (Perugia, Italy) 18

Development, validation and implementation of a receptor based bioassay capable of detecting a broad range of β -agonist drugs in animal feedingstuffs
S. Boyd (Belfast, Northern Ireland), H.H. Heskamp, T.F.H. Bovee, M.W.F. Nielsen (Wageningen, The Netherlands) and
C.T. Elliott (Belfast, Northern Ireland) 24

Concentrations of danofloxacin 18% solution in plasma, milk and tissues after subcutaneous injection in dairy cows
N. Mestorino, M.L. Marchetti, E. Turic, J. Pesoa and J. Errecalde (Buenos Aires, Argentina) 33

A confirmatory method for the determination of tetracyclines in muscle using high-performance liquid chromatography with diode-array detection
E. Cristofani, C. Antonini, G. Tovo, L. Fioroni, A. Piersanti and R. Galarini (Perugia, Italy) 40

Confirmatory method for the determination of resorcylic acid lactones in urine sample using immunoaffinity cleanup and liquid chromatography-tandem mass spectrometry
G. Dusi, E. Bozzoni, W. Assini, N. Tognoli, M. Gasparini and E. Ferretti (Brescia, Italy) 47

Multi-functional sample preparation procedure for measuring phytoestrogens in milk, cereals, and baby-food by liquid-chromatography tandem mass spectrometry with subsequent determination of their estrogenic activity using transcriptomic assay
J.-P. Antignac, I. Gaudin-Hirret (Nantes, France), H. Naegeli (Zürich, Switzerland), R. Cariou (Nantes, France), C. Elliott (Belfast, Ireland) and B. Le Bizec (Nantes, France) 55

Determination of streptomycin residues in honey by liquid chromatography-tandem mass spectrometry
R.H.M.M. Granja, A.M.M. Niño, R.A.M. Zucchetti, R.E.M. Niño (Campinas, Brazil), R. Patel (Egham, UK) and A.G. Salerno (Campinas, Brazil) 64

The development and validation of a multiclass liquid chromatography tandem mass spectrometry (LC-MS/MS) procedure for the determination of veterinary drug residues in animal tissue using a QuEChERS (QUick, Easy, CHeap, Effective, Rugged and Safe) approach
G. Stubbings and T. Bigwood (York, UK) 68

Determination of 5-nitroimidazoles and corresponding hydroxy metabolites in swine kidney by ultra-performance liquid chromatography coupled to electrospray tandem mass spectrometry X. Xia, X. Li, S. Ding, S. Zhang, H. Jiang, J. Li and J. Shen (Beijing, China)	79
Mass Spectrometry	
Rapid multi-residue analysis of antibiotics in muscle by liquid chromatography-tandem mass spectrometry K. Granelli, C. Elgerud, Å. Lundström, A. Ohlsson and P. Sjöberg (Uppsala, Sweden)	87
Desorption electrospray ionisation mass spectrometry: A rapid screening tool for veterinary drug preparations and forensic samples from hormone crime investigations M.W.F. Nielsen, H. Hooijerink, F.C. Claassen, M.C. van Engelen and T.A. van Beek (Wageningen, The Netherlands)	92
Determination of cyanuric acid residues in catfish, trout, tilapia, salmon and shrimp by liquid chromatography-tandem mass spectrometry C.M. Karbiwnyk, W.C. Andersen, S.B. Turnipseed, J.M. Storey, M.R. Madson (Denver, CO, USA), K.E. Miller (Laurel, MD, USA), C.M. Giesecker, R.A. Miller, N.G. Rummel and R. Reimschuessel (Denver, CO, USA)	101
Development of a rapid method for the analysis of synthetic growth promoters in bovine muscle using liquid chromatography tandem mass spectrometry E.M. Malone (Kildare, Ireland and Belfast, UK), C.T. Elliott, D.G. Kennedy (Belfast, UK) and L. Regan (Kildare, Ireland)	112
Elimination kinetic of recombinant somatotropin in bovine M.-H. Le Breton (Nantes, France and Lausanne, Switzerland), S. Rochereau-Roulet, G. Pinel, N. Cesbron and B. Le Bizec (Nantes, France)	121
A determinative and confirmatory method for residues of the metabolites of carbadox and olaquindox in porcine tissues J.O. Boison, S.C. Lee and R.G. Gedir (Saskatoon, Canada)	128
A generic method for the quantitative analysis of aminoglycosides (and spectinomycin) in animal tissue using methylated internal standards and liquid chromatography tandem mass spectrometry F.L. van Holthoorn, M.L. Essers, P.J. Mulder (Wageningen, The Netherlands), S.L. Stead, M. Caldow, H.M. Ashwin and M. Sharman (York, UK)	135
Validation of multiresidue methods for veterinary drug residues; related problems and possible solutions A. Kaufmann (Zürich, Switzerland)	144
In-house validation and factorial effect analysis of a liquid chromatography-tandem mass spectrometry method for the determination of steroids in bovine muscle K.S. Schmidt, C.S. Stachel and P. Gowik (Berlin, Germany)	156
Determination of hormonal growth promoters in bovine hair: Comparison of liquid chromatography-mass spectrometry and gas chromatography-mass spectrometry methods for estradiol benzoate and nortestosterone decanoate E. Duffy (Dublin, Ireland), L. Rambaud, B. Le Bizec (Nantes, France) and M. O'Keeffe (Dublin, Ireland)	165
Depletion of chloramphenicol in trout after a hypothetic therapeutic treatment G. Bianotto, L. Contiero, C. Benetti (Legnaro, Italy), M. Calligaris, E. Tibaldi (Pagnacco, Italy), L. Cerni and M. Francese (Trieste, Italy)	173
Detection and identification of 20-hydroxyecdysone metabolites in calf urine by liquid chromatography-high resolution or tandem mass spectrometry measurements and establishment of their kinetics of elimination after 20-hydroxyecdysone administration B. Destrez, G. Pinel, F. Monteau (Nantes, France), R. Lafont (Paris, France) and B. Le Bizec (Nantes, France)	178
Rapid method for the determination of tranquilizers and a beta-blocker in porcine and bovine kidney by liquid chromatography with tandem mass spectrometry K. Mitrowska, A. Posyniak and J. Zmudzki (Pulawy, Poland)	185
How to decrease ion suppression in a multiresidue determination of β-agonists in animal liver and urine by liquid chromatography-mass spectrometry with ion-trap detector F. Moragues and C. Igualada (Valencia, Spain)	193
New method for the analysis of flukicide and other anthelmintic residues in bovine milk and liver using liquid chromatography-tandem mass spectrometry B. Kinsella (Dublin, Ireland), S.J. Lehota, K. Mastovska, A.R. Lightfield (Wyndmoor, PA, USA), A. Furey (Cork, Ireland) and M. Danaher (Dublin, Ireland)	196
Simple and rapid determination of enrofloxacin and ciprofloxacin in edible tissues by turbulent flow chromatography/tandem mass spectrometry (TFC-MS/MS) R. Krebber, F.-J. Hoffend and F. Ruttmann (Monheim am Rhein, Germany)	208
Determination of ten sulphonamides in egg by liquid chromatography-tandem mass spectrometry A.F. Forti and G. Scorticini (Teramo, Italy)	214
Determination of dapsone in meat and milk by liquid chromatography tandem mass spectrometry M. Hadjigeorgiou, C. Papachrysostomou, Z. Theodorou, P. Kanari and S. Constantinou (Nicosia, Cyprus)	220
Sensors and Bioselective Reagents	
Validation and application of a yeast bioassay for screening androgenic activity in calf urine and feed T.F.H. Bovee, G. Bor, H.H. Heskamp, J.J.P. Lasaroms, M.B. Sanders and M.W.F. Nielsen (Wageningen, The Netherlands)	225

Biosensor-based detection of reduced sex hormone-binding globulin binding capacities in response to growth-promoter administrations	
M.H. Mooney (Belfast, UK), A.A. Bergwerff, J.A. van Meeuwen (Utrecht, The Netherlands), P.B. Luppa (Munich, Germany) and C.T. Elliott (Belfast, UK)	235
A rapid microbial inhibition-based screening strategy for fluoroquinolone and quinolone residues in foods of animal origin	
H. Ashwin, S. Stead, M. Caldow, M. Sharman (York, UK), J. Stark, A. de Rijk (Delft, The Netherlands) and B.J. Keely (York, UK)	241
Detection of anabolic androgenic steroid abuse in doping control using mammalian reporter gene bioassays	
C.J. Houtman (Amsterdam, The Netherlands), S.S. Sterk (Bilthoven, The Netherlands), M.P.M. van de Heijning, A. Brouwer (Amsterdam, The Netherlands), R.W. Stephany (Bilthoven, The Netherlands), B. van der Burg and E. Sonneveld (Amsterdam, The Netherlands)	247
Improved screening method for the detection of a range of nitroimidazoles in various matrices by optical biosensor	
C.S. Thompson, I.M. Traynor, T.L. Fodey and S.R.H. Crooks (Northern Ireland, UK)	259
Inter-laboratory comparison of a yeast bioassay for the determination of estrogenic activity in biological samples	
T.F.H. Bovee, G. Bor (Wageningen, The Netherlands), I. Becue (Gent, Belgium), F.E.J. Daamen, M.B.M. van Duursen (Utrecht, The Netherlands), S. Lehmann, G. Vollmer (Dresden, Germany), R. De Maria (Grugliasco, Italy), J.E. Fox (Eugene, OR, USA), H. Witters (Mol, Belgium), S. Bernhöft, K.-W. Schramm (Neuherberg, Germany), R.L.A.P. Hoogenboom and M.W.F. Nielsen (Wageningen, The Netherlands)	265
Validation of an enzyme-linked immunosorbent assay screening for quinolones in egg, poultry muscle and feed samples	
G. Scorticchini, L. Annunziata (Teramo, Italy), V. Di Girolamo, R. Buratti and R. Galarini (Perugia, Italy)	273
Separation Methods	
Multiresidue method for the triphenylmethane dyes in fish: Malachite green, crystal (gentian) violet, and brilliant green	
W.C. Andersen, S.B. Turnipseed, C.M. Karbiwnyk, R.H. Lee, S.B. Clark, W.D. Rowe, M.R. Madson and K.E. Miller (Denver, CO, USA)	279
Other Techniques	
Comparison of screening methods for antibiotics in beef kidney juice and serum	
M.J. Schneider, K. Mastovska, S.J. Lehotay, A.R. Lightfield (Wyndmoor, PA, USA), B. Kinsella (Dublin, Ireland) and C.E. Shultz (Washington, DC, USA)	290
Comparison of three microbial screening methods for antibiotics using routine monitoring samples	
M.G. Pikkemaat, M.L.B.A. Rapallini, S.O.-v. Dijk and J.W.A. Elferink (Wageningen, The Netherlands)	298
Detection of anabolic steroids in dietary supplements: The added value of an androgen yeast bioassay in parallel with a liquid chromatography-tandem mass spectrometry screening method	
J.C.W. Rijk, T.F.H. Bovee, S. Wang (Wageningen, The Netherlands), C. Van Poucke, C. Van Peteghem (Ghent, Belgium) and M.W.F. Nielsen (Wageningen, The Netherlands)	305
Study of the depletion of lincomycin residues in honey extracted from treated honeybee (<i>Apis mellifera</i> L.) colonies and the effect of the shook swarm procedure	
S.J. Adams, R.J. Fussell, M. Dickinson, S. Wilkins and M. Sharman (York, UK)	315
Detection of glucocorticoid bioactivity in bovine urine samples using a reporter gene assay	
L. Connolly, K. Cai (Belfast, UK), E. Van der Heiden, M.-L. Scippo, M. Muller (Liege, Belgium), J. Tarbin (Sand Hutton, UK) and C. Elliott (Belfast, UK)	321
Antibody production: Low dose immunogen vs. low incorporation haptens using salmeterol as a model	
T.L. Fodey, N.M. Greer and S.R.H. Crooks (Belfast, UK)	328
A National Residue Control Plan from the analytical perspective—The Brazilian case	
A. de Q. Mauricio, E.S. Lins and M.B. Alvarenga (Brasília, Brazil)	333
Food flavonoid aryl hydrocarbon receptor-mediated agonistic/antagonistic/synergistic activities in human and rat reporter gene assays	
E. Van der Heiden, N. Bechoux (Liège, Belgium), M. Muller (Sart-Tilman, Belgium), T. Sergeant, Y.-J. Schneider, Y. Larondelle (Louvain-la-Neuve, Belgium), G. Maghuin-Rogister and M.-L. Scippo (Liège, Belgium)	337
In-house reference materials: 5-hydroxyflunixin and meloxicam in cow milk—preparation and evaluation	
P. Jedziniak, T. Szprengier-Juskiewicz and M. Olejnik (Pulawy, Poland)	346
Effect of growth-promoting 17 β -estradiol, 19-nortestosterone and dexamethasone on circulating levels of nine potential biomarker candidates in veal calves	
G. Cacciatore, S.W.F. Eisenberg (Utrecht, The Netherlands), C. Situ, M.H. Mooney (Belfast, UK), P. Delahaut (Marloie, Belgium), S. Klarenbeek (Utrecht, The Netherlands), A.-C. Huet (Marloie, Belgium), A.A. Bergwerff (Utrecht, The Netherlands) and C.T. Elliott (Belfast, UK)	351
Calendar of forthcoming meetings	I