

# Index

## A

absorption spectroscopy, 218  
 accuracy  
   defined, 10, 12  
 active-experiment approach, 266  
 active experiments, presumptions for, 265  
   control of values and, 265  
   measurement of output variables and, 265  
   noise variables and, 265  
 algorithms, 4  
 algorithms for realizable optimal experimental designs  
   exact (or N-Point) D-Optimal designs, 307–309  
   sequential composite D-Optimal designs, 313–315  
   sequential D-Optimal designs, 310–311  
 aliasing, 388  
 all possible subsets, 138  
 ALS algorithm  
   PARAFAC and, 491  
 alternative hypothesis, 48–49, 122  
 alternating least-squares (ALS) methods, 491  
 alternating least-squares (ALS) methods, solution constraints, 493  
   known profiles, 494  
   nonnegativity, 494  
   unimodality, 494  
 alternating least-squares methods  
   PARAFAC / CANDECOMP, 491–494  
 American Society for Testing Materials (ASTM), 107, 114  
 analysis of emission spectroscopic data, 241  
 analysis of multivariate data in kinetics, 228  
 analysis of variance (ANOVA), 28  
   analysis of residuals, 30–32  
   ANOVA to test for differences between means, 28  
   between-sample variation (between-treatment variation), 29–30  
   within-sample variation (within-treatment variation), 29  
 analyte concentrations in unknown solutions, 476  
 analytical chemistry, 2, 512  
 ANOVA (analysis of variance), 125  
   calculations, summary approach, 31  
   table, 32

apodization  
   transfer function and, 397  
 application of DOE in multivariate calibration, 321–329  
   improving quality from historical data, 330–335  
 application of DOE in multivariate calibration, construction of a calibration set, 321–329  
   defining the bounds of the factor space, 327–328  
   estimating extinction coefficients, 329–330  
   identifying the number of significant factors, 322–324  
 application of DOE in multivariate calibration, improving quality from historical data  
   improving the numerical stability of the data set, 333  
   prediction ability, 334  
 applications, chemometric  
   biochemical processes, 449, 449–459  
   environmental data, 454  
 archaeological artifacts  
   obsidian, composition of, 356  
 artificial neural networks (ANN), 477  
 Association of Official Analytical Chemists (AOAC), 114  
 AT&T Bell Labs, 294  
 autocatalysis, 246–248  
 autocatalytic reaction scheme, 246  
 autoscaled data with E-optimal subset (ref Fig 8.31), 335  
 autoscaling, 79, 458

## B

backward elimination, 137  
   Efroymson's stepwise regression algorithm and, 137  
   forward selection and, 137  
 bandwidth-limited discrete Fourier transform, 388  
 barycentric coordinate systems, 319  
 baseline correction, 80  
   of Raman emission spectra, 80  
 basis vectors, 71–72, 96, 141, 189  
   column basis vectors, 71  
   row basis vectors, 71  
 Beer-Lambert's Law, 219, 237, 241, 324, 328

matrix notation, result of data-fitting procedure and, 238  
 spectroscopy and, 462  
 Beer's law, 107  
 Belousov-Zhabotinsky (BZ) Reaction, 251–253, 256  
   FKN mechanism, 252  
   four stages of, 252  
   Oregonator model, 252  
   oscillating reactions and, 251–252  
   oxidation of organic species in, 251  
   represented by the Oregonator model, 263  
 between-run, defined, 10  
 bias  
   laboratory, 18  
   method, 18  
   systematic error and, 18  
 biased methods of calibration, 139  
 bilinear data matrix, 71  
 bivariate normal distribution  
   scatter plots of, 56–58  
 “black box” principle, 265, 265, 266, 269  
 BP Amoco naphtha processing plant, 330  
 Butterworth filter equation, 396

## C

calculation of the concentration profiles: case I,  
   simple mechanisms, 220  
 calculation of the concentration profiles: case II,  
   complex mechanisms, 241–252  
   Fourth-Order Runge-Kutta Method in Excel,  
   242–245  
   Interesting kinetic examples, 246–252  
 calculation of the concentration profiles: case III,  
   very complex mechanisms, 253  
   Arrhenius theory and, 254  
   changes in ionic strength and, 254–255  
   changes in pH and, 254–255  
   experimental inconsistencies and, 253  
   Eyring theory and, 254  
   temperature fluctuations and, 253  
 calibration, 105–166  
   curvilinear, 112  
   data sets and, 107–108  
   internal standards, 156  
   introduction to, 109  
   measurement error and measures of prediction  
   error, 114  
   multivariate, 111  
   non-zero intercepts and, 110  
   preprocessing techniques, 156, 156–157  
   sample sets, 264  
   selection of calibration and validation samples,  
   113  
   sets, 322  
   software, 159  
   statistical evaluation of calibration models  
     obtain by least squares, 121–134  
   univariate, 109, 211  
 calibration, a practical example, 116  
   graphical survey of NIR water-methanol data,  
   116  
   multivariate, 119–120  
   univariate, 118, 118–119  
 calibration, biased methods of, 139–153  
   example regularization results, 153  
   other calibration methods, 150–151  
   partial least squares, 147–149  
   principal component regression, 140–153  
   regularization, 151  
 calibration line, fulcrum and, 129  
 calibration models, 330  
   spectral data and, 324  
   spectroscopic, 321  
 calibration models, statistical evaluation obtained  
   by least squares, 121–134  
   coefficient of determination and multiple  
   correlation coefficient, 130  
   confidence interval and hypothesis tests for  
   regression coefficients, 126  
   hypothesis testing, 122  
   interference effects and selectivity, 234  
   interpreting regression in ANOVA tables, 125  
   leverage and influence, 128  
   model departures and outliers, 129  
   partitioning of variance in least-squares  
   solutions, 123–124  
   prediction confidence intervals, 127  
   sensitivity and limit of detection, 131–132  
 calibration, multivariate, 119–120  
   chemometrics and, 264  
 calibration, software and, 159  
   DeLight, 159  
   GRAMS, 159  
   MATLAB and, 159–160  
   Pirouete, 159  
   PLS\_Toolbox, 159  
   SIMCA, 159  
   Unscrambler, 159  
 calibration, standard addition method, 153–154,  
   153–155  
   multivariate standard addition method, 155  
   univariate standard addition method, 154,  
   154–155  
 calibration, standardization, 157, 157–159  
   alternative standardization (transfer) methods,  
   157  
   calibration-transfer methods, 157  
   of instrument response, 158

- of predicted values, 157–158
- with preprocessing techniques, 159
- calibration steps
  - calibration prediction step, 143
  - compute a regression vector using calibration samples, 143
  - compute the projected calibration spectra, 142
  - estimate the RMSEC, 143
- calibration, three-way, 4
- hyphenated data and, 475–508
- calibration, variable selection, 135–139
  - all possible subsets, 138
  - backward elimination, 137
  - Efroymson's stepwise regression algorithm, 136–137
  - forward selection, 135
  - recommendations and precautions, 138
  - sequential-replacement algorithms, 138
  - simulated annealing and genetic algorithm, 138
- calibration work, UV range and, 330
- cascade of wavelet coefficient vectors from
  - wavelet tree filter banks defining the discrete wavelet transform, 411
- catalog of four exact *D*-Optimal experimental designs
  - for the spectroscopic calibration problem, 327
- central composite designs (CCD), 293
  - three-factor with axial values and four center points, 294
  - for two factors, 293
- central limit theorem, 13, 45
  - implications of, 45
- Chebyshev polynomials, 396, 404
- chemical analysis
  - interferences and, 134–135
- chemical measurements
  - chemical properties and, 2
  - decision making and, 1, 210
  - physical properties and, 2
  - statistical properties and, 2
  - as three-legged platform, 2
- chemical properties, 2
  - equilibria, 2
  - kinetics, 2
  - mass balance, 2
  - stoichiometry, 2
- chemometrics, 1–6, 509, 510, 516
  - analytic chemistry and, 510
  - biomedical applications of, 514
  - calibration, validation, and significance testing, 2
  - in *Chemical Abstracts*, 511
  - chemical and biological sciences, 511, 514
  - chemical data and, 510
  - chemical measurements and, 2, 509
  - computing and, 2, 510, 512, 513
  - environmental forensics and, 512
  - extraction of chemical information from analytical data, 2
  - future trends, 509–520
  - gas chromatography (GC) and, 510
  - gas chromatography-mass spectrometry (GC-MS), 510
  - historical development of, 50, 509
  - hyper spectral images, 514
  - large data sets and, 514
  - live cell hyperspectral imaging, 514
  - mathematical modeling and, 509
  - as maturing discipline, 511
  - multivariate calibration methods, 510
  - multivariate methods of analysis, 510
  - multivariate statistics and, 509
  - near infrared spectroscopy and, 510
  - new instruments and, 513
  - NIR spectroscopy and, 510, 514
  - partial least-squares, 510
  - partial least-squares, 509
  - pattern recognition and, 509
  - principal component analysis (PCA), 509
  - principle component analysis (PCA) and, 510
  - quantitative biology and, 514
  - Raman spectroscopy and, 514
  - smart sensors and, 512
- chemometrics citations in *Chemical Abstracts* from 1974–2004, 511
- chemometrics, drivers of growth in, 513
  - chemical and biological sciences, 514
  - large data sets, 514
- chemometrics, reviews of and future trends, 511
  - food and feed chemistry, 512–513
  - interesting application areas, 511, 513
  - process analytical chemistry, 512
  - spectroscopy, 512
- chi-square distribution, 47, 58–59
  - degrees of freedom and, 48
- chromatography, 140, 477
- chromatographic profile optimized by ITTFA, 438
- classical and robust tolerance ellipse of the phosphorus data set, 173
- classical PCA (CPCA), 186, 188, 194
- classical PCR (CPCR), 196
- classical quadratic discriminant analysis (CQDA), 207
- classification, 207, 351–354
  - discriminant analysis and, 207
  - K-Nearest neighbor, 352
  - partial least squares (PLS), 352
  - SIMCA, 353
  - supervised learning and, 207
- classification and pattern recognition, 339–378

- clustering and, 347
- data preprocessing, 341
- introduction to, 339–340
- mapping and display, 342–346
- practical considerations, 354
- classification in high dimensions, 211
  - dimension-reduction procedure (PCA), 211
  - MCD as uncomputable, 211
  - SIMCA method (soft independent modeling of class analogy), 211
- classification in low dimensions, 207–210
  - classical and robust discriminant rules, 207
  - evaluating the discriminant rules, 208
  - an example, 209
- classification methods, 351
  - K-nearest neighbor (KNN), 351
  - partial least squares (PLS), 351
  - soft independent modeling by class analogy (SIMCA), 351
- classification of jet fuels, 364, 365
- closure, 434
- cluster analysis, 347
- clustering, 345, 347, 363
  - classification with PCA score plots and, 98
  - hierarchical, 351
  - single linkage and, 351
- clustering algorithms, 348
  - FCV, 348
  - hierarchical, 348
  - K-means, 348
  - Patrick Jarvis, 348
  - similarity matrix, 348
- clusters
  - determining numbers of, 348
  - elliptical, 351
  - identifying, 347
  - linear, 351
  - spherical, 351
  - validity, 347
- CM-estimators, 183
- coefficient of determination, 131
  - multiple correlation coefficient and, 130
- combined canonical models
  - independent (process) variables and, 284
  - mutually dependent (mixture variables) and, 284
- combined mixture and process factor space
  - constructed from three mixture variables and one process variable, 286
- components included in product P1, with UV-Inactive components shaded, 328
- composition (loading) profiles of resolved components in the MCR-ALS analysis of raw augmented data matrix, 460
- compound maps, 462
- compounds, kinetic or thermodynamic process and, 435
- computation for distance between data cluster and a point, 350
- concentration of Components C1 to C9 used to measure pure-component spectra, 324
- concentration profiles
  - for a first-order reaction at constant and increasing temperature, 254
  - of an HPLC-DAD data set, 425
  - for a reaction as calculated by MATLAB, 221
  - for a two-component system, 421
- concentration windows, 426
- confidence intervals, 14, 299
  - compared with nonoptimal design and D-optimal design, 301
  - critical value of the Student statistic, 300
  - estimate of the measurement error and, 300
  - hypothesis tests for regression coefficients and, 126–127
  - influence of on the error of the prediction, 300
  - of the mean, equation for, 46
  - for the regression line and for predictions, illustration of, 128
  - variance of the prediction, 300
- confidence level, 133
- constrained region
  - in three-component mixtures, 281
  - of three mixture variables with lower bounds only, 280
- constrained mixture region with upper and lower bounds, 282
- constrained mixture spaces, 279–280
- constraints, 433–437
  - application of, 433
  - chemical properties and, 433
  - closure, 434
  - effect on the shape of resolved profiles, 433
  - equality and inequality and, 433
  - hard-modeling constraints: physicochemical models, 435
  - implementation of, 433
  - iterative optimization and, 433
  - known profiles, 435
  - local rank, 435
  - mathematical properties and, 433
  - nonnegativity, 433, 434, 465
  - selectivity, 435
  - unimodality, 434
  - zero-concentration, 435
- constraints, hard-modeling, 435
  - concentration profiles and, 436
  - physicochemical models, 435
- construction and decomposition of a three-way array via the trilinear PARAFAC model, 480

- construction of a three-way array according to the unconstrained Tucker model, 480
  - continuous sine-wave signal with aliased curve reconstructed, 387
  - controlled and noise variables, 297
    - examples of, 295
  - convergence progress for PARAFAC from DTLD
    - initialization and random initialization, 499
  - convolution, 397
  - Cook's distance
    - distance measures and, 129
  - coordinate systems
    - barycentric, 271
    - barycentric and Descartes, three-dimensional example, 271
    - barycentric and Descartes, two-dimensional example, 270
    - Descartes, 271
    - simplex, 271
  - correlation coefficients between absorbance at selected pairs of wavelengths for sulfamethoxazole training set, 63
  - CPCR
    - RPCR and, 199
  - CQDR, 209
  - critical values for  $F$ 
    - for a one-tailed test, 20
    - for a two-tailed test, 21
  - curvature effect, 113
  - curve resolution (CR) method, 421, 423
  - curvilinear calibration, 112–113
  - cutoff frequency, 395
- D**
- D-optimal design, 301
  - data analysis algorithms, 481
    - DTLD Algorithm, 481
    - GRAM algorithm, 481
    - MATLAB programming language and, 481
    - PARAFAC algorithm, 481
  - data matrix
    - augmentation arrangement and bilinear model for PCA or MCR-ALS decompositions, 455
    - plotting, 431
  - data points in subspace of original measurement space, 344
  - data preprocessing, 341
    - normalization, 342
    - raw data into computer ready form, 341
    - scaling, 341
  - data sets, 55
    - fundamental modes of vibration, overtones, and combinations, 108
    - near infrared spectroscopy and, 107
    - solvent interactions, 108
    - water-methanol mixtures, 108
  - data sets from multicomponent systems, 418
  - data space
    - dividing, 351
  - data vector, 78
  - DATAN data analysis method, 495
  - datascope, 431
  - Daubechies wavelets, 407
  - decision limits, graphical representation of, 132
  - degrees of freedom, 46, 123–124, 125, 235
  - dendrogram, single-linkage
    - of the obsidian data, 357
  - dendograms, 351
    - as intuitive and, 349
    - using single-linkage method developed from gas chromatograms of scent marks of Marmoset monkeys, 349
  - denoised signal, 414
  - denoising, 413
    - of wavelet components using entropy threshold, 413
  - Descartes coordinate systems, 319
  - design efficiency, 303
  - Design Expert, 319
  - design measures, 303
  - design of experiment (DOE), 264
    - software, 320
  - designer transfer function, 395
  - designs, full-factorial and simplex-lattice, 283
  - determination limit, 134
  - DETMAX algorithm, 308
  - DFT, 400
  - direct exponential curve-resolution algorithm (DECRA), 495
  - direct methods, simplex, 225
    - simplex algorithm and, 225
  - direct trilinear decomposition (DTLD), 489
    - GRAM algorithm and, 489
  - discrete, bandwidthlimited Fourier transform of noisy spectrum, 390
  - discrete convolution of two functions, 393
  - discrete Fourier transforms (DFT), 389, 409
  - discrete wavelet, 409
  - discrete wavelet transform (DWT), 409, 410
  - discriminant analysis
    - classical quadratic discriminant analysis (CQDA), 207
    - cross-validation and, 208
    - evaluating the discriminant rules, 208
    - probability of misclassification, 208
  - distribution
    - functions, 41–68
    - normal, 14
    - volume of, 53

distribution curves  
 normal and standard normal, 44

distribution of error in Eigenvalues, 93

Dixon Q test, 33, 33

DTLD, 498  
 algorithm (Appendix 12.2), 503  
 application, 490, 494–495  
 application, GRAM and, 490

## E

E-Optimal calibrations, 336

effect of using a 30-point E-optimal subset during  
 the latent-variable extraction, 334

effects of constrains in feasible bands, 447

Efroymson's stepwise regression algorithm,  
 136–137  
 convergence of algorithm, 137  
 variable-addition step, 136, 136–137  
 variable-deletion step, 137

eigenvalues, 70, 74, 75, 76, 141  
 information and, 345  
 primary, 74  
 secondary, 75

eigenvectors, 70, 74, 75, 76, 141, 189  
 complete set of, 74  
 eigenspectra and, 73

environmental data  
 environmental monitoring and, 454  
 MCR-ALS and, 454  
 multivariate curve resolution, 454  
 PCA and, 454  
 principle component analysis and, 454

environmental monitoring program in Portugal,  
 455

environmental study of SVOc in surface waters of  
 Portugal, 460

error, type I, 50, 123, 133

error, type II, 50, 123

error, types of  
 gross error, 10  
 random error (noise), 10  
 systematic error, 10

errors, estimating, 235

estimated concentration in mixture samples from  
 GRAM with concatenated matrices, 486

estimated concentrations in mixture samples from  
 GRAM with added matrices, 486

estimated factors  
 visual inspection and, 498

estimating extinction coefficients, 329

Euclidean distance between two data points, 350

Euler's method for numerical integration, 242  
 simple, 242

evolving factor analysis (EFA), 423, 426

backward EFA, 424

emergence and decay of compounds and, 424

forward EFA, 424

plot, 424

exact (or N-Point) D-Optimal designs  
 DETMAX algorithm, 308  
 Fedorov's algorithm, 307  
 MD Galil and Kiefer's algorithm, 309  
 Wynn-MITtchell and van Schalkwyk algorithms,  
 308

Excel, 218

Excel's Solver, 227  
 algorithm, 234  
 using for nonlinear fitting of a first-order  
 reaction, 228

experimental design, 4, 265  
 catalogs of, 320  
 controlled factors and, 294  
 desired properties for, 298–299  
 full factorial design (FFD), 285  
 incomplete lattice, 285  
 for lithium lubricant study, 278  
 mixture and process design, 285  
 noise factors and, 294  
 simplex-lattice designs, 272

experimental design, off-the-shelf software  
 packages, 316–320  
 Design Expert, 319  
 MATLAB, 316–318  
 other packages, 319

experimental error  
 Monte Carlo simulation methods and, 448  
 resampling methods and, 448

experimental region, 268

experiments, passive, 330

explained variances by PCA for three data  
 pretreatment methods, 458

extensions of three-way methods, 495

extraction of analyte, 476

extreme values  
 outliers, 34  
 stragglers, 34

## F

F-test for determining the number of factors, 93

*Factor Analysis in Chemistry*, Malinowsky  
 history of chemometrics and, 422

factor spaces  
 constrained mixture spaces, 279–282  
 defining bonds of, 327  
 mixture and process spaces, 283–286  
 one-dimensional, 269  
 process, 268  
 simplex-centroid designs, 275–278

simplex-lattice designs, 272–274  
 two and three-dimensional in coded variables, 269  
 two-dimensional, 268

factor spaces, mixture, 269–271  
 Euclidean space and, 269  
 mixture variables and, 269

factorial designs  
 advantages of, 290  
 disadvantages of, 290

factorial designs, full, 290  
 three-level, three factor, 292  
 three or more levels in, 291  
 two-level three factor, **291**

fast Fourier transforms (FFT), 389  
 algorithms, 389  
 Cooley-Tukey transform and, 389  
 Sande-Tukey transform and, 389

FAST-MCD method, 203

Fearn's algorithm for OSC smoothing, 385

Fedorov's algorithm, 307

FFT, 400

figures of merit, analytical, 496  
 "net analyte signal" (NAS) and, 496

filter  
 lag, 400  
 window, 402

filtering, 380, 400

finite impulse response (FIR) filter, 401  
 signal processing with a simple moving average filter, 401

first and second derivatives, 82  
 spectra of water-methanol mixtures, 117

first order kinetic single-wavelength experiment  
 and result of a least-squares fit, 223

fitting parameters to a given measurement, 225  
 direct methods, 225  
 Newton-Gauss methods, 225

fitting, with Excel's Solver, 239

fixed-size image window-evolving-factor  
 analysis (FSIW-EFA), 463

fixed-size moving-window--evolving-factor  
 analysis (FSMW-EFA), 423, 426

flash photolysis, 218

flow reactors, 256

food and feed chemistry, 512

Ford Motor CO., 294

forecasting, 127

forward selection, 136

Fourier domain, 388, 399

Fourier series, 385

Fourier transform, 386, 397, 400  
 convolution and, 391  
 correlation and, 391  
 Parseval's theorem and, 391

properties of, 389–394, **391**  
 window function and real part, 393

Fourier transform infrared spectrometer  
 (FTIR), 70

frequency-domain signal processing, 385–393  
 bandwidth-limited discrete Fourier  
 transform, 388  
 Fourier transform, 386  
 properties of the Fourier transform, 389  
 sampling theorem and aliasing, 386–387

frequency-domain smoothing, 394, 394–397  
 with designer transfer functions, 395–397

frequency response of noisy signal and its  
 matched smoother, 398

FSIW-EFA analysis, 464

fuel spill identification, 358  
 gas chromatograms, 358

full factorial design (FFD), 283–284

Fundamental Reviews in Chemometrics, 512

fundamental modes of vibration, overtones, and  
 combinations, 108

## G

gas chromatography (GC), 70, 330

general equivalence theorem, 304

generalized eigenvalue-eigenvector problem  
 (GEP), 482

generalized M-estimators (GM-estimators), 183

generalized multivariate squared distance, 52

generalized or mahalanobis distances, 52

generalized rank annihilation method (GRAM),  
 4, 475, 485

generic plot of a variance indicator against a bias  
 measure, 145

genetic algorithm (GA), 138

genome sequences, chemometrics and, 514

GRAM, 489, 495, 498  
 algorithm (Appendix 12.1), 502  
 application, 486–488  
 trends in predictive performance, 487

graphical survey of NIR water-methanol data, 116

Grubbs' tests  
 application to chemical data, 35  
 critical values of G for, 35  
 outliers, 34

## H

Hadamard transform function, 317

hard-modeling constraints: physicochemical  
 models, 435

harmonious PCR plot for methanol, 149

Hawkins-Bradu-Kass data set, 187  
 artificial data set, 186

Heisenberg uncertainty principle, 409  
 Hessian matrix, 234  
 heuristic evolving latent projections (HELP), 429–430  
   HELP method, 431  
 hierarchical models, 514  
 high-performance liquid chromatography (HPLC), 70  
 high-speed gas chromatographic profiles of jet fuels, 360  
 histogram showing absorbance of sulfamethoxazole training set, 62  
 Hotelling, 510  
 HPLC-DAD  
   data set, 431  
   system, 430  
 “hyphenated” mass spectrometry-chromatographic methods, 423  
 hyphenated methods, 477  
 hypothesis test  
   comparison of multivariate means, 59  
 hypothesis testing, 122–123  
   alternative hypothesis, 48–49  
   null hypothesis, 48–49  
   principles of, 48–49  
   statistical inference and, 48–49

**I**

impulse response function, 399  
 impulse response function of the seven-point rectangular smoother window function, 396  
 incept, nonzero, 119, 462  
 initial estimates, chemometric methods and, 432  
 input variables, 268  
   mixture variables, 268  
   process variables, 268  
 interferences, 134–135  
   chemical, 134–135  
   physical, 134–135  
   spectral, 134–135  
 International Chemometric Society, 509  
 International Union of Pure and Applied Chemistry (IUPAC), 12  
 Interpreting regression ANOVA tables, 125  
 iterative resolution methods, 431  
 iterative target transformation factor analysis (ITFA), 432, 437–438, 477  
   process and, 438–439  
   target factor analysis (TFA) and, 437  
   target vector and, 439

**J**

Jacobian J  
   representation of, 232  
 Jacobian J and Marquardt parameter, 234  
 Jacobian J matrix, 232–233

**K**

K-nearest neighbor (KNN), 352, 354  
   Bayes classifier and, 352  
   Euclidean distance and, 352  
 kinetic data, analysis of, 246–253  
   additional methods, 258  
   autocatalysis, 246  
   Belousov-Zhabotinsky (BZ) Reaction, 251–253  
   flow reactors, 258  
   globalization of the analysis, 258–259  
   Lotka-Volterra (Sheep and Wolves), 250–251  
   measurement techniques and, 256  
   model parser, 256  
   related issues, 255–258  
   soft-modeling methods, 257–258  
   Zeroth-Order reaction, 248–249  
 kinetic modeling of multivariate measurements,  
   nonlinear regression, 217–262  
   Beer-Lambert’s Law, 219  
   calculation of the concentration profiles:  
     case I, simple mechanisms, 220  
     calculation of the concentration profiles:  
       case II, complex mechanisms, 241–252  
     calculation of the concentration profiles: case  
       III, very complex mechanisms, 253–254  
   matrix notation, 219  
   model-based non-linear fitting and, 222–240  
   multivariate data, 219–220  
   related issues, 255–258  
 kinetic modeling of multivariate measurements,  
   related issues  
   flow reactors, 256  
   globalization of the analysis, 256  
   measurement techniques, 256  
   model parser, 256  
   other methods, 258  
   soft-modeling methods, 257  
 kinetics of chemical processes  
   light-absorption spectroscopy and, 218  
 kinetic reactions, 246–252  
   autocatalysis, 242–246  
   Belousov-Zhabotinsky (BZ) Reaction, 251–252  
   Lotka-Volterra (Sheep and Wolves), 250  
   zeroth-order reaction, 248–249  
 Kjeldahl nitrogen titration method, 107



known profiles, 435  
 Kohonen neural network, 345–346  
 Kowalski, Bruce  
   chemometrics and, 509

## L

least median of squares (LMS), 183  
 least squares, 121  
   classic, 107  
   estimates, 111  
 least-squares fit, 222, 237  
   of emission spectroscopic data, 240  
 leave-multiple-out CV (LM) (CV)  
   Monte Carlo CV (MCCV), 115  
 leave-one-out cross-validation (LOOCV), 115  
 leverage and influence, 128  
   calibration samples and, 128  
 leverage, in statistics, 128  
 leverage points, 184, 191  
 light-absorption spectroscopy  
   CD (circular dichroism), 218  
   NIR (near-infrared\_ reflectance spectra, 218  
   UV/Visible spectroscopy, 218  
 limit of detection, 132  
   determination limit, 134  
   multivariate detection limit, 134  
   univariate decision limit, 132  
   univariate detection limit, 133  
 limitation of quantity (LOQ), 114, 134  
 linear and non-linear parameters, 228  
 linear discriminant rule, 209  
 linear effect, 113  
 linear models with  $n$  standards, diagram of three  
   different types, 109  
 linear parameter fitting, 241  
 linear regression in low dimensions, 176  
   linear regression with one response variable, 176  
 linear regression with one response variable, 176  
   the classical least-squares estimator, 177  
   the multiple linear regression model, 176  
   other robust regression estimators, 182,  
   182–183  
   an outlier map, 180, 180–181  
   the robust LTS estimator (least trimmed  
   squares), 178–179  
 linear regression with several response variables,  
   183  
   an example, 185, 194  
   the multivariate linear regression model,  
   183–184  
   principle component regression (PCR), 195  
   the robust MCD-regression estimator, 184  
   selecting the number of principle components,  
   193

local rank analysis methods, 423  
   evolving factor analysis (EFA), 423  
   fixed-size moving-window--evolving-factor  
   analysis (FSMW-EFA), 423  
   images and, 463  
 local rank constraints, 435–436  
 local rank maps, 426  
   of pill images, 464  
 location and covariance estimation in low  
   dimensions, 173  
   the empirical mean and covariance matrix, 173  
   other robust estimators of location and  
   covariance, 176  
   the robust MCD estimator, 174–176  
 location and scale estimation  
   mean and standard deviation, 169–171  
   median and median absolute deviation, 171  
   other robust estimators of location and scale,  
   171–173  
 logarithm of the square sum (ssq) of the residual  
   vector  $r$  as a function of the rate constant  $k$ ,  
   224  
 Lotka-Volterra (Sheep and Wolves), 250–251  
   dynamics of population of predators and preys  
   in closed systems, 250  
   kinetic population profiles and, 250  
   MATLAB's Runge-Kuta solver and, 250  
   predator and prey kinetics, 250  
 LTS estimator  
   regression, scale, and affine equivariant, 179  
 LTS regression, 183, 197, 198  
   errors and, 179

## M

M-estimators, 183  
 Mahalanobis, 52, 192  
   distances, 52, 59, 64  
 Mahalanobis Distances and Probability Densities  
   from Hotelling's  $T^2$  Statistic for Test Samples  
   of Sulfamethoxazole compared with the  
   Training Set, 64, 66  
 Malinowski's empirical indicator function (IND),  
   91  
 Malinowski's RE function, 91  
 Manne's resolution theorems, 421, 427  
 mapping and display, 341  
   data compression, 343  
   dimensionality reduction with PCA and, 343  
   principle component analysis (PCA) and, 342,  
   343  
   variance and, 342  
 Marquardt parameter mp, 234  
 mass spectrometer (MS), 70  
 matched smoothing of noisy signal, 398

- mathematical handling of multiwave absorption data sets, 218
- MATLAB, 74, 202, 232, 241, 456, 465, 481
- backslash notation, 234
  - chemometrics and, 218
  - code, 59, 218
  - corrocoef functions, 62
  - cov functions, 62
  - D-optimal designs and, 317–318
  - diff command, 60
  - example 3.1, 61
  - example 3.2, 61
  - example 3.3, 62
  - example 3.4, 63
  - example 3.5, 65
  - example 3.6, 66
  - example 4.1: eigenvectors and eigenvalues, 75
  - example 4.2: principal component analysis of a spectroscopic data set, 75–76
  - example 4.3: principal component analysis using the SVD, 76–77
  - example 4.4 autoscale of a matrix, 79–80
  - example 4.5: PCA procedure, 86
  - example 4.6: function for calculating Malinowski's RE, IND, and REV functions, 92–93
  - example 4.7: determining the number of significant principal components in a data matrix, 95–96
  - example 4.8: computation of residual spectra, 100
  - example 4.9: residual variance analysis, 102
  - example 7.1, 222, 223, 254
  - example 7.2, 222
  - example 7.3a, 226, 227
  - example 7.3b, 226, 227
  - example 7.4a, 229
  - example 7.4b, 230
  - example 7.5a, 235, 236
  - example 7.5b, 235, 236–237
  - example 7.5c, 235, 237
  - example 7.6a (autocatalysis), 247
  - example 7.6b, 248
  - example 7.7a (Zeroth-order kinetics), 249
  - example 7.7b (Zeroth-order kinetics), 249
  - example 7.8a, 250
  - example 7.8a lotka volterra, 251
  - example 7.8b ode lotka volterra, 251
  - find command, 61, 65
  - functions for DECA, 495
  - hist command, 61
  - m\_dist.m function, 64
  - mean functions, 62
  - plot command, 60
  - Runge-Kutta method and, 242, 246
  - Statistics Toolbox, 317
  - zoom command, 60, 65
- matrix notation, 218, 219
- maximum-likelihood rule, 207
- MCD estimates, 184, 187, 193
- MCD estimator, 198
- MCD regression, 197, 198, 203
- an example, 185
- MCR-ALS decomposition method, 448
- biochemical processes and, 449–452
- MD Galil and Kiefer's algorithm, 309
- mean
- arithmetic, 11
  - defined, 10
  - inferences about, 49
  - population, 13
- mean absolute deviation (MAD), 37
- mean centering, 78, 159
- effect on a bivariate distribution of data points, graphical illustration, 78
- measurement error and measures of prediction error, 114
- measurement techniques, 256
- median absolute deviation (MAD), 171
- membership probability, 207, 208, 209
- mesh representation of pure distribution maps, 465
- methanol model at without an intercept term, graphical displays, 120–121
- methanol model with a non-zero intercept, 124–125
- methanol model with a nonzero incept, graphical displays, 122–123
- methanol result from univariate and multivariate calibration of water-methanol mixture, 118
- middle point of the window, 402
- misclassification probabilities for RQDR and CQDR applied to the fruit data set, 209
- misclassification probability(MP), 208, 209
- misclassification rates, 210
- mixture and process factor space for incomplete cubic simplex-lattice design combined with a full factorial design, 284
- mixture and process factor spaces, 283
- symmetric experimental designs and, 283
- mixture factor spaces
- coordinates in, shown in barycentric and Descartes coordinate systems, 271
- model
- calibration and validation, 198
  - parser, 256
  - validation, 114
- model-based fitting of kinetic data, 218
- methods for nonlinear least-squares fitting, 218

- modeling the concentration profiles of the reacting components, 218
- model-based non-linear fitting, 222
  - direct methods, simplex, 225–226
  - linear and non-linear parameters, 228–229
  - Newton-Gauss-Levenberg/Marquardt (NGLM), 230
  - non-linear fitting using Excel's Solver, 227
  - nonwhite noise, 237–239
- model departures and outliers, 129
  - residual plot and, 129
- model determination showing the bias/variance tradeoff with selection of the metaparameter, 144
- Monte Carlo CV (MCCV)
  - leave-multiple-out CV (LM)(CV), 115
- mother Daubechies wavelet and two shifted and scaled children fitting a noisy signal, 407
- mother wavelet functions, 407, 407
- moving-average filter, noise reduction and, 403
- multiblock models, large data sets and, 514
- multicollinearity, 194
- multicomponent systems
  - mixtures, 448
  - processes, 448
- multiple correlation coefficient, 131
- multiple linear regression (MLR), 169
- multiple overlapping absorption bands in the NIR spectrum of water, 89
- multiple rank analyses, 423
- multiplicative scatter correction (MSC), 83, 85
- multivariate
  - distribution, 52
  - measurements, 225
- multivariate absorbance data
  - Beer-Lambert's Law and, 241
- multivariate adaptive regression splines (MARS), 477
- multivariate advantages, 3, 52
- multivariate calibration, 3, 111–112, 195
  - methods, 512
- multivariate classification and pattern recognition, 4
- multivariate curve resolution, 4, 417–467, 417–474
  - ambiguities and resolution theorems, 418–422
  - environmental data, 454–460
  - error in resolution, 446–447
  - historical background of, 422
  - local rank and resolution: evolving factor analysis and related techniques, 423–425
  - MCR-ALS Simultaneous analysis of multiple correlated data matrices, 440–445
  - range of feasible solutions, 446–447
  - resolving diverse chemical problems and, 448
  - self-modeling curve resolution extended to multiway data, 440–445
  - software, 465–467
  - spectroscopic images, 461–465
  - uncertainty in resolution results, 446–447
  - useful methods and, 441
- multivariate curve resolution-alternating least squares (MCR-ALS), 432
- multivariate curve resolution-alternating least squares MCR-ALS, 439
- multivariate curve resolution-alternating least squares (MCR-ALS), 439
- multivariate curve resolution-alternating least squares MCR-ALS, 456–459
  - procedure, 439
- multivariate curve resolution, applications, 448–453
  - biochemical processes, 449–450
  - global description of the protein folding process, 453
  - study of changes in the protein secondary structure, 451–452
- multivariate curve resolution, iterative methods, 431–439
  - constraints based on chemical or mathematical properties, 433–437
  - constraints, classification, 433–437
  - constraints, definition, 433–437
  - constraints equality and inequality, 433–437
  - generation of initial estimates, 432
  - iterative target transformation factor analysis (ITFA), 437–438
  - multivariate curve resolution-alternating least squares (MCR-ALS), 439
- multivariate curve resolution, noniterative resolution methods, 426–429
  - Heuristic Evolving Latent Projections (HELP), 429–430
  - other techniques, 429–430
  - Subwindow Factor Analysis (SFA), 429–430
  - window factor analysis (WFA), 427–428
- multivariate data, 219
- parameters and, 229
- multivariate dataset, 218
  - matrix form and, 54
- multivariate detection limit, 134
- multivariate distances, 59
  - “acceptable” and “unacceptable” objects, 64
  - calculate Mahalanobis distances and probability densities, 64
  - compute training set mean and variance-covariance matrix, 62
  - example of, 59–69
  - graphical review of smx.mat data file, 69

histograms of variables (wavelengths), 69  
 selection of, 60

multivariate distances, example

- step 1: graphical review of smx.mat data file, 60
- step 2: selection of variables (wavelengths), 61
- step 3: view histograms of selected variables, 61–62
- step 4: compute the training set mean and variance-covariance matrix, 62
- step 5: calculate Mahalanobis distances and probability densities, 64
- step 6: find “acceptable” and “unacceptable” objects, 65

multivariate limit of detection (LOD), 134

multivariate normal distribution, 3, 41–68, 51, 51–52

- assumptions, 55
- bell-shaped curve and, 51
- chi-square distribution, 56
- estimation of population parameters from small samples, 54
- generalized or Mahalanobis distances, 52
- generalized sample variance, 55
- graphic illustration of selected bivariate normal distributions, 56
- variance-covariance matrix, 53

multiway data analysis approaches, 461

## N

N-way data, 478

“net analyte signal” (NAS)

- calibration and, 496–497

net analyte signal (NAS) vector, 134

Newton-Gauss algorithm, 233

- flow diagram of, 233

Newton-Gauss-Levenberg/Marquardt (NGL/M), 230–237

- algorithm, 230, 234, 235, 235
- fitting procedure, 235
- routine, 235

Newton-Gauss method, 233

NIR absorbance spectra of water-methanol mixtures, 109

NIR (near-infrared-reflectance spectra), 60, 510

- of sulfamethoxazole powder, 59, 60

NIR spectra projected into a three-dimensional space formed by the first three principle components, 335

NIR water-methanol data set, 141

noise

- amplitude, 380
- variables, 266

noise removal and the problem of prior information, 380–381

- signal estimation and signal detection, 381
- noise removal, matched smoothers and, 397
- noisy spectrum to be enhanced, 381
- nomadization applied to Raman spectra, 83
- non-zero intercepts, 109, 110
- non-zero mean noise effects, 399
- noniterative resolution methods, HELP method, 431
- nonlinear fitting, using Excel’s Solver, 227
- nonlinear multivariate model estimation, chemical kinetics and, 4
- nonlinear parameter fitting, 241
- nonlinear regression, 477
- nonnegativity, 433
- nonsymmetric optimal designs, 298–306

  - optimal vx. equally distanced designs, 299–301
  - optimality criteria, 298

- nonsymmetric optimal designs, design optimality and design efficiency criteria, 302–302

  - A-Optimality, 306
  - D-Optimality and D-Efficiency, 304
  - design measures, 303
  - E-Optimality, 306
  - G-Optimality and G-Efficiency, 305

- nonwhite noise, 237
- normal distribution

  - plot of, 14

- normalization, 83
- null hypothesis, 19, 48–49, 122, 133

  - acceptance and rejection errors, 122

- numerical approximation of the first derivative on near-infrared spectra of water-methanol mixtures, 82
- Nyquist criterion, 388

## O

octane data set with NIR absorbance spectra

- outlier maps and, 205

one variable at a time approach, 288–290

- advantages of, 290
- applied to a two-factor problem, 289
- disadvantages of, 290
- illustration of, 289

one variable at a time vs. optimal design

- bivariate (multivariate) example, 288–289
- one-variable at a time approach, advantages of, 290
- one-variable at a time approach, disadvantages, 290

one-way data, 476

operating region, 268

optimal composite sequential designs (OCSD), 314

- schematic representation of, 314

- for three mixture variables and one process variable, optimal for models having structures in equations, **321**
  - for two process variables and two polynomial models, **315**
  - optimal sequential designs (OSD), **314**
  - ordinary differential equation solvers (ODE-solvers), **245**
  - orthogonal projection approach (OPA), **432**
  - orthogonal signal correction (OSC), **384**
  - other calibration methods, **150**
    - common basis vectors and a generic model, **150, 150–151**
    - continuum regression (CSR), **150**
    - cykuc subspace regression (CSR), **150**
    - generalized RR (GRR), **150**
    - ridge regression (RR), **150**
    - ridge variations of PCR, PLS, etc., **150**
  - outlier maps, **185, 194, 199**
    - of artificial regression data, **181**
    - of the biscuit dough data set obtained with RPCR, **201**
    - of glass data set based on three principal components computed with CPCA and ROBPCA, **195**
  - PCA analysis results and, **191–192**
    - of robust residuals vs. robust distances of the carriers for the foam data set, **186**
    - for the stars data set, **182**
  - outlier maps, three-dimensional
    - PCA and regression outlier maps and, **202**
  - outliers, **32, 168–169**
    - experimental, **33**
    - Grubbs' tests, **34**
    - orthogonal, **191, 194**
    - and outlier map of a three-dimensional data set projected on a robust two-dimensional PCA subspace, **192**
    - vertical, **184**
- P**
- PARAFAC, **481, 494–495, 495, 496**
    - Algorithm (Appendix 12.3), **504**
    - application, **494**
    - relative speed of, **499**
  - PARAFAC / CANDECOMP, **491–494**
    - algorithm, **491**
    - PARAFAC application, **494**
    - solution constraints, **493**
    - tuckals, **493**
  - PARAFAC model, **491, 498**
    - as trilinear, **495**
  - parallel factor analysis (PARAFAC), **4, 455, 475, 478–480**
  - parallel vector analysis (PVA), **428**
  - parameters
    - linear, **228**
    - molar absorptivities, **228**
    - non-linear, **228**
    - rate-constants, **228**
  - Parseval's theorem, **391**
  - partial least squares (PLS), **112, 140, 147, 147–149, 196**
    - applications, **352**
    - comparison with PCR, **149, 149–150**
    - mathematical procedures, **148–149**
    - method summarized, **352**
    - modeling method, **352**
    - number of basis vectors selection, **149**
    - regression method, **352**
    - software, **352**
  - partial least squares (PLS) regression, **202, 477**
    - classical PLSR and, **202**
    - an example, **204**
    - model calibration and validation, **204**
    - robust PLSR and, **202**
  - partitioning of variance in least-squares solutions, **123**
  - pattern-recognition, **355**
  - pattern-recognition methodology, **341**
    - classification, **341**
    - clustering, **341**
    - mapping and display, **341**
  - pattern-recognition techniques, applications of, **355, 355–371**
    - archaeological artifacts, **356–357**
    - class-membership problem, **355**
    - classification, **355**
    - data preprocessing, **355**
    - feature selection, **355**
    - fuel spill identification, **358–364**
    - mapping and display, **355**
    - PLS models, **355**
    - principle component analysis, **355**
    - self-organizing map (SOM), **355**
    - sorting plastics for recycling, **365–370**
    - taxonomy based on chemical constitution, **371–373**
  - PCR, PLS, and RR harmonious plots for methanol, **152**
  - PCR, PLS, and simplex harmonious plots for NI analysis of moisture in soy samples, **153**
  - peak-matching program, **361**
  - pharmaceutical manufacturing process, **43**
  - piecewise direct standardization (PDS), **158**
  - pixels, **461, 462**
    - PCA analysis and, **463**
  - plots

- log 19 of singular values vs. estimated concentration for RAFA, 484
- NIR spectra of water-methanol mixtures, 86
- residual spectra from the training set and the residual spectrum for an unknown w spectrum, 99
- of second and third largest principal components for jet fuel samples, 362
- simulated two-component mixture spectra (solid line) and a spectrum contaminated with a third unknown component, 99
- two largest PLS components in the training set (bees), 373
- two largest principle components for jet fuel samples, 361
- two largest principle components of 10 metals in obsidian samples, 356
- two largest principle components of the training set, 369
- PLS, 514
  - map from training-set data (honeybees), 374
  - models, 336
- point, measurement and, 302
- polynomial least-squares filtering, 404
- polynomial moving-average (Savitsky-Golay) filters, 403
- polynomial smoothing, 81–82
  - polynomial smoothing on near-infrared spectra of water-methanol mixtures, 81
- polynomials
  - canonical form of, 272
- population, 42
  - in chemistry, 43
  - finite, 42
  - infinite, 43
  - mean, 42
  - mean vector, 54
  - standard deviation, 42
- population parameters, estimation of
  - from small samples, 54
- population variance, 47
- population variance-covariance matrix, 54
- possible residual plots with the estimated residual
  - on the y-axis and the estimated concentration on the x-axis, 130
- practical considerations in experiment design, 354
  - choice of classifier, 354
  - choice of training set, 354
  - controlling experimental variables, 354
  - feature selection, 354–355
  - validity set, 354
- precision and accuracy
  - four common scenarios, 13
- precision, defined, 10, 12
- predicted calibration spectra for product P1, 330
- predicted standard and mixture concentrating from DTLD and PARAFAC, 490
- predicted sum of squares (PRESS) statistic, 193–194
- prediction confidence intervals, 127–128
- prediction set
  - for honeybee example, 373
  - for jet fuel example, 359
  - for plastics sample, 368
- prediction-set results (F-values) for Jet Fuel sample, 366
- prediction-set samples projected onto the principle component map, 369
- predictive alternating conditional expectations (PACE), 477
- preprocessing options, 77
  - baseline correction, 80
  - first and second derivatives, 82
  - mean centering, 77, 78
  - multiplicative scatter correction (MSC) and, 83
  - normalization, 83
  - smoothing and filtering, 81
  - standard normal variate (SNV) transforms, 83–85
  - variance scaling, 77, 78
- PRESS statistic, 198
- principal component analysis (PCA), 69–104, 70
  - chemometrics and, 3
  - principal component regression (PCR), 112
  - principle component analysis (PCA), 70–102, 140–153, 189, 347, 351, 422, 456–459, 514
  - alternative formulations of, 77
  - applied to chromatographic-spectroscopic data, 70
  - axes defining basis vectors, 344
  - basis vectors, 141, 144
  - classical PCA (CPCA), 185
  - conclusions, 102
  - data exploration procedures, 86
  - example PCR results, 145–146
  - introduction to, 70
  - loadings for raw mean-centered augmented data matrix, 459
  - mathematical procedures, 142
  - outlier map, 199
  - outliers, 193, 198, 199
  - PCA data exploration procedure, 86
  - preprocessing options and, 77–85
  - score plots of the water-methanol mixtures, 88
  - scores for raw mean-centered augmented data matrix, 460
- principle component analysis (PCA), basic vectors, 96–98
  - clustering and classification with PCA score plots, 98

- principle component analysis (PCA), influencing factors, 87–88
    - distribution of error in Eigenvalues, 93
    - F-test for determining the number of factors, 93
    - variance and residual variance, 89–92
  - principle component analysis (PCA) outlier map
    - applying RPCR to the biscuit dough data set
    - regression outlier map and, 201
  - principle component analysis (PCA) outlier map of octane data set obtained with RSIMPLS and SIMPLS
    - regression outlier map obtained with RSIMPLS and SIMPLS, 205–206
  - principle component analysis (PCA), residual spectra, 98–101
    - residual variance analysis, 100–101
  - principle component map, jet fuel data, 362, 363
  - principle component model, 73–76
    - alternative formulations of, 77
    - Eigenvectors and Eigenvalues, 74–75
    - singular-value decomposition, 76
  - principle component model, diagram for the chromatographic-spectroscopic data set shown in Figure 4.1, 73
  - principle component regression (PCR), 140–141, 195, 477
    - basic vectors, 141
    - classical PCR, 194
    - linear regression models and, 194
    - methanol, 145–146
    - method, 196
    - model calibration and validation, 198
    - number of principal components and, 198
  - principle component regression (PCR)
    - applications, multivariate calibration and, 195
  - principle component regression (PCR), mathematical procedures, 141–142
    - calibration steps, 142
    - example PCR results, 145
    - number of basic vectors, 144
    - unknown prediction steps, 143
  - principle component regression (PCR) models, 336
    - prediction error, estimating, 198
    - validation of, 198
  - principle components, selecting the number of, 193
  - probability distribution function, 43
  - process analytical chemistry, applications, 512
  - process variables, transformation formula, 269
  - products with the respective quantities of the components in each, 323
  - projection of a three-dimensional vector onto a two-dimensional subspace, 96
  - projection pursuit (PP) algorithm
    - RAPCA (relection algorithm for principal components analysis), 188–189
  - projection pursuit (PP) techniques, 188
  - protein folding, 451, 451
    - changes in tertiary structure and, 453
    - global description of the protein folding process, 453
  - protocols for planning experiments, 3
  - pulse radiolysis, 218
  - pulsed gradient spin echo (PGSE), 495
  - pure-component NIR spectra for water and methanol, 116
- ## Q
- quadratic discriminant rule (RQDR), 209
  - quality, illustrated with distributions of important quality characteristics, 296
  - QZ algorithm, 486
- ## R
- RAFA, 489
    - application, 483
  - Raman
    - images, 462
    - spectra of plastic, 367
  - rank annihilation factor analysis (RAFA), 476
  - rank annihilation methods, 482–490
    - direct trilinear decomposition (DTLD), 489
    - direct trilinear decomposition (DTLD) application, 490
    - generalized rank annihilation method (GRAM), 485
    - generalized rank annihilation method (GRAM) application, 486–488
    - rank annihilation factor analysis (RAFA), 482
    - rank annihilation factor analysis (RAFA) application, 482–484
  - RAPCA (relection algorithm for principal components analysis), 189
  - rate constants, 228
  - reactions, physical properties and, 2
    - energy transfer, 2
    - phase transitions, 2
    - temperature, 2
  - recovery of concentration compound by window factor analysis, 428
  - “reduced” product formulations after removal of UV-Inactive components, 326
  - reexpressing data in alternate bases to analyze structure, 381–384
  - projection-based signal analysis as signal processing, 383–384

- regression
  - deepest, 183
  - depth, 183
  - outliers, 198
- regression analysis notation, 286
- regression coefficients, 326
- regression model
  - identifying the type of, 325–326
- regularization, 151–152
  - Tikhonov regularization, 152, 153
- regularization results: example, 153
- relative speed (in GigaFLOPs) for the discussed three-way methods), 498
- repeatability, defined, 10
- reproducibility, defined, 10
- residual spectra, 98
- residual sum of squares (RSS), 136
- residual variance analysis, 100
  - example, **102**
- residuals, 128
  - studentized, 129
- resolution of a multicomponent chromatographic HPLC-DAD, **420**
- resolution results, ambiguity in, 447
- response-surface modeling (RSM), 264, 265, 265–286
  - design of experiments (DOE) and, 265
  - general scheme of, 265–267
  - methodology, 267
  - model fitting and, 265
  - process or product optimization, 265
  - regression-analysis-related notation and, 286–287
- response-surface modeling (RSM) and experimental design, 256, 263–339
  - algorithms for realizable optimal experimental designs, 306–315
  - example: application of DOE in multivariate calibration, 321–329
  - nonsymmetric optimal designs, 298–306
  - off-the-shelf software and catalogs of designs of experiments, 316–320
  - one variable at a time vs. optimal design, 288–290
  - symmetric optimal designs, 290–294
  - Taguchi Experimental Design Approach, 294
- response-surface modeling (RSM), factor spaces, 268–285
  - constrained mixture spaces, 279–282
  - mixture and process spaces, 283–286
  - mixture factor spaces, 269–271
  - process factor spaces, 268
  - simplex-centroid designs, 275–278
  - simplex-lattice designs, 272–274
- results from factor analysis of simulated chromatographic data, **94**
- results from univariate calibration of the water-methanol mixture, **118**
- ridge regression (RR), 112, 140
- RMSEC values, 118, 119
- RMSEV values, 118, 119
- ROBPCA, 193, 197, 199, 202, 203, 211
  - algorithm, 190, 198
  - outlier map and, 194
  - projection pursuit and robust covariance estimation, 189
- robust calibration, 159–160, 166–216
  - introduction to, 168
  - principle component analysis, 185
  - software availability, 211
- robust calibration, classification, 207–208, 207–211
  - classification in high dimensions, 211
  - classification in low dimensions, 207–208
  - an example (spectra of three cultivars of cantaloupe), 209
  - in high dimensions, 211
  - in low dimensions, 207–210
- robust calibration, linear regression in low dimensions
  - linear regression with one response variable, 176
  - linear regression with several response variables, 183
- robust calibration, location and covariance estimation in low dimensions, 173–176
  - the empirical mean and covariance matrix, 173
  - other robust estimators of location and covariance, 176
  - the robust MCD estimator, 174
- robust calibration, location and scale estimation, 162–172
  - the mean and standard deviation, 169
  - the median and the median absolute deviation, 169
  - other robust estimators of location and scale, 169
- robust calibration, partial least-squares regression, 202–206
  - classical PLSR, 202
  - an example, 204
  - robust PLSR, 203
- robust calibration, principle components analysis, 185–194
  - classical principle components analysis, 185
  - an example, 194
  - an outlier map, 191
  - robust PCA based on a robust covariance estimator, 187



robust PCA based on projection pursuit, 188  
 robust PCA based on projection pursuit and he MCD, 189  
 selecting the number of principle components, 193

robust calibration, principle components regression, 194–199  
 classical PCR, 194  
 an example, 199  
 model calibration and validation, 198  
 robust PCR, 197

robust calibration, software availability, 211  
 FAST-LTS, 211  
 FAST-MCD, 211  
 LIBRA (Library for Robust Analysis), 211

robust discriminant analysis, 210

robust linear discriminant analysis, 208  
 MCD estimates and, 208

robust LTS-subspace estimator, 191

robust multivariate methods, 4

robust PCA, 4  
 based on a robust covariance estimator, 187  
 based on projection pursuit, 188  
 based on projection pursuit and he MCD, 189–190

robust PCA methods  
 outliers and, 187

robust PCR, an example (biscuit data set), 198

robust PCR (RPCR) method, 197  
 regression techniques and, 197  
 robust PCA and, 197

robust PLC, 4

robust principal component analysis (RAPCA), 209

robust quadratic discriminant analysis (RQDA), 207

robust R-RMSECV curve for biscuit dough data set, 199

robust tolerance ellipses for the fruit data, 210

root mean square error of cross-validation (RMSECV), 114, 198

root mean square error of calibration (RMSEC), 114

root mean square error of prediction (RMSEP) computing, 114  
 RMSEV for validation and, 114

root mean squared error of estimation (RMSEE), 382

RPCR algorithm, 198

RSIMPLS  
 ROBPCA and, 202

RSIMPLS approach, 203  
 octane data set with NIR absorbance spectra, 204

Runge-Kutta algorithms

chemical kinetics and, 242

Runge-Kutta method, fourth order  
 in Excel, 242–245, 244  
 in Excel, concentration profiles modeled, 246

## S

S-estimators, 183

S-PLUS software, 202

sample variance, 47  
 generalized, 55

sample variance-covariance matrix, 54

samples, 43  
 influential, 128  
 projected onto two-dimensional space, 343

sampling and sampling distributions, 42, 42–44  
 normal distribution, 43–45  
 simple random sampling, 42  
 standard normal distribution, 45

sampling theorem, 387

sampling theorem and aliasing, 386–388

sampling theory, 41–68

Savitsky-Golay filtering, 404

Savitzky-Golay smoothing, 81

scale plots, 410

scatter plot  
 of absorbance at 1912 nm vs. 2264 for he sulfamethoxazole training set, 63  
 of the principal component (PC) scores from the SVD analysis of the water-methanol NIR mixture data, 142  
 of the principal component scores from the analysis of the HPLC-UV/visible data set, 96

score plot and 97.5% tolerance ellipse of the Hawkins-Bradu-Kass data obtained with CPCA and MCD, 187

second loading vector and calibration vector of sucrose for the biscuit dough data set computed with CPCR and RPCR, 200

selected bivariate normal distributions  
 graphical illustration of, 56

selection of calibration and validation samples, 113

selectivity, 135

self-modeling curve resolution extended to multiway data  
 MCR-ALS Simultaneous analysis of multiple correlated data matrices, 440–441

self-modeling curve resolution (SMCR), 477, 479

self-modeling methods, 257–258

self-organizing map (SOM), 345, 351  
 disadvantages of, 347  
 distance not preserved in, 347  
 generated from the training set, 370

- semivolatile organic compounds (SVOC), 455, 456
- sensitivity, 131
  - analytic, 131
  - calibration, 131
  - effective, 131
  - and limit of detection, 131
- sequential D-Optimal experimental design for two process variables and a second-order polynomial, 312
- sequential-replacement algorithms, 138
  - forward-selection algorithm and, 138
  - obtaining, 138
- Shell foam data, 185
- short-term Fourier transform (STFT), 406
- signal estimation and signal detection, 381
  - known signal and, 381
  - root mean squared error of estimation (RMSEE), 382
- signal processing
  - convolution property and, 391
  - digital filtering and, 4, 379–416
  - serial, 400
- signal processing and digital filtering, 379–416
  - frequency-domain signal processing, 385–393
  - frequency-domain smoothing, 394–397
  - introduction to, 379
  - noise removal and the problem of prior information, 380–381
  - reexpressing data in alternate bases to analyze structure, 382–384
  - time-domain filtering and smoothing, 398–405
  - wavelet-based signal processing, 406–413
- signal-to-noise ratio (SNR), 297, 322, 381, 411
  - formulas for, 298
- significance level, 50
- significance testing, 18
  - comparison of a sample mean with a certified value, 24–25
  - comparison of the means from two samples, 24, 25
  - comparison of two methods with different test objects or specimens, 26
  - F-test for comparison of variance (precision), 19–21
  - one-tailed or two-tailed test, 24
  - student t-test, 21, 22–23
- significant factors, identifying the number of, 322
- SIMCA method (soft independent modeling of class analogy), 211
- SIMCA training-set results for Jet Fuel Sample, 365
- similarity values, 348
- simple moving average filter (“boxcar averager”), 401
- simple-to-use interactive self-modeling analysis (SIMPLISMA), 432
- simplex, 280
  - algorithm, 226, 228, 234
  - defined, 226
  - unconstrained, 281
- simplex-centroid designs, 275–276
  - advantages of, 277
  - disadvantages of, 277
  - example (lithium lubricant study), 278
- simplex-lattice designs, 272
  - advantages of, 274
  - component contour plots and, 274
  - as composite designs, 274
  - as D-optimal, 274
  - disadvantages of, 274
  - example, 275
  - examples of for linear, quadratic, full cubic, and special cubic models, 273
  - homogenous mixtures required for, 275
- simplex-lattice mixture design, four-component for a cubic polynomial model, 276
- simplex minimization with three parameters, principle of, 226
- SIMPLISMA, 463, 465
- SIMPLS algorithm, 202
- simulated annealing (SA) and genetic algorithm, 138
- simulated HPLC-UV/visible chromatographic data set showing two overlapping peaks with different UV/visible spectra, 72
- singular-value decomposition (SVD), 75–76, 188, 430
- “slow-fast” ambiguity, 237
- small sample distributions, 46
  - chi-square distribution, 47
  - t-distribution, 46
  - t-distribution, the, 46
- smoothed data
  - after inverse Fourier transformation, 394
  - from application of trapezoidal smoother, 397
- smoothing, 380, 394
  - with designer transfer functions, 395–398
  - in the frequency domain with a trapezoidal smoothing function, 396
  - of noisy signal with wavelet transform, 412
  - operation, 383
  - time-domain and, 398
  - transfer function and, 394
- smoothing and filtering, 81
  - signal-to-noise ratio and, 81
- soft independent modeling by class analogy (SIMCA), 354, 364
  - classification in high dimensions, 353
  - quadratic discriminant analysis and, 354

- soft modeling, 345
- software, 465
  - curve-resolution algorithms and methods, 465
- software applications, 4
  - ANOVA using Excel, 39–40
  - MATLAB and, 4
  - Microsoft Excel, 37
- solution for the integration of ODEs, 222
- solvent interactions, 108–109
- sorting plastics for recycling, 365
  - pattern recognition techniques and, 365
  - principle component analysis, 368
  - Raman spectroscopy, 365
  - self-organizing map (SOM), 368
- spectra, 462
  - first-derivative, 116–117
  - pixels and, 463
  - second derivative, 116–117
  - of the UV-active components, 324
  - of the UV-inactive components, 325
- spectral profiles, 476
- spectral windows, 423
- spectroscopic-chromatographic data, 70–72
  - basic vectors, 71
- spectroscopic images, 461–462, 462
  - multivariate resolution and, 462
  - pharmaceutical quality control and, 462
  - resolution of, 463
  - structure of, 462
- spectroscopic measurement, 256
- spectroscopy, 477
  - applications of, 512
  - chemometrics and, 512
  - near infrared spectroscopy and, 512
- spectrum, 78, 195
- spectrum, techniques for obtaining, 195
  - energy-dispersive x-ray fluorescence spectrometry (ED-XRF), 195
  - fluorescence spectrometry, 195
  - NIR (near-infrared reflectance spectrometry), 195
  - nuclear magnetic resonance (NMR), 195
  - ultraviolet spectrometry (UV), 195
- square sum  $ssq$  of the residuals  $r$  as a function of two parameters, 224
- squared correlation coefficient (angle cosign)
  - between True and Estimated Profiles from DTLD and PARAFAC, 491
- squared correlation coefficient (angle cosign) of predicted and true X-Way Spectra from GRAM with added matrices, 488
- squared correlation coefficient (angle cosign) of predicted and true X-Way Spectra from GRAM with concatenated matrices, 488
- squared correlation coefficient (angle cosign) of predicted and true Y-Way Spectra from GRAM with added matrices, 488
- squared correlation coefficient (angle cosign) of predicted and true Y-Way spectra from GRAM with concatenated matrices, 487
- standard deviation, 10
  - population, 11, 13
  - relative standard (or coefficient of variation), 11
  - sample, 11
- standard normal distribution, 45
- standard normal variate (SNV) preprocessing, 85, 87
- standard normal variate (SNV) transforms, 83
- standard residuals of the stars data set based on a classical MLR and a robust LTS estimator, 179
- stars data, 180–181
- stars regression data set with classical and robust fit, 177
- statistical evaluation of data, 7–40
  - analysis of variance and, 27–30
  - common terms, 10
  - introduction to, 8
  - normal distribution, properties of, 14
  - outliers, 33–35
  - precision and accuracy, 12
  - robust estimates of central tendency and spread, 36–38
  - significance testing, 18–26
  - software and, 38–40
  - sources of error, 9
- statistical evolution of calibration models
  - obtained by least squares, 121
- statistical modeling, 265
- statistical properties, 3
  - calibration of response signals, 2
  - control of interfering factors, 2
  - measurement process errors, 2
  - modeling of complex multivariate signals, 2
- strong filtering of noisy data using quadratic polynomials, 405
- structure of the extended design matrix  $F$  for a second-order model with two process variables, 287
- structure of protein, 450
- student t-test
  - paired t-test, 21
  - standard t-test, 21
  - t-test with nonequal variance, 21
- subwindow factor analysis (SFA), 428–430, 429–430
  - algorithm, 431
  - application of, 430
- sum of squares ( $ssq$ ), 123–124, 234

summary data for the analysis of Pb in orchard leaves, 49  
 summary statistics for NIR calibration of water in water-methanol mixtures using one wavelength and a nonzero intercept, 125  
 sums of squares, 125  
 symmetric optimal designs, 290  
 symmetric optimal designs, central composite designs, 293–294  
 symmetric optimal designs, three or more levels in full factorial designs, 291–292  
   advantages, 290  
   symmetric optimal designs, two-level factorial designs, 290–294  
   disadvantages, 291  
 symmetry matrix, 349  
 systematic error or bias, 17

## T

t-distribution, 16  
   degrees of freedom and, 47  
 t-test  
   comparison of a sample mean with a certified value, 24  
   comparison of the means from two samples, 24  
   comparison of the means of two methods with different samples, 24  
 Taguchi, 296  
   Experimental Design Approach, 294  
   methods, 295  
   orthogonal designs, 319  
 Taguchi design, 297  
   for two controlled factors and three noise factors, 297  
 Taguchi, Genichi, 294  
   quality engineering and, 294  
 target factor analysis (TFA) and, 437  
 taxonomy based on chemical constitution  
   Africanized and European honeybees, 371  
   gas chromatography and, 371  
   principle component analysis, 372  
 three-factor principal component model for NIR spectra of water-methanol mixtures, diagram of, 142  
 three-way calibration, advantages of, 475  
 three-way calibration with hyphenated data, 475–499  
   alternating least-squares methods, 491–494  
   background, 476  
   caveats, 497  
   extensions of three-way methods, 495  
   figures of merit, 496  
   introduction to, 475  
   nomenclature of three-way data, 478  
   rank annihilation methods, 482–490  
   three-way data, 477  
     models of, 478–479  
 three-way data analysis, 443–444, 464, 477  
   calibration with, 477  
   constraints and, 445  
   examples of, 481  
   MCR-ALS decomposition method, 445  
   nontrilinear, 442  
   trilinear, 442  
   unfolding and, 441  
   unfolding or matricization, 441  
 three-way data, bilinear models for unfolded PCA and unfolded MCR, 444  
 three-way data, nomenclature of  
   data as orders, 478  
   data as ways, 478  
 time-domain filtering and smoothing, 398–405  
   filtering, 400–402  
   polynomial moving-average (Savitsky-Golay) filters, 403–405  
   smoothing, 398–399  
 time-domain processing of noisy data with moving-average filter, 402  
 time-domain smoothing of noisy data, 399  
 training set  
   for the honeybee sample, 372  
   jet fuel sample, 358  
   for plastics sample, 368  
 transfer functions for FIR filters as a function of window size, 403  
 translated experimental design for product P1, 329  
 translation of coded factor levels to real experimental levels for product P1, 328  
 trilinearity constraint in MCR bilinear models for three-way data, 446  
 true, best, and worst estimated X-way and Y-way analyte profiles from GRAM, 487  
 “true values”, 114  
 TUCK alternating least squares, 493  
 Tuckals, 493  
 Tucker3 model, 479–480, 493  
 two-way data, 476  
 two-way data matrix, augmentation of, 441  
 types of observations based on their robust distance (RD) and their Residual Distance (RD), 185  
 types of observations based on score distance (SD) and orthogonal distance (OD), 193

## U

uncertainty in resolution results, experimental error and, 448  
 uncertainty in resolution results, sources of, 446

ambiguity of recovered profiles, 446  
 experimental noise of the data, 446  
 unimodality, 434  
 Union of Pure and Applied Chemistry (IUPAC), 131  
 univariate calibration, 109, 118, 118  
   with an intercept term, 119  
   without an intercept term, 118, 118–119  
 univariate decision limit, 132  
 univariate detection limit, 133  
 univariate hypotheses testing, 48  
   inferences about means, 49  
   inferences about variance and the  
     F-distribution, 51  
 univariate MCD, 193  
 univariate normal distribution, 3  
 unknown prediction steps  
   prediction step, 143  
   validation step, 143  
 UV absorption, 323  
 UV-inactive components, 323  
 UV spectra, 323  
   of all nine pure components, 324  
 UV spectroscopy, 322  
 UV/visible spectrometer, 70

## V

v-fold CV, 115  
 validation set, 208  
 variables, 10, 11, 44  
   between-sample means, 28  
   collinear, 344  
   continuous random, 43  
   discrete random, 43  
   heteroscedastic, 130  
   homoscedastic, 130  
   residual variance and, 89  
   scaling, 78–79  
   selection, 135  
   within-sample means, 28  
 variance and the f-distribution  
   inferences about, 51  
 variance-covariance matrix, 53  
 viscose data set and standardized residuals  
   obtained with different estimators of location  
   and scale, 170  
 Visual Basic, 218

## W

water-methanol mixtures, 108  
 wavelengths, 462

wavelet, 406  
   components resulting for discrete wavelet  
     transform, 410  
   sine basis functions for orthogonal transforms  
     and, 408  
   truncation and, 412  
 wavelet analysis  
   Fourier analysis and, 408  
 wavelet-based signal processing, 406–413  
   discrete wavelet transform, 409–410  
   smoothing and denoising with wavelets,  
     411–413  
   wavelet function, 406–407  
   wavelet function, time and frequency  
     localizations of, 408  
 wavelet family, 407  
   vanishing moments and, 407  
 wavelet functions, 411  
   coiflet wavelet families, 411  
   Daubechies wavelet families, 411  
   symlet wavelet families, 411  
   time and frequency localizations of,  
     408  
 weak filtering of noisy data using quadratic  
   polynomials, 405  
 white noise, 222, 397  
   defined, 237  
 Wilson-Hilferty transformation, 192  
 window factor analysis (WFA), 427–428  
   steps of, 427–428  
 within-run, defined, 10  
 Wold, Herman  
   partial least squares (PLS) and, 352, 510  
 Wold, Svante  
   chemometrics and, 509  
   partial least squares (PLS) and, 510  
 Wynn-Mitchell and van Schalkwyk algorithms,  
   308

## X

X-way and Y-way profiles used to construct data  
 sets for each example, 481  
 Xerox, 294

## Z

Zeroth-Order reaction, 248–249  
   catalyst and, 248  
   macroscopically observed, 248