

Subject Index

Page references to *figures* and *tables* are shown in *italics*.

- Abbott, S. 24
- acetone-water
 pressure and temperature *vs.* column dimensions 73
 pure acetone, physical properties 3
 static permittivity 77, 201–4
 vapour pressure isothermal P-x-data 196
 viscosity and temperature 200
- acetonitrile-water
 pressure and temperature *vs.* column dimensions 73
 pure acetonitrile, physical properties 3
 static permittivity 77, 201–4
 TOF-MS 61–2, 62, 63
 vapour pressure isothermal P-x-data 57, 58–9, 193
 viscosity and temperature 65, 197
- alcohols
 HPLC-FID separation 162, 163
 longer chain 72–3, 78
- aldehydes and ketones, separation 78–80, 79, 80, 82, 82–3
- analyte stability
 critical criteria 154–7
 Damköhler number (Da) 149–50
 evaluation 150–2
 influence of the stationary phase 152–4
- Antia and Horváth form of the van Deemter equation 128
- antibiotic drugs 136, 139, 152
- back-pressure regulation, phase transition prevention 22–3, 54, 56–60
- capillaries
 polyetheretherketone (PEEK) 36, 106
 stainless-steel 28, 36, 36–7, 39
- capillary and nano HPLC 184–9, 187
- capillary-TOF-MS 184–9
- carbon columns *see* pure graphitized carbon columns
- carbon-clad zirconia 100–1
- Carr, P. W. 20
 Carr and Thompson rules, analyte stability 154–6
- Chester, T. L. 69
- chromatographic techniques
 gel-permeation chromatography (GPC) 30–1, 67, 104–5
 high-pressure liquid chromatography (HPLC) 1, 55
 high-temperature liquid chromatography (HTLC) 1–8
 reversed-phase liquid chromatography (RP-HPLC) 2–3
 size-exclusion chromatography (SEC) 104–5
 two-dimensional liquid chromatography 189–91
 unified chromatography 69
- column ageing test procedure 88
- column bleed 88–91

- column design *see* stationary phases
- column dimensions, effect on mobile phase pressure 72–4, 73
- column heating
- air-bath ovens 17, 24, 40, 46–8
 - block-heating ovens 41–2, 42, 48–9
 - water-jacket ovens 24, 40–1, 41, 48
 - see also* stationary phase, general aspects
- column systems 20–1
- cytostatic drugs 136, 139, 152
- Damköhler number (Da) 149–50
- detectors 21–2, 90
- optimization 143–6
 - see also* specific detector system
- dielectric constant *see* static permittivity
- diffusion coefficient 71–2
- Dortmund Data Bank (DDB) 83–5
- drug screening 177–80
- DryLab[®] software 80, 118, 120
- efficiency, temperature effect on 128–31
- plate height *vs.* velocity 129
- electrospray-ionization mass spectrometry (ESI-MS) 177–80, 184–5
- eluent cooling 18, 42–3, 44
- Engelhardt, H. 1
- Engelhardt test 69
- equations
- Hagen–Poiseuille 67–8
 - Knox and Thijssen, resolution 118
 - peak area 150
 - selectivity, temperature effects 125
 - van Deemter equation and derivatives 6, 71, 128–9
 - Antia and Horváth form 128
 - van't Hoff 119–23
- ethanol-water
- pressure and temperature *vs.* column dimensions 73
 - pure, physical properties 3
 - static permittivity 77, 201–4
 - supercritical conditions 117
 - vapour pressure isothermal P-x-data 195
 - viscosity and temperature 199
- evaporative light-scattering detection 22, 61, 90
- flame ionization detection (FID) 159–64
- fluorescence detection
- and eluent temperature 18, 21–2
 - vs.* UV 17, 59
- frictional heating *see* viscous heat dissipation
- future prospects
- capillary and nano HPLC 184–9, 187
 - pellicular particles 182–3
 - two-dimensional liquid chromatography 189–91
- gel-permeation chromatography (GPC) 30–1, 67, 104–5
- graphitized carbon columns *see* pure graphitized carbon columns
- Hagen–Poiseuille equation 67–8
- heating and cooling systems
- basic aspects 15–20
 - block temperature *vs.* heating rate 46
 - column heating and oven systems 40–2, 46–50
 - mobile phase
 - column considerations 116–18
 - post-column cooling 18, 42–3, 44, 47
 - preheating 17–18, 19, 25–39 - special requirements 114–16
 - temperature programming 43–6, 135–40
 - see also* column heating; temperature
- height equivalent to a theoretical plate $H(u)$ (HETP) 6–7, 70–1
- vs.* linear velocity 7, 70
- Heinisch, S. 2
- Hesse, G. 1
- high-pressure liquid chromatography (HPLC)
- definition 1
 - typical system 55
- high-temperature liquid chromatography (HTLC)
- definitions 1–4
 - rational for 5–8
 - terminologies, alternative 2

- honey, adulterated, HPLC-IRMS separation 174
- HPLC-FID *see* flame ionization detection
- hyphenation techniques 8, 42, 77, 158–9
- capillary-TOF-MS 184–9
- electrospray-ionization mass spectrometry (ESI-MS) 177–80, 184–5
- flame ionization detection (FID) 159–64
- isotope ratio mass spectrometry (IRMS) 168–74
- LC Taste[®] 174–7, 176
- nuclear magnetic resonance (NMR) spectroscopy 164–8
- two-dimensional liquid chromatography 189–91
- isocratic separation, temperature gradient in 135–40
- isopropanol-water
- heptane–isopropanol separation 102
- isopropanol-water system 72
- pressure and temperature *vs.* column dimensions 73
- pure isopropanol, physical properties 3
- static permittivity 77, 201–4
- vapour pressure isothermal P-x-data 195
- viscosity and temperature 199
- isothermal and isocratic separation, steroids 131–5, 132–4
- isotope ratio mass spectrometry (IRMS) 168–74
- ketones and aldehydes, separation 78–80, 79, 80, 82, 82–3
- Knox and Thijssen resolution equation 118
- LC Taste[®] 174–7, 176
- metal oxide columns *see* zirconium dioxide-based stationary phases
- methanol-water
- high temperature column purging 92
- pressure and temperature *vs.* column dimensions 73
- pure methanol, physical properties 3
- static permittivity 77, 201–4
- vapour pressure isothermal P-x-data 58, 194
- viscosity and temperature 66, 198
- mobile phase
- post-column cooling 42–3
- preheating 17–18, 19, 25–39
- system pressure, temperature and column dimensions 52–64
- temperature and vapour pressure 52–64
- Molnár Institute for Applied Chromatography 123
- nano HPLC 184–9, 187
- Neue test 69, 92
- non random two liquids (NRTL) model 84
- nuclear magnetic resonance (NMR) spectroscopy 164–8
- on-flow NMR 165
- ovens systems *see* column heating
- peak area
- analyte stability evaluation 150–2
- equation 150
- vs.* temperature 151, 152
- see also* van't Hoff equation
- peak splitting 33
- pellicular particles 182–3
- Peltier element, cooling 42–3
- pesticides, capillary HPLC separation 187–9, 188, 189
- pharmaceutical compounds, separation 68
- phase transition prevention
- back-pressure regulation 22–3, 54, 56–60
- using a restriction capillary 61–4
- phenols, HPLC-FID separation 162
- plate height (H)
- plate height *vs.* velocity 129
- see* height equivalent to a theoretical plate $H(u)$ (HETP)

- PLRP-S column
 on-column degradation of thalidomide 155
 retention factors 122
- polar retention effect on graphite (PREG) 106–7
- polycyclic aromatic hydrocarbon (PAH)
 effect of eluent preheating 17–18, 19
 peak area vs. eluent temperature 42–3, 44, 144
 retention times 121
 separation conditions 16
- polyetheretherketone (PEEK)
 capillaries 36, 106
- polymeric stationary phases 104–5
- predictive Soave-Robinson-Kwong (PSRK) model 84
- preheating, mobile phase
 effects 17–18, 19
 efficiency assessment 37–9
 eluent temperature measurement 38–9
 technical considerations 35–7
 thermal mismatch broadening 25–31
 viscous heat dissipation 26, 31–5
see also heating systems; temperature
- propanol *see* isopropanol-water
- pure graphitized carbon columns 105–7
 polar retention effect on graphite (PREG) 106–7
- resolution
 Knox and Thijssen equation 118
 temperature effect on 15–17, 118–19
- restriction capillary
 kinetics and column pressure, practical implications 69–74
 for phase transition prevention 61–4
 viscosity, practical implications 67–9
- retention
 enthalpy 119–25
 experimental vs. simulated retention times 121, 124
 linear/non-linear responses 120
 retention factor 122
 temperature effects 119–25
 temperature-gradient separation 124, 137, 139
- reversed-phase liquid chromatography (RP-HPLC), definition 2–3
- selectivity, temperature effects 125–7
 equation 125
 van't Hoff plots 127
- silica-based stationary phases 93–7
 characteristics 87
 pellicular particles 182–3
 reversed-phase 5, 21
 temperature stability 5, 20–1, 94–6
 temperature/pH interaction 93–7
- size-exclusion chromatography (SEC) 104–5
- Smith, Roger M. 2, 8
- solute retention 33–4, 34
- solvents, pure
 physical properties 3
 static permittivity 76
- solvents, static permittivity of mixtures 77
- 'sputtering' 62
- stainless-steel capillaries 28, 36, 36–7, 39
- static permittivity (DEC) 8, 201–4
 influence of temperature 8, 75–7, 75–80, 84, 158, 201–4
- stationary phase, general aspects
 batch-to-batch reproducibility 87
 column ageing test procedure 88
 column bleed 88–91
 columns and temperature maxima 110
 design difficulties 88–9
 detector response in different columns 90–1
 high temperature degradation 91–3
 influence on analyte stability 152–4
see also column heating; specific stationary phase components
- steroids
 isothermal and isocratic separation 131–5, 132–4
 peak areas 151
 thermal mismatch broadening 25–6
- stop-flow NMR 165
- sulfonamides
 isothermal and isocratic separation 140–3, 141–2

- retention times 120, 123
- temperature-gradient separation 124, 126
- system
 - basic set-up 15–23
 - capillary and nano HPLC 186–9, 187
 - continuous-flow for ESI-MS 177–80, 178
 - general requirements 8–9
 - HPLC-FID 160–1
 - HPLC-IRMS 169
 - HPLC-NMR 165–6, 167
 - LC Taste[®] 174–7, 176
- temperature
 - columns and temperature maxima 110
 - effect on analyte stability 149–57
 - effect on efficiency 128–31
 - effect on resolution 15–17, 118–19
 - effect on retention 119–25
 - effect on selectivity 125–7
 - high temperature column
 - degradation 91–3
 - mobile phase temperature and vapour pressure 52–64
 - range 4–5
 - temperature programming 43–6, 135–40
 - see also* heating and cooling systems
- temperature-gradient separation
 - simultaneous temperature and solvent gradient separation 140, 143
 - sulfonamides and uracil 124
 - using HPLC-IRMS 171–2, 173
- testosterone and related hormones
 - HPLC-IRMS separation 172, 173
 - retention and peak shape 27
 - thermo-responsive separation 108
- tetrahydrofuran (THF)-water
 - miscibility gap 116–17
 - pressure and temperature *vs.* column dimensions 73
 - pure tetrahydrofuran, physical properties 3
 - solvent system 81, 81–3
 - static permittivity 77, 201–4
 - vapour pressure isothermal P-x-data 60, 194
 - viscosity and temperature 198
- thermal mismatch 25
 - broadening 25–31
 - column thermal effects 26
 - contact heating 28–9
 - preheating coil length and flow rate 26–31, 27, 28, 29
- thermo-responsive stationary phases 107–9, 108–9
- Thompson, J. D., Carr and Thompson rules, analyte stability 154–6
- time-of-flight mass spectrometer (TOF-MS) 61–2, 62, 63, 68
- titanium dioxide stationary phases
 - temperature and chemical stability 101–4, 102–3
 - see also* zirconium dioxide-based stationary phases
- trimethoprim, non-linear retention 120
- two-dimensional liquid chromatography 189–91
- unified chromatography 69
- UV *vs.* fluorescence detection 17, 59
- van Deemter equation and derivatives 6, 71, 128–9
- van't Hoff equation 119–23
- vapour pressure
 - data 193–6
 - liquid water 53
 - and temperature 52–64
- viscosity
 - data 197–200
 - influence of temperature 7–8, 64–74
 - maximum 7–8
- viscous heat dissipation
 - definition 31–2
 - effect on flow profile 26, 32–5
- water
 - binary mixes
 - pressure and temperature *vs.* column dimensions 73
 - static permittivity 77, 201–4
 - tetrahydrofuran (THF)-water
 - miscibility gap 116–17

- vapour pressure isothermal P-x-data
193–6
- viscosity and temperature
197–200
- pure
 - physical properties 3
 - static permittivity and temperature
75
 - vapour pressure 53
 - viscosity and temperature 64
 - water only separation 21
- zirconium dioxide-based stationary phases
 - as alternative to silicates 20
 - and analyte stability 150–2
 - elution of sulfathiazole on 153
 - on-column degradation of thalidomide 154
 - polymer-coated 98–101
 - retention mechanisms 87–8
 - steroids, isothermal and isocratic separation 131–5
 - temperature stability 97–101, 99