



- Aerosols, 167–71
 future trends, 187
 identification procedures, 184–6
 particle phase analysis, 175–82
 sources of, 170–1
- Air, 167–71
 future trends, 187
 identification procedures, 184–5
 in situ analyses, 182–4
 sources and fate of organic compounds, 170–1
 volatile organic compound analysis, 172–5
See also Aerosols
- Allergens, 202–4
- Analysis of variance (ANOVA), 103, 119
- Analytical Data Interchange (ANDI) standards, 81
- Analytical Information Markup Language (AnIML), 81
- Arabica coffee beans, 216–17, 233–6
- Aromatic organohalogenated pollutants, 244–59
- Atmosphere, *See* Air
- Atomic emission detector (AED), 36–8
- Autocovariance functions (ACVF), 109
- Baseline correction, 90–2
- Beef aroma, 236–7
- Blob detection, 92–5
- Bubble plot, 185
- Carbazoles, 161
- Chemical identification, 95–100
 air and aerosol analyses, 184–6
 by retention time, 95–7
 multivariate methods, 97–9
 Smart Templates, 99–100
- Chemometrics, 107–8, 164
 classification, 116–18
 computational considerations, 121
 data mining, 118–21
 deconvolution, 111–15
 feature selection, 119–21
 peak finding, 109–11
 retention calculation and prediction, 108–9
 separation optimization, 109
- Chiral compounds analysis, 204–10
 pollutants, 274–7
- Chlorfenvinphos, 270
- Chlorpyrifos, 10
- ChromaTOF, 120
- Chromatographic response parameters, 52–64
 estimation of retention parameters, 57–62
 holdup time, 62–3
 peak width, 63–4
 retention time, 52–7
- Chromatographic structure, 11, 51, 71–4
- Cod liver oil, 217, 225, 226–8, 229
- Coding, 80–1
- Coffee beans, 216–17, 233–6
- Colorization, 82–4
- Colour plot, 51
- Column combinations, 20–6, 43
 GC column dimensions, 23–4
 optimisation, 24
 orthogonality principle, 20–3
 stationary phases, 24–6

- Complex samples, 124
Comprehensive, 51
Comprehensive three-dimensional (3D) systems, 125
 additional dimensions prior to GC×GC, 135–6
 analysis time, 126–9
 coarse prefractionation methods, 136–7
 hardware solutions, 133–5
 LC dimensions, 137–44
 operational approaches, 129–33
 separations in space and time, 125–6
Comprehensive two-dimensional chromatography, 8–13, 15, 51, 123–5
Computer Language for Identifying Chemicals (CLIC), 99
Cosmetics, 202–4
Crude oil, 157, 158
 See also Petrochemicals
Cryogenic modulators, 29–34
Data acquisition, 13, 79–81
 digitization and coding, 80–1
 file formats, 81
 modulation, 79–80
 sampling, 79–80
Data processing, 89–95
 baseline correction, 90–2
 peak detection, 92–5
 phase correction, 89–90
Data visualization, 13, 78, 82–9, 101
 flicker visualization, 101–2
 graphical overlays and annotations, 87–9
 image visualizations, 82–5
 one-dimensional visualizations, 86
 text and tabular visualizations, 86–7
 three-dimensional visualizations, 85–6
Databases, 104
Deconvolution, 110, 111–15
Detectors, 36–43, 44
 element selective detectors, 36–8
 flame-ionization detector (FID), 36, 161
 mass spectrometer (MS), 38–43
Di-naphthenes, 151
Diesel, 92, 114–15, 157, 158, 159, 161
 See also Petrochemicals
Digitization, 80–1
4,6-Dimethyl-dibenzothiophene, 161
Dimethyltetrabromobisphenol-A (Me-TBBP-A), 257–9
Direct thermal desorption (DTD) system, 175–9
Dissolved inorganic carbon (DIC), 158
Dolphin blubber, 259, 260
Dual-stage liquid CO₂ cryogenic modulator, 30–1
Edible oils, 142
 See also Olive oil
Egyptian geranium essential oil, 199
Electron-capture detector (ECD), 36, 110
 environmental applications, 245, 249, 256–9, 261–3, 264, 276, 277
Electron ionisation (EI), 42–3
Elongated clusters, 72, 74
Enantiomeric composition, *See* Chiral compounds analysis
Environmental samples, 243–4
 group-type analysis, 263–5
 See also Pollutants
Essential oils, 189–90
 chiral compounds analysis, 205–10
 Egyptian geranium, 199, 201
 future trends, 210–11
 lavender, 73, 197, 205–7
 lemon, 196
 peppermint, 205–7
 use of retention indices, 200–2
 vetiver, 195
 See also Plant extracts
Fatty acid methyl esters (FAMES), 125, 142–4, 217, 223–7, 238
Fatty acids, 218–23
Feature selection, 109, 119–21

- File formats, 81
- Fingerprinting, 103–4, 125
- Fisher ratio (*f*-ratio), 119–20
- Flame-ionization detector (FID), 36
 - air and aerosol analysis, 172, 175, 182
 - environmental applications, 245–9, 269, 272–3
 - food constituent analysis, 225–6
 - petrochemical analysis, 161
 - plant extract analysis, 190–4
- Flame retardants, 257–9
- Flavours, 230–7
- Flicker visualization, 102
- Fly ash, 256
- Food constituents, 215–40
 - food flavour applications, 230–7
 - lipids, 218–30
- Fragrances, 190
 - allergens, 202–4
 - future trends, 210–11
 - use of retention indices, 200–2*See also* Essential oils
- Fungicides, 269
- Fuzzy differences, 102

- Gasoline, 98, 161, 162
 - See also* Petrochemicals
- GC×GC, *See* Two-dimensional gas chromatography
- Generalized rank annihilation (GRAM) method, 111–13, 117
- Geranium essential oil, 199
- Globular clusters, 72
- Gradient-Based Value Mapping (GBVM), 83–4
- Graphical overlays, 87–9
- Group separation, 131–2
 - petrochemicals, 151–8
- Group-type analysis, 124–5, 136
 - air and aerosols, 185
 - environmental applications, 263–5

- Heating modulators, 27–9
- Heptadecanoic acid, 269
- Herring oil, 223, 224

- Hexabromocyclododecane (HBDE), 257
- Hexyl cinnamaldehyde, 204
- High-resolution mass spectrometry (HRMS), 43
 - environmental applications, 252
- High-resolution time-of-flight mass spectrometry (HRTof MS), 176–80
 - plant extract analysis, 197–9
- Holdup time, 62–3
- Honey, 237
- Hydrodesulfurization (HDS) feed, 161, 163

- Identification, *See* Chemical identification
- Image visualizations, 82–5
 - colorization, 82–4
 - navigation, 84
 - qualitative analysis, 84–5
 - rasterization, 82
- Indoles, 161
- Informational similarity (IS), 68
- Information systems, 104
- Intermediate collection of fractions, 130
- Isotope ratio mass spectrometry (IRMS), 207
- Isovolatility curves, 56–7

- Lanolin, 229–30, 231
- Lavender essential oil, 73, 197, 205–7
- LC×GC×GC, *See* Comprehensive three-dimensional (3D) systems
- Lemon essential oil, 196
- Linear retention index (LRI), 54–5, 96
- Lipids, food, 218–30
 - See also* Fatty acid methyl esters (FAMES)
- Liquid chromatography (LC):
 - food analysis, 238–9
 - silver phase LC (AgLC), 142–4*See also* Comprehensive three-dimensional (3D) systems
- Longitudinal Modulated Cryogenic System (LMCS), 29, 30

- Ma Huang, 209
- Mass-spectrometric (MS) detector, 10, 38–43, 110
- high-resolution mass spectrometer (HRMS), 43
 - plant extract analysis, 190–4, 197–9
 - quadrupole mass spectrometer (qMS), 42–3, 176–80, 199
 - rapid-scanning quadrupole MS instruments, 11
- See also* Time-of-flight mass spectrometric (ToF MS) detector
- Mass-to-charge ratio, 99
- Mathematical resolution, 111–15
- Maximum residue levels (MRLs), 245
- Melaleuca alternifolia*, 208–9
- Menhaden oil, 224–5
- 1'Methyl-1,2'bipyrroles (MBPs), 259
- Micro electron-capture detector (ECD), 9
- Micropollutants, *See* Pollutants
- Mineral oil, 139–42
- Modulation, 9, 62, 79–80
- Modulation ratio (M_R), 207
- Modulator block, 18
- Modulators, 9, 16–19, 26–35, 44
- thermal, 27–34
 - valve-based, 34–5
- Mono-naphthenes, 151
- Multi-dataset analyses, 100–4
- databases and information systems, 104
 - sample comparison, classification and recognition, 101–3
- Multidimensional gas
- chromatography (MDGC), 4, 5–8
 - chiral compound analysis, 205–7, 274
 - environmental applications, 274
- See also* Comprehensive three-dimensional (3D) systems; Two-dimensional gas chromatography (GC×GC)
- Multidimensionality, 4–5, 51
- Multi-way partial least squares (N-PLS), 118–19
- Multi-way principal components analysis (MPCA), 118
- Multiplex gas chromatography, 27
- National Institute of Standards and Technology (NIST), 97
- Navigation, 84
- Negative chemical ionisation (NCI), 42–3
- Nitrogen-phosphorus detector (NPD), 38
- air and aerosol analysis, 176–80
 - environmental applications, 269
 - food constituent analysis, 236
- Nitrogen-specific chemiluminescence detectors (NCDs), 38, 161
- Noise, 124
- Nomenclature, 50–2
- Nonaromatic organohalogenated pollutants, 259–63
- Nonhalogenated pollutants, 266–73
- pesticides, 266–72
- Nonylphenolpolyethoxilates, 273
- Oil spill, 102, 103
- Olive oil, 21, 22, 223, 230–3
- One-dimensional (1D) gas chromatography, 3–4, 5
- One-dimensional visualizations, 86
- Organohalogenated pollutants, 244–65
- aromatic, 244–59
 - nonaromatic, 259–63
- Orthogonality, 20–3, 67–71, 74–5
- measurement of, 68–71
- Oversegmentation, 94
- Paraffins, 151
- Parallel factor analysis (PARAFAC), 111–15, 117, 120
- Partial least squares (PLS), 118
- multi-way extension (N-PLS), 118–19

- Particulate matter (PM), 168–9
analyses, 175–82
fates of, 171
sources of, 170–1
- PBDEs, 257–9, 265
- Peak capacity, 65–7, 69, 126–8
practical, 69
- Peak detection, 92–5, 109–11
- Peak width, 63–4
- Pelargonium graveolens*, 199, 201
- Pentachlorophenol, 269
- Pepper, 236
- Peppermint essential oil, 205–7
- Percent synentropy (PS), 69
- Perfumes, *See* Fragrances
- Pesticides, 266–72, 278
- Petrochemicals, 149
group-type separation, 151–8
overview of studies, 154–6
PIONA analyses, 164–5
quantification, 161–4
sample dimensionality, 150–1
target analysis, 161
- Phase correction, 89–90
- PIONA analyses, 164–5
- Plant extracts:
chromatography applications, 190–7
mass spectrometric detection, 197–9
See also Essential oils
- Pogostemon cablin Benth.*, 197
- Pollutants, 244–77
aromatic organohalogenated
pollutants, 244–59
chiral pollutant analysis, 274–7
environmental samples, 243–4
group-type analysis, 263–5
nonaromatic organohalogenated
pollutants, 259–63
nonhalogenated pollutants, 266–73
- Polychlorinated biphenyls (PCBs), 6,
7, 12, 21, 245, 249–57, 274–7
- Polychlorinated dibenzo-*p*-dioxins
and furans (PCDD/Fs), 43, 245,
250–7, 277
- Polychlorinated n-alkanes (PCAs),
262–3, 265, 277
- Polycyclic aromatic hydrocarbons
(PAHs), 176–80, 272–3, 278
- Practical peak capacity, 69
- Prefractionation, 136–7
- Principal component analysis (PCA),
103, 116–18, 121
multi-way (MPCA), 118
- Principal-component discriminant
analysis (PCDA), 104
- Principal component regression
(PCR), 118
- Profiling, 125
- Pseudocolorization, 83
- Pyrazines, 235
- Quad-jet dual-stage modulator, 31–4
- Quadrupole mass spectrometer
(qMS), 42–3, 199
air and aerosol analysis, 176–80
environmental applications, 256–9,
272, 273
food constituent analysis, 230–2,
235
fragrance analysis, 204
- Qualitative analysis, 84–5
- Quantification, 100–2
petrochemicals, 161–4
- Rapid-scanning quadrupole mass
spectrometric instruments, 11
See also Quadrupole mass
spectrometer (qMS)
- Rasterization, 82
- Real-time operation, 130–2
- Reformulyzer, 164, 165
- Resolution, 64–5
- Response, 52
See also Chromatographic response
parameters
- Retention index (RI), 53–4
determination of, 55–7
use in essential oil and fragrance
analysis, 200–2
- Retention time, 52–7
calculation and prediction, 108–9
chemical identification by, 95–7

- dependence on chromatographic conditions, 59–62
- estimation of retention parameters, 57–62
- retention-time windows, 96–7
- Reversed selectivities, 141
- Reverse-type column combinations, 1, 21–3, 157
- Roast beef aroma, 236–7
- Sample classification, 100, 102–4, 116
- Sample comparison, 100, 102
- Sample dimensionality, 124, 150
 - apparent, 150
 - petrochemicals, 150–1
 - required, 150
 - See also* Multidimensionality
- Sample prefractionation, 136–7
- Sample query, 100
- Sample recognition, 100
- Sampling, 79–80
- Secondary organic aerosol (SOA), 168–9
- Selectivity, 124
- Semivolatiles organic compounds (SVOCs), 168, 171
- Separation:
 - in space, 126
 - in time, 126
 - ordered, 150–1
- Separation efficiency parameters, 64–74
 - chromatographic structure, 71–4
 - orthogonality, 67–71
 - peak capacity, 65–7
 - resolution, 64–5
- Signal ratio method, 120–1
- Signal-to-noise (S/N) ratio, 113
- Silver phase LC (AgLC), 142–4
- Smart Templates, 99–100
- Split-flow GC×GC, 226–8
- Stationary phases, 24–6
- Stop-flow operation, 130, 132
- Strawberry, 236
- Structured chromatograms, 11, 21
- Sulphur chemiluminescence detector (SCD), 10, 36–8, 161
- Sweeper modulator, 28–9, 161
- Synentropy, 69
- System dimensionality, 124
- Tabular text visualizations, 86–7
- Target compound analysis, 124
 - petrochemicals, 161
- Tea tree, 208–9
- Template matching, 96–7
 - Smart Templates, 99–100
- Terpenoids, 204
- Tetrabromobisphenol-A (TBBP-A), 257–9
- Text visualizations, 86–7
- Thermal desorption modulation (TDM), 27–9
- Thermal modulators, 27–34
 - cryogenic modulators, 29–34
 - heating modulators, 27–9
- Thin-layer chromatography (TLC), 126
- Three-dimensional visualizations, 85–6
 - See also* Comprehensive three-dimensional (3D) systems
- Time-of-flight mass spectrometric (ToF MS) detector, 10, 38–41
 - air and aerosol analysis, 170, 175–6, 180–2, 185, 187
 - environmental applications, 252–6, 261–3, 270–2, 277–8
 - food constituent analysis, 228–30, 233–7
 - high-resolution (HRToF MS), 176–80
 - plant extract analysis, 197–9, 201
- Total intensity count (TIC), 82
- Toxaphene, 259–61, 265, 274, 277
- Triacylglycerides (TAGs), 141, 142–4, 238
- Two-dimensional gas
 - chromatography (GC×GC), 5–8
 - comprehensive two-dimensional chromatography, 8–13, 15, 51, 123–5

- split-flow GC×GC, 226–8
- use in 3D systems, 125–9
- Undersegmentation, 94
- Valve-based modulators, 34–5
- Vanilla, 233, 234
- Variance, 117–18
- Vetiver essential oil, 195
- Visualization, *See* Data visualization
- Volatile organic compounds (VOCs), 167–8
 - fates of, 171
 - oxygenated (oVOCs), 176
 - sampling methods, 171–5
 - sources of, 170–1
- Wine, 237
- Wrap-around, 11, 18, 23–4, 51