

## INDEX

## A

- Acenaphthene, 553
- Acetone  
sorption isotherm for, 382f  
volume of absorbed, 383r
- Acid retardation, 475-480, 476f-479f  
LiCl/HCl in test for, 477-479, 478f  
mechanism of, 447  
NaCl/HCl in test for, 475-476, 476f  
process in ion exchangers of, 448
- Activated carbons  
advantages of, 371  
shortcomings of, 372
- Activity coefficients of electrolytes,  
446-449
- Adsorption isotherms  
BDVT classification for, 374  
BET equation for, 374-377  
hysteresis loop in, 77-81  
Kelvin equation for, 76-77  
sorption isotherms for nitrogen, 374,  
375f, 376f, 379, 380  
specific surface area calculations from, 377
- Aerosol OT (AOT)  
formation of gigaporous texture with,  
119-123, 121f-124f  
molar ratio to water of, 119-120  
specific inner surface area as function of  
molar ratio to water, 120, 121f  
pore diameter and volume as function of  
molar ratio to water, 120, 121f
- Ag<sup>+</sup>, 443, 444f  
radius for, 462t
- Al<sup>3+</sup>, radius for, 460t, 462t
- Aldehydes, wine quality with, 546
- Aliphatic alkylamines, 551
- Amberchrom GC-161, pre-concentration  
of phenolic compounds with, 530, 532
- Amberlite XAD-2  
cephalosporin C sorption with, 436  
phenols sorption with, 427
- Amberlite XAD-4, 418  
caffeine extraction with, 432, 432t  
cephalosporin C sorption with, 436  
chlorogenic acid removal with, 432, 432t  
furfural sorption with, 436  
gasoline sorption, 423, 423t  
hydrogen storage capacity of, 606t  
hypercrosslinked sorbents v., 439-440  
lecithin sorption with, 422, 422t  
lipophilic marine toxins accumulated  
by, 552  
phenols sorption with, 424, 424f, 426f, 427  
pre-concentration of phenolic  
compounds with, 531, 532, 533t  
sorption isotherms for *n*-valeric acid  
onto, 417, 417f, 418t
- Amberlite XAD-7  
deswelling, 291-293, 292f
- Amberlite XAD-16  
caffeine extraction with, 432, 432t  
cephalosporin C sorption with, 436  
citrus juice de-bittering with, 434  
hydrogen storage capacity of, 606t
- Amberlite XAD-180, cephalosporin C  
sorption with, 436
- $\alpha$ -Amino acids, 418-419, 421  
hydrolysis of methyl esters of, 596t
- Aminophenols, separation of  
dihydroxybenzenes and  
phenylenediamine in RP mode,  
515, 516
- Aminosine, 438
- Ammonium acetate, resolution of, 500, 501f
- Amperozide, 509, 509
- Anion exchange resins Styrosorbs,  
595-597, 596r
- Anions, 448-449
- Anthraquinone sulfonates, 547, 547t
- AOT, *See* Aerosol OT (AOT)
- Aromatic amines, 550t
- Aromatic compounds, separation in RP  
mode, 515, 516f, 519f
- Aromatic sulfonates  
anthraquinone sulfonates, 547, 547t  
benzene sulfonates, 547, 547t  
naphthalene sulfonates, 547, 547t  
stilbene sulfonates, 547, 547t

Artificial extracorporeal kidney, 582–583, 583f

Atomic force microscopy (AFM),  
nanosponges in, 312–313, 313f

## B

Ba<sup>2+</sup>, radius for, 460t, 462t

Base retardation, 475–480, 476f–479f

LiCl/LiOH in test for, 477–479, 479f

NaCl/NaOH in test for, 476–477, 477f

Bayer AG, 355

BDDT classification, *See* Adsorption isotherms

Be<sup>2+</sup>, radius for, 460t, 462t

BET equation, *See* Adsorption isotherms

Benzene sulfonates, 547, 547t

BetaSorb<sup>TM</sup>, 579–580

Bio-Gel A, 448

Bio-Gel P, 449

Biporous hypercrosslinked polystyrene  
networks, 268–274, 269t, 271f

1,4-Bis(*p*-chloromethylphenyl)-butane  
(CMB), 171, 203t, 241t, 206f

1,4-Bis-(chloromethyl)-diphenyl (CMDP),  
82t, 171–172, 197f, 203t, 214f, 217

Bismuth, determination in lead-lithium  
alloy, 512

Bond Elut-ENV, food analysis with, 546

Branched macromolecules, 297, 298f

Brunauer, Emmett, and Teller equation  
(BET equation), 73, 74f, 374–377, 378t

## C

Ca<sup>2+</sup>, 461

radius for, 460t, 462t

Caffeine extraction, coffee beans, 432–433,  
432t

Calcium chloride (CaCl<sub>2</sub>) elution profiles  
for, 452–453, 452f, 453f

CaCl<sub>2</sub>/HCl separation, 467t

phase distribution for, 470–474, 470f, 472f

Capillary electrochromatography (CEC),  
138

mobile phase electroosmotic flow,  
141–142, 142f

Capillary monolithic columns, 128–146,  
621

flow-through transport pores in, 138

gas adsorption chromatography with, 144  
glycidyl methacrylate rods/disks in, 143  
hypercrosslinked polystyrene monoliths,  
621

porous molded monolithic rods as separating  
media, 135, 138–145, 140f, 142f

separation of polystyrene standards in,  
139, 140f

silica-based monolithic columns, 144

Carbon dioxide, sorption parameters, 378t

Carbonaceous sorbents, 611–615, 612t, 613t  
porous structure of, 612t

Cations, 448–449

Cauliflower texture, 98, 99f, 101

*n*-C<sub>8</sub>F<sub>18</sub>, sorption isotherm, 387–388, 389f,  
390f

Cephalosporin C, sorption of, 435–436, 436f

Cd<sup>2+</sup>, 441, 444

Zn<sup>2+</sup>/Cd<sup>2+</sup>/Pb<sup>2+</sup> separation, 511, 512f

Ce<sup>3+</sup>, radius for, 460t, 462t

CHA-101, 438

CHA-111, sorption of organic compounds  
with, 436

Chemical groups in hypercrosslinked  
polystyrene, 188–192

Chemviron F-400, pesticides sorption with,  
431

Chloroform, removal from industrial  
wastewaters, 428, 429f

Chlorogenic acid removal, Amberlite  
XAD-4 for, 432, 432t

Chloromethylated polystyrene (CMPS)  
properties of, 303t

Chromalite 5HGN, 514f, 515, 516f, 517,  
518f, 519f

Chromatography

gas, 144, 402–407, 403t, 404t, 406t,  
gel permeation, 455

high performance liquid  
chromatography, 455

inversed size-exclusion, 82–83, 251–255,  
252t, 253t, 254f, 256f

mixed mode chromatography, 518, 519f  
perfusion, 126

quasi-normal phase, 515–518, 517f

reversed phase, 513–515, 514f, 516f

size-exclusion, 306–308, 308f, 445–501

- Citrus juice de-bittering, 434–435, 435t  
 MN-100 for, 434  
 MN-102 for, 434–435, 435t  
 MN-150 for, 434
- CMM, *See* Tris-(chloromethyl)-mesitylene
- $\text{Co}^{2+}$ , 441, 444  
 radius for, 460t, 462t
- COF, *See* Covalent organic frameworks
- Coils macromolecular, 297, 298f
- Column packing material for HPLC,  
 503–521  
 macroporous polystyrene v. silica-based  
 503–507, 506f, 507f  
 benefits of polystyrene-based, 505  
 drawbacks to silica-based materials, 503  
 hypercrosslinked polystyrene as  
 restricted-access adsorption material,  
 507–510, 509f
- Copolymerization anionic, 29–56, 34f–39f,  
 42f, 43f, 45t, 46t, 48f, 49t, 50f, 51f, 56  
 free radical, 3–28, 5f, 8t, 9t, 10f, 11f  
 Schwan and Price equations for  $r$ -values  
 $m$ , 8–9  
 monomer reactivity ratio in, 4–12, 5f, 8t,  
 9t, 10f, 11f
- Covalent organic frameworks (COF),  
 347–354  
 composition of, 347–348  
 interpenetrating MOF, 349–350  
 structure of, 348
- Cross-linking agents,  
 1,4-bis(*p*-chloromethylphenyl)-butane,  
 171  
 bis-chloromethyl derivatives of aromatic  
 hydrocarbons, 171  
 bis-(chloromethyl)-biphenyl, 171–172  
 carbon tetrachloride, 178  
 chloroform, 178  
 cyanuric chloride, 178  
 dimethylformal, 175  
 double bonds in styrene-DVB  
 copolymers for, 175–177  
 halogen anhydrides of dicarboxylic acids,  
 177–178  
 methylene dichloride, 174  
 monochlorodimethyl ether, 172–174  
 paraformaldehyde, 178  
 polyfunctional compounds, 178  
*p*-xylylene dichloride, 171–172, 175  
 sulfonyl chloride, 178  
 thionyl chloride, 178  
 tris-(chloromethyl)-mesitylene, 171–172
- $\text{Cr}^{3+}$ , radius for, 460t, 462t
- $\text{Cs}^+$ , radius for, 462t
- $\text{Cu}^{2+}$ , 441, 444  
 radius for, 460t, 462t
- CytoSorb<sup>TM</sup>, 580–581
- ## D
- Deminerization, gel-type ion exchange  
 resins for, 57
- Dendrimers, 297, 298f
- Deformation of networks, 231–238, 233t,  
 234f–238f  
 photoelasticity phenomenon in, 231–233  
 visualization of inner stresses in network,  
 233–238, 233t, 234f–238f
- Density,  
 styrene-divinylbenzene networks, 64, 64f  
 hypercrosslinked polystyrene networks,  
 238–240, 239f, 239t
- Desorption,  
 equation for process of, 399  
 hydrocarbons, 398–401, 399f  
 isotherm for methanol on Styrosorb 1, 385f  
 isotherm for  $n\text{-C}_8\text{F}_{18}$  on Styrosorb 1BP,  
 387–388, 389f, 390f  
 isotherm for  $n\text{-C}_8\text{F}_{18}$  on Styrosorb 2, 388f  
 kinetics of, 399, 399f  
 Styrosorb 2 in cycles of sorption-, 400
- Deswelling, porous network polymers,  
 291–295, 292f, 294f, 295, 295f
- Diaion HP-20, lipophilic marine toxins  
 accumulated by, 552
- Dialysis-related amyloidosis (DRA), 569
- Diblock copolymers, formation of  
 nanospheres/micelles from, 313
- Differential thermal analysis, 289, 289f
- Diisopropenylbenzene (DIPB), 11
- Dimethoxy methane, 182t, 193, 193t, 214f,  
 215f, 241t, 333–334
- Dow Chemical Company, 355–356
- Dow Optipore<sup>TM</sup>, 355
- Dowex MSA-1, 433, 434t
- Duolite A-7, 107t
- Duolite A-561, 107t

Duolite C-464, 107t  
 Duolite ES-771, 107t  
 Duolite ES-863, 107t  
 Duolite S-37, 107t  
 Duolite S-861, 107t  
 Dyes synthetic, 411–415, 412t, 414f  
   Azo dye Acid Red 14, 415  
   Cresyl Violet, 415  
   Direct Bordeaux, 413–415, 414f  
   Malachite Green, 413–415, 414f  
   Methyl Orange, 415  
   Methylene Blue, 413–415, 414f  
   Neutral Red, 415  
   Rhodamine B, 415  
   Rhodamine C, 415  
 Dynamic ion hydration numbers, 458, 459t

## E

Elastomer, 325  
 Electron microscopy, macroporous styrene-divinylbenzene copolymers texture, 98  
   polymer texture investigated with, 259–266, 260f–262f, 266f, 266t  
   pore size distribution determined by, 83  
 End-stage renal disease (ESRD), 568, 578, 580  
 Envi-Chrom P, 530, 532, 533t  
 EPR spectra, 290, 290f

## F

$\text{Fe}^{2+}$ , 441  
   radius for, 460t, 462t  
 $\text{Fe}^{3+}$ , 441, 444  
   radius for, 460t, 462t  
 Fermentation liquid clarification, 418–421, 419f, 421t  
 Fick's second law, 390  
 Flory-Dušek-Prins approach, 36, 42  
 Flory-Huggins parameter, 84, 85, 87, 272  
 Flory-Rener equation, 29, 67  
 Fluoranthene, 437  
 Fluorine, 437  
 Fourier transform infrared (FTIR)  
   spectroscopy, 184–189, 185f–187f  
 Friedel-Crafts catalyst, 148, 149, 192–193, 193t  
 FTIR spectra of polymers, 184–189, 185f–187f  
 Furfural, 436–437

## G

Gas/organic vapor sorption, 371–409  
   isotherms for nitrogen, 374, 375f, 376f, 379, 380  
   free energy of organic solvent vapors sorption, 212t  
   capillary condensation, 77  
   carbon dioxide, sorption parameters 378t  
   carbon SKT-6A, cyclohexanone adsorbed on, 402  
   column protection time, 396, 397f  
   diffusion coefficients, 391–392, 392f  
   hydrocarbon sorption/desorption, 398–401, 399f  
   hysteresis loop in, 77–81  
   heat effect, 394–395  
   organic compounds retention volumes, 403–404, 403t  
   pore size determined by, 76–80  
   pore volume determined by, 74–75  
 Gasoline emulsion, 423, 423t  
 Gigaporous polymeric separating media, 117–145  
 Glass transition temperature, 55–56, 56t  
 Glauicine, 542t

## H

Hammet-Taft constant, 517  
 Halogenated hydrocarbons, 406r  
 Hemodialysis, hemoperfusion v., 567–570  
 Hemoperfusion, 567–570  
 Hemosorbents, 567–583  
   b<sub>2</sub>M-microglobulin removal with, 570–574, 573f  
 Henry constant, 403  
 Heterogeneous membranes, 601–603  
 Hg<sup>2+</sup>, 441, 443, 444f  
 High performance liquid chromatography,  
   column packing material in, 503–521  
   ion-exchanging ability of, 510–512, 512f  
   macroporous polystyrene v. silica-based, 503–507, 506f, 507f  
   metal-complexing ability of, 510–512, 512f  
   polystyrene as restricted-access adsorption material, 507–510, 509f

- $\pi$ - $\pi$ -interaction selectivity in, 513-524, 514f, 516f, 517f, 519f
  - quasi-normal phase, 515-518, 517f
  - high-temperature, 520
  - mixed mode chromatography, 518, 519f
  - other modes of HPLC separations, 518-521
  - reversed phase, 504, 513-515, 514f, 516f
  - HR-P., 357t, 556, 557t, 558t
  - n-Hydrocarbons, 382f, 383t, 385f, 387f, 392f, 403t
    - desorption of, 398-401, 399f
    - kinetics of sorption of, 389-393, 391f, 392f
  - Hydrogen chloride gas, purification of, 617
  - Hydrogen storage, 605-610, 606t, 609t
  - Hydroquinone, hypercrosslinking of, 620
  - Hypercrosslinked polyamide, 336
  - Hypercrosslinked polyaniline, 333-336
  - Hypercrosslinked polyarylates, 330-331
  - Hypercrosslinked polydivinylbenzene, 328
  - Hypercrosslinked polyimides, 336
  - Hypercrosslinked polypyrrole, 333-336
  - Hypercrosslinked polysilsesquioxane, 346-347
  - Hypercrosslinked polysulfone, 329-33
  - Hypercrosslinked polyxylylene, 331-333, 332t
  - Hypercrosslinked pyridine-containing polymers, 337-338, 338t
  - Hypercrosslinked spirobisindane-based, 341
  - Hypermer 1070, 125
  - Hayesep B, 606t
  - Hayesep N, 606t
  - Hayesep S 606t
  - HYSphere-1, 357t
  - Hysteresis loop, 77-81
- I**
- Ideal network, 29
  - Ideal separation process, 482-486
  - Inner stresses of hypercrosslinked network, 202-213, 202f, 203t, 204t, 206f-208f, 210t, 212t
  - Integral heat of solvation, 210t
  - International Sorbent Technology, 357t
  - Interpenetrating polystyrene networks (IPN), 63-70
  - Intramolecular crosslinking, 297-303
  - Intramolecular cyclization, 12-19
  - Ion,
    - dynamic hydration numbers for, 458, 459t
    - electrophoretic mobility, 459, 460t
    - hydroxyl ions and protons, 461-463, 462t
    - radius of, 460-461, 460t, 462
    - Stokes-Einstein relationship, 458
    - structure making/structure breaking effects, 457
    - van der Waals radius, 460, 460t
  - Ion exchange resins, anion-exchange resins
    - Styrosorbs, 595-597, 596t
    - gel-type resins, 56-62, 59t, 60t, 61t
    - hypercrosslinked resins, 585-600
    - hypercrosslinked sulfonates, 586t, 587t, 590, 595, 596t
    - isoporous anion-exchange resins, 147
  - Isolute ENV, 357t, 532, 533t, 535, 546
  - Isotope isomers of water, 407-409, 408f
- J**
- Jiangsu N&G Environmental Technology Co. Ltd., 355
- K**
- K<sup>+</sup>, radius for, 462t
  - Kelvin equation, 76-77
  - Kinetics of swelling in toluene, 224f
- L**
- La<sup>3+</sup>, radius for, 460t, 462t
  - Lecithin, 422, 422t
  - Lewatit EP63, 606t
  - Lewatit S-7768, 355
  - Lewatit VP OC 1163, 355
  - LiChrolut EN, 357t, 534, 538-540, 545-546, 553-556, 555t-558t, 559
  - LiCl separation of mixture LiCl/HCl, 477-479, 478f
    - separation of mixture LiCl/LiOH, 477-479, 479f
  - Limits of detection (LOD), 528-529
  - Limits of quantification (LOQ), 528-529
  - Limonin, 434-435, 435t
  - Lipids, sorption of, 422, 422t
    - lecithin, 422, 422t
    - oleic acid, 422, 422t
  - Liquid-liquid extraction (LLE), 524

**M**

- Macherey Nagel, 357t
- Macro-phase separation, 320
- Macroporous copolymers, 71-116
  - cauliflower texture of, 98, 99f, 101
  - pores classified in, 99
  - porosity formation via nucleation 99-101, 100f
- Macrosyneresis, 84, 94, 98, 320
- Melanoidins, 419-420, 419f
- Mg<sup>2+</sup>, radius for, 460t, 462t
- Memory term, 33, 41
- Mercury intrusion porosimetry, 75-76, 81-82, 134, 250-251
- Mesitylene tricarboxic acid, 177-178
- Mesopores, 72
- Metal-organic frameworks (MOF), 347-354, 621
  - interpenetrating MOF, 349-350
- Methane, storage on hypercrosslinked polystyrene, 605-610
- Methanol,
  - sorption isotherm for, 382f, 594, 594f
  - volume of absorbed, 383t
- Methylstyrene, 30
- Mg<sup>2+</sup>, radius for, 460t, 462t
- Micelles, formation from diblock copolymers of, 313
- β<sub>2</sub>-Microglobulin removal, 570-574, 573f
- Microbeads of hypercrosslinked polystyrene, 503-513, 618-619
- Micrographs in polarized light, 234-236, 234f-236f
- Micro-phase separation, 316
- Microsyneresis, 84, 96, 98, 101, 316
- Mn<sup>2+</sup>, radius for, 460t, 462t
- MN-100,
  - cephalosporin C sorption with, 436, 436f
  - citrus juice de-bittering with, 434
  - hydrogen storage capacity of, 606t
  - hexane/benzene retention with, 405
  - phenols sorption by, 426
- MN-102, citrus juice de-bittering with, 434-435, 435t
- MN-150,
  - cephalosporin C sorption with, 436, 436f
  - citrus juice de-bittering with, 434
  - phenols sorption by, 425-426, 425f
  - MN-200,
    - acenaphthene sorption with, 437
    - anthracene sorption with, 437
    - bismuth determination in lead-lithium alloys with, 512
    - cephalosporin C sorption with, 436, 436f
    - chloroform removal with, 428
    - clarification of riboxin solutions with, 420, 421t
    - fluoranthene sorption with, 437
    - fluorine sorption with, 437
    - halogenated hydrocarbons retention volumes on, 406t
    - hydrogen storage capacity of, 606t
    - mercury acetate and mercury nitrate sorption with, 441, 442f
    - naphthalene sorption with, 437
    - n*-valeric acid sorption with, 418t
    - pesticides sorption with, 429-431, 43
    - phenols sorption with, 425-426, 425f, 426f
    - pyrene sorption with, 437
    - tryptophan cultural liquid, 420
    - Zn<sup>2+</sup>/Cd<sup>2+</sup>/Pb<sup>2+</sup> separation with, 512f
  - MN-202, 450, 488
    - electrolyte separation with, 467f
  - MN-250, 402f
  - MN-270, 450
    - electrolyte separation with, 467f
  - MN-400, cephalosporin C sorption with, 436, 436f
  - MN-500, 488
    - electrolyte separation, 467f
    - hydrogen storage capacity of, 606
- Model networks, desinterspersion effect, 51f
  - distance between junction points, 44-45, 45t
  - gyration radii of polystyrene chains, 48, 48f
  - network equilibrium with cyclohexane, 38
  - presumptive mechanism of swelling, 50-52, 50f, 51f
  - swelling theory verification with, 32-43, 34f-39f, 42f, 43f
  - synthesis of model "ideal networks," 30-32
  - characterization by SANS technique, 43-50, 45t, 46t, 48f, 49t

Monochlorodimethyl ether (MCDE),  
172–174  
Monoliths, 128–146, 621

## N

Na<sup>+</sup>, radius for, 460t, 462t  
NanoNet 357, 445, 450  
Naphthalene sulfonates, 547, 547t  
NDA-100, phenols sorption, 427–428  
NDA-150, 1-naphthylamine sorption,  
438–439  
NG-99, 355  
NG-100, 355  
Nanocomposite catalysts of organic  
reactions, 603–605, 618  
NanoNet, 381, 357, 445, 450, 781  
Nanosponge, 303–313, 313f  
Naphthalene, 30, 437  
Naphthalene sulfonates, 547, 547t  
1-Naphthylamine, 438–439  
Network topology, 26–27  
Ni<sup>2+</sup>, 441, 444  
NMR of water protons in pores, 256, 256f  
NN-381, 357, 450, 452, 488  
NN-781, 357  
Nucleosides, 420

## O

Oasis HL3, 534, 552–560, 557t, 555t–558t,  
Oestradiol removal, 618  
Oleic acid, 422, 422t  
Optipore L-493, 552  
Ortho-para-spin isomers of water,  
407–409, 408f

## P

Pb<sup>2+</sup>, 443, 444f  
Pd-1,10-phenanthroline cluster, 313  
Pendent vinyl group (PVG), 12–15  
Pentamidine, 143  
Perfluorocarbons, 387  
Pervaporation, 524–525  
Pesticides, 429–432, 430f, 535–539, 536t  
antifouling pesticides, 538  
Atrazine, 429–431, 430f, 536  
Benazolin, MN-200 sorbing, 431  
Bendazol, 542t

Bentazone, MN-200 sorbing, 431  
Chlorotoluron, MN-200 sorbing,  
429–430, 430f  
Deisopropylatrazine (DIA), 536, 536t  
Desethylatrazine, 539, 536t  
Desethyl-terbutylazine (DTBA), 536t  
Desethyltriazine (DEA), 536t  
Didealkylatrazine (DDA), 536t  
Diprazine, 438  
Diuron, 429–430, 430f  
Fenamiphos, 539  
Fenitrothion, 539  
Fonofos, 539  
Hydroxyatrazine, 536t  
Imazapyr, 431  
Isoproturon, 429–431, 430f  
Propazine, 536t  
Simazine, 429–430, 430f, 536f  
triazine pesticides, 431  
Terbutylazine, 536t  
Triclopyr, 431  
Pharmaceuticals  
bendazol, 542t  
carbamazepine, 555t  
clofibric acid, 555t  
diclofenac, 556t  
glaucine, 542t  
ibuprofen, 555t  
ketoprofen, 555t  
metoprolol, 543t  
nadolol, 543t  
naproxen, 555t  
oxytetracycline, 553–554  
pentamidine, 143  
prednisalon, 542t  
propranolol, 541t  
sanorine, 541t  
seduxen, 542t  
verapamil, 541t  
Phloroglucinol, 530  
Phosphatidylcholine, 438  
Photoelasticity phenomenon, 231–233  
PLRP-S, 504, 506f, 530, 532, 533t  
Polybrominated diphenyl ethers (Flame  
retardant agents), 543  
PolyHIPE, 125t, 125–126,  
Polysorb-1, 404t, 422, 422t

- Popcorns, 21–22, 56  
 Porapak T, 404t  
 Porolas SG-2T, 404t  
 Porolas V-2T, specific interactions/total interaction energy with, 404t  
 POROS, 126, 127f  
 Porous graphitic carbon (PGC), 530, 533t  
 Positronium annihilation lifetime spectroscopy (PAL), 255  
 $\pi$ - $\pi$ -interaction selectivity in HPLC, 513–524  
 Protons, radii of, 461–463, 462t  
 Purosep 200 for SPE, 525f, 534  
 Pyrene, 437  
 Pyridine-containing hypercrosslinked polymers, 338
- Q**
- Quasi-normal phase chromatography, 515–518, 517f
- R**
- $Ra^{2+}$ , radius, 460t, 462t  
 $Rb^+$ , radius for, 462t  
 Reference state, 41  
 Rhenium extraction with, 601  
 Riboxin, 420, 421t  
 Rod-like macromolecules, 297, 298f
- S**
- Salt retardation, 475–480, 476f–479f  
    $MgCl_2/K_2SO_4$  separation  
    $(NH_4)_2SO_4/NH_4Cl$  separation  
 SANS, small-angle neutron scattering, 38, 44–48, 54–55, 45t–46t, 47f–48f, 49t  
 SAXS, small-angle X-ray scattering, 266–268, 267f, 268f  
 Schroeder's paradox, 439  
 Self-assembling to regular clusters, 306–314, 307f, 308f, 311f, 313f  
 Sepabeads SP850, 552  
 Sephadex G-10, 448  
 Shallow shell resins, 58  
 Shilov' constant, 396, 397  
 Simulated moving bed (SMB), 492–493, 493f  
 Solid-phase microextraction (SPME), 525, 524  
 Spark Holland, 357t  
 Stilbene sulfonates, 547, 547t
- Stir bar sorptive extraction (SBSE), 524  
 Styrosorb 1, 373  
 Styrosorb 1BP, 373  
 Styrosorb 2, 373–374, 423, 423t  
 Sulfachloropyridazine, 553–554  
 Svedberg's equation, 312
- T**
- Telogens, gel-type ion exchange resins with, 61  
 Tenax GC, 404t  
 Tetrabutylammonium iodide resolution, 499, 499f  
 Thermal stability of copolymers, 287–291, 288f–290f  
 Thermoporosimetry, pore size distribution determined by, 83  
 $Tl^+$ , radius for, 462t  
 $Tm^{3+}$ , radius for, 460t, 462t  
 Topology of networks, 26–27  
 Tributyl phosphate (TBP), 415–416, 416f  
 Tris-(chloromethyl) mesitylene (CMM), 171–172, 182t, 203t, 205, 206f, 214f, 217, 241t
- V**
- Van der Waals radius, 460, 460t  
 Veterinary antibiotics recovered with SPE 553–554  
 Volatile organic compounds in air, 564t
- W**
- Water isotope isomers, 407–409, 408f  
 Water vapor, sorption of, 379, 400  
   sorption isotherm for, 382f  
 Wofatit Y77, 606t
- X**
- $p$ -Xylylene dichloride (XDC), 171–172, 175, 318
- Y**
- Young's modulus, 67
- Z**
- Zimm's diagram, 306, 307f  
 $Zn^{2+}$ , 441, 444  
   radius for, 460t, 462t  
    $Zn^{2+}/Cd^{2+}/Pb^{2+}$  separation, 511, 512f