

# Index

Page numbers followed by *f* indicate figures.

## A

- Abiotical stresses induce Pd callose, 447–448
- Abiotic stresses, callose in, 446–447
- Acer pseudoplatanus*, 449
- Acetobacter xylinum*, 331
- Achlya*
  - cytoplasmic (1,3)- $\beta$ -glucan of, 366–367
- Achlya bisexualis*, 241, 367
- Achnanthes brevipes*, 362
- Achyla bisexualis*, 15
- Acidic fibroblast growth factor (aFGF), 597
- Acidic terpenoids, 271
- Acremonium*, 73
- Acremonium persicinum*, 139
- Actin filaments, 428
- Actinidia deliciosa*, 441
- AFM. *See* Atomic force microscopy (AFM)
- Agaricus bisporus*, 394
- Agrobacterium*, 328, 329
  - 10C3K strain of, 328
  - growth of, 331*f*
  - LTU50 strain of, 333, 333*f*
  - scanning electron microscopy on, 331
- Agrobacterium curdlan synthase*, 288
- Agrobacterium* LTU50, 202

- Agrobacterium tumefaciens*, 202, 329
- Alaria angusta*, 370
- Alkaline peeling, 11
- Allium cepa*, 441, 503
- Alternaria brassicicola*, 535
- Aluminium
  - induced callose formation, 505–507
  - induced inhibition of root elongation, 506
  - reducing K<sup>+</sup> efflux, 500
  - toxicity, 504
  - treated plants, 447
- Alveolation
  - in syncytial systems, 429–432
- Alveoli, 429, 430*f*
- 1-Aminocyclopropane-1-carboxylic acid, 507
- Amphibolus griffithi*, 474
- Amphiobolus*, 476
- Amphotericin B, 270, 271, 300, 305
- AMP-producing genes, 564
- AMP synthesis, 563
  - induction of, 564–565
- Anarthriaceae, 633
- Angiosperms, 465
- Aniline Blue, 11, 265, 267, 295, 309, 397
- fluorochrome, 283
- fluorophore in, 61

- Anoxia, 510–511
- Anticlinal walls (AW), 429, 430*f*, 431
- callose and, 431–432
- Antimicrobial peptides (AMPs), 563
- Aphanomyces astaci*, 240–241, 243, 246
- Apodachlya*, 408
- Apomictic embryo sacs, callose in, 487
  - events of embryo sac formation in, 488
- Arabidopsis*, 286
  - pollen grains, 471–472
- Arabidopsis thaliana*, 441, 501, 526
- Archaeabacteria, 655
- Aromatic cradle, 179
- Arum alpinum*, 476
- Ascomycete side-chain-branched (1,3;1,6)- $\beta$ -glucans, 16
- Ash
  - seasonal variation of, in *Laminaria hyperborea*, 372*f*
- Aspergillus fumigatus*, 261–262, 388, 392, 394, 587, 632, 658
  - cell wall glucan, 18
  - wall composition of, 393*f*
- Aspergillus nidulans*, 261–262, 394
- Aspergillus niger*, 141
- Astasia ocellata*, 355

- AtBG\_pap* gene, 453  
*AtBG* T-DNA insertion mutants, 540  
*AtCalS1* gene, 501  
*AtGSL6*, 302  
*AtGSL8* and *AtGSL10* proteins, 481  
*Atgsl10* mutants, 482  
 Atomic force microscopy (AFM), 60  
*Avena sativa*, 91  
 AW. *See* Anticlinal walls (AW)  
*Azorhizobium caulinodans*, 19  
*Azospirillum brasilense*, 19, 336  
 Azuki bean, 303
- B**
- Bacillariophyceae*, 236. *See also* Diatoms  
*Bacillus*, 329  
*Bacillus amyloliquefaciens*, 12  
*Bacillus circulans*, 188  
*Bacillus halodurans*, 182  
*Bacillus subtilis*, 136  
 Bacitracin, 224  
 Barley endotransglycosylase/hydrolase, 661  
 Barley grain, 64–1  
 Basic fibroblast growth factor (bFGF), 596  
 Basidiomycete side-chain-branched (1,3;1,6)- $\beta$ -glucans, 16  
*Bathycoccus sp.* BAN7, 311  
*Bcl10-Malt1*-mediated NF- $\kappa$ B activation, 585  
(1,3)- $\beta$ -D-glucan endohydrolases  
 GH17 family, 122–128  
 GH26 family, 132–133  
(1,6)- $\beta$ -D-glucan endohydrolases, 139  
 $\beta$ -D-glucan glucohydrolases  
 in auxin-mediated cell, 145  
 in pollen development, 146  
(1,3)- $\beta$ -D-glucan phosphorylases, 153–154
- (1,3;1,4)- $\beta$ -D-glucans, 140, 141, 622  
 in commercial processes, 622  
 fine structure of, 624–625  
 cellobiosyl and cellobetaosyl residues, 624  
 molar ratios of, cellobiosyl to cellobetaosyl units, 625  
 from grasses, 623  
 $\beta$ -D-glucosidases of family GH1  
 action pattern, 142–143  
 catalytic mechanism, 143  
 classification of, 139–140  
 3D structures, 143–145  
 properties of, 140  
 substrate specificity, 140–142  
 $\beta$ -D-glycoside exo-hydrolases of family GH3  
 action pattern, 147–148  
 catalytic mechanism, 148–150  
 classification of, 145  
 3D structures, 150–151  
 properties of, 145–146  
 substrate specificity, 146–147  
 structural determinants of, 151–153  
*Betula pubescens*, 441  
(1,3)- $\beta$ -glucanase  
 to break down fungal cell walls, 543  
 degradation of transient callose wall, 471  
 in endoplasmic reticulum membrane system, 542  
 expressed by tapetum, 470  
 gene into TMV genome, 541  
 in regulation of cell-to-cell transport, 502  
 roles in plasmodesmatal trafficking and, 540  
 transgenes expression in tapetum, 475  
 $\beta$ -glucan biosynthesis in protozoans and chromatins, molecular biology of, 248–249
- (1,3)- $\beta$ -glucan endo-hydrolases  
 activity in early spring, 449  
 catalysing callose degradation, 443  
 genes encoding, 452  
 importance of, 452  
 limited/absent expression of gene, 452  
 localization, 444, 451  
 plasmodesmal associated, 453  
*Ulva lactuca*, polysaccharide hydrolysed by, 630  
 vesicles transportation to, 453  
 virus recruiting and/or activating, 452
- (1,3)- $\beta$ -glucan in embryophytes, synthesis of, 285  
 biochemical identification of GSL proteins, 296–298  
 callose synthase complexes, 298–303  
 functional analysis of GSL genes, 294–296  
 GSL gene family, 288, 292–294  
 GSL proteins, 286–288, 287f  
 $\beta$ -glucan recognition proteins (BGRP), 189
- (1,3)- $\beta$ -glucans, 563  
 anti-inflammatory activity, 597–598  
 anti-tumor activity of, 594–595  
 on arthropod immune system, 572  
 bacterial infections, 591–592  
 biosensor for, 568  
 chemical characteristics of, 5–10  
 chemical structure variants of, 62–67  
 conformations of, 48–52, 65–67  
 derivatives of, 24–27  
 detection of, 11  
 dietary supplementation with, 599  
 dissolution of, 11–12  
 distribution, 655–656

- in embryophytes, 284–285  
 evolutionary significance of, 660–661  
 fluorescence spectroscopy, 61–62  
 forms and functions of, 657  
 fungal infections, 592–593  
 in fungi. *See Fungi*  
 gelation behaviour of, 93–103  
 induced AMP synthesis, in *Drosophila*, 565f  
 induced coagulation, in horseshoe crabs, 567f  
 induced proPO activation, in crustaceans, 569f  
 ischemia/reperfusion injury, 595–596  
 isolation of, from cell walls, 389–391  
 labelling of, using Dectin1-Fc chimeric protein, 398f  
 in lichens, 406–407  
 linear, 399, 407  
 localization of, 399f  
 location of, in fungi, 409f  
 mammalian receptors for, 580  
   complement receptor 3, 581–582  
   dectin-1, 582–587  
   lactosylceramide, 580–581  
   scavenger receptors, 581  
 microscopy, 60–61  
 occurrence of, 658  
 in oomycetes  
   oomycete cell wall, 407–408  
   storage (1,3)- $\beta$ -glucans, 408–409  
 SAXS, 61  
 sedimentation analysis, 61  
 solid state  $^{13}\text{C}$  NMR of, 58–60  
 solid state conformations in, 52–67  
 to stimulate innate immunity and, 579  
 and stimulation, of innate immunity, 598–600  
 structural variations in, 396–405  
 structure/activity relationships with, 587–588. *See also Dectin-1*  
 structure and molecular organization of, in yeast and module cell walls, 391–396  
 structures of, 12  
 synthases, 241–242  
 triggering release of immune components from, 571  
 and viral infections, 593–594  
*in vivo*  
   effect, *administration on*, 590  
   effects, in animal models of disease, 590–591  
 pharmacokinetics,  
   pharmacodynamics and bioavailability, 588–589  
 and wound repair, 596–597  
 x-ray diffraction of, 54–58  
 yeast cell wall structure, 260–261, 260f  
(1,3;1,2)- $\beta$ -glucans  
   side-chain-branched, 20  
(1,3;1,4)- $\beta$ -glucans, 178, 621, 622, 629. *See also* (1,3;1,4)- $\beta$ -D-glucans  
   in aleurone-rich fraction, 643  
   among different taxa, 659  
   biological role of, 625  
   concentrations of, 629  
   effects of fine structure on, 627–628  
   formation of gel-like matrices in, 625–627  
   as source of metabolizable energy, 629  
   in cell walls of mature wheat grains, 642  
   cereal, 21–22  
   from *Cetraria islandica*, 628  
   chlorophyte, 23–24  
   chromalveolate, 23–24  
 chromistan, 23–24  
 concentrations and locations, in cereal grains, 637–639  
 in barley, 638–639  
 cereals other than oats and barley, 639  
 developing oat genotypes with, 639  
 quantification of, 638  
 reducing concentrations of, 638  
 in whole grains, 637–638  
 in cultivars of oats, 642  
 distribution of, 629, 631  
   in angiosperms, 632–634  
   in cell walls, of mature grains, 640–643  
 chromalveolates, 632  
 lichens and fungi, 630–632  
 plants other than angiosperms, 630  
 prokaryotes, 632  
 effects of environment on, 639–640  
 environmental factor implicated in, 640  
*Equisetum arvense*, 22  
 evolutionary significance of, 660–661  
 fungal cell wall, 23  
 genes, in biosynthesis of, 660  
 in grasses, 21–22, 622  
 interactions, 627  
 lichen, 23  
 liverwort, 23  
 location vegetative organs, of Poaceae, 634–636  
   transmission electron micrographs, 635  
 occurrence of, 658–659  
 within plant variation in, 640  
 rhodophyte sulfated, 24  
 role in cell-wall architecture, of Poaceae, 637  
 stock feed formulations for, 623

- (1,3;1,4)- $\beta$ -glucans (*Continued*)  
 in walls of *E. arvense*, 630  
 in walls of starchy endosperm, 629  
 in wheat grain, 628
- (1,3;1,6)- $\beta$ -glucans  
 branch-on-branch, 16–19, 392  
 chemical structure of, 357, 358f  
 cyclic, 19–20  
 side-chain-branched, 14–16
- (1,6)- $\beta$ -glucans  
 localization of, 399f
- $\beta$ -glucans biosynthesis,  
 hypothetical model for, 234f
- $\beta$ -glucans biosynthesis in  
 Protozoans and  
 Chromistans, biochemistry of  
 assay conditions and kinetic parameters, 239–240
- biosynthesis of side-chain-branched (1,3;1,6)- $\beta$ -glucans, 248
- model organisms, 235–236
- modulation of (1,3)- $\beta$ -glucan synthase activities in vitro, 240–244
- product characterization, 246–248
- purification of (1,3)- $\beta$ -glucan synthases, 244–246
- subcellular localization of  $\beta$ -glucan synthases and preparation of cell-free extracts, 236–239
- (1,3)- $\beta$ -glucan synthases  
 movement on cell wall surface, 267f  
 fungal *FKS* genes, role in, 442–443  
 genes encoding, 452, 657–658  
 and Pd conductivity, 445
- (1,3;1,4)- $\beta$ -glucan synthases, 659–660
- (1,3)- $\beta$ -glucosidic linkages, 176
- (1,4)- $\beta$ -glycosidic linkages, 393
- BGRP. *See*  $\beta$ -glucan recognition proteins (BGRP)
- BhCBM6*, 182
- (1,3)- $\beta$ -homopolymers, 175, 176
- Bioassays, 341
- Biofilm matrix, 388
- Biosynthesis of linear  $\beta$ -glucans, hypothetical model for, 234
- Biosynthetic enzymes for (1-3)- $\beta$ -glucans  
 catalytic subunit, 261–263  
 cell wall integrity checkpoint in yeast, 269–270  
 cross talk in the regulation of other cell wall components, 270–271  
 glucan synthase inhibitors, 271–272  
 glucan synthesis in spore formation, 267–269  
 other proteins associated with glucan synthase activity, 264
- regulation of (1,3)- $\beta$ -glucan synthesis in yeast  
 factors involved in the regulation of glucan synthase activity via Rho1p, 264–265  
 location of (1,3)- $\beta$ -glucan synthase, 266  
 movement of Fks1p in the plasma membrane, 266–267
- post-translational modification of Rho1p, 265
- transcriptional regulation of the FKS genes, 265–266
- regulatory subunit, 263–264
- Biosynthetic enzymes for (1-6)- $\beta$ -glucans, 272–273
- Biotic stresses, callose in, 525–531  
 herbivorous attack, 451–452  
 viral infection, 452–453
- hydrolysis of callose and gating of Pd, 454
- Blaberus craniifer*, 570
- Blaberus discoidalis*, 571
- Blastomyces dermatitidis*, 400
- (1,3)- $\beta$ -linked glucosidic linkages, 243
- Blumeria graminis*, 293, 532
- Boltzmann constant, 70
- Botrychium virginianum*, 448
- Botryosphaeria rhodina*, 16
- Botrytis cinerea*, 388, 536
- Bradyrhizobium japonicum*, 5, 19, 336  
 nodulation and, 338–339  
 strains, 19
- Brevicoryne brassicae*, 143
- Bromeliaceae, 634
- Brown algae. *See* Phaeophyceae
- Bryophyllum daigremontiana*, 202
- Bryophytes, 23
- BspCBM28*, 185
- C**
- $\text{Ca}^{2+}$  influx, 543
- Calcofluor white, 271
- Callose, 11, 14, 283, 284, 364  
 accumulation, 294f  
 AW and, 431–432  
 deposition, 305–306, 432–434  
 after microbial attack, 528  
 function of, in cytokinesis, 434–435  
 periclinal wall and, 431–432  
 plant cytokinesis and, 425  
 in pollen tubes, 473  
 and role in, flowering plant, 465
- Callose during development, 448  
 cotton fibre elongation, 451  
 definitive and dormancy callose, 449–451  
 sieve plate pore development, 448
- Callose formation  
 and anoxia, 510–511

- and gravitropic response, 513  
kinetics A1-induced and  
digitonin-induced, 503  
metal toxicity and, 502–508  
aluminium, 504–507  
manganese (Mn), 507–508  
in response to abiotic stress,  
499–502  
and temperature stress, 511–513  
TTSS suppressing, 538  
wounding, osmotic stress and,  
508–510
- Callose localization  
in Pd and sieve plates, 441–442  
localization of callose, 442  
raised collar structures, 442
- Callose SCW, during microgamete development, 476
- Callose synthase (CalS), 300, 432  
cellulose and, 512  
components of, 530  
gene *GSL5*, 538  
genes coding for, 514  
inhibition of, 536  
in plants, 442  
preparations, 299  
molecular weight of major polypeptides present in,  
297–298  
purify and characterization, 529  
regulatory mechanisms, 300f,  
543  
role in regulating of, 531
- Callose synthesis, regulation of,  
303–304, 306–309  
synthesis of (1,3)- $\beta$ -glucans in chlorophytes, 309–311  
synthesis of (1,3)- $\beta$ -glucans in rhodophytes, 311–312
- Calmodulin, 302
- Calnexin, 301
- CalS. *See* Callose synthase (CalS)
- Candida albicans*, 153, 261, 262,  
271, 272, 388, 400, 599  
cell wall glucan, 18
- Candida* enzyme, 153  
*Candida glabrata*, 396  
Capsular polysaccharide (CPS),  
342
- Carbohydrate-Active enZymes (CAZy) database, 121,  
139
- Carbohydrate-binding modules (CBM), 172–174  
biochemical analysis of  
CBM13, 188–189  
CBM39, 189–191  
classification, 173–174  
ligand binding profiles for,  
172  
structural comparison of,  
191–192  
structure-function of  
CBM4, 176–179  
CBM6, 179–182  
CBM11, 182–183  
CBM17, 183–185  
CBM22, 186  
CBM28, 185  
CBM43, 186–187  
three-dimensional structures for,  
177
- Carboxymethylation, 72
- $\kappa$ -carrageenan, 134
- Caspofungin, 270, 271
- Catalytic acid, 123
- Catalytic base, 123
- CAZy. *See* Carbohydrate-Active enZymes (CAZy) database
- CBM4, 176–179, 191  
aromatic cradle, 179  
*Cf*CBM4-1, 179  
*Tm*CBM4-2, 177–178, 179
- CBM6, 179–182, 191
- CBM11, 182–183
- CBM13, 188–189
- CBM17, 183–185
- CBM22, 186
- CBM28, 185
- CBM43, 186–187
- Cellotriosyl, 63
- Cell plate, formation of, 428–429
- Cell plate assembly matrix (CPAM), 428
- Cellular immunity, 571–572
- Cellular metabolism, global and curdlan, 335
- Cellulins, 408
- Cellulomonas*, 329, 335
- Cellulomonas falvigena*, 13
- Cellulomonas fimi*, 149, 177
- Cellulomonas flavigena*  
floc formation and, 334  
KU, 332
- Cellulose colocalization, with callose, 472
- Cellulose synthase-like (CSL) proteins, 287–288
- Cellulosimicrobium cellulans*, 188
- Cellvibrio mixtus* endoglucanase 5 A (*Cm*CBM6-2), 180
- Cell wall glucan  
*Aspergillus fumigatus*, 18  
*Candida albicans*, 18  
*Pythium aphanidermatum*, 18–19  
*Saccharomyces cerevisiae*, 16–18
- Cell wall polysaccharides, 367–368
- Cell walls, fungal  
(1,3)- $\beta$ -glucans, 388  
isolation of (1,3)- $\beta$ -glucans from cell walls, 389–391  
structural variations, 396–405  
structure and molecular organization of, in yeast and module cell walls, 391–396  
chemical composition of yeasts and fungi, 389t  
fractionation. *See* Fractionation, of cell walls  
protocol for extraction of components, 390, 391f
- Cell walls and  $\beta$ -glucans in yeast, 260–261, 260f
- Centrolepidaceae, 633

- Cereal β-glucans  
behaviour of concentrated solutions, 91–93  
gelation of, 100–103  
physical properties of, 22  
solution properties of, 81–82
- Cerk1* mutants, 535
- Cetraria islandica*, 407, 630, 658
- Cf* CBM4-1, 179
- Chaetoceros brevis*, 362
- Chaetoceros mulleri*, 16
- Chain association, 49–52
- Chain shape, 49
- Chain stiffness, 48
- CHAPS. *See* 3-[*(3-cholamidopropyl)-dimethylammonio]-1-propane sulfonate (CHAPS)*
- CHAPSO. *See* 3-[*(3-cholamidopropyl)-dimethylammonio]-2-hydroxy-1-propane sulfonate (CHAPSO)*
- Charophytes, 23
- Chitin, 11, 261
- Chitinases, 535
- Chitin hydrolases, 527
- Chitin oligosaccharide elicitor binding protein, 534
- Chlamydomonas reinhardtii*, 309, 310
- Chlorarachniophyceae. *See* Chlorarachniophytes
- Chlorarachniophyta. *See* Chlorarachniophytes
- Chlorarachniophytes, 371–372
- (3-[*(3-cholamidopropyl)-dimethylammonio]-2-hydroxy-1-propane sulfonate (CHAPSO)*), 238
- 3-[*(3-cholamidopropyl)-dimethylammonio]-1-propane sulfonate (CHAPS)*, 238, 245, 296
- detergent extraction, 297
- Chondrus crispus*, 312
- Chorda filum*, 370
- Chromatin immunoprecipitation (ChIP), 265
- Chromistans  
types and structures of polysaccharides in, 354*t*
- Chryschromulina*, 357
- Chrysolaminarin and spores, 362
- Chrysolaminarins, 15, 236, 357, 359, 364  
diel changes in, 361*f*, 362  
formation of resting cells and, 362  
formation of spores and, 362
- Chrysophyceae. *See* Chrysophytes
- Chrysophytes, 364–365
- Clostridium cellulovorans* cellulase (CccBM17), 183–185
- Clostridium stercorarium* xylanase (CsCBM6-3), 180
- Clostridium thermocellum*, 132
- Clostridium thermocellum* xylanase (CtCBM6), 180
- Coat protein, 452
- Coccidioides immitis*, 400
- Coleochaete scutata*, 660
- Collema leptosporum*, 407
- Colletotrichum lagenarium*, 535
- Colletotrichum lindemutianum*, 154
- Complement receptor 3, 581–582
- Conduritol B epoxide, 149
- Congo Red, 244, 271
- Conidia  
comparison of wall composition between, and Hyphae, 399  
antifungal compounds, 401–402  
cell wall mutants, 402  
dimorphic forms, 400–401  
environmental conditions, 401
- Coomassie Blue, 243, 245
- Coprinus cinereus*, 394
- Coscinodiscus concinnus*, 362
- Coscinodiscus nobilis*, 359
- Covalently bound wall proteins (CWP)  
GPI-anchored, 396  
schematic representation of, in yeast, 395*f*
- COX activation, 584
- Cox-Merz rule, 83
- CP. *See* Cross polarization (CP)
- CPAM. *See* Cell plate assembly matrix (CPAM)
- CPS. *See* Capsular polysaccharide (CPS)
- CR3. *See* Complement receptor 3
- Craspedostaurops australis*, 359
- CrdASC gene, 203–205
- CR3-deficient leukocytes, 582
- CrdR gene, 205–206
- CrdS, 288
- C-reactive proteins, 567
- Critical gelation point, 93
- CR3-mediated phagocytosis, 582
- Cross polarization (CP), 53
- Cryogelation  
of water-soluble polymers, 103
- Cryptococcus neoformans*, 388, 400, 589, 593
- CsCBM6-3, 191
- CsF cellulose synthase-like genes, 644
- CtCBM11, 183
- CtCBM22-2, 186
- C-type lectin-like receptors, 584
- Curdlan, 13  
behaviour of concentrated solutions, 84–85  
biosynthesis locus, 202  
chain, 214–216  
crdASC gene, 203–205  
crdR gene, 205–206  
enzymology, 211  
formation and functional roles, 331–336  
gelation of, 93–96  
global cellular metabolism and, 335

- membrane typology, 211–212  
*ntrBC*, 206  
*ntrC*, 207–208  
*ntrYX* gene, 208–209  
 occurrence, 328–330, 330t  
*pSSA* gene, 205  
*relA* gene, 209–211  
 Curdlan sulfate, 593  
 CWP. *See* Covalently bound wall proteins (CWP)  
*Cyanidioschyzon merolae*, 311  
 Cyclic (1,3)- $\beta$ -glucans  
   formation and functional roles  
 hypo-osmotic conditions,  
   adaptation to, 337–338  
 nodulation and N<sub>2</sub>-fixation,  
   338–340  
 suppression of plant defense  
   response, 340–342  
 occurrence, 336–337  
 Cyclic (1,3;1,6)- $\beta$ -glucans, 19–20,  
   216–222  
   bioassays and, 341  
   formation and functional roles  
 hypo-osmotic conditions,  
   adaptation to, 337–338  
 nodulation and N<sub>2</sub>-fixation,  
   338–340  
 suppression of plant defense  
   response, 340–342  
 molecular genetics of, 216–219  
*ndvB* genes, 219–221  
*ndvC* genes, 221–222  
*ndvD* genes, 221–222  
*ndv* genes, 216–219  
 occurrence, 336–337  
 structure of, 19  
*Cyclotella cryptica*, 236, 246, 362  
*Cylindrotheca closterium*, 362–363  
*Cylindrotheca fusiformis*, 359  
 Cyperaceae, 634  
*Cystophora scalaris*, 370  
*Cystoseira barbata*, 371  
*Cystoseira crinita*, 371  
*CYTI* gene, 499  
 Cytokines, 571  
 Cytokinesis, 398, 470  
   function of callose in, 434–435  
   plant, callose and, 425  
 Cytoplasmic (1,3)- $\beta$ -glucan  
   of *Achlya*, 366–367  
**D**  
 Daylily. *See* *Hemerocallis fulva*  
 Dectin-1, 581–587, 593, 595. *See also* (1,3)- $\beta$ -glucans  
   with CD37, 585  
   to enhance TLR-mediated cytokine, 586  
   expressing in leukocytes, 583  
   inducing autoimmune diseases, 586  
   in mammalian systems, 587–588  
   mediating cellular responses, 584  
   mediating IL-6 production, 586  
   receptor for (1,3)- $\beta$ -glucans, 590  
   recognising synthetic oligomeric, 587  
   role in, innate immune response, 592  
   signalling, 585  
 Dectin-1, 190–191  
 Dectin1-Fc chimeric protein  
   labelling of (1,3)- $\beta$ -glucans using, 398f  
 Degree of polymerization (DP), 11, 355  
 Denaturation – renaturation (DR) process, 88  
 Dentritic cell-associated C-type lectin-1 (CLEC7A). *See* Dectin-1  
 2-Deoxy-D-glucose (DDG), 446  
*Deschampsia antarctica*, 621  
*DEX1* gene, 475  
*Dianthus caryophyllus*, 441  
 Diatoms, 358  
   callose, 364  
   chrysotaminarin, 359–364  
 transmission electron micrographs of, 360f  
*Dictyonema glabratum*, 407  
*Dictyostelium discoideum*, 246–247  
 Digitonin, 296  
 Dilute solution  
   definitions of molecular parameters, 69–80  
 Dimethyl sulfoxide (DMSO), 13, 47  
 Dinoflagellates, 371  
 Dinophyceae. *See* Dinoflagellates  
 Dinophyta. *See* Dinoflagellates  
 Disaccharides, 342  
 Dithiothreitol, 244, 270  
 Dityrosine, 267  
 DMSO. *See* Dimethyl sulfoxide (DMSO)  
 DP. *See* Degree of polymerization (DP)  
 2D-PAGE analysis, 245  
 DR. *See* Denaturation – renaturation (DR) process  
*Drosophila GNBP3*, 570  
*Dunaliella salina*, 311  
 Dye  
   phenoxyzone, 11  
   triphenylmethane, 11  
 Dynactin, 269  
**E**  
 Ecdeiocoleaceae, 633  
*Ecdeiocolea monostachya*, 633  
 Echinocandin drugs, 271  
 Echinocandins, 271  
   LY303366, 264  
*Ecklonia radiata*, 370  
 EGTA, 302  
*Eisenia bicyclis*, 12, 139, 370  
*Eisenia foetida*, 570  
*Emiliania*, 357  
*Emiliania huxleyi*, 357  
 Endoplasmic reticulum (ER), 439  
*Equisetum arvense*  
   (1,3;1,4)- $\beta$ -glucans, 22

- Equisetum* spp, 630, 660  
*Erwinia amilovora*, 539  
*Escherichia coli*, 303, 390  
EST databases, 249  
Eubacterial (1,3)- $\beta$ -glucan synthase genes, 658  
*Euglena gracilis*, 153, 235–236, 238, 241–242, 244, 245, 246, 354  
biosynthesis of paramylon in, 356  
Euglenids, 354–356  
Euglenophyceae. *See* Euglenids  
Euglenophyta. *See* Euglenids  
*Evernia*, 632  
ExoI, 151
- F**  
FAB. *See* Fast atom bombardment (FAB)  
Factor C, 566  
Factor G, 566  
Fast atom bombardment (FAB), 370  
FK506, 266  
FKS gene, 443, 530, 658  
Fks1p-GFP, 267  
FKS proteins, 658  
Flagellariaceae, 633, 634  
*Flagellaria guineensis*, 633  
*Flagellaria indica*, 633  
Flagellin sensitive 2 (FLS2), 538  
*Flavobacterium meningosepticum*, 149  
Flg22 receptor, 538  
Flory universal viscosity constant, 71  
Fluorescence resonance energy transfer (FRET), 62  
Fluorescence spectroscopy, 61–62  
Fractionation, of cell walls, 390  
protocols for, 390, 391f  
*Fraxinus excelsior*, 449  
Free fatty acids (FFAs), 536
- FRET. *See* Fluorescence resonance energy transfer (FRET)  
Fungi  
(1,3)- $\beta$ -glucans in cell walls. *See* Cell walls, fungal  
fungal and yeast, extracellular polysaccharides, 387–388  
location of, 409f  
biofilms, 388  
*Fusarium graminearum*, 536
- G**  
Galactomannan, 18  
*Galdieria sulphuraria*, 311–312  
Gelation behaviour of (1,3)- $\beta$ -glucans, 93–103  
*Gemini pollen 1 (gem1)* mutants, 482  
Gentiobiose – enzyme models, 152  
GH16, 189–190  
GH17 (1,3)- $\beta$ -D-glucan endohydrolases, 122–128  
catalytic mechanism, 124–128  
classification of, 122  
properties of, 122  
structure of, 123–124  
substrate specificities of, 122–123  
transglycosylation reactions, 128  
GH17 (1,3;1,4)- $\beta$ -D-glucan endohydrolases, 128–132  
catalytic mechanism, 131  
classification of, 128–129  
3D structure of, 131–132  
properties of, 129–130  
GH26 (1,3;1,4)- $\beta$ -D-glucan endohydrolases, 132–133  
GH17 enzymes, 121  
GH16 family evolution of, 138–139  
*GhGSL1* in cotton fibres, 443  
Gigartinales, 630  
GIP-1 protein, 155  
GLS genes, 479
- Glucanase, 453  
Glucans, 259  
Glucan synthase-like(GSL) gene, 530  
*Gluconacetobacter xylinus*, 242  
Glucuronoxylmannan, 589  
Glyceollins, soybean, 341  
*Glycine max*, 302, 512  
Glycine-rich protein, cdiGRP, 542  
Glycogens, 366–367, 408–409  
Glycoprotein, 243  
Glycosidic oxygen protonation of, 125  
Glycosylation, 309  
Glycosylphosphatidylinositol, 17  
Glycosylphosphatidylinositol-anchored protein, 444  
Glycosyl-phosphatidylinositol (GPI), 261  
Glycosylphosphatidylinositol (GPI)-anchor, 395–396  
GNBP3 protein, 564  
GNBP proteins, 565  
Golden algae. *See* Chrysophytes  
*Golovinomyces cichoracearum*, 532, 534  
*Gossypium barbadense*, 441  
*Gossypium hirsutum*, 285, 292, 441  
GPI-anchor. *See* Glycosyl phosphatidylinositol (GPI)-anchor  
Gram-negative bacteria, 332  
Gram-positive *Cellulomonas*, 120  
Gravitropism, 513  
Green fluorescent protein (GFP), 443  
*Grifola frondosa*, 588  
GSL gene family members, in *Arabidopsis*, 531  
*Gsl1/gls1gsl5/* + plants, 479  
GSL (Glucan Synthase-Like) genes, 442, 443, 531  
*Gsl1* mutation, 479  
GSL5 null mutants, 542  
GSL proteins, 442, 543

- GT2 cellulose synthase-like (CSL) proteins, 659
- GT2 enzyme, 224
- GT48 enzymes, 309
- GT48 family, in angiosperms, 658
- GT2 protein-encoding gene, 658
- Guanine nucleotide-binding proteins (GNBPs), in *Drosophila*, 564
- Gut associated lymphoid tissue (GALT) expression, 600
- Gymnosperm pollen tubes, slow growth, 660
- Gyration mean square radius of, 70
- H**
- Haemolymph clotting, 565–567
- Haptophyceae, 357–358
- Haptophyta. *See* Haptophyceae
- Haramonas dimorpha*, 15, 372
- Hemerocallis fulva* meristematic cells of cell plate in, 426f cytokinetic apparatus in, 426f
- Heterokontophyta. *See* Diatoms
- Histoplasma capsulatum*, 400
- Hordeum vulgare*, 292, 298, 441, 633
- Horseshoe crab TGase, 567
- Hrp* mutants, 538
- Huggins constant, 89
- Human isoform (hDectin-1E), 583
- Hydrodynamic volume of polysaccharide, 48
- p-hydroxymercuribenzoate, 244
- Hyphae** comparison of wall composition between, and Conidia, 399 antifungal compounds, 401–402 cell wall mutants, 402 dimorphic forms, 400–401 environmental conditions, 401
- Hyphal, cell walls chemical composition of, 368t
- I**
- I.C. streptococcal type 37 (1,3;1,2)- $\beta$ -glucan, 342
- IgE responses, 598
- IMC carcinoma, 598
- IMD pathway, 564
- Immunoreceptor tyrosine-based activation-like motif (ITAM), 583
- Inositol 1,4,5-trisphosphate (IP3), 513
- Ionomycin, 501
- Ishige okamurai*, 370
- J**
- Joinvilleaceae, 633
- Juncaceae, 634
- K**
- Kappapheus*, containing sulfated form glucan, 659
- Kappaphycus alvarezii*, 630
- Keratin sulfate, 134
- Kinesin, 479
- K1 killer toxin, 271
- KRE6, 248
- KRE genes mutations in, 402
- Kre5p, 272
- Kuhn's length, 69
- L**
- Lactosylceramide, 580–581
- Lamellibrachia satsuma*, 248
- Laminaria cichorioides*, 370
- Laminaria digitata*, 122, 370
- Laminaria hyperborea*, 369, 370 seasonal variation of laminarin and ash in, 372f
- Laminaria longicrusis*, 371
- Laminaria saccharina*, 176
- Laminaribiose phosphorylase, 153
- Laminarin, 14–15, 236, 370, 371, 589 seasonal variation of, in *Laminaria hyperborea*, 372f
- Larix laricina*, 14
- Lectin, 570
- Lemna gibba*, 508
- Lentinan, 78 gelation mechanism proposed for, 97
- Lentinus edodes*, 78
- Leptomitus lacteus*, 367
- Leucine-rich repeats (LRRs), 526
- Leucosin, 16, 359
- Lewis lung carcinoma, 598
- Lichens (1,3)- $\beta$ -glucans in, 406–407
- Light-induced paramylon kinetics of, 355, 356f
- Lignin, 140
- Limpet pollen (lip)* mutants, 482
- Linear (1,3)- $\beta$ -glucans, 12–14, 234f, 399, 407. *See also* Curdlan dilute solution properties of, 70–73 in walls, gametophytes and gametes, 466
- Linear (1,3;1,4)- $\beta$ -glucans, 20–24
- Lipopolysaccharides (LPS), 331, 342, 566
- Lipoteichoic acids, 567
- Lolium multiflorum*, 292
- Lolium perenne*, 633
- Lophocolea bidentata*, 23
- LPS-induced haemocyte exocytosis, 566. *See also* Lipopolysaccharides (LPS)
- LRR transmembrane receptor kinase, 538
- Lycopersicon esculentum*, 441

**M**

- Macrocystis*, 368  
 Magic angle spinning (MAS), 18  
*Magnolia kobus*, 449  
 MALDI. *See* Matrix-assisted laser desorption/ionization (MALDI)  
 MALS. *See* Multi-angle laser-light scattering (MALS)  
*Malus domestica*, 441  
 MAMP-induced signalling, 538  
*Manduca sexta*, 570  
 Mannan, 587  
 Mannoproteins, 16, 261  
 Marine diatom. *See* Diatoms  
 Mark-Houwink-Sakurada (MHS), 81  
   of cereal, 82  
 MAS. *See* Magic angle spinning (MAS)  
 Matrix-assisted laser desorption/ionization (MALDI), 370  
 Megagamete development, in *Arabidopsis*, 483–487  
   different stages of, 485  
   ovule development, 484  
 Melanin production, 568  
 Melanization, of microbial pathogen, 568  
 Melanization cascade, 568–571  
 Meristematic cells, 425  
   cell plate formation in, 428  
   cell plate in, of daylily, 426f  
   cytokinetic apparatus in, of daylily, 426f  
 Metal toxicity, 502  
 Methanol, 54  
 α-methyl-D-glucoside, 242  
 MHS. *See* Mark-Houwink-Sakurada (MHS)  
*Micrasterias*, 630, 659  
 Microgamete development, 466  
   appearance of callose during, 468  
   in *Arabidopsis*, 466, 469

- meiosis to cytokinesis, 469–470  
 premeiosis I and II, 469  
 released microspore I, 470  
 released microspore II, 470–471  
 second mitotic division to, 471  
 cell wall compositions and changes, 467  
 functions for callose during, 473  
   during cytokinesis, 476–480  
   megagamete development, 482–483  
   in post-meiotic male gamete development, 480–482  
   in special cell wall, 474–478  
 pollen development, 467  
*Micromonas pusilla*, 311  
 Microsporogenesis, 466  
 Microtubules, 428  
 MLG:xyloglucan  
   endotransglucosylase (MXE), 660  
*Mlo* mutants, 536  
 MLO protein, 537  
 Mn toxicity, 507–508  
*Monodus subterraneus*, 23, 372, 632  
 Moth bean (*Vigna aconitifolia*), 308  
 Movement proteins (MPs), 452  
   interactions with microtubule and microfilaments, 453  
 MS/MS proteomic analysis, 443  
 Multi-angle laser-light scattering (MALS), 11  
 Multinucleate cytoplasm, 429  
 Mung beans (*Vigna radiata*), 301, 307  
 Mutants  
   cell wall, 402  
 Mycolaminarin, 15, 236, 237, 365, 366–367, 408  
 MyD88-coupled TLRs, 585

**N**

- N-acetyl-β-glucosaminidase*, 146, 149, 151, 388  
*N-acetylchitooligosaccharides*, 534  
*N-acetylglucosamine* (GlcNAc), 261  
*N-acetylhexosamine* residues, 15  
*NaGSL1* gene, 443, 530  
*NaGSL1* protein, 295–296, 297, 308  
 NBS-LRR proteins, 526  
 NCD. *See* Nuclear-cytoplasmic domain (NCD)  
*NdvB* genes, 219–221  
*NdvC* genes, 221–222  
*NdvD* genes, 221–222  
*Ndv* genes, 216–219  
 Nebenlinien, 364  
 Neo β-glucans, 28  
*Nereocystis*, 368  
*Neurospora crassa*, 262, 263, 264, 288, 394  
 Neurotrophin 3 (NT-3), 596  
*Newropogon*, 632  
 Newtonian plateau, 82  
 Newtonian viscosity, 90  
*Nicotiana*, 284  
*Nicotiana alata*, 14, 287, 297, 304, 308  
*Nicotiana benthamiana*, 441, 539  
*Nicotiana plumbaginifolia*, 541  
*Nicotiana sylvestris*, 541  
*Nicotiana tabacum*, 441, 452  
   pollen tube, 473  
*Nilaparvata lugens*, 452  
 NMR. *See* Nuclear magnetic resonance (NMR)  
 N-myristoylation, 309  
 NO, in primary signals of anoxia, 511  
 Nodulation, 338–340  
 Non-Newtonian fluids, 82  
*NtrBC* gene, 206  
*NtrB* gene, 206–207  
*NtrC* gene, 207–208  
*NtrYX* gene, 208–209

- Nuclear – cytoplasmic domain (NCD), 426
- Nuclear factor of activated T-cells (NFAT), 585
- Nuclear magnetic resonance (NMR), 47, 390  
in solid state, 53–54
- Nystatin, 271
- O**
- Ochromonas malhamensis*, 365
- Ochromonas tuberculatus*, 365
- OeCBM43A*, 186–187
- OeCBM43B*, 187
- Olea europaea*, 186
- (1,3;1,4)-oligoglucosides, 24
- Oligomers, 65
- Oligosaccharide, 5
- Oomycetes  
(1,3)- $\beta$ -glucans in  
oomycete cell wall, 407–408  
storage (1,3)- $\beta$ -glucans,  
408–409
- Oomycota, 365  
cellulin, 367  
cell wall polysaccharides,  
367–368  
chemical structure of, 366f  
mycolaminarin, 366–367
- Opsonins, 571
- Optical rotatory dispersion (ORD)  
in curdlan, 72
- ORD. *See* Optical rotatory  
dispersion (ORD)
- Oryza sativa*, 292  
cross-sections of, 430f  
wall development in endosperm  
of, 431f
- Ostreococcus tauri*, 310
- Oxidation  
of scleroglucan, 80
- P**
- Pachyman, 13
- Paenibacillus*, 390
- Papulacandin B, 244
- Papulacandin B-resistant mutant,  
262
- Papulacandins, 271
- Paracoccidioides brasiliensis*, 396,  
400
- Paramylon, 13, 235, 242
- Paramylon granules, 354, 355  
biosynthesis of, in *E. gracilis*,  
356
- freeze-etch electron micrographs  
of, 355f
- light-induced, kinetics of, 355,  
356f
- Paramylon synthase, 245
- Parmelia*, 632
- Parmeliaceae, 632
- Parmotrema*, 632
- Pathogenesis-related (PR) proteins,  
443
- Pavlova mesolychnon*, 13, 357
- Pavlova prymnesiida*, 357
- Pd. *See* Plasmodesmata
- Pd callose synthase-associated  
protein, 443
- Pd regulation, by callose turnover,  
444–446
- Pellicle, 331
- Pen1-1* mutant, 536
- PEN1/ROR2 syntaxins, 537
- Peranema trichophorum*, 13, 355
- Pergularia daemia*, 476
- Periclinal wall  
callose and, 431–432
- Peridinium westii*, 23, 371, 632
- Periodate-Schiff reaction, 11
- Peronospora parasitica*, 293, 532
- PGG-(1,3)- $\beta$ -glucan, 591
- Phaeocystis*, 357, 358
- Phaeocystis globosa*, 357
- Phaeodactylum tricornutum*, 248,  
249, 359
- Phaeophyceae, 368–371
- Phaseolus vulgaris*, 441, 443, 512
- Phenoloxidase (PO), 568
- Phenoxyzone dye, 11
- Phragmoplastin, 295, 302
- Phragmoplasts, 425  
development in syncytial  
systems, 426–428, 427f
- Phythium debaryanum*, 408
- Phytoalexins, 527
- Phytophthora*, 235
- Phytophthora cinnamomi*, 235, 243,  
246, 247
- Phytophthora* genus, 248
- Phytophthora infestans*, 249, 368
- Phytophthora palmivora*, 237, 241,  
256, 366  
spores, 408
- Phytophthora parasitica*, 15, 408
- Phytophthora ramorum*, 249
- Phytophthora* species, 248
- Phytophthora sojae*, 155, 238, 240,  
246, 249, 340
- Pichia pastoris*, 390
- PI3K/Akt signalling, 592, 596
- PI3K inhibitor, 592
- PI3K pathway, 597
- Pinnularia*, 359, 364
- PIR. *See* Putative proteins with  
internal repeats (PIR)
- Pir proteins, 17
- Pisum sativum*, 441, 512, 535
- Plant–bacteria interactions  
callose in, 537–539  
extracellular enzymes for, 539  
MAMP elicitors, 538  
producing negatively charged  
exopolysaccharides in, 539
- Plant defense response, suppression  
of, 340–342
- Plant–fungi interactions  
and callose-containing papilla  
formation in, 533  
callose in, 532–537
- Plant–microbe interactions  
enzymes of general cell wall,  
536  
function of callose in, 525

- Plant-pathogen interactions, 543  
papillae, as defence mechanism, 529  
role of callose in, 543
- Plant-virus interactions  
callose in, 539–542  
AtBG T-DNA insertion mutants, 540  
barrier to viral spread, 542  
callose deposition, at  
plasmodesmata, 540  
cdiGRP, role of, 542  
expression antisense (1,3)- $\beta$ -glucanase, 541  
localization of, 541  
movement proteins, role of, 541
- Plasmalemma, 443
- Plasmodesmata, 439. *See also*  
(1,3)-glucanase  
AI treatment induced callose deposition around, 506  
callose deposition at, 531  
callose levels, 540  
callose turnover at, 540  
cell-to-cell transport processes through, 502  
coefficient of conductivity of, 540  
conductivity, 440–441  
electron micrographs of, 440  
flow resistance due to, 512  
intercellular communication through, 507  
intercellular trafficking via, 541  
plant viruses replication through, 539  
structural changes in, 441  
structure of, 439–440  
synthesis and hydrolysis of callose, 441  
of virus-infected plant tissue, 539
- Platelet-derived growth factor A (PDGF-A), 597
- Platelet-derived growth factor B (PDGF-B), 597
- Pleurochrysis haptonemofera*, 357
- Pleurotus tuber-regium*, 80
- Plodia interpunctella*, 189, 570
- Pmr4* mutants, 534, 542, 543
- Pneumococcal type 37 (1,3;1,2)- $\beta$ -glucan  
enzymology, 224  
membrane topology, 223–224  
molecular genetics, 222–223
- Pneumocystis*, 586
- Poaceae, 658, 660
- Polygonum*-type development, 486
- Polygonum*-type embryo sacs, 485
- Polypodium vulgare*, 448
- Polysaccharide chain network formation, 52
- Polysaccharide conformation principles of, 48–52
- Polysaccharides, 660  
cell walls, of oomycetes, 367–368  
chrysotolaminarin and, 359–364  
degradation of, 365  
extracellular, fungal and yeast (1,3)- $\beta$ -glucan, 387–388  
storage, in euglenids, 354  
types and structures of, in protozoans and chromistans, 354t  
x-ray fibre diffraction of, 53
- Populus tremula*, 137
- Populus trichocarpa*, 288
- Poria cocos*, 13, 72
- Porphyra purpurea*, 312
- Poterioochromonas malhamensis*, 365
- PPB. *See* Predictive preprophase band (PPB)
- Predictive preprophase band (PPB), 425
- Prenylation, 309
- Product entrapment, 245, 261, 262, 263, 297
- ProPO-activating enzyme, 571
- ProPO system, 568
- (1,3)- $\beta$ -glucan-induced activation of, 570
- Proteinaceous inhibitors of plant, 154–155
- Protein disulfide isomerase (PDI), 301
- Protein kinase B (PKB), 592
- Proteins  
biochemical analysis of, 188–192  
CBM. *See* Carbohydrate-binding modules (CBM)  
GPI-anchored, 395–396  
overview, 171–172  
PIR. *See* Putative proteins with internal repeats (PIR)  
structural properties of, 172–174
- Protozoans  
types and structures of polysaccharides in, 354t
- Proxin, 567
- Prymnesiophyceae. *See*  
Haptophyceae
- Pseudoalteromonas carragenovora*, 135
- Pseudomonas aeruginosa*, 596
- Pseudomonas syringae*, 536
- Psilotum nudum*, 448
- PSSA gene, 205
- Pullulan, 587
- Putative proteins with internal repeats (PIR), 394
- Saccharomyces cerevisiae* and, 395
- Pythium aphanidermatum*, 408  
cell wall glucan, 18–19
- Q**
- Quantazyme, 390
- Quartet (*qrt*) mutations ( *qrt1-3* ), 480
- Quercus rubra*, 505
- R**
- Ralstonia solanacearum*, 539
- Ramalina celastri*, 407
- Ramalina usnea*, 407

- RelA gene, 209–211  
 Resorcin blue, 11  
 Resorcinol blue, 16  
 Resting cells  
     chrysolaminarin and, 362  
 Restionaceae, 633, 634  
*Reviews of Pure and Applied Chemistry*, 1  
*Rhizobium*, 99, 329  
*Rhizobium loti*, 19  
*Rhizopus arrhizus*, 135  
 Rhodophyte sulfated (1,3;1,4)- $\beta$ -glucans, 24  
*Rhodothermus marinus* (*RmCBM4-2*), 176  
 Rho1 GTPase, 531  
 Rho1p, 263  
 Rice. *See* *Oryza sativa*  
*Rimelia*, 632  
 R/K-X-G-G, 263  
 ROP1, 302  
 R protein, 527
- S**  
*Saccharomyces cerevisiae*, 12, 16, 121, 251, 261, 262, 271, 272, 285, 389, 391–392, 394, 587  
     cell wall glucan, 16–18  
     PIR proteins and, 395  
     wall composition of, 393f  
 Salicylic acid signal transduction, 543  
 Saponins, 300  
*Saprolegnia*, 235  
*Saprolegnia monoica*, 237–238, 241, 242, 243, 244, 245–246, 247, 248–249  
*Sarcina ventriculi*, 24, 632  
 SAXS. *See* Small angle X-ray scattering (SAXS)  
 Scanning electron microscopy, on *agrobacterium*, 331  
 Scavenger receptors, 581  
 S-celluloside – enzyme complexes, 152
- Schizophyllum commune*, 66, 73  
 Schizophyllum (SPG) glucan, 595  
*Schizophyllum commune*, 394  
*Schizosaccharomyces pombe*, 263, 389, 392  
     localization of (1,3)- $\beta$ -glucans in septum of, 399f  
     localization of (1,6)- $\beta$ -glucans in septum of, 399f  
 Scleroglucan, 75, 589  
 Scleroglucan – borax complexes  
     gelation of, 96  
*Sclerospora graminicola*, 535  
*Sclerotium glucanicum*, 75  
 SDS-PAGE analysis, 245  
 SDS-PAGE chromatography, 154  
 SDS-PAGE gels, 243, 297  
 SEC-MALS/RI. *See* Size-exclusion chromatography with multi-angle light scattering/refractive index technique (SEC-MALS/RI), 81  
 Sedimentation analysis, 61  
*Selaginella kraussiana*, 448  
*Sensu lato*, 660  
*Serratia marcescens*, 390  
*Sesbania rostrata*, 336  
 Shear thinning, 85  
*Sidecar* mutant, 481  
 Side-chain-branched (1,3;1,6)- $\beta$ -glucans  
     behavior of concentrated solutions, 85–88  
     concentrated solution properties of, 88–90  
     dilute solution properties of, 73–80  
     gelation of, 96–99  
     solution properties of charge, 80–81  
 Sieve plates. *See also* Plasmodesmata  
     callose deposited in, 446, 512  
     chemical stresses inducing callose in, 447  
     cytokinin promoting callose, 449
- localization of callose at Pd and, 441–442  
 plamodesmatal transport and in, 657  
 pores, 439  
 sensitive response of phloem, 513  
*Sinapsis alba*, 143  
*Sinorhizobium*, 338  
*Sinorhizobium meliloti*, 20, 339  
 Size-exclusion chromatography, 154  
 Size-exclusion chromatography with multi-angle light scattering/refractive index technique (SEC-MALS/RI), 81  
*Skeletonema costatum*, 359  
 S-laminaribioside – enzyme complexes, 152  
 Small angle X-ray scattering (SAXS), 61  
*Solanum lycopersicum*, 512  
 Solid state  $^{13}\text{C}$  NMR  
     of (1,3)- $\beta$ -glucans, 58–60  
 Sophorose – enzyme models, 152  
 Sorbitol, 98  
 S-palmitoylation, 309  
 Sparganiaceae, 634  
 Specific cell types in cereal grains, composition, 623t  
 Spectroscopy, 54  
     fluorescence, 58, 61–62, 183–184  
     mobility-resolved, 54  
     near-infrared, 638  
     peptide mass fingerprinting and tandem mass, 288  
 Saturation Transfer Difference NMR, 142  
*Sphaerotiltrichia divaricata*, 370  
*Spinacea oleracea*, 298  
 Spindle pole body (SPB), 269  
*Sporobolus africanus*, 441  
*Sporothrix schenckii*, 401  
*Sporotrichum dimorphosporum*, 12

- STAND ATPases, 526  
*Staphylococcus aureus*, 599  
*Stauroneis amphioxys*, 359  
 Steady shear viscosity  
   of curdlan, 72  
   of scleroglucan, 87  
*Stereocaulon ramulosum*, 407  
*Streptococcus pneumoniae*, 5,  
   20, 222, 342. *See also*  
   Pneumococcal type 37  
   (1,3;1,2)- $\beta$ -glucanTts gene  
*Streptococcus pyogenes*, 215, 342  
*Streptococcus suis*, 599  
*Strongylocentrotus purpuratus*, 135  
*Suberites domuncula*, 572  
 Sucrose synthase (SuSy), 301–302,  
   433, 500–501  
 Sulfated heteropolysaccharides, 593  
 SuSy. *See* Sucrose synthase (SuSy)  
 SuSy activity in submerged potato  
   roots, 511  
 Swi4-Swi6 cell cycle box-binding  
   factor (SBF), 265  
 Syk independent signalling  
   pathways, 585  
 Synaptosome-associated protein  
   receptor, 536  
 Syncytial systems  
   alveolation in, 429–432  
   phragmoplast development in,  
     426–428, 427f  
 Synthetic  $\beta$ -oligoglucosides, 24
- T**  
*Tachylepus tridentatus*, 565  
*Thalassiosira*, 364  
*Thalassiosira pseudonana*, 248,  
   249, 359  
*Thalassiosira weissflogii*, 16, 248  
*Thermoplasma acidophilum*, 505  
*Thermotoga neapolitana*, 390  
 TLR2/MYD88 pathways, 585  
*TmCBM4-2*, 177–178, 179, 191  
 TMV-induced lesions, 541  
 TMV infection, 452  
 TNF  $\alpha$  production, 598
- Toll interleukin-1 receptor  
   nucleotide binding (TIR-NB) proteins, 537  
 Toll-like receptors (TLR), 585  
 Transforming growth factor alpha  
   (TGF $\alpha$ ), 597  
 Transforming growth factor beta  
   (TGF $\beta$ ), 597  
 Transglutaminase (TGase), 566  
 Trichloroacetic acid, 243  
*Trichoderma harzianum*, 390  
 Triphenylmethane dye, 11  
*Triticum aestivum*, 441, 639  
*Triticum durum*, 639  
*Tropaeolum majus*, 137  
 Trypsin, 308  
 Tts gene, 222  
   enzymology, 224  
   membrane topology of, 223–224  
 TTSS effectors, 538  
 Tubular – vesicular network (TVN),  
   428  
 TVN. *See* Tubular-vesicular  
   network (TVN)  
*Two-in-one pollen (tio)* mutants,  
   482  
 Type III secretion system (TTSS),  
   537–538  
*Typaceae*, 634
- U**  
 UDP- $\alpha$ -D-[ $^{14}$ C]glucose, 246  
 UDP- $\alpha$ -D-glucose, 239–240  
 UDP- $\alpha$ -D-[ $^3$ H]glucose, 246  
 UDP degrading enzyme, 241  
 UDP-Glc, 297  
 $\alpha$ -[ $^{32}$ P]UDPglucse, 245  
 UDPglucose: glycoprotein  
   glucosyltransferase  
   (UGGT), 272  
 UDP-glucose-binding proteins,  
   245  
 UDP-glucose pyrophosphorylases,  
   362–363  
 UDP-glucose transferase (UGT1),  
   435
- UGT1. *See* UDP-glucose  
   transferase (UGT1)  
*Ulva lactuca*, 23, 630  
*Usnea*, 632
- V**  
 Vascular endothelial growth factor  
   (VEGF), 597  
*Vibrio cholerae*, 151  
*Vicia faba*, 510  
*Vigna radiata*, 301  
*Vigna sinensis*, 512  
*Vigna unguiculata*, 503  
 Viscometers, 74  
   Zimm-Crothers type, 85  
*Vitis vinifera*, 14  
*Volvox carteri f. nagariensis*, 311
- W**  
 Winter-Chambon criterion  
   for gelation, 97, 98  
 Wortmannin, 592
- X**  
*Xanthomonas campestris*, 538, 539  
 XEH. *See* Xyloglucan  
   endohydrolases (XEH)  
 X-ray diffraction  
   (1,3)- $\beta$ -glucans, 54–58  
   in polysaccharide chain  
   conformation, 52–53  
 XTH. *See* Xyloglucan  
   endotransglycosylases/  
   hydrolase (XTH) group  
 Xylanase Z-like domain, 566  
 Xyloglucan endohydrolases (XEH),  
   137  
 Xyloglucan endotransglycosylases/  
   hydrolase (XTH) group,  
   133, 137–138  
 Xyloglucans, 137  
*Xyridaceae*, 634
- Y**  
*Yarrowia lipolytica*, 396  
 Yeasts

- (1,3)- $\beta$ -glucans extracellular polysaccharides, 387–388
- chemical composition of cell wall of, 389*t*
- KRE genes. *See* KRE genes
- protocol for extraction of, 390, 391*f*
- schematic representation of CWP in, 395*f*
- structure and molecular organization of (1,3)- $\beta$ -glucans, 391–396
- Yeast cell wall
- localization of (1,3)- $\beta$ -glucan in, 261
- structure and (1,3)- $\beta$ -glucan, 260
- Young's modulus, 103

**Z**

- Zea mays*, 302, 441, 512, 633
- Zero shear viscosity, 82
- Zimm plot analysis, 82