

- Abietic acid, correlation with lanosterol, 67.
- Absolute configuration, of steroids and triterpenes, 18.
- Acetic acid, as biogenetic triterpene precursor, 116-118, 120, 124.
- Acetoin, as degradation product, 87.
- "Aglycone 157", and derivatives, 173.  
— natural occurrence, 173.
- "Aglycone 178", and derivatives, 173.  
— natural occurrence, 173.
- Agnoesterol, and derivatives, 139.  
— natural occurrence, 139.  
— total synthesis of, 109.
- Allobetulone, bromination of, 34-35.
- Alnusfolienediolone, and derivatives, 165.  
— natural occurrence, 165.
- Alnus glutinosa*, as triterpene source, 165.
- Ambrein, correlation with lanosterol, 67.
- Androstanes, 4,4,14-trimethyl derivatives from eburicoic acid, 130.
- Animal tissue, as triterpene source, 135, 136, 137.
- Anthothecol, and derivatives, 169.  
— natural occurrence, 169.
- Antibiotics, 39, 80, 82, 100, 101, 130.
- Artocarpus integrifolia*, as triterpene source, 145, 155.
- Artostenone (see cycloartenone).
- Aspergillus fumigatus*, as triterpene source, 178.
- Aster tataricus*, as triterpene source, 178.

## B

- Baeyer-Villiger oxidation, in side-chain degradation, 86.
- Betula alba*, as triterpene source, 164.
- Betulafolienetetraol, 164.
- Betulafolienetriol, and derivatives, 164.  
— dehydrogenation of, 104.  
— natural occurrence, 164.
- Biogenesis, of terpenes, 115-125.
- "Biogenetic isoprene rule" (see isoprene rule).

(\*) Bold face numbers indicate catalogue pages.

- Biological activity, of steroid hormone analogues, 46, 130.  
— of tetracyclic triterpenes, 129-130.
- Birch leaves, as triterpene source, 164.
- Bis (lanost-8-en-3,7,11-trione-2 $\xi$ -yl) methane, 36.
- Black Ash leaves, as triterpene source, 165.
- Black fungus of birch, as triterpene source, 177.
- Boat form, in biogenesis of dammaranes, 122.  
— in cyclohexane derivatives, 24.  
— in dehydration involving C-3 hydroxyl groups, 36.  
— of ring A in 2 $\beta$ -bromolanostanone derivatives, 34-35.  
— of rings B or C in eupheol, 50.  
— of ring C in euphanes, 62.
- 2 $\alpha$ -Bromolanostanone, preparation of, 34.
- 2 $\alpha$ - and 2 $\beta$ -Bromolanost-8-en-3-one, preparation of, 34.
- Bryonia dioica*, as triterpene source, 173.
- Bryogenin (see "Aglycone 157").
- "Buttressing effect", 31, 36.
- Butyrospermol, and derivatives, 155.  
— and configuration at C-9, 49.  
— correlation with iso-euphenol, 70.  
— dihydro derivative, oxidation of, 69-70.  
— —rearrangement of double bond, 48.  
— — C-3 acetate, oxidation of, 70, 125.  
— natural occurrence, 21, 155.  
— oxidation of, 50.  
— position of double bond in, 48-49.  
— rearrangement of double bonds in, 94-95.
- Butyrospermone, 155.
- Butyrospermum parkii*, as triterpene source, 138, 155.
- Buxus sempervirens*, as triterpene source, 153.

## C

- Camphor, microbiological degradation of, 125.

- Canarium* spp., as triterpene source, 157, 158.
- Carotenoids, 117.
- mevalonic acid in biosynthesis of, 118.
- Casimiroa edulis*, as triterpene source, 170.
- Casimiroside (see obacunone).
- Cedrela toona*, as triterpene source, 172.
- Cedrelone, 23.
- and derivatives, 172.
  - biogenesis of, 126.
  - natural occurrence, 172.
  - ring B contraction, 56, 76.
- Cephalosporin P<sub>1</sub>, absence of C-4 methyl in, 30.
- and derivatives, 179.
  - isolation of, 21.
  - natural occurrence, 179.
  - physiological properties of, 130.
- Cephalosporium* sp., as triterpene source, 179.
- Cholesta-8,24-dien-3 $\beta$ -ol, 4,4-dimethyl derivative (see 4,4-dimethylcholestadienol).
- Cholestan-3 $\beta$ -ol, 4 $\alpha$ ,14 $\alpha$ -dimethyl-24-methylene-9,19-cyclo derivative (see cycloeucaenol).
- Cholestanone, steric hindrance of C-3 keto group, 40.
- Cholest-8-en-3 $\beta$ ,6 $\alpha$ -diol (see Peniocerol).
- 14 $\alpha$ -methyl derivative (see Macdougallin).
- $\Delta^7$ -Cholestenol, in cholesterol biosynthesis, 126, 127.
- $\Delta^8$ -Cholestenol, in cholesterol biosynthesis, 126, 127.
- Cholestenol, 4 $\alpha$ -methyl derivative (see methylcholest-7-en-3-ol).
- Cholest-8-en-3 $\beta$ -ol, 4 $\alpha$ -methyl derivative (see methylcholest-8-en-3-ol).
- Cholest-7-en-3 $\beta$ -ol, 4 $\alpha$ -methyl derivative (see methylcholest-7-en-3-ol).
- Cholest-7-en-3 $\beta$ -ol, 4 $\alpha$ -methyl-24-ethylidene derivative (see citrostadienol).
- $\Delta^7$ -Cholesten-3-one, 4-hydroxymethylene derivative, in cholesterol biosynthesis, 127.
- Cholesterol, biogenetic relationship with lanosterol, 120, 126-129.
- conversion to 7-dehydro derivative, 109.
  - 7-dehydro derivative in cholesterol biosynthesis, 126.
  - 24-dehydro derivative in cholesterol biosynthesis, 126.
  - — side chain in, 81.
  - mevalonic acid in biogenesis of, 117, 120.
  - side chain in, 81.
  - — structure determination of, 100.
  - structural similarity to tetracyclic triterpenes, 18.
  - total synthesis, 107.
- Circular dichroism, use in structure elucidation, 106.
- Citrostadienol, 39.
- absence of C-4 methyl in, 30.
  - and derivatives, 135-136.
  - correlation with steroids, 40.
  - natural occurrence, 21, 135-136.
  - rearrangement of double bond in, 47.
  - side chain in, 83.
  - — biogenesis of, 125.
- Citrostanone, physical properties and structure of, 136.
- reduction of C-3 keto group, 32.
- Citrullus* spp., as triterpene source, 176.
- Citrus seed oil, as triterpene source, 170, 171.
- Citrus* spp., as triterpene source, 171.
- Classification of tetracyclic triterpenes, 17 ff.
- Conformational analysis, of tetracyclic triterpenes, 24 ff.
- "Conformational transmission", 36, 38.
- Cruciferae, as triterpene source, 176.
- Cryptosterol (see lanosterol).
- Cucumis* spp., as triterpene source, 174, 175.
- Cucurbitaceae, as triterpene source, 128, 174, 175, 176.
- Cucurbitacin A, natural occurrence, 20, 174.
- physical properties and structure of, 174.
  - reaction with base, 72-73.
- Cucurbitacin B, 128.
- natural occurrence, physical properties and structure of, 174-175.
  - nuclear magnetic resonance study of, 44.
  - preparation of, 43.
- Cucurbitacin C, natural occurrence, 20, 175.
- physical properties and structure of, 175.

- reaction with base of C-19 aldehyde derivative, 72-73.
  - Cucurbitacin D, natural occurrence, physical properties and structure of, 174-175.
  - Cucurbitacin E, 72, 128.
    - anti-cancer activity of, 130.
    - natural occurrence, physical properties, and structure of, 176.
    - oxidation of, 43.
    - tetrahydro derivative, methyl ether, 89.
  - Cucurbitacin F, natural occurrence, 20.
    - physical properties, 176.
  - Cucurbitacin G, physical properties, 176.
  - Cucurbitacin I, natural occurrence, physical properties, and structure of, 176.
    - preparation of, 43.
  - Cucurbitacin J, physical properties, 176.
  - Cucurbitacin K, physical properties, 176.
  - Cucurbitacin L, physical properties, 176.
  - Cucurbitacins (see also individual cucurbitacins), 74.
    - biogenesis of, 123-124.
    - dehydrogenation of, 102.
    - enzymic hydrolysis, 20.
    - natural occurrence, 20, 128.
    - oxidations of rings B and C, 71-73.
    - physical properties and structures of, 173 ff.
    - physiological properties of, 129-130.
    - reactions of ring A, 43-44.
    - rearranged methyl groups in, 31.
    - ring A contraction in, 39.
    - side-chain degradation, 87-89.
    - side-chains in, 82.
    - structure determinations of, 71-74.
    - structures of ring A, 29.
  - Cucurbita pepo*, as triterpene source, 175, 176.
  - Cucurbitone B, 7-oxo derivative, 72.
    - reaction with HCl, 72.
  - Cucurbits, origin of bitterness in, 128.
  - 9,19-Cyclolanostan-3 $\beta$ -ol (see cycloartanol).
  - 9,19-Cyclolanost-24-en-3 $\beta$ -ol (see cycloartenol).
  - Cycloartane, total synthesis of, 109.
  - Cycloartanol, 63, 92.
    - and derivatives, 144.
    - 4 $\beta$ -demethyl-24-methylene derivative (see cycloeucaleanol).
    - 24-methylene derivative, natural occurrence, 152.
    - — structures and physical properties of derivatives of, 152.
    - natural occurrence, 144, 145.
  - Cycloartanone, bromination of, 35.
    - preparation of, 41.
  - Cycloart-23-ene-3 $\beta$ ,25-diol, and derivatives, 146.
    - natural occurrence, 146.
  - Cycloart-25-ene-3 $\beta$ ,24 $\xi$ -diol, and derivatives, 145-146.
    - natural occurrence, 146.
  - Cycloartenol, acetate, 52.
    - and derivatives, 144-145.
    - correlation with cycloeucaleanol, 41.
    - degradation of side-chain, 94.
    - 23-dihydro-25-hydroxy derivative, natural occurrence, 145.
    - natural occurrence, 128, 145, 146.
    - occurrence as 3-ferulate, 21, 145.
    - structure determination, 52.
  - Cycloartenone, natural occurrence, 145, 146.
    - physical properties and structure of, 145.
  - Cycloart-1-en-3-one, preparation of, 35.
  - Cyclolaudenol, 95.
    - and derivatives, 151.
    - natural occurrence, 151.
    - side-chain in, 83.
    - — structure of, 94.
  - Cyclobuxine, and derivatives, 152-153.
    - natural occurrence, 153.
  - Cycloeucaleanol, 39.
    - absence of C-4 methyl in, 30.
    - and derivatives, 150-151.
    - correlation with cycloartenol, 41.
    - natural occurrence, 151.
  - Cycloeucalenone, reduction of C-3 keto group, 32.
  - $\beta$ -Cyclo-orysterol, rearrangement of side-chain double bond in, 95.
- D
- Dammadienol (see dammaradienol).
  - Dammadienolic acid (see dammarenic acid).
  - Dammar resins, as triterpene source, 21, 161, 162, 166.
  - Dammaradienol, and derivatives, 161.
    - identification by mass spectrum, 105.

- natural occurrence, 105, **161**.
- side-chain in, 82.
- Dammara-20,24-dien-3 $\beta$ -ol (see dammaradienol).
- Dammarane, ring A distortion in, 28.
  - structure of ring D, 74.
  - use of basic skeleton in naming, 20.
- (13 $\xi$ ,17 $\xi$ )-Dammarane, 3 $\beta$ ,12 $\xi$ -dihydroxy-20 $\xi$ ,25-epoxy derivative (see panaxadiol).
- Dammaranediols, 17, 122.
- Dammaranes, biogenesis of, 121-122.
  - boat form in, 31.
  - bromination of, 34-35.
  - dehydrogenation of, 103.
  - natural occurrence, 128, **161 ff.**
  - physical properties and structures of, **161 ff.**
  - preparation from euphanes, 50.
  - side-chain in, 81-86.
  - — rearrangement of double bonds in, 96.
- Dammara-3,4-seco-4(30),24-dien-3-oic acid, 20 $\xi_2$ -hydroxy derivative (see dammarenolic acid).
- Dammarenediol I, and derivatives, **161-162**.
  - natural occurrence, **162**.
- Dammarenediol II, and derivatives, **162**.
  - natural occurrence, **162**.
- Dammarenediols, 122.
  - side-chain in, 82.
- Dammara-24-ene-3 $\beta$ ,20 $\xi_1$ -diol (see dammarenediol I).
- Dammara-24-ene-3 $\beta$ ,20 $\xi_2$ -diol (see dammarenediol II).
- 13 $\xi$ ,17 $\xi$ -Dammara-24-ene, 3 $\beta$ ,12 $\xi$ ,20 $\xi$ -trihydroxy derivative, natural occurrence, **166**.
- Dammara-24-ene-3 $\alpha$ ,12 $\beta$ ,17 $\alpha$ (?),20 $\beta$ -tetraol (see betulafolienetetraol).
- Dammara-24-ene-3 $\alpha$ ,12 $\beta$ ,20 $\beta$ -triol (see betulafolienetriol).
- (20 $\alpha$ H)-Dammara-13(17)-en-3 $\beta$ -ol (see isotrucallanol).
- (20 $\beta$ H)-Dammara-13(17)-en-3 $\beta$ -ol (see isoeuphenol).
- Dammarenolic acid, and derivatives, **166**.
  - biogenesis of, 125.
  - correlation with dipterocarpol, 41.
  - natural occurrence, **166**.
  - structure of ring A, 29.
- use of mass spectrum in structural determination, 105.
- Dammara-25-en-3-one, 12 $\beta$ ,20 $\beta$ (?)-dihydroxy derivative (see alnusfolienediolone).
- Dehydroabiatic acid, correlation with  $\beta$ -onocerin, 53-54.
- Dehydrobromination, of 2-bromo-3-ketones, 35.
- 23-Dehydro-3 $\beta$ ,25-dihydroxycycloartane (see cycloart-23-ene-3 $\beta$ ,25-diol).
- 25-Dehydro-3 $\beta$ ,24 $\xi$ -dihydroxycycloartane (see cycloart-25-ene-3 $\beta$ ,24 $\xi$ -diol).
- Dehydroeburonic acid, physical properties and structure of, **148**.
- $\Delta^{7,9(11)}$ -Dehydroeburicoic acid, preparation of, 98.
- Dehydrogenation, of tetracyclic triterpenes, 102-104.
- Deoxylimonin, ring B cleavage of, 56.
- Desmosterol (see cholesterol, 24-dehydro derivative).
- Dictamnolactone (see limonin).
- Diels hydrocarbon, 104.
- Dienes, as contaminants, 22, 57.
  - preparation of  $\Delta^{7,9(11)}$ , 57-59, 60.
  - reactions of  $\Delta^{7,9(11)}$ , 64.
  - ultraviolet absorption of, 57, 59.
- Dihydroagnosterol, and derivatives, 139.
  - natural occurrence, **139-140**.
- Dihydroelemolic acid, 86.
- Dihydrolanosterol (see lanostenol).
- Dihydrotirucallol, 90, 91, 92.
- 3 $\beta$ ,16 $\alpha$ -Dihydroxyeburico-8,24(28)-dien-21-oic acid (see tumulosic acid).
- 3 $\alpha$ ,12 $\alpha$ -Dihydroxy-24-methylenelanost-8-en-26-oic acid (see polyporenic acid A).
- 3 $\beta$ ,16 $\alpha$ -Dihydroxy-24-methylenelanost-8-en-21-oic acid (see tumulosic acid).
- 2,3-Diketones, preparation of, 38.
- 4,4-Dimethylcholesta-8,24-dien-3 $\beta$ -ol (see 4,4-dimethyleolestadienol).
- 4,4-Dimethylcholestadienol, and derivatives, **136**.
  - natural occurrence, **136**.
- D-(—)-Dimethylheptanoic acid, methyl ester, obtained as degradation product, 78.
- 8,14-Dimethyl-18-nor-testosterone, biological activity, 46, 130.
  - preparation of, 46.
- 1,8-Dimethylphcnanthrene, as dehydrogenation product, 104.

Dipterocarpaceae, as triterpene source, **161, 162, 166.**

Dipteroearpol, conversion to 8,14-dimethyl-18-nor-testosterone, 46.

— correlation with dammadienolic acid, 41.

— degradation of side-chain, 85-86.

— isolation of, 21.

— natural occurrence, 128, **162.**

— physical properties and structure of, **162.**

*Dipterocarpus* spp., triterpene source, 21, 128.

Diterpenes, mevalonic acid in biogenesis of, 117.

## E

Eburic-25-en-3 $\beta$ -ol, 9,19-cyclo derivative (see cyclolaudenol).

Eburicadienonic acid, physical properties and structure of, **147.**

Eburicoic acid, 98.

— and derivatives, **147.**

— as source for 14-methyl-pregnanes, 130.

— biogenesis of, 125.

— C-3 acetate, natural occurrence, 21.

— — position of carboxyl group in, 97.

— dehydro derivative, natural occurrence, **148.**

— dehydro derivatives, **148.**

— natural occurrence, 21, 147.

— rearrangement of side-chain double bond in, 96.

— structure in side-chain, 92.

Eburiconic acid (see Eburicadienonic acid).

Ecballic acid, methyl ester, oxidation of, 88, 89.

*Ecballium elaterium*, as triterpene source, **175, 176.**

*Echinocystis fabeacea*, as triterpene source, **175.**

Elatericin A (see cucurbitacin D).

Elatericin B (see cucurbitacin I).

Elaterin (see cucurbitacin E).

$\alpha$ -Elaterin (see cucurbitacin E).

Elaterium, as triterpene source, **175, 176.**

Elecampane root oil, as triterpene source, **161.**

Elemadienolic acid (see  $\alpha$ -elemolic acid).

Elemadienolic acids, dehydrogenation of, 102.

Elemadienonic acid (see  $\beta$ -elemonic acid).

Elemi acids, determination of side-chain structures, 86.

— oxidation in rings B and C, 59.

— position of carboxyl group in, 97.

— structure of side-chain, 90.

Elemic acid (see  $\beta$ -elemonic acid).

Elemolic acid (see  $\alpha$ -elemolic acid).

$\alpha$ -Elemolic acid, 28.

— and derivatives, **157.**

— C-3 acetate, side-chain structure, 97.

— natural occurrence, **157.**

— reduction of, 68.

— side-chain structure in, 82, 86-87.

$\beta$ -Elemonic acid, methyl ester, physical properties and structures of, **158.**

— natural occurrence, **158.**

— reduction of methyl ester, 33.

— side-chain structure in, 86-87, 90-91.

$\beta$ -Elimination, in cyclohexyl derivatives, 26.

— of C-3 hydroxyl groups, 36.

*Entandrophragma angolense*, as triterpene source, **168.**

2-Epicucurbitacin B, natural occurrence, physical properties and structure of, **174-175.**

Epielemenol (see tirucallenol).

Epilimonol, structure and physical properties, **171.**

Ergostane, side-chain correlation, 94.

Ergosterol, biogenesis of, 124.

*Erythrophloeum guineense*, as triterpene source, **151.**

*Eucalyptus microcorys*, as triterpene source, **151.**

Euphadienol (see euphol).

Eupha-7,24-dien-3-ol (see butyrospermol).

Eupha-8,24-dien-3-ol (see euphol).

Euphane, ring A distortion in, 28.

— structure of ring D, 74.

— use as a basic name, 20.

Euphanes, biogenesis of, 122-123.

— bond isomerizations in, 60-62.

— conversion to dammaranes, 50.

— oxidation in rings B and C, 59.

— physical properties and structures of, **153 ff.**

- rearrangement of double bond in, 47.
  - side-chain in, 81-83.
  - Euphene, dehydrogenation of, 104.
  - Euphenol, and derivatives, 154.
    - conformational strain in, 50.
    - dehydrogenation of, 102.
    - preparation of, 48.
    - rearrangement of, 49-50, 104.
  - $\Delta^8$ -Euphenol-7-d, 48.
  - Euphen-8-en-3-one, reduction of, 33.
  - Euphol, 17, 22.
    - and derivatives, 153-154.
    - biogenesis of, 122.
    - conformational analysis in structure proof, 27.
    - $\Delta^8$  derivative, 122.
    - dihydro derivative (see euphenol).
    - natural occurrence, 128, 154.
    - side-chain structure in, 81-84, 90, 91, 100.
    - stereochemistry of, 68, 78.
    - structure determination of, 50.
  - Euphorbadienol (see euphorbol).
  - Euphorbia* spp., as triterpene source, 21, 128, 137, 145, 154, 156, 160.
  - Euphorbol, and derivatives, 160.
    - natural occurrence, 128, 160.
    - side-chain structure of, 92.
  - $\alpha$ -Euphorbol (see euphorbol).
  - Evodia rutaecarpa*, as triterpene source, 171.
  - Evodine (see limonin).
  - Evodol, natural occurrence, 171.
- F
- Fabacein, natural occurrence, 175.
    - physical properties, 176.
  - Farnesol, as sesquiterpene precursor, 116.
    - pyrophosphate derivative, in squalene biogenesis, 119.
  - Ferulic acid, methyl ester (see oryzanol A).
  - Flindevsia* spp., as triterpene source, 160.
  - Flindissol, natural occurrence, 160.
    - physical properties and structure of, 159.
    - side-chain in, 82.
  - Flindissone, physical properties and structure of, 159.
  - 2-Fluoromevalonic acid, in squalene biogenesis, 120.
- Fomes* spp., as triterpene source, 128, 141, 142, 147, 148.
  - Formate, role in triterpene biogenesis, 124-125.
  - Friedelin, biogenesis of, 124.
  - Fnmigacin, as triterpene source, 178.
  - Fusidic acid, 74, 100.
    - absence of C-4 methyl in, 30.
    - and derivatives, 179-180.
    - degradation of side-chain, 80.
    - dehydrogenation of, 104.
    - 24,25-dihydro methyl ester, side-chain degradation of, 101-102.
    - isolation of, 21.
    - natural occurrence, 21, 180.
  - Fusidium, physiological properties of, 130.
  - Fusidium coccineum*, as triterpene source, 180.
- G
- Gedunin, 23.
    - and derivatives, 168.
    - dihydro derivative, natural occurrence of, 168.
    - natural occurrence, 168.
  - Gedunohor wood, as triterpene source, 168.
  - Geraniol, as monoterpene precursor, 116.
    - pyrophosphate derivative, in squalene biogenesis, 119.
  - Geranyl-geraniol, as diterpene precursor, 116.
  - Geranyl-linalool, as diterpene precursor, 116.
  - Gliotoxin, natural occurrence, 178.
  - Grapefruit, as triterpene source, 136.
  - Guarea thomsonii*, as triterpene source, 168.
  - Gum mastic, as triterpene source, 156, 158, 159.
  - Gurjun, as triterpene source, 21.
- H
- Haudianol (see cycloartenol).
  - Helvolic acid, absence of C-4 methyl in, 30.
    - and derivatives, 178.
    - isolation of, 21.
    - natural occurrence, 21, 178.
    - physiological properties of, 130.
  - Hopea odorata* resin, as triterpene source, 162.

Hydrolysis, accompanied by other reactions in cucurbitacins, 87-88.  
 — of C-3 acetates as structure proof of onocerins, 33.  
 — of ester groups, effects of more distant substituents on, 36.  
 Hydroxyelemadienic acid (see  $\alpha$ -elemolic acid).  
 $\alpha$ -Hydroxyisobutyraldehyde, as degradation product, 87.  
*trans*-Hydroxyisohexenoic acid, as degradation product, 87-88.  
 3 $\beta$ -Hydroxylanosta-8,24-dien-21-oic acid (see trametenolic acid B).  
 21-Hydroxylanosta-8,24-dien-3-one, and derivatives, 140.  
 — natural occurrence, 140.  
 3 $\beta$ -Hydroxy-24-methylenelanosta-7,9(11)-dien-21-oic acid (see eburicoic acid, dehydro derivative).  
 3 $\beta$ -Hydroxy-24-methylenelanost-8-en-21-oic acid (see eburicoic acid).  
 2-Hydroxy-6-methylheptanoic acid, methyl ester as side-chain degradation product, 101.

## I

Ibamarin (see cucurbitacin E).  
*Iberis amara*, as triterpene source, 176.  
 Infrared absorption, of ring A bromo-ketones, 34.  
 — of tetracyclic triterpenes, 181 ff.  
 — use in structure elucidation, 76, 79, 106.  
 Iutodiol, and derivatives, 177.  
 — natural occurrence, 177.  
*Inonotus* sp., as triterpene source, 128, 137, 141, 177.  
*Inula helenium*, as triterpene source, 105, 161.  
 Isocholesterol (see lanosterol).  
 20-Isocholesterol, 81.  
 "Isocholesterol" fraction, as triterpene source, 138, 139.  
 Isoeuphene, dehydrogenation of, 104.  
 Isophenol, and derivatives, 74, 96, 163.  
 — correlation with butyrospermol, 70.  
 — degradation of, 76-78.  
 — oxidation of derivatives of, 76-78.  
 — preparation of, 49-50, 70.  
 — steric effects in, 50.  
 20-Isoeuphol (see tirucallol).

Isohexanoic acid, as degradation product, 87.  
 Isoinotodiol, physical properties and structure of, 177.  
 Isolanostenol, 49.  
 — acetate, preparation of, 47.  
 Isolanosterol, 48, 94-95, 123.  
 Isomasticadienonic acid, and derivatives, 159.  
 — natural occurrence, 159.  
 Iso-obacunoic acid, preparation of as methyl ester, 43.  
 Isopentenol, 116.  
 — role in terpene biogenesis, 117-119.  
 Isopentenyl pyrophosphate, synthesis and role in terpene biogenesis, 116, 118-119.  
 Isoprene rule, 117.  
 Isopropylidene derivatives, degradation of, 38.  
 Isotirucallenol, and 3 $\beta$ -acetate, 163.  
 — degradation of, 76-78.  
 — derivatives of, 96.  
 — — oxidation of, 76-78.  
 — preparation of, 51.

## K

Kan-Zui, 156.  
 Kanzuiol (see tirucallol).  
 11-Ketolanosterol, as intermediate in lanosterol synthesis, 109.  
 — degradation of ring A, 45.  
 11-Keto-14-methylprogesterone, biological activity of, 46, 130.  
 — preparation of, 46.  
*Khaya* spp., as triterpene source, 169.  
 Khivorin, and 7-deacetoxy-7-keto derivative, 169.  
 — natural occurrence, 169.

## L

$\delta$ -Lactones, conversion to cyclopentenones, 97.  
*Lagenaria leucantha*, as triterpene source, 175.  
 Lagos wood, as triterpene source, 169.  
 Lanosta-8,24-diene-3 $\xi$ ,(?)-diol (see inotodiol).  
 Lanosta-7,9(11)-dien-21-oic acid, 16 $\alpha$ -hydroxy-3-oxo-24-methylene derivative (see polyoprenic acid C).

- Lanosta-8,24-dien-21-oic acid, 3 $\alpha$ -acetoxy, methyl ester, natural occurrence, **144**.  
 — 3 $\alpha$ -hydroxy and other derivatives, **143**.  
 — 3 $\alpha$ -hydroxy derivative, methyl ester, natural occurrence, **141**.  
 — 3 $\beta$ -hydroxy and other derivatives, **141** (see also trametenolic acid B).  
 — 3-oxo derivative (see pinicollic acid A).  
 (13 $\alpha$ ,14 $\beta$ ,17 $\beta$ H,20 $\alpha$ H)-Lanosta-7,24-dien-26-oic acid, 3-oxo derivative (see masticadienonic acid).  
 (13 $\alpha$ ,14 $\beta$ ,17 $\beta$ H,20 $\alpha$ H)-Lanosta-8,24-dien-21-oic acid, 3 $\alpha$ -hydroxy derivative (see  $\alpha$ -elemolic acid).  
 — 3-oxo derivative (see  $\beta$ -elemonic acid).  
 (13 $\alpha$ ,14 $\beta$ ,17 $\beta$ H,20 $\alpha$ H)-Lanosta-8,24-dien-26-oic acid, 3-oxo derivative (see isomasticadienonic acid).  
 Lanostadienol (see lanosterol).  
 Lanosta-5,7-dien-3 $\beta$ -ol, **130**.  
 — preparation of, **69**.  
 Lanosta-8,24-dien-3 $\beta$ -ol (see lanosterol).  
 Lanosta-9(11),24-dien-3 $\beta$ -ol (see parkeol).  
 (13 $\alpha$ ,14 $\beta$ ,17 $\beta$ H)-Lanosta-7,24-dien-3 $\beta$ -ol (see butyrospermol).  
 (13 $\alpha$ ,14 $\beta$ ,17 $\beta$ H)-Lanosta-8,24-dien-3 $\beta$ -ol (see euphol).  
 (13 $\alpha$ ,14 $\beta$ ,17 $\beta$ H,20 $\alpha$ H)-Lanosta-8,24-dien-3 $\beta$ -ol (see tirucallol).  
 Lanosta-8,24-dien-3-one, 21-hydroxy derivative (see 21-hydroxylanosta-8,24-dien-3-one).  
 — reduction of, **33**.  
 Lanostane, cucurbitacin skeleton different from, **44**.  
 — 3 $\beta$ -hydroxy-24-methylene-9,19-cyclo derivative (see cycloartanol, 24-methylene derivative).  
 — preparation of, **62**.  
 — structure of ring D, **74**.  
 — use as basic name, **19,20**.  
 Lanostane-3,11-dione, Clemmensen reduction of, **62**.  
 Lanostanes, biogenesis of, **122-123**.  
 — bond isomerizations in, **47, 60-62**.  
 — lack of euphane-type rearrangement in, **51**.  
 — natural occurrence, **128, 134 ff**.  
 — oxidation in rings B and C, **59**.  
 — physical properties and structures of, **134 ff**.  
 — reactions of diene derivatives, **64**.  
 — side-chain degradation, **78**.  
 — side-chain in, **81-83**.  
 — unreactivity of C-11 keto groups in, **45**.  
 Lanostanol (see lanostan-3 $\beta$ -ol).  
 Lanostan-3 $\beta$ -ol, 9,19-cyclo derivative (see cycloartanol).  
 — preparation of, **62**.  
 Lanostanone, bromination of, **34**.  
 Lanosta-7,9(11),24-triene-3 $\beta$ ,21-diol, and derivatives, **140-141**.  
 — natural occurrence, **141**.  
 Lanosta-7,9(11),24-trien-3 $\beta$ -ol (see agnosterol).  
 Lanosta-7,9(11),24-trien-3-one, 21-hydroxy derivative, **140**.  
 Lanostene, **66, 92**.  
 Lanost-8-en-21-oic acid, 3 $\beta$ -hydroxy-24-methylene derivative (see eburicoic acid).  
 Lanostenol, acetate, conversion to isolanostenyl acetate, **47**.  
 — — in synthesis of lanosterol, **109**.  
 — — oxidation of, **60**.  
 — — side-chain oxidation of, **100**.  
 — and derivatives, **137-138**.  
 — as intermediate in cholesterol biosynthesis, **126**.  
 — isomer of, **94**.  
 — natural occurrence, **137-138**.  
 — total synthesis, **107-109**.  
 Lanost-7-en-3 $\beta$ -ol, oxidation of, **69**.  
 — preparation of, **109**.  
 Lanost-8-en-3 $\beta$ -ol (see lanostenol).  
 Lanost-9(11)-en-3 $\beta$ -ol, preparation of, **62**.  
 Lanost-9(11)-en-3 $\beta$ -ol-19-d, acetate, **53**.  
 Lanost-24-en-3 $\beta$ -ol, 9,19-cyclo derivative (see cycloartenol).  
 Lanost-25-en-3 $\beta$ -ol, 24 $\beta$ -methyl-9,19-cyclo derivative (see cyclolaudenol).  
 (13 $\alpha$ ,14 $\beta$ ,17 $\beta$ H)-Lanost-8-en-3 $\beta$ -ol (see euphenol).  
 (13 $\alpha$ ,14 $\beta$ ,17 $\beta$ H,20 $\alpha$ H)-Lanost-8-en-3 $\beta$ -ol (see tirucallenol).  
 (13 $\alpha$ ,14 $\beta$ ,17 $\beta$ H,20 $\alpha$ H)-Lanost-8-en-3 $\beta$ -ol, 24-methylene derivative (see euphorbol).  
 Lanost-8-en-3-one, bromination of, **34**.  
 — reduction of derivatives of, **33**.



- Lanost-9(11)-en-12-one-19-d, 3 $\beta$ -acetate, 53.
- (5 $\beta$ ,8 $\alpha$ ,9 $\beta$ ,10 $\alpha$ ,13 $\alpha$ ,14 $\beta$ ,20 $\xi$ H)-Lanost-24-en-3-one, 4 $\beta$ ,5 $\beta$ ,9 $\beta$ -trimethyl-19-nor derivative (see shionone).
- Lanost-8-en-3,7,11-trione, 36.
- Lanosterol, 15, 17, 18, 22, 98, 122.  
— and derivatives, determination of side-chain structure, 100-101.  
— physical properties and structures of, 187.  
— biogenetic relationship with cholesterol, 120, 126-129.  
— C-20 epimer of, 123.  
— conformational analysis in structure proof, 27.  
— conversion to 14-methylcholestan-3 $\beta$ -ol, 46.  
— correlation with other terpenes, 67.  
— dehydrogenation of, 102.  
— 24,25-dihydro derivative (see lanostenol).  
— isolation, 22.  
— natural occurrence, 21, 22, 128, 137.  
— 11-oxo derivative (see 11-ketolanosterol).  
— physiological properties of, 129.  
— rearrangement of side-chain double bond in, 94-95.  
— side-chain correlation of, 91, 94.  
— side-chain in, 81-86.  
— stereochemistry of, 68.  
— structure determination of, 28, 64-65.  
— total synthesis, 41, 107-109.  
— x-ray analysis of, 28.
- Lanolin, 129.
- Lentinus dactyloides*, as triterpene source, 147, 148.
- Limoclastic acid, preparation of, 43.
- Limonilic acid, 54.
- Limonin, 23.  
— and derivatives, 74, 75, 170.  
— autoxidation of, 55.  
— natural occurrence, 171.  
— reactions of ring A, 42, 43.  
— reactions of rings B and C, 54-56.  
— reactions of ring D, 43, 75-76.  
— tetrahydro derivative, 54.  
— degradation of, 55-56.
- Limon diosphenol, natural occurrence, 171.
- Limoninic acid, hexahydro derivative, 54-55.  
— rearrangement of methyl group in, 76.
- Limonins, biogenesis of, 125-126.  
— natural occurrence, 128, 168 ff.  
— physical properties and structures of, 168 ff.  
— side-chain in, 82.  
— structures of ring A, 29.
- Limonol, degradation of ring D, 76.  
— structure and physical properties, 171.
- Linaloöl, as monoterpene precursor, 116.
- Liver, as source for lanostanes, 128.
- Lophenol, 21, 39.  
— absence of C-4 methyl in, 30.  
— and derivatives, 134-135.  
— as intermediate in cholesterol biosynthesis, 127.  
— correlation with steroids, 40.  
— determination of side-chain structure, 101.  
— mass spectrum of, 105.  
— natural occurrence, 21, 134-135.  
— rearrangement of double bond in, 47.
- Lophenone, physical properties and structure of, 185.  
— reduction of C-3 keto group, 32.  
— steric hindrance of C-3 keto group, 40.
- Lophocereus schottii*, as source for triterpenes, 135.
- Luffa echinata*, as triterpene source, 175.
- Lycopodium clavatum*, as triterpene source, 23, 167.

## M

- Maclura pomifera*, as triterpene source, 155.
- MacDougallin, and derivatives, 134.  
— natural occurrence, 134.
- Mahogany wood, as triterpene source, 151.
- Manila elemi* resin, as triterpene source, 21, 157, 158.
- Manoöl, correlation with lanosterol, 67.
- Mass spectroscopy, use in structure elucidation, 104-105.
- Masticadienonic acid, and derivatives, 158.

- natural occurrence, 158.
  - side-chain correlation of, 91.
  - side-chain in, 82.
- Mechanism of, biogenetic cyclization of squalene, 121-125.
- chromic acid oxidation of cyclohexanols, 26.
  - cyclopropane ring opening, 51.
  - double bond migration in Wolff-Kishner reaction, 60-62.
  - ester hydrolysis in cyclohexyl derivatives, 26.
  - halogenation of cyclohexenes, 27.
  - lactone formation in polyprenate side-chains, 99-100.
  - limonin-limonilic acid conversion, 55.
  - Meerwein-Verley-Ponndorf reduction, 32.
  - methyl group migration in euphanes, 50.
  - one-carbon methylation in triterpene biogenesis, 125.
  - reduction of C-3 keto groups, 31, 32.
- Meliaceae, as triterpene source, 128, 168.
- Merolimonol, 76.
- Methionine, role in triterpene biogenesis, 125.
- Methostenol (see lophenol).
- 6-Methoxy-1-methyltetralone-2 as starting material for total synthesis of onocerin, 109.
- $\alpha$ -Methylacrylic aldehyde, as degradation product, 87.
- Methylation, in lanosterol and onocerin total synthesis, 107-109.
- of cholesta-4,7-dien-3-one, 109.
  - of citrostadienol, 40.
  - of cycloecalenone, 41.
  - of  $\Delta^7$ -3-keto steroids, 40.
  - of lophenol, 40.
- 14-Methylcholestan-3 $\beta$ -ol, preparation of, 46.
- Methylcholest-7-en-3-ol (see lophenol).
- Methylcholest-8-en-3-ol, and derivatives, 135.
- natural occurrence, 135.
- Methyl derivatives (see also parent compounds).
- 24-Methylenecycloartanol, occurrence as 3-ferulate, 21.
- side-chain correlation of, 92.
- 24-Methylenecycloartenol, side-chain in, 83.
- 5-Methylhexanoic acid, as a side-chain degradation product, 101.
- Methyl isohexyl ketone, as side-chain degradation product, 100.
- Methyl isopropyl ketone, as side-chain degradation product, 96.
- Methyl migration (see rearrangement).
- 4-Methyl steroids, 39 ff.
- Mevaldic acid, in terpene biogenesis, 117.
- Mevalonic acid, and lactone, in terpene biogenesis, 117-120.
- synthesis of (D,L)-phospho derivative, 119.
- Meystre-Miescher method for side-chain degradation, 84, 85, 90.
- Molecular rotation differences, in side-chain correlations, 94.
- Monoterpenes, mevalonic acid in biogenesis of, 117.
- Myrtaceae, as triterpene source, 152.

## N

- Natural rubber, 117.
- mevalonic acid in biogenesis of, 118.
- Nerolidol, as sesquiterpene precursor, 116.
- Nomilin, natural occurrence, physical properties and structure, 171.
- reactions of ring A, 42.
  - reactions of ring D, 75.
- A-Nor-ketones, of cucurbitacins, 44, 72.
- preparation of, 38, 39, 45.
- A-Nor-diketones, preparation of, 39.
- A-Nor- $\alpha$ -hydroxy acids, preparation of, 38.
- A-Nor-3-isopropyl-3(5)-enes, preparation of, 37.
- A-Nor-3-isopropylidene derivatives, preparation of, 37.
- A-Nor-3-isopropyl-5(6)-lanostenes, preparation of, 37.
- A-Norpregnanes, 14-methyl derivatives, 130.
- B-Nor-A-homo structures, preparation of, 67-68.
- B-Nor-C-homo structures, preparation of, 60.
- B-Nor derivative of cedrelone, 56, 76.
- Nuclear magnetic resonance, of cucurbitacin B, 44.

- of cyclopropyl groups, 53.
  - use in structure elucidation, 106.
- Nyctanthic acid, biogenesis of, 125.

## O

- Obaculactone (see limonin).
- Obacunonic acid, methyl ester, 43.
- preparation of, 42.
- $\alpha$ -Obacunol, structure and physical properties, 170.
- $\beta$ -Obacunol, structure and physical properties, 170.
- Obacunone, and derivatives, 170.
- natural occurrence, 170.
  - preparation from nomilin, 42.
  - reactions of ring A, 42.
  - reactions of ring D, 75.
- Obliquol, 177.
- Obtusifoliosin, correlation of side-chain, 94.
- Oleanolic acid, correlation with lanosterol, 67.
- $\beta$ -Onoceradiene, total synthesis of, 112-113.
- $\alpha$ -Onoceradienediol (see  $\alpha$ -onocerin).
- $\beta$ -Onoceradienediol (see  $\beta$ -onocerin).
- Onocera-8,13-diene-3 $\beta$ ,21 $\alpha$ -diol (see  $\beta$ -onocerin).
- Onocera-8(26),14(27)-diene-3 $\beta$ ,21 $\alpha$ -diol (see  $\alpha$ -onocerin).
- Onocerane, bisnordione derivative, synthesis of, 111.
- dinor derivative, synthesis of, 112.
  - mass spectrum of, 105.
  - total synthesis of ring system, 109-113.
  - use as basic name, 20.
- Onoceranones, biogenesis of, 124.
- dehydrogenation of, 103.
  - physical properties and structures of, 167 ff.
- Onocerin (see  $\alpha$ -onocerin).
- $\alpha$ -Onocerin, 18, 124.
- and derivatives, 111, 167.
  - equilibration with  $\beta$ -onocerin, 47-48.
  - natural occurrence, 167.
  - oxidation of, 53.
  - total synthesis of, 107, 109.
- $\beta$ -Onocerin, 18.
- and derivatives, 168.
  - correlation with dehydroabiatic acid, 53-54.
  - natural occurrence, 168.
  - oxidation of derivatives of, 53.
  - stereochemistry, 54.
- $\gamma$ -Onocerin, preparation of, 48.
- Onocerin, rearrangement of double bonds in, 47.
- Onocol (see  $\alpha$ -onocerin).
- Ononis spinosa*, as triterpene source, 23, 167, 168.
- Opinin, as triterpene source, 21, 151.
- Optical rotatory dispersion, absence of a Cotton effect in 18-nor-20-keto compounds, 97.
- effect of  $\Delta^7$  and  $\Delta^8$  unsaturation in euphanes and lanostanes, 49.
  - in cucurbitacin structure determination, 74.
  - in dammarane structure determination, 79.
  - in determining ring A conformations, 35.
  - in measuring hindrance of C-3 keto groups, 40.
  - in structure elucidation, 106, 181 ff.
- Oranges, as triterpene source, 136, 155.
- $\gamma$ -Orysterol (see cycloartanol, 24-methylene derivative).
- Oryzanol A, natural occurrence, 21, 145.
- side-chain, 125.
- Oryzanol C, natural occurrence, 21, 152.
- physical properties and structure of, 152.
  - side-chain, 125.
- Oxidation (see also under individual compounds).
- biogenetic of C-30 or C-31, 30.
  - of butyrospermol, 50.
  - of cucurbitacin rings B and C, 71-73.
  - of  $\Delta^7$ ,  $\Delta^8$  and  $\Delta^{7,9(11)}$  derivatives, 56-60.
  - of dihydrobutyrospermol and acetate, 69-70, 125.
  - of 8(9)-double bond with ruthenium tetroxide, 60.
  - of euphane derivatives, 62.
  - of fusidic acid side-chain, 80.
  - of isoeuphenol derivatives, 76-78.
  - of isotirucallenol derivatives, 76-78.
  - of lanostane derivatives, 68, 69.
  - of limonin derivatives, 55-56.
  - of methyl ecballate, 88-89.
  - of polypropenic acid A, 71.

- of side-chains, 83-89.
- of tetrahydroelaterin methyl ether, 89.
- positions of, in tetracyclic triterpenes, 125.
- rearrangement of double bond during, 50.
- Oxoelmadienic acid (see  $\beta$ -elemenic acid).
- 3-Oxolanosta-8,24-dien-21-oic acid (see pinicolic acid A).

## P

- Pachyma hoelen*, as triterpene source, 149.
- Pachymic acid, 21.
  - natural occurrence, 149.
  - physical properties and structure, 149.
- Panax* spp., as triterpene source, 166.
- Panaxadiol, and derivatives, 165-166.
  - natural occurrence, 166.
  - side-chain in, 82.
- Parkeol, and derivatives, 63, 64, 138.
  - dihydro derivative, structure of, 63.
  - natural occurrence, 138.
- Peniocereus macdougalii*, as source for triterpenes, 134.
- Peniocerol, natural occurrence, 134.
- Pentacyclic triterpenes, dehydrogenation of, 103.
- Phellodendron* spp., as triterpene source, 170, 171.
- Photochemical reactions, cleavage of ring A, 41.
  - of 4,4-dimethyl-10-isosteroids, 47.
  - of lanostane derivatives, 67-68.
- Pinicolic acid A, and derivatives, 143.
  - side-chain correlations of, 91-92.
  - natural occurrence, 143.
- $\alpha$ -Pinicolic Acid (see pinicolic acid A).
- Pinicolic acids, natural occurrence, 128, 143.
  - side-chain in, 82.
- Pistacia lentiscus* resin, as triterpene source, 158.
- Plants, as source for lanostanes, 128.
- Polyporenic acid A, and derivatives, 148.
  - C-3  $\alpha$ -hydroxyl group in, 28.
  - dehydrogenation of, 102.
  - natural occurrence, 128, 149.
  - oxidation of, 71.
  - side-chain correlation of, 94.

- Polyporenic acid B, 3-acetate, 21.
- Polyporenic acid C, 3-acetoxy-16-keto derivative, 98.
  - and derivatives, 150.
  - configuration of C-16 hydroxyl group, 100.
  - dihydro derivative methylester, 98.
  - methyl ester, rearrangement of side-chain double bond, 96.
  - methyl ester of 16-keto derivative, reduction of, 100.
  - natural occurrence, 128, 150.
  - position of carboxyl group in, 97-100.
- Polyporus* spp., as triterpene source, 21, 128, 140, 141, 142, 143, 144, 147, 148, 149, 150.
- Polyprenoids, definition of, 115.
- Poria cocos*, as triterpene source, 147, 148, 149.
- 5 $\alpha$ -Pregnane, 3 $\beta$ ,20 $\alpha$ -di-(methylamino)-16 $\alpha$ -hydroxy-4-methylene-14 $\alpha$ -methyl-9 $\beta$ ,19-cyclo derivative (see cyclobuxine).
- Pregnanes, 14-methyl and 4,4,14-trimethyl derivatives from eburicoic acid, 130.
- Provitamin D<sub>3</sub> (see cholesterol, 7-dehydro derivative).
- Pyrolysis, in structure determination, 76, 101.
  - of, 2,3-seco diacids.

## R

- Radioactive tracer techniques, in biogenesis studies, 116 ff.
- Rearrangement, benzilic acid, of rings A or B, 38, 56.
  - in cucurbitacins and derivatives, 73, 87-89.
  - in oxidation of dihydrobutyrospermyl acetate, 125.
  - of acyl group in fusidic acid, 80.
  - of double bond, and methyl groups in euphanes, 49-51.
  - — in biogenesis of terpene precursors, 119.
  - — in lanostane derivatives, 65.
  - — in onocerins, 47.
  - — in side-chain degradations, 87-88, 94-97.
  - — in steroid biogenesis, 127.

- — in Wolff-Kishner reduction, 60-62.
  - of double bonds in lanostenes, 109.
  - — in rings B and C, 47 ff.
  - of lanostane skeleton, 67-68.
  - of methyl group, in cedrelone, 56, 76.
  - — during oxidation, 70, 76.
  - — in dehydrogenation, 102-103.
  - — in hexahydrolimoninic acid, 76.
  - of methyl groups in triterpene and steroid biogenesis, 120, 122-124.
  - retropinacol, of ring A, 36-38.
  - Reduction, control at C-17 by C-14 methyl group, 80.
  - of C-3 ketones, 31 ff.
  - of C-11 ketones, 62.
  - of  $\Delta^{7,9(11)}$  dienes, 59.
  - of elemolic acid, 68.
  - of groups in euphane rings B and C, 60-62.
  - of groups in lanostane rings B and C, 60-62.
  - Reverse aldol reaction, in degradation of cucurbitacins, 87-88.
  - Rice-bran oil, as triterpene source, 144, 145, 152.
  - Ring contraction, cedrelone ring B, 56, 76.
  - during dehydration of C-3 hydroxyl groups, 30-31, 37.
  - during solvolysis of 3 $\beta$ -sulphonate esters, 37.
  - in lanostane derivatives, 30-31, 37, 65.
  - of ring A, in cucurbitacins, 39, 44, 88-89.
  - — of 2,3-diketones, 38.
  - Rotation differences, of euphanes isomeric at C-9, 61.
  - of mono- and dienes, 57, 59.
  - Rotatory dispersion (see optical rotatory dispersion).
  - Rutaceae, as triterpene source, 128, 170, 171.
- S
- Schinol, natural occurrence, 158.
  - Schinus molle* resin, as triterpene source, 158.
  - Sclareol, 113.
  - 2,3-Seco diacids, 36.
  - preparation of, 38.
  - pyrolysis of, 38.
  - Sesquiterpenes, mevalonic acid in biogenesis of, 117.
  - Shea nut butter (see *Butyrospermum parkii*).
  - Sheep's wool fat, as triterpene source, 128, 137, 138, 139, 140.
  - Shionone, and derivatives, 177-178.
  - biogenesis of, 123-124.
  - natural occurrence, 178.
  - rearranged methyl group in, 29, 31.
  - structure of ring A, 29.
  - Shorea vulgaris* resin, as triterpene source, 162.
  - Side-chain, position of attachment in lanosterol, 64-65.
  - reactions involving a carboxyl group in, 97.
  - Side-chains, degradation of, 64, 80, 83 ff.
  - interconversions, 89-94.
  - Spanish inoss, as triterpene source, 145, 146.
  - Spinasterol, side-chain biogenesis, 125.
  - Squalene, 15.
  - as cholesterol precursor, 120-121.
  - as triterpene precursor, 116, 120-121.
  - biogenesis of 119-120.
  - mevalonic acid in biogenesis of, 117, 118, 120.
  - Stabilities, of isomeric 2-bromo-3-ketones, 34.
  - Stereochemistry, of A-B ring junction in terpenes, 31.
  - of metal hydride reductions, 31-32.
  - of  $\beta$ -onocerin, 54.
  - of reduction of C-3 keto groups, 31-33.
  - of the tetracyclic triterpenes, 24 ff.
  - requirements for contraction of ring A, 37.
  - Steric effects, and oxidation in rings B and C, 57.
  - in euphane derivatives, 60, 62.
  - in isoeuphenol, 50.
  - in lanostane derivatives, 60, 62.
  - in lophcnone, 40.
  - in polyprenate side-chains, 99-100.
  - in ring D of dammaranes, 79-80.
  - in side-chain reactions, 96-97.
  - in six-membered rings, 25-26.

- in synthesis of ouoceranes, 113.
- of C-4 methyl groups, 40.
- of 8 $\beta$ -methyl group, 34-35.
- Steroids,  $\alpha,\beta$ -rule in naming triterpenes, 19.
- biogenesis of, 126-127.
- Stigmasta-7,24(28)-dien-3 $\beta$ -ol, 4 $\alpha$ -methyl derivative (see citrostadienol).
- Strychnos nux-vomica*, as triterpene source, 145.
- Succinic acid, as a side-chain degradation product, 101.
- Swietenia mahagoni*, as triterpene source, 151.

## T

- Tallow tree, as triterpene source, 151.
- Terebinthone, natural occurrence, 158.
- Tetracyclol triterpenes, antibiotics, 39.
- 80, 100, 101, 130, side-chain in, 82.
- 2-benzylidene derivatives of, 36.
- biochemistry of, 115 ff.
- biogenesis of, 115-125.
- bromination of ring A, 34-35.
- catabolism of, 125-127.
- cyclopropane ring, reactions of, 51.
- dehydration involving C-3 hydroxyl groups, 36.
- dehydrogenations of, 102-104.
- derivatives as steroid hormone analogues, 46, 130.
- 2-furfurylidene derivatives, 36.
- 2-hydroxy methylene derivatives, 36, 38.
- isolation methods, 21.
- naming and numbering, 19.
- occurrence as glucosides, 20.
- physiological activity, 129-130.
- syntheses of, 107-113.
- types, of ring A in, 28.
- — of rings B and C in, 46-47.
- — of ring D in, 74.
- — of side-chains in, 81-83.
- 1,2,5,6-Tetramethylnaphthalene, as dehydrogenation product, 104.
- 1,2,7,8-Tetramethylphenanthrene, as dehydrogenation product, 104.
- Tillandsia usneoides*, as triterpene source, 145, 146.
- side-chain in substances from, 82.
- Tirucalladienol (see tirucallol).
- Tirucallanes, biogenesis of, 122-123.
- Tirucallenol, and derivatives, 156.
- 3 $\alpha$ -hydroxy epimer, preparation of, 68.
- methyl migration in, 50-51.
- Tirucall-7-en-3 $\beta$ -ol (see masticadienonic acid).
- Tirucall-8-en-3 $\beta$ -ol (see isomasticadienonic acid).
- Tirucallol, 17.
- and derivatives, 155-156.
- biogenesis of, 122.
- $\Delta^8$  derivative, 122.
- dihydro derivative (see tirucallenol).
- natural occurrence, 128, 156.
- oxidation derivatives of, 59.
- side-chain in, 81-82, 90, 91.
- stereochemistry of, 68, 78.
- TPNH, in squalene biogenesis, 120.
- in steroid biogenesis, 127.
- Trametenonic acid, structure and physical properties, 142.
- Trametenonic acid A, and derivatives, 142.
- dehydrogenation of, 104.
- natural occurrence, 141-142.
- Trametenonic acid B, 141.
- and derivatives, 141, 142.
- natural occurrence, 142.
- Trametes odorata*, as triterpene source, 128, 141, 142.
- 1,2,5-Trimethyl-6-hydroxynaphthalene, as dehydrogenation product, 104.
- 1,2,5-Trimethylnaphthalene, as dehydrogenation product, 103, 104.
- 1,2,8-Trimethylphenanthrene, as degradation product, 102, 104.
- Tristania conferta*, as triterpene source, 151.
- Triterpenes, mevalonic acid in biogenesis of, 117-118 (see also tetracyclol triterpenes).
- Tumour, as triterpene source, 135, 138.
- Tumulosic acid, and derivatives, 74, 149.
- natural occurrence, 149.
- Tun wood, as triterpene source, 172.
- Twist form, in cyclohexane derivatives, 24.

## U

- Ultraviolet absorption, effect of conjugated cyclopropane ring on, 35.
- in lanosterol structure determination, 65.

- of  $\Delta^{7,9(11)}$  dienes, 57, 59.
  - of isolated double bonds, 105.
  - of tetracyclic triterpenes, **181 ff.**
  - Ungulina betulina*, as triterpene source, **149.**
  - Ungulinic acid (see polyporenic acid A).
- W
- Wolff-Kishner reduction, C-20-epimerization during, 98.
  - in side-chain interconversion, 90, 92, 98.
- X
- X-ray analysis, in lanosterol structure proof, 27.
- Y
- Yeast, as triterpene source, **137.**
- Z
- Zymosterol, side-chain in, 81.