

Index

A

- Acetonitrile covalent-adduct chemical ionization (CACI) MS-MS
 - UFA methyl ester isomers
 - acetonitrile CACI for FAME analysis, 162–170
 - characterization of UFA double bond position, 158–162
 - covalent adduct chemical ionization (CACI), 162
- Acetyl number, of waxes, 423–424
- Acid value of waxes, 422–423
- Actinomadura* sp., 97
- Acylcarnitine
 - EI-MS and, 55
 - shotgun lipodomics and, 65
- Acyl-CoA
 - EI-MS and, 55
 - shotgun lipodomics and, 65
- Acylglycerols, extraction methods and, 3
- Adipocytes, shotgun lipodomics and, 66–67
- Alkyl glycosides FA esters, 390
 - as fat replacers, 383–384
- α -hydroxy migration, HPLC on silica-based matrices and, 52–53
- Amalean I, 385
- American Type Culture Collection, 273
- Amylodextrin, 386
- Anabolic steroids, APPI and, 42
- Anionic lipids, EI-MS and, 55–56
- Arachidonic acid
 - chiral-phase HPLC enantiomere separation and, 77
 - dual detection and, 217
 - HETE/HODE synthesis and, 75
- Aston, Francis William, 29
- Atherosclerosis, metabolic syndrome and, 51
- Atmospheric pressure chemical ionization (APCI)
 - analyte detection after HPLC using, 10

- development of, 33–35

- GC/MS for double bond position UFAs and, 161

- Atmospheric pressure photoionization (APPI)

- development of, 34–35

- soft ionization via, 38–42

- Attenuated total reflectance (ATR-FTIR) spectroscopy, 305–306

- Automation of lipid analysis, 5

- 2D identification of individual species and, 63

- Avicel, 386

B

- Bacteria. *See also* Food-borne pathogens; specific bacteria

- fast GC for cellular FAME analysis

- bacteria used/preparation, 273–274

- bacterial identification, 277, 280

- chromatographic methods, 274–277

- future research opportunities, 281–282

- GC use, 272–273

- microbial identification by FA analysis, 272

- sample preparation, 277, 281

- β -glucans, 385–386

- Bicyclic endoperoxides, 126

- Bile acids, EI-MS and, 55

- Biopolymer/fiber-based products, 385–386

- Bis(monoacylglycerol)phosphate (BMP),
 - chiral-phase HPLC enantiomeric separation and, 97, 99–100

C

- Calcium sulfate, with adsorbent silica gel G, 6–7

- Calorie Control council National (1998), 379

- Calorie free labeling, 381

- Candida antarctica*, 94
- Capillary exit voltage (CapEx), stereoisomeric lipids and, 80
- Capillary gas chromatography (GC). *See also* Gas chromatography (GC)
- foodborne pathogen identification via ATR-FTIR spectroscopy protocol, 295
 - foodborne pathogen identification with FTIR spectroscopy
 - cultivation of cells, 289–290
 - fatty acid structures in cellular lipids, 288–289
 - GC-FID limitations, 291
 - GC-FID methodology, 290–294
 - general discussion, 287–288
 - methodology, 294–296
 - preparation of cellular FAMES for analysis, 290
 - profiling cellular FAMES by GC, 289
- Cardiolipin
- EI-MS and, 55
 - LC/MS (recent developments) and, 139, 141
 - shotgun lipidomics and, 65
- Catechin, APPI and, 42
- Cavendish Laboratory (University of Cambridge), 29
- Cellular lipidomes. *See* Shotgun lipidomics
- Celluloses, 385–386
- Ceramide phosphorylethanolamines
- EI-MS and, 55
 - FAB-MS and, 36
 - shotgun lipidomics and, 65
- Cerebrosides
- EI-MS and, 55
 - shotgun lipidomics and, 66
- Chemical ionization (CI), 33–35
- Chemluminescent assays, determination of peroxide value (PV) by, 6
- Chemometrics, 337–338
- Chevreul, Michel Eugène, 424–425
- Chiral-phase HPLC applications
- analysis advantages for
 - enantiomers/diastereomers, 73
 - derivitization of enantiomeric hydroxy FA, 75
 - materials/methods
 - derivitization of enantiomeric DAGs/MAGs, 75–76
 - diastereomeric DAG derivatives, 78–79
 - diastereomeric derivative preparation, 76
 - enantiomeric epoxides, 77
 - enantiomeric glycerolipids, 77–78
 - enantiomeric glycerophospholipids, 79
 - enantiomeric hydroxy FA, 76–77
 - LC/ESI-MS and MS/MS of stereoisomeric lipids and, 79–81
- reagents/columns/suppliers, 74–75
- results/discussion
- steric analysis of DAGs, 91–95
 - steric analysis of dihydroxy FA, 88–89
 - steric analysis of epoxy FA, 89
 - steric analysis of glycerophospholipids, 95–101
 - steric analysis of MAGs, 89–91
 - steric analysis of monohydroxy FA, 81–88
- Cholesterol
- dual detection and, 217
 - low density lipoprotein (LDL), 380
 - silica gel G and separation of, 6
- Cholesteryl esters of FAs
- EI-MS and, 55
 - LC/MS (recent developments), 137–149
 - supercritical fluid extraction (SFE) and, 17
- Choline glycerophospholipid (PD)
- EI-MS and, 55
 - shotgun lipidomics and, 65
- Chromarods, 261, 263, 265
- Chromatographic methods of lipid analysis
- counter current chromatography (CCC), 16
 - gas chromatography (GC), 13–16
 - high performance liquid chromatography (HPLC), 8–13
 - size exclusion chromatography (SEC), 16
 - supercritical fluid chromatography

- (SFC), 16–17
thin-layer chromatography (TLC), 6–8
- Classical chemical applications of lipid analysis, 5–6
- Collisionally induced dissociation (CID)
ESI and, 37
MALDI-TOF MS and, 42
stereoisomeric lipids and, 79–80
- Columns
supercritical fluid chromatography (SFC)
packed vs. capillary, 241–244
PAG/sterols/tocopherols/seed oils, 250–254
TAG/FFA/glyceride mixtures, 245–250
for supercritical fluid extraction (SFE), 17
- Concentrated fraction applications
HPSEC
crude/refined oils, 223–224
digestibility coefficients in animals, 226
intact oil samples, 207–208, 218–219
oxidized fats/oils, 224–227
used frying fats/oils, 220–223
- Conjugated C18 trienes, acetonitrile CACI for FAME analysis and, 168, 170
- Conjugated linoleic acids (CLAs)
acetonitrile CACI for FAME analysis and, 168–171
bioactivities of, 158
exclusion from *trans*-FA methods, 304–306
FT-NIR and, 306–330
quantification with IR spectroscopy chemometrics, 337–338
experimental
standards/measurements/data handling, 343–344
general discussion, 335–337
PLS technique of calibration, 338–342
spectra of triglycerides, 344–350
RP-HPLC and, 9
- Corn bran/fiber lipid extracts, 31
- Cornstarch, 385
- Coronaô CADô (charged aerosol detector), 39
- Counter current chromatography (CCC) as chromatographic methods of lipid analysis
general discussion, 16
- Coumaroylferuloylputrescine (CFP), 31
- Covalent-adduct chemical ionization (CACI). *See also* Acetonitrile covalent-adduct chemical ionization (CACI) MS-MS
- GC/MS for double bond position UFAs and
FAME analysis of acetonitrile CACE, 162–166
future applications of conjugated C18 trienes (CLnA), 168, 170
homoallylic/diagnostic ion structures, 163, 167–169
UFA methyl ester isomers and, 162
- Crude/refined oils, HPSEC and, 223–224.
See also Fats/oils
- CUBO solvent, 19
- D**
- Dairy products, extraction methods and, 4
- Databases, GC-MS-EI comparisons and, 33
- Densitometric measurement for
quantification of lipid analytes, 7–8
- Desorption, development of, 34
- Diabetes, metabolic syndrome and, 51
- Diacylglycerol isomer determination, 9
- Diacylglycerol lipid class (DAG), HPLC-MS-APCI and, 39–40
- Diacylglycerols (DAG)
and 2D identification of individual species, 59–60
chiral-phase steric analysis of, 91–96
conversion to NEUs or PEUs, 94–95
diastereomeric derivatives of, 76, 78–79
EI-MS and, 55
shotgun lipodomics and, 65
silica gel G and separation of, 6
size exclusion chromatography (SEC) and, 16
- Diastereomers. *See also* Chiral-phase HPLC

- applications
 - advantages of chiral separation LC/MS and, 73
 - DAG derivatives, 78–79, 94–95
 - Diferuloylputrescine (DFP), 31
 - Dihydroxy FA
 - chiral-phase steric analysis of, 88–89
 - steric analysis of, 88–89
 - Dihydroxyecosatrienoic acid (DiHETrE),
 - steric analysis of, 88–89
 - Dihydroxypropyl propyl ether, as diol phase of HPLC, 8
 - Dimethylloxazoline (DMOX)
 - GC/MS for double bond position UFAs and, 160–161
 - use of derivatives of, 14–15
 - Dimyristoylphosphatidic acid (DMPA), 366–368
 - Dimyristoylphosphatidylcholine (DMPC- d_{54}), 367
 - Diol phase of HPLC, 8
 - Dipalmitoyl (GroPSer), 101
 - diradylglycerol
 - hydroxides/hydroperoxides/isoprostanes/core aldehydes
 - LC/MS (recent developments), 128–131
 - Direct APPI, 41
 - Dopant APPI, 41
 - Dual detection (UV diode-array/refractive), 217
 - Dyes for visualization, thin-layer chromatography (TLC) and, 6–7
- E**
- Electrical properties of analytes
 - EI-MS and, 53–56
 - and quantitation of lipid molecular species, 63–64
 - Electrochemical detection devices
 - analyte detection after HPLC using, 10
 - Electron impact (EI)
 - hard GC-MS-EI/LC-MS-EI and, 30–33
 - steryl esters and, 124
 - Electron multiplier, ion detection and, 46
 - Electrospray ionization mass spectrometry (ESI-MS)
 - 2D identification of lipid molecular species, 59–63
 - 2D quantitation of lipid molecular species, 63–65
 - advantages of, 52
 - biological applications of, 65–67
 - initial phospholipid analysis using, 53
 - intrasource separation of lipids and, 55–58
 - intrasource separation of lipid classes (shotgun lipidomics), 53–59
 - Electrospray ionization mass spectrometry (ESI-MS-MS)
 - analyte detection after HPLC using, 10–12
 - development of, 34–35
 - soft ionization via, 37–38
 - Elution order
 - silver-ion HPLC and temperatures, 179, 181–183
 - trans*-18:1 FA analysis by silver ion HPLC and, 195, 197–199
 - Enantiomers. *See also* Chiral-phase HPLC applications
 - advantages of chiral separation LC/MS and, 73
 - derivitization of DAGs/MAGs, 75–76
 - derivitization of enantiomeric DAGs/MAGs, 75–76
 - derivitization of enantiomeric hydroxy FA, 75
 - epoxide chiral separation, 77
 - glycerolipid chiral separation, 77–78
 - glycerophospholipid chiral separation, 79
 - hydroxy FA chiral separation, 76–77
 - Epoxy FA, chiral-phase steric analysis of, 89
 - Epoxyecosatetraenoic acids (EETs), 77, 89
 - Erythro-DiHETErE, 89
 - ESA Inc., 39
 - ESI-MS-MS. *See* Tandem electrospray ionization (ESI-MS-MS)
 - Ester value of waxes, 423
 - Esterified propoxylated glycerols (EPGs), as fat replacers, 382
 - Ethanolamine glycerophospholipid (PE)

- El-MS and, 55
 shotgun lipodomics and, 65
- Ether, extraction methods and, 3
- Evaporative light scattering detector
 (ELSD), analyte detection after
 HPLC using, 10
- Extraction methods, 3–5
- F**
- FAME applications
 HPSEC, 218
 FAME, 219–221
- Faraday cup, ion detection and, 46
- Fast atom bombardment (FAB)
 development of, 34–35
 mechanism of, 36–37
- FAST methods of GC. *See* Bacteria
- Fat free labeling, 381
- Fat replacers
 chemistry
 alkyl glycosides FA esters, 383–384
 carbohydrate FA esters, 382–383
 EPGs, 382
 PEPs, 382
 protein-based fat substitutes, 386–388
 reduced-calorie fat-based fat
 replacers, 388–389
 SPE-Olestra (Olean), 383
 starch-based fat replacers, 384–386
- food applications, 389–391
- health issues
 general discussion, 379–381
 labeling, 381
 toxicology, 391–392
- Fats/oils
 HPSEC
 crude/refined oils, 223–224
 fats/oils replacers, 214–216
 fish oils, 212–213
 minor lipid compounds, 216–217
 oxidized fats/oils, 224–225
 partial glycerides, 213–214
 pesticide analysis, 217–218
 used frying oils, 211–213
- replacers
 HPSEC and, 214–216
- Fatty acid esters (alkyl glycosides), as fat
 replacers, 383–384
- Fatty acid esters (carbohydrate), as fat
 replacers, 382–383
- Fatty acid methyl esters (FAME)
 acetonitrile CACI for, 162–166
 effects on lipid elution (silver-ion
 HPLC) and, 179–183
 with fast GC for bacteria analysis
 bacteria used/preparation, 273–274
 bacterial identification, 277, 280
 chromatographic methods, 274–277
 future research opportunities,
 281–282
 GC use, 272–273
 microbial identification by FA
 analysis, 272
 sample preparation, 277, 281
- FTIR method for identification of, 15–16
- GC determination of, 5, 13–16
- hard GC-MS-EI/LC-MS-EI and, 30–31
- HPSEC
 applications, 207
 HPSEC and, 218–223
 nuclear magnetic resonance
 spectroscopy (NMR) and, 19
 RP-HPLC and, 9
 silver-ion HPLC and temperatures,
 183–186
 trans-18:1 FA analysis and, 191
 trans-18:1 FA separation as, 192–194
- Fatty acid partially esterified polysaccharide
 (PEP), as fat replacers, 382
- Fatty acids (FAs)
 α -oxidation and, 67
 cellular. *See* Capillary gas
 chromatography (GC)
- EI-MS and, 55
 enantiomeric hydroxy, 75
 extraction methods and, 3
 shotgun lipodomics and, 65
 supercritical fluid extraction (SFE) and,
 17
 trans monoethylenic, 13–14
 UV-VIS detection of, 9
- Ferricytochrome c interaction with DMPA,
 366–368

- Field desorption, 34
- Fish. *See* Meat/fish
- Fish oils, HPSEC and, 212–213. *See also*
Fats/oils
- Flame ionization detection (FID)
analyte detection after HPLC using,
8–10
GC/MS for double bond position UFAs
and, 158, 160
with thin-layer chromatography (TLC)
general discussion, 261–262
principles, 262–263
quantitation/applications, 263–268
- Flame photometric detector (FPD), 265
- Flavonoids, APPI and, 42
- Fluorimetric measurement for quantification
of lipid analytes, 7–8
- Food and Drug Administration (FDA)
GRAS and, 381
- Food-borne pathogens
attenuated total reflectance (ATR)-FTIR
spectroscopy
methodology, 294–296
multivariate statistical analysis of IR
spectra, 296–297
protocol, 295
identification via infrared
spectroscopy/capillary GC
cultivation of cells, 289–290
- fatty acid structures in cellular lipids,
288–289
GC-FID limitations, 291
GC-FID methodology, 290–294
general discussion, 287–288
preparation of cellular FAMES for
analysis, 290
profiling cellular FAMES by GC, 289
- Fourier transform infrared spectroscopy
(FTIR)
coupled with GC, 15–16
determination of peroxide value (PV) by,
6
foodborne pathogen identification with
ATR
methodology, 294–296
multivariate statistical analysis of IR
spectra, 296–297
protocol, 295
foodborne pathogen identification with
capillary GC
cultivation of cells, 289–290
fatty acid structures in cellular lipids,
288–289
GC-FID limitations, 291
GC-FID methodology, 290–294
general discussion, 287–288
preparation of cellular FAMES for
analysis, 290
profiling cellular FAMES by GC, 289
HPSEC and, 211
and IR spectroscopy with edible oils/but-
ter/margarine, 360
as spectroscopic method of lipid analysis
general discussion, 17–18
trans-18:1 FA analysis by, 191
- Fourier transform near infrared
spectroscopy (FT-NIR)
determination of *trans* fatty acids (FAs)
applications, 306–307
development of FT-NIR
quantification model, 321–325
FA quantification, 320
FT-NIR classification of fats/oils,
314–318
FT-NIR classification of fats/oils
(typical report), 318–320
FT-NIR spectral measurements,
309–312
FT-NIR/FTIR differences, 312–314
future applications, 328–329
GC procedures, 307–309
GC separation of FAME/CLA
isomers, 325–328
official FTIR methods for TFA,
305–306d
official GC methods for TFA,
304–305
reasoning for use of GC as
primary/FT-NIR as secondary,
320–321
TFA issues, 303–304
- Fraction applications
HPSEC
applications, 207–208

crude/refined oils, 223–224
digestibility coefficients in animals,
226
intact oil samples, 218–219
oxidized fats/oils, 224–227
used frying fats/oils, 220–223

Fractionation, *trans*-18:1 FA analysis by
silver ion HPLC and, 195, 200–201

Fragmentation, ESI and, 37

Frankia, EI-MS and, 32

Free fatty acids (FFAs)

classical applications and, 5

Fourier transform infrared spectroscopy
(FTIR) and, 17

packed-column-SFC and, 245–250

RP-HPLC and, 9

silica gel G and separation of, 6

size exclusion chromatography (SEC)
and, 16

Free sterols, size exclusion chromatography
(SEC) and, 16

Frying oils. *See* Fats/oils

G

Galactocerebroside, shotgun lipodomics
and, 65–66

γ -linolenic acid

EI fragmentation of fatty acid esters of,
32

Gas chromatography (GC)

as chromatographic methods of lipid
analysis

general discussion, 13–16

elution orders of FAME/TAG standards,
177

FAME analysis of bacteria and. *See*
Bacteria

foodborne pathogen identification via
ATR-FTIR spectroscopy

multivariate statistical analysis of IR
spectra, 296–297

protocol, 295

foodborne pathogen identification with
FTIR spectroscopy

cultivation of cells, 289–290

fatty acid structures in cellular lipids,

288–289

GC-FID limitations, 291

GC-FID methodology, 290–294

general discussion, 287–288

methodology, 294–296

preparation of cellular FAMES for
analysis, 290

profiling cellular FAMES by GC, 289
two-dimensional (2D), 14

Gas chromatography-mass spectrometry
(GC-MS)

double bond position UFA and, 158,
160

EI ionization mechanism and, 30–33

emergence of, 29

Generally Recognized As Safe (GRAS), 381

Gluococerebroside

shotgun lipodomics and, 65–66

Glycerides (partial)

HPSEC

partial glycerides, 213–214

packed-column-SFC and, 245–250

Glycerol-based lipids. *See also* specific

glycerol, i.e. Triacylglycerols (TAG)

and 2D identification of individual
species, 59–60

Glycerophospholipids

chiral-phase HPLC enantiomeric separa-
tion and, 79

chiral-phase steric analysis of, 95–101

ESI-MS-MS and, 37

LC/MS (recent developments) and,
109–114, 137–149

Glycolipids. *See* Plant tissues

Gums, 385–386, 390–391

H

Hanus/Kaufmann method for IV
determination, 5

Hard GC-MS-EI/LC-MS-EI, 30–33

Healthy People 2000, 379

Hemicelluloses, 385–386

Hexane

extraction methods and, 3

as solvent in Ag-ion HPLC. *See* Silver-
ion chromatography

- Hg-HPLC. *See* Silver-ion chromatography
- High performance liquid chromatography (HPLC). *See also* Silver-ion chromatography
- as chromatographic methods of lipid analysis
 - general discussion, 8–13
 - emergence of, 29
- High performance size-exclusion chromatography (HPSEC)
- characteristics in organic media (phases), 206–211
 - concentrated fraction applications
 - crude/refined oils, 223–224
 - digestibility coefficients in animals, 226 - FAME, 219–221
 - intact oil samples, 218–219
 - oxidized fats/oils, 224–227
 - used frying fats/oils, 220–223
- FAME applications, 218
- lipid applications
- fats/oils replacers, 214–216
 - fish oils, 212–213
 - minor lipid compounds, 216–217
 - partial glycerides, 213–214
 - pesticide analysis, 217–218
 - used frying oils, 211–213
- High-resolution FAB, soft ionization via, 44
- Homoallylic FAME, 163, 167–169
- Homoallylic PUFAs, 157–158
- Homoconjugated PUFAs, 157
- Hopanoids, 36–37
- HPLC-MS-APCI, TAGs and, 39–40
- Hydroperoxides
- diradylglycerol
 - hydroxides/isoprostane/core aldehydes and, 128–131
 - steryl esters and, 124–128 - TAGs/isoprostane/core aldehydes and, 131–137
- Hydroperoxy cholesterol esters, HPSEC and, 217
- Hydroperoxy TG, HPSEC and, 217
- Hydrophobicity of lipids, 3, 64
- Hydroxyeicosatetraenoic acids (HETEs), 75
- chiral-phase HPLC enantiomeric separation and, 81–88
- Hydroxy/hydroperoxy FA, LC/MS (recent developments) and, 114–119
- Hydroxyoctadecadienoic acids (HODEs), 75
- Hypertension, metabolic syndrome and, 51
- ## I
- Iatroscan, 261–262, 267–268
- Induced protein-lipid complexes, 355–356.
- See also* Protein-lipid interactions
- Infrared spectroscopy. *See also* Fourier transform infrared spectroscopy (FTIR)
- Infrared spectroscopy (IR)
- compared with sampling for Raman technique, 364–365
 - edible oils/butter/margarine and, 360
 - peptide bonds and, 359
 - quantification of *trans*/CLAs
 - chemometrics, 337–338
 - experimental
 - standards/measurements/data handling, 343–344
 - general discussion, 335–337
 - PLS technique of calibration, 338–342
 - spectra of triglycerides, 344–350 - vibrational spectroscopy
 - protein-lipid interactions, 357–362
- Instrument response factors
- ESI-MS and, 52
- Intact lipids
- HPSEC
 - intact oil samples, 207, 218–219
 - mass spectrometry, 33–35
- International Union of Pure and Applied Chemistry (IUPAC) Commission on Oils, Fats and Derivatives, 212
- Intrasource separation of lipid classes (shotgun lipidomics), 53–59
- Intrasource separation of lipids, EI-MS and, 55
- Iodine value (IV)
- determination of, 5
 - of waxes, 423
- Ion analysis/detector of mass spectrometer,

- 29–30
- Ion separation methods, 44–46
- Ion trap, chiral-phase HPLC enantiomeric separation and, 84
- Ion trap (MSⁿ), 37, 46
- Ionization of mass spectrometer, 29–30. *See also* Mass spectrometry
- Irish potato famine, 36
- Isobaric species analysis, 61, 63, 66
- Isomer identification, GC-FTIR and, 15–16
- Isoprostanes
- diradylglycerol
 - hydroxides/hydroperoxides and, 128–131
 - glycerophospholipids and, 137–149
 - prostaglandins and, 119–124
 - steryl esters and, 124–128
 - TAGs/hydroperoxides and, 131–137
- J**
- Journal of Agriculture and Food Chemistry*, 388
- K**
- King Midas (Phrygia-700 B.C.), 38–39
- L**
- Labeling of fat replacers, 381
- Laser desorption, 34
- Less fat labeling, 381
- Light food labeling, 381
- Lignins, 385–386
- Linoleic acid, HETE/HODE synthesis and, 75, 85
- Lipid analysis
- automation of, 5
 - chromatographic methods
 - counter current chromatography (CCC), 16
 - gas chromatography (GC), 13–16
 - high performance liquid chromatography (HPLC), 8–13
 - size exclusion chromatography (SEC), 16
 - supercritical fluid chromatography (SFC), 16–17
 - thin-layer chromatography (TLC), 6–8
 - classical applications, 5–6
 - extraction methods, 3–5
 - influential MS ionization methods, 33–34
 - mass spectrometry
 - future applications, 46
 - hard GC-MS-EI/LC-MS-EI, 30–33
 - high resolution/accurate mass FAB and ESI methods, 44
 - modern instrumental
 - separation/detection approaches, 44–46
 - soft ionization methods (large biomolecules/intact lipids), 33–35
 - soft ionization methods via APCI, 38–40
 - soft ionization methods via APPI, 40–42
 - soft ionization methods via ESI, 37–38
 - soft ionization methods via FAB-MS, 36–37
 - soft ionization methods via MALDI-TOF, 42–43
 - minor lipid compounds
 - HPSEC and, 216–217
 - spectroscopic methods
 - Fourier transform infrared spectroscopy (FTIR), 17–18
 - near infrared spectroscopy (NIR), 18
 - nuclear magnetic resonance spectroscopy (NMR), 18–19
 - structural identification using FAB, 44
- Lipidome, 51–52
- Lipidomics. *See also* Shotgun lipidomics
- ESI-MS-MS and, 37–38
- Lipids
- common characteristics of, 3
 - roles of, 51
- Lipoproteins, extraction methods and, 3
- Liquid chromatography (LC)-MS
- advantages of, 10
 - of PtdCho (with ESI), 139–140
- Liquid chromatography (LC)-MS (recent

- developments)
 - materials/methods
 - oxo-FA/cholesteryl
 - esters/TAGs/glycerophospholipids, 109–114
 - results/discussion
 - diradylglycerol
 - hydroxides/hydroperoxides/isoprostanes/core aldehydes, 128–131
 - glycerophospholipid
 - hydroperoxides/isoprostanes/core aldehydes, 137–149
 - hydroxy/hydroperoxy FA, 114–119
 - prostaglandins/isoprostanes, 119–124
 - steryl ester
 - hydroxides/hydroperoxides/isoprostanes/core aldehydes, 124–128
 - TAG hydroxides/hydroperoxides/isoprostanes/core aldehydes, 131–137
 - Low calorie labeling, 381
 - Low density lipoprotein (LDL) cholesterol, 380
 - Low fat labeling, 381
 - Lycadex, 384
 - LysoPE
 - EI-MS and, 55
 - shotgun lipodomics and, 65
 - Lysophospholipids, EI-MS and, 55
- M**
- MALDI. *See* Matrix-assisted laser desorption/ionization (MALDI) technique
 - MALDI-TOF MS
 - emergence of, 29
 - soft ionization via, 42–43
 - MALDI-TOF-TOF, 42
 - MALDI-TOF-TOF-MS-MS, 42
 - Maltin M040, 384
 - Maltodextrins, 384, 390
 - Marine lipids, TLC-FID and, 261–268
 - Mass analyzer, quadrupole, 44–46
 - Mass spectrometry
 - basic components of, 29–30
 - future applications to lipid chemistry/biochemistry of, 46
 - hard GC-MS-EI/LC-MS-EI, 30–33
 - high resolution/accurate mass FAB and ESI methods, 44
 - modern instrumental separation/detection approaches, 44–46
 - soft ionization methods
 - large biomolecules/intact lipids, 33–35. *See also* High performance size-exclusion chromatography (HPSEC)
 - via APCI, 38–40
 - via APPI, 40–42
 - via ESI, 37–38
 - via FAB-MS, 36–37
 - via MALDI-TOF, 42–43
 - Matrices
 - extraction methods and, 3–5
 - HPLC on silica-based, 52
 - Matrix-assisted laser desorption/ionization (MALDI) technique, 8
 - development of, 34–35
 - with time-of-flight mass spectrometry. *See* MALDI-TOF MS
 - Meat/fish, Weibull-Stoldt extraction method and, 4
 - Membrane properties of phospholipids. *See* Phospholipids
 - Metabolic syndrome, 51
 - Methyl esters. *See also* Fatty acid methyl esters (FAME)
 - chiral-phase HPLC enantiomeric separation and, 77
 - conversion to PFB esters, 75
 - supercritical fluid extraction (SFE) and, 17
 - Micelle formation and quantitation of lipid molecular species, 64
 - Microparticulation, 387
 - Microrod TLC technology for quantification of lipid analytes, 8
 - Midas, King (Phrygia-700 B.C.), 38–39
 - Monoacylglycerol (MAG)
 - and 2D identification of individual species, 59–60

- chiral-phase HPLC enantiomeric separation and, 77–78
- chiral-phase steric analysis of, 89–91
- Monoacylglycerols, size exclusion chromatography (SEC) and, 16
- Monohydroxy FA, chiral-phase steric analysis of, 81–88
- Monounsaturated fatty acids, α -oxidation and, 67
- MSⁿ, 37, 46
- Mycobacteria*, long-chain FA in, 271
- N**
- Naphthylethyl urethane diastereomeric DAG derivatives and, 80–81
- Near infrared spectroscopy (NIR) as spectroscopic method of lipid analysis general discussion, 18
- Nitrogen fixing, EI-MS and, 32
- Nitrogen ions. *See* Atmospheric pressure chemical ionization (APCI)
- Nobel Prize, 29, 37
- N-Oil, 384
- Non-methylene-interrupted PUFA (NMI-PUFA), 158, 168
- Nonpolar solvents, extraction methods and, 3
- Nuclear magnetic resonance (NMR) spectrometry
- 2D identification of individual species and, 61
- Nuclear magnetic resonance spectroscopy (NMR) as spectroscopic method of lipid analysis general discussion, 18–19
- Nutricol, 386
- Nutrio-P-fiber, 386
- O**
- Oatrim, 386
- Obesity. *See also* Fat replacers global, 379–380 metabolic syndrome and, 51
- Octadecylsilane, RP-HPLC and, 8–9
- Octylsilane, RP-HPLC and, 8–9
- Oilseeds
- Fourier transform infrared spectroscopy (FTIR) and, 18
- nuclear magnetic resonance spectroscopy (NMR) and, 18–19
- packed-column-SFC and, 250–254
- Weibull-Stoldt extraction method and, 4
- Olestra, 214–215, 381, 390
- Olestra (Olean), 383
- Oomycetes, high-resolution FAB and, 44
- Oxo-FA, LC/MS (recent developments) and, 109–114
- Oxysterols, 426–427
- P**
- P-150 c/P-285 F, 386
- Partial glycerides
- HPSEC and, 213–215
- Partial least square (PLS) calibration or regression
- quantification of *trans*/CLAs chemometrics, 337–338
- experimental standards/measurements/data handling, 343–344
- general discussion, 335–337
- spectra of triglycerides, 344–350
- technique, 338–342
- Partially esterified polysaccharide (PEP), as fat replacers, 382
- Partially hydrogenated vegetable oils (PHVO), *trans*-18:1 FA analysis and, 191
- Partition chromatography, 8–9
- Paselli SA2, 384
- p*-coumaroylferuloylputrescine (CFP), 31
- Pectins, 385–386
- Pentafluorobenzyl (PFB) esters
- chiral-phase HPLC enantiomeric separation and, 77
- methyl ester conversion to, 75
- Peptide bonds, infrared bands of, 359
- Peroxide value (PV), 5–6
- Pesticide analysis, HPSEC and, 217–218
- Petrochemicals, HPSEC and, 217–218
- pH conditions

- fat replacers and, 389
- lipid classifications and, 55–56
- Phosphatidic acid, EI-MS and, 55
- Phosphatidic acid, shotgun lipodomics and, 65
- Phosphatidylcholines, 12, 36
- Phosphatidylglycerol (PtdGros), 55–58, 65, 95, 97–98
- Phosphatidylinositol
 - EI-MS and, 55
 - shotgun lipodomics and, 65
- Phosphatidylserine
 - EI-MS and, 55
 - shotgun lipodomics and, 65
- Phosphatidylserines (PtdSers), 99–101
- Phosphoacylglycerols (PAG),
 - packed-column-SFC and, 250–254
- Phospholipids
 - bilayer/membrane properties
 - curvature/stress/surface tension, 406
 - dipole potential, 406–407
 - fluidity/diffusion/transbilayer movement, 402
 - heat capacity, 403–404
 - miscibility/solubility, 404–405
 - permeability, 405–406
 - phase transitions, 402–403
 - chemical modification
 - complexation to cholesterol, 411–412
 - complexation to ions, 411
 - complexation to protein, 412
 - conjugation, 414
 - hydration, 409–411
 - hydrogenation, 407–408
 - hydrolysis, 408–409
 - hydroxylation, 408
 - oxidation, 412–414
 - chemical structures, 399–401
 - ELSD and, 10–12
 - ESI-MS-MS and, 37
 - extraction methods and, 3
 - initial ESI-MS studies of, 53
 - physical modification by external forces, 407
 - physical structures, 401–402
 - silica gel G and separation of, 6–7
- Photodiode array detector (PDA), HPLC and, 9
- Photoionization. *See* Atmospheric pressure photoionization (APPI)
- Photomultiplier conversion dynode, ion detection and, 46
- Phytophthora infestans*
 - FAB-MS and, 36
 - high-resolution FAB and, 44
- Picolinyl esters, GC/MS for double bond position UFAs and, 160–161
- Plant tissues, ESI-MS-MS and, 37, 148
- Plasma desorption, 34
- Polydextrose, 385
- Polymerized triacylglycerols (TG), intact oil samples
- Polyunsaturated fatty acids (PUFAs), 157–158
 - counter current chromatography (CCC) and, 16
 - necessity of high sensitivity methodology for, 161–162
 - non-methylene-interrupted (NMI), 168
- Pre-cursor ion scanning, and 2D identification of individual species, 60–62
- Prostaglandins, chiral-phase HPLC enantiomeric separation and, 81–82, 85
- Prostaglandins/isoprostanes, LC/MS (recent developments) and, 119–124
- Protein-based fat substitutes, 386–388
- Protein-lipid interactions
 - infrared bands of peptide bonds, 359
 - infrared frequencies of protein secondary structures, 359
 - vibrational spectroscopy
 - general discussion, 355–357
 - history/development of IR spectroscopy, 357–358
 - IR spectroscopy applications, 358–362
 - principles, 357
 - Raman spectroscopy, 362–371
- Proteins/peptides
 - ESI and, 37
 - MALDI-TOF MS and, 43
- Proteomics, MALDI-TOF MS and, 43
- Pseudomolecular ions, 36

Psoriatic skin scales, chiral-phase HPLC enantiomeric separation and, 81–83
PtdCho, 139–140, 143–144

Q

Quadrupole ion trap mass analyzers, 44–46
2D identification of individual species and, 61
acetonitrile CACI for FAME analysis and, 162–163
Quantification of lipid analytes, 7–8
Quantitation of lipid species by 2D MS, 63–65

R

Racemic glycerolipids, chiral-phase HPLC and, 73–74
Raman spectroscopy
history/development of, 363–365
vs. IR, 362–363
protein-lipid applications
edible oils, 365, 367
food proteins, 365–366
Reduced calorie labeling, 381
Reduced fat labeling, 381
Reduced-calorie fat-based fat replacers, 388–389
Reflectron time-of-flight (RTOF) analyzer, 46
Refractive indexes, analyte detection after HPLC using, 9
Regiospecific lipid identification, 61, 63
Reproducibility of ESI-MS, 52
Reversed-phase HPLC
chiral-phase HPLC enantiomeric separation and, 82–85
TAGs/hydroperoxides/isoprostanes/core aldehydes and, 131–137
Reversed-phase HPLC (RP-HPLC), 8–9, 38
Rhizomucor miehei, 94
Rhizopus niveus, 97

S

Salatrim, 388
Saponification value (SV)

determination of, 5
of waxes, 423
Saturated fatty acids, α -oxidation and, 67
Selected ion monitoring (SIM)-mode of GC, 14
Shotgun lipidomics
2D identification of lipid molecular species, 59–63
2D quantitation of lipid molecular species, 63–65
biological applications of, 65–67
intrasource separation of lipid classes (shotgun lipidomics), 53–59
Signal-to-noise ratio, ESI-MS and, 52
Silanes, RP-HPLC and, 8–9
Silica gel G as adsorbent in TLC, 6–7
Silica-based matrices, 52–53
Silver nitrate
with adsorbent silica gel G, 6
TLC-FID and, 263
Silver-ion chromatography
advantages of, 9
general discussion, 175–176
temperature effects on lipid elution
discussion, 183–186
HPLC methodology, 176–179
TAG/FAME separations, 179–183
trans-18:1 FA analysis
elution order, 195, 197–199
fractionation, 195, 200–201
history, 191–192
internal standard for separations, 194–196
separation as FAME, 192–194
triacylglycerol isomer determination and, 9
Simplese, 387, 390–391
Size exclusion chromatography (SEC)
as chromatographic methods of lipid analysis
with CCC, 16
general discussion, 16
Soft ionization methods
large biomolecules/intact lipids, 33–35
via APCI, 38–40
via APPI, 40–42
via ESI, 37–38

- via FAB-MS, 36–37
 - via MALDI-TQF, 42–43
 - Software, GC-MS-EI comparisons and, 33
 - Solid phase extraction (SPE) methods, 3, 219
 - Solids in fats determination, FTIR and, 17–18
 - Solvents
 - CUBO, 19
 - extraction methods and, 3
 - fish lipid extraction and, 4
 - lipid solubility in nonpolar, 3
 - Soxhlet-apparatus, Weibull-Stoldt extraction method and, 4
 - Spectroscopic methods of lipid analysis. *See also* Mass spectrometry
 - Fourier transform infrared spectroscopy (FTIR), 17–18
 - near infrared spectroscopy (NIR), 18
 - nuclear magnetic resonance spectroscopy (NMR), 18–19
 - SPE-Olestra (Olean). *See* Olestra
 - Sphingolipids
 - ESI-MS-MS and, 37
 - FAB-MS and, 36
 - Sphingomyelin analysis
 - EI-MS and, 55
 - ESI-MS-MS and, 12
 - shotgun lipidomics and, 65
 - Sphingosine-based lipids. *See also* specific sphingosine, i.e. Sphingomyelin
 - Sphingosine-based lipids, and 2D identification of individual species, 59–60
 - Standards
 - and quantitation of lipid molecular species, 63–65
 - trans*-18:1 FA analysis by silver ion HPLC and, 194–196
 - Starch-based fat replacers, 384–386
 - StoSlim, 385
 - Stellar, 385
 - Stenotrophomonas maltophilia*, 277, 279–280
 - Stereoisomers
 - bis(monoacylglycerol)phosphate (BMP), 97
 - LC/ESI-MS and MS/MS of, 79–81
 - Steroids (anabolic), APPI and, 42
 - Sterols
 - analysis/characterization of, 428–431
 - packed-column-SFC and, 250–254
 - structure, 424–428
 - Steryl ester hydroxides/hydroperoxides/isoprostanes/core aldehydes, LC/MS and, 124–128
 - Streptomyces septatus*, 97–98
 - Stroke, metabolic syndrome and, 51
 - Subcritical fluid chromatography (SFC), 245
 - Sucrose polyesters, 215–216, 390–391
 - as fat replacers
 - SPE-Olestra (Olean), 214–215, 381, 383, 390
 - Sulfatide
 - EI-MS and, 55
 - shotgun lipidomics and, 65
 - Supercritical carbon dioxide extraction methods and, 3
 - Supercritical fluid chromatography (SFC) as chromatographic methods of lipid analysis
 - general discussion, 16–17
 - general discussion, 239–241
 - manufacturers of, 242
 - modifiers, 244–245
 - packed column applications of lipids
 - PAG/sterols/tocopherols/seed oils, 250–254
 - TAG/FFA/glyceride mixtures, 245–250
 - packed column applications of saccharides, 250
 - packed vs. capillary columns, 241–244
 - subcritical fluid chromatography and, 245
 - Supercritical fluid extraction (SFE), advantages of, 4–5
- T**
- TAG hydroxides/hydroperoxides/isoprostanes/core aldehydes, LC/MS and, 131–137

- Tandem electrospray ionization
(ESI-MS-MS). *See also* Acetonitrile
covalent-adduct chemical ionization
(CACI) MS-MS
- 2D identification of individual species
and, 61–62
- epoxy FAs and, 89
- and quantitation of lipid molecular
species, 65
- Tandem mass spectrometry. *See* MSⁿ
- Temperature
- chiral-phase HPLC and, 73
- effects on lipid elution (silver-ion
HPLC)
- discussion, 183–186
- general discussion, 183–186
- HPLC methodology, 176–179
- TAG/FAME separations, 179–183
- for HPLC, 13
- ThermaBeam[®] MS, 31, 44
- Thermolabile analytes
- HPLC and, 13
- supercritical fluid extraction (SFE)
and, 16–17
- Thermospray, 34
- Thin-layer chromatography (TLC)
- as chromatographic methods of lipid
analysis
- general discussion, 6–8
- with flame ionization detector (FID)
- general discussion, 261–262
- principles, 262–263
- quantitation/applications, 263–268
- trans-18:1 FA analysis and, 191, 195,
200
- Thomson, Joseph John, 29
- Threo-DiHETERe, 88–89
- Time-of-flight (TOF) analyzer, 44–46
- Tocopherols, packed-column-SFC
and, 250–254
- Total fat content, 4. *See also* Lipid analysis
- Toxicology of fat replacers, 391–392
- Trans* fatty acids (TFAs). *See also*
Trans-18:1 FA analysis
- exclusion of CLA for, 304–306
- formation of, 158
- Fourier transform infrared spectroscopy
(FTIR) and, 17
- quantification with IR spectroscopy
- chemometrics, 337–338
- experimental
- standards/measurements/data
handling, 343–344
- general discussion, 335–337
- PLS technique of calibration,
338–342
- spectra of triglycerides, 344–350
- use of FT-NIR for determination
- applications, 306–307
- development of FT-NIR
- quantification model, 321–325
- FA quantification, 320
- FT-NIR classification of fats/oils,
314–318
- FT-NIR classification of fats/oils
(typical report), 318–320
- FT-NIR spectral measurements,
309–312
- FT-NIR/FTIR differences, 312–314
- future applications, 328–329
- GC procedures, 307–309
- GC separation of FAME/CLA
isomers, 325–328
- official FTIR methods for TFA,
305–306d
- official GC methods for TFA,
304–305
- reasoning for use of GC
- as primary/FT-NIR as
secondary, 320–321
- TFA issues, 303–304
- Trans* monoethylenic fatty acids
- GC determination of, 13–14
- Trans*-18:1 FA analysis
- silver-ion chromatography
- elution order, 195, 197–199
- fractionation, 195, 200–201
- history, 191–192
- internal standard for separations,
194–196
- separation as FAME, 192–194
- Transesterification reactions, HPSEC and,
214, 226
- Triacylglycerol isomer determination

- RP-HPLC and, 9
silver-ion HPLC and, 9
Triacylglycerols (TAGs)
and 2D identification of individual
species, 59–60
diastereomeric derivatives of, 76
effects on lipid elution (silver-ion
HPLC) and, 179–183
EI-MS and, 55
GC determination of, 13
HPLC-MS-APCI and, 39–40
LC/MS (recent developments), 109–114
LC/MS (recent developments) and,
131–137
long-chain FA in, 388
NEU/PEU derivatives and, 95
packed-column-SFC and, 245–250
short-chain FA in, 388
shotgun lipodomics and, 65–67
silver-ion HPLC and temperatures,
182–186
size exclusion chromatography (SEC)
and, 16
supercritical fluid extraction (SFE)
and, 17
Two-dimensional identification of lipid
molecular species, shotgun
lipodomics and, 59–63
Two-dimensional quantitation of lipid
molecular species, shotgun
lipodomics and, 63–65
- U**
- Ultraviolet detectability
HPLC and, 9
HPSEC and, 206
MALDI-TOF MS and, 42
Unsaturated fatty acid (UFA) methyl ester
isomers
acetonitrile covalent-adduct chemical
ionization (CACI) MS-MS
acetonitrile CACI for FAME
analysis, 162–170
characterization of UFA double bond
position, 158–162
covalent adduct chemical ionization
(CACI), 162
Unsaturated fatty acids (UFAs),
configurations/classes of, 157–158
Urine (human)
HETE/HODE synthesis and, 85–87
- V**
- Vibrational spectroscopy
protein-lipid interactions
general discussion, 355–357
history/development of IR
spectroscopy, 357–358
IR spectroscopy applications,
358–362
principles, 357
Raman spectroscopy, 362–371
- W**
- Waters Corporation, 31
Waxes
analysis/characterization of, 422–424
structure, 421–422
Weibull-Stoldt method, 3–4
Whey protein concentrates, 387
Wijs method for IV determination, 5, 423
World Health Organization (WHO),
on obesity, 379–380
- X**
- Xylenol orange method, determination of
peroxide value (PV) by, 6
- Z**
- Zymomonas mobilis*, FAB-MS and, 36–37