

Index

A

- AACC, *see* American Association of Cereal Chemists
- A- and B-type proanthocyanidins, 156
- Accelerated solvent extraction, 169
- AccQ-Tag, 456
- Acetyl-CoA carboxylase (ACCCase), 11
- Acid-butanol assay, in flavonoids analysis, 170
- AHP, *see* American Herbal Pharmacopoeia
- Ailanthus altissima*, 370
- Amberlite polymeric adsorbents, 255–256
- Amberlite XAD-7 column chromatography, *see* Amberlite polymeric adsorbents
- American Association of Cereal Chemists, 468
- American Herbal Pharmacopoeia, 478
- American Oil Chemists' Society, 105, 171
- Amino acids
- acid hydrolysis of, 443–445
 - alkaline hydrolysis of, 445–446
 - chromatography
 - ion-exchange, 456–460
 - reversed-phase, 460–462
 - detection of
 - 9-Fluorenylmethyl chloroformate precolumn derivatization, 451–453
 - ninhydrin postcolumn derivatization, 451–453
 - o*-Phthalaldehyde postcolumn derivatization, 453–454
 - o*-Phthalaldehyde precolumn derivatization, 455–456
 - extraction, 448
 - internal standards and controls, 449–450
 - in nutritional matrices, 443
 - protein and non-protein, 443–445
 - protein and peptides
 - alkaline hydrolysis, 445–446
 - microwave acid hydrolysis for, 447–448
- Anthocyanins, 155; *see also* Plant flavonoids, chemistry of
- extraction of, 252–253
 - identification of
 - chromatographic techniques, 263
 - hydrolysis, 262–263
 - spectroscopy for, 264–270
 - separation and purification of
 - capillary zone electrophoresis, 260–262
 - column chromatography, 255–256
 - countercurrent chromatography, 257–258
 - high-performance liquid chromatography, 258–260
 - paper chromatography, 253–254
 - solid-phase extraction, 256–257
 - thin-layer chromatography, 254–255
 - structures and chemistry of, 247–251
- Anthocyanins in grapevine, 211–213; *see also* Polyphenolic compounds
- Anticarcinogenic and anti-atherogenic effects, of CLA, 91
- AOCS, *see* American Oil Chemists' Society
- Apigenin, 149–150, 153, 187–188
- Apigenin-7-glucoside, health benefits, 226–227
- Aronia melanocarpa*, 257
- Artificial colorants, 320
- Ascorbic acid (AA), 417–418
- ASE, *see* Accelerated solvent extraction
- Aspergillus niger*, 17
- Association of Official Analytical Chemists (AOAC), 101
- Atmospheric pressure chemical ionization (APCI), 47, 186, 267
- Atmospheric pressure ionization (API), 49
- Attention deficit hyperactivity disorder (ADHD), 89

B

- Beer–Lambert law, 316, 319, 350, 377, 382
- Bioassay system, in phytoestrogens determination, 69
- Biochanin A; *see also* Isoflavones compounds
- fluorescence properties of, 9
 - solid-phase extraction of, 19
 - sources of, 6
 - structure of, 7
- Biotin, 405, 413–414
- Bixa orellana*, 282, 284
- Bovine serum albumin (BSA), 66
- BSTFA, *see* *N,O*-bis-(trimethylsilyl)-trifluoroacetamide
- Buffelute, 459
- Burgen Soy-Lin, health benefits of, 4
- Butylated hydroxytoluene (BHT), 15

C

- Cancer treatment and phytoestrogens, 3
- Capacity factor, 369
- Capillary electrophoresis (CE) technique
 - in OBPs separation, 233
 - in phytoestrogens determination, 60–63
- Capillary zone electrophoresis (CZE)
 - in flavonoids separation, 178
 - in isoflavones separation, 61
 - separation of anthocyanins, 260–262
- Carbohydrate
 - analytical aspects of, 478–480
 - complex, 474–475
 - dietary, 475–477
 - HPAEC-PAD analysis
 - agricultural applications, 510–511
 - food and beverage, 499–510
 - industrial applications, 510–511
 - role in nutrition, 471–473
 - serotonin effect, 471
 - structure of, 472
- Carbowax-templated polydivinylbenzene resin (CW-TPR), 21
- Cardiovascular disease (CVD), 86, 88
- β -Carotene, 324–325; *see also* Provitamin A
 - absorption maxima, 287, 312
 - antioxidant effects, 279
 - description of, 278
 - fine structure, 289
 - quantative determination in foods, 303
 - retinol activity equivalent, 290–291, 324–325
 - ultraviolet-visible spectrophotometric determination, 319
 - uses of, 294
- Carotenoids, 324–325; *see also* Provitamin A
 - absorption coefficients, 316
 - Beer–Lambert law, 316
 - chromatographic techniques
 - column chromatography, 298–301
 - high-performance liquid chromatography, 302–310
 - thin layer chromatography, 301–302
 - classification of, 280
 - description and distribution of, 284–285
 - 5,6-epoxide groups, 296
 - extraction method
 - phase distribution, 298
 - sample and solvent preparation, 297
 - saponification, 298
 - functional groups identification
 - acetylation of hydroxyl groups, 315–316
 - allyl hydroxyl groups, 316
 - 5,6-epoxide group, 314
 - reduction of carbonyl groups, 314–315
 - industrial production of, 294–295
 - isomerization, 296
 - nutritional and beneficial properties of
 - antioxidant effects, 291
 - bioavailability, 292
 - retinol equivalent activity, 290–291
 - oxidative degradation of, 296
 - in plants and animals, 285
 - as protectors, 284
 - quantitative determination
 - provitamin A, 324–325
 - UV-visible spectrophotometry, 319–324
 - separation and isolation method, 298–299
 - spectroscopic properties
 - fine structure, 287–288
 - fluorescence and IR spectrum of, 287
 - UV-visible light absorption spectrum of, 286–289
 - standards preparation, 310
 - structure of, 280–281
 - uses of
 - in clinical and pharmaceuticals, 294
 - food colorants, 292–294
- CEAD. *see* Coulometric electrode array detection
- Centrifugal partition chromatography (CPC), in anthocyanins separation, 258
- Cetyltrimethylammonium bromide (CTAB), 262
- Chlorophyll
 - absorption coefficients of, 379
 - as additives, 360–362
 - allomerization and epimerization, 345–346
 - characteristics of, 339
 - chemical properties, 343–350
 - chromatographic separation techniques
 - column chromatography, 365
 - high-performance liquid chromatography, 366–368
 - paper chromatography, 365
 - thin layer chromatography, 365, 371–375
 - derivatives of, 349
 - description of, 338
 - extinction coefficients of, 377–378
 - extraction techniques
 - chromatographic separation techniques, 364–368
 - isolation and purification, 368–369
 - methodology, 363–364
 - pigment identification and standards, 369–371, 376
 - quantification analysis, 376–387
 - functional properties
 - anticarcinogenic activity, 347–348
 - antimutagenic activity, 346–347
 - antioxidant properties, 349

- chemoprevention of, 348–349
 - digestion, absorption, metabolism, 349–350
 - in higher plants, 343
 - porphyrin system of, 340
 - quantitative analysis
 - Beer–Lambert law, 377
 - extinction coefficients of, 377–378
 - spectroscopic properties
 - fluorescence spectra of, 355–356
 - infrared absorption bands, 350, 356
 - primary allomerization reactions, 353
 - visible/UV electronic absorption spectra, 350–355
 - stability during food processing
 - conservation treatments, 357–359
 - degradation mechanism, 359–360
 - dehydration, 359
 - pheophytinization, 360
 - sterilization process, 359
 - structural modification of
 - chelate modification, 343–345
 - de-esterification, 345
 - isocyclic ring reactions, 345–346
 - macrocycle cleavage of, 346
 - structure and types of, 339–343
 - vs. protein interactions, 342
 - Chrysin, 149–151, 153
 - Cinnamate 4-hydroxylase (C4H), 11
 - Citrus bioflavonoids. *see* Flavanones
 - Cobalamins, 416
 - Collision activated decomposition (CAD), 47
 - Collision-induced dissociation (CID), 47, 186
 - Colorimetric assay, in flavonoids analysis, 171
 - Column chromatography (CC), in anthocyanins separation, 255–256
 - Conjugated linoleic acid (CLA)
 - analysis of, 131–132
 - commercial isomer distribution, 119–124
 - derivation of, 105–107
 - foods with, 97–99
 - health benefits of, 90–92
 - measurement of total, 129–131
 - natural isomer distribution, 124–129
 - Consumer Products and the Environment (COT), 2
 - Correlation spectroscopy (COSY), 189
 - Coulometric array detection, 38–39
 - Coulometric electrode array detection, 182–183
 - 4-Coumarate ligase (4CL), 11
 - Coumestans compounds
 - electrochemical detection of, 37–38
 - fluorescence properties of, 9
 - health benefits of, 2–3
 - structural features of, 7
 - Countercurrent chromatography (CCC)
 - in anthocyanins separation, 257–258
 - in flavonoids separation, 176–178
 - Critical micelle concentration (CMC), 178
 - Cryptocodinium cohnii*, 94
 - Cuminum cyminum*, 172
 - Cyanidin 3-glucoside, isolation of, 252
 - Cyanocobalamins, 416
 - Cysteine
 - description of, 446
 - performic acid oxidation of, 446–447
- ## D
- Dacrycarpus dacrydioides*, 256
 - Daidzein; *see also* Isoflavones compounds
 - electrochemical detection of, 37–38
 - fluorescence properties of, 9
 - HPLC-ESI-MS method for determination of, 52
 - RIA for, 66–67
 - separation by CE method, 60–61
 - solid-phase extraction of, 19
 - sources of, 6
 - structure of, 7
 - Daucus carota*, 268
 - DCCC. *see* Droplet countercurrent chromatography
 - Degree of polymerization (DP), 479
 - Dehydroascorbic acid (DHAA), 417–418
 - Delphinidin 3-glucoside, isolation of, 252
 - Designer food, 467
 - Dietary carbohydrates
 - complex carbohydrates, 474
 - dietary fiber, 476
 - resistant and modified starch, 475–476
 - sugars, 474
 - Dietary fiber
 - analysis, 479–480
 - classification and components of, 476
 - definition, 476
 - Dietary Supplement Health and Education Act, 467
 - Dietary supplements, 467–468
 - Dihomo- γ -linolenic acid (DGLA), 89
 - 2,3-Dihydroflavonols, *see* Flavanonols
 - p-Dimethylaminobenzaldehyde, 445
 - 4,4-Dimethyloxazoline (DMOX), 108
 - GC methods, 114–115
 - Dimethyl sulfoxide, 453
 - Diode array detection (DAD), 182
 - Dioxins, effects of, 348
 - 1,3-Diphenylpropane, 149

2,6-di-*tert*-Butyl-4-methylphenol (BHT), 226
 DMSO, *see* Dimethyl sulfoxide
 Docosahexaenoic acid (DHA), health benefits of, 88–89
 Droplet countercurrent chromatography, 258
 DSHEA, *see* Dietary Supplement Health and Education Act

E

E-141i, *see* Sodium-copper-chlorophyllin
 Eicosapentaenoic acid (EPA), health benefits of, 88–89
 Electrocatalysis
 cyclic voltammetric analysis, 484–486
 detection methods, 484–485
 electrode characteristics, 483
 Electrochemical detection (ED), 37, 182–183
 Electrodes, noble metal, 483
 Electrospray (ES) mass spectrometry, 8
 Electrospray ionization (ESI), 47, 173
 Electrospray ionization mass spectrometry (ESI-MS), 267
Enterococcus hirae, 415
 Enterodiol, structure of, 7
 Enterolactone, structure of, 7
 Enzyme immunoassay (EIA), 68
 Equivalent chain length (ECL), 112
Escherichia coli, 417
 Esterified fatty acids (EFA), 101
 HPLC methods in analyzing, 115
 17- β -Estradiol, health benefits of, 4
 Ethanol, health benefits of, 209
 European Prospective Investigation of Cancer (EPIC), 2
 Evaporative light-scattering detector (ELSD), 115–116

F

FAD, *see* Flavin adenine dinucleotide
 FAO, *see* Food and Agriculture Organization
 Fast atom bombardment (FAB), 186
 Fast atom bombardment (FAB)-MS technique, 265
 Fatty acid methyl esters (FAME), 101
 qualitative GC in analysis of, 109–112
 quantitative GC in analysis of, 113–114
 ω 3 Fatty acids, 109–116
 biosynthetic pathway, 87
 foods with, 94–97
 health benefits of, 88–89
 recommended level of, 93–94

Fatty acids (FA)
 derivation of
 dimethylxozoline derivatives, 108–109
 methyl esters, 104–108
 extraction of
 free fatty acids recovery, 101–102
 total lipids extraction, 102–104
 health benefits of, 87–93
 qualitative and quantitative analysis of
 conjugated linoleic acid, 119–132
 pretreatment in, 109
 trans fatty acids, 116–119
 ω -3 and other fatty acids, *see* ω 3 Fatty acids
 recommended level of, 93–94
 types of, 86
 FDA, *see* Food and Drug Administration
 Federal Food, Drug and Cosmetic Act, 469
 FFDC, *see* Federal Food, Drug and Cosmetic Act
 FFH, *see* Functional Foods for Health program
 Flame ionization detection (FID), 101
 Flavandriols, 155; *see also* Plant flavonoids,
 chemistry of
 Flavan-3,4-diols, 155, 216
 Flavanols, 155, 213–216; *see also* Plant flavonoids,
 chemistry of; Polyphenolic compounds
 Flavan-3-ols in grapes, 213–216
 Flavanones, 151; *see also* Plant flavonoids,
 chemistry of
 Flavanonols in grapevine, 151; *see also*
 Polyphenolic compounds
 Flavin adenine dinucleotide, 409
 Flavin mononucleotide, 409
 Flavones, 149–151, 217; *see also* Plant flavonoids,
 chemistry of; Polyphenolic compounds
 Flavonoid-*O*-glycosides, 189
 Flavonoids
 analysis with spectrophotometric methods,
 170–171
 beneficial effect of, 148, 191
 biosynthesis pathway of, 150
 chemical structure of, 152–154
 classes of, 149–156
 detection systems of
 electrochemical detection, 182–183
 fluorescence detection, 183–185
 mass spectrometry, 186–189
 nuclear magnetic resonance, 189–191
 UV-Visible, 180–182
 effect of food processing and storage, 160–161
 extraction of, 163–166
 intake of, 161–162
 occurrence in plant foods and nutraceuticals,
 156–160
 role in plant physiology, 149
 sample preparation of, 161–162

- solid-phase extraction, 167–168
 - solvent extraction and LLE, 162–167
 - supercritical and subcritical fluid extraction, 168–170
 - separation methods of
 - chromatographic techniques, 172–178
 - electrophoretic techniques, 178–179
 - structure and numbering system of, 151
 - types of food, 155
 - Flavonols, 149–151, 216–217; *see also* Plant flavonoids, chemistry of; Polyphenolic compounds
 - Flow injection analysis (FIA), 59
 - Fluorescence detection, in flavonoids detection, 183–185
 - Fluorescence spectrum, for isoflavonoids, 9
 - FMN, *see* Flavin mononucleotide
 - Folch method, 103
 - Folic acid, 414–416
 - Folin–Ciocalteu method, in flavonoids analysis, 160, 171
 - Food and Agriculture Organization, 289, 324
 - Food and Drug Administration, 292, 470
 - Food carbohydrates, 473
 - Foods and beverages, extraction of flavonoids, 163–165
 - Foods for Specified Health Use, 466
 - Formononetin; *see also* Isoflavones compounds
 - fluorescence properties of, 9
 - LC-PB-MS technique in, 47
 - RIA for, 67–68
 - solid-phase extraction of, 19
 - sources of, 6
 - structure of, 7
 - Forsythia intermedia*, 13
 - FOSHU, *see* Foods for Specified Health Use
 - Fourier transform infrared (FTIR) spectroscopy, 115, 477
 - Fourier transform ion cyclotron resonance (FT-ICR), 269
 - Fragaria ananassa*, 252
 - Free amino acids, 448–449
 - Free fatty acids (FFA), 94
 - HPLC methods in analyzing, 115–116
 - recovery of, 101–102
 - FT-NIR methods, in total trans fatty acid analysis, 119
 - Functional foods, 481–492
 - classification of, 467
 - conferences on, 468–470
 - definition of, 468
 - description of, 466–467
 - designer food, 467
 - history, 482–483
 - noncarbohydrate applications
 - biogenic amines, 512
 - biotin chromatogram, 514
 - thiocompounds detection, 515
 - nutrition, 468
 - prebiotics and probiotics, 467
 - scientific criteria for, 470–471
 - Functional Foods for Health program, 467
 - Functional foods, in carbohydrates
 - agricultural applications, 510
 - analytical aspects
 - dietary fiber, 480
 - starch, 479
 - beverage applications
 - chromatographic analysis, 500
 - HPAEC-PAD method, 501
 - PAD detection, 499
 - food additives and nutriment applications
 - chromatogram analysis, 503
 - HPAEC-PAD method, 503–505
 - fruit juices applications
 - adulteration technique, 502–503
 - chromatogram analysis, 502
 - HPAEC-PAD, 502–503
 - primary sources, 501
 - industrial applications, 511–512
 - pulsed electrochemical detection
 - detection methods, 486–498
 - electrocatalysis, 465, 483
 - electrochemical detection, 482
 - solid food applications
 - acid hydrolysis analysis, 506, 509
 - chromatogram analysis, 505, 509, 511
 - HPAEC-PAD method, 505–510
 - nonstarch polysaccharides analysis, 506
- ## G
- Gas chromatography (GC), 100
 - in anthocyanins identification, 263
 - in CLA separation, 119–120
 - in flavonoids separation, 173–174
 - Gas chromatography-mass spectrometry (GC-MS)
 - in anthocyanins identification, 263
 - in CLA separation, 121–124
 - in fatty acid extraction, 101
 - in phytoestrogens separation, 15
 - Genistein, 6–7; *see also* Isoflavones compounds
 - electrochemical detection, 37–38
 - extraction of, 16
 - fluorescence properties of, 9
 - HPLC-ESI-MS method for
 - determination of, 52
 - RIA for, 67–68
 - separation by CE method, 60–61

- solid-phase extraction of, 19
- sources of, 6
- structure of, 7
- GI, *see* Glycemic index
- Glycemic index, 471
- Glycosidic bonds of lignans, hydrolysis of, 18–19
- Glycyrrhizae glabra*, 6
- Grape pomace
 - extraction of, 222–224
 - uses of, 221
- Grape seeds, antioxidants in, 224–225
- Green extraction processes, 223

H

- Heated nebulizer-atmospheric pressure chemical ionization (HN-APCI), 36
- Helix pomatia*, 17
- Hesperetin, 151–152, 180, 187
- Heteronuclear multiple bond correlation (HMBC), 191
- Heteronuclear single quantum coherence (HSQC), 191
- High-density lipoprotein (HDL), 87
- High-performance anion-exchange chromatography-Pulsed electrochemical detection, 514
- High-performance anion-exchange chromatography with pulsed amperometric detection
 - agricultural applications, 510–511
 - beverage analysis
 - carbohydrates and alcohols, 500–501
 - instant coffee, 501
 - food additives and nutrients, 502–504
 - fruit juices, 501–503
 - industrial applications
 - biofuels, 511
 - paper industry, 511–512
 - noncarbohydrate applications
 - biogenic amines, 512–513
 - biotin chromatogram, 514
 - thiocompounds detection, 515
 - solid foods, 504–510
- High-performance capillary electrophoretic analysis (HPCE), 17, 63–64
- High-performance liquid chromatography
 - advantages of, 366
 - analysis
 - functional foods, 478
 - oligosaccharides, 479
 - in anthocyanins separation, 250, 258–260
- Beer–Lambert law, 382
- carbohydrates applications, 510
- in carotenoids
 - chromatographic techniques, 303–304
 - detection systems, 304–305
 - quantitative determination
 - in animal products, 322–324
 - in green vegetables, 320–321
 - in processed vegetables, 322
 - in ripe fruits, 321–322
 - detection of, 367–368
 - electrochemical detection in, 482
 - electronic absorption spectra, 381–387
 - in fatty acids analysis, 101
 - in flavonoids separation, 172–176
 - herbal products, 478
 - ion-exchange chromatography, 456–460
 - in phytoestrogens analysis, 14
 - pigment separation, 378–380
 - quantitative pigment analysis, 380–387
 - retention factor of, 369
 - reversed-phase chromatography, 460–462
 - types of, 456
- High-performance liquid chromatography-diode array detector analysis, 228
- High-performance liquid chromatography-integrated pulsed amperometric detection, 498–499
- High-performance TLC, 23, 172–173
- High-speed countercurrent chromatography
 - in anthocyanins separation, 258
 - in flavonoids separation, 176
- High temperature-high resolution (HT-HR), in flavonoids separation, 173
- HMRLignan, health benefits of, 5
- Hormone replacement therapy (HRT), 2
- HPAEC-PAD, *see* High-performance anion-exchange chromatography with pulsed amperometric detection
- HPAEC-PED, *see* High-performance anion-exchange chromatography-Pulsed electrochemical detection
- HPLC, *see* High-performance liquid chromatography
- HPLC-DAD, *see* High-performance liquid chromatography-diode array detector analysis
- HPLC-IPAD, *see* High-performance liquid chromatography-Integrated pulsed amperometric detection
- HPTLC, *see* High-performance TLC
- HSCCC, *see* High-speed countercurrent chromatography
- Humulus lupulus*, 5
- Hydrolysis of anthocyanins, 262–263
- Hydroxytyrosol in olive leaves, 226–227

I

- IFIC, *see* International Food Information Council Foundation
- IFT, *see* Institute of Food Technologist
- Immunoassay, for phytoestrogen measurements, 65–68
- INAMVP, *see* Nutraceutical Advancement's Methods Validation Program
- Infrared spectroscopy (IR), 101
- Institute of Food Technologist, 468
- Institute of Medicine's Food and Nutrition Board, 467
- Integrated pulsed amperometric detection, 482
- Integrated voltammetric detection, 482
- International Food Information Council Foundation, 469
- International Union of Pure and Applied Chemistry, 282
- IOM/FNB, *see* Institute of Medicine's Food and Nutrition Board
- Ion-exchange chromatography (IEC),
in amino acids
characteristics, 460
chromatogram of, 458
pKa and pH of, 457–458
uses of, 456
- IPAD, *see* Integrated pulsed amperometric detection
- Isocratic elution systems, in phytoestrogens analysis, 33–34
- Isoflavones compounds, 151; *see also* Plant flavonoids, chemistry of
biosynthesis of, 9–11
chromatographic properties of, 22
CZE separation of, 61–63
electrodriven separation of, 64–65
extraction of, 15–16
gas chromatography in, 24–25
health benefits of, 3–5
HPLC analysis of, 33–36
LC-MS technique for, 39–47
MALDI-TOF-MS method for analysis, 54
metabolism of, 13–14
nomenclature of, 9
RP-HPLC analysis of, 36–37
separation by CE method, 60–61
sources of, 5–6
spectrum of, 7–8
- Isotope dilution gas chromatography-mass spectrometry (ID-GC-MS-SIM), 25–28
- Isotope dilution (ID), 15
- IUPAC, *see* International Union of Pure and Applied Chemistry
- IVD, *see* Integrated voltammetric detection

K

Kloeckera brevis, 409, 413–414

L

- Lactobacilli arabinosus*, 412, 414
- Lactobacilli plantarum*, 412, 414
- Lactobacillus casei*, 409, 412, 414
- Lactobacillus fermentum*, 409
- Lactobacillus lactis*, 417
- Lactobacillus leischmannii*, 417
- Laser-induced fluorescence (LIF), 64
- LC-PAD, *see* Liquid chromatography-pulsed amperometric detection
- Leuconostoc mesenteroides*, 409, 412
- Lignans compounds
analysis of, 54–59
biosynthesis of, 11–13
extraction of, 17–18
gas chromatography in, 24–25
health benefits of, 3–4
HPLC analysis of, 41–46
ionspray mass spectrometry of, 59–60
metabolism of, 13–14
nomenclature of, 8
sources of, 5
- Lignin in grapevine, 220–221; *see also* Polyphenolic compounds
- Linoleic acid (LA), 86
- α -Linolenic acid (ALA), 86
health benefits of, 89
- γ -Linolenic acid (GLA), 92
foods with, 99–100
- Lipoproteins, 291
- Liquid chromatography (LC), 15
- Liquid chromatography-mass spectrometry (LC-MS), 39–40
- Liquid chromatography-nuclear magnetic resonance (LC-NMR), in flavonoids detection, 189, 192
- Liquid chromatography-pulsed amperometric detection, 489
- Liquid-liquid extraction (LLE), 162, 167, 230
- Long-chain fatty acid (LCFA), 86
- Low-density lipoprotein (LDL), 23, 86
- Luteolin, 149, 153, 172, 180, 187–188
- Luteolin-7-glucoside, health benefits, 226–227

M

- Maceration extraction method, 229
- Mass spectrometry (MS)
in anthocyanins separation, 265–269
in flavonoids detection, 186–189

- Matairesinol, structure of, 7
- Matrix-assisted laser desorption/ionization (MALDI), 186, 266
- Matrix-assisted laser desorption-ionization time-of-flight mass spectrometry (MALDI-TOF-MS), 54
- Matrix solid-phase dispersion (MSPD), 20–21
- Matrix solid-phase extraction (MSPE), 167
- Mediterranean diet, health benefits of, 208
- Medium-chain fatty acid (MCFA), 86
- Menopause symptoms and phytoestrogens, 3
- Mental health and fatty acid, 89
- Methionine
 - description of, 446
 - performic acid oxidation of, 446–447
- Methyloxime–trimethylsilyl (MO-TMS), in anthocyanins identification, 263
- 4-Methyl-1,2,4-triazoline-3,5-dione (MTAD), 124
- Micellar electrokinetic chromatography (MEKC)
 - in anthocyanins separation, 262
 - in flavonoids separation, 178
 - in phytoestrogens determination, 64–65
- Microencapsulated marine oils, 95
- MLCCC, *see* Multilayer coil countercurrent chromatography
- Monarda didyma*, 252
- Monosodium glutamate, 449
- Monounsaturated fatty acids (MUFA), 86
- MSG, *see* Monosodium glutamate
- Multilayer coil countercurrent chromatography, 258
- Multiple electrochemical detection technique, 39
- Multiple reaction ion monitoring (MRM), 51
- N**
- Naringenin, 150–152, 187–189
- National Health and Nutrition Examination Survey (NHANES), 36
- National Institute of Standards and Technology, 477
- National Institutes of Health, 477
- Near-infrared spectroscopy, 477
- Neohesperidin dihydrochalcone, 156
- Neurospora crassa*, 409, 414
- NHDC, *see* Neohesperidin dihydrochalcone
- Niacin, 411–412
- Nicotinamide, 411
- Nicotinic acid, 411
- NIH, *see* National Institutes of Health
- NIR, *see* Near-infrared spectroscopy
- NIST, *see* National Institute of Standards and Technology
- Nitzschia closterium*, 366
- NMR, *see* Nuclear magnetic resonance spectroscopy
- NMR spectroscopy, in conjugated linoleic acid, 124
- N,N*-dimethylformamide (DMF), 16
- N,O*-bis-(trimethylsilyl)-trifluoroacetamide, 24, 173, 233, 263
- Noble metal electrodes, 483
- Nonstarch polysaccharides, 474, 476–477; *see also* Dietary fiber
 - analysis, 480–481, 506
 - definition, 474
 - and dietary fiber, 476–477
- Norbixin, 294
- Novasoy daily, health benefits of, 4
- NSP, *see* Nonstarch polysaccharides
- N-(tert-butyl)dimethylsilyl*-*N*-methyltrifluoroacetamide, 28, 173
- Nuclear magnetic resonance spectroscopy, 477
 - in anthocyanins identification and separation, 250, 270
 - in flavonoids detection, 189–191
 - in lipid analysis, 116
- Nuclear overhauser effect spectroscopy (NOESY), 191
- Nutraceutical Advancement's Methods Validation Program, 478
- Nutraceuticals, 468
- O**
- OBPs, *see* Olive biophenols
- Ochromonas danica*, 409, 414
- O*-desmethylangolensin (*O*-DMA), 7
- O*-glycosylated, 149, 156
- OIW, *see* Olive oil industry wastes
- Oleic acid (OA), health benefits of, 93
- Oleuropein in olive leaves, 226–227
- Oligomeric procyanidins, 224–225
- Olive biophenols, 209
 - characteristics of, 226–228
 - extraction of, 228–233
- Olive oil, health benefits of, 209
- Olive oil industry wastes, 232
- Olive trees and olive oil production residues
 - biphenols from, 226–228
 - olive leaves, 228–231
 - olive oil industry wastes, 231–233
- OPCs, *see* Oligomeric procyanidins

- P**
- PAD, *see* Pulsed amperometric detection, in functional foods
- Palmitoleic acid, health benefits of, 93
- Paper chromatography (PC), in anthocyanins separation, 253–254
- Partially hydrogenated vegetable oils (PHVO), 86
- PCD, *see* Pulsed coulometric detection
- PDBA, *see* p-dimethylaminobenzaldehyde
- Pectin, 507
- PED, *see* Pulsed electrochemical detection, in functional foods
- Phaseolus vulgaris* L., 172
- Phenolic acids in grapevine, 217–218; *see also* Polyphenolic compounds
- L-Phenylalanine ammonia-lyase (PAL), 11
- Phenylpropanoid pathway and isoflavonoid biosynthesis, 11
- Pheophytin
absorption coefficients of, 379
degradation of, 359
electronic absorption spectra, 353–355
thin layer chromatography, 371–376
uses of, 347
- Pheophytinization, 343, 351, 360
- Photodiode array (PDA), 47
- Photosynthetic cells, 342
- Photosynthetic organisms, 342
- Phycomyces blakeslekanus*, 409
- Phytoestrogens
bioassays, 69
biosynthesis of, 9–13
capillary electrophoresis, 60–63
chromatographic separation in, 21–36
detection in, 36–39
health benefits of, 3
high-performance capillary electrophoresis, 63–64
immunoassay for measurement of, 65–68
liquid chromatography–mass spectrometry for
ionspray mass spectrometry of
lignans, 59–60
isoflavones, 39–54
lignans, 54–59
metabolism of, 13–14
micellar electrokinetic chromatography, 64–65
products of, 4–5
sample preparation and extraction in, 14–19
sample treatment, 19–21
sources of, 5–6
structural features and chemistry of, 6–9
types of, 2–3
- Picea abies*, 5
- Pico-Tag, 456
- Plant flavonoids, chemistry of, 148–156
- Plant food and nutraceuticals, flavonoids in
food processing and storage effect, 160–161
intake of, 161
occurrence of, 156–160
- Plasma desorption mass spectrometry (PDMS), 266
- PLC, *see* Preparative layer chromatography
- PLE, *see* Pressurized liquid extraction technique
- Polygonum cuspidatum*, 225
- Polyphenolic compounds
from grape pomace, 221
health benefits of, 209
from ligneous organs of vine, 225
from vine shoots and leaves, 221–222
from vineyard and wine production residues
flavonoids, 210–217
nonflavonoids, 217–221
- Polyunsaturated fatty acids (PUFA), 86
- Potential sweep-pulsed coulometric detection, 482
- Prebiotic foods, 467
- 8-Prenylaringinin, sources and beneficial health effects, 5
- Preparative layer chromatography, 172
- Pressurized liquid extraction technique, 223–224
- Pressurized solvent extraction (PSE), 16
- Proanthocyanidins, 155–156; *see also* Plant flavonoids, chemistry of
- Probiotic foods, 468
- Procyanidins in grapes, 216
- Protein amino acids
acid hydrolysis of, 443–445
gas-phase hydrolysis, 444
properties of, 442–443
- Provitamin A
conversion factors, 325
retinol equivalency, 324–325
- Prussian blue assay, in flavonoids analysis, 171
- PS-PCD, *see* Potential sweep-pulsed coulometric detection
- Pueraria* sp., 49
- Pulsed amperometric detection, in functional foods
for aldehyde and alcohol compounds
chronoamperometric analysis, 488, 490
detection potential, 487
pulsed voltammetry analysis, 491–492
quadruple-potential waveform, 487–488
signal-to-noise ratio, 490
three-step potential-time waveform, 487
for amine and sulfur compounds
current–potential response, 493
IPAD waveform and parameter functions, 497
pulsed voltammetric analysis, 494

cation-exchange chromatography, 500
 disadvantages of, 496–497
 and HPAEC
 agricultural applications, 510–511
 beverages, 499–501
 food additives and nutriments, 502–504
 fruit juices, 501–503
 industrial applications, 510–511
 noncarbohydrate applications, 511–514
 solid foods, 504–510
 HPLC-IPAD, 498
 Pulsed coulometric detection, 482
 Pulsed electrochemical detection, in functional foods
 advantages of, 514
 aldehyde and alcohol-containing compounds
 cyclic voltammetric analysis, 484–486
 pulsed amperometric detection, 487–492
 amine and sulfur-containing compounds,
 492–495
 integrated pulsed amperometric detection,
 495–499
 pulsed voltammetric analysis, 494
 ED in HPLC, 482
 in food industry, 481–482
 history, 499
 HPLC-IPAD, 498
 IPAD waveform and parameter functions, 497
 noble metal electrodes
 anodic detection modes, 485–486
 cyclic voltammetric analysis, 484
 using PAD
 for aldehyde and alcohol compounds,
 488–491
 for amine and sulfur compounds,
 492–495
 disadvantages, 495–496
 Pulsed voltammetry (PV), 491
 Pyridoxine, 413–414

Q

Quercetin, 150–152, 180, 187–189

R

Radioimmunoassay (RIA), 23, 66–68
Radix astragalus, 20
Raphanus sativus, 270
 Reconstructed ion chromatograms (RIC), 126
 Resistant starch, 475
 Resveratrol, sources and beneficial health effects, 5

Retinol activity equivalent (RAE)
 in β -carotene, 290–291, 324–325
 in provitamin A, 324–325
 Retro-Diels-Alder (RDA), 173
 Reversed-phase chromatography (RPC),
 in amino acids
 chromatogram of, 461
 hydrophobic and hydrophilic characteristics of,
 460–461
 principle of, 460
 Reversed-Phase High-Performance Liquid
 Chromatography, 152, 174–175,
 180, 189
 in CLA fraction isolation, 127
 in fatty acids analysis, 116
Ribes nigrum, 252, 258
 Riboflavin, 409–411
 RP-HPLC, *see* Reversed-Phase High-Performance
 Liquid Chromatography
 RS, *see* Resistant starch
Rubus idaeus L., 258

S

Saccharomyces uvarum, 413
Sambucus nigra, 258
 Saponification, 298
 Saturated fatty acids (SFA), 86
 Secoisolariciresinol diglucoside (SDG), 13
 Secoisolariciresinol, structure of, 7
 Sefcal process, in grape pomace extraction, 222
 Selected-ion monitoring (SIM), 173
 Selected-ion recording (SIR), 24–25, 126
 Sephadex gel column chromatography, 256
 SEWE, *see* Superheated ethanol–water extraction
 SFE, *see* Supercritical fluid extraction
 Shikimic acid, 149–150
 Shonanin, acid hydrolysis of, 18
 Short-chain fatty acid (SCFA), 86
 derivation of, 107–108
 methods for analysis, 112
 Silver-ion HPLC
 in CLA separation, 120–121
 in fatty acids analysis, 116
 Simple carbohydrates, 472
 SLE, *see* Solid-liquid extraction
 Sodium-copper-chlorophyllin, 361
 Sodium dodecyl sulfate (SDS), 178
 Solid-liquid extraction, 223
 Solid-phase extraction, 162, 167–168
 in anthocyanins separation, 256–257
 in biophenol identification, 230
 in phytoestrogen isolation, 19
 Solid-phase microextraction, 21, 167–168

- Soxhlet extraction method, 104, 229, 232
 SPE, *see* Solid-phase extraction
 SPME, *see* Solid-phase microextraction
 Starch
 analysis, 480, 506
 modified, 476
 NSP, 476–477
 polysaccharides, 474
 resistant, 475
 Static–dynamic superheated extraction method, 229–230
 Stearidonic acid (SA), health benefits of, 93
 Stilbenes in grapevine, 218–220; *see also*
 Polyphenolic compounds
Streptococcus faecalis, 415
 Subcritical water extraction, 169
 Sulfolane, 453
 Supercritical fluid chromatography (SFC), 16, 116
 Supercritical fluid extraction, 103, 168–170, 223
 Superheated ethanol–water extraction, 169–170
 SWE, *see* Subcritical water extraction
 Sweeteners, 505
- T**
- Tangeretin, 149, 153
 TBDMS, *see* *N*-(*tert*-butyldimethylsilyl)-*N*-methyltri-fluoroacetamide
 TDP, *see* Thiamin diphosphate
 Tetrahydrofuran (THF), 460
Tetrahymena pyriformis, 409
 Tetramethylguanidine (TMG), 106
 TFA, *see* Trifluoroacetic acid
 Theoretical correction factor (TRF), 111
 Thermospray mass spectrometry (TSP-MS), 37
 Thermospray (TSP)-LC-MS technique, 40
 Thiamin
 analytical methods, 408–409
 constituents of, 403, 408
 diphosphate, 408
 pyrophosphate, 408
 triphosphate, 408
 Thiaminases, 408
 Thiamin-degrading enzymes, 408
 Thin-layer chromatography (TLC)
 in anthocyanins separation, 251, 254–255
 in flavonoids separation, 172–173
 in isoflavonoids identification, 21–22
 Thylakoid membrane, 342
 Time-resolved fluoroimmunoassay (TR-FIA), 68
 TMAO, *see* Trimethylamine *N*-oxide
 TMAU, *see* Trimethylaminuria
 TMCS, *see* Trimethylchlorosilane
 Total lipids, extraction of, 102–104
 TPP, *see* Thiamin pyrophosphate
 Trans fatty acids, 92
 GC analysis for, 117–118
 IR and FTIR spectroscopy, 118–119
 Trifluoroacetic acid (TFA), 15, 251
 2,4,4'-trihydroxydeoxybenzoic acid (THB), 14
 Trimethylamine *N*-oxide, 349
 Trimethylaminuria, 349
 Trimethylchlorosilane, 233, 263
 Trimethylsilyldiazomethane (TMS-DM), 106
 Trimethylsilyl ether (TMS), 28, 173
 TRIONE, 453
 Tryptophan
 acid hydrolysis, 444–445
 alkaline hydrolysis, 445–446
 TTP, *see* Thiamin triphosphate
 Tyrosol in olive leaves, 226–227
- U**
- Ultrasound-assisted extraction method, 229
Ultraviolet (UV) absorption spectra, for
 isoflavonoids, 7–9
 United States Department of Agriculture, 471
 United States Pharmacopoeia, 478
 Unsaturated fatty acids (UFA), 86
 USDA, *see* United States Department of
 Agriculture
 U.S. Institute of Medicine (IOM), 291
 USP, *see* United States Pharmacopoeia
 UV–Vis adsorption spectroscopy, in anthocyanins
 separation, 264–265
 UV–visible (UV–Vis), in flavonoids detection,
 180–182, 193
- V**
- Vaccinium myrtillus*, 252
 Vanillin assay, in flavonoids analysis, 170–171
 Verbacoside in olive leaves, 226–227
 Very low-density lipoprotein (VLDL), 88
Vicia villosa, 256
 Vine shoots and leaves, uses of, 221–222
 Vineyard and wine production residues
 extraction of, 222–225
 flavonoid polyphenols from, 210–217
 nonflavonoid polyphenols from, 217–221
 uses of, 221–222
 Vitamin analytical methods, classification of,
 402–403
 Vitamin B₁₂, 416–417
 Vitamin C, 417–420
Vitis vinifera, 211

W

Water-soluble vitamins, 404–407

 high-performance liquid chromatography
 methods, 404–407

 biotin, 413–414

 folic acid, 414–415

 niacin, 411–412

 pyridoxine, 412–413

 riboflavin, 409–411

 thiamin, 403–404, 408–409

 vitamin B₁₂, 416–417

 vitamin C, 417–420

Women's Health Initiative (WHI), 2

World Health Organization (WHO),
289, 324