

# Subject Index

- A-1 broth 245  
 4ABC-MRS medium 209  
 Abeyta-Hunt-Bark agar 415  
 absolute growth index 978  
 acetamide broth 496, 499  
 acetic acid dichloran yeast extract  
   sucrose (ADYS) agar 568  
 acidic tomato broth (ATB) 182,  
   413  
*Acinetobacter* spp. 484–5  
   *A. anitratus* 501  
   *A. calcoaceticus-baumannii* 485,  
     498  
   *A. haemolyticus* 485  
   *A. lwoffii* 485, 501  
 enrichment media 495–6, 497  
   *see also Pseudomonadales*  
 aerobic endospore-forming  
   bacteria 92–110  
   acidophilic and  
     acidotolerant 106–8  
   culture media 98–9  
   detection/enumeration 95–8  
   mesophilic 102–6  
   physiological groups 99–110  
   relevance to food 94–9  
   thermophilic and thermotolerant  
     108–9  
 aerobic low peptone (ALP)  
   media 496, 500, 501  
*Aerococcus* spp. 176  
*Aeromonadales* 483  
*Aeromonas* spp. 451–81, 453, 456, 654  
   basal media 457–8  
   culture media 470  
   diagnostic systems 460–2  
     carbohydrates 460–1  
     DNA 461  
      $\beta$ -haemolysis 461  
     Tween 80 461  
   enrichment media 467–8  
   in food 455, 457  
   identification 454–5, 456  
   incubation atmosphere 462  
   incubation temperature 462  
   isolation from foods 468–70  
   plating media 463–7  
   selective media 458–60  
     antibiotics 458–9  
     antiseptics 459–60  
     bile salts 460  
     dyes 458  
     surfactants 458  
     vibriostatics 460  
   standard methods 470  
   taxonomy 452–4  
   water testing 614–15  
 aesculin bile azide agar 156, 162,  
   163–4, 166  
 agar *Listeria* according to Ottaviani  
   and Agosti *see* ALOA  
*Agrobacterium* spp. 483  
*Alcaligenaceae* 483  
 ALI medium 107, 108  
*Alicyclobacillus* spp. 91, 92, 95, 96,  
   100  
   culture media 107, 109  
   *see also* aerobic endospore-forming  
     bacteria  
*Alkalibacillus* spp. 92, 101

- alkaline peptone water (APW)  
381, 383, 384, 452, 462, 467,  
468
- alkaline peptone water polymyxin  
(APWP) 383
- alkaline peptone water salt  
(APWS) 383
- all purpose Tween agar (APT) 176,  
230, 663–5
- Alloiococcus* spp. 176
- ALOA 115, 120, 121, 124,  
658–62
- Alternaria* spp. 569, 570
- Alteromonadales* 483
- ALVC medium 492
- American Association of Analytical  
Chemists *see* AOAC
- Ammoniphilus* spp. 92
- Amphibacillus* spp. 92
- ampicillin dextrin agar (ADA)  
464, 466, 615
- ampicillin dextrin broth (ADB)  
467, 468
- Anacker and Ordal agar 512, 513,  
514
- Aneurinibacillus* spp. 92, 100
- Anoxybacillus* spp. 92
- AOAC standards 4  
*Enterobacteriaceae* 252  
*Salmonella* spp. 262
- apple pulp gelatin medium 570
- Arcobacter* spp. 403–50, 404, 405,  
429, 654  
characteristics of 407–8  
isolation media 428–9, 430
- Arroyo, Martin and Cotton (AMC)  
medium 208, 215
- Arthrobacter* spp. 195, 483
- ASM agar 430
- Aspergillus* differential medium  
567
- Aspergillus flavus* and *parasiticus* agar  
(AFPA) 567, 666–8
- Aspergillus* spp. 559, 565, 567, 568,  
577
- Atopobium* spp. 176
- Bacillus* spp. 90, 95, 96, 100, 101, 106  
*B. cereus* 90, 95, 96, 100, 101,  
102–5, 654  
food poisoning 102  
group members 103–4  
stress damage 41  
*B. coagulans* 95, 96, 100  
culture media 109  
*B. laevolacticus*, culture media  
109  
*B. licheniformis* 72, 95, 96, 100,  
106  
*B. racemilacticus*, culture  
media 109  
*B. subtilis* 95, 96, 100, 105–6  
group members 106  
stress damage 42  
culture media 90–114  
relevance to food 94–9  
systematics 91–4  
*see also* aerobic endospore-forming  
bacteria
- Bacillus acidoterrestris* medium 107
- Bacteriological Analytical Manual*  
(BAM) 262, 266, 271
- Campylobacter* spp. 424
- Escherichia coli* 345
- Salmonella* spp. 262
- Shigella* spp. 289
- V. cholerae* 382
- Yersinia* spp. 311
- Baird-Parker agar 53, 135, 136–7,  
669–73
- Baird-Parker liquid (LBP)  
medium 135, 142, 674–6
- basal agar-maltose 184
- basal agar-sorbitol 184
- BBL Crystal Enteric/Nonfermenter  
(E/NF) system 370
- Beerens medium 210–11
- Bergeyella* spp. 509, 517  
*B. zoohelcum*, fatty acid methyl  
esters 543  
characteristics 515–16  
colony morphology 538–9  
culture media 511

- Bif medium 209, 217
- bifidobacteria selective medium (BSM) 210, 211
- Bifidobacterium* spp. 184, 199–227
- elective media 202–3
  - non-culture-dependent enumeration 220
  - selective media 203–19
    - enumeration from fermented milk products 212–19
    - recovery from natural habitats 205–12
    - stressed cells 219–20
    - taxonomy 201–2
- bile aesculin agar (BAA) 612, 677–9
- bile aesculin azide agar (BAAA) 612
- bile oxalate sorbose (BOS) broth 298, 302, 305
- bile salts brilliant green agar (BGA) 464, 471
- bile salts brilliant green starch agar (BBGS) 464, 466
- bile salts Irgasan brilliant green (BSIBG) agar 452, 464, 466, 470, 680–2
- modified 464, 466–7
- BIM-25 medium 205, 206, 210
- bioluminescent ATP assays 575–6
- Biosynth<sup>®</sup> culture medium (BCM) O157:H7 336
- BioSys optical assay 249
- bismuth sulfite agar 274, 275, 276, 683–6
- Blaser A medium 412
- BlaserWang A medium 412
- blood agar 423
- with ampicillin (BAA) 463, 464, 468
  - with vancomycin, cefixime and cefsulodin (BVCC) 343
- blood liver (BL)-OG medium 209, 216–17
- blood media 416
- Bolton broth 413, 415, 424, 426, 427, 687–9
- boric acid lactose broth 236
- Botrytis* spp. 570
- B. cinerea* 570
- brain heart infusion (BHI) agar 157, 203, 512, 514, 518
- with Isovitalex 433
  - with yeast extract + FCS 433
- Brevibacillus* spp. 92, 95, 100
- Brevibacterium* spp. 195
- Briggs agar 176, 690–2
- Brilliance Salmonella agar 274, 275
- brilliant green 266, 269, 458
- and skim milk 267
- brilliant green agar (BGA) 265, 274, 275, 276
- modified 866–9
- brilliant green bile (BGB) broth 236, 244–5, 250, 251, 693–5
- brilliant green bile lactose broth (BRILA) 244, 326, 330
- brilliant green water 267
- Brochothrix* spp. 195
- B. thermosphacta* 195, 654
- bromothymol blue Teepol (BTBT) agar 384, 386
- Brucella* broth/agar 431, 433
- Bryant Burkey lactose broth 73
- BS-LV medium 207, 214
- BS1-neomycin sulfate, paromomycin sulfate, sodium propionate and lithium (NPPL) medium 205
- BU40 B medium 412
- buffered cefamandole polymyxin anisomycin  $\alpha$ -ketoglutarate (BMPA $\alpha$ ) agar 597, 599, 696–7
- buffered charcoal yeast extract (BCYE) agar 597, 599, 698–700
- buffered peptone water (BPW) 51, 52, 117, 265, 267
- additives 266, 267
  - with casamino acids 330
  - modified, with maltol, cefsulodin, vancomycin and novobiocin 326

- modified with pyruvate, with acriflavine, cefsulodin and vancomycin 326, 332
- with vancomycin, cefixime and cefsulodin 326
- Burkholderia cepacia* 501
- Burkholderiales* 483
- Butzler agar 412, 426, 427
- Butzler medium Oxoid agar 412, 419
- Butzler medium Virion agar 410, 412, 419
- Byssochlamys* spp. 566
- C-2 A medium 412
- C-2 B medium 412
- cadaverine, lysine, ethylamine and nitrate (CLEN) agar 572
- CAK agar 413
- Caldalkalibacillus* spp. 92
- Campy BAP agar 412, 415, 422
- Campy Cefex agar 414, 426
- Campy-CVA agar 406
- Campy-FDA medium 426
- Campylobacter* enrichment broth (CEB) 413
- Campylobacter* spp. 403–50, 403, 405, 654
  - C. jejuni* 403
    - stress damage 44
    - VBNC 55
  - diagnostic features 406
  - enrichment media 412–14, 415, 422
    - incubation atmosphere 425
  - freeze-injured 422–4
  - isolation from faeces 427
  - isolation from foods 426–7
    - standard methods 427–8
  - isolation from surfaces 424
  - membrane filter methods 420–2
  - plating media 418–20
    - storage 416–17
  - selective media 405–28
    - antibiotic activity 417–18
    - antibiotic choice 417
    - basal media 415
      - blood 416
      - ferrous sulfate, sodium metabisulfite and sodium pyruvate (FBP) 416
      - pH 417
      - water testing 605, 616
- Campylosel medium 415, 434
- Candida* spp. 571–2
- CAR broth/agar 413
- Carnobacterium* spp. 175, 176, 181, 654
  - culture media 180–1
- Caulobacterales* 483
- cefixime tellurite sorbitol MacConkey (CT-SMAC) agar 336, 337, 341, 701–3
- cefoperazone amphotericin teicoplanin (CAT) agar/broth 414, 429, 430, 704–9
- cefoperazone cefsulodin Irgasan novobiocin (CIN) agar 298, 305–6, 429, 459, 464, 710–12
  - modified 430
- cellobiose polymyxin B colistin (CPC) agar 386, 389, 391–3, 713–15
- cellobiose-colistin (CC) agar 241
- cephaloridine fucidin cetrimide (CFC) agar 491–2, 494, 615, 716–18
- cephalothin alkaline peptone water (CAPW-10) 467, 468
- cephalothin buffered dextrin broth (BCDB-10) 467, 468
- Cerasibacillus* spp. 92
- CETCH medium 492
- cetrimide agar 490, 493
- charcoal cefoperazone deoxycholate (CCD) agar 413, 415, 722–4
  - modified (mCCD) 413, 415, 419, 420, 426, 428, 434, 719–21
- charcoal yeast extract (CYE) agar 598–9
- chi-squared test 12
  - distribution-free 17–18
- chitin salt agar 514
- Christopher B + A medium 412
- CHROMagar™ *Acinetobacter* 496

- CHROMagar™ Candida** 573  
**CHROMagar™ O157** 336, 338,  
 341–2  
   with potassium tellurite 336, 338  
   with potassium tellurite and  
   novobiocin 336, 338  
**CHROMagar™ Salmonella** 278–9  
**CHROMagar™ Staph. aureus** 140  
**chromID™ 0157:H7 agar** 336  
**ChromoCult™ Agar** 339  
**ChromoCult™ coliforms agar** 241,  
 248  
**ChromoCult™ ESA** 365–6, 368  
 chromogenic *Cronobacter* isolation  
 (CCI) agar 368  
 chromogenic esterase agar 279  
 $\alpha\beta$ -chromogenic medium 274, 275,  
 276, 278  
 chromogenic salmonella esterase 274,  
 275, 276  
 chromogenic *Shigella* spp. plating  
 medium (CSPM) 294–5  
 chromogenic *Yersinia* spp. plating  
 medium 306  
 chromogenic/fluorogenic media  
 140–1, 623–46  
   hydrolysis 632  
   induction 628–9  
   nomenclature 624–6  
   sugars 624, 625  
     glycones and aglycones  
     626–8  
   synthetic substrates 632–40  
     chromogens 635–7  
     fluorogens 633–5  
     trojan horses 637–40  
   transportation 629–32  
   water testing 611  
 chromogens 635–7  
*Chryseobacterium* spp. 501, 509,  
 517–26  
   characteristics 515–16  
   colony morphology 539–40  
   culture media 511–12, 518  
   fatty acid methyl esters 543  
   pigment production 539  
   sources of  
     beer-bottling plants 521  
     dairy 518–19  
     edible plants 521, 526  
     fish 519–20  
     meat and poultry 520  
     water 521  
*Chryseomonas luteola* 501  
*Chryso sporium* spp. 565  
 citrate azide Tween carbonate  
 (CATC) agar 156, 162, 164, 166,  
 725–7  
*Citrobacter* spp. 234  
   characteristics 308  
*Cladosporium* spp. 569  
*Clostridium* spp. 67–8, 72  
   *C. baratii* 654  
   *C. botulinum* 66, 67  
     culture media 79–81  
   *C. botulinum* isolation agar 80  
   *C. butyricum* 72, 654  
   *C. difficile* 66, 67  
     culture media 81–2  
   *C. perfringens* 67, 654  
     culture media 74–6  
       liquid 77  
     stress damage 41  
     water testing 605, 607, 612–13  
   *C. thermosaccharolyticum* 655  
     culture media 66–89  
     isolation of 67–9  
     sulfite-reducing species  
       culture media 70–4  
       water testing 612–13  
 coagulase-positive staphylococci  
 confirmation and rapid  
 identification 143–4  
 culture media 133–43  
   liquid 141–3  
   solid 134–41  
   thermonuclease assay 144–5  
 coconut cream agar (CCA) 568  
*Cohnella* spp. 92  
 Coli ID medium 241  
 coli-aerogenes group 655  
 ColiComplete medium 241, 243, 252

- Coliform Count medium 241  
coliforms 233–60  
  detection/enumeration 240–2  
  faecal 235, 608–10  
  total 235  
  water testing 605, 607, 608–10  
Colilert<sup>®</sup> 18, 241, 607, 610  
Colisure<sup>®</sup> 241  
colony counting  
  comparison of differences 17–18  
  liquid media 7–8, 22–30  
  productivity ratio *see* productivity ratio  
  solid media 6–7, 8–21  
  statistics 9–16  
Columbia agar 431, 433, 434  
  modified 203  
Compact Dry *E. coli* medium 241  
Compact Dry Yeast and Mold plates 574–5  
Compass *Salmonella* agar<sup>®</sup> 278  
confidence intervals 12  
confidence limits 10, 11  
Congo red brain heart infusion agarose 307  
cooked meat medium 80–1  
creatine sucrose agar 569  
cresol red thallium acetate sucrose (CTAS) agar 180, 728–30  
cresol red thallium sucrose inulin agar 180, 728–30  
CHROMID *Salmonella* agar 274  
*Cronobacter* broth 363, 364  
*Cronobacter* screening broth 360, 363, 364  
*Cronobacter* spp. 234, 358, 371  
  confirmation 370–2  
  enrichment 362–3  
  isolation and enumeration 363–70  
    chromogenic media 368  
    fluorogenic media 369  
  pre-enrichment 359–62  
  *see also Enterobacter* spp.  
*Cryptococcus neoformans* 572  
*Cryptosporidium* spp., water testing 617  
CVA agar 430  
cycloserine cefoxitin fructose agar 81–2  
cycloserine hydrochloride, moxalactam and norfloxacin agar and broth 82  
cys-MAR medium 204, 213  
Czapek yeast agar 559, 567  
Czapek–Dox agar 569  
  
de Man, Rogosa and Sharpe *see* MRS *Debaryomyces hansenii* 565, 573  
Dekeyser A medium 412  
*Dekkera* (*Brettanomyces*) *bruxellensis* 573, 577  
*Delftia acidovorans* 501  
deoxycholate agar 238  
deoxycholate citrate agar (Leifson agar) 274, 275, 276, 292  
deoxycholate citrate lactose agar 274, 275, 276  
deoxycholate hydrogen sulfite lactose agar 238, 239  
*Desulfotomaculum nigrificans* 72, 655  
dextrin fuchsin sulfite (DFS) agar 464  
diagnostic *Salmonella* selective semisolid medium (DIASALM) 731–4  
DIASALM 269, 272, 731–4  
dichloran chloramphenicol peptone agar (DCPA) 569  
dichloran creatine (CREAD) agar 569  
dichloran glycerol (DG18) agar 558, 565, 735–7  
dichloran rose bengal chloramphenicol (DRBC) agar 564, 738–40  
dichloran rose bengal yeast extract sucrose (DRYES) agar 568  
dicloxacillin medium 217–18  
Difco heart infusion agar with diamide 492  
differential clostridial agar 70, 71, 73–4, 741–3

- differential reinforced clostridial agar 73  
 dilution error 9  
 DIN standards, *Escherichia coli* 346  
 direct viable count 55  
 disaccharides 625  
 distribution error 9–10, 12  
 DNA stress damage 42–3  
 DNA-containing media 461  
 DNAase ampicillin agar (DNAA) 464  
 dormancy 55  
 Doyle-Roman B medium 412, 413  
 Drake's medium 489, 493  
 Druggan-Forsythe-Iversen formulation (DFI) 365–6, 368  
   modified 368  
 Dynabeads anti-*E. coli* procedure 332  
  
 EBRER medium 181  
 EC broth 236, 245, 250  
 EC-MUG broth 236  
 ecometric streaking method 7  
*Edwardsiella* spp. 234  
 EHEC *see also Escherichia coli*, enterohaemorrhagic  
 EHEC agar 336, 339–40, 343  
 EHEC enrichment broth with vancomycin, cefixime and cefsulodin 326  
 ELISA 344, 576–7  
*Elizabethkingia meningoseptica* 517  
 EMJH P80 broth 428, 430  
*Empedobacter brevis* 509, 526  
   characteristics 515–16  
   culture media 512  
   fatty acid methyl esters 543  
   pigment production 539  
 Endo agar 238, 250, 458  
 enriched tomato juice broth 182  
*Enterobacter sakazakii* agar (ESA) 365–6  
*Enterobacter sakazakii* enrichment broth (ESEB) 363, 364, 369  
*Enterobacter sakazakii* isolation agar (ESIA) 365–6, 368  
  
*Enterobacter sakazakii* plating medium (ESPM) 369  
*Enterobacter sakazakii* screening plate (ESSP) 370  
*Enterobacter sakazakii* selective broth 362–3  
*Enterobacter* spp. 234, 358, 371  
   characteristics 308  
   *see also Cronobacter* spp.  
*Enterobacteriaceae* 234–5  
   culture media 235  
   ISO/AOAC recommendations 252  
   liquid 235–7  
   solid 237–40  
 detection methods 242–9  
   comparative studies 244–9  
   incubation conditions 243–4  
   membrane methods 242–3  
   “total” 233–60  
   *see also* coliforms; *Escherichia coli*; *Shigella* spp.  
*Enterobacteriaceae* enrichment (EE) broth 236, 244, 245, 250, 251, 290, 330, 332, 362, 364, 744–6  
 Enterococcosel (ECS) agar/broth with/without vancomycin 156, 162, 747–50  
*Enterococcus* spp. 156, 157, 159, 176, 655  
   *E. avium* 159  
     in food 157  
   *E. casseliflavus* 156, 157  
     in food 157  
   *E. durans/hirae* 156, 157, 159  
     in food 157  
   *E. faecalis* 156, 157, 159  
     in food 158  
     VBNC 55  
   *E. faecium* 156, 157  
     in food 158  
   *E. gallinarum* 156, 157, 159  
     in food 157  
   *E. mundtii* 156, 159  
     in food 157  
   glycopeptide-resistant 156

- growth factors 161
- indicators 161
- selective cultivation 155–73
  - confirmatory tests 167–9
  - selectivity and productivity 166
  - storage of ready-to-use plates 167
  - sublethally damaged cells 167
- vancomycin-resistant 156, 157
- water testing 610–12
- enzyme-linked immunosorbent assay *see* ELISA
- eosin methylene blue agar 238, 239–40, 292
  - with sorbitol and novobiocin (MEMB) 342
- Erwinia* spp. 234
- ESA *see* *Enterobacter sakazakii* agar
- Escherichia* spp. 234
  - characteristics 308
- Escherichia coli* 233–60, 321–56, 655
  - detection/enumeration 240, 241–2
  - diffusively adherent (DAEC) 322, 323
  - enteroaggregative (EAEC) 322, 323
  - enterohaemorrhagic (EHEC) 322, 323
  - enteroinvasive (EIEC) 322, 323
  - enteropathogenic (EPEC) 322, 323
  - enterotoxigenic (ETEC) 322, 323
    - as food pathogens 322–3
    - isolation from foods 323–5
    - stress damage 41, 43, 44
  - VBNC 55
  - VTEC 321
    - acid treatment 329
    - confirmation and identification 344–5
    - enrichment 325–8, 330–5
    - immunocapture 328–9
    - isolation from foods 324–5
    - plating media 335–44
    - standardised methods 345–6
    - stress damage 327
  - water testing 605, 607, 608–10, 616
  - see also* *Enterobacteriaceae*
- Escherichia coli* broth, modified (mEC), with novobiocin 326, 331
- Eupenicillium brifeldianum* 566
- European Normalisation Committee (CEN) 346
- Eurotium* spp. 565
  - E. chevalieri* 559
  - E. herbariorum* 566
- Exeter agar 413
- Exeter broth 413, 415, 423, 426
  - modified 751–3
- faecal coliform agar 238
- faecal coliforms 235
  - water testing 608–10
- FBP (ferrous sulfate, sodium metabisulfite and sodium pyruvate) 416
- FBP-AM B/A medium 412
- FCS *see* foetal calf serum
- FDA *Listeria* enrichment broth 754–6
- FDA standards *see* *Bacteriological Analytical Manual*
- Feeley–Gorman medium 598
- ferrous sulfate, sodium metabisulfite and sodium pyruvate (*see* FBP)
- Filobacillus* spp. 92
- Fisher's Index of Dispersion 11
- Flagellimonas eckloniae* 509, 527, 535
  - characteristics 515–16
  - colony morphology 539
  - culture media 512–13
  - fatty acid methyl esters 543
  - pigment production 540
- Flavimonas oryzihabitans* 501
- Flavobacteriaceae* 508–56, 655
  - chemotaxonomic characteristics 542–5
  - culture media 510–37
  - differentiation 537–45
  - phenotypic characteristics 538–42
    - capnophilic metabolism 541
    - colony morphology 538–9
    - enzyme production 542
    - gliding motility 540–1



- Flavobacteriaceae* (continued)  
 pigment production 539–40  
 salinity requirement 541  
*see also individual members*
- Flavobacterium* spp. 483, 484, 509, 527–32  
 characteristics 515–16  
 fatty acid methyl esters 543–4  
 pigment production 540
- Flexibacter maritimus* medium 514, 534
- flow cytometry 575
- Fluorocult *E. coli* O157:H7 agar 336
- Fluorocult ECD medium 247
- Fluorocult LMX broth 248
- fluorogenic media *see* chromogenic/fluorogenic media
- fluorogens 633–5
- foetal calf serum 433
- Food and Drug Administration *see* FDA
- formate lactose glutamate medium 757–61
- Fraser broth 762–4
- Fructobacillus* spp. 175
- fructose and Tween 80 medium 182
- Fusarium* spp. 569  
*F. sporotrichoides* 577
- $\beta$ -galactosidase 241
- Gassner's agar 238
- gelatin phosphate saline (GPS) broth 382, 383
- gelatin taurocholate tellurite (GTT) agar 386, 389
- Geobacillus* spp. 92  
*G. stearothermophilus* (*see also Bacillus stearothermophilus*) 95, 96, 109
- GH medium 492
- Giardia* spp., water testing 617
- Giolitti-Cantoni broth with Tween 80 (GCBT) 135, 765–8
- glucose salt Teepol broth (GSTB) 382, 383, 384
- glucose salt tellurite crystal violet (GSTC) agar 388
- glucose yeast extract sucrose agar 565
- glucose-blood-liver agar 203
- $\alpha$ -glucosidase 367
- $\beta$ -D-glucuronidase 240, 247
- Glupczynski's medium 432
- glutamate starch penicillin/phenol red (GSP) agar 463, 464, 491, 494  
 modified 491
- glycerol nitrate agar 565
- glycine, vancomycin, polymyxin B cycloheximide (GVPC) agar 597, 599, 769–71
- GN-modified broth 236
- Gould S1 medium 492
- Gracilibacillus* spp. 92  
*G. dipsosauri* 101  
*G. halotolerans* 101
- Gram-negative (Hajna) broth 269, 290–1, 293
- Gram-positive, catalase-positive food spoilage bacteria 193–8  
 identification 195–6  
 selective isolation media 194–5  
 total colony counts 194
- GSP agar *see* glutamate starch penicillin/phenol red agar
- H. pylori* special peptone agar (HPSPA) 432, 433
- haemorrhagic-colitis agar 336
- Hafnia* spp. 234  
 characteristics 308
- Halalkalibacillus* spp. 92
- half-Fraser broth 117
- Hall's medium 247
- Halobacillus* spp. 92, 101
- Harlequin listeria medium 120
- hazard analysis critical control point 123
- heart infusion agar (HIA) 512, 513, 543, 544
- Hektoen enteric (HE) agar 274, 275, 276, 278, 292, 293, 772–4

- Helicobacter* spp. 403, 404, 405,  
431–434  
*H. pullorum* 404, 405, 429, 432,  
434  
detection 429, 430–4  
selective media 431–2, 433  
HHD medium 178  
HiChrome ESA 365–6, 368  
high performance liquid  
chromatography (HPLC) 576  
Holton medium 497  
Horie's arabinose ethyl violet  
broth 383, 384  
horse blood agar 514  
Houf *Arcobacter* selective agar 429,  
775–7  
Houf *Arcobacter* selective  
broth 778–80  
Howard mould count technique 575  
Hungate technique 54  
Hunt-Radle broth 414, 424  
hydrophobic grid membrane filter  
method 242, 252  
yeasts and moulds 574
- Idexx plate count agar 607  
IDF standards  
*Bifidobacterium* spp. 213, 215  
*Lactobacillus* spp. 183  
*Listeria* spp. 122  
*Salmonella* spp. 262, 265  
immunomagnetic separation  
(IMS) 328, 334  
index organisms 234  
infant formulae, *Cronobacter*  
isolation 358–9  
Inhibigen™ 637, 639, 641  
inoculum standardisation 5–6  
inositol brilliant green bile salts agar  
(IBB) 451, 463, 464, 471  
Institute for Reference Materials and  
Measurements (IRMM) 5–6  
International Dairy Federation *see*  
IDF  
International Organization for  
Standardization *see* ISO
- Irgasan 459–60, 491  
Irgasan ticarcillin chlorate (ITC)  
broth 298, 302, 303, 305, 781–3  
iron milk medium 78  
iron sulfite agar 70, 71, 72, 74, 784–6  
ISO 2001a 252  
ISO 2001b 252  
ISO 2005b 252  
ISO 2005c 252  
ISO 6579 262, 263, 264, 271  
ISO 6785 262, 265, 271  
ISO 9001 4  
ISO 17025 4  
ISO 19250 262  
ISO 22964 359  
ISO standards  
*Escherichia coli* 345  
*Salmonella* spp. 262  
*Yersinia* spp. 311  
Iso-Sensitest broth 186  
*Issatchenkia orientalis* 571
- Jeotgalibacillus* spp. 92  
Johnson and Murano agar/broth  
429, 430, 432
- K agar 107, 108  
kanamycin aesculin azide (KAA)  
agar 156, 162, 164–5, 166, 612,  
787–90  
Karmali *Campylobacter* selective  
agar 413, 415, 419, 426, 427, 430,  
791–3  
King's agar A 493  
King's agar B 487, 490, 493  
Kirin Ohkochi Taguchi medium 179  
*Klebsiella* spp. 234  
characteristics 308  
*Kloeckera apiculata* 572  
*Kluyveromyces* spp. 573  
KPL medium 178  
*Kurthia* spp. 195  
KV202 agar 306
- L-S differential (LSD) agar 811–13  
La agar 176

- LAB *see* lactic acid bacteria  
 LAB susceptibility test medium 186  
*Laceyella* spp. 92  
 lactic acid bacteria 656  
   culture media 174–92  
   enumeration of 182–4  
   probiotic 185  
   susceptibility testing 186  
 lactic bacteria differential (LBD)  
   agar 184  
*Lactobacillus* sorbic acid (LaS)  
   agar 178, 794–6  
*Lactobacillus* spp. 175, 176, 184, 656  
   culture media 177–8  
*Lactococcus* spp. 175, 655, 656  
   culture media 181  
 lactose bile brilliant green broth  
   (LBBG) 331  
 lactose broth 236, 267  
   with cellulase 267  
   with papain 267  
   with Tergitol 7 267  
   with Triton X-100 267  
 lactose formate glutamate broth 607  
 lactose sulfite broth 78  
 lactose Tergitol TT7 agar 607  
 lactose tryptone-MUG broth 247  
 lactose TTC Tergitol 7 agar 797–800  
*Lactosphaera* spp. 176  
 lag period, extended 45, 46  
 Lammerding method 430  
 LAMVAB medium 185  
 Lander B medium 412  
 latex agglutination assay 344–5  
 lauryl sulfate (LST) broth 236, 244,  
   245, 250, 251, 252, 364  
   membrane 607, 609  
 lauryl sulfate-MUG broth 245  
 lauryl sulfate-MUG X-gal (LMX)  
   broth 801–4  
 lauryl tryptose (LTB) broth 236, 330,  
   362, 805–7  
 lauryl tryptose MU broth 236  
 LBDC 365–6  
 least significant difference 15–16, 28  
 Leeds medium 497  
*Legionella* spp. 596–604, 656  
   concentration techniques 600–1  
   confirmation tests 601  
   culture media 597, 598–9  
     VBNC 55  
 legionnaires' disease 597  
*Lentibacillus* spp. 92  
*Leuconostoc* spp. 175, 176, 229, 656  
   *L. gasicomitatum* 228–32  
     control of 229–30  
     culture media 230  
     identification 230–1  
     as niche spoilage  
       organism 228–9  
     spoilage characteristics 229  
     culture media 178–9  
 Levine's EMB agar 249  
 limiting precision 10  
 Line agar 414, 426  
 liquid media  
   quality assurance 7–8, 22–30,  
     983–5  
   MPN 27–30  
   multiple-tube dilution  
     series 25–7  
   single-tube dilution series 22–5  
*Listeria monocytogenes* blood agar  
   (LMBA) 808–10  
*Listeria* repair broth 117, 118  
*Listeria* spp. 656  
   *L. monocytogenes*  
     culture media 115–17  
     enrichment broths 117–19  
     isolation media 119–21  
     stress damage 41  
     stress recovery 51  
     VBNC 55  
 lithium chloride phenylethanol  
   moxalactam agar 119  
 lithium chloride-sodium propionate  
   *see* LP  
 LMX broth 241  
 LP medium 208, 215  
 LP-MRS medium 184, 208  
 LP-TPY medium 208  
 LPSM medium 185

- LS differential (LSD) agar 176, 811–13
- Luria-Bartani agar 512, 518
- Luria-Bertani bile salts Irgasan (LB-BSI) broth 303
- lysine agar 572
- Lysinibacillus* spp. 92, 95
- M 5 medium 178
- M 17 medium 176, 184, 814–16
- 3M Petrifilm medium 241
- m-Colibblue system 241
- M-Endo medium 607, 609, 837–9
- M-Enterococcus (mE) agar 156, 162, 166, 607, 612, 830–2, 840–2
- M-FC medium 607, 843–5
- M-PA agar 607, 846–8
- M104 medium 182
- MacConkey agar 238, 289, 292, 293, 305, 458
- with 1% xylose (Mac-X) 294
- with rhamnose (RMAC) 339
- MacConkey broth 236, 247, 250, 609
- MacConkey mannitol ampicillin (MMA) agar 464, 466, 468
- MacConkey trehalose (non-lactose; MT) agar 464
- MacConkey Tween 80 ampicillin (MAT) agar 465
- MacConkey xylose ampicillin (MXA) agar 465, 466, 468
- MacConkey-ampicillin (MA) agar 465
- malt extract agar 559, 560
- malt salt agar 565
- malt yeast extract sucrose agar (MYSA) 565
- malt yeast salt glucose agar 565
- Mandell medium 497
- Mann-Whitney-Wilcoxon *U*-test 12
- mannitol egg-yolk polymyxin (MEYP) agar 91, 98–9, 823–5
- mannitol lysine crystal violet brilliant green (MLCB) agar 274, 275, 276, 826–9
- mannitol salt agar 136
- marine agar 512, 513, 514, 543
- marine broth 384
- Marinescu broth 430
- Marinibacillus* spp. 92
- M. marinus* 100
- marker organisms 234
- mE agar for Enterococci *see* M-Enterococcus (mE) agar
- Mechercharimyces* spp. 92
- media
- quality assessment 6–8
- future developments 31
- liquid media 7–8, 22–30
- solid media 6–7, 8–16
- medium A 544
- medium K 512
- Melissococcus* spp. 176
- membrane filtration
- Campylobacter* spp. 420–2
- Enterobacteriaceae 242–3
- hydrophobic grid membrane filters 242, 252
- membrane lactose glucuronide agar 607, 833–6
- membrane lauryl sulfate broth (MLSB) 607, 609
- MGV medium 492
- Microbacterium* spp. 195
- Micrococcaceae 194
- Micrococcus luteus*, VBNC 55
- Microval 4
- Miles-Misra method, modified 6, 978
- milk agar 498, 499
- Miller-Mallinson agar 274, 275, 276, 278
- minerals-modified glutamate (MMG) agar 238, 609
- minerals-modified glutamate (MMG) broth 236, 244, 245
- minimal salt acetate (MSA) medium 497
- MIX agar 465
- molds *see* yeasts and moulds
- molybdate agar 572
- Monascus* spp., *M. ruber* 568
- monosaccharides 625

- Monsur's taurocholate tellurite peptone (TTP) broth 382, 383
- Moraxellaceae* 484
- most probable number (MPN) 7, 27–30, 123
- Enterobacteriaceae* 243, 246, 249
- improbable results 27
- resuscitation of injured cells 50
- significance of differences 27–30
- statistics 27
- yeasts and moulds 563
- moulds *see* yeasts and moulds
- MPN *see* most probable number
- MRS agar 176, 178, 181, 230, 817–19
- with acetic acid 184
- with amphotericin 230
- with clindamycin 184
- with dicloxacillin (MRS-Dic) 209, 217–18
- with galactose 184
- modified bile (MB-MRS) 203, 209
- with nalidixic acid, neomycin sulfate and lithium chloride (MRS-NNL) 207
- with nalidixic acid, neomycin sulfate, lithium chloride and paromomycin sulfate (MRS-NPNL) 184, 207, 213
- with oxgall and dicloxacillin (MRS-BCD) 209
- with sorbic acid (MRS-S) 178, 230, 820–2
- with trehalose 184
- with vancomycin 184
- Mucor* spp. 570
- Mueller-Hinton broth/agar 186, 432
- Mueller-Kauffmann tetrathionate broth 849–52
- Mueller-Kauffmann tetrathionate novobiocin broth (MKTTn) 263, 268, 269, 270, 271
- Mueller-Hinton agar with 1% ISOVitaleX and 1% haemoglobin 598
- multiple-tube dilution series 25–7
- MW (modified Wilkins-Chalgren) medium 211
- Myroides* spp. 509, 533
- characteristics 515–16
- colony morphology 538–9
- culture media 513
- fatty acid methyl esters 544
- pigment production 540
- National collection of type cultures (NCTC) 5
- Neisseriales* 483
- Neosartorya fischeri* 566
- nitrite actidione polymyxin B (NAP) agar 178
- p*-nitrophenyl- $\beta$ -D-glucuronide *see* PNPG
- Nordic Committee on Food Analysis (NMKL) standards
- Aeromonas* spp. 470
- Campylobacter* spp. 424
- Escherichia coli* 345
- Yersinia* spp. 311
- novobiocin 269, 290, 459
- nutrient agar 458, 512, 513, 518
- nutrient broth 267
- nutrient broth no. 2 513
- Oceanobacillus* spp. 92
- Oceanospirillales* 483
- Oenococcus* spp. 175, 176
- culture media 181–2
- OF medium 499
- Oh and Kang (OK) agar 365–6, 369
- oleandomycin polymyxin sulfadiazine perfringens agar (OPSPA) 70, 71, 76, 853–5
- ONE Broth-Salmonella 272
- orange serum agar 107, 108
- Ornithinibacillus* spp. 92
- Oxford agar 120, 856–9
- modified 120, 860–2
- oxidative stress 39
- prevention of cell death from 52–4
- sensitivity of injured cells 45, 47–9

- Oxoid *Arcobacter* broth/agar 430  
Oxoid Brilliance agar 411  
Oxoid broth 430  
Oxoid rapid salmonella test 272  
Oxoid Salmonella Chromogenic medium II 274, 275, 279  
Oxyrase 54, 266, 267, 425  
oxytetracycline glucose yeast extract (OGY) agar 564, 571, 863–5
- Paecilomyces varioti* 568  
*Paenibacillus* spp. 92, 95, 96, 100  
*Paralactobacillus* spp. 176  
*Paraliobacillus* spp. 92  
Park B medium 412  
    modified 412  
Park-Sanders broth 413, 423–4  
Park-Stankiewickz B medium 412  
*Pasteuria* spp. 92  
PATHATRIX 328  
*Paucisalibacillus* spp. 92  
pectin agar 463, 465  
*Pediococcus* selective medium 179–80  
*Pediococcus* spp. 175, 176, 656  
    culture media 179–80  
*Pelagibacillus* spp. 92  
penicillin and pimarcin agar (PPA) 492, 494, 498  
*Penicillium* spp. 560, 568, 569  
pentachloronitrobenzene peptone agar 569  
pentachloronitrobenzene rose bengal yeast extract sucrose (PRYES) agar 568  
peptone beef extract glycogen (PBG) agar 463, 465, 468  
peptone saline cellobiose (PNC) enrichment broth 383, 385  
peptone, sorbitol and bile salts (PSB) broth 303  
peptone yeast glucose starch medium 80  
Petrifilm™ Coliform Count Plate 242  
Petrifilm™ *E. coli* Count Plate 242, 247, 248, 252  
Petrifilm™ *Enterobacteriaceae* 242, 252  
Petrifilm™ YM 574  
phenol red brilliant green agar 866–9  
phenol red sorbitol MacConkey agar with MUG 336, 337–8  
2-phenylethanol 179  
phosphate-buffered saline with 0.5% peptone 301  
    with 1% mannitol 301  
    with 1% sorbitol/0.15% bile salts (PBSSB) 298, 301, 305  
    with peptone and cycloheximide 301  
*Pichia membranaefaciens* 571  
*Piscibacillus* spp. 92  
plate count agar (PCA) 492, 512, 513, 514, 518, 870–2  
Plate and Ordal agar 513  
plated media  
    quality assurance 978–82  
    see also individual media  
*Plesiomonas* (PL) agar 471  
*Plesiomonas shigelloides* 451–81  
    culture media 470–2  
    identification 454–5, 456  
Poisson distribution 9–13  
polymerase chain reaction 577  
polymyxin acriflavine lithium chloride ceftazidime aesculin mannitol (PALCAM) agar 120, 122, 873–6  
polymyxin egg-yolk mannitol bromothymol blue agar 91, 98–9  
polymyxin mannose tellurite (PMT) agar 386, 389  
polymyxin pyruvate egg yolk mannitol bromothymol blue agar (PEMBA) 877–80  
Pontiac fever 597  
*Pontibacillus* spp. 92  
potassium thiocyanate actidione sodium azide egg-yolk pyruvate (KRANEP) agar 134, 136  
potato dextrose agar (PDA) 107, 108, 564

- Premier EHEC kit 344  
 presence/absence screening 52  
 Preston agar 412, 415  
 Preston *Campylobacter* selective agar 881–3  
 Preston enrichment broth 412, 415, 426, 427, 884–6  
 prill xylose ampicillin (PXA) agar 465  
 probiotic LAB 185  
 productivity ratio 3, 6, 16–20, 29–30  
   confidence intervals 18–20  
   definition of 977  
   ISO standard 16  
*Propionibacterium* spp. 195  
*Proteus* spp. 234  
   characteristics 308  
*Pseudomonadales* 482–507  
   culture media 487–501  
     differential features 487–9  
     enrichment media 489–90  
     identification 496–501  
     selective plating media 490–6  
*Pseudomonas* agar base with cetrimide and nalidixic acid 887–9  
*Pseudomonas* agar fluorescein (PA-F) 487–8, 499, 513  
*Pseudomonas* agar pyocyanin (PA-P) 487–8, 499  
*Pseudomonas*  
   cetyltrimethylammonium bromide and nalidixic acid (CN) agar 490–1, 493, 607  
*Pseudomonas* selective isolation agar (PSIA) 492  
*Pseudomonas* spp. 485–7, 656, 484, 486, 501  
   *P. aeruginosa* 486, 501  
     water testing 605, 607, 614–15  
   *P. fluorescens* 484  
     resuscitation time 45  
     stress damage 42  
   see also *Pseudomonadales*  
*Psychrobacter* spp. 485  
   see also *Pseudomonadales*  
*Pullulanibacillus* spp. 92, 100  
 quality assessment 3, 4  
   of media 6–8  
     liquid media 7–8, 22–30  
     solid media 6–7, 8–16  
   standardisation of inoculum 5–6  
 quality assurance of culture media 977–85  
   absolute and relative growth indices 978  
   definition of productivity ratios 977  
   definition of selectivity factor 978  
   liquid media 983–5  
   plated media 978–82  
 R&F-*E. coli* O157:H7 336, 343  
 R&F-ECEB medium 333  
 R2A (see Reasoner's 2A) agar 890–2  
 rabbit plasma fibrinogen agar (RPF) 135, 136, 137–40, 893–7  
 raffinose media  
   RAF 5.1 208, 216  
   RAF + gentamicin 208  
   RAF + LiCl 208  
   RAF + nalidixic acid, neomycin sulfate, lithium chloride and paromomycin sulfate (NPNL) 208  
   RAF *Bifidobacterium* (RB) medium 208, 215–16  
 Rainbow<sup>®</sup> Agar O157 336, 338, 339, 341, 343  
   with potassium tellurite and novobiocin 336, 338  
*Ralstonia pickettii* 501  
 Rambach agar 277, 278, 898–900  
 Rapid *E. coli* 2 242  
 rapid perfringens medium 70, 77–8, 901–3  
 RapidChek<sup>®</sup> *Escherichia coli* O157:H7 enrichment broth (REB) 333  
 RAPID'*E. coli* O157:H7 agar 336  
 Rappaport broth, modified (MRB) 302, 305

- Rappaport medium 271
- Rappaport–Vassiliadis medium (MSRV), semisolid  
modification 269, 271, 908–11
- Rappaport–Vassiliadis soya peptone broth 263, 265, 271
- Rappaport–Vassiliadis (RVS) agar 264, 271
- Rappaport–Vassiliadis (RVS) broth 22, 266, 270, 271, 904–7  
with soya 270, 271
- RCAAD medium 209
- reactive oxygen species 39, 47  
enhanced sensitivity to 49  
removal of 52–4  
sources of 47–9
- ReadyCult coliforms medium 241
- Reasoner's 2A (R2A) medium 512, 513, 543, 544, 890–2
- recovery supplements 51
- reinforced clostridial agar (RCA) 70, 77–8, 203, 204
- relative growth index 978
- resuscitant agar 238
- reversed passive latex agglutination (RPLA) test 344
- Rhizobiales* 483
- Rhodocyclales* 483
- Rhodospirillales* 483
- Rimler-Shotts broth 463, 465  
modified 467
- RMAC medium 339
- Rogosa agar 178, 230, 912–14  
modified 915–17
- Rogosa-neomycin sulfate, paromomycin sulfate, sodium propionate and lithium (NPPL) medium 207
- rose bengal chloramphenicol (RBC) agar 918–20
- rose bengal chlortetracycline agar 564
- rose bengal streptomycin agar 567
- Rosef B medium 412
- Rosef-Kapperud A medium 413
- Ryan medium 466, 470
- Sabouraud agar, modified 568
- Saccharomyces cerevisiae* 571
- Saccharomycodes ludwigii* 572
- Salegendibacter* spp. 509, 513, 533–4, 535–6  
characteristics 515–16  
colony morphology 540  
culture media 513  
pigment production 540
- Salimicrobium halophilum* 101
- Salinibacillus* spp. 92
- Salirhabdus* spp. 92
- Sallibacillus salexigens* 101
- Salmonella* identification medium II 274, 275, 277
- Salmonella* identification (SMID) agar 279
- Salmonella* Precis™ 272
- Salmonella Shigella* agar 274, 275, 277, 292, 293
- Salmonella Shigella* deoxycholate calcium (SSDC) agar 298, 305, 921–3
- Salmonella* spp. 234, 261–86, 657  
*S. anatum*, stress damage 44  
*S. bareilly*, stress damage 44  
*S. enteritidis*, VBNC 55  
*S. typhimurium*  
stress damage 41, 42, 44  
stress recovery 51, 53  
enumeration 280–1  
pre-enrichment 264–8  
supplement addition 266–8  
reference methods 262–4  
resuscitation time 45  
selective enrichment 268–72  
modifications 271–2  
Rappaport medium 270  
selective media 270–1  
selenite-based media 270  
tetrathionate media 268–70  
selective plating 272–80  
water testing 605, 616
- Salmosyst® broth 272
- Salsuginibacillus* spp. 92
- salt colistin broth (SCB) 382, 383



- salt polymyxin broth (SPB) 382, 383  
 salt starch xylose lysine deoxycholate (SSXLD) agar 465  
 sampling error 8–9  
*Schizosaccharomyces* spp. 571  
   *S. pombe* 571  
*Scopulariopsis* spp. 565  
 SD-39 agar 336  
*Seinonella* spp. 92  
 selective enrichment media 40, 52, 55, 332–3  
 selectivity factor 3, 20–1  
   definition of 978  
 selenite cystine broth 265, 266, 270, 271, 290, 291, 924–6  
 semi-solid selective motility (SSM) agar/broth 413  
*Serratia* spp. 657  
   characteristics 308  
 SGAP-10C agar 465  
 Shahidi Ferguson perfringens agar 72  
*Shewanella putrefaciens* 483, 498  
 Shieh agar, modified 543, 544  
 Shiga toxins 324  
*Shigella* broth 289, 290, 293  
*Shigella* spp. 287–97, 657  
   *S. boydii* 288  
   *S. dysenteriae* 288  
   *S. flexneri* 288  
     isolation media 293  
     stress damage 43  
   *S. sonnei* 288  
     alternative culture media 294–5  
     enrichment media 289–91  
     isolation media 291–3  
     water testing 616  
*Shimazuella* spp. 92  
 SimPlate Coliform and *E. coli* medium 241, 252  
 SimPlate technique, yeasts and moulds 574  
 simple method *Salmonella* 270  
 single-tube dilution series 22–5  
 skim milk agar 184  
 Skirrow *Campylobacter* selective agar 927–9  
 Skirrow medium 412, 415  
 Slantez and Bartley agar 156, 165  
 sodium dodecyl sulphate polymyxin sucrose (SPS) agar 386, 389, 393  
 solid media  
   quality assessment 6–7, 8–16  
   productivity ratio 16–20  
   selectivity factor and specificity 20–1  
   statistics 9–16  
   technical errors 8–9  
 solid medium repair 50–1  
 sorbitol MacConkey agar (SMAC) 51, 336, 337–8, 341–2  
   with cefixime, potassium tellurite and 8-hydroxy-quinoline- $\beta$ -D-glucuronide (HQG) 336, 337  
   with cefixime and rhamnose 336, 337  
   phenol red, with MUG 336, 337–8  
 specificity 20–1  
*Sphingomonadales* 483  
*Sphingomonas paucimobilis* 501  
 spiral plate technique 6, 978–81  
*Sporolactobacillus* spp. 92  
   culture media 109  
   *S. laevolacticus* 100  
*Sporosarcina* spp. 92  
   *S. globispora* 100  
   *S. pasteurii* 101  
 stab inoculation method 982  
 standard error of mean 14  
 staphylococcal food poisoning 131–2  
*Staphylococcus* spp. 132  
   *S. aureus* 132, 657, 655, 657  
     stress damage 41, 42, 43  
     water testing 605, 615–16  
   *S. aureus* subsp. *anaerobius* 132  
 starch ampicillin agar (SAA) 452, 465, 466, 468, 930–2  
 starch bile salts (SB) agar 465  
 starch gelatin polymyxin (SGP) broth 382, 383  
*Stenotrophomonas maltophilia* 501

- streaking method 981-2  
*Streptococcus* spp. 156, 157, 176,  
 184, 657  
   water testing 610-12  
 streptomycin thallos acetate  
   actidione (STAA) agar 933-5  
 stress  
   effects on cells 41-3  
     cytoplasmic membrane  
       damage 41  
     DNA damage 42-3  
     Gram-negative outer membrane  
       damage 41  
     ribosomal and RNA  
       damage 41-2  
   manifestations of 43-9  
     increased lag period 45, 46  
     sensitivity to oxidative  
       stress 45, 47-9  
     sensitivity to secondary  
       stresses 44-5  
     sensitivity to selective  
       agents 43-4  
 stressed microorganisms, recovery  
   of 49-54  
   MPN 50  
   presence/absence screening 52  
   prevention of death during  
     52-4  
   solid medium repair 50-1  
 Student's *t*-test 12-14  
 sublethal microbial injury 40  
 sucrose Teepol tellurite (STT)  
   agar 386, 389  
 sugars 624, 625  
   glycones and aglycones 626-8  
 sulfite cycloserine azide (SCA)  
   agar 70, 71, 73, 74, 936-8  
 sulfite iron agar 71  
 sulfite polymyxin sulfadiazine  
   agar 71, 75  
 sulfite-reducing bacteria 70, 72  
 sulfite-reducing clostridia 70  
*Sulfobacillus* spp. 92  
   *S. acidophilus* 100  
   *S. thermosulfidooxidans* 100  
 surface spread plate 6  
 surfactants 458  
   *see also* sodium lauryl sulfate  
  
*Talaromyces* spp.  
   *T. bacillisporus* 566  
   *T. macrosporus* 566  
 tannin sucrose agar 569  
 taurocholate tellurite peptone (TTP)  
   broth 382, 383  
 Teknaf enteric agar (TEA) 294  
 Tempo EC medium 241, 249  
*Tenacibaculum* spp. 509, 514,  
 534-6  
   *T. soleae* 514, 535-6  
   characteristics 515-16  
   culture media 514  
   pigment production 540  
*Tenuibacillus* spp. 92  
 Tergitol 7 agar 266, 292  
*Terribacillus* spp. 92  
*Tetragenococcus* spp. 175, 176  
   culture media 181  
 tetrathionate broth 266, 270, 271  
   plus brilliant green 267  
*Thalassobacillus* spp. 92  
 thallos acetate tetrazolium glucose  
   (TITG) agar 156, 162, 165, 166,  
   939-41  
*Thermoactinomyces* spp. 92  
*Thermobacillus* spp. 92, 100  
 thermonuclease assay 144-5  
 thiosulfate chloride iodide (TCI)  
   agar 386, 388  
 thiosulfate citrate bile salts sucrose  
   (TCBS) agar 386, 387-9,  
   942-4  
 TITG *see* thallos acetate tetrazolium  
   glucose agar  
 tomato juice agar (TJA) 178, 230  
 tomato juice-glucose-fructose-malate  
   broth 182  
*Torulaspora delbrueckii* 573  
 TPPY agar 176  
*trans*-galactosylated oligosaccharide  
   *see* TOS

- transoligosaccharide (TOS) agar 203, 214–15  
 with mupirocin and acetic acid (TOS-AM) 206, 212  
 with nalidixic acid, neomycin sulfate, lithium chloride and paromomycin sulfate (TOS-NPNL) 207, 213
- Triclosan 491
- tris-buffered peptone water 301, 302
- Trojan horse 632, 637–40, 641,
- trypticase peptone glucose yeast-extract beef-extract medium 80–1
- trypticase phytone yeast extract-Dic medium 209, 217–18
- trypticase soy agar (TSA) 44, 365–6, 458, 512, 513, 514, 518, 543, 544  
 with sodium pyruvate (TSAP) 367
- trypticase soy broth ampicillin Irgasan (TSBAI) 467, 468
- trypticase soy broth ampicillin (TSBA) 452, 465, 467, 468
- trypticase soy broth plus polymyxin and novobiocin (TSPN) 301
- trypticase soy broth (TSB) 267, 301, 468, 512
- trypticase soya agar triphenyltetrazolium (TSAT) agar 386, 388, 391
- tryptone 10% glucose yeast extract (TGY) agar 566
- tryptone bile agar (TBA) 238, 607, 609, 945–8
- tryptone bile X-glucuronide (TBX) agar 249, 251, 252
- tryptone glucose yeast extract (TGY) agar 203, 520
- tryptone phytone yeast extract 203
- tryptone soy broth 135, 298, 489  
 with 10% NaCl and 1% sodium pyruvate (PTSBS) 949–51  
 with casamino acids 326  
 with cefixime, tellurite and vancomycin 326, 331, 334  
 with novobiocin 326  
 with novobiocin and vancomycin 335
- tryptone sulphite neomycin agar 71
- tryptose sulfite cycloserine (TSC) agar 70, 71, 75, 607, 952–5
- tryptose sulfite cycloserine-MUP agar 75–6
- Tuberibacillus* spp. 92
- Tween 80 230, 461
- Tween agar, all purpose (APT) 663–5
- UK National Collection of Type Cultures 5
- universal beer agar (UBA) 182
- Universal pre-enrichment (UP) broth 267–8
- University of Vermont (UVM) broths I & II 956–8
- Ureibacillus* spp. 92  
*U. thermosphaericus* 100
- Vagococcus* spp. 176
- vancomycin-resistant enterococci 156, 157
- VBNC *see* viable but non-culturable cells
- verocytotoxins 324, 344
- Verotox FTM test 344
- viable but non-culturable cells 54–5
- Vibrio fluvialis* enrichment medium (FEM) 385
- Vibrio fluvialis* medium 383
- Vibrio parahaemolyticus* (VP) agar 386, 389, 391
- Vibrio* spp. 377–402, 454, 657  
*V. cholerae* 377, 378  
 culture media 394  
 enrichment 382  
 occurrence in foods 378–9  
 plating media 388–9  
 VBNC 55  
*V. parahaemolyticus* 377, 378  
 culture media 394  
 enrichment 382–4  
 occurrence in foods 379

- plating media 390–1
- stress damage 42, 45
- V. vulnificus* 377, 378
  - culture media 394
  - enrichment 384–5
  - occurrence in foods 379–80
  - plating media 391–3
  - VBNC 55
- colony size and appearance 390
- enrichment 380–5
  - modes of action 381
- identification 394–5
- ISO standard methods 395–6
- occurrence in foods 378–80
- plating media 385–94
- sucrose reactions 387
- water testing 617
- Vibrio vulnificus* enumeration (VVE)
  - agar 386
- Vibrio vulnificus* (VV) agar 386, 393
- vibriostatics 460
- violet red bile glucose (VRBG)
  - agar 238, 239, 250, 251, 362, 363, 365–6, 962–4
- violet red bile lactose (VRBL)
  - agar 292
- violet red bile MUG agar 238, 249
- violet red bile Redigel 243
- violet red bile (VRB) agar 238, 247, 250, 251, 959–61
- Virgibacillus* spp. 92, 101
- Virion medium 413
- virulent *Yersinia enterocolitica*
  - agar 305, 306
- Vitek Immuno Diagnostic Assay System 329
- Vitellibacter vladivostokensis* 509, 535–6, 537
  - characteristics 515–16
  - colony morphology 540
  - culture media 514
  - enzyme production 542
  - fatty acid methyl esters 544
  - pigment production 540
- Vogel Johnson agar 136
  - modified 136
- VRBL-2 pour-plate procedure 246–7
- VTP-FBP B medium 413
- Wadowsky Yee (MWY) agar,
  - modified 597, 598–9, 965–7
- Walleimia sebi* 565
- Wallerstein Laboratory nutrient medium 572
- washed sheep blood agar with calcium (WSBA-Ca) 340
- water testing 605–22
  - Clostridium* spp. 612–13
  - coliforms 605, 607, 608–10
  - culturable heterotrophic bacteria 613–14
  - enterococci and streptococci 610–12
  - Escherichia coli* 605, 607, 608–10
  - history 606
  - isolation media for indicator organisms 606–8
  - pathogenic organisms 607, 614–17
- Waterman A medium 413
- Waterman B medium 413
- Weeksella* spp. 509, 537
  - characteristics 515–16
  - colony morphology 538–9
  - culture media 514
  - fatty acid methyl esters 544
- Weiler and Radler medium 182
- Weissella* spp. 176
- wey agar with casminoacids and calcium lactate 181
- Wilkins-Chalgren (WC) medium 211
- Working Party for Culture Media of the International Committee for Food Microbiology and Hygiene (WPCM) 4
- Xanthomonadales* 483
- Xeromyces bisporus* 565
- xylose ampicillin agar 465
- xylose galactosidase medium 295
- xylose lysine deoxycholate (XLD) agar 263, 274, 275, 277, 292, 293, 968–71

- xylose lysine Tergitol 4 (XLT4)
  - agar 274, 275, 277, 972-4
- xylose sodium deoxycholate citrate (XDC) agar 465
- Yarrowia lipolytica* 573
- yeast extract agar 565, 607, 975-6
- yeast extract, beef extract, special peptone, FCS plus iron medium 433
- yeast extract glucose chloramphenicol agar (YGCA) 570
- yeast extract lactose tryptone bile agar 238
- yeast glucose peptone medium 181
- yeast starch glucose agar 107, 108
- yeasts and molds 557-95, 656
  - culture media 564-73
    - general purpose 564-5
    - heat-resistant moulds 566-7
    - mycotoxigenic moulds 567-70
    - preservative- and acid-tolerant yeasts 571
    - selective 570-1
    - wild yeast 571-3
    - xerophilic moulds and osmotolerant yeasts 565-6
  - data interpretation 577-8
  - diluent 562
  - immunological methods 576-7
  - incubation 563-4
  - indirect methods 575-6
  - microscopic methods 575
  - modified cultivation
    - methods 573-5
  - molecular techniques 577
  - plating vs. MPN 563
  - precautions 578
  - sample preparation 561-2
- Yersinia* spp., 299, 309, 657
  - Y. enterocolitica* 298-320
    - alkaline treatment 304
    - biotypes 310
    - enrichment media 301-4
    - epidemiology 300
    - identification 307-11
    - pathogenic potential 300, 311
    - plating media 304-7
    - standard methods 311
  - Y. pseudotuberculosis* 298, 299, 301, 309
    - enrichment media 303-4
    - pathogenic potential 300
  - biochemical differentiation 309
  - characteristics 308
  - water testing 617
- YN-6 medium 206
- YN-17 medium 206
- Z-agar 492
- Zygosaccharomyces* spp.
  - Z. bailii* 566, 571, 573, 577
  - Z. bisporus* 573
  - Z. rouxii* 565-6, 571