

INDEX

- Actiote – see annatto
- ADI, of Allura Red 90
- of amaranth 94
 - of annatto 7
 - of anthocyanins 14
 - of beetroot red 33
 - of beta-apo-8'-carotenal 20
 - of beta-carotene 25
 - of Brilliant Black BN 98
 - of Brilliant Blue FCF 100
 - of Brown HT 106
 - of canthaxanthin 37
 - of caramel 40
 - of Carbon Black 109
 - of carmine 45
 - of carmoisine 112
 - of chlorophyll 49
 - of copper complexes of chlorophyllins 55
 - of curcumin 62
 - of erythrosine 114
 - of ethyl ester of beta-apo-8'-carotenoid acid 67
 - of Fast Green FCF 118
 - of Green S 122
 - of Indigotine 126
 - of iron oxides and hydroxides 131
 - of Litholrubine BK 134
 - of lutein 72
 - of lycopene 76
 - of mixed carotenes 29
 - of paprika 81
 - of Patent Blue V 136
 - of Ponceau 4R 140
 - of Quinoline Yellow 144
 - of riboflavin 84
 - of Sunset Yellow FCF 148
 - of tartrazine 153
 - of titanium dioxide 156
- Aleoholic beverages,
- use of amaranth 94
 - use of annatto 6
 - use of beta-carotene 23, 24
 - use of caramel 40
- Allura Red AC 89-91
- ADI 90
- analytical methods 90
 - applications 90
 - chemical structure (fig.) 89
 - colour shade 89
 - physical characteristics 90
 - physiological properties 90
 - solubility 90
 - stability to acid, heat, light and sulphur dioxide 90
 - suppliers 173
- Amaranth 93-5
- ADI 94
 - analytical methods 95
 - applications 94
 - chemical structure (fig.) 93
 - colour shade 93
 - physical characteristics 94
 - physiological properties 94-5
 - solubility in oil, propylene glycol and water 94
 - stability to acid, heat, light and sulphur dioxide 94
 - suppliers 173
- American, use of amaranth 94
- Ammonia caramel – see caramel
- Annatto 1-8
- ADI 7
 - analytical methods 7-8
 - applications 6-7
 - colour shade 4
 - extraction 5
 - physical characteristics 5-6
 - physiological properties 7
 - solubility in ethanol, oil and water 6
 - source 4
 - stability to heat and light 5
 - structural formula (fig.) 3
 - suppliers 171
 - susceptibility to oxidation 5
- Anthocyanins 9-15
- ADI 14
 - analytical methods 14
 - applications 12-3
 - colour shade 10
 - effect of pH 11

- physical characteristics 11-2
- physiological properties 13-4
- solubility in ethanol, oil and water 12
- source 10-1
- stability to heat and light 11-2
- structural formula (fig.) 10
- suppliers 171
- susceptibility to oxidation 12
- Apéritif wines and spirit drinks, use of
amaranth 94
- Azorubine – see carmoisine
- Baked products, use of Allura Red 90
 - use of Brilliant Blue FCF 100
 - use of Brown HT 106
 - use of carmine 44
 - use of carmoisine 112
 - use of curcumin 62-3
 - use of Fast Green FCF 118
 - use of Patent Blue V 136
 - use of Ponceau 4R 140
 - use of Quinoline Yellow 144
 - use of Sunset Yellow FCF 148
 - use of tartrazine 152
 - use of titanium dioxide 156
- Beer caramel – see caramel
- Beers, use of caramel 49
- Beetred – see beetroot red
- Beetroot red 31-3
 - ADI 33
 - analytical methods 33
 - applications 32-3
 - colour shade 32
 - effect of pH 32
 - physical characteristics 32
 - physiological properties 33
 - source 32
 - stability to heat and light 32
 - suppliers 171-2
 - susceptibility to oxidation 32
- Beta-apo-8'-carotenal 17-20
 - ADI 20
 - analytical methods 20
 - applications 19
 - colour shade 17
 - effect of pH 18
 - physical characteristics 18-9
 - physiological properties 19-20
 - solubility in ethanol, oil and water 19
 - source 18
 - stability to heat and light 18
 - structural formula (fig.) 17
 - suppliers 171
 - susceptibility to oxidation 18
- Beta-apo-8'-carotenol ester – see ethyl ester
of beta-apo-8'-carotenol acid
- Beta-carotene 21-6
 - ADI 25
 - analytical methods 25
 - applications 23-5
 - colour shade 21
 - effect of pH 22
 - physical characteristics 22-3
 - physiological properties 25
 - solubility in oil and water 22, 23
 - source 22
 - stability to heat and light 22
 - structural formula (fig.) 21, 27
 - suppliers 171
 - susceptibility to oxidation 223
- Betanin – see beetroot red
 - structural formula (fig.) 31
- Beverages – see Alcoholic beverages, Milk
drinks and Soft drinks
- Biscuits, use of lycopene 77
- Bitter soda, use of amaranth 94
- Bixin – see annatto
 - structural formula (fig.) 4
- Black PN, suppliers 174
- Black shade, of caramel 39
 - of Carbon Black 107
 - of iron oxides and hydroxides 129
- Blue shade, of anthocyanins 10
 - of Brilliant Blue FCF 99
 - of Green S 121-2
 - of Indigo Carmine 125
 - of Patent Blue V 135
- Blue/green shade, of Fast Green FCF 117
- Blue/red shade, of amaranth 93
 - of carmoisine 111
- Blue/violet shade, of Brilliant Black BN 97
- Brandy, use of caramel 40
- Breadcrumbs, use of annatto 7
- Brilliant Black BN 97-8
 - ADI 98
 - applications 98
 - chemical formula (fig.) 97
 - colour shade 97
 - physical characteristics 98
 - physiological properties 98
 - solubility in ethanol, oil and water 98
 - stability to acid, alkaline, heat and light 98
 - suppliers 174
- Brilliant Black, analytical methods 98
 - solubility in propylene glycol 98
 - stability to sulphur dioxide 98
- Brilliant Blue FCF 99-101
 - ADI 100
 - analytical methods 100
 - applications 100
 - chemical structure (fig.) 99
 - colour shade 99
 - physical characteristics 100
 - physiological properties 100

- solubility in ethanol, oil, propylene glycol and water 100
 stability to acids, alkaline, heat, light and sulphur dioxide 100
 suppliers 174
- Brilliant Green BS – see Green S
- Brown FK 103
 applications 103
 colour shade 103
 physical characteristics 103
 solubility in water 103
 stability to acid, alkaline, heat, light and sulphur dioxide 103
 suppliers 174
- Brown HT 105-6
 analytical methods 106
 applications 106
 chemical structure (fig.) 105
 colour shade 105
 physical characteristics 105-6
 physiological properties 106
 solubility in ethanol, oil, propylene glycol and water 106
 stability to acids, alkaline, heat, light and sulphur dioxide 105
 suppliers 174
- Brown shade, of Brown FK 103
 of Brown HT 105
 of caramel 39
- Cake batters, use of beta-carotene 24
 use of lutein 72
 use of lycopene 77
 use of erythrosine 114
- Canned products, use of Brilliant Blue FCF 100
 use of Fast Green FCF 118
 use of Green S 122
 use of Patent Blue V 136
 use of tartrazine 152
- Canthaxanthin 35-8
 ADI 37
 analytical methods 37-8
 applications 37
 colour shade 35
 effect of pH 36
 physical characteristics 36
 physiological properties 37
 solubility in ethanol, oil and water 37
 source 36
 stability to heat and light 36
 structural formula (fig.) 35
 suppliers 172
 susceptibility to oxidation 36
- Capsanthin, structural formula (fig.) 79
- Caramel 39-41
 ADI 40
 analytical methods 40-1
 applications, 40
 colour shade 39
 physical characteristics 39-40
 physiological properties 40
 stability to heat and light 40
 suppliers 172
- Carbon Black 107-10
 ADI 109
 analytical methods 109
 applications 108
 colour shade 107
 effect of pH 108
 physical characteristics 108
 physiological properties 108-9
 solubility in ethanol, oil and water 108
 source 107
 stability to heat and light 108
 suppliers 174
 susceptibility to oxidation 108
- Carmine 43-6
 ADI 45
 analytical methods 46
 applications 44-5
 colour shade 44
 physical characteristics 44
 physiological properties 45
 source 44
 structural formula (fig.) 43
 suppliers 172
- Carmine 6B – see Litholrubine BK
- Carminic acid – see carmine
- Carmoisine 111-2
 ADI 112
 analytical methods 112
 applications 112
 chemical structure (fig.) 111
 colour shade 111
 physical characteristics 111-2
 physiological properties 112
 solubility in ethanol, oil, propylene glycol and water 112
 stability to acids, alkaline, heat, light and sulphur dioxide 111-2
 suppliers 174
- Carotenes, mixed 27-30
 ADI 20
 analytical methods 29
 applications 28, 29
 colour shade 27
 physical characteristics 28
 physiological properties 29
 source 28
 suppliers 171
- Carotenoids (mixed) – see lutein
- Caustic sulphur caramel – see caramel
- Cereal products, use of riboflavin 84
- Cheese, use of annatto 7

- use of copper complexes of chlorophyllins 54
 - use of lycopene 77
 - use of titanium dioxide 156
- Cheese rind, use of Litholrubine BK 134
- use of Ponceau 4R 140
- Chlorophyll 47-9
 - ADI 49
 - analytical methods 49
 - applications 49
 - physical characteristics 48
 - physiological properties 49
 - source 48
 - stability to heat and light 48
 - structural formula (fig.) 48
 - suppliers 172
 - susceptibility to oxidation 48
- Chlorophyllin (sodium copper), structural formula (fig.) 52
- Chlorophyllins (copper complexes) – see Copper complexes of chlorophyllins
- Chocolate Brown HT – see Brown HT
- Coatings, use of Carbon Black 108
 - use of Quinoline Yellow 144
 - use of tartrazine 152
- Cochineal – see carmine and Ponceau 4R
- Cocktail cherries, use of erythrosine 114
- Confectionery, use of Allura Red 90
 - use of anthocyanins 13
 - use of beetroot red 32
 - use of beta-carotene 24
 - use of Brilliant Black BN 98
 - use of Brilliant Blue FCF 100
 - use of Brown HT 106
 - use of Carbon Black 108
 - use of carmine 44
 - use of carmoisine 112
 - use of copper complexes of chlorophyllins 54
 - use of Fast Green FCF 118
 - use of lutein 72
 - use of lycopene 77
 - use of Patent Blue V 136
 - use of Ponceau 4R 140
 - use of Quinoline Yellow 144
 - use of Sunset Yellow FCF 148
 - use of tartrazine 152
 - use of titanium dioxide 156
- Copper chlorophyll, suppliers 172
- Copper complexes of chlorophyllins 51-6
 - ADI 55
 - analytical methods 55
 - applications 53
 - colour shade 52
 - effect of pH 53
 - physical characteristics 53
 - physiological properties 55
 - solubility in ethanol, oil and water 53
 - source 52-3
 - stability to heat and light 53
 - susceptibility to oxidation 53
- Crocin 57-60
 - analytical methods 59
 - applications 58
 - colour shade 57
 - effect of pH 58
 - physical characteristics 58
 - physiological properties 59
 - solubility in ethanol, oil and water 59
 - source 57-8
 - stability to heat and light 58
 - structural formula (fig.) 57
 - suppliers 172
 - susceptibility to oxidation 58
- Curcumin 61-4
 - ADI 62
 - analytical methods 63
 - applications 62-3
 - colour shade 61
 - effect of pH 62
 - physical characteristics 62
 - physiological properties 62
 - source 62
 - stability to heat and light 62
 - structural formula (fig.) 61
 - suppliers 173
- Custard powder, use of annatto 6
- D&C Red No. 7 – see Litholrubine BK
- Dairy products, use of annatto 7
 - use of beetroot red 32
 - use of carmine 44
 - use of carotenes 28
 - use of curcumin 62, 63
 - use of lycopene 77
 - use of riboflavin 84
 - use of beta-carotene 25
 - use of lutein 72
- Desserts, use of Allura Red 90
 - use of Brilliant Blue FCF 100
 - use of Brown HT 106
 - use of carmoisine 112
 - use of copper complexes of chlorophyllins 54
 - use of Ponceau 4R 140
 - use of Quinoline Yellow 144
 - use of Sunset Yellow FCF 148
 - use of tartrazine 151
- Difcrolymethane – see curcumin
- Edible ices, use of Brilliant Blue FCF 100
 - use of carmoisine 112
 - use of Ponceau 4R 140
 - use of Quinoline Yellow 144
 - use of Sunset Yellow FCF 148
 - use of tartrazine 152

- Enocianina – see anthocyanins
- Erythrosine 113-5
 ADI 114
 analytical methods 115
 applications 114
 chemical structure (fig.) 113
 colour shade 113
 physical characteristics 114
 physiological properties 114
 solubility in ethanol, oil, propylene glycol and water 114
 stability to acids, alkaline, heat, light and sulphur dioxide 114
 suppliers 174
- Ethyl ester of beta-apo-8'-carotenoic acid 65-68
 ADI 67
 analytical methods 67
 applications 67
 effect of pH 66
 physical characteristics 66
 physiological properties 67
 solubility in ethanol 66
 solubility in oil and in water 66
 source 66
 stability to heat and light 66
 structural formula (fig.) 65
 suppliers 172
- European Community legislation 161-6
 permitted colours 162-5
 specific labelling requirements for foods containing colours 166
- Fast Green FCF 117-9
 ADI 118
 analytical methods 118
 applications 118
 chemical structure (fig.) 117
 colour shade 117
 physical characteristics 118
 physiological properties 118
 solubility in ethanol, oil propylene glycol and water 118
 stability to acids, alkaline, heat, light and sulphur dioxide 118
 suppliers 175
 susceptibility to oxidation 118
- Fat-based coatings and toppings, use of lutein 71
- FD&C Yellow No 5 – see tartrazine
- FD&C Yellow No 6 – see Sunset Yellow FCF
- Fish products, use of Allura Red 90
 use of amaranth 94
 use of annatto 7
 use of Brilliant Black BN 98
 use of Brown FK 103
 use of curcumin 62, 63
 use of tartrazine 152
- Flavoured drinks – see soft drinks
- Flour confectionery, use of lutein 72
- Food Black 1 – see Brilliant Black BN
- Food Blue – see Indigotine
- Food Blue 2 – see Brilliant Blue FCF
- Food Blue 5 – see Patent Blue V
- Food Brown 1 – see Brown FK
- Food Brown 3 – see Brown HT and Fast Green FCD
- Food Green 4 – see Green S
- Food Orange 5 – see beta-carotene and carotenes
- Food Orange 6 – see beta-apo-8'-carotenal
- Food Orange 7 – see ethyl ester of beta-apo-8'-carotenoic acid
- Food Orange 8 – see eanthaxanthin
- Food Red 7 – see Ponceau 4R
- Food Red 9 – see amaranth
- Food Red 14 – see erythrosine
- Food Red 17 – see Allura Red
- Food Yellow 3 – see Sunset Yellow FCF
- Food Yellow 4 – see Tartrazine
- Food Yellow 5 – see Sunset Yellow FCF
- Food Yellow 13 – see Quinoline Yellow
- Food-intolerance studies, of beetroot red 153
- Fruit preparations, use of beetroot red 32
- Gardenia extract – see crocin
- Grape colour extract – see anthocyanins
- Grape skin extract – see anthocyanins
- Gravies, use of caramel 40
 use of carotenes 28
 use of lycopene 77
- Green S 121-3
 ADI 122
 analytical methods 122
 applications 122
 chemical structure (fig.) 121
 colour shade 121-2
 physical characteristics 122
 physiological properties 122
 solubility in ethanol, oils, propylene glycol and water 122
 stability to acids, alkaline, heat, light and sulphur dioxide 122
 suppliers 175
- Green shade, of chlorophyll 48
 of Fast Green FCF 118
- Green/blue shade, of Brilliant Blue FCF 100
 of copper complexes of chlorophyllins 52
 of Green S 121-2
 of Patent Blue V 135
- Green/yellow shade, of Quinoline Yellow 143
- Grey shade, of Carbon Black 107
- Haddock (smoked), use of tartrazine 152
- Herring, smoked – use of annatto 7
- Ice cream, use of annatto 6

- use of beetroot red 32
- use of beta-carotene 24
- use of carmine 44
- use of carotenes 28
- use of curcumin 62, 63
- use of lutein 71
- use of lycopene 77
- use of titanium dioxide 156
- Icings, use of beta-carotene 24**
- use of lutein 72
- use of titanium dioxide 156
- Indigo Carmine 125-7**
- chemical structure (fig.) 125
- colour shade 125
- suppliers 175
- Indigotine 125-7**
- ADI 126
- analytical methods 126
- applications 126
- effect of pH 125
- physical characteristics 125-6
- physiological properties 126
- solubility in ethanol, oil, propylene glycol and water 126
- stability to acids, heat, light and sulphur dioxide 126
- suppliers 175
- Iron oxides and hydroxides 129-32**
- ADI 131
- analytical methods 131
- applications 130-1
- colour shade 129
- effect of pH 130
- molecular mass (Table) 129
- physical characteristics 130
- physiological properties 131
- solubility in ethanol, oil and water 130
- source 130
- stability to heat and light 130
- suppliers 175
- susceptibility to oxidation 130
- Jams and jellies,**
- use of anthocyanins 13
- use of beetroot red 32
- use of beta-carotene 24
- use of copper complexes of chlorophyllins 54
- use of tartrazine 152
- Kippers, use of Brown FK 103**
- Labelling requirements, in European Community legislation 166**
- in UK legislation 167
- in USA legislation 167-9
- Lactoflavin – see riboflavin**
- Legislation 161-70**
- European Community 161-6
- international 169-70
- UK 166-7
- USA 167-70
- Litholrubine BK 133-6**
- ADI 134
- analytical methods 134
- applications 134
- chemical structure (fig.) 133
- colour shade 133
- physical characteristics 134
- physiological properties 134
- solubility in ethanol, oil and water 134
- stability to heat 134
- suppliers 175
- Lutein 69-74**
- ADI
- analytical methods 72
- applications 71-2
- colour shade 69
- effect of pH 70
- physical characteristics 70
- physiological properties 72
- solubility in ethanol, oil and water 71
- source 70
- stability to heat and light 70, 72
- structural formula (fig.) 69
- suppliers 173
- susceptibility to oxidation 70
- Lycopene 75-8**
- ADI 76
- analytical methods 77
- applications 76, 77
- colour shade 75
- physical characteristics 76
- physiological properties 76
- solubility in oil 76
- source 75
- stability to light 76
- structural formula (fig.) 75
- suppliers 173
- Mackerel, smoked – use of annatto 7**
- Magnesium chlorophyll – see chlorophyll**
- Magnesium phaeophytin – see chlorophyll**
- Meat products, use of Allura Red 90**
- use of earamel 40
- use of carmine 44
- use of lycopene 77
- use of paprika 80
- Milk drinks, use of beta-carotene 25**
- use of earmine 44
- use of lutein 72
- Mint sauce/jelly, use of copper complexes of chlorophyllins 54**
- Mixed carotenes 27-30**
- Mixed carotenoids – see lutein**
- Mozzarella cheese, use of titanium dioxide 155**
- Natural colours 1-85**

- suppliers 171-3
- Natural Green 5 – see copper complexes of chlorophyllins
- Natural Yellow 3 – see curcumin
- Natural Yellow 27 – see lycopene
- New Coccine – see Ponceau 4R
- Non-dairy creamers, use of titanium dioxide 156
- Norbixin – see annatto
structural formula (fig.) 4
- Orange shade, of annatto 4
of beta-carotene 21
of carmine 44
of lycopene 76
- Orange/red shade, of beta-apo-8'-carotenal 17
of canthaxanthin 35
of paprika 79
- Orange Yellow 5 – see Sunset Yellow FCF
- Paprika 79-81
ADI 81
analytical methods 81
applications 80-1
colour shade 79
effect of pH 80
physical characteristics 80
physiological properties 81
solubility in oil and propylene glycol 80
source 79-80
stability to heat and light 80
suppliers 173
- Pasta, use of lycopene 77
- Patent Blue V 135-7
ADI 136
analytical methods 136
applications 136
chemical structure (fig.) 135
colour shade 135
physical characteristics 136
physiological properties 136
solubility in ethanol, oil, propylene glycol and water 136
stability to acids, alkaline, heat, light and sulphur dioxide 136
suppliers 175
- Permitted colours, in European Community legislation 162-5
in UK legislation 166-7
in USA legislation 167-9
- Pet foods, use of caramel 40
use of iron oxides and hydroxides 131
- Pharmaceutical products, use of Ponceau 4R 140
- Pickles, use of curcumin 62, 63
use of Quinoline Yellow 144
use of tartrazine 152
- Pigment Red 57.1 – see Litholrubine BK
- Pigment Rubine, suppliers 175
- Pigment White 6 – see Titanium dioxide
- Pigment Yellow 42 and 43 – see Iron oxides
- Ponceau 4R 139-41
ADI 140
analytical methods 140
applications 140
chemical structure (fig.) 139
colour shade 139
physical characteristics 140
solubility in ethanol 140
solubility in oil 140
solubility in propylene glycol and in water 140
stability to acids, alkaline, heat, light and sulphur dioxide 140
suppliers 175
- Potassium chlorophyllin – see chlorophyll
- Potassium copper chlorophyllin – see copper complexes of chlorophyllins
- Quinoline Yellow 143-5
ADI 144
analytical methods 145
applications 144
chemical structure (fig.) 143
colour shade 143
physical characteristics 143-4
physiological properties 144
solubility in ethanol, oil, propylene glycol and water 144
stability to acids, alkaline, heat, light and sulphur dioxide 143
suppliers 175-6
- Ready meals, use of carmine 44
- Red fruit extract – see anthocyanins
- Red shade, of Allura Red 89
of beetroot red 32
of canthaxanthin 35
of carmine 44
of carmoisine 112
of carotenes 27
of erythrosine 113
of iron oxides and hydroxides 129
of Litholrubine BK 133
of lycopene 75
of Ponceau 4R 139
- Red/blue shade, of anthocyanins 10
- Red/brown shade, of amaranth 93
of Brown FK 103
of Brown HT 105
- Red/orange shade, from anthocyanins 10
- Riboflavin 83-5
ADI 84
analytical methods 84
applications 84
colour shade 83
physical characteristics 84
physiological properties 84

- solubility in ethanol, oil and water 84
- source 84
- stability to acid, heat and light 84
- structural formula (fig.) 83
- suppliers 173
- Rocou – see annatto
- Rubinpigment – see Litholrubine BK
- Saffron extract – see crocin
- Sauces, use of caramel 40
 - use of carotenes 28
 - use of curcumin 62, 63
 - use of lycopenc 77
 - use of paprika 80
 - use of Quinoline Yellow 144
 - use of tartrazine 152
- Saucisses de Strasbourg, use of canthaxanthin 37
- Sausage casings, use of iron oxides and hydroxides 130-1
- Sodium chlorophyllin – see chlorophyll
- Sodium copper chlorophyllin – see copper complexes of chlorophyllins
- Soft drink caramel – see caramel
- Soft drinks, use of Allura Red 90
 - use of annatto 6
 - use of anthocyanins 13
 - use of beta-apo-8'-carotenal 19
 - use of beta-carotene 24
 - use of beetroot red 32
 - use of Brilliant Blue FCF 100
 - use of caramel 40
 - use of carmoisine 112
 - use of carotenes 28, 29
 - use of curcumin 62
 - use of Fast Green FCF 118
 - use of lutein 71
 - use of lycopenc 77
 - use of paprika 79, 80
 - use of Ponceau 4R
 - use of Quinoline Yellow 144
 - use of Sunset Yellow FCF 148
 - use of tartrazine 152
- Sorbets, use of anthocyanins 13
 - use of beta-carotene 24
 - use of lutein 71
- Soups, use of carotenes 28
 - use of lycopenc 77
 - use of paprika 80
- Spices, use of paprika 80
- Spirit caramel – see caramel
- Spreads, use of lycopenc 77
- Sugar confectionery, use of anthocyanins 13
 - use of beetroot red 32
 - use of beta-carotenc 24
 - use of Carbon Black 108
 - use of carmine 44
- use of copper complexes of chlorophyllins 54
- use of curcumin 62, 63
- use of iron oxides and hydroxides 131
- use of lutein 72
- use of lycopenc 77
- Sulphite ammonia caramel – see caramel
- Sunset Yellow FCF 147-9
 - ADI 148
 - analytical methods 148
 - applications 148
 - chemical structure (fig.) 147
 - colour shade 147
 - effect of pH 148
 - physical characteristics 148
 - physiological properties 148
 - solubility in ethanol, oil, propylene glycol and water 148
 - stability to acids, heat, light and sulphur dioxide 148
 - suppliers 176
- Suppliers, of colours 171-80
- Synthetic colours 87-157
 - suppliers 173-80
- Tablet coatings, use of riboflavin 84
- Tartrazine 151-3
 - ADI 153
 - analytical methods 153
 - applications 152
 - chemical structure (fig.) 151
 - colour shade 151
 - effect of pH 151
 - in food-intolerance studies 153
 - physical characteristics 151-2
 - physiological properties 153
 - solubility in ethanol, oil, propylene glycol and water 152
 - stability to acids, heat, light and sulphur dioxide 151-2
 - suppliers 176
 - susceptibility to oxidation 151-2
- Titanium dioxide 155-7
 - ADI 156
 - analytical methods 156-7
 - applications 156
 - colour shade 155
 - effect of pH 156
 - physical characteristics 156
 - physiological properties 156
 - solubility in acids, oil and water 156
 - source 155
 - stability to heat and light 156
 - suppliers 176
- Tomatoes, as source of lycopenc 75
- Turmeric – see curcumin
 - suppliers 173
- UK legislation 166-7

- specific labelling requirements for foods containing colours 167
- USA legislation 167-9
 - specific labelling requirements for foods containing colours 169
- Vegetable Carbon – see Carbon Black
- Vitamin B2 – see riboflavin
- Water ices, use of anthocyanins 13
 - use of beta-carotene 24
 - use of Carbon Black 108
 - use of curcumin 62, 63
 - use of lutein 71
- White shade, of titanium dioxide 157
- Xanthophylls – see lutein
- Yellow 2G 176-80
- Yellow fats, use of beta-carotene 25
 - use of carotenes 28
- Yellow shade, of annatto 4
 - of carotenes 27
 - of copper complexes of chlorophyllins 52, 54
 - of crocin 57
 - of curcumin 61
 - of iron oxides and hydroxides 129
 - of lutein 69
 - of lycopene 75
 - of Quinoline Yellow 143
 - of riboflavin 83
 - of tartrazine 151
- Yellow/orange shade, of beta-carotene 21
 - of ethyl ester of beta-apo-8'carotenoic acid 65
 - of Sunset Yellow FCF 147
- Yoghurt, use of beta-carotene 25
 - use of carmine 44
 - use of copper complexes of chlorophyllins 54