


JOURNAL OF

CHROMATOGRAPHY

INTERNATIONAL JOURNAL ON CHROMATOGRAPHY, ELECTROPHORESIS AND RELATED METHODS


 EDITOR, Michael Lederer (Switzerland)

ASSOCIATE EDITORS, R. W. Frei (Amsterdam), R. W. Giese (Boston, MA), J. K. Haken (Kensington, N.S.W.), K. Macek (Prague), L. R. Snyder (Orinda, CA)

EDITOR, SYMPOSIUM VOLUMES, E. Heftmann (Orinda, CA)

EDITORIAL BOARD

W. A. Aue (Halifax)
 V. G. Berezkin (Moscow)
 V. Betina (Bratislava)
 A. Bevenue (Belmont, CA)
 P. Boček (Brno)
 P. Boulanger (Lille)
 A. A. Boulton (Saskatoon)
 G. P. Cartoni (Rome)
 S. Dilli (Kensington, N.S.W.)
 L. Fishbein (Washington, DC)
 A. Frigerio (Milan)
 C. W. Gehrke (Columbia, MO)
 E. Gil-Av (Rehovot)
 G. Guiochon (Knoxville, TN)
 I. M. Hais (Hradec Králové)
 S. Hjertén (Uppsala)
 E. C. Horning (Houston, TX)
 Cs. Horváth (New Haven, CT)
 J. F. K. Huber (Vienna)
 A. T. James (Harrold)
 J. Janák (Brno)
 E. sz. Kováts (Lausanne)
 K. A. Kraus (Oak Ridge, TN)
 E. Lederer (Gif-sur-Yvette)
 A. Liberti (Rome)
 H. M. McNair (Blacksburg, VA)
 Y. Marcus (Jerusalem)
 G. B. Marini-Bettolo (Rome)
 A. J. P. Martin (Cambridge)
 Č. Michalec (Prague)
 R. Neher (Basel)
 G. Nickless (Bristol)
 N. A. Parris (Wilmington, DE)
 R. L. Patience (Sunbury-on-Thames)
 P. G. Righetti (Milan)
 O. Samuelson (Göteborg)
 R. Schwarzenbach (Dübendorf)
 A. Zlatkis (Houston, TX)

EDITORS, BIBLIOGRAPHY SECTION

Z. Deyl (Prague), J. Janák (Brno), V. Schwarz (Prague), K. Macek (Prague)

ELSEVIER

Scope. The *Journal of Chromatography* publishes papers on all aspects of chromatography, electrophoresis and related methods. Contributions consist mainly of research papers dealing with chromatographic theory, instrumental development and their applications. The section *Biomedical Applications*, which is under separate editorship, deals with the following aspects: developments in and applications of chromatographic and electrophoretic techniques related to clinical diagnosis or alterations during medical treatment; screening and profiling of body fluids or tissues with special reference to metabolic disorders; results from basic medical research with direct consequences in clinical practice; drug level monitoring and pharmacokinetic studies; clinical toxicology; analytical studies in occupational medicine.

Submission of Papers. Papers in English, French and German may be submitted, in three copies. Manuscripts should be submitted to: The Editor of *Journal of Chromatography*, P.O. Box 681, 1000 AR Amsterdam, The Netherlands, or to: The Editor of *Journal of Chromatography, Biomedical Applications*, P.O. Box 681, 1000 AR Amsterdam, The Netherlands. Review articles are invited or proposed by letter to the Editors. An outline of the proposed review should first be forwarded to the Editors for preliminary discussion prior to preparation. Submission of an article is understood to imply that the article is original and unpublished and is not being considered for publication elsewhere. For copyright regulations, see below.

Subscription Orders. Subscription orders should be sent to: Elsevier Science Publishers B.V., P.O. Box 211, 1000 AE Amsterdam, The Netherlands, Tel. 5803 911, Telex 18582 ESPA NL. The *Journal of Chromatography* and the *Biomedical Applications* section can be subscribed to separately.

Publication. The *Journal of Chromatography* (incl. *Biomedical Applications* and *Cumulative Author and Subject Indexes, Vols. 401-450*) has 37 volumes in 1988. The subscription prices for 1988 are:

J. Chromatogr. (incl. *Cum. Indexes, Vols. 401-450*) + *Biomed. Appl.* (Vols. 424-460):

Dfl. 6290.00 plus Dfl. 962.00 (p.p.h.) (total ca. US\$ 3920.00)

J. Chromatogr. (incl. *Cum. Indexes, Vols. 401-450*) only (Vols. 435-460):

Dfl. 5070.00 plus Dfl. 676.00 (p.p.h.) (total ca. US\$ 3106.00)

Biomed. Appl. only (Vols. 424-434):

Dfl. 2145.00 plus Dfl. 286.00 (p.p.h.) (total ca. US\$ 1314.00).

Our p.p.h. (postage, package and handling) charge includes surface delivery of all issues, except to subscribers in Argentina, Australia, Brasil, Canada, China, Hong Kong, India, Israel, Malaysia, Mexico, New Zealand, Pakistan, Singapore, South Africa, South Korea, Taiwan, Thailand and the U.S.A. who receive all issues by air delivery (S.A.L. — Surface Air Lifted) at no extra cost. For Japan, air delivery requires 50% additional charge; for all other countries airmail and S.A.L. charges are available upon request. Back volumes of the *Journal of Chromatography* (Vols. 1 through 423) are available at Dfl. 230.00 (plus postage). Claims for missing issues will be honoured, free of charge, within three months after publication of the issue. Customers in the U.S.A. and Canada wishing information on this and other Elsevier journals, please contact Journal Information Center, Elsevier Science Publishing Co. Inc., 52 Vanderbilt Avenue, New York, NY 10017. Tel. (212) 916-1250.

Abstracts/Contents Lists published in Analytical Abstracts, ASCA, Biochemical Abstracts, Biological Abstracts, Chemical Abstracts, Chemical Titles, Chromatography Abstracts, Current Contents/Physical, Chemical & Earth Sciences, Current Contents/Life Sciences, Deep-Sea Research/Part B: Oceanographic Literature Review, Excerpta Medica, Index Medicus, Mass Spectrometry Bulletin, PASCAL-CNRS, Referativnyi Zhurnal and Science Citation Index.

See inside back cover for Publication Schedule, Information for Authors and information on Advertisements.

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, without the prior written permission of the publisher, Elsevier Science Publishers B.V., P.O. Box 330, 1000 AH Amsterdam, The Netherlands.

Upon acceptance of an article by the journal, the author(s) will be asked to transfer copyright of the article to the publisher. The transfer will ensure the widest possible dissemination of information.

Submission of an article for publication entails the authors' irrevocable and exclusive authorization of the publisher to collect any sums or considerations for copying or reproduction payable by third parties (as mentioned in article 17 paragraph 2 of the Dutch Copyright Act of 1912 and in the Royal Decree of June 20, 1974 (S. 351) pursuant to article 16 b of the Dutch Copyright Act of 1912) and/or to act in or out of Court in connection therewith.

Special regulations for readers in the U.S.A. This journal has been registered with the Copyright Clearance Center, Inc. Consent is given for copying of articles for personal or internal use, or for the personal use of specific clients. This consent is given on the condition that the copier pays through the Center the per-copy fee stated in the code on the first page of each article for copying beyond that permitted by Sections 107 or 108 of the U.S. Copyright Law. The appropriate fee should be forwarded with a copy of the first page of the article to the Copyright Clearance Center, Inc., 27 Congress Street, Salem, MA 01970, U.S.A. If no code appears in an article, the author has not given broad consent to copy and permission to copy must be obtained directly from the author. All articles published prior to 1980 may be copied for a per-copy fee of US\$ 2.25, also payable through the Center. This consent does not extend to other kinds of copying, such as for general distribution, resale, advertising and promotion purposes, or for creating new collective works. Special written permission must be obtained from the publisher for such copying.

No responsibility is assumed by the Publisher for any injury and/or damage to persons or property as a matter of products liability, negligence or otherwise, or from any use or operation of any methods, products, instructions or ideas contained in the materials herein. Because of rapid advances in the medical sciences, the Publisher recommends that independent verification of diagnoses and drug dosages should be made. Although all advertising material is expected to conform to ethical (medical) standards, inclusion in this publication does not constitute a guarantee or endorsement of the quality or value of such product or of the claims made of it by its manufacturer.

CONTENTS

Bibliography Section

Liquid Column Chromatography	B121
Gas Chromatography	B189
Planar Chromatography	B222
Electrophoresis	B242

ห้องสมุดกรมวิทยาศาสตร์บริการ
10 พค ๑๙๘๘

Bibliography Section

Liquid Column Chromatography

1. REVIEWS AND BOOKS

- 784 Belen'kii, B.G., Gankina, E.S. and Ml'tsev, V.G.: *Capillary Liquid Chromatography*. Nauka, Leningrad, 1987, 207 p.; *C.A.*, 107 (1987) 205900t.
- 785 Chretien, J.R.: Chemometrics in chromatography. *TrAC*, 6 (1987) 275-278 - a review with 22 refs.
- 786 Hattori, K.: Host-guest chromatography. *Bio Ind.*, 4 (1987) 327-338; *C.A.*, 107 (1987) 167984p - a review with 75 refs.
- 787 Horvath, C.: Interference by on-column reactions in HPLC. *Symp. Biol. Hung.*, 34 (1986) 1-18; *C.A.*, 107 (1987) 228083t - a review with 26 refs.
- 788 Kasai, H.: (High performance liquid chromatography). *Gendai Kagaku*, 197 (1987) 35-41; *C.A.*, 107 (1987) 194304y - a review with 3 refs.
- 789 Kubeczka, K.-H.: Analyse von Naturstoffen. Neue Entwicklungen. *Dtsch. Apoth.-Ztg.*, 127 (1987) 2443-2450.
- 790 Lawrence, J.G.: Chromatography. *Annu Rep. Prog. Chem., Sect. C*, 83 (1987) 177-196; *C.A.*, 107 (1987) 205621c - a review with 160 refs.
- 791 Maeda, M.: (Purification by extraction with solid phase). *Nippon Koshohin Kagakkaishi*, 11 (1987) 117-123; *C.A.*, 107 (1987) 214239m - a review with 7 refs.
- 792 Maehr, H.: (Practical applications of the Craig distribution. Part 2). *GIT Fachz. Lab.*, 31 (1987) 549-556; *C.A.*, 107 (1987) 162688k - a review with no refs.
- 793 Maehr, H.: (Practical applications of the Craig distribution. Part 3). *GIT Fachz. Lab.*, 31 (1987) 626-633; *C.A.*, 107 (1987) 162687j - a review with no refs.
- 794 Matsushita, T. and Shono, T.: (Simultaneous determination of many trace metals by using dithiocarbamate complexes). *Kagaku (Kyoto)*, 42 (1987) 496-497; *C.A.*, 107 (1987) 167777j - a review with 15 refs.
- 795 Parvez, H., Bastart-Malsot, M., Parvez, S., Nagatsu, T. and Carpentier, G. (Editors): *Electrochemical Detection in Medicine and Chemistry (Progress in HPLC, Vol. 2)*. VNU Science Press, Utrecht, 1987, XI + 472 p.
- 796 Rehak, V.: (Liquid chromatography in analytical practice. I. Instrumentation). *Chem. Prum.*, 37 (1987) 436-439; *C.A.*, 107 (1987) 130117g - a review with 9 refs.
- 797 Takei, N. and Low, D.: (Modern chromatographic methods and future prospects in downstream). *Bio Ind.*, 4 (1987) 407-414; *C.A.*, 107 (1987) 174216t - a review with 7 refs.
- 798 Wankat, P.C.: *Large-Scale Adsorption and Chromatography, Vol. 1*. CRC Press, Boca Raton, 1986, 256 p.; *C.A.*, 107 (1987) 157471e.
- 799 Wankat, P.C.: *Large-Scale Adsorption and Chromatography, Vol. 2*. CRC Press, Boca Raton, 1986, 256 p.; *C.A.*, 107 (1987) 157472f.
- See also 837, 844, 846, 851-853, 878, 879, 886, 889, 910, 911, 915, 919, 922, 924, 927, 1027, 1031, 1068, 1092, 1119, 1155, 1156, 1279, 1280, 1335, 1378, 1385, 1448, 1492, 1505, 1529, 1537, 1576, 1588, 1598, 1636, 1659, 1678, 1806, 1823, 1831.

2. FUNDAMENTALS, THEORY AND GENERAL

2a. General

- 800 Aranovich, G.L.: (Calculation of component distribution in a chromatographic column with incomplete separation of peaks). *Teor. Osn. Khim. Tekhnol.*, 21 (1987) 448-454; *C.A.*, 107 (1987) 184345y.

- 801 Bicking, M.K.L. and Serwon, S.J.: Molecular interactions on polystyrene stationary phases. *J. Liq. Chromatogr.*, 10 (1987) 1369-1382.
- 802 Dolan, J.W., Snyder, L.R. and Quarry, M.A.: HPLC method development and column reproducibility. *Am. Lab. (Fairfield)*, 19 (1987) 43-47; *C.A.*, 107 (1987) 190073z.
- 803 Ganetsos, G.: Prediction of the distribution coefficient (K_d) variation with operating conditions in chromatographic systems. *J. Chromatogr.*, 411 (1987) (1987) 81-94.
- 804 Gorse, J. and Burke, M.F.: The effects of an ionic mobile phase modifier and thermal history on a reversed-phase liquid chromatographic system. *J. Liq. Chromatogr.*, 10 (1987) 1049-1064.
- 805 Hwu, J.R., Robl, J.A. and Khoudary, K.P.: Instrumentation and separation results of medium pressure liquid chromatography. *J. Chromatogr. Sci.*, 25 (1987) 501-505.
- 806 Mulder, R., Hoogerbrugge, P.J. and Smit, H.C.: Deterministic noise in correlation chromatography; origins and properties of ghost peaks. *Chemom. Intell. Lab. Syst.*, 1 (1987) 243-264; *C.A.*, 107 (1987) 168007j.
- 807 Nielen, M.W.F., van Ingen, H.E., Valk, A.J., Frei, R.W. and Brinkman, U.A.T.: Metal-loaded sorbents for selective on-line sample handling and trace enrichment in liquid chromatography. *J. Liq. Chromatogr.*, 10 (1987) 617-633.
- 808 Oscik-Mendyk, B. and Rozylo, J.K.: Analysis of chromatographic parameters in the systems with ternary mobile phases. *J. Liq. Chromatogr.*, 10 (1987) 1399-1415.
- 809 Salov, V.V., Timerbaev, A.R., Petrukhin, O.M. and Zolotov, Yu.A.: (General features of interphase distribution of compounds under conditions of extraction and chromatography). *Dokl. Akad. Nauk SSSR*, 295 (1987) 918-922; *C.A.*, 107 (1987) 184681e.
- 810 Shatts, V.I., Sakkartova, O.V. and Brivkalne, L.: (General principle of chromatographic behavior in systems with different sorption mechanisms). *Latv. PSR Zinat. Akad. Vestis, Kim. Ser.*, (1987) 459-466; *C.A.*, 107 (1987) 184346z.
- For additional information see:
C.A., 107 (1987) 76929h.

See also 839, 980.

2b. Thermodynamics and theoretical relationships

- 811 Antia, F.D. and Horvath, C.: High-performance liquid chromatography at elevated temperatures: examination of conditions for the rapid separation of large molecules. *J. Chromatogr.*, 435 (1988) 1-15.
- 812 Chen, S. and Peng, S.: A generalized equation for HETP in column chromatography. *Set. Sin., Ser. B*, 30 (1987) 572-581; *C.A.*, 107 (1987) 167822j.
- 813 Cheong, W.J. and Carr, P.W.: The surface tension of mixtures of methanol, acetonitrile, tetrahydrofuran, isopropanol, tertiary butanol and dimethylsulfide with water at 25°C. *J. Liq. Chromatogr.*, 10 (1987) 561-581.
- 814 Gago, F., Alvarez-Builla, J. and Elguero, J.: Hydrophobicity measurements by HPLC: a new approach to π constants. *J. Liq. Chromatogr.*, 10 (1987) 1031-1047.
- 815 Jaroniec, M. and Martire, D.E.: Description of non-specific and specific solute-solvent interactions in liquid chromatography with mixed eluents. *J. Liq. Chromatogr.*, 10 (1987) 541-559.
- 816 Kolosov, N.V. and Zolotarev, P.: (Theory of chromatography in capillary columns of arbitrary cross sectional shapes). *Zh. Fiz. Khim.*, 61 (1987) 1981-1985; *C.A.*, 107 (1987) 205707k.
- 817 Lamparczyk, H., Atomura, M. and Jinno, K.: Qualitative description of dispersive and inductive electrostatic interactions in reversed-phase liquid chromatography. *Chromatographia*, 23 (1987) 752-759.
- 818 Moriyasu, M., Kato, A. and Hashimoto, Y.: Kinetic studies of fast equilibrium by means of high-performance liquid chromatography. XVI. Separation of tautomers of β -diketones and β -diketoamides. *J. Chromatogr.*, 411 (1987) 466-471.
- 819 Moriyasu, M., Yamagami, C., Kato, A., Hashimoto, Y. and Takao, N.: Kinetic studies of fast equilibrium by means of HPLC. XIV. Separation of rotamers of phenyl methylcarbamates and their rotational energy barriers. *Bull. Chem. Soc. Jpn.*, 59 (1986) 1539-1543; *C.A.*, 107 (1987) 77028a.
- 820 Petrovic, S.M., Lomic, S.M. and Sefer, I.: Solute retention in the stationary phase of a liquid-solid chromatographic system. *Chromatographia*, 23 (1987) 915-924.

- 821 Soczewinski, E. and Dzido, T.: Adsorption equilibria of polar aromatic solutes in systems consisting of silica and a binary mixture of heptane and a polar modifier. *Symp. Biol. Hung.*, 34 (1986) 485-489; *C.A.*, 107 (1987) 211212m.
- 822 Waldmann, Meyer, H.: Structure parameters of molecules and media evaluated by chromatographic partition. II. Geometrical exclusion in gels. *J. Chromatogr.*, 410 (1987) 233-248.

See also 801, 808, 884, 960, 1145.

2c. Relationship between structure and chromatographic behaviour

- 823 Karunanithy, R., Leung, S.L. and Yeo, S.H.: Use of high-performance liquid chromatography for quantitative structure - activity relationship studies of a series of potential contraceptive steroidal esters. *J. Pharm. Biomed. Anal.*, 5 (1987) 597-604.
- 824 Sates, V.: Relationship between retention and organic solvent molarity in reverse-phase HPLC. *Symp. Biol. Hung.*, 34 (1986) 579-590; *C.A.*, 107 (1987) 228105b.

For additional information see:

C.A., 107 (1987) 133950j, 183668u, 205791h.

See also 821, 926, 947, 955, 1732.

2d. Measurement of physico-chemical and related values

- 825 Cuellar, J. and Bravo, M.C.: Determination of adsorption isotherms by chromatographic methods. *Tec. Lab.*, 11 (1987) 132-138; *C.A.*, 107 (1987) 205670t.
- 826 Lang, P. and Burchard, W.: Determination of branching by gel size-exclusion chromatography in combination with on-line small angle light scattering. *Macromol. Chem., Rapid Commun.*, 8 (1987) 451-455; *C.A.*, 107 (1987) 155101k.

For additional information see:

C.A., 107 (1987) 155090f, 176838w, 176884h, 218381z, 219616s.

See also 1330.

3. GENERAL TECHNIQUES

3a. Apparatus and accessories

- 827 Fujiwara, M., Ishida, Y., Nimura, N., Toyama, A. and Kinoshita, T.: Postcolumn fluorometric detection system for liquid chromatographic analysis of amino and imino acids using *o*-phthalaldehyde/N-acetyl-L-cysteine reagent. *Anal. Biochem.*, 166 (1987) 72-78.
- 828 Hobbs, J.S.: An application-oriented approach to HPLC instrumentation. *Am. Lab. (Fairfield)*, 19 (1987) 48-50; *C.A.*, 107 (1987) 130144p.
- 829 Ito, Y.: Cross-axis synchronous flow-through coil planet centrifuge free of rotary seals for preparative countercurrent chromatography. Part I. Apparatus and analysis of acceleration. *Sep. Sci. Technol.*, 22 (1987) 1971-1987; *C.A.*, 107 (1987) 178744m.
- 830 Jones, W.R., Heckenberg, A.L. and Jandik, P.: Ion chromatography method and apparatus. *Eur. Pat. Appl.* EP 230,307 (Cl. G01N30/96), 29 Jul. 1987, US Appl. 821,154, 21 Jan. 1986; 34 pp.; *C.A.*, 107 (1987) 167961d.
- 831 Morev, S.N., Yanotovskii, M.T. and Klyachko, Yu.A.: (Study on elastic columns used in high-performance liquid chromatography. III. Application of elastic columns in analytical practice). *Khim.-Farm. Zh.*, 21 (1987) 1004-1008; *C.A.*, 107 (1987) 183666s.
- 832 Moskvina, L.N., Bumerov, M.F. and Gorshkov, A.I.: (Multiple-column chromatograph for continuous separation of mixtures). *U.S.S.R. Pat.* SU 1,236,893 (Cl. B01N30/46), 15 Feb. 1987, Appl. 3,504,382, 26 Oct. 1982; *C.A.*, 107 (1987) 167948e.

- 833 Selavka, C.M., Jiao, K.-S. and Krull, I.S.: Construction and comparison of open tubular reactors for postcolumn reaction detection in liquid chromatography. *Anal. Chem.*, 59 (1987) 2221-2224.

For additional information see:

C.A., 107 (1987) 157499v, 168004f, 189980m.

See also 805, 898, 913.

3b. Detectors and detection reagents

- 834 Ciurczak, E.W. and Weis, F.M.B.: Evaluation of a near-infrared detector for high performance liquid chromatography. *Spectroscopy (Eugene)*, 2 (1987) 33-38; *C.A.*, 107 (1987) 228096z.
- 835 De Wit, J.S.M. and Jorgenson, J.W.: Photoionization detector for open-tubular liquid chromatography. *J. Chromatogr.*, 411 (1987) 201-212.
- 836 Ebel, S., Kühnert, H. and Mück, W.: Limits of determination and calibration in HPLC. *Chromatographia*, 23 (1987) 934-938.
- 837 Froehlich, P.: Fluorescence detection in HPLC. *BioChromatography*, 2 (1987) 144-147; *C.A.*, 107 (1987) 190018k - a review with 4 refs.
- 838 Hashimoto, Y. and Maruyama, Y.: Recent advances in electrochemical detector for highly selective and sensitive analysis of biogenic compounds. *Prog. HPLC*, 2 (Electrochem. Detect. Med. Chem.) (1987) 427-466; *C.A.*, 107 (1987) 127258y - a review with 90 refs.
- 839 Hayashi, Y., Shibazaki, T. and Uchiyama, M.: Resolution of overlapped chromatograms by the use of a Kremen filter. III. Effects of overlapping and outstripping of a solvent peak on the analysis. *J. Chromatogr.*, 411 (1987) 95-105.
- 840 Hudson, M.K. and Busch, K.W.: Infrared emission from a flame as the basis for chromatographic detection of organic compounds. *Anal. Chem.*, 59 (1987) 2603-2609.
- 841 Ji, H. and Wang, E.: A thin-layer electrochemical detector coated with nafion film for liquid chromatography. *J. Chromatogr.*, 410 (1987) 111-120.
- 842 Kientz, C.E. and Verweij, A.: The application of a flame photometric detector in packed microcapillary liquid chromatography: detection of organophosphates. *Int. J. Environ. Anal. Chem.*, 30 (1987) 255-263; *C.A.*, 107 (1987) 211269k.
- 843 Koizumi, S., Imato, T. and Ishibashi, N.: Anion-selective electrode for high-performance liquid chromatography detector based on oleophilic anion-exchange resin membrane. *Anal. Sci.*, 3 (1987) 319-325; *C.A.*, 107 (1987) 211113e.
- 844 Kurahashi, T., Nishino, H., Parvez, S., Parvez, H., Kojima, K. and Nagatsu, T.: Biochemical and pharmacological application of electrochemical detection in HPLC. *Prog. HPLC*, 2 (Electrochem. Detect. Med. Chem.) (1987) 3-36; *C.A.*, 107 (1987) 171634s.
- 845 Maris, F.A., Nijenhuis, M., Frei, R.W., DeJong, G.J. and Brinkman, U.A.T.: On-line post-column extraction in column liquid chromatography with electron-capture detection. II. Applications. *J. Chromatogr.*, 435 (1988) 297-306.
- 846 McClintock, S.A. and Purdy, W.C.: Developments in electrochemical detection for liquid chromatography. *Prog. HPLC*, 2 (1987) 37-51; *C.A.*, 107 (1987) 190010b - a review with many refs.
- 847 Nagy, J., Toth, G.K. and Kovacs, K.: A low-cost computing integrator system for HPLC. *Symp. Biol. Hung.*, 34 (1986) 639-646; *C.A.*, 107 (1987) 211222q.
- 848 Plitzko, B.: Detector for liquid and high-pressure liquid chromatography in an indirect operating method. *Ger. (East) Pat.* DD 233,192, 19 Feb. 1986, Appl. 265,436, 19 Jul. 1984, 3 pp.; *C.A.*, 107 (1987) 228042d.
- 849 Ramstad, T. and Weaver, M.J.: Electrosorptive detection in liquid chromatography of simple charged organic compounds. *Chromatographia*, 23 (1987) 883-887.
- 850 Ramstad, T. and Weaver, M.J.: Thin-layer cell with a rapidly dropping mercury electrode for liquid chromatographic detection. *Chromatographia*, 23 (1987) 878-882.
- 851 Sievers, R.E., Nyarady, S.A., Barak, S., Banning, S.S. and Hutte, R.S.: Redox chemiluminescence detector (RCD): Successes in gas chromatography an prospects for liquid chromatography. *Symp. Biol. Hung.*, 34 (1986) 673-677; *C.A.*, 107 (1987) 228084u - a review with 14 refs.

- 852 Slais, K.: Electrochemical detectors for low-dispersion liquid chromatography. *Prog. HPLC*, 2 (1987) 53-94; *C.A.*, 107 190011c - a review with many refs.
- 853 Stulik, K. and Pacakova, V.: Some theoretical and practical aspects of electrochemical detection in high-performance liquid chromatography. *Prog. HPLC*, 2 (1987) 131-178; *C.A.*, 107 (1987) 190012d - a review with many refs.
- 854 Styring, M.G., Armonas, J.E. and Hamielec, A.E.: An experimental evaluation of a new commercial viscometric detector for size-exclusion chromatography (SEC) using linear and branched polymers. *J. Liq. Chromatogr.*, 10 (1987) 783-804.
- 855 Takeuchi, T. and Ishii, D.: Signal enhancement by on-column fluorometric detection in high-performance liquid chromatography using micropacked fused-silica columns. *J. Chromatogr.*, 435 (1988) 319-325.

For additional information see:

C.A., 107 (1987) 190091d, 190092e.

See also 938, 945, 992, 1153, 1164, 1568, 1808, 1817, 1864.

3c. Sorbents, carriers, column and layer performance, packing procedures

- 856 Andreolini, F., Borra, C. and Novotny, M.: Preparation and evaluation of slurry-packed capillary columns for normal-phase liquid chromatography. *Anal. Chem.*, 59 (1987) 2428-2432.
- 857 Antakli, S.C. and Serpinet, J.: Determination of the concentration of silanol groups by a chemical reaction with methylolithium and GC measurements of evolved methane. *Chromatographia*, 23 (1987) 767-769.
- 858 Carr, J.W. and Harris, J.M.: Heterogeneity of reversed-phase chromatographic surfaces: quenching of sorbed pyrene fluorescence. *Anal. Chem.*, 59 (1987) 2546-2550.
- 859 Carunchio, V., Girelli, A.M., Messina, A. and Sinibaldi, M.: Chitosan-coated silica gel as a new support in high-performance liquid chromatography. *Chromatographia*, 23 (1987) 731-735.
- 860 Cortes, H.J., Pfeiffer, C.D., Stevens, T.S. and Richter, B.E.: Chromatography columns with cast porous plugs and methods of fabricating them. *Eur. Pat. Appl. EP 231,684 (Cl. G01N30/56)*, 12 Aug. 1987, US Appl. 807,729, 11 Dec. 1985; 8 pp.; *C.A.*, 107 (1987) 190132t.
- 861 Danielson, N.D. and Kirkland, J.J.: Synthesis and characterization of 2 μ m wide-pore silica microspheres as column packings for the reversed-phase liquid chromatography of peptides and proteins. *Anal. Chem.*, 59 (1987) 2501-2506.
- 862 De Weerd, M., Dewaele, C. and Verzele, M.: On the stability of micro-LC columns (packed fused silica capillary columns). *J. High Resolut. Chromatogr. Chromatogr. Commun.*, 10 (1987) 553-556.
- 863 Delaney, M.F., Papas, A.N. and Walters, M.J.: Chemometric classification of reversed-phase high-performance liquid chromatography columns. *J. Chromatogr.*, 410 (1987) 31-41.
- 864 Fabian, G., Schierhorn, M., Poerschmann, J., Kraus, G., Altmann, H. and Zschke, H.: (Preparation of alkenoyl-*tert.*-butylvalinamides as a chiral chromatography media). *Ger. Pat. DD 241,410 (Cl. C07C103/50)*, 10 Dec. 1986, Appl. 281,258, 01 Oct. 1985; 11 pp.; *C.A.*, 107 (1987) 198927b.
- 865 Gangoda, M., Gilpin, R.K. and Fung, B.M.: Effect of temperature on reversed-phase chromatographic surfaces studied by carbon-13 NMR. *J. Magn. Reson.*, 74 (1987) 134-138; *C.A.*, 107 (1987) 184322p.
- 866 Issaq, H.J., Fox, S.D., Lindsey, K., McConnell, J.H. and Weiss, D.E.: Effect of temperature on HPLC separations using C₁, C₄, C₈ and C₁₈ alkyl chain bonded silica columns. *J. Liq. Chromatogr.*, 10 (1987) 49-70.
- 867 Issaq, H.J., Glennon, M.L., Weiss, D.E. and Fox, S.D.: High-performance liquid chromatography using a β -cyclodextrin-bonded silica column: effect of temperature on retention. *ACS Symp. Ser.*, 342 (1987) 260-271; *C.A.*, 107 (1987) 190064x.
- 868 Iwaki, K., Yoshida, S., Nimura, N., Kinoshita, T., Takeda, K. and Ogura, H.: Preparation of chiral stationary phase via activated carbanate intermediate for liquid chromatographic optical resolution. *Chromatographia*, 23 (1987) 727-730.
- 869 Jost, W., Krebs, K.-F. and Winkler, H.: LiChrospher® 60 RP-select B - ein neuer HPLC-Träger zur Trennung basischer Pharmawirkstoffe. *Kontakte (Darmstadt)*, No. 2 (1987) 32-36.

- 870 Kiniwa, H., Doi, Y., Nishikaji, T. and Ogata, N.: Poly(α -amino acid)-immobilized polymer adsorbents for optical resolution. 2. Effect of crosslinking degree of the support resin on the resolution efficiency. *Macromol. Chem.*, 188 (1987) 1841-1850; *C.A.*, 107 (1987) 168021j.
- 871 Kiniwa, H., Doi, Y., Nishikaji, T. and Ogata, N.: Poly(α -amino acid)-immobilized polymer adsorbents for optical resolution. 3. A total chromatographic optical resolution on poly(N^5 -benzyl-L-glutamine)-immobilized resin. *Macromol. Chem.*, 188 (1987) 1851-1860; *C.A.*, 107 (1987) 190072y.
- 872 Kolosov, N.V. and Zolotarev, P.P.: Efficiency of chromatographic columns of different cross-sectional forms. *Zh. Fiz. Khim.*, 61 (1987) 2228-2232; *C.A.*, 107 (1987) 205879t.
- 873 Mistryukov, E.A., Andreev, S.P. and Rozpravka, J.: Method of manufacturing glass-capillary chromatographic columns. *U.S.S.R. Pat.* SU 1,323,952 (Cl. G01N30/50), 15 Jul. 1987, Appl. 4,015,377, 21 Jan. 1986; *C.A.*, 107 (1987) 211298u.
- 874 Morev, S.N., Yanotovskii, M.T. and Klyachko, Yu.A.: (Study of elastic columns designed for high-performance liquid chromatography. II. Hydrodynamic and chromatographic characterizations). *Khim.-Farm. Zh.*, 21 (1987) 883-887; *C.A.*, 107 (1987) 168000b.
- 875 Morii, N., Suzuki, Y. and Ichimura, K.: (Optically active silane coupling agents). *Jpn. Kokai Tokkyo Koho Pat.* JP 62 51,689 (87 51,689) (Cl. C07F7/18), 06 Mar. 1987, Appl. 85/190,397, 29 Aug. 1985, 3 pp.; *C.A.*, 107 (1987) 198655m.
- 876 Nielsen, R.G. and Wilson, G.S.: Characterization of adsorption on the stationary phase using high-performance immunoaffinity chromatography. *J. Chromatogr.*, 423 (1987) 41-50.
- 877 Nilsson, K.G.C.: Macroporous particles. *Eur. Pat. Appl.* EP 222,718 (Cl. C08J3/12), 20 May 1987, SE Appl. 85/4,764, 15 Oct. 1985; 7 pp.; *C.A.*, 107 (1987) 199807z.
- 878 Okamoto, Y.: (Polymers having optical resolution power). *Kobunshi*, 36 (1987) 418-421; *C.A.*, 107 (1987) 150457w - a review with 27 refs.
- 879 Okuyama, T., Ogawa, T. and Ebihara, M.: (Apatite used for chromatography). *Gypsum Lime*, 210 (1987) 323-331; *C.A.*, 107 (1987) 214230b - a review with no refs.
- 880 Pietrogrande, M.C., Dondi, F., Blo, G., Borea, P.A. and Bigli, C.: Octadecyl, phenyl, and cyano phases comparison for the RP-HPLC prediction of octanol-water partition coefficient. *J. Liq. Chromatogr.*, 10 (1987) 1065-1075.
- 881 Porsch, B. and Novakova, J.: (Silica gel preparation with chemically bonded epoxy group for chromatographic separation of proteins). *Czech. Pat.*, CS 243, 336 (Cl. B01J20/10), 15 May 1987, Appl. 85/1,042, 14 Feb. 1985; 4 pp.; *C.A.*, 107 (1987) 201371y.
- 882 Sadek, P.C., Koester, C.J. and Bowers, L.D.: The investigation of the influence of trace metals in chromatographic retention processes. *J. Chromatogr. Sci.*, 25 (1987) 489-493.
- 883 Shelly, D.C. and Edkins, T.J.: Contribution to the packing of liquid chromatographic microcolumns. *J. Chromatogr.*, 411 (1987) 185-199.
- 884 Smith, P.L. and Cooper, W.T.: Retention and selectivity in amino, cyano and diol normal bonded phase high-performance liquid chromatographic columns. *J. Chromatogr.*, 410 (1987) 249-265.
- 885 Street, K.W., Jr.: Cyclodextrin cavity polarity and chromatographic implications. *J. Liq. Chromatogr.*, 10 (1987) 655-662.
- 886 Szejtli, J.: Application of cyclodextrins in chromatography. *Starch/Staerke*, 39 (1987) 357-362; *C.A.*, 107 (1987) 214263q - a review with 49 refs.
- 887 Tennikova, T.B., Horak, D., Svec, F., Kolar, J., Coupek, J., Trushin, S.A., Maltzev, V.G. and Belenkii, B.G.: Hydrolysed macroporous glycidyl methacrylate-ethylene dimethacrylate copolymer sorbent for size-exclusion high-performance liquid chromatography of synthetic polymers and biopolymers. *J. Chromatogr.*, 435 (1988) 357-362.
- 888 Tindall, G.W.: Bonded cyclodextrin stationary phase columns for the separation of *cis/trans* cyclohexane derivatives. *J. Liq. Chromatogr.*, 10 (1987) 1077-1084.
- 889 Unger, K.K.: Design and characteristics of adsorbents for the separation of biopolymers by chromatographic processes. *Proc. Eng. Found. Conf. Fundam. Adsorpt.*, 2 (1986, Pub. 1987) 579-588; *C.A.*, 107 (1987) 150463v - a review with 24 refs.
- 890 Viska, J., Slavikova, A. and Vykoupil, J.: (Dispersion of organic monomer phase containing hydroxyalkyl methacrylates). *Czech Pat.* CS 233,520 (Cl. C08F2/18), 15 Aug. 1986, Appl. 83/24, 03 Jan. 1983; 5 pp.; *C.A.*, 107 (1987) 218607c.

- 891 Wojcik, A. and Kwietniewski, L.: Preparation of porous bead silica. *J. Chromatogr.*, 435 (1988) 55-62.
- 892 Yasukawa, K., Tamura, Y., Uchida, T., Yanagihara, Y. and Noguchi, K.: Development of a C₁₈-bonded vinyl alcohol copolymer gel for reversed-phase high-performance liquid chromatography. *J. Chromatogr.*, 410 (1987) 129-136.

For additional information see:

C.A., 107 (1987) 130500b, 155732s, 171508d, 176670k, 177150c, 178430f, 178800b, 179286a, 190093f, 190094g, 194590p, 200796d, 201456e, 206029c, 214169p, 218865k, 219566a, 223820w.

See also 905, 906, 918, 926, 943, 1273, 1276, 1378, 1482, 1548.

3d. Quantitative analysis

See 938.

3e. Preparative scale chromatography

- 893 Colin, H.: Simple considerations on column design in preparative-scale liquid chromatography. *Sep. Sci. Technol.*, 22 (1987) 1851-1868; *C.A.*, 107 (1987) 178976p.
- 894 Cretier, G. and Rocca, J.L.: Optimization of injection conditions in preparative liquid chromatography. *Sep. Sci. Technol.*, 22 (1987) 1881-1907; *C.A.*, 107 (1987) 178977q.
- 895 Dewaele, C., De Coninck, M. and Verzele, M.: Preparative high performance liquid chromatography and the H/u curve. *Sep. Sci. Technol.*, 22 (1987) 1919-1931; *C.A.*, 107 (1987) 178979s.
- 896 Hupe, K.P. and Hoffmann, B.: Optimum conditions in preparative liquid chromatography. II. Selection of column dimensions. *Sep. Sci. Technol.*, 22 (1987) 1869-1880; *C.A.*, 107 (1987) 201162f.
- 897 Ito, Y.: Cross-axis synchronous flow-through coil planet centrifuge free of rotary seals for preparative countercurrent chromatography. Part I. Apparatus and analysis of acceleration. *Sep. Sci. Technol.*, 22 (1987) 1971-1978; *C.A.*, 107 (1987) 178744m.
- 898 Ito, Y.: Cross-axis synchronous flow-through coil planet centrifuge free of rotary seals for preparative countercurrent chromatography. Part II. Studies on phase distribution and partition efficiency in coaxial coils. *Sep. Sci. Technol.*, 22 (1987) 1989-2009; *C.A.*, 107 (1987) 201163g.
- 899 Kowalczyk, J.S., Gazda, K., Kaminski, M. and Klawiter, J.: Preparative and industrial LC. *GIT-Suppl.*, (1987) 62-67; *C.A.*, 107 (1987) 201158j.
- 900 Meyer, V.R.: The influence of sample mass and flow velocity on performance in preparative column liquid chromatography. *Sep. Sci. Technol.*, 22 (1987) 1909-1917; *C.A.*, 107 (1987) 178978r.
- 901 Newburger, J., Liebes, L., Colin, H. and Guiochon, G.: Investigation of the influence of particle size on the productivity of preparative HPLC columns. *Sep. Sci. Technol.*, 22 (1987) 1933-1951; *C.A.*, 107 (1987) 178980k.
- 902 Reese, W. and Merck, E.: Optimum control for preparative HPLC units. *Swiss Chem.*, 9 (1987) 29-36; *C.A.*, 107 (1987) 178740g.
- 903 Tice, P.A., Mazsaroff, I., Lin, N.T. and Regnier, F.E.: Effects of large sample loads on column lifetime in preparative-scale liquid chromatography. *J. Chromatogr.*, 410 (1987) 43-51.
- 904 Volkov, S.A., Sakodynskii, K.I. and Dustov, S.I.: The productivity of preparative liquid chromatography with linear sorption isotherm. *Symp. Biol. Hung.*, 31 (Chromatography '84) (1986) 185-194; *C.A.*, 107 (1987) 197625q.

See also 1296, 1327.

3f. Programmed temperature, pressure, vapors, gradients

- 905 Rocklin, R.D., Pohl, C.A. and Schibler, J.A.: Gradient elution in ion chromatography. *J. Chromatogr.*, 411 (1987) 107-119.

3g. High performance procedures

- 906 Shetty, P.H., Youngberg, P.J., Kersten, B.R. and Poole, C.F.: Solvent properties of liquid organic salts used as mobile phases in microcolumn reversed-phase liquid chromatography. *J. Chromatogr.*, 411 (1987) 61-79.

See also 905, 1533.

4. SPECIAL TECHNIQUES

4a. Automation and computerization

- 907 Aust, E., Dahlke, K. and Emig, G.: Simulation and evaluation of chromatographic response curves using various methods. *Chem. Eng. J. (Lausanne)*, 35 (1987) 179-190; *C.A.*, 107 (1987) 179087m.
- 908 Hartley, T.F., Philcox, J.C., Beesley, J.F. and Robson, J.B.: Computerized chromatographic peak detection using the Trigg Tracking Signal. An application devised for use with an online analog to digital converter connected between an amino acid analyser and a personal computer. *Biomed. Chromatogr.*, 2 (1987) 104-109.
- 909 Henshall, A. and Schibler, J.A.: Chromatography automation using the Microsoft Windows® operating system. *Int. Lab.*, 17, No. 9 (1987) 28-42.
- 910 Landon, K.: Filling the information gap. *LC-GC*, 5 (1987) 837-838; *C.A.*, 107 (1987) 227937n - a review with no refs.
- 911 Manabe, T.: (Microcomputer-aided gradient elution). *Bunseki*, (1987) 426-429; *C.A.*, 107 (1987) 150459y - a review with 8 refs.
- 912 Schmidt, G.J., Dong, M.W. and Salit, M.: A general robotic procedure for performing precolumn derivatization and its application to on-line liquid chromatographic analysis. *J. Chromatogr. Sci.*, 25 (1987) 453-459.

See also 828, 847, 902, 948, 1184, 1669.

4b. Combination of various chromatographic techniques

See 948, 949, 951, 1049, 1803.

4c. Combination with other physico-chemical techniques (MS, IR etc.)

- 913 De Wit, J.S.M., Parker, C.E., Tomer, K.B. and Jorgenson, J.W.: Direct coupling of open-tubular liquid chromatography with mass spectrometry. *Anal. Chem.*, 59 (1987) 2400-2404.
- 914 Gustavsson, A. and Nygren, O.: Characterization of a nebulizer interface for flame atomic absorption spectroscopy. *Spectrochim. Acta, Part B*, 42B (1987) 883-888; *C.A.*, 107 (1987) 189728k.

For additional information see:
C.A., 107 (1987) 168027r, 168028s.

See also 834, 949, 986, 1101, 1331, 1475, 1772, 1860.

4d. Affinity chromatography

- 915 Groman, E.V. and Wilchek, M.: Recent developments in affinity chromatography supports. *Trends Biotechnol.*, 5 (1987) 220-224; *C.A.*, 107 (1987) 171646x - a review with 33 refs.
- 916 Hubble, J.: Cooperative binding interactions in affinity chromatography: Theoretical considerations. *Biotechnol. Bioeng.*, 30 (1987) 208-215; *C.A.*, 107 (1987) 132580b.

For additional information see:
C.A., 107 (1987) 73734e, 171773m, 171775p.

See also 876, 1293, 1312, 1318, 1334, 1460, 1622.

4g. Separation of enantiomers

- 917 Akanya, J.N. and Taylor, D.R.: Direct resolution of isomeric and diastereoisomeric mixtures of amides, lactones, alcohols, carbazones and pyrethroid derivatives. *J. Liq. Chromatogr.*, 10 (1987) 805-817.
- 918 Armstrong, D.W., Yang, X., Han, S.M. and Menges, R.A.: Direct liquid chromatographic separation of racemates with an α -cyclodextrin bonded phase. *Anal. Chem.*, 59 (1987) 2594-2596.
- 919 Bojarski, J.: Chromatographic resolution of enantiomers and its pharmaceutical applications. *Symp. Biol. Hung.*, 34 (1986) 343-360; *C.A.*, 107 (1987) 161752q - a review with 140 refs.
- 920 Caldwell, J. and Testa, B.: Criteria for the acceptability of experimental evidence for the enantiomeric composition of xenobiotics and their metabolites. *Drug Metab. Disp.*, 15 (1987) 587-588 - a review.
- 921 Gerding, T.K., Drenth, B.F.H. and de Zeeuw, R.A.: Separation of N-0437 enantiomers by RP-HPLC after pre-column derivatization with D(+)-glucuronic acid. *J. High Resolut. Chromatogr. Chromatogr. Commun.*, 10 (1987) 523-525.
- 922 Gyllenhaal, O. and Vessman, J.: Phosgene as a derivatizing reagent prior to gas and liquid chromatography. *J. Chromatogr.*, 435 (1988) 259-269 - a review with 38 refs.
- 923 Iwaki, K., Yoshida, S., Nimura, N., Kinoshita, T., Takeda, K. and Ogura, H.: Activated carbamate reagent as chiral derivatizing agent for liquid chromatographic optical resolution of enantiomeric amino compounds. *Chromatographia*, 23 (1987) 899-902.
- 924 Karnes, H.T. and Sarkar, M.A.: Enantiomeric resolution of drug compounds by liquid chromatography. *Pharm. Res.*, 4 (1987) 285-292; *C.A.*, 107 (1987) 161754s - a review with 72 refs.
- 925 Krstulovic, A.M., Rossey, G., Porziemsky, J.-P., Long, D. and Chekroun, I.: Direct determination of the enantiomeric purity of (5S)-3-isopropyl-5p-toluene-sulphonyloxymethylloxazolidin-2-one on a cellulose-based chiral stationary phase. In-process control of a chiral intermediate used in the synthesis of enantiomerically pure β -blocking agents. *J. Chromatogr.*, 411 (1987) 461-465.
- 926 Lipkowitz, K.B., Demeter, D.A., Parish, C.A., Landwer, J.M. and Darden, T.: Column design. 3. Theoretical studies of a chiral stationary phase used in column chromatography. *J. Comput. Chem.*, 8 (1987) 753-760; *C.A.*, 107 (1987) 157244h.
- 927 Okamoto, Y.: Separate optical isomers by chiral HPLC. *ChemTech.*, 17 (1987) 176-181; *C.A.*, 107 (1987) 197137g - a review with 52 refs.
- 928 Röder, W., Ruffing, F.-J., Schomburg, G. and Pirkle, W.H.: Chiral SFC-separations using polymer-coated open tubular fused silica columns. Comparison of enantiomeric selectivity in SFC and LC using the same stationary phase of the Pirkle type. *J. High Resolut. Chromatogr. Chromatogr. Commun.*, 10 (1987) 665-667.
- 929 Sidorov, A.P., Belousova, T.A., Belov, Yu.P., Chumachenko, M.N. and Martynov, I.V.: (Separation of proline enantiomers on a soluble stationary chiral phase). *Zh. Anal. Khim.*, 42 (1987) 727-729; *C.A.*, 107 (1987) 190046t.
- 930 Spahn, H.: Formation of diastereomeric derivatives of 2-arylpropionic acids using L-leucinamide. *J. Chromatogr.*, 423 (1987) 334-339.
- 931 Wainer, I.W., Stiffin, R.M. and Shibata, T.: Resolution of enantiomeric aromatic alcohols on a cellulose tribenzoate high-performance liquid chromatography chiral stationary phase. A proposed chiral recognition mechanism. *J. Chromatogr.*, 411 (1987) 139-151.

For additional information see:
C.A., 107 (1987) 190081a.

See also 864, 868, 871, 875, 878, 1004, 1037, 1086, 1508, 1623, 1650, 1652, 1698, 1704, 1742.

4h. Other special techniques

- 932 Burton, D.E.: Development and applications of micellar electrokinetic capillary chromatography. Avail. *Univ. Microfilm Int.*, Order No. DA8713452, 1987, 111 pp.; *C.A.*, 107 (1987) 171953v.

- 933 Burton, D.E., Sephaniak, M.J. and Maskarinec, M.P.: Evaluation of the use of various surfactants in micellar electrokinetic capillary chromatography. *J. Chromatogr. Sci.*, 25 (1987) 514-518.
- 934 Chang, C.A. and Wu, Q.: Facile liquid chromatographic separation of positional isomers with a γ -cyclodextrin bonded phase column. *J. Liq. Chromatogr.*, 10 (1987) 1359-1368.
- 935 Janca, J., Pribylova, D. and Jahnova, V.: Separation and dispersion in sedimentation field-flow fractionation. *J. Liq. Chromatogr.*, 10 (1987) 767-782.
- 936 Tanaka, Y., Sato, H. and Kageyu, A.: (Separation of geometric isomers of 3,7,11,15-tetramethyl-2,6,10,14-hexadecatetraen-1-ol or derivatives). *Jpn. Kokai Tokkyo Koho Pat.* JP 62 84,033 (87 84,033) (Cl. C07C33/02), 17 Apr. 1987, Appl. 85/225,660, 09 Oct. 1985, 6 pp.; *C.A.*, 107 (1987) 198701y.

See also 829, 888, 897, 898, 1157, 1549.

5. HYDROCARBONS AND HALOGEN DERIVATIVES

5a. Aliphatic hydrocarbons

See 1540.

5b. Cyclic hydrocarbons

- 937 Bazylak, G. and Maslowska, J.: An evaluation of commercial HPLC silica gel columns for the separation of pericondensed PAHs mixture. *Fresenius' Z. Anal. Chem.*, 328 (1987) 662-663.
- 938 Lafleur, A.L., Monchamp, P.A., Plummer, E.F. and Wornat, M.J.: Universal calibration method for the determination of polycyclic aromatic hydrocarbons by high performance liquid chromatography with broad-band diode-array detection. *Anal. Lett.*, 20 (1987) 1171-1192.
- 939 McKay, S., Farmer, P.B., Cary, P.D. and Grover, P.L.: The metabolism of 7-ethylbenzo[*a*]anthracene by rat liver microsomal preparations. *Drug Metab. Disp.*, 15 (1987) 682-694.
- 940 Melikian, A.A., Bagheri, K. and Hecht, S.S.: Contrasting disposition and metabolism of topically applied benzo[*a*]pyrene, *trans*-7,8-dihydroxy-7,8-dihydrobenzo[*a*]pyrene, and 7 β ,8 α -dihydroxy-9 α ,10 α -epoxy-7,8,9,10-tetrahydrobenzo[*a*]pyrene in mouse epidermis *in vivo*. *Cancer Res.*, 47 (1987) 5354-5360.
- 941 Ostman, C.E. and Colmsjö, A.L.: Backflush HPLC for the isolation of polycyclic aromatic compounds - a comparative study. *Chromatographia*, 23 (1987) 903-908.
- 942 Plakunov, I., Smolarek, T.A., Fischer, D.L., Wiley, J.C., Jr. and Baird, W.M.: Separation by ion-pair HPLC of the glucuronide, sulfate and glutathione conjugates formed from benzo[*a*]pyrene in cell cultures from rodents, fish and humans. *Carcinogenesis*, 8 (1987) 59-66; *C.A.*, 107 (1987) 192711e.
- 943 Sander, L.C. and Wise, S.A.: Effect of phase length on column selectivity for the separation of polycyclic aromatic hydrocarbons by reversed-phase liquid chromatography. *Anal. Chem.*, 59 (1987) 2309-2313.
- 944 Spitzer, T.: Determination of polyphenylarenes and polynuclear aromatic hydrocarbons by reversed phase HPLC. *J. Liq. Chromatogr.*, 10 (1987) 593-601.
- 945 Van Zoonen, P., Kamminga, D.A., Gooijer, C., Velthorst, N.H. and Frei, R.W.: A simplified method for peroxyoxalate chemiluminescence detection of fluorescers based on solid state reagent addition. *J. Liq. Chromatogr.*, 10 (1987) 819-827.

For additional information see:
C.A., 107 (1987) 77007t.

See also 817, 866, 906, 953, 1813, 1832.

5c. Halogen derivatives

- 946 Munder, A., Buchert, H., Niemczyk, R., Swerev, M. and Ballschmiter, K.: Determination of PCDD/PCDF in motor oil and air particulates. A modular approach in clean-up. *Fresenius' Z. Anal. Chem.*, 328 (1987) 639-643.

5d. Complex hydrocarbon mixtures

- 947 Burke, F.P. and Winschel, R.A.: Correlation of liquid chromatographic retention times with molecular structures of coal liquid components. *Anal. Chem.*, 59 (1987) 2586-2593.
- 948 Davies, I.L., Raynor, M.W., Williams, P.T., Andrews, G.E. and Bartle, K.D.: Application of automated on-line microbore high-performance liquid chromatography/capillary gas chromatography to Diesel exhaust particulates. *Anal. Chem.*, 59 (1987) 2579-2583.
- 949 Philip, C.V. and Anthony, R.G.: A study on the distribution of various chemical species in different coal liquids by size exclusion chromatography-gas chromatography-mass spectrometry. *Am. Chem. Soc., Div. Fuel Chem.*, 32 (1987) 368-376; *C.A.*, 107 (1987) 158022w.
- 950 Platonov, V.V., Klyavina, O.A., Kasimtseva, T.V., Ivleva, L.N., Tabolenko, N.V. and Mazurov, V.V.: (Structure of hydrocarbons from Kansk-Achinsk brown coal tar). *Khim. Tverd. Topl. (Moscow)*, (1987) 19-26; *C.A.*, 107 (1987) 157990y.
- 951 Rustamov, M.I., Basili, I.K., Arustamova, L.G. and Sultanov, N.T.: (Hydrocarbon composition of gasoline fractions (initial boiling point 200°) from crude oil of Rasgarib field (Egypt)). *Dokl. Akad. Nauk Az. SSR*, 42 (1986) 50-55; *C.A.*, 107 (1987) 157901v.
- 952 Takeuchi, T. and Ishii, D.: High-performance microcolumn liquid chromatography of hydrocarbons with indirect photometric detection. *J. High Resolut. Chromatogr. Chromatogr. Commun.*, 10 (1987) 571-573.
- 953 Zoccolillo, L., Carboni, G.P., Rizzo, A.P. and Ronchetti, M.: HPLC determination of the aromatic fraction of gasoline. *Chromatographia*, 23 (1987) 825-828.

6. ALCOHOLS

- 954 Goncalves, J.C.S., Selavons, M., Poupaert, J.H. and Dumont, P.: High-performance liquid chromatography of 1,2-diacyl-rac-glycerols and 1,3-diacylglycerols through chromogenic derivatization with 3,5-dinitrobenzoyl. *J. Chromatogr.*, 411 (1987) 472-475.
- 955 Much, H., Czichocki, G. and Vellhardt, D.: (The effect of alkyl chain length on the determination of alcohol traces in sodium alkyl sulphates by HPLC and surface tension measurement). *Abh. Akad. Wiss. DDR, Abt. Math., Naturwiss., Tech.* 1986, 1N (1987) 323-329; *C.A.*, 107 (1987) 178664k.
- 956 Valdez, D. and Reier, J.C.: Derivatization of glycols, hydroxyamines, and polyols at trace levels with 3,5-dinitrobenzoyl chloride utilizing aqueous to nonaqueous phase transfer on a reverse phase cartridge. *J. Liq. Chromatogr.*, 10 (1987) 863-880.

For additional information see:
C.A., 107 (1987) 211388y.

See also 820, 917, 931, 973, 992, 1052, 1161, 1165, 1783.

7. PHENOLS

- 957 Arai, Y., Hirukawa, M. and Hanai, T.: Retention of phenols on an octadecyl-bonded vinylalcohol copolymer gel. *J. Liq. Chromatogr.*, 10 (1987) 635-642.
- 958 Hatano, T., Yoshida, T. and Okuda, T.: Chromatography of tannins. III. Multiple peaks in high-performance liquid chromatography of some hydrolyzable tannins. *J. Chromatogr.*, 435 (1988) 285-295.

See also 1809, 1820.

8. SUBSTANCES CONTAINING HETEROCYCLIC OXYGEN

8a. *Flavonoids*

- 959 Bokkenheuser, V.D., Shackleton, C.H.L. and Winter, J.: Hydrolysis of dietary flavonoid glycosides by strains of intestinal *Bacteroides* from humans. *Biochem. J.*, 248 (1987) 953-956.
- 960 Dondi, F., Kahie, Y.D., Lodi, G., Reschiglian, P., Pietrogrande, C., Bigli, C. and Carboni, G.P.: Comparison of the sequential simplex method and linear solvent strength theory in HPLC gradient elution optimization of multicomponent flavonoid mixtures. *Chromatographia*, 23 (1987) 844-849.
- 961 Sendra, J.M., Navarro, J.L. and Izquierdo, L.: Determination of polymethoxylated flavones in citrus juices by HPLC. *Int. Fruchtsaft-Union, Wiss.-Tech. Komm.*, 19 (1986) 381-387; *C.A.*, 107 (1987) 196574k.
- 962 Takino, Y., Miyahara, T., Arichi, E., Arichi, S., Hayashi, T. and Karikura, M.: Determination of some flavonoids in *Scutellariae* radix by high-performance liquid chromatography. *Chem. Pharm. Bull.*, 35 (1987) 3494-3497.

See also 950, 1524.

8b. *Aflatoxins and other mycotoxins*

- 963 Leitao, J., De Saint Blanquat, G., Bailly, J.R. and Paillas, C.: Quantitation of aflatoxin from various strains of aspergillus in foodstuffs. *J. Chromatogr.*, 435 (1988) 229-234.
- 964 Sano, A., Matsutani, S., Suzuki, M. and Takitani, S.: High-performance liquid chromatographic method for determining trichothecene mycotoxins by post-column fluorescence derivatization. *J. Chromatogr.*, 410 (1987) 427-436.
- 965 Scott, P.M. and Lawrence, G.A.: Liquid chromatographic determination and stability of the *Fusarium* mycotoxin moniliformin in cereal grains. *J. Assoc. Off. Anal. Chem.*, 70 (1987) 850-853.
- 966 Simonella, A., Torreti, L., Filipponi, C., Falgiani, A. and Ambrosii, L.: Simultaneous determination of aflatoxins G₁, B₁, G₂, B₂, in animal feedstuffs by HPTLC and RP-HPLC. *J. High Resolut. Chromatogr. Chromatogr. Commun.*, 10 (1987) 626-628.
- 967 Van Egmond, H.P., Leussink, A.B. and Paulsch, W.E.: The determination of aflatoxin M₁ in milk powder. A comparison of four methods of analysis. *Bull. Int. Dairy Fed.*, 207 (1986) 150-197; *C.A.*, 107 (1987) 132706x.
- 968 Yousef, A.E. and Marth, E.H.: Kinetics of interaction of aflatoxin M₁ in aqueous solutions irradiated with ultraviolet energy. *J. Agric. Food Chem.*, 35 (1987) 785-789.

For additional information see:

C.A., 107 (1987) 132766s, 132767t.

8c. *Other compounds with heterocyclic oxygen (including tannins)*

- 969 Nonaka, G.-I., Ishimatsu, M., Tanaka, T., Nishioka, I., Nishizawa, M. and Yamagishi, T.: Tannins and related compounds. LVIII. Novel gallotannins possessing an α -glucose core from *Nuphar japonicum* DC. *Chem. Pharm. Bull.*, 35 (1987) 3127-3131.
- 970 Shen, E.S. and Olson, J.R.: Relationship between the murine phenotype and the hepatic uptake and metabolism of 2,3,7,8-tetrachlorodibenzo-*p*-dioxin. *Drug Metab. Disp.*, 15 (1987) 653-660.
- 971 Spencer, G.F., Tjarks, L.W. and Powell, R.G.: Analysis of linear and angular furanocoumarins by dual-column high-performance liquid chromatography. *J. Agric. Food Chem.*, 35 (1987) 803-805.
- 972 Stolk, L.M.L., de Ruiter, R., Saadawi, A., Siddiqui, A.H. and Cormane, R.H.: Determination of psoralen in serum by reversed-phase high-performance liquid chromatography. *J. Chromatogr.*, 423 (1987) 283-286.

For additional information see:

C.A., 107 (1987) 213467x.

See also 789, 917, 941, 1543, 1726.

9. OXO COMPOUNDS, ETHERS, EPOXIDES AND QUINONES

- 973 Albarran, G. and Collins, C.H.: Application of ion moderated-partition chromatography to the separation of a mixture of simple non-ionized compounds. *J. High Resolut. Chromatogr. Chromatogr. Commun.*, 10 (1987) 581-583.
- 974 Amoss, C.W., Slack, R.W. and Taylor, L.R.: On-stream HPLC determination of residual ethylene oxide and propylene oxide in glycerol based polyol synthesis. *J. Liq. Chromatogr.*, 10 (1987) 583-592.
- 975 Buffinton, G.D., Hunt, N.H., Cowden, W.B. and Clark, I.A.: Detection of short-chain carbonyl products of lipid peroxidation from malaria-parasite (*Plasmodium vivax*)-infected red blood cells exposed to oxidative stress. *Biochem. J.*, 249 (1988) 63-68.
- 976 Edlund, P.O.: On-column catalysis of hydratization equilibria during liquid chromatographic analysis of acetaldehyde and formaldehyde. *Chromatographia*, 23 (1987) 709-712.
- 977 Polo, C. and Almy, J.: HPLC analysis of aldehydes and ketones in sherry. *J. Liq. Chromatogr.*, 10 (1987) 1165-1174.
- 978 Possanzini, M., Ciccio, P., Di Palo, V. and Draisci, R.: Determination of low boiling aldehydes in air and exhaust gases by using annular denuders combined with HPLC techniques. *Chromatographia*, 23 (1987) 829-834.
- 979 Reed, J.S. and Ragan, C.I.: The effect of rate limitation by cytochrome c on the redox state of the ubiquinone pool in reconstituted NADH:cytochrome c reductase. *Biochem. J.*, 247 (1987) 657-662.
- 980 Walczak, B., Chretien, J.R., Dreux, M., Morin-Allory, L. and Lafosse, M.: Factor analysis and experiment design in high-performance liquid chromatography. IV. Influence of mobile phase modifications on the selectivity of chalcones on an ODS stationary phase. *Chemom. Intell. Lab. Syst.*, 1 (1987) 177-189; *C.A.*, 107 (1987) 69988y.
- 981 Wang, M.: Analysis of gossypol by HPLC. *J. Ethnopharmacol.*, 20 (1987) 1-11; *C.A.*, 107 (1987) 147486z.

For additional information see:

C.A., 107 (1987) 73593h.

See also 818, 820, 866, 1060, 1813.

10. CARBOHYDRATES

10a. Mono and oligosaccharides. Structural studies

- 982 Abbou, M. and Siouffi, A.-M.: Liquid chromatography of sugars on diol columns. Detection with cuprammonium reagent. *J. Liq. Chromatogr.*, 10 (1987) 95-106.
- 983 Azzi, A. and Niola, I.: (Characterization of a commercial isoglucose). *Ind. Aliment.*, 26 (1987) 550-552; *C.A.*, 107 (1987) 196607y.
- 984 Brooks, J.R. and Griffin, V.K.: Saccharide analysis of corn syrup solids and maltodextrins using HPLC. *Cereal Chem.*, 64 (1987) 253-255; *C.A.*, 107 (1987) 174499n.
- 985 Daniel, P.F.: Separation of benzoylated oligosaccharides by reversed-phase high-pressure liquid chromatography: application to high-mannose type oligosaccharides. *Methods Enzymol.*, 138 (Complex Carbohydr., Pt. E) (1987) 94-116; *C.A.*, 107 (1987) 150511j.
- 986 Galensa, R., Landt, K. and Herres, W.: Application of the flow-cell HPLC-FTIR technique to the analysis of benzoylated saccharides and compounds with similar polarity. *Z. Lebensm.-Unters. Forsch.*, 185 (1987) 36-38; *C.A.*, 107 (1987) 171737c.
- 987 Gross, H.J., Bünsch, A., Paulson, J.C. and Brossmer, R.: Activation and transfer of novel synthetic 9-substituted sialic acids. *Eur. J. Biochem.*, 168 (1987) 595-602.
- 988 Koski, P.G.: Determination of reducing as well as non-reducing carbohydrates using LCEC. *Curr. Sep.*, 8 (1987) 26-28; *C.A.*, 107 (1987) 171752d.

- 989 Machado de Moraes, R.: Chromatography for liquids. *Braz. Pedido Pat.* PI BR 85 04,180 (Cl. GO1N30/04), 07 Apr. 1987, Appl. 85/4,180, 29 Aug. 1985; 19 pp.; C.A., 107 (1987) 168059c.
- 990 Malmberg, A. and Theander, O.: Analysis of sucrose content in beet crops. *Swed. J. Agric. Res.*, 17 (1987) 3-7; C.A., 107 (1987) 216250g.
- 991 McDowell, W., Grier, T.J., Rasmussen, J.R. and Schwarz, R.T.: The role of C-4-substituted mannose analogues in protein glycosylation. Effect of the guanosine diphosphate esters of 4-deoxy-4-fluoro-D-mannose and 4-deoxy-D-mannose on lipid-linked oligosaccharide assembly. *Biochem. J.*, 248 (1987) 523-531.
- 992 McKay, D.B., Tanner, G.P., Maclean, D.J. and Scott, K.J.: Detection of polyols and sugars by cuprammonium ion in the presence of strong base. *Anal. Biochem.*, 165 (1987) 392-398.
- 993 Mentasti, E., Gennaro, M.C., Sarzanini, C. and Porta, V.: Derivatization, separation and determination of oligosaccharides by reversed-phase HPLC and fluorometric detection. *Ann. Chim. (Rome)*, 77 (1987) 579-590; C.A., 107 (1987) 168040q.
- 994 Olechno, J.D., Carter, S.R., Edwards, W.T. and Gillen, D.G.: Developments in the chromatographic determination of carbohydrates. *Am. Biotechnol. Lab.*, 5 (1987) 38-50; C.A., 107 (1987) 214312e.
- 995 Uchiyama, H., Kikuchi, K., Ogamo, A. and Nagasawa, K.: Determination of the distribution of constituent disaccharide units within the chain near the linkage region of shark-cartilage chondroitin sulfate C. *Biochim. Biophys. Acta*, 926 (1987) 239-248.
- 996 Watanabe, N. and Kawasaki, S.: Determination of galactose in human plasma by HPLC with electrochemical detection. *Biomed. Chromatogr.*, 2 (1987) 95-98.
- 997 White, J.W., Jr.: Wiley led the way: a century of federal honey research. *J. Assoc. Off. Anal. Chem.*, 70 (1987) 181-189 - a review with 52 refs.

For additional information see:
C.A., 107 (1987) 95620s, 171772k.

See also 803, 1005, 1006, 1013, 1022, 1154, 1566.

10b. *Polysaccharides, mucopolysaccharides, lipopolysaccharides*

- 998 Altman, E., Brisson, J.-R. and Perry, M.B.: Structure of the capsular polysaccharide of *Haemophilus pleuropneumoniae* serotype 5. *Eur. J. Biochem.*, 170 (1987) 185-192.
- 999 Chlenov, M.A., Yarovaya, S.M., Alekseeva, G.S., Kudrvashov, L.I., Titova, E.V., Grineva, L.P. and Reshetov, A.S.: (Determination of molecular weights of microbial polysaccharides by high-performance size-exclusion chromatography). *Príkl. Biokhím. Mikrobiol.*, 23 (1987) 515-521; C.A., 107 (1987) 150537x.
- 1000 Gherezghiher, T., Koss, M.C., Nordquist, R.E. and Wilkinson, C.P.: Analysis of vitreous and aqueous levels of hyaluronic acid: application of HPLC. *Exp. Eye Res.*, 45 (1987) 347-349; C.A., 107 (1987) 150546z.
- 1001 Hourdet, D. and Muller, G.: Solution properties of pectin polysaccharides. I. Aqueous size exclusion chromatography of flax pectins. *Carbohydr. Polym.*, 7 (1987) 301-312; C.A., 107 (1987) 156842q.
- 1002 Jarzebinski, J. and Gryz, K.: (Heterogeneity of heparin. III. Analytical characteristics of the components of Polish heparin separated by gel column chromatography). *Acta Pol. Pharm.*, 43 (1986) 612-616; C.A., 107 (1987) 161473z.
- 1003 Motohashi, N., Nakamichi, Y., Mori, I., Nishikawa, H. and Umemoto, J.: Analysis by high-performance gel permeation chromatography of hyaluronic acid in animal skins and rabbit synovial fluid. *J. Chromatogr.*, 435 (1988) 335-342.
- 1004 Murata, K. and Yokoyama, Y.: Characterization of the products generated from oversulphated dermatan sulphate isomers with chondroitinase-B by high-performance liquid chromatography. *J. Chromatogr.*, 423 (1987) 51-61.
- 1005 Oeben, M., Keller, R., Stuhlsatz, H.W. and Greiling, H.: Constant and variable domains of different disaccharide structure in corneal keratan sulphate chains. *Biochem. J.*, 248 (1987) 85-93.
- 1006 Pejler, G. and David, G.: Basement-membrane heparan sulphate with high affinity for antithrombin synthesized by normal and transformed mouse mammary epithelial cells. *Biochem. J.*, 248 (1987) 69-77.

- 1007 Steiner, M.: Platelet surface glycosaminoglycans are an effective shield for distinct platelet receptors. *Biochim. Biophys. Acta*, 931 (1987) 286-293.
- 1008 Tang, J., Julian, J., Glasser, S.R. and Carson, D.D.: Heparan sulfate proteoglycan synthesis and metabolism by mouse uterine epithelial cells cultured *in vitro*. *J. Biol. Chem.*, 262 (1987) 12832-12842.
- 1009 Yu, L.P. and Rollings, J.E.: Low-angle laser light scattering-aqueous size exclusion chromatography of polysaccharides: molecular weight distribution and polymer branching determination. *J. Appl. Polym. Sci.*, 33 (1987) 1909-1921; *C.A.*, 107 (1987) 78174g.

For additional information see:

C.A., 107 (1987) 156853u, 215826n.

See also 1828.

10c. Glycoproteins and their components

- 1010 Campbell, M.A. and Handley, C.J.: The effect of retinoic acid on proteoglycan turnover in bovine articular cartilage cultures. *Arch. Biochem. Biophys.*, 258 (1987) 143-155.
- 1011 Castillo, C.J., Colburn, P. and Buonassisi, V.: Characterization and N-terminal sequence of heparan sulphate proteoglycan synthesized by endothelial cells in culture. *Biochem. J.*, 247 (1987) 687-693.
- 1012 De Boeck, H., Lories, V., David, G., Cassiman, J.-J. and van den Berghe, H.: Identification of a 64 kDa heparan sulphate proteoglycan core protein from human lung fibroblast plasma membranes with a monoclonal antibody. *Biochem. J.*, 247 (1987) 765-771.
- 1013 Hughes, R.C., Foddy, L. and Bause, E.: Asparagine-linked oligosaccharides of BHK cells treated with inhibitors of oligosaccharide processing. *Biochem. J.*, 247 (1987) 537-545.
- 1014 Mallet, B., Franc, J.L., Miquel, M. and Arnaud, C.: Effects of severe burns on glycan microheterogeneity of four acute phase proteins. *Clin. Chim. Acta*, 167 (1987) 247-257.
- 1015 Mazars, C., Rossignol, M. and Auriol, P.: (HPLC purification of phytotoxic glycoproteins produced by *Rhynchosporium secalis* (Oud.) Davis). *Bio-Sciences*, 6 (1987) 63-68; *C.A.*, 107 (1987) 214306f.
- 1016 Morris, J.E., Yanagishita, M. and Hascall, V.C.: Proteoglycans synthesized by embryonic chicken retina in culture: composition and compartmentalization. *Arch. Biochem. Biophys.*, 258 (1987) 206-218.
- 1017 Pohl, G., Kenne, L., Nilsson, B. and Einarsson, M.: Isolation and characterization of three different carbohydrate chains from melanoma tissue plasminogen activator. *Eur. J. Biochem.*, 170 (1987) 69-75.
- 1018 Quinn, J.M. and Etzler, M.E.: Isolation and characterization of a lectin from the roots of *Dolichos biflorus*. *Arch. Biochem. Biophys.*, 258 (1987) 535-544.
- 1019 Saxena, S., Shailubhai, K., Dong-Yu, B. and Vijay, I.K.: Purification and characterization of glucosidase II involved in N-linked glycoprotein processing in bovine mammary gland. *Biochem. J.*, 247 (1987) 563-570.
- 1020 Schaller, J., Akiyama, K., Tsuda, R., Hara, M., Marti, T. and Rickli, E.E.: Isolation, characterization and amino-acid sequence of γ -seminoprotein, a glycoprotein from human seminal plasma. *Eur. J. Biochem.*, 170 (1987) 111-120.
- 1021 Schmelzer, C.H. and Poduslo, S.E.: Glycoproteins in the whorls of membrane produced by oligodendroglia in culture. *Biochim. Biophys. Acta*, 903 (1987) 103-111.
- 1022 Scudder, P., Lawson, A.M., Hounsell, E.F., Carruthers, R.A., Childs, R.A. and Feizi, T.: Characterisation of oligosaccharides released from human-blood-group O erythrocyte glycopeptides by the endo- β -galactosidase of *Bacteroides fragilis*. A study of the enzyme susceptibility of branched poly(N-acetyl-lactosamine) structures. *Eur. J. Biochem.*, 168 (1987) 585-593.
- 1023 Shailubhai, K., Pratta, M.A. and Vijay, I.K.: Purification and characterization of glucosidase I involved in N-linked glycoprotein processing in bovine mammary gland. *Biochem. J.*, 247 (1987) 555-562.
- 1024 Shimoda, N., Muramatsu, H., Osawa, T., Yamada, T., Adachi, M. and Muramatsu, T.: Carbohydrate moieties of peanut agglutinin receptors isolated from human gastric cancer cells. *J. Biochem. (Tokyo)*, 102 (1987) 657-664.

- 1025 Smith, B.F.: Human gallbladder mucin binds biliary lipids and promotes cholesterol crystal nucleation in model bile. *J. Lipid Res.*, 28 (1987) 1088-1097.
- 1026 Sturn, A., van Kuik, J.A., Vliegthart, J.F.G. and Chrispeels, M.J.: Structure, position, and biosynthesis of the high mannose and the complex oligosaccharide side chains of the bean storage protein phaseolin. *J. Biol. Chem.*, 262 (1987) 13392-13403.
- 1027 Toyoshima, S. and Tsuji, T.: (Separation of lectins). *Farumashia Rebyu*, 21 (Seibutsu Kassei Busshitsu no Tanri) (1986) 75-90; *C.A.*, 107 (1987) 150455u - a review with 88 refs.
- 1028 Van Kuik, J.A., Sijbesma, R.P., Kamerling, J.P., Vliegthart, J.F.G. and Wood, E.J.: Primary structure of determination of seven novel N-linked carbohydrate chains derived from hemocyanin of *Lymnaea stagnalis*. 3-O-Methyl-D-galactose and N-acetyl-D-galactosamine as constituents of xylose-containing N-linked oligosaccharides in an animal glycoprotein. *Eur. J. Biochem.*, 169 (1987) 399-411.
- 1029 Webber, C., Glant, T.T., Roughley, P.J. and Poole, A.R.: The identification and characterization of two populations of aggregating proteoglycans of high buoyant density isolated from post-natal human articular cartilages of different ages. *Biochem. J.*, 248 (1987) 735-740.

See also 1223, 1298, 1347, 1432.

11. ORGANIC ACIDS AND LIPIDS

11a. Organic acids and simple esters

- 1030 Ansari, M.H. and Ahmad, M.: Reaction of methyl trans-2,3-epoxy-octadecanoate with phenyl isocyanate: synthesis of fatty 3-phenyl-2-oxazolidones. *J. Am. Oil Chem. Soc.*, 64 (1987) 1544-1547.
- 1031 Bedetti, C. and Canraffa, A.: Extraction and purification of arachidonic acid metabolites from cell cultures. *Adv. Biochem. Eng./Biotechnol.*, 35 (1987) 47-81; *C.A.*, 107 (1987) 147443h - a review with 177 refs.
- 1032 Boley, N.P. and Colwell, R.K.: Comparison of methods for the determination of butyric acid in foodstuffs by gas-liquid chromatography and high-performance liquid chromatography. *J. Chromatogr.*, 410 (1987) 190-194.
- 1033 Brash, A.R., Baertschi, S.W., Ingram, C.D. and Harris, T.M.: On non-cyclooxygenase prostaglandin synthesis in the sea whip coral, *Plexaura homomalla*: an 8(R)-lipoxygenase pathway leads to formation of an α -ketol and a racemic prostanoid. *J. Biol. Chem.*, 262 (1987) 15829-15839.
- 1034 Chia, J.S. and Badriah, Z.: Determination of benzoic acid in chili sauce by HPLC. *MARDI Res. Bull.*, 13 (1985) 190-193; *C.A.*, 107 (1987) 174480z.
- 1035 Coleman, R.A. and Haynes, E.B.: Synthesis and release of fatty acids by human trophoblast cells in culture. *J. Lipid Res.*, 28 (1987) 1335-1341.
- 1036 Connelly, J.A. and Siehl, D.L.: Purification of chorismate, prephenate, and agrogenate by HPLC. *Methods Enzymol.*, 142 (Metab. Aromat. Amino Acids Amines) (1987) 422-431; *C.A.*, 107 (1987) 111996b.
- 1037 Dernoncour, R. and Azerad, R.: High-performance liquid chromatographic separation of the enantiomers of substituted 2-aryloxypropionic acid methyl esters. *J. Chromatogr.*, 410 (1987) 355-361.
- 1038 Downey, B.P. and Jenke, D.R.: Determination of oxalate in pharmaceutical matrices by indirect photometric chromatography. *J. Chromatogr. Sci.*, 25 (1987) 519-524.
- 1039 Ertel, K.D. and Carstensen, J.T.: Quantitative determination of octanoic acid by high-performance liquid chromatography following derivatization with 4-bromomethyl-7-methoxycoumarin. *J. Chromatogr.*, 411 (1987) 297-304.
- 1040 Funk, M.O., Jr., Andre, J.C. and Otsuki, T.: Oxygenation of trans polyunsaturated fatty acids by lipoxygenase reveals steric features of the catalytic mechanism. *Biochemistry*, 26 (1987) 6880-6884.
- 1041 Grandgirard, A., Julliar, F., Prevost, J. and Sebedio, J.L.: Preparation of geometrical isomers of linolenic acid. *J. Am. Oil Chem. Soc.*, 64 (1987) 1434-1440.

- 1042 Haller, T. and Lackner, R.: Screening for organic acids in fish tissues with special reference to the distribution of taurine in *Rutilus rutilus* L. *Fish Physiol. Biochem.*, 3 (1987) 145-149; *C.A.*, 107 (1987) 130181y.
- 1043 Hecker, M., Ullrich, V., Fischer, C. and Meese, C.O.: Identification of novel arachidonic acid metabolites formed by prostaglandin H synthase. *Eur. J. Biochem.*, 169 (1987) 113-123.
- 1044 Isseroff, R.R., Ziboh, V.A., Chapkin, R.S. and Martinez, D.T.: Conversion of linoleic acid into arachidonic acid by cultured murine and human keratinocytes. *J. Lipid Res.*, 28 (1987) 1342-1349.
- 1045 Kihara, K., Rokushika, S. and Hatano, H.: Elution behaviour of aliphatic carboxylic acids on a strong cation-exchange resin column. *J. Chromatogr.*, 410 (1987) 103-110.
- 1046 Lydon, J., Duke, S.O. and Hedin, P.A.: Coextraction of 5-(hydroxymethyl)-2-furaldehyde with phenolic acids in acid-hydrolyzed plant extracts. *J. Chromatogr.*, 410 (1987) 474-479.
- 1047 Maeda, Y., Yamamoto, M., Owada, K., Sato, S., Masui, T., Nakazawa, H. and Fujita, M.: High-performance liquid chromatographic determination of six *p*-hydroxybenzoic acid esters in cosmetics using Sep-Pak Florisil cartridges for sample pre-treatment. *J. Chromatogr.*, 410 (1987) 413-418.
- 1048 Mathur, S.N. and Field, F.J.: Effect of cholesterol enrichment on 12-hydroxy-eicosatetraenoic acid metabolism by mouse peritoneal macrophages. *J. Lipid Res.*, 28 (1987) 1166-1176.
- 1049 Mutter, M. and Homan, H.R.: Estimation of amount of *trans,trans*-isomers of octadecadienoic acids in hydrogenated oils and fats. *J. High Resolut. Chromatogr. Chromatogr. Commun.*, 10 (1987) 672-673.
- 1050 Niyazov, A.N., Niyazberdiyeva, E.F. and Vakhobova, Kh.D.: (Ion-exchange chromatography of individual carboxylic acids and their mixtures). *Izv. Akad. Nauk Turk. SSR, Ser. Fiz.-Tekh., Khim. Geol. Nauk*, (1987) 59-66; *C.A.*, 107 (1987) 190044r.
- 1051 Ontiveros, R.R., Shermer, W.D. and Berner, R.A.: An HPLC method for the determination of 2-hydroxy-4-(methylthio)butanoic acid (HMB) in supplemented animal feeds. *J. Agric. Food Chem.*, 35 (1987) 692-694.
- 1052 Oning, L.: ω -Oxidation of cysteine-containing leukotrienes by rat-liver microsomes. Isolation and characterization of ω -hydroxy and ω -carboxy metabolites of leukotriene E₄ and N-acetylleukotriene E₄. *Eur. J. Biochem.*, 170 (1987) 77-85.
- 1053 Philip, T. and Chen, T.-S.: Separation and quantitative analysis of some carotenoid fatty acid esters of fruits by liquid chromatography. *J. Chromatogr.*, 435 (1988) 113-126.
- 1054 Sjöström, J. and Holmbom, B.: Size-exclusion chromatography of deposits in pulp and paper mills. *J. Chromatogr.*, 411 (1987) 363-370.
- 1055 Soberman, R.J., Okita, R.T., Fitzsimmons, B., Rokach, J., Spur, B. and Austen, K.F.: Stereochemical requirements for substrate specificity of LTB₄ 20-hydroxylase. *J. Biol. Chem.*, 262 (1987) 12421-12427.
- 1056 Uji, Y., Noma, A., Shiraki, M., Maeda, M., Tsuji, A. and Okabe, H.: Separation and quantitation of plasma free fatty acids as phenacyl esters by HPLC. *Biomed. Chromatogr.*, 2 (1987) 110-114.

For additional information see:

C.A., 107 (1987) 76220v, 171761f, 174501g, 174502h, 174539a.

See also 820, 866, 912, 917, 973, 1059, 1189.

11b. Prostaglandins

- 1057 Ball, H.A. and Keppler, D.: ω -Oxidation products of leukotriene E₄ in bile and urine of the monkey. *Biochem. Biophys. Res. Commun.*, 148 (1987) 664-670.
- 1058 Dobrowsky, R.T., O'Sullivan, G., Ballas, L.M., Fleisher, L.N. and Olson, C.N.: Formation of isoindole derivatives of sulfidopeptide leukotrienes by reaction with *o*-phthalaldehyde and separation by reverse-phase high performance liquid chromatography. *J. Liq. Chromatogr.*, 10 (1987) 137-160.
- 1059 Hatzelmann, A. and Ullrich, V.: Regulation of 5-lipoxygenase activity by the glutathione status in human polymorphonuclear leukocytes. *Eur. J. Biochem.*, 169 (1987) 175-184.

- 1060 Okamoto, Y., Aburatani, R., Kawashima, M., Hataja, K. and Okamura, N.: Chromatographic resolution. XIII. Resolution of 4-hydroxy-2-cyclopentenone derivatives by HPLC on cellulose tris(phenylcarbamate) derivatives. *Chem. Lett.*, (1986) 1767-1770; *C.A.*, 107 (1987) 197561r.
- 1061 Oliw, E.H., Benthin, G. and Hamberg, M.: Isolation of 19,20-dehydroprostaglandins E₁ and E₂ in human seminal fluid and further studies on 18,19-dehydroprostaglandins E₁ and E₂. *Arch. Biochem. Biophys.*, 258 (1987) 272-286.
- 1062 Palm, C.D. and Levine, B.A.: Simultaneous preparation and extraction of prostaglandins from gastric tissue with separation and quantification by high pressure liquid chromatography. *J. Liq. Chromatogr.*, 10 (1987) 881-897.
- 1063 Shak, S.: Leukotriene B₄ catabolism: quantitation of leukotriene B₄ and its ω -oxidation products by reversed-phase HPLC. *Methods Enzymol.*, 141 (Cell. Regul., Pt. B) (1987) 355-371; *C.A.*, 107 (1987) 132103y.
- 1064 Umeda, M., Amagaya, S. and Ogihara, Y.: Separation of thromboxane B₂ by high-performance liquid chromatography. *J. Chromatogr.*, 422 (1987) 246-252.
- 1065 Uotila, P., Vargas, R., Wroblewska, B., d'Alarcao, M., Matsuda, S.P., Corey, E.J., Cunard, C.M. and Ramwell, P.W.: Relaxing effects of 15-lipoxygenase products of arachidonic acid on rat aorta. *J. Pharmacol. Exp. Ther.*, 242 (1987) 945-949.
- 1066 Verhagen, J., Bel, E.H., Kijne, G.M., Sterk, P.J., Bruynzeel, P.L.B., Veldink, G.A. and Vliegthart, J.F.G.: The excretion of leukotriene E₄ into urine following inhalation of leukotriene D₄ by human individuals. *Biochem. Biophys. Res. Commun.*, 148 (1987) 864-868.
- 1067 Voyksner, R.D., Bush, E.D. and Brent, D.: Derivatization to improve thermospray HPLC/MS sensitivity for the determination of prostaglandins and thromboxane B₂. *Biomed. Environ. Mass Spectrom.*, 14 (1987) 523-531; *C.A.*, 107 (1987) 191111x.

See also 1052.

11c. Lipids and their constituents

- 1068 Ando, S., Waki, H., Kon, K. and Kishimoto, Y.: Up-to-date chromatography of gangliosides. *NATO ASI Ser., Ser. H*, 7 (Gangliosides Modulation Neuronal Funct.) (1987) 167-177; *C.A.*, 107 (1987) 171632q - a review with 33 refs.
- 1069 Bishop, D.G.: Analysis of molecular species of chloroplast lipids by HPLC. *J. Liq. Chromatogr.*, 10 (1987) 1497-1505.
- 1070 Dixon, J.F. and Hokin, L.E.: Inositol 1,4,5-trisphosphate in pancreatic minilobules on stimulation with carbamylcholine in the absence of lithium. Possible role as a second messenger in long- but not short-term responses. *J. Biol. Chem.*, 262 (1987) 13892-13895.
- 1071 Foglia, T.A., Vail, P.D. and Iwama, T.: High performance liquid chromatographic analysis of 1-alkyl-2-acyl- and 1-alkyl-3-acyl-sn-glycerols. *Lipids*, 22 (1987) 362-365; *C.A.*, 107 (1987) 200831m.
- 1072 Hachey, D.L., Thomas, M.R., Emien, E.A., Garza, C., Brown-Booth, L., Adlof, R.O. and Klein, P.D.: Human lactation: maternal transfer of dietary triglycerides labeled with stable isotopes. *J. Lipid Res.*, 28 (1987) 1185-1192.
- 1073 Kaneko, T., Ohta, Y. and Machida, Y.: Sensitive analysis of aminophospholipids with precolumn fluorescent derivatization by HPLC. *Agric. Biol. Chem.*, 51 (1987) 2023-2024; *C.A.*, 107 (1987) 130198j.
- 1074 Kannagi, R., Watanabe, K. and Hakomori, S.: Isolation and purification of glycosphingolipids by HPLC. *Methods Enzymol.*, 138 (Complex Carbohydr., Pt. E) (1987) 3-13; *C.A.*, 107 (1987) 150510h.
- 1075 Lawrence, J.F., Chadha, R.K. and Conacher, B.S.: Characterization of brominated vegetable oils by normal and reversed phase liquid chromatography. *J. Liq. Chromatogr.*, 10 (1987) 205-214.
- 1076 Nakamura, M. and Sweeley, C.C.: Glycosphingolipid glycosyltransferase assay using Sephadex G-50 chromatography in aqueous phase. *Anal. Biochem.*, 166 (1987) 230-234.
- 1077 Nakayama, R., Yasuda, K. and Saito, K.: Existence of endogenous inhibitors of platelet activating factor (PAF) with PAF in rat uterus. *J. Biol. Chem.*, 262 (1987) 13174-13179.
- 1078 Ng, H.T., Hsu, L.M., Leung, W.Y. and Chen, C.Y.: A comparison of HPLC and TLC in the assessment of amniotic fluid lecithin/sphingomyelin ratio. *Proc. Natl. Sci. Council, Repub. China, Part B: Life Sci.*, 11 (1987) 91-96; *C.A.*, 107 (1987) 171743b.

- 1079 Nurmela, K.V.V. and Satama, L.T.: Quantitative analysis of triglycerides by high-performance liquid chromatography using non-linear gradient elution and flame ionization detection. *J. Chromatogr.*, 435 (1988) 139-148.
- 1080 Rahim, M.A.A., Tan, B.K., Mamat and Embong, S.: (Triglyceride analysis of cocoa butter and cocoa butter equivalent and components using reverse HPLC). *Proc. Malays. Biochem. Soc. Conf.*, 11 (1985) 34-40; *C.A.*, 107 (1987) 152967m.
- 1081 Reddy, S.Y. and Prabhakar, J.V.: Effect of diglycerides on the solidification properties of sal (*Shorea robusta*) fat. *Fett Wissen Technol.-Fat Sci. Technol.*, 89 (1987) 394-397.
- 1082 Riviere, M., Fournie, J.-J. and Puzo, G.: A novel mannose containing phenolic glycolipid from *Mycobacterium kansasii*. *J. Biol. Chem.*, 262 (1987) 14879-14884.
- 1083 Sallustio, B.C., Meffin, P.J. and Thompson, M.: High-performance liquid chromatographic quantitation of triacylglycerols containing fenoprofen from biological samples. *J. Chromatogr.*, 422 (1987) 33-41.
- 1084 Sharp, P., Poulos, A., Fellenberg, A. and Johnson, D.: Structure and lipid distribution of polyenoic very-long-chain fatty acids in the brain of peroxisome-deficient patients (Zellweger syndrome). *Biochem. J.*, 248 (1987) 61-67.
- 1085 Stolyho, A., Martin, M. and Guiochon, G.: Analysis of lipid classes by HPLC with the evaporative light scattering detector. *J. Liq. Chromatogr.*, 10 (1987) 1237-1253.
- 1086 Takagi, T. and Itabashi, Y.: Rapid separations of diacyl- and dialkylglycerol enantiomers by HPLC on a chiral stationary phase. *Lipids*, 22 (1987) 596-600; *C.A.*, 107 (1987) 194372u.

For additional information see:

C.A., 107 (1987) 147650y, 150514n, 152982n.

See also 1053.

11d. Lipoproteins and their constituents

- 1087 Fadeeva, I.V., Staroverov, S.M., Lisichkin, G.V., Gaida, A.V., Magerovskii, Yu.V. and Monastyrskii, V.A.: (Gel chromatography of high-molecular-weight complexes of thromboplastin). *Vysokomol. Soedin., Ser. A*, 29 (1987) 1665-1668; *C.A.*, 107 (1987) 150545y.
- 1088 Huff, M.W. and Strong, W.L.P.: Separation and isolation of human apolipoproteins C-II, C-III₀, C-III₁, and C-III₂ by chromatofocusing on the Fast Protein Liquid Chromatography system. *J. Lipid Res.*, 28 (1987) 1118-1123.
- 1089 Manabe, M., Abe, T., Nozawa, M., Maki, A., Hirata, M. and Itakura, H.: New substrate for determination of serum lecithin: cholesterol acyltransferase. *J. Lipid Res.*, 28 (1987) 1206-1215.
- 1090 Rinninger, F. and Pittman, R.C.: Regulation of the selective uptake of high density lipoprotein - associated cholesteryl esters. *J. Lipid Res.*, 28 (1987) 1313-1325.
- 1091 Wilson, D.E., Hejazi, J., Elstad, N.L., Chan, I.-F., Gleeson, J.M. and Iverius, P.-H.: Novel aspects of vitamin A metabolism in the dog: distribution of lipoprotein retinyl esters in vitamin A-deprived and cholesterol-fed animals. *Biochim. Biophys. Acta*, 922 (1987) 247-258.

For additional information see:

C.A., 107 (1987) 150520m.

13. STEROIDS

- 1092 Bastart-Malsot, M., Parvez, H., Parvez, S., Carpentier, G. and Nagatsu, T.: Separation of steroids by HPLC. Tentative applications of electrochemical detection for high sensitivity yield. *Prog. HPLC*, 2 (Electrochem. Detect. Med. Chem.) (1987) 261-290; *C.A.*, 107 (1987) 168824y.

13a. Pregnane and androstane derivatives

- 1093 Belov, A.B., Arzamastsev, A.P., Sadchikova, N.P., Umyashov, A.N. and Sibilev, A.V.: (A microcolumn high-performance liquid chromatography for the identification and quality evaluation of cortisone derivatives). *Farmatsiya (Moscow)*, 36 (1987) 32-34; *C.A.*, 107 (1987) 223374d.
- 1094 Cornarakis-Lentzos, M. and Cowin, P.R.: Dilutions of corticosteroid creams and ointments - a stability study. *J. Pharm. Biomed. Anal.*, 5 (1987) 707-716.
- 1095 Eriksson, C.-G. and Eneroth, P.: Studies on rat liver microsomal steroid metabolism using ^{18}O -labelled testosterone and progesterone. *J. Steroid Biochem.*, 28 (1987) 549-557.
- 1096 Formento, J.L., Moll, J.L., Francoual, M., Krebs, B.P., Milano, G., Rencee, N., Khater, R., Frenay, M. and Namer, M.: HPLC micromethod for simultaneous measurement of estradiol, progesterone, androgen and glucocorticoid receptor levels. Application to breast cancer biopsies. *Eur. J. Cancer Clin. Oncol.*, 23 (1987) 1307-1314; *C.A.*, 107 (1987) 214316j.
- 1097 Kobayashi, Y., Nishiguchi, Y., Kiguchi, T., Ninomiya, I. and Watanabe, F.: An alternative metabolic pathway of 11-deoxycorticosterone in bovine adrenal *in vitro*: evidence for the presence of a pathway of 11-deoxycorticosterone oxidation at 19-position. *J. Steroid Biochem.*, 28 (1987) 759-767.
- 1098 Lacroix, C., Fiet, J., Benais, J.-P., Gueux, B., Bonete, R., Villette, J.-M., Gourmel, B. and Dreux, C.: Simultaneous radioimmunoassay of progesterone, androst-4-enedione, pregnenolone, dehydroepiandrosterone and 17-hydroxyprogesterone in specific regions of human brain. *J. Steroid Biochem.*, 28 (1987) 317-325.
- 1099 Lagana, A., d'Ascenzo, G. and Marino, A.: An efficient procedure for extraction and determination of steroids in the tissue of laboratory animals. *Chromatographia*, 23 (1987) 796-802.
- 1100 Levin, W., Thomas, P.E., Ryan, D.E. and Wood, A.W.: Isozyme specificity of testosterone 7α -hydroxylation in rat hepatic microsomes: Is cytochrome P-450a the sole catalyst? *Arch. Biochem. Biophys.*, 258 (1987) 630-635.
- 1101 Liberato, D.J., Yergey, A.L., Esteban, N., Gomez-Sanchez, C.E. and Shackleton, C.H.L.: Thermospray HPLC/MS: a new mass spectrometric technique for the profiling of steroids. *J. Steroid Biochem.*, 27 (1987) 61-70.
- 1102 Matlin, S.A. and Wu, Z.Y.: Antihormonal agents. Part 2: HPLC of ZK 98734 and its metabolites and ZK 98299. *J. High Resolut. Chromatogr. Chromatogr. Commun.*, 10 (1987) 557-559.
- 1103 Matsuguchi, M., Dehennin, L., Habrioux, G., Matsuguchi-Moreau, L. and Degrelle, H.: Pregnenolone metabolism in Y-1 mouse adrenal cells: HPLC analysis and identification of metabolites by gas chromatography-mass spectrometry. *J. Steroid Biochem.*, 28 (1987) 311-315.
- 1104 Ng, L.L.: Reverse phase liquid chromatographic determination of dexamethasone acetate and cortisone acetate in bulk drug substances and dosage forms: method development. *J. Assoc. Off. Anal. Chem.*, 70 (1987) 829-833.
- 1105 Numazawa, M., Tsuji, M. and Mutsumi, A.: Studies on aromatase inhibition with 4-androstene-3,6,17-trione: its 3β -reduction and time-dependent irreversible binding to aromatase with human placental microsomes. *J. Steroid Biochem.*, 28 (1987) 337-344.
- 1106 Oka, K., Hirano, T. and Noguchi, M.: Combined use of rapid-flow fractionation and high-performance liquid chromatography for the determination of serum testosterone and application to study of stress response to physical exercise. *J. Chromatogr.*, 423 (1987) 285-291.
- 1107 Rao, P.N., Burdett, J.E., Jr., Moore, P.H., Jr. and Horton, R.: Isolation and identification of androstanediol glucuronide from human plasma. *J. Steroid Biochem.*, 28 (1987) 565-569.
- 1108 Senciall, I.R.: Primary and secondary progesterone hydroxylation by the rabbit liver microsomal cytochromes P-450 system. *J. Steroid Biochem.*, 28 (1987) 711-716.
- 1109 Swinney, D.C., Ryan, D.E., Thomas, P.E. and Levin, W.: Regioselective progesterone hydroxylation catalyzed by relevant rat hepatic cytochrome P-450 isozymes. *Biochemistry*, 26 (1987) 7073-7083.

- 1110 Tunn, S., Claus, S., Schulze, H., Braun, B.-E. and Krieg, M.: 5 α -Androstane-3 β ,17 β -diol hydroxylating enzymes in stroma and epithelium of human benign prostatic hyperplasia (BPH). *J. Steroid Biochem.*, 28 (1987) 257-265.

For additional information see:
C.A., 107 (1987) 90167q.

See also 823, 1129, 1366.

13b. Estrogens

- 1111 Carfagnini, G., Di Corcia, A., Marchetti, M. and Samperi, R.: Assay for urinary estriol during the menstrual cycle based on extraction by a graphitized carbon black cartridge followed by high-performance liquid chromatography. *Chromatographia*, 23 (1987) 835-838.
- 1112 Desta, B.: Complete separation of nine equine oestrogens by high-performance liquid chromatography. *J. Chromatogr.*, 435 (1988) 385-390.
- 1113 Fotsis, T.: The multicomponent analysis of estrogens in urine by ion exchange chromatography and GC-MS - II. Fractionation and quantitation of the main groups of estrogen conjugates. *J. Steroid Biochem.*, 28 (1987) 215-226.
- 1114 Fotsis, T. and Adlercreutz, H.: The multicomponent analysis of estrogens in urine by ion exchange chromatography and GC-MS - I. Quantitation of estrogens after initial hydrolysis of conjugates. *J. Steroid Biochem.*, 28 (1987) 203-213.
- 1115 Fullerton, F.R., Greenman, D.L. and Young, J.F.: Influence of a purified diet and route of administration on the metabolism and disposition of estradiol in B6C3F1 mice. *Drug Metab. Disp.*, 15 (1987) 602-607.
- 1116 Murphy, C., Fotsis, T., Adlercreutz, H. and Martin, F.: Analysis of tamoxifen and 4-hydroxytamoxifen levels in immature rat uterine cytoplasm and KCl-nuclear extracts by gas chromatography-mass spectrometry (GC-MS) using selected ion monitoring (SIM). *J. Steroid Biochem.*, 28 (1987) 289-299.
- 1117 Murphy, C., Fotsis, T., Pantzar, P., Adlercreutz, H. and Martin, F.: Analysis of tamoxifen, N-desmethyltamoxifen and 4-hydroxytamoxifen levels in cytosol and KCl-nuclear extracts of breast tumours from tamoxifen treated patients by gas chromatography-mass spectrometry (GC-MS) using selected ion monitoring (SIM). *J. Steroid Biochem.*, 28 (1987) 609-618.
- 1118 Rogerson, G.: Anabolic steroid screening in bovine urine. *Int. Lab.*, 17, No. 9 (1987) 60-62.
- 1119 Shimada, K. and Nambara, T.: Analysis of estrogens by HPLC with electrochemical detection. *Prog. HPLC*, 2 (Electrochem. Detect. Med. Chem.) (1987) 235-260; *C.A.*, 107 (1987) 127254u - a review with 102 refs.
- 1120 Simpson, D.M., Elliston, J.F. and Katzenellenbogen, J.A.: Desmethylnafoxidine aziridine: an electrophilic affinity label for the estrogen receptor with high efficiency and selectivity. *J. Steroid Biochem.*, 28 (1987) 233-245.
- 1121 Smilde, A.K., Bruins, C.H.P., Doornbos, D.A. and Vink, J.: Optimization of the reversed-phase high-performance liquid chromatographic separation of synthetic estrogenic and progestogenic steroids using the multi-criteria decision making method. *J. Chromatogr.*, 410 (1987) 1-12.

For additional information see:
C.A., 107 (1987) 216242f.

See also 1096.

13c. Sterols

- 1122 Armstrong, M.J. and Carey, M.C.: Thermodynamic and molecular determinants of sterol solubilities in bile salt micelles. *J. Lipid Res.*, 28 (1987) 1144-1155.
- 1123 Burkl, E., Logel, J. and Sinensky, M.: Endogenous sterol synthesis is not required for regulation of 3-hydroxy-3-methylglutaryl coenzyme A reductase by low density lipoprotein. *J. Lipid Res.*, 28 (1987) 1199-1205.
- 1124 Gale, D.J., Logan, R.I. and Rivett, D.E.: Detection of desmosterol in the internal lipids of wool fibers. *Text. Res. J.*, 57 (1987) 539-542; *C.A.*, 107 (1987) 200217r.

- 1125 Homberg, E. and Bielefeld, B.: Vergleichende Steringehaltsbestimmungen in Fetten and Ölen. Teil II: Enzymatische und colorimetrische Methodes. *Fett Wissen. Technol.- Fat Sci. Technol.*, 89 (1987) 353-355.
- 1126 Panini, S.R., Gupta, A., Sexton, R.C., Parish, E.J. and Rudney, H.: Regulation of sterol biosynthesis and of 3-hydroxy-3-methyl-glutaryl-coenzyme A reductase activity in cultured cells by progesterone. *J. Biol. Chem.*, 262 (1987) 14435-14440.
- 1127 Reichen, J., Karlaganis, G. and Kern, F., Jr.: Cholesterol synthesis in the perfused liver of pregnant hamsters. *J. Lipid Res.*, 28 (1987) 1046-1052.
- 1128 Saucier, S.E., Kandutsch, A.A., Phirwa, S. and Spencer, T.A.: Accumulation of regulatory oxysterols, 32-oxolanosterol and 32-hydroxylanosterol in mevalonate-treated cell cultures. *J. Biol. Chem.*, 262 (1987) 14056-14062.
- 1129 Yoshimoto, T., Ritani, A., Ohwada, K., Takashi, K., Kodera, Y., Matsushima, A., Saito, Y. and Inada, Y.: Polyethylene glycol derivative-modified cholesterol oxidase soluble and active in benzene. *Biochem. Biophys. Res. Commun.*, 148 (1987) 876-882.

For additional information see:

C.A., 107 (1987) 183672r.

13d. Bile acids and alcohols

- 1130 Baker, P.R., Vitale, G.C. and Siow, Y.F.: Medroxyprogesterone acetate- and ethinylestradiol-induced changes in biliary bile acids of the rat studied by high-performance liquid chromatography. *J. Chromatogr.*, 423 (1987) 63-73.
- 1131 Bellentani, S., Pecorari, M., Cordoma, P., Marchegiano, P., Manenti, F., Bosisio, E., DeFabiani, E. and Galli, G.: Taurine increases bile acid pool size and reduces bile saturation index in the hamster. *J. Lipid Res.*, 28 (1987) 1021-1027.
- 1132 Goto, J., Chikai, T., Makiuchi, K. and Nambara, T.: (Solid phase extraction of bile acids in biological fluids). *Bunseki Kagaku*, 36 (1987) 772-775.
- 1133 Ichimiya, H., Yanagisawa, J. and Nakayama, F.: Altered metabolism of bile alcohol and bile acid in complete extrahepatic cholestasis: qualitative and quantitative aspects. *J. Lipid Res.*, 28 (1987) 1028-1037.
- 1134 Rocchi, E., Bellentani, S., Silingardi, M., Borghi, A., Farina, F. and Casalgrandi, G.: Improved high performance liquid chromatographic determination of conjugated bile acids in serum. *Ital. J. Gastroenterol.*, 19 (1987) 105-109; C.A., 107 (1987) 130188f.
- 1135 Swobodnik, W., Zhang, Y.Y., Klueppelberg, U., Janowitz, P., Wechsler, J.G., Fuerst, P. and Ditschuneit, H.: Analysis of conjugated bile acids in human serum by high-performance liquid chromatography using post-column enzyme reaction and off-line fluorimetric determination. *J. Chromatogr.*, 423 (1987) 75-84.
- 1136 Thompson, M.B., Blair, P.C., Morris, R.W., Neptun, D.A., Deyo, D.F. and Popp, J.A.: Validation and application of a liquid-chromatographic/enzymatic assay for individual bile acids in the serum of rats. *Clin. Chem. (Winston-Salem)*, 33 (1987) 1856-1862.

For additional information see:

C.A., 107 (1987) 171762g, 194561e.

See also 1127.

13e. Ecdysones and other insect steroid hormones

- 1137 Bathori, M., Szendrei, K., Miklos, P., Pelczer, I. and Solymosi, P.: Ecdysteroids from *Silene nutans* L. *Symp. Biol. Hung.*, 34 (Chromatography '85) (1986) 241-250; C.A., 107 (1987) 150528v.
- 1138 O'Hanlon, G.M., Howarth, O.W. and Rees, H.H.: Identification of ecdysone 25-O- β -D-glucopyranoside as a new metabolite of ecdysone in the nematode *Parascaris equorum*. *Biochem. J.*, 248 (1987) 305-307.

13f. *Other steroids*

For additional information see:
C.A., 107 (1987) 205310a.

14. STEROID GLYCOSIDES AND SAPONINS

1139 Dasgupta, A., Yeo, K.-T., Malik, S., Sandu, P., Ahmad, S. and Kenny, M.: Two novel endogenous digoxin-like immunoreactive substances isolated from human plasma ultrafiltrate. *Biochem. Biophys. Res. Commun.*, 148 (1987) 623-628.

1140 Sener, B., Evren, N., Ozguven, M., Bingol, F. and Mutlugil, A.: Determination of some cardiac glycosides by high performance liquid chromatography. *J. Pharm. Belg.*, 42 (1987) 188-190; C.A., 107 (1987) 205268t.

For additional information see:
C.A., 107 (1987) 183644h.

15. TERPENES AND OTHER VOLATILE AROMATIC COMPOUNDS

15a. *Terpenes*

1141 Lin, L.-J. and Shiao, M.-S.: Separation of oxygenated triterpenoids from *Ganoderma lucidum* by high-performance liquid chromatography. *J. Chromatogr.*, 410 (1987) 195-200.

1142 Sezik, E. and Turkoz, S.: Separation of some triterpenic saponins by column chromatography. *Acta Pharm. Turc.*, 29 (1987) 87-93; C.A., 107 (1987) 223107u.

See also 1746.

15b. *Essential oils*

For additional information see:
C.A., 107 (1987) 161358r.

15c. *Bitter substances*

1143 Ono, M., Kakudo, Y., Yamamoto, R., Nagami, K. and Kumada, J.: Simultaneous analysis of hop bittering components by HPLC. II. Evaluation of hop deterioration. *J. Am. Soc. Brew. Chem.*, 45 (1987) 61-69; C.A., 107 (1987) 75998t.

16. NITRO AND NITROSO COMPOUNDS

1144 Bergens, A.: Determination of nitrodiphenylamines by liquid chromatography and dual-electrode amperometric detection. *J. Chromatogr.*, 410 (1987) 437-446.

1145 Bogusz, M. and Aderjan, R.: Improved standardization in reversed-phase high-performance liquid chromatography using 1-nitroalkanes as a retention index scale. *J. Chromatogr.*, 435 (1988) 43-53.

1146 Chiarotti, M., Fucci, N., Lucchini, P. and Carnevale, A.: Thin-layer and HPLC analyses of organic gunshot residues. *Nippon Hoigaku Zasshi*, 41 (1987) 65-70; C.A., 107 (1987) 170054r.

1147 Dong, S. and An, D.: Determination of glycerin trinitrate in nitroglycerin transdermal films by HPLC. *Zhongguo Yaoke Daxue Xuebao*, 18 (1987) 124-125; C.A., 107 (1987) 161779d.

1148 Kunugi, A., Tsuji, C. and Tabei, K.: HPLC of nitrophenols: Determination of products of on-column nitration of phenol. *J. High Resolut. Chromatogr. Chromatogr. Commun.*, 10 (1987) 624-625.

- 1149 Lloyd, J.B.F.: Liquid chromatography with electrochemical detection of explosives and firearms propellant traces. *Anal. Proc. (London)*, 24 (1987) 239-240; *C.A.*, 107 (1987) 179544h.
- 1150 Righezza, M., Murello, M.H. and Siouffi, A.M.: Determination of nitrosamines by liquid chromatography with post-column photolysis and electrochemical detection. *J. Chromatogr.*, 410 (1987) 145-155.
- 1151 Selavka, C.M., Tontarski, R.E., Jr. and Strobel, R.A.: Improved determination of nitrotoluenes using liquid chromatography with photolytically assisted thermal energy analysis (IC-PAT). *J. Forensic Sci.*, 32 (1987) 941-952; *C.A.*, 107 (1987) 128572h.
- 1152 Swynnerton, N., Hamil, H. and Seals, W.O.: Development of assay methods for liquid propellants. *Report*, ARAED-CR-86021 (1986) Order No. AD-A175387/GAR, 14 pp.; *C.A.*, 107 (1987) 201503t.

See also 866, 1183, 1813.

17. AMINES, AMIDES AND RELATED NITROGEN COMPOUNDS

17a. Amines and polyamines

- 1153 Chang, M.-Y., Chen, L.-R., Ding, X.-D., Selavka, C.M. and Krull, I.S.: Pre-column derivatization for improved detection in liquid chromatography-photolysis-electrochemistry. *J. Chromatogr. Sci.*, 25 (1987) 460-467.
- 1154 Dominguez, L.M. and Dunn, R.S.: Analysis of OPA-derivatized amino sugars in tobacco by high-performance liquid chromatography with fluorimetric detection. *J. Chromatogr. Sci.*, 25 (1987) 468-471.
- 1155 Dostal, A.: (Possibilities of analytical determination of stabilizers, based on hindered amines, for polymers). *Chem. Listy*, 81 (1987) 765-770; *C.A.*, 107 (1987) 155244j - a review with 28 refs.
- 1156 Ponzio, F., Achilli, G. and Perego, C.: Coulometric detection of brain monoamines and their metabolites: relevance in the study of functionality of central monoaminergic systems. *Prog. HPLC*, 2 (Electrochem. Detect. Med. Chem.) (1987) 307-328; *C.A.*, 107 (1987) 127255v - a review with 95 refs.
- 1157 Shibukawa, A., Nakagawa, T., Kaihara, A., Yagi, K. and Tanaka, H.: Reversed-phase high-performance liquid chromatography of substituted anilines utilizing molecular-recognizing ability of crown-ether: comparison with ion-pair chromatography. *Anal. Chem.*, 59 (1987) 2496-2501.

For additional information see:

C.A., 107 (1987) 171751c, 171755g.

See also 849, 906, 923, 934, 1146, 1682.

17b. Catecholamines and their metabolites

- 1158 Ahnert-Hilger, G., Bräutigam, M. and Gratzl, M.: Ca²⁺-stimulated catecholamine release from α -toxin-permeabilized PC12 cells: biochemical evidence for exocytosis and its modulation by protein kinase C and G proteins. *Biochemistry*, 26 (1987) 7842-7848.
- 1159 Dutrieu, J. and Delmotte, Y.A.: A rapid preparative procedure for the treatment of free urinary catecholamines. *J. Liq. Chromatogr.*, 10 (1987) 715-729.
- 1160 Elchisak, M.A.: Determination of sulfated biogenic amines and some metabolites by HPLC with electrochemical detection: Emphasis on dopamine sulfate isomers. *Prog. HPLC*, 2 (Electrochem. Detect. Med. Chem.) (1987) 329-375; *C.A.*, 107 (1987) 127256w - a review with 87 refs.
- 1161 Gupta, R.N., Steiner, M. and Lew, M.: Determination of unconjugated 3-methoxy-4-hydroxyphenylglycol by liquid chromatography for monitoring inhibition of monoamine oxidase activity in plasma. *Clin. Chem. (Winston-Salem)*, 33 (1987) 2078-2080.
- 1162 Hanai, J., Kawai, T., Sato, Y., Takasugi, N., Nishi, M. and Takeda, T.: Simple liquid-chromatographic measurement of vanillylmandelic acid and homovanillic acid in urine on filter paper for mass screening of neuroblastoma in infants. *Clin. Chem. (Winston-Salem)*, 33 (1987) 2043-2046.

- 1163 Iwamoto, T., Yoshiura, M. and Iriyama, K.: Liquid chromatographic identification of urinary catecholamine metabolites adsorbed on alumina. *J. Liq. Chromatogr.*, 10 (1987) 1217-1235.
- 1164 Morier, E. and Rips, R.: Automatic assay of biogenic amines and their metabolites in mice using high pressure liquid chromatography and electrochemical detection LCED. *Prog. HPLC*, 2 (Electrochem. Detect. Med. Chem.) (1987) 377-396; *C.A.*, 107 (1987) 127311k.
- 1165 Morier-Teissier, E. and Rips, R.: Determination of 3-hydroxy-4-methoxy-phenylethyleneglycol and other monoamine metabolites in brain areas using LC-EC and a two-column switching technique. *J. Liq. Chromatogr.*, 10 (1987) 1463-1483.
- 1166 Morier-Teissier, E. and Rips, R.: Catecholamine metabolite measurements in mouse brain using HPLC and electrochemical detection: Comparison of a one-column technique with a two-column switching technique. *Methods Enzymol.*, 142 (Metab. Aromat. Amino Acids Amines) (1987) 535-549; *C.A.*, 107 (1987) 168888x.
- 1167 Nyyssönen, K. and Parviainen, M.T.: Practical observations and sources of error in assays of plasma catecholamines by "high-performance" liquid chromatography with electrochemical detection. *Clin. Chem. (Winston-Salem)*, 33 (1987) 1938-1939.
- 1168 Ong, C.N., Lee, B.L., Ong, H.Y. and Jacob, E.: Simple high-performance liquid chromatographic method for the simultaneous measurement of vanillylmandelic acid and homovanillic acid. *J. Chromatogr.*, 423 (1987) 278-284.
- 1169 Pillai, D.N.: Inexpensive alternative to kit methods for extraction of urinary catecholamines. *Clin. Chem. (Winston-Salem)*, 33 (1987) 2122.
- 1170 Reeuwijk, H.J.E.M., Lingeman, H., De Jong, J., Tjaden, U.R. and van der Greef, J.: Automated, bioanalysis of catecholamines with reversed-phase liquid chromatography and amperometric detection. *Anal. Proc.*, 24 (1987) 301-303; *C.A.*, 107 (1987) 211947e.
- 1171 Richards, D.A. and Titheradge, A.C.: Simultaneous liquid chromatographic determination of vanillylmandelic acid, homovanillic acid, and 5-hydroxy-3-indoleacetic acid in urine, using isocratic elution and electrochemical detection. *Biomed. Chromatogr.*, 2 (1987) 115-119.
- 1172 Tippett, P.A., West, R.S., McEwan, A.J., Middleton, J.E. and Ackery, D.M.: A comparison of dopamine and homovanillic acid excretion, as prognostic indicators in malignant pheochromocytoma. *Clin. Chim. Acta*, 166 (1987) 123-133.
- 1173 Trouvin, J.H. and Billaud-Mesguich, E.: Determination of urinary metanephrines in man using liquid chromatography with electrochemical detection. *J. Liq. Chromatogr.*, 10 (1987) 731-747.
- 1174 Wakabayashi, H., Watanabe, K., Shibata, A., Yamato, S., Nakajima, M. and Shimada, K.: (Sample clean-up method for determination of catecholamines in plasma and heart by HPLC). *Bunseki Kagaku*, 36 (1987) 814-817.

For additional information see:

C.A., 107 (1987) 72213x, 147498e, 171757j.

See also 1544.

17c. Urea and guanidine derivatives

- 1175 Kobayashi, Y., Kubo, H. and Kinoshita, T.: Fluorometric determination of guanidino compounds by HPLC using water-soluble 9,10-phenanthrenequinone-3-sulfonate. *Anal. Sci.*, 3 (1987) 363-367; *C.A.*, 107 (1987) 194373v.
- 1176 Tyrefors, N. and Gillberg, P.G.: Determination of acetylcholine and choline in microdialysates from spinal cord of rat using liquid chromatography with electrochemical detection. *J. Chromatogr.*, 423 (1987) 85-91.

See also 1179, 1768.

17d. Other amine derivatives and amides (excluding peptides)

- 1177 Barnes, N.M., Costall, B., Fell, A.F. and Naylor, R.J.: An HPLC assay procedure of sensitivity and stability for measurement of acetylcholine and choline in neuronal tissue. *J. Pharm. Pharmacol.*, 39 (1987) 727-731; *C.A.*, 107 (1987) 168906b.
- 1178 Brondeau, M.T., Simon, P., Bonnet, P., Guenier, J.P. and De Ceaurriz, J.: Reversed-phase HPLC as a method of revealing nitromethane-induced false hypercreatininemia. *J. Liq. Chromatogr.*, 10 (1987) 175-190.

- 1179 Gordon, I.L., Kar, R., Opfell, R.W. and Wile, A.G.: Pharmacokinetics of hexamethylmelamine in intralipid following hepatic regional administration in rabbits. *Cancer Res.*, 47 (1987) 5070-5073.
- 1180 Hashizume, K., Toda, C., Yasui, T. and Nagano, H.: (Determination of tetramine in *Neptunea intersculpta* by HPLC). *Eisei Kagaku*, 33 (1987) 179-184; *C.A.*, 107 (1987) 152976p.
- 1181 Huang, C.B.: Determination of dialkyldimethylammonium salt in rolling oil. *J. Liq. Chromatogr.*, 10 (1987) 1103-1125.
- 1182 Neff, N.H., Meek, J.L., Hadjiconstantinou, M. and Laird, H.E.: Analysis of choline and acetylcholine in tissue by HPLC with electrochemical detection. *Prog. HPLC*, 2 (Electrochem. Detect. Med. Chem.) (1987) 181-191; *C.A.*, 107 (1987) 127253t - a review with 15 refs.
- 1183 Wigfield, Y.Y., Gurprasad, N.P., Lanouette, M. and Ripley, S.: Determination of N-nitrosodiethanolamine in dinoseb formulations by mass spectrometry and thermal energy analyzer detection. *J. Assoc. Off. Anal. Chem.*, 70 (1987) 792-796.
- See also 818, 917, 956, 1144, 1150.

18. AMINO ACIDS AND PEPTIDES; CHEMICAL STRUCTURE OF PROTEINS

18a. Amino acids and their derivatives

- 1184 Al-Najafi, S., Wellington, C.A., Wade, A.P., Sly, T.J. and Betteridge, D.: Computer-assisted optimization of HPLC with post-column reaction for the determination of amino acids. *Talanta*, 34 (1987) 749-756; *C.A.*, 107 (1987) 190089j.
- 1185 Anastasia, M., Colombo, D. and Fiecchi, A.: Quantitative high-performance liquid chromatographic determination of the amino acid spinacine in blood and chow of rats. *J. Chromatogr.*, 410 (1987) 504-508.
- 1186 Araki, A. and Sako, Y.: Determination of free and total homocysteine in human plasma by high-performance liquid chromatography with fluorescence detection. *J. Chromatogr.*, 422 (1987) 43-52.
- 1187 Ashworth, R.B.: Ion-exchange separation of amino acids with postcolumn orthophthalaldehyde detection. *J. Assoc. Off. Anal. Chem.*, 70 (1987) 248-252.
- 1188 Bidlingmeyer, B.A., Cohen, S.A., Tarvin, T.L. and Frost, B.: A new, rapid, high-sensitivity analysis of amino acids in food type samples. *J. Assoc. Off. Anal. Chem.*, 70 (1987) 241-247.
- 1189 Carlson, M. and Thompson, R.D.: Determination of *p*-aminobenzoic and salicylic acid salts in pharmaceuticals by high performance liquid chromatography. *J. Liq. Chromatogr.*, 10 (1987) 997-1009.
- 1190 Cotgreave, I.A. and Moldeus, P.: Methodologies for the analysis of reduced and oxidized N-acetylcysteine in biological systems. *Biopharm. Drug Dispos.*, 8 (1987) 365-375; *C.A.*, 107 (1987) 146688e.
- 1191 Csiba, A., Trezl, L., Rusznak, I. and Szarvas, T.: N^Y-Hydroxymethyl-L-arginines: newly discovered serum and urine components and their characterization by HPLC. *Symp. Biol. Hung.*, 34 (Chromatography '85) (1986) 57-62; *C.A.*, 107 (1987) 194370s.
- 1192 Dupuy, D. and Szabo, S.: Measurement of tissue sulfhydryls and disulfides in tissue protein and nonprotein fractions by high performance liquid chromatography using electrochemical detection. *J. Liq. Chromatogr.*, 10 (1987) 107-119.
- 1193 Eslami, M., Stuart, J.D. and Cohen, K.A.: Improvement in the resolution of *o*-phthalaldehyde derivatized amino acids by applying gradient steepness optimization to five reversed-phase columns of different lengths and particle sizes. *J. Chromatogr.*, 411 (1987) 121-138.
- 1194 Eslami, M., Stuart, J.D. and Dean, R.W.: Analysis of taurine in blood plasma of epileptic patients using an improved isocratic HPLC method for amino acids. *J. Liq. Chromatogr.*, 10 (1987) 977-995.
- 1195 Fuerst, P., Godel, H.G. and Graeser, T.A.: HPLC analysis of free amino acids in biological fluids. Methodological advantages, shortcomings and technical pitfalls. *Symp. Biol. Hung.*, 34 (Chromatography '85) (1986) 29-41; *C.A.*, 107 (1987) 150521n.

- 1196 Fukuhara, T. and Yuasa, S.: Rapid resolution of 2,4-dinitrophenyl-DL-amino acids on a native cellulose column. *J. Chromatogr.*, 411 (1987) 502-506.
- 1197 Goldsmith, R.F., Earl, J.W. and Cunningham, A.M.: Determination of δ -amino-butyric acid and other amino acids in cerebrospinal fluid of pediatric patients by reversed-phase liquid chromatography. *Clin. Chem. (Winston-Salem)*, 33 (1987) 1736-1740.
- 1198 Hayashi, K., Maeda, Y., Toyomizu, M. and Tomita, Y.: HPLC method for the analysis of N_γ-methylhistidine in food, chicken excreta, and rat urine. *J. Nutr. Sci. Vitaminiol.*, 33 (1987) 151-156; *C.A.*, 107 (1987) 76246h.
- 1199 Hearn, M.T.W., Aguilar, M.I., Nguyen, T. and Fridman, M.: High-performance liquid chromatography of amino acids, peptides and proteins. LXXXIV. Application of derivative spectroscopy to the study of column residency effects in the reversed-phase and size exclusion liquid chromatographic separation of proteins. *J. Chromatogr.*, 435 (1988) 271-284.
- 1200 Igarashi, P., Gulyassy, P., Stanfel, L. and Depner, T.: Plasma hippurate in renal failure: HPLC method and clinical application. *Nephron*, 47 (1987) 290-294; *C.A.*, 107 (1987) 214313f.
- 1201 Iinuma, F., Tabara, M. and Watanabe, M.: (Fluorometric determination of 3-hydroxyanthranilic acid in biological fluids by HPLC). *Bunseki Kagaku*, 36 (1987) 776-781.
- 1202 James, L.B.: Amino acid analysis. II. Nuances of integration. *J. Chromatogr.*, 410 (1987) 176-180.
- 1203 Jansen, E.H.J.M. and Both-Miedema, R.: Instability of phenylthiohydantoin amino acids. *J. Chromatogr.*, 435 (1988) 363-367.
- 1204 Kleinmann, I., Svoboda, V. and Vockova, J.: Preparative separation of an enzymatic hydrolysate of ¹⁴C-labelled proteins. *J. Chromatogr.*, 411 (1987) 335-344.
- 1205 Lepri, L., Desideri, P.G., Merlini, L. and Checchini, L.: HPLC of aminoacids on silica coated with ammonium tungstophosphate. *Chromatographia*, 23 (1987) 814-817.
- 1206 Leyrer, J. and Schuster, R.: (HPLC determination of amino acids in parenteral solutions). *LaborPraxis*, 11 (1987) 922-925; *C.A.*, 107 (1987) 183700y.
- 1207 Montanarella, L.: Relative merits of pre- and post-column derivatization in amino acid analysis by HPLC. *Chim. Oggi*, (1987) 27-32; *C.A.*, 107 (1987) 194368x.
- 1208 Muramoto, K., Sunahara, S. and Kamiya, H.: Measurement of tryptophan in peptides by acid hydrolysis in the presence of phenol and its application to the amino acid sequence of a sea anemone toxin. *Agric. Biol. Chem.*, 51 (1987) 1607-1616; *C.A.*, 107 (1987) 73591f.
- 1209 Ohtsuki, K., Kawabata, M., Kokura, H. and Taguchi, K.: Simultaneous determination of S-methylmethionine, vitamin U, and free amino acids in extracts of green tea with an HPLC-amino acid analyzer. *Agric. Biol. Chem.*, 51 (1987) 2479-2484; *C.A.*, 107 (1987) 196615z.
- 1210 Palmerini, C.A., Fini, C., Cantelmi, M.G. and Floridi, A.: Assessment of taurine in plasma and urine by anion-exchange high-performance liquid chromatography with pre-column derivatization. *J. Chromatogr.*, 423 (1987) 292-296.
- 1211 Penney, C.L., Landi, S., Shah, P., Leung, K.H. and Archer, M.C.: Analysis of the immunoadjuvant octadecyl tyrosine hydrochloride. *J. Biol. Stand.*, 14 (1986) 345-349; *C.A.*, 107 (1986) 183681t.
- 1212 Qureshi, G.A., Gutierrez, A. and Bergstroem, J.: Determination of histidine, 1-methylhistidine and 3-methylhistidine in biological samples by HPLC. Clinical application of urinary 3-methylhistidine in evaluating the muscle protein breakdown in uremic patients. *Symp. Biol. Hung.*, 34 (Chromatography '85) (1986) 103-115; *C.A.*, 107 (1987) 150524r.
- 1213 Rajendra, W.: High performance liquid chromatographic determination of amino acids in biological samples by precolumn derivatization with o-phthaldialdehyde. *J. Liq. Chromatogr.*, 10 (1987) 941-955.
- 1214 Sandor, A., Kispal, G., Melegh, B. and Alkonyi, I.: Ester composition of carnitine in the perfusate of liver and in the plasma of donor rats. *Eur. J. Biochem.*, 170 (1987) 443-445.
- 1215 Schrynmackers-Pitance, P. and Schoos-Barbette, S.: Determination of aromatic and neutral aminoacids by HPLC in blood specimens collected on filter paper. *Clin. Chim. Acta*, 166 (1987) 91-97.
- 1216 Walker, T.A. and Pietrzyk, D.J.: Ion-interaction chromatographic separation of free amino acids. *J. Liq. Chromatogr.*, 10 (1987) 161-174.

- 1217 Watanabe, N., Toyo'oka, T. and Imai, K.: HPLC electrochemical fluorometric detection of amino acids including tryptophan using 4-fluoro-7-nitrobenzo-2-oxa-1,3-diazole. *Biomed. Chromatogr.*, 2 (1987) 99-103.
- 1218 Woolfson, A.D., Millership, J.S. and Karin, E.F.I.A.: Determination of the sulphoxide metabolites of S-carboxymethyl-L-cysteine by high-performance liquid chromatography with electrochemical detection. *Analyst (London)*, 112 (1987) 1421-1425.
- 1219 Xue, X. and Yan, P.: (The optimum conditions of amino acid analysis in honey). *Shentwu Huaxue Yu Shengwu Wuli Jinzhan*, (1987) 70-71; *C.A.*, 107 (1987) 152969p.
- 1220 Yamaguchi, Y., Haginaka, J., Kunitomo, M., Yasuda, H. and Bando, Y.: High-performance liquid chromatographic determination of desmosine and isodesmosine in tissues and its application to studies of alteration of elastin induced by atherosclerosis. *J. Chromatogr.*, 422 (1987) 53-59.

For additional information see:

C.A., 107 (1987) 70906b, 73605p, 92878w, 82963v, 95406b, 130197h, 146774e, 150522p, 174498m.

See also 827, 859, 868, 908, 912, 923, 928, 929, 1020, 1153, 1223, 1319, 1365, 1549, 1809.

18b. *Peptides and peptidic and proteinous hormones*

- 1221 Abhold, R.H., Sullivan, M.J., Wright, J.W. and Harding, J.W.: Binding, degradation and pressor activity of angiotensins II and III after aminopeptidase inhibition with amastatin and bestatin. *J. Pharmacol. Exp. Ther.*, 242 (1987) 957-962.
- 1222 Bahouth, S.W. and Malbon, C.C.: Human β -adrenergic receptors. Simultaneous purification of β_1 and β_2 -adrenergic-receptor peptides. *Biochem. J.*, 248 (1987) 557-566.
- 1223 Bak, H.J. and Beintema, J.J.: *Panulirus interruptus* hemocyanin. The elucidation of the complete amino acid sequence of subunit a. *Eur. J. Biochem.*, 169 (1987) 333-348.
- 1224 Brown, D.S. and Jenke, D.R.: Determination of trace levels of oxytocin in pharmaceutical solutions by high-performance liquid chromatography. *J. Chromatogr.*, 410 (1987) 157-168.
- 1225 Burke, T.R., Jr. and Knight, M.: Counter-current chromatographic purification of [D-Ala¹]peptide T amide. *J. Chromatogr.*, 411 (1987) 431-435.
- 1226 Cantor, P. and Rehfeld, J.F.: The molecular nature of cholecystokinin in human plasma. *Clin. Chim. Acta*, 168 (1987) 153-158.
- 1227 Chu, Y., Wang, R., Burke, G.T., Chanley, J.D. and Katsoyannis, P.G.: Possible involvement of the A²⁰-A²¹ peptide bond in the expression of the biological activity of insulin. 1. [21-des-Asparagine, 20-cysteinamide-A]insulin and [21-desasparagine, 20-cysteine isopropylamide-A]insulin. *Biochemistry*, 26 (1987) 6966-6971.
- 1228 Conlon, J.M., Davis, M.S., Falkmer, S. and Thim, L.: Structural characterization of peptides derived from prosomatostatins I and II isolated from the pancreatic islets of two species of teleostean fish: the daddy sculpin and the flounder. *Eur. J. Biochem.*, 168 (1987) 647-652.
- 1229 Conlon, J.M., Eriksson, B., Grimelius, L., Oberg, K. and Thim, L.: Characterization of three peptides derived from prosomatostatin [prosomatostatin-(1-63)-, -(65-76)- and -(79-92)-peptides] in a human pancreatic tumour. *Biochem. J.*, 248 (1987) 123-127.
- 1230 Davies, J.G., Muir, A.V., Rose, K. and Offord, R.E.: Identification of radioactive insulin fragments liberated by insulin proteinase during the degradation of semisynthetic [³H]Gly^{A1}]insulin and [³H]Phe^{B1}]insulin. *Biochem. J.*, 249 (1988) 209-214.
- 1231 DiAugustine, R.P., Gibson, B.W., Aberth, W., Kelly, M., Ferrua, C.M., Tomooka, Y., Brown, C.F. and Walker, M.: Evidence for isoaspartyl (deamidated) forms of mouse epidermal growth factor. *Anal. Biochem.*, 165 (1987) 420-429.
- 1232 Draviam, E.J., Greeley, G.H., Jr., Beauchamp, R.D. and Thompson, J.C.: A high pressure liquid chromatography-radioimmunoassay method for measurement of cholecystokinin-8 and cholecystokinin-33/39 in plasma. *J. Liq. Chromatogr.*, 10 (1987) 1431-1439.

- 1233 Bonzales de Llano, D., Ramos, M. and Polo, C.: Gel filtration and high-performance liquid chromatographic analysis of phosphotungstic acid soluble peptides from blue cheeses. *Chromatographia*, 23 (1987) 764-766.
- 1234 Hillar, M., Santarelli, I., Stolzmann, Z., Wafeeg, W., Allen, S., Chan, J.Y.H., Patt, L.M., Houck, J.C. and Wyborny, L.E.: Nuclear peptides from calf liver: large scale isolation and fractionation: control of gene expression in cell-free systems, and inhibition of growth of cells in culture. *Basic Appl. Histochem.*, 31 (1987) 299-313; *C.A.*, 107 (1987) 214315h.
- 1235 Hirota, M., Shimizu, C., Nishino, T. and Shima, K.: A large molecular form of glucagon-like peptide-1 (CLP-1) immunoreactivity is co-released with glucagon from pancreas by arginine in normal subjects. *Clin. Chim. Acta*, 167 (1987) 293-302.
- 1236 Jones, R.M.L., Rose, K. and Offord, R.E.: Semisynthetic human [$^3\text{H}_2$]Phe 1]pro-insulin. *Biochem. J.*, 247 (1987) 785-788.
- 1237 Kawasaki, H., Imajoh, S. and Suzuki, K.: Separation of peptides on the basis of the difference in positive charge: Simultaneous isolation of C-terminal and blocked N-terminal peptides from tryptic digests. *J. Biochem. (Tokyo)*, 102 (1987) 393-400.
- 1238 Liu, D. and Desiderio, D.M.: Reversed-phase high-performance liquid chromatographic-radioreceptor assay of human cerebrospinal fluid neuropeptides. Maximizing recovery of picomoles of peptides and minimizing memory. *J. Chromatogr.*, 422 (1987) 61-71.
- 1239 Marchetti, P., Benzi, L., Cecchetti, P., Trischitta, V., Brunetti, A., Ciccarone, A.M. and Navalesi, R.: Intracellular processing of A14- ^{125}I -insulin: Analysis in isolated human monocytes by HPLC. *Serono Symp. Publ. Raven Press.*, 37 (Adv. Models Ther. Insulin-Depend. Diabetes) (1986) 129-133; *C.A.*, 107 (1987) 169167y.
- 1240 Mayo, K.H. and Burke, C.: Structural and dynamical comparison of α , β and γ forms of murine epidermal growth factor. *Eur. J. Biochem.*, 169 (1987) 201-207.
- 1241 Miller, K.F., Goldsby, R.A. and Bolt, D.J.: Immunoaffinity chromatography of bovine FSH using monoclonal antibodies. *J. Endocrinol.*, 115 (1987) 283-288; *C.A.*, 107 (1987) 191124d.
- 1242 Millington, W.R., O'Donohue, T.L. and Mueller, G.P.: Dopaminergic agents selectively alter the post-translational processing of β -endorphin in the intermediate pituitary of the rat. *J. Pharmacol. Exp. Ther.*, 243 (1987) 160-170.
- 1243 Mulac-Jericevic, B. and Atassi, M.Z.: Profile of the α -bungarotoxin-binding regions on the extracellular part of the α -chain of *Torpedo californica* acetylcholine receptor. *Biochem. J.*, 248 (1987) 847-852.
- 1244 Nakano, M., Kai, M., Ohno, M. and Ohkura, Y.: High-performance liquid chromatography of N-terminal tyrosine-containing oligopeptides by pre-column fluorescence derivatization with hydroxylamine, cobalt(II) and borate reagents. *J. Chromatogr.*, 411 (1987) 305-311.
- 1245 Ong, H., Lazure, C., Nguyen, T.T., McNicoll, N., Seidah, N., Chretien, M. and de Lean, A.: Bovine adrenal chromaffin granules are a site of synthesis of atrial natriuretic factor. *Biochem. Biophys. Res. Commun.*, 147 (1987) 957-963.
- 1246 Riggan, R.M., Shaar, C.J., Dodrulla, G.K., Lefeber, D.S. and Miner, D.J.: High-performance size-exclusion chromatographic determination of the potency of biosynthetic human growth hormone products. *J. Chromatogr.*, 435 (1988) 307-318.
- 1247 Rose, K., Herrero, C., Proudfoot, A.E.I., Offord, R.E. and Wallace, C.J.A.: Enzyme-assisted semisynthesis of polypeptide active esters and their use. *Biochem. J.*, 249 (1988) 83-88.
- 1248 Savoy, L.-A., Jones, R.M.L., Pochon, S., Davies, J.G., Muir, A.V., Offord, R.E. and Rose, K.: Identification by fast atom bombardment mass spectrometry of insulin fragments produced by insulin proteinase. *Biochem. J.*, 249 (1988) 215-222.
- 1249 Tashiro, Y., Suzuki, A., Noda, K. and Noguchi, H.: (Determination of immunostimulative acylpeptide (FK565) in plasma by HPLC with fluorogenic postcolumn derivatization). *Bunseki Kagaku*, 36 (1987) 787-791.
- 1250 Taylor, T.C. and Kimmel, J.R.: Iodinated derivatives of the hormone avian pancreatic polypeptide. *Anal. Biochem.*, 166 (1987) 194-203.
- 1251 Tode, B., Messeri, G., Bassi, F., Pazzagli, M. and Serio, M.: Chemiluminescence immunoassay for somatomedin C in serum. *Clin. Chem. (Winston-Salem)*, 33 (1987) 1989-1993.

- 1252 Torok, A., Kovacs, K. and Szokan, G.: HPLC of peptides containing basic non-proteinogenic amino acids. *Symp. Biol. Hung.*, 34 (Chromatography '85) (1986) 117-127; *C.A.*, 107 (1987) 176469b.
- 1253 Trethewey, J.: Moving away from animal tests for the standardization of insulin formulations. *Trends Pharmacol. Sci.*, 8 (1987) 287-289; *C.A.*, 107 (1987) 183371s.
- 1254 Van Damme, J., Cayphas, S., van Snick, J., Conings, R., Put, W., Lenaerts, J.-P., Simpson, R.J. and Billiau, A.: Purification and characterization of human fibroblast-derived hybridoma growth factor identical to T-cell-derived B-cell stimulatory factor-2 (interleukin-6). *Eur. J. Biochem.*, 168 (1987) 543-550.
- 1255 Venn, R.F.: Combined high-performance liquid chromatographic-radioimmunoassay method for the analysis of endorphins, enkephalins and other neurotransmitter peptides. *J. Chromatogr.*, 423 (1987) 93-104.
- 1256 Yiangou, Y., Di Marzo, V., Spokes, R.A., Panico, M., Morris, H.R. and Bloom, S.R.: Isolation, characterization, and pharmacological actions of peptides histidine valine 42, a novel prepro-vasoactive intestinal peptidederived peptide. *J. Biol. Chem.*, 262 (1987) 14010-14013.
- 1257 Zhu, Q., Singh, A.V., Bateman, A., Esch, F. and Solomon, S.: The corticostatic (anti-ACTH) and cytotoxic activity of peptides isolated from fetal, adult and tumor-bearing lung. *J. Steroid Biochem.*, 27 (1987) 1017-1022.

For additional information see:

C.A., 107 (1987) 71570z, 90135c, 95420b, 127315q, 128023e, 128559j, 168901w, 191104x.

See also 861, 1199, 1274, 1280, 1310, 1319, 1372, 1471, 1552, 1773.

18c. *Elucidation of structure of proteins and enzymes*

- 1258 Ametani, A., Kaminogawa, S., Shimizu, M. and Yamauchi, K.: Rapid screening of antigenically reactive fragments of α s₁-casein using HPLC and ELISA. *J. Biochem. (Tokyo)*, 102 (1987) 421-425.
- 1259 Anderson, G.P., Sanderson, D.G., Lee, C.H., Anderson, L.B. and Gross, E.L.: The effect of ethylenediamine chemical modification of plastocyanin on the rate of cytochrome oxidation and P-700⁺ reduction. *Biochim. Biophys. Acta*, 894 (1987) 386-398.
- 1260 Cairns, M.T., Alvarez, J., Panico, M., Gibbs, A.F., Morris, H.R., Chapman, D. and Baldwin, S.A.: Investigation of the structure and function of the human erythrocyte glucose transporter by proteolytic dissection. *Biochim. Biophys. Acta*, 905 (1987) 295-310.
- 1261 Hawke, D.H., Lahm, H.-W., Shively, J.E. and Todd, C.W.: Microsequence analysis of peptides and proteins: trimethylsilylisothiocyanate as a reagent for COOH-terminal sequence analysis. *Anal. Biochem.*, 166 (1987) 298-307.
- 1262 Hayashi, T., Hayashi, H. and Iwai, K.: *Tetrahymena* histone H1. Isolation and amino acid sequence lacking the central hydrophobic domain conserved in other H1 histones. *J. Biochem. (Tokyo)*, 102 (1987) 369-376.
- 1263 Ito, W., Nishimura, M., Sakato, N., Fujio, H. and Arata, Y.: A ¹H NMR method for the analysis of antigen-antibody interactions: Binding of a peptide fragment of lysozyme to anti-lysozyme monoclonal antibody. *J. Biochem. (Tokyo)*, 102 (1987) 643-649.
- 1264 Jänig, G.-R., Kraft, R., Blacnk, J., Ristau, O., Rabe, H. and Ruckpaul, K.: Chemical modification of cytochrome P-450 LM4. Identification of functionally linked tyrosine residues. *Biochim. Biophys. Acta*, 916 (1987) 512-523.
- 1265 Lee, F.-T., Ahamad, Z., DePaoli-Roach, A.A. and Roach, P.J.: Multiple phosphorylation of mouse muscle glycogen synthase. *Arch. Biochem. Biophys.*, 258 (1987) 615-620.
- 1266 Minchiotti, L., Galliano, M., Iadarola, P., Stoppini, M., Ferri, G. and Castellani, A.A.: Structural characterization of two genetic variants of human serum albumin. *Biochim. Biophys. Acta*, 916 (1987) 411-418.
- 1267 Niyibizi, C., Wu, J.-J. and Eyre, D.R.: The carboxypropeptide trimer of type II collagen is a prominent component of immature cartilages and intervertebral-disc tissue. *Biochim. Biophys. Acta*, 916 (1987) 493-499.
- 1268 Nomura, K., Mikami, B. and Morita, Y.: Partial amino acid sequences around sulfhydryl groups of soybean β -amylase. *J. Biochem. (Tokyo)*, 102 (1987) 341-349.

- 1269 Reiser, K.M., Hennessy, S.M. and Last, J.A.: Analysis of age-associated changes in collagen crosslinking in the skin and lung in monkeys and rats. *Biochim. Biophys. Acta*, 926 (1987) 339-348.
- 1270 Soeda, S., Kakiki, M., Shimeno, H. and Nagamatsu, A.: Localization of the binding sites of porcine tissue-type plasminogen activator and plasminogen to heparin. *Biochim. Biophys. Acta*, 916 (1987) 279-287.
- 1271 Yoshino, H. and Minari, O.: Heat-induced dissociation of human erythrocyte spectrin dimer into monomers. *Biochim. Biophys. Acta*, 905 (1987) 100-108.

See also 1304, 1310.

19. PROTEINS

19a. General techniques

- 1272 Balen'kii, B.G., Zimina, T.M. and Mal'tsev, V.G.: Study of protein association and micelle formation in surfactants by microcolumn exclusion chromatography. *Symp. Biol. Hung.*, 34 (Chromatography '85) (1986) 193-206; *C.A.*, 107 (1987) 150525s.
- 1273 Berkowitz, S.A., Henry, M.P., Nau, D.R. and Crane, L.J.: A strategic approach to the use of silica-based chromatography media (bonded phases) for protein purification. *Am. Lab. (Fairfield)*, 19 (1987) 33-42; *C.A.*, 107 (1987) 130173x.
- 1274 Hochuli, E., Döbeli, H. and Schacher, A.: New metal chelate adsorbent selective for proteins and peptides containing neighbouring histidine residues. *J. Chromatogr.*, 411 (1987) 177-184.
- 1275 Kim, Y.S., Sands, B.W. and Bass, J.L.: Protein loading capacity and textural properties of column packings in reversed-phase HPLC. *J. Liq. Chromatogr.*, 10 (1987) 839-851.
- 1276 Mazsaroff, I., Rounds, M.A. and Regnier, F.E.: Facile preparation of a non-porous strong anion-exchange column for proteins. *J. Chromatogr.*, 411 (1987) 452-455.
- 1277 McCall, M.N. and Easterbrook-Smith, S.B.: High-performance liquid chromatography-based assay for the quantitation of protein maleylation at the picomole level. *Anal. Biochem.*, 166 (1987) 440-445.
- 1278 Nesterenko, A.A., Nesterenko, M.A., Borukhov, S.I., Izotova, L.S., Gaida, G.Z., Rebentish, B.A., Novikov, A.A., Demyanova, N.G., Bogush, V.G. and Strongin, A.Ya.: (Usage of fast protein liquid chromatography for microbial protein isolation). *Biotehnologiya*, 3 (1987) 139-145; *C.A.*, 107 (1987) 194364t.
- 1279 Ueda, T., Ishida, Y. and Yasui, Y.: (HPLC separation of proteins and nucleic acids). *Shimadzu Hyoron*, 44 (1987) 39-50; *C.A.*, 107 (1987) 171659d - a review with 14 refs.
- 1280 Unger, K.K.: (Phase systems and operation - HPLC of biopolymers). *GIT-Suppl.*, (1987) 14-20; *C.A.*, 107 (1987) 194327h - a review with 18 refs.

For additional information see:

C.A., 107 (1987) 93080s, 171747f, 171748g.

See also 861, 887, 1199.

19b. Proteins of cells, viruses and subcellular particles

- 1281 Bronskill, P.M. and Wong, J.T.-F.: Suppression of fluorescence of tryptophan residues in proteins by replacement with 4-fluorotryptophan. *Biochem. J.*, 249 (1988) 305-308.
- 1282 Cannon, M., Threadgill, G.J., Mount, R. and Conrad, R.C.: Identification and analysis of yeast ribosomal proteins by HPLC. *Biochem. Soc. Trans.*, 15 (1987) 1040-1041; *C.A.*, 107 (1987) 130220k.
- 1283 Hirano, H., Eckart, K., Kimura, M. and Wittmann-Liebold, B.: Semi-preparative HPLC purification of ribosomal proteins from *Bacillus stearothermophilus* and sequence determination of the highly conserved protein S19. *Eur. J. Biochem.*, 170 (1987) 149-157.

- 1284 Hosey, M., Barhanin, J., Schmid, A., Vandaele, S., Ptasienski, J., O'Callahan, C., Cooper, C. and Lazdunski, M.: Photoaffinity labelling and phosphorylation of a 165 kilodalton peptide associated with dihydropyridine and phenylalkyl-amine-sensitive calcium channels. *Biochem. Biophys. Res. Commun.*, 147 (1987) 1137-1145.
- 1285 Mohanty, J.G. and Elazhary, Y.: High-performance liquid chromatographic separation and immunological characterization of soluble bovine viral diarrhea virus antigen. *J. Chromatogr.*, 435 (1988) 149-158.
- 1286 Murphey, R.J. and Gerner, E.W.: Hypusine formation in protein by a two-step process in cell lysates. *J. Biol. Chem.*, 262 (1987) 15033-15036.
- 1287 Szewczyk, A., Nalecz, M.J., Broger, C., Wojtczak, L. and Azzi, A.: Purification by affinity chromatography of the dicarboxylate carrier from bovine heart mitochondria. *Biochim. Biophys. Acta*, 894 (1987) 252-260.

For additional information see:

C.A., 107 (1987) 194497p.

19c. *Proteins synthesized by genetic manipulation*

- 1288 Kikuchi, N., Nagata, K., Horii, T., Miyazaki, S., Shin, M., Takimoto, N., Tsuruta, Y., Tamaki, M., Teraoka, H. and Yoshida, N.: Production of recombinant human pancreatic secretory trypsin inhibitor by *Escherichia coli*. *J. Biochem. (Tokyo)*, 102 (1987) 607-612.
- 1289 Ross, A.H., Herlyn, D. and Koprowski, H.: Purification of monoclonal antibodies from ascites using ABX liquid chromatography column. *J. Immunol. Methods*, 102 (1987) 227-231; C.A., 107 (1987) 215819n.

19d. *Microbial and plant proteins*

- 1290 Berger, G., Girault, G., Andre, F. and Galmiche, J.-M.: Application of HPLC to the purification of coupling factor CF₁ from spinach chloroplasts and of some of its subunits. *J. Liq. Chromatogr.*, 10 (1987) 1507-1517.
- 1291 Georgiou, C.D. and Webster, D.A.: Purification and partial characterization of the membrane-bound cytochrome *c*(561, 564) from *Vitreoscilla*. *Biochemistry*, 26 (1987) 6521-6526.
- 1292 Goodenough, U.W., Gebhardt, B., Mermall, V., Mitchell, D.R. and Heuser, J.E.: High-pressure liquid chromatography fractionation of *Chlamydomonas* dynein extracts and characterization of inner-arm dynein subunits. *J. Mol. Biol.*, 194 (1987) 481-494; C.A., 107 (1987) 73576e.
- 1293 Lambert, N., Plumb, G.W. and Wright, D.J.: A fast protein liquid chromatography procedure for the assessment of subunit heterogeneity of plant 11S storage globulins. *Biochem. Soc. Trans.*, 15 (1987) 538-539; C.A., 107 (1987) 73590e.
- 1294 Lobarzewski, J. and Waksmundzki, A.: The optimization of the separation of fungal proteins by affinity chromatography on controlled-porosity glass modified by boron and with vanillin as a ligand. *Symp. Biol. Hung.*, 34 (Chromatography '85) (1986) 223-232; C.A., 107 (1987) 150527u.
- 1295 Tashiro, M., Hashino, K., Shiozaki, M., Ibuki, F. and Maki, Z.: The complete amino acid sequence of rice bran trypsin inhibitor. *J. Biochem. (Tokyo)*, 102 (1987) 297-306.
- 1296 Tice, P.A., Robinson, G.W. and Regnier, F.E.: Preparative HPLC of soybean trypsin inhibitor using large particle diameter supports. *J. Liq. Chromatogr.*, 10 (1987) 1439-1461.
- 1297 Zanetti, G. and Merati, G.: Interaction between photosystem I and ferredoxin. Identification by chemical cross-linking of the polypeptide which binds ferredoxin. *Eur. J. Biochem.*, 169 (1987) 143-146.

For additional information see:

C.A., 107 (1987) 92884v, 130174y, 130221m, 150507n, 174497k.

See also 1278, 1283, 1327.

19e. *Proteins of blood, serum and blood cells*

- 1298 Akerström, B., Lögdberg, L., Babiker-Mohamed, H., Lohmander, S. and Rask, L.: Structural relationship between α_1 -microglobulin from man, guinea-pig, rat and rabbit. *Eur. J. Biochem.*, 170 (1987) 143-148.
- 1299 Churakova, N.I., Filatova, T.N. and Titova, E.V.: (Preparation of a rabbit serum albumin-free globulin fraction). *Khim.-Farm. Zh.*, 21 (1987) 874-876; *C.A.*, 107 (1987) 161472y.
- 1300 Danze, P.M., Tarjoman, A., Bousseaux, J., Fossati, P. and Dautrevaux, M.: Evidence for an increased glycation of IgG in diabetic patients. *Clin. Chim. Acta*, 166 (1987) 143-153.
- 1301 Folena-Wasserman, G., Inacker, R. and Rosenbloom, J.: Assay, purification and characterization of a recombinant malaria circumsporozoite fusion protein by high-performance liquid chromatography. *J. Chromatogr.*, 411 (1987) 345-354.
- 1302 Hardouin, S., Hossenlopp, P., Segovia, B., Seurin, D., Portolan, G., Lassarre, C. and Binoux, M.: Heterogeneity of insulin-like growth factor binding proteins and relationships between structure and affinity. 1. Circulating forms in man. *Eur. J. Biochem.*, 170 (1987) 121-132.
- 1303 Higashiyama, S., Ohkubo, I., Ishiguro, H., Sasaki, M., Matsuda, T. and Nakamura, R.: Heavy chain of human high molecular weight and low molecular weight kininogens binds calcium ion. *Biochemistry*, 26 (1987) 7450-7458.
- 1304 Hochuli, E., Gillessen, D. and Kocher, H.P.: Specificity of the immunoabsorbent used for large-scale recovery of interferon α -2a. *J. Chromatogr.*, 411 (1987) 371-378.
- 1305 Hossenlopp, P., Seurin, D., Segovia, B., Portolan, G. and Binoux, M.: Heterogeneity of insulin-like growth factor binding proteins and relationships between structure and affinity. 2. Forms released by human and rat liver in culture. *Eur. J. Biochem.*, 170 (1987) 133-142.
- 1306 Hou, K.C., Mandaro, R.M. and Roy, A.: Plasma protein separation by ion exchange and affinity cartridges. *Symp. Biol. Hung.*, 34 (Chromatography '85) (1986) 207-221; *C.A.*, 107 (1987) 150526t.
- 1307 Ikegawa, S., Tanase, S., Araki, S. and Morino, Y.: Quantitative detection of a variant prealbumin associated with type 1 familial amyloidotic polyneuropathy (Japanese type) by high-performance liquid chromatography. *Clin. Chim. Acta*, 167 (1987) 165-172.
- 1308 Kimura, M., Sueyoshi, T., Takada, K., Tanaka, K., Morita, T. and Iwanaga, S.: Isolation and characterization of ornitho-kininogen. *Eur. J. Biochem.*, 168 (1987) 493-501.
- 1309 Kuwajima, S. and Okuda, K.: Affinity chromatography for C-reactive protein using *p*-nitrophenylphosphorylcholine as a ligand. *Clin. Chim. Acta*, 166 (1987) 101-102.
- 1310 Mackinnon, C.M., Carter, P.E., Smyth, S.J., Dunbar, B. and Fothergill, J.E.: Molecular cloning of cDNA for human complement component C1s. The complete amino acid sequence. *Eur. J. Biochem.*, 169 (1987) 547-553.
- 1311 Miribel, L. and Arnaud, P.: Fractionation of plasma proteins using dye-ligand chromatography: elution profiles on immobilized green TSK-AF. *J. Biochem. Biophys. Methods*, 14 (1987) 291-302; *C.A.*, 107 (1987) 171758k.
- 1312 Smith, K.J.: Blood-coagulation factor IX-containing therapeutic blood product, means and methods of preparing same by affinity chromatography. *Eur. Pat. Appl.* EP 229,026 (Cl. CO7K3/18), 15 Jul. 1987, US Appl. 816,291, 06 Jan. 1986; 22 pp.; *C.A.*, 107 (1987) 161669t.
- 1313 Strahler, J.R., Rosenblum, B.B. and Hanash, S.M.: Identification and characterization of a human transthyretin variant. *Biochem. Biophys. Res. Commun.*, 148 (1987) 471-477.
- 1314 Vos, K., Laane, C., Weijers, S.R., van Hoek, A., Veeger, C. and Visser, A.J.W.G.: Time-resolved fluorescence and circular dichroism of porphyrin cytochrome c and Zn-porphyrin cytochrome c incorporated in reversed micelles. *Eur. J. Biochem.*, 169 (1987) 259-268.

For additional information see:

C.A., 107 (1987) 75718b, 132130e, 150285p, 152311t, 152316y, 152317z, 171983e, 171984f, 171985g, 173759k, 173764h, 194395d, 196018g, 213994s, 215808h, 215817k, 215824k, 215833n.

See also 1271, 1364, 1384.

19f. *Structural and muscle proteins*

- 1315 Brazel, D., Oberbäumer, I., Dieringer, H., Babel, W., Glanville, R.W., Deutzmann, R. and Kühn, K.: Completion of the amino acid sequence of the $\alpha 1$ chain of human basement membrane collagen (type IV) reveals 21 non-triplet interruptions located within the collagenous domain. *Eur. J. Biochem.*, 168 (1987) 529-536.
- 1316 Chaufer, B., Loret, C., Seville, B., Candau, S., Hanselin, M., Blain, Y. and Le Hir, A.: (Characterization of modified gelatin by size-exclusion chromatography. Comparison with dextrans and proteins). *Spectra 2000*, 122 (1987) 61; *C.A.*, 107 (1987) 223139f.
- 1317 D'Haese, J., Rutschmann, M., Dahlmann, B. and Hinssen, H.: Activity of a gelsolin-like actin modulator in rat skeletal muscle under proteins catabolic conditions. *Biochem. J.*, 248 (1987) 397-402.
- 1318 Shapiro, S.Z.: Elimination of the detection of an artefactual 65 kDa keratin band from immunoblots. *J. Immunol. Methods*, 102 (1987) 143-146; *C.A.*, 107 (1987) 171877y.
- 1319 Siebold, B., Qian, R.-w., Glanville, R.W., Hofmann, H., Deutzmann, R. and Kühn, K.: Construction of a model for the aggregation and cross-linking region (7S domain) of type IV collagen based upon an evaluation of the primary structure of the $\alpha 1$ and $\alpha 2$ chains in this region. *Eur. J. Biochem.*, 168 (1987) 569-575.
- 1320 Strohsacker, M.W., Minnich, M.D., Clark, M.A., Shorr, R.G.L. and Crooke, S.T.: Selective purification of cardiac myosin by a high-performance salicylate affinity column. *J. Chromatogr.*, 435 (1988) 185-192.
- 1321 Watanabe, T.K., Kuczmarski, E.R. and Reddy, J.K.: Myosin from pancreatic acinar carcinoma cells. Isolation, characterization and demonstration of heavy- and light-chain phosphorylation. *Biochem. J.*, 247 (1987) 513-518.
- 1322 Wu, J.-J., Eyre, D.R. and Slayter, H.S.: Type VI collagen of the intervertebral disc. Biochemical and electron-microscopic characterization of the native protein. *Biochem. J.*, 248 (1987) 373-381.

19g. *Protamines, histones and other chromosomal proteins*

- 1323 Higurashi, M., Adachi, H. and Ohba, Y.: Synthesis and degradation of H1 histone subtypes in mouse lymphoma L5178Y cells. *J. Biol. Chem.*, 262 (1987) 13075-13080.
- 1324 Palvimo, J., Mahonen, A. and Mäenpää, P.H.: Phosphorylation of high-mobility group chromatin proteins by protein kinase C from rat brain. *Biochim. Biophys. Acta*, 931 (1987) 376-383.

19h. *Chromoproteins and metalloproteins*

- 1325 Alter, B.P. and Stump, D.D.: Prenatal diagnosis of hemoglobinopathies: A potential application of HPLC. *Hemoglobin*, 11 (1987) 341-352; *C.A.*, 107 (1987) 130214m.
- 1326 Baudin-Chich, V., Rochette, J. and Wajcman, H.: Strategy for structural characterization of hemoglobin variants. *Acta Haematol.*, 78 (1987) 127-129; *C.A.*, 107 (1987) 171769q.
- 1327 Berger, G., Wollenweber, A.M., Kleo, J., Andrianambintsoa, S. and Mantele, W.G.: A rapid preparative method for purification of bacteriochlorophyll A and B. *J. Liq. Chromatogr.*, 10 (1987) 1519-1531.
- 1328 Chersi, A., Trinca, M.L. and Muratti, E.: Isolation of haem-bearing peptides of haem proteins by the use of adsorption chromatography. *J. Chromatogr.*, 410 (1987) 463-465.
- 1329 Covell, A.M. and Cock, J.D.: Stability of radioiodinated ferritin. Stability of radioiodinated ferritin. *Clin. Chim. Acta*, 168 (1987) 261-272.
- 1330 De Haën, C.: Molecular weight standards of calibration and sodium dodecyl sulfate-polyacrylamide gel electrophoresis: ferritin and apoferritin. *Anal. Biochem.*, 166 (1987) 235-245.
- 1331 Dean, J.R., Munro, S., Ebdon, L., Crews, H.M. and Massey, R.C.: Studies of metalloprotein species by directly coupled HPLC-inductively coupled plasma mass spectrometry. *J. Anal. At. Spectrom.*, 2 (1987) 607-610; *C.A.*, 107 (1987) 194381w.
- 1332 Funae, Y. and Imaoka, S.: Purification and characterization of liver microsomal cytochrome P-450 from untreated male rats. *Biochim. Biophys. Acta*, 926 (1987) 349-358.

- 1333 Gelfi, C., Righetti, P.G., Rovida, E. and Samaja, M.: Purification of human hemoglobin valence intermediates by preparative immobilized pH gradients. *J. Biochem. Biophys. Methods*, 14 (1987) 139-147; *C.A.*, 107 (1987) 150733h.
- 1334 Hinsch, W.: (Reference values for the affinity-chromatographic determination of glycosylated hemoglobins). *AerztL. Lab.*, 33 (1987) 173-176; *C.A.*, 107 (1987) 150536w.
- 1335 Huisman, T.H.J.: HPLC as a method to identify hemoglobin abnormalities. *Acta Haematol.*, 78 (1987) 123-126; *C.A.*, 107 (1987) 171673d - a review with 14 refs.
- 1336 Imaoka, S., Terano, Y. and Funae, Y.: Purification and characterization of two constitutive cytochrome P-450 (F-1 and F-2) from adult female rats: identification of P-450F-1 as the phenobarbital-inducible cytochrome P-450 in male rat liver. *Biochim. Biophys. Acta*, 916 (1987) 358-367.
- 1337 Kleinschmidt, T., Koop, B.F. and Braunitzer, G.: The primary structure of the Pallid Bat (*Antrozous pallidus*, Chiroptera) hemoglobin. *Biol. Chem. Hoppe-Seyler*, 368 (1987) 1197-1202.
- 1338 Kobayashi, S. and Sayato-Suzuki, J.: Zinc and copper accumulation and iso-metallothionein induction in mouse ascites sarcoma S180A cells. *Biochem. J.*, 249 (1988) 69-75.
- 1339 Oremek, C., Seiffert, U.B. and Schmid, G.: Determination of glycosylated hemoglobin by affinity chromatography. *Clin. Chim. Acta*, 163 (1987) 81-86.
- 1340 Schweiger, G., Dutschko, R. and Buckel, W.: Purification of 2-hydroxyglutaryl-CoA dehydratase from *Acidaminococcus fermentans*. An iron-sulfur protein. *Eur. J. Biochem.*, 169 (1987) 441-448.
- 1341 Turanek, J., Kovar, J. and Zabrsova, H.: Rapid purification of the isoforms of rat liver Ca,Zn-metallothionein by high-performance liquid chromatography. *J. Chromatogr.*, 410 (1987) 470-473.
- 1342 Vandenesch, F., Baklouti, F., Francina, A., Vianey-Liaud, C., Bertrand, A., Le Devehat, C. and Delaunay, J.: Hemoglobin J-Baltimore (B16(A13)Gly+Asp): interference with the assay of HbA_{1c}. *Clin. Chim. Acta*, 168 (1987) 121-128.

See also 1146, 1291, 1297, 1798.

19i. *Proteins of glands, gland products and various zymogens (including milk proteins)*

- 1343 Adams, M.E., Carney, R.L., Enderlin, F.E., Fu, E.T., Jarema, M.A., Li, J.P., Miller, C.A., Schooley, D.A., Shapiro, M.J. and Venema, V.J.: Structures and biological activities of three synaptic antagonists from orb weaver spider venom. *Biochem. Biophys. Res. Commun.*, 148 (1987) 678-683.
- 1344 Davies, D.T. and Law, A.J.R.: Quantitative fractionation of casein mixtures by fast protein liquid chromatography. *J. Dairy Res.*, 54 (1987) 369-376; *C.A.*, 107 (1987) 132714y.
- 1345 Govers-Riemslog, J.W.P., Knapen, M.J.H., Tans, G., Zwaal, R.F.A. and Rosing, J.: Structural and functional characterization of a prothrombin activator from the venom of *Bothrops Neuwiedi*. *Biochim. Biophys. Acta*, 916 (1987) 388-401.
- 1346 Haque, Z., Kristjansson, M.M. and Kinsella, J.E.: Interaction between κ -casein and β -lactoglobulin: possible mechanism. *J. Agric. Food Chem.*, 35 (1987) 644-649.
- 1347 Leclercq, Y., Sawatzki, G., Wieruszkeski, J.-M., Montreuil, J. and Spik, G.: Primary structure of the glycans from mouse serum and milk transferrins. *Biochem. J.*, 247 (1987) 571-578.
- 1348 Nendza, R., Digweed, M., Meyer, H.E., Erdmann, V.A. and Mayr, G.W.: 5S-rRNA-Containing ribonucleoproteins from rabbit muscle and liver. Complex and partial primary structures. *Eur. J. Biochem.*, 169 (1987) 85-95.
- 1349 Otting, G., Marchot, P., Bougis, P.E., Rochat, H. and Wüthrich, K.: Monitoring the purification by high-performance liquid chromatography of cardiotoxins from *Naja mossaambica mossaambica* using phase-sensitive two-dimensional nuclear magnetic resonance. *Eur. J. Biochem.*, 168 (1987) 603-607.
- 1350 Tosteson, M.T., Levy, J.J., Caporale, L.H., Rosenblatt, M. and Tosteson, D.C.: Solid-phase synthesis of melittin: purification and functional characterization. *Biochemistry*, 26 (1987) 6627-6631.

- 1351 Van Wormhoudt, A. and Fouchereau-Peron, M.: Partial characterization and amino acid composition of a high molecular weight peptide with salmon calcitonin immunoreactivity in a crustacean: *Nephrops norvegicus*. *Biochem. Biophys. Res. Commun.*, 148 (1987) 463-470.

For additional information see:

C.A., 107 (1987) 95417f, 130190a, 132713x, 215818m.

19j. *Proteins of brain, cerebrospinal fluid and eye*

- 1352 Frappier, T., Regnouf, F. and Pradel, L.A.: Binding of brain spectrin to the 70-kDa neurofilament subunit protein. *Eur. J. Biochem.*, 169 (1987) 651-657.
- 1353 Lang, J. and Costa, T.: Antisera against the 3-17 sequence of rat G α i recognize only a 40 kDa G-protein in brain. *Biochem. Biophys. Res. Commun.*, 148 (1987) 838-848.
- 1354 Vaerøy, H., Nyberg, F., Franzen, H. and Terenius, L.: Characterization of a substance P-Gly¹² amidating enzyme in human cerebrospinal fluid. *Biochem. Biophys. Res. Commun.*, 148 (1987) 24-30.
- 1355 Weitzer, G. and Wiche, G.: Plectin from bovine lenses. Chemical properties, structural analysis and initial identification of interaction partners. *Eur. J. Biochem.*, 169 (1987) 41-52.

For additional information see:

C.A., 107 (1987) 132141j.

See also 1357.

19k. *Proteins of neoplastic tissue and transformed cells*

- 1356 Hnatowich, D.J., Gionet, M., Rusckowski, M., Siebecker, D.A., Roche, J., Shealy, D., Mattis, J.A., Wilson, J., McGann, J., Hunter, R.E., Griffin, T. and Doherty, P.W.: Pharmacokinetics of ¹¹¹In-labeled OC-125 antibody in cancer patients compared with the 19-9 antibody. *Cancer Res.*, 47 (1987) 6111-6117.

For additional information see:

C.A., 107 (1987) 196026h.

19l. *Specific binding and receptor proteins*

- 1357 Abduragimov, A.R., Itkes, A.V., Kochetkov, S.N., Tunitskaya, V.L., Tyurkin, V.V. and Severin, E.S.: (A heat-stable protein inhibitor of protein kinase from pig brain: isolation and interaction with the enzyme). *Biokhimiya (Moscow)*, 52 (1987) 1798-1807.
- 1358 Auricchio, F., Migliaccio, A., Castoria, G., Rotondi, A., Di Domenico, M., Pagano, M. and Nola, E.: Phosphorylation on tyrosine of oestradiol-17 β receptor in uterus and interaction of oestradiol-17 β and glucocorticoid receptors with antiphosphotyrosine antibodies. *J. Steroid Biochem.*, 27 (1987) 245-253.
- 1359 Chang, C. and Liao, S.: Topographic recognition of cyclic hydrocarbons and related compounds by receptors for androgens, estrogens, and glucocorticoids. *J. Steroid Biochem.*, 27 (1987) 123-131.
- 1360 Denner, L.A., Bingman, W.E., III, Greene, G.L. and Weigel, N.L.: Phosphorylation of the chicken progesterone receptor. *J. Steroid Biochem.*, 27 (1987) 235-243.
- 1361 Egawa, T., Yamamoto, H., Kishimoto, S.: Vitamin-D-binding protein synthesized by a human hepatocellular carcinoma cell line. *Clin. Chim. Acta*, 167 (1987) 273-284.
- 1362 Fukai, F., Kase, T., Shidotani, T., Nagai, T. and Katayama, T.: A novel role of fatty acid-binding protein as a vehicle of retinoids. *Biochem. Biophys. Res. Commun.*, 147 (1987) 899-903.
- 1363 Garcia, T., Buchou, T., Jung-Testas, I., Renoir, J.-M. and Baulieu, E.E.: Chick oviduct progesterone receptor phosphorylation: characterization of a copurified kinase and phosphorylation in primary cultures. *J. Steroid Biochem.*, 27 (1987) 227-234.

- 1364 Groener, J.E.M. and Kostner, G.M.: Lipid transfer protein-catalyzed exchange of cholesteryl ester between high density lipoproteins and apoB-containing lipoproteins. *J. Lipid Res.*, 28 (1987) 1053-1056.
- 1365 Guerini, D., Krebs, J. and Carafoli, E.: Stimulation of the erythrocyte Ca^{2+} -ATPase and of bovine brain cyclic nucleotide phosphodiesterase by chemically modified calmodulin. *Eur. J. Biochem.*, 170 (1987) 35-42.
- 1366 Haggood, J.P. and von Holt, C.: Steroid-affinity purification of the rat liver glucocorticoid hormone receptor complex. *J. Steroid Biochem.*, 28 (1987) 769-777.
- 1367 Hasunuma, K., Furukawa, K., Tomita, K., Mukai, C. and Nakamura, T.: GTP-binding proteins in etiolated epicotyls of *Pisum sativum* (Alaska) seedlings. *Biochem. Biophys. Res. Commun.*, 148 (1987) 133-139.
- 1368 Heubner, A., Rde, S., Juchem, M., Grill, H.-J. and Pollow, K.: An enzyme-linked immunosorbent assay (ELISA) for human sex hormone-binding globulin. *J. Steroid Biochem.*, 28 (1987) 647-652.
- 1369 Hutchens, T.W., Li, C.M. and Besch, P.K.: Proteins associated with untransformed estrogen *in vitro*. Perturbation of hydrophobic interactions induces alterations in quaternary structure and exposure of the DNA-binding site. *Biochemistry*, 26 (1987) 5608-5616.
- 1370 Hyder, S.M. and Wittliff, J.L.: High-performance hydrophobic interaction chromatography of a labile regulatory protein: the estrogen receptor. *BioChromatography*, 2 (1987) 121-130; *C.A.*, 107 (1987) 171763h.
- 1371 Kwok, Y.C. and Yip, C.C.: Tyrosine phosphorylation of two cytosolic proteins of 50 kDa and 35 kDa in rat liver by insulin-receptor kinase *in vitro*. *Biochem. J.*, 248 (1987) 27-33.
- 1372 Lewis, R.E. and Czech, M.P.: Phospholipid environment alters hormone-sensitivity of the purified insulin receptor kinase. *Biochem. J.*, 248 (1987) 829-836.
- 1373 Logeat, F., Le Cunff, M., Rauch, M., Brailly, S. and Milgrom, E.: Characterization of a casein kinase which interacts with the rabbit progesterone receptor. Differences with the *in vivo* hormone-dependent phosphorylation. *Eur. J. Biochem.*, 170 (1987) 51-57.
- 1374 Manjunath, P., Sairam, M.R. and Uma, J.: Purification of four gelatin-binding proteins from bovine seminal plasma by affinity chromatography. *BioSci. Rep.*, 7 (1987) 231-238; *C.A.*, 107 (1987) 150556c.
- 1375 Postoenko, V.A., Parkhomenko, Yu.M., Vovk, A.I., Khalmuradov, A.G. and Donchenko, G.V.: (Purification and properties of thiamine-binding protein from rat brain synaptosomes). *Biokhimiya (Moscow)*, 52 (1987) 1792-1797.
- 1376 Prozialeck, W.C., Wallace, T.L. and Weiss, B.: Differential inhibition of calmodulin-sensitive phosphodiesterase and Ca^{++} -adenosine triphosphatase by chlorpromazine-linked calmodulin. *J. Pharmacol. Exp. Ther.*, 243 (1987) 171-179.
- 1377 Rehmus, E.H., Howard, K.J., Janiga, K.E. and Distelhorst, C.W.: Immunochemical comparison of mutant glucocorticoid receptors and wild type receptor fragments produced by neutrophil elastase and chymotrypsin. *J. Steroid Biochem.*, 28 (1987) 167-177.
- 1378 Stephenson, F.A.: Understanding the GABA_A receptor: a chemically gated ion channel. *Biochem. J.*, 249 (1988) 21-32 - a review.
- 1379 Van Schravendijk, M.R., Wilson, R.J.M. and Newbold, C.I.: Possible pitfalls in the identification of glycoprotein-binding proteins of *Plasmodium falciparum*. *J. Exp. Med.*, 166 (1987) 376-390; *C.A.*, 107 (1987) 150519t.

For additional information see:

C.A., 107 (1987) 130176a, 130191b, 150559f, 171734z, 194366v, 194517v.

See also 1260, 1783.

19n. Other proteins

- 1380 Hergenbahn, H.-G., Aspan, A. and Sderhll, K.: Purification and characterization of a high- M_r proteinase inhibitor of pro-phenol oxidase activation from crayfish plasma. *Biochem. J.*, 248 (1987) 223-228.
- 1381 Kato, A. and Takagi, T.: Estimation of the molecular weight distribution of heat-induced ovalbumin aggregates by the low-angle laser light scattering technique combined with high-performance gel chromatography. *J. Agric. Food Chem.*, 35 (1987) 633-637.

- 1382 McDonald, T.P., Clift, R., Cottrell, M. and Long, M.D.: Recovery of thrombopoietin during purification. *Biochem. Med. Metab. Biol.*, 37 (1987) 335-343; *C.A.*, 107 (1987) 70946q.
- 1383 Meuwissen, J.A.T.P., Kinnaert, M., Michiels, G. and Heirwegh, K.P.M.: Principles of a competitive binding assay for the study of the interactions of poorly water-soluble ligands with their soluble binding partners. Application to bilirubin with the use of Sephadex G-10 as a competitive adsorbent. *Biochem. J.*, 249 (1988) 263-270.
- 1384 Sivakoff, S.I.: HPLC analysis and purification of antibody conjugates. *BioChromatography*, 1 (1986) 42-48; *C.A.*, 107 (1987) 132129m.
- 1385 Sullivan, R. and Klagsbrun, M.: Purification and assay of intact human basic fibroblast growth factor using heparin-Sepharose chromatography. *J. Tissue Cult. Methods*, 10 (1986) 125-132; *C.A.*, 107 (1987) 168899b.

For additional information see:

C.A., 107 (1987) 127173s, 130172w, 130194e.

20. ENZYMES AND ENZYME ACTIVITY ESTIMATION

- 1386 Rossomando, E.F.: Strategy for designing and HPLC system for assay of enzyme activity. *LC-GC*, 5 (1987) 880-889; *C.A.*, 107 (1987) 193557w - a review with 9 refs.

20a. Oxidoreductases

- 1387 Bruguera, P., Lopez-Cabrera, A. and Canela, E.I.: Kinetic mechanism of chicken liver xanthine dehydrogenase. *Biochem. J.*, 249 (1988) 171-178.
- 1388 Davis, A.T.: Assay for trimethyllysine hydroxylase by high-performance liquid chromatography. *J. Chromatogr.*, 422 (1987) 253-256.
- 1389 Ehrlich, R.S. and Colman, R.F.: Characterization of an active site peptide modified by the substrate analogue 3-bromo-2-ketoglutarate on a single chain of dimeric NADP⁺-dependent isocitrate dehydrogenase. *J. Biol. Chem.*, 262 (1987) 12614-12619.
- 1390 Frydman, R.B., Rosenfeld, J., Camoretti-Mercado, B. and Peisach, J.: Porphobilinogen oxygenase. Purification and evidence of its hemoprotein structure. *J. Biol. Chem.*, 262 (1987) 15118-15126.
- 1391 Hakvoort, T.B.M., Moolenaar, K., Lankvelt, A.H.M., Sinjorgo, K.M.C., Dekker, H.L. and Muijsers, A.O.: Separation, stability and kinetics of monomeric and dimeric bovine heart cytochrome c oxidase. *Biochim. Biophys. Acta*, 894 (1987) 347-354.
- 1392 Haumont, P.-Y., Thomas, M.-A., Labeyrie, F. and Lederer, F.: Amino-acid sequence of the cytochrome-b₅-like heme-binding domain from *Hansenula anomala* flavocytochrome b₂. *Eur. J. Biochem.*, 169 (1987) 539-546.
- 1393 Hoey, M.E., Allison, N., Scott, A.J. and Fewson, C.A.: Purification and properties of L-mandelate dehydrogenase and comparison with other membrane-bound dehydrogenases from *Acinetobacter calcoaceticus*. *Biochem. J.*, 248 (1987) 871-876.
- 1394 Ishii-Ohba, H., Inano, H. and Tamaoki, B.-I.: Testicular and adrenal 3 β -hydroxy-5-ene-steroid dehydrogenase and 5-ene-4-ene isomerase. *J. Steroid Biochem.*, 27 (1987) 775-779.
- 1395 Ishikawa, T., Hunaiti, A.R., Piechot, G. and Wolf, B.: Isolation and characterization of basic superoxide dismutase consisting of M_r-25000 subunits in rat liver. *Eur. J. Biochem.*, 170 (1987) 317-323.
- 1396 Johansson, G. and Joelson, M.: Effect of polymer structure on affinity partitioning of lactate dehydrogenase in polymer-water two-phase systems. *J. Chromatogr.*, 411 (1987) 161-166.
- 1397 Lin, A.I., Ashley, G.W. and Stubbe, J.: Location of the redox-active thiols of ribonucleotide reductase: sequence similarity between the *Escherichia coli* and *Lactobacillus leichmannii* enzymes. *Biochemistry*, 26 (1987) 6905-6909.
- 1398 Matsushita, K., Shinagawa, E., Adachi, O. and Ameyama, M.: Purification, characterization and reconstitution of cytochrome o-type oxidase from *Gluconobacter suboxydans*. *Biochim. Biophys. Acta*, 894 (1987) 304-312.

- 1399 Muth, E., Mörschel, E. and Klein, A.: Purification and characterization of an 8-hydroxy-5-deazaflavin-reducing hydrogenase from the archaeobacterium *Methanococcus voltae*. *Eur. J. Biochem.*, 169 (1987) 571-577.
- 1400 Nagai, K., Miyamori, I., Takeda, R., Suhara, K. and Katagiri, M.: Effect of ketoconazole, etomidate and other inhibitors of steroidogenesis on cytochrome P-450c11-catalyzed reactions. *J. Steroid Biochem.*, 28 (1987) 333-336.
- 1401 Osawa, Y., Higashiyama, T., Fronckowiak, M., Noshida, N. and Yarborough, C.: Aromatase. *J. Steroid Biochem.*, 27 (1987) 781-789.
- 1402 Samanta, T.B. and Ghosh, D.K.: Characterization of progesterone 11 α -hydroxylase of *Aspergillus ochraceus* TS: A cytochrome P-450 linked monooxygenase. *J. Steroid Biochem.*, 28 (1987) 327-332.
- 1403 Sejlitz, T. and Neujahr, H.Y.: Phenol hydroxylase from yeast. A model for phenol binding and an improved purification procedure. *Eur. J. Biochem.*, 170 (1987) 343-349.
- 1404 Taroni, F., Gellera, C. and Di Donato, S.: Evidence for two distinct mitochondrial malic enzymes in human skeletal muscle: purification and properties of the NAD(P)⁺-dependent enzyme: *Biochim. Biophys. Acta*, 916 (1987) 446-454.
- 1405 Van der Zwaan, J.W., Albracht, S.P.J., Fontijn, R.D. and Mul, P.: On the anomalous temperature behaviour of the EPR signal of monovalent nickel in hydrogenase. *Eur. J. Biochem.*, 169 (1987) 377-384.
- 1406 Wakao, H., Wakagi, T. and Oshima, T.: Purification and properties of NADH dehydrogenase from a thermoacidophilic archaeobacterium, *Sulfolobus acidocaldarius*. *J. Biochem. (Tokyo)*, 102 (1987) 255-262.

For additional information see:

C.A., 107 (1987) 92451b, 92883u, 129682f, 171056e, 193670c, 193739g.

20b. *Transferases (excluding E.C. 2.7.-.)*

- 1407 Burtscher, H.J., Klocker, H., Schneider, R., Auer, B., Hirsch-Kauffmann, M. and Schweiger, M.: ADP-Ribosyltransferase from *Helix pomatia*. Purification and characterization. *Biochem. J.*, 248 (1987) 859-864.
- 1408 Cabrero, C., Puerta, J. and Alemany, S.: Purification and comparison of two forms of S-adenosyl-L-methionine synthetase from rat liver. *Eur. J. Biochem.*, 170 (1987) 299-304.
- 1409 Franklin, R.B., Kukoyi, B.I., Akuffo, V. and Costello, L.C.: Testosterone stimulation of mitochondrial aspartate aminotransferase levels and biosynthesis in rat ventral prostate. *J. Steroid Biochem.*, 28 (1987) 247-256.
- 1410 Hegazy, M.G., Thysseril, T.J., Schlender, K.K. and Reimann, E.M.: Characterization of GSK-M, a glycogen synthase kinase from rat skeletal muscle. *Arch. Biochem. Biophys.*, 258 (1987) 470-481.
- 1411 Hyaniti, A.A. and Abukhalaf, I.K.: A rapid purification procedure for camel kidney glutathione S-transferase. *Prepar. Biochem.*, 17 (1987) 239-259.
- 1412 Kane, R.E. and Chen, L.J.: Influence of gonadal hormones upon rat hepatic acetaminophen sulfotransferases. *Drug Metab. Disp.*, 15 (1987) 725-728.
- 1413 Oda, T., Miyajima, H., Suzuki, Y. and Ichiyama, A.: Nucleotide sequence of the cDNA encoding the precursor for mitochondrial serine: pyruvate aminotransferase of rat liver. *Eur. J. Biochem.*, 168 (1987) 537-542.
- 1414 Otterness, D.M. and Weinshilboum, R.M.: Mouse thiopurine methyltransferase pharmacogenetics: biochemical studies and recombinant inbred strains. *J. Pharmacol. Exp. Ther.*, 243 (1987) 180-186.
- 1415 Padgett, S.R., Huynh, Q.K., Borgmeyer, Shah, D.M., Brand, L.A., Biest Re, D., Bishop, B.F., Rogers, S.G., Fraley, R.T. and Kishore, G.M.: Bacterial expression and isolation of *Petunia hybrida* 5-enol-pyruvylshikimate-3-phosphate synthase. *Arch. Biochem. Biophys.*, 258 (1987) 564-573.
- 1416 Park, M.-S.C., Kudchodkar, B.J., Frohlich, J., Pritchard, H. and Lacko, A.G.: Study of the components of reverse cholesterol transport in lecithin-cholesterol acyltransferase deficiency. *Arch. Biochem. Biophys.*, 258 (1987) 545-554.
- 1417 Pitcher, J., Smuthe, C., Campbell, D.G. and Cohen, P.: Identification of the 38-kDa subunit of rabbit skeletal muscle glycogen synthase as glycogenin. *Eur. J. Biochem.*, 169 (1987) 497-502.
- 1418 Saido, T.C., Toyoshima, S. and Osawa, T.: Protein-O-carboxymethyltransferase from cytosol and membranes of chicken erythrocytes. *J. Biochem. (Tokyo)*, 102 (1987) 319-326.

- 1419 Scott, T.R. and Kirsch, R.E.: The isolation of a fetal liver glutathione S-transferase isoenzyme with high glutathione peroxidase activity. *Biochim. Biophys. Acta*, 926 (1987) 264-269.
- 1420 Suda, M., Watanabe, T., Kobayashi, M. and Matsuda, K.: Two types of phosphorylase from etiolated soybean cotyledons. *J. Biochem. (Tokyo)*, 102 (1987) 471-479.
- 1421 Tateoka, N., Tsuchida, S., Soma, Y. and Sato, K.: Purification and characterization of glutathione S-transferases in human kidney. *Clin. Chim. Acta*, 166 (1987) 207-218.
- 1422 Tomita, Y. and Motokawa, Y.: Characterization and partial purification of a novel mannosylphosphodolichol phosphodiesterase from chicken liver microsomes. *Eur. J. Biochem.*, 170 (1987) 363-368.
- 1423 Tsuchida, S., Izumi, T., Shimizu, T., Ishikawa, T., Hatayama, I., Satoh, K. and Sato, K.: Purification of a new acidic glutathione S-transferase, GST-Yn₁Yn₁, with a high leukotriene-C₄ synthase activity from rat brain. *Eur. J. Biochem.*, 170 (1987) 159-164.

For additional information see:
C.A., 107 (1987) 213952b.

20c. *Transferases transferring phosphorus containing groups (E.C. 2.7.-.-)*

- 1424 Azhar, S., Butte, J. and Reaven, E.: Calcium-activated, phospholipid-dependent protein kinases from rat liver: subcellular distribution, purification, and characterization of multiple forms. *Biochemistry*, 26 (1987) 7047-7057.
- 1425 Bartelt, D.C., Moroney, S. and Wolff, D.J.: Purification, characterization and substrate specificity of calmodulin-dependent myosin light-chain kinase from bovine brain. *Biochem. J.*, 247 (1987) 747-756.
- 1426 Bubis, J. and Taylor, S.S.: Limited proteolysis alters the photoaffinity labeling of adenosine 3',5'-monophosphate dependent protein kinase II with 8-azidoadenosine 3',5'-monophosphate. *Biochemistry*, 26 (1987) 5997-6004.
- 1427 Bubis, J., Vedvick, T.S. and Taylor, S.S.: Antiparallel alignment of the two protomers of the regulatory subunit dimer of cAMP-dependent protein kinase I. *J. Biol. Chem.*, 262 (1987) 14961-14966.
- 1428 Buss, W.C., Stalter, K., Stepanek, J. and Piatt, M.K.: Studies on the mechanism of glucocorticoid hormone induced alterations in rat thymic transcription - II. Partial purification and characterization of RNA polymerases II from hydrocortisone and control vehicle treated animals. *J. Steroid Biochem.*, 28 (1987) 129-138.
- 1429 Dobrowolska, G., Meggio, F. and Pinna, L.A.: Characterization of multiple forms of maize seedling protein kinases reminiscent of animal casein kinases S (type 1) and TS (type 2). *Biochim. Biophys. Acta*, 931 (1987) 188-195.
- 1430 Klucis, E. and Polya, G.M.: Calcium-independent activation of two plant leaf calcium-regulated protein kinases by unsaturated fatty acids. *Biochem. Biophys. Res. Commun.*, 147 (1987) 1041-1047.
- 1431 Kwok, F., Scholz, G. and Churchich, J.E.: Brain pyridoxal kinase dissociation of the dimeric structure and catalytic activity of the monomeric species. *Eur. J. Biochem.*, 168 (1987) 577-583.
- 1432 Little, L., Alcouloumre, M., Drotar, A.M., Herman, S., Robertson, R., Yeh, R.Y. and Miller, A.L.: Properties of N-acetylglucosamine 1-phosphotransferase from human lymphoblasts. *Biochem. J.*, 248 (1987) 151-159.
- 1433 Stephens, L.R., Hawkins, P.T., Morris, A.J. and Downes, P.C.: L-myo-Inositol 1,4,5,6-tetrakisphosphate (3-hydroxy)kinase. *Biochem. J.*, 249 (1988) 283-292.
- 1434 Wol'fson, A.D., Motorin, Yu.A., Orlovsky, A.F. and Gladilin, K.L.: (Isolation, polypeptide content and properties of the aminoacyl-tRNA-synthetase complex from rabbit liver). *Biokhimiya (Moscow)*, 52 (1987) 1847-1854.

See also 987.

20d. *Hydrolases, acting on ester bonds (E.C. 3.1.-.-)*

- 1435 Banno, Y. and Nozawa, Y.: Characterization of partially purified phospholipase C from human platelet membranes. *Biochem. J.*, 248 (1987) 95-101.

- 1436 Gibson, G.J., Saccone, G.T.P., Brooks, D.A., Clements, P.R. and Hopwood, J.J.: Human N-acetylgalactosamine-4-sulphate sulphatase. Purification, monoclonal antibody production and native and subunit M_r values. *Biochem. J.*, 248 (1987) 755-764.
- 1437 Inborn, R.C. and Majerus, P.W.: Inositol polyphosphate 1-phosphatase from calf brain. Purification and inhibition by Li^+ , Ca^{2+} , and Mn^{2+} . *J. Biol. Chem.*, 262 (1987) 15946-15952.
- 1438 Kramer, R.M., Checani, G.C. and Deykin, D.: Stimulation of Ca^{2+} -activated human platelet phospholipase A_2 by diacylglycerol. *Biochem. J.*, 248 (1987) 779-783.
- 1439 Lockridge, O., Adkins, S. and La Du, B.N.: Location of disulfide bonds within the sequence of human serum cholinesterase. *J. Biol. Chem.*, 262 (1987) 12945-12952.
- 1440 Maouri, A. and Georgatsos, J.G.: Specificity and other properties of three ribonucleases of *Tetrahymena pyriformis*. *Eur. J. Biochem.*, 168 (1987) 523-528.
- 1441 Robbi, M. and Beaufay, H.: Biosynthesis of rat liver pI-6.1 esterase, a carboxylesterase of the cisternal space of the endoplasmic reticulum. *Biochem. J.*, 248 (1987) 545-550.
- 1442 Ryu, S.H., Cho, K.S., Lee, K.-Y., Suh, P.-G. and Rhee, S.G.: Purification and characterization of two immunologically distinct phosphoinositide-specific phospholipase C from bovine brain. *J. Biol. Chem.*, 262 (1987) 12511-12518.
- 1443 Schoenau, E., Haezog, K.H. and Boehles, H.J.: (Determination of isoenzymes of alkaline phosphatase in serum and tissue homogenates by HPLC). *Monatsschr. Kinderheilkd.*, 135 (1987) 632-636; *C.A.*, 107 (1987) 193626t.
- 1444 Shriner, C.L. and Brautigan, D.L.: Metal chelate affinity chromatography of zinc-inhibited Tyr(P) phosphatases. *J. Biochem. Biophys. Methods*, 14 (1987) 273-278; *C.A.*, 107 (1987) 193812a.
- 1445 Van den Bergh, C.J., Bekkers, A.C.A.P.A., De Geus, P., Verheij, H.M. and De Haas, G.H.: Secretion of biologically active porcine phospholipase A_2 *Saccharomyces cerevisiae*. Use of the prepro sequence of the α -mating factor. *Eur. J. Biochem.*, 170 (1987) 241-246.
- 1446 Villa-Moruzzi, E. and Heilmeyer, L.M.G., Jr.: Phosphorylase phosphatase from skeletal muscle membranes. *Eur. J. Biochem.*, 169 (1987) 659-667.

For additional information see:

C.A., 107 (1987) 171156n, 193562u.

20e. *Hydrolases, acting on glycosyl compounds (E.C. 3.2.-.-)*

- 1447 Dobransky, T., Polivka, L., Haas, I., Sova, O. and Petrvalsky, E.: Purification of α -amylase by industrial autofocusing. *J. Chromatogr.*, 411 (1987) 486-489.
- 1448 Galich, I.P.: (Affinity chromatography of amylases). *Biokhim. Zhivotn. Chel.*, 11 (1987) 42-52; *C.A.*, 107 (1987) 129542k - a review with 85 refs.
- 1449 Kutty, A.V.M. and Pattabiraman, T.N.: Behavior of plant amylase and trypsin inhibitors during dye-ligand chromatography on blue Sepharose and red Sepharose. *Biochem. Arch.*, 3 (1987) 231-240; *C.A.*, 107 (1987) 193861r.
- 1450 Nicot, G., Lachatre, G., Gonnet, C., Dupuy, J.-L. and Valette, J.-P.: Rapid assay of N-acetyl- β -D-glucosaminidase isoenzymes in urine by ion-exchange chromatography. *Clin. Chem. (Winston-Salem)*, 33 (1987) 1796-1800.
- 1451 Poulsen, O.M. and Petersen, L.W.: Purification of an extracellular cellulose-binding endoglucanase of *Cellulomonas* sp. ATCC 21399 by affinity chromatography on phosphoric acid-swollen cellulose. *Biotechnol. Bioeng.*, 29 (1987) 799-804; *C.A.*, 107 (1987) 73161r.
- 1452 Suzuki, Y. and Shinji, M.: An improved method of the purification of *Bacillus stearothermophilus* exo- α -1,4-glucosidase active on soluble starch by the use of an immunosorbent. *Starch/Staerke*, 39 (1987) 271-273; *C.A.*, (1987) 150000s.

For additional information see:

C.A., 107 (1987) 193740a.

See also 1019, 1023, 1422, 1455.

20f. Other hydrolases

- 1453 Alter, S.C., Metcalfe, D.D., Bradford, T.R. and Schwartz, L.B.: Regulation of human mast cell tryptase. Effects of enzyme concentration, ionic strength and the structure and negative charge density of polysaccharides. *Biochem. J.*, 248 (1987) 821-827.
- 1454 Bradbury, A.F. and Smyth, D.G.: Enzyme-catalysed peptide amidation. Isolation of a stable intermediate formed by reaction of the amidating enzyme with an imino acid. *Eur. J. Biochem.*, 169 (1987) 579-584.
- 1455 Brassart, D., Baussant, T., Wieruszeski, J.-M., Strecker, G., Montreuil, J. and Michalski, J.-C.: Catabolism of N-glycosylprotein glycans: evidence for a degradation pathway of sialylglyco-asparagines resulting from the combined action of the lysosomal aspartylglucosaminidase and endo-N-acetyl- β -D-glucosaminidase. A 400-MHz $^1\text{H-NMR}$ study. *Eur. J. Biochem.*, 169 (1987) 131-136.
- 1456 Dufour, E., Obléd, A., Valin, C., Bechet, D., Ribadeau-Dumas, B. and Huet, J.C.: Purification and amino acid sequence of chicken liver cathepsin L. *Biochemistry*, 26 (1987) 5689-5695.
- 1457 Hayakawa, K. and Oizumi, J.: Determination of lipoamidase activity by liquid chromatography with fluorimetric detection. *J. Chromatogr.*, 423 (1987) 304-307.
- 1458 Jaenicke, L., Kuhne, W., Spessert, R., Wahle, U. and Waffenschmidt, S.: Cell-wall lytic enzymes (autolysins) of *Chlamydomonas reinhardtii* are (hydroxy) proline-specific proteases. *Eur. J. Biochem.*, 170 (1987) 485-491.
- 1459 Moody, D.E. and Hammock, B.D.: Purification of microsomal epoxide hydrolase from liver of rhesus monkey: partial separation of *cis*- and *trans*-stilbene oxide hydrolase. *Arch. Biochem. Biophys.*, 258 (1987) 156-166.
- 1460 Muller, D.: Pseudo affinity chromatography of thrombin on polystyrene substituted by arginine derivatives. *Dev. Hematol. Immunol.*, 14 (Blood Compat. Mater. Their Test.) (1986) 327-336; *C.A.*, 107 (1987) 149838q.
- 1461 Namba, T. and Mukohata, Y.: A membrane-bound ATPase from *Halobacterium halobium*: purification and characterization. *J. Biochem. (Tokyo)*, 102 (1987) 591-598.
- 1462 Nyberg, F., Lyrenaes, S. and Terenius, L.: Assay and biochemical characterization of a dynorphin converting enzyme in human cerebrospinal fluid. *NIDA Res. Monogr.*, 75 (Prog. Opioid Res.) (1986) 251-254; *C.A.*, 107 (1987) 149814d.
- 1463 Pietta, P., Mauri, P. and Pace, M.: High-performance liquid chromatographic assay of chymotrypsin and chymotrypsin-like enzyme activity. *J. Chromatogr.*, 411 (1987) 498-501.
- 1464 Sakharov, I.Y., Danilov, S.M. and Sukhova, N.V.: Isolation of human liver angiotensin-converting enzyme by chromatofocusing. *Anal. Biochem.*, 166 (1987) 14-17.
- 1465 Schäfer, M., Behle, G., Varsanyi, M. and Heilmeyer, L.M.G., Jr.: Ca^{2+} regulation of 1-(3-sn-phosphatidyl)-1D-myo-inositol 4-phosphate formation and hydrolysis on sarcoplasmic-reticular Ca^{2+} -transport ATPase. A new principle of phospholipid turnover regulation. *Biochem. J.*, 247 (1987) 579-587.
- 1466 Subramanian, S.: Separation of chymosin and pepsin in calf rennet by dye-ligand affinity chromatography. *Prepar. Biochem.*, 17 (1987) 297-312.
- 1467 Tamaki, N., Mizutani, N., Kikugawa, M., Fujimoto, S. and Mizota, C.: Purification and properties of β -ureidopropionase from the rat liver. *Eur. J. Biochem.*, 169 (1987) 21-26.

For additional information see:

C.A., 107 (1987) 92450a, 194161z.

20g. Lyases

- 1468 Chiew, Y.E., O'Sullivan, W.J. and Lee, C.S.: Studies on pig liver mevalonate-5-diphosphate decarboxylase. *Biochim. Biophys. Acta*, 916 (1987) 271-278.
- 1469 Crawford, M.S. and Kolattukudy, P.E.: Pectate lyase from *Fusarium solani* f. sp. *psii*: purification, characterization, *in vitro* translation of the mRNA, and involvement in pathogenicity. *Arch. Biochem. Biophys.*, 258 (1987) 196-205.
- 1470 Dominici, P., Tancini, B., Barra, D. and Voltattorni, C.B.: Purification and characterization of rat-liver 3,4-dihydroxyphenylalanine decarboxylase. *Eur. J. Biochem.*, 169 (1987) 209-213.

- 1471 Fernley, R.T., Coghlan, J.P. and Wright, R.D.: Purification and characterization of a high- M_r carbonic anhydrase from sheep parotid gland. *Biochem. J.*, 249 (1988) 201-207.

See also 1340.

20h. Isomerases

See 1394.

20i. Ligases

- 1472 Charlier, J. and Sanchez, R.: Lysyl-tRNA synthetase from *Escherichia coli* K12. Chromatographic heterogeneity and the lysU-gene product. *Biochem. J.*, 248 (1987) 43-51.
- 1473 Fedorov, A.N. and Ovchinnikov, L.P.: Interaction of eukaryotic threonyl-tRNA synthetase with high- M_r RNAs and tRNA^{Thr}. *Eur. J. Biochem.*, 169 (1987) 185-191.
- 1474 Kulmacz, R.J.: Attachment of substrate metabolite to prostaglandin H synthase upon reaction with arachidonic acid. *Biochem. Biophys. Res. Commun.*, 148 (1987) 539-545.

20j. Complex mixtures and incompletely identified enzymes

For additional information see:
C.A., 107 (1987) 214170g.

21. PURINES, PYRIMIDINES, NUCLEIC ACIDS AND THEIR CONSTITUENTS

21a. Purines, pyrimidines, nucleosides, nucleotides

- 1475 Alexander, A.J., Kebarle, P., Fuciarelli, A.F. and Raleigh, J.A.: Characterization of radiation-induced damage to polyadenylic acid using high-performance liquid chromatography/tandem mass spectrometry. *Anal. Chem.*, 59 (1987) 2484-2491.
- 1476 Andersson, M., Christensson, P.I., Lewan, L. and Stenram, U.: Isotachophoretic and HPLC determination of nucleotides in rat liver cell nuclei isolated by non-aqueous technique. *Int. J. Biochem.*, 19 (1987) 641-647; *C.A.*, 107 (1987) 73781t.
- 1477 Au, J.L.-S., Su, M.-H. and Wientjes, M.G.: Isocratic ion-pair reversed-phase high-performance liquid chromatographic analysis of ATP and ADP. *J. Chromatogr.*, 423 (1987) 308-312.
- 1478 Baumann, U., Lehmann, U., Schweltnus, K., van Boom, J.H. and Kuhn, H.: Complexes of DNA hairpins and a single-stranded oligonucleotide detected by affinity chromatography and mung bean nuclease cleavage. *Eur. J. Biochem.*, 170 (1987) 267-272.
- 1479 Eadie, J.S., McBride, L.J., Efcavitch, J.W., Hoff, L.B. and Cathcarf, R.: High-performance liquid chromatographic analysis of oligodeoxyribonucleotide base composition. *Anal. Biochem.*, 165 (1987) 442-447.
- 1480 Galushko, S.V., Shishkina, I.P. and Pilipenko, A.T.: (Reversed-phase high-performance liquid chromatography of atypical antitumor nucleosides). *Zh. Anal. Khim.*, 42 (1987) 1684-1690; *C.A.*, 107 (1987) 183702a.
- 1481 Germann, M.W., Pon, R.T. and van de Sande, J.H.: A general method for the purification of synthetic oligodeoxyribonucleotides containing strong secondary structure by reversed-phase high-performance liquid chromatography on RPR-1 resin. *Anal. Biochem.*, 165 (1987) 399-405.
- 1482 Inaki, Y., Nagae, S., Miyamoto, T., Sugiura, Y. and Takemoto, K.: Nucleic acid analogs for HPLC. *Polym. Mater. Sci. Eng.*, 57 (1987) 286-290; *C.A.*, 107 (1987) 194378a.
- 1483 Kobylinska-Luczko, A., Brzeszkiewicz, A., Cendrowska, I. and Butkiewicz, K.: Investigations of the stability of cytosine arabinoside in injection solutions by HPLC method. *Symp. Biol. Hung.*, 34 (1986) 391-400; *C.A.*, 107 (1987) 161517s.
- 1484 Kojima, T., Nishina, T., Kitamura, M., Kamatani, N. and Nishioka, K.: Liquid chromatography with multichannel ultraviolet detection used for studying disorders of purine metabolism. *Clin. Chem. (Winston-Salem)*, 33 (1987) 2052-2056.

- 1485 Koski, P.G.: Nucleotide separations on the BAS 200. *Curr. Sep.*, 8 (1987) 30-31; *C.A.*, 107 (1987) 171753e.
- 1486 Lennard, L.: Assay of 6-thioinosinic acid and 6-thioguanine nucleotides, active metabolites of 6-mercaptopurine, in human red blood cells. *J. Chromatogr.*, 423 (1987) 169-178.
- 1487 Leoncini, G., Buzzi, E., Maresca, M., Mazzei, M. and Balbi, A.: Alkaline extraction and reverse-phase high-performance liquid chromatography of adenine and pyridine nucleotides in human platelets. *Anal. Biochem.*, 165 (1987) 279-383.
- 1488 Makino, K., Matsumoto, T., Nakatsuji, Y., Nagahara, S. and Takeuchi, T.: Large scale purification of synthesized oligodeoxyribonucleotides by reversed-phase chromatography. *Chromatographia*, 23 (1987) 909-914.
- 1489 Makino, K., Ozaki, H., Imaishi, H., Takeuchi, T. and Fukui, T.: A RPLC C₃₀ column for the separation of oligonucleotides. *Chem. Lett.*, (1987) 1251-1254; *C.A.*, 107 (1987) 128450s.
- 1490 McLaughlin, L.W., Benseler, F., Graeser, E., Piel, N. and Scholtissek, S.: Effects of functional group changes in the EcoRI recognition site on the cleavage reaction catalyzed by the endonuclease. *Biochemistry*, 26 (1987) 7238-7245.
- 1491 Reysz, L.J., Carroll, A.G. and Jarrett, H.W.: An automated assay for adenylyl cyclase using reversed-phase high-performance liquid chromatography. *Anal. Biochem.*, 166 (1987) 107-112.
- 1492 Takenaka, S. and Takagi, M.: (Separation and analysis of DNA and oligonucleotides). *Kagaku*, 42 (1987) 636-637; *C.A.*, 107 (1987) 171651u - a review with 13 refs.
- 1493 Tamura, S., Fujioka, H., Nakano, T., Hada, T. and Higashino, K.: Serum pseudouridine as a biochemical marker in small cell lung cancer. *Cancer Res.*, 47 (1987) 6138-6141.
- 1494 Trujillo, M.A., Roux, D., Fueri, J.P., Samuel, D., Cailla, H.L. and Rickenberg, H.V.: The occurrence of 2'-5'-oligoadenylates in *Escherichia coli*. *Eur. J. Biochem.*, 169 (1987) 167-173.
- 1495 Valera, V., Fung, M., Wessler, A.N. and Richards, W.R.: Synthesis of 4R- and 4S-tritium labeled NADPH for the determination of the coenzyme stereospecificity of NADPH:photochlorophyllide oxidoreductase. *Biochem. Biophys. Res. Commun.*, 148 (1987) 515-520.
- 1496 Vockova, J. and Svoboda, V.: Fast reversed-phase high-performance liquid chromatographic determination of ¹⁴C- or ³H-labelled S-adenosylmethionine in reaction mixtures. *J. Chromatogr.*, 410 (1987) 500-503.
- 1497 Wagner, J., Hirth, Y., Claverie, N. and Danzin, C.: HPLC with fluorescence detection for the analysis of decarboxylated S-adenosylmethionine and its analogs. *Symp. Biol. Hung.*, 34 (Chromatography' 85) (1986) 85-102; *C.A.*, 107 (1987) 150523q.
- 1498 Waranis, R.P. and Sloan, K.B.: Effects of vehicles and prodrug properties and their interactions on the delivery of 6-mercaptopurine through skin: bisacyloxymethyl-6-mercaptopurine prodrugs. *J. Pharm. Sci.*, 76 (1987) 587-595.
- 1499 Zaton, A., Marinez, A. and de Gandarias, J.M.: Study of binding of benzylthiouracil to human serum albumin by gel filtration chromatography. *J. Liq. Chromatogr.*, 10 (1987) 899-908.

For additional information see:

C.A., 107 (1987) 73595k, 76224z, 130208n, 171057f, 187083x.

See also 859, 1567.

21b. Nucleic acids, RNA

- 1500 Hanukoglu, I., Gutfinger, T., Haniu, M. and Shively, J.E.: Isolation of a cDNA for adrenodoxin reductase (ferredoxin-NADP⁺ reductase). Implications for mitochondrial cytochrome P-450 systems. *Eur. J. Biochem.*, 169 (1987) 449-455.
- 1501 Jack, P.L. and Alang, Z.C.: Isolation of messenger RNA from oil palm embryos. *Pertanika*, 10 (1987) 97-102; *C.A.*, 107 (1987) 171754f.

- 1502 Kodama, M., Inoue, F., Nakayama, T. and Nagata, C.: Activation of deoxycholic acid by the lipid peroxy radical and its covalent binding to nucleic acids. *Arch. Biochem. Biophys.*, 258 (1987) 574-578.

For additional information see:
C.A., 107 (1987) 194560d.

See also 1279, 1473, 1482, 1504.

21c. Nucleic acids, DNA

- 1503 Kettleborough, C.A., Parry, M.A.J., Burton, S., Gutteridge, S., Keys, A.J. and Phillips, A.L.: The role of the N-terminus of the large subunit of ribulose-biphosphate carboxylase investigated by construction and expression of chimaeric genes. *Eur. J. Biochem.*, 170 (1987) 335-342.
- 1504 Milligan, J.R., Catz-Biro, L. and Archer, M.C.: Effect of phosphate concentration on the yield and purity of DNA separated from RNA by hydroxyapatite chromatography. *J. Chromatogr.*, 411 (1987) 481-485.
- 1505 Mueller, W.: Liquid-liquid partition chromatography. *Kontakte*, (1987) 45-56; *C.A.*, 107 (1987) 171652w - a review with 18 refs.

See also 1279, 1482, 1492, 1500.

21e. Structural studies on DNA and DNA mapping

- 1506 Jagadeeswaran, P. and Kaul, R.K.: Use of reverse-phase chromatography in the Maxam-Gilbert method of DNA sequencing. A step toward automation. *Gene Anal. Tech.*, 3 (1986) 79-85; *C.A.*, 107 (1987) 72079h.
- 1507 Westman, E., Eriksson, S., Laas, T., Pernemalm, P.-A. and Skold, S.E.: Separation of DNA restriction fragments by ion-exchange chromatography of FPLC columns Mono P and Mono Q. *Anal. Biochem.*, 166 (1987) 158-171.

21f. Complex mixtures of nucleic acids and their fragments

See 1475.

22. ALKALOIDS

- 1508 Armstrong, D.W., Spino, L.A., Han, S.M., Seeman, J.I. and Secor, H.V.: Enantiomeric resolution of racemic nicotine analogues by microcolumn liquid chromatography with β -cyclodextrin inclusion complexes. *J. Chromatogr.*, 411 (1987) 490-493.
- 1509 Attuquayefic, V.K. and Buckle, K.A.: Rapid sample preparation method for HPLC analysis of capsaicinoids in capsicum fruits and oleoresins. *J. Agric. Food Chem.*, 35 (1987) 777-779.
- 1510 Benuck, M., Lajtha, A. and Reith, M.E.A.: Pharmacokinetics of systemically administered cocaine and locomotor stimulation in mice. *J. Pharmacol. Exp. Ther.*, 243 (1987) 144-149.
- 1511 Chiou, R., Stubbs, R.J. and Bayne, W.F.: Improved method for the determination of theophylline in plasma and urine using high-performance liquid chromatography. *J. Chromatogr.*, 422 (1987) 281-287.
- 1512 Hotchkiss, S.A. and Caldwell, J.: High-performance liquid chromatographic assay for theophylline and its major metabolites in human urine. *J. Chromatogr.*, 423 (1987) 179-188.
- 1513 Kapke, G.F. and Franklin, R.B.: A direct liquid chromatography method for serum caffeine analysis. *J. Liq. Chromatogr.*, 10 (1987) 451-463.
- 1514 Kennedy, J.-S., Leduc, B.W., Scavone, J.M., Harmatz, J.S., Shader, R.I. and Greenblatt, D.J.: Pharmacokinetics of intravenous caffeine: comparison of high-performance liquid chromatographic and gas chromatographic methods. *J. Chromatogr.*, 422 (1987) 274-280.
- 1515 Kester, M.B., Saccar, C.L. and Mansmann, H.C., Jr.: A new simplified microassay for the quantitation of theophylline in serum by high-performance liquid chromatography. *J. Liq. Chromatogr.*, 10 (1987) 957-975.

- 1516 Majak, W., McDiarmid, R.E. and Benn, M.H.: Isolation and HPLC determination of methyllycaconitine in a species of low larkspur (*Delphinium nuttalianum*). *J. Agric. Food Chem.*, 35 (1987) 800-802.
- 1517 Martelli, A. and Belliaro, F.: (Effect of the extraction procedure on caffeine determination by HPLC in foods and dietetic products). *Riv. Soc. Ital. Sci. Aliment.*, 16 (1987) 121-128; *C.A.*, 107 (1987) 216259s.
- 1518 Mattammal, M.B., Lakshmi, V.M., Zenser, T.V. and Davis, B.B.: Lung prostaglandin H synthase and mixed-function oxidase metabolism of nicotine. *J. Pharmacol. Exp. Ther.*, 242 (1987) 827-832.
- 1519 Naaranlahti, T., Nordström, M., Huhtikangas, A. and Lounasmaa, M.: Determination of *Catharanthus* alkaloids by reversed-phase high-performance liquid chromatography. *J. Chromatogr.*, 410 (1987) 488-493.
- 1520 Shulgin, A.T., Jacob, P., III, Benowitz, N.L. and Lau, D.: Identification and quantitative analysis of cotinine-N-oxide in human urine. *J. Chromatogr.*, 423 (1987) 365-372.
- 1521 Somani, S.M. and Khalique, A.: Pharmacokinetics and pharmacodynamics of physostigmine in the rat after intravenous administration. *Drug Metab. Disp.*, 15 (1987) 627-633.
- 1522 Spearman, M.E., Goodwin, R.M. and Kau, D.: Disposition of the monoclonal antibody-vinca alkaloid conjugate, DS1/4-DAVLB (LY256787), in Fischer 344 rats and rhesus monkeys. *Drug Metab. Disp.*, 15 (1987) 640-647.
- 1523 Wahbi, A.A.M., Abounassif, M.A. and Gad-Kariem, El Rasheed, A.: HPLC determination of strychnine and brucine in nux vomica liquid extract. *Arch. Pharm. Chemi. Sci. Ed.*, 15 (1987) 87-90; *C.A.*, 107 (1987) 192389f.
- 1524 Zhang, T.-Y., Cai, D.-G. and Ito, Y.: Separations of flavonoids and alkaloids in medicinal herbs by high-speed counter-current chromatography. *J. Chromatogr.*, 435 (1988) 159-166.

See also 866, 1726, 1801.

23. OTHER SUBSTANCES CONTAINING HETEROCYCLIC NITROGEN

23a. Porphyrins and other pyrroles

- 1525 Correia, M.A., Decker, C., Sugiyama, K., Caldera, P., Bornheim, L., Wrighton, S.A., Rettie, A.E. and Trager, W.F.: Degradation of rat hepatic cytochrome P-450 heme by 3,5-dicarbethoxy-2,6-dimethyl-4-ethyl-1,4-dihydropyridine to irreversibly bound protein adducts. *Arch. Biochem. Biophys.*, 258 (1987) 436-451.
- 1526 Dellinger, M. and Brault, D.: Normal-phase high-performance liquid chromatography of free acid dicarboxylic porphyrins and hematoporphyrin derivative on silica. *J. Chromatogr.*, 422 (1987) 73-84.
- 1527 Pfaltz, A., Kobelt, A., Hüster, R. and Thauer, R.K.: Biosynthesis of coenzyme F430 in methanogenic bacteria. Identification of 15,17³-seco-F430-17³-acid as an intermediate. *Eur. J. Biochem.*, 170 (1987) 459-467.
- 1528 Sassa, S.: (Fractionation and determination of porphyrin). *Rinsho Kensa*, 31 (1987) 259-267; *C.A.*, 107 (1987) 150453s - a review with 11 refs.
- 1529 Sun, X.: Porphyrin analysis by reversed-phase HPLC and its biomedical applications. *Shenyang Yaowuexueyan Xuebao*, 4 (1987) 144-148; *C.A.*, 107 (1987) 171672c - a review with 16 refs.
- 1530 Takayama, M.: Reversed-phase high-performance liquid chromatography of urinary porphyrin free acids. *J. Chromatogr.*, 423 (1987) 313-318.

For additional information see:

C.A., 107 (1987) 150551x.

23b. Bile pigments

- 1531 Yoshida, H., Inagaki, T., Hirano, M. and Sugimoto, T.: Analyses of azopigments obtained from the delta fraction of bilirubin from mammalian plasma (mammalian biliprotein). *Biochem. J.*, 248 (1987) 79-84.

23c. Indole derivatives

- 1532 Cardinali, D.P., Vacas, M.I., Keller-Sarmiento, M.I., Etchegoven, G.S., Pereyra, E.N. and Chuluyan, H.E.: Neuroendocrine integrative mechanisms in mammalian pineal gland: effects of steroid and adenohipophysal hormones on melatonin synthesis *in vitro*. *J. Steroid Biochem.*, 27 (1987) 565-571.
- 1533 Frankenberger, W.T., Jr. and Poth, M.: Determination of substituted indole derivatives by ion suppression-reverse-phase high-performance liquid chromatography. *Anal. Biochem.*, 165 (1987) 300-308.
- 1534 Marasini, B., Biondi, M.L. and Agostini, A.: Liquid chromatographic determination of free serotonin in human plasma. *Riv. Clin. Lab.*, 17 (1987) 149-154; *C.A.*, 107 (1987) 211934y.
- 1535 Niles, L.P., Pickering, D.S. and Sayer, B.G.: HPLC-purified 2-[¹²⁵I]iodomelatonin labels multiple binding sites in hamster brain. *Biochem. Biophys. Res. Commun.*, 147 (1987) 949-956.
- 1536 Sharp, T., Zetterstroem, T., Series, H.G., Carlsson, A., Grahame-Smith, D.G. and Ungerstedt, U.: HPLC-EC analysis of catechols and indoles in rat brain dialyzates. *Life Sci.*, 41 (1987) 869-872; *C.A.*, 107 (1987) 70908d.
- 1537 Tan, F.: (HPLC of plant hormones). *Zhijiu Shenglixue Tongxun*, (1986) 15-23; *C.A.*, 107 (1987) 73522j - a review with 37 refs.
- 1538 Trouvin, J.H., Gardier, A., El Gemayel, G. and Jacquot, C.: Rapid and specific determination of serotonin, its precursors and metabolites in rat brain tissue by liquid chromatography with electrochemical detection. *J. Liq. Chromatogr.*, 10 (1987) 121-136.

See also 1571, 1732.

23d. Pyridine derivatives

- 1539 Shibata, K., Kawada, T. and Iwai, K.: High-performance liquid chromatographic determination of nicotinamide in rat tissue samples and blood after extraction with diethyl ether. *J. Chromatogr.*, 422 (1987) 257-262.

See also 906, 1508.

23e. Other N-heterocyclic compounds

- 1540 D'Avila, L.A., Colin, H.M. and Guiochon, G.: In-column derivatization for the separation of heavy aza-arenes and olefins by liquid chromatography. *J. Liq. Chromatogr.*, 10 (1987) 71-82.
- 1541 Feierman, D.E. and Cederbaum, A.I.: Oxidation of the alcohol dehydrogenase inhibitor pyrazole to 4-hydroxypyrazole by microsomes. Effect of cytochrome P-450 inducing agents. *Drug Metab. Disp.*, 15 (1987) 634-639.
- 1542 Frolov, E.B., Vanyukova, N.A. and Sanin, P.I.: (Selective isolation of carbazoles from crude petroleum). *Neftekhimiya*, 27 (1987) 328-333; *C.A.*, 107 (1987) 157831x.
- 1543 Gazdzik, W.R., Filipek, M. and Kmiotek, W.: Rapid and simple HPLC determination of 5,5-dimethyl-2,4-oxazolidinedione in rabbit serum. *Biomed. Chromatogr.*, 2 (1987) 137-138.
- 1544 Gouygou, J.P., Siquin, C. and Durand, P.: High pressure liquid chromatography determination of histamine in fish. *J. Food Sci.*, 52 (1987) 925-927; *C.A.*, 107 (1987) 132761m.
- 1545 Javor, J., Kiss, I. and Kovacs, A.: Thin-layer chromatography and high-pressure liquid chromatography of different acridine-glycosides. *Symp. Biol. Hung.*, 34 (1986) 263-288; *C.A.*, 107 (1987) 161765w.
- 1546 Manabe, S., Yanagisawa, H., Ishikawa, S., Kitagawa, Y., Kanai, Y. and Wada, O.: Accumulation of 2-amino-6-methyldipyrido[1,2-a:3',2'-d]imidazole and 2-amino-dipyrido[1,2-a:3',2'-d]imidazole, carcinogenic glutamic acid pyrolysis products, in plasma of patients with uremia. *Cancer Res.*, 47 (1987) 6150-6155.
- 1547 Soto-Otero, R., Mendez-Alvarez, E., Sierra-Paredes, G., Galan-Valiente, J., Aguilar-Veiga, E. and Sierra-Marcuno, G.: High-performance liquid chromatographic procedure for quantitative determination of pentylenetetrazol in serum and discrete areas of rat brain. *Anal. Biochem.*, 165 (1987) 331-336.

See also 917, 941, 1610.

24. ORGANIC SULPHUR COMPOUNDS

- 1548 Andersson, J.T.: Retention properties of a palladium-chloride/silica sorbent for the liquid chromatographic separation of polycyclic aromatic sulfur heterocycles. *Anal. Chem.*, 59 (1987) 2207-2209.
- 1549 Baeyens, W., van der Weken, G. and De Moerloose, P.: Effects of reducing agents on the determination of thiolic compounds in the presence of their disulfides using bimane pre-column derivatization. *Chromatographia*, 23 (1987) 717-721.
- 1550 Buchner, R.: Comparison of procedures for optimum extraction, purification and analysis of desulfo indolyl glucosinolates. *World Crops: Prod., Util., Descr.*, 13 (Glucosinolates, Rapeseeds) (1987) 76-89; *C.A.*, 107 (1987) 214445a.
- 1551 Claycamp, H.G. and Ludwig, B.: Simplified monitoring of reduced and oxidized dithiothreitol using high-performance liquid chromatography. *J. Chromatogr.*, 422 (1987) 239-245.
- 1552 Kim, D.-H. and Kobashi, K.: Immobilized arylsulfotransferase. *J. Biochem. (Tokyo)*, 102 (1987) 487-491.
- 1553 Matissek, R.: High-performance liquid chromatography determination of methyl isothiazolones in cosmetics. *Seifen, Oele, Fette, Wachse*, 113 (1987) 474-478; *C.A.*, 107 (1987) 204888h.
- 1554 Moeckel, H.J., Melzer, H., Hoefler, F. and Fojtik, A.T.: Comparison of the RPLC retention of organic sulphur, selenium and tellurium compounds. *Symp. Biol. Hung.*, 34 (1986) 525-539; *C.A.*, 107 (1987) 228103z.
- 1555 Quinsac, A. and Ribaillier, D.: Optimization of glucosinolate desulfation before HPLC. *World Crops: Prod., Util., Descr.*, 13 (Glucosinolates, Rapeseeds) (1987) 90-108; *C.A.*, 107 (1987) 214287a.
- 1556 Subba Rao, G.N.: Liquid chromatographic determination of taurine in vitamin premix formulations. *J. Assoc. Off. Anal. Chem.*, 70 (1987) 799-801.

See also 849, 941, 1042, 1160, 1596.

25. ORGANIC PHOSPHORUS COMPOUNDS (INCLUDING SUGAR PHOSPHATES)

- 1557 Balla, T., Guillemette, G., Baukal, A.J. and Catt, K.J.: Formation of inositol 1,3,4,6-tetrakisphosphate during angiotensin II action in bovine adrenal glomerulosa cells. *Biochem. Biophys. Res. Commun.*, 148 (1987) 199-205.
- 1558 Bird, I.M., Smith, A.D. and Schulster, D.: HPLC analysis of inositol monophosphate isomers formed on angiotensin II stimulation of rat adrenal glomerulosa cells. *Biochem. J.*, 248 (1987) 203-208.
- 1559 Erneux, C., Delvaux, A., Moreau, C. and Dumont, J.E.: The dephosphorylation pathway of D-myo-inositol 1,3,4,5-tetrakisphosphate in rat brain. *Biochem. J.*, 247 (1987) 635-639.
- 1560 Hughes, P.J. and Drummond, A.H.: Formation of inositol phosphate isomers in GH₃ pituitary tumour cells stimulated with thyrotropin-releasing hormone. Acute effects of lithium ions. *Biochem. J.*, 248 (1987) 463-470.
- 1561 Imboden, J.B. and Weiss, A.: The T-cell antigen receptor regulates sustained increases in cytoplasmic free Ca²⁺ through extracellular Ca²⁺ influx and ongoing intracellular Ca²⁺ mobilization. *Biochem. J.*, 247 (1987) 695-700.
- 1562 Kalant, N., Parniak, M. and Lemieux, M.: Compartmentation of glucose 6-phosphate in hepatocytes. *Biochem. J.*, 248 (1987) 927-931.
- 1563 Kiesel, L., Helm, K., Bertges, K., Maier, C., Rabe, T. and Runnebaum, B.: Contraceptive progestins and gonadotropin secretion *in vitro*. *J. Steroid Biochem.*, 27 (1987) 991-1002.
- 1564 Morris, A.J., Downes, C.P., Harden, T.K. and Michell, R.H.: Turkey erythrocytes possess a membrane-associated inositol 1,4,5-triphosphate 3-kinase that is activated by Ca²⁺ in the presence of calmodulin. *Biochem. J.*, 248 (1987) 489-493.
- 1565 Ragan, C.I., Watling, K.J., Gee, N.S., Aspley, S., Jackson, R.G., Reid, G.G., Baker, R., Billington, D.C., Barnaby, R.J. and Leeson, P.D.: The dephosphorylation of inositol 1,4-biphosphate to inositol in liver and brain involves two distinct Li⁺-sensitive enzymes and proceeds via inositol 4-phosphate. *Biochem. J.*, 249 (1988) 143-148.

- 1566 Stephens, L.R., Hawkins, P.T., Carter, N., Chahwala, S.B., Morris, A.J., Whetton, A.D. and Downes, P.C.: L-myo-Inositol 1,4,5,6-tetrakisphosphate is present in both mammalian and avian cells. *Biochem. J.*, 249 (1988) 271-282.
- 1567 Woodcock, E.A., Smith, A.I., Wallace, C.A. and Schmauk White, L.B.: Evidence for a lack of inositol-(1,4,5)trisphosphate kinase activity in norepinephrine-perfused rat hearts. *Biochem. Biophys. Res. Commun.*, 148 (1987) 68-77.
- See also 842, 1477, 1772.

26. ORGANOMETALLIC AND RELATED COMPOUNDS

26a. Organometallic compounds

- 1568 Blair, W.R., Parks, E.J., Olson, G.J., Brinckman, F.F., Valeiras-Price, M.C. and Bellama, J.M.: Characterization of organotin species using microbore and capillary liquid chromatographic techniques with an epifluorescence microscope as a novel imaging detector. *J. Chromatogr.*, 410 (1987) 383-394.
- 1569 Blotcky, A.J., Hansen, G.T., Borkar, N., Ebrahim, A. and Rack, E.P.: Simultaneous determination of selenite and trimethylselenonium ions in urine by anion exchange chromatography and molecular neutron activation analysis. *Anal. Chem.*, 59 (1987) 2063-2066.
- 1570 Ebdon, L. and Alonso, J.I.G.: Determination of tributyltin ions in estuarine waters by high-performance liquid chromatography with fluorimetric detection using morin in micellar solution. *Analyst (London)*, 112 (1987) 1551-1554.

26b. Boranes, silanes and related non-metallic compounds

- 1571 Bertrand, C., Felix, G., Biran, C. and Fourtignon, M.: Preparative HPLC of indole silyl compounds on 3,5-dinitrobenzamide silica gel. *J. Liq. Chromatogr.*, 10 (1987) 853-861.

26c. Coordination compounds

- 1572 Baiocchi, C., Cantone, F., Marchetto, A., Gennaro, M.C., Mentasti, E. and Sarzanini, C.: Reversed-phase HPLC behaviour of metal complexes with 4-(2-pyridylazo)resorcinol. A detailed study in the light of solvophobic theory. *Chromatographia*, 23 (1987) 736-746.
- 1573 Grigorova, B., Wright, S.A. and Josephson, M.: Separation and determination of stable metallo-cyanide complexes in metallurgical plant solutions and effluents by reversed-phase ion-pair chromatography. *J. Chromatogr.*, 410 (1987) 419-426.
- 1574 Noda, H., Saitoh, K. and Suzuki, N.: Elution recovery of tris(acetylacetonato)aluminium(III), -gallium(III) and -indium(III) in size-exclusion chromatography. *J. Chromatogr.*, 435 (1988) 368-373.
- 1575 Palmieri, M.D. and Fritz, J.S.: Determination of metal ions by high-performance liquid chromatographic separation of their hydroxamic acid chelates. *Anal. Chem.*, 59 (1987) 2226-2231.
- 1576 Steinbrech, B.: Thin layer chromatography and high pressure liquid chromatography of metal chelates. *J. Liq. Chromatogr.*, 10 (1987) 1-48 - a review with 77 refs.

See also 1848.

27. VITAMINS AND VARIOUS GROWTH REGULATORS (NON-PEPTIDIC)

- 1577 Aihara, M. and Kouguchi, Y.: Food additives. I. Electrochemical behavior of tocopherols and antioxidants and their analysis by HPLC. *Fukuoka Joshi Daigaku Kaseigakubu Kiyo*, 18 (1987) 67-72; *C.A.*, 107 (1987) 76236e.
- 1578 Bhat, P.V. and Jetten, A.M.: Metabolism of all-trans-retinol and all-trans-retinoic acid in rabbit tracheal epithelial cells in culture. *Biochim. Biophys. Acta*, 922 (1987) 18-27.

- 1579 Bishop, P.D. and Griswold, M.D.: Uptake and metabolism of retinol in cultured Sertoli cells: evidence for a kinetic model. *Biochemistry*, 26 (1987) 7511-7518.
- 1580 Bui, M.H.: Sample preparation and liquid chromatographic determination of vitamin D in food products. *J. Assoc. Off. Anal. Chem.*, 70 (1987) 802-805.
- 1581 Gardner, R.M., Allison, M.J., Hartman, P.A., Reinhardt, T.A. and Horst, R.L.: Factors influencing production of 5(E)-19-nor-10-keto-vitamin D₃ by rumen bacteria. *J. Steroid Biochem.*, 28 (1987) 189-192.
- 1582 Gatautis, V.J. and Pearson, K.H.: Separation of plasma carotenoids and quantitation of beta-carotene using HPLC. *Clin. Chim. Acta*, 166 (1987) 195-206.
- 1583 Gill, H.S., Londowski, J.M., Corradino, R. and Kumar, R.: Synthesis, biologic activity, and protein binding characteristics of a new vitamin D analog, 22-hydroxyvitamin D₃. *J. Steroid Biochem.*, 28 (1987) 147-153.
- 1584 Howells, D.W. and Hyland, K.: Direct analysis of tetrahydrobiopterin in cerebrospinal fluid by high-performance liquid chromatography with redox electrochemistry: prevention of autooxidation during storage and analysis. *Clin. Chim. Acta*, 167 (1987) 23-30.
- 1585 Hung, T.H.T., Seib, P.A. and Kramer, K.J.: Determination of L-ascorbyl 6-palmitate in bread using reverse-phase HPLC with electrochemical detection. *J. Food Sci.*, 52 (1987) 948-974; *C.A.*, 107 (1987) 132763p.
- 1586 Ishizuka, S., Matsui, T., Nakao, Y., Fujita, T., Reichel, H. and Norman, A.W.: Metabolism of the vitamin D₃ derivative (24R)-hydroxycalcidiol by human promyelocytic leukemia cells (HL-60). Isolation and identification of (5Z) and (5E)-(24R)-19-nor-10-oxo-24-hydroxycalcidiol. *Eur. J. Biochem.*, 170 (1987) 475-483.
- 1587 Jensen, R.K., Cullum, M.E., Deyo, J. and Zile, M.H.: Vitamin A metabolism in rats chronically treated with 3,3',4,4',5,5'-hexabromobiphenyl. *Biochim. Biophys. Acta*, 926 (1987) 310-320.
- 1588 Kamiyama, S., Tajika, K. and Ito, Y.: (Measurement of carotene). *Rinsho Kensa*, 31 (1987) 268-274; *C.A.*, 107 (1987) 150454t - a review with 7 refs.
- 1589 Kurumaya, K., Sakamoto, T., Okada, Y. and Kaziwara, M.: Application of droplet counter-current chromatography to the isolation of vitamin B₁₂. *J. Chromatogr.*, 435 (1988) 235-240.
- 1590 Kutnink, M.A., Hawkes, W., Schaus, E.E. and Omaye, S.T.: An internal standard method for the unattended high-performance liquid chromatographic analysis of ascorbic acid in blood components. *Anal. Biochem.*, 166 (1987) 424-430.
- 1591 Lavigne, C., Zee, J.A., Simard, R.E. and Gosselin, C.: High-performance liquid chromatographic-diode-array determination of ascorbic acid, thiamine and riboflavin in goats' milk. *J. Chromatogr.*, 410 (1987) 201-205.
- 1592 Lohnes, D. and Jones, G.: Side chain metabolism of vitamin D₃ in osteosarcoma cell line UMR-106. Characterization of products. *J. Biol. Chem.*, 262 (1987) 14394-14401.
- 1593 Lopez-Anaya, A. and Mayersohn, M.: Quantification of riboflavin, riboflavin 5'-phosphate and flavin adenine dinucleotide in plasma and urine by high-performance liquid chromatography. *J. Chromatogr.*, 423 (1987) 105-113.
- 1594 Lopez-Anaya, A. and Mayersohn, M.: Ascorbic and dehydroascorbic acids simultaneously quantified in biological fluids by liquid chromatography with fluorescence detection, and comparison with a colorimetric assay. *Clin. Chem. (Winston-Salem)*, 33 (1987) 1874-1878.
- 1595 Lucock, M., Hartley, R. and Wild, N.J.: HPLC determination of vitamin K₁ in neonatal plasma following oral or parenteral supplementation with Konakion. *J. Liq. Chromatogr.*, 10 (1987) 191-203.
- 1596 Margolis, S.A. and Black, I.: Stabilization of ascorbic acid and its measurement by liquid chromatography in nonfat dry milk. *J. Assoc. Off. Anal. Chem.*, 70 (1987) 806-809.
- 1597 Nakasato, S., Hoshida, H., Maruyama, T., Ujiie, T., Ohta, Y., Kawaguchi, Y., Shimbo, K., Watanabe, S., Wada, S. et al.: (HPLC determination of tocopherols in vegetable oils and fats collaborative study). *Yukagaku*, 36 (1987) 506-514; *C.A.*, 107 (1987) 52973k.
- 1598 Porteous, C.E., Coldwell, R.D., Trafford, D.J.H. and Makin, H.L.J.: Recent developments in the measurement of vitamin D and its metabolites in human body fluids. *J. Steroid Biochem.*, 28 (1987) 785-801 - a review with 146 refs.
- 1599 Quackenbush, F.W.: Reverse phase HPLC separation of *cis*- and *trans*-carotenoids and its application to β -carotenes in food materials. *J. Liq. Chromatogr.*, 10 (1987) 643-653.

- 1600 Rabe, H. and Volk, D.: Analysis of urinary neopterin in autoimmune diseases with HPLC. *Symp. Biol. Hung.*, 34 (Chromatography '85) (1986) 337-341; *C.A.*, 107 (1987) 150529w.
- 1601 Rebello, T.: Trace enrichment of biological folates on solid-phase adsorption cartridges and analysis by high-pressure liquid chromatography. *Anal. Biochem.*, 166 (1987) 55-64.
- 1602 Rundhaug, J., Gubler, M.L., Sherman, M.I., Blaner, W.S. and Bertram, J.S.: Differential uptake, binding, and metabolism of retinol and retinoic acid by 10T1/2 cells. *Cancer Res.*, 47 (1987) 5637-5643.
- 1603 Samokyszyn, V.M. and Marnett, L.J.: Hydroperoxide-dependent cooxidation of 13-*cis*-retinoic acid by prostaglandin H synthase. *J. Biol. Chem.*, 262 (1987) 14119-14133.
- 1604 Schwartz, S.J.: *Cis/trans* isomerization of retinyl palmitate in foods. *J. Agric. Food Chem.*, 35 (1987) 748-751.
- 1605 Stancher, B., Zonta, F. and Bogoni, P.: Determination of olive oil carotenoids by HPLC. *J. Micronutr. Anal.*, 3 (1987) 97-106; *C.A.*, 107 (1987) 174488h.
- 1606 Tuan, S.H. and Wyatt, J.: The separation of erythroic acid by HPLC. *J. Micronutr. Anal.*, 3 (1987) 119-128; *C.A.*, 107 (1987) 174489j.
- 1607 Ueda, T. and Igarashi, O.: New solvent system for extraction of tocopherols from biological specimens for HPLC determination and the evaluation of 2,2,5,7,8-pentamethyl-6-chromanol as an internal standard. *J. Micronutr. Anal.*, 3 (1987) 185-198; *C.A.*, 107 (1987) 171771j.
- 1608 Vuorela, H., Hinkkanen, R. and Hiltunen, R.: Rapid determination of ascorbic acid in pharmaceuticals and biological samples by HPLC. *Acta Pharm. Fenn.*, 95 (1986) 141-149; *C.A.*, 107 (1987) 205275t.
- 1609 Wall, J.S., Young, M.R. and Carpenter, K.J.: Transformation of niacin-containing compounds in corn during grain development: relationship to niacin nutritional availability. *J. Agric. Food Chem.*, 35 (1987) 752-758.
- 1610 Werner, E.R., Fuchs, D., Hausen, A., Reibnegger, G. and Wachter, H.: Simultaneous determination of neopterin and creatinine in serum with solid-phase extraction and on-line elution liquid chromatography. *Clin. Chem. (Winston-Salem)*, 33 (1987) 2028-2033.
- 1611 Yamada, S., Yamamoto, K., Takayama, H., Hayashi, T., Miyaura, C., Tanaka, H., Abe, E. and Suda, T.: Isolation, identification, and chemical synthesis of 8 α ,25-dihydroxy-9,10-seco-4,6,10(9)-cholestatrien-3-one. A new metabolite of 25-hydroxyvitamin D₃ produced by mouse myeloid leukemia cells (M1). *J. Biol. Chem.*, 262 (1987) 12939-12944.
- 1612 Yamaguchi, T., Uchimura, K., Takenaka, H., Fukumoto, K., Kurokawa, Y., Hamada, M., Sugimoto, S., Matsukura, S., Inoue, K. and Shibuya, N.: Quantitative determination of thiamin and its phosphate esters in human whole blood by HPLC and clinical application. *J. Clin. Biochem. Nutr.*, 2 (1987) 203-216; *C.A.*, 107 (1987) 73589m.

For additional information see:

C.A., 107 (1987) 154169b, 161771v, 174496j, 174512m, 205270n, 223380c.

See also 831, 1091, 1656, 1762, 1809.

28. ANTIBIOTICS

- 1613 Barends, D.M., Brouwers, J.C.A.M. and Hulshoff, A.: Fast pre-column derivatization of aminoglycosides with 1-fluoro-2,4-dinitrobenzene and its application to pharmaceutical analysis. *J. Pharm. Biomed. Anal.*, 5 (1987) 613-617.
- 1614 Bowers, L.D. and Singh, J.: A gradient HPLC method for quantitation of cyclosporine and its metabolites in blood and bile. *J. Liq. Chromatogr.*, 10 (1987) 411-420.
- 1615 Buice, R.G., Stentz, F.B. and Gurley, B.J.: Analytical methodologies for cyclosporine pharmacokinetics: a comparison of radioimmunoassay with high performance liquid chromatography. *J. Liq. Chromatogr.*, 10 (1987) 421-438.
- 1616 Dubourg, D., Saux, M.C., Lafevre, M.A. and Fourtillan, J.B.: Assay of chloramphenicol in biological media by high performance liquid chromatography with UV absorbance as the detection mode. *J. Liq. Chromatogr.*, 10 (1987) 921-940.

- 1617 El-Yazigi, A., Yusuf, A. and Al-Humaidan, A.: Direct, simultaneous measurement of chloramphenicol and its monosuccinate ester in microsamples of plasma by radial-compression liquid chromatography. *Clin. Chem. (Winston-Salem)*, 33 (1987) 1814-1816.
- 1618 Falkowski, A.J., Look, Z.M., Noguchi, H. and Silber, B.M.: Determination of cefixime in biological samples by reversed-phase high-performance liquid chromatography. *J. Chromatogr.*, 422 (1987) 145-152.
- 1619 Ikai, Y., Oka, H., Kawamura, N., Yamada, M., Harada, K.-I. and Suzuki, M.: Improvement of chemical analysis of antibiotics. XIII. Systematic simultaneous analysis of residual tetracyclines in animal tissues using thin-layer and high-performance liquid chromatography. *J. Chromatogr.*, 411 (1987) 313-323.
- 1620 Inman, E.L.: Determination of vancomycin related substances by gradient high-performance liquid chromatography. *J. Chromatogr.*, 410 (1987) 363-372.
- 1621 Kabra, P.M. and Wall, J.H.: Improved liquid chromatographic analysis of cyclosporine in whole blood with solid phase (Bond-ElutTM) extraction. *J. Liq. Chromatogr.*, 10 (1987) 477-490.
- 1622 Katrukha, G.S., Trifonova, Zh.P., Smirnova, I.G., Banini, K. and Vasil'kova, E.A.: (Analytical biospecific chromatography of glycopeptide antibiotics of the vancomycin group). *Antibiot. Med. Biotekhnol.*, 32 (1987) 664-668; *C.A.*, 107 (1987) 205271p.
- 1623 Kekesy, I. and Simonyi, I.: HPLC investigation of chloramphenicol stereoisomers. *Symp. Biol. Hung.*, 34 (1986) 383-389; *C.A.*, 107 (1987) 183642f.
- 1624 Knupp, G., Bugl-Kreickmann, G. and Commichau, C.: (A method for the verification of chloramphenicol residues in animal tissues and eggs by HPLC with radioimmunological detection). *Z. Lebensm.-Unters. Forsch.*, 184 (1987) 390-391; *C.A.*, 107 (1987) 76225a.
- 1625 Lensmeyer, G.L., Wiebe, D.A. and Carlson, I.H.: Identification and analysis of nine metabolites of cyclosporine in whole blood by liquid chromatography. 1. Purification of analytical standards and optimization of the assay. *Clin. Chem. (Winston-Salem)*, 33 (1987) 1841-1850.
- 1626 Lensmeyer, G.L., Wiebe, D.A. and Carlson, I.H.: Identification and analysis of nine metabolites of cyclosporine in whole blood by liquid chromatography. 2. Comparison of patients' results. *Clin. Chem. (Winston-Salem)*, 33 (1987) 1851-1855.
- 1627 Mahdadi, R., Pommery, N., Pommery, J. and Lhermitte, M.: Quantitative determination of adriamycin in rat hepatocytes using a volatile extraction buffer, HPLC and fluorescence detection. *Biomed. Chromatogr.*, 2 (1987) 91-94.
- 1628 Margosis, M.: Quantitative liquid chromatography of ampicillin: collaborative study. *J. Assoc. Off. Anal. Chem.*, 70 (1987) 206-212.
- 1629 McAteer, J.A., Hiltke, M.F., Silber, B.M. and Faulkner, R.D.: Liquid-chromatographic determination of five orally active cephalosporins - cefixime, cefaclor, cefadroxil, cephalixin and cepharadine - in human serum. *Clin. Chem. (Winston-Salem)*, 33 (1987) 1788-1790.
- 1630 Morse, G.D., Nairn, D.K., Bertino, J.S., Jr. and Walshe, J.J.: Overestimation of vancomycin concentrations utilizing a fluorescence polarization immunoassay in patients on peritoneal dialysis. *Ther. Drug Monit.*, 9 (1987) 212-215; *C.A.*, 107 (1987) 70131v.
- 1631 Nilsson, L.-G., Walldorf, B. and Paulsen, O.: Determination of erythromycin in human plasma, using column liquid chromatography with a polymeric packing material, alkaline mobile phase and amperometric detection. *J. Chromatogr.*, 423 (1987) 189-197.
- 1632 Nordlander, I., Johnsson, H. and Osterdahl, B.: Oxytetracycline residues in rainbow trout analyzed by a rapid HPLC method. *Food Addit. Contam.*, 4 (1987) 291-296; *C.A.*, 107 (1987) 152996v.
- 1633 Okada, J. and Kondo, S.: Liquid chromatographic method for determining the macrolide antibiotic sedecamycin and its major metabolites in swine plasma and tissues. *J. Assoc. Off. Anal. Chem.*, 70 (1987) 818-824.
- 1634 Oldfield, N., Chang, D., Garland, W. and Town, C.: Quantitation of ceftetrame in human plasma and urine by high-performance liquid chromatography. *J. Chromatogr.*, 422 (1987) 135-143.
- 1635 Prieto, J.G., Santos, L., Barrio, J.P. and Alonso, M.L.: Intestinal absorption of acidic β -lactam antibiotics: contribution of ionized and un-ionized forms. *J. Pharm. Sci.*, 76 (1987) 596-598.
- 1636 Ristuccia, P.A.: Liquid chromatographic assays of antimicrobial agents. *J. Liq. Chromatogr.*, 10 (1987) 241-276 - a review with 98 refs.

- 1637 Shibata, N., Minouchi, T., Hayashi, Y., Ono, T. and Shimakawa, H.: Quantitative determination of cyclosporin A in whole blood and plasma by HPLC. *Res. Commun. Chem. Pathol. Pharmacol.*, 57 (1987) 261-271; *C.A.*, 107 (1987) 168142z.
- 1638 Sitrin, R.D., DePhillips, P.A. and Dingerdissen, J.J.: Preparative reversed-phase HPLC of polar fermentation products. *Dev. Ind. Microbiol.*, 27 (1987) 65-75; *C.A.*, 107 (1987) 75992m.
- 1639 Van de Water, C. and Haagsma, N.: Determination of chlormaphenicol in swine muscle tissue using a monoclonal antibody-mediated clean-up procedure. *J. Chromatogr.*, 411 (1987) 415-421.
- 1640 Yoshida, K.: (Newly developed HPLC method for simultaneous assay of cefmetazole and gentamicin in combination therapy). *Chemotherapy*, 35 (1987) 482-488; *C.A.*, 107 (1987) 168144b.

For additional information see:

C.A., 107 (1987) 76221w, 89189s, 89190k, 174272h, 190249m.

See also 1549, 1745, 1767.

29. INSECTICIDES, PESTICIDES AND OTHER AGROCHEMICALS

29a. Chlorinated insecticides

- 1641 Noren, K. and Sjövall, J.: Analysis of organochlorine pesticides, polychlorinated biphenyls, dibenzo-*p*-dioxins and dibenzofurans in human milk by extraction with the lipophilic gel Lipidex 5000. *J. Chromatogr.*, 422 (1987) 103-115.

For additional information see:

C.A., 107 (1987) 192383z.

See also 1814.

29b. Phosphorus insecticides

- 1642 Sultatos, L.G.: The role of the liver in mediating the acute toxicity of the pesticide methyl parathion in the mouse. *Drug Metab. Disp.*, 15 (1987) 613-617.
- 1643 Szabo-Ravasz, B.: High-performance liquid chromatographic methods for the determination of contaminants in technical phosmethylan. *J. Chromatogr.*, 435 (1988) 380-384.

29c. Carbamates

- 1644 Michels, K.: (Determination of carbadox, olaquinox, and coyden in mixed feeds and premixes). *Landwirtsch. Forsch.*, 39 (1986) 298-306; *C.A.*, 107 (1987) 95429m.

29d. Herbicides

- 1645 Goewie, C.E. and Hogendoorn, E.A.: Analysis of bromacil, diuron and 3,4-dichloroaniline in contaminated well water, using a high-performance liquid chromatographic column-switching procedure. *J. Chromatogr.*, 410 (1987) 211-216.
- 1646 Lesser, J.H., Shustina, R. and Ovadia, D.: Liquid chromatographic behaviour of chlorinated carbanilides. *J. Chromatogr.*, 410 (1987) 95-102.
- 1647 Simon, V.A.: HPLC assay with confirmation of low levels of paraquat and diquat in well water. *LC-GC*, 5 (1987) 899-903; *C.A.*, 107 (1987) 204703u.
- 1648 Wheeler, J.R. and McNally, M.E.: Comparison of packed column and capillary column supercritical fluid chromatography and high-performance liquid chromatography using representative herbicides and pesticides as typical moderate polarity and molecular weight range molecules. *J. Chromatogr.*, 410 (1987) 343-353.

- 1649 Worobey, B.L.: Analytical method for the simultaneous determination of diquat and paraquat residues in potatoes by high-pressure liquid chromatography. *Pestic. Sci.*, 18 (1987) 245-257; *C.A.*, 107 (1987) 76228d.

See also 1183.

29e. Fungicides

See 1784.

29f. Other types of pesticides and various agrochemicals

- 1650 Abidi, S.L.: Enantiomeric separation of rotenone and rotenolone on chiral stationary phases. *J. Liq. Chromatogr.*, 10 (1987) 1085-1102.
- 1651 Goto, T., Shinshi, E., Tanaka, K. and Manabe, M.: Analysis of cyclopiazonic acid by normal phase HPLC. *Agric. Biol. Chem.*, 51 (1987) 2581-2582; *C.A.*, 107 (1987) 213074s.
- 1652 Hunter, K., Sharp, E.A. and Newton, A.: Determination of diastereoisomers of bromadiolone, an anticoagulant rodenticide, in animal tissues by high-performance liquid chromatography. *J. Chromatogr.*, 435 (1988) 83-95.
- 1653 Pepperman, A.B., Jr. and Wartelle, L.H.: Quantitative HPLC assay of strigol and epistrigol - witchweed germination stimulants. *J. Liq. Chromatogr.*, 10 (1987) 1485-1495.

See also 1648, 1820.

30. SYNTHETIC AND NATURAL DYES

30a. Synthetic dyes

- 1654 Burton, S.J., McLoughlin, S.B., Stead, C.V. and Lowe, C.R.: Design and application of biomimetic anthraquinone dyes. I. Synthesis and characterisation of terminal ring isomers of C.I. Reactive Blue 2. *J. Chromatogr.*, 435 (1988) 127-137.
- 1655 Sweatman, T.W., Larussa, R.I., Seshadri, R. and Israel, M.: An analytical system for the detection and quantitation of rhodamine-123 in biological samples. *J. Liq. Chromatogr.*, 10 (1987) 1417-1429.

For additional information see:

C.A., 107 (1987) 174533u, 200378u.

30b. Chloroplast and other natural pigments

- 1656 Khachik, F. and Beecher, G.R.: Application of a C-45- β -carotene as an internal standard for the quantification of carotenoids in yellow/orange vegetables by liquid chromatography. *J. Agric. Food Chem.*, 35 (1987) 732-738.
- 1657 Pukhila, J., Peltonen, K. and Savolainen, T.: Determination of melamine in air by high-performance liquid chromatography with ultraviolet detection. *J. Chromatogr.*, 411 (1987) 409-414.

See also 1605, 1809.

31. PLASTICS AND THEIR INTERMEDIATES

- 1658 Froment, P. and Revillon, A.: Some aspects of analysis of polymers by steric exclusion chromatography and on-line low angle laser light scattering (SEC-LALLS). *J. Liq. Chromatogr.*, 10 (1987) 1383-1397.
- 1659 Gur'yanova, V.V., Prudskova, T.N. and Pavlov, A.V.: Use of gel-permeation and liquid chromatography in the chemistry of condensation polymers. *Plast. Massy*, (1987) 54-56; *C.A.*, 107 (1987) 199220c - a review with 18 refs.

- 1660 Maurer, J.J. and Klemann, L.P.: Characterization of water soluble polymers by ion chromatography. I. Poly(acrylamide-co-sodium acrylate). *J. Liq. Chromatogr.*, 10 (1987) 83-94.
- 1661 Mogilevich, I.M., Podosenova, N.G. and Chudnova, V.M.: (Monitoring and control of molecular characteristics of vinyl acetate-N-vinylphthalimide random copolymers). *Zh. Prikl. Khim. (Leningrad)*, 60 (1987) 1902-1906; *C.A.*, 107 (1987) 199273x.
- 1662 Mori, S.: Comparison between size-exclusion chromatography and liquid adsorption chromatography in the determination of the chemical heterogeneity of copolymers. *J. Chromatogr.*, 411 (1987) 355-362.
- 1663 Senak, L., Wu, C.S. and Malawer, E.G.: Size exclusion chromatography of poly(vinylpyrrolidone). II. Absolute molecular weight distribution by SEC/LALLS and SEC with universal calibration. *J. Liq. Chromatogr.*, 10 (1987) 1127-1150.
- 1664 Swanson, C.L., Carr, M.E. and Nielsen, H.C.: Calibration of GPC columns for rubber analysis: polystyrene vs. polyisoprene molecular weight standards. *J. Polym. Mater.*, 3 (1986) 211-216; *C.A.*, 107 (1987) 200059r.

For additional information see:

C.A., 107 (1987) 155094k, 155098g.

See also 854, 887.

32. DRUG ANALYSIS

See 920.

32a. Drug analysis. General techniques

- 1665 Biemer, T.A.: Simultaneous analysis of acetaminophen, pseudoephedrine hydrochloride and chlorpheniramine maleate in a cold tablet using an isocratic, mixed micellar high-performance liquid chromatographic mobile phase. *J. Chromatogr.*, 410 (1987) 206-210.
- 1666 Bogusz, M., Klys, M., De Zeeuw, R.A. and Franke, J.P.: Preliminary experience concerning the applicability of the high-speed spectrophotometric HPLC detector in toxicological screening. *Symp. Biol. Hung.*, 34 (Chromatography '85) (1986) 647-657; *C.A.*, 107 (1987) 170051n.
- 1667 De Schutter, J.A., van den Bossche, W. and De Moerloose, P.: Simultaneous effects of amines and alkylsulphonates in reversed-phase ion-pair liquid chromatography - application to the separation of 2-imidazoline drugs. *J. Pharm. Biomed. Anal.*, 5 (1987) 559-576.
- 1668 De Smet, M. and Massart, D.L.: Retention behaviour of acidic, neutral and basic drugs on a CN column using phosphate buffers in the mobile phase. *J. Chromatogr.*, 410 (1987) 77-94.
- 1669 De Smet, M. and Massart, D.L.: Automated method selection for HPLC analysis of drugs. *TrAC*, 6 (1987) 266-271.
- 1670 Doyle, E., Pearce, J.C., Picot, V.S. and Lee, R.M.: Analysis of drugs from biological fluids using disposable solid phase columns. *J. Chromatogr.*, 411 (1987) 325-333.
- 1671 Gant, J.R. and Dong, M.W.: Practical benefit of high-speed liquid chromatography in pharmaceutical analysis. *Pharm. Technol.*, 11 (1987) 44-58; *C.A.*, 107 (1987) 205253j - a review with 35 refs.
- 1672 Henry, M.P.: (Wide-pore bonded phases in the chromatography of pharmaceuticals). *GIT-Suppl.*, (1987) 33-41; *C.A.*, 107 (1987) 205254k.
- 1673 Ruane, R.J. and Wilson, I.D.: The use of C18 bonded silica in the solid phase extraction of basic drugs - possible role for ionic interactions with residual silanols. *J. Pharm. Biomed. Anal.*, 5 (1987) 723-727.
- 1674 Tamai, G., Yoshida, H. and Imai, H.: High-performance liquid chromatographic drug analysis by direct injection of whole blood samples. I. Determination of moderately hydrophobic drugs incorporated into blood corpuscles. *J. Chromatogr.*, 423 (1987) 147-153.

- 1675 Tamai, G., Yoshida, H. and Imai, H.: High-performance liquid chromatographic drug analysis by direct injection of whole blood samples. II. Determination of hydrophilic drugs. *J. Chromatogr.*, 423 (1987) 155-161.
- 1676 Tamai, G., Yoshida, H. and Imai, H.: High-performance liquid chromatographic drug analysis by direct injection of whole blood samples. III. Determination of hydrophobic drugs adsorbed on blood cell membranes. *J. Chromatogr.*, 423 (1987) 163-168.
- 1677 Taylor, H.E. and Ackerman, B.H.: Free drug monitoring by liquid chromatography and implications for therapeutic drug monitoring. *J. Liq. Chromatogr.*, 10 (1987) 323-343 - a review with 45 refs.
- 1678 Wilson, J.M.: Drug assays for the clinical pharmacokinetics laboratory. *J. Liq. Chromatogr.*, 10 (1987) 277-292 - a review with 38 refs.
- 1679 Yoshida, Y., Yamauchi, S., Yokoyama, T. and Iwasa, A.: (Simultaneous determination of pharmaceutical components). *Jpn. Kokai Tokkyo Koho Pat.* JP 61,184,460 (86,184,460) (Cl. G01N30/46), 18 Aug. 1986, Appl. 85/24,963, 12 Feb. 1985; 11 pp.; *C.A.*, 107 (1987) 205320d.

For additional information see:

C.A., 107 (1987) 223353w.

See also 844, 869, 919, 924, 930, 1801.

32b. Antirheumatics and antiinflammatory drugs

See 1094.

32c. Automatic and cardiovascular drugs

- 1680 Ahmed, I., Gokhale, R.D., Shah, M.V. and Patton, T.F.: Physicochemical determinants of drug diffusion across the conjunctiva, sclera, and cornea. *J. Pharm. Sci.*, 76 (1987) 583-586.
- 1681 Ascalone, V. and Dal Bo', L.: Automated high-performance liquid chromatographic and column-switching technique for on-line clean-up and analysis of diltiazem in human plasma. *J. Chromatogr.*, 423 (1987) 239-249.
- 1682 Brown, D.S. and Jenke, D.R.: Application of fast LC to pharmaceutical analysis: dopamine HCl. *J. Chromatogr. Sci.*, 25 (1987) 494-500.
- 1683 Dai, X. and Li, S.: (Determination of verapamil in plasma by HPLC). *Seppu*, 5 (1987) 129-130; *C.A.*, 107 (1987) 126391z.
- 1684 Das Gupta, V.: Quantitation of baclofen in tablets using high-performance liquid chromatography. *J. Liq. Chromatogr.*, 10 (1987) 749-755.
- 1685 Duche, J.-C., Barre, J. and Tillement, J.-P.: Rapid liquid chromatographic determination of debrisoquine and its hydroxy metabolite in human urine to define hydroxylation phenotypes. *J. Chromatogr.*, 423 (1987) 340-343.
- 1686 Erhard, K.F.: Separation of chlorobenzazepine isomers by preparative liquid chromatography. *J. Liq. Chromatogr.*, 10 (1987) 701-714.
- 1687 Giachetti, C., Poletti, P. and Zanolo, G.: Analysis of calcium blocker drugs in plasma, HRGC and HPLC analytical conditions for pharmacokinetic studies. *J. High Resolut. Chromatogr. Commun.*, 10 (1987) 654-658.
- 1688 Golden, M.H. and Zoutendam, P.H.: The determination of minoxidil in human serum by high-performance liquid chromatography with amperometric detection. *J. Pharm. Biomed. Anal.*, 5 (1987) 543-551.
- 1689 Hitscherich, M.E., Rydberg, E.M., Tsilifonis, D.C. and Daly, R.E.: Simultaneous determination of hydrochlorothiazide and propranolol hydrochloride in tablets by high-performance liquid chromatography. *J. Liq. Chromatogr.*, 10 (1987) 1011-1021.
- 1690 Humbert, H., Denouel, J. and Keller, H.P.: Column liquid chromatographic determination of hydrolysed bopindolol, in the picogram per millilitre range in plasma, using cartridge extraction and dual electrochemical detection. *J. Chromatogr.*, 422 (1987) 205-215.
- 1691 Hussain, M.A., Aungst, B.J., Lam, G. and Shefter, E.: Phenylpropranolamine pharmacokinetics in dogs after intravenous, oral, and oral controlled-release doses. *Biopharm. Drug Dispos.*, 8 (1987) 497-505; *C.A.*, 107 (1987) 190304a.
- 1692 Jarvie, D.R., Thompson, A.M. and Dyson, E.H.: Laboratory and clinical features of self-poisoning with salbutamol and terbutaline. *Clin. Chim. Acta*, 168 (1987) 313-322.

- 1693 Johnson, S.M. and Wahba Khalil, S.K.: An HPLC method for the determination of diltiazem and desacetyldiltiazem in human plasma. *J. Liq. Chromatogr.*, 10 (1987) 673-685.
- 1694 Johnson, S.M. and Wahba Khalil, S.K.: An HPLC method for the determination of verapamil and norverapamil in human plasma. *J. Liq. Chromatogr.*, 10 (1987) 1187-1201.
- 1695 Koshakji, R.P. and Wood, A.J.J.: Improved high-performance liquid chromatographic method for the simultaneous determination of propranolol and 4-hydroxypropranolol in plasma with fluorescence detection. *J. Chromatogr.*, 422 (1987) 294-300.
- 1696 Li, T., Yang, D., Liu, Q. and Li, F.: (Determination of trimethoprim in serum by reversed-phase HPLC). *Sepu*, 5 (1987) 198-200; *C.A.*, 107 (1987) 126394c.
- 1697 McLean, A.M., Babcock, -Atkinson, E., Rein, K., Ruggirello, D.A., Conzaes, M.A. and Noonan, P.K.: HPLC assay using fluorescence detection for the simultaneous determination of gallopamil and norgallopamil in human plasma. *Pharm. Res.*, 4 (1987) 327-331; *C.A.*, 107 (1987) 190215x.
- 1698 Meese, C.O., Thalheimer, P. and Eichelbaum, M.: High-performance liquid chromatographic method for the analysis of debrisoquine and its S-(+)- and R-(-)-hydroxy metabolites in urine. *J. Chromatogr.*, 423 (1987) 344-350.
- 1699 Moore, E.S.: Liquid chromatographic determination of coumarin anticoagulants in tablets: collaborative study. *J. Assoc. Off. Anal. Chem.*, 70 (1987) 834-836.
- 1700 Okuda, T., Itoh, S., Yamazaki, M., Nakahama, H., Fukuhara, Y. and Orita, Y.: Biopharmaceutical studies of thiazide diuretics. III. *In vivo* formation of 2-amino-4-chloro-m-benzenedisulfonamide as a metabolite of hydrochlorothiazide in a patient. *Chem. Pharm. Bull.*, 35 (1987) 3516-3518.
- 1701 Reiff, K.: High-performance liquid chromatographic method for the determination of adibendan in plasma and urine. *J. Chromatogr.*, 422 (1987) 346-352.
- 1702 Straka, R.J., Hoon, T.J., Lalonde, R.L., Pieper, J.A. and Botorff, M.B.: Liquid chromatography and fluorescence polarization immunoassay methods compared for measuring flecainide acetate in serum. *Clin. Chem. (Winston-Salem)*, 33 (1987) 1898-1900.
- 1703 Tam, Y.K., Tawfik, S.R., Ke, J., Coutts, R.T., Gray, M.R. and Wyse, D.G.: High-performance liquid chromatography of lidocaine and nine of its metabolites in human plasma and urine. *J. Chromatogr.*, 423 (1987) 199-206.
- 1704 Tank, Y.K. and Soldin, S.J.: Analysis of salbutamol enantiomers in human urine by chiral high-performance liquid chromatography and preliminary studies related to the stereoselective disposition kinetics in man. *J. Chromatogr.*, 422 (1987) 187-195.
- 1705 Wannerberg, O. and Persson, B.: Liquid chromatographic method for the determination of bambuterol hydrochloride and related compounds. *J. Chromatogr.*, 435 (1988) 199-203.
- 1706 Yamada, M. and Uda, Y.: Enhancement of percutaneous absorption of molsidomine. *Chem. Pharm. Bull.*, 35 (1987) 3390-3398.

For additional information see:

C.A., 107 (1987) 89168j, 89193p, 89195r, 211351f, 223385h.

See also 923, 1140, 1521, 1667, 1670, 1673, 1711, 1801.

32d. Central nervous system drugs

- 1707 Alkaysi, H.N., Salem, M.S. and Badwan, A.A.: Reversed-phase high-performance liquid chromatographic analysis of terfenadine. *J. Pharm. Biomed. Anal.*, 5 (1987) 729-733.
- 1708 Anonymous: Acetaminophen in drug tablets. Liquid chromatographic method. Changes in official methods of analysis. *J. Assoc. Off. Anal. Chem.*, 70 (1987) 392.
- 1709 Berg, J.D. and Buckely, B.M.: Rapid measurement of anticonvulsant drug concentrations in the out-patient clinic, using HPLC with direct injection of plasma. *Ann. Clin. Biochem.*, 24 (1987) 488-493; *C.A.*, 107 (1987) 190233b.
- 1710 Cyr, T.D., Matsui, F., Sears, R.W., Curran, N.M. and Lovering, E.G.: Liquid chromatographic methods for assay of carbamazepine, 10,11-dihydrocarbamazepine, and related compounds in carbamazepine drug substance and tablets. *J. Assoc. Off. Anal. Chem.*, 70 (1987) 836-840.

- 1711 Daoud, N., Arvidsson, T. and Wahlund, K.-G.: The determination of lidocaine and its metabolites by a direct injection LC method using large sample volumes of untreated plasma. *J. Pharm. Biomed. Anal.*, 5 (1987) 533-542.
- 1712 Fazio, A., Spina, E. and Pisani, F.: Tricyclic antidepressants analysis by liquid chromatography. *J. Liq. Chromatogr.*, 10 (1987) 223-240.
- 1713 Furlanut, M., Perosa, A., Benetello, P. and Colombo, G.: Electrochemical detection of benperidol in serum for drug monitoring in humans. *Ther. Drug Monit.*, 9 (1987) 343-346; *C.A.*, 107 (1987) 211362k.
- 1714 Ganes, D.A. and Midha, K.K.: Sensitive and specific high-performance liquid chromatographic assay for the quantification of sulforidazine and two diastereomeric sulforidazine-5-sulfoxide metabolites in plasma. *J. Chromatogr.*, 423 (1987) 227-237.
- 1715 Inoue, T. and Suzuki, S.-I.: High-performance liquid chromatographic determination of triazolam and its metabolites in human urine. *J. Chromatogr.*, 422 (1987) 197-204.
- 1716 Juergens, U.: HPLC analysis of antiepileptic drugs in blood samples: microbore separation of fourteen compounds. *J. Liq. Chromatogr.*, 10 (1987) 507-532.
- 1717 Klockowski, P.M. and Levy, G.: Simultaneous determination of diazepam and its active metabolites in rat serum, brain and cerebrospinal fluid by high-performance liquid chromatography. *J. Chromatogr.*, 422 (1987) 334-339.
- 1718 Krieger, D.J.: Liquid chromatographic determination of acetaminophen in tablets: collaborative study. *J. Assoc. Off. Anal. Chem.*, 70 (1987) 212-214.
- 1719 Kwong, T.C.: Analysis of acetylsalicylic acid and its metabolites by liquid chromatography. *J. Liq. Chromatogr.*, 10 (1987) 305-321.
- 1720 Lau, C.E., Falk, J.L., Dolan, S. and Tang, M.: Simultaneous determination of flurazepam and five metabolites in serum by high-performance liquid chromatography and its application to pharmacokinetic studies in rats. *J. Chromatogr.*, 423 (1987) 251-259.
- 1721 Lopez, F.G., Marino, E.L. and Dominguez-Gil, A.: Determination of bentazepam in plasma by high-performance liquid chromatography. *J. Pharm. Biomed. Anal.*, 5 (1987) 553-558.
- 1722 Lovett, L.J., Nygard, G.A. and Wahba Khalil, S.K.: A simple HPLC method for the determination of trazodone in human serum. *J. Liq. Chromatogr.*, 10 (1987) 909-919.
- 1723 Lovett, L.J., Nygard, G.A., Erdmann, G.R., Burley, C.Z. and Wahba Khalil, S.K.: HPLC determination of valproic acid in human serum using ultraviolet detection. *J. Liq. Chromatogr.*, 10 (1987) 687-699.
- 1724 Marunaka, T. and Maniwa, M.: High-performance liquid chromatographic determination of [3,4-di-(4-methoxyphenyl)-5-isoxazolyl]acetic acid and its metabolites in human plasma and urine. *J. Chromatogr.*, 422 (1987) 227-233.
- 1725 Matsunaga, H., Suehiro, T., Saita, T., Nakano, Y., Mori, M., Takata, K. and Oda, K.: High-performance liquid chromatographic determination of pyridostigmine in plasma. *J. Chromatogr.*, 422 (1987) 353-355.
- 1726 Mays, D.C., Nawoot, S., Hilliard, J.B., Pacula, C.M. and Gerber, N.: Inhibition and induction of drug biotransformation *in vivo* by 8-methoxy psoralen: studies of caffeine, phenytoin and hexobarbital metabolism in the rat. *J. Pharmacol. Exp. Ther.*, 243 (1987) 227-233.
- 1727 McMillian, C.L., DeRuiter, J. and Clark, C.R.: Liquid chromatographic optimization of reaction conditions for the synthesis of 2,2-dimethyl-N-benzylmalonamide, a new anticonvulsant. *J. Chromatogr. Sci.*, 25 (1987) 510-513.
- 1728 Mendez-Alvarez, E., Soto-Otero, R. and Sierra-Marcuno, G.: Simultaneous determination of carbamazepine, phenobarbital and their major metabolites in serum, brain, tissue, and urine by high-performance liquid chromatography. *Anal. Lett.*, 20 (1987) 1275-1292.
- 1729 Michotte, Y., Moors, M., Deleu, D., Herregodts, P. and Ebinger, G.: Simultaneous determination of levodopa, carbidopa, 3-O-methyldopa and dopamine in plasma using high-performance liquid chromatography with electrochemical detection. *J. Pharm. Biomed. Anal.*, 5 (1987) 659-664.
- 1730 Mroszczak, E.J., Lee, F.W., Combs, D., Sarnquist, F.H., Juang, B.-L., Wu, A.T., Tokes, L.G., Maddox, M.L. and Cho, D.K.: Ketorolac tromethamine absorption, distribution, metabolism, excretion, and pharmacokinetics in animals and humans. *Drug Metab. Disp.*, 15 (1987) 618-626.

- 1731 Ni, P., Guyon, F., Caude, M. and Rosset, R.: (Automatic determination of clomipramine, imipramine and their demethylated metabolites in plasma by high performance liquid-phase chromatography). *Analisis*, 15 (1987) 77-83; *C.A.*, 107 (1987) 70117v.
- 1732 Pietrogrande, M.C., Borea, P.A., Lodi, G. and Bighi, C.: Reversed-phase ion-pair systems to predict partition coefficients of β -carbolines by HPLC and TLC. *Chromatographia*, 23 (1987) 713-716.
- 1733 Reszka, I. and Tyfczynska, J.: (Quick isolation and detection of diazepam and its main metabolite in blood serum). *Farm. Pol.*, 42 (1986) 472-474; *C.A.*, 107 (1987) 168164h.
- 1734 Rio, J., Hodnett, N. and Bidanset, J.H.: The determination of propoxyphene, norpropoxyphene, and methadone in *postmortem* blood and tissues by HPLC. *J. Anal. Toxicol.*, 11 (1987) 222-224; *C.A.*, 107 (1987) 148709m.
- 1735 Shihabi, Z.K., Dyer, R.D. and Scaro, J.: Serum injection on the HPLC column for pentobarbital assay. *J. Liq. Chromatogr.*, 10 (1987) 663-672.
- 1736 Smith, C.S., Morgan, S.L., Greene, S.V. and Abramson, R.K.: Solid-phase extraction and high-performance liquid chromatographic method for chlorpromazine and thirteen metabolites. *J. Chromatogr.*, 423 (1987) 207-216.
- 1737 Soine, P.J. and Soine, W.H.: High-performance liquid chromatographic determination of the diastereomers of 1-(β -D-glucopyranosyl)amobarbital in urine. *J. Chromatogr.*, 422 (1987) 309-314.
- 1738 Suckow, R.F.: Problems in sampling techniques for psychotropic drug assays. *J. Liq. Chromatogr.*, 10 (1987) 293-304.
- 1739 Sun, J.X.S., Embil, K., Chow, D.S.L. and Lee, C.C.S.: High-performance liquid chromatographic analysis, plasma protein binding and red blood cell partitioning of phenprobamate. *Biopharm. Drug Dispos.*, 8 (1987) 341-351; *C.A.*, 107 (1987) 126458b.
- 1740 Tada, K., Miyahira, A. and Moroji, T.: Liquid chromatographic assay of nitrazepam and its main metabolites in serum, and its application to pharmacokinetic study in the elderly. *J. Liq. Chromatogr.*, 10 (1987) 465-476.
- 1741 Van den Eeckhout, E., Jenjintanant, P., Bens, G. and De Moerloose, P.: Influence of column temperature, water content, polarity, and acidity of isohydric solvents on the separation of butyrophenones on silica gel. *Chromatographia*, 23 (1987) 893-898.
- 1742 Wainer, I.W., Alembik, M.C. and Fishcer, L.J.: The determination of (R)- and (S)-glutethimide and the corresponding 4-hydroxyglutethimide metabolites in human serum and urine using a Pirkle-type HPLC chiral stationary phase. *J. Pharm. Biomed. Anal.*, 5 (1987) 735-740.

For additional information see:

C.A., 107 (1987) 89192n, 89196s, 146700c, 168129a, 168170g, 211343e.

See also 1376, 1521, 1691, 1796, 1799, 1800, 1801.

32e. Chemotherapeutics (except cytostatics and antibiotics)

- 1743 Aerts, M.M.L., Beek, W.M.J. and Brinkman, U.A.T.: Monitoring of veterinary drug residues by a combination of continuous flow techniques and columns-switching high-performance liquid chromatography. I. Sulphonamides in egg, meat and milk using post-column derivatization with dimethylaminobenzaldehyde. *J. Chromatogr.*, 435 (1988) 97-112.
- 1744 Bergqvist, Y., Hjelm, E. and Rombo, L.: Sulfadoxine assay using capillary blood samples dried on filter paper-suitable for monitoring of blood concentrations in the field. *Ther. Drug Monit.*, 9 (1987) 203-207; *C.A.*, 107 (1987) 70129a.
- 1745 DeBarbeyrac, B., Penouil, F., Gardere, J.J., Saux, M.C., Brachet-Liermain, A. and Bebear, C.: (Estimation of serum ceftriaxone. Comparison of microbiological and HPLC methods). *Pathol. Biol.*, 35 (1987) 713-716; *C.A.*, 107 (1987) 211359q.
- 1746 ElSohly, H.N., Croom, E.M. and ElSohly, M.A.: Analysis of the antimalarial sesquiterpene artemisin in *Artemisia annua* by high-performance liquid chromatography (HPLC) with postcolumn derivatization and ultraviolet detection. *Pharm. Res.*, 4 (1987) 258-260; *C.A.*, 107 (1987) 83987d.
- 1747 Gomaa, Z.S.: Determination of diiodohydroxyquin in dosage forms by high performance liquid chromatography. *J. Liq. Chromatogr.*, 10 (1987) 1175-1185.

- 1748 Hennig, U.G.G., Chatten, J.G., Pon, R.A. and von Borstel, R.C.: The detection of chemical impurities by high pressure liquid chromatography, and the genetic activity of medical grades of pyruvium pamoate in *Saccharomyces cerevisiae* and *Salmonella typhimurium*. *Arch. Toxicol.*, 60 (1987) 278-286; *C.A.*, 107 (1987) 183673s.
- 1749 Jehl, F., Monteil, H. and Brogard, J.M.: (Direct determination of clavulanic acid in human biological fluids by HPLC). *Pathol. Biol.*, 35 (1987) 702-706; *C.A.*, 107 (1987) 211358p.
- 1750 Joos, B. and Luethy, R.: Determination of teicoplanin concentrations in serum by high-pressure liquid chromatography. *Antimicrob. Agents Chemother.*, 31 (1987) 1222-1224; *C.A.*, 107 (1987) 168131v.
- 1751 Kelley, J.A., Litterst, C.L., Roth, J.S., Vistica, D.T., Poplack, D.G., Cooney, D.A., Nadkarni, M., Balis, F.M., Broder, S. and Johns, D.G.: The disposition and metabolism of 2',3'-dideoxycytidine, an *in vitro* inhibitor of human T-lymphotrophic virus type III infectivity, in mice and monkeys. *Drug Metab. Disp.*, 15 (1987) 595-601.
- 1752 Konaka, R., Hashimoto, H., Nishimura, R., Kuruma, K. and Hirauchi, K.: (Assays of 6315-S (Flomoxef) and its metabolites in body fluids by HPLC). *Chemotherapy*, 35 (Suppl. 1) (1987) 137-144; *C.A.*, 107 (1987) 126382x.
- 1753 Lacroix, C., Poncet, P., Laine, G., Guyonnaud, C., Ray, M., Menager, S. and Lafont, O.: Microdétermination du pyrazinamide et de ses métabolites (acide pyrazinoïque, acide hydroxy-5-pyrazinoïque, hydroxy-5-pyrazinamide, acide pyrazinurique) plasmatiques et urinaires par chromatographie en phase liquide. *J. Chromatogr.*, 422 (1987) 217-225.
- 1754 Molokhia, A.M., El-Hoofy, S. and Dardiri, M.: Liquid chromatographic assay for amodiaquine in tablets and biological fluids. *J. Liq. Chromatogr.*, 10 (1987) 1203-1215.
- 1755 Mourot, D., Dagorn, M. and Delepine, B.: Liquid chromatographic determination of depletion of bithionol sulfoxide and its two major metabolites in bovine milk. *J. Assoc. Off. Anal. Chem.*, 70 (1987) 810-812.
- 1756 Myers, C.M. and Blumer, J.L.: High-performance liquid chromatography of ciprofloxacin and its metabolites in serum, urine and sputum. *J. Chromatogr.*, 422 (1987) 153-164.
- 1757 Paulson, G.D. and Feil, V.J.: The disposition of N-(4,6-dimethyl-2-pyrimidinyl) benzene [^{14}C] sulfomanide in the rat. *Drug Metab. Disp.*, 15 (1987) 671-675.
- 1758 Petz, M.: Fluorimetrische Bestimmung von Sulfanilamid- und Sulfaguanidinrückständen in Tierischen Matrices durch Ionenaustausch-Hochleistungsflüssigchromatographie. *J. Chromatogr.*, 423 (1987) 217-225.
- 1759 Pizzimenti, F.C., Naccari, F., Alonzo, V. and Rotiroti, D.: Evaluation of cefotaxime concentrations in rat cerebral areas and serum by microbiological method and HPLC. *Chemioterapia*, 6 (1987) 101-104; *C.A.*, 107 (1987) 126370s.
- 1760 Sadana, G.S. and Parikh, G.G.: A high-performance liquid chromatographic method for the determination of mebendazole from its pharmaceutical preparations. *Indian Drugs*, 24 (1987) 533-534; *C.A.*, 107 (1987) 205284v.
- 1761 Staniforth, D.H., Coates, P. and Clarke, J.G.N.: An HPLC assay for sulfapyridine in plasma and its use to assess small bowel transit time after the administration of sulfasalazine. *Int. J. Clin. Pharmacol., Ther. Toxicol.*, 25 (1987) 406-409; *C.A.*, 107 (1987) 171750b.
- 1762 Vanderslice, J.T. and Huang, M.-H.A.: Liquid chromatographic determination of amprolium in poultry feed and premixes using postcolumn chemistry with fluorometric detection. *J. Assoc. Off. Anal. Chem.*, 70 (1987) 920-922.
- 1763 Waggoner, T.B. and Bowman, M.C.: Spectrofluorometric determination of BAY V_p 2674 residues in poultry tissues. *J. Assoc. Off. Anal. Chem.*, 70 (1987) 813-818.
- 1764 Zou, A.-Q., Xie, M. and Luo, X.-D.: Determination of artesunic acid after chemical derivatization with O-p-nitrobenzyl-N,N'-diisopropylisourea by high-performance liquid chromatography and ultraviolet absorption. *J. Chromatogr.*, 410 (1987) 217-221.

For additional information see:

C.A., 107 (1987) 76213v, 223400j.

See also 1636, 1779.

32f. Cytostatics

- 1765 Brons, P.P.T., Wessels, J.M.C., Linssen, P.C.M., Haanen, C. and Speth, P.A.J.: Determination of amacrine in human nucleated hematopoietic cells. *J. Chromatogr.*, 422 (1987) 175-185.
- 1766 Dawson, M.I., O'Krongly, D., Hobbs, P.D., Barrueco, J.R. and Sirotnak, F.M.: Synthesis of the 7-hydroxy metabolites of methotrexate and 10-ethyl-10-deaza-aminopterin. *J. Pharm. Sci.*, 76 (1987) 635-638.
- 1767 Egorin, M.J., Conley, B.A., Forrest, A., Zuhowski, E.G., Sinibaldi, V. and van Echo, D.A.: Phase I study and pharmacokinetics of menogaril (NSC 269148) in patients with hepatic dysfunction. *Cancer Res.*, 47 (1987) 6104-6110.
- 1768 Ehniger, G., Klingebiel, T., Kumbier, I., Schuler, U., Feine, U., Treuner, J. and Waller, H.D.: Stability and pharmacokinetics of m-[¹³¹I]iodobenzylguanidine in patients. *Cancer Res.*, 47 (1987) 6147-6149.
- 1769 El-Yazigi, A. and Al-Humaidan, A.: Rapid analysis of 5-fluorouracil in plasma or formulations by high-performance liquid chromatography. *J. Pharm. Biomed. Anal.*, 5 (1987) 747-753.
- 1770 Formelli, F., Carsana, R. and Pollini, C.: Pharmacokinetics of 4'-deoxy-4'-iodo-doxorubicin in plasma and tissues of tumor-bearing mice compared with doxorubicin. *Cancer Res.*, 47 (1987) 5401-5406.
- 1771 Haim, N., Nemeč, J., Roman, J. and Sinha, B.K.: Peroxidase-catalyzed metabolism of etoposide (VP-16-213) and covalent binding of reactive intermediates to cellular macromolecules. *Cancer Res.*, 47 (1987) 5835-5840.
- 1772 Hardy, R.W., Fraser, P.E., Deber, C.M. and Soldin, S.J.: ³¹P NMR and HPLC investigations of the cyclophosphamide metabolite, phosphoramidate mustard. *Anal. Lett.*, 20 (1987) 1303-1314.
- 1773 Kenley, R.A., Hamme, K.J., Lee, M.O. and Tom, J.: Silica-based size exclusion chromatography to characterize the decapeptide nafarelin in a controlled-release pharmaceutical formulation. *Anal. Chem.*, 59 (1987) 2050-2054.
- 1774 Lien, E.A., Ueland, P.M., Solheim, E. and Kvinnsland, S.: Determination of tamoxifen and four metabolites in serum by low-dispersion liquid chromatography. *Clin. Chem. (Winston-Salem)*, 33 (1987) 1608-1614.
- 1775 Malecki, F., Crawhall, J.C. and Staroscik, R.: Determination of melphalan in the perfusate from patients treated by regional perfusion chemotherapy. *J. Pharm. Biomed. Anal.*, 5 (1987) 717-721.
- 1776 Rustum, A.M. and Hoffman, N.E.: Determination of cyclophosphamide in whole blood and plasma by reversed-phase high-performance liquid chromatography. *J. Chromatogr.*, 422 (1987) 125-134.
- 1777 Umprayn, K., Kennedy, P.E., Lee, J.C., Waugh, W., Riley, C.M. and Stella, V.J.: Development of reversed-phase HPLC assays for cyclodisone and clomesone. *J. Pharm. Biomed. Anal.*, 5 (1987) 625-633.
- 1778 Umprayn, K., Stella, V.J. and Riley, C.M.: Stability indicating assay for fetindomide (NSC 373965), a potential prodrug of mitindomide (NSC 284356), employing high-performance liquid chromatography. *J. Pharm. Biomed. Anal.*, 5 (1987) 675-685.
- 1779 Van der Steuijt, K. and Sonneveld, P.: Concurrent analysis of methotrexate, trimethoprim, sulphamethoxazole and their major metabolites in plasma by high-performance liquid chromatography. *J. Chromatogr.*, 422 (1987) 328-333.
- 1780 Workman, P., Oppitz, M., Donaldson, J. and Lee, F.Y.F.: High-performance liquid chromatography of chlorambucil analogues. *J. Chromatogr.*, 422 (1987) 315-321.

For additional information see:

C.A., 107 (1987) 89191m, 126420h, 161794e.

See also 1116, 1117, 1120, 1179, 1627.

32g. Other drug categories

- 1781 Albani, F., Riva, R., Contin, M. and Baruzzi, A.: Liquid chromatographic analysis of metoclopramide with fluorescence detection in cirrhotic patients. *Biomed. Chromatogr.*, 2 (1987) 135-136.
- 1782 Alvarez, M. and Kusumoto, I.T.: (Quantitative analysis of glycosidic sweeteners from *Stevia rebaudiana* and their hydrolysis products by HPLC). *Arq. Biol. Tecnol.*, 30 (1987) 337-348; C.A., 107 (1987) 196610u.

- 1783 Del Grosso, A.V. and May, J.C.: Gas chromatographic, liquid chromatographic, and titrimetric procedures for determination of glycerin in allergenic extracts and diagnostic antigens: comparative study. *J. Assoc. Off. Anal. Chem.*, 70 (1987) 825-828.
- 1784 Hardee, G.E., Tshabalala, M.A., Moore, J.N. and Gokhale, R.D.: HPLC determination and pharmacokinetics of thiabendazole and its major metabolite 5-hydroxythiabendazole in equine plasma. *Res. Vet. Sci.*, 43 (1987) 13-17; *C.A.*, 107 (1987) 211390t.
- 1785 Huang, S.-M. and Tran-Phan, H.: High-performance liquid chromatographic determination of a new H₂ blocker (ORF 17 910) in plasma. *J. Chromatogr.*, 423 (1987) 358-364.
- 1786 Karnes, H.T., Opong-Mensah, K., Farthing, D. and Beightol, L.A.: Automated solid-phase extraction and high-performance liquid chromatographic determination of ranitidine from urine, plasma and peritoneal dialysate. *J. Chromatogr.*, 422 (1987) 165-173.
- 1787 Kidouchi, K., Sugiyama, N., Morishita, H., Wada, Y., Nagai, S. and Sakakibara, J.: Analytical method for urinary glutarylcarnitine, acetylcarnitine and propionylcarnitine with a carboxylic acid analyser and a reversed-phase column. *J. Chromatogr.*, 423 (1987) 297-303.
- 1788 Motassim, N., Decolin, D., Le Dinh, T., Nicolas, A. and Siest, G.: Direct determination of dextromethorphan and its three metabolites in urine by high-performance liquid chromatography using a precolumn switching system for sample clean-up. *J. Chromatogr.*, 422 (1987) 340-345.
- 1789 Nieder, M. and Jaeger, H.: High performance liquid chromatographic determination of metoclopramide in plasma and urine and its application to biopharmaceutical investigations. *J. High Resolut. Chromatogr. Chromatogr. Commun.*, 10 (1987) 659-664.
- 1790 Palosi-Szantho, V. and Kurcz, M.: HPLC determination of furosemide in blood. *Symp. Biol. Hung.*, 34 (Chromatography '85) (1986) 401-411; *C.A.*, 107 (1987) 146687d.
- 1791 Pieters, F.A.J.M., Vincken, B.J. and Zuidema, J.: Dapsone and monoacetyldapsone determined in serum and saliva by a sensitive high-performance liquid chromatographic method with a single extraction step. *J. Chromatogr.*, 422 (1987) 322-327.
- 1792 Shaw, L.M. and Bonner, H.S.: Detection and measurement of the disulfide WR33278 $[\text{NH}_2(\text{CH}_2)_3\text{NHCH}_2\text{CH}_2\text{S}]_2$ in blood and tissues. *J. Liq. Chromatogr.*, 10 (1987) 439-450.
- 1793 Strong, H.A. and Spino, M.: Highly sensitive determination of cimetidine and its metabolites in serum and urine by high-performance liquid chromatography. *J. Chromatogr.*, 422 (1987) 301-308.
- 1794 Van der Meer, M.J. and Brown, L.W.: Simultaneous determination of amiloride and hydrochlorothiazide in plasma by reversed-phase high-performance liquid chromatography. *J. Chromatogr.*, 423 (1987) 351-357.
- 1795 Venkataram, S. and Rahman, Y.E.: High-performance liquid chromatographic analysis of desferrioxamine using a metal-free system. *J. Chromatogr.*, 411 (1987) 494-497.

For additional information see:

C.A., 107 (1987) 90150d, 168154e, 168155f, 171528k, 211352g.

See also 823, 1047, 1102, 1116, 1117, 1120, 1201, 1224, 1515, 1614, 1615, 1621, 1625, 1626, 1637, 1770, 1807, 1875 - 1877.

32h. Toxicological and forensic applications

- 1796 Fraser, A.D., Isner, A.F. and Perry, R.A.: Distribution of trimipramine and its major metabolites in a fatal overdose case. *J. Anal. Toxicol.*, 11 (1987) 168-170; *C.A.*, 107 (1987) 128569n.
- 1797 Hill, D.W. and Langner, K.J.: HPLC photodiode array UV detection for toxicological drug analysis. *J. Liq. Chromatogr.*, 10 (1987) 377-409.
- 1798 Ito, S., Suzuki, K., Matsui, K. and Matsumoto, H.: Determination of bloodstain origin by HPLC of the hemoglobin peptides. *Hansaigaku Zasshi*, 53 (1987) 68-75; *C.A.*, 107 (1987) 170053g.

- 1799 Selavka, C.M. and Krull, I.S.: The forensic determination of drugs of abuse using liquid chromatography with electrochemical detection: a review. *J. Liq. Chromatogr.*, 10 (1987) 345-375.
- 1800 Sood, S.P. and Green, V.I.: Routine methods in toxicology and therapeutic drug monitoring by HPLC. I. Rapid method for determination of acetaminophen in plasma, including a STAT procedure. *Ther. Drug Monit.*, 9 (1987) 248-254; *C.A.*, 107 (1987) 70133x.
- 1801 Wong, S.H.Y., Marzouk, N. Aziz, O. and Sheeran, S.: Microbore liquid chromatography for therapeutic drug monitoring and toxicology: clinical analyses of theophylline, caffeine, procainamide, and N-acetyl procainamide. *J. Liq. Chromatogr.*, 10 (1987) 491-506.

For additional information see:
C.A., 107 (1987) 72214y.

See also 1151, 1666, 1692.

32i. Plant extracts

- 1802 Besset, S. and Nemes, M.: (Aqueous glycol extract of *Hamamelis virginica* L. Identification of drugs by HPLC). *Bull. Tech./Gattefosse Rep.*, 79 (1986) 71-76; *C.A.*, 107 (1987) 161757n.
- 1803 Glowniak, K., Wawrzynowicz, T. and Soczewinski, E.: Isolation of coumarins from fruits of *Pastinaca sativa* L. by column chromatography and modified thin-layer chromatography. *Symp. Biol. Hung.*, 34 (1986) 251-261; *C.A.*, 107 (1987) 161764v.
- 1804 Lee, S.M. and Klocke, J.A.: Combined florisil, droplet countercurrent, and high performance liquid chromatographies for preparative isolation and purification of azadirachtin from neem (*Azadirachta indica*) seeds. *J. Liq. Chromatogr.*, 10 (1987) 1151-1163.
- 1805 Sheu, S.J., Ho, Y.S., Chen, Y.P. and Hsu, H.Y.: Analysis and processing of Chinese herbal drugs; VI. The study of *Angelicae radix*. *Planta Med.*, 53 (1987) 377-378; *C.A.*, 107 (1987) 161766x.
- 1806 Soediro, I., Kusmardiyani, S. and Sukrasno: (Rapid chromatography as a method for fractionation of plant extracts). *Acta Pharm. Indones.*, 11 (1986) 28-34; *C.A.*, 107 (1987) 171617p - a review with 5 refs.
- 1807 Yamagishi, T., Nishizawa, M., Ikura, M., Hikichi, K., Nonaka, G.-I. and Nishioka, I.: New laxative constituents of rhubarb. Isolation and characterization of rheinosides A, B, C and D. *Chem. Pharm. Bull.*, 35 (1987) 3132-3138.

For additional information see:
C.A., 107 (1987) 76320c, 223376f.

See also 1746.

33. CLINICO-CHEMICAL APPLICATIONS

33a. General papers and reviews

See 1136, 1335, 1529, 1594.

33b. Complex mixtures and profiling (single compounds by cross ref. only)

See 908, 961, 996, 1014, 1056, 1078, 1084, 1101, 1110, 1133, 1134, 1159, 1161-1163, 1169, 1171-1173, 1197, 1215, 1226, 1229, 1300, 1307, 1309, 1325, 1338, 1339, 1342, 1356, 1432, 1443, 1450, 1484, 1582, 1584, 1598, 1617, 1625, 1626, 1629, 1771.

34. FOOD ANALYSES

34a. *General papers and reviews*

1808 Lawrence, J.F.: Recent developments in derivatization for the chromatographic determination of food additives. *J. Chromatogr. Sci.*, 25 (1987) 444-452 - a review with 97 refs.

34b. *Complex mixtures (single compounds by cross ref. only)*

1809 Bricout, J. and Koziat, J.: (Analytical progress in the evaluation of orange juice quality). *Int. Fruchtsaft.-Union, Wiss.-Tech. Komm.*, 19 (1986) 367-375; *C.A.*, 107 (1987) 132707y.

1810 Uhlig, J.W., Chang, A. and Jen, J.J.: Effect of phthalides on celery flavor. *J. Food Sci.*, 52 (1987) 658-660; *C.A.*, 107 (1987) 132933u.

1811 Witte, L.: Analytical methods in the structure elucidation of fermentation products: low molecular weight compounds. *Food Sci. Technol.*, 21 (Food Biotechnol.) (1987) 163-191; *C.A.*, 107 (1987) 196409k.

See also 966, 967, 977, 984, 997, 1034, 1042, 1049, 1053, 1208, 1344, 1509, 1544, 1577, 1580, 1585, 1596, 1604, 1605, 1624, 1632, 1649, 1656, 1743, 1755, 1758, 1821.

35. ENVIRONMENTAL ANALYSIS

35a. *General papers and reviews*

1812 Petrick, G., Schulz, D.E. and Duinker, J.C.: Clean-up of environmental samples by high-performance liquid chromatography for analysis of organochlorine compounds by gas chromatography with electron-capture detection. *J. Chromatogr.*, 435 (1988) 241-248.

35b. *Air pollution (complex mixtures; single compounds by cross ref. only)*

1813 Draisci, R., Cecinato, A., Brancaloni, E. and Ciccioli, P.: Preparation and performance of small bore columns packed with reversed-phase materials and their use for the evaluation of air pollutants. *Chromatographia*, 23 (1987) 803-810.

1814 Fuchs, G. and Baechmann, K.: Evaluation of analytical methods for the determination of degradation products of chlorinated hydrocarbons. *Comm. Eur. Communities*, EUR 10832 (1987) 126-132; *C.A.*, 107 (1987) 183138w.

For additional information see:
C.A., 107 (1987) 168036r.

See also 978.

35c. *Water pollution (complex mixtures; single compounds by cross ref. only)*

1815 Pozsgai, A.: (Ion chromatograph in environmental protection). *Meres Auton.*, 35 (1987) 103-104; *C.A.*, 107 (1987) 183110f.

1816 Schwedt, G. and Hippe, H.: (Combined procedures for characterization of organic compounds in water with regard to metal complexation (element species). *Vom Wasser*, 68 (1987) 151-164; *C.A.*, 107 (1987) 222905r.

See also 1645, 1840, 1858, 1866.

35d. *Soil pollution (complex mixtures; single compounds by cross ref. only)*

See 1850.

36. SOME TECHNICAL PRODUCTS AND COMPLEX MIXTURES

36a. Surfactants

- 1817 Boiani, J.A.: Spectator ion indirect photometric detection of aliphatic anionic surfactants separated by reversed-phase high-performance liquid chromatography. *Anal. Chem.*, 59 (1987) 2583-2586.
- 1818 Hwang, R.J. and Stauffer, M.: HPLC analysis of cosurfactants used in enhanced oil recovery. *J. Liq. Chromatogr.*, 10 (1987) 603-615.
- 1819 Kaplan, B. and Pras, M.: Removal of sodium dodecyl sulphate from proteins by gel permeation chromatography. *J. Chromatogr.*, 423 (1987) 376-379.
- 1820 Schreuder, R.H. and Marijn, A.: Determination of dodecylbenzenesulphonates and ethoxylated alkylphenols in liquid pesticide formulations by high-performance liquid chromatography. *J. Chromatogr.*, 435 (1988) 73-82.

For additional information see:

C.A., 107 (1987) 157006f, 223032r.

36b. Antioxidants and preservatives

- 1821 Bui, L.V. and Cooper, C.: Reverse-phase liquid chromatographic determination of benzoic and sorbic acids in foods. *J. Assoc. Off. Anal. Chem.*, 70 (1987) 892-896.
- 1822 Verhagen, H., Thijssen, H.H.W. and Kleinjans, J.C.S.: Determination of butylated hydroxytoluene in plasma by high-performance liquid chromatography. *J. Chromatogr.*, 422 (1987) 288-293.

For additional information see:

C.A., 107 (1987) 183298y.

See also 1038.

36c. Various technical products

- 1823 Bartl, P.: (Evaluation of the quality of synthetic oils for aircraft turbines using chromatographic, spectroscopic and thermal analysis methods). *Tribol. Schmierungstech.*, 34 (1987) 208-216; *C.A.*, 107 (1987) 179605d - a review with 13 refs.
- 1824 Chase, H.A.: Rapid chromatographic monitoring of bioprocesses. *Biosensors*, 2 (1986) 269-286; *C.A.*, 107 (1987) 152724e.
- 1825 Garrick, N.W. and Wood, L.E.: Relationship between high-pressure gel permeation chromatography data and the rheological properties of asphalts. *Transp. Res. Rec.*, 1096 (1986) 35-41; *C.A.*, 107 (1987) 182238s.
- 1826 Gorbunova, L.V., Filimonova, T.A. and Kam'yanov, V.F.: (Chromatographic separation and structural analysis of resinous components of west Siberia crude oils). *Neftekhimiya*, 27 (1987) 462-468; *C.A.*, 107 (1987) 179758f.
- 1827 Platonov, V.V., Tabolenko, N.V., Klyavina, O.A., Ivleva, L.N., Gomez, N. and Vol-Epshtein, A.B.: (Composition and structure of compounds of organic bases in Kansk-Achinsk brown coal tars). *Khim. Tverd. Topl. (Moscow)*, (1987) 27-32; *C.A.*, 107 (1987) 220237a.
- 1828 Rantanen, T. and Sunquist, J.: The use of size-exclusion chromatography in optimization of dissolving pulp and viscose manufacture. *Tappi J.*, 70 (1987) 109-111; *C.A.*, 107 (1987) 156720v.
- 1829 Rodgers, P.A., Creagh, A.L., Prange, M.M. and Prausnitz, J.M.: Molecular weight distribution for heavy fossil fuels from gel-permeation chromatography and characterization data. *Ind. Eng. Chem. Res.*, 26 (1987) 2312-2318; *C.A.*, 107 (1987) 201584v.
- 1830 Rusin, A. and Modrzejewski, Z.: (Use of column chromatography and ultraviolet spectrometry for determination of basic and acidic compounds in products of coal hydrogenation). *Chem. Anal. (Warsaw)*, 31 (1986) 419-430; *C.A.*, 107 (1987) 220258h.

- 1831 Stock, A.F.: Review of the application of high-pressure liquid chromatography to pavements. *Trans. Res. Rec.*, 1096 (1986) 100-105; *C.A.*, 107 (1987) 182169v - a review with 14 refs.

For additional information see:

C.A., 107 (1987) 179680z.

See also 1054, 1144, 1811.

36d. Complex mixtures and unidentified compounds

- 1832 Egestad, B., Pettersson, P., Sjövall, J., Rafter, J., Hyvönen, K. and Gustafsson, J.-A.: Studies on the chromatographic fractionation of metabolites of benzo[*a*]pyrene in faeces and urine from germfree and conventional rats. *Biomed. Chromatogr.*, 2 (1987) 120-134.
- 1833 Gallice, P., Monti, J.-P., Le Gall, J.-M., Crevat, A. and Muriasco, A.: Ion-pair high-performance liquid chromatography profiling of a uremic toxin fraction. *J. Chromatogr.*, 422 (1987) 263-266.
- 1834 Krausse, R. and Ullmann, U.: Identification of anaerobic bacteria using HPLC. *Zentralbl. Bacteriol., Microbiol. Hyg., Ser. A*, 265 (1987) 340-352; *C.A.*, 107 (1987) 214317k.

37. CELLS, CELLULAR PARTICLES AND SUPRAMOLECULAR STRUCTURES

- 1835 Ferrer, E., Martin, M., Alonso, R. and Luque, J.: Separation by counter-current distribution of young-chick erythrocyte populations and numerical resolution of distribution curves. *J. Chromatogr.*, 411 (1987) 167-176.
- 1836 Hammer, D.A., Linderman, J.J., Graves, D.J. and Lauffenburger, D.A.: Affinity chromatography for cell separation: Mathematical model and experimental analysis. *Biotechnol. Prog.*, 3 (1987) 189-204; *C.A.*, 107 (1987) 194391z.
- 1837 Howard, G.A.: Skeletal growth factor affinity columns to separate bone cells by activity. *Int. Congr. Ser.-Excerpta Med.*, 735 (Calcium Regul. Bone Metab.) (1987) 295-300; *C.A.*, 107 (1987) 130196g.
- 1838 Kataoka, K., Sakurai, Y. and Tsuruta, T.: Affinity selection of cells on solid-phase matrices with immobilized proteins. *ACS Symp. Ser.*, 343 (Proteins Interfaces) (1987) 603-614; *C.A.*, 107 (1987) 171738d.
- 1839 Privat, J.-P., Erget-Charlier, M. and Ptak, M.: Instability of bovine brain clathrin-coated vesicles on Sephacryl S-1000 gel chromatography. *Anal. Biochem.*, 166 (1987) 18-26.

38. INORGANIC COMPOUNDS

38a. Cations

- 1840 Akaiwa, H., Kawamoto, H. and Kinoshita, H.: (Preconcentration of copper(II) with 4,7-diphenyl-1,10-phenanthroline disulfonic acid-loaded resin). *Bunseki Kagaku*, 36 (1987) 662-665.
- 1841 Akseli, A.: Use of radiochemical methods for cation-exchange separation of rare earths in a monazite. *Chim. Acta Turc.*, 14 (1986) 101-107; *C.A.*, 107 (1987) 189746q.
- 1842 Aoyama, E., Nakagawa, T., Hasegawa, N., Tanaka, T., Chikuma, M., Nakayama, M. and Tanaka, H.: (Preconcentration of selenium using micro flow-column packed with anion-exchange resin loaded with Bismuthiol-II sulfonic acid). *Bunseki Kagaku*, 36 (1987) 801-803.
- 1843 Dufer, P., Vobecky, M., Holik, J. and Valasek, J.: Separation of some rare earths and transplutonium elements by high-performance liquid chromatography on Ostion cation exchanger. *J. Chromatogr.*, 435 (1988) 249-252.
- 1844 Fukuda, H., Tsunoda, J., Matsumoto, K. and Terada, K.: (Determination of trace amounts of antimony in water by hybriide generation AAS after preconcentration with thionalide loaded silica gel). *Bunseki Kagaku*, 36 (1987) 683-687.

- 1845 Guseva, L.I., Tikhomirova, G.S. and Stepushikina, V.V.: (Isolation and separation of transplutonium elements from other actinides on ion exchangers from aqueous and aqueous-ethanolic sulphuric acid solutions). *Radiokhimiya*, 29 (1987) 211-216; *C.A.*, 107 (1987) 189744n.
- 1846 Heberling, S.S., Riviello, J.M., Shifen, M. and Ip, A.W.: Separate lanthanides by ion chromatography. *Res. Dev.*, 29 (1987) 74-77; *C.A.*, 107 (1987) 189661h - a review with 4 refs.
- 1847 Hosoe, M., Oi, T., Kawada, K. and Kakihana, H.: Separation of rubidium isotopes by ion-exchange chromatography. *J. Chromatogr.*, 435 (1988) 253-255.
- 1848 Hsueh, Y.-M., Miyamura, K., Sawada, T. and Gohshi, Y.: (Determination of cadmium using preconcentration with adsorbent followed by photoacoustic examination of one particle). *Bunseki Kagaku*, 36 (1987) 698-701.
- 1849 Karcher, B.D. and Krull, I.S.: Fluorescence detection of metal ions separated on a silica-based HPLC reversed-phase support. *J. Chromatogr. Sci.*, 25 (1987) 472-478.
- 1850 Kuroda, R., Hayashibe, Y. and Oguma, K.: (A combined anion-exchange/spectrophotometric method for the determination of traces of tin in environmental materials). *Bunseki Kagaku*, 36 (1987) 652-655.
- 1851 Mooney, J.P., Meaney, M., Smyth, M.R., Leonard, R.G. and Wallace, G.G.: Determination of copper(II) and iron(III) in some anaerobic adhesive formulation using high-performance liquid chromatography. *Analyst (London)*, 112 (1987) 1555-1558.
- 1852 Razbash, A.A., Bykhovskii, D.N. and Sevast'yanov, Yu.G.: (Ion-exchange behavior of a set of elements on phosphonate group-containing cation exchanger in hydrochloric acid solutions). *Radiokhimiya*, 29 (1987) 189-193; *C.A.*, 107 (1987) 189743m.
- 1853 Ross, J.W., Jr., Goldring, L.S. and Riseman, J.H.: Method and apparatus for chromatographic analysis of ion mixture. *U.S. Pat.* US 4,672,042 (Cl. 436-161; G01N30/14), 09 Jun. 1987, US Appl. 332,370, 21 Dec. 1981; 8 pp.; *C.A.*, 107 (1987) 167957g.
- 1854 Ryzhkov, N.T.: (Spectrochemical determination of a group of metals in solutions with preliminary extraction-chromatographic separation of iron). *Gidrokhim. Mater.*, 93 (1985) 87-97; *C.A.*, 107 (1987) 161174c.
- 1855 Singh, D.K. and Mehrotra, P.: Separation of transition metal ions and preconcentration of platinum(IV) and chromium(III) on alumina-immobilized diethylene-triaminepentaacetic acid by ligand exchange. *Chromatographia*, 23 (1987) 747-751.
- 1856 Strelow, F.W.E.: Separation of traces of indium from gram amounts of tellurium, mercury and cadmium by cation-exchange chromatography on a single column. *S. Afr. J. Chem.*, 40 (1987) 179-182; *C.A.*, 107 (1987) 189751n.
- 1857 Strelow, F.W.E. and van der Walt, T.N.: Separation of trace amounts of indium, gallium and aluminum from each other by cation-exchange chromatography on AG50W-X4 resin. *Talanta*, 34 (1987) 895-897; *C.A.*, 107 (1987) 227966w.
- 1858 Tanaka, T., Higashi, K., Kawahara, A., Wakida, S.-i., Yamane, M. and Hiroy, K.: (Determination of trace amounts of iron(III) in environmental waters by ion exchange preconcentration and UV detection). *Bunseki Kagaku*, 36 (1987) 647-651.
- 1859 Usuda, S.: Anion exchange behavior of the transplutonium and rare earth elements in nitric acid-methyl alcohol media at elevated temperature. *J. Radioanal. Nucl. Chem.*, 111 (1987) 399-410; *C.A.*, 107 (1987) 184313m.
- 1860 Victor, A.H.: Accurate determination of nickel in South African primary and secondary geochemical reference materials by cation exchange chromatography and atomic absorption spectrometry. *Geostand. Newsl.*, 11 (1987) 187-192; *C.A.*, 107 (1987) 228037f.
- 1861 Yamada, H. and Hattori, T.: Determination of molybdenum in soils and plants by high-performance liquid chromatography. *J. Chromatogr.*, 411 (1987) 401-408.
- 1862 Yoshimura, K.: (On-line concentration and flow analysis of trace amounts of bismuth with anion-exchange method and ion-exchanger absorptiometry). *Bunseki Kagaku*, 36 (1987) 656-661.

For additional information see:

C.A., 107 (1987) 167862x, 189749t, 211066s, 227964u, 227987d.

See also 794, 1572, 1867.

38b. Anions

- 1863 Diop, A., Jardy, A., Caude, M. and Rosset, R.: (Ion chromatography with benzene-polycarboxylate buffers as a mobile phase. II. Indirect spectrophotometric detection of UV transparent anions and quantitative analysis). *Analusis*, 15 (1987) 168-178; *C.A.*, 107 (1987) 227963t.
- 1864 Michlewicz, K.G. and Carnahan, J.W.: A directly coupled microwave-induced plasma atomic emission liquid chromatography detector for nonmetals: preliminary characterization with halides and oxohalogen salts. *Anal. Lett.*, 20 (1987) 1193-1205.
- 1865 Nonomura, M.: Indirect determination of cyanide compounds by ion chromatography with conductivity measurement. *Anal. Chem.*, 59 (1987) 2073-2076.
- 1866 Okada, T.: (Determination of trace nitrite by ion chromatography with large volume injection; Elimination of major interferences anions with high-capacity ion exchange resin column). *Bunseki Kagaku*, 36 (1987) 702-706.
- 1867 Polite, L.N., McNair, H.M. and Rocklin, R.D.: Linearity in ion chromatography. *J. Liq. Chromatogr.*, 10 (1987) 829-838.
- 1868 Takahashi, Y.: (Determination of trace amounts of boron by ICP-AES after ion exchange preconcentration and separation). *Bunseki Kagaku*, 36 (1987) 693-697.
- 1869 Talmage, J.M. and Biemer, T.A.: Determination of potassium nitrate and sodium monofluorophosphate in the presence of phosphate and sulfate by high-resolution ion chromatography. *J. Chromatogr.*, 410 (1987) 494-499.
- 1870 Tanaka, A., Imasaka, Y.-i., Hayashi, K. and Deguchi, T.: (Gel chromatographic deproteinization for fluorometric catalytic determination of iodide and thiocyanate). *Bunseki Kagaku*, 36 (1987) 811-814.
- 1871 Vadhavavikit, S., Kraus, R.J. and Ganther, H.E.: Metabolism of selenocyanate in the rat. *Arch. Biochem. Biophys.*, 258 (1987) 1-6.
- 1872 Watkinson, J.H., Lee, A. and Lauren, D.R.: Measurement of elemental sulfur in soil and sediments: Field sampling, sample storage, pretreatment, extraction and analysis by HPLC. *Aust. J. Soil Res.*, 25 (1987) 167-178; *C.A.*, 107 (1987) 153384f.
- 1873 Yarita, T., Masui, A. and Noshiro, M.: (Analysis of total- and sulphide-sulphur in silicate glasses by using strong phosphoric acid in combination with ion chromatography). *Asahi-Garasu Kenkyu Hokoku*, 37 (1987) 37-42; *C.A.*, 107 (1987) 181777e.

For additional information see:

C.A., 107 (1987) 189762s, 189969q, 204800y, 228073q.

See also 830, 843, 1853.

38d. Volatile inorganic compounds

- 1874 Hamano, T., Mitsuhashi, Y. and Yamamoto, S.: Determination of hydrogen peroxide in beverages by high-performance liquid chromatography with fluorescence detection. *J. Chromatogr.*, 411 (1987) 423-429.

39. RADIOACTIVE AND OTHER ISOTOPE COMPOUNDS

- 1875 Huigen, Y.M., Gelsema, W.J. and De Ligny, C.L.: Gel chromatographic analysis of the bone seeking radiopharmaceutical $^{99m}\text{Tc}(\text{Sn})\text{-EHDP}$: the influence of pH and EHDP-concentration on the size of its constituent. *Appl. Radiat. Isot.*, 38 (1987) 615-618; *C.A.*, 107 (1987) 161655k.
- 1876 Stylli, C. and Theobald, A.E.: Determination of ionization constants of radiopharmaceuticals in mixed solvents by HPLC. *Appl. Radiat. Isot.*, 38 (1987) 701-708; *C.A.*, 107 (1987) 205148d.
- 1877 Torres de Toledo e Souza, I., Sosa de Pereira, N. and Goncalves da Silva, C.P.: Quality control procedures for iodinated radiopharmaceuticals ^{131}I -Hippuran and ^{131}I -Risa. *Publ. IPEN* 1986, Publ. IPEN 97, 6 pp.; *C.A.*, 107 (1987) 183674t.

See also 1841, 1845, 1859.

Gas Chromatography

1. REVIEWS AND BOOKS

- 437 Desty, D.H.: The history and potentiality of capillary columns in gas chromatography. *J. Chromatogr. Sci.*, 25 (1987) 552-563.
- 438 Grob, K.: *On-Column Injection in Capillary Gas Chromatography: Basic Technique, Retention Gaps, Solvent Effects*. Huethig Verlag, Heidelberg, 1987, 591 p.
- 439 Katz, E.: *Quantitative Analysis Using Chromatographic Techniques*. Wiley, New York, 1987.
- 440 Kuksis, A. (Editor): *Chromatography of Lipids in Biochemical Research and Clinical Diagnosis*. *J. Chromatogr. Libr.*, Vol. 37, Elsevier, Amsterdam 1987.
- 441 Pavlov, Yu. N., Rutenberg, O.L. and Fatkudinova, S.R.: (Introduction of GOST 8.485-83. GSI. Analytical gas chromatography laboratory. Methods and means of control). *Izmer. Tekh.*, No. 7 (1987) 55-56; *C.A.*, 107 (1987) 205613b.
- 442 Traitler, H.: Recent advances in capillary gas chromatography applied to lipid analysis. *Prog. Lipid Res.*, 26 (1987) 257-280; *C.A.*, 107 (1987) 194320a - a review with 164 refs.
- 443 Warner, M.: New promise for an old technique. *Anal. Chem.*, 59 (1987) 1403A-1405A.
- 444 Zumwalt, R.W. (Editor): *Amino Acid Analysis by Gas Chromatography*, Vols. 1, 2, 3. CRC Press, Boca Raton 1987, 630 p.

See also 445, 519, 550, 571, 744, 789.

2. FUNDAMENTALS, THEORY AND GENERAL

2a. General

- 445 Etre, L.S.: Twenty-five years ago - a personal reminiscence. *J. Chromatogr. Sci.*, 25 (1987) 530-535.
- 446 Roth, M.: Effect of column inlet concentration profile on the sensitivity of analysis. A quantitative assessment. *J. Chromatogr.*, 409 (1987) 360-364.

See also 548, 594.

2b. Thermodynamics and theoretical relationships

- 447 Chen Songying and Peng Shaoyi: (A generalized equation for HETP in column chromatography). *Sci. Sin., Ser. B*, 30 (1987) 572-581; *C.A.*, 107 (1987) 167822j.
- 448 Frey, D.D.: Free energy consumption at self-sharpening concentration fronts in isothermal fixed-bed adsorption. Theory for isotherms of arbitrary shape and applications in displacement chromatography. *J. Chromatogr.*, 409 (1987) 1-13.
- 449 Kolosov, N.V. and Zolotarev, P.P.: (Theory of chromatographic peak broadening in capillary column of rectangular cross section I. Formulation of the problem and basic equations). *Zh. Fiz. Khim.*, 61 (1987) 2733-2738.
- 450 Kolosov, N.V. and Zolotarev, P.P.: (Theory of chromatographic peak broadening in capillary section II. Analysis of obtained results). *Zh. Fiz. Khim.*, 61 (1987) 2739-2743.
- 451 Zenkovich, I.G., Maevskii, G.A. and Ioffe, B.V.: (On the theory of gas chromatographic analysis in the nonisothermal regimes). *Dokl. Akad. Nauk SSSR*, 297 (1987) 101-105.

See also 535.

2c. Relationship between structure and chromatographic behaviour

- 452 Dimov, N.: (Identification possibilities by using gas chromatography). *Izv. Khim.*, 19 (1986) 437-443; *C.A.*, 107 (1987) 146537e.
- 453 Golovnya, R.V. and Grigor'eva, D.N.: (Principle of nonadditivity of energy contributions to the energy of sorption and its use for identification in chromatography). *Izv. Akad. Nauk SSSR, Ser. Khim.*, (1987) 596-602.
- 454 Poole, C.F., Poole, S.K., Pomaville, R.M. and Kersten, B.R.: Importance of the retention index standards in determining the magnitude of McReynolds phase constants. *J. High Resolut. Chromatogr. Chromatogr. Commun.*, 10 (1987) 670-671.
- 455 Wang Fuan and Wang Gang: (Prediction of the retention indexes of alkanes on nonsqualane stationary liquids). *Shiyou Huagong*, 16 (1987) 439-441; *C.A.*, 107 (1987) 146559p.

See also 508, 517, 521, 598, 599, 609, 776, 787, 819.

2d. Measurement of physico-chemical and related values

- 456 Abraham, M.H., Buist, G.J., Grellier, P.L., McGill, R.A., Doherty, R.M., Kamlet, M.J., Taft, R.W. and Maroldo, S.G.: Solubility properties in polymers and biological media. II. A new method for the characterisation of the adsorption of gases and vapours on solids. *J. Chromatogr.*, 409 (1987) 15-27.
- 457 Below, E., Moeller, E., Pscheidl, H. and Haberland, D.: (Chromatographic reaction process. Gas chromatographic determination of kinetics processes of association of hydrocarbons in a dissolved aluminum chloride catalyst). *Z. Chem.*, 27 (1987) 184-185; *C.A.*, 107 (1987) 217024y.
- 458 Berezkin, V.G. and Korolev, A.A.: (Effect of adsorption in a coated capillary column on the determination of the heat of solution by gas chromatography). *Zh. Fiz. Khim.*, 61 (1987) 1402-1404.
- 459 Bischof, J.G., Haeusler, K.G., Popov, G. and Schwachula, G.: (Determining the pore volume of plastics and coal). *Ger. (East) Pat.* DD 243,766 (Cl.G01N15/08), 11 Mar. 1987, Appl. 283,014, 20 Nov. 1985; 5 pp.; *C.A.*, 107 (1987) 155234f.
- 460 Burdick, D.S. and Rayens, W.S.: Using discriminant analysis to estimate linear mixing proportions. *J. Chemom.*, 1 (1987) 157-173; *C.A.*, 107 (1987) 211283k.
- 461 Fan Wo, Mu Wantong, Sun Xiurong, Ma Jiayu, Yang Weifan, Xie Xiujun and Zhao Lili: (Determination of adsorption enthalpies of thallium-204 oxides by isothermal gas chromatographic and thermochromatographic methods). *He Huaxue Yu Fangshe Huaxue*, 9 (1987) 109-114; *C.A.*, 107 (1987) 162901z.
- 462 Filimonov, V.N., Milyaev, Yu.F. and Balyatinskaya, L.N.: (Distribution coefficients of alcohols in the air-butanol system). *Zh. Prikl. Khim.*, 60 (1987) 1399-1401.
- 463 Gurnagul, N. and Gray, D.G.: Surface adsorption at low coverage of *n*-alkanes on bleached kraft papers using gas chromatography. *Can. J. Chem.*, 65 (1987) 1935-1939; *C.A.*, 107 (1987) 156737j.
- 464 Ioffe, B.V., Katsanos, N.A., Kokovina, L.A., Marinichev, A.N. and Stolyarov, B.V.: (Reverse flow gas chromatography study of the kinetics of liquid-phase chemical reactions complicated by mass transfer). *Kinet. Katal.*, 28 (1987) 805-810; *C.A.*, 107 (1987) 206000m.
- 465 Isono, K., Yamamoto, H. and Tokunaga, J.: (Gas chromatographic determination of solubilities of carbon dioxide in *n*-propyl alcohol, *sec*-butyl alcohol, and *n*-amyl alcohol). *Chem. Express*, 1 (1986) 627-630; *C.A.*, 107 (1987) 197147k.
- 466 Ito, M.M., Kato, J., Takamura, I., Wakatsuki, K. and Endo, T.: Unusual effect of methylation of a benzene ring on the strength of phenyl-phenyl interactions. *Chem. Lett.*, (1987) 241-244; *C.A.*, 107 (1987) 175195x.
- 467 Itsuki, H., Teresawa, S., Yamana, N. and Ohtaka, S.: Gas-liquid chromatographic determination of partial molar free energies at infinite dilution of 18 organic solutes in *n*-hexane solvent at 25.0°C. *Anal. Chem.*, 59 (1987) 2918-2921.
- 468 Jönsson, J.A. and Lökvist, P.: Determination of adsorption isotherms from chromatographic measurements, using the peak maxima method. *J. Chromatogr.*, 408 (1987) 1-7.
- 469 Kersten, B.R., Poole, C.F. and Furton, K.G.: Ambiguities in the determination of McReynolds stationary phase constants. *J. Chromatogr.*, 411 (1987) 43-59.

- 470 Li Fuhua, He Jiajun, Wen Xiaolin, Lu Suhuan and Peng Yincheng: (Determination of activity coefficients of toluene and *n*-decane at infinite dilution in polystyrene solutions by gas chromatography). *Gaofenzi Xuebao*, (1987) 161-165; *C.A.*, 107 (1987) 218456c.
- 471 Lin Weizhong, Tang Xinshuo and Jin Songshou: Studies on adsorption models of reactants on catalysts by pulse gas chromatography. *Shiyou Huagong*, 16 (1987) 511-517; *C.A.*, 107 (1987) 178572d.
- 472 Nawrocki, J.: How to count strong adsorption sites by gas chromatography? *Chromatographia*, 23 (1987) 722-726.
- 473 Tian Bingshou, Chen Fan'geng and Yang Jie: (Study of thermodynamic properties of chelating resin by inverse gas chromatography). *Sepu*, 5 (1987) 339-343.
- 474 Voelkel, A., Szymanowski, J., Berger, J. and Ebert, K.: Polarity of 1,3-bis(ω -alkoxyoligo(oxyethylene)) propan-2-ols as measured by reversed-phase gas chromatography. *J. Chromatogr.*, 409 (1987) 29-41.
- 475 Weidlich, U., Roehm, H.J. and Gmehling, J.: Measurement of γ using GLC. 2. Results for the stationary phases *N*-formylmorpholine and *N*-methylpyrrolidone. *J. Chem. Eng. Data*, 32 (1987) 450-453; *C.A.*, 107 (1987) 142394h.
- 476 Zainel, H.A., Bender, A.O., Manook, T. and Talib, E.K.: Solution thermodynamic study of several hydrocarbons in γ -butyrolactone and ethylene glycol mixtures by GLC. *J. Chem. Technol. Biotechnol.*, 37 (1987) 229-232; *C.A.*, 107 (1987) 142418u.

See also 448, 508, 525, 585, 591, 762, 766, 819, 822, 823, 825, 831, 833.

3. GENERAL TECHNIQUES

3a. Apparatus and accessories

- 477 Boev, A.I. and Klok, L.N.: (Sample metering pump for a gas chromatograph). *U.S.S.R. Pat.* SU 1,308,893 (Cl. G01N30/24), 7 May 1987, Appl. 3,915,311, 25 Jun. 1985; *C.A.*, 107 (1987) 168058b.
- 478 Deripasko, N.I., Pedchenko, V.N., Parfenenkova, L.R., Ukhabin, M.M., Lipavskii, V.N., Sakodynskii, K.I., Pavlushkov, G.G., Khomyakov, V.P. and Strizhkov, V.N.: (Carrier-gas humidifier for gas chromatographic analysis of trace contaminants of moisture). *U.S.S.R. Pat.* SU 1,330,553 (Cl.G01N30/06), 15 Aug. 1987, Appl. 3,973,147, 4 Nov. 1985; *C.A.*, 107 (1987) 228132h.
- 479 Gong Jiang and Li Haitao: (Changes on the 102-G gas chromatograph and its combination with a microreactor). *Hunan Shifan Daxue Ziran Kexue Xuebao*, 10, No. 1 (1987). 50-54; *C.A.*, 107 (1987) 136284f.
- 480 Herraiz, M., Roglero, G., Loyola, E. and Herraiz, T.: Sampling of volatile components using a PTV in the solvent split mode. *J. High Resolut. Chromatogr. Chromatogr. Commun.*, 10 (1987) 598-602.
- 481 Hinshaw, J.: GC troubleshooting. Inlets. *LC,GC*, 5 (1987) 954-960.
- 482 Hopper, M.L.: Modified inlet for injecting a bonded phase wide-bore fused silica capillary column with chlorinated pesticides in a lipid extract. *J. High Resolut. Chromatogr. Chromatogr. Commun.*, 10 (1987) 620-622.
- 483 Lapin, V.A., Musin, A.R. and Isaev, A.G.: (Gas chromatograph). *U.S.S.R. Pat.* SU 1,323,947 (Cl.G01N30/00), 15 Jul. 1987, Appl. 4,025,716, 21 Feb. 1986; *C.A.*, 107 (1987) 211297t.
- 484 Luu, D.: Thermal desorption and chemical modulators for gas chromatography. Available *Diss. Abstr. Int. B*, 48 (1987) 115; *Univ. Microfilms*, Order No. DA8709265 (1986) 136 pp.; *C.A.*, 107 (1987) 142072b.
- 485 Nishikawa, Y. and Shiomi, K.: (Capillary columns for analysis of basic compounds by gas chromatography). *Jpn. Kokai Tokkyo Koho Pat.* JP 62,168,048 (87,168,048) (Cl. G01N30/60), 25 Jul. 1987, Appl. 86/8,658, 17 Jan. 1986; 3 pp.; *C.A.*, 107 (1987) 211301q.
- 486 Oka, S., Tawara, O. and Kobayashi, J.: (Apparatus for measuring water content in liquid). *Jpn. Kokai Tokkyo Koho Pat.* JP 62,167,474 (87,167,474) G01N30/88), 23 Jul. 1987, Appl. 85/299,553, 28 Dec. 1985; 6 pp., *C.A.*, 107 (1987) 178877g.
- 487 Pankow, J.F. and Isabelle, L.M.: Fused silica open-split GC/MS interface. *J. High Resolut. Chromatogr. Chromatogr. Commun.*, 10 (1987) 617-619.

- 488 Popl, M., Voznakova, Z. and Sandera, J.: (Chromatographic gas-mixture analyzer). *Czech. Pat. CS 231,397* (Cl.G01N31/08), 1 Mar. 1987, Appl. 83/4,558, 21 Jun. 1983; 8 pp.; *C.A.*, 107 (1987) 190128w.
- 489 Schaller, H.-G.: Säulenschalten in der Kapillar-GC. *LaborPraxis*, 11 (1987) 1426-1431.
- 490 Schilling, B., Grob, K., Pichler, P., Dubs, R. and Brechbühler, B.: Fused-silica ends glass capillaries: widening glass capillary butts by a heated tungsten point for preparing press-fit connections. *J. Chromatogr.*, 435 (1988) 204-209.
- 491 Wright, M.D.: A dual-capillary column system for automated analysis of workplace contaminants by thermal desorption-gas chromatography. *Anal. Proc.*, 24 (1987) 309-311.

See also 551, 588, 781.

3b. Detectors and detection reagents

- 492 Biehl, K.E., Roedern, C. and Tyssen, E.: (Method and apparatus for determining trace gases or vapors in gas mixtures such as air). *Ger. Offen. Pat. DE 3,604,893* (Cl. G01N27/66), 20 Aug. 1987, Appl. 15 Feb. 1986; 10 pp.; *C.A.*, 107 (1987) 222456v.
- 493 Bruno, T.J.: Catalytic cracking as the basis a potential detector for gas chromatography. *J. Res. Natl. Bur. Stand. (U.S.)*, 92 (1987) 261-265; *C.A.*, 107 (1987) 167995t.
- 494 Hudson, M.K. and Busch, K.W.: Infrared emission from a flame as the basis for chromatographic detection of organic compounds. *Anal. Chem.*, 59 (1987) 2603-2609.
- 495 Jones, C.S. and Grimsrud, E.P.: Mass spectrometric measurements of the negative ions formed in the thermionic ionization detector with nitrogen carrier gas. *J. Chromatogr.*, 409 (1987) 139-146.
- 496 Koeller, J.: (Inert-gas ionization detector for gas chromatography). *Ger (East) Pat. DD 237,222* (Cl. G01N27/66), 2 Jul. 1986, Appl. 276,101, 9 May 1985; 4 pp.; *C.A.*, 107 (1987) 211296s.
- 497 Koponen, J.: (New detector for gas chromatography). *Kem.-Kemi*, 14 (1987) 535-540; *C.A.*, 107 (1987) 178894k - a review with 26 refs.
- 498 Mierzwinski, A. and Witkiewicz, Z.: Piezoelectric detectors coated with liquid-crystal materials. *Talanta*, 34 (1987) 865-871.
- 499 Riviere, B., Mermet, J.M. and Deruaz, D.: Study of a microwave-induced plasma (surfatron) as a detector in capillary-column gas chromatography with reference to pesticides. *J. Anal. At. Spectrom.*, 2 (1987) 705-709; *C.A.*, 107 (1987) 192852b.
- 500 Siu, K.W.M., Berman, S.S. and Aue, W.A.: Correlations and constants in unipolar and bipolar electron-capture detection. *J. Chromatogr.*, 408 (1987) 53-67.
- 501 Tang You Zhi and Aue, W.A.: General photometric detector for gas chromatography. *J. Chromatogr.*, 409 (1987) 125-137.
- 502 Tang You Zhi and Aue, W.A.: Spectra and response linearization of chlorine and bromine compounds in a copper-sensitized flame photometric detector. *J. Chromatogr.*, 408 (1987) 69-79.

See also 557, 571, 641, 784, 913, 928.

3c. Sorbents, carriers, column and layer performance, packing procedures

- 503 Andreev, E.L., Novikov, V.F., Vigdergauz, M.S., Zabirow, N.G. and Valetdinov, R.K.: (Highly polar stationary phase for gas chromatography). *U.S.S.R. Pat. SU 1,315,898* (Cl.G01N30/02), 7 Jun. 1987, Appl. 3,988,625, 16 Oct. 1985; *C.A.*, 107 (1987) 168062y.
- 504 Berezkin, V.G., Gavrichev, V.S. and Malik, A.: High-efficiency fused-silica capillary micro-packed columns in gas chromatography. *J. Liq. Chromatogr.*, 10 (1987) 1707-1726.
- 505 Bulanova, A.V., Vigdergauz, M.S. and Lazareva, L.S.: (Binary liquid stationary phase for gas-liquid chromatography). *U.S.S.R. Pat. SU 1,330,552* (Cl. G01N30/02), 15 Aug. 1987, Appl. 3,984,160, 6 Dec. 1985; *C.A.*, 107 (1987) 228137p.

- 506 Castello, G., D'Amato, G.: Effects of solute polarity on the performance of Porapak type porous polymers. *Chromatographia*, 23 (1987) 839-843.
- 507 Chukwunenye, C. and McAtee, J.L.: Organo-modified smectite for the gas chromatographic separation of various compounds. *J. Chromatogr.*, 410 (1987) 121-128.
- 508 Fernandez-Sanchez, E., Fernandez-Torres, A., Garcia-Dominguez, J.A., Garcia-Munoz, J., Menendez, V., Molera, M.J., Santiuste, J.M. and Pertierra-Rimada, E.: Mixed stationary phases in gas-liquid chromatography. Partition coefficients and retention indices in OV-101 - OV-25, OV-101 - Carbowax 20 M and OV-225 - SP-2340 mixtures. *J. Chromatogr.*, 410 (1987) 13-29.
- 509 Gao Yifei and Zhang Guiqin: (Crosslinked SE-54 flexible glass capillary column). *Fenxi Huaxue*, 15 (1987) 520-523; *C.A.*, 107 (1987) 211218t.
- 510 Horka, M., Janak, K., Kahle, V. and Tesarik, K.: Properties of immobilized Carbowax. *J. High Resolut. Chromatogr. Chromatogr. Commun.*, 10 (1987) 678-679.
- 511 Hoshino, T., Hibi, K. and Soma, M.: Treatment of carriers for chromatography. *Jpn. Kokai Tokkyo Koho Pat.* JP 62,151,754 (87,151,754) (Cl. G01N30/48), 6 Jul. 1987, Appl. 85/295,537, 26 Dec. 1985; 5 pp.; *C.A.*, 107 (1987) 214484n.
- 512 Howery, D.G. and Soroka, J.M.: Target factor analysis of parameters influencing solute-monomeric stationary phase interactions. *J. Chemom.*, 1 (1987) 91-101; *C.A.*, 107 (1987) 205778j.
- 513 Karavanova, T.M., Koemets, L.A., Legotkina, T.F. and Adon'eva, N.N.: (Method of producing silica sorbent for gas chromatography). *U.S.S.R. Pat.* SU 1,329,812 (Cl. B01J20/10), 15 Aug. 1987, Appl. 3,975, 484, 23 Sep. 1985; *C.A.*, 107 (1987) 219811b.
- 514 Kirsh, S.I., Pomicheva, T.N. and Karabanov, N.T.: (Stationary phase for gas chromatography). *U.S.S.R. Pat.* SU 1,278,701 (Cl. G01N30/48), 23 Dec. 1986, Appl. 3,722,514, 5 Apr. 1984; *C.A.*, 107 (1987) 190130r.
- 515 Kolarz, B.N., Luczynski, J., Trochimczuk, A. and Wojaczynska, M.: Gas chromatographic properties of porous polyacrylonitrile and divinylbenzene copolymers. *J. Chromatogr.*, 408 (1987) 308-315.
- 516 Lekova, K.: (Trends in the deactivation of the support surface in gas chromatography). *Izv. Khim.*, 19 (1986) 503-516; *C.A.*, 107 (1987) 146524y.
- 517 Lin Gwo Chung and Chang, C. Allen: Heavy alkylated benzene stationary phase for the gas-liquid chromatographic separation of ethylbenzene, xylenes, styrene, iso- and n-propylbenzene. *J. Chromatogr.*, 409 (1987) 371-376.
- 518 Murata, T.: (Separation and analysis of compounds with high boiling points by gas chromatography using a silicone oil liquid stationary phase). *Jpn. Kokai Tokkyo Koho Pat.* JP 62 52,460 (87 52,460) (Cl. G01N30/48), 7 Mar. 1987, Appl. 85/192,232, 30 Aug. 1985; 5 pp.; *C.A.*, 107 (1987) 228134k.
- 519 Ottenstein, D.M.: Column support materials for use in gas chromatography. *J. Chromatogr. Sci.*, 25 (1987) 536-546 - a review with 77 refs.
- 520 Panse, D.G., Bhale, A., Gumaste, V.K., Mane, M.V., Nikhite, S.M. and Bapat, B.V.: Liquid crystals VI. Comparative study of the supercooling effect of some liquid crystalline stationary phases using gas-liquid chromatography. *J. Chromatogr.*, 411 (1987) 456-460.
- 521 Poole, S.K. and Poole, C.F.: Chemical interactions as a possible limitation on the useful solvent properties of liquid alkylammonium salts. *J. Chromatogr.*, 435 (1988) 17-28.
- 522 Purygin, P.P., Belyaev, N.F., Vigdergauz, M.S., Papchikhin, A.V., Belousova, Z.P., Yakimova, I.A. and Onuchak, V.N.: (Stationary phase for gas chromatography). *U.S.S.R. Pat.* SU 1,310,717 (Cl. G01N30/02), 15 May 1987, Appl. 4,003,116, 6 Dec. 1985; *C.A.*, 107 (1987) 190131s.
- 523 Sakagami, S.: (Preparation of carriers for high-performance gas chromatography). *Jpn. Kokai Tokkyo Koho Pat.* JP 62,124,459 (87,124,459) (Cl. G01N30/48), 5 Jun. 1987, Appl. 85/265,615, 26 Nov. 1985; 4 pp.; *C.A.*, 107 (1987) 157505u.
- 524 Shah, P.A.: Characterization of gas chromatographic and liquid chromatographic stationary phases. Available *Diss. Abstr. Int. B*, 48 (1987) 414-415; *Univ. Microfilms*, Order No. DA8706894 (1986) 128 pp.; *C.A.*, 107 (1987) 146561h.
- 525 Voelkel, A.: Dispersive forces parameters as a measure of the polarity of surfactants used as stationary phase in gas-liquid chromatography. *J. Chromatogr.*, 435 (1988) 29-41.
- 526 Wu Ningsheng and Qiu Aimin: (Evaluation of chromatographic column efficiency). *Zhongguo Kexue Jishu Daxue Xuebao*, 17 (1987) 65-68; *C.A.*, 107 (1987) 190059z.

- 527 Xu Hong, Kang Xiaojuan, Dong Zhenwen and Sun Zhuolian: (Polymerization in the capillary column for the gas chromatographic stationary phase). *Dalian Gongxueyuan Xuebao*, 26, No. 1 (1987) 62; *C.A.*, 107 (1987) 134739j.
- 528 Zelikman, A.M., Fedoseev, D.V., Vnukov, S.P., Davydenkov, A.K., Lipavskii, V.N., Anikin, B.A. and Silakov, Yu.K.: (Method of preparing a carbon adsorbent for gas chromatographic analysis). *U.S.S.R. Pat.* SU 1,325,356 (Cl.G01N30/50), 23 Jul. 1987, Appl. 3,987,602, 9 Dec. 1985; *C.A.*, 107 (1987) 211302r.
- 529 Zheivot, V.I., Bubenshchikova, V.M., Kovaleva, N.V. and Malakhov, V.V.: (Application of graphitized thermal carbon black modified by Carbowax 20M to gas-chromatographic analysis of catalytic reaction products). *Kolloidn. Zh.*, 49 (1987) 563-567; *C.A.*, 107 (1987) 146542c.
- 530 Zhou Liangmo, Huang Mingxian, Wang Guanghua and Zhu Daoqian: (Preparation and study of polar crosslinking PEG 20M capillary column). *Fenxi Huaxue*, 15 (1987) 542-544; *C.A.*, 107 (1987) 211219u.

See also 437, 469, 563, 574, 704, 739.

3d. Quantitative analysis

- 531 Fan Tony Yu Wen: Quantitative determination of severely overlapping gas chromatographic peaks with identical peak shapes. Available *Diss. Abstr. Int. B*, 48 (1987) 411; *Univ. Microfilms*, Order No. DA8711464 (1987) 147 pp.; *C.A.*, 107 (1987) 168026q.
- 532 Lu Zhichun: (Quantitative determination of high-concentration components by gas chromatography. (I).) *Shiyou Huagong*, 16 (1987) 370-375; *C.A.*, 107 (1987) 190104k.
- 533 Wampler, T.P. and Levy, E.J.: Reproducibility in pyrolysis. Recent developments. XX. *J. Anal. Appl. Pyrolysis*, 12 (1987) 75-82; *C.A.*, 107 (1987) 190013e.

See also 480.

3e. Preparative scale chromatography

- 534 Wilson, D.J.: Line shapes in gas chromatography. III. Solute-solute interactions in preparative columns. *Sep. Sci. Technol.*, 22 (1987) 1835-1849.

See also 584.

3f. Programmed temperature, pressure, vapors, gradients

- 535 Chen Bangjie, Guo Xijian and Peng Shaoyi: (The calculation of retention indices in temperature-programmed capillary gas chromatography). *Sepu*, 5 (1987) 335-338.
- 536 Wang Tiansong and Sun Yiliang: On the influence of the solute sample size on temperature-programmed retention indices. *J. High Resolut. Chromatogr. Chromatogr. Commun.*, 10 (1987) 603-606.

See also 625.

3g. High performance procedures

- 537 Kalmanovskii, V.I., Knyazeva, A.A., Yashin, E.Ya. and Yashin, Ya.I.: (Increase of the information content of the analysis of petroleum fractions by the capillary gas chromatography with a signal differentiation). *Neftekhimiya*, 27 (1987) 585-592.

4. SPECIAL TECHNIQUES

4a. Automation and computerization

- 538 Arnold, M.E. and Passalacqua, P.V.: An optimized PC-based workstation for chromatography data handling. *Am. Lab. (Fairfield)*, 19, No. 9 (1987) 112-119.
- 539 Chretien, J.R.: Chemometrics in chromatography. *TrAC*, 6 (1987) 275-278.

- 540 Crilly, P.B.: Evaluation of Jansson's method for resolving overlapped gas chromatographic peaks. *J. Chemom.*, 1 (1987) 175-183; *C.A.*, 107 (1987) 190069c.
- 541 Crilly, P.B.: Numerical deconvolution of gas chromatograph peaks using Jansson's method. *J. Chemom.*, 1 (1987) 79-90; *C.A.*, 107 (1987) 228112b.
- 542 Fluck, E.R. and Hofmann, J.A.: Development of pseudomultitasking data acquisition network for the chromatography laboratory. *Am. Lab. (Fairfield)*, 19, No. 9 (1987) 34-54.
- 543 Henshall, A. and Schibler, J.A.: Chromatography automation using the Microsoft Windows[®] operating system. *Am. Lab. (Fairfield)*, 19, No. 9 (1987) 42-54.
- 544 Judd, R.: How to specify a data acquisition system for a PC. *Am. Lab. (Fairfield)*, 19, No. 9 (1987) 90-97.
- 545 Kraft, L.: (Computer simulation of a chromatographic separation). *Prax. Naturwiss., Chem.*, 36, No. 6 (1987) 24-25; *C.A.*, 107 (1987) 197048d.
- 546 Sobolev, D.M. and Kozhenkov, A.V.: (Calculation of chromatograms with overlapped peaks). *Zh. Anal. Khim.*, 42 (1987) 1874-1876.

See also 555.

4b. Combination of various chromatographic techniques

- 547 Burke, F.P. and Winschel, R.A.: Correlation of liquid chromatographic retention times with molecular structures of coal liquid components. *Anal. Chem.*, 59 (1987) 2586-2593.
- 548 Ettore, L.S.: Is capillary gas chromatography actually supercritical-fluid chromatography? *J. High Resolut. Chromatogr. Chromatogr. Commun.*, 10 (1987) 637-640.
- 549 Lochmüller, C.H.: The future in chromatography. *J. Chromatogr. Sci.*, 25 (1987) 583-586 - some comments on GC and SFC.
- 550 Middleditch, B.S. and Zlatkis, A.: Artifacts in chromatography: An overview. *J. Chromatogr. Sci.*, 25 (1987) 547-551 - a review with 38 refs., some on GC.
- 551 Raglione, T.V., Troskosky, J.A. and Hartwick, R.A.: On-line microbore high-performance liquid chromatography-capillary gas chromatography-mass spectrometry. I. Development of isotachic eluent splitters. *J. Chromatogr.*, 409 (1987) 205-212.
- 552 Raglione, T.V., Troskosky, J.A. and Hartwick, R.A.: On-line microbore high-performance liquid chromatography-capillary gas chromatography-mass spectrometry. II. Application to the analysis of solvent refined coal. *J. Chromatogr.*, 409 (1987) 213-221.
- 553 Regina, F.J., Boyle, W.J., Jr., Diamond, S.E. and Ladic, I.: Highly sensitive immunoassays with gas-liquid chromatography. *Clin. Chem. (Winston-Salem)*, 33 (1987) 1562-1563.
- 554 Sadek, P.C., Koester, C.J. and Bowers, L.D.: The investigation of the influence of trace metals in chromatographic retention processes. *J. Chromatogr. Sci.*, 25 (1987) 489-493.
- 555 Schmidt, G.J., Dong, M.W. and Salit, M.: A general robotic procedure for preforming precolumn derivatization and its application in on-line liquid chromatographic analysis. *J. Chromatogr. Sci.*, 25 (1987) 453-459.
- 556 Wheeler, J.R. and McNally, M.E.: Comparison of packed column and capillary column supercritical fluid chromatography and high-performance liquid chromatography using representative herbicides and pesticides as typical moderate polarity and molecular weight range molecules. *J. Chromatogr.*, 410 (1987) 343-353.

See also 439, 440, 453, 524, 570, 574, 595, 606, 639, 643, 648, 651, 668, 691, 697, 712, 724, 725, 727, 735, 742, 758, 760, 854, 867, 872, 875, 883, 911.

4c. Combination with other physico-chemical techniques (MS, IR etc.)

- 557 Bilhorn, R.B., Sweedler, J.V., Epperson, P.M. and Denton, M.B.: Charge transfer device detectors for analytical optical spectroscopy - operation and characteristics. *Appl. Spectr.*, 41 (1987) 1114-1125.
- 558 Chen Bangjie and Dou Lingqi: (Application of absorbance subtraction technique to improve resolution in GC/FTIR (gas chromatography/Fourier-transform IR spectrometry)). *Sepu*, 5 (1987) 253-256.

- 559 Eichelberger, J.W. and Budde, W.L.: Studies in mass spectrometry with the ion trap detector. *Biomed. Environ. Mass Spectrom.*, 14 (1987) 357-362; *C.A.*, 107 (1987) 168049z.
- 560 Fields, R.E., III. and White, R.L.: Real-time library search of vapor-phase spectra for gas chromatography/Fourier transform infrared spectrometry. *Anal. Chem.*, 59 (1987) 2709-2716.
- 561 Fujimoto, C., Yoshida, H. and Jinno, K.: Interfacing of inductively coupled plasma-atomic emission spectrometry to super critical fluid chromatography for elemental detection. *J. Chromatogr.*, 411 (1987) 213-220.
- 562 Mehran, M.F. and Cooper, W.J.: Observations on the GC/MS analysis of active solutes: acetylenic thiophenes. *J. High Resolut. Chromatogr. Chromatogr. Commun.*, 10 (1987) 676-677.

See also 605, 633, 657, 700, 701, 705, 734, 738, 752, 817.

4e. Functional analysis

- 563 Antakli, S.C. and Serpinet, J.: Determination of the concentration of silanol groups by a chemical reaction with methylolithium and GC measurements of evolved methane. *Chromatographia*, 23 (1987) 767-769.

See also 555, 561, 571, 685, 784, 928.

4f. Trace analysis and preseparation techniques

- 564 Dix, K.D. and Fritz, J.S.: Simple steam distillation for sample preparation prior to gas chromatographic determination of organic compounds. *J. Chromatogr.*, 408 (1987) 201-210.
- 565 Fabbri, A., Crescentini, G., Mangani, F., Mastrogiacomo, A.R. and Brunner, F.: Advances in the determination of volatile organic solvents and other organic pollutants by gas chromatography with thermal desorption sampling and injection. *Chromatographia*, 23 (1987) 856-860.
- 566 Hu Ye and Zou Jianyun: (Study on the release rate of microencapsulated fragrances by gas chromatography). *Sepu*, 5 (1987) 240-243.
- 567 Kawamoto, K. and Urano, K.: (A sampling method for monitoring organic halogen compounds in air). *Nippon Kagaku Kaishi*, (1987) 1746-1752; *C.A.*, 107 (1987) 160558u.
- 568 Kok, M.F., Yong, F.M. and Lim, G.: Rapid extraction method for reproducible analysis of aroma volatiles. *J. Agric. Food Chem.*, 35 (1987) 779-781.
- 569 Newman, K.A. and Gschwend, P.M.: (A method for quantitative determination of volatile organic compounds in marine macroalgae). *Limnol. Oceanogr.*, 32 (1987) 702-708; *C.A.*, 107 (1987) 171924m.
- 570 Petrick, G., Schulz, D.E. and Duinker, J.C.: Clean-up of environmental samples by high-performance liquid chromatography for analysis of organochlorine compounds by gas chromatography with electron-capture detection. *J. Chromatogr.*, 435 (1988) 241-248.
- 571 Poole, C.F. and Poole, S.K.: Derivatization as an approach to trace analysis by gas chromatography with electron-capture detection. *J. Chromatogr. Sci.*, 25 (1987) 434-443 - a review with 83 refs.
- 572 Pretorius, V. and Lawson, K.H.: Analysis of trace components of vapor samples using the static solvent effect and capillary gas-liquid chromatography. *S. Afr. J. Chem.*, 40 (1987) 169-171; *C.A.*, 107 (1987) 190074a.

See also 491, 492, 603, 780, 802, 849, 892, 896, 904, 906, 920, 927, 928, 934.

4g. Separation of enantiomers

- 573 Gyllenhaal, O., Lamm, B. and Vessman, J.: Enantiomer separation of tocinide and some of its analogues and derivatives on a Chirasil-Val capillary column. *J. Chromatogr.*, 411 (1987) 285-295.
- 574 Röder, W., Ruffing, F.-J., Schomburg, G. and Pirkle, W.H.: Chiral SFC-separations using polymer-coated open tubular fused silica columns. Comparison of enantiomers selectivity in SFC and LC using the same stationary phase of the Pirkle type. *J. High Resolut. Chromatogr. Chromatogr. Commun.*, 10 (1987) 665-667.

- 575 Schmidt, N., Gercken, G. and König, W.A.: Enantioselective capillary gas chromatography of 1,2-isopropylidene glycerol and O-alkylated glycerol derivatives. *J. Chromatogr.*, 410 (1987) 458-462.
- 576 Wang-Iverson, D., Jemal, M. and Cohen, A.I.: Determination of enantiomeric purity of 3-mercapto-2-methylpropionic acid. *J. Chromatogr.*, 408 (1987) 365-371.
- 577 Watabe, K., Chang, S.C., Gil-Av, E. and Koppenhoefer, B.: The determination of small amounts of enantiomeric impurities in α -halo carboxylic acids. *Synthesis*, (1987) 225-228; *C.A.*, 107 (1987) 190099n.

4h. Other special techniques

- 578 Berezkin, V.G., Zakirov, N.S., Lipavskii, V.N. and Koshevnik, M.A.: (Thin-layer gas chromatography). *Dokl. Akad. Nauk SSSR*, 294 (1987) 1144-1146.
- 579 Chen Yiwen, Li Qinghong, Yang Meiyang and Peng Xiahui: (Determination of lead by homogeneous catalysis and gas chromatography). *Fenxi Huaxue*, 15 (1987) 354-356; *C.A.*, 107 (1987) 204791w.
- 580 Fritsch, H.: Pyrolyse-GC bringt wessentliche Arbeitserleichterung. *LaborPraxis*, 11 (1987) 1092-1098.

4i. Supercritical fluid chromatography

- 581 Bartle, K.D., Barker, I.K., Clifford, A.A., Kithinji, J.P., Raynor, M.W. and Shilstone, G.F.: Supercritical fluid chromatography. *Anal. Proc.*, 24 (1987) 299-301.
- 582 Goates, S.R., Zabriskie, N.A., Simons, J.K. and Khoobehi, B.: Detection of aerosol formation in the effluent of a supercritical fluid chromatography. *Anal. Chem.*, 59 (1987) 2927-2930.
- 583 Huau, J. and Tinet, C.: (Supercritical-phase chromatography. New developments). *Inf. Chim.*, 285 (1987) 151-154; *C.A.*, 107 (1987) 190008g - a review with 6 refs.
- 584 Jusforques, P., Berger, C. and Perrut, M.: (New separation process: preparative supercritical liquid chromatography). *Chem.-Ing.-Tech.*, 59 (1987) 666-667; *C.A.*, 107 (1987) 178933x.
- 585 Kopner, A., Hamm, A., Ellert, J., Feist, R. and Schneider, G.M.: Determination of binary diffusion coefficients in supercritical chlorotrifluoromethane and sulfur hexafluoride with supercritical fluid chromatography (SFC). *Chem. Eng. Sci.*, 42 (1987) 2213-2218; *C.A.*, 107 (1987) 178974m.
- 586 Kuei, J.C.H.: Capillary supercritical fluid chromatography with ammonia-based mobile phases. Available *Diss. Abstr. Int. B*, 48 (1987) 784; *Univ. Microfilms*, Order. No. DA8712620 (1987) 106 pp.; *C.A.*, 107 (1987) 178969p.
- 587 Leaseburge, E.J. and Melda, K.J.: Sample dilution system for supercritical fluid chromatography. *U.S. Pat.* US 4,681,678 (Cl.210-101; B01D15/08), 21 Jul. 1987, Appl. 917,628, 10 Oct. 1986; 6 pp.; *C.A.*, 107 (1987) 146585u.
- 588 Leaseburge, E.J. and Thomas, T.J.: Supercritical fluid chromatograph with pneumatically controlled pump. *U.S. Pat.* US 4,684,465 (Cl.210-198.2; B01D15/08), 4 Aug. 1987, Appl. 917,466, 10 Oct. 1986; 6 pp.; *C.A.*, 107 (1987) 136553t.
- 589 Lee, C. and Ellington, R.T.: Density-based correlation for solid solubility in supercritical solvents. *Separ. Sci. Technol.*, 22 (1987) 1557-1576.
- 590 Leyendecker, D., Leyendecker, D., Schmitz, F.P. and Klesper, E.: Comparison of eluents in supercritical fluid chromatography. *J. Liq. Chromatogr.*, 10 (1987) 1917-1947; *C.A.*, 107 (1987) 211268j.
- 591 Lochmüller, C.H. and Mink, L.P.: Adsorption isotherms of ethyl acetate modifier on silica from supercritical carbon dioxide. *J. Chromatogr.*, 409 (1987) 55-60.
- 592 Sanagi, M.M. and Smith, R.W.: Applications of a packed column supercritical fluid chromatograph with flame ionization detector. *Anal. Proc.*, 24 (1987) 304-306.
- 593 Schmitz, F.P.: Comparison of effluents in supercritical fluid chromatography as based on density and free volume. *J. High Resolut. Chromatogr. Chromatogr. Commun.*, 10 (1987) 650-653.
- 594 Schmitz, F.P.: Ernst Klesper and the beginnings of supercritical fluid chromatography. *J. High Resolut. Chromatogr. Chromatogr. Commun.*, 10 (1987) 636.

- 595 Schneiderman, M.A., Sharma, A.K. and Locke, D.C.: Determination of anthraquinone in paper and wood using supercritical fluid extraction and high-performance liquid chromatography with electrochemical detection. *J. Chromatogr.*, 409 (1987) 343-353.
- 596 Schwartz, H.E., Barthel, P.J. and Moring, S.E.: Pump filling techniques in SFC: Do they affect precision? *J. High Resolut. Chromatogr. Chromatogr. Commun.*, 10 (1987) 668-669.

See also 548, 556, 561, 574, 811.

5. HYDROCARBONS AND HALOGEN DERIVATIVES

5a. Aliphatic hydrocarbons

- 597 Bauer, M.: (The chemical-toxicological analysis of certain paraffin - hydrocarbons in autopsy materials). *Arch. Med. Sadowej Kryminol.*, 37 (1987) 14-16; *C.A.*, 107 (1987) 213083u.
- 598 Boneva, S.: (Comparison of retention index calculation methods of alkenes on squalane). *Izv. Khim.*, 19 (1986) 460-464; *C.A.*, 107 (1987) 146523x.
- 599 Dimov, N. and Boneva, S.: (An equation for calculation of gas chromatographic retention of alkenes on dimethylsilicone OV-101). *Izv. Khim.*, 19 (1986) 456-459; *C.A.*, 107 (1987) 146539g.
- 600 Giles, J.J.: The analysis of waxes and greases using high resolution gas chromatography. *J. Forensic Sci. Soc.*, 27 (1987) 231-239; *C.A.*, 107 (1987) 213094y.
- 601 Reed, L.E.: A rapid method for the analysis of mixtures of hydrocarbons and inorganic gases. *J. Chromatogr. Sci.*, 25 (1987) 485-488.
- 602 Shiomi, K. and Takahashi, S.: Capillary gas chromatography of saturated hydrocarbons in gasoline by removing unsaturated components by precolumn sulfonation. *Anal. Sci.*, 3 (1987) 379-383; *C.A.*, 107 (1987) 201626k.
- 603 Takayama, Y., Yamaguchi, T. and Kawai, S.: (Determination of trace amounts of n-alkanes by GC with precolumn concentration technique). *Bunseki Kagaku*, 36 (1987) 761-765.
- 604 Zinkovskaya, Z.I. and Rybakova, L.A.: (Chromatographic determination of dicyclopentadiene in the presence of gasoline vapors in the air of industrial facilities). *Neftepererab. Neftekhim. (Moscow)*, No. 8 (1987) 29-30; *C.A.*, 107 (1987) 182476t.

5b. Cyclic hydrocarbons

- 605 Buchanan, M.V. and Wise, M.B.: (Negative ion processes for the unambiguous identification of polycyclic aromatic compounds). *Fuel*, 66 (1987) 954-959; *C.A.*, 107 (1987) 220236z.
- 606 Davies, I.L., Reynor, M.W., Williams, P.T., Andrews, G.E. and Bartle, K.D.: Application of automated on-line microbore high-performance liquid chromatography/capillary gas chromatography to Diesel exhaust particulates. *Anal. Chem.*, 59 (1987) 2579-2583.
- 607 Lee, H.B., Szawiola, R. and Chau, A.S.Y.: Solvent effects on response factors for polynuclear aromatic hydrocarbons determined by capillary gas chromatography using splitless injections. *J. Assoc. Off. Anal. Chem.*, 70 (1987) 929-930.
- 608 Succar, S.D., Roca, A.E. and Catoggio, J.A.: Detection, identification and evaluation (by GLC and HPLC) of the first terms of the polynuclear aromatic hydrocarbons' family in industrial by-products used as raw matter due to their high contents in benzene. *Rev. Latinoam. Ing. Quim. Quim. Apl.*, 17 (1987) 31-40; *C.A.*, 107 (1987) 157918f.
- 609 Topalova, I., Chanev, Kh., Petsev, N. and Dimitrov, Kh.: (Structure-retention correlation of some alkenylbenzenes using gas-solid chromatography). *Izv. Khim.*, 19 (1986) 444-450; *C.A.*, 107 (1987) 146538f.
- 610 Yang Zedong and Cai Zhengying: (Concentration of trace dimethylbenzene in water with microextractant and sampling of Curie point evaporation for gas chromatographic analysis). *Fenxi Huaxue*, 14, No. 1 (1986) 13-15; *C.A.*, 107 (1987) 140743x.

See also 517, 547, 914.

5c. Halogen derivatives

- 611 Benchev, I., Rizov, N. and Kolarska, A.: (Determination of volatile toxic substances in biological products by gas chromatography-head space analysis. III. Determination of trichloroethylene and tetrachloroethylene (perchloroethylene) in blood). *Khig. Zdraveopaz.*, 30, No. 2 (1987) 77-79; *C.A.*, 107 (1987) 170044n.
- 612 Gabrio, T. and Galetzka, G.: (Gas-chromatographic testing of halothane purity). *Pharmazie*, 42 (1987) 306-307; *C.A.*, 107 (1987) 183670p.
- 613 Suzuki, S., Nagano, S. and Satoh, S.: (Measurement method for chlorobenzenes in air). *Kawasaki-shi Kogai Kenkyusho Nempo*, No. 13 (1986) 14-23; *C.A.*, 107 (1987) 140025q.
- 614 Zhang Lin: (Quantitative studies on chromatographic peaks of monobromination products of *o*-chlorotoluene). *Hunan Shifan Daxue Ziran Kexue Xuebao*, 10 (1987) 57-61; *C.A.*, 107 (1987) 211272f.

See also 565, 567, 798, 815, 904, 909, 910.

5d. Complex hydrocarbon mixtures

- 615 Albaiges, J. and Grimalt, J.: A quality assurance study for the analysis of hydrocarbons in sediments. *Int. J. Environ. Anal. Chem.*, 31 (1987) 281-293.
- 616 Wang Yujie and Zhao Qingyue: (Gas chromatographic analysis of propylene oxidative dehydrogenation and aromatization reoxygenation products). *Fenxi Huaxue*, 15 (1987) 189-190; *C.A.*, 107 (1987) 168038v.

See also 537, 552, 818, 905, 921, 922.

6. ALCOHOLS

- 617 Del Grosso, A.V. and May, J.C.: Gas chromatographic liquid chromatographic, and titrimetric procedures for determination of glycerin in allergenic extracts and diagnostic antigens: comparative study. *J. Assoc. Off. Anal. Chem.*, 70 (1987) 825-828.
- 618 Fenton, M. and Aherne, F.X.: Determination of glycerol in pig plasma by capillary gas chromatography. *J. Chromatogr.*, 410 (1987) 480-483.
- 619 Molever, K.: Quantitative determination of glycerin in soap by capillary GC. *J. Am. Oil Chem. Soc.*, 64 (1987) 1356-1357.
- 620 Penton, Z.: Gas-chromatographic determination of ethanol in blood with 0.53-mm fused-silica open tubular columns. *Clin. Chem. (Winston-Salem)*, 33 (1987) 2094-2095.
- 621 Prochazka, L. and Culik, J.: (Alcohol analysis using gas chromatography). *Kvasny Prum.*, 33 (1987) 7-10; *C.A.*, 107 (1987) 152734h.
- 622 Profumo, A., Riolo, C., Pesavento, M. and Francoli, A.: Direct gas-chromatographic determination of methanol in alcoholic beverages. *Ann. Chim. (Rome)*, 77 (1987) 509-514; *C.A.*, 107 (1987) 132492z.
- 623 Xu Shuying: (Determination of higher alcohols from carbon monoxide + hydrogen by GC/MS). *Ranliao Huaxue Xuebao*, 15 (1987) 84-89; *C.A.*, 107 (1987) 178554z.

See also 575, 640, 676, 695, 901, 921.

7. PHENOLS

- 624 Jobst, H.: Bestimmung von Isomeren 4-Nonylphenolen in Böden. *Fresenius' Z. Anal. Chem.*, 328 (1987) 644-647.
- 625 Masi, O.H. and Gulick, W.M., Jr.: An optimized gas chromatographic determination of priority pollutant phenols. *J. High Resolut. Chromatogr. Chromatogr. Commun.*, 10 (1987) 647-649.

8. SUBSTANCES CONTAINING HETEROCYCLIC OXYGEN

8a. Flavonoids

- 626 Fritz, J.O. and Moore, K.J.: Separation and quantification of lignin-derived phenolic monomers using high-resolution gas chromatography. *J. Agric. Food Chem.*, 35 (1987) 710-713.

8b. Aflatoxins and other mycotoxins

- 627 Sugimoto, T., Yamamoto, K. and Minamisawa, M.: (Efficient method for the determination of deoxynivalenol and nivalenol in wheat and barley by gas chromatography). *Shokuhin Eiseigaku Zasshi*, 28 (1987) 169-174; *C.A.*, 107 (1987) 152997w.
- 628 Sydenham, E.W. and Thiel, P.G.: The simultaneous determination of diacetoxyscirpenol and T-2 toxin in fungal cultures and grain samples by capillary gas chromatography. *Food Addit. Contam.*, 4 (1987) 277-284; *C.A.*, 107 (1987) 152995u.

8c. Other compounds with heterocyclic oxygen (including tannins)

- 629 Donnelly, J.R., Munslow, W.D., Vonnahme, T.L., Nunn, N.J., Hedin, C.M., Sovocool, G.W. and Mitchum, R.K.: The chemistry and mass spectrometry of brominated dibenzo-*p*-dioxins and dibenzofurans. *Biomed. Environ. Mass Spectrom.*, 14 (1987) 465-472; *C.A.*, 107 (1987) 190097k.
- 630 Grainger, J. and Gelbaum, L.T.: Tetrachlorodibenzo-*p*-dioxin isomer differentiation by capillary gas chromatography Fourier transform-infrared spectroscopy. *Appl. Spectrosc.*, 41 (1987) 809-820; *C.A.*, 107 (1987) 146562j.
- 631 Kuehl, D.W., Butterworth, B.C., DeVita, W.M. and Sauer, C.P.: Environmental contamination by polychlorinated dibenzo-*p*-dioxins and dibenzofurans associated with pulp and paper mill discharge. *Biomed. Environ. Mass Spectrom.*, 14 (1987) 443-447; *C.A.*, 107 (1987) 211274h.
- 632 Munder, A., Buchert, H., Niemczyk, R., Swerev, M. and Ballschmitter, K.: Determination of PCDD/PCDF in motor oil and air particulates. A modular approach in clean-up. *Fresenius' Z. Anal. Chem.*, 328 (1987) 639-643.
- 633 Tondeur, Y., Niederhut, W.N., Campana, J.E. and Missler, S.R.: A hybrid HRGC/MS/MS method for the characterization of tetrachlorinated-*p*-dioxins in environmental samples. *Biomed. Environ. Mass Spectrom.*, 14 (1987) 449-456; *C.A.*, 107 (1987) 168050t.
- 634 Waddell, D.S., McKinnon, H.S., Chittim, B.G., Safe, S. and Boyd, R.K.: Gas chromatographic/mass spectrometric characteristics of purified synthetic isomers of tetrachlorodibenzofuran. *Biomed. Environ. Mass Spectrom.*, 14 (1987) 457-464; *C.A.*, 107 (1987) 168022k.
- 635 Yasuhara, A., Ito, H. and Morita, M.: Isomer-specific determination of polychlorinated dibenzo-*p*-dioxins and dibenzofurans in incinerator-related environmental samples. *Environ. Sci. Technol.*, 21 (1987) 971-979; *C.A.*, 107 (1987) 146579v.

See also 799, 861.

9. OXO COMPOUNDS, ETHERS, EPOXIDES AND QUINONES

- 636 Boneva, S. and Dimov, N.: Gas chromatographic separation of C₄-C₅ epoxides on a glass capillary column coated with OV-101 methylsilicone. *Chromatographia*, 23 (1987) 770-772 - retention data for 10 epoxides.
- 637 Liu Yuhua and Yin Zhigang: (Gas-liquid chromatographic determination of synthetic 1,11-dichloro-3,6,9-trioxadecane). *Sepu*, 5 (1987) 262-263.
- 638 Ono, A.: Thermal decomposition and dehalogenation of α -halo ketones by GLC. *Thermochim. Acta*, 104 (1986) 373-376; *C.A.*, 107 (1987) 175591y.
- 639 Sagara, K., Oshima, T., Yoshida, T., Tong Yu Yi, Zhang Guande and Chen Yu Heng: Determination of Cinnamomi Cortex by high-performance liquid chromatography. *J. Chromatogr.*, 409 (1987) 365-370.

- 640 Szewczyk, H., Dziwinski, E. and Szymanowski, J.: Analysis and identification of some aminoether alcohols and their ethers. *J. Chromatogr.*, 410 (1987) 447-452.
- 641 Tang You Zhi and Aue, W.A.: Gas chromatographic determination of benzaldehyde by gas-phase luminescence. *J. Chromatogr.*, 409 (1987) 243-250.

See also 595, 876, 912.

10. CARBOHYDRATES

10a. Mono and oligosaccharides. Structural studies

- 642 Biodi, P.A., Manca, F., Negri, A., Secchi, C. and Montana, M.: Gas chromatographic analysis of neutral monosaccharides as their O-pentafluorobenzoyloxime acetates. *J. Chromatogr.*, 411 (1987) 275-284.
- 643 Lehrfeld, J.: Simultaneous gas-liquid chromatographic determination of aldoses and alduronic acids. *J. Chromatogr.*, 408 (1987) 245-253.
- 644 Li Han Ping: Capillary gas chromatography of neutral sugars as their aldonitrile acetates from the hydrolyzate of corn bran residues. *J. Chromatogr.*, 410 (1987) 484-487.
- 645 Malmberg, A. and Theander, O.: Analysis of sucrose content in beet crops. *Swed. J. Agric. Res.*, 17 (1987) 3-7; *C.A.*, 107 (1987) 216250g.
- 646 Mateo, R., Bosch, F., Pastor, A. and Jimenez, M.: Capillary column gas chromatographic identification of sugars in honey as trimethylsilyl derivatives. *J. Chromatogr.*, 410 (1987) 319-328.
- 647 Minor, J.L. and Petterson, R.C.: Mass spectrometric method to determine the chain length of oligosaccharides attached to phenolic polymers by nonglycosidic linkages. *J. Agric. Food Chem.*, 35 (1987) 993-996.

See also 673.

11. ORGANIC ACIDS AND LIPIDS

11a. Organic acids and simple esters

- 648 Abd El Aal, M.H., Gomaa, E.G. and Karara, H.A.: Bitter almond, plum and mango kernels as sources of lipids. *Fett Wiss. Technol.-Fat Sci. Technol.*, 89 (1987) 304-306.
- 649 Ahmed, F.A., Ghali, Y., Osman, O. and Ali, M.S.: Effect of some herbicides on lipid composition and agronomic characters of corn grains. *Grasas Aceites*, 38 (1987) 149-153.
- 650 Audisio, G., Rossini, A., Bianchi, G. and Avato, P.: CGC-MS determination of mixtures of long chain aliphatic esters. *J. High Resolut. Chromatogr. Chromatogr. Commun.*, 10 (1987) 594-597.
- 651 Boley, N.P. and Colwell, R.K.: Comparison of methods for the determination of butyric acid in foodstuffs by gas-liquid chromatography and high-performance liquid chromatography. *J. Chromatogr.*, 410 (1987) 190-194.
- 652 Calorini, L., Fallani, A., Tombaccini, D., Mugnai, G. and Ruggieri, S.: Lipid composition of cultured B 16 melanoma cell variants with different lung-colonizing potential. *Lipids*, 22 (1987) 651-656.
- 653 Canonica, F.P.: Gas-liquid chromatographic analysis of fatty acid methyl esters of *Aeromonas hydrophila*, *Aeromonas sobria* and *Aeromonas caviae*. Available *Diss. Abstr. Int. B*, 48 (1987) 45; *Univ. Microfilms*, Order No. DA8709619 (1987) 93 pp.; *C.A.*, 107 (1987) 130605q.
- 654 Carballeira, N.M., Maldonado, L. and Porras, B.: Isoprenoid fatty acids from marine sponges. Are sponges selective? *Lipids*, 22 (1987) 767-769.
- 655 Chen Yiwen, Li Qingheng, He Ling and Zheng Liqin: (Rapid gas chromatographic determination of sorbic and benzoic acids in food). *Sepu*, 5 (1987) 238-240.
- 656 Chirouze, V., Entressangles, B. and Helme, J.-P.: Le placenta humain comme source d'acides gras polyinsaturés à chaînes longues. I. Composition lipidique du placenta et potentialités en acide arachidonique. *Rev. Fr. Corps Gras*, 34 (1987) 275-280.

- 657 Christie, W.W., Brechany, E.Y., Gunstone, F.D., Lie Ken Jie, M.S.F. and Holman, R.T.: Mass spectra of the picolinyl ester derivatives of some non-methylene-interrupted octadecadienoic acids. *Lipids*, 22 (1987) 664-666.
- 658 Cillard, J. and Cillard, P.: Antioxidant activity of associated alpha-tocopherol and ascorbic acid in aqueous media. *Rev. Fr. Corps Gras*, 34 (1987) 271-274.
- 659 Cohen, J.D., Bausher, M.G., Bialek, K., Buta, J.G., Gocal, G.F.W., Janzen, L.M., Pharis, R.P., Reed, A.N. and Slovin, J.P.: Comparison of commercial ELISA assay for indole-3-acetic acid at several stages of purification and analysis by gas chromatography-selected ion monitoring-mass spectrometry using a $^{13}\text{C}_6$ -labeled internal standard. *Plant Physiol.*, 84 (1987) 982-986; *C.A.*, 107 (1987) 130477z.
- 660 Craske, J.D. and Bannon, C.D.: Gas liquid chromatography analysis of the fatty acid composition of fats and oils: A total system for high accuracy. *J. Am. Oil Chem. Soc.*, 64 (1987) 1413-1417.
- 661 Croft, K.D., Beilin, L.J., Legge, F.M. and Vandongen, R.: Effects of diets enriched in eicospentaenoic or docosahexaenoic acids on prostanoid metabolism in the rat. *Lipids*, 22 (1987) 647-650.
- 662 Daulatabad, C.D., Hosamani, K.M., Desai, V.A. and Alagawadi, K.R.: Cyclopropenoid fatty acids in *Leguminosae* oils. *J. Am. Oil Chem. Soc.*, 64 (1987) 1423 only.
- 663 Dave, G.R., Patel, M.J., Patel, R.M. and Patel, R.J.: Characteristics and composition of seeds and oil of *Diplocyclos palmatus* L., C. Jaffery from Gujarat, India. *Fett Wiss. Technol.-Fat. Sci. Technol.*, 89 (1987) 331.
- 664 Fuerst, P., Krueger, C., Habersaat, K. and Groebel, W.: (Halogenated carboxylic acids in beverages - gas chromatographic determination and confirmation by gas chromatography/mass spectrometry with negative chemical ionization). *Z. Lebensm.-Unters. Forsch.*, 185 (1987) 17-20; *C.A.*, 107 (1987) 132490x.
- 665 Fujii, R., Arimoto, K. and Zinkel, D.F.: Dimeric components from the dimerization of abietic acid. *J. Am. Oil Chem. Soc.*, 64 (1987) 1144-1146.
- 666 Fussell, R.J. and McCalley, D.V.: Determination of volatile fatty acid (C_2 - C_5) and lactic acid in silage by gas chromatography. *Analyst (London)*, 112 (1987) 1213-1216.
- 667 Gangadhar, A., Subbarao, R. and Lakshminarayana, G.: Synthesis of 18-octadecanolide, a major constituent of Dufour's gland secretion of *Colletes* bees. *J. Am. Oil Chem. Soc.*, 64 (1987) 1135-1137.
- 668 Gertz, C.: Bestimmung von *cis,cis*-9,12-Octadecadiensäure in Lebensmitteln. *Fett Wiss. Technol.-Fat Sci. Technol.*, 89 (1987) 320-324.
- 669 Grandgirard, A., Julliard, F., Prevost, J. and Sebedio, L.: Preparation of geometrical isomers of linolenic acid. *J. Am. Oil Chem. Soc.*, 64 (1987) 1434-1440.
- 670 Haslbeck, F. and Grosch, W.: (Studies on fat determination by lipolysis. III. Quantification of free fatty acids). *Z. Lebensm.-Unters. Forsch.*, 185 (1987) 31-35; *C.A.*, 107 (1987) 132738j.
- 671 Hiyama, K.: (Analysis of volatile fatty acids and lactic acid in silage by gas chromatography). *Chikusan no Kenkyu*, 40 (1986) 1410-1412; *C.A.*, 107 (1987) 174513n.
- 672 Hoverstad, T. and Midtvedt, T.: Short-chain fatty acids in seven sterile thioglycollate media. *Acta Pathol., Microbiol. Immunol. Scand., Sect. B*, 95B (1987) 211-212; *C.A.*, 107 (1987) 130195f.
- 673 Hughes, W.J. and Thorpe, T.M.: Determination of organic acids and sucrose in roasted coffee by capillary gas chromatography. *J. Food Sci.*, 52 (1987) 1078-1083; *C.A.*, 107 (1987) 132764q.
- 674 Hu Manling, Huang Songling and Xiao Bo: (Determination of magnesium valproate in serum by gas chromatography). *Sepu*, 5 (1987) 362-364.
- 675 Joshi, S. and Mathur, J.M.S.: Effect of growth temperature on the physico-chemical characteristics of fungal oil. *Fett Wiss. Technol.-Fat Sci. Technol.*, 89 (1987) 311-313.
- 676 Jyoyama, Y., Michitsuji, H., Ohara, A., Yamaguchi, K. and Fujiki, Y.: (On determination of trichloroacetic acid (TCA) and trichloro-ethanol (TCE) by headspace gas chromatography). *Shojinkai Igakushi*, 25 (1986) 225-231; *C.A.*, 107 (1987) 170028k.
- 677 Katagiri, M., Shimizu, S., Kaihara, H. and Katagiri, C.: (Studies on the components of miso. III. Determination of esterified and free carboxylic acids in natto by gas chromatography). *Nippon Shokuhin Kogyo Gakkaishi*, 34 (1987) 305-308; *C.A.*, 107 (1987) 174685v.

- 678 Kobayashi, K., Suginaka, H. and Yano, I.: Analysis of fatty acid composition of *Candida* species by gas-liquid chromatography using a polar column. *Microbios*, 51 (1987) 37-42; *C.A.*, 107 (1987) 130601k.
- 679 Lamkin, W.M., Unruh, N.C. and Pomeranz, Y.: Gas chromatographic determination of calcium propionate added as preservative to bread. *J. Assoc. Off. Anal. Chem.*, 70 (1987) 763-767.
- 680 Lee, T.W.: Quantitative determination of linoleic acid in infant formulas by gas chromatography. *J. Assoc. Off. Anal. Chem.*, 70 (1987) 702-705.
- 681 Liu Yongxing and Zhang Shuming: (Analysis of fatty acids in oil shale oxidation products by gas chromatography). *Sepu*, 5 (1987) 384-387.
- 682 Lognay, G., Marlier, M., Severin, M. and Wathelet, J.P.: Etude des lipides du lierre (*Hedera helix* L.). Identification d'acides gras monoéniques isomères. *Rev. Fr. Corps Gras*, 34 (1987) 407-411.
- 683 Losada, A.V. and Peluffo, R.O.: Effect of cold environment on hepatic microsomal Δ^6 and Δ^9 desaturase activity of male rats. *Lipids*, 22 (1987) 583-588.
- 684 Mazumder, C., Kundu, M., Basu, J. and Chakrabarti, P.: Lipid composition and amino acid transport in a nustatin-resistant mutant of *Aspergillus niger*. *Lipid*, 22 (1987) 609-612.
- 685 Molnar-Perl, I. and Morvai, M.: Esterification of aliphatic hydroxy acids to *n*-propyl esters in aqueous solutions for their gas chromatographic analysis. *Chromatographia*, 23 (1987) 760-763.
- 686 Morgan, E.D. and Ollett, D.G.: Methyl 6-methylsalicylate, trail pheromone of the ant *Tetramorium impurum*. *Naturwissenschaften*, 74 (1987) 596-597.
- 687 Murata Takeshi, Chen Zhihang and Zou Liansheng: (Analysis of royal jelly by gas chromatography-mass spectrometry). *Huaxue Tongbao*, No. 5 (1987) 31-32; *C.A.*, 107 (1987) 195198d.
- 688 Mutter, M. and Homan, H.R.: Estimation of amount of *trans,trans*-isomer of octadecadienoic acids in hydrogenated oils and fats. *J. High Resolut. Chromatogr. Chromatogr. Commun.*, 10 (1987) 672-673.
- 689 Nishikawa, H., Sakai, T. and Kidani, Y.: Gas chromatographic determination of benzoic acid and toluic acid in airborne particulates. *Anal. Sci.*, 3 (1987) 463-466; *C.A.*, 107 (1987) 222293q.
- 690 Ozawa, H. and Tsukioka, T.: Gas chromatographic determination of sodium mono-fluoroacetate in water by derivatization with dicyclohexylcarbodiimide. *Anal. Chem.*, 59 (1987) 2914-2917.
- 691 Perkins, E.G. and Smick, C.: Octadecatrenoic fatty acid isomers of partially hydrogenated soybean oil. *J. Am. Oil Chem. Soc.*, 64 (1987) 1150-1155.
- 692 Prasad, R.B.N., Rao, Y.N. and Rao, S.V.: Phospholipids of Palash (*Butea monosperma*), Papaya (*Carica papaya*), Jangli Badam (*Sterculia foetida*), Coriander (*Coriandrum sativum*), and carrot (*Daucus carota*) seeds. *J. Am. Oil Chem. Soc.*, 64 (1987) 1424-1427.
- 693 Qi Fenglan, Han Yingsu and Zhao Jinhai: (Gas chromatographic estimation of free fatty acids of fermentation with direct sample injection). *Sepu*, 5 (1987) 382-384.
- 694 Rao, K.S. and Lakshminarayana, G.: Characteristics and composition of six newer seeds and the oils. *Fett Wiss. Technol.-Fat Sci. Technol.*, 89 (1987) 324-326.
- 695 Rao, Y.R., Pantulu, A.J. and Lakshminarayana, G.: Reduction of fatty acid methyl esters with sodium borohydride-*t*-butanol - methanol. *Fett Wiss. Technol.-Fat Sci. Technol.*, 89 (1987) 398-400.
- 696 Reddy, Y. and Prabhakar, J.V.: Effect of diglycerides on the solidification of sal (*Shorea robusta*) fat. *Fett Wiss. Technol.-Fat Sci. Technol.*, 89 (1987) 394-395.
- 697 Rezanika, T. and Mares, P.: Unusual and very long-chain fatty acids produced by *Basidiomycetes*. *J. Chromatogr.*, 409 (1987) 390-395.
- 698 Rogel, A.M. and Watkins, B.A.: Liver subcellular fatty profiles of chicks fed diets containing hydrogenated fats and varying linoleate levels. *Lipids*, 22 (1987) 637-642.
- 699 Rutledge, D.N., Phan, P.A. and Dubacq, J.P.: Methode rapide de dosage de la cire dans des graines individuelles de jojoba par RMN impulsioneel a bass resolution. Identification de souches a haut rendement. *Rev. Fr. Corps Gras*, 34 (1987) 391-396.
- 700 Schatowitz, B. and Gercken, G.: Benzyl esters of C_2-C_{20} fatty acids and metabolically relevant carboxylic acids. Preparation and gas chromatography-mass spectrometry. *J. Chromatogr.*, 409 (1987) 43-54.

- 701 Sebedio, J.L., Quere, J.L., Semon, E., Morin, O., Prevost, J. and Grandgirard, A.: Heat treatment of vegetable oils. II. GC-MS and GC-FTIR spectra of some isolated cyclic fatty acid monomers. *J. Am. Oil Chem. Soc.*, 64 (1987) 1324-1333.
- 702 Siguel, E.N., Chee, K.M., Gong Junxian and Schraefer, E.J.: Criteria for essential fatty acid deficiency in plasma as assessed by capillary column gas-liquid chromatography. *Clin. Chem. (Winston-Salem)*, 33 (1987) 1869-1873.
- 703 Tokagi, T., Wakasa, N. and Miyashita, K.: Formation of conjugated diene and triene products in lipoxygenase oxidation of C₁₈, C₂₀, and C₂₂ polyunsaturated fatty acids. *J. Am. Oil Chem. Soc.*, 64 (1987) 1320-1323.
- 704 Vannieuwenhuyze, F. and Sandra, P.: Selectivity optimization for the capillary gas chromatographic analysis of bacteria cellular fatty acids. *Chromatographia*, 23 (1987) 850-855.
- 705 Viden, I. and Rezanka, T.: Capillary gas chromatography-mass spectrometry of aliphatic saturated α,ω -dicarboxylic acid dimethyl esters and direct inlet mass spectrometry of the corresponding free acids. *J. Chromatogr.*, 408 (1987) 145-155.
- 706 Yanagita, T., Satoh, M., Nomura, H., Enomoto, N. and Sugano, M.: Alteration of hepatic phospholipids in rats and mice by feeding di-(2-ethylhexyl)adipate and di-(2-ethylhexyl)phthalate. *Lipids*, 22 (1987) 572-577.
- 707 Yasuhara, A.: Determination of trace lower fatty acids by capillary column gas chromatography-mass spectrometry. *Agric. Biol. Chem.*, 51 (1987) 2259-2261; *C.A.*, 107 (1987) 190114p.
- 708 Zorin, A.E. and Rudenko, B.A.: (Determination of a spectrum of fatty acids in a biological fluid). *U.S.S.R. Pat. SU 1,320,756 (Cl.G01N33/92)*, 30 Jun. 1987, Appl. 3,922,948, 3 Jul. 1985; *C.A.*, 107 (1987) 150771u.
- See also 442, 576, 577, 714, 716, 717, 718, 719, 733, 772, 790, 807, 856, 868, 869, 870, 873, 877, 880, 883, 884, 893, 916, 921.

11b. Prostaglandins

- 709 Herold, D.A., Savory, J., Kinter, M., Ross, R. and Wills, M.R.: Determination of urinary prostanoids by capillary gas chromatography/high-resolution mass spectrometry. *Anal. Chim. Acta*, 197 (1987) 149-158.
- 710 Ishibashi, M., Watanabe, K., Miyazaki, H. and Krolik, S.: (Diethylhydrogensilyl-cyclic diethylsilylene (DEHS-DES) derivatives in gas chromatography-mass spectrometry (GC/MS) of prostanoids. II. GC and GC/MS properties of DEHS-DES derivative of thromboxane B₂ methyl ester-methyloxime). *Yakugaku Zasshi*, 106 (1986) 1118-1125; *C.A.*, 107 (1987) 147480t.
- 711 Pace-Asciak, C.R.: Prostaglandins and related products profiled by GC/MS with negative-ion chemical-ionization detection. *Clin. Invest. Med.*, 10 (1987) 117-120; *C.A.*, 107 (1987) 211933x.
- 712 Umeda, M., Amagaya, S. and Ogiwara, Y.: Separation of thromboxane B₂ by high-performance liquid chromatography. *J. Chromatogr.*, 422 (1987) 246-252.

11c. Lipids and their constituents

- 713 Fukatsu, M. and Tamura, T.: (Gas chromatography-mass spectrometry analysis of triglyceride. V. Application to the analysis of hardened coconut oil). *Yakugaku*, 36 (1987) 581-583; *C.A.*, 107 (1987) 174509r.
- 714 Kanzaki, T., Ishikawa, Y., Morisaki, N., Shirai, K., Saito, Y. and Yoshida, S.: Abnormal metabolism of polyunsaturated fatty acids and phospholipids in diabetic glomeruli. *Lipids*, 22 (1987) 704-710.
- 715 Khalafy, E., Fadel, H.M. and Solima, M.M.A.: Adulterated beef fat: the structure of triglycerides and 2-monoglycerides. *Grasas Aceites*, 38 (1987) 176-179.
- 716 Mori, T.A., Codde, J.P., Vandongen, R. and Beilin, L.J.: New findings in the fatty acid composition of individual platelet phospholipids in man after dietary fish oil supplementation. *Lipids*, 22 (1987) 744-750.
- 717 Nikolopoulou, M. and Vary, J.C.: Lipid composition of *Bacillus megaterium* spores and spore membranes. *Lipids*, 22 (1987) 698-703.

718 Zwingelstein, G., Brichon, G., Meister, R., Maury, C. and Gresser, I.: Interferon α/β induces changes in the metabolism of polyenoic phospholipids and diacylglycerols in the livers of suckling mice. *Lipids*, 22 (1987) 736-743.

See also 440, 442, 648, 652, 660, 663, 675, 683, 692, 694, 696, 698, 727, 728, 881, 883, 884.

12. ORGANIC PEROXIDES

719 Awl, R.A., Frankel, E.N. and Weisleder, D.: Cyclic fatty esters: Hydroperoxides from autoxidation of methyl 9-(6-propyl-3-cyclohexenyl)-(Z)8-nonenolate. *Lipids*, 22 (1987) 721-730.

720 Bogomolov, V.I., Etinger, A.G., Poponova, R.V. and Chupakhin, M.: (Chromatographic-mass spectrometric study of impurity composition of high-purity solvents. III. Impurity composition of tetrahydrofuran. Identification of peroxides). *Vysokochist. Veshchestva*, (1987) 203-208; *C.A.*, 107 (1987) 228119j.

721 Cai Hanlin: (Gas chromatographic determination of cyclohexyl hydroperoxide). *Sepu*, 5 (1987) 245-247.

722 Frank, H., Wiegand, M., Strecker, M. and Thiel, D.: Monohydroperoxides of linoleic acid in endoplasmic lipids of rats exposed to tetrachloromethane. *Lipids*, 22 (1987) 689-697.

See also 736.

13. STEROIDS

13a. Pregnane and androstane derivatives

723 Rolland, D., Patricot, M.C., Mathian, B. and Revol, A.: (Determination of the urinary excretion of tetrahydroaldosterone by gas chromatography. Normal adult values). *Ann. Biol. Clin. (Paris)*, 45 (1987) 70-73; *C.A.*, 107 (1987) 191103w.

13b. Estrogens

724 Fotsis, T.: The multicomponent analysis of estrogens in urine by ion exchange chromatography and GC-MS. II. Fractionation and quantitation of the main groups of estrogen conjugates. *J. Steroid Biochem.*, 28 (1987) 215-236; *C.A.*, 107 (1987) 191096w.

725 Fotsis, T. and Adlercreutz, H.: The multicomponent analysis of estrogens in urine by ion exchange chromatography and GC-MS. I. Quantitation of estrogens after initial hydrolysis of conjugates. *J. Steroid Biochem.*, 28 (1987) 203-213; *C.A.*, 107 (1987) 191095v.

13c. Sterols

726 Deng Feng and Zhou Yingtian: (Gas chromatographic determination of cholesterol content in egg yolk). *Yingyang Xuebao*, 9 (1987) 158-163; *C.A.*, 107 (1987) 174510j.

727 Horstmann, P. and Montag, A.: Sterinanalytik zum Nachweis eines Zusatzes von Sonnenblumenöl zu Safrolöl. *Fett Wiss. Technol.-Fat Sci. Technol.*, 89 (1987) 381-388.

728 Low, C. and Parks, L.W.: Sterol and phospholipid acyl chain alterations in *Saccharomyces cerevisiae* secretion mutants as a function of temperature stress. *Lipids*, 22 (1987) 715-720.

729 Parish, E.J., Wei Tsao Yi and Livant, P.: A facile synthesis and carbon-13 nuclear magnetic resonance spectral properties of 7-ketocholesteryl benzoate. *Lipids*, 22 (1987) 760-763.

730 Smith, R.L., Sullivan, D.M. and Richter, E.F.: Determination of phytosterols in butter samples by using capillary column gas chromatography. *J. Assoc. Off. Anal. Chem.*, 70 (1987) 912-915.

See also 649, 868, 881.

13d. Bile acids and alcohols

- 731 Chikai, T., Nakao, H. and Uchida, K.: Deconjugation of bile acids by human intestinal bacteria implanted in germ-free rats. *Lipids*, 22 (1987) 669-671.
- 732 Yamaga, N., Adachi, K., Kohara, H. and Shimizu, K.: Internal standards for quantitative gas chromatography of individual bile acids after group separation of bile acids in urine. *J. Chromatogr.*, 422 (1987) 25-32.

14. STEROID GLYCOSIDES AND SAPONINS

- 733 Rao, D. and Bories, G.: Simple gas chromatographic method for the determination of medicagenic acid in alfalfa (*Medicago sativa*). *J. Chromatogr.*, 410 (1987) 169-175.

15. TERPENES AND OTHER VOLATILE AROMATIC COMPOUNDS

15a. Terpenes

- 734 Biougne, J., Chalchat, J.-C., Garry, R.-P. and Michet, A.: Characterisation des cadinenes dans les huiles essentielles. *Parf. Cosmet. Aromes*, No. 73 (1987) 59-61.
- 735 Leven, W. and Willuhn, G.: Sesquiterpene lactones from *Arnica chamissonis* Less. VI. Identification and quantitative determination by high-performance liquid and gas chromatography. *J. Chromatogr.*, 410 (1987) 329-342.
- 736 Schieberle, P., Maier, W., Firl, J. and Grosch, W.: HRGC separation of hydroperoxides formed during the photosensitized oxidation of (R)-(+)-limonene. *J. High Resolut. Chromatogr. Chromatogr. Commun.*, 10 (1987) 588-593.

See also 536, 881.

15b. Essential oils

- 737 Bicchi, C., D'Amato, A. and Frattini, C.: Selective sampling and capillary gas chromatography in the analysis of essential oils. *Chromatographia*, 23 (1987) 818-824.
- 738 Bicchì, C., Frattini, C. and Raverdino, V.: Considerations and remarks on the analysis of *Athemis nobilis* L. essential oil by capillary gas chromatography and "hephenated" techniques. *J. Chromatogr.*, 411 (1987) 237-249.
- 739 Cartoni, G.P., Goretti, G. and Russo, V.M.: Capillary columns in series for the gas chromatographic analysis of essential oils. *Chromatographia*, 23 (1987) 790-795.
- 740 Curro, P., Lanuzza, F. and Micali, G.: (Evaluation of the volatile fraction of saffron by headspace gas chromatography). *Rass. Chim.*, 38 (1986) 331-334; *C.A.*, 107 (1987) 132735f.
- 741 Gao Hai, Sha Zhenfang and Sun Wenji: (Gas chromatography of eucalyptole in eucalyptus oil). *Yaowu Fenxi Zazhi*, 7 (1987) 168-169; *C.A.*, 107 (1987) 161758w.
- 742 Gu Yuecui, Zhong Mingfang and Chen Shaohuai: (Comparison of the essential oils from Sichuan commodity Xixin (Asarum) by TLC and GC). *Zhongcaoyao*, 18 (1987) 299-301; *C.A.*, 107 (1987) 205265q.
- 743 Hethelyi, E., Danos, B., Tetenyi, P. and Koczka, I.: GC-MS analysis of the essential oils of four *Tagetes* species and the antimicrobial activity of *Tagetes minuta*. *Flavour Fragrance J.*, 1 (1986) 169-173; *C.A.*, 107 (1987) 161480z.
- 744 Klyuev, N.A., Zamurenko, V.A. and Evtushenko, N.S.: (Use of chromatography-mass spectrometry to detect constituents of volatile oils). *Farmatsiya (Moscow)*, 36, No. 5 (1987) 70-77; *C.A.*, 107 (1987) 223025r - a review with 87 refs.
- 745 Krajewska, A.M. and Powers, J.J.: Gas chromatography of methyl derivatives of naturally occurring capsaicinoids. *J. Chromatogr.*, 409 (1987) 223-233.
- 746 Li Chongfang, Li Huifen, Zhang Qingwei, Zhang Wenying, Zhou Jing and Chen Qimeng: (Determination of borneol in Niu Huang Guanxin pills). *Zhongcaoyao*, 18 (1987) 353-354; *C.A.*, 107 (1987) 183665r.

- 747 Mossa, J.S., Hifnawy, M.S., Al-Yahya, M.A., Al-Mesha, I.A. and Mekkawi, A.G.: Aromatic plants of Saudi Arabia. Part 8 - GC/MS analysis of essential oils of *Pulicaria arabica* and *P. undulata*. *Int. J. Crude Drug Res.*, 25 (1987) 113-119; *C.A.*, 107 (1987) 204891d.
- 748 Zhu Yeping, Huang Qin, Wang Zhihua and Hong Xiaokun: (The gas chromatographic comparison of essential oils from leaves of *Chimonanthus* species in China during the flowering and vegetative stages). *Shanghai Yike Daxue Xuebao*, 14, No. 2 (1987) 112; *C.A.*, 107 (1987) 140857n.

See also 727, 734, 886.

16. NITRO AND NITROSO COMPOUNDS

- 749 Dawson, B.A. and Lawrence, R.C.: Analysis of piperazine drug formulations for N-nitrosamines. *J. Assoc. Off. Anal. Chem.*, 70 (1987) 840-841.
- 750 Henning, B. and Benecke, R.: (Gas-chromatographic assay of nitroglycerin in human plasma). *Pharmazie*, 42 (1987) 507-510; *C.A.*, 107 (1987) 211364n.
- 751 Jaeger, H., Lutz, D., Michaelis, K. and Salama, Z.B.: Determination of nitrates in plasma. *Drugs*, 33, (Suppl. 4) (1987) 9-22; *C.A.*, 107 (1987) 146714k.
- 752 Korfmacher, W.A., Rushing, L.G., Arey, J., Zielinska, B. and Pitts, J.N., Jr.: Identification of mononitropyrenes and mononitrofluoranthenes in air particulate matter *via* fused silica gas chromatography combined with negative ion atmospheric pressure ionization mass spectrometry. *J. High Resolut. Chromatogr. Chromatogr. Commun.*, 10 (1987) 641-646.
- 753 Wang Delin, Wang Pendi, Xu Xiaoyong, Liu Jinbao and Jin Jing: (Determination of trace nitro-compounds in biological samples by acid digestion capillary gas chromatography). *Sepu*, 5 (1987) 380-382.
- 754 Zielinska-Psuja, B. and Budziak, A.: (Determination of *p*-nitroaniline in the urine by gas chromatography). *Bromatol. Chem. Toksykol.*, 20 (1987) 59-62; *C.A.*, 107 (1987) 213069u.

See also 838, 907, 913.

17. AMINES, AMIDES AND RELATED NITROGEN COMPOUNDS

17a. Amines and polyamines

- 755 Bakumenko, T.T., Usenko, V.A. and Pryadko, Z.N.: (Analysis of aliphatic amines by gas chromatography and thermoionic detector). *Fermentn. Spirt. Prom-st.*, No. 5 (1987) 39-42; *C.A.*, 107 (1987) 216076e.
- 756 Hill, R.H., Jr., Todd, G.D., Kilbourne, E.M., Cline, R.E., McCraw, J., Orti, D.L., Bailey, S.L. and Needham, L.L.: Gas chromatographic/mass spectrometric determination of aniline in food oils associated with the Spanish toxic oil syndrome. *Bull. Environ. Contam. Toxicol.*, 39 (1987) 511-515; *C.A.*, 107 (1987) 196586r.
- 757 Iwasaki, M., Nakasono, I., Ogata, M., Kubo, S. and Suyama, H.: (Relation between methamphetamine levels in the marrow and serum). *Eisei Kagaku*, 33 (1987) 136-139; *C.A.*, 107 (1987) 148698g.
- 758 Keller, M.A. and Saba, C.S.: Chromatographic analysis of phenyl-1-naphthylamine and 4,4'-dioctyldiphenylamine and their intermediate oxidation products in oxidized lubricants. *J. Chromatogr.*, 409 (1987) 325-335.
- 759 Lynn, B.C. and Alley, E.G.: Derivatization of diaminocyclohexanes for determination of *cis* and *trans* isomers b- gas chromatography. *J. Chromatogr.*, 408 (1987) 211-215.
- 760 Sitaram, B.R., Lockett, L., McLeish, M., Hayasaka, Y., Blackman, G.L. and McLeod, W.R.: Gas chromatographic-mass spectroscopic characterisation of the psychotomimetic indolealkylamines and their *in vivo* metabolites. *J. Chromatogr.*, 422 (1987) 13-23.

See also 640, 875.

17c. Urea and guanidine derivatives

See 745.

17d. Other amine derivatives and amides (excluding peptides)

- 761 Wright, D., Jr.: New method for the determination of 1,1-dimethylhydrazine residues in apples and peaches. *J. Assoc. Off. Anal. Chem.*, 70 (1987) 718-720.

18. AMINO ACIDS AND PEPTIDES; CHEMICAL STRUCTURE OF PROTEINS

18a. Amino acids and their derivatives

- 762 Baer, W. and Fricke, G.: Rapid and improved gas-liquid chromatography technique for detection of hippurate hydrolysis by *Campylobacter jejuni* and *Campylobacter coli*. *J. Clin. Microbiol.*, 25 (1987) 1776-1778; *C.A.*, 107 (1987) 130217q.
- 763 Cabrys, J., Konecki, J., Plaza, J., Wesolek, W. and Shani, J.: Free amino acids in the cytosol of human atrium and ventricle of the heart, as analyzed by gas-liquid chromatography. *Cor. Vasa*, 29 (1987) 152-158; *C.A.*, 107 (1987) 173030r.
- 764 Greenaway, W. and Whatley, F.R.: Formation of N-carboxy compounds during trimethylsilylation of amino acids in the presence of carbon dioxide. *J. Chromatogr.*, 409 (1987) 383-389.
- 765 Sakamoto, M., Nishimoto, M., Kohara, N. and Masuko, F.: Gas chromatographic and mass spectrometric behaviour of O-alkylserines as their N-trifluoroacetyl butyl esters. *J. Chromatogr.*, 411 (1987) 259-273.
- 766 Wallace, P.L., Patton, C.M. and Moss, C.W.: Determination of hippurate hydrolysis by gas-liquid chromatography. *J. Clin. Microbiol.*, 25 (1987) 1766-1768; *C.A.*, 107 (1987) 130216p.
- 767 Wu Jiang and Ouyang Zheng: Determination of selenomethionine by reaction with cyanogen bromide and gas chromatography). *Sepu*, 5 (1987) 236-238; *C.A.*, 107 (1987) 150547a.

See also 444, 573, 868.

20. ENZYMES AND ENZYME ACTIVITY ESTIMATION

20a. Oxidoreductases

See 553.

21. PURINES, PYRIMIDINES, NUCLEIC ACIDS AND THEIR CONSTITUENTS

21a. Purines, pyrimidines, nucleosides, nucleotides

- 768 Minnetian, O., Saha, M. and Giese, R.W.: Oxidation-elimination of a DNA base from its nucleoside to facilitate determination of alkyl chemical damage to DNA by gas chromatography with electrophore detection. *J. Chromatogr.*, 410 (1987) 453-457.
- 769 Palni, L.M.S., Tay, S.A.B. and MacLeod, J.K.: Determination by gas chromatography-mass spectrometry of (¹⁵N₅)adenine incorporation into endogenous cytokinins and the effect of tissue age on cytokinin biosynthesis in *Datura innoxia* crown gall tissue. *Plant Physiol.*, 84 (1987) 1158-1165; *C.A.*, 107 (1987) 172542x.

See also 854.

22. ALKALOIDS

- 770 Akhmed-Zade, F.D., Dement'eva, N.N. and Stepnykh, Yu.P.: (Gas-liquid chromatographic determination of tropine and tropinone). *Khim.-Farm. Zh.*, 21 (1987) 1012-1017; *C.A.*, 107 (1987) 183697c.
- 771 Chen Jinhua: (Determination of caffeine in tea, coffee and cola drink by gas chromatography with nitrogen-phosphorus detector). *Sepu*, 5 (1987) 367-369.
- 772 Moore, J.M., Cooper, D.A., Lurie, I.S., Kram, T.C., Carr, S., Harper, C. and Yeh, J.: Capillary gas chromatographic-electron capture detection of coca-leaf-related impurities in illicit cocaine: 2,4-diphenylcyclobutane-1,3-dicarboxylic acids, 1,4-diphenylcyclobutane-2,3-dicarboxylic acids and their alkaloidal precursors, the truxillines. *J. Chromatogr.*, 410 (1987) 297-318.
- 773 Taylor, R.W., Jain, N.C. and George, M.P.: Simultaneous identification of cocaine and benzoylecgonine using solid phase extraction and gas chromatography/mass spectrometry. *J. Anal. Toxicol.*, 11 (1987) 233-234; *C.A.*, 107 (1987) 213085w.
- 774 Timor, C.E., Perez, R., Mena, I., Pena, A. and Corrales, I.: (Determination of the scopolamine content in leaves of *Datura candida*, staff, by gas chromatography). *Rev. Cubana Farm.*, 21 (1987) 103-110; *C.A.*, 107 (1987) 183660k.

See also 856, 859.

23. OTHER SUBSTANCES CONTAINING HETEROCYCLIC NITROGEN

23c. Indole derivatives

- 775 Murray, S., Gooderham, N.J., Barnes, V.F., Boobis, A.R. and Davies, D.S.: Trp-P-2 is not detectable in cooked meat and fish. *Carcinogenesis (London)*, 8 (1987) 937-940; *C.A.*, 107 (1987) 132772r.

See also 659, 790.

23d. Pyridine derivatives

- 776 Golovnya, R.V., Samusenko, A.L. and Dmitriev, L.B.: (Prediction of methyl pyridine retention indices in capillary gas chromatography based on the sorption energy non-additivity principle). *Izv. Akad. Nauk SSSR*, (1987) 2234-2239.
- 777 Krasavin, V.Yu., Egorenko, G.A., Kanaeva, O.A., Zel'venskii, V. Yu., Samolyuk, S.I., Zhadanov, B.V., Olikova, V.A., Parkhaeva, G.N. and Stepanova, A.G.: (Study of the composition of impurities in purified pyridine). *Koks Khim.*, No. 7 (1987) 34-35; *C.A.*, 107 (1987) 136287j.

See also 872.

24. ORGANIC SULPHUR COMPOUNDS

- 778 Rycaj, B.: (Determination of ethylenethiourea in the form of trimethylsilyl derivative by gas chromatography). *Bromatol. Chem. Toksykol.*, 20 (1987) 76-82; *C.A.*, 107 (1987) 213070n.
- 779 Schrementi, J. and Meganathan, R.: A gas chromatographic method for the determination of tetrahydrothiophene produced by enzymic reduction of tetrahydrothiophene-1-oxide. *Microbios. Lett.*, 35 (1987) 79-85; *C.A.*, 107 (1987) 171744c.
- 780 Talyzin, V.V., Anisimov, V.Ya., Yakovleva, O.I., Misharina, T.A. and Golovnya, R.V.: (Extraciton and preconcentration of volatile dialkylpolysulphides - onion odour components). *Zh. Anal. Khim.*, 42 (1987) 2076-2079.

See also 562, 576.

26. ORGANOMETALLIC AND RELATED COMPOUNDS

26a. Organometallic compounds

- 781 Baugh, P.J., Casson, A., Jones, M.W. and Jones, A.C.: Sampling and injection system for the analysis of metallorganic chemicals by gas chromatography-mass spectrometry. *J. Chromatogr.*, 411 (1987) 445-451.
- 782 Ohhira, S. and Matsui, H.: (Determination of triphenyltin compounds in biological materials by gas chromatography). *Pokkyo Igakkai Zasshi*, 2 (1987) 177-183; *C.A.*, 107 (1987) 192394d.
- 783 Short, J.W.: Measuring tri-*n*-butyltin in salmon by atomic absorption: analysis with and without gas chromatography. *Bull. Environ. Contam. Toxicol.*, 39 (1987) 412-416; *C.A.*, 107 (1987) 148690y.

26b. Boranes, silanes and related non-metallic compounds

- 784 Gholson, A.R., Louis, R.S. and Hill, H.H.: Determination of silicon number in silylated organic compounds using the hydrogen atmosphere flame ionization detector. *J. Chromatogr.*, 408 (1987) 329-334.
- 785 Hasegawa, I., Sakka, S., Kuroda, K. and Kato, C.: Trimethylsilylation of the hydrolysed and polycondensed products of methyltriethoxysilane. *J. Chromatogr.*, 410 (1987) 137-143.
- 786 Oyamada, N., Kikuchi, M. and Ishizaki, M.: Determination of dimethyl selenide in breath air of mice by gas chromatography. *Anal. Sci.*, 3 (1987) 373-376; *C.A.*, 107 (1987) 148688d.

See also 767.

26c. Coordination compounds

- 787 Bazhenova, L.N., Kirichenko, V.E., Pashkevich, K.I., Sereda, I.P. and Mikhailenko, V.P.: (Interaction of nickel β -aminovinylketonates and stationary liquid phases). *Izv. Akad. Nauk SSSR, Ser. Khim.*, (1987) 741-745.
- 788 Khuhawar, M.Y. and Vasandani, A.G.M.: Preparation, thermogravimetric and gas-chromatographic studies of copper(II) and nickel(II) complexes of some new ketoamine Schiff bases. *J. Chem. Soc. Pak.*, 9 (1987) 43-52; *C.A.*, 107 (1987) 167666m.
- 789 Robards, K., Patsalides, E. and Dilli, S.: Gas chromatography of metal β -diketonates and their analogues. *J. Chromatogr.*, 411 (1987) 1-41 - a review with 360 refs.

27. VITAMINS AND VARIOUS GROWTH REGULATORS (NON-PEPTIDIC)

- 790 Sanderson, K.J., Jameson, P.E. and Zabkiewicz, J.A.: Auxin in a seaweed extract: identification and quantitation of indole-3-acetic acid by gas chromatography-mass spectrometry. *J. Plant Physiol.*, 129 (1987) 363-367; *C.A.*, 107 (1987) 129053b.
- 791 Sastry, C.S.P., Singh, N.R., Rao, Y.R., Bapuji, M., Reddy, M.N. and Shankar, D.G.: Gas chromatographic determination of menadione. *Indian Drugs*, 24 (1987) 438-440; *C.A.*, 107 (1987) 205286x.
- 792 Staroverov, V.M., Chuev, V.P. and Romanenko, V.P.: (Chromatographic analysis of intermediates in vitamin A synthesis). *Khim.-Farm. Zh.*, 20 (1986) 1379-1382; *C.A.*, 107 (1987) 190020e.

See also 658, 761, 849.

28. ANTIBIOTICS

- 793 Al Ashmawi, M.I. and El Sadek, M.: Gas chromatographic and gas chromatographic-mass spectrometric analysis of penicillin V benzathine oral suspension. *Egypt. J. Pharm. Sci.*, 28 (1987) 361-370; *C.A.*, 107 (1987) 223367d.
- 794 Takatsuki, K., Suzuki, S., Sato, N., Ushizawa, I. and Shoji, T.: Gas chromatographic/mass spectrometric determination of erythromycin in beef and pork. *J. Assoc. Off. Anal. Chem.*, 70 (1987) 708-713.

29. INSECTICIDES, PESTICIDES AND OTHER AGROCHEMICALS

29a. Chlorinated insecticides

- 795 Glastrup, J.: Insecticide analysis by gas chromatography in the stores of the Danish National Museum's Ethnographic collection. *Stud. Conserv.*, 32 (1987) 59-64; *C.A.*, 107 (1987) 129080h.
- 796 Hirai, Y. and Tomokuni, K.: (Simple sample-pretreatment for GC/MS determination of residual chlordanes in river sediments). *Bunseki Kagaku*, 36 (1987) 817-819.
- 797 Huschek, G. and Schieck, K.: (Tracer study of the behavior of HCH isomers during sample preparation for gas chromatography). *Acta Hydrochim. Hydrobiol.*, 15 (1987) 351-355; *C.A.*, 107 (1987) 183155z.
- 798 Kapila, S., Duebelbeis, D., Malhotra, R., Yanders, A.F. and Manahan, S.E.: A two-dimensional reaction gas chromatographic system for congener specific determination of polychlorinated organics. In: R. Greenhalgh and T.R. Roberts (Editors), *Pesticide Science Biotechnology*, Blackwell, Oxford, 1987, pp. 325-328; *C.A.*, 107 (1987) 128537a.
- 799 Noren, K. and Sjövall, J.: Analysis of organochlorine pesticides, polychlorinated biphenyls, dibenzo-*p*-dioxins and dibenzofurans in human milk by extraction with the lipophilic gel Lipidex® 5000. *J. Chromatogr.*, 422 (1987) 103-115.
- 800 Sabbah, S., Jemaa, Z. and Bouguerra, M.L.: Etude chromatographique des résidus de pesticides organochlores dans le lait maternel, le sang de cordon et le sang d'adulte. *Analisis*, 15 (1987) 399-403.

See also 482, 499, 570, 805, 883, 908.

29b. Phosphorus insecticides

- 801 Basile, M. and Lamberti, F.: (Gas-chromatographic determination of fenamiphos residues in peaches). *Essenze Deriv. Agrum.*, 57 (1987) 112-115; *C.A.*, 107 (1987) 174596s.
- 802 Gyorfı, L., Ambrus, A. and Bolygo, E.: Optimization on determination and clean-up parameters for sensitive multiresidue analysis of pesticides. In: R. Greenhalgh and T.R. Roberts (Editors), *Pesticide Science Biotechnology*, Blackwell, Oxford, 1987, pp. 353-356; *C.A.*, 107 (1987) 132730a.
- 803 Hang Changmian: (Study on separation of four organophosphorous pesticides by gas chromatography). *Sepu*, 5 (1987) 390-392.
- 804 Li Yuzhen and Wang Gendi: (Determination of organophosphorus pesticides in wastewater). *Shanghai Huanjing Kesue*, 6, No. 4 (1987) 29-31, 42; *C.A.*, 107 (1987) 222884h.
- 805 Prinsloo, S.M. and De Beer, P.R.: Gas chromatographic relative retention data for pesticides on nine packed columns: II. Organophosphorus and organochlorine pesticides, using electron-capture detection. *J. Assoc. Off. Anal. Chem.*, 70 (1987) 878-888.

See also 499, 903.

29c. Carbamates

- 806 Gerasimenko, V.A. and Voskoboinikov, V.V.: (Thermostability and gas chromatographic retention of substituted alkylacrylcarbamates). *Zh. Anal. Khim.*, 42 (1987) 2043-2047.

- 807 Meemken, H.A., Rudolph, P. and Fuerst, P.: (Detection and determination of chloro-phenoxy carboxylic acids by capillary GC/MS). *Dtsch. Lebensm.-Rundsch.*, 83 (1987) 239-245; *C.A.*, 107 (1987) 196616a.
- 808 Richardson, G.M. and Qadri, S.U.: Extraction of aminocarb and a metabolite from whole fish and derivatization for electron-capture gas chromatography. *J. Agric. Food Chem.*, 35 (1987) 877-878.
- 809 Worobey, B.L. and Sun, W.F.: Isolation and identification of chlorpropham and two of its metabolites in potatoes by GC-MS. *Chemosphere*, 16 (1987) 1457-1462; *C.A.*, 107 (1987) 152975n.

See also 556, 816, 903.

29d. Herbicides

- 810 Greenberg, R.S.: Gas chromatographic determination of mecarbam and its metabolites mecarboxon, diethoate, and diethoxon in cottonseeds. *J. Assoc. Off. Anal. Chem.*, 70 (1987) 870-874.
- 811 McNally, M.E.P. and Wheeler, J.R.: Supercritical fluid extraction coupled with supercritical fluid chromatography for the separation of sulfonylurea herbicides and their metabolites from complex matrices. *J. Chromatogr.*, 435 (1988) 63-71.
- 812 Mitsui, T. and Minami, Y.: (Determination of herbicide Paraquat using pyrolysis GC). *Bunseki Kagaku*, 36 (1987) 821-824.
- 813 Shiraishi, H. and Otsuki, A.: Identification and determination of 4-(chloro-methylsulfonyl)bromobenzene, a new herbicide additive, in lake waters. *Water Res.*, 21 (1987) 843-847; *C.A.*, 107 (1987) 140754b.
- 814 Viden, I., Rathouska, Z., Davidek, J. and Hajslova, J.: Use of gas liquid chromatography/mass spectrometry for triazine herbicide residues analysis in forage and milk. *Z. Lebensm.-Unters. Forsch.*, 185 (1987) 98-105; *C.A.*, 107 (1987) 152983p.

29e. Fungicides

- 815 Daft, J.: Determining multifungigants in whole grains and legumes, milled and low-fat grain products, spices, citrus fruit, and beverages. *J. Assoc. Off. Anal. Chem.*, 70 (1987) 734-739.

29f. Other types of pesticides and various agrochemicals

- 816 Forbes, S.: Capillary gas chromatography in pesticide residue analysis. *Anal. Proc.*, 24 (1987) 306-308.
- 817 Toth, J.P. and Bardalaye, P.C.: Capillary gas chromatographic separation and mass spectrometric detection of cyromazine and its metabolite melamine. *J. Chromatogr.*, 408 (1987) 335-340.

See also 849.

31. PLASTICS AND THEIR INTERMEDIATES

- 818 Alajbeg, A., Jarm, V. and Kovac-Filipovic, M.: (Evaluation of pyrocondensates as a polymerization raw material). *Croat. Chem. Acta*, 60 (1987) 337-350; *C.A.*, 107 (1987) 134700q.
- 819 Becker, H. and Gnauck, R.: Inversgaschromatographische Untersuchungen von Polyetheralkoholen. *J. Chromatogr.*, 410 (1987) 267-278.
- 820 Chaffee, A.L., Cavell, K.J., Masters, A.F. and Western, R.J.: Detailed gas chromatography/mass spectrometric structural determination of olefin oligomerization products. *Ind. Eng. Chem. Res.*, 26 (1987) 1822-1824; *C.A.*, 107 (1987) 156886g.
- 821 Cwiek, K. and Mazur, H.: (Determination of residues of free styrene in polystyrene for food packaging). *Rocz. Panstw. Zakl. Hig.*, 37 (1986) 551-554; *C.A.*, 107 (1987) 196596u.

- 822 Demertzis, P.G., Kontominas, M.G. and Gilbert, S.G.: Gas chromatographic determination of sorption isotherms of vinylidene chloride on vinylidene chloride copolymers. *J. Food Sci.*, 52 (1987) 747-750; *C.A.*, 107 (1987) 174542w.
- 823 Freytag, U. and Radusch, H.J.: (Study of the thermodynamic compatibility of high-density polyethylene and low-density polyethylene in the melt state. I. Determination of solubility parameters). *Angew. Makromol. Chem.*, 152 (1987) 1-13; *C.A.*, 107 (1987) 135060t.
- 824 Furukawa, M., Yoshitake, N. and Yokoyama, T.: Identification of polymer glycols in polyurethane elastomers by pyrolysis-gas chromatography. *J. Chromatogr.*, 435 (1988) 219-224.
- 825 Grenier-Loustalot, M.F., Mouline, G., Grenier, P., Serrano, D., Harran, D. and Horny, P.: (Study of gelation phenomena in DGEBA- and TGMDA-DDS systems by inverse gas chromatography, viscosimetry, and rheological methods). *Makromol. Chem., Macromol. Symp.*, 9 (1987) 143-154; *C.A.*, 107 (1987) 177074f.
- 826 Huh, D.S., Kim, J.S., Kim, K.J., Ahn, B.K., Suh, S.K. and Han, O.K.: (Analysis of vulcanized rubber by pyrolysis-gas chromatography (I) (Vulcanizates of NR (natural rubber), BR (butadiene rubber), and SBR)). *Komu Hakhoehi*, 22 (1987) 11-19; *C.A.*, 107 (1987) 155983z.
- 827 Kotsev, B.Kh., Belousov, V.N. and Mikitaev, A.K.: (Study of the molecular mobility of polystyrene near the glass transition by inverse gas chromatography). *Vysokomol. Soedin., Ser. B*, 29 (1987) 565-568; *C.A.*, 107 (1987) 199235m.
- 828 Nakagawa, H., Tsuge, S. and Koyama, T.: Studies on thermal degradation of epoxy resins by high-resolution pyrolysis-gas chromatography. *J. Anal. Appl. Pyrolysis*, 12 (1987) 97-113; *C.A.*, 107 (1987) 176996w.
- 829 Ohtani, H., Kimura, T., Okamoto, K., Tsuge, S., Nagataki, Y. and Miyata, K.: Characterization of polyurethanes by high-resolution pyrolysis-capillary gas chromatography. *J. Anal. Appl. Pyrolysis*, 12 (1987) 115-133; *C.A.*, 107 (1987) 176837v.
- 830 Osipchik, V.S., Romyantseva, N.D. and Novichenkova, V.N.: Gas-chromatographic determination of the content of low-molecular-weight products and residual quantities of dimethyldichlorosilane and trimethylchlorosilane in poly(methylsiloxane). *Lakokras. Mater. Ikh Primen.*, No. 4 (1987) 50-52; *C.A.*, 107 (1987) 200439q.
- 831 Sanetra, R., Kolarz, B.N. and Wlochowicz, A.: Determination of thermodynamic data for the interaction of aliphatic alcohols with poly(styrene-co-divinylbenzene) using inverse gas chromatography. *Polymer*, 28 (1987) 1753-1757; *C.A.*, 107 (1987) 135079f.
- 832 Sotnikov, E.E. and Bubchikova, L.A.: (Identification of the elemental composition of additives in a polymer by ion and pyrolytic gas chromatography). *Zh. Anal. Khim.*, 42 (1987) 1896-1900.
- 833 Tseng, H.S., Wong, P.C., Lloyd, D.R. and Barlow, J.W.: Thermodynamic interaction in polybutadiene/solute systems by inverse gas chromatography. *Polym. Eng. Sci.*, 27 (1987) 1141-1147; *C.A.*, 107 (1987) 155221z.
- 834 Tsotsis, T.K., Turchi, C. and Glover, C.J.: Comparison of multicomponent gas chromatography measurements of vapor-liquid equilibrium with static measurements using a polymer/two solvent ternary system. *Macromolecules*, 20 (1987) 2445-2452; *C.A.*, 107 (1987) 176848z.
- 835 Usova, E.P., Mitina, L.I., Sergeeva, G.S., Sizova, G.C. and Znamenskaya, A.P.: (Gas chromatographic determination of impurities in caprolactam produced from toluene). *Zh. Anal. Khim.*, 42 (1987) 2084-2087.
- 836 Xu Zhengyan and Li Ming: (Study of an epoxy urethane by pyrolysis-gas chromatography). *Sepu*, 5 (1987) 247-250.
- 837 Yosida, R. and Watanabe, Y.: (Determination of vinyl acetate in ethylenevinyl acetate copolymers by pyrolysis GC). *Bunseki Kagaku*, 36 (1987) 862-866.
- 838 Zwickenpflug, W. and Richter, E.: Rapid method for the detection and quantification of N-nitrosodibutylamine in rubber products. *J. Chromatogr. Sci.*, 25 (1987) 506-509.

See also 456, 470, 486, 533, 915.

32. DRUG ANALYSIS

32a. Drug analysis, general techniques

- 839 Finkle, B.S.: Drug-analysis technology: Overview and state of art. *Clin. Chem. (Winston-Salem)*, 33 (1987) 13B-18B.
- 840 Krzek, J.: (Gas-chromatographic determination of the components in prescription drugs). *Acta Pol. Pharm.*, 43 (1986) 250-255; *C.A.*, 107 (1987) 205312c.
- 841 Rao, C.R., Murthy, S.S.N. and Khadgpathi, P.: Recent trends and applications of gas chromatography to pharmaceutical analysis. *East Pharm.*, 30 (1987) 35-41; *C.A.*, 107 (1987) 183637h - a review with 33 refs.

See also 612, 749, 772.

32c. Cardiovascular drugs

- 842 Marko, V.: Selective solid-phase extraction of mexiletine from human serum prior to its GLC analysis. *Pharmazie*, 42 (1987) 387-389; *C.A.*, 107 (1987) 146726r.

See also 751.

32d. Central nervous system drugs

- 843 Jindal, S.P., Lutz, T. and Bagchi, S.P.: Gas-liquid chromatographic-mass spectrometric determination of 1-methyl-4-phenyl-4-propionoxypiperidine using a stable isotope-labeled analogue as an internal standard. *J. Chromatogr.*, 408 (1987) 356-359.
- 844 Karnes, H.T. and Farthing, D.: Improved method for the determination of oxazepam in plasma using capillary gas chromatography and nitrogen-phosphorus detection. *LC-GC*, 5 (1987) 978-979.

32e. Chemotherapeutics (except cytostatics)

- 845 Jemal, M. and Cohen, A.I.: Simultaneous determination of the acid chloride S-benzoyl-3-mercapto-2-methylpropanoyl chloride and residual carboxylic acid by capillary gas chromatography. *J. Chromatogr.*, 408 (1987) 360-364.
- 846 Kaniewska, T., Kublin, E. and Weclawowicz, E.: (Identification and determination of minoxidil by gas chromatography). *Acta Pol. Pharm.*, 43 (1986) 588-592; *C.A.*, 107 (1987) 161800d.
- 847 Karlaganis, G. and Bircher, J.: Bioavailability determination of lidocaine by capillary gas chromatography ammonia chemical ionization mass spectrometry. *Biomed. Environ. Mass Spectrom.*, 14 (1987) 513-516; *C.A.*, 107 (1987) 211423f.
- 848 Liu Xiaodong, Huang Shengkai and Yuan Huinan: (Determination of sophoramine in rabbit plasma by capillary column gas chromatography and pharmacokinetics of sophoramine and sophocarpine). *Zhongguo YaoLi Xuebao*, 8 (1987) 474-477; *C.A.*, 107 (1987) 168156g.
- 849 Malisch, R.: (Method for the simultaneous determination of residues of chemotherapeutics, antiparasitics, and growth promoters in foods of animal origin. Part III. Determination of chloramphenicol and meticlorpindol). *Z. Lebensm.-Unters. Forsch.*, 184 (1987) 467-477; *C.A.*, 107 (1987) 132711v.
- 850 Marzo, A., Pasotti, V. and Maggi, G.C.: Gas chromatographic evaluation of trapidil and its desethyl metabolite in biological fluids for pharmacokinetic and bioavailability investigations. *Arzneim.-Forsch.*, 37 (1987) 947-950; *C.A.*, 107 (1987) 168138c.
- 851 Mauer, H. and Pflieger, K.: Identification and differentiation of benzodiazepines and their metabolites in urine by computerized gas chromatography-mass spectrometry. *J. Chromatogr.*, 422 (1987) 85-101.
- 852 Morris, W.J., Nandrea, G.J., Roybal, J.E., Munns, R.K., Shimoda, W. and Skinner, H.R., Jr.: Quantitative confirmation of dimetridazole and ipronidazole in swine feed by capillary gas chromatography/mass spectrometry with multiple ion detection. *J. Assoc. Off. Anal. Chem.*, 70 (1987) 630-634.

- 853 Zeng Su, Yan Jixiang, Shi Huizhen and Liu Zhiqiang: (Determination of hypnotic and sedative drugs in the blood and liver by using activated enzymic digestion methods). *Yaowu Fenxi Zazhi*, 7 (1987) 209-212; *C.A.*, 107 (1987) 211360h.

See also 750, 874.

32g. Other drug categories

- 854 Kennedy, J.S., Leduc, B.W., Scavone, J.M., Harmatz, J.S., Shader, R.I. and Greenblatt, D.J.: Pharmacokinetics of an intravenous caffeine: comparison of high-performance liquid chromatographic and gas chromatographic methods. *J. Chromatogr.*, 422 (1987) 274-280.

32h. Toxicological and forensic applications

- 855 Anderson, W.H. and Fuller, D.C.: A simplified procedure for the isolation, characterization, and identification of weak acid and neutral drugs from whole blood. *J. Anal. Toxicol.*, 11 (1987) 198-204; *C.A.*, 107 (1987) 213084v.
- 856 Beckstead, H.D. and Neville, G.A.: A novel application of headspace and GC-FT-IR analysis to identify the ethyl acetate complex of O⁶-acetylmorphine. *J. Can. Soc. Forensic Sci.*, 20 (1987) 71-76; *C.A.*, 107 (1987) 192403f.
- 857 Christophersen, A.S., Biseth, A., Skuterud, B. and Gadeholt, G.: Identification of opiates in urine by capillary column gas chromatography of two different derivatives. *J. Chromatogr.*, 422 (1987) 117-124.
- 858 Hodgson, B.T. and Taylor, M.D.: Evaluation of Fisher Toxicchem whole blood alcohol controls (human). *J.-Can. Soc. Forensic Sci.*, 20 (1987) 87-91; *C.A.*, 107 (1987) 148681w.
- 859 Lora-Tamayo, C., Tena, T. and Tena, G.: Concentration of free and conjugated morphine in blood in twenty cases of heroin-related deaths. *J. Chromatogr.*, 422 (1987) 267-273.
- 860 Mueller, G., Verkoyen, C., Soton, N. and Norpoth, K.: Urinary excretion of acrylonitrile and its metabolites in rats. *Arch. Toxicol.*, 60 (1987) 464-466; *C.A.*, 107 (1987) 192561f.
- 861 Ritchie, L.K., Caplan, Y.H. and Park, J.: Δ^9 -Tetrahydrocannabinol analysis in forensic blood specimens using capillary column gas chromatography with nitrogen sensitive detection. *J. Anal. Toxicol.*, 11 (1987) 205-209; *C.A.*, 107 (1987) 192405h.
- 862 Spiehler, V. and Brown, R.: Unconjugated morphine in blood by radioimmunoassay and gas chromatography/mass spectrometry. *J. Forensic Sci.*, 32 (1987) 906-916; *C.A.*, 107 (1987) 148699h.
- 863 Watts, V.W. and Simonick, T.F.: (A retention index library for commonly encountered drugs and metabolites using tri-n-alkylamines as reference compounds, nitrogen-phosphorus detectors, and dual capillary chromatography). *J. Anal. Toxicol.*, 11 (1987) 210-214; *C.A.*, 107 (1987) 170058v.

See also 597, 600, 611, 756, 757, 773, 925.

32i. Plant extracts

- 864 Krzek, J.: (Determination of volatile oil in composed formulations by gas chromatography). *Farm. Pol.*, 42 (1986) 479-483; *C.A.*, 107 (1987) 183663p.
- 865 Kuraishi, S., Ohara, T. and Sakurai, N.: Gas chromatography-flame thermionic detector analysis of cytokinins in etiolated squash seedlings using ¹⁴C-BA as an internal standard. *Plant Cell Physiol.*, 28 (1987) 1033-1041; *C.A.*, 107 (1987) 194389e.

See also 790.

33. CLINICO-CHEMICAL APPLICATIONS

33a. General papers and reviews

- 866 Wolthers, B.G., Muskier, F.A.J., Pavel, S., Keijzer, J.J., Koopman, B.J., van den Berg, G.A., Dikkeschei, L.D., Berger, R. and Reijngoud, D.J.: Use of gas chromatography-mass spectrometry in clinical chemistry: a review of experiences gained during the past twelve years at the University Hospital of Groningen. *Tijdschr. Ned. Ver. Klin. Chem.*, 12 (1987) 137-153; *C.A.*, 107 (1987) 171619r.

See also 440, 839.

33b. Complex mixtures and profiling

- 867 Bennett, M.J., Worthy, E. and Pollitt, R.J.: Chromatographic diagnosis of inherited diseases of organic acid metabolism. *Anal. Proc.*, 24 (1987) 322-333.
- 868 Dietel, P., Hölzel, C., Schneider, K. and Spitteller, G.: Excretion of urinary acids during inverted sleep-waking rhythm. *J. Chromatogr.*, 422 (1987) 1-12.
- 869 Estrangin, E., Thiers, B. and Peloux, Y.: (Micromethods and gaseous phase chromatography-mass spectrometry of carboxylic acids, products of glucose fermentation, in the identification of corynebacteria). *Ann. Biol. Clin. (Paris)*, 45 (1987) 285-289; *C.A.*, 107 (1987) 194839b.
- 870 French, G.L., Chan, C.Y., Cheung, S.W. and Oo, K.T.: Diagnosis of pulmonary tuberculosis by detection of tuberculostearic acid in sputum by using gas chromatography-mass spectrometry with selected ion monitoring. *J. Infect. Dis.*, 156 (1987) 356-362; *C.A.*, 107 (1987) 130444m.
- 871 Garrigues, M.L. and Veron, M.: (Contribution of gas-liquid chromatography to microbiological diagnosis). *Ann. Biol. Clin. (Paris)*, 45 (1987) 135-143; *C.A.*, 107 (1987) 194318f.
- 872 Giachetti, C., Poletti, P. and Zanolo, G.: Analysis of calcium blocker drugs in plasma, HRGC and HPLC analytical conditions for pharmacokinetic studies. *J. High Resolut. Chromatogr. Chromatogr. Commun.*, 10 (1987) 654-658.
- 873 Han Liwei and Yu Gue: (Quantitative analysis of free fatty acids for sclerema neomatorum by gas chromatography). *Sepu*, 5 (1987) 365-366.
- 874 Maylin, G.A., Dewey, E.A. and Henion, J.D.: Determination of drug in equine urine by gas chromatography-Fourier transform infrared spectroscopy. *LC-GC*, 5 (1987) 904-914.
- 875 Parnetti, L., Gottfries, J., Karlsson, I., Laengstroem, G., Gottfries, C.G. and Svennerholm, L.: Monoamines and their metabolites in cerebrospinal fluid of patients with senile dementia of Alzheimer type using high performance liquid chromatography and gas chromatography-mass spectrometry. *Acta Psychiatr. Scand.*, 75 (1987) 542-548; *C.A.*, 107 (1987) 131973v.
- 876 Phillips, M. and Greenberg, J.: Detection of endogenous acetone in normal human breath. *J. Chromatogr.*, 422 (1987) 235-238.
- 877 Taylor, A.J., Jennings, P.E., Barnett, A.H., Pandov, H.I. and Lawson, N.: An alternative explanation for the change in erythrocyte fatty acids observed in diabetes. *Clin. Chem. (Winston-Salem)*, 33 (1987) 2083-2085.
- 878 Tuchman, M., Aury-Blais, C., Ramnaraine, M.L.R., Neglia, J., Krivit, W. and Lemieux, B.: Determination of urinary homovanillic and vanillylmandelic acids from dried filter paper samples: assessment of potential methods for neuroblastoma screening. *Clin. Biochem. (Ottawa)*, 20 (1987) 173-177; *C.A.*, 107 (1987) 171749h.
- 879 Uehori, R., Nagata, T., Kimura, K., Kudo, K. and Noda, M.: Screening of volatile compounds present in human blood using retention indices in gas chromatography. *J. Chromatogr.*, 411 (1987) 251-257.
- 880 Zhou Fang, Zhu Houchu, Tang Guangjiang, Wang Yizheng, Zhang Xutuan, Gao Shude, Qiu Mingqing, Zhong Wenpeng and Zhao Lin: Discrimination of *Moraxella* and *Legionella pneumophila* by gas chromatography of whole-cell fatty acids. *Sci. Sin., Ser. B*, 30 (1987) 616-624; *C.A.*, 107 (1987) 194524v.

See also 617, 620, 652, 656, 702, 723.

34. FOOD ANALYSIS

34a. General papers and reviews

- 881 Firestone, D.: Control of olive oil adulteration and misbranding in the United States. *Riv. Ital. Sost. Grasse*, 64 (1987) 293-297.
- 882 Lawrence, J.F.: Recent development in derivatization for the chromatographic determination of food additives. *J. Chromatogr. Sci.*, 25 (1987) 444-452 - a review with 97 refs.
- 883 Mariani, C.: (The problems of quality control). *Riv. Ital. Sost. Grasse*, 64, No. 8 (1987) 325-333.
- 884 Seher, A. and Arens, M.: Gemeinschaftsarbeiten der DGF. 100. Mitt.: Deutsche Einheitsmethoden zur Untersuchung von Fetten, Fettprodukten und verwandten Stoffen, 77. Mitt.: Analyse von Fetten. XXIV. Gas chromatographie: Analyse von Fettsäureestern. *Fett Wiss. Technol.-Fat Sci. Technol.*, 89 (1987) 307-311.

34b. Complex mixtures

- 885 Aishima, T. and Nakai, S.: Pattern recognition of GC profiles for classification of cheese variety. *J. Food Sci.*, 52 (1987) 939-942; *C.A.*, 107 (1987) 132762n.
- 886 Bocchio, F. and Fellous, R.: Analyse de donnees appliquee aux profils chromatographiques de fraises. *Parf. Cosmet. Aromes*, No. 76 (1987) 71-78.
- 887 Hu Guodong: (Headspace analysis and its application to determination of flavor components in alcoholic beverages). *Shipin Yu Fajiao Gongye*, No. 4 (1987) 51-57; *C.A.*, 107 (1987) 152711y.
- 888 Krajewska, A.M. and Powers, J.J.: Gas chromatographic determination of capsaicinoids in green *Capsicum* fruits. *J. Assoc. Off. Anal. Chem.*, 70 (1987) 926-928.
- 889 Li Maosheng, Shao Lingxian and Jin Xinrong: (Investigation of the volatile constituents of grape must and grape wine by GC/MS/computer). *Seppu*, 5 (1987) 359-362.
- 890 Nijssen, L.M. and Maarse, H.: Authenticity studies of fruit juice concentrates using a GC-fingerprinting technique. *Int. Fruchtsaft-Union, Wiss.-Tech. Komm.*, (Ber.), 19 (1986) 333-355; *C.A.*, 107 (1987) 216243g.
- 891 Ohta, H., Kinjo, S. and Osajima, Y.: Glass capillary gas chromatographic analysis of volatile components of canned Philippine pineapple juice. *J. Chromatogr.*, 409 (1987) 409-412.
- 892 Ortega, J., Martin, I. and Martin, A.: (Porous polymers use for analysis of wines' volatile compounds). *An. Bromatol.*, 39 (1987) 147-160; *C.A.*, 107 (1987) 196433p.
- 893 Reddy, B.S. and Sarojini, G.: Chemical and nutritional evaluation of chili (*Capsicum annum*) seed oil. *J. Am. Oil Chem. Soc.*, 64 (1987) 1419-1423.
- 894 Sugiyama, N., Saito, K. and Sato, H.: (Discrimination of natural food thickeners with the pyrolysis capillary gas chromatograph-mass spectrometer). *Nippon Shokuhin Kogyo Gakkaishi*, 34 (1987) 370-375; *C.A.*, 107 (1987) 196580j.
- 895 Tonogai, Y., Ando, T., Tsumura, A. and Ito, Y.: Determination of rosin ester gum emulsifiers in fruit juices by gas chromatography. *J. Food Prot.*, 50 (1987) 464-467; *C.A.*, 107 (1987) 174487g.
- 896 Vercellotti, J.R., Kuan, J.C.W., Liu, R.H., Legendre, M.G., St. Angelo, A.J. and Dupuy, H.P.: Analysis of volatile heteroatomic meat flavor principles by purge-and-trap/gas chromatography-mass spectrometry. *J. Agric. Food Chem.*, 35 (1987) 1030-1035.
- 897 Waimaleongora-Ek, C., Chen, T.C. and Alley, E.G.: (Headspace volatiles of heated chicken-drying shortening). *Poult. Sci.*, 66 (1987) 1071-1074; *C.A.*, 107 (1987) 153084h.
- 898 Yasuhara, A.: Comparison of volatile components between fresh and rotten mussels by gas chromatography-mass spectrometry. *J. Chromatogr.*, 409 (1987) 251-258.
- See also 622, 645, 646, 648, 651, 655, 664, 668, 673, 679, 680, 688, 691, 701, 713, 716, 745, 756, 761, 771, 775, 794, 801, 807, 809, 814, 815, 852, 915.

35. ENVIRONMENTAL ANALYSIS

35b. Air pollution (complex mixtures)

- 899 Babina, M.D.: (Gas chromatographic determination of isophen (acrex) in workplace air). *Gig. Tr. Prof. Zabol.*, No. 8 (1987) 57-58; *C.A.*, 107 (1987) 222263e.
- 900 Chmil, V.D., Pogorelyi, V.K., Samodumova, I.M. and Kiseleva, L.I.: (Gas-chromatographic determination of Counter in air). *U.S.S.R. Pat.* SU 1,314,257 (Cl. G01N30/04), 30 May 1987, Appl. 3,940,025, 26 Jul. 1985; *C.A.*, 107 (1987) 129120w.
- 901 Choudhary, G.: Determination of 2,3-dibromopropanol in air. *Am. Ind. Hyg. Assoc. J.*, 48 (1987) 809-813; *C.A.*, 107 (1987) 204155s.
- 902 Fukabori, S. and Nakaaki, K.: (Gas chromatographic determination of 2,4- and 2,6-tolylene diisocyanate). *Rodo Kazaku*, 62 (1986) 591-610; *C.A.*, 107 (1987) 140024p.
- 903 Williams, D.T., Shewchuck, C., Lebel, G.L. and Muir, N.: Diazinon levels in indoor air after periodic application for insect control. *Am. Ind. Hyg. Assoc. J.*, 48 (1987) 780-785; *C.A.*, 107 (1987) 204154r.
- 904 Zaranski, M.T. and Bidleman, T.F.: High-volume elution chromatography of dichlorobenzenes on a polyurethane foam-Tenax sandwich cartridge. *J. Chromatogr.*, 409 (1987) 235-242.

See also 491, 565, 567, 604, 606, 613, 632, 689, 752.

35c. Soil pollution (complex mixtures)

- 905 Butler, A.C. and Sibbald, R.R.: Sampling and GC-FID, GC/MS analysis of petroleum hydrocarbons in the ocean surface microlayer off Richards Bay, South Africa. *Estuarine, Coastal Shelf Sci.*, 25 (1987) 27-42; *C.A.*, 107 (1987) 183044n.
- 906 Johnsen, P.B. and Kuan Jui Chang, W.: Simplified method to quantify geosmin and 2-methylisoborneol concentrations in water and microbiological cultures. *J. Chromatogr.*, 409 (1987) 337-342.
- 907 Kozeiko, T.A., Aizenshtadt, M.A., Maktaz, E.D. and Kasman, V.S.: Comparative assessment of gas chromatographic and photometric methods for determination of chloropicrin in water. *Gig. Sanit.*, No. 7 (1987) 41-42; *C.A.*, 107 (1987) 204787z.
- 908 Mestres, J.P., Spiliopoulos, C., Mestres, R. and Cooper, J.F.: Transport of organic contaminants in groundwater: gas chromatography model to forecast the significance as applied to aldicarb sulfone residues. *Arch. Environ. Contam. Toxicol.*, 16 (1987) 649-656; *C.A.*, 107 (1987) 149203d.
- 909 Onodera, S., Nishikawa, T. and Suzuki, S.: Chemical changes of organic compounds in chlorinated water. XIV. Characterization and determination of halogenated organics formed during chlorination of water from the Tama river. *J. Chromatogr.*, 409 (1987) 259-270.
- 910 Rossel, D., Honsberger, P. and Tarradellas, J.: Bioaccumulative behaviour of some PCB congeners in Lake Geneva brown trout (*Salmo trutta lacustris* L.). *Int. J. Environ. Anal. Chem.*, 31 (1987) 219-233.
- 911 Subra, P., Hennion, M.C., Foucault, A. and Rosset, R.: Analysis of organic mater in water of low TOC (total organic carbon) content by chromatographic techniques. *Int. J. Environ. Anal. Chem.*, 31 (1987) 197-217.

See also 610, 625, 804, 813, 923, 934, 939.

35d. Soil pollution (complex mixtures)

See 615, 624, 796.

36. SOME TECHNICAL PRODUCTS AND COMPLEX MIXTURES

36a. *Surfactants*

- 912 Dahlgran, J.R. and Shingleton, C.R.: Determination of ethylene oxide in ethoxylated surfactants and demulsifiers by headspace gas chromatography. *J. Assoc. Off. Anal. Chem.*, 70 (1987) 796-798.

See also 525.

36b. *Antioxidants and preservatives*

See 679.

36c. *Various technical products*

- 913 Douse, J.M.F.: Improved method for the trace analysis of explosives by silica capillary column gas chromatography with thermal energy analysis detection. *J. Chromatogr.*, 410 (1987) 181-189.
- 914 Gizzatullin, R.R., Tukmanov, R.G. and Matrosova, N.N.: (Determination of impurities in ethylbenzene feed by an improved method). *Neftepererab. Neftekhim. (Moscow)*, No. 9 (1987) 31-32; *C.A.*, 107 (1987) 219449q.
- 915 Gramiccioni, L., Milana, M.R. and Di Marzio, S.: (PET containers for carbonated beverages: a gas chromatography determination of free dimethyl terephthalate (DMT)). *Rass. Chim.*, 38 (1986) 255-258; *C.A.*, 107 (1987) 132736g.
- 916 Harvey, T.G., Matheson, T.W. and Pratt, K.C.: Extraction and identification of nitrile compounds in Rundle shale oil. *J. Chromatogr.*, 435 (1988) 193-198.
- 917 Kholostova, G.G., Shimonae, G.S. and Burova, M.O.: (Quality control of petroleum products by chromatographic distillation). *Khim. Tekhnol. Topl. Masel*, No. 8 (1987) 3-5; *C.A.*, 107 (1987) 157833z.
- 918 Tatsuka, K., Kohama, M., Suekane, S. and Mori, D.: Volatile components in kamaboko (seasoned steamed minced fish paste) and its supporting wooden plate by gas chromatography mass spectrometry. *Nippon Noei Kagaku Kaishi*, 61 (1987) 587-598; *C.A.*, 107 (1987) 196685x.
- 919 Thoma, H. and Hutzinger, O.: Pyrolysis and GC/MS analysis of brominated flame retardants in online operation. *Chemosphere*, 16 (1987) 1353-1360; *C.A.*, 107 (1987) 135211t.

See also 463, 552, 595, 602, 619, 631, 632, 644, 659, 690, 720, 777, 821.

36d. *Complex mixtures and unidentified compounds*

- 920 Bicchi, C., D'Amato, A., Nano, G.M. and Lorenzi, M.C.: (Direct-capture determination of the volatile compounds emitted by plants and animals). *Essenze Deriv. Agrum.*, 57 (1987) 100-111; *C.A.*, 107 (1987) 214304d.
- 921 Cechi, G. and Bonfand, A.: Conversion des huiles vegetales en carburants potentiels. Essais preliminaires. *Rev. Fr. Corps Gras*, 34 (1987) 397-401.

See also 608, 681.

37. CELLS, CELLULAR PARTICLES AND SUPRAMOLECULAR STRUCTURES

- 922 Sllers, C.S., Mar, T., Bertsch, W. and Crewe, R.M.: Investigation of African honeybee races. *J. High Resolut. Chromatogr. Chromatogr. Commun.*, 10 (1987) 674-675.

See also 626, 653, 672, 678, 684, 704, 869, 880.

38. INORGANIC COMPOUNDS

38a. Cations

See 579.

38b. Anions

- 923 Hashimoto, S., Fujiwara, K. and Fuwa, K.: Determination of phosphorus in natural water using hydride generation and gas chromatography. *Limnol. Oceanogr.*, 32 (1987) 729-735; *C.A.*, 107 (1987) 183131p.
- 924 Mitsuhashi, T., Adechi, F. and Kaneda, Y.: (Determination of total bromine in cereals by gas chromatography). *Shokuhin Eiseigaku Zasshi*, 28 (1987) 130-135; *C.A.*, 107 (1987) 174514p.
- 925 Yamano, Y., Ito, I., Nagao, N. and Ishizu, S.: (A simple determination method of bromide ion in plasma of methyl bromide workers by head space gas chromatography). *Sengyo Igaku*, 29 (1987) 196-201; *C.A.*, 107 (1987) 182625r.

38c. Permanent and rare gases

- 926 Huang, D.: (Simultaneous determination of carbon monoxide and total hydrocarbon by gas chromatography). *Huanjing Huaxue*, 6, No. 3 (1987) 56-61; *C.A.*, 107 (1987) 222296t.
- 927 Lei Wenmin and Wu Daxin: (Improved variable-temperature enrichment gas chromatography for determination of trace impurities in ultrapure hydrogen). *Seppu*, 5 (1987) 266-267.
- 928 Saprykin, Yu.A., Golovko, B.M., Pazderskii, Yu.A. and Zorin, A.D.: (Chromatographic determination of hydrogen in silicon chlorides using a high-frequency emission detector). *Zh. Anal. Khim.*, 42 (1987) 2017-2021.
- 929 Semiletov, I.P. and Radaev, E.F.: (Device and method for analyzing gas inclusions in ice). *U.S.S.R. Pat. SU 1,323,948 (Cl.G01N30/06)*, 15 Jul. 1987, Appl. 3,815,113, 20 Nov. 1984; *C.A.*, 107 (1987) 211192e.
- 930 Sugisaki, R. and Taki, K.: Simplified determination of helium, neon, and argon dissolved in natural waters. *Geochem. J.*, 21 (1987) 23-27; *C.A.*, 107 (1987) 183116n.

See also 488, 601, 941.

38d. Volatile inorganic compounds

- 931 Cheng Weiwei, Ma Guilan and Zhu Xinlong: (Determination of microamount of sulfur fluoride oxide (S₂F₁₀O) in industrial sulfur hexafluoride by gas chromatography). *Fenxi Huaxue*, 15 (1987) 588-591; *C.A.*, 107 (1987) 189882f.
- 932 Gao Xiuying, Ji Lige, Wu Jingke, Tan Zhiyi, Yun Zihou and Jiao Shuping: (A gas chromatographic device used in the determination of carbon dioxide and some application in plant physiology). *Beijing Linye Daxue Xuebao*, 9 (1987) 182-185; *C.A.*, 107 (1987) 150489h.
- 933 He Luqing, Cao Lei and Gao Guangjin: (Gas chromatographic determination of microamount of elemental sulfur in minerals). *Zhongnan Kuangye Xueyuan Xuebao*, No. 6 (1986) 81-85; *C.A.*, 107 (1987) 211100y.
- 934 Kim Ki-Hyun and Andreae, M.O.: Determination of carbon disulfide in natural waters by adsorbent preconcentration and gas chromatography with flame photometric detection. *Anal. Chem.*, 59 (1987) 2670-2673.
- 935 Krylov, V.A., Lazarev, V.A., Sokolova, G.V. and Krasotskii, S.G.: (Gas-chromatographic analysis of phosphorus trichloride and oxychloride of high purities). *Vysokochist. Veshchestva*, No. 2 (1987) 159-162; *C.A.*, 107 (1987) 189782y.
- 936 Scheffrahn, R.H., Osbrink, W.L.A., Hsu Ruei Ching and Su Nan Yao: Desorption of residual sulfonyl fluoride from structural and household commodities by headspace analysis using gas chromatography. *Bull. Environ. Contam. Toxicol.*, 39 (1987) 769-775; *C.A.*, 107 (1987) 213578j.
- 937 Tanaka, E., Nakamura, T., Terada, M., Misawa, S., Suzuki, Y. and Kuroiwa, Y.: (Determination of hydrogen sulfide in fluid and tissue samples by gas chromatography with a flame photometric detector). *Eisei Kagaku*, 33 (1987) 149-152; *C.A.*, 107 (1987) 170038p.

- 938 Vishnevskii, V.Yu., Beskova, G.S. and Butuzova, A.I.: (Determination of nitrogen dioxide by gas chromatography). *Zh. Anal. Khim.*, 42 (1987) 2022-2026.
- 939 Yamamoto, K. and Tsuji, H.: (Determination of hydrogen sulfide in water by stripping head-space GC). *Bunseki Kagaku*, 36 (1987) 830-835.

See also 488, 601.

39. RADIOACTIVE AND OTHER ISOTOPE COMPOUNDS

- 940 Lo Jiunn Guang, Chao Jiunn Hsing and Chuang Jui Tang: Preparation of labelled ^{38}Cl hydrocarbons by recoil ^{38}Cl atom reaction and detection by radio-gas chromatographic techniques. *Appl. Radiat. Isot.*, 38 (1987) 685-688; *C.A.*, 107 (1987) 185247e.
- 941 Parbuzin, V.C., Leshchev, Yu.A., Luk'yanov, A.A. and Shcherbakova, K.D.: (Gas chromatographic method of investigation of selective adsorption of hydrogen isotope at higher pore filling). *Zh. Fiz. Khim.*, 61 (1987) 3077-3078.

Planar Chromatography

1. REVIEWS AND BOOKS

- 263 Brinkman, U.A.T.: Thin-layer chromatography on chemically bonded phases. In: R.E. Kaiser (Editor), *Proceedings of the 3rd International Symposium on Instrumental High Performance Thin-Layer Chromatography, Wuerzburg, April 17-19, 1985*, Inst. Chromatogr., Bad Duerkheim, 1985, pp. 25-35; *C.A.*, 107 (1987) 197139j - a review with 21 refs.
- 264 Hattori, K.: (Host-guest chromatography). *Bio Ind.*, 4 (1987) 327-338; *C.A.*, 107 (1987) 167984p - a review with 75 refs.
- 265 Kubeczka, K.-H.: Analyse von Naturstoffen. Neue Entwicklungen. *Dtsch. Apoth.-Ztg.*, 127 (1987) 2443-2450 - a review with 45 refs.
- 266 Szejtli, J., Zsados, B. and Cserhati, T.: Cyclodextrin use in separations. *ACS Symp. Ser.*, 342 (Ordered Media Chem. Sep.) (1987) 200-217; *C.A.*, 107 (1987) 161442p - a review with 71 refs.
- 267 Zhang, Z.: (Thin-layer chromatography on chemically bonded phases). *Sepu*, 5 (1987) 225-231; *C.A.*, 107 (1987) 190014f - a review with 43 refs.

See also 277, 307, 328, 402, 429, 431, 496.

2. FUNDAMENTALS, THEORY AND GENERAL

2a. General

- 268 Steinbrunner, J.E., Malik, D.J. and Nurok, D.: Contour diagrams for the presentation of separation quality in two-dimensional thin layer chromatography. *J. High Resolut. Chromatogr. Chromatogr. Commun.*, 10 (1987) 560-563.

2b. Thermodynamics and theoretical relationships

- 269 Nurok, D., Habibi-Goudarzi, S. and Kleyle, R.: Statistical approach to solvent selection as applied to two-dimensional thin-layer chromatography. *Anal. Chem.*, 59 (1987) 2424-2428.
- 270 Rozylo, J.K. and Kolodziejczyk, H.: Evaluation of the separation obtained in thin-layer chromatographic systems using binary and ternary mobile phases. *Symp. Biol. Hung.*, 34 (Chromatography '85) (1986) 629-637; *C.A.*, 107 (1987) 211244y.

See also 271, 415.

2d. Measurement of physico-chemical and related values

- 271 Kubota, T., Kano, K., Kawata, E., Hashizume, H. and Uno, B.: (New method for determining true partition coefficients by the technique of chemically modified reversed-phase high-performance thin-layer chromatography). *Gifu Yakka Daigaku Kiyo*, No. 36 (1987) 32-41; *C.A.*, 107 (1987) 184358e.
- 272 Wu, F.: (R_s value of the solution and its application). *Sepu*, 5 (1987) 12-16; *C.A.*, 107 (1987) 168009m.

See also 492.

3. GENERAL TECHNIQUES

3c. Sorbents, carriers, column and layer performance, packing procedures

- 273 Janos, E. and Cserhati, T.: Preadsorbent salts to modify the retention behavior of silica. *Symp. Biol. Hung.*, 34 (Chromatography '85) (1986) 461-472; *C.A.*, 107 (1987) 228102y.
- 274 Ohta, M., Okamoto, M., Futamura, A., Gotoh, S. and Tamaoki, S.: (Evaluation of 3-cyanopropyltrichlorosilane treated thin layer chromatographic plate. Application of analysis for typical steroids in ointment). *Byoin Yakugaku*, 13 (1987) 111-115; *C.A.*, 107 (1987) 161776a.
- 275 Oliveira dos Santos, W.: (Polyamide particles for use in thin-layer chromatography). *Braz. Pedido PI Pat.* BR 85 04,803 (Cl. CO8J3/12), 05 May 1987, Appl. 85/4,803, 30 Sep. 1985; *C.A.*, 107 (1987) 218866m.

See also 263, 508.

3g. High performance procedures

See 265, 271.

4. SPECIAL TECHNIQUES

4a. Automation and computerization

- 276 De la Vigne, U.: Computer controlled evaluation of thin-layer chromatograms and electrophoresis objects. *Symp. Biol. Hung.*, 34 (Chromatography '85) (1986) 591-603; *C.A.*, 107 (1987) 150724f.

See also 269.

4g. Separation of enantiomers

- 277 Caldwell, J. and Testa, B.: Criteria for the acceptability of experimental evidence for the enantiomeric composition of xenobiotics and their metabolites. *Drug Metab. Disp.*, 15 (1987) 587-588 - a review.

See also 264, 323.

4h. Other special techniques

- 278 Tyihak, E.: Overpressured layer chromatography and its applicability in pharmaceutical and biomedical analysis. *J. Pharm. Biomed. Anal.*, 5 (1987) 191-203; *C.A.*, 107 (1987) 150479e.
- 279 Vajda, J., Pick, J. and Leisztner, L.: On the additivity of isocratic OPLC data for retention prediction in multistep gradient OPLC. *Symp. Biol. Hung.*, 34 (Chromatography '85) (1986) 605-616; *C.A.*, 107 (1987) 211243x.

5. HYDROCARBONS AND HALOGEN DERIVATIVES

5a. Aliphatic hydrocarbons

See 339.

5b. Cyclic hydrocarbons

- 280 Egestad, B., Pettersson, P., Sjövall, J., Rafter, J., Hyvönen, K. and Gustafsson, J.Å.: Studies on the chromatographic fractionation of metabolites of benzo[a]pyrene in faeces and urine from germfree and conventional rats. *Biomed. Chromatogr.*, 2 (1987) 120-134.

- 281 McKay, S., Farmer, P.B., Cary, P.D. and Grover, P.L.: The metabolism of 7-ethylbenz[*a*]anthracene by rat liver microsomal preparations. *Drug Metab. Disp.*, 15 (1987) 682-694.
- 282 Vo Dinh, T. and Alak, A.: Enhanced room-temperature phosphorescence of anthracene on cyclodextrin-treated filter paper. *Appl. Spectrosc.*, 41 (1987) 963-966; *C.A.*, 107 (1987) 168034r.

See also 270, 273, 292.

5d. *Complex hydrocarbon mixtures*

- 283 Dickakian, G.B.: Improved chromatographic method for determining fouling tendency of liquid hydrocarbons. *Eur. Pat. Appl. EP 234,857 (Cl. GOIN30/90)*, 02 Sep. 1987, US Appl. 830,386, 18 Feb. 1986; *C.A.*, 107 (1987) 179882s.

6. ALCOHOLS

See 320, 321, 339, 501.

7. PHENOLS

- 284 Akkuratova, O.L., Kuk'yanova, L.K., Spitsyna, A.V. and Bryukhovetskaya, L.V.: (Analysis of phenolic alcohols by thin-layer chromatography). *Plast. Massy*, No. 8 (1987) 50; *C.A.*, 107 (1987) 218483j.

See also 498.

8. SUBSTANCES CONTAINING HETEROCYCLIC OXYGEN

8a. *Flavonoids*

See 265, 456, 498, 501.

8b. *Aflatoxins and other mycotoxins*

- 285 Adensam, L., Lebedova, M. and Turek, B.: (The determination of very low aflatoxin concentrations). *Cesk. Hyg.*, 32 (1987) 362-367; *C.A.*, 107 (1987) 234954u.
- 286 Francis, O.J., Jr., Ware, G.M., Carman, A.S., Kirschenheuter, G.P., Kuan, S.S. and Newell, R.F.: Thin-layer chromatographic determination of sterigmatocystin in cheese: interlaboratory study. *J. Assoc. Off. Anal. Chem.*, 70 (1987) 842-844.
- 287 Kawano, M.: (Forensic toxicological studies on *Lampteromyces* poisoning. Identification of illudin S (lampterol)). *Igaku Kenkyu*, 56, No. 1 (1986) 31-34; *C.A.*, 107 (1987) 148703e.
- 288 Rati, E.R., Prema, V. and Shantha, T.: Modification of Pons's method of estimating aflatoxin B₁ in corn, groundnut and groundnut cake. *J. Food Sci. Technol.*, 24, No. 2 (1987) 90-91; *C.A.*, 107 (1987) 152962f.
- 289 Simonella, A., Torreti, L., Filipponi, C., Falgiani, A. and Ambrosii, L.: Simultaneous determination of aflatoxins G₁, B₁, G₂, B₂, in animal feedstuffs by HPTLC and RP-HPLC. *J. High Resolut. Chromatogr. Chromatogr. Commun.*, 10 (1987) 626-628.

8c. *Other compounds with heterocyclic oxygen (including tannins)*

- 290 Glowniak, K., Wawrzynowicz, T. and Soczewinski, E.: Isolation of coumarins from the fruits of *Patinaca sativa* L. by column chromatography and modified thin-layer chromatography. *Symp. Biol. Hung.*, 34 (Chromatography '85) (1986) 251-261; *C.A.*, 107 (1987) 161764v.

- 291 Kernoczi, Z., Hethelyi, E., Danos, B. and Tetenyi, P.: Presence of carlina oxide in plants of Hungary. Stabilization and antimicrobial effect. *Acta Pharm. Hung.*, 57 (1987) 171-181; *C.A.*, 107 (1987) 223119z.
- 292 Shen, E.S. and Olson, J.R.: Relationship between the murine phenotype and the hepatic uptake and metabolism of 2,3,7,8-tetrachlorodibenzo-*p*-dioxin. *Drug. Metab. Disp.*, 15 (1987) 653-660.

See also 447.

9. OXO COMPOUNDS, ETHERS EPOXIDES AND QUINONES

See 309, 385, 498.

10. CARBOHYDRATES

10a. Mono and oligosaccharides. Structural studies

- 293 Allegretti, M., Ambrosoli, G. and Cantoni, C.: (Detection of isoglucose in honey). *Ind. Aliment. (Pinerolo)*, 26 (1987) 566-573; *C.A.*, 107 (1987) 216258r.
- 294 Altman, E., Brisson, J.-R. and Perry, M.B.: Structure of the capsular polysaccharide of *Haemophilus pleuropneumoniae* serotype 5. *Eur. J. Biochem.*, 170 (1987) 185-192.
- 295 Brassart, D., Baussant, T., Wieruszeski, J.-M., Strecker, G., Montreuil, J. and Michalski, J.-C.: Catabolism of N-glycosylprotein glycans: evidence for a degradation pathway of sialylglyco-asparagines resulting from the combined action of the lysosomal aspartylglucosaminidase and endo-N-acetyl- β -D-glucosaminidase. A 400-MHz $^1\text{H-NMR}$ study. *Eur. J. Biochem.*, 169 (1987) 131-136.
- 296 Gross, H.J., Bünsch, A., Paulson, J.C. and Brossmer, R.: Activation and transfer of novel synthetic 9-substituted sialic acids. *Eur. J. Biochem.*, 168 (1987) 595-602.
- 297 Hughes, R.C., Foddy, L. and Bause, E.: Asparagine-linked oligosaccharides of BHK cells treated with inhibitors of oligosaccharide processing. *Biochem. J.*, 247 (1987) 537-545.
- 298 Leclercq, Y., Sawatzki, G., Wieruszeski, J.-M., Montreuil, J. and Spik, G.: Primary structure of the glycans from mouse serum and milk transferrins. *Biochem. J.*, 247 (1987) 571-578.
- 299 Mann, B.: Identification of carbohydrate gums in dental cream by thin-layer chromatography. *J. Chromatogr.*, 407 (1987) 369-376.
- 300 McDowell, W., Grier, T.J., Rasmussen, J.R. and Schwarz, R.T.: The role of C-4-substituted mannose analogues in protein glycosylation. Effect of the guanosine diphosphate esters of 4-deoxy-4-fluoro-D-mannose and 4-deoxy-D-mannose on lipid-linked oligosaccharide assembly. *Biochem. J.*, 248 (1987) 523-531.
- 301 Mikheeva, M.N., Grigor'eva, T.V., Shmel'kova, S.A. and Brutko, L.I.: (Identification of glucose and mannitol in cardioplegic solutions). *Farmatsiya (Moscow)*, 36, No. 5 (1987) 66-67; *C.A.*, 107 (1987) 223375e.
- 302 Ray, T.C., Smith, A.R.W., Wait, R. and Hignett, R.C.: Structure of the sidechain of lipopolysaccharide from *Erwinia amylovora* T. *Eur. J. Biochem.*, 170 (1987) 357-361.
- 303 Scudder, P., Lawson, A.M., Hounsell, E.F., Carruthers, R.A., Childs, R.A. and Feizi, T.: Characterisation of oligosaccharides released from human-blood-group O erythrocyte glycopeptides by the endo- β -galactosidase of *Bacteroides fragilis*. A study of the enzyme susceptibility of branched poly(N-acetylglucosamine) structures. *Eur. J. Biochem.*, 168 (1987) 585-593.
- 304 Shailubhai, K., Pratta, M.A. and Vijay, I.K.: Purification and characterization of glucosidase I involved in N-linked glycoprotein processing in bovine mammary gland. *Biochem. J.*, 247 (1987) 555-562.
- 305 Song, Z., Li, S.-C. and Li, Y.-T.: Absence of endo- β -N-acetylglucosaminidase activity in the kidneys of sheep, cattle and pig. *Biochem. J.*, 248 (1987) 145-149.

- 306 Tomita, Y. and Motokawa, Y.: Characterization and partial purification of a novel mannosylphosphodolichol phosphodiesterase from chicken liver microsomes. *Eur. J. Biochem.*, 170 (1987) 363-368.
- 307 White, J.W., Jr.: Wiley led the way: a century of federal honey research. *J. Assoc. Off. Anal. Chem.*, 70 (1987) 181-189 - a review with 52 refs.
- 308 Zhang, C. and Zhang, Y.: (Thin layer chromatography of unsaturated sulfated disaccharides). *Shengwu Huaxue Yu Shengwu Wuli Jinzhan*, No. 2 (1987) 64-66; *C.A.*, 107 (1987) 150513m.

See also 450, 473.

11. ORGANIC ACIDS AND LIPIDS

11a. Organic acids and simple esters

- 309 Afaque, S., Siddiqui, M.M., Ahmad, I., Siddiqui, M.S. and Osman, S.M.: *Kigelia pinnata* seed oil - a moderate source of vernolic acid. *Fett Wissen. Technol.-Fat Sci. Technol.*, 89 (1987) 433-435.
- 310 Bradbury, A.F. and Smyth, D.G.: Enzyme-catalysed peptide amidation. Isolation of a stable intermediate formed by reaction of the amidating enzyme with an imino acid. *Eur. J. Biochem.*, 169 (1987) 579-584.
- 311 Daulatabad, C.D., Hosamani, K.M., Desai, V.A. and Alagawadi, K.R.: Cyclopropenoid fatty acids in leguminosae oil. *J. Am. Oil Chem. Soc.*, 64 (1987) 1423.
- 312 Demidkina, T.V., Myagkikh, I.V. and Azhayev, A.V.: Transamination catalysed by tyrosine phenol-lyase from *Citrobacter intermedius*. *Eur. J. Biochem.*, 170 (1987) 311-316.
- 313 Grandgirard, A., Julliard, F., Prevost, J. and Sebedio, J.L.: Preparation of geometrical isomers of linolenic acid. *J. Am. Oil Chem. Soc.*, 64 (1987) 1434-1440.
- 314 Gu, Z. and He, L.: (Quantitative determination of organic acids in Guangdong antivenom drug by TLC-densitometry). *Zhongcaovao*, 18 (1987) 399-401; *C.A.*, 107 (1987) 223370z.
- 315 Hatzelmann, A. and Ullrich, V.: Regulation of 5-lipoxygenase activity by the glutathione status in human polymorphonuclear leukocytes. *Eur. J. Biochem.*, 169 (1987) 175-184.
- 316 Hecker, M., Ullrich, V., Fischer, C. and Meese, C.O.: Identification of novel arachidonic acid metabolites formed by prostaglandin H synthase. *Eur. J. Biochem.*, 169 (1987) 113-123.
- 317 Henry, D.Y., Gueritte-Voegelein, F., Insel, P.A., Ferry, N., Bouquet, J., Potier, P., Sevenet, T. and Hanoune, J.: Isolation and characterization of 9-hydroxy-10-*trans*, 12-*cis*-octadecadienoic acid, a novel regulator of platelet adenylate cyclase from *Glechoma hederacea* L. *Labiatae*. *Eur. J. Biochem.*, 170 (1987) 389-394.
- 318 Ichimura, T. and Maeda, T.: (Determination of vanillylmandelic acid in urine by thin-layer chromatography). *Shimadzu Hyoron*, 44 (1987) 243-246; *C.A.*, 107 (1987) 230684h.
- 319 James, M.O. and Pritchard, J.B.: *In vivo* and *in vitro* renal metabolism and excretion of benzoic acid by a marine teleost, the southern flounder. *Drug Metab. Disp.*, 15 (1987) 665-670.
- 320 Kim, R.S. and LaBella, F.S.: Comparison of analytical methods for monitoring autoxidation profiles in authentic lipids. *J. Lipid Res.*, 28 (1987) 1110-1117.
- 321 Rao, Y.R., Pantulu, A.J. and Lakshminarayana, G.: Reduction of fatty acid methyl esters with sodium borohydride-*t*-butanol-methanol. *Fett Wissen. Technol.-Fat Sci. Technol.*, 89 (1987) 398-400.
- 322 Reddy, S.Y. and Prabhakar, J.V.: Isolation of 9,10-dihydroxystearic acid from sal (*Shorea robusta*) fat. *J. Am. Oil Chem. Soc.*, 64 (1987) 97-99.
- 323 Slegel, P., Vereczkey-Donath, G., Ladanyi, L. and Toth-Lauritz, M.: Enantiomeric separation of chiral carboxylic acids, as their diastereomeric carboxamides, by thin-layer chromatography. *J. Pharm. Biomed. Anal.*, 5 (1987) 665-673.
- 324 Tuchman, M., Auray-Blais, C., Ramnaraine, M.L.R., Neglia, J., Krivit, W. and Lemieux, B.: Determination of urinary homovanillic and vanillylmandelic acids from dried filter paper samples: assessment of potential methods for neuroblastoma screening. *Clin. Biochem. (Ottawa)*, 20 (1987) 173-177; *C.A.*, 107 (1987) 171749h.

325 Ushiyama, H., Nishijima, M., Yasuda, K., Kamimura, H., Tabata, S. and Nishima, T.: (Determination of isopropyl citrate in edible oil). *Shokuhin Eiseigaku Zasshi*, 28 (1987) 200-204; *C.A.*, 107 (1987) 174501g.

See also 264, 330, 335, 345, 356, 358, 380, 464, 473, 501.

11b. Prostaglandins

326 Bonicamp, J.M. and Wheeler, C.: Purification of tritium-labeled prostaglandins by high performance thin-layer chromatography. *J. Tenn. Acad. Sci.*, 62, No. 3 (1987) 55-59; *C.A.*, 107 (1987) 191098y.

327 Nagase, H. and Karaki, H.: Palytoxin-induced contraction and release of prostaglandins and norepinephrine in the aorta. *J. Pharmacol. Exp. Ther.*, 242 (1987) 1120-1125.

See also 317.

11c. Lipids and their constituents

328 Ando, S., Waki, H., Kon, K. and Kishimoto, Y.: Up-to-date chromatography of gangliosides. *NATO ASI Ser., Ser. H*, 7 (1987) 167-177; *C.A.*, 107 (1987) 171632q - a review with 33 refs.

329 Ariga, T., Kobayashi, K., Kuroda, Y., Yu, R.K., Suzuki, M., Kitagawa, H., Inagaki, F. and Miyatake, T.: Characterization of tumor-associated fucogangliosides from PC 12 pheochromocytoma cells. *J. Biol. Chem.*, 262 (1987) 14146-14153.

330 Bae, M.J.: (Lipids and fatty acid composition of herba capsellae (*Capsella bursa-pastoris* Linne)). *Han'guk Yongyang Siklyong Hakhoechi*, 16, No. 2 (1987) 83-86; *C.A.*, 107 (1987) 161481a.

331 Beaudoin, A.R., Piotte, M., Jolicoeur, C., Grenier, G., Grondin, G., Rother, N., Gaudreau, J. and St.-Jean, P.: Lipid analysis of a novel type of cell secretion in the exocrine pancreas: the pancreasomes. *Biochim. Biophys. Acta*, 922 (1987) 62-66.

332 Bhakoo, M., Lewis, R.N.A.H. and McElhaney, R.N.: Isolation and characterization of a novel monoacetylated glucopyranosyl neutral lipid from the plasma membrane of *Acholeplasma laidlawii* B. *Biochim. Biophys. Acta*, 922 (1987) 34-45.

333 Blahova, M., Ranny, M., Sedlacek, J. and Ruzicka, V.: Determination of native phospholipids by TLC-FID. *Acta Univ. Carol. Med.*, 32, No. 1-2 (1986) 33-37; *C.A.*, 107 (1987) 232413t.

334 Brasitus, T.A., Dudeja, P.K., Dahiya, R. and Halline, A.: Dexamethasone-induced alterations in lipid composition and fluidity of rat proximal-small-intestinal brush-border membranes. *Biochem. J.*, 248 (1987) 455-461.

335 Chalova, L.I., Avdiyushko, S.A., Ozeretskoykaya, O.L., Khotimchenko, S.V. and Romashina, N.A.: (The composition of lipids of *Phytophthora infestans* and their ability to induce potato phytoalexin accumulation). *Biokhimiya (Moscow)*, 52 (1987) 1445-1453.

336 Clausen, H., Levery, S.B., Nudelman, E.D., Stroud, M., Salyan, M.E.K. and Hakomori, S.-i.: Isolation and characterization of novel glycolipids with blood group A-related structures: galastosyl-A and sialosylgalactosyl-A. *J. Biol. Chem.*, 262 (1987) 14228-14234.

337 Farag, R.S., Ghali, Y., Hewedi, M.M. and Fatah Abdel, O.M.: Biochemical studies on whey lipid substances. *Fett. Wissen. Technol.-Fat Sci. Technol.*, 89 (1987) 450-452.

338 Fowler, S.D., Brown, W.J., Warfel, J. and Greenspan, P.: Use of Nile red for the rapid *in situ* quantitation of lipids on thin-layer chromatograms. *J. Lipid Res.*, 28 (1987) 1225-1232.

339 Gülz, P.-G., Scora, R.W., Müller, E. and Marner, F.-J.: Epicuticular leaf waxes of *Citrus halimii* stone. *J. Agric. Food Chem.*, 35 (1987) 716-720.

340 Hachey, D.L., Thomas, M.R., Emken, E.A., Garza, C., Brown-Booth, L., Adlof, R.O. and Klein, P.D.: Human lactation: maternal transfer of dietary triglycerides labeled with stable isotopes. *J. Lipid Res.*, 28 (1987) 1185-1192.

341 Hara, A. and Taketomi, T.: Occurrence of sulfatide as a major glycosphingolipid in WHHL rabbit serum lipoproteins. *J. Biochem. (Tokyo)*, 102 (1987) 83-92.

- 342 Havekes, L.M., De Wit, E.C.M. and Princen, H.M.G.: Cellular free cholesterol in Hep G2 cells is only partially available for down-regulation of low-density-lipoprotein receptor activity. *Biochem. J.*, 247 (1987) 739-746.
- 343 Jovall, P.-A., Lindström, K., Pascher, I., Pimlott, W. and Samuelsson, B.E.: Identification of a blood group A active hexaglycosylceramide with a type 1 carbohydrate chain in plasma of an A₁ Le(a-b-) secretor. *Arch. Biochem. Biophys.*, 257 (1987) 409-415.
- 344 Kennerly, D.A.: Molecular species analysis of lysophospholipids using high-performance liquid chromatography and argentation thin-layer chromatography. *J. Chromatogr.*, 409 (1987) 291-297.
- 345 Mathur, S.N. and Field, F.J.: Effect of cholesterol enrichment on 12-hydroxy-eicosatetraenoic acid metabolism by mouse peritoneal macrophages. *J. Lipid Res.*, 28 (1987) 1166-1176.
- 346 Michalec, C. and Ledvinova, J.: TLC of phospholipids. Application in biological material. *Acta Univ. Carol. Med.*, 32, No. 1-2 (1986) 47-52; *C.A.*, 107 (1987) 232415v.
- 347 Ng, H.T., Hsu, L.M., Lung, W.Y. and Chen, C.Y.: A comparison of HPLC and TLC in the assessment of amniotic fluid lecithin/sphingomyelin ration. *Proc. Natl. Sci. Council, Republ. China, Part B: Life Sci.*, 11, No. 2 (1987) 91-96; *C.A.*, 107 (1987) 171743b.
- 348 O'Flaherty, J.T., Redman, J.F., Jr., Schmitt, J.D., Ellis, J.M., Surles, J.R., Marx, M.H., Piantadosi, C. and Wykle, R.L.: 1-O-Alkyl-2-N-methylcarbamyl-glycerophosphocholine: a biologically potent, non-metabolizable analog of platelet-activating factor. *Biochem. Biophys. Res. Commun.*, 147 (1987) 18-24.
- 349 Prasad, R.B.N., Rao, Y.N. and Rao, S.V.: Phospholipids of palash (*Butea monosperma*), papaya (*Carica papaya*), jangli badam (*Sterculia foetida*), coriander (*Coriandrum sativum*) and carrot (*Daucus carota*) seeds. *J. Am. Oil Chem. Soc.*, 64 (1987) 1424-1427.
- 350 Reddy, S.Y. and Prabhakar, J.V.: Effect of diglycerides on the solidification properties of sal (*Shorea robusta*) fat. *Fett. Wissen. Technol.-Fat Sci. Technol.*, 89 (1987) 394-397.
- 351 Ritter, G., Krause, W., Geyer, R., Stirn, S. and Wiegandt, H.: Glycosphingo-lipid composition of human semen. *Arch. Biochem. Biophys.*, 257 (1987) 370-378.
- 352 Salvayre, R., Negre, A., Maret, A., Radom, J. and Douste-Blazy, L.: Extracellular origin of the lipid lysosomal storage in cultured fibroblasts from Wolman's disease. *Eur. J. Biochem.*, 170 (1987) 453-458.
- 353 Samanta, T.B. and Ghosh, D.K.: Characterization of progesterone 11 α -hydroxylase of *Aspergillus ochraceus* TS: a cytochrome P-450 linked monooxygenase. *J. Steroid Biochem.*, 28 (1987) 327-332.
- 354 Schäfer, M., Behle, G., Varsanyi, M. and Heilmeyer, L.M.G., Jr.: Ca²⁺ regulation of 1-(3-sn-phosphatidyl)-1D-myo-inositol 4-phosphate formation and hydrolysis on sarcoplasmic-reticular Ca²⁺-transport ATPase. A new principle of phospholipid turnover regulation. *Biochem. J.*, 247 (1987) 579-587.
- 355 Schuch, R. and Ahmad, F.: Structure and biological significance of triacyl-glycerols containing cyclopropene acyl moieties. *Fett Wissen. Technol.-Fat Sci. Technol.*, 89 (1987) 338-339.
- 356 Sharp, P., Poulos, A., Fellenberg, A. and Johnson, D.: Structure and lipid distribution of polyenoic very-long-chain fatty acids in the brain of peroxisome-deficient patients (Zellweger syndrome). *Biochem. J.*, 248 (1987) 61-67.
- 357 Su, S.N. and Jeng, I.: Selective effects of counter ions on the chromatographic behavior of gangliosides. *J. Clin. Biochem. Soc.*, 15, No. 1 (1986) 50-52; *C.A.*, 107 (1987) 214310c.
- 358 Tanaka, T., Yamaguchi, H., Kishimoto, Y. and Gould, R.M.: Lipid metabolism in various regions of squid giant nerve fiber. *Biochim. Biophys. Acta*, 922 (1987) 85-94.
- 359 Tettenborn, C.S. and Mueller, G.C.: Phorbol esters activate the pathway for phosphatidylethanol synthesis in differentiating HL-60 cells. *Biochim. Biophys. Acta*, 931 (1987) 242-250.
- 360 Toth, M. and Hertelendy, F.: Differential effect of progesterone on the labeling of phosphatidylinositol with [³H]inositol and [³²P]phosphate in the uterus of the estrogen-treated ovariectomized rat. *J. Steroid Biochem.*, 28 (1987) 629-635.
- 361 Watarai, S., Handa, S., Tadakuma, T. and Yasuda, T.: Application of liposomes to generation of monoclonal antibody to glycosphingolipid: production of monoclonal antibody to GgOse₄Cer. *J. Biochem. (Tokyo)*, 102 (1987) 59-67.

- 362 Yamaguchi, H., Tanaka, T., Ichioka, T., Stoskopf, M., Kishimoto, Y. and Gould, R.: Characterization and comparison of lipids in different squid nervous tissues. *Biochim. Biophys. Acta*, 922 (1987) 78-84.

See also 279, 315, 321, 379, 380, 382, 451, 501.

13. STEROIDS

See 269.

13a. Pregnane and androstane derivatives

- 363 Ayub, M. and Levell, M.J.: Inhibition of testicular 17 α -hydroxylase and 17,20-lyase but not 3 β -hydroxysteroid dehydrogenase-isomerase or 17 β -hydroxysteroid oxidoreductase by ketoconazole and other imidazole drugs. *J. Steroid Biochem.*, 28 (1987) 521-531.
- 364 Bourget, C., Femino, A., Franz, C., Hastings, S. and Longcope, C.: The effects of l-thyroxine and dexamethasone on steroid dynamics in male cynomolgous monkeys. *J. Steroid Biochem.*, 28 (1987) 575-579.
- 365 Eriksson, C.-G. and Eneroth, P.: Studies on rat liver microsomal steroid metabolism using ¹⁸O-labelled testosterone and progesterone. *J. Steroid Biochem.*, 28 (1987) 549-557.
- 366 Hapgood, J.P. and von Holt, C.: Steroid-affinity purification on the rat liver glucocorticoid hormone receptor complex. *J. Steroid Biochem.*, 28 (1987) 769-777.
- 367 Kobayashi, Y., Nishiguchi, Y., Kiguchi, T., Ninomiya, I. and Watanabe, F.: An alternative metabolic pathway of 11-deoxycorticosterone in bovine adrenal *in vitro*: evidence for the presence of a pathway of 11-deoxycorticosterone oxidation at 19-position. *J. Steroid Biochem.*, 28 (1987) 759-767.
- 368 Numazawa, M., Tsuji, M. and Mutsumi, A.: Studies on aromatase inhibition with 4-androstene-3,6-17-trione: its 3 β -reduction and time-dependent irreversible binding to aromatase with human placental microsomes. *J. Steroid Biochem.*, 28 (1987) 337-344.
- 369 Stupnicki, R., Kowalczyk-Przeewloka, T., Kula, E. and Kasongo, L.Y.: HME - a novel derivative for direct iodination in steroid radioimmunoassays. *J. Steroid Biochem.*, 28 (1987) 663-667.
- 370 Vanderstichele, H., Eechaute, W., Lacroix, E. and Leusen, I.: Influence of neonatal androgenization on the testicular steroidogenesis in the adult rat. *J. Steroid Biochem.*, 28 (1987) 421-427.
- 371 Vazquez, M.H., Tezon, J.G. and Blaquier, J.A.: Studies on the mechanism of the antiandrogenic effect of a putative 5 α -reductase inhibitor. *J. Steroid Biochem.*, 28 (1987) 227-231.
- 372 Zakelj-Mavric, M. and Belic, I.: Hydroxylation of steroids with 11 α -hydroxylase of *Rhizopus nigricans*. *J. Steroid Biochem.*, 28 (1987) 197-201.

See also 274, 381.

13b. Estrogens

- 373 Bindal, R.D., Carlson, K.E., Reiner, G.C.A. and Katzenellenbogen, J.A.: 11 β -Chloromethyl-³H]estradiol-17 β : a very high affinity, reversible ligand for the estrogen receptor. *J. Steroid Biochem.*, 28 (1987) 361-370.
- 374 Simpson, D.M., Elliston, J.F. and Katzenellenbogen, J.A.: Desmethylnafoxidine aziridine: an electrophilic affinity label for the estrogen receptor with high efficiency and selectivity. *J. Steroid Biochem.*, 28 (1987) 233-245.
- 375 Weniger, J.-P. and Zeis, A.: Oestrogen synthesis by the foetal rat testis in organ culture. *J. Steroid Biochem.*, 28 (1987) 307-310.

See also 364.

13c. Sterols

- 376 Armstrong, M.J. and Carey, M.C.: Thermodynamic and molecular determinants of sterol solubilities in bile salt micelles. *J. Lipid Res.*, 28 (1987) 1144-1155.
- 377 Björkhem, I., Miettinen, T., Reihner, E., Ewerth, S., Angelin, B. and Einarsson, K.: Correlation between serum levels of some cholesterol precursors and activity of HMG-CoA reductase in human liver. *J. Lipid Res.*, 28 (1987) 1137-1143.
- 378 Doi, S.Q., Meinertz, H., Nilausen, K., Faria, E.C. and Quintao, E.C.R.: Intestinal cholesterol absorption in the chyluria model. *J. Lipid Res.*, 28 (1987) 1129-1136.
- 379 Field, F.J., Albright, E. and Mathur, S.N.: Regulation of cholesterol esterification by micellar cholesterol in CaCo-2 cells. *J. Lipid Res.*, 28 (1987) 1057-1066.
- 380 Gebhard, R.L., Clayman, R.V., Prigge, W.F., Figenshau, R., Staley, N.A., Reese, C. and Bear, A.: Abnormal cholesterol metabolism in renal clear cell carcinoma. *J. Lipid Res.*, 28 (1987) 1177-1184.
- 381 Monnier, N., Defaye, G. and Chambaz, E.M.: Phosphorylation of bovine adrenodoxin. Structural study and enzymatic activity. *Eur. J. Biochem.*, 169 (1987) 147-153.
- 382 Pratihari, H.K., Mohanty, N. and Das, P.B.: Studies on tallow in vegetable ghee by TLC and GLC. *J. Inst. Chem. (India)*, 59 (1987) 124; *C.A.*, 107 (1987) 196604v.
- 383 Reichen, J., Karlaganis, G. and Kern, F., Jr.: Cholesterol synthesis in the perfused liver of pregnant hamsters. *J. Lipid Res.*, 28 (1987) 1046-1052.
- 384 Slotte, J.P. and Bierman, E.L.: Movement of plasma-membrane sterols to the endoplasmic reticulum in cultured cells. *Biochem. J.*, 248 (1987) 237-242.

See also 335, 339, 342, 345, 358.

15d. Bile acids and alcohols

- 385 Guo, Y., Meng, X. and Zhang, E.: (Determination of cholic acid and emodin by TLC in Niu Huang Jie Du Pian). *Zhongcaoyao*, 18 (1987) 253-254; *C.A.*, 107 (1987) 205266r.
- 386 Jagarinec, N. and Parag, G.: Separation of unconjugated cholic, chenodeoxycholic, deoxycholic and lithocholic acids by thin-layer chromatography. *Period. Biol.*, 89, No. 1 (1987) 21-26; *C.A.*, 107 (1987) 171745d.
- 387 Liu, L., Xu, D. and Xie, D.: (TLC-scanning determination of cholic and deoxycholic acids in fast-acting cold capsules). *Yiyao Bongye*, 18 (1987) 361-362; *C.A.*, 107 (1987) 183699e.
- 388 Ni, K. and Lin, S.: (Preparation and determination of ursodeoxycholic acid and its epimer by TLC). *Zhongguo Yaoke Daxue Xuebao*, 18, No. 2 (1987) 91-94; *C.A.*, 107 (1987) 161778c.
- 389 Vamecq, J., Schepers, L., Parmentier, G. and Mannaerts, G.P.: Inhibition of peroxisomal fatty acyl-CoA oxidase by antimycin A. *Biochem. J.*, 248 (1987) 603-607.

See also 383.

13e. Ecdysones and other insect steroid hormones

- 390 Bathori, M., Szendrei, K., Miklos, P., Pelczer, I. and Solymosi, P.: Ecdysteroids from *Silene nutans* L. *Symp. Biol. Hung.*, 34 (Chromatography '85) (1986) 241-250; *C.A.*, 107 (1987) 150528v.
- 391 Wilson, I.D. and Lewis, S.: Separation of ecdysteroids by high-performance thin-layer chromatography using automated multiple development. *J. Chromatogr.*, 408 (1987) 445-448.

13f. Other steroids

See 374.

14. STEROID GLYCOSIDES AND SAPONINS

- 392 Blum, L.M. and Rieders, F.: Oleandrin distribution in a fatality from rectal and oral *Nerium Oleander* extract administration. *J. Anal. Toxicol.*, 11 (1987) 219-221; *C.A.*, 107 (1987) 192406j.
- 393 Kamat, S.S. and Kawale, G.B.: Anthrone as a reagent for the detection of glycosides and saponins by thin layer chromatography. *Curr. Sci.*, 56 (1987) 818-821; *C.A.*, 107 (1987) 213082t.
- 394 Lalitha, T., Seshadri, R. and Venkataraman, L.V.: Isolation and properties of saponins from *Madhuca butyracea* seeds. *J. Agric. Food Chem.*, 35 (1987) 744-748.
- 395 Sezik, E. and Turkoz, S.: Separation of some triterpenic saponins by column chromatography. *Acta Pharm. Turc.*, 29, No. 3 (1987) 87-93; *C.A.*, 107 (1987) 223107u.
- 396 Yu, B., Xu, G., Jin, R. and Xu, L.: (Determination of ruscogenin in Maidong). *Zhongguo Yaokexue Xuebao*, 18, No. 2 (1987) 117-119; *C.A.*, 107 (1987) 161761s.
- 397 Zhao, G., Huang, Q., Peng, G., Mei, S. and Yu, J.: (A study on factors affecting the contents of saponins of Jiegeng (*Platycodon grandiflorum*)). *Zhongcaoyao*, 18 (1987) 257-258, 254; *C.A.*, 107 (1987) 205010c.

See also 432, 473, 497.

15. TERPENES AND OTHER VOLATILE AROMATIC COMPOUNDS

15a. Terpenes

- 398 Alam, M., Umar, M. and Dar, F.: Quantitative determination of glycyrrhizin in licorice roots. *J. Pharm. (Lahore)*, 7, No. 1-2 (1986) 21-24; *C.A.*, 107 (1987) 223358b.
- 399 Li, S. and Zhang, R.: (Determination of betulinic acid in Suanzaoren (Semen *Ziziphi spinosae*) by TLC). *Zhongcaoyao*, 18 (1987) 255-256; *C.A.*, 107 (1987) 205267s.
- 400 Satterwhite, D.M. and Croteau, R.B.: Resolution of monoterpene enantiomers by gas chromatography. *J. Chromatogr.*, 407 (1987) 243-252.

See also 339, 497.

15b. Essential oils

- 401 Gu, Y., Zhong, M. and Chen, S.: (Comparison of the essential oils from Sichuan commodity Xixin (*Asarum*) by TLC and GC). *Zhongcaoyao*, 18 (1987) 299-301; *C.A.*, 107 (1987) 205265q.

See also 432.

15c. Bitter substances

- 402 Hayashi, T.: (Analysis of bitter components of *Swertiae herba*, *Gentianae radix* and *Gentianae scabrae radix*). *Pharm. Tech. Jpn.*, 3 (1987) 777-784; *C.A.*, 107 (1987) 161753r - a review with 29 refs.

16. NITRO AND NITROSO COMPOUNDS

See 267, 505.

17. AMINES, AMIDES AND RELATED NITROGEN COMPOUNDS

17a. Amines and polyamines

- 403 Nishikata, M.: Separation of β -amino alcohols as their dansyl derivatives by thin-layer chromatography on a polyamide layer sheet. *J. Chromatogr.*, 408 (1987) 449-452.
- 404 Polotebnova, N.A. and Esmond, N.A.A.A.: (Separation and determination of aniline and its derivatives using heteropoly compounds). *Metallkkelaty Ikh Svoistva*, (1985) 48-53; *C.A.*, 107 (1987) 183111g.

See also 264, 502, 505.

17d. Other amine derivatives and amides (excluding peptides)

- 405 Izokun-Etiobhio, B.O., Oraedu, A.C.I. and Igochukwu, E.N.: A simple procedure for the quantification of intact linamarin in cassava and its products. *J. Sci. Food Agric.*, 41 (1987) 81-86; *C.A.*, 107 (1987) 196620x.

See also 272, 310, 323, 502.

18. AMINO ACIDS AND PEPTIDES; CHEMICAL STRUCTURE OF PROTEINS

18a. Amino acids and their derivatives

- 406 Bartelt, D.C., Moroney, S. and Wolff, D.J.: Purification, characterization and substrate specificity of calmodulin-dependent myosin light-chain kinase from bovine brain. *Biochem. J.*, 247 (1987) 747-756.
- 407 Bhusan, R. and Reddy, G.P.: Thin-layer chromatography of phenyl thiohydantoin derivatives of amino acids on silica gel thin layers impregnated with zinc salts and some metal sulphates. *Analyst (London)*, 112 (1987) 1467-1469.
- 408 Cole, S.C.J. and Yon, R.J.: Active-site-directed inactivation of wheat-germ aspartate transcarbamoylase by pyridoxal 5'-phosphate. *Biochem. J.*, 348 (1987) 403-408.
- 409 Gullner, G., Cserhati, T. and Szogyi, M.: Lipophilicity determination of amino acids by reversed-phase thin-layer chromatography. *Symp. Biol. Hung.*, 34 (Chromatography '85) (1986) 73-83; *C.A.*, 107 (1987) 171742a.
- 410 Hirschberg, K., Faust, H. and Balint, A.: (Thin-layer chromatography of amino acid mixtures on ion exchange foils for the determination of ^{15}N -abundances in the separated amino acids). *Isotopenpraxis*, 23 (1987) 153-155; *C.A.*, 107 (1987) 232392k.
- 411 Kusano, G., Ogawa, H., Takahashi, A., Nozoe, S. and Yokoyama, K.: A new amino acid (2S,3R)-(-)-3-hydroxybaikaiin from *Russula subnigricans* Hongo. *Chem. Pharm. Bull.*, 35 (1987) 3482-3486.
- 412 Lepri, L., Desideri, P.G., Merlini, L. and Checchini, L.: HPLC of amino acids on silica coated with ammonium tubgstoposphate. *Chromatographia*, 23 (1987) 814-817.
- 413 Ramis Ramos, G., Mauri Aucejo, A.R., Garcia Alvarez-Coque, M.C. and Mongay Fernandez, C.: Determination of tyrosine and phenylalanine by derivatization with nitric acid and differential pulse polarography. *Microchem. J.*, 36 (1987) 113-117; *C.A.*, 107 (1987) 168048y.
- 414 Rieul, C., Cortay, J.-C., Bleicher, F. and Cozzone, A.J.: Effect of bacteriophage M13 infection on phosphorylation of dnaK protein and other *Escherichia coli* proteins. *Eur. J. Biochem.*, 168 (1987) 621-627.
- 415 Rutan, S.C. and Motley, C.B.: Factor analysis and Kalman filter studies of severely overlapped amino acid derivatives in thin-layer chromatography. *Anal. Chem.*, 59 (1987) 2045-2050.
- 416 Sandor, A., Kispal, G., Melegh, B. and Alkonyi, I.: Ester composition of carnitine in the perfusate of liver and in the plasma of donor rats. *Eur. J. Biochem.*, 170 (1987) 443-445.

- 417 Watanabe, T.K., Kuczmarski, E.R. and Reddy, J.K.: Myosin from pancreatic acinar carcinoma cells. Isolation, characterization and demonstration of heavy- and light-chain phosphorylation. *Biochem. J.*, 247 (1987) 513-518.
- 418 Wu, Y., Xiong, L. and Nan, G.: (Quantitative determination of amino acids by two-dimensional thin-layer chromatography and *in situ* scanning. *Sepu*, 5 (1987) 256-259; *C.A.*, 107 (1987) 150548b.

See also 264, 278, 312, 403, 423.

18b. Peptides and peptidic and proteinous hormones

- 419 Davis, S.S.: Advanced delivery systems for peptides and proteins - pharmaceutical considerations. *NATO ASI Ser., Ser. A*, 125 (Delivery Syst. Pept. Drugs) (1986) 1-21; *C.A.*, 107 (1987) 204974h - a review with 84 refs.
- 420 Levine, H., III and Sahyoun, N.E.: Characterization of a soluble M_r -30000 catalytic fragment of the neuronal calmodulin-dependent protein kinase II. *Eur. J. Biochem.*, 168 (1987) 481-486.
- 421 Pearson, R.K., Powers, S.P., Hadac, E.M., Gaisano, H. and Miller, L.J.: Establishment of a new short, protease-resistant, affinity labeling reagent for the cholecystokinin receptor. *Biochem. Biophys. Res. Commun.*, 147 (1987) 346-353.
- 422 Wolff, J.M., Rathjen, F.G., Frank, R. and Roth, S.: Biochemical characterization of polypeptide components involved in neurite fasciculation and elongation. *Eur. J. Biochem.*, 168 (1987) 551-561.

See also 278, 403.

19. PROTEINS

See 419.

21. PURINES, PYRIMIDINES, NUCLEIC ACIDS AND THEIR CONSTITUENTS

21a. Purines, pyrimidines, nucleosides, nucleotides

- 423 Freist, W., Sternbach, H. and Cramer, F.: Isoleucyl-tRNA synthetase from baker's yeast and from *Escherichia coli* MRE 600. Discrimination of 20 amino acids in aminoacylation of tRNA^{Ile}-C-C-A(3'NH₂). *Eur. J. Biochem.*, 169 (1987) 33-39.
- 424 Kwok, F., Scholz, G. and Churchich, J.E.: Brain pyridoxal kinase dissociation of the dimeric structure and catalytic activity of the monomeric species. *Eur. J. Biochem.*, 168 (1987) 577-583.
- 425 L thje, J. and Ogilvie, A.: Catabolism of A_{p4}A and A_{p3}A in human serum. Identification of isoenzymes and their partial characterization. *Eur. J. Biochem.*, 169 (1987) 285-288.
- 426 Rabin, B.A. and Chase, J.W.: DNA ligase from *Drosophila melanogaster* embryos. Substrate specificity and mechanism of action. *J. Biol. Chem.*, 262 (1987) 14105-14111.
- 427 Shcherbakov, D.Yu., Kit, Yu.Ya., Koldashova, Z.T. and Sidorov, V.N.: (Phosphorylation of proteins on the cell surface of *Drosophila melanogaster* larvae salivary glands). *Biokhimiya (Moscow)*, 52 (1987) 1608-1613.

See also 296.

22. ALKALOIDS

- 428 Bergisadi, N. and Ozsoy, Y.: Determination of codeine and dionin in codeine-dionin combined tablets. *Acta Pharm. Turc.*, 29, No. 3 (1987) 81-83; *C.A.*, 107 (1987) 223390f.

- 429 Chen, R., Li, T.M., Merrick, H., Parrish, R.F., Bruno, V., Kwong, A., Stiso, C. and Litman, D.J.: An internal clock reaction used in a one-step enzyme immuno-chromatographic assay of theophylline in whole blood. *Clin. Chem. (Winston-Salem)*, 33 (1987) 1521-1525.
- 430 Gaitonde, R.V. and Rivankar, U.: Analysis of a drug preparation containing ephedrine hydrochloride, theophylline, chlorpheniramine maleate and diazepam by thin layer chromatography. *Indian Drugs*, 24 (1987) 486-488; *C.A.*, 107 (1987) 161783a.
- 431 Horvath, P.: Analysis of alkaloids. *Chromatogr. Sci.*, 36 (Quant. Thin-Layer Chromatogr. Its Ind. Appl) (1987) 217-243; *C.A.*, 107 (1987) 154554y - a review with 84 refs.
- 432 Hou, S., Liao, G., Yang, H. and Li, S.: (Quality control method of compound "Sansheng" injection). *Zhongcaoyao*, 18 (1987) 349-352; *C.A.*, 107 (1987) 183664q.
- 433 Jain, S.C. and Sharma, R.: Antimicrobial activity of pyrrolizidine alkaloids from *Heliotropium ellipticum*. *Chem. Pharm. Bull.*, 35 (1987) 3487-3489.
- 434 Li, A., Wang, X. and Feng, H.: (Quantitative determination of sinomenine in qingfengteng (*Sinomenium acutum* and *S. acutum* var. *cinereum*) by TLC densitometry and relation between the content and the seasons). *Zhongyao Tongbao*, 12 (1987) 332-334; *C.A.*, 107 (1987) 151241h.
- 435 Mistry, M. and Houston, J.B.: Glucuronidation *in vitro* and *in vivo*. Comparison of intestinal and hepatic conjugation of morphine, naloxone, and buprenorphine. *Drug Metab. Disp.*, 15 (1987) 710-717.
- 436 Mousinho, M.C. and De Oliveira, F.: (Detection of xanthine alkaloids of guarana (*Paullinia cupana* KBK) in urine samples by thin-layer chromatography). *Rev. Bras. Farmacogn.*, 1, No. 1 (1986) 35-44; *C.A.*, 107 (1987) 196581k.
- 437 Muhamad bin Zakaria and Jeffreys, J.A.D.: The isolation of β -carboline from *Lolium perenne* L. *Malays. J. Sci.*, 8 (1986) 115-119; *C.A.*, 107 (1987) 205001a.
- 438 Tencheva, J., Yamboliev, I. and Zhivkova, Z.: Reversed-phase liquid chromatography for the determination of galanthamine and its metabolites in human plasma and urine. *J. Chromatogr.*, 421 (1987) 396-400.
- 439 Wang, L. and Zhang, B.: (TLC-scanning determination of lappaconitine in tall monkshood (*Aconitum sinomontanum*)). *Zhongcaoyao*, 18 (1987) 305-306; *C.A.*, 107 (1987) 183653k.
- 440 Zhang, X. and Tian, X.: (Quantitative determination of berberine in Goldthread (*Coptis chinensis*) by densitometry). *Zhongcaoyao*, 18 (1987) 394-396; *C.A.*, 107 (1987) 223368e.

See also 485, 493.

23. OTHER SUBSTANCES CONTAINING HETEROCYCLIC NITROGEN

23a. Porphyrins and other pyrroles

- 441 Bobarevic, B., Lovren, F. and Gaon, I.D.: Thin layer chromatography of some pyrrolinone derivatives. *Folia Med. Fac. Med. Univ. Sarajevensis*, 20, No. 1 (1985) 25-30; *C.A.*, 107 (1987) 161775z.
- 442 Pfaltz, A., Kobelt, A., Hüster, R. and Thauer, R.K.: Biosynthesis of coenzyme F430 in methanogenic bacteria. Identification of 15,17³-seco-F430-17³-acid as an intermediate. *Eur. J. Biochem.*, 170 (1987) 459-467.

23b. Bile pigments

- 443 Meisel, P., Jährig, D. and Meisel, M.: Detection of photobilirubin in urine of jaundiced infants supporting the diagnosis of 'bronze baby syndrome'. *Clin. Chim. Acta*, 166 (1987) 61-65.
- 444 Yoshida, H., Inagaki, T., Hirano, M. and Sugimoto, T.: Analyses of azopigments obtained from the delta fraction of bilirubin from mammalian plasma (mammalian biliprotein). *Biochem. J.*, 248 (1987) 79-84.

23c. Indole derivatives

See 485.

23d. Pyridine derivatives

See 502.

23e. Other N-heterocyclic compounds

- 445 Javor, J., Kiss, I. and Kovacs, A.: Thin-layer chromatography and high-pressure liquid chromatography of different acridine-glycosides. *Symp. Biol. Hung.*, 34 (Chromatography '85) (1986) 263-288; *C.A.*, 107 (1987) 161765w.
- 446 Saito, M.L. and De Oliveira, F.: (Physical and chemical characteristics of fluid extract of *Cordia ealyculata* Vell-Boraginaceae). *Rev. Bras. Farmacogn.*, 1, No. 1 (1986) 3-11; *C.A.*, 107 (1987) 183394b.

See also 502.

24. ORGANIC SULPHUR COMPOUNDS

- 447 Bottura, G. and Pavesi, M.A.: Thin-layer chromatography of some organic selenium compounds. *Microchem. J.*, 35 (1987) 223-226; *C.A.*, 107 (1987) 236451b.
- 448 Succar, S.D. and Catoggio, J.A.: (Separation of thiocarbamates by thin layer chromatography. Detection limits with different reagents). *Rev. Soc. Quim. Mex.*, 30, No. 2 (1986) 70-72; *C.A.*, 107 (1987) 168003e.
- 449 Toader, O.V., Trifan, V., Mazare, M. and Budrugaec, S.A.: (Determination of trace α -naphthylthiourea in the presence of colored extracts). *Rom. Pat. RO 89,225* (Cl. G01N31/08), 30 Apr. 1986, Appl. 114,438, 03 May 1984; *C.A.*, 107 (1987) 190133u.

25. ORGANIC PHOSPHORUS COMPOUNDS (INCLUDING SUGAR PHOSPHATES)

- 450 Rietschel, E.T., Brade, L., Schade, U., Galanos, C., Freudenberg, M., Lüderitz, O., Kusumoto, S. and Shiba, T.: Endotoxic properties of synthetic pentaacyl lipid A precursor Ib and a structural isomer. *Eur. J. Biochem.*, 169 (1987) 27-31.
- 451 Sedlacek, J., Ranny, M., Blahova, M. and Ruzicka, V.: TLC-FID of phosphatidic and biophosphatidic acids. *Acta Univ. Carol. Med.*, 32, No. 1-2 (1986) 39-45; *C.A.*, 107 (1987) 232414u.

See also 306.

26. ORGANOMETALLIC AND RELATED COMPOUNDS

26a. Organometallic compounds

- 452 Li, S., Yang, X., Li, Y., Du, Y. and Chen, S.: Analysis of organometallic compounds. IX. Paper chromatography of bis(alkenylcyclopentadienyl)titanium, -zirconium, and -hafnium dichlorides, dithiocyanato and dimercapto derivatives. *Septu*, 5 (1987) 77-80; *C.A.*, 107 (1987) 228099c.

26c. Coordination compounds

- 453 Galli, B., Gasparrini, F., Misiti, D., Natile, G. and Palmieri, G.: Thin-layer chromatography of metal complexes on glycidoxypropyl functionalized silica plates. *J. Chromatogr.*, 409 (1987) 377-382.

27. VITAMINS AND VARIOUS GROWTH REGULATORS (NON-PEPTIDIC)

- 454 Surai, P.F. and Ionov, I.A.: (Practical use of silufole plates for the analysis of fat-soluble vitamins). *Vopr. Pitan.*, No. 2 (1987) 66-70; *C.A.*, 107 (1987) 150550w.
- 455 Wall, J.S., Young, M.R. and Carpentier, K.J.: Transformation of niacin-containing compounds in corn during grain development: relationship to niacin nutritional availability. *J. Agric. Food Chem.*, 35 (1987) 752-758.

See also 267.

28. ANTIBIOTICS

- 456 Dahiya, J.S.: Reversed-phase high-performance liquid chromatography of *Cajanus* phytoalexins. *J. Chromatogr.*, 409 (1987) 355-359.
- 457 Erguven, H. and Mascaro de Piccoli, A.: (Neomycin sulfate: effects of phosphate, borate and hydroxide ions on its stability). *Acta Farm. Bonaerense*, 6, No. 1 (1987) 9-14; *C.A.*, 107 (1987) 223189x.
- 458 Hou, E.W. and Li, S.S.-L.: Cyclic AMP-induced expression of the mouse lactate dehydrogenase-A promoter-cat fusion gene in Chinese hamster ovary wild-type cells, but not in cAMP-dependent protein kinase mutant cells. *Biochem. Biophys. Res. Commun.*, 147 (1987) 501-505.
- 459 Jarzebinski, J., Lugowska, E. and Czaja, K.: (Application of densitometry in determination of drugs in pharmaceuticals. XI. Determination of tetracycline antibiotics in some pharmaceutical preparations). *Acta Pol. Pharm.*, 43 (1986) 260-263; *C.A.*, 107 (1987) 205274s.
- 460 Neidert, E., Saschenbrecker, P.W. and Tittiger, F.: Thin layer chromatographic/bioautographic method for identification of antibiotic residues in animal tissues. *J. Assoc. Off. Anal. Chem.*, 70 (1987) 197-200.
- 461 Sun, C., Yu, R., Yang, Q., Sheng, S. and Zhao, X.: (Quantitative analyses of spiramycin and acetylspectinomycin by thin-layer chromatography). *Yaowu Xuebao*, 22 (1987) 515-519; *C.A.*, 107 (1987) 205259r.

29. INSECTICIDES, PESTICIDES AND OTHER AGROCHEMICALS

29b. *Phosphorus insecticides*

- 462 United Kingdom Dept. of Environment (UK): Organophosphorus pesticides in sewage sludge 1985; organophosphorus pesticides in river and drinking water, an addition, 1985. *Methods Exam. Waters Assoc. Mater.*, 1986, 20 pp.; *C.A.*, 107 (1987) 149096w.

29c. *Carbamates*

- 463 Chmil, V.D. and Vasyagina, R.D.: (Partition thin-layer chromatography of thiocarbamic derivatives). *Zh. Anal. Khim.*, 42 (1987) 1691-1695; *C.A.*, 107 (1987) 192858h.

See also 448, 502.

29e. *Fungicides*

- 464 Davidkova, P. and Gasparic, J.: TLC and HPTLC of alkyl-substituted phenoxy-alkanoic acids. *Symp. Biol. Hung.*, 34 (Chromatography '85) (1986) 567-578; *C.A.*, 107 (1987) 228104a.

29f. Other types of pesticides and various agrochemicals

- 465 Uskova, I.A., Panasyuk, T.D. and Prokhorchuk, E.A.: (Determination of polidim residues in water, soil, and plants by thin-layer chromatography). *Khim. Sel'sk. Khoz.*, No. 8 (1987) 69-70; *C.A.*, 107 (1987) 149100t.

30. SYNTHETIC AND NATURAL DYES

30a. Synthetic dyes

- 466 Nasierowska, Z., Weglowska, W., Suffczynski, J. and Kwiatkowska, B.: (Analytical studies of dyes used for coloring drugs). *Acta Pol. Pharm.*, 43 (1986) 583-587; *C.A.*, 107 (1987) 161799k.
- 467 Patel, A.K., Hallett, M.B. and Campbell, A.K.: Threshold responses in production of reactive oxygen metabolites in individual neutrophils detected by flow cytometry and microfluorimetry. *Biochem. J.*, 248 (1987) 173-180.

See also 272.

30b. Chloroplast and other natural pigments

- 468 Levadoux, W.L., Kalmokoff, M.L., Pickard, M.D. and GrootWassink, J.W.D.: Pigment removal from canola oil using chlorophyllase. *J. Am. Oil Chem. Soc.*, 64 (1987) 139-144.
- 469 Manabe, A., Tokumori, T., Sumida, Y., Yoshida, T., Hatano, T., Yazaki, K. and Okuda, T.: (Application of supercritical fluid extraction to components of crude drugs and plants. III. Extraction of pigments from *Lithospermum* root and licorice root). *Yakugaku Zasshi*, 107 (1987) 506-510; *C.A.*, 107 (1987) 205004d.
- 470 Ni, J., Zhou, H. and Wang, X.: (Determination of pigments in green plant leaves by using a dual-wavelength TLC scanner). *Lanzhou Daxue Xuebao, Ziran Kexueban*, 23 (1987) 109-113; *C.A.*, 107 (1987) 150551x.
- 471 Suzuki, N., Saitoh, K. and Adachi, K.: Reversed-phase high-performance thin-layer chromatography and column liquid chromatography of chlorophylls and their derivatives. *J. Chromatogr.*, 408 (1987) 181-190.

See also 267.

32. DRUG ANALYSIS

- 472 Pachaly, P.: Thin layer chromatographic identity testing in the pharmaceutical laboratory. *OAZ, Oesterr. Apoth.-Ztg.*, 40 (1986) 809-819; *C.A.*, 107 (1987) 183636g - a review.

See also 266, 277, 419.

32a. Drug analysis, general techniques

- 473 Zhang, G., Liu, H. and Hong, W.: (Application of thin-layer stick chromatography in pharmaceutical analysis). *Fenxi Huaxue*, 15 (1987) 367-371. *C.A.*, 107 (1987) 205255m.

See also 278.

32c. Autonomic and cardiovascular drugs

- 474 Bezakova, Z., Blesova, M., Bachrata, M., Cizmarik, J. and Benes, L.: (Analytical profile of stobadin (DH 1011)). *Farm. Obs.*, 56 (1987) 341-351; *C.A.*, 107 (1987) 223388m.

- 475 Blesova, M., Bezakova, Z., Bachrata, M., Cizmarik, J. and Benes, L.: (Analytical evaluation of stobadine dipalmitate (DP 1031)). *Cesk. Farm.*, 36 (1987) 245-249; *C.A.*, 107 (1987) 205278w.
- 476 Jarzebinski, J. and Ciszewska-Hedrasik, M.: (Application of densitometry in determination of active substances in drugs. XII. Determination of substances in compound preparations containing naphazoline nitrate). *Acta Pol. Pharm.*, 43 (1986) 264-269; *C.A.*, 107 (1987) 223391g.
- 477 Wang, Y., Shu, H. and Zeng, Y.: (Determination of changrolin by quantitative TLC scanning technique). *Yaowu Fenxi Zazhi*, 7 (1987) 221-223; *C.A.*, 107 (1987) 211361j.

See also 429, 430, 438, 493.

32d. Central nervous system drugs

- 478 Bezakova, Z., Bachrata, M., Cizmarik, J. and Blesova, M.: Pentacaine hydrochloride, carbisocaine hydrochloride and heptacaine hydrochloride - long-time and short-time tests in study of stability. *Pharmazie*, 42 (1987) 633.
- 479 Chen, D. and Wang, W.: (Indirect determination of chlorpheniramine maleate, diphenhydramine hydrochloride, and carbetapentane citrate by spectrophotometry). *Shenyang Yaoxueyuan Xuebao*, 4, No. 2 (1987) 122-126; *C.A.*, 107 (1987) 205305c.
- 480 Duan, G., Lu, M., Xiong, Y., Dai, X. and Fei, Z.: (TLC densitometric determination of tiapride in serum and urine). *Shanghai Yike Daxue Xuebao*, 14 (1987) 189-193; *C.A.*, 107 (1987) 228322v.
- 481 Gruhl, H.: (Detection of benzo- and thienodiazepines by EMIT ST and TLC as part of a drug screening program). *Z. Rechtsmed.*, 98 (1987) 221-228; *C.A.*, 107 (1987) 213075t.
- 482 Horne, C., Spahn, H. and Mutschler, E.: Fluorimetric determination of levo-protiline in human plasma after thin-layer chromatographic or high performance liquid chromatographic separation. *Arzneim.-Forsch.*, 37 (1987) 1179-1181.
- 483 Mroszczak, E.J., Lee, F.W., Combs, D., Sarnquist, F.H., Huang, B.-L., Wu, A.T., Tokes, L.G., Maddox, M.L. and Cho, D.K.: Ketorolac tromethamine absorption, distribution, metabolism, excretion, and pharmacokinetics in animals and humans. *Drug Metab. Disp.*, 15 (1987) 618-626.
- 484 Pawelczyk, E., Knitter, B. and Majewski, W.: (Kinetics of drug decomposition. LXXXI. Stability of acetylsalicylic acid and paracetamol in polyethylene glycol 1500 and Witepsol H15). *Acta Pol. Pharm.*, 43 (1986) 242-249; *C.A.*, 107 (1987) 205079g.
- 485 Pietrogrande, M.C., Borea, P.A., Lodi, G. and Bigli, C.: Reversed-phase ion-pair systems to predict partition coefficients of β -carbolines by HPLC and TLC. *Chromatographia*, 23 (1987) 713-716.
- 486 Reszka, I. and Tyfczynska, J.: (Quick isolation and detection of diazepam and its main metabolite in blood serum). *Farm. Pol.*, 42 (1986) 472-474; *C.A.*, 107 (1987) 168164h.
- 487 St-Pierre, M.V. and Pang, K.S.: Determination of diazepam and its metabolites by high-performance liquid chromatography and thin-layer chromatography. *J. Chromatogr.*, 421 (1987) 291-307.

See also 430, 435, 476, 493.

32e. Chemotherapeutics (except cytostatics and antibiotics)

- 488 De la Vigne, U.: (HPTLC determination of sulfonamides in veterinary preparations). *LaborPraxis*, 11 (1987) 944 and 946-947; *C.A.*, 107 (1987) 23386j.
- 489 Galy, A.-M., Galy, J.-P., Barbe, J. and Sharples, D.: Preparation of a series of 9-alkylaminoacridines and 9-imino-10-alkylacridines and their binding to deoxyribonucleic acid. *Arzneim.-Forsch.*, 37 (1987) 1095-1098.
- 490 Klimes, J. and Zahradnicek, M.: (Identification of hydrolysis products of selected sulfonamides by TLC). *Cesk. Farm.*, 36 (1987) 250-253; *C.A.*, 107 (1987) 183677w.

See also 477.

32f. *Cytostatics*

See 489.

32g. *Other drug categories*

See 276, 374, 479, 512.

32h. *Toxicological and forensic applications*

- 491 Blakeslay, J., Wood, D., Howse, C. and Spencer-Peet, J.: A simplified thin-layer chromatography system for the detection of commonly abused basic drugs. *Ann. Clin. Biochem.*, 24 (1987) 508-510; *C.A.*, 107 (1987) 192384a.
- 492 Borchert, A., Schneider, W.R. and Schuetz, H.: (Corrected R_{F_c} -value in forensic toxicology study and experiences). *Beitr. Gerichtl. Med.*, 45 (1987) 193-210; *C.A.*, 107 (1987) 230710p.
- 493 Moll, H. and Clerc, J.T.: (Simple, qualitative determination of drugs and/or their metabolites in urine-compliance test and illicit drug screening). *Pharm. Acta Helv.*, 62 (1987) 210-215; *C.A.*, 107 (1987) 168162f.
- 494 Neuninger, H.: (Detection and identification of phenethylamines (stimulants and hallucinogens)). *Sci. Pharm.*, 55 (1987) 1-11; *C.A.*, 107 (1987) 192399j.

See also 287, 392, 393, 481.

32i. *Plant extracts*

- 495 Carr, J.D. and Rogers, C.B.: Chemosystematic studies of the genus *Combretum* (*Combretaceae*) I. A convenient method of identifying species of this genus by a comparison of the polar constituents extracted from leaf material. *S. Afr. J. Bot.*, 53 (1987) 173-176; *C.A.*, 107 (1987) 171736b.
- 496 Hanus, L.: Our experience with identification and separation of compounds from marijuana (*Cannabis sativa* L.) by thin layer chromatography. *Acta Univ. Palacki. Olomouc. Pac. Med.*, 116 (1987) 15-24; *C.A.*, 107 (1987) 230589f - a review with 24 refs.
- 497 Kitagawa, I., Taniyama, T., Shibuya, H., Noda, T. and Yoshikawa, M.: (Chemical studies on crude drug processing. V. On the constituents of ginseng radix rubra (2): comparison of the constituents of white ginseng and red ginseng prepared from the same *Panax ginseng* root). *Yakugaku Zasshi*, 107 (1987) 495-505; *C.A.*, 107 (1987) 205003c.
- 498 Okamoto, M., Jinno, K. and Omori, T.: Preparation and evaluation of mono-, di- and triphenyl-treated plates for the high-performance thin-layer chromatographic analysis of puerarin and paeonol. *J. Chromatogr.*, 407 (1987) 314-318.
- 499 Peishan, X. and Yuzhen, Y.: HPTLC fingerprint identification of commercial ginseng drugs - reinvestigation of HPTLC of ginsenosides. *J. High Resolut. Chromatogr. Chromatogr. Commun.*, 10 (1987) 607-613.
- 500 Petri, G., Kery, A. and Lemberkovics, E.: (Instrumental facilities of standardization of complex plant preparations, paramedicaments). *Acta Pharm. Hung.*, 57, No. 3-4 (1987) 111-132; *C.A.*, 107 (1987) 161767y.
- 501 Stoyanova, A., Popova, M. and Georgiev, E.: (Thin-layer chromatography of *Hypericum perforatum*). *Farmatsiya (Sofia)*, No. 1 (1987) 8-13; *C.A.*, 107 (1987) 205272q.

See also 291, 314, 330, 399, 402, 446.

33. CLINICO-CHEMICAL APPLICATIONS

33b. *Complex mixtures and profiling (single compounds by cross ref. only)*

See 318, 324, 329, 352, 356, 378, 380, 410, 418, 429, 438, 443.

34. FOOD ANALYSIS

34b. *Complex mixtures (single compounds by cross ref. only)*

See 286, 289, 307, 503.

35. ENVIRONMENTAL ANALYSIS

502 Qureshi, M., Ahmad, A., Sulaiman, A.M. and Shakeel, N.A.: Separation of certain organic pollutants containing tertiary nitrogen on stannic molybdosilicate papers. *Anal. Lett.*, 20 (1987) 1157-1169.

35c. *Water pollution (complex mixtures; single compounds by cross ref. only)*

See 404, 462, 465.

35d. *Soil pollution (complex mixtures; single compounds by cross ref. only)*

See 465.

36. SOME TECHNICAL PRODUCTS AND COMPLEX MIXTURES

36b. *Antioxidants and preservatives*

503 Arens, M. and Gertz, C.: Gemeinschaftsarbeiten der DGF, 101. Mitteilung Deutsche Einheitsmethoden zur Untersuchung von Fetten, Fettprodukten und verwandten Stoffen, 78. Mitt.: Analyse von Fetten XXV. Analyse von fettreichen Lebensmitteln III. *Fett Wissen. Technol.-Fat Sci. Technol.*, 89 (1987) 435-438.

504 Lolova, D., Tencheva, Zh. and Iordanova, I.: (Separation and determination of preservatives in cosmetic products by thin-layer chromatography and high-performance liquid chromatography). *Khig. Zdraveopaz.*, 30, No. 3 (1987) 74-79; *C.A.*, 107 (1987) 183298y.

36c. *Various technical products*

505 Chiarotti, M., Fucci, N., Lucchini, P. and Carnevale, A.: Thin-layer and high-performance liquid chromatographic analyses of organic gunshot residues. *Nippon Hoigaku Zasshi*, 41, No. 2 (1987) 65-70; *C.A.*, 107 (1987) 170054r.

38. INORGANIC COMPOUNDS

38a. *Cations*

506 Luo, H. and Wang, H.: (Thin-layer chromatographic separation of partial metal ions. II. Study of *n*-propyl alcohol systems). *Wuhan Daxue Xuebao, Ziran Kexueban*, No. 1 (1987) 79-84; *C.A.*, 107 (1987) 189752p.

507 Ma, G.: (Paper chromatographic separation-spectrophotometric determination of antimony in gold (thallium) ores). *Fenxi Huaxue*, 15 (1987) 478; *C.A.*, 107 (1987) 189890g.

508 Maslowska, J. and Mlodzikowski, Z.: Chitosan as a new adsorbent for thin-layer chromatography. Separation of metal ions contained in "Polfamiks" mixtures. *Chem. Anal. (Warsaw)*, 31 (1986) 193-199; *C.A.*, 107 (1987) 223408t.

509 Matkovskii, K.I., Martynenko, S.S. and Chekhovskaya, L.M.: (Determination of lead, aluminium, zinc and copper by peak paper chromatography). *U.S.S.R. Pat. SU 1,308,894 (Cl. GO1N30/90)*, 07 May 1987, Appl. 4,018,371, 04 Feb. 1986; *C.A.*, 107 (1987) 167973j.

- 510 Wang, J. and Hu, C.: (TLC separation of rare earth elements with P 507 system). *Huaxue Tongbao*, No. 6 (1987) 29-30; *C.A.*, 107 (1987) 189748s.

See also 453.

38b. Anions

- 511 Qin, A.: (Paper chromatographic separation of sodium triphosphate). *Fenxi Huaxue*, 15 (1987) 573; *C.A.*, 107 (1987) 167819p.

39. RADIOACTIVE AND OTHER ISOTOPE COMPOUNDS

- 512 Gennaro, G.P., Schramm, E., Murthy, Y.S., Sulner, J.W., Karesh, S.M. and Rossi, C.: Rapid chromatographic determination of radiochemical impurities in ^{99m}Tc -mebrofenin (Choletec). *Nucl. Med. Biol.*, 14 (1987) 535; *C.A.*, 107 (1987) 205317h.
- 513 Kazikiewicz, J.M., Zimmer, A.M., Spies, S.M. and Rosen, S.T.: Rapid miniaturized chromatography procedures for iodinated monoclonal antibodies: comparison to gel exclusion chromatography. *J. Nucl. Med. Technol.*, 15 (1987) 129-321; *C.A.*, 107 (1987) 214186s.
- 514 Zimmer, A.M., Kazikiewicz, J.M., Rosen, S.T. and Spies, S.M.: Chromatographic evaluation of the radiochemical purity of sodium ^{131}I -iodine: effect on monoclonal antibody labeling. *Nucl. Med. Biol.*, 14 (1987) 533-534; *C.A.*, 107 (1987) 171526h.

Electrophoresis

2. FUNDAMENTALS, THEORY AND GENERAL

2a. General

- 274 Dukhin, S.S., Vidybida, A.K., Dukhin, A.S. and Serikov, A.A.: (Aperiodic electrophoresis. Direct drift of disperse particles in a uniform alternating unharmonic electric field). *Kolloidn. Zh.*, 49 (1987) 853-857; *C.A.*, 107 (1987) 223835e.
- 275 Tondeur, D. and Bailly, M.: Unifying concepts in non-linear unsteady processes part II. Multicomponent waves, competition and diffusion. *Chem. Eng. Process.*, 22 (1987) 91-105; *C.A.*, 107 (1987) 179104q.

2b. Thermodynamics and theoretical relationships

- 276 Kuhn, R., Wagner, H., Mosher, R.A. and Thormann, W.: Experimental and theoretical investigation of the stability of stepwise pH gradients in continuous flow electrophoresis. *Electrophoresis (Weinheim)*, 8 (1987) 503-508.
- 277 Shimao, K.: (Theory of steady state electrophoresis and its application to Tiselius' electrophoresis). *Seibutsu Butsuri Kagaku*, 31 (1987) 241-250; *C.A.*, 107 (1987) 219659h.
- 278 Viovy, J.L.: Theory of resonant electrophoresis. *C.R. Acad. Sci., Ser. 2*, 305 (1987) 181-184; *C.A.*, 107 (1987) 184241u.

2d. Measurement of physico chemical and related values

See 432.

3. GENERAL TECHNIQUES

3a. Apparatus and accessories

- 279 Ffrench-Constant, R.H. and Devonshire, A.L.: A multiple homogenizer for rapid sample preparation in immunoassays and electrophoresis. *Biochem. Genet.*, 25 (1987) 493-499; *C.A.*, 107 (1987) 171700k.
- 280 Gardiner, K.J., Laas, W.A. and Patterson, D.: Pulsed-field gradient gel electrophoretic apparatus. *Eur. Pat. Appl. EP 234,836 (Cl.G01N27/26)*, 02 Sep. 1987, US Appl. 833,691, 27 Feb. 1986; 27 pp.; *C.A.*, 107 (1987) 228133j.
- 281 Hinson, W.G., Pipkin, J.L., Anson, J.F., Casciano, D.A. and Burns, E.R.: Micro-size polyacrylamide gel electrophoresis system. *Rev. Sci. Instrum.*, 58 (1987) 1736-1742; *C.A.*, 107 (1987) 150491c.
- 282 Hjerten, S.: Coating for electrophoresis tube. *U.S. Pat. US 4,680,201 (Cl.427-230; B05D7/22)*, 14 Jul. 1987, Appl. 792,724, 30 Oct. 1985, 6 pp.; *C.A.*, 107 (1987) 194573k.
- 283 Rhodes, P.H. et al.: Moving wall, continuous flow electrophoresis apparatus. *U.S. Pat. Appl. US 904,128*, 15 Jul. 1987, Appl. 05 Sep. 1986; 17 pp.; *C.A.*, 107 (1987) 228231g.
- 284 Serwer, P.: Apparatus for bidimensional electrophoretic separations. *U.S. Pat. US 4,693,804 (Cl.204-182.1; G01N27/26)*, 15 Sep. 1987, 12 pp.; *C.A.*, 107 (1987) 214480h.
- 285 Skorniyakov, E.P., Zolotarev, P.P., Khokhlov, V.N. and Lelikov, Yu.A.: (Method for electrophoretic separation of mixtures of substances). *U.S.S.R. Pat. SU 1,323,937 (Cl.G01N27/26)*, 15 Jul. 1987, Appl. 3,901,265, 27 May, 1985; *C.A.*, 107 (1987) 172001v.

- 286 Tombs, M.P.: Method for electrophoretic separation. *PCT Int. Pat.*, Appl. WO 87 00,635 (Cl.G01N27/26), 29 Jan. 1987, GB Appl. 85/18,030, 17 Jul. 1985, 25 pp.; *C.A.*, 107 (1987) 150781x.

3b. Detection procedures and detectors

- 287 Horowitz, P.M. and Bowman, S.: Ion-enhanced fluorescence staining of sodium dodecyl sulfate-polyacrylamide gels using bis(8-*p*-toluidino-1-naphthalene-sulfonate). *Anal. Biochem.*, 165 (1987) 430-434.
- 288 Pfisterer, J.: (Blotting - a new generation of detection methods). *Bioengineering*, (1987) 84-89; *C.A.*, 107 (1987) 194331e.
- 289 Ricci, G., Caccuri, A.M., Bello, M.L., Solinas, S.P. and Nardini, M.: Ketimine rings: useful detectors of enzymatic activities in solution and on polyacrylamide gel. Detection of L-amino acid oxidase, glutamine transaminase, pantetheinase and acylase I. *Anal. Biochem.*, 165 (1987) 356-361.

See also 306, 333, 348, 352, 498, 501, 522, 529.

3c. Electrophoresis in stabilized media

- 290 Ogawa, M. and Shiraishi, H.: (A medium for electrophoresis). *Jpn. Kokai Tokkyo Koho Pat.* JP 61,296,256 (86,296,256) (Cl.G01N27/26), 27 Dec. 1986, Appl. 85/139,680, 25 Jun. 1985, 9 pp.; *C.A.*, 107 (1987) 171958a.
- 291 Ogawa, M. and Shiraishi, H.: (A medium for electrophoresis). *Jpn. Kokai Tokkyo Koho Pat.* JP 61,296,258 (86,296,258) (Cl.G01N27/26), 27 Dec. 1986, Appl. 85/139,682, 25 Jun. 1985, 10 pp.; *C.A.*, 107 (1987) 171959b.
- 292 Radola, B.J. and Allen, R.C.: Rehydratable polyacrylamide gels. *Eur. Pat.* Appl. EP 225,403 (Cl.C08F220/56), 16 Jun. 1987, Appl. 85/308,438, 20 Nov. 1985, 9 pp.; *C.A.*, 107 (1987) 176708d.
- 293 Sugihara, M. and Sugimoto, N.: (Media for electrophoresis). *Jpn. Kokai Tokkyo Koho Pat.* JP 62 66,153, (Cl.G01N27/26), 25 Mar. 1987, Appl. 85/206,090, 18 Sep. 1985, 7 pp.; *C.A.*, 107 (1987) 17200u.

4. SPECIAL TECHNIQUES

4b. Preparative and continuous procedures

- 294 Margolis, J.: Device and method for preparative gel electrophoresis. *Brit. UK Pat.* Appl. GB 2,177,211 (Cl.C25B7/00), 14 Jan. 1987, AU Appl. 85/347, 29 Apr. 1985, 11 pp.; *C.A.*, 107 (1987) 171960v.

See also 318, 347, 484.

4c. Isoelectric focusing

- 295 Righetti, P.G.: Immobilized pH gradients: the resolving power is here to stay, part II. *LC-GC*, 5 (1987) 796-802; *C.A.*, 107 (1987) 150480y.

See also 328, 392, 541, 591.

4d. Isotachopheresis

- 296 Snopek, J., Jelinek, I. and Smolkova-Keulemansova, E.: Use of cyclodextrins in isotachopheresis. II. α -, β - and γ -cyclodextrins as leading electrolyte additives for the separation of *ortho*-, *meta*- and *para*-substituted halogenobenzoic acids. *J. Chromatogr.*, 411 (1987) 153-159.
- 297 Yagi, T.: (Leading solutions for isotachopheresis). *Jpn. Kokai Tokkyo Koho Pat.* JP 62,167,462 (87,167,462) (Cl.G01N27/26), 23 Jul. 1987, Appl. 85/277,624, 09 Dec. 1985, 7 pp.; *C.A.*, 107 (1987) 214494r.

See also 320, 339, 347, 587, 595.

4e. Two dimensional electrophoresis

- 298 Endler, A.T., Young, D.S. and Tracy, R.P.: Methodology for high-resolution two-dimensional gel electrophoresis. *Cancer Invest.*, 5 (1987) 127-149; *C.A.*, 107 (1987) 171628t.
- 299 Gianazza, E., Rabilloud, T., Quaglia, L., Caccia, P., Astrua-Testori, S., Osio, L., Grazioli, G. and Righetti, P.H.: Additives for immobilized pH gradient two-dimensional separation of particulate material: comparison between commercial and new synthetic detergents. *Anal. Biochem.*, 165 (1987) 247-257.
- See also 284, 325, 328, 353, 361, 366, 367, 369, 371, 372, 379, 411, 416, 477.

4f. Affinity electrophoresis

- 300 Breborowicz, J., Gan, J. and Klosin, J.: Application of monoclonal antibodies for affinity electrophoresis. *J. Immunol. Methods.*, 102 (1987) 101-107; *C.A.*, 107 (1987) 196025g.

4g. Other special techniques

- 301 De la Vigne, U.: Computer controlled evaluation of thin-layer chromatograms and electrophoresis objects. *Symp. Biol. Hung.*, 34 (Chromatography '85) (1986) 591-603; *C.A.*, 107 (1987) 150724f.
- 302 Deutsch, J.M.: Dynamics of pulsed-field electrophoresis. *Phys. Rev. Lett.*, 59 (1987) 1255-1258; *C.A.*, 107 (1987) 171843j.
- 303 Schaper, B.: (Immunofixation-electrophoresis). *Labor.-Med.*, 10 (1987) 351-353; *C.A.*, 107 (1987) 215793z.
- 304 Sofer, W. and Martin, P.: Analysis of densitometric data obtained from electrophoretic analysis. *Comput. Appl. Biosci.*, 3 (1987) 129; *C.A.*, 107 (1987) 171910d.
- 305 Tsuda, T.: Modification of electroosmotic flow with cetyltrimethylammonium bromide in capillary zone electrophoresis. *J. High Resolut. Chromatogr. Chromatogr. Commun.*, 10 (1987) 622-624.

See also 333.

10. CARBOHYDRATES

10a. Mono and oligosaccharides. Structural studies

- 306 Romay, C. and Pascual, C.: Nitroblue tetrazolium staining of serum fructosamine on agarose gel electrophoretograms. *Clin. Chem. (Winston-Salem)*, 33 (1987) 1949.

10b. Polysaccharides, mucopolysaccharides, lipopolysaccharides

- 307 Candiano, G., Ghiggeri, G.M., Piccardo, M.T., Bertelli, R., Barboro, P., Lazzarini, G., Rialdi, G. and Vecchio, G.: Purification of intact concanavalin A in tetramers by isoelectric focusing. *J. Chromatogr.*, 423 (1987) 319-326.
- 308 Sugahara, K., Okamoto, H., Nakamura, M., Shibamoto, S. and Yamashina, I.: Developmental changes in glycosaminoglycan sulfotransferase activities in animal sera. *Arch. Biochem. Biophys.*, 258 (1987) 391-403.
- 309 Vogel, K.G. and Evanko, S.P.: Proteoglycans of fetal bovine tendon. *J. Biol. Chem.*, 262 (1987) 13607-13613.

10c. Glycoproteins and their components

- 310 Epstein, H., Polson, A. and van der Merwe, K.J.: A novel method of electrophoresis for the separation of individual human serum glycoproteins. *Prepar. Biochem.*, 17 (1987) 109-120.
- 311 Gianazza, E., Catto, N., Righetti, P.G. and Bianchi-Bosisio, A.: Alpha-1 acid glycoprotein analysis on immobilized pH gradients. *Electrophoresis (Weinheim)*, 8 (1987) 538-540.

- 312 Heimer, R., Molinaro, L., Jr. and Sampson, P.M.: Detection by ^{125}I -cationized cytochrome *c* of proteoglycans and glycosaminoglycans immobilized on unmodified and on positively charged nylon 66. *Anal. Biochem.*, 165 (1987) 448-455.
- 313 Holt, G.D., Haltiwanger, R.S., Torres, C.-R. and Hart, G.W.: Erythrocytes contain cytoplasmic glycoproteins. O-Linked GlcNAc on band 4.1. *J. Biol. Chem.*, 262 (1987) 14847-14850.
- 314 James, W.M. and Agnew, W.S.: Multiple oligosaccharide chains in the voltage-sensitive Na channel from *Electrophorus electricus*: evidence for α -2,8-linked polysialic acid. *Biochem. Biophys. Res. Commun.*, 148 (1987) 817-826.
- 315 Moule, S.K., Peak, M., Thompson, S. and Turner, G.A.: Studies of the sialylation and microheterogeneity of human serum alpha₁-acid glycoprotein in health and disease. *Clin. Chim. Acta*, 166 (1987) 177-185.
- 316 Pala, A., Di Ruzza, A., Curtilli, L., Strom, R., Moro, M., Reverberi, L. and Carezza, L.: Interference of an α_2 -component in immunological determinations of pregnancy-specific β_1 glycoprotein in serum. *Clin. Chim. Acta*, 168 (1987) 33-46.
- 317 Smith, P.J., Sabbatini, G.P., Grant, K.I. and von Holt, C.: Identification of nuclear envelope proteins and glycoproteins which co-isolate with the nuclear protein matrix. *Biochim. Biophys. Acta*, 904 (1987) 365-372.
- 318 Stanescu, V. and Do Pham, T.: Preparative electrophoresis on agarose submerged gels of two aggregating proteoglycan monomers from articular cartilage. *Prepar. Biochem.*, 17 (1987) 229-238.
- 319 Vogel, K.G., Koob, T.J. and Fisher, L.W.: Characterization and interactions of a fragment of the core protein of the small proteoglycan (PG II) from bovine tendon. *Biochem. Biophys. Res. Commun.*, 148 (1987) 658-663.

11. ORGANIC ACIDS AND LIPIDS

11a. Organic acids and simple esters

- 320 Barth, T.: Quantitative determination of volatile carboxylic acids in formation waters by isotachopheresis. *Anal. Chem.*, 59 (1987) 2232-2237.
- 321 Fanali, S., Ossicini, L. and Sinibaldi, M.: Capillary isotachopheretic study of outer-sphere complex formation. *Chromatographia*, 23 (1987) 811-813.

11d. Lipoproteins and their constituents

- 322 Gurske, W.A.: Electrophoretic technique for separation of lipoproteins and electrophoretic gel for use therein. *U.S. Pat.* US 4,696,958 (Cl.524-21; C08L1/00) 29 Sep. 1987, Appl. 801,113, 22 Nov. 1985, 9 pp.; *C.A.*, 107 (1987) 214495s.
- 323 Huff, M.W. and Strong, W.L.P.: Separation and isolation of human apolipoproteins C-II, C-III₀, C-III₁, and C-III₂ by chromatofocusing on the Fast Protein Liquid Chromatography system. *J. Lipid Res.*, 28 (1987) 1118-1123.
- 324 Levy, E., Marcel, Y., Deckelbaum, R.J., Milne, R., Lepage, G., Seidman, E., Bendayan, M. and Roy, C.C.: Intestinal apoB synthesis, lipids, and lipoproteins in chylomicron retention disease. *J. Lipid Res.*, 28 (1987) 1263-1274.
- 325 McConathy, W.J., McCaffree, M.A. and Lane, D.M.: Polymorphism of human plasma apolipoprotein C-III at birth and in early neonatal life. *J. Lipid Res.*, 28 (1987) 1193-1198.
- 326 Pitas, R.E., Boyles, J.K., Lee, S.H., Hui, D. and Weisgraber, K.H.: Lipoproteins and their receptors in the central nervous system. Characterization of the lipoproteins in cerebrospinal fluid and identification of apolipoprotein B, E (LDL) receptors in the brain. *J. Biol. Chem.*, 262 (1987) 14352-14360.
- 327 Rinninger, F. and Pittman, R.C.: Regulation of the selective uptake of high density lipoprotein - associated cholesteryl esters. *J. Lipid Res.*, 28 (1987) 1313-1325.
- 328 Steinmetz, A.: Phenotyping of human apolipoprotein E from whole blood plasma by immunoblotting. *J. Lipid Res.*, 28 (1987) 1364-1370.
- 329 Swinkels, D.W., Hak-Lemmers, H.L.M. and Demacker, P.N.M.: Single spin density gradient ultracentrifugation method for the detection and isolation of light and heavy low density lipoprotein subfractions. *J. Lipid Res.*, 28 (1987) 1233-1239.

- 330 Traber, M.G., Kayden, H.J. and Rindler, M.J.: Polarized secretion of newly synthesized lipoproteins by the Caco-2 human intestinal cell line. *J. Lipid Res.*, 28 (1987) 1350-1363.
- 331 Yang, L.-Y. and Kuksis, A.: Size and composition of lymph chylomicrons following feeding corn oil or its fatty acid methyl esters. *Biochem. Cell Biol.*, 65 (1987) 514-524.

18. AMINO ACIDS AND PEPTIDES: CHEMICAL STRUCTURE OF PROTEINS

18a. Amino acids and their derivatives

- 332 Strous, G.J., van Kerkhof, P., Fallon, R.J. and Schwartz, A.L.: Golgi galactosyltransferase contains serine-linked phosphate. *Eur. J. Biochem.*, 169 (1987) 307-311.

See also 289, 381.

18b. Peptides and peptidic and proteinous hormones

- 333 Bergman, T. and Jörnvall, H.: Electroblothing of individual polypeptides from SDS/polyacrylamide gels for direct sequence analysis. *Eur. J. Biochem.*, 169 (1987) 9-12.
- 334 Kerr, D.E., Kissinger, L.F., Gentry, L.E., Purchio, A.F. and Shoyab, M.: Structural requirements of diacylglycerols for binding and activating phospholipid-dependent, Ca^{++} -sensitive protein kinase. *Biochem. Biophys. Res. Commun.*, 148 (1987) 776-782.
- 335 Masojidek, J., Droppa, M. and Horvath, G.: Analysis of the polypeptide composition of grana and stroma thylakoids by two-dimensional electrophoresis. Distribution of photosystem II between grana and stroma lamellae. *Eur. J. Biochem.*, 169 (1987) 283-288.
- 336 Ronin, C., Papandreou, M.-J., Canonne, C. and Weintraub, B.D.: Carbohydrate chains of human thyrotropin are differentially susceptible to endoglycosidase removal on combined and free polypeptide subunits. *Biochemistry*, 26 (1987) 5848-5853.
- 337 Ruddon, R.W., Krzesicki, R.F., Norton, S.E., Beebe, J.S., Peters, B.P. and Perini, F.: Detection of a glycosylated, incompletely formed form of chorionic gonadotropin β subunit that is a precursor of hormone assembly in tropoblastic cells. *J. Biol. Chem.*, 262 (1987) 12533-12540.
- 338 Sinha, Y.N. and Jacobsen, B.P.: Glycosylated prolactin in the murine pituitary: detection by a novel assay and alteration of concentrations by physiological and pharmacological stimuli. *Biochem. Biophys. Res. Commun.*, 148 (1987) 505-514.
- 339 Stehle, P. and Fuerst, P.: Isotachophoretic control of peptide synthesis and purification. *Symp. Biol. Hung.*, 34 (Chromatography '85) (1986) 179-191; *C.A.*, 107 (1987) 154749r.

See also 487, 534, 572.

18c. Elucidation of structure of proteins and enzymes

- 340 Larsson, U.K., Sundby, C. and Andersson, B.: Characterization of two different subpopulations of spinach light-harvesting chlorophyll a/b-protein complex (LHC II): polypeptide composition, phosphorylation pattern and association with Photosystem II. *Biochim. Biophys. Acta*, 894 (1987) 59-68.
- 341 Little, M., Quinlan, R.A., Röhrich, C. and Diez, J.: Structural differences between blood-platelet tubulin and other mammalian tubulins. *Biochim. Biophys. Acta*, 916 (1987) 93-98.
- 342 Masojidek, J., Droppa, M. and Horvath, G.: Organization of the photosynthetic membrane in maize mesophyll and bundle sheath chloroplasts studied by two-dimensional gel electrophoresis. *Biochim. Biophys. Acta*, 894 (1987) 49-58.
- 343 Seshagiri, P.B. and Adiga, P.R.: Isolation and characterization of a biotin-binding protein from the pregnant-rat serum and comparison with that from the chicken egg-yolk. *Biochim. Biophys. Acta*, 916 (1987) 474-481.

- 344 Stangl, D., Wiederkehr, F., Suter, F. and Zuber, H.: Structure and function of L-lactate dehydrogenases from thermophilic and mesophilic bacteria. V. The complete amino-acid sequence of the mesophilic L-lactate dehydrogenase from *Bacillus megaterium*. *Biol. Chem. Hoppe-Seyler*, 368 (1987) 1157-1166.

19. PROTEINS

19a. General techniques

- 345 Cookson, E.J. and Beynon, R.J.: Degradation artefacts during sample preparation for sodium dodecyl sulfate polyacrylamide gel electrophoresis. *Biosci. Rep.*, 7 (1987) 209-215; *C.A.*, 107 (1987) 194454x.
- 346 Dunn, M.J. (Editor): *Gel Electrophoresis of Proteins*. J. Wright, Bristol, 1986, 407 pp.; *C.A.*, 107 (1987) 150746q.
- 347 Johansson, G., Ofverstedt, L.-G. and Hjerten, S.: Preparative displacement electrophoresis (isotachopheresis) of proteins on cellulose columns. *Anal. Biochem.*, 166 (1987) 267-275.
- 348 Lee, C., Levin, A. and Branton, D.: Copper staining: a five-minute protein stain for sodium dodecyl sulfate-polyacrylamide gels. *Anal. Biochem.*, 166 (1987) 308-312.
- 349 Luescher, I.F.: Adsorption, immunoadsorption and sodium dodecyl sulfate-polyacrylamide gel electrophoresis of proteins using nitrocellulose membrane filters. *Electrophoresis (Weinheim)*, 8 (1987) 508-514.
- 350 Mitichkin, O.V., Lepskii, A.A., Zemlyakov, S.A. and Stoyalova, L.R.: (Method of separating a protein fraction). *U.S.S.R. Pat. SU 1,296,919 (Cl.G01N27/26)*, 15 Mar. 1987, Appl. 3,811,410, 29 Oct. 1984; *C.A.*, 107 (1987) 150785b.
- 351 Onishi, A. and Kawakatsu, S.: (A highly sensitive detection method for a protein or nucleic acid). *Jpn. Kokai Tokkyo Koho Pat. JP 62 35,262 (87 35,262) (Cl.G01N33/68)*, 16 Feb. 1987, Appl. 85/174,030, 09 Aug. 1985, 10 pp.; *C.A.*, 107 (1987) 171999q.
- 352 Onishi, A. and Kawakatsu, S.: (A highly sensitive detection method for a protein or nucleic acid). *Jpn. Kokai Tokkyo Koho Pat. JP 62 35,261 (87 35,261) (Cl.G01N33/68)*, 16 Feb. 1987, Appl. 85/174,029, 09 Aug. 1985, 9 pp.; *C.A.*, 107 (1987) 194572j.
- 353 Perrett, C.W.: High-resolution two-dimensional gel electrophoresis of proteins: A clinical tool in Brain research. *Anal. Proc.*, 24 (1987) 303-304; *C.A.*, 107 (1987) 194461x.
- 354 Schägger, H. and von Jagow, G.: Tricine-sodium dodecyl sulfate-polyacrylamide gel electrophoresis for the separation of proteins in the range from 1 to 100 kDa. *Anal. Biochem.*, 166 (1987) 368-379.

See also 287, 592.

19b. Proteins of cells, viruses and subcellular particles

- 355 Boulanger, P.A., Yoshinaga, S.K. and Berk, A.J.: DNA-binding properties and characterization of human transcription factor TFIIC2. *J. Biol. Chem.*, 262 (1987) 15098-15105.
- 356 Brodie, A.E. and Reed, D.J.: Reversible oxidation of glyceraldehyde 3-phosphate dehydrogenase thiols in human lung carcinoma cells by hydrogen peroxide. *Biochem. Biophys. Res. Commun.*, 148 (1987) 120-125.
- 357 Christou, M., Wilson, N.M. and Jefcoate, C.R.: Expression and function of three cytochrome P-450 isozymes in rat extrahepatic tissues. *Arch. Biochem. Biophys.*, 258 (1987) 519-534.
- 358 Chu, A., Saito, A. and Fleischer, S.: Preparation and characterization of longitudinal tubules of sarcoplasmic reticulum from fast skeletal muscle. *Arch. Biochem. Biophys.*, 258 (1987) 13-23.
- 359 Kantor, R.R.S., Mattes, M.J., Lloyd, K.O., Old, L.J. and Albino, A.P.: Biochemical analysis of two cell surface glycoprotein complexes, very common antigen 1 and very common antigen 2. Relationship to very late activation T cell antigens. *J. Biol. Chem.*, 262 (1987) 15158-15165.

- 360 Kisilevsky, R. and Gore, J.: A comparison of accessibility of ribosomal proteins on free and membrane-bound ribosomes: the ribosomal proteins potentially involved in ribosome-membrane binding. *Biochim. Biophys. Acta*, 910 (1987) 282-291.
- 361 Klösser-Ziegelmayr, S. and Cleve, H.: Two-dimensional electrophoresis of cellular proteins of human cultured fibroblasts: Search for sex-chromosome-dependent variants. *Electrophoresis (Weinheim)*, 8 (1987) 518-523.
- 362 Kubo, K. and Fukuda, M.: (Biological membrane protein isolation by electrophoresis). *Jpn. Kokai Tokkyo Koho Pat.* JP 62 22,796 (Cl.C07K3/14), 30 Jan. 1987, Appl. 85/163,230, 24 Jul. 1985, 6 pp.; *C.A.*, 107 (1987) 195157q.
- 363 Miki, M., Tamai, H., Mino, M., Yamamoto, Y. and Niki, E.: Free-radical chain oxidation of rat red blood cells by molecular oxygen and its inhibition by α -tocopherol. *Arch. Biochem. Biophys.*, 258 (1987) 373-380.
- 364 Miyazaki, K., Ashida, Y., Kihira, Y., Mashima, K., Yamashita, J. and Horio, T.: Transformation of rat liver cell line by Rous sarcoma virus causes loss of cell surface fibronectin, accompanied by secretion of metallo-proteinase that preferentially digests the fibronectin. *J. Biochem. (Tokyo)*, 102 (1987) 569-582.
- 365 Morii, H., Takisawa, H. and Yamamoto, T.: A possible role of protein phosphorylation in the inactivation of a Ca^{2+} -induced Ca^{2+} release channel from skeletal muscle sarcoplasmic reticulum. *J. Biochem. (Tokyo)*, 102 (1987) 263-271.
- 366 Pankov, R.G., Uschewa, A.A., Tasheva, B.T. and Markov, G.G.: Vertebrate liver cytokeratins: a comparative study. *Biochem. Cell Biol.*, 65 (1987) 547-557.
- 367 Remy, R., Ambard-Bretteville, F. and Colas des Francs, C.: Analysis by two-dimensional gel electrophoresis of the polypeptide composition of pea mitochondria isolated from different tissues. *Electrophoresis (Weinheim)*, 8 (1987) 528-532.
- 368 Sakane, F., Takahashi, K., Takayama, H. and Koyama, J.: Stabilizing effect of glutaraldehyde on the respiratory burst NADPH oxidase of guinea pig polymorphonuclear leukocytes. *J. Biochem. (Tokyo)*, 102 (1987) 247-253.
- 369 Volodina, T.V., Koloskov, E.E., Shishkin, S.S. and Kozel'tsev, V.L.: (Two-dimensional electrophoresis in studies of rat liver mitochondrial proteins). *Vopr. Med. Khim.*, 33 (1987) 25-28; *C.A.*, 107 (1987) 150637e.
- 370 Wang, L.-H., Tsai, S.Y., Sagami, I., Tsai, M. and O'Malley, B.W.: Purification and characterization of chicken ovalbumin upstream promoter transcription factor from HeLa cells. *J. Biol. Chem.*, 262 (1987) 16080-16086.

19c. Proteins synthesized by genetic manipulation

- 371 Olsen, I., Rosseland, S.K., Thorsrud, A.K. and Jellum, E.: Differentiation between *Haemophilus parrophophilus*, *H. aphrophilus*, *H. influenzae*, *Actinobacillus actinomycetemcomitans*, *Pasteurella multocida*, *P. haemolytica*, and *P. ureae* by high resolution two-dimensional protein electrophoresis. *Electrophoresis (Weinheim)*, 8 (1987) 532-535.
- 372 Rademacher, B.E. and Steele, W.J.: Affinity chromatography of reticulocyte lysates on haptoglobin improves two-dimensional mapping of translation products. *Electrophoresis (Weinheim)*, 8 (1987) 524-528.
- 373 Yamada, H., Oshima, N., Mizuno, T., Matsui, H., Kai, Y., Noguchi, H. and Mizushima, S.: Use of a series of ompF-ompC chimeric proteins for locating antigenic determinants recognized by monoclonal antibodies against the ompC and ompF proteins of the *Escherichia coli* outer membrane. *J. Biochem. (Tokyo)*, 102 (1987) 455-464.
- 374 Zhang, Y.: (Experiments in mammalian genetic engineering. Experiment 4: Agarose gel electrophoresis). *Yichuan*, 9 (1987) 42-44; *C.A.*, 107 (1987) 194333g.

19d. Microbial and plant proteins

- 375 Daley, L.S. and Theriot, L.J.: Electrophoretic analysis, redox activity, and other characteristics of proteins similar to purothionins from tomato (*Lycopersicon esculenta*), mango (*Mangifera indica*), papaya (*Careia papaya*) and walnut (*Juglans regia*). *J. Agric. Food Chem.*, 35 (1987) 680-687.
- 376 Freier, T. and Rüdiger, H.: *In vivo* binding partners of the *Lens culinaris* lectin. *Biol. Chem. Hoppe-Seyler*, 368 (1987) 1215-1223.

- 377 He, H. and Zhou, Y.: (Determination of proteins in edible mushrooms by CAM electrophoresis). *Shiyong Jun*, (1987) 39-40; *C.A.*, 107 (1987) 152987t.
- 378 Hussain, A., Ramirez, H., Bushuk, W. and Roca, W.: Identification of cultivars of forage legume (*Desmodium ovalifolium* Guill et Perr.) by their electrophoretic patterns. *Can. J. Plant Sci.*, 67 (1987) 713-717; *C.A.*, 107 (1987) 171842h.
- 379 Krishnan, H.B. and Peuppke, S.G.: Heat shock triggers rapid protein phosphorylation in soybean seedlings. *Biochem. Biophys. Res. Commun.*, 148 (1987) 762-767.
- 380 Reddy, A.S.N. and Poovacah, B.W.: Accumulation of a glycine rich protein in auxin-deprived strawberry fruits. *Biochem. Biophys. Res. Commun.*, 147 (1987) 885-891.
- 381 Rieul, C., Cortay, J.-C., Bleicher, F. and Cozzone, A.J.: Effect of bacteriophage M13 infection on phosphorylation of dnaK protein and other *Escherichia coli* proteins. *Eur. J. Biochem.*, 168 (1987) 621-627.
- 382 Rmbeli, R., Wirth, M. and Zuber, H.: Crosslinking of phycobiliproteins from the cyanobacterium *Mastigocladus laminosus* with bis-imidates. Localization of an intrasubunit and an intersubunit crosslink in C-phycocyanin. *Biol. Chem. Hoppe-Seyler*, 368 (1987) 1179-1191.
- 383 Semino, G.A. and Cerletti, P.: Effect of preliminary thermal treatment on the digestion by trypsin of lupin seed protein. *J. Agric. Food Chem.*, 35 (1987) 656-660.

See also 340, 342.

19e. *Proteins of blood, serum and blood cells*

- 384 Backer, E.T., Harff, G.A. and Beyer, C.: A patient with an IgG paraprotein and complexes of lactate dehydrogenase and IgG in the serum. *Clin. Chem. (Winston-Salem)*, 33 (1987) 1937-1938.
- 385 Baumstark, J.S.: The elastase inhibitors of swine serum: an improved method for the isolation of α_2 -macroglobulin. *Prepar. Biochem.*, 17 (1987) 195-208.
- 386 Btikofer, P., Brodbeck, U. and Ott, P.: Modulation of red cell vesiculation by protease inhibitors. *Biochim. Biophys. Acta*, 904 (1987) 259-267.
- 387 Churchi, F.C., Treanor, R.E., Sherrill, G.B. and Whinna, H.C.: Carboxylate polyanions accelerate inhibition of thrombin by heparin cofactor II. *Biochem. Biophys. Res. Commun.*, 148 (1987) 362-368.
- 388 Conradt, H.S., Egge, H., Peter-Katalinic, J., Reiser, W., Siklosi, T. and Schaper, K.: Structure of the carbohydrate moiety of human interferon- β secreted by a recombinant chinese hamster ovary cell line. *J. Biol. Chem.*, 262 (1987) 14600-14605.
- 389 Erdjument, H., Lane, D.A., Ireland, H., Panico, M., Di Marzo, V., Blench, I. and Morris, H.R.: Formation of a covalent disulfide-linked antithrombin-albumin complex by an antithrombin variant, antithrombin "Northwick Park". *J. Biol. Chem.*, 262 (1987) 13381-13384.
- 390 Fan, P. and Shen, Z.: (Acrylamide gradient SDS slab gel electrophoresis of α_2 -macroglobulin). *Xibao Shengwuxue Zazhi*, 9 (1987) 35-38; *C.A.*, 107 (1987) 214373a.
- 391 Goldberger, G., Paz, M.A., Torrelino, B.M., Okamoto, Y. and Gallop, P.M.: Effect of hydroxyorganoboranes on synthesis, transport and N-linked glycosylation of plasma proteins. *Biochem. Biophys. Res. Commun.*, 148 (1987) 493-499.
- 392 Haas, H. and Schlaak, M.: Conventional isoelectric focusing does not dissociate immune complexes. *J. Immunol. Methods*, 103 (1987) 79-85; *C.A.*, 107 (1987) 215825m.
- 393 Hardouin, S., Hossenlopp, P., Segovia, B., Seurin, D., Portolan, G., Lassarre, C. and Binoux, M.: Heterogeneity of insulin-like growth factor binding proteins and relationships between structure and affinity. 1. Circulating forms in man. *Eur. J. Biochem.*, 170 (1987) 121-132.
- 394 Heeb, M.J., Espana, F., Geiger, M., Collen, D., Stump, D.C. and Griffin, J.H.: Immunological identity of heparin-dependent plasma and urinary protein C inhibitor and plasminogen activator inhibitor-3. *J. Biol. Chem.*, 262 (1987) 15813-15816.
- 395 Hossenlopp, P., Seurin, D., Segovia, B., Portolan, G. and Binoux, M.: Heterogeneity of insulin-like growth factor binding proteins and relationships between structure and affinity. 2. Forms released by human and rat liver in culture. *Eur. J. Biochem.*, 170 (1987) 133-142.

- 396 Joles, J.A. and Kortlandt, W.: Massive overestimation of albumin in analbuminemic rat serum by bromocresol green. *Clin. Chem. (Winston-Salem)*, 33 (1987) 2121.
- 397 Knight, E., Jr.: Purification of the interferons. *J. Interferon Res.*, 7 (1987) 501-503; *C.A.*, 107 (1987) 215796c.
- 398 Leahy, D.T. and McLaughlin, H.: A second albumin variant in an Irish population. *Clin. Chim. Acta*, 168 (1987) 273-279.
- 399 Lollar, P., Parker, C.G., Kajenski, P.J., Litwiller, R.D. and Fass, D.N.: Degradation of coagulation proteins by an enzyme from Malayan pit viper (*Akistrodon rhodostoma*) venom. *Biochemistry*, 26 (1987) 7627-7636.
- 400 Meijers, J.C.M., Tijburg, P.N.M. and Bouma, B.N.: Inhibition of human blood coagulation factor Xa by α_2 -macroglobulin. *Biochemistry*, 26 (1987) 5932-5937.
- 401 Minchiotti, L., Galliano, M., Iadarola, P., Stoppini, M., Ferri, G. and Castellani, A.A.: Structural characterization of two genetic variants of human serum albumins. *Biochim. Biophys. Acta*, 916 (1987) 411-418.
- 402 Pepys, M.B. and Butler, P.J.G.: Serum amyloid P component is the major calcium-dependent specific DNA binding protein of the serum. *Biochem. Biophys. Res. Commun.*, 148 (1987) 308-313.
- 403 Pezzoli, A. and Pascali, E.: Bence Jones proteinuria in multiple sclerosis. (*Clin. Chem. (Winston-Salem)*), 33 (1987) 1923-1924.
- 404 Pratt, C.W., Roche, P.A. and Pizzo, S.V.: The role of inter- α -trypsin inhibitor and other proteinase inhibitors in the plasma clearance of neutrophil elastase and plasmin. *Arch. Biochem. Biophys.*, 258 (1987) 591-599.
- 405 Sasaki, T., Kobayashi, K. and Ozeki, T.: Interaction of silkworm larval hemolymph antitrypsin and bovine trypsin. *J. Biochem. (Tokyo)*, 102 (1987) 433-441.
- 406 Thornton, R.D. and Kirby, E.P.: Isolation and characterization of an inhibitor of factor XIIIa from bovine plasma. *J. Biol. Chem.*, 262 (1987) 12714-12721.
- 407 Turpeinen, U., Koivunen, E. and Stenman, U.-H.: Liquid-chromatographic determination of β_2 -microglobulin, α_1 -acid glycoprotein, and albumin in urine. *Clin. Chem. (Winston-Salem)*, 33 (1987) 1756-1760.
- 408 Vertosick, F.T., Jr. and Kelly, R.H.: Stability of model immune complexes during zone electrophoresis and isoelectric focusing in agarose gels. *Clin. Chem. (Winston-Salem)*, 33 (1987) 1741-1746.
- 409 Widdershoven, J., van Munster, P., De Abreu, R., Bosman, H., van Lith, T., van der Putten-van Meyel, M., Motohara, K. and Matsuda, I.: Four methods compared for measuring des-carboxy-prothrombin (PIVKA II). *Clin. Chem. (Winston-Salem)*, 33 (1987) 2074-2078.
- 410 Wilhelm, O., Hafter, R. and Graeff, H.: Fibrinolysis in ascitic fluid: isolation and characterization of fibrin derivatives, their interaction with albumin. *Thromb. Res.*, 47 (1987) 673-681; *C.A.*, 107 (1987) 171952u.
- 411 Yokohama, M., Gawahara, H., Manabe, T., Okuyama, T. and Mogi, K.: On the identification map of thoroughbred horse plasma proteins by microscale multi-sample two-dimensional electrophoresis. *Nippon Chikusan Gakkaiho*, 58 (1987) 245-252; *C.A.*, 107 (1987) 214368c.

See also 311, 341, 460, 489.

19f. Structural and muscle proteins

- 412 Bag, J. and Pramanik, S.: Attachment of mRNA to the cytoskeletal framework and translational control of gene expression in rat L6 muscle cells. *Biochem. Cell Biol.*, 65 (1987) 565-575.
- 413 Colombatti, A. and Bonaldo, P.: Biosynthesis of chick type VI collagen. II. Processing and secretion in fibroblasts and smooth muscle cells. *J. Biol. Chem.*, 262 (1987) 14461-14466.
- 414 Colombatti, A., Bonaldo, P., Ainger, K., Bressan, G.M. and Volpin, D.: Biosynthesis of chick type VI collagen. I. Intracellular assembly and molecular structure. *J. Biol. Chem.*, 262 (1987) 14454-14460.
- 415 Diano, M., LeBivic, A. and Hirn, M.: A method for the production of highly specific polyclonal antibodies. *Anal. Biochem.*, 166 (1987) 224-229.
- 416 Gianazza, E., Osio, L., Grazioli, G., Astrua-Testori, S., Righetti, P.G., Accinni, R., Renoldi, I. and Repossini, A.: An examination of heart proteins by two-dimensional electrophoresis. *Clin. Chem. (Winston-Salem)*, 33 (1987) 2011-2018.

- 417 Meng, J., Li, P. and Chen, S.: (Analysis of the composition of wool keratin by two-dimensional polyacrylamide gel electrophoresis). *Fangzhi Xuebao*, 8 (1987) 95-97; *C.A.*, 107 (1987) 177868t.
- 418 Pipoly, D.J. and Crouch, E.C.: Degradation of native type IV procollagen by human neutrophil elastase. Implications for leukocyte-mediated degradation of basement membranes. *Biochemistry*, 26 (1987) 5748-5754.
- 419 Samarel, A.M., Ferguson, A.G. and Burgess, L.: Quantitative analysis of myofibrillar protein subunits: demonstration of large molar excesses of myosin light chains in rabbit ventricular myocardium. *J. Mol. Cell. Cardiol.*, 19 (1987) 699-707; *C.A.*, 107 (1987) 195465p.
- 420 Shapiro, S.Z.: Elimination of the detection of an artefactual 65 kDa keratin band from immunoblots. *J. Immunol. Methods*, 102 (1987) 143-146; *C.A.*, 107 (1987) 171877y.
- 421 Strohsacker, M.W., Minnich, M.D., Clark, M.A., Shorr, R.G.L. and Crooke, S.T.: Selective purification of cardiac myosin by a high-performance salicylate affinity column. *J. Chromatogr.*, 435 (1988) 185-192.
- 422 Wu, J.-J., Eyre, D.R. and Slayter, H.S.: Type VI collagen of the intervertebral disc. Biochemical and electron-microscopic characterization of the native protein. *Biochem. J.*, 248 (1987) 373-381.

See also 410.

19g. Protamines, histones and other chromosomal proteins

- 423 Ali, Z. and Singh, N.: Binding of linker histones to the core nucleosome. *J. Biol. Chem.*, 262 (1987) 12989-12993.
- 424 Davie, J.R., Delcuve, G.P., Nickel, B.E., Moirier, R. and Bailey, G.: Reduced levels of histones H1^o and H1b, and unaltered content of methylated DNA in rainbow trout hepatocellular carcinoma chromatin. *Cancer Res.*, 47 (1987) 5407-5410.
- 425 Duschak, V.G.: Modified techniques for histone H1 identification in polyacrylamide gels. *Commun. Biol.*, 5 (1987) 415-424; *C.A.*, 107 (1987) 194456z.
- 426 Giacotti, V., Pani, B., D'Andrea, P., Berlingieri, M.T., Di Fiore, P.P., Fusco, A., Vecchio, G., Crane-Robinson, C. and Goodwin, G.H.: Histone and nonhistone proteins from normal and virus-transformed rat thyroid epithelial cells. *Basic Appl. Histochem.*, 31 (1987) 229-238; *C.A.*, 107 (1987) 214462d.
- 427 Holst, M. and Egyhazi, E.: Differential kinase systems are involved in the rapidly turning over phosphorylation of prominent nuclear proteins. *Biochim. Biophys. Acta*, 931 (1987) 224-233.
- 428 Karpenchuk, K.G. and Klimyuk, V.I.: (Identification of plant histones by non-ionic detergent binding: evolutionary argument). *Mol. Biol.*, 21 (1987) 981-991; *C.A.*, 107 (1987) 214370x.
- 429 Shimidzu, M., Shindo, H., Takahashi, K., Taniguchi, S. and Matsumoto, U.: *In vivo* phosphorylation of histones H1 and H5 in calf thymus and chicken erythrocyte as studied by ³¹P nuclear magnetic resonance spectroscopy. *J. Biochem. (Tokyo)*, 102 (1987) 351-358.
- 430 Spiker, S., Hopkins, R., Fischer, R. and Quatrano, R.S.: Synthesis of nucleosomal histone variants during wheat grain development. *Biochim. Biophys. Acta*, 910 (1987) 157-162.

19h. Chromoproteins and metalloproteins

- 431 Constantinidis, I. and Satterlee, J.D.: Isoelectric focusing purity criteria and ¹NMR detectable spectroscopic heterogeneity in the major isolated monomer hemoglobins from *Glycera dibranchiata*. *Biochemistry*, 26 (1987) 7779-7786.
- 432 De Haën, C., Molecular weight standards for calibration of gel filtration and sodium dodecyl sulfate-polyacrylamide gel electrophoresis: ferritin and apoferritin. *Anal. Biochem.*, 166 (1987) 235-245.
- 433 Gelfi, C., Righetti, P.G., Roviida, E. and Samaja, M.: Purification of human hemoglobin valence intermediates by preparative immobilized pH gradients. *J. Biochem. Biophys. Methods*, 14 (1987) 139-147; *C.A.*, 107 (1987) 150733h.
- 434 Hase, T., Harabayashi, M., Kawai, K. and Matsubara, H.: A carboxyl-terminal hydrophobic region of yeast cytochrome c₁ is necessary for functional assembly into complex III of the respiratory chain. *J. Biochem. (Tokyo)*, 102 (1987) 411-419.

- 435 Kawano, S., Kamataki, T., Yasumori, T., Yamazoe, Y. and Kato, R.: Purification of human liver cytochrome P-450 catalyzing testosterone 6 β -hydroxylation. *J. Biochem. (Tokyo)*, 102 (1987) 493-501.
- 436 Lasker, J.M., Raucy, J., Kubota, S., Blowski, B.P., Black, M. and Lieber, C.S.: Purification and characterization of human liver cytochrome P-450-ALC. *Biochem. Biophys. Res. Commun.*, 148 (1987) 232-238.
- 437 Petren, S.: Determination of transferrin isoproteins in human cerebrospinal fluid using isoelectric focusing and zone immunoelectrophoresis assay. *Electrophoresis (Weinheim)*, 8 (1987) 515-517.
- 438 Sanchez-Cabezudo, M., Martin, C. and Rodenas, S.: (New technique for determination of the glycosylated hemoglobins by isoelectric focusing. Comparative analysis with the colorimetric method). *An. R. Acad. Farm.*, 53 (1987) 103-114; *C.A.*, 107 (1987) 171850j.
- 439 Sczekan, S.R. and Joshi, J.G.: Isolation and characterization of ferritin from soybeans (*Glycine max*). *J. Biol. Chem.*, 262 (1987) 13780-13788.

19i. *Proteins of glands, gland products and various zymogens (including milk proteins)*

- 440 Cheng, C.Y. and Bardin, C.W.: Identification of two testosterone-responsive testicular proteins in Sertoli cell-enriched culture medium whose secretion is suppressed by cells of the intact seminiferous tubule. *J. Biol. Chem.*, 262 (1987) 12768-12779.
- 441 Haque, Z., Kristjansson, M.M. and Kinsella, J.E.: Interaction between K-casein and β -lactoglobulin: possible mechanism. *J. Agric. Food Chem.*, 35 (1987) 644-649.
- 442 Pandey, K.N., Inagami, T., Girard, P.R., Kuo, J.F. and Misono, K.S.: New signal transduction mechanisms of atrial natriuretic factor: inhibition of phosphorylation of protein kinase C and A 240 kDa protein in adrenal cortical plasma membrane by cGMP dependent and independent mechanisms. *Biochem. Biophys. Res. Commun.*, 148 (1987) 589-595.

See also 369, 547.

19j. *Proteins of brain, cerebrospinal fluid and eye*

- 443 Bizozero, O.A., McGarry, J.F. and Lees, M.B.: Autoacylation of myelin proteolipid protein with acyl coenzyme A. *J. Biol. Chem.*, 262 (1987) 13550-13557.
- 444 Frappier, T., Regnouf, F. and Pradel, L.A.: Binding of brain spectrin to the 70-kDa neurofilament subunit protein. *Eur. J. Biochem.*, 169 (1987) 651-657.
- 445 Gardner, E.E. and Dahl, D.: Immunological study of a neurofilament protein variant (S150) with polyclonal and monoclonal antibodies. *Biochim. Biophys. Acta*, 926 (1987) 280-286.
- 446 Jackson, T.R., Patterson, S.I., Wong, Y.H. and Hanley, M.R.: Bradykinin stimulation of inositol phosphate and calcium responses is insensitive to pertussis toxin in NG115-401L neuronal cells. *Biochem. Biophys. Res. Commun.*, 148 (1987) 412-416.
- 447 Sakihara, H.: (A biochemical study on neurofilament proteins in soft tissue tumors). *Nippon Seikeigeka Gakkai Zasshi*, 6 (1987) 553-561; *C.A.*, 107 (1987) 171911e.
- 448 Shinozawa, T., Sokabe, M., Terada, S., Matsusaka, H. and Yoshizawa, T.: Detection of cyclic GMP binding protein and ion channel activity in frog rod outer segments. *J. Biochem. (Tokyo)*, 102 (1987) 281-290.
- 449 Townsend, L.E., Gilroy, J., LeWitt, P., Wolfe, D.E., Pomara, N., Weinbraub, J. and Reitz, D.: Comparison of methods for analysis of CSF proteins in patients with Alzheimer's disease. *Neurochem. Pathol.*, 6 (1987) 213-229; *C.A.*, 107 (1987) 150719h.
- 450 Van den Oetelaar, P.J.M. and Hoenders, H.J.: Electrophoretic titration curve in 6 M urea of the bovine eye lens protein α -crystallin. *J. Chromatogr.*, 411 (1987) 507-509.
- 451 Wang, C., Reen, C., Chiu, H.C. and Lin, K.T.: Characterization of multiple gamma globulin bands of cerebrospinal fluid on electrophoresis. *T'ai-wan I Hsueh Hui Tsa Chih*, 86 (1987) 480-483; *C.A.*, 107 (1987) 150626a.

- 452 Wollner, D.A., Messner, D.J. and Catterall, W.A.: $\beta 2$ Subunits of sodium channels from vertebrate brain. Studies with subunit-specific antibodies. *J. Biol. Chem.*, 262 (1987) 14709-14715.

See also 553.

19k. *Proteins of neoplastic tissue and transformed cells*

- 453 Bray, K.R., Koda, J.E. and Gaur, P.K.: Serum levels and biochemical characteristics of cancer-associated antigen CA-549, a circulating breast cancer marker. *Cancer Res.*, 47 (1987) 5853-5860.
- 454 Burchell, J., Gendler, S., Taylor-Papadimitriou, J., Girling, A., Lewis, A., Millis, R. and Lampert, D.: Development and characterization of breast cancer reactive monoclonal antibodies direct to the core protein of the human milk mucin. *Cancer Res.*, 47 (1987) 5476-5482.
- 455 Clair, T., Miller, W.R. and Cho-Chung, Y.S.: Prognostic significance of the expression of a ras protein with a molecular weight of 21,000 by human breast cancer. *Cancer Res.*, 47 (1987) 5290-5293.
- 456 Furukawa, J., Shimano, T., Inaji, H., Saito, M., Matsuura, N., Monden, T. and Mori, T.: Purification and characterization of a pancreas cancer-related antigen. *Clin. Chim. Acta*, 168 (1987) 219-230.
- 457 Giometti, C.S., Gemmell, M.A., Nance, S.L., Tollaksen, S.L. and Taylor, J.: Detection of heritable mutations as quantitative changes in protein expression. *J. Biol. Chem.*, 262 (1987) 12764-12767.
- 458 Keyse, S.M. and Tyrrell, R.M.: Both near ultraviolet radiation and the oxidizing agent hydrogen peroxide induce a 32-kDa stress protein in normal human skin fibroblasts. *J. Biol. Chem.*, 262 (1987) 14821-14825.
- 459 Mackay, A.M., Tracy, R.P. and Craighead, J.E.: Intermediate filament proteins in asbestos-induced mesotheliomas of the rat. *Cancer Res.*, 47 (1987) 5461-5468.
- 460 Morrow, B. and Rubin, Ch.S.: Biogenesis of glycophorin A in K562 human erythroleukemia cells. *J. Biol. Chem.*, 262 (1987) 13812-13820.
- 461 Oh, S.L., Very, D.L., Walker, J., Raam, S. and Ju, S.: An analogy between fetal haptoglobin and a potent immunosuppressant in cancer. *Cancer Res.*, 47 (1987) 5120-5126.
- 462 Ternell, M., Moldawer, L.L., Lönnroth, Ch., Gelin, J. and Lundholm, K.G.: Plasma protein synthesis in experimental cancer compared to paraneoplastic conditions, including monokine administration. *Cancer Res.*, 47 (1987) 5825-5830.

See also 356.

19l. *Specific binding and receptor proteins*

- 463 Cieplak, W., Gaudin, H.M. and Eidels, L.: Diphtheria toxin receptor. Identification of specific diphtheria toxin-binding proteins on the surface of Vero and BS-C-1 cells. *J. Biol. Chem.*, 262 (1987) 13246-13253.
- 464 Danzo, B.J., Eller, B.C. and Bell, B.W.: The apparent molecular weight of androgen-binding protein (ABP) in the blood of immature rats differs from that of ABP in the epididymis. *J. Steroid Biochem.*, 28 (1987) 411-419.
- 465 Estes, P.A., Suba, E.J., Lawler-Heavner, J., Elashry-Stowers, D., Wei, L.L., Toft, D.O., Sullivan, W.P., Horwitz, K.B. and Edwards, D.P.: Immunologic analysis of human breast cancer progesterone receptors. 1. Immunoaffinity purification of transformed receptors and production of monoclonal antibodies. *Biochemistry*, 26 (1987) 6250-6262.
- 466 Fishman, J.B., Dickey, B.F. and Fine, R.E.: Purification and characterization of the rat liver vasopressin (V1) receptor. *J. Biol. Chem.*, 262 (1987) 14049-14055.
- 467 Fowler, V.M.: Identification and purification of a novel M_r 43,000 tropomyosin-binding protein from human erythrocyte membranes. *J. Biol. Chem.*, 262 (1987) 12792-12800.
- 468 Fricker, G., Schneider, S., Gerok, W. and Kurz, G.: Identification of different transport systems for bile salts in sinusoidal and canalicular membranes of hepatocytes. *Biol. Chem. Hoppe-Seyler*, 368 (1987) 1143-1150.

- 469 Goldsmith, P., Gierschik, P., Milligan, G., Unson, C.G., Vinitsky, R., Malech, H.L. and Spiegel, A.M.: Antibodies directed against synthetic peptides distinguish between GTP-binding proteins in neutrophil and brain. *J. Biol. Chem.*, 262 (1987) 14683-14688.
- 470 Hayashi, F., Sumi, M. and Amakawa, T.: Phosphatidylinositol stimulates phosphorylation of protein components I and II in rod outer segments of frog photoreceptors. *Biochem. Biophys. Res. Commun.*, 148 (1987) 54-60.
- 471 Hofmann, K., Romovacek, H., Titus, G., Ridge, K., Raffensperger, J.A. and Finn, F.M.: The rat liver insulin receptor. *Biochemistry*, 26 (1987) 7384-7390.
- 472 Horvath, G., Fernoe, M. and Trope, C.: Comparison of estrogen receptor enzyme immunoassay and a radioligand binding assay (isoelectric focusing) in adenocarcinoma of the uterine body. *Neoplasma*, 34 (1987) 313-317; *C.A.*, 107 (1987) 171938u.
- 473 Karpf, D.B., Arnaud, C.D., King, K., Bambino, T., Winer, J., Nyireddy, K. and Nissenson, R.A.: The canine renal parathyroid hormone receptor is a glycoprotein: characterization and partial purification. *Biochemistry*, 26 (1987) 7825-7833.
- 474 Klausner, R.D., O'Shea, J.J., Luong, H., Ross, P., Bluestone, J.A. and Samelson, L.E.: T cell receptor tyrosine phosphorylation. Variable coupling for different activating ligands. *J. Biol. Chem.*, 262 (1987) 12654-12659.
- 475 Kwok, Y.C. and Yip, C.C.: Tyrosine phosphorylation of two cytosolic proteins of 50 kDa and 35 kDa in rat liver by insulin-receptor kinase *in vitro*. *Biochem. J.*, 248 (1987) 27-33.
- 476 Lee, R.T. and Lee, Y.C.: Affinity labeling of the galactose/N-acetylgalactosamine-specific receptor of rat hepatocytes: preferential labeling of one of the subunits. *Biochemistry*, 26 (1987) 6320-6329.
- 477 Manabe, T., Higuchi, N. and Okuyama, T.: Staining of serum concanavalin A-binding proteins after micro two-dimensional electrophoresis and blotting. *J. Chromatogr.*, 423 (1987) 115-122.
- 478 Nemoto, T., Ohara-Nemoto, Y. and Ota, M.: Purification and characterization of a nonhormone-binding component of the nontransformed glucocorticoid receptor from rat liver. *J. Biochem. (Tokyo)*, 102 (1987) 513-523.
- 479 Nukatsuka, M. and Nagasawa, S.: Characterization of the interaction between human protein S and C4b-binding protein(C4bp). *J. Biochem. (Tokyo)*, 102 (1987) 599-605.
- 480 O'Connor-McCourt, M.D. and Wakefield, L.M.: Latent transforming growth factor- β in serum. A specific complex with α_2 -macroglobulin. *J. Biol. Chem.*, 262 (1987) 14090-14099.
- 481 Prozialeck, W.C., Wallace, T.L. and Weiss, B.: Differential inhibition of calmodulin-sensitive phosphodiesterase and Ca^{++} -adenosine triphosphatase by chlorpromazine-linked calmodulin. *J. Pharmacol. Exp. Ther.*, 243 (1987) 171-179.
- 482 Rehmus, E.H., Howard, K.J., Janiga, K.E. and Distelhorst, C.W.: Immunochemical comparison of mutant glucocorticoid receptors and wild type receptor fragments produced by neutrophil elastase and chymotrypsin. *J. Steroid Biochem.*, 28 (1987) 167-177.
- 483 Seshagiri, P.B. and Adiga, P.R.: Identification and molecular characterization of a biotin-binding protein distinct from avidin of chicken egg white and comparison with yolk biotin-binding protein. *Biochim. Biophys. Acta*, 926 (1987) 321-330.
- 484 Starita-Geribaldi, M., Poiree, J.-C. and Sudaka, P.: Use of mild detergent gel electrophoresis for isolation and characterization of the kidney brush border D-glucose transporter. *Anal. Biochem.*, 165 (1987) 406-413.
- 485 Stephenson, J.D. and Shepherd, V.L.: Purification of the human alveolar macrophage mannose receptor. *Biochem. Biophys. Res. Commun.*, 148 (1987) 883-889.
- 486 Van Driel, I.R., Davis, C.G., Goldstein, J.L. and Brown, M.S.: Self-association of the low density lipoprotein receptor mediated by the cytoplasmic domain. *J. Biol. Chem.*, 262 (1987) 16127-16134.
- 487 Wolff, J.M., Rathjen, F.G., Frank, R. and Roth, S.: Biochemical characterization of polypeptide components involved in neurite fasciculation and elongation. *Eur. J. Biochem.*, 168 (1987) 551-561.

See also 375, 550, 553.

18m. Urinary proteins

- 488 Bilobrov, V.M., Mironov, O.L. and Khil'ko, B.S.: (The urine protein disc-electrophoresis on polyacrylamide gel). *Lab. Delo*, (1987) 678-680; *C.A.*, 107 (1987) 194457a.
- 489 Marshall, T. and Williams, K.M.: Simplified sodium dodecyl sulfate-polyacrylamide gel electrophoresis of unconcentrated urine with enhanced resolution and detection sensitivity. *Clin. Chem. (Winston-Salem)*, 33 (1987) 1886-1887.
- 490 Norden, A.G.W., Fulcher, L.M. and Flynn, F.V.: Immunoglobulin light-chain immunoblots of urine proteins from patients with tubular and Bence-Jones proteinuria. *Clin. Chim. Acta*, 166 (1987) 307-315.

19n. Other proteins

- 491 Bolton, D.C., Bendheim, P.E., Marmorstein, A.D. and Potempska, A.: Isolation and structural studies of the intact scrapie agent protein. *Arch. Biochem. Biophys.*, 258 (1987) 579-590.
- 492 Mitchell, H.K. and Petersen, N.S.: Spontaneous fragmentation of several proteins in *Drosophila* pupae. *J. Biol. Chem.*, 262 (1987) 14298-14304.
- 493 Weidman, P.J., Teller, D.C. and Shapiro, B.M.: Purification and characterization of proteoliasin, a coordinating protein in fertilization envelope assembly. *J. Biol. Chem.*, 262 (1987) 15076-15084.

20. ENZYMES AND ENZYME ACTIVITY ESTIMATION

- 494 Cheng, M.L. and VandeBerg, J.L.: Cryopreservation of erythrocytes in small aliquots for isozyme electrophoresis. *Biochem. Genet.*, 25 (1987) 411-435; *C.A.*, 107 (1987) 171095s.
- 495 Naka, K.: (Lectin-affinity electrophoresis. Isoenzyme analysis). *Kensa to Gijutsu*, 15 (1987) 881-885; *C.A.*, 107 (1987) 213847w.

20a. Oxidoreductases

- 496 Akao, T., Akao, T. and Kobashi, K.: Purification and characterization of 7 β -hydroxyteroid dehydrogenase from *Ruminococcus* sp. of human intestine. *J. Biochem. (Tokyo)*, 102 (1987) 613-619.
- 497 Dehal, S.S. and Croteau, R.: Metabolism of monoterpenes: specificity of the dehydrogenases responsible for the biosynthesis of camphor, 3-thujone, and 3-isothujone. *Arch. Biochem. Biophys.*, 258 (1987) 287-291.
- 498 Eberman, R. and Pichorner, H.: Improved method for the detection of peroxidase isoenzymes after electrophoretic separation with chlorophyllin, cysteamine and light as a hydrogen peroxide generating system. *J. Chromatogr.*, 410 (1987) 466-469.
- 499 Finel, M.: Studies on the oligomeric state of isolated cytochrome oxidase using cross-linking reagents. *Biochim. Biophys. Acta*, 894 (1987) 174-179.
- 500 Haumont, P.Y., Thomas, M.-A., Labeyrie, F. and Lederer, F.: Amino-acid sequence of the cytochrome-b₅-like heme-binding domain from *Hansenula anomala* flavocytochrome b₂. *Eur. J. Biochem.*, 169 (1987) 539-546.
- 501 Isono, H., Horie, K., Katoh, H. and Miyaura, S.: Detection of alcohol dehydrogenase by using NAD after separation by gel electrophoresis. *Eisei Kagaku*, 33 (1987) 191-199; *C.A.*, 107 (1987) 193612k.
- 502 Kanemitsu, F.: Isoenzyme specificities of immunoglobulins isolated from lactate dehydrogenase-immunoglobulin complexes. *Clin. Chim. Acta*, 167 (1987) 147-154.
- 503 Kuppe, A., Mrsny, R.J., Shimizu, M., Firsan, S.J., Keana, J.F.W. and Griffith, O.H.: Labeling of bovine heart cytochrome c oxidase with analogues of phospholipids. Synthesis and reactivity of a new cardiolipin benzaldehyde probe. *Biochemistry*, 26 (1987) 7693-7701.
- 504 Moses, G.C. and Henderson, A.R.: Increased serum lactate dehydrogenase isoenzyme 1 and "flipped" LD-1/LD-2 ratio in myopathy associated with partial carnitine palmitoyltransferase deficiency. *Clin. Chem. (Winston-Salem)*, 33 (1987) 2111-2113.

- 505 Penning, T.M. and Sharp, R.B.: Prostaglandin dehydrogenase activity of purified rat liver 3 α -hydroxysteroid dehydrogenase. *Biochem. Biophys. Res. Commun.*, 148 (1987) 646-652.
- 506 Taroni, F., Gellera, C. and Di Domato, S.: Evidence for two distinct mitochondrial malic enzymes in human skeletal muscle: purification and properties of the NAD(P)⁺-dependent enzyme. *Biochim. Biophys. Acta*, 916 (1987) 446-454.
- 507 Van Pee, K.-H., Sury, G. and Lingens, F.: Purification and properties of a nonheme bromoperoxidase from *Streptocymes aureofaciens*. *Biol. Chem. Hoppe-Seyler*, 368 (1987) 1225-1232.

See also 289.

20b. *Transferases (excluding E.C. 2.7.-.-)*

- 508 Baer, W. and Biedermann, V.: (Glutamate-pyruvate-transaminase: a new preparation by means of agarose-electrophoresis). *Beitr. Gerichtl. Med.*, 45 (1987) 181-183; *C.A.*, 107 (1987) 193625s.
- 509 Burlina, A., Secchiero, S., Bertorelle, R. and Zaninotto, M.: An improved electrophoretic method for γ -glutamyltransferase isoenzyme demonstration. *Med. Sci. Res.*, 15 (1987) 499-500; *C.A.*, 107 (1987) 171062d.
- 510 Franklin, R.B., Kukoyi, B.I., Akuffo, V. and Costello, L.C.: Testosterone stimulation of mitochondrial aspartate aminotransferase levels and biosynthesis in rat ventral prostate. *J. Steroid Biochem.*, 28 (1987) 247-256.
- 511 Horowitz, P.M. and Bowman, S.: Oxidation increases the proteolytic susceptibility of a localized region in rhodanese. *J. Biol. Chem.*, 262 (1987) 14544-14548.
- 512 Kanemitsu, F.: Electrophoretic and kinetic characterization of two phenotypes of soluble cytoplasmic L-aspartate:2-oxoglutarate aminotransferase in human liver tissues. *Electrophoresis (Weinheim)*, 8 (1987) 541-543.
- 513 Otterness, D.M. and Weinshilboum, R.M.: Mouse thiopurine methyltransferase pharmacogenetics: biochemical studies and recombinant inbred strains. *J. Pharmacol. Exp. Ther.*, 243 (1987) 180-186.
- 514 Scherzer, E., Auer, B. and Schweiger, M.: Identification, purification, and characterization of *Escherichia coli* virus T1 DNA methyltransferase. *J. Biol. Chem.*, 262 (1987) 15225-15231.
- 515 Schiele, F., Muller, J., Colinet, E. and Siest, G.: Production and certification of an enzyme reference material for γ -glutamyltransferase (CRM 319). Part 1. Preparation and characterization. *Clin. Chem. (Winston-Salem)*, 33 (1987) 1971-1977.

See also 289.

20c. *Transferases transferring phosphorus containing groups (E.C. 2.7.-.-)*

- 516 Azhar, S., Butte, J. and Reaven, E.: Calcium-activated, phospholipid-dependent protein kinases from rat liver: subcellular distribution, purification, and characterization of multiple forms. *Biochemistry*, 26 (1987) 7047-7057.
- 517 Blenis, J., Kuo, C.J. and Erikson, R.L.: Identification of a ribosomal protein S6 kinase regulated by transformation and growth-promoting stimuli. *J. Biol. Chem.*, 262 (1987) 14373-14376.
- 518 Cadena, D.L. and Dahmus, M.E.: Messenger RNA synthesis in mammalian cells is catalyzed by the phosphorylated form of RNA polymerase II. *J. Biol. Chem.*, 262 (1987) 12468-12474.
- 519 Caizergues-Ferrer, M., Belenguer, P., Lapeyre, B., Amalric, F., Wallace, M.O. and Olson, M.O.J.: Phosphorylation of nucleolin by a nucleolar type NII protein kinase. *Biochemistry*, 26 (1987) 7876-7883.
- 520 Coughlan, S.J. and Hind, G.: A protein kinase that phosphorylates light-harvesting complex is autophosphorylated and is associated with photosystem II. *Biochemistry*, 26 (1987) 6515-6521.
- 521 Levine, H., III and Sahyoun, N.E.: Characterization of a soluble M_r-30000 catalytic fragment of the neuronal calmodulin-dependent protein kinase II. *Eur. J. Biochem.*, 168 (1987) 481-486.
- 522 Mochizuki, Y.: (Methods of electrophoresis and staining of starfish hexokinase). *Igaku to Seibutsugaku*, 114 (1987) 47-49; *C.A.*, 107 (1987) 149807d.

- 523 Nealon, D.: Isoelectric focusing of the cytosolic human creatine kinase isoenzymes. *Clin. Chim. Acta*, 166 (1987) 73-78.
- 524 Ohmura, Y.: (Detection by SDS-PAGE of DNA kinase and protein tyrosine kinase in mammalian cell nucleus). *Ochanomizu Igaku Zasshi*, 35 (1987) 37-47; *C.A.*, 107 (1987) 193744e.
- 525 Schneider, G.J., Tumer, N.E., Richaud, C., Borbely, G. and Haselkorn, R.: Purification and characterization of RNA polymerase from the cyanobacterium *Anabaena* 7120. *J. Biol. Chem.*, 262 (1987) 14633-14639.
- 526 Shin, Y.S., Niedermeier, H.P., Endres, W., Schaub, J., Weidinger, S.: Agarose gel isoelectrofocusing of UDP-galactose pyrophosphorylase and galactose-1-phosphate uridyltransferase. Developmental aspect of UDP-galactose pyrophosphorylase. *Clin. Chim. Acta*, 166 (1987) 27-35.
- 527 Staal, G.E.J., Kalff, A., Heesbeen, E.C., van Veelen, C.W.M. and Rijkssen, G.: Subunit composition, regulatory properties, and phosphorylation of phosphofruktokinase from human gliomas. *Cancer Res.*, 47 (1987) 5047-5051.
- 528 Tombes, R.M. and Shapiro, B.M.: Enzyme termini of a phosphocreatine shuttle. Purification and characterization of two creatine kinase isozymes from sea urchin sperm. *J. Biol. Chem.*, 47 (1987) 16011-16019.
- 529 Williams, J. and Marshall, T.: Improved Nitro Blue Tetrazolium staining of creatine kinase isoenzymes following agarose gel electrophoresis or isoelectric focusing in polyacrylamide gels. *Electrophoresis (Weinheim)*, 8 (1987) 536-537.
- 530 Wol'fson, A.D., Motorin, Yu.A., Orlovsky, A.F. and Gladilin, K.L.: (Isolation, polypeptide content and properties of the aminoacyl-tRNA-synthetase complex from rabbit liver). *Biokhimiya (Moscow)*, 52 (1987) 1847-1854.

See also 569.

20d. *Hydrolases, acting on ester bonds (E.C. 3.1.-.-)*

- 531 Asano, R. and Hokari, S.: Isoelectric focusing of nonspecific esterases in polyacrylamide microgels. *Jpn. J. Vet. Sci.*, 48 (1986) 1015-1019; *C.A.*, 107 (1987) 149804a.
- 532 Azhayeva, E.V., Severin, S.E. and Kondratyev, A.D.: (cGMP-Dependent phosphodiesterase of cyclic nucleotides from human lymphoid cells). *Biokhimiya (Moscow)*, 52 (1987) 1781-1785.
- 533 Bennett, C.F. and Crooke, S.T.: Purification and characterization of a phosphoinositide-specific phospholipase C from guinea pig uterus. Phosphorylation by protein kinase C *in vivo*. *J. Biol. Chem.*, 262 (1987) 13789-13797.
- 534 Doolittle, M.H., Wong, H., Davis, R.C. and Schotz, M.C.: Synthesis of hepatic lipase in liver and extrahepatic tissues. *J. Lipid Res.*, 28 (1987) 1326-1334.
- 535 Holm, C., Belfrage, P. and Fredrikson, G.: Immunological evidence for the presence of hormone-sensitive lipase in rat tissues other than adipose tissue. *Biochem. Biophys. Res. Commun.*, 148 (1987) 99-105.
- 536 Kinnunen, L., Hukkanen, V., Ewart, K. and Hovi, T.: Improved sensitivity of restriction endonuclease analysis of herpes simplex virus type 2 DNA with polyacrylamide gradient gel electrophoresis. *J. Virol. Methods*, 16 (1987) 187-193; *C.A.*, 107 (1987) 150721c.
- 537 Ozols, J.: Isolation and characterization of a 60-kilodalton glycoprotein esterase from liver microsomal membranes. *J. Biol. Chem.*, 262 (1987) 15316-15321.
- 538 Rosendahl, K., Waldenlind, L. and Onica, D.: Microheterogeneity of serum alkaline phosphatase isoenzymes as revealed by isoelectric focusing. *Clin. Chim. Acta*, 168 (1987) 297-306.
- 539 Zlotnick, G.W., Mackow, M.C. and Gottlieb, M.: Renaturation of *Leishmania donovani* 3'-nucleotidase following sodium dodecyl sulfate-polyacrylamide gel electrophoresis. *Comp. Biochem. Physiol., B: Comp. Biochem.*, 87B (1987) 629-635; *C.A.*, 107 (1987) 149983h.

See also 559.

20e. *Hydrolases, acting on glycosyl compounds (E.C. 3.2.-.-)*

- 540 Brown, R.K.: Purification of N-acetylgalactosaminidase by isoelectric focusing. Avail. *Univ. Microfilm Int.*, Order No. AD-A 173478/9/GAR, 1987, 21 pp.; *C.A.*, 107 (1987) 149885c.

- 541 Farkas, V., Kolarova, N. and Labudova, I.: Complexation with carrier ampholytes as the possible source of artifacts during isoelectric focusing of cellulase. *Biologia*, 42 (1987) 327-333; *C.A.*, 107 (1987) 213854w.
- 542 Robbi, M. and Beaufay, H.: Biosynthesis of rat liver pI-6.1 esterase, a carboxylesterase of the cisternal space of the endoplasmic reticulum. *Biochem. J.*, 248 (1987) 545-550.
- 543 Tsuji, A., Yang, R.C., Omura, K., Imabayashi, T. and Suzuki, Y.: A simple differential immunoprecipitation assay of urinary acid and neutral alpha-glucosidases for glycogenosis II. *Clin. Chim. Acta*, 167 (1987) 313-320.
- 544 Uhm, T-B., Jeon, D-Y., Buyn, S.M., Hong, J-S. and Groot Wassink, J.W.D.: Purification and properties of β -fructofuranosidase from *Aspergillus niger*. *Biochim. Biophys. Acta*, 926 (1987) 119-126.
- 545 White, J.W., Jr.: Wiley led the way: a century of federal honey research. *J. Assoc. Off. Anal. Chem.*, 70 (1987) 181-189 - a review with 52 refs.

20f. Other hydrolases

- 546 Ievleva, E.V., Zimacheva, A.V. and Mosolov, V.V.: (Effect of free and immobilized cystatin on proteinases of *Carica papaya* latex). *Biokhimiya (Moscow)*, 52 (1987) 1786-1791.
- 547 Kisiel, W., Kondo, S., Smith, K.J., McMullen, B.A. and Smith, L.F.: Characterization of a protein C activator from *Agkistrodon contortrix contortrix* venom. *J. Biol. Chem.*, 262 (1987) 12607-12613.
- 548 Leger, R.J.S., Cooper, R.M. and Charnley, A.K.: Distribution of chymoelastases and trypsin-like enzymes in five species of entomopathogenic Deuteromycetes. *Arch. Biochem. Biophys.*, 258 (1987) 123-131.
- 549 Lunardi, J., Garin, J., Issartel, J.-P. and Vignais, P.V.: Mapping of nucleotide-depleted mitochondrial F_1 -ATPase with 2-azido- $[\alpha\text{-}^{32}\text{P}]$ adenosine diphosphate. Evidence for two nucleotide binding sites in the β subunit. *J. Biol. Chem.*, 262 (1987) 15172-15181.
- 550 Mileykovskaya, E.I., Abuladze, A.N. and Ostrovsky, D.N.: Subunit composition of the H^+ -ATPase complex from an anaerobic bacterium *Lactobacillus casei*. *Eur. J. Biochem.*, 168 (1987) 703-708.
- 551 Nishimura, Y., Higaki, M. and Kato, K.: Identification of a precursor form of cathepsin D in microsomal lumen: characterization of enzymatic activation and proteolytic processing *in vitro*. *Biochem. Biophys. Res. Commun.*, 148 (1987) 335-343.
- 552 Patterson, S.D. and Bell, K.: ISO-DALT characterization of 12 "new" equine plasma protease inhibitor alleles. *Anim. Genet.*, 18 (1987) 167-180; *C.A.*, 107 (1987) 150629d.
- 553 Voeikov, V.L., Pronin, A.N., Slepak, V.Z., Chestukhin, A.N. and Shlensky, A.B.: (High molecular weight forms of GTP-binding regulatory proteins from bovine cerebellum). *Biokhimiya (Moscow)*, 52 (1987) 1766-1769.
- 554 Yamamoto, K., Ueno, E., Uemura, H. and Kato, Y.: Biochemical and immunological similarity between erythrocyte membrane aspartic proteinase and cathepsin E. *Biochem. Biophys. Res. Commun.*, 148 (1987) 267-272.
- 555 Yoshimoto, T., Sattar, A.K.M.A., Hirose, W. and Tsuru, D.: Studies on prolyl endopeptidase from carrot (*Daucus carota*): purification and enzymatic properties. *Biochim. Biophys. Acta*, 916 (1987) 29-37.

20g. Lyases

- 556 Husain, I., Sancar, G.B., Holbrook, S.R. and Sancar, A.: Mechanism of damage recognition by *Escherichia coli* DNA photolyase. *J. Biol. Chem.*, 262 (1987) 13188-13197.
- 557 Urbina, J.A.: The phosphoenolpyruvate carboxykinase of *Trypanosoma (Schizotrypanum) cruzi* epimastigotes: molecular, kinetic, and regulatory properties. *Arch. Biochem. Biophys.*, 258 (1987) 186-195.

See also 559.

20i. Ligases

- 558 Lim, F., Rohde, M., Morris, C.P. and Wallace, J.C.: Pyruvate carboxylase in the yeast *pyc* mutant. *Arch. Biochem. Biophys.*, 258 (1987) 259-264.

20j. Complex mixtures and incompletely identified enzymes

- 559 Inoue, Y., Rhee, H.-i., Watanabe, K., Murata, K. and Kimura, A.: Metabolism of 2-ketoaldehydes in mold: Purification and characterization of glyoxalase I from *Aspergillus niger*. *J. Biochem. (Tokyo)*, 102 (1987) 583-589.

21. PURINES, PYRIMIDINES, NUCLEIC ACIDS AND THEIR CONSTITUENTS

21a. Purines, pyrimidines, nucleosides, nucleotides

- 560 Goebel, U., Maas, R. and Clad, A.: Quantitative electroelution of oligonucleotides and large DNA fragments from gels and purification by electro dialysis. *J. Biochem. Biophys. Methods*, 14 (1987) 245-260; *C.A.*, 107 (1987) 194343k.
- 561 Mishra, A.P., Mishra, S.K., Singh, S. and Yadava, K.L.: Study of binary complexes of lead(II) and manganese(II) with nucleic acids constituent bases. *Proc. Indian Natl. Sci. Acad., Part A*, 53 (1987) 421-426; *C.A.*, 107 (1987) 206138n.
- 562 Rahmouni, A. and Leng, M.: Reaction of nucleic acids with *cis*-diamine-dichloro-platinum(II): interstrand cross-links. *Biochemistry*, 26 (1987) 7229-7234.
- 563 Züllli, F., Weber, H. and Zuber, H.: Structure and function of L-lactate dehydrogenases from thermophilic and mesophilic bacteria. VI. Nucleotide sequences of lactate dehydrogenase genes from the thermophilic bacteria *Bacillus stearothermophilus*, *B. caldolyticus* and *B. caldotenax*. *Biol. Chem. Hoppe-Seyler*, 368 (1987) 1167-1177.

See also 569, 579.

21b. Nucleic acids, RNA

- 564 Arai, H., Nakao, K., Saito, Y., Morii, N., Sugawara, A., Yamada, T., Itoh, H., Shiono, S., Mukoyama, M., Ohkubo, H., Nakanishi, S. and Imura, H.: Simultaneous measurement of atrial natriuretic polypeptide (ANP) messenger RNA and ANP in rat heart: evidence for a preferentially increased synthesis and secretion of ANP in left atrium of spontaneously hypertensive rats (SHR). *Biochem. Biophys. Res. Commun.*, 148 (1987) 239-245.
- 565 Bird, C.R. and Sells, B.H.: Small cytoplasmic RNAs and their location within the cytoplasm. *Biochem. Cell Biol.*, 65 (1987) 582-587.
- 566 Davison, P.J., Stalenhoef, A.F.H. and Humphries, S.E.: Apolipoprotein CII (apo CII) gene expression defect in an individual with familial apo CII deficiency. *Biochem. Biophys. Res. Commun.*, 148 (1987) 320-328.
- 567 Inagaki, Y., Tsunokawa, Y., Sakamoto, H., Hirohashi, S., Kobayashi, K., Hattori, N., Ramirez, F., Terada, M. and Sugimura, T.: Presence of different types of procollagen messenger RNAs in human hepatoma cell lines. *Biochem. Biophys. Res. Commun.*, 148 (1987) 869-875.
- 568 Lo, A.C., Liu, W., Culham, D.E. and Nazar, R.N.: Effects of ribosome dissociation on the structure of the ribosome-associated 5.8S RNA. *Biochem. Cell Biol.*, 65 (1987) 536-542.
- 569 Meienhofer, M.-C., De Medicis, E., Cognet, M. and Kahn, A.: Regulation of genes for glycolytic enzymes in cultured rat hepatoma cell lines. *Eur. J. Biochem.*, 169 (1987) 237-243.
- 570 Miki, K., Sugimoto, E. and Kitagawa, Y.: Reversible interconversion between primitive endoderm- and parietal endoderm-like F9 cells demonstrated by mRNA expression. *J. Biochem. (Tokyo)*, 102 (1987) 385-392.
- 571 Pompidou, A., Corral, M., Michel, P., Defer, N., Kruh, J. and Curran, T.: The effects of phorbol ester and Ca ionophore on c-fos and c-myc expression and on DNA synthesis in human lymphocytes are not directly related. *Biochem. Biophys. Res. Commun.*, 148 (1987) 435-442.
- 572 Pramanik, S.K. and Bag, J.: Translation of an mRNA in rat L6 muscle cells is regulated within the cell cycle. *Eur. J. Biochem.*, 170 (1987) 59-67.

See also 351, 352.

21c. *Nucleic acids, DNA*

- 573 Aronovitch, J., Godinger, D., Goldstein, S. and Czapski, G.: Substitution reactions of iron-bleomycin-DNA with EDTA, DETAPAC or desferrioxamine. *J. Antibiot.*, 40 (1987) 1344-1348.
- 574 Curran, J. and Webster, J.M.: Identification of nematodes using restriction fragment length differences and species-specific DNA probes. *Can. J. Plant Pathol.*, 9 (1987) 162-166; *C.A.*, 107 (1987) 215113c.
- 575 Diekmann, S. and Lilley, D.M.J.: The anomalous gel migration of a stable cruciform: temperature and salt dependence, and some comparisons with craved DNA. *Nucleic Acids Res.*, 15 (1987) 5765-5774; *C.A.*, 107 (1987) 193247v.
- 576 Jones, C. and Su, R.T.: Association of viral and plasmid DNA with the nuclear matrix during productive infection. *Biochim. Biophys. Acta*, 910 (1987) 52-62.
- 577 Murasugi, A., Takemori, N. and Hashimoto, S.: Discrimination and quantitative analysis of wild type and point mutant early region 1B genes of adenovirus using oligodeoxyribonucleotide hybridization probes. *J. Biochem. (Tokyo)*, 102 (1987) 627-633.
- 578 Okino, S.T., Quattrochi, L.C., Pendurthi, U.R., McBride, O.W. and Tukey, R.H.: Characterization of multiple human cytochrome P-450 1 cDNAs. The chromosomal localization of the gene and evidence for alternate RNA splicing. *J. Biol. Chem.*, 262 (1987) 16072-16079.
- 579 Takenaka, S. and Takagi, M.: (Separation and analysis of DNA and oligonucleotides). *Kagaku*, 42 (1987) 636-637; *C.A.*, 107 (1987) 171651v.
- 580 Thibodeau, S.N.: Use of restriction fragment length polymorphism analysis for detecting carriers of "Fragile X" syndrome. *Clin. Chem. (Winston-Salem)*, 33 (1987) 1726-1730.

See also 351, 352, 569, 570, 572.

21d. *Structural studies on RNA and RNA mapping*

- 581 Ueno, T., Takahashi, K., Matsuguchi, T., Endo, H. and Yamamoto, M.: A new leader exon identified in the rat insulin-like growth factor II gene. *Biochem. Biophys. Res. Commun.*, 148 (1987) 344-349.

21e. *Structural studies on DNA and DNA mapping*

- 582 Morohashi, K.-i., Yoshioka, H., Gotoh, O., Okada, Y., Yamamoto, K., Miyata, T., Sogawa, K., Fujii-Kuriyama, Y. and Omura, T.: Molecular cloning and nucleotide sequence of DNA of mitochondrial cytochrome P-450(118) of bovine adrenal cortex. *J. Biochem. (Tokyo)*, 102 (1987) 559-568.
- 583 Westman, W., Eriksson, S., Laas, T., Pernemalm, P.-A. and Sköld, S.E.: Separation of DNA restriction fragments by ion-exchange chromatography on FPLC columns Mono P and Mono Q. *Anal. Biochem.*, 166 (1987) 158-171.

See also 560, 563.

21f. *Complex mixtures of nucleic acids and their fragments*

- 584 Kiryanov, G.I., Smirnova, T.A., Manamshyan, G.A. and Khodosovskaya, A.M.: (Internucleosomal interaction: detection of dinucleosomal fragmentation of chromatin by micrococcal nuclease. Analysis of products of rat liver chromatin nuclei and L-cell cleavage by micrococcal nuclease). *Biokhimiya (Moscow)*, 52 (1987) 1855-1866.
- 585 Skolnik-David, H., Moore, C.L. and Sharp, P.A.: Electrophoretic separation of polyadenylation-specific complexes. *Genes Dev.*, 1 (1987) 672-682; *C.A.*, 107 (1987) 212956u.

26. ORGANOMETALLIC AND RELATED COMPOUNDS

26c. *Coordination compounds*

See 321.

27. VITAMINS AND VARIOUS GROWTH REGULATORS (NON-PEPTIDIC)

- 586 Hansen, M. and Nexø, E.: The interaction of human transcobalamin isopeptides in cerebrospinal fluid and plasma with cobalamin and the cellular acceptor. *Biochim. Biophys. Acta*, 926 (1987) 359-364.

28. ANTIBIOTICS

- 587 Kong, Y., Zhu, Y., Wang, Y. and Gao, X.: Separation and analysis of streptomycin by isotachopheresis. *Yaowu Fenxi Zazhi*, 7 (1987) 170-171; *C.A.*, 107 (1987) 161759x.

30. SYNTHETIC AND NATURAL DYES

30b. Chloroplast and other natural pigments

- 588 Bassi, R., Hoyer-Hansen, G., Barbato, R., Giacometti, G.M. and Simpson, D.J.: Chlorophyll-proteins of the photosystem II antenna system. *J. Biol. Chem.*, 262 (1987) 13333-13341.

See also 340, 342.

32. DRUG ANALYSIS

32d. Central nervous system drugs

See 481.

32e. Chemotherapeutics (except cytostatics)

- 589 Volkova, Z. and Jelinek, I.: (Methanamine hippurate determination in tablets by the capillary isotachopheresis method). *Cesk. Farm.*, 36 (1987) 254-256; *C.A.*, 107 (1987) 161788f.

33. CLINICO-CHEMICAL APPLICATIONS

33b. Complex mixtures and profiling (single compounds by cross ref. only)

See 306, 315, 316, 324, 325, 384, 396, 398, 403, 407-409, 449, 456, 489, 490, 502, 504, 523, 526, 538, 543, 580, 593.

34. FOOD ANALYSIS

34b. Complex mixtures (single compounds by cross ref. only)

See 545.

35. ENVIRONMENTAL ANALYSIS

35c. Water pollution (complex mixtures; single compounds by cross ref. only)

See 594.

36. SOME TECHNICAL PRODUCTS AND COMPLEX MIXTURES

36d. *Complex mixtures and unidentified compounds*

- 590 Kawano, M., Tsuda, R. and Hara, M.: (Agar gel electrophoresis of mushroom extracts for identifying *Lampteromyces Japonicus*). *Igaku Kenkyu.* 56 (1986) 66-71; *C.A.*, 107 (1987) 214809x.
- 591 Padyukov, L.N. and Tsvetkova, N.V.: (Use of isoelectric focusing for evaluating the composition of pollen allergen preparations). *Zh. Mikrobiol., Epidemiol. Immunobiol.*, (1987) 94-97; *C.A.*, 107 (1987) 215809j.

37. CELLS, CELLULAR PARTICLES AND SUPRAMOLECULAR STRUCTURES

- 592 Mullon, C.J.P.: Forced-flow electrophoresis for separation of proteins and viruses. Avail. *Univ. Microfilms Int.*, Order No. DA 8709279, 1986, 187 pp.; *C.A.*, 107 (1987) 171839n.
- 593 Tabaqchali, S., Silman, R. and Holland, D.: Automation in clinical microbiology: a new approach to identifying microorganisms by automated pattern matching of proteins labeled with ^{35}S -methionine. *J. Clin. Pathol.*, 40 (1987) 1070-1087; *C.A.*, 107 (1987) 171844k.

38. INORGANIC COMPOUNDS

38b. *Anions*

- 594 Fukushi, K. and Hiro, K.: (Determination of bromide ion in seawater by capillary type isotachopheresis). *Bunseki Kagaku*, 36 (1987) 712-716.
- 595 Hirokawa, T., Kobayashi, S. and Kiso, Y.: Complex-forming equilibria in isotachopheresis. VII. Evaluation of the ion-pair formation constants of phosphorus oxo acids with histidine and assessment of their separability. *J. Chromatogr.*, 410 (1987) 279-295.
- 596 Sen, A.K., Das, S.B. and Ghosh, U.D.: Studies on hydrous zirconium oxide. Part III. Electrophoretic behavior of anions on hydrous zirconium oxide impregnated papers. *J. Indian Chem. Soc.*, 64 (1987) 425-427; *C.A.*, 107 (1987) 223840c.

MONTH	J	F	M	A	M	J	J	A	S	O	N	D
Journal of Chromatography	435/1 435/2 435/3 436/1	436/2 436/3	437/1 437/2	438/1 438/2	439/1 439/2 440 441/1	441/2		The publication schedule for further issues will be published later.				
Bibliography Section		460/1		460/2		460/3						
Cumulative Indexes, Vols. 401-450												
Biomedical Applications	424/1	424/2	425/1 425/2	426/1 426/2	427/1	427/2 428/1	428/2 429					

INFORMATION FOR AUTHORS

(Detailed *Instructions to Authors* were published in Vol. 396, pp. 451-454. A free reprint can be obtained by application to the publisher, Elsevier Science Publishers B.V., P.O. Box 330, 1000 AH Amsterdam, The Netherlands.)

Types of Contributions. The following types of papers are published in the *Journal of Chromatography* and the section on *Biomedical Applications*: Regular research papers (Full-length papers), Notes, Review articles and Letters to the Editor. Notes are usually descriptions of short investigations and reflect the same quality of research as Full-length papers, but should preferably not exceed six printed pages. Letters to the Editor can comment on (parts of) previously published articles, or they can report minor technical improvements of previously published procedures; they should preferably not exceed two printed pages. For review articles, see inside front cover under Submission of Papers.

Submission. Every paper must be accompanied by a letter from the senior author, stating that he is submitting the paper for publication in the *Journal of Chromatography*. Please do not send a letter signed by the director of the institute or the professor unless he is one of the authors.

Manuscripts. Manuscripts should be typed in double spacing on consecutively numbered pages of uniform size. The manuscript should be preceded by a sheet of manuscript paper carrying the title of the paper and the name and full postal address of the person to whom the proofs are to be sent. Authors of papers in French or German are requested to supply an English translation of the title of the paper. As a rule, papers should be divided into sections, headed by a caption (e.g., Summary, Introduction, Experimental, Results, Discussion, etc.). All illustrations, photographs, tables, etc., should be on separate sheets.

Introduction. Every paper must have a concise introduction mentioning what has been done before on the topic described, and stating clearly what is new in the paper now submitted.

Summary. Full-length papers and Review articles should have a summary of 50-100 words which clearly and briefly indicates what is new, different and significant. In the case of French or German articles an additional summary in English, headed by an English translation of the title, should also be provided. (Notes and Letters to the Editor are published without a summary.)

Illustrations. The figures should be submitted in a form suitable for reproduction, drawn in Indian ink on drawing or tracing paper. Each illustration should have a legend, all the legends being typed (with double spacing) together on a separate sheet. If structures are given in the text, the original drawings should be supplied. Coloured illustrations are reproduced at the author's expense, the cost being determined by the number of pages and by the number of colours needed. The written permission of the author and publisher must be obtained for the use of any figure already published. Its source must be indicated in the legend.




References. References should be numbered in the order in which they are cited in the text, and listed in numerical sequence on a separate sheet at the end of the article. Please check a recent issue for the layout of the reference list. Abbreviations for the titles of journals should follow the system used by *Chemical Abstracts*. Articles not yet published should be given as "in press" (journal should be specified), "submitted for publication" (journal should be specified), "in preparation" or "personal communication".

Dispatch. Before sending the manuscript to the Editor please check that the envelope contains three copies of the paper complete with references, legends and figures. One of the sets of figures must be the originals suitable for direct reproduction. Please also ensure that permission to publish has been obtained from your institute.

Proofs. One set of proofs will be sent to the author to be carefully checked for printer's errors. Corrections must be restricted to instances in which the proof is at variance with the manuscript. "Extra corrections" will be inserted at the author's expense.

Reprints. Fifty reprints of Full-length papers, Notes and Letters to the Editor will be supplied free of charge. Additional reprints can be ordered by the authors. An order form containing price quotations will be sent to the authors together with the proofs of their article.

Advertisements. Advertisement rates are available from the publisher on request. The Editors of the journal accept no responsibility for the contents of the advertisements.



Three pumps for a thousand jobs.

These microprocessor-controlled pumps let you set any flow rate directly from the keypad in ml/min (or any other unit) with 1% accuracy. If you're using Wiz as a dispenser or diluter, any selected volume can be dispensed at the press of a button. Choose from two peristaltic models for flow rates from less than 1 ml/hr to 75 liters/hr. There's also a reciprocating piston model that operates at up to 100 psi.

Send now for an Isco general catalog describing these outstandingly useful instruments. In the U.S., phone toll-free (800) 228-4250. Or write:

Isco, Inc.
P.O. Box 5347
Lincoln, NE 68505
USA.

