

ISSN 0378-4355

PUBLISHED
WEEKLY

VOL. 271 NO. 2 JUNE 16, 1983

JOURNAL OF

CHROMATOGRAPHY

INTERNATIONAL JOURNAL ON CHROMATOGRAPHY, ELECTROPHORESIS AND RELATED METHODS

CHROMATOGRAPHIC REVIEWS (Vol. 27, No. 2)

edited by

Michael Lederer

ELSEVIER

JOURNAL OF CHROMATOGRAPHY

Scope. The *Journal of Chromatography* publishes papers on all aspects of chromatography, electrophoresis and related methods. Contributions consist mainly of research papers dealing with chromatographic theory, instrumental development and their applications. The section *Biomedical Applications*, which is under separate editorship, deals with the following aspects: developments in and applications of chromatographic and electrophoretic techniques related to clinical diagnosis (including the publication of normal values); screening and profiling procedures with special reference to metabolic disorders; results from basic medical research with direct consequences in clinical practice; combinations of chromatographic and electrophoretic methods with other physicochemical techniques such as mass spectrometry. In *Chromatographic Reviews*, reviews on all aspects of chromatography, electrophoresis and related methods are published.

Submission of Papers. Papers in English, French and German may be submitted, in three copies. Manuscripts should be submitted to: The Editor of *Journal of Chromatography*, P.O. Box 681, 1000 AR Amsterdam, The Netherlands, or to: The Editor of *Journal of Chromatography, Biomedical Applications*, P.O. Box 681, 1000 AR Amsterdam, The Netherlands. Review articles are invited or proposed by letter to the Editors and will appear in *Chromatographic Reviews* or *Biomedical Applications*. An outline of the proposed review should first be forwarded to the Editors for preliminary discussion prior to preparation. Submission of an article is understood to imply that the article is original and unpublished and is not being considered for publication elsewhere. For copyright regulations, see below.

Subscription Orders. Subscription orders should be sent to: Elsevier Scientific Publishing Company, P.O. Box 211, 1000 AE Amsterdam, The Netherlands. The *Journal of Chromatography* and the *Biomedical Applications* section can be subscribed to separately.

Publication. The *Journal of Chromatography* (incl. *Biomedical Applications*, *Chromatographic Reviews* and *Cumulative Author and Subject Indexes*, Vols. 251-260 and 261-270) has 27 volumes in 1983. The subscription prices for 1983 are:

J. Chromatogr. (incl. *Chromatogr. Rev.* and *Cum. Indexes Vols. 251-260 and 261-270*) + *Biomed. Appl.* (Vols. 252-278):

Dfl. 4185.00 plus Dfl. 540.00 (postage) (total ca. US\$ 1890.00)

J. Chromatogr. (incl. *Chromatogr. Rev.* and *Cum. Indexes Vols. 251-260 and 261-270*) only (Vols. 252-271):

Dfl. 3400.00 plus Dfl. 400.00 (postage) (total ca. US\$ 1520.00)

Biomed. Appl. only (Vols. 272-278):

Dfl. 1120.00 plus Dfl. 140.00 (postage) (total ca. US\$ 504.00).

Journals are automatically sent by airmail to the U.S.A. and Canada at no extra costs, and to Japan, Australia and New Zealand, with a small additional postal charge. Back volumes of the *Journal of Chromatography* (Vols. 1 through 251) are available at Dfl. 173.00 (plus postage). Claims for issues not received should be made within three months of publication of the issue. If not, they cannot be honoured free of charge. For customers in the U.S.A. and Canada, wishing additional bibliography information on this and other Elsevier journals, please contact Elsevier Science Publishing Company Inc., Journal Information Center, 52 Vanderbilt Avenue, New York, NY 10017. Tel: (212) 867-9040.

Abstracts/Contents Lists published in Analytical Abstracts, Biochemical Abstracts, Biological Abstracts, Chemical Abstracts, Chemical Titles, Current Contents/Physical, Chemical & Earth Sciences, Current Contents/Life Sciences, Index Medicus, Mass Spectrometry Bulletin, and Science Citation Index.

See page 3 of cover for Publication Schedule, Information for Authors, and information on the News Section and Advertisements.

© ELSEVIER SCIENCE PUBLISHERS B.V. — 1983

0378-4355/83/\$00.30 per page (minimum \$03.00)

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, without the prior written permission of the publisher, Elsevier Science Publishers B.V., P.O. Box 330, 1000 AH Amsterdam, The Netherlands.

Submission of an article for publication implies the transfer of the copyright from the author(s) to the publisher and entails the authors' irrevocable and exclusive authorization of the publisher to collect any sums of considerations for copying or reproduction payable by third parties (as mentioned in article 17 paragraph 2 of the Dutch Copyright Act of 1912 and in the Royal Decree of June 20, 1974 (S. 351) pursuant to article 16 b of the Dutch Copyright Act of 1912) and/or to act in or out of Court in connection therewith.

Special regulations for readers in the U.S.A. This journal has been registered with the Copyright Clearance Center, Inc. Consent is given for copying of articles for personal or internal use, or for the personal use of specific clients. This consent is given on the condition that the copier pays through the Center the per-copy fee stated in the code on the first page of each article for copying beyond that permitted by Sections 107 or 108 of the U.S. Copyright Law. The appropriate fee should be forwarded with a copy of the first page of the article to the Copyright Clearance Center, Inc., 21 Congress Street, Salem, MA 01970, U.S.A. If no code appears in an article, the author has not given broad consent to copy and permission to copy must be obtained directly from the author. All articles published prior to 1980 may be copied for a per-copy fee of US\$ 2.25, also payable through the Center. This consent does not extend to other kinds of copying, such as for general distribution, resale, advertising and promotional purposes, or for creating new collective works.

Special written permission must be obtained from the publisher for such copying.

Special regulations for authors in the U.S.A. Upon acceptance of an article by the journal, the author(s) will be asked to transfer copyright of the article to the publisher. This transfer will ensure the widest possible dissemination of information under the U.S. Copyright Law.

Printed in The Netherlands

CONTENTS

(Abstracts/Contents Lists published in Analytical Abstracts, Biochemical Abstracts, Biological Abstracts, Chemical Abstracts, Chemical Titles, Current Contents/Physical, Chemical & Earth Sciences, Current Contents/Life Sciences, Index Medicus, Mass Spectrometry Bulletin and Science Citation Index)

Applications of two-dimensional thin-layer chromatography

by M. Zakaria, M.-F. Gonnord and G. Guiochon (Palaiseau, France) 127

Statistical approach to chromatographic theory

by J. S. Fritz (Ames, IA, U.S.A.) and D. M. Scott (Stanford, CA, U.S.A.) 193


Chromatographic Data

Table of isotachophoretic indices. I. Simulated qualitative and quantitative indices of 287 anionic substances in the range pH 3-10

by T. Hirokawa, M. Nishino, N. Aoki and Y. Kiso (Higashi-hiroshima, Japan) and Y. Sawamoto, T. Yagi and J.-I. Akiyama (Kyoto, Japan) D1

VYDAC

COMPLEMENTARY REVERSE PHASE MATERIALS NO RESEARCH LAB CAN AFFORD TO BE WITHOUT.



SEM photomicrograph
of VYDAC HS silica.

VYDAC TP 300 Angstroms and/or
Spherical Silica based reverse phase

Applications


Polyaromatic hydrocarbons
Vitamins D₂ and D₃
Large Peptides
Proteins
Anti-oxidants in plastics
Vitamin A Isomers

VYDAC HS 80 Angstroms
Spherical Silica based reverse phase

Applications

Nucleosides, purines, pyrimidines
Catecholamines
Small Peptides
Vitamin Bs and C
Organic Acids

Improved PAH Column




Vitamin Ds




Alpha and Beta Carotenes



Vitamin Bs



Organic Acids



Only the Separations Group offers C₁₈ reverse phase materials bonded to distinctly different spherical silica substrates yielding uniquely different selectivities for the HPLC chromatographer.

For your free copy of our "Comprehensive Guide to Reverse Phase Materials for HPLC," contact the Separations Group. Do not hesitate to call collect.

THE SEPARATIONS GROUP

MANUFACTURERS OF VYDAC

Box 867 • 16695 Spruce St. • Hesperia, CA 92345 USA
Telephone (619) 244-6107

SPHERICAL SILICAS and separation materials for high performance liquid chromatography since 1971



*use
this card
for more
information
on
advertisements
and news
items*

Reader service card

JOURNAL OF

CHROMATOGRAPHY

INTERNATIONAL JOURNAL ON CHROMATOGRAPHY, ELECTROPHORESIS AND RELATED METHODS

I would like to receive, without any obligation, further information on

News items: _____

Advertisements: _____

I am especially interested in the fields as indicated below:

- | | |
|--|--|
| ☐ 6040 Analytical Chemistry | ☐ 6240 Pharmaceutical Chemistry |
| ☐ 6050 Chromatography | ☐ 6241 Medicinal Chemistry |
| ☐ 6150 Mass Spectrometry | ☐ 6242 Clinical Chemistry |
| ☐ 6140 Spectroscopy | ☐ 6260 Environmental Chemistry |

Name: _____

Position: _____

Address: _____

please use
envelope;
mail as
printed
matter

ELSEVIER SCIENCE PUBLISHERS

Advertising department

P.O. BOX 211,

1000 AE AMSTERDAM – THE NETHERLANDS



please use
envelope;
mail as
printed
matter

ELSEVIER SCIENCE PUBLISHERS

Advertising department

P.O. BOX 211,

1000 AE AMSTERDAM – THE NETHERLANDS



*use
this card
for more
information
on
advertisements
and news
items*

Reader service card



I would like to receive, without any obligation, further information on

News items: _____

Advertisements: _____

I am especially interested in the fields as indicated below:


- | | |
|--|--|
| ☐ 6040 Analytical Chemistry | ☐ 6240 Pharmaceutical Chemistry |
| ☐ 6050 Chromatography | ☐ 6241 Medicinal Chemistry |
| ☐ 6150 Mass Spectrometry | ☐ 6242 Clinical Chemistry |
| ☐ 6140 Spectroscopy | ☐ 6260 Environmental Chemistry |

Name: _____

Position: _____

Address: _____

YOURS FOR THE ASKING...



Write now for your copy!

**Elsevier
Science Publishers**

P.O. Box 330
1000 AH Amsterdam
The Netherlands
Tel: (20) 5803-449

52 Vanderbilt Avenue
New York, NY 10017
U.S.A.
Tel: (212) 867-9040

7239

Reagents

MERCK

Hibar LiChroCART® HPLC Cartridge System



**Innovation
from
Experience**

Hibar LiChroCART®

- HPLC cartridges
- manu-fix cartridge holder
- auto-fix clamping system

simpler · better · faster · cheaper · neater

Please ask for our special leaflet!

CHREV. 167

APPLICATIONS OF TWO-DIMENSIONAL THIN-LAYER CHROMATOGRAPHY

MONA ZAKARIA, MARIE-FRANCE GONNORD and GEORGES GUIOCHON*

Laboratoire de Chimie Analytique Physique, Ecole Polytechnique, Route de Saclay, 91128, Palaiseau Cedex (France)

(First received January 5th, 1983; revised manuscript received January 25th, 1983)

CONTENTS

1. Introduction	128
2. Stationary phases and retention mechanisms	129
2.1. Inorganic adsorbents	130
2.2. Organic adsorbents	131
2.2.1. Polyamide	131
2.2.2. Cellulose	132
2.3. Reversed phases	132
2.4. Ion exchangers	133
2.5. Thin-layer gel chromatography phases	133
2.6. Stationary phases for two-dimensional thin-layer chromatography	133
3. Detection	134
4. Applications	137
4.1. Lipids	137
4.1.1. Fatty acids and their esters	139
4.1.2. Steroids	139
4.1.3. Phospholipids	141
4.1.4. Glycolipids	144
4.2. Pigments	149
4.3. Alkaloids	153
4.4. Proteins and their constituents	160
4.4.1. Amino acids	160
4.4.2. Dimethylaminonaphthalenesulphonyl (Dns) and dinitrophenyl (DNP) amino acids	163
4.4.3. Peptides and oligopeptides	165
4.4.4. Proteins	165
4.5. Carbohydrates	167
4.6. Glycopeptides	167
4.7. Nucleic acids and their constituents	169
4.7.1. Bases, nucleosides and nucleotides	169
4.7.2. Oligonucleotides and nucleic acids	172
4.8. Environmental pollutants	175
4.9. Pesticides	177
4.10. Inorganic compounds	180
4.11. Miscellaneous	181
5. Conclusion	187
6. Summary	188
References	188

1. INTRODUCTION

Bi-directional chromatography, which is a more specific term than two-dimensional chromatography when referring to flat paper or thin-layer beds¹, is an analytical technique that is becoming increasingly important in the separation of mixtures of varying complexity. This method consists in spotting a sample at the corner of a flat bed and allowing the migration of a solvent in one direction, followed by another migration at right-angles to the first.

When the same solvent is fed successively in the two directions of a uniform adsorbent layer, all chromatographed spots align on a diagonal. The sole advantage of the method is then a very slight increase in resolution corresponding to an increase, by a factor of $\sqrt{2}$, in the distance of migration of the spot. The realization of more efficient systems, by means of which a larger number of spots can be resolved², implies the use of the entire chromatographic area available and requires the selection of appropriate different mobile phases for each direction.

Similar efficiencies are attained when the same mobile phase is fed successively, at an orthogonal angle, to a chromatographic plate that has been coated with two adsorbents. Different retention mechanisms dictate the behaviour of the solutes chromatographed for instance, on a reversed-phase layer in one direction and a normal-phase layer in the other. The solute R_F values are then different in each direction and the spots are dispersed on the chromatographic plate.

The tremendous separation potential afforded by this technique is apparent from the flexibility available in combining various stationary and mobile phases for use in either direction.

In 1944, Consden, Gordon and Martin³ first reported a bi-dimensional separation of amino acids on paper. The eluents used were collidine-water in one direction and phenol-water in the second. The chromatography took 5 days and resulted in the resolution of 15 of the 22 amino acids in the sample mixture. This technique was subsequently described by Overell⁴, Cheftel *et al.*⁵ and Nordmann *et al.*^{6,7}, who successfully achieved the separation of various organic acids.

The importance of this method increased in the early 1950s as bi-directional chromatography became instrumental in the elucidation of several biochemical pathways⁸. Calvin^{9,10}, for instance, used two-dimensional paper chromatography in combination with autoradiography to assay algal extracts kept in air containing $^{14}\text{CO}_2$. After the identification of the various sample components, a kinetic study of ^{14}C incorporation helped to suggest a scheme for carbon reduction during photosynthesis. The same technique enabled Beloff-Chain *et al.*^{11,12} to differentiate between the conversion of glucose into oligosaccharides, polysaccharides and lactate in muscles and that into amino acids and lactate in the brain. Chain *et al.*^{13,14} also studied the effect of insulin on glucose metabolism in various types of animal tissue such as the diaphragm, liver, brown adipose tissue, brain and the pituitary.

Although two-dimensional thin-layer chromatography (2-D TLC) has been used for almost 40 years (Fig. 1), there has been no review of its various applications prior to this paper. Several workers have reported past and present developments of classical TLC and its future trends¹⁵⁻²⁰. The performance and spot capacity of two-dimensional TLC have been described elsewhere²¹, and also recent developments in TLC instrumentation²².




Fig. 1. Histogram showing the number of original two-dimensional TLC publications in the last 18 years.

This paper briefly reviews the different types of stationary phases available for chromatography, and the effective mechanisms of solute retention on each phase, prior to discussing the various applications of the 2-D technique. A short discussion on the methods available for spot detection is also included.

We have tried to cover the literature as thoroughly as possible. Authors who have not been cited are kindly asked not to take offence but rather to inform us of their work. More importantly, this review aims to demonstrate the efficiency of the method in resolving compounds of widely different nature, polarity or class, and the importance of the judicious selection of chromatographic conditions such that different mechanisms may govern solute retention in each direction.

2. STATIONARY PHASES AND RETENTION MECHANISMS

Various powders are commercially available that may be slurried alone or admixed with agents for specific applications, and spread on glass, aluminium foil or plastic supports. Pre-coated TLC plates are also available without or with a binder for increased adhesion and mechanical stability of the layer, or an indicator for solute detection.

The thickness of the layer is often 0.1–0.3 mm for analytical applications and 0.5–2.0 mm for preparative work. The particle size varies, a small diameter being less critical than a narrow size distribution in achieving high efficiencies and good resolution²: the narrow size range, 5 μm "HPTLC" packings have disadvantages such as a slow flow-rate of the mobile phase, a long development time and spot diffusion, whereas packings with the same size distribution and a larger diameter (10–20 μm) afford faster and still adequate separations^{2,23}, as is obvious from results obtained with good but conventional TLC plates made with particles *ca.* 11 μm in diameter²⁴.

The type of stationary phase needed to resolve a given mixture depends, to a large extent, on the properties of the sample components. In general, stationary phases may be inorganic sorbents (silica, alumina, magnesia), organic sorbents (cellulose, polyamide), ion exchangers (ionic moieties substituted on to an organic adsorbent) and reversed phases (non-polar, organic groups chemically bonded often to an in-

organic silica backbone). These thin layers may be impregnated with a specific reagent to enhance or create a particular type of solute-stationary phase interaction, or coated side-by-side on a support, a practice often used in bi-directional TLC.

2.1. Inorganic adsorbents

Inorganic oxides such as silica, alumina and magnesia are polar adsorbents which, according to their degree of activation, may be used to separate polar or moderately polar compounds. Procedures for adsorbent activation by heat or deactivation by addition of a moderator such as water have been standardized to ensure reproducibility of separations.

Heat activation rids TLC plates of physically adsorbed water, thus freeing the adsorptive surface for interactions with various solutes. Adsorptive sites in silica and magnesia are essentially hydroxyl groups. Thus, heating these two adsorbents to temperatures above 200°C for silica and 100°C for magnesia, a process whereby Si-OH and Mg-OH groups are converted into Si-O-Si and Mg-O-Mg, respectively, results in the loss of chromatographic activity²⁵. In alumina, active adsorptive sites are thought to be aluminium atoms, strained Al-O bonds, basic oxide ions and cationic centres. The loss of most of the surface hydroxyl groups early in the heating process does not reduce the activity of alumina, which actually increases with increasing temperature up to about 800°C and is lost only at 1000°C²⁵.

In chromatographic systems that involve weak or moderately strong solvents, adsorption is described primarily by the following relationship²⁵:

$$\log K = \log V_a + \alpha (S^0 - A_s \epsilon^0) \quad (1)$$

where K , the distribution coefficient, is the ratio of the solute concentration in the adsorbed state to that in the mobile phase, V_a is the volume of a monolayer of solvent adsorbed on the surface, α , the activity of the adsorbent, is proportional to its average surface energy and related to its water content, S^0 is the adsorption energy of the solute in the reference solvent and on an adsorbent of standard activity ($\alpha = 1$), A_s is the molecular area required for the adsorption of a solute molecule on the adsorbent surface and ϵ^0 is the solvent strength parameter²⁵.

Hence, the extent of retention by adsorption on a given phase increases primarily with the activity, average surface energy and surface area of the adsorbent. In addition, retention increases with S^0 , the adsorption energy of the solute molecule, which is approximately equal to the sum of the individual adsorption energies contributed by each constituent group. Certain functional groups interact more strongly with polar adsorbents than others. The retention, for the corresponding solutes, increases in the order saturated hydrocarbons < aromatic hydrocarbons $\approx$ halogen derivatives < ethers < esters $\approx$ aldehydes $\approx$ ketones < amines $\approx$ alcohols < carboxylic acids.

Stronger solvents (larger ϵ^0) decrease the solute retention. This effect is more important for larger solute molecules (large A_s , eqn. 1). The polarity of the mobile phase needed to achieve adequate R_F values is determined by that of the solutes studied. Polar compounds often require the presence of water in the mobile phase to ensure the *in situ* deactivation of the plate and prevent the occurrence of tailing or

irreversible chemisorption caused by strong adsorbent-solute interactions. In addition, desorption is ensured by the favoured interactions of the solute with the polar eluent. However, when the amount of water in the eluent exceeds 16%, solute partition is effective to a certain extent²⁶.

Other factors that affect adsorptivity are the acidic or basic functionalities of the adsorbent, as well as steric factors. Each adsorbent is characterized by a definite geometrical arrangement of adsorptive sites, favouring interaction with certain types of solutes. For instance, alumina is better suited than silica for the separation of aromatic hydrocarbons with different functional groups or steric arrangement. Moreover, although the adsorptive properties of alumina are similar to those of magnesia, the latter adsorbent has a much higher affinity for carbon-carbon double bonds and is especially suitable for the separation of planar aromatic hydrocarbons.

In general, solute interactions with a given adsorbent may involve London dispersion forces, orientation or induction dipole forces, hydrogen bonding and weak covalent bonding (acid-base interactions and complex formation). The extent to which any of these occurs depends on the particular chromatographic system used. Adsorbents may often be impregnated with specific reagents that may complex reversibly with the solute or covalently bond to it. Impregnation with boric acid, for instance, is used for the separation of vicinal dihydroxy isomers which are retained by forming cyclic boric acid derivatives. Argentation chromatography or impregnation with silver nitrate allows the separation of unsaturated compounds owing to rapid and reversible complexation of their π double bond electrons with the impregnated silver ions. Similarly, the separation of nitrogenous bases on plates impregnated with Zn, Cd or Mg involves the occurrence of selective interactions between the metal ion used and the free nitrogen electrons¹⁹.

2.2. Organic sorbents

In addition to the aforementioned inorganic sorbents, various organic macromolecular stationary phases are also commercially available. Of these, polyamide and cellulose are the most commonly used, and are particularly suitable for the separation of hydrophilic, strongly polar substances.

2.2.1. Polyamide

With aqueous or polar eluents, this sorbent behaves as a reversed phase, whereupon partition is thought to be involved, at least to some extent, in solute retention. With non-polar developers, polyamide is a normal phase on to which chromatographed substances may adsorb. This adsorption may entail the function of hydrogen bonds between proton-donating groups such as hydroxyl, amino, carboxylic or phosphoric in the solute molecule and carbonyl oxygens in the sorbent. Additional interactions may occur between electrophilic nitro compounds, quinones or aldehydes and amino groups in the polyamide phase. Compounds with delocalized π electron systems are also found to exhibit strong affinity for the sorbent. Clearly, chromatography on polyamide is a process of simultaneous e^- donor- e^- acceptor interactions, the extent of which depends on the chemical nature of the solute, the composition of the desorbing eluent and the presence of free $-NH-$ and $>C=O$ groups on the adsorbent surface²⁶.

2.2.2. Cellulose

In paper chromatography, one often witnesses rapid spreading of substances along cellulose fibres. The powdered sorbent used in TLC, however, is characterized by very short fibres which afford sharper spots and a better resolution. TLC cellulose exists in two forms: native fibrous (fibre length 2–25 μm , average degree of polymerization 400–500) and microcrystalline (particle size 20–40 μm , average degree of polymerization 40–200). In cellulose, most glucose units constitute crystalline regions, joined together as fibrils. These are bound into fibres by amorphous β -glucoside linkages. Whereas the crystalline surface is impermeable to most solvents, the amorphous regions hold imbibed water. Both structural types are responsible for solute retention, the former involving adsorption and the latter partition.

As partition involves the distribution of a solute between two immiscible liquids, it may be surprising at first to evoke this mechanism when solvents miscible with water, if not water itself, are used as developers. However, the water present in the amorphous regions has different properties to bulk water: it is linked by hydrogen bonds to the cellulose, forming a "liquid gel" with high apparent density and low velocity of diffusion.

The partition coefficient, a measure of solute retention, is determined by the extent of solute incorporation in the gel and interaction, by means of hydrogen bonds, with the "liquid gel" water. Molecular size and shape and the position and nature of hydrophilic groups in the chromatographed substance are important in this process²⁶.

Partition is not, however, solely responsible for chromatographic behaviour, as the migration sequence of several solutes differs from that based on partition coefficients alone. Adsorption and, to some extent, ion exchange are thought to participate in the chromatographic process.

2.3. Reversed phases

TLC layers for reversed-phase applications were first obtained by impregnating polar plates with organic solvents such as paraffin²⁷. Normal phases were later chemically modified, *e.g.*, by the *in situ* acetylation of cellulose or silanization of silica plates. The resulting phases could withstand development in a wide variety of solvents, and ensured reproducibility of separations^{28–30}. The number of commercially available reversed-phase plates increased subsequent to the more recent development of non-polar powders, in which different aliphatic chains are chemically bonded to a silica backbone. Halpaap *et al.*³¹ tested such phases with respect to their chemical and physico-chemical characteristics, wettability and chromatographic retention data.

The mechanisms of solute retention on these various packings are diverse: partition is responsible for chromatography on paraffin-coated silica layers; with acetylated cellulose, although adsorption cannot be excluded, solutes are thought to partition, when aqueous eluents are used, between the mobile phase and the water-impregnated cellulose fibres; adsorption due mainly to solvophobic forces is responsible for solute retention on silanized or alkyl-chain bonded silica phases. However, when water is used in large amounts, especially with small-chain reversed phases, mixed-mode mechanisms are believed to occur.

2.4. Ion exchangers

Polystyrene sulphonic (Dowex 50) or amine (Dowex 1) resins have long been used in TLC work. However, polyacrylamide gels or carbohydrate (cellulose and dextran) polymers, whether chemically modified or impregnated with liquid ion exchangers, are more suitable for the separation of bulky solutes, such as proteins and nucleic acids. Compared with resin exchangers, these matrices have larger pores which allow a greater capacity and a better transfer of macromolecules. In addition, the greater spacing between the active sites helps to achieve selective desorption under milder conditions than with resin phases.

The moderately strong polyethyleneimine (PEI) and weak diethylaminoethyl (DEAE) cellulose basic anion exchangers are commonly used in TLC applications, as well as the strong sulphoethyl (SE) and weak carboxymethyl (CM) Sephadex (polydextran) acidic cation exchangers. Inorganic cation exchangers such as zirconium(IV) phosphate and ammonium molybdate, are also commercially available.

The affinity of a compound for an ion exchanger is known to be proportional to its charge. Besides the electrostatic forces which contribute mainly to chromatographic retention in ion exchange, additional interactions are often operative as in the physical adsorption of hydrophobic solute moieties on the skeleton of the exchanger. With weakly acidic and weakly basic phases, the pH of the eluent may affect selectivity as the exchanging groups ion pair with H^+ or OH^- ions in the solution³².

2.5. Thin-layer gel chromatography phases

Dextran gels such as Sephadex G-10, admixed with silica gel or cellulose, have been used as TLC layers for the separation of purine and pyrimidine nucleotides and bases, for instance. While the character of the separations remains that of Sephadex, the layer has mechanical properties similar to those of silica gel or cellulose. Polyacrylamide gels (Biogel P) have also been used in TLC³³.

2.6. Stationary phases for two-dimensional thin-layer chromatography

Different sorbents may be slurried together and spread homogeneously on a plate for two-dimensional TLC purposes. The composition of the eluents is chosen such that, in each direction, a different sorbent interacts primarily with the solutes. Sorbents have also been coated side-by-side, forming coupled, two-phase plates; chromatography is then performed on a different phase in each direction.

The development of a plate in a given solvent is well known to affect the solute R_F values obtained from a subsequent development in another direction (Fig. 2). In addition, if the second solvent is very different from the first, its ascent in the orthogonal direction may be irregular, leading to distorted spots. In order to correct for those problems, Randerath and Randerath³⁵, after developing a PEI-cellulose plate in a buffer solution, washed away the residual salts by immersing the plate in methanol prior to a second elution. Bond³⁶ chose to transfer the spots obtained from the first development of a cellulose sheet on to another plate by clamping both thin layers face to face and developing with the second solvent, in a direction perpendicular to the first elution. The resulting chromatogram showed no interferences. In addition, the solute R_F values were unaffected by the previous run.




Fig. 2. (a) Theoretical and (b) experimental two-dimensional chromatograms of dinitrophenyl (DNP) amino acids on RP-18 plates. Eluents: in the first direction, hexane-ethyl acetate-acetic acid (80:18:2); in the second direction, 1 M ammonia + 3% potassium chloride in 60% methanol. S.P. = starting point. 1 = DNP-Gly; 2 = DNP-Ala; 3 = DNP-Ser; 4 = DNP-Thr; 5 = DNP-Val; 6 = DNP-Leu; 7 = DNP-Ile; 8 = DNP-Pro; 9 = DNP-Met- O_2 ; 10 = DNP-Trp; 11 = DNP-Phe; 12 = Di-DNP-Tyr; 13 = DNP-Asp; 14 = DNP-Glu; 15 = DNP-CySO₃Na; 16 = Di-DNP-Lys; 17 = α -N-DNP-Arg; 18 = Di-DNP-His; 19 = DNP-OH; 20 = DNP-NH₂. Reproduced from ref. 34 with permission.

3. DETECTION

The localization, characterization and identification of spots are essential aspects of TLC. The techniques used in two-dimensional TLC are identical with those used in one-dimensional TLC, except that the whole surface of the plate has to be scanned, which makes the problems much more difficult. Detection may therefore require a very long time and very sophisticated instrumentation, contrasting with the great simplicity of the separation method. Techniques for recovering or displaying the two-dimensional separation have benefitted from recent improvements in image processing made possible through the availability of powerful, yet cheap, computers.

In two-dimensional chromatography the exact position of the spot must be determined for qualitative analysis and the whole spot profile, in both directions, must be integrated for quantitative analysis. The problem is therefore much more difficult than in conventional TLC, where the centre of the spot is on the vertical of the sample spot and the solute concentration profile in the direction perpendicular to that of the development is devoid of interest and can be integrated at once, as is done with the use of a narrow slit perpendicular to the scanning direction in photodensitometers for TLC. Such devices cannot be used in two-dimensional TLC; the spatial resolution of the scanner, whether an optical beam, an electronic beam or any other system, must be small compared with the spot size, about 10–20 times smaller in dimension than the spot base width. Typically, this means that it should not exceed 0.1–0.5 mm in diameter, depending on the TLC conditions. If ω is the width of the sensor beam and L the development length, a number L/ω of parallel scans must be carried out. This number varies from about 100 to around 2000 depending on the conditions, but in any case is large. Convenient software must be available to align these scans, detect the spots and segment and integrate them.

In the following we review the various detection methods used in conventional TLC and discuss how they can be applied to the special problem of two-dimensional TLC.

A laborious detection technique consists in scraping off the parts of the adsorbent layer containing each spot of interest and extraction from the adsorbent or *in situ* elution of the spot. Although some commercial equipment is now available (for example, Eluchrom Automatic Elution System from Camag), which considerably reduces the amount of solvent necessary for recovering the sample, this technique is inconvenient, time consuming and not quantitative. A final measurement step must also be added, such as a spectrophotometric determination. To apply this method the spot must first be located precisely.

One-dimensional TLC has sometimes been coupled with gas detection devices: detection of compounds eluted on a Chromarod can be performed either with a flame-ionization detector³⁷ or by direct insertion into a thermoionic³⁸ or mass spectrometric³⁹ detector. However, these techniques are not suitable for two-dimensional TLC because of their lack of spatial resolution and/or the considerable amount of time necessary for the scans.

Spots may be revealed after performing adequate chemical tests. Reagents for these tests are often sprayed on the layer. The non-uniformity of the spray may drastically affect the detection of the linearity of response. Therefore, whenever possible spots are rendered visible by exposure to gaseous reagents, which act more uniformly.

Coloured or fluorescent spots can be easily detected visually under white or UV light. If the compound itself is not visible or fluorescent, detection can be performed by applying gaseous, colorimetric complexing agents or even radioactive markers, or by using the quenching effect of the compound on the fluorescence of an additive in the chromatographic bed. The compound then appears as a black spot on a brilliant background. This method permits the accurate determination of the position of the spot, but not of its size or concentration. The latter are determined by the use of one of the photometric methods of TLC classified as densitometric and fluorimetric techniques.

Densitometry can be carried out in the transmission or the reflection mode. It measures the intensity of the light transmitted through the TLC plate or reflected by it. Quantitation is based on the decrease in light transmitted or reflected by the spot in comparison with the neighbouring background⁴⁰. It requires background suppression and assumes a relationship between the signal and the concentration of a compound. This relationship is linear only at very low concentrations⁴¹. For higher concentrations it is not given by the conventional Beer's law but by the Kubelka-Munk equation⁴⁰, because of diffusion of light by the particles in the layer. The response factor of a compound depends on its coefficients of light adsorption and scattering⁴⁰. Obviously the densitometric measurements must be made using a monochromatic incident beam, and the wavelength must be optimized depending on the compounds of interest. Transmission densitometry has been shown theoretically⁴² to be more sensitive than reflectometry, although layer uniformity is more critical in the former method. Both non-uniformity of the layer and poor light intensity affect the signal-to-noise ratio. However, such problems may be overcome to some extent with a double-beam instrument. Reflectometry is more widely used because of the higher

optical density of most compounds, especially in the UV range. Frei⁴⁰ published an extensive study on reflectance spectroscopy. Using double-beam instruments, one can also compensate for the background effects due to the layer.

In fluorimetry, the incident radiation excites the various compounds, and those which emit secondary radiation are detected. This emitted light is characterized by a longer wavelength than the incident radiation. The non-spotted, background or base regions of the layer do not fluoresce. The excitation wavelength can be adjusted to optimize the sensitivity of the measurement for a given compound. The most commonly used wavelengths are 254 and 360 nm.

Fluorescence measurements can also be carried out in the so-called quenching mode. A phosphor additive is mixed to the sorbent slurry before coating the plate. When the developed plate is irradiated with UV light the base regions of the plate emit a uniform visible light, while the sample compounds appear as dark spots. This method is more sensitive when the light is transmitted from the back of the plate.

Many commercially available instruments are single-wavelength devices in which a monochromatic beam is used to scan the plate surface. They can be applied to all modes of densitometry or fluorimetry, with the use of suitable mirrors and filters. In the fluorescence mode, better instruments permit the adjustment of both the excitation and emission wavelengths. In these instruments the scanning optical beam is supposed to illuminate on the plate a rectangle with a very large aspect ratio, the long dimension (*ca.* 1 cm) being perpendicular to the scanning direction. This yields a signal proportional to the light intensity perceived, *i.e.*, integrated over the whole slit length. If the relationship between light intensity and spot concentration is non-linear, this results in quantitative errors when the concentration is not constant along the slit. On the other hand, this permits integration of the fluctuations due to layer heterogeneity over a larger area and decreasing the noise. Hence, the signal should be integrated over the slit to linearize the characteristic response of a single slit in terms of concentration⁴³. Knapstein *et al.*⁴⁴ obtained a linear calibration graph for steroids detected on silica gel in the quenching mode.

With dual-wavelength densitometers, fluctuation of the light source is balanced and a flying spot can replace a fixed slit aperture, which leads to greater accuracy in quantitation. The complexity of dual-wavelength densitometers requires computer-assisted processing of the signal⁴⁵. Speed, like resolution, is expensive and rapid evaluation of a one-dimensional thin-layer chromatogram requires the scanner to be connected either to a recorder or to a video screen. Speed can also be achieved at the expense of a lower sensitivity if the signal noise depends on the integration time.

In 1976, Devenyi and co-workers^{46,47} introduced video techniques in the densitometric evaluation of thin-layer chromatograms. The whole chromatogram is irradiated by a suitable light source of relatively homogeneous intensity and an image is taken by a Vidicon camera. This is extremely fast and also reasonably accurate and sensitive, and the rather sophisticated processing such as background subtraction or spot integration is easy to carry out in an interactive mode.

Generally, the HPTLC scanners available for photodensitometric measurements cannot be adapted easily to two-dimensional TLC. The speed of detection is an extremely critical factor as all of the plate surface has to be scanned. Scanning a conventional TLC plate takes a few minutes, but scanning a two-dimensional plate would take from several hours to one day. No software is available to align all these scans and process them to obtain quantitative information.

Two-dimensional detection devices have resulted mainly from two-dimensional gel electrophoretic analyses^{48,49} and from the TLC of radioactive compounds⁸. In both instances an autoradiograph is obtained and then scanned with a very narrow visible light beam (spatial resolution around 0.1 mm or below). Complete data acquisition and processing systems have been developed by several groups⁴⁸⁻⁵⁰. These systems offer possibilities of data acquisition, filtering, background subtraction, calibration and data reduction, and provide for spot detection, segmentation, integration, pairing and data file handling on thousands of spots.

Video-densitometric techniques have been applied to two-dimensional TLC. A two-dimensional matrix of pixels has sometimes been used^{47,51} together with a Vidicon camera to obtain both normal two-dimensional and spectral images. Both fluorescence and absorption measurements can be carried out with this technique. The sensitivity of this type of detector is in the nanomole range. Although its main drawbacks are cost and complexity, it is still more attractive to the analyst than the classical spectrodensitometer, the use of which is tedious and time consuming and necessitates image reconstitution after hundreds of one-dimensional scans of a two-dimensional plate.

Recently, Guiochon and co-workers^{52,53} conceived the use of a detection device on-line with a two-dimensional liquid chromatographic system. Detection is now performed in the elution mode, in the second direction, with a photodiode array Reticon camera used as a series of individual UV cells parallel to the exit edge of the plate. In this system, the solvent is pumped through the chromatographic bed under pressure and no longer moves under the influence of capillary forces. At the exit, the solvent is forced through a slot which constitutes the detector cell.

A monochromatic light passes through this slot and is focused on the diode array of the Reticon camera. A time-dependent signal similar to the line signal of an autoradiography scan is recorded and processed through a data acquisition system, then displayed on a colour TV screen. HPLC elution conditions may thus be reproduced and quantitation is no longer affected by plate characteristics and chromatographic bed diffusivity as in regular TLC.

4. APPLICATIONS

The separations that have been accomplished using two-dimensional TLC are classified in this section under lipids, pigments, alkaloids, proteins and their constituents, carbohydrates, glycopeptides, nucleic acids and their constituents, environmental pollutants, pesticides and miscellaneous. We have tried to emphasize the peculiarities of various assays by analysing the chromatographic behaviour of solutes. The retention and chromatography of compounds may involve adsorption, partition, ion exchange or restricted diffusion (gel permeation). Although one particular chromatographic mechanism may be emphasized for the retention of solutes under given conditions, other concurrent processes should not be forgotten.

4.1. *Lipids*

"Lipids" is a term used to qualify various classes of compounds that are characterized in general by solubility in organic solvents and insolubility in water. In this

TABLE I

SEPARATION OF FATTY ACIDS AND THEIR ESTERS

N is the number of spots resolved in the best published chromatogram. Compounds: name of classes, groups or individual compounds studied. Stationary phase: composition of the thin layer used as the chromatographic bed. Also includes dimensions and activation conditions when available. When two different phases are used, 1 refers to the phase used for the first development, usually a narrow strip along the edge of the plate, and 2 to the second stationary phase. SSP: sample spot position; distance from the edges of the plate. Solvent 1: composition of the solvent used for the first development. Sometimes several successive developments are carried out using different solvents. Their compositions are given in the order a, b, c, etc. *t*/*L*: time (min) or length (cm) of the first (with solvent 1) and second (with solvent 2) developments. Drying: conditions under which the plate is dried after the first development is finished. Solvent 2: same as solvent 1, but for the second development. Ref.: also includes figure numbers, if one is reproduced here.

<i>N</i>	Compounds	Stationary phase	SSP (cm)	Solvent 1	<i>t</i> / <i>L</i>	Drying	Solvent 2	<i>t</i> / <i>L</i>	Ref.
	Alkenyl acyl-, alkyl acyl- + diacyl glycerol acetates	Si gel G, 110°C, 1 h		Skelly F-diethyl ether (88:12)	14 cm	N ₂ dry	Toluene		54
11	Fatty acids, prostaglandins, thromboxanes	Si gel, 20 × 20 cm		Diethyl ether-CH ₃ OH-acetic acid (90:1:2)			Chloroform-methanol-acetic acid-water (90:8:1:0.8)		55
13	Fatty acids esters	Si gel D.B., Camag, 20 × 20 cm, 0.25 mm, wash, 139-142°C, 30 min	2.5	Light petroleum-diethyl ether-acetic acid (75:25:2.5)	70-80 min, 16 cm	15 min, room temp.	Diethyl ether-light petroleum-acetic acid (70:30:1)	65-75 min, 15 cm	56
8	Ginsenosides	Silanized Si gel 60 F254, 20 × 20 cm, 0.25 mm	2.0	(a) Chloroform-methanol-ethyl acetate-butanol-water (4:4:8:1:2) (b) Chloroform-butanol-methanol-water (4:8:3:4)	10 cm	air dry, 10 min	(c) Chloroform-methanol-water (13:7:2)	10 cm	57
13	Mercury(II) adducts of unsat. esters (mono-, di-, tri-, tetra-, penta-, hexa-enes)	(1) Si gel G-Kieselguhr (3:7), 20 × 20 cm (2) Impregnate with 10% paraffin in light petroleum		On (1): isobutanol-formic acid-water (8:8:5)	Top	1 h air, 10 min H ₂ S chamber	On (2): formic acid-acetonitrile-acetone (8:8:5)		58
16	Trialkylglyceryl esters, dialkoxylglycerides, alkoxylglycerides + <i>n</i> -glycerides	(1) Si gel G, 4 × 20 cm, 0.25 mm (2) AgNO ₃ -impregnated Si gel G, 16 × 20 cm, 0.25 mm, 110-120°C, 1-2 h	2.0	On (1): petroleum hydrocarbon-diethyl ether (90:10) (need 50% relative humidity) Benzene-acetone (20:1)			On (2): petroleum hydrocarbon-diethyl ether (90:10) (2 × need 50% rel. humidity)		59 (Fig. 3)
4	Positional isomers of hydrostearic acid	Si gel G, 0.25 mm		Benzene-acetone (20:1)			Benzene-acetone (20:1)		60

section, four main groups of substances are considered: fatty acids and their esters, steroids, phospholipids and glycolipids. The different two-directional assays are briefly discussed in the text and detailed chromatographic conditions are given in Tables 1-5.

4.1.1. Fatty acids and their esters (Table 1)

Separations of various glyceryl acetates⁵⁴ and of some fatty acids from prostaglandins and thromboxanes⁵⁵ have been achieved on silica plates owing to the different selectivity of the solvent system in each direction. Fatty acid esters have also been resolved on silica⁵⁶ with a more polar eluent in the second than in the first direction.

A special category of compounds, the ginsenosides (triterpenoid saponins of the dammarane type), have been assayed on silanized silica gel using a less polar mobile phase in the second than in the first direction⁵⁷.

El-Zeany and Ahmed⁵⁸ devised an original method to resolve fatty acid esters. They first separated the mercury(II) adducts of these unsaturated esters on a 3:7 silica gel-Kieselguhr adsorbent with isobutanol-formic acid-water, based on the different degree of solute unsaturation. Then, the plate was positioned in a hydrogen sulphide chamber; mercury(II) sulphide was deposited as black spots and the esters were liberated. Chromatography in the second direction was accomplished after impregnating the plate with paraffin. The esters partitioned between that phase and the formic acid-acetonitrile-acetone eluent. Their mobility was a function of chain length. Whereas in this method mercury(II) adducts of solutes were chosen to achieve separations based on different degrees of unsaturation, more commonly unsaturated compounds are resolved according to their ability to form coordination complexes with ions (such as silver) which impregnate the stationary phase. Schmid *et al.*⁵⁹, for instance, after separating glycerol esters into four classes according to basic structure, type and number of functional groups on plain silica gel, differentiated each class according to the number and configuration of the double bonds in the constituent compounds, on silica gel impregnated with silver nitrate (Fig. 3).

4.1.2. Steroids (Table 2)

This class of lipids, of which cholesterol is a common representative, includes physiologically important compounds such as the bile acids, the androgens and the oestrogens, and essential metabolic regulators, such as the adrenocortical hormones. Thus, the separation of these substances and their quantitation in biological samples are essential, especially for diagnostic purposes.

Using silica, Bouillé *et al.*⁶¹ first eluted the impurities associated with cortisone in plasma and adrenal extracts. Chromatography in the orthogonal direction was achieved with a more polar mobile phase containing water, which resulted in the migration of corticosterone from its original position on the plate. Also using a silica phase, Bicknell and Gower⁶² succeeded in resolving a mixture of polar and non-polar compounds. Two successive developments in the first direction with benzene-diethyl ether (9:1) permitted the separation of the non-polar C₁₉ 16-unsaturated steroids 5 α -androst-16-en-3 β -ol, 5 β -androst-16-en-3 α -ol, 5 α -androst-16-en-3 α -ol, 1,3,5-[10],16-oestratetraen-3-ol and 5 α -androst-16-en-3-one. The use of a more polar eluent, benzene-methanol (9:1), for development in the second direction permitted




Fig. 3. Two-dimensional chromatogram of a model mixture containing the following compounds: (A) trialkyl glyceryl ethers: trioctadecyloxypropane, 1,2-dioctadecyloxy-3-octadecenyloxypropane, 1-octadecyloxy-2,3-dioctadecenyloxypropane, trioctadecenyloxypropane; (B) dialkoxylglycerides; 2,3-dioctadecyloxy-1-O-octadecanoylpropanol, 2,3-dioctadecyloxy-1-O-octadecenoylpropanol, 2,3-dioctadecenyloxy-1-O-octadecanoylpropanol; (C) alkoxydiglycerides; 3-octadecyloxy-1,2-di-O-octadecanoylpropanediol, 3-octadecenyloxy-1,2-di-O-octadecanoylpropanediol, 3-octadecyloxy-1,2-di-O-octadecenoylpropanediol, 3-octadecenyloxy-1,2-di-O-octadecenoylpropanediol; (D) triglycerides: tri-O-octadecanoylpropanetriol (tristearin), 1,2-di-O-octadecanoyl-3-O-octadecenoylpropanetriol (oleodistearin), 1-O-octadecanoyl-2,3-di-O-octadecenoylpropanetriol (stearodistearin), tri-O-octadecenoylpropanetriol (triolein). Solvents: first direction, petroleum hydrocarbon-diethyl ether (90:10), developed once on plain silica gel G strip; second direction, petroleum hydrocarbon-diethyl ether (90:10) developed twice on silver nitrate-impregnated silica gel G. Indicator: charring after spraying with chromic-sulphuric acid solution. Reproduced from ref. 59 with permission.

the separation of the more polar Δ^4 -3-oxosteroids testosterone and progesterone and of the Δ^5 -3 β -hydroxysteroids pregnenolone and 5,16-androstadien-3 β -ol.

Taylor⁶³ separated thirteen polar Δ^4 -3-oxosteroids and oestrogens on silica with chloroform-methanol-water (94:6:0.5) in the first direction, followed by cyclohexane-ethyl acetate (50:50) in the second direction (Fig. 4). Cavina *et al.*⁶⁴ achieved similar resolutions of oestrogens and progestagens on silica by combining appropriate different developing systems. Clearly, the polarity of the mobile phase, its constituents and the degree of deactivation of the stationary phase achieved by use of water in the eluent are all factors that affect such assays.

Issaq⁶⁵ resolved two isomeric pairs, namely 7 α - from 7 β -hydroxycholesterol and 5 β ,6 β - from 5 α ,6 β -epoxycholesterol on a special plate with two different sorbents coated side-by-side. Chromatography in the first direction was accomplished on silica gel with diethyl ether as the eluent. Orthogonal elution was then achieved for a few centimetres on silica and, for the remaining part, on silanized silica, with a non-polar heptane-HMDS (98:2) as the mobile phase.

Similarly, Levitt and Touchstone⁶⁶ separated six bile acids on a two-phase plate where a 3 × 20 cm reversed-phase strip had been coated at the side of a 17 × 20 cm silica layer. Reversed-phase chromatography was carried out on the thin strip with an acid-organic eluent mixture and resulted in the separation of the bile acids into three classes: cholic, deoxycholic and chenodeoxycholic. Orthogonal development with the mixed aqueous (acetic acid-water) and organic (chloroform-methanol) solvent on the silica phase further separated the taurine from the glycine conjugates in all three classes of compounds. It is important to remember that these bile acids have functionalities that interact strongly with silica unless the adsorbent is modified through bonding with alkyl chains (reversed-phase strip) or deactivated by the presence, in the mobile phase, of components such as acetic acid or water, which compete strongly with the solute for the available adsorptive sites.

Vidrine and Nicholas⁶⁷ resolved two different lipid classes, *viz.*, steroids and unsaturated aliphatic hydrocarbons (terpenoids), on a mixed thin-layer plate coated with a slurry of silica, silanized silica and calcium sulphate. The eluent was relatively non-polar in the first direction, and retention could be attributed to adsorption. In the second direction, however, a more polar eluent was used, containing 25% of water. Under these conditions, a partition-adsorption mechanism is probably in effect.

Finally, Curtius and Müller⁶⁸ achieved an original, truly two-dimensional GC-TLC separation of a hormone mixture. The first dimension involved a separation on a GC column with a split effluent, 10% directed to a flame-ionization detector and 90% spotted on a TLC silica plate which moved linearly with time. The development of this plate with chloroform-ethanol (90:10) represented the second dimension and improved the resolution obtained with GC alone (Fig. 5). Clearly, the second separation is not especially helpful with the twelve components shown here, which are sufficiently well resolved by the GC step for quantitative analysis. It is, however, of greater use with more complex mixtures that cannot be resolved by GC but are resolved by the two-dimensional separation scheme.

4.1.3. Phospholipids (Tables 3 and 4)

This lipid class includes the phosphoglycerides and sphingolipids, all important membrane constituents. Chromatography of these compounds is often accomplished on silica gel, in its pure form or impregnated with either boric acid, ammonium sulphate, magnesium acetate, magnesium silicate or magnesium hydroxycarbonate.

In general, separations of phosphoglycerides (Table 3) require development of the plate with a water-organic eluent containing acetic acid in one direction and a base such as ammonia solution in the other⁶⁹⁻⁷³. Thus, the mobility of the molecules in each direction is affected by the differing degree of ionization, *i.e.*, the extent of amine protonation and phosphate and/or carboxyl dissociation, as well as by the competition of the acid or base in the mobile phase for adsorptive sites on the silica plate. Moreover, Jain and Subrahmanyam⁷⁴ have reported that appropriate amounts of ammonium sulphate in the stationary phase and acetic acid in the eluent are important for obtaining spots with no tailing and good resolution.

It is interesting that, on a phase containing boric acid⁷⁵, phosphatidylinositol, which has vicinal *cis*-hydroxyl groups in its sugar moiety, is retained the longest (Fig. 6). This may exemplify the selective affinity of boric acid for *cis*-diol compounds.

TABLE 2
SEPARATION OF STEROIDS
Column headings as in Table I.

<i>N</i>	<i>Compounds</i>	<i>Stationary phase</i>	<i>SSP (cm)</i>	<i>Solvent 1</i>	<i>t/L</i>	<i>Drying</i>	<i>Solvent 2</i>	<i>t/L</i>	<i>Ref.</i>
	Corticosterone purification	Kieselgel GF 254 (Si), 20 × 20 cm, 0.4 mm	2.0	(a) <i>n</i> -Hexane-ethyl acetate (4:1) (b) Ethyl acetate- cyclohexane-toluene (10:10:1)	20-30 min, 10 cm 30 min 10 cm	room temp.	Ethyl-acetate- chloroform-water (90:10:1)	30 min, 10 cm	61
17	Δ^4 -3-Oxosteroids, Δ^5 - 3 β -hydroxysteroids, C ₁₉ -16-unsat. steroids	Kieselgel G (Merck), 20 × 20 cm, 0.5 mm	2.0	Benzene-diethyl ether (9:1) (2 ×)	16.8 cm		Benzene-methanol (9:1)	16.8 cm	62
13	Δ^4 -3-Oxosteroids, oestrogens	Kieselgel GF 254, 20 × 20 cm, 0.5 mm		Chloroform- methanol-water (94:6:0.5)	14.4 cm from spot		Cyclohexane-ethyl acetate (50:50) (2 ×)	18 cm	63 (Fig. 4)
6 or 8	Estrogens, progesta- gens	Si gel 60 F ₂₅₄ , 20 × 20 cm, 0.25 mm	2.0	Toluene-95% ethanol (90:10)	18 cm	Air dry, 20 min, dark	Butyl acetate-light petroleum acetic acid (70:30:1)	~ 18 cm	64

4	7 α -Hydroxycholesterol, 7 β -hydroxycholesterol, 5 β ,6 β -epoxycholesterol, 5 α ,6 α -epoxycholesterol	(1) 20 × 20 cm Si gel (2) 10 cm dipped in HMDS, 100°C, 2 h	(1) Diethyl ether	13 cm	Dry	(2) Heptane-HMDS (48:2)	~ 17 cm	65
6	Bile acids	2-D plate, 20 × 20 cm, CS Multi K (Whatman) (1) 3 cm KC 18 strip (2) remainder, Si gel	(1) Ethanol-0.01 M KH ₂ PO ₄ (pH 2.5) (1:1)		Dry	(2) Chloroform-methanol-acetic acid-water (75:25:5:5)		66
8	Cholesterol, lanosterol, dihydrolanosterol, oxidosqualene, geraniol, <i>trans</i> , <i>trans</i> -farnesol, <i>cis</i> , <i>trans</i> -farnesol	Si gel G-silanized Si gel H-CaSO ₄ · $\frac{1}{2}$ H ₂ O (50:43:7), 20 × 20 cm	Hexane-ethyl acetate (90:10)	45 min, 14 cm		<i>p</i> -Dioxane-water (75:25)	2 h	67
12	Male and female hormones	(1) GC column impregnated with 3% XE-60 (2) Kieselgel G	(1) N ₂ , 50 ml/min, 210 → 265°C		Spray plate with HCl, split trimethylsilyl-esters	(2) Dichloromethane-acetone (70:30) or chloroform-ethanol (90:10)		68 (Fig. 5)




Fig. 4. Two-dimensional thin-layer chromatogram of steroids on silica gel. Solvent systems: (1) chloroform-methanol-water (94:6:0.5); (2) cyclohexane-ethyl acetate (50:50). 1 = Oestrone [3-hydroxy-1,3,5(10)-estradien-17-one]; 2 = progesterone(4-pregnene-3,20-dione); 3 = oestradiol [1,3,5(10)-estratriene-3,17 β -diol]; 4 = androstenedione (4-androstene-3,17-dione); 5 = 17 α -hydroxyprogesterone (17 α -hydroxy-4-pregnene-3,20-dione); 6 = testosterone (17 β -hydroxy-4-androsten-3-one); 7 = deoxycorticosterone (21-hydroxy-4-pregnene-3,20-dione); 8 = 11-deoxycortisol (17 α ,21-dihydroxy-4-pregnene-3,20-dione); 9 = corticosterone (11 β ,21-dihydroxy-4-pregnene-3,20-dione); 10 = cortisone (17 α ,21-dihydroxy-4-pregnene-3,11,20-trione); 11 = aldosterone (18,11-hemiacetal of 11 β -21-dihydroxy-3,20-dioxo-4-pregnen-18-al); 12 = cortisol (11 β ,17 α ,21-trihydroxy-4-pregnene-3,20-dione); 13 = oestriol [1,3,5(10)-estratriene-3,16 α ,17 β -triol]. Reproduced from ref. 63 with permission.

Pollet *et al.*⁷⁷ achieved the separation of different lipid classes, namely cholesterol, cerebrosides, sulphatides, phospholipids and gangliosides, on plain silica gel by repetitive elution with chloroform-methanol solutions of varying polarity in orthogonal directions (Table 4). Radwan⁷⁸ and Ikawa and Goto⁷⁹ resolved mixtures of fatty acids, steroids, phospholipids and other compounds on silica with an organic-acidic phase in one direction and an organic-basic phase in the other. Hubmann⁸⁰ was able to separate these various substances using different eluents for successive developments in each direction, owing to the different selectivity of each organic modifier with respect to the various solutes (Table 4).

4.1.4. Glycolipids (Table 5)

Glycolipids, such as the cerebrosides and the more complex gangliosides, are sugar-containing membrane lipids. The separation of such compounds was performed on silica gel by Ohashi⁸¹, who used chloroform-methanol-aqueous ammonia to develop the plate in the first direction and *n*-propanol-aqueous ammonia in the second. Ledeen *et al.*⁸², and subsequently Hunter *et al.*⁸³, in assaying for brain gangliosides, modified these conditions to a certain extent: ammonia was included in one dimension only, while salt was added to either or both solvents.




Fig. 5. Two-dimensional chromatography of twelve steroids (trimethylsilyl derivatives) using GC and TLC in combination. The lower part of the figure represents the gas chromatogram and the upper part illustrates the TLC separation after plate elution. It may be noticed that solutes 1, 2 and 3 are completely resolved after the TLC step. 1 = Allopregnandiol; 2 = pregnandiol; 3 = androsterone; 4 = etiocholanolone; 5 = dehydroepiandrosterone; 6 = pregnanolone; 7 = pregrantriol; 8 = 11-ketoandrosterone; 9 = 11-ketoetiocholanolone; 10 = 11-hydroxyandrosterone; 11 = 11-hydroxyetiocholanolone; 12 = pregnantriolone. St = starting point. Reproduced from ref. 68 with permission.

Cardenolides, which are glycosides with a steroid moiety, constitute a different category of sugar-containing lipids. Thirty-two cardenolides were separated by Clarke and Cobbs⁸⁴ on silica with a different solvent system for each direction. A correlation between the chromatographic mobility of the compounds and their chemical structure was observed. The number and position of the hydroxyl substituents played an important role in solute retention (Fig. 7).

TABLE 3
SEPARATION OF PHOSPHOLIPIDS
Column headings as in Table 1.

<i>N</i>	<i>Compounds</i>	<i>Stationary phase</i>	<i>SSP</i> (<i>cm</i>)	<i>Solvent 1</i>	<i>t/L</i>	<i>Drying</i>	<i>Solvent 2</i>	<i>t/L</i>	<i>Ref.</i>
6	Phosphatidyl (P) glycerol (G), pslerine (PS), P-ethanolamine (PE), pinositol (PI), leathin (L), sphingomyelin (S)	Si gel H with 5% ammonium sulphate, 20 × 20 cm		Chloroform-methanol-water-gl. acetic acid (65:25:4:8)	10 cm	70°C, 5 min	THF-methylol-methanol-2 M ammonia (40:28.5:7.8:4.2)	10 cm	69
6	PG, PE, PI, PS, L, S (in amniotic fluid)	Si gel H with 50 g/l ammonium sulphate, 20 × 20 cm, 110°C, 30 min		Chloroform-methanol-water-acetic acid, 17.4 M (162.5:65.0:10.0:0.5)	10 cm	70°C, 5 min	THF-methylol-methanol-2 M ammonia (120:80:35:10)	10 cm	70
13	PE, PC, PS, PI, L, S, and others, cerebro-sides and sulphatides (in brain and mitochondrial inner membrane)	Si gel H (20 g) with magnesium acetate (1.5 g), 0.25 mm, N ₂ , 50% humidity, 30 min		Chloroform-methanol-28% aq. ammonia (65:25:5)		Dry, 10 min, N ₂	Chloroform-acetone-methanol-acetic acid-water (3:4:1:1:0.5)		71

10	PI, PS, PE, PG, etc. (in erythrocytes)	Si gel with 1% alkaline magnesium silicate or 2% magnesium hydroxycarbonate	Chloroform-methanol-7 M ammonia (90:54:11)	Dry, 15 min; if rel. hum. > 65%, dry 1 h <i>in vacuo</i>	Chloroform-methanol-acetic acid-water (90:40:12:1)	72
11	PI, PS, PG, etc. (in yeast)	Si gel H, 20 × 20 cm, 0.25 mm	Chloroform-methanol-acetic acid-water (52:20:7:3)	60-90 min Air dry, 20 min, wash with diethyl ether, air dry, 20 min	Chloroform-methanol-40% aq. methylamine-water (13:7:1:1)	73
9	PG + derivatives, cardiolipin	Si gel G (25 g) with 0.4 M boric acid (50 ml), 20 × 20 cm, 0.25 mm, 120°C, 1 h	Chloroform-methanol-H ₂ O-conc. NH ₃ (70:30:3:2)	Dry, 30 min, N ₂	Chloroform-methanol-water (63:35:5)	75 (Fig. 6)
	L, S from PI, PS, PE, etc.	Si gel-impregnated glass-fibre sheets (ITLC type SG), 100°C, 30 min	Dichloromethane-ethanol-water (100:25:3)	6 cm, 3 min Air dry, 5 min	Chloroform-methanol-ammonia (170:20:3)	4 cm, 2 min 76

TABLE 4
SEPARATION OF PHOSPHOLIPIDS AND OTHER LIPID CLASSES
Column headings as in Table 1.

<i>N</i>	<i>Compound</i>	<i>Stationary phase</i>	<i>SSP (cm)</i>	<i>Solvent 1*</i>	<i>t_f/L</i>	<i>Drying</i>	<i>Solvent 2</i>	<i>t_f/L</i>	<i>Ref.</i>
11	Phospholipids, cerebrosides, sulphatide (plasma, liver, brain)	Si gel impregnated with ammonium sulphate, 20 × 20 cm; 0.5 mm, 110°C, 1 h		Chloroform-methanol-acetone-water (60:25:8:4)	Top	50°C, 5 min	Chloroform-methanol-acetic acid-water (50:35:8:1)	Top	74
11	Cholesterol, cerebrosides, gangliosides, phospholipids (cellular membranes)	Si gel HPTLC 60 F 254 (Merck), 10 × 10 cm	2	(a) Chloroform-methanol-water (70:30:4) (d) Chloroform-methanol (2:1)	Top	Bench dryer, few min Bench dryer, few min	(b) Chloroform-methanol (1:4) (c) Chloroform-methanol (2:1)	2/3 of the plate Up to the top	77
17	Triglycerides, fatty acids, steroids, phospholipids (plant leaves)	Si gel H, 20 × 20 cm, 0.3 mm		Chloroform-methanol 7 N ammonia (65:30:4)			Chloroform-methanol-acetic acid-water (85:10:10:2)		78
17	Triglycerides, fatty acids, steroids, phospholipids (plant leaves)	Si gel H impregnated with 2% aq. ammonium sulphate, 20 × 20 cm, 0.3 mm, 110°C, 1 h		Acetone-benzene-water (91:30:8)			Chloroform-methanol-acetic acid-water (85:10:10:2)		78
15	Bile acids and salts, fatty acids, sterols, phospholipids (gall bladder)	Kieselgel H (type 60), 20 × 20 cm, 0.25 mm, 120°C, 1 h	1.5	Chloroform-ethanol-28% aq. ammonia (25:35:1)	15 cm	Air stream, 30 min	Chloroform-ethanol-ethyl acetate-acetic acid-water (8:6:5:4:1)		79
11	Fatty acids, triglycerides, steroids, phospholipids (serum)	Si gel, 4.5 × 2.6 cm		(a) Chloroform-methanol-water (65:25:4) (b) <i>n</i> -Hexane-diethyl ether-gl. acetic acid (85:20:2)	3 cm 4.5 cm		(c) <i>n</i> -Hexane-diethyl ether-gl. acetic acid (85:20:2) (for lower right-hand side of plate) (d) Butanol-gl. acetic acid-water (60:20:20)	2.6 cm 2.6 cm	80

* (a), (b), (c), (d) indicate the sequence of developments.




Fig. 6. Two-dimensional thin-layer chromatogram of rat lung liquids on silica gel impregnated with boric acid. Abbreviations: NL = neutral lipids; U1, unidentified; BMP, bis(monoacylglyceryl)phosphate; CL = cardiolipin; PE = phosphatidylethanolamine; PC = phosphatidylcholine; PG = phosphatidylglycerol; SM = sphingomyelin; PS = phosphatidylserine; PI = phosphatidylinositol; LPC = lysophosphatidylcholine; O = origin. The solvent for development in the vertical direction was chloroform-methanol-water-concentrated ammonia (70:30:3:2) and in the horizontal direction chloroform-methanol-water (65:35:5). Reproduced from ref. 75 with permission.

4.2. Pigments

This general group contains the various carotenes, xanthophylls, porphyrins and anthocyanidins, *i.e.*, all natural pigments, and the synthetic tannins. Detailed conditions for their two-dimensional separation are reported in Table 6.

Xanthophylls were chromatographed by Knowles and Livingston⁸⁶ on a 1:4 magnesium oxide-Celite strip in the first direction and a 1:6 silica gel G-calcium hydroxide layer in the second. Adsorbability increased with the number of hydroxyl groups on the cyclic moieties of the solutes, the presence of epoxide functionalities especially at the 5,8-position and the length of the conjugated double bond system (Fig. 8). Other separations of carotenoids have been achieved by Tirimanna⁸⁷ and Sherma and Zweig⁸⁸. The hydrocarbonaceous carotenes exhibited the largest R_F values on silica. Owing to its greater number of conjugated double bonds, zeaxanthin was more retained than lutein⁸⁷.

Porphyrin methyl esters were resolved by Elder⁸⁹ on a silica gel plate. A quaternary solvent system was used in the first direction. Retention was increased by carboxylic acid substituents at positions 2 and 4 of the porphyrin ring, as witnessed by the smaller R_F for coporphyrin in comparison with mesoporphyrin. Additional ethanoic acid substituents at positions 1, 3 and 5, as in uroporphyrin, further enhanced the adsorption interactions of the solute with the stationary phase. Of all the

TABLE 5
SEPARATION OF GLYCOLIPIDS
Column headings as in Table 1.

<i>N</i>	<i>Compounds</i>	<i>Stationary phase</i>	<i>SSP (cm)</i>	<i>Solvent 1</i>	<i>t/L</i>	<i>Drying</i>	<i>Solvent 2</i>	<i>t/L</i>	<i>Ref.</i>
	Gangliosides (rabbit brain)	Si gel 60, 20 × 20 cm, 0.25 mm		Chloroform-methanol-0.2% aq. CaCl ₂ (50:40:10) or Chloroform-methanol-0.25% aq. KCl (50:40:10)	4 h		Chloroform-methanol-25 M aq. ammonia (50:40:10) or Chloroform-methanol-0.25% KCl in aq. ammonia (50:40:10)	5 h	82
11	Gangliosides (tissues)	Si gel (Merck), 10 × 10 cm	10 ×	Chloroform-methanol-28% aq. ammonia-water (2 ×)		Over P ₂ O ₅ , 2 h, <i>in vacuo</i>	<i>n</i> -Propanol-28% aq. ammonia-water (75:5:25) or chloroform-methanol water (60:40:9)		81
32	Cardenolides (leaf)	Kieselgel 60 DC Fertigplatten, 20 × 20 cm, 0.25 mm	2	Ethyl acetate-dichloromethane-methanol-water (120:72:7:4)	2 h	Air	Dichloromethane-methanol (9:1)	2 h	84 (Fig. 7)
	Sterioside (leaf diterpene glycoside) purification	Kieselgel G type 60 impregnated with boric acid, 20 × 20 cm, 0.5 mm	60	<i>n</i> -Butanol-ethyl acetate-isopropanol-water (35:100:60:30)			Ethyl acetate-acetic acid-water (8:4:1)		85




Fig. 7. Chromatogram of a mixture of cardenolide standards on silica gel. 1 = Digitoxigenin; 2 = digitoxigenin monodigitoxoside; 3 = digitoxigenin bisdigitoxoside; 4 = digitoxin; 5 = α -acetyldigitoxin; 6 = β -acetyldigitoxin; 7 = lanotoside A; 8 = gitaloxigenin; 9 = gitaloxin; 10 = gitoxigenin; 11 = gitoxigenin monodigitoxoside; 12 = gitoxigenin bisdigitoxoside; 13 = gitoxin; 14 = β -acetylgitoxin; 15 = lanatoside B; 16 = strosposide; 17 = digitalinum verum; 18 = digoxigenin; 19 = digoxigenin monodigitoxoside; 20 = digoxigenin bisdigitoxoside; 21 = digoxin; 22 = digoxoside; 23 = α -acetyldigoxin; 24 = β -acetyldigoxin; 25 = lanatoside C; 26 = deacetyl lanatoside C; 27 = α -anhydrodigoxigenin; 28 = digitatigenin monodigitoxoside; 29 = digitatigenin; 30 = digitatigenin monodigitoxoside; 31 = digitatigenin bisdigitoxoside; 32 = digitatin. Solvent in the first direction, ethyl acetate-dichloromethane-methanol-water (120:72:7:4), and in the second direction dichloromethane-methanol (9:1). Reproduced from ref. 84 with permission.

compounds studied, haematoporphyrin was the only porphyrin with two hydroxyl groups and it was retained the most on the silica plate (Fig. 9).

Saitoh *et al.*⁹⁰ achieved the separation of various metal tetraphenylporphyrin chelates on a reversed-phase plate. The first development was carried out with acetone-propylene carbonate (20:80) and the second with acetone alone. Retention may be related to the ratio of the central metal electronegativity to its ionic radius, En/r_i ⁹¹. Larger En/r_i values correspond to more delocalized chelate-metal bonds and smaller R_F values on the reversed-phase plate. In addition, chromatographic behaviour is affected by the configuration of each complex, whether planar, pyramidal or distorted octahedral, and its stability in each of the eluents used.

Anthocyanidins were resolved on cellulose by Mullick⁹². A methanol-acid-water solvent mixture did not differentiate between the methoxy and hydroxy group substituents on the anthocyanidin aromatic ring. A formic acid-hydrochloric acid or

TABLE 6
SEPARATION OF PIGMENTS
Column headings as in Table 1.

N	Compounds	Stationary phase	SSP (cm)	Solvent 1	t/L	Drying	Solvent 2	t/L	Ref.
14	Xanthophylls	(1) MgO-Celite (1:4) stripe (2) Si gel G-Ca(OH) ₂ (1:6) layer	2.5 cm from end of MgO-Celite stripe	On (1): hexane-acetone (7:3) + ethoxyquin On (2): 2% butanol in benzene + ethoxyquin	40 min	Dry dark, 1-2 min, N ₂		1 h 20 min	86 (Fig. 8)
17	Carotenes, xanthophylls (seeds)	Si gel G, 0.35 mm, 100°C, 30 min		Chloroform-ethyl acetate (40:10)			Diethyl ether		87
7	Carotene, xanthophylls, chlorophyll	Si gel sheets, Eastman K 301 R, 15 × 15 cm		Benzene-acetone (3:1)	Top		Isocane-acetone-diethyl ether (3:1:1)	Top	88
7	Carotene, xanthophylls, chlorophyll	Cellulose sheets		1% n-propanol in light petroleum			Light petroleum-benzene-chloroform-acetone-isopropanol (50:35:10:5:0.17)		88
11	Porphyrin methyl esters	Si gel, Camag D 5, 20 × 20 cm, 0.3 mm, dry at 110°C, 18 h exposure to air	3	Carbon tetrachloride-dichloromethane-methyl acetate-methyl propionate (2:2:1:1)	Top		(a) Benzene-butanone (40:3) (b) Chloroform-kerosene-ethanol (200:100:17)	Top Top	89 (Fig. 9)
7	Metal tetraphenylporphyrin chelates	RP-18 F 254s (Merck), 10 × 10 cm, 0.2 mm, 1 h at 110°C, cool in desiccator	1.5	Acetone-propylene carbonate (20:80) (2 ×)	7.5 cm		Acetone	7.5 cm	90
6	Anthocyanidins	Microcrystalline cellulose, Avicel SF, 20 × 20 cm, 0.25 mm		Formic acid-4 N HCl (2:1)	4.5 h		Methanol-conc. HCl-water (190:1:10)	2 h	92
6	Anthocyanidins	Microcrystalline cellulose, Avicel SF, 20 × 20 cm, 0.25 mm		Methanol-conc. HCl-water (190:1:10)	2 h		n-Amyl alcohol-acetic acid-water (2:1:1)	8 h	92
29	Tannin A or B constituents	Silica (Merck) GF 254, 110°C, 30 min		n-Butanol-acetic acid-water (4:1:5)	2 h, 15 cm		n-Butanol-ethanol-ammonia-water (75:10:15:10)	1.5 h, 15 cm	93




Fig. 8. Two-dimensional, two-adsorbent TLC of a carotenoid mixture on a magnesium oxide-Celite + silica gel G- $\text{Ca}(\text{OH})_2$ plate. C = Carotenes; α = α -cryptoxanthin; β = β -cryptoxanthin; Zeino = zeinoxanthin; L = lutein; Z = zeaxanthin; L_1 and L_2 = lutein *cis*-isomers (corresponding to "neo-A" or "V" and "neo-B" or "U", respectively); V = violaxanthin; V_1 and V_2 = violaxanthin *cis*-isomers; Z_1 and Z_2 = zeaxanthin *cis*-isomers; Luteo = luteoxanthin; N = neoxanthin; N-5,8 = neoxanthin 5,8-epoxide. The first elution is on the magnesium oxide-Celite strip using hexane-acetone (7:3) and the second is on silica gel G- $\text{Ca}(\text{OH})_2$ with 2% butanol in benzene. Reproduced from ref. 86 with permission.

amyl alcohol-acetic acid-water eluent used orthogonally did provide such a differentiation, and the retention increased with the number of hydroxyl substituents on the solute phenyl ring.


Finally, 29 intermediates in the synthesis of tannin A or B were separated on silica using an inorganic-acidic solvent mixture in one direction and an organic-basic eluent in the other. The migration sequence under alkaline conditions corresponded to different degrees of solute condensation or methylation whereas that under acidic conditions depended on the nature of the product⁹³.

4.3. Alkaloids

Heterogeneous nitrogenous substances produced by plants have been grouped under this heading. The two-dimensional chromatography of alkaloids pertaining to the cocaine, harmane, ergot and other families is considered here (Table 7).


Early attempts to separate opium⁹⁴, *Strychnos* and curare⁹⁵ alkaloids by two-directional paper chromatography were time consuming; development in one direction alone could require as long as 17 h.

TLC enabled faster analyses to be achieved. Using a silica gel plate, Munier and Meunier⁹⁶ separated seven cocaine alkaloids. Elution with a non-aqueous neutral mobile phase was followed by an orthogonal development in an acidic solvent



Solvent system A

Fig. 9. Separation of a mixture of porphyrin methyl esters on silica by two-dimensional TLC. 1 = Mesoporphyrin IX; 2 = harderoporphyrin; 3 = coproporphyrin; 4,5,6 = penta-, hexa- and heptacarboxylic porphyrins, respectively; 7 = uroporphyrin; 8 = mono- β -hydroxypropionic monoproponic deuteroporphyrin IX; 9 = bis- β -hydroxypropionic deuteroporphyrin IX; 10 = haematoporphyrin IX; 11 = isohaematoporphyrin IX (all as their methyl esters). Basic structure:



No.	Compound	Substituent in position							
		1	2	3	4	5	6	7	8
1	Mesoporphyrin	CH ₃	C ₂ H ₅	CH ₃	C ₂ H ₅	CH ₃	(CH ₂) ₂ COOH	(CH ₂) ₂ COOH	CH ₃
3	Coproporphyrin	CH ₃	(CH ₂) ₂ COOH	CH ₃	(CH ₂) ₂ COOH	CH ₃	(CH ₂) ₂ COOH	(CH ₂) ₂ COOH	CH ₃
7	Uroporphyrin	CH ₂ COOH	(CH ₂) ₂ COOH	CH ₂ COOH	(CH ₂) ₂ COOH	CH ₂ COOH	(CH ₂) ₂ COOH	(CH ₂) ₂ COOH	CH ₃
10	Haematoporphyrin	CH ₃	CH(OH)CH ₃	CH ₃	CH(OH)CH ₃	CH ₃	(CH ₂) ₂ COOH	(CH ₂) ₂ COOH	CH ₃

Solvent system for the first direction: (A) carbon tetrachloride-dichloromethane-methyl acetate-methyl propionate (2:2:1:1). Solvent systems for the second direction: (B) benzene-butanone (40:3) followed by (C) chloroform-kerosene-methanol (200:100:7). Reproduced from ref. 89 with permission.

TABLE 7

SEPARATION OF ALKALOIDS

Column headings as in Table I.

<i>N</i>	<i>Compounds</i>	<i>Stationary phase</i>	<i>SSP (cm)</i>	<i>Solvent 1</i>	<i>t_R</i>	<i>Drying</i>	<i>Solvent 2</i>	<i>t_R</i>	<i>Ref.</i>
11	Major opium alkaloids	Acid-buffered Whatman paper No. 1 immersed in 2% ammonium sulphate, 28 × 32 cm, 0.3 mm	2.5	Isobutanol-glacial acetic acid (100:20)–water to saturation	16–17 h		90 ml of diethyl ether saturated with aq. phase (water or 0.1 M acetic acid)	2 h	94
7	Cocaine alkaloids (similar polarity and structure)	Si gel 60 (Merck), 20 × 20 cm, 0.25 mm	2.5	Methanol–carbon tetrachloride (50:50)	155 min		Acetone–acetic acid–water (72:8:20)	180 min	96
5	Alkaloids of unequal polarity and structure, morphine, codeine, thebaine, etc.	Si gel 60 (Merck), 20 × 20 cm, 0.25 mm	2.5	Ethyl acetate–2-nitropropane (50:50)	90 min		Methanol–carbon tetrachloride (50:50)	138 min	96
19	Morphine, codeine and derivatives	Activated Si gel G, 20 × 20 cm, 0.25 mm	1.5	Methanol–chloroform–ammonia (85:15:0.7)	15 cm	Air, 10 min	Hydrated diethyl ether–acetone–diethylamine (85:8:7)	15 cm	97 (Fig. 10)
13	Harmaline alkaloids	Si gel 60 (Merck), 20 × 20 cm, 0.25 mm	2.5	Methyl ethyl ketone	98 min		Chloroform–methanol–acetic acid (64:16:20)	185 min	96
15	Alkaloids from <i>Rauwolfia corymbifolia</i> , pseudocinchona	Si gel 60 (Merck), 20 × 20 cm, 0.25 mm		Diisopropyl ether–methanol (85:15)	101 min	Dark, 30 min, 220°C	Acetone–acetic acid (90:10)	74 min	98
37	Ergot alkaloids	Si gel G, 20 × 20 cm, 0.25 mm, 110°C, 30 min	1.5	Methanol–chloroform (2:8)	15 cm	Air, 5 min	Diethylamine–chloroform (1:9)	15 cm	99
10	Ergot alkaloids	Si gel G in high-voltage electrophoresis apparatus, 20 × 20 cm, 0.25 mm	5 cm from anode, 4 cm from bottom edge	Acetic acid–pyridine buffer (pH 5.6), 1500 V, 45 mA	45 min	Fan dry, few minutes	Diethylamine–chloroform (1:9)		99 (Fig. 11)

containing 20% of water. Retention involved adsorption under the former conditions and a mixed partition-adsorption mechanism under the latter. The mobility of each cocaine alkaloid was approximately inversely proportional to the number of functional groups that can be adsorbed simultaneously on the silica gel, and also to the strength of adsorption of each group. Ecgonine, for instance, the only alkaloid in the mixture assayed to possess two strongly adsorptive groups (hydroxyl and carbonyl), on the same side of the six-membered ring, also exhibited the smallest R_F in both directions. Pseudococaine (2 α -carbomethoxy-3 β -benzoxypyrane), with its adsorptive functionalities in different planes although on the same side of the ring, was less retained than cocaine (2 β -carbomethoxy-3 β -benzoxypyrane), where both substituents are co-planar in the β -position.

Munier and Meunier⁹⁶ also achieved the separation of morphine, codeine, thebaine, narcotine and papaverine, alkaloids of different polarity and structure, on a silica gel plate developed with a weakly polar solvent in one direction and a more polar one in the other. Earlier work by Viala and Estadieu⁹⁷ involved the resolution of morphine, codeine and their derivatives by chromatography on activated silica. The first elution with methanol-chloroform-ammonia resulted in three spots, the most retained containing both morphine and codeine, the next corresponding to




Fig. 10. Two-dimensional separation of morphine, codeine and their derivatives on silica. 1 = Morphine; 2 = codeine; 3 = O⁶-monoacetylmorphine; 4 = diacetylmorphine; 5 = acetylcodeine; 6 = nalorphine; 7 = cocaine; 8 = quinine; 9 = nicotine; 10 = cotinine; 11 = methaqualone; 12 = diphenhydramine; 13 = mecloqualone; 14 = hydrochlorobenzethylamine; 15 = N-dimethyldiazepam; 16 = oxazepam; 17 = methadone; 18 = amphetamine; 19 = methylamphetamine. Solvent in the first direction: methanol-chloroform-ammonia (85:15:0.7), and in the second direction: hydrated diethyl ether-acetone-diethylamine (85:8:7). Reproduced from ref. 97 with permission.

heroin and the least retained to O⁶-monoacetylmorphine and acetylcodeine. Orthogonal development in a mixture of hydrated diethyl ether, acetone and diethylamine differentiated all solutes (Fig. 10).

Harmane alkaloids⁹⁶ and alkaloids from *Rauwolfia*, *Corynantha* and *Pseudo-cinchona*⁹⁸ were resolved on silica using a polar neutral mobile phase in the first direction and a more polar, acidic solvent mixture orthogonally. Other than the nature of the solvent itself, which selectively affects the retention order, the pH of the eluent, which determines the ionization state of the molecule as a function of its pK_a value(s), also influences the retention behaviour. Whereas the first development of the silica plate in a neutral solvent did not resolve harmane from norharmane⁹⁶, the subsequent development in the acidic eluent provided such a differentiation; nor-




Fig. 11. Two-dimensional chromatogram of ergot alkaloids separated on silica gel G. Solvents: in direction I, MC = methanol-chloroform (2:8); in direction II, DC = diethylamine-chloroform (1:9). Agro = agroclavine; Chano I = Chanoclavine-(I); Chano II = chanoclavine-(II); Costa = costaclavine; α -Di = α -dihydrolysergol; Dialtry = DL-4-dimethylallyltryptophan; Elymo = elymoclavine; Ecor = Ergocornine; Econi = ergocorninine; Ecris = ergocristine; Ecrisi = ergocristinine; Ecry = ergocryptine; Ecrypti = ergocryptinine; Emet = ergometrine; Emetri = ergometrinine; Esin = ergosine; Esini = ergosinine; Estin = ergostine; Estini = ergostinine; Etam = ergotamine; Etamini = ergotaminine; Festu = festuclavine; Fum A = fumigaclavine A; Fum B = fumigaclavine B; Fum C = fumigaclavine C; Δ -8,9-Lysac = 6-methyl- Δ ^{8,9}-ergolene-8-carboxylic acid; Lysene = lysergene; Lysac = D-lysergic acid; Lysam = D-lysergic acid amide; Isolysam = D-isolysergic acid amide; Lyscaram = D-lysergic acid methylcarbinolamide; Lysine = Lysergine; Lysol = lysergol; Isolysol = isolysergol; Lysval = D-lysergyl-L-valine-methylester; Noragro = noragroclavine; Penni = penniclavine; Isopenni = isopenniclavine; Pyro = pyroclavine; Seto = setoclavine, Isoseto = isosetoclavine. Reproduced from ref. 99 with permission.

TABLE 8
SEPARATION OF AMINO ACIDS
Column headings as in Table 1.

<i>N</i>	<i>Compounds</i>	<i>Stationary phase</i>	<i>SSP</i> (<i>cm</i>)	<i>Solvent 1</i>	<i>t/L</i>	<i>Drying</i>	<i>Solvent 2</i>	<i>t/L</i>	<i>Ref.</i>
13	Cystine, serine, glycine, threonine, alanine, lysosine, valine, phenylalanine, arginine, etc.	Filter-paper sheet, 18 × 22 in.		Collidine	48-72 h		Water-sat. phenol, in atmosphere containing NH ₃		3,100
19	Amino acids	Zeokarb 225 (WR 1.5-2.0) equilibrated with 0.2 <i>M</i> (pH 3.1) buffer		0.2 <i>M</i> buffer (pH 3.1)	1 day		<i>m</i> -Cresol-0.3% ammonia, + over ammonia vapor		102 (Fig. 12)
17	Amino acids	Whatman No. 3 MM filter-paper, 7 × 6 in.	1.5 cm from long paper edge, 4.5 cm from anode	Glacial acetic-formic acid buffer (pH 1.85), 20 kV, 100 mA	7 min	Dry, air, 40°C	2,4/2,5-Lutidine-ethanol-water (11:4:5)	2.5 h	103
18	Amino acids	Whatman No. 3 MM filter paper, 48 × 48 cm		Glacial acetic acid-formic acid, 21. kV (pH 1.85)	70 min	Warm air	<i>n</i> -Butanol-acetic acid water (4:1:5)	16 h	104
20	Amino acids (protein hydrolysate)	Microcrystalline cellulose, Avicel, 20 × 20 cm, 0.5 mm, room temp., 24 h	1.5	1-Butanol-acetic acid-water (3:1:1)	12 cm, 1.5 h	Air dry, room temp., 6 h	Phenol-water (3:1)	12 cm, 2 h	105
12	Amino acids (protein hydrolysate)	Cellulose ECS 6064, 10 × 10 cm, 100°C, 30 min, cool, wash with solvent 1, dry, room temp.	1.5	Methanol-water-pyridine (4:1:2)	25-30 min	Air, ambient temp.	Ethanol-1-butanol-water-propionic acid (5:5:2.5:1)	40-45 min	106

17	Amino acids (urine)	Cellulose + 2 cm ion exchange zone (described)	On ion ex- change zone	Pyridine-dioxane- $\text{NH}_3\text{-H}_2\text{O}$ (35:35:15:15) (2 x)	90 min	<i>n</i> -Butanol-acetone-acetic acid-water (35:35:10:20) (2 x)	60 min	107
26	Polar + non-polar, natural + modified	Cellulose, short fibres, 20 x 20 cm, 0.25 cm		Propanol-formic acid-water (75:5:20)	~20 cm	Methyl ethyl ketone-2-methylpyridine-formic acid-water (31:47:2:20)	16 cm	110 (Fig. 13)
13	Bacterial cell-wall amino acids	Cellulose (Merck), 20 x 20 cm	1.5	Isopropanol-acetic acid-water (75:10:15)	3 h, 15.5 cm	Methanol-pyridine-10 <i>N</i> HCl-water (64:8:2:14)	1.25 h, 15.5 cm	111
10	Amino acids	Si gel G (Merck), 20 x 20 cm, 0.35 mm		<i>n</i> -Propanol-34% ammonia (67:33)	18 cm	0.2% ninhydrin in <i>n</i> -propanol-water glacial acetic acid (64:36:20)	18 cm	112
22	(Not all resolved) amino acids	Silufol (Kavalier), 15 x 15 cm	2	Ethyl acetate-isopropanol-water-ammonia (20:20:25:1.5)	13 cm	Acetone-water-acetic acid-formic acid (50:15:12:3)	13 cm	113
36	Amino acids	Cellulose, 20 x 20 cm	2.5	<i>n</i> -Butanol-acetone-diethylamine-water (10:10:2:5) (pH 12)	Top	Isopropanol-formic acid-water (40:2:10) (pH 2.5)		115
13	Amino acids	Kieselgel G, 20 x 20 cm	20	2 <i>N</i> acetic acid-0.6 <i>N</i> formic acid buffer (1:1), pH 2, 460 V, 12.6 mA	1 h	<i>n</i> -Butanol-glacial acetic acid-water (3:1:1)		114

harmane, with the larger pK_a value, exhibited a higher chromatographic mobility.

Finally, ergot alkaloids were separated on activated silica gel after elution with chloroform-methanol in one direction, followed by an organic solvent-diethylamine in the orthogonal direction (Fig. 11). The strong diethylamine base helped to desorb the alkaloids by competing for adsorption on to silanols⁹⁹. Agurell⁹⁹ also resolved nine ergot alkaloids on silica by combining thin-layer electrophoresis in one direction with chromatography in the other.

4.4. Proteins and their constituents

The two-dimensional chromatographic assays of amino acids, peptides and proteins, important morphological and physiological cellular components, are discussed here. Conditions for their separation are given in Tables 8-10.

4.4.1. Amino acids (Table 8)

Early reports by Consden *et al.*³ and Martin and Synge¹⁰⁰ described the bi-directional separation of several amino acids on paper. The chromatographic behaviour of the solutes was found to be a function of their partition between the mobile phase and the water-saturated cellulose. The analysis time was extremely long, one elution alone requiring 2-3 days. After the pioneering studies of Cohn on ion exchange¹⁰¹, slightly faster separations of amino acids were achieved on sulphonated polystyrene resin-loaded paper, for instance¹⁰². In the assay reported by Knight¹⁰², the first eluent was a 0.2 *M* buffer at a pH of 3.1 and cation exchange determined the elution order: compounds such as lysine, arginine, ornithine, histidine, tryptophan, cysteine, tyrosine and phenylalanine, which contain two amine groups, each proton-




Fig. 12. Two-dimensional separation of amino acids on cellulose. Solvent used for the first development, 0.2 *M* buffer (pH 3.1), and for the second development, 0.3% ammonia in *m*-cresol. Reproduced from ref. 102 with permission.

ated to a certain extent, were retained the most. In the second direction, the cation-exchange phenomenon was suppressed by the eluent, which consisted of 0.3% ammonia in *m*-cresol. The amino acids were chromatographed in their anionic form. Aspartic acid and glutamic acid, solutes with two negative charges, had the smallest R_F values (Fig. 12). Nineteen proteinic amino acids were resolved.

Still faster amino acid separations on paper were achieved by electrophoresis followed by chromatography with a hydro-organic eluent orthogonally^{103,104}.

The development of short-fibre cellulose powder for TLC purposes facilitated the chromatography of amino acids and allowed rapid routine analytical work. In all assays reported on cellulose, the mobile phases in both directions were water-organic mixtures¹⁰⁵⁻¹¹¹. Munier *et al.*^{109,110} achieved the separation of 22 polar and non-polar amino acids, among which certain pairs that were previously unresolved such as leucine-isoleucine, arginine-lysine and serine-glycine. A high degree of resolution is afforded owing to differences between alcohol-solute interactions in the first direction and ketone-solute and tertiary amine-solute interactions in the second, and to the ionic equilibria created by using 2-methylpyridine together with formic acid in the second eluent. The water content of the mobile phase, by affecting solute partition, plays a determining role in these assays (Fig. 13). Munier *et al.*¹¹⁰ found a 20% water content in both directions to be optimal. Smaller percentages did not ensure the elution of the most polar amino acids and higher percentages led to decreased resolution.

Krafczyk *et al.*¹⁰⁷ included a 2 cm wide layer of a strongly acidic ion exchanger at the bottom of the cellulose plate for the desalting of urine samples when analysing




Fig. 13. Two-dimensional separation of amino acids on short-fibre cellulose. Solvent used in the first direction, propanol-formic acid-water (75:5:20), and in the second direction, methyl ethyl ketone-2-methylpyridine-formic acid-water (31:47:2:20). Reproduced from ref. 110 with permission.

TABLE 9
SEPARATION OF Dns- AND DNP-AMINO ACIDS
Column headings as in Table 1.

<i>N</i>	<i>Compounds</i>	<i>Stationary phase</i>	<i>SSP (cm)</i>	<i>Solvent 1</i>	<i>t_R</i>	<i>Drying</i>	<i>Solvent 2</i>	<i>t_R</i>	<i>Ref.</i>
18	Dns-amino acids	Polyamide sheet, 15 × 15 cm		Water-90% formic acid (200:3)			Benzene-acetic acid (9:1)		117
22	Dns-amino acids (re- nal tubular fluid sam- ple)	Polyamide TLC, Schleicher & Schüll, 3 × 3 cm		4% Formic acid			(a) Benzene-glacial acetic acid (8:2) (b) Ethyl acetate- methanol-glacial ace- tic acid (20:1:1)		118 (Fig. 14)
39	Dns-amino acids + others (serum and urine samples)	Micropolyamide plates, F1700, Schleicher + Schüll, 4.2 × 4.2 cm	0.5	Formic acid-water (1.5:98.5)	5 min	Dry, hot air, cool to room temp.	Benzene-acetic acid (4.5:1)	5 min	119
19	Dns-amino acids	Kieselgel G, 0.25 mm		Ethyl acetate- chloroform-ethanol- acetic acid (70:45:4:4)			Chloroform-acetic acid-water (50:45:5)		120
19	Dns-amino acids	HPTLC Fertigplatten RP-8 F 254, 10 × 10 cm		<i>n</i> -Heptane-ethyl acetate-acetic acid (65:33:2)	7 cm		Methanol-0.01 <i>M</i> Na ₂ HPO ₄ (75:25)	7 cm	121
20	DNP-amino acids	RP-18		Hexane-ethyl acetate- acetic acid (80:18:2)	6 cm		1 <i>M</i> ammonia + 3% KCl in 60% methanol	6 cm	34 (Fig. 2)
17	DNP-amino acids	Polyamide, 15 × 15 cm, 100°C, 15 min	2.0	<i>n</i> -Butanol-glacial acetic acid (90:10)	10 cm, 200 min	Dry, hot air, 15 min	90% formic acid- water (50:50)	11 cm, 60 min	122
18 + 23	DNP- + Dns-amino acids	Cellulose powder, Macherey, Nagel & Co., 30 × 20 cm, 0.11 mm		Toluene-ethylene gly- col chlorhydrin- pyridine-0.8 <i>M</i> am- monia (150:90:45:90) + a few drops of <i>n</i> - octanol	30 cm		Water-water satd. with ammonium sulphate-sodium do- decyl sulphate (700 ml:100 ml:567 mg)	20 cm	124

for urinary amino acids. Other separations on silica gel were reported^{112,113}. An organic-basic eluent was used in the first direction followed by an organic-acidic mixture in the second. Solute retention depended on the types of functional groups and their degree of ionization. Ionophoresis followed by chromatography on silica gel has also been applied to the separation of amino acids¹¹⁴.

4.4.2. Dimethylaminonaphthalenesulphonyl (Dns) and dinitrophenyl (DNP) amino acids (Table 9)

Seiler¹¹⁶ has reviewed the separations of various Dns-derivatives, among which Dns-amino acids. Two-directional TLC has been achieved on a polyamide stationary phase¹¹⁷⁻¹¹⁹ (Fig. 14) and has recently been applied to the identification of human acidopathies in serum and urine¹¹⁹. Although Dns-amino acids have also been analysed on silica¹²⁰, assays on polyamide are preferred as they are less time consuming, do not result in spot tailing and do not require previous deionization of urine samples¹¹⁹. An aqueous eluent is generally used in the first direction followed by an organic eluent in the second. This change in the nature of the mobile phase causes alterations in the sorbent properties such that the mechanisms for solute retention are different in each direction. Polyamide is thought to act as a reversed phase in the presence of the aqueous eluent, and it behaves as a hydrophilic sorbent there-




Fig. 14. Two-dimensional chromatogram of some Dns-amino acids on polyamide after autoradiography. 1 = Starting point; 2 = taurine; 3 = Dns-OH; 4 = cystine; 5 = *n*-serotonin; 6 = tryptophan; 7 = bis-tyrosine; 8 = *n*-tyrosine; 9 = ornithine; 10 = bis-lysine; 11 = phenylglycine; 12 = phenylalanine; 13 = leucine and histidine; 14 = isoleucine; 15 = methionine; 16 = norvaline; 17 = valine; 18 = proline; 19 = GABA; 20 = δ -aminolevulinic acid; 21 = alanine and Dns-NH₂; 22 = glycine; 23 = glutamic acid; 24 = aspartic acid; 25 = hydroxyproline; 26 = glutamine; 27 = asparagine; 28 = threonine; 29 = serine; 30 = arginine, ϵ -lysine and α -lysine; 31-33 = unknown dansylation products. Solvent used in the first direction, 4% formic acid, and in the second direction, benzene-glacial acetic acid (8:2) followed by ethyl acetate-methanol-glacial acetic acid (20:1:1). Reproduced from ref. 118 with permission.

TABLE 10
SEPARATION OF PEPTIDES
Column headings as in Table 1.

<i>N</i>	<i>Compounds</i>	<i>Stationary phase</i>	<i>Solvent 1</i>	<i>t/L</i>	<i>Solvent 2</i>	<i>t/L</i>	<i>Ref.</i>
11	Peptides	Cellulose MN 300, 20 × 20 cm, 0.25 mm	<i>tert.</i> -Amyl alcohol-eucalyptol-2-chloroethanol-pyridine-98% formic acid (150:150:200:280:43:200)	Top, 5 h	<i>n</i> -Amyl alcohol-pyridine-H ₂ O (4:4:2)	5 h	125
8	Oligopeptides	Cellulose MN 300, 0.25 mm	Isoamyl alcohol-ethanol-glacial acetic acid-pyridine-water (17.5:5:1.3:17.5:15) (pH 5.75)		Isopropanol-ethanol-ammonia (20:20:15) (pH 12.3)		126
19	Peptides	Cellulose MN 300, 40 × 20 cm, 0.25 mm	Electrophoresis, 1.04 <i>N</i> formic acid, 427 V/40 cm, 8 mA/20 cm × 0.25 mm	3.5 h	<i>n</i> -Butanol-pyridine-98% formic acid-water (479:278:43:200)	To top, 3 h	125
19	Peptides	Cellulose MN 30, 40 × 40 cm, 0.25 mm	Pyridine-ethanol-water (4:4:2)	Top, 4 h	Electrophoresis, 0.025 <i>M</i> aq. borax soln., 380 V/40 cm, 4.5 mA/20 cm × 0.25 mm	4 h	125
	Polypeptides/proteins with acidic or neutral <i>pI</i>	(1) Tube gel, polyacrylamide (2) Slab gel, acrylamide gradient	(1) 9 <i>M</i> urea, Nonidet P-40, pH 3.5-10 Ampholines, 400-700 V	20 h	(2) Electrophoresis in presence of sodium dodecyl sulphate	3 h	128, 129
	Basic proteins	(1) Tube gel, polyacrylamide (2) Slab gel, acrylamide gradient	(1) Dipalmitoyl-L- α -phosphatidylcholine + urea, pH 3.5-10 Ampholines, 800 V	2.6 h	(2) Unchanged, 3200 V/h		133

after¹²¹. Thus, whereas the R_F values for arginine and aspartic acid are relatively large in the first direction, they are much smaller in the second¹¹⁷⁻¹¹⁹.

Exploiting similar solvent changes with a reversed-phase plate, Macek *et al.*¹²¹ succeeded in resolving 19 Dns-amino acids. The RP-8 TLC plate behaved as a normal phase in presence of the non-aqueous eluent used in the first direction and as a true reversed phase when developed with an aqueous solvent in the second direction. Exact reversal of the retention order of the Dns-glycine, -alanine, -valine and -leucine could be observed in the second direction with respect to the first elution. In addition, the aqueous eluent was basic enough to reduce the amino group ionization and hence to increase solute retention on the reversed-phase plate.

DNP-amino acids were separated under similar conditions on an RP-18 thin-layer plate³⁴. It is interesting that the addition of a salt to the non-aqueous solvent did not modify the solute behaviour, whereas it increased the retention when present in the aqueous eluent, a true expression of reversed-phase solvophobic behaviour. Separations of DNP-amino acids were also accomplished on polyamide¹²² and cellulose^{123,124} by development with an organic solvent in the first direction and an aqueous solvent orthogonally. The efficiency of the chromatographic system required judicious selection of the mobile phase components. Munier and Drapier¹²⁴ included a few drops of *n*-octanol in the first eluent in order to suppress the chromatography of the solvent itself. In addition, as impregnation of the plate with the organic mobile phase caused irregularities in the perpendicular progression of the second aqueous solvent, spot tailing and diffusion, sodium dodecyl sulphate was added to this aqueous phase and helped to achieve small and round spots.

4.4.3. Peptides and oligopeptides (Table 10)

Peptides¹²⁵ and oligopeptides¹²⁶ have been separated on cellulose with an acidic solvent in one direction and a basic solvent in the other. In addition to the differences in solute ionization which originate from the use of eluents at different pH values, differences in molecular weight also affect the retention and the partition process. In a given category of compounds, the dipeptides for instance, solutes with a large number of hydrophobic carbon centres, more soluble in the organic portion of the eluent, exhibit larger R_F values¹²⁶.

Other peptide assays on cellulose¹²⁵ have combined electrophoresis in one direction and chromatography with an organic-aqueous mobile phase in the other. The electrophoresis step necessitated the use of binder-free cellulose, sufficiently large plates, a weak gradient potential and an adequate electrolyte solution. Peptides migrated according to their charge, the solute pK_a values and the pH of the electrolyte.

4.4.4. Proteins

Although not related to TLC, the high-resolution, two-dimensional electrophoretic method developed by O'Farrell¹²⁷ and later by Anderson and co-workers¹²⁸⁻¹³⁶ should also be mentioned. Nearly 1000 spots were obtained from such an analysis of lymphocyte proteins using autoradiography (Fig. 15). More sensitive detection techniques are expected to afford the resolution of about 5000 components.

In the first dimension, proteins or polypeptides were separated according to their charge by isoelectric focusing in a narrow gel tube. Next, the gel was extruded and positioned on a polyacrylamide gradient slab, and electrophoresis was performed




Fig. 15. Diagram showing the spots resulting from the two-dimensional gel electrophoresis of lymphocyte proteins. Numbers are the key to information about the respective spots in the lymphocyte protein index data base developed by Anderson *et al.* For further details see ref. 48. Reproduced from ref. 48 with permission.

in the presence of sodium dodecyl sulphate. In this second dimension, compounds were resolved according to their molecular weight, as sodium dodecyl sulphate binds electrostatically to the solutes, thus masking charge differences (Table 10).

The method described was applied successfully to a study of the protein constituents of physiological fluids and tissues in normal and diseased individual.^{49,50, 129-131,134,135,137-140}

Because of the considerable number of proteins that can be separated in one analysis, data handling becomes very complex as huge files have to be stored and monitored. The corresponding software has been developed by Anderson *et al.*⁴⁸ and Lester *et al.*⁴⁹. A discussion of the problems of spot detection, segmentation, integration and pairing is beyond the scope of this review.

4.5. Carbohydrates (Table 11)

Sugars have been separated on cellulose¹⁴¹ and Kieselgel G¹⁴², developed with different water-organic solvents. Hotta and Kurokawa¹⁴³ modified a silica gel plate by dipping it in 0.2 M sodium dihydrogen phosphate solution, a procedure known to enhance sugar retention in proportion to the number of hydroxyl substituents¹⁴⁴. This selective stationary phase was developed with an eluent containing 10% of water in the first direction and 25% in the second.

The effect of water on chromatographic mechanisms can be illustrated by the behaviour of rhamnose and fucose in each eluent. These sugars differ in the steric arrangement of their functional groups: rhamnose has one pair of *cis*-hydroxyl groups in a β -position with respect to a bulky methyl group, compared with two pairs of *cis*-hydroxyl groups in fucose, one β - and one α - to the methyl group. With the first eluent, the adsorption of rhamnose may be more sterically hindered than that of fucose and thus rhamnose migrates faster. These structural differences are no longer important with the second eluent, as solute adsorption on to silica is reduced owing to plate deactivation by water present in the mobile phase. The solutes then are believed to partition between the mobile and the modified stationary phases, which results in similar R_F values for rhamnose and fucose (Fig. 16).

Carbohydrates have also been separated on silica gel impregnated with sodium tetraborate, sodium tungstate¹⁴⁵ and boric acid¹⁴⁴. In the first direction, a neutral solvent was used and sugars were retained according to their capacity to form borate complexes. In the second elution, the impregnating agent was completely deactivated by the acidic solvent. Solute partition was in effect and the heavier disaccharides were retained the longest. Phenylboronic and boric acids have more recently been used in the mobile phase to resolve carbohydrates on silica¹⁴⁶. Other separations have been accomplished on cellulose developed in different mobile phases¹⁴⁷.

4.6. Glycopeptides

Moczar¹⁴⁸ reported the separation of some glycopeptides from a glycoprotein enzymatic digest on silica. Electrophoresis in presence of acetic acid-pyridine-water (10:1:89) (pH 3.8) was performed for 2-3 h at 10-20 V/cm on a 20 × 20 cm plate. This operation was followed by a continuous development with ethanol-nitromethane-acetic acid-water (5:3:3:3) for 18-30 h.

TABLE II
SEPARATION OF CARBOHYDRATES
Column headings as in Table I.

N	Compounds	Stationary phase	SSP (cm)	Solvent 1	t/L	Drying	Solvent 2	t/L	Ref.
12	Sugars	Cellulose MN 300 G, 110°C, 10 min	2	<i>n</i> -Butanol-acetone-diethylamine-water (10:10:2:5)		80°C, 30 min	Vapour phase equilibrated with 25% ammonia, phenol-water (3:1)		141
4	Man, Gal, Glc-NAc, Man, 1-6-Man, Gal- β 1-4-Glc-NAc	Kieselgel G (Merck)		<i>n</i> -Butanol-acetic acid-water (12:3:5)	16 h		<i>n</i> -Propanol-ethyl acetate-water (7:2:1)	6-8 h	142
8	NANA, Gal, Glc, Man, Gal-NAc, Glc-NAc, Fuc, Rham	Si gel F ₂₅₄ dipped in 0.2 <i>M</i> sodium dihydrogen phosphate, 110-120°C, 30 min, 20 $\times$ 20 cm, 0.25 mm		Butanol-acetone-water (4:5:1) (2 $\times$)	15 cm		Phenol-water (3:1) (2 $\times$)		143 (Fig. 16)
40	CHO, including deoxy sugars and methylglycosides	Si gel: (1) 4.2 cm, impregnated with sodium tetraborate-boric acid-sodium tungstate (2) 14.5 cm, impregnated with boric acid		On (1): methyl acetate-isopropanol-water (2:2:1)		550°C < 3 h	On (2): ethyl acetate-acetic acid-methanol-water (60:15:15:10)		144
16	CHO,	Sodium tetraborate-sodium tungstate impregnated silica, 20 $\times$ 20 cm, 0.3 mm, 90°C, 30 min		Ethyl acetate-isopropanol-water (2:2:1)	2 h	70°C, 4-5 h	Ethyl acetate-acetic acid-methanol-water (60:15:15:10)	50 min	145
13	Sugars	Si gel 60 (Merck)		(a) Ethyl methyl ketone-diisopropyl ether-2-propanol-pyridine-water-phenylboronic acid (40:10:30:10:10:1:2)	10 cm		(b) Diisopropyl ether-methanol-acetic acid-water (88:12:2:2) (c) Ethyl methyl ketone-2-propanol-acetonitrile-0.5 <i>M</i> boric acid + 0.25 <i>M</i> isopropylamine-acetic acid (40:30:20:15:0:4)	10 cm	146
15	Sugars	Cellulose, 20 $\times$ 20 cm, 0.1 mm		2-Propanol-90% formic acid-water (80:4:20)	5-6 h		Lutidine (2,6-dimethylpyridine)-water (65:35)	6 h	147




Fig. 16. Two-dimensional TLC of sugar mixture on silica gel dipped in 0.2 *M* sodium dihydrogen phosphate. Rha = rhamnose; Fuc = fucose; GlcNAc = acetylglucosamine; GalNAc = acetylgalactosamine; Man = mannose; Glc = glucose; Gal = galactose; NANA = N-acetylneuraminic acid. First direction: two successive developments in butanol-acetone-water (4:5:1). Second direction: two successive developments in phenol-water (3:1). Reproduced from ref. 143 with permission.

More recently, the soluble inhibitory factor (SIF), a molecule constituted of both protein and glycolipid moieties, has been isolated from serum and purified on a silica plate using chloroform-methanol-water in the proportions 60:38:8 in the first dimension and 50:50:5 in the second¹⁴⁹.

4.7. Nucleic acids and their constituents

In cells, nucleic acids are carriers of genetic information and their constituents, the free bases, nucleosides and nucleotides, actively participate in cellular metabolism. Detailed conditions for the two-dimensional separation of these compounds are given in Tables 12 and 13.

4.7.1. Bases, nucleosides and nucleotides (Table 12)

Randerath and Randerath¹⁵⁰ achieved the first rapid, bi-directional separation of 23 ribonucleotides on PEI-cellulose. Several mono-, di- and trinucleotides were

TABLE 12
SEPARATION OF NUCLEOTIDES, NUCLEOSIDES AND BASES

Column headings as in Table 1.

<i>N</i>	<i>Compounds</i>	<i>Stationary phase</i>	<i>SSP (cm)</i>	<i>Solvent 1</i>	<i>t/L</i>	<i>Drying</i>	<i>Solvent 2</i>	<i>t/L</i>	<i>Ref.</i>
23	Ribonucleotides	PEI, cellulose, 20 × 20 cm, 0.5 mm	20 ×	(a) 0.2 <i>M</i> LiCl (b) 1.0 <i>M</i> LiCl (c) 1.6 <i>M</i> LiCl	2 min 6 min 13 cm or 75 min	Air, < 50°C, wash LiCl in CH ₃ OH	(d) 0.5 <i>M</i> formate buffer (pH 3.4) (e) 2.0 <i>M</i> formate buffer (f) 4.0 <i>M</i> formate buffer	30 sec 2 min 15 cm or 60 min	150 (Fig. 17)
8	2', 3'-G, A, U, CMP	PEI, cellulose MN 300	3	1.0 or 0.1 <i>N</i> formic acid (pH 3.6)	10-15 cm	Air	1 <i>M</i> LiCl (pH 7.0)	8 cm	151
9	Adenine + pyridine phosphates	PEI, cellulose (Brinkmann) 20 × 20 cm, 0.1 mm	3	0.5 <i>M</i> formate buffer (pH 3.6)		Dry, wash in CH ₃ OH, dry	0.2 + 1.0 <i>M</i> LiCl		152
14	Phosphate esters	Cellulose MN 300, 20 × 17.5 cm, 0.25 mm		<i>n</i> -Propanol- ammonia-water (6:3:1) (2 ×)	3 h		<i>n</i> -Propyl acetate-90% formic acid-water (11:5:3) (2 ×) or electrophoresis with 0.28 <i>M</i> acetate buffer (pH 3.6), 1000 V, 35 mA	80 min 16 min	153
22	Nucleosides, nucleotides	Cellulose MN 300		Isobutyric acid-conc. ammonia-water (57:4:39)			Methanol-1 <i>M</i> ammonium acetate (7:3) (2 ×)		154
8	2', 3'-G, A, U, CMP	Cellulose MN 300, 100°C, 30 min, 20 × 20 cm, 0.5 mm	3	<i>n</i> -Butyric acid- ammonia-water (57:4:39)	5-6 h, 12-15 cm	Over- night in hood	Sat. ammonium sulphate-1 <i>M</i> sodium acetate-isopropanol (79:19:2)	3-4 h, top	155

8	5'-O-Methyl-nucleotides	Cellulose		Isobutyric acid-0.5 N ammonia (5:3)		Isopropanol-conc. HCl-water (17:15:15)	156
5	Bases: A, C, G, T, bromo-U	Cellulose (Eastman 6065), 3.5 × 20 cm, 20 × 20 cm (2 layers)	2	Isopropanol-water-HCl (65:18.4:16.6)	14 cm 6 h	1 M disodium hydrogen phosphate-1.2 M ammonium sulphate-isopropanol (123:74:3)	36
37	Purine and pyrimidine nucleosides and bases (not all resolved) in urine	Cellulose DC-Alufolien, 10 × 10 cm, 0.1 mm	1.5	Isopropanol-5% ammonia (8:2) (2 ×)	90 min	Butanol-acetic acid-H ₂ O (8:2:2) (2 ×)	157
44	Bases, nucleosides, nucleotides (not all resolved)	Cellulose MN 300, 20 × 20 cm, 0.5 mm	3	n-Propanol-25% ammonia-water (6:3:1)	3 h, 18 cm	Isopropanol-sat. ammonium sulphate-water (2:79:19)	158 (Fig. 18)
19	Bases, nucleosides, nucleotides	Polygram CEL 400, UV 254 nm, 10 × 10 cm		(a) Methanol (b) Methanol-isopropanol-25% (w/w) ammonia-1% (w/v) EDTA (3:9:2:6) (c) Solvent (b)	Middle Top	(d) Isobutyric acid-25% (w/w) ammonia-1% (w/v) EDTA (200:9:114)	159
22	Methylated tRNA bases	Si-cellulose (40:60), 20 × 20 cm, 0.4 mm, desiccated	2-3	Ethyl acetate methanol-water-88% formic acid (100:25:20:1)	150 min 60 min	Acetonitrile-ethyl acetate-2-propanol-1-butanol-58% ammonia-water (40:30:20:10:22:5)	162
13	Bases, nucleosides, nucleotides	Si gel (Merck) HPTLC F ₂₅₄ , 10 × 10 cm	2	1-Butanol-acetone-acetic acid-5% ammonia-water (10:5:2:3:2)	7 cm Thorough drying	1-Propanol methanol-ammonia water (10:1:5:2)	163

resolved, as well as some nucleotide sugars. Increasing concentrations of lithium chloride solutions and formate buffer (pH 3.4) were used stepwise in the first and second directions, respectively. Compared with elution at constant concentration, step-gradient elution resulted in sharper circular spots. Solutes were resolved according to their charge: the most negatively charged trinucleotide species migrated the least on the PEI anion exchanger (Fig. 17). Similar conditions were later applied by Santini and Ulrich¹⁵¹ to the separation of 2'- and 3'-adenine, guanine, uridine and cytosine monophosphates, and by Barton *et al.*¹⁵² to that of nine pyridine and adenine nucleotides.

Various phosphate esters^{153,154}, 2'- and 3'-monophosphates¹⁵⁵, 2',O-methyl nucleotides¹⁵⁶, purine and pyrimidine bases³⁶ and nucleosides¹⁵⁷ have also been chromatographed on cellulose. The pH of the eluent, which affects the extent of solute dissociation, plays an important role in these assays. Thus, in most instances, development was accomplished with an acidic solvent in one direction and a basic solvent in the other.

The use of a basic solvent has been coupled with that of a salt in order specifically to affect nucleotide retention^{158,159}. A mixture of 44 bases, nucleosides, deoxynucleosides, 2'-, 3'- and 5'-nucleoside monophosphates, nucleoside di- and triphosphates and nucleotide sugars was resolved by Pataki¹⁵⁸ on cellulose. The eluents were *n*-propanol-25% ammonia-water (5:3:1) in the first direction and isopropanol-saturated ammonium sulphate-water (2:79:19) in the second. Triphosphates were retained the most with the first eluent. The second buffer system enhanced their migration whereas it hardly affected the position of nucleosides and bases (Fig. 18). Randerath and Randerath¹⁶⁰ had previously observed these ionic strength effects on the chromatographic behaviour of nucleic acid constituents.

Using a silica gel-cellulose plate with acidic and basic solvents, Munns and co-workers^{161,162} resolved 22 methylated tRNA purine and pyrimidine bases. Solutes with primary amine groups such as adenine, guanine and cytosine exhibited lower R_F values in a formic acid than in an ammonia system. These differences were found to arise from interactions with silica, the adsorption of the primary amines also being decreased by their methylation. At a basic pH, both the bases and the silica phase are negatively charged, which weakens adsorbent-solute interactions.

Finally, Das¹⁶³ separated 13 bases, nucleosides and nucleotides on pure silica by varying the solvent pH and composition in the second direction.

4.7.2. Oligonucleotides and nucleic acids (Table 13)

Miller and Burgess¹⁶⁴ resolved five oligonucleotides on PEI-cellulose by means of lithium chloride-EDTA solution at pH 6.5 in the first direction and at pH 2.0 in the other. The charges on both the phosphate and base moieties affected the separation. Also by varying the pH of the eluent, Mirzabekov and Griffin¹⁶⁵ were able to separate 18 oligonucleotides on PEI-cellulose. The first direction involved successive developments with 1.4 and 1.8 *M* solutions of lithium formate at pH 3.4; the lower concentration ensured the resolution of smaller oligonucleotides and the higher one that of the larger solutes. The second eluent was 0.8 *M* lithium chloride solution in Tris base at pH 8.0. The chromatographic assay was carried out at 60°C for sharper spot formation. Migration in the first direction was according to charge and nucleotide composition whereas in the second direction it was more a function of molecular




Fig. 17. Two-dimensional anion-exchange separation of nucleotides on PEI-cellulose. First dimension: successive developments in (a) 0.2 *M* LiCl, (b) 1.0 *M* LiCl, (c) 1.6 *M* LiCl. Second dimension: successive developments in (d) 0.5 *M* formate buffer (pH 3.4), (e) 2.0 *M* formate buffer, (f) 4.0 *M* formate buffer. CMP = cytosine 5'-monophosphate; AMP = adenosine 5'-monophosphate; CDPG = cytidine diphosphate glucose; UMP = uridine 5'-monophosphate; UDPAG = uridine diphosphate-N-acetyl-glucosamine; ADPG = adenosine diphosphate glucose; IMP = inosine 5'-monophosphate; GMP = guanosine 5'-monophosphate; GDPM = guanosine diphosphate mannose; CDP = cytosine 5'-diphosphate; ADP = adenosine 5'-diphosphate; UDPGA = uridine diphosphate glucuronic acid; UDP = uridine 5'-diphosphate; IDP = inosine 5'-diphosphate; GDP = guanosine 5'-diphosphate; CTP = cytosine 5'-triphosphate; ATP = adenosine 5'-triphosphate; UTP = uridine 5'-triphosphate; ITP = inosine 5'-triphosphate; GTP = guanosine 5'-triphosphate; DPN = nicotinamide adenine dinucleotide; TPN = nicotinamide adenine dinucleotide phosphate. Reproduced from ref. 150 with permission.

weight. More complex mixtures of half or whole 5S-RNA molecules were resolved by continuous gradient elution. The concentrations of the salt in the eluents depended on the fragments to be separated, the chromatography of a whole 5S-RNA molecule requiring 4.5 *M* lithium formate.

The separation of more than seven methylated mRNA 5'-termini, achieved by Gross *et al.*¹⁶⁶ on cellulose, again involved different ionic equilibria in each direction and their effect on solute migration patterns. Faster assays on cellulose have been reported by Bergquist¹⁶⁷, who separated 17 oligo- and mononucleotides by coupling electrophoresis with TLC. The system described is extremely powerful, capable of resolving tetra- or oligonucleotides of similar charge and chain length.

The combination of gel electrophoresis and TLC has more recently been applied to RNA sequencing studies¹⁶⁸. First, 3'-terminal RNA half-molecules are sep-

TABLE 13
SEPARATION OF OLIGONUCLEOTIDES AND NUCLEIC ACIDS
Column headings as in Table 1.

<i>N</i>	<i>Compounds</i>	<i>Stationary phase</i>	<i>SSP (in.)</i>	<i>Solvent 1</i>	<i>t/L</i>	<i>Drying</i>	<i>Solvent 2</i>	<i>t/L</i>	<i>Ref.</i>
5	5'-RNA terminal oligonucleotides	PEI, cellulose MN 300, Brinkmann, 20 × 20 cm	1	2.0 M LiCl + 0.01 M EDTA (pH 6.5)	Top	Wash in CH ₃ OH, dry	3 M LiCl + 0.01 M EDTA-3.6 M formic acid (pH 2.0, adjusted with solid LiOH)		164
18	Oligonucleotides (T1 RNase digest of 5S RNA)	PEI, cellulose MN 300, 20 × 20 cm		(a) 1.4 M lithium formate in 7 M urea (pH 3.4), 60°C (b) Above mixture, 1.8 M (pH 3.4), 60°C	10 cm, top	Wash in CH ₃ OH, dry	(c) 0.8 M LiCl in 0.02 M TMS base and 7 M urea (pH 8)		165
	Partial fragments of 5S RNA	PEI, cellulose MN 300, 20 × 20 cm		4.0 M lithium formate in 7 M urea; gradually introduce 7.5 M above solution (pH 3.4)			1.0 M LiCl in 7 M urea; gradually introduce 3.2 M LiCl (pH 8), 60°C		165
>7	Methylated mRNA 5'-termini	Cellulose (Merck), 20 × 20 cm		Isobutyric acid-ammonia-water (pH 23.7) (66:1:33)	8 h	Hood, overnight	Sat. ammonium sulphide-1 M sodium acetate (pH 5.5)-propan-2-ol (40:9:1)	6-7 h	166
17	Oligonucleotides + mononucleotides	Cellulose MN 300, 20 × 20 cm, 0.25 mm		Electrophoresis with 0.1 M ammonium formate (pH 2.45) containing 0.001 M EDTA, 50V/cm, 16-20 mA	20-25 min	Dry under IR lamp	<i>tert.</i> -Butanol-0.08 M formic acid-isoamyl alcohol (50:50:2)	10 cm, 4 h	167
	(1) Oligonucleotides (2) Nucleotide sequencing	(1) Gel electrophoresis (2) DEAE-cellulose		(1) Electrophoresis			(2) 15 mM sodium citrate buffer (pH 3.1)		168
17	Oligonucleotides	(1) Cellulose acetate paper strips (2) DEAE-cellulose plates		(1) Electrophoresis (pH 3.5)			(2) Homochromatography		169




Fig. 18. Two-dimensional separation of nucleo-derivatives on purified cellulose layers. First dimension, *n*-propanol-25% ammonia-water (6:3:1); second dimension, isopropanol-saturated ammonium sulphate-water (2:79:19). Absorption spots are hatched, fluorescence spots are surrounded by broken lines. Reproduced from ref. 158 with permission.

arated by gel electrophoresis. Then, the entire ladder of bands is excised from the gel, transferred to a thin layer of DEAE-cellulose by blotting and hydrolysed *in situ* with takadiastase RNAase. The thin layer is developed in a direction perpendicular to gel electrophoresis with 15 mM sodium citrate buffer (pH 3.1). The enzyme is supposed to hydrolyse one nucleotide at a time and thus the resulting pNp spots correspond to the sequence of nucleotides in the original RNA molecule.

4.8. Environmental pollutants (Table 14)

It is of interest for the environmental chemist to identify and quantify poly-aromatic hydrocarbons and aza heterocycles in air samples from different sources. This was achieved by Sawicki and co-workers¹⁷⁰⁻¹⁷⁶, who analysed by two-dimensional TLC extracts obtained directly or after column fractionation of airborne particulate and coal-tar pitch fume samples.

TABLE 14
SEPARATION OF ENVIRONMENTAL POLLUTANTS: AZAARENES AND POLYNUCLEAR AROMATIC HYDROCARBONS (PAHs)
Column headings as in Table 1.

N	Compounds	Stationary phase	SSP	Solvent 1	t/L	Drying	Solvent 2	t/L	Ref.
8	Azaarenes + others	Al ₂ O ₃		Cyclohexane-ethyl acetate (1:1)	15 cm		Toluene	15 cm	170
33	Benzene-soluble fraction of airborne particulate sample; acridine, pyrene, benz[a]acridine identified	Alumina-cellulose (2:1), 20 × 20 cm, 0.25 mm		Pentane	15 cm		DMF-water (35:65)	15 cm?	173
41	Benzene-soluble airborne particulate fraction with anthracene + other PAHs	Alumina-cellulose acetate (2:1)		Pentane	15 cm		Ethanol-toluene-water (17:4:4)	15 cm	173
65	Coal-tar pitch basic fraction with various azaarenes	Si gel-cellulose (2:1), 20 × 20 cm	1.5 cm from base	Pentane-diethyl ether (9:1)	13 cm, 3 h		DMF-water (35:65)	13 cm, 4.5 h	177 (Fig. 19)
67 or 15	Tokyo air with various PAHs; PAH reference solutions	(1) Alumina, 4 × 20 cm (2) 26% acetylated cellulose, 16 × 20 cm, 0.3 mm. 1 h, 110°C, store at 20% humidity		On (1): n-hexane-diethyl ether (19:1) with sat. potassium acetate, 2% humidity	15 cm, 35 min	Air, 5 min	Methanol-diethyl ether-water (4:4:1) (3 × for Tokyo air sample)	10 cm, 55 min	178
Numerous	Benzene-enriched airborne particulate extract; no spot identification	Chemically bonded phase, 5 × 5 cm		Methanol-water (97:3)	2 min		Methanol-acetonitrile	4 min	184

Of all of the compounds in an airborne particulate, benzene-soluble, "oxy" fraction, compounds with an aza nitrogen such as 9-acridone and 6(5H)-phenanthridone exhibited the longest retention on an alumina plate. Substitution of the aza nitrogen by an oxygen atom or sulphur weakened adsorption, xanthen-9-one and thioxanthen-9-one migrating faster than 9-acridone¹⁷⁰.

In their assay on an alumina-cellulose (2:1) mixed sorbent plate, Sawicki *et al.*¹⁷³ used an organic eluent for the first elution. Compounds were primarily retained by adsorption; acridine had the smallest and pyrene the largest R_F values. In the orthogonal direction, the plate was developed with a solvent containing 65% of water. Water interfered with solute adsorption on alumina and also impregnated the cellulose fibres, creating a partitioning medium. Under these conditions, acridine migrated further than pyrene.

Similar mechanisms were effective in the two-dimensional separation of coal-tar pitch aza heterocyclic on a silica-cellulose (2:1) plate. Adsorption on to silica prevailed on development with the organic eluent: owing to the steric hindrance of its aza nitrogen, benzo(*h*)quinoline was less retained than benzo(*f*)quinoline (Fig. 19). In the orthogonal direction, compounds were separated according to their partition coefficients: the bulky dibenz(*a,j*)acridine was retained the most whereas acridine, benzo(*h*)- and benzo(*f*)quinoline had the largest R_F values; in general, retention increased with the number of additional aromatic substituents¹⁷⁷.

Matsushita and Suzuki¹⁷⁸ succeeded in resolving the polyaromatic hydrocarbons anthracene, phenanthrene, pyrene, benz(*c*)anthracene, chrysene, perylene, coronene, benz(*a*)- and benz(*c*)pyrene, benzophenylene, dibenzanthracene and dibenzopyrene. The first development was carried out on an alumina strip with an organic eluent. Adsorption increased with aromaticity and was affected by steric factors. The second development was performed on the acetylated cellulose part of the plate with a polar eluent. Compounds were separated by partition, those with the smallest number of rings migrating further than the others.

Polyaromatic hydrocarbons have more recently been separated on acetylated cellulose¹⁷⁹ and on an acetylated cellulose-silica gel-alumina sorbent mixture¹⁸⁰. Other methods had involved the use of GC and TLC in combination^{181,182}. Janák¹⁸² and Kaiser¹⁸³ reviewed in depth both the instrumentation and the applications of this technique.

4.9. Pesticides

Various pesticides, including aldrin and dieldrin¹⁸⁵, DDE and DDT analogues¹⁸⁶, organophosphorus¹⁸⁷ and other compounds¹⁸⁸⁻¹⁹⁰, have been separated on silica plates developed with different solvents in two directions (Table 15).

From the chromatographic behaviour of Planavin and its substituted 4-methylsulphonylaniline derivatives, it could be seen that the propylation of a primary amine weakens its adsorption on to silica, whereas its substitution by a hydroxyl group increases the retention¹⁸⁸. The chlorination of phenol or benzoquinone also decreases their retention on a silica plate¹⁹⁰.

Yasuda's¹⁸⁹ separation of various substituted benzenes, phenols and anilines on a silica gel-zinc dust plate show that retention generally decreases in the order phenol > aniline > N-methylaniline > benzene > chlorobenzene (Fig. 20). In the

TABLE 15
SEPARATION OF PESTICIDES
Column headings as in Table 1.

<i>N</i>	<i>Compounds</i>	<i>Stationary phase</i>	<i>SSP</i> (<i>cm</i>)	<i>Solvent 1</i>	<i>t/L</i>	<i>Drying</i>	<i>Solvent 2</i>	<i>t/L</i>	<i>Ref.</i>
3	Dieldrin, <i>p,p'</i> -TDE, aldrin from interferences	Alumina, 10 × 20 cm	3	<i>n</i> -Heptane	10 cm	Air, 15 min	Acetonitrile saturated with <i>n</i> -heptane		185
10	<i>p,p'</i> -DDE + DDT analogues from PCBs	Kieselgel G-HR HN + AgNO ₃ (chromogenic agent), 20 × 20 cm, 0.25 mm, 80°C, 20 min	3	<i>n</i> -Heptane	10 cm, 15–20 min	Air, 5–10 min	<i>n</i> -Heptane-acetone (98:2)	10 cm	186
16	Group I organophosphorus pesticides	Si gel G, 20 × 20 cm, 0.25 mm, no activation if < 60% humidity; if > 60%, 110°C, 1 h	3.8 cm, bottom	Toluene	10 min	30 sec exposure to bromine vapours, chromatography of oxidation products in other directions	25% heptane in ethyl acetate	5 min	187
10	Group II organophosphorus pesticides	idem		25% Heptane in ethyl acetate	10 min	idem	Ethyl acetate	5 min	187
12	Planavin herbicide + related compounds	Si gel F ₂₅₄ (Merck), 20 × 20 cm, 0.25 mm		Hexane-ethyl acetate-THF (66:30:4) (2 ×)			Benzene-acetonitrile (3:2)		188
23	Tetryl and related compounds	Si gel G-Zn (30:0.5, w/w), 110°C, 2 h		Chloroform	55 min	Air, 10 min	Ethyl acetate-light petroleum (1:3)	55 min	189 (Fig. 20)
9	Chloranil + chlorinated derivatives of <i>p</i> -benzoquinone, hydroquinone + phenol	Reflex Si gel foils (Kavalier), UV 254 mm, 12 × 12 cm, 0.3 mm, 120°C, 60 min	1.5	Chloroform-benzene (1:6.78)	10 cm		<i>n</i> -Heptane-carbon tetrachloride (1:11.56)	10 cm	190




Fig. 19. Two-dimensional thin-layer chromatographic separation on silica gel-cellulose (2:1) of basic fraction equivalent to 2 mg of coal-tar pitch and the characterization of seven spots. Standards: (1) acridine; (2) benz[a]acridine; (3) benz[c]acridine; (4) benzo[f]quinoline; (5) benzo[h]quinoline; (6) dibenz[a,h]acridine; (7) 11H-indeno[1,2-b]quinoline. The plate was sprayed with trifluoroacetic acid fumes and the fluorescence colours were marked. B = blue; Br = brown; G = green; l = light; P = purple; Pk = pink; R = red; Y = yellow; O = orange; V = violet; - = no fluorescence; so located by the quenching of the fluorescent plate. Reproduced from ref 177 with permission.

second direction, differentiation was further made between *meta*-, *ortho*- and *para*-isomers. The elution order may appear surprising at first, as *m*-nitroaniline, a 35-fold stronger base than *p*-nitroaniline, is less retained on the acidic silica phase. Similarly, 3,5-dinitroaniline is less retained than 2,4-dinitroaniline. This may be attributed to the effect of zinc dust in the sorbent mixture.

Gardner's¹⁸⁷ assay of organophosphorus pesticides is one of the few that uses solute derivatization prior to the second plate development. This was accomplished in order to obtain, in the second direction, a widely different migration pattern from the first. The resulting spots are then more separated, which allows better identification of multiple pesticide residues. Thus, after the first development, the plate was exposed to bromine vapour, which converted the thionate and dithioate pesticides into their corresponding oxygen analogues. The plate was then chromatographed with a more polar solvent in a direction perpendicular to the first one.




Fig. 20. Two-dimensional separation of tetryl and related compounds on a silica gel-G-zinc plate (30:0.5, w/w). 1 = 1-Chloro-2,4,6-trinitrobenzene (picryl chloride); 2 = N,N-dimethylaniline; 3 = 1,3,5-trinitrobenzene (TNB); 4 = 2,4,6-trinitroanisole; 5 = 1-chloro-2,4-dinitrobenzene; 6 = N-methylaniline; 7 = N-methyl-2,6-dinitroaniline; 8 = 2,4,6-trinitroaniline (picramide); 9 = N,2,4,6-tetranitroaniline; 10 = N-methyl-2,4,6-trinitroaniline; 11 = *o*-nitroaniline; 12 = 1,3-dinitrobenzene (reference compound); 13 = N-methyl-N,2,4,6-tetranitroaniline (tetryl); 14 = 1-chloro-2,6-dinitrobenzene; 15 = 3,5-dinitroaniline; 16 = *m*-nitroaniline; 17 = 2,4-dinitroanisole; 18 = N-methyl-2,4-dinitroaniline; 19 = 2,4-dinitroaniline; 20 = *p*-nitroaniline; 21 = N-methyl-N,2,4-trinitroaniline; 22 = 2,4-dinitrophenol; 23 = 2,4,6-trinitrophenol (picric acid). Solvent I, chloroform; solvent II, ethyl acetate-light petroleum (1:3). Reproduced from ref. 189 with permission.

4.10. Inorganic compounds

Perhaps the most important application of the two-dimensional separation of inorganic compounds is shown in Fig. 21, where polyphosphates are separated according to both their type and their chain length. This is achieved by combining an alkaline and an acidic solvent¹⁹¹.

In the separation of metal ions, two possibilities have been explored, as follows.

(1) The separation is effected using different kinds of complex formation in two directions. This is only feasible, however, if complexant I is volatile or does not interfere with complexant II. A typical example of the latter is shown in Fig. 22. The first solvent (methanol-ethanol-2 *N* hydrochloric acid) separates metal ions mainly according to hydration differences and complexation with hydrochloric acid at low acid concentrations. The second solvent separates according to the extraction of complexes at high hydrochloric acid concentrations as well as forming complexes with hydrofluoric acid [mainly for Zr(IV) and Nb(V)]¹⁹².




Fig. 21. Two-dimensional separation of oxyacids of phosphorus on paper. Alkaline solvent 1 (Ebel)-acidic solvent 3 (Grunze and Thilo). 1, orthophosphate; 2, pyrophosphate; 3, triphosphate; 4, tetrphosphate; 5, pentaphosphate; 6, hexaphosphate; 7, heptaphosphate; 8, octaphosphate; 9, Graham's salt; 10, trimetaphosphate; 11, tetrametaphosphate; 12, hypophosphite; 13, phosphite; 14, hypophosphate. Reproduced from ref. 191 with permission.

(2) It is also possible to separate in one direction by partition chromatography (butanol-2 *N* hydrochloric acid) and in the other by ion exchange (1 *N* aqueous magnesium chloride) on phosphorylated cellulose paper (Fig. 23). In the presence of hydrochloric acid (in the first dimension) the ion-exchange groups remain practically unionized and the metal ions move as on ordinary cellulose paper. The butanol-hydrochloric acid solvent cannot be removed entirely by evaporation and the remaining hydrochloric acid is neutralized with ammonia vapour¹⁹³.

4.11. Miscellaneous

Several reports have discussed the two-dimensional chromatography of organic acids on paper^{5-7,194} and cellulose¹⁹⁵, of indole derivatives^{196,197} and catechola-




Fig. 22. A two-dimensional chart showing the distribution of the major fission products when chromatographed first with methanol-ethanol-2 *N* HCl (30:30:40) and then with butanol-conc. HCl-water-conc. HF (100:50:48:2). Reproduced from ref. 192 with permission.




Fig. 23. Two-dimensional separation of metal ions on a 14-in. square of phosphorylated cellulose paper containing dihydrogen phosphate groups in the monoammonium salt form. Development is by the downward flow of the organic phase of a mixture of *n*-butanol saturated with 2 *N* HCl (partition chromatography) followed by 1 *N* magnesium chloride (ion-exchange chromatography). After the first run, the sheet is dried over ammonia to neutralize excess acid and to convert the paper to the original form. Reproduced from ref. 193 with permission.

TABLE 16
SEPARATION OF MISCELLANEOUS COMPOUNDS
Column headings as in Table 1.

<i>N</i>	<i>Compounds</i>	<i>Stationary phase</i>	<i>SSP</i> (<i>in.</i>)	<i>Solvent 1</i>	<i>t/L</i>	<i>Drying</i>	<i>Solvent 2</i>	<i>t/L</i>	<i>Ref.</i>
12	Organic acids	Paper (Durieux 122), 45 × 45 cm		Isopropanol- ammonia-water (70:5:25)	12 h, 35-40 cm	Hood	1-Propanol- eucalyptol-formic acid (50:50:20) + water until turbid	5 h, 35-40 cm	5
21	Organic acids	Paper (Whatman No. 1), 40 × 50 cm		Ethanol-ammonia- water (80:5:5)	14 h	8 h	1-Propanol- eucalyptol-formic acid (50:50:20) + water until turbid	14 h	6,7
14	Organic acids	Paper (Whatman 3MM)	3	<i>sec.</i> -Butanol- <i>n</i> - propanol-90% formic acid-water (75:25:100:48); after standing, use upper layer for development while lower layer saturates tank for 48 h prior to run	7 h	Dry, 12 h, wash with acetic acid, dry	<i>n</i> -Propanol- ammonia-water (1000:1.25:2.00)	9 h	194
8	Oxy acids	Cellulose MN 300 HR, 20 × 20 cm, 0.35 mm		Isopropanol-ethyl acetate-water (23.5:65:11.5)	14.5 cm		Formic acid-ethyl acetate-water (1:3:1)	14.5 cm	195
6	Indole derivatives	Si gel G 22, 20 × 10 cm, 0.3 mm, 20°C, 12 h		Isopropanol-25%- ammonia-water (20:1:2)			Butanol-acetic acid- water (15:3:5)		196
2	Indoleacetic and in- dolecarboxylic acids	Si gel G (Merck), 20 × 20 cm, 0.3 mm, 105°C, 20 min		Methyl acetate- isopropanol-ammonia (45:35:20)			Chloroform-acetic acid (95:5)		197

(Continued on p. 184)

TABLE 16 (continued)
SEPARATION OF MISCELLANEOUS COMPOUNDS

Column headings as in Table I.

<i>N</i>	<i>Compounds</i>	<i>Stationary phase</i>	<i>SSP</i> (<i>in.</i>)	<i>Solvent 1</i>	<i>t/L</i>	<i>Drying</i>	<i>Solvent 2</i>	<i>t/L</i>	<i>Ref.</i>
5	Catecholamine metabolites	18% cellulose, 10 × 10 cm, 0.4 mm		Isopropanol-5% ammonia (4:1)	2 h		Benzene-acetic acid-water (146:73:2)	45 min	198
11	Dansylated derivatives of brain catecholamines and their derivatives	Kieselgel G		(a) Diisopropyl ether (b) Butyl acetate-triethylamine (100:20) (2 ×)			Triethylamine-diisopropyl ether (100:20) (2 ×)		120
11	N-Acetylated derivatives of biogenic amines	Kiesel G		(a) Ethyl acetate-butyl acetate (100:20) (b) Benzene-methanol (90:10) (2 ×)			Chloroform-triethylamine (100:20)	13 cm	120
20	Hypnotics (15 barbiturates, 3 bromoutides, phenytoin + glutethimide)	Si gel GF ₂₅₄ , 20 × 20 cm, 0.25 mm, 105°C, 30 min		Chloroform-diethyl ether (75:25)	10 cm		Isopropanol-chloroform-25% ammonia (45:45:10)	10 cm	199
3	Streptomycins	Si gel (Merck), 0.03 mm, 110°C, 60 min	5.0 cm away from cathode	2% Piperidine + 2% <i>p</i> -toluenesulphonic acid in water-saturated <i>n</i> -butanol	16 cm, 3.5 h		Electrophoresis with 1% Na ₂ B ₄ O ₇ solution (pH 9.6), 300 V, 23-28°C	3 h	200
11	1,4-Benzodiazepine drugs (blood, urine)	Si gel G, 20 × 20 cm, 0.25 mm, 110°C, 30 min	1.5	Chloroform-benzene-diethyl ether (20:10:70)	10 cm		Benzene-acetone (90:10)		201, 202
15	Chlorpromazine and metabolites	Si gel (Brinkmann 254F), 0.25 mm, 100°C, 4 h	2	Acetone-methanol-methanolamine (100:30:1)	12 cm	42°C, 10 min	Isopropanol-ethanol-ammonia (2:1:1)	12 cm	203
14	Phthalodinitriles and related metabolites	(1) Si gel HF-cellulose (1:1) on 4 × 20 cm strip (2) Acetylated cellulose F layer, 15 × 20 cm		On (1): Carbon tetrachloride-ethyl acetate-acetic acid (7:2:0.5)	15 cm	Air, 20 min	On (2): Methanol-diethyl ether-water (7:2:0.5)	10 cm	204

4	Aflatoxins b ₁ , b ₂ , g ₁ , g ₂ (figs)	Si gel 60, 105°C, 1 h	Toluene-ethyl acetate-90% formic acid (5:4:1)	17 cm	Dry, dark, room temp., 15 min	Acetone-chloroform (1:9)	17 cm	205
4	Aflatoxins (corn)	SILG-HR-25 (Brinkmann), 10 × 5 cm	Acetone-chloroform (12:88)	8 cm	Air stream	95% Denatured ethanol (Mallinckrodt 7006)	4.5 cm	206
37	Milk constituents from which aflatoxin M ₁ isolated	Si gel 60 (Merck), 20 × 20 cm, 0.2 mm	Chloroform-acetone-2-propanol (80:15:15)	10 cm		Toluene-ethyl acetate-90% formic acid (60:30:10)	10 cm	207
12	Carbonyl dinitrophenylhydrazones	MgO-Microcell T38 (1:1), 60°C, 1 h, 20 × 20 cm	Light petroleum-chloroform (85:35)	12 cm		(a) Chloroform-SVR (80:20) (2 ×) (b) 20% Carbowax 400 in chloroform (impregnates plates) (c) Rotate plate 180° and develop in PE	208 (Fig. 24)	
5	Mixture of Ni(II) bis-diethyl- and bisdiethyl-dithiocarbamate complexes	Silufol Si gel (Kavaliel), 15 × 15 cm, 120°C, 1 h	Chloroform-cyclohexane (3:4)	20 min		Chloroform-cyclohexane (3:4)	209	
7	B ₆ vitamins	Si gel H, 10 × 15 cm, 0.25 mm	Isoamyl alcohol-acetone-water-diet-hylamine (24:18:6:8)		Warm air	2-Butanol-ethanol-1.5 N ammonia-isoamyl alcohol-diethylamine (20:6:7:2:2)	210	
12	1,3,5-Triamino-2,4,6-trinitrobenzene and its impurities	30 g Si gel G + 1 g Zn dust, 20 × 20 cm, 110°C, 2 h	1,2-Dichloroethane	1 h	10 min	Acetone-light petroleum (1:9)	211	




Fig. 24. Diagrammatic representation of the two-dimensional separation of dinitrophenylhydrazones (direction of solvent travel indicated by arrow). (a) Stage 1, class separation; (b) stage 2, first concentration; (c) stage 3, second concentration; (d) stages 4 and 5, impregnation and marker application; (e) stage 6, chain length separation. K = ketones; A = anals; E = enals; D = dienals. Reproduced from ref. 208 with permission.

mine metabolites^{120,198} on silica and cellulose, of barbiturates¹⁹⁹, streptomycins²⁰⁰, benzodiazepine^{201,202} and chlorpromazine²⁰³ drugs on silica and of phthalodinitriles²⁰⁴ on a mixed plate, where a silica-cellulose strip had been coated at the side of an acetylated cellulose layer. In addition, aflatoxins^{205–207}, carbonyldinitrophenylhydrazones²⁰⁸, dithiocarbamate complexes²⁰⁹ and some B₆ vitamers²¹⁰ have also been resolved by two-dimensional TLC. Detailed chromatographic conditions for these separations are given in Table 16.

Craske and Edwards²⁰⁸ developed an original two-dimensional technique for the separation of carbonyl dinitrophenylhydrazones (Fig. 24). Solutes were first resolved into different classes (alkanones, alkanals, alk-2-enals and alka-2,4-dienals) by adsorption, and next on the basis of chain length by partition. Whereas initial class separation required a small sample to prevent overloading, secondary chain length separation following Carbowax 400 impregnation required a large sample to facilitate solute detection. This problem was solved by streaking the sample across the width of the plate, developing in the first solvent and subsequently performing a chromatographic concentration step. Orthogonal elution in chloroform–SVR (80:20) concentrated the spots on one side of the plate. The layer was then impregnated with 20% Carbowax 400 in chloroform. The plate was rotated through 180° and chain resolution was effected by development in light petroleum.

5. CONCLUSION

In summary, two-dimensional TLC applications may be achieved using:

(1) a one-sorbent layer and solvents with different selectivity for the various solutes;

(2) a mixed-sorbent layer and eluents chosen such that one particular sorbent dictates retention in a given direction;

(3) coupled layers where two different sorbents are coated side-by-side, each effecting a separation according to a different retention mechanism;

(4) modification of the plate (impregnation) prior to the second development;

(5) derivatization of the solute prior to the second development;

(6) coupled separation techniques (TLC-GC, TLC-electrophoresis).

Sample purification may be achieved in one direction and separation in the other. The main advantage of the technique lies in its efficiency, as a large number of spots may be resolved on the chromatographic plate²¹, provided that the initial spot size is small and the sample is not overloaded. In addition, widely different components may be resolved from a given mixture.

Perhaps the main disadvantages of a two-dimensional system are the serious difficulties encountered in quantitating a significant number of spots, the difficulties of predicting R_F values obtained from the second plate development, as these are generally affected by the previous elution, and the possible irregular ascent of the second solvent, leading to spot distortion. It is possible to correct for the latter problem, if not by selecting more "miscible" solvent systems, then by grafting two thin layers and transferring spots from one to the other prior to the second development²¹². As for the prediction of R_F values, a blank plate could be developed in the first eluent prior to spotting the various reference solutions and chromatography in the second direction.

Thus the problem of measuring compound concentrations remains the most critical, at least when working with non-radioactively labelled material or with compounds that cannot all react with a fluorescent or similar tag.

Nevertheless, this method is extremely powerful and rapid. With a total analysis time of about 1 h, a spot capacity well exceeding those currently achieved in conventional TLC or even HPLC can easily be obtained. The spot capacity of the figures reproduced above is in the range 150–300 (except in Fig. 15). We are of the opinion that this method or more generally bi- and tri-directional chromatographic techniques have considerable potential for the separation of complex mixtures; when there are a few ten to a hundred components to resolve, their use may save a considerable time in the selection of a convenient solvent mixture and/or chromatographic system allowing the complete separation in one HPLC run. If the mixture is more complex (several hundred components) they offer the only possibility of a simple, total analysis.

Accordingly, instrumentation for two-dimensional chromatography will be the subject of intense development work in the near future, and solutions will eventually be found for the problems discussed above.

6. SUMMARY

In view of the large spot capacity available, much larger than in conventional thin-layer chromatography (TLC), two-dimensional thin-layer chromatography is an extremely powerful and rapid separation technique. This paper presents a review of the literature up to early 1982 on the analytical applications of two-dimensional TLC classified according to the chemical nature of the compounds separated. Technical aspects of the method (single, mixed or coupled sorbent layers, plate impregnation, solute derivatization, coupling with other analytical techniques), together with effective mechanisms of solute retention are discussed. Available methods for spot detection and quantitation are reported. Future trends of two-dimensional TLC are commented. Over 200 references are cited.

REFERENCES

- 1 I. M. Hais, *J. Chromatogr.*, 187 (1980) 466.
- 2 G. Guiochon, A. Siouffi, H. Engelhardt and I. Halasz, *J. Chromatogr. Sci.*, 16 (1978) 152.
- 3 R. Consden, A. H. Gordon and A. J. P. Martin, *J. Biochem.*, 38 (1944) 224.
- 4 B. T. Overell, *Austr. J. Sci.*, (1952) 28.
- 5 R.-I. Cheftel, R. Munier and M. Macheboeuf, *Bull. Soc. Chim. Biol.*, 34 (1952) 380.
- 6 R. Nordmann, O. Gauchery, J.-P. du Ruisseau, Y. Thomas and J. Nordmann, *Bull. Soc. Chim. Biol.*, 36 (1954) 1461.
- 7 R. Nordmann, O. Gauchery, J.-P. du Ruisseau, Y. Thomas and J. Nordmann, *Bull. Soc. Chim. Biol.*, 36 (1954) 1641.
- 8 F. Pocchiari and C. Rossi, *Chromatogr. Rev.*, 4 (1961) 1.
- 9 M. Calvin, *Harvey Lect.*, 46 (1950-51) 218.
- 10 M. Calvin, *Proc. 3rd Int. Congr. Biochem., Brussels, 1955*, p. 211.
- 11 A. Beloff-Chain, R. Catanzaro, E. B. Chain, I. Masi, F. Pocchiari and C. Rossi, *Proc. Roy. Soc (London), Ser. B*, 143 (1955) 481.
- 12 A. Beloff-Chain, R. Catanzaro, E. B. Chain, I. Masi and F. Pocchiari, *Sel. Sci. Pap. Ist. Super. Sanità*, 2 (1959) 109.
- 13 E. B. Chain, A. Beloff-Chain and F. Pocchiari, *Sel. Sci. Pap. Ist. Super. Sanità*, 1 (1956) 389.
- 14 E. B. Chain, *Brit. Med. J.*, 2 (1959) 709.
- 15 H. J. Issaq and E. W. Barr, *Anal. Chem.*, 49 (1977) 83A.
- 16 E. Stahl, *J. Chromatogr.*, 165 (1979) 59.
- 17 J. A. Perry, *J. Chromatogr.*, 165 (1979) 117.
- 18 G. Zweig and J. Sherma, *Anal. Chem.*, 52 (1980) 276R.
- 19 H. J. Issaq, *Separ. Purif. Methods*, 10 (1981) 73.
- 20 D. C. Fenimore and C. M. Davis, *Anal. Chem.*, 53 (1981) 252A.
- 21 G. Guiochon, M. F. Gonnord, A. Siouffi and M. Zakaria, *J. Chromatogr.*, 250 (1982) 1.
- 22 G. Guiochon, L. A. Beaver, M. F. Gonnord, A. Siouffi and M. Zakaria, *J. Chromatogr.*, 255 (1983) 415.
- 23 U. A. Th. Brinkman, G. Devries and R. Cuperus, *J. Chromatogr.*, 198 (1980) 421.
- 24 R. Delley and G. Szekely, *Chimia*, 32 (1978) 261.
- 25 L. R. Snyder, in E. Heftman (Editor), *Chromatography: A Laboratory Handbook of Chromatographic and Electrophoresis Methods*, Van Nostrand Reinhold, New York, 3rd ed., 1975, pp. 46-76.
- 26 O. Motl and L. Novotny, in O. Mikès (Editor), *Laboratory Handbook of Chromatographic and Allied Methods*, Ellis Horwood, Chichester, 1979, pp. 150-217.
- 27 K. Randerath, *Chromatographie sur Couches Minces*, Gauthier-Villars, Paris, 2nd ed., 1971, p. 22.
- 28 R. K. Gilpin and W. R. Sisco, *J. Chromatogr.*, 124 (1976) 257.
- 29 M. Ericsson and L. G. Blomberg, *J. High Resolut. Chromatogr. Chromatogr. Commun.*, 3 (1980) 345.
- 30 I. D. Wilson, S. Scalia and E. D. Morgan, *J. Chromatogr.*, 212 (1981) 211.
- 31 H. Halpaap, K.-F. Krebs and H. E. Hauck, *J. High Resolut. Chromatogr. Chromatogr. Commun.*, 3 (1980) 215.

- 32 O. Mikš, J. Štamberg, M. Hejtmánek and K. Šebesta, in O. Mikš (Editor), *Laboratory Handbook of Chromatographic and Allied Methods*, Ellis Horwood, Chichester, 1979, pp. 218-327.
- 33 O. Motl and L. Novotny, in O. Mikš (Editor), *Laboratory Handbook of Chromatographic and Allied Methods*, Ellis Horwood, Chichester, 1979, pp. 462-523.
- 34 L. Lepri, P. G. Desideri and D. Heimler, *J. Chromatogr.*, 235 (1982) 411.
- 35 K. Randerath and E. Randerath, *J. Chromatogr.*, 16 (1964) 111.
- 36 P. S. Bond, *J. Chromatogr.*, 34 (1968) 554.
- 37 J. Pore, J. P. Houis and I. Rasori, *Rev. Fr. Corps Gras*, 72 (1981) 164.
- 38 A. R. Ritchie, S. R. B. Souy and M. H. Clear, *J. Chromatogr.*, 150 (1978) 557.
- 39 Y. V. Zanevskit, A. B. Ivanov, V. D. Peshekhonov, I. A. Tyapkin, S. P. Chernenko, L. B. Kaminir, E. P. Senchenkov and V. N. Kalinin, *Report 1977, JINR-R-14-10934, C.A.*, 90 (1979) 114557a.
- 40 R. W. Frei, *Progr. Thin-Layer Chromatogr. Relat. Methods*, 2 (1971) 1.
- 41 V. Pollak, *J. Chromatogr.*, 152 (1978) 201.
- 42 J. Goldman and R. R. Goodall, *J. Chromatogr.*, 32 (1968) 24.
- 43 J. C. Touchstone, S. S. Levin and T. Murawec, in J. C. Touchstone (Editor), *Quantitative Thin-Layer Chromatography*, Wiley-Interscience, New York, 1979, pp. 1-22.
- 44 P. Knapstein, J. C. Touchstone, P. Menzel and G. W. Oertel, *J. Chromatogr.*, 44 (1969) 190.
- 45 V. Pollak, *J. Liq. Chromatogr.*, 3 (1980) 1881.
- 46 G. Kerenyi, T. Pataki and T. Devenyi, *Hung. Pat.*, 170-287 (1976).
- 47 T. Devenyi and S. Pungor, *Hung. Sci. Instrum.*, 50 (1980) 3.
- 48 N. L. Anderson, J. Taylor, A. E. Scandora, B. P. Coulter and N. G. Anderson, *Clin. Chem.*, 27 (1981) 1807.
- 49 E. P. Lester, P. F. Lemkin and L. E. Lipkin, *Anal. Chem.*, 53 (1981) 390A.
- 50 A. K. Thorsrud, H. F. Haugen and E. Jellum, *Electrophoresis '79, Advan. Methods Biochem. Clin. Appl., Proc. Int. Conf. Electrophoresis*, Walter de Gruyter, Berlin, 1980, pp. 426-435.
- 51 M. L. Gianelli, J. B. Callis, N. H. Anderson and G. D. Christian, *Anal. Chem.*, 53 (1981) 1357.
- 52 L. A. Beaver and G. Guiochon, U.S. patent pending.
- 53 G. Guiochon, L. A. Beaver, M. F. Gonnord, A. M. Siouffi and M. Zakaria, *Chromatographia*, 17 (1983) 21.
- 54 C. V. Viswanathan, F. Phillips and W. O. Lundberg, *J. Chromatogr.*, 38 (1968) 267.
- 55 J. M. Bailey, R. W. Bryant, S. J. Feinmark and A. N. Makheja, *Prostaglandins*, 13 (1977) 479.
- 56 D. C. Palmer, J. A. Kintzios and N. M. Papadopoulos, *J. Chromatogr. Sci.*, 10 (1972) 107.
- 57 T. M. Lee and A. der Marderosian, *J. Pharm. Sci.*, 70 (1981) 89.
- 58 B. A. El-Zeany and A.-K.-S. Ahmed, *Egypt. J. Food Sci.*, 1-2 (1977) 1.
- 59 H. H. O. Schmid, W. J. Baumann, J. M. Cubero and H. K. Mangold, *Biochim. Biophys. Acta*, 125 (1966) 189.
- 60 T. Takatori and A. Yamaoka, *Nippon Hoigaku Zasshi (Jap. J. Legal Med.)*, 30 (1976) 368.
- 61 C. Bouillé, J. Y. Daniel, J. Boissin and I. Assenmacher, *Steroids*, 14 (1969) 7.
- 62 D. C. Bicknell and D. B. Gower, *J. Chromatogr.*, 61 (1971) 358.
- 63 T. Taylor, *J. Chromatogr.*, 66 (1972) 177.
- 64 G. Cavina, G. Moretti and M. Petrella, *J. Chromatogr.*, 103 (1975) 368.
- 65 H. L. Issaq, *J. Liq. Chromatogr.*, 3 (1980) 841.
- 66 R. E. Levitt and J. C. Touchstone, *J. High Resolut. Chromatogr. Chromatogr. Commun.*, 2 (1979) 587.
- 67 D. W. Vidrine and H. J. Nicholas, *J. Chromatogr.*, 89 (1974) 92.
- 68 H.-Ch. Curtius and M. Müller, *J. Chromatogr.*, 32 (1968) 222.
- 69 M. J. Whittle, A. I. Wilson, C. R. Whitfield, R. D. Paton and R. W. Logan, *Brit. Med. J.*, 282 (1981) 428.
- 70 T. L. Gross, M. V. Wilson, P. M. Kuhnert and R. J. Sokol, *Clin. Chem.*, 27 (1981) 486.
- 71 G. Rouser, S. Fleischer and A. Yamamoto, *Lipids*, 5 (1970) 494.
- 72 R. M. Broekhuysse, *Clin. Chim. Acta*, 23 (1969) 457.
- 73 G. S. Getz, S. Jakovcic, J. Heywood, J. Frank and M. Rabinowitz, *Biochim. Biophys. Acta*, 218 (1970) 441.
- 74 S. K. Jain and D. Subrahmanyam, *Ital. J. Biochem.*, 27 (1978) 11.
- 75 B. J. H. M. Poorthuis, P. J. Yazaki and K. Y. Hostettler, *J. Lipid Res.*, 17 (1976) 433.
- 76 K. G. Blass, R. L. Briand, D. S. Ng and S. Harold, *J. Chromatogr.*, 182 (1980) 311.
- 77 S. Pollet, S. Ermidou, F. Le Saux, M. Monge and N. Baumann, *J. Lipid Res.*, 19 (1978) 916.

- 78 S. S. Radwan, *J. Chromatogr. Sci.*, 16 (1978) 538.
79 S. Ikawa and M. Goto, *J. Chromatogr.*, 114 (1975) 237.
80 F.-H. Hubmann, *J. Chromatogr.*, 86 (1973) 197.
81 M. Ohashi, *Lipids*, 14 (1979) 52.
82 R. W. Ledeen, J. E. Haley and J. A. Skrivanek, *Anal. Biochem.*, 112 (1981) 135.
83 G. D. Hunter, V. M. Wiegant and A. J. Dunn, *J. Neurochem.*, 37 (1981) 1025.
84 C. J. Clarke and P. H. Cobb, *J. Chromatogr.*, 168 (1979) 541.
85 J. Metivier and A. M. Viana, *J. Exp. Bot.*, 30 (1979) 805.
86 R. E. Knowles and A. L. Livingston, *J. Chromatogr.*, 61 (1971) 133.
87 A. S. L. Tirimanna, *Mikrochim. Acta*, 2 (1981) 11.
88 J. Sherma and G. Zweig, *J. Chromatogr.*, 31 (1967) 589.
89 G. H. Elder, *J. Chromatogr.*, 59 (1971) 234.
90 K. Saitoh, M. Kobayashi and N. Suzuki, *Anal. Chem.*, 53 (1981) 2309.
91 Y. Onoue, *Bull. Chem. Soc. Jap.*, 54 (1981) 1972.
92 D. B. Mullick, *J. Chromatogr.*, 39 (1969) 291.
93 J. Aurenge, G. Barbe-Richaud and L. Gelpi, *J. Chromatogr.*, 22 (1966) 369.
94 G. Nadeau, G. Sobolewski, L. Fiset and C. G. Farmilo, *J. Chromatogr.*, 1 (1958) 327.
95 G. B. Marini-Bettolo and G. C. Casinovi, *J. Chromatogr.*, 1 (1958) 411.
96 R. L. Munier and S. Meunier, *Chromatographia*, 13 (1980) 259.
97 A. Viala and M. Estadieu, *J. Chromatogr.*, 72 (1972) 127.
98 P. Le Xuan, R. L. Munier and S. Meunier, *Chromatographia*, 13 (1980) 693.
99 S. Agurell, *Acta Pharm. Suecia*, 2 (1965) 357.
100 A. J. P. Martin and R. L. M. Synge, *Anal. Chem. Proteins*, 1 (1945) 1.
101 W. E. Cohn, *J. Amer. Chem. Soc.*, 72 (1950) 1471.
102 C. S. Knight, *Nature (London)*, 188 (1960) 739.
103 M. C. Corfield and E. C. Simpson, *J. Chromatogr.*, 17 (1965) 420.
104 J. H. Buchanan and M. C. Corfield, *J. Chromatogr.*, 31 (1967) 274.
105 D. Horton, A. Tanimura and M. L. Wolfram, *J. Chromatogr.*, 23 (1966) 309.
106 J. Maugenet and M.-L. Dalmeyda, *Ann. Technol. Agr.*, 18 (1969) 237.
107 F. Kraficzky, R. Helger and H. Lang, *Clin. Chem.*, 16 (1970) 662.
108 R. L. Munier, B. Faivre and A. Lebreau, *Chromatographia*, 6 (1973) 71.
109 R. L. Munier, C. Gervais and A.-M. Drapier, *C.R. Acad. Sci., Ser. D*, 281 (1975) 1513.
110 R. L. Munier, C. Gervais and A.-M. Drapier, *C.R. Acad. Sci., Ser. D*, 281 (1975) 1757.
111 J. J. Harper and G. H. G. Davis, *Int. J. Syst. Bacteriol.*, 29 (1979) 56.
112 M. M. Frodyma and R. W. Frei, *J. Chromatogr.*, 17 (1965) 131.
113 K. Chmel, *Collect. Czech. Chem. Commun.*, 37 (1972) 3034.
114 C. G. Honegger, *Helv. Chim. Acta*, 44 (1961) 173.
115 E. Von Arx and R. Neher, *J. Chromatogr.*, 12 (1963) 329.
116 N. Seiler, *Methods Biochem. Anal.*, 18 (1970) 259.
117 P. Fernlund and L. Joseffson, *Biochim. Biophys. Acta*, 158 (1968) 262.
118 M. Weise and D. E. Oken, *J. Chromatogr.*, 152 (1978) 175.
119 D. Biou, N. Queyrel, M. N. Visseaux, I. Collignon and M. Pays, *J. Chromatogr., Biomed. Appl.*, 226 (1981) 477.
120 N. Seiler and M. Wiechmann, *J. Chromatogr.*, 28 (1967) 351.
121 K. Macek, Z. Deyl and M. Smrř, *J. Chromatogr.*, 193 (1980) 421.
122 K.-T. Wang, J. M. K. Huang and I. S. Y. Wang, *J. Chromatogr.*, 22 (1966) 362.
123 R. L. Munier and G. Sarrazin, *J. Chromatogr.*, 22 (1966) 347.
124 R. L. Munier and A. M. Drapier, *Chromatographia*, 5 (1972) 306.
125 R. L. Munier, B. Faivre and A. Lebreau, *Chromatographia*, 6 (1973) 14.
126 I. M. Wainwright and P. Shapshak, *J. Chromatogr. Sci.*, 17 (1979) 535.
127 P. H. O'Farrel, *J. Biol. Chem.*, 250 (1975) 4007.
128 N. G. Anderson and N. L. Anderson, *Anal. Biochem.*, 85 (1978) 331.
129 N. L. Anderson and N. G. Anderson, *Anal. Biochem.*, 85 (1978) 341.
130 N. L. Anderson and N. G. Anderson, *Biochem. Biophys. Res. Commun.*, 88 (1979) 258.
131 N. G. Anderson, N. L. Anderson, S. L. Tollaksen, H. Hahn, F. Giere and J. Edwards, *Anal. Biochem.*, 95 (1979) 48.
132 N. L. Anderson and B. J. Hickman, *Anal. Biochem.*, 93 (1979) 312.

- 133 K. E. Willard, C. S. Giometti, N. L. Anderson, T. E. O'Connor and N. G. Anderson, *Anal. Biochem.*, 100 (1979) 289.
- 134 B. J. Hickman, N. L. Anderson, K. E. Willard and N. G. Anderson, *Electrophoresis '79, Advan. Methods Biochem. Clin. Appl., Proc. Int. Conf. Electrophoresis*, Walter de Gruyter, Berlin, 1980, p. 341.
- 135 N. L. Anderson, *Immunol. Lett.*, 2 (1981) 195.
- 136 T. Pearson and L. Anderson, *Anal. Biochem.*, 101 (1980) 377.
- 137 C. S. Giometti, N. G. Anderson and N. L. Anderson, *Clin. Chem.*, 25 (1979) 1877.
- 138 C. S. Giometti, N. G. Anderson, S. L. Tollaksen, J. J. Edwards and N. L. Anderson, *Anal. Biochem.*, 102 (1980) 47.
- 139 G. Volden, A. K. Thorsrud, I. Bjørson and E. Jellum, *J. Invest. Dermatol.*, 75 (1980) 421.
- 140 E. Jellum, *J. Chromatogr.*, 239 (1982) 29.
- 141 S. M. Petrović and V. D. Canić, *Mikrochim. Acta*, (1969) 599.
- 142 E. Moczar and M. Moczar, *Bull. Soc. Chim. Biol.*, 49 (1967) 1159.
- 143 K. Hotta and M. Kurokawa, *Anal. Biochem.*, 26 (1968) 472.
- 144 T. Mezzetti, M. Ghebregziabher, S. Rufini, G. Ciuffini and M. Lato, *J. Chromatogr.*, 74 (1972) 273.
- 145 M. Ghebregziabher, S. Rufini, G. Ciuffini and M. Lato, *J. Chromatogr.*, 95 (1974) 51.
- 146 M. Ghebregziabher, S. Rufini, G. M. Sapia and M. Lato, *J. Chromatogr.*, 180 (1979) 1.
- 147 E. A. Ryan and A. M. Kropinski, *J. Chromatogr.*, 195 (1980) 127.
- 148 E. Moczar, *J. Chromatogr., Biomed. Appl.*, 146 (1978) 337.
- 149 R. L. Wolf and J. Andreoni, *Cell. Immunol.*, 67 (1982) 299.
- 150 E. Randerath and K. Randerath, *J. Chromatogr.*, 16 (1964) 126.
- 151 G. Santini and V. Ulrich, *J. Chromatogr.*, 49 (1970) 560.
- 152 R. A. Barton, A. Schulman, E. L. Jacobson and M. K. Jacobson, *J. Chromatogr.*, 130 (1977) 145.
- 153 R. L. Bielecki, *Anal. Biochem.*, 12 (1965) 230.
- 154 C. V. Cole and C. Ross, *Anal. Biochem.*, 17 (1966) 526.
- 155 D. S. S. Kumar, L. Pinck and L. Hirth, *Anal. Biochem.*, 48 (1972) 497.
- 156 S. Hashimoto, W. S. M. Wold, K. H. Brackmann and M. Green, *Biochemistry*, 20 (1981) 6640.
- 157 A. H. van Gennip, D. Y. van Noordenburg-Huistra, P. K. de Bree and S. K. Wadman, *Clin. Chim. Acta*, 86 (1978) 7.
- 158 G. Pataki, *J. Chromatogr.*, 29 (1967) 126.
- 159 S. Seipel and U. Reichert, *J. Chromatogr.*, 135 (1977) 485.
- 160 E. Randerath and K. Randerath, *J. Chromatogr.*, 16 (1964) 111.
- 161 T. W. Munns, K. C. Podratz and P. A. Katzman, *J. Chromatogr.*, 76 (1973) 401.
- 162 T. W. Munns and H. F. Sims, *J. Chromatogr.*, 111 (1975) 403.
- 163 I. Das, *Biochem. Soc. Transactions, 576th Meet., London*, 6 (1978) 1047.
- 164 J. S. Miller and R. R. Burgess, *Biochemistry*, 17 (1978) 2054.
- 165 A. D. Mirzabekov and B. E. Griffin, *J. Mol. Biol.*, 72 (1972) 633.
- 166 H. J. Gross, H. Kroath, H.-G. Janda and C. Jungwirth, *Mol. Biol. Rep.*, 4 (1978) 105.
- 167 P. L. Bergquist, *J. Chromatogr.*, 19 (1965) 615.
- 168 L. Diels and R. DeWachter, *Arch. Int. Physiol. Biochim.*, 88 (1980) B26.
- 169 S. Hashimoto, W. S. M. Wold, K. H. Brackmann and M. Green, *Biochemistry*, 20 (1981) 6640.
- 170 E. Sawicki, T. W. Stanley and W. C. Elbert, *Mikrochim. Acta*, (1965) 1110.
- 171 E. Sawicki, T. W. Stanley, W. C. Elbert and M. Morgan, *Talanta*, 2 (1965) 605.
- 172 E. Sawicki and H. Johnson, *J. Chromatogr.*, 23 (1966) 142.
- 173 E. Sawicki, T. W. Stanley, S. McPherson and M. Morgan, *Talanta*, 13 (1966) 619.
- 174 E. Sawicki, C. R. Engel and W. C. Elbert, *Talanta*, 14 (1967) 1169.
- 175 E. Sawicki, T. W. Stanley, W. C. Elbert, J. Meeker and S. McPherson, *Atmos. Environ.*, 1 (1967) 131.
- 176 E. Sawicki, M. Guyer, R. Schumaker, W. C. Elbert and C. R. Engel, *Mikrochim. Acta*, (1968) 1025.
- 177 C. R. Engel and E. Sawicki, *J. Chromatogr.*, 31 (1967) 109.
- 178 H. Matsushita and Y. Suzuki, *Bull. Chem. Soc. Jap.*, 42 (1969) 460.
- 179 J. Kraft, A. Hartung, K.-H. Lies and J. Schulze, *J. High Resolut. Chromatogr. Chromatogr. Commun.*, 5 (1982) 482.
- 180 D. Srzić and L. Klasinc, *Recent Dev. Mass Spectrom. Biochem. Med. Environ. Res.*, 7 (1981) 195.
- 181 J. Janák, *J. Gas Chromatogr.*, 1 (1963) 20.
- 182 J. Janák, *Progr. Thin-Layer Chromatogr. Relat. Methods*, 2 (1971) 63.
- 183 R. Kaiser, *Chromatographie in der Gasphase*, Bibliographisches Institut, Mannheim, 1969.

- 184 R. E. Kaiser and R. Rieder, *J. Chromatogr.*, 142 (1977) 411.
- 185 J. A. Kawatski and D. L. Frasch, *J. Ass. Offic. Anal. Chem.*, 52 (1969) 1108.
- 186 N. V. Fehringier and J. E. Westfall, *J. Chromatogr.*, 57 (1971) 397.
- 187 A. M. Gardner, *J. Ass. Offic. Anal. Chem.*, 54 (1971) 517.
- 188 W. B. Burton, *J. Chromatogr.*, 53 (1970) 397.
- 189 S. K. Yuda, *J. Chromatogr.*, 50 (1970) 453.
- 190 P. Švec and M. Zbiasrovský, *Sb. Vys. Sk. Chem.-Technol. Praze, Org. Chem. Technol.*, 17 (1972) 95.
- 191 J. P. Ebel, *Metodi di Separazione nella Chimica Inorganica*, Vol. 1, CNR, Rome, 1963.
- 192 A. Grand-Clement, Z. Jakovac, M. Lederer and E. Pluchet, *Proceedings of International Symposium on Microchemistry, Birmingham, 1958*, Pergamon Press, Oxford, 1960, p. 231.
- 193 C. S. Knight, *Nature (London)*, 183 (1959) 165.
- 194 F. Stern, G. Grumet, F. Trabal, A. Mennis and H. H. Zinsser, *J. Chromatogr.*, 19 (1965) 130.
- 195 D. Baraldi, *J. Chromatogr.*, 42 (1969) 125.
- 196 J. Opieńska-Blauth, H. Krackowski, H. Brzus-Kiewicz and Z. Zagórski, *J. Chromatogr.*, 17 (1965) 288.
- 197 H. Takaki, *Miyazaki Daigaku Nogakubu, Kenkyu Hokoku*, 27 (1980) 211.
- 198 K. Breebart, A. M. Haan and S. K. Wadman, *Clin. Chim. Acta*, 37 (1972) 157.
- 199 R. A. de Zeeuw, *Chromatogr., Electrophor. Symp. Int., 4th*, (1968) 537.
- 200 T. Katayama and H. Ikeda, *Sci. Pap. Inst. Phys. Chem. Res.*, 63 (1969) 49.
- 201 S. N. Tewari and S. K. Shukla, *Pharmazie*, 32 (1977) 536.
- 202 S. N. Tewari and S. K. Shukla, *J. Indian Acad. Forensic Sci.*, 17 (1978) 80.
- 203 P. Turano and W. J. Turner, *J. Chromatogr.*, 64 (1972) 347.
- 204 H. Matsushita and K. Arashidani, *Bunseki Kagaku (Jap. Anal.)*, 20 (1971) 78.
- 205 L. Allen, *J. Ass. Offic. Anal. Chem.*, 57 (1974) 1398.
- 206 R. J. Alexander and M. C. Baur, *Cereal Chem.*, 54 (1977) 699.
- 207 D. S. P. Patterson, E. M. Glancy and B. A. Roberts, *Food Cosmet. Toxicol.*, 16 (1978) 49.
- 208 J. D. Craske and R. A. Edwards, *J. Chromatogr.*, 51 (1970) 237.
- 209 O. Liška, J. Lehotay, E. Brandsteterová and G. Guiochon, *J. Chromatogr.*, 171 (1979) 153.
- 210 J. D. Mahuren and S. P. Coburn, *Anal. Biochem.*, 82 (1977) 246.
- 211 S. K. Yasuda, *J. Chromatogr.*, 71 (1972) 481.
- 212 A. M. Pumphrey, *Biochem. J.*, 102 (1964) 30P.

CHREV. 166

STATISTICAL APPROACH TO CHROMATOGRAPHIC THEORY

JAMES S. FRITZ*

Ames Laboratory, Iowa State University, Ames, IA 50011 (U.S.A.)*

and

DAN M. SCOTT

Department of Operations Research, Stanford University, Stanford, CA 94305 (U.S.A.)

(First received January 27th, 1982; revised manuscript received August 13th, 1982)

CONTENTS

1. Introduction	193
2. Plate models	196
3. Discrete flow model	196
4. Continuous flow model	199
5. Theoretical plate numbers in the two models	200
6. Some consequences of the discrete flow model	201
7. Calculation of plate numbers from chromatographic data	205
7.1. Diffusion broadening, σ_d^2	205
7.2. Inter-phase broadening, σ_{ip}^2	206
8. Summary	211
References	211

1. INTRODUCTION

The theoretical plate model continues to enjoy widespread popularity as a description of the behavior of chromatographic peaks. The plate model is generally attributed to Martin and Synge¹, but has been developed further and modified by Glueckauf², Mayer and Tompkins³ and others. Giddings⁴ gave an excellent discussion of the plate theory and its limitations.

In the plate theory of chromatography, a column is divided into a number of imaginary sections called plates. In each plate it is assumed that a sample substance is partitioned between the stationary and mobile phases and that equilibrium has been attained. Then the mobile phase in each plate moves forward to the next plate, carrying the sample substance in the mobile phase forward exactly one plate. A new partitioning occurs simultaneously in each plate, and it again is assumed that equilibrium is attained. This process is repeated many times, until sample constituents move along and off the column at different rates and thus are separated.

The number of imaginary sections in a column is generally called the number of theoretical plates, or theoretical plate number, and is designated as N (or n). The numerical value of N for any given column length is maximized by using very small spheres of uniform diameter, thinly coated with a stationary phase and carefully packed, as the column stationary phase. Factors such as column dimensions and

* Operated for the U.S. Department of Energy by Iowa State University under contract No. W-7405-ENG-82. This work was supported by the Director of Energy Research, Office of Basic Energy Sciences.

fittings and proper eluent flow-rate also affect the value of N and hence the separation efficiency of the column.

There is no convenient way of measuring the intrinsic value of N ; it must be calculated from measurable parameters, namely the retention time (or volume) of a peak and the peak width (or σ , which is a function of peak width). The defining equation for this calculation is

$$N = 16 \left(\frac{t}{w} \right)^2 = \frac{t^2}{\sigma^2}$$

where t is the peak retention time, w is the peak width at its base (in the same units as t) and σ^2 is the variance of the peak width.

Some chromatographers prefer to use the number of effective theoretical plates as a measure of the separation power of a column:

$$N_{\text{eff}} = 16 \left(\frac{t - t_0}{w} \right)^2 = \frac{(t - t_0)^2}{\sigma^2}$$

where t_0 is the retention time of a non-sorbed substance, or the "dead time" of the column. As early as 1959 Purnell^{5,6} noted that N_{eff} (for which he used S , rather than N) is critical in determining the separation ability of a chromatographic column. He found that the resolution of alkanes was inconsistent with the observed value of N when the retention volumes are very small. Widely used equations show that resolution of chromatographic peaks is proportional to the number of effective theoretical plates.

The plate theory of chromatography has been criticized as lacking in physical reality because eluent flow in actual columns obviously is continuous and not a discontinuous flow from one plate to the next as portrayed by the plate theory. It is debatable whether this objection is valid because any theory that gives results consistent with experimental behavior would be useful despite differences in the actual and theoretical mechanisms.

Giddings⁴ criticized the plate model on several counts and concluded that it "fails in the most important test of all—the very practical matter of predicting zone dispersion as a function of the numerous variables open to manipulation by the investigator". These variables include the particle size of the solid support, thickness of stationary phase coating, eluent flow-rate and diffusion coefficients in the stationary and mobile phases.

The rate theory of chromatography has tended to supplement, rather than replace, the plate theory. The Van Deemter equation⁷ describes the effects of eluent flow-rate, longitudinal diffusion and multi-path effects within the column on plate height, H (H = column length, divided by N). For capillary columns in gas chromatography, the Golay equation⁸ shows how diffusion coefficients, thickness of stationary phase coating, column diameter, flow-rate, etc., contribute to plate height, H . Giddings⁴ has made major contributions to the dynamics of chromatography and has shown the effect of many experimental parameters on plate height in both gas (GC) and liquid chromatography (LC).

Although a theoretical value for H can be calculated from the diffusion coef-

ficients and other parameters contained in rate equations, the calculations often are not very accurate. For example, Desty and Goldup⁹ compared experimental and calculated values for terms in the Golay equation and found good agreement for the B term but that the experimental C term was 2–6 times the calculated value. The major value of rate equations is in predicting in a qualitative manner (or semi-quantitative at best) how a change in particle size, for example, will affect chromatographic behavior.

Most chromatographers still rely on only two actual measurements to evaluate performance: peak retention time and peak width. The discussion in a recent book on liquid chromatography¹⁰ is fairly typical of the way in which the dynamics of chromatography currently are handled. The authors consider that the theoretical plate number, N , of various chromatographic peaks is approximately constant and therefore they consider both N and H as measures of chromatographic efficiency. Thus, the rate theory and plate theory are in a way still tied together.

There seems to be some confusion as to how N varies (or does not vary) as the capacity factor, k , changes (recall that retention time of a peak varies directly with its capacity factor). While Snyder and Kirkland consider that N is approximately constant with k , Grob¹¹ states that N tends to increase with higher k and Kaiser¹² indicates that N decreases as k becomes larger. Jennings¹³ provides a nice experimental answer to this question. Careful gas chromatographic separation of n -alkanes on a capillary column shows that N is very high at low values of k and decreases with increasing k , rapidly at first when k is small, and then more slowly at larger k values, eventually approaching an asymptotic value. Their data show that n_{eff} starts at or near zero and increases as k becomes larger. N_{eff} appears to approach the same asymptotic values as N . The change in N or N_{eff} with k is particularly great at k values below 1.0.

In this paper we derive simple statistical expressions for calculating the mean and variance of chromatographic peaks that are still on a column (position peaks) and for these peaks as they leave the column (exit peaks). The variances of position and exit peaks as a function of capacity factor, k , are compared. We also show how the major contributions to peak broadening in capillary column GC can be determined at any k value using gaseous diffusion coefficients computed from empirically determined constants for the atoms and chemical groups making up the carrier gas and the compounds separated. First, the peak variance contributions resulting from axial diffusion are calculated from a modified form of the Van Deemter equation and are subtracted from the measured peak variances. Then the remaining variance of each peak is plotted against a function of the capacity factor, k . A plate number is calculated from the slope of this linear plot which is independent of k and thus provides an excellent measure of performance for a column operating under specified, fixed conditions. The intercept of this plot provides an estimate of the extra-column peak broadening of the chromatographic system.

For both GC and LC columns run at a single flow-rate, a similar approach is outlined that gives another plate number that also is independent of k but which includes any peak broadening effects resulting from axial diffusion. This plate number is also useful in describing the separating ability of a column under fixed conditions and in predicting the variance of peaks of different k values from those in the test chromatogram.

The authors feel that the material presented here will correct some of the confusion or misconceptions that many chromatographers seem to have about plate theory. We also think that the approach presented is a simple but practical way of handling some of the dynamic effects that determine the variance of chromatographic peaks.

2. PLATE MODELS

Using a statistical approach, we shall study two classical models falling into the category of plate theories: the first is the discrete flow model and the second the continuous flow model. The characteristic feature of a plate model is that the chromatographic column is considered to be divided into a sequence of disjoint segments, or "plates". In our discussion, we shall refer to these as theoretical segments (abbreviated to TS). The sample molecules proceed from one TS to the next until they leave the last segment and exit from the column.

The purpose here is to derive statistical equations for the mean and variance of a peak that is still on the column (a position peak), and for a peak leaving the column (an exit peak); this will be done for each of the plate models. The statistical approach also leads to equations that describe the entire shape of position and exit peaks, although such equations for counter-current chromatography (CCC) are already available in the literature.

3. DISCRETE FLOW MODEL

The assumptions of the discrete flow plate model are (1) the entire mobile phase moves from each segment to the next at the end of discrete equal time intervals and (2) the sample chemical remains in equilibrium between the stationary and the mobile phases. In our analysis we shall study the movement of a single molecule through the column. In this form the equilibrium condition is expressed by the condition that the probability that the molecule is in the mobile phase is p , while the probability it is in the stationary phase is $1 - p$. Thus $p = \frac{1}{1 + k}$, where k is the well known capacity factor. Thus the position of the molecule is random and depends crucially on the random "decisions" on whether to stick in the stationary phase or be free in the mobile phase during a given time unit. We shall measure the passage of time by the number of transfers (n) of the mobile phase that have occurred. We shall measure the location of the molecule by noting which segment it is in. Two quantities are of particular interest: one is the probability distribution of the position of the particle at time n , and the other the probability distribution of the time when the molecule leaves the column (the exit time).

We shall first examine the position distribution. In each time period the particle either goes to the next segment or stays in its current segment. The first event takes place with probability p and the second with probability $1 - p$. Which of these two events actually occurs in the k th time interval can be represented by the random variable

$$B_k = \begin{cases} 1 & \text{with probability } p \\ 0 & \text{with probability } 1 - p \end{cases}$$

Random variables of this type are known as “Bernoulli” random variables¹⁴. If we assume that our molecule starts in segment zero at time zero, then its position at time n [denoted by $X(n)$] is

$$X(n) = \sum_{k=1}^n B_k$$

As the B_k s are random, $X(n)$ is also random. The Bernoulli random variables in the sum are all independent and identically distributed with mean p and variance $p(1 - p)$. It follows that the expected value of $X(n)$ is

$$E[X(n)] = E\left(\sum_{k=1}^n B_k\right) = \sum_{k=1}^n E(B_k) = np \quad (1)$$

and the variance of $X(n)$ is

$$\text{Var}[X(n)] = \text{Var}\left(\sum_{k=1}^n B_k\right) = \sum_{k=1}^n \text{Var}(B_k) = np(1 - p) \quad (2)$$

These could also be written as

$$E[X(n)] = n \left(\frac{1}{1 + k} \right) \quad (3)$$

$$\text{Var}[X(n)] = n \left[\frac{k}{(1 + k)^2} \right] \quad (4)$$

The central limit theorem of probability¹⁴ tells us that $X(n)$ is asymptotically normal, *i.e.*, the distribution of $X(n)$ approaches closer and closer to a normal, or Gaussian, distribution as n increases ($n \approx 30$ is usually considered large enough for this to be a good approximation).

Probability theory will provide even more detailed information. An elementary result tells us that the sum of n independent identically distributed Bernoulli random variables with parameter p has a binomial distribution with parameters n and p (ref. 14). It follows that

$$\begin{aligned} P\{X(n) = j\} &= \binom{n}{j} p^j (1 - p)^{n-j} = \frac{n!}{(n-j)!j!} \cdot p^j (1 - p)^{n-j} \\ &= \frac{n!}{(n-j)!j!} \frac{k^{n-j}}{(1+k)^n} \end{aligned} \quad (5)$$

where j is some particular TS in the column and n is the number of transfers, or the

number of time intervals that have elapsed for a substance of capacity factor k . Eqn. 5 can be used to calculate the distribution of molecules along the various TS in a column by inserting various values of j into the equation. The result calculated will be the fraction of the molecules in each theoretical segment.

Next we examine the problem of determining the exit time distribution. If we denote the exit time by T , then $T = r + \sum_{k=0}^{r-1} D_k$. In this equation r is the number of theoretical segments in the column and therefore is the minimal amount of time necessary for the molecule to leave the column if it never enters the stationary phase. D_j is the (random) time that the molecule spends in the stationary phase of segment j . D_j can take on the values 0, 1, 2, ..., and it does so with probabilities $p, p(1-p), p(1-p)^2, \dots$, respectively. This is the distribution of a geometric random variable with parameter p (ref. 14). Thus the random variables $D_j, j = 0, \dots, r-1$ are independent, identically distributed, geometric random variables with parameter p . The mean and variance of such a geometric random variable is $(1-p)/p$ and $(1-p)/p^2$, respectively. Thus we obtain

$$E(T) = E\left(r + \sum_{k=0}^{r-1} D_k\right) = r + \sum_{k=0}^{r-1} E(D_k) = r + r\left(\frac{1-p}{p}\right) =$$

$$\frac{r}{p} = r(1+k) \quad (6)$$

$$\text{Var}(T) = \text{Var}\left(r + \sum_{k=0}^{r-1} D_k\right) = \text{Var}\left(\sum_{k=0}^{r-1} D_k\right) = \sum_{k=0}^{r-1} \text{Var}(D_k) =$$

$$r\left(\frac{1-p}{p^2}\right) = rk(1+k) \quad (7)$$

We can once again apply the central limit theorem of probability to conclude that T is asymptotically normal, i.e., as r becomes large the distribution of T approaches the normal distribution. It is even possible to obtain the exact probability distribution of T . To do this we utilize an earlier result: $P\{X(n) = j\} = \binom{n}{j} p^j (1-p)^{n-j}$.

We obtain the following:

$P\{T = n\} = P\{X(n-1) = r-1 \text{ and that the molecule is in the mobile phase in the next time interval}\} =$

$P\{X(n-1) = r-1\} P\{\text{molecule is in mobile phase in next time interval}\} =$

$$\left[\binom{n-1}{r-1} p^{r-1} (1-p)^{n-r}\right] p = \binom{n-1}{r-1} p^r (1-p)^{n-r}$$

Thus

$$P\{T = n\} = \frac{(n-1)!}{(n-r)!(r-1)!} \cdot \frac{k^{n-r}}{(1+k)^n} \quad (8)$$

The result in eqn. 8 is the fraction of molecules of capacity factor k that leaves a column, containing r theoretical segments, during the n th time period.

4. CONTINUOUS FLOW MODEL

The basic assumptions of this model are as follows: (1) eluent flows at a constant rate from one segment to the next; (2) the mobile and stationary phases remain in equilibrium at all times; and (3) there is perfect mixing in the segments. As we shall be looking again at only a single molecule, we interpret condition (2) to mean that the probability that the molecule is in the mobile phase at any given time is $\frac{1}{1+k}$ and the probability that it is in the stationary phase is $\frac{k}{1+k}$. "Units" are set up as follows. The amount of eluent in one segment is one unit of volume. The amount of time it takes for one unit volume to flow from one segment to the next is defined to be one time unit. Thus, in a time interval Δt , an amount of eluent Δt flows from each segment to the next.

Suppose we have r segments (numbered 0, 1, ..., $r-1$). Looking at some molecule of sample as it moves through the segments, we see that:

- (1) It spends a random amount of time, D_i , in segment i .
- (2) These "delay times", D_i , are independent, identically distributed random variables.

- (3) The exit time for the molecule is $\sum_{i=0}^{r-1} D_i$.

- (4) The TS the molecule is in at time t , $N(t)$, is given by

$$N(t) = \left\{ \text{minimum value of } n \text{ such that } \sum_{i=0}^{n-1} D_i \leq t < \sum_{i=0}^n D_i \right\} \quad (9)$$

(Note the above D_i is not to be confused with the one used in the discrete flow model.)

Clearly, our first task must be to determine the distribution of the random variables, D_i . To do this we look at a single segment. We wish to calculate the probability that a molecule (which is in this segment) leaves the segment in the next time interval Δt (Δt is assumed to be small). For this to happen it is necessary that the molecule (a) be in the mobile phase, and (b) be in that particular Δt volume of eluent that is leaving. This probability is

$$P\{a\} \cdot P\{b\} = \left(\frac{1}{1+k} \right) \left(\frac{\Delta t}{1} \right) = \Delta t \left(\frac{1}{1+k} \right)$$

This result is characteristic of an exponential delay time with parameter $\lambda = \frac{1}{1+k}$ (ref. 14). The density of such a random variable is $\lambda e^{-\lambda t}$ ($t \geq 0$).

With this piece of information together with some standard results of probability theory, we can now determine the exit time density and the probability distribution of the molecules' position at time t . First, the distribution of the exit time is the distribution of the sum of r independent identically distributed exponential random

variables. It is known that this distribution is a gamma distribution with parameters r and λ (ref. 14). The density function is

$$f(t) = \frac{\lambda(\lambda t)^{r-1} e^{-\lambda t}}{(r-1)!} \quad (t \geq 0)$$

Now we use the fact that the mean and variance of our exponential random variables are $1/\lambda$ and $1/\lambda^2$ respectively. Let T denote the exit time [which will have the density $f(t)$ above]. Then:

$$E(T) = E\left(\sum_{i=0}^{r-1} D_i\right) = \sum_{i=0}^{r-1} E(D_i) = \frac{r}{\lambda} = r(1+k) \quad (10)$$

$$\text{Var}(T) = \text{Var}\left(\sum_{i=0}^{r-1} D_i\right) = \sum_{i=0}^{r-1} \text{Var}(D_i) = \frac{r}{\lambda^2} = r(1+k)^2 \quad (11)$$

Once again we can apply the central limit theorem to conclude that the distribution of T is asymptotically normal (as $r \rightarrow \infty$).

It will be noted that the mean exit time (eqn. 10) is the same for both models but that the variance of the exit peak is $r(1+k)^2$ for the continuous flow model and $rk(1+k)$ for the discrete flow model.

To obtain the distribution of the position of sample molecules at time t , we note from eqn. 9 that D_i being identically distributed random variables makes $N(t)$ a Poisson process with parameter λ (ref. 15) [for the definition of $N(t)$, see eqn. 9]:

$$P\{N(t) = k\} = \frac{(\lambda t)^k e^{-\lambda t}}{k!} \quad (12)$$

for $k = 0, 1, \dots, r-1$.

It can be shown by a slightly more sophisticated application of the central limit theorem that the distribution of $N(t)$ is asymptotically normal (as $t \rightarrow \infty$) (this result also requires that r be large enough that the molecule will not yet have left the column at values of t large enough to yield a good normal approximation).

5. THEORETICAL PLATE NUMBERS IN THE TWO MODELS

The theoretical plate number, N , and the effective theoretical plate number, N_{eff} , are of course widely used in chromatography. These may be defined by

$$N = \frac{[E(T)]^2}{\text{Var}(T)} \quad (13)$$

and

$$N_{\text{eff}} = \frac{[E(T) - t_0]^2}{\text{Var}(T)} \quad (14)$$

where t_0 is the exit time for a non-sorbed substance and $E(T)$ is the expected exit time for a sample substance.

We can use the result we have obtained to calculate these quantities for both the discrete flow and continuous flow models. For the discrete flow mode:

$$N = \frac{[E(T)]^2}{\text{Var}(T)} = \frac{[r(1 + k)]^2}{rk(1 + k)} = r \left(\frac{1 + k}{k} \right) \quad (15)$$

$$N_{\text{eff}} = \frac{[E(T) - t_0]^2}{\text{Var}(T)} = \frac{(rk)^2}{rk(1 + k)} = r \left(\frac{k}{1 + k} \right) \quad (16)$$

For the continuous flow model:

$$N = \frac{[E(T)]^2}{\text{Var}(T)} = \frac{[r(1 + k)]^2}{r(1 + k)^2} = r \quad (17)$$

$$N_{\text{eff}} = \frac{[E(T) - t_0]^2}{\text{Var}(T)} = \frac{(rk)^2}{r(1 + k)^2} = r \left(\frac{k}{1 + k} \right)^2 \quad (18)$$

Thus, only in one of the four cases is the plate number result independent of k .

6. SOME CONSEQUENCES OF THE DISCRETE FLOW MODEL

In the discrete flow model the mean and variance of the location, $X(n)$, are expressed in units involving TS (here theoretical segments indicate the *position* of the molecule). On the other hand, the mean and variance of the exit time, T , must be expressed in time units. The fundamental time unit is the time needed for eluent to flow from one TS to the next. We wish to show that simple unit conversions can be used to relate results for peak locations to those for peak exit times. We define T_j as the time when a sample molecule enters the j th theoretical segments (note that $T_r = T$). Reasoning by analogy with eqns. 6 and 7, we have $E(T_j) = j(1 + k)$ and $\text{Var}(T_j) = jk(1 + k)$. Using eqns. 3 and 4 to calculate $E[X(n)]$ and $\text{Var}[X(n)]$ when $n \approx j(1 + k)$:

$$E[X(n)] = j(1 + k) \cdot \frac{1}{1 + k} = j \quad (19)$$

$$\text{Var}[X(n)] = j(1 + k) \cdot \frac{k}{(1 + k)^2} = j \cdot \frac{k}{1 + k} \quad (20)$$

Now we see that $E(T_j) = E[X(n)](1 + k)$ and $\text{Var}(T_j) = \text{Var}[X(n)](1 + k)^2$. In other words, we can convert between means and variances of position and exit time random variables simply by multiplying by the conversion fact $(1 + k)$ for the means and $(1 + k)^2$ for the variances.

Counter-current chromatography (CCC) is a good system for illustrating the practical results of the equations derived. In CCC the solute molecules attain equilibrium between the mobile and stationary phases prior to each transfer to the next tube in the series. Thus, each tube is by definition a theoretical plate, according to the plate theory of chromatography, or a theoretical segment in our nomenclature. It has been recognized by several authors that CCC also is a reasonable model for explaining elution behavior in column chromatography, provided that axial diffusion, multipaths and other factors that contribute to peak broadening in column chromatography, but which are non-existent in CCC, are also taken into account. This will be done in the next section, but first let us consider only the peak broadening resulting from inter-phase broadening effects, with CCC as a model.

Fig. 1 shows the distribution of a solute along the series of 36 tubes ($r = 36$) in a CCC apparatus as a function of the number of transfers, n . The peaks were plotted for $k = 1.5$ from eqn. 5 with the aid of a programmable calculator. Note that the peak has become well defined and approximately Gaussian after only a few transfers and that the peak broadens and decreases in height as it moves along the series of tubes. Fig. 2 shows the same peak, calculated from eqn. 8, as it leaves the CCC apparatus and passes through an imaginary detector. The peak that was symmetrical in its position distribution of molecules now has a distinct tail. The explanation for this is that the latter part of the peak continues to broaden after the earlier portions have left the "column". This effect is of some significance because it demonstrates that in columns of low "performance" some peak tailing is predicted mathematically and is not necessarily caused by a poorly functioning column.

The tailing of an exit peak decreases and the peak becomes more Gaussian as the number of theoretical segments (r) in a column becomes larger. The tailing, or skewness, of a peak is predicted by the following equation:

$$\text{Coefficient of skewness} = \frac{1}{\sqrt{r}} \cdot \frac{2k^2 + 3k + 1}{(k + 1)^2 [k/(k + 1)]^{1/2}} \quad (21)$$

(see ref. 14 for a definition of the coefficient of skewness). The second term in this equation has a value of just over 2 from $k = 1$ to $k = 10$, but increases rapidly as the value of k becomes very small.

It is well known that later eluting peaks are much broader than those with shorter retention times. From the exit peak variance (eqn 7) it will be seen that the peak width (4σ) is proportional to $\sqrt{k(1 + k)}$ and thus increases in approximately linear fashion with k (except at very low values of k). It is interesting to compare this with the way the width of a peak still on a column varies with k . This may be done by calculating the peak widths when the peak maximum is located in the j th theoretical segment in a column containing r theoretical segments. We shall choose j to be near the end of the column, say $j = 0.9r$. From eqn. 20 for the variance of a position peak we see that $4\sigma = 4 \left[\frac{jk}{1 + k} \right]^{1/2}$. In Table 1 we see that the widths of peaks centered at j vary only slightly with increasing k , much less than the exit peaks (of course it takes longer for peaks of higher k to arrive at j). The reason that elution peaks are so much broader when k is high is that such peaks are moving more slowly and take longer to pass through the detector.




Fig. 1. Position peaks for a compound with $k = 1.5$ in a CCC apparatus of 36 tubes after n transfers.

Exit peaks calculated for the discrete flow model using eqn. 8 emphasize a major drawback of the plate theory that has been so widely used in chromatography. In no case does either N or N_{eff} (calculated from eqns. 13 and 14) agree with the actual number of plates (tubes) in CCC, which is r . The correct values for N and N_{eff} are given by eqns. 15 and 16, respectively. Fig. 3 shows the expected values of N and N_{eff} as a function of k , calculated from eqns. 15 and 16 for a CCC apparatus where $r = 1000$. The values of N are very high as $k \rightarrow 0$ and become meaningless as a measure of separating power. Similarly, a single value of H (height equivalent to a theoretical plate) is an ambiguous expression of column efficiency when H varies so widely for different peaks.

The application of eqns. 5 and 8 to CCC is well documented and there would seem to be little doubt as to their correctness. It is logical that eqns. 3–8 should apply also to column chromatography. An examination of actual column chromatograms is now in order, to determine whether the equations do or do not apply.




Fig. 2. Exit peak for a compound with $k = 1.5$ from a CCC apparatus of 36 tubes (n = number of transfers).

TABLE I

WIDTHS (4σ) OF POSITION AND EXIT PEAKS AS A FUNCTION OF k

k	Position width* (TS)	Exit width** (TS)	Exit width*** (sec)
0.1	36	42	2.5
0.2	49	62	3.7
0.4	64	95	5.7
0.6	74	124	7.4
0.8	80	152	9.1
1.0	85	179	10.7
1.5	93	245	14.7
2.0	98	310	18.6
3.0	104	438	26.3
4.0	107	566	33.9
6.0	111	820	49.2
8.0	113	1073	64.4
10.0	114	1327	79.6
15.0	116	1960	117.6

* Calculated from $4\sigma = 4 \left[\frac{jk}{1+k} \right]^{1/2}$, where $j = 900$.

** Calculated from $4\sigma = 4[rk(1+k)]^{1/2}$, where $r = 1000$.

*** Calculated from the previous column for an eluent flow-rate of 16.67 TS/sec.




Fig. 3. Expected values of N and N_{eff} for an exit peak from a CCC apparatus where $r = 1000$.

7. CALCULATION OF PLATE NUMBERS FROM CHROMATOGRAPHIC DATA

When working with actual data for column chromatography, it is necessary to consider the various dynamic effects that contribute to the width of chromatographic peaks. The major effect almost always is what is termed "resistance to mass transfer", which we shall call "inter-phase broadening". Additional peak broadening occurs through axial diffusion, multi-paths (in packed columns) and extra-column effects such as broadening in fittings, connecting tubing and the detector. Injection band broadening is apt to be significant also, as was recognized by Keulemans as early as 1959^{16,17}.

If we assume that these broadening effects act independently of one another, the measured peak variance is the sum of the variances of the individual broadening effects. For capillary GC columns (where there is no multi-path effect), we can write

$$\sigma^2 = \sigma_{ec}^2 + \sigma_d^2 + \sigma_{ip}^2 \quad (22)$$

where ec stands for extra column, d for diffusion and ip for inter-phase. The approach taken will be first to estimate the peak variance resulting from axial diffusion and to subtract this from the measured peak variance. Then a plate number representing the inter-phase broadening can be calculated from the slope of a plot of the remaining peak variance against $k(1 + k)$, which comes from the peak variance in the discrete model, $rk(1 + k)$.

7.1. Diffusion broadening, σ_d^2

The effect of diffusion on plate height, H , is given by the Van Deemter equation, which has the form, $H = \frac{B}{u} + Cu$, where B is the diffusion term, u is the linear column flow-rate and C is the resistance to mass transfer. In this equation, $B = 2D_g$, where D_g is the axial diffusion coefficient in the gas phase (cm^2/sec).

According to the classical definition of H and N , the diffusion contribution to H is converted to σ_d^2 as follows:

$$H = \frac{L}{N} = \frac{L \sigma_d^2}{[t_0(1 + k)]^2} = \frac{2D_g}{u} \quad (23)$$

$$\sigma_d^2 = \frac{2D_g}{u} \cdot \frac{(t_0)^2 (1 + k)^2}{L} = \frac{2D_g L (1 + k)^2}{u^3} \quad (24)$$

where L is the column length (cm) and u is the linear flow-rate (cm/sec).

One way to estimate the diffusion broadening variance is to run a chromatogram at two different flow-rates. Diffusion broadening varies as the inverse cube of the flow velocity, but inter-phase broadening varies only as the inverse of the flow velocity. This difference allows us to solve for diffusion broadening by solving two simultaneous equations (the dependence of the broadening on flow velocity will be considered in the next section).

Although this method works, a more convenient way is to use the method for

estimating gaseous diffusion coefficients proposed by Giddings¹⁸:

$$D_g \text{ (for A,B)} = \frac{10^{-3} T^{1.75} \left(\frac{1}{M_A} + \frac{1}{M_B} \right)^{\frac{1}{2}}}{p[(\sum_A v_i)^{1/3} + (\sum_B v_i)^{1/3}]^2} \quad (25)$$

where M_A and M_B are the molecular weights of the carrier gas and a chromatographically separated substance, respectively, T is the temperature (°K) p is the pressure (bar) and v_i are empirically determined values for various atoms and groups making up the carrier gas and the separated chemical. Giddings and co-workers obtained excellent agreement (usually to within $\pm 5\%$) between D_g values calculated this way and experimentally reported values¹⁸.

As will be shown by an actual example, we found that this method gives reasonable results when applied to column chromatography. Although the values of D_g decrease as one proceeds to increasingly higher members of a homologous series, the fact that the D_g values are multiplied by $(1 + k)^2$ means that σ_d^2 (in sec²) actually increases as one proceeds to later eluting peaks. Despite differences in individual D_g values, a linear regression plot of calculated σ_d^2 versus $(1 + k)^2$ is generally obtained with a high correlation coefficient.

7.2. Inter-phase broadening, σ_{ip}^2

The general approach taken is to subtract σ_d^2 from the measured variance of each peak and to plot the difference, σ_{net}^2 , against one of the expressions for stochastic peak variance derived earlier. This should give a straight line with a slope related to a plate number and an intercept indicative of extra-column peak variance.

However, one question is whether $rk(1 + k)$ in the discrete flow model or $r(1 + k)^2$ in the continuous flow model is correct. For many plots, both give straight lines by linear regression. This is because the ratio of $(1 + k)^2/k(1 + k)$ does not change greatly with k at higher k values (say 2.5–15) and also that linear regression is insensitive to points of lower k that might not lie on a straight-line plot.

However, C values calculated from the Golay equation where several of the k values are quite low shows that slopes of individual points (y/x for each point) are nearly constant for plots of σ^2 versus $k(1 + k)$ but they vary considerably for the same k values when σ^2 is plotted versus $(1 + k)^2$. This supports the contention that the discrete flow is correct for column chromatography and that it is correct to plot σ^2 versus $k(1 + k)$ for inter-phase broadening.

Using eqn. 7 and making use of the conversion factor t_0/r (sec/TS) it can be shown that

$$\sigma_{ip}^2 = \frac{(t_0)^2 k(1 + k)}{r} = \frac{L^2 k(1 + k)}{u^2 r} \quad (26)$$

where L is the column length (cm), u is the linear flow-rate (cm/sec) and r is the number of theoretical segments in the column at the flow-rate employed. However, we know that the plate number, r , is proportional to the reciprocal of linear flow-rate.

Introducing a plate number, r' , which is the number of theoretical segments in a column at $u = 1\text{cm/sec}$:

$$r = \frac{r'}{u}$$

$$\sigma_s^2 = \frac{L^2 k(1 + k)}{u r'} \quad (27)$$

This equation for peak variance due to stochastic broadening was tested on data published by Kaiser and Rieder¹⁹ for the separation of alkanes and aliphatic esters by capillary column GC at six different flow-rates. First, the peak variance resulting from axial diffusion was subtracted from the measured peak variance as outlined above. Then the remaining variance (σ_{net}^2) was plotted against $k(1 + k)$ and the plate number r' calculated from the slope of the straight-line plot obtained. A sample calculation is shown in Table 2. The results are summarized in Table 3, which shows a constant value for r' within a reasonable experimental error. The plate number, r , at any flow-rate can be calculated simply by dividing r' by u .

The plate number, r' , seems to be a convenient way of evaluating and comparing column performances. It is computed easily from measured peak variances using linear regression and is independent of peak capacity factor. It is also independent of axial diffusion effects. This is good because diffusion really has nothing to do with the separating ability of a column other than contributing to the broadening of peaks. In some respects r' is similar to the C term in the Van Deemter and Golay equations. However, C is different for each peak whereas r' is not. It would appear that r' can be more easily and accurately measured from ordinary chromatographic data than can C .

At a fixed flow-rate it is convenient to obtain a plate number that includes

TABLE 2

SAMPLE DATA FROM KAISER AND REDIER¹⁹ FOR $t_0 = 146$ sec SHOWING AXIAL DIFFUSION CONTRIBUTION TO PEAK VARIANCE

σ^2 (diffusion) was calculated from eqn. 24 using D_g values calculated from eqn. 25 at an estimated pressure of 1.2 bar. The values of the gaseous diffusion coefficient (D_g) used ranged from 0.0795 for C_9 to 0.0653 for C_{13} .

Compound*	k	σ^2 (measured) (sec ²)	σ^2 (diffusion) (sec ²)	σ^2 (net) (sec ²)
C ₉	1.26	1.52	0.63	0.88
E ₉	1.45	1.96	0.74	1.22
C ₁₀	2.38	4.15	1.33	2.82
E ₁₀	2.75	4.88	1.64	3.24
C ₁₁	4.51	10.41	3.38	7.03
E ₁₁	5.23	13.34	4.32	9.02
C ₁₂	8.55	35.34	9.65	25.69
E ₁₂	9.94	43.88	12.66	31.22
C ₁₃	16.2	112.7	30.0	82.7

* C = *n*-alkane and carbon number; E = alkane ester and carbon number.

TABLE 3

PLATE NUMBER (r') CALCULATED FROM DATA OF KAISER AND RIEDER¹⁹ AFTER SUBTRACTING DIFFUSION PEAK VARIANCE

t_0 (sec)	u (cm/sec)	$r' \cdot 10^5$ (TS)
276	7.25	9.19
154.4	12.95	10.40
146.0	13.70	9.86
106.8	18.73	10.71
88.2	22.68	9.53
87.4	22.88	9.82
Average: 9.92 ($s = 0.56$)		

broadening resulting from axial diffusion as well as inter-phase peak broadening. This can be done simply by measuring the variance of each of the chromatographic peaks and plotting measured σ^2 against peak variance, $k(1 + k)$. The apparent plate number, r , is obtained from the slope of the linear plot. The intercept is indicative of the extra-column peak broadening plus the diffusion broadening of a non-sorbed or a very slightly sorbed peak.

This simple plotting method works because a plot of σ_a^2 versus $(1 + k)^2$ is linear and a plot of σ_a^2 versus $k(1 + k)$ is almost linear. The flow-rates that are normally used in chromatography sufficiently fast that diffusion contribution to peak width usually is not very substantial. Thus, any slight non-linearity of the σ_a^2 part of the plot is insignificant.

The equation relating measured peak variance to r and k is derived by converting the peak variance, $rk(1 + k)$, from TS units to sec^2 by dividing $\frac{r^2}{(t_0)^2}$ (t_0 is the hold-up time):

$$\sigma_{ip}^2 = \frac{rk(1 + k)}{r^2/(t_0)^2} = \frac{(t_0)^2}{r} [k(1 + k)] \quad (28)$$

where σ_{ip}^2 and t_0 are now in minutes or seconds. Substitution into eqn. 22 gives

$$\sigma^2 = \frac{(t_0)^2}{r} [k(1 + k)] + \sigma_{ec}^2 \quad (29)$$

where σ , t_0 and σ_{ec} are in minutes or seconds. This equation predicts that a plot of σ^2 versus $k(1 + k)$ for the various peaks in a chromatogram will give a straight line with slope $(t_0)^2/r$ and intercept $= \sigma_{ec}^2$.

The validity of eqn. 29 was tested on a number of actual chromatograms. Usually the chart speed of the recorder was 4–10 cm/min to facilitate measurement of the peak widths with a ruler and magnifying lens. The peak variance was obtained from the peak width at half of its height, $w_{1/2}$ (where $\sigma = 2.355 w_{1/2}$) in order to minimize the effect of any tailing in the lower part of the peak.

Data for five chromatographic separations are summarized in Table 4. Chro-

TABLE 4
CALCULATION OF r FROM CHROMATOGRAPHIC DATA

Chromatogram	Peak	k	$k(1 + k)$	σ^2	Results
(A) <i>n</i> -Alkanes by capillary GC ¹³ $t_0 = 9.72$ min Oven at 130°C	C ₇	0.14	0.16	0.00032 (min ²)	Correlation coefficient = 0.9999 $\sigma_2^2 = 0.00001$ Slope = 0.00518 $r = 182,500$
	C ₈	0.8	1.44	0.00090	
	C ₉	1.4	3.36	0.00187	
	C ₁₀	2.3	7.59	0.00380	
	C ₁₁	3.9	19.1	0.0096	
	C ₁₂	6.5	48.8	0.0249	
	C ₁₃	10.8	127.4	0.0661	
(B) <i>n</i> -Alkanes by capillary GC at two different temperatures*. First set, 60°C; second set 70°C. $t_0 = 1.20$ min	C ₈	0.97	1.91	$4.62 \cdot 10^{-5}$ (min ²)	Correlation coefficient = 0.9999 $\sigma_2^2 = 2.27 \cdot 10^{-5}$ Slope = $1.676 \cdot 10^{-5}$ $r = 85,900$
	C ₉	2.31	7.65	$14.1 \cdot 10^{-5}$	
	C ₁₀	5.42	34.8	$62.8 \cdot 10^{-5}$	
	C ₁₁	12.33	164.4	$277 \cdot 10^{-5}$	
	C ₈	0.60	0.96	$4.06 \cdot 10^{-5}$	
	C ₉	1.44	3.51	$7.21 \cdot 10^{-5}$	
	C ₁₀	3.26	13.9	$26.0 \cdot 10^{-5}$	
	C ₁₁	7.19	58.9	$101 \cdot 10^{-5}$	
(C) Carboxylic acids by packed-column GC**. $t_0 = 2.0$ mm	Acetic	1.8	5.04	0.0451 (mm ²)	Correlation coefficient = 0.9998 $\sigma_2^2 = 0.0348$ Slope = 0.00196 $r = 2040$
	Propionic	3.4	15.0	0.0649	
	Isobutyric	5.0	30.0	0.0909	
	<i>n</i> -Butyric	6.55	49.4	0.1303	
	Isovaleric	9.7	104	0.2426	
(D) Alkali metal ions by ion exchange***. 50- μ l sample. $t_0 = 42.5$ sec	<i>n</i> -Valeric	13.3	190	0.4057	Correlation coefficient = 0.9988 $\sigma_2^2 = 19.0$ Slope = 1.514 $r = 1190$
	Li ⁺	4.35	23.26	51.8 (sec ²)	
	Na ⁺	6.10	43.2	83.0	
	NH ₄ ⁺	8.47	80.1	148.0	
	K ⁺	11.45	142.6	229.8	
(E) Alkali metals by ion exchange. 20- μ l sample. $t_0 = 39$ sec	Rb ⁺	13.0	182.0	295.8	Correlation coefficient = 0.9979 $\sigma_2^2 = 16.3$ Slope = 1.067 $r = 1430$
	Li ⁺	4.90	28.88	54.6 (sec ²)	
	Na ⁺	6.54	49.30	62.4	
	NH ₄ ⁺	9.44	98.47	117.2	
	K ⁺	12.51	169.0	200.0	
(F) Pesticides by liquid chromatog- raphy. $t_0 =$ 0.445 min [§]	Rb ⁺	14.43	222.7	253.6	Correlation coefficient = 0.9996 $\sigma_2^2 = 2.11 \cdot 10^{-4}$ Slope = $0.7445 \cdot 10^{-4}$ $r = 2660$
	Sevin	0.955	1.87	0.000357 (min ²)	
	Prolate	1.16	2.50	0.000391	
	Ruelene	1.72	4.67	0.000786	
	Coral	1.94	5.72	0.000612	
	Dursban	3.45	15.35	0.001356	
	<i>trans</i> - Permethrin	5.81	39.55	0.003236	
	<i>cis</i> - Permethrin	6.80	53.01	0.004100	

* J. Kaczvinsky, unpublished results (1981).

** Data from chromatogram in an advertising brochure (1981).

*** G. Sevenich, unpublished results (1981).

§ L. Rice, unpublished results (1981).




Fig. 4. Linear regression plot for chromatogram C in Table 2.

matogram A is unique among those tested in that the intercept, σ_{ec}^2 , was essentially zero. Of course there is some error in measuring k and σ^2 for each peak, but the equipment and technique used were both known to be excellent.

The data for B in Table 4 are for two chromatograms run at different temperatures, yet all eight points lie on a straight line within reasonable experimental error. The intercept, σ_{ec}^2 , is significant in this instance and constitutes nearly 50% of the measured variance of the first C₈ peak. Note that the early peaks are far more affected by σ_{ec}^2 than the latter ones if the actual peak widths (4σ) are compared.

Chromatogram C was run on a packed GC column and shows a far greater σ_{ec}^2 relative to the measured peak variances than the chromatograms or capillary columns. Fig. 4 confirms that the experimental points actually do fall on a straight line when plotted according to eqn. 29. It is interesting to calculate the traditional plate number, N , before and after subtraction of σ_{ec}^2 from σ^2 for each peak. Before any correction is applied, N increases from peaks 1 to 6: $N = 695, 1190, 1580, 1750, 1890, 2020$. After subtracting σ_{ec}^2 from the variance of each peak, N decreases from peaks 1 to 6: $N = 3050, 2570, 2390, 2200, 2200$.

Chromatograms D and E are for an ion-exchange separation of alkali metal ions. Again, the data support the validity of eqn. 28 even though some of the peaks were tailed, and accurate measurement of peak width was difficult. The intercept, σ_{ec}^2 , is smaller and the value of r is larger when the sample volume is reduced from 50 to 20 μ l.

Chromatogram F in Table 4 was obtained with a commercial liquid chromatograph with a 10 cm $\times$ 4.6 mm I.D. column. The large value for r confirmed the excellent separating ability of the column, but the additional broadening represented by σ_{ec}^2 was large. Much of this can be attributed to the use of a small "guard" column filled with a coarser reversed-phase absorbent than that used in the chromatographic column. Subsequent replacement of the guard column with a pre-column of slightly smaller diameter packed with 10- μ m absorbent reduced σ_{ec}^2 by almost 50%.

While there is no direct proof that the intercept, σ_{ec}^2 , of our linear regression analysis gives an accurate measure of the sum of in-column and extra-column broad-

ening that applied equally to all peaks, the underlying principle does appear sound and the results obtained from actual data are reasonable. The idea of linear regression analysis of chromatographic data originated with Kaiser and co-workers¹⁹⁻²¹, who plotted $w_{1/2}$ against k . They obtained the "real" number of theoretical plates from the slope and an indication of the extra-column peak broadening from the intercept. We think that the concept pioneered by Kaiser and co-workers is a valuable one but that their method of plotting the data is not completely correct. According to Guiochon²², a plot of $w_{1/2}$ versus k is not linear at all values of k and the intercept of such a plot underestimates the "extra-column" peak broadening.

Smuts *et al.*²³ proposed a plot of peak variance against $(1 + k)^2$, although the purpose of this work was different from ours. This type of plot would follow if the continuous flow model [where the peak variance is $(1 + k)^2$] is correct for column chromatography. However, Smuts *et al.*²⁴ stated that such a plot is not completely linear. Plots of σ^2 versus $(1 + k)^2$ for the data in Table 2 also did not give a completely straight line, although the intercept was often not greatly different from that obtained when using the discrete flow model. All of our results suggest that the discrete flow model is better for column chromatography.

8. SUMMARY

The classical plate theory, and original extensions of it, are derived by using simple concepts from the theory of probability and statistics. Each molecule of a sample chemical substance is examined separately and its motion through the column is described as a stochastic process. Equations for calculating the mean and variance of chromatographic peaks as a function of capacity factor, k , are given for a discrete flow model and a continuous flow model. The variance of position and exit peaks as a function of capacity factor is compared and a simple relationship between the two is derived. The expressions for the mean and variance of chromatographic peaks are used to define plate numbers that describe the separating ability of a chromatographic column under fixed conditions and which, unlike the classical plate numbers N and N_{eff} , are independent of capacity factor, k . In capillary column GC a method is given for determining diffusion coefficients in the gaseous phase that makes it possible to subtract the contribution to peak variance resulting from axial diffusion. When the remaining variance is plotted against $k(1 + k)$, linear regression indicates a straight line with an excellent correlation coefficient and an intercept that is indicative of extra-column peak broadening. A plate number, r' , is computed from the slope that is independent of k and can be used to calculate a plate number, r , for any given linear flow-rate. Another simple plotting method gives a plate number that includes the multi-path contribution to peak broadening for packed columns. Numerous examples are given to demonstrate the applicability of these simple concepts to actual chromatographic data.

REFERENCES

- 1 A. J. P. Martin and R. L. M. Synge, *Biochem. J.*, 35 (1941) 1358.
- 2 E. Glueckauf, *Trans. Faraday Soc.*, 51 (1955) 34.
- 3 S. W. Mayer and E. R. Tompkins, *J. Amer. Chem. Soc.*, 69 (1947) 2866.
- 4 J. C. Giddings, *Dynamics of Chromatography, Part I*, Marcel Dekker, New York, 1965, Ch. 2.

- 5 J. H. Purnell, *Nature (London)*, 184 (1959) 2009.
- 6 J. H. Purnell, *J. Chem. Soc.*, 54 (1950) 1268.
- 7 J. J. van Deemter, F. J. Zuiderweg and A. Klinkenberg, *Chem. Eng. Sci.*, 5 (1956) 271.
- 8 R. L. Grob, *Modern Practice of Gas Chromatography*, Wiley-Interscience, New York, 1977, pp. 71–77 and 117.
- 9 D. H. Desty and A. Goldup, in R. P. W. Scott (Editor), *Gas Chromatography 1960*, Butterworths, London, 1960, p. 162.
- 10 L. R. Snyder and J. J. Kirkland, *Introduction to Modern Liquid Chromatography*, Wiley, New York, 1979, p. 28.
- 11 R. L. Grob, *Modern Practice of Gas Chromatography*, Wiley-Interscience, New York, 1977, p. 65.
- 12 R. E. Kaiser, *Chromatographia*, 10 (1977) 455.
- 13 W. Jennings, *Gas Chromatography with Glass Capillary Columns*, Academic Press, New York, 1980, pp. 7 and 8.
- 14 A. M. Mood, F. A. Graybill and D. C. Boes, *Introduction to the Theory of Statistics*, McGraw-Hill, New York, 1950.
- 15 E. Çinlar, *Introduction to Stochastic Processes*, Prentice-Hall, Englewood Cliffs, NJ, 1975.
- 16 A. I. M. Keulemans, *Gas Chromatography*, Reinhold, New York, 2nd ed., 1959, p. 123.
- 17 A. B. Littlewood, *Gas Chromatography*, Academic Press, New York, 2nd ed., 1970, pp. 35–36.
- 18 E. N. Fuller, K. Ensley and J. C. Giddings, *Ind. Eng. Chem.*, 58 No. 5 (1966) 18.
- 19 R. E. Kaiser and R. Rieder, *Chromatographia*, 10 (1977) 455.
- 20 R. E. Kaiser, *Chromatographia*, 9 (1976) 337 and 463.
- 21 R. E. Kaiser and E. Oelrich, *Optimierung in der HPLC*, Hüthig, Heidelberg, 1979.
- 22 G. Guiochon, *Chromatographia*, 11 (1978) 249.
- 23 T. W. Smuts, T. S. Buys, K. de Clerk and T. G. du Toit, *J. High Resolut. Chromatogr. Chromatogr. Commun.*, 1 (1979) 41.
- 24 T. W. Smuts, T. S. Buys, G. du Toit and J. W. du Toit, *J. High Resolut. Chromatogr. Chromatogr. Commun.*, 4 (1981) 363.

CHROMATOGRAPHIC DATA

SUPPLEMENT TO
CHROMATOGRAPHIC REVIEWS
1983

COORDINATING EDITOR:
J. GASPARIČ (Hradec Králové)

EDITORS:
Z. DEYL (Prague)
M. LEDERER (Switzerland)
J. JANÁK (Brno)



ELSEVIER SCIENCE PUBLISHERS B.V.
AMSTERDAM—OXFORD—NEW YORK

© ELSEVIER SCIENCE PUBLISHERS B.V. — 1983

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, without the prior written permission of the publisher, Elsevier Science Publishers B.V., P.O. Box 330, Amsterdam, The Netherlands.

Printed in The Netherlands

TABLE 1181

TABLE OF ISOTACHOPHORETIC INDICES

I. SIMULATED QUALITATIVE AND QUANTITATIVE INDICES OF 287 ANIONIC SUBSTANCES IN THE RANGE pH 3–10

TAKESHI HIROKAWA*, MAKOTO NISHINO*, NOBUYUKI AOKI and YOSHIYUKI KISO
Applied Physics and Chemistry, Faculty of Engineering, Hiroshima University, Shitami, Saijo, Higashi-hiroshima 724 (Japan)

and

YASUYO SAWAMOTO, TAKAO YAGI and JUN-ICHI AKIYAMA
Shimadzu Corporation, Nakagyo-ku, Kyoto 604 (Japan)

(Received January 31st, 1983)

SUMMARY

A simulation of isotachophoretic equilibria has been applied to 287 anions, evaluating R_E indices for qualitative determination and time-based zone lengths for quantitation under 31 leading electrolyte conditions in the range pH 3–10, using eight different buffers. The R_E values may be useful for correcting asymmetric potentials of sensing electrodes for precise R_E measurement, assessing the separability of samples and estimating the optimum separation conditions. The zone lengths for 10-nmol samples obtained at a driving current of 100 μ A may be useful for quantitation.

INTRODUCTION

A unique feature of capillary isotachophoretic apparatus commonly used is that there are no packing materials in the separation column. The isotachophoretic behaviour in such a free electrolyte system can be analyzed theoretically and the observed isotachopherograms can be simulated using a few physico-chemical constants of the sample components and appropriate separation conditions^{1,2}. In previous papers we have described applications of this simulative technique to analytical problems such as the assessment of the separability of samples at different pH values², estimation of the optimum separation conditions² and quantitation without a calibration line³. In principle, the separability of samples by isotachopheresis is determined by the effective mobilities of the sample components, which are closely related to their absolute mobilities and pK_a values and to the pH of the leading electrolyte, pH_L, buffered by an appropriate counter ion. To ensure that there is sufficient difference between the effective mobilities of the sample components, a complex-forming technique and a solvent effect are also utilized⁴.

* Present address: Shimadzu Corporation, Kyoto, Japan.

In the present work, the qualitative and quantitative indices of 287 anions are simulated under 31 aqueous electrolyte conditions in the range pH 3–10 by considering the pH dependence of the effective mobility. These indices, *i.e.*, R_E and the time-based zone length, l , can be used as a data base for analytical purposes and for correction of asymmetric potential in the measurement of R_E^5 , qualitative identification without addition of a standard and conversion of R_E into other qualitative indices and *vice versa*.

SIMULATION CONDITIONS

The pH of the leading electrolyte, pH_L , was varied in the range 3–10 (0.25 pH units per step) using eight kinds of buffer ions: β -alanine ($pH_L = 3$ –4); ϵ -aminocaproic acid (4–4.75); creatinine (4.75–5.25); histidine (5.5–6.25); imidazole (6.5–7.25); tris(hydroxymethyl)aminomethane (Tris) (7.5–8.25); 2-amino-2-methyl-1,3-propanediol (amediol) (8.5–9.25); ethanolamine (9.5–10). The absolute mobilities, m_0 , and the dissociation constants, pK_a , of the buffers used are summarized in Table I. The absolute mobility of cationic β -alanine ($36.7 \cdot 10^{-5} \text{ cm}^2 \text{ V}^{-1} \text{ sec}^{-1}$) obtained by conductivity measurement was different from the previously used value⁵ of $31.0 \cdot 10^{-5} \text{ cm}^2 \text{ V}^{-1} \text{ sec}^{-1}$ obtained by analysing the step height of anionic β -alanine and assuming the cationic m_0 to be equal to that of the anionic m_0 . The leading ion was 0.01 *M* chloride and the driving current was 100 μA . Table II shows the simulation conditions, the simulated concentration of buffer, the effective mobility of the buffer and leading ion, the specific conductivity and the potential gradient of the leading zones. For simulation of the latter the inner diameter of the separation column was assumed to be 0.5 mm. The isotachophoretic zone velocity at each pH_L condition is also shown in Table II. This velocity can be used to estimate the time needed for detecting the first sample zone.

The m_0 and pK_a values of the 287 substances are mainly literature values. For about 80 substances, isotachophoretically evaluated constants were used. It should be noted that the correction of m_0 of β -alanine and revision of the computer program caused little changes in the m_0 and pK_a values evaluated previously⁵. For a part of the samples, especially for isomers of benzene derivatives, pK_a could be found in the

TABLE I
PHYSICO-CHEMICAL CONSTANTS OF BUFFERS USED IN SIMULATION (25°C)

m_0 = Absolute mobility ($\text{cm}^2 \text{ V}^{-1} \text{ sec}^{-1} \times 10^5$). pK_a = thermodynamic acid dissociation constant. The absolute mobilities were obtained isotachophoretically, and the pK_a values were taken from the literature.

pH_L range	Buffer	pK_a	m_0
3.00–4.00	β -Alanine	3.552	36.7
4.00–4.75	ϵ -Aminocaproic acid	4.373	28.8
4.75–5.25	Creatinine	4.828	37.2
5.50–6.25	Histidine	6.04	29.6
6.50–7.25	Imidazole	7.15	52.0
7.50–8.25	Tris	8.076	29.5
8.50–9.25	Amediol	8.78	29.5
9.50–10.0	Ethanolamine	9.498	44.3

TABLE II

SIMULATED EFFECTIVE MOBILITIES AND CONCENTRATION OF LEADING ZONE CONSTITUENTS (25°C)

pH_L = pH of leading electrolyte; $\bar{m}_L$ = effective mobility ($\text{cm}^2 \text{V}^{-1} \text{sec}^{-1} \times 10^5$) of leading ion (Cl^-), m_0 of Cl^- being $79.08 \cdot 10^{-5} \text{cm}^2 \text{V}^{-1} \text{sec}^{-1}$; $C_{B,L}^t$ = total concentration (mM) of buffer ion; $\bar{m}_{B,L}$ = effective mobility ($\text{cm}^2 \text{V}^{-1} \text{sec}^{-1} \times 10^5$) of buffer ion, the counter direction being indicated by a negative sign; κ_L = specific conductivity (S cm^{-1}) of leading zone; E_L = potential gradient (V cm^{-1}) of leading zone at a driving current of 100 μA (I.D. of narrow bore tube: 0.5 mm); v = isotachophoretic zone velocity ($\text{cm sec}^{-1} \times 10^3$).

Buffer	pH_L	$\bar{m}_L$	$C_{B,L}^t$	$\bar{m}_{B,L}$	κ_L	E_L	v
β -Alanine	3.00	74.64	11.28	-26.48	1.348	37.77	28.20
	3.25	74.61	13.68	-22.88	1.212	41.98	31.32
	3.50	74.57	17.43	-18.43	1.137	44.80	33.40
	3.75	74.51	23.79	-13.70	1.094	46.56	34.69
	4.00	74.44	34.94	-9.40	1.069	47.63	35.46
ϵ -Aminocaproic acid	4.00	74.60	13.68	-18.41	0.997	51.09	38.12
	4.25	74.58	16.70	-15.15	0.982	51.84	38.65
	4.50	74.48	22.01	-11.52	0.974	52.29	38.94
	4.75	74.40	31.42	-8.08	0.969	52.57	39.11
Creatinine	4.75	74.57	17.50	-19.20	1.050	48.51	36.18
	5.00	74.52	23.37	-14.39	1.047	48.65	36.25
	5.25	74.45	33.80	-9.95	1.045	48.74	36.29
Histidine	5.50	74.63	12.59	-20.82	0.974	52.28	39.02
	5.75	74.59	14.62	-17.93	0.973	52.33	39.03
	6.00	74.53	18.22	-14.39	0.972	52.37	39.03
	6.25	74.46	24.62	-10.65	0.972	52.42	39.03
Imidazole	6.50	74.67	12.02	-40.08	1.202	42.97	32.08
	6.75	74.66	13.59	-35.44	1.185	42.97	32.08
	7.00	74.63	16.38	-29.40	1.185	42.98	32.08
	7.25	74.60	21.35	-22.56	1.185	42.99	32.07
Tris	7.50	74.63	12.39	-21.08	0.972	52.38	39.10
	7.75	74.60	14.26	-18.32	0.972	52.40	39.09
	8.00	74.54	17.57	-14.87	0.971	52.43	39.08
	8.25	74.47	23.46	-11.14	0.971	52.45	39.06
Amediol	8.50	74.59	14.74	-17.73	0.972	52.38	39.07
	8.75	74.53	18.42	-14.19	0.972	52.38	39.04
	9.00	74.45	24.99	-10.46	0.973	52.37	38.99
	9.25	74.37	36.67	-7.14	0.973	52.32	38.91
Ethanolamine	9.50	74.58	19.12	-21.31	1.119	45.53	33.95
	9.75	74.53	26.25	-15.56	1.124	45.32	33.77
	10.00	74.47	39.01	-10.51	1.133	44.94	33.46

literature but m_0 could not. In such cases, the m_0 values were assumed equal to those of compounds of similar structure or similar molecular weight.

The calculations were carried using a SORD microcomputer M223 Mk III

equipped with two 1 MB floppy disk drivers. The results were filed on a floppy disk. To calculate 8897 (287×31) R_E values required *ca.* 10 h.

USE OF INDICES

The names of the 287 anionic substances simulated are arranged in alphabetical order in Table III. The simulated results are summarized in Table IV together with values of m_0 and pK_a , the molecular formula and molecular weight. The negative signs of m_0 for anions are omitted except for a few amino acids. Also listed in Table IV are the ratio, R_E , of the potential gradient of the sample zone, E_s , to that of the leading zone, E_L , the time-based zone length, t (sec), the effective charge, $-Z$, and the pH of the sample zone, pH_s , at each pH_L .

I. Correction of asymmetric potential

When a potential gradient detector is used, the actual baseline for the measurement of the step heights of the leading, h_L , and the sample zone, h_s , sometimes drifts due to an asymmetric potential at the sensing electrodes. This drift sometimes results in a significant error in the observed R_E values. In such cases, the simulated R_E values of samples with well defined m_0 and pK_a can be obtained from an evaluation of the drifted step height, Δh . In Table IV, the R_E values of substances marked with (S), R_E (std), are useful as internal standard indices. The Δh can be evaluated as

$$\Delta h = [h_{std} - h_L \cdot R_E(std)]/[R_E(std) - 1] \quad (1)$$

where h_{std} is the apparent step height of the zone of the internal standard and h_L that of the leading zone. Using Δh , the R_E values of the sample components can be corrected

$$R_E = E_s/E_L = (h_s + \Delta h)/(h_L + \Delta h) \quad (2)$$

where h_s is the observed step height of the sample. Examples of this mode of correction were reported in ref. 5.

II. Assessment of separability

From Table IV, one can assess the separability of the given samples by comparing the R_E indices at the different pH_L values. According to our experience, when the R_E values of two samples coincided within *ca.* 0.15, their separation was difficult or impossible. The linear interpolation of adjacent R_E values gives approximate R_E values at any pH_L using the same buffer: *e.g.*, the R_E values of acetic acid at pH_L 3 and 3.25 are 8.876 and 7.444, then the R_E value at pH_L 3.125 can be estimated as $8.876 - (8.876 - 7.444) \times 0.125/0.25 = 8.160$. The exactly simulated value is 8.090 and the discrepancy is thus slight.

For the estimation of pH_L and the selection of a terminator for optimum separation, the pH_L should be selected so as to result in sufficient difference between the R_E values of the samples. A plot of R_E against pH_L is useful for this purpose. Fig. 1 shows the dependence on pH_L of the R_E values of benzoate (No. 12 in Table IV), cacodylate (27), caproate (29), chlorate (32), citrate (57), phenolate (229) and N-tris(hydroxymethyl)methyl-3-aminopropanesulphonic acid (TAPS) (262). In the range




Fig. 1. Effect of pH_L and the buffers on the simulated R_E values of chlorate, citrate, benzoate, acetate, caproate, cacodylate, TAPS and phenolate ions at the isotachophoretically steady state. The leading ion was 10 mM Cl^- . The buffers used were β -alanine ($pH_L = 3-4$), ϵ -aminocaproic acid (4-4.75), creatinine (4.75-5.25), histidine (5.5-6.25), imidazole (6.5-7.25), tris(hydroxymethyl)aminomethane (7.5-8.25), amediol (8.5-9.25), and ethanolamine (9.5-10).

pH_L 5.5-6.25 histidine buffer, chlorate, citrate, acetate, benzoate and caproate ions can easily be separated using cacodylate ion as terminator. At intermediate pH_L values of 4.5-4.75 (ϵ -aminocaproic acid buffer), benzoate and acetate ions may form a mixed zone. In the low pH_L range 3-4 the samples can easily be separated, however, cacodylate ion is not suitable as the terminator as the R_E values are too large; caprylate (30) may be suitable. The separability is strongly dependent on the sample amount. For the separation of large amounts of samples, a high loading of leading ion is necessary, even when the difference between R_E values is larger than *ca.* 0.15 (the latter is only a measure of the separation).

III. Qualitative identification

For the identification of unknown samples, the method of standard addition has usually been employed. Instead qualitative identification can be achieved by comparing the observed and simulated R_E values, if the R_E values are measured carefully using an internal standard. At least, the possible candidates for the components in actual samples can be limited to several kinds, and their number can be reduced further by considering the R_E values measured at other pH_L values. This procedure can be performed rapidly by computer-sorting of samples having similar R_E values to those observed.

IV. Quantitation

When the separated samples are identified, their amounts can be determined




Fig. 2. Effect of pH_L and the buffers on the simulated PU values of chlorate, citrate, lactate, benzoate and acetate ions at the isotachophoretically steady state. The terminator is caproate ion. For the electrolyte systems, see Fig. 1.

using the simulated and observed zone lengths. In Table IV the simulated time-based zone lengths, t (sec), are listed for 10-nmol samples at a driving current of $100 \mu\text{A}$. If the observed time-based zone length is t_{obs} (sec) at a driving current of I (μA), the sample amount, n_s (nmol), can be expressed:

$$n_s = t_{\text{obs}} I / 10 t \quad (3)$$

For example, at a driving current of $50 \mu\text{A}$ and pH_L 6, the time-based zone length of acetic acid was 50 sec. The sample amount can easily be estimated as $50 \times 50 / 10 \times 16.22 = 15.4$ nmol, and the slope of the calibration line, $I/10t$, is 0.308.

V. Conversion of R_E into different qualitative indices

Besides R_E , several qualitative indices have been proposed such as relative step height⁴, IRM^6 , PU^7 , PR^8 , RU ($= \text{IRM} \times 100$)⁹ and SU ($= \text{PU} \times 100$)⁹. The definition of IRM is the same as R_E and $\text{PR} = 1/R_E$. PU is defined as

$$\text{PU} = (h_s - h_L) / (h_T - h_L) \quad (4)$$

where h_T is the step height of the terminating zone. The R_E values of samples can be converted into PU values using the R_E value of the sample, $R_E(\text{S})$, and terminating ion, $R_E(\text{T})$:

$$\text{PU} = [R_E(\text{S}) - 1] / [R_E(\text{T}) - 1] \quad (5)$$

Fig. 2 shows the pH dependence of PU values for acetate (1), benzoate (12), chlorate (32), citrate (57) and lactate (184) using R_E values of caproate ion (29) as the terminator. A similar conversion of the relative step heights of 65 anions⁴ to R_E values has been reported in ref. 5.

Thus, the simulated qualitative and quantitative indices can be utilized for several analytical purposes, although the number of simulated samples is limited. It should be noted that the accuracy of the simulated indices depends mainly on the physico-chemical constants of the samples and buffers used. The present table is as yet in an experimental stage, but by continuous revision it will be used widely in future as a data base for isotachopheresis. We await the critical comments of users.

Similar tables for amino acids and cations will be submitted in due course.

ACKNOWLEDGEMENTS

The authors wish to express their thanks to Shinji Kobayashi and Toshihiko Gojo at Hiroshima University for their help with the experiments.

REFERENCES

- 1 T. Hirokawa and Y. Kiso, *J. Chromatogr.*, 242 (1982) 227.
- 2 T. Hirokawa and Y. Kiso, *J. Chromatogr.*, 257 (1983) 197.
- 3 T. Hirokawa and Y. Kiso, *J. Chromatogr.*, (1983) in press.
- 4 F. M. Everaerts, J. L. Beckers and Th. P. E. M. Verheggen, *Isotachopheresis*, Elsevier, Amsterdam, 1976.
- 5 T. Hirokawa and Y. Kiso, *J. Chromatogr.*, 252 (1982) 49.
- 6 M. Deml, P. Boček and J. Janak, *J. Chromatogr.*, 109 (1975) 49.
- 7 H. Miyazaki and K. Katoh, *J. Chromatogr.*, 119 (1976) 369.
- 8 H. Yoshida, I. Nukatsuka and S. Hikime, *Bunseki Kagaku (Jap. Anal.)*, 28 (1979) 382.
- 9 F. M. Everaerts, F. E. P. Mikkers and Th. P. E. M. Verheggen, *2nd International Symposium on Isotachopheresis*, Eindhoven, 1980.

(For Tables III and IV, see pp. D8–D106; references to Table IV are listed on p. D106)

TABLE III
ANIONIC SUBSTANCES LISTED ALPHABETICALLY IN TABLE IV

No.	Samples	No.	Samples
1	ACETIC ACID (S)	49	2-CHLOROPROPIONIC ACID
2	ACRYLIC ACID	50	3-CHLOROPROPIONIC ACID
3	ADIPIC ACID	51	CHLOROUS ACID
4	ALLYLACETIC ACID	52	5-CHLOROVALERIC ACID
5	o-AMINOBENZOIC ACID	53	CHROMIC ACID
6	p-AMINOBENZOIC ACID	54	CINNAMIC ACID
7	ARSENIC ACID	55	cis-CINNAMIC ACID
8	ASPARTIC ACID	56	CITRACONIC ACID
9	ASPIRIN	57	CITRIC ACID (S)
10	AZELAIC ACID	58	CRESOL
11	BENZENESULFONIC ACID	59	m-CRESOL
12	BENZOIC ACID (S)	60	p-CRESOL
13	BENZYLASPARATIC ACID	61	CROTONIC ACID
14	BROMIC ACID (S)	62	CYANIC ACID
15	BROMOACETIC ACID (S)	63	CYANOACETIC ACID
16	m-BROMOBENZOIC ACID	64	m-CYANOBENZOIC ACID
17	o-BROMOBENZOIC ACID	65	CYCLOBUTANE-1,1-DICARBOXYLIC ACID
18	p-BROMOBENZOIC ACID	66	CYCLOHEXANE-1,1-DICARBOXYLIC ACID
19	2-BROMOBUTYRIC ACID	67	CYCLOPENTANE-1,1-DICARBOXYLIC ACID
20	4-BROMOBUTYRIC ACID	68	DECYLSULFONIC ACID
21	o-BROMOPHENYLACETIC ACID	69	DIBROMOFLUOROACETIC ACID
22	2-BROMOPROPIONIC ACID	70	2,3-DIBROMOPROPIONIC ACID
23	5-BROMOVALERIC ACID	71	DICHLOROACETIC ACID (S)
24	o-t-BUTYLBENZOIC ACID	72	DICHLOROFLUOROACETIC ACID
25	p-t-BUTYLBENZOIC ACID	73	2,4-DICHLOROPHENOL
26	BUTYRIC ACID (S)	74	2,4-DICHLOROPHENOXYACETIC ACID
27	CACODYLIC ACID	75	DICHROMIC ACID
28	CAPRIC ACID	76	DIETHYLACETIC ACID
29	CAPROIC ACID (S)	77	DIETHYLMALONIC ACID
30	CAPRYLIC ACID	78	DIFLUOROACETIC ACID
31	CARBONIC ACID	79	2,4-DIHYDROXYBENZOIC ACID
32	CHLORIC ACID (S)	80	2,5-DIHYDROXYBENZOIC ACID
33	2-CHLORO-3-OH-BUTYRIC ACID	81	2,6-DIHYDROXYBENZOIC ACID
34	CHLOROACETIC ACID	82	3,4-DIHYDROXYBENZOIC ACID
35	m-CHLOROBENZOIC ACID	83	3,5-DIHYDROXYBENZOIC ACID
36	o-CHLOROBENZOIC ACID	84	DIHYDROXYTARTARIC ACID
37	p-CHLOROBENZOIC ACID	85	2,3-DIMETHYLBENZOIC ACID
38	2-CHLOROBUTYRIC ACID	86	2,4-DIMETHYLBENZOIC ACID
39	3-CHLOROBUTYRIC ACID	87	2,5-DIMETHYLBENZOIC ACID
40	2-CHLOROCROTONIC ACID	88	2,6-DIMETHYLBENZOIC ACID
41	3-CHLOROCROTONIC ACID	89	3,4-DIMETHYLBENZOIC ACID
42	CHLORODIBROMOACETIC ACID	90	3,5-DIMETHYLBENZOIC ACID
43	CHLORODIFLUOROACETIC ACID	91	DIMETHYLMALONIC ACID
44	2-CHLOROISOCROTONIC ACID	92	2,6-DINITROBENZOIC ACID
45	m-CHLOROPHENOL	93	3,5-DINITROBENZOIC ACID
46	o-CHLOROPHENOL	94	2,3-DINITROPHENOL
47	p-CHLOROPHENOL	95	2,4-DINITROPHENOL
48	p-CHLOROPHENYLACETIC ACID	96	2,5-DINITROPHENOL

No.	Samples	No.	Samples
97	2,6-DINITROPHENOL	145	3-HYDROXY-o-TOLUIC ACID
98	3,4-DINITROPHENOL	146	3-HYDROXY-p-TOLUIC ACID
99	3,5-DINITROPHENOL	147	4-HYDROXY-m-TOLUIC ACID
100	DIPHENYLACETIC ACID	148	6-HYDROXY-o-TOLUIC ACID
101	DIPROPYLMALONIC ACID	149	m-HYDROXYBENZOIC ACID
102	DITHIONIC ACID	150	p-HYDROXYBENZOIC ACID
103	DITHIONOUS ACID	151	2-HYDROXYBUTYRIC ACID
104	DODECYLSULFONIC ACID	152	3-HYDROXYBUTYRIC ACID
105	ENANTHYLIC ACID	153	4-HYDROXYBUTYRIC ACID
106	o-ETHOXYBENZOIC ACID	154	m-HYDROXYCINNAMIC ACID
107	p-ETHOXYBENZOIC ACID	155	o-HYDROXYCINNAMIC ACID
108	o-ETHYLBENZOIC ACID	156	p-HYDROXYCINNAMIC ACID
109	p-ETHYLBENZOIC ACID	157	2-HYDROXYISOBUTYRIC ACID (S)
110	2-ETHYLBUTYRIC ACID	158	p-HYDROXYPHENYLACETIC ACID
111	ETHYLFUMARIC ACID	159	HYDROXYTRIMETHYLACETIC ACID
112	3-ETHYLGUTARIC ACID	160	2-HYDROXYVALERIC ACID
113	ETHYLMALEIC ACID	161	4-HYDROXYVALERIC ACID
114	ETHYLMALONIC ACID	162	HYPOPHOSPHOROUS ACID (S)
115	ETHYLPROPYLMALONIC ACID	163	IODIC ACID
116	ETHYLSULFONIC ACID	164	IDOACETIC ACID
117	FLUOROACETIC ACID	165	m-IODOBENZOIC ACID
118	m-FLUOROBENZOIC ACID	166	o-IODOBENZOIC ACID
119	o-FLUOROBENZOIC ACID	167	p-IODOBENZOIC ACID
120	p-FLUOROBENZOIC ACID	168	4-IODOBUTYRIC ACID
121	FORMIC ACID (S)	169	m-IODOPHENYLACETIC ACID
122	FUMARIC ACID	170	o-IODOPHENYLACETIC ACID
123	GALLIC ACID	171	p-IODOPHENYLACETIC ACID
124	GLUCONIC ACID	172	3-IODOPROPIONIC ACID
125	GLUCURONIC ACID	173	5-IDOVALERIC ACID
126	GLUTACONIC ACID	174	ISOBUTYRIC ACID
127	GLUTAMIC ACID (S)	175	ISOCAPROIC ACID
128	GLUTARIC ACID	176	ISOCITRIC ACID
129	GLYCERIC ACID	177	ISOPHTHALIC ACID
130	GLYCOLIC ACID	178	o-ISOPROPYLBENZOIC ACID
131	GLYOXYLIC ACID	179	p-ISOPROPYLBENZOIC ACID
132	HIPPURIC ACID	180	ISOPROPYLMALONIC ACID
133	HYDROBROMIC ACID	181	ISOVALERIC ACID
134	HYDROCHLORIC ACID	182	ITACONIC ACID
135	HYDROCYANIC ACID	183	2-KETOGLUTARIC ACID
136	HYDROFLUORIC ACID (S)	184	LACTIC ACID (S)
137	HYDROGEN AZIDE	185	LAURYL SULFONIC ACID
138	HYDROGEN HEXAFLUOROPHOSPHIDE	186	LEVULINIC ACID
139	HYDROGEN SULFIDE	187	MALEIC ACID
140	HYDRIODIC ACID	188	MALIC ACID
141	2-HYDROXY-3-CHLOROBUTYRIC A.	189	MALONIC ACID (S)
142	2-HYDROXY-3-Cl-ISOBUTYRIC A.	190	MANDELIC ACID
143	2-HYDROXY-3-Cl-PROPIONIC A.	191	MES (S)
144	2-HYDROXY-m-TOLUIC ACID	192	MESACONIC ACID

No.	Samples	No.	Samples
193	MESOTARTARIC ACID	241	3-PROPYLGLUTARIC ACID
194	METHACRYLIC ACID	242	PROPYLMALONIC ACID
195	METHOXYACETIC ACID	243	PROPYLSULFONIC ACID
196	m-METHOXYBENZOIC ACID	244	PYRAZOLE-2,3-DICARBOXYLIC ACID
197	o-METHOXYBENZOIC ACID	245	PYROPHOSPHORIC ACID
198	p-METHOXYBENZOIC ACID	246	PYRUVIC ACID
199	p-METHOXYPHENYLACETIC ACID	247	SACCHARIN
200	METHYLETHYLMALONIC ACID	248	SALICYLIC ACID
201	METHYLMALONIC ACID	249	SEBACIC ACID
202	METHYLSULFONIC ACID	250	SELENIC ACID
203	2-NAPHTALENESULFONIC ACID	251	SELENIOS ACID
204	NICOTINIC ACID	252	SELENOCYANIC ACID
205	NITRIC ACID	253	SORBIC ACID
206	2-NITRO-3-BROMOBENZOIC ACID	254	SUBERIC ACID
207	3-NITRO-2-BROMOBENZOIC ACID	255	SUCCINIC ACID (S)
208	3-NITRO-4-BROMOBENZOIC ACID	256	SULFAMIC ACID (S)
209	2-NITRO-3-CHLOROBENZOIC ACID	257	SULFANILIC ACID
210	3-NITRO-4-CHLOROBENZOIC ACID	258	m-SULFANILIC ACID
211	3-NITRO-5-CHLOROBENZOIC ACID	259	o-SULFANILIC ACID
212	m-NITROBENZOIC ACID	260	SULFURIC ACID
213	o-NITROBENZOIC ACID	261	SULFUROUS ACID
214	p-NITROBENZOIC ACID	262	TAPS
215	m-NITROPHENOL	263	TARTARIC ACID (S)
216	o-NITROPHENOL	264	TARTRONIC ACID
217	p-NITROPHENOL	265	TEREPHTHALIC ACID
218	NITROUS ACID	266	THIOACETIC ACID
219	OCTYLSULFONIC ACID	267	THIOCYANIC ACID
220	OROTIC ACID	268	THIOCYANOACETIC ACID
221	OXALIC ACID (S)	269	THIOGLYCOLIC ACID
222	OXALOACETIC ACID	270	THIOSULFURIC ACID
223	PELARGONIC ACID	271	p-TOLUENESULFONIC ACID
224	PERCHLORIC ACID (S)	272	m-TOLUIC ACID
225	PERIODIC ACID	273	o-TOLUIC ACID
226	PERMANGANIC ACID	274	p-TOLUIC ACID
227	PEROXYSULFURIC ACID	275	TRIBROMOACETIC ACID
228	PERRHENIC ACID	276	TRICHLOROACETIC ACID
229	PHENOL	277	TRICHLOROACRYLIC ACID
230	PHENOXYACETIC ACID	278	TRICHLOROLACTIC ACID
231	PHENYLACETIC ACID	279	TRIMETAPHOSPHORIC ACID
232	4-PHENYLBUTYRIC ACID	280	TRIMETHYLACETIC ACID (S)
233	2-PHENYLPROPIONIC ACID	281	TRIMETHYLACRYLIC ACID
234	3-PHENYLPROPIONIC ACID	282	2,4,6-TRIMETHYLBENZOIC ACID
235	PHOSPHORIC ACID (S)	283	TRIMETYL PYRUVIC ACID
236	PHOSPHOROUS ACID	284	VALERIC ACID
237	PTHALIC ACID	285	VANILLIC ACID
238	PICRIC ACID	286	VINYLAACETIC ACID
239	PIMELIC ACID	287	VINYLGLYCOLIC ACID
240	PROPIONIC ACID (S)		

TABLE IV
ISOTACHOPHORETIC INDICES OF 287 ANIONIC SUBSTANCES

1	ACETIC ACID (S)				C2H4O2					
-1	42.4	4.756	60.05	pHL	3.00	3.25	3.50			
				RE	8.876	7.444	6.370			
				t	21.01	20.13	19.51			
				Z	0.206	0.247	0.289			
				pHS	4.151	4.249	4.342			
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
5.428	4.576	3.848	3.485	3.111	2.763	2.650	2.427	2.244	2.087	2.027
19.02	18.64	16.48	16.37	16.26	16.18	18.16	18.12	18.09	16.23	16.22
0.340	0.404	0.485	0.537	0.603	0.681	0.711	0.778	0.843	0.912	0.940
4.443	4.561	4.700	4.789	4.904	5.050	5.112	5.264	5.449	5.733	5.910
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
1.980	1.949	1.925	1.915	1.909	1.904	1.912	1.910	1.909	1.908	1.909
16.22	16.21	21.62	21.61	21.61	21.61	16.18	16.18	16.19	16.20	16.20
0.962	0.978	0.988	0.993	0.996	0.998	0.999	0.999	1.000	1.000	1.000
6.122	6.355	6.651	6.887	7.132	7.380	7.593	7.841	8.091	8.340	8.591
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 1				
1.908	1.907	1.906	1.902	1.901	1.899					
16.23	16.27	16.34	20.09	20.36	20.85					
1.000	1.000	1.000	1.000	1.000	1.000					
8.842	9.092	9.343	9.626	9.878	10.133					
2	ACRYLIC ACID				C3H4O2					
-1	42.7	4.258	72.06	pHL	3.00	3.25	3.50			
				RE	5.663	4.816	4.183			
				t	21.28	20.26	19.56			
				Z	0.324	0.382	0.441			
				pHS	3.914	4.023	4.127			
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
3.635	3.153	2.863	2.638	2.422	2.241	2.206	2.093	2.012	1.963	1.936
19.00	18.59	16.48	16.34	16.21	16.12	18.11	18.06	18.03	16.18	16.17
0.509	0.588	0.653	0.710	0.775	0.839	0.853	0.900	0.937	0.964	0.978
4.244	4.382	4.498	4.611	4.758	4.938	4.983	5.173	5.390	5.650	5.865
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
1.918	1.907	1.896	1.893	1.890	1.888	1.896	1.896	1.895	1.893	1.895
16.16	16.16	21.53	21.53	21.52	21.52	16.13	16.13	16.14	16.15	16.15
0.987	0.993	0.996	0.998	0.999	0.999	1.000	1.000	1.000	1.000	1.000
6.100	6.344	6.634	6.879	7.128	7.377	7.590	7.839	8.089	8.339	8.590
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 2 , 3				
1.894	1.893	1.892	1.889	1.887	1.885					
16.18	16.22	16.29	20.01	20.27	20.76					
1.000	1.000	1.000	1.000	1.000	1.000					
8.840	9.091	9.341	9.624	9.877	10.131					
3	ADIPIC ACID				C6H10O4					
-1	24.6	4.430	146.14	pHL	3.00	3.25	3.50			
-2	52.4	5.277		RE	9.612	7.981	6.740			
				t	31.81	30.07	29.04			
				Z	0.335	0.404	0.480			
				pHS	4.042	4.147	4.248			
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
5.657	4.689	4.086	3.601	3.124	2.698	2.587	2.307	2.086	1.918	1.840
28.45	28.28	25.15	25.48	26.02	26.77	30.54	31.29	32.03	29.61	29.95
0.574	0.695	0.807	0.919	1.062	1.234	1.286	1.446	1.602	1.764	1.840
4.359	4.489	4.599	4.705	4.839	5.002	5.055	5.223	5.419	5.692	5.882
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
1.783	1.745	1.708	1.696	1.687	1.680	1.703	1.701	1.698	1.696	1.699
30.22	30.39	40.01	40.06	40.07	40.06	30.61	30.62	30.62	30.63	30.66
1.900	1.941	1.968	1.982	1.990	1.994	1.996	1.998	1.999	1.999	2.000
6.102	6.338	6.625	6.863	7.108	7.356	7.577	7.826	8.076	8.325	8.576
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 10 , 1				
1.697	1.695	1.693	1.682	1.678	1.675					
30.69	30.75	30.85	37.38	37.77	38.49					
2.000	2.000	2.000	2.000	2.000	2.000					
8.826	9.076	9.327	9.603	9.854	10.107					

4	ALLYLACETIC ACID -1			33.9	4.674	C5H8O2 100.12		pHL RE t Z pHS	3.00 10.017 24.29 0.230 4.130	3.25 8.409 23.04 0.275 4.232	3.50 7.202 22.18 0.322 4.328
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75	5.75
6.150	5.208	4.527	4.100	3.673	3.288	3.176	2.930	2.736	2.595	2.530	2.530
21.51	21.01	18.32	18.17	18.03	17.93	20.44	20.39	20.35	18.01	18.00	18.00
0.378	0.448	0.521	0.576	0.645	0.723	0.749	0.814	0.873	0.928	0.952	0.952
4.434	4.558	4.681	4.777	4.902	5.058	5.116	5.281	5.477	5.745	5.936	5.936
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50	8.50
2.483	2.451	2.423	2.413	2.407	2.402	2.418	2.416	2.414	2.412	2.415	2.415
17.99	17.99	24.82	24.82	24.81	24.81	17.95	17.96	17.96	17.97	17.99	17.99
0.971	0.983	0.991	0.995	0.997	0.998	0.999	0.999	1.000	1.000	1.000	1.000
6.157	6.394	6.702	6.942	7.188	7.437	7.635	7.884	8.134	8.384	8.636	8.636
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 1 , 3					
2.413	2.411	2.408	2.402	2.398	2.394						
18.03	18.09	18.19	23.02	23.47	24.34						
1.000	1.000	1.000	1.000	1.000	1.000						
8.886	9.137	9.388	9.686	9.941	10.199						
5	o-AMINOBENZOIC ACID -1			31.6 -31.6	2.108 4.939	C7H7NO2 137.10		pHL RE t Z pHS	3.00 13.704 25.78 0.180 4.282	3.25 11.444 24.32 0.216 4.374	3.50 9.740 23.32 0.254 4.462
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75	5.75
8.245	6.895	5.799	5.208	4.602	4.038	3.835	3.472	3.169	2.909	2.808	2.808
22.54	21.95	19.01	18.84	18.70	18.58	21.27	21.22	21.18	18.67	18.66	18.66
0.301	0.362	0.435	0.485	0.551	0.631	0.665	0.736	0.809	0.890	0.923	0.923
4.559	4.674	4.802	4.888	5.000	5.141	5.206	5.353	5.533	5.810	5.980	5.980
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50	8.50
2.727	2.672	2.623	2.606	2.595	2.588	2.607	2.603	2.601	2.598	2.601	2.601
18.65	18.65	26.00	25.99	25.99	25.98	18.61	18.61	18.62	18.63	18.65	18.65
0.951	0.970	0.985	0.991	0.995	0.997	0.998	0.999	0.999	1.000	1.000	1.000
6.188	6.417	6.730	6.965	7.209	7.457	7.652	7.900	8.149	8.399	8.651	8.651
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 10 , 1					
2.599	2.596	2.593	2.585	2.581	2.576						
18.69	18.76	18.88	24.12	24.66	25.72						
1.000	1.000	1.000	1.000	1.000	1.000						
8.901	9.152	9.404	9.707	9.963	10.223						
6	p-AMINOBENZOIC ACID -1			31.6 -31.6	2.413 4.853	C7H7NO2 137.10		pHL RE t Z pHS	3.00 12.621 26.28 0.196 4.259	3.25 10.570 24.68 0.234 4.346	3.50 9.018 23.57 0.275 4.432
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75	5.75
7.653	6.422	5.459	4.915	4.361	3.849	3.678	3.349	3.081	2.865	2.775	2.775
22.70	22.05	19.07	18.88	18.72	18.59	21.28	21.22	21.18	18.67	18.66	18.66
0.325	0.389	0.463	0.515	0.583	0.663	0.694	0.764	0.833	0.904	0.934	0.934
4.528	4.643	4.767	4.856	4.972	5.117	5.179	5.333	5.518	5.791	5.969	5.969
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50	8.50
2.706	2.659	2.617	2.603	2.593	2.587	2.606	2.603	2.600	2.598	2.601	2.601
18.65	18.65	26.00	25.99	25.99	25.98	18.61	18.61	18.62	18.63	18.65	18.65
0.959	0.975	0.988	0.993	0.996	0.998	0.999	0.999	1.000	1.000	1.000	1.000
6.181	6.414	6.727	6.963	7.209	7.457	7.651	7.900	8.149	8.399	8.651	8.651
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 10 , 1					
2.599	2.596	2.593	2.585	2.581	2.576						
18.69	18.76	18.88	24.12	24.66	25.72						
1.000	1.000	1.000	1.000	1.000	1.000						
8.901	9.152	9.404	9.707	9.963	10.223						

7	ARSENIC ACID				H3AsO4		pHL	3.00	3.25	3.50
	-1	35.2		2.190	141.94		RE	2.463	2.407	2.370
	-2	70.0		6.940			t	27.10	24.32	22.57
							Z	0.936	0.959	0.974
							pHS	3.318	3.513	3.713
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
2.344	2.326	2.336	2.324	2.312	2.297	2.293	2.272	2.238	2.186	2.104
21.43	20.74	18.38	18.03	17.85	17.81	20.28	20.38	20.62	18.92	19.64
0.985	0.992	0.993	0.998	1.003	1.009	1.010	1.020	1.036	1.069	1.114
3.930	4.162	4.171	4.396	4.634	4.879	4.905	5.152	5.400	5.690	5.928
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
1.993	1.857	1.619	1.527	1.440	1.367	1.321	1.291	1.270	1.257	1.251
20.64	21.86	30.76	31.62	32.45	33.13	26.77	27.04	27.24	27.36	27.44
1.182	1.277	1.479	1.577	1.682	1.779	1.866	1.914	1.948	1.970	1.983
6.162	6.392	6.768	6.935	7.127	7.340	7.598	7.811	8.044	8.287	8.537
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3 , 4				
1.246	1.243	1.241	1.234	1.232	1.231					
27.50	27.56	27.64	32.36	32.58	32.97					
1.990	1.994	1.997	1.998	1.999	1.999					
8.784	9.032	9.282	9.541	9.792	10.042					
8	ASPARTIC ACID				C4H7NO4		pHL	3.00	3.25	3.50
	1	31.6		1.990	133.10		RE	5.614	4.865	4.305
	-1	-31.6		3.900			t	26.69	24.86	23.61
	-2	-51.8		10.002			Z	0.449	0.520	0.589
							pHS	3.807	3.922	4.038
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
3.832	3.437	3.311	3.107	2.932	2.801	2.781	2.703	2.653	2.640	2.623
22.67	22.00	19.19	18.92	18.71	18.57	21.32	21.24	21.20	18.68	18.66
0.664	0.743	0.777	0.830	0.881	0.923	0.929	0.956	0.974	0.985	0.991
4.173	4.334	4.412	4.555	4.734	4.943	4.982	5.201	5.436	5.674	5.909
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.611	2.603	2.589	2.585	2.581	2.576	2.594	2.587	2.574	2.553	2.523
18.65	18.65	26.02	26.02	26.04	26.06	18.72	18.81	18.96	19.23	19.71
0.995	0.997	0.999	1.000	1.002	1.003	1.006	1.010	1.018	1.032	1.057
6.153	6.400	6.709	6.957	7.206	7.456	7.653	7.903	8.153	8.403	8.669
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 11 , 1				
2.470	2.391	2.287	2.131	2.014	1.911					
20.45	21.57	23.09	31.05	33.26	35.62					
1.098	1.161	1.253	1.400	1.531	1.661					
8.917	9.162	9.405	9.699	9.925	10.157					
9	ASPIRIN				C9H8O4		pHL	3.00	3.25	3.50
	-1	26.7		3.485	180.16		RE	4.917	4.415	4.042
							t	30.70	28.20	26.51
							Z	0.617	0.689	0.755
							pHS	3.663	3.800	3.942
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
3.740	3.506	3.488	3.359	3.260	3.193	3.180	3.142	3.117	3.127	3.119
25.29	24.45	21.20	20.81	20.53	20.36	23.65	23.55	23.49	20.49	20.47
0.818	0.875	0.886	0.921	0.950	0.970	0.972	0.984	0.991	0.994	0.997
4.105	4.296	4.340	4.517	4.725	4.954	4.991	5.226	5.470	5.698	5.941
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
3.112	3.106	3.088	3.086	3.083	3.079	3.112	3.110	3.107	3.104	3.109
20.46	20.45	29.23	29.22	29.21	29.20	20.41	20.42	20.42	20.44	20.47
0.998	0.999	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
6.189	6.438	6.755	7.005	7.254	7.504	7.688	7.937	8.187	8.438	8.690
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 11 , 3				
3.106	3.102	3.098	3.084	3.078	3.069					
20.53	20.62	20.80	27.20	28.05	29.79					
1.000	1.000	1.000	1.000	1.000	1.000					
8.941	9.192	9.444	9.762	10.020	10.287					

10	AZELAIC ACID			C9H16O4			pHL	3.00	3.25	3.50
	-1	22.0	4.550	188.22			RE	11.739	9.723	8.186
	-2	45.9	5.333				t	34.49	32.49	31.28
							Z	0.307	0.373	0.444
							pHS	4.110	4.213	4.311
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
6.845	5.647	4.905	4.307	3.719	3.195	3.048	2.705	2.434	2.233	2.138
30.56	30.31	26.70	27.00	27.52	28.26	32.55	33.33	34.13	31.30	31.69
0.534	0.651	0.761	0.871	1.014	1.189	1.245	1.410	1.575	1.745	1.827
4.420	4.546	4.652	4.754	4.883	5.041	5.096	5.260	5.452	5.720	5.908
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.067	2.020	1.973	1.958	1.947	1.939	1.971	1.968	1.964	1.961	1.965
31.98	32.18	43.29	43.33	43.35	43.34	32.44	32.44	32.45	32.46	32.50
1.892	1.936	1.966	1.980	1.989	1.994	1.996	1.998	1.999	1.999	2.000
6.127	6.362	6.657	6.896	7.141	7.389	7.602	7.850	8.100	8.349	8.601
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 10 , 1				
1.963	1.960	1.956	1.941	1.936	1.931					
32.54	32.61	32.74	40.34	40.86	41.85					
2.000	2.000	2.000	2.000	2.000	2.000					
8.851	9.101	9.352	9.636	9.888	10.142					
11	BENZENESULFONIC ACID			C6H6O3S			pHL	3.00	3.25	3.50
	-1	38.7	-2.000	158.17			RE	2.093	2.093	2.093
							t	25.93	23.10	21.37
							Z	1.000	1.000	1.000
							pHS	3.239	3.453	3.669
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
2.092	2.092	2.100	2.100	2.099	2.097	2.096	2.094	2.093	2.101	2.101
20.29	19.64	17.51	17.16	16.95	16.83	19.05	18.99	18.95	16.90	16.89
1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
3.896	4.134	4.137	4.369	4.611	4.857	4.878	5.126	5.375	5.609	5.858
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.100	2.098	2.093	2.092	2.091	2.090	2.102	2.101	2.100	2.098	2.101
16.89	16.88	22.83	22.83	22.83	22.82	16.85	16.85	16.86	16.87	16.88
1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
6.107	6.357	6.651	6.901	7.151	7.401	7.607	7.857	8.107	8.357	8.608
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 2 , 12				
2.099	2.098	2.096	2.092	2.090	2.087					
16.91	16.95	17.04	21.19	21.52	22.14					
1.000	1.000	1.000	1.000	1.000	1.000					
8.859	9.109	9.360	9.650	9.903	10.159					
12	BENZOIC ACID			C7H6O2			pHL	3.00	3.25	3.50
	-1	33.6	4.203	122.12			RE	6.683	5.705	4.971
							t	24.91	23.46	22.47
							Z	0.352	0.413	0.476
							pHS	3.913	4.025	4.134
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
4.339	3.791	3.525	3.255	3.004	2.801	2.762	2.637	2.551	2.509	2.480
21.72	21.16	18.49	18.29	18.12	18.00	20.56	20.50	20.46	18.09	18.08
0.547	0.628	0.681	0.739	0.803	0.863	0.875	0.917	0.949	0.971	0.982
4.256	4.400	4.500	4.622	4.778	4.967	5.012	5.212	5.455	5.683	5.905
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.460	2.448	2.432	2.428	2.425	2.422	2.440	2.438	2.436	2.434	2.437
18.07	18.07	24.97	24.96	24.96	24.95	18.03	18.04	18.04	18.06	18.07
0.990	0.994	0.997	0.998	0.999	0.999	1.000	1.000	1.000	1.000	1.000
6.144	6.389	6.694	6.940	7.189	7.438	7.636	7.886	8.136	8.386	8.637
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 2 , 1				
2.436	2.433	2.431	2.424	2.421	2.417					
18.11	18.17	18.28	23.16	23.62	24.51					
1.000	1.000	1.000	1.000	1.000	1.000					
8.888	9.138	9.390	9.689	9.944	10.202					

CHROMATOGRAPHIC DATA (1983)

D15

13	BENZYLASPARATIC ACID				C11H13NO4		pHL	3.00	3.25	3.50
	1	25.8	-3.000		223.23		RE	14.983	12.525	10.669
	-1	-25.8	4.855				t	29.33	27.56	26.35
							Z	0.203	0.243	0.287
							pHS	4.243	4.344	4.439
	3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50
	9.051	7.605	6.621	5.944	5.269	4.659	4.463	4.070	3.758	3.531
	25.44	24.77	21.23	21.03	20.86	20.73	24.07	24.01	23.98	20.89
	0.339	0.406	0.471	0.527	0.597	0.678	0.708	0.778	0.845	0.909
	4.544	4.666	4.779	4.874	4.996	5.147	5.209	5.369	5.560	5.822
	6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25
	3.342	3.288	3.233	3.217	3.206	3.198	3.231	3.227	3.224	3.219
	20.86	20.86	29.96	29.95	29.94	29.93	20.82	20.83	20.84	20.85
	0.962	0.978	0.989	0.994	0.996	0.998	0.999	0.999	1.000	1.000
	6.224	6.459	6.782	7.021	7.267	7.515	7.698	7.946	8.196	8.446
	8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 10			
	3.221	3.217	3.212	3.197	3.190	3.180				
	20.95	21.05	21.24	27.92	28.85	30.80				
	1.000	1.000	1.000	1.000	1.000	1.000				
	8.950	9.201	9.453	9.774	10.033	10.301				
14	BROMIC ACID (S)				HBrO3		pHL	3.00	3.25	3.50
	-1	57.8	-2.000		128.91		RE	1.381	1.381	1.381
							t	20.82	18.65	17.38
							Z	1.000	1.000	1.000
							pHS	3.101	3.333	3.567
	3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50
	1.380	1.380	1.382	1.382	1.382	1.381	1.381	1.381	1.380	1.383
	16.62	16.19	14.80	14.55	14.41	14.34	15.80	15.76	15.73	14.38
	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
	3.806	4.051	4.053	4.295	4.542	4.790	4.799	5.048	5.298	5.541
	6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25
	1.382	1.382	1.381	1.380	1.380	1.380	1.383	1.382	1.382	1.382
	14.37	14.37	18.28	18.28	18.29	18.29	14.34	14.35	14.35	14.36
	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
	6.040	6.290	6.559	6.809	7.059	7.309	7.540	7.790	8.040	8.290
	8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 1 , 12			
	1.382	1.382	1.381	1.380	1.380	1.379				
	14.38	14.40	14.44	17.11	17.25	17.51				
	1.000	1.000	1.000	1.000	1.000	1.000				
	8.791	9.041	9.291	9.556	9.807	10.059				
15	BROMOACETIC ACID (S)				C2H3BrO2		pHL	3.00	3.25	3.50
	-1	40.6	2.901		138.95		RE	2.556	2.374	2.247
							t	23.93	21.93	20.62
							Z	0.773	0.834	0.882
							pHS	3.398	3.565	3.738
	3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50
	2.154	2.089	2.090	2.054	2.030	2.014	2.012	2.003	1.997	2.002
	19.72	19.13	17.07	16.77	16.58	16.47	18.58	18.52	18.49	16.54
	0.922	0.951	0.954	0.971	0.983	0.990	0.990	0.995	0.997	0.998
	3.933	4.150	4.176	4.384	4.613	4.854	4.876	5.119	5.366	5.603
	6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25
	1.998	1.996	1.991	1.990	1.989	1.988	1.999	1.998	1.997	1.996
	16.52	16.52	22.18	22.18	22.17	22.17	16.49	16.49	16.50	16.51
	0.999	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
	6.099	6.348	6.639	6.889	7.139	7.389	7.598	7.848	8.098	8.348
	8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3 , 1			
	1.997	1.995	1.994	1.990	1.988	1.986				
	16.54	16.58	16.66	20.60	20.89	21.44				
	1.000	1.000	1.000	1.000	1.000	1.000				
	8.850	9.100	9.351	9.637	9.890	10.145				

16	m-BROMOBENZOIC ACID -1 31.5 3.812			C7H5BrO2 201.02		pHL RE t Z pHS	3.00 5.249 26.63 0.483 3.755	3.25 4.582 24.82 0.555 3.879	3.50 4.082 23.59 0.625 4.003	
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
3.662	3.316	3.222	3.040	2.886	2.775	2.758	2.692	2.650	2.642	2.628
22.67	22.02	19.22	18.95	18.74	18.60	21.36	21.28	21.23	18.71	18.69
0.698	0.774	0.802	0.852	0.898	0.936	0.940	0.964	0.979	0.987	0.993
4.145	4.314	4.385	4.536	4.722	4.936	4.974	5.197	5.434	5.671	5.908
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.618	2.611	2.598	2.595	2.593	2.590	2.611	2.610	2.608	2.606	2.609
18.68	18.68	26.06	26.05	26.05	26.04	18.64	18.64	18.65	18.66	18.68
0.996	0.998	0.999	0.999	1.000	1.000	1.000	1.000	1.000	1.000	1.000
6.153	6.401	6.709	6.957	7.206	7.456	7.650	7.899	8.149	8.399	8.651
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 2 , 1				
2.607	2.605	2.602	2.593	2.589	2.584					
18.72	18.79	18.92	24.17	24.72	25.78					
1.000	1.000	1.000	1.000	1.000	1.000					
8.902	9.153	9.404	9.708	9.964	10.224					

17	o-BROMOBENZOIC ACID -1 31.5 2.854			C7H5BrO2 201.02		pHL RE t Z pHS	3.00 3.183 28.37 0.810 3.449	3.25 2.993 25.76 0.862 3.617	3.50 2.860 24.03 0.904 3.791	
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
2.763	2.695	2.705	2.667	2.641	2.624	2.619	2.609	2.602	2.614	2.611
22.85	22.08	19.40	19.01	18.75	18.60	21.37	21.29	21.24	18.71	18.69
0.936	0.961	0.962	0.976	0.986	0.992	0.992	0.996	0.998	0.999	0.999
3.987	4.205	4.225	4.434	4.663	4.904	4.933	5.177	5.424	5.654	5.901
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.609	2.606	2.595	2.594	2.592	2.590	2.611	2.610	2.608	2.605	2.609
18.68	18.68	26.06	26.05	26.05	26.04	18.64	18.64	18.65	18.66	18.68
1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
6.150	6.399	6.707	6.956	7.206	7.456	7.649	7.899	8.149	8.399	8.651
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 2 , 1				
2.607	2.605	2.602	2.593	2.589	2.584					
18.72	18.79	18.92	24.17	24.72	25.78					
1.000	1.000	1.000	1.000	1.000	1.000					
8.902	9.153	9.404	9.708	9.964	10.224					

18	p-BROMOBENZOIC ACID -1 31.5 3.971			C7H5BrO2 201.02		pHL RE t Z pHS	3.00 5.890 26.38 0.429 3.821	3.25 5.093 24.67 0.497 3.939	3.50 4.493 23.51 0.566 4.057	
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
3.985	3.555	3.406	3.184	2.990	2.843	2.819	2.731	2.673	2.655	2.635
22.63	22.00	19.18	18.93	18.74	18.60	21.35	21.28	21.23	18.71	18.69
0.640	0.720	0.757	0.812	0.866	0.912	0.919	0.949	0.970	0.982	0.990
4.190	4.349	4.431	4.571	4.745	4.950	4.991	5.206	5.439	5.679	5.911
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.622	2.614	2.599	2.596	2.593	2.590	2.611	2.610	2.608	2.606	2.609
18.68	18.68	26.06	26.05	26.05	26.04	18.64	18.64	18.65	18.66	18.68
0.994	0.997	0.998	0.999	0.999	1.000	1.000	1.000	1.000	1.000	1.000
6.154	6.401	6.709	6.957	7.207	7.456	7.650	7.899	8.149	8.399	8.651
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 2 , 1				
2.607	2.605	2.602	2.593	2.589	2.584					
18.72	18.79	18.92	24.17	24.72	25.78					
1.000	1.000	1.000	1.000	1.000	1.000					
8.902	9.153	9.404	9.708	9.964	10.224					

CHROMATOGRAPHIC DATA (1983)

D17

19	2-BROMOBUTYRIC ACID -1			32.3	2.987	C4H7BrO2 167.00		pHL RE t Z pHS	3.00 3.263 27.66 0.768 3.474	3.25 3.033 25.22 0.828 3.635	3.50 2.870 23.61 0.876 3.803
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75	
2.747	2.661	2.667	2.618	2.585	2.563	2.558	2.545	2.537	2.547	2.544	
22.48	21.75	19.12	18.75	18.51	18.36	21.06	20.98	20.93	18.46	18.45	
0.917	0.947	0.950	0.968	0.981	0.989	0.990	0.994	0.997	0.998	0.999	
3.992	4.206	4.231	4.435	4.661	4.900	4.929	5.171	5.418	5.649	5.896	
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50	
2.541	2.538	2.528	2.527	2.525	2.522	2.543	2.541	2.540	2.537	2.541	
18.44	18.44	25.62	25.62	25.61	25.61	18.40	18.40	18.41	18.42	18.44	
0.999	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	
6.144	6.394	6.700	6.949	7.199	7.449	7.644	7.894	8.144	8.394	8.646	
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3					
2.539	2.537	2.534	2.526	2.523	2.518						
18.48	18.54	18.66	23.77	24.28	25.27						
1.000	1.000	1.000	1.000	1.000	1.000						
8.896	9.147	9.399	9.701	9.956	10.215						
20	4-BROMOBUTYRIC ACID -1			34.4	4.585	C4H7BrO2 167.00		pHL RE t Z pHS	3.00 9.113 24.13 0.250 4.086	3.25 7.670 22.87 0.297 4.190	3.50 6.586 22.01 0.347 4.288
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75	
5.643	4.802	4.224	3.839	3.456	3.116	3.025	2.809	2.643	2.529	2.473	
21.34	20.84	18.20	18.04	17.90	17.80	20.28	20.22	20.19	17.88	17.87	
0.407	0.480	0.551	0.607	0.677	0.753	0.776	0.837	0.891	0.938	0.960	
4.396	4.523	4.643	4.743	4.873	5.034	5.090	5.261	5.463	5.728	5.925	
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50	
2.433	2.407	2.383	2.375	2.369	2.365	2.381	2.379	2.377	2.375	2.378	
17.86	17.86	24.59	24.58	24.58	24.57	17.82	17.83	17.83	17.84	17.86	
0.976	0.986	0.993	0.996	0.998	0.999	0.999	1.000	1.000	1.000	1.000	
6.150	6.389	6.695	6.937	7.184	7.433	7.632	7.881	8.131	8.381	8.632	
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3					
2.376	2.374	2.372	2.365	2.362	2.358						
17.89	17.95	18.06	22.80	23.24	24.08						
1.000	1.000	1.000	1.000	1.000	1.000						
8.883	9.133	9.385	9.682	9.936	10.195						
21	o-BROMOPHENYLACETIC ACID -1			31.4	4.054	C8H7BrO2 215.05		pHL RE t Z pHS	3.00 6.292 26.31 0.402 3.857	3.25 5.416 24.65 0.468 3.973	3.50 4.757 23.51 0.535 4.087
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75	
4.195	3.715	3.529	3.284	3.067	2.897	2.868	2.766	2.698	2.673	2.650	
22.65	22.04	19.20	18.96	18.77	18.63	21.39	21.32	21.27	18.74	18.72	
0.609	0.690	0.732	0.788	0.846	0.897	0.906	0.940	0.964	0.979	0.988	
4.217	4.370	4.457	4.591	4.760	4.960	5.002	5.213	5.443	5.684	5.914	
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50	
2.634	2.624	2.609	2.605	2.602	2.599	2.620	2.619	2.617	2.614	2.618	
18.71	18.71	26.11	26.11	26.10	26.09	18.67	18.67	18.68	18.69	18.72	
0.993	0.996	0.998	0.999	0.999	1.000	1.000	1.000	1.000	1.000	1.000	
6.156	6.402	6.711	6.959	7.208	7.457	7.650	7.900	8.150	8.400	8.652	
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 2					
2.616	2.613	2.610	2.602	2.598	2.593						
18.75	18.82	18.95	24.23	24.78	25.85						
1.000	1.000	1.000	1.000	1.000	1.000						
8.903	9.153	9.405	9.709	9.965	10.225						

22	2-BROMOPROPIONIC ACID -1 33.4 2.975				C3H5BrO2 152.98		pHL	3.00	3.25	3.50
							RE	3.149	2.927	2.769
							t	27.05	24.69	23.12
							Z	0.768	0.828	0.877
							pHS	3.463	3.625	3.793
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
2.651	2.568	2.573	2.527	2.495	2.474	2.470	2.457	2.449	2.459	2.455
22.03	21.32	18.79	18.43	18.20	18.06	20.66	20.58	20.53	18.15	18.14
0.917	0.948	0.951	0.969	0.981	0.989	0.990	0.994	0.997	0.998	0.999
3.983	4.197	4.223	4.427	4.654	4.893	4.921	5.163	5.410	5.642	5.888
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.453	2.450	2.441	2.440	2.438	2.436	2.454	2.453	2.451	2.450	2.453
18.13	18.12	25.06	25.06	25.05	25.05	18.09	18.09	18.10	18.11	18.13
0.999	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
6.137	6.387	6.691	6.940	7.190	7.440	7.637	7.887	8.137	8.387	8.639
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3				
2.451	2.449	2.446	2.439	2.436	2.432					
18.16	18.22	18.33	23.25	23.72	24.62					
1.000	1.000	1.000	1.000	1.000	1.000					
8.889	9.140	9.391	9.691	9.945	10.204					
23	5-BROMOVALERIC ACID -1 32.3 4.721				C5H9BrO2 181.03		pHL	3.00	3.25	3.50
							RE	10.913	9.150	7.825
							t	25.06	23.74	22.83
							Z	0.222	0.265	0.311
							pHS	4.157	4.258	4.354
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
6.670	5.635	4.888	4.418	3.947	3.521	3.393	3.120	2.905	2.745	2.673
22.12	21.60	18.77	18.61	18.47	18.36	21.01	20.95	20.92	18.46	18.45
0.366	0.435	0.507	0.562	0.631	0.710	0.737	0.804	0.865	0.923	0.949
4.459	4.582	4.704	4.800	4.923	5.077	5.136	5.299	5.493	5.761	5.949
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.618	2.583	2.550	2.539	2.532	2.526	2.545	2.543	2.540	2.538	2.541
18.44	18.44	25.62	25.62	25.61	25.61	18.40	18.40	18.41	18.42	18.44
0.968	0.981	0.991	0.995	0.997	0.998	0.999	0.999	1.000	1.000	1.000
6.169	6.405	6.716	6.955	7.202	7.450	7.646	7.895	8.144	8.394	8.646
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3				
2.539	2.537	2.534	2.526	2.523	2.518					
18.48	18.54	18.66	23.77	24.28	25.27					
1.000	1.000	1.000	1.000	1.000	1.000					
8.896	9.147	9.399	9.701	9.956	10.215					
24	o-t-BUTYLBenzoic ACID -1 23.2 3.535				C11H14O2 178.23		pHL	3.00	3.25	3.50
							RE	5.717	5.137	4.704
							t	34.09	31.20	29.25
							Z	0.614	0.685	0.751
							pHS	3.709	3.845	3.985
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
4.352	4.076	4.062	3.909	3.791	3.711	3.693	3.647	3.617	3.634	3.623
27.83	26.87	23.10	22.65	22.33	22.13	25.94	25.83	25.76	22.29	22.26
0.814	0.872	0.883	0.919	0.948	0.969	0.971	0.983	0.990	0.994	0.997
4.147	4.336	4.378	4.554	4.761	4.990	5.030	5.265	5.509	5.733	5.976
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
3.615	3.607	3.581	3.578	3.574	3.570	3.615	3.612	3.609	3.604	3.611
22.25	22.24	32.42	32.41	32.39	32.38	22.20	22.21	22.22	22.24	22.28
0.998	0.999	0.999	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
6.224	6.473	6.799	7.048	7.298	7.547	7.723	7.973	8.223	8.473	8.727
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 11 , 1				
3.607	3.602	3.596	3.575	3.565	3.551					
22.36	22.49	22.73	30.36	31.65	34.47					
1.000	1.000	1.000	1.000	1.000	1.000					
8.978	9.229	9.482	9.813	10.075	10.350					

CHROMATOGRAPHIC DATA (1983)

D19

25	p-t-BUTYLBENZOIC -1	ACID 23.2	4,400	C11H14O2 178.23	pHL RE t Z pHS	3.00 10.773 32.41 0.318 4.049	3.25 9.164 30.15 0.376 4.158	3.50 7.940 28.63 0.435 4.264		
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
6.882	5.958	5.554	5.080	4.640	4.278	4.193	3.969	3.813	3.746	3.691
27.50	26.69	22.80	22.51	22.27	22.10	25.88	25.80	25.74	22.28	22.26
0.505	0.586	0.636	0.697	0.766	0.834	0.849	0.899	0.937	0.963	0.977
4.383	4.524	4.613	4.732	4.884	5.067	5.120	5.317	5.537	5.775	5.995
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
3.654	3.630	3.592	3.584	3.577	3.572	3.616	3.613	3.609	3.605	3.611
22.25	22.24	32.42	32.40	32.39	32.38	22.20	22.21	22.22	22.24	22.28
0.987	0.992	0.996	0.998	0.999	0.999	1.000	1.000	1.000	1.000	1.000
6.232	6.477	6.804	7.050	7.298	7.548	7.723	7.973	8.223	8.473	8.727
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 11 , 1				
3.607	3.602	3.596	3.575	3.565	3.551					
22.36	22.49	22.73	30.36	31.65	34.47					
1.000	1.000	1.000	1.000	1.000	1.000					
8.978	9.229	9.482	9.813	10.075	10.350					
26	BUTYRIC ACID (S) -1	33.8	4,820	C4H8O2 88.11	pHL RE t Z pHS	3.00 11.526 24.23 0.200 4.200	3.25 9.640 23.01 0.240 4.299	3.50 8.224 22.18 0.282 4.392		
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
6.987	5.874	4.999	4.507	4.007	3.546	3.394	3.097	2.856	2.659	2.578
21.53	21.03	18.33	18.18	18.06	17.95	20.47	20.42	20.38	18.04	18.03
0.333	0.397	0.471	0.524	0.592	0.671	0.702	0.771	0.838	0.908	0.937
4.495	4.614	4.743	4.833	4.950	5.097	5.159	5.313	5.499	5.775	5.953
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.516	2.474	2.438	2.425	2.417	2.411	2.427	2.424	2.422	2.420	2.422
18.02	18.02	24.87	24.86	24.86	24.85	17.98	17.98	17.99	18.00	18.02
0.960	0.976	0.988	0.993	0.996	0.998	0.999	0.999	1.000	1.000	1.000
6.166	6.399	6.708	6.945	7.190	7.438	7.637	7.885	8.135	8.384	8.636
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 2 , 1				
2.421	2.418	2.416	2.409	2.406	2.402					
18.05	18.11	18.22	23.07	23.52	24.40					
1.000	1.000	1.000	1.000	1.000	1.000					
8.887	9.137	9.388	9.687	9.942	10.200					
27	CACODYLIC ACID (S) -1	29.9	6,184	C2H7AsO2 138.00	pHL RE t Z pHS	3.00 53.330 25.77 0.048 4.877	3.25 43.793 24.52 0.058 4.967	3.50 36.666 23.67 0.070 5.049		
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
30.417	24.720	18.467	16.251	13.887	11.544	10.054	8.598	7.202	4.876	4.509
23.00	22.48	19.41	19.30	19.19	19.10	21.87	21.84	21.81	19.21	19.21
0.084	0.104	0.141	0.160	0.188	0.226	0.261	0.306	0.367	0.552	0.598
5.137	5.236	5.382	5.447	5.530	5.631	5.712	5.807	5.924	6.246	6.327
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
4.097	3.701	3.170	3.042	2.932	2.851	2.832	2.801	2.781	2.767	2.765
19.20	19.20	26.95	26.95	26.96	26.96	19.16	19.17	19.17	19.19	19.21
0.661	0.734	0.859	0.896	0.931	0.957	0.973	0.984	0.991	0.995	0.997
6.442	6.591	6.934	7.085	7.278	7.498	7.704	7.929	8.169	8.414	8.666
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 10				
2.759	2.754	2.750	2.739	2.735	2.728					
19.25	19.33	19.47	25.06	25.68	26.92					
0.998	0.999	0.999	1.000	1.000	1.000					
8.915	9.165	9.417	9.725	9.981	10.243					

28	CAPRIC ACID -1	22.1	5.000	C10H20O2 172.27	pHL RE t Z pHS	3.00 19.575 32.90 0.181 4.330	3.25 16.318 30.75 0.218 4.430	3.50 13.849 29.30 0.258 4.524		
3.75 11.694 28.21 0.307 4.627	4.00 9.766 27.43 0.370 4.747	4.00 8.486 23.30 0.431 4.855	4.25 7.568 23.07 0.485 4.948	4.50 6.655 22.87 0.554 5.067	4.75 5.826 22.73 0.636 5.214	4.75 5.540 26.65 0.669 5.278	5.00 5.004 26.59 0.744 5.433	5.25 4.571 26.56 0.817 5.618	5.50 4.246 22.95 0.891 5.878	5.75 4.095 22.94 0.925 6.056
6.00 3.978 22.93 0.953 6.269	6.25 3.900 22.92 0.972 6.501	6.50 3.817 33.63 0.987 6.833	6.75 3.793 33.62 0.992 7.071	7.00 3.777 33.61 0.996 7.316	7.25 3.766 33.60 0.997 7.564	7.50 3.813 22.89 0.998 7.738	7.75 3.808 22.90 0.999 7.987	8.00 3.803 22.91 0.999 8.236	8.25 3.797 22.93 1.000 8.486	8.50 3.804 22.98 1.000 8.740
8.75 3.800 23.06 1.000 8.991	9.00 3.794 23.21 1.000 9.243	9.25 3.787 23.49 1.000 9.496	9.50 3.762 31.60 1.000 9.832	9.75 3.752 33.11 1.000 10.096	10.00 3.735 36.50 1.000 10.375	(Ref.) : 10 , 12				
29	CAPROIC ACID (S) -1	30.2	4.857	C6H12O2 116.16	pHL RE t Z pHS	3.00 13.156 26.15 0.197 4.228	3.25 10.992 24.73 0.236 4.327	3.50 9.364 23.76 0.278 4.421		
3.75 7.945 23.02 0.328 4.524	4.00 6.670 22.46 0.393 4.644	4.00 5.711 19.43 0.464 4.767	4.25 5.136 19.27 0.517 4.858	4.50 4.555 19.12 0.585 4.976	4.75 4.023 19.01 0.665 5.123	4.75 3.845 21.84 0.696 5.186	5.00 3.503 21.79 0.767 5.341	5.25 3.224 21.75 0.835 5.527	5.50 3.003 19.12 0.905 5.798	5.75 2.909 19.11 0.935 5.977
6.00 2.837 19.10 0.959 6.191	6.25 2.788 19.10 0.976 6.424	6.50 2.743 26.81 0.988 6.738	6.75 2.729 26.80 0.993 6.976	7.00 2.719 26.80 0.996 7.221	7.25 2.712 26.79 0.998 7.469	7.50 2.734 19.06 0.999 7.661	7.75 2.731 19.06 0.999 7.910	8.00 2.728 19.07 1.000 8.159	8.25 2.725 19.09 1.000 8.409	8.50 2.729 19.11 1.000 8.661
8.75 2.726 19.15 1.000 8.912	9.00 2.723 19.23 1.000 9.162	9.25 2.720 19.36 1.000 9.414	9.50 2.710 24.89 1.000 9.721	9.75 2.706 25.49 1.000 9.977	10.00 2.700 26.69 1.000 10.239	(Ref.) : 3 , 1				
30	CAPRYLIC ACID -1	27.4	4.894	C8H16O2 144.21	pHL RE t Z pHS	3.00 14.801 28.02 0.193 4.255	3.25 12.356 26.40 0.232 4.355	3.50 10.515 25.30 0.273 4.449		
3.75 8.909 24.46 0.323 4.552	4.00 7.469 23.83 0.388 4.672	4.00 6.428 20.50 0.455 4.790	4.25 5.767 20.32 0.509 4.883	4.50 5.103 20.16 0.578 5.001	4.75 4.497 20.03 0.658 5.148	4.75 4.293 23.17 0.690 5.211	5.00 3.902 23.11 0.762 5.367	5.25 3.586 23.08 0.831 5.553	5.50 3.340 20.17 0.902 5.820	5.75 3.232 20.16 0.933 6.000
6.00 3.149 20.15 0.958 6.213	6.25 3.093 20.15 0.975 6.446	6.50 3.039 28.69 0.988 6.766	6.75 3.022 28.68 0.993 7.004	7.00 3.011 28.67 0.996 7.250	7.25 3.003 28.67 0.998 7.498	7.50 3.031 20.11 0.999 7.684	7.75 3.028 20.12 0.999 7.932	8.00 3.024 20.12 1.000 8.182	8.25 3.020 20.14 1.000 8.431	8.50 3.025 20.17 1.000 8.684
8.75 3.022 20.22 1.000 8.935	9.00 3.018 20.31 1.000 9.186	9.25 3.014 20.48 1.000 9.438	9.50 3.001 26.68 1.000 9.753	9.75 2.995 27.47 1.000 10.011	10.00 2.988 29.08 1.000 10.276	(Ref.) : 10 , 7				

CHROMATOGRAPHIC DATA (1983)

D21

31	CARBONIC ACID					H2CO3		pHL	3.00	3.25	3.50
	-1	46.1		6.352		62.03		RE	43.836	36.281	30.593
	-2	71.8		10.329				t	19.64	19.01	18.56
								Z	0.038	0.045	0.054
								pHS	4.934	5.020	5.097
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75	
25.532	20.840	14.611	12.986	11.188	9.343	7.956	6.854	5.745	3.518	3.267	
18.18	17.88	15.88	15.82	15.76	15.70	17.46	17.44	17.42	15.72	15.72	
0.065	0.079	0.114	0.128	0.149	0.179	0.211	0.245	0.294	0.487	0.525	
5.180	5.275	5.446	5.504	5.579	5.672	5.759	5.843	5.947	6.299	6.364	
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50	
2.966	2.653	2.146	2.051	1.955	1.875	1.823	1.793	1.770	1.753	1.739	
15.71	15.70	20.61	20.61	20.62	20.63	15.67	15.70	15.76	15.87	16.05	
0.579	0.650	0.808	0.846	0.889	0.928	0.960	0.977	0.991	1.004	1.019	
6.459	6.586	6.937	7.053	7.213	7.412	7.657	7.862	8.092	8.334	8.592	
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 1 , 4					
1.720	1.694	1.655	1.589	1.519	1.443						
16.37	16.90	17.75	22.86	24.69	26.87						
1.039	1.070	1.120	1.210	1.317	1.447						
8.839	9.088	9.338	9.625	9.861	10.097						
32	CHLORIC ACID (S)					HClO3		pHL	3.00	3.25	3.50
	-1	67.0		-2.700		84.46		RE	1.186	1.186	1.186
								t	19.42	17.43	16.28
								Z	1.000	1.000	1.000
								pHS	3.052	3.293	3.534
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75	
1.186	1.185	1.186	1.186	1.186	1.186	1.186	1.186	1.186	1.187	1.186	
15.63	15.25	14.06	13.85	13.73	13.66	14.92	14.88	14.86	13.70	13.69	
1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	
3.778	4.026	4.027	4.272	4.521	4.770	4.775	5.024	5.274	5.520	5.770	
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50	
1.186	1.186	1.186	1.186	1.185	1.185	1.186	1.186	1.186	1.186	1.186	
13.69	13.69	17.05	17.05	17.05	17.05	13.67	13.67	13.67	13.68	13.68	
1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	
6.020	6.270	6.530	6.780	7.030	7.280	7.520	7.770	8.020	8.270	8.520	
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 1 , 4					
1.186	1.186	1.186	1.186	1.185	1.185						
13.69	13.71	13.75	16.03	16.13	16.32						
1.000	1.000	1.000	1.000	1.000	1.000						
8.770	9.020	9.270	9.528	9.779	10.029						
33	2-CHLORO-3-OH-BUTYRIC ACID					C4H7ClO3		pHL	3.00	3.25	3.50
	-1	34.4		2.602		138.55		RE	2.721	2.600	2.518
								t	27.09	24.49	22.81
								Z	0.866	0.907	0.938
								pHS	3.377	3.557	3.743
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75	
2.460	2.421	2.431	2.409	2.394	2.384	2.381	2.375	2.370	2.381	2.379	
21.68	20.97	18.53	18.16	17.93	17.79	20.31	20.24	20.19	17.88	17.87	
0.960	0.976	0.977	0.986	0.992	0.995	0.996	0.997	0.999	0.999	1.000	
3.949	4.175	4.189	4.406	4.640	4.884	4.909	5.154	5.402	5.634	5.882	
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50	
2.377	2.375	2.367	2.365	2.364	2.362	2.379	2.378	2.376	2.375	2.378	
17.86	17.86	24.59	24.58	24.58	24.57	17.82	17.83	17.83	17.84	17.86	
1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	
6.131	6.381	6.682	6.932	7.182	7.432	7.630	7.881	8.131	8.381	8.632	
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3					
2.376	2.374	2.372	2.365	2.362	2.358						
17.89	17.95	18.06	22.80	23.24	24.08						
1.000	1.000	1.000	1.000	1.000	1.000						
8.883	9.133	9.385	9.682	9.936	10.195						

34	CHLOROACETIC ACID -1 43.7			2.865	C2H3ClO2 94.50			pHL RE t Z pHS	3.00 2.356 22.92 0.778 3.373	3.25 2.189 21.04 0.838 3.542	3.50 2.073 19.81 0.886 3.717
	3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
	1.989	1.931	1.932	1.900	1.878	1.864	1.863	1.855	1.849	1.853	1.851
	18.96	18.42	16.51	16.23	16.06	15.95	17.91	17.85	17.82	16.02	16.01
	0.924	0.953	0.956	0.972	0.983	0.990	0.991	0.995	0.997	0.998	0.999
	3.914	4.133	4.159	4.369	4.599	4.841	4.860	5.104	5.351	5.589	5.836
	6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
	1.850	1.848	1.845	1.844	1.843	1.842	1.850	1.850	1.849	1.848	1.850
	16.00	16.00	21.24	21.24	21.24	21.24	15.97	15.97	15.98	15.99	15.99
	0.999	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
	6.085	6.335	6.621	6.871	7.121	7.371	7.585	7.835	8.085	8.335	8.586
	8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 1				
	1.849	1.848	1.846	1.844	1.842	1.840					
	16.02	16.05	16.12	19.75	20.00	20.46					
	1.000	1.000	1.000	1.000	1.000	1.000					
	8.836	9.086	9.337	9.619	9.871	10.125					
35	m-CHLOROBENZOIC ACID -1 32.0			3.830	C7H5ClO2 156.57			pHL RE t Z pHS	3.00 5.244 26.31 0.475 3.760	3.25 4.571 24.55 0.547 3.883	3.50 4.066 23.34 0.617 4.007
	3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
	3.642	3.290	3.191	3.006	2.851	2.737	2.720	2.652	2.609	2.600	2.585
	22.45	21.81	19.06	18.79	18.59	18.45	21.16	21.09	21.04	18.55	18.54
	0.691	0.767	0.796	0.847	0.895	0.933	0.938	0.962	0.978	0.987	0.992
	4.147	4.314	4.388	4.537	4.722	4.935	4.972	5.194	5.431	5.668	5.905
	6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
	2.575	2.568	2.555	2.553	2.550	2.548	2.568	2.567	2.565	2.563	2.566
	18.53	18.52	25.78	25.78	25.77	25.77	18.49	18.49	18.50	18.51	18.53
	0.996	0.998	0.999	0.999	1.000	1.000	1.000	1.000	1.000	1.000	1.000
	6.149	6.397	6.704	6.953	7.202	7.452	7.646	7.896	8.146	8.396	8.648
	8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 2 , 1				
	2.564	2.562	2.559	2.551	2.547	2.542					
	18.57	18.64	18.76	23.92	24.44	25.46					
	1.000	1.000	1.000	1.000	1.000	1.000					
	8.898	9.149	9.401	9.704	9.959	10.219					
36	o-CHLOROBENZOIC ACID -1 33.8			2.943	C7H5ClO2 156.57			pHL RE t Z pHS	3.00 3.077 26.89 0.777 3.452	3.25 2.866 24.53 0.836 3.615	3.50 2.717 22.96 0.883 3.785
	3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
	2.606	2.528	2.534	2.491	2.461	2.441	2.437	2.426	2.418	2.428	2.425
	21.88	21.17	18.67	18.32	18.09	17.95	20.52	20.44	20.39	18.04	18.03
	0.922	0.951	0.954	0.971	0.982	0.990	0.990	0.994	0.997	0.998	0.999
	3.977	4.193	4.217	4.423	4.650	4.890	4.917	5.160	5.407	5.639	5.886
	6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
	2.422	2.419	2.411	2.409	2.408	2.406	2.424	2.422	2.421	2.419	2.422
	18.02	18.02	24.87	24.87	24.86	24.86	17.98	17.98	17.99	18.00	18.02
	0.999	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
	6.135	6.384	6.688	6.937	7.187	7.437	7.634	7.884	8.134	8.384	8.636
	8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 2 , 1				
	2.420	2.418	2.416	2.409	2.406	2.402					
	18.05	18.11	18.22	23.07	23.52	24.40					
	1.000	1.000	1.000	1.000	1.000	1.000					
	8.886	9.137	9.388	9.687	9.942	10.200					

37	p-CHLOROBENZOIC ACID -1 33.4 3.977			C7H5ClO2 156.57			pHL RE t Z pHS	3.00 5.625 25.32 0.423 3.815	3.25 4.857 23.74 0.491 3.934	3.50 4.280 22.66 0.559 4.050
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
3.791	3.375	3.223	3.011	2.824	2.682	2.659	2.574	2.517	2.498	2.478
21.84	21.26	18.60	18.37	18.18	18.06	20.64	20.57	20.53	18.15	18.14
0.633	0.713	0.752	0.807	0.862	0.909	0.916	0.947	0.968	0.982	0.989
4.183	4.340	4.426	4.563	4.736	4.940	4.980	5.194	5.426	5.667	5.899
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.466	2.457	2.445	2.442	2.439	2.437	2.455	2.453	2.452	2.450	2.453
18.13	18.12	25.06	25.06	25.05	25.05	18.09	18.09	18.10	18.11	18.13
0.994	0.996	0.998	0.999	0.999	1.000	1.000	1.000	1.000	1.000	1.000
6.142	6.389	6.693	6.941	7.190	7.440	7.637	7.887	8.137	8.387	8.639
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 2 , 1				
2.451	2.449	2.446	2.439	2.436	2.432					
18.16	18.22	18.33	23.25	23.72	24.62					
1.000	1.000	1.000	1.000	1.000	1.000					
8.889	9.140	9.391	9.691	9.945	10.204					
38	2-CHLOROBUTYRIC ACID -1 34.4 2.839			C4H7ClO2 122.55			pHL RE t Z pHS	3.00 2.916 26.74 0.806 3.423	3.25 2.738 24.33 0.860 3.592	3.50 2.613 22.75 0.902 3.767
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
2.522	2.459	2.467	2.431	2.407	2.392	2.388	2.379	2.373	2.382	2.380
21.66	20.96	18.52	18.16	17.93	17.79	20.31	20.24	20.19	17.88	17.87
0.936	0.960	0.962	0.976	0.986	0.992	0.992	0.996	0.998	0.999	0.999
3.964	4.183	4.205	4.414	4.644	4.886	4.912	5.155	5.403	5.635	5.882
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.377	2.375	2.367	2.365	2.364	2.362	2.379	2.378	2.376	2.375	2.378
17.86	17.86	24.59	24.58	24.58	24.57	17.82	17.83	17.83	17.84	17.86
1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
6.131	6.381	6.682	6.932	7.182	7.432	7.631	7.881	8.131	8.381	8.632
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3				
2.376	2.374	2.372	2.365	2.362	2.358					
17.89	17.95	18.06	22.80	23.24	24.08					
1.000	1.000	1.000	1.000	1.000	1.000					
8.883	9.133	9.385	9.682	9.936	10.195					
39	3-CHLOROBUTYRIC ACID -1 34.4 4.056			C4H7ClO2 122.55			pHL RE t Z pHS	3.00 5.824 24.70 0.395 3.846	3.25 5.006 23.23 0.461 3.961	3.50 4.391 22.22 0.527 4.075
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
3.867	3.418	3.230	3.005	2.802	2.643	2.617	2.521	2.456	2.430	2.408
21.45	20.89	18.30	18.09	17.92	17.79	20.30	20.23	20.19	17.88	17.87
0.601	0.682	0.727	0.783	0.842	0.894	0.902	0.937	0.962	0.978	0.987
4.203	4.355	4.447	4.579	4.745	4.943	4.984	5.193	5.422	5.667	5.895
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.394	2.384	2.371	2.368	2.365	2.363	2.380	2.378	2.377	2.375	2.378
17.86	17.86	24.59	24.58	24.58	24.57	17.82	17.83	17.83	17.84	17.86
0.992	0.996	0.998	0.999	0.999	1.000	1.000	1.000	1.000	1.000	1.000
6.137	6.383	6.686	6.934	7.182	7.432	7.631	7.881	8.131	8.381	8.632
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3				
2.376	2.374	2.372	2.365	2.362	2.358					
17.89	17.95	18.06	22.80	23.24	24.08					
1.000	1.000	1.000	1.000	1.000	1.000					
8.883	9.133	9.385	9.682	9.936	10.195					

40	2-CHLOROCROTONIC -1	35.4	ACID 3.155		C4H5ClO2 120.54		pHL RE t Z pHS	3.00 3.225 25.70 0.704 3.499	3.25 2.947 23.63 0.773 3.652	3.50 2.746 22.24 0.831 3.811
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
2.592	2.479	2.476	2.414	2.369	2.340	2.336	2.319	2.308	2.315	2.311
21.27	20.61	18.21	17.89	17.68	17.55	19.99	19.92	19.88	17.63	17.62
0.882	0.923	0.929	0.953	0.971	0.983	0.984	0.991	0.995	0.997	0.998
3.991	4.197	4.232	4.427	4.648	4.885	4.911	5.152	5.398	5.631	5.877
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.308	2.305	2.297	2.296	2.294	2.293	2.308	2.307	2.306	2.304	2.307
17.61	17.61	24.14	24.14	24.13	24.13	17.57	17.58	17.58	17.59	17.61
0.999	0.999	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
6.126	6.375	6.675	6.925	7.174	7.424	7.625	7.875	8.125	8.375	8.626
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3				
2.305	2.304	2.301	2.296	2.293	2.289					
17.64	17.70	17.79	22.39	22.80	23.57					
1.000	1.000	1.000	1.000	1.000	1.000					
8.877	9.127	9.379	9.674	9.928	10.186					
41	3-CHLOROCROTONIC -1	35.4	ACID 3.848		C4H5ClO2 120.54		pHL RE t Z pHS	3.00 4.865 24.52 0.461 3.753	3.25 4.227 22.98 0.532 3.875	3.50 3.750 21.93 0.602 3.997
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
3.348	3.013	2.905	2.732	2.585	2.475	2.459	2.394	2.352	2.340	2.326
21.14	20.57	18.08	17.85	17.67	17.55	19.98	19.92	19.87	17.63	17.62
0.676	0.754	0.787	0.838	0.887	0.928	0.933	0.958	0.975	0.986	0.992
4.135	4.300	4.380	4.525	4.707	4.917	4.953	5.173	5.408	5.649	5.884
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.316	2.310	2.300	2.297	2.295	2.293	2.309	2.307	2.306	2.304	2.307
17.61	17.61	24.14	24.14	24.13	24.13	17.57	17.58	17.58	17.59	17.61
0.995	0.997	0.999	0.999	1.000	1.000	1.000	1.000	1.000	1.000	1.000
6.129	6.376	6.677	6.925	7.175	7.424	7.625	7.875	8.125	8.375	8.626
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3				
2.305	2.304	2.301	2.296	2.293	2.289					
17.64	17.70	17.79	22.39	22.80	23.57					
1.000	1.000	1.000	1.000	1.000	1.000					
8.877	9.127	9.379	9.674	9.928	10.186					
42	CHLORODIBROMOACETIC -1	36.5	ACID 1.000		C2HBr2ClO2 252.29		pHL RE t Z pHS	3.00 2.235 26.84 0.995 3.264	3.25 2.231 23.91 0.997 3.474	3.50 2.228 22.11 0.998 3.686
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
2.226	2.225	2.234	2.233	2.232	2.230	2.228	2.227	2.225	2.235	2.234
20.97	20.28	18.02	17.64	17.42	17.29	19.66	19.59	19.55	17.37	17.36
0.999	0.999	0.999	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
3.911	4.148	4.152	4.382	4.623	4.869	4.892	5.139	5.388	5.620	5.869
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.233	2.231	2.224	2.224	2.222	2.221	2.235	2.234	2.233	2.231	2.234
17.35	17.35	23.68	23.67	23.67	23.66	17.32	17.32	17.33	17.34	17.35
1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
6.119	6.368	6.666	6.916	7.166	7.416	7.619	7.869	8.119	8.369	8.620
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3 , 12				
2.233	2.231	2.229	2.223	2.221	2.218					
17.38	17.43	17.52	21.96	22.34	23.06					
1.000	1.000	1.000	1.000	1.000	1.000					
8.870	9.121	9.372	9.665	9.919	10.176					

CHROMATOGRAPHIC DATA (1983)

D25

43	CHLORODIFLUOROACETIC -1	38.6	1.000	C2HC1F2O2 130.48	pHL RE t Z pHS	3.00 2.109 25.94 0.995 3.244	3.25 2.105 23.13 0.997 3.456	3.50 2.102 21.40 0.998 3.671		
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
2.100	2.099	2.107	2.106	2.105	2.103	2.102	2.100	2.099	2.107	2.106
20.32	19.67	17.53	17.18	16.97	16.85	19.08	19.01	18.97	16.92	16.91
0.999	0.999	0.999	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
3.897	4.135	4.139	4.370	4.611	4.858	4.879	5.127	5.376	5.609	5.858
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.105	2.104	2.098	2.098	2.097	2.095	2.107	2.107	2.106	2.104	2.106
16.91	16.90	22.87	22.86	22.86	22.86	16.87	16.87	16.88	16.89	16.90
1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
6.108	6.358	6.652	6.902	7.152	7.401	7.608	7.858	8.108	8.358	8.609
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3 , 12				
2.105	2.104	2.102	2.098	2.095	2.093					
16.93	16.97	17.06	21.22	21.56	22.18					
1.000	1.000	1.000	1.000	1.000	1.000					
8.859	9.110	9.361	9.650	9.903	10.159					
44	2-CHLOROISOCROTONIC -1	35.4	2.810	C4H5C1O2 120.54	pHL RE t Z pHS	3.00 2.812 26.28 0.811 3.409	3.25 2.643 23.92 0.865 3.580	3.50 2.526 22.36 0.906 3.757		
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
2.441	2.382	2.390	2.357	2.334	2.320	2.317	2.308	2.302	2.311	2.309
21.30	20.62	18.25	17.90	17.68	17.55	19.99	19.92	19.88	17.63	17.62
0.938	0.962	0.964	0.977	0.987	0.992	0.993	0.996	0.998	0.999	0.999
3.955	4.175	4.196	4.407	4.638	4.879	4.904	5.149	5.396	5.629	5.876
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.307	2.304	2.297	2.296	2.294	2.292	2.308	2.307	2.306	2.304	2.307
17.61	17.61	24.14	24.14	24.13	24.13	17.57	17.58	17.58	17.59	17.61
1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
6.125	6.375	6.675	6.924	7.174	7.424	7.625	7.875	8.125	8.375	8.626
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3				
2.305	2.304	2.301	2.296	2.293	2.289					
17.64	17.70	17.79	22.39	22.80	23.57					
1.000	1.000	1.000	1.000	1.000	1.000					
8.877	9.127	9.379	9.674	9.928	10.186					
45	m-CHLOROPHENOL -1	33.4	9.023	C6H5C1O 128.56	pHL RE t Z pHS	3.00 23.86 0.002 6.286	3.25 22.87 0.002 6.373	3.50 22.17 0.003 6.451		
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
21.62	21.18	18.44	18.35	18.27	18.20	20.61	20.58	20.56	18.30	18.30
0.003	0.004	0.006	0.007	0.008	0.010	0.012	0.014	0.017	0.037	0.041
6.534	6.626	6.786	6.843	6.914	7.001	7.089	7.164	7.255	7.602	7.644
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
48.789	41.096	18.793	17.288	15.352	13.145	7.802	7.185	6.417	5.579	4.249
18.29	18.29	24.92	24.93	24.94	24.95	18.27	18.27	18.28	18.29	18.33
0.047	0.056	0.123	0.134	0.151	0.176	0.302	0.328	0.369	0.425	0.564
7.704	7.782	8.155	8.197	8.257	8.337	8.637	8.689	8.765	8.866	9.106
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3 , 1				
3.850	3.456	3.119	2.800	2.659	2.562					
18.37	18.43	18.54	23.51	23.98	24.91					
0.625	0.698	0.776	0.866	0.912	0.947					
9.213	9.354	9.527	9.798	10.005	10.237					

46	o-CHLOROPHENOL -1			33.4	8.477	C6H5C10 128.56		pHL RE t Z pHS	3.00 RE t Z pHS	3.25 RE t Z pHS	3.50 RE t Z pHS
	3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
	21.54	21.11	18.37	18.28	18.20	18.13	20.53	20.51	20.48	18.22	18.22
	0.006	0.008	0.011	0.012	0.015	0.018	0.022	0.026	0.032	0.068	0.075
	6.260	6.353	6.513	6.569	6.641	6.729	6.816	6.892	6.984	7.330	7.374
	6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
	26.879	22.697	11.020	10.152	9.048	7.807	5.171	4.789	4.330	3.852	3.255
	18.21	18.21	24.87	24.88	24.89	24.90	18.17	18.17	18.17	18.18	18.20
	0.085	0.101	0.211	0.230	0.258	0.300	0.461	0.499	0.553	0.624	0.744
	7.435	7.516	7.887	7.933	7.999	8.088	8.382	8.447	8.540	8.666	8.907
	8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3 , 1				
	3.026	2.826	2.678	2.559	2.506	2.471					
	18.23	18.29	18.40	23.33	23.80	24.70					
	0.803	0.861	0.910	0.951	0.971	0.983					
	9.051	9.232	9.443	9.729	9.964	10.214					
47	p-CHLOROPHENOL -1			33.4	9.378	C6H5C10 128.60		pHL RE t Z pHS	3.00 RE t Z pHS	3.25 RE t Z pHS	3.50 RE t Z pHS
	3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
	21.76	21.32	18.56	18.48	18.40	18.33	20.74	20.72	20.70	18.43	18.43
	0.002	0.003	0.004	0.004	0.005	0.006	0.008	0.009	0.011	0.025	0.028
	6.712	6.805	6.965	7.022	7.093	7.179	7.267	7.342	7.432	7.781	7.822
	6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
	72.299	60.855	27.121	24.943	22.127	18.897	10.628	9.768	8.685	7.481	5.343
	18.43	18.42	25.07	25.07	25.08	25.10	18.42	18.43	18.44	18.45	18.52
	0.031	0.037	0.085	0.092	0.104	0.122	0.220	0.240	0.270	0.315	0.445
	7.881	7.959	8.332	8.372	8.430	8.507	8.809	8.857	8.925	9.017	9.256
	8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3 , 1				
	4.788	4.211	3.687	3.120	2.884	2.705					
	18.56	18.63	18.75	23.80	24.30	25.26					
	0.498	0.569	0.652	0.773	0.838	0.894					
	9.347	9.468	9.618	9.878	10.059	10.271					
48	p-CHLOROPHENYLACETIC ACID -1			34.5	4.190	C8H7C102 170.60		pHL RE t Z pHS	3.00 RE t Z pHS	3.25 RE t Z pHS	3.50 RE t Z pHS
	3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
	4.202	3.673	3.414	3.155	2.914	2.718	2.682	2.562	2.479	2.439	2.411
	21.38	20.84	18.24	18.05	17.89	17.77	20.26	20.20	20.16	17.86	17.84
	0.550	0.631	0.684	0.742	0.805	0.865	0.876	0.919	0.950	0.971	0.982
	4.247	4.392	4.493	4.615	4.771	4.960	5.005	5.205	5.429	5.677	5.900
	6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
	2.392	2.380	2.366	2.362	2.359	2.356	2.372	2.371	2.370	2.367	2.370
	17.84	17.83	24.54	24.54	24.53	24.53	17.80	17.80	17.81	17.82	17.83
	0.990	0.994	0.997	0.998	0.999	0.999	1.000	1.000	1.000	1.000	1.000
	6.138	6.384	6.687	6.933	7.182	7.431	7.630	7.880	8.130	8.380	8.632
	8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 2				
	2.369	2.367	2.364	2.358	2.355	2.351					
	17.87	17.93	18.03	22.76	23.20	24.02					
	1.000	1.000	1.000	1.000	1.000	1.000					
	8.882	9.133	9.384	9.681	9.936	10.194					

CHROMATOGRAPHIC DATA (1983)

D27

49	2-CHLOROPROPIONIC -1	ACID 36.9	2.796	C3H5ClO2 108.52	pHL RE t Z pHS	3.00 2.692 25.60 0.812 3.396	3.25 2.530 23.32 0.865 3.567	3.50 2.418 21.82 0.906 3.746		
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
2.337	2.281	2.287	2.256	2.234	2.221	2.218	2.210	2.204	2.212	2.210
20.80	20.15	17.88	17.54	17.33	17.20	19.54	19.47	19.43	17.28	17.27
0.939	0.962	0.964	0.978	0.987	0.992	0.993	0.996	0.998	0.999	0.999
3.945	4.165	4.186	4.398	4.629	4.871	4.895	5.139	5.387	5.620	5.868
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.208	2.206	2.200	2.199	2.197	2.195	2.210	2.209	2.208	2.206	2.208
17.26	17.26	23.52	23.51	23.51	23.50	17.23	17.23	17.24	17.25	17.26
1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
6.117	6.366	6.664	6.913	7.163	7.413	7.616	7.866	8.116	8.366	8.618
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3				
2.207	2.205	2.203	2.198	2.196	2.193					
17.29	17.34	17.43	21.82	22.18	22.88					
1.000	1.000	1.000	1.000	1.000	1.000					
8.868	9.119	9.370	9.662	9.916	10.173					
50	3-CHLOROPROPIONIC -1	ACID 36.9	3.867	C3H5ClO2 108.52	pHL RE t Z pHS	3.00 4.756 23.82 0.452 3.756	3.25 4.125 22.38 0.522 3.877	3.50 3.653 21.39 0.592 3.997		
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
3.255	2.922	2.806	2.636	2.489	2.379	2.364	2.299	2.256	2.242	2.227
20.64	20.10	17.71	17.49	17.32	17.20	19.53	19.47	19.43	17.28	17.27
0.666	0.744	0.780	0.831	0.882	0.924	0.929	0.956	0.974	0.985	0.991
4.134	4.297	4.380	4.524	4.703	4.911	4.947	5.165	5.400	5.643	5.877
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.218	2.212	2.202	2.200	2.198	2.196	2.210	2.209	2.208	2.206	2.208
17.27	17.26	23.51	23.51	23.51	23.50	17.23	17.23	17.24	17.25	17.26
0.995	0.997	0.999	0.999	1.000	1.000	1.000	1.000	1.000	1.000	1.000
6.121	6.368	6.666	6.914	7.164	7.413	7.617	7.866	8.116	8.367	8.618
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3				
2.207	2.205	2.203	2.198	2.196	2.193					
17.29	17.34	17.43	21.82	22.18	22.88					
1.000	1.000	1.000	1.000	1.000	1.000					
8.868	9.119	9.370	9.662	9.916	10.173					
51	CHLOROUS -1	ACID 53.9	1.960	HC102 68.46	pHL RE t Z pHS	3.00 1.566 21.27 0.946 3.166	3.25 1.534 19.19 0.967 3.380	3.50 1.514 17.93 0.980 3.598		
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
1.501	1.493	1.496	1.492	1.489	1.487	1.487	1.485	1.484	1.487	1.487
17.15	16.69	15.18	14.93	14.78	14.69	16.27	16.22	16.20	14.74	14.74
0.988	0.993	0.993	0.996	0.998	0.999	0.999	0.999	1.000	1.000	1.000
3.828	4.068	4.075	4.310	4.554	4.801	4.812	5.061	5.310	5.552	5.801
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
1.486	1.485	1.484	1.484	1.483	1.483	1.487	1.486	1.486	1.486	1.486
14.73	14.73	18.94	18.94	18.94	18.94	14.71	14.71	14.71	14.72	14.72
1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
6.051	6.300	6.573	6.823	7.073	7.323	7.550	7.800	8.050	8.300	8.551
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 2 , 4				
1.486	1.485	1.485	1.484	1.483	1.482					
14.74	14.77	14.81	17.70	17.86	18.15					
1.000	1.000	1.000	1.000	1.000	1.000					
8.801	9.051	9.302	9.571	9.822	10.074					

52	5-CHLOROVALERIC ACID				C5H9ClO2					
	-1	32.3	4.697		136.58		pHL	3.00	3.25	3.50
							RE	10.670	8.952	7.661
							t	25.08	23.75	22.84
							Z	0.227	0.271	0.318
							pHS	4.145	4.247	4.343
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
6.536	5.529	4.812	4.354	3.894	3.481	3.359	3.095	2.887	2.736	2.666
22.13	21.60	18.78	18.62	18.47	18.36	21.01	20.95	20.92	18.46	18.45
0.374	0.444	0.515	0.571	0.640	0.718	0.745	0.811	0.871	0.926	0.951
4.449	4.573	4.694	4.791	4.915	5.071	5.130	5.294	5.490	5.756	5.947
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.615	2.581	2.549	2.539	2.532	2.526	2.545	2.543	2.540	2.538	2.541
18.44	18.44	25.62	25.62	25.61	25.61	18.40	18.40	18.41	18.42	18.44
0.970	0.982	0.991	0.995	0.997	0.998	0.999	0.999	1.000	1.000	1.000
6.168	6.405	6.715	6.955	7.202	7.450	7.646	7.895	8.144	8.394	8.646
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3				
2.539	2.537	2.534	2.526	2.523	2.518					
18.48	18.54	18.66	23.77	24.28	25.27					
1.000	1.000	1.000	1.000	1.000	1.000					
8.896	9.147	9.399	9.701	9.956	10.215					
53	CHROMIC ACID				H2CrO4					
	-1	59.3	0.745		118.01		pHL	3.00	3.25	3.50
	-2	81.1	6.490				RE	1.350	1.347	1.346
							t	20.55	18.43	17.20
							Z	0.997	0.999	1.000
							pHS	3.095	3.328	3.562
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
1.344	1.343	1.345	1.343	1.340	1.336	1.335	1.327	1.315	1.294	1.266
16.49	16.10	14.74	14.55	14.52	14.62	16.08	16.36	16.88	16.45	17.62
1.002	1.004	1.004	1.008	1.015	1.026	1.027	1.047	1.080	1.146	1.230
3.802	4.047	4.050	4.293	4.540	4.789	4.800	5.049	5.299	5.587	5.825
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
1.230	1.190	1.140	1.114	1.092	1.076	1.074	1.068	1.065	1.062	1.062
19.10	20.75	28.02	29.31	30.35	31.08	25.71	25.95	26.10	26.19	26.24
1.342	1.475	1.647	1.748	1.835	1.898	1.940	1.965	1.980	1.988	1.993
6.062	6.299	6.601	6.809	7.037	7.275	7.527	7.768	8.015	8.263	8.513
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 10 , 2				
1.061	1.060	1.060	1.055	1.055	1.054					
26.29	26.33	26.40	30.41	30.58	30.88					
1.996	1.998	1.999	1.999	2.000	2.000					
8.763	9.013	9.262	9.515	9.765	10.015					
54	CINNAMIC ACID				C9H8O2					
	-1	28.3	4.438		148.16		pHL	3.00	3.25	3.50
							RE	9.443	7.998	6.906
							t	27.82	26.11	24.95
							Z	0.296	0.350	0.407
							pHS	4.040	4.147	4.250
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
5.962	5.130	4.685	4.278	3.890	3.562	3.484	3.279	3.131	3.052	3.001
24.07	23.43	20.23	20.01	19.83	19.69	22.75	22.68	22.64	19.82	19.80
0.473	0.552	0.611	0.671	0.740	0.811	0.828	0.882	0.924	0.956	0.973
4.366	4.501	4.604	4.716	4.860	5.036	5.088	5.276	5.491	5.739	5.952
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.966	2.943	2.916	2.909	2.903	2.899	2.926	2.924	2.921	2.918	2.923
19.79	19.79	28.04	28.04	28.03	28.02	19.75	19.75	19.76	19.78	19.80
0.984	0.991	0.995	0.997	0.999	0.999	0.999	1.000	1.000	1.000	1.000
6.186	6.429	6.745	6.990	7.238	7.487	7.674	7.924	8.174	8.424	8.676
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 11 , 1				
2.920	2.917	2.913	2.901	2.896	2.889					
19.85	19.94	20.09	26.06	26.78	28.24					
1.000	1.000	1.000	1.000	1.000	1.000					
8.927	9.178	9.430	9.742	10.000	10.263					

55	cis-CINNAMIC -1	ACID 28.3	3.879		C9H8O2 148.16		pHL RE t Z pHS	3.00 6.035 28.66 0.469 3.799	3.25 5.258 26.63 0.540 3.921	3.50 4.673 25.26 0.609 4.043	
	3.75 4.181 24.23 0.683 4.183	4.00 3.771 23.51 0.760 4.349	4.00 3.662 20.38 0.789 4.419	4.25 3.444 20.08 0.841 4.568	4.50 3.261 19.85 0.890 4.751	4.75 3.125 19.70 0.930 4.963	4.75 3.103 22.78 0.935 5.004	5.00 3.023 22.70 0.960 5.226	5.25 2.972 22.65 0.977 5.462	5.50 2.963 19.82 0.986 5.696	5.75 2.945 19.80 0.992 5.932
	6.00 2.933 19.79 0.995 6.177	6.25 2.925 19.79 0.997 6.425	6.50 2.907 28.05 0.999 6.739	6.75 2.904 28.04 0.999 6.988	7.00 2.900 28.03 1.000 7.237	7.25 2.897 28.02 1.000 7.487	7.50 2.925 19.75 1.000 7.674	7.75 2.923 19.75 1.000 7.924	8.00 2.921 19.76 1.000 8.174	8.25 2.918 19.78 1.000 8.424	8.50 2.923 19.80 1.000 8.676
	8.75 2.920 19.85 1.000 8.927	9.00 2.917 19.94 1.000 9.178	9.25 2.913 20.09 1.000 9.430	9.50 2.901 26.06 1.000 9.742	9.75 2.896 26.78 1.000 10.000	10.00 2.889 28.24 1.000 10.263	(Ref.) : 11 , 1				
56	CITRACONIC -1 -2	ACID 28.0 56.0	2.458 6.146		C5H6O4 130.10		pHL RE t Z pHS	3.00 3.215 31.42 0.910 3.419	3.25 3.113 28.21 0.941 3.602	3.50 3.041 26.16 0.964 3.790	
	3.75 2.985 24.82 0.983 3.997	4.00 2.938 24.05 1.000 4.224	4.00 2.954 21.03 1.001 4.240	4.25 2.910 20.76 1.017 4.458	4.50 2.852 20.80 1.038 4.692	4.75 2.770 21.16 1.071 4.934	4.75 2.750 24.40 1.077 4.974	5.00 2.631 25.14 1.130 5.212	5.25 2.474 26.20 1.208 5.447	5.50 2.234 25.14 1.362 5.772	5.75 2.087 26.17 1.467 5.956
	6.00 1.946 27.17 1.584 6.156	6.25 1.825 28.03 1.699 6.370	6.50 1.700 37.63 1.826 6.680	6.75 1.651 37.99 1.884 6.883	7.00 1.615 38.24 1.928 7.110	7.25 1.591 38.39 1.957 7.349	7.50 1.599 29.67 1.975 7.577	7.75 1.591 29.73 1.985 7.819	8.00 1.585 29.77 1.992 8.066	8.25 1.581 29.80 1.995 8.315	8.50 1.582 29.83 1.997 8.566
	8.75 1.580 29.86 1.998 8.815	9.00 1.577 29.92 1.999 9.065	9.25 1.575 30.01 2.000 9.315	9.50 1.566 36.07 2.000 9.587	9.75 1.563 36.40 2.000 9.838	10.00 1.560 37.03 2.000 10.091	(Ref.) : 12 , 2				
57	CITRIC -1 -2 -3	ACID (S) 28.7 54.7 74.4	3.128 4.761 6.396		C6H8O7 192.13		pHL RE t Z pHS	3.00 3.601 31.06 0.796 3.529	3.25 3.244 28.91 0.889 3.683	3.50 2.958 27.81 0.984 3.842	
	3.75 2.698 27.49 1.090 4.021	4.00 2.454 27.90 1.215 4.220	4.00 2.396 25.15 1.260 4.286	4.25 2.212 26.00 1.384 4.465	4.50 2.037 27.08 1.527 4.664	4.75 1.883 28.28 1.679 4.878	4.75 1.859 31.80 1.701 4.912	5.00 1.744 33.01 1.843 5.126	5.25 1.649 34.33 1.984 5.354	5.50 1.573 32.98 2.156 5.632	5.75 1.511 34.50 2.296 5.849
	6.00 1.452 36.15 2.450 6.078	6.25 1.400 37.73 2.601 6.314	6.50 1.341 49.56 2.749 6.599	6.75 1.314 50.49 2.837 6.825	7.00 1.294 51.13 2.900 7.063	7.25 1.281 51.52 2.941 7.307	7.50 1.302 41.22 2.965 7.546	7.75 1.297 41.35 2.980 7.793	8.00 1.293 41.43 2.989 8.041	8.25 1.290 41.48 2.994 8.291	8.50 1.293 41.53 2.996 8.541
	8.75 1.291 41.57 2.998 8.791	9.00 1.289 41.63 2.999 9.041	9.25 1.286 41.73 2.999 9.291	9.50 1.273 48.97 3.000 9.552	9.75 1.270 49.29 3.000 9.802	10.00 1.266 49.89 3.000 10.052	(Ref.) : 10 , 1				

58	CRESOL -1	33.4	7.377	C7H8O 108.14			pHL RE t Z pHS	3.00 RE t Z pHS	3.25 RE t Z pHS	3.50 RE t Z pHS
3.75 0.021 5.712	4.00 21.52 0.027 5.806	4.00 60.064 0.038 5.964	4.25 52.789 0.043 6.022	4.50 44.885 0.051 6.096	4.75 36.899 0.062 6.186	4.75 30.724 0.075 6.272	5.00 25.974 0.088 6.351	5.25 21.256 0.108 6.447	5.50 10.974 0.213 6.790	5.75 10.027 0.187 6.841
6.00 8.860 18.17 0.265 6.912	6.25 7.590 18.16 0.310 7.007	6.50 4.643 24.94 0.513 7.372	6.75 4.319 24.95 0.553 7.441	7.00 3.931 24.96 0.609 7.540	7.25 3.533 24.98 0.679 7.672	7.50 3.047 18.10 0.798 7.938	7.75 2.897 18.11 0.841 8.063	8.00 2.752 18.11 0.886 8.232	8.25 2.637 18.12 0.926 8.436	8.50 2.558 18.14 0.957 8.685
8.75 2.514 18.17 0.974 8.910	9.00 2.485 18.23 0.985 9.149	9.25 2.467 18.34 0.991 9.396	9.50 2.450 23.25 0.996 9.694	9.75 2.442 23.72 0.998 9.947	10.00 2.435 24.63 0.999 10.205	(Ref.) : 3				

59	m-CRESOL -1	33.4	7.770	C7H8O 108.14			pHL RE t Z pHS	3.00 RE t Z pHS	3.25 RE t Z pHS	3.50 RE t Z pHS
3.75 0.014 5.908	4.00 21.52 0.017 6.001	4.00 93.326 18.34 0.024 6.160	4.25 81.981 18.26 0.028 6.217	4.50 69.643 18.17 0.033 6.290	4.75 57.161 18.10 0.040 6.378	4.75 47.317 20.52 0.048 6.465	5.00 39.934 20.49 0.057 6.543	5.25 32.584 20.47 0.070 6.637	5.50 16.066 18.19 0.144 6.981	5.75 14.652 18.18 0.158 7.028
6.00 12.898 18.18 0.180 7.095	6.25 10.969 18.18 0.213 7.182	6.50 6.067 24.90 0.389 7.551	6.75 5.616 24.91 0.421 7.608	7.00 5.058 24.93 0.469 7.690	7.25 4.457 24.94 0.534 7.802	7.50 3.512 18.12 0.688 8.081	7.75 3.298 18.12 0.734 8.178	8.00 3.068 18.12 0.791 8.314	8.25 2.861 18.13 0.850 8.488	8.50 2.686 18.15 0.910 8.735
8.75 2.597 18.18 0.941 8.936	9.00 2.535 18.24 0.964 9.163	9.25 2.496 18.35 0.979 9.403	9.50 2.464 23.26 0.989 9.699	9.75 2.450 23.73 0.994 9.949	10.00 2.439 24.64 0.997 10.206	(Ref.) : 3				

60	p-CRESOL -1	33.4	7.959	C7H8O 108.14			pHL RE t Z pHS	3.00 RE t Z pHS	3.25 RE t Z pHS	3.50 RE t Z pHS
3.75 0.011 6.002	4.00 21.52 0.014 6.095	4.00 18.34 0.020 6.254	4.25 18.26 0.022 6.311	4.50 86.177 18.18 0.026 6.384	4.75 70.693 18.10 0.032 6.472	4.75 58.401 20.52 0.039 6.559	5.00 49.261 20.49 0.046 6.636	5.25 40.153 20.47 0.057 6.729	5.50 19.470 18.19 0.119 7.074	5.75 17.746 18.19 0.130 7.120
6.00 15.601 18.19 0.149 7.185	6.25 13.234 18.18 0.175 7.270	6.50 7.023 24.89 0.335 7.639	6.75 6.490 24.90 0.363 7.692	7.00 5.824 24.91 0.406 7.769	7.25 5.095 24.93 0.465 7.873	7.50 3.830 18.13 0.629 8.157	7.75 3.580 18.13 0.675 8.243	8.00 3.299 18.13 0.734 8.365	8.25 3.034 18.14 0.800 8.525	8.50 2.785 18.16 0.876 8.770
8.75 2.667 18.19 0.916 8.957	9.00 2.579 18.25 0.947 9.174	9.25 2.522 18.35 0.969 9.409	9.50 2.478 23.27 0.984 9.703	9.75 2.458 23.74 0.991 9.951	10.00 2.444 24.65 0.995 10.207	(Ref.) : 3				

61	CROTONIC ACID -1	33.5	4.705	C4H6O2 86.09	pHL RE t Z pHS	3.00 10.417 24.46 0.224 4.146	3.25 8.738 23.20 0.268 4.247	3.50 7.476 22.33 0.314 4.343		
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
6.377	5.392	4.671	4.226	3.779	3.374	3.254	2.995	2.790	2.638	2.569
21.66	21.15	18.43	18.27	18.14	18.03	20.58	20.52	20.49	18.12	18.11
0.369	0.438	0.511	0.566	0.635	0.713	0.740	0.806	0.867	0.924	0.949
4.448	4.571	4.695	4.790	4.913	5.067	5.127	5.289	5.484	5.753	5.941
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.518	2.485	2.454	2.444	2.437	2.432	2.449	2.447	2.445	2.442	2.445
18.10	18.10	25.01	25.01	25.00	25.00	18.06	18.06	18.07	18.08	18.10
0.969	0.982	0.991	0.995	0.997	0.998	0.999	0.999	1.000	1.000	1.000
6.161	6.398	6.705	6.946	7.192	7.440	7.638	7.887	8.136	8.386	8.638
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 11 , 7				
2.443	2.441	2.438	2.432	2.428	2.424					
18.13	18.20	18.30	23.20	23.67	24.56					
1.000	1.000	1.000	1.000	1.000	1.000					
8.888	9.139	9.390	9.690	9.944	10.203					
62	CYANIC ACID -1	67.0	3.470	HCNO 43.03	pHL RE t Z pHS	3.00 2.136 17.37 0.548 3.519	3.25 1.878 16.54 0.625 3.655	3.50 1.690 15.98 0.695 3.791		
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
1.538	1.416	1.379	1.318	1.269	1.235	1.233	1.214	1.202	1.195	1.192
15.55	15.24	13.96	13.83	13.73	13.67	14.92	14.88	14.86	13.70	13.69
0.766	0.833	0.857	0.898	0.933	0.959	0.960	0.976	0.986	0.992	0.996
3.946	4.127	4.207	4.372	4.571	4.795	4.812	5.041	5.282	5.536	5.776
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
1.189	1.188	1.187	1.186	1.186	1.185	1.187	1.186	1.186	1.186	1.186
13.69	13.69	17.05	17.05	17.05	17.05	13.67	13.67	13.67	13.68	13.68
0.997	0.999	0.999	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
6.022	6.271	6.532	6.780	7.030	7.280	7.520	7.770	8.020	8.270	8.520
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 2 , 4				
1.186	1.186	1.186	1.186	1.185	1.185					
13.69	13.71	13.75	16.03	16.13	16.32					
1.000	1.000	1.000	1.000	1.000	1.000					
8.770	9.020	9.270	9.528	9.779	10.029					
63	CYANOACETIC ACID -1	45.0	2.469	C3H3NO2 85.06	pHL RE t Z pHS	3.00 2.035 23.06 0.876 3.281	3.25 1.947 20.93 0.916 3.470	3.50 1.889 19.58 0.945 3.667		
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
1.850	1.824	1.828	1.814	1.804	1.798	1.797	1.794	1.791	1.796	1.794
18.70	18.16	16.33	16.04	15.86	15.76	17.66	17.60	17.57	15.82	15.81
0.966	0.980	0.980	0.988	0.993	0.996	0.996	0.998	0.999	0.999	1.000
3.881	4.112	4.126	4.349	4.587	4.832	4.849	5.096	5.344	5.582	5.830
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
1.794	1.792	1.789	1.789	1.788	1.787	1.794	1.794	1.793	1.792	1.794
15.81	15.81	20.89	20.89	20.88	20.88	15.78	15.78	15.78	15.79	15.80
1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
6.080	6.330	6.614	6.864	7.114	7.364	7.580	7.830	8.080	8.330	8.580
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 1				
1.793	1.792	1.791	1.788	1.787	1.785					
15.82	15.86	15.92	19.43	19.67	20.10					
1.000	1.000	1.000	1.000	1.000	1.000					
8.831	9.081	9.332	9.611	9.864	10.117					

64	m-CYANO BENZOIC ACID				C8H5NO2		pHL	3.00			3.25	3.50		
	-1	31.3	3.598		147.13			RE	4.581	4.058		3.668		
							t	27.13	25.14	23.80		23.80		
							Z	0.559	0.633	0.703		0.703		
							pHS	3.673	3.805	3.940		3.940		
	3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75			
	3.349	3.094	3.051	2.913	2.805	2.729	2.718	2.674	2.647	2.648	2.638			
	22.81	22.12	19.34	19.03	18.81	18.67	21.44	21.36	21.31	18.77	18.75			
	0.772	0.838	0.855	0.897	0.932	0.959	0.961	0.977	0.987	0.992	0.996			
	4.094	4.276	4.333	4.500	4.700	4.924	4.959	5.190	5.431	5.665	5.906			
	6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50			
	2.631	2.626	2.614	2.612	2.609	2.607	2.629	2.627	2.626	2.623	2.627			
	18.74	18.74	26.17	26.16	26.16	26.15	18.70	18.71	18.71	18.73	18.75			
	0.997	0.999	0.999	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000			
	6.153	6.401	6.709	6.959	7.208	7.458	7.651	7.901	8.151	8.401	8.653			
	8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3 , 7							
	2.625	2.622	2.619	2.610	2.607	2.601								
	18.79	18.86	18.98	24.28	24.83	25.91								
	1.000	1.000	1.000	1.000	1.000	1.000								
	8.903	9.154	9.406	9.710	9.966	10.226								
65	CYCLOBUTANE-1,1-DICARBOXYLIC ACID				C6H8O4		pHL	3.00			3.25	3.50		
	-1	25.5	3.127		144.13			RE	4.258	3.935		3.696		
	-2	51.1	5.879				t	32.64	29.71	27.80		27.80		
							Z	0.753	0.817	0.872		0.872		
							pHS	3.567	3.721	3.881		3.881		
	3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75			
	3.502	3.343	3.343	3.223	3.090	2.932	2.894	2.708	2.503	2.253	2.119			
	26.52	25.81	22.46	22.34	22.59	23.23	26.90	27.93	29.16	27.68	28.54			
	0.922	0.968	0.976	1.015	1.061	1.122	1.136	1.220	1.329	1.506	1.611			
	4.064	4.272	4.308	4.505	4.726	4.959	5.005	5.230	5.454	5.760	5.941			
	6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50			
	1.999	1.905	1.815	1.780	1.756	1.739	1.758	1.753	1.748	1.745	1.747			
	29.30	29.90	40.12	40.34	40.48	40.54	30.88	30.91	30.93	30.95	30.98			
	1.717	1.810	1.895	1.934	1.961	1.977	1.986	1.992	1.996	1.998	1.999			
	6.145	6.366	6.667	6.886	7.122	7.366	7.587	7.832	8.081	8.330	8.581			
	8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 2 , 1							
	1.745	1.742	1.740	1.728	1.724	1.720								
	31.01	31.08	31.19	37.90	38.31	39.08								
	1.999	2.000	2.000	2.000	2.000	2.000								
	8.831	9.081	9.331	9.609	9.860	10.113								
66	CYCLOHEXANE-1,1-DICARBOXYLIC ACID				C8H12O4		pHL	3.00			3.25	3.50		
	-1	24.0	3.451		172.18			RE	3.966	3.398		2.971		
	-2	48.0	4.110				t	37.68	35.67	34.61		34.61		
							Z	0.877	1.032	1.191		1.191		
							pHS	3.607	3.737	3.871		3.871		
	3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75			
	2.623	2.348	2.309	2.160	2.046	1.970	1.958	1.914	1.886	1.891	1.881			
	34.13	34.08	30.53	30.70	30.91	31.10	35.02	35.14	35.22	31.78	31.80			
	1.362	1.535	1.581	1.701	1.804	1.880	1.886	1.932	1.961	1.977	1.987			
	4.025	4.206	4.260	4.432	4.636	4.863	4.887	5.120	5.362	5.603	5.846			
	6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50			
	1.874	1.869	1.852	1.849	1.846	1.842	1.873	1.871	1.869	1.866	1.871			
	31.81	31.82	42.27	42.26	42.23	42.20	31.80	31.80	31.81	31.81	31.84			
	1.993	1.996	1.998	1.999	1.999	2.000	2.000	2.000	2.000	2.000	2.000			
	6.093	6.342	6.628	6.877	7.127	7.376	7.591	7.841	8.091	8.341	8.592			
	8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 2 , 1							
	1.868	1.866	1.863	1.849	1.845	1.840								
	31.88	31.95	32.07	39.28	39.75	40.64								
	2.000	2.000	2.000	2.000	2.000	2.000								
	8.842	9.092	9.343	9.624	9.876	10.130								

CHROMATOGRAPHIC DATA (1983)

D33

67	CYCLOPENTANE-1,1-DICARBOXYLIC ACID C7H10O4						pHL			
	-1	25.0	3.230			158.15	RE	3.00	3.25	3.50
	-2	50.0	4.080				t	36.94	34.89	33.84
							Z	0.953	1.104	1.256
							pHS	3.540	3.682	3.828
	3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50
	2.421	2.196	2.170	2.041	1.943	1.877	1.867	1.827	1.802	1.807
	33.40	33.40	30.00	30.18	30.39	30.58	34.32	34.44	34.52	31.22
	1.416	1.575	1.612	1.723	1.818	1.888	1.893	1.936	1.963	1.979
	3.994	4.185	4.234	4.415	4.624	4.853	4.876	5.110	5.353	5.595
	6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25
	1.792	1.787	1.772	1.769	1.766	1.763	1.790	1.789	1.787	1.785
	31.25	31.26	41.26	41.24	41.22	41.19	31.24	31.24	31.24	31.25
	1.993	1.996	1.998	1.999	1.999	2.000	2.000	2.000	2.000	2.000
	6.085	6.334	6.618	6.867	7.117	7.366	7.584	7.834	8.084	8.333
	8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 2 , 1			
	1.786	1.784	1.781	1.769	1.765	1.761				
	31.31	31.37	31.49	38.37	38.80	39.61				
	2.000	2.000	2.000	2.000	2.000	2.000				
	8.834	9.085	9.335	9.614	9.866	10.119				
68	DECYLSULFONIC ACID C10H22O3S						pHL			
	-1	26.9	1.000			222.30	RE	3.00	3.25	3.50
							t	32.83	29.17	26.85
							Z	0.996	0.998	0.998
							pHS	3.377	3.577	3.778
	3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50
	3.065	3.063	3.084	3.083	3.080	3.077	3.071	3.068	3.065	3.086
	25.33	24.39	21.26	20.77	20.46	20.28	23.54	23.44	23.38	20.40
	0.999	0.999	0.999	1.000	1.000	1.000	1.000	1.000	1.000	1.000
	3.995	4.227	4.232	4.454	4.691	4.936	4.968	5.214	5.463	5.688
	6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25
	3.082	3.079	3.062	3.061	3.058	3.055	3.087	3.085	3.082	3.079
	20.37	20.36	29.07	29.06	29.06	29.05	20.32	20.33	20.34	20.35
	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
	6.185	6.435	6.752	7.002	7.252	7.502	7.686	7.936	8.186	8.436
	8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 2 , 12			
	3.081	3.078	3.073	3.060	3.054	3.045				
	20.44	20.53	20.71	27.05	27.88	29.58				
	1.000	1.000	1.000	1.000	1.000	1.000				
	8.939	9.190	9.443	9.759	10.018	10.284				
69	DIBROMOFLUOROACETIC ACID C2HBr2FO2						pHL			
	-1	38.6	1.000			235.84	RE	3.00	3.25	3.50
							t	25.94	23.13	21.40
							Z	0.995	0.997	0.998
							pHS	3.244	3.456	3.671
	3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50
	2.100	2.099	2.107	2.106	2.105	2.103	2.102	2.100	2.099	2.107
	20.32	19.67	17.53	17.18	16.97	16.85	19.08	19.01	18.97	16.92
	0.999	0.999	0.999	1.000	1.000	1.000	1.000	1.000	1.000	1.000
	3.897	4.135	4.139	4.370	4.611	4.858	4.879	5.127	5.376	5.609
	6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25
	2.105	2.104	2.098	2.098	2.097	2.095	2.107	2.107	2.106	2.104
	16.91	16.90	22.87	22.86	22.86	22.86	16.87	16.87	16.88	16.89
	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
	6.108	6.358	6.652	6.902	7.152	7.401	7.608	7.858	8.108	8.358
	8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3 , 12			
	2.105	2.104	2.102	2.098	2.095	2.093				
	16.93	16.97	17.06	21.22	21.56	22.18				
	1.000	1.000	1.000	1.000	1.000	1.000				
	8.859	9.110	9.361	9.650	9.903	10.159				

70	2,3-DIBROMOPROPIONIC ACID -1 32.3				C3H4Br202 231.87		pHL RE t Z pHS	3.00 2.688 28.66 0.938 3.347	3.25 2.630 25.68 0.960 3.540	3.50 2.593 23.78 0.974 3.737	
	3.75 2.567 22.53 0.984 3.952	4.00 2.549 21.76 0.991 4.183	4.00 2.563 19.17 0.991 4.191	4.25 2.553 18.77 0.994 4.413	4.50 2.546 18.51 0.997 4.651	4.75 2.540 18.37 0.998 4.895	4.75 2.537 21.06 0.998 4.922	5.00 2.533 20.98 0.999 5.168	5.25 2.530 20.93 0.999 5.417	5.50 2.543 18.46 1.000 5.646	5.75 2.542 18.45 1.000 5.895
	6.00 2.540 18.44 1.000 6.144	6.25 2.537 18.44 1.000 6.394	6.50 2.528 25.62 1.000 6.700	6.75 2.526 25.62 1.000 6.949	7.00 2.525 25.61 1.000 7.199	7.25 2.522 25.61 1.000 7.449	7.50 2.543 18.40 1.000 7.644	7.75 2.541 18.40 1.000 7.894	8.00 2.540 18.41 1.000 8.144	8.25 2.537 18.42 1.000 8.394	8.50 2.541 18.44 1.000 8.646
	8.75 2.539 18.48 1.000 8.896	9.00 2.537 18.54 1.000 9.147	9.25 2.534 18.66 1.000 9.399	9.50 2.526 23.77 1.000 9.701	9.75 2.523 24.28 1.000 9.956	10.00 2.518 25.27 1.000 10.215	(Ref.) : 3				
71	DICHLOROACETIC ACID (S) -1 39.7				C2H2Cl202 128.94		pHL RE t Z pHS	3.00 2.057 25.48 0.991 3.237	3.25 2.050 22.74 0.994 3.449	3.50 2.045 21.06 0.996 3.664	
	3.75 2.042 20.01 0.998 3.891	4.00 2.039 19.37 0.999 4.129	4.00 2.047 17.30 0.999 4.133	4.25 2.046 16.96 0.999 4.364	4.50 2.044 16.75 1.000 4.606	4.75 2.043 16.63 1.000 4.853	4.75 2.041 18.80 1.000 4.873	5.00 2.040 18.74 1.000 5.121	5.25 2.038 18.70 1.000 5.370	5.50 2.046 16.71 1.000 5.604	5.75 2.045 16.70 1.000 5.853
	6.00 2.044 16.69 1.000 6.102	6.25 2.043 16.69 1.000 6.352	6.50 2.038 22.48 1.000 6.645	6.75 2.037 22.48 1.000 6.895	7.00 2.036 22.47 1.000 7.145	7.25 2.035 22.47 1.000 7.394	7.50 2.046 16.65 1.000 7.602	7.75 2.046 16.66 1.000 7.852	8.00 2.044 16.66 1.000 8.102	8.25 2.043 16.67 1.000 8.352	8.50 2.045 16.68 1.000 8.604
	8.75 2.044 16.71 1.000 8.854	9.00 2.043 16.75 1.000 9.104	9.25 2.041 16.83 1.000 9.355	9.50 2.037 20.87 1.000 9.643	9.75 2.035 21.18 1.000 9.896	10.00 2.033 21.76 1.000 10.151	(Ref.) : 2 , 7				
72	DICHLOROFLUOROACETIC ACID -1 36.5				C2HCl2F02 146.93		pHL RE t Z pHS	3.00 2.235 26.84 0.995 3.264	3.25 2.231 23.91 0.997 3.474	3.50 2.228 22.11 0.998 3.686	
	3.75 2.226 20.97 0.999 3.911	4.00 2.225 20.28 0.999 4.148	4.00 2.234 18.02 0.999 4.152	4.25 2.233 17.64 1.000 4.382	4.50 2.232 17.42 1.000 4.623	4.75 2.230 17.29 1.000 4.869	4.75 2.228 19.66 1.000 4.892	5.00 2.227 19.59 1.000 5.139	5.25 2.225 19.55 1.000 5.388	5.50 2.235 17.37 1.000 5.620	5.75 2.234 17.36 1.000 5.869
	6.00 2.233 17.35 1.000 6.119	6.25 2.231 17.35 1.000 6.368	6.50 2.224 23.68 1.000 6.666	6.75 2.224 23.67 1.000 6.916	7.00 2.222 23.67 1.000 7.166	7.25 2.221 23.66 1.000 7.416	7.50 2.235 17.32 1.000 7.619	7.75 2.234 17.32 1.000 7.869	8.00 2.233 17.33 1.000 8.119	8.25 2.231 17.34 1.000 8.369	8.50 2.234 17.35 1.000 8.620
	8.75 2.233 17.38 1.000 8.870	9.00 2.231 17.43 1.000 9.121	9.25 2.229 17.52 1.000 9.372	9.50 2.223 21.96 1.000 9.665	9.75 2.221 22.34 1.000 9.919	10.00 2.218 23.06 1.000 10.176	(Ref.) : 3 , 12				

73	2,4-DICHLOROPHENOL -1	31.3	7.699	C6H4C120	163.00	pHL RE t Z pHS	3.00 24.84 0.008 5.628	3.25 23.72 0.010 5.715	3.50 22.95 0.012 5.794
	3.75 4.00 4.00 4.25 4.50 4.75 4.75 5.00 5.25 5.50 5.75	91.443 80.177 67.993 55.733 46.287 38.990 31.774 15.933 14.509	18.94 18.84 18.75 18.67 21.26 21.23 21.21 18.78 18.77 18.77 18.77	0.015 0.019 0.027 0.030 0.036 0.044 0.053 0.063 0.077 0.156 0.171	5.878 5.972 6.127 6.186 6.259 6.349 6.435 6.513 6.609 6.949 6.997				
	6.00 6.25 6.50 6.75 7.00 7.25 7.50 7.75 8.00 8.25 8.50	12.756 10.843 6.132 5.670 5.106 4.509 3.641 3.423 3.196 2.997 2.839	18.77 18.77 25.98 25.99 26.00 26.03 18.72 18.72 18.73 18.74 18.76	0.195 0.230 0.413 0.447 0.498 0.566 0.712 0.758 0.814 0.870 0.922	7.066 7.155 7.521 7.582 7.669 7.786 8.058 8.162 8.306 8.488 8.735				
	8.75 9.00 9.25 9.50 9.75 10.00	2.756 2.699 2.663 2.632 2.619 2.608	18.80 18.87 18.99 24.29 24.85 25.93	0.950 0.970 0.983 0.991 0.995 0.997	8.943 9.173 9.415 9.717 9.969 10.228	(Ref.) : 3			
74	2,4-DICHLOROPHENOXYACETIC -1	25.0	1.000	ACID C8H6C1203	221.00	pHL RE t Z pHS	3.00 3.320 34.59 0.996 3.405	3.25 3.316 30.71 0.998 3.602	3.50 3.313 28.24 0.999 3.801
	3.75 4.00 4.00 4.25 4.50 4.75 4.75 5.00 5.25 5.50 5.75	3.311 3.309 3.333 3.332 3.329 3.326 3.319 3.315 3.311 3.337 3.335	26.62 25.60 22.22 21.69 21.36 21.16 24.69 24.58 24.51 21.30 21.27	0.999 0.999 0.999 1.000 1.000 1.000 1.000 1.000 1.000 1.000 1.000	4.016 4.248 4.252 4.473 4.709 4.954 4.988 5.234 5.483 5.706 5.954				
	6.00 6.25 6.50 6.75 7.00 7.25 7.50 7.75 8.00 8.25 8.50	3.332 3.328 3.308 3.306 3.303 3.299 3.337 3.335 3.332 3.328 3.334	21.26 21.25 30.66 30.65 30.64 30.63 21.21 21.22 21.23 21.25 21.28	1.000 1.000 1.000 1.000 1.000 1.000 1.000 1.000 1.000 1.000 1.000	6.203 6.453 6.775 7.024 7.274 7.524 7.704 7.954 8.204 8.454 8.707				
	8.75 9.00 9.25 9.50 9.75 10.00	3.331 3.326 3.322 3.305 3.297 3.286	21.35 21.46 21.66 28.60 29.63 31.79	1.000 1.000 1.000 1.000 1.000 1.000	8.958 9.209 9.462 9.785 10.045 10.315	(Ref.) : 2 , 12			
75	DICHROMIC ACID -1 -2	58.9 82.1	0.745 6.723	H2Cr207	218.00	pHL RE t Z pHS	3.00 1.359 20.62 0.996 3.097	3.25 1.357 18.48 0.998 3.330	3.50 1.355 17.24 1.000 3.564
	3.75 4.00 4.00 4.25 4.50 4.75 4.75 5.00 5.25 5.50 5.75	1.354 1.353 1.355 1.354 1.352 1.349 1.348 1.343 1.334 1.320 1.298	16.51 16.11 14.74 14.53 14.46 14.49 15.94 16.10 16.41 15.65 16.49	1.001 1.002 1.002 1.005 1.009 1.015 1.016 1.028 1.048 1.089 1.146	3.803 4.048 4.051 4.293 4.540 4.789 4.799 5.048 5.299 5.576 5.821				
	6.00 6.25 6.50 6.75 7.00 7.25 7.50 7.75 8.00 8.25 8.50	1.266 1.227 1.166 1.133 1.103 1.079 1.070 1.017 1.054 1.051 1.049	17.69 19.20 26.43 27.92 29.25 30.28 25.29 14.52 25.89 26.03 26.11	1.232 1.347 1.528 1.643 1.750 1.838 1.904 1.006 1.966 1.980 1.989	6.064 6.305 6.624 6.826 7.046 7.280 7.536 4.350 8.015 8.263 8.513				
	8.75 9.00 9.25 9.50 9.75 10.00	1.048 1.047 1.046 1.042 1.041 1.040	26.17 26.23 26.29 30.26 30.43 30.72	1.994 1.996 1.998 1.999 1.999 2.000	8.762 9.011 9.261 9.513 9.763 10.013	(Ref.) : 10			

76	DIETHYLACETIC ACID -1 30.3 4.734				C6H12O2 160.17		pHL RE t Z pHS	3.00 11.668 26.19 0.221 4.169	3.25 9.780 24.74 0.265 4.271	3.50 8.360 23.75 0.311 4.367
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
7.122	6.016	5.240	4.731	4.223	3.766	3.628	3.335	3.104	2.938	2.859
22.99	22.43	19.42	19.25	19.09	18.98	21.81	21.75	21.71	19.09	19.08
0.366	0.435	0.505	0.561	0.630	0.710	0.737	0.804	0.865	0.922	0.949
4.472	4.596	4.715	4.811	4.935	5.090	5.149	5.313	5.508	5.773	5.962
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.801	2.763	2.726	2.715	2.707	2.701	2.723	2.721	2.718	2.715	2.719
19.07	19.06	26.75	26.74	26.74	26.73	19.03	19.03	19.04	19.05	19.07
0.968	0.981	0.991	0.995	0.997	0.998	0.999	0.999	1.000	1.000	1.000
6.183	6.419	6.733	6.973	7.220	7.468	7.660	7.909	8.158	8.408	8.660
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3 , 1				
2.717	2.714	2.710	2.701	2.697	2.691					
19.12	19.19	19.32	24.83	25.43	26.62					
1.000	1.000	1.000	1.000	1.000	1.000					
8.911	9.162	9.413	9.720	9.976	10.238					
77	DIETHYLMALONIC ACID -1 25.0 2.211 -2 49.5 7.290				C7H12O4 116.16		pHL RE t Z pHS	3.00 3.483 34.27 0.948 3.435	3.25 3.421 30.58 0.966 3.623	3.50 3.380 28.20 0.978 3.814
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
3.351	3.330	3.354	3.341	3.328	3.314	3.305	3.285	3.256	3.239	3.169
26.62	25.62	22.23	21.72	21.42	21.27	24.81	24.79	24.89	21.95	22.37
0.987	0.993	0.993	0.997	1.000	1.004	1.005	1.010	1.019	1.034	1.059
4.024	4.251	4.260	4.477	4.712	4.955	4.991	5.236	5.483	5.734	5.978
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
3.062	2.912	2.576	2.413	2.250	2.105	2.010	1.941	1.888	1.852	1.835
23.04	24.00	35.80	36.93	38.06	39.06	30.02	30.48	30.83	31.06	31.23
1.100	1.162	1.325	1.425	1.539	1.657	1.774	1.846	1.902	1.941	1.966
6.221	6.461	6.855	7.037	7.232	7.440	7.685	7.885	8.109	8.347	8.597
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 2 , 7				
1.822	1.812	1.806	1.791	1.786	1.781					
31.34	31.45	31.59	38.57	39.02	39.85					
1.980	1.989	1.994	1.997	1.998	1.999					
8.841	9.089	9.338	9.617	9.869	10.122					
78	DIFLUOROACETIC ACID -1 40.6 1.000				C2H2F2O2 96.03		pHL RE t Z pHS	3.00 2.002 25.17 0.995 3.226	3.25 1.997 22.45 0.997 3.440	3.50 1.995 20.80 0.998 3.657
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
1.993	1.991	1.998	1.998	1.996	1.995	1.994	1.992	1.991	1.998	1.998
19.76	19.14	17.12	16.78	16.58	16.47	18.59	18.52	18.49	16.54	16.53
0.999	0.999	0.999	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
3.885	4.123	4.127	4.359	4.602	4.848	4.868	5.116	5.365	5.600	5.849
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
1.997	1.996	1.991	1.990	1.989	1.988	1.999	1.998	1.997	1.996	1.998
16.52	16.52	22.18	22.18	22.17	22.17	16.49	16.49	16.50	16.51	16.52
1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
6.098	6.348	6.639	6.889	7.139	7.389	7.598	7.848	8.098	8.348	8.599
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 2 , 12				
1.997	1.995	1.994	1.990	1.988	1.986					
16.54	16.58	16.66	20.60	20.89	21.44					
1.000	1.000	1.000	1.000	1.000	1.000					
8.850	9.100	9.351	9.637	9.890	10.145					

CHROMATOGRAPHIC DATA (1983)

D37

79	2,4-DIHYDROXYBENZOIC ACID -1	32.0	3.395	C7H6O4 154.12			pHL RE t Z pHS	3.00 3.987 27.09 0.630 3.597	3.25 3.585 24.99 0.703 3.738	3.50 3.288 23.57 0.768 3.883
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
3.052	2.870	2.853	2.754	2.679	2.629	2.621	2.592	2.574	2.580	2.573
22.55	21.85	19.15	18.83	18.60	18.45	21.17	21.09	21.04	18.55	18.54
0.830	0.884	0.895	0.928	0.955	0.973	0.975	0.985	0.991	0.995	0.997
4.049	4.243	4.289	4.470	4.681	4.911	4.943	5.179	5.423	5.656	5.900
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.569	2.565	2.553	2.552	2.550	2.547	2.568	2.567	2.565	2.563	2.566
18.53	18.52	25.78	25.78	25.77	25.77	18.49	18.49	18.50	18.51	18.53
0.998	0.999	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
6.147	6.396	6.703	6.952	7.202	7.452	7.646	7.896	8.146	8.396	8.648
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 10				
2.564	2.562	2.559	2.551	2.547	2.542					
18.57	18.64	18.76	23.92	24.44	25.46					
1.000	1.000	1.000	1.000	1.000	1.000					
8.898	9.149	9.401	9.704	9.959	10.219					
80	2,5-DIHYDROXYBENZOIC ACID -1	34.4	2.975	C7H6O4 154.12			pHL RE t Z pHS	3.00 3.065 26.52 0.766 3.456	3.25 2.845 24.22 0.826 3.618	3.50 2.690 22.70 0.875 3.786
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
2.574	2.492	2.496	2.451	2.419	2.399	2.395	2.383	2.375	2.383	2.380
21.65	20.96	18.50	18.16	17.93	17.79	20.31	20.24	20.19	17.88	17.87
0.916	0.947	0.950	0.968	0.981	0.989	0.990	0.994	0.997	0.998	0.999
3.976	4.190	4.217	4.421	4.648	4.887	4.914	5.157	5.403	5.636	5.882
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.378	2.375	2.367	2.366	2.364	2.362	2.379	2.378	2.376	2.375	2.378
17.86	17.86	24.59	24.58	24.58	24.57	17.82	17.83	17.83	17.84	17.86
0.999	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
6.131	6.381	6.682	6.932	7.182	7.432	7.631	7.881	8.131	8.381	8.632
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3				
2.376	2.374	2.372	2.365	2.362	2.358					
17.89	17.95	18.06	22.80	23.24	24.08					
1.000	1.000	1.000	1.000	1.000	1.000					
8.883	9.133	9.385	9.682	9.936	10.195					
81	2,6-DIHYDROXYBENZOIC ACID -1	37.2	1.301	C7H6O4 154.12			pHL RE t Z pHS	3.00 2.202 26.50 0.990 3.261	3.25 2.194 23.63 0.994 3.470	3.50 2.189 21.86 0.996 3.682
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
2.185	2.182	2.192	2.190	2.188	2.186	2.185	2.183	2.181	2.191	2.190
20.74	20.07	17.85	17.48	17.26	17.14	19.46	19.39	19.35	17.22	17.21
0.998	0.999	0.999	0.999	1.000	1.000	1.000	1.000	1.000	1.000	1.000
3.907	4.144	4.148	4.378	4.619	4.865	4.887	5.135	5.384	5.616	5.865
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.188	2.187	2.181	2.180	2.179	2.177	2.191	2.190	2.189	2.187	2.190
17.20	17.20	23.40	23.39	23.39	23.39	17.16	17.17	17.17	17.18	17.19
1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
6.115	6.365	6.661	6.911	7.161	7.411	7.615	7.865	8.115	8.365	8.616
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3				
2.188	2.187	2.185	2.180	2.177	2.174					
17.22	17.27	17.36	21.71	22.07	22.75					
1.000	1.000	1.000	1.000	1.000	1.000					
8.867	9.117	9.368	9.660	9.914	10.170					

82	3,4-DIHYDROXYBENZOIC -1	ACID 34.4	4.509	C7H6O4 154.12			pHL RE t Z pHS	3.00 8.509 24.19 0.268 4.051	3.25 7.179 22.92 0.318 4.156	3.50 6.180 22.04 0.371 4.255
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
5.312	4.540	4.039	3.681	3.329	3.021	2.944	2.749	2.603	2.508	2.459
21.36	20.85	18.21	18.05	17.91	17.80	20.28	20.23	20.19	17.88	17.87
0.433	0.508	0.577	0.634	0.703	0.778	0.798	0.856	0.906	0.946	0.966
4.366	4.496	4.613	4.717	4.851	5.017	5.071	5.248	5.455	5.716	5.919
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.424	2.402	2.380	2.373	2.368	2.364	2.381	2.379	2.377	2.375	2.378
17.86	17.86	24.59	24.58	24.58	24.57	17.82	17.83	17.83	17.84	17.86
0.979	0.988	0.994	0.997	0.998	0.999	0.999	1.000	1.000	1.000	1.000
6.147	6.388	6.693	6.936	7.184	7.432	7.632	7.881	8.131	8.381	8.632
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3				
2.376	2.374	2.372	2.365	2.362	2.358					
17.89	17.95	18.06	22.80	23.24	24.08					
1.000	1.000	1.000	1.000	1.000	1.000					
8.883	9.133	9.385	9.682	9.936	10.195					
83	3,5-DIHYDROXYBENZOIC -1	ACID 34.4	4.051	C7H6O4 154.12			pHL RE t Z pHS	3.00 5.802 24.71 0.397 3.843	3.25 4.988 23.23 0.463 3.959	3.50 4.376 22.22 0.529 4.073
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
3.855	3.409	3.224	2.999	2.798	2.640	2.614	2.519	2.455	2.429	2.408
21.45	20.89	18.30	18.09	17.92	17.79	20.30	20.23	20.19	17.88	17.87
0.602	0.684	0.728	0.785	0.843	0.895	0.903	0.938	0.962	0.978	0.987
4.201	4.354	4.446	4.577	4.744	4.943	4.984	5.193	5.422	5.667	5.895
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.393	2.384	2.371	2.368	2.365	2.363	2.380	2.378	2.377	2.375	2.378
17.86	17.86	24.59	24.58	24.58	24.57	17.82	17.83	17.83	17.84	17.86
0.993	0.996	0.998	0.999	0.999	1.000	1.000	1.000	1.000	1.000	1.000
6.137	6.383	6.686	6.934	7.182	7.432	7.631	7.881	8.131	8.381	8.632
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3				
2.376	2.374	2.372	2.365	2.362	2.358					
17.89	17.95	18.06	22.80	23.24	24.08					
1.000	1.000	1.000	1.000	1.000	1.000					
8.883	9.133	9.385	9.682	9.936	10.195					
84	DIHYDROXYTARTARIC -1 -2	ACID 32.0 60.0	1.947 4.004	C4H6O8 182.09			pHL RE t Z pHS	3.00 2.262 32.61 1.174 3.310	3.25 2.111 30.62 1.276 3.507	3.50 1.967 29.92 1.389 3.702
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
1.832	1.713	1.706	1.627	1.567	1.525	1.519	1.492	1.476	1.478	1.472
29.90	30.22	27.50	27.84	28.15	28.38	31.45	31.60	31.70	28.99	29.02
1.514	1.641	1.663	1.760	1.842	1.902	1.905	1.943	1.967	1.981	1.989
3.906	4.122	4.162	4.363	4.584	4.819	4.835	5.073	5.317	5.563	5.807
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
1.468	1.465	1.455	1.454	1.452	1.449	1.467	1.466	1.465	1.464	1.466
29.04	29.05	37.26	37.25	37.24	37.22	29.03	29.03	29.04	29.04	29.06
1.994	1.996	1.998	1.999	1.999	2.000	2.000	2.000	2.000	2.000	2.000
6.055	6.304	6.576	6.825	7.075	7.324	7.554	7.803	8.053	8.303	8.554
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 12 , 7				
1.465	1.463	1.462	1.454	1.451	1.449					
29.09	29.13	29.22	34.82	35.11	35.64					
2.000	2.000	2.000	2.000	2.000	2.000					
8.804	9.054	9.304	9.572	9.823	10.075					

85	2,3-DIMETHYLBENZOIC -1	27.1	ACID 3.738		C9H10O2 150.18		pHL RE t Z pHS	3.00 5.679 29.86 0.523 3.751	3.25 5.001 27.61 0.596 3.878	3.50 4.491 26.09 0.665 4.007
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
4.067	3.722	3.657	3.471	3.319	3.212	3.193	3.130	3.090	3.091	3.077
24.97	24.19	20.94	20.60	20.35	20.19	23.42	23.32	23.27	20.31	20.29
0.737	0.808	0.829	0.875	0.917	0.948	0.952	0.971	0.983	0.990	0.994
4.155	4.330	4.390	4.550	4.743	4.963	5.002	5.230	5.470	5.700	5.940
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
3.067	3.060	3.041	3.038	3.035	3.032	3.063	3.061	3.058	3.055	3.060
20.28	20.28	28.92	28.91	28.90	28.89	20.24	20.24	20.25	20.27	20.30
0.997	0.998	0.999	0.999	1.000	1.000	1.000	1.000	1.000	1.000	1.000
6.186	6.434	6.752	7.000	7.250	7.500	7.684	7.934	8.184	8.434	8.687
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 11 , 1				
3.057	3.054	3.049	3.036	3.030	3.022					
20.35	20.44	20.61	26.90	27.72	29.38					
1.000	1.000	1.000	1.000	1.000	1.000					
8.937	9.188	9.441	9.757	10.015	10.281					
86	2,4-DIMETHYLBENZOIC -1	27.1	ACID 4.182		C9H10O2 150.18		pHL RE t Z pHS	3.00 7.891 29.08 0.372 3.932	3.25 6.765 27.14 0.436 4.045	3.50 5.912 25.82 0.500 4.156
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
5.181	4.552	4.303	3.980	3.687	3.455	3.409	3.269	3.174	3.139	3.106
24.83	24.12	20.81	20.55	20.33	20.18	23.39	23.31	23.26	20.31	20.29
0.573	0.655	0.700	0.758	0.821	0.878	0.889	0.928	0.956	0.974	0.985
4.282	4.431	4.518	4.646	4.809	5.004	5.050	5.257	5.484	5.723	5.950
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
3.084	3.069	3.046	3.041	3.037	3.033	3.064	3.061	3.059	3.055	3.060
20.28	20.27	28.92	28.91	28.90	28.89	20.24	20.24	20.25	20.27	20.30
0.991	0.995	0.998	0.999	0.999	1.000	1.000	1.000	1.000	1.000	1.000
6.190	6.436	6.754	7.001	7.250	7.500	7.684	7.934	8.184	8.434	8.687
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 11 , 1				
3.057	3.054	3.049	3.036	3.030	3.022					
20.35	20.44	20.61	26.90	27.72	29.38					
1.000	1.000	1.000	1.000	1.000	1.000					
8.937	9.188	9.441	9.757	10.015	10.281					
87	2,5-DIMETHYLBENZOIC -1	27.1	ACID 3.977		C9H10O2 150.18		pHL RE t Z pHS	3.00 6.727 29.42 0.439 3.845	3.25 5.832 27.35 0.508 3.965	3.50 5.156 25.94 0.576 4.083
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
4.583	4.100	3.956	3.702	3.482	3.317	3.287	3.189	3.125	3.111	3.089
24.90	24.15	20.87	20.57	20.34	20.18	23.41	23.32	23.27	20.31	20.29
0.651	0.731	0.764	0.818	0.872	0.916	0.923	0.952	0.972	0.983	0.990
4.219	4.379	4.454	4.596	4.774	4.981	5.024	5.242	5.476	5.710	5.944
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
3.074	3.064	3.043	3.039	3.036	3.032	3.063	3.061	3.058	3.055	3.060
20.28	20.28	28.92	28.91	28.90	28.89	20.24	20.24	20.25	20.27	20.30
0.994	0.997	0.998	0.999	1.000	1.000	1.000	1.000	1.000	1.000	1.000
6.188	6.435	6.753	7.001	7.250	7.500	7.684	7.934	8.184	8.434	8.687
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 11 , 1				
3.057	3.054	3.049	3.036	3.030	3.022					
20.35	20.44	20.61	26.90	27.72	29.38					
1.000	1.000	1.000	1.000	1.000	1.000					
8.937	9.188	9.441	9.757	10.015	10.281					

88	2,6-DIMETHYLBENZOIC -1	27.1	ACID 3.246		C9H10O2 150.18		pHL RE t Z pHS	3.00 4.280 30.85 0.701 3.585	3.25 3.917 28.18 0.768 3.734	3.50 3.653 26.38 0.825 3.888
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
3.446	3.292	3.295	3.210	3.147	3.106	3.097	3.073	3.057	3.072	3.066
25.10	24.24	21.06	20.65	20.36	20.19	23.43	23.33	23.27	20.31	20.29
0.877	0.919	0.925	0.950	0.969	0.982	0.983	0.990	0.994	0.997	0.998
4.064	4.267	4.299	4.490	4.709	4.944	4.979	5.218	5.464	5.691	5.936
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
3.061	3.056	3.040	3.037	3.035	3.031	3.063	3.061	3.058	3.055	3.060
20.28	20.28	28.92	28.91	28.90	28.89	20.24	20.24	20.25	20.27	20.30
0.999	0.999	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
6.184	6.434	6.751	7.000	7.250	7.499	7.684	7.934	8.184	8.434	8.687
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 11 , 1				
3.057	3.054	3.049	3.036	3.030	3.022					
20.35	20.44	20.61	26.90	27.72	29.38					
1.000	1.000	1.000	1.000	1.000	1.000					
8.937	9.188	9.441	9.757	10.015	10.281					
89	3,4-DIMETHYLBENZOIC -1	27.1	ACID 4.408		C9H10O2 150.18		pHL RE t Z pHS	3.00 9.540 28.76 0.306 4.032	3.25 8.094 26.94 0.362 4.140	3.50 6.999 25.70 0.420 4.244
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
6.053	5.224	4.806	4.394	4.005	3.679	3.604	3.402	3.258	3.185	3.135
24.76	24.08	20.75	20.51	20.32	20.17	23.38	23.30	23.26	20.31	20.29
0.488	0.568	0.623	0.684	0.753	0.822	0.838	0.890	0.930	0.959	0.975
4.361	4.499	4.597	4.712	4.859	5.038	5.090	5.282	5.498	5.744	5.960
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
3.101	3.079	3.051	3.044	3.038	3.033	3.064	3.062	3.059	3.055	3.060
20.28	20.27	28.92	28.91	28.90	28.89	20.24	20.24	20.25	20.27	20.30
0.985	0.992	0.996	0.998	0.999	0.999	1.000	1.000	1.000	1.000	1.000
6.195	6.439	6.757	7.002	7.251	7.500	7.685	7.934	8.184	8.434	8.687
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 11 , 1				
3.057	3.054	3.049	3.036	3.030	3.022					
20.35	20.44	20.61	26.90	27.72	29.38					
1.000	1.000	1.000	1.000	1.000	1.000					
8.937	9.188	9.441	9.757	10.015	10.281					
90	3,5-DIMETHYLBENZOIC -1	27.1	ACID 4.301		C9H10O2 150.18		pHL RE t Z pHS	3.00 8.706 28.91 0.336 3.984	3.25 7.420 27.03 0.396 4.094	3.50 6.448 25.76 0.457 4.202
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
5.610	4.880	4.550	4.182	3.841	3.563	3.503	3.332	3.213	3.161	3.120
24.79	24.10	20.78	20.53	20.32	20.18	23.39	23.31	23.26	20.31	20.29
0.528	0.609	0.660	0.720	0.786	0.850	0.864	0.910	0.944	0.967	0.980
4.323	4.466	4.558	4.680	4.834	5.021	5.070	5.269	5.491	5.733	5.955
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
3.092	3.074	3.048	3.042	3.037	3.033	3.064	3.061	3.059	3.055	3.060
20.28	20.27	28.92	28.91	28.90	28.89	20.24	20.24	20.25	20.27	20.30
0.988	0.993	0.997	0.998	0.999	0.999	1.000	1.000	1.000	1.000	1.000
6.193	6.437	6.755	7.002	7.250	7.500	7.684	7.934	8.184	8.434	8.687
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 11 , 1				
3.057	3.054	3.049	3.036	3.030	3.022					
20.35	20.44	20.61	26.90	27.72	29.38					
1.000	1.000	1.000	1.000	1.000	1.000					
8.937	9.188	9.441	9.757	10.015	10.281					

91	DIMETHYLMALONIC ACID			C5H8O4		pHL		3.00	3.25	3.50
	-1	25.5	3.166		132.12	RE	4.339	4.000	3.750	
	-2	51.1	6.059			t	32.53	29.62	27.71	
						Z	0.738	0.803	0.858	
						pHS	3.577	3.729	3.888	
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
3.551	3.392	3.394	3.284	3.170	3.039	3.007	2.845	2.653	2.388	2.234
26.40	25.62	22.26	22.04	22.13	22.58	26.19	27.03	28.17	26.80	27.78
0.909	0.953	0.960	0.994	1.032	1.079	1.089	1.156	1.246	1.412	1.519
4.069	4.275	4.309	4.505	4.726	4.960	5.005	5.235	5.465	5.779	5.962
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.090	1.971	1.850	1.804	1.771	1.748	1.764	1.756	1.750	1.746	1.748
28.70	29.46	39.86	40.17	40.37	40.48	30.84	30.89	30.92	30.94	30.97
1.634	1.742	1.854	1.905	1.942	1.966	1.980	1.988	1.993	1.996	1.998
6.162	6.378	6.687	6.896	7.127	7.368	7.590	7.834	8.081	8.330	8.581
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 2 , 7				
1.745	1.743	1.740	1.728	1.724	1.720					
31.01	31.07	31.18	37.90	38.31	39.08					
1.999	1.999	2.000	2.000	2.000	2.000					
8.831	9.081	9.331	9.609	9.860	10.113					
92	2,6-DINITROBENZOIC ACID			C7H4N2O6		pHL		3.00	3.25	3.50
	-1	31.3	1.140		212.12	RE	2.628	2.622	2.618	
						t	29.60	26.34	24.30	
						Z	0.994	0.996	0.998	
						pHS	3.321	3.525	3.731	
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
2.616	2.613	2.628	2.626	2.624	2.622	2.618	2.616	2.613	2.628	2.627
22.98	22.18	19.51	19.08	18.82	18.67	21.45	21.37	21.32	18.77	18.75
0.999	0.999	0.999	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
3.952	4.186	4.191	4.417	4.656	4.901	4.929	5.175	5.424	5.653	5.901
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.625	2.623	2.612	2.611	2.609	2.607	2.629	2.627	2.626	2.623	2.627
18.74	18.74	26.17	26.16	26.16	26.15	18.70	18.71	18.71	18.73	18.75
1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
6.151	6.400	6.708	6.958	7.208	7.458	7.651	7.901	8.151	8.401	8.653
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3 , 1				
2.625	2.622	2.619	2.610	2.607	2.601					
18.79	18.86	18.98	24.28	24.83	25.91					
1.000	1.000	1.000	1.000	1.000	1.000					
8.903	9.154	9.406	9.710	9.966	10.226					
93	3,5-DINITROBENZOIC ACID			C7H4N2O6		pHL		3.00	3.25	3.50
	-1	29.3	2.824		212.12	RE	3.373	3.187	3.058	
						t	29.90	27.07	25.19	
						Z	0.824	0.874	0.912	
						pHS	3.462	3.631	3.806	
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
2.963	2.897	2.910	2.873	2.847	2.830	2.825	2.814	2.807	2.822	2.819
23.91	23.08	20.19	19.77	19.50	19.33	22.32	22.23	22.17	19.45	19.43
0.942	0.964	0.966	0.979	0.987	0.993	0.993	0.996	0.998	0.999	0.999
4.003	4.222	4.240	4.449	4.679	4.920	4.950	5.194	5.442	5.670	5.917
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.816	2.813	2.800	2.798	2.796	2.793	2.819	2.818	2.815	2.813	2.817
19.42	19.41	27.38	27.37	27.36	27.35	19.37	19.38	19.39	19.40	19.43
1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
6.166	6.415	6.727	6.977	7.227	7.477	7.666	7.916	8.166	8.416	8.668
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 2 , 1				
2.815	2.811	2.808	2.797	2.793	2.786					
19.47	19.55	19.69	25.42	26.08	27.39					
1.000	1.000	1.000	1.000	1.000	1.000					
8.919	9.169	9.421	9.731	9.988	10.250					

94	2,3-DINITROPHENOL -1	31.3	4.018	C6H4N2O5 184.11	pHL RE t Z pHS	3.00 6.137 26.43 0.414 3.842	3.25 5.293 24.73 0.481 3.959	3.50 4.658 23.58 0.549 4.075		
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
4.118	3.658	3.490	3.254	3.046	2.886	2.859	2.763	2.699	2.678	2.656
22.71	22.08	19.24	18.99	18.80	18.66	21.43	21.36	21.31	18.77	18.75
0.623	0.704	0.743	0.799	0.855	0.904	0.912	0.944	0.967	0.981	0.988
4.206	4.362	4.446	4.583	4.754	4.956	4.998	5.211	5.442	5.682	5.913
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.642	2.632	2.617	2.614	2.611	2.608	2.629	2.627	2.626	2.623	2.627
18.74	18.74	26.17	26.16	26.16	26.15	18.70	18.71	18.71	18.73	18.75
0.993	0.996	0.998	0.999	0.999	1.000	1.000	1.000	1.000	1.000	1.000
6.156	6.403	6.711	6.959	7.208	7.458	7.651	7.901	8.151	8.401	8.653
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3				
2.625	2.622	2.619	2.610	2.607	2.601					
18.79	18.86	18.98	24.28	24.83	25.91					
1.000	1.000	1.000	1.000	1.000	1.000					
8.903	9.154	9.406	9.710	9.966	10.226					
95	2,4-DINITROPHENOL -1	31.3	4.110	C6H4N2O5 184.11	pHL RE t Z pHS	3.00 6.592 26.29 0.384 3.881	3.25 5.658 24.65 0.449 3.996	3.50 4.955 23.53 0.515 4.108		
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
4.354	3.837	3.624	3.363	3.127	2.941	2.907	2.795	2.719	2.689	2.663
22.68	22.07	19.22	18.98	18.80	18.66	21.43	21.36	21.31	18.77	18.75
0.588	0.670	0.715	0.772	0.832	0.887	0.896	0.933	0.959	0.976	0.986
4.235	4.386	4.476	4.606	4.770	4.967	5.010	5.218	5.446	5.689	5.916
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.646	2.634	2.618	2.614	2.611	2.608	2.629	2.628	2.626	2.623	2.627
18.74	18.74	26.17	26.16	26.16	26.15	18.70	18.71	18.71	18.73	18.75
0.992	0.995	0.998	0.999	0.999	1.000	1.000	1.000	1.000	1.000	1.000
6.157	6.403	6.712	6.960	7.209	7.458	7.651	7.901	8.151	8.401	8.653
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3 , 1				
2.625	2.622	2.619	2.610	2.607	2.601					
18.79	18.86	18.98	24.28	24.83	25.91					
1.000	1.000	1.000	1.000	1.000	1.000					
8.903	9.154	9.406	9.710	9.966	10.226					
96	2,5-DINITROPHENOL -1	31.3	5.216	C6H4N2O5 184.11	pHL RE t Z pHS	3.00 18.229 25.28 0.136 4.397	3.25 15.122 24.01 0.164 4.493	3.50 12.792 23.14 0.194 4.581		
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
10.752	8.904	7.235	6.453	5.638	4.860	4.523	4.022	3.581	3.123	2.980
22.47	21.95	19.02	18.88	18.76	18.66	21.36	21.32	21.29	18.76	18.75
0.232	0.281	0.350	0.393	0.451	0.526	0.566	0.639	0.720	0.835	0.876
4.677	4.787	4.923	5.002	5.104	5.232	5.302	5.433	5.594	5.885	6.030
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.854	2.762	2.679	2.651	2.632	2.620	2.637	2.632	2.628	2.624	2.628
18.74	18.74	26.16	26.16	26.15	26.15	18.70	18.71	18.71	18.73	18.75
0.916	0.947	0.974	0.984	0.991	0.995	0.997	0.998	0.999	0.999	1.000
6.217	6.434	6.752	6.976	7.215	7.461	7.656	7.903	8.152	8.401	8.653
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3 , 1				
2.625	2.622	2.619	2.611	2.607	2.601					
18.79	18.86	18.98	24.28	24.83	25.91					
1.000	1.000	1.000	1.000	1.000	1.000					
8.904	9.154	9.406	9.710	9.966	10.226					

CHROMATOGRAPHIC DATA (1983)

D43

97	2,6-DINITROPHENOL -1			31.3	3.706	C6H4N2O5 184.11		pHL RE t Z pHS	3.00 4.911 26.94 0.521 3.714	3.25 4.317 25.03 0.983 3.842	3.50 3.873 23.74 0.664 3.971
	3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
	3.505	3.206	3.140	2.980	2.850	2.758	2.744	2.690	2.656	2.653	2.641
	22.79	22.11	19.31	19.02	18.81	18.66	21.44	21.36	21.31	18.77	18.75
	0.736	0.807	0.830	0.876	0.917	0.948	0.952	0.971	0.983	0.990	0.994
	4.119	4.294	4.359	4.518	4.711	4.930	4.966	5.194	5.433	5.668	5.907
	6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
	2.633	2.627	2.614	2.612	2.610	2.607	2.629	2.627	2.626	2.623	2.627
	18.74	18.74	26.17	26.16	26.16	26.15	18.70	18.71	18.71	18.73	18.75
	0.997	0.998	0.999	0.999	1.000	1.000	1.000	1.000	1.000	1.000	1.000
	6.153	6.402	6.710	6.959	7.208	7.458	7.651	7.901	8.151	8.401	8.653
	8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3 , 1				
	2.625	2.622	2.619	2.610	2.607	2.601					
	18.79	18.86	18.98	24.28	24.83	25.91					
	1.000	1.000	1.000	1.000	1.000	1.000					
	8.903	9.154	9.406	9.710	9.966	10.226					
98	3,4-DINITROPHENOL -1			31.3	5.424	C6H4N2O5 184.11		pHL RE t Z pHS	3.00 22.598 25.19 0.109 4.498	3.25 18.690 23.95 0.132 4.592	3.50 15.762 23.10 0.157 4.679
	3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
	13.196	10.865	8.613	7.652	6.642	5.664	5.187	4.560	3.991	3.316	3.139
	22.44	21.94	19.00	18.87	18.75	18.66	21.35	21.31	21.28	18.76	18.75
	0.188	0.229	0.292	0.330	0.381	0.449	0.491	0.561	0.643	0.784	0.830
	4.772	4.879	5.018	5.093	5.189	5.308	5.382	5.501	5.649	5.950	6.078
	6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
	2.971	2.838	2.715	2.674	2.646	2.628	2.642	2.635	2.630	2.626	2.628
	18.75	18.74	26.16	26.16	26.15	26.15	18.70	18.71	18.71	18.73	18.75
	0.875	0.921	0.960	0.975	0.985	0.992	0.995	0.997	0.998	0.999	0.999
	6.247	6.451	6.774	6.986	7.220	7.463	7.660	7.904	8.152	8.401	8.654
	8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3 , 1				
	2.625	2.623	2.619	2.611	2.607	2.601					
	18.79	18.86	18.98	24.28	24.83	25.91					
	1.000	1.000	1.000	1.000	1.000	1.000					
	8.904	9.154	9.406	9.710	9.966	10.226					
99	3,5-DINITROPHENOL -1			31.3	6.699	C6H4N2O5 184.11		pHL RE t Z pHS	3.00 91.107 24.90 0.027 5.129	3.25 74.708 23.76 0.032 5.218	3.50 62.465 22.97 0.039 5.298
	3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
	51.713	41.886	30.334	26.656	22.695	18.729	15.913	13.504	11.147	6.575	6.033
	22.35	21.88	18.94	18.84	18.75	18.66	21.28	21.25	21.23	18.76	18.76
	0.047	0.058	0.081	0.092	0.108	0.132	0.156	0.184	0.223	0.386	0.421
	5.383	5.480	5.632	5.693	5.770	5.865	5.949	6.035	6.139	6.472	6.536
	6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
	5.390	4.722	3.536	3.333	3.123	2.941	2.826	2.755	2.702	2.667	2.652
	18.75	18.75	26.08	26.10	26.11	26.12	18.71	18.71	18.72	18.73	18.75
	0.473	0.542	0.729	0.775	0.829	0.882	0.927	0.952	0.970	0.983	0.990
	6.625	6.743	7.098	7.204	7.351	7.537	7.766	7.955	8.175	8.412	8.663
	8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3				
	2.639	2.630	2.624	2.613	2.608	2.602					
	18.79	18.86	18.98	24.28	24.83	25.92					
	0.994	0.997	0.998	0.999	1.000	1.000					
	8.908	9.156	9.407	9.711	9.966	10.227					

100	DIPHENYLACETIC ACID				C14H12O2		pHL	3.00	3.25	3.50
	-1	26.2	3.939		212.20		RE	6.729	5.853	5.190
							t	30.24	28.06	26.57
							Z	0.455	0.525	0.594
							pHS	3.835	3.956	4.076
	3.75	4.00	4.00	4.25	4.50	4.75	5.00	5.25	5.50	5.75
	4.629	4.160	4.034	3.784	3.571	3.413	3.290	3.230	3.221	3.199
	25.47	24.69	21.31	20.99	20.74	20.58	23.92	23.77	20.71	20.69
	0.668	0.747	0.777	0.830	0.881	0.923	0.930	0.974	0.985	0.991
	4.214	4.378	4.448	4.594	4.775	4.985	5.027	5.483	5.716	5.951
	6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.50
	3.185	3.175	3.153	3.149	3.145	3.141	3.175	3.173	3.170	3.172
	20.68	20.67	29.63	29.62	29.61	29.60	20.64	20.64	20.65	20.70
	0.995	0.997	0.999	0.999	1.000	1.000	1.000	1.000	1.000	1.000
	6.196	6.443	6.763	7.011	7.260	7.510	7.692	7.942	8.192	8.695
	8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 2 , 7			
	3.169	3.165	3.160	3.145	3.139	3.130				
	20.76	20.86	21.04	27.59	28.49	30.34				
	1.000	1.000	1.000	1.000	1.000	1.000				
	8.946	9.197	9.449	9.769	10.027	10.295				
101	DIPROPYLMALONIC ACID				C9H16O4		pHL	3.00	3.25	3.50
	-1	23.0	2.067		188.22		RE	3.743	3.697	3.666
	-2	46.0	7.510				t	36.55	32.53	29.94
							Z	0.964	0.976	0.985
							pHS	3.458	3.646	3.837
	3.75	4.00	4.00	4.25	4.50	4.75	5.00	5.25	5.50	5.75
	3.645	3.630	3.658	3.648	3.638	3.626	3.616	3.578	3.579	3.526
	28.21	27.12	23.42	22.86	22.51	22.33	26.19	26.13	22.82	23.09
	0.991	0.995	0.995	0.998	1.000	1.003	1.003	1.006	1.012	1.037
	4.047	4.274	4.282	4.498	4.733	4.976	5.013	5.259	5.745	5.991
	6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.50
	3.442	3.316	2.992	2.803	2.605	2.417	2.275	2.175	2.095	2.007
	23.55	24.27	36.53	37.70	38.92	40.09	30.52	31.12	31.59	32.18
	1.064	1.108	1.231	1.322	1.432	1.554	1.693	1.779	1.854	1.947
	6.237	6.481	6.879	7.073	7.272	7.480	7.728	7.918	8.134	8.617
	8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 12 , 7			
	1.985	1.970	1.960	1.941	1.934	1.928				
	32.34	32.48	32.66	40.26	40.79	41.79				
	1.969	1.982	1.990	1.995	1.997	1.998				
	8.858	9.104	9.353	9.636	9.888	10.142				
102	DITHIONIC ACID				H2S2O6		pHL	3.00	3.25	3.50
	-1	48.2	0.200		162.13		RE	1.208	1.114	1.039
	-2	96.4	3.400				t	28.62	27.30	26.79
							Z	1.421	1.550	1.671
							pHS	3.118	3.340	3.557
	3.75	4.00	4.00	4.25	4.50	4.75	5.00	5.25	5.50	5.75
	0.983	0.944	0.943	0.920	0.904	0.895	0.894	0.888	0.885	0.884
	26.62	26.60	24.87	24.88	24.91	24.93	26.76	26.78	26.79	25.11
	1.775	1.855	1.862	1.914	1.948	1.970	1.970	1.983	1.990	1.997
	3.780	4.013	4.036	4.264	4.503	4.748	4.994	5.242	5.496	5.744
	6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.50
	0.884	0.883	0.880	0.880	0.879	0.879	0.883	0.883	0.883	0.883
	25.12	25.12	30.06	30.06	30.05	30.05	25.09	25.09	25.10	25.11
	1.998	1.999	1.999	2.000	2.000	2.000	2.000	2.000	2.000	2.000
	5.994	6.243	6.487	6.736	6.986	7.236	7.493	7.743	7.993	8.493
	8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 2 , 8			
	0.883	0.883	0.883	0.880	0.880	0.879				
	25.12	25.15	25.19	28.52	28.64	28.86				
	2.000	2.000	2.000	2.000	2.000	2.000				
	8.743	8.993	9.243	9.487	9.737	9.987				

CHROMATOGRAPHIC DATA (1983)

D45

103	DITHIONOUS -1 -2	ACID 34.5 68.9	0.300 2.500		H2S2O4 130.13		pHL RE t Z pHS	3.00 1.349 38.20 1.852 3.110	3.25 1.313 34.76 1.906 3.334	3.50 1.290 32.73 1.943 3.563
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
1.276	1.267	1.274	1.269	1.265	1.263	1.260	1.258	1.256	1.263	1.263
31.53	30.83	28.34	27.95	27.72	27.60	30.24	30.17	30.13	27.71	27.70
1.966	1.980	1.981	1.989	1.994	1.996	1.996	1.998	1.999	1.999	2.000
3.800	4.043	4.048	4.289	4.535	4.784	4.790	5.039	5.288	5.534	5.783
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
1.262	1.261	1.254	1.254	1.252	1.251	1.263	1.263	1.262	1.261	1.262
27.69	27.69	34.75	34.74	34.73	34.72	27.65	27.65	27.65	27.66	27.67
2.000	2.000	2.000	2.000	2.000	2.000	2.000	2.000	2.000	2.000	2.000
6.033	6.283	6.546	6.796	7.046	7.295	7.533	7.783	8.033	8.283	8.533
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 2 , 8				
1.262	1.261	1.260	1.254	1.253	1.251					
27.69	27.73	27.80	32.60	32.82	33.23					
2.000	2.000	2.000	2.000	2.000	2.000					
8.783	9.034	9.284	9.544	9.794	10.045					
104	DODECYLSULFONIC -1	ACID 24.9	1.000		C12H26O3S 250.40		pHL RE t Z pHS	3.00 3.334 34.69 0.996 3.407	3.25 3.330 30.80 0.998 3.604	3.50 3.327 28.32 0.999 3.803
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
3.325	3.323	3.348	3.346	3.344	3.340	3.333	3.329	3.326	3.351	3.349
26.69	25.67	22.28	21.74	21.41	21.21	24.76	24.64	24.57	21.35	21.32
0.999	0.999	0.999	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
4.018	4.249	4.254	4.474	4.711	4.955	4.989	5.235	5.484	5.707	5.955
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
3.346	3.342	3.322	3.320	3.317	3.313	3.352	3.349	3.346	3.343	3.349
21.31	21.30	30.75	30.74	30.73	30.72	21.26	21.27	21.28	21.30	21.34
1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
6.204	6.454	6.776	7.026	7.275	7.525	7.705	7.955	8.205	8.455	8.708
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 2 , 12				
3.345	3.341	3.336	3.318	3.311	3.300					
21.40	21.51	21.72	28.69	29.73	31.93					
1.000	1.000	1.000	1.000	1.000	1.000					
8.959	9.210	9.463	9.787	10.047	10.317					
105	ENANTHYLIC -1	ACID 28.4	4.893		C7H14O2 130.19		pHL RE t Z pHS	3.00 14.349 27.30 0.192 4.251	3.25 11.979 25.76 0.230 4.351	3.50 10.194 24.71 0.271 4.444
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
8.638	7.241	6.209	5.574	4.933	4.346	4.147	3.769	3.462	3.219	3.114
23.91	23.31	20.09	19.92	19.76	19.64	22.66	22.61	22.57	19.77	19.76
0.321	0.385	0.454	0.507	0.575	0.656	0.688	0.759	0.829	0.901	0.932
4.547	4.667	4.787	4.879	4.996	5.142	5.206	5.360	5.545	5.815	5.993
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
3.033	2.979	2.927	2.911	2.900	2.892	2.918	2.915	2.911	2.908	2.912
19.75	19.75	27.97	27.97	27.96	27.95	19.71	19.71	19.72	19.74	19.76
0.957	0.974	0.988	0.993	0.996	0.998	0.998	0.999	1.000	1.000	1.000
6.206	6.438	6.756	6.994	7.239	7.487	7.675	7.924	8.173	8.423	8.676
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 2 , 7				
2.909	2.906	2.902	2.890	2.885	2.878					
19.81	19.90	20.05	25.99	26.71	28.15					
1.000	1.000	1.000	1.000	1.000	1.000					
8.926	9.177	9.429	9.741	9.998	10.262					

106	o-ETHOXYBENZOIC ACID -1 26.6 4.208				C9H10O3 166.18		pHL RE t Z pHS	3.00 8.187 29.45 0.365 3.946	3.25 7.011 27.48 0.428 4.058	3.50 6.121 26.13 0.492 4.169
	3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50
	5.357	4.697	4.433	4.095	3.787	3.542	3.493	3.344	3.243	3.205
	25.13	24.41	21.03	20.76	20.55	20.39	23.67	23.59	23.54	20.53
	0.565	0.647	0.692	0.751	0.815	0.873	0.885	0.925	0.954	0.973
	4.294	4.442	4.529	4.656	4.818	5.011	5.058	5.264	5.490	5.729
	6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25
	3.147	3.131	3.106	3.101	3.097	3.092	3.125	3.123	3.120	3.116
	20.50	20.49	29.31	29.30	29.29	29.28	20.45	20.46	20.47	20.48
	0.991	0.995	0.997	0.999	0.999	1.000	1.000	1.000	1.000	1.000
	6.195	6.441	6.760	7.007	7.256	7.505	7.689	7.938	8.188	8.438
	8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 11 , 2			
	3.118	3.115	3.110	3.096	3.090	3.081				
	20.57	20.67	20.85	27.28	28.14	29.90				
	1.000	1.000	1.000	1.000	1.000	1.000				
	8.942	9.193	9.445	9.763	10.022	10.288				
107	p-ETHOXYBENZOIC ACID -1 26.6 4.796				C9H10O3 166.18		pHL RE t Z pHS	3.00 13.806 28.73 0.214 4.212	3.25 11.558 27.01 0.256 4.314	3.50 9.861 25.84 0.301 4.410
	3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50
	8.384	7.064	6.179	5.560	4.945	4.393	4.221	3.867	3.588	3.391
	24.95	24.30	20.87	20.67	20.50	20.37	23.61	23.56	23.52	20.52
	0.355	0.424	0.490	0.546	0.616	0.697	0.725	0.794	0.858	0.917
	4.515	4.639	4.752	4.848	4.972	5.126	5.187	5.351	5.544	5.805
	6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25
	3.223	3.176	3.128	3.113	3.104	3.096	3.127	3.124	3.120	3.117
	20.50	20.49	29.30	29.30	29.29	29.28	20.45	20.46	20.47	20.48
	0.966	0.980	0.990	0.994	0.997	0.998	0.999	0.999	1.000	1.000
	6.214	6.450	6.771	7.011	7.258	7.506	7.690	7.939	8.189	8.439
	8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 11 , 2			
	3.119	3.115	3.110	3.096	3.090	3.081				
	20.57	20.67	20.85	27.28	28.14	29.90				
	1.000	1.000	1.000	1.000	1.000	1.000				
	8.942	9.193	9.445	9.763	10.022	10.288				
108	o-ETHYLBENZOIC ACID -1 26.5 3.793				C9H10O2 150.18		pHL RE t Z pHS	3.00 6.008 30.26 0.505 3.776	3.25 5.274 27.99 0.577 3.901	3.50 4.721 26.46 0.647 4.028
	3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50
	4.258	3.879	3.801	3.596	3.428	3.307	3.286	3.215	3.170	3.169
	25.33	24.53	21.21	20.87	20.61	20.44	23.75	23.66	23.60	20.58
	0.720	0.793	0.816	0.864	0.908	0.942	0.947	0.968	0.981	0.989
	4.173	4.345	4.407	4.563	4.754	4.971	5.012	5.238	5.477	5.707
	6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25
	3.142	3.134	3.114	3.111	3.108	3.104	3.137	3.135	3.132	3.129
	20.54	20.54	29.39	29.38	29.37	29.36	20.50	20.50	20.51	20.53
	0.996	0.998	0.999	0.999	1.000	1.000	1.000	1.000	1.000	1.000
	6.192	6.440	6.759	7.007	7.257	7.506	7.689	7.939	8.189	8.439
	8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 11 , 1			
	3.131	3.127	3.123	3.108	3.102	3.093				
	20.62	20.72	20.90	27.35	28.22	30.01				
	1.000	1.000	1.000	1.000	1.000	1.000				
	8.943	9.194	9.446	9.765	10.023	10.290				

CHROMATOGRAPHIC DATA (1983)

D47

109	p-ETHYLBENZOIC ACID				C9H10O2	150.18	pHL	3.00	3.25	3.50
	-1	26.5	4.353							
							RE	9.270	7.886	6.837
							t	29.32	27.41	26.12
							Z	0.323	0.381	0.441
							pHS	4.010	4.119	4.225
	3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50
	5.933	5.142	4.772	4.374	4.004	3.698	3.630	3.441	3.308	3.247
	25.14	24.44	21.04	20.79	20.58	20.43	23.72	23.64	23.59	20.57
	0.510	0.591	0.644	0.704	0.772	0.838	0.853	0.901	0.938	0.964
	4.345	4.486	4.580	4.698	4.850	5.033	5.084	5.280	5.500	5.743
	6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25
	3.170	3.150	3.122	3.115	3.110	3.105	3.138	3.135	3.132	3.129
	20.54	20.54	29.39	29.38	29.37	29.36	20.50	20.50	20.51	20.53
	0.987	0.993	0.996	0.998	0.999	0.999	1.000	1.000	1.000	1.000
	6.199	6.443	6.763	7.009	7.257	7.507	7.690	7.939	8.189	8.439
	8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 11 , 1			
	3.131	3.127	3.123	3.108	3.102	3.093				
	20.62	20.72	20.90	27.35	28.22	30.01				
	1.000	1.000	1.000	1.000	1.000	1.000				
	8.943	9.194	9.446	9.765	10.023	10.290				
110	2-ETHYLBUTYRIC ACID				C6H12O6	180.16	pHL	3.00	3.25	3.50
	-1	30.2	4.734							
							RE	11.701	9.808	8.383
							t	26.25	24.80	23.80
							Z	0.222	0.265	0.311
							pHS	4.169	4.271	4.367
	3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50
	7.143	6.033	5.257	4.746	4.237	3.778	3.639	3.346	3.114	2.948
	23.04	22.47	19.45	19.28	19.13	19.01	21.85	21.79	21.76	19.12
	0.366	0.436	0.505	0.561	0.631	0.710	0.737	0.804	0.866	0.923
	4.473	4.596	4.715	4.811	4.936	5.090	5.150	5.314	5.509	5.773
	6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25
	2.811	2.773	2.736	2.724	2.716	2.711	2.733	2.730	2.728	2.725
	19.10	19.10	26.81	26.80	26.80	26.79	19.06	19.06	19.07	19.09
	0.968	0.981	0.991	0.995	0.997	0.998	0.999	0.999	1.000	1.000
	6.183	6.420	6.734	6.974	7.221	7.469	7.661	7.909	8.159	8.409
	8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 12 , 7			
	2.726	2.723	2.720	2.710	2.706	2.700				
	19.15	19.23	19.36	24.89	25.49	26.69				
	1.000	1.000	1.000	1.000	1.000	1.000				
	8.912	9.162	9.414	9.721	9.977	10.239				
111	ETHYLFUMARIC ACID				C6H8O4	144.13	pHL	3.00	3.25	3.50
	-1	29.2	3.325							
							RE	4.166	3.778	3.493
							t	29.05	26.66	25.05
							Z	0.665	0.735	0.797
							pHS	3.592	3.737	3.886
	3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50
	3.268	3.098	3.092	2.998	2.928	2.882	2.873	2.847	2.830	2.834
	23.89	23.11	20.16	19.79	19.53	19.37	22.36	22.27	22.22	19.48
	0.854	0.902	0.910	0.940	0.962	0.978	0.979	0.988	0.993	0.996
	4.058	4.256	4.295	4.481	4.696	4.929	4.962	5.200	5.445	5.675
	6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25
	2.829	2.824	2.811	2.808	2.806	2.803	2.829	2.828	2.826	2.823
	19.45	19.45	27.44	27.43	27.43	27.42	19.41	19.42	19.42	19.44
	0.999	0.999	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
	6.167	6.416	6.729	6.978	7.228	7.478	7.666	7.916	8.166	8.416
	8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3			
	2.825	2.822	2.818	2.807	2.803	2.796				
	19.51	19.59	19.73	25.48	26.15	27.47				
	1.000	1.000	1.000	1.000	1.000	1.000				
	8.919	9.170	9.422	9.732	9.989	10.252				

112	3-ETHYLGLUTARIC ACID										
	-1	25.0	4.287		C7H12O4		pHL	3.00	3.25	3.50	
	-2	50.0	5.329		160.17		RE	8.621	7.242	6.187	
							t	31.56	29.74	28.64	
							Z	0.371	0.443	0.522	
							pHS	3.980	4.089	4.195	
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75	
5.266	4.443	3.998	3.561	3.132	2.749	2.654	2.391	2.179	2.018	1.938	
28.00	27.83	24.66	25.01	25.64	26.53	30.41	31.35	32.29	29.92	30.36	
0.617	0.737	0.831	0.939	1.077	1.239	1.284	1.437	1.589	1.748	1.828	
4.313	4.452	4.554	4.670	4.817	4.991	5.043	5.222	5.424	5.696	5.889	
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50	
1.877	1.837	1.798	1.784	1.775	1.768	1.793	1.791	1.788	1.785	1.789	
30.70	30.93	41.07	41.13	41.16	41.16	31.22	31.23	31.24	31.25	31.27	
1.891	1.935	1.965	1.980	1.988	1.993	1.996	1.998	1.999	1.999	2.000	
6.110	6.346	6.636	6.874	7.120	7.367	7.586	7.834	8.084	8.334	8.585	
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 12 , 7					
1.787	1.784	1.781	1.769	1.765	1.761						
31.31	31.37	31.49	38.37	38.80	39.61						
2.000	2.000	2.000	2.000	2.000	2.000						
8.835	9.085	9.335	9.614	9.866	10.119						
113	ETHYLMALIC ACID										
	-1	29.2	2.959		C6H8O4		pHL	3.00	3.25	3.50	
					144.13		RE	3.543	3.313	3.149	
							t	29.75	27.02	25.21	
							Z	0.786	0.843	0.888	
							pHS	3.492	3.654	3.823	
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75	
3.027	2.941	2.952	2.904	2.870	2.848	2.841	2.828	2.819	2.834	2.830	
23.95	23.13	20.22	19.80	19.53	19.37	22.36	22.27	22.22	19.48	19.46	
0.925	0.953	0.955	0.972	0.983	0.990	0.991	0.995	0.997	0.998	0.999	
4.014	4.229	4.251	4.455	4.682	4.922	4.953	5.196	5.443	5.671	5.918	
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50	
2.827	2.823	2.810	2.808	2.806	2.803	2.829	2.828	2.826	2.823	2.827	
19.45	19.45	27.44	27.43	27.43	27.42	19.41	19.42	19.42	19.44	19.46	
0.999	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	
6.167	6.416	6.729	6.978	7.228	7.478	7.666	7.916	8.166	8.416	8.669	
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3					
2.825	2.822	2.818	2.807	2.803	2.796						
19.51	19.59	19.73	25.48	26.15	27.47						
1.000	1.000	1.000	1.000	1.000	1.000						
8.919	9.170	9.422	9.732	9.989	10.252						
114	ETHYLMALONIC ACID										
	-1	25.5	2.988		C5H8O4		pHL	3.00	3.25	3.50	
	-2	51.0	5.833		132.12		RE	4.032	3.769	3.575	
							t	32.91	29.86	27.87	
							Z	0.797	0.854	0.903	
							pHS	3.534	3.695	3.862	
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75	
3.418	3.285	3.291	3.182	3.054	2.896	2.857	2.670	2.465	2.225	2.096	
26.57	25.87	22.53	22.43	22.72	23.42	27.11	28.17	29.43	27.90	28.72	
0.946	0.987	0.993	1.029	1.075	1.138	1.153	1.241	1.353	1.530	1.634	
4.051	4.264	4.296	4.499	4.723	4.956	5.003	5.227	5.451	5.754	5.936	
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50	
1.982	1.894	1.811	1.779	1.756	1.741	1.761	1.756	1.752	1.748	1.751	
29.45	30.02	40.22	40.42	40.54	40.60	30.91	30.94	30.96	30.97	31.00	
1.737	1.825	1.903	1.940	1.964	1.979	1.988	1.993	1.996	1.998	1.999	
6.140	6.363	6.663	6.884	7.121	7.366	7.586	7.832	8.081	8.330	8.581	
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 2 , 7					
1.749	1.746	1.743	1.732	1.728	1.724						
31.04	31.10	31.21	37.94	38.36	39.13						
1.999	2.000	2.000	2.000	2.000	2.000						
8.831	9.081	9.332	9.609	9.861	10.114						

CHROMATOGRAPHIC DATA (1983)

D49

115	ETHYLPROPYLMALONIC -1 -2	23.5 47.0	ACID 2.145 7.430	C8H14O4 174.20	pHL RE t Z pHS	3.00 3.685 35.92 0.957 3.454	3.25 3.631 32.01 0.971 3.641	3.50 3.595 29.48 0.982 3.832			
	3.75 3.570 27.79 0.989 4.041	4.00 3.552 26.72 0.994 4.269	4.00 3.579 23.10 0.994 4.276	4.25 3.567 22.56 0.997 4.493	4.50 3.556 22.22 1.000 4.727	4.75 3.543 22.05 1.003 4.971	4.75 3.532 25.83 1.004 5.007	5.00 3.515 25.78 1.008 5.253	5.25 3.490 25.83 1.014 5.500	5.50 3.485 22.60 1.025 5.743	5.75 3.425 22.92 1.044 5.988
	6.00 3.331 23.45 1.076 6.233	6.25 3.193 24.26 1.126 6.475	6.50 2.855 36.44 1.265 6.874	6.75 2.672 37.60 1.360 7.063	7.00 2.484 38.78 1.472 7.260	7.25 2.310 39.88 1.593 7.467	7.50 2.185 30.44 1.725 7.713	7.75 2.096 30.98 1.806 7.906	8.00 2.027 31.40 1.873 8.126	8.25 1.978 31.69 1.922 8.360	8.50 1.953 31.91 1.955 8.610
	8.75 1.934 32.05 1.974 8.852	9.00 1.922 32.17 1.985 9.099	9.25 1.913 32.34 1.991 9.348	9.50 1.896 39.75 1.995 9.631	9.75 1.889 40.26 1.997 9.882	10.00 1.883 41.20 1.999 10.136	(Ref.) : 12 , 7				
116	ETHYLSULFONIC ACID -1	41.0	-2.000	C2H6O3S 110.13	pHL RE t Z pHS	3.00 1.971 25.06 1.000 3.219	3.25 1.971 22.34 1.000 3.435	3.50 1.971 20.69 1.000 3.653			
	3.75 1.970 19.66 1.000 3.882	4.00 1.970 19.04 1.000 4.121	4.00 1.977 17.04 1.000 4.124	4.25 1.976 16.71 1.000 4.357	4.50 1.975 16.51 1.000 4.600	4.75 1.974 16.40 1.000 4.847	4.75 1.973 18.49 1.000 4.866	5.00 1.972 18.43 1.000 5.113	5.25 1.971 18.39 1.000 5.363	5.50 1.978 16.47 1.000 5.598	5.75 1.977 16.46 1.000 5.847
	6.00 1.976 16.45 1.000 6.096	6.25 1.975 16.45 1.000 6.346	6.50 1.971 22.05 1.000 6.637	6.75 1.970 22.05 1.000 6.887	7.00 1.969 22.04 1.000 7.137	7.25 1.968 22.04 1.000 7.386	7.50 1.978 16.42 1.000 7.596	7.75 1.978 16.42 1.000 7.846	8.00 1.977 16.43 1.000 8.096	8.25 1.975 16.44 1.000 8.346	8.50 1.977 16.44 1.000 8.597
	8.75 1.976 16.47 1.000 8.848	9.00 1.975 16.51 1.000 9.098	9.25 1.973 16.59 1.000 9.349	9.50 1.970 20.48 1.000 9.634	9.75 1.968 20.77 1.000 9.887	10.00 1.966 21.31 1.000 10.142	(Ref.) : 2 , 12				
117	FLUOROACETIC ACID -1	46.0	2.584	C2H3FO2 78.04	pHL RE t Z pHS	3.00 2.051 22.62 0.849 3.296	3.25 1.946 20.62 0.896 3.481	3.50 1.875 19.34 0.931 3.672			
	3.75 1.826 18.49 0.956 3.882	4.00 1.793 17.96 0.974 4.110	4.00 1.797 16.17 0.975 4.128	4.25 1.779 15.89 0.985 4.348	4.50 1.767 15.72 0.991 4.585	4.75 1.760 15.62 0.995 4.829	4.75 1.759 17.48 0.995 4.846	5.00 1.754 17.42 0.997 5.092	5.25 1.751 17.39 0.998 5.340	5.50 1.755 15.68 0.999 5.579	5.75 1.754 15.67 0.999 5.827
	6.00 1.753 15.67 1.000 6.076	6.25 1.752 15.66 1.000 6.326	6.50 1.749 20.63 1.000 6.609	6.75 1.748 20.63 1.000 6.859	7.00 1.748 20.63 1.000 7.109	7.25 1.747 20.63 1.000 7.359	7.50 1.754 15.63 1.000 7.576	7.75 1.753 15.64 1.000 7.826	8.00 1.753 15.64 1.000 8.076	8.25 1.752 15.65 1.000 8.326	8.50 1.753 15.66 1.000 8.577
	8.75 1.752 15.68 1.000 8.827	9.00 1.751 15.71 1.000 9.077	9.25 1.750 15.77 1.000 9.328	9.50 1.748 19.20 1.000 9.606	9.75 1.747 19.42 1.000 9.858	10.00 1.745 19.83 1.000 10.112	(Ref.) : 1				

118	m-FLUOROBENZOIC ACID				C7H5FO2		pHL	3.00	3.25	3.50
	-1	33.4	3.865		140.11		RE	5.181	4.502	3.993
							t	25.48	23.83	22.71
							Z	0.460	0.531	0.600
							pHS	3.768	3.890	4.011
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
3.565	3.207	3.095	2.909	2.751	2.633	2.615	2.546	2.501	2.488	2.473
21.87	21.27	18.62	18.38	18.19	18.06	20.64	20.57	20.53	18.15	18.14
0.674	0.752	0.785	0.836	0.886	0.927	0.932	0.958	0.975	0.986	0.992
4.150	4.314	4.392	4.538	4.719	4.929	4.967	5.187	5.422	5.661	5.896
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.463	2.456	2.444	2.441	2.439	2.436	2.455	2.453	2.452	2.450	2.453
18.13	18.12	25.06	25.06	25.05	25.05	18.09	18.09	18.10	18.11	18.13
0.995	0.997	0.999	0.999	1.000	1.000	1.000	1.000	1.000	1.000	1.000
6.141	6.388	6.693	6.941	7.190	7.440	7.637	7.887	8.137	8.387	8.639
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3 , 1				
2.451	2.449	2.446	2.439	2.436	2.432					
18.16	18.22	18.33	23.25	23.72	24.62					
1.000	1.000	1.000	1.000	1.000	1.000					
8.889	9.140	9.391	9.691	9.945	10.204					
119	o-FLUOROBENZOIC ACID				C7H5FO2		pHL	3.00	3.25	3.50
	-1	33.4	3.267		140.11		RE	3.588	3.255	3.012
							t	26.53	24.42	23.00
							Z	0.672	0.742	0.804
							pHS	3.546	3.693	3.845
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
2.821	2.679	2.671	2.593	2.536	2.498	2.492	2.470	2.456	2.463	2.458
21.99	21.31	18.75	18.42	18.19	18.06	20.65	20.58	20.53	18.15	18.14
0.860	0.907	0.915	0.943	0.965	0.979	0.980	0.989	0.993	0.996	0.998
4.019	4.220	4.260	4.448	4.665	4.899	4.928	5.167	5.412	5.645	5.890
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.454	2.451	2.441	2.440	2.438	2.436	2.454	2.453	2.451	2.450	2.453
18.13	18.12	25.06	25.06	25.05	25.05	18.09	18.09	18.10	18.11	18.13
0.999	0.999	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
6.138	6.387	6.691	6.940	7.190	7.440	7.637	7.887	8.137	8.387	8.639
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3 , 1				
2.451	2.449	2.446	2.439	2.436	2.432					
18.16	18.22	18.33	23.25	23.72	24.62					
1.000	1.000	1.000	1.000	1.000	1.000					
8.889	9.140	9.391	9.691	9.945	10.204					
120	p-FLUOROBENZOIC ACID				C7H5FO2		pHL	3.00	3.25	3.50
	-1	34.2	4.142		140.11		RE	6.264	5.362	4.684
							t	24.69	23.24	22.26
							Z	0.369	0.432	0.496
							pHS	3.884	3.997	4.108
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
4.103	3.601	3.370	3.121	2.893	2.710	2.677	2.566	2.490	2.455	2.429
21.50	20.95	18.34	18.13	17.96	17.85	20.36	20.30	20.26	17.94	17.92
0.568	0.650	0.700	0.758	0.819	0.876	0.887	0.926	0.955	0.974	0.984
4.232	4.380	4.477	4.602	4.763	4.955	4.999	5.202	5.428	5.675	5.900
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.412	2.401	2.387	2.383	2.380	2.377	2.394	2.393	2.391	2.389	2.392
17.91	17.91	24.68	24.68	24.67	24.67	17.87	17.88	17.88	17.90	17.91
0.991	0.995	0.997	0.999	0.999	1.000	1.000	1.000	1.000	1.000	1.000
6.139	6.385	6.689	6.935	7.184	7.434	7.632	7.882	8.132	8.382	8.634
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 2 , 1				
2.391	2.389	2.386	2.380	2.377	2.372					
17.95	18.00	18.11	22.89	23.33	24.18					
1.000	1.000	1.000	1.000	1.000	1.000					
8.884	9.135	9.386	9.684	9.938	10.197					

CHROMATOGRAPHIC DATA (1983)

D51

121	FORMIC ACID -1	(S) 56.6	3.752	CH2O2 46.03	pHL RE t Z pHS	3.00 2.988 18.60 0.464 3.659	3.25 2.593 17.73 0.536 3.782	3.50 2.300 17.12 0.605 3.904			
	3.75 2.055 16.66 0.679 4.042	4.00 1.850 16.32 0.756 4.205	4.00 1.766 14.77 0.795 4.301	4.25 1.666 14.63 0.844 4.445	4.50 1.579 14.52 0.891 4.626	4.75 1.514 14.45 0.930 4.836	4.75 1.508 15.93 0.934 4.863	5.00 1.469 15.90 0.959 5.080	5.25 1.445 15.87 0.976 5.315	5.50 1.432 14.49 0.986 5.569	5.75 1.424 14.48 0.992 5.803
	6.00 1.419 14.48 0.995 6.047	6.25 1.416 14.48 0.997 6.295	6.50 1.413 18.48 0.999 6.566	6.75 1.412 18.48 0.999 6.814	7.00 1.411 18.48 1.000 7.063	7.25 1.410 18.48 1.000 7.313	7.50 1.413 14.45 1.000 7.543	7.75 1.413 14.45 1.000 7.793	8.00 1.413 14.46 1.000 8.043	8.25 1.412 14.46 1.000 8.293	8.50 1.413 14.47 1.000 8.544
	8.75 1.413 14.48 1.000 8.794	9.00 1.412 14.51 1.000 9.044	9.25 1.412 14.55 1.000 9.294	9.50 1.411 17.28 1.000 9.561	9.75 1.410 17.43 1.000 9.812	10.00 1.409 17.69 1.000 10.063	(Ref.) : 2 , 1				
122	FUMARIC ACID -1 -2	31.0 61.2	3.019 4.384	C4H4O4 116.07	pHL RE t Z pHS	3.00 2.995 30.64 0.886 3.471	3.25 2.666 28.88 1.002 3.630	3.50 2.400 28.13 1.122 3.792			
	3.75 2.164 28.06 1.255 3.971	4.00 1.956 28.50 1.402 4.169	4.00 1.912 25.95 1.449 4.232	4.25 1.777 26.51 1.571 4.411	4.50 1.663 27.10 1.691 4.614	4.75 1.577 27.60 1.794 4.836	4.75 1.566 30.62 1.802 4.859	5.00 1.511 30.97 1.874 5.084	5.25 1.475 31.21 1.924 5.321	5.50 1.464 28.65 1.956 5.572	5.75 1.451 28.74 1.974 5.809
	6.00 1.443 28.78 1.985 6.054	6.25 1.438 28.81 1.992 6.302	6.50 1.426 36.86 1.995 6.573	6.75 1.424 36.86 1.997 6.821	7.00 1.422 36.85 1.999 7.071	7.25 1.419 36.84 1.999 7.320	7.50 1.436 28.81 2.000 7.550	7.75 1.435 28.82 2.000 7.800	8.00 1.434 28.82 2.000 8.050	8.25 1.433 28.83 2.000 8.300	8.50 1.435 28.85 2.000 8.551
	8.75 1.434 28.87 2.000 8.801	9.00 1.432 28.92 2.000 9.051	9.25 1.431 29.00 2.000 9.301	9.50 1.423 34.48 2.000 9.568	9.75 1.421 34.76 2.000 9.819	10.00 1.419 35.27 2.000 10.070	(Ref.) : 11 , 1				
123	GALLIC ACID -1	34.4	4.469	C7H6O5 170.12	pHL RE t Z pHS	3.00 8.212 24.23 0.278 4.032	3.25 6.938 22.94 0.330 4.138	3.50 5.980 22.05 0.384 4.238			
	3.75 5.149 21.37 0.447 4.350	4.00 4.412 20.85 0.524 4.482	4.00 3.948 18.22 0.590 4.597	4.25 3.604 18.05 0.649 4.703	4.50 3.267 17.91 0.717 4.840	4.75 2.975 17.80 0.790 5.009	4.75 2.905 20.28 0.809 5.062	5.00 2.721 20.23 0.866 5.242	5.25 2.584 20.19 0.913 5.451	5.50 2.499 17.88 0.950 5.710	5.75 2.452 17.87 0.968 5.916
	6.00 2.420 17.86 0.981 6.146	6.25 2.399 17.86 0.989 6.387	6.50 2.379 24.59 0.995 6.692	6.75 2.372 24.58 0.997 6.936	7.00 2.368 24.58 0.998 7.183	7.25 2.364 24.57 0.999 7.432	7.50 2.380 17.82 0.999 7.632	7.75 2.379 17.83 1.000 7.881	8.00 2.377 17.83 1.000 8.131	8.25 2.375 17.84 1.000 8.381	8.50 2.378 17.86 1.000 8.632
	8.75 2.376 17.89 1.000 8.883	9.00 2.374 17.95 1.000 9.133	9.25 2.372 18.06 1.000 9.385	9.50 2.365 22.80 1.000 9.682	9.75 2.362 23.24 1.000 9.936	10.00 2.358 24.08 1.000 10.195	(Ref.) : 3				

124	GLUCONIC ACID -1 27.2	3.662	C6H12O7 196.16	pHL RE t Z pHS	3.00 5.387 29.92 0.550 3.722	3.25 4.769 27.63 0.623 3.852	3.50 4.306 26.07 0.692 3.984			
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
3.924	3.617	3.569	3.402	3.268	3.175	3.159	3.105	3.071	3.075	3.062
24.93	24.14	20.92	20.57	20.31	20.14	23.36	23.27	23.21	20.27	20.25
0.762	0.830	0.847	0.890	0.928	0.956	0.959	0.975	0.986	0.992	0.995
4.137	4.317	4.372	4.537	4.735	4.958	4.996	5.226	5.467	5.697	5.938
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
3.054	3.047	3.029	3.026	3.023	3.020	3.051	3.049	3.047	3.043	3.048
20.24	20.23	28.84	28.84	28.83	28.82	20.20	20.20	20.21	20.22	20.25
0.997	0.998	0.999	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
6.185	6.433	6.750	6.999	7.249	7.498	7.683	7.933	8.183	8.433	8.686
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3				
3.045	3.042	3.037	3.024	3.018	3.010					
20.31	20.40	20.57	26.83	27.63	29.28					
1.000	1.000	1.000	1.000	1.000	1.000					
8.936	9.188	9.440	9.756	10.014	10.279					
125	GLUCURONIC -1 26.7	3.677	C6H10O7 194.14	pHL RE t Z pHS	3.00 5.527 30.31 0.546 3.731	3.25 4.890 27.98 0.619 3.860	3.50 4.412 26.40 0.689 3.992			
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
4.017	3.700	3.650	3.477	3.338	3.241	3.224	3.168	3.132	3.137	3.124
25.24	24.43	21.15	20.79	20.53	20.36	23.64	23.55	23.49	20.49	20.47
0.759	0.827	0.845	0.888	0.926	0.955	0.958	0.975	0.985	0.991	0.995
4.144	4.323	4.378	4.543	4.741	4.963	5.001	5.232	5.472	5.702	5.943
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
3.115	3.108	3.089	3.086	3.083	3.079	3.112	3.110	3.107	3.104	3.109
20.46	20.45	29.23	29.22	29.21	29.20	20.41	20.42	20.42	20.44	20.47
0.997	0.998	0.999	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
6.189	6.438	6.756	7.005	7.254	7.504	7.688	7.938	8.187	8.438	8.690
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 10				
3.106	3.102	3.098	3.084	3.078	3.069					
20.53	20.62	20.80	27.20	28.05	29.79					
1.000	1.000	1.000	1.000	1.000	1.000					
8.941	9.192	9.444	9.762	10.020	10.287					
126	GLUTACONIC -1 -2 56.0	3.767 5.077	C5H6O4 130.10	pHL RE t Z pHS	3.00 5.272 29.98 0.546 3.747	3.25 4.542 28.17 0.637 3.872	3.50 3.978 27.13 0.731 3.998			
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
3.486	3.048	2.879	2.610	2.352	2.125	2.081	1.920	1.793	1.709	1.661
26.62	26.65	23.88	24.42	25.23	26.21	29.64	30.52	31.27	28.97	29.28
0.840	0.967	1.036	1.150	1.286	1.436	1.466	1.600	1.722	1.833	1.891
4.140	4.307	4.394	4.542	4.718	4.917	4.959	5.157	5.374	5.641	5.850
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
1.627	1.605	1.581	1.573	1.567	1.563	1.583	1.581	1.579	1.577	1.580
29.50	29.64	38.57	38.60	38.61	38.61	29.80	29.80	29.81	29.82	29.84
1.934	1.961	1.979	1.988	1.993	1.996	1.998	1.999	1.999	2.000	2.000
6.080	6.322	6.602	6.845	7.092	7.340	7.565	7.815	8.064	8.314	8.565
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 12 , 1				
1.578	1.577	1.575	1.565	1.563	1.560					
29.87	29.92	30.01	36.07	36.40	37.03					
2.000	2.000	2.000	2.000	2.000	2.000					
8.815	9.065	9.315	9.587	9.838	10.091					

CHROMATOGRAPHIC DATA (1983)

D53

127	GLUTAMIC ACID (S)				C5H9NO4		pHL	3.00	3.25	3.50
	1	28.9	2.155		147.13		RE	8.843	7.502	6.489
	-1	-28.9	4.376				t	27.92	26.10	24.85
	-2	-49.6	9.960				Z	0.309	0.366	0.424
							pHS	4.030	4.132	4.232
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
5.614	4.846	4.447	4.072	3.717	3.420	3.353	3.169	3.038	2.971	2.926
23.90	23.20	20.05	19.81	19.62	19.47	22.47	22.40	22.35	19.59	19.58
0.492	0.573	0.630	0.690	0.758	0.827	0.843	0.893	0.932	0.961	0.976
4.345	4.481	4.581	4.695	4.842	5.021	5.072	5.264	5.481	5.728	5.944
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.895	2.875	2.850	2.842	2.836	2.828	2.850	2.839	2.821	2.792	2.748
19.57	19.57	27.65	27.66	27.68	27.71	19.65	19.75	19.93	20.24	20.80
0.986	0.992	0.997	0.999	1.001	1.004	1.006	1.012	1.021	1.036	1.066
6.180	6.423	6.738	6.984	7.232	7.481	7.675	7.924	8.174	8.424	8.693
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 11 , 5				
2.675	2.571	2.438	2.249	2.116	2.001					
21.63	22.84	24.43	32.87	34.99	37.27					
1.111	1.180	1.278	1.432	1.562	1.688					
8.938	9.182	9.421	9.715	9.938	10.169					
128	GLUTARIC ACID				C5H8O4		pHL	3.00	3.25	3.50
	-1	26.6	4.343		132.12		RE	8.432	7.032	5.967
	-2	55.6	5.272				t	30.12	28.53	27.59
							Z	0.353	0.425	0.503
							pHS	3.994	4.102	4.205
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
5.037	4.205	3.699	3.275	2.858	2.486	2.392	2.144	1.946	1.793	1.722
27.07	26.96	24.09	24.45	25.04	25.85	29.39	30.20	30.99	28.81	29.18
0.598	0.720	0.828	0.939	1.081	1.248	1.297	1.452	1.605	1.764	1.840
4.319	4.453	4.563	4.673	4.813	4.981	5.033	5.205	5.404	5.679	5.871
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
1.670	1.635	1.602	1.591	1.583	1.577	1.596	1.594	1.592	1.590	1.593
29.46	29.65	38.68	38.73	38.75	38.75	29.88	29.89	29.89	29.90	29.92
1.900	1.940	1.968	1.981	1.989	1.994	1.996	1.998	1.999	1.999	2.000
6.091	6.327	6.611	6.849	7.095	7.342	7.567	7.816	8.066	8.315	8.566
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 10 , 1				
1.591	1.589	1.587	1.578	1.575	1.572					
29.95	30.01	30.10	36.20	36.55	37.18					
2.000	2.000	2.000	2.000	2.000	2.000					
8.816	9.066	9.317	9.589	9.840	10.092					
129	GLYCERIC ACID				C3H6O4		pHL	3.00	3.25	3.50
	-1	36.4	3.737		106.08		RE	4.395	3.843	3.430
							t	24.23	22.68	21.62
							Z	0.497	0.570	0.640
							pHS	3.703	3.830	3.956
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
3.087	2.806	2.727	2.580	2.458	2.370	2.359	2.307	2.274	2.266	2.255
20.83	20.26	17.85	17.62	17.43	17.31	19.68	19.62	19.58	17.40	17.38
0.714	0.787	0.815	0.863	0.907	0.941	0.945	0.967	0.980	0.989	0.993
4.100	4.271	4.345	4.499	4.687	4.903	4.937	5.161	5.399	5.639	5.877
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.247	2.242	2.233	2.231	2.229	2.227	2.242	2.241	2.239	2.238	2.240
17.38	17.37	23.72	23.71	23.71	23.71	17.34	17.34	17.35	17.36	17.37
0.996	0.998	0.999	0.999	1.000	1.000	1.000	1.000	1.000	1.000	1.000
6.122	6.370	6.669	6.918	7.167	7.417	7.619	7.869	8.119	8.369	8.621
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 10				
2.239	2.237	2.235	2.230	2.227	2.224					
17.40	17.46	17.55	22.00	22.38	23.10					
1.000	1.000	1.000	1.000	1.000	1.000					
8.871	9.122	9.373	9.666	9.920	10.177					

130	GLYCOLIC ACID -1	42.4	3.886	C2H4O3 76.05	pHL RE t Z pHS	3.00 4.264 21.78 0.437 3.747	3.25 3.687 20.58 0.506 3.867	3.50 3.256 19.76 0.574 3.985		
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
2.892	2.585	2.463	2.309	2.175	2.072	2.059	1.998	1.958	1.940	1.927
19.13	18.67	16.59	16.41	16.26	16.16	18.18	18.12	18.09	16.23	16.22
0.649	0.728	0.768	0.821	0.873	0.917	0.923	0.951	0.971	0.983	0.990
4.119	4.278	4.370	4.509	4.684	4.890	4.924	5.139	5.372	5.620	5.852
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
1.918	1.913	1.906	1.904	1.903	1.901	1.910	1.909	1.908	1.907	1.909
16.21	16.21	21.62	21.61	21.61	21.61	16.18	16.18	16.19	16.20	16.20
0.994	0.997	0.998	0.999	0.999	1.000	1.000	1.000	1.000	1.000	1.000
6.095	6.342	6.632	6.879	7.129	7.378	7.591	7.840	8.090	8.340	8.591
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 10 , 1				
1.908	1.907	1.905	1.902	1.901	1.899					
16.23	16.27	16.34	20.09	20.36	20.85					
1.000	1.000	1.000	1.000	1.000	1.000					
8.841	9.092	9.343	9.626	9.878	10.133					
131	GLYOXYLIC ACID -1	39.6	3.337	C2H4O4 92.05	pHL RE t Z pHS	3.00 3.196 23.57 0.631 3.538	3.25 2.867 21.89 0.705 3.681	3.50 2.628 20.75 0.771 3.829		
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
2.438	2.293	2.275	2.197	2.139	2.100	2.096	2.074	2.060	2.061	2.056
19.93	19.37	17.21	16.95	16.77	16.65	18.82	18.76	18.72	16.73	16.72
0.832	0.886	0.898	0.930	0.956	0.974	0.975	0.986	0.992	0.995	0.997
3.997	4.192	4.242	4.424	4.637	4.868	4.894	5.130	5.375	5.613	5.856
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.053	2.050	2.044	2.043	2.042	2.040	2.052	2.051	2.050	2.049	2.051
16.71	16.71	22.51	22.51	22.51	22.51	16.67	16.68	16.68	16.69	16.70
0.998	0.999	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
6.104	6.353	6.646	6.896	7.145	7.395	7.603	7.853	8.103	8.353	8.604
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3				
2.050	2.048	2.046	2.042	2.040	2.038					
16.73	16.77	16.85	20.90	21.21	21.80					
1.000	1.000	1.000	1.000	1.000	1.000					
8.854	9.105	9.356	9.643	9.896	10.152					
132	HIPPURIC ACID -1 -2	25.3 55.1	2.700 7.271	C9H9NO3 179.18	pHL RE t Z pHS	3.00 3.744 33.47 0.868 3.484	3.25 3.586 30.10 0.907 3.656	3.50 3.477 27.88 0.937 3.835		
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
3.397	3.341	3.362	3.329	3.302	3.278	3.268	3.241	3.203	3.170	3.084
26.37	25.41	22.05	21.56	21.27	21.13	24.62	24.60	24.69	21.81	22.23
0.960	0.976	0.977	0.987	0.995	1.001	1.002	1.009	1.019	1.036	1.062
4.035	4.257	4.271	4.482	4.713	4.954	4.990	5.234	5.481	5.739	5.982
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.954	2.778	2.390	2.224	2.060	1.913	1.806	1.739	1.687	1.653	1.635
22.89	23.79	34.85	35.69	36.52	37.25	28.95	29.31	29.58	29.76	29.89
1.104	1.168	1.340	1.435	1.545	1.659	1.780	1.848	1.903	1.942	1.966
6.222	6.459	6.863	7.033	7.220	7.424	7.677	7.872	8.093	8.329	8.579
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 10				
1.622	1.614	1.608	1.596	1.592	1.588					
29.98	30.07	30.19	36.36	36.72	37.37					
1.981	1.989	1.994	1.997	1.998	1.999					
8.823	9.070	9.319	9.592	9.843	10.095					

CHROMATOGRAPHIC DATA (1983)

D55

133	HYDROBROMIC -1	ACID 81.0	-2.000	HBr 80.91		pHL RE t Z pHS	3.00 0.976 17.88 1.000 2.993	3.25 0.976 16.11 1.000 3.244	3.50 0.976 15.11 1.000 3.495	
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
0.976	0.976	0.976	0.976	0.976	0.976	0.976	0.976	0.976	0.976	0.976
14.56	14.25	13.27	13.09	12.99	12.94	13.97	13.94	13.93	12.97	12.96
1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
3.746	3.996	3.996	4.247	4.497	4.747	4.747	4.997	5.247	5.497	5.747
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
0.976	0.976	0.976	0.976	0.976	0.976	0.976	0.976	0.976	0.976	0.976
12.96	12.97	15.72	15.72	15.72	15.73	12.94	12.95	12.95	12.95	12.95
1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
5.997	6.247	6.496	6.746	6.996	7.246	7.497	7.747	7.997	8.247	8.497
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 2 , 12				
0.976	0.976	0.976	0.976	0.976	0.976					
12.96	12.98	13.00	14.87	14.94	15.08					
1.000	1.000	1.000	1.000	1.000	1.000					
8.747	8.997	9.247	9.496	9.746	9.996					
134	HYDROCHLORIC -1	ACID 79.1	-2.000	HCl 36.46		pHL RE t Z pHS	3.00 1.000 18.06 1.000 3.000	3.25 1.000 16.26 1.000 3.250	3.50 1.000 15.25 1.000 3.500	
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
14.68	14.36	13.36	13.18	13.08	13.02	14.08	14.05	14.03	13.05	13.05
1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
3.750	4.000	4.000	4.250	4.500	4.750	4.750	5.000	5.250	5.500	5.750
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
13.05	13.05	15.87	15.87	15.87	15.88	13.03	13.03	13.03	13.04	13.04
1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
6.000	6.250	6.500	6.750	7.000	7.250	7.500	7.750	8.000	8.250	8.500
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 1 , 12				
1.000	1.000	1.000	1.000	1.000	1.000					
13.05	13.06	13.09	15.00	15.08	15.22					
1.000	1.000	1.000	1.000	1.000	1.000					
8.750	9.000	9.250	9.500	9.750	10.000					
135	HYDROCYANIC -1	ACID 80.9	9.400	HCN 27.03		pHL RE t Z pHS	3.00 0.000 15.14 0.001 6.443	3.25 0.000 14.91 0.001 6.519	3.50 0.000 14.72 0.002 6.588	
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
14.55	14.40	13.27	13.24	13.22	13.19	14.18	14.17	14.16	13.23	13.23
0.002	0.002	0.004	0.004	0.005	0.005	0.007	0.008	0.010	0.023	0.025
6.663	6.749	6.950	6.997	7.059	7.138	7.237	7.300	7.381	7.772	7.805
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
33.026	28.390	11.914	11.207	10.217	8.972	4.575	4.271	3.864	3.380	2.314
13.22	13.22	15.91	15.91	15.91	15.90	13.17	13.16	13.16	13.16	13.15
0.028	0.033	0.079	0.084	0.092	0.105	0.208	0.223	0.246	0.282	0.415
7.855	7.922	8.319	8.347	8.391	8.453	8.797	8.834	8.890	8.969	9.220
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 2 , 4				
2.099	1.859	1.624	1.356	1.245	1.152					
13.15	13.16	13.18	15.05	15.12	15.24					
0.458	0.518	0.594	0.715	0.780	0.845					
9.295	9.398	9.530	9.761	9.910	10.093					

136	HYDROFLUORIC ACID (S)					HF		pHL	3.00	3.25	3.50
	-1	57.4	3.173			20.01		RE	2.095	1.885	1.734
								t	19.19	17.99	17.19
								Z	0.657	0.732	0.797
								pHS	3.420	3.572	3.729
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75	
1.618	1.532	1.519	1.474	1.441	1.420	1.419	1.407	1.399	1.398	1.395	
16.61	16.23	14.76	14.57	14.45	14.37	15.84	15.80	15.78	14.42	14.41	
0.855	0.904	0.914	0.943	0.965	0.979	0.980	0.988	0.993	0.996	0.998	
3.905	4.108	4.159	4.350	4.569	4.804	4.818	5.057	5.303	5.549	5.794	
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50	
1.394	1.393	1.391	1.391	1.390	1.390	1.393	1.392	1.392	1.392	1.392	
14.41	14.41	18.35	18.35	18.35	18.35	14.38	14.38	14.39	14.39	14.40	
0.999	0.999	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	
6.042	6.292	6.561	6.810	7.060	7.310	7.541	7.791	8.041	8.291	8.541	
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 1 , 7					
1.392	1.392	1.391	1.390	1.390	1.389						
14.41	14.44	14.48	17.17	17.31	17.57						
1.000	1.000	1.000	1.000	1.000	1.000						
8.792	9.042	9.292	9.558	9.809	10.060						
137	HYDROGEN AZIDE					HN3		pHL	3.00	3.25	3.50
	-1	72.0	4.720			43.00		RE	5.240	4.433	3.818
								t	16.00	15.60	15.30
								Z	0.204	0.242	0.281
								pHS	4.107	4.200	4.287
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75	
3.269	2.763	2.239	2.041	1.831	1.629	1.558	1.430	1.319	1.209	1.175	
15.05	14.84	13.57	13.51	13.46	13.41	14.55	14.52	14.50	13.41	13.41	
0.329	0.390	0.484	0.532	0.594	0.669	0.701	0.765	0.831	0.909	0.936	
4.383	4.497	4.660	4.742	4.850	4.989	5.052	5.193	5.370	5.677	5.843	
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50	
1.147	1.128	1.115	1.110	1.106	1.104	1.103	1.102	1.102	1.102	1.102	
13.40	13.40	16.51	16.51	16.51	16.52	13.38	13.38	13.38	13.39	13.39	
0.959	0.976	0.987	0.992	0.996	0.997	0.999	0.999	1.000	1.000	1.000	
6.049	6.278	6.548	6.779	7.021	7.268	7.514	7.762	8.011	8.261	8.511	
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 2					
1.101	1.101	1.101	1.101	1.101	1.101						
13.40	13.42	13.45	15.56	15.65	15.82						
1.000	1.000	1.000	1.000	1.000	1.000						
8.761	9.011	9.261	9.516	9.766	10.016						
138	HYDROGEN HEXAFLUOROPHOSPHIDE					HPF6		pHL	3.00	3.25	3.50
	-1	59.0	0.000			146.00		RE	1.353	1.352	1.352
								t	20.61	18.47	17.21
								Z	0.999	1.000	1.000
								pHS	3.094	3.328	3.562
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75	
1.352	1.351	1.353	1.353	1.353	1.352	1.352	1.352	1.351	1.353	1.353	
16.47	16.05	14.69	14.45	14.31	14.24	15.67	15.63	15.60	14.28	14.27	
1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	
3.802	4.042	4.049	4.292	4.539	4.787	4.795	5.044	5.294	5.538	5.787	
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50	
1.353	1.353	1.352	1.352	1.351	1.351	1.353	1.353	1.353	1.353	1.353	
14.27	14.27	18.10	18.10	18.10	18.10	14.24	14.25	14.25	14.26	14.26	
1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	
6.037	6.287	6.555	6.804	7.054	7.304	7.537	7.787	8.037	8.287	8.538	
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 2 , 12					
1.353	1.353	1.352	1.352	1.351	1.350						
14.27	14.30	14.34	16.95	17.09	17.33						
1.000	1.000	1.000	1.000	1.000	1.000						
8.788	9.038	9.288	9.552	9.803	10.055						

CHROMATOGRAPHIC DATA (1983)

D57

139	HYDROGEN SULFIDE -1 67.4	6.960	H2S 34.08	pHL RE t Z pHS	3.00 60.809 16.25 0.018 5.226	3.25 50.836 15.92 0.022 5.305	3.50 43.172 15.66 0.026 5.377
	3.75 4.00 4.00 4.25 4.50 4.75 4.75 5.00 5.25 5.50 5.75						
	36.218 29.649 19.491 17.435 15.090 12.608 10.373 8.960 7.477 3.801 3.530						
	15.44 15.25 13.88 13.85 13.81 13.78 14.97 14.95 14.94 13.78 13.77						
	0.031 0.038 0.058 0.065 0.075 0.089 0.109 0.126 0.152 0.302 0.326						
	5.455 5.545 5.736 5.787 5.854 5.938 6.033 6.104 6.194 6.572 6.618						
	6.00 6.25 6.50 6.75 7.00 7.25 7.50 7.75 8.00 8.25 8.50						
	3.182 2.787 1.850 1.754 1.635 1.510 1.347 1.300 1.258 1.226 1.206						
	13.77 13.76 17.03 17.02 17.02 17.02 13.66 13.66 13.66 13.66 13.66						
	0.362 0.414 0.630 0.665 0.714 0.775 0.873 0.905 0.936 0.960 0.977						
	6.686 6.780 7.155 7.221 7.320 7.458 7.754 7.895 8.081 8.298 8.549						
	8.75 9.00 9.25 9.50 9.75 10.00 (Ref.) : 2 , 4						
	1.195 1.188 1.184 1.181 1.180 1.179						
	13.67 13.69 13.72 15.99 16.09 16.28						
	0.987 0.992 0.996 0.998 0.999 0.999						
	8.782 9.025 9.272 9.529 9.778 10.029						
140	HYDROIODIC ACID -1 79.6	-2.000	HI 127.91	pHL RE t Z pHS	3.00 0.993 18.01 1.000 2.998	3.25 0.993 16.22 1.000 3.248	3.50 0.993 15.21 1.000 3.499
	3.75 4.00 4.00 4.25 4.50 4.75 4.75 5.00 5.25 5.50 5.75						
	0.993 0.993 0.993 0.993 0.993 0.993 0.993 0.993 0.993 0.993 0.993						
	14.65 14.33 13.34 13.15 13.05 13.00 14.05 14.02 14.00 13.03 13.03						
	1.000 1.000 1.000 1.000 1.000 1.000 1.000 1.000 1.000 1.000 1.000						
	3.749 3.999 3.999 4.249 4.499 4.749 4.749 4.999 5.249 5.499 5.749						
	6.00 6.25 6.50 6.75 7.00 7.25 7.50 7.75 8.00 8.25 8.50						
	0.993 0.993 0.993 0.993 0.993 0.993 0.993 0.993 0.993 0.993 0.993						
	13.02 13.03 15.83 15.83 15.83 15.84 13.00 13.01 13.01 13.02 13.01						
	1.000 1.000 1.000 1.000 1.000 1.000 1.000 1.000 1.000 1.000 1.000						
	5.999 6.249 6.499 6.749 6.999 7.249 7.499 7.749 7.999 8.249 8.499						
	8.75 9.00 9.25 9.50 9.75 10.00 (Ref.) : 1 , 12						
	0.993 0.993 0.993 0.993 0.993 0.993 0.993 0.993 0.993 0.993 0.993						
	13.02 13.04 13.07 14.96 15.04 15.18						
	1.000 1.000 1.000 1.000 1.000 1.000						
	8.749 8.999 9.249 9.499 9.749 9.999						
141	2-HYDROXY-3-CHLOROBUTYRIC -1 34.4	3.092	A. C4H7ClO3 138.55	pHL RE t Z pHS	3.00 3.217 26.31 0.728 3.487	3.25 2.958 24.12 0.794 3.643	3.50 2.772 22.66 0.848 3.805
	3.75 4.00 4.00 4.25 4.50 4.75 4.75 5.00 5.25 5.50 5.75						
	2.631 2.528 2.529 2.472 2.432 2.406 2.402 2.387 2.377 2.385 2.381						
	21.63 20.96 18.49 18.15 17.93 17.79 20.31 20.24 20.19 17.88 17.87						
	0.896 0.933 0.937 0.959 0.975 0.986 0.986 0.992 0.996 0.997 0.999						
	3.989 4.198 4.230 4.428 4.651 4.889 4.916 5.158 5.404 5.637 5.883						
	6.00 6.25 6.50 6.75 7.00 7.25 7.50 7.75 8.00 8.25 8.50						
	2.378 2.375 2.367 2.366 2.364 2.362 2.379 2.378 2.376 2.375 2.378						
	17.86 17.86 24.59 24.58 24.58 24.57 17.82 17.83 17.83 17.84 17.86						
	0.999 1.000 1.000 1.000 1.000 1.000 1.000 1.000 1.000 1.000 1.000						
	6.131 6.381 6.683 6.932 7.182 7.432 7.631 7.881 8.131 8.381 8.632						
	8.75 9.00 9.25 9.50 9.75 10.00 (Ref.) : 3						
	2.376 2.374 2.372 2.365 2.362 2.358						
	17.89 17.95 18.06 22.80 23.24 24.08						
	1.000 1.000 1.000 1.000 1.000 1.000						
	8.883 9.133 9.385 9.682 9.936 10.195						

142	2-HYDROXY-3- -1	Cl- 34.4	ISOBUTYRIC 3.208	A.	C4H7ClO3 138.55	pHL RE t Z pHS	3.00 3.394 26.11 0.689 3.522	3.25 3.090 24.02 0.759 3.672	3.50 2.870 22.61 0.818 3.827
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50
2.700	2.574	2.569	2.499	2.449	2.416	2.411	2.392	2.380	2.387
21.62	20.95	18.47	18.15	17.93	17.79	20.31	20.24	20.19	17.88
0.872	0.916	0.922	0.949	0.968	0.981	0.983	0.990	0.994	0.997
4.004	4.208	4.245	4.437	4.656	4.891	4.919	5.159	5.405	5.638
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25
2.379	2.376	2.367	2.366	2.364	2.362	2.379	2.378	2.377	2.375
17.86	17.86	24.59	24.58	24.58	24.57	17.82	17.83	17.83	17.84
0.999	0.999	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
6.132	6.381	6.683	6.932	7.182	7.432	7.631	7.881	8.131	8.381
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3			
2.376	2.374	2.372	2.365	2.362	2.358				
17.89	17.95	18.06	22.80	23.24	24.08				
1.000	1.000	1.000	1.000	1.000	1.000				
8.883	9.133	9.385	9.682	9.936	10.195				
143	2-HYDROXY-3- -1	Cl- 36.5	PROPIONIC 3.124	A.	C3H5ClO3 124.52	pHL RE t Z pHS	3.00 3.093 25.24 0.712 3.484	3.25 2.830 23.21 0.780 3.638	3.50 2.642 21.85 0.837 3.798
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50
2.497	2.392	2.390	2.332	2.291	2.264	2.261	2.245	2.235	2.241
20.90	20.26	17.94	17.63	17.42	17.29	19.66	19.59	19.55	17.37
0.887	0.927	0.932	0.956	0.973	0.984	0.985	0.991	0.995	0.997
3.980	4.188	4.222	4.419	4.641	4.878	4.904	5.145	5.391	5.625
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25
2.235	2.232	2.225	2.224	2.222	2.221	2.235	2.234	2.233	2.231
17.35	17.35	23.68	23.67	23.67	23.66	17.32	17.32	17.33	17.34
0.999	0.999	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
6.119	6.369	6.667	6.916	7.166	7.416	7.619	7.869	8.119	8.369
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3			
2.233	2.231	2.229	2.223	2.221	2.218				
17.38	17.43	17.52	21.96	22.34	23.06				
1.000	1.000	1.000	1.000	1.000	1.000				
8.870	9.121	9.372	9.665	9.919	10.176				
144	2-HYDROXY-m- -1	TOLUIC 31.3	ACID 3.000		C8H8O3 152.15	pHL RE t Z pHS	3.00 3.376 28.25 0.767 3.485	3.25 3.138 25.74 0.827 3.645	3.50 2.969 24.08 0.875 3.812
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50
2.842	2.751	2.758	2.708	2.673	2.650	2.645	2.631	2.622	2.633
22.92	22.16	19.45	19.07	18.82	18.67	21.45	21.36	21.32	18.77
0.916	0.947	0.950	0.968	0.981	0.989	0.989	0.994	0.997	0.998
4.001	4.214	4.239	4.442	4.668	4.907	4.937	5.179	5.426	5.656
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25
2.627	2.624	2.613	2.611	2.609	2.607	2.629	2.627	2.626	2.623
18.74	18.74	26.17	26.16	26.16	26.15	18.70	18.71	18.71	18.73
0.999	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
6.151	6.401	6.709	6.958	7.208	7.458	7.651	7.901	8.151	8.401
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3			
2.625	2.622	2.619	2.610	2.607	2.601				
18.79	18.86	18.98	24.28	24.83	25.91				
1.000	1.000	1.000	1.000	1.000	1.000				
8.903	9.154	9.406	9.710	9.966	10.226				

CHROMATOGRAPHIC DATA (1983)

D59

145	3-HYDROXY-o-TOLUIC -1	31.3	ACID 2.983		C8H8O3 152.15		pHL RE t Z pHS	3.00 3.354 28.28 0.772 3.480	3.25 3.121 25.76 0.831 3.642	3.50 2.957 24.08 0.879 3.809
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
2.834	2.746	2.754	2.705	2.671	2.648	2.643	2.630	2.621	2.633	2.630
22.92	22.16	19.45	19.07	18.82	18.67	21.45	21.36	21.32	18.77	18.75
0.918	0.949	0.951	0.969	0.981	0.989	0.990	0.994	0.997	0.998	0.999
3.999	4.213	4.238	4.441	4.668	4.907	4.936	5.179	5.426	5.656	5.903
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.627	2.623	2.613	2.611	2.609	2.607	2.629	2.627	2.626	2.623	2.627
18.74	18.74	26.17	26.16	26.16	26.15	18.70	18.71	18.71	18.73	18.75
0.999	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
6.151	6.401	6.709	6.958	7.208	7.458	7.651	7.901	8.151	8.401	8.653
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3				
2.625	2.622	2.619	2.610	2.607	2.601					
18.79	18.86	18.98	24.28	24.83	25.91					
1.000	1.000	1.000	1.000	1.000	1.000					
8.903	9.154	9.406	9.710	9.966	10.226					
146	3-HYDROXY-p-TOLUIC -1	31.3	ACID 3.143		C8H8O3 152.15		pHL RE t Z pHS	3.00 3.585 27.99 0.721 3.523	3.25 3.293 25.61 0.786 3.676	3.50 3.082 24.02 0.842 3.835
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
2.920	2.802	2.804	2.738	2.691	2.660	2.655	2.637	2.625	2.635	2.631
22.90	22.15	19.43	19.06	18.82	18.67	21.45	21.36	21.31	18.77	18.75
0.890	0.929	0.934	0.957	0.974	0.985	0.986	0.992	0.995	0.997	0.998
4.017	4.224	4.255	4.451	4.673	4.910	4.940	5.181	5.427	5.657	5.903
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.627	2.624	2.613	2.611	2.609	2.607	2.629	2.627	2.626	2.623	2.627
18.74	18.74	26.17	26.16	26.16	26.15	18.70	18.71	18.71	18.73	18.75
0.999	0.999	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
6.151	6.401	6.709	6.958	7.208	7.458	7.651	7.901	8.151	8.401	8.653
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3				
2.625	2.622	2.619	2.610	2.607	2.601					
18.79	18.86	18.98	24.28	24.83	25.91					
1.000	1.000	1.000	1.000	1.000	1.000					
8.903	9.154	9.406	9.710	9.966	10.226					
147	4-HYDROXY-m-TOLUIC -1	31.3	ACID 3.066		C8H8O3 152.15		pHL RE t Z pHS	3.00 3.467 28.13 0.746 3.502	3.25 3.205 25.68 0.809 3.659	3.50 3.018 24.05 0.861 3.822
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
2.875	2.773	2.778	2.721	2.681	2.654	2.649	2.633	2.623	2.634	2.630
22.91	22.16	19.44	19.06	18.82	18.67	21.45	21.36	21.32	18.77	18.75
0.905	0.939	0.943	0.963	0.978	0.987	0.988	0.993	0.996	0.998	0.999
4.008	4.218	4.246	4.446	4.670	4.908	4.938	5.180	5.426	5.656	5.903
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.627	2.624	2.613	2.611	2.609	2.607	2.629	2.627	2.626	2.623	2.627
18.74	18.74	26.17	26.16	26.16	26.15	18.70	18.71	18.71	18.73	18.75
0.999	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
6.151	6.401	6.709	6.958	7.208	7.458	7.651	7.901	8.151	8.401	8.653
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3				
2.625	2.622	2.619	2.610	2.607	2.601					
18.79	18.86	18.98	24.28	24.83	25.91					
1.000	1.000	1.000	1.000	1.000	1.000					
8.903	9.154	9.406	9.710	9.966	10.226					

148	6-HYDROXY-o-TOLUIC -1 31.3	ACID 3.321	C8H8O3 152.15	pHL RE t Z pHS	3.00 3.908 27.65 0.659 3.577	3.25 3.537 25.43 0.730 3.721	3.50 3.265 23.94 0.793 3.871			
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
3.051	2.888	2.879	2.790	2.724	2.680	2.672	2.647	2.631	2.639	2.633
22.87	22.14	19.40	19.05	18.81	18.67	21.45	21.36	21.31	18.77	18.75
0.851	0.900	0.908	0.938	0.962	0.977	0.979	0.988	0.993	0.996	0.998
4.042	4.240	4.280	4.466	4.681	4.914	4.945	5.183	5.428	5.659	5.904
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.628	2.625	2.613	2.611	2.609	2.607	2.629	2.627	2.626	2.623	2.627
18.74	18.74	26.17	26.16	26.16	26.15	18.70	18.71	18.71	18.73	18.75
0.999	0.999	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
6.152	6.401	6.709	6.958	7.208	7.458	7.651	7.901	8.151	8.401	8.653
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3				
2.625	2.622	2.619	2.610	2.607	2.601					
18.79	18.86	18.98	24.28	24.83	25.91					
1.000	1.000	1.000	1.000	1.000	1.000					
8.903	9.154	9.406	9.710	9.966	10.226					
149	m-HYDROXYBENZOIC -1 -2 34.0 62.0	ACID 4.082 9.609	C7H6O3 138.12	pHL RE t Z pHS	3.00 6.004 24.86 0.388 3.858	3.25 5.154 23.38 0.453 3.973	3.50 4.516 22.36 0.518 4.086			
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
3.971	3.503	3.303	3.068	2.856	2.689	2.661	2.559	2.491	2.462	2.439
21.59	21.03	18.40	18.19	18.02	17.90	20.43	20.37	20.33	17.99	17.98
0.592	0.673	0.719	0.776	0.835	0.889	0.898	0.934	0.960	0.977	0.986
4.213	4.364	4.457	4.587	4.752	4.948	4.991	5.198	5.426	5.671	5.899
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.423	2.413	2.398	2.393	2.386	2.377	2.385	2.365	2.332	2.280	2.191
17.97	17.98	24.81	24.83	24.87	24.94	18.19	18.39	18.73	19.29	20.29
0.992	0.996	0.999	1.001	1.004	1.008	1.014	1.025	1.043	1.075	1.135
6.140	6.386	6.691	6.938	7.187	7.436	7.648	7.897	8.146	8.395	8.680
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3 , 2				
2.084	1.955	1.821	1.674	1.583	1.513					
21.47	22.89	24.42	31.18	32.51	33.81					
1.213	1.317	1.443	1.600	1.715	1.812					
8.914	9.145	9.374	9.646	9.866	10.098					
150	p-HYDROXYBENZOIC -1 -2 34.0 62.0	ACID 4.530 9.310	C7H6O3 138.12	pHL RE t Z pHS	3.00 8.759 24.36 0.263 4.062	3.25 7.385 23.07 0.313 4.166	3.50 6.353 22.19 0.365 4.265			
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
5.455	4.658	4.135	3.765	3.401	3.081	2.999	2.797	2.644	2.544	2.492
21.50	20.99	18.32	18.15	18.01	17.90	20.41	20.36	20.32	17.99	17.98
0.426	0.501	0.570	0.628	0.697	0.772	0.793	0.852	0.903	0.945	0.965
4.375	4.505	4.622	4.725	4.858	5.024	5.078	5.254	5.459	5.721	5.923
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.455	2.430	2.404	2.393	2.381	2.364	2.361	2.324	2.266	2.181	2.046
17.98	17.99	24.84	24.88	24.96	25.11	18.46	18.84	19.44	20.35	21.85
0.979	0.989	0.997	1.002	1.008	1.016	1.028	1.049	1.085	1.140	1.243
6.151	6.391	6.700	6.942	7.189	7.438	7.661	7.909	8.155	8.400	8.689
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3 , 2				
1.921	1.793	1.677	1.567	1.505	1.462					
23.22	24.65	25.98	32.39	33.38	34.37					
1.350	1.475	1.606	1.740	1.828	1.893					
8.908	9.129	9.355	9.620	9.847	10.085					

CHROMATOGRAPHIC DATA (1983)

D61

151	2-HYDROXYBUTYRIC -1	34.3	ACID 3.979		C4H8O3 104.11		pHL RE t Z pHS	3.00 5.505 24.86 0.420 3.813	3.25 4.751 23.33 0.488 3.931	3.50 4.185 22.29 0.556 4.047
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
3.704	3.296	3.142	2.934	2.752	2.611	2.589	2.505	2.450	2.429	2.410
21.50	20.93	18.34	18.12	17.94	17.82	20.33	20.27	20.22	17.91	17.90
0.630	0.710	0.750	0.805	0.860	0.908	0.915	0.946	0.968	0.981	0.989
4.179	4.336	4.423	4.560	4.732	4.935	4.975	5.188	5.420	5.663	5.894
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.398	2.390	2.378	2.375	2.372	2.370	2.387	2.385	2.384	2.382	2.385
17.89	17.88	24.63	24.63	24.62	24.62	17.85	17.85	17.86	17.87	17.89
0.994	0.996	0.998	0.999	0.999	1.000	1.000	1.000	1.000	1.000	1.000
6.136	6.383	6.686	6.934	7.183	7.433	7.632	7.881	8.131	8.381	8.633
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 11 , 3				
2.383	2.381	2.379	2.372	2.369	2.366					
17.92	17.98	18.08	22.85	23.29	24.13					
1.000	1.000	1.000	1.000	1.000	1.000					
8.883	9.134	9.385	9.683	9.937	10.196					
152	3-HYDROXYBUTYRIC -1	34.3	ACID 4.519		C4H8O3 104.11		pHL RE t Z pHS	3.00 8.608 24.23 0.266 4.056	3.25 7.260 22.95 0.316 4.160	3.50 6.247 22.07 0.368 4.260
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
5.367	4.585	4.074	3.711	3.354	3.041	2.962	2.765	2.615	2.518	2.468
21.39	20.88	18.24	18.07	17.93	17.82	20.31	20.26	20.22	17.91	17.90
0.430	0.505	0.573	0.631	0.700	0.775	0.795	0.854	0.904	0.945	0.965
4.370	4.500	4.617	4.720	4.854	5.020	5.074	5.250	5.456	5.718	5.920
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.433	2.410	2.388	2.381	2.376	2.372	2.388	2.386	2.384	2.382	2.385
17.89	17.88	24.63	24.63	24.62	24.62	17.85	17.85	17.86	17.87	17.89
0.979	0.988	0.994	0.997	0.998	0.999	0.999	1.000	1.000	1.000	1.000
6.148	6.389	6.694	6.937	7.184	7.433	7.632	7.882	8.131	8.381	8.633
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 11 , 3				
2.383	2.381	2.379	2.372	2.369	2.366					
17.92	17.98	18.08	22.85	23.29	24.13					
1.000	1.000	1.000	1.000	1.000	1.000					
8.883	9.134	9.385	9.683	9.937	10.196					
153	4-HYDROXYBUTYRIC -1	34.3	ACID 4.721		C4H8O3 104.11		pHL RE t Z pHS	3.00 10.360 24.06 0.220 4.151	3.25 8.687 22.85 0.263 4.252	3.50 7.430 22.01 0.308 4.347
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
6.334	5.350	4.614	4.173	3.729	3.325	3.202	2.943	2.737	2.581	2.512
21.36	20.87	18.21	18.06	17.93	17.82	20.30	20.25	20.22	17.91	17.90
0.362	0.431	0.504	0.559	0.627	0.706	0.734	0.800	0.862	0.921	0.947
4.452	4.574	4.700	4.794	4.916	5.068	5.128	5.289	5.481	5.752	5.939
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.460	2.426	2.396	2.385	2.378	2.373	2.389	2.387	2.385	2.382	2.385
17.89	17.89	24.63	24.63	24.62	24.62	17.85	17.85	17.86	17.87	17.89
0.968	0.981	0.990	0.994	0.997	0.998	0.999	0.999	1.000	1.000	1.000
6.157	6.393	6.700	6.939	7.185	7.434	7.633	7.882	8.131	8.381	8.633
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 11 , 3				
2.384	2.381	2.379	2.372	2.369	2.366					
17.92	17.98	18.08	22.85	23.29	24.13					
1.000	1.000	1.000	1.000	1.000	1.000					
8.883	9.134	9.385	9.683	9.937	10.196					

154	m-HYDROXYCINNAMIC -1	27.0	ACID 4.397		C9H8O3 164.16		pHL RE t Z pHS	3.00 9.479 28.86 0.309 4.027	3.25 8.045 27.02 0.366 4.135	3.50 6.961 25.77 0.424 4.240
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
6.025	5.203	4.795	4.386	4.001	3.680	3.606	3.407	3.265	3.195	3.146
24.83	24.14	20.80	20.56	20.36	20.22	23.43	23.36	23.31	20.35	20.34
0.492	0.572	0.627	0.688	0.757	0.825	0.841	0.892	0.932	0.960	0.975
4.358	4.496	4.593	4.709	4.857	5.037	5.088	5.281	5.498	5.744	5.960
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
3.112	3.091	3.062	3.055	3.050	3.045	3.076	3.074	3.071	3.067	3.072
20.32	20.32	28.99	28.99	28.98	28.97	20.28	20.28	20.29	20.31	20.34
0.986	0.992	0.996	0.998	0.999	0.999	1.000	1.000	1.000	1.000	1.000
6.196	6.439	6.758	7.004	7.252	7.501	7.686	7.935	8.185	8.435	8.688
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 11 , 1				
3.069	3.066	3.061	3.048	3.042	3.034					
20.39	20.49	20.66	26.97	27.80	29.48					
1.000	1.000	1.000	1.000	1.000	1.000					
8.938	9.189	9.442	9.758	10.016	10.282					
155	o-HYDROXYCINNAMIC -1	27.0	ACID 4.613		C9H8O3 164.16		pHL RE t Z pHS	3.00 11.491 28.60 0.254 4.125	3.25 9.674 26.85 0.302 4.229	3.50 8.302 25.66 0.354 4.329
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
7.110	6.052	5.420	4.911	4.416	3.983	3.867	3.592	3.386	3.261	3.189
24.76	24.10	20.74	20.52	20.34	20.21	23.41	23.35	23.30	20.35	20.33
0.415	0.489	0.552	0.611	0.683	0.759	0.782	0.844	0.897	0.940	0.961
4.440	4.569	4.676	4.780	4.914	5.079	5.136	5.313	5.518	5.772	5.974
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
3.139	3.106	3.070	3.060	3.052	3.046	3.077	3.074	3.071	3.067	3.072
20.32	20.32	28.99	28.99	28.98	28.97	20.28	20.28	20.29	20.31	20.34
0.977	0.987	0.993	0.996	0.998	0.999	0.999	1.000	1.000	1.000	1.000
6.202	6.442	6.762	7.005	7.252	7.501	7.686	7.935	8.185	8.435	8.688
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 11 , 1				
3.069	3.066	3.061	3.048	3.042	3.034					
20.39	20.49	20.66	26.97	27.80	29.48					
1.000	1.000	1.000	1.000	1.000	1.000					
8.938	9.189	9.442	9.758	10.016	10.282					
156	p-HYDROXYCINNAMIC -1	27.0	ACID 4.678		C9H8O3 164.16		pHL RE t Z pHS	3.00 12.202 28.53 0.239 4.156	3.25 10.251 26.80 0.285 4.259	3.50 8.777 25.63 0.334 4.357
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
7.496	6.356	5.642	5.100	4.567	4.096	3.963	3.663	3.434	3.286	3.206
24.74	24.09	20.72	20.52	20.34	20.21	23.41	23.34	23.30	20.35	20.33
0.393	0.465	0.530	0.588	0.659	0.737	0.762	0.827	0.884	0.932	0.956
4.465	4.592	4.702	4.803	4.934	5.094	5.153	5.324	5.526	5.782	5.979
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
3.149	3.112	3.073	3.061	3.053	3.047	3.077	3.074	3.071	3.067	3.072
20.32	20.32	28.99	28.99	28.98	28.97	20.28	20.28	20.29	20.31	20.34
0.973	0.984	0.992	0.996	0.998	0.999	0.999	0.999	1.000	1.000	1.000
6.205	6.444	6.763	7.006	7.253	7.501	7.686	7.935	8.185	8.435	8.688
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 11 , 7				
3.069	3.066	3.061	3.048	3.042	3.034					
20.39	20.49	20.66	26.97	27.80	29.48					
1.000	1.000	1.000	1.000	1.000	1.000					
8.938	9.189	9.442	9.758	10.016	10.282					

CHROMATOGRAPHIC DATA (1983)

D63

157	2-HYDROXYISOBUTYRIC -1	33.5	ACID (S) 3.971	C4H8O3 104.11			pHL RE t Z pHS	3.00 5.585 25.27 0.424 3.812	3.25 4.824 23.69 0.493 3.931	3.50 4.252 22.62 0.561 4.048
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
3.767	3.356	3.206	2.996	2.812	2.671	2.648	2.564	2.509	2.489	2.470
21.81	21.22	18.57	18.34	18.16	18.03	20.61	20.54	20.49	18.12	18.11
0.635	0.715	0.754	0.809	0.863	0.910	0.917	0.948	0.969	0.982	0.989
4.180	4.338	4.423	4.561	4.735	4.939	4.978	5.193	5.425	5.666	5.898
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.458	2.450	2.437	2.434	2.432	2.429	2.447	2.446	2.444	2.442	2.445
18.10	18.10	25.02	25.01	25.01	25.00	18.06	18.06	18.07	18.08	18.10
0.994	0.997	0.998	0.999	0.999	1.000	1.000	1.000	1.000	1.000	1.000
6.141	6.388	6.693	6.940	7.190	7.439	7.637	7.886	8.136	8.386	8.638
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 11 , 4				
2.443	2.441	2.438	2.432	2.428	2.424					
18.13	18.20	18.30	23.20	23.67	24.56					
1.000	1.000	1.000	1.000	1.000	1.000					
8.888	9.139	9.390	9.690	9.944	10.203					
158	p-HYDROXYPHENYLACETIC -1	26.9	ACID 3.171	C8H8O3 152.15			pHL RE t Z pHS	3.00 4.162 31.17 0.727 3.566	3.25 3.835 28.40 0.791 3.718	3.50 3.597 26.55 0.845 3.876
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
3.414	3.279	3.287	3.212	3.158	3.122	3.113	3.092	3.078	3.094	3.089
25.23	24.36	21.17	20.74	20.45	20.27	23.54	23.44	23.38	20.40	20.38
0.892	0.930	0.935	0.957	0.974	0.985	0.986	0.992	0.995	0.997	0.998
4.056	4.263	4.292	4.487	4.708	4.944	4.979	5.219	5.465	5.692	5.938
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
3.084	3.080	3.063	3.061	3.058	3.055	3.087	3.085	3.082	3.079	3.084
20.37	20.36	29.07	29.06	29.06	29.05	20.32	20.33	20.34	20.35	20.38
0.999	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
6.186	6.435	6.753	7.002	7.252	7.502	7.686	7.936	8.186	8.436	8.688
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 11 , 2				
3.081	3.078	3.073	3.060	3.054	3.045					
20.44	20.53	20.71	27.05	27.88	29.58					
1.000	1.000	1.000	1.000	1.000	1.000					
8.939	9.190	9.443	9.759	10.018	10.284					
159	HYDROXYTRIMETHYLACETIC -1	31.3	ACID 4.863	C5H10O3 118.13			pHL RE t Z pHS	3.00 12.834 25.50 0.194 4.227	3.25 10.722 24.16 0.233 4.327	3.50 9.134 23.23 0.274 4.420
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
7.749	6.503	5.543	4.986	4.422	3.902	3.727	3.393	3.119	2.898	2.806
22.52	21.98	19.06	18.91	18.77	18.66	21.38	21.33	21.30	18.76	18.75
0.324	0.388	0.460	0.513	0.580	0.660	0.692	0.762	0.831	0.903	0.933
4.522	4.641	4.767	4.857	4.974	5.120	5.183	5.336	5.521	5.794	5.971
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.735	2.687	2.643	2.629	2.619	2.612	2.632	2.630	2.627	2.624	2.627
18.74	18.74	26.17	26.16	26.16	26.15	18.70	18.71	18.71	18.73	18.75
0.958	0.975	0.988	0.993	0.996	0.998	0.999	0.999	1.000	1.000	1.000
6.184	6.416	6.729	6.966	7.211	7.459	7.653	7.902	8.151	8.401	8.653
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3				
2.625	2.622	2.619	2.611	2.607	2.601					
18.79	18.86	18.98	24.28	24.83	25.91					
1.000	1.000	1.000	1.000	1.000	1.000					
8.903	9.154	9.406	9.710	9.966	10.226					

160	2-HYDROXYVALERIC -1	ACID 32.3	3.886	C5H10O3 118.13	pHL RE t Z pHS	3.00 5.414 26.05 0.455 3.782	3.25 4.702 24.34 0.526 3.903	3.50 4.168 23.18 0.595 4.024		
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
3.717	3.340	3.222	3.026	2.858	2.733	2.714	2.640	2.592	2.579	2.562
22.31	21.69	18.95	18.69	18.50	18.36	21.04	20.97	20.93	18.46	18.45
0.669	0.748	0.781	0.833	0.883	0.925	0.930	0.957	0.974	0.985	0.991
4.161	4.325	4.403	4.548	4.728	4.937	4.976	5.195	5.430	5.669	5.904
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.551	2.544	2.531	2.528	2.526	2.523	2.543	2.542	2.540	2.537	2.541
18.44	18.44	25.62	25.62	25.61	25.61	18.40	18.40	18.41	18.42	18.44
0.995	0.997	0.999	0.999	1.000	1.000	1.000	1.000	1.000	1.000	1.000
6.148	6.395	6.702	6.950	7.199	7.449	7.644	7.894	8.144	8.394	8.646
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3				
2.539	2.537	2.534	2.526	2.523	2.518					
18.48	18.54	18.66	23.77	24.28	25.27					
1.000	1.000	1.000	1.000	1.000	1.000					
8.896	9.147	9.399	9.701	9.956	10.215					
161	4-HYDROXYVALERIC -1	ACID 32.3	4.699	C5H10O3 118.13	pHL RE t Z pHS	3.00 10.690 25.08 0.227 4.146	3.25 8.968 23.75 0.271 4.248	3.50 7.674 22.83 0.317 4.344		
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
6.547	5.538	4.819	4.359	3.899	3.484	3.362	3.097	2.888	2.737	2.667
22.13	21.60	18.78	18.62	18.47	18.36	21.01	20.95	20.92	18.46	18.45
0.373	0.443	0.514	0.570	0.639	0.718	0.744	0.810	0.870	0.926	0.951
4.450	4.574	4.695	4.791	4.916	5.071	5.130	5.294	5.490	5.757	5.947
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.615	2.581	2.549	2.539	2.532	2.526	2.545	2.543	2.540	2.538	2.541
18.44	18.44	25.62	25.62	25.61	25.61	18.40	18.40	18.41	18.42	18.44
0.970	0.982	0.991	0.995	0.997	0.998	0.999	0.999	1.000	1.000	1.000
6.168	6.405	6.715	6.955	7.202	7.450	7.646	7.895	8.144	8.394	8.646
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3				
2.539	2.537	2.534	2.526	2.523	2.518					
18.48	18.54	18.66	23.77	24.28	25.27					
1.000	1.000	1.000	1.000	1.000	1.000					
8.896	9.147	9.399	9.701	9.956	10.215					
162	HYPOPHOSPHOROUS -1	ACID (S) 45.2	1.230	H3PO2 66.00	pHL RE t Z pHS	3.00 1.798 23.64 0.990 3.192	3.25 1.791 21.13 0.994 3.409	3.50 1.787 19.61 0.996 3.629		
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
1.784	1.782	1.787	1.786	1.785	1.784	1.783	1.782	1.781	1.786	1.786
18.68	18.12	16.32	16.01	15.83	15.73	17.62	17.57	17.54	15.79	15.78
0.998	0.999	0.999	0.999	1.000	1.000	1.000	1.000	1.000	1.000	1.000
3.860	4.100	4.104	4.339	4.582	4.829	4.845	5.093	5.343	5.580	5.829
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
1.785	1.784	1.781	1.780	1.780	1.779	1.786	1.786	1.785	1.784	1.785
15.78	15.78	20.83	20.83	20.83	20.83	15.75	15.75	15.76	15.76	15.77
1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
6.079	6.329	6.613	6.863	7.113	7.363	7.579	7.829	8.079	8.329	8.580
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 11 , 4				
1.785	1.784	1.783	1.780	1.779	1.777					
15.79	15.83	15.89	19.38	19.62	20.04					
1.000	1.000	1.000	1.000	1.000	1.000					
8.830	9.080	9.331	9.610	9.862	10.116					

CHROMATOGRAPHIC DATA (1983)

D65

163	IODIC ACID -1	42.0	0.770	H102 159.91		pHL RE t Z pHS	3.00 1.928 24.69 0.997 3.213	3.25 1.926 22.03 0.998 3.429	3.50 1.924 20.41 0.999 3.647	
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
1.923	1.922	1.928	1.928	1.927	1.925	1.924	1.923	1.922	1.929	1.928
19.41	18.81	16.86	16.53	16.34	16.22	18.27	18.21	18.17	16.29	16.28
0.999	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
3.876	4.116	4.119	4.352	4.595	4.842	4.861	5.108	5.358	5.593	5.842
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
1.927	1.926	1.922	1.921	1.920	1.919	1.929	1.928	1.927	1.926	1.928
16.28	16.28	21.74	21.74	21.73	21.73	16.24	16.25	16.25	16.26	16.27
1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
6.092	6.342	6.631	6.881	7.130	7.380	7.592	7.842	8.092	8.342	8.593
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 1 , 4				
1.927	1.926	1.924	1.921	1.920	1.917					
16.29	16.33	16.41	20.20	20.47	20.98					
1.000	1.000	1.000	1.000	1.000	1.000					
8.843	9.094	9.344	9.628	9.881	10.136					
164	IODOACETIC -1	ACID 42.1	3.174	C2H3102 185.95		pHL RE t Z pHS	3.00 2.782 22.95 0.682 3.471	3.25 2.524 21.25 0.753 3.623	3.50 2.338 20.11 0.814 3.780	
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
2.195	2.089	2.081	2.024	1.982	1.956	1.953	1.938	1.928	1.930	1.927
19.31	18.77	16.76	16.49	16.32	16.21	18.25	18.19	18.15	16.28	16.27
0.869	0.914	0.921	0.948	0.968	0.981	0.982	0.990	0.994	0.997	0.998
3.959	4.163	4.204	4.397	4.617	4.853	4.875	5.115	5.360	5.599	5.844
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
1.924	1.922	1.918	1.917	1.916	1.915	1.924	1.923	1.923	1.921	1.923
16.26	16.26	21.71	21.70	21.70	21.70	16.23	16.23	16.24	16.25	16.25
0.999	0.999	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
6.092	6.342	6.631	6.880	7.130	7.380	7.591	7.841	8.091	8.342	8.593
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 1				
1.922	1.921	1.920	1.916	1.915	1.913					
16.28	16.32	16.39	20.17	20.44	20.95					
1.000	1.000	1.000	1.000	1.000	1.000					
8.843	9.093	9.344	9.628	9.880	10.135					
165	m-IODOBENZOIC ACID -1	33.4	3.851	C7H5102 248.02		pHL RE t Z pHS	3.00 5.129 25.50 0.465 3.762	3.25 4.461 23.85 0.536 3.885	3.50 3.960 22.72 0.605 4.007	
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
3.539	3.188	3.080	2.898	2.742	2.627	2.610	2.542	2.499	2.487	2.472
21.87	21.27	18.63	18.38	18.19	18.06	20.64	20.57	20.53	18.15	18.14
0.679	0.757	0.789	0.840	0.889	0.929	0.934	0.959	0.976	0.986	0.992
4.146	4.311	4.388	4.535	4.717	4.928	4.965	5.186	5.422	5.661	5.896
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.462	2.455	2.444	2.441	2.439	2.436	2.455	2.453	2.452	2.450	2.453
18.13	18.12	25.06	25.06	25.05	25.05	18.09	18.09	18.10	18.11	18.13
0.995	0.997	0.999	0.999	1.000	1.000	1.000	1.000	1.000	1.000	1.000
6.141	6.388	6.693	6.941	7.190	7.440	7.637	7.887	8.137	8.387	8.639
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3 , 1				
2.451	2.449	2.446	2.439	2.436	2.432					
18.16	18.22	18.33	23.25	23.72	24.62					
1.000	1.000	1.000	1.000	1.000	1.000					
8.889	9.140	9.391	9.691	9.945	10.204					

166	o-iodobenzoic acid -1 33.4			2.863	C7H5IO2 248.02			pH RE t Z pHS	3.00 3.023 27.24 0.802 3.436	3.25 2.835 24.78 0.856 3.603	3.50 2.703 23.16 0.899 3.778
	3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
	2.607	2.540	2.548	2.510	2.485	2.468	2.464	2.454	2.447	2.458	2.455
	22.04	21.32	18.80	18.43	18.20	18.06	20.66	20.58	20.53	18.15	18.14
	0.933	0.959	0.961	0.975	0.985	0.992	0.992	0.995	0.997	0.998	0.999
	3.973	4.191	4.213	4.421	4.651	4.892	4.919	5.163	5.410	5.641	5.888
	6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
	2.452	2.450	2.441	2.440	2.438	2.436	2.454	2.453	2.451	2.450	2.453
	18.13	18.12	25.06	25.06	25.05	25.05	18.09	18.09	18.10	18.11	18.13
	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
	6.137	6.387	6.691	6.940	7.190	7.440	7.637	7.887	8.137	8.387	8.639
	8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3 , 1				
	2.451	2.449	2.446	2.439	2.436	2.432					
	18.16	18.22	18.33	23.25	23.72	24.62					
	1.000	1.000	1.000	1.000	1.000	1.000					
	8.889	9.140	9.391	9.691	9.945	10.204					
167	p-iodobenzoic acid -1 33.4			3.930	C7H5IO2 248.02			pH RE t Z pHS	3.00 5.432 25.38 0.438 3.795	3.25 4.702 23.78 0.507 3.915	3.50 4.155 22.68 0.576 4.034
	3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
	3.692	3.301	3.166	2.966	2.792	2.660	2.640	2.561	2.510	2.493	2.476
	21.85	21.26	18.61	18.37	18.18	18.06	20.64	20.57	20.53	18.15	18.14
	0.650	0.730	0.766	0.820	0.872	0.917	0.923	0.952	0.971	0.983	0.990
	4.169	4.329	4.411	4.552	4.729	4.935	4.974	5.191	5.424	5.665	5.898
	6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
	2.464	2.457	2.444	2.441	2.439	2.436	2.455	2.453	2.452	2.450	2.453
	18.13	18.12	25.06	25.06	25.05	25.05	18.09	18.09	18.10	18.11	18.13
	0.994	0.997	0.998	0.999	0.999	1.000	1.000	1.000	1.000	1.000	1.000
	6.141	6.389	6.693	6.941	7.190	7.440	7.637	7.887	8.137	8.387	8.639
	8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3 , 1				
	2.451	2.449	2.446	2.439	2.436	2.432					
	18.16	18.22	18.33	23.25	23.72	24.62					
	1.000	1.000	1.000	1.000	1.000	1.000					
	8.889	9.140	9.391	9.691	9.945	10.204					
168	4-iodobutyric acid -1 34.4			4.638	C4H7IO2 214.00			pH RE t Z pHS	3.00 9.566 24.08 0.238 4.111	3.25 8.039 22.85 0.283 4.214	3.50 6.892 22.00 0.332 4.310
	3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
	5.893	5.000	4.364	3.958	3.553	3.189	3.086	2.855	2.674	2.545	2.484
	21.34	20.84	18.19	18.04	17.90	17.80	20.28	20.22	20.19	17.88	17.87
	0.389	0.460	0.532	0.588	0.658	0.735	0.760	0.823	0.881	0.932	0.955
	4.417	4.542	4.665	4.763	4.889	5.047	5.104	5.271	5.469	5.737	5.930
	6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
	2.440	2.411	2.385	2.376	2.370	2.365	2.381	2.379	2.377	2.375	2.378
	17.86	17.86	24.59	24.58	24.58	24.57	17.82	17.83	17.83	17.84	17.86
	0.973	0.984	0.992	0.995	0.997	0.999	0.999	0.999	1.000	1.000	1.000
	6.153	6.391	6.696	6.938	7.184	7.433	7.632	7.881	8.131	8.381	8.632
	8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3				
	2.376	2.374	2.372	2.365	2.362	2.358					
	17.89	17.95	18.06	22.80	23.24	24.08					
	1.000	1.000	1.000	1.000	1.000	1.000					
	8.883	9.133	9.385	9.682	9.936	10.195					

CHROMATOGRAPHIC DATA (1983)

D67

169	m-iodophenylacetic -1	34.3	ACID 4.159	C8H7IO2 262.05		pHL RE t Z pHS	3.00 6.334 24.62 0.364 3.891	3.25 5.418 23.19 0.426 4.004	3.50 4.729 22.21 0.490 4.114	
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
4.138	3.627	3.386	3.133	2.901	2.713	2.679	2.565	2.486	2.450	2.423
21.46	20.92	18.30	18.10	17.94	17.82	20.33	20.26	20.22	17.91	17.90
0.562	0.643	0.694	0.752	0.814	0.872	0.883	0.923	0.953	0.973	0.984
4.238	4.384	4.482	4.607	4.766	4.957	5.001	5.203	5.428	5.676	5.900
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.405	2.394	2.380	2.376	2.373	2.370	2.387	2.386	2.384	2.382	2.385
17.89	17.88	24.63	24.63	24.63	24.62	17.85	17.85	17.86	17.87	17.89
0.991	0.995	0.997	0.998	0.999	1.000	1.000	1.000	1.000	1.000	1.000
6.139	6.384	6.688	6.935	7.183	7.433	7.632	7.881	8.131	8.381	8.633
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 2 , 1				
2.383	2.381	2.379	2.372	2.369	2.366					
17.92	17.98	18.08	22.85	23.29	24.13					
1.000	1.000	1.000	1.000	1.000	1.000					
8.883	9.134	9.385	9.683	9.937	10.196					
170	o-iodophenylacetic -1	34.3	ACID 4.038	C8H7IO2 262.05		pHL RE t Z pHS	3.00 5.758 24.77 0.401 3.838	3.25 4.954 23.28 0.467 3.954	3.50 4.350 22.26 0.534 4.069	
3.75	4.00	4.00	4.25	4.50	4.75	5.00	5.25	5.50	5.75	
3.835	3.395	3.216	2.994	2.796	2.641	2.616	2.522	2.460	2.414	
21.49	20.93	18.33	18.12	17.94	17.82	20.33	20.27	20.22	17.91	
0.608	0.689	0.733	0.789	0.846	0.897	0.905	0.940	0.964	0.987	
4.198	4.351	4.442	4.575	4.742	4.942	4.983	5.193	5.422	5.896	
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.400	2.391	2.378	2.375	2.373	2.370	2.387	2.386	2.384	2.382	2.385
17.89	17.88	24.63	24.63	24.63	24.62	17.85	17.85	17.86	17.87	17.89
0.993	0.996	0.998	0.999	0.999	1.000	1.000	1.000	1.000	1.000	1.000
6.137	6.384	6.687	6.934	7.183	7.433	7.632	7.881	8.131	8.381	8.633
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 2 , 1				
2.383	2.381	2.379	2.372	2.369	2.366					
17.92	17.98	18.08	22.85	23.29	24.13					
1.000	1.000	1.000	1.000	1.000	1.000					
8.883	9.134	9.385	9.683	9.937	10.196					
171	p-iodophenylacetic -1	34.3	ACID 4.178	C8H7IO2 262.05		pHL RE t Z pHS	3.00 6.432 24.59 0.358 3.900	3.25 5.497 23.17 0.420 4.012	3.50 4.793 22.20 0.483 4.121	
3.75	4.00	4.00	4.25	4.50	4.75	5.00	5.25	5.50	5.75	
4.190	3.666	3.415	3.157	2.919	2.726	2.690	2.572	2.491	2.425	
21.46	20.92	18.30	18.10	17.94	17.82	20.33	20.26	20.22	17.91	
0.555	0.636	0.688	0.746	0.809	0.868	0.879	0.921	0.951	0.972	
4.244	4.390	4.489	4.612	4.770	4.959	5.004	5.205	5.429	5.677	
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.406	2.395	2.380	2.376	2.373	2.370	2.387	2.386	2.384	2.382	2.385
17.89	17.88	24.63	24.63	24.63	24.62	17.85	17.85	17.86	17.87	17.89
0.990	0.994	0.997	0.998	0.999	0.999	1.000	1.000	1.000	1.000	1.000
6.139	6.385	6.688	6.935	7.183	7.433	7.632	7.881	8.131	8.381	8.633
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 2 , 1				
2.383	2.381	2.379	2.372	2.369	2.366					
17.92	17.98	18.08	22.85	23.29	24.13					
1.000	1.000	1.000	1.000	1.000	1.000					
8.883	9.134	9.385	9.683	9.937	10.196					

172	3-iodopropionic -1	ACID 36.5	4.056	C3H5IO2 199.98	pHL RE t Z pHS	3.00 5.531 23.74 0.391 3.839	3.25 4.750 22.38 0.457 3.954	3.50 4.164 21.44 0.523 4.067		
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
3.665	3.236	3.048	2.834	2.642	2.490	2.465	2.373	2.311	2.284	2.263
20.73	20.21	17.76	17.57	17.41	17.29	19.65	19.58	19.55	17.37	17.36
0.596	0.677	0.724	0.780	0.839	0.891	0.900	0.936	0.961	0.978	0.987
4.194	4.345	4.441	4.571	4.736	4.933	4.974	5.181	5.410	5.656	5.884
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.249	2.241	2.229	2.226	2.224	2.221	2.236	2.235	2.233	2.231	2.234
17.36	17.35	23.68	23.67	23.67	23.66	17.32	17.32	17.33	17.34	17.35
0.992	0.996	0.998	0.999	0.999	1.000	1.000	1.000	1.000	1.000	1.000
6.125	6.371	6.671	6.918	7.167	7.416	7.619	7.869	8.119	8.369	8.620
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3				
2.233	2.231	2.229	2.223	2.221	2.218					
17.38	17.43	17.52	21.96	22.34	23.06					
1.000	1.000	1.000	1.000	1.000	1.000					
8.870	9.121	9.372	9.665	9.919	10.176					
173	5-iodovaleric -1	ACID 32.3	4.770	C5H9IO2 228.03	pHL RE t Z pHS	3.00 11.429 25.02 0.212 4.180	3.25 9.570 23.71 0.253 4.280	3.50 8.173 22.81 0.297 4.375		
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
6.955	5.862	5.048	4.556	4.059	3.607	3.465	3.175	2.943	2.765	2.686
22.12	21.59	18.77	18.61	18.47	18.36	21.00	20.95	20.92	18.46	18.45
0.351	0.418	0.490	0.544	0.613	0.692	0.721	0.789	0.853	0.916	0.943
4.479	4.601	4.725	4.818	4.939	5.090	5.150	5.309	5.500	5.771	5.955
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.627	2.588	2.553	2.541	2.533	2.527	2.546	2.543	2.541	2.538	2.541
18.44	18.44	25.62	25.62	25.61	25.61	18.40	18.40	18.41	18.42	18.44
0.965	0.979	0.990	0.994	0.997	0.998	0.999	0.999	1.000	1.000	1.000
6.172	6.407	6.717	6.956	7.202	7.450	7.646	7.895	8.144	8.394	8.646
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3				
2.539	2.537	2.534	2.526	2.523	2.518					
18.48	18.54	18.66	23.77	24.28	25.27					
1.000	1.000	1.000	1.000	1.000	1.000					
8.896	9.147	9.399	9.701	9.956	10.215					
174	isobutyric -1	ACID 34.4	4.860	C4H8O2 88.11	pHL RE t Z pHS	3.00 11.795 23.92 0.192 4.217	3.25 9.857 22.74 0.230 4.316	3.50 8.402 21.93 0.271 4.409		
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
7.130	5.984	5.054	4.552	4.040	3.564	3.401	3.095	2.842	2.629	2.545
21.30	20.82	18.16	18.02	17.90	17.80	20.27	20.22	20.19	17.88	17.87
0.320	0.383	0.458	0.509	0.576	0.655	0.687	0.757	0.826	0.901	0.931
4.510	4.628	4.759	4.847	4.962	5.106	5.169	5.320	5.502	5.781	5.955
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.479	2.435	2.396	2.383	2.374	2.367	2.382	2.380	2.378	2.375	2.378
17.86	17.86	24.59	24.58	24.58	24.57	17.82	17.83	17.83	17.84	17.86
0.957	0.974	0.987	0.992	0.996	0.998	0.998	0.999	1.000	1.000	1.000
6.166	6.397	6.705	6.941	7.186	7.433	7.633	7.882	8.131	8.381	8.632
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3 , 7				
2.376	2.374	2.372	2.365	2.362	2.358					
17.89	17.95	18.06	22.80	23.24	24.08					
1.000	1.000	1.000	1.000	1.000	1.000					
8.883	9.133	9.385	9.682	9.936	10.195					

CHROMATOGRAPHIC DATA (1983)

D69

175	ISOCAPROIC -1	ACID 30.3	4.845		C6H12O2 116.16		pHL RE t Z pHS	3.00 12.968 26.10 0.199 4.222	3.25 10.838 24.69 0.238 4.322	3.50 9.236 23.72 0.281 4.416	
	3.75 7.839 22.97 0.332 4.519	4.00 6.585 22.41 0.397 4.639	4.00 5.646 19.40 0.467 4.761	4.25 5.080 19.24 0.521 4.853	4.50 4.508 19.09 0.589 4.972	4.75 3.984 18.98 0.669 5.119	4.75 3.812 21.80 0.700 5.182	5.00 3.475 21.75 0.770 5.337	5.25 3.202 21.71 0.838 5.524	5.50 2.987 19.09 0.906 5.795	5.75 2.895 19.08 0.936 5.975
	6.00 2.824 19.07 0.960 6.189	6.25 2.777 19.06 0.976 6.422	6.50 2.733 26.75 0.988 6.737	6.75 2.719 26.74 0.993 6.975	7.00 2.709 26.74 0.996 7.220	7.25 2.702 26.73 0.998 7.468	7.50 2.724 19.03 0.999 7.660	7.75 2.721 19.03 0.999 7.909	8.00 2.718 19.04 1.000 8.158	8.25 2.715 19.05 1.000 8.408	8.50 2.719 19.07 1.000 8.660
	8.75 2.717 19.12 1.000 8.911	9.00 2.714 19.19 1.000 9.162	9.25 2.710 19.32 1.000 9.413	9.50 2.701 24.83 1.000 9.720	9.75 2.697 25.43 1.000 9.976	10.00 2.691 26.62 1.000 10.238	(Ref.) : 3 , 1				
176	ISOCITRIC -1 -2 -3	ACID 29.0 55.0 74.0	3.250 4.720 6.380		C6H8O7 192.13		pHL RE t Z pHS	3.00 3.728 30.79 0.761 3.559	3.25 3.316 28.83 0.862 3.707	3.50 2.989 27.84 0.965 3.858	
	3.75 2.698 27.59 1.082 4.030	4.00 2.432 28.03 1.218 4.222	4.00 2.364 25.33 1.269 4.291	4.25 2.178 26.16 1.399 4.465	4.50 2.004 27.21 1.545 4.661	4.75 1.854 28.38 1.699 4.875	4.75 1.831 31.88 1.720 4.908	5.00 1.722 33.07 1.860 5.122	5.25 1.632 34.38 1.999 5.351	5.50 1.563 33.02 2.167 5.627	5.75 1.505 34.56 2.308 5.846
	6.00 1.449 36.24 2.461 6.076	6.25 1.400 37.83 2.610 6.313	6.50 1.344 49.70 2.756 6.597	6.75 1.319 50.64 2.841 6.824	7.00 1.300 51.28 2.903 7.063	7.25 1.287 51.66 2.943 7.308	7.50 1.309 41.30 2.967 7.547	7.75 1.305 41.43 2.981 7.794	8.00 1.301 41.51 2.989 8.042	8.25 1.298 41.56 2.994 8.292	8.50 1.300 41.61 2.997 8.542
	8.75 1.298 41.65 2.998 8.792	9.00 1.296 41.71 2.999 9.042	9.25 1.294 41.81 2.999 9.292	9.50 1.280 49.10 3.000 9.553	9.75 1.277 49.42 3.000 9.803	10.00 1.274 50.02 3.000 10.054	(Ref.) : 12 , 4				
177	ISOPHTHALIC -1 -2	ACID 28.0 53.0	3.720 4.600		C8H6O4 166.13		pHL RE t Z pHS	3.00 4.711 31.55 0.625 3.712	3.25 4.007 30.03 0.741 3.835	3.50 3.469 29.28 0.866 3.959	
	3.75 3.007 29.09 1.012 4.099	4.00 2.611 29.42 1.183 4.263	4.00 2.481 26.59 1.265 4.342	4.25 2.261 27.16 1.405 4.490	4.50 2.070 27.86 1.555 4.670	4.75 1.923 28.56 1.694 4.877	4.75 1.902 32.09 1.711 4.908	5.00 1.810 32.63 1.811 5.123	5.25 1.749 33.03 1.884 5.354	5.50 1.727 30.16 1.933 5.604	5.75 1.706 30.30 1.960 5.836
	6.00 1.692 30.39 1.977 6.079	6.25 1.683 30.44 1.987 6.326	6.50 1.667 39.85 1.993 6.607	6.75 1.663 39.85 1.996 6.854	7.00 1.660 39.84 1.998 7.103	7.25 1.656 39.82 1.999 7.352	7.50 1.680 30.47 1.999 7.574	7.75 1.678 30.48 2.000 7.823	8.00 1.677 30.48 2.000 8.073	8.25 1.675 30.49 2.000 8.323	8.50 1.678 30.51 2.000 8.574
	8.75 1.676 30.54 2.000 8.824	9.00 1.674 30.60 2.000 9.074	9.25 1.672 30.70 2.000 9.325	9.50 1.661 37.15 2.000 9.600	9.75 1.658 37.52 2.000 9.851	10.00 1.654 38.23 2.000 10.104	(Ref.) : 12 , 7				

D70

CHROMATOGRAPHIC DATA (1983)

178	o-ISOPROPYLBENZOIC -1			ACID 24.7	3.635	C10H12O2 164.20		pHL RE t Z pHS	3.00 5.753 32.26 0.570 3.730	3.25 5.118 29.66 0.643 3.862	3.50 4.642 27.90 0.711 3.997
	3.75 4.251 26.61 0.779 4.152	4.00 3.939 25.73 0.844 4.335	4.00 3.903 22.18 0.858 4.384	4.25 3.731 21.78 0.899 4.553	4.50 3.596 21.49 0.934 4.754	4.75 3.502 21.30 0.960 4.979	4.75 3.483 24.87 0.963 5.018	5.00 3.429 24.77 0.978 5.251	5.25 3.395 24.70 0.987 5.493	5.50 3.404 21.45 0.992 5.720	5.75 3.392 21.43 0.996 5.961
	6.00 3.383 21.41 0.998 6.208	6.25 3.375 21.40 0.999 6.457	6.50 3.352 30.93 0.999 6.779	6.75 3.349 30.92 1.000 7.029	7.00 3.346 30.91 1.000 7.278	7.25 3.342 30.90 1.000 7.528	7.50 3.381 21.37 1.000 7.707	7.75 3.379 21.37 1.000 7.957	8.00 3.375 21.38 1.000 8.207	8.25 3.371 21.40 1.000 8.457	8.50 3.378 21.44 1.000 8.710
	8.75 3.374 21.50 1.000 8.961	9.00 3.370 21.62 1.000 9.212	9.25 3.364 21.83 1.000 9.465	9.50 3.346 28.87 1.000 9.790	9.75 3.339 29.94 1.000 10.050	10.00 3.328 32.19 1.000 10.320	(Ref.) : 11 , 1				
179	p-ISOPROPYLBENZOIC -1			ACID 24.7	4.354	C10H12O2 164.20		pHL RE t Z pHS	3.00 9.836 30.93 0.327 4.020	3.25 8.376 28.84 0.386 4.130	3.50 7.267 27.42 0.446 4.236
	3.75 6.310 26.36 0.516 4.357	4.00 5.476 25.59 0.598 4.499	4.00 5.108 21.94 0.647 4.588	4.25 4.683 21.67 0.708 4.709	4.50 4.289 21.44 0.776 4.862	4.75 3.966 21.29 0.842 5.047	4.75 3.893 24.82 0.857 5.098	5.00 3.694 24.74 0.904 5.296	5.25 3.556 24.69 0.940 5.518	5.50 3.496 21.44 0.965 5.757	5.75 3.448 21.42 0.979 5.978
	6.00 3.415 21.41 0.987 6.216	6.25 3.394 21.40 0.993 6.460	6.50 3.361 30.93 0.997 6.784	6.75 3.354 30.92 0.998 7.030	7.00 3.349 30.91 0.999 7.279	7.25 3.343 30.90 0.999 7.528	7.50 3.382 21.37 1.000 7.707	7.75 3.379 21.37 1.000 7.957	8.00 3.376 21.38 1.000 8.207	8.25 3.372 21.40 1.000 8.457	8.50 3.378 21.44 1.000 8.710
	8.75 3.374 21.50 1.000 8.961	9.00 3.370 21.62 1.000 9.212	9.25 3.364 21.83 1.000 9.465	9.50 3.346 28.87 1.000 9.790	9.75 3.339 29.94 1.000 10.050	10.00 3.328 32.19 1.000 10.320	(Ref.) : 11 , 1				
180	ISOPROPYLMALONIC -1 -2			ACID 26.0 52.0	2.943 5.879	C6H10O4 146.14		pHL RE t Z pHS	3.00 3.901 32.50 0.807 3.519	3.25 3.658 29.47 0.863 3.682	3.50 3.480 27.49 0.909 3.851
	3.75 3.337 26.19 0.950 4.043	4.00 3.218 25.48 0.987 4.257	4.00 3.226 22.21 0.992 4.287	4.25 3.127 22.08 1.025 4.492	4.50 3.012 22.33 1.067 4.718	4.75 2.866 22.97 1.125 4.952	4.75 2.831 26.57 1.138 4.998	5.00 2.654 27.60 1.221 5.225	5.25 2.455 28.84 1.328 5.450	5.50 2.211 27.43 1.505 5.757	5.75 2.079 28.29 1.610 5.938
	6.00 1.962 29.06 1.716 6.142	6.25 1.869 29.67 1.809 6.363	6.50 1.781 39.70 1.894 6.663	6.75 1.747 39.93 1.933 6.882	7.00 1.723 40.07 1.960 7.118	7.25 1.707 40.14 1.977 7.362	7.50 1.725 30.65 1.986 7.584	7.75 1.719 30.68 1.992 7.829	8.00 1.715 30.70 1.996 8.078	8.25 1.712 30.72 1.997 8.327	8.50 1.714 30.75 1.999 8.578
	8.75 1.712 30.79 1.999 8.828	9.00 1.709 30.84 2.000 9.078	9.25 1.707 30.95 2.000 9.328	9.50 1.696 37.54 2.000 9.604	9.75 1.692 37.93 2.000 9.856	10.00 1.689 38.67 2.000 10.109	(Ref.) : 12 , 2				

CHROMATOGRAPHIC DATA (1983)

D71

181	ISOVALERIC -1	ACID 32.1	4.780	C5H10O2 102.13		pHL RE t Z pHS	3.00 11.600 25.12 0.210 4.185	3.25 9.710 23.80 0.251 4.286	3.50 8.291 22.90 0.295 4.380	
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
7.053	5.942	5.112	4.612	4.107	3.647	3.501	3.206	2.969	2.787	2.707
22.20	21.67	18.83	18.67	18.53	18.42	21.08	21.03	20.99	18.52	18.51
0.348	0.415	0.487	0.541	0.610	0.689	0.718	0.787	0.851	0.915	0.942
4.484	4.605	4.729	4.823	4.943	5.093	5.154	5.313	5.503	5.773	5.957
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.646	2.606	2.570	2.558	2.549	2.544	2.562	2.560	2.557	2.555	2.558
18.50	18.49	25.73	25.72	25.72	25.71	18.46	18.46	18.47	18.48	18.50
0.964	0.979	0.990	0.994	0.997	0.998	0.999	0.999	1.000	1.000	1.000
6.174	6.408	6.719	6.958	7.204	7.452	7.647	7.896	8.146	8.395	8.647
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3 , 1				
2.556	2.554	2.551	2.543	2.539	2.534					
18.54	18.60	18.72	23.87	24.39	25.39					
1.000	1.000	1.000	1.000	1.000	1.000					
8.898	9.149	9.400	9.703	9.958	10.218					
182	ITACONIC -1 -2	ACID 28.0 56.0	3.815 5.449	C5H6O4 130.10		pHL RE t Z pHS	3.00 5.643 29.41 0.509 3.771	3.25 4.892 27.48 0.589 3.895	3.50 4.317 26.26 0.670 4.020	
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
3.820	3.385	3.230	2.959	2.690	2.434	2.377	2.170	1.989	1.827	1.747
25.52	25.25	22.33	22.63	23.28	24.26	27.71	28.80	29.89	28.12	28.66
0.761	0.864	0.915	1.003	1.111	1.237	1.267	1.397	1.536	1.702	1.788
4.161	4.329	4.412	4.563	4.745	4.947	4.996	5.195	5.406	5.691	5.881
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
1.684	1.640	1.600	1.584	1.574	1.567	1.585	1.582	1.580	1.578	1.580
29.09	29.39	38.42	38.51	38.56	38.58	29.78	29.79	29.81	29.82	29.84
1.861	1.914	1.954	1.972	1.984	1.991	1.995	1.997	1.998	1.999	1.999
6.097	6.331	6.616	6.851	7.094	7.341	7.567	7.815	8.065	8.314	8.565
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 12 , 2				
1.579	1.577	1.575	1.565	1.563	1.560					
29.87	29.92	30.01	36.07	36.40	37.03					
2.000	2.000	2.000	2.000	2.000	2.000					
8.815	9.065	9.315	9.587	9.838	10.091					
183	2-KETOGLUTARIC -1 -2	ACID 33.6 59.0	2.800 5.006	C5H6O5 146.10		pHL RE t Z pHS	3.00 2.869 27.86 0.849 3.418	3.25 2.675 25.63 0.916 3.590	3.50 2.523 24.40 0.980 3.768	
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
2.387	2.251	2.232	2.106	1.972	1.843	1.822	1.718	1.634	1.582	1.548
23.93	24.19	21.77	22.59	23.74	25.02	28.13	29.28	30.23	28.22	28.60
1.048	1.130	1.152	1.247	1.364	1.497	1.517	1.642	1.754	1.850	1.904
3.967	4.184	4.235	4.438	4.652	4.874	4.908	5.123	5.350	5.613	5.832
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
1.524	1.509	1.491	1.485	1.481	1.477	1.495	1.494	1.492	1.491	1.493
28.86	29.02	37.46	37.51	37.53	37.53	29.20	29.21	29.22	29.23	29.24
1.942	1.966	1.982	1.990	1.994	1.997	1.998	1.999	1.999	2.000	2.000
6.067	6.311	6.588	6.832	7.079	7.328	7.557	7.806	8.056	8.306	8.556
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 10				
1.492	1.490	1.488	1.480	1.478	1.475					
29.27	29.32	29.40	35.11	35.41	35.97					
2.000	2.000	2.000	2.000	2.000	2.000					
8.806	9.057	9.307	9.576	9.827	10.078					

184	LACTIC ACID -1	(S) 36.5	3.860	C3H6O3 90.08	pHL RE t Z pHS	3.00 4.778 24.01 0.455 3.754	3.25 4.146 22.53 0.526 3.876	3.50 3.673 21.53 0.595 3.996			
	3.75 3.276 20.77 0.669 4.134	4.00 2.943 20.22 0.747 4.297	4.00 2.830 17.80 0.782 4.379	4.25 2.659 17.58 0.834 4.524	4.50 2.513 17.41 0.884 4.703	4.75 2.403 17.29 0.925 4.913	4.75 2.388 19.65 0.930 4.949	5.00 2.323 19.59 0.957 5.167	5.25 2.281 19.55 0.974 5.402	5.50 2.267 17.37 0.985 5.644	5.75 2.253 17.36 0.991 5.879
	6.00 2.243 17.35 0.995 6.123	6.25 2.237 17.35 0.997 6.370	6.50 2.228 23.68 0.999 6.669	6.75 2.225 23.67 0.999 6.917	7.00 2.223 23.67 1.000 7.167	7.25 2.221 23.66 1.000 7.416	7.50 2.235 17.32 1.000 7.619	7.75 2.234 17.32 1.000 7.869	8.00 2.233 17.33 1.000 8.119	8.25 2.231 17.34 1.000 8.369	8.50 2.234 17.35 1.000 8.620
	8.75 2.233 17.38 1.000 8.870	9.00 2.231 17.43 1.000 9.121	9.25 2.229 17.52 1.000 9.372	9.50 2.223 21.96 1.000 9.665	9.75 2.221 22.34 1.000 9.919	10.00 2.218 23.06 1.000 10.176	(Ref.) : 3 , 1				
185	LAURYL SULFONIC -1	ACID 22.9	1.000	C12H26O3S 250.40	pHL RE t Z pHS	3.00 3.641 36.88 0.997 3.439	3.25 3.637 32.73 0.998 3.634	3.50 3.635 30.07 0.999 3.830			
	3.75 3.633 28.30 0.999 4.043	4.00 3.631 27.19 1.000 4.273	4.00 3.660 23.48 1.000 4.278	4.25 3.658 22.90 1.000 4.497	4.50 3.656 22.53 1.000 4.732	4.75 3.652 22.32 1.000 4.976	4.75 3.642 26.19 1.000 5.013	5.00 3.638 26.07 1.000 5.259	5.25 3.634 25.99 1.000 5.507	5.50 3.665 22.47 1.000 5.729	5.75 3.662 22.45 1.000 5.977
	6.00 3.658 22.43 1.000 6.226	6.25 3.654 22.42 1.000 6.475	6.50 3.629 32.74 1.000 6.802	6.75 3.626 32.73 1.000 7.052	7.00 3.623 32.71 1.000 7.302	7.25 3.619 32.70 1.000 7.551	7.50 3.665 22.38 1.000 7.726	7.75 3.663 22.39 1.000 7.976	8.00 3.659 22.40 1.000 8.226	8.25 3.655 22.42 1.000 8.476	8.50 3.662 22.47 1.000 8.730
	8.75 3.657 22.54 1.000 8.981	9.00 3.652 22.68 1.000 9.233	9.25 3.646 22.93 1.000 9.486	9.50 3.624 30.68 1.000 9.818	9.75 3.614 32.02 1.000 10.081	10.00 3.600 34.99 1.000 10.357	(Ref.) : 2 , 12				
186	LEVULINIC ACID -1	33.4	4.602	C5H8O3 116.12	pHL RE t Z pHS	3.00 9.495 24.60 0.247 4.097	3.25 7.988 23.30 0.294 4.200	3.50 6.855 22.40 0.344 4.298			
	3.75 5.870 21.71 0.403 4.406	4.00 4.991 21.20 0.476 4.533	4.00 4.391 18.47 0.546 4.652	4.25 3.987 18.31 0.603 4.752	4.50 3.586 18.17 0.672 4.882	4.75 3.229 18.06 0.749 5.043	4.75 3.132 20.62 0.772 5.099	5.00 2.906 20.56 0.835 5.270	5.25 2.732 20.52 0.889 5.471	5.50 2.612 18.15 0.937 5.735	5.75 2.553 18.14 0.959 5.932
	6.00 2.511 18.13 0.975 6.157	6.25 2.484 18.12 0.986 6.396	6.50 2.458 25.06 0.993 6.703	6.75 2.449 25.06 0.996 6.945	7.00 2.443 25.05 0.998 7.192	7.25 2.439 25.05 0.999 7.441	7.50 2.456 18.09 0.999 7.638	7.75 2.454 18.09 1.000 7.887	8.00 2.452 18.10 1.000 8.137	8.25 2.450 18.11 1.000 8.387	8.50 2.453 18.13 1.000 8.639
	8.75 2.451 18.16 1.000 8.889	9.00 2.449 18.22 1.000 9.140	9.25 2.446 18.33 1.000 9.391	9.50 2.439 23.25 1.000 9.691	9.75 2.436 23.72 1.000 9.945	10.00 2.432 24.62 1.000 10.204	(Ref.) : 10 , 3				

CHROMATOGRAPHIC DATA (1983)

D73

187	MALEIC ACID										
	-1	41.3	1.921		C4H4O4		pHL	3.00	3.25	3.50	
	-2	62.4	6.225		116.07		RE	2.037	2.005	1.984	
							t	24.74	22.20	20.64	
							Z	0.960	0.976	0.987	
							pHS	3.246	3.452	3.662	
	3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
	1.969	1.957	1.964	1.953	1.939	1.919	1.916	1.884	1.837	1.765	1.696
	19.69	19.17	17.16	16.96	16.99	17.26	19.45	20.07	21.07	20.73	22.21
	0.996	1.005	1.005	1.015	1.029	1.052	1.055	1.094	1.157	1.275	1.386
	3.886	4.123	4.132	4.362	4.604	4.851	4.875	5.121	5.368	5.671	5.887
	6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
	1.624	1.557	1.484	1.452	1.427	1.411	1.418	1.412	1.408	1.405	1.406
	23.77	25.22	34.13	34.95	35.54	35.91	28.34	28.46	28.53	28.58	28.61
	1.514	1.643	1.783	1.856	1.910	1.947	1.968	1.982	1.990	1.994	1.997
	6.108	6.336	6.637	6.850	7.081	7.323	7.557	7.801	8.049	8.298	8.548
	8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 10 , 1				
	1.405	1.403	1.401	1.394	1.392	1.390					
	28.65	28.70	28.78	34.15	34.42	34.91					
	1.998	1.999	1.999	2.000	2.000	2.000					
	8.798	9.048	9.298	9.564	9.815	10.066					
188	MALIC ACID										
	-1	32.6	3.460		C4H6O5		pHL	3.00	3.25	3.50	
	-2	59.0	5.050		134.09		RE	3.897	3.436	3.082	
							t	27.43	25.72	24.76	
							Z	0.640	0.730	0.821	
							pHS	3.610	3.748	3.889	
	3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
	2.777	2.508	2.430	2.247	2.068	1.906	1.878	1.755	1.659	1.595	1.557
	24.35	24.53	22.12	22.81	23.82	25.01	28.16	29.26	30.19	28.20	28.58
	0.921	1.034	1.081	1.188	1.317	1.459	1.482	1.613	1.732	1.838	1.895
	4.050	4.236	4.311	4.481	4.676	4.887	4.924	5.133	5.355	5.621	5.836
	6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
	1.530	1.512	1.492	1.486	1.481	1.478	1.495	1.494	1.492	1.491	1.493
	28.85	29.02	37.46	37.51	37.53	37.53	29.20	29.21	29.22	29.23	29.24
	1.937	1.963	1.980	1.988	1.993	1.996	1.998	1.999	1.999	2.000	2.000
	6.069	6.312	6.589	6.832	7.080	7.329	7.557	7.806	8.056	8.306	8.556
	8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 10 , 7				
	1.492	1.490	1.488	1.480	1.478	1.475					
	29.27	29.32	29.40	35.11	35.41	35.97					
	2.000	2.000	2.000	2.000	2.000	2.000					
	8.806	9.057	9.307	9.576	9.827	10.078					
189	MALONIC ACID (S)										
	-1	40.7	2.847		C3H4O4		pHL	3.00	3.25	3.50	
	-2	67.0	5.696		104.06		RE	2.489	2.317	2.195	
							t	24.10	22.12	20.87	
							Z	0.795	0.856	0.907	
							pHS	3.384	3.554	3.730	
	3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
	2.102	2.028	2.026	1.971	1.910	1.836	1.824	1.737	1.640	1.528	1.464
	20.11	19.78	17.73	17.83	18.29	19.16	21.57	22.89	24.44	24.18	25.23
	0.951	0.992	0.998	1.037	1.087	1.157	1.168	1.264	1.386	1.563	1.671
	3.928	4.148	4.182	4.394	4.625	4.864	4.897	5.129	5.362	5.664	5.860
	6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
	1.409	1.367	1.330	1.314	1.303	1.296	1.306	1.304	1.302	1.300	1.301
	26.13	26.81	34.51	34.78	34.96	35.05	27.83	27.87	27.89	27.91	27.93
	1.772	1.853	1.917	1.949	1.970	1.983	1.990	1.994	1.997	1.998	1.999
	6.076	6.308	6.588	6.816	7.058	7.304	7.541	7.788	8.038	8.287	8.538
	8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 10 , 1				
	1.300	1.299	1.298	1.292	1.291	1.289					
	27.95	28.00	28.07	33.02	33.25	33.68					
	1.999	2.000	2.000	2.000	2.000	2.000					
	8.787	9.037	9.288	9.549	9.800	10.051					

190	MANDELIC ACID -1	28.3	3.411	C8H8O3 152.15		pHL RE t Z pHS	3.00 4.483 29.55 0.637 3.626	3.25 4.041 27.15 0.709 3.767	3.50 3.715 25.53 0.773 3.912	
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
3.454	3.253	3.241	3.130	3.047	2.991	2.981	2.949	2.928	2.938	2.930
24.36	23.56	20.50	20.13	19.87	19.70	22.80	22.70	22.65	19.82	19.80
0.834	0.887	0.897	0.930	0.956	0.973	0.975	0.986	0.992	0.995	0.997
4.079	4.273	4.315	4.496	4.708	4.938	4.973	5.210	5.454	5.683	5.927
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.925	2.920	2.904	2.902	2.900	2.897	2.925	2.923	2.921	2.918	2.923
19.79	19.79	28.05	28.04	28.03	28.02	19.75	19.75	19.76	19.78	19.80
0.998	0.999	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
6.175	6.424	6.738	6.987	7.237	7.487	7.674	7.924	8.174	8.424	8.676
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 11 , 7				
2.920	2.917	2.913	2.901	2.896	2.889					
19.85	19.94	20.09	26.06	26.78	28.24					
1.000	1.000	1.000	1.000	1.000	1.000					
8.927	9.178	9.430	9.742	10.000	10.263					
191	MES (S) -1	28.0	6.095	C6H13NO4S 195.23		pHL RE t Z pHS	3.00 51.093 27.00 0.053 4.838	3.25 41.949 25.62 0.065 4.929	3.50 35.109 24.68 0.078 5.012	
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
29.119	23.671	17.940	15.770	13.471	11.209	9.832	8.414	7.069	4.950	4.581
23.94	23.39	20.11	19.98	19.86	19.77	22.74	22.70	22.68	19.91	19.90
0.094	0.116	0.155	0.177	0.207	0.250	0.286	0.335	0.401	0.584	0.632
5.101	5.201	5.343	5.410	5.495	5.598	5.678	5.776	5.897	6.212	6.300
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
4.177	3.799	3.323	3.200	3.099	3.028	3.021	2.993	2.975	2.963	2.963
19.90	19.90	28.21	28.21	28.22	28.22	19.87	19.87	19.88	19.89	19.92
0.695	0.767	0.878	0.913	0.944	0.966	0.978	0.987	0.993	0.996	0.998
6.422	6.580	6.921	7.083	7.285	7.511	7.711	7.940	8.182	8.429	8.681
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 11 , 9				
2.957	2.952	2.947	2.934	2.929	2.921	2-(N-Morpholino)ethane sulfonic acid				
19.97	20.06	20.22	26.26	27.01	28.51					
0.999	0.999	1.000	1.000	1.000	1.000					
8.930	9.181	9.433	9.746	10.003	10.268					
192	MESACONIC ACID -1 -2	28.0 56.0	3.098 4.750	C5H6O4 130.10		pHL RE t Z pHS	3.00 3.612 31.58 0.809 3.527	3.25 3.257 29.33 0.902 3.682	3.50 2.968 28.16 0.995 3.843	
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
2.704	2.453	2.392	2.206	2.031	1.884	1.861	1.759	1.684	1.647	1.620
27.76	28.05	25.26	25.94	26.76	27.57	30.94	31.56	32.03	29.40	29.57
1.099	1.220	1.264	1.380	1.510	1.640	1.657	1.762	1.847	1.910	1.945
4.022	4.221	4.288	4.466	4.663	4.876	4.909	5.123	5.351	5.607	5.833
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
1.602	1.590	1.573	1.569	1.565	1.562	1.582	1.581	1.579	1.577	1.580
29.68	29.75	38.63	38.64	38.63	38.62	29.80	29.81	29.81	29.82	29.84
1.967	1.981	1.990	1.994	1.997	1.998	1.999	1.999	2.000	2.000	2.000
6.072	6.318	6.596	6.842	7.091	7.340	7.565	7.814	8.064	8.314	8.565
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 12 , 2				
1.578	1.577	1.575	1.565	1.563	1.560					
29.87	29.92	30.02	36.07	36.40	37.03					
2.000	2.000	2.000	2.000	2.000	2.000					
8.815	9.065	9.315	9.587	9.838	10.091					

CHROMATOGRAPHIC DATA (1983)

D75

193	MESOTARTARIC ACID				C4H6O6		pH			
-1	32.4		3.222		150.09		RE	3.00	3.25	3.50
-2	60.6		4.815				t	3.392	3.029	2.743
							Z	28.30	26.49	25.57
							pHS	0.743	0.839	0.934
								3.531	3.680	3.835
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
2.491	2.261	2.201	2.035	1.878	1.743	1.723	1.626	1.555	1.517	1.490
25.32	25.71	23.32	24.10	25.09	26.09	29.16	29.97	30.60	28.35	28.58
1.040	1.162	1.208	1.325	1.459	1.595	1.614	1.728	1.822	1.895	1.935
4.009	4.205	4.275	4.453	4.651	4.865	4.898	5.110	5.338	5.598	5.822
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
1.472	1.461	1.446	1.442	1.438	1.435	1.452	1.451	1.449	1.448	1.450
28.74	28.83	37.00	37.02	37.02	37.02	28.91	28.92	28.93	28.94	28.95
1.961	1.978	1.988	1.993	1.996	1.998	1.999	1.999	2.000	2.000	2.000
6.060	6.306	6.579	6.825	7.073	7.322	7.552	7.802	8.052	8.302	8.552
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 12 , 7				
1.449	1.448	1.446	1.438	1.436	1.434					
28.98	29.02	29.11	34.64	34.93	35.45					
2.000	2.000	2.000	2.000	2.000	2.000					
8.802	9.052	9.303	9.570	9.821	10.072					
194	METHACRYLIC ACID				C4H6O2		pH			
-1	36.7		4.483		86.09		RE	3.00	3.25	3.50
							t	7.858	6.634	5.716
							Z	23.20	22.03	21.22
							pHS	0.272	0.323	0.375
								4.033	4.137	4.237
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
4.918	4.208	3.738	3.411	3.089	2.808	2.739	2.561	2.427	2.340	2.295
20.59	20.12	17.64	17.49	17.36	17.25	19.58	19.52	19.49	17.33	17.32
0.438	0.513	0.583	0.640	0.709	0.782	0.802	0.859	0.908	0.948	0.967
4.347	4.478	4.597	4.701	4.835	5.002	5.055	5.232	5.439	5.702	5.905
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.263	2.243	2.224	2.218	2.214	2.210	2.224	2.222	2.221	2.219	2.221
17.31	17.31	23.59	23.59	23.59	23.58	17.27	17.28	17.28	17.29	17.30
0.980	0.988	0.994	0.997	0.998	0.999	0.999	1.000	1.000	1.000	1.000
6.134	6.375	6.676	6.919	7.166	7.415	7.619	7.868	8.118	8.368	8.619
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 10				
2.220	2.218	2.216	2.211	2.208	2.205					
17.33	17.39	17.48	21.89	22.26	22.97					
1.000	1.000	1.000	1.000	1.000	1.000					
8.869	9.120	9.371	9.664	9.918	10.174					
195	METHOXYACETIC ACID				C3H6O3		pH			
-1	36.5		3.570		90.08		RE	3.00	3.25	3.50
							t	3.934	3.478	3.139
							Z	24.46	22.79	21.66
							pHS	0.556	0.630	0.700
								3.637	3.770	3.905
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
2.863	2.642	2.597	2.480	2.387	2.323	2.315	2.277	2.254	2.252	2.244
20.83	20.24	17.86	17.60	17.41	17.29	19.65	19.59	19.55	17.37	17.36
0.769	0.836	0.855	0.896	0.932	0.958	0.961	0.977	0.987	0.992	0.995
4.059	4.241	4.303	4.469	4.669	4.893	4.923	5.154	5.395	5.633	5.874
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.238	2.234	2.226	2.224	2.223	2.221	2.235	2.234	2.233	2.231	2.234
17.35	17.35	23.68	23.67	23.67	23.66	17.32	17.32	17.33	17.34	17.35
0.997	0.999	0.999	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
6.121	6.369	6.668	6.917	7.166	7.416	7.619	7.869	8.119	8.369	8.620
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3 , 7				
2.233	2.231	2.229	2.223	2.221	2.218					
17.38	17.43	17.52	21.96	22.34	23.06					
1.000	1.000	1.000	1.000	1.000	1.000					
8.870	9.121	9.372	9.665	9.919	10.176					

196	m-METHOXYBENZOIC -1	ACID 28.3	4.088	C8H8O3 152.15		pHL RE t Z pHS	3.00 7.060 28.31 0.399 3.885	3.25 6.077 26.42 0.465 4.001	3.50 5.335 25.13 0.531 4.115	
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
4.701	4.159	3.961	3.682	3.433	3.240	3.205	3.088	3.011	2.986	2.959
24.17	23.48	20.33	20.06	19.85	19.70	22.77	22.69	22.64	19.82	19.80
0.605	0.687	0.727	0.784	0.843	0.895	0.904	0.939	0.963	0.978	0.987
4.245	4.398	4.482	4.615	4.784	4.984	5.028	5.239	5.469	5.707	5.937
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.941	2.929	2.909	2.905	2.901	2.898	2.925	2.923	2.921	2.918	2.923
19.79	19.79	28.05	28.04	28.03	28.02	19.75	19.75	19.76	19.78	19.80
0.993	0.996	0.998	0.999	0.999	1.000	1.000	1.000	1.000	1.000	1.000
6.179	6.426	6.741	6.988	7.237	7.487	7.674	7.924	8.174	8.424	8.676
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 11 , 1				
2.920	2.917	2.913	2.901	2.896	2.889					
19.85	19.94	20.09	26.06	26.78	28.24					
1.000	1.000	1.000	1.000	1.000	1.000					
8.927	9.178	9.430	9.742	10.000	10.263					
197	o-METHOXYBENZOIC -1	ACID 28.3	4.094	C8H8O3 152.15		pHL RE t Z pHS	3.00 7.093 28.30 0.397 3.888	3.25 6.103 26.41 0.463 4.003	3.50 5.356 25.13 0.529 4.117	
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
4.718	4.173	3.971	3.690	3.439	3.244	3.208	3.091	3.013	2.986	2.959
24.17	23.48	20.32	20.06	19.85	19.70	22.77	22.69	22.64	19.82	19.80
0.603	0.684	0.725	0.783	0.842	0.894	0.903	0.938	0.963	0.978	0.987
4.246	4.400	4.483	4.617	4.785	4.984	5.029	5.240	5.470	5.708	5.938
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.941	2.929	2.909	2.905	2.901	2.898	2.925	2.923	2.921	2.918	2.923
19.79	19.79	28.05	28.04	28.03	28.02	19.75	19.75	19.76	19.78	19.80
0.993	0.996	0.998	0.999	0.999	1.000	1.000	1.000	1.000	1.000	1.000
6.179	6.426	6.741	6.988	7.237	7.487	7.674	7.924	8.174	8.424	8.676
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 11 , 1				
2.920	2.917	2.913	2.901	2.896	2.889					
19.85	19.94	20.09	26.06	26.78	28.24					
1.000	1.000	1.000	1.000	1.000	1.000					
8.927	9.178	9.430	9.742	10.000	10.263					
198	p-METHOXYBENZOIC -1	ACID 30.0	4.471	C8H8O3 152.15		pHL RE t Z pHS	3.00 9.253 26.64 0.284 4.048	3.25 7.823 25.07 0.337 4.154	3.50 6.745 24.01 0.392 4.256	
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
5.810	4.985	4.511	4.114	3.731	3.403	3.324	3.118	2.967	2.880	2.829
23.20	22.60	19.58	19.38	19.21	19.08	21.95	21.89	21.85	19.19	19.18
0.456	0.534	0.596	0.656	0.725	0.797	0.816	0.872	0.917	0.952	0.970
4.369	4.503	4.610	4.719	4.859	5.032	5.085	5.269	5.480	5.733	5.942
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.793	2.770	2.744	2.737	2.732	2.727	2.751	2.749	2.746	2.744	2.747
19.17	19.17	26.93	26.93	26.92	26.91	19.13	19.13	19.14	19.15	19.18
0.982	0.990	0.995	0.997	0.998	0.999	0.999	1.000	1.000	1.000	1.000
6.174	6.416	6.729	6.973	7.221	7.470	7.661	7.911	8.160	8.410	8.663
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 2 , 1				
2.745	2.742	2.739	2.729	2.725	2.719					
19.22	19.30	19.43	25.00	25.62	26.84					
1.000	1.000	1.000	1.000	1.000	1.000					
8.913	9.164	9.416	9.723	9.980	10.242					

199	p-METHOXYPHENYLACETIC ACID				C9H10O3			pHL	3.00	3.25	3.50
-1	29.7	4.361	166.18					RE	8.480	7.200	6.235
								t	26.96	25.33	24.21
								Z	0.314	0.371	0.429
								pHS	3.999	4.108	4.213
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75	
5.402	4.671	4.293	3.934	3.595	3.312	3.249	3.074	2.948	2.884	2.841	
23.37	22.75	19.71	19.50	19.32	19.18	22.10	22.03	21.98	19.30	19.28	
0.497	0.578	0.635	0.694	0.762	0.829	0.845	0.895	0.934	0.962	0.976	
4.330	4.469	4.570	4.685	4.833	5.013	5.064	5.256	5.473	5.721	5.938	
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50	
2.812	2.793	2.770	2.764	2.760	2.756	2.780	2.778	2.776	2.773	2.777	
19.27	19.27	27.12	27.11	27.11	27.10	19.23	19.24	19.24	19.26	19.28	
0.986	0.992	0.996	0.998	0.999	0.999	1.000	1.000	1.000	1.000	1.000	
6.173	6.417	6.730	6.976	7.224	7.473	7.663	7.913	8.163	8.413	8.665	
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 2 , 7					
2.774	2.772	2.768	2.758	2.753	2.747						
19.33	19.40	19.54	25.18	25.81	27.07						
1.000	1.000	1.000	1.000	1.000	1.000						
8.915	9.166	9.418	9.727	9.983	10.245						
200	METHYLETHYLMALONIC ACID				C6H10O4			pHL	3.00	3.25	3.50
-1	25.0	2.860	146.14					RE	3.954	3.743	3.593
-2	50.0	6.413						t	33.54	30.28	28.13
								Z	0.831	0.879	0.917
								pHS	3.514	3.680	3.853
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75	
3.479	3.393	3.411	3.352	3.292	3.221	3.201	3.103	2.965	2.745	2.566	
26.66	25.76	22.34	21.96	21.83	21.97	25.57	26.03	26.81	25.10	26.23	
0.949	0.974	0.976	0.994	1.013	1.036	1.041	1.076	1.129	1.240	1.336	
4.048	4.265	4.285	4.492	4.721	4.960	5.001	5.241	5.480	5.796	5.994	
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50	
2.380	2.207	2.001	1.925	1.864	1.821	1.825	1.809	1.799	1.791	1.792	
27.40	28.50	39.61	40.12	40.52	40.77	31.00	31.10	31.16	31.21	31.25	
1.451	1.576	1.743	1.819	1.882	1.928	1.956	1.974	1.985	1.992	1.995	
6.199	6.412	6.740	6.930	7.148	7.382	7.605	7.842	8.087	8.335	8.586	
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 2					
1.788	1.785	1.782	1.769	1.766	1.761						
31.30	31.36	31.48	38.37	38.80	39.61						
1.997	1.998	1.999	2.000	2.000	2.000						
8.835	9.085	9.335	9.614	9.866	10.119						
201	METHYLMALONIC ACID				C4H6O4			pHL	3.00	3.25	3.50
-1	29.3	3.072	118.09					RE	3.665	3.384	3.177
-2	58.5	5.787						t	29.59	27.03	25.36
								Z	0.757	0.822	0.877
								pHS	3.518	3.675	3.839
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75	
3.010	2.871	2.866	2.758	2.636	2.491	2.459	2.293	2.114	1.901	1.793	
24.28	23.72	20.84	20.82	21.18	21.92	25.13	26.22	27.48	26.38	27.17	
0.928	0.975	0.983	1.024	1.075	1.142	1.157	1.248	1.362	1.545	1.647	
4.024	4.234	4.273	4.474	4.696	4.929	4.974	5.197	5.421	5.730	5.911	
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50	
1.698	1.626	1.559	1.533	1.514	1.502	1.515	1.511	1.508	1.506	1.507	
27.87	28.41	37.24	37.43	37.55	37.62	29.25	29.28	29.29	29.31	29.33	
1.748	1.833	1.907	1.942	1.966	1.980	1.988	1.993	1.996	1.998	1.999	
6.116	6.339	6.632	6.852	7.089	7.334	7.563	7.810	8.058	8.308	8.558	
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 2 , 1					
1.506	1.504	1.502	1.494	1.491	1.489						
29.36	29.41	29.50	35.26	35.57	36.14						
1.999	2.000	2.000	2.000	2.000	2.000						
8.808	9.058	9.308	9.578	9.829	10.080						

202	METHYLSULFONIC ACID				CH403S		pHL	3.00	3.25	3.50
	-1	50.6	-2.000		96.10		RE	1.584	1.584	1.584
							t	22.29	19.92	18.52
							Z	1.000	1.000	1.000
							pHS	3.145	3.371	3.598
	3.75	4.00	4.00	4.25	4.50	4.75	5.00	5.25	5.50	5.75
	1.584	1.583	1.587	1.587	1.586	1.586	1.585	1.584	1.588	1.587
	17.67	17.17	15.57	15.29	15.13	15.04	16.72	16.68	16.65	15.10
	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
	3.834	4.077	4.079	4.317	4.562	4.810	4.823	5.072	5.321	5.561
	6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25
	1.587	1.586	1.584	1.584	1.583	1.583	1.588	1.587	1.587	1.587
	15.09	15.09	19.58	19.58	19.58	19.58	15.06	15.06	15.06	15.07
	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
	6.060	6.310	6.587	6.837	7.087	7.337	7.560	7.810	8.060	8.310
	8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 2 , 12			
	1.587	1.586	1.585	1.584	1.583	1.582				
	15.09	15.12	15.17	18.26	18.45	18.78				
	1.000	1.000	1.000	1.000	1.000	1.000				
	8.811	9.061	9.312	9.584	9.836	10.089				
203	2-NAPHTALENESULFONIC ACID				C10H803S		pHL	3.00	3.25	3.50
	-1	31.3	0.167		208.23		RE	2.614	2.613	2.613
							t	29.64	26.35	24.30
							Z	0.999	1.000	1.000
							pHS	3.318	3.523	3.730
	3.75	4.00	4.00	4.25	4.50	4.75	5.00	5.25	5.50	5.75
	2.612	2.611	2.626	2.625	2.624	2.621	2.618	2.616	2.613	2.627
	22.99	22.18	19.51	19.08	18.82	18.67	21.45	21.37	21.32	18.77
	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
	3.951	4.186	4.190	4.416	4.656	4.901	4.929	5.175	5.424	5.653
	6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25
	2.625	2.623	2.612	2.611	2.609	2.607	2.629	2.627	2.626	2.623
	18.74	18.74	26.17	26.16	26.16	26.15	18.70	18.71	18.71	18.73
	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
	6.151	6.400	6.708	6.958	7.208	7.458	7.651	7.901	8.151	8.401
	8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 10 , 2			
	2.625	2.622	2.619	2.610	2.607	2.601				
	18.79	18.86	18.98	24.28	24.83	25.91				
	1.000	1.000	1.000	1.000	1.000	1.000				
	8.903	9.154	9.406	9.710	9.966	10.226				
204	NICOTINIC ACID				C6H5N02		pHL	3.00	3.25	3.50
	-1	34.6	2.070		123.11		RE	11.277	9.448	8.068
		-34.6	4.819				t	24.32	23.01	22.10
							Z	0.200	0.239	0.280
							pHS	4.216	4.309	4.398
	3.75	4.00	4.00	4.25	4.50	4.75	5.00	5.25	5.50	5.75
	6.856	5.763	4.887	4.408	3.920	3.468	3.318	3.028	2.791	2.596
	21.38	20.84	18.16	18.01	17.87	17.76	20.21	20.16	20.12	17.83
	0.331	0.395	0.471	0.523	0.590	0.669	0.700	0.770	0.837	0.907
	4.497	4.614	4.743	4.832	4.948	5.094	5.156	5.309	5.495	5.771
	6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25
	2.456	2.415	2.379	2.367	2.359	2.353	2.368	2.365	2.363	2.361
	17.81	17.81	24.49	24.49	24.49	24.48	17.77	17.78	17.78	17.79
	0.960	0.976	0.988	0.993	0.996	0.998	0.999	0.999	1.000	1.000
	6.162	6.394	6.702	6.939	7.184	7.432	7.632	7.880	8.130	8.380
	8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 10			
	2.362	2.360	2.357	2.351	2.348	2.344				
	17.84	17.90	18.00	22.72	23.15	23.97				
	1.000	1.000	1.000	1.000	1.000	1.000				
	8.882	9.132	9.383	9.680	9.935	10.193				

205	NITRIC ACID -1	74.1	-1.370	HN03 63.01		pHL RE t Z pHS	3.00 1.069 18.57 1.000 3.020	3.25 1.069 16.69 1.000 3.267	3.50 1.069 15.63 1.000 3.513		
	3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
	1.069	1.069	1.069	1.069	1.069	1.069	1.069	1.069	1.069	1.069	1.069
	15.03	14.69	13.62	13.43	13.32	13.26	14.39	14.36	14.34	13.29	13.29
	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
	3.761	4.010	4.010	4.259	4.508	4.758	4.759	5.009	5.259	5.508	5.758
	6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
	1.069	1.069	1.069	1.069	1.069	1.069	1.069	1.069	1.069	1.069	1.069
	13.29	13.29	16.31	16.31	16.31	16.31	13.26	13.27	13.27	13.28	13.28
	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
	6.008	6.258	6.511	6.761	7.011	7.261	7.507	7.758	8.008	8.258	8.508
	8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 1 , 2				
	1.069	1.069	1.069	1.069	1.069	1.069					
	13.29	13.30	13.33	15.38	15.47	15.63					
	1.000	1.000	1.000	1.000	1.000	1.000					
	8.758	9.008	9.258	9.511	9.761	10.011					
206	2-NITRO-3-BROMOBENZOIC ACID -1	28.2	2.495	C7H4BrNO4 246.02		pHL RE t Z pHS	3.00 3.222 31.18 0.901 3.421	3.25 3.116 27.99 0.933 3.603	3.50 3.045 25.92 0.955 3.790		
	3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
	2.995	2.961	2.979	2.960	2.946	2.936	2.931	2.924	2.919	2.937	2.935
	24.53	23.65	20.66	20.20	19.92	19.74	22.85	22.76	22.70	19.86	19.84
	0.972	0.983	0.984	0.990	0.994	0.997	0.997	0.998	0.999	0.999	1.000
	3.996	4.222	4.234	4.450	4.684	4.927	4.958	5.203	5.451	5.677	5.925
	6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
	2.932	2.929	2.914	2.913	2.910	2.907	2.936	2.934	2.932	2.929	2.934
	19.83	19.83	28.12	28.11	28.10	28.09	19.79	19.79	19.80	19.82	19.84
	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
	6.175	6.424	6.738	6.988	7.238	7.488	7.674	7.924	8.174	8.425	8.677
	8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3				
	2.931	2.928	2.924	2.912	2.907	2.899					
	19.89	19.98	20.13	26.13	26.86	28.33					
	1.000	1.000	1.000	1.000	1.000	1.000					
	8.928	9.179	9.431	9.744	10.001	10.265					
207	3-NITRO-2-BROMOBENZOIC ACID -1	28.2	2.979	C7H4BrNO4 246.02		pHL RE t Z pHS	3.00 3.686 30.47 0.784 3.506	3.25 3.445 27.66 0.840 3.667	3.50 3.274 25.79 0.886 3.835		
	3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
	3.145	3.054	3.066	3.015	2.979	2.955	2.949	2.934	2.925	2.941	2.937
	24.49	23.64	20.62	20.19	19.91	19.74	22.85	22.76	22.70	19.86	19.84
	0.923	0.952	0.954	0.971	0.982	0.990	0.991	0.995	0.997	0.998	0.999
	4.025	4.239	4.261	4.464	4.691	4.930	4.963	5.205	5.452	5.679	5.926
	6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
	2.933	2.929	2.915	2.913	2.910	2.908	2.936	2.934	2.932	2.929	2.934
	19.83	19.83	28.12	28.11	28.10	28.09	19.79	19.79	19.80	19.82	19.84
	0.999	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
	6.175	6.424	6.739	6.988	7.238	7.488	7.674	7.924	8.174	8.425	8.677
	8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3				
	2.931	2.928	2.924	2.912	2.907	2.899					
	19.89	19.98	20.13	26.13	26.86	28.33					
	1.000	1.000	1.000	1.000	1.000	1.000					
	8.928	9.179	9.431	9.744	10.001	10.265					

D80

CHROMATOGRAPHIC DATA (1983)

208	3-NITRO-4-BROMOBENZOIC ACID -1	28.2	3.377	C7H4BrNO4 246.02	pHL RE t Z pHS	3.00 4.415 29.70 0.650 3.616	3.25 3.991 27.25 0.721 3.758	3.50 3.679 25.61 0.784 3.905		
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
3.431	3.241	3.232	3.127	3.049	2.997	2.987	2.957	2.937	2.948	2.941
24.42	23.62	20.55	20.17	19.91	19.74	22.85	22.75	22.70	19.86	19.84
0.843	0.894	0.903	0.934	0.959	0.975	0.977	0.987	0.992	0.995	0.997
4.074	4.270	4.310	4.493	4.706	4.938	4.973	5.210	5.454	5.683	5.928
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.935	2.931	2.915	2.913	2.911	2.908	2.936	2.934	2.932	2.929	2.934
19.83	19.83	28.12	28.11	28.10	28.09	19.79	19.79	19.80	19.82	19.84
0.999	0.999	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
6.175	6.425	6.739	6.988	7.238	7.488	7.675	7.924	8.174	8.425	8.677
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3				
2.931	2.928	2.924	2.912	2.907	2.899					
19.89	19.98	20.13	26.13	26.86	28.33					
1.000	1.000	1.000	1.000	1.000	1.000					
8.928	9.179	9.431	9.744	10.001	10.265					
209	2-NITRO-3-CHLOROBENZOIC ACID -1	31.3	2.398	C7H4ClNO4 201.57	pHL RE t Z pHS	3.00 2.857 29.10 0.912 3.377	3.25 2.771 26.14 0.941 3.563	3.50 2.714 24.23 0.961 3.755		
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
2.674	2.648	2.662	2.646	2.636	2.628	2.625	2.619	2.615	2.630	2.628
22.97	22.17	19.49	19.08	18.82	18.67	21.45	21.37	21.32	18.77	18.75
0.976	0.986	0.986	0.991	0.995	0.997	0.997	0.998	0.999	0.999	1.000
3.965	4.194	4.204	4.423	4.659	4.903	4.931	5.176	5.425	5.654	5.902
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.625	2.623	2.612	2.611	2.609	2.607	2.629	2.627	2.626	2.623	2.627
18.74	18.74	26.17	26.16	26.16	26.15	18.70	18.71	18.71	18.73	18.75
1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
6.151	6.401	6.708	6.958	7.208	7.458	7.651	7.901	8.151	8.401	8.653
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3				
2.625	2.622	2.619	2.610	2.607	2.601					
18.79	18.86	18.98	24.28	24.83	25.91					
1.000	1.000	1.000	1.000	1.000	1.000					
8.903	9.154	9.406	9.710	9.966	10.226					
210	3-NITRO-4-CHLOROBENZOIC ACID -1	31.3	3.357	C7H4ClNO4 201.57	pHL RE t Z pHS	3.00 3.983 27.58 0.646 3.588	3.25 3.595 25.40 0.718 3.731	3.50 3.309 23.92 0.782 3.879		
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
3.082	2.909	2.897	2.802	2.732	2.685	2.677	2.650	2.633	2.640	2.634
22.86	22.14	19.39	19.05	18.81	18.67	21.45	21.36	21.31	18.77	18.75
0.841	0.893	0.902	0.934	0.958	0.975	0.977	0.987	0.992	0.995	0.997
4.048	4.244	4.286	4.470	4.683	4.915	4.946	5.184	5.428	5.660	5.904
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.629	2.625	2.613	2.612	2.609	2.607	2.629	2.627	2.626	2.623	2.627
18.74	18.74	26.17	26.16	26.16	26.15	18.70	18.71	18.71	18.73	18.75
0.999	0.999	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
6.152	6.401	6.709	6.958	7.208	7.458	7.651	7.901	8.151	8.401	8.653
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3				
2.625	2.622	2.619	2.610	2.607	2.601					
18.79	18.86	18.98	24.28	24.83	25.91					
1.000	1.000	1.000	1.000	1.000	1.000					
8.903	9.154	9.406	9.710	9.966	10.226					

CHROMATOGRAPHIC DATA (1983)

D81

211	3-NITRO-5-CHLOROBENZOIC ACID					C7H4ClNO4	pHL	3.00	3.25	3.50
	-1	31.3	3.130	201.57			RE	3.564	3.277	3.070
							t	28.01	25.62	24.03
							Z	0.725	0.790	0.845
							pHS	3.519	3.673	3.833
	3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50
	2.912	2.796	2.799	2.735	2.689	2.659	2.654	2.636	2.625	2.635
	22.90	22.15	19.43	19.06	18.82	18.67	21.45	21.36	21.31	18.77
	0.893	0.931	0.935	0.958	0.974	0.985	0.986	0.992	0.995	0.997
	4.015	4.223	4.254	4.450	4.672	4.909	4.939	5.180	5.427	5.657
	6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25
	2.627	2.624	2.613	2.611	2.609	2.607	2.629	2.627	2.626	2.623
	18.74	18.74	26.17	26.16	26.16	26.15	18.70	18.71	18.71	18.73
	0.999	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
	6.151	6.401	6.709	6.958	7.208	7.458	7.651	7.901	8.151	8.401
	8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3			
	2.625	2.622	2.619	2.610	2.607	2.601				
	18.79	18.86	18.98	24.28	24.83	25.91				
	1.000	1.000	1.000	1.000	1.000	1.000				
	8.903	9.154	9.406	9.710	9.966	10.226				
212	m-NITROBENZOIC ACID					C7H5NO4	pHL	3.00	3.25	3.50
	-1	32.3	3.493	167.12			RE	4.182	3.730	3.395
							t	26.73	24.73	23.38
							Z	0.594	0.668	0.736
							pHS	3.629	3.766	3.906
	3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50
	3.125	2.913	2.884	2.769	2.680	2.620	2.611	2.576	2.554	2.550
	22.40	21.72	19.04	18.73	18.51	18.36	21.05	20.97	20.93	18.46
	0.802	0.862	0.876	0.913	0.944	0.966	0.969	0.981	0.989	0.994
	4.066	4.254	4.306	4.480	4.686	4.913	4.945	5.179	5.422	5.656
	6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25
	2.544	2.540	2.529	2.527	2.525	2.523	2.543	2.542	2.540	2.537
	18.44	18.44	25.62	25.62	25.61	25.61	18.40	18.40	18.41	18.42
	0.998	0.999	0.999	1.000	1.000	1.000	1.000	1.000	1.000	1.000
	6.146	6.394	6.700	6.950	7.199	7.449	7.644	7.894	8.144	8.394
	8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3 , 1			
	2.539	2.537	2.534	2.526	2.523	2.518				
	18.48	18.54	18.66	23.77	24.28	25.27				
	1.000	1.000	1.000	1.000	1.000	1.000				
	8.896	9.147	9.399	9.701	9.956	10.215				
213	o-NITROBENZOIC ACID					C7H5NO4	pHL	3.00	3.25	3.50
	-1	33.6	2.173	167.12			RE	2.575	2.521	2.486
							t	27.94	25.04	23.20
							Z	0.940	0.961	0.975
							pHS	3.332	3.526	3.725
	3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50
	2.461	2.446	2.457	2.448	2.442	2.437	2.434	2.430	2.427	2.439
	22.00	21.26	18.78	18.39	18.14	18.00	20.59	20.51	20.46	18.10
	0.985	0.991	0.991	0.995	0.997	0.998	0.998	0.999	0.999	1.000
	3.941	4.173	4.181	4.404	4.642	4.887	4.912	5.159	5.407	5.638
	6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25
	2.436	2.434	2.425	2.424	2.423	2.421	2.439	2.438	2.436	2.437
	18.07	18.07	24.97	24.96	24.96	24.95	18.03	18.04	18.04	18.06
	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
	6.136	6.385	6.689	6.938	7.188	7.438	7.636	7.886	8.136	8.386
	8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 2 , 1			
	2.436	2.433	2.431	2.424	2.421	2.417				
	18.11	18.17	18.28	23.16	23.62	24.51				
	1.000	1.000	1.000	1.000	1.000	1.000				
	8.888	9.138	9.390	9.689	9.944	10.202				

214	p-NITROBENZOIC ACID				C7H5NO4		pHL		3.00	3.25	3.50
	-1	32.3	3.523		167.12		RE	4.257	3.788	3.441	
							t	26.67	24.700	23.37	
							Z	0.583	0.657	0.726	
							pHS	3.640	3.775	3.913	
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75	
3.158	2.936	2.903	2.783	2.690	2.626	2.617	2.580	2.556	2.555	2.551	
22.39	21.72	19.03	18.73	18.50	18.36	21.05	20.97	20.93	18.46	18.45	
0.793	0.855	0.870	0.909	0.941	0.964	0.967	0.980	0.989	0.993	0.996	
4.072	4.258	4.312	4.484	4.688	4.914	4.947	5.180	5.423	5.656	5.899	
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50	
2.545	2.540	2.529	2.527	2.525	2.523	2.543	2.542	2.540	2.537	2.541	
18.44	18.44	25.62	25.62	25.61	25.61	18.40	18.40	18.41	18.42	18.44	
0.998	0.999	0.999	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	
6.146	6.394	6.700	6.950	7.199	7.449	7.644	7.894	8.144	8.394	8.646	
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3 , 1					
2.539	2.537	2.534	2.526	2.523	2.518						
18.48	18.54	18.66	23.77	24.28	25.27						
1.000	1.000	1.000	1.000	1.000	1.000						
8.896	9.147	9.399	9.701	9.956	10.215						
215	m-NITROPHENOL				C6H5NO3		pHL		3.00	3.25	3.50
	-1	33.4	8.399		139.11		RE	23.80	22.79	22.09	
							t	0.004	0.005	0.005	
							Z	5.973	6.060	6.138	
							pHS				
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75	
21.54	21.10	18.36	18.28	18.19	18.12	20.53	20.50	20.48	18.21	18.21	
0.007	0.008	0.012	0.014	0.016	0.019	0.024	0.028	0.035	0.074	0.081	
6.221	6.314	6.474	6.530	6.602	6.690	6.777	6.853	6.945	7.291	7.335	
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50	
24.723	20.887	10.255	9.451	8.430	7.285	4.914	4.555	4.128	3.689	3.161	
18.21	18.21	24.87	24.88	24.89	24.90	18.16	18.16	18.16	18.17	18.19	
0.093	0.110	0.227	0.247	0.278	0.322	0.486	0.525	0.581	0.653	0.767	
7.397	7.478	7.849	7.896	7.964	8.054	8.347	8.414	8.511	8.641	8.883	
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3 , 1					
2.951	2.773	2.643	2.541	2.494	2.464						
18.22	18.28	18.39	23.31	23.78	24.69						
0.824	0.878	0.922	0.958	0.975	0.986						
9.033	9.220	9.435	9.723	9.961	10.213						
216	o-NITROPHENOL				C6H5NO3		pHL		3.00	3.25	3.50
	-1	33.4	7.234		139.11		RE	23.81	22.79	22.09	
							t	0.014	0.017	0.021	
							Z	5.391	5.479	5.558	
							pHS				
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75	
89.568	72.459	51.238	45.044	38.317	31.525	26.323	22.272	18.254	9.625	8.802	
21.52	21.09	18.34	18.25	18.16	18.09	20.52	20.49	20.47	18.17	18.17	
0.025	0.031	0.045	0.051	0.060	0.073	0.087	0.103	0.126	0.244	0.267	
5.641	5.736	5.893	5.952	6.026	6.116	6.202	6.282	6.379	6.722	6.774	
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50	
7.792	6.700	4.268	3.979	3.640	3.300	2.928	2.798	2.678	2.588	2.531	
18.17	18.16	24.95	24.96	24.97	24.99	18.10	18.10	18.11	18.12	18.13	
0.302	0.352	0.560	0.601	0.659	0.729	0.832	0.872	0.912	0.944	0.968	
6.848	6.946	7.310	7.384	7.490	7.632	7.892	8.030	8.211	8.424	8.673	
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3 , 1					
2.497	2.475	2.461	2.447	2.440	2.434						
18.17	18.23	18.34	23.25	23.72	24.62						
0.981	0.989	0.994	0.997	0.998	0.999						
8.904	9.147	9.395	9.693	9.946	10.205						

217	p-NITROPHENOL -1			33.4	7.149	C6H5NO3 139.11		pHL RE t Z pHS	3.00 23.81 0.016 5.349	3.25 22.80 0.019 5.436	3.50 98.291 22.09 0.023 5.515
	3.75 81.358 21.52 0.028 5.599	4.00 65.835 21.09 0.034 5.694	4.00 46.641 18.34 0.049 5.851	4.25 41.010 18.25 0.056 5.910	4.50 34.896 18.16 0.066 5.984	4.75 28.725 18.09 0.080 6.075	4.75 24.031 20.52 0.096 6.161	5.00 20.344 20.49 0.113 6.241	5.25 16.691 20.47 0.138 6.339	5.50 8.923 18.17 0.263 6.681	5.75 8.165 18.17 0.288 6.734
	6.00 7.237 18.16 0.326 6.810	6.25 6.238 18.16 0.379 6.910	6.50 4.073 24.96 0.587 7.273	6.75 3.803 24.97 0.630 7.351	7.00 3.490 24.98 0.688 7.462	7.25 3.182 24.99 0.757 7.611	7.50 2.867 18.10 0.850 7.867	7.75 2.748 18.10 0.888 8.012	8.00 2.643 18.11 0.925 8.200	8.25 2.565 18.12 0.953 8.418	8.50 2.518 18.13 0.973 8.668
	8.75 2.489 18.17 0.984 8.902	9.00 2.470 18.23 0.991 9.146	9.25 2.458 18.34 0.995 9.394	9.50 2.445 23.25 0.997 9.693	9.75 2.439 23.72 0.999 9.946	10.00 2.434 24.62 0.999 10.205	(Ref.) : 3 , 1				
218	NITROUS ACID -1			74.6	3.220	HNO2 47.01		pHL RE t Z pHS	3.00 1.684 16.80 0.625 3.404	3.25 1.501 15.94 0.702 3.554	3.50 1.370 15.36 0.770 3.706
	3.75 1.268 14.94 0.833 3.878	4.00 1.192 14.65 0.888 4.076	4.00 1.175 13.52 0.901 4.138	4.25 1.136 13.39 0.933 4.324	4.50 1.107 13.29 0.958 4.539	4.75 1.088 13.24 0.975 4.772	4.75 1.087 14.36 0.976 4.782	5.00 1.077 14.33 0.986 5.019	5.25 1.070 14.31 0.992 5.263	5.50 1.067 13.27 0.995 5.516	5.75 1.065 13.26 0.997 5.760
	6.00 1.063 13.26 0.999 6.008	6.25 1.063 13.26 0.999 6.257	6.50 1.062 16.26 1.000 6.511	6.75 1.062 16.26 1.000 6.761	7.00 1.062 16.27 1.000 7.010	7.25 1.062 16.27 1.000 7.260	7.50 1.062 13.24 1.000 7.507	7.75 1.062 13.24 1.000 7.757	8.00 1.062 13.25 1.000 8.007	8.25 1.062 13.25 1.000 8.257	8.50 1.062 13.25 1.000 8.507
	8.75 1.062 13.26 1.000 8.757	9.00 1.062 13.28 1.000 9.007	9.25 1.062 13.31 1.000 9.257	9.50 1.062 15.34 1.000 9.510	9.75 1.062 15.43 1.000 9.760	10.00 1.062 15.58 1.000 10.010	(Ref.) : 2 , 7				
219	OCTYLSULFONIC ACID -1			30.1	-2.000	C8H18O3S 194.30		pHL RE t Z pHS	3.00 2.722 30.42 1.000 3.332	3.25 2.722 27.04 1.000 3.536	3.50 2.722 24.92 1.000 3.742
	3.75 2.721 23.55 1.000 3.962	4.00 2.720 22.71 1.000 4.196	4.00 2.736 19.94 1.000 4.200	4.25 2.736 19.49 1.000 4.426	4.50 2.734 19.22 1.000 4.664	4.75 2.732 19.06 1.000 4.910	4.75 2.728 21.96 1.000 4.938	5.00 2.725 21.87 1.000 5.185	5.25 2.723 21.82 1.000 5.434	5.50 2.739 19.16 1.000 5.662	5.75 2.738 19.15 1.000 5.910
	6.00 2.736 19.14 1.000 6.159	6.25 2.733 19.13 1.000 6.409	6.50 2.722 26.87 1.000 6.720	6.75 2.720 26.87 1.000 6.969	7.00 2.718 26.86 1.000 7.219	7.25 2.715 26.85 1.000 7.469	7.50 2.740 19.09 1.000 7.659	7.75 2.738 19.10 1.000 7.909	8.00 2.736 19.11 1.000 8.159	8.25 2.734 19.12 1.000 8.409	8.50 2.738 19.14 1.000 8.662
	8.75 2.735 19.18 1.000 8.912	9.00 2.733 19.26 1.000 9.163	9.25 2.729 19.40 1.000 9.415	9.50 2.720 24.94 1.000 9.722	9.75 2.715 25.56 1.000 9.979	10.00 2.709 26.77 1.000 10.240	(Ref.) : 3 , 12				

220	OROTIC ACID -1	33.1	2.803	C5H4N2O4 156.10	pHL RE t Z pHS	3.00 2.988 27.50 0.819 3.425	3.25 2.817 24.97 0.870 3.595	3.50 2.697 23.31 0.910 3.773		
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
2.611	2.551	2.560	2.527	2.504	2.489	2.485	2.476	2.470	2.481	2.478
22.17	21.44	18.89	18.52	18.28	18.14	20.76	20.68	20.64	18.23	18.22
0.941	0.964	0.965	0.978	0.987	0.993	0.993	0.996	0.998	0.999	0.999
3.971	4.191	4.210	4.421	4.652	4.893	4.920	5.164	5.412	5.643	5.890
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.476	2.473	2.464	2.463	2.461	2.459	2.478	2.477	2.475	2.473	2.476
18.21	18.21	25.21	25.21	25.20	25.20	18.17	18.17	18.18	18.19	18.21
1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
6.139	6.389	6.693	6.943	7.192	7.442	7.639	7.889	8.139	8.389	8.641
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 10				
2.474	2.472	2.469	2.462	2.459	2.455					
18.25	18.31	18.42	23.39	23.87	24.79					
1.000	1.000	1.000	1.000	1.000	1.000					
8.891	9.142	9.393	9.693	9.948	10.207					
221	OXALIC ACID -1 -2	(S) 42.4 77.0	1.271 4.266	C2H2O4 90.04	pHL RE t Z pHS	3.00 1.785 26.36 1.095 3.213	3.25 1.709 24.63 1.159 3.434	3.50 1.619 24.23 1.244 3.651		
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
1.517	1.414	1.397	1.318	1.252	1.203	1.198	1.166	1.145	1.137	1.130
24.63	25.43	23.55	24.31	25.00	25.54	27.88	28.25	28.50	26.52	26.61
1.353	1.481	1.513	1.629	1.738	1.828	1.832	1.895	1.937	1.964	1.979
3.871	4.094	4.146	4.348	4.564	4.794	4.808	5.040	5.280	5.535	5.775
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
1.126	1.123	1.116	1.115	1.114	1.112	1.121	1.121	1.120	1.120	1.121
26.66	26.69	32.97	32.98	32.98	32.97	26.68	26.69	26.70	26.71	26.71
1.988	1.993	1.996	1.998	1.999	1.999	2.000	2.000	2.000	2.000	2.000
6.021	6.270	6.527	6.775	7.025	7.274	7.519	7.769	8.019	8.269	8.519
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 10 , 1				
1.120	1.120	1.119	1.115	1.114	1.113					
26.73	26.76	26.82	31.07	31.25	31.57					
2.000	2.000	2.000	2.000	2.000	2.000					
8.769	9.019	9.269	9.524	9.774	10.024					
222	OXALOACETIC -1 -2	ACID 28.0 56.0	2.555 4.370	C4H4O5 132.07	pHL RE t Z pHS	3.00 2.924 33.32 1.011 3.414	3.25 2.698 30.94 1.102 3.593	3.50 2.492 29.88 1.200 3.775		
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
2.291	2.100	2.069	1.932	1.816	1.727	1.715	1.658	1.620	1.611	1.597
29.68	30.04	27.15	27.67	28.21	28.66	31.99	32.30	32.50	29.67	29.74
1.315	1.446	1.482	1.597	1.710	1.806	1.814	1.882	1.928	1.959	1.976
3.969	4.176	4.231	4.417	4.624	4.848	4.872	5.099	5.337	5.585	5.822
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
1.588	1.582	1.569	1.566	1.564	1.561	1.581	1.580	1.579	1.577	1.580
29.78	29.81	38.66	38.66	38.64	38.62	29.81	29.81	29.81	29.82	29.84
1.986	1.992	1.996	1.998	1.999	1.999	2.000	2.000	2.000	2.000	2.000
6.068	6.316	6.593	6.841	7.090	7.340	7.564	7.814	8.064	8.314	8.565
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 12 , 1				
1.578	1.577	1.575	1.565	1.563	1.560					
29.87	29.92	30.02	36.07	36.40	37.03					
2.000	2.000	2.000	2.000	2.000	2.000					
8.815	9.065	9.315	9.587	9.838	10.091					

CHROMATOGRAPHIC DATA (1983)

D85

223	PELARGONIC -1	ACID 26.7	4.955	C9H18O2 158.24	pHL RE t Z pHS	3.00 16.050 28.51 0.182 4.287	3.25 13.381 26.84 0.219 4.386	3.50 11.369 25.71 0.259 4.479		
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
9.613	8.038	6.880	6.160	5.433	4.766	4.534	4.103	3.750	3.467	3.347
24.85	24.22	20.79	20.61	20.45	20.32	23.54	23.49	23.45	20.47	20.46
0.307	0.369	0.436	0.489	0.557	0.637	0.671	0.744	0.816	0.892	0.925
4.582	4.700	4.819	4.909	5.026	5.170	5.234	5.386	5.569	5.838	6.013
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
3.253	3.189	3.128	3.108	3.095	3.087	3.116	3.113	3.108	3.105	3.109
20.45	20.45	29.22	29.22	29.21	29.20	20.41	20.42	20.42	20.44	20.47
0.953	0.972	0.986	0.992	0.995	0.997	0.998	0.999	0.999	1.000	1.000
6.223	6.454	6.776	7.013	7.258	7.505	7.690	7.938	8.188	8.438	8.691
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 10 , 7				
3.106	3.102	3.098	3.084	3.078	3.069					
20.53	20.62	20.80	27.20	28.05	29.79					
1.000	1.000	1.000	1.000	1.000	1.000					
8.941	9.192	9.444	9.762	10.020	10.287					
224	PERCHLORIC -1	ACID (S) 69.8	-2.000	HC104 100.46	pHL RE t Z pHS	3.00 1.137 19.06 1.000 3.039	3.25 1.137 17.12 1.000 3.282	3.50 1.137 16.01 1.000 3.525		
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
1.137	1.137	1.137	1.137	1.137	1.137	1.137	1.137	1.137	1.137	1.137
15.38	15.02	13.88	13.67	13.56	13.49	14.69	14.66	14.64	13.53	13.52
1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
3.771	4.019	4.020	4.267	4.515	4.765	4.768	5.018	5.268	5.515	5.765
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
1.137	1.137	1.137	1.137	1.137	1.137	1.137	1.137	1.137	1.137	1.137
13.52	13.52	16.74	16.74	16.74	16.74	13.50	13.50	13.51	13.51	13.51
1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
6.015	6.265	6.522	6.772	7.022	7.272	7.515	7.765	8.015	8.265	8.515
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 7 , 12				
1.137	1.137	1.137	1.137	1.137	1.136					
13.52	13.54	13.57	15.76	15.85	16.03					
1.000	1.000	1.000	1.000	1.000	1.000					
8.765	9.015	9.265	9.521	9.771	10.022					
225	PERIODIC ACID -1	ACID 56.5	1.550	HI04 191.91	pHL RE t Z pHS	3.00 1.447 20.93 0.976 3.127	3.25 1.433 18.81 0.986 3.351	3.50 1.425 17.55 0.992 3.578		
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
1.420	1.417	1.419	1.417	1.416	1.415	1.415	1.414	1.413	1.416	1.416
16.79	16.35	14.92	14.67	14.53	14.45	15.95	15.90	15.88	14.50	14.49
0.995	0.997	0.997	0.998	0.999	0.999	0.999	1.000	1.000	1.000	1.000
3.814	4.057	4.061	4.300	4.546	4.794	4.803	5.052	5.302	5.544	5.794
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
1.415	1.415	1.414	1.413	1.413	1.412	1.416	1.415	1.415	1.415	1.415
14.49	14.49	18.49	18.49	18.49	18.49	14.46	14.46	14.47	14.47	14.48
1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
6.043	6.293	6.564	6.813	7.063	7.313	7.543	7.793	8.043	8.293	8.544
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 2 , 4				
1.415	1.415	1.414	1.413	1.413	1.412					
14.49	14.52	14.56	17.30	17.45	17.71					
1.000	1.000	1.000	1.000	1.000	1.000					
8.794	9.044	9.294	9.561	9.812	10.064					

226	PERMANGANIC ACID -1 63.4			-2.000	HMn04 119.90		pHL RE t Z pHS	3.00 1.255 19.92 1.000 3.070	3.25 1.255 17.86 1.000 3.308	3.50 1.255 16.67 1.000 3.546	
	3.75 1.255 15.98 1.000 3.789	4.00 1.255 15.58 1.000 4.035	4.00 1.256 14.32 1.000 4.036	4.25 1.256 14.10 1.000 4.281	4.50 1.256 13.97 1.000 4.528	4.75 1.256 13.90 1.000 4.777	4.75 1.255 15.23 1.000 4.783	5.00 1.255 15.19 1.000 5.033	5.25 1.255 15.17 1.000 5.282	5.50 1.256 13.94 1.000 5.528	5.75 1.256 13.94 1.000 5.777
	6.00 1.256 13.93 1.000 6.027	6.25 1.256 13.93 1.000 6.277	6.50 1.255 17.49 1.000 6.540	6.75 1.255 17.49 1.000 6.790	7.00 1.255 17.49 1.000 7.040	7.25 1.255 17.49 1.000 7.290	7.50 1.256 13.91 1.000 7.527	7.75 1.256 13.91 1.000 7.777	8.00 1.256 13.92 1.000 8.027	8.25 1.256 13.92 1.000 8.277	8.50 1.256 13.92 1.000 8.528
	8.75 1.256 13.94 1.000 8.778	9.00 1.256 13.96 1.000 9.028	9.25 1.255 13.99 1.000 9.278	9.50 1.255 16.41 1.000 9.538	9.75 1.255 16.53 1.000 9.789	10.00 1.254 16.74 1.000 10.040	(Ref.) : 2 , 12				
227	PEROXYSULFURIC ACID -1 44.6 -2 89.1			-2.000 -1.000	H2S208 194.13		pHL RE t Z pHS	3.00 0.958 35.24 2.000 2.987	3.25 0.958 31.74 2.000 3.242	3.50 0.958 29.79 2.000 3.496	
	3.75 0.957 28.71 2.000 3.748	4.00 0.957 28.11 2.000 3.999	4.00 0.960 26.23 2.000 3.999	4.25 0.960 25.88 2.000 4.250	4.50 0.960 25.69 2.000 4.501	4.75 0.960 25.59 2.000 4.751	4.75 0.958 27.59 2.000 4.750	5.00 0.957 27.53 2.000 5.000	5.25 0.957 27.50 2.000 5.250	5.50 0.960 25.65 2.000 5.501	5.75 0.960 25.64 2.000 5.751
	6.00 0.959 25.64 2.000 6.001	6.25 0.959 25.64 2.000 6.251	6.50 0.956 31.01 2.000 6.499	6.75 0.955 31.00 2.000 6.749	7.00 0.955 31.00 2.000 6.999	7.25 0.954 31.00 2.000 7.249	7.50 0.960 25.60 2.000 7.502	7.75 0.960 25.61 2.000 7.752	8.00 0.960 25.61 2.000 8.001	8.25 0.959 25.62 2.000 8.251	8.50 0.960 25.62 2.000 8.501
	8.75 0.960 25.64 2.000 8.751	9.00 0.959 25.67 2.000 9.001	9.25 0.959 25.71 2.000 9.251	9.50 0.956 29.34 2.000 9.499	9.75 0.955 29.48 2.000 9.749	10.00 0.955 29.73 2.000 9.999	(Ref.) : 2 , 12				
228	PERRHENIC ACID -1 56.7			0.000	HRe04 251.20		pHL RE t Z pHS	3.00 1.409 21.02 0.999 3.108	3.25 1.409 18.82 1.000 3.339	3.50 1.409 17.53 1.000 3.571	
	3.75 1.408 16.77 1.000 3.810	4.00 1.408 16.32 1.000 4.055	4.00 1.410 14.90 1.000 4.057	4.25 1.410 14.65 1.000 4.298	4.50 1.410 14.51 1.000 4.545	4.75 1.409 14.43 1.000 4.793	4.75 1.409 15.92 1.000 4.802	5.00 1.409 15.88 1.000 5.051	5.25 1.408 15.86 1.000 5.301	5.50 1.410 14.48 1.000 5.544	5.75 1.410 14.47 1.000 5.793
	6.00 1.410 14.47 1.000 6.043	6.25 1.409 14.47 1.000 6.293	6.50 1.408 18.46 1.000 6.563	6.75 1.408 18.46 1.000 6.813	7.00 1.408 18.46 1.000 7.063	7.25 1.407 18.46 1.000 7.313	7.50 1.411 14.44 1.000 7.543	7.75 1.410 14.44 1.000 7.793	8.00 1.410 14.45 1.000 8.043	8.25 1.410 14.46 1.000 8.293	8.50 1.410 14.46 1.000 8.543
	8.75 1.410 14.47 1.000 8.793	9.00 1.409 14.50 1.000 9.044	9.25 1.409 14.54 1.000 9.294	9.50 1.408 17.27 1.000 9.560	9.75 1.407 17.42 1.000 9.811	10.00 1.407 17.68 1.000 10.063	(Ref.) : 2 , 12				

CHROMATOGRAPHIC DATA (1983)

D87

229	PHENOL -1	34.4	9.998	C6H6O 94.11	pHL RE t Z pHS	3.00 24.19 0.001 6.781	3.25 23.31 0.001 6.867	3.50 22.69 0.001 6.945		
	3.75	4.00	4.00	4.25	4.50	4.75	5.00	5.25	5.50	5.75
	22.18	21.76	19.01	18.94	18.87	18.80	21.20	21.17	21.15	18.90
	0.001	0.001	0.002	0.002	0.003	0.003	0.004	0.005	0.006	0.013
	7.026	7.118	7.283	7.338	7.408	7.494	7.583	7.657	7.746	8.099
	6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25
	18.90	18.90	50.561	46.612	41.437	35.408	18.542	17.040	15.112	12.921
	0.016	0.019	0.044	0.047	0.053	0.063	0.121	0.132	0.149	0.174
	8.197	8.272	8.650	8.687	8.740	8.813	9.122	9.164	9.224	9.305
	8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3 , 7			
	7.457	6.419	5.417	4.083	3.612	3.199				
	19.23	19.34	19.51	24.78	25.39	26.52				
	0.306	0.356	0.424	0.566	0.642	0.727				
	9.620	9.717	9.839	10.086	10.222	10.392				
230	PHENOXYACETIC ACID -1	27.8	3.171	C8H8O3 152.15	pHL RE t Z pHS	3.00 4.040 30.42 0.724 3.558	3.25 3.718 27.75 0.788 3.710	3.50 3.484 25.96 0.843 3.868		
	3.75	4.00	4.00	4.25	4.50	4.75	5.00	5.25	5.50	5.75
	3.304	3.172	3.178	3.104	3.051	3.016	3.007	2.987	2.988	2.983
	24.69	23.85	20.76	20.35	20.07	19.90	23.05	22.96	22.90	20.02
	0.891	0.929	0.934	0.957	0.974	0.984	0.986	0.992	0.995	0.997
	4.048	4.255	4.284	4.479	4.700	4.937	4.970	5.211	5.457	5.930
	6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25
	2.979	2.974	2.959	2.957	2.954	2.951	2.981	2.979	2.977	2.974
	19.99	19.98	28.40	28.39	28.38	28.37	19.95	19.95	19.96	19.97
	0.999	0.999	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
	6.178	6.428	6.743	6.993	7.242	7.492	7.678	7.928	8.178	8.428
	8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 11 , 1			
	2.976	2.972	2.968	2.956	2.950	2.943				
	20.05	20.14	20.30	26.40	27.16	28.69				
	1.000	1.000	1.000	1.000	1.000	1.000				
	8.931	9.182	9.434	9.748	10.006	10.270				
231	PHENYLACETIC ACID -1	31.7	4.405	C8H8O2 136.15	pHL RE t Z pHS	3.00 8.332 25.69 0.298 4.011	3.25 7.058 24.22 0.353 4.119	3.50 6.099 23.21 0.410 4.222		
	3.75	4.00	4.00	4.25	4.50	4.75	5.00	5.25	5.50	5.75
	5.270	4.539	4.126	3.774	3.437	3.151	3.086	2.907	2.777	2.660
	22.44	21.87	19.02	18.83	18.66	18.54	21.25	21.19	21.15	18.64
	0.476	0.555	0.616	0.675	0.744	0.814	0.831	0.883	0.925	0.957
	4.337	4.472	4.580	4.691	4.834	5.010	5.061	5.248	5.462	5.715
	6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25
	2.630	2.610	2.589	2.583	2.578	2.574	2.595	2.593	2.591	2.588
	18.62	18.61	25.95	25.94	25.93	25.93	18.58	18.58	18.59	18.60
	0.984	0.991	0.995	0.997	0.999	0.999	0.999	1.000	1.000	1.000
	6.160	6.403	6.713	6.957	7.205	7.455	7.649	7.898	8.148	8.398
	8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 10 , 1			
	2.590	2.587	2.584	2.576	2.572	2.567				
	18.66	18.73	18.85	24.07	24.61	25.65				
	1.000	1.000	1.000	1.000	1.000	1.000				
	8.900	9.151	9.403	9.706	9.962	10.222				

232	4-PHENYLBUTYRIC -1	ACID 24.7	4.757	C10H12O2 164.20			pHL RE t Z pHS	3.00 14.144 30.42 0.225 4.203	3.25 11.861 28.50 0.270 4.305	3.50 10.133 27.20 0.317 4.402
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
8.629	7.289	6.458	5.816	5.184	4.624	4.457	4.098	3.822	3.641	3.544
26.22	25.50	21.82	21.60	21.41	21.27	24.78	24.71	24.67	21.43	21.42
0.373	0.444	0.507	0.565	0.636	0.717	0.743	0.811	0.872	0.925	0.951
4.510	4.636	4.743	4.843	4.971	5.129	5.189	5.358	5.556	5.811	6.005
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
3.474	3.428	3.377	3.364	3.354	3.346	3.384	3.380	3.376	3.372	3.378
21.41	21.40	30.93	30.92	30.91	30.90	21.37	21.37	21.38	21.40	21.44
0.970	0.982	0.992	0.995	0.997	0.998	0.999	0.999	1.000	1.000	1.000
6.229	6.467	6.791	7.033	7.280	7.529	7.708	7.957	8.207	8.457	8.710
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 11 , 7				
3.374	3.370	3.364	3.346	3.339	3.328					
21.50	21.62	21.83	28.87	29.94	32.19					
1.000	1.000	1.000	1.000	1.000	1.000					
8.961	9.212	9.465	9.790	10.050	10.320					
233	2-PHENYLPROPIONIC -1	ACID 26.5	2.237	C9H10O2 150.20			pHL RE t Z pHS	3.00 3.294 32.83 0.942 3.417	3.25 3.230 29.33 0.962 3.604	3.50 3.187 27.08 0.975 3.797
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
3.158	3.138	3.159	3.147	3.138	3.132	3.125	3.120	3.115	3.137	3.134
25.57	24.63	21.44	20.95	20.63	20.45	23.77	23.67	23.60	20.58	20.56
0.985	0.991	0.991	0.995	0.997	0.998	0.998	0.999	0.999	1.000	1.000
4.007	4.235	4.244	4.462	4.697	4.940	4.973	5.219	5.467	5.692	5.940
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
3.132	3.128	3.111	3.109	3.107	3.103	3.136	3.135	3.132	3.129	3.134
20.54	20.54	29.39	29.38	29.37	29.36	20.50	20.50	20.51	20.53	20.56
1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
6.189	6.439	6.757	7.007	7.256	7.506	7.689	7.939	8.189	8.439	8.692
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 11 , 3				
3.131	3.127	3.123	3.108	3.102	3.093					
20.62	20.72	20.90	27.35	28.22	30.01					
1.000	1.000	1.000	1.000	1.000	1.000					
8.943	9.194	9.446	9.765	10.023	10.290					
234	3-PHENYLPROPIONIC -1	ACID 26.5	4.664	C9H10O2 150.20			pHL RE t Z pHS	3.00 12.233 28.94 0.243 4.151	3.25 10.283 27.17 0.290 4.255	3.50 8.810 25.96 0.339 4.353
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
7.528	6.390	5.693	5.148	4.614	4.144	4.013	3.714	3.488	3.344	3.265
25.05	24.38	20.95	20.74	20.56	20.42	23.69	23.62	23.58	20.57	20.55
0.399	0.472	0.536	0.594	0.665	0.744	0.768	0.832	0.888	0.935	0.958
4.462	4.590	4.698	4.801	4.932	5.094	5.152	5.326	5.528	5.783	5.982
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
3.209	3.173	3.133	3.121	3.113	3.107	3.139	3.136	3.133	3.129	3.134
20.54	20.54	29.38	29.38	29.37	29.36	20.50	20.50	20.51	20.53	20.56
0.974	0.985	0.993	0.996	0.998	0.999	0.999	1.000	1.000	1.000	1.000
6.208	6.448	6.768	7.011	7.258	7.507	7.691	7.940	8.189	8.439	8.692
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 11 , 1				
3.131	3.127	3.123	3.108	3.102	3.093					
20.62	20.72	20.90	27.35	28.22	30.01					
1.000	1.000	1.000	1.000	1.000	1.000					
8.943	9.194	9.446	9.765	10.023	10.290					

CHROMATOGRAPHIC DATA (1983)

D89

235	PHOSPHORIC	ACID (S)			H3P04			pHL	3.00	3.25	3.50
	-1	35.1	2.120		98.00			RE	2.449	2.401	2.369
	-2	61.5	7.470					t	27.20	24.38	22.60
	-3	71.5	12.360					Z	0.944	0.964	0.977
								pHS	3.313	3.510	3.711
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75	
2.348	2.333	2.344	2.336	2.328	2.322	2.319	2.312	2.302	2.300	2.278	
21.45	20.74	18.38	18.01	17.79	17.68	20.16	20.13	20.18	18.06	18.31	
0.986	0.992	0.993	0.996	0.999	1.002	1.002	1.005	1.010	1.019	1.034	
3.929	4.162	4.170	4.394	4.633	4.878	4.903	5.150	5.398	5.645	5.893	
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50	
2.241	2.183	2.037	1.936	1.825	1.715	1.619	1.559	1.510	1.475	1.456	
18.74	19.42	28.10	29.47	30.97	32.47	26.37	27.12	27.72	28.15	28.43	
1.058	1.099	1.209	1.300	1.411	1.536	1.686	1.775	1.852	1.909	1.947	
6.141	6.389	6.770	6.975	7.183	7.397	7.665	7.860	8.080	8.314	8.564	
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 11 , 4					
1.443	1.434	1.429	1.419	1.415	1.412						
28.62	28.76	28.90	34.40	34.74	35.32						
1.969	1.983	1.991	1.997	2.002	2.007						
8.806	9.053	9.302	9.568	9.818	10.070						
236	PHOSPHOROUS	ACID			H3P03			pHL	3.00	3.25	3.50
	-1	40.0	2.000		82.00			RE	2.120	2.083	2.059
	-2	65.4	7.000					t	25.15	22.56	20.95
								Z	0.952	0.970	0.982
								pHS	3.262	3.465	3.673
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75	
2.043	2.032	2.040	2.032	2.026	2.018	2.016	2.006	1.990	1.972	1.933	
19.94	19.33	17.26	16.95	16.79	16.74	18.91	18.99	19.19	17.61	18.23	
0.990	0.995	0.995	0.999	1.003	1.008	1.009	1.016	1.030	1.054	1.093	
3.895	4.130	4.138	4.366	4.607	4.853	4.874	5.122	5.371	5.638	5.884	
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50	
1.875	1.796	1.648	1.573	1.501	1.440	1.407	1.380	1.361	1.349	1.344	
19.17	20.44	29.42	30.83	32.18	33.32	26.94	27.39	27.70	27.90	28.02	
1.152	1.240	1.423	1.532	1.648	1.756	1.848	1.903	1.941	1.966	1.980	
6.127	6.368	6.730	6.918	7.123	7.344	7.596	7.815	8.051	8.295	8.546	
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 11 , 6					
1.339	1.336	1.334	1.327	1.324	1.322						
28.11	28.19	28.28	33.38	33.63	34.08						
1.989	1.994	1.996	1.998	1.999	1.999						
8.793	9.042	9.292	9.555	9.805	10.056						
237	PTHALIC ACID				C8H6O4			pHL	3.00	3.25	3.50
	-1	28.1	2.950		166.13			RE	3.608	3.360	3.173
	-2	52.9	5.408					t	30.89	28.17	26.48
								Z	0.807	0.869	0.924
								pHS	3.499	3.661	3.831
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75	
3.013	2.866	2.857	2.722	2.565	2.390	2.353	2.186	2.033	1.903	1.829	
25.50	25.19	22.20	22.54	23.34	24.50	28.03	29.31	30.55	28.70	29.29	
0.977	1.034	1.047	1.108	1.190	1.297	1.318	1.441	1.572	1.721	1.805	
4.022	4.236	4.278	4.481	4.700	4.925	4.969	5.183	5.403	5.681	5.879	
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50	
1.772	1.733	1.695	1.681	1.671	1.664	1.686	1.683	1.681	1.679	1.682	
29.75	30.07	39.66	39.76	39.81	39.82	30.47	30.49	30.50	30.51	30.54	
1.874	1.923	1.958	1.975	1.986	1.992	1.995	1.997	1.998	1.999	2.000	
6.101	6.337	6.625	6.862	7.106	7.354	7.576	7.825	8.074	8.324	8.575	
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 10 , 1					
1.680	1.678	1.675	1.665	1.661	1.658						
30.57	30.62	30.73	37.18	37.56	38.27						
2.000	2.000	2.000	2.000	2.000	2.000						
8.825	9.075	9.325	9.600	9.852	10.105						

D90

CHROMATOGRAPHIC DATA (1983)

238	PICRIC ACID -1 31.5			0.708	C6H3N3O7 229.11		pHL RE t Z pHS	3.00 2.601 29.50 0.998 3.317	3.25 2.598 26.24 0.999 3.522	3.50 2.597 24.20 0.999 3.729	
	3.75 2.596 22.90 0.999 3.950	4.00 2.594 22.09 1.000 4.184	4.00 2.609 19.45 1.000 4.189	4.25 2.608 19.02 1.000 4.415	4.50 2.606 18.76 1.000 4.654	4.75 2.604 18.60 1.000 4.900	4.75 2.601 21.37 1.000 4.927	5.00 2.598 21.29 1.000 5.174	5.25 2.596 21.24 1.000 5.423	5.50 2.611 18.71 1.000 5.651	5.75 2.609 18.69 1.000 5.900
	6.00 2.607 18.68 1.000 6.149	6.25 2.605 18.68 1.000 6.399	6.50 2.595 26.06 1.000 6.707	6.75 2.593 26.05 1.000 6.956	7.00 2.592 26.05 1.000 7.206	7.25 2.589 26.04 1.000 7.456	7.50 2.611 18.64 1.000 7.649	7.75 2.610 18.64 1.000 7.899	8.00 2.608 18.65 1.000 8.149	8.25 2.605 18.66 1.000 8.399	8.50 2.609 18.68 1.000 8.651
	8.75 2.607 18.72 1.000 8.902	9.00 2.605 18.79 1.000 9.153	9.25 2.602 18.92 1.000 9.404	9.50 2.593 24.17 1.000 9.708	9.75 2.589 24.72 1.000 9.964	10.00 2.584 25.78 1.000 10.224	(Ref.) : 2 , 7				
239	PIMELIC ACID -1 -2 24.0 49.9			4.509 5.312	C7H12O4 160.17		pHL RE t Z pHS	3.00 10.544 32.34 0.313 4.080	3.25 8.738 30.57 0.379 4.184	3.50 7.364 29.52 0.452 4.283	
	3.75 6.166 28.90 0.542 4.392	4.00 5.095 28.72 0.660 4.519	4.00 4.412 25.48 0.773 4.629	4.25 3.882 25.80 0.882 4.731	4.50 3.359 26.34 1.026 4.861	4.75 2.892 27.10 1.199 5.020	4.75 2.764 31.04 1.254 5.074	5.00 2.458 31.83 1.418 5.239	5.25 2.215 32.63 1.580 5.433	5.50 2.031 30.12 1.750 5.704	5.75 1.947 30.50 1.830 5.892
	6.00 1.884 30.80 1.894 6.111	6.25 1.842 31.00 1.937 6.347	6.50 1.802 41.15 1.967 6.637	6.75 1.788 41.20 1.981 6.875	7.00 1.779 41.22 1.989 7.120	7.25 1.772 41.21 1.996 7.368	7.50 1.797 31.25 1.996 7.586	7.75 1.794 31.26 1.998 7.835	8.00 1.792 31.26 1.999 8.084	8.25 1.789 31.27 1.999 8.334	8.50 1.793 31.30 2.000 8.585
	8.75 1.791 31.34 2.000 8.835	9.00 1.788 31.40 2.000 9.085	9.25 1.785 31.51 2.000 9.336	9.50 1.773 38.41 2.000 9.614	9.75 1.769 38.85 2.000 9.866	10.00 1.765 39.66 2.000 10.120	(Ref.) : 10 , 1				
240	PROPIONIC ACID (S) -1 37.1			4.874	C3H6O2 74.08		pHL RE t Z pHS	3.00 11.190 22.76 0.187 4.218	3.25 9.352 21.71 0.224 4.316	3.50 7.973 20.98 0.264 4.407	
	3.75 6.766 20.41 0.312 4.507	4.00 5.675 19.97 0.373 4.624	4.00 4.747 17.50 0.450 4.760	4.25 4.278 17.38 0.501 4.846	4.50 3.796 17.27 0.566 4.959	4.75 3.344 17.17 0.645 5.100	4.75 3.185 19.45 0.678 5.164	5.00 2.895 19.41 0.748 5.312	5.25 2.653 19.37 0.818 5.491	5.50 2.439 17.24 0.897 5.775	5.75 2.359 17.23 0.928 5.945
	6.00 2.296 17.22 0.954 6.153	6.25 2.253 17.22 0.972 6.383	6.50 2.217 23.43 0.986 6.688	6.75 2.203 23.43 0.992 6.922	7.00 2.195 23.43 0.995 7.166	7.25 2.189 23.42 0.997 7.413	7.50 2.201 17.18 0.998 7.618	7.75 2.198 17.19 0.999 7.866	8.00 2.196 17.19 0.999 8.116	8.25 2.194 17.20 1.000 8.366	8.50 2.196 17.22 1.000 8.617
	8.75 2.195 17.24 1.000 8.867	9.00 2.193 17.30 1.000 9.118	9.25 2.191 17.39 1.000 9.369	9.50 2.186 21.74 1.000 9.661	9.75 2.184 22.11 1.000 9.914	10.00 2.180 22.79 1.000 10.171	(Ref.) : 2 , 1				

CHROMATOGRAPHIC DATA (1983)

D91

241	3-PROPYLGLUTARIC ACID				C8H14O4		pHL	3.00	3.25	3.50
	-1	23.5	4.312		174.20		RE	9.319	7.837	6.702
	-2	47.0	5.386				t	32.96	30.96	29.74
							Z	0.365	0.437	0.513
							pHS	4.000	4.109	4.215
	3.75	4.00	4.00	4.25	4.50	4.75	5.00	5.25	5.50	5.75
	5.710	4.824	4.360	3.883	3.416	2.998	2.890	2.599	2.363	2.181
	29.00	28.75	25.31	25.63	26.23	27.11	31.24	32.21	33.18	30.66
	0.606	0.724	0.812	0.918	1.053	1.212	1.257	1.410	1.564	1.729
	4.333	4.472	4.571	4.688	4.835	5.009	5.062	5.241	5.442	5.713
	6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25
	2.019	1.972	1.926	1.910	1.899	1.891	1.920	1.917	1.914	1.911
	31.50	31.75	42.61	42.68	42.70	42.70	32.08	32.09	32.10	32.11
	1.881	1.928	1.962	1.978	1.987	1.993	1.996	1.998	1.999	1.999
	6.123	6.359	6.652	6.890	7.135	7.383	7.597	7.846	8.095	8.345
	8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 12 , 7			
	1.912	1.909	1.906	1.892	1.887	1.882				
	32.18	32.25	32.38	39.77	40.27	41.20				
	2.000	2.000	2.000	2.000	2.000	2.000				
	8.846	9.096	9.347	9.629	9.882	10.136				
242	PROPYLMALONIC ACID				C6H10O4		pHL	3.00	3.25	3.50
	-1	26.0	2.971		146.14		RE	3.937	3.683	3.496
	-2	52.0	5.845				t	32.46	29.45	27.50
							Z	0.799	0.857	0.905
							pHS	3.525	3.687	3.855
	3.75	4.00	4.00	4.25	4.50	4.75	5.00	5.25	5.50	5.75
	3.345	3.219	3.225	3.120	2.998	2.846	2.809	2.627	2.427	2.188
	26.22	25.52	22.25	22.15	22.43	23.12	26.72	27.78	29.03	27.59
	0.947	0.986	0.992	1.028	1.072	1.134	1.148	1.234	1.345	1.522
	4.045	4.258	4.290	4.494	4.719	4.952	4.998	5.223	5.448	5.753
	6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25
	1.947	1.859	1.776	1.743	1.721	1.706	1.724	1.719	1.715	1.711
	29.16	29.74	39.75	39.96	40.08	40.15	30.65	30.68	30.70	30.72
	1.731	1.820	1.901	1.938	1.963	1.979	1.987	1.993	1.996	1.998
	6.138	6.361	6.660	6.880	7.117	7.361	7.583	7.829	8.078	8.327
	8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 12 , 2			
	1.712	1.709	1.707	1.696	1.692	1.689				
	30.79	30.84	30.95	37.54	37.93	38.67				
	1.999	2.000	2.000	2.000	2.000	2.000				
	8.828	9.078	9.328	9.604	9.856	10.109				
243	PROPYLSULFONIC ACID				C3H8O3S		pHL	3.00	3.25	3.50
	-1	38.5	-2.000		124.15		RE	2.104	2.104	2.104
							t	26.01	23.17	21.44
							Z	1.000	1.000	1.000
							pHS	3.241	3.455	3.670
	3.75	4.00	4.00	4.25	4.50	4.75	5.00	5.25	5.50	5.75
	2.104	2.103	2.111	2.111	2.110	2.109	2.107	2.106	2.104	2.113
	20.35	19.69	17.55	17.20	16.99	16.87	19.11	19.04	19.00	16.94
	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
	3.897	4.135	4.139	4.370	4.612	4.858	4.879	5.127	5.376	5.610
	6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25
	2.111	2.110	2.104	2.103	2.102	2.101	2.113	2.112	2.111	2.110
	16.93	16.92	22.90	22.90	22.90	22.89	16.89	16.89	16.90	16.91
	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
	6.108	6.358	6.653	6.902	7.152	7.402	7.608	7.858	8.108	8.358
	8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 2 , 12			
	2.111	2.109	2.108	2.103	2.101	2.098				
	16.95	16.99	17.08	21.26	21.59	22.22				
	1.000	1.000	1.000	1.000	1.000	1.000				
	8.860	9.110	9.361	9.651	9.904	10.160				

244	PYRAZOLE-2,3-DICARBOXYLIC ACID	C5H5N2O4	pH	3.00	3.25	3.50
-1	33.8	1.000	RE	2.258	2.180	2.086
-2	56.0	4.228	t	30.82	28.74	28.14
			Z	1.126	1.194	1.283
			pHS	3.286	3.495	3.702
3.75	4.00	4.00	4.25	4.50	4.75	4.75
1.980	1.871	1.866	1.782	1.712	1.660	1.652
28.43	29.23	26.53	27.30	28.00	28.56	31.83
1.396	1.525	1.549	1.663	1.767	1.850	1.855
3.916	4.137	4.176	4.380	4.600	4.833	4.853
5.00	5.25	5.50	5.75	6.00	6.25	6.50
1.584	1.580	1.568	1.566	1.563	1.560	1.581
29.78	29.81	38.66	38.65	38.64	38.62	29.81
1.990	1.994	1.997	1.998	1.999	1.999	2.000
6.066	6.315	6.591	6.840	7.090	7.339	7.564
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 10
1.578	1.577	1.575	1.565	1.563	1.560	
29.87	29.92	30.02	36.07	36.40	37.03	
2.000	2.000	2.000	2.000	2.000	2.000	
8.815	9.065	9.315	9.587	9.838	10.091	
245	PYRROPHOSPHORIC ACID	H4P2O7	pH	3.00	3.25	3.50
-1	24.9	1.520	RE	1.604	1.568	1.546
-2	57.9	2.360	t	42.47	38.28	35.77
-3	75.3	6.750	Z	1.895	1.935	1.962
-4	88.8	9.290	pHS	3.163	3.379	3.601
3.75	4.00	4.00	4.25	4.50	4.75	4.75
1.531	1.521	1.533	1.527	1.522	1.516	1.511
34.26	33.38	30.38	29.92	29.70	29.66	32.88
1.979	1.990	1.991	1.999	2.007	2.018	2.018
3.834	4.075	4.080	4.317	4.562	4.810	4.822
5.00	5.25	5.50	5.75	6.00	6.25	6.50
1.441	1.405	1.350	1.319	1.292	1.272	1.286
33.32	34.94	46.74	48.38	49.73	50.78	41.24
2.279	2.408	2.579	2.705	2.814	2.901	2.971
6.067	6.312	6.605	6.831	7.066	7.308	7.552
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 11 , 4
1.230	1.215	1.201	1.174	1.164	1.157	
47.45	49.26	50.91	60.49	61.66	62.82	
3.463	3.609	3.737	3.835	3.900	3.942	
8.793	9.040	9.288	9.544	9.792	10.041	
246	PYRUVIC ACID	C3H4O3	pH	3.00	3.25	3.50
-1	42.3	2.490	RE	2.171	2.078	2.016
			t	23.90	21.67	20.25
			Z	0.876	0.916	0.945
			pHS	3.301	3.489	3.683
3.75	4.00	4.00	4.25	4.50	4.75	4.75
1.973	1.946	1.951	1.935	1.925	1.918	1.917
19.31	18.73	16.78	16.47	16.28	16.18	18.20
0.965	0.979	0.980	0.988	0.993	0.996	0.996
3.896	4.125	4.139	4.361	4.599	4.843	4.862
5.00	5.25	5.50	5.75	6.00	6.25	6.50
1.913	1.912	1.908	1.907	1.906	1.905	1.915
16.23	16.23	21.65	21.64	21.64	21.64	16.19
1.000	1.000	1.000	1.000	1.000	1.000	1.000
6.091	6.341	6.629	6.879	7.129	7.379	7.591
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 11 , 7
1.913	1.912	1.910	1.907	1.906	1.903	
16.24	16.28	16.35	20.12	20.38	20.88	
1.000	1.000	1.000	1.000	1.000	1.000	
8.842	9.092	9.343	9.627	9.879	10.134	

CHROMATOGRAPHIC DATA (1983)

D93

247	SACCHARIN -1	31.3	2.432	C7H5NO3S 183.20	pHL RE t Z pHS	3.00 2.874 29.06 0.906 3.381	3.25 2.783 26.12 0.937 3.566	3.50 2.722 24.23 0.958 3.757			
	3.75 2.679 22.96 0.974 3.966	4.00 2.651 22.17 0.984 4.194	4.00 2.665 19.49 0.985 4.205	4.25 2.648 19.08 0.991 4.424	4.50 2.637 18.82 0.995 4.659	4.75 2.629 18.67 0.997 4.903	4.75 2.625 21.45 0.997 4.931	5.00 2.620 21.37 0.998 5.176	5.25 2.615 21.32 0.999 5.425	5.50 2.630 18.77 0.999 5.654	5.75 2.628 18.75 1.000 5.902
	6.00 2.625 18.74 1.000 6.151	6.25 2.623 18.74 1.000 6.401	6.50 2.612 26.17 1.000 6.709	6.75 2.611 26.16 1.000 6.958	7.00 2.609 26.16 1.000 7.208	7.25 2.607 26.15 1.000 7.458	7.50 2.629 18.70 1.000 7.651	7.75 2.627 18.71 1.000 7.901	8.00 2.626 18.71 1.000 8.151	8.25 2.623 18.73 1.000 8.401	8.50 2.627 18.75 1.000 8.653
	8.75 2.625 18.79 1.000 8.903	9.00 2.622 18.86 1.000 9.154	9.25 2.619 18.98 1.000 9.406	9.50 2.610 24.28 1.000 9.710	9.75 2.607 24.83 1.000 9.966	10.00 2.601 25.91 1.000 10.226	(Ref.) : 3				
248	SALICYLIC ACID -1	35.4	3.107	C7H6O3 138.12	pHL RE t Z pHS	3.00 3.155 25.79 0.721 3.485	3.25 2.895 23.67 0.787 3.641	3.50 2.708 22.26 0.843 3.802			
	3.75 2.565 21.27 0.891 3.985	4.00 2.461 20.61 0.930 4.193	4.00 2.460 18.22 0.935 4.226	4.25 2.403 17.89 0.958 4.423	4.50 2.363 17.68 0.974 4.646	4.75 2.336 17.55 0.985 4.884	4.75 2.332 19.99 0.986 4.910	5.00 2.317 19.92 0.992 5.151	5.25 2.307 19.88 0.995 5.397	5.50 2.314 17.63 0.997 5.631	5.75 2.310 17.62 0.998 5.877
	6.00 2.308 17.61 0.999 6.125	6.25 2.305 17.61 1.000 6.375	6.50 2.297 24.14 1.000 6.675	6.75 2.296 24.14 1.000 6.925	7.00 2.294 24.13 1.000 7.174	7.25 2.292 24.13 1.000 7.424	7.50 2.308 17.57 1.000 7.625	7.75 2.307 17.58 1.000 7.875	8.00 2.306 17.58 1.000 8.125	8.25 2.304 17.59 1.000 8.375	8.50 2.307 17.61 1.000 8.626
	8.75 2.305 17.64 1.000 8.877	9.00 2.304 17.70 1.000 9.127	9.25 2.301 17.79 1.000 9.379	9.50 2.296 22.39 1.000 9.674	9.75 2.293 22.80 1.000 9.928	10.00 2.289 23.57 1.000 10.186	(Ref.) : 10 , 1				
249	SEBACIC ACID -1 -2	20.7 44.9	4.530 5.380	C10H18O4 202.25	pHL RE t Z pHS	3.00 12.173 36.02 0.315 4.111	3.25 10.107 33.79 0.380 4.214	3.50 8.526 32.42 0.452 4.314			
	3.75 7.143 31.57 0.541 4.425	4.00 5.907 31.21 0.657 4.553	4.00 5.173 27.33 0.759 4.655	4.25 4.540 27.58 0.867 4.760	4.50 3.919 28.05 1.008 4.892	4.75 3.364 28.74 1.177 5.051	4.75 3.206 33.18 1.232 5.107	5.00 2.839 33.91 1.394 5.272	5.25 2.545 34.65 1.556 5.464	5.50 2.318 31.68 1.731 5.733	5.75 2.212 32.04 1.815 5.918
	6.00 2.132 32.33 1.883 6.135	6.25 2.079 32.52 1.930 6.368	6.50 2.026 43.90 1.963 6.665	6.75 2.008 43.94 1.978 6.902	7.00 1.996 43.95 1.988 7.147	7.25 1.987 43.93 1.993 7.395	7.50 2.020 32.77 1.996 7.606	7.75 2.016 32.78 1.998 7.854	8.00 2.013 32.79 1.999 8.104	8.25 2.009 32.79 1.999 8.354	8.50 2.014 32.83 2.000 8.605
	8.75 2.011 32.87 2.000 8.855	9.00 2.008 32.95 2.000 9.105	9.25 2.004 33.09 2.000 9.356	9.50 1.988 40.88 2.000 9.642	9.75 1.983 41.43 2.000 9.894	10.00 1.978 42.48 2.000 10.149	(Ref.) : 10 , 12				

250	SELENIC ACID				H ₂ SeO ₄		pH _L	3.00	3.25	3.50
	-1	41.0	0.000		144.97		RE	1.100	1.085	1.077
	-2	80.5	2.050				t	36.21	32.80	30.84
							Z	1.933	1.961	1.977
							pH _S	3.038	3.278	3.521
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
1.072	1.068	1.073	1.071	1.070	1.069	1.067	1.066	1.065	1.069	1.069
29.72	29.09	26.99	26.63	26.42	26.31	28.54	28.48	28.45	26.39	26.38
1.987	1.993	1.993	1.996	1.998	1.999	1.999	1.999	2.000	2.000	2.000
3.768	4.016	4.017	4.265	4.514	4.763	4.765	5.015	5.264	5.513	5.763
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
1.069	1.068	1.063	1.063	1.062	1.061	1.069	1.069	1.069	1.068	1.069
26.38	26.38	32.36	32.35	32.35	32.34	26.34	26.34	26.35	26.36	26.36
2.000	2.000	2.000	2.000	2.000	2.000	2.000	2.000	2.000	2.000	2.000
6.013	6.263	6.516	6.766	7.016	7.266	7.513	7.763	8.013	8.263	8.513
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 11 , 4				
1.069	1.068	1.068	1.064	1.063	1.062					
26.38	26.41	26.46	30.51	30.68	30.97					
2.000	2.000	2.000	2.000	2.000	2.000					
8.763	9.013	9.263	9.516	9.766	10.016					
251	SELENIOUS ACID				H ₂ SeO ₃		pH _L	3.00	3.25	3.50
	-1	41.2	2.620		128.97		RE	2.301	2.184	2.105
	-2	60.5	8.450				t	24.12	21.93	20.52
							Z	0.848	0.895	0.929
							pH _S	3.331	3.512	3.701
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
2.049	2.012	2.018	1.997	1.983	1.974	1.973	1.967	1.963	1.968	1.966
19.58	18.99	16.98	16.67	16.48	16.37	18.45	18.40	18.37	16.47	16.48
0.955	0.973	0.974	0.984	0.991	0.995	0.995	0.998	0.999	1.001	1.003
3.908	4.135	4.152	4.370	4.605	4.849	4.869	5.114	5.363	5.599	5.848
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
1.963	1.958	1.946	1.935	1.915	1.885	1.847	1.790	1.723	1.653	1.587
16.52	16.59	22.41	22.72	23.24	24.07	19.31	20.67	22.28	23.96	25.64
1.006	1.010	1.020	1.035	1.061	1.104	1.180	1.272	1.390	1.523	1.667
6.097	6.347	6.648	6.897	7.145	7.393	7.660	7.889	8.117	8.348	8.607
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 11				
1.541	1.506	1.482	1.460	1.449	1.442					
26.78	27.66	28.30	34.15	34.66	35.32					
1.772	1.854	1.912	1.950	1.971	1.984					
8.832	9.068	9.311	9.577	9.824	10.075					
252	SELENOCYANIC ACID				HSeCN		pH _L	3.00	3.25	3.50
	-1	67.1	0.000		105.99		RE	1.185	1.184	1.184
							t	19.40	17.41	16.27
							Z	0.999	1.000	1.000
							pH _S	3.052	3.293	3.534
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
1.184	1.184	1.185	1.184	1.184	1.184	1.184	1.184	1.184	1.185	1.185
15.62	15.24	14.05	13.84	13.72	13.66	14.91	14.87	14.85	13.69	13.69
1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
3.778	4.026	4.027	4.272	4.521	4.770	4.774	5.024	5.274	5.520	5.770
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
1.184	1.184	1.184	1.184	1.184	1.184	1.185	1.185	1.184	1.184	1.185
13.69	13.69	17.04	17.04	17.04	17.04	13.66	13.66	13.67	13.67	13.67
1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
6.020	6.270	6.529	6.779	7.029	7.279	7.520	7.770	8.020	8.270	8.520
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 2 , 12				
1.184	1.184	1.184	1.184	1.184	1.183					
13.69	13.71	13.74	16.02	16.12	16.31					
1.000	1.000	1.000	1.000	1.000	1.000					
8.770	9.020	9.270	9.528	9.778	10.029					

CHROMATOGRAPHIC DATA (1983)

D95

253	SORBIC ACID -1	33.4	4.770	C6H8O2 112.13	pHL RE t Z pHS	3.00 11.103 24.46 0.210 4.177	3.25 9.298 23.21 0.252 4.277	3.50 7.942 22.35 0.296 4.372		
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
6.758	5.696	4.888	4.413	3.933	3.494	3.356	3.074	2.847	2.671	2.595
21.68	21.18	18.45	18.30	18.16	18.06	20.61	20.56	20.52	18.15	18.14
0.349	0.415	0.489	0.543	0.611	0.690	0.719	0.787	0.852	0.915	0.943
4.475	4.596	4.722	4.815	4.935	5.085	5.146	5.304	5.493	5.766	5.949
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.538	2.499	2.465	2.454	2.446	2.440	2.457	2.455	2.453	2.450	2.453
18.13	18.12	25.06	25.06	25.05	25.05	18.09	18.09	18.10	18.11	18.13
0.964	0.979	0.990	0.994	0.997	0.998	0.999	0.999	1.000	1.000	1.000
6.165	6.400	6.709	6.947	7.193	7.441	7.639	7.888	8.137	8.387	8.639
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3				
2.451	2.449	2.446	2.439	2.436	2.432					
18.16	18.22	18.33	23.25	23.72	24.62					
1.000	1.000	1.000	1.000	1.000	1.000					
8.889	9.140	9.391	9.691	9.945	10.204					
254	SUBERIC ACID -1 -2	22.5 47.2	4.517 5.403	C8H14O4 174.20	pHL RE t Z pHS	3.00 11.348 33.79 0.311 4.095	3.25 9.435 31.79 0.375 4.199	3.50 7.974 30.58 0.446 4.299		
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
6.697	5.554	4.857	4.276	3.703	3.188	3.042	2.698	2.420	2.199	2.099
29.83	29.55	26.00	26.27	26.78	27.54	31.72	32.54	33.37	30.75	31.16
0.533	0.646	0.749	0.854	0.991	1.158	1.213	1.373	1.537	1.718	1.804
4.409	4.538	4.643	4.748	4.880	5.041	5.096	5.262	5.454	5.726	5.910
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.022	1.970	1.920	1.903	1.891	1.883	1.911	1.908	1.905	1.902	1.906
31.50	31.72	42.52	42.58	42.60	42.60	32.02	32.03	32.04	32.05	32.08
1.875	1.925	1.960	1.977	1.987	1.992	1.995	1.997	1.999	1.999	2.000
6.126	6.359	6.654	6.890	7.134	7.382	7.596	7.845	8.094	8.344	8.595
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 10 , 7				
1.903	1.900	1.897	1.883	1.879	1.874					
32.12	32.19	32.32	39.67	40.16	41.09					
2.000	2.000	2.000	2.000	2.000	2.000					
8.845	9.096	9.346	9.628	9.880	10.135					
255	SUCCINIC ACID (S) -1 -2	33.0 60.9	4.207 5.638	C4H6O4 118.09	pHL RE t Z pHS	3.00 6.629 25.62 0.363 3.914	3.25 5.619 24.26 0.430 4.025	3.50 4.856 23.41 0.500 4.133		
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
4.193	3.604	3.280	2.972	2.667	2.385	2.316	2.104	1.916	1.728	1.643
22.86	22.62	20.16	20.38	20.86	21.69	24.71	25.82	27.11	26.16	26.93
0.583	0.683	0.761	0.846	0.954	1.082	1.118	1.251	1.399	1.600	1.702
4.254	4.397	4.504	4.625	4.779	4.962	5.015	5.201	5.407	5.704	5.885
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
1.571	1.520	1.474	1.455	1.443	1.435	1.448	1.445	1.443	1.441	1.443
27.60	28.09	36.49	36.67	36.78	36.84	28.82	28.84	28.86	28.88	28.89
1.797	1.871	1.929	1.957	1.975	1.986	1.992	1.995	1.997	1.998	1.999
6.095	6.323	6.608	6.837	7.078	7.324	7.555	7.803	8.052	8.301	8.552
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 10 , 1				
1.441	1.440	1.439	1.431	1.429	1.426					
28.92	28.97	29.05	34.56	34.84	35.36					
2.000	2.000	2.000	2.000	2.000	2.000					
8.802	9.052	9.302	9.569	9.820	10.071					

256	SULFAMIC ACID (S)				HSO ₃ NH ₂			pHL	3.00	3.25	3.50
-1	50.3	-2.000			97.09			RE	1.594	1.594	1.594
								t	22.36	19.98	18.57
								Z	1.000	1.000	1.000
								pHS	3.147	3.373	3.600
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75	
1.594	1.593	1.597	1.597	1.596	1.596	1.595	1.595	1.594	1.598	1.597	
17.72	17.22	15.61	15.33	15.17	15.08	16.77	16.72	16.69	15.13	15.12	
1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	
3.835	4.078	4.080	4.318	4.563	4.811	4.824	5.073	5.322	5.562	5.812	
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50	
1.597	1.596	1.594	1.594	1.593	1.592	1.598	1.597	1.597	1.596	1.597	
15.12	15.12	19.64	19.64	19.64	19.64	15.09	15.09	15.10	15.11	15.11	
1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	
6.061	6.311	6.589	6.838	7.088	7.338	7.561	7.811	8.061	8.311	8.562	
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 10 , 12					
1.597	1.596	1.595	1.594	1.593	1.591						
15.13	15.16	15.21	18.32	18.50	18.84						
1.000	1.000	1.000	1.000	1.000	1.000						
8.812	9.062	9.313	9.586	9.837	10.090						
257	SULFANILIC ACID				C ₆ H ₇ N ₃ S			pHL	3.00	3.25	3.50
-1	33.7	3.227			173.19			RE	3.489	3.175	2.946
								t	26.44	24.32	22.89
								Z	0.685	0.754	0.814
								pHS	3.532	3.681	3.835
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75	
2.768	2.636	2.630	2.558	2.505	2.471	2.465	2.445	2.432	2.439	2.434	
21.88	21.20	18.67	18.34	18.11	17.98	20.55	20.47	20.43	18.07	18.05	
0.868	0.913	0.920	0.947	0.967	0.981	0.982	0.990	0.994	0.996	0.998	
4.011	4.214	4.252	4.443	4.661	4.896	4.925	5.164	5.410	5.642	5.888	
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50	
2.431	2.428	2.419	2.417	2.415	2.413	2.431	2.430	2.429	2.427	2.430	
18.05	18.04	24.92	24.91	24.91	24.90	18.01	18.01	18.02	18.03	18.05	
0.999	0.999	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	
6.136	6.385	6.688	6.938	7.188	7.437	7.635	7.885	8.135	8.385	8.637	
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 10 , 1					
2.428	2.426	2.423	2.417	2.413	2.409						
18.08	18.14	18.25	23.11	23.57	24.45						
1.000	1.000	1.000	1.000	1.000	1.000						
8.887	9.138	9.389	9.688	9.943	10.201						
258	m-SULFANILIC ACID				C ₆ H ₇ N ₃ S			pHL	3.00	3.25	3.50
-1	33.4	3.740			173.19			RE	4.748	4.158	3.717
								t	25.68	23.95	22.77
								Z	0.503	0.576	0.646
								pHS	3.717	3.844	3.971
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75	
3.350	3.049	2.973	2.815	2.684	2.589	2.576	2.521	2.486	2.480	2.468	
21.90	21.28	18.65	18.39	18.19	18.06	20.65	20.57	20.53	18.15	18.14	
0.719	0.792	0.818	0.866	0.909	0.943	0.947	0.968	0.981	0.989	0.994	
4.116	4.288	4.358	4.513	4.703	4.920	4.955	5.180	5.419	5.656	5.894	
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50	
2.460	2.454	2.443	2.441	2.439	2.436	2.454	2.453	2.452	2.450	2.453	
18.13	18.12	25.06	25.06	25.05	25.05	18.09	18.09	18.10	18.11	18.13	
0.996	0.998	0.999	0.999	1.000	1.000	1.000	1.000	1.000	1.000	1.000	
6.140	6.388	6.692	6.941	7.190	7.440	7.637	7.887	8.137	8.387	8.639	
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3					
2.451	2.449	2.446	2.439	2.436	2.432						
18.16	18.22	18.33	23.25	23.72	24.62						
1.000	1.000	1.000	1.000	1.000	1.000						
8.889	9.140	9.391	9.691	9.945	10.204						

CHROMATOGRAPHIC DATA (1983)

D97

259	o-SULFANILIC ACID			C6H7NO3S			pHL	3.00	3.25	3.50
	-1	33.4	7.488	173.19			RE	□□□□□	□□□□□	□□□□□
							t	23.80	22.79	22.08
							Z	0.011	0.013	0.016
							pHS	5.518	5.605	5.684
	3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50
	□□□□□	96.602	67.988	59.744	50.784	41.726	34.677	29.299	23.954	12.186
	21.52	21.08	18.34	18.25	18.17	18.09	20.52	20.49	20.47	18.18
	0.019	0.023	0.033	0.038	0.045	0.055	0.066	0.078	0.096	0.191
	5.767	5.861	6.019	6.077	6.151	6.240	6.326	6.405	6.501	6.893
	6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25
	9.820	8.393	4.981	4.626	4.196	3.748	3.156	2.989	2.822	2.685
	18.17	18.17	24.93	24.94	24.95	24.97	18.11	18.11	18.11	18.12
	0.238	0.280	0.477	0.515	0.569	0.639	0.769	0.814	0.863	0.909
	6.963	7.055	7.422	7.487	7.580	7.706	7.976	8.092	8.252	8.448
	8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3			
	2.531	2.495	2.473	2.453	2.443	2.436				
	18.17	18.23	18.34	23.26	23.72	24.63				
	0.967	0.981	0.989	0.994	0.997	0.998				
	8.915	9.152	9.397	9.695	9.947	10.205				
260	SULFURIC ACID			H2SO4			pHL	3.00	3.25	3.50
	-1	52.0	-3.000	98.07			RE	1.052	1.044	1.039
	-2	82.9	1.921				t	35.73	32.38	30.47
							Z	1.948	1.970	1.983
							pHS	3.020	3.266	3.513
	3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50
	1.036	1.034	1.038	1.037	1.036	1.036	1.034	1.033	1.032	1.036
	29.39	28.78	26.74	26.39	26.20	26.09	28.25	28.20	28.16	26.17
	1.990	1.994	1.994	1.997	1.998	1.999	1.999	1.999	2.000	2.000
	3.761	4.011	4.011	4.260	4.510	4.760	4.760	5.010	5.260	5.510
	6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25
	1.036	1.035	1.031	1.031	1.030	1.029	1.036	1.036	1.036	1.036
	26.16	26.16	31.95	31.94	31.94	31.93	26.11	26.12	26.13	26.13
	2.000	2.000	2.000	2.000	2.000	2.000	2.000	2.000	2.000	2.000
	6.010	6.260	6.511	6.761	7.011	7.261	7.510	7.760	8.010	8.260
	8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 1 , 4			
	1.036	1.035	1.035	1.031	1.030	1.030				
	26.15	26.19	26.24	30.16	30.31	30.59				
	2.000	2.000	2.000	2.000	2.000	1.999				
	8.760	9.010	9.260	9.511	9.761	10.011				
261	SULFUROUS ACID			H2SO3			pHL	3.00	3.25	3.50
	-1	51.8	1.780	82.07			RE	1.603	1.580	1.566
	-2	74.6	6.991				t	21.82	19.62	18.29
							Z	0.964	0.978	0.987
							pHS	3.165	3.382	3.603
	3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50
	1.557	1.551	1.555	1.551	1.548	1.544	1.544	1.539	1.531	1.523
	17.49	17.01	15.45	15.20	15.08	15.05	16.71	16.78	16.96	15.77
	0.993	0.997	0.997	1.000	1.003	1.008	1.008	1.015	1.028	1.050
	3.834	4.075	4.080	4.316	4.561	4.808	4.821	5.070	5.319	5.583
	6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25
	1.476	1.435	1.360	1.313	1.267	1.227	1.206	1.188	1.176	1.168
	17.18	18.41	25.97	27.63	29.24	30.63	25.49	26.05	26.42	26.65
	1.143	1.228	1.394	1.511	1.634	1.747	1.843	1.901	1.940	1.965
	6.078	6.324	6.664	6.866	7.082	7.309	7.566	7.791	8.031	8.276
	8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 2 , 7			
	1.162	1.160	1.159	1.153	1.152	1.151				
	26.89	26.97	27.05	31.46	31.66	32.01				
	1.989	1.994	1.996	1.998	1.999	1.999				
	8.774	9.023	9.273	9.529	9.779	10.030				

262	TAPS -1	25.0	8.300	C7H17NO6S 243.27				pHL RE t Z pHS	3.00 29.08 0.004 5.946	3.25 27.52 0.005 6.034	3.50 26.46 0.007 6.115
	3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
	25.63	25.01	21.36	21.23	21.11	21.01	24.24	91.527	73.907	36.649	33.079
	0.008	0.010	0.014	0.016	0.019	0.023	0.028	0.033	0.041	0.084	0.093
	6.200	6.296	6.438	6.499	6.575	6.666	6.748	6.829	6.925	7.252	7.301
	6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
	28.756	24.110	12.350	11.273	9.964	8.562	6.149	5.678	5.143	4.622	4.077
	21.16	21.17	30.19	30.21	30.24	30.29	21.21	21.22	21.24	21.27	21.34
	0.108	0.129	0.255	0.280	0.317	0.371	0.527	0.572	0.634	0.708	0.809
	7.369	7.455	7.816	7.870	7.947	8.049	8.320	8.398	8.509	8.654	8.894
	8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 11 , 9				
	3.833	3.640	3.507	3.394	3.348	3.314	N-Tris(hydroxymethyl)methyl-3-				
	21.40	21.51	21.72	28.69	29.72	31.90	aminopropanesulfonic acid				
	0.862	0.909	0.944	0.972	0.984	0.991					
	9.063	9.266	9.493	9.806	10.056	10.321					
263	TARTARIC ACID (S)							pHL	3.00	3.25	3.50
	-1	32.6						RE	2.907	2.590	2.336
	-2	60.7	3.036					t	29.86	28.25	27.62
			4.366					Z	0.878	0.997	1.120
								pHS	3.464	3.622	3.784
	3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
	2.114	1.921	1.883	1.758	1.653	1.575	1.565	1.515	1.482	1.474	1.462
	27.67	28.23	25.73	26.37	27.03	27.59	30.62	31.01	31.28	28.72	28.81
	1.257	1.407	1.452	1.577	1.698	1.800	1.808	1.878	1.927	1.958	1.975
	3.964	4.163	4.223	4.404	4.610	4.833	4.855	5.083	5.321	5.571	5.809
	6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
	1.455	1.450	1.439	1.437	1.434	1.432	1.449	1.448	1.447	1.445	1.447
	28.87	28.90	37.01	37.01	37.01	36.99	28.90	28.90	28.91	28.92	28.93
	1.986	1.992	1.996	1.998	1.999	1.999	2.000	2.000	2.000	2.000	2.000
	6.055	6.303	6.575	6.823	7.072	7.322	7.552	7.802	8.051	8.301	8.552
	8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 10 , 1				
	1.446	1.445	1.444	1.436	1.434	1.431					
	28.96	29.01	29.09	34.62	34.90	35.42					
	2.000	2.000	2.000	2.000	2.000	2.000					
	8.802	9.052	9.302	9.570	9.820	10.072					
264	TARTRONIC ACID							pHL	3.00	3.25	3.50
	-1	36.0						RE	2.374	2.253	2.141
	-2	68.3	2.366					t	27.42	25.20	24.17
			4.735					Z	0.957	1.013	1.073
								pHS	3.327	3.520	3.718
	3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
	2.021	1.886	1.858	1.733	1.611	1.505	1.491	1.414	1.358	1.326	1.306
	24.01	24.55	22.42	23.31	24.32	25.27	27.97	28.71	29.25	27.25	27.45
	1.148	1.246	1.277	1.386	1.511	1.638	1.653	1.758	1.843	1.908	1.943
	3.929	4.152	4.212	4.411	4.621	4.840	4.868	5.085	5.315	5.576	5.802
	6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
	1.292	1.283	1.272	1.268	1.266	1.264	1.276	1.275	1.274	1.273	1.274
	27.58	27.66	34.83	34.85	34.86	34.86	27.72	27.73	27.73	27.74	27.75
	1.967	1.981	1.989	1.994	1.997	1.998	1.999	1.999	2.000	2.000	2.000
	6.042	6.288	6.554	6.800	7.048	7.298	7.535	7.785	8.034	8.284	8.535
	8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 10 , 1				
	1.274	1.273	1.272	1.266	1.264	1.263					
	27.77	27.81	27.88	32.73	32.96	33.37					
	2.000	2.000	2.000	2.000	2.000	2.000					
	8.785	9.035	9.285	9.546	9.796	10.047					

265	TEREPHTHALIC ACID				C8H6O4		pHL	3.00	3.25	3.50
	-1	28.0			166.13		RE	4.141	3.556	3.106
	-2	53.0	3.540				t	32.28	30.69	29.94
			4.460				Z	0.716	0.843	0.976
							pHS	3.642	3.772	3.904
	3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50
	2.722	2.399	2.314	2.130	1.975	1.860	1.844	1.773	1.726	1.715
	29.78	30.14	27.23	27.77	28.40	28.98	32.49	32.92	33.21	30.27
	1.129	1.302	1.369	1.505	1.643	1.762	1.774	1.856	1.913	1.950
	4.054	4.229	4.299	4.458	4.649	4.864	4.892	5.113	5.349	5.596
	6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25
	1.688	1.681	1.665	1.662	1.659	1.656	1.680	1.678	1.677	1.675
	30.43	30.46	39.86	39.86	39.85	39.83	30.48	30.48	30.48	30.49
	1.983	1.990	1.995	1.997	1.998	1.999	1.999	2.000	2.000	2.000
	6.077	6.325	6.605	6.854	7.103	7.352	7.574	7.823	8.073	8.323
	8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 11 , 7			
	1.676	1.674	1.672	1.661	1.658	1.654				
	30.54	30.60	30.70	37.15	37.52	38.23				
	2.000	2.000	2.000	2.000	2.000	2.000				
	8.824	9.074	9.325	9.600	9.851	10.104				
266	THIOACETIC ACID				C2H4OS		pHL	3.00	3.25	3.50
	-1	43.7	3.337		76.11		RE	2.925	2.617	2.393
							t	22.16	20.66	19.65
							Z	0.623	0.697	0.764
							pHS	3.521	3.665	3.812
	3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50
	2.216	2.080	2.059	1.987	1.933	1.897	1.893	1.873	1.860	1.859
	18.91	18.41	16.46	16.22	16.06	15.95	17.91	17.86	17.82	16.02
	0.827	0.882	0.895	0.928	0.955	0.973	0.974	0.985	0.991	0.995
	3.979	4.174	4.227	4.408	4.620	4.851	4.874	5.110	5.354	5.595
	6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25
	1.852	1.849	1.845	1.844	1.843	1.842	1.850	1.850	1.849	1.848
	16.00	16.00	21.24	21.24	21.24	21.24	15.97	15.97	15.98	15.99
	0.998	0.999	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
	6.086	6.335	6.622	6.871	7.121	7.371	7.585	7.835	8.085	8.335
	8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3			
	1.849	1.848	1.846	1.844	1.842	1.840				
	16.02	16.05	16.12	19.75	20.00	20.46				
	1.000	1.000	1.000	1.000	1.000	1.000				
	8.836	9.086	9.337	9.619	9.871	10.125				
267	THIOCYANIC ACID				HSCN		pHL	3.00	3.25	3.50
	-1	68.5	0.848		59.09		RE	1.166	1.163	1.161
							t	19.19	17.25	16.13
							Z	0.994	0.997	0.998
							pHS	3.050	3.290	3.531
	3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50
	1.160	1.159	1.160	1.160	1.160	1.159	1.159	1.159	1.159	1.160
	15.49	15.12	13.96	13.75	13.63	13.57	14.79	14.76	14.74	13.61
	0.999	0.999	0.999	1.000	1.000	1.000	1.000	1.000	1.000	1.000
	3.775	4.023	4.024	4.270	4.518	4.767	4.771	5.021	5.271	5.518
	6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25
	1.159	1.159	1.159	1.159	1.159	1.159	1.160	1.160	1.160	1.159
	13.60	13.60	16.88	16.88	16.88	16.88	13.58	13.58	13.58	13.59
	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
	6.017	6.267	6.525	6.775	7.025	7.275	7.517	7.767	8.017	8.267
	8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 2			
	1.159	1.159	1.159	1.159	1.159	1.158				
	13.60	13.62	13.65	15.88	15.98	16.16				
	1.000	1.000	1.000	1.000	1.000	1.000				
	8.767	9.017	9.268	9.524	9.775	10.025				

D100

CHROMATOGRAPHIC DATA (1983)

268	THIOCYANOACETIC ACID -1	40.6	2.585	C3H3NOS 101.10	pHL RE t Z pHS	3.00 2.311 24.38 0.858 3.328	3.25 2.200 22.14 0.902 3.511	3.50 2.126 20.70 0.934 3.701		
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
2.074	2.039	2.045	2.025	2.013	2.004	2.003	1.997	1.994	2.000	1.999
19.74	19.14	17.09	16.78	16.58	16.47	18.59	18.52	18.49	16.54	16.53
0.958	0.975	0.976	0.985	0.991	0.995	0.995	0.997	0.998	0.999	1.000
3.910	4.137	4.153	4.372	4.607	4.851	4.872	5.117	5.366	5.601	5.849
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
1.997	1.996	1.991	1.990	1.989	1.988	1.999	1.998	1.997	1.996	1.998
16.52	16.52	22.18	22.18	22.17	22.17	16.49	16.49	16.50	16.51	16.52
1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
6.098	6.348	6.639	6.889	7.139	7.389	7.598	7.848	8.098	8.348	8.599
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3				
1.997	1.995	1.994	1.990	1.988	1.986					
16.54	16.58	16.66	20.60	20.89	21.44					
1.000	1.000	1.000	1.000	1.000	1.000					
8.850	9.100	9.351	9.637	9.890	10.145					
269	THIOGLYCOLIC ACID -1	38.6	3.545	C2H4O2S 92.11	pHL RE t Z pHS	3.00 3.687 23.62 0.560 3.619	3.25 3.260 22.04 0.635 3.753	3.50 2.945 20.97 0.704 3.888		
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
2.687	2.483	2.441	2.332	2.247	2.188	2.181	2.147	2.125	2.123	2.115
20.18	19.63	17.39	17.14	16.96	16.85	19.07	19.01	18.97	16.92	16.91
0.773	0.839	0.858	0.899	0.934	0.960	0.962	0.977	0.987	0.992	0.996
4.044	4.226	4.289	4.456	4.657	4.881	4.910	5.141	5.383	5.622	5.863
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.110	2.107	2.100	2.099	2.097	2.095	2.108	2.107	2.106	2.104	2.106
16.91	16.90	22.87	22.86	22.86	22.86	16.87	16.87	16.88	16.89	16.90
0.998	0.999	0.999	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
6.110	6.358	6.653	6.902	7.152	7.402	7.608	7.858	8.108	8.358	8.609
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3				
2.105	2.104	2.102	2.098	2.095	2.093					
16.93	16.97	17.06	21.22	21.56	22.18					
1.000	1.000	1.000	1.000	1.000	1.000					
8.859	9.110	9.361	9.650	9.903	10.159					
270	THIOSULFURIC ACID -1 -2	44.0 88.1	0.600 1.720	H2S2O3 114.13	pHL RE t Z pHS	3.00 0.986 35.11 1.965 3.001	3.25 0.979 31.74 1.980 3.250	3.50 0.974 29.85 1.989 3.501		
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
0.972	0.970	0.973	0.972	0.972	0.971	0.969	0.969	0.968	0.971	0.971
28.79	28.20	26.29	25.95	25.77	25.66	27.69	27.63	27.60	25.73	25.72
1.994	1.996	1.996	1.998	1.999	1.999	1.999	2.000	2.000	2.000	2.000
3.751	4.002	4.002	4.252	4.503	4.753	4.752	5.002	5.252	5.503	5.753
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
0.971	0.971	0.967	0.967	0.966	0.965	0.971	0.971	0.971	0.971	0.971
25.72	25.72	31.15	31.14	31.14	31.14	25.68	25.68	25.69	25.70	25.70
2.000	2.000	2.000	2.000	2.000	2.000	2.000	2.000	2.000	2.000	2.000
6.003	6.253	6.501	6.751	7.001	7.251	7.503	7.753	8.003	8.253	8.503
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 2 , 4				
0.971	0.971	0.971	0.967	0.967	0.966					
25.72	25.75	25.79	29.46	29.60	29.86					
2.000	2.000	2.000	2.000	2.000	2.000					
8.753	9.003	9.253	9.501	9.751	10.001					

CHROMATOGRAPHIC DATA (1983)

D101

271	p-TOLUENESULFONIC -1	31.1	ACID -2.000		C7H8O3S 172.20		pHL RE t Z pHS	3.00 2.630 29.77 1.000 3.320	3.25 2.630 26.46 1.000 3.525	3.50 2.630 24.40 1.000 3.732	
	3.75 2.630 23.08 1.000 3.953	4.00 2.628 22.26 1.000 4.188	4.00 2.643 19.58 1.000 4.192	4.25 2.643 19.15 1.000 4.418	4.50 2.641 18.88 1.000 4.657	4.75 2.639 18.73 1.000 4.902	4.75 2.636 21.53 1.000 4.930	5.00 2.633 21.45 1.000 5.177	5.25 2.631 21.40 1.000 5.426	5.50 2.646 18.83 1.000 5.654	5.75 2.645 18.82 1.000 5.903
	6.00 2.643 18.81 1.000 6.152	6.25 2.640 18.80 1.000 6.402	6.50 2.630 26.28 1.000 6.710	6.75 2.629 26.28 1.000 6.960	7.00 2.627 26.27 1.000 7.210	7.25 2.624 26.26 1.000 7.459	7.50 2.646 18.77 1.000 7.652	7.75 2.645 18.77 1.000 7.902	8.00 2.643 18.78 1.000 8.152	8.25 2.641 18.79 1.000 8.402	8.50 2.645 18.81 1.000 8.654
	8.75 2.643 18.85 1.000 8.905	9.00 2.640 18.92 1.000 9.156	9.25 2.637 19.05 1.000 9.407	9.50 2.628 24.39 1.000 9.712	9.75 2.624 24.95 1.000 9.968	10.00 2.619 26.05 1.000 10.229	(Ref.) : 3 , 12				
272	m-TOLUIC -1	29.1	ACID 4.272		C8H8O2 136.15		pHL RE t Z pHS	3.00 8.002 27.47 0.340 3.962	3.25 6.822 25.75 0.400 4.072	3.50 5.932 24.58 0.462 4.180	
	3.75 5.166 23.69 0.532 4.301	4.00 4.499 23.05 0.614 4.445	4.00 4.188 19.96 0.666 4.540	4.25 3.854 19.73 0.725 4.661	4.50 3.545 19.53 0.791 4.816	4.75 3.292 19.40 0.854 5.003	4.75 3.240 22.38 0.867 5.051	5.00 3.085 22.30 0.911 5.250	5.25 2.977 22.26 0.945 5.472	5.50 2.928 19.52 0.968 5.716	5.75 2.891 19.50 0.981 5.938
	6.00 2.866 19.49 0.989 6.176	6.25 2.850 19.48 0.994 6.421	6.50 2.828 27.51 0.997 6.734	6.75 2.823 27.50 0.998 6.981	7.00 2.819 27.49 0.999 7.229	7.25 2.815 27.48 0.999 7.479	7.50 2.840 19.45 1.000 7.668	7.75 2.839 19.45 1.000 7.917	8.00 2.836 19.46 1.000 8.167	8.25 2.833 19.47 1.000 8.417	8.50 2.838 19.50 1.000 8.670
	8.75 2.835 19.54 1.000 8.920	9.00 2.832 19.63 1.000 9.171	9.25 2.828 19.77 1.000 9.423	9.50 2.817 25.54 1.000 9.733	9.75 2.813 26.21 1.000 9.990	10.00 2.806 27.55 1.000 10.253	(Ref.) : 11 , 1				
273	o-TOLUIC -1	29.1	ACID 3.908		C8H8O2 136.15		pHL RE t Z pHS	3.00 6.018 28.03 0.456 3.806	3.25 5.231 26.09 0.527 3.927	3.50 4.638 24.78 0.596 4.048	
	3.75 4.138 23.80 0.670 4.186	4.00 3.719 23.10 0.748 4.350	4.00 3.598 20.06 0.780 4.423	4.25 3.378 19.77 0.832 4.569	4.50 3.190 19.55 0.883 4.750	4.75 3.051 19.40 0.925 4.960	4.75 3.027 22.39 0.931 5.000	5.00 2.945 22.31 0.957 5.221	5.25 2.891 22.26 0.975 5.456	5.50 2.880 19.52 0.985 5.691	5.75 2.861 19.50 0.991 5.927
	6.00 2.849 19.49 0.995 6.171	6.25 2.840 19.48 0.997 6.419	6.50 2.823 27.51 0.999 6.732	6.75 2.820 27.50 0.999 6.980	7.00 2.817 27.49 1.000 7.229	7.25 2.814 27.48 1.000 7.479	7.50 2.840 19.45 1.000 7.667	7.75 2.838 19.45 1.000 7.917	8.00 2.836 19.46 1.000 8.167	8.25 2.833 19.47 1.000 8.417	8.50 2.837 19.50 1.000 8.670
	8.75 2.835 19.54 1.000 8.920	9.00 2.832 19.63 1.000 9.171	9.25 2.828 19.77 1.000 9.423	9.50 2.817 25.54 1.000 9.733	9.75 2.813 26.21 1.000 9.990	10.00 2.806 27.55 1.000 10.253	(Ref.) : 11 , 1				

274	p-TOLUIC ACID -1	29.1	4.373	C8H8O2 136.15			pHL RE t Z pHS	3.00 8.717 27.34 0.312 4.007	3.25 7.400 25.67 0.368 4.115	3.50 6.406 24.53 0.427 4.220
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
5.547	4.794	4.406	4.035	3.685	3.392	3.327	3.145	3.015	2.949	2.904
23.67	23.03	19.93	19.71	19.53	19.40	22.37	22.30	22.26	19.52	19.50
0.495	0.575	0.631	0.691	0.759	0.827	0.843	0.893	0.933	0.961	0.976
4.337	4.476	4.576	4.692	4.839	5.019	5.070	5.262	5.479	5.727	5.943
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.874	2.855	2.830	2.824	2.819	2.815	2.841	2.839	2.836	2.833	2.838
19.49	19.48	27.50	27.50	27.49	27.48	19.45	19.45	19.46	19.47	19.50
0.986	0.992	0.996	0.998	0.999	0.999	1.000	1.000	1.000	1.000	1.000
6.178	6.422	6.736	6.981	7.230	7.479	7.668	7.917	8.167	8.417	8.670
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 11 , 1				
2.835	2.832	2.828	2.817	2.813	2.806					
19.54	19.63	19.77	25.54	26.21	27.55					
1.000	1.000	1.000	1.000	1.000	1.000					
8.920	9.171	9.423	9.733	9.990	10.253					
275	TRIBROMOACETIC ACID -1	36.5	1.000	C2HBr3O2 296.74			pHL RE t Z pHS	3.00 2.235 26.84 0.995 3.264	3.25 2.231 23.91 0.997 3.474	3.50 2.228 22.11 0.998 3.686
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
2.226	2.225	2.234	2.233	2.232	2.230	2.228	2.227	2.225	2.235	2.234
20.97	20.28	18.02	17.64	17.42	17.29	19.66	19.59	19.55	17.37	17.36
0.999	0.999	0.999	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
3.911	4.148	4.152	4.382	4.623	4.869	4.892	5.139	5.388	5.620	5.869
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.233	2.231	2.224	2.224	2.222	2.221	2.235	2.234	2.233	2.231	2.234
17.35	17.35	23.68	23.67	23.67	23.66	17.32	17.32	17.33	17.34	17.35
1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
6.119	6.368	6.666	6.916	7.166	7.416	7.619	7.869	8.119	8.369	8.620
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3 , 12				
2.233	2.231	2.229	2.223	2.221	2.218					
17.38	17.43	17.52	21.96	22.34	23.06					
1.000	1.000	1.000	1.000	1.000	1.000					
8.870	9.121	9.372	9.665	9.919	10.176					
276	TRICHLOROACETIC ACID -1	37.9	0.635	C2HCl3O2 163.39			pHL RE t Z pHS	3.00 2.144 26.25 0.998 3.249	3.25 2.142 23.39 0.999 3.461	3.50 2.141 21.63 0.999 3.675
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
2.139	2.138	2.147	2.146	2.145	2.144	2.142	2.141	2.139	2.148	2.147
20.53	19.86	17.69	17.33	17.11	16.99	19.27	19.20	19.16	17.07	17.06
1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
3.901	4.139	4.143	4.373	4.615	4.862	4.883	5.131	5.380	5.613	5.862
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.146	2.145	2.139	2.138	2.137	2.135	2.148	2.148	2.146	2.145	2.147
17.05	17.05	23.13	23.12	23.12	23.12	17.01	17.02	17.02	17.03	17.04
1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
6.111	6.361	6.657	6.906	7.156	7.406	7.611	7.861	8.111	8.361	8.613
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 2 , 7				
2.146	2.144	2.143	2.138	2.136	2.133					
17.07	17.12	17.21	21.46	21.81	22.46					
1.000	1.000	1.000	1.000	1.000	1.000					
8.863	9.113	9.364	9.655	9.909	10.165					

CHROMATOGRAPHIC DATA (1983)

D103

277	TRICHLOROACRYLIC -1	ACID 33.4	0.000	C3HC1302 175.40	pHL RE t Z pHS	3.00 2.442 28.41 1.000 3.294	3.25 2.441 25.28 1.000 3.501	3.50 2.441 23.33 1.000 3.711		
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
2.440	2.439	2.452	2.451	2.450	2.448	2.445	2.443	2.441	2.454	2.453
22.09	21.34	18.85	18.44	18.20	18.06	20.66	20.58	20.53	18.15	18.14
1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
3.934	4.169	4.173	4.401	4.641	4.887	4.913	5.160	5.409	5.639	5.887
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.451	2.449	2.441	2.439	2.438	2.436	2.454	2.453	2.451	2.450	2.453
18.13	18.12	25.06	25.06	25.05	25.05	18.09	18.09	18.10	18.11	18.13
1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
6.137	6.387	6.690	6.940	7.190	7.440	7.637	7.887	8.137	8.387	8.639
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3 , 12				
2.451	2.449	2.446	2.439	2.436	2.432					
18.16	18.22	18.33	23.25	23.72	24.62					
1.000	1.000	1.000	1.000	1.000	1.000					
8.889	9.140	9.391	9.691	9.945	10.204					
278	TRICHLOROLACTIC -1	ACID 34.2	2.347	C3H3C1303 193.41	pHL RE t Z pHS	3.00 2.595 27.48 0.915 3.343	3.25 2.519 24.70 0.943 3.533	3.50 2.469 22.93 0.963 3.728		
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
2.434	2.411	2.422	2.409	2.400	2.394	2.391	2.387	2.383	2.395	2.393
21.77	21.04	18.60	18.22	17.98	17.84	20.38	20.30	20.26	17.94	17.92
0.977	0.986	0.987	0.992	0.995	0.997	0.997	0.999	0.999	1.000	1.000
3.941	4.171	4.181	4.402	4.639	4.884	4.909	5.155	5.403	5.634	5.883
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.391	2.389	2.381	2.380	2.378	2.376	2.394	2.393	2.391	2.389	2.392
17.91	17.91	24.68	24.68	24.67	24.67	17.87	17.88	17.88	17.90	17.91
1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
6.132	6.382	6.684	6.934	7.183	7.433	7.632	7.882	8.132	8.382	8.634
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3				
2.391	2.389	2.386	2.380	2.377	2.372					
17.95	18.00	18.11	22.89	23.33	24.18					
1.000	1.000	1.000	1.000	1.000	1.000					
8.884	9.135	9.386	9.684	9.938	10.197					
279	TRIMETAPHOSPHORIC -1 -2 -3	ACID 28.9 57.8 87.6	-2.000 -1.000 2.050	H3P309 239.94	pHL RE t Z pHS	3.00 1.088 54.45 2.946 3.029	3.25 1.081 49.14 2.969 3.275	3.50 1.075 46.13 2.982 3.522		
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
1.072	1.069	1.080	1.078	1.077	1.076	1.070	1.068	1.067	1.078	1.077
44.42	43.45	40.37	39.82	39.50	39.33	42.66	42.56	42.49	39.47	39.45
2.990	2.994	2.994	2.997	2.998	2.999	2.999	2.999	3.000	3.000	3.000
3.771	4.020	4.021	4.269	4.519	4.769	4.770	5.020	5.270	5.519	5.769
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
1.076	1.075	1.064	1.063	1.061	1.059	1.078	1.078	1.077	1.076	1.077
39.44	39.43	48.34	48.32	48.29	48.26	39.39	39.39	39.39	39.40	39.41
3.000	3.000	3.000	3.000	3.000	3.000	3.000	3.000	3.000	3.000	3.000
6.019	6.269	6.522	6.772	7.021	7.271	7.519	7.769	8.019	8.269	8.519
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 11 , 4				
1.077	1.075	1.074	1.064	1.062	1.061					
39.44	39.48	39.55	45.57	45.80	46.24					
3.000	3.000	3.000	3.000	3.000	3.000					
8.769	9.019	9.269	9.521	9.771	10.021					

280	TRIMETHYLACETIC ACID (S)				C5H10O2		pHL	3.00	3.25	3.50
-1	31.8	5.037			102.13		RE	15.015	12.497	10.607
							t	25.11	23.83	22.96
							Z	0.163	0.196	0.231
							pHS	4.309	4.406	4.497
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
8.954	7.461	6.196	5.549	4.883	4.256	4.014	3.610	3.265	2.947	2.833
22.28	21.76	18.88	18.74	18.61	18.51	21.18	21.13	21.10	18.61	18.60
0.275	0.331	0.403	0.451	0.515	0.593	0.630	0.702	0.779	0.872	0.908
4.596	4.710	4.841	4.926	5.034	5.170	5.237	5.378	5.550	5.834	5.995
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.738	2.672	2.615	2.595	2.581	2.573	2.590	2.587	2.584	2.581	2.584
18.59	18.59	25.89	25.88	25.88	25.87	18.55	18.55	18.56	18.57	18.59
0.940	0.963	0.982	0.989	0.994	0.997	0.998	0.999	0.999	1.000	1.000
6.195	6.420	6.735	6.966	7.209	7.456	7.651	7.899	8.148	8.398	8.650
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3 , 7				
2.582	2.579	2.576	2.568	2.564	2.559					
18.63	18.70	18.82	24.02	24.55	25.59					
1.000	1.000	1.000	1.000	1.000	1.000					
8.900	9.151	9.402	9.705	9.961	10.221					
281	TRIMETHYLACRYLIC ACID				C6H10O2		pHL	3.00	3.25	3.50
-1	30.3	4.420			114.14		RE	8.773	7.431	6.419
							t	26.51	24.94	23.88
							Z	0.297	0.351	0.408
							pHS	4.023	4.131	4.234
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
5.543	4.772	4.344	3.971	3.614	3.310	3.240	3.051	2.914	2.838	2.791
23.06	22.47	19.48	19.28	19.11	18.98	21.83	21.76	21.72	19.09	19.08
0.474	0.553	0.613	0.673	0.742	0.812	0.829	0.882	0.925	0.956	0.973
4.349	4.484	4.590	4.701	4.845	5.020	5.072	5.259	5.473	5.725	5.937
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.759	2.738	2.714	2.708	2.703	2.699	2.722	2.720	2.718	2.715	2.719
19.07	19.06	26.75	26.74	26.74	26.73	19.03	19.03	19.04	19.05	19.07
0.984	0.991	0.995	0.997	0.999	0.999	0.999	1.000	1.000	1.000	1.000
6.170	6.413	6.725	6.970	7.218	7.467	7.659	7.908	8.158	8.408	8.660
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3				
2.717	2.714	2.710	2.701	2.696	2.691					
19.12	19.19	19.32	24.83	25.43	26.62					
1.000	1.000	1.000	1.000	1.000	1.000					
8.911	9.162	9.413	9.720	9.976	10.238					
282	2,4,6-TRIMETHYLBENZOIC ACID				C10H12O2		pHL	3.00	3.25	3.50
-1	24.7	3.437			164.20		RE	5.123	4.630	4.265
							t	32.68	29.90	28.02
							Z	0.643	0.713	0.777
							pHS	3.664	3.804	3.948
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
3.971	3.746	3.739	3.613	3.518	3.455	3.440	3.403	3.380	3.395	3.386
26.67	25.75	22.23	21.80	21.50	21.31	24.88	24.77	24.70	21.45	21.43
0.837	0.889	0.898	0.930	0.956	0.974	0.976	0.986	0.992	0.995	0.997
4.115	4.309	4.348	4.529	4.740	4.971	5.009	5.246	5.490	5.716	5.960
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
3.380	3.374	3.351	3.349	3.345	3.342	3.381	3.379	3.375	3.371	3.378
21.41	21.40	30.93	30.92	30.91	30.90	21.37	21.37	21.38	21.40	21.44
0.998	0.999	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
6.208	6.457	6.779	7.028	7.278	7.528	7.707	7.957	8.207	8.457	8.710
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 11 , 7				
3.374	3.370	3.364	3.346	3.339	3.328					
21.50	21.62	21.83	28.87	29.94	32.19					
1.000	1.000	1.000	1.000	1.000	1.000					
8.961	9.212	9.465	9.790	10.050	10.320					

CHROMATOGRAPHIC DATA (1983)

D105

283	TRIMETILPYRUVIC ACID						C6H10O3				
	-1	29.2	1.469				130.14	pHL	3.00	3.25	3.50
								RE	2.842	2.830	2.823
								t	30.97	27.56	25.41
								Z	0.988	0.992	0.995
								pHS	3.351	3.551	3.754
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75	
2.817	2.813	2.830	2.828	2.825	2.822	2.817	2.814	2.811	2.829	2.828	
24.01	23.15	20.28	19.82	19.54	19.37	22.37	22.27	22.22	19.48	19.46	
0.997	0.998	0.998	0.999	0.999	1.000	1.000	1.000	1.000	1.000	1.000	
3.972	4.205	4.210	4.434	4.672	4.917	4.946	5.193	5.442	5.669	5.917	
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50	
2.825	2.822	2.809	2.808	2.806	2.803	2.829	2.828	2.826	2.823	2.827	
19.45	19.45	27.44	27.43	27.43	27.42	19.41	19.42	19.42	19.44	19.46	
1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	
6.166	6.416	6.728	6.978	7.228	7.477	7.666	7.916	8.166	8.416	8.669	
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3					
2.825	2.822	2.818	2.807	2.803	2.796						
19.51	19.59	19.73	25.48	26.15	27.47						
1.000	1.000	1.000	1.000	1.000	1.000						
8.919	9.170	9.422	9.732	9.989	10.252						
284	VALERIC ACID						C5H10O2				
	-1	31.6	4.842				102.13	pHL	3.00	3.25	3.50
								RE	12.475	10.427	8.888
								t	25.35	24.02	23.10
								Z	0.198	0.237	0.279
								pHS	4.216	4.316	4.410
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75	
7.545	6.338	5.413	4.873	4.326	3.823	3.657	3.333	3.070	2.860	2.771	
22.39	21.86	18.97	18.82	18.68	18.57	21.27	21.22	21.18	18.67	18.66	
0.330	0.395	0.467	0.520	0.588	0.667	0.698	0.768	0.836	0.906	0.936	
4.512	4.632	4.757	4.848	4.966	5.112	5.175	5.329	5.516	5.788	5.967	
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50	
2.703	2.657	2.616	2.602	2.593	2.586	2.606	2.603	2.600	2.598	2.601	
18.65	18.65	26.00	25.99	25.99	25.98	18.61	18.61	18.62	18.63	18.65	
0.960	0.976	0.988	0.993	0.996	0.998	0.999	0.999	1.000	1.000	1.000	
6.180	6.413	6.726	6.963	7.209	7.456	7.651	7.900	8.149	8.399	8.651	
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 10 , 1					
2.599	2.596	2.593	2.585	2.581	2.576						
18.69	18.76	18.88	24.12	24.66	25.72						
1.000	1.000	1.000	1.000	1.000	1.000						
8.901	9.152	9.404	9.707	9.963	10.223						
285	VANILLIC ACID						C8H8O4				
	-1	27.1	4.523				168.15	pHL	3.00	3.25	3.50
								RE	10.559	8.918	7.677
								t	28.62	26.85	25.64
								Z	0.276	0.328	0.382
								pHS	4.084	4.189	4.291
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75	
6.601	5.651	5.121	4.658	4.213	3.830	3.734	3.494	3.317	3.218	3.156	
24.73	24.06	20.72	20.49	20.31	20.17	23.37	23.30	23.25	20.31	20.29	
0.446	0.523	0.583	0.643	0.714	0.788	0.808	0.865	0.913	0.949	0.968	
4.404	4.537	4.640	4.749	4.889	5.060	5.114	5.298	5.508	5.758	5.967	
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50	
3.114	3.087	3.054	3.046	3.039	3.034	3.064	3.062	3.059	3.055	3.060	
20.28	20.27	28.92	28.91	28.90	28.89	20.24	20.24	20.25	20.27	20.30	
0.981	0.989	0.995	0.997	0.998	0.999	0.999	1.000	1.000	1.000	1.000	
6.198	6.440	6.759	7.003	7.251	7.500	7.685	7.934	8.184	8.434	8.687	
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 11 , 2					
3.057	3.054	3.049	3.036	3.030	3.022						
20.35	20.44	20.61	26.90	27.72	29.38						
1.000	1.000	1.000	1.000	1.000	1.000						
8.937	9.188	9.441	9.757	10.015	10.281						

286	VINYLAACETIC -1	ACID 38.6	4.342	C4H6O2 86.09	pHL RE t Z pHS	3.00 6.648 22.58 0.306 3.962	3.25 5.639 21.44 0.361 4.070	3.50 4.882 20.66 0.418 4.173		
3.75	4.00	4.00	4.25	4.50	4.75	5.00	5.25	5.50	5.75	
4.227	3.649	3.297	3.026	2.765	2.542	2.494	2.355	2.254	2.193	
20.04	19.58	17.24	17.09	16.95	16.85	19.06	18.97	16.92	16.91	
0.485	0.563	0.629	0.686	0.753	0.821	0.837	0.888	0.928	0.959	
4.287	4.423	4.538	4.648	4.791	4.967	5.016	5.202	5.416	5.675	
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	
2.136	2.121	2.107	2.103	2.099	2.097	2.108	2.107	2.106	2.104	
16.91	16.90	22.87	22.86	22.86	22.86	16.87	16.87	16.88	16.89	
0.985	0.991	0.996	0.998	0.999	0.999	1.000	1.000	1.000	1.000	
6.120	6.363	6.660	6.905	7.153	7.402	7.609	7.858	8.108	8.358	
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3 , 7				
2.105	2.104	2.102	2.098	2.095	2.093					
16.93	16.97	17.06	21.22	21.56	22.18					
1.000	1.000	1.000	1.000	1.000	1.000					
8.859	9.110	9.361	9.650	9.903	10.159					
287	VINYLGOLIC -1	ACID 35.4	3.319	C4H6O3 102.09	pHL RE t Z pHS	3.00 3.499 25.41 0.648 3.552	3.25 3.155 23.48 0.720 3.696	3.50 2.903 22.18 0.784 3.845		
3.75	4.00	4.00	4.25	4.50	4.75	4.75	5.00	5.25	5.50	5.75
2.704	2.554	2.540	2.458	2.397	2.357	2.352	2.328	2.314	2.318	2.313
21.24	20.61	18.18	17.88	17.68	17.55	19.99	19.92	19.88	17.63	17.62
0.844	0.895	0.904	0.935	0.960	0.976	0.977	0.987	0.992	0.996	0.997
4.016	4.213	4.257	4.442	4.656	4.889	4.917	5.154	5.399	5.634	5.878
6.00	6.25	6.50	6.75	7.00	7.25	7.50	7.75	8.00	8.25	8.50
2.309	2.306	2.298	2.296	2.295	2.293	2.308	2.307	2.306	2.304	2.307
17.61	17.61	24.14	24.14	24.13	24.13	17.57	17.58	17.58	17.59	17.61
0.999	0.999	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
6.126	6.375	6.675	6.925	7.174	7.424	7.625	7.875	8.125	8.375	8.626
8.75	9.00	9.25	9.50	9.75	10.00	(Ref.) : 3				
2.305	2.304	2.301	2.296	2.293	2.289					
17.64	17.70	17.79	22.39	22.80	23.57					
1.000	1.000	1.000	1.000	1.000	1.000					
8.877	9.127	9.379	9.674	9.928	10.186					

REFERENCES TO TABLE IV

- 1 R. A. Robinson and R. H. Stokes, *Electrolyte Solutions*, Butterworths, London, 2nd ed., 1959.
- 2 Landholt-Bornstein, *Zahlenwerte und Funktionen*, 6 Aufl. Bd. II, Teil 7, Springer, Berlin, 1960. Tokyo, 1966.
- 3 *International Critical Tables of Numerical Data, Physics, Chemistry and Technology*, Vol. VI, McGraw-Hill, New York, London, 1933.
- 4 L. G. Sillen and A. E. Martell (Editors), *Stability Constants of Metal-Ion Complexes*, Special publication No. 17, The Chemical Society, London, 1964.
- 5 Cyril Long (Editor), *Biochemist's Handbook*, E. & F. N. Spon Ltd., London, 1961.
- 6 R. C. Weast (Editor), *Handbook of Chemistry and Physics*, Chemical Rubber Co., Cleveland, OH, 52nd ed., 1971.
- 7 S. Nagai (Editor), *Kagaku Binran*, Chemical Society of Japan, Maruzen Publ., Tokyo, 1966.
- 8 G. Charlot, *Les Reactions Chimiques en Solution*, Masson, Paris, 1969.
- 9 T. Imamura and K. Saito, *Kagaku No Ryoiki*, 30, No. 2 (1966) 79.
- 10 T. Hirokawa, M. Nishino and Y. Kiso, *J. Chromatogr.*, 252 (1982) 49; m_0 and pK_a were recalculated.
- 11 Unpublished data obtained in our laboratory.
- 12 Assumed values.

PUBLICATION SCHEDULE FOR 1983

Journal of Chromatography (incorporating *Chromatographic Reviews*) and *Journal of Chromatography, Biomedical Applications*

MONTH	D 1982	J	F	M	A	M	J	The publication schedule for further issues will be published later.
Journal of Chromatography	252 253/1 253/2	254 255 256/1	256/2 256/3 257/1	257/2 258 259/1	259/2 259/3 260/1	260/2 261/1 261/2	261/3 262 263	
Chromatographic Reviews					271/1		271/2	
Biomedical Applications		272/1	272/2	273/1	273/2	274	275/1	

INFORMATION FOR AUTHORS

(Detailed *Instructions to Authors* were published in Vol. 244, No. 2, pp. 401-404. A free reprint can be obtained by application to the publisher.)

Types of Contributions. The following types of papers are published in the *Journal of Chromatography* and the section on *Biomedical Applications*: Regular research papers (Full-length papers), Short communications and Notes. Short communications are preliminary announcements of important new developments and will, whenever possible, be published with maximum speed. Notes are usually descriptions of short investigations and reflect the same quality of research as Full-length papers, but should preferably not exceed four printed pages. For review articles, see page 2 of cover under Submission of Papers.

Submission. Every paper must be accompanied by a letter from the senior author, stating that he is submitting the paper for publication in the *Journal of Chromatography*. Please do not send a letter signed by the director of the institute or the professor unless he is one of the authors.

Manuscripts. Manuscripts should be typed in double spacing on consecutively numbered pages of uniform size. The manuscript should be preceded by a sheet of manuscript paper carrying the title of the paper and the name and full postal address of the person to whom the proofs are to be sent. Authors of papers in French or German are requested to supply an English translation of the title of the paper. As a rule, papers should be divided into sections, headed by a caption (e.g., Summary, Introduction, Experimental, Results, Discussion, etc.). All illustrations, photographs, tables, etc., should be on separate sheets.

Introduction. Every paper must have a concise introduction mentioning what has been done before on the topic described, and stating clearly what is new in the paper now submitted.

Summary. Full-length papers and Review articles should have a summary of 50-100 words which clearly and briefly indicates what is new, different and significant. In the case of French or German articles an additional summary in English, headed by an English translation of the title, should also be provided. (Short communications and Notes are published without a summary.)

Illustrations. The figures should be submitted in a form suitable for reproduction, drawn in Indian ink on drawing or tracing paper. Each illustration should have a legend, all the legends being typed (with double spacing) together on a separate sheet. If structures are given in the text, the original drawings should be supplied. Coloured illustrations are reproduced at the author's expense, the cost being determined by the number of pages and by the number of colours needed. The written permission of the author and publisher must be obtained for the use of any figure already published. Its source must be indicated in the legend.

References. References should be numbered in the order in which they are cited in the text, and listed in numerical sequence on a separate sheet at the end of the article. Please check a recent issue for the layout of the reference list. Abbreviations for the titles of journals should follow the system used by *Chemical Abstracts*. Articles not yet published should be given as "in press", "submitted for publication", "in preparation" or "personal communication".

Proofs. One set of proofs will be sent to the author to be carefully checked for printer's errors. Corrections must be restricted to instances in which the proof is at variance with the manuscript. "Extra corrections" will be inserted at the author's expense.

Reprints. Fifty reprints of Full-length papers, Short communications and Notes will be supplied free of charge. Additional reprints can be ordered by the authors. An order form containing price quotations will be sent to the authors together with the proofs of their article.

News. News releases of new products and developments, and information leaflets of meetings should be addressed to: The Editor of the News Section, *Journal of Chromatography*/*Journal of Chromatography, Biomedical Applications*, Elsevier Science Publishers B.V., P.O. Box 330, 1000 AH Amsterdam, The Netherlands.

Advertisements. Advertisement rates are available from the publisher on request. The Editors of the journal accept no responsibility for the contents of the advertisements.

ELECTROPHORESIS

A Survey of Techniques and Applications

edited by Z. DEYL, Czechoslovak Academy of Sciences, Prague

JOURNAL OF CHROMATOGRAPHY LIBRARY, 18

PART A: TECHNIQUES

Z. DEYL (editor)
F.M. EVERAERTS, Z. PRUSÍK and
P.J. SVENDSEN (co-editors)

"... provides a sound, state-of-the-art survey of its subject".

— Chemistry in Britain

"... the editors have set out to bring everything together into a coherent whole... they have succeeded remarkably well... the book is bound to be well liked and appreciated by readers".

— Journal of Chromatography

This first part deals with the principles, theory and instrumentation of modern electromigration methods. Both standard procedures and newer developments are discussed and hints are included to help the reader overcome difficulties frequently arising from the lack of suitable equipment. Adequate theoretical background of the individual techniques is given and a theoretical approach to the deteriorative processes is presented to facilitate further development of a particular technique and its application to a special problem. In each chapter practical realisations of different techniques are described and examples are presented to demonstrate the limits of each method.

CONTENTS:

Introduction. Chapters: 1. Theory of electromigration processes (J. Vacík). 2. Classification of electromigration methods (J. Vacík). 3. Evaluation of the results of electrophoretic separations (J. Vacík). 4. Molecular size and shape in electrophoresis (Z. Deyl). 5. Zone electrophoresis (except gel-type techniques and immunoelectrophoresis) (W. Ostrowski). 6. Gel-type techniques (Z. Hrkál). 7. Quantitative immunoelectrophoresis (P.J. Svendsen). 8. Moving boundary electrophoresis in narrow-bore tubes (F.M. Everaerts and J.L. Beckers). 9. Isoelectric focusing (N. Catsimpoolas). 10. Analytical isotachopheresis (J. Vacík and F.M. Everaerts). 11. Continuous flow-through electrophoresis (Z. Prusík). 12. Continuous flow deviation electrophoresis (A. Kolín). 13. Preparative electrophoresis in gel media (Z. Hrkál). 14. Preparative electrophoresis in columns (P.J. Svendsen). 15. Preparative isoelectric focusing (P. Blanický). 16. Preparative isotachopheresis (P.J. Svendsen). 17. Preparative isotachopheresis on the micro scale (L. Arlinger). List of frequently occurring symbols. Subject Index.

1979 xvi + 390 pp. US \$83.00/Dfl. 195.00
ISBN 0-444-41721-4

PART B: APPLICATIONS

Z. DEYL (editor)
A. CHRAMBACH, F.M. EVERAERTS and
Z. PRUSÍK (co-editors)

Part B is an exhaustive survey of the present status of the application of electrophoretic techniques to many diverse compounds. Those categories of compounds most suited to these separations, such as proteins and peptides, are dealt with in detail, while the perspectives of the applications of these techniques to other categories of compounds less commonly electrophoresed are given. Special attention is paid to naturally occurring mixtures of compounds and their treatment. This is the first attempt to cover the field on such a broad scale and the book will be valuable to separation chemists, pharmacologists, organic chemists and those involved in biomedical research.

CONTENTS: 1. Alcohols and phenolic compounds (Z. Deyl). 2. Aldehydes and ketones (Z. Deyl). 3. Carbohydrates (Z. Deyl). 4. Carboxylic acids (F.M. Everaerts). 5. Steroids and steroid conjugates (Z. Deyl). 6. Amines (Z. Deyl). 7. Amino acids and their derivatives (Z. Deyl). 8. Peptides and structural analysis of proteins (Z. Prusík). 9. Gel electrophoresis and electrofocusing of proteins (edited by A. Chrambach). Usefulness of second-generation gel electrophoretic tools in protein fractionation (A. Chrambach). Membrane proteins, native (L.M. Hjelmeland). Membrane proteins, denatured (H. Baumann, D. Doyle). Protein membrane receptors (U. Lang). Steroid receptors (S. Ben-Or). Cell surface antigens (R.A. Reisfeld, M.A. Pellegrino). Lysosomal glycosidases and sulphatases (A.L. Fluharty). Haemocyanins (M. Brenowitz et al.). Human haemoglobins (A.B. Schneider, A.N. Schechter). Isoelectric focusing of immunoglobulins (M.H. Freedman). Contractile and cytoskeletal proteins (P. Rubenstein). Proteins of connective tissue (Z. Deyl, M. Horáková). Microtubular proteins (K.F. Sullivan, L. Wilson). Protein hormones (A.D. Rogol). Electrophoresis of plasma proteins: a contemporary clinical approach (M. Engliš). Allergens (H. Baer, M.C. Anderson). 10. Glycoproteins and glycopeptides (affinity electrophoresis) (T.C. Bøg-Hansen, J. Hau.). 11. Lipoproteins (H. Peeters). 12. Lipopolysaccharides (P.F. Coleman, O. Gabriel). 13. Electrophoretic examination of enzymes (W. Ostrowski). 14. Nucleotides, nucleosides, nitrogenous constituents of nucleic acids (S. Zdražil). 15. Nucleic acids (S. Zdražil). 16. Alkaloids (Z. Deyl). 17. Vitamins (Z. Deyl). 18. Antibiotics (V. Betina). 19. Dyes and pigments (Z. Deyl). 20. Inorganic compounds (F.M. Everaerts, Th. P.E.M. Verheggen). Contents of "Electrophoresis, Part A: Techniques". Subject Index. Index of compounds separated.

1982 xiii + 462 pp. US \$95.75/Dfl. 225.00
ISBN 0-444-42114-9



ELSEVIER
P.O. Box 211, Amsterdam
The Netherlands
52 Vanderbilt Avenue
New York, NY 10017, U.S.A.