


ISSN 0221-9673

VOL. 451 DECEMBER 31, 1988

COMPLETE IN ONE ISSUE

CUMULATIVE  
AUTHOR AND SUBJECT INDEXES  
VOLUMES 401-450



EDITOR, Michael Lederer (Switzerland)  
ASSOCIATE EDITORS, R. W. Giese (Boston, MA.),  
J. K. Haken (Kensington, N.S.W.),  
K. Macek (Prague), L. R. Snyder (Orinda, CA)  
EDITOR, SYMPOSIUM VOLUMES, F. Heftmann (Orinda, CA)  
EDITORIAL BOARD

W. A. Aue (Halifax)  
V. G. Berezkin (Moscow)  
V. Betina (Bratislava)  
A. Bevenue (Belmont, CA)  
P. Bucek (Brno)  
P. Boulanger (Lille)  
A. A. Bouiton (Saskatoon)  
G. P. Cartoni (Rome)  
S. Dilli (Kensington, N.S.W.)  
L. Fishbein (Washington, DC)  
A. Frigerio (Milan)  
C. W. Gehrke (Columbia, MO)  
E. Gil-av (Rehovot)  
G. Guiochon (Knoxville, TN)  
I. A. Hais (Hradec Králové)  
S. Hjertén (Uppsala)  
E. C. Horning (Houston, TX)  
Cs. Horváth (New Haven, CT)  
J. F. K. Huber (Vienna)  
A. T. James (Harrold)  
J. Janák (Brno)  
E. sz. Kováts (Lausanne)  
K. A. Kraus (Oak Ridge, TN)  
A. Liberti (Rome)  
H. M. McNair (Blacksburg, VA)  
Y. Marcus (Jerusalem)  
G. B. Marini-Bettolo (Pinerolo)  
A. J. P. Martin (Cambridge)  
Č. Michalec (Prague)  
R. Neher (Basel)  
G. Nickless (Bristol)  
N. A. Parris (Wilmington, DE)  
R. L. Patience (Sunbury-on-Thames)  
P. G. Rizzotti (Milan)  
O. Sauerbion (Göteborg)  
R. Schwarzenbach (Dübendorf)  
A. Taitis (Houston, TX)

EDITORS, BIBLIOGRAPHY SECTION  
Z. Deyl (Prague), J. Janák (Brno), V. Schwarz (Prague), K. Macek (Prague)

ELSEVIER

## JOURNAL OF CHROMATOGRAPHY

**Scope.** The *Journal of Chromatography* publishes papers on all aspects of chromatography, electrophoresis and related methods. Contributions consist mainly of research papers dealing with chromatographic theory, instrumental development and their applications. The section *Biomedical Applications*, which is under separate editorship, deals with the following aspects: developments in and applications of chromatographic and electrophoretic techniques related to clinical diagnosis or alterations during medical treatment; screening and profiling of body fluids or tissues with special reference to metabolic disorders; results from basic medical research with direct consequences in clinical practice; drug level monitoring and pharmacokinetic studies; clinical toxicology; analytical studies in occupational medicine.

**Submission of Papers.** Papers in English, French and German may be submitted, in three copies. Manuscripts should be submitted to: The Editor of *Journal of Chromatography*, P.O. Box 681, 1000 AR Amsterdam, The Netherlands, or to: The Editor of *Journal of Chromatography, Biomedical Applications*, P.O. Box 681, 1000 AR Amsterdam, The Netherlands. Review articles are invited or proposed by letter to the Editors. An outline of the proposed review should first be forwarded to the Editors for preliminary discussion prior to preparation. Submission of an article is understood to imply that the article is original and unpublished and is not being considered for publication elsewhere. For copyright regulations, see below.

**Subscription Orders.** Subscription orders should be sent to: Elsevier Science Publishers B.V., P.O. Box 211, 1000 AE Amsterdam, The Netherlands, Tel. 5803 911, Telex 18582 ESPA NL. The *Journal of Chromatography* and the *Biomedical Applications* section can be subscribed to separately.

**Publication.** The *Journal of Chromatography* (incl. *Biomedical Applications*) has 37 volumes in 1989. The subscription prices for 1989 are:

*J. Chromatogr.* + *Biomed. Appl.* (Vols. 461–497):

Dfl. 6475.00 plus Dfl. 999.00 (p.p.h.) (total ca. US\$ 3933.75)

*J. Chromatogr.* only (Vols. 461–486):

Dfl. 5200.00 plus Dfl. 702.00 (p.p.h.) (total ca. US\$ 3106.25)

*Biomed. Appl.* only (Vols. 487–497):

Dfl. 2200.00 plus Dfl. 297.00 (p.p.h.) (total ca. US\$ 1314.25).

Our p.p.h. (postage, package and handling) charge includes surface delivery of all issues, except to subscribers in Argentina, Australia, Brazil, Canada, China, Hong Kong, India, Israel, Malaysia, Mexico, New Zealand, Pakistan, Singapore, South Africa, South Korea, Taiwan, Thailand and the U.S.A. who receive all issues by air delivery (S.A.L. — Surface Air Lifted) at no extra cost. For Japan, air delivery requires 50% additional charge; for all other countries airmail and S.A.L. charges are available upon request. Back volumes of the *Journal of Chromatography* (Vols. 1–460) are available at Dfl. 230.00 (plus postage). Claims for missing issues will be honoured, free of charge, within three months after publication of the issue. Customers in the U.S.A. and Canada wishing information on this and other Elsevier journals, please contact Journal Information Center, Elsevier Science Publishing Co. Inc., 655 Avenue of the Americas, New York, NY 10010. Tel. (212) 989-5800.

**Abstracts/Contents Lists** published in Analytical Abstracts, ASCA, Biochemical Abstracts, Biological Abstracts, Chemical Abstracts, Chemical Titles, Chromatography Abstracts, Current Contents/Physical, Chemical & Earth Sciences, Current Contents/Life Sciences, Deep-Sea Research/Part B: Oceanographic Literature Review, Excerpta Medica, Index Medicus, Mass Spectrometry Bulletin, PASCAL-CNRS, Referativnyi Zhurnal and Science Citation Index.

**See inside back cover** for Publication Schedule, Information for Authors and information on Advertisements.

---

© ELSEVIER SCIENCE PUBLISHERS B.V. — 1989

0021-9673/89/003.00

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, without the prior written permission of the publisher, Elsevier Science Publishers B.V., P.O. Box 330, 1000 AH Amsterdam, The Netherlands.

Upon acceptance of an article by the journal, the author(s) will be asked to transfer copyright of the article to the publisher. The transfer will ensure the widest possible dissemination of information.

Submission of an article for publication entails the authors' irrevocable and exclusive authorization of the publisher to collect any sums or considerations for copying or reproduction payable by third parties (as mentioned in article 17 paragraph 2 of the Dutch Copyright Act of 1912 and the Royal Decree of June 20, 1974 (S. 351) pursuant to article 16 b of the Dutch Copyright Act of 1912) and/or to act in or out of Court in connection therewith.

**Special regulations for readers in the U.S.A.** This journal has been registered with the Copyright Clearance Center, Inc. Consent is given for copying of articles for personal or internal use, or for the personal use of specific clients. This consent is given on the condition that the copier pays through the Center the per-copy fee stated in the code on the first page of each article for copying beyond that permitted by Sections 107 or 108 of the U.S. Copyright Law. The appropriate fee should be forwarded with a copy of the first page of the article to the Copyright Clearance Center Inc., 27 Congress Street, Salem, MA 01970, U.S.A. If no code appears in an article, the author has not given broad consent to copy and permission to copy must be obtained directly from the author. All articles published prior to 1980 may be copied for a per-copy fee of US\$ 2.25, also payable through the Center. This consent does not extend to other kinds of copying, such as for general distribution, resale, advertising and promotion purposes, or for creating new collective works. Special written permission must be obtained from the publisher for such copying.

No responsibility is assumed by the Publisher for any injury and/or damage to persons or property as a matter of products liability, negligence or otherwise, or from any use or operation of any methods, products, instructions or ideas contained in the materials herein. Because of rapid advances in the medical sciences, the Publisher recommends that independent verification of diagnoses and drug dosages should be made. Although all advertising material is expected to conform to ethical (medical) standards, inclusion in this publication does not constitute a guarantee or endorsement of the quality or value of such product or of the claims made of it by its manufacturer.

This issue is printed on acid-free paper.

Printed in The Netherlands

---

# JOURNAL OF CHROMATOGRAPHY

## Cumulative Author and Subject Indexes

*An invaluable tool for locating published work, the CUMULATIVE AUTHOR AND SUBJECT INDEXES make the vast amount of information in the journal more easily accessible.*

*Supplied automatically to subscribers to the JOURNAL OF CHROMATOGRAPHY, the Indexes are also available separately for desk use.*

**Vols. 1 - 50 (1972)**  
US\$ 87.50 / Dfl. 175.00

**Vols. 51 - 100 (1975)**  
US\$ 112.50 / Dfl. 225.00

**Vols. 101 - 150**  
(Published as *J. Chromatogr.*  
*Vol. 293*, 1984)  
US\$ 127.50 / Dfl. 255.00

**Vols. 151 - 250**  
(Published as *J. Chromatogr.*  
*Vol. 263*, 1983)  
US\$ 217.50 / Dfl. 435.00

**Vols. 251 - 350**  
(Published as *J. Chromatogr.*  
*Vol. 453*, 1988)  
US\$ 226.50 / Dfl. 453.00

**Vols. 351 - 400**  
(Published as *J. Chromatogr.*  
*Vol. 401*, 1987)  
US\$ 147.50 / Dfl. 295.00

PRICES QUOTED INCLUDE POSTAGE



### Elsevier Science Publishers

Back Volumes Journal Department  
P.O. Box 211, 1000 AE Amsterdam, The Netherlands

# ANALYTICA CHIMICA ACTA

## Cumulative Author, Subject and Technique Indexes

*An invaluable tool for locating published work, the CUMULATIVE AUTHOR, SUBJECT AND TECHNIQUE INDEXES make the vast amount of information in the journal more accessible.*

*Supplied automatically to subscribers to ANALYTICA CHIMICA ACTA, the Indexes are also available separately for desk use.*

**Vols. 1 - 100**  
(Published as *Analytica Chimica Acta* Vol. 137, 1982)  
US\$ 155.00 / Dfl. 310.00

**Vols. 101 - 150**  
(Published as *Analytica Chimica Acta* Vol. 168, 1985)  
US\$ 120.00 / Dfl. 240.00

**Vols. 151 - 200**  
(Published as *Analytica Chimica Acta* Vol. 203, No. 2, 1988)  
US\$ 78.00 / Dfl. 156.00

**Vols. 201 - 210**  
(Published as *Analytica Chimica Acta* Vol. 210, No. 2, 1989)

PRICES QUOTED INCLUDE  
POSTAGE



**Elsevier Science Publishers**

Back Volumes Journal Department  
P.O. Box 211, 1000 AE Amsterdam, The Netherlands

# JOURNAL *of* CHROMATOGRAPHY

INTERNATIONAL JOURNAL ON CHROMATOGRAPHY,  
ELECTROPHORESIS AND RELATED METHODS

EDITOR

MICHAEL LEDERER (Switzerland)

ASSOCIATE EDITORS

R. W. GIESE (Boston, MA), J. K. HAKEN (Kensington, N.S.W.),  
K. MACEK (Prague), L. R. SNYDER (Orinda, CA)

EDITOR, SYMPOSIUM VOLUMES

E. HEFTMANN (Orinda)

EDITORIAL BOARD

W. A. Aue (Halifax), V. G. Berezkin (Moscow), V. Betina (Bratislava), A. Bevenue (Belmont, CA), P. Boček (Brno), P. Boulanger (Lille), A. A. Boulton (Saskatoon), G. P. Cartoni (Rome), S. Dilli (Kensington, N.S.W.), L. Fishbein (Washington, DC), A. Frigerio (Milan), C. W. Gehrke (Columbia, MO), E. Gil-Av (Rehovot), G. Guiochon (Knoxville, TN), I. M. Hais (Hradec Králové), S. Hjertén (Uppsala), E. C. Horning (Houston, TX), Cs. Horváth (New Haven, CT), J. F. K. Huber (Vienna), A. T. James (Harrold), J. Janák (Brno), E. sz. Kováts (Lausanne), K. A. Kraus (Oak Ridge, TN), A. Liberti (Rome), H. M. McNair (Blacksburg, VA), Y. Marcus (Jerusalem), G. B. Marini-Bettolo (Rome), A. J. P. Martin (Cambridge), Č. Michalec (Prague), R. Neher (Basel), G. Nickless (Bristol), N. A. Parris (Wilmington, DE), R. L. Patience (Sunbury-on-Thames), P. G. Righetti (Milan), O. Samuelson (Göteborg), R. Schwarzenbach (Dü-bendorf), A. Zlatkis (Houston, TX)

EDITORS, BIBLIOGRAPHY SECTION

Z. Deyl (Prague), J. Janák (Brno), V. Schwarz (Prague), K. Macek (Prague)



ELSEVIER

AMSTERDAM — OXFORD — NEW YORK — TOKYO

---

*J. Chromatogr.*, Vol. 451 (1988)

ห้องสมุด.จุฬาลงกรณ์มหาวิทยาลัย

๒๐ ก.ค. ๒๕๓๒

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, without the prior written permission of the publisher, Elsevier Science Publishers B.V., P.O. Box 330, 1000 AH Amsterdam, The Netherlands.

Upon acceptance of an article by the journal, the author(s) will be asked to transfer copyright of the article to the publisher. The transfer will ensure the widest possible dissemination of information.

Submission of an article for publication entails the authors' irrevocable and exclusive authorization of the publisher to collect any sums or considerations for copying or reproduction payable by third parties (as mentioned in article 17 paragraph 2 of the Dutch Copyright Act of 1912 and the Royal Decree of June 20, 1974 (S. 351) pursuant to article 16 b of the Dutch Copyright Act of 1912) and/or to act in or out of Court in connection therewith.

**Special regulations for readers in the U.S.A.** This journal has been registered with the Copyright Clearance Center, Inc. Consent is given for copying of articles for personal or internal use, or for the personal use of specific clients. This consent is given on the condition that the copier pays through the Center the per-copy fee stated in the code on the first page of each article for copying beyond that permitted by Sections 107 or 108 of the U.S. Copyright Law. The appropriate fee should be forwarded with a copy of the first page of the article to the Copyright Clearance Center, Inc., 27 Congress Street, Salem, MA 01970, U.S.A. If no code appears in an article, the author has not given broad consent to copy and permission to copy must be obtained directly from the author. All articles published prior to 1980 may be copied for a per-copy fee of US\$ 2.25, also payable through the Center. This consent does not extend to other kinds of copying, such as for general distribution, resale, advertising and promotion purposes, or for creating new collective works. Special written permission must be obtained from the publisher for such copying.

No responsibility is assumed by the Publisher for any injury and/or damage to persons or property as a matter of products liability, negligence or otherwise, or from any use or operation of any methods, products, instructions or ideas contained in the materials herein. Because of rapid advances in the medical sciences, the Publisher recommends that independent verification of diagnoses and drug dosages should be made.

Although all advertising material is expected to conform to ethical (medical) standards, inclusion in this publication does not constitute a guarantee or endorsement of the quality or value of such product or of the claims made of it by its manufacturer.

This issue is printed on acid-free paper.

SPECIAL VOLUME

*Cumulative Author and Subject Indexes*  
*Volumes 401–450 (1987–1988)*





JOURNAL OF CHROMATOGRAPHY

VOL. 451 (1988)



## AUTHOR INDEX

## VOLUMES 401–450

## A

- Aarons, L., see Adedoyin, A. 416(1987)160
- Abbiati, A., see Guiso, G. 413(1987)363
- Abbott, F.S., see Chan, G.L.-Y. 417(1987)295
- Abdalla, T., see Greene, S.V. 417(1987)121
- Abdallah, M.A., see Demange, P. 438(1988)291
- Abdennebi, E.H., see Singh, A.K. 404(1987)223
- Abe, I., Tsujioka, H. and Wasa, T.  
Solvent extraction clean-up for pre-treatment  
in amino acid analysis by GC. Application to  
age estimation from the D/L ratio of aspartic  
acid in human dentine 449(1988)165
- Abe, J.-I., see Koizumi, K. 437(1988)47
- Abe, S., see Yamada, K. 424(1988)367
- Abe, T., see Ogawa, K. 426(1988)188
- Abel, S., Krauss, G.-J. and Glund, K.  
Single-step HPLC analysis of nucleoside  
monophosphates for ribonuclease specificity  
446(1988)187
- Abels, J., see Lindemans, J. 432(1988)103
- Ábermann, M., see Szederkényi, F. 446(1988)253
- Abernethy, D.R., see Montamat, S.C.  
415(1987)203
- Abián, J., see Cárdenas, A. 426(1988)83
- Abidi, S.L.  
Chiral-phase HPLC of rotenoid racemates  
404(1987)133
- Abidi, S.L.  
HPLC separation of subcomponents of  
antimycin A 447(1988)65
- Abou-Khalil, S., Abou-Khalil, W.H., Masoud,  
A.N. and Yunis, A.A.  
HPLC determination of chloramphenicol and  
four analogues using reductive and oxidative  
electrochemical and UV detection  
417(1987)111
- Abou-Khalil, W.H., see Abou-Khalil, S.  
417(1987)111
- Abraham, F., see Haken, J.K. 440(1988)329
- Abraham, M.H., Buist, G.J., Grellier, P.L.,  
McGill, R.A., Doherty, R.M., Kamlet, M.J.,  
Taft, R.W. and Maroldo, S.G.  
Solubility properties in polymers and  
biological media. II. A new method for the  
characterisation of the adsorption of gases  
and vapours on solids 409(1987)15
- Abramson, R.K., see Smith, C.S. 423(1987)207
- Abumrad, N.N., see Brown, L.L. 426(1988)370
- Aburatani, R., see Okamoto, Y. 448(1988)454
- Abusteit, E.O., see Parker, C.E. 438(1988)359
- Aceves, M., Grimalt, J., Albaigés, J., Broto, F.,  
Comellas, L. and Gassiot, M.  
Analysis of hydrocarbons in aquatic  
sediments. II. Evaluation of common  
preparative procedures for petroleum and  
chlorinated hydrocarbons 436(1988)503
- Achari, R., Drissel, D., Thomas, D., Shin, K. and  
Look, Z.  
Analysis of esmolol in human blood by HPLC  
and its application to pharmacokinetic studies  
424(1988)430
- Acheson, C.M., see Coppella, S.J. 402(1987)189
- Achtert, G., see Dahmen, W. 427(1988)157
- Achtert, G., see Gielsdorf, W. 419(1987)351
- Achtert, G., see Gielsdorf, W. 419(1987)357
- Ackermann, R., see Kaiser, G. 419(1987)123
- Ackermann, T., see Rockstroh, H. 442(1988)401
- Ackermans, M.T., see Gladdines, M.M.  
431(1988)317
- Acree, Jr., W.E.  
Thermochemical investigations of GLC. IV.  
Partition coefficients of solutes on binary  
mixtures containing two complexing solvents  
402(1987)41
- Adachi, K., Kim, J.Y. and Asakura, T.  
Effects of differences in charge and  
hydrophobicity of surface amino acids of  
hemoglobins on high-performance GPC  
428(1988)247
- Adachi, K., see Suzuki, N. 408(1987)181
- Adachi, K., see Yamaga, N. 422(1987)25

- Adachi, K. and Asakura, T.  
Separation of asymmetrical hybrid hemoglobins by hydrophobic interaction chromatography 419(1987)303
- Adachi, S., see Hashimoto, K. 448(1988)241
- Adachi, T., see Yoshida, S. 430(1988)156
- Adam, A., see Souvannavong, V. 440(1988)95
- Adam, M., see Macek, J. 414(1987)156
- Adam, M., see Macek, J. 434(1988)423
- Adam, M., see Štol, M. 419(1987)308
- Adamcová, E., see Knox, J.H. 447(1988)13
- Adams, P.C., Roberts, F.D., Powell, L.W. and Halliday, J.W.  
Chromatographic detergent exchange in the preparation of membrane receptor protein 427(1988)341
- Adams, R.F., see Murray, K.E. 431(1988)143
- Adams, R.W., see Griffin, R.M.E. 445(1988)441
- Adaya Godoy, A., see Guevara, O.L. 403(1987)350
- Adedoyin, A., Aarons, L. and Houston, J.B.  
HPLC determination of  $\beta$ -naphthoflavone in biological fluids 416(1987)160
- Ader, J.C., see Dobrev, A.A. 445(1988)268
- Aderjan, R., see Bogusz, M. 435(1988)43
- Adinarayana, M., Singh, U.S. and Dwivedi, T.S.  
A biochromatographic technique for the quantitative estimation of triphenyltin fungicides 435(1988)210
- Adlard, M.W., see Shah, A.J. 424(1988)325
- Aducci, P., see Marra, M. 440(1988)47
- Advenier, C., see Naline, E. 419(1987)177
- Advenier, C., see Spreux-Varoquaux, O. 416(1987)311
- Aerts, M.M.L., Beek, W.M.J. and Brinkman, U.A.T.  
Monitoring of veterinary drug residues by a combination of continuous flow techniques and column-switching HPLC. I. Sulphonamides in egg, meat and milk using post-column derivatization with dimethylaminobenzaldehyde 435(1988)97
- Aetake, A., see Yoshida, T. 445(1988)175
- Agarwal, G.S., see Ramachandran, P.K. 426(1988)239
- Ageta, T., Mizobuchi, N. and Kodama, H.  
Isotachophoretic analysis for the determination of urinary  $\gamma$ -carboxyglutamic acid 419(1987)334
- Aggarwal, S.K., see Bradshaw, J.S. 405(1987)169
- Aggazzotti, G., Predieri, G., Fantuzzi, G. and Benedetti, A.  
Headspace GC analysis for determining low levels of chloroform in human plasma 416(1987)125
- Agostini, O., Manzini, S. and Fedi, M.  
Determination of 1,3-dimethyl-7-isobutylxanthine in guinea pig plasma by HPLC 433(1988)288
- Agoston, S., see Biessels, P.T.M. 421(1987)392
- Agrapart, M., see Jaureguiberry, G. 440(1988)385
- Agrawal, O.P., see Kharbade, B.V. 439(1988)430
- Agris, P.F., see Guenther, R.H. 444(1988)79
- Aguilar, M.B., see Huberman, A. 443(1988)337
- Aguilar, M.I., Hart, S.J. and Calder, I.C.  
Complete separation of urinary metabolites of paracetamol and substituted paracetamols by reversed-phase ion-pair HPLC 426(1988)315
- Aguilar, M.I., see Hearn, M.T.W. 435(1988)271
- Aguilar, M.I., see Hearn, M.T.W. 438(1988)197
- Aguilar, M.I., see Hearn, M.T.W. 443(1988)97
- Ahel, M., see Czuczwa, J. 403(1987)233
- Aherne, F.X., see Fenton, M. 410(1987)480
- Ahmad, S., Khan, H.M., Mahdi, A.A., Kumar, H. and Khan, N.  
*Entamoeba histolytica* antigens as possible vaccinoagens? A short review 440(1988)467
- Airaud, C.B., Gayte-Sorbier, A., Ajuoulat, P. and Mercier, V.  
TLC of amine antioxidants and antiozonants used in elastomers 437(1988)59
- Akai, H., see Oi, T. 450(1988)343
- Akane, A., see Matsubara, K. 424(1988)49
- Akbari, A., Gordon, B.J., Bush, P.B. and Moore, J.N.  
Determination of xylazine in blood components using HPLC 426(1988)207
- Åkesson, B., Forslind, K. and Wollheim, F.  
Analysis of homogentisic acid in body fluids by HPLC 413(1987)233
- Akiyama, S., see Nakashima, K. 414(1987)11
- Akporhonor, E.E., Le Vent, S. and Taylor, D.R.  
Calculation of programmed temperature GC characteristics from isothermal data. I. Theory and computational procedures 405(1987)67
- Alakhov, Y.B., see Prisyazhnoy, V.S. 424(1988)243
- Alarcon De La Lastra, C., see Marhuenda, E. 436(1988)103
- Al-Attar, A.F. and Nickless, G.  
1,2-Diaminobenzene derivatives for the determination of selenium by GC 440(1988)333
- Albaigés, J., see Aceves, M. 436(1988)503
- Albaigés, J., see Alexander, G. 446(1988)87
- Albaigés, J., see Porte, C. 442(1988)386
- Albertsson, A.-C., see Karlsson, S. 442(1988)267
- Albertsson, P.-Å., see Sellergren, B. 450(1988)277
- Albin, H., see Demotes-Mainard, F. 419(1987)388

- Albroscheit, G., see Schulz, H. 442(1988)353
- Alcazar, B., see Rubio, N. 403(1987)312
- Alderweireldt, F.C., see Engelmann, G.J. 432(1988)29
- Aldinger, S., see Preti, G. 432(1988)1
- Aldous, S., see Ismail, Z. 404(1987)372
- Aldrich, J.R., see Rudmann, A.A. 407(1987)324
- Aleotti, M., see Fedeli, G. 447(1988)263
- Aleschko-Ozhevskii, Y.P., see Zaslavsky, B.Y. 439(1988)267
- Alessandri, J.-M., Christon, R., Arfi, T.S., Riaz, A. and Leger, C.L.  
Comparative chromatographic study of modifications of brush-border membrane vesicles induced by an essential fatty acid-deficient diet 432(1988)75
- Alexander, G., Hazai, I., Grimalt, J. and Albaiges, J.  
GC-MS study of  $\alpha$ -phyllocladane dehydrogenation 446(1988)87
- Alfredsson, G., Wiesel, F.-A. and Lindberg, M.  
Glutamate and glutamine in cerebrospinal fluid and serum from healthy volunteers – analytical aspects 424(1988)378
- Aliberti, L.M., see Severini, G. 421(1987)344
- Alick, B., Bridges, C., Cox, Jr., T., Earl, V. and Thedford, R.  
Separation of synthetic cycloalkylated bases, nucleosides and nucleotides by reversed-phase HPLC 430(1988)309
- Al-Khalidi, A.H.E., see Girard, J.E. 448(1988)355
- Allain, P., see Premel-Cabic, A. 434(1988)187
- Allenmark, S., Andersson, S. and Bojarski, J.  
Direct LC separation of enantiomers on immobilized protein stationary phases. VI. Optical resolution of a series of racemic barbiturates: studies of substituent and mobile phase effects 436(1988)479
- Alley, E.G., see Lynn, B.C. 408(1987)211
- Allmendinger, H., see Koppenhoefer, B. 441(1988)89
- Al-Mallah, N.R., Bun, H., Coassolo, P., Aubert, C. and Cano, J.P.  
Determination of the aromatic retinoids (etretin and isoetretin) in biological fluids by HPLC 421(1987)177
- Aloe, M., see Child, P. 415(1987)13
- Alonso, R., see Ferrer, E. 411(1987)167
- Alonso, R., see Ferrer, E. 445(1988)327
- Alpert, A.J.  
Hydrophobic interaction chromatography of peptides as an alternative to reversed-phase chromatography 444(1988)269
- Alpert, A.J. and Andrews, P.C.  
Cation-exchange chromatography of peptides on poly(2-sulfoethyl aspartamide)-silica 443(1988)85
- Alstine, J.M. Van, see Karr, L.J. 442(1988)219
- Al-Thamir, W.K.  
GC determination of water in crude oils 403(1987)288
- Altmann, K. and Mukkur, T.K.S.  
Alternative procedure for the purification of the heat-stable enterotoxin of enterotoxigenic *Escherichia coli* pathogenic for calves 420(1987)253
- Altmayer, B., see Schwenk, S. 448(1988)424
- Alton, K.B., Petrucci, R.F. and Patrick, J.E.  
HPLC assay for dilevalol in human plasma and urine using a PRP-1 column and fluorimetric detection 425(1988)363
- Alvarez-Builla, J., see Gago, F. 449(1988)95
- Alvarez-Gonzalez, R.  
Synthesis and purification of deoxyribose analogues of NAD<sup>+</sup> by affinity chromatography and strong-anion-exchange HPLC 444(1988)89
- Alvarez, J.G., see Touchstone, J.C. 429(1988)359
- Alvarez, J.G. and Touchstone, J.C.  
Separation of acidic and neutral glycolipids by diphasic TLC 436(1988)515
- Amacher, D.E. and Smith, D.J.  
Differences in the constitutive forms of hepatic cytochrome P-450 in male and female adult beagle dogs 419(1987)61
- Amagaya, S., see Umeda, M. 422(1987)246
- Amantea, M.A. and Narang, P.K.  
Improved procedure for quantitation of omeprazole and metabolites using reversed-phase HPLC 426(1988)216
- Amato, A., see Gagliardi, L. 404(1987)267
- Amato, A., see Gagliardi, L. 408(1987)409
- Amato, A., see Gagliardi, L. 448(1988)296
- Ambrus, G., see Szederkényi, F. 446(1988)253
- Ames, M.M., see Frank, S.K. 419(1987)225
- Amici, M., see Moretti, G. 442(1988)459
- Amicosante, G., Marinucci, M.C., Franceschini, N., Tizzani, M.I., Oliva, B. and Oratore, A.  
Fractionation and characterization of two  $\beta$ -lactamases in *Citrobacter diversus* ULA-27 strain by chromatofocusing 403(1987)366
- Amidzhin, B., see Nikolova-Damyanova, B. 446(1988)283
- Amidzhin, B. and Nikolova-Damyanova, B.  
Densitometric identification of triglycerides separated by reversed-phase TLC 446(1988)259
- Amin, M.  
Quantitative determination of some

- prostaglandin derivatives of the sub-groups E<sub>2</sub>, A<sub>2</sub> and B<sub>2</sub> by HPLC 404(1987)385
- Amor, J.-C., see Pedroso, E. 409(1987)281
- Amsden, W.B., see Wilson, T.D. 445(1988)299
- Anastasia, M., Colombo, D. and Fiecchi, A.  
Quantitative HPLC determination of the amino acid spinacine in blood and chow of rats 410(1987)504
- Anazawa, N., see Jinno, K. 402(1987)173
- Anders, G., see Krupčik, J. 407(1987)65
- Anders, J.C., see Zhou, Z.M. 414(1987)77
- Andersen, J.P., see Vilsen, B. 442(1988)229
- Anderson, G.M., Durkin, T.A., Chakraborty, M. and Cohen, D.J.  
LC determination of taurine in whole blood, plasma and platelets 431(1988)400
- Anderson, G.M., Durkin, T.A., Morton, J.B. and Cohen, D.J.  
LC determination of urinary catecholamines after one-step alumina extraction 424(1988)373
- Anderson, G.M., see Durkin, T.A. 428(1988)9
- Anderson, S.D., see Hayes, Jr., P.C. 437(1988)365
- Andersson, B.  
Thermal degradation of weldable poly(vinyl chloride) samples at low temperatures 445(1988)353
- Andersson, E., Håkanson, E., Larsson, J. and Mårtensson, J.  
Rapid and sensitive method for the determination of arterial-venous differences and leg efflux of 3-methylhistidine using ion-pair HPLC and post-column fluorescence derivatization 414(1987)174
- Andersson, L., see Belew, M. 403(1987)197
- Andersson, L., Sulkowski, E. and Porath, J.  
Purification of commercial human albumin on immobilized IDA-Ni<sup>2+</sup> 421(1987)141
- Andersson, S., see Allenmark, S. 436(1988)479
- Andersson, T., see Kågedal, B. 417(1987)261
- Ando, S., see Hirabayashi, Y. 445(1988)377
- Ando, S., Waki, H. and Kon, K.  
New solvent system for HPTLC and HPLC of gangliosides 405(1987)125
- Ando, S., Waki, H. and Kon, K.  
HPLC of underivatized gangliosides 408(1987)285
- Andrejak, M., see Hary, L. 419(1987)396
- Andreolini, F., Borra, C., Wiesler, D. and Novotny, M.  
High-efficiency separation and characterization of high-molecular-weight heterocyclic sulfur compounds in fossil fuels by microcolumn LC and related techniques 406(1987)375
- Andreolini, F., see Flurer, C.L. 448(1988)73
- Andrews, A.T., see Chaplin, L.C. 450(1988)420
- Andrews, P.A., Bachur, N.R., Musser, S.M., Wright, J. and Callery, P.S.  
Silylated N,O-ketals from the reaction of ketones with N-trimethylsilylimidazole 419(1987)271
- Andrews, P.C., see Alpert, A.J. 443(1988)85
- Andrikopoulos, N.K., see Moschidis, M.C. 403(1987)363
- Andronikashvili, T.G., see Soják, L. 436(1988)47
- Andronikashvili, T.G., Tsitsishvili, G.V., Kvernadze, T.K. and Okudzhava, N.G.  
Potential solid supports for GC prepared on the basis of certain rocks 406(1987)145
- Ang, S.B., see Lee, E.J.D. 413(1987)300
- Ang, S.B., see Lee, E.J.D. 420(1987)203
- Angenot, L., see Brasseur, T. 437(1988)260
- Angerer, E. Von, see Birnböck, H. 414(1987)235
- Anjaneyulu, B., see Bhatia, S.C. 434(1988)288
- Anlér, E.-L., see Johansson, M. 427(1988)55
- Ansari, G.A.S., see Bandi, Z.L. 435(1988)513
- Ansari, K.A. and Shoeman, D.  
Rapid separation of free fatty acids from neutral lipids in brain extracts 439(1988)453
- Ansorgová, A., see Vejrosta, J. 447(1988)170
- Anspach, B., Gierlich, H.U. and Unger, K.K.  
Comparative study of Zorbax Bio Series GF 250 and GF 450 and TSK-GEL 3000 SW and SWXL columns in SEC of proteins 443(1988)45
- Antia, F.D. and Horváth, C.  
HPLC at elevated temperatures: examination of conditions for the rapid separation of large molecules 435(1988)1
- Antle, P.E., see Cox, G.B. 444(1988)325
- Antle, P.E., see Eble, J.E. 405(1987)1
- Antle, P.E., see Eble, J.E. 405(1987)31
- Antle, P.E., see Eble, J.E. 405(1987)51
- Antle, P.E., see Snyder, L.R. 444(1988)303
- Antoniadou-Vyza, A., see Tsantili-Kakoulidou, A. 445(1988)317
- Antonopoulos, C.A., see Karamanos, N.K. 405(1987)221
- Antonucci, A., see Pecci, L. 426(1988)183
- Anzai, T., Jaffe, N. and Wang, Y.-M.  
Separation and identification of methotrexate and its metabolites, 7-hydroxymethotrexate and polyglutamates, in human tissues by reversed-phase HPLC coupled with radioimmunoassay 415(1987)445
- Anzion, R.B.M., see Jongeneelen, F.J. 413(1987)227
- Aoyama, T., Kotaki, H., Saitoh, Y. and Nakagawa, F.

- Determination of pemoline in plasma, urine and tissues by reversed-phase HPLC 430(1988)351
- Arai, K., see Koyama, I. 420(1987)275
- Arai, T., Koike, H., Hirata, K. and Ōizumi, H. Separation of pyridone carboxylic acid enantiomers by HPLC using copper(II)-L-amino acid as the eluent 448(1988)439
- Arakawa, Y., see Shibasaki, S. 415(1987)365
- Araki, A. and Sako, Y.  
Determination of free and total homocysteine in human plasma by HPLC with fluorescence detection 422(1987)43
- Araki, M., see Barnhart, E.R. 445(1988)145
- Araki, M., see Tanaka, N. 406(1987)247
- Araki, M., see Tanaka, N. 448(1988)95
- Araki, T., see Sakai, T. 433(1988)73
- Arbin, A., see Jacobsson, S. 447(1988)329
- Arcelloni, C., see Fermo, I. 433(1988)53
- Archer, A.W.  
Determination of safrole and myristicin in nutmeg and mace by HPLC 438(1988)117
- Archer, A.W.  
Determination of cinnamaldehyde, coumarin and cinnamyl alcohol in cinnamon and cassia by HPLC 447(1988)272
- Archer, M.C., see Milligan, J.R. 411(1987)481
- Arco, A. Del, see Usero, J.L. 407(1987)355
- Arfi, T.S., see Alessandri, J.-M. 432(1988)75
- Argilés, J.M., see Peinado, J. 415(1987)372
- Argoudelis, C.J.  
Simple isocratic HPLC method for the separation of the six vitamers of vitamin B<sub>6</sub> 424(1988)315
- Arkesteyn, G.A., see Odink, J. 431(1988)309
- Arm, H., see Brügger, R.R. 440(1988)197
- Armani, E., Barazzoni, L., Dossena, A. and Marchelli, R.  
Bis(L-amino acid amidato)copper(II) complexes as chiral eluents in the enantiomeric separation of D,L-dansylamino acids by reversed-phase HPLC 441(1988)287
- Armani, E., Dossena, A., Marchelli, R. and Virgili, R.  
Copper(II) complexes of diamino-diamido-type ligands as chiral eluents in the enantiomeric separation of D,L-dansylamino acids by reversed-phase HPLC 441(1988)275
- Armstrong, D.W., He, F.-Y. and Han, S.M.  
Planar chromatographic separation of enantiomers and diastereomers with cyclodextrin mobile phase additives 448(1988)345
- Armstrong, D.W., see Armstrong, R.D. 414(1987)192
- Armstrong, D.W., see Han, S.M. 441(1988)376
- Armstrong, D.W., see Spino, L.A. 409(1987)147
- Armstrong, D.W., Spino, L.A., Han, S.M., Seeman, J.I. and Secor, H.V.  
Enantiomeric resolution of racemic nicotine and nicotine analogues by microcolumn LC with  $\beta$ -cyclodextrin inclusion complexes 411(1987)490
- Armstrong, R.D., Ward, T.J., Pattabiraman, N., Benz, C. and Armstrong, D.W.  
Separation of tamoxifen geometric isomers and metabolites by bonded-phase  $\beta$ -cyclodextrin chromatography 414(1987)192
- Arnaud, P., see Miribel, L. 405(1987)337
- Arnemann, W., see Gielsdorf, W. 419(1987)351
- Arnemann, W., see Gielsdorf, W. 419(1987)357
- Arnoux, P., see Payet, B. 424(1988)337
- Aronson, A.L., see Tyczkowska, K. 427(1988)103
- Aronsson, E., see Odham, G. 434(1988)31
- Arrendale, R.F., Stewart, J.T. and Tackett, R.L.  
Determination of clonidine in human plasma by cold on-column injection capillary GC-selected-ion monitoring-MS 432(1988)165
- Arvidsson, T.  
Effects of system peaks in a column-switching LC system for bioanalysis 407(1987)49
- Arvidsson, T.  
Effects of drug-protein binding on trace enrichment of drugs in blood plasma on short precolumns 439(1988)353
- Arwert, F., see Lindemans, J. 432(1988)103
- Arwood, L.J., see Spiker, S. 405(1987)327
- Asai, H., see Takeuchi, T. 407(1987)151
- Asakura, T., see Adachi, K. 419(1987)303
- Asakura, T., see Adachi, K. 428(1988)247
- Asano, O., see Yamada, K. 433(1988)243
- Ascalone, V., Catalani, P., Dal Bo', L., Deschamps, N. and Guinebault, P.  
Determination of alpidem and its metabolites in human plasma by HPLC and fluorimetric detection 414(1987)101
- Ascalone, V. and Dal Bo', L.  
Automated HPLC and column-switching technique for on-line clean-up and analysis of diltiazem in human plasma 423(1987)239
- Ascencio-Valle, F., see Ochoa, J.L. 449(1988)322
- Ashoor, S.H. and Knox, M.J.  
Determination of vitamin A derivatives in fortified foods and commercial vitamin supplements by HPLC 409(1987)419
- Ashraf, M., see Singh, A.K. 404(1987)223
- Ashraf, M., see Singh, A.K. 425(1988)245
- Asotra, S., Mladenov, P.V. and Burke, R.D.  
Improved method for benzoyl chloride

- derivatization of polyamines for HPLC  
408(1987)227
- Assan, R., see Ducrocq, R. 419(1987)75
- Assenza, S.P., see Koenigbauer, M.J.  
413(1987)161
- Assmann, B. and Haas, H.J.  
Determination of the age dependency of the  
creatinine-related pseudouridine excretion in  
children's urine by HPLC 434(1988)202
- Assmann, G., see Jabs, H.-U. 414(1987)323
- Astbury, C. and Dixon, J.S.  
Rapid method for the determination of either  
plasma sulphapyridine or sulphamethoxazole  
and their acetyl metabolites using HPLC  
414(1987)223
- Astorg, P.O., see S eb edio, J.L. 405(1987)371
- Au, J.L.-S., Su, M.-H. and Wientjes, M.G.  
Isocratic ion-pair reversed-phase HPLC  
analysis of ATP and ADP 423(1987)308
- Au-Yeung, C.Y., see Leung, C.P. 449(1988)341
- Aubel, M.T. and Rogers, L.B.  
Effects of pretreatment on the  
enantioselectivity of silica-bound proteins  
used as HPLC stationary phases 408(1987)99
- Aubert, C., see Al-Mallah, N.R. 421(1987)177
- Aue, W.A., see Siu, K.W.M. 406(1987)81
- Aue, W.A., see Siu, K.W.M. 408(1987)53
- Aue, W.A., see Tang, Y.-Z. 408(1987)69
- Aue, W.A., see Tang, Y.-Z. 409(1987)125
- Aue, W.A., see Tang, Y.-Z. 409(1987)243
- Auer, B., see Haring, C. 428(1988)160
- Aujoulat, P., see Airaud, C.B. 437(1988)59
- Aussel, C., Cynober, L. and Giboudeau, J.  
GC of branched-chain keto acids analysed as  
their silylated oxime derivatives 423(1987)270
- Autelitano, D.J., see Smith, A.I. 416(1987)255
- Auwera, L. Van der, see De Bondt, M.  
442(1988)165
- Avery, M.J. and Junk, G.A.  
Procedure for the determination of amines in  
small urine samples 420(1987)379
- Avgerinos, A. and Hutt, A.J.  
Determination of the enantiomeric  
composition of ibuprofen in human plasma  
by HPLC 415(1987)75
- Avnir, D., see Farin, D. 406(1987)317
- Avram, M.J. and Krejcie, T.C.  
Determination of sodium pentobarbital and  
either sodium methohexital or sodium  
thiopental in plasma by HPLC with  
ultraviolet detection 414(1986)484
- Awni, W.M., Clarkson, J. and Guay, D.R.P.  
Determination of ciprofloxacin and its 7-  
ethylenediamine metabolite in human serum  
and urine by HPLC 419(1987)414
- Awni, W.M. and Maloney, J.A.  
Optimized HPLC method for the analysis of  
cyclosporine and three of its metabolites in  
blood and urine 425(1988)233
- Axelsson, J.E., see Chan, G.L.-Y. 417(1987)295
- Ayyangar, N.R. and Bhide, S.R.  
Separation of eight alkaloids and meconic  
acid and quantitation of five principal  
alkaloids in gum opium by gradient  
reversed-phase HPLC 436(1988)455
- Azeemuddin, S., Israili, Z.H. and Bharmal, F.M.  
Rapid method for evaluating compliance of  
6-mercaptopurine therapy in children with  
leukemia 430(1988)163
- Azerad, R., see Dernoncour, R. 410(1987)355
- Azerad, R., see Maurs, M. 440(1988)209
- Azuma, I., see Sugimura, K. 440(1988)131
- Azuma, R., see Marunaka, T. 433(1988)177

## B

- Baars, A.J., see Van Beck, H. 442(1988)345
- Baars, A.M., see Vree, T.B. 417(1987)458
- Baars, A.M., see Vree, T.B. 424(1988)440
- Baba, S., see Furuta, T. 413(1987)1
- Baba, S., see Sasaki, Y. 426(1988)93
- Babashak, J.V. and Phillips, T.M.  
Use of avidin-coated glass beads as a support  
for high-performance immunoaffinity  
chromatography 444(1988)21
- Babuřikova, O., see Bohacek, J. 434(1988)439
- Bachur, N.R., see Andrews, P.A. 419(1987)271
- Badarani, M.M., see Dubruc, C. 417(1987)208
- Badcock, N.R. and Zoanetti, G.D.  
Micro-determination of clobazam and N-  
desmethylclobazam in plasma or serum by  
electron-capture GC 421(1987)147
- Badreldin, M.M., see Moneam, N.M.A.  
437(1988)316
- Badreldin, M.M., see Moneam, N.M.A.  
438(1988)438
- Baert, L., see Der Meeren, P.V. 447(1988)436
- Bagchi, S.P.  
Single-step method for derivatization of  
dopamine and some related compounds in  
aqueous media for GC 421(1987)227
- Bagchi, S.P., see Jindal, S.P. 408(1987)356
- Bagirov, T.O., see Zaslavsky, B.Y. 439(1988)267
- Bailey, E., Brooks, A.G.F., Purchase, R.,



- Meakings, M., Davies, M. and Walters, D.G.  
Improved method for the determination of the major neutral steroids and unconjugated bile acids in human faeces using capillary GC 421(1987)21
- Bailey, E., Farmer, P.B., Peal, J.A., Hotchkiss, S.A. and Caldwell, J.  
Analytical methodology to determine stable isotopically labelled and unlabelled theophylline in human plasma using capillary GC-MS 416(1987)81
- Bailey, E., Peal, J.A. and Philbert, M.  
Determination of 1,3-dinitrobenzene and its metabolites in rat blood by capillary GC with electron-capture detection 425(1988)187
- Bailey, Jr., J.E., see Richfield-Fratz, N.  
405(1987)283
- Bailly, E., see Jaureguierry, G. 440(1988)385
- Bailly, J.R., see Leita, J. 435(1988)229
- Bailon, P., see Weber, D.V. 431(1988)55
- Baird-Lambert, J.A., see Gross, A.S.  
416(1987)400
- Baker, G.B., see Drebit, R. 432(1988)334
- Baker, G.B., see Wong, J.T.F. 428(1988)140
- Baker, J.O., Tucker, M.Y., Lastick, S.M. and Himmel, M.E.  
Degradation of ketoses during aqueous HPLC on lead-form cation-exchange resins 437(1988)387
- Baker, P.R., Vitale, G.C. and Siow, Y.F.  
Medroxyprogesterone acetate- and ethinylestradiol-induced changes in biliary bile acids of the rat studied by HPLC 423(1987)63
- Bakker, F.P., see Keuken, M.P. 439(1988)13
- Bakos, J., see Lecomte, L. 408(1987)416
- Balachandran Nayar, M.S., see Egorin, M.J.  
415(1987)148
- Balásperi, L., see Péter, A. 408(1987)43
- Baldessarini, R.J., see Lampen, P. 426(1988)283
- Baldwin, R.P., see Santos, L.M. 414(1987)161
- Balíková, M., Nováková, E. and Kohlíček, J.  
Identification of polychlorinated phenols in urine by GC and TLC 431(1988)431
- Balíková, M. and Kohlíček, J.  
Rapid determination of ethylene glycol at toxic levels in serum and urine 434(1988)469
- Ballard, K.D., see Wilson, M.J. 431(1988)222
- Ballery, N., see Yon, J.M. 440(1988)421
- Ballester, P., see García-Raso, A. 402(1987)323
- Ballet, J.-J., see Jaureguierry, G. 440(1988)385
- Ballio, A., see Marra, M. 440(1988)47
- Balmér, K., Zhang, Y., Lagerström, P.-O. and Persson, B.-A.  
Determination of metoprolol and two major metabolites in plasma and urine by column LC and fluorometric detection 417(1987)357
- Baltensperger, U. and Kern, S.  
Determination of mono- and divalent cations and anions in small fog samples by ion chromatography 439(1988)121
- Baltes, E., Coupez, R., Brouwers, L. and Gobert, J.  
GC method for the determination of cetirizine in plasma 430(1988)149
- Bamonte, F., see Dionisotti, S. 434(1988)169
- Banavalikar, M.M., see Bhatia, S.C. 434(1988)288
- Bandi, Z.L. and Ansari, G.A.S.  
Separation of rat liver phospholipids from polar non-phospholipids by TLC 435(1988)513
- Bandó, Y., see Yamaguchi, Y. 422(1987)53
- Banhidi, Z.G., see Karlsson, S. 442(1988)267
- Bannon, C.D., Craske, J.D., Felder, D.L., Garland, I.J. and Norman, L.M.  
Analysis of fatty acid methyl esters with high accuracy and reliability. VI. Rapid analysis by split injection capillary GLC 407(1987)231
- Bannon, C.D., Craske, J.D., Felder, D.L. and Norman, L.M.  
Investigation of pre-set and actual splitting ratios in split injection capillary GLC 404(1987)340
- Bannon, C.D., Craske, J.D. and Norman, L.M.  
Effect of overload of capillary GLC columns on the equivalent chain lengths of C<sub>18</sub> unsaturated fatty acid methyl esters 447(1988)43
- Bannow, C.A., see Kaluzny, B.D. 414(1987)228
- Bapat, B.V., see Panse, D.G. 411(1987)456
- Baraldi, R., Chen, K.-H. and Cohen, J.D.  
Microscale isolation technique for quantitative GC-MS analysis of indole-3-acetic acid from cherry (*Prunus cerasus* L.) 442(1988)301
- Barascut, J.-L., see Maggio, A.-F. 436(1988)23
- Barazzoni, L., see Armani, E. 441(1988)287
- Barbalas, M., see Rubio, F. 421(1987)281
- Barbhaiya, R.H., see Forgue, S.T. 414(1987)343
- Barbier, M.  
Interactions between ascorbic acid and linear and cyclized bile pigments. A TLC study 440(1988)275
- Barbieri, L., Stoppa, C. and Bolognesi, A.  
Large scale chromatographic purification of ribosome-inactivating proteins 408(1987)235
- Barboro, P., see Candiano, G. 423(1987)319
- Barcelo, A.R., Muñoz, R. and Sabater, F.  
Reaction of Fast Blue BB salt and *p*-

- dimethylaminocinnamaldehyde with indole-3-acetic acid degradation products.  
Evaluation of the coupling properties of the reagents used in the zymographic screening of indole-3-acetic acid oxidases 411(1987)476
- Barceló, D., see Porte, C. 442(1988)386
- Bardalaye, P.C., see Toth, J.P. 408(1987)335
- Bareggi, S.R., see Zecca, L. 420(1987)417
- Barford, R.A., see Parris, N. 444(1988)219
- Bargar, E.M.  
Application of column switching in HPLC analysis of medroxoalol in plasma 417(1987)143
- Barnoux, H., see Bougerolle, A.M. 434(1988)232
- Barker, I.K., see Raynor, M.W. 436(1988)497
- Barkia, A., see Mezdoor, H. 414(1987)35
- Barkley, R.M., see DeAngelis, J.J. 441(1988)125
- Barlow, R.D., see Parviainen, M.T. 431(1988)216
- Barlow, R.D., Thompson, P.A. and Stone, R.B.  
Simultaneous determination of nicotine, cotinine and five additional nicotine metabolites in the urine of smokers using pre-column derivatisation and HPLC 419(1987)375
- Barnas, C., see Haring, C. 428(1988)160
- Barnby, M.A., see Yamasaki, R.B. 447(1988)277
- Barnhart, E.R., Patterson, Jr., D.G., Tanaka, N. and Araki, M.  
Reversed-phase LC of the twenty-two tetrachlorodibenzo-*p*-dioxin isomers on pyrenylethyl- and octadecylsilylated silica gel columns 445(1988)145
- Barot-Ciorbaru, R., see Moroux, Y. 440(1988)119
- Barr, S.W., see Larson, J.R. 405(1987)163
- Barre, J., see Duche, J.-C. 423(1987)340
- Barrientos, A., see Chan, K.K. 428(1988)331
- Barry, H.D., see Ismail, Z. 404(1987)372
- Bartek, M.J., Vekshteyn, M., Boarman, M.P. and Gallo, D.G.  
LC determination of sotalol in plasma and urine employing solid-phase extraction and fluorescence detection 421(1987)309
- Barter, P.J., see Clifton, P.M. 414(1987)25
- Bartle, K.D., see Raynor, M.W. 436(1988)497
- Barton, D.H.R., see Porziemsky, J.-P. 440(1988)183
- Barton, R.N., see Passingham, B.J. 416(1987)25
- Bartosevich, J.F., see Kapetanovic, I.M. 419(1987)458
- Basak, B., see Laskar, S. 436(1988)341
- Basili, A., see Gagliardi, L. 408(1987)409
- Bassoli, A., Chioccare, F., Di Gregorio, G., Rindone, B., Tollari, S., Falagiani, P., Riva, G. and Bolzacchini, E.  
Analysis of allergenic components of a *Parietaria judaica* pollen extract by chromatographic methods for the evaluation of purification procedures 444(1988)209
- Bassoli, A., Chioccare, F., Di Gregorio, G., Rindone, B., Tollari, S., Falagiani, P., Riva, G. and Bolzacchini, E.  
Micropreparative-scale enrichment of some allergenic components of *Parietaria* pollen extract 446(1988)179
- Basta, M.T., see Brooks, J.B. 430(1988)209
- Basu, P.S., see Datta, P.K. 431(1988)37
- Bates, E.J., see Ferrante, A. 440(1988)105
- Bates, M., see Spiker, S. 405(1987)327
- Bathelt, E.R., see Haagsma, N. 436(1988)73
- Batten, C.F., see Wentworth, W.E. 441(1988)45
- Baty, J.D., see Lindsay, R.M. 433(1988)292
- Baty, J.D. and Sharp, S.  
Prediction of HPLC capacity factors following column calibration. Influence of different solvents and columns on the separation of a series of aromatic amides 437(1988)13
- Bauchet, J., Pussard, E. and Garaud, J.J.  
Determination of vancomycin in serum and tissues by column LC using solid-phase extraction 414(1987)472
- Baudin-Chich, V., Marden, M. and Wajcman, H.  
Investigation of the tetramer-dimer equilibrium in haemoglobin solutions by high-performance SEC on a diol column 437(1988)193
- Bauer, G., see Menge, G.P. 414(1987)477
- Bauer, J.  
Electrophoretic separation of cells (Review) 418(1987)359
- Bauer, J., Linton, C. and Norris, B.  
Stability-indicating HPLC analysis of tin protoporphyrin and other free acid metalloporphyrins 445(1988)429
- Bauer, J.H., see Souther, M.E. 417(1987)27
- Baugh, P.J., Casson, A., Jones, M.W. and Jones, A.C.  
Sampling and injection system for the analysis of metalorganic chemicals by GC-MS. Some initial results 411(1987)445
- Baumann, R.A., Kamminga, D.A., Derlagen, H., Gooijer, C., Velthorst, N.H. and Frei, R.W.  
Dynamic quenching of europium(III) and terbium(III) luminescence: a potential detection method in ion chromatography 439(1988)165
- Baumann, R.J., see Robbins, D.K. 415(1987)208
- Baumhoer, G., see Blaszkewicz, M. 439(1988)109
- Bauwens, R.M., see De Leenheer, A.P. 429(1988)3
- Baveja, S.K., see Padmalatha Devi, K. 426(1988)229

- Baveja, S.K., see Padmalatha Devi, K. 434(1988)265
- Baverel, G., see Durozard, D. 414(1987)460
- Bayer, E., see Koppenhoefer, B. 406(1987)157
- Baykousheva, S.  
Secretion of proteins by *Bacillus subtilis* 168 grown in the presence of membrane active agents (alcohols) 440(1988)379
- Bayliss, M.A.J., Homer, R.B. and Shepherd, M.J.  
Anthracene-9-carbonyl chloride as a fluorescence and UV derivatising reagent for the HPLC analysis of hydroxy compounds 445(1988)393
- Bayliss, M.A.J., Homer, R.B. and Shepherd, M.J.  
Determination of diethylene glycol in wine by HPLC using anthracene-9-carbonyl chloride as a derivatising reagent 445(1988)403
- Bayne, W.F., see Chiou, R. 416(1987)196
- Bayne, W.F., see Chiou, R. 422(1987)281
- Bayne, W.F., see Constanzer, M.L. 413(1987)277
- Bayne, W.F., see Hsieh, J.Y.-K. 430(1988)175
- Bayne, W.F., see Matuszewski, B.K. 426(1988)295
- Bayne, W.F., see Musson, D.G. 414(1987)121
- Bayne, W.F., see Rothwell, C.M. 417(1987)447
- Bayne, W.F., see Stubbs, R.J. 413(1987)171
- Bažant, L., Wurst, M. and Smolková-Keulemansová, E.  
 $\beta$ -Cyclodextrin as a selective agent for separation of selected aromatic acids by HPLC 445(1988)337
- Beasley, T.M., see Zaromb, S. 438(1988)100
- Beasley, V.R., see Harada, K.-I. 448(1988)275
- Beaty, N.B. and Mello, R.J.  
Extracellular mammalian polysaccharides: glycosaminoglycans and proteoglycans (Review) 418(1987)187
- Beaudoin, N., see Dubé, L.M. 427(1988)113
- Beaune, P.H., Cresteil, T., Flinois, J.-P. and Leroux, J.-P.  
Hydrophobic chromatography: a one-step method for the purification of human liver microsomal epoxide hydrolase 426(1988)169
- Beauvallet, C., Hountondji, C. and Schmitter, J.-M.  
Analytical strategy for determination of active site sequences in aminoacyl-tRNA synthetases 438(1988)347
- Beck, K., see Radeck, W. 432(1988)297
- Beck, R.H.F. and Uchiyama, T.  
Activity staining of inulin fructotransferase II on polyacrylamide gels 445(1988)282
- Becker, H. and Gnauck, R.  
Inverse GC study of polyether alcohols (in German) 410(1987)267
- Becker, J.U., see Krämer, B.K. 427(1988)351
- Becker, M., see Hartley, R. 415(1987)357
- Becker, T.A., see Brown, L.L. 426(1988)370
- Becue, T., see Cõng, H.N.G. 407(1987)349
- Beecher, G.R., see Khachik, F. 449(1988)119
- Beek, H. Van and Baars, A.J.  
Isolation and quantitation of cadmium-, zinc- and copper-metallothioneins by HPLC atomic absorption spectrometry 442(1988)345
- Beek, W.M.J., see Aerts, M.M.L. 435(1988)97
- Beeksmá, S., see De Jong, J. 414(1987)285
- Beger, J., see Voelkel, A. 409(1987)29
- Beger, J., see Voelkel, A. 448(1988)219
- Begley, P., Foulger, B. and Simmonds, P.  
Femtogram detection of perfluorocarbon tracers using capillary GC-electron-capture negative ion chemical ionisation MS 445(1988)119
- Begley, P. and Foulger, B.E.  
Detection of trace levels of triclopyr using capillary GC-electron-capture negative-ion chemical ionization MS 438(1988)45
- Behar, N., see Sébille, B. 409(1987)61
- Beightol, L.A., see Karnes, H.T. 422(1987)165
- Beightol, L.A., see Karnes, H.T. 424(1988)398
- Beijsterveldt, L.E.C. Van, see Wuis, E.W. 420(1987)212
- Bekersky, I., see Puglisi, C.V. 417(1987)347
- Bekersky, I., see Puglisi, C.V. 419(1987)253
- Belanger, K., Collins, J.M. and Klecker, Jr., R.W.,  
Technique for detection of DNA nucleobases by reversed-phase HPLC optimized for quantitative determination of thymidine substitution by iododeoxyuridine 417(1987)57
- Belbeke, F. and DeBackere, M.  
High-sensitivity determination of nadolol in plasma with mass-selective detection 416(1987)443
- Belenkii, B.G.  
Use of laser detectors in capillary LC (Review) 434(1988)337
- Belenkii, B.G., see Maltsev, V.G. 416(1987)45
- Belenkii, B.G., see Tennikova, T.B. 435(1988)357
- Belew, M., Yip, T.-T., Andersson, L. and Porath, J.  
Interaction of proteins with immobilized  $\text{Cu}^{2+}$ . Quantitation of adsorption capacity, adsorption isotherms and equilibrium constants by frontal analysis 403(1987)197
- Belfroid, R.D.M., see Van der Krogt, J.A. 427(1988)9
- Bellama, J.M., see Blair, W.R. 410(1987)393
- Bellama, J.M., see Eng, G. 403(1987)1
- Bellama, J.M., see Tierney, E.J. 441(1988)229
- Belle, S.J.P. Van, see De Smet, M. 416(1987)375
- Belleville, F., see Brossat, B. 413(1987)141

- Bellon, G. and Borel, J.P.  
Quantitative evaluation of collagen fractions separated by acrylamide gel electrophoresis: its application for collagen typing in normal and pathological tissues 433(1988)81
- Belsner, K., see Weber, C. 426(1988)345
- Belvito, M.L., see Fóti, G. 440(1988)315
- Ben-Hur, E., see Zuk, M.M. 433(1988)367
- Benanchi, P.L., Gazzei, G. and Giannozzi, A.  
Purification of specific heterologous F(ab)<sub>2</sub> fragments with DEAE-Zeta-Prep cartridges for ion-exchange chromatography 450(1988)133
- Benbow, R.M., see Stowers, D.J. 444(1988)47
- Benčekroun, Y., Ribon, B., Desage, M. and Brazier, J.L.  
Simultaneous quantitation of phenobarbitone and *p*-hydroxyphenobarbitone and their perdeuterioethyl analogues in biological fluids by combined GC-MS 420(1987)287
- Benecke, R., Vetter, B. and De Zecuw, R.A.  
Rapid micromethod for the analysis of mitotane and its metabolite in plasma by GC with electron-capture detection 417(1987)287
- Benedek, K., Hughes, B., Seaman, M.B. and Swadesh, J.K.  
Reversed-phase LC and sodium dodecyl sulfate polyacrylamide gel electrophoresis characteristics of a recombinant DNA derived malaria antigen 444(1988)191
- Benedetti, A., see Aggazzotti, G. 416(1987)125
- Beneken Kolmer, E.W.J., see Wuis, E.W. 415(1987)419
- Beneš, L., see Štefek, M. 415(1987)163
- Benet, L.Z., see Gawienowski, M. 430(1988)412
- Benet, L.Z., see Hyneck, M.L. 420(1987)349
- Benet, L.Z., see Lee, F.W. 426(1988)259
- Benet, L.Z., see Spahn, H. 433(1988)331
- Benfenati, E., see Latini, R. 424(1988)211
- Benická, E., see Krupčík, J. 448(1988)203
- Beninati, S., Martinet, N. and Folk, J.E.  
HPLC method for the determination of  $\epsilon$ -( $\gamma$ -glutamyl)lysine and mono- and bis- $\gamma$ -glutamyl derivatives of putrescine and spermidine 443(1988)329
- Bennekom, W.P. Van, see Van Opstal, M.A.J. 432(1988)395
- Benny, A.G. and Brimer, C.M.  
Chromatographic procedure for preparation of a high-purity C 1 esterase inhibitor concentrate 433(1988)363
- Benowitz, N.L., see Jacob, III, P. 417(1987)277
- Benowitz, N.L., see Shulgin, A.T. 423(1987)365
- Benveniste, J., see Bossant, M.J. 423(1987)23
- Benz, C., see Armstrong, R.D. 414(1987)192
- Beran, M., Jizba, J., Blumauerová, M., Němeček, J., Novák, J., Zima, J., Kandybin, N.V. and Samoukina, G.V.  
Preparative HPLC of macrotretolides 442(1988)431
- Beránková, E., see Rauch, P. 438(1988)451
- Berek, D., see Buszewski, B. 446(1988)191
- Berenson, G.S., see Lee, Y.J. 419(1987)275
- Beresford, A.P., Macrae, P.V., Stopher, D.A. and Wood, B.A.  
Analysis of amlodipine in human plasma by GC 420(1987)178
- Beretta, E., Botturi, S., Ferrari, P., Tuan, G. and Zerilli, L.F.  
Identification of 5-hydroxypyrazinamide isolated from urine of subjects given pyrazinamide 416(1987)144
- Berezkin, V.G. and Korolev, A.A.  
"Superdynamic" coating of the inner surface of a capillary column with a liquid stationary phase 440(1988)323
- Berezkina, L.G., Borisova, S.I. and Melnikov, S.E.  
Chromatographic method for kinetic study of heterogeneous reactions 446(1988)65
- Berg, D., see Bruns, A. 450(1988)111
- Berg, G.A. Van den, Kingma, A.W. and Muskiet, F.A.J.  
Determination of polyamines in human erythrocytes by capillary GC with nitrogen-phosphorus detection 415(1987)27
- Berg, R.H. Van den, see Jansen, E.H.J.M. 437(1988)268
- Bergenheim, N., Carlsson, U., Ellow, L.-E. and Rosén, S.  
Rapid ion-exchange chromatography for preparative separation of proteins. III. Application to purification of bovine factor X and prothrombin 403(1987)383
- Bergens, A.  
Determination of nitrodiphenylamines by LC and dual-electrode amperometric detection 410(1987)437
- Berger, J.A., see Bougerolle, A.M. 434(1988)232
- Berger, Y., see Fels, J.-P. 430(1988)73
- Bergeron, M.G., see Croteau, D. 419(1987)205
- Bergqvist, Y., Churchill, F.C. and Mount, D.L.  
Determination of mefloquine by electron-capture GC after phosgene derivatization in biological samples and in capillary blood collected on filter paper 428(1988)281
- Bergqvist, Y., Eckerbom, S., Larsson, N. and Churchill, F.C.  
HPLC separation and extraction investigation for the simultaneous determination of

- mefloquine and its carboxylic acid metabolite 427(1988)295
- Bergqvist, Y., Hellgren, U. and Churchill, F.C. HPLC assay for the simultaneous monitoring of mefloquine and its acid metabolite in biological samples using protein precipitation and ion-pair extraction 432(1988)253
- Bergqvist, Y. and Churchill, F.C. Detection and determination of antimalarial drugs and their metabolites in body fluids (Review) 434(1988)1
- Bergueiro, R., see García-Raso, A. 402(1987)323
- Berman, S.S., see Siu, K.W.M. 408(1987)53
- Bernard, M., see Ricard, B. 428(1988)203
- Bernstein, J.R., see Miller, B.J. 426(1988)438
- Berridge, J.C. Analysis of tioconazole using HPLC with a porous graphitic carbon column 449(1988)317
- Berry, D.J., see Streete, J.M. 416(1987)281
- Bertaux, O., see Công, H.N.G. 407(1987)349
- Bertelli, R., see Candiano, G. 423(1987)319
- Berthod, A. and Roussel, A. The rôle of the stationary phase in micellar liquid chromatography. Adsorption and efficiency 449(1988)349
- Bertrand, M.J., see Steffenrud, S. 416(1987)219
- Bertrand, M.J., see Steffenrud, S. 423(1987)1
- Bertrand, O., see Krovianski, Y. 449(1988)403
- Bertrand, O., see Krovianski, Y. 449(1988)413
- Bertucci, C., see Salvadori, P. 450(1988)163
- Betteridge, R.F., Bosanquet, A.G., Hunt, P. and Gilby, E.D. Determination of diaziridinyl-3,6-bis(2-hydroxyethylamino)-1,4-benzoquinone in plasma by solid-phase extraction and HPLC 432(1988)352
- Betti, A., see Coppi, S. 442(1988)97
- Betton, J.-M., see Yon, J.M. 440(1988)421
- Betts, T.J. Characterisation of volatile oil constituents with relatively long GC retention times on two stationary phases 449(1988)312
- Betts, T.J. and Sufredo, D. Requirements of mutually miscible liquids for ternary chromatographic solvents with reference to monosaccharide PC 436(1988)53
- Bevan, R., see Sommadossi, J.-P. 414(1987)429
- Beveridge, D.J., Harrison, D.M. and McGrath, R.M. Liquid CC using TLC silica gel 450(1988)443
- Beyec, Y. Le, see Van, T.D. 440(1988)71
- Bezard, J., see Narce, M. 448(1988)249
- Bhale, A., see Panse, D.G. 411(1987)456
- Bhamra, R.K., Ward, A.E. and Holt, D.W. Column LC measurement of betaxolol in plasma or serum 417(1987)229
- Bhargava, V.O. Quantitation of 1-( $\beta$ -D-glucopyranosyl)-phenobarbital in plasma and bile 419(1987)421
- Bhargava, V.O., Emodi, S. and Hirate, J. Quantitation of acetaminophen and its metabolites in rat plasma after a toxic dose 426(1988)212
- Bharmal, F.M., see Azeemuddin, S. 430(1988)163
- Bharucha, E.D., see Bhatia, S.C. 434(1988)296
- Bharucha, E.D., see Bhatia, S.C. 434(1988)303
- Bhatia, A., see Jain, R. 441(1988)454
- Bhatia, K.L., see Kumar, S. 434(1988)228
- Bhatia, S.C., Revankar, S.N., Anjaneyulu, B., Desai, N.D., Banavalikar, M.M., Sahajwalla, C.G. and Subrahmanyam, D. HPLC of antifilarials. I. Simultaneous determination of 1-4-(4-nitrophenyl-aminophenyl thiocarbamido-4-methyl)piperazine and three of its metabolites in biological fluids 434(1988)288
- Bhatia, S.C., Revankar, S.N., Bharucha, E.D., Desai, N.D. and Sahajwalla, C.G. HPLC of antifilarials. III. Determination of 2-*tert*-butyl-6-isothiocyanato-5-piperidin-1-yl)benzthiazole in biological fluids 434(1988)303
- Bhatia, S.C., Revankar, S.N., Doshi, K.J., Desai, N.D., Bharucha, E.D. and Sahajwalla, C.G. HPLC of antifilarials. II. Determination of S-(2-carboxyethyl)-N-(2-*tert*-butyl-5-methoxybenzthiazol-6-yl) dithiocarbamate and its active isothiocyanate metabolite in biological fluids 434(1988)296
- Bhatnagar, D., see McCormick, S.P. 441(1988)400
- Bhavnagri, V.P., see Osman, M.A. 419(1987)452
- Bhide, M.V., see Dunphy, M.J. 420(1987)394
- Bhide, S.R., see Ayyangar, N.R. 436(1988)455
- Bhide, S.V., see Talib, Y.Y. 403(1987)373
- B'Hymer, C. Static headspace GC method for the determination of residual solvents in vigabatrin drug substance 438(1988)103
- Biagi, G.L., see Pietrogrande, M.C. 447(1988)404
- Bialer, M., see Samara, E. 416(1987)370
- Bianchetti, G., see Ganansia, J. 421(1987)83
- Bianchi, A. and Cook, H.A. Modified method for analysis of C<sub>2</sub>-C<sub>5</sub> hydrocarbons in an aromatic-alkane matrix using an automated thermal desorber 449(1988)175

- Bianchi, G. and Caccia, S.  
Simultaneous determination of buspirone, gepirone, ipsapirone and their common metabolite 1-(2-pyrimidinyl)piperazine in rat plasma and brain by HPLC 431(1988)477
- Bianco, O., see Petrarulo, M. 432(1988)37
- Bicchi, C., Frattini, C. and Raverdino, V.  
Considerations and remarks on the analysis of *Anthemis nobilis* L. essential oil by capillary GC and "hyphenated" techniques 411(1987)237
- Bickel, M.H., see Moor, M.J. 431(1988)455
- Bickley, H.M., see Laing, D.K. 442(1988)187
- Bidault, R., see Le Coguic, A. 434(1988)320
- Bidleman, T.F., see Zaranski, M.T. 409(1987)235
- Biedermann, M., see Grob, K. 448(1988)387
- Biemer, T.A.  
Simultaneous analysis of acetaminophen, pseudoephedrine hydrochloride and chlorpheniramine maleate in a cold tablet using an isocratic, mixed micellar HPLC mobile phase 410(1987)206
- Biemer, T.A., see Talmage, J.M. 410(1987)494
- Bier, M., see Mosher, R.A. 436(1988)191
- Bies, J.M., see Schneebaum, S. 427(1988)166
- Biessels, P.T.M., Agoston, S. and Horn, A.S.  
HPLC determination of 2,4-diaminopyridine in plasma and urine 421(1987)392
- Biggs, W.R., see Jinno, K. 402(1987)173
- Biggs, W.R., see Jinno, K. 436(1988)1
- Bighi, C., see Coppi, S. 442(1988)97
- Billard, V., Bruneteau, M., Bonnet, P., Ricci, P., Pernollet, J.C., Huet, J.C., Vergne, A., Richard, G. and Michel, G.  
Chromatographic purification and characterization of elicitors of necrosis on tobacco produced by incompatible *Phytophthora* species 440(1988)87
- Billiet, H.A.H., see Erkelens, C. 404(1987)67
- Billing, U., see Engewald, W. 446(1988)71
- Binst, G. Van, see De Bondt, M. 442(1988)165
- Biondi, P.A., Manca, F., Negri, A., Secchi, C. and Montana, M.  
GC analysis of neutral monosaccharides as their O-pentafluorobenzyloxime acetates 411(1987)275
- Birckel, P., see Jehl, F. 413(1987)109
- Birk, I.M., Matern, U., Kaiser, I., Martin, C. and Weckesser, J.  
The peptide toxin of the cyanobacterium *Microcystis aeruginosa* PCC 7941. Isolation and analysis by NMR and fast atom bombardment MS 449(1988)423
- Birkett, D.J., see Chan, K. 426(1988)103
- Birkett, D.J., see Lelo, A. 430(1988)203
- Birkholz, D.A., Coutts, R.T. and Hrudey, S.E.  
Determination of polycyclic aromatic compounds in fish tissue 449(1988)251
- Birnböck, H., Ringshandl, R. and Von Angerer, E.  
Chromatographic analysis of the new antiestrogen zindoxifene and its metabolites in biological material 414(1987)235
- Birsner, U., Gilles, U., Nielsen, P. and McMaster, G.K.  
Reversed-phase HPLC system for the rapid, automated purification of oligonucleotides 402(1987)381
- Bischoff, R., see Mazsaroff, I. 437(1988)429
- Bischoff, R., see McLaughlin, L.W. 418(1987)51
- Biseth, A., see Christophersen, A.S. 422(1987)117
- Bisse, E. and Wieland, H.  
HPLC separation of human haemoglobins. Simultaneous quantitation of foetal and glycosylated haemoglobins 434(1988)95
- Bisseret, F., see Rihn, B. 428(1988)408
- Bizzi, B., see De Cristofaro, R. 426(1988)376
- Björklund, H.  
Determination of oxytetracycline in fish by HPLC 432(1988)381
- Björkman, S.  
Determination of the enantiomers of ketoprofen in blood plasma by ion-pair extraction and HPLC of leucinamide derivatives 414(1987)465
- Björkman, S., see Jonung, M. 415(1987)35
- Björkman, S. and Elisson, L.O.  
Determination of quinacrine (mepacrine) in plasma by HPLC with fluorimetric detection 420(1987)341
- Björkman, S. and Stanski, D.R.  
Simultaneous determination of fentanyl and alfentanil in rat tissues by capillary column GC 433(1988)95
- Björkqvist, B. and Hase, A.  
Separation and determination of intact glucosinolates in rapeseed by HPLC 435(1988)501
- Blaauboer, B.J., see 't Klooster, G.A.E.V. 432(1988)223
- Black, B.H., Wechter, M.A. and Hardy, D.R.  
Determination of corrosion inhibitor/lubricity enhancer additives in jet fuels by SEC 437(1988)203
- Black, R.M. and Read, R.W.  
Detection of trace levels of thiodiglycol in blood, plasma and urine using GC-electron-capture negative-ion chemical ionisation MS 449(1988)261
- Blackett, A., Dhillon, S., Cromarty, J.A., Horne, R. and Richards, G.  
Rapid and sensitive HPLC assay for

- midazolam and 1-hydroxymidazolam, the major metabolite, in human serum 433(1988)326
- Blacklaws, C., see Laing, D.K. 442(1988)187
- Blackman, G.L., see Sitaram, B.R. 422(1987)13
- Bładek, J.  
New method of visualizing thin-layer chromatograms by using liquid crystals 405(1987)203
- Bładek, J.  
Parameters of the liquid crystal method of visualizing thin-layer chromatograms 437(1988)131
- Blair, W.R., Parks, E.J., Olson, G.J., Brinckman, F.E., Valeiras-Price, M.C. and Bellama, J.M. Characterization of organotin species using microbore and capillary LC techniques with an epifluorescence microscope as a novel imaging detector 410(1987)393
- Blaise, A., see Larroque, M. 407(1987)384
- Blaise, A., see Larroque, M. 445(1988)107
- Blanc-Continsouza, D., see Chabenat, C. 414(1987)417
- Blanchin, M.D. and Rondot-Dudragne, M.L.  
Determination of cefaloglycine and cefroxadine in biological media by TLC with fluorimetric detection (in French) 432(1988)407
- Blanquat, G.D. Saint, Lamboeuf, Y. and Fritsch, P.  
Determination of catecholamines in biological tissues by LC with coulometric detection 415(1987)388
- Błaszkwicz, M., Baumhoer, G., Neidhart, B., Ohlendorf, R. and Linscheid, M.  
Identification of trimethyllead in urine by HPLC with column switching and chemical reaction detection and by LC-MS 439(1988)109
- Blau, P.A., Hines, J.W. and Voyksner, R.D.  
Application of HPLC and thermospray HPLC-MS to the analysis and identification of 2',3'-dideoxyadenosine and its metabolite in biological media 420(1987)1
- Bleau, G., see Vézina, D. 426(1988)385
- Blitterswijk, H. Van, see Van Ginkel, L.A. 445(1988)385
- Blouin, F.A. and Zarins, Z.M.  
HPLC of cottonseed flavonoids 441(1988)443
- Blouin, R.A., see Huang Chandler, M.H. 419(1987)426
- Blumauerová, M., see Beran, M. 442(1988)431
- Blumer, J.L., see Myers, C.M. 422(1987)153
- Bo', L. Dal, see Ascalone, V. 414(1987)101
- Bo', L. Dal, see Ascalone, V. 423(1987)239
- Boarman, M.P., see Bartek, M.J. 421(1987)309
- Bobbitt, D.R., see Rice, P.D. 437(1988)3
- Bochner, F., see Chen, Z.R. 424(1988)170
- Bochner, F., see Siebert, D.M. 420(1987)425
- Boer, G. De, see Kubo, I. 402(1987)354
- Boersma, E.R., see Van der Steege, G. 415(1987)1
- Boese, F.G.  
Contributions to a mathematical theory of free-flow electrophoresis 438(1988)145
- Bogaards, J.J.P., see Odink, J. 431(1988)309
- Bogar, R.G., see Krahn, M.M. 437(1988)161
- Bogusz, M. and Aderjan, R.  
Improved standardization in reversed-phase HPLC using 1-nitroalkanes as a retention index scale 435(1988)43
- Boháček, J., Hošek, B., Babušíková, O. and Hrivňáková, A.  
Determination of adenosine deaminase and purine nucleoside phosphorylase activities using HPLC and radiochromatographic methods 434(1988)439
- Bohbot, M., Doare, L. and Diquet, B.  
Determination of a new benzamide, amisulpride, in human plasma by reversed-phase ion-pair HPLC 416(1987)414
- Boismare, F., see Chabenat, C. 414(1987)417
- Boivin, P., see Kroviarski, Y. 449(1988)403
- Boivin, P., see Kroviarski, Y. 449(1988)413
- Bojarski, J., see Allenmark, S. 436(1988)479
- Bojarski, J., see Żukowski, J. 436(1988)381
- Bojkova, N., see Lekova, K. 446(1988)31
- Bojkova, S.V., see Rachev, R.C. 437(1988)287
- Bökens, H., Bourscheidt, C. and Müller, R.F.  
Determination of bumetanide in plasma by HPLC 434(1988)327
- Bolasco, A., see Gagliardi, L. 404(1987)267
- Boldizsár, M., see Kremmer, T. 415(1987)53
- Boley, N.P. and Colwell, R.K.  
Comparison of methods for the determination of butyric acid in foodstuffs by GLC and HPLC 410(1987)190
- Bolognesi, A., see Barbieri, L. 408(1987)235
- Bolzacchini, E., see Bassoli, A. 444(1988)209
- Bolzacchini, E., see Bassoli, A. 446(1988)179
- Bolzacchini, E., see Fermo, I. 433(1988)53
- Bonakdar, M., see Wang, J. 415(1987)432
- Bondt, M. De, Couder, J., Van der Auwera, L., Van Marsenille, M., Elseviers, M., Delaet, N., Laus, G., Tourwé, D. and Van Binst, G.  
Determination of the chiral purity of dipeptide isosteres containing a reduced peptide bond by GC analysis 442(1988)165
- Bonini, P.A., see Fermo, I. 433(1988)53
- Bonn, G., see Derler, H. 440(1988)281
- Bonnet, P., see Billard, V. 440(1988)87
- Bonniec, B. Le, see Ducrocq, R. 419(1987)75
- Booij, L.H.D.J., see Vree, T.B. 424(1988)440

- Boomsma, F., Van der Hoorn, F.A.J.,  
Man in 't Veld, A.J. and Schalekamp,  
M.A.D.H.  
Determination of D,L-threo-3,4-  
dihydroxyphenylserine and of the D- and L-  
enantiomers in human plasma and urine  
427(1988)219
- Boot, J.H., see 't Klooster, G.A.E.V.  
432(1988)223
- Booy, L.H.D.J., see Koopman-Kimenai, P.M.  
416(1987)382
- Boppana, V.K., see Rhodes, G.R. 444(1988)123
- Borchardt, L.G., see Crockett, T.D. 407(1987)330
- Borchardt, R.T., see Scriba, G.K.E. 433(1988)31
- Borea, P.A., see Pietrogrande, M.C. 447(1988)404
- Borel, C., see Marston, A. 450(1988)91
- Borel, J.P., see Bellon, G. 433(1988)81
- Borgeat, P., see Steffenrud, S. 416(1987)219
- Borgeat, P., see Steffenrud, S. 423(1987)1
- Borger, F.R., see Lee, J.K. 444(1988)141
- Borghini, C., see Paroni, R. 420(1987)189
- Bories, G., see Rao, D. 410(1987)169
- Borisova, S.I., see Berezkina, L.G. 446(1988)65
- Borisova, V., see Krasilnikov, I. 446(1988)211
- Borner, S. and Brenneisen, R.  
Determination of tryptamine derivatives in  
hallucinogenic mushrooms using HPLC with  
photodiode array detection 408(1987)402
- Borra, C., see Andreolini, F. 406(1987)375
- Borra, C., see Flurer, C.L. 448(1988)73
- Bos, O.J.M., Fischer, M.J.E., Wilting, J. and  
Janssen, L.H.M.  
Procedure for the isolation of a large peptic  
fragment of human serum albumin  
424(1988)13
- Bosanquet, A.G., see Betteridge, R.F.  
432(1988)352
- Bosch, D.V. Den, see Van Ginkel, L.A.  
445(1988)385
- Bosch, F., see Mateo, R. 410(1987)319
- Boschetti, E., see Moroux, Y. 440(1988)119
- Bosin, T.R. and Faull, K.F.  
Measurement of  $\beta$ -carbolines by HPLC with  
fluorescence detection 428(1988)229
- Bossant, M.J., Farinotti, R., Mencia-Huerta,  
J.M., Benveniste, J. and Mahuzier, G.  
Characterization and quantification of PAF-  
acether (platelet-activating factor) as a  
heptafluorobutyrate derivative of 1-O-alkyl-  
2-acetyl-sn-glycerol by capillary column GC  
with electron-capture detection 423(1987)23
- Bossche, W. Van den, see De Schutter, J.A.  
428(1988)301
- Bossinger, P., see Räder, K. 408(1987)341
- Bossle, P.C., Reutter, D.J. and Sarver, E.W.  
Analysis of alkyl methylphosphonic acids in  
aqueous matrices by ion-pair reversed-phase  
ion chromatography 407(1987)399
- Boswell, R.T., see Sonnet, P.E. 436(1988)205
- Both-Miedema, R., see Jansen, E.H.J.M.  
435(1988)363
- Botha, H., see Swart, K.J. 413(1987)315
- Bothwell, B.E., see Shah, J.A. 414(1987)1
- Böttcher, A., see Schmitz, G. 431(1988)327
- Botterblom, M.H.A., Janssen, T.J., Guelen,  
P.J.M. and Vree, T.B.  
Rapid and sensitive determination of  
ambroxol in human plasma and urine by  
HPLC 421(1987)211
- Botturi, S., see Beretta, E. 416(1987)144
- Boucly, P., see Chabenat, C. 414(1987)417
- Boucly, P., see Tharasse-Bloch, C. 421(1987)407
- Boudinot, F.D., see Satterwhite, J.H.  
431(1988)444
- Boudinot, F.D. and Ibrahim, S.S.  
HPLC assay for piroxicam in human plasma  
430(1988)424
- Bougerolle, A.M., Chabard, J.L., Jbilou, M.,  
Bagnoux, H., Petit, J., Berger, J.A. and  
Dordain, G.  
Simultaneous determination of trimipramine  
and its demethylated metabolites in plasma  
by GC-MS 434(1988)232
- Boulton, A.A., see Durden, D.A. 440(1988)253
- Bourne, D.W.A., see Hakim, L. 424(1988)111
- Bourquin, D. and Brenneisen, R.  
Confirmation of cannabis abuse by the  
determination of 11-nor- $\Delta^9$ -  
tetrahydrocannabinol-9-carboxylic acid in  
urine with HPLC and electrochemical  
detection 414(1987)187
- Bours, J., Dempewolf, J., Schaake, S. and Rink,  
H.  
Determination of the electrophoretic order of  
26 nucleotides by isotachopheresis  
403(1987)336
- Bourscheidt, C., see Bökens, H. 434(1988)327
- Boussetta, H., see Créminon, C. 440(1988)439
- Bouthillier, F., Carmel, M., Elhilali, M., Radwan,  
F. and Lehoux, J.-G.  
Application of high-performance liquid  
chromatofocusing to the study of prostatic  
androgen receptors. Effect of stabilizing  
agents on the heterogeneity of receptor  
structure 403(1987)171
- Bouvet, J.-P., see Moroux, Y. 440(1988)119
- Bowers, E., see McCormick, S.P. 441(1988)400
- Bowman, P.B., see Clark, C.P. 408(1987)275
- Bradshaw, J.S., Aggarwal, S.K., Rouse, C.A.,  
Tarbet, B.J., Markides, K.E. and Lee, M.L.  
Polysiloxanes containing thermally stable  
chiral amide side-chains for capillary GC and



- supercritical fluid chromatography  
405(1987)169
- Brady, P.H., see Cornell, H.J. 421(1987)61
- Brain, K.R., see Pourgholami, M.H. 424(1988)163
- Braithwaite, I.M., see Coleman, M.D.  
414(1987)242
- Bramer, A.M., see Hughes, D.E. 408(1987)296
- Brandin, S., see Polacek, J. 425(1988)424
- Brandšterová, E., Lehotay, J., Garaj, J. and  
Leclerq, P.A.  
LC of dibutylthiocarbamate degradation  
products 404(1987)359
- Brandts, P.M., Gelsema, W.J. and De Ligny, C.L.  
Influence of additives to the eluent on the  
hydrophobic interaction chromatography of  
simple compounds. IV. Influence of  
electrolytes on the retention of normal  
alkanols at 25 °C 437(1988)337
- Brandts, P.M., Gelsema, W.J. and De Ligny, C.L.  
Influence of additives to the eluent on  
hydrophobic interaction chromatography of  
simple compounds. V. Influence of pH and  
electrolytes on the retention of *n*-  
alkylammonium ions and *n*-alkyl  
carboxylates 438(1988)181
- Brashear, W.T., see Kuhnert, B.R. 426(1988)392
- Brassat, B., see König, W.A. 447(1988)193
- Brasseur, T., Wauters, J.-N. and Angenot, L.  
Densitometric determination of quercetin  
heterosides in plant extracts (in French)  
437(1988)260
- Brassinne, C., Laduron, C., Coune, A., Sculier,  
J.P., Hollaert, C., Collette, N. and Meunier,  
F.  
HPLC determination of amphotericin B in  
human serum 419(1987)401
- Braut, D., see Dellinger, M. 422(1987)73
- Brayan, J.G., Haddad, P.R., Sharp, G.J., Dilli, S.  
and Desmarchelier, J.M.  
Determination of organophosphate pesticides  
and carbaryl on paddy rice by reversed-phase  
HPLC 447(1988)249
- Brazier, J.L., see Benchekroun, Y. 420(1987)287
- Brechbühler, B., see Schilling, B. 435(1988)204
- Breckenridge, A.M., see Coleman, M.D.  
414(1987)242
- Bredeweg, R.A., see Larson, J.R. 405(1987)163
- Bredow, J. Von, see Kapetanovic, I.M.  
419(1987)458
- Breen, C., see Meaney, M. 449(1988)241
- Breimer, D.D., see Schellens, J.H.M.  
431(1988)203
- Breimer, D.D., see Soons, P.A. 428(1988)362
- Breimer, D.D., see Van Harten, J. 423(1987)327
- Bremm, K.D., see Knöller, J. 427(1988)257
- Brendel, E., Meineke, I., Henne, E.-M., Zschunke,  
M. and De Mey, C.  
Sensitive HPLC determination of  
pseudoephedrine in plasma and urine  
426(1988)406
- Brendel, E., Meineke, I., Stüwe, E. and Osterwald,  
H.  
Stability of 5-aminosalicylic acid and 5-  
acetylaminosalicylic acid in plasma  
432(1988)358
- Brendel, S., see Kleuss, C. 407(1987)281
- Brenneisen, R., see Borner, S. 408(1987)402
- Brenneisen, R., see Bourquin, D. 414(1987)187
- Breslow, E., see Burman, S. 443(1988)285
- Bresson, M., see Rop, P.P. 427(1988)172
- Bretschneider, W., see Werkhoff, P. 405(1987)87
- Bretschneider, W., see Werkhoff, P. 405(1987)99
- Brett, M.A., Dierdorf, H.-D., Zussman, B.D. and  
Coates, P.E.  
Determination of paroxetine in human  
plasma, using HPLC with fluorescence  
detection 419(1987)438
- Breuer, J., see Haas, R.H. 425(1988)47
- Brewster, M.A., see Turley, C.P. 421(1987)430
- Brezgunov, V.N., see Shukun, S.A. 432(1988)137
- Bridges, C., see Alick, B. 430(1988)309
- Briele, H.A., see Schneebaum, S. 427(1988)166
- Brieva, A., see Rubio, N. 403(1987)312
- Brieva, A., see Rubio, N. 404(1987)378
- Brieva, A., see Rubio, N. 407(1987)408
- Brieva, A., see Rubio, N. 408(1987)420
- Brimer, C.M., see Benny, A.G. 433(1988)363
- Brimmell, P.A. and Sams, D.J.  
Rapid and simple assay for the measurement  
of methotrexate in serum, urine and red blood  
cells by reversed-phase HPLC 413(1987)320
- Brinckman, F.E., see Blair, W.R. 410(1987)393
- Brinckman, F.E., see Eng, G. 403(1987)1
- Brinckman, F.E., see Tierney, E.J. 441(1988)229
- Brindle, P.A., see Brooks, C.J.W. 438(1988)108
- Brinkman, U.A.T., see Aerts, M.M.L.  
435(1988)97
- Brinkman, U.A.T., see Broersen, J.J. 436(1988)39
- Brinkman, U.A.T., see De Ruiter, C.  
436(1988)429
- Brinkman, U.A.T., see Irth, H. 424(1988)95
- Brinkman, U.A.T., see Irth, H. 439(1988)63
- Brinkman, U.A.T., see Jansen, H. 440(1988)217
- Brinkman, U.A.T., see Maris, F.A. 435(1988)297
- Brinkman, U.A.T., see Maris, F.A. 445(1988)129
- Brinkman, U.A.T., see Nielen, M.W.F.  
417(1987)159
- Brodfehrer, J.I. and Powis, G.  
GC assay for the new antitumor agent  
sulfamic acid diester (NSC 329680) and its

- stability in buffer, blood and plasma  
427(1988)247
- Brodie, M.J., see Forrest, G. 428(1988)123
- Brodie, R.R., Chasseaud, L.F. and Robbins, A.D.  
Determination of physostigmine in plasma  
by HPLC 415(1987)423
- Broersen, J.J., Jansen, H., De Ruiter, C.,  
Brinkman, U.A.T., Frei, R.W., Buijtenhuijs,  
F.A. and Van der Maeden, F.P.B.  
Derivatization of some polymers with *o*-  
phthalaldehyde prior to high-performance  
SEC 436(1988)39
- Broly, F., Libersa, C. and Lhermitte, L.  
HPLC assay for mexiletine hydroxylation in  
microsomes of human liver 431(1988)369
- Bromet-Petit, M., see Duchene, P. 424(1988)205
- Bron, J., see De Jong, A.P. 419(1987)233
- Brons, P.P.T., Wessels, J.M.C., Linssen, P.C.M.,  
Haanen, C. and Speth, P.A.J.  
Determination of amsacrine in human  
nucleated hematopoietic cells 422(1987)175
- Brooks, A.G.F., see Bailey, E. 421(1987)21
- Brooks, C.J.W., Brindle, P.A., Cole, W.J. and  
Watson, D.G.  
Analytical separations of enantiomeric  
primary amines as pyrrole derivatives formed  
by reaction with polygodial 438(1988)108
- Brooks, C.J.W. and Cole, W.J.  
Cyclic di-*tert*.-butylsilylene derivatives of  
substituted salicylic acids and related  
compounds. A study by GC-MS 441(1988)13
- Brooks, D.E., see Skuse, D.R. 432(1988)127
- Brooks, J.B., Basta, M.T. and El Kholy, A.M.  
Studies of metabolites in diarrheal stool  
specimens positive for *Klebsiella*, *Serratia*,  
and *Proteus* spp. by frequency-pulsed  
electron-capture GC 430(1988)209
- Brooks, J.B., see Daneshvar, M.I. 433(1988)248
- Brooks, P.M., see Dawson, M. 420(1987)129
- Brossat, B., Straczek, J., Heulin, M.-H.,  
Herbeuval, X., Belleville, F., Nabet, P. and  
Lokiec, F.  
LC determination of cyclosporin A in serum  
with use of a solid-phase extraction.  
Comparison between HPLC and  
radioimmunoassay levels in clinical  
investigations 413(1987)141
- Broto, F., see Aceves, M. 436(1988)503
- Brouwer, E., see Irth, H. 439(1988)63
- Brouwers, L., see Baltés, E. 430(1988)149
- Brown, D.S. and Jenke, D.R.  
Determination of trace levels of oxytocin in  
pharmaceutical solutions by HPLC  
410(1987)157
- Brown, D.W., see Krahn, M.M. 437(1988)161
- Brown, E.M., Dower, H.J. and Greenberg, R.  
Effects of reductive alkylation on peptide  
retention times 443(1988)247
- Brown, H., see Holland, O.B. 426(1988)148
- Brown, J.W., Carballeira, A. and Fishman, L.M.  
Rapid two-dimensional TLC system for the  
separation of multiple steroids of secretory  
and neuroendocrine interest 439(1988)441
- Brown, K.F., see Gross, A.S. 416(1987)400
- Brown, L.L., Williams, P.E., Becker, T.A., Ensley,  
R.J., May, M.E. and Abumrad, N.N.  
Rapid HPLC method to measure plasma  
leucine: importance in the study of leucine  
kinetics *in vivo* 426(1988)370
- Brown, L.W., see Van der Meer, M.J.  
423(1987)351
- Brown, S.Y., see Miwa, B.J. 431(1988)343
- Browne, T.R., see Mikati, M.A. 433(1988)305
- Brügger, R.R., Marti, A.R., Meyer, V.R. and  
Arm, H.  
Optical resolution of samples with weak  
interactions on chiral "brush-type" LC  
stationary phases 440(1988)197
- Bruijn, E.A. De, see Reeuwijk, H.J.E.M.  
428(1988)93
- Bruijn, E.A. De, see Tjaden, U.R. 420(1987)53
- Bruins, A.P., see Hendriks, H. 428(1988)352
- Bruins, C.H.P., see Smilde, A.K. 410(1987)1
- Bruinsma, J., see Vermeer, E. 404(1987)346
- Brummelen, P. Van, see Van Harten, J.  
423(1987)327
- Brun, B., see Rudziński, W. 406(1987)295
- Brun, S., see Larroque, M. 445(1988)107
- Brundrett, R.B., see Cheng, T.-L. 425(1988)303
- Bruneau, P., see Meinard, C. 450(1988)169
- Bruneteau, M., see Billard, V. 440(1988)87
- Brüning, T., see Schmitz, G. 431(1988)327
- Bruns, A., Berg, D. and Werner-Busse, A.  
Isolation of tocopherol homologues by  
preparative HPLC 450(1988)111
- Bruns, M.B., see Lakings, D.B. 428(1988)131
- Bruyn, P.C. De, see Claessens, H.A. 436(1988)345
- Bryan, C.S., see Greene, S.V. 417(1987)121
- Buchanan, M.R., see Haas, T.A. 430(1988)1
- Buchberger, W.  
Investigations into lactoperoxidase-catalysed  
bromination of tyrosine and thyroglobulin  
432(1988)57
- Buchberger, W.  
Determination of iodide and bromide by ion  
chromatography with post-column reaction  
detection 439(1988)129
- Bucheli, F., see Wyss, R. 424(1988)303
- Bucheli, F., see Wyss, R. 430(1988)81
- Bucheli, F., see Wyss, R. 431(1988)297

- Buck, W.B., see Swanson, S.P. 414(1987)335
- Buckley, D.G., see Whelpton, R. 426(1988)223
- Budd, R.D.  
Validity of *post mortem* chest cavity blood ethanol determinations 449(1988)337
- Budd, R.D., see Mathis, D.F. 439(1988)466
- Budevsky, O., see Dimitrova, B. 409(1987)81
- Buell, P.E., see Girard, J.E. 448(1988)355
- Buffham, B.A., see Rathor, M.N. 404(1987)33
- Buffham, B.A., see Rathor, M.N. 445(1988)37
- Buijtenhuijs, F.A., see Broersen, J.J. 436(1988)39
- Buist, G.J., see Abraham, M.H. 409(1987)15
- Bulbena, O., see Cárdenas, A. 426(1988)83
- Bult, A., see Van Opstal, M.A.J. 432(1988)395
- Bun, H., see Al-Mallah, N.R. 421(1987)177
- Burckart, G.J., see D'Souza, M.J. 421(1987)198
- Burford, R.P., see Haken, J.K. 441(1988)207
- Burger, B.V. and Munro, Z.  
Headspace gas analysis. Liquid desorption of headspace volatiles trapped on activated carbon open tubular traps 402(1987)95
- Burger, G., see Pochard, M.-F. 409(1987)315
- Burger, K., see Péter, A. 408(1987)43
- Burgers, R.C.M., see Wuis, E.W. 415(1987)419
- Burke, Jr., T.R., see Knight, M. 444(1988)345
- Burke, J.T., see Krstulović, A.M. 426(1988)417
- Burke, R.D., see Asotra, S. 408(1987)227
- Burke, Jr., T.R. and Knight, M.  
Counter-current chromatographic purification of [D-Ala<sup>1</sup>]peptide T amide 411(1987)431
- Burke, T.W.L., see Hodges, R.S. 444(1988)349
- Burleigh, B.D., see Meng, H. 443(1988)183
- Burman, S., Breslow, E., Chait, B.T. and Chaudhary, T.  
Application of HPLC in neurophysin disulfide assignment 443(1988)285
- Burnett, D., see Ladds, G. 414(1987)355
- Burrows, J.L.  
Determination of oxpentifylline and three metabolites in plasma by automated capillary GC using nitrogen-selective detection 423(1987)139
- Burrows, J.L., Jolley, K.W. and Sullivan, D.J.  
Determination of roxatidine in human plasma, urine and milk by capillary GC using nitrogen-selective detection 432(1988)199
- Burrows, R., Gillespie, D.G., Cooke, M. and Eddib, O.  
Simple preparation of a Poly-S 179 capillary column for the analysis of polycyclic aromatic hydrocarbons and triglycerides 404(1987)248
- Burstein, E.S., Friedman, H. and Greenblatt, D.J.  
Simplified measurement of haloperidol by GC with nitrogen-phosphorus detection 423(1987)380
- Bursukova, M., see Lekova, K. 446(1988)23
- Bursukova, M., see Lekova, K. 446(1988)31
- Burton, L.C. and James, C.A.  
Rapid method for the determination of ifosfamide and cyclophosphamide in plasma by HPLC with solid-phase extraction 431(1988)450
- Burton, S.J., McLoughlin, S.B., Stead, C.V. and Lowe, C.R.  
Design and applications of biomimetic anthraquinone dyes. I. Synthesis and characterisation of terminal ring isomers of C.I. Reactive Blue 2 435(1988)127
- Burton, W.G., Nugent, K.D., Slattery, T.K., Summers, B.R. and Snyder, L.R.  
Separation of proteins by reversed-phase HPLC. I. Optimizing the column 443(1988)363
- Burton, W.G., see Nugent, K.D. 443(1988)381
- Bury, R.W., see Fullinlaw, R.O. 415(1987)347
- Bury, R.W., see Fullinlaw, R.O. 433(1988)131
- Buschauer, A., see Büyüktimkin, N. 450(1988)281
- Bush, P.B., see Akbari, A. 426(1988)207
- Busson, R., see Cachet, T. 445(1988)290
- Buszewski, B., Berek, D., Garaj, J., Novák, I. and Suprynowicz, Z.  
Influence of the porous silica gel structure on the coverage density of a chemically bonded C<sub>18</sub> phase for HPLC 446(1988)191
- Buszewski, B., see Nawrocki, J. 449(1988)1
- Buszewski, B., see Suprynowicz, Z. 446(1988)347
- Butler, E.C.V.  
Determination of inorganic arsenic species in aqueous samples by ion-exclusion chromatography with electrochemical detection 450(1988)353
- Buttar, H.S., see Qureshi, S.A. 431(1988)465
- Büyüktimkin, N. and Buschauer, A.  
Separation and determination of some amino acid ester enantiomers by TLC after derivatization with (*S*)-(+)-naproxen 450(1988)281
- Bystrický, Ľ., Veselá, Z. and Sohler, E.  
Study of the side-reaction products of phenylhydrazine production by GC-MS 447(1988)202
- Bystrický, S., see Strašák, M. 403(1987)331
- Bystrický, S., see Strašák, M. 450(1988)284

## C

- Caccamo, F., Carfagnini, G., Di Corcia, A. and Samperi, R.  
Measurement of urinary estriol glucuronides during the menstrual cycle by HPLC 434(1988)61
- Caccia, S., see Bianchi, G. 431(1988)477
- Caccia, S., see Guiso, G. 413(1987)363
- Cacciapuoti, G., see Porcelli, M. 440(1988)151
- Cache, P., see Capon, C. 425(1988)35
- Cachet, T., Haest, G., Busson, R., Janssen, G. and Hoogmartens, J.  
Isolation of erythromycin E from commercial erythromycin 445(1988)290
- Cachet, T., Kibwage, I.O., Roets, E., Hoogmartens, J. and Vanderhaeghe, H.  
Optimization of the separation of erythromycin and related substances by HPLC 409(1987)91
- Cachet, T., Roets, E., Hoogmartens, J. and Vanderhaeghe, H.  
Separation of novel derivatives from commercial erythromycin samples by TLC 403(1987)343
- Cagniant, E., see Tillier, C. 441(1988)406
- Cai, D.-G., see Zhang, T.-Y. 435(1988)159
- Cai, D.-G., see Zhang, T.-Y. 442(1988)455
- Cai, H., see Yamaguchi, J. 441(1988)183
- Calder, I.C., see Aguilar, M.I. 426(1988)315
- Caldwell, J., see Bailey, E. 416(1987)81
- Caldwell, J., see Hotchkiss, S.A. 423(1987)179
- Callery, P.S., see Andrews, P.A. 419(1987)271
- Callery, P.S., see Egorin, M.J. 415(1987)148
- Callery, P.S., see Vicchio, D. 415(1987)104
- Calvert, R.T., see Mehta, A.C. 421(1987)377
- Cammarata, P., see Moretti, G. 442(1988)459
- Camp, R.D.R., see Mallet, A.I. 432(1988)278
- Campeau, D., Gruda, I., Thibeault, Y. and Legendre, F.  
Analysis of amphoteric surfactants of the alkylaminopropylglycine type by GC 405(1987)305
- Campos, L.G., see Jham, G.N. 436(1988)100
- Canafax, D.M., see Erdmann, G.R. 433(1988)187
- Canafax, D.M., see Remmel, R.P. 432(1988)388
- Candiano, G., Ghiggeri, G.M., Piccardo, M.T., Bertelli, R., Barboro, P., Lazzarini, G., Rialdi, G. and Vecchio, G.  
Purification of intact concanavalin A in tetramers by isoelectric focusing 423(1987)319
- Candito, M., Lapalus, P. and Chambon, P.  
Determination of free and sulphate-conjugated 3-methoxy-4-hydroxyphenylethylene glycol in human plasma by LC with electrochemical detection 430(1988)198
- Canfell, P.C., see Furuta, T. 427(1988)41
- Cann-Moisan, C., Caroff, J. and Girin, E.  
Determination of flavin adenine dinucleotide in biological tissues by HPLC with electrochemical detection 442(1988)441
- Cannell, G.R., see Hooper, W.D. 416(1987)347
- Cano, J.P., see Al-Mallah, N.R. 421(1987)177
- Cano, J.P., see Payet, B. 424(1988)337
- Cant, N.W., see Yong, Y.-S. 450(1988)399
- Cantelmi, M.G., see Palmerini, C.A. 417(1987)378
- Cantelmi, M.G., see Palmerini, C.A. 423(1987)292
- Capon, C., Cache, P., Leroy, Y., Strecker, G., Montreuil, J. and Fournet, B.  
Isolation of the major O-glycosidically linked oligosaccharides obtained by alkaline borohydride degradation of human meconium glycoproteins 425(1988)35
- Caponecchi, G., see Polcaro, C. 448(1988)127
- Cappiello, P.E., see Kling, G.J. 407(1987)377
- Capri, S., see Marcomini, A. 403(1987)243
- Caprioli, R.M., Moore, W.T., DaGue, B. and Martin, M.  
Microbore HPLC-MS for the analysis of proteolytic digests by continuous-flow fast-atom bombardment MS 443(1988)355
- Carballeira, A., see Brown, J.W. 439(1988)441
- Cárdenas, A., Abián, J., Bulbena, O., Roselló, J. and Gelpi, E.  
Determination of oxidation products of N-phenyllinoleamide: Spanish toxic oil syndrome studies 426(1988)83
- Carfagnini, G., see Caccamo, F. 434(1988)61
- Carilli, C.T., Wallace, L.C., Smith, L.M., Wong, M.A. and Lewicki, J.A.  
Semi-preparative purification of recombinant human renin and prorenin 444(1988)203
- Carlier, J.-P. and Sellier, N.  
Identification by GC-MS of short-chain hydroxy acids produced by *Fusobacterium* species and *Clostridium innocuum* 420(1987)121
- Carlier, O., see Ducrocq, R. 419(1987)75
- Carlin, A.S., Prasad, V.K., Sager, A.O., Simmons, J.E. and Skelly, J.P.  
Analysis of prednisolone acetate and related corticoids in swine plasma by reversed-phase HPLC 425(1988)162

- Carlin, J.R., Christofalo, P. and VandenHeuvel, W.J.A.  
HPLC determination of N-(2-methyl-2-propyl)-3-oxo-4-aza-5 $\alpha$ -androst-1-ene-17 $\beta$ -carboxamide, a 4-azasteroid, in human plasma from a phase I study 427(1988)79
- Carlson, R.M., see Xue, W. 447(1988)81
- Carlsson, M., see Kågedal, B. 417(1987)261
- Carlsson, U., see Bergenhem, N. 403(1987)383
- Carlucci, G., see Palumbo, G. 424(1988)201
- Carmel, M., see Bouthillier, F. 403(1987)171
- Carmichael, W.W., see Harada, K.-I. 448(1988)275
- Caroff, J., see Cann-Moisan, C. 442(1988)441
- Carpita, N.C., see Shea, E.M. 445(1988)424
- Carr, P.W., see Wade, J.L. 449(1988)53
- Carr, S., see Moore, J.M. 410(1987)297
- Carroll, D.L., see Tavss, E.A. 438(1988)273
- Carroll, D.L., see Tavss, E.A. 438(1988)281
- Carrott, P.J.M. and Sing, K.S.W.  
GC study of microporous carbons 406(1987)139
- Carstensen, J.T., see Ertel, K.D. 411(1987)297
- Carter, R.H., see Firestone, M.A. 407(1987)363
- Cartoni, G.P., see Coppi, S. 442(1988)97
- Carty, P. and O'Kennedy, R.  
Use of HPLC for the purification of antibodies and antibody conjugates and the study of antibody-antigen interactions 442(1988)279
- Casado, J., see Usero, J.L. 407(1987)355
- Casella, M.L., see Castells, R.C. 402(1987)65
- Casoli, A., Mangia, A., Predieri, G. and Sappa, E.  
Behaviour of tri- and tetranuclear iron and nickel clusters in HPLC 447(1988)187
- Caspi, E., see Njar, V.C.O. 440(1988)415
- Cassada, D., see Raha, C. 450(1988)325
- Cassiano, L., see Castagnola, M. 440(1988)231
- Casson, A., see Baugh, P.J. 411(1987)445
- Castagnola, M., Cassiano, L., De Cristofaro, R., Landolfi, R., Rossetti, D.V. and Marini Bettolo, G.B.  
HPLC in protein sequence determinations 440(1988)231
- Castagnola, M., see De Cristofaro, R. 426(1988)376
- Castagnoli, K.P., see Furuta, T. 427(1988)41
- Castañeda-Hernández, G., see Flores-Murrieta, F.J. 428(1988)167
- Castello, G., Gerbino, T.C., Giacosa, A., Frascio, F. and Sukkar, S.G.  
GC evaluation of hydrogen in the breath as a diagnostic method for the study of intestinal digestion and absorption of dietary carbohydrates in man 416(1987)119
- Castello, G. and Gerbino, T.C.  
Effect of temperature on the GC separation of halogenated compounds on polar and non-polar stationary phases 437(1988)33
- Castells, R.C. and Casella, M.L.  
Evaporation rates of solvent blends measured by GC 402(1987)65
- Castle, L., Gilbert, J., Jickells, S.M. and Gramshaw, J.W.  
Analysis of the plasticiser acetyltributyl citrate in foods by stable isotope dilution GC-MS 437(1988)281
- Castle, L., Sharman, M. and Gilbert, J.  
Analysis of the epoxidised soya bean oil additive in plastics by GC 437(1988)247
- Castoldi, D., Monzani, V. and Tofanetti, O.  
Simultaneous determination of nimesulide and hydroxynimesulide in human plasma and urine by HPLC 425(1988)413
- Castro, M.D.L. De, see Lázaro, F. 448(1988)173
- Catalani, P., see Ascalone, V. 414(1987)101
- Catalin, J., see Payet, B. 424(1988)337
- Catellani, P., see Fanali, S. 405(1987)385
- Catterick, T., see White, P.C. 402(1987)135
- Catterick, T. and Russell, J.  
Development of an inexpensive flowmeter for HPLC using a programmable pocket computer 407(1987)310
- Catz-Biro, L., see Milligan, J.R. 411(1987)481
- Caude, M., see Jadaud, P. 439(1988)195
- Caude, M., see Janicot, J.L. 437(1988)351
- Caude, M., see Jardy, A. 439(1988)137
- Caude, M., see Lienne, M. 448(1988)55
- Caude, M., see Macaudiere, P. 405(1987)135
- Caude, M., see Macaudière, P. 450(1988)255
- Caude, M., see Morin, P. 407(1987)87
- Caude, M., see Ni, P. 424(1988)255
- Cautreels, W., see Fels, J.-P. 430(1988)73
- Cavallini, D., see Pecci, L. 426(1988)183
- Cavazzutti, G., see Gagliardi, L. 404(1987)267
- Cavazzutti, G., see Gagliardi, L. 408(1987)409
- Cavazzutti, G., see Gagliardi, L. 448(1988)296
- Cavin, J.C. and Rodriguez, E.  
HPLC identification of simple  $\beta$ -carboline alkaloids in specimens of *Heliconiini* butterflies 447(1988)432
- Cazau, M., see Oliveros, L. 409(1987)183
- Cazenave, J.-P., see Leray, C. 420(1987)411
- Cecchini, D.J., Guan, K.L. and Giese, R.W.  
Staphylococcal nuclease HPLC characterization of diamino-octane-modified DNA and its biotin and fluorescein derivatives 444(1988)97

- Cedarbaum, J.M., Williamson, R. and Kutt, H.  
Simultaneous determination of levodopa, its metabolites and carbidopa in clinical samples 415(1987)393
- Čelap, M.B., see Janjić, T.J. 404(1987)307
- Černý, M., see Jegorov, A. 434(1988)417
- Cerutti, F., see Lacroix, C. 415(1987)85
- Ceva, P., see Pietta, P. 404(1987)279
- Chabard, J.L., see Bougerolle, A.M. 434(1988)232
- Chabenat, C., Ladure, P., Blanc-Continsouza, D., Boismare, F. and Boucly, P.  
Determination of calcium acetylhomotaurinate by LC with fluorimetric and electrochemical detection 414(1987)417
- Chabenat, C., see Tharasse-Bloch, C. 421(1987)407
- Chae, K., Levy, L.A. and Korach, K.S.  
Chromatographic separation and isolation of the enantiomers of diethylstilbestrol metabolites 439(1988)484
- Chai, B., Minkler, P.E. and Hoppel, C.L.  
Determination of ibuprofen and its major metabolites in human urine by HPLC 430(1988)93
- Chai, M., see Emm, T.A. 427(1988)162
- Chain, B.M., see Wallace, G.R. 427(1988)239
- Chait, B.T., see Burman, S. 443(1988)285
- Chakrabarti, S., see Colin, P. 413(1987)151
- Chakraborty, M., see Anderson, G.M. 431(1988)400
- Chambers, J.G., see Hubbard, R.W. 431(1988)163
- Chambon, P., see Candito, M. 430(1988)198
- Chan, G.L.-Y., Axelson, J.E., Abbott, F.S., Kerr, C.R. and McErlane, K.M.  
Determination of propafenone in biological fluids by fused-silica capillary GC using electron-capture detection 417(1987)295
- Chan, J.A., see Poehland, B.L. 439(1988)459
- Chan, K.  
Comparison of GC and HPLC assays for the determination of debrisoquine and its 4-hydroxy metabolite in human fluids 425(1988)311
- Chan, K., Miners, J.O. and Birkett, D.J.  
Direct and simultaneous HPLC assay for the determination of *p*-aminobenzoic acid and its conjugates in human urine 426(1988)103
- Chan, K., see Woo, J. 420(1987)73
- Chan, K.K. and Barrientos, A.  
Analysis of clomesone in plasma by GC-electrolytic conductivity detection 428(1988)331
- Chan, L.T.F., Chhuy, L.H. and Crowley, R.J.  
GC analysis of verapamil and norverapamil in *post mortem* specimens using a 530- $\mu$ m fused-silica column with nitrogen-phosphorus detection 402(1987)361
- Chan, S.-L., see Krahn, M.M. 437(1988)161
- Chandler, M.H. Huang, Guttendorf, R.J., Blouin, R.A. and Wedlund, P.J.  
Simultaneous quantitation of *d*- and *l*-hexobarbital in rat blood by HPLC 419(1987)426
- Chang, C.A., see Lin, G. 404(1987)328
- Chang, C.A., see Lin, G.-C. 409(1987)371
- Chang, C.A., Wu, Q. and Sheu, C.  
Determination of cations at the parts-per-billion level with non-suppressed ion chromatography and indirect photometric detection 404(1987)282
- Chang, D., see Oldfield, N. 422(1987)135
- Chang, F.C., see Iida, T. 438(1988)423
- Chang, S.-L., see Robbins, D.K. 415(1987)208
- Chang, S.F., see Welscher, T.M. 431(1988)438
- Chao, W.-F., Kai, M. and Ohkura, Y.  
HPLC determination of forphenicine in mouse serum and muscle by pre-column fluorescence derivatization using 1,2-diamino-4,5-ethylenedioxybenzene as fluorogenic reagent 430(1988)361
- Chaplin, L.C. and Andrews, A.T.  
Rapid preparative-scale purification of peptides derived from caseins using mass ion-exchange chromatography 450(1988)420
- Charbonneau, C.F., see Lawrence, J.F. 403(1987)379
- Charbonneau, C.F., see Lawrence, J.F. 407(1987)405
- Charbonneau, C.F., see Lawrence, J.F. 445(1988)189
- Charpin, P., see Coleman, A.W. 450(1988)175
- Charreau, E.H., see Santa-Coloma, T.A. 415(1987)297
- Chartier, A., Gonnet, C., Morel, D., Rocca, J.L. and Serpinet, J.  
Mixed bonded phases for HPLC. I. Synthesis and physico-chemical characterization 438(1988)263
- Chasseaud, L.F., see Brodie, R.R. 415(1987)423
- Chasseaud, L.F., see Wood, S.G. 416(1987)154
- Châtinier, W.M. Du, see De Jong, A.P. 419(1987)233
- Chaudhary, T., see Burman, S. 443(1988)285
- Chaumont, A.J., see Kintz, P. 432(1988)329
- Chawla, R.P., see Singh, P.P. 450(1988)452
- Cheam, V. and Li, E.X.  
Ion chromatographic determination of low level cadmium(II), cobalt(II) and manganese(II) in water 450(1988)361
- Cheetham, P.S.J., see Cookson, D. 402(1987)265

- Chekroun, I., see Krstulović, A.M. 411(1987)461
- Chen, A.J.C., see Hjertén, S. 403(1987)47
- Chen, E.C.M., see Wentworth, W.E. 441(1988)45
- Chen, F.-M., Naeve, G.S. and Epstein, A.L.  
Comparison of Mono O, Superose-6, and ABx fast protein LC for the purification of IgM monoclonal antibodies 444(1988)153
- Chen, K.-H., see Baraldi, R. 442(1988)301
- Chen, S., see Puglisi, C.V. 419(1987)253
- Chen, S., see Rubio, F. 421(1987)281
- Chen, S.S., see Hsu, D.-S. 420(1987)398
- Chen, T.-L., see Hsu, J.T. 404(1987)1
- Chen, T.-S., see Philip, T. 435(1988)113
- Chen, T.-S., see Philip, T. 442(1988)249
- Chen, T.K. and Thornton, D.  
Regio-selective derivatization of 1-[(3,5-difluorophenyl)methyl]-1,3-dihydro-2H-imidazole for GC analysis 450(1988)403
- Chen, Y.-F., see Meng, Q.C. 445(1988)29
- Chen, Y.-H., see Sagara, K. 409(1987)365
- Chen, Y.-Y., see Chriswell, C.D. 405(1987)213
- Chen, Z.M., see Latini, R. 424(1988)211
- Chen, Z.R., Siebert, D.M., Somogyi, A.A. and Bochner, F.  
Determination of pholcodine in biological fluids by HPLC with fluorescence detection 424(1988)170
- Cheng, B. and Kowal, J.  
Specific reversed-phase LC method for analysis of steroids in Y-1 adrenal cell cultures 432(1988)302
- Cheng, L.K., see Osman, M.A. 419(1987)452
- Cheng, M.C., see Smith, A.I. 416(1987)255
- Cheng, T.-L., Grochow, L.B., Hurowitz, L.A. and Brundrett, R.B.  
Determination of busulfan in human plasma by GC with electron-capture detection 425(1988)303
- Cheng, W. and Hollis, D.  
Flow-rate effect on elution volume in SEC 408(1987)9
- Chenieux, J.C., see Montagu, M. 409(1987)426
- Chernoborodova, L.S., see Manoilov, S.E. 440(1988)53
- Chersi, A., Trinca, M.L. and Muratti, E.  
Isolation of haem-bearing peptides of haem proteins by the use of adsorption chromatography 410(1987)463
- Chervet, J.P., see Humbert, H. 417(1987)319
- Cheung, S.-T. and Lin, W.-N.  
Simultaneous determination of methanol, ethanol, acetone, isopropanol and ethylene glycol in plasma by GC 414(1987)248
- Chevalier, M., see Pochard, M.-F. 409(1987)315
- Cheymol, G., see Garnier-Moiroux, A. 416(1987)183
- Cheymol, G., see Poirier, J.-M. 426(1988)431
- Cheymol, G., see Poirier, J.M. 421(1987)223
- Chhabra, B.R., see Dhillon, R.S. 435(1988)256
- Chhuy, L.H., see Chan, L.T.F. 402(1987)361
- Chiarotti, M., see De Giovanni, N. 428(1988)321
- Chichila, T.M.P., see Weidolf, L.O.G. 433(1988)9
- Chicz, R.M. and Regnier, F.E.  
Surface-mediated retention effects of subtilisin site-specific variants in cation-exchange chromatography 443(1988)193
- Chijiwa, K. and Nakayama, F.  
Simultaneous microanalysis of bile acids and cholesterol in bile by glass capillary GC 431(1988)17
- Chikai, T., see Goto, J. 415(1987)45
- Child, P., Aloe, M. and Mee, D.  
Separation and quantitation of fatty acids, sterols and bile acids in feces by GC as the butyl ester-acetate derivatives 415(1987)13
- Chimenti, F., see Gagliardi, L. 404(1987)267
- Chin, J.L.  
Melatonin and other indoles in rat pineal 428(1988)206
- Chioccare, F., see Bassoli, A. 444(1988)209
- Chioccare, F., see Bassoli, A. 446(1988)179
- Chiou, R., Stubbs, R.J. and Bayne, W.F.  
Determination of ivermectin in human plasma and milk by HPLC with fluorescence detection 416(1987)196
- Chiou, R., Stubbs, R.J. and Bayne, W.F.  
Improved method for the determination of theophylline in plasma and urine using HPLC 422(1987)281
- Chiou, S.-H. and Wang, K.-T.  
Simplified protein hydrolysis with methanesulphonic acid at elevated temperature for the complete amino acid analysis of proteins 448(1988)404
- Chiu, S.-H.L., Sestokas, E. and Taub, R.  
Application of two-dimensional HPLC in metabolism studies of ivermectin 433(1988)217
- Chiu, W.-Y., Yamada, K., Saito, A., Ogawa, H., Takagi, T. and Chung, T.G.  
Comparison of serum protein removal in haemodialysis therapy by partly protein-permeable haemodialysers 428(1988)25
- Cho, S.Y., see Nam, H.S. 448(1988)445
- Choi, K.E., Sinkule, J.A., Han, D.S., MacGrath, S.C., Daly, K.M. and Larson, R.A.  
HPLC assay for mitoxantrone in plasma using electrochemical detection 420(1987)81

- Choma, N., see Miwa, B.J. 431(1988)343
- Chouliotis, A., see Noël, D. 408(1987)129
- Chrétien, J.R., see Szymoniak, K. 404(1987)11
- Chrétien, J.R. and Szymoniak, K.  
Hierarchical ascending classification of Kováts retention indices as a means of indicating siloxane stationary phase specificities towards aliphatic carboxylic esters 406(1987)95
- Christensen, J.M., Rasmussen, K. and Køppen, B.  
Automatic headspace GC method for the simultaneous determination of trichloroethylene and metabolites in blood and urine 442(1988)317
- Christensen, J.M., see Køppen, B. 442(1988)325
- Christians, U., Zimmer, K.-O., Wonigeit, K. and Sewing, K.-F.  
Measurement of cyclosporin A and of four metabolites in whole blood by HPLC 413(1987)121
- Christie, R.B., see Muirhead, D.C. 416(1987)420
- Christie, W.W.  
Equivalent chain-lengths of methyl ester derivatives of fatty acids on GC. A reappraisal 447(1988)305
- Christie, W.W. and Morrison, W.R.  
Separation of complex lipids of cereals by HPLC with mass detection 436(1988)510
- Christman, R.F., see Jacobs, J.S.A. 450(1988)433
- Christofalo, P., see Carlin, J.R. 427(1988)79
- Christon, R., see Alessandri, J.-M. 432(1988)75
- Christophersen, A.S., Biseth, A., Skuterud, B. and Gadeholt, G.  
Identification of opiates in urine by capillary column GC of two different derivatives 422(1987)117
- Christophersen, A.S., Dahlin, E. and Pettersen, G.  
GC quantification of underivatized amphetamine in whole blood. Blood stored in plastic containers—a source of error 432(1988)290
- Christopherson, R.J., see Early, R.J. 416(1987)15
- Chriswell, C.D., Chen, Y.-Y., McGowan, C.W. and Markuszewski, R.  
Use of silicalite molecular sieve for the GC determination of permanent gases and volatile hydrocarbons emitted during coal processing 405(1987)213
- Chu, L.V., see Coutts, R.T. 421(1987)267
- Chuiko, A.A., see Lysjuk, L.S. 407(1987)189
- Chukwunye, C. and McAtee, Jr., J.L.  
Organo-modified smectite for the GC separation of various compounds 410(1987)121
- Chung, C.H., see Kee, Y. 428(1988)345
- Chung, H., see Zhou, Z.M. 414(1987)77
- Chung, T.G., see Chiu, W.-Y. 428(1988)25
- Churchill, F.C., see Bergqvist, Y. 427(1988)295
- Churchill, F.C., see Bergqvist, Y. 428(1988)281
- Churchill, F.C., see Bergqvist, Y. 432(1988)253
- Churchill, F.C., see Bergqvist, Y. 434(1988)1
- Churchill, F.C., see Mount, D.L. 423(1987)261
- Churchill, F.C., see Mount, D.L. 428(1988)196
- Churchill, F.C., see Williams, S.B. 433(1988)197
- Cingolani, L.  
Rapid GC determination of phytanic acid from serum of a patient suffering from Refsum's disease 419(1987)475
- Ciucanu, I., Pop, D. and Timoceanu, V.  
Derivatization of prostaglandins and related compounds to (methoxime) alkyl ester alkyl ether derivatives for GC analysis 436(1988)219
- Claessens, H.A., De Haan, J.W., Van de Ven, L.J.M., De Bruyn, P.C. and Cramers, C.A.  
Chromatographic and solid state NMR study of the changes in reversed-phase packings for HPLC at different eluent compositions 436(1988)345
- Claperon, N., see Laham, A. 440(1988)455
- Clark, B.J., see Fell, A.F. 434(1988)377
- Clark, C.P., Snider, B.G. and Bowman, P.B.  
HPLC method for determining the enantiomeric purity of a benzindene prostaglandin by a diastereomeric separation 408(1987)275
- Clark, M.A., see Strohsacker, M.W. 435(1988)185
- Clarke, G.S., see Sheridan, M.E. 434(1988)253
- Clarke, M.L., see Ivin, P.C. 408(1987)393
- Clarkson, J., see Awni, W.M. 419(1987)414
- Clavier, F., see Pussard, E. 421(1987)192
- Claycamp, H.G. and Ludwig, B.  
Simplified monitoring of reduced and oxidized dithiothreitol using HPLC 422(1987)239
- Clayton, B.E., see Powers, A.G. 432(1988)321
- Cli
- Clifton, P.M., MacKinnon, A.M. and Barter, P.J.  
Separation and characterization of high-density lipoprotein subpopulations by GPC 414(1987)25
- Clonis, Y.D.  
Matrix evaluation for preparative high-performance affinity chromatography 407(1987)179
- Coas, V., see Lepri, L. 405(1987)394



- Coassolo, P., see Al-Mallah, N.R. 421(1987)177
- Coates, P.E., see Brett, M.A. 419(1987)438
- Coccioli, F., see Coppi, S. 442(1988)97
- Cochet, S., see Kroviarski, Y. 449(1988)403
- Cochet, S., see Kroviarski, Y. 449(1988)413
- Coen, C.W., see Opacka-Juffry, J. 433(1988)41
- Coguic, A. Le, Bidault, R., Farinotti, R. and Dauphin, A.  
Determination of ofloxacin in plasma and urine by LC 434(1988)320
- Cohen, A.I., Jemal, M., Ivashkiv, E. and Ribick, M.  
Rationale for application of low-mass selected-ion detection in the quantitative body fluid assay of nadolol 416(1987)445
- Cohen, A.I., see Jemal, M. 408(1987)360
- Cohen, A.I., see Jemal, M. 428(1988)81
- Cohen, A.I., see Wang-Iverson, D. 408(1987)365
- Cohen, A.S., see Guttman, A. 448(1988)41
- Cohen, D.J., see Anderson, G.M. 424(1988)373
- Cohen, D.J., see Anderson, G.M. 431(1988)400
- Cohen, D.J., see Durkin, T.A. 428(1988)9
- Cohen, I.C., see Lumley, I.D. 408(1987)115
- Cohen, J.D., see Baraldi, R. 442(1988)301
- Cohen, K.A., see Eslami, M. 411(1987)121
- Cohen, P., see Créminon, C. 440(1988)439
- Cohen, S.M., see Tibbels, T.S. 441(1988)448
- Colacchio, T.A., see Nierenberg, D.W. 413(1987)79
- Colburn, W.A., see Ferry, J.J. 421(1987)187
- Colca, J.R., see McCroskey, M.C. 442(1988)307
- Cole, K.C., see Noël, D. 408(1987)129
- Cole, K.C., see Noël, D. 447(1988)141
- Cole, M.D., Fellows, L.E. and Romeo, J.T.  
Separation of hydroxy and polyhydroxy derivatives of 2-carboxy and 2-hydroxymethyl piperidine and pyrrolidine by HPLC 445(1988)295
- Cole, W.J., see Brooks, C.J.W. 438(1988)108
- Cole, W.J., see Brooks, C.J.W. 441(1988)13
- Coleman, A.W., Tsoucaris, G., Parrot, H., Galons, H., Miocque, M., Perly, B., Keller, N. and Charpin, P.  
Cyclodextrin-chiral substrate interactions. High field NMR and molecular graphics studies 450(1988)175
- Coleman, M.D., Edwards, G., Braithwaite, I.M. and Breckenridge, A.M.  
HPLC method for the determination of ampyroquinone in biological fluids 414(1987)242
- Colin, P., Sirois, G. and Chakrabarti, S.  
Rapid HPLC assay of acetaminophen in serum and tissue homogenates 413(1987)151
- Colistra, C., see Del Principe, D. 419(1987)329
- Collette, L.M., see Wenzel, T.J. 433(1988)149
- Collette, L.M., see Wenzel, T.J. 436(1988)299
- Collette, N., see Brassinne, C. 419(1987)401
- Collins, A.J., see Notarianni, L.J. 413(1987)305
- Collins, C.H., see Valente, A.L.P. 402(1987)349
- Collins, F.W. and Mullin, W.J.  
HPLC determination of avenanthramides, N-aroylanthranilic acid alkaloids from oats 445(1988)363
- Collins, J.M., see Belanger, K. 417(1987)57
- Collins-Nakai, R.L., see Pereira, C.M. 425(1988)208
- Colombo, D., see Anastasia, M. 410(1987)504
- Colussi, D., see Sioufi, A. 414(1987)131
- Colwell, R.K., see Boley, N.P. 410(1987)190
- Comellas, L., see Aceves, M. 436(1988)503
- Cometti, A., see Riva, E. 421(1987)99
- Čomor, J.J., see Todorović, M. 442(1988)105
- Compton, G., see Griffin, M. 431(1988)285
- Cone, E.J., see Thompson, L.K. 421(1987)91
- Công, H.N.G., Valencia, R., Becue, T. and Bertaux, O.  
Identification by HPLC and MS of a guanine derivative arising from acid hydrolysis of DNA 407(1987)349
- Coninck, M. De, see Verzele, M. 450(1988)47
- Connolly, K.M., see Pruden, D.J. 437(1988)399
- Connor, M., see Meaney, M. 449(1988)241
- Constanzer, M.L., see Matuszewski, B.K. 426(1988)295
- Constanzer, M.L. and Bayne, W.F.  
Determination of picogram levels of (*E*)-3-(9-chloro-6,11-dihydro-5H-pyrrolo(2,1-B) (3) benzazepin-11-ylidene)-N,N-dimethyl-1-propanamine (*Z*)-2-butenedioate (1:1) in plasma using capillary GC with nitrogen-selective detection 413(1987)277
- Conte, Jr., J.E. and Zurlinden, E.  
Column LC determination of cefpiramide in human serum and urine 417(1987)452
- Cook, C.E., see Nakahara, Y. 434(1988)247
- Cook, H.A., see Bianchi, A. 449(1988)175
- Cook, S., see Mazsaroff, I. 443(1988)119
- Cooke, M., see Burrows, R. 404(1987)248
- Cookson, D., Cheetham, P.S.J. and Rathbone, E.B.  
Preparative HPLC purification and structural determination of 1-O- $\alpha$ -D-glucopyranosyl-D-fructose (trehalulose) 402(1987)265
- Cooper, D.A., see Lurie, I.S. 405(1987)273
- Cooper, D.A., see Moore, J.M. 410(1987)297
- Cooper, J.K., see Srinivas, N.R. 433(1988)105
- Cooper, R., see Mierzwa, R. 436(1988)259
- Cooper, T.B., see Suckow, R.F. 427(1988)287
- Cooper, W.T., see Smith, P.L. 410(1987)249

- Coppella, S.J., Acheson, C.M. and Dhurjati, P.  
Isolation of high-molecular-weight nucleic acids for copy number analysis using HPLC 402(1987)189
- Coppi, G., see Springolo, V. 428(1988)173
- Coppi, S., Betti, A., Bigli, C., Cartoni, G.P. and Caccioli, F.  
Characterization of some styrene-divinylbenzene sorbents for reversed-phase chromatography 442(1988)97
- Corbett, M.E., Scrimgeour, C.M. and Watt, P.W.  
Use of *tert*-butyldimethylsilyl derivatives for GC-MS analysis of dipeptides 419(1987)263
- Corbin, F.T., see Parker, C.E. 438(1988)359
- Corcia, A. Di, Marchetti, M. and Samperi, R.  
Rapid determination of simazine and atrazine in water at the parts per trillion ( $10^{12}$ ) level. Extraction by a miniaturized Carbo-pack B trap followed by HPLC 405(1987)357
- Cormane, R.N., see Stolk, L.M.L. 423(1987)383
- Cornell, H.J. and Brady, P.H.  
Application of preparative reversed-phase HPLC to the isolation of insulin-like growth factor II from human serum 421(1987)61
- Corre, P. Le, Gibassier, D., Sado, P. and Le Verge, R.  
Simultaneous assay of disopyramide and monodesisopropylidisopyramide enantiomers in biological samples by LC 424(1988)424
- Corre, P. Le, Gibassier, D., Sado, P. and Le Verge, R.  
Direct enantiomeric resolution of disopyramide and its metabolite using chiral HPLC. Application to stereoselective metabolism and pharmacokinetics of racemic disopyramide in man 450(1988)211
- Corse, J., see Stafford, A.E. 436(1988)93
- Corti, F., Luzzani, F. and Ventura, P.  
Sulphate quantification in rabbit tracheobronchial mucus by reversed-phase ion-pair chromatography 424(1988)147
- Costa, M., see Murphy, R. 408(1987)388
- Costa, M., see Pecci, L. 426(1988)183
- Cote, L.M., see Swanson, S.P. 414(1987)335
- Cotroneo, A., Dugo, G., Favretto, L. and Favretto, L.G.  
GC separation and chemometric analysis of mandarin essential oils 449(1988)183
- Couder, J., see De Bondt, M. 442(1988)165
- Coughlin, R.T., see Samu, J. 435(1988)167
- Coune, A., see Brassinne, C. 419(1987)401
- Counis, R., see Starzec, A. 440(1988)353
- Čoupek, J., see Mikeš, O. 440(1988)287
- Čoupek, J., see Sanitrák, J. 415(1987)129
- Čoupek, J., see Tennikova, T.B. 435(1988)357
- Coupez, R., see Baltes, E. 430(1988)149
- Courtot, D., see Jaussaud, P. 423(1987)123
- Coutts, R.T., see Birkholz, D.A. 449(1988)251
- Coutts, R.T., see Mozayani, A. 423(1987)131
- Coutts, R.T., see Tam, Y.K. 423(1987)199
- Coutts, R.T., see Wong, J.T.F. 428(1988)140
- Coutts, R.T., Torok-Both, G.A., Chu, L.V., Tam, Y.K. and Pasutto, F.M.  
In vivo metabolism of lidocaine in the rat. Isolation of urinary metabolites as pentafluorobenzoyl derivatives and their identification by combined GC-MS 421(1987)267
- Couvreur, P., see Laham, A. 440(1988)455
- Cowman, M.K., see Hittner, D.M. 402(1987)149
- Cox, G.B., Antle, P.E. and Snyder, L.R.  
Preparative separation of peptide and protein samples by HPLC with gradient elution. II. Experimental examples compared with theory 444(1988)325
- Cox, G.B., see Eble, J.E. 405(1987)31
- Cox, G.B., see Snyder, L.R. 444(1988)303
- Cox, J.W. and Pullen, R.H.  
Irregular retention properties of 21-aminosteroid antioxidants in octylsilane bonded-phase HPLC 424(1988)285
- Cox, J.W. and Pullen, R.H.  
HPLC determination of a 21-aminosteroid antioxidant in plasma 424(1988)293
- Cox, Jr., T., see Alick, B. 430(1988)309
- Cozzi, R., see Polcaro, C. 448(1988)127
- Craft, A.W., see Seviour, J.A. 432(1988)273
- Cramer, S.M., see Subramanian, G. 439(1988)341
- Cramers, C.A., see Claessens, H.A. 436(1988)345
- Crane, S.W., see Zuk, M.M. 433(1988)367
- Crankshaw, D.P., see Kumar, K. 419(1987)464
- Craske, J.D., see Bannon, C.D. 404(1987)340
- Craske, J.D., see Bannon, C.D. 407(1987)231
- Craske, J.D., see Bannon, C.D. 447(1988)43
- Créminon, C., Rholam, M., Boussetta, H., Marrakchi, N. and Cohen, P.  
Synthetic peptide substrates as models to study a pro-ocytocin/neurophysin converting enzyme 440(1988)439
- Cress-Tijhuis, M.W., see Koopman-Kimenai, P.M. 416(1987)382
- Cresteil, T., see Beaune, P.H. 426(1988)169
- Crevat, A., see Gallice, P. 422(1987)263
- Crews, T., see Rubio, F. 421(1987)281
- Crimmins, D.L., Gorka, J., Thoma, R.S. and Schwartz, B.D.  
Peptide characterization with a sulfoethyl aspartamide column 443(1988)63
- Cristalli, M., see Fanali, S. 405(1987)385

- Cristofaro, R. De, Landolfi, R., Bizzi, B. and Castagnola, M.  
Human platelet glycolicin purification by phenyl boronate affinity chromatography coupled to anion-exchange HPLC 426(1988)376
- Cristofaro, R. De, see Castagnola, M. 440(1988)231
- Crockett, T.D., Webb, A.A., Borchardt, L.G. and Easty, D.B.  
Identification of polymers in paper by pyrolysis GC. Reduction of paper interference by prepyrolysis 407(1987)330
- Crolla, T., see Visconti, M. 416(1987)433
- Cromarty, J.A., see Blackett, A. 433(1988)326
- Crooke, S.T., see Strohsacker, M.W. 435(1988)185
- Crosby, S.S., see Unadkat, J.D. 430(1988)420
- Croteau, D., Vallée, F., Bergeron, M.G. and LeBel, M.  
HPLC assay of erythromycin and its esters using electrochemical detection 419(1987)205
- Croteau, R.B., see Satterwhite, D.M. 407(1987)243
- Crouch, Jr., F.W., Riley, C.M. and Stobaugh, J.F.  
HPLC of bis[1,2-bis(diphenylphosphino)ethane]gold(I) chloride, a potential antineoplastic agent 448(1988)333
- Crouzette, J., see Girre, C. 419(1987)363
- Crowley, R.J., see Chan, L.T.F. 402(1987)361
- Csákvári, B., see Szederkényi, F. 446(1988)253
- Cserhádi, T., Szejtli, J. and Fenyvesi, É.  
Reversed-phase TLC of some chlorophenols in the presence of a soluble  $\beta$ -cyclodextrin polymer 439(1988)393
- Cserhádi, T. and Somogyi, A.  
TLC separation of some tributylphenyl ethylene oxide oligomers according to the length of the ethylene oxide chain 446(1988)17
- Cserhádi, T. and Szógyi, M.  
Amino acid-amino acid interactions studied by charge-transfer chromatography 434(1988)455
- Csipo, I., see Nagy, E. 425(1988)214
- Cucchi, M.L., see Schinelli, S. 431(1988)150
- Cuisinaud, G., see Riva, E. 421(1987)99
- Cukrová, V. and Hrkál, Z.  
Purification and characterization of granulocyte-macrophage colony stimulating factor from human placenta 413(1987)242
- Cummings, J., Kerr, D.J., Kaye, S.B. and Smyth, J.F.  
Optimisation of a reversed-phase HPLC method for the determination of flavone acetic acid and its major human metabolites in plasma and urine 431(1988)77
- Cunico, R.L., see Kunitani, M.G. 443(1988)205
- Curini, R., see Laganá, A. 417(1987)135
- Currò, P., Micali, G. and Lanuzza, F.  
Determination of  $\beta$ -asarone, safrole, isosafrole and anethole in alcoholic drinks by HPLC 404(1987)273
- Curtis, M., see Pullen, R.H. 434(1988)271
- Curtis, M.A., see Koenigbauer, M.J. 413(1987)161
- Curtis, M.A., see Koenigbauer, M.J. 427(1988)277
- Curvale, C., see Jardy, A. 439(1988)137
- Cutié, S.S., Kallos, G.J. and Smith, P.B.  
Separation and characterization of acrylic acid oligomers by NMR spectroscopy and thermospray ion-exchange LC-MS 408(1987)349
- Cynober, L., see Aussel, C. 423(1987)270
- Czejka, M.J. and Georgopoulos, A.  
HPLC determination of adriablastin in human plasma, urine, saliva and liver punctuate by column switching for drug monitoring studies 424(1988)182
- Czejka, M.J. and Georgopoulos, A.  
Mitoxantron determination using HPLC: improved sensitivity by loop-column injection for dual-dose pharmacokinetic studies 425(1988)429
- Czejka, M.J. and Koch, H.P.  
Determination of thalidomide and its major metabolites by HPLC 413(1987)181
- Czuczwa, J., Leuenberger, C., Tremp, J., Giger, W. and Ahel, M.  
Determination of trace levels of phenol and cresols in rain by continuous liquid-liquid extraction and HPLC 403(1987)233
- ## D
- Dadgar, D. and Power, A.  
Applications of column-switching technique in biopharmaceutical analysis. I. HPLC determination of amitriptyline and its metabolites in human plasma 416(1987)99
- Dadgar, D. and Power, A.  
Applications of column-switching techniques

- in biopharmaceutical analysis. II. HPLC determination of tripeleennamine in bovine plasma and milk 421(1987)216
- D'Agostino, P.A., Provost, L.R. and Visentini, J. Analysis of O-ethyl S-[2-(diisopropylamino)ethyl] methyl-phosphonothiolate (VX) by capillary column GC-MS 402(1987)221
- D'Agostino, P.A. and Provost, L.R. GC retention indices of sulfur vesicants and related compounds 436(1988)399
- Dague, B., see Caprioli, R.M. 443(1988)355
- Dahiya, J.S. Reversed-phase HPLC of *Cajanus cajan* phytoalexins 409(1987)355
- Dahiya, J.S. and Rimmer, S.R. HPLC of phytoalexins in stem callus tissues of rapeseed 448(1988)448
- Dahlem, A.M., see Harada, K.-I. 448(1988)275
- Dahlen, D.T., see Wenzel, T.J. 433(1988)149
- Dahlin, E., see Christophersen, A.S. 432(1988)290
- Dahmen, W., Gielsdorf, W., Jeremic, B., Achtert, G. and Hausleiter, H.-J. Determination of eproxindine hydrochloride in human plasma by capillary GC 427(1988)157
- Daimon, E., see Nakano, K. 438(1988)61
- Dal Bo', L., see Ascalone, V. 414(1987)101
- Dal Bo', L., see Ascalone, V. 423(1987)239
- Dale, G., see Seviour, J.A. 432(1988)273
- Dale, G.L. Use of the dye Janus Green to facilitate quantitative analysis of phosphoinositides 424(1988)445
- Dalene, M., Mathiasson, L., Skarping, G., Sangö, C. and Sandström, J.F. Trace analysis of airborne aromatic isocyanates and related aminoisocyanates and diamines using HPLC with UV and electrochemical detection 435(1988)469
- Dalene, M., see Skarping, G. 435(1988)453
- Dalferes, Jr., E.R., see Lee, Y.J. 419(1987)275
- Dalgaard, L., see Hansen-Møller, J. 420(1987)99
- Dalgaard, L., see Køppen, B. 442(1988)325
- Dallenbach-Toelke, K., Nyiredy, S. and Sticher, O. Application of various planar chromatographic techniques for the separation of iridoid glycosides from *Veronica officinalis* 404(1987)365
- Dallenbach-Toelke, K., see Nyiredy, S. 450(1988)241
- Daly, K.M., see Choi, K.E. 420(1987)81
- Daly, R.E., see Geria, T. 450(1988)407
- Damm, P., see Lehr, K.-H. 425(1988)153
- Damsma, G. and Flentge, F. LC with electrochemical detection for the determination of choline and acetylcholine in plasma and red blood cells. Failure to detect acetylcholine in blood of humans and mice 428(1988)1
- Daneshvar, M.I. and Brooks, J.B. Improved procedure for preparation of pentafluorobenzyl derivatives of carboxylic acids for analysis by GC with electron-capture detection 433(1988)248
- Danhauser, L., see Gandhi, V. 413(1987)293
- Daniel, R.M., see Patchett, M.L. 425(1988)269
- Daniele, R., see Preti, G. 432(1988)1
- Danielson, T.J., see Mozayani, A. 423(1987)131
- Dann, O., see Gluth, W.P. 417(1987)331
- Däppen, R., see Pirkle, W.H. 404(1987)107
- Däppen, R., see Pirkle, W.H. 407(1987)211
- Dareer, S.M. El, see Rose, L.M. 425(1988)419
- Das, D.K., see Prasad, M.R. 428(1988)221
- D'Ascenzo, G., see Laganá, A. 417(1987)135
- DasGupta, B.R., see Woody, M.A. 430(1988)279
- Datta, P.K., Basu, P.S. and Datta, T.K. Purification of human erythrocytes specific lectins from rice bean, *Phaseolus calcaratus* syn. *Vigna umbellata*, by HPLC 431(1988)37
- Datta, T.K., see Datta, P.K. 431(1988)37
- Daučík, P., Rizzi, A.M. and Huber, J.F.K. Separation of isomeric mono-, di- and tributylphenols by HPLC on octadecylsilica with methanol-water using alkyl nitrile and alkylamine as surface modifiers 442(1988)53
- Daugherty, K.E., see Jen, J.-F. 436(1988)86
- Dauphin, A., see Le Coguic, A. 434(1988)320
- Daussant, J., see Lecommandeur, D. 441(1988)436
- Dautrevaux, M., see Maes, P. 445(1988)409
- Dautrevaux, M., see Sablonniere, B. 403(1987)183
- Davalli, P., see Desiderio, M.A. 419(1987)285
- Davídek, J., see Hajšlová, J. 438(1988)55
- Davídek, J., see Hajšlová, J. 439(1988)307
- Davídková, P. and Gasparič, J. Separation efficiency of adsorption and partition TLC and reversed-phase HPTLC with chemically bonded phases for alkylphenoxyalkanoic acids 410(1987)53
- Davidson, A.G., see Fadiran, E.O. 442(1988)363
- Davidson, C., see Lee, J.K. 444(1988)141
- Davies, M., see Bailey, E. 421(1987)21
- Davies, N.W. Open split interfaces in capillary GC-MS. Yield and quantitative aspects 450(1988)388
- Davis, A., see Dhingra, B.S. 447(1988)284

- Davis, A.T.  
Assay for trimethyllysine hydroxylase by HPLC 422(1987)253
- Davis, B.A.  
Effects of long-term storage on the concentrations of the unconjugated acidic metabolites of the trace amines, indoleamines and catecholamines 433(1988)23
- Davis, J.M.  
Experimental affirmation of the statistical model of overlap 449(1988)41
- Davis, L.J., see Hoskins, J.A. 426(1988)155
- Davis, L.J., see Pomaville, R.M. 438(1988)1
- Davis, P.C.  
Determination of xamoterol in human plasma by HPLC with electrochemical detection 417(1987)233
- Davis, P.J., see Miski, M. 437(1988)436
- Davoudi, H., see Szabo, G.K. 430(1988)112
- Dawidowicz, A.L., see Supryniewicz, Z. 446(1988)347
- Dawson, M., MacGee, C.M., Brooks, P.M., Vine, J.H., Lacey, E. and Watson, T.R.  
Quantitation of biphenylacetic acid in plasma and synovial fluid by GC-MS-MS 420(1987)129
- Dayer, P., see Padmalatha Devi, K. 434(1988)265
- D'Costa, M., see Diamandis, E.P. 426(1988)25
- DeAngelis, J.J., Barkley, R.M. and Sievers, R.E.  
Coupling LC with redox chemiluminescence detection 441(1988)125
- DeBackere, M., see Belbeke, F. 416(1987)443
- Debackere, M., see Delbeke, F.T. 426(1988)194
- DeBethizy, J.D., see Kyerematen, G.A. 419(1987)191
- De Boer, G., see Kubo, I. 402(1987)354
- De Bondt, M., Couder, J., Van der Auwera, L., Van Marsenille, M., Elseviers, M., Delaet, N., Laus, G., Tourwé, D. and Van Binst, G.  
Determination of the chiral purity of dipeptide isosteres containing a reduced peptide bond by GC analysis 442(1988)165
- De Bruijn, E.A., see Reeuwijk, H.J.E.M. 428(1988)93
- De Bruijn, E.A., see Tjaden, U.R. 420(1987)53
- De Bruyn, P.C., see Claessens, H.A. 436(1988)345
- De Castro, M.D.L., see Lázaro, F. 448(1988)173
- Decolin, D., Nicolas, A., Motassim, N. and Siest, G.  
Direct determination of debrisoquine and 4-hydroxydebrisoquine using a HPLC switching technique 419(1987)469
- Decolin, D., Nicolas, A. and Siest, G.  
Determination of flumequine and its 7-hydroxy metabolite by reversed-phase HPLC 414(1987)499
- Decolin, D., see Motassim, N. 422(1987)340
- De Coninck, M., see Verzele, M. 450(1988)47
- Décosterd, L.A., Dorsaz, A.-C. and Hostettmann, K.  
Application of semi-preparative HPLC to difficult natural product separations 406(1987)367
- De Cristofaro, R., Landolfi, R., Bizzi, B. and Castagnola, M.  
Human platelet glycoalcin purification by phenyl boronate affinity chromatography coupled to anion-exchange HPLC 426(1988)376
- De Cristofaro, R., see Castagnola, M. 440(1988)231
- De Galan, L., see Erkelens, C. 404(1987)67
- De Giovanni, N. and Chiarotti, M.  
Analysis of benzodiazepines. II. HPLC fluorescence detection after molecular rearrangement to acridanones 428(1988)321
- De Graaf, G.J. and Spierenburg, T.J.  
LC determination of cyadox in medicated feeds and in the contents of the porcine gastrointestinal tract with fluorescence detection 447(1988)244
- De Grazia, F., see Rubio, F. 421(1987)281
- Degrell, I., see Nagy, E. 425(1988)214
- De Good, P.M.R.M., see Vree, T.B. 417(1987)458
- De Haan, J.W., see Claessens, H.A. 436(1988)345
- De Jesus, A.E., Gorst-Allman, C.P., Horak, R.M. and Vleggaar, R.  
Large-scale purification of the mycotoxins aflatoxin B<sub>1</sub>, B<sub>2</sub> and G<sub>1</sub> 450(1988)101
- De Jong, A.P., Wittebrood, A.J., Du Châtinier, W.M. and Bron, J.  
LC analysis of alizapride and metoclopramide in human plasma and urine using solid-phase extraction 419(1987)233
- De Jong, G.J., see Irth, H. 424(1988)95
- De Jong, G.J., see Irth, H. 439(1988)63
- De Jong, G.J., see Maris, F.A. 435(1988)297
- De Jong, G.J., see Maris, F.A. 445(1988)129
- De Jong, J., Point, A.J.F., Tjaden, U.R., Beeksmas, S. and Kraak, J.C.  
Determination of catecholamines in urine (and plasma) by LC after on-line sample pretreatment on small alumina or dihydroxyborylsilica columns 414(1987)285
- De Jong, J., Tjaden, U.R., Visser, E. and Meijer, W.H.  
Determination of serotonin and 5-

- hydroxyindoleacetic acid in urine by reversed-phase ion-pair partition chromatography with fluorimetric detection 419(1987)85
- De Jong, P., see Odink, J. 431(1988)309
- De Korte, D., Marijnen, Y.M.T., Haverkort, W.A., Van Gennip, A.H. and Roos, D. Sensitive on-line radioactivity measurements with a heterogeneous flow-cell: application to lymphoid cell ribonucleotides separated by HPLC 415(1987)383
- De Kruijf, N., Rijk, M.A.H., Pranoto-Soetardhi, L.A. and Schouten, A. Determination of preservatives in cosmetic products. I. TLC procedure for the identification of preservatives in cosmetic products 410(1987)395
- Delaet, N., see De Bondt, M. 442(1988)165
- Delaney, M.F., Papas, A.N. and Walters, M.J. Chemometric classification of reversed-phase HPLC columns 410(1987)31
- Del Arco, A., see Usero, J.L. 407(1987)355
- Delattre, J., see Laham, A. 440(1988)455
- Delbeke, F.T., Debackere, M., Desmet, N. and Maertens, F. Qualitative GC and GC-MS screening for  $\beta$ -blockers in urine after solid-phase extraction using Extrelut-1 columns 426(1988)194
- Delee, E., Jullien, I., Le Garrec, L., Loupy, A., Sansoulet, J. and Zaparucha, A. Chiral Michael additions of acetamidomalonate to  $\alpha$ -enones. NMR and HPLC resolution of enantiomers 450(1988)183
- Delee, E., Jullien, I. and Le Garrec, L. Direct HPLC resolution of dihydropyridine enantiomers 450(1988)191
- De Leenheer, A.P., Nelis, H.J., Lambert, W.E. and Bauwens, R.M. Chromatography of fat-soluble vitamins in clinical chemistry (Review) 429(1988)3
- De Leenheer, A.P., see Lambert, W.E. 427(1988)195
- De Leer, E.W.B., see Erkelens, C. 404(1987)67
- De Leeuw, J.W., see Sinninghe Damsté, J.S. 435(1988)435
- De Ligny, C.L., see Brandts, P.M. 437(1988)337
- De Ligny, C.L., see Brandts, P.M. 438(1988)181
- Della Ragione, F., Oliva, A., Gragnaniello, V., Fioretti, M., Fioretti, A., Menna, L.F., Papparella, V. and Zappia, V. Chromatographic and radioimmunological methods for the determination of 5'-deoxy-5'-methylthioadenosine in biological fluids 440(1988)141
- Dellinger, M. and Brault, D. Normal-phase HPLC of free acid dicarboxylic porphyrins and hematoporphyrin derivative on silica 422(1987)73
- Del Principe, D., Colistra, C., Fiorentino, A.M., Spagnolo, G., Mancuso, G. and Menichelli, A. Detection of fibrinogen heterogeneity by capillary isotachopheresis 419(1987)329
- Delrieu, P., see Inchauspe, G. 404(1987)53
- Delucas, L.J., see Meng, Q.C. 445(1988)29
- Deluccia, F.J., see Lee, J.K. 444(1988)141
- De Maack, F., see Guichardant, M. 425(1988)25
- Demange, P., Abdallah, M.A. and Frank, H. Assignment of the configurations of the amino acids in peptidic siderophores 438(1988)291
- Dembitsky, V.M. Quantification of plasmalogen, alkylacyl and diacyl glycerophospholipids by micro-TLC 436(1988)467
- Demby, L., see Mathes, L.E. 432(1988)346
- De Mey, C., see Brendel, E. 426(1988)406
- De Miguel, I., Puech-Costes, E. and Samain, D. Use of mixed perfluorinated ion-pairing agents as solvents in ion-pair HPLC for the preparative purification of aminoglycoside antibiotics 407(1987)109
- De Miranda, P., see Good, S.S. 431(1988)123
- De Moerloose, P., see De Schutter, J.A. 428(1988)301
- De Moerloose, P., see De Schutter, J.A. 437(1988)83
- De Moerloose, P., see De Schutter, J.A. 450(1988)337
- Demole, E., see Ohloff, G. 406(1987)181
- Demotes-Mainard, F., Vinçon, G., Jarry, C., Necciari, J. and Albin, H. Micromethod for the determination of cefpiramide in human plasma and urine by HPLC using automated column switching 419(1987)388
- Dempewolf, J., see Bours, J. 403(1987)336
- Den, T.G. and Lee, S.L. Stationary phases. XV. HPLC behaviour of quinones on a saffrole stationary phase 408(1987)323
- Den Bosch, D.V., see Van Ginkel, L.A. 445(1988)385
- Denneberg, T., see Kågedal, B. 417(1987)261
- Dennis, M.J., see Mallett, D.N. 428(1988)190
- Denouel, J., see Humbert, H. 417(1987)319

- Denouel, J., see Humbert, H. 422(1987)205
- DePace, A., see Verebey, K. 427(1988)151
- De Pol, H.V., see Veenstra, J. 431(1988)413
- Derendorf, H., see Townsend, R.W. 450(1988)414
- Der Haar, J.V., Kip, J. and Kraak, J.C.  
Effect of the mobile phase composition and ligand structure on the separation of D- and L-dansylamino acids, as mixed metal complexes, by reversed-phase HPLC 445(1988)219
- Derks, H.J.G.M., Van Heiningen, A., Klaassen, R. and Olling, M.  
HPLC method for the determination of 2-nitropropane in rat plasma 442(1988)436
- Derlagen, H., see Baumann, R.A. 439(1988)165
- Derler, H., Hörmeyer, H.F. and Bonn, G.  
HPLC analysis of oligosaccharides. I. Separation on an ion-exchange stationary phase of low cross-linking 440(1988)281
- Der Linden, P.J.Q.V., see Gladdines, M.M. 431(1988)317
- Der Meeren, P.V., Vanderdeelen, J., Huys, M. and Baert, L.  
Simple and rapid method for HPLC separation and quantification of soybean phospholipids 447(1988)436
- Dernoncour, R. and Azerad, R.  
HPLC separation of the enantiomers of substituted 2-aryloxypropionic acid methyl esters 410(1987)355
- De Robertis, E., see Peña, C. 431(1988)192
- Der Torre, H.V., see Veenstra, J. 431(1988)413
- De Ruiter, C., Otten, R.R., Brinkman, U.A.T. and Frei, R.W.  
Rapid and simple dansylation of phenolic steroids using a two-phase system and phase transfer catalysis 436(1988)429
- De Ruiter, C., see Broersen, J.J. 436(1988)39
- De Ruiter, R., see Stolk, L.M.L. 423(1987)383
- Der Wart, H.H.F.V., see Schellens, J.H.M. 431(1988)203
- Desage, M., see Benckekroun, Y. 420(1987)287
- Desai, N.D., see Bhatia, S.C. 434(1988)288
- Desai, N.D., see Bhatia, S.C. 434(1988)296
- Desai, N.D., see Bhatia, S.C. 434(1988)303
- De Saint Blanquat, G., see Leitao, J. 435(1988)229
- Deschamps, N., see Ascalone, V. 414(1987)101
- De Schutter, J.A., Van der Weken, G., Van den Bossche, W. and De Moerloose, P.  
Determination of S-carboxymethylcysteine in serum by reversed-phase ion-pair LC with column switching following pre-column derivatization with  $\alpha$ -phthalaldehyde 428(1988)301
- De Schutter, J.A. and De Moerloose, P.  
Polar contributions of the stationary phase to the reversed-phase ion-pair HPLC separation of quaternary ammonium drugs 437(1988)83
- De Schutter, J.A. and De Moerloose, P.  
Determination of oxybutinin chloride in pharmaceuticals by reversed-phase ion-pair LC with two counter-ions in the eluent 450(1988)337
- Deshusses, J., see Nobile, S. 449(1988)331
- Desideri, P.G., see Lepri, L. 405(1987)394
- Desiderio, D.M., see Liu, D. 422(1987)61
- Desiderio, M.A., Davalli, P. and Perin, A.  
Simultaneous determination of  $\gamma$ -aminobutyric acid and polyamines by HPLC 419(1987)285
- Desmadril, M., see Yon, J.M. 440(1988)421
- Desmarchelier, J.M., see Brayon, J.G. 447(1988)249
- De Smet, M., Van Belle, S.J.P., Seneca, V., Storme, G.A. and Massart, D.L.  
Determination of vinblastine in MO<sub>4</sub> mouse fibrosarcoma cells by HPLC 416(1987)375
- De Smet, M. and Massart, D.L.  
Retention behaviour of acidic, neutral and basic drugs on a CN column using phosphate buffers in the mobile phase 410(1987)77
- Desmet, N., see Delbeke, F.T. 426(1988)194
- Desta, B.  
Separation of digoxin from dihydrodigoxin and the other metabolites by HPLC with post-column derivatization 421(1987)381
- Desta, B.  
Complete separation of nine equine oestrogens by HPLC 435(1988)385
- DeVane, C.L., see Miller, R.L. 430(1988)180
- Devaux, P., see Tillier, C. 441(1988)406
- De Vries, J.X., Walter-Sack, I. and Ittensohn, A.  
Analysis of benzbromarone in human plasma and urine by HPLC and GC-MS 417(1987)420
- De Waal, W.A.J., Maessen, F.J.M.J. and Kraak, J.C.  
Analysis of platinum species originating from *cis*-diamminedichloroplatinum(II) (cisplatin) in human and rat plasma by HPLC with on-line inductively coupled plasma atomic emission spectrometric detection 407(1987)253
- Dewaele, C., see Verzele, M. 450(1988)47
- DeWald, D.B., see Pearson, J.D. 418(1987)245
- Dewhurst, W.G., see Drebit, R. 432(1988)334
- De Wit, J.S.M. and Jorgenson, J.W.  
Photoionization detector for open-tubular LC 411(1987)201

- Dey, P.M., see Šecová, E. 436(1988)59
- Deyl, Z., Horáková, M. and Macek, K.  
Column LC separation of collagen  $\alpha$ -chain  
polymers on sepiolite 426(1988)162
- Deyl, Z., see Štol, M. 419(1987)308
- Deyman, M.E., see Kuhn, R.W. 421(1987)123
- De Zeeuw, R.A., see Benecke, R. 417(1987)287
- Dhariwal, K.R., see Goren, M.B. 440(1988)487
- Dharmaratne, H.R.W., see Njar, V.C.O.  
440(1988)415
- Dhillon, R.S., Singh, J., Gautam, V.K. and  
Chhabra, B.R.  
Detection of aminophenols and aromatic  
amines on thin-layer plates containing nitrite  
435(1988)256
- Dhillon, S., see Blackett, A. 433(1988)326
- Dhingra, B.S. and Davis, A.  
Determination of free ellagic acid by  
reversed-phase HPLC 447(1988)284
- Dhurjati, P., see Coppella, S.J. 402(1987)189
- Diamandis, E.P. and D'Costa, M.  
Selective determination of urinary free  
cortisol by LC after solid-state extraction  
426(1988)25
- Dibussolo, J.M., see Sagliano, Jr., N.  
443(1988)155
- Dickinson, R.G., see Hooper, W.D. 416(1987)347
- Di Corcia, A., Marchetti, M. and Samperi, R.  
Rapid determination of simazine and atrazine  
in water at the parts per trillion ( $10^{12}$ ) level.  
Extraction by a miniaturized Carboxpack B  
trap followed by HPLC 405(1987)357
- Di Corcia, A., see Caccamo, F. 434(1988)61
- Diehl, H.A., see Düren, R. 445(1988)49
- Diemann, E., see Weiss, J. 439(1988)91
- Dierdorf, H.-D., see Brett, M.A. 419(1987)438
- Dietel, P., Hölzel, C., Schneider, K. and Spittler,  
G.  
Excretion of urinary acids during inverted  
sleep-waking rhythm 422(1987)1
- Dieterle, W., see Kaiser, G. 419(1987)123
- Dietrich, A.M., Millington, D.S. and Seo, Y.-H.  
Specific identification of synthetic organic  
chemicals in river water using liquid-liquid  
extraction and resin adsorption coupled with  
electron impact, chemical ionization and  
accurate mass measurement GC-MS analyses  
436(1988)229
- DiGiovanni, J.H., see Kapetanovic, I.M.  
419(1987)458
- Di Gregorio, G., see Bassoli, A. 444(1988)209
- Di Gregorio, G., see Bassoli, A. 446(1988)179
- Di Iorio, M., see Porcelli, G. 414(1987)423
- Dijck, J. Van, see Vindevogel, J. 447(1988)297
- Dilli, S., see Brayán, J.G. 447(1988)249
- Dilli, S., see Robards, K. 411(1987)1
- Dily, J. Le, see Duchene, P. 424(1988)205
- Dimitrov, C., see Gavrilova, T.B. 402(1987)87
- Dimitrova, B. and Budevsky, O.  
Investigation of an alkylamine modified and  
pH-controlled mobile phase for separation of  
pyrazolone derivatives by HPLC 409(1987)81
- Dimitrova, M.P., see Rachev, R.C. 437(1988)287
- Dinan, L.  
TLC of ecdysone acyl esters and their 2,3-  
acetonide derivatives 411(1987)379
- Dinan, L.  
Separation of ecdysone acyl esters and their  
2,3-acetonide derivatives by HPLC  
436(1988)279
- Ding, S.F., see Peng, C.T. 436(1988)137
- Dinh, T. Le, see Motassim, N. 422(1987)340
- Dionisotti, S., Bamonte, F., Gamba, M. and  
Ongini, E.  
HPLC determination of netilmicin in guinea-  
pig and human serum by  
fluorodinitrobenzene derivatization with  
spectrophotometric detection 434(1988)169
- Diop, A., see Jardy, A. 439(1988)137
- Dios, G.C., see Sánchez-Rasero, F. 447(1988)426
- Diquet, B., see Bohbot, M. 416(1987)414
- Diquet, B., see Joulin, Y. 425(1988)402
- Dirks, R.J.M., see Wuis, E.W. 420(1987)212
- Ditschuneit, H., see Swobodnik, W. 423(1987)75
- Dix, K.D. and Fritz, J.S.  
Simple steam distillation for sample  
preparation prior to GC determination of  
organic compounds 408(1987)201
- Dixon, J.S., see Astbury, C. 414(1987)223
- Dmitrenko, A.V., see Shadrina, N.E.  
404(1987)183
- Doare, L., see Bohbot, M. 416(1987)414
- Dobbs, N.A. and James, C.A.  
Estimation of doxorubicin and doxorubicinol  
by HPLC and advanced automated sample  
processor 420(1987)184
- Döbeli, H., see Hochuli, E. 411(1987)177
- Dobranksý, T., Polívka, Ľ., Haas, I., Sova, O.  
and Petrálský, E.  
Purification of  $\alpha$ -amylase by industrial  
autofocusing 411(1987)486
- Dobrev, A.A., Ader, J.C. and Lattes, A.  
HPLC analysis of the *endo*-*exo* isomer ratio  
in the 5-substituted 2-norbornene series  
445(1988)268
- Doherty, R.M., see Abraham, M.H. 409(1987)15
- Dohnal, J., see Jelinek, I. 435(1988)496
- Dohnal, J., see Snopek, J. 450(1988)373
- Dolan, S., see Lau, C.E. 416(1987)212
- Dolan, S., see Lau, C.E. 423(1987)251



- Dollery, C.T., see Richmond, R. 417(1987)241
- Dombrovskis, D., see Remmel, R.P. 432(1988)388
- Dombrovskis, D., see Singh, A.K. 404(1987)223
- Donaldson, J., see Workman, P. 422(1987)315
- Doner, L.W., Irwin, P.L. and Kurantz, M.J.  
Preparative chromatography of oligogalacturonic acids 449(1988)229
- Dong, M.W., Lepore, J. and Tarumoto, T.  
Factors affecting the ion-pair chromatography of water-soluble vitamins 442(1988)81
- Donnelly, D.M.X., Martin, M. and Polonsky, J.  
HPLC determination of fomannonin in European strains of *Heterobasidion annosum* (Fr.) Bref. 440(1988)65
- Doormaal, J.J. Van, see Hoving, E.B. 434(1988)395
- Doornbos, D.A., see Smilde, A.K. 410(1987)1
- Dordain, G., see Bougerolle, A.M. 434(1988)232
- Dorsaz, A.-C., see Décosterd, L.A. 406(1987)367
- Dorulla, G.K., see Riggin, R.M. 435(1988)307
- Doshi, K.J., see Bhatia, S.C. 434(1988)296
- Dossena, A., see Armani, E. 441(1988)275
- Dossena, A., see Armani, E. 441(1988)287
- Douse, J.M.F.  
Improved method for the trace analysis of explosives by silica capillary column GC with thermal energy analysis detection 410(1987)181
- Douse, J.M.F.  
Trace analysis of explosives by capillary supercritical fluid chromatography with thermal energy analysis detection 445(1988)244
- Dower, H.J., see Brown, E.M. 443(1988)247
- Doyle, E., Pearce, J.C., Picot, V.S. and Lee, R.M.  
Analysis of drugs from biological fluids using disposable solid phase columns 411(1987)325
- Drabowicz, J., Kotyński, A. and Kudzin, Z.H.  
Selective detection of sulphoxides and sulphimides by TLC using trifluoroacetic anhydride-sodium iodide as a reagent 447(1988)225
- Drager, R.R. and Regnier, F.E.  
Retention mechanism of lactate dehydrogenase in anion-exchange chromatography 406(1987)237
- Draviam, E.J., Greeley, Jr., G.H., Lluís, F., Uchida, T., Trowbridge, J. and Thompson, J.C.  
Characterization of fat-induced neurotensin-like immunoreactivity in plasma using column LC and radioimmunoassay 414(1987)180
- Drayer, D.E., see Wu, D.-F. 433(1988)141
- Drebit, R., Baker, G.B. and Dewhurst, W.G.  
Determination of maprotiline and desmethylmaprotiline in plasma and urine by GC with nitrogen-phosphorus detection 432(1988)334
- Drenth, B.F.H., see Frijlink, H.W. 415(1987)325
- Dreux, M., see Lafosse, M. 404(1987)95
- Drijkoningen, G., see Koopman-Kimenai, P.M. 416(1987)382
- Driscoll, J.N., Duffy, M. and Pappas, S.  
Capillary GC analysis with the far-UV absorbance detector 441(1988)63
- Driscoll, J.N., see Duffy, M. 441(1988)73
- Driskell, J.A., see McChrisley, B. 428(1988)35
- Driskell, W.J., see Sowell, A.L. 431(1988)424
- Drissel, D., see Achari, R. 424(1988)430
- Drozd, J., see Koupil, P. 425(1988)99
- Drozd, J., see Vejrosta, J. 447(1988)170
- Drozdowicz, B., see Rotocki, P. 446(1988)329
- D'Souza, M.J., Zemaitis, M.A., Burckart, G.J. and Venkataraman, R.  
Simple HPLC method for simultaneous determination of aminopyrine and its metabolites 421(1987)198
- Dubé, L.M., Beaudoin, N., Lalande, M. and McGilveray, I.J.  
Determination of nalbuphine by HPLC with electrochemical detection: application to clinical samples from postoperative patients 427(1988)113
- Dubé, L.M., Mousseau, N. and McGilveray, I.J.  
HPLC determination of diltiazem and four of its metabolites in plasma: evaluation of their stability 430(1988)103
- Dube, P., see Holland, O.B. 426(1988)148
- Dubiel, W., see Toguzov, R.T. 434(1988)447
- Dubois, J.-P., see Kaiser, G. 419(1987)123
- Dubois, J.P., see Godbillon, J. 427(1988)269
- Dubois, J.P., see Menge, G.P. 414(1987)477
- Dubois, J.P., see Sioufi, A. 413(1987)101
- Dubois, J.P., see Sioufi, A. 414(1987)131
- Dubois, J.P., see Sioufi, A. 428(1988)71
- Dubois, J.P., see Sioufi, A. 434(1988)239
- Dubois, J.P., see Sioufi, A. 450(1988)221
- Dubruc, C., Hermann, P., Haddouche, A., Badarani, M.M. and Thénot, J.P.  
Determination of S-carboxymethyl-L-cysteine in plasma by HPLC with column switching following precolumn derivatization 417(1987)208
- Dubs, R., see Schilling, B. 435(1988)204
- Du Châtinier, W.M., see De Jong, A.P. 419(1987)233
- Duche, J.-C., Barre, J. and Tillement, J.-P.  
Rapid LC determination of debrisoquine and its hydroxy metabolite in human urine to

- define hydroxylation phenotypes  
423(1987)340
- Duchene, P., Le Dily, J., Bromet-Petit, M.,  
Mosser, J. and Feniou, C.  
HPLC assay of the metabolites of  
thymoxamine 424(1988)205
- Ducrocq, R., Le Bonniec, B., Carlier, O., Assan,  
R. and Elion, J.  
Measurement of glycosylated albumin in diabetic  
patients by biospecific affinity  
chromatography 419(1987)75
- Duda, C.T., see Pascoe, G.A. 414(1987)440
- Duerden, J.M., see Harvey, D.J. 414(1987)253
- Dufek, P., Vobecký, M., Holík, J. and Valášek, J.  
Separation of some rare earths and  
transplutonium elements by HPLC on Ostion  
cation exchanger 435(1988)249
- Duff, G.A., Yeager, S.A., Singhal, A.K., Pestel,  
B.C., Ressler, J.M. and Foster, N.  
Separation of metalloporphyrins from  
metallation reactions by LC and  
electrophoresis 416(1987)71
- Duffy, M., Driscoll, J.N., Pappas, S. and Sanford,  
W.  
Analysis of ppb levels of organics in water by  
means of purge-and-trap, capillary GC and  
selective detectors 441(1988)73
- Duffy, M., see Driscoll, J.N. 441(1988)63
- Dugo, G., see Cotroneo, A. 449(1988)183
- Duinker, J.C., see Petrick, G. 435(1988)241
- Duke, C.C. and Holder, G.M.  
*endo*-1,4,5,6,7,7-  
Hexachlorobicyclo[2.2.1]hept-5-ene-2-carboxylic  
acid, a superior resolving agent for the HPLC  
separation of enantiomers of hydroxylated  
derivatives of two azaaromatic hydrocarbons  
430(1988)53
- Duke, S.O., see Lydon, J. 410(1987)474
- Dumont, M., see Lemaire, S. 425(1988)77
- Dumont, P., see Goncalves, J.C.S. 411(1987)472
- Duncan, G.F., see Sciacca, M.A. 428(1988)265
- Duncan, J.R., see Perkins, D.M. 405(1987)319
- Dunn, M.J.  
Two-dimensional gel electrophoresis of  
proteins (Review) 418(1987)145
- Dunphy, M.J., Bhide, M.V. and Smith, D.J.  
Determination of hydroxyproline in tissue  
collagen hydrolysate by derivatization and  
isocratic reversed-phase HPLC 420(1987)394
- Dupin, P., see Inchauspe, G. 404(1987)53
- Dupont, A.G., see Gerlo, E. 414(1987)301
- Duprat, F., see Gassend, R. 404(1987)87
- Durden, D.A. and Boulton, A.A.  
Analysis of tryptamine at the femtomole level  
in tissue using negative ion chemical  
ionization GC-MS 440(1988)253
- Düren, R. and Diehl, H.A.  
Quantitative structure-activity relationship of  
coumarin derivatives. Involvement of  
partition between aqueous and membrane  
phase 445(1988)49
- Durkin, T.A., Anderson, G.M. and Cohen, D.J.  
HPLC analysis of neurotransmitter amino  
acids in brain 428(1988)9
- Durkin, T.A., see Anderson, G.M. 424(1988)373
- Durkin, T.A., see Anderson, G.M. 431(1988)400
- Durozard, D. and Baverel, G.  
Gas chromatographic method for the  
measurement of sodium valproate utilization  
by kidney tubules 414(1987)460
- Durso, R., see Szabo, G.K. 430(1988)112
- Durussel, J.J., see Laham, A. 440(1988)455
- Dutrizac, J.E., see Tan, L.K. 405(1987)247
- Dutta, I., O'Donovan, G.A. and Smith, D.W.  
Nucleotide extraction and quantitation from  
tomato roots by HPLC 444(1988)183
- Duval, M., see Godbillon, J. 427(1988)269
- Dwivedi, T.S., see Adinarayana, M. 435(1988)210
- Dyr, J.E. and Suttmar, J.  
On the increased retention volume of human  
haemoglobin in high-performance gel  
filtration 408(1987)303
- Dysken, M.W., see Vatassery, G.T. 433(1988)312
- Dzido, T.  
Modification of retention of some alkaloids  
in the system silanized silica/methanol +  
water + di(2-ethylhexyl)orthophosphoric acid  
439(1988)257
- Dziwiński, E., see Szewczyk, H. 410(1987)447
- Dziwiński, E., see Szewczyk, H. 446(1988)109
- Dziwiński, E. and Hetper, J.  
GC-MS of C<sub>1</sub>-C<sub>10</sub> alkyl benzyl succinates  
446(1988)103

## E

- Earl, V., see Alick, B. 430(1988)309
- Early, R.J., Thompson, J.R., Sedgwick, G.W.,  
Kelly, J.M. and Christopherson, R.J.  
Capillary GC analysis of amino acids in blood  
and protein hydrolysates as *tert*-  
butyldimethylsilyl derivatives 416(1987)15

- Easton-Taylor, M., see Ferry, J.J. 421(1987)187
- Easty, D.B., see Crockett, T.D. 407(1987)330
- Ebermann, R. and Pichorner, H.  
Improved method for the detection of peroxidase isoenzymes after electrophoretic separation with chlorophyllin, cysteamine and light as a hydrogen peroxide generating system 410(1987)466
- Ebert, K., see Voelkel, A. 409(1987)29
- Ebert, R.F. and Schmelzer, C.H.  
Isolation and comparative peptide mapping of fibrinogen subunits by reversed-phase HPLC 443(1988)309
- Ebinger, G., see Herregodts, P. 421(1987)51
- Eble, J.E., Grob, R.L., Antle, P.E., Cox, G.B. and Snyder, L.R.  
Simplified description of HPLC separation under overload conditions, based on the Craig distribution model. IV. Comparison of the model with experimental data for samples containing two or more solutes 405(1987)31
- Eble, J.E., Grob, R.L., Antle, P.E. and Snyder, L.R.  
Simplified description of HPLC separation under overload conditions, based on the Craig distribution model. III. Computer simulations for two co-eluting bands assuming a Langmuir isotherm 405(1987)1
- Eble, J.E., Grob, R.L., Antle, P.E. and Snyder, L.R.  
Simplified description of HPLC separation under overload conditions, based on the Craig distribution model. V. Gradient elution separation 405(1987)51
- Echizen, H., Itoh, R. and Ishizaki, T.  
Microassay of free 3-methoxy-4-hydroxyphenylglycol in plasma using HPLC with electrochemical detection 426(1988)351
- Echizen, H., see Takahashi, H. 419(1987)243
- Eckenhausen, M., see Powell, D.R. 420(1987)163
- Eckerbom, S., see Bergqvist, Y. 427(1988)295
- Eckerlin, R.H., see Hsu, S.-H. 424(1988)219
- Eckhardt-Wallasch, H., see Wallasch, T.M. 425(1988)175
- Eddib, O., see Burrows, R. 404(1987)248
- Edholm, L.-E., Lindberg, C., Paulson, J. and Walhagen, A.  
Determination of drug enantiomers in biological samples by coupled column LC and LC-MS 424(1988)61
- Edkins, R.D., see Hoffman, D.W. 414(1987)504
- Edkins, T.J., see Shelly, D.C. 411(1987)185
- Edlund, P.O.  
Determination of coenzyme Q<sub>10</sub>,  $\alpha$ -tocopherol and cholesterol in biological samples by coupled-column LC with coulometric and UV detection 425(1988)87
- Edwards, G., see Coleman, M.D. 414(1987)242
- Edwards, G., see Milton, K.A. 433(1988)339
- Efthymiopoulos, C., Staveris, S., Weber, F., Koffel, J.C. and Jung, L.  
Simultaneous quantitative determination of tertatolol and its hydroxylated metabolite in human plasma and urine by GC-MS 421(1987)360
- Egeler, E., see Jacob, K. 441(1988)171
- Egger, H.-J. and Fischer, G.  
Determination of the monocyclic  $\beta$ -lactam antibiotic carumonam in plasma and urine by ion-pair and ion-suppression reversed-phase HPLC 420(1987)357
- Egger, R.J., see Odink, J. 431(1988)309
- Eggers, H., see Geschke, R. 420(1987)111
- Egorin, M.J., Zuhowski, E.G., Balachandran Nayar, M.S. and Callery, P.S.  
GC analysis of metabolites of the cell differentiating agent hexamethylene bisacetamide 415(1987)148
- Ehrsson, H., see Hassan, M. 434(1988)
- Eibel, G., see Scharpf, F. 413(1987)91
- Eichelbaum, M., see Meese, C.O. 423(1987)344
- Eichhorn, K.W., see Schwenk, S. 448(1988)424
- Eichinger, A., see Nitsche, V. 420(1987)207
- Eichinger, K., see Kienzl, E. 430(1988)263
- Eisen, O., see Rang, S. 406(1987)75
- Eisenberg, B.A., see Kling, G.J. 407(1987)377
- Eisenhofer, G., Kirk, K.L., Kopin, I.J. and Goldstein, D.S.  
Simultaneous determination of endogenous catechols and exogenous 2- and 6-fluorinated catechols in tissue and plasma using LC with electrochemical detection 431(1988)156
- Ekberg, B., see Sellergren, B. 450(1988)277
- Ekundayo, O., Laakso, I., Hiltunen, R. and Seppänen, T.  
Rapid analyses of chromenes (benzopyrans) by GC-MS 403(1987)358
- Elazhary, Y., see Mohanty, J.G. 435(1988)149
- Elchisak, M.A., see Scott, M.C. 413(1987)17
- El Dareer, S.M., see Rose, L.M. 425(1988)419
- Elenbring, K., see Hjertén, S. 403(1987)47
- Elguero, J., see Gago, F. 449(1988)95
- Elhilali, M., see Bouthillier, F. 403(1987)171
- Elias-Baker, B.A., see Jacob, III, P. 417(1987)277
- Elion, J., see Ducrocq, R. 419(1987)75
- Elisss, L.O., see Björkman, S. 420(1987)341
- El Kholi, A.M., see Brooks, J.B. 430(1988)209
- Ellard, G.A., see Jenner, P.J. 415(1987)188
- Eller, T.D., see Vrbanac, J.J. 425(1988)1

- Ellinwood, Jr., E.H., see Gupta, S.K.  
445(1988)310
- Ellis, P.P., see Riegel, M.A. 424(1988)177
- Ellow, L.-E., see Bergenhem, N. 403(1987)383
- Elseviers, M., see De Bondt, M. 442(1988)165
- Elsharaky, A.S., see Moneam, N.M.A.  
437(1988)316
- El Sharaky, A.S., see Moneam, N.M.A.  
438(1988)438
- Elson, C.M., see Sim, P.G. 445(1988)239
- El Tayar, N., Tsantili-Kakoulidou, A.,  
Roethlisberger, T., Testa, B. and Gal, J.  
Different partitioning behaviour of  
sulphonyl-containing compounds in  
reversed-phase HPLC and octanol-water  
systems 439(1988)237
- Elton, T.S., see Reeves, R. 418(1987)73
- Ema, M., see Itami, T. 417(1987)168
- Embil, K., see Townsend, R.W. 450(1988)414
- Emery, P.W., see Woodward, C.J.H.  
419(1987)280
- Emilsson, H.  
High-performance liquid chromatographic  
determination of glipizide in human plasma  
and urine 421(1987)319
- Emilsson, H.  
HPLC determination of cadralazine in human  
plasma and urine 424(1988)119
- Emm, T., Lesko, L.J., Leslie, J. and Perkal, M.B.  
Determination of albuterol in human serum  
by reversed-phase HPLC with electrochemical  
detection 427(1988)188
- Emm, T., Lesko, L.J. and Perkal, M.B.  
Simultaneous determination of doxepin and  
nordoxepin in serum using HPLC  
419(1987)445
- Emm, T.A., Leslie, J., Chai, M., Lesko, L.J. and  
Perkal, M.B.  
HPLC assay of cephalixin in serum and urine  
427(1988)162
- Emodi, S., see Bhargava, V.O. 426(1988)212
- Endert, E., see Van Ingen, H.E. 430(1988)233
- Endoh, Y.S., Yoshimura, H. and Sasaki, N.  
HPLC determination of bromofenofos and  
its metabolite in rat plasma 426(1988)202
- Eng, G., Johannesen, R.B., Tierney, E.J., Bellama,  
J.M. and Brinckman, F.E.  
Holistic molecular conformation and total  
surface area calculations as predictors for  
solution properties and chromatographic  
parameters 403(1987)1
- Eng, G., see Tierney, E.J. 441(1988)229
- Engelen, D.L. Van, Thomas, L.C. and Piepmeier,  
E.H.  
UV spectrometric detection for GC of  
polynuclear aromatic compounds with  
repetitive spectral scans using absorbance and  
concurrent fluorescence measurements  
405(1987)191
- Engelmann, G.J., Esmans, E.L., Alderweireldt,  
F.C. and Rillaerts, E.  
Rapid method for the analysis of red blood  
cell fatty acids by reversed-phase HPLC  
432(1988)29
- Engels, W., Kamps, M.A.F., Lemmens, P.J.M.R.,  
Van der Vusse, G.J. and Reneman, R.S.  
Determination of prostaglandins and  
thromboxane in whole blood by HPLC with  
fluorimetric detection 427(1988)209
- Engelsma, J.W., see Haagsma, N. 436(1988)73
- Engewald, W., Billing, U., Topalova, I. and  
Petsev, N.  
Structure-retention correlations of  
alkylphenols in GLC and GSC 446(1988)71
- Enkelaar-Willemsen, C., see Jansen, E.H.J.M.  
437(1988)268
- Ennachachibi, A., Nicolas, P., Fauvelle, F.,  
Perret, G. and Petitjean, O.  
Effective HPLC determination of glafenine  
in plasma: pharmacokinetic application  
427(1988)307
- Ensley, R.J., see Brown, L.L. 426(1988)370
- Entwistle, L.A., see Stubbs, R.J. 413(1987)171
- Epstein, A.L., see Chen, F.-M. 444(1988)153
- Er, S.S., see McGarrity, M.J. 413(1987)53
- Er, S.S., see McGarrity, M.J. 415(1987)136
- Er, S.S., see McGarrity, M.J. 419(1987)37
- Erdélyi-Tóth, V., see Vendrig, D.E.M.M.  
414(1987)91
- Erdmann, G.R., Canafax, D.M. and Giebink,  
G.S.  
HPLC analysis of trimethoprim and  
sulfamethoxazole in microliter volumes of  
chinchilla middle ear effusion and serum  
433(1988)187
- Erdmann, G.R., see Nygard, G.A. 415(1987)438
- Eriksson, A., see Qureshi, G.A. 441(1988)19723
- Eriksson, J.E., see Meriluoto, J.A.O. 438(1988)93
- Erkelens, C., Billiet, H.A.H., De Galan, L. and  
De Leer, E.W.B.  
Origin of system peaks in single-column ion  
chromatography of inorganic anions using  
high pH borate-gluconate buffers and  
conductivity detection 404(1987)67
- Erni, F., see Steuer, W. 447(1988)287
- Ertel, K.D. and Carstensen, J.T.  
Quantitative determination of octanoic acid  
by HPLC following derivatization with 4-

- bromomethyl-7-methoxycoumarin  
411(1987)297
- Escribano, J., Matas, R. and Méndez, E.  
HPLC and photodiode-array detection of the  
human protein HC (human complex-forming  
glycoprotein heterogeneous in charge), a  
chromophore-associated protein  
444(1988)165
- Eslami, M., Stuart, J.D. and Cohen, K.A.  
Improvement in the resolution of *o*-  
phthalaldehyde derivatized amino acids by  
applying gradient steepness optimization to  
five reversed-phase columns of different  
lengths and particle sizes 411(1987)121
- Esmans, E.L., see Engelmann, G.J. 432(1988)29
- Euerby, M.R., Partridge, L.Z. and Gibbons, W.A.  
HPLC determination of lombricine and N-  
phosphoryl lombricine in the earthworm by  
pre-column fluorescence derivatisation with  
*o*-phthalaldehyde-ethanethiol 445(1988)433
- Euerby, M.R., Partridge, L.Z. and Rajani, P.  
Resolution of lombricine enantiomers by  
HPLC utilising pre-column derivatisation  
with *o*-phthalaldehyde-chiral thiols  
447(1988)392
- Evans, M.B. and Haken, J.K.  
Classification of GC stationary phases by a  
new variant of Kováts retention index  
406(1987)105
- Evans, M.J., see Steffenrud, S. 416(1987)219
- Evans, M.J., see Steffenrud, S. 423(1987)1
- Evelo, C.T.A., Kon, T.C., Fijneman, P.H.S. and  
Henderson, P.T.  
Separation of human haemoglobin alkylated  
at  $\beta$ 93 cysteine from its native form by fast  
protein LC 427(1988)335
- Evelo, C.T.A., Niessen, H.J.J.M., Roelofs, H.M.J.  
and Henderson, P.T.  
Cysteine adducts of human haemoglobin  
measured by isoelectric focusing in  
polyacrylamide gels with a non-linear pH  
gradient 420(1987)35
- Everaerts, F.M., see Gladdines, M.M.  
431(1988)317
- Everaerts, F.M., see Lemmens, A.A.G.  
439(1988)423
- Evershed, R.P., Spooner, N., Prescott, M.C. and  
Goad, L.J.  
Isolation and characterisation of intact steryl  
ferulates from seeds 440(1988)23
- Ewing, A.G., see Wallingford, R.A. 441(1988)299
- Excoffier, J.L., see Schlabach, T.D. 439(1988)173
- Ezra, I., see Massil, S.E. 439(1988)476
- F**
- Faber, J.P.M., see Ugalde, T.D. 449(1988)207
- Fábián-Vonsik, V., see Molnár-Perl, I.  
446(1988)231
- Fábián-Vonsik, V., see Molnár-Perl, I.  
446(1988)237
- Fadeev, A.Y. and Staroverov, S.M.  
Geometric structural properties of bonded  
layers of chemically modified silicas  
447(1988)103
- Fadiran, E.O. and Davidson, A.G.  
Determination of low levels of phenothiazine  
sulphoxides in phenothiazine drug substances  
and formulations by TLC-second-derivative  
spectrofluorimetry 442(1988)363
- Fainaru, M., see Knobler, H. 421(1987)136
- Fairhead, A.P., see Forrest, G. 428(1988)123
- Falagiani, P., see Bassoli, A. 444(1988)209
- Falagiani, P., see Bassoli, A. 446(1988)179
- Falardeau, P., see Martineau, A. 417(1987)1
- Falk, J.L., see Lau, C.E. 423(1987)251
- Falkowski, A.J., Look, Z.M., Noguchi, H. and  
Silber, B.M.  
Determination of cefixime in biological  
samples by reversed-phase HPLC  
422(1987)145
- Familletti, P.C., see Weber, D.V. 431(1988)55
- Fanali, S., Cristalli, M. and Catellani, P.  
Determination of minoxidil in pharmaceutical  
forms by capillary isotachopheresis  
405(1987)385
- Fanali, S., Lederer, M., Masia, P. and Ossicini,  
L.  
Paper electrophoretic study of ion-pair  
formation. XVII. Influence of the  
enantiomeric purity of the sodium tartrate  
used as electrolyte 440(1988)361
- Fanali, S., Masia, P. and Ossicini, L.  
Paper electrophoretic study of ion-pair  
formation. XVI. Resolution of optical isomers  
of some cobalt(III) complexes with amino  
acid ligands 403(1987)388
- Fanali, S., Sinibaldi, M. and Quaglia, M.G.  
Isotachopheretic analysis of amoxicillin and  
carboxymethylcysteine in pharmaceuticals  
408(1987)441
- Fanali, S. and Sinibaldi, M.  
Host-guest complexation in capillary  
isotachopheresis. I.  $\alpha$ -,  $\beta$ - and  $\gamma$ -cyclodextrins  
as complexing agents for the resolution of  
substituted benzoic acid isomers 442(1988)371
- Fantuzzi, G., see Aggazzotti, G. 416(1987)125

- Fanucchi, M.P., see Stevens, Y.-W. 416(1987)203
- Fanucchi, M.P., see Walsh, T.D. 414(1987)510
- Farin, D. and Avnir, D.  
Accessibility of irregular (fractal) surfaces of porous silicas to silylation reactions 406(1987)317
- Farinotti, R., see Bossant, M.J. 423(1987)23
- Farinotti, R., see Le Coguic, A. 434(1988)320
- Farjam, A., see Nielen, M.W.F. 417(1987)159
- Farkas, M., see Papp, O. 403(1987)19
- Farmen, R.H., see Sciacca, M.A. 428(1988)265
- Farmer, P.B., see Bailey, E. 416(1987)81
- Farrant, M., Zia-Gharib, F. and Webster, R.A.  
Automated pre-column derivatization with *o*-phthalaldehyde for the determination of neurotransmitter amino acids using reversed-phase LC 417(1987)385
- Farré, R., Font, G. and Fuster, A.  
Determination of 5-nitrofurylacrylic acid in wines by HPLC 445(1988)264
- Farthing, D., see Karnes, H.T. 422(1987)165
- Farthing, D., see Karnes, H.T. 424(1988)398
- Fasolato, C. and Galzigna, L.  
Assay of  $\gamma$ -L-glutamylcyclotransferase activity in rat brain synaptosomes by HPLC 426(1988)381
- Fathi, M., see Padmalatha Devi, K. 426(1988)229
- Fattal, E., see Laham, A. 440(1988)455
- Faulkner, III, H.C., see Sciacca, M.A. 428(1988)265
- Faull, K.F., see Bosin, T.R. 428(1988)229
- Fausnaugh, J., see Tous, G. 444(1988)67
- Fausnaugh-Pollitt, J., Thevenon, G., Janis, L. and Regnier, F.E.  
Chromatographic resolution of lysozyme variants 443(1988)221
- Fauvelle, F., see Ennachachibi, A. 427(1988)307
- Favretto, L., see Cotroneo, A. 449(1988)183
- Favretto, L.G., see Cotroneo, A. 449(1988)183
- Favretto, L.G., see Stancher, B. 440(1988)37
- Fayolle, C. and Fillion, G.  
Reversed-phase LC determination of cyclic adenosine 3',5'-monophosphate in rat brain cortex 426(1988)177
- Fazio, M., see Righetti, P.G. 440(1988)367
- Fedeli, F., see Musu, C. 449(1988)432
- Fedeli, G., Moltrasio, D., Aleotti, M. and Gazzani, G.  
HPLC determination of sulphur and captan in a mixture 447(1988)263
- Fedi, M., see Agostini, O. 433(1988)288
- Feely, J., see O'Reilly, T. 417(1987)190
- Fehérvári, A., see Papp, E. 447(1988)315
- Feibush, B.  
Chiral separation through a single substituent association 436(1988)517
- Feibush, B., see Gisch, D.J. 433(1988)264
- Feibush, B., see Shieh, C.H. 406(1987)343
- Feige, M., see Terlinden, R. 430(1988)438
- Feit, M.J., see Pesselman, R.L. 439(1988)448
- Felber, E., see Winkler, E. 436(1988)447
- Feldberg, R.S. and Reppucci, L.M.  
Rapid separation of anomeric purine nucleosides by TLC on a chiral stationary phase 410(1987)226
- Felder, D.L., see Bannon, C.D. 404(1987)340
- Felder, D.L., see Bannon, C.D. 407(1987)231
- Felice, L.J., see Igwe, O.J. 432(1988)113
- Fell, A.F., Noctor, T.A.G., Mama, J.E. and Clark, B.J.  
Computer-aided optimisation of drug enantiomer separation in chiral HPLC 434(1988)377
- Fellows, L.E., see Cole, M.D. 445(1988)295
- Fels, J.-P., Guyonnet, J., Berger, Y. and Cautreels, W.  
Determination of (4-chlorophenyl)thiomethylene biphosphonic acid, a new bisphosphonate, in biological fluids by HPLC 430(1988)73
- Femia, R.A. and Weinberger, R.  
Determination of reducing and non-reducing carbohydrates in food products by LC with post-column catalytic hydrolysis and derivatization. Comparison with refractive index detection 402(1987)127
- Feniou, C., see Duchene, P. 424(1988)205
- Fennessey, P.V., see Scriba, G.K.E. 433(1988)31
- Fenton, M. and Aherne, F.X.  
Determination of glycerol in pig plasma by capillary GC 410(1987)480
- Fenyvesi, É., see Cserhádi, T. 439(1988)393
- Fermaint, A., see Zaromb, S. 439(1988)283
- Fermo, I., Rubino, F.M., Bolzacchini, E., Arcelloni, C., Paroni, R. and Bonini, P.A.  
Pre-column derivatization of amino acids with N,N-diethyl-2,4-dinitro-5-fluoroaniline and reversed-phase LC separation 433(1988)53
- Fernández-Sánchez, E., Fernández-Torres, A., García-Domínguez, J.A., García-Muñoz, J., Menéndez, V., Molera, M.J., Santiuste, J.M. and Pertierra-Rimada, E.  
Mixed stationary phases in GLC. Partition coefficients and retention indices in OV-101-OV-25, OV-101-carbowax 20M and OV-225-SP-2340 mixtures 410(1987)13

- Fernández-Sánchez, E., García-Domínguez, J.A., Menéndez, V., Santiuste, J.M. and Pertierra-Rimada, E.  
Reduction of the number of characterizing probes in GC by factor analysis 402(1987)318
- Fernández-Torres, A., see Fernández-Sánchez, E. 410(1987)13
- Ferrante, A., Bates, E.J. and Nandoskar, M.  
Neutrophil stimulating activity released by *Staphylococcus*-stimulated mononuclear leukocyte conditioned medium. Further characterization and partial purification 440(1988)105
- Ferrara, F.J., see Puglisi, C.V. 417(1987)347
- Ferrari, M. and Verotta, L.  
Centrifugal TLC of alkaloids from vegetable sources using an aluminium oxide layer 437(1988)328
- Ferrari, P., see Beretta, E. 416(1987)144
- Ferrario, P., see Zecca, L. 420(1987)417
- Ferrer, E., Martin, M., Alonso, R. and Luque, J.  
Separation by counter-current distribution of young chick erythrocyte populations and numerical resolution of distribution curves 411(1987)167
- Ferrer, E., Martin, M., Alonso, R. and Luque, J.  
Partitioning behaviour in aqueous two-phase systems and fractionation by counter-current distribution of chick embryo erythrocytes with numerical resolution of distribution curves 445(1988)327
- Ferri, G., see Iadarola, P. 443(1988)317
- Ferry, J.J., Horvath, A.M., Easton-Taylor, M., Toothaker, R.D. and Colburn, W.A.  
Determination of quinapril and its active metabolite in human plasma and urine by GC with electron-capture detection 421(1987)187
- Ferry, N., see Riva, E. 421(1987)99
- Fetizon, M., see Van, T.D. 440(1988)71
- Fetzer, J.C., see Jinno, K. 402(1987)173
- Fetzer, J.C., see Jinno, K. 436(1988)1
- Fiecchi, A., see Anastasia, M. 410(1987)504
- Fields, S.M., Markides, K.E. and Lee, M.L.  
Retention in capillary supercritical fluid chromatography with carbon dioxide mobile phases containing polar modifiers 406(1987)223
- Fife, P.C., see Su, Y. 405(1987)77
- Fijneman, P.H.S., see Evelo, C.T.A. 427(1988)335
- Filipek, M., Pączkowski, D. and Podleśny, J.  
HPLC method for the determination of *p*-hydroxylated and hydroxymethylated metabolites of mexiletine in human serum 430(1988)406
- Fillion, G., see Fayolle, C. 426(1988)177
- Findenegg, G.H., see Koch, C.S. 406(1987)257
- Fini, C., see Palmerini, C.A. 417(1987)378
- Fini, C., see Palmerini, C.A. 423(1987)292
- Fiore, M., see Polcaro, C. 448(1988)127
- Fiorentino, A.M., see Del Principe, D. 419(1987)329
- Fioretti, A., see Della Ragione, F. 440(1988)141
- Fioretti, M., see Della Ragione, F. 440(1988)141
- Firestone, M.A., Michaud, J.-P., Carter, R.H. and Thormann, W.  
Capillary zone electrophoresis and isotachopheresis as alternatives to chromatographic methods for purity control of synthetic peptides 407(1987)363
- Firestone, M.A. and Thormann, W.  
Capillary isoelectric focusing using an LKB 2127 Tachophor isotachopheretic analyzer 436(1988)309
- Fischer, G., see Egger, H.-J. 420(1987)357
- Fischer, M.J.E., see Bos, O.J.M. 424(1988)13
- Fischer Ross, L.  
GC technique to simultaneously quantitate the gases produced by intestinal microorganisms from fermentation mixtures 414(1987)405
- Fisher, E., see Kruck, T.P.A. 433(1988)207
- Fisher, H., see Kasziba, E. 432(1988)315
- Fishman, L.M., see Brown, J.W. 439(1988)441
- Fishman, R.M., see Moore, G.L. 419(1987)95
- Fitos, I. and Simonyi, M.  
Selective effect of clonazepam and (*S*)-uxepam on the binding of warfarin enantiomers to human serum albumin 450(1988)217
- Fitzpatrick, J.C., see Kasziba, E. 432(1988)315
- Flaherty, M.D., see Henner, W.D. 416(1987)426
- Flamini, G., Ria, F. and Scuderi, F.  
Size-exclusion HPLC of immune complexes isolated from patients with acute leukaemia. Preliminary observations 421(1987)434
- Flamini, G., Ria, F., Pizzigallo, E. and Gignina, S.  
Chromatographic and electrophoretic studies of immune complexes in non-A, non-B hepatitis 426(1988)55
- Gherezghiher, T., Koss, M.C., Nordquist, R.E. and Wilkinson, C.P.  
Rapid and sensitive method for measurement of hyaluronic acid and isomeric chondroitin sulfates using HPLC 413(1987)9
- Flanbaum, L., see Kasziba, E. 432(1988)315
- Flanigan, E., see Heinitz, M.L. 443(1988)229
- Flann, B. and Lodge, B.  
Analysis of estrogen sulphate mixtures in

- pharmaceutical formulations by reversed-phase chromatography 402(1987)273
- Flatmark, T., see Kvannes, J. 419(1987)291
- Fleckenstein, L., see Gawienowski, M. 430(1988)412
- Fleischhacker, W., see Haring, C. 428(1988)160
- Fleming, L.H. and Reynolds, Jr., N.C.  
Simultaneous analysis of methionine- and leucine-enkephalin from rat brain: quantification by LC-electrochemistry 431(1988)65
- Flentge, F., see Damsma, G. 428(1988)1
- Flinois, J.-P., see Beaune, P.H. 426(1988)169
- Floberg, S., see Lindström, B. 428(1988)156
- Florance, J., Galdes, A., Konteatis, Z., Kosarych, Z., Langer, K. and Martucci, C.  
HPLC separation of peptide and amino acid stereoisomers 414(1987)313
- Flores-Murrieta, F.J., Hong, E. and Castañeda-Hernández, G.  
Sensitive HPLC assay of pelanserine, a novel antihypertensive agent, in plasma samples 428(1988)167
- Florida, A., see Palmerini, C.A. 417(1987)378
- Florida, A., see Palmerini, C.A. 423(1987)292
- Flouvat, B., see Naline, E. 419(1987)177
- Floyd, T.R., see Sagliano, Jr., N. 443(1988)155
- Flückiger, R. and Mortensen, H.B.  
Glycated haemoglobins (Review) 429(1988)279
- Flurer, C.L., Borra, C., Andreolini, F. and Novotny, M.  
Separation of proteins by microcolumn LC based on the reversed-phase and size-exclusion principles 448(1988)73
- Foglietti, M.J., see Ricard, B. 428(1988)203
- Folena-Wasserman, G., Inacker, R. and Rosenbloom, J.  
Assay, purification and characterization of a recombinant malaria circumsporozoite fusion protein by HPLC 411(1987)345
- Foley, J.P.  
Separation of *m*- and *p*-cresol and other methylated phenols in shale oil by reversed-phase LC with electrochemical detection 441(1988)347
- Foley, R.C.L., see Haddad, P.R. 407(1987)133
- Folk, J.E., see Beninati, S. 443(1988)329
- Follette, G. La, Gambertoglio, J., White, J.A., Knuth, D.W. and Lin, E.T.  
Determination of clindamycin in plasma or serum by HPLC with UV detection 431(1988)379
- Fölsch, U.R., see Löser, C. 430(1988)249
- Font, G., see Farré, R. 445(1988)264
- Ford, C.W. and Hartley, R.D.  
Identification of phenols, phenolic acid dimers, and monosaccharides by GLC on a capillary column 436(1988)484
- Ford, J.M., see Forsyth, D.R. 433(1988)352
- Forgione, A., see Pedrazzini, S. 413(1987)338
- Forgione, A., see Pedrazzini, S. 415(1987)214
- Forgue, S.T., Pittman, K.A. and Barbhuiya, R.H.  
HPLC analysis of a novel carbapenem antibiotic in human plasma and urine 414(1987)343
- Formstecher, P., see Maes, P. 445(1988)409
- Formstecher, P., see Sablonniere, B. 403(1987)183
- Forrest, G., McInnes, G.T., Fairhead, A.P., Thompson, G.G. and Brodie, M.J.  
Simple HPLC method for the measurement of amiloride in body fluids 428(1988)123
- Forslind, K., see Åkesson, B. 413(1987)233
- Forsmo-Bruce, H., see Johansson, M. 432(1988)265
- Förster, H.-G., Siems, W., Rathmann, J. and Müller, M.  
New method of purification of perchloric acid extracts for quantitative GC of some Krebs cycle acids 424(1988)357
- Forsyth, D.R., Ford, J.M., Truman, C.A., Roberts, C.J.C. and Wilcock, G.K.  
Determination of tacrine hydrochloride in human serum by chloroform extraction, reversed-phase HPLC and fluorimetric detection 433(1988)352
- Foster, N., see Duff, G.A. 416(1987)71
- Foster, R.T. and Jamali, F.  
HPLC assay of ketoprofen enantiomers in human plasma and urine 416(1987)388
- Foster, W.S. and Jarrett, H.W.  
Melittin-silica, a high-pressure affinity chromatography resin for calmodulin 403(1987)99
- Foth, H., see Michaelis, H.C. 416(1987)176
- Fóti, G., Belvito, M.L. and Kováts, E.  
Packing density of organyldimethylsiloxy-covered silica gel and hydrolytic stability of the coating 440(1988)315
- Fóti, G., see Nagy, L.G. 406(1987)311
- Fouda, H.G., Ronfeld, R.A. and Weidler, D.J.  
GC-MS analysis and preliminary human pharmacokinetics of sertraline, a new antidepressant drug 417(1987)197
- Foulger, B., see Begley, P. 445(1988)119
- Foulger, B.E., see Begley, P. 438(1988)45
- Fournet, B., see Capon, C. 425(1988)35
- Fournié, J.-J., see Rivière, M. 445(1988)87
- Fournier, P.E., see Girre, C. 419(1987)363
- Fourtillan, J.B., see Mignot, A. 430(1988)192



- Fox, A., see Gilbart, J. 441(1988)323
- Fox, A., see Sonesson, A. 431(1988)1
- Fox, K.E., see Sullivan, W.R. 425(1988)396
- Fracassi, F., see Moretti, G. 442(1988)459
- Fracasso, C., see Guiso, G. 413(1987)363
- Fraillon, P., see Mura, P. 416(1987)303
- France, N.C., Holland, P.T., McGhie, T.K. and Wallace, M.R.  
Measurement of plasma oxalate by capillary GC and its validation by isotope dilution MS 433(1988)1
- Franceschini, N., see Amicosante, G. 403(1987)366
- Frank, H., see Demange, P. 438(1988)291
- Frank, S.K., Mathiesen, D.A., Whitfield, L.R. and Ames, M.M.  
HPLC assay for the experimental anticancer agent oxantrazole 419(1987)225
- Frank, G. and Verillon, F.  
Economic laboratory practice in preparative column HPLC 450(1988)81
- Franklin, R.B., see Miller, B.J. 426(1988)438
- Frantz, S.C., Henriques, H.F. and Phillips, T.M.  
Immunoaffinity isolation of the Lewis blood group antigens 444(1988)41
- Frantz, S.C., see Phillips, T.M. 444(1988)13
- Frascio, F., see Castello, G. 416(1987)119
- Fraser, A.J. and Taggart, C.T.  
TLC-flame ionization detection calibration with natural and synthetic standards 439(1988)404
- Frattini, C., see Bicchi, C. 411(1987)237
- Frattini, P., see Schinelli, S. 431(1988)150
- Frei, R.W., see Baumann, R.A. 439(1988)165
- Frei, R.W., see Broersen, J.J. 436(1988)39
- Frei, R.W., see De Ruiter, C. 436(1988)429
- Frei, R.W., see Irth, H. 424(1988)95
- Frei, R.W., see Irth, H. 439(1988)63
- Frei, R.W., see Jansen, H. 440(1988)217
- Frei, R.W., see Kok, W.T. 436(1988)421
- Frei, R.W., see Maris, F.A. 435(1988)297
- Frei, R.W., see Nielen, M.W.F. 417(1987)159
- Freitag, W.  
Determination of the polymeric light stabilizer Chimassorb 944 in polyolefins by reversed-phase HPLC 450(1988)430
- Freitag, W., Wurster, R. and Mady, N.  
Determination of the polymeric light stabilizer Tinuvin 622 in polyolefins 450(1988)426
- Freudenberg, M.A., see Galanos, C. 440(1988)397
- Frey, D.D.  
Free energy consumption at self-sharpening concentration fronts in isothermal fixed-bed adsorption. Theory for isotherms of arbitrary shape and applications in displacement chromatography 409(1987)1
- Frič, P., see Kocna, P. 434(1988)429
- Fridman, M., see Hearn, M.T.W. 435(1988)271
- Friedel, H.D. and Matusch, R.  
Separation of non-polar sesquiterpene olefins from tolu balsam by HPLC; silver perchlorate impregnation of a prepacked preparative silica gel column 407(1987)343
- Friedman, H., see Burstein, E.S. 423(1987)380
- Friedman, H., see Kennedy, J.S. 423(1987)373
- Friedman, H., see Scavone, J.M. 419(1987)339
- Friesen, W.T., see Owen, S.G. 416(1987)293
- Friis, J.M., see Lakings, D.B. 428(1988)131
- Frijlink, H.W., Visser, J. and Drenth, B.F.H.  
Determination of cyclodextrins in biological fluids by HPLC with negative colorimetric detection using post-column complexation with phenolphthalein 415(1987)325
- Frisvad, J.C. and Thrane, U.  
Standardized HPLC of 182 mycotoxins and other fungal metabolites based on alkylphenone retention indices and UV-VIS spectra (diode array detection) 404(1987)195
- Fritsch, P., see Saint Blanquat, G.D. 415(1987)388
- Fritz, J.S.  
A look at contemporary ion chromatography 439(1988)3
- Fritz, J.S., see Dix, K.D. 408(1987)201
- Fritz, J.S., see Tanaka, K. 409(1987)271
- Frommel, D., see Ogunkolade, B.W. 440(1988)459
- Frommer-Filep, M., see Gál, S. 403(1987)217
- Fruchart, J.-C., see Mezdour, H. 414(1987)35
- Frueh, F., see Graf, P. 425(1988)353
- Frydman, A., see Le Liboux, A. 417(1987)151
- Fuerst, P., see Swobodnik, W. 423(1987)75
- Fujii, J., see Inotsume, N. 428(1988)402
- Fujii, Y., Ikeda, Y. and Yamazaki, M.  
Micro HPLC determination of cardiac glycosides in  $\beta$ -methyl digoxin and digoxin tablets 448(1988)157
- Fujimaki, Y., Hokusui, H. and Takegoshi, T.  
Quantitative analysis of budralazine in human plasma and urine by GC-MS 421(1987)367
- Fujimaki, Y., Sudo, K. and Tachizawa, H.  
High-performance liquid chromatographic determination of a new nootropic, N-(2,6-dimethyl-phenyl)-2-(2-oxo-1-pyrrolidinyl)-acetamide, in human serum and urine 433(1988)235
- Fujimori, K. and Yamamoto, K.  
Determination of acetylcholine and choline in perchlorate extracts of brain tissue using

- LC-electrochemistry with an immobilized-enzyme reactor 414(1987)167
- Fujimoto, C., Morita, T. and Jinno, K.  
Microcolumn HPLC-TLC-Fourier transform IR spectrometry 438(1988)329
- Fujimoto, C., Yoshida, H. and Jinno, K.  
Interfacing of inductively coupled plasma-atomic emission spectrometry to SFC for elemental detection 411(1987)213
- Fujimoto, T., see Tawa, R. 425(1988)143
- Fujimoto, Y., Ishi, K., Nishi, H., Tsumagari, N., Kakimoto, T. and Shimizu, R.  
New derivatization reagents for the resolution of carboxylic acid enantiomers by HPLC 402(1987)344
- Fujita, K., see Imai, Y. 420(1987)404
- Fujiwara, K., see Kobayashi, K. 417(1987)439
- Fujiwara, S., see Honda, S. 404(1987)313
- Fujiwara, T., see Tokuma, Y. 415(1987)156
- Fujiwara, Y., see Ogawa, K. 426(1988)188
- Fukasawa, K., see Harada, M. 424(1988)129
- Fukasawa, K.M., see Harada, M. 424(1988)129
- Fukuda, E.K., see Miwa, B.J. 431(1988)343
- Fukuda, H., see Ohta, M. 402(1987)392
- Fukuda, K., see Yamaguchi, M. 414(1987)275
- Fukuhara, K. and Matsuki, Y.  
Simple method for the determination of pipemic acid in biological fluids by HPLC 416(1987)409
- Fukuhara, T. and Yuasa, S.  
Rapid resolution of 2,4-dinitrophenyl-DL-amino acids on a native cellulose column 411(1987)502
- Fukui, M., see Igarashi, K. 415(1987)407
- Fukui, M., see Kasuya, F. 416(1987)189
- Fukui, Y., see Matsubara, K. 424(1988)49
- Fukunaga, Y., see Okamoto, T. 430(1988)11
- Fukushige, S., see Kato, Y. 447(1988)212
- Fullinaw, R.O., Bury, R.W. and Moulds, R.F.W.  
LC screening of diuretics in urine 415(1987)347
- Fullinaw, R.O., Bury, R.W. and Moulds, R.F.W.  
Screening procedure for stimulant laxatives in urine using HPLC with diode array detection 433(1988)131
- Fulton, S., see Kopaciewicz, W. 409(1987)111
- Fumagalli, L., see Musu, C. 449(1988)432
- Funazo, K., see Tanaka, M. 404(1987)175
- Funazo, K., see Tanaka, M. 438(1988)253
- Furlong, E.A., see Henner, W.D. 416(1987)426
- Furness, J.B., see Murphy, R. 408(1987)388
- Fürst, P., Krüger, C., Meemken, H.-A. and Groebel, W.  
Determination of the polychlorinated biphenyl substitute Ugilec (tetrachlorobenzyltoluenes) in fish 405(1987)311
- Fürst, P., see Stehle, P. 449(1988)299
- Furton, K.G., see Kersten, B.R. 411(1987)43
- Furukawa, M., Yoshitake, N. and Yokoyama, T.  
Identification of polymer glycols in polyurethane elastomers by pyrolysis-GC 435(1988)219
- Furuta, T., Canfell, P.C., Castagnoli, K.P., Sharma, M.L. and Miller, R.D.  
Quantitation of pancuronium, 3-desacetylpancuronium, vecuronium, 3-desacetylvecuronium, pipecuronium and 3-desacetylpipecuronium in biological fluids by capillary GC using nitrogen-sensitive detection 427(1988)41
- Furuta, T., Kasuya, Y., Shibasaki, H. and Baba, S.  
Determination of stable isotopically substituted histidine in human plasma by GC-MS 413(1987)1
- Furuta, T., see Hirota, N. 425(1988)237
- Fusek, M., see Prisyazhnoy, V.S. 424(1988)243
- Fuster, A., see Farré, R. 445(1988)264

## G

- Gachályi, B., see Róna, K. 420(1987)228
- Gadeholt, G., see Christophersen, A.S. 422(1987)117
- Gagliardi, L., Amato, A., Basili, A., Cavazzutti, G. and Tonelli, D.  
Determination of sun-screen agents in cosmetic products by reversed-phase HPLC 408(1987)409
- Gagliardi, L., Amato, A., Cavazzutti, G., Chimenti, F., Bolasco, A. and Tonelli, D.  
Identification and quantification of hydroquinone and some of its ethers in cosmetic products by reversed-phase HPLC 404(1987)267
- Gagliardi, L., Amato, A., Cavazzutti, G., Tonelli, D. and Montanarella, L.  
Identification of xanthene dyes in lipsticks by reversed-phase HPLC 448(1988)296
- Gago, F., Alvarez-Builla, J. and Elguero, J.  
Group contributions to hydrophobicity and elution behaviour of pyridine derivatives in reversed-phase HPLC 449(1988)95

- Gagosian, R.B., see Kawamura, K. 438(1988)299
- Gagosian, R.B., see Kawamura, K. 438(1988)309
- Gaida, A.V., Monastyrskii, V.A., Magerovskii, Y.V., Staroverov, S.M. and Lisichkin, G.V.  
Affinity chromatography of human thrombin on modified silica 424(1988)385
- Gaillard, S., see Yon, J.M. 440(1988)421
- Gaillot, J., see Le Liboux, A. 417(1987)151
- Gal, J., see El Tayar, N. 439(1988)237
- Gál, S., see Kiss, L. 448(1988)109
- Gál, S., Tar, A., Frommer-Filep, M., Toth-Martinez, B.L., Hernádi, F.J. and Kiss, L.  
Use of chromatofocusing for separation of  $\beta$ -lactamases. VIII. Analytical chromatofocusing of chromosomal cephalosporinases from four *Klebsiella* strains 403(1987)217
- Galan, L. De, see Erkelens, C. 404(1987)67
- Galanos, C., Jiao, B., Komuro, T., Freudenberg, M.A. and Lüderitz, O.  
Large-scale fractionation of S-form lipopolysaccharide from *Salmonella abortus equi*. Chemical and serological characterization of the fractions 440(1988)397
- Galanos, C., see Komuro, T. 450(1988)381
- Galdes, A., see Florance, J. 414(1987)313
- Galensa, R., see Heimhuber, B. 439(1988)481
- Galensa, R., see Krause, M. 441(1988)417
- Galensa, R., see Risch, B. 448(1988)291
- Galindo, I.C. and Tarter, J.G.  
Causes for the loss of resolution in an ion chromatography resin 445(1988)211
- Gall, J.-M. Le, see Gallice, P. 422(1987)263
- Galli, B., Gasparrini, F., Misiti, D., Natile, G. and Palmieri, G.  
TLC of metal complexes on glycidoxypropyl functionalized silica plates 409(1987)377
- Galliano, M., see Iadarola, P. 443(1988)317
- Gallice, P., Monti, J.-P., Le Gall, J.-M., Crevat, A. and Murisasco, A.  
Ion-pair HPLC profiling of a uremic toxin fraction 422(1987)263
- Gallo, D.G., see Bartek, M.J. 421(1987)309
- Gallo, D.G., see Kerns, E.H. 416(1987)357
- Gallo, G.G., see Riva, E. 421(1987)99
- Gallo, P., Sidén, Å. and Tavalato, B.  
Anion-exchange chromatography of normal and monoclonal serum immunoglobulins 416(1987)53
- Galons, H., see Coleman, A.W. 450(1988)175
- Galushko, S.V., Shishkina, I.P. and Pilipenko, A.T.  
Reversed-phase HPLC of pyrimidine bases and nucleosides. Application of solvophobic theory 445(1988)59
- Galzigna, L., see Fasolato, C. 426(1988)381
- Gamba, M., see Dionisotti, S. 434(1988)169
- Gambertoglio, J., see La Follette, G. 431(1988)379
- Games, D.E., see Morgan, E.D. 441(1988)165
- Games, D.E., see Parker, C.E. 438(1988)359
- Gan, D.-C. and Tarter, J.G.  
Simultaneous determination of sodium and chloride ions by ion chromatography at the  $\mu\text{g/l}$  level 404(1987)285
- Ganansia, J., Bianchetti, G. and Thénot, J.P.  
Correlation between reversed-phase HPLC and plasma protein binding 421(1987)83
- Gancheva, V.K., see Rachev, R.C. 437(1988)287
- Gandhe, B.R., see Ramachandran, P.K. 426(1988)239
- Gandhi, V., Danhauser, L. and Plunkett, W.  
Separation of 1- $\beta$ -D-arabinofuranosylcytosine 5'-triphosphate and 9- $\beta$ -D-arabinofuranosyl-2-fluoroadenine 5'-triphosphate in human leukemia cells by HPLC 413(1987)293
- Ganes, D.A. and Midha, K.K.  
Sensitive and specific HPLC assay for the quantification of sulforidazine and two diastereomeric sulforidazine-5-sulfoxide metabolites in plasma 423(1987)227
- Ganes, D.A. and Wagner, J.G.  
Improved HPLC assay for the quantification of 5-bromo-2'-deoxyuridine and 5-bromouracil in plasma 432(1988)233
- Ganetsos, G.  
Prediction of the distribution coefficient ( $K_d$ ) variation with operating conditions in chromatographic systems 411(1987)81
- Gangoda, M.E. and Gilpin, R.K.  
Comparative study of alkyl and fluoroalkyl N-(2-hydroxy-ethoxyethyl)-amides as reversed-phase LC modifiers 448(1988)365
- Garaev, E.S., see Zaslavsky, B.Y. 439(1988)267
- Garaj, J., see Brandšteterová, E. 404(1987)359
- Garaj, J., see Buszewski, B. 446(1988)191
- Garaj, J., see Krupčík, J. 406(1987)117
- Garaud, J.J., see Bauchet, J. 414(1987)472
- García-Domínguez, J.A., see Fernández-Sánchez, E. 402(1987)318
- García-Domínguez, J.A., see Fernández-Sánchez, E. 410(1987)13
- García-Muñoz, J., see Fernández-Sánchez, E. 410(1987)13
- García-Pérez, A.I., Recio, M.N., Sancho, P. and Luque, J.  
Partitioning behaviour of rat bone marrow cells in aqueous two-phase systems. Dependence of cell partition on the interfacial

- tension and electrical potential difference between the phases 403(1987)131
- García-Raso, A., Ballester, P., Bergueiro, R., Martínez, I., Sanz, J. and Jimeno, M.L. Estimation of the polarity of stationary phases by proton nuclear magnetic resonance spectroscopy. Application to phenyl and methyl silicones (OV and SE series) 402(1987)323
- Gareil, P. and Rosset, R. Analysis of the information in a preparative chromatogram for further optimization of the operating conditions 450(1988)13
- Garland, I.J., see Bannon, C.D. 407(1987)231
- Garland, W., see Oldfield, N. 422(1987)135
- Garland, W.A., see Miwa, B.J. 431(1988)343
- Garland, W.A., see Rubio, F. 421(1987)281
- Garnier-Moiroux, A., Poirier, J.-M., Cheymol, G. and Jaillon, P. HPLC determination of pentoxifylline and its hydroxy metabolite in human plasma 416(1987)183
- Garraffo, H.M., see Santa-Coloma, T.A. 415(1987)297
- Garrec, L. Le, see Delee, E. 450(1988)183
- Garrec, L. Le, see Delee, E. 450(1988)191
- Garside, D.R., see Smith, R.M. 407(1987)19
- Garthwaite, K., see Japp, M. 439(1988)317
- Garty, M., Steinmetz, Y., Rosenfeld, J.B. and Goldstein, D.S. Species-dependent differences in recovery of 3,4-dihydroxybenzylamine in assays of plasma catecholamines 430(1988)123
- Garvie, C.T., Straub, K.M. and Lynn, R.K. Quantitative LC determination of disulfide-containing peptide analogues of vasopressin with dual Hg/Au electrochemical detection 413(1987)43
- Gasanova, G.Z., see Zaslavsky, B.Y. 403(1987)123
- Gasparič, J., see Davidková, P. 410(1987)53
- Gasparrini, F., see Galli, B. 409(1987)377
- Gassend, R., Duprat, F. and Gau, G. Inductive adsorption as a new approach to amine enantiomer resolution 404(1987)87
- Gassiot, M., see Aceves, M. 436(1988)503
- Gateau-Olesker, A., see Porziemsky, J.-P. 440(1988)183
- Gathercole, P.S. and Thiel, P.G. LC determination of the cyanoginosins, toxins produced by the cyanobacterium *Microcystis aeruginosa* 408(1987)435
- Gattrell, M.A. and Kirk, D.W. Interconversion of hydroquinone and benzoquinone during styrene-divinylbenzene resin-based ion-exclusion chromatography 409(1987)404
- Gau, G., see Gassend, R. 404(1987)87
- Gauron, S., see Godbillon, J. 427(1988)269
- Gautam, V.K., see Dhillon, R.S. 435(1988)256
- Gavinelli, M., see Latini, R. 424(1988)211
- Gavrilova, T.B., Parshina, I.V., Dimitrov, C., Ivanov, S., Petsev, N. and Topalova, I. Carbon black adsorbents modified with *n*-hydrocarbons and treated with an high-frequency low-temperature plasma 402(1987)87
- Gavrilova, T.B., Roshchina, T.M. and Vasil'eva, E.V. Use of modified tungsten selenide as an adsorbent for GC 446(1988)79
- Gavryushkin, A.V., see Shukun, S.A. 432(1988)137
- Gawdzik, B. and Patrykiewicz, A. GC on porous polymers. IV. Influence of the geometric structure of porous copolymers of 1,4-di(methacryloyloxymethyl)naphthalene with 1,4-divinylbenzene on their chromatographic behaviour 448(1988)233
- Gawienowski, M., Benet, L.Z., Fleckenstein, L. and Lin, E.T. Ion-paired LC method for the analysis of blood and plasma for the antimalarial drug halofantrine and its putative mono-debutylated metabolite 430(1988)412
- Gayte-Sorbier, A., see Airaud, C.B. 437(1988)59
- Gazdag, M., Szepesi, G. and Huszár, L.  $\alpha$ -,  $\beta$ - and  $\gamma$ -cyclodextrins as mobile phase additives in the HPLC separation of enantiomeric compounds. II. Optimization of the separation method by using  $\alpha$ -,  $\beta$ - and  $\gamma$ -cyclodextrins in mixture 436(1988)31
- Gazdag, M., Szepesi, G. and Mihályfi, K. Some aspects of the selection of HPLC methods for the separation of chiral compounds in pharmaceutical analysis 450(1988)145
- Gazzani, G., see Fedeli, G. 447(1988)263
- Gazzei, G., see Benanchi, P.L. 450(1988)133
- Geerdink, R.B. Determination of aniline derivatives by HPLC with fluorescence detection 445(1988)273
- Geeson, A.V., see Japp, M. 439(1988)317
- Geeson, A.V., see Parker, C.E. 438(1988)359
- Gehrke, C.W., see Souther, M.E. 417(1987)27
- Gelber, L.R., see Shieh, C.H. 406(1987)343
- Gelder, W.M.J. Van, Jonker, H.H., Huizing, H.J. and Scheffer, J.J.C. Capillary GC of steroidal alkaloids from Solanaceae. Retention indices and

- simultaneous flame ionization/nitrogen-specific detection 442(1988)133
- Gelfi, C., see Rabilloud, T. 402(1987)105
- Gelpi, E., see Cárdenas, A. 426(1988)83
- Gelsema, W.J., see Brandts, P.M. 437(1988)337
- Gelsema, W.J., see Brandts, P.M. 438(1988)181
- Gengo, F.M., see Schuster, D. 433(1988)318
- Gennaro, M.C.  
Role of the alkyl chain length on the ion interaction reagent, flow-rate, column packing and detection in ion interaction reversed-phase HPLC in separations of anions using amine salicylates 449(1988)103
- Gennip, A.H. Van, see De Korte, D. 415(1987)383
- Georgakopoulos-Gregoriades, E., see Satsmadjis, J. 437(1988)254
- Georgopoulos, A., see Czejka, M.J. 424(1988)182
- Georgopoulos, A., see Czejka, M.J. 425(1988)429
- Gerber, G., see Toguzov, R.T. 434(1988)447
- Gerber, G., see Werner, A. 421(1987)257
- Gerbino, T.C., see Castello, G. 416(1987)119
- Gerbino, T.C., see Castello, G. 437(1988)33
- Gercken, G., see Schatowitz, B. 409(1987)43
- Gercken, G., see Schatowitz, B. 425(1988)257
- Gercken, G., see Schmidt, N. 410(1987)458
- Geria, T., Hong, W.-H. and Daly, R.E.  
HPLC assay of imazodan, methylparaben and propylparaben in imazodan injection 450(1988)407
- Gerlo, E., Malfait, R. and Dupont, A.G.  
Identification and quantification of N-methylepinephrine in human urine 414(1987)301
- Gero, S.D., see Porziemsky, J.-P. 440(1988)183
- Geschke, R., Körner, J. and Eggers, H.  
Determination of the new monoamine oxidase inhibitor moclobemide and three of its metabolites in biological fluids by HPLC 420(1987)111
- Gether, U., Vendelbo Nielsen, H. and Schwartz, T.W.  
Tyrosylation and purification of peptides for radioiodination 447(1988)341
- Geyer, R., see Mc Kerrell, P.J. 408(1987)399
- Ghiggeri, G.M., see Candiano, G. 423(1987)319
- Ghodbane, S. and Guiochon, G.  
Influence of the relative concentrations of two components of a binary mixture on their overloaded elution profiles 440(1988)9
- Ghodbane, S. and Guiochon, G.  
Optimization of concentration overload in preparative LC 444(1988)275
- Ghodbane, S. and Guiochon, G.  
Column loading and relative retention in overloaded elution chromatography 450(1988)27
- Gholson, A.R., St. Louis, R. and Hill, H.H.  
Determination of silicon number in silylated organic compounds using the hydrogen atmosphere flame ionization detector 408(1987)329
- Ghosh, P., Sil, P. and Thakur, S.  
Spray reagent for the detection of coumarins and flavonoids on thin-layer plates 403(1987)285
- Giacosa, A., see Castello, G. 416(1987)119
- Giannozzi, A., see Benanchi, P.L. 450(1988)133
- Gianviti, J.M., see Krstulović, A.M. 426(1988)417
- Gibassier, D., see Le Corre, P. 424(1988)424
- Gibassier, D., see Le Corre, P. 450(1988)211
- Gibassier, D., see Sado, P.A. 434(1988)157
- Gibbons, W.A., see Euerby, M.R. 445(1988)433
- Giboudeau, J., see Aussel, C. 423(1987)270
- Gibson, D.M., see Wright, D.S. 417(1987)223
- Giebink, G.S., see Erdmann, G.R. 433(1988)187
- Gielsdorf, W., Arnemann, W., Aichert, G., Hausleiter, H.J. and Li, P.-L.  
Determination of timelotem dihydrogenmaleate and its main metabolite N-desmethyltimelotem in plasma by capillary GC 419(1987)357
- Gielsdorf, W., Hausleiter, H.J., Arnemann, W., Lichtenberger, A. and Aichert, G.  
Determination of revenast hydrochloride in human plasma by GC with nitrogen-selective detection 419(1987)351
- Gielsdorf, W., see Dahmen, W. 427(1988)157
- Gierlich, H.U., see Anspach, B. 443(1988)45
- Giese, R.W., see Cecchini, D.J. 444(1988)97
- Giese, R.W., see Minnetian, O. 410(1987)453
- Gietl, Y., Spahn, H. and Mutschler, E.  
Simultaneous determination of R- and S-prenylamine in plasma and urine by reversed-phase HPLC 426(1988)305
- Giger, W., see Czuczwa, J. 403(1987)233
- Giger, W., see Marcomini, A. 403(1987)243
- Gignina, S., see Flamini, G., 426(1988)55
- Gilbart, J., Harrison, J., Parks, C. and Fox, A.  
Analysis of the amino acid and sugar composition of streptococcal cell walls by GC-MS 441(1988)323
- Gilbert, B.E., see Smith, R.H.A. 414(1987)202
- Gilbert, J., see Castle, L. 437(1988)247
- Gilbert, J., see Castle, L. 437(1988)281
- Gilbert, J., see Mortimer, D.N. 407(1987)393
- Gilby, E.D., see Betteridge, R.F. 432(1988)352
- Gill, R., see Laing, D.K. 442(1988)187
- Gillberg, P.G., see Tyrefors, N. 423(1987)85
- Gilles, U., see Birsner, U. 402(1987)381

- Gillespie, D.G., see Burrows, R. 404(1987)248
- Gillessen, D., see Hochuli, E. 411(1987)371
- Gilpin, R.K., see Gangoda, M.E. 448(1988)365
- Gilpin, R.K., see Yang, S.S. 408(1987)93
- Gilpin, R.K., see Yang, S.S. 439(1988)414
- Gilpin, R.K., see Yang, S.S. 449(1988)115
- Gimenez-Gallego, G. and Thomas, K.A.  
HPLC of phenylthiocarbonyl-amino acids.  
Application to carboxyl-terminal sequencing  
of proteins 409(1987)299
- Ginkel, L.A. Van, Van Blitterswijk, H., Zootjes,  
P.W., Den Bosch, D.V. and Stephany, R.W.  
Assay for trenbolone and its metabolite 17 $\alpha$ -  
Trenbolone in bovine urine based on  
immunoaffinity chromatographic clean-up  
and off-line HPLC-TLC 445(1988)385
- Gioeli, C., see Pettersson, C. 407(1987)217
- Gioeli, C., see Pettersson, C. 435(1988)225
- Giovanni, N. De and Chiarotti, M.  
Analysis of benzodiazepines. II. HPLC  
fluorescence detection after molecular  
rearrangement to acridanones 428(1988)321
- Giralt, E., see Pedroso, E. 409(1987)281
- Girard, J.E., Rebbani, N., Buell, P.E. and Al-  
Khalidi, A.H.E.  
Carbohydrate-borate eluents for anion  
chromatography 448(1988)355
- Girardot, R., see Rihn, B. 428(1988)408
- Girin, E., see Cann-Moisan, C. 442(1988)441
- Girre, C., Koch, H., Crouzette, J., Palombo, S.  
and Fournier, P.E.  
GC determination of chlorproethazine in  
plasma 419(1987)363
- Gisch, D.J., Hunter, B.T. and Feibush, B.  
Shielded hydrophobic phase; a new concept  
for direct injection analysis of biological fluids  
by HPLC 433(1988)264
- Gisch, D.J. and Pearson, J.D.  
Determination of monosaccharides in  
glycoproteins by reversed-phase HPLC on  
2.1-mm narrowbore columns 443(1988)299
- Gjerde, D.T.  
Eluent selection for the determination of  
cations in ion chromatography 439(1988)49
- Gladdines, M.M., Ackermans, M.T., Everaerts,  
F.M., Der Linden, P.J.Q.V., Vader, H.L. and  
Wiegerinck, M.A.H.M.  
Analysis of the aqueous phase of human  
cervical mucus by reversed-phase HPLC and  
capillary isotachopheresis 431(1988)317
- Glatz, Z., see Škabrahová, Z. 408(1987)372
- Gleizes, E., see Pochard, M.-F. 409(1987)315
- Glöckner, G.  
Normal- and reversed-phase separation of  
copolymers prepared from styrene and ethyl  
methacrylate 403(1987)280
- Glover, M.R., see Tribble, D.L. 414(1987)411
- Glund, K., see Abel, S. 446(1988)187
- Gluth, W.P. and Dann, O.  
Determination of fluorescent diamidines in  
plasma of experimental animals by HPLC  
417(1987)331
- Gmur, D.J., Meier, P. and Yee, G.C.  
HPLC column-switching method for two  
cyclosporine metabolites in blood  
425(1988)343
- Gnauck, R., see Becker, H. 410(1987)267
- Goad, L.J., see Evershed, R.P. 440(1988)23
- Gobert, J., see Baltés, E. 430(1988)149
- Goca, B., see Muniak, W. 436(1988)323
- Goca, B., see Witkiewicz, Z. 402(1987)73
- Godbillon, J., Duval, M., Gauron, S. and Dubois,  
J.P.  
HPLC determination of the penem antibiotic  
(5*R*,6*S*)-2-aminomethyl-6-[(1*R*)-  
hydroxyethyl]-2-penem-3-carboxylic acid in  
human plasma and urine 427(1988)269
- Godeneche, D., see Labarre, P. 419(1987)381
- Godoy, A. Adaya, see Guevara, O.L.  
403(1987)350
- Goewie, C.E. and Hogendoorn, E.A.  
Pre-column clean-up and LC determination  
of residues of N-methylcarbamate pesticides  
in extracts of total diet 404(1987)352
- Goewie, C.E. and Hogendoorn, E.A.  
Analysis of bromacil, diuron and 3,4-  
dichloroaniline in contaminated well water,  
using a HPLC column-switching procedure  
410(1987)211
- Goldberg, A.L., see Kee, Y. 428(1988)345
- Goldstein, D.S., see Eisenhofer, G. 431(1988)156
- Goldstein, D.S., see Garty, M. 430(1988)123
- Goldstein, D.S., see Hovevey-Sion, D.  
426(1988)141
- Goldstein, D.S., see Kågedal, B. 429(1988)177
- Gomaa, K.S., see Marzouki, Z.M.H. 425(1988)11
- Gomez-Sanchez, C., see Holland, O.B.  
426(1988)148
- Goncalves, J.C.S., Sclavons, M., Poupaert, J.H.  
and Dumont, P.  
High-performance liquid chromatography of  
1,2-diacyl-*rac*-glycerols and 1,3-  
diacylglycerols through chromogenic  
derivatization with 3,5-dinitrobenzoyl  
411(1987)472
- Gonnet, C., see Chartier, A. 438(1988)263
- González-Martin, G., see Paulos, C. 433(1988)359
- Gonzalez, M.D., Grant, S.K., Williams, H.J. and  
Scott, A.I.

- Analytical and preparative separation of uroporphyrin I and III by HPLC 437(1988)311
- Good, S.S., Reynolds, D.J. and De Miranda, P. Simultaneous quantification of zidovudine and its glucuronide in serum by HPLC 431(1988)123
- Goodwin, K.R., see Schmerr, M.J.F. 425(1988)379
- Gooijer, C., see Baumann, R.A. 439(1988)165
- Gopal, D.H., see Guenther, R.H. 444(1988)79
- Gopalan, N., see Ramachandran, P.K. 426(1988)239
- Gorbunov, A.A., Solovyova, L.Y. and Pasechnik, V.A. Fundamentals of the theory and practice of polymer GPC as a method of chromatographic porosimetry (Review) 448(1988)307
- Gordon, B.J., see Akbari, A. 426(1988)207
- Gordon, M.J., see Huang, X. 425(1988)385
- Goren, M.B., Dhariwal, K.R. and Jenkins, I.D. Concerning hydrolysis of mycolate esters, of phthiocerol dimycocerosates and of related mycobacterial lipids: an anecdotal account 440(1988)487
- Gorka, J., see Crimmins, D.L. 443(1988)63
- Gorrod, J.W., see Jatoe, S.D. 442(1988)394
- Gorst-Allman, C.P., see De Jesus, A.E. 450(1988)101
- Gosnell, A.B., see Holder, C.L. 419(1987)113
- Gosselin, C., see Lavigne, C. 410(1987)201
- Gotelli, G.R. and Kabra, P.M. Simultaneous determination of urinary vanillylmandelic acid, 5-hydroxyindoleacetic acid and homovanillic acid by LC 428(1988)147
- Goto, J., Chikai, T. and Nambara, T. Studies on steroids. CCXXVII. Separation and determination of bile acid 7- and 12-sulphates in urine by HPLC with fluorescence labelling 415(1987)45
- Goto, J., Sano, Y., Tsuchiya, K. and Nambara, T. Studies on steroids. CCXXXIII. Separation and characterization of bile acid disulphates in human urine by HPLC 425(1988)59
- Goto, J., see Iida, T. 438(1988)423
- Goto, K., see Kawai, S. 442(1988)451
- Goto, M., see Takeuchi, T. 435(1988)482
- Goto, T., Matsui, M. and Kitsuwu, T. Determination of aflatoxins by capillary column GC 447(1988)410
- Gottfries, J., see Wester, P. 415(1987)261
- Gottfries, J., see Wester, P. 415(1987)275
- Gottvaldova, M., Hrabova, H., Sillinger, V. and Kucera, J. Biospecific sorption of cyclodextrin glucosyltransferase on physically modified starch 427(1988)331
- Gouyette, A., see Vassal, G. 428(1988)357
- Gozel, P., see Roach, M.C. 426(1988)129
- Graaf, G.J. De and Spierenburg, T.J. LC determination of cyadox in medicated feeds and in the contents of the porcine gastrointestinal tract with fluorescence detection 447(1988)244
- Graf, P., Frueh, F. and Schmid, K. Determination of the angiotensin converting enzyme inhibitor benazeprilat in plasma and urine by an enzymic method 425(1988)353
- Gragnaniello, V., see Della Ragione, F. 440(1988)141
- Graham, G., see Rauch, K. 430(1988)170
- Gramshaw, J.W., see Castle, L. 437(1988)281
- Grandas, A., see Pedroso, E. 409(1987)281
- Grandgirard, A., see Sébédio, J.L. 405(1987)371
- Granley, K., see Singh, A.K. 404(1987)223
- Granneman, G.R. and Sennello, L.T. HPLC procedures for the determination of difloxacin and its metabolites in biological matrices 413(1987)199
- Grant, S.K., see Gonzalez, M.D. 437(1988)311
- Grasela, D.M. and Rocci, Jr., M.L. HPLC analysis of pentoxifylline and 1-(5'-hydroxyhexyl)-3,7-dimethylxanthine in whole blood 419(1987)368
- Gravellese, D., see Musson, D.G. 414(1987)121
- Gray, M.R., see Tam, Y.K. 423(1987)199
- Grazia, F. De, see Rubio, F. 421(1987)281
- Grech-Belanger, O., Leboeuf, E. and Langlois, S. Assay of diltiazem and deacetyldiltiazem by capillary GC 417(1987)89
- Greef, J. Van der, see Reeuwijk, H.J.E.M. 428(1988)93
- Greef, J. Van der, see Tjaden, U.R. 420(1987)53
- Greeley, Jr., G.H., see Draviam, E.J. 414(1987)180
- Green, M.R., see Haken, J.K. 403(1987)145
- Greenaway, W., Wollenweber, E., Scaysbrook, T. and Whatley, F.R. Novel isoferulate esters identified by GC-MS in bud exudate of *Populus nigra* 448(1988)284
- Greenaway, W. and Whatley, F.R. Formation of N-carboxy compounds during trimethylsilylation of amino acids in the presence of carbon dioxide 409(1987)383
- Greenberg, J., see Phillips, M. 422(1987)235
- Greenberg, R., see Brown, E.M. 443(1988)247
- Greenblatt, D.J., see Burstein, E.S. 423(1987)380

- Greenblatt, D.J., see Kennedy, J.S. 422(1987)274  
 Greenblatt, D.J., see Kennedy, J.S. 423(1987)373  
 Greenblatt, D.J., see Locniskar, A. 424(1988)215  
 Greenblatt, D.J., see Mikati, M.A. 433(1988)305  
 Greenblatt, D.J., see Scavone, J.M. 419(1987)339  
 Greene, S.V., Abdalla, T., Morgan, S.L. and Bryan, C.S.  
 HPLC analysis of vancomycin in plasma, bone, atrial appendage tissue and pericardial fluid 417(1987)121  
 Greene, S.V., see Smith, C.S. 423(1987)207  
 Gref, R.  
 GC analysis of underivatized resin acids 448(1988)428  
 Gregorio, G. Di, see Bassoli, A. 444(1988)209  
 Gregorio, G. Di, see Bassoli, A. 446(1988)179  
 Grellier, P.L., see Abraham, M.H. 409(1987)15  
 Gresti, J., see Narce, M. 448(1988)249  
 Grgurinovich, N.  
 HPLC of sulindac and its sulphone and sulphide metabolites in plasma 414(1987)211  
 Grgurinovich, N. and Matthews, A.  
 Analysis of erythromycin and roxithromycin in plasma or serum by HPLC using electrochemical detection 433(1988)298  
 Griest, W.H., see Row, K.H. 409(1987)193  
 Griffin, M., Leah, J., Mould, N. and Compton, G.  
 Construction of an ion-exchange amino acid analyser kit for use with HPLC apparatus 431(1988)285  
 Griffin, R.L., see Minkler, P.E. 413(1987)33  
 Griffin, R.M.E., Kee, T.G. and Adams, R.W.  
 HPLC system for the separation of basic dyes 445(1988)441  
 Griffith, P.E., see Robinson, J.L. 402(1987)201  
 Grigorova, B., Wright, S.A. and Josephson, M.  
 Separation and determination of stable metallo-cyanide complexes in metallurgical plant solutions and effluents by reversed-phase ion-pair chromatography 410(1987)419  
 Grimalt, J., see Aceves, M. 436(1988)503  
 Grimalt, J., see Alexander, G. 446(1988)87  
 Grimsrud, E.P., see Jones, C.S. 409(1987)139  
 Grinberg, N., see Guttman, A. 448(1988)41  
 Grizzle, W., see Jenkins, R.L. 426(1988)249  
 Grob, K., Biedermann, M. and Li, Z.  
 Checking of capacity of a splitless injector — a simple test 448(1988)387  
 Grob, K., see Schilling, B. 435(1988)204  
 Grob, K. and Müller, E.  
 Sample reconcentration by column-external solvent evaporation or injection of large volumes into GC capillary columns? 404(1987)297  
 Grob, R.L., see Eble, J.E. 405(1987)1  
 Grob, R.L., see Eble, J.E. 405(1987)31  
 Grob, R.L., see Eble, J.E. 405(1987)51  
 Grochow, L.B., see Cheng, T.-L. 425(1988)303  
 Groebel, W., see Fürst, P. 405(1987)311  
 Groenland, W.P.T., see Verhaar, L.A.T. 437(1988)185  
 Grønhøj Larsen, C., see Jakobsen, P. 415(1987)413  
 Grønhøj Larsen, F., see Jakobsen, P. 415(1987)413  
 Grood, P.M.R.M. De, see Vree, T.B. 417(1987)458  
 Gros, E.G., see Santa-Coloma, T.A. 415(1987)297  
 Gross, A.S., Brown, K.F., Baird-Lambert, J.A. and Nation, R.L.  
 Determination of ritodrine in blood and plasma by HPLC with fluorescence detection 416(1987)400  
 Gross, M., see Klemm, D. 438(1988)122  
 Gruber, H.A., see Parris, N. 444(1988)219  
 Gruda, I., see Campeau, D. 405(1987)305  
 Grushka, E., see Massil, S.E. 439(1988)476  
 Grynspan, F., see Young, P.R. 421(1987)130  
 Guan, K.L., see Cecchini, D.J. 444(1988)97  
 Guay, D.R.P., see Awani, W.M. 419(1987)414  
 Gübitz, G. and Juffmann, F.  
 Resolution of the enantiomers of thyroid hormones by high-performance ligand-exchange chromatography using a chemically bonded chiral stationary phase 404(1987)391  
 Guelen, P.J.M., see Botterblom, M.H.A. 421(1987)211  
 Guenther, R.H., Gopal, D.H. and Agris, P.F.  
 Purification of transfer RNA species by single-step ion-exchange HPLC 444(1988)79  
 Guevara, O.L., Adaya Godoy, A. and Nava, C.C.  
 Quantification of halothane, enflurane and nitrous oxide by means of a simple GC method 403(1987)350  
 Guha, O.K. and Roy, J.  
 GC grade silica gel from paddy husk ash 437(1988)244  
 Guichardant, M., Lagarde, M., Lesieur, M. and De Maack, F.  
 Thermospray-MS analysis of underivatized monohydroxy fatty acids: application to stimulated platelets 425(1988)25  
 Guieze, P., see Varotsis, N. 403(1987)159  
 Guillemin, C.L.  
 New concept in chromatographic instrumentation. The probe-process liquid chromatograph 441(1988)1  
 Guimbard, J.P., Person, M. and Vergnaud, J.P.  
 Determination of residual solvents in



- pharmaceutical products by GC coupled to a head-space injection system and using an external standard 403(1987)109
- Guinebault, P., see Ascalone, V. 414(1987)101
- Guiochon, G., see Ghodbane, S. 440(1988)9
- Guiochon, G., see Ghodbane, S. 444(1988)275
- Guiochon, G., see Ghodbane, S. 450(1988)27
- Guiochon, G.A., see Katti, A.M. 449(1988)25
- Guiochon, G. and Gutierrez, J.E.N.  
Theoretical prediction of plate height in serially coupled capillary columns 406(1987)3
- Guiso, G., Fracasso, C., Caccia, S. and Abbiati, A.  
Determination of ranitidine in rat plasma and brain by HPLC 413(1987)363
- Gulaid, A.A., see Mallett, D.N. 428(1988)190
- Gumaste, V.K., see Panse, D.G. 411(1987)456
- Gunawan, S. and Treiman, D.M.  
HPLC determination of nafimidone and its major metabolite nafimidone alcohol in human plasma and urine 414(1987)389
- Gunnarsson, P.O., see Polacek, J. 425(1988)424
- Gunter, E.W., see Sowell, A.L. 431(1988)424
- Günther, K.  
TLC enantiomeric resolution via ligand exchange 448(1988)11
- Gupta, M.B., Hubbard, J.W. and Midha, K.K.  
Separation of enantiomers of derivatised or underderivatised propranolol by means of HPLC 424(1988)189
- Gupta, R.N.  
Determination of catecholamines in body fluids 427(1988)
- Gupta, R.N. and Lew, M.  
Determination of isoniazid in plasma by LC 425(1988)441
- Gupta, S.K. and Ellinwood, Jr., E.H.  
LC determination of halazepam in commercial tablets 445(1988)310
- Gurevich, V.A., see Rodchenkov, G.M. 432(1988)283
- Gusovsky, F., Jacobson, K.A., Kirk, K.L., Marshall, T. and Linnoila, M.  
New HPLC procedure for the detection and quantification of  $\beta$ -phenylethylamine 415(1987)124
- Guthrie, R., see Ho, J. 417(1987)269
- Gutierrez, J.E.N., see Guiochon, G. 406(1987)3
- Gutmann, M., see Zobel, H. 446(1988)305
- Guttendorf, R.J., see Huang Chandler, M.H. 419(1987)426
- Guttman, A., Paulus, A., Cohen, A.S., Grinberg, N. and Karger, B.L.  
Use of complexing agents for selective separation in high-performance capillary electrophoresis. Chiral resolution via cyclodextrins incorporated within polyacrylamide gel columns 448(1988)41
- Guyon, F., see Ni, P. 424(1988)255
- Guyonnaud, C., see Lacroix, C. 422(1987)217
- Guyonnet, J., see Fels, J.-P. 430(1988)73
- Guyot, J.L., see Jaussaud, P. 423(1987)123
- Guyt-Scholten, J.W., see Van Harten, J. 423(1987)327
- Gyenge, R., see Patthy, M. 449(1988)191
- Gyllenhaal, O.  
Derivatization of 2-amino alcohols with phosgene in aqueous media: limitations of the reaction selectivity as found in the presence of O-glucuronides of alprenolol in urine 413(1987)270
- Gyllenhaal, O., Lamm, B. and Vessman, J.  
Enantiomer separation of tocainide and some of its analogues and derivatives on a Chirasil-Val capillary column 411(1987)285
- Gyllenhaal, O. and Vessman, J.  
Phosgene as a derivatizing reagent prior to GC and LC (Review) 435(1988)259
- ## H
- Haagsma, N., Bathelt, E.R. and Engelsma, J.W.  
TLC screening method for the tranquilizers azaperone, propiopromazine and carazolol in pig tissues 436(1988)73
- Haagsma, N., see Van de Water, C. 411(1987)415
- Haan, J.W. De, see Claessens, H.A. 436(1988)345
- Haanen, C., see Brons, P.P.T. 422(1987)175
- Haar, J.V. Der, Kip, J. and Kraak, J.C.  
Effect of the mobile phase composition and ligand structure on the separation of D- and L-dansylamino acids, as mixed metal complexes, by reversed-phase HPLC 445(1988)219
- Haard, P.M.M. Van and Pavel, S.  
Chromatography of urinary indole derivatives (Review) 429(1988)59
- Haas, H.J., see Assmann, B. 434(1988)202
- Haas, I., see Dobránsky, T. 411(1987)486
- Haas, R.H., Breuer, J. and Hammen, M.  
HPLC measurement of selected blood citric acid cycle intermediates 425(1988)47
- Haas, T.A. and Buchanan, M.R.  
Automated HPLC extraction and quantification procedure for lipoxigenase metabolites 430(1988)1

- Habermann, R., see Josić, D. 444(1988)29
- Habu, Y., see Sugimura, K. 440(1988)131
- Hack, M.H. and Helmy, F.M.  
Application of methylamine O-deacylation and Vitride reduction to the analysis of neutral lipids: a TLC evaluation 434(1988)215
- Hada, T., see Yamamoto, T. 413(1987)342
- Haddad, P.R., see Brayan, J.G. 447(1988)249
- Haddad, P.R., see Jackson, P.E. 439(1988)37
- Haddad, P.R. and Foley, R.C.L.  
Contamination of aqueous eluents due to corrosion of stainless-steel chromatographic components 407(1987)133
- Haddad, P.R. and Heckenberg, A.L.  
Trace determination of sulfide by reversed-phase ion-interaction chromatography using pre-column derivatization 447(1988)415
- Haddad, P.R. and Jackson, P.E.  
Studies on sample preconcentration in ion chromatography. VI. Use of concentrator columns with "permanently" bound ion-interaction reagents 407(1987)121
- Haddad, P.R. and Jackson, P.E.  
Studies on sample preconcentration in ion chromatography. VIII. Preconcentration of carboxylic acids prior to ion-exclusion separation 447(1988)155
- Haddad, P.R. and Rochester, N.E.  
Ion-interaction reversed-phase chromatographic method for the determination of gold(I) cyanide in mine process liquors using automated sample preconcentration 439(1988)23
- Haddouche, A., see Dubruc, C. 417(1987)208
- Hadidi, A.-H.F.A. and Idle, J.R.  
Combined TLC-photography-densitometry for the quantitation of cyclophosphamide and its four principal urinary metabolites 427(1988)121
- Hadžija, O., Iskrić, S. and Tonković, M.  
Mobility of Fe(III)-phenolic acid complexes in TLC 402(1987)358
- Haerdi, W., see Veuthey, J.L. 445(1988)183
- Haest, G., see Cachet, M. 445(1988)290
- Hage, D.S. and Walters, R.R.  
Non-linear elution effects in split-peak chromatography. I. Computer simulations for the cases of irreversible diffusion- and adsorption-limited kinetics 436(1988)111
- Hagemeier, C.J., see Martin, C.F. 402(1987)376
- Hagen, M., see Rieck, M. 439(1988)301
- Haginaka, J., see Nakagawa, T. 420(1987)297
- Haginaka, J., see Yamaguchi, Y. 422(1987)53
- Haginaka, J., Wakai, J., Nishimura, Y. and Yasuda, H.  
LC determination of penicillins by postcolumn degradation with sodium hypochlorite using an hollow-fibre membrane reactor 447(1988)365
- Haginaka, J., Wakai, J., Yasuda, H. and Nomura, T.  
Ion-exclusion chromatography of carboxylic acids with conductivity detection. Peak enhancement using a cation-exchange hollow-fibre membrane and an alkaline solution 447(1988)373
- Haginaka, J. and Nomura, T.  
LC determination of carbohydrates with pulsed amperometric detection and a membrane reactor 447(1988)268
- Haginaka, J. and Wakai, J.  
LC determination of amoxicillin and its metabolites in human urine by postcolumn degradation with sodium hypochlorite 413(1987)219
- Hagman, A., see Jacobsson, S. 447(1988)329
- Hagman, A. and Jacobsson, S.  
Trapping efficiency of capillary cold traps 448(1988)117
- Hais, I.M.  
Tswett's letters to Claparède 440(1988)509
- Hais, I.M.  
PC 445(1988)314
- Hais, I.M., see Nobilis, M. 434(1988)363
- Hajšlová, J., Kocourek, V., Zemanová, I., Pudil, F. and Davídek, J.  
GC determination of chlorinated phenols in the form of various derivatives 439(1988)307
- Hajšlová, J., Pudil, F., Jehličková, Z., Víden, I. and Davídek, J.  
GC-MS investigation of phenoxypropanoic acid derivatives possessing herbicidal activity 438(1988)55
- Håkanson, E., see Andersson, E. 414(1987)174
- Haken, J.K.  
Use of acid or alkaline fusion for the quantitative degradation of condensation polymers 406(1987)167
- Haken, J.K., Harahap, N. and Burford, R.P.  
GC procedure for the qualitative analysis of silicone polyester resins after alkaline fusion 441(1988)207
- Haken, J.K., see Evans, M.B. 406(1987)105
- Haken, J.K. and Abraham, F.  
Demonstration of intermolecular and intramolecular hydrogen bonding by GC measurements 440(1988)329
- Haken, J.K. and Green, M.R.  
Cleavage of urea-formaldehyde resin in cross-linked alkyd systems 403(1987)145

- Hakim, L., Bourne, D.W.A. and Triggs, E.J.  
HPLC assay of cefotaxime,  
desacetylcefotaxime and ceftriaxone in rat  
plasma 424(1988)111
- Hakusui, H., see Fujimaki, Y. 421(1987)367
- Hal, H.J.M. Van, see Paanakker, J.E.  
417(1987)203
- Hall, L.K., see Martin, C.F. 402(1987)376
- Halliday, J.W., see Adams, P.C. 427(1988)341
- Halling, P.J., see McLellan, D.W. 445(1988)251
- Hamada, S., see Horikoshi, T. 416(1987)353
- Hamano, T., Mitsuhashi, Y. and Yamamoto, S.  
Determination of hydrogen peroxide in  
beverages by HPLC with fluorescence  
detection 411(1987)423
- Hamazaki, H., see Iwase, H. 416(1987)37
- Hamazaki, H., see Iwase, H. 417(1987)403
- Hamilton, M.  
LC determination of indolidan and its  
dehydro metabolite in plasma. Use of the  
Varian AASP for solid-phase extraction  
431(1988)103
- Hammen, M., see Haas, R.H. 425(1988)47
- Hammes, W. and Weyhenmeyer, R.  
Quantitative determination of vinpocetine in  
human plasma by capillary GC-MS  
413(1987)264
- Hamon, M., see Pradeau, D. 447(1988)234
- Hamper, B.C., see Pirkle, W.H. 450(1988)199
- Han, D.S., see Choi, K.E. 420(1987)81
- Han, S.M., Han, Y.I. and Armstrong, D.W.  
Structural factors affecting chiral recognition  
and separation on  $\beta$ -cyclodextrin bonded  
phases 441(1988)376
- Han, S.M., see Armstrong, D.W. 411(1987)490
- Han, S.M., see Armstrong, D.W. 448(1988)345
- Han, Y.I., see Han, S.M. 441(1988)376
- Hanai, T., see Yamaguchi, J. 441(1988)183
- Hanaoka, Y., see Murayama, T. 435(1988)417
- Hann, J.T., see Mawer, E.B. 415(1987)305
- Hansen, B., see Welinder, B.S. 408(1987)191
- Hansen-Møller, J., Dalgaard, L. and Hansen,  
S.H.  
Reversed-phase HPLC assay for the  
simultaneous determination of diflunisal and  
its glucuronides in serum and urin.  
Rearrangement of the 1-O-acylglucuronide  
420(1987)99
- Hansen, S.H., Helboe, P. and Thomsen, M.  
Separation of basic drug substances by  
reversed-phase HPLC on dynamically  
modified silica and on bonded-phase  
materials 409(1987)71
- Hansen, S.H., Helboe, P. and Thomsen, M.  
HPLC on dynamically modified silica. VIII.  
Gradient elution using eluents containing  
cetyltrimethylammonium bromide  
447(1988)182
- Hansen, S.H., see Hansen-Møller, J. 420(1987)99
- Hansson, C., Kågedal, B. and Källberg, M.  
Determination of 5-S-cysteinyldopa in human  
urine by direct injection in coupled-column  
HPLC 420(1987)146
- Hara, I., see Okazaki, M. 430(1988)135
- Hara, S., see Ijitsu, T. 427(1988)29
- Hara, T., see Shimizu, K. 432(1988)21
- Harada, H., Kamei, M., Sunada, K., Yui, S. and  
Koyama, F.  
Assay of glycosidase by lectin affinity HPLC  
407(1987)299
- Harada, K.-I., Matsuura, K., Suzuki, M., Oka,  
H., Watanabe, M.F., Oishi, S., Dahlem, A.M.,  
Beasley, V.R. and Carmichael, W.W.  
Analysis and purification of toxic peptides  
from cyanobacteria by reversed-phase HPLC  
448(1988)275
- Harada, K.-I., see Ikai, Y. 411(1987)313
- Harada, K.-I., see Oka, H. 449(1988)448
- Harada, M., Hiraoka, B.Y., Mogi, M., Fukasawa,  
K. and Fukasawa, K.M.  
HPLC separation of peptides possessing a  
proline residue in the amino-terminal  
penultimate position, and their products  
generated by enzymatic hydrolysis  
424(1988)129
- Harahap, N., see Haken, J.K. 441(1988)207
- Harduf, Z.  
Integrators as a source of inaccurate results  
404(1987)326
- Harduf, Z., Nir, T. and Juven, B.J.  
HPLC of biogenic amines, derivatized with  
9-fluorenylmethyl chloroformate  
437(1988)379
- Hardy, D.R., see Black, B.H. 437(1988)203
- Hardy, K.J., see Mihaly, G.W. 415(1987)177
- Haring, C., Humpel, C., Auer, B., Saria, A.,  
Barnas, C., Fleischhacker, W. and  
Hinterhuber, H.  
Clozapine plasma levels determined by HPLC  
with UV detection 428(1988)160
- Harkness, R.A.  
Hypoxanthine, xanthine and uridine in body  
fluids, indicators of ATP depletion (Review)  
429(1988)255
- Harmatz, J.S., see Kennedy, J.S. 422(1987)274
- Harmatz, J.S., see Kennedy, J.S. 423(1987)373
- Harper, C., see Moore, J.M. 410(1987)297
- Harrington, M.G. and Merrill, C.R.  
Cerebrospinal fluid protein analysis in

- diseases of the nervous system (Review)  
429(1988)345
- Harris, J.M., see Karr, L.J. 442(1988)219
- Harrison, D.M., see Beveridge, D.J. 450(1988)443
- Harrison, J., see Gilbert, J. 441(1988)323
- Harrison, P.M., see Keech, A.C. 426(1988)234
- Hart, B.J. 't and Wilting, J.  
Sensitive GC method for determining  
nitrazepam in serum and saliva 424(1988)403
- Hart, R.J., see Sworn, G. 403(1987)307
- Hart, S.J., see Aguilar, M.I. 426(1988)315
- Harten, J. Van, Lodewijks, M.T.M., Guyt-  
Scholten, J.W., Van Brummelen, P. and  
Breimer, D.D.  
GC determination of nisoldipine and one of  
its metabolites in plasma 423(1987)327
- Hartley, R., Luccock, M. and Becker, M.  
Improved HPLC procedure for the  
determination of chlormethiazole levels  
following solid-phase extraction from plasma  
415(1987)357
- Hartley, R.D., see Ford, C.W. 436(1988)484
- Hartman, C.A., see Pivonka, J. 420(1987)89
- Hartman, P.A. and Stodola, J.D.  
Reversed-phase HPLC of fertirelin acetate  
and related compounds 444(1988)177
- Hartmann, D.J., see Ricard-Blum, S.  
431(1988)474
- Hartmann, M., see Klemm, D. 438(1988)122
- Hartwick, R.A., see Raglione, T.V. 409(1987)205
- Hartwick, R.A., see Raglione, T.V. 409(1987)213
- Hartwick, R.A., see Sagliano, Jr., N.  
443(1988)155
- Harvey, D.J., Tiffany, J.M., Duerden, J.M.,  
Pandher, K.S. and Mengher, L.S.  
Identification by combined GC-MS of  
constituent long-chain fatty acids and  
alcohols from the meibomian glands of the  
rat and a comparison with human meibomian  
lipids 414(1987)253
- Harvey, T.G., Matheson, T.W. and Pratt, K.C.  
Extraction and identification of nitrile  
compounds in Rundle shale oil 435(1988)193
- Harvey-White, J., see Hovevey-Sion, D.  
426(1988)141
- Hary, L. and Andrejak, M.  
Determination of serum cefotaxime and  
desacetylcefotaxime by ion-pair HPLC (in  
French) 419(1987)396
- Hase, A., see Björkqvist, B. 435(1988)501
- Hase, S., Ikenaka, K., Mikoshiba, K. and  
Ikenaka, T.  
Analysis of tissue glycoprotein sugar chains  
by two-dimensional HPLC mapping  
434(1988)51
- Hasegawa, I., Sakka, S., Kuroda, K. and Kato,  
C.  
Trimethylsilylation of the hydrolysed and  
polycondensed products of  
methyltriethoxysilane 410(1987)137
- Hashida, N., see Seki, T. 424(1988)410
- Hashidzume, K., see Tanaka, N. 448(1988)95
- Hashimoto, H., Koyama, Y. and Shimamura, T.  
Isolation of *cis-trans* isomers of  
canthaxanthin by HPLC using a calcium  
hydroxide column and identification of their  
configurations by <sup>1</sup>H NMR spectroscopy  
448(1988)182
- Hashimoto, H., see Koyama, Y. 439(1988)417
- Hashimoto, H., see Yamashita, J. 403(1987)275
- Hashimoto, K., Sekine, T., Omori, T. and  
Yoshihara, K.  
Adsorption behaviour of tris( $\beta$ -  
diketonato)metal(III) complexes on silica gel  
438(1988)401
- Hashimoto, K., Shirai, Y., Adachi, S. and Horie,  
M.  
Effect of the content of divinylbenzene in  
ion-exchange resins on the chromatographic  
separation of  $\alpha$ -cyclodextrin and glucose  
448(1988)241
- Hashimoto, T., see Kato, Y. 404(1987)333
- Hashimoto, T., see Kato, Y. 447(1988)212
- Hashimoto, Y., see Moriyasu, M. 411(1987)466
- Hashimoto, Y., see Takeuchi, T. 402(1987)328
- Hashimoto, Y., see Yamashita, K. 424(1988)39
- Hashizume, Y., see Sasaoka, T. 428(1988)152
- Hassan, M. and Ehrsson, H.  
Determination of busulfan in human plasma  
by GC with electron-capture detection  
434(1988)
- Hatada, K., see Okamoto, Y. 448(1988)454
- Hatano, H., see Kihara, K. 410(1987)103
- Hatano, H., see Murayama, T. 435(1988)417
- Hatano, H., see Yamamoto, F.M. 408(1987)21
- Hatano, T., Yoshida, T. and Okuda, T.  
Chromatography of tannins. III. Multiple  
peaks in HPLC of some hydrolyzable tannins  
435(1988)285
- Hatsumura, H., see Sayo, H. 417(1987)129
- Hatsumura, H., see Sayo, H. 426(1988)449
- Hatta, M., see Hirano, T. 417(1987)371
- Hattori, T., see Yamada, H. 411(1987)401
- Haughey, D.B. and Jusko, W.J.  
Analysis of methylprednisolone,  
methylprednisone and corticosterone for  
assessment of methylprednisolone disposition  
in the rat 430(1988)241
- Hausleiter, H.-J., see Dahmen, W. 427(1988)157
- Hausleiter, H.J., see Gielsdorf, W. 419(1987)351

- Hausleiter, H.J., see Gielsdorf, W. 419(1987)357
- Haverkort, W.A., see De Korte, D. 415(1987)383
- Hawkes, S.J.  
Does the obstruction factor change with flow-rate? 436(1988)
- Hawthorne, S.B. and Miller, D.J.  
Directly coupled supercritical fluid extraction-GC analysis of polycyclic aromatic hydrocarbons and polychlorinated biphenyls from environmental solids 403(1987)63
- Hayakawa, K. and Oizumi, J.  
Determination of free biotin in plasma by LC with fluorimetric detection 413(1987)247
- Hayakawa, K. and Oizumi, J.  
Determination of lipoamidase activity by LC with fluorimetric detection 423(1987)304
- Hayasaka, Y., see Sitaram, B.R. 422(1987)13
- Hayashi, N., see Suzuki, Y. 426(1988)33
- Hayashi, S., see Matsukura, H. 434(1988)196
- Hayashi, T., Naruse, H., Matsuda, F. and Iida, Y.  
Sensitive determination of deuterated and non-deuterated indole-3-acetic acid and 5-hydroxyindole-3-acetic acid by combined capillary GC negative-ion chemical ionization MS 428(1988)209
- Hayashi, Y., see Matsuda, R. 438(1988)319
- Hayashi, Y., Shibasaki, T., Matsuda, R. and Uchiyama, M.  
Real-time quantitative analysis of unresolved LC peaks with the use of the Kalman filter 407(1987)59
- Hayashi, Y., Shibasaki, T. and Uchiyama, M.  
Resolution of overlapped chromatograms by the use of a Kalman filter. III. Effects of overlapping and outstripping of a solvent peak on the analysis 411(1987)95
- Hayashibara, M., see Katagiri, Y. 431(1988)135
- Hayashida, M., see Jinno, K. 436(1988)11
- Hayes, Jr., P.C. and Anderson, S.D.  
Paraffins, olefins, naphthenes and aromatics analysis of selected hydrocarbon distillates using on-line column switching HPLC with dielectric constant detection 437(1988)365
- Hazai, I., see Alexander, G. 446(1988)87
- He, F.-Y., see Armstrong, D.W. 448(1988)345
- Healey, M.C., see Hwang, H.H. 430(1988)329
- Hearn, M.T.W.  
General strategies in the separation of proteins by HPLC methods (Review) 418(1987)3
- Hearn, M.T.W., Aguilar, M.I., Mant, C.T. and Hodges, R.S.  
HPLC of amino acids, peptides and proteins. LXXXV. Evaluation of the use of hydrophobicity coefficients for the prediction of peptide elution profiles 438(1988)197
- Hearn, M.T.W., Aguilar, M.I., Nguyen, T. and Fridman, M.  
HPLC of amino acids, peptides and proteins. LXXXIV. Application of derivative spectroscopy to the study of column residency effects in the reversed-phase and size-exclusion LC separation of proteins 435(1988)271
- Hearn, M.T.W., Hodder, A.N. and Aguilar, M.I.  
HPLC of amino acids, peptides and proteins. LXXXVI. The influence of different displacer salts on the retention and bandwidth properties of proteins separated by isocratic anion-exchange chromatography 443(1988)97
- Heavey, D., see Richmond, R. 417(1987)241
- Hechler, J.-J., see Noël, D. 408(1987)129
- Hechler, J.-J., see Noël, D. 447(1988)141
- Heckenberg, A.L., see Haddad, P.R. 447(1988)415
- Hecker, R. and Riesner, D.  
Chromatographic separation of DNA restriction fragments (Review) 418(1987)97
- Hedin, P.A., see Lydon, J. 410(1987)474
- Heiffer, M.H., see Kapetanovic, I.M. 419(1987)458
- Heil, B., see Lecomte, L. 408(1987)416
- Heilmann, P., see Schöneshöfer, M. 417(1987)434
- Heimhuber, B., Galensa, R. and Herrmann, K.  
HPLC determination of polymethoxylated flavones in orange juice after solid-phase extraction 439(1988)481
- Heimler, D. and Vidrich, V.  
TLC characterization of essential oils 448(1988)301
- Hein, S.J., see Ramus, T.L. 404(1987)155
- Heiningen, A. Van, see Derks, H.J.G.M. 442(1988)436
- Heinitz, M.L., Flanigan, E., Orłowski, R.C. and Regnier, F.E.  
Correlation of calcitonin structure with chromatographic retention in HPLC 443(1988)229
- Heinitz, M.L., Kennedy, L., Kopaciewicz, W. and Regnier, F.E.  
Chromatography of proteins on hydrophobic interaction and ion-exchange chromatographic matrices: mobile phase contributions to selectivity 443(1988)173
- Heirwegh, K.P.M., see Rothuizen, J. 427(1988)19
- Heizmann, P., Jordan, J.-C. and Ludwig, B.  
Determination of climazolam in the plasma of farm animals by GC 415(1987)170

- Hekkens, W.T.J.M., see Kocna, P. 434(1988)429
- Helboe, P., see Hansen, S.H. 409(1987)71
- Helboe, P., see Hansen, S.H. 447(1988)182
- Hellberg, U., see Johansson, B.-L. 403(1987)85
- Hellgren, U., see Bergqvist, Y. 432(1988)253
- Helliger, W., see Lindner, H. 450(1988)309
- Helms, M., see Zobel, H. 446(1988)305
- Helmy, F.M., see Hack, M.H. 434(1988)215
- Hemmendinger, S., see Leray, C. 420(1987)411
- Henatsch, J.J. and Jüttner, F.  
Capillary GC analysis of low-boiling organic sulphur compounds in anoxic lake-water by cryoadsorption 445(1988)97
- Henderson, P.T., see Evelo, C.T.A. 420(1987)35
- Henderson, P.T., see Evelo, C.T.A. 427(1988)335
- Henderson, P.T., see Jongeneelen, F.J. 413(1987)227
- Hendrickson, S.M., see Wenzel, T.J. 433(1988)149
- Hendriks, H., Huizing, H.J. and Bruins, A.P.  
Ammonium positive-ion and hydroxide negative-ion chemical ionization GC-MS for the identification of pyrrolizidine alkaloids in *Eupatorium rotundifolium* L. var. *ovatum* 428(1988)352
- Hengartner, H., see Schott, H. 441(1988)115
- Henion, J.D., see Hsu, S.-H. 424(1988)219
- Henion, J.D., see Weidolf, L.O.G. 433(1988)9
- Henne, E.-M., see Brendel, E. 426(1988)406
- Henner, W.D., Furlong, E.A., Flaherty, M.D., Shea, T.C. and Peters, W.P.  
Measurement of busulfan in plasma by HPLC 416(1987)426
- Henning, K., see Wallasch, T.M. 425(1988)175
- Henriques, H.F., see Frantz, S.C. 444(1988)41
- Hensler, T., see Knöller, J. 427(1988)199
- Henton, D., see Vreman, H.J. 427(1988)315
- Henzel, W.J., Rodriguez, H. and Watanabe, C.  
Computer analysis of automated Edman degradation and amino acid analysis data 404(1987)41
- Hepp, M.A. and Klee, M.S.  
Characterization of porous polymers by polar strength and selectivity 404(1987)145
- Herbeval, X., see Brossat, B. 413(1987)141
- Herbranson, D.E., Kriss-Danziger, P., Rosenberg, L.S. and Hostetler, C.  
Development of a TLC method for the evaluation of antiarrhythmic N-{[3,5-di(pyrrolidinylmethyl)-4-hydroxy]benzoyl}anilin (ACC-9358) 410(1987)222
- Herche, L.R., see Parrish, C.C. 435(1988)350
- Hermann, P., see Dubruc, C. 417(1987)208
- Hermann, W., see Larsson, K. 450(1988)71
- Hermez, I., see Papp, O. 403(1987)11
- Hermez, I., see Papp, O. 403(1987)19
- Hernádi, F.J., see Gál, S. 403(1987)217
- Hernádi, F.J., see Kiss, L. 448(1988)109
- Herraiz, M., see Reglero, G. 438(1988)243
- Herraiz, T., see Reglero, G. 438(1988)243
- Herregodts, P., Michotte, Y. and Ebinger, G.  
Combined ion-pair extraction and HPLC for the determination of the biogenic amines and their major metabolites in single brain tissue samples 421(1987)51
- Herrero, P. and Roselli, R.J.  
Computer analysis of proteins separated by polyacrylamide gradient pore gel electrophoresis 416(1987)265
- Herring, S.W., see Shitanishi, K.T. 444(1988)107
- Herrle, I., see Kauert, G. 428(1988)291
- Herrmann, K., see Heimhuber, B. 439(1988)481
- Herrmann, K., see Klick, S. 441(1988)335
- Herrmann, K., see Risch, B. 448(1988)291
- Herrmann, T., see Knospe, J. 442(1988)444
- Herrmann, T., Steinhilber, D., Knospe, J. and Roth, H.J.  
Determination of picogram amounts of lipoxin A<sub>4</sub> and lipoxin B<sub>4</sub> by HPLC with electrochemical detection 428(1988)237
- Herrmann, T., Steinhilber, D. and Roth, H.J.  
Determination of leukotriene B<sub>4</sub> by HPLC with electrochemical detection 416(1987)170
- Hertel, C.C., see Unadkat, J.D. 430(1988)420
- Herzan, L.A., see Vatassery, G.T. 433(1988)312
- Hetper, J., see Dziwiński, E. 446(1988)103
- Heulin, M.-H., see Brossat, B. 413(1987)141
- Hevesi, T., see Krupčík, J. 406(1987)117
- Hevesi, T., see Krupčík, J. 407(1987)65
- Hevesi, T., see Krupčík, J. 448(1988)203
- Hewetson, D., see Singh, A.K. 404(1987)223
- Heyes, M.P. and Quearry, B.J.  
Quantification of 3-hydroxykynurenine in brain by HPLC and electrochemical detection 428(1988)340
- Heykants, J., see Woestenborghs, R. 413(1987)332
- Heykants, J., see Woestenborghs, R. 424(1988)195
- Heys, J.R.  
HPLC studies of pH dependence in the isotopic fractionation of deuterated benzazepines 407(1987)37
- Hicks, K.B. and Hotchkiss, Jr., A.T.  
HPLC of plant-derived oligosaccharides on a new cation-exchange resin stationary phase: HPX-22H 441(1988)382
- Higashi, A., see Inotsume, N. 428(1988)402
- Higashino, K., see Yamamoto, T. 413(1987)342
- Higgins, H.W., see Mackey, D.J. 436(1988)243
- Higuchi, N., see Manabe, T. 423(1987)115

- Higuchi, N., see Manabe, T. 431(1988)45
- Hildebrand, M. and Schütt, A.  
Column LC of the novel aldosterone antagonist, mespirenone, and its active metabolite in plasma 414(1987)217
- Hill, D.L., see Kalin, J.R. 431(1988)184
- Hill, D.L., see Rose, L.M. 425(1988)419
- Hill, D.L., see Waud, W.R. 425(1988)220
- Hill, E., Humphreys, E. and Malcolme-Lawes, D.J.  
Electrochemiluminescence as a detection technique for reversed-phase HPLC. IV. Detection of fluorescent derivatives 441(1988)394
- Hill, H.H., see Gholson, A.R. 408(1987)329
- Hill, H.M., see Selinger, K. 434(1988)259
- Hiltunen, R., see Ekundayo, O. 403(1987)358
- Himmel, M.E., see Baker, J.O. 437(1988)387
- Hines, J.W., see Blau, P.A. 420(1987)1
- Hinterhuber, H., see Haring, C. 428(1988)160
- Hintz, R.L., see Powell, D.R. 420(1987)163
- Hirabayashi, Y., Nakao, T., Matsumoto, M., Obata, K. and Ando, S.  
Improved method for large-scale purification of brain gangliosides by Q-Sepharose CC. Immunochemical detection of C-series polysialogangliosides in adult bovine brains 445(1988)377
- Hiraga, Y., see Iwaki, K. 407(1987)273
- Hirai, K., see Kominami, G. 417(1987)216
- Hirai, Y., see Kitano, H. 420(1987)13
- Hiraiki, H., Kimura, M. and Itokawa, Y.  
Determination of K vitamins (phyloquinone and menaquinones) in umbilical cord plasma by a platinum-reduction column 430(1988)143
- Hirano, T., Hatta, M., Oka, K. and Wakasugi, K.  
LC determination of serum lipid peroxides as diphenylthiobarbituric acid-reactive substance to collagen disease patients—a preliminary study 417(1987)371
- Hirano, T., see Oka, K. 423(1987)285
- Hiraoka, B.Y., see Harada, M. 424(1988)129
- Hiraoka, E., see Yamaji, A. 415(1987)143
- Hirata, K., see Arai, T. 448(1988)439
- Hirata, N., Kasai, M., Yanagihara, Y. and Noguchi, K.D.  
Separation of peptides by multimode chromatography on a column packed with vinyl alcohol copolymer gel 434(1988)71
- Hirate, J., see Bhargava, V.O. 426(1988)212
- Hirauchi, K., Sakano, T., Nagaoka, T. and Morimoto, A.  
Simultaneous determination of vitamin K<sub>1</sub>, vitamin K<sub>1</sub> 2,3-epoxide and menaquinone-4 in human plasma by HPLC with fluorimetric detection 430(1988)21
- Hirayama, C., Ihara, H., Shiba, M., Nakamura, M., Motozato, Y. and Kunitake, T.  
Macroporous glucomannan spheres for the size-exclusion separation of aqueous macromolecules 409(1987)175
- Hirayama, C., Ihara, H. and Tanaka, K.  
Chromatographic resolution of dipeptide enantiomers and diastereomers on chiral stationary phases from poly(L-leucine) or poly(L-phenylalanine) 450(1988)271
- Hirayama, N. and Kuwamoto, T.  
Non-suppressed ion chromatography of arsenic anions with potassium hydroxide–aromatic salt mixed eluents 447(1988)323
- Hirokawa, T., Kobayashi, S. and Kiso, Y.  
Complex-forming equilibria in isotachopheresis. VII. Evaluation of the ion-pair formation constants of phosphorus oxo acids with histidine and assessment of their separability 410(1987)279
- Hirokawa, T., Tsuyoshi, T. and Kiso, Y.  
Isotachopheretic determination of mobility and pK<sub>a</sub> by means of computer simulation. VI. Mobility and pK<sub>a</sub> of sixty anions in methanol 408(1987)27
- Hirooka, T., see Naoi, M. 426(1988)75
- Hirose, A. and Ishii, D.  
Multiple detection system for the high-resolution microcolumn SEC separation of proteins with a slurry-packed capillary column 411(1987)21
- Hirose, A. and Ishii, D.  
Microcolumn chromatofocusing. III. Effect of the analysis time on the resolution and the use of a semi-disposable microcolumn 438(1988)15
- Hirose, S., see Tawa, R. 425(1988)143
- Hirose, S., see Yoshida, S. 430(1988)156
- Hirose, S., see Yoshida, S. 431(1988)27
- Hirota, N., Furuta, T. and Kasuya, Y.  
Determination of cortisol in human plasma by capillary GC–MS using [<sup>2</sup>H<sub>5</sub>] cortisol as an internal standard 425(1988)237
- Hirota, T., see Ohmori, S. 415(1987)221
- Hittner, D.M. and Cowman, M.K.  
High-performance GPC of glycosaminoglycans. Column calibration by gel electrophoresis 402(1987)149
- Hirada, K., see Takada, Y. 421(1987)350
- Hizukuri, S., see Koizumi, K. 409(1987)396
- Hizukuri, S., see Koizumi, K. 437(1988)47
- Hjertén, S., Elenbring, K., Kilár, F., Liao, J.-L.,

- Chen, A.J.C., Siebert, C.J. and Zhu, M.-D. Carrier-free zone electrophoresis, displacement electrophoresis and isoelectric focusing in a high-performance electrophoresis apparatus 403(1987)47
- Hjertén, S., Lindeberg, J. and Shopova, B. High-performance adsorption chromatography of proteins on 2- $\mu$ m spherical beads of hydroxyapatite 440(1988)305
- Ho, J., Guthrie, R. and Tieckelmann, H. Quantitative determination of porphyrins, their precursors and zinc protoporphyrin in whole blood and dried blood by HPLC with fluorimetric detection 417(1987)269
- Hochuli, E. Large-scale chromatography of recombinant proteins 444(1988)293
- Hochuli, E., Döbeli, H. and Schacher, A. New metal chelate adsorbent selective for proteins and peptides containing neighbouring histidine residues 411(1987)177
- Hochuli, E., Gillessen, D. and Kocher, H.P. Specificity of the immunoabsorbent used for large-scale recovery of interferon  $\alpha$ -2a 411(1987)371
- Hodder, A.N., see Hearn, M.T.W. 443(1988)97
- Hodges, R.S., Burke, T.W.L. and Mant, C.T. Preparative purification of peptides by reversed-phase chromatography. Sample displacement mode *versus* gradient elution mode 444(1988)349
- Hodges, R.S., see Hearn, M.T.W. 438(1988)197
- Hodges, R.S., see Mant, C.T. 409(1987)155
- Hodr, R., see Knobloch, E. 428(1988)255
- Hoenders, H.J., see Van den Oetelaar, P.J.M. 411(1987)507
- Hoerr, D.C., see Weaner, L.E. 437(1988)109
- Hoeven, R.A.M. Van der, see Tjaden, U.R. 420(1987)53
- Hoevers, J.W., see Schellens, J.H.M. 431(1988)203
- Hoffman, D.W., Edkins, R.D., Shillcutt, S.D. and Salama, A. New HPLC method for fluphenazine and metabolites in human plasma 414(1987)504
- Hoffman, N.E., see Rahman, A. 428(1988)395
- Hoffman, N.E., see Rahman, A. 433(1988)159
- Hoffman, N.E., see Rustum, A.M. 421(1987)387
- Hoffman, N.E., see Rustum, A.M. 421(1987)418
- Hoffman, N.E., see Rustum, A.M. 422(1987)125
- Hoffman, N.E., see Rustum, A.M. 426(1988)121
- Hoffmann, G. and Sweetman, L. O-(2,3,4,5,6-Pentafluorobenzyl)oxime-trimethylsilyl ester derivatives for quantitative GC and GC-MS studies of aldehydes, ketones and oxoacids 421(1987)336
- Hoffmann, T.J., see Thompson, R.D. 438(1988)369
- Hofmann, W., see Josić, D. 444(1988)29
- Hofstetter, M., Nyiredy, S., Meier, B. and Sticher, O. Reversed-phase HPLC separation of the tetrameric lectin concanavalin A from *Canavalia ensiformis* 407(1987)319
- Hogan, E.L., see Leskawa, K.C. 411(1987)393
- Hogan, T.F., see Mehta, R.C. 430(1988)341
- Hogendoorn, E.A., see Goewie, C.E. 404(1987)352
- Hogendoorn, E.A., see Goewie, C.E. 410(1987)211
- Höglund, P. and Nilsson, L.-G. Liquid chromatographic determination of diltiazem and its metabolites using *trans* isomers as internal standards, with dynamic modification of the solid phase by addition of an amine to the mobile phase 414(1987)109
- Höglund, U. Use of perfluorinated carboxylic acids in the separation of metabolites of vasopressin prior to radioimmunoassay 430(1988)128
- Hogstrand, C., see Olsson, P.-E. 402(1987)293
- Hohaus, E., see Winkler, E. 436(1988)447
- Holzinger, L., see Kremmer, T. 415(1987)53
- Holder, C.L., Korfmacher, W.A., Rushing, L.G., Thompson, Jr., H.C., Slikker, Jr., W. and Gosnell, A.B. Formation of artifactual metabolites of doxylamine following acid hydrolysis 419(1987)113
- Holder, G.M., see Duke, C.C. 430(1988)53
- Holík, J., see Dufek, P. 435(1988)249
- Hollaert, C., see Brassinne, C. 419(1987)401
- Holland, M.L., Uetz, J.A. and Ng, K.T. Automated capillary GC assay using flame ionization detection for the determination of topiramate in plasma 433(1988)276
- Holland, O.B., Brown, H., Risk, M., Paone, R., Dube, P. and Gomez-Sanchez, C. Simplified method for the determination of plasma 18-hydroxycorticosterone 426(1988)148
- Holland, P.T., see France, N.C. 433(1988)1
- Höllner, M., see Weber, C. 426(1988)345
- Holley, J.H., see Kitzman, J.V. 437(1988)306
- Hollis, D., see Cheng, W. 408(1987)9
- Holmbom, B., see Sjöström, J. 411(1987)363
- Holt, D.L., Wehling, R.L. and Zeece, M.G. Determination of native folates in milk and other dairy products by HPLC 449(1988)271



- Holt, D.W., see Bhamra, R.K. 417(1987)229
- Holthuis, J.J.M., see Van der Horst, F.A.L. 426(1988)267
- Holthuis, J.J.M., see Van Opstal, M.A.J. 432(1988)395
- Holthuis, J.J.M., see Vendrig, D.E.M.M. 414(1987)91
- Holthuis, J.J.M., see Vendrig, D.E.M.M. 424(1988)83
- Holthuis, J.J.M., see Vendrig, D.E.M.M. 434(1988)145
- Hölzel, C., see Dietel, P. 422(1987)1
- Homer, R.B., see Bayliss, M.A.J. 445(1988)393
- Homer, R.B., see Bayliss, M.A.J. 445(1988)403
- Honda, A., see Ukai, S. 435(1988)374
- Honda, M., see Inotsume, N. 428(1988)402
- Honda, S., Iwase, S. and Fujiwara, S.  
Evaluation of an automatic siphonic sampler for capillary zone electrophoresis 404(1987)313
- Honda, S., Ohmura, T., Segawa, Y., Minematsu, T. and Kakehi, K.  
LC assay of the relative activities of serum pancreatic and salivary  $\alpha$ -amylase using reductively pyridylaminated maltopentaose as a fluorescent substrate 419(1987)51
- Honda, S., see Iwase, S.F.S. 447(1988)133
- Honda, S., Suzuki, S., Nitta, T. and Kakehi, K.  
Analytical high-performance affinity chromatography of ovalbumin-derived glycopeptides on columns of concanavalin A- and wheat germ agglutinin-immobilized gels 438(1988)73
- Hondo, T., see Jinno, K. 436(1988)11
- Hong, E., see Flores-Murrieta, F.J. 428(1988)167
- Hong, S.-Y., see Park, K.S. 440(1988)225
- Hong, W.-H., see Geria, T. 450(1988)407
- Honkasalo, T. and Nissinen, E.  
Determination of phenol sulphotransferase activity by HPLC 424(1988)136
- Hoogmartens, J., see Cachet, T. 403(1987)343
- Hoogmartens, J., see Cachet, T. 409(1987)91
- Hoogmartens, J., see Cachet, T. 445(1988)290
- Hoogmartens, J., see Khan, N.H. 405(1987)229
- Hoogmartens, J., see Kibwage, I.O. 434(1988)177
- Hooper, W.D., Cannell, G.R. and Dickinson, R.G.  
Sensitive and selective assay of danazol in plasma by HPLC 416(1987)347
- Hoorn, F.A.J. Van der, see Boomsma, F. 427(1988)219
- Hoppe, A., see Kågedal, B. 417(1987)261
- Hoppel, C.L., see Chai, B. 430(1988)93
- Hoppel, C.L., see Minkler, P.E. 413(1987)33
- Hoppel, C.L., see Minkler, P.E. 420(1987)385
- Hoppel, C.L., see Minkler, P.E. 428(1988)388
- Hoque, E.  
Isocratic reversed-phase HPLC analysis of pigments in Norway spruce 448(1988)417
- Horai, Y., see Kubota, K. 425(1988)67
- Horai, Y., see Oshima, J. 414(1987)381
- Horák, D., see Tennikova, T.B. 435(1988)357
- Horak, R.M., see De Jesus, A.E. 450(1988)101
- Horáková, M., see Deyl, Z. 426(1988)162
- Horie, M., Saito, K., Hoshino, Y., Nose, N., Mochizuki, E. and Nakazawa, H.  
Simultaneous determination of nalidixic acid, oxolinic acid and piromidic acid in fish by HPLC with fluorescence and UV detection 402(1987)301
- Horie, M., see Hashimoto, K. 448(1988)241
- Horikoshi, T., Koga, T. and Hamada, S.  
TLC using multiple development for analysis of reaction products of sucrases 416(1987)353
- Horitsu, H., see Idaka, E. 420(1987)373
- Hörmeyer, H.F., see Derler, H. 440(1988)281
- Horn, A.S., see Biessels, P.T.M. 421(1987)392
- Horne, R., see Blackett, A. 433(1988)326
- Horst, F.A.L. Van der and Holthuis, J.J.M.  
Study of the derivatization of *n*-alkylamines with 1-fluoro-2,4-dinitrobenzene in the presence of aqueous cetyltrimethylammonium bromide micelles 426(1988)267
- Horvath, A.M., see Ferry, J.J. 421(1987)187
- Horváth, C., see Antia, F.D. 435(1988)1
- Horváth, C., see Huang, J.-X. 406(1987)275
- Horváth, C., see Huang, J.-X. 406(1987)285
- Horváth, C., see Kalghatgi, K. 443(1988)343
- Horváth, C., see Lee, A.L. 443(1988)31
- Horváth, C., see Maa, Y.-F. 445(1988)71
- Horváth, C., see Velayudhan, A. 435(1988)397
- Horváth, C., see Velayudhan, A. 443(1988)13
- Horváth, C., see Viscomi, G. 440(1988)157
- Horváth, G., see Szederkényi, F. 446(1988)253
- Horwood, J.F. and Urbach, G.E.  
Modification to the chromatographic behaviour of Porapak Q by chemical reaction 437(1988)238
- Hošek, B., see Boháček, J. 434(1988)439
- Hoshino, Y., see Horie, M. 402(1987)301
- Hoskins, J.A. and Davis, L.J.  
Analysis of the isomeric tyrosines in mammalian and avian systems using HPLC with fluorescence detection 426(1988)155
- Hosoe, M., Oi, T., Kawada, K. and Kakihana, H.  
Separation of rubidium isotopes by ion-exchange chromatography 435(1988)253
- Hosoe, M., Oi, T., Kawaka, K. and Kakihana, H.  
Fractionation of rubidium isotopes in

- cation-exchange chromatography 438(1988)225
- Hosoe, M., see Oi, T. 450(1988)343
- Hosokawa, M., see Sayo, H. 417(1987)129
- Hosokawa, M., see Sayo, H. 426(1988)449
- Hosomi, M., see Koyama, Y. 439(1988)417
- Hosotsubo, H., Takahara, S. and Taenaka, N. Simplified HPLC method for determination of mizoribine in human serum 432(1988)340
- Hostetler, C., see Herbranson, D.E. 410(1987)222
- Hostettmann, K., see Décosterd, L.A. 406(1987)367
- Hostettmann, K., see Marston, A. 450(1988)3
- Hostettmann, K., see Marston, A. 450(1988)91
- Hostettmann, K., see Slacanin, I. 448(1988)265
- Hostomská, Z., see Mikeš, O. 440(1988)287
- Hotchkiss, Jr., A.T., see Hicks, K.B. 441(1988)382
- Hotchkiss, S.A., see Bailey, E. 416(1987)81
- Hotchkiss, S.A. and Caldwell, J. HPLC assay for theophylline and its major metabolites in human urine 423(1987)179
- Hotta, K., see Iwase, H. 416(1987)37
- Hotta, K., see Iwase, H. 417(1987)403
- Hou, W., see Wang, E. 447(1988)256
- Hountondji, C., see Beauvallet, C. 438(1988)347
- Houston, J.B., see Adedoyin, A. 416(1987)160
- Hovevey-Sion, D., Harvey-White, J., Kopin, I.J. and Goldstein, D.S. Measurement of homovanillic acid in small volumes of plasma using LC with electrochemical detection 426(1988)141
- Hoving, E.B., Jansen, G., Volmer, M., Van Doormaal, J.J. and Muskiet, F.A.J. Profiling of plasma cholesterol ester and triglyceride fatty acids as their methyl esters by capillary GC, preceded by a rapid aminopropyl-silica column chromatographic separation of lipid classes 434(1988)395
- Hoving, E.B., Prins, J., Rutgers, H.M. and Muskiet, F.A.J. Behaviour of plasmalogens during HPLC on a silica column with a mobile phase containing phosphoric acid 434(1988)411
- Howard, II, R.R., Peterson, T. and Kastl, P.R. HPLC determination of ascorbic acid in human tears 414(1987)434
- Howell, S., see Smith, R.E. 439(1988)83
- Howell, S.B., see Velury, S. 424(1988)141
- Howells, D.W., see Hyland, K. 429(1988)95
- Hoyer, G.L. Improved HPLC method for the analysis of serum sotalol 427(1988)181
- Hrabova, H., see Gottvaldova, M. 427(1988)331
- Hrivňáková, A., see Boháček, J. 434(1988)439
- Hrkal, Z., see Cukrová, V. 413(1987)242
- Hrudey, S.E., see Birkholz, D.A. 449(1988)251
- Hsia, J.C., see McGarrity, M.J. 413(1987)53
- Hsia, J.C., see McGarrity, M.J. 415(1987)136
- Hsia, J.C., see McGarrity, M.J. 419(1987)37
- Hsieh, J.Y.-K., see Rothwell, C.M. 417(1987)447
- Hsieh, J.Y.-K., Thornton, T.J. and Bayne, W.F. HPLC determination of enprofylline in human plasma 430(1988)175
- Hsu, D.-S. and Chen, S.S. Assay of 5'-nucleotidase and simultaneous detection of interfering enzymes by anion-exchange CC 420(1987)398
- Hsu, J.T. and Chen, T.-L. Theoretical analysis of the asymmetry in chromatographic peaks 404(1987)1
- Hsu, S.-H., Eckerlin, R.H. and Henion, J.D. Identification and quantitation of trenbolone in bovine tissue by GC-MS 424(1988)219
- Hu, R.Z., see Tian, Z.W. 439(1988)151
- Hu, R.Z., see Tian, Z.W. 439(1988)159
- Hu, W.L., see Tian, Z.W. 439(1988)151
- Hua, R.L., see Peng, C.T. 436(1988)137
- Huang, J.-X. and Horváth, C. Adsorption isotherms on HPLC sorbents. I. Peptides and nucleic acid constituents on octadecyl-silica 406(1987)275
- Huang, J.-X. and Horváth, C. Adsorption isotherms on HPLC sorbents. II. Proteins on cation exchangers with silica support 406(1987)285
- Huang, S.-M. and Huang, Y.-C. Chromatography and electrophoresis of creatinine and other guanidino compounds (Review) 429(1988)235
- Huang, S.-M. and Tran-Phan, H. High-performance liquid chromatographic determination of a new H<sub>2</sub> blocker (ORF 17910) in plasma 423(1987)358
- Huang, X., Gordon, M.J. and Zare, R.N. Quantitation of Li<sup>+</sup> in serum by capillary zone electrophoresis with an on-column conductivity detector 425(1988)385
- Huang, Y.-C., see Huang, S.-M. 429(1988)235
- Huang Chandler, M.H., Guttendorf, R.J., Blouin, R.A. and Wedlund, P.J. Simultaneous quantitation of *d*- and *l*-hexobarbital in rat blood by HPLC 419(1987)426
- Hubbard, J.W., see Gupta, M.B. 424(1988)189
- Hubbard, J.W., see Srinivas, N.R. 433(1988)105
- Hubbard, R.W., Chambers, J.G., Sanchez, A., Slocum, R. and Lee, P. Amino acid analysis of plasma: studies in sample preparation 431(1988)163
- Huber, J.F.K., see Daučík, P. 442(1988)53

- Huber, W.G., see Kitzman, J.V. 437(1988)306
- Huberman, A. and Aguilar, M.B.  
Single-step purification of two hyperglycaemic neurohormones from the sinus gland of *Procambarus bowieri*.  
Comparative peptide mapping by means of HPLC 443(1988)337
- Hudecz, F., see Szókán, G. 444(1988)115
- Huet, J.C., see Billard, V. 440(1988)87
- Huff, D.L., see Sowell, A.L. 431(1988)424
- Hughes, B., see Benedek, K. 444(1988)191
- Hughes, D.E. and Bramer, A.M.  
Anomalous LC results originating from the presence of soap residues and the use of nylon filters 408(1987)296
- Huhtikangas, A., see Naaranlahti, T. 410(1987)488
- Huisman, T.H.J.  
Separation of hemoglobins and hemoglobin chains by HPLC (Review) 418(1987)277
- Huizing, H.J., see Hendriks, H. 428(1988)352
- Huizing, H.J., see Van Gelder, W.M.J. 442(1988)133
- Hulshoff, A., see Vendrig, D.E.M.M. 414(1987)91
- Humbert, H., Denouel, J., Chervet, J.P., Lavene, D. and Kiechel, J.R.  
Determination of sub-nanogram amounts of dihydroergotamine in plasma and urine using LC and fluorimetric detection with off-line and on-line solid-phase drug enrichment 417(1987)319
- Humbert, H., Denouel, J. and Keller, H.P.  
Column LC determination of hydrolysed bopindolol, in the picogram per millilitre range in plasma, using cartridge extraction and dual electrochemical detection 422(1987)205
- Humpel, C., see Haring, C. 428(1988)160
- Humphreys, E., see Hill, E. 441(1988)394
- Hundt, H.K.L., see Steyn, J.M. 432(1988)177
- Hung, C.T., Lim, J.K.C., Zoest, A.R. and Lam, F.C.  
Optimization of HPLC analysis for isoxazolyl penicillins using factorial design 425(1988)331
- Hunt, P., see Betteridge, R.F. 432(1988)352
- Hunter, B.T., see Gisch, D.J. 433(1988)264
- Hunter, K., Sharp, E.A. and Newton, A.  
Determination of diastereoisomers of bromadiolone, an anticoagulant rodenticide, in animal tissues by HPLC 435(1988)83
- Hunter, K. and Sharp, E.A.  
Modification to procedures for the determination of chlorophacinone and for multi-residue analysis of rodenticides in animal tissues 437(1988)301
- Hupe, K.P., see Kok, W.T. 436(1988)421
- Hurlbut, J.A., see Munns, R.K. 442(1988)209
- Hurowitz, L.A., see Cheng, T.-L. 425(1988)303
- Hurst, W.J., Snyder, K.P. and Martin, Jr., R.A.  
Determination of aflatoxins in peanut products using disposable bonded-phase columns and post-column reaction detection 409(1987)413
- Hurtubise, R.J., see Scolla, P.L. 405(1987)107
- Hussénius, A., Isaksson, R. and Matsson, O.  
Separation of alkyl-substituted indene enantiomers by chromatography on microcrystalline triacetylcellulose 405(1987)155
- Huszár, L., see Gazdag, M. 436(1988)31
- Hutt, A.J., see Avgerinos, A. 415(1987)75
- Hutter, N.H., see Van der Steege, G. 415(1987)1
- Huys, M., see Der Meeren, P.V. 447(1988)436
- Hwang, H.H., Healey, M.C. and Johnston, A.V.  
Purification of ascitic fluid-derived murine monoclonal antibodies by anion-exchange and size-exclusion HPLC 430(1988)329
- Hwang, S.S., see Musson, D.G. 414(1987)121
- Hyánek, J., see Verner, P. 416(1987)439
- Hyder, S.M. and Wittliff, J.L.  
High-performance hydrophobic interaction chromatography as a means of identifying estrogen receptors expressing different binding domains 444(1988)225
- Hyland, K. and Howells, D.W.  
Analysis and clinical significance of pterins (Review) 429(1988)95
- Hyman, K.M., see Mihaly, G.W. 415(1987)177
- Hyneck, M.L., Smith, P.C., Unseld, E. and Benet, L.Z.  
HPLC determination of tolmetin, tolmetin flucuronide and its isomeric conjugates in plasma and urine 420(1987)349
- Hynninen, P.H., see Sainio, E.-L. 420(1987)263

## I

- Iadarola, P., Zapponi, M.C., Stoppini, M., Meloni, M.L., Minchiotti, L., Galliano, M. and Ferri, G.  
HPLC of complex mixtures of cyanogen bromide-produced peptides from different proteins 443(1988)317
- Ibe, K., see Köppel, C. 427(1988)144

- Ibrahim, R.K., see Khouri, H.E. 407(1987)291  
 Ibrahim, S.S., see Boudinot, F.D. 430(1988)424  
 Ichikawa, N., see Katagiri, Y. 431(1988)135  
 Ichinose, H., see Naoi, M. 434(1988)209  
 Ickrath, O., see Ress, K. 417(1987)465  
 Ida, Y., see Okamoto, T. 430(1988)11  
 Idaka, E., Kaiya, H. and Horitsu, H.  
   Determination of prostaglandin E<sub>2</sub> in plasma  
   of schizophrenics and healthy controls by  
   GC-negative-ion chemical ionization MS and  
   radioimmunoassay 420(1987)373  
 Idle, J.R., see Hadidi, A.-H.F.A. 427(1988)121  
 Idle, J.R., see Schmid, B.J. 425(1988)107  
 Idvall, J., see Jonung, M. 415(1987)35  
 Igarashi, K., Kasuya, F., Mori, E. and Fukui, M.  
   Determination of dibucaine in biological  
   samples by GC with a nitrogen-phosphorus  
   detector 415(1987)407  
 Igarashi, K., see Kasuya, F. 416(1987)189  
 Igwe, O.J., Felice, L.J., Seybold, V.S. and Larson,  
 A.A.  
   Optimization of HPLC-radioimmunoassay  
   protocols for the analyses of substance P and  
   some of its metabolic fragments 432(1988)113  
 Igwemezie, L., see McErlane, K.M. 415(1987)335  
 Ihara, H., see Hirayama, C. 409(1987)175  
 Ihara, H., see Hirayama, C. 450(1988)271  
 Iida, T., Shinohara, T., Momose, T., Chang, F.C.,  
 Goto, J. and Nambara, T.  
   Separation of keto and keto-hydroxy bile acid  
   isomers by HPLC 438(1988)423  
 Iida, Y., see Hayashi, T. 428(1988)209  
 Ijiri, I., see Tomita, M. 414(1987)454  
 Ijitsu, T., Ueno, M. and Hara, S.  
   Two-dimensional HPLC system for the direct  
   determination of tocopherol derivatives in  
   plasma 427(1988)29  
 Ikai, Y., Oka, H., Kawamura, N., Yamada, M.,  
 Harada, K.-I. and Suzuki, M.  
   Improvement of chemical analysis of  
   antibiotics. XIII. Systematic simultaneous  
   analysis of residual tetracyclines in animal  
   tissues using TLC and HPLC 411(1987)313  
 Ikai, Y., see Oka, H. 411(1987)437  
 Ikai, Y., see Oka, H. 449(1988)448  
 Ikebuchi, J., Suenaga, K., Kotoku, S., Yuasa, I.  
 and Inagaki, O.  
   TLC with flame ionization detection for the  
   determination of tetrodotoxin in biological  
   fluids 432(1988)401  
 Ikeda, Y., see Fujii, Y. 448(1988)157  
 Ikeda, Y., see Okuyama, S. 431(1988)389  
 Ikenaka, K., see Hase, S. 434(1988)51  
 Ikenaka, T., see Hase, S. 434(1988)51  
 Ikenaka, T., see Omichi, K. 420(1987)158  
 Ikenaka, T., see Omichi, K. 428(1988)415  
 Ikezawa, H., see Koyama, I. 420(1987)275  
 Ilkőy, É., see Szederkényi, F. 446(1988)253  
 Imai, H., see Tamai, G. 423(1987)147  
 Imai, H., see Tamai, G. 423(1987)155  
 Imai, H., see Tamai, G. 423(1987)163  
 Imai, K., see Kotaniguchi, H. 420(1987)141  
 Imai, Y., Ito, S. and Fujita, K.  
   Determination of natural thiols by LC after  
   derivatization with 3,5-di-*tert*-butyl-1,2-  
   benzoquinone 420(1987)404  
 Imaizumi, K., see Nakano, K. 438(1988)61  
 Imamura, T., see Kudo, K. 431(1988)353  
 Imamura, Y., see Shibasaki, S. 415(1987)365  
 Imamura, Y., see Shibasaki, S. 421(1987)425  
 Imashuku, S., see Tsunamoto, K. 417(1987)414  
 Imbach, J.-L., see Maggio, A.-F. 436(1988)23  
 Inaba, T., see Nicoll-Griffith, D.A. 428(1988)103  
 Inacker, R., see Folena-Wasserman, G.  
 411(1987)345  
 Inagaki, K., see Oka, H. 411(1987)437  
 Inagaki, O., see Ikebuchi, J. 432(1988)401  
 Inchauspe, G., Delrieu, P., Dupin, P., Laurent,  
 M. and Samain, D.  
   Mechanism of selectivity in ion-pair HPLC  
   of aminoglycoside antibiotics using  
   perfluorinated pairing ions 404(1987)53  
 Ingalls, S.T., see Minkler, P.E. 413(1987)33  
 Ingalls, S.T., see Minkler, P.E. 420(1987)385  
 Ingen, H.E. Van, see Nielen, M.W.F.  
 417(1987)159  
 Ingen, H.E. Van and Endert, E.  
   Improved method for the determination of  
   the cortisol production rate using HPLC and  
   liquid scintillation counting 430(1988)233  
 Inman, E.L.  
   Determination of vancomycin related  
   substances by gradient HPLC 410(1987)363  
 Inman, E.L. and Rickard, E.C.  
   Chromatographic detection limits in  
   pharmaceutical method development  
   447(1988)1  
 Inotsume, N., Fujii, J., Honda, M., Nakano, M.,  
 Higashi, A. and Matsuda, I.  
   Stereoselective analysis of the enantiomers of  
   ethotoxin in human serum using chiral  
   stationary phase LC and GC-MS  
   428(1988)402  
 Inoue, H., see Oka, H. 411(1987)437  
 Inoue, T. and Suzuki, S.-I.  
   HPLC determination of triazolam and its  
   metabolites in human urine 422(1987)197  
 Ioffe, B.V., see Zenkevich, I.G. 439(1988)185  
 Iorio, M. Di, see Porcelli, G. 414(1987)423  
 Iri, H., see Matsumoto, K. 425(1988)323

- Irth, H., Brouwer, E., De Jong, G.J., Brinkman, U.A.T. and Frei, R.W.  
Trace enrichment and separation of As<sup>III</sup>, Sb<sup>III</sup> and Bi<sup>III</sup> as diethyldithiophosphate complexes by reversed-phase HPLC 439(1988)63
- Irth, H., De Jong, G.J., Brinkman, U.A.T. and Frei, R.W.  
Determination of disulfiram and two of its metabolites in urine by reversed-phase LC and spectrophotometric detection after post-column complexation 424(1988)95
- Irvine, G.B.  
High-performance SEC of polypeptides on a TSK G2000SW column in acidic mobile phases 404(1987)215
- Irwin, P.L., see Doner, L.W. 449(1988)229
- Isaksson, K.  
Evaluation of a chemical reaction detector with electrochemical production of bromine for the LC determination of physostigmine 411(1987)229
- Isaksson, K. and Kissinger, P.T.  
Metabolism of physostigmine in mouse liver microsomal incubations studied by LC with dual-electrode amperometric detection 419(1987)165
- Isaksson, R., see Hussénius, A. 405(1987)155
- Ise, N., see Kitano, H. 420(1987)13
- Ishi, K., see Fujimoto, Y. 402(1987)344
- Ishibashi, K., see Terada, T. 430(1988)388
- Ishibashi, M., see Matsuda, R. 438(1988)319
- Ishibashi, M., see Yamashita, K. 424(1988)1
- Ishida, J., Kai, M. and Ohkura, Y.  
Determination of oestrogens in pregnancy urine by HPLC with fluorescence detection 431(1988)249
- Ishida, J., see Kai, M. 430(1988)271
- Ishida, T., see Kumagai, Y. 421(1987)155
- Ishii, D., see Hirose, A. 411(1987)221
- Ishii, D., see Hirose, A. 438(1988)15
- Ishii, D., see Takeuchi, T. 402(1987)328
- Ishii, D., see Takeuchi, T. 405(1987)117
- Ishii, D., see Takeuchi, T. 407(1987)141
- Ishii, D., see Takeuchi, T. 407(1987)151
- Ishii, D., see Takeuchi, T. 435(1988)319
- Ishii, D., see Takeuchi, T. 435(1988)482
- Ishii, D., see Takeuchi, T. 445(1988)139
- Ishii, D., see Takeuchi, T. 447(1988)221
- Ishii, D., see Takeuchi, T. 403(1987)324
- Ishii, I., see Iwase, H. 416(1987)37
- Ishii, I., see Iwase, H. 417(1987)403
- Ishii, M. and Ogata, H.  
Determination of rifampicin and its main metabolites in human plasma by HPLC 426(1988)412
- Ishikawa, K., see Shibasaki, S. 415(1987)365
- Ishikawa, K., see Shibasaki, S. 421(1987)425
- Ishiwatari, R., see Morinaga, S. 403(1987)225
- Ishizaki, T., see Echizen, H. 426(1988)351
- Ishizaki, T., see Kubota, K. 425(1988)67
- Ishizaki, T., see Oshima, J. 414(1987)381
- Ishizaki, T., see Takahashi, H. 419(1987)243
- Iskrić, S., see Hadžija, O. 402(1987)358
- Ismail, Z., Aldous, S., Triggs, E.J., Smithurst, B.A. and Barry, H.D.  
GC analysis of Didronel tablets 404(1987)372
- Israili, Z.H., see Azeemuddin, S. 430(1988)163
- Issaq, H.J., see Mathes, L.E. 432(1988)346
- Itabashi, Y. and Takagi, T.  
HPLC separation of diacylglycerol enantiomers on a chiral stationary phase 402(1987)257
- Italia, M.P. and Uden, P.C.  
Multiple element emission spectral detection GC profiles of halogenated products from chlorination of humic acid and drinking water 438(1988)35
- Italia, M.P. and Uden, P.C.  
Comparison of volatile halogenated compounds formed in the chloramination and chlorination of humic acid by GC-electron-capture detection 449(1988)326
- Itami, T., Ema, M., Sakamoto, J. and Kawasaki, H.  
Microheterogeneity of rat  $\alpha$ -fetoprotein on a copper chelate column 417(1987)168
- Ito, S., see Imai, Y. 420(1987)404
- Ito, T., Matsuki, Y., Kurihara, H. and Nambara, T.  
Sensitive method for determination of captopril in biological fluids by GC-MS 417(1987)79
- Ito, Y.  
Foam counter-current chromatography with the cross-axis synchronous flow-through coil planet centrifuge 403(1987)77
- Ito, Y., see Zhang, T.-Y. 435(1988)159
- Ito, Y., see Zhang, T.-Y. 442(1988)455
- Ito, Y., see Zhang, T.-Y. 445(1988)199
- Ito, Y. and Zhang, T.-Y.  
Multi-stage mixer-settler planet centrifuge. Preliminary studies on partition of macromolecules with organic-aqueous and aqueous-aqueous two-phase solvent systems 437(1988)121
- Ito, Y. and Zhang, T.-Y.  
Cross-axis synchronous flow-through coil planet centrifuge for large-scale preparative

- counter-current chromatography. I. Apparatus and studies on stationary phase retention in short coils 449(1988)135
- Ito, Y. and Zhang, T.-Y. Cross-axis synchronous flow-through coil planet centrifuge for large-scale preparative counter-current chromatography. II. Studies on partition efficiency in short coils and preparative separations with multilayer coils 449(1988)153
- Itoh, R., see Echizen, H. 426(1988)351
- Itoh, T., see Shibasaki, S. 421(1987)425
- Itokawa, Y., see Hiraike, H. 430(1988)143
- Ittensohn, A., see De Vries, J.X. 417(1987)420
- Ivanchev, S.S., see Shadrina, N.E. 404(1987)183
- Ivanichkova, S., see Lekova, K. 446(1988)31
- Ivanov, S., see Gavrilova, T.B. 402(1987)87
- Ivashkiv, E., see Cohen, A.I. 416(1987)445
- Ivashkiv, E., see Jemal, M. 428(1988)81
- Iversen, P., see Raha, C. 450(1988)325
- Ivin, P.C. and Clarke, M.L. Isolation of kestoses and nystose from enzyme digests by HPLC 408(1987)393
- Iwabuchi, H., Nakagawa, A. and Nakamura, K.-I. Application of molecular-secondary-ion MS for drug metabolism studies. I. Direct analysis of conjugates by GLC-secondary-ion mass spectrometry 414(1987)139
- Iwaguchi, K., see Tanaka, N. 448(1988)95
- Iwai, K., see Shibata, K. 417(1987)173
- Iwai, K., see Shibata, K. 422(1987)257
- Iwai, K., see Shibata, K. 424(1988)23
- Iwai, K., see Shibata, K. 430(1988)381
- Iwaki, K., Nimura, N., Hiraga, Y., Kinoshita, T., Takeda, K. and Ogura, H. Amino acid analysis by reversed-phase HPLC. Automatic pre-column derivatization with activated carbamate reagent 407(1987)273
- Iwaki, K., see Nimura, N. 402(1987)387
- Iwaki, K., Yoshida, S., Nimura, N., Kinoshita, T., Takeda, K. and Ogura, H. Optical resolution of enantiomeric amino acid derivatives on a naphthylethylurea multiple-bonded chiral stationary phase prepared via an activated carbamate intermediate 404(1987)117
- Iwamoto, K., see Katagiri, Y. 431(1988)135
- Iwamoto, T., see Ohmori, S. 431(1988)239
- Iwamura, S. and Tsukamoto, K. Highly sensitive determination of 2-hydroxy-N-(3-{*m*-(1-piperidinylmethyl)phenoxy}propyl)-acetamide (desacetyl TZU-0460) in rat plasma by HPLC 413(1987)370
- Iwase, H., Ishii, I., Hamazaki, H. and Hotta, K. Detection of uridine 5'-diphosphate glucose-collagen glucosyltransferase activity by HPLC and possible participation of a novel intermediate as a glucose donor in collagen biosynthesis 417(1987)403
- Iwase, H., Ishii, I., Kato, Y., Hamazaki, H., Hotta, K., Umezawa, A. and Kanzaki, T. Subfractionation of the dansylated derivatives of glucosyl galactosyl hydroxylysine by LC and its application to a specific  $\alpha$ -1,2-glucosidase assay 416(1987)37
- Iwase, S., see Honda, S. 404(1987)313
- Iwase, S.F.S. and Honda, S. Analysis of water-soluble vitamins by micellar electrokinetic capillary chromatography 447(1988)133
- Iwata, H., Matsuda, T. and Tonomura, H. Improved HPLC determination of thiamine and its phosphate esters in animal tissues 450(1988)317
- Iwata, T., Yamaguchi, M. and Nakamura, M. Highly sensitive and simple determination of cholesterol and cholestanol in human serum by HPLC with fluorescence detection 421(1987)43
- Iyengar, J.R., see Lawrence, J.F. 404(1987)261
- Izquierdo, C., see Usero, J.L. 407(1987)355

## J

- Jabs, H.-U. and Assmann, G. Characterization of an apolipoprotein C-III mutant by HPLC and time-of-flight secondary ion MS 414(1987)323
- Jackson, L.S. and Stafford, J.E.H. HPLC assay for the simultaneous measurement of two spironolactone analogues, 17 $\beta$ -hydroxy-3-oxo-17-pregn-4-ene-21-carboxylic acid- $\gamma$ -lactone 7 $\alpha$ -methyl ethyl ester and its 6 $\beta$ -hydroxylated derivative 428(1988)377
- Jackson, P.E., see Haddad, P.R. 407(1987)121
- Jackson, P.E., see Haddad, P.R. 447(1988)155
- Jackson, P.E. and Haddad, P.R. Studies on sample preconcentration in ion chromatography. VII. Review of the methodology and applications of anion preconcentration 439(1988)37

- Jacob, K., Kossien, I., Egeler, E. and Knedel, M.  
Detection of non-typical porphyrin isomers  
in human urines by ion-pair reversed-phase  
HPLC 441(1988)171
- Jacob, III, P., Elias-Baker, B.A., Jones, R.T. and  
Benowitz, N.L.  
Determination of benzoylecgonine and  
cocaine in biologic fluids by automated GC  
417(1987)277
- Jacob III, P., see Shulgin, A.T. 423(1987)365
- Jacobs, J.S.A., Christman, R.F. and Johnson,  
J.D.  
High-pressure SEC of chlorinated and  
unchlorinated aquatic fulvic acids using a  
polyacrylamide gel 450(1988)433
- Jacobsen, D., see Øvrebø, S. 416(1987)111
- Jacobson, K.A., see Gusovsky, F. 415(1987)124
- Jacobson, K.B., see Yim, J.J. 419(1987)296
- Jacobsson, S., Larsson, A., Arbin, A. and  
Hagman, A.  
Extractive pentafluorobenzoylation of formic,  
acetic, levulinic, benzoic and phthalic acids,  
studied by LC and dual-oven capillary GC  
447(1988)329
- Jacobsson, S., see Hagman, A. 448(1988)117
- Jadaud, P., Caude, M. and Rosset, R.  
Main parameters governing the retention of  
polynuclear aromatic hydrocarbons on  
tetrachlorophthalimidopropyl-bonded silica  
439(1988)195
- Jacob, E., see Ong, C.N. 423(1987)278
- Jaeger, H., see Nieder, M. 413(1987)207
- Jaeger, H., see Nieder, M. 414(1987)492
- Jaeger, H., see Nieder, M. 424(1988)73
- Jaffe, N., see Anzai, T. 415(1987)445
- Jaillon, P., see Garnier-Moiroux, A. 416(1987)183
- Jaillon, P., see Poirier, J.-M. 426(1988)431
- Jain, R. and Bhatia, A.  
TLC separation of some sulpha drugs using  
acetoacetanilide as a coupling agent  
441(1988)454
- Jakobsen, P., Grønhej Larsen, F. and  
Grønhej Larsen, C.  
Simultaneous determination of the aromatic  
retinoids etretin and etretinate and their main  
metabolites by reversed-phase liquid  
chromatography 415(1987)413
- Jamali, F., see Foster, R.T. 416(1987)388
- Jamali, F., see Mehvar, R. 425(1988)135
- Jamali, F., see Mehvar, R. 431(1988)
- James, C.A., see Burton, L.C. 431(1988)450
- James, C.A., see Dobbs, N.A. 420(1987)184
- James, L.B.  
Amino acid analysis: the histidine hangover  
403(1987)271
- James, L.B.  
Amino acid analysis. I. Nuances of integration  
404(1987)321
- James, L.B.  
Amino acid analysis: a fall-off in performance  
408(1987)291
- James, L.B.  
Amino acid analysis. II. Nuances of  
integration 410(1987)176
- James, L.B.  
Amino acid analysis: buffers and artifacts  
436(1988)80
- James, L.B.  
Amino acid analysis. III. Nuances of  
integration 436(1988)474
- Janák, J., see Soják, L. 406(1987)43
- Janča, J.  
Flow velocity profile in channels with  
modulated cross-sectional permeability for  
sedimentation-flotation focusing field-flow  
fractionation using step density gradients  
404(1987)23
- Jandera, P.  
Methods for characterization of selectivity in  
reversed-phase LC. IV. Retention behaviour  
of oligomeric series 449(1988)361
- Jandke, J. and Spittler, G.  
Unusual conjugates in biological profiles  
originating from consumption of onions and  
garlic 421(1987)1
- Janeček, M., Kahle, V. and Krejčí, M.  
UV fibre optic photometric cell with an  
effective volume of 30 nl for micro-scale  
HPLC 438(1988)409
- Janicot, J.L., Caude, M. and Rosset, R.  
Separation of opium alkaloids by carbon  
dioxide sub- and supercritical fluid  
chromatography with packed columns.  
Application to the quantitative analysis of  
poppy straw extracts 437(1988)351
- Janis, L., see Fausnaugh-Pollitt, J. 443(1988)221
- Janis, L.J. and Regnier, F.E.  
Immunological-chromatographic analysis  
444(1988)1
- Janjić, T.J., Tešić, Ž.L., Vučković, G.N. and  
Čelap, M.B.  
Effect of the geometrical configuration of  
facial and meridional isomers of cobalt(III)  
and chromium(III) complexes on their  $R_F$   
values 404(1987)307
- Janoušek, V., see Sanitřák, J. 415(1987)129
- Janoušek, V., see Tvrzická, E. 431(1988)231
- Janowitz, P., see Swobodnik, W. 423(1987)75

- Jansen, E.H.J.M., Van den Berg, R.H.,  
Enkelaar-Willemsen, C. and Zomer, G.  
HPLC of chemiluminogenic labels with  
sensitive photoplate detection 437(1988)268
- Jansen, E.H.J.M. and Both-Miedema, R.  
Instability of phenylthiohydantoin amino  
acids 435(1988)363
- Jansen, G., see Hoving, E.B. 434(1988)395
- Jansen, H., Brinkman, U.A.T. and Frei, R.W.  
Stereoselective determination of L-amino  
acids using column LC with an enzymatic  
solid-phase reactor and chemiluminescence  
detection 440(1988)217
- Jansen, H., see Broersen, J.J. 436(1988)39
- Janssen, G., see Cachet, T. 445(1988)290
- Janssen, L.H.M., see Bos, O.J.M. 424(1988)13
- Janssen, T.J., see Botterblom, M.H.A.  
421(1987)211
- Japp, M., Garthwaite, K., Geeson, A.V. and  
Osselton, M.D.  
Collection of analytical data for  
benzodiazepines and benzophenones  
439(1988)317
- Jardy, A., Caude, M., Diop, A., Curvale, C. and  
Rosset, R.  
Single-column anion chromatography with  
indirect UV detection using pyromellitate  
buffers as eluents 439(1988)137
- Jarrett, H.W.  
Development of N-hydroxysuccinimide ester  
silica, a novel support for high-performance  
affinity chromatography 405(1987)179
- Jarrett, H.W., see Foster, W.S. 403(1987)99
- Jarry, C., see Demotes-Mainard, F. 419(1987)388
- Jatoo, S.D., Kajbaf, M. and Gorrod, J.W.  
HPLC with on-line radioactivity monitoring  
for metabolic studies of 2- and 4-  
[<sup>14</sup>C]aminobiphenyls 442(1988)394
- Jaureguiberry, G., Ogunkolade, W., Bailly, E.,  
Rhodes-Feuillette, A., Agrapart, M. and  
Ballet, J.-J.  
*Plasmodium falciparum* exoprotein  
stimulation of human T-lymphocytes  
unsensitized to malaria 440(1988)385
- Jaussaud, P., Courtot, D., Guyot, J.L. and Paris,  
J.  
Identification of a flunixin metabolite in the  
horse by GC-MS 423(1987)123
- Jbilou, M., see Bougerolle, A.M. 434(1988)232
- Jean, C. and Laplanche, R.  
Assay of isradipine and of its major  
metabolites in biological fluids by capillary  
GC and chemical ionization MS 428(1988)61
- Jegorov, A., Tříska, J., Trnka, T. and Černý, M.  
Separation of  $\alpha$ -amino acid enantiomers by  
reversed-phase HPLC after derivatization  
with *o*-phthaldialdehyde and a sodium salt of  
1-thio- $\beta$ -D-glucose 434(1988)417
- Jehl, F., Birckel, P. and Monteil, H.  
Hospital routine analysis of penicillins,  
third-generation cephalosporins and  
aztreonam by conventional and high-speed  
HPLC 413(1987)109
- Jehl, F., see Koechlin, C. 425(1988)197
- Jehličková, Z., see Hajšlová, J. 438(1988)55
- Jelínek, I., Dohnal, J., Snopek, J. and Smolková-  
Keulemansová, E.  
Use of cyclodextrins in isotachopheresis. III.  
Purity control of naftidrofuryl  
hydrogenoxalate and some of its synthesis  
intermediates 435(1988)496
- Jelínek, I., see Snopek, J. 411(1987)153
- Jelínek, I., see Snopek, J. 438(1988)211
- Jelínek, I., see Snopek, J. 450(1988)373
- Jelínek, I., Snopek, J. and Smolková-  
Keulemansová, E.  
Use of cyclodextrins in isotachopheresis. I.  
Effect of cyclodextrin on the isotachopheretic  
separation of related penicillins 405(1987)379
- Jelínek, I., Snopek, J. and Smolková-  
Keulemansová, E.  
Use of cyclodextrins in isotachopheresis. V.  
The separation of ketotifen and its polar  
intermediate enantiomers 439(1988)386
- Jemal, M., Ivashkiv, E., Teitz, D. and Cohen,  
A.I.  
Simultaneous determination of the prodrug  
zofenopril and its active drug in plasma by  
capillary GC-MS 428(1988)81
- Jemal, M., see Cohen, A.I. 416(1987)445
- Jemal, M., see Wang-Iverson, D. 408(1987)365
- Jemal, M. and Cohen, A.I.  
Simultaneous determination of the acid  
chloride S-benzoyl-3-mercapto-2-  
methylpropanoyl chloride and residual  
carboxylic acid by capillary GC 408(1987)360
- Jen, J.-F., Daugherty, K.E. and Tarter, J.G.  
Use of first derivative conductometric  
detection in ion chromatography 436(1988)86
- Jenke, D.R.  
Quantitation of oxalate and citrate by ion  
chromatography with a buffered, strong acid  
eluent 437(1988)231
- Jenke, D.R., see Brown, D.S. 410(1987)157
- Jenkins, I.D., see Goren, M.B. 440(1988)487
- Jenkins, R.L., McDaniel, H.G., Grizzle, W.,  
Parrish, S.W. and McDaniel, H.B.  
Demonstration of the adenosine reservoirs  
with nitrobenzylthioinosine in liver and  
diaphragm by HPLC 426(1988)249



- Jenner, C.F., see Ugalde, T.D. 449(1988)207
- Jenner, P.J. and Ellard, G.A.  
Determination of the isoniazid metabolite monoacetylhydrazine in urine by HPLC 415(1987)188
- Jenny, H.B. and Peter, H.H.  
Determination of desferrioxamine B, its complexes with iron and aluminium and selected derivatives by HPLC using a polymer column 438(1988)433
- Jeremic, B., see Dahmen, W. 427(1988)157
- Jesus, A.E. De, Gorst-Allman, C.P., Horak, R.M. and Vlegaar, R.  
Large-scale purification of the mycotoxins aflatoxin B<sub>1</sub>, B<sub>2</sub> and G<sub>1</sub> 450(1988)101
- Jham, G.N. and Campos, L.G.  
Use of chloroform as a solvent for the formation of N-trifluoroacetyl-*n*-butylamino acid derivatives for GC analysis 436(1988)100
- Ji, H. and Wang, E.  
Thin-layer electrochemical detector coated with Nafion film for LC 410(1987)111
- Jiao, B., see Galanos, C. 440(1988)397
- Jickells, S.M., see Castle, L. 437(1988)281
- Jimenez, J., see Sonesson, A. 417(1987)366
- Jimenez, M., see Mateo, R. 410(1987)319
- Jimeno, M.L., see García-Raso, A. 402(1987)323
- Jindal, S.P., Lutz, T. and Bagchi, S.P.  
GLC-MS determination of 1-methyl-4-phenyl-4-propionoxypiperidine using a stable isotope-labeled analogue as an internal standard 408(1987)356
- Jinno, K., Anazawa, N., Okamoto, M., Tanaka, N., Fetzer, J.C. and Biggs, W.R.  
Elution behaviour of peropyrene-type polycyclic aromatic hydrocarbons in LC 402(1987)173
- Jinno, K., Kuwajima, M., Hayashida, M., Watanabe, T. and Hondo, T.  
Automated identification of toxic substances in poisoned human fluids by a retention prediction system in reversed-phase LC 436(1988)11
- Jinno, K., Nagoshi, T., Tanaka, N., Okamoto, M., Fetzer, J.C. and Biggs, W.R.  
Effect of column temperature on the retention of peropyrene-type polycyclic aromatic hydrocarbons on various chemically bonded stationary phases in reversed-phase LC 436(1988)1
- Jinno, K., see Fujimoto, C. 411(1987)213
- Jinno, K., see Fujimoto, C. 438(1988)329
- Jinno, K., see Okamoto, M. 407(1987)314
- Jinno, K., see Wahyuni, W.T. 448(1988)398
- Jizba, J., see Beran, M. 442(1988)431
- Joel, S.P., Osborne, R.J. and Slevin, M.L.  
Improved method for the simultaneous determination of morphine and its principal glucuronide metabolites 430(1988)394
- Joelsson, M., see Johansson, G. 411(1987)161
- Johannesen, R.B., see Eng, G. 403(1987)1
- Johannesen, R.B., see Tierney, E.J. 441(1988)229
- Johansson, B.-L., Hellberg, U. and Wennberg, O.  
Determination of the leakage from Phenyl-Sepharose CL-4B, Phenyl-Sepharose FF and Phenyl-Superose in bulk and column experiments 403(1987)85
- Johansson, G. and Joelsson, M.  
Effect of polymer structure on affinity partitioning of lactate dehydrogenase in polymer-water two-phase systems 411(1987)161
- Johansson, K., see Wester, P. 415(1987)261
- Johansson, M. and Anlér, E.-L.  
GC analysis of flunixin in equine urine after extractive methylation 427(1988)55
- Johansson, M. and Forsmo-Bruce, H.  
Determination of atenolol in plasma by dual-column LC and fluorimetric detection 432(1988)265
- Johansson, M. and Lenngren, S.  
Determination of cysteine, glutathione and N-acetylcysteine in plasma by ion-pair reversed-phase LC and post-column derivatization 432(1988)65
- Johansson, M. and Lenngren, S.  
Determination of bromhexine in plasma by reversed-phase LC. Interference of lipoproteins on extraction 432(1988)243
- John, J., see Robinson, J.L. 402(1987)201
- Johno, I., see Suzuki, K. 425(1988)435
- Johnsen, P.B. and Kuan, J.-C.W.  
Simplified method to quantify geosmin and 2-methylisoborneol concentrations in water and microbiological cultures 409(1987)337
- Johnsen, S. and Kolset, K.  
The mass-selective detector as a chlorine-selective detector 438(1988)233
- Johnson, B.C.  
The vitamin K dependent reaction 440(1988)499
- Johnson, B.F., see Nugent, K.D. 443(1988)381
- Johnson, C.A., see Vanscoik, K.G. 417(1987)183
- Johnson, J.D., see Jacobs, J.S.A. 450(1988)433
- Johnston, A.V., see Hwang, H.H. 430(1988)329
- Jokuszies, G., see Štulík, K. 436(1988)334
- Jol, C., see Tjaden, U.R. 420(1987)53
- Jolley, K.W., see Burrows, J.L. 432(1988)199
- Joly, J.P., see Moll, N. 405(1987)347
- Jonas, G., see Whelpton, R. 426(1988)223

- Jones, A.C., see Baugh, P.J. 411(1987)445
- Jones, A.D., see Smidt, C.R. 434(1988)21
- Jones, C.S. and Grimsrud, E.P.  
MS measurements of the negative ions formed in the thermionic ionization detector with nitrogen carrier gas 409(1987)139
- Jones, K. and Malcolme-Lawes, D.J.  
High-performance UV absorption detector for LC. II. A prototype detector 441(1988)387
- Jones, M.D., see Robinson, P.R. 432(1988)153
- Jones, M.W., see Baugh, P.J. 411(1987)445
- Jones, R.M., see Prasad, M.R. 428(1988)221
- Jones, R.T., see Jacob, III, P. 417(1987)277
- Jones, R.W., see Notarianni, L.J. 431(1988)461
- Jong, A.P. De, Wittebrood, A.J., Du Châtinier, W.M. and Bron, J.  
LC analysis of alizapride and metoclopramide in human plasma and urine using solid-phase extraction 419(1987)233
- Jong, G.J. De, see Irth, H. 424(1988)95
- Jong, G.J. De, see Irth, H. 439(1988)63
- Jong, G.J. De, see Maris, F.A. 435(1988)297
- Jong, G.J. De, see Maris, F.A. 445(1988)129
- Jong, J. De, Point, A.J.F., Tjaden, U.R., Beeksmas, S. and Kraak, J.C.  
Determination of catecholamines in urine (and plasma) by LC after on-line sample pretreatment on small alumina or dihydroxyborylsilica columns 414(1987)285
- Jong, J. De, Tjaden, U.R., Visser, E. and Meijer, W.H.  
Determination of serotonin and 5-hydroxyindoleacetic acid in urine by reversed-phase ion-pair partition chromatography with fluorimetric detection 419(1987)85
- Jong, P. De, see Odink, J. 431(1988)309
- Jongejan, P.A.C., see Keuken, M.P. 439(1988)13
- Jongeneelen, F.J., Anzion, R.B.M. and Henderson, P.T.  
Determination of hydroxylated metabolites of polycyclic aromatic hydrocarbons in urine 413(1987)227
- Jonker, H.D., see Lemmens, A.A.G. 439(1988)423
- Jonker, H.H., see Van Gelder, W.M.J. 442(1988)133
- Jönsson, J.Å. and Lövkvist, P.  
Determination of adsorption isotherms from chromatographic measurements, using the peak maxima method 408(1987)1
- Jonung, M., Björkman, S. and Idvall, J.  
Determination of methylprednisolone in rat tissue by HPLC 415(1987)35
- Jordan, J.-C., see Heizmann, P. 415(1987)170
- Jordan, R.A., see Wright, D.S. 417(1987)223
- Jorgenson, J.W., see De Wit, J.S.M. 411(1987)201
- Jorgenson, J.W., see Rose, D.J. 438(1988)23
- Jorgenson, J.W., see Rose, Jr., D.J. 447(1988)117
- Jork, H. and Kunz, F.-R.  
Isolation of gentamicin C compounds from culture filtrates of *Micromonospora purpurea* 450(1988)115
- Josephson, M., see Grigorova, B. 410(1987)419
- Joshi, G.P., see Kharbade, B.V. 439(1988)430
- Josić, D., Hofmann, W., Habermann, R. and Reutter, W.  
High-performance concanavalin A affinity chromatography of liver and hepatoma membrane proteins 444(1988)29
- Joulin, Y., Moretti, J.L. and Diquet, B.  
Highly sensitive method for determination of isopropyl-*p*-iodoamphetamine by GC with electron-capture detection 425(1988)402
- Joyce, D.A. and Wade, D.N.  
Assay for D-penicillamine-proton conjugate in human plasma utilising chemical reduction followed by HPLC with gold/mercury electrochemical detection 430(1988)319
- Józan, M., see Papp, O. 403(1987)11
- Jozefonvicz, J., see Santarelli, X. 443(1988)55
- Juffmann, F., see Gübitz, G. 404(1987)391
- Julkunen-Tiitto, R., see Meier, B. 442(1988)175
- Jullien, I., see Delee, E. 450(1988)183
- Jullien, I., see Delee, E. 450(1988)191
- Jumar, A., see Klemm, D. 438(1988)122
- Jung, L., see Efthymiopoulos, C. 421(1987)360
- Jung, S.S. and Wittliff, J.L.  
Influence of column length and pore size on high-performance ion-exchange chromatography of estrogen and progesterin receptors 425(1988)293
- Junginger, H.E., see Roorda, W.E. 403(1987)355
- Junk, G.A., see Avery, M.J. 420(1987)379
- Junker, A., see Sturm, E. 430(1988)43
- Jurková, M. and Wurst, M.  
Chromatography of microbial metabolites of aromatic amino acids 446(1988)117
- Jusko, W.J., see Haughey, D.B. 430(1988)241
- Jusko, W.J., see Kong, A.-N. 432(1988)308
- Jutisz, M., see Starzec, A. 440(1988)353
- Jüttner, F.  
A cryotrap technique for the quantitation of monoterpenes in humid and ozone-rich forest air 442(1988)157
- Jüttner, F., see Henatsch, J.J. 445(1988)97
- Juven, B.J., see Harduf, Z. 437(1988)379

# K

- Kabra, P.M.  
Clinical analysis of individual steroids by column LC (Review) 429(1988)155
- Kabra, P.M., see Gotelli, G.R. 428(1988)147
- Kadofuku, T. and Sato, T.  
Detection of the changes in serum proteins during chemical hepatocarcinogenesis by two-dimensional electrophoresis under non-denaturing conditions 419(1987)315
- Kagawa, M., see Matsubara, K. 424(1988)49
- Kågedal, B., Andersson, T., Carlsson, M., Denneberg, T. and Hoppe, A.  
HPLC of 2-mercaptopyruvate and its metabolite 2-mercaptopyruvate in plasma and urine after treatment with thiopronine 417(1987)261
- Kågedal, B., see Hansson, C. 420(1987)146
- Kågedal, B. and Goldstein, D.S.  
Catecholamines and their metabolites (Review) 429(1988)177
- Kahl, G.F., see Michaelis, H.C. 416(1987)176
- Kahl, H.-G., see Schmitz, G. 431(1988)327
- Kahle, V., see Janeček, M. 438(1988)409
- Kahle, V., see Škabrahová, Z. 408(1987)372
- Kahn, T.P.  
Simple instrument for teaching the basics of chromatography in schools 446(1988)173
- Kai, M., Ishida, J. and Ohkura, Y.  
HPLC determination of leucine-enkephalin-like peptide in rat brain by pre-column fluorescence derivatization involving formylation followed by reaction with 1,2-diamino-4,5-dimethoxybenzene 430(1988)271
- Kai, M., see Chao, W.-F. 430(1988)361
- Kai, M., see Ishida, J. 431(1988)249
- Kai, M., see Nakano, M. 411(1987)305
- Kai, M., see Ohno, M. 421(1987)245
- Kai, M., see Ohno, M. 430(1988)291
- Kaiser, D.G., Vangiessen, G.J., Shah, J.A. and Weber, D.J.  
GC determination of guanadrel in plasma and urine 434(1988)135
- Kaiser, G., Ackermann, R., Dieterle, W. and Dubois, J.-P.  
Determination of a new angiotensin converting enzyme inhibitor and its active metabolite in plasma and urine by GC-MS 419(1987)123
- Kaiser, G., see Sioufi, A. 434(1988)239
- Kaiser, G., see Sioufi, A. 450(1988)221
- Kaiser, I., see Birk, I.M. 449(1988)423
- Kaiya, H., see Idaka, E. 420(1987)373
- Kajbaf, M., see Jatoo, S.D. 442(1988)394
- Kajiwara, M., see Kurumaya, K. 435(1988)235
- Takehi, K., see Honda, S. 419(1987)51
- Takehi, K., see Honda, S. 438(1988)73
- Kakihana, H., see Hosoe, M. 435(1988)253
- Kakihana, H., see Hosoe, M. 438(1988)225
- Kakihana, H., see Oi, T. 450(1988)343
- Kakimoto, T., see Fujimoto, Y. 402(1987)344
- Kalabina, L.L., see Shmidel, E.B. 446(1988)163
- Kale, N.R., see Talib, Y.Y. 403(1987)373
- Kalghatgi, K. and Horváth, C.  
Rapid peptide mapping by HPLC 443(1988)343
- Kalin, J.R. and Hill, D.L.  
Determination of 2',3'-dideoxyadenosine, 2',3'-dideoxyinosine and 2',3'-dideoxycytidine in biological samples 431(1988)184
- Källberg, M., see Hansson, C. 420(1987)146
- Kallos, G.J., see Cutié, S.S. 408(1987)349
- Kalow, W., see Kruck, T.P.A. 433(1988)207
- Kalow, W., see Nicoll-Griffith, D.A. 428(1988)103
- Kaluzny, B.D. and Bannow, C.A.  
HPLC determination of pimeprofen and its metabolite ibuprofen in sheep plasma and lymph 414(1987)228
- Kamata, S., see Yonekura, T. 427(1988)320
- Kamei, M., see Harada, H. 407(1987)299
- Kamino, M., see Tanaka, M. 438(1988)253
- Kaminski, M. and Reusch, J.F.  
Some problems associated with the selection of separation conditions in preparative LC 436(1988)367
- Kamlet, M.J., see Abraham, M.H. 409(1987)15
- Kamma, D.A., see Baumann, R.A. 439(1988)165
- Kamohara, K., see Yoshikawa, S. 421(1987)111
- Kamoshita, K., see Takahashi, J. 436(1988)316
- Kamps, M.A.F., see Engels, W. 427(1988)209
- Kanamori, N., see Yamaji, A. 415(1987)143
- Kanamori, N., see Yamaji, A. 421(1987)372
- Kanazawa, T., see Seki, T. 424(1988)410
- Kandybin, N.V., see Beran, M. 442(1988)431
- Kaneda, N., see Sasaoka, T. 424(1988)392
- Kaneda, N., see Sasaoka, T. 428(1988)152
- Kanemitsu, F. and Okigaki, T.  
Creatine kinase isoenzymes (Review) 429(1988)399
- Kaneshima, H., see Kanetoshi, A. 442(1988)289
- Kaneta, T., see Tanaka, S. 447(1988)383
- Kanetoshi, A., Ogawa, H., Katsura, E., Kaneshima, H. and Miura, T.  
Formation of polychlorinated dibenzo-*p*-

- dioxins upon combustion of commercial textile products containing 2,4,4'-trichloro-2'-hydroxydiphenyl ether (Irgasan DP300) 442(1988)289
- Kanfer, I., see Stubbs, C. 427(1988)93
- Kaniwa, M.-A.  
HPLC identification and determination of dithiocarbamates in vulcanized rubber products 405(1987)263
- Kano, K., see Kawai, S. 442(1988)451
- Kanzaki, T., see Iwase, H. 416(1987)37
- Kapel, J. Van, see Lindemans, J. 432(1988)103
- Kapetanovic, I.M., DiGiovanni, J.H., Bartosevich, J.F., Melendez, V., Schroeder, A.C., Von Bredow, J. and Heiffer, M.H. Analysis of DL-threo- $\alpha$ -(2-piperidyl)-2-trifluoromethyl-6-(4-trifluoromethylphenyl)-4-pyridinemethanol phosphate (enpiroline), a new candidate antimalarial, in blood using HPLC 419(1987)458
- Kapetanovic, I.M., see Torchin, C.D. 426(1988)444
- Kapetanovic, I.M., Yonekawa, W.D. and Kupferberg, H.J.  
Determination of 4-aminobutyric acid, aspartate, glutamate and glutamine and their  $^{13}\text{C}$  stable-isotopic enrichment in brain tissue by GC-MS 414(1987)265
- Kaplan, B. and Pras, M.  
Removal of sodium dodecyl sulphate from proteins by GPC 423(1987)376
- Kaplan, E.D., see Siezen, R.J. 444(1988)239
- Kaplan, L., see Metz, P.A. 441(1988)31
- Kaplinsky, L.B., see Prasad, M.R. 428(1988)221
- Karamanos, N.K., Tseggenidis, T. and Antonopoulos, C.A.  
Analysis of neutral sugars as dinitrophenyl-hydrazones by HPLC 405(1987)221
- Karger, B.L., see Guttman, A. 448(1988)41
- Karger, B.L., see Shieh, C.H. 406(1987)343
- Karhu, I., Mahonen, A. and Palvimo, J.  
Separation and quantification of histone H1 subtypes and high-mobility-group proteins by reversed-phase LC: protein levels in rat tissues during postnatal development 426(1988)65
- Karlaganis, G., Peretti, E. and Lauterburg, B.H. Analysis of isoniazid, acetylhydrazine and [ $^{15}\text{N}_2$ ]acetylhydrazine in serum by capillary GC-ammonia chemical ionization MS 420(1987)171
- Karle, J.M. and Olmeda, R.  
Rapid and sensitive quantitative analysis of the new antimalarial N<sup>4</sup>-[2,6-dimethoxy-4-methyl-5-[(3-trifluoromethyl)phenoxy]-8-quinolinyl]-1,4-pentanediamine in plasma by LC and electrochemical detection 424(1988)347
- Karlson, A., see Pettersson, C. 407(1987)217
- Karlsson, L.  
Direct injection of urine on an HPLC column-switching system for determination of  $\Delta^9$ -tetrahydrocannabinol-11-oic acid with both UV and electrochemical detection 417(1987)309
- Karlsson, S., Banhidi, Z.G. and Albertsson, A.-C. Detection by HPLC of polyamines formed by clostridal putrefaction of caseins 442(1988)267
- Karmen, A., see Lam, S. 441(1988)81
- Karnes, H.T., Beightol, L.A., Serafin, R.J. and Farthing, D.  
Improved method for the determination of diazepam and N-desmethyldiazepam in plasma using capillary GC and nitrogen-phosphorus detection 424(1988)398
- Karnes, H.T., Opong-Mensah, K., Farthing, D. and Beightol, L.A.  
Automated solid-phase extraction and HPLC determination of ranitidine from urine, plasma and peritoneal dialysate 422(1987)165
- Karnes, H.T., see Sarkar, M.A. 434(1988)324
- Karol, P.J.  
Chromatogram statistical moments from columns of finite length 445(1988)207
- Karr, L.J., Van Alstine, J.M., Snyder, R.S., Shafer, S.G. and Harris, J.M.  
Cell separation by immunoaffinity partitioning with polyethylene glycol-modified protein A in aqueous polymer two-phase systems 442(1988)219
- Karve, M.S., see Talib, Y.Y. 403(1987)373
- Káš, J., see Rauch, P. 438(1988)451
- Kasafirek, E., see Kocna, P. 434(1988)429
- Kasai, M., see Hirata, N. 434(1988)71
- Kasai, M., see Sawada, S. 402(1987)283
- Kasai, R., Yamaguchi, H. and Tanaka, O.  
HPLC of glycosides on a new type of hydroxyapatite column 407(1987)205
- Kaspersen, F.M., see Paanakker, J.E. 421(1987)327
- Kastl, P.R., see Howard, II, R.R. 414(1987)434
- Kasuya, F., Igarashi, K. and Fukui, M.  
Determination of six metabolites of oxybuprocaine in urine by HPLC 416(1987)189
- Kasuya, F., see Igarashi, K. 415(1987)407
- Kasuya, Y., see Furuta, T. 413(1987)1
- Kasuya, Y., see Hirota, N. 425(1988)237
- Kasziba, E., Flancbaum, L., Fitzpatrick, J.C., Schneiderman, J. and Fisher, H.

- Simultaneous determination of histidine-containing dipeptides, histamine, methylhistamine and histidine by HPLC 432(1988)315
- Katae, H., see Kojima, K. 413(1987)326
- Katagiri, Y., Naora, K., Ichikawa, N., Hayashibara, M. and Iwamoto, K. Simultaneous determination of ofloxacin, fenbufen and felbinac in rat plasma by HPLC 431(1988)135
- Kataoka, H., Ohishi, K., Sumida, Y., Ohmori, M. and Makita, M. GC assay for cysteine sulphinate decarboxylase activity in animal tissues 420(1987)135
- Kataoka, H., Sakiyama, N. and Makita, M. GC analysis of aminoalkylphosphonic acids and aminoalkyl phosphates 436(1988)67
- Kataoka, K., see Yamaji, A. 415(1987)143
- Kataoka, K., see Yamaji, A. 421(1987)372
- Katayama, K., see Yamada, K. 424(1988)367
- Katayama, K., see Yamada, K. 433(1988)243
- Kato, A., see Moriyasu, M. 411(1987)466
- Kato, C., see Hasegawa, I. 410(1987)137
- Kato-Jippo, T., see Koyama, Y. 439(1988)417
- Kato, Y., Kitamura, T., Mitsui, A., Yamasaki, Y., Hashimoto, T., Murotsu, T., Fukushige, S. and Matsubara, K. Separation of oligonucleotides by high-performance ion-exchange chromatography on a non-porous ion exchanger 447(1988)212
- Kato, Y., see Iwase, H. 416(1987)37
- Kato, Y., see Moriyama, H. 445(1988)225
- Kato, Y., see Nakamura, K. 445(1988)234
- Kato, Y., Yamasaki, Y., Moriyama, H., Tokunaga, K. and Hashimoto, T. New high-performance gel filtration columns for protein separation 404(1987)333
- Katori, M., see Yamashita, K. 424(1988)1
- Katsanos, N.A. Reversed-flow GC applied to physico-chemical measurements 446(1988)39
- Katsura, E., see Kanetoshi, A. 442(1988)289
- Katti, A.M. and Guiochon, G.A. Prediction of band profiles in displacement chromatography by numerical integration of a semi-ideal model 449(1988)25
- Kauret, G., Herrle, I. and Schoppek, B. Detection of oxilofrine in plasma and urine by HPLC with electrochemical detection and comparison with GC-MS 428(1988)291
- Kauffman, G.B., see Ray, R.K. 442(1988)381
- Kaufmann, S.H.E., see Schoel, B. 448(1988)165
- Kaul, R., Olsson, U. and Mattiasson, B. Affinity chromatography on a hydrophobic matrix using a heterobifunctional ligand 438(1988)339
- Kaul, R.K., see Manja, K.S. 415(1987)377
- Kausch, H.-H., see Nguyen, T.Q. 449(1988)63
- Kaushik, M.P., see Ramachandran, P.K. 426(1988)239
- Kawada, K., see Hosoe, M. 435(1988)253
- Kawada, T., see Shibata, K. 417(1987)173
- Kawada, T., see Shibata, K. 422(1987)257
- Kawada, T., see Shibata, K. 424(1988)23
- Kawada, T., see Shibata, K. 430(1988)381
- Kawai, S., Goto, K., Kano, K. and Kubota, T. Determination of carbaryl by HPLC with electrochemical detection 442(1988)451
- Kawai, S., see Tomita, M. 414(1987)454
- Kawaka, K., see Hosoe, M. 438(1988)225
- Kawakami, N., see Sakamoto, Y. 442(1988)69
- Kawakatsu, M., see Kotaniguchi, H. 420(1987)141
- Kawamura, K. and Gagosian, R.B. Identification of aliphatic keto carboxylic acids in marine aerosols using capillary GC-MS 438(1988)299
- Kawamura, K. and Gagosian, R.B. Identification of isomeric hydroxy fatty acids in aerosol samples by capillary GC-MS 438(1988)309
- Kawamura, N., see Ikai, Y. 411(1987)313
- Kawamura, N., see Oka, H. 411(1987)437
- Kawamura, N., see Oka, H. 449(1988)448
- Kawanabe, K., see Tanabe, S. 424(1988)29
- Kawasaki, H., see Itami, T. 417(1987)168
- Kawase, M., see Ohmori, S. 414(1987)149
- Kawase, M., see Ohmori, S. 415(1987)221
- Kawase, M., see Ohmori, S. 426(1988)15
- Kawashima, T., see Nakagawa, T. 420(1987)297
- Kayano, M., see Miwa, T. 408(1987)316
- Kaye, S.B., see Cummings, J. 431(1988)77
- Kazemier, B., see Welling-Wester, S. 443(1988)255
- Ke, J., see Tam, Y.K. 423(1987)199
- Kee, T.G., see Griffin, R.M.E. 445(1988)441
- Kee, Y., Lee, Y.S., Chung, C.H., Waxman, L. and Goldberg, A.L. Improved methods for purification and assay of glycerol kinase from *Escherichia coli* 428(1988)345
- Keech, A.C., Harrison, P.M. and McLean, A.J. Simple extraction of atenolol from urine and its determination by HPLC 426(1988)234
- Keeling, P.W.N., see O'Reilly, T. 417(1987)190
- Keeney, R.F., see Weber, D.V. 431(1988)55
- Kehrer, J.P., see Murphy, M.E. 421(1987)71
- Keigher, N., see Miwa, B.J. 431(1988)343
- Keim, J.M.B., see Stowers, D.J. 444(1988)47

- Keizer, H.J., see Reeuwijk, H.J.E.M. 428(1988)93
- Keler, T., see Samu, J. 435(1988)167
- Keller, H.P., see Humbert, H. 422(1987)205
- Keller, M.A. and Saba, C.S.  
Chromatographic analysis of phenyl-1-naphthylamine and 4,4'-dioctyldiphenylamine and their intermediate oxidation products in oxidized lubricants 409(1987)325
- Keller, N., see Coleman, A.W. 450(1988)175
- Kelley, V., see Kelly, R.A. 416(1987)247
- Kelly, E.L., see Lee, J.K. 444(1988)141
- Kelly, G.M., see Meng, H. 443(1988)183
- Kelly, J.G., see Kinney, C.D. 419(1987)433
- Kelly, J.M., see Early, R.J. 416(1987)15
- Kelly, P.M.  
Separation of amphotericin B components by combined TLC and HPLC 437(1988)221
- Kelly, R.A., O'Hara, D.S. and Kelley, V.  
HPLC separation of femtomolar quantities of endogenous carboxylic acids, including arachidonic acid metabolites, as 4-bromomethyl-7-acetoxycoumarin derivatives 416(1987)247
- Kennedy, C.M., see Pullen, R.H. 434(1988)271
- Kennedy, E.E., see Taylor, E.H. 416(1987)365
- Kennedy, J.F., see Lloyd, L.L. 437(1988)447
- Kennedy, J.S., Friedman, H., Scavone, J.M., Harmatz, J.S., Shader, R.I. and Greenblatt, D.J.  
Effect of blood collection tubes on antidepressant concentrations 423(1987)373
- Kennedy, J.S., Leduc, B.W., Scavone, J.M., Harmatz, J.S., Shader, R.I. and Greenblatt, D.J.  
Pharmacokinetics of intravenous caffeine: comparison of HPLC and GC methods 422(1987)274
- Kennedy, L., see Heinitz, M.L. 443(1988)173
- Kennerly, D.A.  
Molecular species analysis of lysophospholipids using HPLC and argentation TLC 409(1987)291
- Kenney, W.C., see Watson, E. 436(1988)289
- Kennish, J.M., see Whitsett, J.F. 435(1988)343
- Kern, J.R., Lokensgard, D.M., Manes, L.V., Matsuo, M. and Nakamura, K.  
Separation of the stereoisomers of an allenic E-type prostaglandin 450(1988)233
- Kern, S., see Baltensperger, U. 439(1988)121
- Kerns, E.H., Russell, J.W. and Gallo, D.G.  
Determination of anagrelide in plasma by capillary GC-MS 416(1987)357
- Kerr, C.R., see Chan, G.L.-Y. 417(1987)295
- Kerr, C.R., see McErlane, K.M. 415(1987)335
- Kerr, D.J., see Cummings, J. 431(1988)77
- Kersten, B.R., Poole, C.F. and Furton, K.G.  
Ambiguities in the determination of McReynolds stationary phase constants 411(1987)43
- Kersten, B.R., see Shetty, P.H. 411(1987)61
- Kéry, Á., Turiák, G. and Tétényi, P.  
Isolation of parthenolide by droplet counter-current chromatography 446(1988)157
- Kester, M.B., Sacchar, C.L. and Mansmann, Jr., H.C.  
Microassay for the simultaneous determination of theophylline and dyphylline in serum by HPLC 416(1987)91
- Keuken, M.P., Slanina, J., Jongejan, P.A.C. and Bakker, F.P.  
Optimization of ion chromatography using commercially available detection systems and software 439(1988)13
- Keuth, V., see Townsend, R.W. 450(1988)414
- Kežić, S. and Monster, A.C.  
Determination of methyl ethyl ketone and its metabolites in urine using capillary GC 428(1988)275
- Khac, D.D., see Van, T.D. 440(1988)71
- Khachik, F. and Beecher, G.R.  
Separation of carotenol fatty acid esters by HPLC 449(1988)119
- Khalil, S.K.W., see Nygard, G.A. 415(1987)438
- Khan, H.M., see Ahmad, S. 440(1988)467
- Khan, N., see Ahmad, S. 440(1988)467
- Khan, N.H., Roets, E., Hoogmartens, J. and Vanderhaeghe, H.  
Quantitative analysis of oxytetracycline and related substances by HPLC 405(1987)229
- Khambade, B.V., Srivastava, N., Joshi, G.P. and Agrawal, O.P.  
TLC analysis of some Indian natural resins occurring in art objects 439(1988)430
- Khlebnikova, N.S., see Rogoskin, V.A. 423(1987)33
- Kholy, A.M. El, see Brooks, J.B. 430(1988)209
- Khoury, H.E. and Ibrahim, R.K.  
Resolution of five position-specific flavonoid O-methyltransferases by fast protein liquid chromatofocusing 407(1987)291
- Khvatov, A.B., see Maltsev, V.G. 416(1987)45
- Kibwage, I.O., Roets, E., Verbruggen, A., Hoogmartens, J. and Vanderhaeghe, H.  
TLC study of the metabolites of erythromycins in the Wistar rat 434(1988)177
- Kibwage, I.O., see Cachet, T. 409(1987)91
- Kidouchi, K., Sugiyama, N., Morishita, H., Wada, Y., Nagai, S. and Sakakibara, J.  
Analytical method for urinary

- glutarylcarntine, acetylcarntine and propionylcarntine with a carboxylic acid analyser and a reversed-phase column 423(1987)297
- Kiechel, J.R., see Humbert, H. 417(1987)319
- Kienle, M.G., see Paroni, R. 420(1987)189
- Kientz, C.E. and Verweij, A.  
Decomposition of trifluoroacetyl derivatives of some trichothecenes on fused-silica capillary columns during GC analysis 407(1987)340
- Kienzl, E. and Eichinger, K.  
Preparation of dopamine 3-O-sulphate and dopamine 4-O-sulphate as reference substances and HPLC trace determination 430(1988)263
- Kihara, K., Rokushika, S. and Hatano, H.  
Elution behaviour of aliphatic carboxylic acids on a strong cation-exchange resin column 410(1987)103
- Kihara, K., see Murayama, T. 435(1988)417
- Kiho, T., see Ukai, S. 435(1988)374
- Kikuchi, H., see Matsumoto, K. 425(1988)323
- Kikuchi, J., Nakamura, K., Nakata, O. and Morikawa, Y.  
Determination of monosaccharides constituting the glycosides in saponins by HPLC 403(1987)319
- Kikuchi, M., see Miwa, H. 416(1987)237
- Kilár, F., see Hjertén, S. 403(1987)47
- Kim, J.Y., see Adachi, K. 428(1988)247
- Kim, M., see Kubo, I. 402(1987)354
- Kim, N., see Yim, J.J. 419(1987)296
- Kim, S., see Park, K.S. 440(1988)225
- Kimura, K., see Kudo, K. 431(1988)353
- Kimura, K., see Uehori, R. 411(1987)251
- Kimura, M., see Hiraike, H. 430(1988)143
- Kimura, S., see Utsumi, J. 421(1987)355
- Kimura, T., see Ohta, M. 402(1987)392
- Kingma, A.W., see Van den Berg, G.A. 415(1987)27
- Kinjo, S., see Ohta, H. 409(1987)409
- Kinney, C.D. and Kelly, J.G.  
LC determination of paracetamol and dextropropoxyphene in plasma 419(1987)433
- Kinoshita, M., see Okazaki, M. 430(1988)135
- Kinoshita, T., see Iwaki, K. 404(1987)117
- Kinoshita, T., see Iwaki, K. 407(1987)273
- Kinoshita, T., see Kobayashi, Y. 430(1988)65
- Kinoshita, T., see Nimura, N. 402(1987)387
- Kinoshita, T., see Yoshida, T. 445(1988)175
- Kintz, P., Mangin, P., Lugnier, A.A. and Chaumont, A.J.  
GC assay for dextromoramide in human plasma 432(1988)329
- Kintz, P., see Tracqui, A. 432(1988)412
- Kip, J., see Der Haar, J.V. 445(1988)219
- Kirch, W., see Santos, S.R.C.J. 419(1987)155
- Kirillova, N.V., see Manoilov, S.E. 440(1988)53
- Kirino, O., see Takahashi, J. 436(1988)316
- Kiriyama, T., see Sagara, Y. 431(1988)170
- Kirjanen, I., see Löhmus, M. 449(1988)77
- Kirk, D.W., see Gattrell, M.A. 409(1987)404
- Kirk, K.L., see Eisenhofer, G. 431(1988)156
- Kirk, K.L., see Gusovsky, F. 415(1987)124
- Kirkes, K.A., see Robinson, J.L. 402(1987)201
- Kishi, T., see Okamoto, T. 430(1988)11
- Kiso, Y., see Hirokawa, T. 408(1987)27
- Kiso, Y., see Hirokawa, T. 410(1987)279
- Kiss, F., see Šalamoun, J. 419(1987)213
- Kiss, L., see Gál, S. 403(1987)217
- Kiss, L., Tar, A., Gál, S., Toth-Martinez, B.L. and Hernádi, F.J.  
Modified general affinity adsorbent for large-scale purification of penicillinases 448(1988)109
- Kissinger, P.T., see Isaksson, K. 419(1987)165
- Kissmeyer-Nielsen, A.-M.  
Determination of 6 $\beta$ -bromopenicillanic acid (broactam) in human serum and urine by HPLC 425(1988)425
- Kita, H., see Yamashita, J. 403(1987)275
- Kitagawa, N.  
Ion-exchange chromatography of proteins on a polyethyleneimine-grafted hydrophilic polymer for HPLC 443(1988)133
- Kitahara, M., see Tanabe, S. 424(1988)29
- Kitamura, T., see Kato, Y. 447(1988)212
- Kitano, H., Hirai, Y., Okada, Y., Nakamura, K. and Ise, N.  
Pulse injection analysis of the binding of serum proteins to porous polymer gels modified with formyl groups 420(1987)13
- Kitazawa, S., see Suzuki, K. 425(1988)435
- Kitazawa, Y., see Takeyama, N. 424(1988)361
- Kithinji, J.P., see Raynor, M.W. 436(1988)497
- Kitsuwa, T., see Goto, T. 447(1988)410
- Kitzman, J.V., Martin, J.F., Holley, J.H. and Huber, W.G.  
Determination of antipyrine in catfish plasma by HPLC 437(1988)306
- Klaassen, R., see Derks, H.J.G.M. 442(1988)436
- Klecker, Jr., R.W., see Belanger, K. 417(1987)57
- Klee, M.S., see Hepp, M.A. 404(1987)145
- Klein, M.L., see Lu, H.S. 447(1988)351
- Kleinjans, J.C.S., see Verhagen, H. 413(1987)282
- Kleinjans, J.C.S., see Verhagen, H. 422(1987)288
- Kleinmann, I., see Plicka, J. 442(1988)237
- Kleinmann, I., Svoboda, V. and Vočková, J.  
Preparative separation of an enzymatic

- hydrolysate of  $^{14}\text{C}$ -labelled proteins  
411(1987)335
- Klemm, D., Hartmann, M., Stein, A., Schnabelrauch, M., Spangenberg, S., Gross, M., Klepel, M. and Jumar, A.  
TLC studies of biocidal 2-bromo-2-nitropropane derivatives (in German)  
438(1988)122
- Klepel, M., see Klemm, D. 438(1988)122
- Klesper, E. and Schmitz, F.P.  
Single and multiple gradients in supercritical fluid chromatography (Review) 402(1987)1
- Kleuss, C., Pallast, M., Brendel, S., Rosenthal, W. and Schultz, G.  
Resolution of transducin subunits by chromatography on Blue Sepharose  
407(1987)281
- Kleyn, E. Van der, see Wuis, E.W. 415(1987)419
- Kleyn, E. Van der, see Wuis, E.W. 420(1987)212
- Klick, S. and Herrmann, K.  
Analysis of naturally occurring hydroxybenzoic acid glucosides and hydroxybenzoyl glucoses and their behaviour on a polyamide column 441(1988)335
- Kling, G.J., Perkins, L.M., Cappiello, P.E. and Eisenberg, B.A.  
Improved method for the determination of indole-3-acetic acid in plant tissue  
407(1987)377
- Klinot, J., see Snopek, J. 450(1988)373
- Klinotová, E., see Snopek, J. 450(1988)373
- Klintebäck, F., see Wester, P. 415(1987)261
- Klocke, J.A., see Yamasaki, R.B. 447(1988)277
- Klockowski, P.M. and Levy, G.  
Simultaneous determination of diazepam and its active metabolites in rat serum, brain and cerebrospinal fluid by HPLC 422(1987)334
- Klooster, G.A.E.V. 't, Boot, J.H., Mennes, W.C. and Blaauboer, B.J.  
Rapid method for the determination and quantification of bromosulphophthalein and metabolites in cultured hepatocytes, culture media and bile by HPLC 432(1988)223
- Klueppelberg, U., see Swobodnik, W. 423(1987)75
- Knapp, D.R., see Vrbanac, J.J. 425(1988)1
- Knausz, D., see Szederkényi, F. 446(1988)253
- Knedel, M., see Jacob, K. 441(1988)171
- Knegt, E., see Vermeer, E. 404(1987)346
- Knight, M., Mack, G.D., Perkins, R. and Burke, Jr., T.R.  
Reversed-phase preparative chromatography of [D-Ala<sup>1</sup>]-peptide T amide 444(1988)345
- Knight, M., see Burke, Jr., T.R. 411(1987)431
- Knijff, J.H., see Odink, J. 424(1988)273
- Knobler, H., Fainaru, M. and Sklan, D.  
Single-stage evaluation of serum lipoproteins by gel permeation using a specific fluorescent lipid probe 421(1987)136
- Knobloch, E., Mandys, F. and Hodr, R.  
Study of the mechanism of the photochemical oxidation of bilirubin by HPLC 428(1988)255
- Knöller, J., König, W., Schönfeld, W., Bremm, K.D. and Köller, M.  
Application of HPLC of some antibiotics in clinical microbiology 427(1988)257
- Knöller, J., Schönfeld, W., Köller, M., Hensler, T. and König, W.  
Arachidonic acid metabolites from polymorphonuclear leukocytes of healthy donors, severely burned patients and children with cystic fibrosis – routine monitoring by HPLC 427(1988)199
- Knospe, J., see Herrmann, T. 428(1988)237
- Knospe, J., Steinhilber, D., Herrmann, T. and Roth, H.J.  
Picomole determination of 2,4-dimethoxyanilides of prostaglandins by HPLC with electrochemical detection 442(1988)444
- Knox, J.H., Kříž, J. and Adamcová, E.  
Retention relationships for aromatic hydrocarbons eluted from capped and uncapped octadecyl silica gels 447(1988)13
- Knox, M.J., see Ashoor, S.H. 409(1987)419
- Knudsen, J.J. and McGown, E.L.  
GC analysis of urinary dimercaptosuccinic acid 424(1988)231
- Knuth, D.W., see La Follette, G. 431(1988)379
- Kobayashi, A., see Sugita, S. 421(1987)412
- Kobayashi, K., Sakoguchi, T., Fujiwara, K., Taniuchi, K., Kohri, K. and Matsuoka, A.  
HPLC determination of amphotericin B in human urine 417(1987)439
- Kobayashi, S.-I.  
Simple method for the determination of nicardipine in plasma using HPLC 420(1987)439
- Kobayashi, S., see Hirokawa, T. 410(1987)279
- Kobayashi, Y., Kubo, H., Kinoshita, T. and Nishikawa, T.  
Fluorometric determination of biguanides in serum by HPLC with reagent-containing mobile phase 430(1988)65
- Koch, C.S., Köster, F. and Findenegg, G.H.  
Adsorption of binary solvent mixtures in reversed-phase chromatographic systems. Surface excess isotherms and enthalpies of displacement 406(1987)257
- Koch, E.M., see Koppenhoefer, B. 406(1987)157



- Koch, H., see Girre, C. 419(1987)363  
 Koch, H.P., see Czejka, M.J. 413(1987)181  
 Kocher, H.P., see Hochuli, E. 411(1987)371  
 Kock-Van Dalen, A.C., see Sinninghe Damsté, J.S. 435(1988)435  
 Kocna, P., Frič, P., Kočová-Holáková, M., Slabý, J., Kasafirek, E. and Hekkens, W.T.J.M. Isolation and analysis of peptidic fragments of  $\alpha$ -gliadin using reversed-phase HPLC 434(1988)429  
 Kocourek, J., see Šecová, E. 436(1988)59  
 Kocourek, V., see Hajšlová, J. 439(1988)307  
 Kočová-Holáková, M., see Kocna, P. 434(1988)429  
 Kodama, C., Kodama, T. and Yosizawa, Z. Methods for analysis of urinary glycosaminoglycans (Review) 429(1988)293  
 Kodama, H., see Ageta, T. 419(1987)334  
 Kodama, T., see Kodama, C. 429(1988)293  
 Koechlin, C., Jehl, F. and Monteil, H. Determination of PI<sub>A</sub> and PII<sub>A</sub>, the two main components of pristinamycin, by HPLC in plasma 425(1988)197  
 Koel, M., see Nebinger, P. 427(1988)326  
 Koel, M., see Nebinger, P. 434(1988)43  
 Koenigbauer, M., see Ravichandran, K. 439(1988)213  
 Koenigbauer, M.J., Assenza, S.P., Willoughby, R.C. and Curtis, M.A. Trace analysis of diazepam in serum using microbore HPLC and on-line preconcentration 413(1987)161  
 Koenigbauer, M.J. and Curtis, M.A. Use of micellar mobile phases and microbore column switching for the assay of drugs in physiological fluids 427(1988)277  
 Koerner, Jr., P.J. and Nieman, T.A. HPLC determination of glucosides (glucose conjugates) with post-column reaction detection combining immobilized enzyme reactors and luminol chemiluminescence 449(1988)217  
 Koffel, J.C., see Efthymiopoulos, C. 421(1987)360  
 Koffigan, M., see Mezdour, H. 414(1987)35  
 Koga, R., see Yamashita, K. 424(1988)39  
 Koga, T., see Horikoshi, T. 416(1987)353  
 Kohara, H., see Shimizu, K. 432(1988)21  
 Kohara, H., see Yamaga, N. 422(1987)25  
 Kohara, N., see Sakamoto, M. 411(1987)259  
 Kohashi, K., see Tsuruta, Y. 416(1987)63  
 Kohda, Y., see Yamamoto, K. 433(1988)167  
 Kohlíček, J., see Balíková, M. 431(1988)431  
 Kohlíček, J., see Balíková, M. 434(1988)469  
 Kohout, J., see Kučerová, Z. 419(1987)323  
 Kohri, K., see Kobayashi, K. 417(1987)439  
 Kohse, K.P. and Wisser, H. Quantitative monitoring by HPLC of the dissociation of human serum cholinesterase by limited proteolysis 420(1987)25  
 Koike, H., see Arai, T. 448(1988)439  
 Koizumi, K., Kubota, Y., Okada, Y., Utamura, T., Hizukuri, S. and Abe, J.-I. Retention behaviour of cyclodextrins and branched cyclodextrins on reversed-phase columns in HPLC 437(1988)47  
 Koizumi, K., Utamura, T., Kubota, Y. and Hizukuri, S. Two HPLC columns for analyses of malto-oligosaccharides 409(1987)396  
 Koizumi, K. and Utamura, T. HPLC of gluco-oligomers on C<sub>18</sub>-bonded vinyl alcohol copolymer gel with alkaline eluents 436(1988)328  
 Kojima, C., see Matsubayashi, K. 433(1988)225  
 Kojima, K., see Sakai, T. 416(1987)131  
 Kojima, K., Uezono, Y., Takahashi, T. and Nakanishi, Y. HPLC method for the determination of a cyclandelate metabolite, mandelic acid, in human plasma 425(1988)203  
 Kojima, K., Yamamoto, K., Nakanishi, Y. and Katae, H. Determination of 22,23-dihydroavermectin B<sub>1a</sub> in dog plasma using solid-phase extraction and HPLC 413(1987)326  
 Kojima, T., see Naoi, M. 426(1988)75  
 Kok, W.T., Hupe, K.P. and Frei, R.W. Zone-electrophoretic sample treatment for LC 436(1988)421  
 Kokubu, T., see Takada, Y. 421(1987)350  
 Kolář, J., see Mikeš, O. 440(1988)287  
 Kolář, J., see Tennikova, T.B. 435(1988)357  
 Kolarz, B.N., Łuczyński, J., Trochimczuk, A. and Wojaczyńska, M. GC properties of porous polyacrylonitrile and divinylbenzene copolymers 408(1987)308  
 Köller, M., see Knöller, J. 427(1988)199  
 Köller, M., see Knöller, J. 427(1988)257  
 Kolmer, E.W.J. Beneken, see Wuis, E.W. 415(1987)419  
 Kolset, K., see Johnsen, S. 438(1988)233  
 Komatsu, S., see Marunaka, T. 433(1988)177  
 Kominami, G., Matsumoto, H., Nishimura, R., Yamaguchi, T., Konaka, R., Sugeno, K., Hirai, K. and Kono, M. Combined HPLC and enzyme immunoassay for active metabolites of a sleep inducer (rilmazafone), a ring-opened derivative of benzodiazepines, in human plasma 417(1987)216

- Komoda, T., see Koyama, I. 413(1987)65  
 Komoda, T., see Koyama, I. 420(1987)275  
 Komov, V.P., see Manoilov, S.E. 440(1988)53  
 Komuro, T., see Galanos, C. 440(1988)397  
 Komuro, T. and Galanos, C.  
 Analysis of Salmonella lipopolysaccharides by sodium deoxycholate-polyacrylamide gel electrophoresis 450(1988)381  
 Kon, K., see Ando, S. 405(1987)125  
 Kon, K., see Ando, S. 408(1987)285  
 Kon, T.C., see Evelo, C.T.A. 427(1988)335  
 Konaka, R., see Kominami, G. 417(1987)216  
 Kondo, N., see Takeuchi, T. 435(1988)482  
 Kondoh, M., see Naoi, M. 426(1988)75  
 Kondoh, Y. and Takano, S.  
 Determination of zinc pyrrithione in cosmetic products by HPLC with pre-labelling 408(1987)255  
 Konečná, V., see Verner, P. 416(1987)439  
 Kong, A.-N., Slaughter, R.L. and Jusko, W.J.  
 Simultaneous analysis of methylprednisolone hemisuccinate, cortisol and methylprednisolone by normal-phase HPLC in human plasma 432(1988)308  
 König, W., see Knöller, J. 427(1988)199  
 König, W., see Knöller, J. 427(1988)257  
 König, W.A., Lutz, S., Mischnick-Lübbecke, P., Brassat, B. and Wenz, G.  
 Cyclodextrins as chiral stationary phases in capillary GC. I. Pentylated  $\alpha$ -cyclodextrin 447(1988)193  
 König, W.A., see Rieck, M. 439(1988)301  
 König, W.A., see Schmidt, N. 410(1987)458  
 König, W.A., see Weller, O. 403(1987)263  
 Kono, M., see Kominami, G. 417(1987)216  
 Kononenko, V.L., see Semyonov, S.N. 446(1988)141  
 Konteatis, Z., see Florance, J. 414(1987)313  
 Kooistra, C. and Sluyterman, L.A.Æ.  
 Feasibility of isotachochromatography as a method for the preparative separation of weak acids and weak bases. I. Theoretical considerations 436(1988)173  
 Kooistra, C. and Sluyterman, L.A.Æ.  
 Feasibility of isotachochromatography as a method for the preparative separation of weak acids and weak bases. II. Experimental verification 436(1988)181  
 Koopman, B.J., Van der Molen, J.C., Wolthers, B.G. and Vanderpas, J.B.  
 Determination of some hydroxycholesterols in human serum samples 416(1987)1  
 Koopman-Kimenai, P.M., see Vree, T.B. 428(1988)311  
 Koopman-Kimenai, P.M., Vree, T.B., Cress-  
 Tijhuis, M.W., Booy, L.H.D.J. and Drijkoningen, G.  
 HPLC and preliminary pharmacokinetics of nicomorphine and its metabolites 3-nicotinoyl- and 6-nicotinoylmorphine and morphine 416(1987)382  
 Kopaciewicz, W., Fulton, S. and Lee, S.Y.  
 Influence of pore and particle size on the frontal uptake of proteins. Implications for preparative anion-exchange chromatography 409(1987)111  
 Kopaciewicz, W., see Heinitz, M.L. 443(1988)173  
 Kopečni, M.M., see Todorović, M. 442(1988)105  
 Kopin, I.J., see Eisenhofer, G. 431(1988)156  
 Kopin, I.J., see Hovevey-Sion, D. 426(1988)141  
 Köppel, C., Tenczer, J. and Ibe, K.  
 GC-MS study of urinary metabolism of melperone 427(1988)144  
 Köppel, C. and Tenczer, J.  
 GC-MS study of the urinary metabolism of trimipramine 431(1988)197  
 Koppele, J.M. Te, Van der Mark, E.J. and Mulder, G.J.  
 LC determination of diastereomeric glutathione conjugates and further derivatives of  $\alpha$ -bromoisovalerylurea in rat bile and urine by electrochemically generated bromine 427(1988)67  
 Køppen, B., Dalgaard, L. and Christensen, J.M.  
 Determination of trichloroethylene metabolites in rat liver homogenate using headspace GC 442(1988)325  
 Køppen, B., see Christensen, J.M. 442(1988)317  
 Koppenhoefer, B., Allmendinger, H., Lu, P.C. and Lin, B.C.  
 Resolution of stereoisomers of dipeptides by GC on Chirasil-Val 441(1988)89  
 Koppenhoefer, B., Koch, E.M., Nicholson, G.J. and Bayer, E.  
 Chiral recognition of halo carboxylic acid amides studied by GC on Chirasil-Val 406(1987)157  
 Kopycki, W., see Kudzin, Z.H. 449(1988)306  
 Kora, V.C.I., see Mezdour, H. 414(1987)35  
 Korach, K.S., see Chae, K. 439(1988)484  
 Korbová, L., see Kučerová, Z. 419(1987)323  
 Korf, J., see Wolf, J.H. 436(1988)437  
 Korfmacher, W.A., see Holder, C.L. 419(1987)113  
 Körner, J., see Geschke, R. 420(1987)111  
 Korolev, A.A., see Berezkin, V.G. 440(1988)323  
 Koronelli, T.V.  
 Investigation of the lipids of saprophytic mycobacteria in the U.S.S.R. (Review) 440(1988)479

- Korpela, T., see Mäkelä, M. 448(1988)391
- Korpela, T., see Mattsson, P. 447(1988)398
- Korte, D. De, Marijnen, Y.M.T., Haverkort, W.A., Van Gennip, A.H. and Roos, D.  
Sensitive on-line radioactivity measurements with a heterogeneous flow-cell: application to lymphoid cell ribonucleotides separated by HPLC 415(1987)383
- Korthals, H., see Odink, J. 424(1988)273
- Kosarych, Z., see Florance, J. 414(1987)313
- Koshakji, R.P. and Wood, A.J.J.  
Improved HPLC method for the simultaneous determination of propranolol and 4-hydroxypropranolol in plasma with fluorescence detection 422(1987)294
- Koshida, K., see Tawa, R. 425(1988)143
- Koss, M.C., see Gherezghiher, T. 413(1987)9
- Kossien, I., see Jacob, K. 441(1988)171
- Kostelc, J.G., see Preti, G. 432(1988)1
- Köster, F., see Koch, C.S. 406(1987)257
- Kotaki, H., see Aoyama, T. 430(1988)351
- Kotaniguchi, H., Kawakatsu, M., Toyooka, T. and Imai, K.  
Automatic amino acid analysis utilizing 4-fluoro-7-nitrobenzo-2-oxa-1,3-diazole 420(1987)141
- Kotoku, S., see Ikebuchi, J. 432(1988)401
- Köttgen, E., see Meyer-Sabellek, W. 429(1988)419
- Kotyński, A., see Drabowicz, J. 447(1988)225
- Kotzé, J.M., see Thomas, A.C. 403(1987)292
- Koupil, P., Novák, J. and Drozd, J.  
Comparison of the absolute calibration method with the method of standard addition for the determination of halothane in blood by GC headspace analysis 425(1988)99
- Kourteva, I.  
Chromatographic separation and partial characterization of microbial low-molecular-weight proteinase inhibitors 446(1988)203
- Kouwen, A.M. Van, see Rothuizen, J. 427(1988)19
- Kovacheva, E., see Neicheva, A. 437(1988)249
- Kováčová, D., see Svobodová, X. 425(1988)391
- Kovář, J., see Škabrahová, Z. 408(1987)372
- Kovář, J., see Turánek, J. 410(1987)470
- Kováts, E.  
Composition of essential oils. Part 7. Bulgarian oil of rose (*Rosa damascena* Mill.) 406(1987)185
- Kováts, E., see Fóti, G. 440(1988)315
- Kováts, E., see Samu, J. 435(1988)167
- Kowal, J., see Cheng, B. 432(1988)302
- Koyama, F., see Harada, H. 407(1987)299
- Koyama, I., Arai, K., Sakagishi, Y., Ikezawa, H. and Komoda, T.  
Blood appearance of rat alkaline phosphatase originating from the duodenum in vitro 420(1987)275
- Koyama, I., Miura, M., Matsuzaki, H., Sakagishi, Y. and Komoda, T.  
Sugar-chain heterogeneity of human alkaline phosphatases: differences between normal and tumour-associated isozymes 413(1987)65
- Koyama, Y., Hosomi, M., Miyata, A., Hashimoto, H., Reames, S.A., Nagayama, K., Kato-Jippo, T. and Shimamura, T.  
Supplementary and revised assignment of the peaks of the 7,9-, 9,9'-, 13,13'-, 9,13'-di-cis and 9,9',13-tri-cis isomers of  $\beta$ -carotene in HPLC using a column of calcium hydroxide 439(1988)417
- Koyama, Y., see Hashimoto, H. 448(1988)182
- Kraak, J.C., see De Jong, J. 414(1987)285
- Kraak, J.C., see Der Haar, J.V. 445(1988)219
- Kraak, J.C., see De Waal, W.A.J. 407(1987)253
- Krabbenborg, P., see Van Opstal, M.A.J. 432(1988)395
- Krahn, M.M., Moore, L.K., Bogar, R.G., Wigren, C.A., Chan, S.-L. and Brown, D.W.  
HPLC method for isolating organic contaminants from tissue and sediment extracts 437(1988)161
- Krajewska, A.M. and Powers, J.J.  
GC of methyl derivatives of naturally occurring capsaicinoids 409(1987)223
- Krajičková, J., see Macek, J. 414(1987)156
- Kral, T., see Mennicke, W.H. 414(1987)19
- Kram, T.C., see Lurie, I.S. 405(1987)273
- Kram, T.C., see Moore, J.M. 410(1987)297
- Krämer, B., see Röss, K. 417(1987)465
- Krämer, B.K., Mayer, F., Liebich, H.M., Röss, K.M., Kühnkamp, V., Becker, J.U., Risler, T. and Seipel, L.  
Rapid and inexpensive HPLC method for the quantification of flecainide in serum 427(1988)351
- Krasilnikov, I. and Borisova, V.  
Adsorption and chromatographic properties of modified silica sorbents for the production of viral preparations 446(1988)211
- Kraus, A., see Matisová, E. 439(1988)381
- Kraus, E., see Kuchař, M. 449(1988)391
- Kraus, G., see Matisová, E. 439(1988)381
- Kraus, G., see Soják, L. 436(1988)47
- Kraus, R., Wuthe, J. and Rüfer, R.  
Method for determination of different phospholipids by circular HPTLC 413(1987)257

- Krause, M. and Galensa, R.  
Direct enantiomeric separation of racemic flavanones by HPLC using cellulose triacetate as a chiral stationary phase 441(1988)417
- Krause, R.T.  
HPLC determination of aryl N-methylcarbamate residues using post-column hydrolysis electrochemical detection 442(1988)333
- Krauss, G.-J., see Abel, S. 446(1988)187
- Krejčí, M., see Janeček, M. 438(1988)409
- Krejcie, T.C., see Avram, M.J. 414(1986)484
- Kremmer, T., Boldizsár, M. and Holczinger, L.  
Comparison of high-performance ion-exchange and ion-pair LC methods for the separation of tumour cell nucleotides 415(1987)53
- Kreutzmann, D.J. and Silink, M.  
Determination of 17-oxosteroid glucuronides and sulphates in urine by LC using 2,4-dinitrophenylhydrazine as a prelabelling reagent for spectrophotometric detection 415(1987)253
- Krijt, J., see Samitrák, J. 415(1987)129
- Krijt, J., see Tvrzická, E. 431(1988)231
- Kriss-Danziger, P., see Herbranson, D.E. 410(1987)222
- Kříž, J., Punčochářová, J., Vodička, L. and Vařeka, J.  
HPLC of alkylbenzenes on alumina 437(1988)177
- Kříž, J., see Knox, J.H. 447(1988)13
- Krob, E.J., see Walter, H. 441(1988)261
- Krogt, J.A. Van der, Van Valkenburg, C.F.M. and Belfroid, R.D.M.  
Tritium isotope effect in reversed-phase HPLC analysis of dopamine and dihydroxyphenylacetic acid 427(1988)9
- Król, P., see Szewczyk, H. 446(1988)109
- Kroviarski, Y., Cochet, S., Vadon, C., Truskolaski, A., Boivin, P. and Bertrand, O.  
New strategies for the screening of a large number of immobilized dyes for the purification of enzymes. Application to the purification of enzymes from human haemolysate 449(1988)403
- Kroviarski, Y., Cochet, S., Vadon, C., Truskolaski, A., Boivin, P. and Bertrand, O.  
Purification of human 6-phosphogluconate dehydrogenase from human haemolysate with chromatography on an immobilized dye as the essential step and use of automation. Simultaneous purification of lactate dehydrogenase 449(1988)413
- Krstulović, A.M., Gianviti, J.M., Burke, J.T. and Mompon, B.  
Enantiomeric analysis of a new anti-inflammatory agent in rat plasma using a chiral  $\beta$ -cyclodextrin stationary phase 426(1988)417
- Krstulović, A.M., Rossey, G., Porziemsky, J.-P., Long, D. and Chekroun, I.  
Direct determination of the enantiomeric purity of (5*S*)-3-isopropyl-5-*p*-toluenesulphonyloxymethylloxazolidin-2-one on a cellulose-based chiral stationary phase. In-process control of a chiral intermediate used in the synthesis of enantiomerically pure  $\beta$ -blocking agents 411(1987)461
- Krstulović, A.M., see Porziemsky, J.-P. 440(1988)183
- Kruck, T.P.A., Teichert-Kuliszewska, K., Fisher, E., Kalow, W. and McLachlan, D.R.  
HPLC analysis of desferrioxamine. Pharmacokinetic and metabolic studies 433(1988)207
- Krüger, C., see Fürst, P. 405(1987)311
- Kruijff, N. De, Rijk, M.A.H., Pranoto-Soetardhi, L.A. and Schouten, A.  
Determination of preservatives in cosmetic products. I. TLC procedure for the identification of preservatives in cosmetic products 410(1987)395
- Krull, I.S., Stuting, H.H. and Krzysko, S.C.  
Conformational studies of bovine alkaline phosphatase in hydrophobic interaction and size-exclusion chromatography with linear diode array and low-angle laser light scattering detection 442(1988)29
- Krumbiegel, G., see Mennicke, W.H. 414(1987)19
- Krungkrai, J., Yuthavong, Y. and Webster, H.K.  
HPLC assay for pteroylpolyglutamate hydrolase 417(1987)47
- Krupčík, J., Repka, D., Benická, E., Hevesi, T., Nolte, J., Paschold, B. and Mayer, H.  
Use of retention temperatures for the identification of phenols separated on OV-1701 capillary columns using linear temperature-programmed GC 448(1988)203
- Krupčík, J., Repka, D., Hevesi, T. and Anders, G.  
Use of Kováts indices for prediction of retention temperatures in linear temperature-programmed capillary GC 407(1987)65
- Krupčík, J., Repka, D., Hevesi, T. and Garaj, J.  
Use of Kováts retention indices for characterizing solutes in complex samples,

- separated by linear temperature-programmed capillary GLC 406(1987)117
- Krylov, A.I., see Rogoskin, V.A. 423(1987)33
- Krzysko, S.C., see Krull, I.S. 442(1988)29
- Krzyżanowski, L., see Sikora, E. 442(1988)420
- Kuan, J.-C.W., see Johnsen, P.B. 409(1987)337
- Kubo, H., see Kobayashi, Y. 430(1988)65
- Kubo, I., Kim, M. and De Boer, G.  
Efficient isolation of insect growth inhibitory macrolide alkaloids using recycle high-performance GPC 402(1987)354
- Kubota, K., Horai, Y., Kushida, K. and Ishizaki, T.  
Determination of benzoic acid and hippuric acid in human plasma and urine by HPLC 425(1988)67
- Kubota, T., see Kawai, S. 442(1988)451
- Kubota, T., see Murayama, T. 435(1988)417
- Kubota, Y., see Koizumi, K. 409(1987)396
- Kubota, Y., see Koizumi, K. 437(1988)47
- Kucera, J., see Gottvaldova, M. 427(1988)331
- Kučerová, Z., Korbová, L. and Kohout, J.  
Isolation and characterization of a protein detected in inflamed human gastric mucosa 419(1987)323
- Kuchař, M., Kraus, E., Rejholec, V. and Miller, V.  
Enthalpy-entropy compensation in reversed-phase HPLC of series of aryloxoalkanoic and arylhydroxyalkanoic acids 449(1988)391
- Kucharczyk, N., see Pivonka, J. 420(1987)89
- Kudo, K., Nagata, T., Kimura, K., Imamura, T. and Noda, M.  
Sensitive determination of diazepam and N-desmethyldiazepam in human material using capillary GC-MS 431(1988)353
- Kudo, K., see Uehori, R. 411(1987)251
- Kudzin, Z.H., see Drabowicz, J. 447(1988)225
- Kudzin, Z.H. and Kopycki, W.  
Extraction chromatography of alkanethiols 449(1988)306
- Kühlkamp, V., see Krämer, B.K. 427(1988)351
- Kuhn, A.O. and Lederer, M.  
Adsorption chromatography on cellulose. I. Salting-out chromatography of organic compounds 406(1987)389
- Kuhn, A.O. and Lederer, M.  
Adsorption chromatography on cellulose. II. Separations of aromatic amino acids, biogenic amines, alkaloids, dyes and phenols and determination of hydrophobic constants 440(1988)165
- Kuhn, R.W. and Deyman, M.E.  
Simultaneous quantification of natural glycocorticoids and progestins in serum 421(1987)123
- Kuhn, W., see Wallasch, T.M. 425(1988)175
- Kuhnert, B.R., Brashear, W.T. and Syracuse, C.D.  
Measurement of alphaprodine by selected-ion monitoring 426(1988)392
- Kuhnz, W.  
Automated HPLC assay for cyproterone acetate and 15 $\beta$ -hydroxycyproterone acetate in plasma 420(1987)432
- Kulanthaivel, P. and Pelletier, S.W.  
HPLC analysis of C<sub>19</sub>-diterpenoid alkaloids 402(1987)366
- Kumagai, Y., Ishida, T. and Toki, S.  
Method for determination of morphinone in urine and bile of guinea pig by HPLC 421(1987)155
- Kumanotani, J., see Yamauchi, Y. 408(1987)171
- Kumar, H., see Ahmad, S. 440(1988)467
- Kumar, K., Morgan, D.J. and Crankshaw, D.P.  
Determination of fentanyl and alfentanil in plasma by HPLC with UV detection 419(1987)464
- Kumar, S. and Bhatia, K.L.  
Characterization of buffalo lactotransferrin by polyacrylamide gel electrophoresis in 6 M urea 434(1988)228
- Kuningas, K., see Rang, S. 406(1987)75
- Kunitake, T., see Hirayama, C. 409(1987)175
- Kunitani, M.G., Cunico, R.L. and Staats, S.J.  
Reversible subunit dissociation of tumor necrosis factor during hydrophobic interaction chromatography 443(1988)205
- Kunitomo, M., see Yamaguchi, Y. 422(1987)53
- Kuno, A., see Oka, H. 411(1987)437
- Kunz, F.-R., see Jork, H. 450(1988)115
- Kunze, G., see Paulos, C. 433(1988)359
- Kuo, K.C., see Souther, M.E. 417(1987)27
- Kupferberg, H.J., see Kapetanovic, I.M. 414(1987)265
- Kupferberg, H.J., see Torchin, C.D. 426(1988)444
- Kura, G.  
LC study of the hydrolysis reactions of cyclic and linear polyphosphates in aqueous solution 447(1988)91
- Kurachi, M., see Kurata, K. 434(1988)278
- Kurantz, M.J., see Doner, L.W. 449(1988)229
- Kurata, K., Kurachi, M. and Tani, Y.  
HPLC determination of mianserin in plasma and brain and its application to pharmacokinetic studies in the rat 434(1988)278
- Kurihara, H., see Ito, T. 417(1987)79
- Kuroda, K., see Hasegawa, I. 410(1987)137

- Kurosawa, T., see Tohma, M. 421(1987)9
- Kurotori, M., see Marunaka, T. 433(1988)177
- Kurumaya, K., Sakamoto, T., Okada, Y. and Kajiwara, M.  
Application of droplet counter-current chromatography to the isolation of vitamin B<sub>12</sub> 435(1988)235
- Kuš, P.  
TLC separation of some dithia[3.3]phanes. I. 436(1988)338
- Kuš, P.  
Thin-layer separation of some dithia[3.3]phanes. II. 442(1988)378
- Kusaka, K., see Nakano, K. 438(1988)61
- Kusala, S., see Ostrovská, V. 446(1988)323
- Kusala, S., see Svobodová, X. 425(1988)391
- Kushida, K., see Kubota, K. 425(1988)67
- Kushinsky, S., see Wu, A.T. 415(1987)65
- Kushwaha, R.S., see Williams, M.C. 433(1988)257
- Kusińska, E., see Muniak, W. 436(1988)323
- Kutt, H., see Cedarbaum, J.M. 415(1987)393
- Kuwajima, M., see Jinno, K. 436(1988)11
- Kuwamoto, T., see Hirayama, N. 447(1988)323
- Kuwamoto, T., see Okada, T. 403(1987)35
- Kuwata, K. and Tanaka, S.  
LC determination of traces of phenols in air 442(1988)407
- Kuwatsuka, S., see Spitzer, T. 435(1988)489
- Kvannes, J. and Flatmark, T.  
Rapid and sensitive assay of ornithine decarboxylase activity by HPLC of the *o*-phthalaldehyde derivative of putrescine 419(1987)291
- Kvernadze, T.K., see Andronikashvili, T.G. 406(1987)145
- Kwietniewski, L., see Wójcik, A. 435(1988)55
- Kwong, E.C. and Shen, D.D.  
Versatile isocratic HPLC assay for propranolol and its basic, neutral and acidic metabolites in biological fluids 414(1987)365
- Kyerematen, G.A., Taylor, L.H., DeBethizy, J.D. and Vesell, E.S.  
Radiometric-HPLC assay for nicotine and twelve of its metabolites 419(1987)191
- Kysilka, R. and Wurst, M.  
Optimized determination of indole derivatives by HPLC with selective detection 446(1988)315
- Laakso, I., see Ekundayo, O. 403(1987)358
- Labadarios, D., see Shephard, G.S. 416(1987)138
- Labarre, P., Godeneche, D., Madelmont, J.C., Triana, K., Mathe, G. and Veyre, A.  
Qualitative and quantitative analysis of a new sulphur-containing nitrosourea in blood by HPLC 419(1987)381
- Labows, J.N., see Preti, G. 432(1988)1
- Labrude, P., see Menu, P. 417(1987)397
- Lacey, E., see Dawson, M. 420(1987)129
- LaCreta, F.P., Olszewski, J.J. and Tew, K.  
Purification of glutathione S-transferases from rat liver and Walker 256 mammary carcinoma cells by HPLC and a glutathione affinity column 434(1988)83
- LaCreta, F.P. and Williams, W.M.  
HPLC analysis of fluoropyrimidine nucleosides and fluorouracil in plasma 414(1987)197
- Lacroix, C., Cerutti, F., Nouveau, J., Menager, S. and Lafont, O.  
Determination of ethambutol in plasma by LC and UV spectrophotometric detection (in French) 415(1987)85
- Lacroix, C., Poncet, P., Laine, G., Guyonnaud, C., Ray, M., Menager, S. and Lafont, O.  
Microdetermination of pyrazinamide and its metabolites (2-pyrazinoic acid, 5-hydroxypyrazinoic acid, 5-hydroxypyrazinamide and pyrazinuric acid) in plasma and urine by LC (in French) 422(1987)217
- Ladds, G., Wilson, K. and Burnett, D.  
Automated LC method for the determination of paracetamol and six metabolites in human urine 414(1987)355
- Ladešić, B., see Tomašić, J. 440(1988)405
- Ladure, P., see Chabenat, C. 414(1987)417
- Laduron, C., see Brassinne, C. 419(1987)401
- Laffort, P. and Patte, F.  
Solubility factors established by GLC. A balance sheet 406(1987)51
- La Follette, G., Gambertoglio, J., White, J.A., Knuth, D.W. and Lin, E.T.  
Determination of clindamycin in plasma or serum by HPLC with UV detection 431(1988)379
- Lafont, O., see Lacroix, C. 415(1987)85
- Lafont, O., see Lacroix, C. 422(1987)217
- Lafosse, M., Dreux, M. and Morin-Allory, L.  
Application fields of a new evaporative light

- scattering detector for HPLC and supercritical fluid chromatography (in French) 404(1987)95
- Laganá, A., Curini, R., D'Ascenzo, G., Marino, A. and Rotatori, M.  
HPLC determination of norfloxacin in human tissues and plasma with fluorescence detection 417(1987)135
- Lagarde, M., see Guichardant, M. 425(1988)25
- Lagerström, P.-O., see Balmér, K. 417(1987)357
- Laham, A., Claperon, N., Durussel, J.J., Fattal, E., Delattre, J., Puisieux, F., Couvreur, P. and Rossignol, P.  
Liposomally entrapped adenosine triphosphate. Improved efficiency against experimental brain ischaemia in the rat 440(1988)455
- Lai, P.-H., see Lu, H.S. 447(1988)351
- Lai, Y.-H., see Marriott, P.J. 447(1988)29
- Laine, G., see Lacroix, C. 422(1987)217
- Laing, D.K., Gill, R., Blacklaws, C. and Bickley, H.M.  
Characterisation of acid dyes in forensic fibre analysis by HPLC using narrow-bore columns and diode array detection 442(1988)187
- Lakings, D.B., Friis, J.M. and Bruns, M.B.  
HPLC method for the determination of eclanamine and its N-desmethyl and N,N-didesmethyl metabolites in urine 428(1988)131
- Lalande, M., Mills, P. and Peterson, R.G.  
Determination of dantrolene and its reduced and oxidized metabolites in plasma by HPLC 430(1988)187
- Lalande, M., see Dubé, L.M. 427(1988)113
- Lalka, D., see Schuster, D. 433(1988)318
- Lam, F.C., see Hung, C.T. 425(1988)331
- Lam, G., Williams, R.M. and Whitney, C.C.  
Electrochemical determination of oxymorphone in rat plasma by ion-pair reversed-phase HPLC 413(1987)309
- Lam, H.R. and Tarding, F.  
Specific HPLC method for the determination of creatinine in rat plasma 426(1988)358
- Lam, S., Malikin, G. and Karmen, A.  
HPLC of hydroxysteroids detected with post-column immobilized enzyme reactors 441(1988)81
- Lambert, N., Plumb, G.W. and Wright, D.J.  
Application of HPLC to the assessment of subunit heterogeneity in plant 11S storage globulins 402(1987)159
- Lambert, W.E., De Leenheer, A.P., Walgraeve, H.R. and Tassaneevakul, W.  
Monitoring of cyclosporin in whole blood by reversed-phase liquid chromatography on a butyl column 427(1988)195
- Lambert, W.E., see De Leenheer, A.P. 429(1988)3
- Lambeth, J.D., see Tribble, D.L. 414(1987)411
- Lamboeuf, Y., see Saint Blanquat, G.D. 415(1987)388
- Lamiable, D., Vistelle, R., Nguyen-Khac, M. and Millart, H.  
HPLC determination of dexamethasone in cerebrospinal fluid and plasma in the rabbit 434(1988)315
- Lamm, B., see Gyllenhaal, O. 411(1987)285
- Lampen, P., Neumeyer, J.L. and Baldessarini, R.J.  
HPLC separation and electrochemical or spectrophotometric determination of *R*(-)-*N*-*n*-propylnorapomorphine and *R*(-)-10,11-methylenedioxy-*N*-*n*-propylnorapomorphine in primate plasma 426(1988)283
- Lande, S., see Viscomi, G. 440(1988)157
- Landers, G.M. and Olson, J.A.  
Rapid, simultaneous determination of isomers of retinal, retinal oxime and retinol by HPLC 438(1988)383
- Landolfi, R., see Castagnola, M. 440(1988)231
- Landolfi, R., see De Cristofaro, R. 426(1988)376
- Lang, J.F., see Walker, B.J. 416(1987)150
- Lange, U., see Wallasch, T.M. 425(1988)175
- Lange, W., see Masihi, K.N. 440(1988)473
- Langer, K., see Florance, J. 414(1987)313
- Langlois, S., see Grech-Belanger, O. 417(1987)89
- Langseth, W.  
Separation of inorganic and organomercury chelates on a polystyrene-divinylbenzene copolymer column 438(1988)414
- Lanuzza, F., see Currò, P. 404(1987)273
- Lapalus, P., see Candito, M. 430(1988)198
- Laplanche, R., see Jean, C. 428(1988)61
- Larking, P.W.  
Quantitative GC measurement of glycosaminoglycan hexosamines in urine and plasma 420(1987)231
- Larroque, M., Vian, L., Blaise, A. and Brun, S.  
Chromatographic determination of residual monomers of epoxy resins in wine simulants (in French) 445(1988)107
- Larroque, M., Vian, L. and Blaise, A.  
Determination of phenol by HPLC with fluorimetric detection. Application to wines (in French) 407(1987)384
- Larsen, C. Grønhoj, see Jakobsen, P. 415(1987)413
- Larsen, F. Grønhoj, see Jakobsen, P. 415(1987)413

- Larsen, L.M., Nielsen, T.H., Plöger, A. and Sørensen, H.  
Rapid and efficient method for the isolation and characterization of plant aromatic choline esterases 450(1988)121
- Larson, A.A., see Igwe, O.J. 432(1988)113
- Larson, J.R., Barr, S.W. and Bredeweg, R.A.  
Flow programming system for process capillary GC 405(1987)163
- Larson, R.A., see Choi, K.E. 420(1987)81
- Larsson, A., see Jacobsson, S. 447(1988)329
- Larsson, J., see Andersson, E. 414(1987)174
- Larsson, K., Hermann, W., Möller, P. and Sanchez, D.  
Preparative HPLC of peptides on a new reversed-phase packing material, Kromasil C<sub>18</sub> 450(1988)71
- Larsson, L., see Sonesson, A. 417(1987)11
- Larsson, L., see Sonesson, A. 417(1987)366
- Larsson, L., see Sonesson, A. 431(1988)1
- Larsson, N., see Bergqvist, Y. 427(1988)295
- Laskar, S. and Basak, B.  
Detection of amino acids on thin-layer plates 436(1988)341
- Lastick, S.M., see Baker, J.O. 437(1988)387
- László, K., see Nagy, L.G. 406(1987)311
- Latini, R., Sica, A., Marchi, S., Chen, Z.M., Gavinelli, M. and Benfenati, E.  
HPLC separation and MS identification of propafenone, 5-hydroxypropafenone and N-depropylpropafenone 424(1988)211
- Lattes, A., see Dobrev, A.A. 445(1988)268
- Lau, B.P.-Y., Weber, D. and Page, B.D.  
GC-MS determination of ethyl carbamate in alcoholic beverages 402(1987)233
- Lau, C.E., Dolan, S. and Tang, M.  
Microsample determination of diazepam and its three metabolites in serum by reversed-phase HPLC 416(1987)212
- Lau, C.E., Falk, J.L., Dolan, S. and Tang, M.  
Simultaneous determination of flurazepam and five metabolites in serum by HPLC and its application to pharmacokinetic studies in rats 423(1987)251
- Lau, D., see Shulgin, A.T. 423(1987)365
- Laub, R.J., see Todorović, M. 442(1988)105
- Laufen, H., see Scharpf, F. 413(1987)91
- Lauff, J.J.  
Ion-pair HPLC procedure for the quantitative analysis of theophylline in serum samples 417(1987)99
- Lauren, D.R., Taylor, H.J. and Rahman, A.  
Analysis of the herbicides dicamba, clopyralid and bromacil in asparagus by HPLC 439(1988)470
- Laurent, M., see Inchauspe, G. 404(1987)53
- Laus, G., see De Bondt, M. 442(1988)165
- Lauterburg, B.H., see Karlaganis, G. 420(1987)171
- Lavene, D., see Humbert, H. 417(1987)319
- Lavigne, C., Zee, J.A., Simard, R.E. and Gosselin, C.  
HPLC-diode-array determination of ascorbic acid, thiamine and riboflavine in goats' milk 410(1987)201
- Law, B.  
Analysis of basic compounds on a silica column with an aqueous methanol eluent. The use of quantitative structure-retention relationships in metabolite identification 407(1987)1
- Lawrence, J.F. and Charbonneau, C.F.  
Separation of sulphite adducts by ion chromatography with oxidative amperometric detection 403(1987)379
- Lawrence, J.F. and Charbonneau, C.F.  
Direct method for the determination of 2-acetyl-4(5)-tetrahydroxybutylimidazole in caramel colours and beer by HPLC with absorbance detection 407(1987)405
- Lawrence, J.F. and Charbonneau, C.F.  
Direct, sensitive and selective detection of free fatty acids by HPLC with post-column ion-pair extraction and absorbance detection 445(1988)189
- Lawrence, J.F. and Iyengar, J.R.  
LC determination of beta-aspartame in diet soft drinks, beverage powders and pudding mixes 404(1987)261
- Lawson, D.E.M., see Okano, T. 448(1988)145
- Lázaro, F., De Castro, M.D.L. and Valcárcel, M.  
Fluorimetric determination of aflatoxins in foodstuffs by HPLC with flow injection analysis 448(1988)173
- Lazarov, L., see Stefanova, M. 446(1988)293
- Lazzarini, G., see Candiano, G. 423(1987)319
- Leah, J., see Griffin, M. 431(1988)285
- Leahey, W.J. and McMeekin, J.R.  
Determination of cyclopenthiiazide in plasma by HPLC 421(1987)401
- LeBel, M., see Croteau, D. 419(1987)205
- Le Beyec, Y., see Van, T.D. 440(1988)71
- Leboeuf, E., see Grech-Belanger, O. 417(1987)89
- Le Bonniec, B., see Ducrocq, R. 419(1987)75
- Leclercq, P.A., see Brandšterová, E. 404(1987)359
- Le Coguic, A., Bidault, R., Farinotti, R. and Dauphin, A.  
Determination of ofloxacin in plasma and urine by LC 434(1988)320



- Lecommandeur, D., MacGregor, A.W. and Daussant, J.  
Purification of germinated barley  $\alpha$ -amylase isozymes and limit-dextrinase by chromatofocusing and affinity chromatography 441(1988)436
- Lecomte, L., Triolet, J., Sinou, D., Bakos, J. and Heil, B.  
HPLC separation of sulphonated phosphines 408(1987)416
- Le Corre, P., Gibassier, D., Sado, P. and Le Verge, R.  
Simultaneous assay of disopyramide and monodesisopropylidopyramide enantiomers in biological samples by LC 424(1988)424
- Le Corre, P., Gibassier, D., Sado, P. and Le Verge, R.  
Direct enantiomeric resolution of disopyramide and its metabolite using chiral HPLC. Application to stereoselective metabolism and pharmacokinetics of racemic disopyramide in man 450(1988)211
- Lederer, M., see Fanali, S. 440(1988)361
- Lederer, M., see Kuhn, A.O. 406(1987)389
- Lederer, M., see Kuhn, A.O. 440(1988)165
- Le Dily, J., see Duchene, P. 424(1988)205
- Le Dinh, T., see Motassim, N. 422(1987)340
- Leduc, B.W., see Kennedy, J.S. 422(1987)274
- LeDuc, B.W., see Mikati, M.A. 433(1988)305
- Lee, A.L., Liao, A.W. and Horváth, C.  
Tandem separation schemes for preparative HPLC of proteins 443(1988)31
- Lee, B.L., see Ong, C.N. 423(1987)278
- Lee, D.P.  
Chromatographic evaluation of large-pore and non-porous polymeric reversed phases 443(1988)143
- Lee, E.J.D., Ang, S.B. and Lee, T.L.  
Stereoselective HPLC assay for bupivacaine enantiomers 420(1987)203
- Lee, E.J.D. and Ang, S.B.  
Simple and sensitive HPLC assay for 5-aminosalicylic acid and acetyl-5-aminosalicylic acid in serum 413(1987)300
- Lee, F.W., Watari, N., Rigod, J. and Benet, L.Z.  
Simultaneous determination of nitroglycerin and its dinitrate metabolites by capillary GC with electron-capture detection 426(1988)259
- Lee, F.Y.F., see Workman, P. 422(1987)315
- Lee, H.K., Li, S.F.Y. and Tay, Y.H.  
HPLC of eleven priority substituted phenols by isocratic elution 438(1988)429
- Lee, H.W., see Park, K.S. 440(1988)225
- Lee, J.K., Deluccia, F.J., Kelly, E.L., Davidson, C. and Borger, F.R.  
Determination of the molecular size distribution of immunoglobulin G (IgG) in intravenous IgG-albumin formulations by HPLC 444(1988)141
- Lee, M.-K., Nohta, H. and Ohkura, Y.  
Assay for dopamine  $\beta$ -hydroxylase in human plasma and rat serum by HPLC with fluorimetric detection 421(1987)237
- Lee, M., Nohta, H., Umegae, Y. and Ohkura, Y.  
Assay for tyrosine hydroxylase by HPLC with fluorescence detection 415(1987)289
- Lee, M.L., see Bradshaw, J.S. 405(1987)169
- Lee, M.L., see Fields, S.M. 406(1987)223
- Lee, P., see Hubbard, R.W. 431(1988)163
- Lee, P.D.K., see Powell, D.R. 420(1987)163
- Lee, R.M., see Doyle, E. 411(1987)325
- Lee, S.L., see Den, T.G. 408(1987)323
- Lee, S.Y., see Kopaciewicz, W. 409(1987)111
- Lee, T.L., see Lee, E.J.D. 420(1987)203
- Lee, Y.J., Radhakrishnamurthy, B., Dalferes, Jr., E.R. and Berenson, G.S.  
Fractionation of aorta glycosaminoglycans by HPLC 419(1987)275
- Lee, Y.S., see Kee, Y. 428(1988)345
- Leemann, T., see Padmalatha Devi, K. 434(1988)265
- Leenheer, A.P. De, Nelis, H.J., Lambert, W.E. and Bauwens, R.M.  
Chromatography of fat-soluble vitamins in clinical chemistry (Review) 429(1988)3
- Leer, E.W.B. De, see Erkelens, C. 404(1987)67
- Leeuw, J.W. De, see Sinnighe Damsté, J.S. 435(1988)435
- Lefeber, D.S., see Riggin, R.M. 435(1988)307
- Lefebvre, M.A., see Mignot, A. 430(1988)192
- Lefebvre, P., see Sablonniere, B. 403(1987)183
- Le Gall, J.-M., see Gallice, P. 422(1987)263
- Le Garrec, L., see Delee, E. 450(1988)183
- Le Garrec, L., see Delee, E. 450(1988)191
- Legendre, F., see Campeau, D. 405(1987)305
- Leger, C.L., see Alessandri, J.-M. 432(1988)75
- Lehmann, A., see Zobel, H. 446(1988)305
- Lehotay, J., see Brandšterová, E. 404(1987)359
- Lehoux, J.-G., see Bouthillier, F. 403(1987)171
- Lehr, K.-H. and Damm, P.  
Quantification of the enantiomers of ofloxacin in biological fluids by HPLC 425(1988)153
- Lehrfeld, J.  
Simultaneous GLC determination of aldoses and alduronic acids 408(1987)245
- Leitao, J., De Saint Blanquat, G., Bailly, J.R. and Paillas, C.  
Quantitation of aflatoxins from various

- strains of *Aspergillus* in foodstuffs  
435(1988)229
- Leitner, B., see Schott, H. 441(1988)115
- Lekova, K., Bursukova, M., Ivanichkova, S.,  
Bojkova, N. and Petkova, R.  
Polyester deactivation of pink supports and  
its effect on the GC of some polar, closely  
boiling and isomeric analytes 446(1988)31
- Lekova, K., Radulova, M., Bursukova, M. and  
Rasheva, E.  
GLC analysis of isophorone byproducts  
446(1988)23
- Le Liboux, A., Frydman, A. and Gaillot, J.  
Simultaneous determination of zopiclone and  
its two major metabolites (N-oxide and N-  
desmethyl) in human biological fluids by  
reversed-phase HPLC 417(1987)151
- Lelo, A., Birkett, D.J. and Miners, J.O.  
Radiometric HPLC procedure for the  
determination of theobromine metabolites in  
microsomal incubations 430(1988)203
- Lemaire, S., Dumont, M. and Nolet, S.  
Sensitive method of detection, quantitation  
and purification of peptides using pre-column  
derivatization with phenyl isothiocyanate  
425(1988)77
- Lemberkovic, E. and Petri, G.  
GC characterization of frequently occurring  
aromatic compounds in essential oils  
446(1988)267
- Lemmens, A.A.G., Everaerts, F.M., Venema,  
J.W. and Jonker, H.D.  
Isotachophoretic determination of the  
cationic constituents in mixtures of raw  
materials for glass manufacturing  
439(1988)423
- Lemmens, P.J.M.R., see Engels, W. 427(1988)209
- Lemonnier, F., see Therasse, J. 413(1987)237
- Lennard, L.  
Assay of 6-thioinosinic acid and 6-  
thioguanine nucleotides, active metabolites of  
6-mercaptopurine, in human red blood cells  
423(1987)169
- Lenngren, S., see Johansson, M. 432(1988)65
- Lenngren, S., see Johansson, M. 432(1988)243
- Lepore, J., see Dong, M.W. 442(1988)81
- Lepri, L., Desideri, P.G. and Coas, V.  
HPTLC of dinitropyridyl-amino acids and  
sulphonamides on silanized silica gel  
405(1987)394
- Leray, C., Pelletier, X., Hemmendinger, S. and  
Cazenave, J.-P.  
TLC of human platelet phospholipids with  
fatty acid analysis 420(1987)411
- Leroux, F., see Sioufi, A. 450(1988)221
- Leroux, J.-P., see Beaune, P.H. 426(1988)169
- Leroy, Y., see Capon, C. 425(1988)35
- Lesieur, M., see Guichardant, M. 425(1988)25
- Leskawa, K.C., Schulte, B.A. and Hogan, E.L.  
*In situ* visualization of glycosphingolipids on  
thin-layer chromatograms using lectin-  
peroxidase conjugates 411(1987)393
- Lesko, L.J., see Emm, T. 419(1987)445
- Lesko, L.J., see Emm, T. 427(1988)188
- Lesko, L.J., see Emm, T.A. 427(1988)162
- Leslie, J., see Emm, T. 427(1988)188
- Leslie, J., see Emm, T.A. 427(1988)162
- Lesne, M., see Wallemacq, P.E. 413(1987)131
- Lesne, M. and Pellegrin, P.L.  
Rapid HPLC method for the determination  
of amiodarone and desethylamiodarone in  
human plasma and serum 415(1987)197
- Lesser, J.H., Shustina, R. and Ovadia, D.  
LC behaviour of chlorinated carbanilides  
410(1987)95
- Lester, D.C., see Nierenberg, D.W. 413(1987)79
- Leuenberger, C., see Czuczwa, J. 403(1987)233
- Leung, C.P. and Au-Yeung, C.Y.  
HPLC determination of loperamide  
hydrochloride in pharmaceutical preparations  
449(1988)341
- Leven, W. and Willuhn, G.  
Sesquiterpene lactones from *Arnica  
chamissonis* Less. VI. Identification and  
quantitative determination by HPLC and GC  
410(1987)329
- Le Vent, S., see Akporhonor, E.E. 405(1987)67
- Le Verge, R., see Le Corre, P. 424(1988)424
- Le Verge, R., see Le Corre, P. 450(1988)211
- Leverge, R., see Sado, P.A. 434(1988)157
- Levillain, P., see Montagu, M. 409(1987)426
- Levy, G., see Klockowski, P.M. 422(1987)334
- Levy, L.A., see Chae, K. 439(1988)484
- Lew, M., see Gupta, R.N. 425(1988)441
- Lewicki, J.A., see Carilli, C.T. 444(1988)203
- Lewis, C.A., see Philp, R.P. 446(1988)3
- Lewis, J.J., see Ravichandran, K. 439(1988)213
- Lewis, S., see Wilson, I.D. 408(1987)445
- Lhermitte, L., see Broly, F. 431(1988)369
- Li, E.X., see Cheam, V. 450(1988)361
- Li, F., see Lim, C.K. 429(1988)123
- Li, G. and Liu, C.  
Chromatographic study of charge-transfer  
complexes 406(1987)353
- Li, H.-P.  
Capillary GC of neutral sugars as their  
aldononitrile acetates from the hydrolyzate of  
corn bran residues 410(1987)484
- Li, P.-L., see Gielsdorf, W. 419(1987)357

- Li, S.F.Y., see Lee, H.K. 438(1988)429
- Li, Z., see Grob, K. 448(1988)387
- Liang, W., see Zeng, W. 408(1987)426
- Liao, A.W., see Lee, A.L. 443(1988)31
- Liao, J.-L., see Hjertén, S. 403(1987)47
- Libersa, C., see Broly, F. 431(1988)369
- Liboux, A. Le, Frydman, A. and Gaillot, J.  
Simultaneous determination of zopiclone and its two major metabolites (N-oxide and N-desmethyl) in human biological fluids by reversed-phase HPLC 417(1987)151
- Lichtenberger, A., see Gielsdorf, W. 419(1987)351
- Lichý, A., see Macek, J. 434(1988)423
- Liebich, H.M., see Krämer, B.K. 427(1988)351
- Liebich, H.M., see Röss, K. 417(1987)465
- Lienne, M., Caude, M., Rosset, R. and Tambute, A.  
Optimization of the direct chiral separation of potential cytotoxic  $\alpha$ -methylene- $\gamma$ -butyrolactones and  $\alpha$ -methylene- $\gamma$ -butyrolactams by LC 448(1988)55
- Lierop, J.B.H. Van and Van Veen, R.M.  
Determination of plasticizers in fat by GC-MS 447(1988)230
- Liesch, J.M., see Metz, P.A. 441(1988)31
- Ligny, C.L. De, see Brandts, P.M. 437(1988)337
- Ligny, C.L. De, see Brandts, P.M. 438(1988)181
- Likhite, S.M., see Panse, D.G. 411(1987)456
- Lille, Ü., see Löhmus, M. 449(1988)77
- Lille, Ü., see Löhmus, M. 450(1988)105
- Lillig, B., see Velayudhan, A. 435(1988)397
- Lim, C.K., Li, F. and Peters, T.J.  
HPLC of porphyrins (Review) 429(1988)123
- Lim, H.K., see Roy, S.D. 431(1988)210
- Lim, J.K.C., see Hung, C.T. 425(1988)331
- Lim, K.T., see Yim, J.J. 419(1987)296
- Limero, T., see Wentworth, W.E. 441(1988)45
- Lin, B., see Lin, B. 438(1988)171
- Lin, B., Wang, J. and Lin, B.  
Influence of the partition isotherm on the elution curve 438(1988)171
- Lin, B.C., see Koppenhoefer, B. 441(1988)89
- Lin, E.T., see Gawienowski, M. 430(1988)412
- Lin, E.T., see La Follette, G. 431(1988)379
- Lin, G.-C. and Chang, C.A.  
Heavy alkylated benzene stationary phase for the GLC separation of ethylbenzene, xylenes, styrene, iso- and *n*-propylbenzene 409(1987)371
- Lin, G. and Chang, C.A.  
Heavy alkylated benzene stationary phase for the GLC separation of light hydrocarbons 404(1987)328
- Lin, H.S., see Tian, Z.W. 439(1988)151
- Lin, H.S., see Tian, Z.W. 439(1988)159
- Lin, J.H., see Los, L.E. 437(1988)442
- Lin, L.-J. and Shiao, M.-S.  
Separation of oxygenated triterpenoids from *Ganoderma lucidum* by HPLC 410(1987)195
- Lin, N.T., see Tice, P.A. 410(1987)43
- Lin, W.-N., see Cheung, S.-T. 414(1987)248
- Linari, F., see Mentasti, E. 417(1987)253
- Linari, F., see Petrarulo, M. 432(1988)37
- Lindberg, C., see Edholm, L.-E. 424(1988)61
- Lindberg, M., see Alfredsson, G. 424(1988)378
- Lindeberg, J., see Hjertén, S. 440(1988)305
- Lindemans, J., Van Kapel, J., Arwert, F. and Abels, J.  
Separation of transcobalamin II isoproteins by means of chromatofocusing 432(1988)103
- Linden, P.J.Q.V. Der, see Gladdines, M.M. 431(1988)317
- Lindgren, K.  
Determination of cefadroxil in serum by HPLC with cephradine as internal standard 413(1987)347
- Lindgren, K.  
Determination of cefaclor and cephradine in serum by ion-pair reversed-phase chromatography 413(1987)351
- Lindner, H., Helliger, W. and Puschendorf, B.  
Separation of Friend erythroleukaemic cell histones and high-mobility-group proteins by reversed-phase HPLC 450(1988)309
- Lindsay, R.M., McLaren, A.M. and Baty, J.D.  
Reversed-phase HPLC assay for the determination of the *in vitro* acetylation of *p*-aminobenzoic acid by human whole blood 433(1988)292
- Lindström, B., Sjöberg, P. and Floberg, S.  
GC-MS method for the determination of dimethylacetamide and metabolites in whole blood 428(1988)156
- Ling, D.-K., see Yang, Q. 447(1988)208
- Lingeman, H., see Reeuwijk, H.J.E.M. 428(1988)93
- Lingeman, H., see Tjaden, U.R. 420(1987)53
- Linnoila, M., see Gusovsky, F. 415(1987)124
- Linscheid, M., see Blaszkewicz, M. 439(1988)109
- Linssen, P.C.M., see Brons, P.P.T. 422(1987)175
- Linton, C., see Bauer, J. 445(1988)429
- Lipinski, J. and Stan, H.-J.  
CAPA—computer aided pesticide analysis. Computer program for the automated evaluation of chromatographic data for residue analysis of foods 441(1988)213
- Lisichkin, G.V., see Gaida, A.V. 424(1988)385
- Liu, C., see Li, G. 406(1987)353
- Liu, D. and Desiderio, D.M.  
Reversed-phase HPLC-radioreceptor assay of

- human cerebrospinal fluid neuropeptides.  
Maximizing recovery of picomoles of peptides and minimizing memory 422(1987)61
- Liu, G.  
Oxidation of thiols within GC columns 441(1988)372
- Liu, G., Luo, X. and Zhang, S.  
Determination of sulphur compounds at low parts per billion levels by GC with flame photometric detection 435(1988)327
- Lloyd, J.B.F. and Parry, D.A.  
Detection and determination of common benzodiazepines and their metabolites in blood samples of forensic science interest. Microcolumn cleanup and HPLC with reductive electrochemical detection at a pendent mercury drop electrode 449(1988)281
- Lloyd, L.L., Warner, F.P., Kennedy, J.F. and White, C.A.  
Ion suppression reversed-phase HPLC method for the separation of L-ascorbic acid in fresh fruit juice 437(1988)447
- Luis, F., see Draviam, E.J. 414(1987)180
- Lochmüller, C.H. and Mink, L.P.  
Adsorption isotherms of ethyl acetate modifier on silica from supercritical carbon dioxide 409(1987)55
- Locke, D.C., see Schneiderman, M.A. 409(1987)343
- Lockett, L., see Sitaram, B.R. 422(1987)13
- Locniskar, A. and Greenblatt, D.J.  
HPLC assay of adinazolam and its desmethyl metabolite 424(1988)215
- Lodewijks, M.T.M., see Van Harten, J. 423(1987)327
- Lodge, B., see Flann, B. 402(1987)273
- Lodkowski, R., see Supryniewicz, Z. 446(1988)347
- Lohkamp, R., see Weber, C. 426(1988)345
- Löhmus, M., Kirjanen, I., Lopp, M. and Lille, Ü.  
Solvent selectivity in the resolution of some regioisomeric and diastereomeric prostaglandin intermediates on silica 449(1988)77
- Löhmus, M., Lopp, M., Lille, Ü. and Veisserik, J.  
Preparative LC separation of isomers of prostacyclin carba-analogues and their intermediates 450(1988)105
- Lokensgard, D.M., see Kern, J.R. 450(1988)233
- Lokiec, F., see Brossat, B. 413(1987)141
- Long, D., see Krstulović, A.M. 411(1987)461
- Long, J.W., Snyder, R.W. and Spalik, J.  
Purge-and-trap GC method for residual volatiles in high-temperature polymers 450(1988)394
- Look, Z., see Achari, R. 424(1988)430
- Look, Z.M., see Falkowski, A.J. 422(1987)145
- Lopez-Anaya, A. and Mayersohn, M.  
Quantification of riboflavin, riboflavin 5'-phosphate and flavin adenine dinucleotide in plasma and urine by HPLC 423(1987)105
- López-Soriano, F.J., see Peinado, J. 415(1987)372
- Lopp, M., see Löhmus, M. 449(1988)77
- Lopp, M., see Löhmus, M. 450(1988)105
- Lora-Tamayo, C., Tena, T. and Tena, G.  
Concentrations of free and conjugated morphine in blood in twenty cases of heroin-related deaths 422(1987)267
- Lorreyne, W., see Woestenborghs, R. 413(1987)332
- Lorreyne, W., see Woestenborghs, R. 424(1988)195
- Los, L.E., Lin, J.H. and Ulm, E.H.  
Quantitative determination of L-364,718, a novel cholecystokinin antagonist, by solid-phase extraction and HPLC 437(1988)442
- Löser, C., Wunderlich, U. and Fölsch, U.R.  
Reversed-phase LC separation and simultaneous fluorimetric detection of polyamines and their monoacetyl derivatives in human and animal urine, serum and tissue samples: an improved, rapid and sensitive method for routine application 430(1988)249
- Lough, W.J., see Matlin, S.A. 450(1988)450(1988)157
- Louis, R. St., see Gholson, A.R. 408(1987)329
- Lounasmaa, M., see Naaranlahti, T. 410(1987)488
- Loupy, A., see Delee, E. 450(1988)183
- Louw, M.E.J., see Shephard, G.S. 416(1987)138
- Lovett, L.J., see Nygard, G.A. 415(1987)438
- Lövkvist, P., see Jönsson, J.Å. 408(1987)1
- Lowe, C.R., see Burton, S.J. 435(1988)127
- Loyola, E., see Reglero, G. 438(1988)243
- Lu, H.S., Klein, M.L. and Lai, P.-H.  
Narrow-bore HPLC of phenylthiocarbamyl amino acids and carboxypeptidase P digestion for protein C-terminal sequence analysis 447(1988)351
- Lu, P.C., see Koppenhoefer, B. 441(1988)89
- Lucarelli, C., see Marzo, A. 427(1988)345
- Lucas, M., see Maggio, A.-F. 436(1988)23
- Lucock, M., see Hartley, R. 415(1987)357
- Łuczyński, J., see Kolarz, B.N. 408(1987)308
- Lüderitz, O., see Galanos, C. 440(1988)397
- Ludwig, B., see Claycamp, H.G. 422(1987)239
- Ludwig, B., see Heizmann, P. 415(1987)170
- Lugnier, A.A., see Kintz, P. 432(1988)329
- Lugnier, A.A., see Tracqui, A. 432(1988)412
- Luhmann, I., see Szathmary, S.C. 425(1988)193
- Lumley, I.D., Patel, I. and Cohen, I.C.

- Evaluation of laboratory-prepared high-performance size-exclusion columns.  
Applications to food analysis 408(1987)115
- Lumpkin, R.H., see Souther, M.E. 417(1987)27
- Lundahl, P. and Mascher, E.  
Use of term "detergent" in biochemistry 449(1988)
- Luo, X.-D., see Zou, A.-Q. 410(1987)217
- Luo, X., see Liu, G. 435(1988)327
- Luque, J., see Ferrer, E. 411(1987)167
- Luque, J., see Ferrer, E. 445(1988)327
- Luque, J., see García-Pérez, A.I. 403(1987)131
- Lurie, I.S., Moore, J.M., Cooper, D.A. and Kram, T.C.  
Analysis of manufacturing by-products and impurities in illicit cocaine via HPLC and photodiode array detection 405(1987)273
- Lurie, I.S., see Moore, J.M. 410(1987)297
- Lustenberger, P., see Maes, P. 445(1988)409
- Lutz, S., see König, W.A. 447(1988)193
- Lutz, S., see Rieck, M. 439(1988)301
- Lutz, T., see Jindal, S.P. 408(1987)356
- Luzzani, F., see Corti, F. 424(1988)147
- Lydon, J., Duke, S.O. and Hedin, P.A.  
Coextraction of 5-(hydroxymethyl)-2-furaldehyde with phenolic acids in acid-hydrolyzed plant extracts 410(1987)474
- Lyman, M., see Stafford, A.E. 436(1988)93
- Lynn, B.C. and Alley, E.G.  
Derivatization of diaminocyclohexanes for determination of *cis* and *trans* isomers by GC 408(1987)211
- Lynn, R.K., see Garvie, C.T. 413(1987)43
- Lyoo, Y.S., see Stowers, D.J. 444(1988)47
- Lysjuk, L.S. and Chuiko, A.A.  
Effect of zirconia on the sorption properties of silica gels 407(1987)189
- Mabuchi, H. and Nakahashi, H.  
Determination of 3-carboxy-4-methyl-5-propyl-2-furanpropanoic acid, a major endogenous ligand substance in uremic serum, by HPLC with UV detection 415(1987)110
- Macaudiere, P., Caude, M., Rosset, R. and Tambuté, A.  
Resolution of racemic amides and phosphine oxides on a  $\beta$ -cyclodextrin-bonded stationary phase by subcritical fluid chromatography 405(1987)135
- Macaudière, P., Caude, M., Rosset, R. and Tambuté, A.  
Chiral resolution of a series of 3-thienylcyclohexylglycolic acids by liquid or subcritical fluid chromatography. A mechanistic study 450(1988)255
- MacCrehan, W.A. and Schönberger, E.  
Reversed-phase HPLC separation and electrochemical detection of retinol and its isomers 417(1987)65
- MacCroskey, M.C., see Pearson, J.D. 418(1987)245
- Macek, J., Krájičková, J. and Adam, M.  
Rapid determination of disaccharides from chondroitin and dermatan sulphates by HPLC 414(1987)156
- Macek, J., Lichý, A., Tesařová, E. and Adam, M.  
Separation of type IX collagen from other cartilage collagens by hydrophobic interaction chromatography 434(1988)423
- Macek, J. and Vácha, J.  
Rapid and sensitive method for determination of piroxicam in human plasma by HPLC 420(1987)445
- Macek, K., see Deyl, Z. 426(1988)162
- MacGee, C.M., see Dawson, M. 420(1987)129
- MacGrath, S.C., see Choi, K.E. 420(1987)81
- MacGregor, A.W., see Lecommandeur, D. 441(1988)436
- Mack, G.D., see Knight, M. 444(1988)345
- Mackey, D.J. and Higgins, H.W.  
Reversed-phase chromatographic separation and analysis of marine metal-organic complexes 436(1988)243
- MacKinnon, A.M., see Clifton, P.M. 414(1987)25
- Mac Mathúna, P., see O'Reilly, T. 417(1987)190
- MacQuarrie, R.A., see Smith, R.E. 439(1988)83
- Macrae, P.V., see Beresford, A.P. 420(1987)178
- Maddock, J., see Robinson, P.R. 432(1988)153
- Madelmont, J.C., see Labarre, P. 419(1987)381
- Mady, N., see Freitag, W. 450(1988)426
- Maeda, K., see Niwa, N.T.T. 431(1988)418
- Maeda, O., see Yoshikawa, S. 421(1987)111

## M

- Ma, J.K.H., see Mehta, R.C. 430(1988)341
- Maa, Y.-F. and Horváth, C.  
Rapid analysis of proteins and peptides by reversed-phase chromatography with polymeric micropellicular sorbents 445(1988)71
- Maack, F. De, see Guichardant, M. 425(1988)25
- Maaten, M.J. Van der, see Schmerr, M.J.F. 425(1988)379

- Maeda, Y., Yamamoto, M., Owada, K., Sato, S. and Masui, T.  
HPLC determination of six *p*-hydroxybenzoic acid esters in cosmetics using Sep-Pak Florisil cartridges for sample pre-treatment 410(1987)413
- Maeden, F.P.B. Van der, see Broersen, J.J. 436(1988)39
- Maekawa, M.  
Lactate dehydrogenase isoenzymes (Review) 429(1988)373
- Maertens, F., see Delbeke, F.T. 426(1988)194
- Maes, P., Formstecher, P., Lustenberger, P. and Dautrevaux, M.  
HPLC in the evaluation of the lipophilicity of 17 $\beta$ -carboxamide steroid derivatives 445(1988)409
- Maessen, F.J.M.J., see De Waal, W.A.J. 407(1987)253
- Maessen, P.A., Mross, K.B., Pinedo, H.M. and Van der Vijgh, W.J.F.  
Improved method for the determination of 3'-epidoxorubicin and seven metabolites in plasma by HPLC 417(1987)339
- Maessen, P.A., Pinedo, H.M., Mross, K.B. and Van der Vijgh, W.J.F.  
New method for the determination of doxorubicin, 4'-epidoxorubicin and all known metabolites in cardiac tissue 424(1988)103
- Magerovskii, Y.V., see Gaida, A.V. 424(1988)385
- Maggio, A.-F., Pompon, A., Lucas, M., Barascut, J.-L. and Imbach, J.-L.  
DNA-nitrosourea interactions. HPLC of crosslinked dinucleosides and substituted deoxynucleosides 436(1988)23
- Maglietto, S.M., see Musson, D.G. 414(1987)121
- Magnus, I.A., see Sanitrák, J. 415(1987)129
- Mahara, R., see Tohma, M. 421(1987)9
- Mahdi, A.A., see Ahmad, S. 440(1988)467
- Mahmudov, A.U., see Zaslavsky, B.Y. 403(1987)123
- Mahmudov, A.U., see Zaslavsky, B.Y. 439(1988)267
- Mahonen, A., see Karhu, I. 426(1988)65
- Mahuzier, G., see Bossant, M.J. 423(1987)23
- Mai, W., see Riis, V. 445(1988)285
- Maibaum, J.  
Indirect HPLC resolution of racemic tertiary amines as their diastereomeric urea derivatives after N-dealkylation 436(1988)269
- Mäkelä, M., Mattsson, P., Pintamo-Kenttälä, K. and Korpela, T.  
Cyclodextrin-ligand interaction as a simplified model of biospecific affinity chromatography 448(1988)391
- Mäkelä, M., see Mattsson, P. 447(1988)398
- Makino, I., see Tohma, M. 421(1987)9
- Makita, M., see Kataoka, H. 420(1987)135
- Makita, M., see Kataoka, H. 436(1988)67
- Malcolme-Lawes, D.J., see Hill, E. 441(1988)394
- Malcolme-Lawes, D.J., see Jones, K. 441(1988)387
- Malfait, R., see Gerlo, E. 414(1987)301
- Malikin, G., see Lam, S. 441(1988)81
- Mallet, A.I., McLaughlin-Flynn, C.A. and Camp, R.D.R.  
MS analysis of endogenous diacylglycerols in normal and psoriatic skin 432(1988)278
- Mallett, D.N., Gulaid, A.A. and Dennis, M.J.  
HPLC method with electrochemical detection for the concomitant assay of sulphadiazine, sulphamerazine and sulphapyridine in plasma 428(1988)190
- Maloney, J.A., see Awni, W.M. 425(1988)233
- Maloney, T., see Wilson, T.D. 445(1988)299
- Maltby, N., see Richmond, R. 417(1987)241
- Maltsev, V.G., Zimina, T.M., Khvatov, A.B. and Belenkii, B.G.  
Determination of myoglobin in human serum by HPLC with chemiluminescence detection 416(1987)45
- Maltzev, V.G., see Tennikova, T.B. 435(1988)357
- Mama, J.E., see Fell, A.F. 434(1988)377
- Manabe, T., Higuchi, N., Okuyama, T. and Mukaiyama, Y.  
High-performance affinity chromatography of human serum concanavalin A binding proteins 431(1988)45
- Manabe, T., Higuchi, N. and Okuyama, T.  
Staining of serum concanavalin A-binding proteins after micro two-dimensional electrophoresis and blotting 423(1987)115
- Manca, F., see Biondi, P.A. 411(1987)275
- Mancuso, G., see Del Principe, D. 419(1987)329
- Mandys, F., see Knobloch, E. 428(1988)255
- Mane, M.V., see Panse, D.G. 411(1987)456
- Manera, E., see Pietta, P. 404(1987)279
- Manes, L.V., see Kern, J.R. 450(1988)233
- Mangani, F. and Ninfali, P.  
GC determination of acetaldehyde and acetone in human blood by purge and trap, using permeation tubes for calibration 437(1988)294
- Mangia, A., see Casoli, A. 447(1988)187
- Mangin, P., see Kintz, P. 432(1988)329
- Mangin, P., see Tracqui, A. 432(1988)412
- Man in 't Veld, A.J., see Boomsma, F. 427(1988)219

- Maniwa, M., see Marunaka, T. 420(1987)43
- Maniwa, M., see Marunaka, T. 420(1987)329
- Maniwa, M., see Marunaka, T. 422(1987)227
- Maniwa, M., see Marunaka, T. 431(1988)87
- Maniwa, M., see Marunaka, T. 433(1988)177
- Maniwa, M., see Umeno, Y. 434(1988)123
- Maniwa, M., Umeno, Y. and Marunaka, T.  
Determination of a new hypolipidemic agent, 2-(*trans*-4-isobutylcyclohexyl)-2-oxoethyl benzenesulphonate, and its metabolite by HPLC and GC-mass fragmentography in human biological materials 433(1988)282
- Manja, K.S. and Kaul, R.K.  
GC detection of bacteria causing enteric fever 415(1987)377
- Mann, B.  
Identification of carbohydrate gums in dental cream by TLC 407(1987)369
- Manoilov, S.E., Komov, V.P., Kirillova, N.V., Strelkova, M.A., Samolin, Y.A., Chernoborodova, L.S. and Nicolaeva, L.A.  
Chromatographic isolation of primary and secondary metabolites from plant cell cultures 440(1988)53
- Mansmann, Jr., H.C., see Kester, M.B. 416(1987)91
- Mant, C.T., see Hearn, M.T.W. 438(1988)197
- Mant, C.T., see Hodges, R.S. 444(1988)349
- Mant, C.T. and Hodges, R.S.  
On-line derivatization of silica supports for regeneration and preparation of reversed-phase columns 409(1987)155
- Manzini, S., see Agostini, O. 433(1988)288
- Marangella, M., see Mentasti, E. 417(1987)253
- Marangella, M., see Petrarulo, M. 432(1988)37
- Marchand, J., see Tharasse-Bloch, C. 421(1987)407
- Marchelli, R., see Armani, E. 441(1988)275
- Marchelli, R., see Armani, E. 441(1988)287
- Marchetti, M., see Di Corcia, A. 405(1987)357
- Marchi, S., see Latini, R. 424(1988)211
- Marcomini, A., Capri, S. and Giger, W.  
Determination of linear alkylbenzenesulphonates, alkylphenol polyethoxylates and nonylphenol in waste water by HPLC after enrichment on octadecylsilica 403(1987)243
- Marden, M., see Baudin-Chich, V. 437(1988)193
- Mardmomen, S., see Mehta, R.C. 430(1988)341
- Mardon, M., see Villanueva, V.R. 440(1988)261
- Mareš, P., see Řezanka, T. 409(1987)390
- Marfil, F., see Sioufi, A. 414(1987)131
- Marhuenda, E., Menendez, M. and Alarcon De La Lastra, C.  
Constituents of essential oil of *Thymus carnosus* Boiss. 436(1988)103
- Marijnen, Y.M.T., see De Korte, D. 415(1987)383
- Marini Bettolo, G.B., see Castagnola, M. 440(1988)231
- Marino, A., see Laganá, A. 417(1987)135
- Marinucci, M.C., see Amicosante, G. 403(1987)366
- Maris, F.A., Nijenhuis, M., Frei, R.W., De Jong, G.J. and Brinkman, U.A.T.  
On-line post-column extraction in column LC with electron-capture detection. II. Applications 435(1988)297
- Maris, F.A., Stäb, J.A., De Jong, G.J. and Brinkman, U.A.T.  
On-line trace enrichment on a reversed-phase pre-column for normal-phase LC with electron-capture detection 445(1988)129
- Mark, E.J. Van der, see Te Koppele, J.M. 427(1988)67
- Markides, K.E., see Bradshaw, J.S. 405(1987)169
- Markides, K.E., see Fields, S.M. 406(1987)223
- Marko, V.  
Determination of stobadine, a novel cardioprotective drug, using capillary GC with nitrogen-phosphorus detection after its selective solid-phase extraction from serum 433(1988)269
- Marko-Varga, G.  
HPLC separation of some mono- and disaccharides with detection by a post-column enzyme reactor and a chemically modified electrode 408(1987)157
- Markuszewski, R., see Chriswell, C.D. 405(1987)213
- Marletta, G.P., see Zonta, F. 403(1987)207
- Maroldo, S.G., see Abraham, M.H. 409(1987)15
- Marra, M., Ballio, A. and Aducci, P.  
Immunoaffinity chromatography of fusicoccin 440(1988)47
- Marrakchi, N., see Créminon, C. 440(1988)439
- Marriott, P.J. and Lai, Y.-H.  
Capillary column GC method for the study of dynamic intramolecular interconversion behaviour 447(1988)29
- Marrs, W.M., see Sworn, G. 403(1987)307
- Marsaioli, A.J., see Wenkert, E. 440(1988)449
- Marsenille, M. Van, see De Bondt, M. 442(1988)165
- Marshall, T., see Gusovsky, F. 415(1987)124
- Marston, A., Borel, C. and Hostettmann, K.  
Separation of natural products by centrifugal partition chromatography 450(1988)91
- Marston, A., Potterat, O. and Hostettmann, K.

- Isolation of biologically active plant constituents by LC 450(1988)3
- Marston, A., see Slacanin, I. 448(1988)265
- Mårtensson, E., see Svensson, C. 432(1988)363
- Mårtensson, J.  
Method for determination of free and total glutathione and  $\gamma$ -glutamylcysteine concentrations in human leukocytes and plasma 420(1987)152
- Mårtensson, J., see Andersson, E. 414(1987)174
- Marti, A.R., see Brügger, R.R. 440(1988)197
- Martijn, A., see Schreuder, R.H. 435(1988)73
- Martin, B.M., see Mussar, K.J. 408(1987)378
- Martin, C., see Birk, I.M. 449(1988)423
- Martin, C.F., Takahashi, L.T., Worsley, J.L., Hagemeyer, C.J. and Hall, L.K.  
Analysis of sodium cefazolin in ophthalmic ointment by HPLC 402(1987)376
- Martin, J.F., see Kitzman, J.V. 437(1988)306
- Martin, M., see Caprioli, R.M. 443(1988)355
- Martin, M., see Donnelly, D.M.X. 440(1988)65
- Martin, M., see Ferrer, E. 411(1987)167
- Martin, M., see Ferrer, E. 445(1988)327
- Martin, Jr., R.A., see Hurst, W.J. 409(1987)413
- Martin, S.A., see Tomašić, J. 440(1988)405
- Martineau, A. and Falardeau, P.  
Purification and quantitative analysis of urinary prostanoids in human and in rat by packed and capillary GC-negative-ion chemical-ionization MS 417(1987)1
- Martinet, N., see Beninati, S. 443(1988)329
- Martínez, I., see García-Raso, A. 402(1987)323
- Martini, I.A., see Van der Steege, G. 415(1987)1
- Martire, D.E.  
Molecular theory of selectivity based on solute structure for nematic liquid-crystalline phases in GLC 406(1987)27
- Martire, D.E., see Stalcup, A.M. 442(1988)1
- Martucci, C., see Florance, J. 414(1987)313
- Marudov, G., see Neicheva, A. 437(1988)249
- Marunaka, T., Maniwa, M., Matsushima, E., Yoshida, K.-I., Azuma, R., Kurotori, M. and Komatsu, S.  
HPLC determination of 6-amidino-2-naphthyl [4-(4,5-dihydro-1*H*-imidazol-2-yl)amino]benzoate dimethanesulphonate and its metabolites in biological fluids 433(1988)177
- Marunaka, T., Maniwa, M., Matsushima, E. and Minami, Y.  
HPLC determination of a new  $\beta$ -lactamase inhibitor and its metabolite in combination therapy with piperacillin in biological materials 431(1988)87
- Marunaka, T., Matsushima, E. and Maniwa, M.  
Determination of cefodizime in biological materials by HPLC 420(1987)329
- Marunaka, T., see Maniwa, M. 433(1988)282
- Marunaka, T., see Umeno, Y. 434(1988)123
- Marunaka, T., Umeno, Y., Minami, Y., Matsushima, E., Maniwa, M., Yoshida, K. and Nagamachi, M.  
GC-mass fragmentographic determination of propiverine and its metabolites in plasma and urine 420(1987)43
- Marunaka, T. and Maniwa, M.  
HPLC determination of [3,4-di-(4-methoxyphenyl)-5-isoxazolyl]acetic acid and its metabolites in human plasma and urine 422(1987)227
- Marzo, A. and Lucarelli, C.  
GC evaluation of bufloxedil in biological samples employing the thermoionic specific detector 427(1988)345
- Marzouki, Z.M.H., Taha, A.M. and Gomaa, K.S.  
Fatty acid profiles of sebaceous triglycerides by capillary GC with mass-selective detection 425(1988)11
- Mascher, E., see Lundahl, P. 449(1988)
- Mascher, H.  
HPLC determination of dextrorphan and 3-hydroxymorphinan in human plasma based on a selective pre-column sample clean-up 420(1987)217
- Maseda, C., see Matsubara, K. 424(1988)49
- Masia, P., see Fanali, S. 403(1987)388
- Masia, P., see Fanali, S. 440(1988)361
- Masihi, K.N., Lange, W. and Özel, M.  
Biomedical applications of chromatographic fraction containing trehalose dimycolate in squalane emulsion (Review) 440(1988)473
- Maskarinec, M.P., see Row, K.H. 409(1987)193
- Maslow, K.I., see Semyonov, S.N. 446(1988)151
- Mason, G., see Rathor, M.N. 404(1987)33
- Mason, G., see Rathor, M.N. 445(1988)37
- Masoud, A.N., see Abou-Khalil, S. 417(1987)111
- Massart, D.L., see De Smet, M. 410(1987)77
- Massart, D.L., see De Smet, M. 416(1987)375
- Massart, D.L., see Musch, G. 432(1988)209
- Massey, I.J., see Wu, A.T. 415(1987)65
- Massil, S.E., Ezra, I. and Grushka, E.  
Determination of amidines by reversed-phase ion-pair HPLC 439(1988)476
- Masuda, H., see Mihara, S. 402(1987)309
- Masui, T., see Maeda, Y. 410(1987)413
- Masuko, F., see Sakamoto, M. 411(1987)259
- Matas, R., see Escribano, J. 444(1988)165
- Matejovič, I. and Polonský, J.  
Capillary isotachopheresis in the study of the



- absorption of inorganic ions by roots of intact plants 438(1988)454
- Mateo, R., Bosch, F., Pastor, A. and Jimenez, M. Capillary column GC identification of sugars in honey as trimethylsilyl derivatives 410(1987)319
- Matern, U., see Birk, I.M. 449(1988)423
- Mathe, G., see Labarre, P. 419(1987)381
- Matheou, D., see Selinger, K. 434(1988)259
- Mathes, L.E., Muschik, G., Demby, L., Polas, P., Mellini, D.W., Issaq, H.J. and Sams, R. HPLC determination of 2',3'-dideoxycytidine and 3'-azido-3'-deoxythymidine in plasma using a column-switching technique 432(1988)346
- Matheson, T.W., see Harvey, T.G. 435(1988)193
- Mathiasson, L., see Dalene, M. 435(1988)469
- Mathiasson, L., see Skarping, G. 435(1988)453
- Mathiesen, D.A., see Frank, S.K. 419(1987)225
- Mathis, D.F. and Budd, R.D. Extraction of acetaminophen and theophylline from post-mortem tissues and urine for HPLC analysis 439(1988)466
- Mathúna, P. Mac, see O'Reilly, T. 417(1987)190
- Matisová, E. Identification of alkylbenzenes by capillary GC and combined GC-MS (Review) 438(1988)131
- Matisová, E., Kraus, G. and Kraus, A. New liquid crystalline stationary phase for capillary GC 439(1988)381
- Matlin, S.A., Stacey, V.E. and Lough, W.J. Hexahelicene chiral stationary phase. I. Phase synthesis and use in HPLC resolution of enantiomers 450(1988)450(1988)157
- Matsanaga, H., Suehiro, T., Saita, T., Nakano, Y., Mori, M., Takata, K. and Oda, K. HPLC determination of pyridostigmine in plasma 422(1987)353
- Matsson, O., see Hussénus, A. 405(1987)155
- Matsubara, K., Akane, A., Takahashi, S., Shiono, H., Fukui, Y., Kagawa, M. and Maseda, C. GC determination for forensic purposes of petroleum fuel inhaled just before fatal burning 424(1988)49
- Matsubara, K., see Kato, Y. 447(1988)212
- Matsubayashi, K., Kojima, C. and Tachizawa, H. Determination of tranexamic acid in human serum by HPLC using selective pre-column derivatization with phenyl isothiocyanate 433(1988)225
- Matsuda, F., see Hayashi, T. 428(1988)209
- Matsuda, I., see Inotsume, N. 428(1988)402
- Matsuda, R., Hayashi, Y., Ishibashi, M. and Takeda, Y. Totally automated robotic system for LC analysis of solid dosage formulations with the aid of a reduced Kalman filter 438(1988)319
- Matsuda, R., see Hayashi, Y. 407(1987)59
- Matsuda, S., see Utsumi, J. 421(1987)355
- Matsuda, T., see Iwata, H. 450(1988)317
- Matsui, M., see Goto, T. 447(1988)410
- Matsuki, Y., see Fukuhara, K. 416(1987)409
- Matsuki, Y., see Ito, T. 417(1987)79
- Matsukura, H., Suzuki, Y., Okada, T., Naiki, S., Hayashi, S. and Yamaji, Y. HPLC determination of acid and alkaline phosphatases in urine 434(1988)196
- Matsukura, H., Suzuki, Y., Okada, T., Naiki, S. and Sakuragawa, N. Automated separation and measurement of urinary isoenzymes and protein by ion-exchange LC 414(1987)47
- Matsumoto, H., see Kominami, G. 417(1987)216
- Matsumoto, K., Kikuchi, H., Iri, H., Takahasi, H. and Umino, M. Automated determination of drugs in serum by column-switching HPLC. II. Separation of theophylline and its metabolites 425(1988)323
- Matsumoto, M., see Hirabayashi, Y. 445(1988)377
- Matsumoto, S., see Togari, A. 417(1987)41
- Matsunaga, R., see Yamaguchi, M. 414(1987)275
- Matsuo, M., see Kern, J.R. 450(1988)233
- Matsuoka, A., see Kobayashi, K. 417(1987)439
- Matsushima, E., see Marunaka, T. 420(1987)43
- Matsushima, E., see Marunaka, T. 420(1987)329
- Matsushima, E., see Marunaka, T. 431(1988)87
- Matsushima, E., see Marunaka, T. 433(1988)177
- Matsushima, E., see Umeno, Y. 434(1988)123
- Matsushita, A., see Suzuki, S.-I. 437(1988)322
- Matsutani, S., see Sano, A. 410(1987)427
- Matsuura, K., see Harada, K.-I. 448(1988)275
- Matsuzaki, H., see Koyama, I. 413(1987)65
- Matthee, F.N., see Thomas, A.C. 403(1987)292
- Matthews, A., see Grgurinovich, N. 433(1988)298
- Mattiasson, B., see Kaul, R. 438(1988)339
- Mattsson, P., Mäkelä, M. and Korpela, T. Chromatographic determination of cyclodextrins on benzoylated polyacrylamide gels 447(1988)398
- Mattsson, P., see Mäkelä, M. 448(1988)391
- Matusch, R., see Friedel, H.D. 407(1987)343
- Matuszewski, B.K., Constanzer, M.L. and Bayne, W.F. Determination of 2,3-dihydro-6-[3-(2-hydroxymethyl)-phenyl-2-propenyl]-5-benzofuranol

- in plasma using LC with electrochemical detection 426(1988)295
- Matysik, G. and Soczewiński, E.  
Stepwise gradient development in TLC. IV. Miniaturized generators for continuous and stepwise gradients 446(1988)275
- Mauk, A.G., see Mauk, M.R. 439(1988)408
- Mauk, M.R. and Mauk, A.G.  
Characterization of wild-type and mutant forms of yeast iso-1-cytochrome *c* by reversed-phase HPLC tryptic peptide mapping 439(1988)408
- Maurer, H. and Pflieger, K.  
Identification and differentiation of benzodiazepines and their metabolites in urine by computerized GC-MS 422(1987)85
- Maurer, H. and Pflieger, K.  
Screening procedure for the detection of alkanolamine antihistamines and their metabolites in urine using computerized GC-MS 428(1988)43
- Maurer, H. and Pflieger, K.  
Identification and differentiation of alkylamine antihistamines and their metabolites in urine by computerized GC-MS 430(1988)31
- Mauri, P., see Pietta, P. 411(1987)498
- Mauri, P., see Pietta, P. 437(1988)453
- Mauri, P., see Pietta, P. 441(1988)431
- Maus, M., Trigalo, F. and Azerad, R.  
Resolution of  $\alpha$ -substituted amino acid enantiomers by HPLC after derivatization with a chiral adduct of *o*-phthalaldehyde. Application to glutamic acid analogues 440(1988)209
- Mawer, E.B. and Hann, J.T.  
Rapid automated HPLC assay for ercalcidiol and calcidiol (25-hydroxyvitamins D<sub>2</sub> and D<sub>3</sub>) using *trans*-calcidiol as an UV-absorbing internal standard 415(1987)305
- May, M.E., see Brown, L.L. 426(1988)370
- Mayer, F., see Krämer, B.K. 427(1988)351
- Mayer, H., see Krupčík, J. 448(1988)203
- Mayersohn, M., see Lopez-Anaya, A. 423(1987)105
- Mazsaroff, I., Bischoff, R., Tice, P.A. and Regnier, F.E.  
Influence of mobile phase pH on HPLC column loading capacity. Implications for the design of preparative protein separations 437(1988)429
- Mazsaroff, I., Cook, S. and Regnier, F.E.  
Molecular orientation of immunoglobulin G at high concentration on an ion-exchange sorbent 443(1988)119
- Mazsaroff, I., Rounds, M.A. and Regnier, F.E.  
Facile preparation of a non-porous strong anion-exchange column for proteins 411(1987)452
- Mazsaroff, I., see Tice, P.A. 410(1987)43
- Mazsaroff, I. and Regnier, F.E.  
Phase ratio determination in an ion-exchange column having pores partially accessible to proteins 442(1988)15
- Mazzo, D.J. and Snyder, P.A.  
HPLC of timolol and potential degradates on dynamically modified silica 438(1988)85
- McAtee, Jr., J.L., see Chukwunyenye, C. 410(1987)121
- McChrisley, B., Thye, F.W., McNair, H.M. and Driskell, J.A.  
Plasma B<sub>6</sub> vitamer and 4-pyridoxic acid concentrations of men fed controlled diets 428(1988)35
- McClain, C.J., see Schmidt, J. 419(1987)1
- McCormick, S.P., Bowers, E. and Bhatnagar, D.  
HPLC procedure for determining the profiles of aflatoxin precursors in wildtype and mutant strains of *Aspergillus parasiticus* 441(1988)400
- McCroskey, M.C., Colca, J.R. and Pearson, J.D.  
Determination of [<sup>32</sup>P]phosphoamino acids in protein hydrolysates by isocratic anion-exchange HPLC 442(1988)307
- McCune, J.E., see Pirkle, W.H. 441(1988)311
- McDaniel, D.M. and Snider, B.G.  
Resolution of  $\alpha$ -arylacetic acid enantiomers on two chiral stationary phases 404(1987)123
- McDaniel, H.B., see Jenkins, R.L. 426(1988)249
- McDaniel, H.G., see Jenkins, R.L. 426(1988)249
- McDowall, M., see Odham, G. 434(1988)31
- McEldowney, A.M. and Menary, R.C.  
Analysis of pyrethrins in pyrethrum extracts by HPLC 447(1988)239
- McErlane, K.M., Igwemezie, L. and Kerr, C.R.  
Stereoselective analysis of the enantiomers of mexiletine by HPLC using fluorescence detection and study of their stereoselective disposition in man 415(1987)335
- McErlane, K.M., see Chan, G.L.-Y. 417(1987)295
- McGarrity, M.J., Er, S.S., Nightingale, K.A. and Hsia, J.C.  
Isolation and partial characterization of pyridoxal 5'-phosphate hemoglobins by HPLC as a quality-control method for hemoglobin-based blood substitutes 413(1987)53

- McGarrity, M.J., Er, S.S., Nightingale, K.A. and Hsia, J.C.  
ATP-hemoglobin: anomalous oxygen binding properties 415(1987)136
- McGarrity, M.J., Er, S.S. and Hsia, J.C.  
Hemoglobin-based blood substitutes: characterization of five pyridoxal 5'-phosphate derivatives of hemoglobin 419(1987)37
- McGhie, T.K., see France, N.C. 433(1988)1
- McGill, A.C., see Seviour, J.A. 432(1988)273
- McGill, R.A., see Abraham, M.H. 409(1987)15
- McGilveray, I.J., see Dubé, L.M. 427(1988)113
- McGilveray, I.J., see Dubé, L.M. 430(1988)103
- McGowan, C.W., see Chriswell, C.D. 405(1987)213
- McGown, E.L., see Knudsen, J.J. 424(1988)231
- McGrath, R.M., see Beveridge, D.J. 450(1988)443
- McInnes, G.T., see Forrest, G. 428(1988)123
- McKerrell, P.J., Saunders, G.A. and Geyer, R.  
Determination of 4,4'-methylenebis(2-chloroaniline) in urine by HPLC 408(1987)399
- McLachlan, D.R., see Kruck, T.P.A. 433(1988)207
- McLaren, A.M., see Lindsay, R.M. 433(1988)292
- McLaughlin-Flynn, C.A., see Mallet, A.I. 432(1988)278
- McLaughlin, L.W. and Bischoff, R.  
Resolution of RNA using HPLC (Review) 418(1987)51
- McLean, A.J., see Keech, A.C. 426(1988)234
- McLean, J.D., see Nair, J.B. 416(1987)340
- McLeish, M., see Sitaram, B.R. 422(1987)13
- McLellan, D.W. and Halling, P.J.  
Preparation and chromatographic analysis of poly(3-hydroxybutyrate) hydrolysis products 445(1988)251
- McLeod, A., see Wallace, G.R. 427(1988)239
- McLeod, W.R., see Sitaram, B.R. 422(1987)13
- McLoughlin, S.B., see Burton, S.J. 435(1988)127
- McMaster, G.K., see Birsner, U. 402(1987)381
- McMeekin, J.R., see Leahey, W.J. 421(1987)401
- McNair, H.M., see McChrisley, B. 428(1988)35
- McNally, M.E., see Wheeler, J.R. 410(1987)343
- McNally, M.E.P. and Wheeler, J.R.  
Supercritical fluid extraction coupled with SFC for the separation of sulfonylurea herbicides and their metabolites from complex matrices 435(1988)63
- McNally, M.E.P. and Wheeler, J.R.  
Increasing extraction efficiency in supercritical fluid extraction from complex matrices. Predicting extraction efficiency of diuron and linuron in supercritical fluid extraction using supercritical fluid chromatographic retention 447(1988)53
- McQuinn, R.L., see Welscher, T.M. 431(1988)438
- Meakings, M., see Bailey, E. 421(1987)21
- Meaney, M., Connor, M., Breen, C. and Smyth, M.R.  
Comparison of HPLC and atomic spectrometric methods for the determination of Fe(III) and Al(III) in soil and clay samples 449(1988)241
- Medina, J.H., see Peña, C. 431(1988)192
- Mee, D., see Child, P. 415(1987)13
- Meemken, H.-A., see Fürst, P. 405(1987)311
- Meer, M.J. Van der and Brown, L.W.  
Simultaneous determination of amiloride and hydrochlorothiazide in plasma by reversed-phase HPLC 423(1987)351
- Meeren, P.V. Der, Vanderdeelen, J., Huys, M. and Baert, L.  
Simple and rapid method for HPLC separation and quantification of soybean phospholipids 447(1988)436
- Meese, C.O., Thalheimer, P. and Eichelbaum, M.  
HPLC method for the analysis of debrisoquine and its *S*-(+)- and *R*-(-)-hydroxy metabolites in urine 423(1987)344
- Meffin, P.J., see Sallustio, B.C. 422(1987)33
- Mefford, I.N.  
Determination of catecholamines in body fluids 427(1988)
- Mefford, I.N., Ota, M., Stipetic, M. and Singleton, W.  
Application of a novel cation-exchange reagent, Igepon T-77 N-methyl oleoyl taurate, to microbore separations of alumina extracts of catecholamines from cerebrospinal fluid, plasma, urine and brain tissue with amperometric detection 420(1987)241
- Mehta, A.C.  
Direct separation of drug enantiomers by HPLC with chiral stationary phases (Review) 426(1988)1
- Mehta, A.C. and Calvert, R.T.  
Determination of ifosfamide in plasma by GC with nitrogen detector 421(1987)377
- Mehta, R.C., Hogan, T.F., Mardmomen, S. and Ma, J.K.H.  
Chromatographic studies of mitomycin C degradation in albumin microspheres 430(1988)341
- Mehvar, R., Jamali, F. and Pasutto, F.M.  
Rapid and sensitive HPLC assay of tiaprofenic acid enantiomers in human plasma and urine 425(1988)135

- Mehvar, R. and Jamali, F.  
Formation of diastereomeric derivatives of 2-arylpropionic acids using L-leucinamide: lack of generality 431(1988)
- Meiborg, G., see Volmer, M. 434(1988)385
- Meier, B., Julkunen-Tiitto, R., Tahvanainen, J. and Sticher, O.  
Comparative HPLC and GLC determination of phenolic glucosides in Salicaceae species 442(1988)175
- Meier, B., see Hofstetter, M. 407(1987)319
- Meier, P., see Gmur, D.J. 425(1988)343
- Meijer, W.H., see De Jong, J. 419(1987)85
- Meinard, C. and Bruneau, P.  
Separation and identification of enantiomers by HPLC with a chiral column and a polarimetric detector as applied to deltamethrin 450(1988)169
- Meineke, I., see Brendel, E. 426(1988)406
- Meineke, I., see Brendel, E. 432(1988)358
- Melendez, V., see Kapetanovic, I.M. 419(1987)458
- Mellini, D.W., see Mathes, L.E. 432(1988)346
- Mello, C.M., see Smith, J.D. 431(1988)395
- Mello, R.J., see Beaty, N.B. 418(1987)187
- Mellström, B.  
HPLC quantitation of 4-hydroxydebrisoquine in microsomal incubates by use of silica columns, aqueous mobile phase and automated column switching 424(1988)435
- Melnikov, S.E., see Berezkina, L.G. 446(1988)65
- Meloni, M.L., see Iadarola, P. 443(1988)317
- Menager, S., see Lacroix, C. 415(1987)85
- Menager, S., see Lacroix, C. 422(1987)217
- Ménard, H., see Noel, L. 436(1988)391
- Menary, R.C., see McEldowney, A.M. 447(1988)239
- Mencia-Huerta, J.M., see Bossant, M.J. 423(1987)23
- Méndez, E., see Escribano, J. 444(1988)165
- Meneilly, G.P., see Scavone, J.M. 419(1987)339
- Menendez, M., see Marhuenda, E. 436(1988)103
- Menéndez, V., see Fernández-Sánchez, E. 402(1987)318
- Menéndez, V., see Fernández-Sánchez, E. 410(1987)13
- Meng, H., Burleigh, B.D. and Kelly, G.M.  
Reduction studies on bacterial recombinant somatomedin C/insulin-like growth factor-1 443(1988)183
- Meng, Q.C., Chen, Y.-F., Delucas, L.J. and Oparil, S.  
Effect of steric exclusion on the separation of proteins by hydrophilic SEC 445(1988)29
- Menge, G.P., Dubois, J.P. and Bauer, G.  
Simultaneous determination of carbamazepine, oxcarbazepine and their main metabolites in plasma by LC 414(1987)477
- Mengher, L.S., see Harvey, D.J. 414(1987)253
- Menichelli, A., see Del Principe, D. 419(1987)329
- Menna, L.F., see Della Ragione, F. 440(1988)141
- Mennes, W.C., see 't Klooster, G.A.E.V. 432(1988)223
- Mennicke, W.H., Kral, T., Krumbiegel, G. and Rittmann, N.  
Determination of N-acetyl-S-(N-alkylthiocarbamoyl)-L-cysteine, a principal metabolite of alkyl isothiocyanates, in rat urine 414(1987)19
- Mentasti, E., Savigliano, M., Marangella, M., Petrarulo, M. and Linari, F.  
HPLC determination of glyoxylic acid and other carbonyl compounds in urine 417(1987)253
- Mentasti, E., see Petrarulo, M. 432(1988)37
- Menu, P., Pauly, F., Labrude, P. and Vigneron, C.  
Haemoglobin pyridoxylation: optimization and observation of an additional new pyridoxal phosphate-haemoglobin species by HPLC resolution 417(1987)397
- Mercier, V., see Airaud, C.B. 437(1988)59
- Meriluoto, J.A.O. and Eriksson, J.E.  
Rapid analysis of peptide toxins in cyanobacteria 438(1988)93
- Merion, M., see Stowers, D.J. 444(1988)47
- Merrill, C.R., see Harrington, M.G. 429(1988)345
- Merwe, K.J. Van der, see Swart, P. 442(1988)424
- Metz, P.A., Omstead, D.R., Kaplan, L., Liesch, J.M., Stearns, R.A. and VandenHeuvel, W.J.A.  
Characterization of a lipid-rich fraction synthesized by *Streptomyces avermitilis* 441(1988)31
- Metzner, K.E., see Stafford, A.E. 436(1988)93
- Meuldijk, J., see Verhaar, L.A.T. 437(1988)185
- Meunier, F., see Brassinne, C. 419(1987)401
- Mey, C. De, see Brendel, E. 426(1988)406
- Meyer-Sabellek, W., Sinha, P. and Köttgen, E.  
Alkaline phosphatase. Laboratory and clinical implications (Review) 429(1988)419
- Meyer, V.R., see Brügger, R.R. 440(1988)197
- Mezdour, H., Kora, V.C.I., Koffigan, M., Barkia, A. and Fruchart, J.-C.  
Anion-exchange fast protein LC characterization and purification of apolipoproteins A-I, A-II, C-I, C-II, C-III<sub>0</sub>, C-III<sub>1</sub>, C-III<sub>2</sub> and E from human plasma 414(1987)35
- Mező, G., see Szókán, G. 444(1988)115

- Mezzetti, T., Proietti, G. and Sbordoni, A.  
Simple method for the determination of  
hormonal iodine and exogenous iodine from  
X-ray contrast media drugs 434(1988)308
- Micali, G., see Currò, P. 404(1987)273
- Michaelis, H.C., Foth, H. and Kahl, G.F.  
Determination of 5-fluoro-2'-deoxyuridine in  
human plasma by HPLC with pre-column  
fluorimetric derivatization 416(1987)176
- Michałek, J., see Rudziński, W. 406(1987)295
- Michaud, J.-P., see Firestone, M.A. 407(1987)363
- Michel, G., see Billard, V. 440(1988)87
- Michelsen, P., see Odham, G. 434(1988)31
- Michida, T., see Sayo, H. 417(1987)129
- Michotte, Y., see Herregodts, P. 421(1987)51
- Midha, K.K., see Ganes, D.A. 423(1987)227
- Midha, K.K., see Gupta, M.B. 424(1988)189
- Midha, K.K., see Srinivas, N.R. 433(1988)105
- Midskov, C.  
HPLC assay of fenflurazole and its demethyl  
metabolites in biological samples  
413(1987)189
- Mierzwa, R., Cooper, R. and Pramanik, B.  
Photodiode array detection of  
peptide-nucleoside antibiotics 436(1988)259
- Mies, H.H. and Zahn, H.  
Chromatographic and electrophoretic  
investigations of the properties of unprotected  
low-sulphur wool keratins 405(1987)365
- Mignot, A., Lefebvre, M.A. and Fourtillan, J.B.  
HPLC determination of ofloxacin in plasma  
and urine 430(1988)192
- Miguel, I. De, Puech-Costes, E. and Samain, D.  
Use of mixed perfluorinated ion-pairing  
agents as solvents in ion-pair HPLC for the  
preparative purification of aminoglycoside  
antibiotics 407(1987)109
- Mihaly, G.W., Hyman, K.M., Smallwood, R.A.  
and Hardy, K.J.  
HPLC analysis of quinine and its  
diastereoisomer quinidine 415(1987)177
- Mihályfi, K., see Gazdag, M. 450(1988)145
- Mihara, S. and Masuda, H.  
Correlation between molecular structures and  
retention indices of pyrazines 402(1987)309
- Miheeva, L.M., see Zaslavsky, B.Y. 403(1987)123
- Miheeva, L.M., see Zaslavsky, B.Y. 439(1988)267
- Mikati, M.A., Szabo, G.K., Pylilo, R.J., LeDuc,  
B.W., Browne, T.R. and Greenblatt, D.J.  
Improved HPLC assay of antipyrine,  
hydroxymethylantipyrine, 4-  
hydroxyantipyrine and norantipyrine in urine  
433(1988)305
- Mikeš, O., Hostomská, Z., Štrop, P., Smrž, M.,  
Slováková, S., Vrátný, P., Rexová-Benková,  
L., Kolář, J. and Čoupek, J.  
Ion-exchange derivatives of Spheron. VI.  
Strongly basic derivatives 440(1988)287
- Mikešová, M., see Vejrosta, J. 447(1988)170
- Mikoshiba, K., see Hase, S. 434(1988)51
- Miksic, J.R., Rivel, E.R. and Rush, K.  
Analysis of a new H<sub>2</sub> receptor antagonist,  
3-amino-5-[3-[4-(1-piperidinoindanyloxy)]-  
propylamino]-1-methyl-1H-1,2,4-triazole, in  
human plasma and urine by HPLC  
428(1988)113
- Milborrow, B.V., see Rhodes, S.H. 442(1988)412
- Millart, H., see Lamiable, D. 434(1988)315
- Miller, B.J., Tenbarger, J.B., Bernstein, J.R. and  
Franklin, R.B.  
HPLC assay of *trans*-(-)-5,5a,6,7,8,9,9a,10-  
octahydro-6-propylpyrimido[4,5-g]quinolin-2-amine:  
dihydrochloride (LY163502), a potent D-2  
agonist, in plasma 426(1988)438
- Miller, C., see Slegowski, M.B. 425(1988)227
- Miller, D.J., see Hawthorne, S.B. 403(1987)63
- Miller, N.T., see Sagliano, Jr., N. 443(1988)155
- Miller, R.D., see Furuta, T. 427(1988)41
- Miller, R.L. and DeVane, C.L.  
Alprazolam,  $\alpha$ -hydroxy- and 4-  
hydroxylprazolam analysis in plasma by  
HPLC 430(1988)180
- Miller, V., see Kuchař, M. 449(1988)391
- Milligan, J.R., Catz-Biro, L. and Archer, M.C.  
Effect of phosphate concentration on the yield  
and purity of DNA separated from RNA by  
hydroxyapatite chromatography  
411(1987)481
- Millington, D.S., see Dietrich, A.M. 436(1988)229
- Millot, M.-C. and Sebillé, B.  
Rapid preparation of bovine mercaptalbumin  
by means of covalent chromatography on  
silica-based materials 408(1987)263
- Mills, P., see Lalande, M. 430(1988)187
- Milton, K.A., Ward, S.A. and Edwards, G.  
Determination of halofantrine and its  
principal metabolite desbutylhalofantrine in  
biological fluids by reversed-phase HPLC  
433(1988)339
- Mimaki, T., see Yamaji, A. 421(1987)372
- Mina, M. and Rossomando, E.F.  
Assay of ATP-sulfurylase activity from rat  
liver by HPLC 433(1988)63
- Minami, Y., see Marunaka, T. 420(1987)43
- Minami, Y., see Marunaka, T. 431(1988)87
- Minard, P., see Yon, J.M. 440(1988)421
- Minchiotti, L., see Iadarola, P. 443(1988)317
- Minder, C.E., see Minder, E.I. 419(1987)135
- Minder, E.I., Schaubhut, R., Minder, C.E. and  
Vonderschmitt, D.J.

- Identification of drugs in human serum by HPLC with photodiode array detection and a search algorithm for UV spectra 419(1987)135
- Minder, E.I., Schaubhut, R. and Vonderschmitt, D.J.  
Screening for drugs in clinical toxicology by HPLC: identification of barbiturates by post-column ionization and detection by a multiplex photodiode array spectrophotometer 428(1988)369
- Minelli, A., see Palmerini, C.A. 417(1987)378
- Minematsu, T., see Honda, S. 419(1987)51
- Miner, D.J., see Riggin, R.M. 435(1988)307
- Miners, J.O., see Chan, K. 426(1988)103
- Miners, J.O., see Lelo, A. 430(1988)203
- Mingjia, W., Pacáková, V., Štulík, K. and Sacchetto, G.A.  
Ion-pair HPLC of inorganic anions with photometric, conductometric and amperometric detection 439(1988)363
- Mink, L.P., see Lochmüller, C.H. 409(1987)55
- Minkler, P.E., Ingalls, S.T., Griffin, R.L. and Hoppel, C.L.  
Rapid HPLC of 3-methylhistidine in human urine 413(1987)33
- Minkler, P.E., Ingalls, S.T. and Hoppel, C.L.  
Determination of total carnitine in human urine by HPLC 420(1987)385
- Minkler, P.E., see Chai, B. 430(1988)93
- Minkler, P.E. and Hoppel, C.L.  
Determination of ibuprofen in human plasma by HPLC 428(1988)388
- Minnietian, O., Saha, M. and Giese, R.W.  
Oxidation-elimination of a DNA base from its nucleoside to facilitate determination of alkyl chemical damage to DNA by GC with electrophore detection 410(1987)453
- Minnich, M.D., see Strohsacker, M.W. 435(1988)185
- Miocque, M., see Coleman, A.W. 450(1988)175
- Miranda, P. De, see Good, S.S. 431(1988)123
- Miribel, L. and Arnaud, P.  
Purification of human  $\alpha_1$ -beta glycoprotein and study of its microheterogeneity and molecular variants 405(1987)337
- Misawa, S., see Tanaka, E. 413(1987)376
- Mischnick-Lübbecke, P., see König, W.A. 447(1988)193
- Mishima, M., see Yamaji, A. 415(1987)143
- Mishra, U., see Singh, A.K. 404(1987)223
- Misiti, D., see Galli, B. 409(1987)377
- Miski, M. and Davis, P.J.  
GC determination of racemic *cis*- and *trans*-chrysanthemols and their potential aldehyde and carboxylic acid microbial metabolites 437(1988)436
- Missiakas, D., see Yon, J.M. 440(1988)421
- Mita, H., Yui, Y., Yasueda, H. and Shida, T.  
Isocratic determination of arachidonic acid 5-lipoxygenase products in human neutrophils by HPLC 430(1988)299
- Mitchell, J.R., see Montamat, S.C. 415(1987)203
- Mitnaul, L.J., see Zaromb, S. 439(1988)283
- Mitraki, A., see Yon, J.M. 440(1988)421
- Mitsubishi, Y., see Hamano, T. 411(1987)423
- Mitsui, A., see Kato, Y. 447(1988)212
- Mitsutani, K., see Sato, H. 447(1988)387
- Miura, M., see Koyama, I. 413(1987)65
- Miura, T., see Kanetoshi, A. 442(1988)289
- Miwa, B.J., Choma, N., Brown, S.Y., Keigher, N., Garland, W.A. and Fukuda, E.K.  
Quantitation of the enantiomers of rimantadine in human plasma and urine by GC-MS 431(1988)343
- Miwa, H., Yamamoto, M., Nishida, T., Nunoi, K. and Kikuchi, M.  
HPLC analysis of serum long-chain fatty acids by direct derivatization method 416(1987)237
- Miwa, H. and Yamamoto, M.  
HPLC analysis of serum short-chain fatty acids by direct derivatization 421(1987)33
- Miwa, T., Miyakawa, T., Kayano, M. and Miyake, Y.  
Application of an ovomucoid-conjugated column for the optical resolution of some pharmaceutically important compounds 408(1987)316
- Miyakawa, T., see Miwa, T. 408(1987)316
- Miyake, K., Mizuno, N. and Terada, H.  
Effect of hydrogen bonding on the HPLC behaviour of organic compounds.  
Relationship between capacity factors and partition coefficients 439(1988)227
- Miyake, Y., see Miwa, T. 408(1987)316
- Miyamoto, Y.  
Highly sensitive determination of chlorpheniramine as fluorescence derivative by HPLC 420(1987)63
- Miyano, S., see Nakano, K. 438(1988)61
- Miyano, S., see Yamashita, J. 403(1987)275
- Miyata, A., see Koyama, Y. 439(1988)417
- Miyazaki, H., see Yamashita, K. 424(1988)1
- Miyazaki, K., see Tanaka, Y. 407(1987)197
- Mizobuchi, N., see Ageta, T. 419(1987)334
- Mizuguchi, S., see Yamashita, K. 424(1988)39
- Mizuno, N., see Miyake, K. 439(1988)227
- Mladenov, P.V., see Asotra, S. 408(1987)227
- Mochizuki, E., see Horie, M. 402(1987)301

- Möckel, H.J., see Weiss, J. 439(1988)91
- Modak, M.J., see Pandey, V.N. 444(1988)133
- Moeller, P.D.R., see Wenkert, E. 440(1988)449
- Moerloose, P. De, see De Schutter, J.A. 428(1988)301
- Moerloose, P. De, see De Schutter, J.A. 437(1988)83
- Moerloose, P. De, see De Schutter, J.A. 450(1988)337
- Mogi, M., see Harada, M. 424(1988)129
- Mohanty, J.G. and Elazhary, Y.  
HPLC separation and immunological characterization of soluble bovine viral diarrhoea virus antigen 435(1988)149
- Molen, J.C. Van der, see Koopman, B.J. 416(1987)1
- Molera, M.J., see Fernández-Sánchez, E. 410(1987)13
- Molinaro, N.C., see Smith, H.T. 424(1988)416
- Moll, N. and Joly, J.P.  
Determination of ascorbic acid in beers by HPLC with electrochemical detection 405(1987)347
- Möller, P., see Larsson, K. 450(1988)71
- Molnár-Perl, I., Pintér-Szakács, M., Morvai, M. and Fábíán-Vonsik, V.  
GC analysis of different homologous series of acids esterified in aqueous solutions with butyl and propyl alcohols 446(1988)237
- Molnár-Perl, I. and Fábíán-Vonsik, V.  
GC analysis of histidine: effect on other amino acids present 446(1988)231
- Moltrasio, D., see Fedeli, G. 447(1988)263
- Momose, T., see Iida, T. 438(1988)423
- Mompon, B., see Krstulović, A.M. 426(1988)417
- Monastyrskii, V.A., see Gaida, A.V. 424(1988)385
- Moncrieff, J.  
Assay of debrisoquine and 4-hydroxydebrisoquine in urine by reversed-phase HPLC using on-line sample clean-up on a standard isocratic chromatograph 428(1988)178
- Moneam, N.M.A., Elsharaky, A.S. and Badreldin, M.M.  
Biological effect of guava oestrone on uterine tissue of mice 437(1988)316
- Moneam, N.M.A., El Sharaky, A.S. and Badreldin, M.M.  
Oestrogen content of pomegranate seeds 438(1988)438
- Monjour, L., see Ogunkolade, B.W. 440(1988)459
- Monk, C.R., see Patchett, M.L. 425(1988)269
- Monsarrat, B., see Rivière, M. 445(1988)87
- Monster, A.C., see Kežić, S. 428(1988)275
- Montagu, M., Levillain, P., Chenieux, J.C. and Rideau, M.  
Fluorodensitometric assay of 6*H*-pyrido[4,3-*b*]-5,11-dimethylcarbazoles (ellipticine and derivatives) biosynthesized by *Ochrosia elliptica* cultures *in vitro* 409(1987)426
- Montamat, S.C., Abernethy, D.R. and Mitchell, J.R.  
High-performance liquid chromatographic determination of diltiazem and its major metabolites, N-monodemethyl diltiazem and desacetyldiltiazem, in plasma 415(1987)203
- Montana, M., see Biondi, P.A. 411(1987)275
- Montanarella, L., see Gagliardi, L. 448(1988)296
- Monteil, H., see Jehl, F. 413(1987)109
- Monteil, H., see Koechlin, C. 425(1988)197
- Monteil, H., see Rihn, B. 428(1988)408
- Montes, M., see Usero, J.L. 407(1987)355
- Monti, J.-P., see Gallice, P. 422(1987)263
- Montreuil, J., see Capon, C. 425(1988)35
- Monzani, V., see Castoldi, D. 425(1988)413
- Moody, R.R., see Taylor, R.B. 416(1987)394
- Moor, M.J., Wyss, P.A. and Bickel, M.H.  
New procedure for the HPLC determination of amiodarone and desethylamiodarone with solid-phase extraction of rat plasma and tissue samples 431(1988)455
- Moore, G.L. and Fishman, R.M.  
HPLC analysis of ascorbate-2-phosphate, adenine and hypoxanthine in stored human blood 419(1987)95
- Moore, J.M., Cooper, D.A., Lurie, I.S., Kram, T.C., Carr, S., Harper, C. and Yeh, J.  
Capillary GC-electron capture detection of coca-leaf-related impurities in illicit cocaine: 2,4-diphenylcyclobutane-1,3-dicarboxylic acids, 1,4-diphenylcyclobutane-2,3-dicarboxylic acids and their alkaloidal precursors, the truxillines 410(1987)297
- Moore, J.M., see Lurie, I.S. 405(1987)273
- Moore, J.N., see Akbari, A. 426(1988)207
- Moore, L.K., see Krahn, M.M. 437(1988)161
- Moore, W.T., see Caprioli, R.M. 443(1988)355
- Moran, E., O'Kennedy, R. and Thornes, R.D.  
Analysis of coumarin and its urinary metabolites by HPLC 416(1987)165
- Morel, D., see Chartier, A. 438(1988)263
- Moretti, G., Amici, M., Cammarata, P. and Fracassi, F.  
Identification of thyrostatic drug residues in animal thyroids by HPTLC and fluorescence reaction detection 442(1988)459
- Moretti, J.L., see Joulin, Y. 425(1988)402
- Moretto, R., see Padriani, R. 415(1987)183
- Morgan, D.J., see Kumar, K. 419(1987)464

- Morgan, E.D., Murphy, S.J., Games, D.E. and Mylchreest, I.C.  
Analysis of ecdysteroids by supercritical-fluid chromatography 441(1988)165
- Morgan, H.W., see Patchett, M.L. 425(1988)269
- Morgan, S.L., see Greene, S.V. 417(1987)121
- Morgan, S.L., see Smith, C.S. 423(1987)207
- Mori, A., see Watanabe, Y. 425(1988)373
- Mori, E., see Igarashi, K. 415(1987)407
- Mori, I., see Motohashi, N. 435(1988)335
- Mori, M., see Matsanaga, H. 422(1987)353
- Mori, M., see Ohmori, S. 414(1987)149
- Mori, M., see Ohmori, S. 415(1987)221
- Mori, S.  
Comparison between SEC and liquid adsorption chromatography in the determination of the chemical heterogeneity of copolymers 411(1987)355
- Morikawa, N., see Takagi, K. 432(1988)47
- Morikawa, Y., see Kikuchi, J. 403(1987)319
- Morimoto, A., see Hirauchi, K. 430(1988)21
- Morin-Allory, L., see Lafosse, M. 404(1987)95
- Morin, D., see Spreux-Varoquaux, O. 416(1987)311
- Morin, P., Caude, M. and Rosset, R.  
Super- and subcritical fluid chromatography-Fourier transform IR spectrometry of medium polar compounds on polymeric styrene-divinylbenzene packed columns 407(1987)87
- Morinaga, S. and Ishiwatari, R.  
GC determination of C<sub>1</sub>-C<sub>5</sub> low-molecular-weight organic acids in alkaline permanganate oxidation products of humic substances 403(1987)225
- Morishita, H., see Kidouchi, K. 423(1987)297
- Morita, T., see Fujimoto, C. 438(1988)329
- Moriwaki, Y., see Yamamoto, T. 413(1987)342
- Moriyama, H., see Kato, Y. 404(1987)333
- Moriyama, H. and Kato, Y.  
New reversed-phase HPLC column for oligonucleotide separation 445(1988)225
- Moriyasu, M., Kato, A. and Hashimoto, Y.  
Kinetic studies of fast equilibrium by means of HPLC. XVI. Separation of tautomers of  $\beta$ -diketones and  $\beta$ -diketoamides 411(1987)466
- Moroux, Y., Boschetti, E., Bouvet, J.-P. and Barot-Ciorbaru, R.  
Application of chromatographic methods to the analysis of macrophage factors induced by *Nocardia opaca* cell walls 440(1988)119
- Morris, M.D., see Peck, K. 448(1988)193
- Morrison, W.R., see Christie, W.W. 436(1988)510
- Mortensen, H.B., see Flückiger, R. 429(1988)279
- Mortimer, D.N., Gilbert, J. and Shepherd, M.J.  
Rapid and highly sensitive analysis of aflatoxin M<sub>1</sub> in liquid and powdered milks using an affinity column cleanup 407(1987)393
- Morton, J.B., see Anderson, G.M. 424(1988)373
- Morvai, M., see Molnár-Perl, I. 446(1988)237
- Mosbach, K., see Sellergren, B. 450(1988)277
- Moschidis, M.C.  
Silicic acid CC of phosphonolipids. IX. Silicic acid CC study of some phosphono analogues of bis(diacylglycero)phosphate and glycerophosphoric acid 435(1988)508
- Moschidis, M.C. and Andrikopoulos, N.K.  
Separation of the phosphono analogues of phosphatidylcholine and phosphatidylethanolamine from related phospholipids by HPLC 403(1987)363
- Mosher, R.A., Thormann, W. and Bier, M.  
Experimental and theoretical dynamics of isoelectric focusing. II. Elucidation of the impact of the electrode assembly 436(1988)191
- Mosser, J., see Duchene, P. 424(1988)205
- Motassim, N., Decolin, D., Le Dinh, T., Nicolas, A. and Siest, G.  
Direct determination of dextromethorphan and its three metabolites in urine by HPLC using a precolumn switching system for sample clean-up 422(1987)340
- Motassim, N., see Decolin, D. 419(1987)469
- Motohashi, N., Nakamichi, Y., Mori, I., Nishikawa, H. and Umemoto, J.  
Analysis by high-performance GPC of hyaluronic acid in animal skins and rabbit synovial fluid 435(1988)335
- Motorin, Y.A.  
Determination of binding constants by high-performance gel filtration 404(1987)254
- Motozato, Y., see Hirayama, C. 409(1987)175
- Mould, N., see Griffin, M. 431(1988)285
- Moulds, R.F.W., see Fullinfaw, R.O. 415(1987)347
- Moulds, R.F.W., see Fullinfaw, R.O. 433(1988)131
- Mount, D.L., Nahlen, B.L., Patchen, L.C. and Churchill, F.C.  
Field-adapted method for HPTLC detection and estimation of chloroquine and desethylchloroquine in urine 423(1987)261
- Mount, D.L., Patchen, L.C. and Churchill, F.C.  
Field-adapted method for HPTLC detection and estimation of chloroquine in finger-stick blood 428(1988)196



- Mount, D.L., see Bergqvist, Y. 428(1988)281
- Mousseau, N., see Dubé, L.M. 430(1988)103
- Mozayani, A., Coutts, R.T. and Danielson, T.J.  
GC analysis of monoalkylhydrazines  
423(1987)131
- Mozersky, S.M., see Parris, N. 444(1988)219
- Mráz, J.  
GC method for the determination of N-acetyl-S-(N-methylcarbamoyl)cysteine, a metabolite of N,N-dimethylformamide and N-methylformamide, in human urine  
431(1988)361
- Mráz, J. and Tureček, F.  
Identification of N-acetyl-S-(N-methylcarbamoyl)cysteine, a human metabolite of N,N-dimethylformamide and N-methylformamide 414(1987)399
- Mross, K.B., see Maessen, P.A. 417(1987)339
- Mross, K.B., see Maessen, P.A. 424(1988)103
- Mucha, I. and Tóth, G.  
Separation of <sup>125</sup>I-labelled prostaglandin E<sub>2</sub>-tyrosine methyl ester by reversed-phase HPLC 438(1988)111
- Mücke, H.-W., see Räder, K. 408(1987)341
- Muda, M., Pelizzoni, F., Sello, G. and Mussini, E.  
Determination of ergothioneine in red blood cells by HPLC 434(1988)191
- Muirhead, D.C. and Christie, R.B.  
Simple, sensitive and selective HPLC method for analysis of chlorthalidone in whole blood  
416(1987)420
- Mukaiyama, Y., see Manabe, T. 431(1988)45
- Mukherjee, K.D., see Schuch, R. 450(1988)448
- Mukkur, T.K.S., see Altmann, K. 420(1987)253
- Mulder, G.J., see Te Koppele, J.M. 427(1988)67
- Müller, A., see Weiss, J. 439(1988)91
- Muller, D., see Santarelli, X. 443(1988)55
- Müller, E., see Grob, K. 404(1987)297
- Müller, M., see Förster, H.-G. 424(1988)357
- Müller, R.F., see Bökens, H. 434(1988)327
- Müller, S., see Reiter, C. 413(1987)251
- Müller, T., see Reiter, C. 413(1987)251
- Mullersman, G., see Townsend, R.W. 450(1988)414
- Mullin, W.J., see Collins, F.W. 445(1988)363
- Munavalli, S. and Pannella, M.  
TLC of mustard and its metabolites  
437(1988)423
- Muniak, W., Witkiewicz, Z., Woryna, E., Kusińska, E., Twardowski, A. and Goca, B.  
Preparation of standard gaseous mixtures by a manometric method using the Benedict-Webb-Rubin equation  
436(1988)323
- Munk, M.N., see Nair, J.B. 416(1987)340
- Munns, R.K., Roybal, J.E., Shimoda, W. and Hurlbut, J.A.  
1-(4-Hydroxyphenyl)-, 1-(2,4-dihydroxyphenyl)- and 1-(2,5-dihydroxyphenyl)-2-bromoethanones: new labels for determination of carboxylic acids by HPLC with electrochemical and UV detection 442(1988)209
- Muñoz, R., see Barcelo, A.R. 411(1987)476
- Munro, Z., see Burger, B.V. 402(1987)95
- Mura, P., Piriou, A., Fraillon, P., Papet, Y. and Reiss, D.  
Screening procedure for benzodiazepines in biological fluids by HPLC using a rapid-scanning multichannel detector 416(1987)303
- Murai, H., see Yoshida, S. 431(1988)27
- Murakita, H.  
Simultaneous determination of creatine and creatinine in serum by HPLC 431(1988)471
- Muramatsu, I., see Tamaki, M. 413(1987)287
- Murase, K., see Takeuchi, T. 445(1988)139
- Murata, K. and Yokoyama, Y.  
LC assay for constituent disaccharides of hyaluronic acid and chondroitin sulphate isomers 415(1987)231
- Murata, K. and Yokoyama, Y.  
Characterization of the products generated from oversulphated dermatan sulphate isomers with chondroitinase-B by HPLC 423(1987)51
- Muratti, E., see Chersi, A. 410(1987)463
- Murayama, T., Kubota, T., Hanaoka, Y., Rokushika, S., Kihara, K. and Hatano, H.  
Peak enhancement of carboxylic acids in ion-exclusion chromatography with a cation-exchange hollow-fibre suppressor using neutral and alkaline enhancers 435(1988)417
- Murello, M.H., see Righezza, M. 410(1987)145
- Muriasco, A., see Gallice, P. 422(1987)263
- Murotsu, T., see Kato, Y. 447(1988)212
- Murphy, M.E. and Kehrer, J.P.  
Simultaneous measurement of tocopherols and tocopheryl quinones in tissue fractions using HPLC with redox-cycling electrochemical detection 421(1987)71
- Murphy, P.A., see O'Keefe, S.F. 445(1988)305
- Murphy, R., Furness, J.B. and Costa, M.  
High-sensitivity phenylthiohydantoin amino acid analysis on-line to a gas phase protein sequencer 408(1987)388
- Murphy, R., see Turner, C.A. 428(1988)383
- Murphy, S.J., see Morgan, E.D. 441(1988)165
- Murray, G.J., see Mussar, K.J. 408(1987)378
- Murray, K.E. and Adams, R.F.

- Determination of simple phenols in faeces and urine by HPLC 431(1988)143
- Musch, G. and Massart, D.L.  
Isolation of basic drugs from plasma using solid-phase extraction with a cyanopropyl-bonded phase 432(1988)209
- Muschik, G., see Mathes, L.E. 432(1988)346
- Muskiet, F.A.J., see Hoving, E.B. 434(1988)395
- Muskiet, F.A.J., see Hoving, E.B. 434(1988)411
- Muskiet, F.A.J., see Van den Berg, G.A. 415(1987)27
- Muskiet, F.A.J., see Van der Steege, G. 415(1987)1
- Muskiet, F.A.J., see Volmer, M. 434(1988)385
- Mussar, K.J., Murray, G.J., Martin, B.M. and Viswanatha, T.  
Rapid chromatographic assay procedure for peptide-N-glycosidase activity 408(1987)378
- Musser, S.M., see Andrews, P.A. 419(1987)271
- Mussini, E., see Muda, M. 434(1988)191
- Musson, D.G., Maglietto, S.M., Hwang, S.S., Gravellese, D. and Bayne, W.F.  
Simultaneous quantification of cycloserine and its prodrug acetylacetyl cycloserine in plasma and urine by HPLC using UV absorbance and fluorescence after post-column derivatization 414(1987)121
- Musu, C., Fumagalli, L., Perego, F., Fedeli, F. and Uggeri, F.  
Quantitative determination of 2-amino-1,3-propanediol and its impurities by capillary GC 449(1988)432
- Mutoh, T., see Naoi, M. 426(1988)75
- Mutschler, E., see Gietl, Y. 426(1988)305
- Mutschler, E., see Pflugmann, G. 416(1987)331
- Mutschler, E., see Pflugmann, G. 421(1987)161
- Mutschler, E., see Spahn, H. 433(1988)331
- Myers, C.M. and Blumer, J.L.  
HPLC of ciprofloxacin and its metabolites in serum, urine and sputum 422(1987)153
- Mylchreest, I.C., see Morgan, E.D. 441(1988)165
- Naeve, G.S., see Chen, F.-M. 444(1988)153
- Nagai, K., see Ukai, S. 435(1988)374
- Nagai, S., see Kidouchi, K. 423(1987)297
- Nagamachi, M., see Marunaka, T. 420(1987)43
- Nagao, T., see Suzuki, S.-I. 437(1988)322
- Nagaoka, T., see Hirauchi, K. 430(1988)21
- Nagata, T., see Kudo, K. 431(1988)353
- Nagata, T., see Uehori, R. 411(1987)251
- Nagatsu, T., see Naoi, M. 426(1988)75
- Nagatsu, T., see Naoi, M. 427(1988)229
- Nagatsu, T., see Naoi, M. 434(1988)209
- Nagatsu, T., see Sakai, T. 416(1987)131
- Nagatsu, T., see Sasaoka, T. 424(1988)392
- Nagatsu, T., see Sasaoka, T. 428(1988)152
- Nagatsu, T., see Togari, A. 417(1987)41
- Nagayama, K., see Koyama, Y. 439(1988)417
- Nagoshi, T., see Jinno, K. 436(1988)1
- Nagy, E., Csipo, I., Degrell, I. and Szabo, G.  
HPLC assay of 5-aminosalicylic acid and its acetylated metabolite in biological fluids using electrochemical detection 425(1988)214
- Nagy, L.G., László, K., Fóti, G. and Schay, G.  
Investigation of HPLC packings by liquid mixture adsorption 406(1987)311
- Nagy, L.G., Vajda, N., Vodicska, M., Zagyvai, P. and Solymosi, J.  
Automated separation process for radioanalytical purposes at nuclear power plants 406(1987)359
- Nahlen, B.L., see Mount, D.L. 423(1987)261
- Naiki, S., see Matsukura, H. 414(1987)47
- Naiki, S., see Matsukura, H. 434(1988)196
- Nair, J.B., Munk, M.N. and McLean, J.D.  
Improved HPLC method with electrochemical detection for the determination of urinary catecholamines using a new electrochemical detector 416(1987)340
- Nakagawa, A., see Iwabuchi, H. 414(1987)139
- Nakagawa, F., see Aoyama, T. 430(1988)351
- Nakagawa, F., see Yamamoto, K. 433(1988)167
- Nakagawa, T., Shibukawa, A., Shimono, N., Kawashima, T., Tanaka, H. and Haginaka, J.  
Retention properties of internal-surface reversed-phase silica packing and recovery of drugs from human plasma 420(1987)297
- Nakahara, Y. and Cook, C.E.  
Confirmation of cannabis use. III.  
Simultaneous quantitation of six metabolites of  $\Delta^9$ -tetrahydrocannabinol in plasma by HPLC with electrochemical detection 434(1988)247
- Nakahashi, H., see Mabuchi, H. 415(1987)110
- Nakajima, K., see Suzuki, S.-I. 437(1988)322
- Nakamichi, Y., see Motohashi, N. 435(1988)335
- Nakamura, K.-I., see Iwabuchi, H. 414(1987)139
- N**
- Naaranlahti, T., Nordström, M., Huhtikangas, A. and Lounasmaa, M.  
Determination of *Catharanthus* alkaloids by reversed-phase HPLC 410(1987)488
- Nabet, P., see Brossat, B. 413(1987)141

- Nakamura, K., see Kern, J.R. 450(1988)233
- Nakamura, K., see Kikuchi, J. 403(1987)319
- Nakamura, K., see Kitano, H. 420(1987)13
- Nakamura, K., Toyoda, K. and Kato, Y.  
High-performance affinity chromatography of proteins on TSKgel Heparin-5PW 445(1988)234
- Nakamura, M., see Hirayama, C. 409(1987)175
- Nakamura, M., see Iwata, T. 421(1987)43
- Nakamura, M., see Yamaguchi, M. 414(1987)275
- Nakamura, T., see Yamashita, K. 424(1988)39
- Nakanishi, Y., see Kojima, K. 413(1987)326
- Nakanishi, Y., see Kojima, K. 425(1988)203
- Nakano, K., Sekino, Y., Yomo, N., Wakayama, S., Miyano, S., Kusaka, K., Daimon, E., Imaizumi, K., Totsuka, Y., Oda, S. and Yamada, Y.  
HPLC of carthamin, safflor yellow A and a precursor of carthamin. Application to the investigation of an unknown red pigment produced in cultured cells of safflower 438(1988)61
- Nakano, M., Kai, M., Ohno, M. and Ohkura, Y.  
HPLC of N-terminal tyrosine-containing oligopeptides by pre-column fluorescence derivatization with hydroxylamine, cobalt(II) and borate reagents 411(1987)305
- Nakano, M., see Inotsume, N. 428(1988)402
- Nakano, Y., see Matsunaga, H. 422(1987)353
- Nakao, T., see Hirabayashi, Y. 445(1988)377
- Nakashima, K., Umekawa, C., Yoshida, H., Nakatsuji, S. and Akiyama, S.  
HPLC fluorometry for the determination of thiols in biological samples using N-[4-(6-dimethylamino-2-benzofuranyl)phenyl]-maleimide 414(1987)11
- Nakata, K., see Ohmori, S. 426(1988)15
- Nakata, O., see Kikuchi, J. 403(1987)319
- Nakatsuji, S., see Nakashima, K. 414(1987)11
- Nakayama, F., see Chijiwa, K. 431(1988)17
- Nakazawa, H.  
Fluorimetric determination of rat growth hormone in pituitary cell culture by HPLC with post-column derivatization 417(1987)409
- Nakazawa, H., see Horie, M. 402(1987)301
- Nakazawa, K., see Sugita, S. 421(1987)412
- Naline, E., Flouvat, B., Advenier, C. and Pays, M.  
Determination of theophylline and its metabolites in plasma and urine by reversed-phase LC using an amine modifier 419(1987)177
- Nam, H.S., Cho, S.Y. and Rhee, J.S.  
HPLC analysis of major carotenoids from *Rhodotorula glutinis* 448(1988)445
- Nambara, T., see Goto, J. 415(1987)45
- Nambara, T., see Goto, J. 425(1988)59
- Nambara, T., see Iida, T. 438(1988)423
- Nambara, T., see Ito, T. 417(1987)79
- Nambara, T., see Shimada, K. 419(1987)17
- Nancollas, G.H., see Sing, R.P. 433(1988)373
- Nandoskar, M., see Ferrante, A. 440(1988)105
- Naoi, M., Ichinose, H., Takahashi, T. and Nagatsu, T.  
Sensitive assay for determination of pyridoxal-5-phosphate in enzymes using HPLC after derivatization with cyanide 434(1988)209
- Naoi, M., Kondoh, M., Mutoh, T., Takahashi, T., Kojima, T., Hirooka, T. and Nagatsu, T.  
Microassay for GM<sub>1</sub> ganglioside  $\beta$ -galactosidase activity using HPLC 426(1988)75
- Naoi, M., Takahashi, T. and Nagatsu, T.  
Simple assay procedure for tyrosine hydroxylase activity by HPLC employing coulometric detection with minimal sample preparation 427(1988)229
- Naora, K., see Katagiri, Y. 431(1988)135
- Narang, P.K., see Amantea, M.A. 426(1988)216
- Narce, M., Gresti, J. and Bezdard, J.  
Method for evaluating the bioconversion of radioactive polyunsaturated fatty acids by use of reversed-phase LC 448(1988)249
- Narita, O., see Takagi, K. 432(1988)47
- Naruse, H., see Hayashi, T. 428(1988)209
- Natile, G., see Galli, B. 409(1987)377
- Nation, R.L., see Gross, A.S. 416(1987)400
- Nau, H., see Wittfoht, W. 448(1988)433
- Nava, C.C., see Guevara, O.L. 403(1987)350
- Nawata, M., see Tanabe, S. 424(1988)29
- Nawrocki, J.  
Small populations of highly retentive silanols on trimethylsilylated silica surfaces 407(1987)171
- Nawrocki, J. and Buszewski, B.  
Influence of silica surface chemistry and structure on the properties, structure and coverage of alkyl-bonded phases for HPLC (Review) 449(1988)1
- Nayar, M.S. Balachandran, see Egorin, M.J. 415(1987)148
- Neale, M.E.  
Rapid HPLC method for determination of ethanol and fusel oil in the alcoholic beverage industry 447(1988)443
- Nebinger, P. and Koel, M.  
Automatic determination of serotonin in biological fluids by LC 427(1988)326
- Nebinger, P. and Koel, M.

- Automated analysis of urinary 3-methoxy-4-hydroxy-mandelic acid using ion-pair chromatography and fluorimetric detection 434(1988)43
- Neča, J., Stehlík, F. and Vespalec, R.  
Characterization of mobile phases for the investigation of electrokinetic phenomena in LC 447(1988)177
- Necciari, J., see Demotes-Mainard, F. 419(1987)388
- Negri, A., see Biondi, P.A. 411(1987)275
- Nehmer, U.  
Simultaneous determination of atracurium besylate and its major decomposition products and related impurities by reversed-phase HPLC 435(1988)425
- Neicheva, A., Kovacheva, E. and Marudov, G.  
Determination of organophosphorus pesticides in apples and water by GLC with electron-capture detection 437(1988)249
- Neidhart, B., see Blaszkewicz, M. 439(1988)109
- Nelis, H.J., see De Leenheer, A.P. 429(1988)3
- Nelson, D.B., see Philip, T. 442(1988)249
- Němeček, J., see Beran, M. 442(1988)431
- Netting, A.G., see Rhodes, S.H. 442(1988)412
- Netting, A.G., see Rhodes, S.H. 448(1988)135
- Neuhoff, V., see Taube, D. 437(1988)411
- Neumeyer, J.L., see Lampen, P. 426(1988)283
- Neurath, G.B. and Pein, F.G.  
GC determination of *trans*-3'-hydroxycotinine, a major metabolite of nicotine in smokers 415(1987)400
- Nevejans, F. and Verzele, M.  
On the structure and chromatographic behaviour of polystyrene phases 406(1987)325
- Newland, P., see Thomas, A.H. 410(1987)373
- Newman, H.C., see Zuk, M.M. 433(1988)367
- Newton, A., see Hunter, K. 435(1988)83
- Newton, C.J., Roberts, N.B. and Taylor, W.H.  
Separation of human pepsins in gastric juice by high-performance ion-exchange chromatography 417(1987)391
- Ng, K.T., see Holland, M.L. 433(1988)276
- Ng, L.L., see Stubbs, R.J. 413(1987)171
- Ng, P., see Pereira, C.M. 425(1988)208
- Nguyen-Khac, M., see Lamiable, D. 434(1988)315
- Nguyen, T., see Hearn, M.T.W. 435(1988)271
- Nguyen, T.Q. and Kausch, H.-H.  
Absolute peak broadening calibration in SEC using a polymer-bound chromophore 449(1988)63
- Nguyen, V., see Samu, J. 435(1988)167
- Nguyen, V.K., see Rihn, B. 428(1988)408
- Ni, P., Guyon, F., Caude, M. and Rosset, R.  
Automatic determination of urinary 4-hydroxy-3-methoxymandelic (vanillylmandelic) acid by LC with electrochemical detection 424(1988)255
- Nicholls, P.J., see Pourgholami, M.H. 424(1988)163
- Nicholson, G.J., see Koppenhoefer, B. 406(1987)157
- Nickless, G., see Al-Attar, A.F. 440(1988)333
- Nicolaeva, L.A., see Manoilov, S.E. 440(1988)53
- Nicolas, A., see Decolin, D. 414(1987)499
- Nicolas, A., see Decolin, D. 419(1987)469
- Nicolas, A., see Motassim, N. 422(1987)340
- Nicolas, P., see Ennachachibi, A. 427(1988)307
- Nicoletti, I., see Polcaro, C. 448(1988)127
- Nicoletti, J., see Swanson, S.P. 414(1987)335
- Nicoll-Griffith, D.A.  
Stereoelectronic model to explain the resolution of enantiomeric ibuprofen amides on the Pirkle chiral stationary phase 402(1987)179
- Nicoll-Griffith, D.A., Inaba, T., Tang, B.K. and Kalow, W.  
Method to determine the enantiomers of ibuprofen from human urine by HPLC 428(1988)103
- Nieder, M. and Jaeger, H.  
Quantification of tamoxifen and N-desmethyltamoxifen in human plasma by HPLC, photochemical reaction and fluorescence detection, and its application to biopharmaceutical investigations 413(1987)207
- Nieder, M. and Jaeger, H.  
Quantification of gallopamil and norgallopamil in human plasma by HPLC with fluorescence detection 414(1987)492
- Nieder, M. and Jaeger, H.  
Sensitive quantification of pseudoephedrine in human plasma and urine by HPLC 424(1988)73
- Nielen, M.W.F., Van Soest, R.E.J., Van Ingen, H.E., Farjam, A., Frei, R.W. and Brinkman, U.A.T.  
Selective on-line trace enrichment for the determination of ethynyl steroids in urine by LC with precolumn technology 417(1987)159
- Nielsen, P., see Birsner, U. 402(1987)381
- Nielsen, R.G. and Wilson, G.S.  
Characterization of adsorption on the stationary phase using high-performance immunoaffinity chromatography 423(1987)41
- Nielsen, T.H., see Larsen, L.M. 450(1988)121
- Nieman, T.A., see Koerner, Jr., P.J. 449(1988)217
- Niemelä, K.  
GLC-MS studies on pine Kraft black liquors.

- III. The liberation of carboxylic acids in the initial phase of pulping 446(1988)247
- Nierenberg, D.W., Lester, D.C. and Colacchio, T.A.  
Determination of tocopherol and tocopherol acetate concentrations in human feces using HPLC 413(1987)79
- Niessen, H.J.J.M., see Evelo, C.T.A. 420(1987)35
- Nightingale, K.A., see McGarrity, M.J. 413(1987)53
- Nightingale, K.A., see McGarrity, M.J. 415(1987)136
- Nijenhuis, M., see Maris, F.A. 435(1988)297
- Nikitin, Y.S., Ro, R.T. and Shonia, N.K.  
GC study of the temperature dependence of the hydrogen bond energy of adsorption on silica 446(1988)55
- Nikolov, R.N.  
Evaluation of interfacial surface areas in ultra-low-loaded GC columns 446(1988)221
- Nikolova-Damyanova, B., see Amidzhin, B. 446(1988)259
- Nikolova-Damyanova, B. and Amidzhin, B.  
Separation of triglyceride groups by reversed-phase TLC on silanized Kieselguhr 446(1988)283
- Nilsson-Ehle, I.  
Assay of ciprofloxacin and norfloxacin in serum and urine by HPLC 416(1987)207
- Nilsson, L.-G., see Höglund, P. 414(1987)109
- Nilsson, L.-G., Walldorf, B. and Paulsen, O.  
Determination of erythromycin in human plasma, using column LC with a polymeric packing material, alkaline mobile phase and amperometric detection 423(1987)189
- Nilsson, L.B.  
Reversed-phase ion-pair LC method for the determination of low concentrations of haloperidol in plasma 431(1988)113
- Nimura, N., Iwaki, K. and Kinoshita, T.  
Separation of norepinephrine enantiomers, DOPA enantiomers and dopamine derivatized with *o*-phthalaldehyde-*N*-acetyl-L-cysteine by HPLC 402(1987)387
- Nimura, N., see Iwaki, K. 404(1987)117
- Nimura, N., see Iwaki, K. 407(1987)273
- Nimura, N., see Yoshida, T. 445(1988)175
- Ninfali, P., see Mangani, F. 437(1988)294
- Nir, T., see Harduf, Z. 437(1988)379
- Nishi, H., see Fujimoto, Y. 402(1987)344
- Nishida, T., see Miwa, H. 416(1987)237
- Nishikata, M.  
Separation of  $\beta$ -amino alcohols as their dansyl derivatives by TLC on a polyamide layer sheet 408(1987)449
- Nishikawa, H., see Motohashi, N. 435(1988)335
- Nishikawa, T., see Kobayashi, Y. 430(1988)65
- Nishikawa, T., see Onodera, S. 409(1987)259
- Nishimoto, M., see Sakamoto, M. 411(1987)259
- Nishimura, K., see Shinbo, T. 405(1987)145
- Nishimura, R., see Kominami, G. 417(1987)216
- Nishimura, Y., see Haginaka, J. 447(1988)365
- Nissinen, E., see Honkasalo, T. 424(1988)136
- Nitsche, V., Schütz, H. and Eichinger, A.  
Rapid HPLC determination of nifedipine in plasma with on-line precolumn solid-phase extraction 420(1987)207
- Nitta, T., see Honda, S. 438(1988)73
- Niwa, N.T.T., Maeda, K., Shibata, M. and Tatematsu, A.  
Rapid assay for indoxyl sulphate in uremic serum by internal-surface reversed-phase HPLC 431(1988)418
- Niwa, T., see Sasaoka, T. 428(1988)152
- Niwa, T., see Takeuchi, T. 405(1987)117
- Niwa, T., see Takeuchi, T. 407(1987)141
- Njar, V.C.O., Dharmaratne, H.R.W. and Caspi, E.  
Synthesis of [16,16,19-<sup>2</sup>H<sub>3</sub>; 19-<sup>3</sup>H]19-oxo-androst-4-ene-3,17-dione 440(1988)415
- Nobile, S. and Deshusses, J.  
Detection of ligand-protein binding by direct electrophoresis of the complex 449(1988)331
- Nobilis, M. and Hais, I.M.  
Direct absorptiometric quantitation in HPLC in the absence of concomitant reference standards 434(1988)363
- Noctor, T.A.G., see Fell, A.F. 434(1988)377
- Noda, H., Saitoh, K. and Suzuki, N.  
Elution recovery of tris(acetylacetonato)aluminium(III), -gallium(III) and -indium(III) in SEC 435(1988)368
- Noda, M., see Kudo, K. 431(1988)353
- Noda, M., see Uehori, R. 411(1987)251
- Noda, Y., see Shibasaki, S. 421(1987)425
- Noël, D., Cole, K.C., Hechler, J.-J., Chouliotis, A. and Overbury, K.C.  
High-performance SEC of fiber-reinforced organic-matrix composites 408(1987)129
- Noël, D., Cole, K.C. and Hechler, J.-J.  
Quantitative analysis of resins used in fiber-reinforced composites by reversed-phase LC 447(1988)141
- Noel, L. and Ménard, H.  
Improved method for determining adsorption isotherms by LC on long and short fibrous material 436(1988)391
- Noguchi, H., see Falkowski, A.J. 422(1987)145
- Noguchi, H., see Terada, T. 430(1988)388

- Noguchi, H., see Tokuma, Y. 415(1987)156  
 Noguchi, K., see Yasukawa, K. 410(1987)129  
 Noguchi, K.D., see Hirata, N. 434(1988)71  
 Noguchi, M., see Oka, K. 423(1987)285  
 Nohta, H., see Lee, M. 415(1987)289  
 Nohta, H., see Lee, M.-K. 421(1987)237  
 Nojima, K., see Shimizu, K. 432(1988)21  
 Nolet, S., see Lemaire, S. 425(1988)77  
 Nolte, J., see Krupčík, J. 448(1988)203  
 Nomura, A., Yamada, J. and Tsunoda, K.-I.  
 Effect of the pore structure of silica gel on retention behaviour in pressure programming supercritical fluid chromatography 448(1988)87  
 Nomura, M., see Yamamoto, S. 409(1987)101  
 Nomura, T., see Haginaka, J. 447(1988)268  
 Nomura, T., see Haginaka, J. 447(1988)373  
 Nonhebel, H.M.  
 Direct separation of (*S*)- and (*R*)-abscisic acid on a commercially available chiral HPLC column 402(1987)374  
 Norbury, C.G.  
 Simplified method for the determination of plasma cotinine using GC-MS 414(1987)449  
 Nordquist, R.E., see Gherezghiher, T. 413(1987)9  
 Nordström, M., see Naaranlahti, T. 410(1987)488  
 Norén, K. and Sjövall, J.  
 Analysis of pentachlorophenol in water and urine by enrichment with Lipidex 5000 414(1987)55  
 Norén, K. and Sjövall, J.  
 Analysis of organochlorine pesticides, polychlorinated biphenyls, dibenzo-*p*-dioxins and dibenzofurans in human milk by extraction with the lipophilic gel Lipidex 5000 422(1987)103  
 Norman, L.M., see Bannon, C.D. 404(1987)340  
 Norman, L.M., see Bannon, C.D. 407(1987)231  
 Norman, L.M., see Bannon, C.D. 447(1988)43  
 Norris, B., see Bauer, J. 445(1988)429  
 Nose, N., see Horie, M. 402(1987)301  
 Notarianni, L.J. and Collins, A.J.  
 Method for the determination of acemetacin, a non-steroidal anti-inflammatory drug, in plasma HPLC 413(1987)305  
 Notarianni, L.J. and Jones, R.W.  
 Method for the determination of ofloxacin, a quinolone carboxylic acid antimicrobial, by HPLC 431(1988)461  
 Nouveau, J., see Lacroix, C. 415(1987)85  
 Novák, I., see Buszewski, B. 446(1988)191  
 Novák, J., see Beran, M. 442(1988)431  
 Novák, J., see Koupil, P. 425(1988)99  
 Nováková, E., see Balíková, M. 431(1988)431  
 Novas, M.L., see Peña, C. 431(1988)192  
 Novotny, M., see Andreolini, F. 406(1987)375  
 Novotny, M., see Flurer, C.L. 448(1988)73  
 Nowotny, A., see Samu, J. 435(1988)167  
 Nowotny, H.-P., see Schurig, V. 441(1988)155  
 Nozaki, S., see Tamaki, M. 413(1987)287  
 Nugent, K.D., Burton, W.G., Slattery, T.K., Johnson, B.F. and Snyder, L.R.  
 Separation of proteins by reversed-phase HPLC. II. Optimizing sample pretreatment and mobile phase conditions 443(1988)381  
 Nugent, K.D., see Burton, W.G. 443(1988)363  
 Numakura, T., see Yamashita, J. 403(1987)275  
 Nunoi, K., see Miwa, H. 416(1987)237  
 Nurmela, K.V.V. and Satama, L.T.  
 Quantitative analysis of triglycerides by HPLC using non-linear gradient elution and flame ionization detection 435(1988)139  
 Nyberg, G., see Svensson, C. 432(1988)363  
 Nygard, G.A., Lovett, L.J., Erdmann, G.R. and Khalil, S.K.W.  
 Analysis of danazol in serum using HPLC 415(1987)438  
 Nyiredy, S., Dallenbach-Toelke, K. and Sticher, O.  
 Applicability of forced-flow planar chromatographic methods for the separation of enantiomers on Chiralplate 450(1988)241  
 Nyiredy, S., see Dallenbach-Toelke, K. 404(1987)365  
 Nyiredy, S., see Hofstetter, M. 407(1987)319

## O

- Obata, K., see Hirabayashi, Y. 445(1988)377  
 Očekpe, N.A., see Taylor, R.B. 416(1987)394  
 Ochoa, J.L. and Ascencio-Valle, F.  
 Difference in cell surface hydrophobicity of *Halobacterium salinarum* strains 449(1988)322  
 Ockhuizen, T., see Veenstra, J. 431(1988)413  
 Oda, K., see Matsanaga, H. 422(1987)353  
 Oda, S., see Nakano, K. 438(1988)61  
 Odham, G., see Sonesson, A. 417(1987)11  
 Odham, G., see Sonesson, A. 431(1988)1  
 Odham, G., Valeur, A., Michelsen, P., Aronsson, E. and McDowall, M.  
 Highly sensitive determination and characterization of intact cellular ester-linked phospholipids using LC-plasma spray MS 434(1988)31

- Odink, J., Bogaards, J.J.P., Sandman, H., Egger, R.J., Arkesteyn, G.A. and De Jong, P.  
Excretion of iodide in 24-h urine as determined by ion-pair reversed-phase LC with electrochemical detection 431(1988)309
- Odink, J., Korthals, H. and Knijff, J.H.  
Simultaneous determination of the major acidic metabolites of catecholamines and serotonin in urine by LC with electrochemical detection after a one-step sample clean-up on Sephadex G-10; influence of vanilla and banana ingestion 424(1988)273
- O'Donovan, G.A., see Dutta, I. 444(1988)183
- Oe, T., see Shimada, K. 419(1987)17
- Oetelaar, P.J.M. Van den and Hoenders, H.J.  
Electrophoretic titration curve in 6 M urea of the bovine eye lens protein  $\alpha$ -crystallin 411(1987)507
- Ogata, H., see Ishii, M. 426(1988)412
- Ogata, H., see Takahashi, H. 419(1987)243
- Ogawa, H., see Chiu, W.-Y. 428(1988)25
- Ogawa, H., see Kanetoshi, A. 442(1988)289
- Ogawa, K., Fujiwara, Y., Sugamata, K. and Abe, T.  
TLC of neutral glycosphingolipids: an improved simple method for the resolution of GlcCer, GalCer, LacCer and Ga<sub>2</sub>Cer 426(1988)188
- Ogihara, Y., see Umeda, M. 422(1987)246
- Ogino, S., see Yanagi, H. 417(1987)178
- Ogunkolade, B.W., Monjour, L., Vouldoukis, I., Rhodes-Feuillette, A. and Frommel, D.  
Inoculation of BALB/c mice against *Leishmania major* infection with *Leishmania*-derived antigens isolated by gel filtration 440(1988)459
- Ogunkolade, W., see Jaureguiberry, G. 440(1988)385
- Ogura, H., see Iwaki, K. 404(1987)117
- Ogura, H., see Iwaki, K. 407(1987)273
- O'Hara, D.S., see Kelly, R.A. 416(1987)247
- Ohe, T., see Yoshikawa, S. 421(1987)111
- Ohishi, K., see Kataoka, H. 420(1987)135
- Ohkura, Y., see Chao, W.-F. 430(1988)361
- Ohkura, Y., see Ishida, J. 431(1988)249
- Ohkura, Y., see Kai, M. 430(1988)271
- Ohkura, Y., see Lee, M. 415(1987)289
- Ohkura, Y., see Lee, M.-K. 421(1987)237
- Ohkura, Y., see Nakano, M. 411(1987)305
- Ohkura, Y., see Ohno, M. 421(1987)245
- Ohkura, Y., see Ohno, M. 430(1988)291
- Ohkura, Y., see Tsuruta, Y. 416(1987)63
- Ohkura, Y., see Wang, Z.-J. 430(1988)223
- Ohlendorf, R., see Blaszkewicz, M. 439(1988)109
- Ohloff, G. and Demole, E.  
Importance of the odoriferous principle of Bulgarian rose oil in flavour and fragrance chemistry 406(1987)181
- Ohmori, M., see Kataoka, H. 420(1987)135
- Ohmori, S., Kawase, M., Mori, M. and Hirota, T.  
Simple and sensitive determination of methylglyoxal in biological samples by GC with electron-capture detection 415(1987)221
- Ohmori, S., Mori, M., Kawase, M. and Tsuboi, S.  
Determination of methylglyoxal as 2-methylquinoxaline by HPLC and its application to biological samples 414(1987)149
- Ohmori, S., Sumii, I., Toyonaga, Y., Nakata, K. and Kawase, M.  
HPLC determination of formate as benzimidazole in biological samples 426(1988)15
- Ohmori, S. and Iwamoto, T.  
Sensitive determination of D-lactic acid in biological samples by HPLC 431(1988)239
- Ohmura, T., see Honda, S. 419(1987)51
- Ohnhaus, E.E., see Santos, S.R.C.J. 419(1987)155
- Ohnishi, S., see Tanaka, N. 448(1988)95
- Ohno, M., Kai, M. and Ohkura, Y.  
On-line post-column fluorescence detection for N-terminal tyrosine-containing peptides in HPLC 421(1987)245
- Ohno, M., Kai, M. and Ohkura, Y.  
Assay for enkephalin-degrading peptidases in rat brain tissues by HPLC with on-line post-column fluorescence detection 430(1988)291
- Ohno, M., see Nakano, M. 411(1987)305
- Ohno, T., see Oka, H. 411(1987)437
- Ohta, H., Kinjo, S. and Osajima, Y.  
Glass capillary GC analysis of volatile components of canned Philippine pineapple juice 409(1987)409
- Ohta, M., Fukuda, H., Kimura, T. and Tanaka, A.  
Quantitative analysis of oxytocin in pharmaceutical preparations by HPLC 402(1987)392
- Ōi, S., see Yamashita, J. 403(1987)275
- Oi, T., see Hosoe, M. 435(1988)253
- Oi, T., see Hosoe, M. 438(1988)225
- Oi, T., Tsukamoto, T., Akai, H., Kakihana, H. and Hosoe, M.  
Boron isotope separation by ion-exchange chromatography using an anion-exchange resin in halide forms. Separation factors at 25 °C 450(1988)343
- Oishi, M., see Yamaji, A. 415(1987)143
- Oishi, M., see Yamaji, A. 421(1987)372

- Oishi, S., see Harada, K.-I. 448(1988)275
- Ōizumi, H., see Arai, T. 448(1988)439
- Oizumi, J., see Hayakawa, K. 413(1987)247
- Oizumi, J., see Hayakawa, K. 423(1987)304
- Oka, H., Ikai, Y., Kawamura, N., Yamada, M., Harada, K.-I., Yamazaki, Y. and Suzuki, M. Improvement of chemical analysis of antibiotics. XIV. Identification of the components of bacitracin using normal- and reversed-phase TLC 449(1988)448
- Oka, H., Ikai, Y., Kawamura, N., Yamada, M., Inoue, H., Ohno, T., Inagaki, K., Kuno, A. and Yamamoto, N. Simple method for the analysis of food dyes on reversed-phase thin-layer plates 411(1987)437
- Oka, H., see Harada, K.-I. 448(1988)275
- Oka, H., see Ikai, Y. 411(1987)313
- Oka, K., Hirano, T. and Noguchi, M. Combined use of rapid-flow fractionation and HPLC for the determination of serum testosterone and application to study of stress response to physical exercise 423(1987)285
- Oka, K., see Hirano, T. 417(1987)371
- Okada, A., see Yonekura, T. 427(1988)320
- Okada, T. Elution characteristics of polyol-borate eluents in anion-exchange chromatography. I. Eluent strength of sugar-, uronate-borate eluents 403(1987)27
- Okada, T., see Matsukura, H. 414(1987)47
- Okada, T., see Matsukura, H. 434(1988)196
- Okada, T., see Tan, S. 447(1988)198
- Okada, T. and Kuwamoto, T. Elution characteristics of polyol-borate eluents in anion-exchange chromatography. II. Tartrate-borate eluent for non-suppressed anion chromatography 403(1987)35
- Okada, Y., see Kitano, H. 420(1987)13
- Okada, Y., see Koizumi, K. 437(1988)47
- Okada, Y., see Kurumaya, K. 435(1988)235
- Okamoto, M., Jinno, K. and Omori, T. Preparation and evaluation of mono-, di- and triphenyl-treated plates for the HPTLC analysis of puerarin and paeonol 407(1987)314
- Okamoto, M., see Jinno, K. 402(1987)173
- Okamoto, M., see Jinno, K. 436(1988)1
- Okamoto, T., Fukunaga, Y., Ida, Y. and Kishi, T. Determination of reduced and total ubiquinones in biological materials by LC with electrochemical detection 430(1988)11
- Okamoto, Y., Aburatani, R. and Hatada, K. Chromatographic resolution. XXI. Direct optical resolution of abscisic acid by HPLC on cellulose tris(3,5-dimethylphenylcarbamate) 448(1988)454
- Okano, T. and Lawson, D.E.M. Chick 28000 *M<sub>r</sub>* vitamin D-dependent calcium-binding protein in intestine, kidney and cerebellum. Purification using chromatofocusing 448(1988)145
- Okatani, Y., see Sagara, Y. 431(1988)170
- Okazaki, M., Kinoshita, M. and Hara, I. Improved HPLC method for the determination of apolipoproteins in serum high-density lipoproteins 430(1988)135
- O'Keefe, S.F. and Murphy, P.A. Rapid determination of vitamin D in fortified skim milk 445(1988)305
- O'Kennedy, R., see Carty, P. 442(1988)279
- O'Kennedy, R., see Moran, E. 416(1987)165
- Okerholm, R.A., see Walker, B.J. 416(1987)150
- Okigaki, T., see Kanemitsu, F. 429(1988)399
- Okoyama, A. Fluorimetric determination of urinary  $\delta$ -aminolevulinic acid by HPLC and post-column derivatization 426(1988)365
- Okuda, S., see Yoshikawa, S. 421(1987)111
- Okuda, T., see Hatano, T. 435(1988)285
- Okudzhava, N.G., see Andronikashvili, T.G. 406(1987)145
- Okumura, K., see Takagi, K. 432(1988)47
- Okuno, A., see Tanaka, N. 448(1988)95
- Okuyama, S., see Takagi, K. 432(1988)47
- Okuyama, S. and Ikeda, Y. Determination of acetylcholine and choline in human cerebrospinal fluid using HPLC combined with an immobilized enzyme reactor: ageing-induced change of acetylcholine level 431(1988)389
- Okuyama, T., see Manabe, T. 423(1987)115
- Okuyama, T., see Manabe, T. 431(1988)45
- Oldfield, N., Chang, D., Garland, W. and Town, C. Quantitation of cefetrame in human plasma and urine by HPLC 422(1987)135
- Olek, M. Determination of diphenylamine residues in apples, and 4-aminobiphenyl residues in diphenylamine, by HPLC and electrochemical detection 447(1988)421
- Oliva, A., see Della Ragione, F. 440(1988)141
- Oliva, A., see Porcelli, M. 440(1988)151
- Oliva, B., see Amicosante, G. 403(1987)366
- Olivares, J. and Verdys, M. Isocratic HPLC method for studying the metabolism of blood plasma pyrimidine



- nucleosides and bases: concentration and radioactivity measurements 434(1988)111
- Oliveros, L. and Cazau, M.  
Resolution of a series of racemic phosphine oxides by LC on two structurally similar chiral stationary phases (in French) 409(1987)183
- Oliw, E.H.  
Chromatography of B prostaglandins on  $\beta$ -cyclodextrin silica: application to analysis of major E prostaglandins in human seminal fluid 421(1987)117
- Olling, M., see Derks, H.J.G.M. 442(1988)436
- Olmeda, R., see Karle, J.M. 424(1988)347
- Olson, G.J., see Blair, W.R. 410(1987)393
- Olson, J.A., see Landers, G.M. 438(1988)383
- Olsson, P.-E. and Hogstrand, C.  
Improved separation of perch liver metallothionein by fast protein LC 402(1987)293
- Olsson, U., see Kaul, R. 438(1988)339
- Olszewski, J.J., see LaCreta, F.P. 434(1988)83
- Omichi, K. and Ikenaka, T.  
Use of ion-exchange chromatography and HPLC to prepare a substrate for differential assay of human pancreatic and salivary  $\alpha$ -amylases in serum 420(1987)158
- Omichi, K. and Ikenaka, T.  
Use of a post-column enzymatic reaction for detection of macroamylasemia by HPLC 428(1988)415
- Omori, T., see Hashimoto, K. 438(1988)401
- Omori, T., see Okamoto, M. 407(1987)314
- Omstead, D.R., see Metz, P.A. 441(1988)31
- Ong, C.N., Lee, B.L., Ong, H.Y. and Jaecob, E.  
Simple HPLC method for the simultaneous measurement of vanillylmandelic acid and homovanillic acid 423(1987)278
- Ong, H.Y., see Ong, C.N. 423(1987)278
- Ongini, E., see Dionisotti, S. 434(1988)169
- Ono, K., see Tanaka, M. 426(1988)111
- Onodera, M., see Shibata, K. 430(1988)381
- Onodera, S., Nishikawa, T. and Suzuki, S.  
Chemical changes of organic compounds in chlorinated water. XIV. Characterization and determination of halogenated organics formed during chlorination of water from the Tama River 409(1987)259
- Opacka-Juffry, J., Tacconelli, F. and Coen, C.W.  
Sensitive method for determination of picogram amounts of epinephrine and other catecholamines in microdissected samples of rat brain using LC with electrochemical detection 433(1988)41
- Oparil, S., see Meng, Q.C. 445(1988)29
- Opong-Mensah, K., see Karnes, H.T. 422(1987)165
- Oppitz, M., see Workman, P. 422(1987)315
- Opstal, M.A.J. Van, Krabbenborg, P., Holthuis, J.J.M., Van Bennekom, W.P. and Bult, A.  
Comparison of flow-injection analysis with HPLC for the determination of etoposide in plasma 432(1988)395
- Oratore, A., see Amicosante, G. 403(1987)366
- Orav, A., see Rang, S. 406(1987)75
- O'Reilly, D.J., see Smith, J.D. 431(1988)395
- O'Reilly, T., Mac Mathúna, P., Keeling, P.W.N. and Feely, J.  
Comparison of the spectrophotometric and HPLC analysis of Indocyanine Green in estimating systemic clearance in patients with chronic liver disease 417(1987)190
- Orlowski, R.C., see Heinitz, M.L. 443(1988)229
- Örvell, C., see Welling-Wester, S. 443(1988)255
- Osajima, Y., see Ohta, H. 409(1987)409
- Osborne, R.J., see Joel, S.P. 430(1988)394
- Oshima, J., Horai, Y. and Ishizaki, T.  
HPLC assay of a new non-steroidal anti-inflammatory agent, 2-(10,11-dihydro-10-oxodibenzo[*b,f*]thiepin-2-yl)propionic acid, in human plasma and urine 414(1987)381
- Oshima, R., see Yamauchi, Y. 408(1987)171
- Oshima, T., see Sagara, K. 403(1987)253
- Oshima, T., see Sagara, K. 409(1987)365
- Osman, M.A., Bhavnagri, V.P., Pinchbeck, F.M., Cheng, L.K. and Wright, G.J.  
Determination of duoperone in plasma by HPLC 419(1987)452
- Oss, G.E.C.J.M. Van, see Vree, T.B. 424(1988)440
- Osselton, M.D., see Japp, M. 439(1988)317
- Ossicini, L., see Fanali, S. 403(1987)388
- Ossicini, L., see Fanali, S. 440(1988)361
- Ossicini, L., see Polcaro, C. 448(1988)127
- Osterwald, H., see Brendel, E. 432(1988)358
- Ostrovská, V., see Svobodová, X. 425(1988)391
- Ostrovská, V., Svobodová, X., Pechová, A., Kusala, S. and Svoboda, M.  
Optimization of fluorescence detection for the determination of labetalol in plasma by HPLC 446(1988)323
- Ostrovský, I., see Soják, L. 406(1987)43
- Ostrovský, I., see Soják, L. 436(1988)47
- Ostrovský, I., see Soják, L. 446(1988)339
- Ota, M., see Mefford, I.N. 420(1987)241
- Otagawa, T., Zaromb, S. and Stetter, J.R.  
Cryogenic sampling for aniline vapor with subsequent LC-electrochemical detection analysis 408(1987)81
- Otten, R.R., see De Ruiter, C. 436(1988)429

- Oung, J., see Philp, R.P. 446(1988)3  
 Ovardia, D., see Lesser, J.H. 410(1987)95  
 Overbury, K.C., see Noël, D. 408(1987)129  
 Øvrebø, S., Jacobsen, D. and Sejersted, O.M.  
 Determination of ionic metabolites from  
 ethylene glycol in human blood by  
 isotachopheresis 416(1987)111  
 Owada, K., see Maeda, Y. 410(1987)413  
 Owen, S.G., Roberts, M.S. and Friesen, W.T.  
 Rapid HPLC assay for the simultaneous  
 analysis of non-steroidal anti-inflammatory  
 drugs in plasma 416(1987)293  
 Özcel, M., see Masihi, K.N. 440(1988)473

## P

- Paanakker, J.E., Thio, J.M.S.L.,  
 Van den Wildenberg, H.M. and Kaspersen,  
 F.M.  
 Assay of vecuronium in plasma using solid-  
 phase extraction, HPLC and post-column  
 ion-pair extraction with fluorimetric detection  
 421(1987)327  
 Paanakker, J.E. and Van Hal, H.J.M.  
 Capillary GC assay for the routine monitoring  
 of the antidepressant mepirzepine in human  
 plasma 417(1987)203  
 Pacáková, V., see Mingjia, W. 439(1988)363  
 Pacáková, V., see Štulík, K. 436(1988)334  
 Pacáková, V., Štulík, K. and Pfiřhoda, M.  
 HPLC of *s*-triazines and their degradation  
 products using UV photometric and  
 amperometric detection 442(1988)147  
 Pace, M., see Pietta, P. 411(1987)498  
 Pace, M., see Pietta, P. 441(1988)431  
 Pachla, L.A., see Wright, D.S. 417(1987)223  
 Pączkowski, D., see Filipek, M. 430(1988)406  
 Padmalatha Devi, K., Ranga Rao, K.V., Baveja,  
 S.K., Fathi, M. and Roth, M.  
 Determination of oxprenolol in plasma by  
 column LC 426(1988)229  
 Padmalatha Devi, K., Ranga Rao, K.V., Baveja,  
 S.K., Leemann, T. and Dayer, P.  
 Determination of alprenolol and metoprolol  
 in plasma by column LC 434(1988)265  
 Padrini, R., Piovan, D. and Moretto, R.  
 Column LC determination of prajmalin in  
 plasma and urine by direct sample injection  
 415(1987)183  
 Page, B.D., see Lau, B.P.-Y. 402(1987)233  
 Paik, W.K., see Park, K.S. 440(1988)225  
 Paillas, C., see Leitao, J. 435(1988)229  
 Paladini, A.C., see Peña, C. 431(1988)192  
 Palamareva, M.D.  
 Chromatographic behaviour of  
 diastereoisomers. IX. Application of a  
 microcomputer program to the TLC  
 separation of some diastereoisomers on silica  
 438(1988)219  
 Palitti, F., see Polcaro, C. 448(1988)127  
 Pallast, M., see Kleuss, C. 407(1987)281  
 Palmerini, C.A., Cantelmi, M.G., Minelli, A.,  
 Fini, C., Zampino, M. and Floridi, A.  
 Determination of plasma serotonin by HPLC  
 with pre-column sample enrichment and  
 fluorimetric detection 417(1987)378  
 Palmerini, C.A., Fini, C., Cantelmi, M.G. and  
 Floridi, A.  
 Assessment of taurine in plasma and urine by  
 anion-exchange HPLC with pre-column  
 derivatization 423(1987)292  
 Palmieri, G., see Galli, B. 409(1987)377  
 Palombo, S., see Girre, C. 419(1987)363  
 Palumbo, G., Pantaleoni, G.C. and Carlucci, G.  
 Determination of furprofen in human plasma  
 by solid-phase extraction and reversed-phase  
 HPLC 424(1988)201  
 Palusinski, O.A., see Su, Y. 405(1987)77  
 Palvimo, J., see Karhu, I. 426(1988)65  
 Pan, P.W., see Tan, H.S.I. 404(1987)233  
 Pandey, V.N. and Modak, M.J.  
 Purification of nucleotide-linked peptide  
 444(1988)133  
 Pandher, K.S., see Harvey, D.J. 414(1987)253  
 Pang, K.S., see St. Pierre, M.V. 421(1987)291  
 Pang, K.S., see Xu, X. 420(1987)313  
 Pannell, L.K., see Zhang, T.-Y. 442(1988)455  
 Pannell, L.K., see Zhang, T.-Y. 445(1988)199  
 Pannella, M., see Munavalli, S. 437(1988)423  
 Panse, D.G., Bhale, A., Gumaste, V.K., Mane,  
 M.V., Likhite, S.M. and Bapat, B.V.  
 Liquid crystals. VI. Comparative study of the  
 supercooling effect of some liquid crystalline  
 stationary phases using GLC 411(1987)456  
 Pantaleoni, G.C., see Palumbo, G. 424(1988)201  
 Pao, J., see Puglisi, C.V. 419(1987)253  
 Paone, R., see Holland, O.B. 426(1988)148  
 Papas, A.N., see Delaney, M.F. 410(1987)31  
 Papet, Y., see Mura, P. 416(1987)303  
 Papp, E.  
 System peaks in ion chromatography using  
 cetrimide-coated columns and indirect  
 photometric detection 402(1987)211  
 Papp, E. and Fehérvári, A.

- Anion chromatography using a coated PRP-1 column and eluents of pH > 7 447(1988)315
- Papp, O., Józán, M., Szász, G. and Hermecz, I.  
Correlation between molar refraction and GC retention data 403(1987)11
- Papp, O., Szász, G., Farkas, M., Simon, G. and Hermecz, I.  
Some problems in the correlation of molecular parameters and the connectivity index 403(1987)19
- Papparella, V., see Della Ragione, F. 440(1988)141
- Pappas, A.A., see Taylor, E.H. 416(1987)365
- Pappas, S., see Driscoll, J.N. 441(1988)63
- Pappas, S., see Duffy, M. 441(1988)73
- Paredes, J., see Paulos, C. 433(1988)359
- Paris, J., see Jaussaud, P. 423(1987)123
- Park, K.S., Lee, H.W., Hong, S.-Y., Shin, S., Kim, S. and Paik, W.K.  
Determination of methylated amino acids in human serum by HPLC 440(1988)225
- Parker, C.E., Geeson, A.V., Games, D.E., Ramsey, E.D., Abusteit, E.O., Corbin, F.T. and Tomer, K.B.  
Thermospray LC-MS method for the analysis of metribuzin and its metabolites 438(1988)359
- Parkin, J.E.  
Assay of phenylmercury salts in pharmaceutical products by HPLC of the morpholinedithiocarbamate derivative 407(1987)389
- Parks, C., see Gilbert, J. 441(1988)323
- Parks, E.J., see Blair, W.R. 410(1987)393
- Paroni, R., see Fermo, I. 433(1988)53
- Paroni, R., Sirtori, C.R., Borghi, C. and Kienle, M.G.  
HPLC determination of ribavirin in serum and urine and of its urinary metabolite 1,2,4-triazole-3-carboxamide 420(1987)189
- Parris, N., Gruber, H.A., Mozersky, S.M. and Barford, R.A.  
Purification of cytochrome *c* oxidase from corn root mitochondria by hydrophobic interaction chromatography 444(1988)219
- Parrish, C.C., Zhou, X. and Herche, L.R.  
Flame ionization and flame thermionic detection of carbon and nitrogen in aquatic lipid and humic-type classes with an Iatroscan Mark IV 435(1988)350
- Parrish, S.W., see Jenkins, R.L. 426(1988)249
- Parrot, H., see Coleman, A.W. 450(1988)175
- Parry, D.A., see Lloyd, J.B.F. 449(1988)281
- Parshina, I.V., see Gavrilova, T.B. 402(1987)87
- Partridge, L.Z., see Euerby, M.R. 445(1988)433
- Partridge, L.Z., see Euerby, M.R. 447(1988)392
- Partyka, S., see Rudziński, W. 406(1987)295
- Parviainen, M.T. and Barlow, R.D.  
Assessment of exposure to environmental tobacco smoke using a HPLC method for the simultaneous determination of nicotine and two of its metabolites in urine 431(1988)216
- Paschold, B., see Krupčík, J. 448(1988)203
- Pascoe, G.A., Duda, C.T. and Reed, D.J.  
Determination of  $\alpha$ -tocopherol and  $\alpha$ -tocopherylquinone in small biological samples by HPLC with electrochemical detection 414(1987)440
- Pasechnik, V.A., see Gorbunov, A.A. 448(1988)307
- Passingham, B.J. and Barton, R.N.  
Application of HPLC to the measurement of cortisol secretion rate 416(1987)25
- Pastor, A., see Mateo, R. 410(1987)319
- Pasutto, F.M., see Coutts, R.T. 421(1987)267
- Pasutto, F.M., see Mehvar, R. 425(1988)135
- Patchen, L.C., see Mount, D.L. 423(1987)261
- Patchen, L.C., see Mount, D.L. 428(1988)196
- Patchen, L.C., see Williams, S.B. 433(1988)197
- Patchett, M.L., Monk, C.R., Daniel, R.M. and Morgan, H.W.  
Determination of agmatine, arginine, citrulline and ornithine by reversed-phase LC using automated pre-column derivatization with *o*-phthalaldehyde 425(1988)269
- Patel, I., see Lumley, I.D. 408(1987)115
- Paterson, S.  
Measurement of dipipanone using capillary GC 424(1988)152
- Patrick, J.E., see Alton, K.B. 425(1988)363
- Patrykiewicz, A., see Gawdzik, B. 448(1988)233
- Patsalides, E., see Robards, K. 411(1987)1
- Pattabiraman, N., see Armstrong, R.D. 414(1987)192
- Patte, F., see Laffort, P. 406(1987)51
- Patterson, Jr., D.G., see Barnhart, E.R. 445(1988)145
- Patthy, M. and Gyenge, R.  
Perfluorinated acids as ion-pairing agents in the determination of monoamine transmitters and some prominent metabolites in rat brain by HPLC with amperometric detection 449(1988)191
- Paul, P.S., see Stowers, D.J. 444(1988)47
- Paulos, C., Paredes, J., Vasquez, I., Kunze, G. and González-Martín, G.  
HPLC determination of nifurtimox in human serum 433(1988)359
- Paulsen, O., see Nilsson, L.-G. 423(1987)189
- Paulson, J., see Edholm, L.-E. 424(1988)61

- Paulus, A., see Guttman, A. 448(1988)41
- Pauly, F., see Menu, P. 417(1987)397
- Pavel, S., see Van Haard, P.M.M. 429(1988)59
- Pavlova-Rouseva, R., see Rachev, R.C. 437(1988)287
- Pavlova, V.F., see Shadrina, N.E. 404(1987)183
- Payet, B., Arnoux, P., Catalin, J. and Cano, J.P.  
Direct determination of mitoxantrone and its mono- and dicarboxylic metabolites in plasma and urine by HPLC 424(1988)337
- Pays, M., see Naline, E. 419(1987)177
- Pays, M., see Spreux-Varoquaux, O. 416(1987)311
- Peal, J.A., see Bailey, E. 416(1987)81
- Peal, J.A., see Bailey, E. 425(1988)187
- Pearce, J.C., see Doyle, E. 411(1987)325
- Pearson, C.D.  
Determination of phenolic antioxidants in JP-5 jet fuels by GC-mass selective detection 449(1988)440
- Pearson, J.D., MacCroskey, M.C. and DeWald, D.B.  
Separation of protein hormones (Review) 418(1987)245
- Pearson, J.D., see Gisch, D.J. 443(1988)299
- Pearson, J.D., see McCroskey, M.C. 442(1988)307
- Peaston, R.T.  
Routine determination of urinary free catecholamines by HPLC with electrochemical detection 424(1988)263
- Pecci, L., Costa, M., Pinnen, F., Antonucci, A. and Cavallini, D.  
Specific method for the determination of 1,4-thiomorpholine-3,5-dicarboxylic acid and its unsaturated analogue lanthionine ketimine 426(1988)183
- Pechová, A., see Ostrovská, V. 446(1988)323
- Pechová, A., see Svobodová, X. 425(1988)391
- Peck, K. and Morris, M.D.  
Optical errors in an LC absorbance cell 448(1988)193
- Pedrazzini, S., Zanononi-Muciaccia, W., Sacchi, C. and Forgione, A.  
Determination of flunoxaprofen enantiomers in biological fluids by HPLC 415(1987)214
- Pedrazzini, S., Zanononi-Muciaccia, W. and Forgione, A.  
Determination of the non-steroidal anti-inflammatory drug flunoxaprofen, *S*(+)-2-(4-fluorophenyl)- $\alpha$ -methyl-5-benzoxazoleacetic acid, in blood and urine by HPLC 413(1987)338
- Pedroso, E., Grandas, A., Amor, J.-C. and Giralt, E.  
Reversed-phase HPLC of protected peptide segments 409(1987)281
- Pefferkorn, E., see Voegel, J.C. 428(1988)17
- Pehal, F., see Verner, P. 416(1987)439
- Pein, F.G., see Neurath, G.B. 415(1987)400
- Peinado, J., López-Soriano, F.J. and Argilés, J.M.  
GC method for the estimation of acetone and its metabolites in biological samples 415(1987)372
- Pelizzoni, F., see Muda, M. 434(1988)191
- Pellegrin, P.L., see Lesne, M. 415(1987)197
- Pellegrino, S., see Petrarulo, M. 432(1988)37
- Pelletier, S.W., see Kulanthaivel, P. 402(1987)366
- Pelletier, X., see Leray, C. 420(1987)411
- Peltonen, K., see Pukkila, J. 411(1987)409
- Peltonen, K., see Pukkila, J. 438(1988)443
- Peltonen, K. and Pukkila, J.  
Determination of bisphenol A in air by HPLC with electrochemical detection 439(1988)375
- Peña, C., Medina, J.H., Novas, M.L., De Robertis, E. and Paladini, A.C.  
Determination of *n*-butyl- $\beta$ -carboline-3-carboxylate in aqueous brain extracts 431(1988)192
- Peng, C.T., Ding, S.F., Hua, R.L. and Yang, Z.C.  
Prediction of retention indexes. I. Structure-retention index relationship on apolar columns 436(1988)137
- Percheron, F., see Ricard, B. 428(1988)203
- Perego, F., see Musu, C. 449(1988)432
- Pereira, C.M., Tam, Y.K., Collins-Nakai, R.L. and Ng, P.  
Simplified determination of captopril in plasma by HPLC 425(1988)208
- Peretti, E., see Karlaganis, G. 420(1987)171
- Perez, M., see Poirier, J.-M. 426(1988)431
- Perez, M., see Poirier, J.M. 421(1987)223
- Perin, A., see Desiderio, M.A. 419(1987)285
- Perkal, M.B., see Emm, T. 419(1987)445
- Perkal, M.B., see Emm, T. 427(1988)188
- Perkal, M.B., see Emm, T.A. 427(1988)162
- Perkins, D.M. and Duncan, J.R.  
Improved isolation of rat microsomal cytochrome *b*<sub>5</sub> reductase 405(1987)319
- Perkins, L.M., see Kling, G.J. 407(1987)377
- Perkins, R., see Knight, M. 444(1988)345
- Perly, B., see Coleman, A.W. 450(1988)175
- Pernelle, J.-J., see Rabilloud, T. 402(1987)105
- Pernollet, J.C., see Billard, V. 440(1988)87
- Peroni, A.E., see Schinelli, S. 431(1988)150
- Perret, G., see Ennachachibi, A. 427(1988)307
- Perrin, J.H., see Townsend, R.W. 450(1988)414
- Perry, H.E., see Schmid, B.J. 425(1988)107
- Person, M., see Guimbar, J.P. 403(1987)109
- Persson, B.-A., see Balmér, K. 417(1987)357
- Persson, B., see Wannenberg, O. 435(1988)199

- Pertierra-Rimada, E., see Fernández-Sánchez, E. 402(1987)318
- Pertierra-Rimada, E., see Fernández-Sánchez, E. 410(1987)13
- Pesselman, R.L. and Feit, M.J.  
Determination of residual epichlorohydrin and 3-chloropropanediol in water by GC with electron-capture detection 439(1988)448
- Pestel, B.C., see Duff, G.A. 416(1987)71
- Peteghem, C. Van, see Rosier, J. 434(1988)222
- Péter, A., Szepesi, G., Balásperi, L. and Burger, K.  
Coordinative interactions in the separation of insulin and derivatives by HPLC 408(1987)43
- Peter, H.H., see Jenny, H.B. 438(1988)433
- Peters, T.J., see Lim, C.K. 429(1988)123
- Peters, W.P., see Henner, W.D. 416(1987)426
- Peterson, R.E. and Wolf, W.J.  
Scale-up from analytical to preparative reversed-phase HPLC of soybean glycinin 444(1988)263
- Peterson, R.G., see Lalande, M. 430(1988)187
- Peterson, T., see Howard, II, R.R. 414(1987)434
- Petit, J., see Bougerolle, A.M. 434(1988)232
- Petitjean, O., see Ennachachibi, A. 427(1988)307
- Petkova, R., see Lekova, K. 446(1988)31
- Petrarulo, M., Pellegrino, S., Bianco, O., Marangella, M., Linari, F. and Mentasti, E.  
HPLC determination of glyoxylic acid in urine 432(1988)37
- Petrarulo, M., see Mentasti, E. 417(1987)253
- Petri, G., see Lemberkovic, É. 446(1988)267
- Petric, G., Schulz, D.E. and Duinker, J.C.  
Clean-up of environmental samples by HPLC for analysis of organochlorine compounds by GC with electron-capture detection 435(1988)241
- Petrucci, R.F., see Alton, K.B. 425(1988)363
- Petrvalský, E., see Dobránsky, T. 411(1987)486
- Petsev, N., see Engewald, W. 446(1988)71
- Petsev, N., see Gavrilova, T.B. 402(1987)87
- Pettersen, G., see Christophersen, A.S. 432(1988)290
- Petterson, C., Karlson, A. and Gioeli, C.  
Influence of enantiomeric purity of a chiral selector on stereoselectivity 407(1987)217
- Petterson, C. and Gioeli, C.  
Improved resolution of enantiomers of naproxen by the simultaneous use of a chiral stationary phase and a chiral additive in the mobile phase 435(1988)225
- Petz, M.  
Fluorimetric determination of sulphanilamide and sulphguanidine residues in animal tissues and products by ion-exchange HPLC (in German) 423(1987)217
- Pfadenhauer, E.H.  
HPLC assay for urinary metabolites of a drug after extraction by antiserum to the drug: an example using *erythro*-9-(2-hydroxy-3-nonyl)hypoxanthine 425(1988)407
- Pfleger, K., see Maurer, H. 422(1987)85
- Pfleger, K., see Maurer, H. 428(1988)43
- Pfleger, K., see Maurer, H. 430(1988)31
- Pflugmann, G., see Spahn, H. 433(1988)331
- Pflugmann, G., Spahn, H. and Mutschler, E.  
Rapid determination of the enantiomers of metoprolol, oxprenolol and propranolol in urine 416(1987)331
- Pflugmann, G., Spahn, H. and Mutschler, E.  
Determination of metoprolol enantiomers in plasma and urine using (S)-(-)-phenylethyl isocyanate as a chiral reagent 421(1987)161
- Philbert, M., see Bailey, E. 425(1988)187
- Philip, T., Chen, T.-S. and Nelson, D.B.  
LC profiles of major carotenoid esters in commercially processed California navel and Valencia orange juice concentrates 442(1988)249
- Philip, T. and Chen, T.-S.  
Separation and quantitative analysis of some carotenoid fatty acid esters of fruits by LC 435(1988)113
- Philipp, E.  
Assay for tyrosine hydroxylase in hypothalamic homogenates using HPLC with electrochemical detection 419(1987)27
- Phillips, M. and Greenberg, J.  
Detection of endogenous acetone in normal human breath 422(1987)235
- Phillips, M.W., see Subramanian, G. 439(1988)341
- Phillips, T.M., see Babashak, J.V. 444(1988)21
- Phillips, T.M., see Frantz, S.C. 444(1988)41
- Phillips, T.M. and Frantz, S.C.  
Isolation of specific lymphocyte receptors by high-performance immunoaffinity chromatography 444(1988)13
- Philp, R.P., Oung, J. and Lewis, C.A.  
Biomarker determinations in crude oils using a triple-stage quadrupole mass spectrometer 446(1988)3
- Piccardo, M.T., see Candiano, G. 423(1987)319
- Pichler, P., see Schilling, B. 435(1988)204
- Pichorner, H., see Ebermann, R. 410(1987)466
- Picot, V.S., see Doyle, E. 411(1987)325
- Piepmeyer, E.H., see Van Engelen, D.L. 405(1987)191
- Pierre, M.V. St. and Pang, K.S.

- Determination of diazepam and its metabolites by HPLC and TLC 421(1987)291
- Pieters, F.A.J.M., Vincken, B.J. and Zuidema, J.  
Dapsone and monoacetyldapsone determined in serum and saliva by a sensitive HPLC method with a single extraction step 422(1987)322
- Pietrogrande, M.C., Borea, P.A. and Biagi, G.L.  
Lipophilicity measurement of benzodiazepine-receptor ligands by reversed-phase LC. Comparison between HPLC and TLC 447(1988)404
- Pietta, P., Manera, E. and Ceva, P.  
Simultaneous isocratic HPLC determination of flavones and coumarins in *Matricaria chamomilla* extracts 404(1987)279
- Pietta, P., Mauri, P. and Pace, M.  
HPLC assay of chymotrypsin and chymotrypsin-like enzyme activity 411(1987)498
- Pietta, P., Mauri, P. and Pace, M.  
HPLC method for the determination of the esterase activity of subtilisin and kallikrein 441(1988)431
- Pietta, P., Mauri, P. and Rava, A.  
Reversed-phase HPLC method for the analysis of biflavones in *Ginkgo biloba* L. extracts 437(1988)453
- Pifferi, G., see Visconti, M. 416(1987)433
- Pilipenko, A.T., see Galushko, S.V. 445(1988)59
- Pimenov, A.M., see Toguzov, R.T. 434(1988)447
- Pimenov, A.M., see Werner, A. 421(1987)257
- Pinchbeck, F.M., see Osman, M.A. 419(1987)452
- Pinedo, H.M., see Maessen, P.A. 417(1987)339
- Pinedo, H.M., see Maessen, P.A. 424(1988)103
- Pini, D., see Salvadori, P. 450(1988)163
- Pinnen, F., see Pecci, L. 426(1988)183
- Pintamo-Kenttälä, K., see Mäkelä, M. 448(1988)391
- Pintér-Szakács, M., see Molnár-Perl, I. 446(1988)237
- Pinza, M., see Visconti, M. 416(1987)433
- Piotrowski, E.G., see Sonnet, P.E. 436(1988)205
- Piovan, D., see Padriani, R. 415(1987)183
- Piquion, J., see Sébille, B. 409(1987)61
- Piriou, A., see Mura, P. 416(1987)303
- Pirkle, W.H., Däppen, R. and Reno, D.S.  
Applications for racemic versions of chiral stationary phases 407(1987)211
- Pirkle, W.H. and Däppen, R.  
Reciprocity in chiral recognition. Comparison of several chiral stationary phases 404(1987)107
- Pirkle, W.H. and Hamper, B.C.  
Chromatographic separation of the enantiomers of 1,3-dithiolane-1-oxides 450(1988)199
- Pirkle, W.H. and McCune, J.E.  
An improved chiral stationary phase for the facile separation of enantiomers 441(1988)311
- Pirola, R., see Zecca, L. 420(1987)417
- Pittman, K.A., see Forgue, S.T. 414(1987)343
- Pittman, K.A., see Sciacca, M.A. 428(1988)265
- Pivonka, J., Segelman, F.H., Hartman, C.A., Segl, W.E., Kucharczyk, N. and Sofia, R.D.  
Determination of azelastine and desmethylazelastine in human plasma by HPLC 420(1987)89
- Pizzigallo, E., see Flamini, G. 426(1988)55
- Platt, D., see Rieck, W. 421(1987)206
- Platt, D., see Rieck, W. 425(1988)121
- Plicka, J., Kleinmann, I. and Svoboda, V.  
Preparative isolation of <sup>14</sup>C-labelled saccharose from a partly purified fraction of *Chlorella vulgaris* 442(1988)237
- Plöger, A., see Larsen, L.M. 450(1988)121
- Plumb, G.W., see Lambert, N. 402(1987)159
- Plummer, G.F.  
Improved method for the determination of propofol in blood by HPLC with fluorescence detection 421(1987)171
- Plunkett, W., see Gandhi, V. 413(1987)293
- Pochard, M.-F., Burger, G., Chevalier, M. and Gleizes, E.  
Determination of chloramphenicol residues by reversed-phase HPLC. Application to a pharmacokinetic study in rainbow trout with confirmation by MS (in French) 409(1987)315
- Podlaha, O.  
Letter symbols in the chemistry of glyceridic fats and oils 410(1987)509
- Podlešny, J., see Filipek, M. 430(1988)406
- Poehland, B.L. and Chan, J.A.  
Direct broth assay for sparsomycin and related nucleoside antitumor antibiotics using reversed-phase HPLC 439(1988)459
- Pohl, C.A., see Rocklin, R.D. 411(1987)107
- Point, A.J.F., see De Jong, J. 414(1987)285
- Poirier, J.-M., Perez, M., Cheymol, G. and Jaillon, P.  
Rapid and sensitive HPLC determination of bisoprolol in plasma and urine 426(1988)431
- Poirier, J.-M., see Garnier-Moiroux, A. 416(1987)183
- Poirier, J.M., Perez, M. and Cheymol, G.  
HPLC determination of gliclazide in human plasma 421(1987)223
- Pol, H.V. De, see Veenstra, J. 431(1988)413
- Polacek, J., Gunnarsson, P.O. and Brandin, S.  
Determination of taumoustine, a

- nitrosourea-based antitumour agent, in plasma by HPLC 425(1988)424
- Poláčiková, O., see Svobodová, X. 425(1988)391
- Polas, P., see Mathes, L.E. 432(1988)346
- Polcaro, C., Nicoletti, I., Ossicini, L., Caponecchi, G., Cozzi, R., Fiore, M. and Palitti, F. Chromatographic and cytogenetic analysis of *in vivo* metabolites of fluoranthene 448(1988)127
- Polívka, Ľ., see Dobránsky, T. 411(1987)486
- Pollak, V.A. and Schulze-Clewing, J. Pros and cons of electronic scanning in densitometry and an experimental device for their verification 437(1988)97
- Polonsky, J., see Donnelly, D.M.X. 440(1988)65
- Polonský, J., see Matejovič, I. 438(1988)454
- Pomaville, R.M., Poole, S.K., Davis, L.J. and Poole, C.F. Solute-solvent interactions in tetra-*n*-butylphosphonium salts studied by GC 438(1988)1
- Pommier, F., see Sioufi, A. 413(1987)101
- Pommier, F., see Sioufi, A. 428(1988)71
- Pommier, F., see Sioufi, A. 434(1988)239
- Pompon, A., see Maggio, A.-F. 436(1988)23
- Poncet, P., see Lacroix, C. 422(1987)217
- Pond, T., see Smith, R.E. 439(1988)83
- Poole, C.F., see Kersten, B.R. 411(1987)43
- Poole, C.F., see Pomaville, R.M. 438(1988)1
- Poole, C.F., see Poole, S.K. 435(1988)17
- Poole, C.F., see Shetty, P.H. 411(1987)61
- Poole, S.K., see Pomaville, R.M. 438(1988)1
- Poole, S.K. and Poole, C.F. Chemical interactions as a possible limitation on the useful solvent properties of liquid alkylammonium salts 435(1988)17
- Pop, D., see Ciucanu, I. 436(1988)219
- Porath, J. High-performance immobilized-metal-ion affinity chromatography of peptides and proteins 443(1988)3
- Porath, J., see Andersson, L. 421(1987)141
- Porath, J., see Belew, M. 403(1987)197
- Porcelli, G., Di Iorio, M. and Volpe, A.R. Determination of kininase I and kininase II activities in human urine by HPLC 414(1987)423
- Porcelli, M., Cacciapuoti, G., Oliva, A. and Zappia, V. HPLC determination of a naturally occurring xylosyl thioether, 9-[5'-deoxy-5'-(methylthio)- $\beta$ -D-xylofuranosyl]adenine 440(1988)151
- Porte, C., Barceló, D. and Albaigés, J. Quantitation of total *versus* selected polychlorinated biphenyl congeners in marine biota samples 442(1988)386
- Porter, R.S., see Slegowski, M.B. 425(1988)227
- Porter, W.R., see Vanscoik, K.G. 417(1987)183
- Porziemsky, J.-P., Krstulović, A.M., Wick, A., Barton, D.H.R., Tachdjian, C., Gateau-Olesker, A. and Gero, S.D. Complementary use of reversed-phase, normal-phase and chiral HPLC for the study of the stereoisomers produced in decarboxylative alkylation of tartaric acid 440(1988)183
- Porziemsky, J.-P., see Krstulović, A.M. 411(1987)461
- Postaire, E., see Pradeau, D. 447(1988)234
- Postaire, M., see Pradeau, D. 447(1988)234
- Postle, A.D. Method for the sensitive analysis of individual molecular species of phosphatidylcholine by HPLC using post-column fluorescence detection 415(1987)241
- Potschka, M. SEC of polyelectrolytes. Experimental evidence for a general mechanism 441(1988)239
- Potter, B.C., see Underwood, R.H. 415(1987)118
- Potterat, O., see Marston, A. 450(1988)3
- Poupaert, J.H., see Goncalves, J.C.S. 411(1987)472
- Pourgholami, M.H., Brain, K.R. and Nicholls, P.J. HPLC determination of imide inhibitors of aromatase in biological samples 424(1988)163
- Powell, D.R., Lee, P.D.K., Shively, J.E., Eckenhausen, M. and Hintz, R.L. Method for purification of an insulin-like growth factor-binding protein produced by human HEP G2 hepatoma cells 420(1987)163
- Powell, L.W., see Adams, P.C. 427(1988)341
- Power, A., see Dadgar, D. 416(1987)99
- Power, A., see Dadgar, D. 421(1987)216
- Powers, A.G., Young, J.H. and Clayton, B.E. Estimation of tetrahydrobiopterin and other pterins in plasma by isocratic LC with electrochemical and fluorimetric detection 432(1988)321
- Powers, J.J., see Krajewska, A.M. 409(1987)223
- Powis, G., see Brodfuehrer, J.I. 427(1988)247
- Powley, C.R., see Ravichandran, K. 439(1988)213
- Pradeau, D., Postaire, M., Postaire, E., Prognon, P. and Hamon, M. GC of carbon monoxide and carbon dioxide. Choice of a calibration method (in French) 447(1988)234
- Pramanik, B., see Mierzwa, R. 436(1988)259

- Pranoto-Soetardhi, L.A., see De Kruijf, N. 410(1987)395
- Pras, M., see Kaplan, B. 423(1987)376
- Prasad, M.R., Jones, R.M., Young, H.S., Kaplinsky, L.B. and Das, D.K.  
Analysis of tissue free fatty acids isolated by aminopropyl bonded-phase columns 428(1988)221
- Prasad, V.K., see Carlin, A.S. 425(1988)162
- Pratt, K.C., see Harvey, T.G. 435(1988)193
- Predieri, G., see Aggazzotti, G. 416(1987)125
- Predieri, G., see Casoli, A. 447(1988)187
- Premel-Cabic, A. and Allain, P.  
Simultaneous LC determination of norepinephrine, 3,4-dihydroxyphenylethyleneglycol, 3,4-dihydroxyphenylalanine and 3,4-dihydroxyphenylacetic acid in human plasma 434(1988)187
- Premkumar, N., see Smith, R.E. 439(1988)83
- Prescott, M.C., see Evershed, R.P. 440(1988)23
- Preti, G., Labows, J.N., Kostelc, J.G., Aldinger, S. and Daniele, R.  
Analysis of lung air from patients with bronchogenic carcinoma and controls using GC-MS 432(1988)1
- Prewitt, M.L., see Wooten, A.L. 445(1988)371
- Přihoda, M., see Pacáková, V. 442(1988)147
- Principe, D. Del, Colistra, C., Fiorentino, A.M., Spagnolo, G., Mancuso, G. and Menichelli, A.  
Detection of fibrinogen heterogeneity by capillary isotachopheresis 419(1987)329
- Prins, J., see Hoving, E.B. 434(1988)411
- Prins, R. and Somorjai, G.A.  
Contributions of GC to the development of heterogeneous catalysis 406(1987)151
- Prisyazhnoy, V.S., Fusek, M. and Alakhov, Y.B.  
Synthesis of high-capacity immunoaffinity sorbents with oriented immobilized immunoglobulins or their  $F_{ab}$  fragments for isolation of proteins 424(1988)243
- Prognon, P., see Pradeau, D. 447(1988)234
- Proietti, G., see Mezzetti, T. 434(1988)308
- Prokudin, V.Y., see Toguzov, R.T. 434(1988)447
- Provost, L.R., see D'Agostino, P.A. 402(1987)221
- Provost, L.R., see D'Agostino, P.A. 436(1988)399
- Pruden, D.J., Connolly, K.M. and Stecher, V.J.  
Single-step purification of rat C-reactive protein and generation of monospecific C-reactive protein antibody 437(1988)399
- Przuntek, H., see Wallasch, T.M. 425(1988)175
- Pu, Q.-L., see Zhang, T.-Y. 442(1988)455
- Puchta, V. and Spittler, G.  
Identification of 3-hydroxydecanoic acid dimethylate in human urine 420(1987)450
- Pudil, F., see Hajšlová, J. 438(1988)55
- Pudil, F., see Hajšlová, J. 439(1988)307
- Puech-Costes, É., see De Miguel, I. 407(1987)109
- Puglisi, C.V., Chen, S., Stelling-Ferrara, M., Pao, J. and Bekersky, I.  
Determination of (*E*)-1,2,3,4-tetrahydro-1,1,4,4-tetramethyl-6-(1-methyl-2-phenylethenyl)naphthalene, an antiacne agent, and its phenolic metabolite in plasma by reversed-phase HPLC 419(1987)253
- Puglisi, C.V., Ferrara, F.J. and Bekersky, I.  
Determination of the agonist-antagonist anxiolytic agent 6-(2-chlorophenyl)-4-hydroxy-4*H*-imidazo[1,5-*a*]-[1,4]benzodiazepine-3-carboxamide and its benzophenone carboxylic acid metabolite in plasma by HPLC 417(1987)347
- Puisieux, F., see Laham, A. 440(1988)455
- Pukkila, J., Peltonen, K. and Savolainen, T.  
Determination of melamine in air by HPLC with UV detection 411(1987)409
- Pukkila, J., Peltonen, K. and Savolainen, T.  
Determination of 2-phenyl-2-imidazoline in air by HPLC with UV detection 438(1988)443
- Pukkila, J., see Peltonen, K. 439(1988)375
- Pullen, R.H., Kennedy, C.M. and Curtis, M.  
Direct plasma injection using internal surface reversed-phase HPLC: feasibility study using propofol as a model compound 434(1988)271
- Pullen, R.H., see Cox, J.W. 424(1988)285
- Pullen, R.H., see Cox, J.W. 424(1988)293
- Punčochářová, J., see Kříž, J. 437(1988)177
- Purchase, R., see Bailey, E. 421(1987)21
- Puschendorf, B., see Lindner, H. 450(1988)309
- Pussard, E., Clavier, F. and Verdier, F.  
LC determination of ampyroquine in rabbit plasma and red blood cells 421(1987)192
- Pussard, E., see Bauchet, J. 414(1987)472
- Putnam, F.W. and Takahashi, N.  
Structural characterization of glycoproteins - 443(1988)267
- Puzo, G., see Rivière, M. 445(1988)87
- Pytilo, R.J., see Mikati, M.A. 433(1988)305

## Q

- Quaglia, M.G., see Fanali, S. 408(1987)441
- Quandt, K., see Zaromb, S. 439(1988)283



- Quearry, B.J., see Heyes, M.P. 428(1988)340  
 Quicke, J., see Rop, P.P. 427(1988)172  
 Quilliam, M.A., see Sim, P.G. 445(1988)239  
 Qureshi, G.A. and Eriksson, A.  
 Determination of clenbuterol and mabuterol in equine plasma by ion-pair LC with electrochemical detection. Chromatographic and electrochemical characteristics 441(1988)19723  
 Qureshi, S.A. and Buttar, H.S.  
 HPLC determination of propranolol and its metabolites in rat serum 431(1988)465  
 Quyen, T.C., see Van, T.D. 440(1988)71

## R

- Rabilloud, T., Pernelle, J.-J., Wahrmann, J.P., Gelfi, C. and Righetti, P.G.  
 Removal of oligomers and *n*-mers from the Immobiline chemicals for isoelectric focusing 402(1987)105  
 Rachev, R.C., Bojkova, S.V., Pavlova-Rouseva, R., Gancheva, V.K. and Dimitrova, M.P.  
 Separation of gibberellins by TLC and GC, and possibilities for their quantitative analysis 437(1988)287  
 Radeck, D., see Zobel, H. 446(1988)305  
 Radeck, W., Beck, K. and Staib, W.  
 Simple method for rapid quantification of branched-chain 2-oxo acids in physiological fluids as quinoxalinol derivatives by HPLC 432(1988)297  
 Räder, K., Wildfeuer, A., Wintersberger, F., Bossinger, P. and Mücke, H.-W.  
 Characterization of bee venom and its main components by HPLC 408(1987)341  
 Radhakrishnamurthy, B., see Lee, Y.J. 419(1987)275  
 Radulova, M., see Lekova, K. 446(1988)23  
 Radwan, F., see Bouthillier, F. 403(1987)171  
 Raglione, T.V., Troskosky, J.A. and Hartwick, R.A.  
 On-line microbore HPLC capillary GC-MS. I. Development of isotachic eluent splitters 409(1987)205  
 Raglione, T.V., Troskosky, J.A. and Hartwick, R.A.  
 On-line microbore HPLC capillary GC-MS. II. Application to the analysis of solvent refined coal 409(1987)213  
 Raha, C., Williamson, D., Thakran, P., Cassada, D. and Iversen, P.  
 Chromatographic and spectral investigations on the *in vivo* metabolites of 6-nitrobenzo[*a*]pyrene 450(1988)325  
 Rahman, A., see Lauren, D.R. 439(1988)470  
 Rahman, A., see Rustum, A.M. 421(1987)418  
 Rahman, A. and Hoffman, N.E.  
 HPLC determination of famotidine in urine 428(1988)395  
 Rahman, A. and Hoffman, N.E.  
 HPLC determination of fusidic acid in plasma 433(1988)159  
 Rahman, Y.E., see Venkataram, S. 411(1987)494  
 Railton, I.D.  
 Resolution of the enantiomers of abscisic acid methyl ester by HPLC using a stationary phase of cellulose tris(3,5-dimethylphenylcarbamate)-coated silica gel 402(1987)371  
 Rajani, P., see Euerby, M.R. 447(1988)392  
 Ramachandran, P.K., Gandhe, B.R., Venkateswaran, K.S., Kaushik, M.P., Vijayaraghavan, R., Agarwal, G.S., Gopalan, N., Suryanarayana, M.V.S., Shinde, S.K. and Sriramachari, S.  
 GC studies of the carbamylation of haemoglobin by methyl isocyanate in rats and rabbits 426(1988)239  
 Ramsey, E.D., see Parker, C.E. 438(1988)359  
 Ramus, T.L., Hein, S.J. and Thomas, L.C.  
 Determinations of polychlorinated biphenyl isomers by response factor calibration 404(1987)155  
 Ramzan, I.M.  
 Determination of the neuromuscular blocking drug gallamine in rat serum using HPLC 417(1987)428  
 Ramzan, I.M.  
 HPLC determination of chemical convulsant pentylenetetrazol in rat serum, cerebrospinal fluid and brain 432(1988)370  
 Rang, S., Kuningas, K., Strenze, T., Orav, A. and Eisen, O.  
 Retention and thermodynamics of solution of *n*-alkenes in OV-101 406(1987)75  
 Ranga Rao, K.V., see Padmalatha Devi, K. 426(1988)229  
 Ranga Rao, K.V., see Padmalatha Devi, K. 434(1988)265  
 Rao, D. and Borjes, G.  
 Simple GC method for the determination of medicagenic acid in alfalfa (*Medicago sativa*) 410(1987)169  
 Rapley, C.H., see Ubbink, J.B. 425(1988)182

- Rapoport, I., see Werner, A. 421(1987)257
- Rasheva, E., see Lekova, K. 446(1988)23
- Rashid, Z.A., see Szymanowski, J. 402(1987)55
- Rasmussen, K., see Christensen, J.M. 442(1988)317
- Rathbone, E.B., see Cookson, D. 402(1987)265
- Rathmann, J., see Förster, H.-G. 424(1988)357
- Rathor, M.N., Buffham, B.A. and Mason, G. Detection of void zones in packed chromatographic columns 404(1987)33
- Rathor, M.N., Buffham, B.A. and Mason, G. Determination of the volumes of void zones in packed chromatographic columns 445(1988)37
- Ratz, R., see Stehle, P. 449(1988)299
- Rau, H., see Stehle, P. 449(1988)299
- Rauch, K., Ray, J. and Graham, G. Improved HPLC method for the determination of quinine in plasma 430(1988)170
- Rauch, P., Beránková, E., Valentová, O., Sajdok, J. and Káš, J. High-performance anion-exchange chromatography of rennet enzymes 438(1988)451
- Rava, A., see Pietta, P. 437(1988)453
- Raverdino, V., see Bicchi, C. 411(1987)237
- Ravichandran, K., Lewis, J.J., Yin, I.-H., Koenigbauer, M., Powley, C.R., Shah, P. and Rogers, L.B. Computer-controlled linear pH gradient for HPLC fractionations of aromatic carboxylic acids and of humic and fulvic acids 439(1988)213
- Ravichandran, K. and Rogers, L.B. Chiral separations of amino acids using di- and tri-peptides for zwitter-ion pair chromatography 402(1987)49
- Ray, J., see Rauch, K. 430(1988)170
- Ray, M., see Lacroix, C. 422(1987)217
- Ray, R.K. and Kauffman, G.B. Chromatographic studies of metal complexes. I. PC of cobalt(III) complexes 442(1988)381
- Raynor, M.W., Kithinji, J.P., Barker, I.K., Bartle, K.D. and Wilson, I.D. Supercritical fluid chromatography of oedysteroids 436(1988)497
- Re, M., see Vassal, G. 428(1988)357
- Read, R.W., see Black, R.M. 449(1988)261
- Reames, S.A., see Koyama, Y. 439(1988)417
- Reams, G.P., see Souther, M.E. 417(1987)27
- Rebbani, N., see Girard, J.E. 448(1988)355
- Rebec, M.V. and Skrinska, V.A. HPLC with fluorescence detection of dazmegrel, a specific thromboxane A<sub>2</sub> synthetase inhibitor 432(1988)375
- Recio, M.N., see García-Pérez, A.I. 403(1987)131
- Reed, A.N. Quantitation of triazole and pyrimidine plant growth retardants 438(1988)393
- Reed, D.J., see Pascoe, G.A. 414(1987)440
- Reeuwijk, H.J.E.M., Lingeman, H., Tjaden, U.R., De Bruijn, E.A., Keizer, H.J. and Van der Greef, J. Bioanalysis of (*E*)-5-(2-bromovinyl)-2'-deoxyuridine 428(1988)93
- Reeves, R. and Elton, T.S. Non-histone chromatin proteins that recognize specific sequences of DNA (Review) 418(1987)73
- Reglero, G., Herraiz, M., Herraiz, T. and Loyola, E. Comparison of the performances of hot and cold sample introduction with a programmed-temperature vaporizer in the split and splitless modes 438(1988)243
- Regnault, V., Rivat, C. and Stoltz, J.F. Affinity purification of human plasma fibronectin on immobilized gelatin 432(1988)93
- Regnier, F.E. Chromatography of complex protein mixtures (Review) 418(1987)115
- Regnier, F.E., see Chicz, R.M. 443(1988)193
- Regnier, F.E., see Drager, R.R. 406(1987)237
- Regnier, F.E., see Fausnaugh-Pollitt, J. 443(1988)221
- Regnier, F.E., see Heinitz, M.L. 443(1988)173
- Regnier, F.E., see Heinitz, M.L. 443(1988)229
- Regnier, F.E., see Janis, L.J. 444(1988)1
- Regnier, F.E., see Mazsaroff, I. 411(1987)452
- Regnier, F.E., see Mazsaroff, I. 437(1988)429
- Regnier, F.E., see Mazsaroff, I. 442(1988)15
- Regnier, F.E., see Mazsaroff, I. 443(1988)119
- Regnier, F.E., see Rounds, M.A. 443(1988)73
- Regnier, F.E., see Stringham, R.W. 409(1987)305
- Regnier, F.E., see Tice, P.A. 410(1987)43
- Reh, E. Determination of digoxin in serum by on-line immuno-adsorptive clean-up HPLC separation and fluorescence-reaction detection 433(1988)119
- Reidenberg, M.M., see Wu, D.-F. 433(1988)141
- Reiff, K. HPLC method for the determination of carvedilol and its desmethyl metabolite in body fluids 413(1987)355
- Reiff, K. HPLC method for the determination of

- adibendan in plasma and urine 422(1987)346  
 Reine, A.H. and Stegink, S.J.  
 Qualitative and quantitative TLC analysis of polar lipids from leaves of glanded and glandless cotton genotypes 437(1988)211
- Reiss, D., see Mura, P. 416(1987)303
- Reiter, C., Müller, S. and Müller, T.  
 Improved method for the determination of sulphate in human serum using ion chromatography 413(1987)251
- Reitsma, B.H. and Yeung, E.S.  
 Reversed-phase HPLC of soybean trypsin inhibitor with optical activity and UV absorbance detection 405(1987)295
- Rejaibi, R., see Schöneshöfer, M. 417(1987)434
- Rejholec, V., see Kuchař, M. 449(1988)391
- Rommel, R.P., Dombrovskis, D. and Canafax, D.M.  
 Assay of itraconazole in leukemic patient plasma by reversed-phase small-bore LC 432(1988)388
- Reneman, R.S., see Engels, W. 427(1988)209
- Reno, D.S., see Pirkle, W.H. 407(1987)211
- Repka, D., see Krupčík, J. 406(1987)117
- Repka, D., see Krupčík, J. 407(1987)65
- Repka, D., see Krupčík, J. 448(1988)203
- Reppucci, L.M., see Feldberg, R.S. 410(1987)226
- Ress, K., Liebich, H.M., Krämer, B., Ickrath, O., Risler, T. and Seipel, L.  
 Determination of amidodarone and its metabolite N-desethylamidodarone in serum by HPLC—comparison of different extraction procedures 417(1987)465
- Ress, K.M., see Krämer, B.K. 427(1988)351
- Ressner, J.M., see Duff, G.A. 416(1987)71
- Reusch, J.F., see Kaminski, M. 436(1988)367
- Reutter, D.J., see Bossle, P.C. 407(1987)399
- Reutter, W., see Josić, D. 444(1988)29
- Revankar, S.N., see Bhatia, S.C. 434(1988)288
- Revankar, S.N., see Bhatia, S.C. 434(1988)296
- Revankar, S.N., see Bhatia, S.C. 434(1988)303
- Rexová-Benková, L., see Mikeš, O. 440(1988)287
- Reynolds, D.J., see Good, S.S. 431(1988)123
- Reynolds, Jr., N.C., see Fleming, L.H. 431(1988)65
- Řezanka, R., see Viden, I. 408(1987)145
- Řezanka, T., see Tvřická, E. 431(1988)231
- Řezanka, T. and Mareš, P.  
 Unusual and very long-chain fatty acids produced by Basidiomycetes 409(1987)390
- Rhee, J.S., see Nam, H.S. 448(1988)445
- Rhodes-Feuillette, A., see Jaureguiberry, G. 440(1988)385
- Rhodes-Feuillette, A., see Ogunkolade, B.W. 440(1988)459
- Rhodes, G.R. and Boppana, V.K.  
 HPLC analysis of arginine-containing peptides in biological fluids by means of a selective post-column reaction with fluorescence detection 444(1988)123
- Rhodes, S.H., Netting, A.G. and Milborrow, B.V.  
 Normal-phase HPLC of carotenes 442(1988)412
- Rhodes, S.H. and Netting, A.G.  
 Normal-phase HPLC of triacylglycerols 448(1988)135
- Rholam, M., see Créminon, C. 440(1988)439
- Ria, F., see Flamini, G. 421(1987)434
- Ria, F., see Flamini, G. 426(1988)55
- Rialdi, G., see Candiano, G. 423(1987)319
- Riazi, A., see Alessandri, J.-M. 432(1988)75
- Riba, M.-L., Tsiropoulos, N. and Torres, L.  
 Efficiency of adsorption-desorption of volatile compounds. Application to isoprene on Tenax GC 437(1988)139
- Ribick, M., see Cohen, A.I. 416(1987)445
- Ribon, B., see Benckekroun, Y. 420(1987)287
- Ricard, B., Bernard, M., Foglietti, M.J. and Percheron, F.  
 Human urinary  $\beta$ -mannosidase: behaviour towards two chromatographic supports 428(1988)203
- Ricard-Blum, S., Ville, G. and Hartmann, D.J.  
 Use of the Pharmacia Phast System for sodium dodecyl sulphate polyacrylamide gel electrophoresis of non-globular proteins: application to collagens 431(1988)474
- Ricci, P., see Billard, V. 440(1988)87
- Rice, L.M., see Zaromb, S. 439(1988)283
- Rice, P.D. and Bobbitt, D.R.  
 Linear model for the determination of solvent strength parameters in LSC 437(1988)3
- Richard, G., see Billard, V. 440(1988)87
- Richards, G., see Blackett, A. 433(1988)326
- Richards, M.P. and Steele, N.C.  
 Isolation and quantitation of metallothionein isoforms using reversed-phase HPLC 402(1987)243
- Richfield-Fratz, N. and Bailey, Jr., J.E.  
 Determination of *p*-cresidine in FD&C Red No. 40 by the diazotization and coupling procedure followed by reversed-phase HPLC 405(1987)283
- Richmond, R., Turner, N.C., Maltby, N., Heavey, D., Vial, J., Dollery, C.T. and Taylor, G.W.  
 Single-step procedure for the extraction and purification of leukotrienes B<sub>4</sub>, C<sub>4</sub> and D<sub>4</sub> 417(1987)241
- Richter, K.  
 Determination of clozapine in human serum by capillary GC 434(1988)465

- Rickard, E.C., see Inman, E.L. 447(1988)1
- Rideau, M., see Montagu, M. 409(1987)426
- Rieck, M., Hagen, M., Lutz, S. and König, W.A.  
Enantiomer separation of acyloins and cyanohydrins by enantioselective capillary GC 439(1988)301
- Rieck, W. and Platt, D.  
Determination of 2-(5-benzoyl-2-thienyl)propionic acid (tiaprofenic acid) in the plasma of elderly patients with multiple diseases using HPLC 421(1987)206
- Rieck, W. and Platt, D.  
HPLC method for the determination of  $\alpha$ -amanitin and phalloidin in human plasma using the column-switching technique and its application in suspected cases of poisoning by the green species of amanita mushroom (*Amanita phalloides*) 425(1988)121
- Riegel, M.A. and Ellis, P.P.  
HPLC assay for piperacillin in aqueous humor of the eye 424(1988)177
- Riemens, L., see Vree, T.B. 428(1988)311
- Riesner, D., see Hecker, R. 418(1987)97
- Riggin, R.M., Shaar, C.J., Dorulla, G.K., Lefeber, D.S. and Miner, D.J.  
High-performance SEC determination of the potency of biosynthetic human growth hormone products 435(1988)307
- Righetti, P.G., Fazio, M. and Tonani, C.  
Buffer isoelectric focusing revisited 440(1988)367
- Righetti, P.G., see Rabilloud, T. 402(1987)105
- Righezza, M., Murello, M.H. and Siouffi, A.M.  
Determination of nitrosamines by LC with post-column photolysis and electrochemical detection 410(1987)145
- Rigod, J., see Lee, F.W. 426(1988)259
- Rihn, B., Bisseret, F., Girardot, R., Scheffel, J.M., Nguyen, V.K. and Monteil, H.  
Fast protein purification of *Clostridium difficile* cytotoxin 428(1988)408
- Riis, V. and Mai, W.  
GC determination of poly- $\beta$ -hydroxybutyric acid in microbial biomass after hydrochloric acid propanolysis 445(1988)285
- Rijk, M.A.H., see De Kruijf, N. 410(1987)395
- Riley, C.M., see Crouch, Jr., F.W. 448(1988)333
- Rillaerts, E., see Engelmann, G.J. 432(1988)29
- Rimmer, S.R., see Dahiya, J.S. 448(1988)448
- Rindone, B., see Bassoli, A. 444(1988)209
- Rindone, B., see Bassoli, A. 446(1988)179
- Ringshandl, R., see Birnböck, H. 414(1987)235
- Rink, H., see Bours, J. 403(1987)336
- Risch, B., Galensa, R. and Herrmann, K.  
Analysis of catechin and epicatechin by HPLC after benzylation 448(1988)291
- Risk, M., see Holland, O.B. 426(1988)148
- Risler, T., see Krämer, B.K. 427(1988)351
- Risler, T., see Ress, K. 417(1987)465
- Ritland, T.G., see Yamasaki, R.B. 447(1988)277
- Rittmann, N., see Mennicke, W.H. 414(1987)19
- Riva, E., Ferry, N., Cometti, A., Cuisinaud, G., Gallo, G.G. and Sassard, J.  
Determination of teicoplanin in human plasma and urine by affinity and reversed-phase HPLC 421(1987)99
- Riva, G., see Bassoli, A. 444(1988)209
- Riva, G., see Bassoli, A. 446(1988)179
- Rivat, C., see Regnault, V. 432(1988)93
- Rivel, E.R., see Miksic, J.R. 428(1988)113
- Rivière, M., Fournié, J.-J., Monsarrat, B. and Puzo, G.  
Identification of partially methylated methyl glycosides by GC-MS of trimethylsilyl derivatives. Application to mycobacterial glycolipid antigen analysis 445(1988)87
- Rizzi, A.M., see Daučík, P. 442(1988)53
- Ro, R.T., see Nikitin, Y.S. 446(1988)55
- Roach, M.C., Gozel, P. and Zare, R.N.  
Determination of methotrexate and its major metabolite, 7-hydroxymethotrexate, using capillary zone electrophoresis and laser-induced fluorescence detection 426(1988)129
- Robards, K., Patsalides, E. and Dilli, S.  
GC of metal  $\beta$ -diketonates and their analogues (Review) 411(1987)1
- Robbins, A.D., see Brodie, R.R. 415(1987)423
- Robbins, D.K., Chang, S.-L., Baumann, R.J. and Wedlund, P.J.  
Quantitation of *trans*-10,11-dihydroxy-10,11-dihydrocarbamazepine in human urine by HPLC 415(1987)208
- Robertis, E. De, see Peña, C. 431(1988)192
- Roberts, C.J.C., see Forsyth, D.R. 433(1988)352
- Roberts, F.D., see Adams, P.C. 427(1988)341
- Roberts, M.S., see Owen, S.G. 416(1987)293
- Roberts, M.S., see Rumble, R.H. 419(1987)408
- Roberts, N.B., see Newton, C.J. 417(1987)391
- Robinson, J.L., John, J., Safa, A.I., Kirkes, K.A. and Griffith, P.E.  
Evaluation of disposable cartridges for trace enrichment from aqueous solutions 402(1987)201
- Robinson, P.R., Jones, M.D. and Maddock, J.  
Determination of clebopride in plasma by capillary GC-negative-ion chemical ionization MS 432(1988)153
- Robinson, R.S., see Tavss, E.A. 438(1988)273
- Robinson, R.S., see Tavss, E.A. 438(1988)281
- Rocca, J.L., see Chartier, A. 438(1988)263
- Rocci, Jr., M.L., see Grasela, D.M. 419(1987)368

- Rochester, N.E., see Haddad, P.R. 439(1988)23
- Rocklin, R.D., Pohl, C.A. and Schibler, J.A.  
Gradient elution in ion chromatography  
411(1987)107
- Rockstroh, H., Wangler, W. and Ackermann, T.  
Preparative fractionation of  
oligoriboguanylates by anion-exchange  
chromatography on TSK DEAE-5PW  
442(1988)401
- Rodchenkov, G.M., Uralets, V.P., Semenov, V.A.  
and Gurevich, V.A.  
GC and MS study of betamethasone synthetic  
corticosteroid metabolism 432(1988)283
- Rodchenkov, G.M., Uralets, V.P. and Semenov,  
V.A.  
Determination of methylprednisolone  
metabolites in human urine by GC-MS  
423(1987)15
- Rodchenkov, G.M., Uralets, V.P. and Semenov,  
V.A.  
GC and mass spectral study of synthetic  
corticosteroid metabolism: fluorometholone  
426(1988)399
- Rodkey, L.S.  
Ionic binding properties of carrier ampholytes  
437(1988)147
- Rodriguez, E., see Cavin, J.C. 447(1988)432
- Rodriguez, H., see Henzel, W.J. 404(1987)41
- Roelofs, H.M.J., see Evelo, C.T.A. 420(1987)35
- Roeraade, J.  
Factors affecting sample transfer from  
microlitre syringes 441(1988)367
- Roethlisberger, T., see El Tayar, N. 439(1988)237
- Roets, E., see Cachet, T. 403(1987)343
- Roets, E., see Cachet, T. 409(1987)91
- Roets, E., see Khan, N.H. 405(1987)229
- Roets, E., see Kibwage, I.O. 434(1988)177
- Rogers, L.B., see Aubel, M.T. 408(1987)99
- Rogers, L.B., see Ravichandran, K. 402(1987)49
- Rogers, L.B., see Ravichandran, K. 439(1988)213
- Rogoskin, V.A., Krylov, A.I. and Khlebnikova,  
N.S.  
GC determination of 1- and 3-methylhistidine  
in biological fluids 423(1987)33
- Rohrbaugh, D.K., Yang, Y.-C. and Ward, J.R.  
Identification of degradation products of 2-  
chloroethyl ethyl sulfide by GC-MS  
447(1988)165
- Rokushika, S., see Kihara, K. 410(1987)103
- Rokushika, S., see Murayama, T. 435(1988)417
- Rokushika, S., see Yamamoto, F.M. 408(1987)21
- Romeo, J.T., see Cole, M.D. 445(1988)295
- Römer, A., see Terlinden, R. 430(1988)438
- Romijn, J.C., see Verkoelen, C.F. 426(1988)41
- Rompaey, F. Van, see Woestenborghs, R.  
424(1988)195
- Róna, K. and Gachályi, B.  
Simple LC method for the determination of  
ornidazole and metronidazole in human  
serum 420(1987)228
- Rondot-Dudragne, M.L., see Blanchin, M.D.  
432(1988)407
- Ronfeld, R.A., see Fouda, H.G. 417(1987)197
- Rood, Jr., H.D., see Swanson, S.P. 414(1987)335
- Roorda, W.E., Verbeek, J. and Junginger, H.E.  
Simple GC determination of the purity of  
hydroxyethyl methacrylate 403(1987)355
- Roos, D., see De Korte, D. 415(1987)383
- Rop, P.P., Spinazzola, J., Zahra, A., Bresson, M.,  
Quicke, J. and Viala, A.  
Column LC analysis of barbiturates in  
biological fluids 427(1988)172
- Rose, Jr., D.J. and Jorgenson, J.W.  
Post-capillary fluorescence detection in  
capillary zone electrophoresis using *o*-  
phthalaldehyde 447(1988)117
- Rose, D.J. and Jorgenson, J.W.  
Fraction collector for capillary zone  
electrophoresis 438(1988)23
- Rose, L.M., Tillery, K.F., El Dareer, S.M. and  
Hill, D.L.  
HPLC determination of doxorubicin and its  
metabolites in plasma and tissue 425(1988)419
- Roselli, R.J., see Herrero, P. 416(1987)265
- Roselló, J., see Cárdenas, A. 426(1988)83
- Rosén, S., see Bergenhem, N. 403(1987)383
- Rosenberg, L.S., see Herbranson, D.E.  
410(1987)222
- Rosenbloom, J., see Folena-Wasserman, G.  
411(1987)345
- Rosenfeld, J.B., see Garty, M. 430(1988)123
- Rosenthal, I., see Zuk, M.M. 433(1988)367
- Rosenthal, P., see Vreman, H.J. 427(1988)315
- Rosenthal, W., see Kleuss, C. 407(1987)281
- Roshchina, T.M., see Gavrilova, T.B. 446(1988)79
- Rosier, J. and Van Peteghem, C.  
Detection of oxidatively modified 2'-  
deoxyguanosine-3'-monophosphate, using  
<sup>32</sup>P-postlabelling and anion-exchange TLC  
434(1988)222
- Rosini, C., see Salvadori, P. 450(1988)163
- Ross, L. Fischer  
GC technique to simultaneously quantitate  
the gases produced by intestinal  
microorganisms from fermentation mixtures  
414(1987)405
- Rosset, R., see Gareil, P. 450(1988)13
- Rosset, R., see Jadaud, P. 439(1988)195
- Rosset, R., see Janicot, J.L. 437(1988)351

- Rosset, R., see Jardy, A. 439(1988)137
- Rosset, R., see Lienne, M. 448(1988)55
- Rosset, R., see Macaudiere, P. 405(1987)135
- Rosset, R., see Macaudière, P. 450(1988)255
- Rosset, R., see Morin, P. 407(1987)87
- Rosset, R., see Ni, P. 424(1988)255
- Rossetti, D.V., see Castagnola, M. 440(1988)231
- Rossey, G., see Krstulović, A.M. 411(1987)461
- Rossignol, P., see Laham, A. 440(1988)455
- Rossomando, E.F., see Mina, M. 433(1988)63
- Rotatori, M., see Laganá, A. 417(1987)135
- Roth, H.J., see Herrmann, T. 416(1987)170
- Roth, H.J., see Herrmann, T. 428(1988)237
- Roth, H.J., see Knospe, J. 442(1988)444
- Roth, M.  
Effect of column inlet concentration profile on the sensitivity of analysis. A quantitative assessment 409(1987)360
- Roth, M., see Padmalatha Devi, K. 426(1988)229
- Rothuizen, J., Heirwegh, K.P.M. and Van Kouwen, A.M.  
Novel method for HPLC of azo derivatives of conjugated and unconjugated bilirubin 427(1988)19
- Rothwell, C.M., Hsieh, J.Y.-K. and Bayne, W.F.  
HPLC determination of 4-[(3-(4-acetyl-3-hydroxy-2-propylphenoxy)propyl)sulfonyl]- $\gamma$ -oxobenzenebutanoic acid and metabolites in human plasma 417(1987)447
- Rotocki, P. and Drozdowicz, B.  
Influence of the carrier gas flow-rate on the pulsed electron-capture detector 446(1988)329
- Rouan, M.C.  
Microbore LC determination of cadralazine and cephalixin in plasma with large-volume injection 426(1988)335
- Rounds, M.A., see Mazsaroff, I. 411(1987)452
- Rounds, M.A. and Regnier, F.E.  
Synthesis of a non-porous, polystyrene-based strong anion-exchange packing material and its application to fast HPLC of proteins 443(1988)73
- Rouse, C.A., see Bradshaw, J.S. 405(1987)169
- Roussel, A., see Berthod, A. 449(1988)349
- Row, K.H., Griest, W.H. and Mäskarinec, M.P.  
Separation of modified nucleic acid constituents by micellar electrokinetic capillary chromatography 409(1987)193
- Roy, J., see Guha, O.K. 437(1988)244
- Roy, S.D. and Lim, H.K.  
Separation of optical isomers of methoxyphenamine and its metabolites as N-heptafluorobutyl-L-prolyl derivatives by fused-silica capillary GC 431(1988)210
- Roybal, J.E., see Munns, R.K. 442(1988)209
- Ruane, R.J. and Wilson, I.D.  
Ion-pair reversed-phase TLC of basic drugs using sulphonic acids 441(1988)355
- Rubino, F.M., see Fermo, I. 433(1988)53
- Rubio, F., Chen, S., Crews, T., De Grazia, F., Garland, W.A. and Barbalas, M.  
Determination of 6-(2'-chlorophenyl)-4-hydroxy-4*H*-imidazo[1,5-*a*][1,4]benzodiazepine-3-carboxamide, a mixed agonist-antagonist anxiolytic agent, in human plasma and urine by GC-negative-ion chemical-ionization MS 421(1987)281
- Rubio, N., Brieva, A. and Alcazar, B.  
Purification of allergens by HPLC. IV. Purification of the allergen of olive pollen (*Olea europea*) 403(1987)312
- Rubio, N. and Brieva, A.  
Purification of allergens by HPLC. V. Purification of cat and dog epithelial allergens 404(1987)378
- Rubio, N. and Brieva, A.  
Purification of allergens by HPLC. VI. Purification of the major allergen from *Parietaria judaica* pollen 407(1987)408
- Rubio, N. and Brieva, A.  
Purification of allergens by HPLC. VII. Purification of the major allergen of birch pollen (*Betula verrucosa*) 408(1987)420
- Rudmann, A.A. and Aldrich, J.R.  
Chirality determinations for a tertiary alcohol. Ratios of linalool enantiomers in insects and plants 407(1987)324
- Rudziński, W., Michalek, J., Brun, B. and Partyka, S.  
Further comments on the preparation of mixed organic layers chemically bonded on silicon dioxide. Adsorption of modifiers from strongly adsorbed solvents 406(1987)295
- Rüfer, R., see Kraus, R. 413(1987)257
- Ruiter, C. De, Otten, R.R., Brinkman, U.A.T. and Frei, R.W.  
Rapid and simple dansylation of phenolic steroids using a two-phase system and phase transfer catalysis 436(1988)429
- Ruiter, C. De, see Broersen, J.J. 436(1988)39
- Ruiter, R. De, see Stolk, L.M.L. 423(1987)383
- Rumble, R.H. and Roberts, M.S.  
HPLC assay of the major components of gentamicin in serum 419(1987)408
- Rush, K., see Miksic, J.R. 428(1988)113
- Rushing, L.G., see Holder, C.L. 419(1987)113
- Russell, J., see Catterick, T. 407(1987)310
- Russell, J.W., see Kerns, E.H. 416(1987)357
- Rüstig, H., see Voelkel, A. 448(1988)219
- Rustum, A.M., Rahman, A. and Hoffman, N.E.

- HPLC determination of ranitidine in whole blood and plasma by using a short polymeric column 421(1987)418
- Rustum, A.M. and Hoffman, N.E.  
HPLC determination of 5-azacytidine in plasma 421(1987)387
- Rustum, A.M. and Hoffman, N.E.  
Determination of cyclophosphamide in whole blood and plasma by reversed-phase HPLC 422(1987)125
- Rustum, A.M. and Hoffman, N.E.  
Determination of 5-fluorouracil in plasma and whole blood by ion-pair HPLC 426(1988)121
- Rutgers, H.M., see Hoving, E.B. 434(1988)411
- Rutledge, D.R.  
Rapid HPLC method for the measurement of gallopamil in plasma 421(1987)165
- Rzeszotarska, B., see Sikora, E. 442(1988)420
- S**
- Saadawi, A., see Stolk, L.M.L. 423(1987)383
- Saba, C.S., see Keller, M.A. 409(1987)325
- Sabater, F., see Barcelo, A.R. 411(1987)476
- Sablonniere, B., Lefebvre, P., Formstecher, P. and Dautrevaux, M.  
Improved Stokes radius measurement of the glucocorticoid receptor using TSK G4000SW and TSK G3000SW high-performance size-exclusion columns. Analytical and preparative applications 403(1987)183
- Saccar, C.L., see Kester, M.B. 416(1987)91
- Sacchetto, G.A., see Mingjia, W. 439(1988)363
- Sacchi, C., see Pedrazzini, S. 415(1987)214
- Sado, P., see Le Corre, P. 424(1988)424
- Sado, P., see Le Corre, P. 450(1988)211
- Sado, P.A., Gibassier, D. and Leverage, R.  
Simultaneous determination of vinburnine and 6-hydroxyvinburnine in human plasma by HPLC (in French) 434(1988)157
- Safa, A.I., see Robinson, J.L. 402(1987)201
- Sagara, K., Oshima, T., Yoshida, T., Tong, Y.-Y., Zhang, G. and Chen, Y.-H.  
Determination of Cinnamomi cortex by HPLC 409(1987)365
- Sagara, K., Oshima, T. and Yoshida, T.  
Rapid and simple determination of sennoside A and B in Rhei Rhizoma by ion-pair HPLC 403(1987)253
- Sagara, Y., Okatani, Y., Yamanaka, S. and Kiriya, T.  
Determination of plasma 5-hydroxytryptophan, 5-hydroxytryptamine, 5-hydroxyindoleacetic acid, tryptophan and melatonin by HPLC with electrochemical detection 431(1988)170
- Sager, A.O., see Carlin, A.S. 425(1988)162
- Sagliano, Jr., N., Floyd, T.R., Hartwick, R.A., Dibussolo, J.M. and Miller, N.T.  
Studies on the stabilization of reversed phases for LC 443(1988)155
- Saha, M., see Minnetian, O. 410(1987)453
- Sahajwalla, C.G., see Bhatia, S.C. 434(1988)288
- Sahajwalla, C.G., see Bhatia, S.C. 434(1988)296
- Sahajwalla, C.G., see Bhatia, S.C. 434(1988)303
- Sainio, E.-L. and Hynninen, P.H.  
High-voltage electrophoretic separation of tryptophan metabolites of the kynurenine pathway 420(1987)263
- Saint Blanquat, G.D., Lamboeuf, Y. and Fritsch, P.  
Determination of catecholamines in biological tissues by LC with coulometric detection 415(1987)388
- Saint Blanquat, G. De, see Leitao, J. 435(1988)229
- Saita, T., see Matsanaga, H. 422(1987)353
- Saito, A., see Chiu, W.-Y. 428(1988)25
- Saito, K., see Horie, M. 402(1987)301
- Saito, M., see Sugiyama, K. 442(1988)121
- Saitoh, K., see Noda, H. 435(1988)368
- Saitoh, K., see Suzuki, N. 408(1987)181
- Saitoh, K., Tsukahara, S. and Suzuki, N.  
LC retention of  $\beta$ -diketonato complexes of chromium(III) and palladium(II) in octadecyl-bonded silica/water-methanol systems. Comparison with liquid-liquid partition using dodecane as a model alkyl-bonded phase 439(1988)245
- Saitoh, Y., see Aoyama, T. 430(1988)351
- Saitoh, Y., see Yamamoto, K. 433(1988)167
- Sajdok, J., see Rauch, P. 438(1988)451
- Sakagishi, Y., see Koyama, I. 413(1987)65
- Sakagishi, Y., see Koyama, I. 420(1987)275
- Sakai, J., see Togari, A. 417(1987)41
- Sakai, T., Kojima, K. and Nagatsu, T.  
Rapid chromatographic purification of dipeptidyl-aminopeptidase II from human kidney 416(1987)131
- Sakai, T., Takeuchi, Y., Araki, T. and Ushio, K.  
Determination of erythrocyte porphyrins by reversed-phase HPLC using capsule-type silica gels coated with silicone polymer 433(1988)73

- Sakakibara, J., see Kidouchi, K. 423(1987)297
- Sakamoto, J., see Itami, T. 417(1987)168
- Sakamoto, M., Nishimoto, M., Kohara, N. and Masuko, F.  
GC and MS behaviour of O-alkylserines as their N-trifluoroacetyl butyl esters  
411(1987)259
- Sakamoto, T., see Kurumaya, K. 435(1988)235
- Sakamoto, Y., Kawakami, N. and Sasagawa, T.  
Prediction of peptide retention times  
442(1988)69
- Sakano, T., see Hirauchi, K. 430(1988)21
- Sakata, C., see Terada, T. 430(1988)388
- Sakiyama, N., see Kataoka, H. 436(1988)67
- Sakka, S., see Hasegawa, I. 410(1987)137
- Sako, Y., see Araki, A. 422(1987)43
- Sakoguchi, T., see Kobayashi, K. 417(1987)439
- Sakuragawa, N., see Matsukura, H. 414(1987)47
- Salama, A., see Hoffman, D.W. 414(1987)504
- Šalamoun, J., Smrž, M., Kiss, F. and Šalamounová, A.  
Column LC of methotrexate and its metabolites using a post-column photochemical reactor and fluorescence detection  
419(1987)213
- Šalamoun, J., Šprta, V., Sládek, T. and Smrž, M.  
Determination of acyclovir in plasma by column LC with fluorescence detection  
420(1987)197
- Šalamounová, A., see Šalamoun, J. 419(1987)213
- Salari, H.  
Comparative study of solid-phase and liquid-phase extraction techniques for isolation of phospholipids from plasma  
419(1987)103
- Salari, H., see Steffenrud, S. 416(1987)219
- Salari, H., see Steffenrud, S. 427(1988)1
- Sallustio, B.C., Meffin, P.J. and Thompson, M.  
HPLC quantitation of triacylglycerols containing fenoprofen from biological samples  
422(1987)33
- Salvadori, P., Pini, D., Rosini, C., Uccello-Barretta, G. and Bertucci, C.  
Chiral stationary phase derived from L-lactic acid for the optical resolution of N-(3,5-dinitrobenzoyl)amino acid methyl esters  
450(1988)163
- Samain, D., see De Miguel, I. 407(1987)109
- Samain, D., see Inchauspe, G. 404(1987)53
- Samara, E. and Bialer, M.  
Rapid HPLC assay with pharmacokinetic applications for monitoring cannabidiol in plasma  
416(1987)370
- Samolin, Y.A., see Manoilov, S.E. 440(1988)53
- Samoukina, G.V., see Beran, M. 442(1988)431
- Samperi, R., see Caccamo, F. 434(1988)61
- Samperi, R., see Di Corcia, A. 405(1987)357
- Sams, D.J., see Brimmell, P.A. 413(1987)320
- Sams, R., see Mathes, L.E. 432(1988)346
- Samu, J., Kováts, E., Nguyen, V., Keler, T., Nowotny, A. and Coughlin, R.T.  
TLC of endotoxins, their derivatives and contaminants  
435(1988)167
- Sanchez, A., see Hubbard, R.W. 431(1988)163
- Sanchez, D., see Larsson, K. 450(1988)71
- Sánchez-Rasero, F. and Dios, G.C.  
LC method for the determination of cyanazine in the presence of some normal soil constituents  
447(1988)426
- Sancho, P., see García-Pérez, A.I. 403(1987)131
- Sandman, H., see Odink, J. 431(1988)309
- Sandnes, K., see Waagboe, R. 427(1988)138
- Sandström, J.F., see Dalene, M. 435(1988)469
- Sanford, W., see Duffy, M. 441(1988)73
- Sangö, C., see Dalene, M. 435(1988)469
- Sanitrák, J., Krijt, J., Čoupek, J., Janoušek, V. and Magnus, I.A.  
Determination of porphyrins in tissue: pre-adsorption followed by HPLC  
415(1987)129
- Sano, A., Matsutani, S., Suzuki, M. and Takitani, S.  
HPLC method for determining trichothecene mycotoxins by post-column fluorescence derivatization  
410(1987)427
- Sano, Y., see Goto, J. 425(1988)59
- Sano, Y., see Yamamoto, S. 409(1987)101
- Sansoulet, J., see Delee, E. 450(1988)183
- Santa-Coloma, T.A., Garraffo, H.M., Gros, E.G. and Charreau, E.H.  
Cyclic biospecific affinity chromatographic method for the purification of the sex steroid binding protein (SBP): application to the purification of SBP from toad  
415(1987)297
- Santagostino, G., see Schinelli, S. 431(1988)150
- Santalucia, J., see Tavss, E.A. 438(1988)273
- Santalucia, J., see Tavss, E.A. 438(1988)281
- Santarelli, X., Muller, D. and Jozefonvicz, J.  
Dextran-coated silica packings for high-performance SEC of proteins  
443(1988)55
- Santiuste, J.M., see Fernández-Sánchez, E. 402(1987)318
- Santiuste, J.M., see Fernández-Sánchez, E. 410(1987)13
- Santos, L.M. and Baldwin, R.P.  
Profiling of oxalic acid and  $\alpha$ -keto acids in blood and urine by LC with electrochemical detection at a chemically modified electrode  
414(1987)161
- Santos, S.R.C.J., Kirch, W. and Ohnhaus, E.E.  
Simultaneous analysis of digitoxin and its



- clinically relevant metabolites using HPLC and radioimmunoassay 419(1987)155
- Sanz, J., see García-Raso, A. 402(1987)323
- Sappa, E., see Casoli, A. 447(1988)187
- Saria, A., see Haring, C. 428(1988)160
- Sarkar, M.A. and Karnes, H.T.  
Comparison of solid-phase extraction and liquid-liquid extraction for HPLC analysis of antipyrine in plasma 434(1988)324
- Sarver, E.W., see Bossle, P.C. 407(1987)399
- Sasagawa, T., see Sakamoto, Y. 442(1988)69
- Sasaki, N., see Endoh, Y.S. 426(1988)202
- Sasaki, Y. and Baba, S.  
Simultaneous determination of imipramine, desipramine and their deuterium-labelled analogues in biological fluids by capillary GC-MS 426(1988)93
- Sasaoka, T., Kaneda, N., Niwa, T., Hashizume, Y. and Nagatsu, T.  
Analysis of salsolinol in human brain using HPLC with electrochemical detection 428(1988)152
- Sasaoka, T., Kaneda, N. and Nagatsu, T.  
Highly sensitive assay for xanthine oxidase activity by HPLC with fluorescence detection 424(1988)392
- Sassard, J., see Riva, E. 421(1987)99
- Satama, L.T., see Nurmela, K.V.V. 435(1988)139
- Sato, H., Mitsutani, K., Shimizu, I. and Tanaka, Y.  
Determination of chemical composition distribution of styrene-methyl methacrylate copolymers by reversed-phase HPLC 447(1988)387
- Sato, H., see Tanaka, Y. 407(1987)197
- Sato, S., see Maeda, Y. 410(1987)413
- Sato, T., see Kadofuku, T. 419(1987)315
- Sato, T., see Yamada, K. 424(1988)367
- Satsmadjis, J., Georgakopoulos-Gregoriades, E. and Voutsinou-Taliadouri, F.  
Separation of organochlorines on alumina 437(1988)254
- Satterwhite, D.M. and Croteau, R.B.  
Resolution of monoterpene enantiomers by GC 407(1987)243
- Satterwhite, J.H. and Boudinot, F.D.  
HPLC determination of ketoprofen and naproxen in rat plasma 431(1988)444
- Saunders, G.A., see McKerrell, P.J. 408(1987)399
- Savigliano, M., see Mentasti, E. 417(1987)253
- Savolainen, T., see Pukkila, J. 411(1987)409
- Savolainen, T., see Pukkila, J. 438(1988)443
- Sawada, S., Kasai, M., Yamada, T. and Takahashi, T.  
Determination of photosynthetic intermediates in leaves by HPLC 402(1987)283
- Sawada, T., see Yoshikawa, S. 421(1987)111
- Sayo, H., Hatsumura, H., Hosokawa, M. and Michida, T.  
Electrochemical immunoassay combined with column LC: determination of phenytoin in human serum 417(1987)129
- Sayo, H., Hatsumura, H. and Hosokawa, M.  
Simultaneous determination of phenytoin and phenobarbital in human serum by electrochemical immunoassay combined with HPLC 426(1988)449
- Sbordoni, A., see Mezzetti, T. 434(1988)308
- Scalia, S.  
Simultaneous determination of free and conjugated bile acids in human gastric juice by HPLC 431(1988)259
- Scavone, J.M., Meneilly, G.P., Greenblatt, D.J. and Friedman, H.  
GC analysis of underivatized tocainide 419(1987)339
- Scavone, J.M., see Kennedy, J.S. 422(1987)274
- Scavone, J.M., see Kennedy, J.S. 423(1987)373
- Scaysbrook, T., see Greenaway, W. 448(1988)284
- Schaafsma, G., see Veenstra, J. 431(1988)413
- Schaake, S., see Bours, J. 403(1987)336
- Schacher, A., see Hochuli, E. 411(1987)177
- Schafer, M.P.  
Reversed-phase HPLC separation and quantitation of reticulocyte  $\alpha$ - and  $\beta$ -globin polypeptide chains from normal and  $\beta$ -thalassemic mice 431(1988)177
- Schafer, R., see Tedesco, J.L. 403(1987)299
- Schalekamp, M.A.D.H., see Boomsma, F. 427(1988)219
- Schalkwijk, W.P. Van, see Verkoelen, C.F. 426(1988)41
- Scharpf, F., Yeates, R.A., Laufen, H. and Eibel, G.  
GC assay of glycerol mononitrates in biological samples 413(1987)91
- Schatowitz, B. and Gercken, G.  
Benzyl esters of  $C_2$ - $C_{20}$  fatty acids and metabolically relevant carboxylic acids. Preparation and GC-MS 409(1987)43
- Schatowitz, B. and Gercken, G.  
Simultaneous determination of  $C_2$ - $C_{22}$  non-esterified fatty acids and other metabolically relevant carboxylic acids in biological material by GC of their benzyl esters 425(1988)257
- Schaubhut, R., see Minder, E.I. 419(1987)135
- Schaubhut, R., see Minder, E.I. 428(1988)369
- Schay, G., see Nagy, L.G. 406(1987)311

- Scheffer, J.J.C., see Van Gelder, W.M.J. 442(1988)133
- Scheftel, J.M., see Rihn, B. 428(1988)408
- Schellens, J.H.M., Der Wart, H.H.F.V., Hoevers, J.W. and Breimer, D.D.  
GC determination of sparteine and 2- and 5-dehydrosparteine in plasma and urine 431(1988)203
- Schenck, P.A., see Sinninghe Damsté, J.S. 435(1988)435
- Schibler, J.A., see Rocklin, R.D. 411(1987)107
- Schill, G., see Steuer, W. 447(1988)287
- Schilling, B., Grob, K., Pichler, P., Dubs, R. and Brechbühler, B.  
Fused-silica ends for glass capillaries: widening glass capillary butts by a heated tungsten point for preparing press-fit connections 435(1988)204
- Schindler, M., see Steuer, W. 447(1988)287
- Schinelli, S., Frattini, P., Cucchi, M.L., Peroni, A.E. and Santagostino, G.  
Determination of vanillylmandelic acid in plasma by HPLC with electrochemical detection 431(1988)150
- Schlabach, T.D. and Excoffier, J.L.  
Multi-variate ranking function for optimizing separations 439(1988)173
- Schmelzer, C.H., see Ebert, R.F. 443(1988)309
- Schmerr, M.J.F., Goodwin, K.R. and Van der Maaten, M.J.  
High-performance GPC of the surface glycoprotein from bovine leukemia virus 425(1988)379
- Schmid, B.J., Perry, H.E. and Idle, J.R.  
Determination of nifedipine and its three principal metabolites in plasma and urine by automated electron-capture capillary GC 425(1988)107
- Schmid, J.  
Assay of ambroxol in biological fluids by capillary GLC 414(1987)65
- Schmid, K., see Graf, P. 425(1988)353
- Schmidt, H., see Werner, A. 421(1987)257
- Schmidt, J. and McClain, C.J.  
Separation strategies for *o*-phthalaldehyde-mercaptoethanol derivatives of amino acids for reversed-phase HPLC 419(1987)1
- Schmidt, K., see Scholl, H. 416(1987)321
- Schmidt, N., Gercken, G. and König, W.A.  
Enantioselective capillary GC of 1,2-isopropylidene glycerol and O-alkylated glycerol derivatives 410(1987)458
- Schmitt, A., see Voegel, J.C. 428(1988)17
- Schmitter, J.-M., see Beauvallet, C. 438(1988)347
- Schmitz, F.P., see Klesper, E. 402(1987)1
- Schmitz, G., Böttcher, A., Kahl, H.-G. and Brüning, T.  
Purification of monoclonal antibodies against the low-density lipoprotein receptor by preparative isotachopheresis 431(1988)327
- Schnabelrauch, M., see Klemm, D. 438(1988)122
- Schneebaum, S., Bies, J.M. and Briele, H.A.  
Reversed-phase HPLC of dactinomycin 427(1988)166
- Schneider, K., see Dietel, P. 422(1987)1
- Schneiderman, J., see Kasziba, E. 432(1988)315
- Schneiderman, M.A., Sharma, A.K. and Locke, D.C.  
Determination of anthraquinone in paper and wood using supercritical fluid extraction and HPLC with electrochemical detection 409(1987)343
- Schnell, A.M., see Ubbink, J.B. 425(1988)182
- Schnell, A.M., see Ubbink, J.B. 431(1988)406
- Schoel, B. and Kaufmann, S.H.E.  
Isolation of recombinant mycobacterial antigens by an automatic and generally applicable purification method for  $\beta$ -galactosidase fusion proteins 448(1988)165
- Scholl, H., Schmidt, K. and Weber, B.  
Sensitive and selective determination of picogram amounts of ciprofloxacin and its metabolites in biological samples using HPLC and photothermal post-column derivatization 416(1987)321
- Schönberger, E., see MacCrehan, W.A. 417(1987)65
- Schöneshöfer, M., Heilmann, P. and Rejaibi, R.  
Automated column LC determination of polythiazide in human serum 417(1987)434
- Schönfeld, W., see Knöller, J. 427(1988)199
- Schönfeld, W., see Knöller, J. 427(1988)257
- Schoppek, B., see Kauert, G. 428(1988)291
- Schott, H.  
Preparative preparation of homologues of oligoribouridylic acid by selective partial hydrolysis of RNA (in German) 445(1988)155
- Schott, H., Leitner, B., Schwendener, R.A. and Hengartner, H.  
Chromatography of functionalized liposomes and their components 441(1988)115
- Schouten, A., see De Kruijff, N. 410(1987)395
- Schrader, P.A. and Sisco, W.R.  
HPLC analysis of fenretinide in rodent feed 408(1987)430
- Schreuder, R.H. and Martijn, A.  
Determination of dodecylbenzenesulphonates and ethoxylated alkylphenols in liquid pesticide formulations by HPLC 435(1988)73

- Schroeder, A.C., see Kapetanovic, I.M. 419(1987)458
- Schroeder, F.H., see Verkoelen, C.F. 426(1988)41
- Schuch, R. and Mukherjee, K.D.  
Rapid radio TLC for assay of lipase-catalyzed esterification and interesterification reactions 450(1988)448
- Schuddinck, G. and Verzele, M.  
Avoidance of trace metal activity with micro-LC in the analysis of hop and beer bitter acids 407(1987)159
- Schulte, B.A., see Leskawa, K.C. 411(1987)393
- Schultz, G., see Kleuss, C. 407(1987)281
- Schulz, D.E., see Petrick, G. 435(1988)241
- Schulz, H. and Albroseheit, G.  
HPLC characterization of some medical plant extracts used in cosmetic formulas 442(1988)353
- Schulze-Clewing, J., see Pollak, V.A. 437(1988)97
- Schulze, J., see Weller, O. 403(1987)263
- Schurig, V.  
Enantiomer analysis by complexation GC. Scope, merits and limitations 441(1988)135
- Schurig, V. and Nowotny, H.-P.  
Separation of enantiomers on diluted permethylated  $\beta$ -cyclodextrin by high-resolution GC 441(1988)155
- Schuster, D., Woodruff Modi, M., Lalka, D. and Gengo, F.M.  
Reversed-phase HPLC assay to quantitate diastereomeric derivatives of metoprolol enantiomers in plasma 433(1988)318
- Schuster, R.  
Determination of amino acids in biological, pharmaceutical, plant and food samples by automated precolumn derivatization and HPLC 431(1988)271
- Schütt, A., see Hildebrand, M. 414(1987)217
- Schutter, J.A. De, Van der Weken, G., Van den Bossche, W. and De Moerloose, P.  
Determination of S-carboxymethylcysteine in serum by reversed-phase ion-pair LC with column switching following pre-column derivatization with *o*-phthalaldehyde 428(1988)301
- Schutter, J.A. De and De Moerloose, P.  
Polar contributions of the stationary phase to the reversed-phase ion-pair HPLC separation of quaternary ammonium drugs 437(1988)83
- Schutter, J.A. De and De Moerloose, P.  
Determination of oxybutinin chloride in pharmaceuticals by reversed-phase ion-pair LC with two counter-ions in the eluent 450(1988)337
- Schütz, H., see Nitsche, V. 420(1987)207
- Schwartz, B.D., see Crimmins, D.L. 443(1988)63
- Schwartz, T.W., see Gether, U. 447(1988)341
- Schwendener, R.A., see Schott, H. 441(1988)115
- Schwenk, S., Altmayer, B. and Eichhorn, K.W.  
Simultaneous detection of trichothecenes and rosenonolactone in grape juice and wine by capillary GC 448(1988)424
- Sciacca, M.A., Duncan, G.F., Shea, J.P., Faulkner, III, H.C., Farmen, R.H. and Pittman, K.A.  
Simultaneous quantitation of buspirone and 1-(2-pyrimidinyl)piperazine in human plasma and urine by capillary GC-MS 428(1988)265
- Sclavons, M., see Goncalves, J.C.S. 411(1987)472
- Scolla, P.L. and Hurtubise, R.J.  
Application of the Snyder displacement model to aromatic nitrogen compounds on an aminopropyl bonded-phase column 405(1987)107
- Scott, A.I., see Gonzalez, M.D. 437(1988)311
- Scott, M.C. and Elchisak, M.A.  
Direct measurement of dopamine O-sulfate in plasma and cerebrospinal fluid 413(1987)17
- Scott, W.J., see Wittfoht, W. 448(1988)433
- Scriba, G.K.E., Borchardt, R.T., Zirrolli, J.A. and Fennessey, P.V.  
Selected-ion monitoring GC-MS analysis of catecholamines: enhancement of sensitivity by a simple clean-up step on Sephadex G-10 433(1988)31
- Scrimgeour, C.M., see Corbett, M.E. 419(1987)263
- Scuderi, F., see Flamini, G. 421(1987)434
- Sculier, J.P., see Brassinne, C. 419(1987)401
- Seaman, M.B., see Benedek, K. 444(1988)191
- Sébédio, J.L., Astorg, P.O., Septier, C. and Grandgirard, A.  
Quantitative analyses of polar components in frying oils by the Iatroscan TLC-flame ionization detection technique 405(1987)371
- Sebille, B., see Millot, M.-C. 408(1987)263
- Sébille, B., Thuaud, N., Piquion, J. and Behar, N.  
Determination of association constants of  $\beta$ -cyclodextrin and  $\beta$ -cyclodextrin-bearing polymers with drugs by HPLC 409(1987)61
- Secchi, C., see Biondi, P.A. 411(1987)275
- Secor, H.V., see Armstrong, D.W. 411(1987)490
- Šecová, E., Tichá, M., Kocourek, J. and Dey, P.M.  
Electrophoretic study of  $\alpha$ -D-galactosidases from seeds of Glycine soja and Vigna radiata possessing erythroagglutinating activity 436(1988)59

- Sedgwick, G.W., see Early, R.J. 416(1987)15
- Seeman, J.I., see Armstrong, D.W. 411(1987)490
- Segawa, Y., see Honda, S. 419(1987)51
- Segelman, F.H., see Pivonka, J. 420(1987)89
- Segl, W.E., see Pivonka, J. 420(1987)89
- Segura, R.  
Preparation of fatty acid methyl esters by direct transesterification of lipids with aluminium chloride-methanol 441(1988)99
- Seipel, L., see Krämer, B.K. 427(1988)351
- Seipel, L., see Röss, K. 417(1987)465
- Sejersted, O.M., see Øvrebø, S. 416(1987)111
- Seki, T., Hashida, N. and Kanazawa, T.  
Determination of tetraprenylacetone in human plasma by HPLC with fluorescence derivatization using dansylhydrazine 424(1988)410
- Sekiba, K., see Suzuki, Y. 426(1988)33
- Sekiguchi, M., see Tokuma, Y. 415(1987)156
- Sekine, T., see Hashimoto, K. 438(1988)401
- Sekino, Y., see Nakano, K. 438(1988)61
- Selinger, K., Matheou, D. and Hill, H.M.  
HPLC method for the determination of miconazole in vaginal fluid 434(1988)259
- Sellergren, B., Ekberg, B., Albertsson, P.-Å. and Mosbach, K.  
Preparative chiral separation in an aqueous two-phase system by a few counter-current extractions 450(1988)277
- Sellers, Jr, T., see Wooten, A.L. 445(1988)371
- Sellier, N., see Cartier, J.-P. 420(1987)121
- Sello, G., see Muda, M. 434(1988)191
- Semba, J.-I., Watanabe, A. and Takahashi, R.  
Determination of plasma homovanillic acid by two-step solid-phase extraction and HPLC with electrochemical detection 430(1988)118
- Semenov, V.A., see Rodchenkov, G.M. 423(1987)15
- Semenov, V.A., see Rodchenkov, G.M. 426(1988)399
- Semenov, V.A., see Rodchenkov, G.M. 432(1988)283
- Semmes, R.L.O. and Shen, D.D.  
Capillary GC assay for valproic acid and its 2-desaturated metabolite in brain and plasma 432(1988)185
- Semyonov, S.N.  
Near-wall field-flow fractionation. Application of surface forces and transversal voltage to fractionation in laminar flow 446(1988)131
- Semyonov, S.N., Kononenko, V.L. and Shimkus, Y.K.  
Integral Doppler anemometry and analytical field-flow fractionation 446(1988)141
- Semyonov, S.N. and Maslow, K.I.  
Acoustic field-flow fractionation 446(1988)151
- Seneca, V., see De Smet, M. 416(1987)375
- Senn, J.P., see Veuthey, J.L. 445(1988)183
- Sennello, L.T., see Granneman, G.R. 413(1987)199
- Senō, M., see Yamauchi, Y. 408(1987)171
- Seo, Y.-H., see Dietrich, A.M. 436(1988)229
- Seppänen, T., see Ekundayo, O. 403(1987)358
- Septier, C., see Sébédio, J.L. 405(1987)371
- Serafin, R.J., see Karnes, H.T. 424(1988)398
- Serpinet, J., see Chartier, A. 438(1988)263
- Serwer, P.  
Agarose gel electrophoresis of bacteriophages and related particles (Review) 418(1987)345
- Sesartić, L., see Tomašić, J. 440(1988)405
- Sestokas, E., see Chiu, S.-H.L. 433(1988)217
- Severini, G. and Aliberti, L.M.  
Biochemical indices of renal damage: separation of urinary alanine aminopeptidase by LC 421(1987)344
- Seviour, J.A., McGill, A.C., Dale, G. and Craft, A.W.  
Method of measurement of urinary homovanillic acid and vanillylmandelic acid by GC-MS suitable for neuroblastoma screening 432(1988)273
- Sewing, K.-F., see Christians, U. 413(1987)121
- Seybold, V.S., see Igwe, O.J. 432(1988)113
- Shaar, C.J., see Riggan, R.M. 435(1988)307
- Shader, R.I., see Kennedy, J.S. 422(1987)274
- Shader, R.I., see Kennedy, J.S. 423(1987)373
- Shadrina, N.E., Dmitrenko, A.V., Pavlova, V.F. and Ivanchev, S.S.  
Investigation by pyrolysis-GC of the composition of multicomponent polymeric microheterogeneous systems based on some vinyl monomers 404(1987)183
- Shafer, S.G., see Karr, L.J. 442(1988)219
- Shah, A.J. and Adlard, M.W.  
Determination of  $\beta$ -lactams and their biosynthetic intermediates in fermentation media by pre-column derivatization followed by fluorescence detection 424(1988)325
- Shah, J.A., see Kaiser, D.G. 434(1988)135
- Shah, J.A., Weber, D.J. and Bothwell, B.E.  
HPLC determination of methylprednisolone and methylprednisolone 21-[8-[methyl-(2-sulfoethyl)amino]-8-oxooctanoate] sodium salt in human plasma 414(1987)1
- Shah, P., see Ravichandran, K. 439(1988)213
- Shaner, D.L., see Singh, B.K. 444(1988)251
- Sharma, A.K., see Schneiderman, M.A. 409(1987)343

- Sharma, M.L., see Furuta, T. 427(1988)41
- Sharma, R., see Subbaramaiah, K. 408(1987)385
- Sharman, M., see Castle, L. 437(1988)247
- Sharp, E.A., see Hunter, K. 435(1988)83
- Sharp, E.A., see Hunter, K. 437(1988)301
- Sharp, G.J., see Brayan, J.G. 447(1988)249
- Sharp, S., see Baty, J.D. 437(1988)13
- Shea, E.M. and Carpita, N.C.  
Separation of partially methylated alditol acetates on SP-2330 and HP-1 vitreous silica capillary columns 445(1988)424
- Shea, J.P., see Sciacca, M.A. 428(1988)265
- Shea, T.C., see Henner, W.D. 416(1987)426
- Shelly, D.C. and Edkins, T.J.  
Contribution to the packing of LC microcolumns 411(1987)185
- Shen, D.D., see Kwong, E.C. 414(1987)365
- Shen, D.D., see Semmes, R.L.O. 432(1988)185
- Shephard, G.S., Louw, M.E.J. and Labadarios, D.  
Analysis of vitamin B<sub>6</sub> vitamers in plasma by cation-exchange HPLC 416(1987)138
- Shepherd, M.J., see Bayliss, M.A.J. 445(1988)393
- Shepherd, M.J., see Bayliss, M.A.J. 445(1988)403
- Shepherd, M.J., see Mortimer, D.N. 407(1987)393
- Sh Sheridan, M.E. and Clarke, G.S.  
Improved HPLC determination of doxycycline in serum and urine using solid-phase extraction columns 434(1988)253
- Shetty, P.H., Youngberg, P.J., Kersten, B.R. and Poole, C.F.  
Solvent properties of liquid organic salts used as mobile phases in microcolumn reversed-phase LC 411(1987)61
- Sheu, C., see Chang, C.A. 404(1987)282
- Shiao, M.-S., see Lin, L.-J. 410(1987)195
- Shiba, M., see Hirayama, C. 409(1987)175
- Shibanoki, S., Imamura, Y., Arakawa, Y. and Ishikawa, K.  
Determination of dosulepin and its metabolite: application of HPLC with electrochemical detection 415(1987)365
- Shibanoki, S., Imamura, Y., Itoh, T., Ishikawa, K., Noda, Y., Yazaki, S. and Suzuki, H.  
Application of HPLC with electrochemical detection for monitoring the concentration of pentazocine in human blood 421(1987)425
- Shibasaki, H., see Furuta, T. 413(1987)1
- Shibata, K.  
Fluorimetric micro-determination of kynurenic acid, an endogenous blocker of neurotoxicity, by HPLC 430(1988)376
- Shibata, K., Kawada, T. and Iwai, K.  
HPLC determination of nicotinamide in rat tissue samples and blood after extraction with diethyl ether 422(1987)257
- Shibata, K., Kawada, T. and Iwai, K.  
Simultaneous micro-determination of nicotinamide and its major metabolites, N<sup>1</sup>-methyl-2-pyridone-5-carboxamide and N<sup>1</sup>-methyl-4-pyridone-3-carboxamide, by HPLC 424(1988)23
- Shibata, K., Kawada, T. and Iwai, K.  
Microdetermination of N<sup>1</sup>-methyl-2-pyridone-5-carboxamide, a major metabolite of nicotinic acid and nicotinamide, in urine by HPLC 417(1987)173
- Shibata, K., Onodera, M., Kawada, T. and Iwai, K.  
Simultaneous micro-determinations of serotonin and 5-hydroxyindole-3-acetic acid with 5-hydroxy-N<sub>w</sub>-methyltryptamine, as an internal standard, in biological materials by HPLC with electrochemical detection 430(1988)381
- Shibata, M., see Niwa, N.T.T. 431(1988)418
- Shibata, T., see Wainer, I.W. 411(1987)139
- Shibazaki, T., see Hayashi, Y. 407(1987)59
- Shibazaki, T., see Hayashi, Y. 411(1987)95
- Shibukawa, A., see Nakagawa, T. 420(1987)297
- Shida, T., see Mita, H. 430(1988)299
- Shieh, C.H., Karger, B.L., Gelber, L.R. and Feibush, B.  
Ligand exchange chromatography of amino alcohol enantiomers as Schiff bases 406(1987)343
- Shillcutt, S.D., see Hoffman, D.W. 414(1987)504
- Shimada, K., Oe, T. and Nambara, T.  
Sensitive ferrocene reagents for derivatization of thiol compounds in HPLC with dual-electrode coulometric detection 419(1987)17
- Shimamura, T., see Hashimoto, H. 448(1988)182
- Shimamura, T., see Koyama, Y. 439(1988)417
- Shimizu, H., see Utsumi, J. 421(1987)355
- Shimizu, I., see Sato, H. 447(1988)387
- Shimizu, K., Hara, T., Yamaga, N., Kohara, H. and Nojima, K.  
Determination of 17-hydroxyprogesterone in plasma by GC-MS with high-resolution selected-ion monitoring 432(1988)21
- Shimizu, K., see Yamaga, N. 422(1987)25
- Shimizu, R., see Fujimoto, Y. 402(1987)344
- Shimizu, T., see Shioya, Y. 434(1988)283
- Shimkus, Y.K., see Semyonov, S.N. 446(1988)141
- Shimoda, W., see Munns, R.K. 442(1988)209
- Shimono, N., see Nakagawa, T. 420(1987)297
- Shimosato, K., see Tomita, M. 414(1987)454
- Shin, K., see Achari, R. 424(1988)430
- Shin, S., see Park, K.S. 440(1988)225

- Shinbo, T., Yamaguchi, T., Nishimura, K. and Sugiura, M.  
Chromatographic separation of racemic amino acids by use of chiral crown ether-coated reversed-phase packings 405(1987)145
- Shinde, S.K., see Ramachandran, P.K. 426(1988)239
- Shinohara, T., see Iida, T. 438(1988)423
- Shiono, H., see Matsubara, K. 424(1988)49
- Shioya, Y. and Shimizu, T.  
HPLC procedure for the determination of a new anti-gastric ulcer agent, 2-(4-chlorobenzoylamino)-3-[2(1H)-quinolinon-4-yl]-propionic acid, in human plasma and urine 434(1988)283
- Shirai, Y., see Hashimoto, K. 448(1988)241
- Shishkina, I.P., see Galushko, S.V. 445(1988)59
- Shitanishi, K.T. and Herring, S.W.  
Separation of human factor X from factor Xa by reversed-phase HPLC 444(1988)107
- Shively, J.E., see Powell, D.R. 420(1987)163
- Shmidel, E.B. and Kalabina, L.I.  
Mechanism governing the sensitivity of a three-electrode ionization chamber 446(1988)163
- Shoeman, D., see Ansari, K.A. 439(1988)453
- Shonia, N.K., see Nikitin, Y.S. 446(1988)55
- Shono, T., see Tanaka, M. 404(1987)175
- Shono, T., see Tanaka, M. 438(1988)253
- Shopova, B., see Hjertén, S. 440(1988)305
- Shorr, R.G.L., see Strohsacker, M.W. 435(1988)185
- Shukun, S.A., Gavryushkin, A.V., Brezgunov, V.N. and Zav'yalov, V.P.  
Free-flow electrophoretic apparatus for separation and concentration of proteins 432(1988)137
- Shulgin, A.T., Jacob III, P., Benowitz, N.L. and Lau, D.  
Identification and quantitative analysis of cotinine-N-oxide in human urine 423(1987)365
- Shustina, R., see Lesser, J.H. 410(1987)95
- Sica, A., see Latini, R. 424(1988)211
- Siddiqui, A.H., see Stolk, L.M.L. 423(1987)383
- Sidén, Å., see Gallo, P. 416(1987)53
- Siebert, C.J., see Hjertén, S. 403(1987)47
- Siebert, D.M., see Chen, Z.R. 424(1988)170
- Siebert, D.M. and Bochner, F.  
Determination of plasma aspirin and salicylic acid concentrations after low aspirin doses by HPLC with post-column hydrolysis and fluorescence detection 420(1987)425
- Siefermann-Harms, D.  
HPLC of chloroplast pigments. One-step separation of carotene and xanthophyll isomers, chlorophylls and pheophytins 448(1988)411
- Siegfried, B., see Vachta, J. 428(1988)183
- Siems, W., see Förster, H.-G. 424(1988)357
- Siems, W., see Werner, A. 421(1987)257
- Siest, G., see Decolin, D. 414(1987)499
- Siest, G., see Decolin, D. 419(1987)469
- Siest, G., see Motassim, N. 422(1987)340
- Sievers, R.E., see DeAngelis, J.J. 441(1988)125
- Siezen, R.J. and Kaplan, E.D.  
Optimal resolution of eye lens  $\gamma$ -crystallins by cation-exchange HPLC on Synchropak CM300 444(1988)239
- Sikora, E., Krzyżanowski, L. and Rzeszutarska, B.  
Separation of N-(4-aminobenzoyl)- $\gamma$ -oligo(L-glutamic acid)s on a thin-layer plate and their quantitation using fluorescamine 442(1988)420
- Sil, P., see Ghosh, P. 403(1987)285
- Silber, B.M., see Falkowski, A.J. 422(1987)145
- Silink, M., see Kreutzmann, D.J. 415(1987)253
- Sillinger, V., see Gottvaldova, M. 427(1988)331
- Sim, P.G., Elson, C.M. and Quilliam, M.A.  
Use of the photoionisation detector in packed-column supercritical fluid chromatography 445(1988)239
- Šimánek, V., see Válka, I. 445(1988)258
- Šimánek, V., see Walterová, D. 405(1987)389
- Simard, R.E., see Lavigne, C. 410(1987)201
- Simmonds, P., see Begley, P. 445(1988)119
- Simmons, J.E., see Carlin, A.S. 425(1988)162
- Simon, G., see Papp, O. 403(1987)19
- Simonyi, M., see Fitos, I. 450(1988)217
- Sing, K.S.W., see Carrott, P.J.M. 406(1987)139
- Sing, R.P. and Nancollas, G.H.  
Determination of phosphate, sulfate and oxalate in urine by ion chromatography 433(1988)373
- Singh, A.K., Mishra, U., Ashraf, M., Abdennebi, E.H., Granley, K., Dombrovskis, D., Hewetson, D. and Stowe, C.M.  
Analysis of detomidine in horse blood, plasma and urine samples utilizing a sensitive GC-MS method 404(1987)223
- Singh, A.K. and Ashraf, M.  
Analysis of amino acids in brain and plasma samples by sensitive GC-MS 425(1988)245
- Singh, B.K., Stidham, M.A. and Shaner, D.L.  
Separation and characterization of two forms of acetohydroxy acid synthase from black Mexican sweet corn cells 444(1988)251
- Singh, J., see Dhillon, R.S. 435(1988)256

- Singh, P.P. and Chawla, R.P.  
TLC of flucythrinate residues 450(1988)452
- Singh, U.S., see Adinarayana, M. 435(1988)210
- Singhal, A.K., see Duff, G.A. 416(1987)71
- Singleton, W., see Mefford, I.N. 420(1987)241
- Sinha, P., see Meyer-Sabellek, W. 429(1988)419
- Sinibaldi, M., see Fanali, S. 408(1987)441
- Sinibaldi, M., see Fanali, S. 442(1988)371
- Sinkule, J.A., see Choi, K.E. 420(1987)81
- Sinninghe Damsté, J.S., Kock-Van Dalen, A.C.,  
De Leeuw, J.W. and Schenck, P.A.  
Identification of homologous series of  
alkylated thiophenes, thiolanes, thianes and  
benzothiophenes present in pyrolysates of  
sulphur-rich kerogens 435(1988)435
- Sinou, D., see Lecomte, L. 408(1987)416
- Siouffi, A.M., see Righezza, M. 410(1987)145
- Sioufi, A., Colussi, D., Marfil, F. and Dubois,  
J.P.  
Determination of the (+)- and (-)-  
enantiomers of pirprofen in human plasma  
by HPLC 414(1987)131
- Sioufi, A., Kaiser, G., Leroux, F. and Dubois,  
J.P.  
Determination of the *S*(+)- and *R*(-)-  
enantiomers of baclofen in plasma and urine  
by GC using a chiral fused-silica capillary  
column and an electron-capture detector  
450(1988)221
- Sioufi, A., Pommier, F., Kaiser, G. and Dubois,  
J.P.  
Determination of benazepril, a new  
angiotensin-converting enzyme inhibitor, and  
its active metabolite, benazeprilat, in plasma  
and urine by capillary GC-mass-selective  
detection 434(1988)239
- Sioufi, A., Pommier, F. and Dubois, J.P.  
Determination of the two dinitrate  
metabolites of nitroglycerin in human plasma  
by capillary GC with electron-capture  
detection 413(1987)101
- Sioufi, A., Pommier, F. and Dubois, J.P.  
Simultaneous determination of clomipramine  
and its *N*-desmethyl metabolite in human  
whole blood by capillary GC with mass-  
selective detection 428(1988)71
- Siow, Y.F., see Baker, P.R. 423(1987)63
- Sirois, G., see Colin, P. 413(1987)151
- Sirtori, C.R., see Paroni, R. 420(1987)189
- Sisco, W.R., see Schrader, P.A. 408(1987)430
- Sitaram, B.R., Lockett, L., McLeish, M.,  
Hayasaka, Y., Blackman, G.L. and McLeod,  
W.R.  
GC-MS characterisation of the  
psychotomimetic indolealkylamines and their  
*in vivo* metabolites 422(1987)13
- Siu, K.W.M., Berman, S.S. and Aue, W.A.  
Correlations and constants in unipolar and  
bipolar electron-capture detection  
408(1987)53
- Siu, K.W.M. and Aue, W.A.  
An experiment to distinguish between  
homogeneous and heterogeneous response  
mechanisms in a parallel-plate electron-  
capture detector 406(1987)81
- Sjöberg, P., see Lindström, B. 428(1988)156
- Sjöström, J. and Holmbom, B.  
SEC of deposits in pulp and paper mills  
411(1987)363
- Sjövall, J., see Norén, K. 414(1987)55
- Sjövall, J., see Norén, K. 422(1987)103
- Škabrahová, Z., Kovář, J., Glatz, Z., Turánek, J.  
and Kahle, V.  
HPLC purification of the H4 and MH3  
isoenzymes of lactate dehydrogenase from  
beef heart 408(1987)372
- Skarping, G., Dalene, M. and Mathiasson, L.  
Trace analysis of airborne 1,6-  
hexamethylenediisocyanate and the related  
aminoisocyanate and diamine by glass  
capillary GC 435(1988)453
- Skarping, G., see Dalene, M. 435(1988)469
- Skelly, J.P., see Carlin, A.S. 425(1988)162
- Sklan, D., see Knobler, H. 421(1987)136
- Skrinska, V.A., see Rebec, M.V. 432(1988)375
- Skuse, D.R. and Brooks, D.E.  
Column-based separation of erythrocytes  
using aqueous polymeric two-phase systems  
432(1988)127
- Skuterud, B., see Christophersen, A.S.  
422(1987)117
- Slabý, J., see Kocna, P. 434(1988)429
- Slacanin, I., Marston, A. and Hostettmann, K.  
HPLC determination of molluscicidal  
saponins from *Phytolacca dodecandra*  
(Phytolaccaceae) 448(1988)265
- Sládek, T., see Šalamoun, J. 420(1987)197
- Šlais, K.  
Simultaneous conductivity and amperometric  
detector with a single cell for microbore LC  
436(1988)413
- Šlais, K.  
Amperometric detector with a manganin  
electrode for microcolumn LC 442(1988)111
- Slanina, J., see Keuken, M.P. 439(1988)13
- Slattery, T.K., see Burton, W.G. 443(1988)363
- Slattery, T.K., see Nugent, K.D. 443(1988)381
- Slaughter, R.L., see Kong, A.-N. 432(1988)308
- Slegowski, M.B., Miller, C. and Porter, R.S.

- Simplified HPLC determination of flosequinan and its metabolite in plasma, serum and urine 425(1988)227
- Slevin, M.L., see Joel, S.P. 430(1988)394
- Slikker, Jr., W., see Holder, C.L. 419(1987)113
- Slocum, R., see Hubbard, R.W. 431(1988)163
- Slováková, S., see Mikeš, O. 440(1988)287
- Sluyterman, L.A.Æ., see Kooistra, C. 436(1988)173
- Sluyterman, L.A.Æ., see Kooistra, C. 436(1988)181
- Smallwood, R.A., see Mihaly, G.W. 415(1987)177
- Smet, M. De, Van Belle, S.J.P., Seneca, V., Storme, G.A. and Massart, D.L.  
Determination of vinblastine in MO<sub>4</sub> mouse fibrosarcoma cells by HPLC 416(1987)375
- Smet, M. De and Massart, D.L.  
Retention behaviour of acidic, neutral and basic drugs on a CN column using phosphate buffers in the mobile phase 410(1987)77
- Smidt, C.R., Jones, A.D. and Clifford, A.J.  
Gas chromatography of retinol and  $\alpha$ -tocopherol without derivatization 434(1988)21
- Smilde, A.K., Bruins, C.H.P., Doornbos, D.A. and Vink, J.  
Optimization of the reversed-phase HPLC separation of synthetic estrogenic and progestogenic steroids using the multi-criteria decision making method 410(1987)1
- Smith, A.I., Wallace, C.A., Cheng, M.C. and Autelitano, D.J.  
Combined use of size-exclusion and reversed-phase HPLC for characterization of  $\beta$ -endorphin processing pathways 416(1987)255
- Smith, C.S., Morgan, S.L., Greene, S.V. and Abramson, R.K.  
Solid-phase extraction and HPLC method for chlorpromazine and thirteen metabolites 423(1987)207
- Smith, D.J., see Amacher, D.E. 419(1987)61
- Smith, D.J., see Dunphy, M.J. 420(1987)394
- Smith, D.W., see Dutta, I. 444(1988)183
- Smith, H.T.  
HPLC method for the determination of pindolol in human plasma 415(1987)93
- Smith, H.T. and Molinaro, N.C.  
HPLC method for the determination of methysergide and methylergonovine in human plasma 424(1988)416
- Smith, J.D., Mello, C.M. and O'Reilly, D.J.  
Separation of phospholipids and phosphonolipids of *Tetrahymena* by HPLC 431(1988)395
- Smith, L.M., see Carilli, C.T. 444(1988)203
- Smith, P.B., see Cutié, S.S. 408(1987)349
- Smith, P.C., see Hyneck, M.L. 420(1987)349
- Smith, P.L. and Cooper, W.T.  
Retention and selectivity in amino, cyano and diol normal bonded phase HPLC columns 410(1987)249
- Smith, R.A., see Tibbels, T.S. 441(1988)448
- Smith, R.E., Howell, S., Yourtee, D., Premkumar, N., Pond, T., Sun, G.Y. and MacQuarrie, R.A.  
Ion chromatographic determination of sugar phosphates in physiological samples 439(1988)83
- Smith, R.H.A. and Gilbert, B.E.  
Quantification of ribavirin in biological fluids and tissues by HPLC 414(1987)202
- Smith, R.M. and Garside, D.R.  
Effect of organic modifier on the separation and efficiencies of homologues on porous polymer columns 407(1987)19
- Smithurst, B.A., see Ismail, Z. 404(1987)372
- Smolková-Keulemansová, E., see Bažant, L. 445(1988)337
- Smolková-Keulemansová, E., see Jelínek, I. 405(1987)379
- Smolková-Keulemansová, E., see Jelínek, I. 435(1988)496
- Smolková-Keulemansová, E., see Jelínek, I. 439(1988)386
- Smolková-Keulemansová, E., see Snopek, J. 411(1987)153
- Smolková-Keulemansová, E., see Snopek, J. 438(1988)211
- Smolková-Keulemansová, E., see Snopek, J. 450(1988)373
- Smrž, M., see Mikeš, O. 440(1988)287
- Smrž, M., see Šalamoun, J. 419(1987)213
- Smrž, M., see Šalamoun, J. 420(1987)197
- Smyth, J.F., see Cummings, J. 431(1988)77
- Smyth, M.R., see Meaney, M. 449(1988)241
- Snider, B.G., see Clark, C.P. 408(1987)275
- Snider, B.G., see McDaniel, D.M. 404(1987)123
- Snopek, J., Jelínek, I. and Smolková-Keulemansová, E.  
Use of cyclodextrins in isotachopheresis. II.  $\alpha$ -,  $\beta$ - and  $\gamma$ -cyclodextrins as leading electrolyte additives for the separation of *ortho*-, *meta*- and *para*-substituted halogenobenzoic acids 411(1987)153
- Snopek, J., Jelínek, I. and Smolková-Keulemansová, E.  
Use of cyclodextrins in isotachopheresis. IV. The influence of cyclodextrins on the chiral



- resolution of ephedrine alkaloid enantiomers 438(1988)211
- Snopek, J., see Jelínek, I. 405(1987)379
- Snopek, J., see Jelínek, I. 435(1988)496
- Snopek, J., see Jelínek, I. 439(1988)386
- Snopek, J., Smolková-Keulemansová, E., Jelínek, I., Dohnal, J., Klinot, J. and Klinotová, E. Use of cyclodextrins in isotachopheresis. VI. Cyclodextrins as leading electrolyte additives for the separation of bile acids 450(1988)373
- Snyder, K.P., see Hurst, W.J. 409(1987)413
- Snyder, L.R., Cox, G.B. and Antle, P.E. Preparative separation of peptide and protein samples by HPLC with gradient elution. I. The Craig model as a basis for computer simulations 444(1988)303
- Snyder, L.R., see Burton, W.G. 443(1988)363
- Snyder, L.R., see Cox, G.B. 444(1988)325
- Snyder, L.R., see Eble, J.E. 405(1987)1
- Snyder, L.R., see Eble, J.E. 405(1987)31
- Snyder, L.R., see Eble, J.E. 405(1987)51
- Snyder, L.R., see Nugent, K.D. 443(1988)381
- Snyder, P.A., see Mazzo, D.J. 438(1988)85
- Snyder, R.S., see Karr, L.J. 442(1988)219
- Snyder, R.W., see Long, J.W. 450(1988)394
- Soczewiński, E., see Matysik, G. 446(1988)275
- Soerensen, K.J., see Yong, Y.-S. 450(1988)399
- Soest, R.E.J. Van, see Nielen, M.W.F. 417(1987)159
- Sofia, R.D., see Pivonka, J. 420(1987)89
- Soga, A., see Yanagi, H. 417(1987)178
- Sohler, E., see Bystrický, Ľ. 447(1988)202
- Soine, P.J. and Soine, W.H. HPLC determination of the diastereomers of 1-( $\beta$ -D-glucopyranosyl)amobarbital in urine 422(1987)309
- Soine, W.H., see Soine, P.J. 422(1987)309
- Soják, L., Ostrovský, I., Kraus, G. and Andronikashvili, T.G. High-resolution GC with liquid crystal glass capillaries. X. Separation of diastereomeric alkanes 436(1988)47
- Soják, L., Ostrovský, I. and Janák, J. Propyl effect and retention-structure correlation as a means of GC identification 406(1987)43
- Soják, L. and Ostrovský, I. Shape selectivity of stationary phases for the separation of isomeric hydrocarbons by capillary GC 446(1988)339
- Soldin, S.J., see Tan, Y.K. 422(1987)187
- Solovyova, L.Y., see Gorbunov, A.A. 448(1988)307
- Solymsi, J., see Nagy, L.G. 406(1987)359
- Sommadossi, J.-P. and Bevan, R. HPLC method for the determination of 9-(1,3-dihydroxy-2-propoxymethyl)guanine in human plasma 414(1987)429
- Somogyi, A., see Cserháti, T. 446(1988)17
- Somogyi, A.A., see Chen, Z.R. 424(1988)170
- Somorjai, G.A., see Prins, R. 406(1987)151
- Sonesson, A., Larsson, L., Fox, A., Westerdahl, G. and Odham, G. Determination of environmental levels of peptidoglycan and lipopolysaccharide using GC with negative-ion chemical-ionization MS utilizing bacterial amino acids and hydroxy fatty acids as biomarkers 431(1988)1
- Sonesson, A., Larsson, L., Westerdahl, G. and Odham, G. Determination of endotoxins by GC: evaluation of electron-capture and negative-ion chemical-ionization MS detection of halogenated derivatives of  $\beta$ -hydroxymyristic acid 417(1987)11
- Sonesson, A., Larsson, L. and Jimenez, J. Use of pentafluorobenzyl and pentafluoropropionyl-pentafluorobenzyl esters of bacterial fatty acids for GC analysis with electron-capture detection 417(1987)366
- Sonnet, P.E., Piotrowski, E.G. and Boswell, R.T. Chromatographic methods for determining the configuration of 1,2- and 1,3-(unsymmetrically substituted) dialkylglycerol ethers 436(1988)205
- Sonneveld, P., see Van der Steuijt, K. 422(1987)328
- Soons, P.A. and Breimer, D.D. GC analysis of nitrendipine and its pyridine metabolite in human plasma 428(1988)362
- Sørensen, H. LC determination of fusidic acid in serum 430(1988)400
- Sørensen, H., see Larsen, L.M. 450(1988)121
- Sørensen, H.H., see Welinder, B.S. 408(1987)191
- Souther, M.E., Lumpkin, R.H., Kuo, K.C., Reams, G.P., Bauer, J.H. and Gehrke, C.W. HPLC-radioimmunoassay method for the measurement of angiotensin II peptides in human plasma 417(1987)27
- Souvannavong, V. and Adam, A. N-Acetylmuramyl-L-alanyl-D-isoglutamine stimulates the immune response of B cells. Use of fast protein LC to purify a novel monokine involved in this effect 440(1988)95
- Sova, O., see Dobránský, T. 411(1987)486
- Sowell, A.L., Huff, D.L., Gunter, E.W. and Driskell, W.J. Identification of *cis*-carotenoids in human

- sera analyzed by reversed-phase HPLC with diode array detection 431(1988)424
- Spagnolo, G., see Del Principe, D. 419(1987)329
- Spahn, H.  
Formation of diastereomeric derivatives of 2-arylpropionic acids using L-leucinamide 423(1987)334
- Spahn, H.  
S-(+)-Flunoxapfen chloride as chiral fluorescent reagent 427(1988)131
- Spahn, H.  
Assay method for product formation in *in vitro* enzyme kinetic studies of uridine diphosphate glucuronyltransferases: 2-arylpropionic acid enantiomers 430(1988)368
- Spahn, H.  
Formation of diastereomeric derivatives of 2-arylpropionic acids using  $\alpha$ -leucinamide: the derivatization principle and its applicability. A reply 431(1988)229
- Spahn, H., see Gietl, Y. 426(1988)305
- Spahn, H., see Pflugmann, G. 416(1987)331
- Spahn, H., see Pflugmann, G. 421(1987)161
- Spahn, H., Spahn, I., Pflugmann, G., Mutschler, E. and Benet, L.Z.  
Measurement of carprofen enantiomer concentrations in plasma and urine using L-leucinamide as the chiral coupling component 433(1988)331
- Spahn, I., see Spahn, H. 433(1988)331
- Spalik, J., see Long, J.W. 450(1988)394
- Spalluto, R., see Visconti, M. 416(1987)433
- Spangenberg, S., see Klemm, D. 438(1988)122
- Speedie, M.K., see Vicchio, D. 415(1987)104
- Speth, P.A.J., see Brons, P.P.T. 422(1987)175
- Spierenburg, T.J., see De Graaf, G.J. 447(1988)244
- Spiker, S., Bates, M. and Arwood, L.J.  
Fractionation of plant and animal high mobility group chromosomal proteins by ion-exchange and reversed-phase HPLC 405(1987)327
- Spinazzola, J., see Rop, P.P. 427(1988)172
- Spino, L.A., Armstrong, D.W. and Vo-Dinh, T.  
Direct synchronous luminescence detection of co-eluting solutes in pseudophase LC 409(1987)147
- Spino, L.A., see Armstrong, D.W. 411(1987)490
- Spino, M., see Strong, H.A. 422(1987)301
- Spiteller, G., see Dietel, P. 422(1987)1
- Spiteller, G., see Jandke, J. 421(1987)1
- Spiteller, G., see Puchta, V. 420(1987)450
- Spitzer, T. and Kuwatsuka, S.  
Simultaneous clean-up of nitroarenes and polycyclic aromatic ketones from soil and particulate matter on XAD-2 435(1988)489
- Splinter, T.A.W., see Verkoelen, C.F. 426(1988)41
- Spooner, N., see Evershed, R.P. 440(1988)23
- Spreux-Varoquaux, O., Morin, D., Advenier, C. and Pays, M.  
Determination of clomipramine and its hydroxylated and demethylated metabolites in plasma and urine by LC with electrochemical detection 416(1987)311
- Springolo, V. and Coppi, G.  
Simple method for the determination of rhein in biological fluids by HPLC 428(1988)173
- Šprta, V., see Šalamoun, J. 420(1987)197
- Srinivas, N.R., Hubbard, J.W., Cooper, J.K. and Midha, K.K.  
Enantioselective GC assay with electron-capture detection for *dl*-fenfluramine and *dl*-norfenfluramine in plasma 433(1988)105
- Sriramachari, S., see Ramachandran, P.K. 426(1988)239
- Srivastava, N., see Kharbade, B.V. 439(1988)430
- Staats, S.J., see Kunitani, M.G. 443(1988)205
- Stäb, J.A., see Maris, F.A. 445(1988)129
- Stacey, V.E., see Matlin, S.A. 450(1988)450(1988)157
- Stafford, A.E., Corse, J., Lyman, M. and Metzner, K.E.  
*E*-Isozeatin and *cis*-norzeatin riboside as internal standards for cytokinin analyses 436(1988)93
- Stafford, J.E.H., see Jackson, L.S. 428(1988)377
- Staub, W., see Radeck, W. 432(1988)297
- Stalcup, A.M., Martire, D.E. and Wise, S.A.  
Thermodynamic comparison of monomeric and polymeric C<sub>18</sub> bonded phases using aqueous methanol and acetonitrile mobile phases 442(1988)1
- Stan, H.-J., see Lipinski, J. 441(1988)213
- Stancher, B., see Zonta, F. 403(1987)207
- Stancher, B., Zonta, F. and Favretto, L.G.  
HPLC of carotenoids from some marine shellfish 440(1988)37
- Stanski, D.R., see Björkman, S. 433(1988)95
- Staroverov, S.M., see Fadeev, A.Y. 447(1988)103
- Staroverov, S.M., see Gajda, A.V. 424(1988)385
- Starzec, A., Counis, R. and Jutisz, M.  
Several intermediate forms in the processing of rat lutropin subunits as shown by sodium dodecyl sulphate polyacrylamide gel electrophoresis 440(1988)353
- Staveris, S., see Efthymiopoulos, C. 421(1987)360
- Stead, C.V., see Burton, S.J. 435(1988)127
- Stearns, R.A., see Metz, P.A. 441(1988)31
- Stecher, V.J., see Pruden, D.J. 437(1988)399

- Steege, G. Van der, Muskiet, F.A.J., Martini, I.A., Hutter, N.H. and Boersma, E.R.  
Simultaneous quantification of total medium- and long-chain fatty acids in human milk by capillary GC with split injection 415(1987)1
- Steele, N.C., see Richards, M.P. 402(1987)243
- Stefanova, M. and Lazarov, L.  
GC-MS study of coal-derived asphaltenes. II. Carbazole-type neutral nitrogen compounds 446(1988)293
- Štefek, M. and Beneš, L.  
GC determination of stobadine, N-desmethylstobadine and stobadine N-oxide in rat liver microsomal incubations 415(1987)163
- Steffenrud, S., Borgeat, P., Evans, M.J. and Bertrand, M.J.  
MS evaluation of the *tert*-butyldimethylsilyl derivatives of monohydroxyeicosatetraenoic acids and leukotrienes 423(1987)1
- Steffenrud, S., Borgeat, P., Salari, H., Evans, M.J. and Bertrand, M.J.  
GC-MS of monohydroxyeicosatetraenoic acids as their methyl esters trimethylsilyl, allyldimethylsilyl and *tert*-butyldimethylsilyl ethers 416(1987)219
- Steffenrud, S. and Salari, H.  
Reversed-phase ion-interaction chromatography of leukotrienes, lipoxins and related compounds 427(1988)1
- Stegink, S.J., see Reine, A.H. 437(1988)211
- Stelke, P., Fürst, P., Ratz, R. and Rau, H.  
Isotachophoresis of quaternary 4,4'-bipyridylum salts. Analytical control of synthesis and purification procedures 449(1988)299
- Stehlik, F., see Neča, J. 447(1988)177
- Stein, A., see Klemm, D. 438(1988)122
- Stein, S., see Tous, G. 444(1988)67
- Steinhilber, D., see Herrmann, T. 416(1987)170
- Steinhilber, D., see Herrmann, T. 428(1988)237
- Steinhilber, D., see Knospe, J. 442(1988)444
- Steinmetz, Y., see Garty, M. 430(1988)123
- Stelling-Ferrara, M., see Puglisi, C.V. 419(1987)253
- Stephany, R.W., see Van Ginkel, L.A. 445(1988)385
- Stephens, R.E.  
Separation of tubulin subunits by reversed-phase HPLC 448(1988)371
- Stetter, J.R., see Otagawa, T. 408(1987)81
- Steuer, W., Schindler, M., Schill, G. and Erni, F.  
Supercritical fluid chromatography with ion-pairing modifiers. Separation of enantiomeric 1,2-aminoalcohols as diastereomeric ion pairs 447(1988)287
- Steuijt, K. Van der and Sonneveld, P.  
Concurrent analysis of methotrexate, trimethoprim, sulphamethoxazole and their major metabolites in plasma by HPLC 422(1987)328
- Stevens, Y.-W., Walsh, T.D. and Fanucchi, M.P.  
Quantitation of merbarone in human plasma by HPLC 416(1987)203
- Stevenson, D.K., see Vreman, H.J. 427(1988)315
- Stewart, J.T., see Arrendale, R.F. 432(1988)165
- Steyn, J.M. and Hundt, H.K.L.  
GC-MS method for the quantitation of cantharidin in human serum 432(1988)177
- Sticher, O., see Dallenbach-Toelke, K. 404(1987)365
- Sticher, O., see Hofstetter, M. 407(1987)319
- Sticher, O., see Meier, B. 442(1988)175
- Sticher, O., see Nyireddy, S. 450(1988)241
- Stidham, M.A., see Singh, B.K. 444(1988)251
- Stiffin, R.M., see Wainer, I.W. 411(1987)139
- Stiffin, R.M., see Wainer, I.W. 424(1988)158
- Stipetic, M., see Mefford, I.N. 420(1987)241
- St. Louis, R., see Gholson, A.R. 408(1987)329
- Stobaugh, J.F., see Crouch, Jr., F.W. 448(1988)333
- Stodola, J.D., see Hartman, P.A. 444(1988)177
- Štol, M., Adam, M. and Deyl, Z.  
Biospecific affinity chromatography of plasma fibronectin on cross-linked gelatin microbeads 419(1987)308
- Stolk, L.M.L., De Ruiter, R., Saadawi, A., Siddiqui, A.H. and Cormane, R.N.  
Determination of psoralen in serum by reversed-phase HPLC 423(1987)383
- Stoltz, J.F., see Regnault, V. 432(1988)93
- Stone, R.B., see Barlow, R.D. 419(1987)375
- Stopher, D.A., see Beresford, A.P. 420(1987)178
- Stoppa, C., see Barbieri, L. 408(1987)235
- Stoppini, M., see Iadarola, P. 443(1988)317
- Storme, G.A., see De Smet, M. 416(1987)375
- Stover, F.S.  
Cationic capillary isotachophoresis of proteins 445(1988)417
- Stowe, C.M., see Singh, A.K. 404(1987)223
- Stowers, D.J., Keim, J.M.B., Paul, P.S., Lyoo, Y.S., Merion, M. and Benbow, R.M.  
High-resolution chromatography of nucleic acids on the Gen-Pak FAX column 444(1988)47
- St. Pierre, M.V. and Pang, K.S.  
Determination of diazepam and its metabolites by HPLC and TLC 421(1987)291
- Straczek, J., see Brossat, B. 413(1987)141

- Strašák, M. and Bystrický, S.  
Partial resolution of cobalt(III) chelate complexes by GPC 403(1987)331
- Strašák, M. and Bystrický, S.  
Rapid optical resolution of anionic metal complexes by GPC 450(1988)284
- Straub, K.M., see Garvie, C.T. 413(1987)43
- Strecker, G., see Capon, C. 425(1988)35
- Streete, J.M. and Berry, D.J.  
GC analysis of phenylethylmalonamide in human plasma 416(1987)281
- Strelkova, M.A., see Manoilov, S.E. 440(1988)53
- Strenze, T., see Rang, S. 406(1987)75
- Stringham, R.W. and Regnier, F.E.  
Selective non-adsorption preparative chromatography of bovine immunoglobulin G1 409(1987)305
- Strohsacker, M.W., Minnich, M.D., Clark, M.A., Shorr, R.G.L. and Crooke, S.T.  
Selective purification of cardiac myosin by a high-performance salicylate affinity column 435(1988)185
- Stromberg, R.  
Statistical optimization of a reversed-phase ion-pair LC method for the analysis of tolmetin sodium in dosage forms 448(1988)1
- Strong, H.A. and Spino, M.  
Highly sensitive determination of cimetidine and its metabolites in serum and urine by HPLC 422(1987)301
- Štrop, P., see Mikeš, O. 440(1988)287
- Stuart, J.D., see Eslami, M. 411(1987)121
- Stubbs, C. and Kanfer, I.  
HPLC of erythromycin propionyl ester and erythromycin base in biological fluids 427(1988)93
- Stubbs, R.J., Ng, L.L., Entwistle, L.A. and Bayne, W.F.  
Analysis of sulindac and metabolites in plasma and urine by HPLC 413(1987)171
- Stubbs, R.J., see Chiou, R. 416(1987)196
- Stubbs, R.J., see Chiou, R. 422(1987)281
- Štulík, K., Pacáková, V. and Jokuszies, G.  
Analysis of dipeptides by reversed-phase HPLC without derivatization using amperometric detection on a copper electrode 436(1988)334
- Štulík, K., see Mingjia, W. 439(1988)363
- Štulík, K., see Pacáková, V. 442(1988)147
- Sturm, E. and Junker, A.  
Determination of telenezepine in human serum by GC-MS 430(1988)43
- Stuting, H.H., see Krull, I.S. 442(1988)29
- Stüwe, E., see Brendel, E. 432(1988)358
- Su, M.-H., see Au, J.L.-S. 423(1987)308
- Su, Y., Palusinski, O.A. and Fife, P.C.  
Isotachopheresis: analysis and computation of the structure of the ionic species interface 405(1987)77
- Subbaramaiah, K. and Sharma, R.  
Immunoaffinity purification of  $\beta$ -amylase from mustard (*Sinapis alba* L.) cotyledons 408(1987)385
- Subrahmanyam, D., see Bhatia, S.C. 434(1988)288
- Subramanian, G., Phillips, M.W. and Cramer, S.M.  
Displacement chromatography of biomolecules 439(1988)341
- Suckow, R.F. and Cooper, T.B.  
Isolation and identification of the glucuronide conjugate of 2-hydroxydesipramine by preparative LC 427(1988)287
- Sudo, K., see Fujimaki, Y. 433(1988)235
- Suehiro, T., see Matsunaga, H. 422(1987)353
- Suenaga, K., see Ikebuchi, J. 432(1988)401
- Sufredo, D., see Betts, T.J. 436(1988)53
- Sugamata, K., see Ogawa, K. 426(1988)188
- Sugeno, K., see Kominami, G. 417(1987)216
- Sugi, H., see Watanabe, Y. 425(1988)373
- Sugimura, K., Tsukahara, K., Ueda, Y., Takeda, K., Habu, Y. and Azuma, I.  
Fast protein LC of lymphocyte blastogenesis inhibitory factor produced by the human macrophage-like cell line U937 440(1988)131
- Sugita, S., Kobayashi, A., Suzuki, S., Yoshida, T. and Nakazawa, K.  
HPLC determination of imipramine and its metabolites in rat brain 421(1987)412
- Sugiura, M., see Shinbo, T. 405(1987)145
- Sugiyama, K. and Saito, M.  
Simple microscale supercritical fluid extraction system and its application to GC-MS of lemon peel oil 442(1988)121
- Sugiyama, N., see Kidouchi, K. 423(1987)297
- Sukkar, S.G., see Castello, G. 416(1987)119
- Sulkowski, E., see Andersson, L. 421(1987)141
- Sullivan, D.J., see Burrows, J.L. 432(1988)199
- Sullivan, N.W. and Weiner, L.  
Simple, rapid HPLC determination of bupivacaine in human plasma 417(1987)236
- Sullivan, W.R. and Fox, K.E.  
GC determination of the high-ceiling diuretic ethacrynic acid and its thiol adducts in plasma of guinea pigs 425(1988)396
- Sumida, Y., see Kataoka, H. 420(1987)135
- Sumii, I., see Ohmori, S. 426(1988)15
- Summers, B.R., see Burton, W.G. 443(1988)363
- Sun, G.Y., see Smith, R.E. 439(1988)83
- Sun, Y., see Wang, T. 407(1987)79

- Sun, Z.-P., see Yang, Q. 447(1988)208
- Sunada, K., see Harada, H. 407(1987)299
- Suprynowicz, Z., Buszewski, B., Lodkowski, R. and Dawidowicz, A.L.  
Comparison of mesoporous controlled-porosity glasses to silica gel as solid supports for C<sub>18</sub> chemically bonded phases 446(1988)347
- Suprynowicz, Z., see Buszewski, B. 446(1988)191
- Suryanarayana, M.V.S., see Ramachandran, P.K. 426(1988)239
- Suttnar, J., see Dyr, J.E. 408(1987)303
- Suzuki, E., see Takeuchi, T. 447(1988)221
- Suzuki, H., see Shibasaki, S. 421(1987)425
- Suzuki, K., John, I. and Kitazawa, S.  
HPLC determination of nitrazepam in plasma and its application to pharmacokinetic studies in the rat 425(1988)435
- Suzuki, M., see Harada, K.-I. 448(1988)275
- Suzuki, M., see Ikai, Y. 411(1987)313
- Suzuki, M., see Oka, H. 449(1988)448
- Suzuki, M., see Sano, A. 410(1987)427
- Suzuki, N., Saitoh, K. and Adachi, K.  
Reversed-phase HPLC and column LC of chlorophylls and their derivatives 408(1987)181
- Suzuki, N., see Noda, H. 435(1988)368
- Suzuki, N., see Saitoh, K. 439(1988)245
- Suzuki, S.-I., see Inoue, T. 422(1987)197
- Suzuki, S.-I., Tsuchihashi, H., Nakajima, K., Matsushita, A. and Nagao, T.  
Analyses of impurities in methamphetamine by inductively coupled plasma MS and ion chromatography 437(1988)322
- Suzuki, S., see Honda, S. 438(1988)73
- Suzuki, S., see Onodera, S. 409(1987)259
- Suzuki, S., see Sugita, S. 421(1987)412
- Suzuki, T., see Yamashita, J. 403(1987)275
- Suzuki, Y.  
Anion-exchange HPLC of water-soluble chromium(VI) and chromium(III) complexes in biological materials 415(1987)317
- Suzuki, Y., Hayashi, N. and Sekiba, K.  
Automated direct assay system for the measurement of sex steroid hormones in serum using HPLC 426(1988)33
- Suzuki, Y., see Matsukura, H. 414(1987)47
- Suzuki, Y., see Matsukura, H. 434(1988)196
- Suzuki, Y., see Tani, K. 404(1987)242
- Švec, F., see Tennikova, T.B. 435(1988)357
- Svensden, J.S., Sydnese, L.K. and Whist, J.E.  
Identification of 3-oxodicarboxylic acids in human urine 432(1988)13
- Svensson, C., Nyberg, G. and Mårtensson, E.  
HPLC quantitation of amitriptyline and nortriptyline in dialysate from plasma or serum using on-line solid-phase extraction 432(1988)363
- Svoboda, M., see Ostrovská, V. 446(1988)323
- Svoboda, M., see Svobodová, X. 425(1988)391
- Svoboda, V., see Kleinmann, I. 411(1987)335
- Svoboda, V., see Plicka, J. 442(1988)237
- Svoboda, V., see Vočková, J. 410(1987)500
- Svobodová, X., Kováčová, D., Ostrovská, V., Pechová, A., Poláček, O., Kusala, S. and Svoboda, M.  
Determination of nitroglycerin in human plasma using bonded-phase capillary column GC with electron-capture detection 425(1988)391
- Svobodová, X., see Ostrovská, V. 446(1988)323
- Swadesh, J.K., see Benedek, K. 444(1988)191
- Swanson, S.P., Nicoletti, J., Rood, Jr., H.D., Buck, W.B., Cote, L.M. and Yoshizawa, T.  
Metabolism of three trichothecene mycotoxins, T-2 toxin, diacetoxyscirpenol and deoxynivalenol, by bovine rumen microorganisms 414(1987)335
- Swart, A.C., see Swart, P. 442(1988)424
- Swart, K.J. and Botha, H.  
Rapid method for the determination of the diuretic triamterene and its metabolites in plasma and urine HPLC 413(1987)315
- Swart, P., Todres, P.C., Swart, A.C. and Van der Merwe, K.J.  
Micro-assay for sheep 11 $\beta$ -hydroxylase activity using HPLC for steroid analysis 442(1988)424
- Sweetman, L., see Hoffmann, G. 421(1987)336
- Svobodnik, W., Zhang, Y.Y., Klueppelberg, U., Janowitz, P., Wechsler, J.G., Fuerst, P. and Ditschuneit, H.  
Analysis of conjugated bile acids in human serum by HPLC using post-column enzyme reaction and off-line fluorimetric determination 423(1987)75
- Sworn, G., Marrs, W.M. and Hart, R.J.  
Characterisation of carrageenans by high-performance size-exclusion chromatography using a LiChrospher 1000 DIOL column 403(1987)307
- Sybilska, D., see Żukowski, J. 436(1988)381
- Sydnese, L.K., see Svendsen, J.S. 432(1988)13
- Syracuse, C.D., see Kuhnert, B.R. 426(1988)392
- Szabo, G., see Nagy, E. 425(1988)214
- Szabo, G.K., Davoudi, H. and Durso, R.  
HPLC method for measuring homovanillic acid in cerebrospinal fluid using electrochemical detection with internal standardization 430(1988)112

Szabo, G.K., see Mikati, M.A. 433(1988)305  
Szabó-Ravasz, B.

HPLC methods for the determination of  
contaminants in technical phosphomethylan  
435(1988)380

Szász, G., see Papp, O. 403(1987)11

Szász, G., see Papp, O. 403(1987)19

Szathmary, S.C. and Luhmann, I.

Sensitive and automated GC method for the  
determination of miconazol in plasma  
samples 425(1988)193

Szederkényi, F., Ambrus, G., Horváth, G., Ilkőy,  
É., Knausz, D., Csákvári, B., Ábermann, M.  
and Ujszászy, K.

N-Methoxy-N,O-bis(trimethylsilyl) carbamate  
as a derivatizing reagent for the GC of  
sitosterol degradation products 446(1988)253

Szejtli, J., see Cserhádi, T. 439(1988)393

Szejtli, J., see Żukowski, J. 436(1988)381

Szepesi, G., see Gazdag, M. 436(1988)31

Szepesi, G., see Gazdag, M. 450(1988)145

Szepesi, G., see Péter, A. 408(1987)43

Szewczyk, H., Dziwiński, E. and Król, P.

Analysis of 2,2-di-(*p*-  
ligooxyethylenophenyl)propane by GC and  
GC-MS 446(1988)109

Szewczyk, H., Dziwinski, E. and Szymanowski, J.

Analysis and identification of some  
aminoether alcohols and their ethers  
410(1987)447

Szógyi, M., see Cserhádi, T. 434(1988)455

Szókán, G., Mezö, G. and Hudecz, F.

Application of Marfey's reagent in  
racemization studies of amino acids and  
peptides 444(1988)115

Szymanowski, J., see Szewczyk, H. 410(1987)447

Szymanowski, J., see Voelkel, A. 409(1987)29

Szymanowski, J., see Voelkel, A. 448(1988)219

Szymanowski, J., Voelkel, A. and Rashid, Z.A.

Polarity and hydrophile-lipophile balance of  
model hydroxyoximes and their intermediates  
as determined by reversed-phase GC  
402(1987)55

Szymoniak, K., see Chrétien, J.R. 406(1987)95

Szymoniak, K. and Chrétien, J.R.

Factor analysis of trends in selectivity in  
GLC of 175 saturated and unsaturated esters  
on 14 polysiloxane stationary phases  
404(1987)11

## T

Tabet, J.-C., see Van, T.D. 440(1988)71

Tacconelli, F., see Opacka-Juffry, J. 433(1988)41

Tachdjian, C., see Porziemsky, J.-P. 440(1988)183

Tachizawa, H., see Fujimaki, Y. 433(1988)235

Tachizawa, H., see Matsubayashi, K.

433(1988)225

Tackett, R.L., see Arrendale, R.F. 432(1988)165

Taenaka, N., see Hosotsubo, H. 432(1988)340

Taft, R.W., see Abraham, M.H. 409(1987)15

Tagawa, T., see Yamaji, A. 421(1987)372

Taggart, C.T., see Fraser, A.J. 439(1988)404

Taha, A.M., see Marzouki, Z.M.H. 425(1988)11

Tahvanainen, J., see Meier, B. 442(1988)175

Takács, J.M., see Tekler, V. 406(1987)131

Takada, M., see Yamauchi, Y. 408(1987)171

Takada, Y., Hiwada, K. and Kokubu, T.

Purification of human renin by renin inhibitor  
(ES-305) affinity chromatography  
421(1987)350

Takagi, D., see Takeyama, N. 424(1988)361

Takagi, K., Okumura, K., Morikawa, N.,

Okuyama, S., Wu, M.-C. and Narita, O.

Simultaneous assay for individual sulphated  
 $3\alpha$ - and  $\beta$ -hydroxysteroids in serum using  
HPLC combined with  $3\alpha$ - and  $\beta$ -  
hydroxysteroid dehydrogenases immobilized  
on one column 432(1988)47

Takagi, T., see Chiu, W.-Y. 428(1988)25

Takagi, T., see Itabashi, Y. 402(1987)257

Takahara, S., see Hosotsubo, H. 432(1988)340

Takahashi, H., Ogata, H., Echizen, H. and

Ishizaki, T.

Determination of metoclopramide and its  
glucuronide and sulphate conjugates in  
human biological fluids (plasma, urine and  
bile) by ion-pair HPLC 419(1987)243

Takahashi, J., Kirino, O., Takayama, C. and

Kamoshita, K.

Studies on fungicidal activity of N-  
phenylcarbamates. IV. Determination of the  
hydrophobicity of N-phenylcarbamates by  
HPLC 436(1988)316

Takahashi, L.T., see Martin, C.F. 402(1987)376

Takahashi, N., see Putnam, F.W. 443(1988)267

Takahashi, R., see Semba, J.-I. 430(1988)118

Takahashi, S., see Matsubara, K. 424(1988)49

Takahashi, S., see Yamamoto, T. 413(1987)342

Takahashi, T., see Kojima, K. 425(1988)203

Takahashi, T., see Naoi, M. 426(1988)75

Takahashi, T., see Naoi, M. 427(1988)229

Takahashi, T., see Naoi, M. 434(1988)209

- Takahashi, T., see Sawada, S. 402(1987)283
- Takahasi, H., see Matsumoto, K. 425(1988)323
- Takai, N., see Tanaka, N. 448(1988)95
- Takai, N., see Yamashita, J. 403(1987)275
- Takano, S., see Kondoh, Y. 408(1987)255
- Takata, K., see Matsunaga, H. 422(1987)353
- Takayama, C., see Takahashi, J. 436(1988)316
- Takayama, M.  
Reversed-phase HPLC of urinary porphyrin free acids 423(1987)313
- Takeda, K., see Iwaki, K. 404(1987)117
- Takeda, K., see Iwaki, K. 407(1987)273
- Takeda, K., see Sugimura, K. 440(1988)131
- Takeda, T., see Yoshida, S. 431(1988)27
- Takeda, Y., see Matsuda, R. 438(1988)319
- Takegoshi, T., see Fujimaki, Y. 421(1987)367
- Takegoshi, T., see Tanaka, M. 426(1988)111
- Takehita, H., see Tohma, M. 421(1987)9
- Takeuchi, T., Asai, H. and Ishii, D.  
Enantiomeric separation of amino acids by micro HPLC on an L-proline-bonded stationary phase 407(1987)151
- Takeuchi, T., Hashimoto, Y. and Ishii, D.  
Temperature effect in packed microcolumn chromatography using a carbon dioxide mobile phase 402(1987)328
- Takeuchi, T., Murase, K. and Ishii, D.  
Determination of alcohols in alcoholic beverages by micro HPLC with indirect photometric detection 445(1988)139
- Takeuchi, T., Niwa, T. and Ishii, D.  
Stepwise gradient elution using switching valves in micro HPLC 405(1987)117
- Takeuchi, T., Niwa, T. and Ishii, D.  
Precolumn concentration and/or stepwise-gradient elution by switching valve techniques in micro HPLC 407(1987)141
- Takeuchi, T., Suzuki, E. and Ishii, D.  
Indirect photometric detection of inorganic anions in micro HPLC with permanently coated columns 447(1988)221
- Takeuchi, T., Watanabe, S., Kondo, N., Ishii, D. and Goto, M.  
Improvement of the interface for coupling of fast atom bombardment MS and micro HPLC 435(1988)482
- Takeuchi, T. and Ishii, D.  
Signal enhancement by on-column fluorometric detection in HPLC using micropacked fused-silica columns 435(1988)319
- Takeuchi, Y., see Sakai, T. 433(1988)73
- Takeyama, N., Takagi, D., Kitazawa, Y. and Tanaka, T.  
Determination of  $\alpha$ -keto acids in rat hindquarter muscle by HPLC with fluorescence detection 424(1988)361
- Takigawa, H., see Tanaka, M. 404(1987)175
- Takimoto, M., see Tamaki, M. 413(1987)287
- Takitani, S., see Sano, A. 410(1987)427
- Takuechi, T. and Ishii, D.  
Micro HPLC of poly(ethylene glycols) with indirect photometric detection 403(1987)324
- Talib, Y.Y., Karve, M.S., Bhide, S.V. and Kale, N.R.  
Separation of starch components based on ligand-induced adsorption of amylose on cellulose 403(1987)373
- Talmage, J.M. and Biemer, T.A.  
Determination of potassium nitrate and sodium monofluorophosphate in the presence of phosphate and sulfate by high-resolution ion chromatography 410(1987)494
- Tam, Y.K., see Coutts, R.T. 421(1987)267
- Tam, Y.K., see Pereira, C.M. 425(1988)208
- Tam, Y.K., Tawfik, S.R., Ke, J., Coutts, R.T., Gray, M.R. and Wyse, D.G.  
HPLC of lidocaine and nine of its metabolites in human plasma and urine 423(1987)199
- Tamai, G., Yoshida, H. and Imai, H.  
HPLC drug analysis by direct injection of whole blood samples. I. Determination of moderately hydrophobic drugs incorporated into blood corpuscles 423(1987)147
- Tamai, G., Yoshida, H. and Imai, H.  
HPLC drug analysis by direct injection of whole blood samples. II. Determination of hydrophilic drugs 423(1987)155
- Tamai, G., Yoshida, H. and Imai, H.  
HPLC drug analysis by direct injection of whole blood samples. III. Determination of hydrophobic drugs adsorbed on blood cell membranes 423(1987)163
- Tamaki, M., Takimoto, M., Nozaki, S. and Muramatsu, I.  
Adsorption of cyclic peptides analogous to gramicidin S and graptisin onto octadecylsilica stationary phase and bacterial cells 413(1987)287
- Tambute, A., see Lienne, M. 448(1988)55
- Tambute, A., see Macaudiere, P. 405(1987)135
- Tambuté, A., see Macaudière, P. 450(1988)255
- Tamura, Y., see Yasukawa, K. 410(1987)129
- Tan, H.S.I. and Pan, P.W.  
HPLC separation of noreximide and norendimide in bulk material, tablets and animal feed 404(1987)233
- Tan, L.K. and Dutrizac, J.E.  
Simultaneous determination of arsenic(III)

- and arsenic(V) in metallurgical processing media by ion chromatography 405(1987)247
- Tan, S., Tatsuno, T. and Okada, T.  
GC determination of calcium stearate in polyethylene food packaging sheets 447(1988)198
- Tan, Y.K. and Soldin, S.J.  
Analysis of salbutamol enantiomers in human urine by chiral HPLC and preliminary studies related to the stereoselective disposition kinetics in man 422(1987)187
- Tanabe, S., Kitahara, M., Nawata, M. and Kawanabe, K.  
Determination of oxidizable inorganic anions by HPLC with fluorescence detection and application to the determination of salivary nitrite and thiocyanate and serum thiocyanate 424(1988)29
- Tanaka, A., see Ohta, M. 402(1987)392
- Tanaka, E. and Misawa, S.  
Simultaneous determination of serum trimethadione and its metabolite by GC 413(1987)376
- Tanaka, H., see Nakagawa, T. 420(1987)297
- Tanaka, K., see Hirayama, C. 450(1988)271
- Tanaka, K. and Fritz, J.S.  
Ion-exclusion chromatography of non-ionic substances with conductivity detection 409(1987)271
- Tanaka, M., Ono, K. and Takegoshi, T.  
Determination of the thromboxane synthetase inhibitor 6-(1-imidazolylmethyl)-5,6,7,8-tetrahydronaphthalene-2-carboxylic acid (DP-1904) in human plasma and urine using solid-phase extraction and HPLC 426(1988)111
- Tanaka, M., Takigawa, H., Yasaka, Y., Shono, T., Funazo, K. and Wu, H.-L.  
Derivatization of inorganic anions with pentafluorobenzyl methanesulphonate for GC 404(1987)175
- Tanaka, M., Yasaka, Y., Kamino, M., Shono, T., Funazo, K. and Wu, H.-L.  
Polymeric pentafluorobenzylating reagent for GC of inorganic anions with electron-capture detection 438(1988)253
- Tanaka, N., Hashizume, K., Araki, M., Tsuchiya, H., Okuno, A., Iwaguchi, K., Ohnishi, S. and Takai, N.  
Internal structures of wide-pore packing materials for HPLC studied by transmission electron microscopy 448(1988)95
- Tanaka, N., see Barnhart, E.R. 445(1988)145
- Tanaka, N., see Jinno, K. 402(1987)173
- Tanaka, N., see Jinno, K. 436(1988)1
- Tanaka, N., Yoshimura, T. and Araki, M.  
Effect of pressure on solute retention in the ionization control mode in reversed-phase LC 406(1987)247
- Tanaka, O., see Kasai, R. 407(1987)205
- Tanaka, S., Kaneta, T. and Yoshida, H.  
Capillary tube isotachophoretic separation of heavy metal ions using complex-forming equilibria between cyanide as terminating ion and the metal ions 447(1988)383
- Tanaka, S., see Kuwata, K. 442(1988)407
- Tanaka, T., see Takeyama, N. 424(1988)361
- Tanaka, Y., Sato, H., Miyazaki, K. and Yamada, Y.  
Synthesis and characterization of stearyl methacrylate gel for reversed-phase LC 407(1987)197
- Tanaka, Y., see Sato, H. 447(1988)387
- Tang, B.K., see Nicoll-Griffith, D.A. 428(1988)103
- Tang, M., see Lau, C.E. 416(1987)212
- Tang, M., see Lau, C.E. 423(1987)251
- Tang, Y.-Z. and Aue, W.A.  
Spectra and response linearization of chlorine and bromine compounds in a copper-sensitized flame photometric detector 408(1987)69
- Tang, Y.-Z. and Aue, W.A.  
General photometric detector for GC 409(1987)125
- Tang, Y.-Z. and Aue, W.A.  
GC determination of benzaldehyde by gas-phase luminescence 409(1987)243
- Tani, K. and Suzuki, Y.  
Retention behaviour of aromatic aldehyde derivatives in reversed-phase LC 404(1987)242
- Tanii, Y., see Kurata, K. 434(1988)278
- Taniuchi, K., see Kobayashi, K. 417(1987)439
- Tar, A., see Gál, S. 403(1987)217
- Tar, A., see Kiss, L. 448(1988)109
- Tarbet, B.J., see Bradshaw, J.S. 405(1987)169
- Tarding, F., see Lam, H.R. 426(1988)358
- Tarján, G., see Tekler, V. 406(1987)131
- Tarter, J.G., see Galindo, I.C. 445(1988)211
- Tarter, J.G., see Gan, D.-C. 404(1987)285
- Tarter, J.G., see Jen, J.-F. 436(1988)86
- Tarumoto, T., see Dong, M.W. 442(1988)81
- Tarutani, T., see Yokoyama, T. 403(1987)151
- Tassaneevakul, W., see Lambert, W.E. 427(1988)195
- Tatematsu, A., see Niwa, N.T.T. 431(1988)418
- Tatsuno, T., see Tan, S. 447(1988)198
- Taub, R., see Chiu, S.-H.L. 433(1988)217



- Taube, D. and Neuhoff, V.  
Quantitative and automatic digital evaluation of one- and two-dimensional thin-layer chromatograms using vertical surface absorption at an optimum wavelength 437(1988)411
- Tavolato, B., see Gallo, P. 416(1987)53
- Tavss, E.A., Santalucia, J., Robinson, R.S. and Carroll, D.L.  
Analysis of flavor absorption into plastic packaging materials using multiple headspace extraction GC 438(1988)281
- Tavss, E.A., Wiet, S.G., Robinson, R.S., Santalucia, J. and Carroll, D.L.  
Flavor characterization of dentifrices using kinetic headspace GC 438(1988)273
- Tawa, R., Koshide, K., Hirose, S. and Fujimoto, T.  
Pre-column derivatization of sisomicin with *o*-phthalaldehyde- $\beta$ -mercaptopyronic acid and its application to sensitive HPLC determination with fluorimetric detection 425(1988)143
- Tawfik, S.R., see Tam, Y.K. 423(1987)199
- Tay, Y.H., see Lee, H.K. 438(1988)429
- Tayar, N. El, Tsantili-Kakoulidou, A., Roethlisberger, T., Testa, B. and Gal, J.  
Different partitioning behaviour of sulphonyl-containing compounds in reversed-phase HPLC and octanol-water systems 439(1988)237
- Taylor, D.R., see Akporhonor, E.E. 405(1987)67
- Taylor, E.H., Kennedy, E.E. and Pappas, A.A.  
Determination of flecainide by GC-MS 416(1987)365
- Taylor, G.W., see Richmond, R. 417(1987)241
- Taylor, H.J., see Lauren, D.R. 439(1988)470
- Taylor, L.H., see Kyerematen, G.A. 419(1987)191
- Taylor, R.B., Moody, R.R. and Ochekepe, N.A.  
Determination of proguanil and its metabolites cycloguanil and 4-chlorophenylbiguanide in plasma, whole blood and urine by HPLC 416(1987)394
- Taylor, W.H., see Newton, C.J. 417(1987)391
- Tedesco, J.L. and Schafer, R.  
Increased yield of the phenylthiocarbamyl-amino acids tyrosine, cysteine and phenylalanine using diethylether-based liquid-liquid extraction 403(1987)299
- Teeuwssen, J., see Vendrig, D.E.M.M. 424(1988)83
- Teeuwssen, J., see Vendrig, D.E.M.M. 434(1988)145
- Teichert-Kuliszewska, K., see Kruck, T.P.A. 433(1988)207
- Teitz, D., see Jemal, M. 428(1988)81
- Tekler, V., Tarján, G. and Takács, J.M.  
Effect of primary and secondary atomic environments on the interactions of some alkane C-H bonds on a squalane stationary phase at different column temperatures in GLC 406(1987)131
- Te Koppele, J.M., Van der Mark, E.J. and Mulder, G.J.  
LC determination of diastereomeric glutathione conjugates and further derivatives of  $\alpha$ -bromoisovalerylurea in rat bile and urine by electrochemically generated bromine 427(1988)67
- Teller, D.C., see Wooten, A.L. 445(1988)371
- Tena, G., see Lora-Tamayo, C. 422(1987)267
- Tena, T., see Lora-Tamayo, C. 422(1987)267
- Tenbarger, J.B., see Miller, B.J. 426(1988)438
- Tencheva, J., Yamboliev, I. and Zhivkova, Z.  
Reversed-phase LC for the determination of galanthamine and its metabolites in human plasma and urine 421(1987)396
- Tenczer, J., see Köppel, C. 427(1988)144
- Tenczer, J., see Köppel, C. 431(1988)197
- Tennikova, T.B., Horák, D., Švec, F., Kolář, J., Čoupek, J., Trushin, S.A., Maltzev, V.G. and Belenkii, B.G.  
Hydrolysed macroporous glycidyl methacrylate-ethylene dimethacrylate copolymer sorbent for size-exclusion HPLC of synthetic polymers and biopolymers 435(1988)357
- Teoh, R., see Woo, J. 420(1987)73
- Terada, H., see Miyake, K. 439(1988)227
- Terada, T., Sakata, C., Ishibashi, K., Tsuchiya, T. and Noguchi, H.  
Determination of the anti-anginal drug 6-chloro-2-pyridylmethyl nitrate in plasma and urine by fused-silica open-tubular column GC with electron-capture detection 430(1988)388
- Terlinden, R., Feige, M. and Römer, A.  
Determination of the two major metabolites of ebselen in human plasma by HPLC 430(1988)438
- Tesařová, E., see Macek, J. 434(1988)423
- Tešić, Ž.L., see Janjić, T.J. 404(1987)307
- Testa, B., see El Tayar, N. 439(1988)237
- Tétényi, P., see Kéry, Á. 446(1988)157
- Tew, K., see LaCreta, F.P. 434(1988)83
- Thakran, P., see Raha, C. 450(1988)325
- Thakur, S., see Ghosh, P. 403(1987)285
- Thalheimer, P., see Meese, C.O. 423(1987)344
- Tharasse-Bloch, C., Chabenat, C., Boucly, P. and Marchand, J.  
Determination of ambenonium chloride in

- serum by reversed-phase ion-pair LC  
421(1987)407
- 't Hart, B.J. and Wilting, J.  
Sensitive GC method for determining  
nitrazepam in serum and saliva 424(1988)403
- Thedford, R., see Alick, B. 430(1988)309
- Theis, D.L.  
Ion-pairing LC method for the determination  
of pirlimycin hydrochloride 402(1987)335
- Thénot, J.P., see Dubruc, C. 417(1987)208
- Thénot, J.P., see Ganansia, J. 421(1987)83
- Theoharides, A.D., see Zhou, Z.M. 414(1987)77
- Therasse, J. and Lemonnier, F.  
Determination of plasma lipoperoxides by  
HPLC 413(1987)237
- Thevenon, G., see Fausnaugh-Pollitt, J.  
443(1988)221
- Thibeault, Y., see Campeau, D. 405(1987)305
- Thiel, P.G., see Gathercole, P.S. 408(1987)435
- Thijssen, H.H.W., see Verhagen, H. 413(1987)282
- Thijssen, H.H.W., see Verhagen, H. 422(1987)288
- Thio, J.M.S.L., see Paanakker, J.E. 421(1987)327
- Thoma, R.S., see Crimmins, D.L. 443(1988)63
- Thomas, A.C., Kotzé, J.M. and Matthee, F.N.  
Characterization of *Botrytis* by GLC  
403(1987)292
- Thomas, A.H. and Newland, P.  
Chromatographic methods for the analysis of  
vancomycin 410(1987)373
- Thomas, D., see Achari, R. 424(1988)430
- Thomas, J.  
Relationships between retention behaviour  
and structure of N-substituted 1,2,4-triazoles  
in TLC on silica gel 404(1987)73
- Thomas, K.A., see Gimenez-Gallego, G.  
409(1987)299
- Thomas, L.C., see Ramus, T.L. 404(1987)155
- Thomas, L.C., see Van Engelen, D.L.  
405(1987)191
- Thompson, G.G., see Forrest, G. 428(1988)123
- Thompson, J.C., see Draviam, E.J. 414(1987)180
- Thompson, Jr., H.C., see Holder, C.L.  
419(1987)113
- Thompson, J.R., see Early, R.J. 416(1987)15
- Thompson, L.K. and Cone, E.J.  
Determination of  $\Delta^9$ -tetrahydrocannabinol in  
human blood and saliva by HPLC with  
amperometric detection 421(1987)91
- Thompson, M., see Sallustio, B.C. 422(1987)33
- Thompson, P.A., see Barlow, R.D. 419(1987)375
- Thompson, R.D. and Hoffmann, T.J.  
Determination of coumarin as an adulterant  
in vanilla flavoring products by HPLC  
438(1988)369
- Thomsen, M., see Hansen, S.H. 409(1987)71
- Thomsen, M., see Hansen, S.H. 447(1988)182
- Thongnopnua, P. and Zimmerman, C.L.  
Simultaneous microassay for etretinate and  
its active metabolite, etretin, by reversed-  
phase HPLC 433(1988)345
- Thormann, W., see Firestone, M.A. 407(1987)363
- Thormann, W., see Firestone, M.A. 436(1988)309
- Thormann, W., see Mosher, R.A. 436(1988)191
- Thornes, R.D., see Moran, E. 416(1987)165
- Thornton, D., see Chen, T.K. 450(1988)403
- Thornton, T.J., see Hsieh, J.Y.-K. 430(1988)175
- Thrane, U., see Frisvad, J.C. 404(1987)195
- Thuaud, N., see Sébille, B. 409(1987)61
- Thye, F.W., see McChrisley, B. 428(1988)35
- Tian, Z.W., Hu, R.Z., Lin, H.S. and Hu, W.L.  
Effect of suppression efficiency on sensitivity  
in ion chromatography 439(1988)151
- Tian, Z.W., Hu, R.Z., Lin, H.S. and Wu, J.T.  
High-performance electrochemical suppressor  
for ion chromatography 439(1988)159
- Tibbels, T.S., Smith, R.A. and Cohen, S.M.  
Determination of saccharin in diet and  
biological materials 441(1988)448
- Tice, P.A., Mazsaroff, I., Lin, N.T. and Regnier,  
F.E.  
Effects of large sample loads on column  
lifetime in preparative-scale LC 410(1987)43
- Tice, P.A., see Mazsaroff, I. 437(1988)429
- Tichá, M., see Šecová, E. 436(1988)59
- Tieckelmann, H., see Ho, J. 417(1987)269
- Tierney, E.J., Bellama, J.M., Eng, G., Brinckman,  
F.E. and Johannesen, R.B.  
Total molecular surface areas as a predictor  
for reversed-phase HPLC in various organotin  
systems. Importance and applications of  
carbon hybridization summaries  
441(1988)229
- Tierney, E.J., see Eng, G. 403(1987)1
- Tietz, D.  
Gel electrophoresis of intact subcellular  
particles (Review) 418(1987)305
- Tiffany, J.M., see Harvey, D.J. 414(1987)253
- Tikhonov, Y.V., see Toguzov, R.T. 434(1988)447
- Tikhonov, Y.V., see Werner, A. 421(1987)257
- Tillement, J.-P., see Duche, J.-C. 423(1987)340
- Tillery, K.F., see Rose, L.M. 425(1988)419
- Tillier, C., Cagniant, E. and Devaux, P.  
Determination of halofuginone in poultry  
feeds by HPLC 441(1988)406
- Timoceanu, V., see Ciucanu, I. 436(1988)219
- Tizzani, M.I., see Amicosante, G. 403(1987)366
- Tjaden, U.R., De Bruijn, E.A., Van der Hoeven,  
R.A.M., Jol, C., Van der Greef, J. and  
Lingeman, H.

- Automated analysis of mitomycin C in body fluids by HPLC with on-line sample pre-treatment 420(1987)53
- Tjaden, U.R., see De Jong, J. 414(1987)285
- Tjaden, U.R., see De Jong, J. 419(1987)85
- Tjaden, U.R., see Reeuwijk, H.J.E.M. 428(1988)93
- 't Klooster, G.A.E.V., Boot, J.H., Mennes, W.C. and Blaauboer, B.J.  
Rapid method for the determination and quantification of bromosulphophtalein and metabolites in cultured hepatocytes, culture media and bile by HPLC 432(1988)223
- Todo, S., see Tsunamoto, K. 417(1987)414
- Todorović, M., Kopečni, M.M., Čomor, J.J. and Laub, R.J.  
Reversibility of solvent-induced surface modification of alumina adsorbents 442(1988)105
- Todres, P.C., see Swart, P. 442(1988)424
- Tofanetti, O., see Castoldi, D. 425(1988)413
- Togari, A., Sakai, J., Matsumoto, S. and Nagatsu, T.  
Highly sensitive assay for alkaline and acid phosphatase activities by HPLC with electrochemical detection 417(1987)41
- Toguzov, R.T., see Werner, A. 421(1987)257
- Toguzov, R.T., Tikhonov, Y.V., Pimenov, A.M., Prokudin, V.Y., Dubiel, W., Ziegler, M. and Gerber, G.  
Optimization of the ion-pair HPLC separation of purine derivatives in erythrocytes, thymocytes and liver mitochondria 434(1988)447
- Tohma, M., Takeshita, H., Mahara, R., Kurosawa, T. and Makino, I.  
Determination of  $3\beta,12\alpha$ -dihydroxy-5- $\alpha$ -cholestan-24-oic acid and related bile acids in human serum by GC-MS 421(1987)9
- Toki, S., see Kumagai, Y. 421(1987)155
- Tokuma, Y., Fujiwara, T., Sekiguchi, M. and Noguchi, H.  
Determination of nilvadipine in plasma and urine by capillary column GC with electron-capture detection 415(1987)156
- Tokunaga, K., see Kato, Y. 404(1987)333
- Tollari, S., see Bassoli, A. 444(1988)209
- Tollari, S., see Bassoli, A. 446(1988)179
- Tomašić, J., Sesartić, L., Martin, S.A., Valinger, Z. and Ladešić, B.  
Comparative susceptibility of a peptidoglycan monomer from *Brevibacterium divaricatum* and its anhydromuramyl analogue to hydrolysis with N-acetylmuramyl-L-alanine amidase. Isolation and characterization of anhydromuramyl-peptidoglycan monomer 440(1988)405
- Tomer, K.B., see Parker, C.E. 438(1988)359
- Tomida, H., see Tsuruta, Y. 416(1987)63
- Tomita, M., Ijiri, I., Shimosato, K. and Kawai, S.  
Simple and sensitive method for the determination of acetaldehyde in blood by GC 414(1987)454
- Tonani, C., see Righetti, P.G. 440(1988)367
- Tonelli, D., see Gagliardi, L. 404(1987)267
- Tonelli, D., see Gagliardi, L. 408(1987)409
- Tonelli, D., see Gagliardi, L. 448(1988)296
- Tong, Y.-Y., see Sagara, K. 409(1987)365
- Tonković, M., see Hadžija, O. 402(1987)358
- Tonomura, H., see Iwata, H. 450(1988)317
- Toothaker, R.D., see Ferry, J.J. 421(1987)187
- Topalova, I., see Engewald, W. 446(1988)71
- Topalova, I., see Gavrilova, T.B. 402(1987)87
- Torchin, C.D., Kapetanovic, I.M., Yonekawa, W.D. and Kupferberg, H.J.  
HPLC assay of flunarizine, (*E*)-1-[bis(4-fluorophenyl)methyl]-4-(3-phenyl-2-propenyl)-piperazine, in plasma of epileptic patients 426(1988)444
- Torok-Both, G.A., see Coutts, R.T. 421(1987)267
- Torre, H.V. Der, see Veenstra, J. 431(1988)413
- Torres, L., see Riba, M.-L. 437(1988)139
- Tóth, G.  
Adsorption chromatographic behaviour of  $^{125}$ I-labelled progesterone-11 $\alpha$ - and -12 $\alpha$ -succinyltyrosine methyl ester 404(1987)258
- Tóth, G., see Mucha, I. 438(1988)111
- Toth, J.P. and Bardalaye, P.C.  
Capillary GC separation and MS detection of cyromazine and its metabolite melamine 408(1987)335
- Toth-Martinez, B.L., see Gál, S. 403(1987)217
- Toth-Martinez, B.L., see Kiss, L. 448(1988)109
- Totsuka, Y., see Nakano, K. 438(1988)61
- Touchstone, J.C., see Alvarez, J.G. 436(1988)515
- Touchstone, J.C. and Alvarez, J.G.  
Phospholipids in amniotic fluid with special reference to the lecithin; Sphingomyelin ratio (Review) 429(1988)359
- Tourwé, D., see De Bondt, M. 442(1988)165
- Tous, G., Fausnaugh, J., Vieira, P. and Stein, S.  
Preparation and chromatographic use of 5'-fluorescent-labelled DNA probes 444(1988)67
- Town, C., see Oldfield, N. 422(1987)135
- Townsend, R.W., Keuth, V., Embil, K., Mullersman, G., Perrin, J.H. and Derendorf, H.  
HPLC determination of conjugated estrogens in tablets 450(1988)414

- Toyoda, K., see Nakamura, K. 445(1988)234  
 Toyonaga, Y., see Ohmori, S. 426(1988)15  
 Toyo'oka, T., see Kotaniguchi, H. 420(1987)141  
 Tracqui, A., Kintz, P., Mangin, P. and Lugnier, A.A.  
 Simultaneous determination of floctafenine and its main metabolites, floctafenic and hydroxyfloctafenic acid, using high-performance liquid chromatography with gradient elution 432(1988)412  
 Tran-Phan, H., see Huang, S.-M. 423(1987)358  
 Treiman, D.M., see Gunawan, S. 414(1987)389  
 Tremp, J., see Czuczwa, J. 403(1987)233  
 Treutter, D.  
 Separation of naturally occurring mixtures of phenolic compounds from various Prunus tissues by reversed-phase HPLC 436(1988)490  
 Triana, K., see Labarre, P. 419(1987)381  
 Tribble, D.L., Glover, M.R. and Lambeth, J.D.  
 Pregnenolone production by adrenal mitochondria: a new HPLC analytical method for cholesterol side-chain cleavage 414(1987)411  
 Trigalo, F., see Maurs, M. 440(1988)209  
 Triggs, E.J., see Hakim, L. 424(1988)111  
 Triggs, E.J., see Ismail, Z. 404(1987)372  
 Trinca, M.L., see Chersi, A. 410(1987)463  
 Triolet, J., see Lecomte, L. 408(1987)416  
 Trfiska, J., see Jegorov, A. 434(1988)417  
 Trnka, T., see Jegorov, A. 434(1988)417  
 Trochimczuk, A., see Kolarz, B.N. 408(1987)308  
 Troskosky, J.A., see Raglione, T.V. 409(1987)205  
 Troskosky, J.A., see Raglione, T.V. 409(1987)213  
 Trowbridge, J., see Draviam, E.J. 414(1987)180  
 Truman, C.A., see Forsyth, D.R. 433(1988)352  
 Trushin, S.A., see Tennikova, T.B. 435(1988)357  
 Truskolaski, A., see Kroviarski, Y. 449(1988)403  
 Truskolaski, A., see Kroviarski, Y. 449(1988)413  
 Tsantili-Kakoulidou, A., see El Tayar, N. 439(1988)237  
 Tsantili-Kakoulidou, A. and Antoniadou-Vyza, A.  
 Hydrophobicity studies on diadamantanyl derivatives. Comparison between calculated partition coefficients and experimental lipophilicity indices determined by reversed-phase HPLC and TLC 445(1988)317  
 Tsegendis, T., see Karamanos, N.K. 405(1987)221  
 Tsiropoulos, N., see Riba, M.-L. 437(1988)139  
 Tsitsishvili, G.V., see Andronikashvili, T.G. 406(1987)145  
 Tsoucaris, G., see Coleman, A.W. 450(1988)175  
 Tsuboi, S., see Ohmori, S. 414(1987)149  
 Tsuchihashi, H., see Suzuki, S.-I. 437(1988)322  
 Tsuchiya, H., see Tanaka, N. 448(1988)95  
 Tsuchiya, K., see Goto, J. 425(1988)59  
 Tsuchiya, T., see Terada, T. 430(1988)388  
 Tsuchiya, T., see Ukai, S. 435(1988)374  
 Tsujioka, H., see Abe, I. 449(1988)165  
 Tsukahara, K., see Sugimura, K. 440(1988)131  
 Tsukahara, S., see Saitoh, K. 439(1988)245  
 Tsukamoto, K., see Iwamura, S. 413(1987)370  
 Tsukamoto, T., see Oi, T. 450(1988)343  
 Tsumagari, N., see Fujimoto, Y. 402(1987)344  
 Tsunamoto, K., Todo, S. and Imashuku, S.  
 Separation of prostaglandins and thromboxane by two-dimensional TLC 417(1987)414  
 Tsunoda, K.-I., see Nomura, A. 448(1988)87  
 Tsuruta, Y., Tomida, H., Kohashi, K. and Ohkura, Y.  
 Simultaneous determination of imidazoleacetic acid and N<sup>r</sup>- and N<sup>w</sup>-methylimidazoleacetic acids in human urine by HPLC with fluorescence detection 416(1987)63  
 Tsuyoshi, T., see Hirokawa, T. 408(1987)27  
 Tu, G., see Zeng, W. 408(1987)426  
 Tu, G., see Zhu, Y. 437(1988)265  
 Tu, G., see Zhu, Y. 438(1988)447  
 Tuan, G., see Beretta, E. 416(1987)144  
 Tucker, M.Y., see Baker, J.O. 437(1988)387  
 Turánek, J., Kovář, J. and Zábřřřová, H.  
 Rapid purification of the isoforms of rat liver Cd,Zn-metallothionein by HPLC 410(1987)470  
 Turánek, J., see Škabrahová, Z. 408(1987)372  
 Tureček, F., see Mráz, J. 414(1987)399  
 Turiák, G., see Kéry, Á. 446(1988)157  
 Turley, C.P. and Brewster, M.A.  
 LC analysis of cyclotrimethylenetrinitramine in biological fluids using solid-phase extraction 421(1987)430  
 Turner, C.A. and Murphy, R.  
 Rapid, high-sensitivity method for measurement of morphine in guinea-pig serum 428(1988)383  
 Turner, N.C., see Richmond, R. 417(1987)241  
 Tvrzická, E., Řezanka, T., Krijt, J. and Janoušek, V.  
 Identification of very-long-chain fatty acids in rat and mouse harderian gland lipids by capillary GC-MS 431(1988)231  
 Twardowski, A., see Muniak, W. 436(1988)323  
 Tyczkowska, K., see Zuk, M.M. 433(1988)367  
 Tyczkowska, K. and Aronson, A.L.  
 Analysis of cephalixin from canine skin biopsy by LC with UV-visible photodiode-

- array detection 427(1988)103
- Työppönen, J.T.  
Rapid and sensitive determination of Dns-amino acids in plasma using high-speed octadecyl LC columns 413(1987)25
- Tyrefors, N. and Gillberg, P.G.  
Determination of acetylcholine and choline in microdialysates from spinal cord of rat using LC with electrochemical detection 423(1987)85
- ## U
- Ubbink, J.B., Schnell, A.M. and Rapley, C.H.  
Quantification of urinary xanthurenic acid excretion by anion-exchange solid-phase extraction and HPLC 425(1988)182
- Ubbink, J.B. and Schnell, A.M.  
HPLC assay of erythrocyte enzyme activity levels involved in vitamin B<sub>6</sub> metabolism 431(1988)406
- Uccello-Barretta, G., see Salvadori, P. 450(1988)163
- Uchida, T., see Draviam, E.J. 414(1987)180
- Uchida, T., see Yasukawa, K. 410(1987)129
- Uchiyama, M., see Hayashi, Y. 407(1987)59
- Uchiyama, M., see Hayashi, Y. 411(1987)95
- Uchiyama, T., see Beck, R.H.F. 445(1988)282
- Uden, P.C., see Italia, M.P. 438(1988)35
- Uden, P.C., see Italia, M.P. 449(1988)326
- Uebis, V.  
Determination of a new thromboxane A<sub>2</sub> receptor blocker in biological fluids by capillary GC with electron-capture detection 419(1987)345
- Ueda, Y., see Sugimura, K. 440(1988)131
- Uehori, R., Nagata, T., Kimura, K., Kudo, K. and Noda, M.  
Screening of volatile compounds present in human blood using retention indices in GC 411(1987)251
- Ueno, M., see Ijitsu, T. 427(1988)29
- Uetz, J.A., see Holland, M.L. 433(1988)276
- Uezono, Y., see Kojima, K. 425(1988)203
- Ugalde, T.D., Faber, J.P.M. and Jenner, C.F.  
Optimizing copper-bicinchoninate carbohydrate detection for use with water-elution HPLC: a technique to measure the major mono- and oligosaccharides in small pieces of wheat endosperm 449(1988)207
- Uggeri, F., see Musu, C. 449(1988)432
- Ujszászy, K., see Szederkényi, F. 446(1988)253
- Ukai, S., Honda, A., Nagai, K., Kiho, T. and Tsuchiya, T.  
GLC of acid hydrolysates of hydroxyethylstarch and hydroxyethylcellulose 435(1988)374
- Ulm, E.H., see Los, L.E. 437(1988)442
- Umeda, M., Amagaya, S. and Ogihara, Y.  
Separation of thromboxane B<sub>2</sub> by HPLC 422(1987)246
- Umegae, Y., see Lee, M. 415(1987)289
- Umekawa, C., see Nakashima, K. 414(1987)11
- Umemoto, J., see Motohashi, N. 435(1988)335
- Umeno, Y., Matsushima, E., Maniwa, M. and Marunaka, T.  
GC-negative-ion chemical ionization MS determination of a new dihydropyridine calcium antagonist (MPC-1304) and its metabolites in human plasma and urine 434(1988)123
- Umeno, Y., see Maniwa, M. 433(1988)282
- Umeno, Y., see Marunaka, T. 420(1987)43
- Umezawa, A., see Iwase, H. 416(1987)37
- Umezawa, H., see Yamashita, K. 424(1988)39
- Umino, M., see Matsumoto, K. 425(1988)323
- Unadkat, J.D., Crosby, S.S., Wang, J.P. and Hertel, C.C.  
Simple and rapid HPLC assay for zidovudine (azidothymidine) in plasma and urine 430(1988)420
- Underwood, R.H., Potter, B.C. and Williams, G.H.  
Three-hour assay for urinary free cortisol: historical comparison of three methods each using a different chromatographic procedure 415(1987)118
- Unger, K.K., see Anspach, B. 443(1988)45
- Unsold, E., see Hyneck, M.L. 420(1987)349
- Uralets, V.P., see Rodchenkov, G.M. 423(1987)15
- Uralets, V.P., see Rodchenkov, G.M. 426(1988)399
- Uralets, V.P., see Rodchenkov, G.M. 432(1988)283
- Urbach, G.  
Dynamic headspace GC of volatile compounds in milk 404(1987)163
- Urbach, G.E., see Horwood, J.F. 437(1988)238
- Usero, J.L., Del Arco, A., Izquierdo, C., Montes, M. and Casado, J.  
Comparative study of HPLC with UV detection and GC with the Hall electrolytic conductivity detector in the analysis of N-nitroso compounds 407(1987)355

- Ushio, K., see Sakai, T. 433(1988)73  
 Utamura, T., see Koizumi, K. 409(1987)396  
 Utamura, T., see Koizumi, K. 436(1988)328  
 Utamura, T., see Koizumi, K. 437(1988)47  
 Utsumi, J., Kimura, S., Matsuda, S. and Shimizu, H.  
 Purification of murine macrophage colony-stimulating factor using hydroxyapatite HPLC in the presence of sodium dodecyl sulphate 421(1987)355
- V**
- Vaara, M.  
 Analytical and preparative HPLC of the papain-cleaved derivative of polymyxin B 441(1988)423  
 Vácha, J., see Macek, J. 420(1987)445  
 Vachta, J. and Siegfried, B.  
 HPLC determination of 1-(3-butoxy-2-carbamoyloxypropyl)-5-ethyl-5-phenyl-(1*H*,3*H*,5*H*)-pyrimidine-2,4,6-trione in human plasma 428(1988)183  
 Vader, H.L., see Gladdines, M.M. 431(1988)317  
 Vadon, C., see Kroviarski, Y. 449(1988)403  
 Vadon, C., see Kroviarski, Y. 449(1988)413  
 Vajda, N., see Nagy, L.G. 406(1987)359  
 Valášek, J., see Dufek, P. 435(1988)249  
 Valcárcel, M., see Lázaro, F. 448(1988)173  
 Valeiras-Price, M.C., see Blair, W.R. 410(1987)393  
 Valencia, R., see Cõng, H.N.G. 407(1987)349  
 Valente, A.L.P. and Collins, C.H.  
 GC quantification of bromine-containing compounds using a flame ionization detector 402(1987)349  
 Valentová, O., see Rauch, P. 438(1988)451  
 Valeur, A., see Odham, G. 434(1988)31  
 Valinger, Z., see Tomašić, J. 440(1988)405  
 Válka, I. and Šimánek, V.  
 Determination of alkaloids of *Fumaria parviflora* and *Fumaria capreolata* by HPLC and capillary isotachopheresis 445(1988)258  
 Valkenburg, C.F.M. Van, see Van der Krogt, J.A. 427(1988)9  
 Vallée, F., see Croteau, D. 419(1987)205  
 Van, T.D., Khac, D.D., Fetizon, M., Tabet, J.-C., Le Beyec, Y. and Quyen, T.C.  
 Identification of dekamycin antibiotics 440(1988)71  
 Van Alstine, J.M., see Karr, L.J. 442(1988)219  
 Van Beek, H. and Baars, A.J.  
 Isolation and quantitation of cadmium-, zinc- and copper-metallothioneins by HPLC atomic absorption spectrometry 442(1988)345  
 Van Beijsterveldt, L.E.C., see Wuis, E.W. 415(1987)419  
 Van Beijsterveldt, L.E.C., see Wuis, E.W. 420(1987)212  
 Van Belle, S.J.P., see De Smet, M. 416(1987)375  
 Van Bennekom, W.P., see Van Opstal, M.A.J. 432(1988)395  
 Van Binst, G., see De Bondt, M. 442(1988)165  
 Van Blitterswijk, H., see Van Ginkel, L.A. 445(1988)385  
 Van Brummelen, P., see Van Harten, J. 423(1987)327  
 Van den Berg, G.A., Kingma, A.W. and Muskiet, F.A.J.  
 Determination of polyamines in human erythrocytes by capillary GC with nitrogen-phosphorus detection 415(1987)27  
 Van den Berg, R.H., see Jansen, E.H.J.M. 437(1988)268  
 Van den Bossche, W., see De Schutter, J.A. 428(1988)301  
 VandenHeuvel, W.J.A., see Carlin, J.R. 427(1988)79  
 VandenHeuvel, W.J.A., see Metz, P.A. 441(1988)31  
 Van den Oetelaar, P.J.M. and Hoenders, H.J.  
 Electrophoretic titration curve in 6 *M* urea of the bovine eye lens protein  $\alpha$ -crystallin 411(1987)507  
 Van den Wildenberg, H.M., see Paanakker, J.E. 421(1987)327  
 Van der Auwera, L., see De Bondt, M. 442(1988)165  
 Vanderdeelen, J., see Der Meeren, P.V. 447(1988)436  
 Van der Greef, J., see Reeuwijk, H.J.E.M. 428(1988)93  
 Van der Greef, J., see Tjaden, U.R. 420(1987)53  
 Vanderhaeghe, H., see Cachet, T. 403(1987)343  
 Vanderhaeghe, H., see Cachet, T. 409(1987)91  
 Vanderhaeghe, H., see Khan, N.H. 405(1987)229  
 Vanderhaeghe, H., see Kibwage, I.O. 434(1988)177  
 Van der Hoeven, R.A.M., see Tjaden, U.R. 420(1987)53  
 Van der Hoorn, F.A.J., see Boomsma, F. 427(1988)219  
 Van der Horst, F.A.L. and Holthuis, J.J.M.

- Study of the derivatization of *n*-alkylamines with 1-fluoro-2,4-dinitrobenzene in the presence of aqueous cetyltrimethylammonium bromide micelles 426(1988)267
- Van der Kleyn, E., see Wuis, E.W. 415(1987)419
- Van der Kleyn, E., see Wuis, E.W. 420(1987)212
- Van der Krogt, J.A., Van Valkenburg, C.F.M. and Belfroid, R.D.M.  
Tritium isotope effect in reversed-phase HPLC analysis of dopamine and dihydroxyphenylacetic acid 427(1988)9
- Van der Maaten, M.J., see Schmerr, M.J.F. 425(1988)379
- Van der Maeden, F.P.B., see Broersen, J.J. 436(1988)39
- Van der Mark, E.J., see Te Koppele, J.M. 427(1988)67
- Van der Meer, M.J. and Brown, L.W.  
Simultaneous determination of amiloride and hydrochlorothiazide in plasma by reversed-phase HPLC 423(1987)351
- Van der Merwe, K.J., see Swart, P. 442(1988)424
- Van der Molen, J.C., see Koopman, B.J. 416(1987)1
- Vanderpas, J.B., see Koopman, B.J. 416(1987)1
- Van der Steege, G., Muskiet, F.A.J., Martini, I.A., Hutter, N.H. and Boersma, E.R.  
Simultaneous quantification of total medium- and long-chain fatty acids in human milk by capillary GC with split injection 415(1987)1
- Van der Steuijt, K. and Sonneveld, P.  
Concurrent analysis of methotrexate, trimethoprim, sulphamethoxazole and their major metabolites in plasma by HPLC 422(1987)328
- Van der Vijgh, W.J.F., see Maessen, P.A. 417(1987)339
- Van der Vijgh, W.J.F., see Maessen, P.A. 424(1988)103
- Van der Vusse, G.J., see Engels, W. 427(1988)209
- Van der Walt, L.A. and Wittliff, J.L.  
High-resolution separation of molybdate-stabilized progesterin receptors using HPLC 425(1988)277
- Van der Weken, G., see De Schutter, J.A. 428(1988)301
- Van der Wiele, K., see Verhaar, L.A.T. 437(1988)185
- Van der Zee, R., see Welling, G.W. 418(1987)223
- Van de Ven, L.J.M., see Claessens, H.A. 436(1988)345
- Van de Water, C. and Haagsma, N.  
Determination of chloramphenicol in swine muscle tissue using a monoclonal antibody-mediated clean-up procedure 411(1987)415
- Van Dijck, J., see Vindevogel, J. 447(1988)297
- Van Doormaal, J.J., see Hoving, E.B. 434(1988)395
- Van Engelen, D.L., Thomas, L.C. and Piepmeier, E.H.  
UV spectrometric detection for GC of polynuclear aromatic compounds with repetitive spectral scans using absorbance and concurrent fluorescence measurements 405(1987)191
- Van Gelder, W.M.J., Jonker, H.H., Huizing, H.J. and Scheffer, J.J.C.  
Capillary GC of steroidal alkaloids from Solanaceae. Retention indices and simultaneous flame ionization/nitrogen-specific detection 442(1988)133
- Van Gennip, A.H., see De Korte, D. 415(1987)383
- Vangiessen, G.J., see Kaiser, D.G. 434(1988)135
- Van Ginkel, L.A., Van Blitterswijk, H., Zoontjes, P.W., Den Bosch, D.V. and Stephany, R.W.  
Assay for trenbolone and its metabolite 17 $\alpha$ -Trenbolone in bovine urine based on immunoaffinity chromatographic clean-up and off-line HPLC-TLC 445(1988)385
- Van Haard, P.M.M. and Pavel, S.  
Chromatography of urinary indole derivatives (Review) 429(1988)59
- Van Hal, H.J.M., see Paanakker, J.E. 417(1987)203
- Van Harten, J., Lodewijks, M.T.M., Guyt-Scholten, J.W., Van Brummelen, P. and Breimer, D.D.  
GC determination of nisoldipine and one of its metabolites in plasma 423(1987)327
- Van Heiningen, A., see Derks, H.J.G.M. 442(1988)436
- Van Ingen, H.E., see Nielen, M.W.F. 417(1987)159
- Van Ingen, H.E. and Endert, E.  
Improved method for the determination of the cortisol production rate using HPLC and liquid scintillation counting 430(1988)233
- Van Kapel, J., see Lindemans, J. 432(1988)103
- Van Kouwen, A.M., see Rothuizen, J. 427(1988)19
- Van Lierop, J.B.H. and Van Veen, R.M.  
Determination of plasticizers in fat by GC-MS 447(1988)230
- Van Marsenille, M., see De Bondt, M. 442(1988)165
- Van Opstal, M.A.J., Krabbenborg, P., Holthuis, J.J.M., Van Bennekom, W.P. and Bult, A.  
Comparison of flow-injection analysis with

- HPLC for the determination of etoposide in plasma 432(1988)395
- Van Oss, G.E.C.J.M., see Vree, T.B. 424(1988)440
- Van Peteghem, C., see Rosier, J. 434(1988)222
- Van Rompaey, F., see Woestenborghs, R. 424(1988)195
- Van Schalkwijk, W.P., see Verkoelen, C.F. 426(1988)41
- Vanscoik, K.G., Johnson, C.A. and Porter, W.R. Sensitive and specific HPLC assay for 6-thiouric acid 417(1987)183
- Van Soest, R.E.J., see Nielsen, M.W.F. 417(1987)159
- Van Valkenburg, C.F.M., see Van der Krogt, J.A. 427(1988)9
- Van Veen, R.M., see Van Lierop, J.B.H. 447(1988)230
- Vařeka, J., see Kříž, J. 437(1988)177
- Varotsis, N. and Guizez, P. Microdistillation: a method combining GC and LC to characterize petroleum liquids 403(1987)159
- Vasil'eva, E.V., see Gavrilova, T.B. 446(1988)79
- Vasquez, I., see Paulos, C. 433(1988)359
- Vassal, G., Re, M. and Gouyette, A. GC-MS assay for busulfan in biological fluids using a deuterated internal standard 428(1988)357
- Vatassery, G.T., Herzan, L.A. and Dysken, M.W. Simultaneous determination of very low concentrations of haloperidol and reduced haloperidol in human serum by an LC method 433(1988)312
- Vecchio, G., see Candiano, G. 423(1987)319
- Veen, R.M. Van, see Van Lierop, J.B.H. 447(1988)230
- Veenstra, J., De Pol, H.V., Der Torre, H.V., Schaafsma, G. and Ockhuizen, T. Rapid and simple methods for the investigation of lipoxygenase pathways in human granulocytes 431(1988)413
- Veisserik, J., see Löhmus, M. 450(1988)105
- Vejrosta, J., Mikešová, M., Ansorgová, A. and Drozd, J. Sorption of benzene on Tenax 447(1988)170
- Vekshteyn, M., see Bartek, M.J. 421(1987)309
- Velayudhan, A., Lillig, B. and Horváth, C. Analysis of multiple front formation in the wetting of thin-layer plates 435(1988)397
- Velayudhan, A. and Horváth, C. Preparative chromatography of proteins. Analysis of the multivalent ion-exchange formalism 443(1988)13
- Veld, A.J. Man in 't, see Boomsma, F. 427(1988)219
- Velthorst, N.H., see Baumann, R.A. 439(1988)165
- Velury, S. and Howell, S.B. Measurement of plasma thiols after derivatization with monobromobimane 424(1988)141
- Ven, L.J.M. Van de, see Claessens, H.A. 436(1988)345
- Vendelbo Nielsen, H., see Gether, U. 447(1988)341
- Vendrig, D.E.M.M., Holthuis, J.J.M., Erdélyi-Tóth, V. and Hulshoff, A. Solid-phase extraction of vinblastine and vincristine from plasma and urine: variable drug recoveries due to non-reproducible column packings 414(1987)91
- Vendrig, D.E.M.M., Teeuwssen, J. and Holthuis, J.J.M. Analysis of vinca alkaloids in plasma and urine using HPLC with electrochemical detection 424(1988)83
- Vendrig, D.E.M.M., Teeuwssen, J. and Holthuis, J.J.M. Determination of vinca alkaloids in plasma and urine using ion-exchange chromatography on silica gel and fluorescence detection 434(1988)145
- Venema, J.W., see Lemmens, A.A.G. 439(1988)423
- Venkataram, S. and Rahman, Y.E. HPLC analysis of desferrioxamine using a metal-free system 411(1987)494
- Venkataramanan, R., see D'Souza, M.J. 421(1987)198
- Venkateswaran, K.S., see Ramachandran, P.K. 426(1988)239
- Venn, R.F. Combined HPLC-radioimmunoassay method for the analysis of endorphins, enkephalins and other neurotransmitter peptides 423(1987)93
- Vent, S. Le, see Akporhonor, E.E. 405(1987)67
- Ventura, P., see Corti, F. 424(1988)147
- Verbeek, J., see Roorda, W.E. 403(1987)355
- Verbruggen, A., see Kibwage, I.O. 434(1988)177
- Verdier, F., see Pussard, E. 421(1987)192
- Verdys, M., see Olivares, J. 434(1988)111
- Verebey, K. and DePace, A. Rapid confirmation of enzyme-multiplied immunoassay test (EMIT) for phencyclidine-positive urine samples with capillary gas chromatography-nitrogen-phosphorus detection 427(1988)151
- Verge, R. Le, see Le Corre, P. 424(1988)424
- Verge, R. Le, see Le Corre, P. 450(1988)211



- Vergnaud, J.P., see Guimbard, J.P. 403(1987)109
- Vergne, A., see Billard, V. 440(1988)87
- Verhaar, L.A.T., Meuldijk, J., Groenland, W.P.T. and Van der Wiele, K.  
HPLC of mixtures of phenol or guaiacol and their reaction products with glyoxylic acid 437(1988)185
- Verhagen, H., Thijssen, H.H.W. and Kleinjans, J.C.S.  
Sensitive HPLC method for the routine determination of butylated hydroxyanisole in plasma 413(1987)282
- Verhagen, H., Thijssen, H.H.W. and Kleinjans, J.C.S.  
Determination of butylated hydroxytoluene in plasma by HPLC 422(1987)288
- Verillon, F., see Franke, G. 450(1988)81
- Verkoelen, C.F., Romijn, J.C., Schroeder, F.H., Van Schalkwijk, W.P. and Splinter, T.A.W.  
Quantitation of polyamines in cultured cells and tissue homogenates by reversed-phase HPLC of their benzoyl derivatives 426(1988)41
- Vermeer, E., Knecht, E. and Bruinsma, J.  
Determination of abscisic acid in small amounts of plant material 404(1987)346
- Vermeulen, N.P.E., see Xu, B. 445(1988)1
- Verner, P., Pehal, F., Konečná, V., Zeman, J. and Hyánek, J.  
Quantification of some glycols in urinary organic acid profiles of acutely ill newborns 416(1987)439
- Verotta, L., see Ferrari, M. 437(1988)328
- Verweij, A., see Kientz, C.E. 407(1987)340
- Verzele, M., De Coninck, M., Vindevogel, J. and Dewaele, C.  
Column hardware in preparative LC with axial flow 450(1988)47
- Verzele, M., see Nevejans, F. 406(1987)325
- Verzele, M., see Schuddinck, G. 407(1987)159
- Verzele, M., see Vindevogel, J. 447(1988)297
- Veselá, Z., see Bystrický, Ľ. 447(1988)202
- Vesell, E.S., see Kyerematen, G.A. 419(1987)191
- Vespalec, R., see Neča, J. 447(1988)177
- Vessman, J., see Gyllenhaal, O. 411(1987)285
- Vessman, J., see Gyllenhaal, O. 435(1988)259
- Vetter, B., see Benecke, R. 417(1987)287
- Veuthey, J.L., Senn, J.P. and Haerdi, W.  
Low-cost rapid separation and determination of ionic species by single-column ion chromatography 445(1988)183
- Veyre, A., see Labarre, P. 419(1987)381
- Vézina, D. and Bleau, G.  
HPLC of selenium in biological samples 426(1988)385
- Vial, J., see Richmond, R. 417(1987)241
- Viala, A., see Rop, P.P. 427(1988)172
- Vian, L., see Larroque, M. 407(1987)384
- Vian, L., see Larroque, M. 445(1988)107
- Vicchio, D., Speedie, M.K. and Callery, P.S.  
GC-MS determination of tryptophan transaminase-catalyzed deuterium exchange 415(1987)104
- Viden, I., see Hajšlová, J. 438(1988)55
- Viden, I. and Řezanka, R.  
Capillary GC-MS of aliphatic saturated  $\alpha, \omega$ -dicarboxylic acid dimethyl esters and direct inlet MS of the corresponding free acids 408(1987)145
- Vidrich, V., see Heimler, D. 448(1988)301
- Vieira, P., see Tous, G. 444(1988)67
- Vigneron, C., see Menu, P. 417(1987)397
- Vijayaraghavan, R., see Ramachandran, P.K. 426(1988)239
- Vijgh, W.J.F. Van der, see Maessen, P.A. 417(1987)339
- Vijgh, W.J.F. Van der, see Maessen, P.A. 424(1988)103
- Villanueva, V.R. and Mardon, M.  
Chromatography, flow injection analysis and electrophoresis in computer-assisted comparative biochemistry: its application and possibilities in clinical research. Preliminary studies on Crohn's disease 440(1988)261
- Ville, G., see Ricard-Blum, S. 431(1988)474
- Villermaux, J.  
Chemical engineering approach to dynamic modelling of linear chromatography. A flexible method for representing complex phenomena from simple concepts 406(1987)11
- Vilsen, B. and Andersen, J.P.  
Use of a small TSK GSW HPLC column for large-zone chromatographic studies of monomer-oligomer equilibria of membrane protein 442(1988)229
- Vincken, B.J., see Pieters, F.A.J.M. 422(1987)322
- Vinçon, G., see Demotes-Mainard, F. 419(1987)388
- Vindevogel, J., see Verzele, M. 450(1988)47
- Vindevogel, J., Van Dijck, J. and Verzele, M.  
Micro-LC and the chiral recognition mechanism on albumin-coated silica gel. Large selectivity changes with sample size 447(1988)297
- Vine, J.H., see Dawson, M. 420(1987)129
- Vink, J., see Smilde, A.K. 410(1987)1
- Virgili, R., see Armani, E. 441(1988)275
- Viscomi, G., Lande, S. and Horváth, C.  
High-performance displacement

- chromatography of melanotropins and their derivatives 440(1988)157
- Visconti, M., Spalluto, R., Crolla, T., Pifferi, G. and Pinza, M.  
Determination of oxiracetam in human serum and urine by HPLC 416(1987)433
- Visentini, J., see D'Agostino, P.A. 402(1987)221
- Visser, E., see De Jong, J. 419(1987)85
- Visser, J., see Frijlink, H.W. 415(1987)325
- Vistelle, R., see Lamiable, D. 434(1988)315
- Viswanatha, T., see Mussar, K.J. 408(1987)378
- Vitale, G.C., see Baker, P.R. 423(1987)63
- Vleggaar, R., see De Jesus, A.E. 450(1988)101
- Vo-Dinh, T., see Spino, L.A. 409(1987)147
- Vobecký, M., see Dufek, P. 435(1988)249
- Vočková, J., see Kleinmann, I. 411(1987)335
- Vočková, J. and Svoboda, V.  
Fast reversed-phase HPLC determination of <sup>14</sup>C- or <sup>3</sup>H-labelled S-adenosylmethionine in reaction mixtures 410(1987)500
- Vodička, L., see Kříž, J. 437(1988)177
- Vodicska, M., see Nagy, L.G. 406(1987)359
- Voegel, J.C., Pefferkorn, E. and Schmitt, A.  
Steric protection of a modified glass surface with adsorbed polyacrylamide: interactions with blood proteins 428(1988)17
- Voelkel, A.  
Dispersive force parameters as a measure of the polarity of surfactants used as stationary phases in GLC 435(1988)29
- Voelkel, A.  
Polarity of oligooxyethylene derivatives of alcohols, thio alcohols and alkylamines as measured by GC 450(1988)291
- Voelkel, A., see Szymanowski, J. 402(1987)55
- Voelkel, A., Szymanowski, J., Beger, J. and Ebert, K.  
Polarity of 1,3-bis[ $\omega$ -alkoxyoligo(oxyethylene)]propan-2-ols as measured by reversed-phase GC 409(1987)29
- Voelkel, A., Szymanowski, J., Beger, J. and Rüstig, H.  
Polarity of some individual derivatives of  $\alpha, \omega$ -diaminoaligoethers as measured by GC 448(1988)219
- Volmer, M., Meiborg, G. and Muskiet, F.A.J.  
Simultaneous capillary GC profiling of medium- and long-chain fatty acid methyl esters with split injection. Correction for injection-related discrimination by the 'bracketing' method 434(1988)385
- Volmer, M., see Hoving, E.B. 434(1988)395
- Volpe, A.R., see Porcelli, G. 414(1987)423
- Von Angerer, E., see Birnböck, H. 414(1987)235
- Von Bredow, J., see Kapetanovic, I.M. 419(1987)458
- Vonderschmitt, D.J., see Minder, E.I. 419(1987)135
- Vonderschmitt, D.J., see Minder, E.I. 428(1988)369
- Vouldoukis, I., see Ogunkolade, B.W. 440(1988)459
- Vouros, P., see Wasilchuk, B.A. 407(1987)305
- Voutsinou-Taliadouri, F., see Satsmadjis, J. 437(1988)254
- Voyksner, R.D., see Blau, P.A. 420(1987)1
- Vrátný, P., see Mikeš, O. 440(1988)287
- Vrbanac, J.J., Eller, T.D. and Knapp, D.R.  
Quantitative analysis of 6-keto-prostaglandin F<sub>1 $\alpha$</sub>  using immunoaffinity purification and GC-MS 425(1988)1
- Vree, T.B., Baars, A.M., Van Oss, G.E.C.J.M. and Booij, L.H.D.J.  
HPLC and preliminary pharmacokinetics of articaïne and its 2-carboxy metabolite in human serum and urine 424(1988)440
- Vree, T.B., Baars, A.M. and De Grood, P.M.R.M.  
HPLC determination and preliminary pharmacokinetics of propofol and its metabolites in human plasma and urine 417(1987)458
- Vree, T.B., Riemens, L. and Koopman-Kimenai, P.M.  
HPLC of xanthines. Effects of N-methyl and C-8 hydroxyl substitution on the retention times 428(1988)311
- Vree, T.B., see Botterblom, M.H.A. 421(1987)211
- Vree, T.B., see Koopman-Kimenai, P.M. 416(1987)382
- Vree, T.B., see Wuis, E.W. 415(1987)419
- Vree, T.B., see Wuis, E.W. 420(1987)212
- Vreman, H.J., Stevenson, D.K., Henton, D. and Rosenthal, P.  
Correlation of carbon monoxide and bilirubin production by tissue homogenates 427(1988)315
- Vries, J.X. De, Walter-Sack, I. and Ittensohn, A.  
Analysis of benzbromarone in human plasma and urine by HPLC and GC-MS 417(1987)420
- Vučković, G.N., see Janjić, T.J. 404(1987)307
- Vusse, G.J. Van der, see Engels, W. 427(1988)209

# W

- Waagboe, R. and Sandnes, K.  
Determination of vitellogenin in serum of rainbow trout (*Salmo gairdneri*) by HPLC 427(1988)138
- Waal, W.A.J. De, Maessen, F.J.M.J. and Kraak, J.C.  
Analysis of platinum species originating from *cis*-diamminedichloroplatinum(II) (cisplatin) in human and rat plasma by HPLC with on-line inductively coupled plasma atomic emission spectrometric detection 407(1987)253
- Wada, H., see Yonekura, T. 427(1988)320
- Wada, Y., see Kidouchi, K. 423(1987)297
- Wade, D.N., see Joyce, D.A. 430(1988)319
- Wade, J.L. and Carr, P.W.  
Split peaks in non-linear chromatography and their effect on sample throughput in large scale separations 449(1988)53
- Wagner, J.G., see Ganes, D.A. 432(1988)233
- Wahrmann, J.P., see Rabilloud, T. 402(1987)105
- Wahyuni, W.T. and Jinno, K.  
Separation of tocopherols on various chemically bonded phases in microcolumn LC 448(1988)398
- Wainer, I.W., Stiffin, R.M. and Shibata, T.  
Resolution of enantiomeric aromatic alcohols on a cellulose tribenzoate HPLC chiral stationary phase. A proposed chiral recognition mechanism 411(1987)139
- Wainer, I.W. and Stiffin, R.M.  
Direct resolution of the stereoisomers of leucovorin and 5-methyltetrahydrofolate using a bovine serum albumin HPLC chiral stationary phase coupled to an achiral phenyl column 424(1988)158
- Wajcman, H., see Baudin-Chich, V. 437(1988)193
- Wakai, J., see Haginaka, J. 413(1987)219
- Wakai, J., see Haginaka, J. 447(1988)365
- Wakai, J., see Haginaka, J. 447(1988)373
- Wakasugi, K., see Hirano, T. 417(1987)371
- Wakayama, S., see Nakano, K. 438(1988)61
- Waki, H., see Ando, S. 405(1987)125
- Waki, H., see Ando, S. 408(1987)285
- Walberg, H.-J., see Weiss, J. 439(1988)91
- Waldmann-Meyer, H.  
Structure parameters of molecules and media evaluated by chromatographic partition. II. Geometrical exclusion in gels 410(1987)233
- Walgraeve, H.R., see Lambert, W.E. 427(1988)195
- Walhagen, A., see Edholm, L.-E. 424(1988)61
- Walker, B.J., Lang, J.F. and Okerholm, R.A.  
Quantitative analysis of dicyclomine in human plasma by capillary GC and nitrogen-selective detection 416(1987)150
- Wallace, C.A., see Smith, A.I. 416(1987)255
- Wallace, G.R., McLeod, A. and Chain, B.M.  
Chromatographic analysis of the trinitrophenyl derivatives of insulin 427(1988)239
- Wallace, L.C., see Carilli, C.T. 444(1988)203
- Wallace, M.R., see France, N.C. 433(1988)1
- Wallasch, T.M., Henning, K., Lange, U., Kuhn, W., Eckhardt-Wallasch, H. and Przuntek, H.  
Comparison of two methods for the determination of substance P immunoreactivity in human cerebrospinal fluid: sulphopropyl-Sephadex ion-exchange chromatography and/or reversed-phase HPLC combined with radiomimmunoassay 425(1988)175
- Walldorf, B., see Nilsson, L.-G. 423(1987)189
- Walle, T., see Wilson, M.J. 431(1988)222
- Wallemacq, P.E. and Lesne, M.  
New automated HPLC analysis of cyclosporin A and G in human serum 413(1987)131
- Wallingford, R.A. and Ewing, A.G.  
Retention of ionic and non-ionic catechols in capillary zone electrophoresis with micellar solutions 441(1988)299
- Walser, P.  
New ion-exclusion phase for organic acids 439(1988)71
- Walsh, T.D., see Stevens, Y.-W. 416(1987)203
- Walsh, T.D. and Fanucchi, M.P.  
Determination of hexamethylene bisacetamide by capillary GC 414(1987)510
- Walt, L.A. Van der and Wittliff, J.L.  
High-resolution separation of molybdate-stabilized progesterin receptors using HPLC 425(1988)277
- Walter, H. and Krob, E.J.  
Partitioning behavior of erythrocytes in aqueous two-phase systems containing hydroxypropyl starch and polyethylene glycol 441(1988)261
- Walterová, D. and Šimánek, V.  
Determination of  $\beta$ -phenylalkylamines in pharmaceutical preparations by analytical capillary isotachopheresis 405(1987)389
- Walters, D.G., see Bailey, E. 421(1987)21
- Walters, M.J., see Delaney, M.F. 410(1987)31
- Walters, R.R., see Hage, D.S. 436(1988)111
- Walter-Sack, I., see De Vries, J.X. 417(1987)420
- Walton, M.I. and Workman, P.

- HPLC assay for the benzotriazine di-N-oxide (SR 4233) and its reduced metabolites in biological materials 430(1988)429
- Wang, E., see Ji, H. 410(1987)111
- Wang, E. and Hou, W.  
Determination of water-soluble vitamins using HPLC and electrochemical or absorbance detection 447(1988)256
- Wang, J., see Lin, B. 438(1988)171
- Wang, J. and Bonakdar, M.  
Sensitive measurements of warfarin by HPLC with electrochemical detection 415(1987)432
- Wang, J.P., see Unadkat, J.D. 430(1988)420
- Wang, K.-T., see Chiou, S.-H. 448(1988)404
- Wang, T. and Sun, Y.  
Reproducibility of temperature-programmed retention indices on several OV-101 columns 407(1987)79
- Wang, Y.-M., see Anzai, T. 415(1987)445
- Wang, Z.-J., Zaitso, K. and Ohkura, Y.  
HPLC determination of  $\alpha$ -keto acids in human serum and urine using 1,2-diamino-4,5-methylenedioxybenzene as a precolumn fluorescence derivatization reagent 430(1988)223
- Wang-Iverson, D., Jemal, M. and Cohen, A.I.  
Determination of enantiomeric purity of 3-mercapto-2-methylpropionic acid 408(1987)365
- Wangler, W., see Rockstroh, H. 442(1988)401
- Wannerberg, O. and Persson, B.  
LC method for the determination of bambuterol hydrochloride and related compounds 435(1988)199
- Ward, A.E., see Bhamra, R.K. 417(1987)229
- Ward, J.R., see Rohrbaugh, D.K. 447(1988)165
- Ward, R. and Workman, P.  
Gradient HPLC method for simultaneous assay of the radiosensitizers etanidazole (SR 2508) and pimonidazole (Ro 03-8799) in biological materials 420(1987)223
- Ward, S.A., see Milton, K.A. 433(1988)339
- Ward, T.J., see Armstrong, R.D. 414(1987)192
- Warner, F.P., see Lloyd, L.L. 437(1988)447
- Wart, H.H.F.V. Der, see Schellens, J.H.M. 431(1988)203
- Wasa, M., see Yonekura, T. 427(1988)320
- Wasa, T., see Abe, I. 449(1988)165
- Wasilchuk, B.A. and Vouros, P.  
Design of an inexpensive on-column injector for capillary GC 407(1987)305
- Watanabe, A., see Semba, J.-I. 430(1988)118
- Watanabe, C., see Henzel, W.J. 404(1987)41
- Watanabe, K., see Yamashita, K. 424(1988)1
- Watanabe, K., see Yamashita, K. 424(1988)39
- Watanabe, M.F., see Harada, K.-I. 448(1988)275
- Watanabe, S., see Takeuchi, T. 435(1988)482
- Watanabe, S., see Watanabe, Y. 425(1988)373
- Watanabe, T., see Jinno, K. 436(1988)11
- Watanabe, T., see Yonekura, T. 427(1988)320
- Watanabe, Y., Sugi, H., Watanabe, S. and Mori, A.  
Identification of 1-guanidino-2-(4-imidazole)propionic acid in human urine by HPLC and field desorption MS 425(1988)373
- Watari, N., see Lee, F.W. 426(1988)259
- Water, C. Van de and Haagsma, N.  
Determination of chloramphenicol in swine muscle tissue using a monoclonal antibody-mediated clean-up procedure 411(1987)415
- Watson, D.G., see Brooks, C.J.W. 438(1988)108
- Watson, E. and Kenney, W.C.  
High-performance SEC of recombinant derived proteins and aggregated species 436(1988)289
- Watson, T.R., see Dawson, M. 420(1987)129
- Watt, P.W., see Corbett, M.E. 419(1987)263
- Waud, W.R. and Hill, D.L.  
HPLC determination of terephthalamidine in plasma 425(1988)220
- Wauters, J.-N., see Brasseur, T. 437(1988)260
- Waxman, L., see Kee, Y. 428(1988)345
- Weaner, L.E. and Hoerr, D.C.  
Separation of fatty acid ester and amide enantiomers by HPLC on chiral stationary phases 437(1988)109
- Webb, A.A., see Crockett, T.D. 407(1987)330
- Weber, B., see Scholl, H. 416(1987)321
- Weber, C., Belsner, K., Lohkamp, R. and Höller, M.  
Determination of prostaglandins  $D_2$ ,  $E_2$ ,  $F_{2\alpha}$  and 6-ketoprostaglandin  $F_{1\alpha}$  as well as thromboxane  $B_2$  in rat brain by GC-MS 426(1988)345
- Weber, D., see Lau, B.P.-Y. 402(1987)233
- Weber, D.J., see Kaiser, D.G. 434(1988)135
- Weber, D.J., see Shah, J.A. 414(1987)1
- Weber, D.V., Keeney, R.F., Familletti, P.C. and Bailon, P.  
Medium-scale ligand-affinity purification of two soluble forms of human interleukin-2 receptor 431(1988)55
- Weber, F., see Efthymiopoulos, C. 421(1987)360
- Weber, M.H.  
Urinary protein analysis (Review) 429(1988)315
- Webster, H.K., see Krungkrai, J. 417(1987)47
- Webster, R.A., see Farrant, M. 417(1987)385
- Wechsler, J.G., see Swobodnik, W. 423(1987)75
- Wechter, M.A., see Black, B.H. 437(1988)203

- Weckesser, J., see Birk, I.M. 449(1988)423
- Wedlund, P.J., see Huang Chandler, M.H. 419(1987)426
- Wedlund, P.J., see Robbins, D.K. 415(1987)208
- Wehling, R.L., see Holt, D.L. 449(1988)271
- Wehr, C.T.  
Sample preparation and column regeneration in biopolymer separations (Review) 418(1987)27
- Weidler, D.J., see Fouda, H.G. 417(1987)197
- Weidolf, L.O.G., Chichila, T.M.P. and Henion, J.D.  
Screening, confirmation and quantification of boldenone sulfate in equine urine after administration of boldenone undecylenate (Equipoise) 433(1988)9
- Weinberger, R., see Femia, R.A. 402(1987)127
- Weiner, L., see Sullivan, N.W. 417(1987)236
- Weinstein, L.H., see Wettlaufer, S.H. 441(1988)361
- Weiss, J., Möckel, H.J., Müller, A., Diemann, E. and Walberg, H.-J.  
Retention of thio- and selenometalates in mobile-phase ion chromatography 439(1988)91
- Weisweiler, P.  
Rapid quantification of apolipoprotein E enriched very-low-density lipoproteins by heparin-Sepharose chromatography 425(1988)169
- Weken, G. Van der, see De Schutter, J.A. 428(1988)301
- Welinder, B.S., Sørensen, H.H. and Hansen, B.  
Recovery of polypeptides after reversed-phase HPLC 408(1987)191
- Weller, O., Schulze, J. and König, W.A.  
Separation of racemic pharmaceuticals by HPLC on silica gel modified with carbohydrate residues 403(1987)263
- Welling, G.W., see Welling-Wester, S. 443(1988)255
- Welling, G.W., Van der Zee, R. and Welling-Wester, S.  
Column LC of integral membrane proteins (Review) 418(1987)223
- Welling-Wester, S., Kazemier, B., Örvell, C. and Welling, G.W.  
Effect of detergents on the structure of integral membrane proteins of Sendai virus studied with size-exclusion HPLC and monoclonal antibodies 443(1988)255
- Welling-Wester, S., see Welling, G.W. 418(1987)223
- Welscher, T.M., McQuinn, R.L. and Chang, S.F.  
HPLC procedure with fluorescence detection for the *m*-O-dealkylated lactam metabolite of flecainide acetate in human plasma 431(1988)438
- Wenkert, E., Marsaioli, A.J. and Moeller, P.D.R.  
Formal total synthesis of rosefuran and eldanolide 440(1988)449
- Wennberg, O., see Johansson, B.-L. 403(1987)85
- Wentworth, W.E., Limero, T., Batten, C.F. and Chen, E.C.M.  
Non-radioactive electron-capture detector operating in the pulsed mode 441(1988)45
- Wenz, G., see König, W.A. 447(1988)193
- Wenzel, T.J., Collette, L.M., Dahlen, D.T., Hendrickson, S.M. and Yarmaloff, L.W.  
LC and flow injection analysis of tetracycline using sensitized europium(III) luminescence detection 433(1988)149
- Wenzel, T.J. and Collette, L.M.  
Lanthanide ions as luminescent chromophores for the LC detection of polynucleotides and nucleic acids 436(1988)299
- Werkhoff, P. and Bretschneider, W.  
Dynamic headspace GC: concentration of volatile components after thermal desorption by intermediate cryofocusing in a cold trap. I. Principle and applications 405(1987)87
- Werkhoff, P. and Bretschneider, W.  
Dynamic headspace GC: concentration of volatile components after thermal desorption by intermediate cryofocusing in a cold trap. II. Effect of sampling and desorption parameters on recovery 405(1987)99
- Werner, A., Siems, W., Schmidt, H., Rapoport, I., Gerber, G., Toguzov, R.T., Tikhonov, Y.V. and Pimenov, A.M.  
Determination of nucleotides, nucleosides and nucleobases in cells of different complexity by reversed-phase and ion-pair HPLC 421(1987)257
- Werner-Busse, A., see Bruns, A. 450(1988)111
- Wessels, J.M.C., see Brons, P.P.T. 422(1987)175
- Wester, P., Gottfries, J., Johansson, K., Klintebäck, F. and Winblad, B.  
Simultaneous LC determination of seventeen of the major monoamine neurotransmitters, precursors and metabolites. I. Optimization of the mobile phase using factorial designs and a computer program to predict chromatograms 415(1987)261
- Wester, P., Gottfries, J. and Winblad, B.  
Simultaneous LC determination of seventeen of the major monoamine neurotransmitters, precursors and metabolites. II. Assessment of

- human brain and cerebrospinal fluid concentrations 415(1987)275
- Westerdahl, G., see Sonesson, A. 417(1987)11
- Westerdahl, G., see Sonesson, A. 431(1988)1
- Wettlaufer, S.H. and Weinstein, L.H.  
Quantitation of polyamines using TLC and image analysis 441(1988)361
- Weyhenmeyer, R., see Hammes, W. 413(1987)264
- Whately, F.R., see Greenaway, W. 409(1987)383
- Whately, F.R., see Greenaway, W. 448(1988)284
- Wheals, B.B.  
Ion chromatography of inorganic anions on a dynamically modified polystyrene-divinylbenzene packing material and its application to anion screening by sequential UV absorbance and electrochemical detection 402(1987)115
- Wheeler, J.R., see McNally, M.E.P. 435(1988)63
- Wheeler, J.R., see McNally, M.E.P. 447(1988)53
- Wheeler, J.R. and McNally, M.E.  
Comparison of packed column and capillary column SFC and HPLC using representative herbicides and pesticides as typical moderate polarity and molecular weight range molecules 410(1987)343
- Whelpton, R., Jonas, G. and Buckley, D.G.  
HPLC resolution of the enantiomers of thioridazine, its metabolites and related compounds 426(1988)223
- Whist, J.E., see Svendsen, J.S. 432(1988)13
- White, C.A., see Lloyd, L.L. 437(1988)447
- White, J.A., see La Follette, G. 431(1988)379
- White, P.C. and Catterick, T.  
Evaluation of absorbance ratioing for solute identification in HPLC using a diode array detector 402(1987)135
- Whitfield, L.R., see Frank, S.K. 419(1987)225
- Whitney, C.C., see Lam, G. 413(1987)309
- Whitsett, J.F. and Kennish, J.M.  
Calibration of TLC with flame ionization detection for the analysis of natural lipid samples 435(1988)343
- Wick, A., see Porziemsky, J.-P. 440(1988)183
- Wiegerinck, M.A.H.M., see Gladdines, M.M. 431(1988)317
- Wieland, H., see Bisse, E. 434(1988)95
- Wiele, K. Van der, see Verhaar, L.A.T. 437(1988)185
- Wientjes, M.G., see Au, J.L.-S. 423(1987)308
- Wiesel, F.-A., see Alfredsson, G. 424(1988)378
- Wiesler, D., see Andreolini, F. 406(1987)375
- Wiet, S.G., see Tavss, E.A. 438(1988)273
- Wigren, C.A., see Krahn, M.M. 437(1988)161
- Wilcock, G.K., see Forsyth, D.R. 433(1988)352
- Wildenberg, H.M. Van den, see Paanakker, J.E. 421(1987)327
- Wildfeuer, A., see Räder, K. 408(1987)341
- Wilkinson, C.P., see Gherezghiher, T. 413(1987)9
- Williams, G.H., see Underwood, R.H. 415(1987)118
- Williams, H.J., see Gonzalez, M.D. 437(1988)311
- Williams, M.C. and Kushwaha, R.S.  
Fractionation of baboon chylomicrons and very-low-density lipoproteins by HPLC 433(1988)257
- Williams, P.E., see Brown, L.L. 426(1988)370
- Williams, R.M., see Lam, G. 413(1987)309
- Williams, S.B., Patchen, L.C. and Churchill, F.C.  
Analysis of blood and urine samples for hydroxychloroquine and three major metabolites by HPLC with fluorescence detection 433(1988)197
- Williams, W.M., see LaCreta, F.P. 414(1987)197
- Williamson, D., see Raha, C. 450(1988)325
- Williamson, R., see Cedarbaum, J.M. 415(1987)393
- Willis, D.E.  
Automated pre-column derivatization of amino acids with *o*-phthalaldehyde by a reagent sandwiching technique 408(1987)217
- Willoughby, R.C., see Koenigbauer, M.J. 413(1987)161
- Willuhn, G., see Leven, W. 410(1987)329
- Wilson, G.S., see Nielsen, R.G. 423(1987)41
- Wilson, I.D.  
Comparison of non-hydrophobic C<sub>18</sub> bonded TLC and HPTLC plates for reversed-phase TLC of ecdysteroids 438(1988)419
- Wilson, I.D., see Raynor, M.W. 436(1988)497
- Wilson, I.D., see Ruane, R.J. 441(1988)355
- Wilson, I.D. and Lewis, S.  
Separation of ecdysteroids by HPTLC using automated multiple development 408(1987)445
- Wilson, K., see Ladds, G. 414(1987)355
- Wilson, M.J., Ballard, K.D. and Walle, T.  
Preparative resolution of the enantiomers of the  $\beta$ -blocking drug atenolol by chiral derivatization and HPLC 431(1988)222
- Wilson, T.D.  
Sample solvent effects in an apparent chiral HPLC separation on  $\beta$ -cyclodextrin 448(1988)31
- Wilson, T.D., Maloney, T. and Amsden, W.B.  
HPLC determination of fentanyl citrate in a parenteral dosage form 445(1988)299
- Wilting, J., see Bos, O.J.M. 424(1988)13
- Wilting, J., see 't Hart, B.J. 424(1988)403
- Winblad, B., see Wester, P. 415(1987)261

- Winblad, B., see Wester, P. 415(1987)275
- Winkler, E., Hohaus, E. and Felber, E.  
Quantitative determination of free primary amines in polymeric amine hardeners. HPLC separation of ethylenediamine and triethylenetetramine after derivatization with salicylaldehyde-diphenylboron chelate and separation of 4,4'-diaminodiphenylmethane by HPLC 436(1988)447
- Wintersberger, F., see Räder, K. 408(1987)341
- Wise, S.A., see Stalcup, A.M. 442(1988)1
- Wisser, H., see Kohse, K.P. 420(1987)25
- Wit, J.S.M. De and Jorgenson, J.W.  
Photoionization detector for open-tubular LC 411(1987)201
- Witkiewicz, Z., see Muniak, W. 436(1988)323
- Witkiewicz, Z. and Goca, B.  
Disc-like liquid crystal stationary phases derived from hexahydroxybenzene 402(1987)73
- Wittebrood, A.J., see De Jong, A.P. 419(1987)233
- Wittfoht, W., Scott, W.J. and Nau, H.  
Assay of methoxyacetic acid in body fluids and tissues by GC-MS following *tert*-butyldimethylsilylation 448(1988)433
- Wittliff, J.L., see Hyder, S.M. 444(1988)225
- Wittliff, J.L., see Jung, S.S. 425(1988)293
- Wittliff, J.L., see Van der Walt, L.A. 425(1988)277
- Woestenborghs, R., Lorreyne, W., Van Rompaey, F. and Heykants, J.  
Determination of cisapride in plasma and animal tissues by HPLC 424(1988)195
- Woestenborghs, R., Lorreyne, W. and Heykants, J.  
Determination of itraconazole in plasma and animal tissues by HPLC 413(1987)332
- Wojaczyńska, M., see Kolarz, B.N. 408(1987)308
- Wójcik, A. and Kwietniewski, L.  
Preparation of porous bead silica 435(1988)55
- Wolf, J.H. and Korf, J.  
Automated solid-phase catalyzed pre-column derivatization of fatty acids for reversed-phase HPLC analysis with fluorescence detection 436(1988)437
- Wolf, W.J., see Peterson, R.E. 444(1988)263
- Wollenweber, E., see Greenaway, W. 448(1988)284
- Wollheim, F., see Åkesson, B. 413(1987)233
- Wolthers, B.G., see Koopman, B.J. 416(1987)1
- Wong, C.L., see Woo, J. 420(1987)73
- Wong, J.T.F., Baker, G.B. and Coutts, R.T.  
Rapid and simple procedure for the determination of urinary phenylacetic acid using derivatization in aqueous medium followed by electron-capture GC 428(1988)140
- Wong, M.A., see Carilli, C.T. 444(1988)203
- Wonigeit, K., see Christians, U. 413(1987)121
- Woo, C.S., see Zaromb, S. 439(1988)283
- Woo, J., Wong, C.L., Teoh, R. and Chan, K.  
LC assay for the simultaneous determination of pyrazinamide and rifampicin in serum samples from patients with tuberculous meningitis 420(1987)73
- Wood, A.J.J., see Koshakji, R.P. 422(1987)294
- Wood, B.A., see Beresford, A.P. 420(1987)178
- Wood, S.G. and Chasseaud, L.F.  
Determination of tulobuterol in human plasma by capillary GC 416(1987)154
- Woodruff Modi, M., see Schuster, D. 433(1988)318
- Woodward, C.J.H. and Emery, P.W.  
Determination of plasma corticosterone using HPLC 419(1987)280
- Woody, M.A. and DasGupta, B.R.  
Fast protein LC of botulinum neurotoxin types A, B and E 430(1988)279
- Wooten, A.L., Prewitt, M.L., Sellers, Jr, T. and Teller, D.C.  
Gel filtration chromatography of resolole phenolic resins 445(1988)371
- Workman, P., Oppitz, M., Donaldson, J. and Lee, F.Y.F.  
HPLC of chlorambucil analogues 422(1987)315
- Workman, P., see Walton, M.I. 430(1988)429
- Workman, P., see Ward, R. 420(1987)223
- Worsley, J.L., see Martin, C.F. 402(1987)376
- Woryna, E., see Muniak, W. 436(1988)323
- Wright, D.J., see Lambert, N. 402(1987)159
- Wright, D.S., Pachla, L.A., Gibson, D.M. and Jordan, R.A.  
Sensitive HPLC method for procaterol in human urine 417(1987)223
- Wright, G.J., see Osman, M.A. 419(1987)452
- Wright, J., see Andrews, P.A. 419(1987)271
- Wright, S.A., see Grigороva, B. 410(1987)419
- Wu, A.T., Massey, I.J. and Kushinsky, S.  
Capillary column GC method using electron-capture detection for the simultaneous determination of nicardipine and its pyridine metabolite II in plasma 415(1987)65
- Wu, D.-F., Reidenberg, M.M. and Drayer, D.E.  
Determination of gossypol enantiomers in plasma after administration of racemate using HPLC with precolumn chemical derivatisation 433(1988)141
- Wu, H.-L., see Tanaka, M. 404(1987)175

- Wu, H.-L., see Tanaka, M. 438(1988)253  
 Wu, J., see Zhu, Y. 438(1988)447  
 Wu, J.T., see Tian, Z.W. 439(1988)159  
 Wu, M.-C., see Takagi, K. 432(1988)47  
 Wu, Q., see Chang, C.A. 404(1987)282  
 Wuis, E.W., Beneken Kolmer, E.W.J.,  
 Van Beijsterveldt, L.E.C., Burgers, R.C.M.,  
 Vree, T.B. and Van der Kleyn, E.  
 Enantioselective HPLC determination of  
 baclofen after derivatization with a chiral  
 adduct of *o*-phthaldialdehyde 415(1987)419  
 Wuis, E.W., Van Beijsterveldt, L.E.C., Dirks,  
 R.J.M., Vree, T.B. and Van der Kleyn, E.  
 Rapid simultaneous determination of  
 baclofen and its  $\gamma$ -hydroxy metabolite in  
 urine by HPLC with UV detection  
 420(1987)212  
 Wunderlich, U., see Löser, C. 430(1988)249  
 Wurst, M., see Bažant, L. 445(1988)337  
 Wurst, M., see Jurková, M. 446(1988)117  
 Wurst, M., see Kysilka, R. 446(1988)315  
 Wurster, R., see Freitag, W. 450(1988)426  
 Wuthe, J., see Kraus, R. 413(1987)257  
 Wyse, D.G., see Tam, Y.K. 423(1987)199  
 Wyss, P.A., see Moor, M.J. 431(1988)455  
 Wyss, R. and Bucheli, F.  
 Quantitative analysis of retinoids in biological  
 fluids by HPLC using column switching. I.  
 Determination of isotretinoin and tretinoin  
 and their 4-oxo metabolites in plasma  
 424(1988)303  
 Wyss, R. and Bucheli, F.  
 Determination of cefetamet and its orally  
 active ester, cefetamet pivoxyl, in biological  
 fluids by HPLC 430(1988)81  
 Wyss, R. and Bucheli, F.  
 Quantitative analysis of retinoids in biological  
 fluids by HPLC using column switching. II.  
 Simultaneous determination of etretinate,  
 acitretin and 13-*cis*-acitretin in plasma  
 431(1988)297

## X

- Xiao, R., see Zhang, T.-Y. 445(1988)199  
 Xiao, Z.-Y., see Zhang, T.-Y. 445(1988)199  
 Xie, M., see Zou, A.-Q. 410(1987)217  
 Xu, B. and Vermeulen, N.P.E.  
 Preparation of wall-coated open-tubular

capillary columns for GC (Review)  
 445(1988)1

- Xu, X. and Pang, K.S.  
 HPLC method for the quantitation of  
 salicylamide and its metabolites in biological  
 fluids 420(1987)313  
 Xue, W. and Carlson, R.M.  
 Separation characteristics of alkylated  
 guanines in HPLC 447(1988)81

## Y

- Yamada, H. and Hattori, T.  
 Determination of molybdenum in soils and  
 plants by HPLC 411(1987)401  
 Yamada, J., see Nomura, A. 448(1988)87  
 Yamada, K., Abe, S., Katayama, K. and Sato, T.  
 Sensitive HPLC method for the determination  
 of phosphatidic acid 424(1988)367  
 Yamada, K., Asano, O., Yoshimura, T. and  
 Katayama, K.  
 Highly sensitive GC-MS method for the  
 determination of platelet-activating factor in  
 human blood 433(1988)243  
 Yamada, K., see Chiu, W.-Y. 428(1988)25  
 Yamada, M., see Ikai, Y. 411(1987)313  
 Yamada, M., see Oka, H. 411(1987)437  
 Yamada, M., see Oka, H. 449(1988)448  
 Yamada, T., see Sawada, S. 402(1987)283  
 Yamada, Y., see Nakano, K. 438(1988)61  
 Yamada, Y., see Tanaka, Y. 407(1987)197  
 Yamaga, N., Adachi, K., Kohara, H. and  
 Shimizu, K.  
 Internal standards for quantitative GC of  
 individual bile acids after group separation of  
 bile acids in urine 422(1987)25  
 Yamaga, N., see Shimizu, K. 432(1988)21  
 Yamaguchi, H., see Kasai, R. 407(1987)205  
 Yamaguchi, H., see Yoshida, T. 445(1988)175  
 Yamaguchi, J., Hanai, T. and Cai, H.  
 Selectivity related to carbon loading and  
 end-capping of octadecyl-bonded silica gels  
 in reversed-phase LC 441(1988)183  
 Yamaguchi, M., Matsunaga, R., Fukuda, K. and  
 Nakamura, M.  
 Determination of free and total phenylacetic  
 and *p*- and *m*-hydroxyphenylacetic acids in  
 human urine by LC with fluorimetric  
 detection 414(1987)275



- Yamaguchi, M., see Iwata, T. 421(1987)43
- Yamaguchi, T., see Kominami, G. 417(1987)216
- Yamaguchi, T., see Shinbo, T. 405(1987)145
- Yamaguchi, Y., Haginaka, J., Kunitomo, M., Yasuda, H. and Bandô, Y.  
HPLC determination of desmosine and isodesmosine in tissues and its application to studies of alteration of elastin induced by atherosclerosis 422(1987)53
- Yamaji, A., Kataoka, K., Oishi, M., Kanamori, N., Hiraoka, E. and Mishima, M.  
Simultaneous determination of procainamide and N-acetylprocainamide in serum by GC with nitrogen-selective detection 415(1987)143
- Yamaji, A., Kataoka, K., Oishi, M., Kanamori, N., Tagawa, T. and Mimaki, T.  
Simple method for determination of flunarizine in serum by GC 421(1987)372
- Yamaji, Y., see Matsukura, H. 434(1988)196
- Yamamoto, F.M., Rokushika, S. and Hatano, H.  
Prediction of thermodynamic retention behaviour in reversed-phase LC with aqueous eluents and various concentrations of methanol by using the solubility concept model 408(1987)21
- Yamamoto, K., Kohda, Y., Saitoh, Y. and Nakagawa, F.  
Determination of ambenonium in biological samples by reversed-phase ion-pair LC 433(1988)167
- Yamamoto, K., see Fujimori, K. 414(1987)167
- Yamamoto, K., see Kojima, K. 413(1987)326
- Yamamoto, M., see Maeda, Y. 410(1987)413
- Yamamoto, M., see Miwa, H. 416(1987)237
- Yamamoto, M., see Miwa, H. 421(1987)33
- Yamamoto, N., see Oka, H. 411(1987)437
- Yamamoto, S., Nomura, M. and Sano, Y.  
Resolution of proteins in linear gradient elution ion-exchange and hydrophobic interaction chromatography 409(1987)101
- Yamamoto, S., see Hamano, T. 411(1987)423
- Yamamoto, T., Moriwaki, Y., Takahashi, S., Hada, T. and Higashino, K.  
Rapid and simultaneous determination of pyrazinamide and its major metabolites in human plasma by HPLC 413(1987)342
- Yamanaka, C., see Yokoyama, T. 403(1987)151
- Yamanaka, S., see Sagara, Y. 431(1988)170
- Yamasaki, R.B., Ritland, T.G., Barnby, M.A. and Klocke, J.A.  
Isolation and purification of salannin from neem seeds and its quantification in neem and chinaberry seeds and leaves 447(1988)277
- Yamasaki, Y., see Kato, Y. 404(1987)333
- Yamasaki, Y., see Kato, Y. 447(1988)212
- Yamashita, J., Numakura, T., Kita, H., Suzuki, T., Ôi, S., Miyano, S., Hashimoto, H. and Takai, N.  
HPLC separation of enantiomers on axially chiral binaphthalene derivatives bonded to silica gel 403(1987)275
- Yamashita, K., Watanabe, K., Ishibashi, M., Katori, M. and Miyazaki, H.  
Dimethylisopropylsilyl ether derivatives in GC-MS of some prostanoids. Application to the microdetermination of 15-ketoprostaglandin  $F_{2\alpha}$  formed by enzymatic conversion of  $PGF_{2\alpha}$  in vitro 424(1988)1
- Yamashita, K., Watanabe, K., Koga, R., Mizuguchi, S., Hashimoto, Y., Nakamura, T. and Umezawa, H.  
Determination of deoxyspergualin by GC-selected-ion monitoring 424(1988)39
- Yamatodani, A., see Yonekura, T. 427(1988)320
- Yamauchi, Y., Takada, M., Oshima, R., Kumanotani, J. and Senô, M.  
LC analysis of products of reaction between methyl linoleate and 4-*tert*-butyl-*o*-benzoquinone 408(1987)171
- Yamazaki, M., see Fujii, Y. 448(1988)157
- Yamazaki, Y., see Oka, H. 449(1988)448
- Yamboliev, I., see Tencheva, J. 421(1987)396
- Yan, K., see Zhu, Y. 437(1988)265
- Yan, K., see Zhu, Y. 438(1988)447
- Yanagi, H., Soga, A. and Ogino, S.  
Purification of human urinary erythropoietin 417(1987)178
- Yanagihara, Y., see Hirata, N. 434(1988)71
- Yanagihara, Y., see Yasukawa, K. 410(1987)129
- Yang, Q., Sun, Z.-P. and Ling, D.-K.  
Resolution of enantiomeric drugs of some  $\beta$ -amino alcohols as their urea derivatives by HPLC on a chiral stationary phase 447(1988)208
- Yang, S.S. and Gilpin, R.K.  
Reordering/resolution of silica immobilized hydrogen bonding ligands used as stationary phases in HPLC 408(1987)93
- Yang, S.S. and Gilpin, R.K.  
Reversed-phase separation of a standard mixture of polycyclic aromatic hydrocarbons on an adamantyl surface 439(1988)414
- Yang, S.S. and Gilpin, R.K.  
Influence of bonding chemistry on the reordering/resolution of silica immobilized alkyl chains 449(1988)115
- Yang, X.-H. and Zhang, H.  
Determination of trace amounts of iodine in

- sedimentary rocks by ion chromatography with amperometric detection 436(1988)107
- Yang, Y.-C., see Rohrbaugh, D.K. 447(1988)165
- Yang, Z.C., see Peng, C.T. 436(1988)137
- Yarmaloff, L.W., see Wenzel, T.J. 433(1988)149
- Yasaka, Y., see Tanaka, M. 404(1987)175
- Yasaka, Y., see Tanaka, M. 438(1988)253
- Yasuda, H., see Haginaka, J. 447(1988)365
- Yasuda, H., see Haginaka, J. 447(1988)373
- Yasuda, H., see Yamaguchi, Y. 422(1987)53
- Yasueda, H., see Mita, H. 430(1988)299
- Yasuhara, A.  
Comparison of volatile components between fresh and rotten mussels by GC-MS 409(1987)251
- Yasukawa, K., Tamura, Y., Uchida, T., Yanagihara, Y. and Noguchi, K.  
Development of a C<sub>18</sub>-bonded vinyl alcohol copolymer gel for reversed-phase HPLC 410(1987)129
- Yazaki, S., see Shibanoki, S. 421(1987)425
- Yeager, S.A., see Duff, G.A. 416(1987)71
- Yeates, R.A., see Scharpf, F. 413(1987)91
- Yee, G.C., see Gmur, D.J. 425(1988)343
- Yeh, J., see Moore, J.M. 410(1987)297
- Yeung, E.S., see Reitsma, B.H. 405(1987)295
- Yim, J.J., Lim, K.T., Kim, N. and Jacobson, K.B.  
Semi-micro quantitative assay for tryptophan oxygenase by HPLC 419(1987)296
- Yin, I.-H., see Ravichandran, K. 439(1988)213
- Yip, T.-T., see Belew, M. 403(1987)197
- Yokoyama, T., see Furukawa, M. 435(1988)219
- Yokoyama, T., Yamanaka, C. and Tarutani, T.  
Formation of silicate complexes of aluminium in aqueous solution 403(1987)151
- Yokoyama, Y., see Murata, K. 415(1987)231
- Yokoyama, Y., see Murata, K. 423(1987)51
- Yomo, N., see Nakano, K. 438(1988)61
- Yon, J.M., Betton, J.-M., Desmadril, M., Mitraki, A., Minard, P., Gaillard, S., Ballery, N. and Missiakas, D.  
Survey of the folding pathway of a two-domain protein phosphoglycerate kinase 440(1988)421
- Yonekawa, W.D., see Kapetanovic, I.M. 414(1987)265
- Yonekawa, W.D., see Torchin, C.D. 426(1988)444
- Yonekura, T., Kamata, S., Wasa, M., Okada, A., Yamatodani, A., Watanabe, T. and Wada, H.  
Simultaneous determination of plasma phenethylamine, phenylethanolamine, tyramine and octopamine by HPLC using derivatization with fluorescamine 427(1988)320
- Yong, Y.-S., Soerensen, K.J. and Cant, N.W.  
Automated analysis of the gas mixture from ethylene epoxidation by oxygen and nitrous oxide mixtures using GC 450(1988)399
- Yoshida, H., see Fujimoto, C. 411(1987)213
- Yoshida, H., see Nakashima, K. 414(1987)11
- Yoshida, H., see Tamai, G. 423(1987)147
- Yoshida, H., see Tamai, G. 423(1987)155
- Yoshida, H., see Tamai, G. 423(1987)163
- Yoshida, H., see Tanaka, S. 447(1988)383
- Yoshida, K.-I., see Marunaka, T. 433(1988)177
- Yoshida, K., see Marunaka, T. 420(1987)43
- Yoshida, S., Adachi, T. and Hirose, S.  
4-Bromomethyl-6,7-dimethoxycoumarin as a fluorescence reagent for precolumn derivatization of 5-fluorouracil compounds in HPLC 430(1988)156
- Yoshida, S., Murai, H., Hirose, S. and Takeda, T.  
Rapid and sensitive on-line precolumn purification and HPLC assay for bile acids in serum 431(1988)27
- Yoshida, S., see Iwaki, K. 404(1987)117
- Yoshida, T., Aetake, A., Yamaguchi, H., Nimura, N. and Kinoshita, T.  
Determination of carnitine by HPLC using 9-anthryldiazomethane 445(1988)175
- Yoshida, T., see Hatano, T. 435(1988)285
- Yoshida, T., see Sagara, K. 403(1987)253
- Yoshida, T., see Sagara, K. 409(1987)365
- Yoshida, T., see Sugita, S. 421(1987)412
- Yoshihara, K., see Hashimoto, K. 438(1988)401
- Yoshikawa, S., Okuda, S., Ohe, T., Kamohara, K., Maeda, O. and Sawada, T.  
LC determination of vanillylmandelic acid and homovanillic acid by a column-switching technique involving direct injection of urine 421(1987)111
- Yoshimura, H., see Endoh, Y.S. 426(1988)202
- Yoshimura, T., see Tanaka, N. 406(1987)247
- Yoshimura, T., see Yamada, K. 433(1988)243
- Yoshitake, N., see Furukawa, M. 435(1988)219
- Yoshizawa, T., see Swanson, S.P. 414(1987)335
- Yosizawa, Z., see Kodama, C. 429(1988)293
- Young, H.S., see Prasad, M.R. 428(1988)221
- Young, J.H., see Powers, A.G. 432(1988)321
- Young, P.R. and Grynspan, F.  
Analysis of methylated amino acids by HPLC: methylation of myelin basic protein 421(1987)130
- Youngberg, P.J., see Shetty, P.H. 411(1987)61
- Yourtee, D., see Smith, R.E. 439(1988)83
- Yuasa, I., see Ikebuchi, J. 432(1988)401
- Yuasa, S., see Fukuhara, T. 411(1987)502
- Yui, S., see Harada, H. 407(1987)299

Yui, Y., see Mita, H. 430(1988)299  
 Yunis, A.A., see Abou-Khalil, S. 417(1987)111  
 Yuthavong, Y., see Krungkrai, J. 417(1987)47

## Z

- Zábršová, H., see Turánek, J. 410(1987)470  
 Zagyvai, P., see Nagy, L.G. 406(1987)359  
 Zahn, H., see Mies, H.H. 405(1987)365  
 Zahra, A., see Rop, P.P. 427(1988)172  
 Zaitso, K., see Wang, Z.-J. 430(1988)223  
 Zampino, M., see Palmerini, C.A. 417(1987)378  
 Zanoboni-Muciaccia, W., see Pedrazzini, S. 413(1987)338  
 Zanoboni-Muciaccia, W., see Pedrazzini, S. 415(1987)214  
 Zaparucha, A., see Delee, E. 450(1988)183  
 Zappia, V., see Della Ragione, F. 440(1988)141  
 Zappia, V., see Porcelli, M. 440(1988)151  
 Zapponi, M.C., see Iadarola, P. 443(1988)317  
 Zaranski, M.T. and Bidleman, T.F.  
 High-volume elution chromatography of dichlorobenzenes on a polyurethane foam-Tenax sandwich cartridge 409(1987)235  
 Zare, R.N., see Huang, X. 425(1988)385  
 Zare, R.N., see Roach, M.C. 426(1988)129  
 Zarins, Z.M., see Blouin, F.A. 441(1988)443  
 Zaromb, S., see Otagawa, T. 408(1987)81  
 Zaromb, S., Woo, C.S., Quandt, K., Rice, L.M., Fermaint, A. and Mitnaul, L.J.  
 Simple permeation absorber for sampling and preconcentrating hazardous air contaminants 439(1988)283  
 Zaromb, S. and Beasley, T.M.  
 Technique for calibrating air samplers and certain other analytical devices 438(1988)100  
 Zaslavsky, B.Y., Miheeva, L.M., Aleschko-Ozhevskii, Y.P., Mahmudov, A.U., Bagirov, T.O. and Garaev, E.S.  
 Distribution of inorganic salts between the coexisting phases of aqueous polymer two-phase systems. Interrelationship between the ionic and polymer composition of the phases 439(1988)267  
 Zaslavsky, B.Y., Miheeva, L.M., Gasanova, G.Z. and Mahmudov, A.U.  
 Effect of polymer composition on the relative hydrophobicity of the phases of the biphasic system aqueous dextran-poly(ethylene glycol) 403(1987)123  
 Zav'yalov, V.P., see Shukun, S.A. 432(1988)137  
 Zecca, L., Ferrario, P., Pirola, R. and Bareggi, S.R.  
 Analysis of chlorodesmethyl diazepam and its metabolites in plasma and urine 420(1987)417  
 Zee, J.A., see Lavigne, C. 410(1987)201  
 Zee, R. Van der, see Welling, G.W. 418(1987)223  
 Zeece, M.G., see Holt, D.L. 449(1988)271  
 Zeeuw, R.A. De, see Benecke, R. 417(1987)287  
 Zemaitis, M.A., see D'Souza, M.J. 421(1987)198  
 Zeman, I.  
 Misuse of the term "detergent" 426(1988)452  
 Zeman, J., see Verner, P. 416(1987)439  
 Zemanová, I., see Hajšlová, J. 439(1988)307  
 Zeng, W., Liang, W. and Tu, G.  
 Assay of five corynoline-type alkaloids in *Corydalis bungeana* by reversed-phase HPLC 408(1987)426  
 Zenkevich, I.G. and Ioffe, B.V.  
 System of retention indices for a linear temperature programming regime 439(1988)185  
 Zerilli, L.F., see Beretta, E. 416(1987)144  
 Zhang, G., see Sagara, K. 409(1987)365  
 Zhang, H., see Yang, X.-H. 436(1988)107  
 Zhang, S., see Liu, G. 435(1988)327  
 Zhang, T.-Y., Cai, D.-G. and Ito, Y.  
 Separations of flavonoids and alkaloids in medicinal herbs by high-speed counter-current chromatography 435(1988)159  
 Zhang, T.-Y., Pannell, L.K., Pu, Q.-L., Cai, D.-G. and Ito, Y.  
 Separation of hydroxyanthraquinone derivatives extracted from rheum with analytical high-speed counter-current chromatography 442(1988)455  
 Zhang, T.-Y., see Ito, Y. 437(1988)121  
 Zhang, T.-Y., see Ito, Y. 449(1988)135  
 Zhang, T.-Y., see Ito, Y. 449(1988)153  
 Zhang, T.-Y., Xiao, R., Xiao, Z.-Y., Pannell, L.K. and Ito, Y.  
 Rapid separation of flavonoids by analytical high-speed counter-current chromatography 445(1988)199  
 Zhang, Y., see Balmér, K. 417(1987)357  
 Zhang, Y.Y., see Swobodnik, W. 423(1987)75  
 Zhivkova, Z., see Tencheva, J. 421(1987)396  
 Zhou, X., see Parrish, C.C. 435(1988)350  
 Zhou, Z.M., Anders, J.C., Chung, H. and Theoharides, A.D.  
 Analysis of artesunic acid and dihydroqinghaosu in blood by HPLC with

- reductive electrochemical detection  
414(1987)77
- Zhu, M.-D., see Hjertén, S. 403(1987)47
- Zhu, Y., Yan, K., Wu, J. and Tu, G.  
Assay of lignans of *Schizandra chinensis* in  
Sheng Mai San by HPLC 438(1988)447
- Zhu, Y., Yan, K. and Tu, G.  
Separation and determination of  
homoisoflavonoids in *Ophiopogon japonicus*  
by reversed-phase HPLC 437(1988)265
- Zia-Gharib, F., see Farrant, M. 417(1987)385
- Ziegler, M., see Toguzov, R.T. 434(1988)447
- Zima, J., see Beran, M. 442(1988)431
- Zimina, T.M., see Maltsev, V.G. 416(1987)45
- Zimmer, K.-O., see Christians, U. 413(1987)121
- Zimmerman, C.L., see Thongnopnua, P.  
433(1988)345
- Zirrolli, J.A., see Scriba, G.K.E. 433(1988)31
- Zoanetti, G.D., see Badcock, N.R. 421(1987)147
- Zobel, H., Helms, M., Lehmann, A., Radeck, D.  
and Gutmann, M.  
Analytical characterisation of hydrocarbon  
and hetero compounds mixtures from  
reactions of brown coal (in German)  
446(1988)305
- Zoest, A.R., see Hung, C.T. 425(1988)331
- Zomer, G., see Jansen, E.H.J.M. 437(1988)268
- Zonta, F., see Stancher, B. 440(1988)37
- Zonta, F., Stancher, B. and Marletta, G.P.  
Simultaneous HPLC analysis of free  
carotenoids and carotenoid esters  
403(1987)207
- Zoontjes, P.W., see Van Ginkel, L.A.  
445(1988)385
- Zou, A.-Q., Xie, M. and Luo, X.-D.  
Determination of artesunic acid after  
chemical derivatization with O-*p*-nitrobenzyl-  
N,N'-diisopropylisourea by HPLC and UV  
absorption 410(1987)217
- Zschunke, M., see Brendel, E. 426(1988)406
- Zuhowski, E.G., see Egorin, M.J. 415(1987)148
- Zuidema, J., see Pieters, F.A.J.M. 422(1987)322
- Zuk, M.M., Tyczkowska, K., Ben-Hur, E.,  
Newman, H.C., Rosenthal, I. and Crane, S.W.  
Reversed-phase LC determination of  
chloroaluminum phthalocyanine  
tetrasulfonate in canine tissues and fluids  
433(1988)367
- Żukowski, J., Sybilska, D., Bojarski, J. and Szejtli,  
J.  
Resolution of chiral barbiturates into  
enantiomers by reversed-phase HPLC using  
methylated  $\beta$ -cyclodextrins 436(1988)381
- Zurlinden, E., see Conte, Jr., J.E. 417(1987)452
- Zussman, B.D., see Brett, M.A. 419(1987)438

SUBJECT INDEX  
VOLUMES 401-450

# A

## Abietic acid

GC analysis of underivatized resin acids  
448(1988)428

## Abscisic acid

Direct separation of (*S*)- and (*R*)-abscisic acid  
on a commercially available chiral HPLC column  
402(1987)374

Determination of abscisic acid in small amounts  
of plant material 404(1987)346

Chromatographic resolution. XXI. Direct optical  
resolution of abscisic acid by HPLC on cellulose  
tris(3,5-dimethylphenylcarbamate) 448(1988)454

## Abscisic acid methyl ester

Resolution of the enantiomers of abscisic acid  
methyl ester by HPLC using a stationary phase  
of cellulose tris(3,5-dimethylphenylcarbamate)-  
coated silica gel 402(1987)371

## Absolute calibration method

Comparison of the absolute calibration method  
with the method of standard addition for the  
determination of halothane in blood by GC  
headspace analysis 425(1988)99

## Absorbance cell

Optical errors in an LC absorbance cell  
448(1988)193

## Absorbance ratioing

Evaluation of absorbance ratioing for solute  
identification in HPLC using a diode array  
detector 402(1987)135

## Absorptiometric quantitation

Direct absorptiometric quantitation in HPLC in

the absence of concomitant reference standards  
434(1988)363

## Acemetacin

Method for the determination of acemetacin, a  
non-steroidal anti-inflammatory drug, in plasma  
HPLC 413(1987)305

## Acetaldehyde

Dynamic headspace GC of volatile compounds  
in milk 404(1987)163

Simple and sensitive method for the  
determination of acetaldehyde in blood by GC  
414(1987)454

GC determination of acetaldehyde and acetone  
in human blood by purge and trap, using  
permeation tubes for calibration 437(1988)294

## Acetamide

GC-MS method for the determination of  
dimethylacetamide and metabolites in whole  
blood 428(1988)156

## *p*-Acetamidobenzoic acid

Direct and simultaneous HPLC assay for the  
determination of *p*-aminobenzoic acid and its  
conjugates in human urine 426(1988)103

## *p*-Acetamidohippuric acid

Direct and simultaneous HPLC assay for the  
determination of *p*-aminobenzoic acid and its  
conjugates in human urine 426(1988)103

## Acetamidomalonic acid

Chiral Michael additions of acetamidomalonic acid  
to  $\alpha$ -enones. NMR and HPLC resolution of  
enantiomers 450(1988)183

## Acetic acid

Studies on sample preconcentration in ion  
chromatography. VIII. Preconcentration of  
carboxylic acids prior to ion-exclusion separation  
447(1988)155

Extractive pentafluorobenzoylation of formic, acetic, levulinic, benzoic and phthalic acids, studied by LC and dual-oven capillary GC 447(1988)329

#### **Acetoacetanilide**

TLC separation of some sulphur drugs using acetoacetanilide as a coupling agent 441(1988)454

#### **Acetohydroxy acid synthase**

Separation and characterization of two forms of acetohydroxy acid synthase from black Mexican sweet corn cells 444(1988)251

#### **Acetone**

Simultaneous determination of methanol, ethanol, acetone, isopropanol and ethylene glycol in plasma by GC 414(1987)248

GC method for the estimation of acetone and its metabolites in biological samples 415(1987)372

Detection of endogenous acetone in normal human breath 422(1987)235

GC determination of acetaldehyde and acetone in human blood by purge and trap, using permeation tubes for calibration 437(1988)294

Static headspace GC method for the determination of residual solvents in vigabatrin drug substance 438(1988)103

#### **Acetonitrile**

Retention in capillary supercritical fluid chromatography with carbon dioxide mobile phases containing polar modifiers 406(1987)223

#### **Acetylacetoncycloserine**

Simultaneous quantification of cycloserine and its prodrug acetylacetoncycloserine in plasma and urine by HPLC using UV absorbance and fluorescence after post-column derivatization 414(1987)121

#### **3-Acetylaconitine**

HPLC analysis of C<sub>19</sub>-diterpenoid alkaloids 402(1987)366

#### **N-Acetyl-S-(N-alkylthiocarbamoyl)-L-cysteine**

Determination of N-acetyl-S-(N-alkylthiocarbamoyl)-L-cysteine, a principal metabolite of alkyl isothiocyanates, in rat urine 414(1987)19

#### **N-Acetyl-S-allylcysteine**

Unusual conjugates in biological profiles originating from consumption of onions and garlic 421(1987)1

#### **Acetyl-5-aminosalicylic acid**

Simple and sensitive HPLC assay for 5-aminosalicylic acid and acetyl-5-aminosalicylic acid in serum 413(1987)300

HPLC assay of 5-aminosalicylic acid and its acetylated metabolite in biological fluids using electrochemical detection 425(1988)214

#### **5-Acetylaminosalicylic acid**

Stability of 5-aminosalicylic acid and 5-acetylaminosalicylic acid in plasma 432(1988)358

#### **N-Acetyl-S-(2-carboxypropyl)cysteine**

Unusual conjugates in biological profiles originating from consumption of onions and garlic 421(1987)1

#### **Acetylcarnitine**

Analytical method for urinary glutarylcarnitine, acetylcarnitine and propionylcarnitine with a carboxylic acid analyser and a reversed-phase column 423(1987)297

#### **Acetylcholine**

Determination of acetylcholine and choline in perchlorate extracts of brain tissue using LC-electrochemistry with an immobilized-enzyme reactor 414(1987)167

Determination of acetylcholine and choline in microdialysates from spinal cord of rat using LC with electrochemical detection 423(1987)85

LC with electrochemical detection for the determination of choline and acetylcholine in plasma and red blood cells. Failure to detect acetylcholine in blood of humans and mice 428(1988)1

Determination of acetylcholine and choline in human cerebrospinal fluid using HPLC combined with an immobilized enzyme reactor: ageing-induced change of acetylcholine level 431(1988)389

#### **N-Acetylcysteine**

HPLC fluorometry for the determination of thiols in biological samples using N-[4-(6-dimethylamino-2-benzofuranyl)phenyl]maleimide 414(1987)11

Determination of cysteine, glutathione and N-acetylcysteine in plasma by ion-pair reversed-phase LC and post-column derivatization 432(1988)65

#### **N-Acetyldekamycins**

Identification of dekamycin antibiotics 440(1988)71

**$\alpha$ -N-Acetylendorphin**

Combined use of size-exclusion and reversed-phase HPLC for characterization of  $\beta$ -endorphin processing pathways 416(1987)255

**N-Acetylgalactosamine-binding lectin**

Purification of human erythrocytes specific lectins from rice bean, *Phaseolus calcaratus* syn. *Vigna umbellata*, by HPLC 431(1988)37

**Acetylhomotaurinate**

Determination of calcium acetylhomotaurinate by LC with fluorimetric and electrochemical detection 414(1987)417

**Acetylhydrazine**

Analysis of isoniazid, acetylhydrazine and [ $^{15}\text{N}_2$ ]acetylhydrazine in serum by capillary GC-ammonia chemical ionization MS 420(1987)171

**4-[(3-(4-Acetyl-3-hydroxy-2-propylphenoxy)-propyl)sulphonyl]- $\gamma$ -oxobenzenebutanoic acid**

HPLC determination of 4-[(3-(4-acetyl-3-hydroxy-2-propylphenoxy)propyl)sulfonyl]- $\gamma$ -oxobenzenebutanoic acid and metabolites in human plasma 417(1987)447

**N-Acetyl-S-(N-methylcarbamoyl)cysteine**

Identification of N-acetyl-S-(N-methylcarbamoyl)cysteine, a human metabolite of N,N-dimethylformamide and N-methylformamide 414(1987)399

GC method for the determination of N-acetyl-S-(N-methylcarbamoyl)cysteine, a metabolite of N,N-dimethylformamide and N-methylformamide, in human urine 431(1988)361

**N-Acetylmuramyl-L-alanine amidase**

Comparative susceptibility of a peptidoglycan monomer from *Brevibacterium divaricatum* and its anhydromuramyl analogue to hydrolysis with N-acetylmuramyl-L-alanine amidase. Isolation and characterization of anhydromuramyl-peptidoglycan monomer 440(1988)405

**N-Acetylmuramyl-L-alanyl-D-isoglutamine**

N-Acetylmuramyl-L-alanyl-D-isoglutamine stimulates the immune response of B cells. Use of fast protein LC to purify a novel monokine involved in this effect 440(1988)95

**N-Acetylprocainamide**

Simultaneous determination of procainamide and

N-acetylprocainamide in serum by GC with nitrogen-selective detection 415(1987)143

HPLC drug analysis by direct injection of whole blood samples. II. Determination of hydrophilic drugs 423(1987)155

**Acetylsalicylic acid**

Rapid HPLC assay for the simultaneous analysis of non-steroidal anti-inflammatory drugs in plasma 416(1987)293

Determination of plasma aspirin and salicylic acid concentrations after low aspirin doses by HPLC with post-column hydrolysis and fluorescence detection 420(1987)425

**Acetylsulphamethoxazole**

Rapid method for the determination of either plasma sulphapyridine or sulphamethoxazole and their acetyl metabolites using HPLC 414(1987)223

**N-Acetylsulphamethoxazole**

Concurrent analysis of methotrexate, trimethoprim, sulphamethoxazole and their major metabolites in plasma by HPLC 422(1987)328

**2-Acetyl-4(5)-tetrahydroxybutylimidazole**

Direct method for the determination of 2-acetyl-4(5)-tetrahydroxybutylimidazole in caramel colours and beer by HPLC with absorbance detection 407(1987)405

**Acetyltributyl citrate**

Analysis of the plasticiser acetyltributyl citrate in foods by stable isotope dilution GC-MS 437(1988)281

**Acid dyes**

Characterisation of acid dyes in forensic fibre analysis by HPLC using narrow-bore columns and diode array detection 442(1988)187

**Acid fusion**

Use of acid or alkaline fusion for the quantitative degradation of condensation polymers 406(1987)167

 **$\alpha_1$ -Acid glycoprotein**

Direct HPLC resolution of dihydropyridine enantiomers 450(1988)191

**Acidic fibroblast growth factor**

HPLC of phenylthiocarbamyl-amino acids. Application to carboxyl-terminal sequencing of proteins 409(1987)299

**Acid phosphatase**

Highly sensitive assay for alkaline and acid phosphatase activities by HPLC with electrochemical detection 417(1987)41

Assay of 5'-nucleotidase and simultaneous detection of interfering enzymes by anion-exchange CC 420(1987)398

HPLC determination of acid and alkaline phosphatases in urine 434(1988)196

**Acid Red**

Simple method for the analysis of food dyes on reversed-phase thin-layer plates 411(1987)437

**Acids**

Glass capillary GC analysis of volatile components of canned Philippine pineapple juice 409(1987)409

Feasibility of isotachochromatography as a method for the preparative separation of weak acids and weak bases. I. Theoretical considerations 436(1988)173

Feasibility of isotachochromatography as a method for the preparative separation of weak acids and weak bases. II. Experimental verification 436(1988)181

 **$\alpha$ -Acids**

Avoidance of trace metal activity with micro-LC in the analysis of hop and beer bitter acids 407(1987)159

 **$\beta$ -Acids**

Avoidance of trace metal activity with micro-LC in the analysis of hop and beer bitter acids 407(1987)159

**Acitretin**

Quantitative analysis of retinoids in biological fluids by HPLC using column switching. II. Simultaneous determination of etretinate, acitretin and 13-*cis*-acitretin in plasma 431(1988)297

***cis*-Aconitate**

HPLC measurement of selected blood citric acid cycle intermediates 425(1988)47

**Aconitine**

HPLC analysis of C<sub>19</sub>-diterpenoid alkaloids 402(1987)366

**Acoustic field-flow fractionation**

Acoustic field-flow fractionation 446(1988)151

**Acridanones**

Analysis of benzodiazepines. II. HPLC fluorescence detection after molecular rearrangement to acridanones 428(1988)321

**Acridines**

Determination of quinacrine (mepacrine) in plasma by HPLC with fluorimetric detection 420(1987)341

**Acrylic acid oligomers**

Separation and characterization of acrylic acid oligomers by NMR spectroscopy and thermospray ion-exchange LC-MS 408(1987)349

**Acrylonitrile**

Investigation by pyrolysis-GC of the composition of multicomponent polymeric microheterogeneous systems based on some vinyl monomers 404(1987)183

**Actin**

Selective purification of cardiac myosin by a high-performance salicylate affinity column 435(1988)185

**Activity coefficients**

Reversed-flow GC applied to physico-chemical measurements 446(1988)39

**Activity staining**

Activity staining of inulin fructotransferase II on polyacrylamide gels 445(1988)282

**Acyclovir**

Determination of acyclovir in plasma by column LC with fluorescence detection 420(1987)197

**Acylaminopenicillins**

Application of HPLC of some antibiotics in clinical microbiology 427(1988)257

**Acylolins**

Enantiomer separation of acylolins and cyanohydrins by enantioselective capillary GC 439(1988)301

**Acyl triglycerides**

Characterization of a lipid-rich fraction synthesized by *Streptomyces avermitilis* 441(1988)31

**Adenine**

HPLC analysis of ascorbate-2-phosphate, adenine and hypoxanthine in stored human blood 419(1987)95



**Adenine nucleotides**

Demonstration of the adenosine reservoirs with nitrobenzylthioinosine in liver and diaphragm by HPLC 426(1988)249

**Adenosine**

Rapid separation of anomeric purine nucleosides by TLC on a chiral stationary phase 410(1987)226

**Adenosine deaminase**

Determination of adenosine deaminase and purine nucleoside phosphorylase activities using HPLC and radiochromatographic methods 434(1988)439

**Adenosine 5'-diphosphate**

Isocratic ion-pair reversed-phase HPLC analysis of ATP and ADP 423(1987)308

**Adenosine 3':5'-monophosphate**

Reversed-phase LC determination of cyclic adenosine 3',5'-monophosphate in rat brain cortex 426(1988)177

**Adenosine 5'-triphosphate**

Isocratic ion-pair reversed-phase HPLC analysis of ATP and ADP 423(1987)308

Liposomally entrapped adenosine triphosphate. Improved efficiency against experimental brain ischaemia in the rat 440(1988)455

**S-Adenosyl-L-methionine**

Fast reversed-phase HPLC determination of <sup>14</sup>C- or <sup>3</sup>H-labelled S-adenosylmethionine in reaction mixtures 410(1987)500

**Adenylate deaminase**

Assay of 5'-nucleotidase and simultaneous detection of interfering enzymes by anion-exchange CC 420(1987)398

**Adibendan**

HPLC method for the determination of adibendan in plasma and urine 422(1987)346

**Adinazolam**

HPLC assay of adinazolam and its desmethyl metabolite 424(1988)215

**Adlumidicine**

Determination of alkaloids of *Fumaria parviflora* and *Fumaria capreolata* by HPLC and capillary isotachopheresis 445(1988)258

**Adrenaline**

Determination of  $\beta$ -phenylalkylamines in

pharmaceutical preparations by analytical capillary isotachopheresis 405(1987)389

Simultaneous LC determination of seventeen of the major monoamine neurotransmitters, precursors and metabolites. I. Optimization of the mobile phase using factorial designs and a computer program to predict chromatograms 415(1987)261

Simultaneous LC determination of seventeen of the major monoamine neurotransmitters, precursors and metabolites. II. Assessment of human brain and cerebrospinal fluid concentrations 415(1987)275

Determination of catecholamines in biological tissues by LC with coulometric detection 415(1987)388

Improved HPLC method with electrochemical detection for the determination of urinary catecholamines using a new electrochemical detector 416(1987)340

Combined ion-pair extraction and HPLC for the determination of the biogenic amines and their major metabolites in single brain tissue samples 421(1987)51

Assay for dopamine  $\beta$ -hydroxylase in human plasma and rat serum by HPLC with fluorimetric detection 421(1987)237

Routine determination of urinary free catecholamines by HPLC with electrochemical detection 424(1988)263

LC determination of urinary catecholamines after one-step alumina extraction 424(1988)373

Species-dependent differences in recovery of 3,4-dihydroxybenzylamine in assays of plasma catecholamines 430(1988)123

Selected-ion monitoring GC-MS analysis of catecholamines: enhancement of sensitivity by a simple clean-up step on Sephadex G-10 433(1988)31

Sensitive method for determination of picogram amounts of epinephrine and other catecholamines in microdissected samples of rat brain using LC with electrochemical detection 433(1988)41

Perfluorinated acids as ion-pairing agents in the determination of monoamine transmitters and some prominent metabolites in rat brain by HPLC with amperometric detection 449(1988)191

**$\beta$ -Adrenergic receptor-blocking agents**

Qualitative GC and GC-MS screening for  $\beta$ -blockers in urine after solid-phase extraction using Extrelut-1 columns 426(1988)194

**Adrenocorticotrophic hormone-related peptides**

Separation of protein hormones (Review) 418(1987)245

**Adriamycin**

Estimation of doxorubicin and doxorubicinol by HPLC and advanced automated sample processor 420(1987)184

New method for the determination of doxorubicin, 4'-epidoxorubicin and all known metabolites in cardiac tissue 424(1988)103

HPLC determination of doxorubicin and its metabolites in plasma and tissue 425(1988)419

**Adriblastin**

HPLC determination of adriblastin in human plasma, urine, saliva and liver punctuate by column switching for drug monitoring studies 424(1988)182

**Adsorbents**

Carbon black adsorbents modified with *n*-hydrocarbons and treated with an high-frequency low-temperature plasma 402(1987)87

Characterization of porous polymers by polar strength and selectivity 404(1987)145

GC study of microporous carbons 406(1987)139

New metal chelate adsorbent selective for proteins and peptides containing neighbouring histidine residues 411(1987)177

Steric protection of a modified glass surface with adsorbed polyacrylamide: interactions with blood proteins 428(1988)17

Use of modified tungsten selenide as an adsorbent for GC 446(1988)79

**Adsorption**

GC study of microporous carbons 406(1987)139

Adsorption of binary solvent mixtures in reversed-phase chromatographic systems. Surface excess isotherms and enthalpies of displacement 406(1987)257

Adsorption chromatography on cellulose. I. Salting-out chromatography of organic compounds 406(1987)389

Effect of zirconia on the sorption properties of silica gels 407(1987)189

Solubility properties in polymers and biological media. II. A new method for the characterisation of the adsorption of gases and vapours on solids 409(1987)15

Characterization of adsorption on the stationary phase using high-performance immunoaffinity chromatography 423(1987)41

Non-linear elution effects in split-peak chromatography. I. Computer simulations for the cases of irreversible diffusion- and adsorption-limited kinetics 436(1988)111

Adsorption behaviour of tris( $\beta$ -diketonato)metal(III) complexes on silica gel 438(1988)401

Adsorption chromatography on cellulose. II. Separations of aromatic amino acids, biogenic amines, alkaloids, dyes and phenols and determination of hydrophobic constants 440(1988)165

GC study of the temperature dependence of the hydrogen bond energy of adsorption on silica 446(1988)55

Adsorption and chromatographic properties of modified silica sorbents for the production of viral preparations 446(1988)211

Split peaks in non-linear chromatography and their effect on sample throughput in large scale separations 449(1988)53

**Adsorption capacity**

Interaction of proteins with immobilized  $\text{Cu}^{2+}$ . Quantitation of adsorption capacity, adsorption isotherms and equilibrium constants by frontal analysis 403(1987)197

**Adsorption-desorption**

Efficiency of adsorption-desorption of volatile compounds. Application to isoprene on Tenax GC 437(1988)139

**Adsorption isotherms**

Interaction of proteins with immobilized  $\text{Cu}^{2+}$ . Quantitation of adsorption capacity, adsorption isotherms and equilibrium constants by frontal analysis 403(1987)197

Simplified description of HPLC separation under overload conditions, based on the Craig distribution model. III. Computer simulations for two co-eluting bands assuming a Langmuir isotherm 405(1987)1

Adsorption of binary solvent mixtures in reversed-phase chromatographic systems. Surface

excess isotherms and enthalpies of displacement 406(1987)257

Adsorption isotherms on HPLC sorbents. I. Peptides and nucleic acid constituents on octadecyl-silica 406(1987)275

Adsorption isotherms on HPLC sorbents. II. Proteins on cation exchangers with silica support 406(1987)285

Further comments on the preparation of mixed organic layers chemically bonded on silicon dioxide. Adsorption of modifiers from strongly adsorbed solvents 406(1987)295

Investigation of HPLC packings by liquid mixture adsorption 406(1987)311

Determination of adsorption isotherms from chromatographic measurements, using the peak maxima method 408(1987)1

Free energy consumption at self-sharpening concentration fronts in isothermal fixed-bed adsorption. Theory for isotherms of arbitrary shape and applications in displacement chromatography 409(1987)1

Adsorption isotherms of ethyl acetate modifier on silica from supercritical carbon dioxide 409(1987)55

Improved method for determining adsorption isotherms by LC on long and short fibrous material 436(1988)391

Reversibility of solvent-induced surface modification of alumina adsorbents 442(1988)105

Preparative chromatography of proteins. Analysis of the multivalent ion-exchange formalism 443(1988)13

Molecular orientation of immunoglobulin G at high concentration on an ion-exchange sorbent 443(1988)119

#### **Affinity adsorbents**

Development of N-hydroxysuccinimide ester silica, a novel support for high-performance affinity chromatography 405(1987)179

Matrix evaluation for preparative high-performance affinity chromatography 407(1987)179

Affinity chromatography of human thrombin on modified silica 424(1988)385

Selective purification of cardiac myosin by a high-performance salicylate affinity column 435(1988)185

Analytical high-performance affinity chromatography of ovalbumin-derived glycopeptides on columns of concanavalin A- and wheat germ agglutinin-immobilized gels 438(1988)73

Affinity chromatography on a hydrophobic matrix using a heterobifunctional ligand 438(1988)339

High-performance affinity chromatography of proteins on TSKgel Heparin-5PW 445(1988)234

Modified general affinity adsorbent for large-scale purification of penicillinases 448(1988)109

#### **Affinity partitioning**

Effect of polymer structure on affinity partitioning of lactate dehydrogenase in polymer-water two-phase systems 411(1987)161

#### **Aflatoxin B<sub>1</sub>**

HPLC procedure for determining the profiles of aflatoxin precursors in wildtype and mutant strains of *Aspergillus parasiticus* 441(1988)400

#### **Aflatoxins**

Standardized HPLC of 182 mycotoxins and other fungal metabolites based on alkylphenone retention indices and UV-VIS spectra (diode array detection) 404(1987)195

Rapid and highly sensitive analysis of aflatoxin M<sub>1</sub> in liquid and powdered milks using an affinity column cleanup 407(1987)393

Determination of aflatoxins in peanut products using disposable bonded-phase columns and post-column reaction detection 409(1987)413

Quantitation of aflatoxins from various strains of *Aspergillus* in foodstuffs 435(1988)229

HPLC procedure for determining the profiles of aflatoxin precursors in wildtype and mutant strains of *Aspergillus parasiticus* 441(1988)400

Determination of aflatoxins by capillary column GC 447(1988)410

Fluorimetric determination of aflatoxins in foodstuffs by HPLC with flow injection analysis 448(1988)173

Large-scale purification of the mycotoxins aflatoxin B<sub>1</sub>, B<sub>2</sub> and G<sub>1</sub> 450(1988)101

#### **Agarose gel electrophoresis**

Agarose gel electrophoresis of bacteriophages and related particles (Review) 418(1987)345

**Aggregated species**

High-performance SEC of recombinant derived proteins and aggregated species 436(1988)289

**Agmatine**

Determination of agmatine, arginine, citrulline and ornithine by reversed-phase LC using automated pre-column derivatization with *o*-phthalaldehyde 425(1988)269

Quantitation of polyamines using TLC and image analysis 441(1988)361

Detection by HPLC of polyamines formed by clostridal putrefaction of caseins 442(1988)267

**Air samplers**

Technique for calibrating air samplers and certain other analytical devices 438(1988)100

**Ajmalin**

Chromatographic isolation of primary and secondary metabolites from plant cell cultures 440(1988)53

**D-Alanine**

Determination of environmental levels of peptidoglycan and lipopolysaccharide using GC with negative-ion chemical-ionization MS utilizing bacterial amino acids and hydroxy fatty acids as biomarkers 431(1988)1

**Alanine aminopeptidase**

Biochemical indices of renal damage: separation of urinary alanine aminopeptidase by LC 421(1987)344

**Albumin**

Recovery of polypeptides after reversed-phase HPLC 408(1987)191

Rapid preparation of bovine mercaptalbumin by means of covalent chromatography on silica-based materials 408(1987)263

Influence of pore and particle size on the frontal uptake of proteins. Implications for preparative anion-exchange chromatography 409(1987)111

Measurement of glycated albumin in diabetic patients by biospecific affinity chromatography 419(1987)75

Purification of commercial human albumin on immobilized IDA-Ni<sup>2+</sup> 421(1987)141

Procedure for the isolation of a large peptic fragment of human serum albumin 424(1988)13

Steric protection of a modified glass surface with

adsorbed polyacrylamide: interactions with blood proteins 428(1988)17

HPLC of complex mixtures of cyanogen bromide-produced peptides from different proteins 443(1988)317

Preparative chiral separation in an aqueous two-phase system by a few counter-current extractions 450(1988)277

**Albumin-coated silica gel**

Micro-LC and the chiral recognition mechanism on albumin-coated silica gel. Large selectivity changes with sample size 447(1988)297

**Albumins**

Interaction of proteins with immobilized Cu<sup>2+</sup>. Quantitation of adsorption capacity, adsorption isotherms and equilibrium constants by frontal analysis 403(1987)197

**Albuterol**

Determination of albuterol in human serum by reversed-phase HPLC with electrochemical detection 427(1988)188

**Alcohols**

HPLC separation of enantiomers on axially chiral binaphthalene derivatives bonded to silica gel 403(1987)275

Potential solid supports for GC prepared on the basis of certain rocks 406(1987)145

Comparison of volatile components between fresh and rotten mussels by GC-MS 409(1987)251

Ion-exclusion chromatography of non-ionic substances with conductivity detection 409(1987)271

Glass capillary GC analysis of volatile components of canned Philippine pineapple juice 409(1987)409

Organo-modified smectite for the GC separation of various compounds 410(1987)121

Resolution of enantiomeric aromatic alcohols on a cellulose tribenzoate HPLC chiral stationary phase. A proposed chiral recognition mechanism 411(1987)139

Identification by combined GC-MS of constituent long-chain fatty acids and alcohols from the meibomian glands of the rat and a comparison with human meibomian lipids 414(1987)253

Studies of metabolites in diarrheal stool specimens positive for *Klebsiella*, *Serratia*, and *Proteus* spp. by frequency-pulsed electron-capture GC 430(1988)209

Secretion of proteins by *Bacillus subtilis* 168 grown in the presence of membrane active agents (alcohols) 440(1988)379

Determination of alcohols in alcoholic beverages by micro HPLC with indirect photometric detection 445(1988)139

Anthracene-9-carbonyl chloride as a fluorescence and UV derivatising reagent for the HPLC analysis of hydroxy compounds 445(1988)393

Polarity of oligoxyethylene derivatives of alcohols, thio alcohols and alkylamines as measured by GC 450(1988)291

#### **Aldehydes**

Retention behaviour of aromatic aldehyde derivatives in reversed-phase LC 404(1987)242

Comparison of volatile components between fresh and rotten mussels by GC-MS 409(1987)251

HPLC determination of glyoxylic acid and other carbonyl compounds in urine 417(1987)253

O-(2,3,4,5,6-Pentafluorobenzyl)oxime-trimethylsilyl ester derivatives for quantitative GC and GC-MS studies of aldehydes, ketones and oxoacids 421(1987)336

Multiple element emission spectral detection GC profiles of halogenated products from chlorination of humic acid and drinking water 438(1988)35

#### **Alditol acetates, partially methylated**

Separation of partially methylated alditol acetates on SP-2330 and HP-1 vitreous silica capillary columns 445(1988)424

#### **Aldolase**

Chromatographic isolation of primary and secondary metabolites from plant cell cultures 440(1988)53

#### **Aldoses**

Simultaneous GLC determination of aldoses and alduronic acids 408(1987)245

#### **Alduronic acids**

Simultaneous GLC determination of aldoses and alduronic acids 408(1987)245

#### **Alfentanil**

Determination of fentanyl and alfentanil in

plasma by HPLC with UV detection 419(1987)464

Simultaneous determination of fentanyl and alfentanil in rat tissues by capillary column GC 433(1988)95

#### **Alimemazine**

Indirect HPLC resolution of racemic tertiary amines as their diastereomeric urea derivatives after N-dealkylation 436(1988)269

#### **Alizapride**

LC analysis of alizapride and metoclopramide in human plasma and urine using solid-phase extraction 419(1987)233

#### **Alkali fusion**

Use of acid or alkaline fusion for the quantitative degradation of condensation polymers 406(1987)167

#### **Alkaline phosphatase**

Sugar-chain heterogeneity of human alkaline phosphatases: differences between normal and tumour-associated isozymes 413(1987)65

Automated separation and measurement of urinary isoenzymes and protein by ion-exchange LC 414(1987)47

Highly sensitive assay for alkaline and acid phosphatase activities by HPLC with electrochemical detection 417(1987)41

Blood appearance of rat alkaline phosphatase originating from the duodenum in vitro 420(1987)275

Assay of 5'-nucleotidase and simultaneous detection of interfering enzymes by anion-exchange CC 420(1987)398

Alkaline phosphatase. Laboratory and clinical implications (Review) 429(1988)419

HPLC determination of acid and alkaline phosphatases in urine 434(1988)196

Conformational studies of bovine alkaline phosphatase in hydrophobic interaction and size-exclusion chromatography with linear diode array and low-angle laser light scattering detection 442(1988)29

#### **Alkaloids**

Efficient isolation of insect growth inhibitory macrolide alkaloids using recycle high-performance GPC 402(1987)354

HPLC analysis of C<sub>19</sub>-diterpenoid alkaloids 402(1987)366

- Standardized HPLC of 182 mycotoxins and other fungal metabolites based on alkylphenone retention indices and UV-VIS spectra (diode array detection) 404(1987)195
- Assay of five corynoline-type alkaloids in *Corydalis bungeana* by reversed-phase HPLC 408(1987)426
- Fluorodensitometric assay of 6*H*-pyrido[4,3-*b*]-5,11-dimethylcarbazoles (ellipticine and derivatives) biosynthesized by *Ochrosia elliptica* cultures *in vitro* 409(1987)426
- Determination of *Catharanthus* alkaloids by reversed-phase HPLC 410(1987)488
- Determination of physostigmine in plasma by HPLC 415(1987)423
- Metabolism of physostigmine in mouse liver microsomal incubations studied by LC with dual-electrode amperometric detection 419(1987)165
- Analysis of vinca alkaloids in plasma and urine using HPLC with electrochemical detection 424(1988)83
- Ammonium positive-ion and hydroxide negative-ion chemical ionization GC-MS for the identification of pyrrolizidine alkaloids in *Eupatorium rotundifolium* L. var. *ovatum* 428(1988)352
- Determination of vinca alkaloids in plasma and urine using ion-exchange chromatography on silica gel and fluorescence detection 434(1988)145
- Separations of flavonoids and alkaloids in medicinal herbs by high-speed counter-current chromatography 435(1988)159
- Separation of eight alkaloids and meconic acid and quantitation of five principal alkaloids in gum opium by gradient reversed-phase HPLC 436(1988)455
- Centrifugal TLC of alkaloids from vegetable sources using an aluminium oxide layer 437(1988)328
- Separation of opium alkaloids by carbon dioxide sub- and supercritical fluid chromatography with packed columns. Application to the quantitative analysis of poppy straw extracts 437(1988)351
- Modification of retention of some alkaloids in the system silanized silica/methanol + water + di(2-ethylhexyl)orthophosphoric acid 439(1988)257
- Chromatographic isolation of primary and secondary metabolites from plant cell cultures 440(1988)53
- Adsorption chromatography on cellulose. II. Separations of aromatic amino acids, biogenic amines, alkaloids, dyes and phenols and determination of hydrophobic constants 440(1988)165
- Capillary GC of steroidal alkaloids from Solanaceae. Retention indices and simultaneous flame ionization/nitrogen-specific detection 442(1988)133
- Determination of alkaloids of *Fumaria parviflora* and *Fumaria capreolata* by HPLC and capillary isotachopheresis 445(1988)258
- HPLC determination of avenanthramides, N-aroylanthranilic acid alkaloids from oats 445(1988)363
- HPLC identification of simple  $\beta$ -carboline alkaloids in specimens of Heliconiini butterflies 447(1988)432
- Alkanes**
- Flow programming system for process capillary GC 405(1987)163
- GC study of microporous carbons 406(1987)139
- Super- and subcritical fluid chromatography-Fourier transform IR spectrometry of medium polar compounds on polymeric styrene-divinylbenzene packed columns 407(1987)87
- High-resolution GC with liquid crystal glass capillaries. X. Separation of diastereomeric alkanes 436(1988)47
- Paraffins, olefins, naphthenes and aromatics analysis of selected hydrocarbon distillates using on-line column switching HPLC with dielectric constant detection 437(1988)365
- GLC analysis of isophorone byproducts 446(1988)23
- Trapping efficiency of capillary cold traps 448(1988)117
- GC on porous polymers. IV. Influence of the geometric structure of porous copolymers of 1,4-di(methacryloyloxymethyl)naphthalene with 1,4-divinylbenzene on their chromatographic behaviour 448(1988)233
- Alkanethiols**
- Extraction chromatography of alkanethiols 449(1988)306

**Alkanoic acids**

Influence of additives to the eluent on hydrophobic interaction chromatography of simple compounds. V. Influence of pH and electrolytes on the retention of *n*-alkylammonium ions and *n*-alkyl carboxylates 438(1988)181

GC on porous polymers. IV. Influence of the geometric structure of porous copolymers of 1,4-di(methacryloyloxymethyl)naphthalene with 1,4-divinylbenzene on their chromatographic behaviour 448(1988)233

**Alkanolamine antihistamines**

Screening procedure for the detection of alkanolamine antihistamines and their metabolites in urine using computerized GC-MS 428(1988)43

**Alkanols**

Influence of additives to the eluent on the hydrophobic interaction chromatography of simple compounds. IV. Influence of electrolytes on the retention of normal alkanols at 25 °C 437(1988)337

GLC analysis of isophorone byproducts 446(1988)23

GC on porous polymers. IV. Influence of the geometric structure of porous copolymers of 1,4-di(methacryloyloxymethyl)naphthalene with 1,4-divinylbenzene on their chromatographic behaviour 448(1988)233

**Alkenes**

Retention and thermodynamics of solution of *n*-alkenes in OV-101 406(1987)75

**7-Alkoxy-4-alkylcoumarins**

Quantitative structure-activity relationship of coumarin derivatives. Involvement of partition between aqueous and membrane phase 445(1988)49

**7-Alkoxy coumarins**

Quantitative structure-activity relationship of coumarin derivatives. Involvement of partition between aqueous and membrane phase 445(1988)49

**Alkyd components**

Cleavage of urea-formaldehyde resin in cross-linked alkyd systems 403(1987)145

**1-O-Alkyl-2-acetyl-*sn*-glycero-3-phosphocholine**

Highly sensitive GC-MS method for the

determination of platelet-activating factor in human blood 433(1988)243

**Alkylamine**

Separation of isomeric mono-, di- and tributylphenols by HPLC on octadecylsilica with methanol-water using alkyl nitrile and alkylamine as surface modifiers 442(1988)53

**Alkylamine antihistamines**

Identification and differentiation of alkylamine antihistamines and their metabolites in urine by computerized GC-MS 430(1988)31

**Alkylamines**

Analysis of amphoteric surfactants of the alkylaminopropylglycine type by GC 405(1987)305

Study of the derivatization of *n*-alkylamines with 1-fluoro-2,4-dinitrobenzene in the presence of aqueous cetyltrimethylammonium bromide micelles 426(1988)267

Influence of additives to the eluent on hydrophobic interaction chromatography of simple compounds. V. Influence of pH and electrolytes on the retention of *n*-alkylammonium ions and *n*-alkyl carboxylates 438(1988)181

Polarity of oligooxyethylene derivatives of alcohols, thio alcohols and alkylamines as measured by GC 450(1988)291

**N-Alkylaminopropylamines**

Analysis of amphoteric surfactants of the alkylaminopropylglycine type by GC 405(1987)305

**N-Alkylaminopropylglycines**

Analysis of amphoteric surfactants of the alkylaminopropylglycine type by GC 405(1987)305

**Alkylammonium nitrate salts**

Solvent properties of liquid organic salts used as mobile phases in microcolumn reversed-phase LC 411(1987)61

**Alkylammonium salts**

Chemical interactions as a possible limitation on the useful solvent properties of liquid alkylammonium salts 435(1988)17

**N-Alkylanilines**

Effect of organic modifier on the separation and efficiencies of homologues on porous polymer columns 407(1987)19

**Alkyl aryl ketones**

Effect of organic modifier on the separation and efficiencies of homologues on porous polymer columns 407(1987)19

**O-Alkylated glycerol derivatives**

Enantioselective capillary GC of 1,2-isopropylidene glycerol and O-alkylated glycerol derivatives 410(1987)458

**Alkylated guanines**

Separation characteristics of alkylated guanines in HPLC 447(1988)81

**Alkylation**

Effects of reductive alkylation on peptide retention times 443(1988)247

**Alkylbenzenes**

Effect of organic modifier on the separation and efficiencies of homologues on porous polymer columns 407(1987)19

HPLC of alkylbenzenes on alumina 437(1988)177

Identification of alkylbenzenes by capillary GC and combined GC-MS (Review) 438(1988)131

GC study of the temperature dependence of the hydrogen bond energy of adsorption on silica 446(1988)55

Retention relationships for aromatic hydrocarbons eluted from capped and uncapped octadecyl silica gels 447(1988)13

**Alkylbenzenesulphonates**

Determination of linear alkylbenzenesulphonates, alkylphenol polyethoxylates and nonylphenol in waste water by HPLC after enrichment on octadecylsilica 403(1987)243

**2-Alkylbenzo[*b*]thiophenes**

Identification of homologous series of alkylated thiophenes, thiolanes, thianes and benzothiophenes present in pyrolysates of sulphur-rich kerogens 435(1988)435

***n*-Alkyl benzyl succinates**

GC-MS of C<sub>1</sub>-C<sub>10</sub> alkyl benzyl succinates 446(1988)103

**Alkyl chain length**

Role of the alkyl chain length on the ion interaction reagent, flow-rate, column packing and detection in ion interaction reversed-phase HPLC in separations of anions using amine salicylates 449(1988)103

**Alkyl chains**

Influence of bonding chemistry on the reordering/resolution of silica immobilized alkyl chains 449(1988)115

**2-Alkyl-5-ethylthiophenes**

Identification of homologous series of alkylated thiophenes, thiolanes, thianes and benzothiophenes present in pyrolysates of sulphur-rich kerogens 435(1988)435

**Alkylindenes**

Separation of alkyl-substituted indene enantiomers by chromatography on microcrystalline triacetylcellulose 405(1987)155

**Alkyl isothiocyanates**

Determination of N-acetyl-S-(N-alkylthiocarbamoyl)-L-cysteine, a principal metabolite of alkyl isothiocyanates, in rat urine 414(1987)19

**2-Alkyl-4-methylbenzo[*b*]thiophenes**

Identification of homologous series of alkylated thiophenes, thiolanes, thianes and benzothiophenes present in pyrolysates of sulphur-rich kerogens 435(1988)435

Identification of homologous series of alkylated thiophenes, thiolanes, thianes and benzothiophenes present in pyrolysates of sulphur-rich kerogens 435(1988)435

**Alkyl methylphosphonic acids**

Analysis of alkyl methylphosphonic acids in aqueous matrices by ion-pair reversed-phase ion chromatography 407(1987)399

**2-Alkyl-5-methylthiolanes**

Identification of homologous series of alkylated thiophenes, thiolanes, thianes and benzothiophenes present in pyrolysates of sulphur-rich kerogens 435(1988)435

**2-Alkyl-5-methylthiophenes**

Identification of homologous series of alkylated thiophenes, thiolanes, thianes and benzothiophenes present in pyrolysates of sulphur-rich kerogens 435(1988)435

**Alkyl nitrile**

Separation of isomeric mono-, di- and tributylphenols by HPLC on octadecylsilica with methanol-water using alkyl nitrile and alkylamine as surface modifiers 442(1988)53



**Alkylphenol polyethoxylates**

Determination of linear alkylbenzenesulphonates, alkylphenol polyethoxylates and nonylphenol in waste water by HPLC after enrichment on octadecylsilica 403(1987)243

**Alkylphenols**

Structure-retention correlations of alkylphenols in GLC and GSC 446(1988)71

**Alkylphenoxyalkanoic acids**

Separation efficiency of adsorption and partition TLC and reversed-phase HPTLC with chemically bonded phases for alkylphenoxyalkanoic acids 410(1987)53

**O-Alkylserines**

GC and MS behaviour of O-alkylserines as their N-trifluoroacetyl butyl esters 411(1987)259

**2-Alkylthianes**

Identification of homologous series of alkylated thiophenes, thiolanes, thianes and benzothiophenes present in pyrolysates of sulphur-rich kerogens 435(1988)435

**2-Alkylthiolanes**

Identification of homologous series of alkylated thiophenes, thiolanes, thianes and benzothiophenes present in pyrolysates of sulphur-rich kerogens 435(1988)435

**2-Alkylthiophenes**

Identification of homologous series of alkylated thiophenes, thiolanes, thianes and benzothiophenes present in pyrolysates of sulphur-rich kerogens 435(1988)435

**Allergens**

Purification of allergens by HPLC. IV.

Purification of the allergen of olive pollen (*Olea europea*) 403(1987)312

Purification of allergens by HPLC. V.

Purification of cat and dog epithelial allergens 404(1987)378

Purification of allergens by HPLC. VI.

Purification of the major allergen from *Parietaria judaica* pollen 407(1987)408

Purification of allergens by HPLC. VII.

Purification of the major allergen of birch pollen (*Betula verrucosa*) 408(1987)420

Analysis of allergenic components of a *Parietaria judaica* pollen extract by chromatographic methods for the evaluation of purification procedures 444(1988)209

Micropreparative-scale enrichment of some allergenic components of *Parietaria* pollen extract 446(1988)179

**Alloxanthin**

HPLC of carotenoids from some marine shellfish 440(1988)37

**Allura Red AC**

Simple method for the analysis of food dyes on reversed-phase thin-layer plates 411(1987)437

**Alpha<sub>1</sub>-beta glycoprotein**

Purification of human alpha<sub>1</sub>-beta glycoprotein and study of its microheterogeneity and molecular variants 405(1987)337

**Alphaprodine**

Measurement of alphaprodine by selected-ion monitoring 426(1988)392

**Alpidem**

Determination of alpidem and its metabolites in human plasma by HPLC and fluorimetric detection 414(1987)101

Correlation between reversed-phase HPLC and plasma protein binding 421(1987)83

**Alprazolam**

Alprazolam,  $\alpha$ -hydroxy- and 4-hydroxyalprazolam analysis in plasma by HPLC 430(1988)180

**Alprenolol**

Derivatization of 2-amino alcohols with phosgene in aqueous media: limitations of the reaction selectivity as found in the presence of O-glucuronides of alprenolol in urine 413(1987)270

Determination of alprenolol and metoprolol in plasma by column LC 434(1988)265

**Alternariol**

Standardized HPLC of 182 mycotoxins and other fungal metabolites based on alkylphenone retention indices and UV-VIS spectra (diode array detection) 404(1987)195

**Alumina**

Determination of catecholamines in urine (and plasma) by LC after on-line sample pretreatment on small alumina or dihydroxyborylsilica columns 414(1987)285

Separation of organochlorines on alumina 437(1988)254

Reversibility of solvent-induced surface modification of alumina adsorbents 442(1988)105

**Aluminium**

Formation of silicato complexes of aluminium in aqueous solution 403(1987)151

Isotachophoretic determination of the cationic constituents in mixtures of raw materials for glass manufacturing 439(1988)423

Comparison of HPLC and atomic spectrometric methods for the determination of Fe(III) and Al(III) in soil and clay samples 449(1988)241

**Aluminium chloride**

Preparation of fatty acid methyl esters by direct transesterification of lipids with aluminium chloride-methanol 441(1988)99

**Aluminosilicates**

Formation of silicato complexes of aluminium in aqueous solution 403(1987)151

 **$\alpha$ -Amanitin**

HPLC method for the determination of  $\alpha$ -amanitin and phalloidin in human plasma using the column-switching technique and its application in suspected cases of poisoning by the green species of amanita mushroom (*Amanita phalloides*) 425(1988)121

**Amaranth**

Simple method for the analysis of food dyes on reversed-phase thin-layer plates 411(1987)437

**Ambenonium**

Determination of ambenonium in biological samples by reversed-phase ion-pair LC 433(1988)167

**Ambenonium chloride**

Determination of ambenonium chloride in serum by reversed-phase ion-pair LC 421(1987)407

**Ambroxol**

Assay of ambroxol in biological fluids by capillary GLC 414(1987)65

Rapid and sensitive determination of ambroxol in human plasma and urine by HPLC 421(1987)211

**Amides**

Resolution of racemic amides and phosphine oxides on a  $\beta$ -cyclodextrin-bonded stationary phase by subcritical fluid chromatography 405(1987)135

Prediction of HPLC capacity factors following column calibration. Influence of different

solvents and columns on the separation of a series of aromatic amides 437(1988)13

**Amidines**

Determination of amidines by reversed-phase ion-pair HPLC 439(1988)476

**6-Amidino-2-naphthyl [4-(4,5-dihydro-1H-imidazol-2-yl)amino]benzoate dimethanesulphonate**

HPLC determination of 6-amidino-2-naphthyl [4-(4,5-dihydro-1H-imidazol-2-yl)amino]benzoate dimethanesulphonate and its metabolites in biological fluids 433(1988)177

**Amiloride**

Simultaneous determination of amiloride and hydrochlorothiazide in plasma by reversed-phase HPLC 423(1987)351

Simple HPLC method for the measurement of amiloride in body fluids 428(1988)123

**Amine hardeners**

Quantitative determination of free primary amines in polymeric amine hardeners. HPLC separation of ethylenediamine and triethylenetetramine after derivatization with salicylaldehyde-diphenylboron chelate and separation of 4,4'-diaminodiphenylmethane by HPLC 436(1988)447

**Amines**

Separation of racemic pharmaceuticals by HPLC on silica gel modified with carbohydrate residues 403(1987)263

HPLC separation of enantiomers on axially chiral binaphthalene derivatives bonded to silica gel 403(1987)275

Inductive adsorption as a new approach to amine enantiomer resolution 404(1987)87

Cryogenic sampling for aniline vapor with subsequent LC-electrochemical detection analysis 408(1987)81

Simultaneous determination of  $\gamma$ -aminobutyric acid and polyamines by HPLC 419(1987)285

Procedure for the determination of amines in small urine samples 420(1987)379

Studies of metabolites in diarrheal stool specimens positive for *Klebsiella*, *Serratia*, and *Proteus* spp. by frequency-pulsed electron-capture GC 430(1988)209

Effects of long-term storage on the concentrations of the unconjugated acidic metabolites of the trace amines, indoleamines and catecholamines 433(1988)23

Detection of aminophenols and aromatic amines on thin-layer plates containing nitrite 435(1988)256

Indirect HPLC resolution of racemic tertiary amines as their diastereomeric urea derivatives after N-dealkylation 436(1988)269

Quantitative determination of free primary amines in polymeric amine hardeners. HPLC separation of ethylenediamine and triethylenetetramine after derivatization with salicylaldehyde–diphenylboron chelate and separation of 4,4'-diaminodiphenylmethane by HPLC 436(1988)447

Quantitative determination of free primary amines in polymeric amine hardeners. HPLC separation of ethylenediamine and triethylenetetramine after derivatization with salicylaldehyde–diphenylboron chelate and separation of 4,4'-diaminodiphenylmethane by HPLC 436(1988)447

TLC of amine antioxidants and antiozonants used in elastomers 437(1988)59

HPLC of biogenic amines, derivatized with 9-fluorenylmethyl chloroformate 437(1988)379

Analytical separations of enantiomeric primary amines as pyrrole derivatives formed by reaction with polygodial 438(1988)108

Simple permeation absorber for sampling and preconcentrating hazardous air contaminants 439(1988)283

Adsorption chromatography on cellulose. II. Separations of aromatic amino acids, biogenic amines, alkaloids, dyes and phenols and determination of hydrophobic constants 440(1988)165

Determination of aniline derivatives by HPLC with fluorescence detection 445(1988)273

#### **Amino acid–amino acid interactions**

Amino acid–amino acid interactions studied by charge-transfer chromatography 434(1988)455

#### **Amino acid analyzer**

Amino acid analysis: a fall-off in performance 408(1987)291

Amino acid analysis. II. Nuances of integration 410(1987)176

Construction of an ion-exchange amino acid analyser kit for use with HPLC apparatus 431(1988)285

#### **Amino acid esters**

Separation and determination of some amino acid ester enantiomers by TLC after derivatization with (*S*)-(+)-naproxen 450(1988)281

#### **Amino acids**

Chiral separations of amino acids using di- and tri-peptides for zwitter-ion pair chromatography 402(1987)49

Amino acid analysis: the histidine hangover 403(1987)271

HPLC separation of enantiomers on axially chiral binaphthalene derivatives bonded to silica gel 403(1987)275

Paper electrophoretic study of ion-pair formation. XVI. Resolution of optical isomers of some cobalt(III) complexes with amino acid ligands 403(1987)388

Computer analysis of automated Edman degradation and amino acid analysis data 404(1987)41

Amino acid analysis. I. Nuances of integration 404(1987)321

Chromatographic separation of racemic amino acids by use of chiral crown ether-coated reversed-phase packings 405(1987)145

Effects of pretreatment on the enantioselectivity of silica-bound proteins used as HPLC stationary phases 408(1987)99

Amino acid analysis. II. Nuances of integration 410(1987)176

Quantitative HPLC determination of the amino acid spinacine in blood and chow of rats 410(1987)504

Preparative separation of an enzymatic hydrolysate of <sup>14</sup>C-labelled proteins 411(1987)335

HPLC separation of peptide and amino acid stereoisomers 414(1987)313

GC–MS determination of tryptophan transaminase-catalyzed deuterium exchange 415(1987)104

Simultaneous LC determination of seventeen of the major monoamine neurotransmitters, precursors and metabolites. I. Optimization of

the mobile phase using factorial designs and a computer program to predict chromatograms 415(1987)261

Simultaneous LC determination of seventeen of the major monoamine neurotransmitters, precursors and metabolites. II. Assessment of human brain and cerebrospinal fluid concentrations 415(1987)275

Determination of free and total homocysteine in human plasma by HPLC with fluorescence detection 422(1987)43

HPLC determination of desmosine and isodesmosine in tissues and its application to studies of alteration of elastin induced by atherosclerosis 422(1987)53

Glutamate and glutamine in cerebrospinal fluid and serum from healthy volunteers – analytical aspects 424(1988)378

Analysis of amino acids in brain and plasma samples by sensitive GC-MS 425(1988)245

HPLC analysis of neurotransmitter amino acids in brain 428(1988)9

Determination of environmental levels of peptidoglycan and lipopolysaccharide using GC with negative-ion chemical-ionization MS utilizing bacterial amino acids and hydroxy fatty acids as biomarkers 431(1988)1

Amino acid analysis of plasma: studies in sample preparation 431(1988)163

Amino acid analysis: buffers and artifacts 436(1988)80

Detection of amino acids on thin-layer plates 436(1988)341

Amino acid analysis. III. Nuances of integration 436(1988)474

Adsorption chromatography on cellulose. II. Separations of aromatic amino acids, biogenic amines, alkaloids, dyes and phenols and determination of hydrophobic constants 440(1988)165

Stereoselective determination of L-amino acids using column LC with an enzymatic solid-phase reactor and chemiluminescence detection 440(1988)217

Chromatography, flow injection analysis and electrophoresis in computer-assisted comparative biochemistry: its application and possibilities in clinical research. Preliminary studies on Crohn's disease 440(1988)261

Copper(II) complexes of diamino-diamido-type ligands as chiral eluents in the enantiomeric separation of D,L-dansylamino acids by reversed-phase HPLC 441(1988)275

Application of Marfey's reagent in racemization studies of amino acids and peptides 444(1988)115

Chromatography of microbial metabolites of aromatic amino acids 446(1988)117

TLC enantiomeric resolution via ligand exchange 448(1988)11

Simplified protein hydrolysis with methanesulphonic acid at elevated temperature for the complete amino acid analysis of proteins 448(1988)404

#### **Amino acids, *p*-bromophenylcarbonyl derivatives**

Optical resolution of enantiomeric amino acid derivatives on a naphthylethylurea multiple-bonded chiral stationary phase prepared via an activated carbamate intermediate 404(1987)117

#### **Amino acids, *tert*-butyldimethylsilyl derivatives**

Capillary GC analysis of amino acids in blood and protein hydrolysates as *tert*-butyldimethylsilyl derivatives 416(1987)15

#### **Amino acids, butyl heptafluorobutyl derivatives**

Analysis of the amino acid and sugar composition of streptococcal cell walls by GC-MS 441(1988)323

#### **Amino acids, *N,N*-diethyl-2,4-dinitro-5-fluoroaniline derivatives**

Pre-column derivatization of amino acids with *N,N*-diethyl-2,4-dinitro-5-fluoroaniline and reversed-phase LC separation 433(1988)53

#### **Amino acid, dinitrophenyl derivatives**

Cross-axis synchronous flow-through coil planet centrifuge for large-scale preparative counter-current chromatography. II. Studies on partition efficiency in short coils and preparative separations with multilayer coils 449(1988)153

#### **Amino acids, dinitropyridyl derivatives**

HPTLC of dinitropyridyl-amino acids and sulphonamides on silanized silica gel 405(1987)394

#### **Amino acids, DNP derivatives**

Effect of polymer composition on the relative

hydrophobicity of the phases of the biphasic system aqueous dextran-poly(ethylene glycol) 403(1987)123

Rapid resolution of 2,4-dinitrophenyl-DL-amino acids on a native cellulose column 411(1987)502

Stepwise gradient elution using switching valves in micro HPLC 405(1987)117

Precolumn concentration and/or stepwise-gradient elution by switching valve techniques in micro HPLC 407(1987)141

Enantiomeric separation of amino acids by micro HPLC on an L-proline-bonded stationary phase 407(1987)151

#### **Amino acids, Dns derivatives**

Rapid and sensitive determination of Dns-amino acids in plasma using high-speed octadecyl LC columns 413(1987)25

Bis(L-amino acid amidato)copper(II) complexes as chiral eluents in the enantiomeric separation of D,L-dansylamino acids by reversed-phase HPLC 441(1988)287

Effect of the mobile phase composition and ligand structure on the separation of D- and L-dansylamino acids, as mixed metal complexes, by reversed-phase HPLC 445(1988)219

Use of complexing agents for selective separation in high-performance capillary electrophoresis. Chiral resolution via cyclodextrins incorporated within polyacrylamide gel columns 448(1988)41

#### **Amino acids, 9-fluorenylmethylchloroformate derivatives**

Determination of amino acids in biological, pharmaceutical, plant and food samples by automated precolumn derivatization and HPLC 431(1988)271

#### **Amino acids, isopropyl esters**

Solvent extraction clean-up for pre-treatment in amino acid analysis by GC. Application to age estimation from the D/L ratio of aspartic acid in human dentine 449(1988)165

#### **Amino acids, $\beta$ -naphthylamide-substituted**

Planar chromatographic separation of enantiomers and diastereomers with cyclodextrin mobile phase additives 448(1988)345

#### **Amino acids, naphthylcarbamyl derivatives**

Amino acid analysis by reversed-phase HPLC. Automatic pre-column derivatization with activated carbamate reagent 407(1987)273

#### **Amino acids, NBD derivatives**

Automatic amino acid analysis utilizing 4-fluoro-7-nitrobenzo-2-oxa-1,3-diazole 420(1987)141

#### **Amino acids, OPA derivatives**

Stepwise gradient elution using switching valves in micro HPLC 405(1987)117

Automated pre-column derivatization of amino acids with *o*-phthalaldehyde by a reagent sandwiching technique 408(1987)217

Improvement in the resolution of *o*-phthalaldehyde derivatized amino acids by applying gradient steepness optimization to five reversed-phase columns of different lengths and particle sizes 411(1987)121

Rapid and sensitive method for the determination of arterial-venous differences and leg efflux of 3-methylhistidine using ion-pair HPLC and post-column fluorescence derivatization 414(1987)174

Automated pre-column derivatization with *o*-phthalaldehyde for the determination of neurotransmitter amino acids using reversed-phase LC 417(1987)385

Separation strategies for *o*-phthalaldehyde-mercaptoethanol derivatives of amino acids for reversed-phase HPLC 419(1987)1

Determination of agmatine, arginine, citrulline and ornithine by reversed-phase LC using automated pre-column derivatization with *o*-phthalaldehyde 425(1988)269

Rapid HPLC method to measure plasma leucine: importance in the study of leucine kinetics *in vivo* 426(1988)370

Determination of amino acids in biological, pharmaceutical, plant and food samples by automated precolumn derivatization and HPLC 431(1988)271

Construction of an ion-exchange amino acid analyser kit for use with HPLC apparatus 431(1988)285

LC determination of taurine in whole blood, plasma and platelets 431(1988)400

Separation of  $\alpha$ -amino acid enantiomers by reversed-phase HPLC after derivatization with *o*-phthalaldehyde and a sodium salt of 1-thio- $\beta$ -D-glucose 434(1988)417

Resolution of  $\alpha$ -substituted amino acid enantiomers by HPLC after derivatization with a

chiral adduct of *o*-phthalaldehyde. Application to glutamic acid analogues 440(1988)209

Post-capillary fluorescence detection in capillary zone electrophoresis using *o*-phthalaldehyde 447(1988)117

#### **Amino acids, pentafluoropropionyl butyl esters**

Determination of 4-aminobutyric acid, aspartate, glutamate and glutamine and their <sup>13</sup>C stable-isotopic enrichment in brain tissue by GC-MS 414(1987)265

#### **Amino acids, N-pentafluoropropionyl isopropyl esters**

Polysiloxanes containing thermally stable chiral amide side-chains for capillary GC and supercritical fluid chromatography 405(1987)169

#### **Amino acids, N-pentafluoropropionyl *n*-propyl esters**

Assignment of the configurations of the amino acids in peptidic siderophores 438(1988)291

#### **Amino acids, phenylthiocarbamyl derivatives**

HPLC of phenylthiocarbamyl-amino acids. Application to carboxyl-terminal sequencing of proteins 409(1987)299

Narrow-bore HPLC of phenylthiocarbamyl amino acids and carboxypeptidase P digestion for protein C-terminal sequence analysis 447(1988)351

#### **Amino acids, PTC derivatives**

Increased yield of the phenylthiocarbamyl-amino acids tyrosine, cysteine and phenylalanine using diethylether-based liquid-liquid extraction 403(1987)299

Determination of hydroxyproline in tissue collagen hydrolysate by derivatization and isocratic reversed-phase HPLC 420(1987)394

#### **Amino acids, PTH derivatives**

High-sensitivity phenylthiohydantoin amino acid analysis on-line to a gas phase protein sequencer 408(1987)388

Instability of phenylthiohydantoin amino acids 435(1988)363

#### **Amino acids, TCB derivatives**

Determination of stable isotopically substituted histidine in human plasma by GC-MS 413(1987)1

#### **Amino acids, N-TFA *n*-butyl esters**

Use of chloroform as a solvent for the formation

of N-trifluoroacetyl-*n*-butylamino acid derivatives for GC analysis 436(1988)100

#### **Amino acids, N(O,S)-TFA isobutyl esters**

GC analysis of histidine: effect on other amino acids present 446(1988)231

#### **Amino acids, TMS esters**

Formation of N-carboxy compounds during trimethylsilylation of amino acids in the presence of carbon dioxide 409(1987)383

#### **Aminoacyl-tRNA synthetases**

Analytical strategy for determination of active site sequences in aminoacyl-tRNA synthetases 438(1988)347

#### **Amino alcohols**

Separation of racemic pharmaceuticals by HPLC on silica gel modified with carbohydrate residues 403(1987)263

Ligand exchange chromatography of amino alcohol enantiomers as Schiff bases 406(1987)343

Separation of  $\beta$ -amino alcohols as their dansyl derivatives by TLC on a polyamide layer sheet 408(1987)449

Derivatization of 2-amino alcohols with phosgene in aqueous media: limitations of the reaction selectivity as found in the presence of O-glucuronides of alprenolol in urine 413(1987)270

Resolution of enantiomeric drugs of some  $\beta$ -amino alcohols as their urea derivatives by HPLC on a chiral stationary phase 447(1988)208

Supercritical fluid chromatography with ion-pairing modifiers. Separation of enantiomeric 1,2-aminoalcohols as diastereomeric ion pairs 447(1988)287

#### **Aminoalkylindoles**

Sample solvent effects in an apparent chiral HPLC separation on  $\beta$ -cyclodextrin 448(1988)31

#### **Aminoalkyl phosphates**

GC analysis of aminoalkylphosphonic acids and aminoalkyl phosphates 436(1988)67

#### **Aminoalkylphosphonic acids**

GC analysis of aminoalkylphosphonic acids and aminoalkyl phosphates 436(1988)67

#### **Aminoantipyrine**

Investigation of an alkylamine modified and pH-controlled mobile phase for separation of pyrazolone derivatives by HPLC 409(1987)81

Simple HPLC method for simultaneous

determination of aminopyrine and its metabolites 421(1987)198

#### ***p*-Aminobenzoic acid**

Direct and simultaneous HPLC assay for the determination of *p*-aminobenzoic acid and its conjugates in human urine 426(1988)103

Reversed-phase HPLC assay for the determination of the *in vitro* acetylation of *p*-aminobenzoic acid by human whole blood 433(1988)292

Determination of water-soluble vitamins using HPLC and electrochemical or absorbance detection 447(1988)256

#### **6-Aminobenzof[a]pyrene**

Chromatographic and spectral investigations on the *in vivo* metabolites of 6-nitrobenzo[a]pyrene 450(1988)325

#### **4-Aminobenzoic acid hydrazide**

Determination of reducing and non-reducing carbohydrates in food products by LC with post-column catalytic hydrolysis and derivatization. Comparison with refractive index detection 402(1987)127

#### **3-Amino-1,2,4-benzotriazine-1,4-dioxide**

HPLC assay for the benzotriazine di-N-oxide (SR 4233) and its reduced metabolites in biological materials 430(1988)429

#### **N-(4-Aminobenzoyl)- $\gamma$ -oligo(L-glutamic acid)s**

Separation of N-(4-aminobenzoyl)- $\gamma$ -oligo(L-glutamic acid)s on a thin-layer plate and their quantitation using fluorescamine 442(1988)420

#### **4-Aminobiphenyl**

Determination of diphenylamine residues in apples, and 4-aminobiphenyl residues in diphenylamine, by HPLC and electrochemical detection 447(1988)421

#### **Aminobiphenyls**

HPLC with on-line radioactivity monitoring for metabolic studies of 2- and 4-[<sup>14</sup>C]aminobiphenyls 442(1988)394

#### **4-Aminobutyric acid**

Determination of 4-aminobutyric acid, aspartate, glutamate and glutamine and their <sup>13</sup>C stable-isotopic enrichment in brain tissue by GC-MS 414(1987)265

Simultaneous determination of  $\gamma$ -aminobutyric acid and polyamines by HPLC 419(1987)285

HPLC analysis of neurotransmitter amino acids in brain 428(1988)9

#### **Aminochloramphenicol**

HPLC determination of chloramphenicol and four analogues using reductive and oxidative electrochemical and UV detection 417(1987)111

#### **Aminodantrolene**

Determination of dantrolene and its reduced and oxidized metabolites in plasma by HPLC 430(1988)187

#### **2-Amino-2-deoxyglucose**

Separation of racemic pharmaceuticals by HPLC on silica gel modified with carbohydrate residues 403(1987)263

#### **1-Amino-1,3-dicarboxycyclohexane**

Resolution of  $\alpha$ -substituted amino acid enantiomers by HPLC after derivatization with a chiral adduct of *o*-phthalaldehyde. Application to glutamic acid analogues 440(1988)209

#### **1-Amino-1,3-dicarboxy-2-cyclohexene**

Resolution of  $\alpha$ -substituted amino acid enantiomers by HPLC after derivatization with a chiral adduct of *o*-phthalaldehyde. Application to glutamic acid analogues 440(1988)209

#### **1-Amino-1,3-dicarboxycyclopentane**

Resolution of  $\alpha$ -substituted amino acid enantiomers by HPLC after derivatization with a chiral adduct of *o*-phthalaldehyde. Application to glutamic acid analogues 440(1988)209

#### **Amino esters**

Chromatographic behaviour of diastereoisomers. IX. Application of a microcomputer program to the TLC separation of some diastereoisomers on silica 438(1988)219

#### **Aminoether alcohols**

Analysis and identification of some aminoether alcohols and their ethers 410(1987)447

#### **Aminoglycosides**

Mechanism of selectivity in ion-pair HPLC of aminoglycoside antibiotics using perfluorinated pairing ions 404(1987)53

Use of mixed perfluorinated ion-pairing agents as solvents in ion-pair HPLC for the preparative purification of aminoglycoside antibiotics 407(1987)109

Pre-column derivatization of sisomicin with *o*-phthalaldehyde- $\beta$ -mercaptopropionic acid and

its application to sensitive HPLC determination with fluorimetric detection 425(1988)143

#### ***p*-Aminohippuric acid**

Direct and simultaneous HPLC assay for the determination of *p*-aminobenzoic acid and its conjugates in human urine 426(1988)103

#### **Aminoisocyanates**

Trace analysis of airborne aromatic isocyanates and related aminoisocyanates and diamines using HPLC with UV and electrochemical detection 435(1988)469

#### **$\delta$ -Aminolevulinic acid**

Quantitative determination of porphyrins, their precursors and zinc protoporphyrin in whole blood and dried blood by HPLC with fluorimetric detection 417(1987)269

Fluorimetric determination of urinary  $\delta$ -aminolevulinic acid by HPLC and post-column derivatization 426(1988)365

#### **Aminolink reagent**

Preparation and chromatographic use of 5'-fluorescent-labelled DNA probes 444(1988)67

#### **(5*R*,6*S*)-2-Aminomethyl-6-[(1*R*)-hydroxyethyl]-2-penam-3-carboxylic acid**

HPLC determination of the penem antibiotic (5*R*,6*S*)-2-aminomethyl-6-[(1*R*)-hydroxyethyl]-2-penam-3-carboxylic acid in human plasma and urine 427(1988)269

#### **N-(4-Amino-1-oxo-1,2,5-thiadiazol-3-yl)-3-[3-[(4-methylpiperidinyl)methyl]phenoxy]-*n*-propylamine**

High-performance liquid chromatographic determination of a new H<sub>2</sub> blocker (ORF 17910) in plasma 423(1987)358

#### **Amino peptidase-P**

HPLC separation of peptides possessing a proline residue in the amino-terminal penultimate position, and their products generated by enzymatic hydrolysis 424(1988)129

#### **Aminophenols**

Detection of aminophenols and aromatic amines on thin-layer plates containing nitrite 435(1988)256

#### **3-(4'-Aminophenyl)-3-ethyl-1-methylpyrrolidine-2,5-dione**

HPLC determination of imide inhibitors of aromatase in biological samples 424(1988)163

#### **3-(4'-Aminophenyl)-3-ethylpyrrolidine-2,5-dione**

HPLC determination of imide inhibitors of aromatase in biological samples 424(1988)163

#### **3-Amino-5-[3-[4-(piperidinoindanyloxy)]propylamino]-1-methyl-1*H*-1,2,4-triazole**

Analysis of a new H<sub>2</sub> receptor antagonist, 3-amino-5-[3-[4-(1-piperidinoindanyloxy)]propylamino]-1-methyl-1*H*-1,2,4-triazole, in human plasma and urine by HPLC 428(1988)113

#### **2-Amino-1,3-propanediol**

Quantitative determination of 2-amino-1,3-propanediol and its impurities by capillary GC 449(1988)432

#### **Aminopyrine**

Simple HPLC method for simultaneous determination of aminopyrine and its metabolites 421(1987)198

#### **5-Aminosalicylic acid**

Simple and sensitive HPLC assay for 5-aminosalicylic acid and acetyl-5-aminosalicylic acid in serum 413(1987)300

HPLC assay of 5-aminosalicylic acid and its acetylated metabolite in biological fluids using electrochemical detection 425(1988)214

Stability of 5-aminosalicylic acid and 5-acetylaminosalicylic acid in plasma 432(1988)358

#### **21-Aminosteroids**

Irregular retention properties of 21-aminosteroid antioxidants in octylsilane bonded-phase HPLC 424(1988)285

HPLC determination of a 21-aminosteroid antioxidant in plasma 424(1988)293

#### **Amiodarone**

Rapid HPLC method for the determination of amiodarone and desethylamiodarone in human plasma and serum 415(1987)197

Determination of amiodarone and its metabolite N-desethylamiodarone in serum by HPLC—comparison of different extraction procedures 417(1987)465

New procedure for the HPLC determination of amiodarone and desethylamiodarone with solid-phase extraction of rat plasma and tissue samples 431(1988)455

#### **Amisulpride**

Determination of a new benzamide, amisulpride,



in human plasma by reversed-phase ion-pair HPLC 416(1987)414

#### **Amitriptyline**

Applications of column-switching technique in biopharmaceutical analysis. I. HPLC determination of amitriptyline and its metabolites in human plasma 416(1987)99

HPLC quantitation of amitriptyline and nortriptyline in dialysate from plasma or serum using on-line solid-phase extraction 432(1988)363

Effects of drug-protein binding on trace enrichment of drugs in blood plasma on short precolumns 439(1988)353

#### **Amlodipine**

Analysis of amlodipine in human plasma by GC 420(1987)178

#### **Ammonium phosphates**

Chromatographic method for kinetic study of heterogeneous reactions 446(1988)65

#### **Ammonium sulphate**

Adsorption chromatography on cellulose. I. Salting-out chromatography of organic compounds 406(1987)389

#### **Amobarbital**

HPLC determination of the diastereomers of 1-( $\beta$ -D-glucopyranosyl)amobarbital in urine 422(1987)309

#### **Amodiaquine**

Detection and determination of antimalarial drugs and their metabolites in body fluids (Review) 434(1988)1

#### **Amoeba antigens**

*Entamoeba histolytica* antigens as possible vaccinogens? A short review 440(1988)467

#### **Amopyroquine**

HPLC method for the determination of amopyroquine in biological fluids 414(1987)242

LC determination of amopyroquine in rabbit plasma and red blood cells 421(1987)192

#### **Amoxicillin**

LC determination of amoxicillin and its metabolites in human urine by postcolumn degradation with sodium hypochlorite 413(1987)219

#### **Amoxicilloic acid**

LC determination of amoxicillin and its

metabolites in human urine by postcolumn degradation with sodium hypochlorite 413(1987)219

#### **Amoxycillin**

Isotachophoretic analysis of amoxycillin and carboxymethylcysteine in pharmaceuticals 408(1987)441

#### **Amperometric detector**

Simultaneous conductivity and amperometric detector with a single cell for microbore LC 436(1988)413

Amperometric detector with a manganin electrode for microcolumn LC 442(1988)111

#### **Amphetamine**

Determination of  $\beta$ -phenylalkylamines in pharmaceutical preparations by analytical capillary isotachopheresis 405(1987)389

GC quantification of underivatized amphetamine in whole blood. Blood stored in plastic containers—a source of error 432(1988)290

#### **Ampholytes**

Ionic binding properties of carrier ampholytes 437(1988)147

#### **Amphotericin B**

HPLC determination of amphotericin B in human urine 417(1987)439

HPLC determination of amphotericin B in human serum 419(1987)401

Separation of amphotericin B components by combined TLC and HPLC 437(1988)221

#### **Ampicillin**

LC determination of penicillins by postcolumn degradation with sodium hypochlorite using an hollow-fibre membrane reactor 447(1988)365

#### **Amsacrine**

Determination of amsacrine in human nucleated hematopoietic cells 422(1987)175

#### **Amyl alcohols**

Rapid HPLC method for determination of ethanol and fusel oil in the alcoholic beverage industry 447(1988)443

#### **$\alpha$ -Amylase**

Purification of  $\alpha$ -amylase by industrial autofocusing 411(1987)486

LC assay of the relative activities of serum pancreatic and salivary  $\alpha$ -amylase using

reductively pyridylaminated maltopentaose as a fluorescent substrate 419(1987)51

Use of ion-exchange chromatography and HPLC to prepare a substrate for differential assay of human pancreatic and salivary  $\alpha$ -amylases in serum 420(1987)158

Purification of germinated barley  $\alpha$ -amylase isozymes and limit-dextrinase by chromatofocusing and affinity chromatography 441(1988)436

### **$\beta$ -Amylase**

Immunoaffinity purification of  $\beta$ -amylase from mustard (*Sinapis alba* L.) cotyledons 408(1987)385

### **Amylose**

Separation of starch components based on ligand-induced adsorption of amylose on cellulose 403(1987)373

### **Anabolics**

Identification and quantitation of trenbolone in bovine tissue by GC-MS 424(1988)219

### **Anagrelide**

Determination of anagrelide in plasma by capillary GC-MS 416(1987)357

### **Analysis time**

Microcolumn chromatofocusing. III. Effect of the analysis time on the resolution and the use of a semi-disposable microcolumn 438(1988)15

### **Androgen receptors**

Application of high-performance liquid chromatofocusing to the study of prostatic androgen receptors. Effect of stabilizing agents on the heterogeneity of receptor structure 403(1987)171

### **Androstenedione**

Automated direct assay system for the measurement of sex steroid hormones in serum using HPLC 426(1988)33

### **Anethole**

Determination of  $\beta$ -asarone, safrole, isosafrole and anethole in alcoholic drinks by HPLC 404(1987)273

Characterisation of volatile oil constituents with relatively long GC retention times on two stationary phases 449(1988)312

### **Angelates**

Considerations and remarks on the analysis of

*Anthemis nobilis* L. essential oil by capillary GC and "hyphenated" techniques 411(1987)237

### **Angiotensin converting enzyme inhibitor**

Determination of a new angiotensin converting enzyme inhibitor and its active metabolite in plasma and urine by GC-MS 419(1987)123

### **Angiotensin peptides**

HPLC-radioimmunoassay method for the measurement of angiotensin II peptides in human plasma 417(1987)27

### **Angiotensins**

Separation of protein hormones (Review) 418(1987)245

### **Anhydroerythromycin A**

TLC study of the metabolites of erythromycins in the Wistar rat 434(1988)177

### **Anhydromuramyl-peptidoglycan monomer**

Comparative susceptibility of a peptidoglycan monomer from *Brevibacterium divaricatum* and its anhydromuramyl analogue to hydrolysis with N-acetylmuramyl-L-alanine amidase. Isolation and characterization of anhydromuramyl-peptidoglycan monomer 440(1988)405

### **Anhydroxytetracycline**

Quantitative analysis of oxytetracycline and related substances by HPLC 405(1987)229

### **Anilides**

Determination of oxidation products of N-phenyllinoleamide: Spanish toxic oil syndrome studies 426(1988)83

### **Aniline**

Cryogenic sampling for aniline vapor with subsequent LC-electrochemical detection analysis 408(1987)81

Analysis of lung air from patients with bronchogenic carcinoma and controls using GC-MS 432(1988)1

Simple permeation absorber for sampling and preconcentrating hazardous air contaminants 439(1988)283

### **Anilines**

Determination of aniline derivatives by HPLC with fluorescence detection 445(1988)273

### **Anion exchangers**

Facile preparation of a non-porous strong anion-exchange column for proteins 411(1987)452

## Anions

Ion chromatography of inorganic anions on a dynamically modified polystyrene-divinylbenzene packing material and its application to anion screening by sequential UV absorbance and electrochemical detection 402(1987)115

Origin of system peaks in single-column ion chromatography of inorganic anions using high pH borate-gluconate buffers and conductivity detection 404(1987)67

Derivatization of inorganic anions with pentafluorobenzyl methanesulphonate for GC 404(1987)175

Contamination of aqueous eluents due to corrosion of stainless-steel chromatographic components 407(1987)133

Isotachophoretic determination of mobility and  $pK_a$  by means of computer simulation. VI. Mobility and  $pK_a$  of sixty anions in methanol 408(1987)27

Coordinative interactions in the separation of insulin and derivatives by HPLC 408(1987)43

Determination of oxidizable inorganic anions by HPLC with fluorescence detection and application to the determination of salivary nitrite and thiocyanate and serum thiocyanate 424(1988)29

Peak enhancement of carboxylic acids in ion-exclusion chromatography with a cation-exchange hollow-fibre suppressor using neutral and alkaline enhancers 435(1988)417

Determination of mono- and divalent cations and anions in small fog samples by ion chromatography 439(1988)121

Ion-pair HPLC of inorganic anions with photometric, conductometric and amperometric detection 439(1988)363

Low-cost rapid separation and determination of ionic species by single-column ion chromatography 445(1988)183

Indirect photometric detection of inorganic anions in micro HPLC with permanently coated columns 447(1988)221

Anion chromatography using a coated PRP-1 column and eluents of pH > 7 447(1988)315

Carbohydrate-borate eluents for anion chromatography 448(1988)355

Role of the alkyl chain length on the ion interaction reagent, flow-rate, column packing

and detection in ion interaction reversed-phase HPLC in separations of anions using amine salicylates 449(1988)103

## Anserine

Simultaneous determination of histidine-containing dipeptides, histamine, methylhistamine and histidine by HPLC 432(1988)315

## Anthracene-9-carbonyl chloride

Anthracene-9-carbonyl chloride as a fluorescence and UV derivatising reagent for the HPLC analysis of hydroxy compounds 445(1988)393

Determination of diethylene glycol in wine by HPLC using anthracene-9-carbonyl chloride as a derivatising reagent 445(1988)403

## Anthracyclines

Improved method for the determination of 3'-epidoxorubicin and seven metabolites in plasma by HPLC 417(1987)339

## Anthranilic acid

High-voltage electrophoretic separation of tryptophan metabolites of the kynurenine pathway 420(1987)263

## Anthraquinone

Determination of anthraquinone in paper and wood using supercritical fluid extraction and HPLC with electrochemical detection 409(1987)343

## Anthraquinones

Separation of natural products by centrifugal partition chromatography 450(1988)91

## 9-Anthryldiazomethane

Determination of carnitine by HPLC using 9-anthryldiazomethane 445(1988)175

## Antibiotics

Ion-pairing LC method for the determination of pirlimycin hydrochloride 402(1987)335

Separation of novel derivatives from commercial erythromycin samples by TLC 403(1987)343

Mechanism of selectivity in ion-pair HPLC of aminoglycoside antibiotics using perfluorinated pairing ions 404(1987)53

Quantitative analysis of oxytetracycline and related substances by HPLC 405(1987)229

Use of cyclodextrins in isotachopheresis. I. Effect

- of cyclodextrin on the isotachophoretic separation of related penicillins 405(1987)379
- Use of mixed perfluorinated ion-pairing agents as solvents in ion-pair HPLC for the preparative purification of aminoglycoside antibiotics 407(1987)109
- Isotachophoretic analysis of amoxycillin and carboxymethylcysteine in pharmaceuticals 408(1987)441
- Optimization of the separation of erythromycin and related substances by HPLC 409(1987)91
- Determination of chloramphenicol residues by reversed-phase HPLC. Application to a pharmacokinetic study in rainbow trout with confirmation by MS (in French) 409(1987)315
- Determination of vancomycin related substances by gradient HPLC 410(1987)363
- Chromatographic methods for the analysis of vancomycin 410(1987)373
- Improvement of chemical analysis of antibiotics. XIII. Systematic simultaneous analysis of residual tetracyclines in animal tissues using TLC and HPLC 411(1987)313
- Determination of chloramphenicol in swine muscle tissue using a monoclonal antibody-mediated clean-up procedure 411(1987)415
- Hospital routine analysis of penicillins, third-generation cephalosporins and aztreonam by conventional and high-speed HPLC 413(1987)109
- Adsorption of cyclic peptides analogous to gramicidin S and gratisin onto octadecylsilica stationary phase and bacterial cells 413(1987)287
- Determination of cefadroxil in serum by HPLC with cephadrine as internal standard 413(1987)347
- Determination of cefaclor and cephadrine in serum by ion-pair reversed-phase chromatography 413(1987)351
- HPLC analysis of a novel carbapenem antibiotic in human plasma and urine 414(1987)343
- Determination of vancomycin in serum and tissues by column LC using solid-phase extraction 414(1987)472
- HPLC determination of chloramphenicol and four analogues using reductive and oxidative electrochemical and UV detection 417(1987)111
- HPLC analysis of vancomycin in plasma, bone, atrial appendage tissue and pericardial fluid 417(1987)121
- HPLC determination of amphotericin B in human urine 417(1987)439
- Column LC determination of cefpiramide in human serum and urine 417(1987)452
- HPLC assay of erythromycin and its esters using electrochemical detection 419(1987)205
- HPLC assay of the major components of gentamicin in serum 419(1987)408
- Automated analysis of mitomycin C in body fluids by HPLC with on-line sample pre-treatment 420(1987)53
- Estimation of doxorubicin and doxorubicinol by HPLC and advanced automated sample processor 420(1987)184
- Determination of the monocyclic  $\beta$ -lactam antibiotic carumonam in plasma and urine by ion-pair and ion-suppression reversed-phase HPLC 420(1987)357
- Determination of teicoplanin in human plasma and urine by affinity and reversed-phase HPLC 421(1987)99
- Quantitation of ceftetrame in human plasma and urine by HPLC 422(1987)135
- Determination of cefixime in biological samples by reversed-phase HPLC 422(1987)145
- Determination of erythromycin in human plasma, using column LC with a polymeric packing material, alkaline mobile phase and amperometric detection 423(1987)189
- HPLC assay of cefotaxime, desacetylcefotaxime and ceftriaxone in rat plasma 424(1988)111
- HPLC assay for piperacillin in aqueous humor of the eye 424(1988)177
- HPLC determination of adriblastin in human plasma, urine, saliva and liver punctuate by column switching for drug monitoring studies 424(1988)182
- Determination of  $\beta$ -lactams and their biosynthetic intermediates in fermentation media by pre-column derivatisation followed by fluorescence detection 424(1988)325
- Pre-column derivatization of isomycin with *o*-phthalaldehyde- $\beta$ -mercaptopropionic acid and its application to sensitive HPLC determination with fluorimetric detection 425(1988)143
- Optimization of HPLC analysis for isoxazolyl penicillins using factorial design 425(1988)331
- HPLC of erythromycin propionyl ester and

- erythromycin base in biological fluids  
427(1988)93
- Analysis of cephalixin from canine skin biopsy  
by LC with UV-visible photodiode-array  
detection 427(1988)103
- HPLC assay of cephalixin in serum and urine  
427(1988)162
- Reversed-phase HPLC of dactinomycin  
427(1988)166
- Application of HPLC of some antibiotics in  
clinical microbiology 427(1988)257
- HPLC determination of the penem antibiotic  
(5*R*,6*S*)-2-aminomethyl-6-[(1*R*)-hydroxyethyl]-  
2-penem-3-carboxylic acid in human plasma and  
urine 427(1988)269
- Determination of cefetamet and its orally active  
ester, cefetamet pivoxyl, in biological fluids by  
HPLC 430(1988)81
- Chromatographic studies of mitomycin C  
degradation in albumin microspheres  
430(1988)341
- HPLC determination of a new  $\beta$ -lactamase  
inhibitor and its metabolite in combination  
therapy with piperacillin in biological materials  
431(1988)87
- Determination of oxytetracycline in fish by  
HPLC 432(1988)381
- LC and flow injection analysis of tetracycline  
using sensitized europium(III) luminescence  
detection 433(1988)149
- Analysis of erythromycin and roxithromycin in  
plasma or serum by HPLC using electrochemical  
detection 433(1988)298
- HPLC determination of netilmicin in guinea-pig  
and human serum by fluorodinitrobenzene  
derivatization with spectrophotometric detection  
434(1988)169
- TLC study of the metabolites of erythromycins  
in the Wistar rat 434(1988)177
- Improved HPLC determination of doxycycline  
in serum and urine using solid-phase extraction  
columns 434(1988)253
- Photodiode array detection of  
peptide-nucleoside antibiotics 436(1988)259
- Displacement chromatography of biomolecules  
439(1988)341
- Direct broth assay for sparsomycin and related  
nucleoside antitumor antibiotics using reversed-  
phase HPLC 439(1988)459
- Identification of dekamycin antibiotics  
440(1988)71
- Analytical and preparative HPLC of the  
papain-cleaved derivative of polymyxin B  
441(1988)423
- Preparative HPLC of macrotetrolides  
442(1988)431
- Isolation of erythromycin E from commercial  
erythromycin 445(1988)290
- HPLC separation of subcomponents of antimycin  
A 447(1988)65
- LC determination of penicillins by postcolumn  
degradation with sodium hypochlorite using an  
hollow-fibre membrane reactor 447(1988)365
- Improvement of chemical analysis of antibiotics.  
XIV. Identification of the components of  
bacitracin using normal- and reversed-phase TLC  
449(1988)448
- Isolation of gentamicin C compounds from  
culture filtrates of *Micromonospora purpurea*  
450(1988)115
- Antibodies**
- Use of HPLC for the purification of antibodies  
and antibody conjugates and the study of  
antibody-antigen interactions 442(1988)279
- Effect of detergents on the structure of integral  
membrane proteins of Sendai virus studied with  
size-exclusion HPLC and monoclonal antibodies  
443(1988)255
- Antibody conjugates**
- Use of HPLC for the purification of antibodies  
and antibody conjugates and the study of  
antibody-antigen interactions 442(1988)279
- Anti-bovine immunoglobulin G**
- Characterization of adsorption on the stationary  
phase using high-performance immunoaffinity  
chromatography 423(1987)41
- Antidepressants**
- Identification of drugs in human serum by HPLC  
with photodiode array detection and a search  
algorithm for UV spectra 419(1987)135
- Effect of blood collection tubes on antidepressant  
concentrations 423(1987)373
- Antigens**
- Inoculation of BALB/c mice against *Leishmania*  
*major* infection with *Leishmania*-derived antigens  
isolated by gel filtration 440(1988)459

*Entamoeba histolytica* antigens as possible vaccinogens? A short review 440(1988)467

#### **Antimalarial drugs**

Detection and determination of antimalarial drugs and their metabolites in body fluids (Review) 434(1988)1

#### **Antimony**

Trace enrichment and separation of As<sup>III</sup>, Sb<sup>III</sup> and Bi<sup>III</sup> as diethyldithiophosphate complexes by reversed-phase HPLC 439(1988)63

#### **Antimycin A**

HPLC separation of subcomponents of antimycin A 447(1988)65

#### **Antioxidants**

Chromatographic analysis of phenyl-1-naphthylamine and 4,4'-dioctyldiphenylamine and their intermediate oxidation products in oxidized lubricants 409(1987)325

Irregular retention properties of 21-aminosteroid antioxidants in octylsilane bonded-phase HPLC 424(1988)285

HPLC determination of a 21-aminosteroid antioxidant in plasma 424(1988)293

TLC of amine antioxidants and antiozonants used in elastomers 437(1988)59

Determination of phenolic antioxidants in JP-5 jet fuels by GC-mass selective detection 449(1988)440

#### **Antiozonants**

TLC of amine antioxidants and antiozonants used in elastomers 437(1988)59

#### **Antipyrine**

Improved HPLC assay of antipyrine, hydroxymethylantipyrine, 4-hydroxyantipyrine and norantipyrine in urine 433(1988)305

Comparison of solid-phase extraction and liquid-liquid extraction for HPLC analysis of antipyrine in plasma 434(1988)324

Determination of antipyrine in catfish plasma by HPLC 437(1988)306

#### **Apamine**

Characterization of bee venom and its main components by HPLC 408(1987)341

#### **Apigenin**

Separations of flavonoids and alkaloids in medicinal herbs by high-speed counter-current chromatography 435(1988)159

#### **Apolipoprotein E**

Rapid quantification of apolipoprotein E enriched very-low-density lipoproteins by heparin-Sepharose chromatography 425(1988)169

#### **Apolipoproteins**

Separation and characterization of high-density lipoprotein subpopulations by GPC 414(1987)25

Anion-exchange fast protein LC characterization and purification of apolipoproteins A-I, A-II, C-I, C-II, C-III<sub>0</sub>, C-III<sub>1</sub>, C-III<sub>2</sub> and E from human plasma 414(1987)35

Characterization of an apolipoprotein C-III mutant by HPLC and time-of-flight secondary ion MS 414(1987)323

Improved HPLC method for the determination of apolipoproteins in serum high-density lipoproteins 430(1988)135

#### **Apooxytetracycline**

Quantitative analysis of oxytetracycline and related substances by HPLC 405(1987)229

#### **1-β-D-Arabinofuranosylcytosine**

Chromatography of functionalized liposomes and their components 441(1988)115

#### **1-β-D-Arabinofuranosylcytosine 5'-triphosphate**

Separation of 1-β-D-arabinofuranosylcytosine 5'-triphosphate and 9-β-D-arabinofuranosyl-2-fluoroadenine 5'-triphosphate in human leukemia cells by HPLC 413(1987)293

#### **9-β-D-Arabinofuranosyl-2-fluoroadenine 5'-triphosphate**

Separation of 1-β-D-arabinofuranosylcytosine 5'-triphosphate and 9-β-D-arabinofuranosyl-2-fluoroadenine 5'-triphosphate in human leukemia cells by HPLC 413(1987)293

#### **Arabinose**

Simultaneous GLC determination of aldoses and alduronic acids 408(1987)245

Requirements of mutually miscible liquids for ternary chromatographic solvents with reference to monosaccharide PC 436(1988)53

#### **Arachidonic acid**

Analysis of tissue free fatty acids isolated by aminopropyl bonded-phase columns 428(1988)221

Rapid and simple methods for the investigation of lipoxygenase pathways in human granulocytes 431(1988)413

**Arachidonic acid metabolites**

HPLC separation of femtomolar quantities of endogenous carboxylic acids, including arachidonic acid metabolites, as 4-bromomethyl-7-acetoxycoumarin derivatives 416(1987)247

**Arachidonic 5-lipoxygenase products**

Isocratic determination of arachidonic acid 5-lipoxygenase products in human neutrophils by HPLC 430(1988)299

**Arginine**

Determination of agmatine, arginine, citrulline and ornithine by reversed-phase LC using automated pre-column derivatization with *o*-phthalaldehyde 425(1988)269

**Arginine-containing peptides**

HPLC analysis of arginine-containing peptides in biological fluids by means of a selective post-column reaction with fluorescence detection 444(1988)123

**Aromatase inhibitors**

HPLC determination of imide inhibitors of aromatase in biological samples 424(1988)163

**Aromatic acids**

Ion chromatography of inorganic anions on a dynamically modified polystyrene-divinylbenzene packing material and its application to anion screening by sequential UV absorbance and electrochemical detection 402(1987)115

Simultaneous determination of C<sub>2</sub>-C<sub>22</sub> non-esterified fatty acids and other metabolically relevant carboxylic acids in biological material by GC of their benzyl esters 425(1988)257

$\beta$ -Cyclodextrin as a selective agent for separation of selected aromatic acids by HPLC 445(1988)337

Chromatography of microbial metabolites of aromatic amino acids 446(1988)117

GC analysis of different homologous series of acids esterified in aqueous solutions with butyl and propyl alcohols 446(1988)237

**Aromatic alcohols**

Resolution of enantiomeric aromatic alcohols on a cellulose tribenzoate HPLC chiral stationary phase. A proposed chiral recognition mechanism 411(1987)139

**Aromatic choline esterases**

Rapid and efficient method for the isolation and characterization of plant aromatic choline esterases 450(1988)121

**Aromatic hydrocarbons**

HPLC method for isolating organic contaminants from tissue and sediment extracts 437(1988)161

Paraffins, olefins, naphthenes and aromatics analysis of selected hydrocarbon distillates using on-line column switching HPLC with dielectric constant detection 437(1988)365

**N-Aroylanthranilic acids**

HPLC determination of avenanthramides, N-aroylanthranilic acid alkaloids from oats 445(1988)363

**Arsenic**

Simultaneous determination of arsenic(III) and arsenic(V) in metallurgical processing media by ion chromatography 405(1987)247

Trace enrichment and separation of As<sup>III</sup>, Sb<sup>III</sup> and Bi<sup>III</sup> as diethyldithiophosphate complexes by reversed-phase HPLC 439(1988)63

Determination of inorganic arsenic species in aqueous samples by ion-exclusion chromatography with electrochemical detection 450(1988)353

**Arsenic anions**

Non-suppressed ion chromatography of arsenic anions with potassium hydroxide-aromatic salt mixed eluents 447(1988)323

**Artesunic acid**

Determination of artesunic acid after chemical derivatization with *O-p*-nitrobenzyl-*N,N'*-diisopropylisourea by HPLC and UV absorption 410(1987)217

Analysis of artesunic acid and dihydroqinghaosu in blood by HPLC with reductive electrochemical detection 414(1987)77

**Articaine**

HPLC and preliminary pharmacokinetics of articaine and its 2-carboxy metabolite in human serum and urine 424(1988)440

**Artifacts**

Amino acid analysis: buffers and artifacts 436(1988)80

**Arylacetic acids**

Enthalpy-entropy compensation in reversed-phase HPLC of series of aryloxoalkanoic and arylhydroxyalkanoic acids 449(1988)391

 **$\alpha$ -Arylacetic acids**

Resolution of  $\alpha$ -arylacetic acid enantiomers on two chiral stationary phases 404(1987)123

**Arylalkylamines**

Analysis of basic compounds on a silica column with an aqueous methanol eluent. The use of quantitative structure-retention relationships in metabolite identification 407(1987)1

**Arylhydroxyalkanoic acids**

Enthalpy-entropy compensation in reversed-phase HPLC of series of aryloxoalkanoic and arylhydroxyalkanoic acids 449(1988)391

**Aryl N-methylcarbamates**

HPLC determination of aryl N-methylcarbamate residues using post-column hydrolysis electrochemical detection 442(1988)333

**Aryloxoalkanoic acids**

Enthalpy-entropy compensation in reversed-phase HPLC of series of aryloxoalkanoic and arylhydroxyalkanoic acids 449(1988)391

**2-Aryloxypropionic acids**

HPLC separation of the enantiomers of substituted 2-aryloxypropionic acid methyl esters 410(1987)355

**1-Arylpiperazines**

Adsorption chromatography on cellulose. I. Salting-out chromatography of organic compounds 406(1987)389

**2-Arylpropionic acids**

Formation of diastereomeric derivatives of 2-arylpropionic acids using L-leucinamide 423(1987)334

Assay method for product formation in *in vitro* enzyme kinetic studies of uridine diphosphate glucuronyltransferases: 2-arylpropionic acid enantiomers 430(1988)368

Formation of diastereomeric derivatives of 2-arylpropionic acids using L-leucinamide: lack of generality 431(1988)228

Formation of diastereomeric derivatives of 2-arylpropionic acids using  $\alpha$ -leucinamide: the derivatization principle and its applicability. A reply 431(1988)229

 **$\beta$ -Asarone**

Determination of  $\beta$ -asarone, safrole, isosafrole and anethole in alcoholic drinks by HPLC 404(1987)273

**Asbestos**

Improved method for determining adsorption

isotherms by LC on long and short fibrous material 436(1988)391

**Ascorbate-2-phosphate**

HPLC analysis of ascorbate-2-phosphate, adenine and hypoxanthine in stored human blood 419(1987)95

**Ascorbic acid**

Determination of ascorbic acid in beers by HPLC with electrochemical detection 405(1987)347

HPLC-diode-array determination of ascorbic acid, thiamine and riboflavine in goats' milk 410(1987)201

HPLC determination of ascorbic acid in human tears 414(1987)434

Ion suppression reversed-phase HPLC method for the separation of L-ascorbic acid in fresh fruit juice 437(1988)447

Interactions between ascorbic acid and linear and cyclized bile pigments. A TLC study 440(1988)275

**Aspartame**

LC determination of beta-aspartame in diet soft drinks, beverage powders and pudding mixes 404(1987)261

**Aspartic acid**

Determination of 4-aminobutyric acid, aspartate, glutamate and glutamine and their  $^{13}\text{C}$  stable-isotopic enrichment in brain tissue by GC-MS 414(1987)265

HPLC analysis of neurotransmitter amino acids in brain 428(1988)9

Solvent extraction clean-up for pre-treatment in amino acid analysis by GC. Application to age estimation from the D/L ratio of aspartic acid in human dentine 449(1988)165

**Asphaltenes**

GC-MS study of coal-derived asphaltenes. II. Carbazole-type neutral nitrogen compounds 446(1988)293

**Aspoxicillin**

Application of HPLC of some antibiotics in clinical microbiology 427(1988)257

**Association constants**

Determination of association constants of  $\beta$ -cyclodextrin and  $\beta$ -cyclodextrin-bearing polymers with drugs by HPLC 409(1987)61



**Atenolol**

Simple extraction of atenolol from urine and its determination by HPLC 426(1988)234

Preparative resolution of the enantiomers of the  $\beta$ -blocking drug atenolol by chiral derivatization and HPLC 431(1988)222

Determination of atenolol in plasma by dual-column LC and fluorimetric detection 432(1988)265

**Atomic absorption spectrometry**

Isolation and quantitation of cadmium-, zinc- and copper-metallothioneins by HPLC atomic absorption spectrometry 442(1988)345

Comparison of HPLC and atomic spectrometric methods for the determination of Fe(III) and Al(III) in soil and clay samples 449(1988)241

**Atomic environments**

Effect of primary and secondary atomic environments on the interactions of some alkane C-H bonds on a squalane stationary phase at different column temperatures in GLC 406(1987)131

**ATP-haemoglobin**

ATP-hemoglobin: anomalous oxygen binding properties 415(1987)136

**ATP-sulphurylase**

Assay of ATP-sulphurylase activity from rat liver by HPLC 433(1988)63

**Atracurium besylate**

Simultaneous determination of atracurium besylate and its major decomposition products and related impurities by reversed-phase HPLC 435(1988)425

**Atrazine**

Rapid determination of simazine and atrazine in water at the parts per trillion ( $10^{12}$ ) level. Extraction by a miniaturized Carboxpack B trap followed by HPLC 405(1987)357

Specific identification of synthetic organic chemicals in river water using liquid-liquid extraction and resin adsorption coupled with electron impact, chemical ionization and accurate mass measurement GC-MS analyses 436(1988)229

**Atrial natriuretic factor**

Separation of protein hormones (Review) 418(1987)245

**Aucubin**

Application of various planar chromatographic techniques for the separation of iridoid glycosides from *Veronica officinalis* 404(1987)365

**Aurocyanide**

Ion-interaction reversed-phase chromatographic method for the determination of gold(I) cyanide in mine process liquors using automated sample preconcentration 439(1988)23

**Auroxanthin**

LC profiles of major carotenoid esters in commercially processed California navel and Valencia orange juice concentrates 442(1988)249

Separation of carotenol fatty acid esters by HPLC 449(1988)119

**Autofocusing**

Purification of  $\alpha$ -amylase by industrial autofocusing 411(1987)486

**Automated multiple development**

Separation of ecdysteroids by HPTLC using automated multiple development 408(1987)445

**Automation**

Totally automated robotic system for LC analysis of solid dosage formulations with the aid of a reduced Kalman filter 438(1988)319

**Avenanthramides**

HPLC determination of avenanthramides, N-aroylanthranilic acid alkaloids from oats 445(1988)363

**Averantin**

HPLC procedure for determining the profiles of aflatoxin precursors in wildtype and mutant strains of *Aspergillus parasiticus* 441(1988)400

**Averufanin**

HPLC procedure for determining the profiles of aflatoxin precursors in wildtype and mutant strains of *Aspergillus parasiticus* 441(1988)400

**Averufin**

HPLC procedure for determining the profiles of aflatoxin precursors in wildtype and mutant strains of *Aspergillus parasiticus* 441(1988)400

**Avidin-coated glass beads**

Use of avidin-coated glass beads as a support for high-performance immunoaffinity chromatography 444(1988)21

**Axial column flow**

Column hardware in preparative LC with axial flow 450(1988)47

**Axial dispersion**

Theoretical analysis of the asymmetry in chromatographic peaks 404(1987)1

**Azaaromatic hydrocarbons**

*endo*-1,4,5,6,7,7-Hexachlorobicyclo[2.2.1]hept-5-ene-2-carboxylic acid, a superior resolving agent for the HPLC separation of enantiomers of hydroxylated derivatives of two azaaromatic hydrocarbons 430(1988)53

**5-Azacytidine**

HPLC determination of 5-azacytidine in plasma 421(1987)387

**Azaperone**

TLC screening method for the tranquillizers azaperone, propiopromazine and carazolol in pig tissues 436(1988)73

**4-Azasteroids**

HPLC determination of N-(2-methyl-2-propyl)-3-oxo-4-aza-5 $\alpha$ -androst-1-ene-17 $\beta$ -carboxamide, a 4-azasteroid, in human plasma from a phase I study 427(1988)79

**Azathioprine**

Sensitive and specific HPLC assay for 6-thiouric acid 417(1987)183

**Azelastine**

Determination of azelastine and desmethylazelastine in human plasma by HPLC 420(1987)89

**3'-Azido-3'-deoxythymidine**

Simultaneous quantification of zidovudine and its glucuronide in serum by HPLC 431(1988)123

HPLC determination of 2',3'-dideoxycytidine and 3'-azido-3'-deoxythymidine in plasma using a column-switching technique 432(1988)346

**Azidothymidine**

Simple and rapid HPLC assay for zidovudine (azidothymidine) in plasma and urine 430(1988)420

**Azlocillin**

Application of HPLC of some antibiotics in clinical microbiology 427(1988)257

**Azotobactins**

Assignment of the configurations of the amino acids in peptidic siderophores 438(1988)291

**Aztreonam**

Hospital routine analysis of penicillins, third-generation cephalosporins and aztreonam by conventional and high-speed HPLC 413(1987)109

**B****Bacitracin**

Improvement of chemical analysis of antibiotics. XIV. Identification of the components of bacitracin using normal- and reversed-phase TLC 449(1988)448

**Baclofen**

Enantioselective HPLC determination of baclofen after derivatization with a chiral adduct of *o*-phthalaldehyde 415(1987)419

Rapid simultaneous determination of baclofen and its  $\gamma$ -hydroxy metabolite in urine by HPLC with UV detection 420(1987)212

Determination of the *S*(+)- and *R*(-)-enantiomers of baclofen in plasma and urine by GC using a chiral fused-silica capillary column and an electron-capture detector 450(1988)221

**Bacteria**

GC detection of bacteria causing enteric fever 415(1987)377

**Bacteriophages**

Agarose gel electrophoresis of bacteriophages and related particles (Review) 418(1987)345

**Baeocystin**

Determination of tryptamine derivatives in hallucinogenic mushrooms using HPLC with photodiode array detection 408(1987)402

**Bambuterol**

LC method for the determination of bambuterol hydrochloride and related compounds 435(1988)199

**Band broadening**

Effect of steric exclusion on the separation of proteins by hydrophilic SEC 445(1988)29

Absolute peak broadening calibration in SEC

using a polymer-bound chromophore  
449(1988)63

### **Band profiles**

Prediction of band profiles in displacement chromatography by numerical integration of a semi-ideal model 449(1988)25

### **Bandwidth**

HPLC of amino acids, peptides and proteins. LXXXVI. The influence of different displacer salts on the retention and bandwidth properties of proteins separated by isocratic anion-exchange chromatography 443(1988)97

### **Barbiturates**

Determination of sodium pentobarbital and either sodium methohexital or sodium thiopental in plasma by HPLC with ultraviolet detection 414(1986)484

Column LC analysis of barbiturates in biological fluids 427(1988)172

Screening for drugs in clinical toxicology by HPLC: identification of barbiturates by post-column ionization and detection by a multiplace photodiode array spectrophotometer 428(1988)369

Resolution of chiral barbiturates into enantiomers by reversed-phase HPLC using methylated  $\beta$ -cyclodextrins 436(1988)381

Direct LC separation of enantiomers on immobilized protein stationary phases. VI. Optical resolution of a series of racemic barbiturates: studies of substituent and mobile phase effects 436(1988)479

### **Barium**

Isotachophoretic determination of the cationic constituents in mixtures of raw materials for glass manufacturing 439(1988)423

### **BAS111 00W**

Quantitation of triazole and pyrimidine plant growth retardants 438(1988)393

### **Baseline shift**

Gradient elution in ion chromatography 411(1987)107

### **Bases**

Feasibility of isotachochromatography as a method for the preparative separation of weak acids and weak bases. I. Theoretical considerations 436(1988)173

Feasibility of isotachochromatography as a method for the preparative separation of weak acids and weak bases. II. Experimental verification 436(1988)181

### **Basic compounds**

Analysis of basic compounds on a silica column with an aqueous methanol eluent. The use of quantitative structure-retention relationships in metabolite identification 407(1987)1

### **Basic drugs**

Separation of basic drug substances by reversed-phase HPLC on dynamically modified silica and on bonded-phase materials 409(1987)71

Ion-pair reversed-phase TLC of basic drugs using sulphonic acids 441(1988)355

### **Basic dyes**

HPLC system for the separation of basic dyes 445(1988)441

### **Bee venom**

Characterization of bee venom and its main components by HPLC 408(1987)341

### **Benazepril**

Determination of benazepril, a new angiotensin-converting enzyme inhibitor, and its active metabolite, benazeprilat, in plasma and urine by capillary GC-mass-selective detection 434(1988)239

### **Benazeprilat**

Determination of the angiotensin converting enzyme inhibitor benazeprilat in plasma and urine by an enzymic method 425(1988)353

Determination of benazepril, a new angiotensin-converting enzyme inhibitor, and its active metabolite, benazeprilat, in plasma and urine by capillary GC-mass-selective detection 434(1988)239

### **Bencyclane**

Screening procedure for the detection of alkanolamine antihistamines and their metabolites in urine using computerized GC-MS 428(1988)43

### **Bendrofluazide**

LC screening of diuretics in urine 415(1987)347

### **Benedict-Webb-Rubin equation**

Preparation of standard gaseous mixtures by a manometric method using the Benedict-Webb-Rubin equation 436(1988)323

**Benflurene**

Direct absorptiometric quantitation in HPLC in the absence of concomitant reference standards 434(1988)363

**Benoxaprofen**

Assay method for product formation in *in vitro* enzyme kinetic studies of uridine diphosphate glucuronyltransferases: 2-arylpropionic acid enantiomers 430(1988)368

**Benzaldehyde**

Retention behaviour of aromatic aldehyde derivatives in reversed-phase LC 404(1987)242

GC determination of benzaldehyde by gas-phase luminescence 409(1987)243

**Benzamides**

Determination of a new benzamide, amisulpride, in human plasma by reversed-phase ion-pair HPLC 416(1987)414

**Benzazepines**

HPLC studies of pH dependence in the isotopic fractionation of deuterated benzazepines 407(1987)37

**Benzbromarone**

Analysis of benzbromarone in human plasma and urine by HPLC and GC-MS 417(1987)420

**Benzene**

GC procedure for the qualitative analysis of silicone polyester resins after alkaline fusion 441(1988)207

Sorption of benzene on Tenax 447(1988)170

**Benzenepolycarboxylate buffers**

Single-column anion chromatography with indirect UV detection using pyromellitate buffers as eluents 439(1988)137

**Benzimidazole**

HPLC determination of formate as benzimidazole in biological samples 426(1988)15

**Benzindene prostaglandin**

HPLC method for determining the enantiomeric purity of a benzindene prostaglandin by a diastereomeric separation 408(1987)275

**Benzoates**

Host-guest complexation in capillary isotachopheresis. I.  $\alpha$ -,  $\beta$ - and  $\gamma$ -cyclodextrins as complexing agents for the resolution of substituted benzoic acid isomers 442(1988)371

**Benzocarbazoles**

GC-MS study of coal-derived asphaltenes. II. Carbazole-type neutral nitrogen compounds 446(1988)293

**Benzodiazepine-receptor ligands**

Lipophilicity measurement of benzodiazepine-receptor ligands by reversed-phase LC. Comparison between HPLC and TLC 447(1988)404

**Benzodiazepines**

Screening procedure for benzodiazepines in biological fluids by HPLC using a rapid-scanning multichannel detector 416(1987)303

Combined HPLC and enzyme immunoassay for active metabolites of a sleep inducer (rilamazafone), a ring-opened derivative of benzodiazepines, in human plasma 417(1987)216

Identification of drugs in human serum by HPLC with photodiode array detection and a search algorithm for UV spectra 419(1987)135

Identification and differentiation of benzodiazepines and their metabolites in urine by computerized GC-MS 422(1987)85

Simultaneous determination of diazepam and its active metabolites in rat serum, brain and cerebrospinal fluid by HPLC 422(1987)334

Analysis of benzodiazepines. II. HPLC fluorescence detection after molecular rearrangement to acridanones 428(1988)321

Collection of analytical data for benzodiazepines and benzophenones 439(1988)317

Detection and determination of common benzodiazepines and their metabolites in blood samples of forensic science interest. Microcolumn cleanup and HPLC with reductive electrochemical detection at a pendent mercury drop electrode 449(1988)281

**Benzoic acid**

Analysis of manufacturing by-products and impurities in illicit cocaine via HPLC and photodiode array detection 405(1987)273

Determination of benzoic acid and hippuric acid in human plasma and urine by HPLC 425(1988)67

1-(4-Hydroxyphenyl)-, 1-(2,4-dihydroxyphenyl)- and 1-(2,5-dihydroxyphenyl)-2-bromoethanones: new labels for determination of carboxylic acids by HPLC with electrochemical and UV detection 442(1988)209

Extractive pentafluorobenylation of formic, acetic, levulinic, benzoic and phthalic acids, studied by LC and dual-oven capillary GC 447(1988)329

#### **Benzonal**

Direct LC separation of enantiomers on immobilized protein stationary phases. VI. Optical resolution of a series of racemic barbiturates: studies of substituent and mobile phase effects 436(1988)479

#### **Benzo[*lm*]phenanthro[4,5,6-*abcd*]perylene**

Effect of column temperature on the retention of peropyrene-type polycyclic aromatic hydrocarbons on various chemically bonded stationary phases in reversed-phase LC 436(1988)1

#### **Benzophenones**

Collection of analytical data for benzodiazepines and benzophenones 439(1988)317

#### **Benzopyrans**

Rapid analyses of chromenes (benzopyrans) by GC-MS 403(1987)358

#### **Benzoquinone**

Interconversion of hydroquinone and benzoquinone during styrene-divinylbenzene resin-based ion-exclusion chromatography 409(1987)404

#### **Benzothiophenes**

Identification of homologous series of alkylated thiophenes, thiolanes, thianes and benzothiophenes present in pyrolysates of sulphur-rich kerogens 435(1988)435

#### **Benzotriazine di-N-oxide**

HPLC assay for the benzotriazine di-N-oxide (SR 4233) and its reduced metabolites in biological materials 430(1988)429

#### **Benzoxycarbonylglycyl-L-proline**

Influence of enantiomeric purity of a chiral selector on stereoselectivity 407(1987)217

Supercritical fluid chromatography with ion-pairing modifiers. Separation of enantiomeric 1,2-aminoalcohols as diastereomeric ion pairs 447(1988)287

#### **14-Benzoylaconine**

HPLC analysis of C<sub>19</sub>-diterpenoid alkaloids 402(1987)366

#### **Benzoylated polyacrylamide**

Chromatographic determination of cyclodextrins on benzoylated polyacrylamide gels 447(1988)398

#### **Benzoyl chloride**

Improved method for benzoyl chloride derivatization of polyamines for HPLC 408(1987)227

#### **Benzoyllecgonine**

Analysis of manufacturing by-products and impurities in illicit cocaine via HPLC and photodiode array detection 405(1987)273

Determination of benzoyllecgonine and cocaine in biologic fluids by automated GC 417(1987)277

#### **S-Benzoyl-3-mercapto-2-methylpropanoyl chloride**

Simultaneous determination of the acid chloride S-benzoyl-3-mercapto-2-methylpropanoyl chloride and residual carboxylic acid by capillary GC 408(1987)360

#### **Benzylhydrazine**

GC analysis of monoalkylhydrazines 423(1987)131

#### **N-Benzyl-N-methylephedrinium**

Chiral Michael additions of acetamidomalonnate to  $\alpha$ -enones. NMR and HPLC resolution of enantiomers 450(1988)183

#### **Benzyloxyphenylacetic acids**

Enthalpy-entropy compensation in reversed-phase HPLC of series of aryloxoalkanoic and aryhydroxyalkanoic acids 449(1988)391

#### **Betamethasone**

GC and MS study of betamethasone synthetic corticosteroid metabolism 432(1988)283

#### **Betaxolol**

Column LC measurement of betaxolol in plasma or serum 417(1987)229

Correlation between reversed-phase HPLC and plasma protein binding 421(1987)83

#### **Biaryls**

HPLC separation of enantiomers on axially chiral binaphthalene derivatives bonded to silica gel 403(1987)275

#### **Bidesmosidic saponins**

HPLC determination of molluscicidal saponins

from *Phytolacca dodecandra* (Phytolaccaceae) 448(1988)265

### **Biflavones**

Reversed-phase HPLC method for the analysis of biflavones in *Ginkgo biloba* L. extracts 437(1988)453

### **Biguanides**

Fluorometric determination of biguanides in serum by HPLC with reagent-containing mobile phase 430(1988)65

### **Bile acid disulphates**

Studies on steroids. CCXXXIII. Separation and characterization of bile acid disulphates in human urine by HPLC 425(1988)59

### **Bile acids**

Separation and quantitation of fatty acids, sterols and bile acids in feces by GC as the butyl ester-acetate derivatives 415(1987)13

Determination of  $3\beta,12\alpha$ -dihydroxy-5-cholen-24-oic acid and related bile acids in human serum by GC-MS 421(1987)9

Improved method for the determination of the major neutral steroids and unconjugated bile acids in human faeces using capillary GC 421(1987)21

Internal standards for quantitative GC of individual bile acids after group separation of bile acids in urine 422(1987)25

Medroxyprogesterone acetate- and ethinylestradiol-induced changes in biliary bile acids of the rat studied by HPLC 423(1987)63

Analysis of conjugated bile acids in human serum by HPLC using post-column enzyme reaction and off-line fluorimetric determination 423(1987)75

Simultaneous microanalysis of bile acids and cholesterol in bile by glass capillary GC 431(1988)17

Rapid and sensitive on-line precolumn purification and HPLC assay for bile acids in serum 431(1988)27

Simultaneous determination of free and conjugated bile acids in human gastric juice by HPLC 431(1988)259

Separation of keto and keto-hydroxy bile acid isomers by HPLC 438(1988)423

Use of cyclodextrins in isotachopheresis. VI.

Cyclodextrins as leading electrolyte additives for the separation of bile acids 450(1988)373

### **Bile acid sulphates**

Studies on steroids. CCXXVII. Separation and determination of bile acid 7- and 12-sulphates in urine by HPLC with fluorescence labelling 415(1987)45

### **Bile pigments**

Interactions between ascorbic acid and linear and cyclized bile pigments. A TLC study 440(1988)275

### **Bilirubin**

Novel method for HPLC of azo derivatives of conjugated and unconjugated bilirubin 427(1988)19

Correlation of carbon monoxide and bilirubin production by tissue homogenates 427(1988)315

Study of the mechanism of the photochemical oxidation of bilirubin by HPLC 428(1988)255

### **Biliverdin**

Study of the mechanism of the photochemical oxidation of bilirubin by HPLC 428(1988)255

Interactions between ascorbic acid and linear and cyclized bile pigments. A TLC study 440(1988)275

### **Binaphthalene**

HPLC separation of enantiomers on axially chiral binaphthalene derivatives bonded to silica gel 403(1987)275

### **Binary mixtures**

Influence of the relative concentrations of two components of a binary mixture on their overloaded elution profiles 440(1988)9

### **Binary solvents**

Thermochemical investigations of GLC. IV. Partition coefficients of solutes on binary mixtures containing two complexing solvents 402(1987)41

### **Binding constants**

Determination of binding constants by high-performance gel filtration 404(1987)254

### **Biomarkers**

Determination of environmental levels of peptidoglycan and lipopolysaccharide using GC with negative-ion chemical-ionization MS

utilizing bacterial amino acids and hydroxy fatty acids as biomarkers 431(1988)1

Biomarker determinations in crude oils using a triple-stage quadrupole mass spectrometer 446(1988)3

#### **Biomimetic dyes**

Design and applications of biomimetic anthraquinone dyes. I. Synthesis and characterisation of terminal ring isomers of C.I. Reactive Blue 2 435(1988)127

#### **Biopolymers**

Hydrolysed macroporous glycidyl methacrylate-ethylene dimethacrylate copolymer sorbent for size-exclusion HPLC of synthetic polymers and biopolymers 435(1988)357

#### **Biopolymer separations**

Sample preparation and column regeneration in biopolymer separations (Review) 418(1987)27

#### **Biopterin**

Estimation of tetrahydrobiopterin and other pterins in plasma by isocratic LC with electrochemical and fluorimetric detection 432(1988)321

#### **Biotin**

Determination of free biotin in plasma by LC with fluorimetric detection 413(1987)247

#### **Biphenylacetic acid**

Quantitation of biphenylacetic acid in plasma and synovial fluid by GC-MS-MS 420(1987)129

#### **4,4'-Bipyridylum salts**

Isotachopheresis of quaternary 4,4'-bipyridylum salts. Analytical control of synthesis and purification procedures 449(1988)299

#### **Birch pollen**

Purification of allergens by HPLC. VII. Purification of the major allergen of birch pollen (*Betula verrucosa*) 408(1987)420

#### **Bisacodyl**

Screening procedure for stimulant laxatives in urine using HPLC with diode array detection 433(1988)131

#### **Bisacodyl diphenol**

Screening procedure for stimulant laxatives in urine using HPLC with diode array detection 433(1988)131

#### **1,3-Bis[ $\omega$ -alkoxyoligo(oxyethylene)]propan-2-ols**

Polarity of 1,3-bis[ $\omega$ -alkoxyoligo(oxyethylene)]propan-2-ols as measured by reversed-phase GC 409(1987)29

#### **Bis(L-amino acid amidato)copper complexes**

Bis(L-amino acid amidato)copper(II) complexes as chiral eluents in the enantiomeric separation of D,L-dansylamino acids by reversed-phase HPLC 441(1988)287

#### **Bis[1,2-bis(diphenylphosphino)ethane]-gold(I)**

HPLC of bis[1,2-bis(diphenylphosphino)ethane]-gold(I) chloride, a potential antineoplastic agent 448(1988)333

#### **2,5-Bis-[5-*tert.*-butylbenzoxazolyl(2')]-thiophene**

Method for determination of different phospholipids by circular HPTLC 413(1987)257

#### **Bis(2-chloroethyl)amine**

Combined TLC-photography-densitometry for the quantitation of cyclophosphamide and its four principal urinary metabolites 427(1988)121

#### **1,1-Bis(4-chlorophenyl)-2,2,2-trichloroethane**

Analysis of pentachlorophenol in water and urine by enrichment with Lipidex 5000 414(1987)55

#### **Bisdesethylchloroquine**

Analysis of blood and urine samples for hydroxychloroquine and three major metabolites by HPLC with fluorescence detection 433(1988)197

#### **Bismuth**

Trace enrichment and separation of As<sup>III</sup>, Sb<sup>III</sup> and Bi<sup>III</sup> as diethyldithiophosphate complexes by reversed-phase HPLC 439(1988)63

#### **Bis[(1-naphthyl)methyl]acetylhistidylstatine 2(S)-methylbutylamide**

Purification of human renin by renin inhibitor (ES-305) affinity chromatography 421(1987)350

#### **Bisoprolol**

Rapid and sensitive HPLC determination of bisoprolol in plasma and urine 426(1988)431

#### **Bisoxatin**

Screening procedure for stimulant laxatives in

urine using HPLC with diode array detection 433(1988)131

#### **Bisoxatin diphenol**

Screening procedure for stimulant laxatives in urine using HPLC with diode array detection 433(1988)131

#### **Bisphenol A**

Determination of bisphenol A in air by HPLC with electrochemical detection 439(1988)375

Chromatographic determination of residual monomers of epoxy resins in wine simulants (in French) 445(1988)107

#### **Bisphosphonates**

Determination of (4-chlorophenyl)thiomethylene biphosphonic acid, a new bisphosphonate, in biological fluids by HPLC 430(1988)73

#### **2,8-Bis(trifluoromethyl)-4-quinoline carboxylic acid**

HPLC separation and extraction investigation for the simultaneous determination of mefloquine and its carboxylic acid metabolite 427(1988)295

#### **Bitter acids**

Avoidance of trace metal activity with micro-LC in the analysis of hop and beer bitter acids 407(1987)159

#### **Blasticidin S**

Photodiode array detection of peptide-nucleoside antibiotics 436(1988)259

#### **Blockage effect**

Simplified description of HPLC separation under overload conditions, based on the Craig distribution model. III. Computer simulations for two co-eluting bands assuming a Langmuir isotherm 405(1987)1

Simplified description of HPLC separation under overload conditions, based on the Craig distribution model. IV. Comparison of the model with experimental data for samples containing two or more solutes 405(1987)31

#### **Blood collection tubes**

Effect of blood collection tubes on antidepressant concentrations 423(1987)373

#### **Blood substitutes**

Hemoglobin-based blood substitutes: characterization of five pyridoxal 5'-phosphate derivatives of hemoglobin 419(1987)37

#### **Boiling point**

Factor analysis of trends in selectivity in GLC of 175 saturated and unsaturated esters on 14 polysiloxane stationary phases 404(1987)11

#### **Boldenone sulphate**

Screening, confirmation and quantification of boldenone sulfate in equine urine after administration of boldenone undecylenate (Equipoise) 433(1988)9

#### **Boldenone undecylenate**

Screening, confirmation and quantification of boldenone sulfate in equine urine after administration of boldenone undecylenate (Equipoise) 433(1988)9

#### **Bonded layer structure**

Geometric structural properties of bonded layers of chemically modified silicas 447(1988)103

#### **Bonding chemistry**

Influence of bonding chemistry on the reordering/resolution of silica immobilized alkyl chains 449(1988)115

#### **Bone marrow cells**

Partitioning behaviour of rat bone marrow cells in aqueous two-phase systems. Dependence of cell partition on the interfacial tension and electrical potential difference between the phases 403(1987)131

#### **Bopindolol**

Column LC determination of hydrolysed bopindolol, in the picogram per millilitre range in plasma, using cartridge extraction and dual electrochemical detection 422(1987)205

#### **Borate**

Elution characteristics of polyol-borate eluents in anion-exchange chromatography. I. Eluent strength of sugar-, uronate-borate eluents 403(1987)27

Elution characteristics of polyol-borate eluents in anion-exchange chromatography. II. Tartrate-borate eluent for non-suppressed anion chromatography 403(1987)35

Origin of system peaks in single-column ion chromatography of inorganic anions using high pH borate-gluconate buffers and conductivity detection 404(1987)67

HPLC of N-terminal tyrosine-containing oligopeptides by pre-column fluorescence



derivatization with hydroxylamine, cobalt(II) and borate reagents 411(1987)305

Carbohydrate-borate eluents for anion chromatography 448(1988)355

### **Boron**

Boron isotope separation by ion-exchange chromatography using an anion-exchange resin in halide forms. Separation factors at 25 °C 450(1988)343

### **Botrytis**

Characterization of *Botrytis* by GLC 403(1987)292

### **Botulinum neurotoxins**

Fast protein LC of botulinum neurotoxin types A, B and E 430(1988)279

### **Bovine leukemia virus**

High-performance GPC of the surface glycoprotein from bovine leukemia virus 425(1988)379

### **Bovine serum albumin**

Rapid preparation of bovine mercaptalbumin by means of covalent chromatography on silica-based materials 408(1987)263

Influence of pore and particle size on the frontal uptake of proteins. Implications for preparative anion-exchange chromatography 409(1987)111

### **Bovine viral diarrhea virus**

HPLC separation and immunological characterization of soluble bovine viral diarrhea virus antigen 435(1988)149

### **Branched-chain amino acids**

Rapid HPLC method to measure plasma leucine: importance in the study of leucine kinetics *in vivo* 426(1988)370

### **Branched-chain keto acids**

GC of branched-chain keto acids analysed as their silylated oxime derivatives 423(1987)270

### **Branched-chain 2-oxo acids**

Simple method for rapid quantification of branched-chain 2-oxo acids in physiological fluids as quinoxalinol derivatives by HPLC 432(1988)297

### **Brilliant Blue FCF**

Simple method for the analysis of food dyes on reversed-phase thin-layer plates 411(1987)437

### **Brobactam**

Determination of 6 $\beta$ -bromopenicillanic acid (brobactam) in human serum and urine by HPLC 425(1988)425

### **Bromacil**

Analysis of bromacil, diuron and 3,4-dichloroaniline in contaminated well water, using a HPLC column-switching procedure 410(1987)211

Analysis of the herbicides dicamba, clopyralid and bromacil in asparagus by HPLC 439(1988)470

### **Bromadiolone**

Determination of diastereoisomers of bromadiolone, an anticoagulant rodenticide, in animal tissues by HPLC 435(1988)83

### **Bromhexine**

Determination of bromhexine in plasma by reversed-phase LC. Interference of lipoproteins on extraction 432(1988)243

### **Bromide**

Derivatization of inorganic anions with pentafluorobenzyl methanesulphonate for GC 404(1987)175

Polymeric pentafluorobenzylating reagent for GC of inorganic anions with electron-capture detection 438(1988)253

Determination of iodide and bromide by ion chromatography with post-column reaction detection 439(1988)129

### **Bromination**

Investigations into lactoperoxidase-catalysed bromination of tyrosine and thyroglobulin 432(1988)57

### **Bromine**

Evaluation of a chemical reaction detector with electrochemical production of bromine for the LC determination of physostigmine 411(1987)229

### **Bromine-containing compounds**

GC quantification of bromine-containing compounds using a flame ionization detector 402(1987)349

Spectra and response linearization of chlorine and bromine compounds in a copper-sensitized flame photometric detector 408(1987)69

### **5-Bromo-2'-deoxyuridine**

Improved HPLC assay for the quantification of

5-bromo-2'-deoxyuridine and 5-bromouracil in plasma 432(1988)233

### **2-Bromoethanones**

1-(4-Hydroxyphenyl)-, 1-(2,4-dihydroxyphenyl)- and 1-(2,5-dihydroxyphenyl)-2-bromoethanones: new labels for determination of carboxylic acids by HPLC with electrochemical and UV detection 442(1988)209

### **Bromofenofos**

HPLC determination of bromofenofos and its metabolite in rat plasma 426(1988)202

### **Bromoform**

GC quantification of bromine-containing compounds using a flame ionization detector 402(1987)349

### **$\alpha$ -Bromoisovalerylurea**

LC determination of diastereomeric glutathione conjugates and further derivatives of  $\alpha$ -bromoisovalerylurea in rat bile and urine by electrochemically generated bromine 427(1988)67

### **4-Bromomethyl-6,7-dimethoxycoumarin**

4-Bromomethyl-6,7-dimethoxycoumarin as a fluorescence reagent for precolumn derivatization of 5-fluorouracil compounds in HPLC 430(1988)156

### **4-Bromomethyl-7-methoxycoumarin**

Quantitative determination of octanoic acid by HPLC following derivatization with 4-bromomethyl-7-methoxycoumarin 411(1987)297

### **2-Bromo-2-nitropropanes**

TLC studies of biocidal 2-bromo-2-nitropropane derivatives (in German) 438(1988)122

### **6 $\beta$ -Bromopenicillanic acid**

Determination of 6 $\beta$ -bromopenicillanic acid (brobactam) in human serum and urine by HPLC 425(1988)425

### **Bromosulphophthalein**

Rapid method for the determination and quantification of bromosulphophthalein and metabolites in cultured hepatocytes, culture media and bile by HPLC 432(1988)223

### **5-Bromouracil**

Improved HPLC assay for the quantification of 5-bromo-2'-deoxyuridine and 5-bromouracil in plasma 432(1988)233

### **(E)-5-(2-Bromovinyl)-2'-deoxyuridine**

Bioanalysis of (E)-5-(2-bromovinyl)-2'-deoxyuridine 428(1988)93

### **Brown coal**

Analytical characterisation of hydrocarbon and hetero compounds mixtures from reactions of brown coal (in German) 446(1988)305

### **Brush-border membranes**

Comparative chromatographic study of modifications of brush-border membrane vesicles induced by an essential fatty acid-deficient diet 432(1988)75

### **BTS 53554**

Simplified HPLC determination of flosequinan and its metabolite in plasma, serum and urine 425(1988)227

### **Budralazine**

Quantitative analysis of budralazine in human plasma and urine by GC-MS 421(1987)367

### **Bufencarb**

HPLC determination of aryl N-methylcarbamate residues using post-column hydrolysis electrochemical detection 442(1988)333

### **Buffers**

Amino acid analysis: buffers and artifacts 436(1988)80

Buffer isoelectric focusing revisited 440(1988)367

### **Buflovedil**

GC evaluation of buflovedil in biological samples employing the thermoionic specific detector 427(1988)345

### **Buformin**

Fluorometric determination of biguanides in serum by HPLC with reagent-containing mobile phase 430(1988)65

### **Bulgarian rose oil**

Importance of the odoriferous principle of Bulgarian rose oil in flavour and fragrance chemistry 406(1987)181

Composition of essential oils. Part 7. Bulgarian oil of rose (*Rosa damascena* Mill.) 406(1987)185

### **Bumetanide**

LC screening of diuretics in urine 415(1987)347

Determination of bumetanide in plasma by HPLC 434(1988)327

### **Bupivacaine**

Simple, rapid HPLC determination of bupivacaine in human plasma 417(1987)236

Stereoselective HPLC assay for bupivacaine enantiomers 420(1987)203

**Buspirone**

Simultaneous quantitation of buspirone and 1-(2-pyrimidinyl)piperazine in human plasma and urine by capillary GC-MS 428(1988)265

Simultaneous determination of buspirone, gepirone, ipsapirone and their common metabolite 1-(2-pyrimidinyl)piperazine in rat plasma and brain by HPLC 431(1988)477

**Busulfan**

Measurement of busulfan in plasma by HPLC 416(1987)426

Determination of busulfan in human plasma by GC with electron-capture detection 425(1988)303

GC-MS assay for busulfan in biological fluids using a deuterated internal standard 428(1988)357

Determination of busulfan in human plasma by GC with electron-capture detection 434(1988)

**2,3-Butanediol**

GC method for the estimation of acetone and its metabolites in biological samples 415(1987)372

Determination of methyl ethyl ketone and its metabolites in urine using capillary GC 428(1988)275

**Butanol**

Determination of methyl ethyl ketone and its metabolites in urine using capillary GC 428(1988)275

Static headspace GC method for the determination of residual solvents in vigabatrin drug substance 438(1988)103

**Butanols**

Determination of alcohols in alcoholic beverages by micro HPLC with indirect photometric detection 445(1988)139

**1-(3-Butoxy-2-carbamoyloxypropyl)-5-ethyl-5-phenyl-(1*H*,3*H*,5*H*)-pyrimidine-2,4,6-trione**

HPLC determination of 1-(3-butoxy-2-carbamoyloxypropyl)-5-ethyl-5-phenyl-(1*H*,3*H*,5*H*)-pyrimidine-2,4,6-trione in human plasma 428(1988)183

**Butyl acrylate**

Investigation by pyrolysis-GC of the composition of multicomponent polymeric

microheterogeneous systems based on some vinyl monomers 404(1987)183

**Butylated hydroxyanisole**

Sensitive HPLC method for the routine determination of butylated hydroxyanisole in plasma 413(1987)282

**Butylated hydroxytoluene**

Determination of butylated hydroxytoluene in plasma by HPLC 422(1987)288

**4-*tert*.-Butyl-*o*-benzoquinone**

LC analysis of products of reaction between methyl linoleate and 4-*tert*.-butyl-*o*-benzoquinone 408(1987)171

***n*-Butyl- $\beta$ -carboline-3-carboxylate**

Determination of *n*-butyl- $\beta$ -carboline-3-carboxylate in aqueous brain extracts 431(1988)192

**2-*tert*.-Butyl-6-isothiocyanato-5-(piperidin-1-yl)benzthiazole**

HPLC of antifilarials. III. Determination of 2-*tert*.-butyl-6-isothiocyanato-5-piperidin-1-yl)-benzthiazole in biological fluids 434(1988)303

**Butylphenols**

Separation of isomeric mono-, di- and tributylphenols by HPLC on octadecylsilica with methanol-water using alkyl nitrile and alkylamine as surface modifiers 442(1988)53

**Butyltin compounds**

Characterization of organotin species using microbore and capillary LC techniques with an epifluorescence microscope as a novel imaging detector 410(1987)393

**Butyric acid**

Comparison of methods for the determination of butyric acid in foodstuffs by GLC and HPLC 410(1987)190

Studies on sample preconcentration in ion chromatography. VIII. Preconcentration of carboxylic acids prior to ion-exclusion separation 447(1988)155

 **$\gamma$ -Butyrobetaine-binding protein**

Detection of ligand-protein binding by direct electrophoresis of the complex 449(1988)331

## C

**Cadaverine**

Improved method for benzoyl chloride derivatization of polyamines for HPLC 408(1987)227

Determination of polyamines in human erythrocytes by capillary GC with nitrogen-phosphorus detection 415(1987)27

Simultaneous determination of  $\gamma$ -aminobutyric acid and polyamines by HPLC 419(1987)285

Reversed-phase LC separation and simultaneous fluorimetric detection of polyamines and their monoacetyl derivatives in human and animal urine, serum and tissue samples: an improved, rapid and sensitive method for routine application 430(1988)249

Quantitation of polyamines using TLC and image analysis 441(1988)361

Detection by HPLC of polyamines formed by clostridal putrefaction of caseins 442(1988)267

**Cadmium**

Ion chromatographic determination of low level cadmium(II), cobalt(II) and manganese(II) in water 450(1988)361

**Cadralazine**

HPLC determination of cadralazine in human plasma and urine 424(1988)119

Microbore LC determination of cadralazine and cephalixin in plasma with large-volume injection 426(1988)335

**Caffeine**

Pharmacokinetics of intravenous caffeine: comparison of HPLC and GC methods 422(1987)274

Use of micellar mobile phases and microbore column switching for the assay of drugs in physiological fluids 427(1988)277

HPLC of xanthines. Effects of N-methyl and C-8 hydroxyl substitution on the retention times 428(1988)311

**Cajanol**

Reversed-phase HPLC of *Cajanus cajan* phytoalexins 409(1987)355

**Cajanol**

Reversed-phase HPLC of *Cajanus cajan* phytoalexins 409(1987)355

**Calcidiol**

Rapid automated HPLC assay for ercalcidiol and calcidiol (25-hydroxyvitamins D<sub>2</sub> and D<sub>3</sub>) using *trans*-calcidiol as an UV-absorbing internal standard 415(1987)305

**Calcitonin**

Separation of protein hormones (Review) 418(1987)245

Correlation of calcitonin structure with chromatographic retention in HPLC 443(1988)229

**Calcium**

Isotachophoretic determination of the cationic constituents in mixtures of raw materials for glass manufacturing 439(1988)423

**Calcium acetylhomotaurinate**

Determination of calcium acetylhomotaurinate by LC with fluorimetric and electrochemical detection 414(1987)417

**Calcium-binding protein**

Chick 28 000 *M<sub>r</sub>* vitamin D-dependent calcium-binding protein in intestine, kidney and cerebellum. Purification using chromatofocusing 448(1988)145

**Calcium dodecylbenzenesulphonate**

Determination of dodecylbenzenesulphonates and ethoxylated alkylphenols in liquid pesticide formulations by HPLC 435(1988)73

**Calcium hydroxide**

Supplementary and revised assignment of the peaks of the 7,9-, 9,9'-, 13,13'-, 9,13'-di-*cis* and 9,9',13-tri-*cis* isomers of  $\beta$ -carotene in HPLC using a column of calcium hydroxide 439(1988)417

**Calcium-pump protein**

Use of a small TSK GSW HPLC column for large-zone chromatographic studies of monomer-oligomer equilibria of membrane protein 442(1988)229

**Calibration methods**

Technique for calibrating air samplers and certain other analytical devices 438(1988)100

TLC-flame ionization detection calibration with natural and synthetic standards 439(1988)404

#### **Calcium stearate**

GC determination of calcium stearate in polyethylene food packaging sheets 447(1988)198

#### **Calmodulin**

Melittin-silica, a high-pressure affinity chromatography resin for calmodulin 403(1987)99

#### **Camphene**

Resolution of monoterpene enantiomers by GC 407(1987)243

#### **Cannabidiol**

Rapid HPLC assay with pharmacokinetic applications for monitoring cannabidiol in plasma 416(1987)370

#### **Cannabinoids**

Confirmation of cannabis use. III. Simultaneous quantitation of six metabolites of  $\Delta^9$ -tetrahydrocannabinol in plasma by HPLC with electrochemical detection 434(1988)247

#### **Cantharidin**

GC-MS method for the quantitation of cantharidin in human serum 432(1988)177

#### **Canthaxanthin**

Isolation of *cis-trans* isomers of canthaxanthin by HPLC using a calcium hydroxide column and identification of their configurations by  $^1\text{H}$  NMR spectroscopy 448(1988)182

#### **Capacity factor**

Theoretical prediction of plate height in serially coupled capillary columns 406(1987)3

Prediction of HPLC capacity factors following column calibration. Influence of different solvents and columns on the separation of a series of aromatic amides 437(1988)13

Effect of hydrogen bonding on the HPLC behaviour of organic compounds. Relationship between capacity factors and partition coefficients 439(1988)227

LC retention of  $\beta$ -diketonato complexes of chromium(III) and palladium(II) in octadecyl-bonded silica/water-methanol systems. Comparison with liquid-liquid partition using dodecane as a model alkyl-bonded phase 439(1988)245

#### **Capillary columns**

see also Columns and microbore columns

Theoretical prediction of plate height in serially coupled capillary columns 406(1987)3

"Superdynamic" coating of the inner surface of a capillary column with a liquid stationary phase 440(1988)323

Preparation of wall-coated open-tubular capillary columns for GC (Review) 445(1988)1

#### **Capillary isoelectric focusing**

Capillary isoelectric focusing using an LKB 2127 Tachophor isotachophoretic analyzer 436(1988)309

#### **Capillary wall pre-treatment**

Preparation of wall-coated open-tubular capillary columns for GC (Review) 445(1988)1

#### **Capillary zone electrophoresis**

Evaluation of an automatic siphonic sampler for capillary zone electrophoresis 404(1987)313

Capillary zone electrophoresis and isotachopheresis as alternatives to chromatographic methods for purity control of synthetic peptides 407(1987)363

Fraction collector for capillary zone electrophoresis 438(1988)23

Retention of ionic and non-ionic catechols in capillary zone electrophoresis with micellar solutions 441(1988)299

#### **Capsaicin**

GC of methyl derivatives of naturally occurring capsaicinoids 409(1987)223

#### **Capsaicinoids**

GC of methyl derivatives of naturally occurring capsaicinoids 409(1987)223

#### **Capsanthin**

Separation and quantitative analysis of some carotenoid fatty acid esters of fruits by LC 435(1988)113

#### **Capsicin**

Chromatographic purification and characterization of elicitors of necrosis on tobacco produced by incompatible *Phytophthora* species 440(1988)87

#### **Capsorubin**

Separation and quantitative analysis of some

carotenoid fatty acid esters of fruits by LC  
435(1988)113

#### **Captan**

HPLC determination of sulphur and captan in a mixture 447(1988)263

#### **Captopril**

Sensitive method for determination of captopril in biological fluids by GC-MS 417(1987)79

Simplified determination of captopril in plasma by HPLC 425(1988)208

#### **Carazolol**

TLC screening method for the tranquillizers azaperone, propiopromazine and carazolol in pig tissues 436(1988)73

#### **Carbamates**

HPLC determination of aryl N-methylcarbamate residues using post-column hydrolysis electrochemical detection 442(1988)333

#### **Carbamazepine**

Simultaneous determination of carbamazepine, oxcarbazepine and their main metabolites in plasma by LC 414(1987)477

HPLC drug analysis by direct injection of whole blood samples. I. Determination of moderately hydrophobic drugs incorporated into blood corpuscles 423(1987)147

#### **Carbanilides**

LC behaviour of chlorinated carbanilides 410(1987)95

#### **Carbapenem**

HPLC analysis of a novel carbapenem antibiotic in human plasma and urine 414(1987)343

#### **Carbaryl**

Pre-column clean-up and LC determination of residues of N-methylcarbamate pesticides in extracts of total diet 404(1987)352

HPLC determination of aryl N-methylcarbamate residues using post-column hydrolysis electrochemical detection 442(1988)333

Determination of carbaryl by HPLC with electrochemical detection 442(1988)451

Determination of organophosphate pesticides and carbaryl on paddy rice by reversed-phase HPLC 447(1988)249

#### **Carbazoles**

GC-MS study of coal-derived asphaltenes. II.

Carbazole-type neutral nitrogen compounds 446(1988)293

#### **Carbidopa**

Simultaneous determination of levodopa, its metabolites and carbidopa in clinical samples 415(1987)393

#### **Carbinoxamine**

Screening procedure for the detection of alkanolamine antihistamines and their metabolites in urine using computerized GC-MS 428(1988)43

#### **Carbocyclic acids**

Analysis of manufacturing by-products and impurities in illicit cocaine via HPLC and photodiode array detection 405(1987)273

#### **Carbofuran**

Pre-column clean-up and LC determination of residues of N-methylcarbamate pesticides in extracts of total diet 404(1987)352

HPLC determination of aryl N-methylcarbamate residues using post-column hydrolysis electrochemical detection 442(1988)333

#### **Carbohydrate gums**

Identification of carbohydrate gums in dental cream by TLC 407(1987)369

#### **Carbohydrates**

see also Sugars and saccharides

Determination of reducing and non-reducing carbohydrates in food products by LC with post-column catalytic hydrolysis and derivatization. Comparison with refractive index detection 402(1987)127

Capillary column GC identification of sugars in honey as trimethylsilyl derivatives 410(1987)319

GC evaluation of hydrogen in the breath as a diagnostic method for the study of intestinal digestion and absorption of dietary carbohydrates in man 416(1987)119

Extracellular mammalian polysaccharides: glycosaminoglycans and proteoglycans (Review) 418(1987)187

HPLC analysis of oligosaccharides. I. Separation on an ion-exchange stationary phase of low cross-linking 440(1988)281

Preparative isolation of <sup>14</sup>C-labelled saccharose from a partly purified fraction of *Chlorella vulgaris* 442(1988)237

LC determination of carbohydrates with pulsed amperometric detection and a membrane reactor 447(1988)268

Carbohydrate-borate eluents for anion chromatography 448(1988)355

#### **$\beta$ -Carboline-3-carboxylic acid**

Determination of *n*-butyl- $\beta$ -carboline-3-carboxylate in aqueous brain extracts 431(1988)192

#### **$\beta$ -Carbolines**

Measurement of  $\beta$ -carbolines by HPLC with fluorescence detection 428(1988)229

Lipophilicity measurement of benzodiazepine-receptor ligands by reversed-phase LC. Comparison between HPLC and TLC 447(1988)404

HPLC identification of simple  $\beta$ -carboline alkaloids in specimens of *Heliconiini* butterflies 447(1988)432

#### **Carbon**

Flame ionization and flame thermionic detection of carbon and nitrogen in aquatic lipid and humic-type classes with an Iatroscan Mark IV 435(1988)350

#### **Carbon adsorbents**

GC study of microporous carbons 406(1987)139

#### **Carbon black**

Carbon black adsorbents modified with *n*-hydrocarbons and treated with an high-frequency low-temperature plasma 402(1987)87

#### **Carbon dioxide**

Temperature effect in packed microcolumn chromatography using a carbon dioxide mobile phase 402(1987)328

Resolution of racemic amides and phosphine oxides on a  $\beta$ -cyclodextrin-bonded stationary phase by subcritical fluid chromatography 405(1987)135

Retention in capillary supercritical fluid chromatography with carbon dioxide mobile phases containing polar modifiers 406(1987)223

Adsorption isotherms of ethyl acetate modifier on silica from supercritical carbon dioxide 409(1987)55

Formation of N-carboxy compounds during trimethylsilylation of amino acids in the presence of carbon dioxide 409(1987)383

GC technique to simultaneously quantitate the gases produced by intestinal microorganisms from fermentation mixtures 414(1987)405

GC of carbon monoxide and carbon dioxide. Choice of a calibration method (in French) 447(1988)234

#### **Carbon hybridization summaries**

Total molecular surface areas as a predictor for reversed-phase HPLC in various organotin systems. Importance and applications of carbon hybridization summaries 441(1988)229

#### **Carbon loading**

Selectivity related to carbon loading and end-capping of octadecyl-bonded silica gels in reversed-phase LC 441(1988)183

#### **Carbon monoxide**

Correlation of carbon monoxide and bilirubin production by tissue homogenates 427(1988)315

GC of carbon monoxide and carbon dioxide. Choice of a calibration method (in French) 447(1988)234

#### **Carbonyl compounds**

HPLC determination of glyoxylic acid and other carbonyl compounds in urine 417(1987)253

Automated solid-phase catalyzed pre-column derivatization of fatty acids for reversed-phase HPLC analysis with fluorescence detection 436(1988)437

#### **17 $\beta$ -Carboxamide steroids**

HPLC in the evaluation of the lipophilicity of 17 $\beta$ -carboxamide steroid derivatives 445(1988)409

#### **2-Carboxyarticaïne**

HPLC and preliminary pharmacokinetics of articaïne and its 2-carboxy metabolite in human serum and urine 424(1988)440

#### **N-Carboxy compounds**

Formation of N-carboxy compounds during trimethylsilylation of amino acids in the presence of carbon dioxide 409(1987)383

#### **S-(2-Carboxyethyl)-N-(2-*tert.*-butyl-5-methoxybenzthiazol-6-yl) dithiocarbamate**

HPLC of antifilarials. II. Determination of S-(2-carboxyethyl)-N-(2-*tert.*-butyl-5-methoxybenzthiazol-6-yl) dithiocarbamate and its active isothiocyanate metabolite in biological fluids 434(1988)296

**$\gamma$ -Carboxyglutamic acid**

Isotachopheretic analysis for the determination of urinary  $\gamma$ -carboxyglutamic acid 419(1987)334

**Carboxyhaemoglobin**

Effects of differences in charge and hydrophobicity of surface amino acids of hemoglobins on high-performance GPC 428(1988)247

**Carboxylic acids**

see also Organic acids

New derivatization reagents for the resolution of carboxylic acid enantiomers by HPLC 402(1987)344

Simultaneous determination of the acid chloride S-benzoyl-3-mercapto-2-methylpropanoyl chloride and residual carboxylic acid by capillary GC 408(1987)360

Benzyl esters of  $C_2$ - $C_{20}$  fatty acids and metabolically relevant carboxylic acids. Preparation and GC-MS 409(1987)43

Comparison of volatile components between fresh and rotten mussels by GC-MS 409(1987)251

Ion-exclusion chromatography of non-ionic substances with conductivity detection 409(1987)271

Elution behaviour of aliphatic carboxylic acids on a strong cation-exchange resin column 410(1987)103

HPLC separation of femtomolar quantities of endogenous carboxylic acids, including arachidonic acid metabolites, as 4-bromomethyl-7-acetoxycoumarin derivatives 416(1987)247

New method of purification of perchloric acid extracts for quantitative GC of some Krebs cycle acids 424(1988)357

Simultaneous determination of  $C_2$ - $C_{22}$  non-esterified fatty acids and other metabolically relevant carboxylic acids in biological material by GC of their benzyl esters 425(1988)257

Studies of metabolites in diarrheal stool specimens positive for *Klebsiella*, *Serratia*, and *Proteus* spp. by frequency-pulsed electron-capture GC 430(1988)209

Improved procedure for preparation of pentafluorobenzyl derivatives of carboxylic acids for analysis by GC with electron-capture detection 433(1988)248

Peak enhancement of carboxylic acids in ion-

exclusion chromatography with a cation-exchange hollow-fibre suppressor using neutral and alkaline enhancers 435(1988)417

Identification of aliphatic keto carboxylic acids in marine aerosols using capillary GC-MS 438(1988)299

Ion chromatographic determination of sugar phosphates in physiological samples 439(1988)83

Computer-controlled linear pH gradient for HPLC fractionations of aromatic carboxylic acids and of humic and fulvic acids 439(1988)213

GC procedure for the qualitative analysis of silicone polyester resins after alkaline fusion 441(1988)207

1-(4-Hydroxyphenyl)-, 1-(2,4-dihydroxyphenyl)- and 1-(2,5-dihydroxyphenyl)-2-bromoethanones: new labels for determination of carboxylic acids by HPLC with electrochemical and UV detection 442(1988)209

$\beta$ -Cyclodextrin as a selective agent for separation of selected aromatic acids by HPLC 445(1988)337

Chromatography of microbial metabolites of aromatic amino acids 446(1988)117

GC analysis of different homologous series of acids esterified in aqueous solutions with butyl and propyl alcohols 446(1988)237

GLC-MS studies on pine Kraft black liquors. III. The liberation of carboxylic acids in the initial phase of pulping 446(1988)247

Studies on sample preconcentration in ion chromatography. VIII. Preconcentration of carboxylic acids prior to ion-exclusion separation 447(1988)155

Extractive pentafluorobenzoylation of formic, acetic, levulinic, benzoic and phthalic acids, studied by LC and dual-oven capillary GC 447(1988)329

Ion-exclusion chromatography of carboxylic acids with conductivity detection. Peak enhancement using a cation-exchange hollow-fibre membrane and an alkaline solution 447(1988)373

**Carboxylic acids, perfluorinated**

Use of perfluorinated carboxylic acids in the separation of metabolites of vasopressin prior to radioimmunoassay 430(1988)128

**Carboxylic esters**

Hierarchical ascending classification of Kováts retention indices as a means of indicating siloxane



stationary phase specificities towards aliphatic carboxylic esters 406(1987)95

### **S-Carboxymethylcysteine**

Isotachophoretic analysis of amoxicillin and carboxymethylcysteine in pharmaceuticals 408(1987)441

Determination of S-carboxymethyl-L-cysteine in plasma by HPLC with column switching following precolumn derivatization 417(1987)208

Determination of S-carboxymethylcysteine in serum by reversed-phase ion-pair LC with column switching following pre-column derivatization with *o*-phthalaldehyde 428(1988)301

### **3-Carboxy-4-methyl-5-propyl-2-furanpropanoic acid**

Determination of 3-carboxy-4-methyl-5-propyl-2-furanpropanoic acid, a major endogenous ligand substance in uremic serum, by HPLC with UV detection 415(1987)110

### **Carboxypeptidase P digestion**

Narrow-bore HPLC of phenylthiocarbamyl amino acids and carboxypeptidase P digestion for protein C-terminal sequence analysis 447(1988)351

### **Carboxyphosphamide**

Combined TLC-photography-densitometry for the quantitation of cyclophosphamide and its four principal urinary metabolites 427(1988)121

### **2-Carboxypiperidine**

Separation of hydroxy and polyhydroxy derivatives of 2-carboxy and 2-hydroxymethyl piperidine and pyrrolidine by HPLC 445(1988)295

### **2-[4-(2-Carboxypropyl)phenyl]propionic acid**

Determination of ibuprofen and its major metabolites in human urine by HPLC 430(1988)93

### **Cardiolipin**

TLC of human platelet phospholipids with fatty acid analysis 420(1987)411

### **3-Carene**

Resolution of monoterpene enantiomers by GC 407(1987)243

### **Carnitine**

Determination of total carnitine in human urine by HPLC 420(1987)385

Analytical method for urinary glutarylcarnitine, acetylcarnitine and propionylcarnitine with a carboxylic acid analyser and a reversed-phase column 423(1987)297

Determination of carnitine by HPLC using 9-anthryldiazomethane 445(1988)175

### **Carnosine**

Simultaneous determination of histidine-containing dipeptides, histamine, methylhistamine and histidine by HPLC 432(1988)315

### **$\beta$ -Carotene**

Chromatography of fat-soluble vitamins in clinical chemistry (Review) 429(1988)3

Supplementary and revised assignment of the peaks of the 7,9-, 9,9'-, 13,13'-, 9,13'-di-*cis* and 9,9',13-tri-*cis* isomers of  $\beta$ -carotene in HPLC using a column of calcium hydroxide 439(1988)417

### **Carotenes**

Normal-phase HPLC of carotenes 442(1988)412

### **Carotenoid esters**

Simultaneous HPLC analysis of free carotenoids and carotenoid esters 403(1987)207

### **Carotenoids**

Simultaneous HPLC analysis of free carotenoids and carotenoid esters 403(1987)207

Identification of *cis*-carotenoids in human sera analyzed by reversed-phase HPLC with diode array detection 431(1988)424

Separation and quantitative analysis of some carotenoid fatty acid esters of fruits by LC 435(1988)113

HPLC of carotenoids from some marine shellfish 440(1988)37

LC profiles of major carotenoid esters in commercially processed California navel and Valencia orange juice concentrates 442(1988)249

Isolation of *cis-trans* isomers of canthaxanthin by HPLC using a calcium hydroxide column and identification of their configurations by  $^1\text{H}$  NMR spectroscopy 448(1988)182

HPLC of chloroplast pigments. One-step separation of carotene and xanthophyll isomers, chlorophylls and pheophytins 448(1988)411

HPLC analysis of major carotenoids from *Rhodotorula glutinis* 448(1988)445

**Carotenol fatty acid esters**

Separation of carotenol fatty acid esters by HPLC 449(1988)119

**Carprofen**

Assay method for product formation in *in vitro* enzyme kinetic studies of uridine diphosphate glucuronyltransferases: 2-arylpropionic acid enantiomers 430(1988)368

Measurement of carprofen enantiomer concentrations in plasma and urine using L-leucinamide as the chiral coupling component 433(1988)331

**Carrageenans**

Characterisation of carrageenans by high-performance size-exclusion chromatography using a LiChrospher 1000 DIOL column 403(1987)307

**Carrier ampholytes**

Ionic binding properties of carrier ampholytes 437(1988)147

**Carrier gas flow-rate**

Influence of the carrier gas flow-rate on the pulsed electron-capture detector 446(1988)329

**Carthamin**

HPLC of carthamin, safflor yellow A and a precursor of carthamin. Application to the investigation of an unknown red pigment produced in cultured cells of safflower 438(1988)61

**Cartridges**

Evaluation of disposable cartridges for trace enrichment from aqueous solutions 402(1987)201

**Carumonam**

Determination of the monocyclic  $\beta$ -lactam antibiotic carumonam in plasma and urine by ion-pair and ion-suppression reversed-phase HPLC 420(1987)357

**Carvedilol**

HPLC method for the determination of carvedilol and its desmethyl metabolite in body fluids 413(1987)355

**Casein**

Detection by HPLC of polyamines formed by clostridal putrefaction of caseins 442(1988)267

**Caseins**

Rapid preparative-scale purification of peptides

derived from caseins using mass ion-exchange chromatography 450(1988)420

**Catalpol**

Application of various planar chromatographic techniques for the separation of iridoid glycosides from *Veronica officinalis* 404(1987)365

**Catalysis**

Contributions of GC to the development of heterogeneous catalysis 406(1987)151

**Catechin**

Analysis of catechin and epicatechin by HPLC after benzylation 448(1988)291

**Catecholamines**

Determination of catecholamines in urine (and plasma) by LC after on-line sample pretreatment on small alumina or dihydroxyborylsilica columns 414(1987)285

Identification and quantification of N-methylepinephrine in human urine 414(1987)301

Simultaneous LC determination of seventeen of the major monoamine neurotransmitters, precursors and metabolites. I. Optimization of the mobile phase using factorial designs and a computer program to predict chromatograms 415(1987)261

Simultaneous LC determination of seventeen of the major monoamine neurotransmitters, precursors and metabolites. II. Assessment of human brain and cerebrospinal fluid concentrations 415(1987)275

Determination of catecholamines in biological tissues by LC with coulometric detection 415(1987)388

Improved HPLC method with electrochemical detection for the determination of urinary catecholamines using a new electrochemical detector 416(1987)340

Application of a novel cation-exchange reagent, Igepon T-77 N-methyl oleoyl taurate, to microbore separations of alumina extracts of catecholamines from cerebrospinal fluid, plasma, urine and brain tissue with amperometric detection 420(1987)241

Combined ion-pair extraction and HPLC for the determination of the biogenic amines and their major metabolites in single brain tissue samples 421(1987)51

Single-step method for derivatization of

dopamine and some related compounds in aqueous media for GC 421(1987)227

Simple HPLC method for the simultaneous measurement of vanillylmandelic acid and homovanillic acid 423(1987)278

Routine determination of urinary free catecholamines by HPLC with electrochemical detection 424(1988)263

Simultaneous determination of the major acidic metabolites of catecholamines and serotonin in urine by LC with electrochemical detection after a one-step sample clean-up on Sephadex G-10; influence of vanilla and banana ingestion 424(1988)273

LC determination of urinary catecholamines after one-step alumina extraction 424(1988)373

Determination of catecholamines in body fluids 427(1988)359

Determination of catecholamines in body fluids 427(1988)360

Catecholamines and their metabolites (Review) 429(1988)177

Species-dependent differences in recovery of 3,4-dihydroxybenzylamine in assays of plasma catecholamines 430(1988)123

Simultaneous determination of endogenous catechols and exogenous 2- and 6-fluorinated catechols in tissue and plasma using LC with electrochemical detection 431(1988)156

Effects of long-term storage on the concentrations of the unconjugated acidic metabolites of the trace amines, indoleamines and catecholamines 433(1988)23

Selected-ion monitoring GC-MS analysis of catecholamines: enhancement of sensitivity by a simple clean-up step on Sephadex G-10 433(1988)31

Sensitive method for determination of picogram amounts of epinephrine and other catecholamines in microdissected samples of rat brain using LC with electrochemical detection 433(1988)41

Simultaneous LC determination of norepinephrine, 3,4-dihydroxyphenylethyleneglycol, 3,4-dihydroxyphenylalanine and 3,4-dihydroxyphenylacetic acid in human plasma 434(1988)187

### **Catechols**

Simultaneous determination of endogenous

catechols and exogenous 2- and 6-fluorinated catechols in tissue and plasma using LC with electrochemical detection 431(1988)156

Retention of ionic and non-ionic catechols in capillary zone electrophoresis with micellar solutions 441(1988)299

### **Catharanthine**

Determination of *Catharanthus* alkaloids by reversed-phase HPLC 410(1987)488

### **Cathedulines**

Efficient isolation of insect growth inhibitory macrolide alkaloids using recycle high-performance GPC 402(1987)354

### **Cation exchangers**

Adsorption isotherms on HPLC sorbents. II.

Proteins on cation exchangers with silica support 406(1987)285

Degradation of ketoses during aqueous HPLC on lead-form cation-exchange resins 437(1988)387

### **Cations**

Determination of cations at the parts-per-billion level with non-suppressed ion chromatography and indirect photometric detection 404(1987)282

Contamination of aqueous eluents due to corrosion of stainless-steel chromatographic components 407(1987)133

Coordinative interactions in the separation of insulin and derivatives by HPLC 408(1987)43

Eluent selection for the determination of cations in ion chromatography 439(1988)49

Determination of mono- and divalent cations and anions in small fog samples by ion chromatography 439(1988)121

### **Cefaclor**

Determination of cefaclor and cephadrine in serum by ion-pair reversed-phase chromatography 413(1987)351

Retention properties of internal-surface reversed-phase silica packing and recovery of drugs from human plasma 420(1987)297

### **Cefadroxil**

Determination of cefadroxil in serum by HPLC with cephradine as internal standard 413(1987)347

### **Cefaloglycine**

Determination of cefaloglycine and cefroxadine

in biological media by TLC with fluorimetric detection (in French) 432(1988)407

#### **Cefazolin**

Analysis of sodium cefazolin in ophthalmic ointment by HPLC 402(1987)376

#### **Cefetamet**

Determination of cefetamet and its orally active ester, cefetamet pivoxyl, in biological fluids by HPLC 430(1988)81

#### **Cefetamet pivoxyl**

Determination of cefetamet and its orally active ester, cefetamet pivoxyl, in biological fluids by HPLC 430(1988)81

#### **Cefixime**

Determination of cefixime in biological samples by reversed-phase HPLC 422(1987)145

Application of HPLC of some antibiotics in clinical microbiology 427(1988)257

#### **Cefodizime**

Determination of cefodizime in biological materials by HPLC 420(1987)329

#### **Cefotaxime**

Determination of serum cefotaxime and desacetylcefotaxime by ion-pair HPLC (in French) 419(1987)396

HPLC assay of cefotaxime, desacetylcefotaxime and ceftriaxone in rat plasma 424(1988)111

#### **Cefpiramide**

Column LC determination of cefpiramide in human serum and urine 417(1987)452

Micromethod for the determination of cefpiramide in human plasma and urine by HPLC using automated column switching 419(1987)388

Retention properties of internal-surface reversed-phase silica packing and recovery of drugs from human plasma 420(1987)297

#### **Cefroxadine**

Determination of cefaloglycine and cefroxadine in biological media by TLC with fluorimetric detection (in French) 432(1988)407

#### **Ceftetrame**

Quantitation of ceftetrame in human plasma and urine by HPLC 422(1987)135

#### **Ceftriaxone**

HPLC assay of cefotaxime, desacetylcefotaxime and ceftriaxone in rat plasma 424(1988)111

#### **Cellobiose**

HPLC separation of some mono- and disaccharides with detection by a post-column enzyme reactor and a chemically modified electrode 408(1987)157

#### **Cells**

Electrophoretic separation of cells (Review) 418(1987)359

#### **Cell separation**

Cell separation by immunoaffinity partitioning with polyethylene glycol-modified protein A in aqueous polymer two-phase systems 442(1988)219

#### **Cell surface hydrophobicity**

Difference in cell surface hydrophobicity of *Halobacterium salinarum* strains 449(1988)322

#### **Cellulose**

Adsorption chromatography on cellulose. I. Salting-out chromatography of organic compounds 406(1987)389

Adsorption chromatography on cellulose. II. Separations of aromatic amino acids, biogenic amines, alkaloids, dyes and phenols and determination of hydrophobic constants 440(1988)165

#### **Cellulose triacetate**

Computer-aided optimisation of drug enantiomer separation in chiral HPLC 434(1988)377

Direct enantiomeric separation of racemic flavanones by HPLC using cellulose triacetate as a chiral stationary phase 441(1988)417

#### **Centrifugal partition chromatography**

Separation of natural products by centrifugal partition chromatography 450(1988)91

#### **Centrifugal thin-layer chromatography**

Centrifugal TLC of alkaloids from vegetable sources using an aluminium oxide layer 437(1988)328

#### **Centrifuge**

Multi-stage mixer-settler planet centrifuge. Preliminary studies on partition of macromolecules with organic-aqueous and aqueous-aqueous two-phase solvent systems 437(1988)121

#### **Cephalexin**

Microbore LC determination of cadralazine and cephalixin in plasma with large-volume injection 426(1988)335

Analysis of cephalixin from canine skin biopsy by LC with UV-visible photodiode-array detection 427(1988)103

HPLC assay of cephalixin in serum and urine 427(1988)162

### **Cephalosporinases**

Use of chromatofocusing for separation of  $\beta$ -lactamases. VIII. Analytical chromatofocusing of chromosomal cephalosporinases from four *Klebsiella* strains 403(1987)217

### **Cephalosporin C**

Displacement chromatography of biomolecules 439(1988)341

### **Cephalosporins**

Hospital routine analysis of penicillins, third-generation cephalosporins and aztreonam by conventional and high-speed HPLC 413(1987)109

Determination of cefadroxil in serum by HPLC with cephadrine as internal standard 413(1987)347

Determination of cefaclor and cephadrine in serum by ion-pair reversed-phase chromatography 413(1987)351

Column LC determination of cefpiramide in human serum and urine 417(1987)452

Micromethod for the determination of cefpiramide in human plasma and urine by HPLC using automated column switching 419(1987)388

Determination of serum cefotaxime and desacetylcefotaxime by ion-pair HPLC (in French) 419(1987)396

Quantitation of cefetrame in human plasma and urine by HPLC 422(1987)135

Determination of cefixime in biological samples by reversed-phase HPLC 422(1987)145

HPLC assay of cefotaxime, desacetylcefotaxime and ceftriaxone in rat plasma 424(1988)111

Determination of  $\beta$ -lactams and their biosynthetic intermediates in fermentation media by pre-column derivatization followed by fluorescence detection 424(1988)325

Analysis of cephalixin from canine skin biopsy by LC with UV-visible photodiode-array detection 427(1988)103

Application of HPLC of some antibiotics in clinical microbiology 427(1988)257

Determination of cefetamet and its orally active

ester, cefetamet pivoxyl, in biological fluids by HPLC 430(1988)81

Determination of cefaloglycine and cefroxadine in biological media by TLC with fluorimetric detection (in French) 432(1988)407

### **Cephadrine**

Determination of cefaclor and cephadrine in serum by ion-pair reversed-phase chromatography 413(1987)351

### **Cereal proteins**

Chromatography of complex protein mixtures (Review) 418(1987)115

### **Cerebrospinal fluid proteins**

Cerebrospinal fluid protein analysis in diseases of the nervous system (Review) 429(1988)345

### **Cervical mucus**

Analysis of the aqueous phase of human cervical mucus by reversed-phase HPLC and capillary isotachopheresis 431(1988)317

### **C 1 esterase inhibitor**

Chromatographic procedure for preparation of a high-purity C 1 esterase inhibitor concentrate 433(1988)363

### **Cetirizine**

GC method for the determination of cetirizine in plasma 430(1988)149

### **Cetrimide**

System peaks in ion chromatography using cetrimide-coated columns and indirect photometric detection 402(1987)211

Anion chromatography using a coated PRP-1 column and eluents of pH > 7 447(1988)315

### **Cetyltrimethylammonium bromide**

Study of the derivatization of *n*-alkylamines with 1-fluoro-2,4-dinitrobenzene in the presence of aqueous cetyltrimethylammonium bromide micelles 426(1988)267

HPLC on dynamically modified silica. VIII. Gradient elution using eluents containing cetyltrimethylammonium bromide 447(1988)182

### **Characterizing probes**

Reduction of the number of characterizing probes in GC by factor analysis 402(1987)318

### **Charge-transfer complexes**

Chromatographic study of charge-transfer complexes 406(1987)353

**Chemical engineering**

Chemical engineering approach to dynamic modelling of linear chromatography. A flexible method for representing complex phenomena from simple concepts 406(1987)11

**Chemical heterogeneity**

Comparison between SEC and liquid adsorption chromatography in the determination of the chemical heterogeneity of copolymers 411(1987)355

**Chemical interactions**

Chemical interactions as a possible limitation on the useful solvent properties of liquid alkylammonium salts 435(1988)17

**Chemical reaction detector**

Evaluation of a chemical reaction detector with electrochemical production of bromine for the LC determination of physostigmine 411(1987)229

**Chemiluminescence detector**

Coupling LC with redox chemiluminescence detection 441(1988)125

**Chemiluminogenic labels**

HPLC of chemiluminogenic labels with sensitive photoplate detection 437(1988)268

**Chenodeoxycholate**

Studies on steroids. CCXXXIII. Separation and characterization of bile acid disulphates in human urine by HPLC 425(1988)59

**Chenodeoxycholic acid**

Analysis of conjugated bile acids in human serum by HPLC using post-column enzyme reaction and off-line fluorimetric determination 423(1987)75

**Chimassorb 944**

Determination of the polymeric light stabilizer Chimassorb 944 in polyolefins by reversed-phase HPLC 450(1988)430

**Chiral counter ion**

Influence of enantiomeric purity of a chiral selector on stereoselectivity 407(1987)217

**Chiral recognition**

Reciprocity in chiral recognition. Comparison of several chiral stationary phases 404(1987)107

Chiral recognition of halo carboxylic acid amides studied by GC on Chirasil-Val 406(1987)157

Structural factors affecting chiral recognition and

separation on  $\beta$ -cyclodextrin bonded phases 441(1988)376

Micro-LC and the chiral recognition mechanism on albumin-coated silica gel. Large selectivity changes with sample size 447(1988)297

Chromatographic separation of the enantiomers of 1,3-dithiolane-1-oxides 450(1988)199

**Chiral recognition mechanism**

Resolution of enantiomeric aromatic alcohols on a cellulose tribenzoate HPLC chiral stationary phase. A proposed chiral recognition mechanism 411(1987)139

**Chiral recognition model**

Stereoelectronic model to explain the resolution of enantiomeric ibuprofen amides on the Pirkle chiral stationary phase 402(1987)179

**Chiral stationary phases, GC**

see also Stationary phases

Polysiloxanes containing thermally stable chiral amide side-chains for capillary GC and supercritical fluid chromatography 405(1987)169

Chiral recognition of halo carboxylic acid amides studied by GC on Chirasil-Val 406(1987)157

Enantiomer analysis by complexation GC. Scope, merits and limitations 441(1988)135

**Chiral stationary phases, LC**

Stereoelectronic model to explain the resolution of enantiomeric ibuprofen amides on the Pirkle chiral stationary phase 402(1987)179

HPLC separation of diacylglycerol enantiomers on a chiral stationary phase 402(1987)257

Separation of racemic pharmaceuticals by HPLC on silica gel modified with carbohydrate residues 403(1987)263

HPLC separation of enantiomers on axially chiral binaphthalene derivatives bonded to silica gel 403(1987)275

Reciprocity in chiral recognition. Comparison of several chiral stationary phases 404(1987)107

Optical resolution of enantiomeric amino acid derivatives on a naphthylethylurea multiple-bonded chiral stationary phase prepared via an activated carbamate intermediate 404(1987)117

Resolution of  $\alpha$ -arylacetic acid enantiomers on two chiral stationary phases 404(1987)123

Chiral-phase HPLC of rotenoid racemates 404(1987)133

- Chromatographic separation of racemic amino acids by use of chiral crown ether-coated reversed-phase packings 405(1987)145
- Enantiomeric separation of amino acids by micro HPLC on an L-proline-bonded stationary phase 407(1987)151
- Applications for racemic versions of chiral stationary phases 407(1987)211
- Effects of pretreatment on the enantioselectivity of silica-bound proteins used as HPLC stationary phases 408(1987)99
- Application of an ovomucoid-conjugated column for the optical resolution of some pharmaceutically important compounds 408(1987)316
- Resolution of a series of racemic phosphine oxides by LC on two structurally similar chiral stationary phases (in French) 409(1987)183
- Resolution of enantiomeric aromatic alcohols on a cellulose tribenzoate HPLC chiral stationary phase. A proposed chiral recognition mechanism 411(1987)139
- Direct resolution of the stereoisomers of leucovorin and 5-methyltetrahydrofolate using a bovine serum albumin HPLC chiral stationary phase coupled to an achiral phenyl column 424(1988)158
- Direct separation of drug enantiomers by HPLC with chiral stationary phases (Review) 426(1988)1
- Enantiomeric analysis of a new anti-inflammatory agent in rat plasma using a chiral  $\beta$ -cyclodextrin stationary phase 426(1988)417
- Computer-aided optimisation of drug enantiomer separation in chiral HPLC 434(1988)377
- Improved resolution of enantiomers of naproxen by the simultaneous use of a chiral stationary phase and a chiral additive in the mobile phase 435(1988)225
- Chiral separation through a single substituent association 436(1988)517
- Separation of fatty acid ester and amide enantiomers by HPLC on chiral stationary phases 437(1988)109
- Optical resolution of samples with weak interactions on chiral "brush-type" LC stationary phases 440(1988)197
- An improved chiral stationary phase for the facile separation of enantiomers 441(1988)311
- Structural factors affecting chiral recognition and separation on  $\beta$ -cyclodextrin bonded phases 441(1988)376
- Micro-LC and the chiral recognition mechanism on albumin-coated silica gel. Large selectivity changes with sample size 447(1988)297
- Hexahelicene chiral stationary phase. I. Phase synthesis and use in HPLC resolution of enantiomers 450(1988)450(1988)157
- Chiral stationary phase derived from L-lactic acid for the optical resolution of N-(3,5-dinitrobenzoyl)amino acid methyl esters 450(1988)163
- Separation and identification of enantiomers by HPLC with a chiral column and a polarimetric detector as applied to deltamethrin 450(1988)169
- Direct HPLC resolution of dihydropyridine enantiomers 450(1988)191
- Chiral resolution of a series of 3-thienylcyclohexylglycolic acids by liquid or subcritical fluid chromatography. A mechanistic study 450(1988)255
- Chromatographic resolution of dipeptide enantiomers and diastereomers on chiral stationary phases from poly(L-leucine) or poly(L-phenylalanine) 450(1988)271
- Chiral stationary phases, SFC**  
Polysiloxanes containing thermally stable chiral amide side-chains for capillary GC and supercritical fluid chromatography 405(1987)169
- Chiral stationary phases, TLC**  
Rapid separation of anomeric purine nucleosides by TLC on a chiral stationary phase 410(1987)226
- Applicability of forced-flow planar chromatographic methods for the separation of enantiomers on Chiralplate 450(1988)241
- Chloral hydrate**  
Determination of trichloroethylene metabolites in rat liver homogenate using headspace GC 442(1988)325
- Chlorambucil**  
HPLC of chlorambucil analogues 422(1987)315
- Chloramination**  
Comparison of volatile halogenated compounds formed in the chloramination and chlorination of humic acid by GC-electron-capture detection 449(1988)326

**Chloramphenicol**

Determination of chloramphenicol residues by reversed-phase HPLC. Application to a pharmacokinetic study in rainbow trout with confirmation by MS (in French) 409(1987)315

Determination of chloramphenicol in swine muscle tissue using a monoclonal antibody-mediated clean-up procedure 411(1987)415

HPLC determination of chloramphenicol and four analogues using reductive and oxidative electrochemical and UV detection 417(1987)111

**Chlorbenzoxamine**

Screening procedure for the detection of alkanolamine antihistamines and their metabolites in urine using computerized GC-MS 428(1988)43

**Chloride**

Simultaneous determination of sodium and chloride ions by ion chromatography at the  $\mu\text{g}/\text{l}$  level 404(1987)285

Optimization of ion chromatography using commercially available detection systems and software 439(1988)13

**Chlorinated alkylbenzenes**

The mass-selective detector as a chlorine-selective detector 438(1988)233

**Chlorinated carbanilides**

LC behaviour of chlorinated carbanilides 410(1987)95

**Chlorinated hydrocarbons**

Analysis of hydrocarbons in aquatic sediments. II. Evaluation of common preparative procedures for petroleum and chlorinated hydrocarbons 436(1988)503

HPLC method for isolating organic contaminants from tissue and sediment extracts 437(1988)161

Separation of organochlorines on alumina 437(1988)254

**Chlorinated organic compounds**

High-pressure SEC of chlorinated and unchlorinated aquatic fulvic acids using a polyacrylamide gel 450(1988)433

**Chlorination**

Comparison of volatile halogenated compounds formed in the chloramination and chlorination of humic acid by GC-electron-capture detection 449(1988)326

**Chlorine-containing compounds**

Spectra and response linearization of chlorine and bromine compounds in a copper-sensitized flame photometric detector 408(1987)69

**Chlormethiazole**

Improved HPLC procedure for the determination of chlormethiazole levels following solid-phase extraction from plasma 415(1987)357

**Chloroaluminum phthalocyanine tetrasulphonate**

Reversed-phase LC determination of chloroaluminum phthalocyanine tetrasulfonate in canine tissues and fluids 433(1988)367

**Chlorobenzaldehyde**

Retention behaviour of aromatic aldehyde derivatives in reversed-phase LC 404(1987)242

**2-(4-Chlorobenzoylamino)-3-[2(1H)-quinolin-4-yl]propionic acid**

HPLC procedure for the determination of a new anti-gastric ulcer agent, 2-(4-chlorobenzoylamino)-3-[2(1H)-quinolin-4-yl]propionic acid, in human plasma and urine 434(1988)283

**2-Chlorobutyroanilide**

HPLC methods for the determination of contaminants in technical phosmethylan 435(1988)380

**Chlorodesmethyldiazepam**

Analysis of chlorodesmethyldiazepam and its metabolites in plasma and urine 420(1987)417

**(E)-3-(9-Chloro-6,11-dihydro-5H-pyrrolo(2,1-B)(3) benzazepin-11-ylidene)-N,N-dimethyl-1-propana**

Determination of picogram levels of (E)-3-(9-chloro-6,11-dihydro-5H-pyrrolo(2,1-B)(3) benzazepin-11-ylidene)-N,N-dimethyl-1-propanamine (Z)-2-butenedioate (1:1) in plasma using capillary GC with nitrogen-selective detection 413(1987)277

**2-Chloroethyl ethyl sulphide**

Identification of degradation products of 2-chloroethyl ethyl sulfide by GC-MS 447(1988)165

**N'-(2-Chloroethyl)-N-[2-(methylsulphinyl)ethyl]-N'-nitrosoarea**

Qualitative and quantitative analysis of a new



sulphur-containing nitrosoarea in blood by HPLC 419(1987)381

**N'-(2-Chloroethyl)-N-[2-(methylsulphonyl)ethyl]-N'-nitrosoarea**

Qualitative and quantitative analysis of a new sulphur-containing nitrosoarea in blood by HPLC 419(1987)381

**Chloroform**

Headspace GC analysis for determining low levels of chloroform in human plasma 416(1987)125

Use of chloroform as a solvent for the formation of N-trifluoroacetyl-*n*-butylamino acid derivatives for GC analysis 436(1988)100

**Chlorophacinone**

Modification to procedures for the determination of chlorophacinone and for multi-residue analysis of rodenticides in animal tissues 437(1988)301

**Chlorophenols**

On-line post-column extraction in column LC with electron-capture detection. II. Applications 435(1988)297

GC determination of chlorinated phenols in the form of various derivatives 439(1988)307

Reversed-phase TLC of some chlorophenols in the presence of a soluble  $\beta$ -cyclodextrin polymer 439(1988)393

On-line trace enrichment on a reversed-phase pre-column for normal-phase LC with electron-capture detection 445(1988)129

**4-Chlorophenylbiguanide**

Determination of proguanil and its metabolites cycloguanil and 4-chlorophenylbiguanide in plasma, whole blood and urine by HPLC 416(1987)394

**6-(2-Chlorophenyl)-4-hydroxy-4H-imidazo[1,5-a]-[1,4]benzodiazepine-3-carboxamide**

Determination of the agonist-antagonist anxiolytic agent 6-(2-chlorophenyl)-4-hydroxy-4H-imidazo[1,5-a]-[1,4]benzodiazepine-3-carboxamide and its benzophenone carboxylic acid metabolite in plasma by HPLC 417(1987)347

**6-(2'-Chlorophenyl)-4-hydroxy-4H-imidazo[1,5-a] [1,4]benzodiazepine-3-carboxamide**

Determination of 6-(2'-chlorophenyl)-4-hydroxy-

4H-imidazo[1,5-a] [1,4]benzodiazepine-3-carboxamide, a mixed agonist-antagonist anxiolytic agent, in human plasma and urine by GC-negative-ion chemical-ionization MS 421(1987)281

**4-[2-(4-Chlorophenylsulfonylamino)ethyl]-phenylacetic acid**

Determination of a new thromboxane A<sub>2</sub> receptor blocker in biological fluids by capillary GC with electron-capture detection 419(1987)345

**(4-Chlorophenyl)thiomethylene bisphosphonic acid**

Determination of (4-chlorophenyl)thiomethylene bisphosphonic acid, a new bisphosphonate, in biological fluids by HPLC 430(1988)73

**Chlorophyllin, Cysteamine**

Improved method for the detection of peroxidase isoenzymes after electrophoretic separation with chlorophyllin, cysteamine and light as a hydrogen peroxide generating system 410(1987)466

**Chlorophylls**

Reversed-phase HPLC and column LC of chlorophylls and their derivatives 408(1987)181

HPLC of chloroplast pigments. One-step separation of carotene and xanthophyll isomers, chlorophylls and pheophytins 448(1988)411

**3-Chloropropanediol**

Determination of residual epichlorohydrin and 3-chloropropanediol in water by GC with electron-capture detection 439(1988)448

**6-Chloro-2-pyridylmethyl nitrate**

Determination of the anti-anginal drug 6-chloro-2-pyridylmethyl nitrate in plasma and urine by fused-silica open-tubular column GC with electron-capture detection 430(1988)388

**Chloroquine**

Field-adapted method for HPTLC detection and estimation of chloroquine and desethylchloroquine in urine 423(1987)261

Field-adapted method for HPTLC detection and estimation of chloroquine in finger-stick blood 428(1988)196

Detection and determination of antimalarial drugs and their metabolites in body fluids (Review) 434(1988)1

**Chlorosulphonic acid**

Spray reagent for the detection of coumarins and flavonoids on thin-layer plates 403(1987)285

**Chlorothiazide**

LC screening of diuretics in urine 415(1987)347

**Chlorphenesin**

Application of an ovomucoid-conjugated column for the optical resolution of some pharmaceutically important compounds 408(1987)316

**Chlorpheniramine**

Application of an ovomucoid-conjugated column for the optical resolution of some pharmaceutically important compounds 408(1987)316

Simultaneous analysis of acetaminophen, pseudoephedrine hydrochloride and chlorpheniramine maleate in a cold tablet using an isocratic, mixed micellar HPLC mobile phase 410(1987)206

Highly sensitive determination of chlorpheniramine as fluorescence derivative by HPLC 420(1987)63

**Chlorphenoxamine**

Screening procedure for the detection of alkanolamine antihistamines and their metabolites in urine using computerized GC-MS 428(1988)43

**Chlorprenaline**

Application of an ovomucoid-conjugated column for the optical resolution of some pharmaceutically important compounds 408(1987)316

**Chlorproethazine**

GC determination of chlorproethazine in plasma 419(1987)363

**Chlorpromazine**

GC determination of chlorpromazine in plasma 419(1987)363

HPLC drug analysis by direct injection of whole blood samples. III. Determination of hydrophobic drugs adsorbed on blood cell membranes 423(1987)163

Solid-phase extraction and HPLC method for chlorpromazine and thirteen metabolites 423(1987)207

**Chlorthalidone**

LC screening of diuretics in urine 415(1987)347

Simple, sensitive and selective HPLC method for analysis of chlorthalidone in whole blood 416(1987)420

**Cholate**

Studies on steroids. CCXXXIII. Separation and characterization of bile acid disulphates in human urine by HPLC 425(1988)59

**Cholecystokinin**

Peptide characterization with a sulfoethyl aspartamide column 443(1988)63

**Cholecystokinin antagonist**

Quantitative determination of L-364,718, a novel cholecystokinin antagonist, by solid-phase extraction and HPLC 437(1988)442

**Cholecystokinins**

Separation of protein hormones (Review) 418(1987)245

**Cholestanol**

Improved method for the determination of the major neutral steroids and unconjugated bile acids in human faeces using capillary GC 421(1987)21

Highly sensitive and simple determination of cholesterol and cholestanol in human serum by HPLC with fluorescence detection 421(1987)43

**Cholesterol**

Pregnenolone production by adrenal mitochondria: a new HPLC analytical method for cholesterol side-chain cleavage 414(1987)411

Improved method for the determination of the major neutral steroids and unconjugated bile acids in human faeces using capillary GC 421(1987)21

Highly sensitive and simple determination of cholesterol and cholestanol in human serum by HPLC with fluorescence detection 421(1987)43

Determination of coenzyme Q<sub>10</sub>,  $\alpha$ -tocopherol and cholesterol in biological samples by coupled-column LC with coulometric and UV detection 425(1988)87

Simultaneous microanalysis of bile acids and cholesterol in bile by glass capillary GC 431(1988)17

**Cholesterol esters**

Profiling of plasma cholesterol ester and triglyceride fatty acids as their methyl esters by capillary GC, preceded by a rapid aminopropyl-silica column chromatographic separation of lipid classes 434(1988)395

**Cholic acid**

Analysis of conjugated bile acids in human serum by HPLC using post-column enzyme reaction and off-line fluorimetric determination 423(1987)75

**Choline**

Determination of acetylcholine and choline in perchlorate extracts of brain tissue using LC-electrochemistry with an immobilized-enzyme reactor 414(1987)167

Determination of acetylcholine and choline in microdialysates from spinal cord of rat using LC with electrochemical detection 423(1987)85

LC with electrochemical detection for the determination of choline and acetylcholine in plasma and red blood cells. Failure to detect acetylcholine in blood of humans and mice 428(1988)1

Determination of acetylcholine and choline in human cerebrospinal fluid using HPLC combined with an immobilized enzyme reactor: ageing-induced change of acetylcholine level 431(1988)389

**Cholinesterases**

Quantitative monitoring by HPLC of the dissociation of human serum cholinesterase by limited proteolysis 420(1987)25

Rapid and efficient method for the isolation and characterization of plant aromatic choline esterases 450(1988)121

**Chondroitin sulphate**

High-performance GPC of glycosaminoglycans. Column calibration by gel electrophoresis 402(1987)149

Rapid and sensitive method for measurement of hyaluronic acid and isomeric chondroitin sulfates using HPLC 413(1987)9

Rapid determination of disaccharides from chondroitin and dermatan sulphates by HPLC 414(1987)156

LC assay for constituent disaccharides of hyaluronic acid and chondroitin sulphate isomers 415(1987)231

Extracellular mammalian polysaccharides: glycosaminoglycans and proteoglycans (Review) 418(1987)187

Fractionation of aorta glycosaminoglycans by HPLC 419(1987)275

Methods for analysis of urinary glycosaminoglycans (Review) 429(1988)293

**Chromarods**

Flame ionization and flame thermionic detection of carbon and nitrogen in aquatic lipid and humic-type classes with an Iatroscan Mark IV 435(1988)350

**Chromatofocusing**

Application of high-performance liquid chromatofocusing to the study of prostatic androgen receptors. Effect of stabilizing agents on the heterogeneity of receptor structure 403(1987)171

Use of chromatofocusing for separation of  $\beta$ -lactamases. VIII. Analytical chromatofocusing of chromosomal cephalosporinases from four *Klebsiella* strains 403(1987)217

Fractionation and characterization of two  $\beta$ -lactamases in *Citrobacter diversus* ULA-27 strain by chromatofocusing 403(1987)366

Resolution of five position-specific flavonoid O-methyltransferases by fast protein liquid chromatofocusing 407(1987)291

Separation of transcobalamin II isoproteins by means of chromatofocusing 432(1988)103

Microcolumn chromatofocusing. III. Effect of the analysis time on the resolution and the use of a semi-disposable microcolumn 438(1988)15

Purification of germinated barley  $\alpha$ -amylase isozymes and limit-dextrinase by chromatofocusing and affinity chromatography 441(1988)436

Chick 28000 *M*, vitamin D-dependent calcium-binding protein in intestine, kidney and cerebellum. Purification using chromatofocusing 448(1988)145

**Chromatophoresis**

Separation of proteins by reversed-phase HPLC. I. Optimizing the column 443(1988)363

**Chromenes**

Rapid analyses of chromenes (benzopyrans) by GC-MS 403(1987)358

Application of semi-preparative HPLC to

difficult natural product separations  
406(1987)367

#### **Chromium complexes**

Effect of the geometrical configuration of facial and meridional isomers of cobalt(III) and chromium(III) complexes on their  $R_F$  values  
404(1987)307

Anion-exchange HPLC of water-soluble chromium(VI) and chromium(III) complexes in biological materials 415(1987)317

LC retention of  $\beta$ -diketonato complexes of chromium(III) and palladium(II) in octadecyl-bonded silica/water-methanol systems. Comparison with liquid-liquid partition using dodecane as a model alkyl-bonded phase  
439(1988)245

#### **Chromophore, polymer-bound**

Absolute peak broadening calibration in SEC using a polymer-bound chromophore  
449(1988)63

#### **Chromosorb 101**

Characterization of some styrene-divinylbenzene sorbents for reversed-phase chromatography  
442(1988)97

#### **Chromosorb Century Series**

Characterization of porous polymers by polar strength and selectivity 404(1987)145

#### **Chromosorb P**

Polyester deactivation of pink supports and its effect on the GC of some polar, closely boiling and isomeric analytes 446(1988)31

#### **Chrysanthemols**

GC determination of racemic *cis*- and *trans*-chrysanthemols and their potential aldehyde and carboxylic acid microbial metabolites  
437(1988)436

#### **Chylomicrons**

Fractionation of baboon chylomicrons and very-low-density lipoproteins by HPLC 433(1988)257

#### **Chymotrypsin**

HPLC assay of chymotrypsin and chymotrypsin-like enzyme activity 411(1987)498

Fraction collector for capillary zone electrophoresis 438(1988)23

#### **CI 45350**

Identification of xanthene dyes in lipsticks by reversed-phase HPLC 448(1988)296

#### **Cicloprofen**

Assay method for product formation in *in vitro* enzyme kinetic studies of uridine diphosphate glucuronyltransferases: 2-arylpropionic acid enantiomers 430(1988)368

#### **Cimetidine**

Highly sensitive determination of cimetidine and its metabolites in serum and urine by HPLC  
422(1987)301

#### **Cimetidine sulphoxide**

Highly sensitive determination of cimetidine and its metabolites in serum and urine by HPLC  
422(1987)301

#### **Cineole**

Characterisation of volatile oil constituents with relatively long GC retention times on two stationary phases 449(1988)312

#### **Cinnamadehyde**

Determination of Cinnamomi cortex by HPLC  
409(1987)365

#### **Cinnamaldehyde**

Determination of cinnamaldehyde, coumarin and cinnamyl alcohol in cinnamon and cassia by HPLC 447(1988)272

#### **Cinnamic acid**

Analysis of manufacturing by-products and impurities in illicit cocaine via HPLC and photodiode array detection 405(1987)273

Determination of Cinnamomi cortex by HPLC  
409(1987)365

#### **Cinnamomi cortex**

Determination of Cinnamomi cortex by HPLC  
409(1987)365

#### **Cinnamomin**

Chromatographic purification and characterization of elicitors of necrosis on tobacco produced by incompatible *Phytophthora* species 440(1988)87

#### **Cinnamoylcocaine**

Analysis of manufacturing by-products and impurities in illicit cocaine via HPLC and photodiode array detection 405(1987)273

#### **Cinnamyl acetate**

Determination of Cinnamomi cortex by HPLC  
409(1987)365

**Cinnamyl alcohol**

Determination of Cinnamomi cortex by HPLC 409(1987)365

Determination of cinnamaldehyde, coumarin and cinnamyl alcohol in cinnamon and cassia by HPLC 447(1988)272

**Ciprofloxacin**

Assay of ciprofloxacin and norfloxacin in serum and urine by HPLC 416(1987)207

Sensitive and selective determination of picogram amounts of ciprofloxacin and its metabolites in biological samples using HPLC and photothermal post-column derivatization 416(1987)321

Determination of ciprofloxacin and its 7-ethylenediamine metabolite in human serum and urine by HPLC 419(1987)414

HPLC of ciprofloxacin and its metabolites in serum, urine and sputum 422(1987)153

**Ciprofloxacin**

Application of HPLC of some antibiotics in clinical microbiology 427(1988)257

**Circumsporozoite protein**

Assay, purification and characterization of a recombinant malaria circumsporozoite fusion protein by HPLC 411(1987)345

**C.I. Reactive Blue 2**

Design and applications of biomimetic anthraquinone dyes. I. Synthesis and characterisation of terminal ring isomers of C.I. Reactive Blue 2 435(1988)127

**Cisapride**

Determination of cisapride in plasma and animal tissues by HPLC 424(1988)195

**Cisplatin**

Analysis of platinum species originating from *cis*-diamminedichloroplatinum(II) (cisplatin) in human and rat plasma by HPLC with on-line inductively coupled plasma atomic emission spectrometric detection 407(1987)253

**Citral**

Characterisation of volatile oil constituents with relatively long GC retention times on two stationary phases 449(1988)312

**Citrate**

New method of purification of perchloric acid

extracts for quantitative GC of some Krebs cycle acids 424(1988)357

HPLC measurement of selected blood citric acid cycle intermediates 425(1988)47

Quantitation of oxalate and citrate by ion chromatography with a buffered, strong acid eluent 437(1988)231

 **$\beta$ -Citraurin**

Separation and quantitative analysis of some carotenoid fatty acid esters of fruits by LC 435(1988)113

**Citreoviridin**

Standardized HPLC of 182 mycotoxins and other fungal metabolites based on alkylphenone retention indices and UV-VIS spectra (diode array detection) 404(1987)195

**Citric acid**

Determination of ionic metabolites from ethylene glycol in human blood by isotachopheresis 416(1987)111

Excretion of urinary acids during inverted sleep-waking rhythm 422(1987)1

**Citrinin**

Standardized HPLC of 182 mycotoxins and other fungal metabolites based on alkylphenone retention indices and UV-VIS spectra (diode array detection) 404(1987)195

**Citrulline**

Determination of agmatine, arginine, citrulline and ornithine by reversed-phase LC using automated pre-column derivatization with *o*-phthalaldehyde 425(1988)269

Assignment of the configurations of the amino acids in peptidic siderophores 438(1988)291

**Claparède, E.**

Tswett's letters to Claparède 440(1988)509

**Clean-up methods**

Determination of chloramphenicol in swine muscle tissue using a monoclonal antibody-mediated clean-up procedure 411(1987)415

Direct determination of dextromethorphan and its three metabolites in urine by HPLC using a precolumn switching system for sample clean-up 422(1987)340

HPLC method for isolating organic contaminants from tissue and sediment extracts 437(1988)161

**Clebopride**

Determination of clebopride in plasma by capillary GC–negative-ion chemical ionization MS 432(1988)153

**Clemastine**

Screening procedure for the detection of alkanolamine antihistamines and their metabolites in urine using computerized GC–MS 428(1988)43

**Clenbuterol**

Determination of clenbuterol and mabuterol in equine plasma by ion-pair LC with electrochemical detection. Chromatographic and electrochemical characteristics 441(1988)197

**Climazolam**

Determination of climazolam in the plasma of farm animals by GC 415(1987)170

**Clindamycin**

Determination of clindamycin in plasma or serum by HPLC with UV detection 431(1988)379

**Clobazam**

Micro-determination of clobazam and N-desmethylclobazam in plasma or serum by electron-capture GC 421(1987)147

**Clomesone**

Analysis of clomesone in plasma by GC–electrolytic conductivity detection 428(1988)331

**Clomipramine**

Determination of clomipramine and its hydroxylated and demethylated metabolites in plasma and urine by LC with electrochemical detection 416(1987)311

Simultaneous determination of clomipramine and its N-desmethyl metabolite in human whole blood by capillary GC with mass-selective detection 428(1988)71

**Clonazepam**

Selective effect of clonazepam and (*S*)-uxepam on the binding of warfarin enantiomers to human serum albumin 450(1988)217

**Clonidine**

Determination of clonidine in human plasma by cold on-column injection capillary GC–selected-ion monitoring–MS 432(1988)165

**Cloпамide**

LC screening of diuretics in urine 415(1987)347

**Clopyralid**

Analysis of the herbicides dicamba, clopyralid and bromacil in asparagus by HPLC 439(1988)470

***Clostridium difficile* toxin B**

Fast protein purification of *Clostridium difficile* cytotoxin 428(1988)408

***Clostridium innocuum***

Identification by GC–MS of short-chain hydroxy acids produced by *Fusobacterium* species and *Clostridium innocuum* 420(1987)121

**Cloxacillin**

Optimization of HPLC analysis for isoxazolyl penicillins using factorial design 425(1988)331

**Clozapine**

Clozapine plasma levels determined by HPLC with UV detection 428(1988)160

Determination of clozapine in human serum by capillary GC 434(1988)465

**Cluster analysis**

Chemometric classification of reversed-phase HPLC columns 410(1987)31

**Clycoproteins**

Purification of human alpha<sub>1</sub>-beta glycoprotein and study of its microheterogeneity and molecular variants 405(1987)337

**Coagulation factors**

Separation of human factor X from factor Xa by reversed-phase HPLC 444(1988)107

**Coating methods**

“Superdynamic” coating of the inner surface of a capillary column with a liquid stationary phase 440(1988)323

Preparation of wall-coated open-tubular capillary columns for GC (Review) 445(1988)1

**Cobalt**

HPLC of N-terminal tyrosine-containing oligopeptides by pre-column fluorescence derivatization with hydroxylamine, cobalt(II) and borate reagents 411(1987)305

Ion chromatographic determination of low level cadmium(II), cobalt(II) and manganese(II) in water 450(1988)361

**Cobalt complexes**

Partial resolution of cobalt(III) chelate complexes by GPC 403(1987)331

Paper electrophoretic study of ion-pair formation. XVI. Resolution of optical isomers of some cobalt(III) complexes with amino acid ligands 403(1987)388

Determination of cations at the parts-per-billion level with non-suppressed ion chromatography and indirect photometric detection 404(1987)282

Effect of the geometrical configuration of facial and meridional isomers of cobalt(III) and chromium(III) complexes on their  $R_F$  values 404(1987)307

Paper electrophoretic study of ion-pair formation. XVII. Influence of the enantiomeric purity of the sodium tartrate used as electrolyte 440(1988)361

Chromatographic studies of metal complexes. I. PC of cobalt(III) complexes 442(1988)381

Rapid optical resolution of anionic metal complexes by GPC 450(1988)284

### **Cocaine**

Analysis of manufacturing by-products and impurities in illicit cocaine via HPLC and photodiode array detection 405(1987)273

Capillary GC-electron capture detection of coca-leaf-related impurities in illicit cocaine: 2,4-diphenylcyclobutane-1,3-dicarboxylic acids, 1,4-diphenylcyclobutane-2,3-dicarboxylic acids and their alkaloidal precursors, the truxillines 410(1987)297

Determination of benzoylecgonine and cocaine in biologic fluids by automated GC 417(1987)277

### **Codeine**

Identification of opiates in urine by capillary column GC of two different derivatives 422(1987)117

Separation of eight alkaloids and meconic acid and quantitation of five principal alkaloids in gum opium by gradient reversed-phase HPLC 436(1988)455

Separation of opium alkaloids by carbon dioxide sub- and supercritical fluid chromatography with packed columns. Application to the quantitative analysis of poppy straw extracts 437(1988)351

### **Co-eluting bands**

Simplified description of HPLC separation under overload conditions, based on the Craig distribution model. III. Computer simulations for two co-eluting bands assuming a Langmuir isotherm 405(1987)1

Simplified description of HPLC separation under overload conditions, based on the Craig distribution model. IV. Comparison of the model with experimental data for samples containing two or more solutes 405(1987)31

### **Coenzyme A**

HPLC fluorometry for the determination of thiols in biological samples using N-[4-(6-dimethylamino-2-benzofuranyl)phenyl]maleimide 414(1987)11

### **Coenzyme Q<sub>10</sub>**

Determination of coenzyme Q<sub>10</sub>,  $\alpha$ -tocopherol and cholesterol in biological samples by coupled-column LC with coulometric and UV detection 425(1988)87

### **Cold traps**

Trapping efficiency of capillary cold traps 448(1988)117

### **Collagen**

Detection of uridine 5'-diphosphate glucose-collagen glucosyltransferase activity by HPLC and possible participation of a novel intermediate as a glucose donor in collagen biosynthesis 417(1987)403

Column LC separation of collagen  $\alpha$ -chain polymers on sepiolite 426(1988)162

Quantitative evaluation of collagen fractions separated by acrylamide gel electrophoresis: its application for collagen typing in normal and pathological tissues 433(1988)81

Separation of type IX collagen from other cartilage collagens by hydrophobic interaction chromatography 434(1988)423

### **Collagens**

Use of the Pharmacia Phast System for sodium dodecyl sulphate polyacrylamide gel electrophoresis of non-globular proteins: application to collagens 431(1988)474

### **Colony-stimulating factor**

Purification and characterization of granulocyte-macrophage colony stimulating factor from human placenta 413(1987)242

Purification of murine macrophage colony-stimulating factor using hydroxyapatite HPLC in the presence of sodium dodecyl sulphate 421(1987)355

**Column blockage**

Simplified description of HPLC separation under overload conditions, based on the Craig distribution model. III. Computer simulations for two co-eluting bands assuming a Langmuir isotherm 405(1987)1

Simplified description of HPLC separation under overload conditions, based on the Craig distribution model. IV. Comparison of the model with experimental data for samples containing two or more solutes 405(1987)31

**Column calibration**

High-performance GPC of glycosaminoglycans. Column calibration by gel electrophoresis 402(1987)149

**Column capacity**

Influence of mobile phase pH on HPLC column loading capacity. Implications for the design of preparative protein separations 437(1988)429

**Column efficiency**

Effect of organic modifier on the separation and efficiencies of homologues on porous polymer columns 407(1987)19

Influence of the relative concentrations of two components of a binary mixture on their overloaded elution profiles 440(1988)9

Sample solvent effects in an apparent chiral HPLC separation on  $\beta$ -cyclodextrin 448(1988)31

**Column hardware**

Column hardware in preparative LC with axial flow 450(1988)47

**Column inlet concentration profile**

Effect of column inlet concentration profile on the sensitivity of analysis. A quantitative assessment 409(1987)360

**Column length**

Influence of column length and pore size on high-performance ion-exchange chromatography of estrogen and progesterin receptors 425(1988)293

**Column lifetime**

Effects of large sample loads on column lifetime in preparative-scale LC 410(1987)43

**Column loading**

Column loading and relative retention in overloaded elution chromatography 450(1988)27

**Column optimization**

Separation of proteins by reversed-phase HPLC. I. Optimizing the column 443(1988)363

**Column packing**

Contribution to the packing of LC microcolumns 411(1987)185

**Column preparation**

Evaluation of laboratory-prepared high-performance size-exclusion columns. Applications to food analysis 408(1987)115

"Superdynamic" coating of the inner surface of a capillary column with a liquid stationary phase 440(1988)323

**Column regeneration**

On-line derivatization of silica supports for regeneration and preparation of reversed-phase columns 409(1987)155

Sample preparation and column regeneration in biopolymer separations (Review) 418(1987)27

**Column residency time**

HPLC of amino acids, peptides and proteins. LXXXIV. Application of derivative spectroscopy to the study of column residency effects in the reversed-phase and size-exclusion LC separation of proteins 435(1988)271

**Columns**

see also Capillary columns and microbore columns

Detection of void zones in packed chromatographic columns 404(1987)33

Evaluation of laboratory-prepared high-performance size-exclusion columns. Applications to food analysis 408(1987)115

On-line derivatization of silica supports for regeneration and preparation of reversed-phase columns 409(1987)155

Chemometric classification of reversed-phase HPLC columns 410(1987)31

Comparison of packed column and capillary column SFC and HPLC using representative herbicides and pesticides as typical moderate polarity and molecular weight range molecules 410(1987)343

Prediction of HPLC capacity factors following column calibration. Influence of different solvents and columns on the separation of a series of aromatic amides 437(1988)13

Microcolumn chromatofocusing. III. Effect of the analysis time on the resolution and the use of a semi-disposable microcolumn 438(1988)15

Determination of the volumes of void zones in packed chromatographic columns 445(1988)37



**Column switching**

Effects of system peaks in a column-switching LC system for bioanalysis 407(1987)49

Use of micellar mobile phases and microbore column switching for the assay of drugs in physiological fluids 427(1988)277

Monitoring of veterinary drug residues by a combination of continuous flow techniques and column-switching HPLC. I. Sulphonamides in egg, meat and milk using post-column derivatization with dimethylaminobenzaldehyde 435(1988)97

**Column temperature**

Temperature effect in packed microcolumn chromatography using a carbon dioxide mobile phase 402(1987)328

Prediction of thermodynamic retention behaviour in reversed-phase LC with aqueous eluents and various concentrations of methanol by using the solubility concept model 408(1987)21

Effect of column temperature on the retention of peropyrene-type polycyclic aromatic hydrocarbons on various chemically bonded stationary phases in reversed-phase LC 436(1988)1

Effect of temperature on the GC separation of halogenated compounds on polar and non-polar stationary phases 437(1988)33

Some aspects of the selection of HPLC methods for the separation of chiral compounds in pharmaceutical analysis 450(1988)145

**Complex-forming equilibria**

Complex-forming equilibria in isotachopheresis. VII. Evaluation of the ion-pair formation constants of phosphorus oxo acids with histidine and assessment of their separability 410(1987)279

**Complexing agents**

Use of complexing agents for selective separation in high-performance capillary electrophoresis. Chiral resolution via cyclodextrins incorporated within polyacrylamide gel columns 448(1988)41

**Computer programmes**

CAPA—computer aided pesticide analysis. Computer program for the automated evaluation of chromatographic data for residue analysis of foods 441(1988)213

**Computer programs**

Computer analysis of automated Edman

degradation and amino acid analysis data 404(1987)41

**Concanavalin A**

Reversed-phase HPLC separation of the tetrameric lectin concanavalin A from *Canavalia ensiformis* 407(1987)319

Purification of intact concanavalin A in tetramers by isoelectric focusing 423(1987)319

Analytical high-performance affinity chromatography of ovalbumin-derived glycopeptides on columns of concanavalin A- and wheat germ agglutinin-immobilized gels 438(1988)73

High-performance concanavalin A affinity chromatography of liver and hepatoma membrane proteins 444(1988)29

**Concanavalin A-binding glycoproteins**

Staining of serum concanavalin A-binding proteins after micro two-dimensional electrophoresis and blotting 423(1987)115

**Concanavalin A binding proteins**

High-performance affinity chromatography of human serum concanavalin A binding proteins 431(1988)45

**Concanavalin A-Sepharose**

Human urinary  $\beta$ -mannosidase: behaviour towards two chromatographic supports 428(1988)203

**Concentration overload**

Optimization of concentration overload in preparative LC 444(1988)275

**Concentration profiles**

Isotachopheresis: analysis and computation of the structure of the ionic species interface 405(1987)77

**Concentrator columns**

Studies on sample preconcentration in ion chromatography. VI. Use of concentrator columns with "permanently" bound ion-interaction reagents 407(1987)121

**Conductivity**

Characterization of mobile phases for the investigation of electrokinetic phenomena in LC 447(1988)177

**Conductivity detection**

Ion-exclusion chromatography of non-ionic

substances with conductivity detection  
409(1987)271

#### **Conductivity detector**

Simultaneous conductivity and amperometric detector with a single cell for microbore LC  
436(1988)413

#### **Conductometric detector**

Use of first derivative conductometric detection in ion chromatography 436(1988)86

#### **Conjugates**

Application of molecular-secondary-ion MS for drug metabolism studies. I. Direct analysis of conjugates by GLC-secondary-ion mass spectrometry 414(1987)139

Unusual conjugates in biological profiles originating from consumption of onions and garlic 421(1987)1

#### **Connectivity index**

Some problems in the correlation of molecular parameters and the connectivity index  
403(1987)19

#### **Continuous flow techniques**

Monitoring of veterinary drug residues by a combination of continuous flow techniques and column-switching HPLC. I. Sulphonamides in egg, meat and milk using post-column derivatization with dimethylaminobenzaldehyde 435(1988)97

#### **Controlled-porosity glasses**

Comparison of mesoporous controlled-porosity glasses to silica gel as solid supports for C<sub>18</sub> chemically bonded phases 446(1988)347

#### **Copolymers**

Comparison between SEC and liquid adsorption chromatography in the determination of the chemical heterogeneity of copolymers  
411(1987)355

#### **Copper-bicinchoninate detection**

Optimizing copper-bicinchoninate carbohydrate detection for use with water-elution HPLC: a technique to measure the major mono- and oligosaccharides in small pieces of wheat endosperm 449(1988)207

#### **Copper**

Interaction of proteins with immobilized Cu<sup>2+</sup>. Quantitation of adsorption capacity, adsorption isotherms and equilibrium constants by frontal analysis 403(1987)197

#### **Copper diethyldithiocarbamate**

Determination of disulfiram and two of its metabolites in urine by reversed-phase LC and spectrophotometric detection after post-column complexation 424(1988)95

#### **Copper-sensitized flame photometric detector**

Spectra and response linearization of chlorine and bromine compounds in a copper-sensitized flame photometric detector 408(1987)69

#### **Coproporphyrin**

Reversed-phase HPLC of urinary porphyrin free acids 423(1987)313

#### **Coptisine**

Determination of alkaloids of *Fumaria parviflora* and *Fumaria capreolata* by HPLC and capillary isotachopheresis 445(1988)258

#### **Copy number analysis**

Isolation of high-molecular-weight nucleic acids for copy number analysis using HPLC  
402(1987)189

#### **Cordia goetzei**

Isolation of biologically active plant constituents by LC 450(1988)3

#### **Coronene**

Effect of column temperature on the retention of peropyrene-type polycyclic aromatic hydrocarbons on various chemically bonded stationary phases in reversed-phase LC  
436(1988)1

#### **Corrosion inhibitors**

Determination of corrosion inhibitor/lubricity enhancer additives in jet fuels by SEC  
437(1988)203

#### **Corrosion products**

Contamination of aqueous eluents due to corrosion of stainless-steel chromatographic components 407(1987)133

#### **Corticosteroids**

Three-hour assay for urinary free cortisol: historical comparison of three methods each using a different chromatographic procedure  
415(1987)118

Application of HPLC to the measurement of cortisol secretion rate 416(1987)25

Determination of plasma corticosterone using HPLC 419(1987)280

Determination of methylprednisolone metabolites in human urine by GC-MS 423(1987)15

Analysis of prednisolone acetate and related corticoids in swine plasma by reversed-phase HPLC 425(1988)162

Determination of cortisol in human plasma by capillary GC-MS using [ $^2\text{H}_5$ ] cortisol as an internal standard 425(1988)237

Selective determination of urinary free cortisol by LC after solid-state extraction 426(1988)25

Simplified method for the determination of plasma 18-hydroxycorticosterone 426(1988)148

GC and mass spectral study of synthetic corticosteroid metabolism: fluorometholone 426(1988)399

Clinical analysis of individual steroids by column LC (Review) 429(1988)155

Improved method for the determination of the cortisol production rate using HPLC and liquid scintillation counting 430(1988)233

Analysis of methylprednisolone, methylprednisone and corticosterone for assessment of methylprednisolone disposition in the rat 430(1988)241

GC and MS study of betamethasone synthetic corticosteroid metabolism 432(1988)283

#### **Corticosterone**

Determination of plasma corticosterone using HPLC 419(1987)280

Analysis of methylprednisolone, methylprednisone and corticosterone for assessment of methylprednisolone disposition in the rat 430(1988)241

#### **Cortisol**

Three-hour assay for urinary free cortisol: historical comparison of three methods each using a different chromatographic procedure 415(1987)118

Application of HPLC to the measurement of cortisol secretion rate 416(1987)25

Determination of cortisol in human plasma by capillary GC-MS using [ $^2\text{H}_5$ ] cortisol as an internal standard 425(1988)237

Selective determination of urinary free cortisol by LC after solid-state extraction 426(1988)25

Clinical analysis of individual steroids by column LC (Review) 429(1988)155

Improved method for the determination of the cortisol production rate using HPLC and liquid scintillation counting 430(1988)233

Simultaneous analysis of methylprednisolone hemisuccinate, cortisol and methylprednisolone by normal-phase HPLC in human plasma 432(1988)308

#### **Corynoline-type alkaloids**

Assay of five corynoline-type alkaloids in *Corydalis bungeana* by reversed-phase HPLC 408(1987)426

#### **Cosmetics**

Determination of preservatives in cosmetic products. I. TLC procedure for the identification of preservatives in cosmetic products 410(1987)395

HPLC determination of six *p*-hydroxybenzoic acid esters in cosmetics using Sep-Pak Florisil cartridges for sample pre-treatment 410(1987)413

#### **Cotinine**

Simplified method for the determination of plasma cotinine using GC-MS 414(1987)449

Simultaneous determination of nicotine, cotinine and five additional nicotine metabolites in the urine of smokers using pre-column derivatisation and HPLC 419(1987)375

Assessment of exposure to environmental tobacco smoke using a HPLC method for the simultaneous determination of nicotine and two of its metabolites in urine 431(1988)216

#### **Cotinine-N-oxide**

Identification and quantitative analysis of cotinine-N-oxide in human urine 423(1987)365

#### ***p*-Coumaric acid**

Identification of phenols, phenolic acid dimers, and monosaccharides by GLC on a capillary column 436(1988)484

#### **Coumarin**

Determination of Cinnamomi cortex by HPLC 409(1987)365

Analysis of coumarin and its urinary metabolites by HPLC 416(1987)165

Determination of coumarin as an adulterant in vanilla flavoring products by HPLC 438(1988)369

Determination of cinnamaldehyde, coumarin and cinnamyl alcohol in cinnamon and cassia by HPLC 447(1988)272

**Coumarins**

Spray reagent for the detection of coumarins and flavonoids on thin-layer plates 403(1987)285

Simultaneous isocratic HPLC determination of flavones and coumarins in *Matricaria chamomilla* extracts 404(1987)279

Quantitative structure-activity relationship of coumarin derivatives. Involvement of partition between aqueous and membrane phase 445(1988)49

Separation of natural products by centrifugal partition chromatography 450(1988)91

**Counter-current chromatography**

Foam counter-current chromatography with the cross-axis synchronous flow-through coil planet centrifuge 403(1987)77

Counter-current chromatographic purification of [D-Ala<sup>1</sup>]peptide T amide 411(1987)431

Separations of flavonoids and alkaloids in medicinal herbs by high-speed counter-current chromatography 435(1988)159

Application of droplet counter-current chromatography to the isolation of vitamin B<sub>12</sub> 435(1988)235

Multi-stage mixer-settler planet centrifuge. Preliminary studies on partition of macromolecules with organic-aqueous and aqueous-aqueous two-phase solvent systems 437(1988)121

Separation of hydroxyanthraquinone derivatives extracted from rheum with analytical high-speed counter-current chromatography 442(1988)455

Rapid separation of flavonoids by analytical high-speed counter-current chromatography 445(1988)199

Cross-axis synchronous flow-through coil planet centrifuge for large-scale preparative counter-current chromatography. I. Apparatus and studies on stationary phase retention in short coils 449(1988)135

Cross-axis synchronous flow-through coil planet centrifuge for large-scale preparative counter-current chromatography. II. Studies on partition efficiency in short coils and preparative separations with multilayer coils 449(1988)153

**Counter-current distribution**

Separation by counter-current distribution of youngchick erythrocyte populations and numerical resolution of distribution curves 411(1987)167

Partitioning behaviour in aqueous two-phase systems and fractionation by counter-current distribution of chick embryo erythrocytes with numerical resolution of distribution curves 445(1988)327

**Counter-current extraction**

Preparative chiral separation in an aqueous two-phase system by a few counter-current extractions 450(1988)277

**Coverage density**

Influence of the porous silica gel structure on the coverage density of a chemically bonded C<sub>18</sub> phase for HPLC 446(1988)191

**Craig distribution model**

Simplified description of HPLC separation under overload conditions, based on the Craig distribution model. III. Computer simulations for two co-eluting bands assuming a Langmuir isotherm 405(1987)1

Simplified description of HPLC separation under overload conditions, based on the Craig distribution model. IV. Comparison of the model with experimental data for samples containing two or more solutes 405(1987)31

Simplified description of HPLC separation under overload conditions, based on the Craig distribution model. V. Gradient elution separation 405(1987)51

**Craig model**

Preparative separation of peptide and protein samples by HPLC with gradient elution. I. The Craig model as a basis for computer simulations 444(1988)303

**C-reactive protein**

Single-step purification of rat C-reactive protein and generation of monospecific C-reactive protein antibody 437(1988)399

**Creatine**

Simultaneous determination of creatine and creatinine in serum by HPLC 431(1988)471

**Creatine kinase**

Creatine kinase isoenzymes (Review) 429(1988)399

**Creatinine**

Specific HPLC method for the determination of creatinine in rat plasma 426(1988)358

Chromatography and electrophoresis of

creatinine and other guanidino compounds  
(Review) 429(1988)235

Simultaneous determination of creatine and  
creatinine in serum by HPLC 431(1988)471

Zone-electrophoretic sample treatment for LC  
436(1988)421

#### ***p*-Cresidine**

Determination of *p*-cresidine in FD&C Red No.  
40 by the diazotization and coupling procedure  
followed by reversed-phase HPLC 405(1987)283

#### **Cresol**

Comparative study of alkyl and fluoroalkyl N-  
(2-hydroxy-ethoxyethyl)-amides as reversed-  
phase LC modifiers 448(1988)365

#### ***p*-Cresol**

Determination of simple phenols in faeces and  
urine by HPLC 431(1988)143

#### **Cresols**

Determination of trace levels of phenol and  
cresols in rain by continuous liquid-liquid  
extraction and HPLC 403(1987)233

Separation of *m*- and *p*-cresol and other  
methylated phenols in shale oil by reversed-phase  
LC with electrochemical detection 441(1988)347

#### **Crohn's disease**

Chromatography, flow injection analysis and  
electrophoresis in computer-assisted comparative  
biochemistry: its application and possibilities in  
clinical research. Preliminary studies on Crohn's  
disease 440(1988)261

#### **Cross-axis synchronous flow-through coil planet centrifuge**

Cross-axis synchronous flow-through coil planet  
centrifuge for large-scale preparative counter-  
current chromatography. I. Apparatus and  
studies on stationary phase retention in short  
coils 449(1988)135

Cross-axis synchronous flow-through coil planet  
centrifuge for large-scale preparative counter-  
current chromatography. II. Studies on partition  
efficiency in short coils and preparative  
separations with multilayer coils 449(1988)153

#### **Crotonates**

Considerations and remarks on the analysis of  
*Anthemis nobilis* L. essential oil by capillary GC  
and "hyphenated" techniques 411(1987)237

#### **Crown ethers**

Planar chromatographic separation of

enantiomers and diastereomers with cyclodextrin  
mobile phase additives 448(1988)345

#### **Cryoadsorption**

Capillary GC analysis of low-boiling organic  
sulphur compounds in anoxic lake-water by  
cryoadsorption 445(1988)97

#### **Cryogenic sampler**

Cryogenic sampling for aniline vapor with  
subsequent LC-electrochemical detection  
analysis 408(1987)81

#### **Cryogenic trapping**

Trapping efficiency of capillary cold traps  
448(1988)117

#### **Cryotrap technique**

A cryotrap technique for the quantitation of  
monoterpenes in humid and ozone-rich forest air  
442(1988)157

#### **Cryptogein**

Chromatographic purification and  
characterization of elicitors of necrosis on  
tobacco produced by incompatible *Phytophthora*  
species 440(1988)87

#### **Cryptopine**

Separation of eight alkaloids and meconic acid  
and quantitation of five principal alkaloids in  
gum opium by gradient reversed-phase HPLC  
436(1988)455

Determination of alkaloids of *Fumaria parviflora*  
and *Fumaria capreolata* by HPLC and capillary  
isotachopheresis 445(1988)258

#### **$\beta$ -Cryptoxanthin**

Separation and quantitative analysis of some  
carotenoid fatty acid esters of fruits by LC  
435(1988)113

Separation of carotenol fatty acid esters by HPLC  
449(1988)119

#### **Crystallins**

Electrophoretic titration curve in 6 *M* urea of  
the bovine eye lens protein  $\alpha$ -crystallin  
411(1987)507

Optimal resolution of eye lens  $\gamma$ -crystallins by  
cation-exchange HPLC on Synchropak CM300  
444(1988)239

#### **Cyadox**

LC determination of cyadox in medicated feeds  
and in the contents of the porcine gastrointestinal  
tract with fluorescence detection 447(1988)244

**Cyanazine**

LC method for the determination of cyanazine in the presence of some normal soil constituents 447(1988)426

**Cyanide**

Derivatization of inorganic anions with pentafluorobenzyl methanesulphonate for GC 404(1987)175

Sensitive assay for determination of pyridoxal-5-phosphate in enzymes using HPLC after derivatization with cyanide 434(1988)209

Polymeric pentafluorobenzylating reagent for GC of inorganic anions with electron-capture detection 438(1988)253

Capillary tube isotachophoretic separation of heavy metal ions using complex-forming equilibria between cyanide as terminating ion and the metal ions 447(1988)383

**Cyanoginosins**

LC determination of the cyanoginosins, toxins produced by the cyanobacterium *Microcystis aeruginosa* 408(1987)435

**Cyanohydrins**

Enantiomer separation of acyloins and cyanohydrins by enantioselective capillary GC 439(1988)301

**Cyclandelate**

HPLC method for the determination of a cyclandelate metabolite, mandelic acid, in human plasma 425(1988)203

**Cycloalkylated bases**

Separation of synthetic cycloalkylated bases, nucleosides and nucleotides by reversed-phase HPLC 430(1988)309

 **$\beta$ -Cyclodextrin**

Determination of association constants of  $\beta$ -cyclodextrin and  $\beta$ -cyclodextrin-bearing polymers with drugs by HPLC 409(1987)61

**Cyclodextrin glucosyltransferase**

Biospecific sorption of cyclodextrin glucosyltransferase on physically modified starch 427(1988)331

**Cyclodextrins**

Use of cyclodextrins in isotachopheresis. I. Effect of cyclodextrin on the isotachopheretic separation of related penicillins 405(1987)379

Use of cyclodextrins in isotachopheresis. II.  $\alpha$ -

$\beta$ - and  $\gamma$ -cyclodextrins as leading electrolyte additives for the separation of *ortho*-, *meta*- and *para*-substituted halogenobenzoic acids 411(1987)153

Enantiomeric resolution of racemic nicotine and nicotine analogues by microcolumn LC with  $\beta$ -cyclodextrin inclusion complexes 411(1987)490

Determination of cyclodextrins in biological fluids by HPLC with negative colorimetric detection using post-column complexation with phenolphthalein 415(1987)325

Computer-aided optimisation of drug enantiomer separation in chiral HPLC 434(1988)377

Use of cyclodextrins in isotachopheresis. III. Purity control of naftidrofuryl hydrogenoxalate and some of its synthesis intermediates 435(1988)496

$\alpha$ -,  $\beta$ - and  $\gamma$ -cyclodextrins as mobile phase additives in the HPLC separation of enantiomeric compounds. II. Optimization of the separation method by using  $\alpha$ -,  $\beta$ - and  $\gamma$ -cyclodextrins in mixture 436(1988)31

Resolution of chiral barbiturates into enantiomers by reversed-phase HPLC using methylated  $\beta$ -cyclodextrins 436(1988)381

Retention behaviour of cyclodextrins and branched cyclodextrins on reversed-phase columns in HPLC 437(1988)47

Use of cyclodextrins in isotachopheresis. IV. The influence of cyclodextrins on the chiral resolution of ephedrine alkaloid enantiomers 438(1988)211

Use of cyclodextrins in isotachopheresis. V. The separation of ketotifen and its polar intermediate enantiomers 439(1988)386

Reversed-phase TLC of some chlorophenols in the presence of a soluble  $\beta$ -cyclodextrin polymer 439(1988)393

Separation of enantiomers on diluted permethylated  $\beta$ -cyclodextrin by high-resolution GC 441(1988)155

Structural factors affecting chiral recognition and separation on  $\beta$ -cyclodextrin bonded phases 441(1988)376

Host-guest complexation in capillary isotachopheresis. I.  $\alpha$ -,  $\beta$ - and  $\gamma$ -cyclodextrins as complexing agents for the resolution of substituted benzoic acid isomers 442(1988)371

$\beta$ -Cyclodextrin as a selective agent for separation of selected aromatic acids by HPLC 445(1988)337

Cyclodextrins as chiral stationary phases in capillary GC. I. Pentylated  $\alpha$ -cyclodextrin 447(1988)193

Chromatographic determination of cyclodextrins on benzoylated polyacrylamide gels 447(1988)398

Sample solvent effects in an apparent chiral HPLC separation on  $\beta$ -cyclodextrin 448(1988)31

Use of complexing agents for selective separation in high-performance capillary electrophoresis.

Chiral resolution via cyclodextrins incorporated within polyacrylamide gel columns 448(1988)41

Effect of the content of divinylbenzene in ion-exchange resins on the chromatographic separation of  $\alpha$ -cyclodextrin and glucose 448(1988)241

Planar chromatographic separation of enantiomers and diastereomers with cyclodextrin mobile phase additives 448(1988)345

Cyclodextrin–ligand interaction as a simplified model of biospecific affinity chromatography 448(1988)391

Some aspects of the selection of HPLC methods for the separation of chiral compounds in pharmaceutical analysis 450(1988)145

Use of cyclodextrins in isotachopheresis. VI. Cyclodextrins as leading electrolyte additives for the separation of bile acids 450(1988)373

#### **Cycloguanil**

Determination of proguanil and its metabolites cycloguanil and 4-chlorophenylbiguanide in plasma, whole blood and urine by HPLC 416(1987)394

#### **Cyclopenthiamide**

LC screening of diuretics in urine 415(1987)347

Determination of cyclopenthiamide in plasma by HPLC 421(1987)401

#### **Cyclophosphamide**

Determination of cyclophosphamide in whole blood and plasma by reversed-phase HPLC 422(1987)125

Combined TLC–photography–densitometry for the quantitation of cyclophosphamide and its four principal urinary metabolites 427(1988)121

Rapid method for the determination of ifosfamide and cyclophosphamide in plasma by HPLC with solid-phase extraction 431(1988)450

#### **Cyclopiazone acid**

Standardized HPLC of 182 mycotoxins and other

fungal metabolites based on alkylphenone retention indices and UV–VIS spectra (diode array detection) 404(1987)195

#### **Cyclopyrrolones**

Lipophilicity measurement of benzodiazepine-receptor ligands by reversed-phase LC.

Comparison between HPLC and TLC 447(1988)404

#### **Cycloserine**

Simultaneous quantification of cycloserine and its prodrug acetylacetyl cycloserine in plasma and urine by HPLC using UV absorbance and fluorescence after post-column derivatization 414(1987)121

#### **Cyclosporin**

Monitoring of cyclosporin in whole blood by reversed-phase liquid chromatography on a butyl column 427(1988)195

#### **Cyclosporins**

Measurement of cyclosporin A and of four metabolites in whole blood by HPLC 413(1987)121

New automated HPLC analysis of cyclosporin A and G in human serum 413(1987)131

LC determination of cyclosporin A in serum with use of a solid-phase extraction. Comparison between HPLC and radioimmunoassay levels in clinical investigations 413(1987)141

Optimized HPLC method for the analysis of cyclosporine and three of its metabolites in blood and urine 425(1988)233

HPLC column-switching method for two cyclosporine metabolites in blood. 425(1988)343

#### **Cyclotrimethylenetrinitramine**

LC analysis of cyclotrimethylenetrinitramine in biological fluids using solid-phase extraction 421(1987)430

#### **Cytochrome c**

Displacement chromatography of biomolecules 439(1988)341

#### **Cyproterone acetate**

Automated HPLC assay for cyproterone acetate and 15 $\beta$ -hydroxycyproterone acetate in plasma 420(1987)432

#### **Cyromazine**

Capillary GC separation and MS detection of cyromazine and its metabolite melamine 408(1987)335

**Cysteamine**

HPLC fluorometry for the determination of thiols in biological samples using N-[4-(6-dimethylamino-2-benzofuranyl)phenyl]maleimide 414(1987)11

**Cysteine**

Increased yield of the phenylthiocarbonyl-amino acids tyrosine, cysteine and phenylalanine using diethylether-based liquid-liquid extraction 403(1987)299

HPLC fluorometry for the determination of thiols in biological samples using N-[4-(6-dimethylamino-2-benzofuranyl)phenyl]maleimide 414(1987)11

Measurement of plasma thiols after derivatization with monobromobimane 424(1988)141

Determination of cysteine, glutathione and N-acetylcysteine in plasma by ion-pair reversed-phase LC and post-column derivatization 432(1988)65

**Cysteine sulphinate decarboxylase**

GC assay for cysteine sulphinate decarboxylase activity in animal tissues 420(1987)135

**5-S-Cysteinyl-dopa**

Determination of 5-S-cysteinyl-dopa in human urine by direct injection in coupled-column HPLC 420(1987)146

**Cytidine**

Isocratic HPLC method for studying the metabolism of blood plasma pyrimidine nucleosides and bases: concentration and radioactivity measurements 434(1988)111

**Cytochrome *b*<sub>5</sub> reductase**

Improved isolation of rat microsomal cytochrome *b*<sub>5</sub> reductase 405(1987)319

**Cytochrome *c***

Multi-stage mixer-settler planet centrifuge. Preliminary studies on partition of macromolecules with organic-aqueous and aqueous-aqueous two-phase solvent systems 437(1988)121

**Cytochrome *c* oxidase**

Purification of cytochrome *c* oxidase from corn root mitochondria by hydrophobic interaction chromatography 444(1988)219

**Cytochrome P-450**

Differences in the constitutive forms of hepatic

cytochrome P-450 in male and female adult beagle dogs 419(1987)61

**Cytokinins**

*E*-Isozeatin and *cis*-norzeatin riboside as internal standards for cytokinin analyses 436(1988)93

**Cytosine**

Isocratic HPLC method for studying the metabolism of blood plasma pyrimidine nucleosides and bases: concentration and radioactivity measurements 434(1988)111

**D****Dactinomycin**

Reversed-phase HPLC of dactinomycin 427(1988)166

**Damascenone**

Composition of essential oils. Part 7. Bulgarian oil of rose (*Rosa damascena* Mill.) 406(1987)185

**Danazol**

Analysis of danazol in serum using HPLC 415(1987)438

Sensitive and selective assay of danazol in plasma by HPLC 416(1987)347

**N-Dansylaziridine**

Determination of zinc pyrithione in cosmetic products by HPLC with pre-labelling 408(1987)255

**Dansylhydrazine**

Determination of tetraprenylacetone in human plasma by HPLC with fluorescence derivatization using dansylhydrazine 424(1988)410

**Danthron**

Screening procedure for stimulant laxatives in urine using HPLC with diode array detection 433(1988)131

**Dantrolene**

Determination of dantrolene and its reduced and oxidized metabolites in plasma by HPLC 430(1988)187



**Dapsone**

Dapsone and monoacetyldapsone determined in serum and saliva by a sensitive HPLC method with a single extraction step 422(1987)322

**Dazmegrel**

HPLC with fluorescence detection of dazmegrel, a specific thromboxane A<sub>2</sub> synthetase inhibitor 432(1988)375

***o,p'*-DDD**

Rapid micromethod for the analysis of mitotane and its metabolite in plasma by GC with electron-capture detection 417(1987)287

***o,p'*-DDE**

Rapid micromethod for the analysis of mitotane and its metabolite in plasma by GC with electron-capture detection 417(1987)287

**Deacetyl-N,O-demethyldiltiazem**

HPLC determination of diltiazem and four of its metabolites in plasma: evaluation of their stability 430(1988)103

**Deacetyl-O-demethyldiltiazem**

HPLC determination of diltiazem and four of its metabolites in plasma: evaluation of their stability 430(1988)103

**Deacetyldiltiazem**

Assay of diltiazem and deacetyldiltiazem by capillary GC 417(1987)89

HPLC determination of diltiazem and four of its metabolites in plasma: evaluation of their stability 430(1988)103

**Deacetyl-N-monodesmethyl diltiazem**

HPLC determination of diltiazem and four of its metabolites in plasma: evaluation of their stability 430(1988)103

**Deacetylthymoxamine**

HPLC assay of the metabolites of thymoxamine 424(1988)205

**DEAE-Zeta-Prep**

Purification of specific heterologous F(ab)<sub>2</sub> fragments with DEAE-Zeta-Prep cartridges for ion-exchange chromatography 450(1988)133

***m*-O-Dealkylated lactam of flecainide**

HPLC procedure with fluorescence detection for the *m*-O-dealkylated lactam metabolite of flecainide acetate in human plasma 431(1988)438

**Debrisoquine**

Direct determination of debrisoquine and 4-hydroxydebrisoquine using a HPLC switching technique 419(1987)469

Rapid LC determination of debrisoquine and its hydroxy metabolite in human urine to define hydroxylation phenotypes 423(1987)340

HPLC method for the analysis of debrisoquine and its *S*-(+)- and *R*-(-)-hydroxy metabolites in urine 423(1987)344

HPLC quantitation of 4-hydroxydebrisoquine in microsomal incubates by use of silica columns, aqueous mobile phase and automated column switching 424(1988)435

Comparison of GC and HPLC assays for the determination of debrisoquine and its 4-hydroxy metabolite in human fluids 425(1988)311

Assay of debrisoquine and 4-hydroxydebrisoquine in urine by reversed-phase HPLC using on-line sample clean-up on a standard isocratic chromatograph 428(1988)178

**Decarboxylative alkylation**

Complementary use of reversed-phase, normal-phase and chiral HPLC for the study of the stereoisomers produced in decarboxylative alkylation of tartaric acid 440(1988)183

**Dehydroabiatic acid**

GC analysis of underivatized resin acids 448(1988)428

**Dehydrochloramphenicol**

HPLC determination of chloramphenicol and four analogues using reductive and oxidative electrochemical and UV detection 417(1987)111

**Dehydrodiferulic acid**

Identification of phenols, phenolic acid dimers, and monosaccharides by GLC on a capillary column 436(1988)484

**Dehydroindolidan**

LC determination of indolidan and its dehydro metabolite in plasma. Use of the Varian AASP for solid-phase extraction 431(1988)103

**Dehydrosparteines**

GC determination of sparteine and 2- and 5-dehydrosparteine in plasma and urine 431(1988)203

**Dekamycin**

Identification of dekamycin antibiotics 440(1988)71

**Delphinine**

HPLC analysis of C<sub>19</sub>-diterpenoid alkaloids 402(1987)366

**Deltamethrin**

Separation and identification of enantiomers by HPLC with a chiral column and a polarimetric detector as applied to deltamethrin 450(1988)169

**Demethylclomipramine**

Determination of clomipramine and its hydroxylated and demethylated metabolites in plasma and urine by LC with electrochemical detection 416(1987)311

**N-Demethylerythromycin A**

TLC study of the metabolites of erythromycins in the Wistar rat 434(1988)177

**Demethylfenflumizole**

HPLC assay of fenflumizole and its demethyl metabolites in biological samples 413(1987)189

**Densitometers**

Pros and cons of electronic scanning in densitometry and an experimental device for their verification 437(1988)97

**Dentifrices**

Flavor characterization of dentifrices using kinetic headspace GC 438(1988)273

**Deoxy- $\alpha$ -acids**

Avoidance of trace metal activity with micro-LC in the analysis of hop and beer bitter acids 407(1987)159

**Deoxyaconitine**

HPLC analysis of C<sub>19</sub>-diterpenoid alkaloids 402(1987)366

**Deoxyadenosine**

Rapid separation of anomeric purine nucleosides by TLC on a chiral stationary phase 410(1987)226

**Deoxycholate**

Studies on steroids. CCXXXIII. Separation and characterization of bile acid disulphates in human urine by HPLC 425(1988)59

**Deoxycholic acid**

Improved method for the determination of the major neutral steroids and unconjugated bile acids in human faeces using capillary GC 421(1987)21

Analysis of conjugated bile acids in human serum by HPLC using post-column enzyme reaction

and off-line fluorimetric determination 423(1987)75

**Deoxycytidine**

Isocratic HPLC method for studying the metabolism of blood plasma pyrimidine nucleosides and bases: concentration and radioactivity measurements 434(1988)111

**2-Deoxyglucose**

HPLC separation of some mono- and disaccharides with detection by a post-column enzyme reactor and a chemically modified electrode 408(1987)157

**2'-Deoxyguanosine**

Separation of modified nucleic acid constituents by micellar electrokinetic capillary chromatography 409(1987)193

**2'-Deoxyguanosine-3'-monophosphate**

Detection of oxidatively modified 2'-deoxyguanosine-3'-monophosphate, using <sup>32</sup>P-postlabelling and anion-exchange TLC 434(1988)222

**5'-Deoxy-5'-methylthioadenosine**

Chromatographic and radioimmunological methods for the determination of 5'-deoxy-5'-methylthioadenosine in biological fluids 440(1988)141

**9-[5'-Deoxy-5'-(methylthio)- $\beta$ -D-xylofuranosyl]adenine**

HPLC determination of a naturally occurring xylosyl thioether, 9-[5'-deoxy-5'-(methylthio)- $\beta$ -D-xylofuranosyl]adenine 440(1988)151

**Deoxynivalenol**

HPLC method for determining tricothecene mycotoxins by post-column fluorescence derivatization 410(1987)427

Metabolism of three tricothecene mycotoxins, T-2 toxin, diacetoxyscirpenol and deoxynivalenol, by bovine rumen microorganisms 414(1987)335

**Deoxynucleosides**

DNA-nitrosourea interactions. HPLC of crosslinked dinucleosides and substituted deoxynucleosides 436(1988)23

**Deoxyribomononucleotides**

Separation of modified nucleic acid constituents by micellar electrokinetic capillary chromatography 409(1987)193

**Deoxyribonucleosides**

Separation of modified nucleic acid constituents by micellar electrokinetic capillary chromatography 409(1987)193

**Deoxyspergualin**

Determination of deoxyspergualin by GC-selected-ion monitoring 424(1988)39

**Dephosphate-bromofenofos**

HPLC determination of bromofenofos and its metabolite in rat plasma 426(1988)202

**N-Depropylpropafenone**

HPLC separation and MS identification of propafenone, 5-hydroxypropafenone and N-depropylpropafenone 424(1988)211

**Derivative spectroscopy**

HPLC of amino acids, peptides and proteins. LXXXIV. Application of derivative spectroscopy to the study of column residency effects in the reversed-phase and size-exclusion LC separation of proteins 435(1988)271

**Derivatization**

Preparative resolution of the enantiomers of the  $\beta$ -blocking drug atenolol by chiral derivatization and HPLC 431(1988)222

**Derivatization, electrophoresis**

Post-capillary fluorescence detection in capillary zone electrophoresis using *o*-phthalaldehyde 447(1988)117

**Derivatization, GC**

Derivatization of inorganic anions with pentafluorobenzyl methanesulphonate for GC 404(1987)175

Resolution of monoterpene enantiomers by GC 407(1987)243

Decomposition of trifluoroacetyl derivatives of some trichothecenes on fused-silica capillary columns during GC analysis 407(1987)340

Derivatization of diaminocyclohexanes for determination of *cis* and *trans* isomers by GC 408(1987)211

Benzyl esters of  $C_2$ - $C_{20}$  fatty acids and metabolically relevant carboxylic acids. Preparation and GC-MS 409(1987)43

GC of methyl derivatives of naturally occurring capsaicinoids 409(1987)223

Formation of N-carboxy compounds during

trimethylsilylation of amino acids in the presence of carbon dioxide 409(1987)383

Trimethylsilylation of the hydrolysed and polycondensed products of methyltriethoxysilane 410(1987)137

Capillary column GC identification of sugars in honey as trimethylsilyl derivatives 410(1987)319

GC and MS behaviour of O-alkylserines as their N-trifluoroacetyl butyl esters 411(1987)259

GC analysis of neutral monosaccharides as their O-pentafluorobenzoyloxime acetates 411(1987)275

Enantiomer separation of tocainide and some of its analogues and derivatives on a Chirasil-Val capillary column 411(1987)285

Derivatization of 2-amino alcohols with phosgene in aqueous media: limitations of the reaction selectivity as found in the presence of O-glucuronides of alprenolol in urine 413(1987)270

Separation and quantitation of fatty acids, sterols and bile acids in feces by GC as the butyl ester-acetate derivatives 415(1987)13

Use of pentafluorobenzyl and pentafluoropropionyl-pentafluorobenzyl esters of bacterial fatty acids for GC analysis with electron-capture detection 417(1987)366

Use of *tert.*-butyldimethylsilyl derivatives for GC-MS analysis of dipeptides 419(1987)263

Silylated N,O-ketals from the reaction of ketones with N-trimethylsilylimidazole 419(1987)271

Single-step method for derivatization of dopamine and some related compounds in aqueous media for GC 421(1987)227

O-(2,3,4,5,6-Pentafluorobenzyl)oxime-trimethylsilyl ester derivatives for quantitative GC and GC-MS studies of aldehydes, ketones and oxoacids 421(1987)336

Identification of opiates in urine by capillary column GC of two different derivatives 422(1987)117

Determination of oxpentifylline and three metabolites in plasma by automated capillary GC using nitrogen-selective detection 423(1987)139

Dimethylisopropylsilyl ether derivatives in GC-MS of some prostanoids. Application to the microdetermination of 15-ketoprostaglandin  $F_{2\alpha}$  formed by enzymatic conversion of  $PGF_{2\alpha}$  in vitro 424(1988)1

- Identification and quantitation of trenbolone in bovine tissue by GC-MS 424(1988)219
- GC analysis of urinary dimercaptosuccinic acid 424(1988)231
- Simultaneous determination of C<sub>2</sub>-C<sub>22</sub> non-esterified fatty acids and other metabolically relevant carboxylic acids in biological material by GC of their benzyl esters 425(1988)257
- Determination of busulfan in human plasma by GC with electron-capture detection 425(1988)303
- Rapid and simple procedure for the determination of urinary phenylacetic acid using derivatization in aqueous medium followed by electron-capture GC 428(1988)140
- Determination of mefloquine by electron-capture GC after phosgene derivatization in biological samples and in capillary blood collected on filter paper 428(1988)281
- Separation of optical isomers of methoxyphenamine and its metabolites as N-heptafluorobutyl-L-prolyl derivatives by fused-silica capillary GC 431(1988)210
- Identification of polychlorinated phenols in urine by GC and TLC 431(1988)431
- Improved procedure for preparation of pentafluorobenzyl derivatives of carboxylic acids for analysis by GC with electron-capture detection 433(1988)248
- Phosgene as a derivatizing reagent prior to GC and LC (Review) 435(1988)259
- GC analysis of aminoalkylphosphonic acids and aminoalkyl phosphates 436(1988)67
- Use of chloroform as a solvent for the formation of N-trifluoroacetyl-*n*-butylamino acid derivatives for GC analysis 436(1988)100
- Derivatization of prostaglandins and related compounds to (methoxime) alkyl ester alkyl ether derivatives for GC analysis 436(1988)219
- Analytical separations of enantiomeric primary amines as pyrrole derivatives formed by reaction with polygodial 438(1988)108
- Polymeric pentafluorobenzylating reagent for GC of inorganic anions with electron-capture detection 438(1988)253
- Assignment of the configurations of the amino acids in peptidic siderophores 438(1988)291
- Enantiomer separation of acyloins and cyanohydrins by enantioselective capillary GC 439(1988)301
- GC determination of chlorinated phenols in the form of various derivatives 439(1988)307
- Analysis of tryptamine at the femtomole level in tissue using negative ion chemical ionization GC-MS 440(1988)253
- 1,2-Diaminobenzene derivatives for the determination of selenium by GC 440(1988)333
- GC analysis of different homologous series of acids esterified in aqueous solutions with butyl and propyl alcohols 446(1988)237
- N-Methoxy-N,O-bis(trimethylsilyl) carbamate as a derivatizing reagent for the GC of sitosterol degradation products 446(1988)253
- Extractive pentafluorobenzylation of formic, acetic, levulinic, benzoic and phthalic acids, studied by LC and dual-oven capillary GC 447(1988)329
- Regio-selective derivatization of 1-[(3,5-difluorophenyl)methyl]-1,3-dihydro-2H-imidazole for GC analysis 450(1988)403
- Derivatization, LC**
- Determination of reducing and non-reducing carbohydrates in food products by LC with post-column catalytic hydrolysis and derivatization. Comparison with refractive index detection 402(1987)127
- New derivatization reagents for the resolution of carboxylic acid enantiomers by HPLC 402(1987)344
- Separation of norepinephrine enantiomers, DOPA enantiomers and dopamine derivatized with *o*-phthalaldehyde-N-acetyl-L-cysteine by HPLC 402(1987)387
- Analysis of neutral sugars as dinitrophenyl-hydrazones by HPLC 405(1987)221
- Determination of *p*-cresidine in FD&C Red No. 40 by the diazotization and coupling procedure followed by reversed-phase HPLC 405(1987)283
- Ligand exchange chromatography of amino alcohol enantiomers as Schiff bases 406(1987)343
- Amino acid analysis by reversed-phase HPLC. Automatic pre-column derivatization with activated carbamate reagent 407(1987)273
- Assay of phenylmercury salts in pharmaceutical products by HPLC of the morpholinedithiocarbamate derivative 407(1987)389
- HPLC separation of some mono- and disaccharides with detection by a post-column

- enzyme reactor and a chemically modified electrode 408(1987)157
- Automated pre-column derivatization of amino acids with *o*-phthalaldehyde by a reagent sandwiching technique 408(1987)217
- Improved method for benzoyl chloride derivatization of polyamines for HPLC 408(1987)227
- Determination of zinc pyrithione in cosmetic products by HPLC with pre-labelling 408(1987)255
- On-line derivatization of silica supports for regeneration and preparation of reversed-phase columns 409(1987)155
- Determination of nitrosamines by LC with post-column photolysis and electrochemical detection 410(1987)145
- Determination of artesunic acid after chemical derivatization with *O*-*p*-nitrobenzyl-*N,N'*-diisopropylisourea by HPLC and UV absorption 410(1987)217
- HPLC method for determining trichothecene mycotoxins by post-column fluorescence derivatization 410(1987)427
- Quantitative determination of octanoic acid by HPLC following derivatization with 4-bromomethyl-7-methoxycoumarin 411(1987)297
- HPLC of *N*-terminal tyrosine-containing oligopeptides by pre-column fluorescence derivatization with hydroxylamine, cobalt(II) and borate reagents 411(1987)305
- High-performance liquid chromatography of 1,2-diacyl-*rac*-glycerols and 1,3-diacylglycerols through chromogenic derivatization with 3,5-dinitrobenzoyl 411(1987)472
- LC determination of amoxicillin and its metabolites in human urine by postcolumn degradation with sodium hypochlorite 413(1987)219
- HPLC fluorometry for the determination of thiols in biological samples using *N*-[4-(6-dimethylamino-2-benzofuranyl)phenyl]maleimide 414(1987)11
- Determination of methylglyoxal as 2-methylquinoxaline by HPLC and its application to biological samples 414(1987)149
- Determination of the enantiomers of ketoprofen in blood plasma by ion-pair extraction and HPLC of leucinamide derivatives 414(1987)465
- Determination of 5-fluoro-2'-deoxyuridine in human plasma by HPLC with pre-column fluorimetric derivatization 416(1987)176
- HPLC analysis of serum long-chain fatty acids by direct derivatization method 416(1987)237
- Sensitive and selective determination of picogram amounts of ciprofloxacin and its metabolites in biological samples using HPLC and photothermal post-column derivatization 416(1987)321
- Automated pre-column derivatization with *o*-phthalaldehyde for the determination of neurotransmitter amino acids using reversed-phase LC 417(1987)385
- Sensitive ferrocene reagents for derivatization of thiol compounds in HPLC with dual-electrode coulometric detection 419(1987)17
- Column LC of methotrexate and its metabolites using a post-column photochemical reactor and fluorescence detection 419(1987)213
- Highly sensitive determination of chlorpheniramine as fluorescence derivative by HPLC 420(1987)63
- Automatic amino acid analysis utilizing 4-fluoro-7-nitrobenzo-2-oxa-1,3-diazole 420(1987)141
- Determination of natural thiols by LC after derivatization with 3,5-di-*tert*.-butyl-1,2-benzoquinone 420(1987)404
- HPLC analysis of serum short-chain fatty acids by direct derivatization 421(1987)33
- Determination of metoprolol enantiomers in plasma and urine using (S)-(-)-phenylethyl isocyanate as a chiral reagent 421(1987)161
- Analysis of conjugated bile acids in human serum by HPLC using post-column enzyme reaction and off-line fluorimetric determination 423(1987)75
- Determination of acetylcholine and choline in microdialysates from spinal cord of rat using LC with electrochemical detection 423(1987)85
- Assessment of taurine in plasma and urine by anion-exchange HPLC with pre-column derivatization 423(1987)292
- Formation of diastereomeric derivatives of 2-arylpropionic acids using *L*-leucinamide 423(1987)334
- Determination of disulfiram and two of its metabolites in urine by reversed-phase LC and

- spectrophotometric detection after post-column complexation 424(1988)95
- Measurement of plasma thiols after derivatization with monobromobimane 424(1988)141
- Separation of enantiomers of derivatised or underivatised propranolol by means of HPLC 424(1988)189
- Determination of  $\beta$ -lactams and their biosynthetic intermediates in fermentation media by pre-column derivatisation followed by fluorescence detection 424(1988)325
- Determination of tetraprenylacetone in human plasma by HPLC with fluorescence derivatization using dansylhydrazine 424(1988)410
- Sensitive method of detection, quantitation and purification of peptides using pre-column derivatization with phenyl isothiocyanate 425(1988)77
- Rapid and sensitive HPLC assay of tiaprofenic acid enantiomers in human plasma and urine 425(1988)135
- Pre-column derivatization of sisomicin with *o*-phthalaldehyde- $\beta$ -mercaptopropionic acid and its application to sensitive HPLC determination with fluorimetric detection 425(1988)143
- Quantification of the enantiomers of ofloxacin in biological fluids by HPLC 425(1988)153
- Determination of agmatine, arginine, citrulline and ornithine by reversed-phase LC using automated pre-column derivatization with *o*-phthalaldehyde 425(1988)269
- Quantitation of polyamines in cultured cells and tissue homogenates by reversed-phase HPLC of their benzoyl derivatives 426(1988)41
- Fluorimetric determination of urinary  $\delta$ -aminolevulinic acid by HPLC and post-column derivatization 426(1988)365
- Novel method for HPLC of azo derivatives of conjugated and unconjugated bilirubin 427(1988)19
- Chromatographic analysis of the trinitrophenyl derivatives of insulin 427(1988)239
- Method to determine the enantiomers of ibuprofen from human urine by HPLC 428(1988)103
- Determination of S-carboxymethylcysteine in serum by reversed-phase ion-pair LC with column switching following pre-column derivatization with *o*-phthalaldehyde 428(1988)301
- Analysis of benzodiazepines. II. HPLC fluorescence detection after molecular rearrangement to acridanones 428(1988)321
- 4-Bromomethyl-6,7-dimethoxycoumarin as a fluorescence reagent for precolumn derivatization of 5-fluorouracil compounds in HPLC 430(1988)156
- HPLC determination of  $\alpha$ -keto acids in human serum and urine using 1,2-diamino-4,5-methylenedioxybenzene as a precolumn fluorescence derivatization reagent 430(1988)223
- HPLC determination of leucine-enkephalin-like peptide in rat brain by pre-column fluorescence derivatization involving formylation followed by reaction with 1,2-diamino-4,5-dimethoxybenzene 430(1988)271
- Assay for enkephalin-degrading peptidases in rat brain tissues by HPLC with on-line post-column fluorescence detection 430(1988)291
- HPLC determination of forphenicine in mouse serum and muscle by pre-column fluorescence derivatization using 1,2-diamino-4,5-ethylenedioxybenzene as fluorogenic reagent 430(1988)361
- Formation of diastereomeric derivatives of 2-arylpropionic acids using L-leucinamide: lack of generality 431(1988)228
- Formation of diastereomeric derivatives of 2-arylpropionic acids using *x*-leucinamide: the derivatization principle and its applicability. A reply 431(1988)229
- Determination of amino acids in biological, pharmaceutical, plant and food samples by automated precolumn derivatization and HPLC 431(1988)271
- Construction of an ion-exchange amino acid analyser kit for use with HPLC apparatus 431(1988)285
- Determination of cysteine, glutathione and N-acetylcysteine in plasma by ion-pair reversed-phase LC and post-column derivatization 432(1988)65
- Pre-column derivatization of amino acids with N,N-diethyl-2,4-dinitro-5-fluoroaniline and reversed-phase LC separation 433(1988)53
- Determination of gossypol enantiomers in plasma after administration of racemate using HPLC with precolumn chemical derivatisation 433(1988)141
- Determination of tranexamic acid in human

serum by HPLC using selective pre-column derivatization with phenyl isothiocyanate 433(1988)225

Measurement of carprofen enantiomer concentrations in plasma and urine using L-leucinamide as the chiral coupling component 433(1988)331

HPLC determination of netilmicin in guinea-pig and human serum by fluorodinitrobenzene derivatization with spectrophotometric detection 434(1988)169

Sensitive assay for determination of pyridoxal-5-phosphate in enzymes using HPLC after derivatization with cyanide 434(1988)209

Separation of  $\alpha$ -amino acid enantiomers by reversed-phase HPLC after derivatization with *o*-phthalaldehyde and a sodium salt of 1-thio- $\beta$ -D-glucose 434(1988)417

Monitoring of veterinary drug residues by a combination of continuous flow techniques and column-switching HPLC. I. Sulphonamides in egg, meat and milk using post-column derivatization with dimethylaminobenzaldehyde 435(1988)97

Phosgene as a derivatizing reagent prior to GC and LC (Review) 435(1988)259

Derivatization of some polymers with *o*-phthalaldehyde prior to high-performance SEC 436(1988)39

Indirect HPLC resolution of racemic tertiary amines as their diastereomeric urea derivatives after N-dealkylation 436(1988)269

Rapid and simple dansylation of phenolic steroids using a two-phase system and phase transfer catalysis 436(1988)429

Automated solid-phase catalyzed pre-column derivatization of fatty acids for reversed-phase HPLC analysis with fluorescence detection 436(1988)437

Quantitative determination of free primary amines in polymeric amine hardeners. HPLC separation of ethylenediamine and triethylenetetramine after derivatization with salicylaldehyde-diphenylboron chelate and separation of 4,4'-diaminodiphenylmethane by HPLC 436(1988)447

HPLC of biogenic amines, derivatized with 9-fluorenylmethyl chloroformate 437(1988)379

Determination of iodide and bromide by ion

chromatography with post-column reaction detection 439(1988)129

Resolution of  $\alpha$ -substituted amino acid enantiomers by HPLC after derivatization with a chiral adduct of *o*-phthalaldehyde. Application to glutamic acid analogues 440(1988)209

Stereoselective determination of L-amino acids using column LC with an enzymatic solid-phase reactor and chemiluminescence detection 440(1988)217

1-(4-Hydroxyphenyl)-, 1-(2,4-dihydroxyphenyl)- and 1-(2,5-dihydroxyphenyl)-2-bromoethanones: new labels for determination of carboxylic acids by HPLC with electrochemical and UV detection 442(1988)209

Picomole determination of 2,4-dimethoxyanilides of prostaglandins by HPLC with electrochemical detection 442(1988)444

HPLC method for the determination of  $\epsilon$ -( $\gamma$ -glutamyl)lysine and mono- and bis- $\gamma$ -glutamyl derivatives of putrescine and spermidine 443(1988)329

Application of Marfey's reagent in racemization studies of amino acids and peptides 444(1988)115

HPLC analysis of arginine-containing peptides in biological fluids by means of a selective post-column reaction with fluorescence detection 444(1988)123

Determination of carnitine by HPLC using 9-anthryldiazomethane 445(1988)175

Anthracene-9-carbonyl chloride as a fluorescence and UV derivatising reagent for the HPLC analysis of hydroxy compounds 445(1988)393

Determination of diethylene glycol in wine by HPLC using anthracene-9-carbonyl chloride as a derivatising reagent 445(1988)403

HPLC determination of lombricine and N-phosphoryl lombricine in the earthworm by pre-column fluorescence derivatisation with *o*-phthalaldehyde-ethanethiol 445(1988)433

Resolution of enantiomeric drugs of some  $\beta$ -amino alcohols as their urea derivatives by HPLC on a chiral stationary phase 447(1988)208

Extractive pentafluorobenzoylation of formic, acetic, levulinic, benzoic and phthalic acids, studied by LC and dual-oven capillary GC 447(1988)329

Tyrosylation and purification of peptides for radioiodination 447(1988)341

Resolution of lombricine enantiomers by HPLC utilising pre-column derivatisation with *o*-phthaldialdehyde-chiral thiols 447(1988)392

Trace determination of sulfide by reversed-phase ion-interaction chromatography using pre-column derivatization 447(1988)415

Analysis of catechin and epicatechin by HPLC after benzylation 448(1988)291

Optimizing copper-bicinchoninate carbohydrate detection for use with water-elution HPLC: a technique to measure the major mono- and oligosaccharides in small pieces of wheat endosperm 449(1988)207

HPLC determination of glucosides (glucose conjugates) with post-column reaction detection combining immobilized enzyme reactors and luminol chemiluminescence 449(1988)217

Separation of the stereoisomers of an allenic E-type prostaglandin 450(1988)233

#### Derivatization, MS

MS evaluation of the *tert*.-butyldimethylsilyl derivatives of monohydroxyeicosatetraenoic acids and leukotrienes 423(1987)1

#### Derivatization, TLC

Separation of  $\beta$ -amino alcohols as their dansyl derivatives by TLC on a polyamide layer sheet 408(1987)449

Separation and determination of some amino acid ester enantiomers by TLC after derivatization with (*S*)-(+)-naproxen 450(1988)281

#### Dermatan sulphate

High-performance GPC of glycosaminoglycans. Column calibration by gel electrophoresis 402(1987)149

Rapid determination of disaccharides from chondroitin and dermatan sulphates by HPLC 414(1987)156

LC assay for constituent disaccharides of hyaluronic acid and chondroitin sulphate isomers 415(1987)231

Extracellular mammalian polysaccharides: glycosaminoglycans and proteoglycans (Review) 418(1987)187

Fractionation of aorta glycosaminoglycans by HPLC 419(1987)275

Characterization of the products generated from

oversulphated dermatan sulphate isomers with chondroitinase-B by HPLC 423(1987)51

Methods for analysis of urinary glycosaminoglycans (Review) 429(1988)293

#### Desacetylcefotaxime

Determination of serum cefotaxime and desacetylcefotaxime by ion-pair HPLC (in French) 419(1987)396

HPLC assay of cefotaxime, desacetylcefotaxime and ceftriaxone in rat plasma 424(1988)111

#### Desacetyldiltiazem

High-performance liquid chromatographic determination of diltiazem and its major metabolites, N-monodemethyldiltiazem and desacetyldiltiazem, in plasma 415(1987)203

#### 3-Desacetylpancuronium

Quantitation of pancuronium, 3-desacetylpancuronium, vecuronium, 3-desacetylvecuronium, pipecuronium and 3-desacetylpipecuronium in biological fluids by capillary GC using nitrogen-sensitive detection 427(1988)41

#### 25-Desacetyl rifampicin

Determination of rifampicin and its main metabolites in human plasma by HPLC 426(1988)412

#### Desacetyl TZU-0460

Highly sensitive determination of 2-hydroxy-N-{3-[*m*-(1-piperidinylmethyl)phenoxy]propyl}acetar: (desacetyl TZU-0460) in rat plasma by HPLC 413(1987)370

#### 3-Desacetylvecuronium

Quantitation of pancuronium, 3-desacetylpancuronium, vecuronium, 3-desacetylvecuronium, pipecuronium and 3-desacetylpipecuronium in biological fluids by capillary GC using nitrogen-sensitive detection 427(1988)41

#### Desacetylvinblastine

Analysis of vinca alkaloids in plasma and urine using HPLC with electrochemical detection 424(1988)83

#### N-1-Desalkylflurazepam

Simultaneous determination of flurazepam and five metabolites in serum by HPLC and its application to pharmacokinetic studies in rats 423(1987)251



**N-1-Desalkyl-3-hydroxyflurazepam**

Simultaneous determination of flurazepam and five metabolites in serum by HPLC and its application to pharmacokinetic studies in rats 423(1987)251

**Desbutylhalofantrine**

Determination of halofantrine and its principal metabolite desbutylhalofantrine in biological fluids by reversed-phase HPLC 433(1988)339

**Desethylamiodarone**

Rapid HPLC method for the determination of amiodarone and desethylamiodarone in human plasma and serum 415(1987)197

New procedure for the HPLC determination of amiodarone and desethylamiodarone with solid-phase extraction of rat plasma and tissue samples 431(1988)455

**N-Desethylamiodarone**

Determination of amiodarone and its metabolite N-desethylamiodarone in serum by HPLC-comparison of different extraction procedures 417(1987)465

**Desethylchloroquine**

Field-adapted method for HPTLC detection and estimation of chloroquine and desethylchloroquine in urine 423(1987)261

Field-adapted method for HPTLC detection and estimation of chloroquine in finger-stick blood 428(1988)196

Analysis of blood and urine samples for hydroxychloroquine and three major metabolites by HPLC with fluorescence detection 433(1988)197

**Desethyleneciprofloxacin**

HPLC of ciprofloxacin and its metabolites in serum, urine and sputum 422(1987)153

**Desethylhydroxychloroquine**

Analysis of blood and urine samples for hydroxychloroquine and three major metabolites by HPLC with fluorescence detection 433(1988)197

**Desferrioxamine**

HPLC analysis of desferrioxamine using a metal-free system 411(1987)494

HPLC analysis of desferrioxamine. Pharmacokinetic and metabolic studies 433(1988)207

**Desferrioxamine B**

Determination of desferrioxamine B, its complexes with iron and aluminium and selected derivatives by HPLC using a polymer column 438(1988)433

**Desipramine**

HPLC determination of imipramine and its metabolites in rat brain 421(1987)412

Effect of blood collection tubes on antidepressant concentrations 423(1987)373

Simultaneous determination of imipramine, desipramine and their deuterium-labelled analogues in biological fluids by capillary GC-MS 426(1988)93

Isolation and identification of the glucuronide conjugate of 2-hydroxydesipramine by preparative LC 427(1988)287

**Desisopropylpropranolol**

HPLC determination of propranolol and its metabolites in rat serum 431(1988)465

**N-Desisopropylpropranolol**

Separation of enantiomers of derivatised or underderivatised propranolol by means of HPLC 424(1988)189

**Desmethyladinazolam**

HPLC assay of adinazolam and its desmethyl metabolite 424(1988)215

**Desmethylazelastine**

Determination of azelastine and desmethylazelastine in human plasma by HPLC 420(1987)89

**Desmethylcarvedilol**

HPLC method for the determination of carvedilol and its desmethyl metabolite in body fluids 413(1987)355

**N-Desmethylclobazam**

Micro-determination of clobazam and N-desmethylclobazam in plasma or serum by electron-capture GC 421(1987)147

**N-Desmethylclomipramine**

Simultaneous determination of clomipramine and its N-desmethyl metabolite in human whole blood by capillary GC with mass-selective detection 428(1988)71

**N-Desmethyldiazepam**

Microsample determination of diazepam and its

three metabolites in serum by reversed-phase HPLC 416(1987)212

Simultaneous determination of diazepam and its active metabolites in rat serum, brain and cerebrospinal fluid by HPLC 422(1987)334

Improved method for the determination of diazepam and N-desmethyldiazepam in plasma using capillary GC and nitrogen-phosphorus detection 424(1988)398

Sensitive determination of diazepam and N-desmethyldiazepam in human material using capillary GC-MS 431(1988)353

#### **N-Desmethyldifloxacin**

HPLC procedures for the determination of difloxacin and its metabolites in biological matrices 413(1987)199

#### **N-Desmethyleclanamine**

HPLC method for the determination of eclanamine and its N-desmethyl and N,N-didesmethyl metabolites in urine 428(1988)131

#### **Desmethylmaprotiline**

Determination of maprotiline and desmethylmaprotiline in plasma and urine by GC with nitrogen-phosphorus detection 432(1988)334

#### **N-Desmethylstobadine**

GC determination of stobadine, N-desmethylstobadine and stobadine N-oxide in rat liver microsomal incubations 415(1987)163

#### **N-Desmethyltamoxifen**

Quantification of tamoxifen and N-desmethyltamoxifen in human plasma by HPLC, photochemical reaction and fluorescence detection, and its application to biopharmaceutic investigations 413(1987)207

#### **Desmethylthioridazine**

HPLC resolution of the enantiomers of thioridazine, its metabolites and related compounds 426(1988)223

#### **N-Desmethyltimelotem**

Determination of timelotem dihydrogenmaleate and its main metabolite N-desmethyltimelotem in plasma by capillary GC 419(1987)357

#### **Desmosine**

HPLC determination of desmosine and isodesmosine in tissues and its application to studies of alteration of elastin induced by atherosclerosis 422(1987)53

#### **Desogestrel**

Optimization of the reversed-phase HPLC separation of synthetic estrogenic and progestogenic steroids using the multi-criteria decision making method 410(1987)1

#### **Desorption gas flow-rate**

Dynamic headspace GC: concentration of volatile components after thermal desorption by intermediate cryofocusing in a cold trap. II. Effect of sampling and desorption parameters on recovery 405(1987)99

#### **Desorption temperature**

Dynamic headspace GC: concentration of volatile components after thermal desorption by intermediate cryofocusing in a cold trap. II. Effect of sampling and desorption parameters on recovery 405(1987)99

#### **Detection, LC**

Lanthanide ions as luminescent chromophores for the LC detection of polynucleotides and nucleic acids 436(1988)299

Microcolumn HPLC-TLC-Fourier transform IR spectrometry 438(1988)329

#### **Detection limits**

Chromatographic detection limits in pharmaceutical method development 447(1988)1

#### **Detection, TLC**

*In situ* visualization of glycosphingolipids on thin-layer chromatograms using lectin-peroxidase conjugates 411(1987)393

#### **Detectors, GC**

UV spectrometric detection for GC of polynuclear aromatic compounds with repetitive spectral scans using absorbance and concurrent fluorescence measurements 405(1987)191

An experiment to distinguish between homogeneous and heterogeneous response mechanisms in a parallel-plate electron-capture detector 406(1987)81

Correlations and constants in unipolar and bipolar electron-capture detection 408(1987)53

Spectra and response linearization of chlorine and bromine compounds in a copper-sensitized flame photometric detector 408(1987)69

Determination of silicon number in silylated organic compounds using the hydrogen atmosphere flame ionization detector 408(1987)329

General photometric detector for GC  
409(1987)125

MS measurements of the negative ions formed in the thermionic ionization detector with nitrogen carrier gas 409(1987)139

GC determination of benzaldehyde by gas-phase luminescence 409(1987)243

Improved method for the trace analysis of explosives by silica capillary column GC with thermal energy analysis detection 410(1987)181

Multiple element emission spectral detection GC profiles of halogenated products from chlorination of humic acid and drinking water 438(1988)35

The mass-selective detector as a chlorine-selective detector 438(1988)233

Non-radioactive electron-capture detector operating in the pulsed mode 441(1988)45

Capillary GC analysis with the far-UV absorbance detector 441(1988)63

Analysis of ppb levels of organics in water by means of purge-and-trap, capillary GC and selective detectors 441(1988)73

Influence of the carrier gas flow-rate on the pulsed electron-capture detector 446(1988)329

### Detectors, LC

Evaluation of absorbance ratioing for solute identification in HPLC using a diode array detector 402(1987)135

Application fields of a new evaporative light scattering detector for HPLC and supercritical fluid chromatography (in French) 404(1987)95

Analysis of platinum species originating from *cis*-diamminedichloroplatinum(II) (cisplatin) in human and rat plasma by HPLC with on-line inductively coupled plasma atomic emission spectrometric detection 407(1987)253

Direct synchronous luminescence detection of co-eluting solutes in pseudophase LC 409(1987)147

Ion-exclusion chromatography of non-ionic substances with conductivity detection 409(1987)271

Thin-layer electrochemical detector coated with Nafion film for LC 410(1987)111

Determination of nitrosamines by LC with post-column photolysis and electrochemical detection 410(1987)145

Characterization of organotin species using microbore and capillary LC techniques with an epifluorescence microscope as a novel imaging detector 410(1987)393

Determination of nitrodiphenylamines by LC and dual-electrode amperometric detection 410(1987)437

Photoionization detector for open-tubular LC 411(1987)201

Multiple detection system for the high-resolution microcolumn SEC separation of proteins with a slurry-packed capillary column 411(1987)221

Evaluation of a chemical reaction detector with electrochemical production of bromine for the LC determination of physostigmine 411(1987)229

Identification of drugs in human serum by HPLC with photodiode array detection and a search algorithm for UV spectra 419(1987)135

LC and flow injection analysis of tetracycline using sensitized europium(III) luminescence detection 433(1988)149

Use of laser detectors in capillary LC (Review) 434(1988)337

Quantitative analysis of triglycerides by HPLC using non-linear gradient elution and flame ionization detection 435(1988)139

HPLC of amino acids, peptides and proteins. LXXXIV. Application of derivative spectroscopy to the study of column residency effects in the reversed-phase and size-exclusion LC separation of proteins 435(1988)271

Signal enhancement by on-column fluorometric detection in HPLC using micropacked fused-silica columns 435(1988)319

Use of first derivative conductometric detection in ion chromatography 436(1988)86

Photodiode array detection of peptide-nucleoside antibiotics 436(1988)259

Simultaneous conductivity and amperometric detector with a single cell for microbore LC 436(1988)413

Paraffins, olefins, naphthenes and aromatics analysis of selected hydrocarbon distillates using on-line column switching HPLC with dielectric constant detection 437(1988)365

UV fibre optic photometric cell with an effective volume of 30 nl for micro-scale HPLC 438(1988)409

Single-column anion chromatography with indirect UV detection using pyromellitate buffers as eluents 439(1988)137

Dynamic quenching of europium(III) and terbium(III) luminescence: a potential detection method in ion chromatography 439(1988)165

Ion-pair HPLC of inorganic anions with photometric, conductometric and amperometric detection 439(1988)363

Stereoselective determination of L-amino acids using column LC with an enzymatic solid-phase reactor and chemiluminescence detection 440(1988)217

Coupling LC with redox chemiluminescence detection 441(1988)125

High-performance UV absorption detector for LC. II. A prototype detector 441(1988)387

Conformational studies of bovine alkaline phosphatase in hydrophobic interaction and size-exclusion chromatography with linear diode array and low-angle laser light scattering detection 442(1988)29

Amperometric detector with a manganin electrode for microcolumn LC 442(1988)111

Indirect photometric detection of inorganic anions in micro HPLC with permanently coated columns 447(1988)221

LC determination of carbohydrates with pulsed amperometric detection and a membrane reactor 447(1988)268

### **Detectors, SFC**

Application fields of a new evaporative light scattering detector for HPLC and supercritical fluid chromatography (in French) 404(1987)95

Super- and subcritical fluid chromatography-Fourier transform IR spectrometry of medium polar compounds on polymeric styrene-divinylbenzene packed columns 407(1987)87

Interfacing of inductively coupled plasma-atomic emission spectrometry to SFC for elemental detection 411(1987)213

Trace analysis of explosives by capillary supercritical fluid chromatography with thermal energy analysis detection 445(1988)244

### **Detectors, TLC**

New method of visualizing thin-layer chromatograms by using liquid crystals 405(1987)203

Calibration of TLC with flame ionization detection for the analysis of natural lipid samples 435(1988)343

Parameters of the liquid crystal method of visualizing thin-layer chromatograms 437(1988)131

Quantitative and automatic digital evaluation of one- and two-dimensional thin-layer chromatograms using vertical surface absorption at an optimum wavelength 437(1988)411

TLC-flame ionization detection calibration with natural and synthetic standards 439(1988)404

### **Detergents**

Misuse of the term "detergent" 426(1988)452

Chromatographic detergent exchange in the preparation of membrane receptor protein 427(1988)341

Effect of detergents on the structure of integral membrane proteins of Sendai virus studied with size-exclusion HPLC and monoclonal antibodies 443(1988)255

Use of term "detergent" in biochemistry 449(1988)345

### **Detomidine**

Analysis of detomidine in horse blood, plasma and urine samples utilizing a sensitive GC-MS method 404(1987)223

### **Deuterium exchange**

GC-MS determination of tryptophan transaminase-catalyzed deuterium exchange 415(1987)104

### **Dexamethasone**

GC and MS study of betamethasone synthetic corticosteroid metabolism 432(1988)283

HPLC determination of dexamethasone in cerebrospinal fluid and plasma in the rabbit 434(1988)315

### **Dexamethasone 17 $\beta$ -carboxamides**

HPLC in the evaluation of the lipophilicity of 17 $\beta$ -carboxamide steroid derivatives 445(1988)409

### **Dextromethorphan**

Direct determination of dextromethorphan and its three metabolites in urine by HPLC using a precolumn switching system for sample clean-up 422(1987)340

**Dextromoramide**

GC assay for dextromoramide in human plasma 432(1988)329

**Dextropropoxyphene**

LC determination of paracetamol and dextropropoxyphene in plasma 419(1987)433

**Dextrorphan**

HPLC determination of dextrorphan and 3-hydroxymorphinan in human plasma based on a selective pre-column sample clean-up 420(1987)217

Direct determination of dextromethorphan and its three metabolites in urine by HPLC using a precolumn switching system for sample clean-up 422(1987)340

**Diacerhein**

Simple method for the determination of rhein in biological fluids by HPLC 428(1988)173

**Diacetoxyscirpenol**

Metabolism of three trichothecene mycotoxins, T-2 toxin, diacetoxyscirpenol and deoxynivalenol, by bovine rumen microorganisms 414(1987)335

**Diacylglycerol**

HPLC separation of diacylglycerol enantiomers on a chiral stationary phase 402(1987)257

**Diacylglycerols**

High-performance liquid chromatography of 1,2-diacyl-*rac*-glycerols and 1,3-diacylglycerols through chromogenic derivatization with 3,5-dinitrobenzoyl 411(1987)472

MS analysis of endogenous diacylglycerols in normal and psoriatic skin 432(1988)278

**Dialkylglycerol ethers**

Chromatographic methods for determining the configuration of 1,2- and 1,3- (unsymmetrically substituted) dialkylglycerol ethers 436(1988)205

**Diamidines**

Determination of fluorescent diamidines in plasma of experimental animals by HPLC 417(1987)331

**Diamines**

Quantitation of polyamines in cultured cells and tissue homogenates by reversed-phase HPLC of their benzoyl derivatives 426(1988)41

Trace analysis of airborne aromatic isocyanates and related aminoisocyanates and diamines using

HPLC with UV and electrochemical detection 435(1988)469

**1,2-Diaminobenzenes**

1,2-Diaminobenzene derivatives for the determination of selenium by GC 440(1988)333

**Diaminocyclohexanes**

Derivatization of diaminocyclohexanes for determination of *cis* and *trans* isomers by GC 408(1987)211

**1,2-Diamino-4,5-dimethoxybenzene**

HPLC determination of leucine-enkephalin-like peptide in rat brain by pre-column fluorescence derivatization involving formylation followed by reaction with 1,2-diamino-4,5-dimethoxybenzene 430(1988)271

**4,4'-Diaminodiphenylmethane**

Quantitative determination of free primary amines in polymeric amine hardeners. HPLC separation of ethylenediamine and triethylenetetramine after derivatization with salicylaldehyde-diphenylboron chelate and separation of 4,4'-diaminodiphenylmethane by HPLC 436(1988)447

**1,2-Diamino-4,5-ethylenedioxybenzene**

HPLC determination of forphenicine in mouse serum and muscle by pre-column fluorescence derivatization using 1,2-diamino-4,5-ethylenedioxybenzene as fluorogenic reagent 430(1988)361

**1,2-Diamino-4,5-methylenedioxybenzene**

HPLC determination of  $\alpha$ -keto acids in human serum and urine using 1,2-diamino-4,5-methylenedioxybenzene as a precolumn fluorescence derivatization reagent 430(1988)223

**2,4-Diamino-N<sup>10</sup>-methylpteroic acid**

Column LC of methotrexate and its metabolites using a post-column photochemical reactor and fluorescence detection 419(1987)213

**1,8-Diaminooctane**

Staphylococcal nuclease HPLC characterization of diaminooctane-modified DNA and its biotin and fluorescein derivatives 444(1988)97

 **$\alpha$ , $\omega$ -Diaminoalcoethers**

Polarity of some individual derivatives of  $\alpha$ , $\omega$ -diaminoalcoethers as measured by GC 448(1988)219

**Diaminopimelic acid**

Determination of environmental levels of peptidoglycan and lipopolysaccharide using GC with negative-ion chemical-ionization MS utilizing bacterial amino acids and hydroxy fatty acids as biomarkers 431(1988)1

**1,3-Diaminopropane**

Determination of polyamines in human erythrocytes by capillary GC with nitrogen-phosphorus detection 415(1987)27

Quantitation of polyamines using TLC and image analysis 441(1988)361

**2,4-Diaminopyridine**

HPLC determination of 2,4-diaminopyridine in plasma and urine 421(1987)392

**cis-Diamminedichloroplatinum(II)**

Analysis of platinum species originating from *cis*-diamminedichloroplatinum(II) (cisplatin) in human and rat plasma by HPLC with on-line inductively coupled plasma atomic emission spectrometric detection 407(1987)253

**Diastereoisomers**

Chromatographic behaviour of diastereoisomers. IX. Application of a microcomputer program to the TLC separation of some diastereoisomers on silica 438(1988)219

**Diastereomeric compounds**

High-resolution GC with liquid crystal glass capillaries. X. Separation of diastereomeric alkanes 436(1988)47

**Diatoxanthin**

HPLC of carotenoids from some marine shellfish 440(1988)37

**Diazepam**

Determination of association constants of  $\beta$ -cyclodextrin and  $\beta$ -cyclodextrin-bearing polymers with drugs by HPLC 409(1987)61

Trace analysis of diazepam in serum using microbore HPLC and on-line preconcentration 413(1987)161

Microsample determination of diazepam and its three metabolites in serum by reversed-phase HPLC 416(1987)212

Determination of diazepam and its metabolites by HPLC and TLC 421(1987)291

Simultaneous determination of diazepam and its active metabolites in rat serum, brain and cerebrospinal fluid by HPLC 422(1987)334

Improved method for the determination of diazepam and N-desmethyldiazepam in plasma using capillary GC and nitrogen-phosphorus detection 424(1988)398

Use of micellar mobile phases and microbore column switching for the assay of drugs in physiological fluids 427(1988)277

Sensitive determination of diazepam and N-desmethyldiazepam in human material using capillary GC-MS 431(1988)353

**2,5-Diaziridinyl-3,6-bis(2-hydroxyethylamino)-1,4-benzoquinone**

Determination of diaziridinyl-3,6-bis(2-hydroxyethylamino)-1,4-benzoquinone in plasma by solid-phase extraction and HPLC 432(1988)352

**Dibenz[*a,j*]acridine**

*endo*-1,4,5,6,7,7-Hexachlorobicyclo[2.2.1]hept-5-ene-2-carboxylic acid, a superior resolving agent for the HPLC separation of enantiomers of hydroxylated derivatives of two azaaromatic hydrocarbons 430(1988)53

**Dibucaine**

Determination of dibucaine in biological samples by GC with a nitrogen-phosphorus detector 415(1987)407

**Dibutylamine**

Investigation of an alkylamine modified and pH-controlled mobile phase for separation of pyrazolone derivatives by HPLC 409(1987)81

**3,5-Di-*tert*.-butyl-1,2-benzoquinone**

Determination of natural thiols by LC after derivatization with 3,5-di-*tert*.-butyl-1,2-benzoquinone 420(1987)404

**Dibutyldithiocarbamate**

LC of dibutyldithiocarbamate degradation products 404(1987)359

**3,5-Di-*tert*.-butyl-4-hydroxytoluene**

Determination of butylated hydroxytoluene in plasma by HPLC 422(1987)288

**Dibutylphenols**

Separation of isomeric mono-, di- and tributylphenols by HPLC on octadecylsilica with methanol-water using alkyl nitrile and alkylamine as surface modifiers 442(1988)53

**Di-*n*-butyl sebacate**

Determination of plasticizers in fat by GC-MS 447(1988)230

**Dicamba**

Analysis of the herbicides dicamba, clopyralid and bromacil in asparagus by HPLC 439(1988)470

**Dicarboxylic acids**

Capillary GC-MS of aliphatic saturated  $\alpha,\omega$ -dicarboxylic acid dimethyl esters and direct inlet MS of the corresponding free acids 408(1987)145

Simultaneous determination of  $C_2$ - $C_{22}$  non-esterified fatty acids and other metabolically relevant carboxylic acids in biological material by GC of their benzyl esters 425(1988)257

Identification of 3-oxodicarboxylic acids in human urine 432(1988)13

**Dicarboxylic porphyrins**

Normal-phase HPLC of free acid dicarboxylic porphyrins and hematoporphyrin derivative on silica 422(1987)73

**3,4-Dichloroaniline**

Analysis of bromacil, diuron and 3,4-dichloroaniline in contaminated well water, using a HPLC column-switching procedure 410(1987)211

**Dichlorobenzenes**

High-volume elution chromatography of dichlorobenzenes on a polyurethane foam-Tenax sandwich cartridge 409(1987)235

**Di[(2-chlorobutyroanilido)methyl] sulphide**

HPLC methods for the determination of contaminants in technical phosmethylan 435(1988)380

**Dichloromethane**

Retention in capillary supercritical fluid chromatography with carbon dioxide mobile phases containing polar modifiers 406(1987)223

**6,7-Dichloro-2-methylquinoxaline**

Simple and sensitive determination of methylglyoxal in biological samples by GC with electron-capture detection 415(1987)221

**N,N-Di(2-chlorophenyl)methylenediamine**

HPLC methods for the determination of contaminants in technical phosmethylan 435(1988)380

**Dichromenes**

Application of semi-preparative HPLC to difficult natural product separations 406(1987)367

**Diclofenac**

Rapid HPLC assay for the simultaneous analysis of non-steroidal anti-inflammatory drugs in plasma 416(1987)293

**Dicloxacillin**

Optimization of HPLC analysis for isoxazolyl penicillins using factorial design 425(1988)331

**Dicyclomine**

Quantitative analysis of dicyclomine in human plasma by capillary GC and nitrogen-selective detection 416(1987)150

**Didemethylclomipramine**

Determination of clomipramine and its hydroxylated and demethylated metabolites in plasma and urine by LC with electrochemical detection 416(1987)311

**2',3'-Dideoxyadenosine**

Determination of 2',3'-dideoxyadenosine, 2',3'-dideoxyinosine and 2',3'-dideoxycytidine in biological samples 431(1988)184

**2',3'-Dideoxyadenosine**

Application of HPLC and thermospray HPLC-MS to the analysis and identification of 2',3'-dideoxyadenosine and its metabolite in biological media 420(1987)1

**2',3'-Dideoxycytidine**

Determination of 2',3'-dideoxyadenosine, 2',3'-dideoxyinosine and 2',3'-dideoxycytidine in biological samples 431(1988)184

HPLC determination of 2',3'-dideoxycytidine and 3'-azido-3'-deoxythymidine in plasma using a column-switching technique 432(1988)346

**2',3'-Dideoxyinoside**

Determination of 2',3'-dideoxyadenosine, 2',3'-dideoxyinosine and 2',3'-dideoxycytidine in biological samples 431(1988)184

**2',3'-Dideoxyinosine**

Application of HPLC and thermospray HPLC-MS to the analysis and identification of 2',3'-dideoxyadenosine and its metabolite in biological media 420(1987)1

**2',3'-Dideoxynucleosides**

Determination of 2',3'-dideoxyadenosine, 2',3'-dideoxyinosine and 2',3'-dideoxycytidine in biological samples 431(1988)184

**Didesethylflurazepam**

Simultaneous determination of flurazepam and

five metabolites in serum by HPLC and its application to pharmacokinetic studies in rats 423(1987)251

#### **N,N-Didesmethyl eclanamine**

HPLC method for the determination of eclanamine and its N-desmethyl and N,N-didesmethyl metabolites in urine 428(1988)131

#### **Didronel**

GC analysis of Didronel tablets 404(1987)372

#### **Dielectric constant detection**

Paraffins, olefins, naphthenes and aromatics analysis of selected hydrocarbon distillates using on-line column switching HPLC with dielectric constant detection 437(1988)365

#### **Diethyldithiocarbamate**

Determination of disulfiram and two of its metabolites in urine by reversed-phase LC and spectrophotometric detection after post-column complexation 424(1988)95

#### **Diethyldithiophosphates**

Trace enrichment and separation of As<sup>III</sup>, Sb<sup>III</sup> and Bi<sup>III</sup> as diethyldithiophosphate complexes by reversed-phase HPLC 439(1988)63

#### **Diethylene glycol**

Determination of diethylene glycol in wine by HPLC using anthracene-9-carbonyl chloride as a derivatising reagent 445(1988)403

#### **Diethyl ether**

HPLC determination of nicotinamide in rat tissue samples and blood after extraction with diethyl ether 422(1987)257

#### **Di(2-ethylhexyl)orthophosphoric acid**

Modification of retention of some alkaloids in the system silanized silica/methanol + water + di(2-ethylhexyl)orthophosphoric acid 439(1988)257

#### **Diethylstilboestrol**

Chromatographic separation and isolation of the enantiomers of diethylstilbestrol metabolites 439(1988)484

#### **Diffusion coefficients**

Non-linear elution effects in split-peak chromatography. I. Computer simulations for the cases of irreversible diffusion- and adsorption-limited kinetics 436(1988)111

Effect of steric exclusion on the separation of proteins by hydrophilic SEC 445(1988)29

#### **Difloxacin**

HPLC procedures for the determination of difloxacin and its metabolites in biological matrices 413(1987)199

#### **Difloxacin N-oxide**

HPLC procedures for the determination of difloxacin and its metabolites in biological matrices 413(1987)199

#### **Diffunisal**

Reversed-phase HPLC assay for the simultaneous determination of diflunisal and its glucuronides in serum and urin. Rearrangement of the 1-O-acylglucuronide 420(1987)99

#### **$\beta,\beta$ -Difluorochlorambucil**

HPLC of chlorambucil analogues 422(1987)315

#### **1-[(3,5-Difluorophenyl)methyl]-1,3-dihydro-2H-imidazole**

Regio-selective derivatization of 1-[(3,5-difluorophenyl)methyl]-1,3-dihydro-2H-imidazole for GC analysis 450(1988)403

#### **Digitoxigenin**

Simultaneous analysis of digitoxin and its clinically relevant metabolites using HPLC and radioimmunoassay 419(1987)155

#### **Digitoxigenin bisdigitoxoside**

Simultaneous analysis of digitoxin and its clinically relevant metabolites using HPLC and radioimmunoassay 419(1987)155

#### **Digitoxigenin monodigitoxoside**

Simultaneous analysis of digitoxin and its clinically relevant metabolites using HPLC and radioimmunoassay 419(1987)155

#### **Digitoxin**

Simultaneous analysis of digitoxin and its clinically relevant metabolites using HPLC and radioimmunoassay 419(1987)155

#### **Diglycerides**

Characterization of a lipid-rich fraction synthesized by *Streptomyces avermitilis* 441(1988)31

#### **Digoxigenin**

Separation of digoxin from dihydrodigoxin and the other metabolites by HPLC with post-column derivatization 421(1987)381

#### **Digoxigenin bisdigitoxoside**

Separation of digoxin from dihydrodigoxin and



the other metabolites by HPLC with post-column derivatization 421(1987)381

#### **Digoxigenin monodigitoxoside**

Separation of digoxin from dihydrodigoxin and the other metabolites by HPLC with post-column derivatization 421(1987)381

#### **Digoxin**

Separation of digoxin from dihydrodigoxin and the other metabolites by HPLC with post-column derivatization 421(1987)381

Determination of digoxin in serum by on-line immunoadsorbent clean-up HPLC separation and fluorescence-reaction detection 433(1988)119

Micro HPLC determination of cardiac glycosides in  $\beta$ -methylidigoxin and digoxin tablets 448(1988)157

#### **22,23-Dihydroavermectin B<sub>1a</sub>**

Determination of 22,23-dihydroavermectin B<sub>1a</sub> in dog plasma using solid-phase extraction and HPLC 413(1987)326

#### **Dihydrobiopterin**

Estimation of tetrahydrobiopterin and other pterins in plasma by isocratic LC with electrochemical and fluorimetric detection 432(1988)321

#### **Dihydrocapsaicin**

GC of methyl derivatives of naturally occurring capsaicinoids 409(1987)223

#### **Dihydrodigoxin**

Separation of digoxin from dihydrodigoxin and the other metabolites by HPLC with post-column derivatization 421(1987)381

#### **10,11-Dihydro-10,11-trans-dihydroxycarbamazepine**

Simultaneous determination of carbamazepine, oxcarbazepine and their main metabolites in plasma by LC 414(1987)477

#### **2,3-Dihydrodiol**

Chromatographic and cytogenetic analysis of *in vivo* metabolites of fluoranthene 448(1988)127

#### **17-Dihydroequilenin**

HPLC determination of conjugated estrogens in tablets 450(1988)414

#### **17-Dihydroequilin**

HPLC determination of conjugated estrogens in tablets 450(1988)414

#### **Dihydroergotamine**

Determination of sub-nanogram amounts of dihydroergotamine in plasma and urine using LC and fluorimetric detection with off-line and on-line solid-phase drug enrichment 417(1987)319

#### **10,11-Dihydro-10-hydroxycarbamazepine**

Simultaneous determination of carbamazepine, oxcarbazepine and their main metabolites in plasma by LC 414(1987)477

#### **2,3-Dihydro-6-[3-(2-hydroxymethyl)phenyl-2-propenyl]-5-benzofuranol**

Determination of 2,3-dihydro-6-[3-(2-hydroxymethyl)-phenyl-2-propenyl]-5-benzofuranol in plasma using LC with electrochemical detection 426(1988)295

#### **2-(10,11-Dihydro-10-oxodibenzo[*b, f*]thiepin-2-yl)propionic acid**

HPLC assay of a new non-steroidal anti-inflammatory agent, 2-(10,11-dihydro-10-oxodibenzo[*b, f*]thiepin-2-yl)propionic acid, in human plasma and urine 414(1987)381

#### **Dihydropyridine derivative**

GC-negative-ion chemical ionization MS determination of a new dihydropyridine calcium antagonist (MPC-1304) and its metabolites in human plasma and urine 434(1988)123

#### **Dihydropyridines**

Direct HPLC resolution of dihydropyridine enantiomers 450(1988)191

#### **Dihydroqinghaosu**

Analysis of artesunic acid and dihydroqinghaosu in blood by HPLC with reductive electrochemical detection 414(1987)77

#### **3,4-Dihydroxybenzylamine**

Species-dependent differences in recovery of 3,4-dihydroxybenzylamine in assays of plasma catecholamines 430(1988)123

Sensitive method for determination of picogram amounts of epinephrine and other catecholamines in microdissected samples of rat brain using LC with electrochemical detection 433(1988)41

#### **Dihydroxyborylsilica**

Determination of catecholamines in urine (and plasma) by LC after on-line sample pretreatment on small alumina or dihydroxyborylsilica columns 414(1987)285

**7 $\beta$ ,12 $\beta$ -Dihydroxy-5 $\beta$ -cholan-24-oic acid**

Internal standards for quantitative GC of individual bile acids after group separation of bile acids in urine 422(1987)25

**3 $\beta$ ,12 $\alpha$ -Dihydroxy-5-cholen-24-oic acid**

Determination of 3 $\beta$ ,12 $\alpha$ -dihydroxy-5-cholen-24-oic acid and related bile acids in human serum by GC-MS 421(1987)9

***trans*-10,11-Dihydroxy-10,11-dihydrocarbamazepine**

Quantitation of *trans*-10,11-dihydroxy-10,11-dihydrocarbamazepine in human urine by HPLC 415(1987)208

**1,3-Dimethyl-7-isobutylxanthine**

Determination of 1,3-dimethyl-7-isobutylxanthine in guinea pig plasma by HPLC 433(1988)288

**3,4-Dihydroxymandelic acid**

Simultaneous LC determination of norepinephrine, 3,4-dihydroxyphenylethyleneglycol, 3,4-dihydroxyphenylalanine and 3,4-dihydroxyphenylacetic acid in human plasma 434(1988)187

**3,4-Dihydroxyphenylacetic acid**

Simultaneous LC determination of seventeen of the major monoamine neurotransmitters, precursors and metabolites. I. Optimization of the mobile phase using factorial designs and a computer program to predict chromatograms 415(1987)261

Simultaneous LC determination of seventeen of the major monoamine neurotransmitters, precursors and metabolites. II. Assessment of human brain and cerebrospinal fluid concentrations 415(1987)275

Combined ion-pair extraction and HPLC for the determination of the biogenic amines and their major metabolites in single brain tissue samples 421(1987)51

Single-step method for derivatization of dopamine and some related compounds in aqueous media for GC 421(1987)227

Simultaneous determination of the major acidic metabolites of catecholamines and serotonin in urine by LC with electrochemical detection after a one-step sample clean-up on Sephadex G-10; influence of vanilla and banana ingestion 424(1988)273

Tritium isotope effect in reversed-phase HPLC analysis of dopamine and dihydroxyphenylacetic acid 427(1988)9

Effects of long-term storage on the concentrations of the unconjugated acidic metabolites of the trace amines, indoleamines and catecholamines 433(1988)23

Simultaneous LC determination of norepinephrine, 3,4-dihydroxyphenylethyleneglycol, 3,4-dihydroxyphenylalanine and 3,4-dihydroxyphenylacetic acid in human plasma 434(1988)187

**3,4-Dihydroxyphenylalanine**

Simultaneous LC determination of norepinephrine, 3,4-dihydroxyphenylethyleneglycol, 3,4-dihydroxyphenylalanine and 3,4-dihydroxyphenylacetic acid in human plasma 434(1988)187

**3,4-Dihydroxyphenylethylene glycol**

Simultaneous LC determination of seventeen of the major monoamine neurotransmitters, precursors and metabolites. I. Optimization of the mobile phase using factorial designs and a computer program to predict chromatograms 415(1987)261

Simultaneous LC determination of seventeen of the major monoamine neurotransmitters, precursors and metabolites. II. Assessment of human brain and cerebrospinal fluid concentrations 415(1987)275

Simultaneous LC determination of norepinephrine, 3,4-dihydroxyphenylethyleneglycol, 3,4-dihydroxyphenylalanine and 3,4-dihydroxyphenylacetic acid in human plasma 434(1988)187

**DL-*threo*-3,4-Dihydroxyphenylserine**

Determination of D,L-*threo*-3,4-dihydroxyphenylserine and of the D- and L-enantiomers in human plasma and urine 427(1988)219

**11 $\beta$ ,20 $\alpha$ -Dihydroxy-4-pregnen-3-one**

Specific reversed-phase LC method for analysis of steroids in Y-1 adrenal cell cultures 432(1988)302

**9-(1,3-Dihydroxy-2-propoxymethyl)guanine**

HPLC method for the determination of 9-(1,3-

dihydroxy-2-propoxymethyl)guanine in human plasma 414(1987)429

#### **1,2-Dihydroxypropyl-3-(2-aminoethyl)phosphonate**

Silicic acid CC of phosphonolipids. IX. Silicic acid CC study of some phosphono analogues of bis(diacylglycero)phosphate and glycerophosphoric acid 435(1988)508

#### **1,2-Dihydroxypropyl-3-phosphonic acid**

Silicic acid CC of phosphonolipids. IX. Silicic acid CC study of some phosphono analogues of bis(diacylglycero)phosphate and glycerophosphoric acid 435(1988)508

#### **1,2-Dihydroxypropyl-3-(2-trimethylammoniumethyl)phosphonate**

Silicic acid CC of phosphonolipids. IX. Silicic acid CC study of some phosphono analogues of bis(diacylglycero)phosphate and glycerophosphoric acid 435(1988)508

#### **Diiodothyronine**

Resolution of the enantiomers of thyroid hormones by high-performance ligand-exchange chromatography using a chemically bonded chiral stationary phase 404(1987)391

#### **2,6-Diisopropylphenol**

HPLC determination and preliminary pharmacokinetics of propofol and its metabolites in human plasma and urine 417(1987)458

#### **$\beta$ -Diketoamides**

Kinetic studies of fast equilibrium by means of HPLC. XVI. Separation of tautomers of  $\beta$ -diketones and  $\beta$ -diketoamides 411(1987)466

#### **$\beta$ -Diketonates**

LC retention of  $\beta$ -diketonato complexes of chromium(III) and palladium(II) in octadecyl-bonded silica/water-methanol systems. Comparison with liquid-liquid partition using dodecane as a model alkyl-bonded phase 439(1988)245

#### **$\beta$ -Diketones**

GC of metal  $\beta$ -diketonates and their analogues (Review) 411(1987)1

Kinetic studies of fast equilibrium by means of HPLC. XVI. Separation of tautomers of  $\beta$ -diketones and  $\beta$ -diketoamides 411(1987)466

#### **Dilevalol**

HPLC assay for dilevalol in human plasma and

urine using a PRP-1 column and fluorimetric detection 425(1988)363

#### **Diltiazem**

Liquid chromatographic determination of diltiazem and its metabolites using *trans* isomers as internal standards, with dynamic modification of the solid phase by addition of an amine to the mobile phase 414(1987)109

High-performance liquid chromatographic determination of diltiazem and its major metabolites, N-monodemethyldiltiazem and desacetyldiltiazem, in plasma 415(1987)203

Assay of diltiazem and deacetyldiltiazem by capillary GC 417(1987)89

Automated HPLC and column-switching technique for on-line clean-up and analysis of diltiazem in human plasma 423(1987)239

HPLC determination of diltiazem and four of its metabolites in plasma: evaluation of their stability 430(1988)103

#### **Dimercaptosuccinic acid**

GC analysis of urinary dimercaptosuccinic acid 424(1988)231

#### **1,4-Di(methacryloyloxymethyl)naphthalene**

GC on porous polymers. IV. Influence of the geometric structure of porous copolymers of 1,4-di(methacryloyloxymethyl)naphthalene with 1,4-divinylbenzene on their chromatographic behaviour 448(1988)233

#### **N<sup>4</sup>-[2,6-Dimethoxy-4-methyl-5-[(3-trifluoromethyl)phenoxy]-8-quinolinyl]-1,4-pentanediamine**

Rapid and sensitive quantitative analysis of the new antimalarial N<sup>4</sup>-[2,6-dimethoxy-4-methyl-5-[(3-trifluoromethyl)phenoxy]-8-quinolinyl]-1,4-pentanediamine in plasma by LC and electrochemical detection 424(1988)347

#### **[3,4-Di-(4-methoxyphenyl)-5-isoxazolyl]acetic acid**

HPLC determination of [3,4-di-(4-methoxyphenyl)-5-isoxazolyl]acetic acid and its metabolites in human plasma and urine 422(1987)227

#### **Dimethylacetamide**

GC-MS method for the determination of dimethylacetamide and metabolites in whole blood 428(1988)156

**Dimethylaminoantipyrine**

Investigation of an alkylamine modified and pH-controlled mobile phase for separation of pyrazolone derivatives by HPLC 409(1987)81

**N-[4-(6-Dimethylamino-2-benzofuranyl)phenyl]maleimide**

HPLC fluorometry for the determination of thiols in biological samples using N-[4-(6-dimethylamino-2-benzofuranyl)phenyl]maleimide 414(1987)11

***p*-Dimethylaminocinnamaldehyde**

Reaction of Fast Blue BB salt and *p*-dimethylaminocinnamaldehyde with indole-3-acetic acid degradation products. Evaluation of the coupling properties of the reagents used in the zymographic screening of indole-3-acetic acid oxidases 411(1987)476

**Dimethylarginine**

Analysis of methylated amino acids by HPLC: methylation of myelin basic protein 421(1987)130

**N<sup>G</sup>-Dimethylarginines**

Determination of methylated amino acids in human serum by HPLC 440(1988)225

 **$\beta,\beta$ -Dimethylchlorambucil**

HPLC of chlorambucil analogues 422(1987)315

**Dimethyl dioxane**

Specific identification of synthetic organic chemicals in river water using liquid-liquid extraction and resin adsorption coupled with electron impact, chemical ionization and accurate mass measurement GC-MS analyses 436(1988)229

**N,N-Dimethylformamide**

Identification of N-acetyl-S-(N-methylcarbamoyl)cysteine, a human metabolite of N,N-dimethylformamide and N-methylformamide 414(1987)399

GC method for the determination of N-acetyl-S-(N-methylcarbamoyl)cysteine, a metabolite of N,N-dimethylformamide and N-methylformamide, in human urine 431(1988)361

**Dimethylisopropylidene tartrate**

Complementary use of reversed-phase, normal-phase and chiral HPLC for the study of the stereoisomers produced in decarboxylative alkylation of tartaric acid 440(1988)183

**N-(2,6-Dimethylphenyl)-2-(2-oxo-1-pyrrolidinyl)acetamide**

High-performance liquid chromatographic determination of a new nootropic, N-(2,6-dimethyl-phenyl)-2-(2-oxo-1-pyrrolidinyl)-acetamide, in human serum and urine 433(1988)235

**N,N-Dimethyltryptamine**

GC-MS characterisation of the psychotomimetic indolealkylamines and their *in vivo* metabolites 422(1987)13

**1,3-Dimethyluric acid**

HPLC assay for theophylline and its major metabolites in human urine 423(1987)179

**1,7-Dimethylxanthine**

Determination of theophylline and its metabolites in plasma and urine by reversed-phase LC using an amine modifier 419(1987)177

**Dinactin**

Preparative HPLC of macrotetrolides 442(1988)431

**1,3-Dinitrobenzene**

Determination of 1,3-dinitrobenzene and its metabolites in rat blood by capillary GC with electron-capture detection 425(1988)187

**N-(3,5-Dinitrobenzoyl)amino acid methyl esters**

Chiral stationary phase derived from L-lactic acid for the optical resolution of N-(3,5-dinitrobenzoyl)amino acid methyl esters 450(1988)163

**N-(3,5-Dinitrobenzoyl)phenylglycine silica**

Optical resolution of samples with weak interactions on chiral "brush-type" LC stationary phases 440(1988)197

**2,4-Dinitrophenylhydrazones**

Retention behaviour of aromatic aldehyde derivatives in reversed-phase LC 404(1987)242

**Dinucleosides**

DNA-nitrosourea interactions. HPLC of crosslinked dinucleosides and substituted deoxynucleosides 436(1988)23

**Dinucleotides**

Demonstration of the adenosine reservoirs with nitrobenzylthioinosine in liver and diaphragm by HPLC 426(1988)249

**4,4'-Dioctyldiphenylamine**

Chromatographic analysis of phenyl-1-naphthylamine and 4,4'-dioctyldiphenylamine and their intermediate oxidation products in oxidized lubricants 409(1987)325

**Diode-array detectors**

Evaluation of absorbance ratioing for solute identification in HPLC using a diode array detector 402(1987)135

Identification of drugs in human serum by HPLC with photodiode array detection and a search algorithm for UV spectra 419(1987)135

HPLC of amino acids, peptides and proteins. LXXXIV. Application of derivative spectroscopy to the study of column residency effects in the reversed-phase and size-exclusion LC separation of proteins 435(1988)271

Photodiode array detection of peptide-nucleoside antibiotics 436(1988)259

Conformational studies of bovine alkaline phosphatase in hydrophobic interaction and size-exclusion chromatography with linear diode array and low-angle laser light scattering detection 442(1988)29

HPLC and photodiode-array detection of the human protein HC (human complex-forming glycoprotein heterogeneous in charge), a chromophore-associated protein 444(1988)165

**2,2-Di-(*p*-oligooxyethylenephenyl)propane**

Analysis of 2,2-di-(*p*-oligooxyethylenephenyl)propane by GC and GC-MS 446(1988)109

**Diols**

Anthracene-9-carbonyl chloride as a fluorescence and UV derivatising reagent for the HPLC analysis of hydroxy compounds 445(1988)393

**Dipeptide isosteres**

Determination of the chiral purity of dipeptide isosteres containing a reduced peptide bond by GC analysis 442(1988)165

**Dipeptides**

HPLC separation of peptide and amino acid stereoisomers 414(1987)313

Use of *tert*-butyldimethylsilyl derivatives for GC-MS analysis of dipeptides 419(1987)263

Analysis of dipeptides by reversed-phase HPLC without derivatization using amperometric detection on a copper electrode 436(1988)334

Resolution of stereoisomers of dipeptides by GC on Chirasil-Val 441(1988)89

Determination of the chiral purity of dipeptide isosteres containing a reduced peptide bond by GC analysis 442(1988)165

TLC enantiomeric resolution via ligand exchange 448(1988)11

Chromatographic resolution of dipeptide enantiomers and diastereomers on chiral stationary phases from poly(L-leucine) or poly(L-phenylalanine) 450(1988)271

**Dipeptidyl-aminopeptidase II**

Rapid chromatographic purification of dipeptidyl-aminopeptidase II from human kidney 416(1987)131

**Dipeptidyl peptidase IV**

HPLC separation of peptides possessing a proline residue in the amino-terminal penultimate position, and their products generated by enzymatic hydrolysis 424(1988)129

**Diphenhydramine**

Screening procedure for the detection of alkanolamine antihistamines and their metabolites in urine using computerized GC-MS 428(1988)43

**Diphenyl**

Real-time quantitative analysis of unresolved LC peaks with the use of the Kalman filter 407(1987)59

**Diphenylamine**

Determination of diphenylamine residues in apples, and 4-aminobiphenyl residues in diphenylamine, by HPLC and electrochemical detection 447(1988)421

**Diphenylcyclobutanedicarboxylic acid**

Capillary GC-electron capture detection of coca-leaf-related impurities in illicit cocaine: 2,4-diphenylcyclobutane-1,3-dicarboxylic acids, 1,4-diphenylcyclobutane-2,3-dicarboxylic acids and their alkaloidal precursors, the truxillines 410(1987)297

**Diphenylpyraline**

Screening procedure for the detection of alkanolamine antihistamines and their metabolites in urine using computerized GC-MS 428(1988)43

**Dipipanone**

Measurement of dipipanone using capillary GC 424(1988)152

**N-{{[3,5-Di(pyrrolidinylmethyl)-4-hydroxy]benzoyl}aniline**

Development of a TLC method for the evaluation of antiarrhythmic N-{{[3,5-di(pyrrolidinylmethyl)-4-hydroxy]benzoyl}aniline (ACC-9358) 410(1987)222

**Disaccharides**

Elution characteristics of polyol-borate eluents in anion-exchange chromatography. I. Eluent strength of sugar-, uronate-borate eluents 403(1987)27

HPLC separation of some mono- and disaccharides with detection by a post-column enzyme reactor and a chemically modified electrode 408(1987)157

Rapid determination of disaccharides from chondroitin and dermatan sulphates by HPLC 414(1987)156

LC assay for constituent disaccharides of hyaluronic acid and chondroitin sulphate isomers 415(1987)231

Characterization of the products generated from oversulphated dermatan sulphate isomers with chondroitinase-B by HPLC 423(1987)51

**Disodium glyco-7 $\alpha$ ,12 $\alpha$ -dihydroxy-5 $\beta$ -cholan-24-oate 7 $\alpha$ -sulphate**

Internal standards for quantitative GC of individual bile acids after group separation of bile acids in urine 422(1987)25

**Disopyramide**

Simultaneous assay of disopyramide and monodesisopropylidisopyramide enantiomers in biological samples by LC 424(1988)424

Direct enantiomeric resolution of disopyramide and its metabolite using chiral HPLC.

Application to stereoselective metabolism and pharmacokinetics of racemic disopyramide in man 450(1988)211

**Dispensing speed**

Factors affecting sample transfer from microlitre syringes 441(1988)367

**Dispersive force parameters**

Dispersive force parameters as a measure of the polarity of surfactants used as stationary phases in GLC 435(1988)29

**Displacement chromatography**

Free energy consumption at self-sharpening concentration fronts in isothermal fixed-bed adsorption. Theory for isotherms of arbitrary shape and applications in displacement chromatography 409(1987)1

Displacement chromatography of biomolecules 439(1988)341

High-performance displacement chromatography of melanotropins and their derivatives 440(1988)157

Prediction of band profiles in displacement chromatography by numerical integration of a semi-ideal model 449(1988)25

**Displacer salts**

HPLC of amino acids, peptides and proteins. LXXXVI. The influence of different displacer salts on the retention and bandwidth properties of proteins separated by isocratic anion-exchange chromatography 443(1988)97

**Disposable bonded-phase columns**

Determination of aflatoxins in peanut products using disposable bonded-phase columns and post-column reaction detection 409(1987)413

**Disposable cartridges**

Evaluation of disposable cartridges for trace enrichment from aqueous solutions 402(1987)201

**Dissociation constants**

Isotachophoretic determination of mobility and  $pK_a$  by means of computer simulation. VI. Mobility and  $pK_a$  of sixty anions in methanol 408(1987)27

**Distribution coefficients**

Prediction of the distribution coefficient ( $K_d$ ) variation with operating conditions in chromatographic systems 411(1987)81

**Disulfiram**

Determination of disulfiram and two of its metabolites in urine by reversed-phase LC and spectrophotometric detection after post-column complexation 424(1988)95

**Diterpenes**

HPLC of glycosides on a new type of hydroxyapatite column 407(1987)205

**Diterpenoid alkaloids**

HPLC analysis of  $C_{19}$ -diterpenoid alkaloids 402(1987)366

**Dithia[3.3]phanes**

TLC separation of some dithia[3.3]phanes. I. 436(1988)338

Thin-layer separation of some dithia[3.3]phanes. II. 442(1988)378

**Dithiocarbamates**

HPLC identification and determination of dithiocarbamates in vulcanized rubber products 405(1987)263

 **$\beta$ -Dithioketones**

GC of metal  $\beta$ -diketonates and their analogues (Review) 411(1987)1

**1,3-Dithiolane-1-oxides**

Chromatographic separation of the enantiomers of 1,3-dithiolane-1-oxides 450(1988)199

**Dithiothreitol**

Simplified monitoring of reduced and oxidized dithiothreitol using HPLC 422(1987)239

**Diuretics**

LC screening of diuretics in urine 415(1987)347

**Diuron**

Analysis of bromacil, diuron and 3,4-dichloroaniline in contaminated well water, using a HPLC column-switching procedure 410(1987)211

Increasing extraction efficiency in supercritical fluid extraction from complex matrices.

Predicting extraction efficiency of diuron and linuron in supercritical fluid extraction using supercritical fluid chromatographic retention 447(1988)53

**Divinylbenzene**

GC on porous polymers. IV. Influence of the geometric structure of porous copolymers of 1,4-di(methacryloyloxymethyl)naphthalene with 1,4-divinylbenzene on their chromatographic behaviour 448(1988)233

Effect of the content of divinylbenzene in ion-exchange resins on the chromatographic separation of  $\alpha$ -cyclodextrin and glucose 448(1988)241

**DNA**

Isolation of high-molecular-weight nucleic acids for copy number analysis using HPLC 402(1987)189

Identification by HPLC and MS of a guanine

derivative arising from acid hydrolysis of DNA 407(1987)349

Effect of phosphate concentration on the yield and purity of DNA separated from RNA by hydroxyapatite chromatography 411(1987)481

Non-histone chromatin proteins that recognize specific sequences of DNA (Review) 418(1987)73

High-resolution chromatography of nucleic acids on the Gen-Pak FAX column 444(1988)47

**DNA adducts**

Oxidation-elimination of a DNA base from its nucleoside to facilitate determination of alkyl chemical damage to DNA by GC with electrophore detection 410(1987)453

**DNA, diamino-octane-derivatized**

Staphylococcal nuclease HPLC characterization of diamino-octane-modified DNA and its biotin and fluorescein derivatives 444(1988)97

**DNA polymerase**

Purification of nucleotide-linked peptide 444(1988)133

**DNA probes**

Preparation and chromatographic use of 5'-fluorescent-labelled DNA probes 444(1988)67

**DNA restriction fragments**

Chromatographic separation of DNA restriction fragments (Review) 418(1987)97

**Dns derivatives**

Copper(II) complexes of diamino-diamido-type ligands as chiral eluents in the enantiomeric separation of D,L-dansylamino acids by reversed-phase HPLC 441(1988)275

**Dodecylbenzenesulphonates**

Determination of dodecylbenzenesulphonates and ethoxylated alkylphenols in liquid pesticide formulations by HPLC 435(1988)73

**DOPA**

Separation of norepinephrine enantiomers, DOPA enantiomers and dopamine derivatized with *o*-phthalaldehyde-N-acetyl-L-cysteine by HPLC 402(1987)387

Simultaneous LC determination of seventeen of the major monoamine neurotransmitters, precursors and metabolites. I. Optimization of the mobile phase using factorial designs and a computer program to predict chromatograms 415(1987)261

Simultaneous LC determination of seventeen of the major monoamine neurotransmitters, precursors and metabolites. II. Assessment of human brain and cerebrospinal fluid concentrations 415(1987)275

Assay for tyrosine hydroxylase by HPLC with fluorescence detection 415(1987)289

Simultaneous determination of levodopa, its metabolites and carbidopa in clinical samples 415(1987)393

Assay for tyrosine hydroxylase in hypothalamic homogenates using HPLC with electrochemical detection 419(1987)27

### **Dopamine**

Separation of norepinephrine enantiomers, DOPA enantiomers and dopamine derivatized with *o*-phthalaldehyde-N-acetyl-L-cysteine by HPLC 402(1987)387

Determination of  $\beta$ -phenylalkylamines in pharmaceutical preparations by analytical capillary isotachopheresis 405(1987)389

Simultaneous LC determination of seventeen of the major monoamine neurotransmitters, precursors and metabolites. I. Optimization of the mobile phase using factorial designs and a computer program to predict chromatograms 415(1987)261

Simultaneous LC determination of seventeen of the major monoamine neurotransmitters, precursors and metabolites. II. Assessment of human brain and cerebrospinal fluid concentrations 415(1987)275

Determination of catecholamines in biological tissues by LC with coulometric detection 415(1987)388

Improved HPLC method with electrochemical detection for the determination of urinary catecholamines using a new electrochemical detector 416(1987)340

Combined ion-pair extraction and HPLC for the determination of the biogenic amines and their major metabolites in single brain tissue samples 421(1987)51

Single-step method for derivatization of dopamine and some related compounds in aqueous media for GC 421(1987)227

Assay for dopamine  $\beta$ -hydroxylase in human plasma and rat serum by HPLC with fluorimetric detection 421(1987)237

Routine determination of urinary free catecholamines by HPLC with electrochemical detection 424(1988)263

LC determination of urinary catecholamines after one-step alumina extraction 424(1988)373

Tritium isotope effect in reversed-phase HPLC analysis of dopamine and dihydroxyphenylacetic acid 427(1988)9

Sensitive method for determination of picogram amounts of epinephrine and other catecholamines in microdissected samples of rat brain using LC with electrochemical detection 433(1988)41

Perfluorinated acids as ion-pairing agents in the determination of monoamine transmitters and some prominent metabolites in rat brain by HPLC with amperometric detection 449(1988)191

### **Dopamine $\beta$ -hydroxylase**

Assay for dopamine  $\beta$ -hydroxylase in human plasma and rat serum by HPLC with fluorimetric detection 421(1987)237

### **Dopamine sulphates**

Direct measurement of dopamine O-sulfate in plasma and cerebrospinal fluid 413(1987)17

Preparation of dopamine 3-O-sulphate and dopamine 4-O-sulphate as reference substances and HPLC trace determination 430(1988)263

### **Doppler anemometry**

Integral Doppler anemometry and analytical field-flow fractionation 446(1988)141

### **Dosulepin**

Determination of dosulepin and its metabolite: application of HPLC with electrochemical detection 415(1987)365

### **Double-layer adsorption model**

Influence of additives to the eluent on hydrophobic interaction chromatography of simple compounds. V. Influence of pH and electrolytes on the retention of *n*-alkylammonium ions and *n*-alkyl carboxylates 438(1988)181

### **Doxepin**

Simultaneous determination of doxepin and nordoxepin in serum using HPLC 419(1987)445

### **Doxorubicin**

see Adriamycin

New method for the determination of



doxorubicin, 4'-epidoxorubicin and all known metabolites in cardiac tissue 424(1988)103

#### **Doxorubicinol**

Estimation of doxorubicin and doxorubicinol by HPLC and advanced automated sample processor 420(1987)184

#### **Doxycycline**

Improved HPLC determination of doxycycline in serum and urine using solid-phase extraction columns 434(1988)253

#### **Doxylamine**

Formation of artifactual metabolites of doxylamine following acid hydrolysis 419(1987)113

Screening procedure for the detection of alkanolamine antihistamines and their metabolites in urine using computerized GC-MS 428(1988)43

#### **Drug-protein binding**

Effects of drug-protein binding on trace enrichment of drugs in blood plasma on short precolumns 439(1988)353

#### **Drugs**

Retention behaviour of acidic, neutral and basic drugs on a CN column using phosphate buffers in the mobile phase 410(1987)77

Analysis of drugs from biological fluids using disposable solid phase columns 411(1987)325

Direct separation of drug enantiomers by HPLC with chiral stationary phases (Review) 426(1988)1

Polar contributions of the stationary phase to the reversed-phase ion-pair HPLC separation of quaternary ammonium drugs 437(1988)83

#### **Drugs, hydrophilic**

HPLC drug analysis by direct injection of whole blood samples. II. Determination of hydrophilic drugs 423(1987)155

#### **Drugs, hydrophobic**

HPLC drug analysis by direct injection of whole blood samples. III. Determination of hydrophobic drugs adsorbed on blood cell membranes 423(1987)163

#### **Drugs, moderately hydrophobic**

HPLC drug analysis by direct injection of whole blood samples. I. Determination of moderately hydrophobic drugs incorporated into blood corpuscles 423(1987)147

#### **Duoperone**

Determination of duoperone in plasma by HPLC 419(1987)452

#### **Dyes**

Evaluation of absorbance ratioing for solute identification in HPLC using a diode array detector 402(1987)135

Evaluation of disposable cartridges for trace enrichment from aqueous solutions 402(1987)201

Simple method for the analysis of food dyes on reversed-phase thin-layer plates 411(1987)437

Use of the dye Janus Green to facilitate quantitative analysis of phosphoinositides 424(1988)445

Design and applications of biomimetic anthraquinone dyes. I. Synthesis and characterisation of terminal ring isomers of C.I. Reactive Blue 2 435(1988)127

Adsorption chromatography on cellulose. II. Separations of aromatic amino acids, biogenic amines, alkaloids, dyes and phenols and determination of hydrophobic constants 440(1988)165

Characterisation of acid dyes in forensic fibre analysis by HPLC using narrow-bore columns and diode array detection 442(1988)187

Identification of xanthene dyes in lipsticks by reversed-phase HPLC 448(1988)296

New strategies for the screening of a large number of immobilized dyes for the purification of enzymes. Application to the purification of enzymes from human haemolysate 449(1988)403

Purification of human 6-phosphogluconate dehydrogenase from human haemolysate with chromatography on an immobilized dye as the essential step and use of automation.

Simultaneous purification of lactate dehydrogenase 449(1988)413

#### **Dynamic modelling**

Chemical engineering approach to dynamic modelling of linear chromatography. A flexible method for representing complex phenomena from simple concepts 406(1987)11

#### **Dynamic viscosity**

Characterization of mobile phases for the investigation of electrokinetic phenomena in LC 447(1988)177

**Dynorphins**

Combined HPLC–radioimmunoassay method for the analysis of endorphins, enkephalins and other neurotransmitter peptides 423(1987)93

**Dyphylline**

Microassay for the simultaneous determination of theophylline and dyphylline in serum by HPLC 416(1987)91

**E****Ebselen**

Determination of the two major metabolites of ebselen in human plasma by HPLC 430(1988)438

**Ecdysone acyl esters**

TLC of ecdysone acyl esters and their 2,3-acetonide derivatives 411(1987)379

Separation of ecdysone acyl esters and their 2,3-acetonide derivatives by HPLC 436(1988)279

**Ecdysteroids**

Separation of ecdysteroids by HPTLC using automated multiple development 408(1987)445

TLC of ecdysone acyl esters and their 2,3-acetonide derivatives 411(1987)379

Separation of ecdysone acyl esters and their 2,3-acetonide derivatives by HPLC 436(1988)279

Supercritical fluid chromatography of ecdysteroids 436(1988)497

Comparison of non-hydrophobic  $C_{18}$  bonded TLC and HPTLC plates for reversed-phase TLC of ecdysteroids 438(1988)419

Analysis of ecdysteroids by supercritical-fluid chromatography 441(1988)165

**Eclanamine**

HPLC method for the determination of eclanamine and its N-desmethyl and N,N-didesmethyl metabolites in urine 428(1988)131

**Economic laboratory practice**

Economic laboratory practice in preparative column HPLC 450(1988)81

**Eicosanoids**

HPLC separation of femtomolar quantities of endogenous carboxylic acids, including arachidonic acid metabolites, as 4-bromomethyl-7-acetoxycoumarin derivatives 416(1987)247

Chromatography of B prostaglandins on  $\beta$ -cyclodextrin silica: application to analysis of major E prostaglandins in human seminal fluid 421(1987)117

**Elastin**

HPLC determination of desmosine and isodesmosine in tissues and its application to studies of alteration of elastin induced by atherosclerosis 422(1987)53

**Elastomers**

Identification of polymer glycols in polyurethane elastomers by pyrolysis–GC 435(1988)219

**Eldanolide**

Formal total synthesis of rosefuran and eldanolide 440(1988)449

**Electrical potential**

Partitioning behaviour of rat bone marrow cells in aqueous two-phase systems. Dependence of cell partition on the interfacial tension and electrical potential difference between the phases 403(1987)131

**Electrochemical detector**

Thin-layer electrochemical detector coated with Nafion film for LC 410(1987)111

Determination of nitrodiphenylamines by LC and dual-electrode amperometric detection 410(1987)437

Analysis of dipeptides by reversed-phase HPLC without derivatization using amperometric detection on a copper electrode 436(1988)334

Amperometric detector with a manganin electrode for microcolumn LC 442(1988)111

**Electrochemical suppressor**

High-performance electrochemical suppressor for ion chromatography 439(1988)159

**Electrochemiluminescence detection**

Electrochemiluminescence as a detection technique for reversed-phase HPLC. IV. Detection of fluorescent derivatives 441(1988)394

**Electrode assembly**

Experimental and theoretical dynamics of

isoelectric focusing. II. Elucidation of the impact of the electrode assembly 436(1988)191

#### **Electrokinetic phenomena**

Characterization of mobile phases for the investigation of electrokinetic phenomena in LC 447(1988)177

#### **Electrolytes**

Influence of additives to the eluent on the hydrophobic interaction chromatography of simple compounds. IV. Influence of electrolytes on the retention of normal alkanols at 25 °C 437(1988)337

Influence of additives to the eluent on hydrophobic interaction chromatography of simple compounds. V. Influence of pH and electrolytes on the retention of *n*-alkylammonium ions and *n*-alkyl carboxylates 438(1988)181

#### **Electrolytic conductivity detector**

Analysis of ppb levels of organics in water by means of purge-and-trap, capillary GC and selective detectors 441(1988)73

#### **Electron-capture detector**

An experiment to distinguish between homogeneous and heterogeneous response mechanisms in a parallel-plate electron-capture detector 406(1987)81

Correlations and constants in unipolar and bipolar electron-capture detection 408(1987)53

Non-radioactive electron-capture detector operating in the pulsed mode 441(1988)45

Influence of the carrier gas flow-rate on the pulsed electron-capture detector 446(1988)329

#### **Electronic scanning**

Pros and cons of electronic scanning in densitometry and an experimental device for their verification 437(1988)97

#### **Electrophoresis**

Carrier-free zone electrophoresis, displacement electrophoresis and isoelectric focusing in a high-performance electrophoresis apparatus 403(1987)47

Computer analysis of proteins separated by polyacrylamide gradient pore gel electrophoresis 416(1987)265

Two-dimensional gel electrophoresis of proteins (Review) 418(1987)145

Gel electrophoresis of intact subcellular particles (Review) 418(1987)305

Agarose gel electrophoresis of bacteriophages and related particles (Review) 418(1987)345

Electrophoretic separation of cells (Review) 418(1987)359

#### **Ellagic acid**

Determination of free ellagic acid by reversed-phase HPLC 447(1988)284

#### **Ellipticine**

Fluorodensitometric assay of 6*H*-pyrido[4,3-*b*]-5,11-dimethylcarbazoles (ellipticine and derivatives) biosynthesized by *Ochrosia elliptica* cultures *in vitro* 409(1987)426

#### **Elution profiles**

Influence of the relative concentrations of two components of a binary mixture on their overloaded elution profiles 440(1988)9

#### **Elution temperature**

Calculation of programmed temperature GC characteristics from isothermal data. I. Theory and computational procedures 405(1987)67

#### **Elution volume**

Flow-rate effect on elution volume in SEC 408(1987)9

#### **Enantiomer separation**

Chiral separations of amino acids using di- and tri-peptides for zwitter-ion pair chromatography 402(1987)49

Stereoelectronic model to explain the resolution of enantiomeric ibuprofen amides on the Pirkle chiral stationary phase 402(1987)179

HPLC separation of diacylglycerol enantiomers on a chiral stationary phase 402(1987)257

New derivatization reagents for the resolution of carboxylic acid enantiomers by HPLC 402(1987)344

Resolution of the enantiomers of abscisic acid methyl ester by HPLC using a stationary phase of cellulose tris(3,5-dimethylphenylcarbamate)-coated silica gel 402(1987)371

Direct separation of (*S*)- and (*R*)-abscisic acid on a commercially available chiral HPLC column 402(1987)374

Separation of norepinephrine enantiomers, DOPA enantiomers and dopamine derivatized with *o*-phthalaldehyde-*N*-acetyl-L-cysteine by HPLC 402(1987)387

Separation of racemic pharmaceuticals by HPLC

- on silica gel modified with carbohydrate residues 403(1987)263
- Inductive adsorption as a new approach to amine enantiomer resolution 404(1987)87
- Reciprocity in chiral recognition. Comparison of several chiral stationary phases 404(1987)107
- Optical resolution of enantiomeric amino acid derivatives on a naphthylethylurea multiple-bonded chiral stationary phase prepared via an activated carbamate intermediate 404(1987)117
- Resolution of  $\alpha$ -arylacetic acid enantiomers on two chiral stationary phases 404(1987)123
- Chiral-phase HPLC of rotenoid racemates 404(1987)133
- Resolution of the enantiomers of thyroid hormones by high-performance ligand-exchange chromatography using a chemically bonded chiral stationary phase 404(1987)391
- Resolution of racemic amides and phosphine oxides on a  $\beta$ -cyclodextrin-bonded stationary phase by subcritical fluid chromatography 405(1987)135
- Chromatographic separation of racemic amino acids by use of chiral crown ether-coated reversed-phase packings 405(1987)145
- Separation of alkyl-substituted indene enantiomers by chromatography on microcrystalline triacetylcellulose 405(1987)155
- Polysiloxanes containing thermally stable chiral amide side-chains for capillary GC and, supercritical fluid chromatography 405(1987)169
- Chiral recognition of halo carboxylic acid amides studied by GC on Chirasil-Val 406(1987)157
- Ligand exchange chromatography of amino alcohol enantiomers as Schiff bases 406(1987)343
- Enantiomeric separation of amino acids by micro HPLC on an L-proline-bonded stationary phase 407(1987)151
- Applications for racemic versions of chiral stationary phases 407(1987)211
- Influence of enantiomeric purity of a chiral selector on stereoselectivity 407(1987)217
- Resolution of monoterpene enantiomers by GC 407(1987)243
- Chirality determinations for a tertiary alcohol. Ratios of linalool enantiomers in insects and plants 407(1987)324
- Effects of pretreatment on the enantioselectivity of silica-bound proteins used as HPLC stationary phases 408(1987)99
- HPLC method for determining the enantiomeric purity of a benzindene prostaglandin by a diastereomeric separation 408(1987)275
- Application of an ovomucoid-conjugated column for the optical resolution of some pharmaceutically important compounds 408(1987)316
- Determination of enantiomeric purity of 3-mercapto-2-methylpropionic acid 408(1987)365
- Resolution of a series of racemic phosphine oxides by LC on two structurally similar chiral stationary phases (in French) 409(1987)183
- HPLC separation of the enantiomers of substituted 2-aryloxypropionic acid methyl esters 410(1987)355
- Enantioselective capillary GC of 1,2-isopropylidene glycerol and O-alkylated glycerol derivatives 410(1987)458
- Resolution of enantiomeric aromatic alcohols on a cellulose tribenzoate HPLC chiral stationary phase. A proposed chiral recognition mechanism 411(1987)139
- Enantiomer separation of tocainide and some of its analogues and derivatives on a Chirasil-Val capillary column 411(1987)285
- Direct determination of the enantiomeric purity of (5*S*)-3-isopropyl-5-*p*-toluenesulphonyloxymethylloxazolidin-2-one on a cellulose-based chiral stationary phase. In-process control of a chiral intermediate used in the synthesis of enantiomerically pure  $\beta$ -blocking agents 411(1987)461
- Enantiomeric resolution of racemic nicotine and nicotine analogues by microcolumn LC with  $\beta$ -cyclodextrin inclusion complexes 411(1987)490
- Determination of the (+)- and (-)-enantiomers of pirofen in human plasma by HPLC 414(1987)131
- HPLC separation of peptide and amino acid stereoisomers 414(1987)313
- Determination of the enantiomers of ketoprofen in blood plasma by ion-pair extraction and HPLC of leucinamide derivatives 414(1987)465
- Determination of the enantiomeric composition of ibuprofen in human plasma by HPLC 415(1987)75

- Determination of flunoxapfen enantiomers in biological fluids by HPLC 415(1987)214
- Stereoselective analysis of the enantiomers of mexiletine by HPLC using fluorescence detection and study of their stereoselective disposition in man 415(1987)335
- Enantioselective HPLC determination of baclofen after derivatization with a chiral adduct of *o*-phthaldialdehyde 415(1987)419
- Rapid determination of the enantiomers of metoprolol, oxprenolol and propranolol in urine 416(1987)331
- HPLC assay of ketoprofen enantiomers in human plasma and urine 416(1987)388
- Simultaneous quantitation of *d*- and *l*-hexobarbital in rat blood by HPLC 419(1987)426
- Stereoselective HPLC assay for bupivacaine enantiomers 420(1987)203
- Determination of metoprolol enantiomers in plasma and urine using (S)-(–)-phenylethyl isocyanate as a chiral reagent 421(1987)161
- Analysis of salbutamol enantiomers in human urine by chiral HPLC and preliminary studies related to the stereoselective disposition kinetics in man 422(1987)187
- Formation of diastereomeric derivatives of 2-arylpropionic acids using L-leucinamide 423(1987)334
- Determination of drug enantiomers in biological samples by coupled column LC and LC-MS 424(1988)61
- Direct resolution of the stereoisomers of leucovorin and 5-methyltetrahydrofolate using a bovine serum albumin HPLC chiral stationary phase coupled to an achiral phenyl column 424(1988)158
- Separation of enantiomers of derivatised or underderivatised propranolol by means of HPLC 424(1988)189
- Simultaneous assay of disopyramide and monodesisopropylidisopyramide enantiomers in biological samples by LC 424(1988)424
- Rapid and sensitive HPLC assay of tiaprofenic acid enantiomers in human plasma and urine 425(1988)135
- Quantification of the enantiomers of ofloxacin in biological fluids by HPLC 425(1988)153
- Direct separation of drug enantiomers by HPLC with chiral stationary phases (Review) 426(1988)1
- HPLC resolution of the enantiomers of thioridazine, its metabolites and related compounds 426(1988)223
- Enantiomeric analysis of a new anti-inflammatory agent in rat plasma using a chiral  $\beta$ -cyclodextrin stationary phase 426(1988)417
- S-(+)-Flunoxapfen chloride as chiral fluorescent reagent 427(1988)131
- Determination of D,L-threo-3,4-dihydroxyphenylserine and of the D- and L-enantiomers in human plasma and urine 427(1988)219
- Method to determine the enantiomers of ibuprofen from human urine by HPLC 428(1988)103
- Stereoselective analysis of the enantiomers of ethotoin in human serum using chiral stationary phase LC and GC-MS 428(1988)402
- endo*-1,4,5,6,7,7-Hexachlorobicyclo[2.2.1]hept-5-ene-2-carboxylic acid, a superior resolving agent for the HPLC separation of enantiomers of hydroxylated derivatives of two azaaromatic hydrocarbons 430(1988)53
- Assay method for product formation in *in vitro* enzyme kinetic studies of uridine diphosphate glucuronyltransferases: 2-arylpropionic acid enantiomers 430(1988)368
- Preparative resolution of the enantiomers of the  $\beta$ -blocking drug atenolol by chiral derivatization and HPLC 431(1988)222
- Quantitation of the enantiomers of rimantadine in human plasma and urine by GC-MS 431(1988)343
- Enantioselective GC assay with electron-capture detection for *dl*-fenfluramine and *dl*-norfenfluramine in plasma 433(1988)105
- Determination of gossypol enantiomers in plasma after administration of racemate using HPLC with precolumn chemical derivatisation 433(1988)141
- Reversed-phase HPLC assay to quantitate diastereomeric derivatives of metoprolol enantiomers in plasma 433(1988)318
- Measurement of carprofen enantiomer concentrations in plasma and urine using L-leucinamide as the chiral coupling component 433(1988)331
- Computer-aided optimisation of drug enantiomer separation in chiral HPLC 434(1988)377

- Separation of  $\alpha$ -amino acid enantiomers by reversed-phase HPLC after derivatization with *o*-phthalaldehyde and a sodium salt of 1-thio- $\beta$ -D-glucose 434(1988)417
- Improved resolution of enantiomers of naproxen by the simultaneous use of a chiral stationary phase and a chiral additive in the mobile phase 435(1988)225
- Phosgene as a derivatizing reagent prior to GC and LC (Review) 435(1988)259
- Indirect HPLC resolution of racemic tertiary amines as their diastereomeric urea derivatives after N-dealkylation 436(1988)269
- Resolution of chiral barbiturates into enantiomers by reversed-phase HPLC using methylated  $\beta$ -cyclodextrins 436(1988)381
- Direct LC separation of enantiomers on immobilized protein stationary phases. VI. Optical resolution of a series of racemic barbiturates: studies of substituent and mobile phase effects 436(1988)479
- Chiral separation through a single substituent association 436(1988)517
- Separation of fatty acid ester and amide enantiomers by HPLC on chiral stationary phases 437(1988)109
- Analytical separations of enantiomeric primary amines as pyrrole derivatives formed by reaction with polygodial 438(1988)108
- Use of cyclodextrins in isotachopheresis. IV. The influence of cyclodextrins on the chiral resolution of ephedrine alkaloid enantiomers 438(1988)211
- Enantiomer separation of acyloins and cyanohydrins by enantioselective capillary GC 439(1988)301
- Use of cyclodextrins in isotachopheresis. V. The separation of ketotifen and its polar intermediate enantiomers 439(1988)386
- Chromatographic separation and isolation of the enantiomers of diethylstilbestrol metabolites 439(1988)484
- Optical resolution of samples with weak interactions on chiral "brush-type" LC stationary phases 440(1988)197
- Resolution of  $\alpha$ -substituted amino acid enantiomers by HPLC after derivatization with a chiral adduct of *o*-phthalaldehyde. Application to glutamic acid analogues 440(1988)209
- Paper electrophoretic study of ion-pair formation. XVII. Influence of the enantiomeric purity of the sodium tartrate used as electrolyte 440(1988)361
- Resolution of stereoisomers of dipeptides by GC on Chirasil-Val 441(1988)89
- Enantiomer analysis by complexation GC. Scope, merits and limitations 441(1988)135
- Separation of enantiomers on diluted permethylated  $\beta$ -cyclodextrin by high-resolution GC 441(1988)155
- Copper(II) complexes of diamino-diamido-type ligands as chiral eluents in the enantiomeric separation of D,L-dansylamino acids by reversed-phase HPLC 441(1988)275
- An improved chiral stationary phase for the facile separation of enantiomers 441(1988)311
- Structural factors affecting chiral recognition and separation on  $\beta$ -cyclodextrin bonded phases 441(1988)376
- Direct enantiomeric separation of racemic flavanones by HPLC using cellulose triacetate as a chiral stationary phase 441(1988)417
- Determination of the chiral purity of dipeptide isosteres containing a reduced peptide bond by GC analysis 442(1988)165
- Effect of the mobile phase composition and ligand structure on the separation of D- and L-dansylamino acids, as mixed metal complexes, by reversed-phase HPLC 445(1988)219
- Resolution of enantiomeric drugs of some  $\beta$ -amino alcohols as their urea derivatives by HPLC on a chiral stationary phase 447(1988)208
- Supercritical fluid chromatography with ion-pairing modifiers. Separation of enantiomeric 1,2-aminoalcohols as diastereomeric ion pairs 447(1988)287
- Resolution of lombricine enantiomers by HPLC utilising pre-column derivatisation with *o*-phthalaldehyde-chiral thiols 447(1988)392
- TLC enantiomeric resolution via ligand exchange 448(1988)11
- Planar chromatographic separation of enantiomers and diastereomers with cyclodextrin mobile phase additives 448(1988)345
- Separation of pyridone carboxylic acid enantiomers by HPLC using copper(II)-L-amino acid as the eluent 448(1988)439
- Chromatographic resolution. XXI. Direct optical resolution of abscisic acid by HPLC on cellulose tris(3,5-dimethylphenylcarbamate) 448(1988)454

Solvent extraction clean-up for pre-treatment in amino acid analysis by GC. Application to age estimation from the D/L ratio of aspartic acid in human dentine 449(1988)165

Some aspects of the selection of HPLC methods for the separation of chiral compounds in pharmaceutical analysis 450(1988)145

Hexahelicene chiral stationary phase. I. Phase synthesis and use in HPLC resolution of enantiomers 450(1988)450(1988)157

Chiral stationary phase derived from L-lactic acid for the optical resolution of N-(3,5-dinitrobenzoyl)amino acid methyl esters 450(1988)163

Separation and identification of enantiomers by HPLC with a chiral column and a polarimetric detector as applied to deltamethrin 450(1988)169

Chiral Michael additions of acetamidomalonnate to  $\alpha$ -enones. NMR and HPLC resolution of enantiomers 450(1988)183

Direct HPLC resolution of dihydropyridine enantiomers 450(1988)191

Chromatographic separation of the enantiomers of 1,3-dithiolane-1-oxides 450(1988)199

Direct enantiomeric resolution of disopyramide and its metabolite using chiral HPLC. Application to stereoselective metabolism and pharmacokinetics of racemic disopyramide in man 450(1988)211

Selective effect of clonazepam and (*S*)-uxepam on the binding of warfarin enantiomers to human serum albumin 450(1988)217

Separation of the stereoisomers of an allenic E-type prostaglandin 450(1988)233

Applicability of forced-flow planar chromatographic methods for the separation of enantiomers on Chiralplate 450(1988)241

Chiral resolution of a series of 3-thienylcyclohexylglycolic acids by liquid or subcritical fluid chromatography. A mechanistic study 450(1988)255

Chromatographic resolution of dipeptide enantiomers and diastereomers on chiral stationary phases from poly(L-leucine) or poly(L-phenylalanine) 450(1988)271

Preparative chiral separation in an aqueous two-phase system by a few counter-current extractions 450(1988)277

Separation and determination of some amino

acid ester enantiomers by TLC after derivatization with (*S*)-(+)-naproxen 450(1988)281

Rapid optical resolution of anionic metal complexes by GPC 450(1988)284

### End-capping

Selectivity related to carbon loading and end-capping of octadecyl-bonded silica gels in reversed-phase LC 441(1988)183

### $\beta$ -Endorphin

Combined use of size-exclusion and reversed-phase HPLC for characterization of  $\beta$ -endorphin processing pathways 416(1987)255

### Endorphins

Combined HPLC–radioimmunoassay method for the analysis of endorphins, enkephalins and other neurotransmitter peptides 423(1987)93

### Endotoxins

Determination of endotoxins by GC: evaluation of electron-capture and negative-ion chemical-ionization MS detection of halogenated derivatives of  $\beta$ -hydroxymyristic acid 417(1987)11

TLC of endotoxins, their derivatives and contaminants 435(1988)167

### Enflurane

Quantification of halothane, enflurane and nitrous oxide by means of a simple GC method 403(1987)350

### Enkephalinase

Assay for enkephalin-degrading peptidases in rat brain tissues by HPLC with on-line post-column fluorescence detection 430(1988)291

### Enkephalins

HPLC of N-terminal tyrosine-containing oligopeptides by pre-column fluorescence derivatization with hydroxylamine, cobalt(II) and borate reagents 411(1987)305

Separation of protein hormones (Review) 418(1987)245

On-line post-column fluorescence detection for N-terminal tyrosine-containing peptides in HPLC 421(1987)245

Combined HPLC–radioimmunoassay method for the analysis of endorphins, enkephalins and other neurotransmitter peptides 423(1987)93

HPLC determination of leucine-enkephalin-like

peptide in rat brain by pre-column fluorescence derivatization involving formylation followed by reaction with 1,2-diamino-4,5-dimethoxybenzene 430(1988)271

Assay for enkephalin-degrading peptidases in rat brain tissues by HPLC with on-line post-column fluorescence detection 430(1988)291

Simultaneous analysis of methionine- and leucine-enkephalin from rat brain: quantification by LC-electrochemistry 431(1988)65

#### **Enpiroline**

Analysis of DL-*threo*- $\alpha$ -(2-piperidyl)-2-trifluoromethyl-6-(4-trifluoromethylphenyl)-4-pyridinemethanol phosphate (enpiroline), a new candidate antimalarial, in blood using HPLC 419(1987)458

#### **Enprofylline**

HPLC determination of enprofylline in human plasma 430(1988)175

#### **Enprostil**

Separation of the stereoisomers of an allenic E-type prostaglandin 450(1988)233

#### **Enterotoxin**

Alternative procedure for the purification of the heat-stable enterotoxin of enterotoxigenic *Escherichia coli* pathogenic for calves 420(1987)253

#### **Enthalpies of displacement**

Adsorption of binary solvent mixtures in reversed-phase chromatographic systems. Surface excess isotherms and enthalpies of displacement 406(1987)257

#### **Enthalpy-entropy compensation**

Enthalpy-entropy compensation in reversed-phase HPLC of series of aryloxoalkanoic and arylhydroxyalkanoic acids 449(1988)391

#### **Enzymatic hydrolysate**

Preparative separation of an enzymatic hydrolysate of  $^{14}$ C-labelled proteins 411(1987)335

#### **Enzyme reactors**

HPLC separation of some mono- and disaccharides with detection by a post-column enzyme reactor and a chemically modified electrode 408(1987)157

#### **Enzymes**

see also Proteins

Use of chromatofocusing for separation of  $\beta$ -lactamases. VIII. Analytical chromatofocusing of chromosomal cephalosporinases from four *Klebsiella* strains 403(1987)217

Fractionation and characterization of two  $\beta$ -lactamases in *Citrobacter diversus* ULA-27 strain by chromatofocusing 403(1987)366

Improved isolation of rat microsomal cytochrome  $b_5$  reductase 405(1987)319

Retention mechanism of lactate dehydrogenase in anion-exchange chromatography 406(1987)237

Matrix evaluation for preparative high-performance affinity chromatography 407(1987)179

Resolution of five position-specific flavonoid O-methyltransferases by fast protein liquid chromatofocusing 407(1987)291

Assay of glycosidase by lectin affinity HPLC 407(1987)299

Characterization of bee venom and its main components by HPLC 408(1987)341

HPLC purification of the H4 and MH3 isoenzymes of lactate dehydrogenase from beef heart 408(1987)372

Rapid chromatographic assay procedure for peptide-N-glycosidase activity 408(1987)378

Immunoaffinity purification of  $\beta$ -amylase from mustard (*Sinapis alba* L.) cotyledons 408(1987)385

Improved method for the detection of peroxidase isoenzymes after electrophoretic separation with chlorophyllin, cysteamine and light as a hydrogen peroxide generating system 410(1987)466

Effect of polymer structure on affinity partitioning of lactate dehydrogenase in polymer-water two-phase systems 411(1987)161

Reaction of Fast Blue BB salt and *p*-dimethylaminocinnamaldehyde with indole-3-acetic acid degradation products. Evaluation of the coupling properties of the reagents used in the zymographic screening of indole-3-acetic acid oxidases 411(1987)476

Purification of  $\alpha$ -amylase by industrial autofocusing 411(1987)486

HPLC assay of chymotrypsin and chymotrypsin-like enzyme activity 411(1987)498

Sugar-chain heterogeneity of human alkaline phosphatases: differences between normal and tumour-associated isozymes 413(1987)65



- Automated separation and measurement of urinary isoenzymes and protein by ion-exchange LC 414(1987)47
- Determination of kinase I and kinase II activities in human urine by HPLC 414(1987)423
- GC-MS determination of tryptophan transaminase-catalyzed deuterium exchange 415(1987)104
- Assay for tyrosine hydroxylase by HPLC with fluorescence detection 415(1987)289
- Subfractionation of the dansylated derivatives of glucosyl galactosyl hydroxylysine by LC and its application to a specific  $\alpha$ -1,2-glucosidase assay 416(1987)37
- Rapid chromatographic purification of dipeptidyl-aminopeptidase II from human kidney 416(1987)131
- TLC using multiple development for analysis of reaction products of sucrases 416(1987)353
- Highly sensitive assay for alkaline and acid phosphatase activities by HPLC with electrochemical detection 417(1987)41
- HPLC assay for pteroylpolylglutamate hydrolase 417(1987)47
- Separation of human pepsins in gastric juice by high-performance ion-exchange chromatography 417(1987)391
- Detection of uridine 5'-diphosphate glucose-collagen glucosyltransferase activity by HPLC and possible participation of a novel intermediate as a glucose donor in collagen biosynthesis 417(1987)403
- Non-histone chromatin proteins that recognize specific sequences of DNA (Review) 418(1987)73
- Assay for tyrosine hydroxylase in hypothalamic homogenates using HPLC with electrochemical detection 419(1987)27
- LC assay of the relative activities of serum pancreatic and salivary  $\alpha$ -amylase using reductively pyridylaminated maltopentaose as a fluorescent substrate 419(1987)51
- Differences in the constitutive forms of hepatic cytochrome P-450 in male and female adult beagle dogs 419(1987)61
- Rapid and sensitive assay of ornithine decarboxylase activity by HPLC of the *o*-phthalaldehyde derivative of putrescine 419(1987)291
- Semi-micro quantitative assay for tryptophan oxygenase by HPLC 419(1987)296
- Quantitative monitoring by HPLC of the dissociation of human serum cholinesterase by limited proteolysis 420(1987)25
- GC assay for cysteine sulphinate decarboxylase activity in animal tissues 420(1987)135
- Use of ion-exchange chromatography and HPLC to prepare a substrate for differential assay of human pancreatic and salivary  $\alpha$ -amylases in serum 420(1987)158
- Blood appearance of rat alkaline phosphatase originating from the duodenum in vitro 420(1987)275
- Assay of 5'-nucleotidase and simultaneous detection of interfering enzymes by anion-exchange CC 420(1987)398
- Assay for dopamine  $\beta$ -hydroxylase in human plasma and rat serum by HPLC with fluorimetric detection 421(1987)237
- Biochemical indices of renal damage: separation of urinary alanine aminopeptidase by LC 421(1987)344
- Purification of human renin by renin inhibitor (ES-305) affinity chromatography 421(1987)350
- Assay for trimethyllysine hydroxylase by HPLC 422(1987)253
- Determination of lipoamidase activity by LC with fluorimetric detection 423(1987)304
- HPLC separation of peptides possessing a proline residue in the amino-terminal penultimate position, and their products generated by enzymatic hydrolysis 424(1988)129
- Determination of phenol sulphotransferase activity by HPLC 424(1988)136
- Affinity chromatography of human thrombin on modified silica 424(1988)385
- Highly sensitive assay for xanthine oxidase activity by HPLC with fluorescence detection 424(1988)392
- Determination of agmatine, arginine, citrulline and ornithine by reversed-phase LC using automated pre-column derivatization with *o*-phthalaldehyde 425(1988)269
- Microassay for GM<sub>1</sub> ganglioside  $\beta$ -galactosidase activity using HPLC 426(1988)75
- Hydrophobic chromatography: a one-step

- method for the purification of human liver microsomal epoxide hydrolase 426(1988)169
- Assay of  $\gamma$ -L-glutamylcyclotransferase activity in rat brain synaptosomes by HPLC 426(1988)381
- Simple assay procedure for tyrosine hydroxylase activity by HPLC employing coulometric detection with minimal sample preparation 427(1988)229
- Biospecific sorption of cyclodextrin glucosyltransferase on physically modified starch 427(1988)331
- Human urinary  $\beta$ -mannosidase: behaviour towards two chromatographic supports 428(1988)203
- Improved methods for purification and assay of glycerol kinase from *Escherichia coli* 428(1988)345
- Use of a post-column enzymatic reaction for detection of macroamylasemia by HPLC 428(1988)415
- Lactate dehydrogenase isoenzymes (Review) 429(1988)373
- Creatine kinase isoenzymes (Review) 429(1988)399
- Alkaline phosphatase. Laboratory and clinical implications (Review) 429(1988)419
- Assay for enkephalin-degrading peptidases in rat brain tissues by HPLC with on-line post-column fluorescence detection 430(1988)291
- Assay method for product formation in *in vitro* enzyme kinetic studies of uridine diphosphate glucuronyltransferases: 2-arylpropionic acid enantiomers 430(1988)368
- HPLC assay of erythrocyte enzyme activity levels involved in vitamin B<sub>6</sub> metabolism 431(1988)406
- Simultaneous assay for individual sulphated 3 $\alpha$ - and  $\beta$ -hydroxysteroids in serum using HPLC combined with 3 $\alpha$ - and  $\beta$ -hydroxysteroid dehydrogenases immobilized on one column 432(1988)47
- Assay of ATP-sulfurylase activity from rat liver by HPLC 433(1988)63
- Purification of glutathione S-transferases from rat liver and Walker 256 mammary carcinoma cells by HPLC and a glutathione affinity column 434(1988)83
- HPLC determination of acid and alkaline phosphatases in urine 434(1988)196
- Sensitive assay for determination of pyridoxal-5-phosphate in enzymes using HPLC after derivatization with cyanide 434(1988)209
- Determination of adenosine deaminase and purine nucleoside phosphorylase activities using HPLC and radiochromatographic methods 434(1988)439
- Electrophoretic study of  $\alpha$ -D-galactosidases from seeds of *Glycine soja* and *Vigna radiata* possessing erythroagglutinating activity 436(1988)59
- Analytical strategy for determination of active site sequences in aminoacyl-tRNA synthetases 438(1988)347
- High-performance anion-exchange chromatography of rennet enzymes 438(1988)451
- Chromatographic isolation of primary and secondary metabolites from plant cell cultures 440(1988)53
- Survey of the folding pathway of a two-domain protein phosphoglycerate kinase 440(1988)421
- Synthetic peptide substrates as models to study a pro-ocytocin/neurophysin converting enzyme 440(1988)439
- HPLC method for the determination of the esterase activity of subtilisin and kallikrein 441(1988)431
- Purification of germinated barley  $\alpha$ -amylase isozymes and limit-dextrinase by chromatofocusing and affinity chromatography 441(1988)436
- Conformational studies of bovine alkaline phosphatase in hydrophobic interaction and size-exclusion chromatography with linear diode array and low-angle laser light scattering detection 442(1988)29
- Micro-assay for sheep 11 $\beta$ -hydroxylase activity using HPLC for steroid analysis 442(1988)424
- Surface-mediated retention effects of subtilisin site-specific variants in cation-exchange chromatography 443(1988)193
- Chromatographic resolution of lysozyme variants 443(1988)221
- Separation of human factor X from factor Xa by reversed-phase HPLC 444(1988)107
- Semi-preparative purification of recombinant human renin and prorenin 444(1988)203
- Separation and characterization of two forms of

acetohydroxy acid synthase from black Mexican sweet corn cells 444(1988)251

Activity staining of inulin fructotransferase II on polyacrylamide gels 445(1988)282

Modified general affinity adsorbent for large-scale purification of penicillinases 448(1988)109

New strategies for the screening of a large number of immobilized dyes for the purification of enzymes. Application to the purification of enzymes from human haemolysate 449(1988)403

Purification of human 6-phosphogluconate dehydrogenase from human haemolysate with chromatography on an immobilized dye as the essential step and use of automation. Simultaneous purification of lactate dehydrogenase 449(1988)413

Rapid and efficient method for the isolation and characterization of plant aromatic choline esterases 450(1988)121

#### **Eosin**

Simple method for the analysis of food dyes on reversed-phase thin-layer plates 411(1987)437

#### **Ephedrine**

Determination of  $\beta$ -phenylalkylamines in pharmaceutical preparations by analytical capillary isotachopheresis 405(1987)389

Use of cyclodextrins in isotachopheresis. IV. The influence of cyclodextrins on the chiral resolution of ephedrine alkaloid enantiomers 438(1988)211

#### **Ephithelial allergens**

Purification of allergens by HPLC. V.

Purification of cat and dog epithelial allergens 404(1987)378

#### **Epicatechin**

Analysis of catechin and epicatechin by HPLC after benzylation 448(1988)291

#### **Epichlorohydrin**

Determination of residual epichlorohydrin and 3-chloropropanediol in water by GC with electron-capture detection 439(1988)448

#### **4'-Epidoxorubicin**

Improved method for the determination of 3'-epidoxorubicin and seven metabolites in plasma by HPLC 417(1987)339

New method for the determination of doxorubicin, 4'-epidoxorubicin and all known metabolites in cardiac tissue 424(1988)103

#### **Epifluorescence microscope**

Characterization of organotin species using microbore and capillary LC techniques with an epifluorescence microscope as a novel imaging detector 410(1987)393

#### **Epigalanthamine**

Reversed-phase LC for the determination of galanthamine and its metabolites in human plasma and urine 421(1987)396

#### **Epinephrine**

see Adrenaline

#### **Epinine**

Selected-ion monitoring GC-MS analysis of catecholamines: enhancement of sensitivity by a simple clean-up step on Sephadex G-10 433(1988)31

#### **4-Epioxytetracycline**

Quantitative analysis of oxytetracycline and related substances by HPLC 405(1987)229

#### **Epoxide hydrolase**

Hydrophobic chromatography: a one-step method for the purification of human liver microsomal epoxide hydrolase 426(1988)169

#### **Epoxidised soya bean oil**

Analysis of the epoxidised soya bean oil additive in plastics by GC 437(1988)247

#### **10,11-Epoxy carbamazepine**

Simultaneous determination of carbamazepine, oxcarbazepine and their main metabolites in plasma by LC 414(1987)477

#### **Epoxy resins**

Quantitative determination of free primary amines in polymeric amine hardeners. HPLC separation of ethylenediamine and triethylenetetramine after derivatization with salicylaldehyde-diphenylboron chelate and separation of 4,4'-diaminodiphenylmethane by HPLC 436(1988)447

Chromatographic determination of residual monomers of epoxy resins in wine simulants (in French) 445(1988)107

#### **Eproxindine**

Determination of eproxindine hydrochloride in human plasma by capillary GC 427(1988)157

#### **Equilenin**

HPLC determination of conjugated estrogens in tablets 450(1988)414

**Equilibrium constants**

Interaction of proteins with immobilized  $\text{Cu}^{2+}$ . Quantitation of adsorption capacity, adsorption isotherms and equilibrium constants by frontal analysis 403(1987)197

**Equilin**

HPLC determination of conjugated estrogens in tablets 450(1988)414

**Equilin sulphate**

Analysis of estrogen sulphate mixtures in pharmaceutical formulations by reversed-phase chromatography 402(1987)273

**Equine oestrogens**

Complete separation of nine equine oestrogens by HPLC 435(1988)385

**Equivalent chain-lengths**

Equivalent chain-lengths of methyl ester derivatives of fatty acids on GC. A reappraisal 447(1988)305

**Equivalent chain lengths**

Effect of overload of capillary GLC columns on the equivalent chain lengths of  $\text{C}_{18}$  unsaturated fatty acid methyl esters 447(1988)43

**Equivalent temperature**

Calculation of programmed temperature GC characteristics from isothermal data. I. Theory and computational procedures 405(1987)67

**Ercalcidiol**

Rapid automated HPLC assay for ercalcidiol and calcidiol (25-hydroxyvitamins  $\text{D}_2$  and  $\text{D}_3$ ) using *trans*-calcidiol as an UV-absorbing internal standard 415(1987)305

**Ergothioneine**

Determination of ergothioneine in red blood cells by HPLC 434(1988)191

**Eriodictyol**

Direct enantiomeric separation of racemic flavanones by HPLC using cellulose triacetate as a chiral stationary phase 441(1988)417

**Erythromycin**

HPLC assay of erythromycin and its esters using electrochemical detection 419(1987)205

**Erythrocytes**

Separation by counter-current distribution of youngchick erythrocyte populations and

numerical resolution of distribution curves 411(1987)167

Column-based separation of erythrocytes using aqueous polymeric two-phase systems 432(1988)127

Partitioning behavior of erythrocytes in aqueous two-phase systems containing hydroxypropyl starch and polyethylene glycol 441(1988)261

Cell separation by immunoaffinity partitioning with polyethylene glycol-modified protein A in aqueous polymer two-phase systems 442(1988)219

Partitioning behaviour in aqueous two-phase systems and fractionation by counter-current distribution of chick embryo erythrocytes with numerical resolution of distribution curves 445(1988)327

**Erythromycin**

Separation of novel derivatives from commercial erythromycin samples by TLC 403(1987)343

Optimization of the separation of erythromycin and related substances by HPLC 409(1987)91

Determination of erythromycin in human plasma, using column LC with a polymeric packing material, alkaline mobile phase and amperometric detection 423(1987)189

HPLC of erythromycin propionyl ester and erythromycin base in biological fluids 427(1988)93

Analysis of erythromycin and roxithromycin in plasma or serum by HPLC using electrochemical detection 433(1988)298

TLC study of the metabolites of erythromycins in the Wistar rat 434(1988)177

Isolation of erythromycin E from commercial erythromycin 445(1988)290

**Erythropoietin**

Purification of human urinary erythropoietin 417(1987)178

**Erythrosin**

Simple method for the analysis of food dyes on reversed-phase thin-layer plates 411(1987)437

***Escherichia coli* enterotoxin**

Alternative procedure for the purification of the heat-stable enterotoxin of enterotoxigenic *Escherichia coli* pathogenic for calves 420(1987)253

**Eseroline**

Metabolism of physostigmine in mouse liver microsomal incubations studied by LC with dual-electrode amperometric detection 419(1987)165

**Esmolol**

Analysis of esmolol in human blood by HPLC and its application to pharmacokinetic studies 424(1988)430

**Essential oils**

see also Volatile oils

Importance of the odoriferous principle of Bulgarian rose oil in flavour and fragrance chemistry 406(1987)181

Composition of essential oils. Part 7. Bulgarian oil of rose (*Rosa damascena* Mill.) 406(1987)185

Separation of non-polar sesquiterpene olefins from tolu balsam by HPLC; silver perchlorate impregnation of a prepacked preparative silica gel column 407(1987)343

Considerations and remarks on the analysis of *Anthemis nobilis* L. essential oil by capillary GC and "hyphenated" techniques 411(1987)237

Constituents of essential oil of *Thymus carnosus* Boiss. 436(1988)103

GC characterization of frequently occurring aromatic compounds in essential oils 446(1988)267

TLC characterization of essential oils 448(1988)301

GC separation and chemometric analysis of mandarin essential oils 449(1988)183

**Esterases**

HPLC method for the determination of the esterase activity of subtilisin and kallikrein 441(1988)431

**Esterification**

GC analysis of different homologous series of acids esterified in aqueous solutions with butyl and propyl alcohols 446(1988)237

Rapid radio TLC for assay of lipase-catalyzed esterification and interesterification reactions 450(1988)448

**Esters**

Factor analysis of trends in selectivity in GLC of 175 saturated and unsaturated esters on 14 polysiloxane stationary phases 404(1987)11

Comparison of volatile components between fresh and rotten mussels by GC-MS 409(1987)251

Glass capillary GC analysis of volatile components of canned Philippine pineapple juice 409(1987)409

Organo-modified smectite for the GC separation of various compounds 410(1987)121

**Etanidazole**

Gradient HPLC method for simultaneous assay of the radiosensitizers etanidazole (SR 2508) and pimonidazole (Ro 03-8799) in biological materials 420(1987)223

**Ethacrynic acid**

GC determination of the high-ceiling diuretic ethacrynic acid and its thiol adducts in plasma of guinea pigs 425(1988)396

**Ethambutol**

Determination of ethambutol in plasma by LC and UV spectrophotometric detection (in French) 415(1987)85

**Ethanol**

Dynamic headspace GC of volatile compounds in milk 404(1987)163

Simultaneous determination of methanol, ethanol, acetone, isopropanol and ethylene glycol in plasma by GC 414(1987)248

Static headspace GC method for the determination of residual solvents in vigabatrin drug substance 438(1988)103

Determination of alcohols in alcoholic beverages by micro HPLC with indirect photometric detection 445(1988)139

Rapid HPLC method for determination of ethanol and fusel oil in the alcoholic beverage industry 447(1988)443

Rapid HPLC method for determination of ethanol and fusel oil in the alcoholic beverage industry 447(1988)443

Validity of *post mortem* chest cavity blood ethanol determinations 449(1988)337

**Ethinylestradiol**

Selective on-line trace enrichment for the determination of ethynyl steroids in urine by LC with precolumn technology 417(1987)159

Medroxyprogesterone acetate- and ethinylestradiol-induced changes in biliary bile acids of the rat studied by HPLC 423(1987)63

Rapid and simple dansylation of phenolic steroids using a two-phase system and phase transfer catalysis 436(1988)429

#### **Ethynyl steroids**

Selective on-line trace enrichment for the determination of ethynyl steroids in urine by LC with precolumn technology 417(1987)159

#### **Ethopropazine**

HPLC resolution of the enantiomers of thioridazine, its metabolites and related compounds 426(1988)223

#### **Ethotoin**

Stereoselective analysis of the enantiomers of ethotoin in human serum using chiral stationary phase LC and GC-MS 428(1988)402

#### **3-[1-Ethoxycarbonyl-3-phenyl-(1*S*)-propyl]-amino-2,3,4,5-tetrahydro-2-oxo-1-(3*S*)-benzazepin**

Determination of a new angiotensin converting enzyme inhibitor and its active metabolite in plasma and urine by GC-MS 419(1987)123

#### **Ethoxyiated alkylphenols**

Determination of dodecylbenzenesulphonates and ethoxylated alkylphenols in liquid pesticide formulations by HPLC 435(1988)73

#### **Ethyl acetate**

Adsorption isotherms of ethyl acetate modifier on silica from supercritical carbon dioxide 409(1987)55

#### **Ethylbenzene**

Heavy alkylated benzene stationary phase for the GLC separation of ethylbenzene, xylenes, styrene, iso- and *n*-propylbenzene 409(1987)371

#### **Ethyl carbamate**

GC-MS determination of ethyl carbamate in alcoholic beverages 402(1987)233

#### **O-Ethyl S-[2-(diisopropylamino)ethyl] methylphosphonothiolate**

Analysis of O-ethyl S-[2-(diisopropylamino)ethyl] methyl-phosphonothiolate (VX) by capillary column GC-MS 402(1987)221

#### **Ethylenediamine**

Quantitative determination of free primary amines in polymeric amine hardeners. HPLC separation of ethylenediamine and triethylenetetramine after derivatization with salicylaldehyde-diphenylboron chelate and

separation of 4,4'-diaminodiphenylmethane by HPLC 436(1988)447

#### **Ethylene glycol**

Simultaneous determination of methanol, ethanol, acetone, isopropanol and ethylene glycol in plasma by GC 414(1987)248

Determination of ionic metabolites from ethylene glycol in human blood by isotachopheresis 416(1987)111

Rapid determination of ethylene glycol at toxic levels in serum and urine 434(1988)469

#### **Ethylene oxide**

Automated analysis of the gas mixture from ethylene epoxidation by oxygen and nitrous oxide mixtures using GC 450(1988)399

#### **N-(N-Ethylglycyl)-2,6-xylidine**

In vivo metabolism of lidocaine in the rat. Isolation of urinary metabolites as pentafluorobenzoyl derivatives and their identification by combined GC-MS 421(1987)267

#### **Ethylindenestrol A**

Chromatographic separation and isolation of the enantiomers of diethylstilbestrol metabolites 439(1988)484

#### **Ethyl methacrylate**

Normal- and reversed-phase separation of copolymers prepared from styrene and ethyl methacrylate 403(1987)280

#### **Ethyl methylphosphonic acid**

Analysis of alkyl methylphosphonic acids in aqueous matrices by ion-pair reversed-phase ion chromatography 407(1987)399

#### **Ethylmorphine**

Identification of opiates in urine by capillary column GC of two different derivatives 422(1987)117

#### **4-Ethylphenol**

Determination of simple phenols in faeces and urine by HPLC 431(1988)143

#### **Ethynylloestradiol**

Optimization of the reversed-phase HPLC separation of synthetic estrogenic and progestogenic steroids using the multi-criteria decision making method 410(1987)1

**Etoposide**

Comparison of flow-injection analysis with HPLC for the determination of etoposide in plasma 432(1988)395

**Etretin**

Simultaneous determination of the aromatic retinoids etretin and etretinate and their main metabolites by reversed-phase liquid chromatography 415(1987)413

Determination of the aromatic retinoids (etretin and isotretin) in biological fluids by HPLC 421(1987)177

Simultaneous microassay for etretinate and its active metabolite, etretin, by reversed-phase HPLC 433(1988)345

**Etretinate**

Simultaneous determination of the aromatic retinoids etretin and etretinate and their main metabolites by reversed-phase liquid chromatography 415(1987)413

Quantitative analysis of retinoids in biological fluids by HPLC using column switching. II.

Simultaneous determination of etretinate, acitretin and 13-*cis*-acitretin in plasma 431(1988)297

Simultaneous microassay for etretinate and its active metabolite, etretin, by reversed-phase HPLC 433(1988)345

**Europium**

LC and flow injection analysis of tetracycline using sensitized europium(III) luminescence detection 433(1988)149

Lanthanide ions as luminescent chromophores for the LC detection of polynucleotides and nucleic acids 436(1988)299

Dynamic quenching of europium(III) and terbium(III) luminescence: a potential detection method in ion chromatography 439(1988)165

**Evans blue**

Foam counter-current chromatography with the cross-axis synchronous flow-through coil planet centrifuge 403(1987)77

**Evaporation**

Sample reconcentration by column-external solvent evaporation or injection of large volumes into GC capillary columns? 404(1987)297

**Evaporation rate**

Evaporation rates of solvent blends measured by GC 402(1987)65

**Evaporative light scattering detector**

Application fields of a new evaporative light scattering detector for HPLC and supercritical fluid chromatography (in French) 404(1987)95

**Explosives**

Improved method for the trace analysis of explosives by silica capillary column GC with thermal energy analysis detection 410(1987)181

Trace analysis of explosives by capillary supercritical fluid chromatography with thermal energy analysis detection 445(1988)244

**External film diffusion**

Theoretical analysis of the asymmetry in chromatographic peaks 404(1987)1

**Extraction**

On-line post-column extraction in column LC with electron-capture detection. II. Applications 435(1988)297

**Extraction efficiency**

Increasing extraction efficiency in supercritical fluid extraction from complex matrices.

Predicting extraction efficiency of diuron and linuron in supercritical fluid extraction using supercritical fluid chromatographic retention 447(1988)53

**Extraction methods**

Qualitative GC and GC-MS screening for  $\beta$ -blockers in urine after solid-phase extraction using Extrelut-1 columns 426(1988)194

Simple extraction of atenolol from urine and its determination by HPLC 426(1988)234

Specific identification of synthetic organic chemicals in river water using liquid-liquid extraction and resin adsorption coupled with electron impact, chemical ionization and accurate mass measurement GC-MS analyses 436(1988)229

**Extraction techniques**

Comparative study of solid-phase and liquid-phase extraction techniques for isolation of phospholipids from plasma 419(1987)103

**Eye lens protein**

Electrophoretic titration curve in 6 M urea of the bovine eye lens protein  $\alpha$ -crystallin 411(1987)507

# F

## Factor analysis

Factor analysis of trends in selectivity in GLC of 175 saturated and unsaturated esters on 14 polysiloxane stationary phases 404(1987)11

## Factorial design

Optimization of HPLC analysis for isoxazolyl penicillins using factorial design 425(1988)331

## Factor X

Rapid ion-exchange chromatography for preparative separation of proteins. III. Application to purification of bovine factor X and prothrombin 403(1987)383

Separation of human factor X from factor Xa by reversed-phase HPLC 444(1988)107

## Factor Xa

Separation of human factor X from factor Xa by reversed-phase HPLC 444(1988)107

## Famotidine

HPLC determination of famotidine in urine 428(1988)395

## Far-UV absorbance detector

Capillary GC analysis with the far-UV absorbance detector 441(1988)63

## Fast Blue BB

Reaction of Fast Blue BB salt and *p*-dimethylaminocinnamaldehyde with indole-3-acetic acid degradation products. Evaluation of the coupling properties of the reagents used in the zymographic screening of indole-3-acetic acid oxidases 411(1987)476

## Fast Green FCF

Simple method for the analysis of food dyes on reversed-phase thin-layer plates 411(1987)437

## Fats

Letter symbols in the chemistry of glyceridic fats and oils 410(1987)509

## Fatty acid anilides

Determination of oxidation products of *N*-phenyllinoleamide: Spanish toxic oil syndrome studies 426(1988)83

## Fatty acid methyl esters

Analysis of fatty acid methyl esters with high accuracy and reliability. VI. Rapid analysis by split injection capillary GLC 407(1987)231

Preparation of fatty acid methyl esters by direct transesterification of lipids with aluminium chloride-methanol 441(1988)99

Effect of overload of capillary GLC columns on the equivalent chain lengths of C<sub>18</sub> unsaturated fatty acid methyl esters 447(1988)43

Equivalent chain-lengths of methyl ester derivatives of fatty acids on GC. A reappraisal 447(1988)305

## Fatty acids

Characterization of *Botrytis* by GLC 403(1987)292

Benzyl esters of C<sub>2</sub>-C<sub>20</sub> fatty acids and metabolically relevant carboxylic acids. Preparation and GC-MS 409(1987)43

Unusual and very long-chain fatty acids produced by Basidiomycetes 409(1987)390

Letter symbols in the chemistry of glyceridic fats and oils 410(1987)509

Identification by combined GC-MS of constituent long-chain fatty acids and alcohols from the meibomian glands of the rat and a comparison with human meibomian lipids 414(1987)253

Simultaneous quantification of total medium- and long-chain fatty acids in human milk by capillary GC with split injection 415(1987)1

Separation and quantitation of fatty acids, sterols and bile acids in feces by GC as the butyl ester-acetate derivatives 415(1987)13

HPLC analysis of serum long-chain fatty acids by direct derivatization method 416(1987)237

HPLC separation of femtomolar quantities of endogenous carboxylic acids, including arachidonic acid metabolites, as 4-bromomethyl-7-acetoxycoumarin derivatives 416(1987)247

Use of pentafluorobenzyl and pentafluoropropionyl-pentafluorobenzyl esters of bacterial fatty acids for GC analysis with electron-capture detection 417(1987)366

TLC of human platelet phospholipids with fatty acid analysis 420(1987)411

HPLC analysis of serum short-chain fatty acids by direct derivatization 421(1987)33



Fatty acid profiles of sebaceous triglycerides by capillary GC with mass-selective detection 425(1988)11

Thermospray-MS analysis of underivatized monohydroxy fatty acids: application to stimulated platelets 425(1988)25

Simultaneous determination of C<sub>2</sub>-C<sub>22</sub> non-esterified fatty acids and other metabolically relevant carboxylic acids in biological material by GC of their benzyl esters 425(1988)257

Analysis of tissue free fatty acids isolated by aminopropyl bonded-phase columns 428(1988)221

Identification of very-long-chain fatty acids in rat and mouse harderian gland lipids by capillary GC-MS 431(1988)231

Rapid method for the analysis of red blood cell fatty acids by reversed-phase HPLC 432(1988)29

Comparative chromatographic study of modifications of brush-border membrane vesicles induced by an essential fatty acid-deficient diet 432(1988)75

Simultaneous capillary GC profiling of medium- and long-chain fatty acid methyl esters with split injection. Correction for injection-related discrimination by the 'bracketing' method 434(1988)385

Profiling of plasma cholesterol ester and triglyceride fatty acids as their methyl esters by capillary GC, preceded by a rapid aminopropyl-silica column chromatographic separation of lipid classes 434(1988)395

Calibration of TLC with flame ionization detection for the analysis of natural lipid samples 435(1988)343

Automated solid-phase catalyzed pre-column derivatization of fatty acids for reversed-phase HPLC analysis with fluorescence detection 436(1988)437

Identification of isomeric hydroxy fatty acids in aerosol samples by capillary GC-MS 438(1988)309

Rapid separation of free fatty acids from neutral lipids in brain extracts 439(1988)453

Characterization of a lipid-rich fraction synthesized by *Streptomyces avermitilis* 441(1988)31

Direct, sensitive and selective detection of free fatty acids by HPLC with post-column ion-pair

extraction and absorbance detection 445(1988)189

GC analysis of different homologous series of acids esterified in aqueous solutions with butyl and propyl alcohols 446(1988)237

Method for evaluating the bioconversion of radioactive polyunsaturated fatty acids by use of reversed-phase LC 448(1988)249

#### **F<sub>ab</sub>' fragments**

Synthesis of high-capacity immunoaffinity sorbents with oriented immobilized immunoglobulins or their F<sub>ab</sub>' fragments for isolation of proteins 424(1988)243

#### **FD&C dyes**

Evaluation of disposable cartridges for trace enrichment from aqueous solutions 402(1987)201

#### **Felbinac**

Simultaneous determination of ofloxacin, fenbufen and felbinac in rat plasma by HPLC 431(1988)135

#### **Fenbufen**

Simultaneous determination of ofloxacin, fenbufen and felbinac in rat plasma by HPLC 431(1988)135

#### **Fenflumizole**

HPLC assay of fenflumizole and its demethyl metabolites in biological samples 413(1987)189

#### **Fenfluramine**

Enantioselective GC assay with electron-capture detection for *dl*-fenfluramine and *dl*-norfenfluramine in plasma 433(1988)105

#### **Fenoprofen**

Rapid HPLC assay for the simultaneous analysis of non-steroidal anti-inflammatory drugs in plasma 416(1987)293

HPLC quantitation of triacylglycerols containing fenoprofen from biological samples 422(1987)33

#### **Fenretinide**

HPLC analysis of fenretinide in rodent feed 408(1987)430

#### **Fentanyl**

Determination of fentanyl and alfentanil in plasma by HPLC with UV detection 419(1987)464

Simultaneous determination of fentanyl and

alfentanil in rat tissues by capillary column GC 433(1988)95

#### **Fentanyl citrate**

HPLC determination of fentanyl citrate in a parenteral dosage form 445(1988)299

#### **Ferbam**

LC of dibutylthiocarbamate degradation products 404(1987)359

#### **Ferritin**

Influence of pore and particle size on the frontal uptake of proteins. Implications for preparative anion-exchange chromatography 409(1987)111

#### **Ferritin receptor**

Chromatographic detergent exchange in the preparation of membrane receptor protein 427(1988)341

#### **Ferrocene**

Interfacing of inductively coupled plasma-atomic emission spectrometry to SFC for elemental detection 411(1987)213

#### **Ferrocene reagents**

Sensitive ferrocene reagents for derivatization of thiol compounds in HPLC with dual-electrode coulometric detection 419(1987)17

#### **N-(Ferrocenyl)maleimide**

Sensitive ferrocene reagents for derivatization of thiol compounds in HPLC with dual-electrode coulometric detection 419(1987)17

#### **Fertirelin acetate**

Reversed-phase HPLC of fertirelin acetate and related compounds 444(1988)177

#### **Ferulic acid**

Identification of phenols, phenolic acid dimers, and monosaccharides by GLC on a capillary column 436(1988)484

#### **$\alpha$ -Fetoprotein**

Microheterogeneity of rat  $\alpha$ -fetoprotein on a copper chelate column 417(1987)168

#### **Fetuin**

Determination of monosaccharides in glycoproteins by reversed-phase HPLC on 2.1-mm narrowbore columns 443(1988)299

#### **Fiber-reinforced organic-matrix composites**

High-performance SEC of fiber-reinforced organic-matrix composites 408(1987)129

#### **Fibrinogen**

Detection of fibrinogen heterogeneity by capillary isotachopheresis 419(1987)329

Steric protection of a modified glass surface with adsorbed polyacrylamide: interactions with blood proteins 428(1988)17

Isolation and comparative peptide mapping of fibrinogen subunits by reversed-phase HPLC 443(1988)309

#### **Fibronectin**

Biospecific affinity chromatography of plasma fibronectin on cross-linked gelatin microbeads 419(1987)308

Affinity purification of human plasma fibronectin on immobilized gelatin 432(1988)93

#### **Field-flow fractionation**

Near-wall field-flow fractionation. Application of surface forces and transversal voltage to fractionation in laminar flow 446(1988)131

Integral Doppler anemometry and analytical field-flow fractionation 446(1988)141

Acoustic field-flow fractionation 446(1988)151

#### **First derivative conductometric detector**

Use of first derivative conductometric detection in ion chromatography 436(1988)86

#### **Fischer-Tropsch synthesis**

Contributions of GC to the development of heterogeneous catalysis 406(1987)151

#### **Fission products**

Automated separation process for radioanalytical purposes at nuclear power plants 406(1987)359

#### **Flame ionization detector**

Quantitative analysis of triglycerides by HPLC using non-linear gradient elution and flame ionization detection 435(1988)139

Calibration of TLC with flame ionization detection for the analysis of natural lipid samples 435(1988)343

Flame ionization and flame thermionic detection of carbon and nitrogen in aquatic lipid and humic-type classes with an Iatroscan Mark IV 435(1988)350

TLC-flame ionization detection calibration with natural and synthetic standards 439(1988)404

#### **Flame photometric detector**

Spectra and response linearization of chlorine

and bromine compounds in a copper-sensitized flame photometric detector 408(1987)69

#### **Flame thermionic detector**

Flame ionization and flame thermionic detection of carbon and nitrogen in aquatic lipid and humic-type classes with an Iatroscan Mark IV 435(1988)350

#### **Flavanones**

Direct enantiomeric separation of racemic flavanones by HPLC using cellulose triacetate as a chiral stationary phase 441(1988)417

#### **Flavin adenine dinucleotide**

Quantification of riboflavin, riboflavin 5'-phosphate and flavin adenine dinucleotide in plasma and urine by HPLC 423(1987)105

Determination of flavin adenine dinucleotide in biological tissues by HPLC with electrochemical detection 442(1988)441

#### **Flavone acetic acid**

Optimisation of a reversed-phase HPLC method for the determination of flavone acetic acid and its major human metabolites in plasma and urine 431(1988)77

#### **Flavones**

Simultaneous isocratic HPLC determination of flavones and coumarins in *Matricaria chamomilla* extracts 404(1987)279

HPLC determination of polymethoxylated flavones in orange juice after solid-phase extraction 439(1988)481

#### **Flavonoids**

Spray reagent for the detection of coumarins and flavonoids on thin-layer plates 403(1987)285

HPLC determination of  $\beta$ -naphthoflavone in biological fluids 416(1987)160

Separations of flavonoids and alkaloids in medicinal herbs by high-speed counter-current chromatography 435(1988)159

Densitometric determination of quercetin heterosides in plant extracts (in French) 437(1988)260

HPLC of cottonseed flavonoids 441(1988)443

Rapid separation of flavonoids by analytical high-speed counter-current chromatography 445(1988)199

Separation of natural products by centrifugal partition chromatography 450(1988)91

#### **Flavonol glucosides**

Resolution of five position-specific flavonoid O-methyltransferases by fast protein liquid chromatofocusing 407(1987)291

#### **Flavour**

Flavor characterization of dentifrices using kinetic headspace GC 438(1988)273

Analysis of flavor absorption into plastic packaging materials using multiple headspace extraction GC 438(1988)281

#### **Flavour volatiles**

Dynamic headspace GC: concentration of volatile components after thermal desorption by intermediate cryofocusing in a cold trap. I. Principle and applications 405(1987)87

#### **Flecainide**

Determination of flecainide by GC-MS 416(1987)365

Rapid and inexpensive HPLC method for the quantification of flecainide in serum 427(1988)351

#### **Flecainide acetate**

HPLC procedure with fluorescence detection for the *m*-O-dealkylated lactam metabolite of flecainide acetate in human plasma 431(1988)438

#### **Floctafenic acid**

Simultaneous determination of floctafenine and its main metabolites, floctafenic and hydroxyfloctafenic acid, using high-performance liquid chromatography with gradient elution 432(1988)412

#### **Floctafenine**

Simultaneous determination of floctafenine and its main metabolites, floctafenic and hydroxyfloctafenic acid, using high-performance liquid chromatography with gradient elution 432(1988)412

#### **Flosequinan**

Simplified HPLC determination of flosequinan and its metabolite in plasma, serum and urine 425(1988)227

#### **Flow cell**

Optical errors in an LC absorbance cell 448(1988)193

#### **Flow injection analysis**

Fluorimetric determination of aflatoxins in

foodstuffs by HPLC with flow injection analysis 448(1988)173

#### **Flow-injection analysis**

Comparison of flow-injection analysis with HPLC for the determination of etoposide in plasma 432(1988)395

#### **Flowmeter**

Development of an inexpensive flowmeter for HPLC using a programmable pocket computer 407(1987)310

#### **Flow programming**

Flow programming system for process capillary GC 405(1987)163

#### **Flow-rate**

Flow-rate effect on elution volume in SEC 408(1987)9

Prediction of the distribution coefficient ( $K_d$ ) variation with operating conditions in chromatographic systems 411(1987)81

#### **Flow velocity profile**

Flow velocity profile in channels with modulated cross-sectional permeability for sedimentation-flotation focusing field-flow fractionation using step density gradients 404(1987)23

#### **Fluazifop-butyl**

GC-MS investigation of phenoxypropanoic acid derivatives possessing herbicidal activity 438(1988)55

#### **Flucloxacillin**

Optimization of HPLC analysis for isoxazolyl penicillins using factorial design 425(1988)331

#### **Flucythrinate**

TLC of flucythrinate residues 450(1988)452

#### **Flumequine**

Determination of flumequine and its 7-hydroxy metabolite by reversed-phase HPLC 414(1987)499

#### **Flunarizine**

Simple method for determination of flunarizine in serum by GC 421(1987)372

HPLC assay of flunarizine, (*E*)-1-[bis(4-fluorophenyl)methyl]-4-(3-phenyl-2-propenyl)-piperazine, in plasma of epileptic patients 426(1988)444

#### **Flunixin**

Identification of a flunixin metabolite in the horse by GC-MS 423(1987)123

GC analysis of flunixin in equine urine after extractive methylation 427(1988)55

#### **Flunoxaprofen**

Determination of the non-steroidal anti-inflammatory drug flunoxaprofen, *S*(+)-2-(4-fluorophenyl)- $\alpha$ -methyl-5-benzoxazoleacetic acid, in blood and urine by HPLC 413(1987)338

Determination of flunoxaprofen enantiomers in biological fluids by HPLC 415(1987)214

Assay method for product formation in *in vitro* enzyme kinetic studies of uridine diphosphate glucuronyltransferases: 2-arylpropionic acid enantiomers 430(1988)368

#### **Flunoxaprofen chloride**

*S*(+)-Flunoxaprofen chloride as chiral fluorescent reagent 427(1988)131

#### **Fluoranthene**

Chromatographic and cytogenetic analysis of *in vivo* metabolites of fluoranthene 448(1988)127

#### **9-Fluorenylmethyl chloroformate**

HPLC of biogenic amines, derivatized with 9-fluorenylmethyl chloroformate 437(1988)379

#### **Fluorescamine**

Separation of *N*-(4-aminobenzoyl)- $\gamma$ -oligo(L-glutamic acid)s on a thin-layer plate and their quantitation using fluorescamine 442(1988)420

#### **Fluorescein isothiocyanate**

Detection of amino acids on thin-layer plates 436(1988)341

#### **Fluoride**

Optimization of ion chromatography using commercially available detection systems and software 439(1988)13

#### **Fluoriprimidol**

Quantitation of triazole and pyrimidine plant growth retardants 438(1988)393

#### **5-Fluoro-2'-deoxyuridine**

HPLC analysis of fluoropyrimidine nucleosides and fluorouracil in plasma 414(1987)197

Determination of 5-fluoro-2'-deoxyuridine in human plasma by HPLC with pre-column fluorimetric derivatization 416(1987)176

4-Bromomethyl-6,7-dimethoxycoumarin as a fluorescence reagent for precolumn derivatization of 5-fluorouracil compounds in HPLC 430(1988)156

#### 1-Fluoro-2,4-dinitrobenzene

Study of the derivatization of *n*-alkylamines with 1-fluoro-2,4-dinitrobenzene in the presence of aqueous cetyltrimethylammonium bromide micelles 426(1988)267

#### 1-Fluoro-2,4-dinitrophenyl-5-L-alanine amide

Application of Marfey's reagent in racemization studies of amino acids and peptides 444(1988)115

#### Fluorometholone

GC and mass spectral study of synthetic corticosteroid metabolism: fluorometholone 426(1988)399

#### 4-Fluoro-7-nitrobenzo-2-oxa-1,3-diazole

Automatic amino acid analysis utilizing 4-fluoro-7-nitrobenzo-2-oxa-1,3-diazole 420(1987)141

#### Fluoropyrimidine nucleosides

HPLC analysis of fluoropyrimidine nucleosides and fluorouracil in plasma 414(1987)197

#### 5-Fluorouracil

HPLC analysis of fluoropyrimidine nucleosides and fluorouracil in plasma 414(1987)197

Determination of 5-fluorouracil in plasma and whole blood by ion-pair HPLC 426(1988)121

4-Bromomethyl-6,7-dimethoxycoumarin as a fluorescence reagent for precolumn derivatization of 5-fluorouracil compounds in HPLC 430(1988)156

#### 5-Fluorouridine

HPLC analysis of fluoropyrimidine nucleosides and fluorouracil in plasma 414(1987)197

#### Fluphenazine

New HPLC method for fluphenazine and metabolites in human plasma 414(1987)504

#### Flurazepam

Simultaneous determination of flurazepam and five metabolites in serum by HPLC and its application to pharmacokinetic studies in rats 423(1987)251

#### Flurbiprofen

Resolution of  $\alpha$ -arylacetic acid enantiomers on two chiral stationary phases 404(1987)123

Application of an ovomucoid-conjugated column for the optical resolution of some pharmaceutically important compounds 408(1987)316

Assay method for product formation in *in vitro* enzyme kinetic studies of uridine diphosphate glucuronyltransferases: 2-arylpropionic acid enantiomers 430(1988)368

#### Foam counter-current chromatography

Foam counter-current chromatography with the cross-axis synchronous flow-through coil planet centrifuge 403(1987)77

#### Fog samples

Determination of mono- and divalent cations and anions in small fog samples by ion chromatography 439(1988)121

#### Folates

Separation of N-(4-aminobenzoyl)- $\gamma$ -oligo(L-glutamic acid)s on a thin-layer plate and their quantitation using fluorescamine 442(1988)420

Determination of native folates in milk and other dairy products by HPLC 449(1988)271

#### Folding pathway

Survey of the folding pathway of a two-domain protein phosphoglycerate kinase 440(1988)421

#### Folic acid

Determination of water-soluble vitamins using HPLC and electrochemical or absorbance detection 447(1988)256

#### Fomannoxin

HPLC determination of fomannoxin in European strains of *Heterobasidion annosum* (Fr.) Bref. 440(1988)65

#### Food dyes

Simple method for the analysis of food dyes on reversed-phase thin-layer plates 411(1987)437

#### Forced-flow planar chromatography

Applicability of forced-flow planar chromatographic methods for the separation of enantiomers on Chiralplate 450(1988)241

#### Formamide

GC method for the determination of N-acetyl-S-(N-methylcarbamoyl)cysteine, a metabolite of N,N-dimethylformamide and N-methylformamide, in human urine 431(1988)361

**Formate**

HPLC determination of formate as benzimidazole in biological samples 426(1988)15

**Formic acid**

Determination of ionic metabolites from ethylene glycol in human blood by isotachopheresis 416(1987)111

Studies on sample preconcentration in ion chromatography. VIII. Preconcentration of carboxylic acids prior to ion-exclusion separation 447(1988)155

Extractive pentafluorobenzoylation of formic, acetic, levulinic, benzoic and phthalic acids, studied by LC and dual-oven capillary GC 447(1988)329

**Formylaminoantipyrine**

Investigation of an alkylamine modified and pH-controlled mobile phase for separation of pyrazolone derivatives by HPLC 409(1987)81

Simple HPLC method for simultaneous determination of aminopyrine and its metabolites 421(1987)198

**3-Formyl-25-desacetyl rifamycin**

Determination of rifampicin and its main metabolites in human plasma by HPLC 426(1988)412

**N'-Formylkynurenine**

Semi-micro quantitative assay for tryptophan oxygenase by HPLC 419(1987)296

**3-Formylrifamycin**

Determination of rifampicin and its main metabolites in human plasma by HPLC 426(1988)412

**Forphenicine**

HPLC determination of forphenicine in mouse serum and muscle by pre-column fluorescence derivatization using 1,2-diamino-4,5-ethylenedioxybenzene as fluorogenic reagent 430(1988)361

**Fossil fuels**

High-efficiency separation and characterization of high-molecular-weight heterocyclic sulfur compounds in fossil fuels by microcolumn LC and related techniques 406(1987)375

**Fourier-transform IR spectrometry**

Super- and subcritical fluid chromatography-Fourier transform IR

spectrometry of medium polar compounds on polymeric styrene-divinylbenzene packed columns 407(1987)87

Microcolumn HPLC-TLC-Fourier transform IR spectrometry 438(1988)329

**Fraction collector**

Fraction collector for capillary zone electrophoresis 438(1988)23

**Fragrance volatiles**

Dynamic headspace GC: concentration of volatile components after thermal desorption by intermediate cryofocusing in a cold trap. I. Principle and applications 405(1987)87

**Free-flow electrophoresis**

Contributions to a mathematical theory of free-flow electrophoresis 438(1988)145

**Free-flow electrophoretic apparatus**

Free-flow electrophoretic apparatus for separation and concentration of proteins 432(1988)137

**Free energy consumption**

Free energy consumption at self-sharpening concentration fronts in isothermal fixed-bed adsorption. Theory for isotherms of arbitrary shape and applications in displacement chromatography 409(1987)1

**Frontal chromatography**

Tandem separation schemes for preparative HPLC of proteins 443(1988)31

**Fructose**

Prediction of the distribution coefficient ( $K_d$ ) variation with operating conditions in chromatographic systems 411(1987)81

**Fructose-6-phosphate**

Ion chromatographic determination of sugar phosphates in physiological samples 439(1988)83

**Fructotransferases II**

Activity staining of inulin fructotransferase II on polyacrylamide gels 445(1988)282

**Fruzemide**

LC screening of diuretics in urine 415(1987)347

**Frying oils**

Quantitative analyses of polar components in frying oils by the Iatroscan TLC-flame ionization detection technique 405(1987)371

**Fucose**

Analysis of neutral sugars as dinitrophenylhydrazones by HPLC 405(1987)221

**Fucoanthin**

HPLC of carotenoids from some marine shellfish 440(1988)37

**Fucoanthinol**

HPLC of carotenoids from some marine shellfish 440(1988)37

**Fulvic acid**

Computer-controlled linear pH gradient for HPLC fractionations of aromatic carboxylic acids and of humic and fulvic acids 439(1988)213

**Fulvic acids**

High-pressure SEC of chlorinated and unchlorinated aquatic fulvic acids using a polyacrylamide gel 450(1988)433

**Fumarate**

New method of purification of perchloric acid extracts for quantitative GC of some Krebs cycle acids 424(1988)357

**Fungal secondary metabolites**

Standardized HPLC of 182 mycotoxins and other fungal metabolites based on alkylphenone retention indices and UV-VIS spectra (diode array detection) 404(1987)195

**Furosemide**

Determination of association constants of  $\beta$ -cyclodextrin and  $\beta$ -cyclodextrin-bearing polymers with drugs by HPLC 409(1987)61

**Furprofen**

Determination of furprofen in human plasma by solid-phase extraction and reversed-phase HPLC 424(1988)201

**Fusarenon-X**

HPLC method for determining trichothecene mycotoxins by post-column fluorescence derivatization 410(1987)427

**Fusarin C**

Standardized HPLC of 182 mycotoxins and other fungal metabolites based on alkylphenone retention indices and UV-VIS spectra (diode array detection) 404(1987)195

**Fused-silica ends**

Fused-silica ends for glass capillaries: widening glass capillary butts by a heated tungsten point for preparing press-fit connections 435(1988)204

**Fusel oil**

Rapid HPLC method for determination of ethanol and fusel oil in the alcoholic beverage industry 447(1988)443

**Fusicoccin**

Immunoaffinity chromatography of fusicoccin 440(1988)47

**Fusidic acid**

LC determination of fusidic acid in serum 430(1988)400

HPLC determination of fusidic acid in plasma 433(1988)159

***Fusobacterium***

Identification by GC-MS of short-chain hydroxy acids produced by *Fusobacterium* species and *Clostridium innocuum* 420(1987)121

**G****Galbiosylceramide**

TLC of neutral glycosphingolipids: an improved simple method for the resolution of GlcCer, GalCer, LacCer and Ga<sub>2</sub>Cer 426(1988)188

**Galactose**

Analysis of neutral sugars as dinitrophenylhydrazones by HPLC 405(1987)221

Simultaneous GLC determination of aldoses and alduronic acids 408(1987)245

Requirements of mutually miscible liquids for ternary chromatographic solvents with reference to monosaccharide PC 436(1988)53

**Galactose-binding lectin**

Purification of human erythrocytes specific lectins from rice bean, *Phaseolus calcaratus* syn. *Vigna umbellata*, by HPLC 431(1988)37

 **$\beta$ -Galactosidase fusion proteins**

Isolation of recombinant mycobacterial antigens by an automatic and generally applicable purification method for  $\beta$ -galactosidase fusion proteins 448(1988)165

**$\alpha$ -D-Galactosidases**

Electrophoretic study of  $\alpha$ -D-galactosidases from seeds of *Glycine soja* and *Vigna radiata* possessing erythroagglutinating activity 436(1988)59

**Galactosylceramide**

TLC of neutral glycosphingolipids: an improved simple method for the resolution of GlcCer, GalCer, LacCer and Ga<sub>2</sub>Cer 426(1988)188

**Galacturonic acids**

Simultaneous GLC determination of aldoses and alduronic acids 408(1987)245

**Galanthamine**

Reversed-phase LC for the determination of galanthamine and its metabolites in human plasma and urine 421(1987)396

**Galanthaminone**

Reversed-phase LC for the determination of galanthamine and its metabolites in human plasma and urine 421(1987)396

**Gallamine**

Determination of the neuromuscular blocking drug gallamine in rat serum using HPLC 417(1987)428

**Gallopamil**

Quantification of gallopamil and norgallopamil in human plasma by HPLC with fluorescence detection 414(1987)492

Rapid HPLC method for the measurement of gallopamil in plasma 421(1987)165

**Gangliosides**

New solvent system for HPTLC and HPLC of gangliosides 405(1987)125

HPLC of underivatized gangliosides 408(1987)285

Improved method for large-scale purification of brain gangliosides by Q-Sepharose CC. Immunochemical detection of C-series polysialogangliosides in adult bovine brains 445(1988)377

**Gas-Chrom**

Characterization of porous polymers by polar strength and selectivity 404(1987)145

**Gas chromatograph**

Simple instrument for teaching the basics of chromatography in schools 446(1988)173

**Gaseous deposit**

Efficiency of adsorption-desorption of volatile compounds. Application to isoprene on Tenax GC 437(1988)139

**Gaseous mixtures**

Preparation of standard gaseous mixtures by a manometric method using the Benedict-Webb-Rubin equation 436(1988)323

**Gasoline**

GC determination for forensic purposes of petroleum fuel inhaled just before fatal burning 424(1988)49

**Gas-phase luminescence**

GC determination of benzaldehyde by gas-phase luminescence 409(1987)243

**Gas phase protein sequencer**

High-sensitivity phenylthiohydantoin amino acid analysis on-line to a gas phase protein sequencer 408(1987)388

**Gastric mucosa protein**

Isolation and characterization of a protein detected in inflamed human gastric mucosa 419(1987)323

**Gastric proteases**

Isolation and characterization of a protein detected in inflamed human gastric mucosa 419(1987)323

**Gen-Pak FAX**

High-resolution chromatography of nucleic acids on the Gen-Pak FAX column 444(1988)47

**Gentamicin**

HPLC assay of the major components of gentamicin in serum 419(1987)408

**Gentamicin C**

Isolation of gentamicin C compounds from culture filtrates of *Micromonospora purpurea* 450(1988)115

**Gentisamide**

HPLC method for the quantitation of salicylamide and its metabolites in biological fluids 420(1987)313

**Gentisamide 5-glucuronide**

HPLC method for the quantitation of salicylamide and its metabolites in biological fluids 420(1987)313



**Gentisamide sulphates**

HPLC method for the quantitation of salicylamide and its metabolites in biological fluids 420(1987)313

**Gentisic acid**

Rapid HPLC assay for the simultaneous analysis of non-steroidal anti-inflammatory drugs in plasma 416(1987)293

**Geometrical exclusion**

Structure parameters of molecules and media evaluated by chromatographic partition. II. Geometrical exclusion in gels 410(1987)233

**Geosmin**

Simplified method to quantify geosmin and 2-methylisoborneol concentrations in water and microbiological cultures 409(1987)337

**Gepirone**

Simultaneous determination of buspirone, gepirone, ipsapirone and their common metabolite 1-(2-pyrimidinyl)piperazine in rat plasma and brain by HPLC 431(1988)477

**Geraniol**

Characterisation of volatile oil constituents with relatively long GC retention times on two stationary phases 449(1988)312

**Gibberellins**

Separation of gibberellins by TLC and GC, and possibilities for their quantitative analysis 437(1988)287

**Glafenic acid**

Effective HPLC determination of glafenine in plasma: pharmacokinetic application 427(1988)307

**Glafenine**

Effective HPLC determination of glafenine in plasma: pharmacokinetic application 427(1988)307

**Glass capillary butts**

Fused-silica ends for glass capillaries: widening glass capillary butts by a heated tungsten point for preparing press-fit connections 435(1988)204

**Glasses**

Isotachophoretic determination of the cationic constituents in mixtures of raw materials for glass manufacturing 439(1988)423

 **$\alpha$ -Gliadin**

Isolation and analysis of peptidic fragments of

$\alpha$ -gliadin using reversed-phase HPLC 434(1988)429

**Gliclazide**

HPLC determination of gliclazide in human plasma 421(1987)223

**Glipizide**

High-performance liquid chromatographic determination of glipizide in human plasma and urine 421(1987)319

**Globin polypeptides**

Reversed-phase HPLC separation and quantitation of reticulocyte  $\alpha$ - and  $\beta$ -globin polypeptide chains from normal and  $\beta$ -thalassemic mice 431(1988)177

**Globotriaosylceramide**

TLC of neutral glycosphingolipids: an improved simple method for the resolution of GlcCer, GalCer, LacCer and Ga<sub>2</sub>Cer 426(1988)188

**Globulins**

Application of HPLC to the assessment of subunit heterogeneity in plant 11S storage globulins 402(1987)159

High-performance adsorption chromatography of proteins on 2- $\mu$ m spherical beads of hydroxyapatite 440(1988)305

**Glucocorticoid receptor**

Improved Stokes radius measurement of the glucocorticoid receptor using TSK G4000SW and TSK G3000SW high-performance size-exclusion columns. Analytical and preparative applications 403(1987)183

**Glucocorticoid receptor proteins**

Non-histone chromatin proteins that recognize specific sequences of DNA (Review) 418(1987)73

**Glucocorticoids**

Simultaneous quantification of natural glucocorticoids and progestins in serum 421(1987)123

**Glucomannan**

Macroporous glucomannan spheres for the size-exclusion separation of aqueous macromolecules 409(1987)175

**Gluconate**

Origin of system peaks in single-column ion chromatography of inorganic anions using high pH borate-gluconate buffers and conductivity detection 404(1987)67

Carbohydrate–borate eluents for anion chromatography 448(1988)355

#### **Gluco-oligomers**

HPLC of gluco-oligomers on C<sub>18</sub>-bonded vinyl alcohol copolymer gel with alkaline eluents 436(1988)328

#### **1-(β-D-Glucopyranosyl)amobarbital**

HPLC determination of the diastereomers of 1-(β-D-glucopyranosyl)amobarbital in urine 422(1987)309

#### **1-O-α-D-Glucopyranosyl-D-fructose**

Preparative HPLC purification and structural determination of 1-O-α-D-glucopyranosyl-D-fructose (trehalulose) 402(1987)265

#### **1-(β-D-Glucopyranosyl)phenobarbital**

Quantitation of 1-(β-D-glucopyranosyl)-phenobarbital in plasma and bile 419(1987)421

#### **Glucosamine**

HPLC separation of some mono- and disaccharides with detection by a post-column enzyme reactor and a chemically modified electrode 408(1987)157

#### **Glucose**

Analysis of neutral sugars as dinitrophenyl-hydrazones by HPLC 405(1987)221

HPLC separation of some mono- and disaccharides with detection by a post-column enzyme reactor and a chemically modified electrode 408(1987)157

Simultaneous GLC determination of aldoses and alduronic acids 408(1987)245

Prediction of the distribution coefficient ( $K_d$ ) variation with operating conditions in chromatographic systems 411(1987)81

Measurement of glycated albumin in diabetic patients by biospecific affinity chromatography 419(1987)75

Chromatography, flow injection analysis and electrophoresis in computer-assisted comparative biochemistry: its application and possibilities in clinical research. Preliminary studies on Crohn's disease 440(1988)261

Effect of the content of divinylbenzene in ion-exchange resins on the chromatographic separation of α-cyclodextrin and glucose 448(1988)241

Carbohydrate–borate eluents for anion chromatography 448(1988)355

#### **Glucose esters**

Analysis of naturally occurring hydroxybenzoic acid glucosides and hydroxybenzoyl glucoses and their behaviour on a polyamide column 441(1988)335

#### **Glucose-6-phosphate**

Ion chromatographic determination of sugar phosphates in physiological samples 439(1988)83

#### **Glucose phosphate isomerase**

New strategies for the screening of a large number of immobilized dyes for the purification of enzymes. Application to the purification of enzymes from human haemolysate 449(1988)403

#### **α-1,2-Glucosidase**

Subfractionation of the dansylated derivatives of glucosyl galactosyl hydroxylysine by LC and its application to a specific α-1,2-glucosidase assay 416(1987)37

#### **Glucosides**

Analysis of naturally occurring hydroxybenzoic acid glucosides and hydroxybenzoyl glucoses and their behaviour on a polyamide column 441(1988)335

Comparative HPLC and GLC determination of phenolic glucosides in Salicaceae species 442(1988)175

HPLC determination of glucosides (glucose conjugates) with post-column reaction detection combining immobilized enzyme reactors and luminol chemiluminescence 449(1988)217

#### **Glucosinolates**

Separation and determination of intact glucosinolates in rapeseed by HPLC 435(1988)501

#### **Glucosylceramide**

TLC of neutral glycosphingolipids: an improved simple method for the resolution of GlcCer, GalCer, LacCer and Ga<sub>2</sub>Cer 426(1988)188

#### **Glucuronic acid**

Simultaneous GLC determination of aldoses and alduronic acids 408(1987)245

#### **Glucuronic acid-1-phosphate**

Ion chromatographic determination of sugar phosphates in physiological samples 439(1988)83

#### **Glutamate**

Glutamate and glutamine in cerebrospinal fluid and serum from healthy volunteers – analytical aspects 424(1988)378

HPLC analysis of neurotransmitter amino acids in brain 428(1988)9

#### **Glutamic acid**

Determination of 4-aminobutyric acid, aspartate, glutamate and glutamine and their <sup>13</sup>C stable-isotopic enrichment in brain tissue by GC-MS 414(1987)265

#### **Glutamic acid analogues**

Resolution of  $\alpha$ -substituted amino acid enantiomers by HPLC after derivatization with a chiral adduct of *o*-phthalaldehyde. Application to glutamic acid analogues 440(1988)209

#### **Glutamine**

Determination of 4-aminobutyric acid, aspartate, glutamate and glutamine and their <sup>13</sup>C stable-isotopic enrichment in brain tissue by GC-MS 414(1987)265

Glutamate and glutamine in cerebrospinal fluid and serum from healthy volunteers – analytical aspects 424(1988)378

#### **$\gamma$ -L-Glutamylcyclotransferase**

Assay of  $\gamma$ -L-glutamylcyclotransferase activity in rat brain synaptosomes by HPLC 426(1988)381

#### **$\gamma$ -Glutamylcysteine**

Method for determination of free and total glutathione and  $\gamma$ -glutamylcysteine concentrations in human leukocytes and plasma 420(1987)152

#### **$\epsilon$ -( $\gamma$ -Glutamyl)lysine**

HPLC method for the determination of  $\epsilon$ -( $\gamma$ -glutamyl)lysine and mono- and bis- $\gamma$ -glutamyl derivatives of putrescine and spermidine 443(1988)329

#### **$\gamma$ -Glutamylpolyamines**

HPLC method for the determination of  $\epsilon$ -( $\gamma$ -glutamyl)lysine and mono- and bis- $\gamma$ -glutamyl derivatives of putrescine and spermidine 443(1988)329

#### **$\gamma$ -Glutamyltransferase**

Automated separation and measurement of urinary isoenzymes and protein by ion-exchange LC 414(1987)47

#### **Glutaryl carnitine**

Analytical method for urinary glutaryl carnitine, acetylcarnitine and propionyl carnitine with a carboxylic acid analyser and a reversed-phase column 423(1987)297

#### **Glutathione**

HPLC fluorometry for the determination of thiols in biological samples using N-[4-(6-dimethylamino-2-benzofuranyl)phenyl]maleimide 414(1987)11

Sensitive ferrocene reagents for derivatization of thiol compounds in HPLC with dual-electrode coulometric detection 419(1987)17

Method for determination of free and total glutathione and  $\gamma$ -glutamylcysteine concentrations in human leukocytes and plasma 420(1987)152

Determination of natural thiols by LC after derivatization with 3,5-di-*tert*-butyl-1,2-benzoquinone 420(1987)404

Measurement of plasma thiols after derivatization with monobromobimane 424(1988)141

LC determination of diastereomeric glutathione conjugates and further derivatives of  $\alpha$ -bromoisovalerylurea in rat bile and urine by electrochemically generated bromine 427(1988)67

Determination of cysteine, glutathione and N-acetylcysteine in plasma by ion-pair reversed-phase LC and post-column derivatization 432(1988)65

#### **Glutathione S-transferases**

Purification of glutathione S-transferases from rat liver and Walker 256 mammary carcinoma cells by HPLC and a glutathione affinity column 434(1988)83

#### **Glycated albumin**

Measurement of glycated albumin in diabetic patients by biospecific affinity chromatography 419(1987)75

#### **Glycated haemoglobins**

Glycated haemoglobins (Review) 429(1988)279

#### **Glyceric fats**

Letter symbols in the chemistry of glyceric fats and oils 410(1987)509

#### **Glycerol**

Determination of glycerol in pig plasma by capillary GC 410(1987)480

#### **Glycerol dinitrates**

Determination of the two dinitrate metabolites of nitroglycerin in human plasma by capillary GC with electron-capture detection 413(1987)101

**Glycerol kinase**

Improved methods for purification and assay of glycerol kinase from *Escherichia coli* 428(1988)345

**Glycerol mononitrates**

GC assay of glycerol mononitrates in biological samples 413(1987)91

**Glycerophospholipids**

Quantification of plasmalogen, alkylacyl and diacyl glycerophospholipids by micro-TLC 436(1988)467

**Glycidoxypropyl functionalized silica plates**

TLC of metal complexes on glycidoxypropyl functionalized silica plates 409(1987)377

**Glycidyl methacrylate-ethylene dimethacrylate**

Hydrolysed macroporous glycidyl methacrylate-ethylene dimethacrylate copolymer sorbent for size-exclusion HPLC of synthetic polymers and biopolymers 435(1988)357

**Glycine**

HPLC analysis of neurotransmitter amino acids in brain 428(1988)9

**Glycinin**

Scale-up from analytical to preparative reversed-phase HPLC of soybean glycinin 444(1988)263

**Glycocalicin**

Human platelet glycocalicin purification by phenyl boronate affinity chromatography coupled to anion-exchange HPLC 426(1988)376

**Glycolic acid**

Determination of ionic metabolites from ethylene glycol in human blood by isotachopheresis 416(1987)111

**Glycolipids**

Separation of acidic and neutral glycolipids by diphasic TLC 436(1988)515

Qualitative and quantitative TLC analysis of polar lipids from leaves of glanded and glandless cotton genotypes 437(1988)211

Identification of partially methylated methyl glycosides by GC-MS of trimethylsilyl derivatives. Application to mycobacterial glycolipid antigen analysis 445(1988)87

**Glycols**

Quantification of some glycols in urinary organic acid profiles of acutely ill newborns 416(1987)439

**Glycopeptides**

Subfractionation of the dansylated derivatives of glucosyl galactosyl hydroxylysine by LC and its application to a specific  $\alpha$ -1,2-glucosidase assay 416(1987)37

Analytical high-performance affinity chromatography of ovalbumin-derived glycopeptides on columns of concanavalin A- and wheat germ agglutinin-immobilized gels 438(1988)73

**Glycoproteins**

Rapid chromatographic assay procedure for peptide-N-glycosidase activity 408(1987)378

Purification and characterization of granulocyte-macrophage colony stimulating factor from human placenta 413(1987)242

Microheterogeneity of rat  $\alpha$ -fetoprotein on a copper chelate column 417(1987)168

Purification of human urinary erythropoietin 417(1987)178

Chromatography of complex protein mixtures (Review) 418(1987)115

Biospecific affinity chromatography of plasma fibronectin on cross-linked gelatin microbeads 419(1987)308

Sulphate quantification in rabbit tracheobronchial mucus by reversed-phase ion-pair chromatography 424(1988)147

Isolation of the major O-glycosidically linked oligosaccharides obtained by alkaline borohydride degradation of human meconium glycoproteins 425(1988)35

High-performance GPC of the surface glycoprotein from bovine leukemia virus 425(1988)379

Medium-scale ligand-affinity purification of two soluble forms of human interleukin-2 receptor 431(1988)55

Analysis of tissue glycoprotein sugar chains by two-dimensional HPLC mapping 434(1988)51

Structural characterization of glycoproteins 443(1988)267

Determination of monosaccharides in glycoproteins by reversed-phase HPLC on 2.1-mm narrowbore columns 443(1988)299

High-performance concanavalin A affinity chromatography of liver and hepatoma membrane proteins 444(1988)29

#### **Glycosaminoglycans**

High-performance GPC of glycosaminoglycans. Column calibration by gel electrophoresis 402(1987)149

Rapid and sensitive method for measurement of hyaluronic acid and isomeric chondroitin sulfates using HPLC 413(1987)9

Rapid determination of disaccharides from chondroitin and dermatan sulphates by HPLC 414(1987)156

Extracellular mammalian polysaccharides: glycosaminoglycans and proteoglycans (Review) 418(1987)187

Fractionation of aorta glycosaminoglycans by HPLC 419(1987)275

Quantitative GC measurement of glycosaminoglycan hexosamines in urine and plasma 420(1987)231

Characterization of the products generated from oversulphated dermatan sulphate isomers with chondroitinase-B by HPLC 423(1987)51

Methods for analysis of urinary glycosaminoglycans (Review) 429(1988)293

#### **Glycosidase**

Assay of glycosidase by lectin affinity HPLC 407(1987)299

#### **Glycosides**

Determination of monosaccharides constituting the glycosides in saponins by HPLC 403(1987)319

HPLC of glycosides on a new type of hydroxyapatite column 407(1987)205

#### **Glycosphingolipids**

*In situ* visualization of glycosphingolipids on thin-layer chromatograms using lectin-peroxidase conjugates 411(1987)393

TLC of neutral glycosphingolipids: an improved simple method for the resolution of GlcCer, GalCer, LacCer and Ga<sub>2</sub>Cer 426(1988)188

#### **Glycosyl galactosyl hydroxylysine**

Subfractionation of the dansylated derivatives of glucosyl galactosyl hydroxylysine by LC and its application to a specific  $\alpha$ -1,2-glucosidase assay 416(1987)37

#### **N-Glycyl-2,6-xylidine**

*In vivo* metabolism of lidocaine in the rat. Isolation of urinary metabolites as pentafluorobenzoyl derivatives and their identification by combined GC-MS 421(1987)267

#### **Glyoxylic acid**

Determination of ionic metabolites from ethylene glycol in human blood by isotachopheresis 416(1987)111

HPLC determination of glyoxylic acid and other carbonyl compounds in urine 417(1987)253

HPLC determination of glyoxylic acid in urine 432(1988)37

HPLC of mixtures of phenol or guaiacol and their reaction products with glyoxylic acid 437(1988)185

#### **GM<sub>1</sub> ganglioside $\beta$ -galactosidase**

Microassay for GM<sub>1</sub> ganglioside  $\beta$ -galactosidase activity using HPLC 426(1988)75

#### **Gold**

Ion-interaction reversed-phase chromatographic method for the determination of gold(I) cyanide in mine process liquors using automated sample preconcentration 439(1988)23

#### **Gossypol**

Determination of gossypol enantiomers in plasma after administration of racemate using HPLC with precolumn chemical derivatisation 433(1988)141

#### **Gradient elution**

Single and multiple gradients in supercritical fluid chromatography (Review) 402(1987)1

Simplified description of HPLC separation under overload conditions, based on the Craig distribution model. V. Gradient elution separation 405(1987)51

Stepwise gradient elution using switching valves in micro HPLC 405(1987)117

Precolumn concentration and/or stepwise-gradient elution by switching valve techniques in micro HPLC 407(1987)141

Gradient elution in ion chromatography 411(1987)107

Quantitative analysis of triglycerides by HPLC using non-linear gradient elution and flame ionization detection 435(1988)139

HPLC of amino acids, peptides and proteins. LXXXV. Evaluation of the use of hydrophobicity coefficients for the prediction of peptide elution profiles 438(1988)197

Conformational studies of bovine alkaline phosphatase in hydrophobic interaction and size-exclusion chromatography with linear diode array and low-angle laser light scattering detection 442(1988)29

Preparative separation of peptide and protein samples by HPLC with gradient elution. I. The Craig model as a basis for computer simulations 444(1988)303

Stepwise gradient development in TLC. IV. Miniaturized generators for continuous and stepwise gradients 446(1988)275

HPLC on dynamically modified silica. VIII. Gradient elution using eluents containing cetyltrimethylammonium bromide 447(1988)182

#### **Gradient steepness**

Improvement in the resolution of *o*-phthalaldehyde derivatized amino acids by applying gradient steepness optimization to five reversed-phase columns of different lengths and particle sizes 411(1987)121

#### **Gradient time**

HPLC of amino acids, peptides and proteins. LXXXV. Evaluation of the use of hydrophobicity coefficients for the prediction of peptide elution profiles 438(1988)197

#### **Gramicidin S-1**

Adsorption of cyclic peptides analogous to gramicidin S and gratisin onto octadecylsilica stationary phase and bacterial cells 413(1987)287

#### **Gratisin**

Adsorption of cyclic peptides analogous to gramicidin S and gratisin onto octadecylsilica stationary phase and bacterial cells 413(1987)287

#### **Group VA triphenyl derivatives**

Holistic molecular conformation and total surface area calculations as predictors for solution properties and chromatographic parameters 403(1987)1

#### **Growth factors**

Method for purification of an insulin-like growth factor-binding protein produced by human HEP G2 hepatoma cells 420(1987)163

Application of preparative reversed-phase HPLC

to the isolation of insulin-like growth factor II from human serum 421(1987)61

Reduction studies on bacterial recombinant somatomedin C/insulin-like growth factor-1 443(1988)183

#### **Growth hormone**

Recovery of polypeptides after reversed-phase HPLC 408(1987)191

Fluorimetric determination of rat growth hormone in pituitary cell culture by HPLC with post-column derivatization 417(1987)409

Separation of protein hormones (Review) 418(1987)245

HPLC of amino acids, peptides and proteins. LXXXIV. Application of derivative spectroscopy to the study of column residency effects in the reversed-phase and size-exclusion LC separation of proteins 435(1988)271

High-performance SEC determination of the potency of biosynthetic human growth hormone products 435(1988)307

#### **Guaiacol**

HPLC of mixtures of phenol or guaiacol and their reaction products with glyoxylic acid 437(1988)185

#### **Guanadrel**

GC determination of guanadrel in plasma and urine 434(1988)135

#### **Guanidino compounds**

Chromatography and electrophoresis of creatinine and other guanidino compounds (Review) 429(1988)235

#### **1-Guanidino-2-(4-imidazole)propionic acid**

Identification of 1-guanidino-2-(4-imidazole)propionic acid in human urine by HPLC and field desorption MS 425(1988)373

#### **Guanine derivative**

Identification by HPLC and MS of a guanine derivative arising from acid hydrolysis of DNA 407(1987)349

#### **Guanines**

Separation characteristics of alkylated guanines in HPLC 447(1988)81

#### **Gum opium**

Separation of eight alkaloids and meconic acid and quantitation of five principal alkaloids in gum opium by gradient reversed-phase HPLC 436(1988)455

**Gums**

Identification of carbohydrate gums in dental cream by TLC 407(1987)369

**H****Haem-bearing peptides**

Isolation of haem-bearing peptides of haem proteins by the use of adsorption chromatography 410(1987)463

**Haemodialysers**

Comparison of serum protein removal in haemodialysis therapy by partly protein-permeable haemodialysers 428(1988)25

**Haemoglobin**

On the increased retention volume of human haemoglobin in high-performance gel filtration 408(1987)303

Isolation and partial characterization of pyridoxal 5'-phosphate hemoglobins by HPLC as a quality-control method for hemoglobin-based blood substitutes 413(1987)53

ATP-hemoglobin: anomalous oxygen binding properties 415(1987)136

Haemoglobin pyridoxylation: optimization and observation of an additional new pyridoxal phosphate-haemoglobin species by HPLC resolution 417(1987)397

Chromatography of complex protein mixtures (Review) 418(1987)115

Separation of hemoglobins and hemoglobin chains by HPLC (Review) 418(1987)277

Hemoglobin-based blood substitutes: characterization of five pyridoxal 5'-phosphate derivatives of hemoglobin 419(1987)37

Separation of asymmetrical hybrid hemoglobins by hydrophobic interaction chromatography 419(1987)303

GC studies of the carbamylation of haemoglobin by methyl isocyanate in rats and rabbits 426(1988)239

Separation of human haemoglobin alkylated at

$\beta$ 93 cysteine from its native form by fast protein LC 427(1988)335

Effects of differences in charge and hydrophobicity of surface amino acids of hemoglobins on high-performance GPC 428(1988)247

Glycated haemoglobins (Review) 429(1988)279

HPLC separation of human haemoglobins. Simultaneous quantitation of foetal and glycated haemoglobins 434(1988)95

Non-linear elution effects in split-peak chromatography. I. Computer simulations for the cases of irreversible diffusion- and adsorption-limited kinetics 436(1988)111

Investigation of the tetramer-dimer equilibrium in haemoglobin solutions by high-performance SEC on a diol column 437(1988)193

**Haemoglobin, cysteine adducts**

Cysteine adducts of human haemoglobin measured by isoelectric focusing in polyacrylamide gels with a non-linear pH gradient 420(1987)35

**Halazepam**

LC determination of halazepam in commercial tablets 445(1988)310

**Haloaliphatic acids**

Multiple element emission spectral detection GC profiles of halogenated products from chlorination of humic acid and drinking water 438(1988)35

**Halobacteria**

Difference in cell surface hydrophobicity of *Halobacterium salinarum* strains 449(1988)322

**Halo carboxylic acid amides**

Chiral recognition of halo carboxylic acid amides studied by GC on Chirasil-Val 406(1987)157

**Halofantrine**

Ion-paired LC method for the analysis of blood and plasma for the antimalarial drug halofantrine and its putative mono-debutylated metabolite 430(1988)412

Determination of halofantrine and its principal metabolite desbutylhalofantrine in biological fluids by reversed-phase HPLC 433(1988)339

**Halofuginone**

Determination of halofuginone in poultry feeds by HPLC 441(1988)406

**Halogenated compounds**

Effect of temperature on the GC separation of halogenated compounds on polar and non-polar stationary phases 437(1988)33

Comparison of volatile halogenated compounds formed in the chloramination and chlorination of humic acid by GC-electron-capture detection 449(1988)326

**Halogenated organic compounds**

Chemical changes of organic compounds in chlorinated water. XIV. Characterization and determination of halogenated organics formed during chlorination of water from the Tama River 409(1987)259

**Halogenobenzoic acids**

Use of cyclodextrins in isotachopheresis. II.  $\alpha$ -,  $\beta$ - and  $\gamma$ -cyclodextrins as leading electrolyte additives for the separation of *ortho*-, *meta*- and *para*-substituted halogenobenzoic acids 411(1987)153

**Haloperidol**

Simplified measurement of haloperidol by GC with nitrogen-phosphorus detection 423(1987)380

Reversed-phase ion-pair LC method for the determination of low concentrations of haloperidol in plasma 431(1988)113

Simultaneous determination of very low concentrations of haloperidol and reduced haloperidol in human serum by an LC method 433(1988)312

**Halothane**

Quantification of halothane, enflurane and nitrous oxide by means of a simple GC method 403(1987)350

Comparison of the absolute calibration method with the method of standard addition for the determination of halothane in blood by GC headspace analysis 425(1988)99

**Haloxypop-ethoxyethyl**

GC-MS investigation of phenoxypropanoic acid derivatives possessing herbicidal activity 438(1988)55

**Harman**

Measurement of  $\beta$ -carbolines by HPLC with fluorescence detection 428(1988)229

HPLC identification of simple  $\beta$ -carboline alkaloids in specimens of *Heliconiini* butterflies 447(1988)432

**Harmol**

Measurement of  $\beta$ -carbolines by HPLC with fluorescence detection 428(1988)229

**Headspace analysis**

Headspace gas analysis. Liquid desorption of headspace volatiles trapped on activated carbon open tubular traps 402(1987)95

Dynamic headspace GC of volatile compounds in milk 404(1987)163

Dynamic headspace GC: concentration of volatile components after thermal desorption by intermediate cryofocusing in a cold trap. I. Principle and applications 405(1987)87

Dynamic headspace GC: concentration of volatile components after thermal desorption by intermediate cryofocusing in a cold trap. II. Effect of sampling and desorption parameters on recovery 405(1987)99

Comparison of the absolute calibration method with the method of standard addition for the determination of halothane in blood by GC headspace analysis 425(1988)99

Flavor characterization of dentifrices using kinetic headspace GC 438(1988)273

Analysis of flavor absorption into plastic packaging materials using multiple headspace extraction GC 438(1988)281

Automatic headspace GC method for the simultaneous determination of trichloroethylene and metabolites in blood and urine 442(1988)317

Determination of trichloroethylene metabolites in rat liver homogenate using headspace GC 442(1988)325

Purge-and-trap GC method for residual volatiles in high-temperature polymers 450(1988)394

**Heats of adsorption**

GC study of the temperature dependence of the hydrogen bond energy of adsorption on silica 446(1988)55

**Heat-stable enterotoxin**

Alternative procedure for the purification of the heat-stable enterotoxin of enterotoxigenic *Escherichia coli* pathogenic for calves 420(1987)253

**Heavy alkylated benzene**

Heavy alkylated benzene stationary phase for the GLC separation of light hydrocarbons 404(1987)328



Heavy alkylated benzene stationary phase for the GLC separation of ethylbenzene, xylenes, styrene, iso- and *n*-propylbenzene 409(1987)371

#### Heavy metal ions

Capillary tube isotachophoretic separation of heavy metal ions using complex-forming equilibria between cyanide as terminating ion and the metal ions 447(1988)383

#### Hematoporphyrin

Normal-phase HPLC of free acid dicarboxylic porphyrins and hematoporphyrin derivative on silica 422(1987)73

#### Henry's constant

Solubility properties in polymers and biological media. II. A new method for the characterisation of the adsorption of gases and vapours on solids 409(1987)15

#### Heparan sulphate

Extracellular mammalian polysaccharides: glycosaminoglycans and proteoglycans (Review) 418(1987)187

Fractionation of aorta glycosaminoglycans by HPLC 419(1987)275

Methods for analysis of urinary glycosaminoglycans (Review) 429(1988)293

#### Heptafluorobutyrate

Perfluorinated acids as ion-pairing agents in the determination of monoamine transmitters and some prominent metabolites in rat brain by HPLC with amperometric detection 449(1988)191

#### Heptafluorobutyric acid

Use of perfluorinated carboxylic acids in the separation of metabolites of vasopressin prior to radioimmunoassay 430(1988)128

#### N-Heptafluorobutyryl-L-prolyl chloride

Separation of optical isomers of methoxyphenamine and its metabolites as N-heptafluorobutyryl-L-prolyl derivatives by fused-silica capillary GC 431(1988)210

#### Heptakis(2,3,6-tri-O-methyl)- $\beta$ -cyclodextrin

Use of cyclodextrins in isotachopheresis. VI. Cyclodextrins as leading electrolyte additives for the separation of bile acids 450(1988)373

#### Heroin

Concentrations of free and conjugated morphine

in blood in twenty cases of heroin-related deaths 422(1987)267

#### Hesperetin

Direct enantiomeric separation of racemic flavanones by HPLC using cellulose triacetate as a chiral stationary phase 441(1988)417

#### Heterobifunctional ligands

Affinity chromatography on a hydrophobic matrix using a heterobifunctional ligand 438(1988)339

#### Heterogeneous catalysis

Contributions of GC to the development of heterogeneous catalysis 406(1987)151

#### Heterogeneous reactions

Chromatographic method for kinetic study of heterogeneous reactions 446(1988)65

#### *endo*-1,4,5,6,7,7-

#### Hexachlorobicyclo[2.2.1]hept-5-ene-2-carboxylic acid

*endo*-1,4,5,6,7,7-Hexachlorobicyclo[2.2.1]hept-5-ene-2-carboxylic acid, a superior resolving agent for the HPLC separation of enantiomers of hydroxylated derivatives of two azaaromatic hydrocarbons 430(1988)53

#### 1-O-Hexadecyl-2-O-acetyl-*sn*-glycero-3-phosphocholine

Characterization and quantification of PAF-acether (platelet-activating factor) as a heptafluorobutyrate derivative of 1-O-alkyl-2-acetyl-*sn*-glycerol by capillary column GC with electron-capture detection 423(1987)23

#### Hexahelicene

Hexahelicene chiral stationary phase. I. Phase synthesis and use in HPLC resolution of enantiomers 450(1988)450(1988)157

#### *trans*-6,6a,7,10a,11-Hexahydro-8,9-dimethyl-11-oxodibenz[*b,e*]oxepin-3-acetic acid

Enantiomeric analysis of a new anti-inflammatory agent in rat plasma using a chiral  $\beta$ -cyclodextrin stationary phase 426(1988)417

#### Hexahydrohippuric acid

Unusual conjugates in biological profiles originating from consumption of onions and garlic 421(1987)1

**Hexahydroxybenzene**

Disc-like liquid crystal stationary phases derived from hexahydroxybenzene 402(1987)73

**1,6-Hexamethyleneaminoisocyanate**

Trace analysis of airborne 1,6-hexamethylenediisocyanate and the related aminoisocyanate and diamine by glass capillary GC 435(1988)453

**Hexamethylene bisacetamide**

Determination of hexamethylene bisacetamide by capillary GC 414(1987)510

GC analysis of metabolites of the cell differentiating agent hexamethylene bisacetamide 415(1987)148

**1,6-Hexamethylenediamine**

Trace analysis of airborne 1,6-hexamethylenediisocyanate and the related aminoisocyanate and diamine by glass capillary GC 435(1988)453

**1,6-Hexamethylenediisocyanate**

Trace analysis of airborne 1,6-hexamethylenediisocyanate and the related aminoisocyanate and diamine by glass capillary GC 435(1988)453

**Hexobarbital**

Simultaneous quantitation of *d*- and *l*-hexobarbital in rat blood by HPLC 419(1987)426

**Hexogen**

Improved method for the trace analysis of explosives by silica capillary column GC with thermal energy analysis detection 410(1987)181

**Hexosamines**

Quantitative GC measurement of glycosaminoglycan hexosamines in urine and plasma 420(1987)231

**2-Hexuloses**

Degradation of ketoses during aqueous HPLC on lead-form cation-exchange resins 437(1988)387

**Hierarchical ascending classification**

Hierarchical ascending classification of Kováts retention indices as a means of indicating siloxane stationary phase specificities towards aliphatic carboxylic esters 406(1987)95

**High-density lipoproteins**

Separation and characterization of high-density lipoprotein subpopulations by GPC 414(1987)25

Anion-exchange fast protein LC characterization and purification of apolipoproteins A-I, A-II, C-I, C-II, C-III<sub>0</sub>, C-III<sub>1</sub>, C-III<sub>2</sub> and E from human plasma 414(1987)35

Improved HPLC method for the determination of apolipoproteins in serum high-density lipoproteins 430(1988)135

**High-mobility-group proteins**

Fractionation of plant and animal high mobility group chromosomal proteins by ion-exchange and reversed-phase HPLC 405(1987)327

Non-histone chromatin proteins that recognize specific sequences of DNA (Review) 418(1987)73

Separation and quantification of histone H1 subtypes and high-mobility-group proteins by reversed-phase LC: protein levels in rat tissues during postnatal development 426(1988)65

Separation of Friend erythroleukaemic cell histones and high-mobility-group proteins by reversed-phase HPLC 450(1988)309

**High-performance electrophoresis**

Carrier-free zone electrophoresis, displacement electrophoresis and isoelectric focusing in a high-performance electrophoresis apparatus 403(1987)47

**High-temperature HPLC**

HPLC at elevated temperatures: examination of conditions for the rapid separation of large molecules 435(1988)1

**High-temperature polymers**

Purge-and-trap GC method for residual volatiles in high-temperature polymers 450(1988)394

**Hippuric acid**

Determination of benzoic acid and hippuric acid in human plasma and urine by HPLC 425(1988)67

Determination of the angiotensin converting enzyme inhibitor benazeprilat in plasma and urine by an enzymic method 425(1988)353

**Histamine**

Simultaneous determination of histidine-containing dipeptides, histamine, methylhistamine and histidine by HPLC 432(1988)315

Detection by HPLC of polyamines formed by clostridal putrefaction of caseins 442(1988)267

**Histidine**

Amino acid analysis: the histidine hangover 403(1987)271

Complex-forming equilibria in isotachopheresis. VII. Evaluation of the ion-pair formation constants of phosphorus oxo acids with histidine and assessment of their separability 410(1987)279

Determination of stable isotopically substituted histidine in human plasma by GC-MS 413(1987)1

Simultaneous determination of histidine-containing dipeptides, histamine, methylhistamine and histidine by HPLC 432(1988)315

GC analysis of histidine: effect on other amino acids present 446(1988)231

**Histidine-containing dipeptides**

Simultaneous determination of histidine-containing dipeptides, histamine, methylhistamine and histidine by HPLC 432(1988)315

**Histones**

Chromatography of complex protein mixtures (Review) 418(1987)115

Separation and quantification of histone H1 subtypes and high-mobility-group proteins by reversed-phase LC: protein levels in rat tissues during postnatal development 426(1988)65

Separation of Friend erythroleukaemic cell histones and high-mobility-group proteins by reversed-phase HPLC 450(1988)309

**Hold-up volume**

Phase ratio determination in an ion-exchange column having pores partially accessible to proteins 442(1988)15

**Holistic total surface area**

Holistic molecular conformation and total surface area calculations as predictors for solution properties and chromatographic parameters 403(1987)1

**Hollow-fibre membrane**

Ion-exclusion chromatography of carboxylic acids with conductivity detection. Peak enhancement using a cation-exchange hollow-fibre membrane and an alkaline solution 447(1988)373

**Hollow-fibre membrane reactor**

LC determination of penicillins by postcolumn

degradation with sodium hypochlorite using an hollow-fibre membrane reactor 447(1988)365

**Homocarnosine**

Simultaneous determination of histidine-containing dipeptides, histamine, methylhistamine and histidine by HPLC 432(1988)315

**Homocysteine**

HPLC fluorometry for the determination of thiols in biological samples using N-[4-(6-dimethylamino-2-benzofuranyl)phenyl]maleimide 414(1987)11

Determination of free and total homocysteine in human plasma by HPLC with fluorescence detection 422(1987)43

Measurement of plasma thiols after derivatization with monobromobimane 424(1988)141

**Homoeriodictyol**

Direct enantiomeric separation of racemic flavanones by HPLC using cellulose triacetate as a chiral stationary phase 441(1988)417

**Homogentisic acid**

Analysis of homogentisic acid in body fluids by HPLC 413(1987)233

**Homoisoflavonoids**

Separation and determination of homoisoflavonoids in *Ophiopogon japonicus* by reversed-phase HPLC 437(1988)265

**Homoserine**

Assignment of the configurations of the amino acids in peptidic siderophores 438(1988)291

**Homotaurine**

Determination of calcium acetylhomotaurinate by LC with fluorimetric and electrochemical detection 414(1987)417

**Homovanillic acid**

Simultaneous LC determination of seventeen of the major monoamine neurotransmitters, precursors and metabolites. I. Optimization of the mobile phase using factorial designs and a computer program to predict chromatograms 415(1987)261

Simultaneous LC determination of seventeen of the major monoamine neurotransmitters, precursors and metabolites. II. Assessment of human brain and cerebrospinal fluid concentrations 415(1987)275

Combined ion-pair extraction and HPLC for the determination of the biogenic amines and their major metabolites in single brain tissue samples 421(1987)51

LC determination of vanillylmandelic acid and homovanillic acid by a column-switching technique involving direct injection of urine 421(1987)111

Simple HPLC method for the simultaneous measurement of vanillylmandelic acid and homovanillic acid 423(1987)278

Simultaneous determination of the major acidic metabolites of catecholamines and serotonin in urine by LC with electrochemical detection after a one-step sample clean-up on Sephadex G-10; influence of vanilla and banana ingestion 424(1988)273

Measurement of homovanillic acid in small volumes of plasma using LC with electrochemical detection 426(1988)141

Simultaneous determination of urinary vanillylmandelic acid, 5-hydroxyindoleacetic acid and homovanillic acid by LC 428(1988)147

HPLC method for measuring homovanillic acid in cerebrospinal fluid using electrochemical detection with internal standardization 430(1988)112

Determination of plasma homovanillic acid by two-step solid-phase extraction and HPLC with electrochemical detection 430(1988)118

Method of measurement of urinary homovanillic acid and vanillylmandelic acid by GC-MS suitable for neuroblastoma screening 432(1988)273

Effects of long-term storage on the concentrations of the unconjugated acidic metabolites of the trace amines, indoleamines and catecholamines 433(1988)23

### Hormone receptors

Application of high-performance liquid chromatofocusing to the study of prostatic androgen receptors. Effect of stabilizing agents on the heterogeneity of receptor structure 403(1987)171

Improved Stokes radius measurement of the glucocorticoid receptor using TSK G4000SW and TSK G3000SW high-performance size-exclusion columns. Analytical and preparative applications 403(1987)183

### Hormones

see also Plant hormones and steroids

Analysis of estrogen sulphate mixtures in pharmaceutical formulations by reversed-phase chromatography 402(1987)273

Resolution of the enantiomers of thyroid hormones by high-performance ligand-exchange chromatography using a chemically bonded chiral stationary phase 404(1987)391

Coordinative interactions in the separation of insulin and derivatives by HPLC 408(1987)43

Recovery of polypeptides after reversed-phase HPLC 408(1987)191

Separation of ecdysteroids by HPTLC using automated multiple development 408(1987)445

TLC of ecdysone acyl esters and their 2,3-acetonide derivatives 411(1987)379

Fluorimetric determination of rat growth hormone in pituitary cell culture by HPLC with post-column derivatization 417(1987)409

Separation of protein hormones (Review) 418(1987)245

Combined use of rapid-flow fractionation and HPLC for the determination of serum testosterone and application to study of stress response to physical exercise 423(1987)285

Automated direct assay system for the measurement of sex steroid hormones in serum using HPLC 426(1988)33

Chromatographic analysis of the trinitrophenyl derivatives of insulin 427(1988)239

Determination of oestrogens in pregnancy urine by HPLC with fluorescence detection 431(1988)249

Measurement of urinary estriol glucuronides during the menstrual cycle by HPLC 434(1988)61

Complete separation of nine equine oestrogens by HPLC 435(1988)385

Separation of ecdysone acyl esters and their 2,3-acetonide derivatives by HPLC 436(1988)279

Supercritical fluid chromatography of ecdysteroids 436(1988)497

Comparison of non-hydrophobic C<sub>18</sub> bonded TLC and HPTLC plates for reversed-phase TLC of ecdysteroids 438(1988)419

Rapid two-dimensional TLC system for the

separation of multiple steroids of secretory and neuroendocrine interest 439(1988)441

High-performance displacement chromatography of melanotropins and their derivatives 440(1988)157

Several intermediate forms in the processing of rat lutropin subunits as shown by sodium dodecyl sulphate polyacrylamide gel electrophoresis 440(1988)353

Synthetic peptide substrates as models to study a pro-ocytocin/neurophysin converting enzyme 440(1988)439

Analysis of ecdysteroids by supercritical-fluid chromatography 441(1988)165

Reduction studies on bacterial recombinant somatomedin C/insulin-like growth factor-1 443(1988)183

Single-step purification of two hyperglycaemic neurohormones from the sinus gland of *Procambarus bouvieri*. Comparative peptide mapping by means of HPLC 443(1988)337

#### **Human growth hormone**

Recovery of polypeptides after reversed-phase HPLC 408(1987)191

#### **Human serum albumin**

Recovery of polypeptides after reversed-phase HPLC 408(1987)191

Procedure for the isolation of a large peptic fragment of human serum albumin 424(1988)13

#### **Humic acid**

Multiple element emission spectral detection GC profiles of halogenated products from chlorination of humic acid and drinking water 438(1988)35

Comparison of volatile halogenated compounds formed in the chloramination and chlorination of humic acid by GC-electron-capture detection 449(1988)326

#### **Humic acids**

Computer-controlled linear pH gradient for HPLC fractionations of aromatic carboxylic acids and of humic and fulvic acids 439(1988)213

#### **Humic compounds**

Flame ionization and flame thermionic detection of carbon and nitrogen in aquatic lipid and humic-type classes with an Iatroscan Mark IV 435(1988)350

#### **Humic substances**

Mobility of Fe(III)-phenolic acid complexes in TLC 402(1987)358

GC determination of C<sub>1</sub>-C<sub>5</sub> low-molecular-weight organic acids in alkaline permanganate oxidation products of humic substances 403(1987)225

#### **Hyaluronate**

Extracellular mammalian polysaccharides: glycosaminoglycans and proteoglycans (Review) 418(1987)187

#### **Hyaluronic acid**

High-performance GPC of glycosaminoglycans. Column calibration by gel electrophoresis 402(1987)149

Rapid and sensitive method for measurement of hyaluronic acid and isomeric chondroitin sulfates using HPLC 413(1987)9

LC assay for constituent disaccharides of hyaluronic acid and chondroitin sulphate isomers 415(1987)231

Fractionation of aorta glycosaminoglycans by HPLC 419(1987)275

Methods for analysis of urinary glycosaminoglycans (Review) 429(1988)293

Analysis by high-performance GPC of hyaluronic acid in animal skins and rabbit synovial fluid 435(1988)335

#### **Hyaluronidase**

Characterization of bee venom and its main components by HPLC 408(1987)341

#### **Hydrocarbons**

Evaluation of disposable cartridges for trace enrichment from aqueous solutions 402(1987)201

Heavy alkylated benzene stationary phase for the GLC separation of light hydrocarbons 404(1987)328

Use of silicalite molecular sieve for the GC determination of permanent gases and volatile hydrocarbons emitted during coal processing 405(1987)213

Potential solid supports for GC prepared on the basis of certain rocks 406(1987)145

Use of Kováts indices for prediction of retention temperatures in linear temperature-programmed capillary GC 407(1987)65

Comparison of volatile components between fresh and rotten mussels by GC-MS 409(1987)251

Organo-modified smectite for the GC separation of various compounds 410(1987)121

GC determination for forensic purposes of petroleum fuel inhaled just before fatal burning 424(1988)49

Analysis of hydrocarbons in aquatic sediments. II. Evaluation of common preparative procedures for petroleum and chlorinated hydrocarbons 436(1988)503

HPLC method for isolating organic contaminants from tissue and sediment extracts 437(1988)161

Analytical characterisation of hydrocarbon and hetero compounds mixtures from reactions of brown coal (in German) 446(1988)305

Shape selectivity of stationary phases for the separation of isomeric hydrocarbons by capillary GC 446(1988)339

Modified method for analysis of C<sub>2</sub>-C<sub>5</sub> hydrocarbons in an aromatic-alkane matrix using an automated thermal desorber 449(1988)175

#### **Hydrochlorothiazide**

LC screening of diuretics in urine 415(1987)347

Simultaneous determination of amiloride and hydrochlorothiazide in plasma by reversed-phase HPLC 423(1987)351

#### **Hydrogen**

GC technique to simultaneously quantitate the gases produced by intestinal microorganisms from fermentation mixtures 414(1987)405

GC evaluation of hydrogen in the breath as a diagnostic method for the study of intestinal digestion and absorption of dietary carbohydrates in man 416(1987)119

#### **Hydrogen atmosphere flame ionization detection**

Determination of silicon number in silylated organic compounds using the hydrogen atmosphere flame ionization detector 408(1987)329

#### **Hydrogen bond energy**

GC study of the temperature dependence of the hydrogen bond energy of adsorption on silica 446(1988)55

#### **Hydrogen bonding**

Reordering/resolution of silica immobilized hydrogen bonding ligands used as stationary phases in HPLC 408(1987)93

Effect of hydrogen bonding on the HPLC behaviour of organic compounds. Relationship between capacity factors and partition coefficients 439(1988)227

Demonstration of intermolecular and intramolecular hydrogen bonding by GC measurements 440(1988)329

#### **Hydrogen peroxide**

Determination of hydrogen peroxide in beverages by HPLC with fluorescence detection 411(1987)423

#### **Hydrophase HP-PEI**

Ion-exchange chromatography of proteins on a polyethyleneimine-grafted hydrophilic polymer for HPLC 443(1988)133

#### **Hydrophile-lipophile balance**

Polarity and hydrophile-lipophile balance of model hydroxyoximes and their intermediates as determined by reversed-phase GC 402(1987)55

#### **Hydrophilic compounds**

Ion-exclusion chromatography of non-ionic substances with conductivity detection 409(1987)271

#### **Hydrophobic constant**

Factor analysis of trends in selectivity in GLC of 175 saturated and unsaturated esters on 14 polysiloxane stationary phases 404(1987)11

#### **Hydrophobic interaction chromatography**

Resolution of proteins in linear gradient elution ion-exchange and hydrophobic interaction chromatography 409(1987)101

Hydrophobic interaction chromatography of peptides as an alternative to reversed-phase chromatography 444(1988)269

#### **Hydrophobicity**

see also Lipophilicity

Effect of polymer composition on the relative hydrophobicity of the phases of the biphasic system aqueous dextran-poly(ethylene glycol) 403(1987)123

Effects of differences in charge and hydrophobicity of surface amino acids of

hemoglobins on high-performance GPC  
428(1988)247

Studies on fungicidal activity of N-phenylcarbamates. IV. Determination of the hydrophobicity of N-phenylcarbamates by HPLC  
436(1988)316

HPLC of amino acids, peptides and proteins. LXXXV. Evaluation of the use of hydrophobicity coefficients for the prediction of peptide elution profiles  
438(1988)197

Adsorption chromatography on cellulose. II. Separations of aromatic amino acids, biogenic amines, alkaloids, dyes and phenols and determination of hydrophobic constants  
440(1988)165

Group contributions to hydrophobicity and elution behaviour of pyridine derivatives in reversed-phase HPLC  
449(1988)95

Difference in cell surface hydrophobicity of *Halobacterium salinarium* strains  
449(1988)322

### Hydroquinone

Identification and quantification of hydroquinone and some of its ethers in cosmetic products by reversed-phase HPLC  
404(1987)267

Interconversion of hydroquinone and benzoquinone during styrene-divinylbenzene resin-based ion-exclusion chromatography  
409(1987)404

### Hydroquinones

Adsorption chromatography on cellulose. I. Salting-out chromatography of organic compounds  
406(1987)389

### $\alpha$ -Hydroxyacetic acids

HPLC of mixtures of phenol or guaiacol and their reaction products with glyoxylic acid  
437(1988)185

### Hydroxyacetone

GC method for the estimation of acetone and its metabolites in biological samples  
415(1987)372

### Hydroxy acids

Identification by GC-MS of short-chain hydroxy acids produced by *Fusobacterium* species and *Clostridium innocuum*  
420(1987)121

Thermospray-MS analysis of underivatized monohydroxy fatty acids: application to stimulated platelets  
425(1988)25

Simultaneous determination of C<sub>2</sub>-C<sub>22</sub> non-esterified fatty acids and other metabolically

relevant carboxylic acids in biological material by GC of their benzyl esters  
425(1988)257

Studies of metabolites in diarrheal stool specimens positive for *Klebsiella*, *Serratia*, and *Proteus* spp. by frequency-pulsed electron-capture GC  
430(1988)209

TLC enantiomeric resolution via ligand exchange  
448(1988)11

### Hydroxyalprazolams

Alprazolam,  $\alpha$ -hydroxy- and 4-hydroxyalprazolam analysis in plasma by HPLC  
430(1988)180

### 4-Hydroxyalprenolol

Derivatization of 2-amino alcohols with phosgene in aqueous media: limitations of the reaction selectivity as found in the presence of O-glucuronides of alprenolol in urine  
413(1987)270

### 10-Hydroxyamitriptyline

Applications of column-switching technique in biopharmaceutical analysis. I. HPLC determination of amitriptyline and its metabolites in human plasma  
416(1987)99

### Hydroxyamphetamine

Determination of  $\beta$ -phenylalkylamines in pharmaceutical preparations by analytical capillary isotachopheresis  
405(1987)389

### 3-Hydroxyanthranilic acid

High-voltage electrophoretic separation of tryptophan metabolites of the kynurenine pathway  
420(1987)263

### Hydroxyanthraquinone

Separation of hydroxyanthraquinone derivatives extracted from rheum with analytical high-speed counter-current chromatography  
442(1988)455

### 4-Hydroxyantipyrine

Improved HPLC assay of antipyrine, hydroxymethylantipyrine, 4-hydroxyantipyrine and norantipyrine in urine  
433(1988)305

### 4-Hydroxyantipyrine sulphate

Application of molecular-secondary-ion MS for drug metabolism studies. I. Direct analysis of conjugates by GLC-secondary-ion mass spectrometry  
414(1987)139

### Hydroxyapatite

HPLC of glycosides on a new type of hydroxyapatite column  
407(1987)205

High-performance adsorption chromatography of

proteins on 2- $\mu$ m spherical beads of hydroxyapatite 440(1988)305

#### Hydroxyarachidic acids

GC-MS of monohydroxyeicosatetraenoic acids as their methyl esters trimethylsilyl, allyldimethylsilyl and *tert.*-butyldimethylsilyl ethers 416(1987)219

#### $\beta$ -Hydroxyaspartic acid

Assignment of the configurations of the amino acids in peptidic siderophores 438(1988)291

#### $\gamma$ -Hydroxybaclofen

Rapid simultaneous determination of baclofen and its  $\gamma$ -hydroxy metabolite in urine by HPLC with UV detection 420(1987)212

#### Hydroxybenzaldehyde

Retention behaviour of aromatic aldehyde derivatives in reversed-phase LC 404(1987)242

#### Hydroxybenzoic acid glucosides

Analysis of naturally occurring hydroxybenzoic acid glucosides and hydroxybenzoyl glucoses and their behaviour on a polyamide column 441(1988)335

#### Hydroxybenzoyl glucoses

Analysis of naturally occurring hydroxybenzoic acid glucosides and hydroxybenzoyl glucoses and their behaviour on a polyamide column 441(1988)335

#### 3-Hydroxy-2-butanone

Determination of methyl ethyl ketone and its metabolites in urine using capillary GC 428(1988)275

#### Hydroxybutyric acid

Determination of ionic metabolites from ethylene glycol in human blood by isotachopheresis 416(1987)111

#### 3-Hydroxycarbofuran

HPLC determination of aryl N-methylcarbamate residues using post-column hydrolysis electrochemical detection 442(1988)333

#### Hydroxychloroquine

Analysis of blood and urine samples for hydroxychloroquine and three major metabolites by HPLC with fluorescence detection 433(1988)197

#### Hydroxycholesterols

Determination of some hydroxycholesterols in human serum samples 416(1987)1

#### Hydroxyclo mipramine

Determination of clo mipramine and its hydroxylated and demethylated metabolites in plasma and urine by LC with electrochemical detection 416(1987)311

#### Hydroxy compounds

Anthracene-9-carbonyl chloride as a fluorescence and UV derivatising reagent for the HPLC analysis of hydroxy compounds 445(1988)393

#### 18-Hydroxycorticosterone

Simplified method for the determination of plasma 18-hydroxycorticosterone 426(1988)148

#### *trans*-3'-Hydroxycotinine

GC determination of *trans*-3'-hydroxycotinine, a major metabolite of nicotine in smokers 415(1987)400

#### 7-Hydroxycoumarin

Analysis of coumarin and its urinary metabolites by HPLC 416(1987)165

#### 15 $\beta$ -Hydroxycyproterone acetate

Automated HPLC assay for cyproterone acetate and 15 $\beta$ -hydroxycyproterone acetate in plasma 420(1987)432

#### 5-Hydroxydantrolene

Determination of dantrolene and its reduced and oxidized metabolites in plasma by HPLC 430(1988)187

#### 4-Hydroxydebrisoquine

Direct determination of debrisoquine and 4-hydroxydebrisoquine using a HPLC switching technique 419(1987)469

Rapid LC determination of debrisoquine and its hydroxy metabolite in human urine to define hydroxylation phenotypes 423(1987)340

HPLC method for the analysis of debrisoquine and its *S*-(+)- and *R*-(-)-hydroxy metabolites in urine 423(1987)344

HPLC quantitation of 4-hydroxydebrisoquine in microsomal incubates by use of silica columns, aqueous mobile phase and automated column switching 424(1988)435

Comparison of GC and HPLC assays for the determination of debrisoquine and its 4-hydroxy metabolite in human fluids 425(1988)311

Assay of debrisoquine and 4-hydroxydebrisoquine in urine by reversed-phase HPLC using on-line sample clean-up on a standard isocratic chromatograph 428(1988)178



**3-Hydroxydecanoic acid dimethylate**

Identification of 3-hydroxydecanoic acid dimethylate in human urine 420(1987)450

**Hydroxydemethylclomipramine**

Determination of clomipramine and its hydroxylated and demethylated metabolites in plasma and urine by LC with electrochemical detection 416(1987)311

**2-Hydroxydesipramine**

HPLC determination of imipramine and its metabolites in rat brain 421(1987)412

Isolation and identification of the glucuronide conjugate of 2-hydroxydesipramine by preparative LC 427(1988)287

**Hydroxyeicosatetraenoic acids**

GC-MS of monohydroxyeicosatetraenoic acids as their methyl esters trimethylsilyl, allyldimethylsilyl and *tert.*-butyldimethylsilyl ethers 416(1987)219

MS evaluation of the *tert.*-butyldimethylsilyl derivatives of monohydroxyeicosatetraenoic acids and leukotrienes 423(1987)1

Automated HPLC extraction and quantification procedure for lipoxygenase metabolites 430(1988)1

Isocratic determination of arachidonic acid 5-lipoxygenase products in human neutrophils by HPLC 430(1988)299

**9-Hydroxyellipticine**

Fluorodensitometric assay of 6*H*-pyrido[4,3-*b*]-5,11-dimethylcarbazoles (ellipticine and derivatives) biosynthesized by *Ochrosia elliptica* cultures *in vitro* 409(1987)426

**N-(2-Hydroxyethoxyethyl)-2,2,3,3,4,4,4-heptafluorobutanamide**

Comparative study of alkyl and fluoroalkyl N-(2-hydroxy-ethoxyethyl)-amides as reversed-phase LC modifiers 448(1988)365

**N-(2-Hydroxyethoxyethyl)-heptanamide**

Comparative study of alkyl and fluoroalkyl N-(2-hydroxy-ethoxyethyl)-amides as reversed-phase LC modifiers 448(1988)365

**N-(2-Hydroxyethoxyethyl)-hexanamide**

Comparative study of alkyl and fluoroalkyl N-(2-hydroxy-ethoxyethyl)-amides as reversed-phase LC modifiers 448(1988)365

**Hydroxyethylcellulose**

GLC of acid hydrolysates of hydroxyethylstarch and hydroxyethylcellulose 435(1988)374

**N-1-Hydroxy-ethylflurazepam**

Simultaneous determination of flurazepam and five metabolites in serum by HPLC and its application to pharmacokinetic studies in rats 423(1987)251

**3-Hydroxy-N-(N-ethylglycyl)-2,6-xylidine**

*In vivo* metabolism of lidocaine in the rat. Isolation of urinary metabolites as pentafluorobenzoyl derivatives and their identification by combined GC-MS 421(1987)267

**Hydroxyethyl methacrylate**

Simple GC determination of the purity of hydroxyethyl methacrylate 403(1987)355

**Hydroxyethylstarch**

GLC of acid hydrolysates of hydroxyethylstarch and hydroxyethylcellulose 435(1988)374

 **$\beta$ -Hydroxyethyltheophylline**

Simplified description of HPLC separation under overload conditions, based on the Craig distribution model. IV. Comparison of the model with experimental data for samples containing two or more solutes 405(1987)31

**Hydroxyethylvinyldeuteroporphyrin**

Normal-phase HPLC of free acid dicarboxylic porphyrins and hematoporphyrin derivative on silica 422(1987)73

**Hydroxy fatty acids**

Use of pentafluorobenzyl and pentafluoropropionyl-pentafluorobenzyl esters of bacterial fatty acids for GC analysis with electron-capture detection 417(1987)366

Reversed-phase ion-interaction chromatography of leukotrienes, lipoxins and related compounds 427(1988)1

Determination of environmental levels of peptidoglycan and lipopolysaccharide using GC with negative-ion chemical-ionization MS utilizing bacterial amino acids and hydroxy fatty acids as biomarkers 431(1988)1

Identification of isomeric hydroxy fatty acids in aerosol samples by capillary GC-MS 438(1988)309

**Hydroxyfloctafenic acid**

Simultaneous determination of floctafenine and its main metabolites, floctafenic and hydroxyfloctafenic acid, using high-performance liquid chromatography with gradient elution 432(1988)412

**7-Hydroxyflumequine**

Determination of flumequine and its 7-hydroxy metabolite by reversed-phase HPLC 414(1987)499

**3'-Hydroxygenkwanin**

Separations of flavonoids and alkaloids in medicinal herbs by high-speed counter-current chromatography 435(1988)159

**1-(5'-Hydroxyhexyl)-3,7-dimethylxanthine**

HPLC determination of pentoxifylline and its hydroxy metabolite in human plasma 416(1987)183

HPLC analysis of pentoxifylline and 1-(5'-hydroxyhexyl)-3,7-dimethylxanthine in whole blood 419(1987)368

**2-Hydroxyimipramine**

HPLC determination of imipramine and its metabolites in rat brain 421(1987)412

**5-Hydroxyindole-3-acetic acid**

Simultaneous LC determination of seventeen of the major monoamine neurotransmitters, precursors and metabolites. I. Optimization of the mobile phase using factorial designs and a computer program to predict chromatograms 415(1987)261

Simultaneous LC determination of seventeen of the major monoamine neurotransmitters, precursors and metabolites. II. Assessment of human brain and cerebrospinal fluid concentrations 415(1987)275

Determination of serotonin and 5-hydroxyindoleacetic acid in urine by reversed-phase ion-pair partition chromatography with fluorimetric detection 419(1987)85

Combined ion-pair extraction and HPLC for the determination of the biogenic amines and their major metabolites in single brain tissue samples 421(1987)51

Simultaneous determination of the major acidic metabolites of catecholamines and serotonin in urine by LC with electrochemical detection after a one-step sample clean-up on Sephadex G-10;

influence of vanilla and banana ingestion 424(1988)273

Simultaneous determination of urinary vanillylmandelic acid, 5-hydroxyindoleacetic acid and homovanillic acid by LC 428(1988)147

Sensitive determination of deuterated and non-deuterated indole-3-acetic acid and 5-hydroxyindole-3-acetic acid by combined capillary GC negative-ion chemical ionization MS 428(1988)209

Simultaneous micro-determinations of serotonin and 5-hydroxyindole-3-acetic acid with 5-hydroxy-N<sub>ω</sub>-methyltryptamine, as an internal standard, in biological materials by HPLC with electrochemical detection 430(1988)381

Determination of plasma 5-hydroxytryptophan, 5-hydroxytryptamine, 5-hydroxyindoleacetic acid, tryptophan and melatonin by HPLC with electrochemical detection 431(1988)170

Effects of long-term storage on the concentrations of the unconjugated acidic metabolites of the trace amines, indoleamines and catecholamines 433(1988)23

**3-Hydroxykynurenine**

Quantification of 3-hydroxykynurenine in brain by HPLC and electrochemical detection 428(1988)340

**Hydroxylamine**

HPLC of N-terminal tyrosine-containing oligopeptides by pre-column fluorescence derivatization with hydroxylamine, cobalt(II) and borate reagents 411(1987)305

**11β-Hydroxylase**

Micro-assay for sheep 11β-hydroxylase activity using HPLC for steroid analysis 442(1988)424

**3-Hydroxylidocaine**

In vivo metabolism of lidocaine in the rat. Isolation of urinary metabolites as pentafluorobenzoyl derivatives and their identification by combined GC-MS 421(1987)267

**2-Hydroxymethamphetamine**

Separation of optical isomers of methoxyphenamine and its metabolites as N-heptafluorobutyl-L-prolyl derivatives by fused-silica capillary GC 431(1988)210

**7-Hydroxymethotrexate**

Separation and identification of methotrexate and its metabolites, 7-hydroxymethotrexate and polyglutamates, in human tissues by reversed-phase HPLC coupled with radioimmunoassay 415(1987)445

Column LC of methotrexate and its metabolites using a post-column photochemical reactor and fluorescence detection 419(1987)213

Concurrent analysis of methotrexate, trimethoprim, sulphamethoxazole and their major metabolites in plasma by HPLC 422(1987)328

Determination of methotrexate and its major metabolite, 7-hydroxymethotrexate, using capillary zone electrophoresis and laser-induced fluorescence detection 426(1988)129

**Hydroxymethylantipyrine**

Improved HPLC assay of antipyrine, hydroxymethylantipyrine, 4-hydroxyantipyrine and norantipyrine in urine 433(1988)305

**Hydroxymethylcimetidine**

Highly sensitive determination of cimetidine and its metabolites in serum and urine by HPLC 422(1987)301

**Hydroxymethylmexiletine**

HPLC method for the determination of *p*-hydroxylated and hydroxymethylated metabolites of mexiletine in human serum 430(1988)406

HPLC assay for mexiletine hydroxylation in microsomes of human liver 431(1988)369

**2-Hydroxymethylpiperidine**

Separation of hydroxy and polyhydroxy derivatives of 2-carboxy and 2-hydroxymethyl piperidine and pyrrolidine by HPLC 445(1988)295

**2-[4-(2-Hydroxy-2-methylpropyl)phenyl]propionic acid**

Determination of ibuprofen and its major metabolites in human urine by HPLC 430(1988)93

**1-Hydroxymethyltriazolam**

HPLC determination of triazolam and its metabolites in human urine 422(1987)197

**5-Hydroxy-N<sub>ω</sub>-methyltryptamine**

Simultaneous micro-determinations of serotonin and 5-hydroxyindole-3-acetic acid with 5-hydroxy-N<sub>ω</sub>-methyltryptamine, as an internal

standard, in biological materials by HPLC with electrochemical detection 430(1988)381

**α-Hydroxymetoprolol**

Determination of metoprolol and two major metabolites in plasma and urine by column LC and fluorometric detection 417(1987)357

***p*-Hydroxymexiletine**

HPLC method for the determination of *p*-hydroxylated and hydroxymethylated metabolites of mexiletine in human serum 430(1988)406

HPLC assay for mexiletine hydroxylation in microsomes of human liver 431(1988)369

**1-Hydroxymidazolam**

Rapid and sensitive HPLC assay for midazolam and 1-hydroxymidazolam, the major metabolite, in human serum 433(1988)326

**3-Hydroxymorphinan**

HPLC determination of dextrorphan and 3-hydroxymorphinan in human plasma based on a selective pre-column sample clean-up 420(1987)217

**β-Hydroxymyristic acid**

Determination of endotoxins by GC: evaluation of electron-capture and negative-ion chemical-ionization MS detection of halogenated derivatives of β-hydroxymyristic acid 417(1987)11

**Hydroxynimesulide**

Simultaneous determination of nimesulide and hydroxynimesulide in human plasma and urine by HPLC 425(1988)413

***erythro*-9-(2-Hydroxy-3-nonyl)hypoxanthine**

HPLC assay for urinary metabolites of a drug after extraction by antiserum to the drug: an example using *erythro*-9-(2-hydroxy-3-nonyl)hypoxanthine 425(1988)407

**10-Hydroxynortriptyline**

Applications of column-switching technique in biopharmaceutical analysis. I. HPLC determination of amitriptyline and its metabolites in human plasma 416(1987)99

**N<sup>δ</sup>-Hydroxyornithine**

Assignment of the configurations of the amino acids in peptidic siderophores 438(1988)291

**Hydroxyoximes**

Polarity and hydrophile-lipophile balance of

model hydroxyoximes and their intermediates as determined by reversed-phase GC 402(1987)55

**17 $\beta$ -Hydroxy-3-oxo-17-pregn-4-ene-21-carboxylic acid- $\gamma$ -lactone 7 $\alpha$ -methyl ethyl ester**

HPLC assay for the simultaneous measurement of two spironolactone analogues, 17 $\beta$ -hydroxy-3-oxo-17-pregn-4-ene-21-carboxylic acid- $\gamma$ -lactone 7 $\alpha$ -methyl ethyl ester and its 6 $\beta$ -hydroxylated derivative 428(1988)377

***p*-Hydroxyphenobarbitone**

Simultaneous quantitation of phenobarbitone and *p*-hydroxyphenobarbitone and their perdeuteroethyl analogues in biological fluids by combined GC-MS 420(1987)287

**Hydroxyphenylacetic acid**

Effects of long-term storage on the concentrations of the unconjugated acidic metabolites of the trace amines, indoleamines and catecholamines 433(1988)23

**Hydroxyphenylacetic acids**

Determination of free and total phenylacetic and *p*- and *m*-hydroxyphenylacetic acids in human urine by LC with fluorimetric detection 414(1987)275

**2-Hydroxy-N-**

**{3-[*m*-(1-piperidinylmethyl)phenoxy]propyl}-acetamide**

Highly sensitive determination of 2-hydroxy-N-{3-[*m*-(1-piperidinylmethyl)phenoxy]propyl}acetamide (desacetyl TZU-0460) in rat plasma by HPLC 413(1987)370

**20 $\alpha$ -Hydroxy-4-pregnen-3-one**

Specific reversed-phase LC method for analysis of steroids in Y-1 adrenal cell cultures 432(1988)302

**Hydroxyprogesterone**

Automated direct assay system for the measurement of sex steroid hormones in serum using HPLC 426(1988)33

**17-Hydroxyprogesterone**

Determination of 17-hydroxyprogesterone in plasma by GC-MS with high-resolution selected-ion monitoring 432(1988)21

**17 $\alpha$ -Hydroxyprogesterone-3-carboxymethylloximaminobutylethylisoluminol**

HPLC of chemiluminogenic labels with sensitive photoplate detection 437(1988)268

**Hydroxyproline**

Determination of hydroxyproline in tissue collagen hydrolysate by derivatization and isocratic reversed-phase HPLC 420(1987)394

**5-Hydroxypropafenone**

HPLC separation and MS identification of propafenone, 5-hydroxypropafenone and N-depropylpropafenone 424(1988)211

**4-Hydroxypropranolol**

Improved HPLC method for the simultaneous determination of propranolol and 4-hydroxypropranolol in plasma with fluorescence detection 422(1987)294

HPLC determination of propranolol and its metabolites in rat serum 431(1988)465

**Hydroxypropyl starch**

Partitioning behavior of erythrocytes in aqueous two-phase systems containing hydroxypropyl starch and polyethylene glycol 441(1988)261

**7- $\beta$ -Hydroxypropyltheophylline**

Simplified description of HPLC separation under overload conditions, based on the Craig distribution model. IV. Comparison of the model with experimental data for samples containing two or more solutes 405(1987)31

**5-Hydroxypyrazinamide**

Identification of 5-hydroxypyrazinamide isolated from urine of subjects given pyrazinamide 416(1987)144

Microdetermination of pyrazinamide and its metabolites (2-pyrazinoic acid, 5-hydroxypyrazinoic acid, 5-hydroxypyrazinamide and pyrazinuric acid) in plasma and urine by LC (in French) 422(1987)217

**5-Hydroxypyrazinoic acid**

Microdetermination of pyrazinamide and its metabolites (2-pyrazinoic acid, 5-hydroxypyrazinoic acid, 5-hydroxypyrazinamide and pyrazinuric acid) in plasma and urine by LC (in French) 422(1987)217

**Hydroxysteroid dehydrogenases**

Simultaneous assay for individual sulphated 3 $\alpha$ - and  $\beta$ -hydroxysteroids in serum using HPLC combined with 3 $\alpha$ - and  $\beta$ -hydroxysteroid dehydrogenases immobilized on one column 432(1988)47

**Hydroxysteroids**

Simultaneous assay for individual sulphated 3 $\alpha$ - and  $\beta$ -hydroxysteroids in serum using HPLC

combined with  $3\alpha$ - and  $\beta$ -hydroxysteroid dehydrogenases immobilized on one column 432(1988)47

HPLC of hydroxysteroids detected with post-column immobilized enzyme reactors 441(1988)81

#### 4-Hydroxyteratolol

Simultaneous quantitative determination of teratolol and its hydroxylated metabolite in human plasma and urine by GC-MS 421(1987)360

#### *p*-Hydroxytriamterene

Rapid method for the determination of the diuretic triamterene and its metabolites in plasma and urine HPLC 413(1987)315

#### 4-Hydroxytriazolam

HPLC determination of triazolam and its metabolites in human urine 422(1987)197

#### 5-Hydroxytryptamine

see Serotonin

#### 5-Hydroxytryptophan

Simultaneous LC determination of seventeen of the major monoamine neurotransmitters, precursors and metabolites. I. Optimization of the mobile phase using factorial designs and a computer program to predict chromatograms 415(1987)261

Simultaneous LC determination of seventeen of the major monoamine neurotransmitters, precursors and metabolites. II. Assessment of human brain and cerebrospinal fluid concentrations 415(1987)275

Determination of plasma 5-hydroxytryptophan, 5-hydroxytryptamine, 5-hydroxyindoleacetic acid, tryptophan and melatonin by HPLC with electrochemical detection 431(1988)170

#### 5-Hydroxytryptophol

Simultaneous LC determination of seventeen of the major monoamine neurotransmitters, precursors and metabolites. I. Optimization of the mobile phase using factorial designs and a computer program to predict chromatograms 415(1987)261

Simultaneous LC determination of seventeen of the major monoamine neurotransmitters, precursors and metabolites. II. Assessment of human brain and cerebrospinal fluid concentrations 415(1987)275

#### 6-Hydroxyvinburnine

Simultaneous determination of vinburnine and 6-hydroxyvinburnine in human plasma by HPLC (in French) 434(1988)157

#### 25-Hydroxyvitamin D<sub>2</sub>

Rapid automated HPLC assay for ercalcidiol and calcidiol (25-hydroxyvitamins D<sub>2</sub> and D<sub>3</sub>) using *trans*-calcidiol as an UV-absorbing internal standard 415(1987)305

Rapid automated HPLC assay for ercalcidiol and calcidiol (25-hydroxyvitamins D<sub>2</sub> and D<sub>3</sub>) using *trans*-calcidiol as an UV-absorbing internal standard 415(1987)305

#### 4-Hydroxy-2,6-xylidine

In vivo metabolism of lidocaine in the rat. Isolation of urinary metabolites as pentafluorobenzoyl derivatives and their identification by combined GC-MS 421(1987)267

#### Hyperglycaemic hormone

Single-step purification of two hyperglycaemic neurohormones from the sinus gland of *Procambarus bowieri*. Comparative peptide mapping by means of HPLC 443(1988)337

#### Hypoxanthine

HPLC analysis of ascorbate-2-phosphate, adenine and hypoxanthine in stored human blood 419(1987)95

Hypoxanthine, xanthine and uridine in body fluids, indicators of ATP depletion (Review) 429(1988)255

## II

#### Ibuprofen

Resolution of  $\alpha$ -arylacetic acid enantiomers on two chiral stationary phases 404(1987)123

Application of an ovomucoid-conjugated column for the optical resolution of some pharmaceutically important compounds 408(1987)316

HPLC determination of pimeprofen and its metabolite ibuprofen in sheep plasma and lymph 414(1987)228

Determination of the enantiomeric composition of ibuprofen in human plasma by HPLC 415(1987)75

Rapid HPLC assay for the simultaneous analysis of non-steroidal anti-inflammatory drugs in plasma 416(1987)293

Method to determine the enantiomers of ibuprofen from human urine by HPLC 428(1988)103

Determination of ibuprofen in human plasma by HPLC 428(1988)388

Determination of ibuprofen and its major metabolites in human urine by HPLC 430(1988)93

Formation of diastereomeric derivatives of 2-arylpropionic acids using L-leucinamide: lack of generality 431(1988)228

#### **Ibuprofen amides**

Stereoelectronic model to explain the resolution of enantiomeric ibuprofen amides on the Pirkle chiral stationary phase 402(1987)179

#### **Ifosfamide**

Determination of ifosfamide in plasma by GC with nitrogen detector 421(1987)377

Rapid method for the determination of ifosfamide and cyclophosphamide in plasma by HPLC with solid-phase extraction 431(1988)450

#### **Igepon T-77**

Application of a novel cation-exchange reagent, Igepon T-77 N-methyl oleoyl taurate), to microbore separations of alumina extracts of catecholamines from cerebrospinal fluid, plasma, urine and brain tissue with amperometric detection 420(1987)241

#### **Imazodan**

HPLC assay of imazodan, methylparaben and propylparaben in imazodan injection 450(1988)407

#### **Imidazobenzodiazepines**

Lipophilicity measurement of benzodiazepine-receptor ligands by reversed-phase LC. Comparison between HPLC and TLC 447(1988)404

#### **Imidazoleacetic acid**

Simultaneous determination of imidazoleacetic acid and N<sup>γ</sup>- and N<sup>ω</sup>-methylimidazoleacetic acids in human urine by HPLC with fluorescence detection 416(1987)63

#### **6-(1-Imidazolylmethyl)-5,6,7,8-tetrahydronaphthalene-2-carboxylic acid**

Determination of the thromboxane synthetase inhibitor 6-(1-imidazolylmethyl)-5,6,7,8-tetrahydronaphthalene-2-carboxylic acid (DP-1904) in human plasma and urine using solid-phase extraction and HPLC 426(1988)111

#### **Imides**

HPLC determination of imide inhibitors of aromatase in biological samples 424(1988)163

#### **Imipenem**

Application of HPLC of some antibiotics in clinical microbiology 427(1988)257

#### **Imipramine**

HPLC determination of imipramine and its metabolites in rat brain 421(1987)412

Effect of blood collection tubes on antidepressant concentrations 423(1987)373

Simultaneous determination of imipramine, desipramine and their deuterium-labelled analogues in biological fluids by capillary GC-MS 426(1988)93

Isolation of basic drugs from plasma using solid-phase extraction with a cyanopropyl-bonded phase 432(1988)209

#### **Immobilines**

Removal of oligomers and *n*-mers from the Immobiline chemicals for isoelectric focusing 402(1987)105

#### **Immobilized dyes**

New strategies for the screening of a large number of immobilized dyes for the purification of enzymes. Application to the purification of enzymes from human haemolysate 449(1988)403

Purification of human 6-phosphogluconate dehydrogenase from human haemolysate with chromatography on an immobilized dye as the essential step and use of automation. Simultaneous purification of lactate dehydrogenase 449(1988)413

#### **Immobilized enzyme reactors**

HPLC determination of glucosides (glucose conjugates) with post-column reaction detection combining immobilized enzyme reactors and luminol chemiluminescence 449(1988)217

#### **Immobilized-metal-ion affinity adsorbents**

Interaction of proteins with immobilized Cu<sup>2+</sup>. Quantitation of adsorption capacity, adsorption

isotherms and equilibrium constants by frontal analysis 403(1987)197

#### **Immobilized-metal-ion affinity chromatography**

High-performance immobilized-metal-ion affinity chromatography of peptides and proteins 443(1988)3

#### **Immobilized pH gradients**

Removal of oligomers and *n*-mers from the Immobiline chemicals for isoelectric focusing 402(1987)105

#### **Immune complexes**

Chromatographic and electrophoretic studies of immune complexes in non-A, non-B hepatitis 426(1988)55

#### **Immunoabsorbents**

Specificity of the immunoabsorbent used for large-scale recovery of interferon  $\alpha$ -2a 411(1987)371

Synthesis of high-capacity immunoaffinity sorbents with oriented immobilized immunoglobulins or their  $F_{ab}$ ' fragments for isolation of proteins 424(1988)243

#### **Immunoabsorptive clean-up**

Determination of digoxin in serum by on-line immunoabsorptive clean-up HPLC separation and fluorescence-reaction detection 433(1988)119

#### **Immunoaffinity partitioning**

Cell separation by immunoaffinity partitioning with polyethylene glycol-modified protein A in aqueous polymer two-phase systems 442(1988)219

#### **Immunoglobulin G**

Size-exclusion HPLC of immune complexes isolated from patients with acute leukaemia. Preliminary observations 421(1987)434

Non-linear elution effects in split-peak chromatography. I. Computer simulations for the cases of irreversible diffusion- and adsorption-limited kinetics 436(1988)111

Molecular orientation of immunoglobulin G at high concentration on an ion-exchange sorbent 443(1988)119

Determination of the molecular size distribution of immunoglobulin G (IgG) in intravenous IgG-albumin formulations by HPLC 444(1988)141

#### **Immunoglobulin G1**

Selective non-adsorption preparative chromatography of bovine immunoglobulin G1 409(1987)305

#### **Immunoglobulin light chain**

HPLC of complex mixtures of cyanogen bromide-produced peptides from different proteins 443(1988)317

#### **Immunoglobulin M**

Comparison of Mono O, Superose-6, and ABx fast protein LC for the purification of IgM monoclonal antibodies 444(1988)153

#### **Immunoglobulins**

Facile preparation of a non-porous strong anion-exchange column for proteins 411(1987)452

Anion-exchange chromatography of normal and monoclonal serum immunoglobulins 416(1987)53

Chromatography of complex protein mixtures (Review) 418(1987)115

Characterization of adsorption on the stationary phase using high-performance immunoaffinity chromatography 423(1987)41

Synthesis of high-capacity immunoaffinity sorbents with oriented immobilized immunoglobulins or their  $F_{ab}$ ' fragments for isolation of proteins 424(1988)243

Chromatographic and electrophoretic studies of immune complexes in non-A, non-B hepatitis 426(1988)55

Purification of ascitic fluid-derived murine monoclonal antibodies by anion-exchange and size-exclusion HPLC 430(1988)329

Purification of specific heterologous  $F(ab)_2$  fragments with DEAE-Zeta-Prep cartridges for ion-exchange chromatography 450(1988)133

#### **Immunological-chromatographic analysis**

Immunological-chromatographic analysis 444(1988)1

#### **Immunsorbents**

Characterization of adsorption on the stationary phase using high-performance immunoaffinity chromatography 423(1987)41

#### **Indenes**

Separation of alkyl-substituted indene enantiomers by chromatography on microcrystalline triacetylcellulose 405(1987)155

**Indenestrol A**

Chromatographic separation and isolation of the enantiomers of diethylstilbestrol metabolites 439(1988)484

**Indenestrol B**

Chromatographic separation and isolation of the enantiomers of diethylstilbestrol metabolites 439(1988)484

**Indian natural resins**

TLC analysis of some Indian natural resins occurring in art objects 439(1988)430

**Indigo Carmine**

Simple method for the analysis of food dyes on reversed-phase thin-layer plates 411(1987)437

**Indirect UV detection**

Single-column anion chromatography with indirect UV detection using pyromellitate buffers as eluents 439(1988)137

**Indocyanine Green**

Comparison of the spectrophotometric and HPLC analysis of Indocyanine Green in estimating systemic clearance in patients with chronic liver disease 417(1987)190

**Indole-3-acetic acid**

Improved method for the determination of indole-3-acetic acid in plant tissue 407(1987)377

Reaction of Fast Blue BB salt and *p*-dimethylaminocinnamaldehyde with indole-3-acetic acid degradation products. Evaluation of the coupling properties of the reagents used in the zymographic screening of indole-3-acetic acid oxidases 411(1987)476

Melatonin and other indoles in rat pineal 428(1988)206

Sensitive determination of deuterated and non-deuterated indole-3-acetic acid and 5-hydroxyindole-3-acetic acid by combined capillary GC negative-ion chemical ionization MS 428(1988)209

Effects of long-term storage on the concentrations of the unconjugated acidic metabolites of the trace amines, indoleamines and catecholamines 433(1988)23

Microscale isolation technique for quantitative GC-MS analysis of indole-3-acetic acid from cherry (*Prunus cerasus* L.) 442(1988)301

**Indole alkaloids**

Determination of *Catharanthus* alkaloids by reversed-phase HPLC 410(1987)488

**Indolealkylamines**

GC-MS characterisation of the psychotomimetic indolealkylamines and their *in vivo* metabolites 422(1987)13

**Indoleamines**

Effects of long-term storage on the concentrations of the unconjugated acidic metabolites of the trace amines, indoleamines and catecholamines 433(1988)23

**Indoles**

Melatonin and other indoles in rat pineal 428(1988)206

Chromatography of urinary indole derivatives (Review) 429(1988)59

Optimized determination of indole derivatives by HPLC with selective detection 446(1988)315

**Indolidan**

LC determination of indolidan and its dehydro metabolite in plasma. Use of the Varian AASP for solid-phase extraction 431(1988)103

**Indomethacin**

Determination of association constants of  $\beta$ -cyclodextrin and  $\beta$ -cyclodextrin-bearing polymers with drugs by HPLC 409(1987)61

Rapid HPLC assay for the simultaneous analysis of non-steroidal anti-inflammatory drugs in plasma 416(1987)293

**Indoprofen**

Assay method for product formation in *in vitro* enzyme kinetic studies of uridine diphosphate glucuronyltransferases: 2-arylpropionic acid enantiomers 430(1988)368

**Indoxyl sulphate**

Rapid assay for indoxyl sulphate in uremic serum by internal-surface reversed-phase HPLC 431(1988)418

**Inductive adsorption**

Inductive adsorption as a new approach to amine enantiomer resolution 404(1987)87

**Inductively coupled plasma atomic emission spectrometry**

Analysis of platinum species originating from *cis*-diamminedichloroplatinum(II) (cisplatin) in



human and rat plasma by HPLC with on-line inductively coupled plasma atomic emission spectrometric detection 407(1987)253

Interfacing of inductively coupled plasma-atomic emission spectrometry to SFC for elemental detection 411(1987)213

### **Injection conditions**

Analysis of the information in a preparative chromatogram for further optimization of the operating conditions 450(1988)13

### **Injection techniques**

Investigation of pre-set and actual splitting ratios in split injection capillary GLC 404(1987)340

Analysis of fatty acid methyl esters with high accuracy and reliability. VI. Rapid analysis by split injection capillary GLC 407(1987)231

Design of an inexpensive on-column injector for capillary GC 407(1987)305

Sampling and injection system for the analysis of metalorganic chemicals by GC-MS. Some initial results 411(1987)445

HPLC drug analysis by direct injection of whole blood samples. I. Determination of moderately hydrophobic drugs incorporated into blood corpuscles 423(1987)147

HPLC drug analysis by direct injection of whole blood samples. II. Determination of hydrophilic drugs 423(1987)155

HPLC drug analysis by direct injection of whole blood samples. III. Determination of hydrophobic drugs adsorbed on blood cell membranes 423(1987)163

Mitoxantron determination using HPLC: improved sensitivity by loop-column injection for dual-dose pharmacokinetic studies 425(1988)429

Comparison of the performances of hot and cold sample introduction with a programmed-temperature vaporizer in the split and splitless modes 438(1988)243

Factors affecting sample transfer from microlitre syringes 441(1988)367

### **Injectors**

Design of an inexpensive on-column injector for capillary GC 407(1987)305

Checking of capacity of a splitless injector — a simple test 448(1988)387

### **Inorganic anions**

Ion chromatography of inorganic anions on a dynamically modified polystyrene-divinylbenzene packing material and its application to anion screening by sequential UV absorbance and electrochemical detection 402(1987)115

Origin of system peaks in single-column ion chromatography of inorganic anions using high pH borate-gluconate buffers and conductivity detection 404(1987)67

Derivatization of inorganic anions with pentafluorobenzyl methanesulphonate for GC 404(1987)175

Adsorption chromatography on cellulose. I. Salting-out chromatography of organic compounds 406(1987)389

Contamination of aqueous eluents due to corrosion of stainless-steel chromatographic components 407(1987)133

Determination of oxidizable inorganic anions by HPLC with fluorescence detection and application to the determination of salivary nitrite and thiocyanate and serum thiocyanate 424(1988)29

Simultaneous conductivity and amperometric detector with a single cell for microbore LC 436(1988)413

Polymeric pentafluorobenzylating reagent for GC of inorganic anions with electron-capture detection 438(1988)253

Ion-pair HPLC of inorganic anions with photometric, conductometric and amperometric detection 439(1988)363

Low-cost rapid separation and determination of ionic species by single-column ion chromatography 445(1988)183

Indirect photometric detection of inorganic anions in micro HPLC with permanently coated columns 447(1988)221

### **Inorganic cations**

Contamination of aqueous eluents due to corrosion of stainless-steel chromatographic components 407(1987)133

### **Inorganic ions**

Coordinative interactions in the separation of insulin and derivatives by HPLC 408(1987)43

Capillary isotachopheresis in the study of the absorption of inorganic ions by roots of intact plants 438(1988)454

**Inorganic salts**

Distribution of inorganic salts between the coexisting phases of aqueous polymer two-phase systems. Interrelationship between the ionic and polymer composition of the phases 439(1988)267

**Inositol**

Simultaneous GLC determination of aldoses and alduronic acids 408(1987)245

**Inositol phosphates**

Ion chromatographic determination of sugar phosphates in physiological samples 439(1988)83

**Insulin**

Coordinative interactions in the separation of insulin and derivatives by HPLC 408(1987)43

Recovery of polypeptides after reversed-phase HPLC 408(1987)191

Separation of protein hormones (Review) 418(1987)245

Chromatographic analysis of the trinitrophenyl derivatives of insulin 427(1988)239

**Insulin-like growth factor 1**

Reduction studies on bacterial recombinant somatomedin C/insulin-like growth factor-1 443(1988)183

**Insulin-like growth factor II**

Application of preparative reversed-phase HPLC to the isolation of insulin-like growth factor II from human serum 421(1987)61

**Insulin-like growth factor-binding protein**

Method for purification of an insulin-like growth factor-binding protein produced by human HEP G2 hepatoma cells 420(1987)163

**Integration**

Amino acid analysis. III. Nuances of integration 436(1988)474

**Integrators**

Amino acid analysis. I. Nuances of integration 404(1987)321

Integrators as a source of inaccurate results 404(1987)326

Amino acid analysis. II. Nuances of integration 410(1987)176

**Interfaces, GC-MS**

Open split interfaces in capillary GC-MS. Yield and quantitative aspects 450(1988)388

**Interfaces, LC-MS**

Improvement of the interface for coupling of fast atom bombardment MS and micro HPLC 435(1988)482

**Interfacial surface areas**

Evaluation of interfacial surface areas in ultra-low-loaded GC columns 446(1988)221

**Interfacial tension**

Partitioning behaviour of rat bone marrow cells in aqueous two-phase systems. Dependence of cell partition on the interfacial tension and electrical potential difference between the phases 403(1987)131

**Interferon**

Specificity of the immunoadsorbent used for large-scale recovery of interferon  $\alpha$ -2a 411(1987)371

High-performance SEC of recombinant derived proteins and aggregated species 436(1988)289

**Interleukin-2**

High-performance SEC of recombinant derived proteins and aggregated species 436(1988)289

Application of chromatographic methods to the analysis of macrophage factors induced by *Nocardia opaca* cell walls 440(1988)119

**Interleukin-2 receptor**

Medium-scale ligand-affinity purification of two soluble forms of human interleukin-2 receptor 431(1988)55

**Internal-surface reversed-phase silica**

Retention properties of internal-surface reversed-phase silica packing and recovery of drugs from human plasma 420(1987)297

**Internal standards**

Internal standards for quantitative GC of individual bile acids after group separation of bile acids in urine 422(1987)25

**Intramolecular interconversion**

Capillary column GC method for the study of dynamic intramolecular interconversion behaviour 447(1988)29

**Intraparticle diffusion**

Theoretical analysis of the asymmetry in chromatographic peaks 404(1987)1

**Inulin fructotransferase II**

Activity staining of inulin fructotransferase II on polyacrylamide gels 445(1988)282

**Inverse gas chromatography**

Inverse GC study of polyether alcohols (in German) 410(1987)267

**Iodide**

Derivatization of inorganic anions with pentafluorobenzyl methanesulphonate for GC 404(1987)175

Excretion of iodide in 24-h urine as determined by ion-pair reversed-phase LC with electrochemical detection 431(1988)309

Polymeric pentafluorobenzylating reagent for GC of inorganic anions with electron-capture detection 438(1988)253

Determination of iodide and bromide by ion chromatography with post-column reaction detection 439(1988)129

**Iodine**

Simple method for the determination of hormonal iodine and exogenous iodine from X-ray contrast media drugs 434(1988)308

Determination of trace amounts of iodine in sedimentary rocks by ion chromatography with amperometric detection 436(1988)107

**Iododeoxyuridine**

Technique for detection of DNA nucleobases by reversed-phase HPLC optimized for quantitative determination of thymidine substitution by iododeoxyuridine 417(1987)57

**Ion chromatography**

A look at contemporary ion chromatography 439(1988)3

**Ion-exchange amino acid analyser kit**

Construction of an ion-exchange amino acid analyser kit for use with HPLC apparatus 431(1988)285

**Ion exchangers**

HPLC analysis of oligosaccharides. I. Separation on an ion-exchange stationary phase of low cross-linking 440(1988)281

Ion-exchange derivatives of Spheron. VI. Strongly basic derivatives 440(1988)287

HPLC of plant-derived oligosaccharides on a new cation-exchange resin stationary phase: HPX-22H 441(1988)382

Phase ratio determination in an ion-exchange column having pores partially accessible to proteins 442(1988)15

Peptide characterization with a sulfoethyl aspartamide column 443(1988)63

Synthesis of a non-porous, polystyrene-based strong anion-exchange packing material and its application to fast HPLC of proteins 443(1988)73

Cation-exchange chromatography of peptides on poly(2-sulfoethyl aspartamide)-silica 443(1988)85

Causes for the loss of resolution in an ion chromatography resin 445(1988)211

Effect of the content of divinylbenzene in ion-exchange resins on the chromatographic separation of  $\alpha$ -cyclodextrin and glucose 448(1988)241

Purification of specific heterologous F(ab)<sub>2</sub> fragments with DEAE-Zeta-Prep cartridges for ion-exchange chromatography 450(1988)133

**Ion-exclusion phases**

New ion-exclusion phase for organic acids 439(1988)71

**Ionic species interface**

Isotachopheresis: analysis and computation of the structure of the ionic species interface 405(1987)77

**Ion-interaction reagents**

Studies on sample preconcentration in ion chromatography. VI. Use of concentrator columns with "permanently" bound ion-interaction reagents 407(1987)121

**Ionization chamber**

Mechanism governing the sensitivity of a three-electrode ionization chamber 446(1988)163

**Ion-pair extraction**

Direct, sensitive and selective detection of free fatty acids by HPLC with post-column ion-pair extraction and absorbance detection 445(1988)189

**Ion-pair formation**

Paper electrophoretic study of ion-pair formation. XVII. Influence of the enantiomeric purity of the sodium tartrate used as electrolyte 440(1988)361

**Ion-pair formation constants**

Complex-forming equilibria in isotachopheresis. VII. Evaluation of the ion-pair formation constants of phosphorus oxo acids with histidine and assessment of their separability 410(1987)279

**Ion-pairing reagents**

Mechanism of selectivity in ion-pair HPLC of aminoglycoside antibiotics using perfluorinated pairing ions 404(1987)53

Use of mixed perfluorinated ion-pairing agents as solvents in ion-pair HPLC for the preparative purification of aminoglycoside antibiotics 407(1987)109

Supercritical fluid chromatography with ion-pairing modifiers. Separation of enantiomeric 1,2-aminoalcohols as diastereomeric ion pairs 447(1988)287

Perfluorinated acids as ion-pairing agents in the determination of monoamine transmitters and some prominent metabolites in rat brain by HPLC with amperometric detection 449(1988)191

**Ipsapirone**

Simultaneous determination of buspirone, gepirone, ipsapirone and their common metabolite 1-(2-pyrimidinyl)piperazine in rat plasma and brain by HPLC 431(1988)477

**Irgasan DP300**

Formation of polychlorinated dibenzo-*p*-dioxins upon combustion of commercial textile products containing 2,4,4'-trichloro-2'-hydroxydiphenyl ether (Irgasan DP300) 442(1988)289

**Iridoid glycosides**

Application of various planar chromatographic techniques for the separation of iridoid glycosides from *Veronica officinalis* 404(1987)365

**Iron**

Comparison of HPLC and atomic spectrometric methods for the determination of Fe(III) and Al(III) in soil and clay samples 449(1988)241

**Iron clusters**

Behaviour of tri- and tetranuclear iron and nickel clusters in HPLC 447(1988)187

**Iron complexes**

Mobility of Fe(III)-phenolic acid complexes in TLC 402(1987)358

**Iron(III) dibutyldithiocarbamate**

LC of dibutyldithiocarbamate degradation products 404(1987)359

**Iso- $\alpha$ -acids**

Avoidance of trace metal activity with micro-LC in the analysis of hop and beer bitter acids 407(1987)159

**Isobutanol**

Rapid HPLC method for determination of ethanol and fusel oil in the alcoholic beverage industry 447(1988)443

**2-(trans-4-Isobutylcyclohexyl)-2-oxoethyl benzenesulphonate**

Determination of a new hypolipidemic agent, 2-(trans-4-isobutylcyclohexyl)-2-oxoethyl benzenesulphonate, and its metabolite by HPLC and GC-mass fragmentography in human biological materials 433(1988)282

**Isochromans**

Chromatographic behaviour of diastereoisomers. IX. Application of a microcomputer program to the TLC separation of some diastereoisomers on silica 438(1988)219

**Isocitrate**

HPLC measurement of selected blood citric acid cycle intermediates 425(1988)47

**Isocyanates**

Trace analysis of airborne aromatic isocyanates and related aminoisocyanates and diamines using HPLC with UV and electrochemical detection 435(1988)469

**Iso-1-cytochrome *c***

Characterization of wild-type and mutant forms of yeast iso-1-cytochrome *c* by reversed-phase HPLC tryptic peptide mapping 439(1988)408

**Isodesmosine**

HPLC determination of desmosine and isodesmosine in tissues and its application to studies of alteration of elastin induced by atherosclerosis 422(1987)53

**Isoelectric focusing**

Removal of oligomers and *n*-mers from the Immobiline chemicals for isoelectric focusing 402(1987)105

Carrier-free zone electrophoresis, displacement electrophoresis and isoelectric focusing in a high-performance electrophoresis apparatus 403(1987)47

Cysteine adducts of human haemoglobin measured by isoelectric focusing in polyacrylamide gels with a non-linear pH gradient 420(1987)35

Purification of intact concanavalin A in tetramers by isoelectric focusing 423(1987)319

Experimental and theoretical dynamics of

isoelectric focusing. II. Elucidation of the impact of the electrode assembly 436(1988)191

Capillary isoelectric focusing using an LKB 2127 Tachophor isotachophoretic analyzer 436(1988)309

Ionic binding properties of carrier ampholytes 437(1988)147

Buffer isoelectric focusing revisited 440(1988)367

#### **Isoenzymes**

Automated separation and measurement of urinary isoenzymes and protein by ion-exchange LC 414(1987)47

Chromatography of complex protein mixtures (Review) 418(1987)115

#### **Isoetretin**

Determination of the aromatic retinoids (etretin and isoetretin) in biological fluids by HPLC 421(1987)177

#### **Isoferulate esters**

Novel isoferulate esters identified by GC-MS in bud exudate of *Populus nigra* 448(1988)284

#### **Isoflavonoids**

Reversed-phase HPLC of *Cajanus cajan* phytoalexins 409(1987)355

#### **Isoniazid**

Determination of the isoniazid metabolite monoacetylhydrazine in urine by HPLC 415(1987)188

Analysis of isoniazid, acetylhydrazine and [<sup>15</sup>N<sub>2</sub>]acetylhydrazine in serum by capillary GC-ammonia chemical ionization MS 420(1987)171

Determination of isoniazid in plasma by LC 425(1988)441

#### **Isophorone**

GLC analysis of isophorone byproducts 446(1988)23

#### **Isoimaric acid**

GC analysis of underivatized resin acids 448(1988)428

#### **Isoprene**

Efficiency of adsorption-desorption of volatile compounds. Application to isoprene on Tenax GC 437(1988)139

#### **Isoprenylated genistein**

Reversed-phase HPLC of *Cajanus cajan* phytoalexins 409(1987)355

#### **Isopreterenol**

Determination of  $\beta$ -phenylalkylamines in pharmaceutical preparations by analytical capillary isotachopheresis 405(1987)389

#### **Isoproc carb**

HPLC determination of aryl N-methylcarbamate residues using post-column hydrolysis electrochemical detection 442(1988)333

#### **Isopropanol**

Simultaneous determination of methanol, ethanol, acetone, isopropanol and ethylene glycol in plasma by GC 414(1987)248

Static headspace GC method for the determination of residual solvents in vigabatrin drug substance 438(1988)103

#### **Isopropylbenzene**

Heavy alkylated benzene stationary phase for the GLC separation of ethylbenzene, xylenes, styrene, iso- and *n*-propylbenzene 409(1987)371

#### **1,2-Isopropylidene glycerol**

Enantioselective capillary GC of 1,2-isopropylidene glycerol and O-alkylated glycerol derivatives 410(1987)458

#### **Isopropyl-*p*-iodoamphetamine**

Highly sensitive method for determination of isopropyl-*p*-iodoamphetamine by GC with electron-capture detection 425(1988)402

#### **Isopropyl methylphosphonic acid**

Analysis of alkyl methylphosphonic acids in aqueous matrices by ion-pair reversed-phase ion chromatography 407(1987)399

#### **(5*S*)-3-Isopropyl-5-*p*-toluenesulphonyloxymethylloxazolidin-2-one**

Direct determination of the enantiomeric purity of (5*S*)-3-isopropyl-5-*p*-toluenesulphonyloxymethylloxazolidin-2-one on a cellulose-based chiral stationary phase. In-process control of a chiral intermediate used in the synthesis of enantiomerically pure  $\beta$ -blocking agents 411(1987)461

#### **Isoquinoline alkaloids**

Determination of alkaloids of *Fumaria parviflora* and *Fumaria capreolata* by HPLC and capillary isotachopheresis 445(1988)258

**Isoquinolines**

Assay of five corynoline-type alkaloids in *Corydalis bungeana* by reversed-phase HPLC 408(1987)426

**Isorhamnetin**

Rapid separation of flavonoids by analytical high-speed counter-current chromatography 445(1988)199

**Isosafrole**

Determination of  $\beta$ -asarone, safrole, isosafrole and anethole in alcoholic drinks by HPLC 404(1987)273

**Isosakuranetin**

Direct enantiomeric separation of racemic flavanones by HPLC using cellulose triacetate as a chiral stationary phase 441(1988)417

**Isotachic splitter**

On-line microbore HPLC capillary GC-MS. I. Development of isotachic eluent splitters 409(1987)205

**Isotachochromatography**

Feasibility of isotachochromatography as a method for the preparative separation of weak acids and weak bases. I. Theoretical considerations 436(1988)173

Feasibility of isotachochromatography as a method for the preparative separation of weak acids and weak bases. II. Experimental verification 436(1988)181

**Isotachophoresis**

Determination of the electrophoretic order of 26 nucleotides by isotachophoresis 403(1987)336

Isotachophoresis: analysis and computation of the structure of the ionic species interface 405(1987)77

Use of cyclodextrins in isotachophoresis. I. Effect of cyclodextrin on the isotachophoretic separation of related penicillins 405(1987)379

Determination of minoxidil in pharmaceutical forms by capillary isotachophoresis 405(1987)385

Determination of  $\beta$ -phenylalkylamines in pharmaceutical preparations by analytical capillary isotachophoresis 405(1987)389

Capillary zone electrophoresis and isotachophoresis as alternatives to chromatographic methods for purity control of synthetic peptides 407(1987)363

Isotachophoretic determination of mobility and  $pK_a$  by means of computer simulation. VI. Mobility and  $pK_a$  of sixty anions in methanol 408(1987)27

Complex-forming equilibria in isotachophoresis. VII. Evaluation of the ion-pair formation constants of phosphorus oxo acids with histidine and assessment of their separability 410(1987)279

Use of cyclodextrins in isotachophoresis. II.  $\alpha$ -,  $\beta$ - and  $\gamma$ -cyclodextrins as leading electrolyte additives for the separation of *ortho*-, *meta*- and *para*-substituted halogenobenzoic acids 411(1987)153

Determination of ionic metabolites from ethylene glycol in human blood by isotachophoresis 416(1987)111

Detection of fibrinogen heterogeneity by capillary isotachophoresis 419(1987)329

Isotachophoretic analysis for the determination of urinary  $\gamma$ -carboxyglutamic acid 419(1987)334

Analysis of the aqueous phase of human cervical mucus by reversed-phase HPLC and capillary isotachophoresis 431(1988)317

Purification of monoclonal antibodies against the low-density lipoprotein receptor by preparative isotachophoresis 431(1988)327

Use of cyclodextrins in isotachophoresis. III. Purity control of naftidrofuryl hydrogenoxalate and some of its synthesis intermediates 435(1988)496

Use of cyclodextrins in isotachophoresis. IV. The influence of cyclodextrins on the chiral resolution of ephedrine alkaloid enantiomers 438(1988)211

Capillary isotachophoresis in the study of the absorption of inorganic ions by roots of intact plants 438(1988)454

Use of cyclodextrins in isotachophoresis. V. The separation of ketotifen and its polar intermediate enantiomers 439(1988)386

Isotachophoretic determination of the cationic constituents in mixtures of raw materials for glass manufacturing 439(1988)423

Host-guest complexation in capillary isotachophoresis. I.  $\alpha$ -,  $\beta$ - and  $\gamma$ -cyclodextrins as complexing agents for the resolution of substituted benzoic acid isomers 442(1988)371

Cationic capillary isotachophoresis of proteins 445(1988)417

Capillary tube isotachophoretic separation of heavy metal ions using complex-forming equilibria between cyanide as terminating ion and the metal ions 447(1988)383

Isotachopheresis of quaternary 4,4'-bipyridylum salts. Analytical control of synthesis and purification procedures 449(1988)299

#### **Isotachophoretic analyzer**

Capillary isoelectric focusing using an LKB 2127 Tachophor isotachophoretic analyzer 436(1988)309

#### **Isothermal fixed-bed adsorption**

Free energy consumption at self-sharpening concentration fronts in isothermal fixed-bed adsorption. Theory for isotherms of arbitrary shape and applications in displacement chromatography 409(1987)1

#### **Isotopic fractionation**

HPLC studies of pH dependence in the isotopic fractionation of deuterated benzazepines 407(1987)37

#### **Isotretinoin**

Quantitative analysis of retinoids in biological fluids by HPLC using column switching. I. Determination of isotretinoin and tretinoin and their 4-oxo metabolites in plasma 424(1988)303

#### **Isoxazolyl penicillins**

Optimization of HPLC analysis for isoxazolyl penicillins using factorial design 425(1988)331

#### **E-Isozeatin**

E-Isozeatin and *cis*-norzeatin riboside as internal standards for cytokinin analyses 436(1988)93

#### **Isozeaxanthin**

Separation of carotenol fatty acid esters by HPLC 449(1988)119

#### **Isradipine**

Assay of isradipine and of its major metabolites in biological fluids by capillary GC and chemical ionization MS 428(1988)61

#### **Itraconazole**

Determination of itraconazole in plasma and animal tissues by HPLC 413(1987)332

Assay of itraconazole in leukemic patient plasma by reversed-phase small-bore LC 432(1988)388

#### **Ivermectin**

Determination of 22,23-dihydroavermectin B<sub>1a</sub>

in dog plasma using solid-phase extraction and HPLC 413(1987)326

Determination of ivermectin in human plasma and milk by HPLC with fluorescence detection 416(1987)196

Application of two-dimensional HPLC in metabolism studies of ivermectin 433(1988)217

## **J**

#### **Janus Green**

Use of the dye Janus Green to facilitate quantitative analysis of phosphoinositides 424(1988)445

#### **JP-5**

Determination of phenolic antioxidants in JP-5 jet fuels by GC-mass selective detection 449(1988)440

## **K**

#### **Kallikrein**

HPLC method for the determination of the esterase activity of subtilisin and kallikrein 441(1988)431

#### **Kalman filter**

Real-time quantitative analysis of unresolved LC peaks with the use of the Kalman filter 407(1987)59

Totally automated robotic system for LC analysis of solid dosage formulations with the aid of a reduced Kalman filter 438(1988)319

#### **Kalman filter analysis**

Resolution of overlapped chromatograms by the use of a Kalman filter. III. Effects of overlapping

and outstripping of a solvent peak on the analysis 411(1987)95

### **Keratan sulphate**

Extracellular mammalian polysaccharides: glycosaminoglycans and proteoglycans (Review) 418(1987)187

### **Kerateins**

Chromatographic and electrophoretic investigations of the properties of unprotected low-sulphur wool kerateins 405(1987)365

### **Kerogen**

GC determination of C<sub>1</sub>-C<sub>5</sub> low-molecular-weight organic acids in alkaline permanganate oxidation products of humic substances 403(1987)225

### **Kerosene**

GC determination for forensic purposes of petroleum fuel inhaled just before fatal burning 424(1988)49

### **Kestoses**

Isolation of kestoses and nystose from enzyme digests by HPLC 408(1987)393

### **Keto acids**

Profiling of oxalic acid and  $\alpha$ -keto acids in blood and urine by LC with electrochemical detection at a chemically modified electrode 414(1987)161

HPLC determination of glyoxylic acid and other carbonyl compounds in urine 417(1987)253

GC of branched-chain keto acids analysed as their silylated oxime derivatives 423(1987)270

Determination of  $\alpha$ -keto acids in rat hindquarter muscle by HPLC with fluorescence detection 424(1988)361

Simultaneous determination of C<sub>2</sub>-C<sub>22</sub> non-esterified fatty acids and other metabolically relevant carboxylic acids in biological material by GC of their benzyl esters 425(1988)257

HPLC determination of  $\alpha$ -keto acids in human serum and urine using 1,2-diamino-4,5-methylenedioxybenzene as a precolumn fluorescence derivatization reagent 430(1988)223

Identification of aliphatic keto carboxylic acids in marine aerosols using capillary GC-MS 438(1988)299

### **Keto bile acids**

Separation of keto and keto-hydroxy bile acid isomers by HPLC 438(1988)423

### **$\alpha$ -Ketocaproate**

GC of branched-chain keto acids analysed as their silylated oxime derivatives 423(1987)270

Determination of  $\alpha$ -keto acids in rat hindquarter muscle by HPLC with fluorescence detection 424(1988)361

### **$\beta$ -Ketoenamides**

GC of metal  $\beta$ -diketonates and their analogues (Review) 411(1987)1

### **$\alpha$ -Ketoglutarate**

HPLC measurement of selected blood citric acid cycle intermediates 425(1988)47

### **Keto-hydroxy bile acids**

Separation of keto and keto-hydroxy bile acid isomers by HPLC 438(1988)423

### **$\alpha$ -Ketoisocaproate**

GC of branched-chain keto acids analysed as their silylated oxime derivatives 423(1987)270

Determination of  $\alpha$ -keto acids in rat hindquarter muscle by HPLC with fluorescence detection 424(1988)361

### **$\alpha$ -Ketoisovalerate**

GC of branched-chain keto acids analysed as their silylated oxime derivatives 423(1987)270

Determination of  $\alpha$ -keto acids in rat hindquarter muscle by HPLC with fluorescence detection 424(1988)361

### **$\alpha$ -Keto- $\beta$ -methylvalerate**

GC of branched-chain keto acids analysed as their silylated oxime derivatives 423(1987)270

Determination of  $\alpha$ -keto acids in rat hindquarter muscle by HPLC with fluorescence detection 424(1988)361

### **Ketones**

Potential solid supports for GC prepared on the basis of certain rocks 406(1987)145

Comparison of volatile components between fresh and rotten mussels by GC-MS 409(1987)251

Organo-modified smectite for the GC separation of various compounds 410(1987)121

HPLC determination of glyoxylic acid and other carbonyl compounds in urine 417(1987)253

Silylated N,O-ketals from the reaction of ketones with N-trimethylsilylimidazole 419(1987)271

O-(2,3,4,5,6-Pentafluorobenzyl)oxime-



trimethylsilyl ester derivatives for quantitative GC and GC-MS studies of aldehydes, ketones and oxoacids 421(1987)336

### **Ketoprofen**

Application of an ovomucoid-conjugated column for the optical resolution of some pharmaceutically important compounds 408(1987)316

Determination of the enantiomers of ketoprofen in blood plasma by ion-pair extraction and HPLC of leucinamide derivatives 414(1987)465

Rapid HPLC assay for the simultaneous analysis of non-steroidal anti-inflammatory drugs in plasma 416(1987)293

HPLC assay of ketoprofen enantiomers in human plasma and urine 416(1987)388

Assay method for product formation in *in vitro* enzyme kinetic studies of uridine diphosphate glucuronyltransferases: 2-arylpropionic acid enantiomers 430(1988)368

HPLC determination of ketoprofen and naproxen in rat plasma 431(1988)444

### **6-Ketoprostaglandin F<sub>1α</sub>**

Quantitative analysis of 6-keto-prostaglandin F<sub>1α</sub> using immunoaffinity purification and GC-MS 425(1988)1

### **15-Ketoprostaglandin F<sub>2α</sub>**

Dimethylisopropylsilyl ether derivatives in GC-MS of some prostanoids. Application to the microdetermination of 15-ketoprostaglandin F<sub>2α</sub> formed by enzymatic conversion of PGF<sub>2α</sub> *in vitro* 424(1988)1

### **Ketoses**

Degradation of ketoses during aqueous HPLC on lead-form cation-exchange resins 437(1988)387

### **Ketotifen**

Use of cyclodextrins in isotachopheresis. V. The separation of ketotifen and its polar intermediate enantiomers 439(1988)386

### **Kinetic studies**

Kinetic studies of fast equilibrium by means of HPLC. XVI. Separation of tautomers of β-diketones and β-diketoamides 411(1987)466

### **Kininases**

Determination of kininase I and kininase II activities in human urine by HPLC 414(1987)423

### **Kinins**

Determination of kininase I and kininase II activities in human urine by HPLC 414(1987)423

### ***Klebsiella pneumoniae***

Studies of metabolites in diarrheal stool specimens positive for *Klebsiella*, *Serratia*, and *Proteus* spp. by frequency-pulsed electron-capture GC 430(1988)209

### **Krebs cycle acids**

New method of purification of perchloric acid extracts for quantitative GC of some Krebs cycle acids 424(1988)357

HPLC measurement of selected blood citric acid cycle intermediates 425(1988)47

### **Kromasil C<sub>18</sub>**

Preparative HPLC of peptides on a new reversed-phase packing material, Kromasil C<sub>18</sub> 450(1988)71

### **Kynurenic acid**

High-voltage electrophoretic separation of tryptophan metabolites of the kynurenic pathway 420(1987)263

Fluorimetric micro-determination of kynurenic acid, an endogenous blocker of neurotoxicity, by HPLC 430(1988)376

### **Kynurenine**

Semi-micro quantitative assay for tryptophan oxygenase by HPLC 419(1987)296

High-voltage electrophoretic separation of tryptophan metabolites of the kynurenic pathway 420(1987)263

Preparative chiral separation in an aqueous two-phase system by a few counter-current extractions 450(1988)277

## **L**

### **L-364,718**

Quantitative determination of L-364,718, a novel cholecystokinin antagonist, by solid-phase extraction and HPLC 437(1988)442

**Labetalol**

Optimization of fluorescence detection for the determination of labetalol in plasma by HPLC 446(1988)323

Planar chromatographic separation of enantiomers and diastereomers with cyclodextrin mobile phase additives 448(1988)345

 **$\alpha$ -Lactalbumin**

Influence of pore and particle size on the frontal uptake of proteins. Implications for preparative anion-exchange chromatography 409(1987)111

 **$\beta$ -Lactamase inhibitor**

HPLC determination of a new  $\beta$ -lactamase inhibitor and its metabolite in combination therapy with piperacillin in biological materials 431(1988)87

 **$\beta$ -Lactamases**

Use of chromatofocusing for separation of  $\beta$ -lactamases. VIII. Analytical chromatofocusing of chromosomal cephalosporinases from four *Klebsiella* strains 403(1987)217

Fractionation and characterization of two  $\beta$ -lactamases in *Citrobacter diversus* ULA-27 strain by chromatofocusing 403(1987)366

 **$\beta$ -Lactams**

Determination of  $\beta$ -lactams and their biosynthetic intermediates in fermentation media by pre-column derivatization followed by fluorescence detection 424(1988)325

**Lactate**

Sensitive determination of D-lactic acid in biological samples by HPLC 431(1988)239

Ion chromatographic determination of sugar phosphates in physiological samples 439(1988)83

**Lactate dehydrogenase**

Retention mechanism of lactate dehydrogenase in anion-exchange chromatography 406(1987)237

Matrix evaluation for preparative high-performance affinity chromatography 407(1987)179

HPLC purification of the H4 and MH3 isoenzymes of lactate dehydrogenase from beef heart 408(1987)372

Effect of polymer structure on affinity partitioning of lactate dehydrogenase in polymer-water two-phase systems 411(1987)161

Automated separation and measurement of

urinary isoenzymes and protein by ion-exchange LC 414(1987)47

Lactate dehydrogenase isoenzymes (Review) 429(1988)373

Purification of human 6-phosphogluconate dehydrogenase from human haemolysate with chromatography on an immobilized dye as the essential step and use of automation. Simultaneous purification of lactate dehydrogenase 449(1988)413

**Lactic acid**

Determination of ionic metabolites from ethylene glycol in human blood by isotachopheresis 416(1987)111

 **$\beta$ -Lactoglobulin**

Tandem separation schemes for preparative HPLC of proteins 443(1988)31

**Lactose**

HPLC separation of some mono- and disaccharides with detection by a post-column enzyme reactor and a chemically modified electrode 408(1987)157

**Lactosylceramide**

TLC of neutral glycosphingolipids: an improved simple method for the resolution of GlcCer, GalCer, LacCer and Ga<sub>2</sub>Cer 426(1988)188

**Lactotransferrin**

Characterization of buffalo lactotransferrin by polyacrylamide gel electrophoresis in 6 M urea 434(1988)228

**Ladroside**

Application of various planar chromatographic techniques for the separation of iridoid glycosides from *Veronica officinalis* 404(1987)365

**Laevulinic acid**

Extractive pentafluorobenzoylation of formic, acetic, levulinic, benzoic and phthalic acids, studied by LC and dual-oven capillary GC 447(1988)329

**Lanthanides**

Separation of some rare earths and transplutonium elements by HPLC on Ostion cation exchanger 435(1988)249

Lanthanide ions as luminescent chromophores for the LC detection of polynucleotides and nucleic acids 436(1988)299

**Lanthionine ketimine**

Specific method for the determination of 1,4-thiomorpholine-3,5-dicarboxylic acid and its unsaturated analogue lanthionine ketimine 426(1988)183

**Large molecules**

HPLC at elevated temperatures: examination of conditions for the rapid separation of large molecules 435(1988)1

**Laser detectors**

Use of laser detectors in capillary LC (Review) 434(1988)337

**Laser fluorescence detector**

Use of laser detectors in capillary LC (Review) 434(1988)337

**Laudanosine**

Separation of eight alkaloids and meconic acid and quantitation of five principal alkaloids in gum opium by gradient reversed-phase HPLC 436(1988)455

**LC**

Preparative resolution of the enantiomers of the  $\beta$ -blocking drug atenolol by chiral derivatization and HPLC 431(1988)222

**Lecithin**

Phospholipids in amniotic fluid with special reference to the lecithin; Sphingomyelin ratio (Review) 429(1988)359

**Lectin-peroxidase conjugates**

*In situ* visualization of glycosphingolipids on thin-layer chromatograms using lectin-peroxidase conjugates 411(1987)393

**Lectins**

Purification of human erythrocytes specific lectins from rice bean, *Phaseolus calcaratus* syn. *Vigna umbellata*, by HPLC 431(1988)37

**Leishmania-derived antigens**

Inoculation of BALB/c mice against *Leishmania major* infection with *Leishmania*-derived antigens isolated by gel filtration 440(1988)459

**Lemon peel oil**

Simple microscale supercritical fluid extraction system and its application to GC-MS of lemon peel oil 442(1988)121

**Leucinamide**

Determination of the enantiomers of ketoprofen

in blood plasma by ion-pair extraction and HPLC of leucinamide derivatives 414(1987)465

Formation of diastereomeric derivatives of 2-arylpropionic acids using L-leucinamide 423(1987)334

Formation of diastereomeric derivatives of 2-arylpropionic acids using L-leucinamide: lack of generality 431(1988)228

Formation of diastereomeric derivatives of 2-arylpropionic acids using *x*-leucinamide: the derivatization principle and its applicability. A reply 431(1988)229

Measurement of carprofen enantiomer concentrations in plasma and urine using L-leucinamide as the chiral coupling component 433(1988)331

**Leucine**

Rapid HPLC method to measure plasma leucine: importance in the study of leucine kinetics *in vivo* 426(1988)370

**Leucine-binding protein**

Detection of ligand-protein binding by direct electrophoresis of the complex 449(1988)331

**Leucine-enkephalin**

HPLC determination of leucine-enkephalin-like peptide in rat brain by pre-column fluorescence derivatization involving formylation followed by reaction with 1,2-diamino-4,5-dimethoxybenzene 430(1988)271

Assay for enkephalin-degrading peptidases in rat brain tissues by HPLC with on-line post-column fluorescence detection 430(1988)291

Simultaneous analysis of methionine- and leucine-enkephalin from rat brain: quantification by LC-electrochemistry 431(1988)65

**Leucovorin**

Direct resolution of the stereoisomers of leucovorin and 5-methyltetrahydrofolate using a bovine serum albumin HPLC chiral stationary phase coupled to an achiral phenyl column 424(1988)158

**Leucylphenylalanine methyl ester**

Chromatographic resolution of dipeptide enantiomers and diastereomers on chiral stationary phases from poly(L-leucine) or poly(L-phenylalanine) 450(1988)271

**Leukotriene antagonist**

HPLC determination of 4-[(3-(4-acetyl)-3-

hydroxy-2-propylphenoxy)propyl)sulfonyl]- $\gamma$ -oxobenzenebutanoic acid and metabolites in human plasma 417(1987)447

#### **Leukotriene B<sub>4</sub>**

Determination of leukotriene B<sub>4</sub> by HPLC with electrochemical detection 416(1987)170

#### **Leukotrienes**

Single-step procedure for the extraction and purification of leukotrienes B<sub>4</sub>, C<sub>4</sub> and D<sub>4</sub> 417(1987)241

MS evaluation of the *tert*-butyldimethylsilyl derivatives of monohydroxyeicosatetraenoic acids and leukotrienes 423(1987)1

Reversed-phase ion-interaction chromatography of leukotrienes, lipoxins and related compounds 427(1988)1

Arachidonic acid metabolites from polymorphonuclear leukocytes of healthy donors, severely burned patients and children with cystic fibrosis – routine monitoring by HPLC 427(1988)199

Automated HPLC extraction and quantification procedure for lipoxygenase metabolites 430(1988)1

Isocratic determination of arachidonic acid 5-lipoxygenase products in human neutrophils by HPLC 430(1988)299

Rapid and simple methods for the investigation of lipoxygenase pathways in human granulocytes 431(1988)413

#### **Levodopa**

Simultaneous determination of levodopa, its metabolites and carbidopa in clinical samples 415(1987)393

#### **Levopimaric acid**

GC analysis of underivatized resin acids 448(1988)428

#### **Lewis blood group antigens**

Immunoaffinity isolation of the Lewis blood group antigens 444(1988)41

#### **Lidocaine**

Retention properties of internal-surface reversed-phase silica packing and recovery of drugs from human plasma 420(1987)297

In vivo metabolism of lidocaine in the rat. Isolation of urinary metabolites as pentafluorobenzoyl derivatives and their

identification by combined GC-MS 421(1987)267

HPLC of lidocaine and nine of its metabolites in human plasma and urine 423(1987)199

#### **Ligand leakage**

Determination of the leakage from Phenyl-Sepharose CL-4B, Phenyl-Sepharose FF and Phenyl-Superose in bulk and column experiments 403(1987)85

#### **Ligand-protein binding**

Detection of ligand-protein binding by direct electrophoresis of the complex 449(1988)331

#### **Light stabilizers**

Determination of the polymeric light stabilizer Tinuvin 622 in polyolefins 450(1988)426

Determination of the polymeric light stabilizer Chimassorb 944 in polyolefins by reversed-phase HPLC 450(1988)430

#### **Lignans**

Assay of lignans of *Schizandra chinensis* in Sheng Mai San by HPLC 438(1988)447

#### **Limit-dextrinase**

Purification of germinated barley  $\alpha$ -amylase isozymes and limit-dextrinase by chromatofocusing and affinity chromatography 441(1988)436

#### **Limonene**

Resolution of monoterpene enantiomers by GC 407(1987)243

Characterisation of volatile oil constituents with relatively long GC retention times on two stationary phases 449(1988)312

#### **Linalool**

Chirality determinations for a tertiary alcohol. Ratios of linalool enantiomers in insects and plants 407(1987)324

#### **Lincosaminides**

Ion-pairing LC method for the determination of pirlimycin hydrochloride 402(1987)335

#### **Linear alkylbenzenesulphonates**

Determination of linear alkylbenzenesulphonates, alkylphenol polyethoxylates and nonylphenol in waste water by HPLC after enrichment on octadecylsilica 403(1987)243

**Linear chromatography**

Chemical engineering approach to dynamic modelling of linear chromatography. A flexible method for representing complex phenomena from simple concepts 406(1987)11

**Linoleate**

LC analysis of products of reaction between methyl linoleate and 4-*tert.*-butyl-*o*-benzoquinone 408(1987)171

**Linoleic acid derivatives**

Determination of corrosion inhibitor/lubricity enhancer additives in jet fuels by SEC 437(1988)203

**Linuron**

Increasing extraction efficiency in supercritical fluid extraction from complex matrices. Predicting extraction efficiency of diuron and linuron in supercritical fluid extraction using supercritical fluid chromatographic retention 447(1988)53

**Lipase activity**

Rapid radio TLC for assay of lipase-catalyzed esterification and interesterification reactions 450(1988)448

**Lipidex 5000**

Analysis of pentachlorophenol in water and urine by enrichment with Lipidex 5000 414(1987)55

**Lipid peroxides**

LC determination of serum lipid peroxides as diphenylthiobarbituric acid-reactive substance to collagen disease patients—a preliminary study 417(1987)371

**Lipids**

HPLC separation of diacylglycerol enantiomers on a chiral stationary phase 402(1987)257

Identification by combined GC-MS of constituent long-chain fatty acids and alcohols from the meibomian glands of the rat and a comparison with human meibomian lipids 414(1987)253

HPLC separation of femtomolar quantities of endogenous carboxylic acids, including arachidonic acid metabolites, as 4-bromomethyl-7-acetoxycoumarin derivatives 416(1987)247

Single-step procedure for the extraction and purification of leukotrienes B<sub>4</sub>, C<sub>4</sub> and D<sub>4</sub> 417(1987)241

TLC of human platelet phospholipids with fatty acid analysis 420(1987)411

HPLC quantitation of triacylglycerols containing fenoprofen from biological samples 422(1987)33

Analysis of organochlorine pesticides, polychlorinated biphenyls, dibenzo-*p*-dioxins and dibenzofurans in human milk by extraction with the lipophilic gel Lipidex 5000 422(1987)103

Fatty acid profiles of sebaceous triglycerides by capillary GC with mass-selective detection 425(1988)11

Identification of very-long-chain fatty acids in rat and mouse harderian gland lipids by capillary GC-MS 431(1988)231

MS analysis of endogenous diacylglycerols in normal and psoriatic skin 432(1988)278

Application of methylamine O-deacylation and Vitride reduction to the analysis of neutral lipids: a TLC evaluation 434(1988)215

Profiling of plasma cholesterol ester and triglyceride fatty acids as their methyl esters by capillary GC, preceded by a rapid aminopropyl-silica column chromatographic separation of lipid classes 434(1988)395

Quantitative analysis of triglycerides by HPLC using non-linear gradient elution and flame ionization detection 435(1988)139

Calibration of TLC with flame ionization detection for the analysis of natural lipid samples 435(1988)343

Separation of rat liver phospholipids from polar non-phospholipids by TLC 435(1988)513

Separation of complex lipids of cereals by HPLC with mass detection 436(1988)510

Qualitative and quantitative TLC analysis of polar lipids from leaves of glanded and glandless cotton genotypes 437(1988)211

Rapid separation of free fatty acids from neutral lipids in brain extracts 439(1988)453

Isolation and characterisation of intact steryl ferulates from seeds 440(1988)23

Investigation of the lipids of saprophytic mycobacteria in the U.S.S.R. (Review) 440(1988)479

Concerning hydrolysis of mycolate esters, of phthiocerol dimycocerosates and of related mycobacterial lipids: an anecdotal account 440(1988)487

Characterization of a lipid-rich fraction synthesized by *Streptomyces avermitilis* 441(1988)31

Preparation of fatty acid methyl esters by direct transesterification of lipids with aluminium chloride-methanol 441(1988)99

Densitometric identification of triglycerides separated by reversed-phase TLC 446(1988)259

Separation of triglyceride groups by reversed-phase TLC on silanized Kieselguhr 446(1988)283

Normal-phase HPLC of triacylglycerols 448(1988)135

#### Lipoamidase

Determination of lipoamidase activity by LC with fluorimetric detection 423(1987)304

#### Liperoxides

Determination of plasma liperoxides by HPLC 413(1987)237

#### Lipophilicity

see also Hydrophobicity

Quantitative structure-activity relationship of coumarin derivatives. Involvement of partition between aqueous and membrane phase 445(1988)49

Lipophilicity measurement of benzodiazepine-receptor ligands by reversed-phase LC. Comparison between HPLC and TLC 447(1988)404

#### Lipophilicity indices

Hydrophobicity studies on diadamantanyl derivatives. Comparison between calculated partition coefficients and experimental lipophilicity indices determined by reversed-phase HPLC and TLC 445(1988)317

#### Lipopolysaccharides

Determination of endotoxins by GC: evaluation of electron-capture and negative-ion chemical-ionization MS detection of halogenated derivatives of  $\beta$ -hydroxymyristic acid 417(1987)11

Determination of environmental levels of peptidoglycan and lipopolysaccharide using GC with negative-ion chemical-ionization MS utilizing bacterial amino acids and hydroxy fatty acids as biomarkers 431(1988)1

Large-scale fractionation of S-form lipopolysaccharide from *Salmonella abortus equi*.

Chemical and serological characterization of the fractions 440(1988)397

Analysis of Salmonella lipopolysaccharides by sodium deoxycholate-polyacrylamide gel electrophoresis 450(1988)381

#### Lipoproteins

Separation and characterization of high-density lipoprotein subpopulations by GPC 414(1987)25

Anion-exchange fast protein LC characterization and purification of apolipoproteins A-I, A-II, C-I, C-II, C-III<sub>0</sub>, C-III<sub>1</sub>, C-III<sub>2</sub> and E from human plasma 414(1987)35

Characterization of an apolipoprotein C-III mutant by HPLC and time-of-flight secondary ion MS 414(1987)323

Chromatography of complex protein mixtures (Review) 418(1987)115

Single-stage evaluation of serum lipoproteins by gel permeation using a specific fluorescent lipid probe 421(1987)136

Determination of coenzyme Q<sub>10</sub>,  $\alpha$ -tocopherol and cholesterol in biological samples by coupled-column LC with coulometric and UV detection 425(1988)87

Rapid quantification of apolipoprotein E enriched very-low-density lipoproteins by heparin-Sepharose chromatography 425(1988)169

Improved HPLC method for the determination of apolipoproteins in serum high-density lipoproteins 430(1988)135

Determination of bromhexine in plasma by reversed-phase LC. Interference of lipoproteins on extraction 432(1988)243

Fractionation of baboon chylomicrons and very-low-density lipoproteins by HPLC 433(1988)257

#### Liposomes

Chromatography of functionalized liposomes and their components 441(1988)115

#### $\beta$ -Lipotropin

Separation of protein hormones (Review) 418(1987)245

Combined HPLC-radioimmunoassay method for the analysis of endorphins, enkephalins and other neurotransmitter peptides 423(1987)93

#### Lipoxins

Reversed-phase ion-interaction chromatography of leukotrienes, lipoxins and related compounds 427(1988)1

Determination of picogram amounts of lipoxin A<sub>4</sub> and lipoxin B<sub>4</sub> by HPLC with electrochemical detection 428(1988)237

#### **Lipoxygenase metabolites**

Automated HPLC extraction and quantification procedure for lipoxygenase metabolites 430(1988)1

#### **Lipoxygenase products**

Rapid and simple methods for the investigation of lipoxygenase pathways in human granulocytes 431(1988)413

#### **Liquid alkylammonium salts**

Chemical interactions as a possible limitation on the useful solvent properties of liquid alkylammonium salts 435(1988)17

#### **Liquid chromatography–mass spectrometry**

Improvement of the interface for coupling of fast atom bombardment MS and micro HPLC 435(1988)482

Thermospray LC–MS method for the analysis of metribuzin and its metabolites 438(1988)359

Microbore HPLC–MS for the analysis of proteolytic digests by continuous-flow fast-atom bombardment MS 443(1988)355

#### **Liquid crystals**

Disc-like liquid crystal stationary phases derived from hexahydroxybenzene 402(1987)73

New method of visualizing thin-layer chromatograms by using liquid crystals 405(1987)203

Molecular theory of selectivity based on solute structure for nematic liquid-crystalline phases in GLC 406(1987)27

Liquid crystals. VI. Comparative study of the supercooling effect of some liquid crystalline stationary phases using GLC 411(1987)456

High-resolution GC with liquid crystal glass capillaries. X. Separation of diastereomeric alkanes 436(1988)47

Parameters of the liquid crystal method of visualizing thin-layer chromatograms 437(1988)131

New liquid crystalline stationary phase for capillary GC 439(1988)381

#### **Liquid desorption**

Headspace gas analysis. Liquid desorption of

headspace volatiles trapped on activated carbon open tubular traps 402(1987)95

#### **Liquid organic salts**

Solvent properties of liquid organic salts used as mobile phases in microcolumn reversed-phase LC 411(1987)61

#### **Lithium**

Quantitation of Li<sup>+</sup> in serum by capillary zone electrophoresis with an on-column conductivity detector 425(1988)385

#### **Lithocholic acid**

Improved method for the determination of the major neutral steroids and unconjugated bile acids in human faeces using capillary GC 421(1987)21

#### **Lombricine**

HPLC determination of lombricine and N-phosphoryl lombricine in the earthworm by pre-column fluorescence derivatisation with *o*-phthalaldehyde–ethanethiol 445(1988)433

Resolution of lombricine enantiomers by HPLC utilising pre-column derivatisation with *o*-phthalaldehyde–chiral thiols 447(1988)392

#### **Loperamide**

HPLC determination of loperamide hydrochloride in pharmaceutical preparations 449(1988)341

#### **Lorazepam**

Analysis of chlorodesmethyldiazepam and its metabolites in plasma and urine 420(1987)417

Computer-aided optimisation of drug enantiomer separation in chiral HPLC 434(1988)377

#### **Low-angle laser light scattering**

Conformational studies of bovine alkaline phosphatase in hydrophobic interaction and size-exclusion chromatography with linear diode array and low-angle laser light scattering detection 442(1988)29

#### **Low-density lipoprotein receptor**

Purification of monoclonal antibodies against the low-density lipoprotein receptor by preparative isotachopheresis 431(1988)327

#### **Lubricity enhancers**

Determination of corrosion inhibitor/lubricity enhancer additives in jet fuels by SEC 437(1988)203

**Luminescence quenching**

Dynamic quenching of europium(III) and terbium(III) luminescence: a potential detection method in ion chromatography 439(1988)165

**Luminol chemiluminescence**

HPLC determination of glucosides (glucose conjugates) with post-column reaction detection combining immobilized enzyme reactors and luminol chemiluminescence 449(1988)217

**Lung air**

Analysis of lung air from patients with bronchogenic carcinoma and controls using GC-MS 432(1988)1

**Lutein**

Separation and quantitative analysis of some carotenoid fatty acid esters of fruits by LC 435(1988)113

Separation of carotenol fatty acid esters by HPLC 449(1988)119

**Luteolin**

Separations of flavonoids and alkaloids in medicinal herbs by high-speed counter-current chromatography 435(1988)159

**Lutropin**

Several intermediate forms in the processing of rat lutropin subunits as shown by sodium dodecyl sulphate polyacrylamide gel electrophoresis 440(1988)353

**Lymphocyte blastogenesis inhibitory factor**

Fast protein LC of lymphocyte blastogenesis inhibitory factor produced by the human macrophage-like cell line U937 440(1988)131

**Lymphocyte receptors**

Isolation of specific lymphocyte receptors by high-performance immunoaffinity chromatography 444(1988)13

**Lymph-to-plasma protein concentration ratios**

Computer analysis of proteins separated by polyacrylamide gradient pore gel electrophoresis 416(1987)265

**Lysophosphatidylcholine**

Method for determination of different phospholipids by circular HPTLC 413(1987)257

**Lysophospholipids**

Molecular species analysis of lysophospholipids using HPLC and argentation TLC 409(1987)291

**Lysozyme**

Interaction of proteins with immobilized  $\text{Cu}^{2+}$ . Quantitation of adsorption capacity, adsorption isotherms and equilibrium constants by frontal analysis 403(1987)197

Multi-stage mixer-settler planet centrifuge. Preliminary studies on partition of macromolecules with organic-aqueous and aqueous-aqueous two-phase solvent systems 437(1988)121

Displacement chromatography of biomolecules 439(1988)341

Chromatographic resolution of lysozyme variants 443(1988)221

Immunological-chromatographic analysis 444(1988)1

**M****Mabuterol**

Determination of clenbuterol and mabuterol in equine plasma by ion-pair LC with electrochemical detection. Chromatographic and electrochemical characteristics 441(1988)197

**Macroamylase**

Use of a post-column enzymatic reaction for detection of macroamylasemia by HPLC 428(1988)415

**Macrolide alkaloids**

Efficient isolation of insect growth inhibitory macrolide alkaloids using recycle high-performance GPC 402(1987)354

**Macrophage colony-stimulating factor**

Purification of murine macrophage colony-stimulating factor using hydroxyapatite HPLC in the presence of sodium dodecyl sulphate 421(1987)355

**Macrophage factors**

Application of chromatographic methods to the analysis of macrophage factors induced by *Nocardia opaca* cell walls 440(1988)119



**Macrophage replacing factor**

N-Acetylmuramyl-L-alanyl-D-isoglutamine stimulates the immune response of B cells. Use of fast protein LC to purify a novel monokine involved in this effect 440(1988)95

**Macrotetrolides**

Preparative HPLC of macrotetrolides 442(1988)431

**Magnesium**

Isotachophoretic determination of the cationic constituents in mixtures of raw materials for glass manufacturing 439(1988)423

**Malaria antigen**

Reversed-phase LC and sodium dodecyl sulfate polyacrylamide gel electrophoresis characteristics of a recombinant DNA derived malaria antigen 444(1988)191

**Malaria circumsporozoite protein**

Assay, purification and characterization of a recombinant malaria circumsporozoite fusion protein by HPLC 411(1987)345

**Malate**

New method of purification of perchloric acid extracts for quantitative GC of some Krebs cycle acids 424(1988)357

**Maleimides**

Sensitive ferrocene reagents for derivatization of thiol compounds in HPLC with dual-electrode coulometric detection 419(1987)17

**Malto-oligosaccharides**

Two HPLC columns for analyses of malto-oligosaccharides 409(1987)396

**Maltopentaose**

LC assay of the relative activities of serum pancreatic and salivary  $\alpha$ -amylase using reductively pyridylaminated maltopentaose as a fluorescent substrate 419(1987)51

**Mandarin essential oils**

GC separation and chemometric analysis of mandarin essential oils 449(1988)183

**Mandelic acid**

HPLC method for the determination of a cyclandelate metabolite, mandelic acid, in human plasma 425(1988)203

Effects of long-term storage on the concentrations of the unconjugated acidic

metabolites of the trace amines, indoleamines and catecholamines 433(1988)23

**Manganese**

Ion chromatographic determination of low level cadmium(II), cobalt(II) and manganese(II) in water 450(1988)361

**Manganin electrode**

Amperometric detector with a manganin electrode for microcolumn LC 442(1988)111

**Mannitol**

Carbohydrate-borate eluents for anion chromatography 448(1988)355

**Mannonic acid**

Carbohydrate-borate eluents for anion chromatography 448(1988)355

**Mannose**

Analysis of neutral sugars as dinitrophenyl-hydrazones by HPLC 405(1987)221

HPLC separation of some mono- and disaccharides with detection by a post-column enzyme reactor and a chemically modified electrode 408(1987)157

Simultaneous GLC determination of aldoses and alduronic acids 408(1987)245

 **$\beta$ -D-Mannosidase**

Human urinary  $\beta$ -mannosidase: behaviour towards two chromatographic supports 428(1988)203

**Mannuronic acid**

Simultaneous GLC determination of aldoses and alduronic acids 408(1987)245

**Manoyloxid acid**

GC analysis of underivatized resin acids 448(1988)428

**Maprotiline**

Determination of maprotiline and desmethylmaprotiline in plasma and urine by GC with nitrogen-phosphorus detection 432(1988)334

**Marfey's reagent**

Application of Marfey's reagent in racemization studies of amino acids and peptides 444(1988)115

**Mass-selective detector**

The mass-selective detector as a chlorine-selective detector 438(1988)233

**Mass overload**

Simplified description of HPLC separation under overload conditions, based on the Craig distribution model. III. Computer simulations for two co-eluting bands assuming a Langmuir isotherm 405(1987)1

Simplified description of HPLC separation under overload conditions, based on the Craig distribution model. IV. Comparison of the model with experimental data for samples containing two or more solutes 405(1987)31

Simplified description of HPLC separation under overload conditions, based on the Craig distribution model. V. Gradient elution separation 405(1987)51

Preparative separation of peptide and protein samples by HPLC with gradient elution. I. The Craig model as a basis for computer simulations 444(1988)303

Preparative separation of peptide and protein samples by HPLC with gradient elution. II. Experimental examples compared with theory 444(1988)325

**Mass transfer coefficients**

Reversed-flow GC applied to physico-chemical measurements 446(1988)39

**McReynolds constants**

Ambiguities in the determination of McReynolds stationary phase constants 411(1987)43

**Mecloxamine**

Screening procedure for the detection of alkanolamine antihistamines and their metabolites in urine using computerized GC-MS 428(1988)43

**Meconic acid**

Separation of eight alkaloids and meconic acid and quantitation of five principal alkaloids in gum opium by gradient reversed-phase HPLC 436(1988)455

**Meconium glycoproteins**

Isolation of the major O-glycosidically linked oligosaccharides obtained by alkaline borohydride degradation of human meconium glycoproteins 425(1988)35

**Medicagenic acid**

Simple GC method for the determination of medicagenic acid in alfalfa (*Medicago sativa*) 410(1987)169

**Medical plant extracts**

HPLC characterization of some medical plant extracts used in cosmetic formulas 442(1988)353

**Medium polar compounds**

Super- and subcritical fluid chromatography-Fourier transform IR spectrometry of medium polar compounds on polymeric styrene-divinylbenzene packed columns 407(1987)87

**Medroxalol**

Application of column switching in HPLC analysis of medroxalol in plasma 417(1987)143

**Medroxyprogesterone acetate**

Medroxyprogesterone acetate- and ethinylestradiol-induced changes in biliary bile acids of the rat studied by HPLC 423(1987)63

**Medrylamine**

Screening procedure for the detection of alkanolamine antihistamines and their metabolites in urine using computerized GC-MS 428(1988)43

**Mefloquine**

HPLC separation and extraction investigation for the simultaneous determination of mefloquine and its carboxylic acid metabolite 427(1988)295

Determination of mefloquine by electron-capture GC after phosgene derivatization in biological samples and in capillary blood collected on filter paper 428(1988)281

HPLC assay for the simultaneous monitoring of mefloquine and its acid metabolite in biological samples using protein precipitation and ion-pair extraction 432(1988)253

Detection and determination of antimalarial drugs and their metabolites in body fluids (Review) 434(1988)1

**Mefruside**

LC screening of diuretics in urine 415(1987)347

**Melamine**

Capillary GC separation and MS detection of cyromazine and its metabolite melamine 408(1987)335

Determination of melamine in air by HPLC with UV detection 411(1987)409

**Melanocyte-stimulating hormones**

High-performance displacement chromatography

of melanotropins and their derivatives  
440(1988)157

### **Melanotropins**

High-performance displacement chromatography of melanotropins and their derivatives  
440(1988)157

### **Melatonin**

Melatonin and other indoles in rat pineal  
428(1988)206

Determination of plasma 5-hydroxytryptophan, 5-hydroxytryptamine, 5-hydroxyindoleacetic acid, tryptophan and melatonin by HPLC with electrochemical detection 431(1988)170

### **Melittine**

Characterization of bee venom and its main components by HPLC 408(1987)341

### **Melittin-silica**

Melittin-silica, a high-pressure affinity chromatography resin for calmodulin  
403(1987)99

### **Melperone**

GC-MS study of urinary metabolism of melperone 427(1988)144

### **Membrane proteins**

Chromatography of complex protein mixtures (Review) 418(1987)115

Column LC of integral membrane proteins (Review) 418(1987)223

Chromatographic detergent exchange in the preparation of membrane receptor protein  
427(1988)341

Use of a small TSK GSW HPLC column for large-zone chromatographic studies of monomer-oligomer equilibria of membrane protein 442(1988)229

Effect of detergents on the structure of integral membrane proteins of Sendai virus studied with size-exclusion HPLC and monoclonal antibodies  
443(1988)255

High-performance concanavalin A affinity chromatography of liver and hepatoma membrane proteins 444(1988)29

### **Membrane reactor**

LC determination of carbohydrates with pulsed amperometric detection and a membrane reactor  
447(1988)268

### **Membranes**

Comparison of serum protein removal in haemodialysis therapy by partly protein-permeable haemodialysers 428(1988)25

### **Menaquinone-4**

Simultaneous determination of vitamin K<sub>1</sub>, vitamin K<sub>1</sub> 2,3-epoxide and menaquinone-4 in human plasma by HPLC with fluorimetric detection 430(1988)21

### **Menaquinones**

Determination of K vitamins (phylloquinone and menaquinones) in umbilical cord plasma by a platinum-reduction column 430(1988)143

### **Menthol**

Cyclodextrin-chiral substrate interactions. High field NMR and molecular graphics studies  
450(1988)175

### **Menthyl acetate**

Characterisation of volatile oil constituents with relatively long GC retention times on two stationary phases 449(1988)312

### **Mepacrine**

Determination of quinacrine (mepacrine) in plasma by HPLC with fluorimetric detection  
420(1987)341

### **Mephénytoin**

Planar chromatographic separation of enantiomers and diastereomers with cyclodextrin mobile phase additives 448(1988)345

### **Mepirzepine**

Capillary GC assay for the routine monitoring of the antidepressant mepirzepine in human plasma  
417(1987)203

### **Merbarone**

Quantitation of merbarone in human plasma by HPLC 416(1987)203

### **Mercaptalbumin**

Rapid preparation of bovine mercaptalbumin by means of covalent chromatography on silica-based materials 408(1987)263

### **2-Mercapto-1-methylimidazole**

Identification of thyrostatic drug residues in animal thyroids by HPTLC and fluorescence reaction detection 442(1988)459

### **3-Mercapto-2-methylpropionic acid**

Determination of enantiomeric purity of 3-mercapto-2-methylpropionic acid 408(1987)365

**2-Mercaptopropionic acid**

HPLC of 2-mercaptopropionylglycine and its metabolite 2-mercaptopropionic acid in plasma and urine after treatment with thiopronine 417(1987)261

**2-Mercaptopropionylglycine**

HPLC of 2-mercaptopropionylglycine and its metabolite 2-mercaptopropionic acid in plasma and urine after treatment with thiopronine 417(1987)261

**6-Mercaptopurine**

Sensitive and specific HPLC assay for 6-thiouric acid 417(1987)183

Assay of 6-thioinosinic acid and 6-thioguanine nucleotides, active metabolites of 6-mercaptopurine, in human red blood cells 423(1987)169

Rapid method for evaluating compliance of 6-mercaptopurine therapy in children with leukemia 430(1988)163

***n*-Mers**

Removal of oligomers and *n*-mers from the Immobiline chemicals for isoelectric focusing 402(1987)105

**Mesaconitine**

HPLC analysis of C<sub>19</sub>-diterpenoid alkaloids 402(1987)366

**Mescaline**

Determination of  $\beta$ -phenylalkylamines in pharmaceutical preparations by analytical capillary isotachopheresis 405(1987)389

**Mesprenone**

Column LC of the novel aldosterone antagonist, mesprenone, and its active metabolite in plasma 414(1987)217

**Metabolite identification**

Analysis of basic compounds on a silica column with an aqueous methanol eluent. The use of quantitative structure-retention relationships in metabolite identification 407(1987)1

**Metal chelate adsorbent**

New metal chelate adsorbent selective for proteins and peptides containing neighbouring histidine residues 411(1987)177

**Metal chelates**

GC of metal  $\beta$ -diketonates and their analogues (Review) 411(1987)1

Elution recovery of tris(acetylacetonato)aluminium(III), -gallium(III) and -indium(III) in SEC 435(1988)368

Separation of inorganic and organomercury chelates on a polystyrene-divinylbenzene copolymer column 438(1988)414

**Metal complexes**

TLC of metal complexes on glycidoxypopyl functionalized silica plates 409(1987)377

Adsorption behaviour of tris( $\beta$ -diketonato)metal(III) complexes on silica gel 438(1988)401

Trace enrichment and separation of As<sup>III</sup>, Sb<sup>III</sup> and Bi<sup>III</sup> as diethyldithiophosphate complexes by reversed-phase HPLC 439(1988)63

LC retention of  $\beta$ -diketonato complexes of chromium(III) and palladium(II) in octadecyl-bonded silica/water-methanol systems. Comparison with liquid-liquid partition using dodecane as a model alkyl-bonded phase 439(1988)245

Chromatographic studies of metal complexes. I. PC of cobalt(III) complexes 442(1988)381

Rapid optical resolution of anionic metal complexes by GPC 450(1988)284

**Metal  $\beta$ -diketonates**

GC of metal  $\beta$ -diketonates and their analogues (Review) 411(1987)1

**Metal ions**

Contamination of aqueous eluents due to corrosion of stainless-steel chromatographic components 407(1987)133

Capillary tube isotachopheretic separation of heavy metal ions using complex-forming equilibria between cyanide as terminating ion and the metal ions 447(1988)383

**Metalloenes**

Planar chromatographic separation of enantiomers and diastereomers with cyclodextrin mobile phase additives 448(1988)345

**Metallo-cyanide complexes**

Separation and determination of stable metallo-cyanide complexes in metallurgical plant solutions and effluents by reversed-phase ion-pair chromatography 410(1987)419

**Metalloporphyrins**

Separation of metalloporphyrins from

metallation reactions by LC and electrophoresis 416(1987)71

Stability-indicating HPLC analysis of tin protoporphyrin and other free acid metalloporphyrins 445(1988)429

#### **Metallothioneins**

Isolation and quantitation of metallothionein isoforms using reversed-phase HPLC 402(1987)243

Improved separation of perch liver metallothionein by fast protein LC 402(1987)293

Rapid purification of the isoforms of rat liver Cd,Zn-metallothionein by HPLC 410(1987)470

Isolation and quantitation of cadmium-, zinc- and copper-metallothioneins by HPLC atomic absorption spectrometry 442(1988)345

#### **Metal-organic complexes**

Reversed-phase chromatographic separation and analysis of marine metal-organic complexes 436(1988)243

#### **Metalorganic compounds**

Sampling and injection system for the analysis of metalorganic chemicals by GC-MS. Some initial results 411(1987)445

#### **Metanephrine**

Simultaneous LC determination of seventeen of the major monoamine neurotransmitters, precursors and metabolites. I. Optimization of the mobile phase using factorial designs and a computer program to predict chromatograms 415(1987)261

Simultaneous LC determination of seventeen of the major monoamine neurotransmitters, precursors and metabolites. II. Assessment of human brain and cerebrospinal fluid concentrations 415(1987)275

Combined ion-pair extraction and HPLC for the determination of the biogenic amines and their major metabolites in single brain tissue samples 421(1987)51

#### **Methacrylates**

Considerations and remarks on the analysis of *Anthemis nobilis* L. essential oil by capillary GC and "hyphenated" techniques 411(1987)237

#### **Methamphetamine**

Analyses of impurities in methamphetamine by inductively coupled plasma MS and ion chromatography 437(1988)322

#### **Methane**

GC technique to simultaneously quantitate the gases produced by intestinal microorganisms from fermentation mixtures 414(1987)405

GC procedure for the qualitative analysis of silicone polyester resins after alkaline fusion 441(1988)207

#### **Methanesulphonic acid**

Simplified protein hydrolysis with methanesulphonic acid at elevated temperature for the complete amino acid analysis of proteins 448(1988)404

#### **Methanol**

Simultaneous determination of methanol, ethanol, acetone, isopropanol and ethylene glycol in plasma by GC 414(1987)248

Static headspace GC method for the determination of residual solvents in vigabatrin drug substance 438(1988)103

Preparation of fatty acid methyl esters by direct transesterification of lipids with aluminium chloride-methanol 441(1988)99

#### **Methiocarb**

HPLC determination of aryl N-methylcarbamate residues using post-column hydrolysis electrochemical detection 442(1988)333

#### **Methionine-enkephalin**

On-line post-column fluorescence detection for N-terminal tyrosine-containing peptides in HPLC 421(1987)245

Assay for enkephalin-degrading peptidases in rat brain tissues by HPLC with on-line post-column fluorescence detection 430(1988)291

Simultaneous analysis of methionine- and leucine-enkephalin from rat brain: quantification by LC-electrochemistry 431(1988)63

#### **Methixene**

Indirect HPLC resolution of racemic tertiary amines as their diastereomeric urea derivatives after N-dealkylation 436(1988)269

#### **Method detection limit**

Chromatographic detection limits in pharmaceutical method development 447(1988)1

#### **Methohexital**

Determination of sodium pentobarbital and either sodium methohexital or sodium thiopental in plasma by HPLC with ultraviolet detection 414(1986)484

**Methotrexate**

Rapid and simple assay for the measurement of methotrexate in serum, urine and red blood cells by reversed-phase HPLC 413(1987)320

Separation and identification of methotrexate and its metabolites, 7-hydroxymethotrexate and polyglutamates, in human tissues by reversed-phase HPLC coupled with radioimmunoassay 415(1987)445

Column LC of methotrexate and its metabolites using a post-column photochemical reactor and fluorescence detection 419(1987)213

Concurrent analysis of methotrexate, trimethoprim, sulphamethoxazole and their major metabolites in plasma by HPLC 422(1987)328

Determination of methotrexate and its major metabolite, 7-hydroxymethotrexate, using capillary zone electrophoresis and laser-induced fluorescence detection 426(1988)129

**Methotrimeprazine**

HPLC resolution of the enantiomers of thioridazine, its metabolites and related compounds 426(1988)223

**Methoxamine**

Determination of  $\beta$ -phenylalkylamines in pharmaceutical preparations by analytical capillary isotachopheresis 405(1987)389

**Methoxyacetic acid**

Assay of methoxyacetic acid in body fluids and tissues by GC-MS following *tert*-butyldimethylsilylation 448(1988)433

**2-Methoxyamphetamine**

Separation of optical isomers of methoxyphenamine and its metabolites as N-heptafluorobutyryl-L-prolyl derivatives by fused-silica capillary GC 431(1988)210

**N-Methoxy-N,O-bis(trimethylsilyl) carbamate**

N-Methoxy-N,O-bis(trimethylsilyl) carbamate as a derivatizing reagent for the GC of sitosterol degradation products 446(1988)253

**5-Methoxy-N,N-dimethyltryptamine**

GC-MS characterisation of the psychotomimetic indolealkylamines and their *in vivo* metabolites 422(1987)13

**2-Methoxy-5-hydroxymethamphetamine**

Separation of optical isomers of

methoxyphenamine and its metabolites as N-heptafluorobutyryl-L-prolyl derivatives by fused-silica capillary GC 431(1988)210

**3-Methoxy-4-Hydroxyphenylethylene glycol**

Simultaneous LC determination of seventeen of the major monoamine neurotransmitters, precursors and metabolites. I. Optimization of the mobile phase using factorial designs and a computer program to predict chromatograms 415(1987)261

Simultaneous LC determination of seventeen of the major monoamine neurotransmitters, precursors and metabolites. II. Assessment of human brain and cerebrospinal fluid concentrations 415(1987)275

Combined ion-pair extraction and HPLC for the determination of the biogenic amines and their major metabolites in single brain tissue samples 421(1987)51

Microassay of free 3-methoxy-4-hydroxyphenylglycol in plasma using HPLC with electrochemical detection 426(1988)351

Determination of free and sulphate-conjugated 3-methoxy-4-hydroxyphenylethylene glycol in human plasma by LC with electrochemical detection 430(1988)198

Effects of long-term storage on the concentrations of the unconjugated acidic metabolites of the trace amines, indoleamines and catecholamines 433(1988)23

**5-Methoxyindole-3-acetic acid**

Melatonin and other indoles in rat pineal 428(1988)206

**N-(Methoxymethyl)-2-chlorobutyroanilide**

HPLC methods for the determination of contaminants in technical phosmethylan 435(1988)380

**3-Methoxymorphinan**

Direct determination of dextromethorphan and its three metabolites in urine by HPLC using a precolumn switching system for sample clean-up 422(1987)340

**3-Methoxyparacetamol**

Complete separation of urinary metabolites of paracetamol and substituted paracetamols by reversed-phase ion-pair HPLC 426(1988)315

**Methoxyphenamine**

Separation of optical isomers of

methoxyphenamine and its metabolites as N-heptafluorobutyl-L-prolyl derivatives by fused-silica capillary GC 431(1988)210

#### **$\alpha$ -Methoxyphenylacetic acid**

Resolution of  $\alpha$ -arylacetic acid enantiomers on two chiral stationary phases 404(1987)123

#### **Methoxypsoralens**

Determination of psoralen in serum by reversed-phase HPLC 423(1987)383

#### **5-Methoxytryptophol**

Melatonin and other indoles in rat pineal 428(1988)206

#### **3-Methoxytyramine**

Simultaneous LC determination of seventeen of the major monoamine neurotransmitters, precursors and metabolites. I. Optimization of the mobile phase using factorial designs and a computer program to predict chromatograms 415(1987)261

Simultaneous LC determination of seventeen of the major monoamine neurotransmitters, precursors and metabolites. II. Assessment of human brain and cerebrospinal fluid concentrations 415(1987)275

Combined ion-pair extraction and HPLC for the determination of the biogenic amines and their major metabolites in single brain tissue samples 421(1987)51

#### **Methyclothiazide**

LC screening of diuretics in urine 415(1987)347

#### **N-Methyladrenaline**

Identification and quantification of N-methylepinephrine in human urine 414(1987)301

#### **Methylamine**

Application of methylamine O-deacylation and Vitride reduction to the analysis of neutral lipids: a TLC evaluation 434(1988)215

#### **Methylamino acids**

Rapid HPLC of 3-methylhistidine in human urine 413(1987)33

Rapid and sensitive method for the determination of arterial-venous differences and leg efflux of 3-methylhistidine using ion-pair HPLC and post-column fluorescence derivatization 414(1987)174

Analysis of methylated amino acids by HPLC: methylation of myelin basic protein 421(1987)130

Determination of methylated amino acids in human serum by HPLC 440(1988)225

#### **9-Methylanthracene**

Signal enhancement by on-column fluorometric detection in HPLC using micropacked fused-silica columns 435(1988)319

#### **Methylated phenols**

Separation of *m*- and *p*-cresol and other methylated phenols in shale oil by reversed-phase LC with electrochemical detection 441(1988)347

#### **Methyl atraton**

Specific identification of synthetic organic chemicals in river water using liquid-liquid extraction and resin adsorption coupled with electron impact, chemical ionization and accurate mass measurement GC-MS analyses 436(1988)229

#### **7-Methylbenz[e]acridine**

*endo*-1,4,5,6,7,7-Hexachlorobicyclo[2.2.1]hept-5-ene-2-carboxylic acid, a superior resolving agent for the HPLC separation of enantiomers of hydroxylated derivatives of two azaaromatic hydrocarbons 430(1988)53

#### **Methylbenzaldehyde**

Retention behaviour of aromatic aldehyde derivatives in reversed-phase LC 404(1987)242

#### **8-O-Methyl-14-benzoylaconine**

HPLC analysis of C<sub>19</sub>-diterpenoid alkaloids 402(1987)366

#### **$\alpha$ -Methylbenzylamine**

S-(+)-Flunoxaprofen chloride as chiral fluorescent reagent 427(1988)131

#### **N-Methylcarbamates**

Pre-column clean-up and LC determination of residues of N-methylcarbamate pesticides in extracts of total diet 404(1987)352

#### **N-Methylcarbamylated haemoglobin**

GC studies of the carbamylation of haemoglobin by methyl isocyanate in rats and rabbits 426(1988)239

#### **$\beta$ -Methyl digoxin**

Micro HPLC determination of cardiac glycosides in  $\beta$ -methyl digoxin and digoxin tablets 448(1988)157

#### **3'-Methyl-4-dimethylaminoazobenzene**

Detection of the changes in serum proteins during

chemical hepatocarcinogenesis by two-dimensional electrophoresis under non-denaturing conditions 419(1987)315

#### **Methyldopa**

Assay for tyrosine hydroxylase by HPLC with fluorescence detection 415(1987)289

Assay for tyrosine hydroxylase in hypothalamic homogenates using HPLC with electrochemical detection 419(1987)27

#### **4,4'-Methylenebis(2-chloroaniline)**

Determination of 4,4'-methylenebis(2-chloroaniline) in urine by HPLC 408(1987)399

#### **Methylene blue**

Foam counter-current chromatography with the cross-axis synchronous flow-through coil planet centrifuge 403(1987)77

#### **$\alpha$ -Methylene- $\gamma$ -butyrolactams**

Optimization of the direct chiral separation of potential cytotoxic  $\alpha$ -methylene- $\gamma$ -butyrolactones and  $\alpha$ -methylene- $\gamma$ -butyrolactams by LC 448(1988)55

#### **$\alpha$ -Methylene- $\gamma$ -butyrolactones**

Optimization of the direct chiral separation of potential cytotoxic  $\alpha$ -methylene- $\gamma$ -butyrolactones and  $\alpha$ -methylene- $\gamma$ -butyrolactams by LC 448(1988)55

#### **Methylene chloride**

Static headspace GC method for the determination of residual solvents in vigabatrin drug substance 438(1988)103

#### **Methylenedianiline**

Chromatographic determination of residual monomers of epoxy resins in wine simulants (in French) 445(1988)107

#### **10,11-Methylenedioxy-N-n-propylnoraporphine**

HPLC separation and electrochemical or spectrophotometric determination of *R*(-)-N-n-propylnorapomorphine and *R*(-)-10,11-methylenedioxy-N-n-propylnoraporphine in primate plasma 426(1988)283

#### **Methylergonovine**

HPLC method for the determination of methysergide and methylergonovine in human plasma 424(1988)416

#### **Methyl ethyl ketone**

Determination of methyl ethyl ketone and its

metabolites in urine using capillary GC 428(1988)275

#### **N-Methylformamide**

Identification of N-acetyl-S-(N-methylcarbamoyl)cysteine, a human metabolite of N,N-dimethylformamide and N-methylformamide 414(1987)399

GC method for the determination of N-acetyl-S-(N-methylcarbamoyl)cysteine, a metabolite of N,N-dimethylformamide and N-methylformamide, in human urine 431(1988)361

#### **2-Methylglutamic acid**

Resolution of  $\alpha$ -substituted amino acid enantiomers by HPLC after derivatization with a chiral adduct of *o*-phthalaldehyde. Application to glutamic acid analogues 440(1988)209

#### **Methyl glycosides, partially methylated**

Identification of partially methylated methyl glycosides by GC-MS of trimethylsilyl derivatives. Application to mycobacterial glycolipid antigen analysis 445(1988)87

#### **Methylglyoxal**

Determination of methylglyoxal as 2-methylquinoxaline by HPLC and its application to biological samples 414(1987)149

Simple and sensitive determination of methylglyoxal in biological samples by GC with electron-capture detection 415(1987)221

GC method for the estimation of acetone and its metabolites in biological samples 415(1987)372

#### **Methylhistamine**

Simultaneous determination of histidine-containing dipeptides, histamine, methylhistamine and histidine by HPLC 432(1988)315

#### **Methylhistidine**

Rapid HPLC of 3-methylhistidine in human urine 413(1987)33

Rapid and sensitive method for the determination of arterial-venous differences and leg efflux of 3-methylhistidine using ion-pair HPLC and post-column fluorescence derivatization 414(1987)174

Analysis of methylated amino acids by HPLC: methylation of myelin basic protein 421(1987)130

GC determination of 1- and 3-methylhistidine in biological fluids 423(1987)33



**Methylhydrazine**

GC analysis of monoalkylhydrazines  
423(1987)131

**Methylimidazoleacetic acids**

Simultaneous determination of imidazoleacetic acid and N<sup>7</sup>- and N<sup>8</sup>-methylimidazoleacetic acids in human urine by HPLC with fluorescence detection 416(1987)63

**2-Methylisoborneol**

Simplified method to quantify geosmin and 2-methylisoborneol concentrations in water and microbiological cultures 409(1987)337

**Methyl isocyanate**

GC studies of the carbamylation of haemoglobin by methyl isocyanate in rats and rabbits  
426(1988)239

**n-Methyl ketones**

GC on porous polymers. IV. Influence of the geometric structure of porous copolymers of 1,4-di(methacryloyloxymethyl)naphthalene with 1,4-divinylbenzene on their chromatographic behaviour 448(1988)233

**Methyl methacrylate**

Investigation by pyrolysis-GC of the composition of multicomponent polymeric microheterogeneous systems based on some vinyl monomers 404(1987)183

**1-Methyl-1'-[(3S)(-)-methylpinanyl]-4,4'-bipyridinium dichloride**

Isotachopheresis of quaternary 4,4'-bipyridylum salts. Analytical control of synthesis and purification procedures 449(1988)299

**N-Methyl oleoyl taurate**

Application of a novel cation-exchange reagent, Igepon T-77 N-methyl oleoyl taurate), to microbore separations of alumina extracts of catecholamines from cerebrospinal fluid, plasma, urine and brain tissue with amperometric detection 420(1987)241

**Methylpogonones**

Separation and determination of homoisoflavonoids in *Ophipogon japonicus* by reversed-phase HPLC 437(1988)265

**(±)-Methyl 2-oxopropyl 1,4-dihydro-2,6-dimethyl-4-(2-nitrophenyl)-3,5-pyridinedicarboxylate**

GC-negative-ion chemical ionization MS

determination of a new dihydropyridine calcium antagonist (MPC-1304) and its metabolites in human plasma and urine 434(1988)123

**Methylparaben**

HPLC assay of imazodan, methylparaben and propylparaben in imazodan injection  
450(1988)407

**1-Methyl-4-phenyl-4-propionoxypiperidine**

GLC-MS determination of 1-methyl-4-phenyl-4-propionoxypiperidine using a stable isotope-labeled analogue as an internal standard  
408(1987)356

**Methylphosphonic acid**

Analysis of alkyl methylphosphonic acids in aqueous matrices by ion-pair reversed-phase ion chromatography 407(1987)399

**Methylprednisolone**

HPLC determination of methylprednisolone and methylprednisolone 21-[8-[methyl-(2-sulfoethyl)amino]-8-oxooctanoate] sodium salt in human plasma 414(1987)1

Determination of methylprednisolone in rat tissue by HPLC 415(1987)35

Determination of methylprednisolone metabolites in human urine by GC-MS  
423(1987)15

Analysis of methylprednisolone, methylprednisone and corticosterone for assessment of methylprednisolone disposition in the rat 430(1988)241

Simultaneous analysis of methylprednisolone hemisuccinate, cortisol and methylprednisolone by normal-phase HPLC in human plasma  
432(1988)308

**Methylprednisolone hemisuccinate**

Simultaneous analysis of methylprednisolone hemisuccinate, cortisol and methylprednisolone by normal-phase HPLC in human plasma  
432(1988)308

**Methylprednisolone 21-[8-[methyl-(2-sulphoethyl)amino]-8-oxooctanoate]**

HPLC determination of methylprednisolone and methylprednisolone 21-[8-[methyl-(2-sulfoethyl)amino]-8-oxooctanoate] sodium salt in human plasma 414(1987)1

**Methylprednisone**

Analysis of methylprednisolone,

methylprednisone and corticosterone for assessment of methylprednisolone disposition in the rat 430(1988)241

**N-(2-Methyl-2-propyl)-3-oxo-4-aza-5 $\alpha$ -androst-1-ene-17 $\beta$ -carboxamide**

HPLC determination of N-(2-methyl-2-propyl)-3-oxo-4-aza-5 $\alpha$ -androst-1-ene-17 $\beta$ -carboxamide, a 4-azasteroid, in human plasma from a phase I study 427(1988)79

**N<sup>1</sup>-Methyl-2-pyridone-5-carboxamide**

Microdetermination of N<sup>1</sup>-methyl-2-pyridone-5-carboxamide, a major metabolite of nicotinic acid and nicotinamide, in urine by HPLC 417(1987)173

Simultaneous micro-determination of nicotinamide and its major metabolites, N<sup>1</sup>-methyl-2-pyridone-5-carboxamide and N<sup>1</sup>-methyl-4-pyridone-3-carboxamide, by HPLC 424(1988)23

Simultaneous micro-determination of nicotinamide and its major metabolites, N<sup>1</sup>-methyl-2-pyridone-5-carboxamide and N<sup>1</sup>-methyl-4-pyridone-3-carboxamide, by HPLC 424(1988)23

**2-Methylquinoxaline**

Determination of methylglyoxal as 2-methylquinoxaline by HPLC and its application to biological samples 414(1987)149

**Methyl silicones**

Estimation of the polarity of stationary phases by proton nuclear magnetic resonance spectroscopy. Application to phenyl and methyl silicones (OV and SE series) 402(1987)323

**O-Methylsterigmatocystin**

HPLC procedure for determining the profiles of aflatoxin precursors in wildtype and mutant strains of *Aspergillus parasiticus* 441(1988)400

**5-Methyltetrahydrofolate**

Direct resolution of the stereoisomers of leucovorin and 5-methyltetrahydrofolate using a bovine serum albumin HPLC chiral stationary phase coupled to an achiral phenyl column 424(1988)158

**N-(Methylthiomethyl)-2-chlorobutyroanilide**

HPLC methods for the determination of contaminants in technical phosmethylan 435(1988)380

**6-Methyl-2-thiouracil**

Identification of thyrostatic drug residues in animal thyroids by HPTLC and fluorescence reaction detection 442(1988)459

**Methyltin compounds**

Characterization of organotin species using microbore and capillary LC techniques with an epifluorescence microscope as a novel imaging detector 410(1987)393

**Methyl-p-toluenesulphinate**

Planar chromatographic separation of enantiomers and diastereomers with cyclodextrin mobile phase additives 448(1988)345

**O-Methyltransferases**

Resolution of five position-specific flavonoid O-methyltransferases by fast protein liquid chromatofocusing 407(1987)291

**Methyltriethoxysilane**

Trimethylsilylation of the hydrolysed and polycondensed products of methyltriethoxysilane 410(1987)137

**1-Methyluric acid**

HPLC assay for theophylline and its major metabolites in human urine 423(1987)179

**3-Methylxanthine**

HPLC assay for theophylline and its major metabolites in human urine 423(1987)179

**Methysergide**

HPLC method for the determination of methysergide and methylergonovine in human plasma 424(1988)416

**Metoclopramide**

LC analysis of alizapride and metoclopramide in human plasma and urine using solid-phase extraction 419(1987)233

Determination of metoclopramide and its glucuronide and sulphate conjugates in human biological fluids (plasma, urine and bile) by ion-pair HPLC 419(1987)243

Isolation of basic drugs from plasma using solid-phase extraction with a cyanopropyl-bonded phase 432(1988)209

**Metolazone**

LC screening of diuretics in urine 415(1987)347

**Metoprolol**

Rapid determination of the enantiomers of

metoprolol, oxprenolol and propranolol in urine 416(1987)331

Determination of metoprolol and two major metabolites in plasma and urine by column LC and fluorometric detection 417(1987)357

Correlation between reversed-phase HPLC and plasma protein binding 421(1987)83

Determination of metoprolol enantiomers in plasma and urine using (S)-(-)-phenylethyl isocyanate as a chiral reagent 421(1987)161

Reversed-phase HPLC assay to quantitate diastereomeric derivatives of metoprolol enantiomers in plasma 433(1988)318

Determination of alprenolol and metoprolol in plasma by column LC 434(1988)265

#### **Metribuzin**

Thermospray LC-MS method for the analysis of metribuzin and its metabolites 438(1988)359

#### **Metronidazole**

Simple LC method for the determination of ornidazole and metronidazole in human serum 420(1987)228

#### **Mexiletine**

Stereoselective analysis of the enantiomers of mexiletine by HPLC using fluorescence detection and study of their stereoselective disposition in man 415(1987)335

HPLC method for the determination of *p*-hydroxylated and hydroxymethylated metabolites of mexiletine in human serum 430(1988)406

HPLC assay for mexiletine hydroxylation in microsomes of human liver 431(1988)369

#### **Mezlocillin**

Application of HPLC of some antibiotics in clinical microbiology 427(1988)257

#### **Mianserin**

HPLC determination of mianserin in plasma and brain and its application to pharmacokinetic studies in the rat 434(1988)278

#### **Micellar electrokinetic capillary chromatography**

Separation of modified nucleic acid constituents by micellar electrokinetic capillary chromatography 409(1987)193

Analysis of water-soluble vitamins by micellar electrokinetic capillary chromatography 447(1988)133

#### **Micellar liquid chromatography**

The rôle of the stationary phase in micellar liquid chromatography. Adsorption and efficiency 449(1988)349

#### **Micelles**

Use of micellar mobile phases and microbore column switching for the assay of drugs in physiological fluids 427(1988)277

#### **Michael addition**

Chiral Michael additions of acetamidomalonate to  $\alpha$ -enones. NMR and HPLC resolution of enantiomers 450(1988)183

#### **Miconazole**

Sensitive and automated GC method for the determination of miconazol in plasma samples 425(1988)193

HPLC method for the determination of miconazole in vaginal fluid 434(1988)259

#### **Microbore columns**

see also Capillary columns and columns

#### **Microcolumns**

Contribution to the packing of LC microcolumns 411(1987)185

Separation of proteins by microcolumn LC based on the reversed-phase and size-exclusion principles 448(1988)73

#### **Microcomputer program**

Chromatographic behaviour of diastereoisomers. IX. Application of a microcomputer program to the TLC separation of some diastereoisomers on silica 438(1988)219

#### **Microcystins**

Analysis and purification of toxic peptides from cyanobacteria by reversed-phase HPLC 448(1988)275

#### **Microdistillation**

Microdistillation: a method combining GC and LC to characterize petroleum liquids 403(1987)159

#### **Micro-liquid chromatography**

Micro-LC and the chiral recognition mechanism on albumin-coated silica gel. Large selectivity changes with sample size 447(1988)297

#### **Micropellicular sorbents**

Rapid analysis of proteins and peptides by

reversed-phase chromatography with polymeric micropellicular sorbents 445(1988)71

#### **Midazolam**

Rapid and sensitive HPLC assay for midazolam and 1-hydroxymidazolam, the major metabolite, in human serum 433(1988)326

#### **Mildiomyacin**

Photodiode array detection of peptide-nucleoside antibiotics 436(1988)259

#### **Milk**

Dynamic headspace GC of volatile compounds in milk 404(1987)163

#### **Minecoside**

Application of various planar chromatographic techniques for the separation of iridoid glycosides from *Veronica officinalis* 404(1987)365

#### **Minoxidil**

Determination of minoxidil in pharmaceutical forms by capillary isotachopheresis 405(1987)385

#### **Mitomycin C**

Automated analysis of mitomycin C in body fluids by HPLC with on-line sample pre-treatment 420(1987)53

Chromatographic studies of mitomycin C degradation in albumin microspheres 430(1988)341

#### **Mitotane**

Rapid micromethod for the analysis of mitotane and its metabolite in plasma by GC with electron-capture detection 417(1987)287

#### **Mitoxantrone**

HPLC assay for mitoxantrone in plasma using electrochemical detection 420(1987)81

Direct determination of mitoxantrone and its mono- and dicarboxylic metabolites in plasma and urine by HPLC 424(1988)337

Mitoxantron determination using HPLC: improved sensitivity by loop-column injection for dual-dose pharmacokinetic studies 425(1988)429

#### **Mixed bonded phases**

Mixed bonded phases for HPLC. I. Synthesis and physico-chemical characterization 438(1988)263

#### **Mixed organic layers**

Further comments on the preparation of mixed

organic layers chemically bonded on silicon dioxide. Adsorption of modifiers from strongly adsorbed solvents 406(1987)295

#### **Mizoribine**

Simplified HPLC method for determination of mizoribine in human serum 432(1988)340

#### **Mobile phase composition**

Temperature effect in packed microcolumn chromatography using a carbon dioxide mobile phase 402(1987)328

Elution characteristics of polyol-borate eluents in anion-exchange chromatography. I. Eluent strength of sugar-, uronate-borate eluents 403(1987)27

Elution characteristics of polyol-borate eluents in anion-exchange chromatography. II. Tartrate-borate eluent for non-suppressed anion chromatography 403(1987)35

Origin of system peaks in single-column ion chromatography of inorganic anions using high pH borate-gluconate buffers and conductivity detection 404(1987)67

High-performance SEC of polypeptides on a TSK G2000SW column in acidic mobile phases 404(1987)215

Determination of cations at the parts-per-billion level with non-suppressed ion chromatography and indirect photometric detection 404(1987)282

Retention in capillary supercritical fluid chromatography with carbon dioxide mobile phases containing polar modifiers 406(1987)223

Ligand exchange chromatography of amino alcohol enantiomers as Schiff bases 406(1987)343

Influence of enantiomeric purity of a chiral selector on stereoselectivity 407(1987)217

Prediction of thermodynamic retention behaviour in reversed-phase LC with aqueous eluents and various concentrations of methanol by using the solubility concept model 408(1987)21

Investigation of an alkylamine modified and pH-controlled mobile phase for separation of pyrazolone derivatives by HPLC 409(1987)81

Retention behaviour of acidic, neutral and basic drugs on a CN column using phosphate buffers in the mobile phase 410(1987)77

Solvent properties of liquid organic salts used as mobile phases in microcolumn reversed-phase LC 411(1987)61

- Effect of phosphate concentration on the yield and purity of DNA separated from RNA by hydroxyapatite chromatography 411(1987)481
- Enantiomeric resolution of racemic nicotine and nicotine analogues by microcolumn LC with  $\beta$ -cyclodextrin inclusion complexes 411(1987)490
- Liquid chromatographic determination of diltiazem and its metabolites using *trans* isomers as internal standards, with dynamic modification of the solid phase by addition of an amine to the mobile phase 414(1987)109
- Simultaneous LC determination of seventeen of the major monoamine neurotransmitters, precursors and metabolites. I. Optimization of the mobile phase using factorial designs and a computer program to predict chromatograms 415(1987)261
- Use of micellar mobile phases and microbore column switching for the assay of drugs in physiological fluids 427(1988)277
- Behaviour of plasmalogens during HPLC on a silica column with a mobile phase containing phosphoric acid 434(1988)411
- Improved resolution of enantiomers of naproxen by the simultaneous use of a chiral stationary phase and a chiral additive in the mobile phase 435(1988)225
- Requirements of mutually miscible liquids for ternary chromatographic solvents with reference to monosaccharide PC 436(1988)53
- Direct LC separation of enantiomers on immobilized protein stationary phases. VI. Optical resolution of a series of racemic barbiturates: studies of substituent and mobile phase effects 436(1988)479
- Prediction of HPLC capacity factors following column calibration. Influence of different solvents and columns on the separation of a series of aromatic amides 437(1988)13
- Influence of additives to the eluent on the hydrophobic interaction chromatography of simple compounds. IV. Influence of electrolytes on the retention of normal alkanols at 25 °C 437(1988)337
- Influence of additives to the eluent on hydrophobic interaction chromatography of simple compounds. V. Influence of pH and electrolytes on the retention of *n*-alkylammonium ions and *n*-alkyl carboxylates 438(1988)181
- HPLC of amino acids, peptides and proteins. LXXXV. Evaluation of the use of hydrophobicity coefficients for the prediction of peptide elution profiles 438(1988)197
- Chromatographic behaviour of diastereoisomers. IX. Application of a microcomputer program to the TLC separation of some diastereoisomers on silica 438(1988)219
- Eluent selection for the determination of cations in ion chromatography 439(1988)49
- Main parameters governing the retention of polynuclear aromatic hydrocarbons on tetrachlorophthalimidopropyl-bonded silica 439(1988)195
- Modification of retention of some alkaloids in the system silanized silica/methanol + water + di(2-ethylhexyl)orthophosphoric acid 439(1988)257
- Bis(L-amino acid amidato)copper(II) complexes as chiral eluents in the enantiomeric separation of D,L-dansylamino acids by reversed-phase HPLC 441(1988)287
- Thermodynamic comparison of monomeric and polymeric C<sub>18</sub> bonded phases using aqueous methanol and acetonitrile mobile phases 442(1988)1
- Factors affecting the ion-pair chromatography of water-soluble vitamins 442(1988)81
- Chromatography of proteins on hydrophobic interaction and ion-exchange chromatographic matrices: mobile phase contributions to selectivity 443(1988)173
- Separation of proteins by reversed-phase HPLC. II. Optimizing sample pretreatment and mobile phase conditions 443(1988)381
- Effect of the mobile phase composition and ligand structure on the separation of D- and L-dansylamino acids, as mixed metal complexes, by reversed-phase HPLC 445(1988)219
- HPLC on dynamically modified silica. VIII. Gradient elution using eluents containing cetyltrimethylammonium bromide 447(1988)182
- Anion chromatography using a coated PRP-1 column and eluents of pH > 7 447(1988)315
- Non-suppressed ion chromatography of arsenic anions with potassium hydroxide-aromatic salt mixed eluents 447(1988)323
- Planar chromatographic separation of enantiomers and diastereoisomers with cyclodextrin mobile phase additives 448(1988)345

Carbohydrate–borate eluents for anion chromatography 448(1988)355

Comparative study of alkyl and fluoroalkyl N-(2-hydroxy-ethoxyethyl)-amides as reversed-phase LC modifiers 448(1988)365

Separation of pyridone carboxylic acid enantiomers by HPLC using copper(II)–L-amino acid as the eluent 448(1988)439

Some aspects of the selection of HPLC methods for the separation of chiral compounds in pharmaceutical analysis 450(1988)145

#### **Mobility**

Isotachophoretic determination of mobility and  $pK_a$  by means of computer simulation. VI. Mobility and  $pK_a$  of sixty anions in methanol 408(1987)27

#### **Moclobemide**

Determination of the new monoamine oxidase inhibitor moclobemide and three of its metabolites in biological fluids by HPLC 420(1987)111

#### **Model simulation**

Simplified description of HPLC separation under overload conditions, based on the Craig distribution model. III. Computer simulations for two co-eluting bands assuming a Langmuir isotherm 405(1987)1

#### **Molar refraction**

Correlation between molar refraction and GC retention data 403(1987)11

#### **Molar refraction–retention relationships**

Correlation between molar refraction and GC retention data 403(1987)11

#### **Molecular conformation**

Holistic molecular conformation and total surface area calculations as predictors for solution properties and chromatographic parameters 403(1987)1

#### **Molecular connectivity**

Factor analysis of trends in selectivity in GLC of 175 saturated and unsaturated esters on 14 polysiloxane stationary phases 404(1987)11

#### **Molecular graphics**

Cyclodextrin–chiral substrate interactions. High field NMR and molecular graphics studies 450(1988)175

#### **Molecular size distribution**

Determination of the molecular size distribution of immunoglobulin G (IgG) in intravenous IgG–albumin formulations by HPLC 444(1988)141

#### **Molecular weight determination**

Characterisation of carrageenans by high-performance size-exclusion chromatography using a LiChrospher 1000 DIOL column 403(1987)307

SEC of deposits in pulp and paper mills 411(1987)363

#### **Molybdenum**

Determination of molybdenum in soils and plants by HPLC 411(1987)401

#### **Monactin**

Preparative HPLC of macrotetrolides 442(1988)431

#### **Monoacetyldapsone**

Dapsone and monoacetyldapsone determined in serum and saliva by a sensitive HPLC method with a single extraction step 422(1987)322

#### **Monoacetylhydrazine**

Determination of the isoniazid metabolite monoacetylhydrazine in urine by HPLC 415(1987)188

#### **6-Monoacetylmorphine**

Identification of opiates in urine by capillary column GC of two different derivatives 422(1987)117

#### **Monoalkylhydrazines**

GC analysis of monoalkylhydrazines 423(1987)131

#### **Monoamine transmitters**

Perfluorinated acids as ion-pairing agents in the determination of monoamine transmitters and some prominent metabolites in rat brain by HPLC with amperometric detection 449(1988)191

#### **Monobromobimane**

Measurement of plasma thiols after derivatization with monobromobimane 424(1988)141

#### **Monoclonal antibodies**

Determination of chloramphenicol in swine muscle tissue using a monoclonal antibody-mediated clean-up procedure 411(1987)415

Purification of ascitic fluid-derived murine monoclonal antibodies by anion-exchange and size-exclusion HPLC 430(1988)329

Purification of monoclonal antibodies against the low-density lipoprotein receptor by preparative isotachopheresis 431(1988)327

Comparison of Mono O, Superose-6, and ABx fast protein LC for the purification of IgM monoclonal antibodies 444(1988)153

#### **N-Monodemethyldiltiazem**

High-performance liquid chromatographic determination of diltiazem and its major metabolites, N-monodemethyldiltiazem and desacetyldiltiazem, in plasma 415(1987)203

#### **Monodesethylflurazepam**

Simultaneous determination of flurazepam and five metabolites in serum by HPLC and its application to pharmacokinetic studies in rats 423(1987)251

#### **Monodesisopropyldisopyramide**

Simultaneous assay of disopyramide and monodesisopropyldisopyramide enantiomers in biological samples by LC 424(1988)424

Direct enantiomeric resolution of disopyramide and its metabolite using chiral HPLC.

Application to stereoselective metabolism and pharmacokinetics of racemic disopyramide in man 450(1988)211

#### **Monodesmethyldiltiazem**

HPLC determination of diltiazem and four of its metabolites in plasma: evaluation of their stability 430(1988)103

#### **Monodesmosidic saponins**

HPLC determination of molluscicidal saponins from *Phytolacca dodecandra* (Phytolaccaceae) 448(1988)265

#### **Monohydroxyeicosatetraenoic acids**

GC-MS of monohydroxyeicosatetraenoic acids as their methyl esters trimethylsilyl, allyldimethylsilyl and *tert.*-butyldimethylsilyl ethers 416(1987)219

MS evaluation of the *tert.*-butyldimethylsilyl derivatives of monohydroxyeicosatetraenoic acids and leukotrienes 423(1987)1

Reversed-phase ion-interaction chromatography of leukotrienes, lipoxins and related compounds 427(1988)1

Arachidonic acid metabolites from

polymorphonuclear leukocytes of healthy donors, severely burned patients and children with cystic fibrosis – routine monitoring by HPLC 427(1988)199

#### **Monohydroxy fatty acids**

Thermospray-MS analysis of underivatized monohydroxy fatty acids: application to stimulated platelets 425(1988)25

#### **Monokines**

N-Acetylmuramyl-L-alanyl-D-isoglutamine stimulates the immune response of B cells. Use of fast protein LC to purify a novel monokine involved in this effect 440(1988)95

Application of chromatographic methods to the analysis of macrophage factors induced by *Nocardia opaca* cell walls 440(1988)119

#### **Monomethylacetamide**

GC-MS method for the determination of dimethylacetamide and metabolites in whole blood 428(1988)156

#### **4-Monomethylaminoantipyrine**

Simple HPLC method for simultaneous determination of aminopyrine and its metabolites 421(1987)198

#### **Monomethylarginine**

Analysis of methylated amino acids by HPLC: methylation of myelin basic protein 421(1987)130

Determination of methylated amino acids in human serum by HPLC 440(1988)225

#### **$\beta$ -Monomethylchlorambucil**

HPLC of chlorambucil analogues 422(1987)315

#### **N-Monomethyldeacetylthymoxamine**

HPLC assay of the metabolites of thymoxamine 424(1988)205

#### **Monomethyllysine**

Analysis of methylated amino acids by HPLC: methylation of myelin basic protein 421(1987)130

#### **Monooxygenase**

Quantitative structure-activity relationship of coumarin derivatives. Involvement of partition between aqueous and membrane phase 445(1988)49

#### **Monosaccharides**

Elution characteristics of polyol-borate eluents in anion-exchange chromatography. I. Eluent strength of sugar-, uronate-borate eluents 403(1987)27

Determination of monosaccharides constituting the glycosides in saponins by HPLC 403(1987)319

HPLC separation of some mono- and disaccharides with detection by a post-column enzyme reactor and a chemically modified electrode 408(1987)157

GC analysis of neutral monosaccharides as their O-pentafluorobenzyloxime acetates 411(1987)275

Requirements of mutually miscible liquids for ternary chromatographic solvents with reference to monosaccharide PC 436(1988)53

Identification of phenols, phenolic acid dimers, and monosaccharides by GLC on a capillary column 436(1988)484

Determination of monosaccharides in glycoproteins by reversed-phase HPLC on 2.1-mm narrowbore columns 443(1988)299

Optimizing copper-bicinchoninate carbohydrate detection for use with water-elution HPLC: a technique to measure the major mono- and oligosaccharides in small pieces of wheat endosperm 449(1988)207

#### **Monoterpenes**

Resolution of monoterpene enantiomers by GC 407(1987)243

A cryotrap technique for the quantitation of monoterpenes in humid and ozone-rich forest air 442(1988)157

#### **Monothio- $\beta$ -diketones**

GC of metal  $\beta$ -diketonates and their analogues (Review) 411(1987)1

#### **$\beta$ -Mono(trifluoromethyl)chlorambucil**

HPLC of chlorambucil analogues 422(1987)315

#### **Morphinan-3-ol**

Direct determination of dextromethorphan and its three metabolites in urine by HPLC using a precolumn switching system for sample clean-up 422(1987)340

#### **Morphine**

HPLC and preliminary pharmacokinetics of nicomorphine and its metabolites 3-nicotinoyl- and 6-nicotinoylmorphine and morphine 416(1987)382

Identification of opiates in urine by capillary column GC of two different derivatives 422(1987)117

Concentrations of free and conjugated morphine in blood in twenty cases of heroin-related deaths 422(1987)267

Rapid, high-sensitivity method for measurement of morphine in guinea-pig serum 428(1988)383

Improved method for the simultaneous determination of morphine and its principal glucuronide metabolites 430(1988)394

Isolation of basic drugs from plasma using solid-phase extraction with a cyanopropyl-bonded phase 432(1988)209

Separation of eight alkaloids and meconic acid and quantitation of five principal alkaloids in gum opium by gradient reversed-phase HPLC 436(1988)455

Separation of opium alkaloids by carbon dioxide sub- and supercritical fluid chromatography with packed columns. Application to the quantitative analysis of poppy straw extracts 437(1988)351

#### **Morphine glucuronides**

Improved method for the simultaneous determination of morphine and its principal glucuronide metabolites 430(1988)394

#### **Morphinone**

Method for determination of morphinone in urine and bile of guinea pig by HPLC 421(1987)155

#### **Morpholinedithiocarbamates**

Assay of phenylmercury salts in pharmaceutical products by HPLC of the morpholinedithiocarbamate derivative 407(1987)389

#### **3-O-Morpholinoethylmorphine**

Determination of pholcodine in biological fluids by HPLC with fluorescence detection 424(1988)170

#### **Mucin**

Sulphate quantification in rabbit tracheobronchial mucus by reversed-phase ion-pair chromatography 424(1988)147

Determination of monosaccharides in glycoproteins by reversed-phase HPLC on 2.1-mm narrowbore columns 443(1988)299

#### **Multi-channel microwave-induced plasma**

Multiple element emission spectral detection GC profiles of halogenated products from chlorination of humic acid and drinking water 438(1988)35



**Multiple detection system**

Multiple detection system for the high-resolution microcolumn SEC separation of proteins with a slurry-packed capillary column 411(1987)221

**Multiple fronts**

Analysis of multiple front formation in the wetting of thin-layer plates 435(1988)397

**Multi-stage mixer-settler planet centrifuge**

Multi-stage mixer-settler planet centrifuge. Preliminary studies on partition of macromolecules with organic-aqueous and aqueous-aqueous two-phase solvent systems 437(1988)121

**Multivalent ion exchange**

Preparative chromatography of proteins. Analysis of the multivalent ion-exchange formalism 443(1988)13

**Multi-variate ranking function**

Multi-variate ranking function for optimizing separations 439(1988)173

**Murine monoclonal antibodies**

Purification of ascitic fluid-derived murine monoclonal antibodies by anion-exchange and size-exclusion HPLC 430(1988)329

**Mussaenoside**

Application of various planar chromatographic techniques for the separation of iridoid glycosides from *Veronica officinalis* 404(1987)365

**Mustard**

TLC of mustard and its metabolites 437(1988)423

**Mutatoxanthin**

LC profiles of major carotenoid esters in commercially processed California navel and Valencia orange juice concentrates 442(1988)249

**Mutually miscible liquids**

Requirements of mutually miscible liquids for ternary chromatographic solvents with reference to monosaccharide PC 436(1988)53

**Mycobacterial antigens**

Isolation of recombinant mycobacterial antigens by an automatic and generally applicable purification method for  $\beta$ -galactosidase fusion proteins 448(1988)165

**Mycolate esters**

Concerning hydrolysis of mycolate esters, of phthiocerol dimycocerosates and of related mycobacterial lipids: an anecdotal account 440(1988)487

**Mycotoxins**

Standardized HPLC of 182 mycotoxins and other fungal metabolites based on alkylphenone retention indices and UV-VIS spectra (diode array detection) 404(1987)195

Decomposition of trifluoroacetyl derivatives of some trichothecenes on fused-silica capillary columns during GC analysis 407(1987)340

HPLC method for determining trichothecene mycotoxins by post-column fluorescence derivatization 410(1987)427

Metabolism of three trichothecene mycotoxins, T-2 toxin, diacetoxyscirpenol and deoxynivalenol, by bovine rumen microorganisms 414(1987)335

Simultaneous detection of trichothecenes and rosenonolactone in grape juice and wine by capillary GC 448(1988)424

Large-scale purification of the mycotoxins aflatoxin B<sub>1</sub>, B<sub>2</sub> and G<sub>1</sub> 450(1988)101

**Myelin basic protein**

Analysis of methylated amino acids by HPLC: methylation of myelin basic protein 421(1987)130

Isolation of specific lymphocyte receptors by high-performance immunoaffinity chromatography 444(1988)13

**Myoglobin**

Determination of myoglobin in human serum by HPLC with chemiluminescence detection 416(1987)45

**Myosin**

Selective purification of cardiac myosin by a high-performance salicylate affinity column 435(1988)185

**Myristicin**

Determination of safrole and myristicin in nutmeg and mace by HPLC 438(1988)117

# N

## NAD<sup>+</sup> analogues

Synthesis and purification of deoxyribose analogues of NAD<sup>+</sup> by affinity chromatography and strong-anion-exchange HPLC 444(1988)89

## Nadolol

High-sensitivity determination of nadolol in plasma with mass-selective detection 416(1987)443

Rationale for application of low-mass selected-ion detection in the quantitative body fluid assay of nadolol 416(1987)445

## Nafimidone

HPLC determination of nafimidone and its major metabolite nafimidone alcohol in human plasma and urine 414(1987)389

## Naftidrofuryl hydrogenoxalate

Use of cyclodextrins in isotachopheresis. III. Purity control of naftidrofuryl hydrogenoxalate and some of its synthesis intermediates 435(1988)496

## Nalbuphine

Determination of nalbuphine by HPLC with electrochemical detection: application to clinical samples from postoperative patients 427(1988)113

## Nalidixic acid

Simultaneous determination of nalidixic acid, oxolinic acid and piromidic acid in fish by HPLC with fluorescence and UV detection 402(1987)301

## Naphthalene

Real-time quantitative analysis of unresolved LC peaks with the use of the Kalman filter 407(1987)59

## Naphthenes

Paraffins, olefins, naphthenes and aromatics analysis of selected hydrocarbon distillates using on-line column switching HPLC with dielectric constant detection 437(1988)365

## $\beta$ -Naphthoflavone

HPLC determination of  $\beta$ -naphthoflavone in biological fluids 416(1987)160

Differences in the constitutive forms of hepatic

cytochrome P-450 in male and female adult beagle dogs 419(1987)61

## Naphthoquinones

Separation of natural products by centrifugal partition chromatography 450(1988)91

## Naphthoxyacetic acid

HPLC determination of propranolol and its metabolites in rat serum 431(1988)465

Cyclodextrin–ligand interaction as a simplified model of biospecific affinity chromatography 448(1988)391

## Naphthoxyactic acid

HPLC determination of propranolol and its metabolites in rat serum 431(1988)465

## Naproxen

Rapid HPLC assay for the simultaneous analysis of non-steroidal anti-inflammatory drugs in plasma 416(1987)293

Assay method for product formation in *in vitro* enzyme kinetic studies of uridine diphosphate glucuronyltransferases: 2-arylpropionic acid enantiomers 430(1988)368

HPLC determination of ketoprofen and naproxen in rat plasma 431(1988)444

Improved resolution of enantiomers of naproxen by the simultaneous use of a chiral stationary phase and a chiral additive in the mobile phase 435(1988)225

Effects of drug–protein binding on trace enrichment of drugs in blood plasma on short precolumns 439(1988)353

Separation and determination of some amino acid ester enantiomers by TLC after derivatization with (*S*)-(+)-naproxen 450(1988)281

## Narceine

Separation of eight alkaloids and meconic acid and quantitation of five principal alkaloids in gum opium by gradient reversed-phase HPLC 436(1988)455

## Naringenin

Direct enantiomeric separation of racemic flavanones by HPLC using cellulose triacetate as a chiral stationary phase 441(1988)417

## Narmco 5208

Quantitative analysis of resins used in fiber-

reinforced composites by reversed-phase LC 447(1988)141

#### **Narmco Rigidite 5208**

High-performance SEC of fiber-reinforced organic-matrix composites 408(1987)129

#### **Natural products**

Separation of natural products by centrifugal partition chromatography 450(1988)91

#### **Neamine**

Identification of dekamycin antibiotics 440(1988)71

#### **Nearly ideal binary solvent model**

Thermochemical investigations of GLC. IV. Partition coefficients of solutes on binary mixtures containing two complexing solvents 402(1987)41

#### **Near-wall field-flow fractionation**

Near-wall field-flow fractionation. Application of surface forces and transversal voltage to fractionation in laminar flow 446(1988)131

#### **Neomycins**

Identification of dekamycin antibiotics 440(1988)71

#### **Neopterin**

Estimation of tetrahydrobiopterin and other pterins in plasma by isocratic LC with electrochemical and fluorimetric detection 432(1988)321

#### **Neopteroilins**

Interactions between ascorbic acid and linear and cyclized bile pigments. A TLC study 440(1988)275

#### **Netilmicin**

HPLC determination of netilmicin in guinea-pig and human serum by fluorodinitrobenzene derivatization with spectrophotometric detection 434(1988)169

#### **Neurohormones**

Single-step purification of two hyperglycaemic neurohormones from the sinus gland of *Procambarus bouvieri*. Comparative peptide mapping by means of HPLC 443(1988)337

#### **Neuroleptics**

Identification of drugs in human serum by HPLC with photodiode array detection and a search algorithm for UV spectra 419(1987)135

#### **Neuropeptides**

Reversed-phase HPLC–radioreceptor assay of human cerebrospinal fluid neuropeptides. Maximizing recovery of picomoles of peptides and minimizing memory 422(1987)61

Combined HPLC–radioimmunoassay method for the analysis of endorphins, enkephalins and other neurotransmitter peptides 423(1987)93

#### **Neurophysin**

Application of HPLC in neurophysin disulfide assignment 443(1988)285

#### **Neurotensin**

Characterization of fat-induced neurotensin-like immunoreactivity in plasma using column LC and radioimmunoassay 414(1987)180

#### **Neurotoxins**

Fast protein LC of botulinum neurotoxin types A, B and E 430(1988)279

#### **Neurotransmitters**

Simultaneous LC determination of seventeen of the major monoamine neurotransmitters, precursors and metabolites. I. Optimization of the mobile phase using factorial designs and a computer program to predict chromatograms 415(1987)261

Simultaneous LC determination of seventeen of the major monoamine neurotransmitters, precursors and metabolites. II. Assessment of human brain and cerebrospinal fluid concentrations 415(1987)275

Automated pre-column derivatization with *o*-phthalaldehyde for the determination of neurotransmitter amino acids using reversed-phase LC 417(1987)385

Combined HPLC–radioimmunoassay method for the analysis of endorphins, enkephalins and other neurotransmitter peptides 423(1987)93

Comparison of two methods for the determination of substance P immunoreactivity in human cerebrospinal fluid: sulphopropyl-Sephadex ion-exchange chromatography and/or reversed-phase HPLC combined with radioimmunoassay 425(1988)175

HPLC analysis of neurotransmitter amino acids in brain 428(1988)9

Perfluorinated acids as ion-pairing agents in the determination of monoamine transmitters and some prominent metabolites in rat brain by

HPLC with amperometric detection  
449(1988)191

#### Neutrophil stimulating activity

Neutrophil stimulating activity released by *Staphylococcus*-stimulated mononuclear leukocyte conditioned medium. Further characterization and partial purification 440(1988)105

#### New Coccine

Simple method for the analysis of food dyes on reversed-phase thin-layer plates 411(1987)437

#### Nicardipine

Capillary column GC method using electron-capture detection for the simultaneous determination of nicardipine and its pyridine metabolite II in plasma 415(1987)65

Simple method for the determination of nicardipine in plasma using HPLC 420(1987)439

#### Nickel bis[3-(heptafluorobutanoyl)-(1*R*)-camphorate]

Enantiomer analysis by complexation GC. Scope, merits and limitations 441(1988)135

#### Nickel clusters

Behaviour of tri- and tetranuclear iron and nickel clusters in HPLC 447(1988)187

#### Nicomorphine

HPLC and preliminary pharmacokinetics of nicomorphine and its metabolites 3-nicotinoyl- and 6-nicotinoylmorphine and morphine 416(1987)382

#### Nicotinamide

Microdetermination of N<sup>1</sup>-methyl-2-pyridone-5-carboxamide, a major metabolite of nicotinic acid and nicotinamide, in urine by HPLC 417(1987)173

HPLC determination of nicotinamide in rat tissue samples and blood after extraction with diethyl ether 422(1987)257

Simultaneous micro-determination of nicotinamide and its major metabolites, N<sup>1</sup>-methyl-2-pyridone-5-carboxamide and N<sup>1</sup>-methyl-4-pyridone-3-carboxamide, by HPLC 424(1988)23

#### Nicotine

Enantiomeric resolution of racemic nicotine and nicotine analogues by microcolumn LC with  $\beta$ -cyclodextrin inclusion complexes 411(1987)490

Simplified method for the determination of plasma cotinine using GC-MS 414(1987)449

GC determination of *trans*-3'-hydroxycotinine, a major metabolite of nicotine in smokers 415(1987)400

Radiometric-HPLC assay for nicotine and twelve of its metabolites 419(1987)191

Simultaneous determination of nicotine, cotinine and five additional nicotine metabolites in the urine of smokers using pre-column derivatisation and HPLC 419(1987)375

Assessment of exposure to environmental tobacco smoke using a HPLC method for the simultaneous determination of nicotine and two of its metabolites in urine 431(1988)216

#### Nicotinic acid

Microdetermination of N<sup>1</sup>-methyl-2-pyridone-5-carboxamide, a major metabolite of nicotinic acid and nicotinamide, in urine by HPLC 417(1987)173

High-voltage electrophoretic separation of tryptophan metabolites of the kynurenine pathway 420(1987)263

#### Nicotinoids

Enantiomeric resolution of racemic nicotine and nicotine analogues by microcolumn LC with  $\beta$ -cyclodextrin inclusion complexes 411(1987)490

Planar chromatographic separation of enantiomers and diastereomers with cyclodextrin mobile phase additives 448(1988)345

#### Nicotinoylmorphine

HPLC and preliminary pharmacokinetics of nicomorphine and its metabolites 3-nicotinoyl- and 6-nicotinoylmorphine and morphine 416(1987)382

#### Nifedipine

Rapid HPLC determination of nifedipine in plasma with on-line precolumn solid-phase extraction 420(1987)207

Determination of nifedipine and its three principal metabolites in plasma and urine by automated electron-capture capillary GC 425(1988)107

#### Nifurtimox

HPLC determination of nifurtimox in human serum 433(1988)359

**Nilvadipine**

Determination of nilvadipine in plasma and urine by capillary column GC with electron-capture detection 415(1987)156

**Nimesulide**

Simultaneous determination of nimesulide and hydroxynimesulide in human plasma and urine by HPLC 425(1988)413

**Ninhydrin**

Detection of amino acids on thin-layer plates 436(1988)341

**Nisoldipine**

GC determination of nisoldipine and one of its metabolites in plasma 423(1987)327

**Nitrate**

Derivatization of inorganic anions with pentafluorobenzyl methanesulphonate for GC 404(1987)175

Optimization of ion chromatography using commercially available detection systems and software 439(1988)13

**Nitrazepam**

Sensitive GC method for determining nitrazepam in serum and saliva 424(1988)403

HPLC determination of nitrazepam in plasma and its application to pharmacokinetic studies in the rat 425(1988)435

**Nitrendipine**

GC analysis of nitrendipine and its pyridine metabolite in human plasma 428(1988)362

**Nitriles**

Extraction and identification of nitrile compounds in Rundle shale oil 435(1988)193

Multiple element emission spectral detection GC profiles of halogenated products from chlorination of humic acid and drinking water 438(1988)35

**Nitrilotriacetic acid adsorbent**

New metal chelate adsorbent selective for proteins and peptides containing neighbouring histidine residues 411(1987)177

**Nitrite**

Derivatization of inorganic anions with pentafluorobenzyl methanesulphonate for GC 404(1987)175

Determination of oxidizable inorganic anions by

HPLC with fluorescence detection and application to the determination of salivary nitrite and thiocyanate and serum thiocyanate 424(1988)29

Detection of aminophenols and aromatic amines on thin-layer plates containing nitrite 435(1988)256

Polymeric pentafluorobenzylating reagent for GC of inorganic anions with electron-capture detection 438(1988)253

**1-Nitroalkanes**

Improved standardization in reversed-phase HPLC using 1-nitroalkanes as a retention index scale 435(1988)43

**Nitroarenes**

Simultaneous clean-up of nitroarenes and polycyclic aromatic ketones from soil and particulate matter on XAD-2 435(1988)489

**N-Nitroaroylamino acids**

Micro-LC and the chiral recognition mechanism on albumin-coated silica gel. Large selectivity changes with sample size 447(1988)297

**Nitrobenzaldehyde**

Retention behaviour of aromatic aldehyde derivatives in reversed-phase LC 404(1987)242

**Nitrobenzenes**

MS measurements of the negative ions formed in the thermionic ionization detector with nitrogen carrier gas 409(1987)139

**6-Nitrobenzo[a]pyrene**

Chromatographic and spectral investigations on the *in vivo* metabolites of 6-nitrobenzo[a]pyrene 450(1988)325

**O-*p*-Nitrobenzyl-N,N'-diisopropylisourea**

Determination of artesunic acid after chemical derivatization with O-*p*-nitrobenzyl-N,N'-diisopropylisourea by HPLC and UV absorption 410(1987)217

**Nitrobenzylthioinosine**

Demonstration of the adenosine reservoirs with nitrobenzylthioinosine in liver and diaphragm by HPLC 426(1988)249

**Nitrodiphenylamines**

Determination of nitrodiphenylamines by LC and dual-electrode amperometric detection 410(1987)437

**5-Nitrofurylacrylic acid**

Determination of 5-nitrofurylacrylic acid in wines by HPLC 445(1988)264

**Nitrogen**

Flame ionization and flame thermionic detection of carbon and nitrogen in aquatic lipid and humic-type classes with an Iatroscan Mark IV 435(1988)350

**Nitrogen-containing bases**

Separation of basic drug substances by reversed-phase HPLC on dynamically modified silica and on bonded-phase materials 409(1987)71

**Nitrogen-containing compounds**

Correlation between molar refraction and GC retention data 403(1987)11

Some problems in the correlation of molecular parameters and the connectivity index 403(1987)19

Application of the Snyder displacement model to aromatic nitrogen compounds on an aminopropyl bonded-phase column 405(1987)107

Comparison of volatile components between fresh and rotten mussels by GC-MS 409(1987)251

GC-MS study of coal-derived asphaltenes. II. Carbazole-type neutral nitrogen compounds 446(1988)293

**Nitroglycerin**

Improved method for the trace analysis of explosives by silica capillary column GC with thermal energy analysis detection 410(1987)181

Determination of the two dinitrate metabolites of nitroglycerin in human plasma by capillary GC with electron-capture detection 413(1987)101

Determination of nitroglycerin in human plasma using bonded-phase capillary column GC with electron-capture detection 425(1988)391

Simultaneous determination of nitroglycerin and its dinitrate metabolites by capillary GC with electron-capture detection 426(1988)259

**1-4-(4-Nitrophenylaminophenyl-thiocarbamido-4-methyl)piperazine**

HPLC of antifilarials. I. Simultaneous determination of 1-4-(4-nitrophenyl-aminophenyl thiocarbamido-4-methyl)piperazine and three of its metabolites in biological fluids 434(1988)288

**Nitrophenylaminopropanedione**

HPLC determination of chloramphenicol and

four analogues using reductive and oxidative electrochemical and UV detection 417(1987)111

**p-Nitrophenylhydrazones**

Retention behaviour of aromatic aldehyde derivatives in reversed-phase LC 404(1987)242

**2-Nitropropane**

HPLC method for the determination of 2-nitropropane in rat plasma 442(1988)436

**Nitrosamines**

Determination of nitrosamines by LC with post-column photolysis and electrochemical detection 410(1987)145

**Nitrosochloramphenicol**

HPLC determination of chloramphenicol and four analogues using reductive and oxidative electrochemical and UV detection 417(1987)111

**Nitroso compounds**

Comparative study of HPLC with UV detection and GC with the Hall electrolytic conductivity detector in the analysis of N-nitroso compounds 407(1987)355

**N-Nitrosodimethylamine**

Determination of nitrosamines by LC with post-column photolysis and electrochemical detection 410(1987)145

**Nitrosoareas**

Qualitative and quantitative analysis of a new sulphur-containing nitrosoarea in blood by HPLC 419(1987)381

**Nitrous oxide**

Quantification of halothane, enflurane and nitrous oxide by means of a simple GC method 403(1987)350

Automated analysis of the gas mixture from ethylene epoxidation by oxygen and nitrous oxide mixtures using GC 450(1988)399

**Nivalenol**

HPLC method for determining trichothecene mycotoxins by post-column fluorescence derivatization 410(1987)427

**Nomenclature**

Letter symbols in the chemistry of glyceridic fats and oils 410(1987)509

**Nonactin**

Preparative HPLC of macrotetrolides 442(1988)431

**Non-globular proteins**

Use of the Pharmacia Phast System for sodium dodecyl sulphate polyacrylamide gel electrophoresis of non-globular proteins: application to collagens 431(1988)474

**Non-histone chromatin proteins**

Non-histone chromatin proteins that recognize specific sequences of DNA (Review) 418(1987)73

**Non-linear elution effects**

Non-linear elution effects in split-peak chromatography. I. Computer simulations for the cases of irreversible diffusion- and adsorption-limited kinetics 436(1988)111

**Nonylphenol**

Determination of linear alkylbenzenesulphonates, alkylphenol polyethoxylates and nonylphenol in waste water by HPLC after enrichment on octadecylsilica 403(1987)243

**Noradrenaline**

Separation of norepinephrine enantiomers, DOPA enantiomers and dopamine derivatized with *o*-phthalaldehyde-N-acetyl-L-cysteine by HPLC 402(1987)387

Determination of  $\beta$ -phenylalkylamines in pharmaceutical preparations by analytical capillary isotachopheresis 405(1987)389

Simultaneous LC determination of seventeen of the major monoamine neurotransmitters, precursors and metabolites. I. Optimization of the mobile phase using factorial designs and a computer program to predict chromatograms 415(1987)261

Simultaneous LC determination of seventeen of the major monoamine neurotransmitters, precursors and metabolites. II. Assessment of human brain and cerebrospinal fluid concentrations 415(1987)275

Determination of catecholamines in biological tissues by LC with coulometric detection 415(1987)388

Improved HPLC method with electrochemical detection for the determination of urinary catecholamines using a new electrochemical detector 416(1987)340

Combined ion-pair extraction and HPLC for the determination of the biogenic amines and their major metabolites in single brain tissue samples 421(1987)51

Routine determination of urinary free catecholamines by HPLC with electrochemical detection 424(1988)263

LC determination of urinary catecholamines after one-step alumina extraction 424(1988)373

Species-dependent differences in recovery of 3,4-dihydroxybenzylamine in assays of plasma catecholamines 430(1988)123

Sensitive method for determination of picogram amounts of epinephrine and other catecholamines in microdissected samples of rat brain using LC with electrochemical detection 433(1988)41

Simultaneous LC determination of norepinephrine, 3,4-dihydroxyphenylethylene glycol, 3,4-dihydroxyphenylalanine and 3,4-dihydroxyphenylacetic acid in human plasma 434(1988)187

Perfluorinated acids as ion-pairing agents in the determination of monoamine transmitters and some prominent metabolites in rat brain by HPLC with amperometric detection 449(1988)191

**Norantipyrene**

Improved HPLC assay of antipyrene, hydroxymethylantipyrene, 4-hydroxyantipyrene and norantipyrene in urine 433(1988)305

**2-Norbornenes**

HPLC analysis of the *endo-exo* isomer ratio in the 5-substituted 2-norbornene series 445(1988)268

**Nordiazepam**

Use of micellar mobile phases and microbore column switching for the assay of drugs in physiological fluids 427(1988)277

**Nordihydrocapsaicin**

GC of methyl derivatives of naturally occurring capsaicinoids 409(1987)223

**Nordoxepin**

Simultaneous determination of doxepin and nordoxepin in serum using HPLC 419(1987)445

**Norendimide**

HPLC separation of noreximide and norendimide in bulk material, tablets and animal feed 404(1987)233

**Norepinephrine**

see Noradrenaline

**Norethindrone**

Selective on-line trace enrichment for the determination of ethynyl steroids in urine by LC with precolumn technology 417(1987)159

**Noreximide**

HPLC separation of noreximide and norendimide in bulk material, tablets and animal feed 404(1987)233

**Norfenfluramine**

Enantioselective GC assay with electron-capture detection for *dl*-fenfluramine and *dl*-norfenfluramine in plasma 433(1988)105

**Norfloxacin**

Assay of ciprofloxacin and norfloxacin in serum and urine by HPLC 416(1987)207

HPLC determination of norfloxacin in human tissues and plasma with fluorescence detection 417(1987)135

**Norfloxacin**

Application of HPLC of some antibiotics in clinical microbiology 427(1988)257

**Norgallopamil**

Quantification of gallopamil and norgallopamil in human plasma by HPLC with fluorescence detection 414(1987)492

**Norgestrel**

Selective on-line trace enrichment for the determination of ethynyl steroids in urine by LC with precolumn technology 417(1987)159

Some aspects of the selection of HPLC methods for the separation of chiral compounds in pharmaceutical analysis 450(1988)145

**Norharman**

Measurement of  $\beta$ -carbolines by HPLC with fluorescence detection 428(1988)229

HPLC identification of simple  $\beta$ -carboline alkaloids in specimens of *Heliconiini* butterflies 447(1988)432

**Norharmol**

Measurement of  $\beta$ -carbolines by HPLC with fluorescence detection 428(1988)229

**Normetanephrine**

Simultaneous LC determination of seventeen of the major monoamine neurotransmitters, precursors and metabolites. I. Optimization of the mobile phase using factorial designs and a

computer program to predict chromatograms 415(1987)261

Simultaneous LC determination of seventeen of the major monoamine neurotransmitters, precursors and metabolites. II. Assessment of human brain and cerebrospinal fluid concentrations 415(1987)275

Combined ion-pair extraction and HPLC for the determination of the biogenic amines and their major metabolites in single brain tissue samples 421(1987)51

**Normorphine**

Improved method for the simultaneous determination of morphine and its principal glucuronide metabolites 430(1988)394

**Norphenephrine**

Determination of  $\beta$ -phenylalkylamines in pharmaceutical preparations by analytical capillary isotachopheresis 405(1987)389

**Norpseudoephedrine**

Determination of  $\beta$ -phenylalkylamines in pharmaceutical preparations by analytical capillary isotachopheresis 405(1987)389

**Norsolorinic acid**

HPLC procedure for determining the profiles of aflatoxin precursors in wildtype and mutant strains of *Aspergillus parasiticus* 441(1988)400

**11-Nor- $\Delta^9$ -tetrahydrocannabinol-9-carboxylic acid**

Confirmation of cannabis abuse by the determination of 11-nor- $\Delta^9$ -tetrahydrocannabinol-9-carboxylic acid in urine with HPLC and electrochemical detection 414(1987)187

**Nortriptyline**

Applications of column-switching technique in biopharmaceutical analysis. I. HPLC determination of amitriptyline and its metabolites in human plasma 416(1987)99

HPLC quantitation of amitriptyline and nortriptyline in dialysate from plasma or serum using on-line solid-phase extraction 432(1988)363

Effects of drug-protein binding on trace enrichment of drugs in blood plasma on short precolumns 439(1988)353

**Norverapamil**

GC analysis of verapamil and norverapamil in



*post mortem* specimens using a 530- $\mu$ m fused-silica column with nitrogen-phosphorus detection 402(1987)361

#### ***cis*-Norzeatin riboside**

*E*-Isozeatin and *cis*-norzeatin riboside as internal standards for cytokinin analyses 436(1988)93

#### **Noscapine**

Separation of eight alkaloids and meconic acid and quantitation of five principal alkaloids in gum opium by gradient reversed-phase HPLC 436(1988)455

#### **Nuclear magnetic resonance spectrometry**

Chromatographic and solid state NMR study of the changes in reversed-phase packings for HPLC at different eluent compositions 436(1988)345

Cyclodextrin-chiral substrate interactions. High field NMR and molecular graphics studies 450(1988)175

#### **Nuclease**

Staphylococcal nuclease HPLC characterization of diaminoctane-modified DNA and its biotin and fluorescein derivatives 444(1988)97

#### **Nucleic acids**

Isolation of high-molecular-weight nucleic acids for copy number analysis using HPLC 402(1987)189

Effect of phosphate concentration on the yield and purity of DNA separated from RNA by hydroxyapatite chromatography 411(1987)481

Resolution of RNA using HPLC (Review) 418(1987)51

Non-histone chromatin proteins that recognize specific sequences of DNA (Review) 418(1987)73

Chromatographic separation of DNA restriction fragments (Review) 418(1987)97

Lanthanide ions as luminescent chromophores for the LC detection of polynucleotides and nucleic acids 436(1988)299

High-resolution chromatography of nucleic acids on the Gen-Pak FAX column 444(1988)47

Purification of transfer RNA species by single-step ion-exchange HPLC 444(1988)79

Preparative preparation of homologues of oligoribouridylic acid by selective partial hydrolysis of RNA (in German) 445(1988)155

#### **Nucleobases**

see also Purines and pyrimidines

Identification by HPLC and MS of a guanine derivative arising from acid hydrolysis of DNA 407(1987)349

Oxidation-elimination of a DNA base from its nucleoside to facilitate determination of alkyl chemical damage to DNA by GC with electrophore detection 410(1987)453

Technique for detection of DNA nucleobases by reversed-phase HPLC optimized for quantitative determination of thymidine substitution by iododeoxyuridine 417(1987)57

Determination of nucleotides, nucleosides and nucleobases in cells of different complexity by reversed-phase and ion-pair HPLC 421(1987)257

Demonstration of the adenosine reservoirs with nitrobenzylthioinosine in liver and diaphragm by HPLC 426(1988)249

Separation of synthetic cycloalkylated bases, nucleosides and nucleotides by reversed-phase HPLC 430(1988)309

Isocratic HPLC method for studying the metabolism of blood plasma pyrimidine nucleosides and bases: concentration and radioactivity measurements 434(1988)111

Optimization of the ion-pair HPLC separation of purine derivatives in erythrocytes, thymocytes and liver mitochondria 434(1988)447

Reversed-phase HPLC of pyrimidine bases and nucleosides. Application of solvophobic theory 445(1988)59

#### **Nucleoside monophosphates**

Single-step HPLC analysis of nucleoside monophosphates for ribonuclease specificity 446(1988)187

#### **Nucleosides**

Adsorption isotherms on HPLC sorbents. I. Peptides and nucleic acid constituents on octadecyl-silica 406(1987)275

Rapid separation of anomeric purine nucleosides by TLC on a chiral stationary phase 410(1987)226

Oxidation-elimination of a DNA base from its nucleoside to facilitate determination of alkyl chemical damage to DNA by GC with electrophore detection 410(1987)453

HPLC analysis of fluoropyrimidine nucleosides and fluorouracil in plasma 414(1987)197

HPLC method for the determination of 9-(1,3-dihydroxy-2-propoxymethyl)guanine in human plasma 414(1987)429

Comparison of high-performance ion-exchange and ion-pair LC methods for the separation of tumour cell nucleotides 415(1987)53

Application of HPLC and thermospray HPLC-MS to the analysis and identification of 2',3'-dideoxyadenosine and its metabolite in biological media 420(1987)1

Determination of nucleotides, nucleosides and nucleobases in cells of different complexity by reversed-phase and ion-pair HPLC 421(1987)257

HPLC determination of 5-azacytidine in plasma 421(1987)387

Demonstration of the adenosine reservoirs with nitrobenzylthioinosine in liver and diaphragm by HPLC 426(1988)249

Separation of synthetic cycloalkylated bases, nucleosides and nucleotides by reversed-phase HPLC 430(1988)309

Determination of 2',3'-dideoxyadenosine, 2',3'-dideoxyinosine and 2',3'-dideoxycytidine in biological samples 431(1988)184

Isocratic HPLC method for studying the metabolism of blood plasma pyrimidine nucleosides and bases: concentration and radioactivity measurements 434(1988)111

Optimization of the ion-pair HPLC separation of purine derivatives in erythrocytes, thymocytes and liver mitochondria 434(1988)447

Photodiode array detection of peptide-nucleoside antibiotics 436(1988)259

Reversed-phase HPLC of pyrimidine bases and nucleosides. Application of solvophobic theory 445(1988)59

#### **5'-Nucleotidase**

Assay of 5'-nucleotidase and simultaneous detection of interfering enzymes by anion-exchange CC 420(1987)398

#### **Nucleotide-linked peptide**

Purification of nucleotide-linked peptide 444(1988)133

#### **Nucleotides**

Determination of photosynthetic intermediates in leaves by HPLC 402(1987)283

Reversed-phase HPLC system for the rapid, automated purification of oligonucleotides 402(1987)381

Determination of the electrophoretic order of 26 nucleotides by isotachopheresis 403(1987)336

Adsorption isotherms on HPLC sorbents. I. Peptides and nucleic acid constituents on octadecyl-silica 406(1987)275

Adsorption chromatography on cellulose. I. Salting-out chromatography of organic compounds 406(1987)389

Oxidation-elimination of a DNA base from its nucleoside to facilitate determination of alkyl chemical damage to DNA by GC with electrophore detection 410(1987)453

Comparison of high-performance ion-exchange and ion-pair LC methods for the separation of tumour cell nucleotides 415(1987)53

Sensitive on-line radioactivity measurements with a heterogeneous flow-cell: application to lymphoid cell ribonucleotides separated by HPLC 415(1987)383

Determination of nucleotides, nucleosides and nucleobases in cells of different complexity by reversed-phase and ion-pair HPLC 421(1987)257

Assay of 6-thioinosinic acid and 6-thioguanine nucleotides, active metabolites of 6-mercaptopurine, in human red blood cells 423(1987)169

Isocratic ion-pair reversed-phase HPLC analysis of ATP and ADP 423(1987)308

Reversed-phase LC determination of cyclic adenosine 3',5'-monophosphate in rat brain cortex 426(1988)177

Demonstration of the adenosine reservoirs with nitrobenzylthioinosine in liver and diaphragm by HPLC 426(1988)249

Separation of synthetic cycloalkylated bases, nucleosides and nucleotides by reversed-phase HPLC 430(1988)309

Optimization of the ion-pair HPLC separation of purine derivatives in erythrocytes, thymocytes and liver mitochondria 434(1988)447

Nucleotide extraction and quantitation from tomato roots by HPLC 444(1988)183

New reversed-phase HPLC column for oligonucleotide separation 445(1988)225

Single-step HPLC analysis of nucleoside monophosphates for ribonuclease specificity 446(1988)187

#### **Nylon filters**

Anomalous LC results originating from the presence of soap residues and the use of nylon filters 408(1987)296

**Nystose**

Isolation of kestoses and nystose from enzyme digests by HPLC 408(1987)393

**Obstruction factor**

Does the obstruction factor change with flow-rate? 436(1988)

**Ochratoxin A**

Standardized HPLC of 182 mycotoxins and other fungal metabolites based on alkylphenone retention indices and UV-VIS spectra (diode array detection) 404(1987)195

**cis,cis-Octadecadienoates**

Equivalent chain-lengths of methyl ester derivatives of fatty acids on GC. A reappraisal 447(1988)305

**cis-Octadecenoates**

Equivalent chain-lengths of methyl ester derivatives of fatty acids on GC. A reappraisal 447(1988)305

**Octadecyl polymer gel**

Development of a C<sub>18</sub>-bonded vinyl alcohol copolymer gel for reversed-phase HPLC 410(1987)129

**trans-(−)-5,5a,6,7,8,9,9a,10-octahydro-6-propylpyrimido[4,5-g]quinolin-2-amine**

HPLC assay of *trans*-(−)-5,5a,6,7,8,9,9a,10-octahydro-6-propylpyrimido[4,5-g]quinolin-2-amine dihydrochloride (LY163502), a potent D-2 agonist, in plasma 426(1988)438

**Octanoic acid**

Quantitative determination of octanoic acid by HPLC following derivatization with 4-bromomethyl-7-methoxycoumarin 411(1987)297

**Octopamine**

Simultaneous determination of plasma phenethylamine, phenylethanolamine, tyramine and octopamine by HPLC using derivatization with fluorecamine 427(1988)320

**Oestradiol**

Automated direct assay system for the measurement of sex steroid hormones in serum using HPLC 426(1988)33

Determination of oestrogens in pregnancy urine by HPLC with fluorescence detection 431(1988)249

HPLC determination of conjugated estrogens in tablets 450(1988)414

**Oestriol**

Determination of oestrogens in pregnancy urine by HPLC with fluorescence detection 431(1988)249

**Oestriol glucuronides**

Measurement of urinary estriol glucuronides during the menstrual cycle by HPLC 434(1988)61

**Oestrogen receptors**

Influence of column length and pore size on high-performance ion-exchange chromatography of estrogen and progesterin receptors 425(1988)293

High-performance hydrophobic interaction chromatography as a means of identifying estrogen receptors expressing different binding domains 444(1988)225

**Oestrogens**

Analysis of estrogen sulphate mixtures in pharmaceutical formulations by reversed-phase chromatography 402(1987)273

Clinical analysis of individual steroids by column LC (Review) 429(1988)155

Determination of oestrogens in pregnancy urine by HPLC with fluorescence detection 431(1988)249

Measurement of urinary estriol glucuronides during the menstrual cycle by HPLC 434(1988)61

Complete separation of nine equine oestrogens by HPLC 435(1988)385

Biological effect of guava oestrone on uterine tissue of mice 437(1988)316

Oestrogen content of pomegranate seeds 438(1988)438

HPLC determination of conjugated estrogens in tablets 450(1988)414

**Oestrone**

Automated direct assay system for the measurement of sex steroid hormones in serum using HPLC 426(1988)33

Determination of oestrogens in pregnancy urine by HPLC with fluorescence detection 431(1988)249

Biological effect of guava oestrone on uterine tissue of mice 437(1988)316

Oestrogen content of pomegranate seeds 438(1988)438

HPLC determination of conjugated estrogens in tablets 450(1988)414

#### **Oestrone sulphate**

Analysis of estrogen sulphate mixtures in pharmaceutical formulations by reversed-phase chromatography 402(1987)273

#### **Ofloxacin**

Quantification of the enantiomers of ofloxacin in biological fluids by HPLC 425(1988)153

HPLC determination of ofloxacin in plasma and urine 430(1988)192

Simultaneous determination of ofloxacin, fenbufen and felbinac in rat plasma by HPLC 431(1988)135

Method for the determination of ofloxacin, a quinolone carboxylic acid antimicrobial, by HPLC 431(1988)461

Determination of ofloxacin in plasma and urine by LC 434(1988)320

Separation of pyridone carboxylic acid enantiomers by HPLC using copper(II)-L-amino acid as the eluent 448(1988)439

#### **Ofloxacin**

Application of HPLC of some antibiotics in clinical microbiology 427(1988)257

#### **Oils**

GC determination of water in crude oils 403(1987)288

Letter symbols in the chemistry of glyceridic fats and oils 410(1987)509

Biomarker determinations in crude oils using a triple-stage quadrupole mass spectrometer 446(1988)3

Characterisation of volatile oil constituents with relatively long GC retention times on two stationary phases 449(1988)312

#### **Olefins**

Paraffins, olefins, naphthenes and aromatics analysis of selected hydrocarbon distillates using

on-line column switching HPLC with dielectric constant detection 437(1988)365

#### **Oligoethylene glycol nonylphenyl**

Methods for characterization of selectivity in reversed-phase LC. IV. Retention behaviour of oligomeric series 449(1988)361

#### **Oligoethylene glycols**

Methods for characterization of selectivity in reversed-phase LC. IV. Retention behaviour of oligomeric series 449(1988)361

#### **Oligogalacturonic acids**

Preparative chromatography of oligogalacturonic acids 449(1988)229

#### **Oligomers**

Removal of oligomers and *n*-mers from the Immobiline chemicals for isoelectric focusing 402(1987)105

#### **Oligonucleotides**

Reversed-phase HPLC system for the rapid, automated purification of oligonucleotides 402(1987)381

Resolution of RNA using HPLC (Review) 418(1987)51

Preparative fractionation of oligoriboguanulates by anion-exchange chromatography on TSK DEAE-5PW 442(1988)401

Chromatographic evaluation of large-pore and non-porous polymeric reversed phases 443(1988)143

Preparation and chromatographic use of 5'-fluorescent-labelled DNA probes 444(1988)67

New reversed-phase HPLC column for oligonucleotide separation 445(1988)225

Separation of oligonucleotides by high-performance ion-exchange chromatography on a non-porous ion exchanger 447(1988)212

#### **Oligo(oxyethylene) compounds**

Polarity of oligooxyethylene derivatives of alcohols, thio alcohols and alkylamines as measured by GC 450(1988)291

#### **Oligopeptides**

HPLC of N-terminal tyrosine-containing oligopeptides by pre-column fluorescence derivatization with hydroxylamine, cobalt(II) and borate reagents 411(1987)305

**Oligoriboguanylates**

Preparative fractionation of oligoriboguanylates by anion-exchange chromatography on TSK DEAE-5PW 442(1988)401

**Oligosaccharides**

Isolation of kestoses and nystose from enzyme digests by HPLC 408(1987)393

Isolation of the major O-glycosidically linked oligosaccharides obtained by alkaline borohydride degradation of human meconium glycoproteins 425(1988)35

Analysis of tissue glycoprotein sugar chains by two-dimensional HPLC mapping 434(1988)51

HPLC of gluco-oligomers on C<sub>18</sub>-bonded vinyl alcohol copolymer gel with alkaline eluents 436(1988)328

HPLC analysis of oligosaccharides. I. Separation on an ion-exchange stationary phase of low cross-linking 440(1988)281

HPLC of plant-derived oligosaccharides on a new cation-exchange resin stationary phase: HPX-22H 441(1988)382

Optimizing copper-bicinchoninate carbohydrate detection for use with water-elution HPLC: a technique to measure the major mono- and oligosaccharides in small pieces of wheat endosperm 449(1988)207

**Oligostyrenes**

Methods for characterization of selectivity in reversed-phase LC. IV. Retention behaviour of oligomeric series 449(1988)361

**Oligouridylic acids**

Preparative preparation of homologues of oligoribouridylic acid by selective partial hydrolysis of RNA (in German) 445(1988)155

**Olive pollen**

Purification of allergens by HPLC. IV. Purification of the allergen of olive pollen (*Olea europaea*) 403(1987)312

**Omeprazole**

Improved procedure for quantitation of omeprazole and metabolites using reversed-phase HPLC 426(1988)216

**On-column fluorimetric detection**

Signal enhancement by on-column fluorometric detection in HPLC using micropacked fused-silica columns 435(1988)319

**Open split interface**

Open split interfaces in capillary GC-MS. Yield and quantitative aspects 450(1988)388

**Opiates**

Identification of opiates in urine by capillary column GC of two different derivatives 422(1987)117

**Opium alkaloids**

Separation of opium alkaloids by carbon dioxide sub- and supercritical fluid chromatography with packed columns. Application to the quantitative analysis of poppy straw extracts 437(1988)351

**Optical errors**

Optical errors in an LC absorbance cell 448(1988)193

**Optimization**

Optimization of the reversed-phase HPLC separation of synthetic estrogenic and progestogenic steroids using the multi-criteria decision making method 410(1987)1

Improvement in the resolution of *o*-phthalaldehyde derivatized amino acids by applying gradient steepness optimization to five reversed-phase columns of different lengths and particle sizes 411(1987)121

Simultaneous LC determination of seventeen of the major monoamine neurotransmitters, precursors and metabolites. I. Optimization of the mobile phase using factorial designs and a computer program to predict chromatograms 415(1987)261

Multi-variate ranking function for optimizing separations 439(1988)173

Separation of proteins by reversed-phase HPLC. I. Optimizing the column 443(1988)363

Separation of proteins by reversed-phase HPLC. II. Optimizing sample pretreatment and mobile phase conditions 443(1988)381

Optimization of concentration overload in preparative LC 444(1988)275

Statistical optimization of a reversed-phase ion-pair LC method for the analysis of tolmetin sodium in dosage forms 448(1988)1

Analysis of the information in a preparative chromatogram for further optimization of the operating conditions 450(1988)13

**ORF 17910**

High-performance liquid chromatographic determination of a new H<sub>2</sub> blocker (ORF 17910) in plasma 423(1987)358

**Organic acids**

see also Carboxylic acids

GC determination of C<sub>1</sub>-C<sub>5</sub> low-molecular-weight organic acids in alkaline permanganate oxidation products of humic substances 403(1987)225

Excretion of urinary acids during inverted sleep-waking rhythm 422(1987)1

HPLC measurement of selected blood citric acid cycle intermediates 425(1988)47

New ion-exclusion phase for organic acids 439(1988)71

**Organic compounds**

Adsorption chromatography on cellulose. I. Salting-out chromatography of organic compounds 406(1987)389

Simple steam distillation for sample preparation prior to GC determination of organic compounds 408(1987)201

Analysis of ppb levels of organics in water by means of purge-and-trap, capillary GC and selective detectors 441(1988)73

**Organic contaminants**

HPLC method for isolating organic contaminants from tissue and sediment extracts 437(1988)161

**Organic modifier**

Effect of organic modifier on the separation and efficiencies of homologues on porous polymer columns 407(1987)19

**Organochlorine compounds**

Analysis of organochlorine pesticides, polychlorinated biphenyls, dibenzo-*p*-dioxins and dibenzofurans in human milk by extraction with the lipophilic gel Lipidex 5000 422(1987)103

Clean-up of environmental samples by HPLC for analysis of organochlorine compounds by GC with electron-capture detection 435(1988)241

Separation of organochlorines on alumina 437(1988)254

**Organomercury chelates**

Separation of inorganic and organomercury chelates on a polystyrene-divinylbenzene copolymer column 438(1988)414

**Organometallic compounds**

Holistic molecular conformation and total surface area calculations as predictors for solution properties and chromatographic parameters 403(1987)1

**Organophosphorus compounds**

Analysis of-alkyl methylphosphonic acids in aqueous matrices by ion-pair reversed-phase ion chromatography 407(1987)399

Determination of organophosphorus pesticides in apples and water by GLC with electron-capture detection 437(1988)249

Determination of organophosphate pesticides and carbaryl on paddy rice by reversed-phase HPLC 447(1988)249

**Organo-smectite**

Organo-modified smectite for the GC separation of various compounds 410(1987)121

**Organotin compounds**

Characterization of organotin species using microbore and capillary LC techniques with an epifluorescence microscope as a novel imaging detector 410(1987)393

A biochromatographic technique for the quantitative estimation of triphenyltin fungicides 435(1988)210

Total molecular surface areas as a predictor for reversed-phase HPLC in various organotin systems. Importance and applications of carbon hybridization summaries 441(1988)229

**Organilydimethylsiloxy-covered silica gel**

Packing density of organilydimethylsiloxy-covered silica gel and hydrolytic stability of the coating 440(1988)315

**Ornidazole**

Simple LC method for the determination of ornidazole and metronidazole in human serum 420(1987)228

**Ornithine**

Determination of agmatine, arginine, citrulline and ornithine by reversed-phase LC using automated pre-column derivatization with *o*-phthalaldehyde 425(1988)269

**Ornithine decarboxylase**

Rapid and sensitive assay of ornithine decarboxylase activity by HPLC of the *o*-phthalaldehyde derivative of putrescine 419(1987)291

**Orphenadrine**

Screening procedure for the detection of alkanolamine antihistamines and their metabolites in urine using computerized GC-MS 428(1988)43

**Orthophosphoric acid**

Complex-forming equilibria in isotachopheresis. VII. Evaluation of the ion-pair formation constants of phosphorus oxo acids with histidine and assessment of their separability 410(1987)279

**Ovalbumin**

Interaction of proteins with immobilized  $\text{Cu}^{2+}$ . Quantitation of adsorption capacity, adsorption isotherms and equilibrium constants by frontal analysis 403(1987)197

Analytical high-performance affinity chromatography of ovalbumin-derived glycopeptides on columns of concanavalin A- and wheat germ agglutinin-immobilized gels 438(1988)73

**Overlap**

Experimental affirmation of the statistical model of overlap 449(1988)41

**Overloading**

Influence of the relative concentrations of two components of a binary mixture on their overloaded elution profiles 440(1988)9

**Ovomucoid-conjugated column**

Application of an ovomucoid-conjugated column for the optical resolution of some pharmaceutically important compounds 408(1987)316

**OV phases**

Estimation of the polarity of stationary phases by proton nuclear magnetic resonance spectroscopy. Application to phenyl and methyl silicones (OV and SE series) 402(1987)323

**Oxalate**

Measurement of plasma oxalate by capillary GC and its validation by isotope dilution MS 433(1988)1

Determination of phosphate, sulfate and oxalate in urine by ion chromatography 433(1988)373

Quantitation of oxalate and citrate by ion chromatography with a buffered, strong acid eluent 437(1988)231

**Oxalic acid**

Profiling of oxalic acid and  $\alpha$ -keto acids in blood

and urine by LC with electrochemical detection at a chemically modified electrode 414(1987)161

Determination of ionic metabolites from ethylene glycol in human blood by isotachopheresis 416(1987)111

**Oxaloacetate**

HPLC measurement of selected blood citric acid cycle intermediates 425(1988)47

**Oxamniquine**

Computer-aided optimisation of drug enantiomer separation in chiral HPLC 434(1988)377

**Oxantrazole**

HPLC assay for the experimental anticancer agent oxantrazole 419(1987)225

**Oxazepam**

Microsample determination of diazepam and its three metabolites in serum by reversed-phase HPLC 416(1987)212

Simultaneous determination of diazepam and its active metabolites in rat serum, brain and cerebrospinal fluid by HPLC 422(1987)334

Use of micellar mobile phases and microbore column switching for the assay of drugs in physiological fluids 427(1988)277

Computer-aided optimisation of drug enantiomer separation in chiral HPLC 434(1988)377

**Oxcarbapazine**

Simultaneous determination of carbamazepine, oxcarbapazine and their main metabolites in plasma by LC 414(1987)477

**Oxilofrine**

Detection of oxilofrine in plasma and urine by HPLC with electrochemical detection and comparison with GC-MS 428(1988)291

**Oxiracetam**

Determination of oxiracetam in human serum and urine by HPLC 416(1987)433

**Oxo acids**

O-(2,3,4,5,6-Pentafluorobenzyl)oxime-trimethylsilyl ester derivatives for quantitative GC and GC-MS studies of aldehydes, ketones and oxoacids 421(1987)336

Simple method for rapid quantification of branched-chain 2-oxo acids in physiological fluids as quinoxalinol derivatives by HPLC 432(1988)297

**[16,16,19-<sup>2</sup>H<sub>3</sub>; 9-<sup>3</sup>H]19-Oxo-androst-4-ene-3,17-dione**

Synthesis of [16,16,19-<sup>2</sup>H<sub>3</sub>; 19-<sup>3</sup>H]19-oxo-androst-4-ene-3,17-dione 440(1988)415

**4-Oxocyclophosphamide**

Combined TLC-photography-densitometry for the quantitation of cyclophosphamide and its four principal urinary metabolites 427(1988)121

**3-Oxodecanedioic acid**

Identification of 3-oxodicarboxylic acids in human urine 432(1988)13

**3-Oxodicarboxylic acids**

Identification of 3-oxodicarboxylic acids in human urine 432(1988)13

**3-Oxofloxacin**

HPLC of ciprofloxacin and its metabolites in serum, urine and sputum 422(1987)153

**3-Oxohexanedioic acid**

Identification of 3-oxodicarboxylic acids in human urine 432(1988)13

**4-Oxoisotretinoin**

Quantitative analysis of retinoids in biological fluids by HPLC using column switching. I. Determination of isotretinoin and tretinoin and their 4-oxo metabolites in plasma 424(1988)303

**Oxolinic acid**

Simultaneous determination of nalidixic acid, oxolinic acid and piromidic acid in fish by HPLC with fluorescence and UV detection 402(1987)301

**3-Oxo-octanedioic acid**

Identification of 3-oxodicarboxylic acids in human urine 432(1988)13

**17-Oxosteroid glucuronides**

Determination of 17-oxosteroid glucuronides and sulphates in urine by LC using 2,4-dinitrophenylhydrazine as a prelabelling reagent for spectrophotometric detection 415(1987)253

**17-Oxosteroids**

Determination of 17-oxosteroid glucuronides and sulphates in urine by LC using 2,4-dinitrophenylhydrazine as a prelabelling reagent for spectrophotometric detection 415(1987)253

**17-Oxosteroid sulphates**

Determination of 17-oxosteroid glucuronides and sulphates in urine by LC using 2,4-dinitrophenylhydrazine as a prelabelling reagent for spectrophotometric detection 415(1987)253

**4-Oxotretinoin**

Quantitative analysis of retinoids in biological fluids by HPLC using column switching. I. Determination of isotretinoin and tretinoin and their 4-oxo metabolites in plasma 424(1988)303

**Oxpentifylline**

Determination of oxpentifylline and three metabolites in plasma by automated capillary GC using nitrogen-selective detection 423(1987)139

**Oxprenolol**

Rapid determination of the enantiomers of metoprolol, oxprenolol and propranolol in urine 416(1987)331

Determination of oxprenolol in plasma by column LC 426(1988)229

**Oxybuprocaine**

Determination of six metabolites of oxybuprocaine in urine by HPLC 416(1987)189

**Oxybutinin chloride**

Determination of oxybutinin chloride in pharmaceuticals by reversed-phase ion-pair LC with two counter-ions in the eluent 450(1988)337

**Oxygen**

Automated analysis of the gas mixture from ethylene epoxidation by oxygen and nitrous oxide mixtures using GC 450(1988)399

**Oxymorphone**

Electrochemical determination of oxymorphone in rat plasma by ion-pair reversed-phase HPLC 413(1987)309

**Oxytetracycline**

Quantitative analysis of oxytetracycline and related substances by HPLC 405(1987)229

Determination of oxytetracycline in fish by HPLC 432(1988)381

**Oxytocin**

Quantitative analysis of oxytocin in pharmaceutical preparations by HPLC 402(1987)392

Determination of trace levels of oxytocin in pharmaceutical solutions by HPLC 410(1987)157

Quantitative LC determination of disulfide-containing peptide analogues of vasopressin with dual Hg/Au electrochemical detection 413(1987)43



# P

## Packed columns

Determination of the volumes of void zones in packed chromatographic columns 445(1988)37

## Packing density

Packing density of organyldimethylsiloxy-covered silica gel and hydrolytic stability of the coating 440(1988)315

## Paclobutrazol

Quantitation of triazole and pyrimidine plant growth retardants 438(1988)393

## Paeonol

Preparation and evaluation of mono-, di- and triphenyl-treated plates for the HPTLC analysis of puerarin and paeonol 407(1987)314

## Palladium complexes

LC retention of  $\beta$ -diketonato complexes of chromium(III) and palladium(II) in octadecyl-bonded silica/water-methanol systems. Comparison with liquid-liquid partition using dodecane as a model alkyl-bonded phase 439(1988)245

## Pancuronium

Quantitation of pancuronium, 3-desacetylpancuronium, vecuronium, 3-desacetylvecuronium, pipecuronium and 3-desacetylpipecuronium in biological fluids by capillary GC using nitrogen-sensitive detection 427(1988)41

## Papaverine

Isolation of basic drugs from plasma using solid-phase extraction with a cyanopropyl-bonded phase 432(1988)209

Separation of eight alkaloids and meconic acid and quantitation of five principal alkaloids in gum opium by gradient reversed-phase HPLC 436(1988)455

## Paper

Identification of polymers in paper by pyrolysis GC. Reduction of paper interference by prepyrolysis 407(1987)330

## Paper process impurities

SEC of deposits in pulp and paper mills 411(1987)363

## Parabens

Simplified description of HPLC separation under overload conditions, based on the Craig distribution model. V. Gradient elution separation 405(1987)51

HPLC determination of six *p*-hydroxybenzoic acid esters in cosmetics using Sep-Pak Florisil cartridges for sample pre-treatment 410(1987)413

HPLC assay of imazodan, methylparaben and propylparaben in imazodan injection 450(1988)407

## Paracetamol

Simultaneous analysis of acetaminophen, pseudoephedrine hydrochloride and chlorpheniramine maleate in a cold tablet using an isocratic, mixed micellar HPLC mobile phase 410(1987)206

Rapid HPLC assay of acetaminophen in serum and tissue homogenates 413(1987)151

Automated LC method for the determination of paracetamol and six metabolites in human urine 414(1987)355

LC determination of paracetamol and dextropropoxyphene in plasma 419(1987)433

Quantitation of acetaminophen and its metabolites in rat plasma after a toxic dose 426(1988)212

Complete separation of urinary metabolites of paracetamol and substituted paracetamols by reversed-phase ion-pair HPLC 426(1988)315

Extraction of acetaminophen and theophylline from post-mortem tissues and urine for HPLC analysis 439(1988)466

## Paraffins

Paraffins, olefins, naphthenes and aromatics analysis of selected hydrocarbon distillates using on-line column switching HPLC with dielectric constant detection 437(1988)365

## Parathyroid hormone

Separation of protein hormones (Review) 418(1987)245

## Paraxanthine

HPLC of xanthines. Effects of N-methyl and C-8 hydroxyl substitution on the retention times 428(1988)311

## Parfumine

Determination of alkaloids of *Fumaria parviflora* and *Fumaria capreolata* by HPLC and capillary isotachopheresis 445(1988)258

**Paroxetine**

Determination of paroxetine in human plasma, using HPLC with fluorescence detection 419(1987)438

**Parthenolide**

Isolation of parthenolide by droplet counter-current chromatography 446(1988)157

**Particle size**

Influence of pore and particle size on the frontal uptake of proteins. Implications for preparative anion-exchange chromatography 409(1987)111

**Partition coefficients**

Thermochemical investigations of GLC. IV. Partition coefficients of solutes on binary mixtures containing two complexing solvents 402(1987)41

Effect of polymer composition on the relative hydrophobicity of the phases of the biphasic system aqueous dextran-poly(ethylene glycol) 403(1987)123

Mixed stationary phases in GLC. Partition coefficients and retention indices in OV-101-OV-25, OV-101-carbowax 20M and OV-225-SP-2340 mixtures 410(1987)13

Structure parameters of molecules and media evaluated by chromatographic partition. II. Geometrical exclusion in gels 410(1987)233

Effect of hydrogen bonding on the HPLC behaviour of organic compounds. Relationship between capacity factors and partition coefficients 439(1988)227

Different partitioning behaviour of sulphonyl-containing compounds in reversed-phase HPLC and octanol-water systems 439(1988)237

LC retention of  $\beta$ -diketonato complexes of chromium(III) and palladium(II) in octadecyl-bonded silica/water-methanol systems.

Comparison with liquid-liquid partition using dodecane as a model alkyl-bonded phase 439(1988)245

Hydrophobicity studies on diadamantanyl derivatives. Comparison between calculated partition coefficients and experimental lipophilicity indices determined by reversed-phase HPLC and TLC 445(1988)317

Reversed-flow GC applied to physico-chemical measurements 446(1988)39

**Partition efficiency**

Cross-axis synchronous flow-through coil planet

centrifuge for large-scale preparative counter-current chromatography. II. Studies on partition efficiency in short coils and preparative separations with multilayer coils 449(1988)153

**Partitioning**

Partitioning behaviour of rat bone marrow cells in aqueous two-phase systems. Dependence of cell partition on the interfacial tension and electrical potential difference between the phases 403(1987)131

Effect of polymer structure on affinity partitioning of lactate dehydrogenase in polymer-water two-phase systems 411(1987)161

Different partitioning behaviour of sulphonyl-containing compounds in reversed-phase HPLC and octanol-water systems 439(1988)237

Partitioning behavior of erythrocytes in aqueous two-phase systems containing hydroxypropyl starch and polyethylene glycol 441(1988)261

Cell separation by immunoaffinity partitioning with polyethylene glycol-modified protein A in aqueous polymer two-phase systems 442(1988)219

Partitioning behaviour in aqueous two-phase systems and fractionation by counter-current distribution of chick embryo erythrocytes with numerical resolution of distribution curves 445(1988)327

**Partition isotherm**

Influence of the partition isotherm on the elution curve 438(1988)171

**Patulin**

Standardized HPLC of 182 mycotoxins and other fungal metabolites based on alkylphenone retention indices and UV-VIS spectra (diode array detection) 404(1987)195

**PC**

PC 445(1988)

**PCBs**

Directly coupled supercritical fluid extraction-GC analysis of polycyclic aromatic hydrocarbons and polychlorinated biphenyls from environmental solids 403(1987)63

Determinations of polychlorinated biphenyl isomers by response factor calibration 404(1987)155

Analysis of organochlorine pesticides, polychlorinated biphenyls, dibenzo-*p*-dioxins and

dibenzofurans in human milk by extraction with the lipophilic gel Lipidex 5000 422(1987)103

Clean-up of environmental samples by HPLC for analysis of organochlorine compounds by GC with electron-capture detection 435(1988)241

Quantitation of total *versus* selected polychlorinated biphenyl congeners in marine biota samples 442(1988)386

#### **Peak asymmetry**

Theoretical analysis of the asymmetry in chromatographic peaks 404(1987)1

#### **Peak distribution**

Multi-variate ranking function for optimizing separations 439(1988)173

#### **Peak enhancement**

Peak enhancement of carboxylic acids in ion-exclusion chromatography with a cation-exchange hollow-fibre suppressor using neutral and alkaline enhancers 435(1988)417

Ion-exclusion chromatography of carboxylic acids with conductivity detection. Peak enhancement using a cation-exchange hollow-fibre membrane and an alkaline solution 447(1988)373

#### **Peak maxima method**

Determination of adsorption isotherms from chromatographic measurements, using the peak maxima method 408(1987)1

#### **Peak overlap**

Real-time quantitative analysis of unresolved LC peaks with the use of the Kalman filter 407(1987)59

Resolution of overlapped chromatograms by the use of a Kalman filter. III. Effects of overlapping and outstripping of a solvent peak on the analysis 411(1987)95

Experimental affirmation of the statistical model of overlap 449(1988)41

#### **Peak shape**

Sample solvent effects in an apparent chiral HPLC separation on  $\beta$ -cyclodextrin 448(1988)31

#### **Peak splitting**

Non-linear elution effects in split-peak chromatography. I. Computer simulations for the cases of irreversible diffusion- and adsorption-limited kinetics 436(1988)111

#### **Pelanserin**

Sensitive HPLC assay of pelanserin, a novel antihypertensive agent, in plasma samples 428(1988)167

#### **Pemoline**

Determination of pemoline in plasma, urine and tissues by reversed-phase HPLC 430(1988)351

#### **Penems**

Application of HPLC of some antibiotics in clinical microbiology 427(1988)257

HPLC determination of the penem antibiotic (5*R*,6*S*)-2-aminomethyl-6-[(1*R*)-hydroxyethyl]-2-penem-3-carboxylic acid in human plasma and urine 427(1988)269

#### **D-Penicillamine**

Assay for D-penicillamine-proton conjugate in human plasma utilising chemical reduction followed by HPLC with gold/mercury electrochemical detection 430(1988)319

#### **D-Penicillamine-albumin disulphide**

Assay for D-penicillamine-proton conjugate in human plasma utilising chemical reduction followed by HPLC with gold/mercury electrochemical detection 430(1988)319

#### **Penicillin acid**

Standardized HPLC of 182 mycotoxins and other fungal metabolites based on alkylphenone retention indices and UV-VIS spectra (diode array detection) 404(1987)195

#### **Penicillinases**

Modified general affinity adsorbent for large-scale purification of penicillinases 448(1988)109

#### **Penicillin G**

Standardized HPLC of 182 mycotoxins and other fungal metabolites based on alkylphenone retention indices and UV-VIS spectra (diode array detection) 404(1987)195

#### **Penicillins**

Use of cyclodextrins in isotachopheresis. I. Effect of cyclodextrin on the isotachopheretic separation of related penicillins 405(1987)379

Hospital routine analysis of penicillins, third-generation cephalosporins and aztreonam by conventional and high-speed HPLC 413(1987)109

Determination of  $\beta$ -lactams and their biosynthetic intermediates in fermentation media by pre-column derivatisation followed by fluorescence detection 424(1988)325

Optimization of HPLC analysis for isoxazolyl penicillins using factorial design 425(1988)331

LC determination of penicillins by postcolumn degradation with sodium hypochlorite using an hollow-fibre membrane reactor 447(1988)365

#### **Penitrem A**

Standardized HPLC of 182 mycotoxins and other fungal metabolites based on alkylphenone retention indices and UV-VIS spectra (diode array detection) 404(1987)195

#### **Pentachlorophenol**

Analysis of pentachlorophenol in water and urine by enrichment with Lipidex 5000 414(1987)55

#### **Pentafluorobenzoylation**

Extractive pentafluorobenzoylation of formic, acetic, levulinic, benzoic and phthalic acids, studied by LC and dual-oven capillary GC 447(1988)329

#### **Pentafluorobenzyl methanesulphonate**

Derivatization of inorganic anions with pentafluorobenzyl methanesulphonate for GC 404(1987)175

#### **Pentafluoropropionic acid**

Use of perfluorinated carboxylic acids in the separation of metabolites of vasopressin prior to radioimmunoassay 430(1988)128

#### **Pentazocine**

Application of HPLC with electrochemical detection for monitoring the concentration of pentazocine in human blood 421(1987)425

#### **Pentobarbital**

Determination of sodium pentobarbital and either sodium methohexital or sodium thiopental in plasma by HPLC with ultraviolet detection 414(1986)484

#### **Pentoxifylline**

HPLC determination of pentoxifylline and its hydroxy metabolite in human plasma 416(1987)183

HPLC analysis of pentoxifylline and 1-(5'-hydroxyhexyl)-3,7-dimethylxanthine in whole blood 419(1987)368

#### **2-Pentuloses**

Degradation of ketoses during aqueous HPLC on lead-form cation-exchange resins 437(1988)387

#### **Pentylene-tetrazol**

HPLC determination of chemical convulsant pentylene-tetrazol in rat serum, cerebrospinal fluid and brain 432(1988)370

#### **N-(2-Pentyl)trifluoroacetamide**

Optical resolution of samples with weak interactions on chiral "brush-type" LC stationary phases 440(1988)197

#### **Pentylurofuran acid**

Excretion of urinary acids during inverted sleep-waking rhythm 422(1987)1

#### **Pepsins**

Separation of human pepsins in gastric juice by high-performance ion-exchange chromatography 417(1987)391

#### **Peptidases**

Assay for enkephalin-degrading peptidases in rat brain tissues by HPLC with on-line post-column fluorescence detection 430(1988)291

#### **Peptide-N-glycosidase**

Rapid chromatographic assay procedure for peptide-N-glycosidase activity 408(1987)378

#### **Peptide mapping**

HPLC in protein sequence determinations 440(1988)231

Rapid peptide mapping by HPLC 443(1988)343

#### **Peptide-nucleosides**

Photodiode array detection of peptide-nucleoside antibiotics 436(1988)259

#### **Peptides**

Chiral separations of amino acids using di- and tri-peptides for zwitter-ion pair chromatography 402(1987)49

Isolation and quantitation of metallothionein isoforms using reversed-phase HPLC 402(1987)243

Adsorption isotherms on HPLC sorbents. I. Peptides and nucleic acid constituents on octadecyl-silica 406(1987)275

Capillary zone electrophoresis and isotachopheresis as alternatives to chromatographic methods for purity control of synthetic peptides 407(1987)363

Reversed-phase HPLC of protected peptide segments 409(1987)281

Isolation of haem-bearing peptides of haem

- proteins by the use of adsorption chromatography 410(1987)463
- New metal chelate adsorbent selective for proteins and peptides containing neighbouring histidine residues 411(1987)177
- Counter-current chromatographic purification of [D-Ala<sup>1</sup>]peptide T amide 411(1987)431
- Quantitative LC determination of disulfide-containing peptide analogues of vasopressin with dual Hg/Au electrochemical detection 413(1987)43
- HPLC separation of peptide and amino acid stereoisomers 414(1987)313
- Combined use of size-exclusion and reversed-phase HPLC for characterization of  $\beta$ -endorphin processing pathways 416(1987)255
- HPLC-radioimmunoassay method for the measurement of angiotensin II peptides in human plasma 417(1987)27
- Use of *tert*-butyldimethylsilyl derivatives for GC-MS analysis of dipeptides 419(1987)263
- On-line post-column fluorescence detection for N-terminal tyrosine-containing peptides in HPLC 421(1987)245
- Reversed-phase HPLC-radioreceptor assay of human cerebrospinal fluid neuropeptides. Maximizing recovery of picomoles of peptides and minimizing memory 422(1987)61
- Combined HPLC-radioimmunoassay method for the analysis of endorphins, enkephalins and other neurotransmitter peptides 423(1987)93
- Procedure for the isolation of a large peptic fragment of human serum albumin 424(1988)13
- HPLC separation of peptides possessing a proline residue in the amino-terminal penultimate position, and their products generated by enzymatic hydrolysis 424(1988)129
- Sensitive method of detection, quantitation and purification of peptides using pre-column derivatization with phenyl isothiocyanate 425(1988)77
- HPLC determination of leucine-enkephalin-like peptide in rat brain by pre-column fluorescence derivatization involving formylation followed by reaction with 1,2-diamino-4,5-dimethoxybenzene 430(1988)271
- Simultaneous analysis of methionine- and leucine-enkephalin from rat brain: quantification by LC-electrochemistry 431(1988)65
- Reversed-phase HPLC separation and quantitation of reticulocyte  $\alpha$ - and  $\beta$ -globin polypeptide chains from normal and  $\beta$ -thalassemic mice 431(1988)177
- Optimization of HPLC-radioimmunoassay protocols for the analyses of substance P and some of its metabolic fragments 432(1988)113
- Simultaneous determination of histidine-containing dipeptides, histamine, methylhistamine and histidine by HPLC 432(1988)315
- Quantitative evaluation of collagen fractions separated by acrylamide gel electrophoresis: its application for collagen typing in normal and pathological tissues 433(1988)81
- Separation of peptides by multimode chromatography on a column packed with vinyl alcohol copolymer gel 434(1988)71
- Isolation and analysis of peptidic fragments of  $\alpha$ -gliadin using reversed-phase HPLC 434(1988)429
- HPLC of amino acids, peptides and proteins. LXXXV. Evaluation of the use of hydrophobicity coefficients for the prediction of peptide elution profiles 438(1988)197
- Displacement chromatography of biomolecules 439(1988)341
- Characterization of wild-type and mutant forms of yeast iso-1-cytochrome *c* by reversed-phase HPLC tryptic peptide mapping 439(1988)408
- Synthetic peptide substrates as models to study a pro-oxytocin/neurophysin converting enzyme 440(1988)439
- Inoculation of BALB/c mice against *Leishmania major* infection with *Leishmania*-derived antigens isolated by gel filtration 440(1988)459
- Prediction of peptide retention times 442(1988)69
- Determination of the chiral purity of dipeptide isosteres containing a reduced peptide bond by GC analysis 442(1988)165
- High-performance immobilized-metal-ion affinity chromatography of peptides and proteins 443(1988)3
- Peptide characterization with a sulfoethyl aspartamide column 443(1988)63
- Cation-exchange chromatography of peptides on poly(2-sulfoethyl aspartamide)-silica 443(1988)85
- Chromatographic evaluation of large-pore and

non-porous polymeric reversed phases  
443(1988)143

Effects of reductive alkylation on peptide  
retention times 443(1988)247

HPLC of complex mixtures of cyanogen  
bromide-produced peptides from different  
proteins 443(1988)317

Microbore HPLC-MS for the analysis of  
proteolytic digests by continuous-flow fast-atom  
bombardment MS 443(1988)355

Application of Marfey's reagent in racemization  
studies of amino acids and peptides 444(1988)115

HPLC analysis of arginine-containing peptides  
in biological fluids by means of a selective post-  
column reaction with fluorescence detection  
444(1988)123

Purification of nucleotide-linked peptide  
444(1988)133

Hydrophobic interaction chromatography of  
peptides as an alternative to reversed-phase  
chromatography 444(1988)269

Preparative separation of peptide and protein  
samples by HPLC with gradient elution. I. The  
Craig model as a basis for computer simulations  
444(1988)303

Preparative separation of peptide and protein  
samples by HPLC with gradient elution. II.  
Experimental examples compared with theory  
444(1988)325

Preparative purification of peptides by reversed-  
phase chromatography. Sample displacement  
mode *versus* gradient elution mode 444(1988)349

Rapid analysis of proteins and peptides by  
reversed-phase chromatography with polymeric  
micropellicular sorbents 445(1988)71

Chromatographic separation and partial  
characterization of microbial low-molecular-  
weight proteinase inhibitors 446(1988)203

Tyrosylation and purification of peptides for  
radioiodination 447(1988)341

Analysis and purification of toxic peptides from  
cyanobacteria by reversed-phase HPLC  
448(1988)275

The peptide toxin of the cyanobacterium  
*Microcystis aeruginosa* PCC 7941. Isolation and  
analysis by NMR and fast atom bombardment  
MS 449(1988)423

Preparative HPLC of peptides on a new

reversed-phase packing material, Kromasil C<sub>18</sub>  
450(1988)71

Chromatographic resolution of dipeptide  
enantiomers and diastereomers on chiral  
stationary phases from poly(L-leucine) or  
poly(L-phenylalanine) 450(1988)271

Rapid preparative-scale purification of peptides  
derived from caseins using mass ion-exchange  
chromatography 450(1988)420

### Peptides, cyclic

Adsorption of cyclic peptides analogous to  
gramicidin S and graptisin onto octadecylsilica  
stationary phase and bacterial cells 413(1987)287

### [D-Ala<sup>1</sup>]Peptide T amide

Counter-current chromatographic purification of  
[D-Ala<sup>1</sup>]peptide T amide 411(1987)431

Reversed-phase preparative chromatography of  
[D-Ala<sup>1</sup>]-peptide T amide 444(1988)345

### Peptide toxins

Rapid analysis of peptide toxins in cyanobacteria  
438(1988)93

### Peptidoglycan

Determination of environmental levels of  
peptidoglycan and lipopolysaccharide using GC  
with negative-ion chemical-ionization MS  
utilizing bacterial amino acids and hydroxy fatty  
acids as biomarkers 431(1988)1

### Peptidoglycan monomer

Comparative susceptibility of a peptidoglycan  
monomer from *Brevibacterium divaricatum* and  
its anhydromuramyl analogue to hydrolysis with  
N-acetylmuramyl-L-alanine amidase. Isolation  
and characterization of anhydromuramyl-  
peptidoglycan monomer 440(1988)405

### Peptidoglycan-polysaccharide

Analysis of the amino acid and sugar composition  
of streptococcal cell walls by GC-MS  
441(1988)323

### Perchloric acid extracts

New method of purification of perchloric acid  
extracts for quantitative GC of some Krebs cycle  
acids 424(1988)357

### Perfluorinated acids

Perfluorinated acids as ion-pairing agents in the  
determination of monoamine transmitters and  
some prominent metabolites in rat brain by

HPLC with amperometric detection  
449(1988)191

#### **Perfluorinated carboxylic acids**

Mechanism of selectivity in ion-pair HPLC of aminoglycoside antibiotics using perfluorinated pairing ions 404(1987)53

#### **Perfluorocarbon tracers**

Femtogram detection of perfluorocarbon tracers using capillary GC–electron-capture negative ion chemical ionisation MS 445(1988)119

#### **Perfluoro-1,3-dimethylcyclohexane**

Femtogram detection of perfluorocarbon tracers using capillary GC–electron-capture negative ion chemical ionisation MS 445(1988)119

#### **Perfluoromethylcyclohexane**

Femtogram detection of perfluorocarbon tracers using capillary GC–electron-capture negative ion chemical ionisation MS 445(1988)119

#### **Permanent gases**

Use of silicalite molecular sieve for the GC determination of permanent gases and volatile hydrocarbons emitted during coal processing 405(1987)213

#### **Permeation absorber**

Simple permeation absorber for sampling and preconcentrating hazardous air contaminants 439(1988)283

#### **Permethylated $\beta$ -cyclodextrin**

Resolution of chiral barbiturates into enantiomers by reversed-phase HPLC using methylated  $\beta$ -cyclodextrins 436(1988)381

#### **Peropyrene**

Elution behaviour of peropyrene–type polycyclic aromatic hydrocarbons in LC 402(1987)173

#### **Peroxidases**

Improved method for the detection of peroxidase isoenzymes after electrophoretic separation with chlorophyllin, cysteamine and light as a hydrogen peroxide generating system 410(1987)466

Reaction of Fast Blue BB salt and *p*-dimethylaminocinnamaldehyde with indole-3-acetic acid degradation products. Evaluation of the coupling properties of the reagents used in the zymographic screening of indole-3-acetic acid oxidases 411(1987)476

#### **Peroxides**

LC determination of serum lipid peroxides as

diphenylthiobarbituric acid-reactive substance to collagen disease patients—a preliminary study 417(1987)371

#### **Perylene**

Resolution of overlapped chromatograms by the use of a Kalman filter. III. Effects of overlapping and outstripping of a solvent peak on the analysis 411(1987)95

#### **Pesticides**

Pre-column clean-up and LC determination of residues of *N*-methylcarbamate pesticides in extracts of total diet 404(1987)352

LC of dibutylidithiocarbamate degradation products 404(1987)359

New method of visualizing thin-layer chromatograms by using liquid crystals 405(1987)203

Rapid determination of simazine and atrazine in water at the parts per trillion ( $10^{12}$ ) level. Extraction by a miniaturized Carbo-pack B trap followed by HPLC 405(1987)357

Capillary GC separation and MS detection of cyromazine and its metabolite melamine 408(1987)335

Analysis of bromacil, diuron and 3,4-dichloroaniline in contaminated well water, using a HPLC column-switching procedure 410(1987)211

Comparison of packed column and capillary column SFC and HPLC using representative herbicides and pesticides as typical moderate polarity and molecular weight range molecules 410(1987)343

Analysis of organochlorine pesticides, polychlorinated biphenyls, dibenzo-*p*-dioxins and dibenzofurans in human milk by extraction with the lipophilic gel Lipidex 5000 422(1987)103

Supercritical fluid extraction coupled with SFC for the separation of sulfonyleurea herbicides and their metabolites from complex matrices 435(1988)63

Determination of dodecylbenzenesulphonates and ethoxylated alkylphenols in liquid pesticide formulations by HPLC 435(1988)73

Determination of diastereoisomers of bromadiolone, an anticoagulant rodenticide, in animal tissues by HPLC 435(1988)83

A biochromatographic technique for the

quantitative estimation of triphenyltin fungicides 435(1988)210

Clean-up of environmental samples by HPLC for analysis of organochlorine compounds by GC with electron-capture detection 435(1988)241

Studies on fungicidal activity of N-phenylcarbamates. IV. Determination of the hydrophobicity of N-phenylcarbamates by HPLC 436(1988)316

Determination of organophosphorus pesticides in apples and water by GLC with electron-capture detection 437(1988)249

GC-MS investigation of phenoxypropanoic acid derivatives possessing herbicidal activity 438(1988)55

Thermospray LC-MS method for the analysis of metribuzin and its metabolites 438(1988)359

Analysis of the herbicides dicamba, clopyralid and bromacil in asparagus by HPLC 439(1988)470

CAPA—computer aided pesticide analysis. Computer program for the automated evaluation of chromatographic data for residue analysis of foods 441(1988)213

HPLC of *s*-triazines and their degradation products using UV photometric and amperometric detection 442(1988)147

HPLC determination of aryl N-methylcarbamate residues using post-column hydrolysis electrochemical detection 442(1988)333

Determination of carbaryl by HPLC with electrochemical detection 442(1988)451

Determination of organophosphate pesticides and carbaryl on paddy rice by reversed-phase HPLC 447(1988)249

Determination of diphenylamine residues in apples, and 4-aminobiphenyl residues in diphenylamine, by HPLC and electrochemical detection 447(1988)421

LC method for the determination of cyanazine in the presence of some normal soil constituents 447(1988)426

TLC of flucythrinate residues 450(1988)452

### **Petroleum**

On-line microbore HPLC capillary GC-MS. II. Application to the analysis of solvent refined coal 409(1987)213

### **Petroleum fuel**

GC determination for forensic purposes of petroleum fuel inhaled just before fatal burning 424(1988)49

### **Petroleum hydrocarbons**

Analysis of hydrocarbons in aquatic sediments. II. Evaluation of common preparative procedures for petroleum and chlorinated hydrocarbons 436(1988)503

### **Petroleum liquids**

Microdistillation: a method combining GC and LC to characterize petroleum liquids 403(1987)159

### **Phalloidin**

HPLC method for the determination of  $\alpha$ -amanitin and phalloidin in human plasma using the column-switching technique and its application in suspected cases of poisoning by the green species of amanita mushroom (*Amanita phalloides*) 425(1988)121

### **Phase ratio**

Phase ratio determination in an ion-exchange column having pores partially accessible to proteins 442(1988)15

### **pH effects**

Influence of mobile phase pH on HPLC column loading capacity. Implications for the design of preparative protein separations 437(1988)429

### **Phencyclidine**

Rapid confirmation of enzyme-multiplied immunoassay test (EMIT) for phencyclidine-positive urine samples with capillary gas chromatography-nitrogen-phosphorus detection 427(1988)151

### **Phenelzine**

GC analysis of monoalkylhydrazines 423(1987)131

### **Phenetole**

Resolution of overlapped chromatograms by the use of a Kalman filter. III. Effects of overlapping and outstripping of a solvent peak on the analysis 411(1987)95

### **Phenformin**

Fluorometric determination of biguanides in serum by HPLC with reagent-containing mobile phase 430(1988)65



**Pheniprazine**

GC analysis of monoalkylhydrazines  
423(1987)131

**Phenobarbital**

Differences in the constitutive forms of hepatic cytochrome P-450 in male and female adult beagle dogs 419(1987)61

Quantitation of 1-( $\beta$ -D-glucopyranosyl)-phenobarbital in plasma and bile 419(1987)421

Simultaneous determination of phenytoin and phenobarbital in human serum by electrochemical immunoassay combined with HPLC 426(1988)449

Use of micellar mobile phases and microbore column switching for the assay of drugs in physiological fluids 427(1988)277

**Phenobarbitone**

Simultaneous quantitation of phenobarbitone and *p*-hydroxyphenobarbitone and their perdeuteroethyl analogues in biological fluids by combined GC-MS 420(1987)287

**Phenol**

Determination of trace levels of phenol and cresols in rain by continuous liquid-liquid extraction and HPLC 403(1987)233

Determination of phenol by HPLC with fluorimetric detection. Application to wines (in French) 407(1987)384

Highly sensitive assay for alkaline and acid phosphatase activities by HPLC with electrochemical detection 417(1987)41

HPLC of mixtures of phenol or guaiacol and their reaction products with glyoxylic acid 437(1988)185

Separation of *m*- and *p*-cresol and other methylated phenols in shale oil by reversed-phase LC with electrochemical detection 441(1988)347

**Phenolic acid dimers**

Identification of phenols, phenolic acid dimers, and monosaccharides by GLC on a capillary column 436(1988)484

**Phenolic acids**

Mobility of Fe(III)-phenolic acid complexes in TLC 402(1987)358

Coextraction of 5-(hydroxymethyl)-2-furaldehyde with phenolic acids in acid-hydrolyzed plant extracts 410(1987)474

Separation of natural products by centrifugal partition chromatography 450(1988)91

**Phenolic antioxidants**

Determination of phenolic antioxidants in JP-5 jet fuels by GC-mass selective detection 449(1988)440

**Phenolic compounds**

Rapid and simple dansylation of phenolic steroids using a two-phase system and phase transfer catalysis 436(1988)429

Separation of naturally occurring mixtures of phenolic compounds from various Prunus tissues by reversed-phase HPLC 436(1988)490

**Phenolic glucosides**

Comparative HPLC and GLC determination of phenolic glucosides in Salicaceae species 442(1988)175

**Phenolic resins**

Gel filtration chromatography of resol phenolic resins 445(1988)371

**Phenolic steroids**

Rapid and simple dansylation of phenolic steroids using a two-phase system and phase transfer catalysis 436(1988)429

**Phenolphthalein**

Determination of cyclodextrins in biological fluids by HPLC with negative colorimetric detection using post-column complexation with phenolphthalein 415(1987)325

Screening procedure for stimulant laxatives in urine using HPLC with diode array detection 433(1988)131

**Phenols**

Ion chromatography of inorganic anions on a dynamically modified polystyrene-divinylbenzene packing material and its application to anion screening by sequential UV absorbance and electrochemical detection 402(1987)115

Adsorption chromatography on cellulose. I. Salting-out chromatography of organic compounds 406(1987)389

Comparison of volatile components between fresh and rotten mussels by GC-MS 409(1987)251

Determination of simple phenols in faeces and urine by HPLC 431(1988)143

Identification of polychlorinated phenols in urine by GC and TLC 431(1988)431

Identification of phenols, phenolic acid dimers, and monosaccharides by GLC on a capillary column 436(1988)484

HPLC of eleven priority substituted phenols by isocratic elution 438(1988)429

Adsorption chromatography on cellulose. II. Separations of aromatic amino acids, biogenic amines, alkaloids, dyes and phenols and determination of hydrophobic constants 440(1988)165

LC determination of traces of phenols in air 442(1988)407

Use of retention temperatures for the identification of phenols separated on OV-1701 capillary columns using linear temperature-programmed GC 448(1988)203

#### **Phenol sulphotransferase**

Determination of phenol sulphotransferase activity by HPLC 424(1988)136

#### **Phenothiazines**

New HPLC method for fluphenazine and metabolites in human plasma 414(1987)504

Determination of low levels of phenothiazine sulfoxides in phenothiazine drug substances and formulations by TLC-second-derivative spectrofluorimetry 442(1988)363

#### **Phenothiazine sulfoxides**

Determination of low levels of phenothiazine sulfoxides in phenothiazine drug substances and formulations by TLC-second-derivative spectrofluorimetry 442(1988)363

#### **Phenoxyalkanoic acids**

GC determination of chlorinated phenols in the form of various derivatives 439(1988)307

#### **Phenoxypropanoic acids**

GC-MS investigation of phenoxypropanoic acid derivatives possessing herbicidal activity 438(1988)55

#### **Phenylacetic acid**

Determination of free and total phenylacetic and *p*- and *m*-hydroxyphenylacetic acids in human urine by LC with fluorimetric detection 414(1987)275

Rapid and simple procedure for the determination of urinary phenylacetic acid using

derivatization in aqueous medium followed by electron-capture GC 428(1988)140

Effects of long-term storage on the concentrations of the unconjugated acidic metabolites of the trace amines, indoleamines and catecholamines 433(1988)23

#### **Phenylacetic mustard**

HPLC of chlorambucil analogues 422(1987)315

#### **Phenylalanine**

Increased yield of the phenylthiocarbonyl-amino acids tyrosine, cysteine and phenylalanine using diethylether-based liquid-liquid extraction 403(1987)299

#### **Phenylalkanols**

Effect of organic modifier on the separation and efficiencies of homologues on porous polymer columns 407(1987)19

#### **$\beta$ -Phenylalkylamines**

Determination of  $\beta$ -phenylalkylamines in pharmaceutical preparations by analytical capillary isotachopheresis 405(1987)389

#### **Phenylbutazone**

Determination of association constants of  $\beta$ -cyclodextrin and  $\beta$ -cyclodextrin-bearing polymers with drugs by HPLC 409(1987)61

Rapid HPLC assay for the simultaneous analysis of non-steroidal anti-inflammatory drugs in plasma 416(1987)293

#### **N-Phenylcarbamates**

Studies on fungicidal activity of N-phenylcarbamates. IV. Determination of the hydrophobicity of N-phenylcarbamates by HPLC 436(1988)316

#### **Phenylephrine**

Determination of  $\beta$ -phenylalkylamines in pharmaceutical preparations by analytical capillary isotachopheresis 405(1987)389

#### **Phenylethanolamine**

Simultaneous determination of plasma phenethylamine, phenylethanolamine, tyramine and octopamine by HPLC using derivatization with fluorescamine 427(1988)320

#### **Phenylethylamine**

New HPLC procedure for the detection and quantification of  $\beta$ -phenylethylamine 415(1987)124

Simultaneous determination of plasma phenethylamine, phenylethanolamine, tyramine and octopamine by HPLC using derivatization with fluorescamine 427(1988)320

Rapid and simple procedure for the determination of urinary phenylacetic acid using derivatization in aqueous medium followed by electron-capture GC 428(1988)140

HPLC of biogenic amines, derivatized with 9-fluorenylmethyl chloroformate 437(1988)379

#### **(S)-(-)-Phenylethyl isocyanate**

Determination of metoprolol enantiomers in plasma and urine using (S)-(-)-phenylethyl isocyanate as a chiral reagent 421(1987)161

#### **Phenylethylmalonamide**

GC analysis of phenylethylmalonamide in human plasma 416(1987)281

#### **Phenylhydrazine**

Study of the side-reaction products of phenylhydrazine production by GC-MS 447(1988)202

#### **Phenylhydrazones**

Retention behaviour of aromatic aldehyde derivatives in reversed-phase LC 404(1987)242

#### **2-Phenyl-2-imidazoline**

Determination of 2-phenyl-2-imidazoline in air by HPLC with UV detection 438(1988)443

#### **Phenyl isothiocyanate**

Sensitive method of detection, quantitation and purification of peptides using pre-column derivatization with phenyl isothiocyanate 425(1988)77

#### **N-Phenyllinoleamide**

Determination of oxidation products of N-phenyllinoleamide: Spanish toxic oil syndrome studies 426(1988)83

#### **Phenylmercury salts**

Assay of phenylmercury salts in pharmaceutical products by HPLC of the morpholinedithiocarbamate derivative 407(1987)389

#### **Phenyl-methyl silicones**

Estimation of the polarity of stationary phases by proton nuclear magnetic resonance spectroscopy. Application to phenyl and methyl silicones (OV and SE series) 402(1987)323

#### **Phenyl-1-naphthylamine**

Chromatographic analysis of phenyl-1-naphthylamine and 4,4'-dioctyldiphenylamine and their intermediate oxidation products in oxidized lubricants 409(1987)325

#### **Phenylpropane derivatives**

GC characterization of frequently occurring aromatic compounds in essential oils 446(1988)267

#### **1-Phenyl-1-(2-pyridinyl)ethanol**

Formation of artifactual metabolites of doxylamine following acid hydrolysis 419(1987)113

#### **1-Phenyl-1-(2-pyridinyl)ethylene**

Formation of artifactual metabolites of doxylamine following acid hydrolysis 419(1987)113

#### **Phenyl-Sepharose CL-4B**

Determination of the leakage from Phenyl-Sepharose CL-4B, Phenyl-Sepharose FF and Phenyl-Superose in bulk and column experiments 403(1987)85

#### **Phenyl-Sepharose FF**

Determination of the leakage from Phenyl-Sepharose CL-4B, Phenyl-Sepharose FF and Phenyl-Superose in bulk and column experiments 403(1987)85

#### **Phenyl-Superose**

Determination of the leakage from Phenyl-Sepharose CL-4B, Phenyl-Sepharose FF and Phenyl-Superose in bulk and column experiments 403(1987)85

#### **6-Phenyl-2-thiouracil**

Identification of thyrostatic drug residues in animal thyroids by HPTLC and fluorescence reaction detection 442(1988)459

#### **Phenyltoloxamine**

Screening procedure for the detection of alkanolamine antihistamines and their metabolites in urine using computerized GC-MS 428(1988)43

#### **Phenytol**

Electrochemical immunoassay combined with column LC: determination of phenytol in human serum 417(1987)129

Simultaneous determination of phenytol and phenobarbital in human serum by

electrochemical immunoassay combined with HPLC 426(1988)449

#### **Pheophorbides**

Reversed-phase HPLC and column LC of chlorophylls and their derivatives 408(1987)181

#### **Pheophytins**

Reversed-phase HPLC and column LC of chlorophylls and their derivatives 408(1987)181

HPLC of chloroplast pigments. One-step separation of carotene and xanthophyll isomers, chlorophylls and pheophytins 448(1988)411

#### **pH gradients**

Experimental and theoretical dynamics of isoelectric focusing. II. Elucidation of the impact of the electrode assembly 436(1988)191

Computer-controlled linear pH gradient for HPLC fractionations of aromatic carboxylic acids and of humic and fulvic acids 439(1988)213

#### **Phloxine**

Simple method for the analysis of food dyes on reversed-phase thin-layer plates 411(1987)437

#### **Pholcodine**

Determination of pholcodine in biological fluids by HPLC with fluorescence detection 424(1988)170

#### **Phorcabilin**

Interactions between ascorbic acid and linear and cyclized bile pigments. A TLC study 440(1988)275

#### **Phosgene**

Derivatization of 2-amino alcohols with phosgene in aqueous media: limitations of the reaction selectivity as found in the presence of O-glucuronides of alprenolol in urine 413(1987)270

Phosgene as a derivatizing reagent prior to GC and LC (Review) 435(1988)259

#### **Phosmethylan**

HPLC methods for the determination of contaminants in technical phosmethylan 435(1988)380

#### **Phosphatases**

Highly sensitive assay for alkaline and acid phosphatase activities by HPLC with electrochemical detection 417(1987)41

#### **Phosphate**

Determination of potassium nitrate and sodium

monofluorophosphate in the presence of phosphate and sulfate by high-resolution ion chromatography 410(1987)494

Effect of phosphate concentration on the yield and purity of DNA separated from RNA by hydroxyapatite chromatography 411(1987)481

Determination of phosphate, sulfate and oxalate in urine by ion chromatography 433(1988)373

#### **Phosphatidic acid**

TLC of human platelet phospholipids with fatty acid analysis 420(1987)411

Sensitive HPLC method for the determination of phosphatidic acid 424(1988)367

#### **Phosphatidylcholine**

Separation of the phosphono analogues of phosphatidylcholine and phosphatidylethanolamine from related phospholipids by HPLC 403(1987)363

Method for the sensitive analysis of individual molecular species of phosphatidylcholine by HPLC using post-column fluorescence detection 415(1987)241

Comparative study of solid-phase and liquid-phase extraction techniques for isolation of phospholipids from plasma 419(1987)103

Identification of very-long-chain fatty acids in rat and mouse harderian gland lipids by capillary GC-MS 431(1988)231

Calibration of TLC with flame ionization detection for the analysis of natural lipid samples 435(1988)343

#### **Phosphatidylethanolamine**

Separation of the phosphono analogues of phosphatidylcholine and phosphatidylethanolamine from related phospholipids by HPLC 403(1987)363

Comparative study of solid-phase and liquid-phase extraction techniques for isolation of phospholipids from plasma 419(1987)103

TLC of human platelet phospholipids with fatty acid analysis 420(1987)411

Identification of very-long-chain fatty acids in rat and mouse harderian gland lipids by capillary GC-MS 431(1988)231

Calibration of TLC with flame ionization detection for the analysis of natural lipid samples 435(1988)343

**Phosphatidylglycerol**

TLC of human platelet phospholipids with fatty acid analysis 420(1987)411

**Phosphatidylinositol**

Comparative study of solid-phase and liquid-phase extraction techniques for isolation of phospholipids from plasma 419(1987)103

TLC of human platelet phospholipids with fatty acid analysis 420(1987)411

**Phosphatidylserine**

Comparative study of solid-phase and liquid-phase extraction techniques for isolation of phospholipids from plasma 419(1987)103

TLC of human platelet phospholipids with fatty acid analysis 420(1987)411

**Phosphine oxides**

Resolution of racemic amides and phosphine oxides on a  $\beta$ -cyclodextrin-bonded stationary phase by subcritical fluid chromatography 405(1987)135

Resolution of a series of racemic phosphine oxides by LC on two structurally similar chiral stationary phases (in French) 409(1987)183

**Phosphines**

HPLC separation of sulphonated phosphines 408(1987)416

**Phosphoamino acids**

Determination of [ $^{32}$ P]phosphoamino acids in protein hydrolysates by isocratic anion-exchange HPLC 442(1988)307

**6-Phosphogluconate dehydrogenase**

New strategies for the screening of a large number of immobilized dyes for the purification of enzymes. Application to the purification of enzymes from human haemolysate 449(1988)403

Purification of human 6-phosphogluconate dehydrogenase from human haemolysate with chromatography on an immobilized dye as the essential step and use of automation. Simultaneous purification of lactate dehydrogenase 449(1988)413

**Phosphoglycerate kinase**

Survey of the folding pathway of a two-domain protein phosphoglycerate kinase 440(1988)421

**Phosphoinositides**

Use of the dye Janus Green to facilitate

quantitative analysis of phosphoinositides 424(1988)445

**Phospholipase A**

Characterization of bee venom and its main components by HPLC 408(1987)341

**Phospholipids**

Separation of the phosphono analogues of phosphatidylcholine and phosphatidylethanolamine from related phospholipids by HPLC 403(1987)363

Method for determination of different phospholipids by circular HPTLC 413(1987)257

Method for the sensitive analysis of individual molecular species of phosphatidylcholine by HPLC using post-column fluorescence detection 415(1987)241

Comparative study of solid-phase and liquid-phase extraction techniques for isolation of phospholipids from plasma 419(1987)103

TLC of human platelet phospholipids with fatty acid analysis 420(1987)411

Phospholipids in amniotic fluid with special reference to the lecithin; Sphingomyelin ratio (Review) 429(1988)359

Separation of phospholipids and phosphonolipids of *Tetrahymena* by HPLC 431(1988)395

Highly sensitive GC-MS method for the determination of platelet-activating factor in human blood 433(1988)243

Highly sensitive determination and characterization of intact cellular ester-linked phospholipids using LC-plasma spray MS 434(1988)31

Behaviour of plasmalogens during HPLC on a silica column with a mobile phase containing phosphoric acid 434(1988)411

Separation of rat liver phospholipids from polar non-phospholipids by TLC 435(1988)513

Quantification of plasmalogen, alkylacyl and diacyl glycerophospholipids by micro-TLC 436(1988)467

Qualitative and quantitative TLC analysis of polar lipids from leaves of glanded and glandless cotton genotypes 437(1988)211

Simple and rapid method for HPLC separation and quantification of soybean phospholipids 447(1988)436

**Phosphonolipids**

Separation of phospholipids and phosphonolipids of *Tetrahymena* by HPLC 431(1988)395

**Phosphoramidate mustard**

Combined TLC-photography-densitometry for the quantitation of cyclophosphamide and its four principal urinary metabolites 427(1988)121

**Phosphoric acid**

Behaviour of plasmalogens during HPLC on a silica column with a mobile phase containing phosphoric acid 434(1988)411

**Phosphorous acid**

Complex-forming equilibria in isotachopheresis. VII. Evaluation of the ion-pair formation constants of phosphorus oxo acids with histidine and assessment of their separability 410(1987)279

**Phosphorus oxo acids**

Complex-forming equilibria in isotachopheresis. VII. Evaluation of the ion-pair formation constants of phosphorus oxo acids with histidine and assessment of their separability 410(1987)279

**N-Phosphoryl lombricine**

HPLC determination of lombricine and N-phosphoryl lombricine in the earthworm by pre-column fluorescence derivatisation with *o*-phthalaldehyde-ethanethiol 445(1988)433

**Photoionization detector**

Photoionization detector for open-tubular LC 411(1987)201

Analysis of ppb levels of organics in water by means of purge-and-trap, capillary GC and selective detectors 441(1988)73

Use of the photoionisation detector in packed-column supercritical fluid chromatography 445(1988)239

**Photometric detector**

General photometric detector for GC 409(1987)125

**Photosynthetic intermediates**

Determination of photosynthetic intermediates in leaves by HPLC 402(1987)283

***o*-Phthalaldehyde**

Automated pre-column derivatization of amino acids with *o*-phthalaldehyde by a reagent sandwiching technique 408(1987)217

***o*-Phthalaldehyde-N-acetyl-L-cysteine**

Separation of norepinephrine enantiomers,

DOPA enantiomers and dopamine derivatized with *o*-phthalaldehyde-N-acetyl-L-cysteine by HPLC 402(1987)387

**Phthalate esters**

Evaluation of disposable cartridges for trace enrichment from aqueous solutions 402(1987)201

**Phthalates**

Stepwise gradient elution using switching valves in micro HPLC 405(1987)117

**Phthalic acid**

Extractive pentafluorobenzoylation of formic, acetic, levulinic, benzoic and phthalic acids, studied by LC and dual-oven capillary GC 447(1988)329

**Phthiocerol dimycocerosates**

Concerning hydrolysis of mycolate esters, of phthiocerol dimycocerosates and of related mycobacterial lipids: an anecdotal account 440(1988)487

 **$\alpha$ -Phyllocladane**

GC-MS study of  $\alpha$ -phyllocladane dehydrogenation 446(1988)87

**Phylloquinone**

Determination of K vitamins (phylloquinone and menaquinones) in umbilical cord plasma by a platinum-reduction column 430(1988)143

**Physico-chemical measurements**

Reversed-flow GC applied to physico-chemical measurements 446(1988)39

**Physostigmine**

Evaluation of a chemical reaction detector with electrochemical production of bromine for the LC determination of physostigmine 411(1987)229

Determination of physostigmine in plasma by HPLC 415(1987)423

Metabolism of physostigmine in mouse liver microsomal incubations studied by LC with dual-electrode amperometric detection 419(1987)165

**Phytanic acid**

Rapid GC determination of phytanic acid from serum of a patient suffering from Refsum's disease 419(1987)475

**Phytoalexins**

Reversed-phase HPLC of *Cajanus cajan* phytoalexins 409(1987)355

HPLC of phytoalexins in stem callus tissues of rapeseed 448(1988)448

#### **Phytochemicals**

Isolation of biologically active plant constituents by LC 450(1988)3

#### **Pigments**

HPLC of carthamin, safflor yellow A and a precursor of carthamin. Application to the investigation of an unknown red pigment produced in cultured cells of safflower 438(1988)61

#### **Pimeprofen**

HPLC determination of pimeprofen and its metabolite ibuprofen in sheep plasma and lymph 414(1987)228

#### **Pimonidazole**

Gradient HPLC method for simultaneous assay of the radiosensitizers etanidazole (SR 2508) and pimonidazole (Ro 03-8799) in biological materials 420(1987)223

#### **Pinacolyl methylphosphonic acid**

Analysis of alkyl methylphosphonic acids in aqueous matrices by ion-pair reversed-phase ion chromatography 407(1987)399

#### **Pindolol**

Application of an ovomucoid-conjugated column for the optical resolution of some pharmaceutically important compounds 408(1987)316

HPLC method for the determination of pindolol in human plasma 415(1987)93

#### **$\alpha$ -Pinene**

Resolution of monoterpene enantiomers by GC 407(1987)243

Efficiency of adsorption-desorption of volatile compounds. Application to isoprene on Tenax GC 437(1988)139

#### **$\beta$ -Pinene**

Resolution of monoterpene enantiomers by GC 407(1987)243

#### **Pinifolic acid**

GC analysis of underivatized resin acids 448(1988)428

#### **Pinocembrin**

Direct enantiomeric separation of racemic

flavanones by HPLC using cellulose triacetate as a chiral stationary phase 441(1988)417

#### **Pipecuronium**

Quantitation of pancuronium, 3-desacetylpancuronium, vecuronium, 3-desacetylvecuronium, pipecuronium and 3-desacetylpipecuronium in biological fluids by capillary GC using nitrogen-sensitive detection 427(1988)41

#### **Pipemidic acid**

Simple method for the determination of pipemidic acid in biological fluids by HPLC 416(1987)409

#### **Piperacillin**

HPLC assay for piperacillin in aqueous humor of the eye 424(1988)177

Application of HPLC of some antibiotics in clinical microbiology 427(1988)257

HPLC determination of a new  $\beta$ -lactamase inhibitor and its metabolite in combination therapy with piperacillin in biological materials 431(1988)87

#### **Piperazine-2',5'-dione**

LC determination of amoxicillin and its metabolites in human urine by postcolumn degradation with sodium hypochlorite 413(1987)219

#### **Pirlimycin**

Ion-pairing LC method for the determination of pirlimycin hydrochloride 402(1987)335

#### **Piromidic acid**

Simultaneous determination of nalidixic acid, oxolinic acid and piromidic acid in fish by HPLC with fluorescence and UV detection 402(1987)301

#### **Piroxicam**

Rapid HPLC assay for the simultaneous analysis of non-steroidal anti-inflammatory drugs in plasma 416(1987)293

Rapid and sensitive method for determination of piroxicam in human plasma by HPLC 420(1987)445

HPLC assay for piroxicam in human plasma 430(1988)424

#### **Pirprofen**

Determination of the (+)- and (-)-enantiomers of pirprofen in human plasma by HPLC 414(1987)131

Assay method for product formation in *in vitro* enzyme kinetic studies of uridine diphosphate glucuronyltransferases: 2-arylpropionic acid enantiomers 430(1988)368

#### **Plant extracts**

HPLC characterization of some medical plant extracts used in cosmetic formulas 442(1988)353

#### **Plant growth retardants**

Quantitation of triazole and pyrimidine plant growth retardants 438(1988)393

#### **Plant hormones**

Resolution of the enantiomers of abscisic acid methyl ester by HPLC using a stationary phase of cellulose tris(3,5-dimethylphenylcarbamate)-coated silica gel 402(1987)371

Direct separation of (*S*)- and (*R*)-abscisic acid on a commercially available chiral HPLC column 402(1987)374

Determination of abscisic acid in small amounts of plant material 404(1987)346

Improved method for the determination of indole-3-acetic acid in plant tissue 407(1987)377

Separation of gibberellins by TLC and GC, and possibilities for their quantitative analysis 437(1988)287

Chromatographic resolution. XXI. Direct optical resolution of abscisic acid by HPLC on cellulose tris(3,5-dimethylphenylcarbamate) 448(1988)454

#### **Plant pigments**

Isocratic reversed-phase HPLC analysis of pigments in Norway spruce 448(1988)417

#### **Plasmalogen**

Behaviour of plasmalogens during HPLC on a silica column with a mobile phase containing phosphoric acid 434(1988)411

Quantification of plasmalogen, alkylacyl and diacyl glycerophospholipids by micro-TLC 436(1988)467

#### **Plasma protein binding**

Correlation between reversed-phase HPLC and plasma protein binding 421(1987)83

#### ***Plasmodium falciparum* proteins**

*Plasmodium falciparum* exoprotein stimulation of human T-lymphocytes unsensitized to malaria 440(1988)385

#### **Plastic containers**

GC quantification of underivatized amphetamine

in whole blood. Blood stored in plastic containers—a source of error 432(1988)290

#### **Plasticizers**

Analysis of the epoxidised soya bean oil additive in plastics by GC 437(1988)247

Analysis of the plasticiser acetyltributyl citrate in foods by stable isotope dilution GC-MS 437(1988)281

Determination of plasticizers in fat by GC-MS 447(1988)230

#### **Plastic packaging materials**

Analysis of flavor absorption into plastic packaging materials using multiple headspace extraction GC 438(1988)281

#### **Plate height**

Theoretical prediction of plate height in serially coupled capillary columns 406(1987)3

#### **Platelet-activating factor**

Characterization and quantification of PAF-acether (platelet-activating factor) as a heptafluorobutyrate derivative of 1-O-alkyl-2-acetyl-*sn*-glycerol by capillary column GC with electron-capture detection 423(1987)23

Highly sensitive GC-MS method for the determination of platelet-activating factor in human blood 433(1988)243

#### **Platelets**

Thermospray-MS analysis of underivatized monohydroxy fatty acids: application to stimulated platelets 425(1988)25

#### **Polar compounds**

Quantitative analyses of polar components in frying oils by the Iatroscan TLC-flame ionization detection technique 405(1987)371

#### **Polarity**

Polarity and hydrophile-lipophile balance of model hydroxyoximes and their intermediates as determined by reversed-phase GC 402(1987)55

Estimation of the polarity of stationary phases by proton nuclear magnetic resonance spectroscopy. Application to phenyl and methyl silicones (OV and SE series) 402(1987)323

Polarity of 1,3-bis[ $\omega$ -alkoxyoligo(oxyethylene)]propan-2-ols as measured by reversed-phase GC 409(1987)29

Dispersive force parameters as a measure of the polarity of surfactants used as stationary phases in GLC 435(1988)29



Polarity of some individual derivatives of  $\alpha,\omega$ -diamino oligoethers as measured by GC 448(1988)219

Polarity of oligooxyethylene derivatives of alcohols, thio alcohols and alkylamines as measured by GC 450(1988)291

#### **Polar strength**

Characterization of porous polymers by polar strength and selectivity 404(1987)145

#### **Pollen**

Analysis of allergenic components of a *Parietaria judaica* pollen extract by chromatographic methods for the evaluation of purification procedures 444(1988)209

Micropreparative-scale enrichment of some allergenic components of *Parietaria* pollen extract 446(1988)179

#### **Pollen allergens**

Purification of allergens by HPLC. VI.

Purification of the major allergen from *Parietaria judaica* pollen 407(1987)408

#### **Pollutants**

Specific identification of synthetic organic chemicals in river water using liquid-liquid extraction and resin adsorption coupled with electron impact, chemical ionization and accurate mass measurement GC-MS analyses 436(1988)229

#### **Polyacrylamide**

Steric protection of a modified glass surface with adsorbed polyacrylamide: interactions with blood proteins 428(1988)17

#### **Polyacrylonitrile-divinylbenzene**

GC properties of porous polyacrylonitrile and divinylbenzene copolymers 408(1987)308

#### **Polyamines**

Improved method for benzoyl chloride derivatization of polyamines for HPLC 408(1987)227

Determination of polyamines in human erythrocytes by capillary GC with nitrogen-phosphorus detection 415(1987)27

Simultaneous determination of  $\gamma$ -aminobutyric acid and polyamines by HPLC 419(1987)285

Quantitation of polyamines in cultured cells and tissue homogenates by reversed-phase HPLC of their benzoyl derivatives 426(1988)41

Reversed-phase LC separation and simultaneous fluorimetric detection of polyamines and their monoacetyl derivatives in human and animal urine, serum and tissue samples: an improved, rapid and sensitive method for routine application 430(1988)249

Chromatography, flow injection analysis and electrophoresis in computer-assisted comparative biochemistry: its application and possibilities in clinical research. Preliminary studies on Crohn's disease 440(1988)261

Quantitation of polyamines using TLC and image analysis 441(1988)361

Detection by HPLC of polyamines formed by clostridal putrefaction of caseins 442(1988)267

#### **Polychlorinated biphenyls**

see PCBs

#### **Polychlorinated dibenzo-*p*-dioxins**

Analysis of organochlorine pesticides, polychlorinated biphenyls, dibenzo-*p*-dioxins and dibenzofurans in human milk by extraction with the lipophilic gel Lipidex 5000 422(1987)103

Formation of polychlorinated dibenzo-*p*-dioxins upon combustion of commercial textile products containing 2,4,4'-trichloro-2'-hydroxydiphenyl ether (Irgasan DP300) 442(1988)289

#### **Polychlorinated dibenzofurans**

Analysis of organochlorine pesticides, polychlorinated biphenyls, dibenzo-*p*-dioxins and dibenzofurans in human milk by extraction with the lipophilic gel Lipidex 5000 422(1987)103

#### **Polychlorinated phenols**

Identification of polychlorinated phenols in urine by GC and TLC 431(1988)431

#### **Polycyclic aromatic hydrocarbons**

see Polynuclear aromatic hydrocarbons

#### **Polyelectrolytes**

SEC of polyelectrolytes. Experimental evidence for a general mechanism 441(1988)239

#### **Polyether alcohols**

Inverse GC study of polyether alcohols (in German) 410(1987)267

#### **Polyethylene**

GC determination of calcium stearate in polyethylene food packaging sheets 447(1988)198

**Polyethylene glycol**

Micro HPLC of poly(ethylene glycols) with indirect photometric detection 403(1987)324

Partitioning behavior of erythrocytes in aqueous two-phase systems containing hydroxypropyl starch and polyethylene glycol 441(1988)261

**Poly(ethylene glycol succinate)**

Polyester deactivation of pink supports and its effect on the GC of some polar, closely boiling and isomeric analytes 446(1988)31

**Polyglutamates**

Separation and identification of methotrexate and its metabolites, 7-hydroxymethotrexate and polyglutamates, in human tissues by reversed-phase HPLC coupled with radioimmunoassay 415(1987)445

**Polygodial**

Analytical separations of enantiomeric primary amines as pyrrole derivatives formed by reaction with polygodial 438(1988)108

**Poly(3-hydroxybutyrate)**

Preparation and chromatographic analysis of poly(3-hydroxybutyrate) hydrolysis products 445(1988)251

**Poly- $\beta$ -hydroxybutyric acid**

GC determination of poly- $\beta$ -hydroxybutyric acid in microbial biomass after hydrochloric acid propanolysis 445(1988)285

**Polyketides**

Standardized HPLC of 182 mycotoxins and other fungal metabolites based on alkylphenone retention indices and UV-VIS spectra (diode array detection) 404(1987)195

**Poly(L-leucine)**

Chromatographic resolution of dipeptide enantiomers and diastereomers on chiral stationary phases from poly(L-leucine) or poly(L-phenylalanine) 450(1988)271

**Polymer gels**

Internal structures of wide-pore packing materials for HPLC studied by transmission electron microscopy 448(1988)95

**Polymer glycols**

Identification of polymer glycols in polyurethane elastomers by pyrolysis-GC 435(1988)219

**Polymeric light stabilizers**

Determination of the polymeric light stabilizer Tinuvin 622 in polyolefins 450(1988)426

Determination of the polymeric light stabilizer Chimassorb 944 in polyolefins by reversed-phase HPLC 450(1988)430

**Polymers**

Normal- and reversed-phase separation of copolymers prepared from styrene and ethyl methacrylate 403(1987)280

Investigation by pyrolysis-GC of the composition of multicomponent polymeric microheterogeneous systems based on some vinyl monomers 404(1987)183

Use of acid or alkaline fusion for the quantitative degradation of condensation polymers 406(1987)167

Identification of polymers in paper by pyrolysis GC. Reduction of paper interference by prepolyrolysis 407(1987)330

Hydrolysed macroporous glycidyl methacrylate-ethylene dimethacrylate copolymer sorbent for size-exclusion HPLC of synthetic polymers and biopolymers 435(1988)357

Derivatization of some polymers with *o*-phthalaldehyde prior to high-performance SEC 436(1988)39

Gel filtration chromatography of resole phenolic resins 445(1988)371

Fundamentals of the theory and practice of polymer GPC as a method of chromatographic porosimetry (Review) 448(1988)307

Purge-and-trap GC method for residual volatiles in high-temperature polymers 450(1988)394

**Polymer structure**

Effect of polymer structure on affinity partitioning of lactate dehydrogenase in polymer-water two-phase systems 411(1987)161

**Polymer two-phase systems**

Distribution of inorganic salts between the coexisting phases of aqueous polymer two-phase systems. Interrelationship between the ionic and polymer composition of the phases 439(1988)267

**Polymethoxylated flavones**

HPLC determination of polymethoxylated flavones in orange juice after solid-phase extraction 439(1988)481

**Polyethylbenzenes**

GC study of the temperature dependence of the hydrogen bond energy of adsorption on silica 446(1988)55

**Polymyxins**

Analytical and preparative HPLC of the papain-cleaved derivative of polymyxin B 441(1988)423

**Polynuclear aromatic compounds**

Direct synchronous luminescence detection of co-eluting solutes in pseudophase LC 409(1987)147

**Polynuclear aromatic hydrocarbons**

Elution behaviour of peropyrene-type polycyclic aromatic hydrocarbons in LC 402(1987)173

Directly coupled supercritical fluid extraction-GC analysis of polycyclic aromatic hydrocarbons and polychlorinated biphenyls from environmental solids 403(1987)63

Simple preparation of a Poly-S 179 capillary column for the analysis of polycyclic aromatic hydrocarbons and triglycerides 404(1987)248

UV spectrometric detection for GC of polynuclear aromatic compounds with repetitive spectral scans using absorbance and concurrent fluorescence measurements 405(1987)191

Determination of hydroxylated metabolites of polycyclic aromatic hydrocarbons in urine 413(1987)227

Effect of column temperature on the retention of peropyrene-type polycyclic aromatic hydrocarbons on various chemically bonded stationary phases in reversed-phase LC 436(1988)1

Improved method for determining adsorption isotherms by LC on long and short fibrous material 436(1988)391

Main parameters governing the retention of polynuclear aromatic hydrocarbons on tetrachlorophthalimidopropyl-bonded silica 439(1988)195

Reversed-phase separation of a standard mixture of polycyclic aromatic hydrocarbons on an adamantyl surface 439(1988)414

Thermodynamic comparison of monomeric and polymeric C<sub>18</sub> bonded phases using aqueous methanol and acetonitrile mobile phases 442(1988)1

Determination of polycyclic aromatic compounds in fish tissue 449(1988)251

**Polynuclear aromatic ketones**

Simultaneous clean-up of nitroarenes and

polycyclic aromatic ketones from soil and particulate matter on XAD-2 435(1988)489

**Polynuclear aromatic nitrogen heterocycles**

Determination of polycyclic aromatic compounds in fish tissue 449(1988)251

**Polynuclear aromatic sulphur heterocyclics**

High-efficiency separation and characterization of high-molecular-weight heterocyclic sulfur compounds in fossil fuels by microcolumn LC and related techniques 406(1987)375

Determination of polycyclic aromatic compounds in fish tissue 449(1988)251

**Polynucleotides**

Lanthanide ions as luminescent chromophores for the LC detection of polynucleotides and nucleic acids 436(1988)299

**Polyol compounds**

Elution characteristics of polyol-borate eluents in anion-exchange chromatography. I. Eluent strength of sugar-, uronate-borate eluents 403(1987)27

**Polyolefins**

Determination of the polymeric light stabilizer Tinuvin 622 in polyolefins 450(1988)426

Determination of the polymeric light stabilizer Chimassorb 944 in polyolefins by reversed-phase HPLC 450(1988)430

**Polyols**

GC procedure for the qualitative analysis of silicone polyester-resins after alkaline fusion 441(1988)207

**Poly(pentafluorobenzyl p-styrenesulphonate)**

Polymeric pentafluorobenzylating reagent for GC of inorganic anions with electron-capture detection 438(1988)253

**Polypeptides**

High-performance SEC of polypeptides on a TSK G2000SW column in acidic mobile phases 404(1987)215

Recovery of polypeptides after reversed-phase HPLC 408(1987)191

Application of preparative reversed-phase HPLC to the isolation of insulin-like growth factor II from human serum 421(1987)61

Comparison of two methods for the

determination of substance P immunoreactivity in human cerebrospinal fluid: sulphopropyl-Sephadex ion-exchange chromatography and/or reversed-phase HPLC combined with radiomimmunoassay 425(1988)175

Column LC separation of collagen  $\alpha$ -chain polymers on sepiolite 426(1988)162

Fast protein LC of lymphocyte blastogenesis inhibitory factor produced by the human macrophage-like cell line U937 440(1988)131

Several intermediate forms in the processing of rat lutropin subunits as shown by sodium dodecyl sulphate polyacrylamide gel electrophoresis 440(1988)353

Correlation of calcitonin structure with chromatographic retention in HPLC 443(1988)229

Immunological-chromatographic analysis 444(1988)1

#### **Poly(L-phenylalanine)**

Chromatographic resolution of dipeptide enantiomers and diastereomers on chiral stationary phases from poly(L-leucine) or poly(L-phenylalanine) 450(1988)271

#### **Polyphosphates**

LC study of the hydrolysis reactions of cyclic and linear polyphosphates in aqueous solution 447(1988)91

#### **Polysaccharides**

Separation of starch components based on ligand-induced adsorption of amylose on cellulose 403(1987)373

Extracellular mammalian polysaccharides: glycosaminoglycans and proteoglycans (Review) 418(1987)187

#### **Poly-Sep 47**

Buffer isoelectric focusing revisited 440(1988)367

#### **Polysialogangliosides**

New solvent system for HPTLC and HPLC of gangliosides 405(1987)125

HPLC of underivatized gangliosides 408(1987)285

Improved method for large-scale purification of brain gangliosides by Q-Sepharose CC. Immunochemical detection of C-series polysialogangliosides in adult bovine brains 445(1988)377

#### **Polysiloxanes**

Estimation of the polarity of stationary phases by proton nuclear magnetic resonance spectroscopy. Application to phenyl and methyl silicones (OV and SE series) 402(1987)323

Factor analysis of trends in selectivity in GLC of 175 saturated and unsaturated esters on 14 polysiloxane stationary phases 404(1987)11

Polysiloxanes containing thermally stable chiral amide side-chains for capillary GC and supercritical fluid chromatography 405(1987)169

Hierarchical ascending classification of Kováts retention indices as a means of indicating siloxane stationary phase specificities towards aliphatic carboxylic esters 406(1987)95

#### **Poly(styrene-divinylbenzene)**

On the structure and chromatographic behaviour of polystyrene phases 406(1987)325

Synthesis of a non-porous, polystyrene-based strong anion-exchange packing material and its application to fast HPLC of proteins 443(1988)73

Chromatographic evaluation of large-pore and non-porous polymeric reversed phases 443(1988)143

#### **Poly(styrene-methyl methacrylate)**

Comparison between SEC and liquid adsorption chromatography in the determination of the chemical heterogeneity of copolymers 411(1987)355

#### **Poly(2-sulphoethyl aspartamide)-silica**

Cation-exchange chromatography of peptides on poly(2-sulfoethyl aspartamide)-silica 443(1988)85

#### **Polythiazide**

Automated column LC determination of polythiazide in human serum 417(1987)434

#### **Polyunsaturated fatty acids**

Analysis of tissue free fatty acids isolated by aminopropyl bonded-phase columns 428(1988)221

Comparative chromatographic study of modifications of brush-border membrane vesicles induced by an essential fatty acid-deficient diet 432(1988)75

Method for evaluating the bioconversion of radioactive polyunsaturated fatty acids by use of reversed-phase LC 448(1988)249

**Polyurethane foam**

High-volume elution chromatography of dichlorobenzenes on a polyurethane foam—Tenax sandwich cartridge 409(1987)235

**Polyurethanes**

Identification of polymer glycols in polyurethane elastomers by pyrolysis—GC 435(1988)219

**Poly(vinyl chloride)**

Thermal degradation of weldable poly(vinyl chloride) samples at low temperatures 445(1988)353

**Polyvinylimidazole-coated silica**

Rapid preparation of bovine mercaptalbumin by means of covalent chromatography on silica-based materials 408(1987)263

**Porapak**

Characterization of porous polymers by polar strength and selectivity 404(1987)145

**Porapak Q**

Modification to the chromatographic behaviour of Porapak Q by chemical reaction 437(1988)238

Characterization of some styrene-divinylbenzene sorbents for reversed-phase chromatography 442(1988)97

**Pore size**

Influence of pore and particle size on the frontal uptake of proteins. Implications for preparative anion-exchange chromatography 409(1987)111

Influence of column length and pore size on high-performance ion-exchange chromatography of estrogen and progesterin receptors 425(1988)293

**Pore structure**

On the structure and chromatographic behaviour of polystyrene phases 406(1987)325

Effect of the pore structure of silica gel on retention behaviour in pressure programming supercritical fluid chromatography 448(1988)87

**Porosimetry**

Fundamentals of the theory and practice of polymer GPC as a method of chromatographic porosimetry (Review) 448(1988)307

**Porosity**

Reversed-flow GC applied to physico-chemical measurements 446(1988)39

**Porous glass**

Internal structures of wide-pore packing

materials for HPLC studied by transmission electron microscopy 448(1988)95

**Porous polymer gels**

Pulse injection analysis of the binding of serum proteins to porous polymer gels modified with formyl groups 420(1987)13

**Porous polymers**

Characterization of porous polymers by polar strength and selectivity 404(1987)145

GC on porous polymers. IV. Influence of the geometric structure of porous copolymers of 1,4-di(methacryloyloxymethyl)naphthalene with 1,4-divinylbenzene on their chromatographic behaviour 448(1988)233

**Porphobilinogen**

Quantitative determination of porphyrins, their precursors and zinc protoporphyrin in whole blood and dried blood by HPLC with fluorimetric detection 417(1987)269

**Porphyrins**

Determination of porphyrins in tissue: pre-adsorption followed by HPLC 415(1987)129

Separation of metalloporphyrins from metallation reactions by LC and electrophoresis 416(1987)71

Quantitative determination of porphyrins, their precursors and zinc protoporphyrin in whole blood and dried blood by HPLC with fluorimetric detection 417(1987)269

Normal-phase HPLC of free acid dicarboxylic porphyrins and hematoporphyrin derivative on silica 422(1987)73

Reversed-phase HPLC of urinary porphyrin free acids 423(1987)313

HPLC of porphyrins (Review) 429(1988)123

Determination of erythrocyte porphyrins by reversed-phase HPLC using capsule-type silica gels coated with silicone polymer 433(1988)73

Analytical and preparative separation of uroporphyrin I and III by HPLC 437(1988)311

Detection of non-typical porphyrin isomers in human urines by ion-pair reversed-phase HPLC 441(1988)171

**Post-column extraction**

On-line post-column extraction in column LC with electron-capture detection. II. Applications 435(1988)297

**Post-column reactors**

Determination of reducing and non-reducing carbohydrates in food products by LC with post-column catalytic hydrolysis and derivatization. Comparison with refractive index detection 402(1987)127

HPLC separation of some mono- and disaccharides with detection by a post-column enzyme reactor and a chemically modified electrode 408(1987)157

Determination of nitrosamines by LC with post-column photolysis and electrochemical detection 410(1987)145

HPLC method for determining trichothecene mycotoxins by post-column fluorescence derivatization 410(1987)427

Evaluation of a chemical reaction detector with electrochemical production of bromine for the LC determination of physostigmine 411(1987)229

LC determination of amoxicillin and its metabolites in human urine by postcolumn degradation with sodium hypochlorite 413(1987)219

Column LC of methotrexate and its metabolites using a post-column photochemical reactor and fluorescence detection 419(1987)213

Analysis of conjugated bile acids in human serum by HPLC using post-column enzyme reaction and off-line fluorimetric determination 423(1987)75

Determination of acetylcholine and choline in microdialysates from spinal cord of rat using LC with electrochemical detection 423(1987)85

Determination of disulfiram and two of its metabolites in urine by reversed-phase LC and spectrophotometric detection after post-column complexation 424(1988)95

Fluorimetric determination of urinary  $\delta$ -aminolevulinic acid by HPLC and post-column derivatization 426(1988)365

Use of a post-column enzymatic reaction for detection of macroamylasemia by HPLC 428(1988)415

Lanthanide ions as luminescent chromophores for the LC detection of polynucleotides and nucleic acids 436(1988)299

Determination of iodide and bromide by ion chromatography with post-column reaction detection 439(1988)129

Stereoselective determination of L-amino acids using column LC with an enzymatic solid-phase reactor and chemiluminescence detection 440(1988)217

HPLC of hydroxysteroids detected with post-column immobilized enzyme reactors 441(1988)81

HPLC determination of aryl N-methylcarbamate residues using post-column hydrolysis electrochemical detection 442(1988)333

HPLC analysis of arginine-containing peptides in biological fluids by means of a selective post-column reaction with fluorescence detection 444(1988)123

LC determination of penicillins by postcolumn degradation with sodium hypochlorite using an hollow-fibre membrane reactor 447(1988)365

HPLC determination of glucosides (glucose conjugates) with post-column reaction detection combining immobilized enzyme reactors and luminol chemiluminescence 449(1988)217

**Potassium**

Isotachophoretic determination of the cationic constituents in mixtures of raw materials for glass manufacturing 439(1988)423

**Potassium hydroxide**

Non-suppressed ion chromatography of arsenic anions with potassium hydroxide-aromatic salt mixed eluents 447(1988)323

**Potassium nitrate**

Determination of potassium nitrate and sodium monofluorophosphate in the presence of phosphate and sulfate by high-resolution ion chromatography 410(1987)494

**Potency**

High-performance SEC determination of the potency of biosynthetic human growth hormone products 435(1988)307

**Practolol**

Isolation of basic drugs from plasma using solid-phase extraction with a cyanopropyl-bonded phase 432(1988)209

**Prajalium**

Column LC determination of prajmalium in plasma and urine by direct sample injection 415(1987)183

**Precolumn concentration**

Precolumn concentration and/or stepwise-gradient elution by switching valve techniques in micro HPLC 407(1987)141

**Precolumns**

Effects of drug-protein binding on trace enrichment of drugs in blood plasma on short precolumns 439(1988)353

**Precolumn switching system**

Direct determination of dextromethorphan and its three metabolites in urine by HPLC using a precolumn switching system for sample clean-up 422(1987)340

**Preconcentration methods**

Studies on sample preconcentration in ion chromatography. VI. Use of concentrator columns with "permanently" bound ion-interaction reagents 407(1987)121

Simple permeation absorber for sampling and preconcentrating hazardous air contaminants 439(1988)283

**Prednisolone acetate**

Analysis of prednisolone acetate and related corticoids in swine plasma by reversed-phase HPLC 425(1988)162

**Pregnenolone**

Pregnenolone production by adrenal mitochondria: a new HPLC analytical method for cholesterol side-chain cleavage 414(1987)411

**Pregnenolone-16 $\alpha$ -carbonitrile**

Differences in the constitutive forms of hepatic cytochrome P-450 in male and female adult beagle dogs 419(1987)61

**Prenylamine**

Simultaneous determination of *R*- and *S*-prenylamine in plasma and urine by reversed-phase HPLC 426(1988)305

**Preparative chromatography**

Preparative HPLC purification and structural determination of 1-O- $\alpha$ -D-glucopyranosyl-D-fructose (trehalulose) 402(1987)265

Rapid ion-exchange chromatography for preparative separation of proteins. III. Application to purification of bovine factor X and prothrombin 403(1987)383

Simplified description of HPLC separation under overload conditions, based on the Craig

distribution model. III. Computer simulations for two co-eluting bands assuming a Langmuir isotherm 405(1987)1

Simplified description of HPLC separation under overload conditions, based on the Craig distribution model. IV. Comparison of the model with experimental data for samples containing two or more solutes 405(1987)31

Simplified description of HPLC separation under overload conditions, based on the Craig distribution model. V. Gradient elution separation 405(1987)51

Large scale chromatographic purification of ribosome-inactivating proteins 408(1987)235

Influence of pore and particle size on the frontal uptake of proteins. Implications for preparative anion-exchange chromatography 409(1987)111

Selective non-adsorption preparative chromatography of bovine immunoglobulin G1 409(1987)305

Application of preparative reversed-phase HPLC to the isolation of insulin-like growth factor II from human serum 421(1987)61

Isolation and identification of the glucuronide conjugate of 2-hydroxydesipramine by preparative LC 427(1988)287

Some problems associated with the selection of separation conditions in preparative LC 436(1988)367

Influence of mobile phase pH on HPLC column loading capacity. Implications for the design of preparative protein separations 437(1988)429

Displacement chromatography of biomolecules 439(1988)341

Influence of the relative concentrations of two components of a binary mixture on their overloaded elution profiles 440(1988)9

Analytical and preparative HPLC of the papain-cleaved derivative of polymyxin B 441(1988)423

Preparative isolation of <sup>14</sup>C-labelled saccharose from a partly purified fraction of *Chlorella vulgaris* 442(1988)237

Preparative HPLC of macrotetrolides 442(1988)431

Preparative chromatography of proteins. Analysis of the multivalent ion-exchange formalism 443(1988)13

Tandem separation schemes for preparative HPLC of proteins 443(1988)31

Optimization of concentration overload in preparative LC 444(1988)275

Large-scale chromatography of recombinant proteins 444(1988)293

Reversed-phase preparative chromatography of [D-Ala<sup>1</sup>]-peptide T amide 444(1988)345

Split peaks in non-linear chromatography and their effect on sample throughput in large scale separations 449(1988)53

Cross-axis synchronous flow-through coil planet centrifuge for large-scale preparative counter-current chromatography. I. Apparatus and studies on stationary phase retention in short coils 449(1988)135

Cross-axis synchronous flow-through coil planet centrifuge for large-scale preparative counter-current chromatography. II. Studies on partition efficiency in short coils and preparative separations with multilayer coils 449(1988)153

Preparative chromatography of oligogalacturonic acids 449(1988)229

Isolation of biologically active plant constituents by LC 450(1988)3

Analysis of the information in a preparative chromatogram for further optimization of the operating conditions 450(1988)13

Column loading and relative retention in overloaded elution chromatography 450(1988)27

Column hardware in preparative LC with axial flow 450(1988)47

Preparative HPLC of peptides on a new reversed-phase packing material, Kromasil C<sub>18</sub> 450(1988)71

Economic laboratory practice in preparative column HPLC 450(1988)81

Separation of natural products by centrifugal partition chromatography 450(1988)91

Large-scale purification of the mycotoxins aflatoxin B<sub>1</sub>, B<sub>2</sub> and G<sub>1</sub> 450(1988)101

Preparative LC separation of isomers of prostacyclin carba-analogues and their intermediates 450(1988)105

Isolation of tocopherol homologues by preparative HPLC 450(1988)111

Purification of specific heterologous F(ab)<sub>2</sub>

fragments with DEAE-Zeta-Prep cartridges for ion-exchange chromatography 450(1988)133

Preparative chiral separation in an aqueous two-phase system by a few counter-current extractions 450(1988)277

Rapid preparative-scale purification of peptides derived from caseins using mass ion-exchange chromatography 450(1988)420

### Preservatives

Determination of preservatives in cosmetic products. I. TLC procedure for the identification of preservatives in cosmetic products 410(1987)395

### Press-fit connections

Fused-silica ends for glass capillaries: widening glass capillary butts by a heated tungsten point for preparing press-fit connections 435(1988)204

### Pressure

Effect of pressure on solute retention in the ionization control mode in reversed-phase LC 406(1987)247

### Pressure / density gradient

Single and multiple gradients in supercritical fluid chromatography (Review) 402(1987)1

### Pressure programming

Effect of the pore structure of silica gel on retention behaviour in pressure programming supercritical fluid chromatography 448(1988)87

### Primaquine

Detection and determination of antimalarial drugs and their metabolites in body fluids (Review) 434(1988)1

### Principal components analysis

Chemometric classification of reversed-phase HPLC columns 410(1987)31

### Priority pollutants

HPLC of eleven priority substituted phenols by isocratic elution 438(1988)429

### Pristinamycin

Determination of PI<sub>A</sub> and PII<sub>A</sub>, the two main components of pristinamycin, by HPLC in plasma 425(1988)197

### Probe-process liquid chromatograph

New concept in chromatographic instrumentation. The probe-process liquid chromatograph 441(1988)1



**Probenecid**

Retention properties of internal-surface reversed-phase silica packing and recovery of drugs from human plasma 420(1987)297

**Procainamide**

Simultaneous determination of procainamide and N-acetylprocainamide in serum by GC with nitrogen-selective detection 415(1987)143

HPLC drug analysis by direct injection of whole blood samples. II. Determination of hydrophilic drugs 423(1987)155

**Procaine**

Isolation of basic drugs from plasma using solid-phase extraction with a cyanopropyl-bonded phase 432(1988)209

**Procaterol**

Sensitive HPLC method for procaterol in human urine 417(1987)223

**Process capillary gas chromatography**

Flow programming system for process capillary GC 405(1987)163

**Process liquid chromatograph**

New concept in chromatographic instrumentation. The probe-process liquid chromatograph 441(1988)1

**Progesterone**

Automated direct assay system for the measurement of sex steroid hormones in serum using HPLC 426(1988)33

**Progesterone-11 $\alpha$ -succinyltyrosine methyl ester**

Adsorption chromatographic behaviour of <sup>125</sup>I-labelled progesterone-11 $\alpha$ - and -12 $\alpha$ -succinyltyrosine methyl ester 404(1987)258

Adsorption chromatographic behaviour of <sup>125</sup>I-labelled progesterone-11 $\alpha$ - and -12 $\alpha$ -succinyltyrosine methyl ester 404(1987)258

**Progestin receptors**

High-resolution separation of molybdate-stabilized progestin receptors using HPLC 425(1988)277

Influence of column length and pore size on high-performance ion-exchange chromatography of estrogen and progestin receptors 425(1988)293

**Progestins**

Simultaneous quantification of natural

glycocorticoids and progestins in serum 421(1987)123

**Proglumide**

Application of an ovomucoid-conjugated column for the optical resolution of some pharmaceutically important compounds 408(1987)316

**Proguanil**

Determination of proguanil and its metabolites cycloguanil and 4-chlorophenylbiguanide in plasma, whole blood and urine by HPLC 416(1987)394

Detection and determination of antimalarial drugs and their metabolites in body fluids (Review) 434(1988)1

**Proline**

HPLC separation of peptides possessing a proline residue in the amino-terminal penultimate position, and their products generated by enzymatic hydrolysis 424(1988)129

**Promethazine**

HPLC resolution of the enantiomers of thioridazine, its metabolites and related compounds 426(1988)223

Indirect HPLC resolution of racemic tertiary amines as their diastereomeric urea derivatives after N-dealkylation 436(1988)269

**Promoter-specific transcription factor Sp1**

Non-histone chromatin proteins that recognize specific sequences of DNA (Review) 418(1987)73

**Pro-ocytocin / neurophysin converting enzyme**

Synthetic peptide substrates as models to study a pro-ocytocin/neurophysin converting enzyme 440(1988)439

**Propafenone**

Determination of propafenone in biological fluids by fused-silica capillary GC using electron-capture detection 417(1987)295

**1,2-Propanediol**

GC method for the estimation of acetone and its metabolites in biological samples 415(1987)372

**2-Propanol**

Retention in capillary supercritical fluid chromatography with carbon dioxide mobile phases containing polar modifiers 406(1987)223

Rapid HPLC method for determination of ethanol and fusel oil in the alcoholic beverage industry 447(1988)443

#### ***n*-Propanol**

The rôle of the stationary phase in micellar liquid chromatography. Adsorption and efficiency 449(1988)349

#### **Propanols**

Determination of alcohols in alcoholic beverages by micro HPLC with indirect photometric detection 445(1988)139

#### **Propionic acid**

Studies on sample preconcentration in ion chromatography. VIII. Preconcentration of carboxylic acids prior to ion-exclusion separation 447(1988)155

#### **Propionylcarnitine**

Analytical method for urinary glutarylcarnitine, acetylcarnitine and propionylcarnitine with a carboxylic acid analyser and a reversed-phase column 423(1987)297

#### **Propiopromazine**

TLC screening method for the tranquilizers azaperone, propiopromazine and carazolol in pig tissues 436(1988)73

#### **Propiverine**

GC-mass fragmentographic determination of propiverine and its metabolites in plasma and urine 420(1987)43

#### **Propofol**

HPLC determination and preliminary pharmacokinetics of propofol and its metabolites in human plasma and urine 417(1987)458

Improved method for the determination of propofol in blood by HPLC with fluorescence detection 421(1987)171

Direct plasma injection using internal surface reversed-phase HPLC: feasibility study using propofol as a model compound 434(1988)271

#### **Propoxur**

Pre-column clean-up and LC determination of residues of *N*-methylcarbamate pesticides in extracts of total diet 404(1987)352

#### **Propranolol**

Influence of enantiomeric purity of a chiral selector on stereoselectivity 407(1987)217

Determination of association constants of  $\beta$ -

cyclodextrin and  $\beta$ -cyclodextrin-bearing polymers with drugs by HPLC 409(1987)61

Versatile isocratic HPLC assay for propranolol and its basic, neutral and acidic metabolites in biological fluids 414(1987)365

Rapid determination of the enantiomers of metoprolol, oxprenolol and propranolol in urine 416(1987)331

Improved HPLC method for the simultaneous determination of propranolol and 4-hydroxypropranolol in plasma with fluorescence detection 422(1987)294

Separation of enantiomers of derivatised or underderivatised propranolol by means of HPLC 424(1988)189

HPLC determination of propranolol and its metabolites in rat serum 431(1988)465

Isolation of basic drugs from plasma using solid-phase extraction with a cyanopropyl-bonded phase 432(1988)209

#### **Propranolol glycol**

HPLC determination of propranolol and its metabolites in rat serum 431(1988)465

#### ***n*-Propylbenzene**

Heavy alkylated benzene stationary phase for the GLC separation of ethylbenzene, xylenes, styrene, iso- and *n*-propylbenzene 409(1987)371

#### **Propyl effect**

Propyl effect and retention-structure correlation as a means of GC identification 406(1987)43

#### ***N-n*-Propylnorapomorphine**

HPLC separation and electrochemical or spectrophotometric determination of *R*(-)*N-n*-propylnorapomorphine and *R*(-)*10,11*-methylenedioxy-*N-n*-propylnorapomorphine in primate plasma 426(1988)283

#### **Propylparaben**

HPLC assay of imazodan, methylparaben and propylparaben in imazodan injection 450(1988)407

#### **6-Propyl-2-thiouracil**

Identification of thyrostatic drug residues in animal thyroids by HPTLC and fluorescence reaction detection 442(1988)459

#### **Propylurofuran acid**

Excretion of urinary acids during inverted sleep-waking rhythm 422(1987)1

**Prorenin**

Semi-preparative purification of recombinant human renin and prorenin 444(1988)203

**Prostacyclin**

Preparative LC separation of isomers of prostacyclin carba-analogues and their intermediates 450(1988)105

**Prostaglandin B<sub>2</sub>**

Automated HPLC extraction and quantification procedure for lipoxygenase metabolites 430(1988)1

**Prostaglandin E<sub>2</sub>**

Separation of <sup>125</sup>I-labelled prostaglandin E<sub>2</sub>-tyrosine methyl ester by reversed-phase HPLC 438(1988)111

**Prostaglandins**

Quantitative determination of some prostaglandin derivatives of the sub-groups E<sub>2</sub>, A<sub>2</sub> and B<sub>2</sub> by HPLC 404(1987)385

HPLC method for determining the enantiomeric purity of a benzidine prostaglandin by a diastereomeric separation 408(1987)275

Purification and quantitative analysis of urinary prostanoids in human and in rat by packed and capillary GC-negative-ion chemical-ionization MS 417(1987)1

Separation of prostaglandins and thromboxane by two-dimensional TLC 417(1987)414

Determination of prostaglandin E<sub>2</sub> in plasma of schizophrenics and healthy controls by GC-negative-ion chemical ionization MS and radioimmunoassay 420(1987)373

Chromatography of B prostaglandins on  $\beta$ -cyclodextrin silica: application to analysis of major E prostaglandins in human seminal fluid 421(1987)117

Dimethylisopropylsilyl ether derivatives in GC-MS of some prostanoids. Application to the microdetermination of 15-ketoprostaglandin F<sub>2 $\alpha$</sub>  formed by enzymatic conversion of PGF<sub>2 $\alpha$</sub>  in vitro 424(1988)1

Quantitative analysis of 6-keto-prostaglandin F<sub>1 $\alpha$</sub>  using immunoaffinity purification and GC-MS 425(1988)1

Determination of prostaglandins D<sub>2</sub>, E<sub>2</sub>, F<sub>2 $\alpha$</sub>  and 6-ketoprostaglandin F<sub>1 $\alpha$</sub>  as well as thromboxane B<sub>2</sub> in rat brain by GC-MS 426(1988)345

Arachidonic acid metabolites from polymorphonuclear leukocytes of healthy donors, severely burned patients and children with cystic fibrosis - routine monitoring by HPLC 427(1988)199

Determination of prostaglandins and thromboxane in whole blood by HPLC with fluorimetric detection 427(1988)209

Derivatization of prostaglandins and related compounds to (methoxime) alkyl ester alkyl ether derivatives for GC analysis 436(1988)219

Picomole determination of 2,4-dimethoxyanilides of prostaglandins by HPLC with electrochemical detection 442(1988)444

Solvent selectivity in the resolution of some regioisomeric and diastereomeric prostaglandin intermediates on silica 449(1988)77

Separation of the stereoisomers of an allenic E-type prostaglandin 450(1988)233

**Prostanoids**

Purification and quantitative analysis of urinary prostanoids in human and in rat by packed and capillary GC-negative-ion chemical-ionization MS 417(1987)1

**Protected peptides**

Reversed-phase HPLC of protected peptide segments 409(1987)281

**Proteinase inhibitors**

Chromatographic separation and partial characterization of microbial low-molecular-weight proteinase inhibitors 446(1988)203

**Proteinases**

Chromatographic isolation of primary and secondary metabolites from plant cell cultures 440(1988)53

**Protein HC**

HPLC and photodiode-array detection of the human protein HC (human complex-forming glycoprotein heterogeneous in charge), a chromophore-associated protein 444(1988)165

**Protein hydrolysis**

Simplified protein hydrolysis with methanesulphonic acid at elevated temperature for the complete amino acid analysis of proteins 448(1988)404

**Proteins**

see also Enzymes

- Application of HPLC to the assessment of subunit heterogeneity in plant 11S storage globulins 402(1987)159
- Improved separation of perch liver metallothionein by fast protein LC 402(1987)293
- Melittin-silica, a high-pressure affinity chromatography resin for calmodulin 403(1987)99
- Application of high-performance liquid chromatofocusing to the study of prostatic androgen receptors. Effect of stabilizing agents on the heterogeneity of receptor structure 403(1987)171
- Improved Stokes radius measurement of the glucocorticoid receptor using TSK G4000SW and TSK G3000SW high-performance size-exclusion columns. Analytical and preparative applications 403(1987)183
- Interaction of proteins with immobilized  $\text{Cu}^{2+}$ . Quantitation of adsorption capacity, adsorption isotherms and equilibrium constants by frontal analysis 403(1987)197
- Rapid ion-exchange chromatography for preparative separation of proteins. III. Application to purification of bovine factor X and prothrombin 403(1987)383
- New high-performance gel filtration columns for protein separation 404(1987)333
- Reversed-phase HPLC of soybean trypsin inhibitor with optical activity and UV absorbance detection 405(1987)295
- Fractionation of plant and animal high mobility group chromosomal proteins by ion-exchange and reversed-phase HPLC 405(1987)327
- Chromatographic and electrophoretic investigations of the properties of unprotected low-sulphur wool keratins 405(1987)365
- Adsorption isotherms on HPLC sorbents. II. Proteins on cation exchangers with silica support 406(1987)285
- Contamination of aqueous eluents due to corrosion of stainless-steel chromatographic components 407(1987)133
- Resolution of transducin subunits by chromatography on Blue Sepharose 407(1987)281
- Coordinative interactions in the separation of insulin and derivatives by HPLC 408(1987)43
- Effects of pretreatment on the enantioselectivity of silica-bound proteins used as HPLC stationary phases 408(1987)99
- Recovery of polypeptides after reversed-phase HPLC 408(1987)191
- Large scale chromatographic purification of ribosome-inactivating proteins 408(1987)235
- Rapid preparation of bovine mercaptalbumin by means of covalent chromatography on silica-based materials 408(1987)263
- On the increased retention volume of human haemoglobin in high-performance gel filtration 408(1987)303
- Characterization of bee venom and its main components by HPLC 408(1987)303
- Resolution of proteins in linear gradient elution ion-exchange and hydrophobic interaction chromatography 409(1987)101
- HPLC of phenylthiocarbonyl-amino acids. Application to carboxyl-terminal sequencing of proteins 409(1987)299
- Selective non-adsorption preparative chromatography of bovine immunoglobulin G1 409(1987)305
- Effects of large sample loads on column lifetime in preparative-scale LC 410(1987)43
- Rapid purification of the isoforms of rat liver Cd,Zn-metallothionein by HPLC 410(1987)470
- New metal chelate adsorbent selective for proteins and peptides containing neighbouring histidine residues 411(1987)177
- Multiple detection system for the high-resolution microcolumn SEC separation of proteins with a slurry-packed capillary column 411(1987)221
- Preparative separation of an enzymatic hydrolysate of  $^{14}\text{C}$ -labelled proteins 411(1987)335
- Assay, purification and characterization of a recombinant malaria circumsporozoite fusion protein by HPLC 411(1987)345
- Facile preparation of a non-porous strong anion-exchange column for proteins 411(1987)452
- Electrophoretic titration curve in 6 M urea of the bovine eye lens protein  $\alpha$ -crystallin 411(1987)507
- Isolation and partial characterization of pyridoxal 5'-phosphate hemoglobins by HPLC

- as a quality-control method for hemoglobin-based blood substitutes 413(1987)53
- Automated separation and measurement of urinary isoenzymes and protein by ion-exchange LC 414(1987)47
- ATP-hemoglobin: anomalous oxygen binding properties 415(1987)136
- Cyclic biospecific affinity chromatographic method for the purification of the sex steroid binding protein (SBP): application to the purification of SBP from toad 415(1987)297
- Determination of myoglobin in human serum by HPLC with chemiluminescence detection 416(1987)45
- Anion-exchange chromatography of normal and monoclonal serum immunoglobulins 416(1987)53
- Computer analysis of proteins separated by polyacrylamide gradient pore gel electrophoresis 416(1987)265
- Haemoglobin pyridoxylation: optimization and observation of an additional new pyridoxal phosphate-haemoglobin species by HPLC resolution 417(1987)397
- Fluorimetric determination of rat growth hormone in pituitary cell culture by HPLC with post-column derivatization 417(1987)409
- General strategies in the separation of proteins by HPLC methods (Review) 418(1987)3
- Non-histone chromatin proteins that recognize specific sequences of DNA (Review) 418(1987)73
- Chromatography of complex protein mixtures (Review) 418(1987)115
- Two-dimensional gel electrophoresis of proteins (Review) 418(1987)145
- Column LC of integral membrane proteins (Review) 418(1987)223
- Separation of protein hormones (Review) 418(1987)245
- Separation of asymmetrical hybrid hemoglobins by hydrophobic interaction chromatography 419(1987)303
- Detection of the changes in serum proteins during chemical hepatocarcinogenesis by two-dimensional electrophoresis under non-denaturing conditions 419(1987)315
- Isolation and characterization of a protein detected in inflamed human gastric mucosa 419(1987)323
- Pulse injection analysis of the binding of serum proteins to porous polymer gels modified with formyl groups 420(1987)13
- Cysteine adducts of human haemoglobin measured by isoelectric focusing in polyacrylamide gels with a non-linear pH gradient 420(1987)35
- Method for purification of an insulin-like growth factor-binding protein produced by human HEP G2 hepatoma cells 420(1987)163
- Purification of commercial human albumin on immobilized IDA-Ni<sup>2+</sup> 421(1987)141
- Size-exclusion HPLC of immune complexes isolated from patients with acute leukaemia. Preliminary observations 421(1987)434
- Characterization of adsorption on the stationary phase using high-performance immunoaffinity chromatography 423(1987)41
- Staining of serum concanavalin A-binding proteins after micro two-dimensional electrophoresis and blotting 423(1987)115
- Removal of sodium dodecyl sulphate from proteins by GPC 423(1987)376
- Synthesis of high-capacity immunoaffinity sorbents with oriented immobilized immunoglobulins or their F<sub>ab</sub>' fragments for isolation of proteins 424(1988)243
- High-resolution separation of molybdate-stabilized progesterin receptors using HPLC 425(1988)277
- Influence of column length and pore size on high-performance ion-exchange chromatography of estrogen and progesterin receptors 425(1988)293
- Separation and quantification of histone H1 subtypes and high-mobility-group proteins by reversed-phase LC: protein levels in rat tissues during postnatal development 426(1988)65
- Column LC separation of collagen  $\alpha$ -chain polymers on sepiolite 426(1988)162
- Chromatographic analysis of the trinitrophenyl derivatives of insulin 427(1988)239
- Separation of human haemoglobin alkylated at  $\beta$ 93 cysteine from its native form by fast protein LC 427(1988)335
- Chromatographic detergent exchange in the preparation of membrane receptor protein 427(1988)341
- Steric protection of a modified glass surface with

- adsorbed polyacrylamide: interactions with blood proteins 428(1988)17
- Comparison of serum protein removal in haemodialysis therapy by partly protein-permeable haemodialysers 428(1988)25
- Effects of differences in charge and hydrophobicity of surface amino acids of hemoglobins on high-performance GPC 428(1988)247
- Urinary protein analysis (Review) 429(1988)315
- Cerebrospinal fluid protein analysis in diseases of the nervous system (Review) 429(1988)345
- High-performance affinity chromatography of human serum concanavalin A binding proteins 431(1988)45
- Purification of monoclonal antibodies against the low-density lipoprotein receptor by preparative isotachopheresis 431(1988)327
- Use of the Pharmacia Phast System for sodium dodecyl sulphate polyacrylamide gel electrophoresis of non-globular proteins: application to collagens 431(1988)474
- Affinity purification of human plasma fibronectin on immobilized gelatin 432(1988)93
- Separation of transcobalamin II isoproteins by means of chromatofocusing 432(1988)103
- Free-flow electrophoretic apparatus for separation and concentration of proteins 432(1988)137
- Quantitative evaluation of collagen fractions separated by acrylamide gel electrophoresis: its application for collagen typing in normal and pathological tissues 433(1988)81
- HPLC separation of human haemoglobins. Simultaneous quantitation of foetal and glycated haemoglobins 434(1988)95
- Characterization of buffalo lactotransferrin by polyacrylamide gel electrophoresis in 6 M urea 434(1988)228
- Separation of type IX collagen from other cartilage collagens by hydrophobic interaction chromatography 434(1988)423
- HPLC separation and immunological characterization of soluble bovine viral diarrhoea virus antigen 435(1988)149
- HPLC of amino acids, peptides and proteins. LXXXIV. Application of derivative spectroscopy to the study of column residency effects in the reversed-phase and size-exclusion LC separation of proteins 435(1988)271
- High-performance SEC of recombinant derived proteins and aggregated species 436(1988)289
- Ionic binding properties of carrier ampholytes 437(1988)147
- Investigation of the tetramer-dimer equilibrium in haemoglobin solutions by high-performance SEC on a diol column 437(1988)193
- Single-step purification of rat C-reactive protein and generation of monospecific C-reactive protein antibody 437(1988)399
- Influence of mobile phase pH on HPLC column loading capacity. Implications for the design of preparative protein separations 437(1988)429
- Microcolumn chromatofocusing. III. Effect of the analysis time on the resolution and the use of a semi-disposable microcolumn 438(1988)15
- Displacement chromatography of biomolecules 439(1988)341
- Chromatographic purification and characterization of elicitors of necrosis on tobacco produced by incompatible *Phytophthora* species 440(1988)87
- HPLC in protein sequence determinations 440(1988)231
- Ion-exchange derivatives of Spheron. VI. Strongly basic derivatives 440(1988)287
- High-performance adsorption chromatography of proteins on 2- $\mu$ m spherical beads of hydroxyapatite 440(1988)305
- Secretion of proteins by *Bacillus subtilis* 168 grown in the presence of membrane active agents (alcohols) 440(1988)379
- Plasmodium falciparum* exoprotein stimulation of human T-lymphocytes unsensitized to malaria 440(1988)385
- Survey of the folding pathway of a two-domain protein phosphoglycerate kinase 440(1988)421
- Use of a small TSK GSW HPLC column for large-zone chromatographic studies of monomer-oligomer equilibria of membrane protein 442(1988)229
- Use of HPLC for the purification of antibodies and antibody conjugates and the study of antibody-antigen interactions 442(1988)279
- Isolation and quantitation of cadmium-, zinc-

- and copper-metallothioneins by HPLC atomic absorption spectrometry 442(1988)345
- High-performance immobilized-metal-ion affinity chromatography of peptides and proteins 443(1988)3
- Preparative chromatography of proteins. Analysis of the multivalent ion-exchange formalism 443(1988)13
- Tandem separation schemes for preparative HPLC of proteins 443(1988)31
- Comparative study of Zorbax Bio Series GF 250 and GF 450 and TSK-GEL 3000 SW and SWXL columns in SEC of proteins 443(1988)45
- Dextran-coated silica packings for high-performance SEC of proteins 443(1988)55
- Synthesis of a non-porous, polystyrene-based strong anion-exchange packing material and its application to fast HPLC of proteins 443(1988)73
- HPLC of amino acids, peptides and proteins. LXXXVI. The influence of different displacer salts on the retention and bandwidth properties of proteins separated by isocratic anion-exchange chromatography 443(1988)97
- Molecular orientation of immunoglobulin G at high concentration on an ion-exchange sorbent 443(1988)119
- Ion-exchange chromatography of proteins on a polyethyleneimine-grafted hydrophilic polymer for HPLC 443(1988)133
- Chromatographic evaluation of large-pore and non-porous polymeric reversed phases 443(1988)143
- Chromatography of proteins on hydrophobic interaction and ion-exchange chromatographic matrices: mobile phase contributions to selectivity 443(1988)173
- Reduction studies on bacterial recombinant somatomedin C/insulin-like growth factor-1 443(1988)183
- Reversible subunit dissociation of tumor necrosis factor during hydrophobic interaction chromatography 443(1988)205
- Application of HPLC in neurophysin disulfide assignment 443(1988)285
- Separation of proteins by reversed-phase HPLC. I. Optimizing the column 443(1988)363
- Separation of proteins by reversed-phase HPLC. II. Optimizing sample pretreatment and mobile phase conditions 443(1988)381
- Isolation of specific lymphocyte receptors by high-performance immunoaffinity chromatography 444(1988)13
- Determination of the molecular size distribution of immunoglobulin G (IgG) in intravenous IgG-albumin formulations by HPLC 444(1988)141
- Optimal resolution of eye lens  $\gamma$ -crystallins by cation-exchange HPLC on Synchropak CM300 444(1988)239
- Scale-up from analytical to preparative reversed-phase HPLC of soybean glycinin 444(1988)263
- Large-scale chromatography of recombinant proteins 444(1988)293
- Preparative separation of peptide and protein samples by HPLC with gradient elution. I. The Craig model as a basis for computer simulations 444(1988)303
- Preparative separation of peptide and protein samples by HPLC with gradient elution. II. Experimental examples compared with theory 444(1988)325
- Effect of steric exclusion on the separation of proteins by hydrophilic SEC 445(1988)29
- Rapid analysis of proteins and peptides by reversed-phase chromatography with polymeric micropellicular sorbents 445(1988)71
- High-performance affinity chromatography of proteins on TSKgel Heparin-5PW 445(1988)234
- Cationic capillary isotachopheresis of proteins 445(1988)417
- Post-capillary fluorescence detection in capillary zone electrophoresis using *o*-phthalaldehyde 447(1988)117
- Separation of proteins by microcolumn LC based on the reversed-phase and size-exclusion principles 448(1988)73
- Chick 28 000  $M_r$  vitamin D-dependent calcium-binding protein in intestine, kidney and cerebellum. Purification using chromatofocusing 448(1988)145
- Isolation of recombinant mycobacterial antigens by an automatic and generally applicable purification method for  $\beta$ -galactosidase fusion proteins 448(1988)165
- Separation of tubulin subunits by reversed-phase HPLC 448(1988)371
- Purification of specific heterologous F(ab)<sub>2</sub>

fragments with DEAE-Zeta-Prep cartridges for ion-exchange chromatography 450(1988)133

Separation of Friend erythroleukaemic cell histones and high-mobility-group proteins by reversed-phase HPLC 450(1988)309

#### **Protein secretion**

Secretion of proteins by *Bacillus subtilis* 168 grown in the presence of membrane active agents (alcohols) 440(1988)379

#### **Protein separation strategies**

General strategies in the separation of proteins by HPLC methods (Review) 418(1987)3

#### **Proteoglycans**

Extracellular mammalian polysaccharides: glycosaminoglycans and proteoglycans (Review) 418(1987)187

#### **Proteolytic digests**

Microbore HPLC-MS for the analysis of proteolytic digests by continuous-flow fast-atom bombardment MS 443(1988)355

#### ***Proteus mirabilis***

Studies of metabolites in diarrheal stool specimens positive for *Klebsiella*, *Serratia*, and *Proteus* spp. by frequency-pulsed electron-capture GC 430(1988)209

#### **Prothrombin**

Rapid ion-exchange chromatography for preparative separation of proteins. III. Application to purification of bovine factor X and prothrombin 403(1987)383

#### **Protopine**

Determination of alkaloids of *Fumaria parviflora* and *Fumaria capreolata* by HPLC and capillary isotachopheresis 445(1988)258

#### **Protoporphyrins**

Normal-phase HPLC of free acid dicarboxylic porphyrins and hematoporphyrin derivative on silica 422(1987)73

Determination of erythrocyte porphyrins by reversed-phase HPLC using capsule-type silica gels coated with silicone polymer 433(1988)73

Stability-indicating HPLC analysis of tin protoporphyrin and other free acid metalloporphyrins 445(1988)429

#### **PRP-1**

Characterization of some styrene-divinylbenzene

sorbents for reversed-phase chromatography 442(1988)97

Anion chromatography using a coated PRP-1 column and eluents of pH > 7 447(1988)315

#### **PRP-X300**

New ion-exclusion phase for organic acids 439(1988)71

#### **PR-toxin**

Standardized HPLC of 182 mycotoxins and other fungal metabolites based on alkylphenone retention indices and UV-VIS spectra (diode array detection) 404(1987)195

#### **Pseudodipeptides**

Determination of the chiral purity of dipeptide isosteres containing a reduced peptide bond by GC analysis 442(1988)165

#### **Pseudoephedrine**

Simultaneous analysis of acetaminophen, pseudoephedrine hydrochloride and chlorpheniramine maleate in a cold tablet using an isocratic, mixed micellar HPLC mobile phase 410(1987)206

Sensitive quantification of pseudoephedrine in human plasma and urine by HPLC 424(1988)73

Sensitive HPLC determination of pseudoephedrine in plasma and urine 426(1988)406

Use of cyclodextrins in isotachopheresis. IV. The influence of cyclodextrins on the chiral resolution of ephedrine alkaloid enantiomers 438(1988)211

#### **Pseudouridine**

Determination of the age dependency of the creatinine-related pseudouridine excretion in children's urine by HPLC 434(1988)202

#### **Psilocin**

Determination of tryptamine derivatives in hallucinogenic mushrooms using HPLC with photodiode array detection 408(1987)402

#### **Psilocybin**

Determination of tryptamine derivatives in hallucinogenic mushrooms using HPLC with photodiode array detection 408(1987)402

#### **Psoralen**

Determination of psoralen in serum by reversed-phase HPLC 423(1987)383



***Psorospermum febrifugum***

Isolation of biologically active plant constituents by LC 450(1988)3

**Pterins**

Analysis and clinical significance of pterins (Review) 429(1988)95

Estimation of tetrahydrobiopterin and other pterins in plasma by isocratic LC with electrochemical and fluorimetric detection 432(1988)321

**Pterobilin**

Interactions between ascorbic acid and linear and cyclized bile pigments. A TLC study 440(1988)275

**Pteroylpolyglutamate hydrolase**

HPLC assay for pteroylpolyglutamate hydrolase 417(1987)47

**Puerarin**

Preparation and evaluation of mono-, di- and triphenyl-treated plates for the HPTLC analysis of puerarin and paeonol 407(1987)314

**Pulp**

SEC of deposits in pulp and paper mills 411(1987)363

**Pulsed amperometric detection**

LC determination of carbohydrates with pulsed amperometric detection and a membrane reactor 447(1988)268

**Pulsed electron-capture detector**

Influence of the carrier gas flow-rate on the pulsed electron-capture detector 446(1988)329

**Purge-and-trap methods**

Purge-and-trap GC method for residual volatiles in high-temperature polymers 450(1988)394

**Purine nucleoside phosphorylase**

Determination of adenosine deaminase and purine nucleoside phosphorylase activities using HPLC and radiochromatographic methods 434(1988)439

**Purines**

see also Nucleobases

Demonstration of the adenosine reservoirs with nitrobenzylthioinosine in liver and diaphragm by HPLC 426(1988)249

Optimization of the ion-pair HPLC separation of

purine derivatives in erythrocytes, thymocytes and liver mitochondria 434(1988)447

**Putrescine**

Improved method for benzoyl chloride derivatization of polyamines for HPLC 408(1987)227

Determination of polyamines in human erythrocytes by capillary GC with nitrogen-phosphorus detection 415(1987)27

Simultaneous determination of  $\gamma$ -aminobutyric acid and polyamines by HPLC 419(1987)285

Rapid and sensitive assay of ornithine decarboxylase activity by HPLC of the *o*-phthalaldehyde derivative of putrescine 419(1987)291

Reversed-phase LC separation and simultaneous fluorimetric detection of polyamines and their monoacetyl derivatives in human and animal urine, serum and tissue samples: an improved, rapid and sensitive method for routine application 430(1988)249

Quantitation of polyamines using TLC and image analysis 441(1988)361

Detection by HPLC of polyamines formed by clostridal putrefaction of caseins 442(1988)267

HPLC method for the determination of  $\epsilon$ -( $\gamma$ -glutamyl)lysine and mono- and bis- $\gamma$ -glutamyl derivatives of putrescine and spermidine 443(1988)329

**Pyoverdins**

Assignment of the configurations of the amino acids in peptidic siderophores 438(1988)291

**Pyrazinamide**

Rapid and simultaneous determination of pyrazinamide and its major metabolites in human plasma by HPLC 413(1987)342

Identification of 5-hydroxypyrazinamide isolated from urine of subjects given pyrazinamide 416(1987)144

LC assay for the simultaneous determination of pyrazinamide and rifampicin in serum samples from patients with tuberculous meningitis 420(1987)73

Microdetermination of pyrazinamide and its metabolites (2-pyrazinoic acid, 5-hydroxypyrazinoic acid, 5-hydroxypyrazinamide and pyrazinuric acid) in plasma and urine by LC (in French) 422(1987)217

**Pyrazines**

Correlation between molecular structures and retention indices of pyrazines 402(1987)309

**2-Pyrazinoic acid**

Microdetermination of pyrazinamide and its metabolites (2-pyrazinoic acid, 5-hydroxypyrazinoic acid, 5-hydroxypyrazinamide and pyrazinuric acid) in plasma and urine by LC (in French) 422(1987)217

**Pyrazinuric acid**

Microdetermination of pyrazinamide and its metabolites (2-pyrazinoic acid, 5-hydroxypyrazinoic acid, 5-hydroxypyrazinamide and pyrazinuric acid) in plasma and urine by LC (in French) 422(1987)217

**Pyrazolones**

Investigation of an alkylamine modified and pH-controlled mobile phase for separation of pyrazolone derivatives by HPLC 409(1987)81

**Pyrazoloquinolines**

Lipophilicity measurement of benzodiazepine-receptor ligands by reversed-phase LC. Comparison between HPLC and TLC 447(1988)404

**Pyrene**

Resolution of overlapped chromatograms by the use of a Kalman filter. III. Effects of overlapping and outstripping of a solvent peak on the analysis 411(1987)95

**Pyrethrins**

Analysis of pyrethrins in pyrethrum extracts by HPLC 447(1988)239

**Pyridines**

Group contributions to hydrophobicity and elution behaviour of pyridine derivatives in reversed-phase HPLC 449(1988)95

**Pyridocarbazole alkaloids**

Fluorodensitometric assay of 6*H*-pyrido[4,3-*b*]-5,11-dimethylcarbazoles (ellipticine and derivatives) biosynthesized by *Ochrosia elliptica* cultures *in vitro* 409(1987)426

**Pyridone carboxylic acids**

Separation of pyridone carboxylic acid enantiomers by HPLC using copper(II)-L-amino acid as the eluent 448(1988)439

**Pyridostigmine**

HPLC determination of pyridostigmine in plasma 422(1987)353

**Pyridoxal**

Analysis of vitamin B<sub>6</sub> vitamers in plasma by cation-exchange HPLC 416(1987)138

Simple isocratic HPLC method for the separation of the six vitamers of vitamin B<sub>6</sub> 424(1988)315

**Pyridoxal kinase**

HPLC assay of erythrocyte enzyme activity levels involved in vitamin B<sub>6</sub> metabolism 431(1988)406

**Pyridoxal 5-phosphate**

Sensitive assay for determination of pyridoxal-5-phosphate in enzymes using HPLC after derivatization with cyanide 434(1988)209

**Pyridoxal 5'-phosphate**

Analysis of vitamin B<sub>6</sub> vitamers in plasma by cation-exchange HPLC 416(1987)138

**Pyridoxal 5'-phosphate-haemoglobin**

Isolation and partial characterization of pyridoxal 5'-phosphate hemoglobins by HPLC as a quality-control method for hemoglobin-based blood substitutes 413(1987)53

Haemoglobin pyridoxylation: optimization and observation of an additional new pyridoxal phosphate-haemoglobin species by HPLC resolution 417(1987)397

Hemoglobin-based blood substitutes: characterization of five pyridoxal 5'-phosphate derivatives of hemoglobin 419(1987)37

**Pyridoxamine**

Analysis of vitamin B<sub>6</sub> vitamers in plasma by cation-exchange HPLC 416(1987)138

Simple isocratic HPLC method for the separation of the six vitamers of vitamin B<sub>6</sub> 424(1988)315

**Pyridoxamine 5'-phosphate**

Analysis of vitamin B<sub>6</sub> vitamers in plasma by cation-exchange HPLC 416(1987)138

**Pyridoxamine (pyridoxine-5'-phosphate) oxidase**

HPLC assay of erythrocyte enzyme activity levels involved in vitamin B<sub>6</sub> metabolism 431(1988)406

**Pyridoxic acid**

Analysis of vitamin B<sub>6</sub> vitamers in plasma by cation-exchange HPLC 416(1987)138

**4-Pyridoxic acid**

Plasma B<sub>6</sub> vitamer and 4-pyridoxic acid concentrations of men fed controlled diets 428(1988)35

**Pyridoxine**

Analysis of vitamin B<sub>6</sub> vitamers in plasma by cation-exchange HPLC 416(1987)138

Simple isocratic HPLC method for the separation of the six vitamers of vitamin B<sub>6</sub> 424(1988)315

**Pyrimethamine**

Detection and determination of antimalarial drugs and their metabolites in body fluids (Review) 434(1988)1

**Pyrimidines**

see also Nucleobases

Separation of modified nucleic acid constituents by micellar electrokinetic capillary chromatography 409(1987)193

Determination of 5-fluoro-2'-deoxyuridine in human plasma by HPLC with pre-column fluorimetric derivatization 416(1987)176

Isocratic HPLC method for studying the metabolism of blood plasma pyrimidine nucleosides and bases: concentration and radioactivity measurements 434(1988)111

Quantitation of triazole and pyrimidine plant growth retardants 438(1988)393

Reversed-phase HPLC of pyrimidine bases and nucleosides. Application of solvophobic theory 445(1988)59

**1-(2-Pyrimidinyl)piperazine**

Simultaneous quantitation of buspirone and 1-(2-pyrimidinyl)piperazine in human plasma and urine by capillary GC-MS 428(1988)265

Simultaneous determination of buspirone, gepirone, ipsapirone and their common metabolite 1-(2-pyrimidinyl)piperazine in rat plasma and brain by HPLC 431(1988)477

**Pyromellitate buffers**

Single-column anion chromatography with indirect UV detection using pyromellitate buffers as eluents 439(1988)137

**Pyrophosphoric acid**

Complex-forming equilibria in isotachopheresis. VII. Evaluation of the ion-pair formation constants of phosphorus oxo acids with histidine and assessment of their separability 410(1987)279

**Pyrrrolizidine alkaloids**

Ammonium positive-ion and hydroxide negative-ion chemical ionization GC-MS for the identification of pyrrrolizidine alkaloids in

*Eupatorium rotundifolium* L. var. *ovatum* 428(1988)352

**Pyruvate**

HPLC measurement of selected blood citric acid cycle intermediates 425(1988)47

Ion chromatographic determination of sugar phosphates in physiological samples 439(1988)83

**Pyruvic acid**

GC of branched-chain keto acids analysed as their silylated oxime derivatives 423(1987)270

**Q****Qualitative indices**

Isotachophoretic determination of mobility and  $pK_a$  by means of computer simulation. VI. Mobility and  $pK_a$  of sixty anions in methanol 408(1987)27

**Quaternary ammonium drugs**

Polar contributions of the stationary phase to the reversed-phase ion-pair HPLC separation of quaternary ammonium drugs 437(1988)83

**Quercetine heterosides**

Densitometric determination of quercetine heterosides in plant extracts (in French) 437(1988)260

**Quinacrine**

Determination of quinacrine (mepacrine) in plasma by HPLC with fluorimetric detection 420(1987)341

**Quinapril**

Determination of quinapril and its active metabolite in human plasma and urine by GC with electron-capture detection 421(1987)187

**Quinethazone**

LC screening of diuretics in urine 415(1987)347

**Quinidine**

HPLC analysis of quinine and its diastereoisomer quinidine 415(1987)177

**Quinine**

HPLC analysis of quinine and its diastereoisomer quinidine 415(1987)177

Improved HPLC method for the determination of quinine in plasma 430(1988)170

Detection and determination of antimalarial drugs and their metabolites in body fluids (Review) 434(1988)1

**Quinolinic acid**

High-voltage electrophoretic separation of tryptophan metabolites of the kynurenine pathway 420(1987)263

**Quinolones**

Application of HPLC of some antibiotics in clinical microbiology 427(1988)257

**Quinones**

Stationary phases. XV. HPLC behaviour of quinones on a safrole stationary phase 408(1987)323

**Quinoxaline-2-carboxylic acid**

1-(4-Hydroxyphenyl)-, 1-(2,4-dihydroxyphenyl)- and 1-(2,5-dihydroxyphenyl)-2-bromoethanones: new labels for determination of carboxylic acids by HPLC with electrochemical and UV detection 442(1988)209

**Quizalofop-ethyl**

GC-MS investigation of phenoxypropanoic acid derivatives possessing herbicidal activity 438(1988)55

**R****Racemic chiral stationary phases**

Applications for racemic versions of chiral stationary phases 407(1987)211

**Radioiodination**

Tyrosylation and purification of peptides for radioiodination 447(1988)341

**Radioiodine nuclides**

Automated separation process for radioanalytical purposes at nuclear power plants 406(1987)359

**Radiolabelled compounds**

Fast reversed-phase HPLC determination of  $^{14}\text{C}$ - or  $^3\text{H}$ -labelled S-adenosylmethionine in reaction mixtures 410(1987)500

**Ranitidine**

Determination of ranitidine in rat plasma and brain by HPLC 413(1987)363

HPLC determination of ranitidine in whole blood and plasma by using a short polymeric column 421(1987)418

Automated solid-phase extraction and HPLC determination of ranitidine from urine, plasma and peritoneal dialysate 422(1987)165

**Rare earths**

Separation of some rare earths and transplutonium elements by HPLC on Ostion cation exchanger 435(1988)249

**RDX**

LC analysis of cyclotrimethylenetrinitramine in biological fluids using solid-phase extraction 421(1987)430

**Reaction kinetics**

Contributions of GC to the development of heterogeneous catalysis 406(1987)151

**Reaction rate constants**

Reversed-flow GC applied to physico-chemical measurements 446(1988)39

**Reagent sandwiching**

Automated pre-column derivatization of amino acids with *o*-phthalaldehyde by a reagent sandwiching technique 408(1987)217

**Recombinant proteins**

Large-scale chromatography of recombinant proteins 444(1988)293

**Redox chemiluminescence detector**

Coupling LC with redox chemiluminescence detection 441(1988)125

**Reduced haloperidol**

Simultaneous determination of very low concentrations of haloperidol and reduced haloperidol in human serum by an LC method 433(1988)312

**Reductive alkylation**

Effects of reductive alkylation on peptide retention times 443(1988)247

**Relative permittivity**

Characterization of mobile phases for the investigation of electrokinetic phenomena in LC 447(1988)177

**Relative retention**

Column loading and relative retention in overloaded elution chromatography 450(1988)27

**Renin**

Purification of human renin by renin inhibitor (ES-305) affinity chromatography 421(1987)350

Determination of monosaccharides in glycoproteins by reversed-phase HPLC on 2.1-mm narrowbore columns 443(1988)299

Semi-preparative purification of recombinant human renin and prorenin 444(1988)203

**Rennets**

High-performance anion-exchange chromatography of rennet enzymes 438(1988)451

**Reordering/resolution**

Influence of bonding chemistry on the reordering/resolution of silica immobilized alkyl chains 449(1988)115

**Resin acids**

GC analysis of underivatized resin acids 448(1988)428

**Resins**

TLC analysis of some Indian natural resins occurring in art objects 439(1988)430

Quantitative analysis of resins used in fiber-reinforced composites by reversed-phase LC 447(1988)141

**Resoles**

Gel filtration chromatography of resole phenolic resins 445(1988)371

**Resolution**

Microcolumn chromatofocusing. III. Effect of the analysis time on the resolution and the use of a semi-disposable microcolumn 438(1988)15

Multi-variate ranking function for optimizing separations 439(1988)173

Causes for the loss of resolution in an ion chromatography resin 445(1988)211

**Response factor calibration**

Determinations of polychlorinated biphenyl isomers by response factor calibration 404(1987)155

**Retention behaviour**

Elution behaviour of peropyrene-type polycyclic aromatic hydrocarbons in LC 402(1987)173

Correlation between molar refraction and GC retention data 403(1987)11

Retention behaviour of aromatic aldehyde derivatives in reversed-phase LC 404(1987)242

Effect of pressure on solute retention in the ionization control mode in reversed-phase LC 406(1987)247

Super- and subcritical fluid chromatography-Fourier transform IR spectrometry of medium polar compounds on polymeric styrene-divinylbenzene packed columns 407(1987)87

Prediction of thermodynamic retention behaviour in reversed-phase LC with aqueous eluents and various concentrations of methanol by using the solubility concept model 408(1987)21

Retention behaviour of acidic, neutral and basic drugs on a CN column using phosphate buffers in the mobile phase 410(1987)77

LC behaviour of chlorinated carbanilides 410(1987)95

Elution behaviour of aliphatic carboxylic acids on a strong cation-exchange resin column 410(1987)103

Retention and selectivity in amino, cyano and diol normal bonded phase HPLC columns 410(1987)249

Retention behaviour of cyclodextrins and branched cyclodextrins on reversed-phase columns in HPLC 437(1988)47

HPLC of amino acids, peptides and proteins. LXXXVI. The influence of different displacer salts on the retention and bandwidth properties of proteins separated by isocratic anion-exchange chromatography 443(1988)97

**Retention index**

Correlation between molecular structures and retention indices of pyrazines 402(1987)309

Reduction of the number of characterizing probes in GC by factor analysis 402(1987)318

Some problems in the correlation of molecular parameters and the connectivity index 403(1987)19

Factor analysis of trends in selectivity in GLC of 175 saturated and unsaturated esters on 14 polysiloxane stationary phases 404(1987)11

Calculation of programmed temperature GC characteristics from isothermal data. I. Theory and computational procedures 405(1987)67

Propyl effect and retention-structure correlation as a means of GC identification 406(1987)43

Retention and thermodynamics of solution of *n*-alkenes in OV-101 406(1987)75

Hierarchical ascending classification of Kováts retention indices as a means of indicating siloxane stationary phase specificities towards aliphatic carboxylic esters 406(1987)95

Classification of GC stationary phases by a new variant of Kováts retention index 406(1987)105

Use of Kováts retention indices for characterizing solutes in complex samples, separated by linear temperature-programmed capillary GLC 406(1987)117

GC study of microporous carbons 406(1987)139

Effect of organic modifier on the separation and efficiencies of homologues on porous polymer columns 407(1987)19

Use of Kováts indices for prediction of retention temperatures in linear temperature-programmed capillary GC 407(1987)65

Reproducibility of temperature-programmed retention indices on several OV-101 columns 407(1987)79

#### **Retention index increments**

Effect of primary and secondary atomic environments on the interactions of some alkane C-H bonds on a squalane stationary phase at different column temperatures in GLC 406(1987)131

#### **Retention index scale**

Improved standardization in reversed-phase HPLC using 1-nitroalkanes as a retention index scale 435(1988)43

#### **Retention indices**

Mixed stationary phases in GLC. Partition coefficients and retention indices in OV-101-OV-25, OV-101-carbowax 20M and OV-225-SP-2340 mixtures 410(1987)13

Analysis and identification of some aminoether alcohols and their ethers 410(1987)447

Screening of volatile compounds present in human blood using retention indices in GC 411(1987)251

Prediction of retention indexes. I.

Structure-retention index relationship on apolar columns 436(1988)137

GC retention indices of sulfur vesicants and related compounds 436(1988)399

Effect of temperature on the GC separation of halogenated compounds on polar and non-polar stationary phases 437(1988)33

System of retention indices for a linear temperature programming regime 439(1988)185

Capillary GC of steroidal alkaloids from Solanaceae. Retention indices and simultaneous flame ionization/nitrogen-specific detection 442(1988)133

Use of retention temperatures for the identification of phenols separated on OV-1701 capillary columns using linear temperature-programmed GC 448(1988)203

#### **Retention mechanisms**

Solute-solvent interactions in tetra-*n*-butylphosphonium salts studied by GC 438(1988)1

Main parameters governing the retention of polynuclear aromatic hydrocarbons on tetrachlorophthalimidopropyl-bonded silica 439(1988)195

#### **Retention prediction**

Holistic molecular conformation and total surface area calculations as predictors for solution properties and chromatographic parameters 403(1987)1

Application of the Snyder displacement model to aromatic nitrogen compounds on an aminopropyl bonded-phase column 405(1987)107

Prediction of thermodynamic retention behaviour in reversed-phase LC with aqueous eluents and various concentrations of methanol by using the solubility concept model 408(1987)21

Automated identification of toxic substances in poisoned human fluids by a retention prediction system in reversed-phase LC 436(1988)11

Prediction of HPLC capacity factors following column calibration. Influence of different solvents and columns on the separation of a series of aromatic amides 437(1988)13

HPLC of amino acids, peptides and proteins. LXXXV. Evaluation of the use of hydrophobicity coefficients for the prediction of peptide elution profiles 438(1988)197

Prediction of peptide retention times 442(1988)69

**Retention relationships**

Retention relationships for aromatic hydrocarbons eluted from capped and uncapped octadecyl silica gels 447(1988)13

**Retention temperature**

Use of Kováts indices for prediction of retention temperatures in linear temperature-programmed capillary GC 407(1987)65

Reproducibility of temperature-programmed retention indices on several OV-101 columns 407(1987)79

Use of retention temperatures for the identification of phenols separated on OV-1701 capillary columns using linear temperature-programmed GC 448(1988)203

**Retention time**

Calculation of programmed temperature GC characteristics from isothermal data. I. Theory and computational procedures 405(1987)67

**Retention volume**

On the increased retention volume of human haemoglobin in high-performance gel filtration 408(1987)303

**Retinal**

Rapid, simultaneous determination of isomers of retinal, retinal oxime and retinol by HPLC 438(1988)383

**Retinal oxime**

Rapid, simultaneous determination of isomers of retinal, retinal oxime and retinol by HPLC 438(1988)383

**Retinoids**

Simultaneous determination of the aromatic retinoids etretin and etretinate and their main metabolites by reversed-phase liquid chromatography 415(1987)413

Determination of the aromatic retinoids (etretin and isoetretin) in biological fluids by HPLC 421(1987)177

Quantitative analysis of retinoids in biological fluids by HPLC using column switching. I.

Determination of isotretinoin and tretinoin and their 4-oxo metabolites in plasma 424(1988)303

Quantitative analysis of retinoids in biological fluids by HPLC using column switching. II.

Simultaneous determination of etretinate, acitretin and 13-*cis*-acitretin in plasma 431(1988)297

**Retinol**

Reversed-phase HPLC separation and electrochemical detection of retinol and its isomers 417(1987)65

Gas chromatography of retinol and  $\alpha$ -tocopherol without derivatization 434(1988)21

Rapid, simultaneous determination of isomers of retinal, retinal oxime and retinol by HPLC 438(1988)383

**Revenast hydrochloride**

Determination of revenast hydrochloride in human plasma by GC with nitrogen-selective detection 419(1987)351

**Reversed-phase chromatography**

Hydrophobic interaction chromatography of peptides as an alternative to reversed-phase chromatography 444(1988)269

**Rhamnose**

Requirements of mutually miscible liquids for ternary chromatographic solvents with reference to monosaccharide PC 436(1988)53

**Rhein**

Simple method for the determination of rhein in biological fluids by HPLC 428(1988)173

Screening procedure for stimulant laxatives in urine using HPLC with diode array detection 433(1988)131

**Rhodamine B**

Foam counter-current chromatography with the cross-axis synchronous flow-through coil planet centrifuge 403(1987)77

**Ribavirin**

Quantification of ribavirin in biological fluids and tissues by HPLC 414(1987)202

HPLC determination of ribavirin in serum and urine and of its urinary metabolite 1,2,4-triazole-3-carboxamide 420(1987)189

**Riboflavin**

HPLC-diode-array determination of ascorbic acid, thiamine and riboflavin in goats' milk 410(1987)201

Quantification of riboflavin, riboflavin 5'-phosphate and flavin adenine dinucleotide in plasma and urine by HPLC 423(1987)105

**Riboflavin 5'-phosphate**

Quantification of riboflavin, riboflavin 5'-

phosphate and flavin adenine dinucleotide in plasma and urine by HPLC 423(1987)105

#### **Ribonucleases**

Determination of monosaccharides in glycoproteins by reversed-phase HPLC on 2.1-mm narrowbore columns 443(1988)299

#### **Ribonucleosides**

Separation of modified nucleic acid constituents by micellar electrokinetic capillary chromatography 409(1987)193

#### **Ribonucleotides**

Sensitive on-line radioactivity measurements with a heterogeneous flow-cell: application to lymphoid cell ribonucleotides separated by HPLC 415(1987)383

#### **Ribose**

HPLC separation of some mono- and disaccharides with detection by a post-column enzyme reactor and a chemically modified electrode 408(1987)157

#### **Ribose-binding protein**

Detection of ligand-protein binding by direct electrophoresis of the complex 449(1988)331

#### **Ribosomal proteins**

Chromatography of complex protein mixtures (Review) 418(1987)115

#### **Ribosome-inactivating proteins**

Large scale chromatographic purification of ribosome-inactivating proteins 408(1987)235

#### **Ribostamycin**

Identification of dekamycin antibiotics 440(1988)71

#### **Rifampicin**

LC assay for the simultaneous determination of pyrazinamide and rifampicin in serum samples from patients with tuberculous meningitis 420(1987)73

Determination of rifampicin and its main metabolites in human plasma by HPLC 426(1988)412

#### **Rilmazafone**

Combined HPLC and enzyme immunoassay for active metabolites of a sleep inducer (rilmazafone), a ring-opened derivative of benzodiazepines, in human plasma 417(1987)216

#### **Rimantadine**

Quantitation of the enantiomers of rimantadine in human plasma and urine by GC-MS 431(1988)343

#### **Ritodrine**

Determination of ritodrine in blood and plasma by HPLC with fluorescence detection 416(1987)400

#### **RNA**

Effect of phosphate concentration on the yield and purity of DNA separated from RNA by hydroxyapatite chromatography 411(1987)481

Resolution of RNA using HPLC (Review) 418(1987)51

High-resolution chromatography of nucleic acids on the Gen-Pak FAX column 444(1988)47

#### **rRNA**

Isolation of high-molecular-weight nucleic acids for copy number analysis using HPLC 402(1987)189

#### **tRNA**

Isolation of high-molecular-weight nucleic acids for copy number analysis using HPLC 402(1987)189

Resolution of RNA using HPLC (Review) 418(1987)51

Purification of transfer RNA species by single-step ion-exchange HPLC 444(1988)79

#### **Robotic system**

Totally automated robotic system for LC analysis of solid dosage formulations with the aid of a reduced Kalman filter 438(1988)319

#### **Rodenticides**

Modification to procedures for the determination of chlorofacinone and for multi-residue analysis of rodenticides in animal tissues 437(1988)301

#### **Roquefortines**

Standardized HPLC of 182 mycotoxins and other fungal metabolites based on alkylphenone retention indices and UV-VIS spectra (diode array detection) 404(1987)195

#### **Rose Bengale**

Simple method for the analysis of food dyes on reversed-phase thin-layer plates 411(1987)437

#### **Rosefuran**

Formal total synthesis of rosefuran and eldanolide 440(1988)449



**Rosenonolactone**

Simultaneous detection of trichothecenes and rosenonolactone in grape juice and wine by capillary GC 448(1988)424

**Rose oil**

Importance of the odoriferous principle of Bulgarian rose oil in flavour and fragrance chemistry 406(1987)181

Composition of essential oils. Part 7. Bulgarian oil of rose (*Rosa damascena* Mill.) 406(1987)185

**Rotenoids**

Chiral-phase HPLC of rotenoid racemates 404(1987)133

**Roxatidine**

Determination of roxatidine in human plasma, urine and milk by capillary GC using nitrogen-selective detection 432(1988)199

**Roxithromycin**

Analysis of erythromycin and roxithromycin in plasma or serum by HPLC using electrochemical detection 433(1988)298

**Rubidium**

Fractionation of rubidium isotopes in cation-exchange chromatography 438(1988)225

**Rubidium isotopes**

Separation of rubidium isotopes by ion-exchange chromatography 435(1988)253

**Rubreserine**

Metabolism of physostigmine in mouse liver microsomal incubations studied by LC with dual-electrode amperometric detection 419(1987)165

**S****Sabinene**

Resolution of monoterpene enantiomers by GC 407(1987)243

**Saccharides**

see also Sugars and Carbohydrates

Two HPLC columns for analyses of malto-oligosaccharides 409(1987)396

Rapid and sensitive method for measurement of hyaluronic acid and isomeric chondroitin sulfates using HPLC 413(1987)9

LC assay for constituent disaccharides of hyaluronic acid and chondroitin sulphate isomers 415(1987)231

TLC using multiple development for analysis of reaction products of sucrases 416(1987)353

**Saccharin**

Determination of saccharin in diet and biological materials 441(1988)448

**Saccharose**

Preparative isolation of <sup>14</sup>C-labelled saccharose from a partly purified fraction of *Chlorella vulgaris* 442(1988)237

**Safflor yellow A**

HPLC of carthamin, safflor yellow A and a precursor of carthamin. Application to the investigation of an unknown red pigment produced in cultured cells of safflower 438(1988)61

**Safrole**

Determination of  $\beta$ -asarone, safrole, isosafrole and anethole in alcoholic drinks by HPLC 404(1987)273

Determination of safrole and myristicin in nutmeg and mace by HPLC 438(1988)117

**Safrole stationary phase**

Stationary phases. XV. HPLC behaviour of quinones on a safrole stationary phase 408(1987)323

**Salannin**

Isolation and purification of salannin from neem seeds and its quantification in neem and chinaberry seeds and leaves 447(1988)277

**Salbutamol**

Analysis of salbutamol enantiomers in human urine by chiral HPLC and preliminary studies related to the stereoselective disposition kinetics in man 422(1987)187

**Salicylamide glucuronide**

HPLC method for the quantitation of salicylamide and its metabolites in biological fluids 420(1987)313

**Salicylamide sulphate**

HPLC method for the quantitation of salicylamide and its metabolites in biological fluids 420(1987)313

**Salicylate affinity column**

Selective purification of cardiac myosin by a high-performance salicylate affinity column 435(1988)185

**Salicylic acid**

Rapid HPLC assay for the simultaneous analysis of non-steroidal anti-inflammatory drugs in plasma 416(1987)293

Determination of plasma aspirin and salicylic acid concentrations after low aspirin doses by HPLC with post-column hydrolysis and fluorescence detection 420(1987)425

1-(4-Hydroxyphenyl)-, 1-(2,4-dihydroxyphenyl)- and 1-(2,5-dihydroxyphenyl)-2-bromoethanones: new labels for determination of carboxylic acids by HPLC with electrochemical and UV detection 442(1988)209

**Salicylic acids**

Cyclic di-*tert.*-butylsilylene derivatives of substituted salicylic acids and related compounds. A study by GC-MS 441(1988)13

**Salicyluric acid**

Rapid HPLC assay for the simultaneous analysis of non-steroidal anti-inflammatory drugs in plasma 416(1987)293

**Salicylamide**

HPLC method for the quantitation of salicylamide and its metabolites in biological fluids 420(1987)313

**Salmonella**

GC detection of bacteria causing enteric fever 415(1987)377

**Salsolinol**

Analysis of salsolinol in human brain using HPLC with electrochemical detection 428(1988)152

**Sample clean-up**

Clean-up of environmental samples by HPLC for analysis of organochlorine compounds by GC with electron-capture detection 435(1988)241

Simultaneous clean-up of nitroarenes and polycyclic aromatic ketones from soil and particulate matter on XAD-2 435(1988)489

**Sample load**

Effects of large sample loads on column lifetime in preparative-scale LC 410(1987)43

**Sample preconcentration**

Ion-interaction reversed-phase chromatographic method for the determination of gold(I) cyanide in mine process liquors using automated sample preconcentration 439(1988)23

Studies on sample preconcentration in ion chromatography. VII. Review of the methodology and applications of anion preconcentration 439(1988)37

**Sample preparation**

Simple steam distillation for sample preparation prior to GC determination of organic compounds 408(1987)201

HPLC determination of six *p*-hydroxybenzoic acid esters in cosmetics using Sep-Pak Florisil cartridges for sample pre-treatment 410(1987)413

Analysis of drugs from biological fluids using disposable solid phase columns 411(1987)325

Determination of chloramphenicol in swine muscle tissue using a monoclonal antibody-mediated clean-up procedure 411(1987)415

Sample preparation and column regeneration in biopolymer separations (Review) 418(1987)27

Direct determination of dextromethorphan and its three metabolites in urine by HPLC using a precolumn switching system for sample clean-up 422(1987)340

Automatic determination of urinary 4-hydroxy-3-methoxymandelic (vanillylmandelic) acid by LC with electrochemical detection 424(1988)255

Amino acid analysis of plasma: studies in sample preparation 431(1988)163

Isolation of basic drugs from plasma using solid-phase extraction with a cyanopropyl-bonded phase 432(1988)209

Comparison of solid-phase extraction and liquid-liquid extraction for HPLC analysis of antipyrine in plasma 434(1988)324

Ion-interaction reversed-phase chromatographic method for the determination of gold(I) cyanide in mine process liquors using automated sample preconcentration 439(1988)23

**Sample pretreatment**

Separation of proteins by reversed-phase HPLC. II. Optimizing sample pretreatment and mobile phase conditions 443(1988)381

**Sample reconcentration**

Sample reconcentration by column-external solvent evaporation or injection of large volumes into GC capillary columns? 404(1987)297

**Sample solvent effects**

Sample solvent effects in an apparent chiral HPLC separation on  $\beta$ -cyclodextrin 448(1988)31

**Sample throughput**

Split peaks in non-linear chromatography and their effect on sample throughput in large scale separations 449(1988)53

**Sample transfer**

Factors affecting sample transfer from microlitre syringes 441(1988)367

**Sample treatment**

Zone-electrophoretic sample treatment for LC 436(1988)421

**Sampling**

Evaluation of an automatic siphonic sampler for capillary zone electrophoresis 404(1987)313

Dynamic headspace GC: concentration of volatile components after thermal desorption by intermediate cryofocusing in a cold trap. I. Principle and applications 405(1987)87

Dynamic headspace GC: concentration of volatile components after thermal desorption by intermediate cryofocusing in a cold trap. II. Effect of sampling and desorption parameters on recovery 405(1987)99

**Sampling methods**

Cryogenic sampling for aniline vapor with subsequent LC-electrochemical detection analysis 408(1987)81

High-volume elution chromatography of dichlorobenzenes on a polyurethane foam-Tenax sandwich cartridge 409(1987)235

Sampling and injection system for the analysis of metalorganic chemicals by GC-MS. Some initial results 411(1987)445

Comparison of the performances of hot and cold sample introduction with a programmed-temperature vaporizer in the split and splitless modes 438(1988)243

Simple permeation absorber for sampling and preconcentrating hazardous air contaminants 439(1988)283

**Sampling temperature**

Dynamic headspace GC: concentration of

volatile components after thermal desorption by intermediate cryofocusing in a cold trap. II. Effect of sampling and desorption parameters on recovery 405(1987)99

**Saponins**

Determination of monosaccharides constituting the glycosides in saponins by HPLC 403(1987)319

Application of semi-preparative HPLC to difficult natural product separations 406(1987)367

HPLC of glycosides on a new type of hydroxyapatite column 407(1987)205

HPLC determination of molluscicidal saponins from *Phytolacca dodecandra* (Phytolaccaceae) 448(1988)265

Separation of natural products by centrifugal partition chromatography 450(1988)91

**Sarpedobilin**

Interactions between ascorbic acid and linear and cyclized bile pigments. A TLC study 440(1988)275

**Scanning photometer**

Quantitative and automatic digital evaluation of one- and two-dimensional thin-layer chromatograms using vertical surface absorption at an optimum wavelength 437(1988)411

**Schiff bases**

Ligand exchange chromatography of amino alcohol enantiomers as Schiff bases 406(1987)343

**Sedimentary rocks**

Determination of trace amounts of iodine in sedimentary rocks by ion chromatography with amperometric detection 436(1988)107

**Sedimentation-flotation focusing field-flow fractionation**

Flow velocity profile in channels with modulated cross-sectional permeability for sedimentation-flotation focusing field-flow fractionation using step density gradients 404(1987)23

**Selectivity**

Factor analysis of trends in selectivity in GLC of 175 saturated and unsaturated esters on 14 polysiloxane stationary phases 404(1987)11

Mechanism of selectivity in ion-pair HPLC of aminoglycoside antibiotics using perfluorinated pairing ions 404(1987)53

Characterization of porous polymers by polar strength and selectivity 404(1987)145

Molecular theory of selectivity based on solute structure for nematic liquid-crystalline phases in GLC 406(1987)27

Retention and selectivity in amino, cyano and diol normal bonded phase HPLC columns 410(1987)249

Chromatography of proteins on hydrophobic interaction and ion-exchange chromatographic matrices: mobile phase contributions to selectivity 443(1988)173

Micro-LC and the chiral recognition mechanism on albumin-coated silica gel. Large selectivity changes with sample size 447(1988)297

Methods for characterization of selectivity in reversed-phase LC. IV. Retention behaviour of oligomeric series 449(1988)361

#### **Selectivity coefficients**

Shape selectivity of stationary phases for the separation of isomeric hydrocarbons by capillary GC 446(1988)339

#### **Selectivity index**

Classification of GC stationary phases by a new variant of Kováts retention index 406(1987)105

#### **Selenium**

HPLC of selenium in biological samples 426(1988)385

1,2-Diaminobenzene derivatives for the determination of selenium by GC 440(1988)333

#### **Selenometalates**

Retention of thio- and selenometalates in mobile-phase ion chromatography 439(1988)91

#### **Self-sharpening concentration fronts**

Free energy consumption at self-sharpening concentration fronts in isothermal fixed-bed adsorption. Theory for isotherms of arbitrary shape and applications in displacement chromatography 409(1987)1

#### **Sendai virus proteins**

Effect of detergents on the structure of integral membrane proteins of Sendai virus studied with size-exclusion HPLC and monoclonal antibodies 443(1988)255

#### **Senosides**

Rapid and simple determination of senoside A and B in Rhei Rhizoma by ion-pair HPLC 403(1987)253

#### **Sensitivity**

Effect of column inlet concentration profile on the sensitivity of analysis. A quantitative assessment 409(1987)360

Effect of suppression efficiency on sensitivity in ion chromatography 439(1988)151

#### **Separation conditions**

Some problems associated with the selection of separation conditions in preparative LC 436(1988)367

#### **Separation time**

Multi-variate ranking function for optimizing separations 439(1988)173

#### **SE phases**

Estimation of the polarity of stationary phases by proton nuclear magnetic resonance spectroscopy. Application to phenyl and methyl silicones (OV and SE series) 402(1987)323

#### **Sequencing**

HPLC in protein sequence determinations 440(1988)231

#### **Sequencing methods**

Analytical strategy for determination of active site sequences in aminoacyl-tRNA synthetases 438(1988)347

#### **Serotonin**

Simultaneous LC determination of seventeen of the major monoamine neurotransmitters, precursors and metabolites. I. Optimization of the mobile phase using factorial designs and a computer program to predict chromatograms 415(1987)261

Simultaneous LC determination of seventeen of the major monoamine neurotransmitters, precursors and metabolites. II. Assessment of human brain and cerebrospinal fluid concentrations 415(1987)275

Determination of plasma serotonin by HPLC with pre-column sample enrichment and fluorimetric detection 417(1987)378

Determination of serotonin and 5-hydroxyindoleacetic acid in urine by reversed-phase ion-pair partition chromatography with fluorimetric detection 419(1987)85

Combined ion-pair extraction and HPLC for the determination of the biogenic amines and their major metabolites in single brain tissue samples 421(1987)51

Simultaneous determination of the major acidic metabolites of catecholamines and serotonin in urine by LC with electrochemical detection after a one-step sample clean-up on Sephadex G-10; influence of vanilla and banana ingestion 424(1988)273

Automatic determination of serotonin in biological fluids by LC 427(1988)326

Simultaneous micro-determinations of serotonin and 5-hydroxyindole-3-acetic acid with 5-hydroxy-N<sub>ω</sub>-methyltryptamine, as an internal standard, in biological materials by HPLC with electrochemical detection 430(1988)381

Determination of plasma 5-hydroxytryptophan, 5-hydroxytryptamine, 5-hydroxyindoleacetic acid, tryptophan and melatonin by HPLC with electrochemical detection 431(1988)170

Detection by HPLC of polyamines formed by clostridal putrefaction of caseins 442(1988)267

Perfluorinated acids as ion-pairing agents in the determination of monoamine transmitters and some prominent metabolites in rat brain by HPLC with amperometric detection 449(1988)191

#### **Serpentine**

Determination of *Catharanthus* alkaloids by reversed-phase HPLC 410(1987)488

#### ***Serratia liquefaciens***

Studies of metabolites in diarrheal stool specimens positive for *Klebsiella*, *Serratia*, and *Proteus* spp. by frequency-pulsed electron-capture GC 430(1988)209

#### **Sertraline**

GC-MS analysis and preliminary human pharmacokinetics of sertraline, a new antidepressant drug 417(1987)197

#### **Serum proteins**

Detection of the changes in serum proteins during chemical hepatocarcinogenesis by two-dimensional electrophoresis under non-denaturing conditions 419(1987)315

Pulse injection analysis of the binding of serum proteins to porous polymer gels modified with formyl groups 420(1987)13

#### ***Sesamum angolense***

Isolation of biologically active plant constituents by LC 450(1988)3

#### **Sesquiterpene lactones**

Sesquiterpene lactones from *Arnica chamissonis* Less. VI. Identification and quantitative determination by HPLC and GC 410(1987)329

Isolation of parthenolide by droplet counter-current chromatography 446(1988)157

#### **Sesquiterpenes**

Separation of non-polar sesquiterpene olefins from tolu balsam by HPLC; silver perchlorate impregnation of a prepacked preparative silica gel column 407(1987)343

Glass capillary GC analysis of volatile components of canned Philippine pineapple juice 409(1987)409

#### **Shale oil**

Extraction and identification of nitrile compounds in Rundle shale oil 435(1988)193

#### **Shielded hydrophobic phase**

Shielded hydrophobic phase; a new concept for direct injection analysis of biological fluids by HPLC 433(1988)264

#### **Siderophores**

Assignment of the configurations of the amino acids in peptidic siderophores 438(1988)291

#### **Signal enhancement**

Signal enhancement by on-column fluorometric detection in HPLC using micropacked fused-silica columns 435(1988)319

#### **Silanization**

On-line derivatization of silica supports for regeneration and preparation of reversed-phase columns 409(1987)155

#### **Silanol groups**

Small populations of highly retentive silanols on trimethylsilylated silica surfaces 407(1987)171

Polar contributions of the stationary phase to the reversed-phase ion-pair HPLC separation of quaternary ammonium drugs 437(1988)83

#### **Silica gel**

Investigation of HPLC packings by liquid mixture adsorption 406(1987)311

Accessibility of irregular (fractal) surfaces of porous silicas to silylation reactions 406(1987)317

Small populations of highly retentive silanols on trimethylsilylated silica surfaces 407(1987)171

Effect of zirconia on the sorption properties of silica gels 407(1987)189

Preparation of porous bead silica 435(1988)55

GC grade silica gel from paddy husk ash 437(1988)244

Influence of the porous silica gel structure on the coverage density of a chemically bonded C<sub>18</sub> phase for HPLC 446(1988)191

Comparison of mesoporous controlled-porosity glasses to silica gel as solid supports for C<sub>18</sub> chemically bonded phases 446(1988)347

Effect of the pore structure of silica gel on retention behaviour in pressure programming supercritical fluid chromatography 448(1988)87

Internal structures of wide-pore packing materials for HPLC studied by transmission electron microscopy 448(1988)95

Influence of silica surface chemistry and structure on the properties, structure and coverage of alkyl-bonded phases for HPLC (Review) 449(1988)1

Liquid CC using TLC silica gel 450(1988)443

#### **Silicalite**

Use of silicalite molecular sieve for the GC determination of permanent gases and volatile hydrocarbons emitted during coal processing 405(1987)213

#### **Silica surface chemistry**

Influence of silica surface chemistry and structure on the properties, structure and coverage of alkyl-bonded phases for HPLC (Review) 449(1988)1

#### **Silicates**

GC procedure for the qualitative analysis of silicone polyester resins after alkaline fusion 441(1988)207

#### **Silicato complexes**

Formation of silicato complexes of aluminium in aqueous solution 403(1987)151

#### **Silicone polyester resins**

GC procedure for the qualitative analysis of silicone polyester resins after alkaline fusion 441(1988)207

#### **Silicon number**

Determination of silicon number in silylated organic compounds using the hydrogen atmosphere flame ionization detector 408(1987)329

#### **Silver perchlorate**

Separation of non-polar sesquiterpene olefins from tolu balsam by HPLC; silver perchlorate impregnation of a prepacked preparative silica gel column 407(1987)343

#### **Silylated N,O-ketals**

Silylated N,O-ketals from the reaction of ketones with N-trimethylsilylimidazole 419(1987)271

#### **Silylation reactions**

Accessibility of irregular (fractal) surfaces of porous silicas to silylation reactions 406(1987)317

#### **Simazine**

Rapid determination of simazine and atrazine in water at the parts per trillion (10<sup>12</sup>) level. Extraction by a miniaturized Carbo-pack B trap followed by HPLC 405(1987)357

#### **Siphonic sampler**

Evaluation of an automatic siphonic sampler for capillary zone electrophoresis 404(1987)313

#### **Sisomicin**

Pre-column derivatization of sisomicin with *o*-phthalaldehyde- $\beta$ -mercaptopropionic acid and its application to sensitive HPLC determination with fluorimetric detection 425(1988)143

#### **Sitosterol**

N-Methoxy-N,O-bis(trimethylsilyl) carbamate as a derivatizing reagent for the GC of sitosterol degradation products 446(1988)253

#### **Slurry packing**

Contribution to the packing of LC microcolumns 411(1987)185

#### **Smectite**

Organo-modified smectite for the GC separation of various compounds 410(1987)121

#### **Snyder displacement model**

Application of the Snyder displacement model to aromatic nitrogen compounds on an aminopropyl bonded-phase column 405(1987)107

#### **Soap**

Anomalous LC results originating from the presence of soap residues and the use of nylon filters 408(1987)296

#### **Sodium**

Simultaneous determination of sodium and chloride ions by ion chromatography at the  $\mu\text{g/l}$  level 404(1987)285

Isotachophoretic determination of the cationic constituents in mixtures of raw materials for glass manufacturing 439(1988)423

#### **Sodium deoxycholate**

Analysis of Salmonella lipopolysaccharides by sodium deoxycholate-polyacrylamide gel electrophoresis 450(1988)381

#### **Sodium dodecyl sulphate**

Removal of sodium dodecyl sulphate from proteins by GPC 423(1987)376

#### **Sodium glyco-7 $\alpha$ ,12 $\alpha$ -dihydroxy-5 $\beta$ -cholan-24-oate**

Internal standards for quantitative GC of individual bile acids after group separation of bile acids in urine 422(1987)25

#### **Sodium hypochlorite**

LC determination of penicillins by postcolumn degradation with sodium hypochlorite using an hollow-fibre membrane reactor 447(1988)365

#### **Sodium iodide**

Selective detection of sulphoxides and sulphimides by TLC using trifluoroacetic anhydride-sodium iodide as a reagent 447(1988)225

#### **Sodium monofluorophosphate**

Determination of potassium nitrate and sodium monofluorophosphate in the presence of phosphate and sulfate by high-resolution ion chromatography 410(1987)494

#### **Sodium salicylate**

Non-suppressed ion chromatography of arsenic anions with potassium hydroxide-aromatic salt mixed eluents 447(1988)323

#### **Sodium tartrate**

Paper electrophoretic study of ion-pair formation. XVII. Influence of the enantiomeric purity of the sodium tartrate used as electrolyte 440(1988)361

#### **Sodium tauro-7 $\beta$ ,12 $\beta$ -dihydroxy-5 $\beta$ -cholan-24-oate**

Internal standards for quantitative GC of individual bile acids after group separation of bile acids in urine 422(1987)25

#### **Sodium valproate**

Gas chromatographic method for the measurement of sodium valproate utilization by kidney tubules 414(1987)460

#### **Software**

Optimization of ion chromatography using commercially available detection systems and software 439(1988)13

#### **Solid-phase adsorption kinetics**

Theoretical analysis of the asymmetry in chromatographic peaks 404(1987)1

#### **Solid-phase extraction**

Analysis of drugs from biological fluids using disposable solid phase columns 411(1987)325

Qualitative GC and GC-MS screening for  $\beta$ -blockers in urine after solid-phase extraction using Extrelut-1 columns 426(1988)194

Comparison of solid-phase extraction and liquid-liquid extraction for HPLC analysis of antipyrine in plasma 434(1988)324

#### **Solubility factors**

Solubility factors established by GLC. A balance sheet 406(1987)51

#### **Solubility properties**

Solubility properties in polymers and biological media. II. A new method for the characterisation of the adsorption of gases and vapours on solids 409(1987)15

#### **Solute-solvent association constant**

Thermochemical investigations of GLC. IV. Partition coefficients of solutes on binary mixtures containing two complexing solvents 402(1987)41

#### **Solute-solvent interactions**

Solute-solvent interactions in tetra-*n*-butylphosphonium salts studied by GC 438(1988)1

#### **Solvent evaporation**

Sample reconcentration by column-external solvent evaporation or injection of large volumes into GC capillary columns? 404(1987)297

#### **Solvent localization parameters**

Solvent selectivity in the resolution of some regioisomeric and diastereomeric prostaglandin intermediates on silica 449(1988)77

#### **Solvent peak**

Resolution of overlapped chromatograms by the use of a Kalman filter. III. Effects of overlapping and outstripping of a solvent peak on the analysis 411(1987)95

**Solvent refined coal**

On-line microbore HPLC capillary GC-MS. II. Application to the analysis of solvent refined coal 409(1987)213

**Solvents**

Determination of residual solvents in pharmaceutical products by GC coupled to a head-space injection system and using an external standard 403(1987)109

Chemical interactions as a possible limitation on the useful solvent properties of liquid alkylammonium salts 435(1988)17

Prediction of HPLC capacity factors following column calibration. Influence of different solvents and columns on the separation of a series of aromatic amides 437(1988)13

Static headspace GC method for the determination of residual solvents in vigabatrin drug substance 438(1988)103

**Solvent selectivity**

Solvent selectivity in the resolution of some regioisomeric and diastereomeric prostaglandin intermediates on silica 449(1988)77

**Solvent strength parameters**

Linear model for the determination of solvent strength parameters in LSC 437(1988)3

**Solvent systems**

New solvent system for HPTLC and HPLC of gangliosides 405(1987)125

**Solvophobic theory**

Reversed-phase HPLC of pyrimidine bases and nucleosides. Application of solvophobic theory 445(1988)59

**Somatomedin C**

Reduction studies on bacterial recombinant somatomedin C/insulin-like growth factor-1 443(1988)183

**Somatomedins**

Application of preparative reversed-phase HPLC to the isolation of insulin-like growth factor II from human serum 421(1987)61

**Sorbents**

Hydrolysed macroporous glycidyl methacrylate-ethylene dimethacrylate copolymer sorbent for size-exclusion HPLC of synthetic polymers and biopolymers 435(1988)357

Fundamentals of the theory and practice of

polymer GPC as a method of chromatographic porosimetry (Review) 448(1988)307

**Sotalol**

LC determination of sotalol in plasma and urine employing solid-phase extraction and fluorescence detection 421(1987)309

Improved HPLC method for the analysis of serum sotalol 427(1988)181

**Soya bean oil**

Analysis of the epoxidised soya bean oil additive in plastics by GC 437(1988)247

**Sparsomycin**

Direct broth assay for sparsomycin and related nucleoside antitumor antibiotics using reversed-phase HPLC 439(1988)459

**Sparteine**

GC determination of sparteine and 2- and 5-dehydrosparteine in plasma and urine 431(1988)203

**Spermidine**

Improved method for benzoyl chloride derivatization of polyamines for HPLC 408(1987)227

Determination of polyamines in human erythrocytes by capillary GC with nitrogen-phosphorus detection 415(1987)27

Simultaneous determination of  $\gamma$ -aminobutyric acid and polyamines by HPLC 419(1987)285

Reversed-phase LC separation and simultaneous fluorimetric detection of polyamines and their monoacetyl derivatives in human and animal urine, serum and tissue samples: an improved, rapid and sensitive method for routine application 430(1988)249

Chromatography, flow injection analysis and electrophoresis in computer-assisted comparative biochemistry: its application and possibilities in clinical research. Preliminary studies on Crohn's disease 440(1988)261

Quantitation of polyamines using TLC and image analysis 441(1988)361

HPLC method for the determination of  $\epsilon$ -( $\gamma$ -glutamyl)lysine and mono- and bis- $\gamma$ -glutamyl derivatives of putrescine and spermidine 443(1988)329

**Spermine**

Improved method for benzoyl chloride



derivatization of polyamines for HPLC 408(1987)227

Determination of polyamines in human erythrocytes by capillary GC with nitrogen-phosphorus detection 415(1987)27

Simultaneous determination of  $\gamma$ -aminobutyric acid and polyamines by HPLC 419(1987)285

Reversed-phase LC separation and simultaneous fluorimetric detection of polyamines and their monoacetyl derivatives in human and animal urine, serum and tissue samples: an improved, rapid and sensitive method for routine application 430(1988)249

Chromatography, flow injection analysis and electrophoresis in computer-assisted comparative biochemistry: its application and possibilities in clinical research. Preliminary studies on Crohn's disease 440(1988)261

Quantitation of polyamines using TLC and image analysis 441(1988)361

#### **Spheron derivatives**

Ion-exchange derivatives of Spheron. VI. Strongly basic derivatives 440(1988)287

#### **Sphingolipids**

Behaviour of plasmalogens during HPLC on a silica column with a mobile phase containing phosphoric acid 434(1988)411

#### **Sphingomyelin**

Phospholipids in amniotic fluid with special reference to the lecithin; Sphingomyelin ratio (Review) 429(1988)359

#### **Spinacine**

Quantitative HPLC determination of the amino acid spinacine in blood and chow of rats 410(1987)504

#### **Spironolactones**

HPLC assay for the simultaneous measurement of two spironolactone analogues, 17 $\beta$ -hydroxy-3-oxo-17-pregn-4-ene-21-carboxylic acid- $\gamma$ -lactone 7 $\alpha$ -methyl ethyl ester and its 6 $\beta$ -hydroxylated derivative 428(1988)377

#### **Split-peak effect**

Non-linear elution effects in split-peak chromatography. I. Computer simulations for the cases of irreversible diffusion- and adsorption-limited kinetics 436(1988)111

#### **Splitless injector**

Checking of capacity of a splitless injector — a simple test 448(1988)387

#### **Split peaks**

Split peaks in non-linear chromatography and their effect on sample throughput in large scale separations 449(1988)53

#### **Splitter**

On-line microbore HPLC capillary GC-MS. I. Development of isotachic eluent splitters 409(1987)205

#### **Splitting ratios**

Investigation of pre-set and actual splitting ratios in split injection capillary GLC 404(1987)340

#### **Spray reagents**

Spray reagent for the detection of coumarins and flavonoids on thin-layer plates 403(1987)285

Detection of amino acids on thin-layer plates 436(1988)341

#### **Squalane**

Biomedical applications of chromatographic fraction containing trehalose dimycolate in squalane emulsion (Review) 440(1988)473

#### **Squalene**

Characterization of a lipid-rich fraction synthesized by *Streptomyces avermitilis* 441(1988)31

#### **Standard addition**

Comparison of the absolute calibration method with the method of standard addition for the determination of halothane in blood by GC headspace analysis 425(1988)99

#### **Starch**

Separation of starch components based on ligand-induced adsorption of amylose on cellulose 403(1987)373

#### **Stationary phase retention**

Cross-axis synchronous flow-through coil planet centrifuge for large-scale preparative counter-current chromatography. I. Apparatus and studies on stationary phase retention in short coils 449(1988)135

#### **Stationary phases, GC**

see also Chiral stationary phases

Disc-like liquid crystal stationary phases derived from hexahydroxybenzene 402(1987)73

- Carbon black adsorbents modified with *n*-hydrocarbons and treated with an high-frequency low-temperature plasma 402(1987)87
- Reduction of the number of characterizing probes in GC by factor analysis 402(1987)318
- Estimation of the polarity of stationary phases by proton nuclear magnetic resonance spectroscopy. Application to phenyl and methyl silicones (OV and SE series) 402(1987)323
- Characterization of porous polymers by polar strength and selectivity 404(1987)145
- Simple preparation of a Poly-S 179 capillary column for the analysis of polycyclic aromatic hydrocarbons and triglycerides 404(1987)248
- Heavy alkylated benzene stationary phase for the GLC separation of light hydrocarbons 404(1987)328
- Use of silicalite molecular sieve for the GC determination of permanent gases and volatile hydrocarbons emitted during coal processing 405(1987)213
- Molecular theory of selectivity based on solute structure for nematic liquid-crystalline phases in GLC 406(1987)27
- Hierarchical ascending classification of Kováts retention indices as a means of indicating siloxane stationary phase specificities towards aliphatic carboxylic esters 406(1987)95
- Classification of GC stationary phases by a new variant of Kováts retention index 406(1987)105
- Potential solid supports for GC prepared on the basis of certain rocks 406(1987)145
- Small populations of highly retentive silanols on trimethylsilylated silica surfaces 407(1987)171
- GC properties of porous polyacrylonitrile and divinylbenzene copolymers 408(1987)308
- Heavy alkylated benzene stationary phase for the GLC separation of ethylbenzene, xylenes, styrene, iso- and *n*-propylbenzene 409(1987)371
- Organo-modified smectite for the GC separation of various compounds 410(1987)121
- Development of a C<sub>18</sub>-bonded vinyl alcohol copolymer gel for reversed-phase HPLC 410(1987)129
- Ambiguities in the determination of McReynolds stationary phase constants 411(1987)43
- Liquid crystals. VI. Comparative study of the supercooling effect of some liquid crystalline stationary phases using GLC 411(1987)456
- Dispersive force parameters as a measure of the polarity of surfactants used as stationary phases in GLC 435(1988)29
- Preparation of porous bead silica 435(1988)55
- High-resolution GC with liquid crystal glass capillaries. X. Separation of diastereomeric alkanes 436(1988)47
- Modification to the chromatographic behaviour of Porapak Q by chemical reaction 437(1988)238
- GC grade silica gel from paddy husk ash 437(1988)244
- New liquid crystalline stationary phase for capillary GC 439(1988)381
- Polyester deactivation of pink supports and its effect on the GC of some polar, closely boiling and isomeric analytes 446(1988)31
- Use of modified tungsten selenide as an adsorbent for GC 446(1988)79
- Evaluation of interfacial surface areas in ultra-low-loaded GC columns 446(1988)221
- Shape selectivity of stationary phases for the separation of isomeric hydrocarbons by capillary GC 446(1988)339
- Cyclodextrins as chiral stationary phases in capillary GC. I. Pentylated  $\alpha$ -cyclodextrin 447(1988)193
- GC on porous polymers. IV. Influence of the geometric structure of porous copolymers of 1,4-di(methacryloyloxymethyl)naphthalene with 1,4-divinylbenzene on their chromatographic behaviour 448(1988)233
- Stationary phases, LC**
- Determination of the leakage from Phenyl-Sepharose CL-4B, Phenyl-Sepharose FF and Phenyl-Superose in bulk and column experiments 403(1987)85
- Melittin-silica, a high-pressure affinity chromatography resin for calmodulin 403(1987)99
- Interaction of proteins with immobilized Cu<sup>2+</sup>. Quantitation of adsorption capacity, adsorption isotherms and equilibrium constants by frontal analysis 403(1987)197
- Factor analysis of trends in selectivity in GLC of 175 saturated and unsaturated esters on 14 polysiloxane stationary phases 404(1987)11
- New high-performance gel filtration columns for protein separation 404(1987)333

- Development of N-hydroxysuccinimide ester silica, a novel support for high-performance affinity chromatography 405(1987)179
- Investigation of HPLC packings by liquid mixture adsorption 406(1987)311
- Accessibility of irregular (fractal) surfaces of porous silicas to silylation reactions 406(1987)317
- On the structure and chromatographic behaviour of polystyrene phases 406(1987)325
- Matrix evaluation for preparative high-performance affinity chromatography 407(1987)179
- Synthesis and characterization of stearyl methacrylate gel for reversed-phase LC 407(1987)197
- Separation of non-polar sesquiterpene olefins from tolu balsam by HPLC; silver perchlorate impregnation of a prepacked preparative silica gel column 407(1987)343
- Reordering/resolution of silica immobilized hydrogen bonding ligands used as stationary phases in HPLC 408(1987)93
- Rapid preparation of bovine mercaptalbumin by means of covalent chromatography on silica-based materials 408(1987)263
- Stationary phases. XV. HPLC behaviour of quinones on a saffrole stationary phase 408(1987)323
- Macroporous glucomannan spheres for the size-exclusion separation of aqueous macromolecules 409(1987)175
- Two HPLC columns for analyses of malto-oligosaccharides 409(1987)396
- Chemometric classification of reversed-phase HPLC columns 410(1987)31
- Effects of large sample loads on column lifetime in preparative-scale LC 410(1987)43
- Retention and selectivity in amino, cyano and diol normal bonded phase HPLC columns 410(1987)249
- Improvement in the resolution of *o*-phthalaldehyde derivatized amino acids by applying gradient steepness optimization to five reversed-phase columns of different lengths and particle sizes 411(1987)121
- New metal chelate adsorbent selective for proteins and peptides containing neighbouring histidine residues 411(1987)177
- Facile preparation of a non-porous strong anion-exchange column for proteins 411(1987)452
- Pulse injection analysis of the binding of serum proteins to porous polymer gels modified with formyl groups 420(1987)13
- Retention properties of internal-surface reversed-phase silica packing and recovery of drugs from human plasma 420(1987)297
- Synthesis of high-capacity immunoaffinity sorbents with oriented immobilized immunoglobulins or their F<sub>ab</sub>' fragments for isolation of proteins 424(1988)243
- Affinity chromatography of human thrombin on modified silica 424(1988)385
- Shielded hydrophobic phase; a new concept for direct injection analysis of biological fluids by HPLC 433(1988)264
- Separation of peptides by multimode chromatography on a column packed with vinyl alcohol copolymer gel 434(1988)71
- Direct plasma injection using internal surface reversed-phase HPLC: feasibility study using propofol as a model compound 434(1988)271
- Selective purification of cardiac myosin by a high-performance salicylate affinity column 435(1988)185
- Hydrolysed macroporous glycidyl methacrylate-ethylene dimethacrylate copolymer sorbent for size-exclusion HPLC of synthetic polymers and biopolymers 435(1988)357
- Effect of column temperature on the retention of peropyrene-type polycyclic aromatic hydrocarbons on various chemically bonded stationary phases in reversed-phase LC 436(1988)1
- Reversed-phase chromatographic separation and analysis of marine metal-organic complexes 436(1988)243
- Chromatographic and solid state NMR study of the changes in reversed-phase packings for HPLC at different eluent compositions 436(1988)345
- Retention behaviour of cyclodextrins and branched cyclodextrins on reversed-phase columns in HPLC 437(1988)47
- Polar contributions of the stationary phase to the reversed-phase ion-pair HPLC separation of quaternary ammonium drugs 437(1988)83
- Analytical high-performance affinity

- chromatography of ovalbumin-derived glycopeptides on columns of concanavalin A- and wheat germ agglutinin-immobilized gels 438(1988)73
- Mixed bonded phases for HPLC. I. Synthesis and physico-chemical characterization 438(1988)263
- New ion-exclusion phase for organic acids 439(1988)71
- Reversed-phase separation of a standard mixture of polycyclic aromatic hydrocarbons on an adamantyl surface 439(1988)414
- HPLC analysis of oligosaccharides. I. Separation on an ion-exchange stationary phase of low cross-linking 440(1988)281
- Ion-exchange derivatives of Spheron. VI. Strongly basic derivatives 440(1988)287
- High-performance adsorption chromatography of proteins on 2- $\mu$ m spherical beads of hydroxyapatite 440(1988)305
- Packing density of organodimethylsiloxy-covered silica gel and hydrolytic stability of the coating 440(1988)315
- Selectivity related to carbon loading and end-capping of octadecyl-bonded silica gels in reversed-phase LC 441(1988)183
- HPLC of plant-derived oligosaccharides on a new cation-exchange resin stationary phase: HPX-22H 441(1988)382
- Thermodynamic comparison of monomeric and polymeric C<sub>18</sub> bonded phases using aqueous methanol and acetonitrile mobile phases 442(1988)1
- Phase ratio determination in an ion-exchange column having pores partially accessible to proteins 442(1988)15
- Characterization of some styrene-divinylbenzene sorbents for reversed-phase chromatography 442(1988)97
- Use of a small TSK GSW HPLC column for large-zone chromatographic studies of monomer-oligomer equilibria of membrane protein 442(1988)229
- Comparative study of Zorbax Bio Series GF 250 and GF 450 and TSK-GEL 3000 SW and SWXL columns in SEC of proteins 443(1988)45
- Peptide characterization with a sulfoethyl aspartamide column 443(1988)63
- Synthesis of a non-porous, polystyrene-based strong anion-exchange packing material and its application to fast HPLC of proteins 443(1988)73
- Cation-exchange chromatography of peptides on poly(2-sulfoethyl aspartamide)-silica 443(1988)85
- Ion-exchange chromatography of proteins on a polyethyleneimine-grafted hydrophilic polymer for HPLC 443(1988)133
- Chromatographic evaluation of large-pore and non-porous polymeric reversed phases 443(1988)143
- Studies on the stabilization of reversed phases for LC 443(1988)155
- Chromatography of proteins on hydrophobic interaction and ion-exchange chromatographic matrices: mobile phase contributions to selectivity 443(1988)173
- Use of avidin-coated glass beads as a support for high-performance immunoaffinity chromatography 444(1988)21
- High-resolution chromatography of nucleic acids on the Gen-Pak FAX column 444(1988)47
- Comparison of Mono O, Superose-6, and ABx fast protein LC for the purification of IgM monoclonal antibodies 444(1988)153
- Rapid analysis of proteins and peptides by reversed-phase chromatography with polymeric micropellicular sorbents 445(1988)71
- New reversed-phase HPLC column for oligonucleotide separation 445(1988)225
- High-performance affinity chromatography of proteins on TSKgel Heparin-5PW 445(1988)234
- Influence of the porous silica gel structure on the coverage density of a chemically bonded C<sub>18</sub> phase for HPLC 446(1988)191
- Adsorption and chromatographic properties of modified silica sorbents for the production of viral preparations 446(1988)211
- Comparison of mesoporous controlled-porosity glasses to silica gel as solid supports for C<sub>18</sub> chemically bonded phases 446(1988)347
- Geometric structural properties of bonded layers of chemically modified silicas 447(1988)103
- Internal structures of wide-pore packing materials for HPLC studied by transmission electron microscopy 448(1988)95
- Modified general affinity adsorbent for large-scale purification of penicillinases 448(1988)109
- Influence of silica surface chemistry and structure

on the properties, structure and coverage of alkyl-bonded phases for HPLC (Review) 449(1988)1

Influence of bonding chemistry on the reordering/resolution of silica immobilized alkyl chains 449(1988)115

The rôle of the stationary phase in micellar liquid chromatography. Adsorption and efficiency 449(1988)349

Preparative HPLC of peptides on a new reversed-phase packing material, Kromasil C<sub>18</sub> 450(1988)71

Purification of specific heterologous F(ab)<sub>2</sub> fragments with DEAE-Zeta-Prep cartridges for ion-exchange chromatography 450(1988)133

Liquid CC using TLC silica gel 450(1988)443

#### Stationary phases, SFC

Effect of the pore structure of silica gel on retention behaviour in pressure programming supercritical fluid chromatography 448(1988)87

#### Stationary phase stability

Studies on the stabilization of reversed phases for LC 443(1988)155

#### Stationary phases, TLC

Preparation and evaluation of mono-, di- and triphenyl-treated plates for the HPTLC analysis of puerarin and paeonol 407(1987)314

TLC of metal complexes on glycidoxypropyl functionalized silica plates 409(1987)377

Comparison of non-hydrophobic C<sub>18</sub> bonded TLC and HPTLC plates for reversed-phase TLC of ecdysteroids 438(1988)419

#### Statistical model of overlap

Experimental affirmation of the statistical model of overlap 449(1988)41

#### Statistical moments

Chromatogram statistical moments from columns of finite length 445(1988)207

#### Steam distillation

Simple steam distillation for sample preparation prior to GC determination of organic compounds 408(1987)201

#### Stearyl methacrylate

Synthesis and characterization of stearyl methacrylate gel for reversed-phase LC 407(1987)197

#### Sterigmatocystin

Standardized HPLC of 182 mycotoxins and other fungal metabolites based on alkylphenone retention indices and UV-VIS spectra (diode array detection) 404(1987)195

HPLC procedure for determining the profiles of aflatoxin precursors in wildtype and mutant strains of *Aspergillus parasiticus* 441(1988)400

#### Steroidal alkaloids

Capillary GC of steroidal alkaloids from Solanaceae. Retention indices and simultaneous flame ionization/nitrogen-specific detection 442(1988)133

#### Steroid binding protein

Cyclic biospecific affinity chromatographic method for the purification of the sex steroid binding protein (SBP): application to the purification of SBP from toad 415(1987)297

#### Steroid hormone receptors

Chromatography of complex protein mixtures (Review) 418(1987)115

#### Steroid receptor proteins

High-resolution separation of molybdate-stabilized progesterin receptors using HPLC 425(1988)277

Influence of column length and pore size on high-performance ion-exchange chromatography of estrogen and progesterin receptors 425(1988)293

High-performance hydrophobic interaction chromatography as a means of identifying estrogen receptors expressing different binding domains 444(1988)225

#### Steroid receptors

Application of high-performance liquid chromatofocusing to the study of prostatic androgen receptors. Effect of stabilizing agents on the heterogeneity of receptor structure 403(1987)171

#### Steroids

see also Hormones

Analysis of estrogen sulphate mixtures in pharmaceutical formulations by reversed-phase chromatography 402(1987)273

Adsorption chromatographic behaviour of <sup>125</sup>I-labelled progesterone-11 $\alpha$ - and -12 $\alpha$ -succinyltyrosine methyl ester 404(1987)258

Separation of ecdysteroids by HPTLC using automated multiple development 408(1987)445

Optimization of the reversed-phase HPLC separation of synthetic estrogenic and progestogenic steroids using the multi-criteria decision making method 410(1987)1

TLC of ecdysone acyl esters and their 2,3-acetonide derivatives 411(1987)379

HPLC determination of methylprednisolone and methylprednisolone 21-[8-[methyl-(2-sulfoethyl)amino]-8-oxooctanoate] sodium salt in human plasma 414(1987)1

Column LC of the novel aldosterone antagonist, mespirenone, and its active metabolite in plasma 414(1987)217

Pregnenolone production by adrenal mitochondria: a new HPLC analytical method for cholesterol side-chain cleavage 414(1987)411

Determination of methylprednisolone in rat tissue by HPLC 415(1987)35

Studies on steroids. CCXXVII. Separation and determination of bile acid 7- and 12-sulphates in urine by HPLC with fluorescence labelling 415(1987)45

Three-hour assay for urinary free cortisol: historical comparison of three methods each using a different chromatographic procedure 415(1987)118

Determination of 17-oxosteroid glucuronides and sulphates in urine by LC using 2,4-dinitrophenylhydrazine as a prelabelling reagent for spectrophotometric detection 415(1987)253

Cyclic biospecific affinity chromatographic method for the purification of the sex steroid binding protein (SBP): application to the purification of SBP from toad 415(1987)297

Determination of some hydroxycholesterols in human serum samples 416(1987)1

Application of HPLC to the measurement of cortisol secretion rate 416(1987)25

Sensitive and selective assay of danazol in plasma by HPLC 416(1987)347

Selective on-line trace enrichment for the determination of ethynyl steroids in urine by LC with precolumn technology 417(1987)159

Determination of plasma corticosterone using HPLC 419(1987)280

Improved method for the determination of the major neutral steroids and unconjugated bile acids in human faeces using capillary GC 421(1987)21

Highly sensitive and simple determination of cholesterol and cholestanol in human serum by HPLC with fluorescence detection 421(1987)43

Simultaneous quantification of natural glycocorticoids and progestins in serum 421(1987)123

Determination of methylprednisolone metabolites in human urine by GC-MS 423(1987)15

Combined use of rapid-flow fractionation and HPLC for the determination of serum testosterone and application to study of stress response to physical exercise 423(1987)285

Identification and quantitation of trenbolone in bovine tissue by GC-MS 424(1988)219

Irregular retention properties of 21-aminosteroid antioxidants in octylsilane bonded-phase HPLC 424(1988)285

HPLC determination of a 21-aminosteroid antioxidant in plasma 424(1988)293

Studies on steroids. CCXXXIII. Separation and characterization of bile acid disulphates in human urine by HPLC 425(1988)59

Analysis of prednisolone acetate and related corticoids in swine plasma by reversed-phase HPLC 425(1988)162

Determination of cortisol in human plasma by capillary GC-MS using [<sup>2</sup>H<sub>5</sub>] cortisol as an internal standard 425(1988)237

Selective determination of urinary free cortisol by LC after solid-state extraction 426(1988)25

Automated direct assay system for the measurement of sex steroid hormones in serum using HPLC 426(1988)33

Simplified method for the determination of plasma 18-hydroxycorticosterone 426(1988)148

GC and mass spectral study of synthetic corticosteroid metabolism: fluorometholone 426(1988)399

HPLC determination of N-(2-methyl-2-propyl)-3-oxo-4-aza-5 $\alpha$ -androst-1-ene-17 $\beta$ -carboxamide, a 4-azasteroid, in human plasma from a phase I study 427(1988)79

Clinical analysis of individual steroids by column LC (Review) 429(1988)155

Improved method for the determination of the cortisol production rate using HPLC and liquid scintillation counting 430(1988)233

Analysis of methylprednisolone, methylprednisone and corticosterone for assessment of methylprednisolone disposition in the rat 430(1988)241

Determination of oestrogens in pregnancy urine by HPLC with fluorescence detection 431(1988)249

Determination of 17-hydroxyprogesterone in plasma by GC-MS with high-resolution selected-ion monitoring 432(1988)21

Simultaneous assay for individual sulphated 3 $\alpha$ - and  $\beta$ -hydroxysteroids in serum using HPLC combined with 3 $\alpha$ - and  $\beta$ -hydroxysteroid dehydrogenases immobilized on one column 432(1988)47

GC and MS study of betamethasone synthetic corticosteroid metabolism 432(1988)283

Specific reversed-phase LC method for analysis of steroids in Y-1 adrenal cell cultures 432(1988)302

Simultaneous analysis of methylprednisolone hemisuccinate, cortisol and methylprednisolone by normal-phase HPLC in human plasma 432(1988)308

Screening, confirmation and quantification of boldenone sulfate in equine urine after administration of boldenone undecylenate (Equipoise) 433(1988)9

Measurement of urinary estriol glucuronides during the menstrual cycle by HPLC 434(1988)61

Complete separation of nine equine oestrogens by HPLC 435(1988)385

Separation of ecdysone acyl esters and their 2,3-acetonide derivatives by HPLC 436(1988)279

Rapid and simple dansylation of phenolic steroids using a two-phase system and phase transfer catalysis 436(1988)429

Supercritical fluid chromatography of ecdysteroids 436(1988)497

Biological effect of guava oestrone on uterine tissue of mice 437(1988)316

Comparison of non-hydrophobic C<sub>18</sub> bonded TLC and HPTLC plates for reversed-phase TLC of ecdysteroids 438(1988)419

Rapid two-dimensional TLC system for the separation of multiple steroids of secretory and neuroendocrine interest 439(1988)441

Synthesis of [16,16,19-<sup>2</sup>H<sub>3</sub>; 19-<sup>3</sup>H]19-oxo-androst-4-ene-3,17-dione 440(1988)415

HPLC of hydroxysteroids detected with post-column immobilized enzyme reactors 441(1988)81

Analysis of ecdysteroids by supercritical-fluid chromatography 441(1988)165

Micro-assay for sheep 11 $\beta$ -hydroxylase activity using HPLC for steroid analysis 442(1988)424

HPLC determination of conjugated estrogens in tablets 450(1988)414

### **Sterols**

Separation and quantitation of fatty acids, sterols and bile acids in feces by GC as the butyl ester-acetate derivatives 415(1987)13

Anthracene-9-carbonyl chloride as a fluorescence and UV derivatising reagent for the HPLC analysis of hydroxy compounds 445(1988)393

### **Steryl ferulates**

Isolation and characterisation of intact steryl ferulates from seeds 440(1988)23

### **Stobadine**

GC determination of stobadine, N-desmethylstobadine and stobadine N-oxide in rat liver microsomal incubations 415(1987)163

Determination of stobadine, a novel cardioprotective drug, using capillary GC with nitrogen-phosphorus detection after its selective solid-phase extraction from serum 433(1988)269

### **Stobadine N-oxide**

GC determination of stobadine, N-desmethylstobadine and stobadine N-oxide in rat liver microsomal incubations 415(1987)163

### **Streaming current**

Characterization of mobile phases for the investigation of electrokinetic phenomena in LC 447(1988)177

### **Streaming potential**

Characterization of mobile phases for the investigation of electrokinetic phenomena in LC 447(1988)177

### **Structure-activity relationships**

Quantitative structure-activity relationship of coumarin derivatives. Involvement of partition between aqueous and membrane phase 445(1988)49

Hydrophobicity studies on diadamantanyl derivatives. Comparison between calculated partition coefficients and experimental

lipophilicity indices determined by reversed-phase HPLC and TLC 445(1988)317

#### **Structure–enantioselectivity relationships**

Chromatographic separation of the enantiomers of 1,3-dithiolane-1-oxides 450(1988)199

#### **Structure parameters**

Structure parameters of molecules and media evaluated by chromatographic partition. II. Geometrical exclusion in gels 410(1987)233

#### **Structure–retention relationships**

Correlation between molecular structures and retention indices of pyrazines 402(1987)309

Some problems in the correlation of molecular parameters and the connectivity index 403(1987)19

Relationships between retention behaviour and structure of N-substituted 1,2,4-triazoles in TLC on silica gel 404(1987)73

Propyl effect and retention–structure correlation as a means of GC identification 406(1987)43

Analysis of basic compounds on a silica column with an aqueous methanol eluent. The use of quantitative structure–retention relationships in metabolite identification 407(1987)1

Inverse GC study of polyether alcohols (in German) 410(1987)267

Prediction of retention indexes. I.

Structure–retention index relationship on apolar columns 436(1988)137

Effect of temperature on the GC separation of halogenated compounds on polar and non-polar stationary phases 437(1988)33

HPLC of alkylbenzenes on alumina 437(1988)177

#### **Styrene**

Normal- and reversed-phase separation of copolymers prepared from styrene and ethyl methacrylate 403(1987)280

Investigation by pyrolysis–GC of the composition of multicomponent polymeric microheterogeneous systems based on some vinyl monomers 404(1987)183

Heavy alkylated benzene stationary phase for the GLC separation of ethylbenzene, xylenes, styrene, iso- and *n*-propylbenzene 409(1987)371

#### **Styrene–divinylbenzene sorbents**

Characterization of some styrene–divinylbenzene sorbents for reversed-phase chromatography 442(1988)97

#### **Styrene–methyl methacrylate**

Determination of chemical composition distribution of styrene–methyl methacrylate copolymers by reversed-phase HPLC 447(1988)387

#### **Subcellular particles**

Gel electrophoresis of intact subcellular particles (Review) 418(1987)305

#### **Subcritical fluid chromatography**

Resolution of racemic amides and phosphine oxides on a  $\beta$ -cyclodextrin-bonded stationary phase by subcritical fluid chromatography 405(1987)135

Chiral resolution of a series of 3-thienylcyclohexylglycolic acids by liquid or subcritical fluid chromatography. A mechanistic study 450(1988)255

#### **Substance P**

Comparison of two methods for the determination of substance P immunoreactivity in human cerebrospinal fluid: sulphopropyl-Sephadex ion-exchange chromatography and/or reversed-phase HPLC combined with radiomimmunoassay 425(1988)175

Optimization of HPLC–radioimmunoassay protocols for the analyses of substance P and some of its metabolic fragments 432(1988)113

#### **Subtilisin**

HPLC method for the determination of the esterase activity of subtilisin and kallikrein 441(1988)431

Surface-mediated retention effects of subtilisin site-specific variants in cation-exchange chromatography 443(1988)193

#### **Subunit dissociation**

Reversible subunit dissociation of tumor necrosis factor during hydrophobic interaction chromatography 443(1988)205

#### **Succinate**

New method of purification of perchloric acid extracts for quantitative GC of some Krebs cycle acids 424(1988)357

#### **Succinimido $\alpha$ -naphthylcarbamate**

Amino acid analysis by reversed-phase HPLC. Automatic pre-column derivatization with activated carbamate reagent 407(1987)273



**Succinyl-1-(succinimidyl-4-(3-amidopropyl-silica)**

Development of N-hydroxysuccinimide ester silica, a novel support for high-performance affinity chromatography 405(1987)179

**Sucrases**

TLC using multiple development for analysis of reaction products of sucrases 416(1987)353

**Sugar phosphates**

Determination of photosynthetic intermediates in leaves by HPLC 402(1987)283

Ion chromatographic determination of sugar phosphates in physiological samples 439(1988)83

**Sugars**

see also Carbohydrates and saccharides

Elution characteristics of polyol-borate eluents in anion-exchange chromatography. I. Eluent strength of sugar-, uronate-borate eluents 403(1987)27

Ion-exclusion chromatography of non-ionic substances with conductivity detection 409(1987)271

Capillary column GC identification of sugars in honey as trimethylsilyl derivatives 410(1987)319

Capillary GC of neutral sugars as their aldonitrile acetates from the hydrolyzate of corn bran residues 410(1987)484

Prediction of the distribution coefficient ( $K_d$ ) variation with operating conditions in chromatographic systems 411(1987)81

GC analysis of neutral monosaccharides as their O-pentafluorobenzyloxime acetates 411(1987)275

Sugar-chain heterogeneity of human alkaline phosphatases: differences between normal and tumour-associated isozymes 413(1987)65

Analysis of tissue glycoprotein sugar chains by two-dimensional HPLC mapping 434(1988)51

Electrophoretic study of  $\alpha$ -D-galactosidases from seeds of *Glycine soja* and *Vigna radiata* possessing erythroagglutinating activity 436(1988)59

Degradation of ketoses during aqueous HPLC on lead-form cation-exchange resins 437(1988)387

Large-scale fractionation of S-form lipopolysaccharide from *Salmonella abortus equi*. Chemical and serological characterization of the fractions 440(1988)397

Analysis of the amino acid and sugar composition of streptococcal cell walls by GC-MS 441(1988)323

Optimizing copper-bicinchoninate carbohydrate detection for use with water-elution HPLC: a technique to measure the major mono- and oligosaccharides in small pieces of wheat endosperm 449(1988)207

**Sulforidazine**

Sensitive and specific HPLC assay for the quantification of sulforidazine and two diastereomeric sulforidazine-5-sulfoxide metabolites in plasma 423(1987)227

**Sulforidazine-5-sulphoxide**

Sensitive and specific HPLC assay for the quantification of sulforidazine and two diastereomeric sulforidazine-5-sulfoxide metabolites in plasma 423(1987)227

**Sulindac**

Analysis of sulindac and metabolites in plasma and urine by HPLC 413(1987)171

HPLC of sulindac and its sulphone and sulphide metabolites in plasma 414(1987)211

Rapid HPLC assay for the simultaneous analysis of non-steroidal anti-inflammatory drugs in plasma 416(1987)293

**Sulphadiazine**

HPLC method with electrochemical detection for the concomitant assay of sulphadiazine, sulphamerazine and sulphapyridine in plasma 428(1988)190

**Sulphadoxine**

Detection and determination of antimalarial drugs and their metabolites in body fluids (Review) 434(1988)1

**Sulpha drugs**

TLC separation of some sulpha drugs using acetoacetanilide as a coupling agent 441(1988)454

**Sulphaguanidine**

Fluorimetric determination of sulphanilamide and sulphaguanidine residues in animal tissues and products by ion-exchange HPLC (in German) 423(1987)217

**Sulphamerazine**

HPLC method with electrochemical detection for the concomitant assay of sulphadiazine, sulphamerazine and sulphapyridine in plasma 428(1988)190

**Sulphamethoxazole**

Rapid method for the determination of either plasma sulphapyridine or sulphamethoxazole and their acetyl metabolites using HPLC 414(1987)223

Concurrent analysis of methotrexate, trimethoprim, sulphamethoxazole and their major metabolites in plasma by HPLC 422(1987)328

HPLC analysis of trimethoprim and sulfamethoxazole in microliter volumes of chinchilla middle ear effusion and serum 433(1988)187

**Sulphamic acid 1,7-heptanediy ester**

GC assay for the new antitumor agent sulfamic acid diester (NSC 329680) and its stability in buffer, blood and plasma 427(1988)247

**Sulphaminoantipyrine**

Investigation of an alkylamine modified and pH-controlled mobile phase for separation of pyrazolone derivatives by HPLC 409(1987)81

**Sulphanilamide**

Fluorimetric determination of sulphanilamide and sulphaguanidine residues in animal tissues and products by ion-exchange HPLC (in German) 423(1987)217

**Sulphapyridine**

Rapid method for the determination of either plasma sulphapyridine or sulphamethoxazole and their acetyl metabolites using HPLC 414(1987)223

HPLC method with electrochemical detection for the concomitant assay of sulphadiazine, sulphamerazine and sulphapyridine in plasma 428(1988)190

**Sulphate**

Determination of potassium nitrate and sodium monofluorophosphate in the presence of phosphate and sulfate by high-resolution ion chromatography 410(1987)494

Improved method for the determination of sulphate in human serum using ion chromatography 413(1987)251

Sulphate quantification in rabbit tracheobronchial mucus by reversed-phase ion-pair chromatography 424(1988)147

Determination of phosphate, sulfate and oxalate in urine by ion chromatography 433(1988)373

Optimization of ion chromatography using

commercially available detection systems and software 439(1988)13

**Sulphide**

Derivatization of inorganic anions with pentafluorobenzyl methanesulphonate for GC 404(1987)175

Determination of oxidizable inorganic anions by HPLC with fluorescence detection and application to the determination of salivary nitrite and thiocyanate and serum thiocyanate 424(1988)29

Polymeric pentafluorobenzylating reagent for GC of inorganic anions with electron-capture detection 438(1988)253

Trace determination of sulfide by reversed-phase ion-interaction chromatography using pre-column derivatization 447(1988)415

**Sulphimides**

Selective detection of sulphoxides and sulphimides by TLC using trifluoroacetic anhydride-sodium iodide as a reagent 447(1988)225

**Sulphite**

Determination of mono- and divalent cations and anions in small fog samples by ion chromatography 439(1988)121

**Sulphite adducts**

Separation of sulphite adducts by ion chromatography with oxidative amperometric detection 403(1987)379

**4-Sulphociprofloxacin**

HPLC of ciprofloxacin and its metabolites in serum, urine and sputum 422(1987)153

**Sulphoethyl aspartamide**

Peptide characterization with a sulfoethyl aspartamide column 443(1988)63

**Sulphonamides**

HPTLC of dinitropyridyl-amino acids and sulphonamides on silanized silica gel 405(1987)394

Monitoring of veterinary drug residues by a combination of continuous flow techniques and column-switching HPLC. I. Sulphonamides in egg, meat and milk using post-column derivatization with dimethylaminobenzaldehyde 435(1988)97

**Sulphonated phosphines**

HPLC separation of sulphonated phosphines 408(1987)416

**Sulphonylureas**

Supercritical fluid extraction coupled with SFC for the separation of sulphonylurea herbicides and their metabolites from complex matrices 435(1988)63

**Sulphonyl-containing compounds**

Different partitioning behaviour of sulphonyl-containing compounds in reversed-phase HPLC and octanol-water systems 439(1988)237

**Sulphoxides**

Determination of low levels of phenothiazine sulphoxides in phenothiazine drug substances and formulations by TLC-second-derivative spectrofluorimetry 442(1988)363

Selective detection of sulphoxides and sulphimides by TLC using trifluoroacetic anhydride-sodium iodide as a reagent 447(1988)225

**Sulphur**

HPLC determination of sulphur and captan in a mixture 447(1988)263

**Sulphur-containing compounds**

High-efficiency separation and characterization of high-molecular-weight heterocyclic sulfur compounds in fossil fuels by microcolumn LC and related techniques 406(1987)375

Comparison of volatile components between fresh and rotten mussels by GC-MS 409(1987)251

Determination of sulphur compounds at low parts per billion levels by GC with flame photometric detection 435(1988)327

Capillary GC analysis of low-boiling organic sulphur compounds in anoxic lake-water by cryoadsorption 445(1988)97

**Sulphur heterocycles**

High-efficiency separation and characterization of high-molecular-weight heterocyclic sulfur compounds in fossil fuels by microcolumn LC and related techniques 406(1987)375

**Sulphur mustard**

TLC of mustard and its metabolites 437(1988)423

**Sulphur vesicants**

GC retention indices of sulfur vesicants and related compounds 436(1988)399

**Sun-screen agents**

Determination of sun-screen agents in cosmetic products by reversed-phase HPLC 408(1987)409

**Sunset Yellow FCF**

Simple method for the analysis of food dyes on reversed-phase thin-layer plates 411(1987)437

**Supercooling**

Liquid crystals. VI. Comparative study of the supercooling effect of some liquid crystalline stationary phases using GLC 411(1987)456

**Supercritical fluid chromatography**

Single and multiple gradients in supercritical fluid chromatography (Review) 402(1987)1

Temperature effect in packed microcolumn chromatography using a carbon dioxide mobile phase 402(1987)328

Application fields of a new evaporative light scattering detector for HPLC and supercritical fluid chromatography (in French) 404(1987)95

Retention in capillary supercritical fluid chromatography with carbon dioxide mobile phases containing polar modifiers 406(1987)223

Super- and subcritical fluid chromatography-Fourier transform IR spectrometry of medium polar compounds on polymeric styrene-divinylbenzene packed columns 407(1987)87

Comparison of packed column and capillary column SFC and HPLC using representative herbicides and pesticides as typical moderate polarity and molecular weight range molecules 410(1987)343

Interfacing of inductively coupled plasma-atomic emission spectrometry to SFC for elemental detection 411(1987)213

Supercritical fluid extraction coupled with SFC for the separation of sulphonylurea herbicides and their metabolites from complex matrices 435(1988)63

Supercritical fluid chromatography of ecdysteroids 436(1988)497

Separation of opium alkaloids by carbon dioxide sub- and supercritical fluid chromatography with packed columns. Application to the quantitative analysis of poppy straw extracts 437(1988)351

Analysis of ecdysteroids by supercritical-fluid chromatography 441(1988)165

Use of the photoionisation detector in packed-column supercritical fluid chromatography 445(1988)239

Trace analysis of explosives by capillary supercritical fluid chromatography with thermal energy analysis detection 445(1988)244

Increasing extraction efficiency in supercritical fluid extraction from complex matrices.

Predicting extraction efficiency of diuron and linuron in supercritical fluid extraction using supercritical fluid chromatographic retention 447(1988)53

Supercritical fluid chromatography with ion-pairing modifiers. Separation of enantiomeric 1,2-aminoalcohols as diastereomeric ion pairs 447(1988)287

Effect of the pore structure of silica gel on retention behaviour in pressure programming supercritical fluid chromatography 448(1988)87

### **Supercritical fluid extraction**

Determination of anthraquinone in paper and wood using supercritical fluid extraction and HPLC with electrochemical detection 409(1987)343

Supercritical fluid extraction coupled with SFC for the separation of sulfonyleurea herbicides and their metabolites from complex matrices 435(1988)63

Simple microscale supercritical fluid extraction system and its application to GC-MS of lemon peel oil 442(1988)121

Increasing extraction efficiency in supercritical fluid extraction from complex matrices.

Predicting extraction efficiency of diuron and linuron in supercritical fluid extraction using supercritical fluid chromatographic retention 447(1988)53

### **"Superdynamic" coating**

"Superdynamic" coating of the inner surface of a capillary column with a liquid stationary phase 440(1988)323

### **Superoxide dismutase**

Chromatographic isolation of primary and secondary metabolites from plant cell cultures 440(1988)53

### **Supports**

Potential solid supports for GC prepared on the basis of certain rocks 406(1987)145

Comparison of mesoporous controlled-porosity glasses to silica gel as solid supports for  $C_{18}$  chemically bonded phases 446(1988)347

### **Suppression efficiency**

Effect of suppression efficiency on sensitivity in ion chromatography 439(1988)151

### **Surface excess isotherms**

Adsorption of binary solvent mixtures in reversed-phase chromatographic systems. Surface excess isotherms and enthalpies of displacement 406(1987)257

### **Surface forces**

Near-wall field-flow fractionation. Application of surface forces and transversal voltage to fractionation in laminar flow 446(1988)131

### **Surface irregularity**

Accessibility of irregular (fractal) surfaces of porous silicas to silylation reactions 406(1987)317

### **Surfactant**

Use of term "detergent" in biochemistry 449(1988)

### **Surfactant micelle**

Use of term "detergent" in biochemistry 449(1988)

### **Surfactants**

Analysis of amphoteric surfactants of the alkylaminopropylglycine type by GC 405(1987)305

### **Sweeteners**

LC determination of beta-aspartame in diet soft drinks, beverage powders and pudding mixes 404(1987)261

### **Switching valves**

Stepwise gradient elution using switching valves in micro HPLC 405(1987)117

Precolumn concentration and/or stepwise-gradient elution by switching valve techniques in micro HPLC 407(1987)141

### **Synchronous luminescence**

Direct synchronous luminescence detection of co-eluting solutes in pseudophase LC 409(1987)147

### **Synthetic organic chemicals**

Specific identification of synthetic organic chemicals in river water using liquid-liquid extraction and resin adsorption coupled with

electron impact, chemical ionization and accurate mass measurement GC-MS analyses 436(1988)229

### System peaks

System peaks in ion chromatography using cetrimide-coated columns and indirect photometric detection 402(1987)211

Origin of system peaks in single-column ion chromatography of inorganic anions using high pH borate-gluconate buffers and conductivity detection 404(1987)67

Effects of system peaks in a column-switching LC system for bioanalysis 407(1987)49

## T

### Tacrine hydrochloride

Determination of tacrine hydrochloride in human serum by chloroform extraction, reversed-phase HPLC and fluorimetric detection 433(1988)352

### Taft electronic constant

Factor analysis of trends in selectivity in GLC of 175 saturated and unsaturated esters on 14 polysiloxane stationary phases 404(1987)11

### Taft steric constant

Factor analysis of trends in selectivity in GLC of 175 saturated and unsaturated esters on 14 polysiloxane stationary phases 404(1987)11

### Tamoxifen

Quantification of tamoxifen and N-desmethyltamoxifen in human plasma by HPLC, photochemical reaction and fluorescence detection, and its application to biopharmaceutical investigations 413(1987)207

Separation of tamoxifen geometric isomers and metabolites by bonded-phase  $\beta$ -cyclodextrin chromatography 414(1987)192

### Tannins

Chromatography of tannins. III. Multiple peaks in HPLC of some hydrolyzable tannins 435(1988)285

### Tartaric acid

Complementary use of reversed-phase, normal-phase and chiral HPLC for the study of the stereoisomers produced in decarboxylative alkylation of tartaric acid 440(1988)183

### Tartrate

Elution characteristics of polyol-borate eluents in anion-exchange chromatography. II. Tartrate-borate eluent for non-suppressed anion chromatography 403(1987)35

### Tartrazine

Simple method for the analysis of food dyes on reversed-phase thin-layer plates 411(1987)437

### Taurine

Assessment of taurine in plasma and urine by anion-exchange HPLC with pre-column derivatization 423(1987)292

HPLC analysis of neurotransmitter amino acids in brain 428(1988)9

LC determination of taurine in whole blood, plasma and platelets 431(1988)400

### Tauromustine

Determination of tauromustine, a nitrosourea-based antitumour agent, in plasma by HPLC 425(1988)424

### Tautomers

Kinetic studies of fast equilibrium by means of HPLC. XVI. Separation of tautomers of  $\beta$ -diketones and  $\beta$ -diketoamides 411(1987)466

### Teaching, chromatography

Simple instrument for teaching the basics of chromatography in schools 446(1988)173

### Teicoplanin

Determination of teicoplanin in human plasma and urine by affinity and reversed-phase HPLC 421(1987)99

### Telenzepine

Determination of telenzepine in human serum by GC-MS 430(1988)43

### Temazepam

Microsample determination of diazepam and its three metabolites in serum by reversed-phase HPLC 416(1987)212

Simultaneous determination of diazepam and its active metabolites in rat serum, brain and cerebrospinal fluid by HPLC 422(1987)334

Use of micellar mobile phases and microbore column switching for the assay of drugs in physiological fluids 427(1988)277

Computer-aided optimisation of drug enantiomer separation in chiral HPLC 434(1988)377

#### **Temperature**

Prediction of the distribution coefficient ( $K_d$ ) variation with operating conditions in chromatographic systems 411(1987)81

Effect of temperature on the GC separation of halogenated compounds on polar and non-polar stationary phases 437(1988)33

#### **Temperature effects**

Temperature effect in packed microcolumn chromatography using a carbon dioxide mobile phase 402(1987)328

HPLC at elevated temperatures: examination of conditions for the rapid separation of large molecules 435(1988)1

#### **Temperature gradient**

Single and multiple gradients in supercritical fluid chromatography (Review) 402(1987)1

#### **Temperature-programmed gas chromatography**

Calculation of programmed temperature GC characteristics from isothermal data. I. Theory and computational procedures 405(1987)67

#### **Temperature-programmed retention index**

Use of Kováts retention indices for characterizing solutes in complex samples, separated by linear temperature-programmed capillary GLC 406(1987)117

#### **Temperature-programmed vaporizer**

Comparison of the performances of hot and cold sample introduction with a programmed-temperature vaporizer in the split and splitless modes 438(1988)243

#### **Temperature programming**

Use of Kováts indices for prediction of retention temperatures in linear temperature-programmed capillary GC 407(1987)65

Reproducibility of temperature-programmed retention indices on several OV-101 columns 407(1987)79

#### **Tenax**

Sorption of benzene on Tenax 447(1988)170

#### **Tenax GC**

High-volume elution chromatography of dichlorobenzenes on a polyurethane foam-Tenax sandwich cartridge 409(1987)235

Efficiency of adsorption-desorption of volatile compounds. Application to isoprene on Tenax GC 437(1988)139

#### **Tenuazonic acid**

Standardized HPLC of 182 mycotoxins and other fungal metabolites based on alkylphenone retention indices and UV-VIS spectra (diode array detection) 404(1987)195

#### **Terbium**

Lanthanide ions as luminescent chromophores for the LC detection of polynucleotides and nucleic acids 436(1988)299

Dynamic quenching of europium(III) and terbium(III) luminescence: a potential detection method in ion chromatography 439(1988)165

#### **Terbutaline**

Determination of drug enantiomers in biological samples by coupled column LC and LC-MS 424(1988)61

#### **Terephthalamidine**

HPLC determination of terephthalamidine in plasma 425(1988)220

#### **Terpenes**

Headspace gas analysis. Liquid desorption of headspace volatiles trapped on activated carbon open tubular traps 402(1987)95

Standardized HPLC of 182 mycotoxins and other fungal metabolites based on alkylphenone retention indices and UV-VIS spectra (diode array detection) 404(1987)195

#### **Tertatolol**

Simultaneous quantitative determination of tertatolol and its hydroxylated metabolite in human plasma and urine by GC-MS 421(1987)360

#### **Testosterone**

Combined use of rapid-flow fractionation and HPLC for the determination of serum testosterone and application to study of stress response to physical exercise 423(1987)285

Automated direct assay system for the measurement of sex steroid hormones in serum using HPLC 426(1988)33

**Tetrabenzoperylene**

Effect of column temperature on the retention of peropyrene-type polycyclic aromatic hydrocarbons on various chemically bonded stationary phases in reversed-phase LC 436(1988)1

**Tetra-*n*-butylphosphonium salts**

Solute-solvent interactions in tetra-*n*-butylphosphonium salts studied by GC 438(1988)1

**Tetrachlorobenzo-*p*-dioxins**

Reversed-phase LC of the twenty-two tetrachlorodibenzo-*p*-dioxin isomers on pyrenylethyl- and octadecylsilylated silica gel columns 445(1988)145

**Tetrachlorobenzyltoluenes**

Determination of the polychlorinated biphenyl substitute Ugilec (tetrachlorobenzyltoluenes) in fish 405(1987)311

**2,3,7,8-Tetrachlorodibenzo-*p*-dioxin**

Analysis of pentachlorophenol in water and urine by enrichment with Lipidex 5000 414(1987)55

**Tetrachlorophthalimidopropyl-bonded silica**

Main parameters governing the retention of polynuclear aromatic hydrocarbons on tetrachlorophthalimidopropyl-bonded silica 439(1988)195

**Tetracyclines**

Quantitative analysis of oxytetracycline and related substances by HPLC 405(1987)229

Improvement of chemical analysis of antibiotics. XIII. Systematic simultaneous analysis of residual tetracyclines in animal tissues using TLC and HPLC 411(1987)313

LC and flow injection analysis of tetracycline using sensitized europium(III) luminescence detection 433(1988)149

**2-Tetradecylglycidic acid**

Separation of fatty acid ester and amide enantiomers by HPLC on chiral stationary phases 437(1988)109

 **$\beta$ , $\beta$ , $\gamma$ , $\gamma$ -Tetrafluorochlorambucil**

HPLC of chlorambucil analogues 422(1987)315

**Tetrafluoroethylene**

Investigation by pyrolysis-GC of the composition of multicomponent polymeric

microheterogeneous systems based on some vinyl monomers 404(1987)183

**Tetrahydrobiopterin**

Estimation of tetrahydrobiopterin and other pterins in plasma by isocratic LC with electrochemical and fluorimetric detection 432(1988)321

 **$\Delta^9$ -Tetrahydrocannabinol**

Determination of  $\Delta^9$ -tetrahydrocannabinol in human blood and saliva by HPLC with amperometric detection 421(1987)91

Confirmation of cannabis use. III. Simultaneous quantitation of six metabolites of  $\Delta^9$ -tetrahydrocannabinol in plasma by HPLC with electrochemical detection 434(1988)247

 **$\Delta^9$ -Tetrahydrocannabinol-11-oic acid**

Direct injection of urine on an HPLC column-switching system for determination of  $\Delta^9$ -tetrahydrocannabinol-11-oic acid with both UV and electrochemical detection 417(1987)309

**Tetrahydrofuran**

Derivatization of some polymers with *o*-phthalaldehyde prior to high-performance SEC 436(1988)39

The rôle of the stationary phase in micellar liquid chromatography. Adsorption and efficiency 449(1988)349

**1-(Tetrahydro-2-furyl)-5-fluorouracil**

4-Bromomethyl-6,7-dimethoxycoumarin as a fluorescence reagent for precolumn derivatization of 5-fluorouracil compounds in HPLC 430(1988)156

**(*E*)-4-[2-(5,6,7,8-Tetrahydro-5,5,8,8-tetramethyl-****2-naphthalenyl)-2-methylethenyl]phenol**

Determination of (*E*)-1,2,3,4-tetrahydro-1,1,4,4-tetramethyl-6-(1-methyl-2-phenylethenyl)-naphthalene, an antiacne agent, and its phenolic metabolite in plasma by reversed-phase HPLC 419(1987)253

**(*E*)-1,2,3,4-Tetrahydro-1,1,4,4-tetramethyl-6-(1-methyl-2-phenylethenyl)naphthalene**

Determination of (*E*)-1,2,3,4-tetrahydro-1,1,4,4-tetramethyl-6-(1-methyl-2-phenylethenyl)-naphthalene, an antiacne agent, and its phenolic metabolite in plasma by reversed-phase HPLC 419(1987)253

**Tetrametaphosphoric acid**

Complex-forming equilibria in isotachopheresis. VII. Evaluation of the ion-pair formation constants of phosphorus oxo acids with histidine and assessment of their separability 410(1987)279

**Tetra-3N-methylpyridyl porphyrin**

Separation of metalloporphyrins from metallation reactions by LC and electrophoresis 416(1987)71

**Tetra-4N-methylpyridyl porphyrin**

Separation of metalloporphyrins from metallation reactions by LC and electrophoresis 416(1987)71

**Tetra-N,N,N-trimethylanilinium porphyrin**

Separation of metalloporphyrins from metallation reactions by LC and electrophoresis 416(1987)71

**Tetraprenylacetone**

Determination of tetraprenylacetone in human plasma by HPLC with fluorescence derivatization using dansylhydrazine 424(1988)410

**Tetrodotoxin**

TLC with flame ionization detection for the determination of tetrodotoxin in biological fluids 432(1988)401

**Thalidomide**

Determination of thalidomide and its major metabolites by HPLC 413(1987)181

**Thebaine**

Separation of eight alkaloids and meconic acid and quantitation of five principal alkaloids in gum opium by gradient reversed-phase HPLC 436(1988)455

Separation of opium alkaloids by carbon dioxide sub- and supercritical fluid chromatography with packed columns. Application to the quantitative analysis of poppy straw extracts 437(1988)351

**Theobromine**

HPLC of xanthines. Effects of N-methyl and C-8 hydroxyl substitution on the retention times 428(1988)311

Radiometric HPLC procedure for the determination of theobromine metabolites in microsomal incubations 430(1988)203

**Theophylline**

Analytical methodology to determine stable isotopically labelled and unlabelled theophylline

in human plasma using capillary GC-MS 416(1987)81

Microassay for the simultaneous determination of theophylline and dyphylline in serum by HPLC 416(1987)91

Ion-pair HPLC procedure for the quantitative analysis of theophylline in serum samples 417(1987)99

Determination of theophylline and its metabolites in plasma and urine by reversed-phase LC using an amine modifier 419(1987)177

Improved method for the determination of theophylline in plasma and urine using HPLC 422(1987)281

HPLC assay for theophylline and its major metabolites in human urine 423(1987)179

Automated determination of drugs in serum by column-switching HPLC. II. Separation of theophylline and its metabolites 425(1988)323

HPLC of xanthines. Effects of N-methyl and C-8 hydroxyl substitution on the retention times 428(1988)311

Extraction of acetaminophen and theophylline from post-mortem tissues and urine for HPLC analysis 439(1988)466

Determination of alcohols in alcoholic beverages by micro HPLC with indirect photometric detection 445(1988)139

**Thermal desorber**

Modified method for analysis of C<sub>2</sub>-C<sub>5</sub> hydrocarbons in an aromatic-alkane matrix using an automated thermal desorber 449(1988)175

**Thermal energy analyzer**

Improved method for the trace analysis of explosives by silica capillary column GC with thermal energy analysis detection 410(1987)181

Trace analysis of explosives by capillary supercritical fluid chromatography with thermal energy analysis detection 445(1988)244

**Thermionic ionization detector**

MS measurements of the negative ions formed in the thermionic ionization detector with nitrogen carrier gas 409(1987)139

**Thermodynamics**

Thermochemical investigations of GLC. IV. Partition coefficients of solutes on binary



mixtures containing two complexing solvents  
402(1987)41

Prediction of thermodynamic retention behaviour  
in reversed-phase LC with aqueous eluents and  
various concentrations of methanol by using the  
solubility concept model 408(1987)21

#### **Thermodynamics of solution**

Retention and thermodynamics of solution of  
*n*-alkenes in OV-101 406(1987)75

#### **Thermo-optical detector**

Use of laser detectors in capillary LC (Review)  
434(1988)337

#### **Thiamine**

HPLC–diode-array determination of ascorbic  
acid, thiamine and riboflavin in goats' milk  
410(1987)201

Improved HPLC determination of thiamine and  
its phosphate esters in animal tissues  
450(1988)317

#### **Thianes**

Identification of homologous series of alkylated  
thiophenes, thiolanes, thianes and  
benzothiophenes present in pyrolysates of  
sulphur-rich kerogens 435(1988)435

#### **3-Thienylcyclohexylglycolic acid**

Chiral resolution of a series of 3-  
thienylcyclohexylglycolic acids by liquid or  
subcritical fluid chromatography. A mechanistic  
study 450(1988)255

#### **Thin-layer electrochemical detector**

Thin-layer electrochemical detector coated with  
Nafion film for LC 410(1987)111

#### **Thin-layer plates**

Comparison of non-hydrophobic C<sub>18</sub> bonded  
TLC and HPTLC plates for reversed-phase TLC  
of ecdysteroids 438(1988)419

#### **Thio alcohols**

Polarity of oligoxyethylene derivatives of  
alcohols, thio alcohols and alkylamines as  
measured by GC 450(1988)291

#### **Thiocyanate**

Derivatization of inorganic anions with  
pentafluorobenzyl methanesulphonate for GC  
404(1987)175

Determination of oxidizable inorganic anions by  
HPLC with fluorescence detection and

application to the determination of salivary  
nitrite and thiocyanate and serum thiocyanate  
424(1988)29

Polymeric pentafluorobenzylating reagent for GC  
of inorganic anions with electron-capture  
detection 438(1988)253

#### **Thiocyanate salts**

Solvent properties of liquid organic salts used as  
mobile phases in microcolumn reversed-phase  
LC 411(1987)61

#### **Thiodiglycol**

Detection of trace levels of thiodiglycol in blood,  
plasma and urine using GC–electron-capture  
negative-ion chemical ionisation MS  
449(1988)261

#### **6-Thioguanine nucleotides**

Assay of 6-thioinosinic acid and 6-thioguanine  
nucleotides, active metabolites of 6-  
mercaptapurine, in human red blood cells  
423(1987)169

#### **6-Thioinosinic acid**

Assay of 6-thioinosinic acid and 6-thioguanine  
nucleotides, active metabolites of 6-  
mercaptapurine, in human red blood cells  
423(1987)169

#### **Thiolanes**

Identification of homologous series of alkylated  
thiophenes, thiolanes, thianes and  
benzothiophenes present in pyrolysates of  
sulphur-rich kerogens 435(1988)435

#### **Thiols**

HPLC fluorometry for the determination of thiols  
in biological samples using N-[4-(6-  
dimethylamino-2-benzofuranyl)phenyl]maleimide  
414(1987)11

Sensitive ferrocene reagents for derivatization of  
thiol compounds in HPLC with dual-electrode  
coulometric detection 419(1987)17

Determination of natural thiols by LC after  
derivatization with 3,5-di-*tert*.-butyl-1,2-  
benzoquinone 420(1987)404

Measurement of plasma thiols after derivatization  
with monobromobimane 424(1988)141

Oxidation of thiols within GC columns  
441(1988)372

#### **Thiometalates**

Retention of thio- and selenometalates in  
mobile-phase ion chromatography 439(1988)91

**3-Thiomethylparacetamol**

Complete separation of urinary metabolites of paracetamol and substituted paracetamols by reversed-phase ion-pair HPLC 426(1988)315

**3-Thiomethylparacetamol sulphoxide**

Complete separation of urinary metabolites of paracetamol and substituted paracetamols by reversed-phase ion-pair HPLC 426(1988)315

**1,4-Thiomorpholine-3,5-dicarboxylic acid**

Specific method for the determination of 1,4-thiomorpholine-3,5-dicarboxylic acid and its unsaturated analogue lanthionine ketimine 426(1988)183

**Thiopental**

Determination of sodium pentobarbital and either sodium methohexital or sodium thiopental in plasma by HPLC with ultraviolet detection 414(1986)484

**Thiophenes**

Identification of homologous series of alkylated thiophenes, thiolanes, thianes and benzothiophenes present in pyrolysates of sulphur-rich kerogens 435(1988)435

**Thiopronine**

HPLC of 2-mercaptopropionylglycine and its metabolite 2-mercaptopropionic acid in plasma and urine after treatment with thiopronine 417(1987)261

**Thioridazine**

HPLC resolution of the enantiomers of thioridazine, its metabolites and related compounds 426(1988)223

Indirect HPLC resolution of racemic tertiary amines as their diastereomeric urea derivatives after N-dealkylation 436(1988)269

**Thiosulphate**

Determination of oxidizable inorganic anions by HPLC with fluorescence detection and application to the determination of salivary nitrite and thiocyanate and serum thiocyanate 424(1988)29

**2-Thiouracil**

Identification of thyrostatic drug residues in animal thyroids by HPTLC and fluorescence reaction detection 442(1988)459

**6-Thiouric acid**

Sensitive and specific HPLC assay for 6-thiouric acid 417(1987)183

**Three-electrode ionization chamber**

Mechanism governing the sensitivity of a three-electrode ionization chamber 446(1988)163

**Thrombin**

Affinity chromatography of human thrombin on modified silica 424(1988)385

**Thromboxane**

Separation of prostaglandins and thromboxane by two-dimensional TLC 417(1987)414

Determination of prostaglandins and thromboxane in whole blood by HPLC with fluorimetric detection 427(1988)209

**Thromboxane A<sub>2</sub>**

Purification and quantitative analysis of urinary prostanoids in human and in rat by packed and capillary GC-negative-ion chemical-ionization MS 417(1987)1

**Thromboxane A<sub>2</sub> receptor blocker**

Determination of a new thromboxane A<sub>2</sub> receptor blocker in biological fluids by capillary GC with electron-capture detection 419(1987)345

**Thromboxane B<sub>2</sub>**

Separation of thromboxane B<sub>2</sub> by HPLC 422(1987)246

Determination of prostaglandins D<sub>2</sub>, E<sub>2</sub>, F<sub>2α</sub> and 6-ketoprostaglandin F<sub>1α</sub> as well as thromboxane B<sub>2</sub> in rat brain by GC-MS 426(1988)345

**Thromboxane synthetase inhibitor**

HPLC with fluorescence detection of dazmegrel, a specific thromboxane A<sub>2</sub> synthetase inhibitor 432(1988)375

**α-Thujene**

Resolution of monoterpene enantiomers by GC 407(1987)243

**Thymidine**

Technique for detection of DNA nucleobases by reversed-phase HPLC optimized for quantitative determination of thymidine substitution by iododeoxyuridine 417(1987)57

**Thymine**

Isocratic HPLC method for studying the metabolism of blood plasma pyrimidine nucleosides and bases: concentration and radioactivity measurements 434(1988)111

**Thymoxamine**

HPLC assay of the metabolites of thymoxamine 424(1988)205

**Thyroglobulin**

Investigations into lactoperoxidase-catalysed bromination of tyrosine and thyroglobulin 432(1988)57

**Thyroid hormones**

Resolution of the enantiomers of thyroid hormones by high-performance ligand-exchange chromatography using a chemically bonded chiral stationary phase 404(1987)391

**Thyroxine**

Resolution of the enantiomers of thyroid hormones by high-performance ligand-exchange chromatography using a chemically bonded chiral stationary phase 404(1987)391

**Tiaprofenic acid**

Determination of 2-(5-benzoyl-2-thienyl)propionic acid (tiaprofenic acid) in the plasma of elderly patients with multiple diseases using HPLC 421(1987)206

Rapid and sensitive HPLC assay of tiaprofenic acid enantiomers in human plasma and urine 425(1988)135

**Tiglates**

Considerations and remarks on the analysis of *Anthemis nobilis* L. essential oil by capillary GC and "hyphenated" techniques 411(1987)237

**Timelotem dihydrogenmaleate**

Determination of timelotem dihydrogenmaleate and its main metabolite N-desmethyltimelotem in plasma by capillary GC 419(1987)357

**Timolol**

HPLC of timolol and potential degradates on dynamically modified silica 438(1988)85

**Tinuvin 622**

Determination of the polymeric light stabilizer Tinuvin 622 in polyolefins 450(1988)426

**Tioconazole**

Analysis of tioconazole using HPLC with a porous graphitic carbon column 449(1988)317

**Titration curves**

Electrophoretic titration curve in 6 M urea of the bovine eye lens protein  $\alpha$ -crystallin 411(1987)507

**Tobacco alkaloids**

Enantiomeric resolution of racemic nicotine and nicotine analogues by microcolumn LC with  $\beta$ -cyclodextrin inclusion complexes 411(1987)490

**Tocainide**

Enantiomer separation of tocinide and some of its analogues and derivatives on a Chirasil-Val capillary column 411(1987)285

GC analysis of underivatized tocinide 419(1987)339

 **$\alpha$ -Tocopherol**

Determination of tocopherol and tocopherol acetate concentrations in human feces using HPLC 413(1987)79

Determination of  $\alpha$ -tocopherol and  $\alpha$ -tocopherylquinone in small biological samples by HPLC with electrochemical detection 414(1987)440

Determination of coenzyme Q<sub>10</sub>,  $\alpha$ -tocopherol and cholesterol in biological samples by coupled-column LC with coulometric and UV detection 425(1988)87

Two-dimensional HPLC system for the direct determination of tocopherol derivatives in plasma 427(1988)29

Gas chromatography of retinol and  $\alpha$ -tocopherol without derivatization 434(1988)21

**Tocopherol acetate**

Determination of tocopherol and tocopherol acetate concentrations in human feces using HPLC 413(1987)79

**Tocopherols**

Simultaneous measurement of tocopherols and tocopheryl quinones in tissue fractions using HPLC with redox-cycling electrochemical detection 421(1987)71

Separation of tocopherols on various chemically bonded phases in microcolumn LC 448(1988)398

Isolation of tocopherol homologues by preparative HPLC 450(1988)111

 **$\alpha$ -Tocopheryl quinone**

Determination of  $\alpha$ -tocopherol and  $\alpha$ -tocopherylquinone in small biological samples by HPLC with electrochemical detection 414(1987)440

**Tocopheryl quinones**

Simultaneous measurement of tocopherols and

tocopheryl quinones in tissue fractions using HPLC with redox-cycling electrochemical detection 421(1987)71

#### **Tolmetin**

HPLC determination of tolmetin, tolmetin flucuronide and its isomeric conjugates in plasma and urine 420(1987)349

Statistical optimization of a reversed-phase ion-pair LC method for the analysis of tolmetin sodium in dosage forms 448(1988)1

#### **Tolmetin glucuronide**

HPLC determination of tolmetin, tolmetin flucuronide and its isomeric conjugates in plasma and urine 420(1987)349

#### **Tolperisone**

Application of an ovomucoid-conjugated column for the optical resolution of some pharmaceutically important compounds 408(1987)316

#### **Toluene**

Static headspace GC method for the determination of residual solvents in vigabatrin drug substance 438(1988)103

#### **Toluenediisocyanates**

Trace analysis of airborne aromatic isocyanates and related aminoisocyanates and diamines using HPLC with UV and electrochemical detection 435(1988)469

#### **o-Toluidine**

Analysis of lung air from patients with bronchogenic carcinoma and controls using GC-MS 432(1988)1

Comparative study of alkyl and fluoroalkyl N-(2-hydroxy-ethoxyethyl)-amides as reversed-phase LC modifiers 448(1988)365

#### **Topiramate**

Automated capillary GC assay using flame ionization detection for the determination of topiramate in plasma 433(1988)276

#### **Topoisomerases**

Non-histone chromatin proteins that recognize specific sequences of DNA (Review) 418(1987)73

#### **Torularhodin**

HPLC analysis of major carotenoids from *Rhodotorula glutinis* 448(1988)445

#### **Torulene**

HPLC analysis of major carotenoids from *Rhodotorula glutinis* 448(1988)445

#### **Total molecular surface area**

Total molecular surface areas as a predictor for reversed-phase HPLC in various organotin systems. Importance and applications of carbon hybridization summaries 441(1988)229

#### **Total surface area**

Holistic molecular conformation and total surface area calculations as predictors for solution properties and chromatographic parameters 403(1987)1

#### **Toxic substances**

Automated identification of toxic substances in poisoned human fluids by a retention prediction system in reversed-phase LC 436(1988)11

#### **Toxins**

LC determination of the cyanoginosins, toxins produced by the cyanobacterium *Microcystis aeruginosa* 408(1987)435

Ion-pair HPLC profiling of a uremic toxin fraction 422(1987)263

HPLC method for the determination of  $\alpha$ -amanitin and phalloidin in human plasma using the column-switching technique and its application in suspected cases of poisoning by the green species of amanita mushroom (*Amanita phalloides*) 425(1988)121

Fast protein purification of *Clostridium difficile* cytotoxin 428(1988)408

Rapid analysis of peptide toxins in cyanobacteria 438(1988)93

Analysis and purification of toxic peptides from cyanobacteria by reversed-phase HPLC 448(1988)275

The peptide toxin of the cyanobacterium *Microcystis aeruginosa* PCC 7941. Isolation and analysis by NMR and fast atom bombardment MS 449(1988)423

#### **Trace analysis**

Sample reconcentration by column-external solvent evaporation or injection of large volumes into GC capillary columns? 404(1987)297

#### **Trace enrichment**

Evaluation of disposable cartridges for trace enrichment from aqueous solutions 402(1987)201

Trace enrichment and separation of As<sup>III</sup>, Sb<sup>III</sup> and Bi<sup>III</sup> as diethyldithiophosphate complexes by reversed-phase HPLC 439(1988)63

Effects of drug-protein binding on trace enrichment of drugs in blood plasma on short precolumns 439(1988)353

A cryotrap technique for the quantitation of monoterpenes in humid and ozone-rich forest air 442(1988)157

On-line trace enrichment on a reversed-phase pre-column for normal-phase LC with electron-capture detection 445(1988)129

#### **Trace metals**

Avoidance of trace metal activity with micro-LC in the analysis of hop and beer bitter acids 407(1987)159

#### **Tranexamic acid**

Determination of tranexamic acid in human serum by HPLC using selective pre-column derivatization with phenyl isothiocyanate 433(1988)225

#### **Transcobalamin II**

Separation of transcobalamin II isoproteins by means of chromatofocusing 432(1988)103

#### **Transcription factor TFIIIA**

Non-histone chromatin proteins that recognize specific sequences of DNA (Review) 418(1987)73

#### **Transducin**

Resolution of transducin subunits by chromatography on Blue Sepharose 407(1987)281

#### **Transferrin**

Assay of glycosidase by lectin affinity HPLC 407(1987)299

#### **Transplutonium elements**

Separation of some rare earths and transplutonium elements by HPLC on Ostion cation exchanger 435(1988)249

#### **Transversal voltage**

Near-wall field-flow fractionation. Application of surface forces and transversal voltage to fractionation in laminar flow 446(1988)131

#### **Tranlylcyclopramide**

S-(+)-Flunoxaprofen chloride as chiral fluorescent reagent 427(1988)131

#### **Trehalose dimycolate**

Biomedical applications of chromatographic fraction containing trehalose dimycolate in squalane emulsion (Review) 440(1988)473

#### **Trehalulose**

Preparative HPLC purification and structural determination of 1-O- $\alpha$ -D-glucopyranosyl-D-fructose (trehalulose) 402(1987)265

#### **Trenbolone**

Identification and quantitation of trenbolone in bovine tissue by GC-MS 424(1988)219

Assay for trenbolone and its metabolite 17 $\alpha$ -Trenbolone in bovine urine based on immunoaffinity chromatographic clean-up and off-line HPLC-TLC 445(1988)385

#### **Tretinoin**

Quantitative analysis of retinoids in biological fluids by HPLC using column switching. I. Determination of isotretinoin and tretinoin and their 4-oxo metabolites in plasma 424(1988)303

#### **Triacetylcellulose**

Separation of alkyl-substituted indene enantiomers by chromatography on microcrystalline triacetylcellulose 405(1987)155

#### **Triacylglycerols**

HPLC quantitation of triacylglycerols containing fenoprofen from biological samples 422(1987)33

TLC-flame ionization detection calibration with natural and synthetic standards 439(1988)404

Normal-phase HPLC of triacylglycerols 448(1988)135

#### **Triamcinolone**

GC and MS study of betamethasone synthetic corticosteroid metabolism 432(1988)283

#### **Triamterene**

Rapid method for the determination of the diuretic triamterene and its metabolites in plasma and urine HPLC 413(1987)315

#### **Triapenthenol**

Quantitation of triazole and pyrimidine plant growth retardants 438(1988)393

#### **Triazines**

HPLC of *s*-triazines and their degradation products using UV photometric and amperometric detection 442(1988)147

**Triazolam**

HPLC determination of triazolam and its metabolites in human urine 422(1987)197

**1,2,4-Triazole-3-carboxamide**

HPLC determination of ribavirin in serum and urine and of its urinary metabolite 1,2,4-triazole-3-carboxamide 420(1987)189

**Triazoles**

Relationships between retention behaviour and structure of N-substituted 1,2,4-triazoles in TLC on silica gel 404(1987)73

Quantitation of triazole and pyrimidine plant growth retardants 438(1988)393

**Triazolopyridazines**

Lipophilicity measurement of benzodiazepine-receptor ligands by reversed-phase LC. Comparison between HPLC and TLC 447(1988)404

**Tri-*n*-butyl citrate**

Determination of plasticizers in fat by GC-MS 447(1988)230

**Tributylphenols**

Separation of isomeric mono-, di- and tributylphenols by HPLC on octadecylsilica with methanol-water using alkyl nitrile and alkylamine as surface modifiers 442(1988)53

**Tributylphenyl ethylene oxide oligomers**

TLC separation of some tributylphenyl ethylene oxide oligomers according to the length of the ethylene oxide chain 446(1988)17

**Tributylphosphate**

Specific identification of synthetic organic chemicals in river water using liquid-liquid extraction and resin adsorption coupled with electron impact, chemical ionization and accurate mass measurement GC-MS analyses 436(1988)229

**Trichloroacetic acid**

Automatic headspace GC method for the simultaneous determination of trichloroethylene and metabolites in blood and urine 442(1988)317

Determination of trichloroethylene metabolites in rat liver homogenate using headspace GC 442(1988)325

**1,2,4-Trichlorobenzene**

Specific identification of synthetic organic chemicals in river water using liquid-liquid

extraction and resin adsorption coupled with electron impact, chemical ionization and accurate mass measurement GC-MS analyses 436(1988)229

**Trichloroethanol**

Automatic headspace GC method for the simultaneous determination of trichloroethylene and metabolites in blood and urine 442(1988)317

Determination of trichloroethylene metabolites in rat liver homogenate using headspace GC 442(1988)325

**Trichloroethylene**

Automatic headspace GC method for the simultaneous determination of trichloroethylene and metabolites in blood and urine 442(1988)317

Determination of trichloroethylene metabolites in rat liver homogenate using headspace GC 442(1988)325

**2,4,4'-Trichloro-2'-hydroxydiphenyl ether**

Formation of polychlorinated dibenzo-*p*-dioxins upon combustion of commercial textile products containing 2,4,4'-trichloro-2'-hydroxydiphenyl ether (Irgasan DP300) 442(1988)289

**Trichothecenes**

Standardized HPLC of 182 mycotoxins and other fungal metabolites based on alkylphenone retention indices and UV-VIS spectra (diode array detection) 404(1987)195

Decomposition of trifluoroacetyl derivatives of some trichothecenes on fused-silica capillary columns during GC analysis 407(1987)340

HPLC method for determining trichothecene mycotoxins by post-column fluorescence derivatization 410(1987)427

Metabolism of three trichothecene mycotoxins, T-2 toxin, diacetoxyscirpenol and deoxynivalenol, by bovine rumen microorganisms 414(1987)335

Anthracene-9-carbonyl chloride as a fluorescence and UV derivatising reagent for the HPLC analysis of hydroxy compounds 445(1988)393

Simultaneous detection of trichothecenes and rosenonolactone in grape juice and wine by capillary GC 448(1988)424

**Triclopyr**

Detection of trace levels of triclopyr using capillary GC-electron-capture negative-ion chemical ionization MS 438(1988)45

**Triethylenetetramine**

Quantitative determination of free primary amines in polymeric amine hardeners. HPLC separation of ethylenediamine and triethylenetetramine after derivatization with salicylaldehyde–diphenylboron chelate and separation of 4,4'-diaminodiphenylmethane by HPLC 436(1988)447

**Triethylphosphate**

Specific identification of synthetic organic chemicals in river water using liquid–liquid extraction and resin adsorption coupled with electron impact, chemical ionization and accurate mass measurement GC–MS analyses 436(1988)229

**Trifluoroacetate**

Perfluorinated acids as ion-pairing agents in the determination of monoamine transmitters and some prominent metabolites in rat brain by HPLC with amperometric detection 449(1988)191

**Trifluoroacetic anhydride**

Selective detection of sulphoxides and sulphimides by TLC using trifluoroacetic anhydride–sodium iodide as a reagent 447(1988)225

**Triglycerides**

Simple preparation of a Poly-S 179 capillary column for the analysis of polycyclic aromatic hydrocarbons and triglycerides 404(1987)248

Fatty acid profiles of sebaceous triglycerides by capillary GC with mass-selective detection 425(1988)11

Profiling of plasma cholesterol ester and triglyceride fatty acids as their methyl esters by capillary GC, preceded by a rapid aminopropyl-silica column chromatographic separation of lipid classes 434(1988)395

Quantitative analysis of triglycerides by HPLC using non-linear gradient elution and flame ionization detection 435(1988)139

Calibration of TLC with flame ionization detection for the analysis of natural lipid samples 435(1988)343

Characterization of a lipid-rich fraction synthesized by *Streptomyces avermitilis* 441(1988)31

Densitometric identification of triglycerides separated by reversed-phase TLC 446(1988)259

Separation of triglyceride groups by reversed-phase TLC on silanized Kieselguhr 446(1988)283

**Trihalomethanes**

Multiple element emission spectral detection GC profiles of halogenated products from chlorination of humic acid and drinking water 438(1988)35

**Triiodothyronine**

Resolution of the enantiomers of thyroid hormones by high-performance ligand-exchange chromatography using a chemically bonded chiral stationary phase 404(1987)391

**Trimeprazine**

HPLC resolution of the enantiomers of thioridazine, its metabolites and related compounds 426(1988)223

**Trimetaphosphoric acid**

Complex-forming equilibria in isotachopheresis. VII. Evaluation of the ion-pair formation constants of phosphorus oxo acids with histidine and assessment of their separability 410(1987)279

**Trimethadione**

Simultaneous determination of serum trimethadione and its metabolite by GC 413(1987)376

**Trimethoprim**

Concurrent analysis of methotrexate, trimethoprim, sulphamethoxazole and their major metabolites in plasma by HPLC 422(1987)328

HPLC analysis of trimethoprim and sulfamethoxazole in microliter volumes of chinchilla middle ear effusion and serum 433(1988)187

**Trimethylaluminium**

Sampling and injection system for the analysis of metalorganic chemicals by GC–MS. Some initial results 411(1987)445

**Trimethylgallium**

Sampling and injection system for the analysis of metalorganic chemicals by GC–MS. Some initial results 411(1987)445

**Trimethylindolinone**

Specific identification of synthetic organic chemicals in river water using liquid–liquid extraction and resin adsorption coupled with electron impact, chemical ionization and accurate

mass measurement GC-MS analyses  
436(1988)229

#### **Trimethyllead**

Identification of trimethyllead in urine by HPLC with column switching and chemical reaction detection and by LC-MS 439(1988)109

#### **Trimethyllysine**

Analysis of methylated amino acids by HPLC: methylation of myelin basic protein 421(1987)130

Determination of methylated amino acids in human serum by HPLC 440(1988)225

#### **Trimethyllysine hydroxylase**

Assay for trimethyllysine hydroxylase by HPLC 422(1987)253

#### **O,O,S-Trimethyl phosphorodithioate**

HPLC methods for the determination of contaminants in technical phosphomethylan 435(1988)380

#### **Trimethylpsoralen**

Determination of psoralen in serum by reversed-phase HPLC 423(1987)383

#### **Trimethylsilylation**

Trimethylsilylation of the hydrolysed and polycondensed products of methyltriethoxysilane 410(1987)137

#### **N-Trimethylsilylimidazole**

Silylated N,O-ketals from the reaction of ketones with N-trimethylsilylimidazole 419(1987)271

Polar contributions of the stationary phase to the reversed-phase ion-pair HPLC separation of quaternary ammonium drugs 437(1988)83

#### **Trimipramine**

HPLC resolution of the enantiomers of thioridazine, its metabolites and related compounds 426(1988)223

GC-MS study of the urinary metabolism of trimipramine 431(1988)197

Simultaneous determination of trimipramine and its demethylated metabolites in plasma by GC-MS 434(1988)232

#### **Trinactin**

Preparative HPLC of macrotetrolides 442(1988)431

#### **2,4,6-Trinitrotoluene**

Improved method for the trace analysis of

explosives by silica capillary column GC with thermal energy analysis detection 410(1987)181

#### **Trioses**

Degradation of ketoses during aqueous HPLC on lead-form cation-exchange resins 437(1988)387

#### **Tripelennamine**

Applications of column-switching techniques in biopharmaceutical analysis. II. HPLC determination of tripelennamine in bovine plasma and milk 421(1987)216

#### **Triphenyltin compounds**

A biochromatographic technique for the quantitative estimation of triphenyltin fungicides 435(1988)210

#### **Triphosphoric acid**

Complex-forming equilibria in isotachopheresis. VII. Evaluation of the ion-pair formation constants of phosphorus oxo acids with histidine and assessment of their separability 410(1987)279

#### **Tris(acetylacetonato)aluminium(III)**

Elution recovery of tris(acetylacetonato)aluminium(III), -gallium(III) and -indium(III) in SEC 435(1988)368

#### **Tris(acetylacetonato)gallium(III)**

Elution recovery of tris(acetylacetonato)aluminium(III), -gallium(III) and -indium(III) in SEC 435(1988)368

#### **Tris(acetylacetonato)indium(III)**

Elution recovery of tris(acetylacetonato)aluminium(III), -gallium(III) and -indium(III) in SEC 435(1988)368

#### **Tris(chloropropyl)phosphate**

Specific identification of synthetic organic chemicals in river water using liquid-liquid extraction and resin adsorption coupled with electron impact, chemical ionization and accurate mass measurement GC-MS analyses 436(1988)229

#### **Tris( $\beta$ -diketonato)metal complexes**

Adsorption behaviour of tris( $\beta$ -diketonato)metal(III) complexes on silica gel 438(1988)401

#### **Triterpenes**

HPLC of glycosides on a new type of hydroxyapatite column 407(1987)205



factor during hydrophobic interaction chromatography 443(1988)205

### **Tungsten**

Determination of molybdenum in soils and plants by HPLC 411(1987)401

### **Tungsten selenide**

Use of modified tungsten selenide as an adsorbent for GC 446(1988)79

### **Two-dimensional gel electrophoresis**

Two-dimensional gel electrophoresis of proteins (Review) 418(1987)145

### **Tyramine**

Simultaneous determination of plasma phenethylamine, phenylethanolamine, tyramine and octopamine by HPLC using derivatization with fluorescamine 427(1988)320

HPLC of biogenic amines, derivatized with 9-fluorenylmethyl chloroformate 437(1988)379

Detection by HPLC of polyamines formed by clostridal putrefaction of caseins 442(1988)267

### **Tyrosine**

Increased yield of the phenylthiocarbonyl-amino acids tyrosine, cysteine and phenylalanine using diethylether-based liquid-liquid extraction 403(1987)299

Assay for tyrosine hydroxylase by HPLC with fluorescence detection 415(1987)289

Analysis of the isomeric tyrosines in mammalian and avian systems using HPLC with fluorescence detection 426(1988)155

Investigations into lactoperoxidase-catalysed bromination of tyrosine and thyroglobulin 432(1988)57

### **Tyrosine-containing oligopeptides**

HPLC of N-terminal tyrosine-containing oligopeptides by pre-column fluorescence derivatization with hydroxylamine, cobalt(II) and borate reagents 411(1987)305

### **Tyrosine-containing peptides**

On-line post-column fluorescence detection for N-terminal tyrosine-containing peptides in HPLC 421(1987)245

### **Tyrosine hydroxylase**

Assay for tyrosine hydroxylase by HPLC with fluorescence detection 415(1987)289

Assay for tyrosine hydroxylase in hypothalamic

homogenates using HPLC with electrochemical detection 419(1987)27

Simple assay procedure for tyrosine hydroxylase activity by HPLC employing coulometric detection with minimal sample preparation 427(1988)229

### **Tyrosylation**

Tyrosylation and purification of peptides for radioiodination 447(1988)341

## **U**

### **Ubiquinol**

Determination of reduced and total ubiquinones in biological materials by LC with electrochemical detection 430(1988)11

### **Ubiquinones**

Determination of reduced and total ubiquinones in biological materials by LC with electrochemical detection 430(1988)11

### **Ugilec**

Determination of the polychlorinated biphenyl substitute Ugilec (tetrachlorobenzyltoluenes) in fish 405(1987)311

### **Ultra-low-loaded columns**

Evaluation of interfacial surface areas in ultra-low-loaded GC columns 446(1988)221

### **Ultraviolet absorbance-fluorescence detector**

UV spectrometric detection for GC of polynuclear aromatic compounds with repetitive spectral scans using absorbance and concurrent fluorescence measurements 405(1987)191

### **Ultraviolet absorption detector**

Capillary GC analysis with the far-UV absorbance detector 441(1988)63

High-performance UV absorption detector for LC. II. A prototype detector 441(1988)387

### **Ultraviolet spectra, search algorithm**

Identification of drugs in human serum by HPLC with photodiode array detection and a search algorithm for UV spectra 419(1987)135

**Triterpenoids**

Separation of oxygenated triterpenoids from *Ganoderma lucidum* by HPLC 410(1987)195

**Tritium isotope effect**

Tritium isotope effect in reversed-phase HPLC analysis of dopamine and dihydroxyphenylacetic acid 427(1988)9

**Truxillic acid**

Analysis of manufacturing by-products and impurities in illicit cocaine via HPLC and photodiode array detection 405(1987)273

Capillary GC–electron capture detection of coca-leaf-related impurities in illicit cocaine: 2,4-diphenylcyclobutane-1,3-dicarboxylic acids, 1,4-diphenylcyclobutane-2,3-dicarboxylic acids and their alkaloidal precursors, the truxillines 410(1987)297

**Truxilline**

Analysis of manufacturing by-products and impurities in illicit cocaine via HPLC and photodiode array detection 405(1987)273

**Truxinic acid**

Analysis of manufacturing by-products and impurities in illicit cocaine via HPLC and photodiode array detection 405(1987)273

Capillary GC–electron capture detection of coca-leaf-related impurities in illicit cocaine: 2,4-diphenylcyclobutane-1,3-dicarboxylic acids, 1,4-diphenylcyclobutane-2,3-dicarboxylic acids and their alkaloidal precursors, the truxillines 410(1987)297

**Trypsin inhibitor**

Reversed-phase HPLC of soybean trypsin inhibitor with optical activity and UV absorbance detection 405(1987)295

**Tryptamine**

HPLC of biogenic amines, derivatized with 9-fluorenylmethyl chloroformate 437(1988)379

Analysis of tryptamine at the femtomole level in tissue using negative ion chemical ionization GC–MS 440(1988)253

Detection by HPLC of polyamines formed by clostridal putrefaction of caseins 442(1988)267

**Tryptamine derivatives**

Determination of tryptamine derivatives in hallucinogenic mushrooms using HPLC with photodiode array detection 408(1987)402

**Tryptophan**

Chiral separations of amino acids using di- and tri-peptides for zwitter-ion pair chromatography 402(1987)49

High-voltage electrophoretic separation of tryptophan metabolites of the kynurenine pathway 420(1987)263

Determination of plasma 5-hydroxytryptophan, 5-hydroxytryptamine, 5-hydroxyindoleacetic acid, tryptophan and melatonin by HPLC with electrochemical detection 431(1988)170

**Tryptophan oxygenase**

Semi-micro quantitative assay for tryptophan oxygenase by HPLC 419(1987)296

**Tryptophan transaminase**

GC–MS determination of tryptophan transaminase-catalyzed deuterium exchange 415(1987)104

**Tryptophol**

Melatonin and other indoles in rat pineal 428(1988)206

**TSKgel Heparin-5PW**

High-performance affinity chromatography of proteins on TSKgel Heparin-5PW 445(1988)234

**TSKgel 3000 SWXL**

Comparative study of Zorbax Bio Series GF 250 and GF 450 and TSK-GEL 3000 SW and SWXL columns in SEC of proteins 443(1988)45

**TSKgel SW<sub>XL</sub>**

New high-performance gel filtration columns for protein separation 404(1987)333

**Tswett, M.S.**

Tswett's letters to Claparède 440(1988)509

**T-2 toxin**

Metabolism of three trichothecene mycotoxins, T-2 toxin, diacetoxyscirpenol and deoxynivalenol, by bovine rumen microorganisms 414(1987)335

**Tubulin**

Separation of tubulin subunits by reversed-phase HPLC 448(1988)371

**Tulobuterol**

Determination of tulobuterol in human plasma by capillary GC 416(1987)154

**Tumor necrosis factor**

Reversible subunit dissociation of tumor necrosis

**Uniconizole**

Quantitation of triazole and pyrimidine plant growth retardants 438(1988)393

**Uracil**

Isocratic HPLC method for studying the metabolism of blood plasma pyrimidine nucleosides and bases: concentration and radioactivity measurements 434(1988)111

**Uraemic toxins**

Ion-pair HPLC profiling of a uremic toxin fraction 422(1987)263

**Urea derivatives**

Simultaneous conductivity and amperometric detector with a single cell for microbore LC 436(1988)413

**Urea-formaldehyde**

Cleavage of urea-formaldehyde resin in cross-linked alkyd systems 403(1987)145

**Uric acids**

Determination of theophylline and its metabolites in plasma and urine by reversed-phase LC using an amine modifier 419(1987)177

**Uridine**

Hypoxanthine, xanthine and uridine in body fluids, indicators of ATP depletion (Review) 429(1988)255

Isocratic HPLC method for studying the metabolism of blood plasma pyrimidine nucleosides and bases: concentration and radioactivity measurements 434(1988)111

**Uridine 5'-diphosphate glucose-collagen glucosyltransferase**

Detection of uridine 5'-diphosphate glucose-collagen glucosyltransferase activity by HPLC and possible participation of a novel intermediate as a glucose donor in collagen biosynthesis 417(1987)403

**Uridine diphosphate glucuronyltransferases**

Assay method for product formation in *in vitro* enzyme kinetic studies of uridine diphosphate glucuronyltransferases: 2-arylpropionic acid enantiomers 430(1988)368

**Urinary acids**

Excretion of urinary acids during inverted sleep-waking rhythm 422(1987)1

**Urinary proteins**

Urinary protein analysis (Review) 429(1988)315

**Uronate**

Elution characteristics of polyol-borate eluents in anion-exchange chromatography. I. Eluent strength of sugar-, uronate-borate eluents 403(1987)27

**Uroporphyrins**

Reversed-phase HPLC of urinary porphyrin free acids 423(1987)313

Analytical and preparative separation of uroporphyrin I and III by HPLC 437(1988)311

Detection of non-typical porphyrin isomers in human urines by ion-pair reversed-phase HPLC 441(1988)171

**Ursodeoxycholate**

Analysis of conjugated bile acids in human serum by HPLC using post-column enzyme reaction and off-line fluorimetric determination 423(1987)75

Studies on steroids. CCXXXIII. Separation and characterization of bile acid disulphates in human urine by HPLC 425(1988)59

**Uxepam**

Selective effect of clonazepam and (*S*)-uxepam on the binding of warfarin enantiomers to human serum albumin 450(1988)217

**V****Valproic acid**

Gas chromatographic method for the measurement of sodium valproate utilization by kidney tubules 414(1987)460

Capillary GC assay for valproic acid and its 2-desaturated metabolite in brain and plasma 432(1988)185

**Vancomycin**

Determination of vancomycin related substances by gradient HPLC 410(1987)363

Chromatographic methods for the analysis of vancomycin 410(1987)373

Determination of vancomycin in serum and

tissues by column LC using solid-phase extraction 414(1987)472

HPLC analysis of vancomycin in plasma, bone, atrial appendage tissue and pericardial fluid 417(1987)121

#### **Vanillylmandelic acid**

Simultaneous LC determination of seventeen of the major monoamine neurotransmitters, precursors and metabolites. I. Optimization of the mobile phase using factorial designs and a computer program to predict chromatograms 415(1987)261

Simultaneous LC determination of seventeen of the major monoamine neurotransmitters, precursors and metabolites. II. Assessment of human brain and cerebrospinal fluid concentrations 415(1987)275

LC determination of vanillylmandelic acid and homovanillic acid by a column-switching technique involving direct injection of urine 421(1987)111

Simple HPLC method for the simultaneous measurement of vanillylmandelic acid and homovanillic acid 423(1987)278

Automatic determination of urinary 4-hydroxy-3-methoxymandelic (vanillylmandelic) acid by LC with electrochemical detection 424(1988)255

Simultaneous determination of the major acidic metabolites of catecholamines and serotonin in urine by LC with electrochemical detection after a one-step sample clean-up on Sephadex G-10; influence of vanilla and banana ingestion 424(1988)273

Simultaneous determination of urinary vanillylmandelic acid, 5-hydroxyindoleacetic acid and homovanillic acid by LC 428(1988)147

Determination of vanillylmandelic acid in plasma by HPLC with electrochemical detection 431(1988)150

Method of measurement of urinary homovanillic acid and vanillylmandelic acid by GC-MS suitable for neuroblastoma screening 432(1988)273

Automated analysis of urinary 3-methoxy-4-hydroxy-mandelic acid using ion-pair chromatography and fluorimetric detection 434(1988)43

#### **Vasopressin**

Quantitative LC determination of disulfide-

containing peptide analogues of vasopressin with dual Hg/Au electrochemical detection 413(1987)43

Use of perfluorinated carboxylic acids in the separation of metabolites of vasopressin prior to radioimmunoassay 430(1988)128

#### **Vasopressin antagonists**

HPLC analysis of arginine-containing peptides in biological fluids by means of a selective post-column reaction with fluorescence detection 444(1988)123

#### **Vecuronium**

Assay of vecuronium in plasma using solid-phase extraction, HPLC and post-column ion-pair extraction with fluorimetric detection 421(1987)327

Quantitation of pancuronium, 3-desacetylpancuronium, vecuronium, 3-desacetylvecuronium, pipecuronium and 3-desacetylpipecuronium in biological fluids by capillary GC using nitrogen-sensitive detection 427(1988)41

#### **Velocity gradient**

Single and multiple gradients in supercritical fluid chromatography (Review) 402(1987)1

#### **Verapamil**

GC analysis of verapamil and norverapamil in *post mortem* specimens using a 530- $\mu\text{m}$  fused-silica column with nitrogen-phosphorus detection 402(1987)361

#### **Verminoside**

Application of various planar chromatographic techniques for the separation of iridoid glycosides from *Veronica officinalis* 404(1987)365

#### **Veronicoside**

Application of various planar chromatographic techniques for the separation of iridoid glycosides from *Veronica officinalis* 404(1987)365

#### **Verproside**

Application of various planar chromatographic techniques for the separation of iridoid glycosides from *Veronica officinalis* 404(1987)365

#### **Verruculogen, Rugulosin**

Standardized HPLC of 182 mycotoxins and other fungal metabolites based on alkylphenone retention indices and UV-VIS spectra (diode array detection) 404(1987)195

**Versamids**

Derivatization of some polymers with *o*-phthalaldehyde prior to high-performance SEC 436(1988)39

**Versicolorin A**

HPLC procedure for determining the profiles of aflatoxin precursors in wildtype and mutant strains of *Aspergillus parasiticus* 441(1988)400

**Versicolorin C**

HPLC procedure for determining the profiles of aflatoxin precursors in wildtype and mutant strains of *Aspergillus parasiticus* 441(1988)400

**Versiconal hemiacetal acetate**

HPLC procedure for determining the profiles of aflatoxin precursors in wildtype and mutant strains of *Aspergillus parasiticus* 441(1988)400

**Very-low-density lipoproteins**

Rapid quantification of apolipoprotein E enriched very-low-density lipoproteins by heparin-Sepharose chromatography 425(1988)169

Fractionation of baboon chylomicrons and very-low-density lipoproteins by HPLC 433(1988)257

**Vigabatrin**

Static headspace GC method for the determination of residual solvents in vigabatrin drug substance 438(1988)103

**Vinblastine**

Determination of *Catharanthus* alkaloids by reversed-phase HPLC 410(1987)488

Solid-phase extraction of vinblastine and vincristine from plasma and urine: variable drug recoveries due to non-reproducible column packings 414(1987)91

Determination of vinblastine in MO<sub>4</sub> mouse fibrosarcoma cells by HPLC 416(1987)375

Analysis of vinca alkaloids in plasma and urine using HPLC with electrochemical detection 424(1988)83

Determination of vinca alkaloids in plasma and urine using ion-exchange chromatography on silica gel and fluorescence detection 434(1988)145

**Vinburnine**

Simultaneous determination of vinburnine and 6-hydroxyvinburnine in human plasma by HPLC (in French) 434(1988)157

**Vinca alkaloids**

Analysis of vinca alkaloids in plasma and urine using HPLC with electrochemical detection 424(1988)83

Determination of vinca alkaloids in plasma and urine using ion-exchange chromatography on silica gel and fluorescence detection 434(1988)145

**Vincristine**

Determination of *Catharanthus* alkaloids by reversed-phase HPLC 410(1987)488

Solid-phase extraction of vinblastine and vincristine from plasma and urine: variable drug recoveries due to non-reproducible column packings 414(1987)91

Analysis of vinca alkaloids in plasma and urine using HPLC with electrochemical detection 424(1988)83

**Vindesine**

Analysis of vinca alkaloids in plasma and urine using HPLC with electrochemical detection 424(1988)83

Determination of vinca alkaloids in plasma and urine using ion-exchange chromatography on silica gel and fluorescence detection 434(1988)145

**Vindoline**

Determination of *Catharanthus* alkaloids by reversed-phase HPLC 410(1987)488

**Vinpocetine**

Quantitative determination of vinpocetine in human plasma by capillary GC-MS 413(1987)264

**Vinyl alcohol**

Development of a C<sub>18</sub>-bonded vinyl alcohol copolymer gel for reversed-phase HPLC 410(1987)129

**Vinyl alcohol copolymer**

Separation of peptides by multimode chromatography on a column packed with vinyl alcohol copolymer gel 434(1988)71

**Violaxanthin**

Separation and quantitative analysis of some carotenoid fatty acid esters of fruits by LC 435(1988)113

Separation of carotenol fatty acid esters by HPLC 449(1988)119

**Viomellein**

Standardized HPLC of 182 mycotoxins and other

fungal metabolites based on alkylphenone retention indices and UV-VIS spectra (diode array detection) 404(1987)195

#### **Viridicatumtoxin**

Standardized HPLC of 182 mycotoxins and other fungal metabolites based on alkylphenone retention indices and UV-VIS spectra (diode array detection) 404(1987)195

#### **Viruses**

Agarose gel electrophoresis of bacteriophages and related particles (Review) 418(1987)345

Adsorption and chromatographic properties of modified silica sorbents for the production of viral preparations 446(1988)211

#### **Virus proteins**

High-performance GPC of the surface glycoprotein from bovine leukemia virus 425(1988)379

HPLC separation and immunological characterization of soluble bovine viral diarrhea virus antigen 435(1988)149

Effect of detergents on the structure of integral membrane proteins of Sendai virus studied with size-exclusion HPLC and monoclonal antibodies 443(1988)255

#### **Vitamin A**

Determination of vitamin A derivatives in fortified foods and commercial vitamin supplements by HPLC 409(1987)419

Simultaneous determination of the aromatic retinoids etretin and etretinate and their main metabolites by reversed-phase liquid chromatography 415(1987)413

Reversed-phase HPLC separation and electrochemical detection of retinol and its isomers 417(1987)65

Determination of the aromatic retinoids (etretin and isoetretin) in biological fluids by HPLC 421(1987)177

Chromatography of fat-soluble vitamins in clinical chemistry (Review) 429(1988)3

Gas chromatography of retinol and  $\alpha$ -tocopherol without derivatization 434(1988)21

Rapid, simultaneous determination of isomers of retinal, retinal oxime and retinol by HPLC 438(1988)383

#### **Vitamin B<sub>1</sub>**

HPLC-diode-array determination of ascorbic

acid, thiamine and riboflavine in goats' milk 410(1987)201

HPLC-diode-array determination of ascorbic acid, thiamine and riboflavine in goats' milk 410(1987)201

Analysis of vitamin B<sub>6</sub> vitamers in plasma by cation-exchange HPLC 416(1987)138

Simple isocratic HPLC method for the separation of the six vitamers of vitamin B<sub>6</sub> 424(1988)315

Plasma B<sub>6</sub> vitamer and 4-pyridoxic acid concentrations of men fed controlled diets 428(1988)35

HPLC assay of erythrocyte enzyme activity levels involved in vitamin B<sub>6</sub> metabolism 431(1988)406

Application of droplet counter-current chromatography to the isolation of vitamin B<sub>12</sub> 435(1988)235

#### **Vitamin B<sub>12</sub>**

Improved HPLC determination of thiamine and its phosphate esters in animal tissues 450(1988)317

#### **Vitamin C**

HPLC-diode-array determination of ascorbic acid, thiamine and riboflavine in goats' milk 410(1987)201

HPLC determination of ascorbic acid in human tears 414(1987)434

Ion suppression reversed-phase HPLC method for the separation of L-ascorbic acid in fresh fruit juice 437(1988)447

Interactions between ascorbic acid and linear and cyclized bile pigments. A TLC study 440(1988)275

#### **Vitamin D**

Rapid automated HPLC assay for ercalcidiol and calcidiol (25-hydroxyvitamins D<sub>2</sub> and D<sub>3</sub>) using *trans*-calcidiol as an UV-absorbing internal standard 415(1987)305

Chromatography of fat-soluble vitamins in clinical chemistry (Review) 429(1988)3

Clinical analysis of individual steroids by column LC (Review) 429(1988)155

Rapid determination of vitamin D in fortified skim milk 445(1988)305

#### **Vitamin D-dependent calcium-binding protein**

Chick 28 000 *M<sub>r</sub>*, vitamin D-dependent calcium-

binding protein in intestine, kidney and cerebellum. Purification using chromatofocusing 448(1988)145

### **Vitamin E**

Determination of tocopherol and tocopherol acetate concentrations in human feces using HPLC 413(1987)79

Determination of  $\alpha$ -tocopherol and  $\alpha$ -tocopherylquinone in small biological samples by HPLC with electrochemical detection 414(1987)440

Simultaneous measurement of tocopherols and tocopheryl quinones in tissue fractions using HPLC with redox-cycling electrochemical detection 421(1987)71

Determination of coenzyme Q<sub>10</sub>,  $\alpha$ -tocopherol and cholesterol in biological samples by coupled-column LC with coulometric and UV detection 425(1988)87

Two-dimensional HPLC system for the direct determination of tocopherol derivatives in plasma 427(1988)29

Chromatography of fat-soluble vitamins in clinical chemistry (Review) 429(1988)3

Separation of tocopherols on various chemically bonded phases in microcolumn LC 448(1988)398

### **Vitamin H**

Determination of free biotin in plasma by LC with fluorimetric detection 413(1987)247

### **Vitamin K**

Chromatography of fat-soluble vitamins in clinical chemistry (Review) 429(1988)3

Determination of K vitamins (phylloquinone and menaquinones) in umbilical cord plasma by a platinum-reduction column 430(1988)143

The vitamin K dependent reaction 440(1988)499

### **Vitamin K<sub>1</sub>**

Simultaneous determination of vitamin K<sub>1</sub>, vitamin K<sub>1</sub> 2,3-epoxide and menaquinone-4 in human plasma by HPLC with fluorimetric detection 430(1988)21

### **Vitamin K<sub>1</sub> 2,3-epoxide**

Simultaneous determination of vitamin K<sub>1</sub>, vitamin K<sub>1</sub> 2,3-epoxide and menaquinone-4 in human plasma by HPLC with fluorimetric detection 430(1988)21

### **Vitamins**

Chromatography of fat-soluble vitamins in clinical chemistry (Review) 429(1988)3

Factors affecting the ion-pair chromatography of water-soluble vitamins 442(1988)81

Analysis of water-soluble vitamins by micellar electrokinetic capillary chromatography 447(1988)133

Determination of water-soluble vitamins using HPLC and electrochemical or absorbance detection 447(1988)256

### **Vitellogenin**

Determination of vitellogenin in serum of rainbow trout (*Salmo gairdneri*) by HPLC 427(1988)138

### **Vitride**

Application of methylamine O-deacylation and Vitride reduction to the analysis of neutral lipids: a TLC evaluation 434(1988)215

### **Void zones**

Detection of void zones in packed chromatographic columns 404(1987)33

Determination of the volumes of void zones in packed chromatographic columns 445(1988)37

### **Volatile acids**

see also Essential oils

### **Volatile compounds**

Headspace gas analysis. Liquid desorption of headspace volatiles trapped on activated carbon open tubular traps 402(1987)95

Dynamic headspace GC of volatile compounds in milk 404(1987)163

Dynamic headspace GC: concentration of volatile components after thermal desorption by intermediate cryofocusing in a cold trap. I. Principle and applications 405(1987)87

Dynamic headspace GC: concentration of volatile components after thermal desorption by intermediate cryofocusing in a cold trap. II. Effect of sampling and desorption parameters on recovery 405(1987)99

Comparison of volatile components between fresh and rotten mussels by GC-MS 409(1987)251

Glass capillary GC analysis of volatile components of canned Philippine pineapple juice 409(1987)409

Screening of volatile compounds present in human blood using retention indices in GC 411(1987)251

Analysis of lung air from patients with bronchogenic carcinoma and controls using GC-MS 432(1988)1

Purge-and-trap GC method for residual volatiles in high-temperature polymers 450(1988)394

#### **Volatile oils**

Characterisation of volatile oil constituents with relatively long GC retention times on two stationary phases 449(1988)312

#### **VX**

Analysis of O-ethyl S-[2-(diisopropylamino)ethyl] methyl-phosphonothiolate (VX) by capillary column GC-MS 402(1987)221

## **W**

#### **Wall-coated open-tubular capillary columns**

Preparation of wall-coated open-tubular capillary columns for GC (Review) 445(1988)1

#### **Warfarin**

Determination of association constants of  $\beta$ -cyclodextrin and  $\beta$ -cyclodextrin-bearing polymers with drugs by HPLC 409(1987)61

Sensitive measurements of warfarin by HPLC with electrochemical detection 415(1987)432

Selective effect of clonazepam and (*S*)-uxepam on the binding of warfarin enantiomers to human serum albumin 450(1988)217

#### **Water**

GC determination of water in crude oils 403(1987)288

#### **Water-soluble vitamins**

Factors affecting the ion-pair chromatography of water-soluble vitamins 442(1988)81

Analysis of water-soluble vitamins by micellar electrokinetic capillary chromatography 447(1988)133

Determination of water-soluble vitamins using HPLC and electrochemical or absorbance detection 447(1988)256

#### **Wax esters**

Identification of very-long-chain fatty acids in rat and mouse harderian gland lipids by capillary GC-MS 431(1988)231

#### **Wheat germ agglutinin**

Analytical high-performance affinity chromatography of ovalbumin-derived glycopeptides on columns of concanavalin A- and wheat germ agglutinin-immobilized gels 438(1988)73

#### **Wide-pore packing materials**

Internal structures of wide-pore packing materials for HPLC studied by transmission electron microscopy 448(1988)95

#### **Wine**

Determination of phenol by HPLC with fluorimetric detection. Application to wines (in French) 407(1987)384

#### **Wood extractives**

SEC of deposits in pulp and paper mills 411(1987)363

## **X**

#### **Xamoterol**

Determination of xamoterol in human plasma by HPLC with electrochemical detection 417(1987)233

#### **Xanthene dyes**

Identification of xanthene dyes in lipsticks by reversed-phase HPLC 448(1988)296

#### **Xanthine**

Hypoxanthine, xanthine and uridine in body fluids, indicators of ATP depletion (Review) 429(1988)255

#### **Xanthine oxidase**

Highly sensitive assay for xanthine oxidase



activity by HPLC with fluorescence detection 424(1988)392

### **Xanthines**

Simplified description of HPLC separation under overload conditions, based on the Craig distribution model. V. Gradient elution separation 405(1987)51

Adsorption chromatography on cellulose. I. Salting-out chromatography of organic compounds 406(1987)389

Ion-pair HPLC procedure for the quantitative analysis of theophylline in serum samples 417(1987)99

Determination of theophylline and its metabolites in plasma and urine by reversed-phase LC using an amine modifier 419(1987)177

HPLC of xanthines. Effects of N-methyl and C-8 hydroxyl substitution on the retention times 428(1988)311

### **Xanthophylls**

HPLC of chloroplast pigments. One-step separation of carotene and xanthophyll isomers, chlorophylls and pheophytins 448(1988)411

### **Xanthurenic acid**

High-voltage electrophoretic separation of tryptophan metabolites of the kynurenine pathway 420(1987)263

Quantification of urinary xanthurenic acid excretion by anion-exchange solid-phase extraction and HPLC 425(1988)182

### **Xylazine**

Determination of xylazine in blood components using HPLC 426(1988)207

### **Xylenes**

Heavy alkylated benzene stationary phase for the GLC separation of ethylbenzene, xylenes, styrene, iso- and *n*-propylbenzene 409(1987)371

### **Xylenols**

Separation of *m*- and *p*-cresol and other methylated phenols in shale oil by reversed-phase LC with electrochemical detection 441(1988)347

### **2,6-Xytilidine**

In vivo metabolism of lidocaine in the rat. Isolation of urinary metabolites as pentafluorobenzoyl derivatives and their identification by combined GC-MS 421(1987)267

### **Xylose**

Analysis of neutral sugars as dinitrophenylhydrazones by HPLC 405(1987)221

HPLC separation of some mono- and disaccharides with detection by a post-column enzyme reactor and a chemically modified electrode 408(1987)157

Simultaneous GLC determination of aldoses and alduronic acids 408(1987)245

### **Xylosyl thioether**

HPLC determination of a naturally occurring xylosyl thioether, 9-[5'-deoxy-5'(methylthio)- $\beta$ -D-xylofuranosyl]adenine 440(1988)151

## **Y**

### **Yohimbine**

Isolation of basic drugs from plasma using solid-phase extraction with a cyanopropyl-bonded phase 432(1988)209

## **Z**

### **Zearalenone**

Standardized HPLC of 182 mycotoxins and other fungal metabolites based on alkylphenone retention indices and UV-VIS spectra (diode array detection) 404(1987)195

### **Zeaxanthin**

Separation of carotenol fatty acid esters by HPLC 449(1988)119

### **Zidovudine**

Simple and rapid HPLC assay for zidovudine (azidothymidine) in plasma and urine 430(1988)420

Simultaneous quantification of zidovudine and its glucuronide in serum by HPLC 431(1988)123

#### **Zinc protoporphyrin**

Quantitative determination of porphyrins, their precursors and zinc protoporphyrin in whole blood and dried blood by HPLC with fluorimetric detection 417(1987)269

#### **Zinc pyrithione**

Determination of zinc pyrithione in cosmetic products by HPLC with pre-labelling 408(1987)255

#### **Zindoxifene**

Chromatographic analysis of the new antiestrogen zindoxifene and its metabolites in biological material 414(1987)235

#### **Zirconia**

Effect of zirconia on the sorption properties of silica gels 407(1987)189

#### **Zirconium phosphate**

Automated separation process for radioanalytical purposes at nuclear power plants 406(1987)359

#### **Zofenopril**

Simultaneous determination of the prodrug

zofenopril and its active drug in plasma by capillary GC-MS 428(1988)81

#### **Zone-electrophoretic sample treatment**

Zone-electrophoretic sample treatment for LC 436(1988)421

#### **Zone electrophoresis**

Carrier-free zone electrophoresis, displacement electrophoresis and isoelectric focusing in a high-performance electrophoresis apparatus 403(1987)47

#### **Zopiclone**

Simultaneous determination of zopiclone and its two major metabolites (N-oxide and N-desmethyl) in human biological fluids by reversed-phase HPLC 417(1987)151

#### **Zorbax Bio Series GF**

Comparative study of Zorbax Bio Series GF 250 and GF 450 and TSK-GEL 3000 SW and SWXL columns in SEC of proteins 443(1988)45

#### **Zwitter-ion pair chromatography**

Chiral separations of amino acids using di- and tri-peptides for zwitter-ion pair chromatography 402(1987)49

## PUBLICATION SCHEDULE FOR 1989

*Journal of Chromatography and Journal of Chromatography, Biomedical Applications*

MONTH	J	F	M	
Journal of Chromatography	461 462 463/1	463/2 464/1	The publication schedule for further issues will be published later	
Bibliography Section		486/1		
Biomedical Applications	487/1	487/2	488/1 488/2	

### INFORMATION FOR AUTHORS

(Detailed *Instructions to Authors* were published in Vol. 445, pp. 453–456. A free reprint can be obtained by application to the publisher, Elsevier Science Publishers B.V., P.O. Box 330, 1000 AH Amsterdam, The Netherlands.)

**Types of Contributions.** The following types of papers are published in the *Journal of Chromatography* and the section on *Biomedical Applications*: Regular research papers (Full-length papers), Notes, Review articles and Letters to the Editor. Notes are usually descriptions of short investigations and reflect the same quality of research as Full-length papers, but should preferably not exceed six printed pages. Letters to the Editor can comment on (parts of) previously published articles, or they can report minor technical improvements of previously published procedures; they should preferably not exceed two printed pages. For review articles, see inside front cover under Submission of Papers.

**Submission.** Every paper must be accompanied by a letter from the senior author, stating that he is submitting the paper for publication in the *Journal of Chromatography*. Please do not send a letter signed by the director of the institute or the professor unless he is one of the authors.

**Manuscripts.** Manuscripts should be typed in double spacing on consecutively numbered pages of uniform size. The manuscript should be preceded by a sheet of manuscript paper carrying the title of the paper and the name and full postal address of the person to whom the proofs are to be sent. Authors of papers in French or German are requested to supply an English translation of the title of the paper. As a rule, papers should be divided into sections, headed by a caption (*e.g.*, Summary, Introduction, Experimental, Results, Discussion, etc.). All illustrations, photographs, tables, etc., should be on separate sheets.

**Introduction.** Every paper must have a concise introduction mentioning what has been done before on the topic described, and stating clearly what is new in the paper now submitted.

**Summary.** Full-length papers and Review articles should have a summary of 50–100 words which clearly and briefly indicates what is new, different and significant. In the case of French or German articles an additional summary in English, headed by an English translation of the title, should also be provided. (Notes and Letters to the Editor are published without a summary.)

**Illustrations.** The figures should be submitted in a form suitable for reproduction, drawn in Indian ink on drawing or tracing paper. Each illustration should have a legend, all the *legends* being typed (with double spacing) together on a *separate sheet*. If structures are given in the text, the original drawings should be supplied. Coloured illustrations are reproduced at the author's expense, the cost being determined by the number of pages and by the number of colours needed. The written permission of the author and publisher must be obtained for the use of any figure already published. Its source must be indicated in the legend.

**References.** References should be numbered in the order in which they are cited in the text, and listed in numerical sequence on a separate sheet at the end of the article. Please check a recent issue for the layout of the reference list. Abbreviations for the titles of journals should follow the system used by *Chemical Abstracts*. Articles not yet published should be given as "in press" (journal should be specified), "submitted for publication" (journal should be specified), "in preparation" or "personal communication".

**Dispatch.** Before sending the manuscript to the Editor please check that the envelope contains three copies of the paper complete with references, legends and figures. One of the sets of figures must be the originals suitable for direct reproduction. Please also ensure that permission to publish has been obtained from your institute.

**Proofs.** One set of proofs will be sent to the author to be carefully checked for printer's errors. Corrections must be restricted to instances in which the proof is at variance with the manuscript. "Extra corrections" will be inserted at the author's expense.

**Reprints.** Fifty reprints of Full-length papers, Notes and Letters to the Editor will be supplied free of charge. Additional reprints can be ordered by the authors. An order form containing price quotations will be sent to the authors together with the proofs of their article.

**Advertisements.** Advertisement rates are available from the publisher on request. The Editors of the journal accept no responsibility for the contents of the advertisements.

# When & Where

*It happens all the time. You need to know - or be reminded of - when and where next year's CHEMISTRY meetings are being held, what BIOTECHNOLOGY meetings are taking place in Germany, or even what other events are being held in or around London (or Chicago or Vancouver) in the month of May. The list of possible questions can be endless.*

*Further information and a comprehensive brochure can be obtained from:*

*In Europe:*

**EVENTLINE**  
P.O. Box 521  
1000 AN Amsterdam  
The Netherlands  
Tel: (31 20) 5862 751

*In USA/Canada:*

**EVENTLINE**  
P.O. Box 57101  
Philadelphia  
PA 19111, USA  
Tel: (215) 572 7424

*In Asia/Pacific:*

**EVENTLINE**  
(Excerpta Medica)  
57 Wyndham St., 8th floor  
Hong Kong  
Tel: 852 5 222566

*In Japan:*

**EVENTLINE**  
3-28-1 Yushima,  
Bunkyo-ku  
Tokyo 113, Japan  
Tel: 81 3 836 0810

- **EVENTLINE** can help you easily with the answers, and without the clutter of events that might be irrelevant to you.
- **EVENTLINE** is a new database which will tell you the when, where (and much more) on all those conferences, trade fairs, exhibitions, and major sporting events, etc., which may have eluded you.
- **EVENTLINE is comprehensive** – Conferences, Conventions, Symposia, Trade Fairs, Exhibitions, Major Sporting Events, and even Public Holidays defined by more than 700 subject classifications.
- **EVENTLINE is up-to-date** – input is done daily, and we can give you information on events through to the year 2000.
- **EVENTLINE gives you only what you ask for** – you specify the search, e.g. by date, city, field of interest, etc. and we give you what you ask for.
- **EVENTLINE is reasonably priced** – Prices start at \$35 plus a small additional charge per selected event.



**EVENTLINE**™ is a part of the ELSEVIER Publishing Group