

VOL. **601** NOS. **1 + 2** APRIL 2, 1993

COMPLETE IN ONE ISSUE

**CUMULATIVE AUTHOR, SUBJECT,
COMPOUND AND SPECIAL ISSUE INDEXES
VOLUMES 551-600**

JOURNAL OF

CHROMATOGRAPHY

INCLUDING ELECTROPHORESIS AND OTHER SEPARATION METHODS

EDITORS

U. A. Th. Brinkman (Amsterdam)
R. W. Giese (Boston, MA)
J. K. Haken (Kensington, N.S.W.)
K. Macek (Prague)
L. R. Snyder (Orinda, CA)

EDITORS, SYMPOSIUM VOLUMES,

E. Heftmann (Orinda, CA), Z. Deyl (Prague)

EDITORIAL BOARD

D. W. Armstrong (Rolla, MO)
W. A. Aue (Halifax)
P. Božek (Brno)
A. A. Boulton (Saskatoon)
P. W. Carr (Minneapolis, MN)
N. H. C. Cooke (San Ramon, CA)
V. A. Davankov (Moscow)
Z. Deyl (Prague)
S. Dilli (Kensington, N.S.W.)
H. Engelhardt (Saarbrücken)
F. Erni (Basle)
M. B. Evans (Hatfield)
J. L. Glajch (N. Billerica, MA)
G. A. Guiochon (Knoxville, TN)
P. R. Haddad (Hobart, Tasmania)
I. M. Hais (Hradec Králové)
W. S. Hancock (San Francisco, CA)
S. Hjertén (Uppsala)
S. Honda (Higashi-Osaka)
Cs. Horváth (New Haven, CT)
J. F. K. Huber (Vienna)
K.-P. Hupe (Waldbronn)
T. W. Hutchens (Houston, TX)
J. Janák (Brno)
P. Jandera (Pardubice)
B. L. Karger (Boston, MA)
J. J. Kirkland (Newport, DE)
E. sz. Kováts (Lausanne)
A. J. P. Martin (Cambridge)
L. W. McLaughlin (Chestnut Hill, MA)
E. D. Morgan (Keele)
J. D. Pearson (Kalamazoo, MI)
H. Poppe (Amsterdam)
F. E. Regnier (West Lafayette, IN)
P. G. Righetti (Milan)
P. Schoenmakers (Eindhoven)
R. Schwarzenbach (Dübendorf)
R. E. Shoup (West Lafayette, IN)
R. P. Singhal (Wichita, KS)
A. M. Siouffi (Marseille)
D. J. Strydom (Boston, MA)
N. Tanaka (Kyoto)
S. Terabe (Hyogo)
K. K. Unger (Mainz)
R. Verpoorte (Leiden)
Gy. Vigh (College Station, TX)
J. T. Watson (East Lansing, MI)
B. D. Westerlund (Uppsala)

EDITORS, BIBLIOGRAPHY SECTION

Z. Deyl (Prague), J. Janák (Brno), V. Schwarz (Prague)

ELSEVIER

JOURNAL OF CHROMATOGRAPHY

INCLUDING ELECTROPHORESIS AND OTHER SEPARATION METHODS

Scope. The *Journal of Chromatography* publishes papers on all aspects of chromatography, electrophoresis and related methods. Contributions consist mainly of research papers dealing with chromatographic theory, instrumental developments and their applications. The section *Biomedical Applications*, which is under separate editorship, deals with the following aspects: developments in and applications of chromatographic and electrophoretic techniques related to clinical diagnosis or alterations during medical treatment; screening and profiling of body fluids or tissues related to the analysis of active substances and to metabolic disorders; drug level monitoring and pharmacokinetic studies; clinical toxicology; forensic medicine; veterinary medicine; occupational medicine; results from basic medical research with direct consequences in clinical practice. In *Symposium volumes*, which are under separate editorship, proceedings of symposia on chromatography, electrophoresis and related methods are published.

Submission of Papers. The preferred medium of submission is on disk with accompanying manuscript (see *Electronic manuscripts* in the Instructions to Authors, which can be obtained from the publisher, Elsevier Science Publishers B.V., P.O. Box 330, 1000 AH Amsterdam, Netherlands). Manuscripts (in English; four copies are required) should be submitted to: Editorial Office of *Journal of Chromatography*, P.O. Box 681, 1000 AR Amsterdam, Netherlands, Telefax (+31-20) 5862 304, or to: The Editor of *Journal of Chromatography, Biomedical Applications*, P.O. Box 681, 1000 AR Amsterdam, Netherlands. Review articles are invited or proposed in writing to the Editors who welcome suggestions for subjects. An outline of the proposed review should first be forwarded to the Editors for preliminary discussion prior to preparation. Submission of an article is understood to imply that the article is original and unpublished and is not being considered for publication elsewhere. For copyright regulations, see below.

Publication. The *Journal of Chromatography* (incl. *Biomedical Applications*) has 40 volumes in 1993. The subscription prices for 1993 are:

J. Chromatogr. (incl. *Cum. Indexes, Vols. 601-650*) + *Biomed. Appl.* (Vols. 612-651):

Dfl. 8520.00 plus Dfl. 1320.00 (p.p.h.) (total ca. US\$ 5622.75)

J. Chromatogr. (incl. *Cum. Indexes, Vols. 601-650*) only (Vols. 623-651):

Dfl. 7047.00 plus Dfl. 957.00 (p.p.h.) (total ca. US\$ 4573.75)

Biomed. Appl. only (Vols. 612-622):

Dfl. 2783.00 plus Dfl. 363.00 (p.p.h.) (total ca. US\$ 1797.75).

Subscription Orders. The Dutch guilder price is definitive. The US\$ price is subject to exchange-rate fluctuations and is given as a guide. Subscriptions are accepted on a prepaid basis only, unless different terms have been previously agreed upon. Subscriptions orders can be entered only by calendar year (Jan.-Dec.) and should be sent to Elsevier Science Publishers, Journal Department, P.O. Box 211, 1000 AE Amsterdam, Netherlands, Tel. (+31-20) 5803 642, Telefax (+31-20) 5803 598, or to your usual subscription agent. Postage and handling charges include surface delivery except to the following countries where air delivery via SAL (Surface Air Lift) mail is ensured: Argentina, Australia, Brazil, Canada, China, Hong Kong, India, Israel, Japan*, Malaysia, Mexico, New Zealand, Pakistan, Singapore, South Africa, South Korea, Taiwan, Thailand, USA. *For Japan air delivery (SAL) requires 25% additional charge of the normal postage and handling charge. For all other countries airmail rates are available upon request. Claims for missing issues must be made within six months of our publication (mailing) date, otherwise such claims cannot be honoured free of charge. Back volumes of the *Journal of Chromatography* (Vols. 1-611) are available at Dfl. 230.00 (plus postage). Customers in the USA and Canada wishing information on this and other Elsevier journals, please contact Journal Information Center, Elsevier Science Publishing Co. Inc., 655 Avenue of the Americas, New York, NY 10010, USA, Tel. (+1-212) 633 3750, Telefax (+1-212) 633 3764.

Abstracts/Contents Lists published in Analytical Abstracts, Biochemical Abstracts, Biological Abstracts, Chemical Abstracts, Chemical Titles, Chromatography Abstracts, Clinical Chemistry Lookout, Current Awareness in Biological Sciences (CABS), Current Contents/Life Sciences, Current Contents/Physical, Chemical & Earth Sciences, Deep-Sea Research/Part B: Oceanographic Literature Review, Excerpta Medica, Index Medicus, Mass Spectrometry Bulletin, PASCAL-CNRS, Pharmaceutical Abstracts, Referativnyi Zhurnal, Research Alert, Science Citation Index and Trends in Biotechnology.

US Mailing Notice. *Journal of Chromatography* (ISSN 0021-9673) is published weekly (total 52 issues) by Elsevier Science Publishers (Sara Burgerhartstraat 25, P.O. Box 211, 1000 AE Amsterdam, Netherlands). Annual subscription price in the USA US\$ 4573.75 (subject to change), including air speed delivery. Second class postage paid at Jamaica, NY 11431. **USA**

POSTMASTERS: Send address changes to *Journal of Chromatography*, Publications Expediting, Inc., 200 Meacham Avenue, Elmont, NY 11003. Airfreight and mailing in the USA by Publications Expediting.

See inside back cover for Publication Schedule, Information for Authors and information on Advertisements.

© 1993 ELSEVIER SCIENCE PUBLISHERS B.V. All rights reserved.

0021-9673/93/906.00

No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, without the prior written permission of the publisher, Elsevier Science Publishers B.V., Copyright and Permissions Department, P.O. Box 521, 1000 AM Amsterdam, Netherlands.

Upon acceptance of an article by the journal, the author(s) will be asked to transfer copyright of the article to the publisher. The transfer will ensure the widest possible dissemination of information.

Special regulations for readers in the USA. This journal has been registered with the Copyright Clearance Center, Inc. Consent is given for copying of articles for personal or internal use, or for the personal use of specific clients. This consent is given on the condition that the copier pays through the Center the per-copy fee stated in the code on the first page of each article for copying beyond that permitted by Sections 107 or 108 of the US Copyright Law. The appropriate fee should be forwarded with a copy of the first page of the article to the Copyright Clearance Center, Inc., 27 Congress Street, Salem, MA 01970, USA. If no code appears in an article, the author has not given broad consent to copy and permission to copy must be obtained directly from the author. All articles published prior to 1980 may be copied for a per-copy fee of US\$ 2.25, also payable through the Center. This consent does not extend to other kinds of copying, such as for general distribution, resale, advertising and promotion purposes, or for creating new collective works. Special written permission must be obtained from the publisher for such copying.

No responsibility is assumed by the Publisher for any injury and/or damage to persons or property as a matter of products liability, negligence or otherwise, or from any use or operation of any methods, products, instructions or ideas contained in the materials herein. Because of rapid advances in the medical sciences, the Publisher recommends that independent verification of diagnoses and drug dosages should be made.

Although all advertising material is expected to conform to ethical (medical) standards, inclusion in this publication does not constitute a guarantee or endorsement of the quality or value of such product or of the claims made of it by its manufacturer.

This issue is printed on acid-free paper.

Printed in the Netherlands

JOURNAL OF CHROMATOGRAPHY

VOL. 601 (1993)

JOURNAL of CHROMATOGRAPHY

INCLUDING ELECTROPHORESIS AND OTHER SEPARATION METHODS

EDITORS

U. A. Th. BRINKMAN (Amsterdam), R. W. GIESE (Boston, MA), J. K. HAKEN (Kensington, N.S.W.), K. MACEK (Prague),
L. R. SNYDER (Orinda, CA)

EDITORS, SYMPOSIUM VOLUMES

E. HEFTMANN (Orinda, CA), Z. DEYL (Prague)

EDITORIAL BOARD

D. W. Armstrong (Rolla, MO), W. A. Aue (Halifax), P. Boček (Brno), A. A. Boulton (Saskatoon), P. W. Carr (Minneapolis, MN),
N. H. C. Cooke (San Ramon, CA), V. A. Davankov (Moscow), Z. Deyl (Prague), S. Dilli (Kensington, N.S.W.), H. Engelhardt
(Saarbrücken), F. Erni (Basle), M. B. Evans (Hatfield), J. L. Glajch (N. Billerica, MA), G. A. Guiochon (Knoxville, TN), P. R.
Haddad (Hobart, Tasmania), I. M. Hais (Hradec Králové), W. S. Hancock (San Francisco, CA), S. Hjertén (Uppsala), S. Honda
(Higashi-Osaka), Cs. Horváth (New Haven, CT), J. F. K. Huber (Vienna), K.-P. Hupe (Waldbronn), T. W. Hutchens (Houston,
TX), J. Janák (Brno), P. Jandera (Pardubice), B. L. Karger (Boston, MA), J. J. Kirkland (Newport, DE), E. sz. Kováts (Lausanne),
A. J. P. Martin (Cambridge), L. W. McLaughlin (Chestnut Hill, MA), E. D. Morgan (Keele), J. D. Pearson (Kalamazoo, MI), H.
Poppe (Amsterdam), F. E. Regnier (West Lafayette, IN), P. G. Righetti (Milan), P. Schoenmakers (Eindhoven), R. Schwarzenbach
(Dübendorf), R. E. Shoup (West Lafayette, IN), R. P. Singhal (Wichita, KS), A. M. Siouffi (Marseille), D. J. Strydom
(Boston, MA), N. Tanaka (Kyoto), S. Terabe (Hyogo), K. K. Unger (Mainz), R. Verpoorte (Leiden), Gy. Vigh (College Station,
TX), J. T. Watson (East Lansing, MI), B. D. Westerlund (Uppsala)

EDITORS, BIBLIOGRAPHY SECTION

Z. Deyl (Prague), J. Janák (Brno), V. Schwarz (Prague)



ELSEVIER
AMSTERDAM — LONDON — NEW YORK — TOKYO

J. Chromatogr., Vol. 601 (1993)

© 1993 ELSEVIER SCIENCE PUBLISHERS B.V. All rights reserved.

0021-9673/93/906.00

No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, without the prior written permission of the publisher, Elsevier Science Publishers B.V., Copyright and Permissions Department, P.O. Box 521, 1000 AM Amsterdam, Netherlands.

Upon acceptance of an article by the journal, the author(s) will be asked to transfer copyright of the article to the publisher. The transfer will ensure the widest possible dissemination of information.

Special regulations for readers in the USA. This journal has been registered with the Copyright Clearance Center, Inc. Consent is given for copying of articles for personal or internal use, or for the personal use of specific clients. This consent is given on the condition that the copier pays through the Center the per-copy fee stated in the code on the first page of each article for copying beyond that permitted by Sections 107 or 108 of the US Copyright Law. The appropriate fee should be forwarded with a copy of the first page of the article to the Copyright Clearance Center, Inc., 27 Congress Street, Salem, MA 01970, USA. If no code appears in an article, the author has not given broad consent to copy and permission to copy must be obtained directly from the author. All articles published prior to 1980 may be copied for a per-copy fee of US\$ 2.25, also payable through the Center. This consent does not extend to other kinds of copying, such as for general distribution, resale, advertising and promotion purposes, or for creating new collective works. Special written permission must be obtained from the publisher for such copying.

No responsibility is assumed by the Publisher for any injury and/or damage to persons or property as a matter of products liability, negligence or otherwise, or from any use or operation of any methods, products, instructions or ideas contained in the materials herein. Because of rapid advances in the medical sciences, the Publisher recommends that independent verification of diagnoses and drug dosages should be made.

Although all advertising material is expected to conform to ethical (medical) standards, inclusion in this publication does not constitute a guarantee or endorsement of the quality or value of such product or of the claims made of it by its manufacturer.

This issue is printed on acid-free paper.

Printed in the Netherlands

SPECIAL VOLUME

*Cumulative Author, Subject,
Compound and Special Issue Indexes
Volumes 551–600 (1991–1992)*

CONTENTS

Publisher's note	VI
Cumulative Author Index, Volumes 551–600	1
Cumulative Subject Index, Volumes 551–600	129
Cumulative Compound Index, Volumes 551–600	233
Special Issue Index, Volumes 551–600	410

Publisher's note

From this cumulative indexes volume onwards, a separate Special Issue index will be included. These special issues comprise symposium issues as well as issues devoted to a certain topic. The new index will facilitate the identification of papers published in such issues.

We hope this new index will prove useful to our readers.

AUTHOR INDEX

VOLUMES 551-600

A

- Aalto, T., see Rosenberg, C. 552(1991)265
- Abbiati, G., Fossati, T., Arrigoni, M., Rolle, P., Dognini, G. L. and Castiglioni, C.
High-performance liquid chromatographic assay of L- α -glycerophosphorylcholine using a two-step enzymic conversion 566(1991)445
- Abbiati, G., see Fossati, T. 577(1992)382
- Abbott, F. S., see Wright, M. R. 565(1991)225
- Abdel-Baky, S., see Li, W. 588(1991)273
- Abderrezag, A., see Desbène, P.-L. 593(1992)321
- Abdul Nour, J., see Goubran-Botros, H. 597(1992)357
- Abdulrahman, S., Harrison, M. E., Welham, K. J., Baldwin, M. A., Phillipson, J. D. and Roberts, M. F.
High-performance liquid chromatographic-mass spectrometric assay of high-value compounds for pharmaceutical use from plant cell tissue culture: *Cinchona* alkaloids 562(1991)713
- Abe, A., see Yamane, M. 568(1991)11
- Abe, A., see Yamane, M. 575(1992)7
- Abe, A., see Yamane, M. 579(1992)25
- Abe, F., see Yasukawa, K. 597(1992)271
- Abe, H., see O'Dowd, A. 577(1992)347
- Abe, I., Ito, N., Noguchi, K., Kazama, M. and Kasai, K.-I.
Immobilized urokinase column as part of a specific detection system for plasminogen species separated by high-performance affinity chromatography 565(1991)183
- Abe, M., see Kanai, M. 567(1991)415
- Abe, S., see Oda, Y. 572(1991)133
- Abete, C., see Colombini, M. P. 592(1992)255
- Abián, J. and Gelpí, E.
Characteristic fragmentation of thromboxane B₂ in thermospray high-performance liquid chromatography-mass spectrometry 562(1991)153
- Abián, J., Pagès, M. and Gelpí, E.
Use of methyl oxime derivatives to enhance structural information in thermospray high-performance liquid chromatography-mass spectrometry. Analysis of linoleic acid lipoxygenase metabolites in maize embryos 554(1991)155
- Abidi, S. L.
High-performance liquid chromatography of phosphatidic acids and related polar lipids 587(1991)193
- Abidi, S. L. and Mounts, T. L.
High-performance liquid chromatographic separation of molecular species of neutral phospholipids 598(1992)209
- Abiko, Y., see Ohsawa, K. 597(1992)285
- Abolfathi, Z., Bélanger, P.-M., Gilbert, M., Rouleau, J. R. and Turgeon, J.
Improved high-performance liquid chromatographic assay for the stereoselective determination of mexiletine in plasma 579(1992)366
- Abou-Rebyeh, H., Josić, D., Gottschall, K., Schubert-Rehberg, K. and Körber, F.
Purification of carbonic anhydrase isoenzymes by high-performance affinity chromatography and hydrophobic interaction chromatography 566(1991)351
- Abou-Rebyeh, H., Körber, F., Schubert-Rehberg, K., Reusch, J. and Josić, D.
Carrier membrane as a stationary phase for affinity chromatography and kinetic studies of membrane-bound enzymes 566(1991)341
- Abouelezz, M., see Talmadge, K. W. 590(1992)83
- Abraham, M. H. and Whiting, G. S.
Hydrogen bonding. XXI. Solvation parameters for alkylaromatic hydrocarbons from gas-liquid chromatographic data 594(1992)229
- Abraham, M. H., Whiting, G. S., Andonian-Haftvan, J., Steed, J. W. and Grate, J. W.
Hydrogen bonding. XIX. The characterisation of two poly(methylphenylsiloxane)s 588(1991)361
- Abraham, M. H., Whiting, G. S., Doherty, R. M. and Shuely, W. J.
Hydrogen bonding. XVI. A new solute solvation parameter, π^H_2 , from gas chromatographic data 587(1991)213
- Abraham, M. H., Whiting, G. S., Doherty, R. M. and Shuely, W. J.
Hydrogen bonding. XVII. The characterisation of 24 gas-liquid chromatographic stationary phases studied by Poole and co-workers, including molten salts, and evaluation of solute-stationary phase interactions 587(1991)229
- Abramowski, V., see Bories, C. 579(1992)143
- Abu-Lafi, S., see Levin, S. 556(1991)277
- Abumrad, N. N., see Donahue, E. P. 571(1991)29

- Aburada, M., see Wakui, Y. 575(1992)131
- Accorsi, C. A. and Blo, G.
Determination of volatile and non-volatile organic acids in technical sugar solutions by ion-exclusion chromatography 555(1991)65
- Ackermans, M., see Takkenberg, B. 565(1991)430
- Ackermans, M. T., Beckers, J. L., Everaerts, F. M., Hoogland, H. and Tomassen, M. J. H.
Determination of sulphonamides in pork meat extracts by capillary zone electrophoresis 596(1992)101
- Ackermans, M. T., Beckers, J. L., Everaerts, F. M. and Seelen, I. G. J. A.
Comparison of isotachopheresis, capillary zone electrophoresis and high-performance liquid chromatography for the determination of salbutamol, terbutaline sulphate and fenoterol hydrobromide in pharmaceutical dosage forms 590(1992)341
- Ackermans, M. T., Everaerts, F. M. and Beckers, J. L.
Determination of some drugs by micellar electrokinetic capillary chromatography. The pseudo-effective mobility as parameter for screening 585(1991)123
- Ackermans, M. T., Everaerts, F. M. and Beckers, J. L.
Relationship between zone length and step height in isotachopheresis 595(1992)327
- Ackman, R. G., see Shantha, N. C. 587(1991)263
- Adam, M., see Lichý, A. 563(1991)153
- Adams, R. N., see Cammack, J. 565(1991)529
- Addison, R. S., see Cannell, G. R. 563(1991)341
- Addison, T. E., see Hopkins, N. K. 577(1992)87
- Ader, J. C., see Halvorsen, K. 557(1991)501
- Adler, C. M., see Scouten, W. H. 597(1992)231
- Advenier, C., see Moreau, J. 576(1992)103
- Advenier, C., see Palette, C. 563(1991)103
- Agache, P., see Makki, S. 563(1991)407
- Agasøster, T. and Rasmussen, K. E.
Automated analysis of oxolinic acid and flumequine in salmon whole blood and plasma using dialysis combined with trace enrichment as on-line sample preparation for high-performance liquid chromatography 564(1991)171
- Agasøster, T. and Rasmussen, K. E.
Fully automated high-performance liquid chromatographic analysis of whole blood and plasma samples using on-line dialysis as sample preparation. Determination of oxytetracycline in bovine and salmon whole blood and plasma 570(1991)99
- Aggarwal, S. K., Kinter, M. and Herold, D. A.
Determination of cobalt in urine by gas chromatography-mass spectrometry employing nickel as an internal standard 576(1992)297
- Agnus, B., Sebille, B. and Gosselet, M.
Effects of β -cyclodextrin in the mobile phase on the retention and indirect detection of non-electrolytes in reversed-phase liquid chromatography. II. Steroids 552(1991)583
- Aguilar, M. I., see Fang, F. W. 599(1992)163
- Aguilar, M. I., see Purcell, A. W. 593(1992)103
- Ahlgren, J. D., see Anderson, L. W. 581(1992)195
- Ahmed, F. and Modrek, B.
Biosep-SEC-S high-performance size-exclusion chromatographic columns for proteins and peptides 599(1992)25
- Ahmad, F., see Awasthi, S. 584(1992)167
- Ahmad, H., see Awasthi, S. 584(1992)167
- Ahnoff, M., Balmér, K. and Lindman, Y.
By-products in the derivatization of amines with the chiral reagent 2,3,4,6-tetra-O-acetyl- β -D-glucopyranosyl isothiocyanate and their elimination 592(1992)323
- Aiba, S., see Kihira, Y. 597(1992)277
- Aiken, J. H. and Huie, C. W.
Use of hematoporphyrin as a fluorescent stain for detection of lipids in high-performance thin-layer chromatography 588(1991)295
- Aiken, J. H., Huie, C. W. and Terzian, J. A.
Use of eluent containing surfactant for the liquid chromatographic analysis of porphyrins by direct serum injection 584(1992)181
- Aitkenhead, A. R., see Mason, J. L. 570(1991)191
- Akabane, H., see Terazawa, K. 565(1991)453
- Akabane, M., see Shibata, N. 566(1991)187
- Akabori, S., see Tamaki, M. 574(1992)65
- Akane, A., see Fukushima, S. 565(1991)35
- Akapo, S. O. and Simpson, C. F.
Structural investigation of oligomeric *n*-octylsilyl reversed phases 557(1991)515
- Akasaka, K., Ijichi, S., Watanabe, K., Ohru, H. and Meguro, H.
High-performance liquid chromatography and post-column derivatization with diphenyl-1-pyrenylphosphine for fluorimetric determination of triacylglycerol hydroperoxides 596(1992)197
- Åkesson, J., see Karlsson, L. 557(1991)99
- Akimenko, V. K., see Golovchenko, N. P. 591(1992)121
- Akimoto, K., Goto, A. and Ohya, K.
Selective and sensitive determination of lactone and hydroxy acid forms of camptothecin and two derivatives (CPT-11 and SN-38) by high-performance liquid chromatography with fluorescence detection 588(1991)165
- Akiyama, H., Shidawara, S., Mada, A., Toyoda, H., Toida, T. and Imanari, T.
Chemiluminescence high-performance liquid chromatography for the determination of hyaluronic acid, chondroitin sulphate and dermatan sulphate 579(1992)203
- Akiyama, H., see Toyoda, H. 565(1991)141
- Akiyama, S., see Nakashima, K. 584(1992)275
- Akiyama, S., see Toyo'oka, T. 588(1991)61
- Akiyama, T.
Separation of neutral mono- and oligosaccharides derivatized with ethyl *p*-aminobenzoate by high-performance liquid chromatography on an amine-bonded vinyl alcohol copolymer column 588(1991)53
- Akiyama, Y., see Takeda, N. 558(1991)175
- Al-Deen, A. N., Cecchini, D. J. and Giese, R. W.
Purification of DNA-derived deoxynucleotides from leukocytes involving nuclease elution of an ion-exchange column 600(1992)229

- Alaiz, M., Navarro, J. L., Girón, J. and Vioque, E.
Amino acid analysis by high-performance liquid chromatography after derivatization with diethyl ethoxymethylenemalonate 591(1992)181
- Alajbeg, A., Todorić, A., Švel-Cerovečki, S. and Šušterčić, M.
Gas chromatography and gas chromatography-mass spectrometry study of hydrocarbons in Vlasta oil (Adriatic Basin) as the basis for geochemical interpretation 557(1991)255
- Albarrán, G., see Navarro-González, R. 587(1991)247
- Albert, K., see Buszewski, B. 552(1991)415
- Albert, K., see Unger, K. K. 556(1991)395
- Alberts, G., Boomsma, F., Man in 't Veld, A. J. and Schalekamp, M. A. D. H.
Simultaneous determination of catecholamines and dobutamine in human plasma and urine by high-performance liquid chromatography with fluorimetric detection 583(1992)236
- Alberts, G., see Boomsma, F. 574(1992)109
- Alderweireldt, F. C., see Maes, P. C. 558(1991)343
- Alessi-Severini, S., Coutts, R. T., Jamali, F. and Pasutto, F. M.
High-performance liquid chromatographic analysis of methocarbamol enantiomers in biological fluids 582(1992)173
- Alexander, P. W., Glód, B. K. and Haddad, P. R.
Mixed potential response at metallic electrodes in potentiometric detection for ion chromatography 589(1992)201
- Alexander, P. W., see Buchberger, W. 558(1991)181
- Alexander, P. W., see Glód, B. K. 589(1992)209
- Alexander, P. W., see Glód, B. K. 595(1992)149
- Alexander, T. G., see Getek, T. A. 554(1991)191
- Algiman, E., see Kong Sing, Y. L. 558(1991)43
- Alhama, J., López-Barea, J. and Toribio, F.
High-performance affinity chromatography of NADP⁺ dehydrogenases from cell-free extracts using a nucleotide analogue as general ligand 586(1991)51
- Alhama, J., López-Barea, J., Toribio, F. and Roldán, J. M.
Purification and determination of glutamine synthetase by high-performance immunoaffinity chromatography 589(1992)121
- Ali, H., see Guzman, N. A. 559(1991)307
- Allegre, F., see Dumas, L. 570(1991)89
- Allen, E., Myers, C., Frank, D., Pettigrew, A., Thompson, S. and Blumer, J. L.
Application of an organic solvent extraction to the determination of catechol-O-methyltransferase activity by high-performance liquid chromatography in human mononuclear cells 578(1992)175
- Allenmark, S., see Chelminska-Bertilsson, M. 575(1992)237
- Allenmark, S. G., see Andersson, S. 591(1992)65
- Alliegro, M. A., Dyer, K. D., Cragoe, Jr., E. J., Glaser, B. M. and Alliegro, M. C.
High-performance liquid chromatographic method for quantitating plasma levels of amiloride and its analogues 582(1992)217
- Alliegro, M. C., see Alliegro, M. A. 582(1992)217
- Allievi, C., Dostert, P. and Strolin Benedetti, M.
Determination of free salsolinol concentrations in human urine using gas chromatography-mass spectrometry 568(1991)271
- Allison, J., see Schultz, G. A. 590(1992)329
- Allmaier, G., see Gasser, H. 562(1991)257
- Alloni, A., see Gelfi, C. 598(1992)277
- Ally, A. and Park, G.
Rapid determination of creatine, phosphocreatine, purine bases and nucleotides (ATP, ADP, AMP, GTP, GDP) in heart biopsies by gradient ion-pair reversed-phase liquid chromatography 575(1992)19
- Almazova, T. V., see Tennikova, T. B. 555(1991)97
- Al Odaib, A., see Leis, H. J. 578(1992)116
- Alonso, R. M., see Rodríguez, F. J. 578(1992)146
- Alpert, A. J., see Boutin, J. A. 583(1992)137
- Alric, R., see Cociglio, M. 572(1991)269
- Altay, C., see Kutlar, F. 567(1991)359
- Altieri, I., see Pichini, S. 568(1991)267
- Altieri, I., see Pichini, S. 577(1992)358
- Altieri, I., see Pichini, S. 581(1992)143
- Altieri, I., see Zuccaro, P. 567(1991)485
- Altomare, C., see El Tayar, N. 556(1991)181
- Alton, K. B., Hernandez, A., Alvarez, N. and Patrick, J. E.
High-performance liquid chromatographic determination of N-[2(S)-(mercaptomethyl)-3-(2-methylphenyl)-1-oxopropyl]-L-methionine, the active plasma metabolite of a prodrug atriopetidase inhibitor (SCH 42495), using a thiol selective (Au/Hg) amperometric detector 579(1992)307
- Altschuh, D., see Dubs, M.-C. 597(1992)391
- Alvarez, E., see Saunders, P. P. 577(1992)37
- Alvarez, J. G. and Touchstone, J. C.
Separation of acidic and neutral lipids by aminopropyl-bonded silica gel column chromatography 577(1992)142
- Alvarez, N., see Alton, K. B. 579(1992)307
- Alvarez-Bujidos, M. L., Ortiz, A., Balaña, R., Cubría, J. C., Ordoñez, D. and Negro, A.
Determination of luxabendazole in biological fluids by high-performance liquid chromatography 578(1992)321
- Alvarez-Bujidos, M. L., see Negro, A. 576(1992)135
- Alvaro, R. F., Rosegay, A. and Chiu, S.-H. L.
Determination of enantiomeric concentrations of a 2,5-diaryltetrahydrofuran (L-668,750), a platelet-activating factor receptor antagonist, in rat plasma using a chiral α_1 -acid glycoprotein high-performance liquid chromatographic column 578(1992)327
- Alvi, S. N., see Husain, S. 558(1991)435
- Alvi, S. N., see Husain, S. 596(1992)127
- Alvi, S. N., see Husain, S. 600(1992)316
- Amari, J. V., Brown, P. R., Pivarnik, P. E., Sehgal, R. K. and Turcotte, J. G.
Isolation of experimental anti-AIDS glycerophospholipids by micro-preparative reversed-phase high-performance liquid chromatography 590(1992)153
- Amidon, G. L., see Janiczek, N. 571(1991)179
- An, D., see Zhou, W. 589(1992)358
- Ancarani, P., see Garibotto, G. 572(1991)111
- Anders, G., see Matucha, M. 588(1991)251

- Andersen, J. V. and Hansen, S. H.
Simultaneous quantitative determination of naproxen, its metabolite 6-O-desmethylnaproxen and their five conjugates in plasma and urine samples by high-performance liquid chromatography on dynamically modified silica 577(1992)325
- Andersen, J. V. and Hansen, S. H.
Simultaneous determination of (*R*)- and (*S*)-naproxen and (*R*)- and (*S*)-6-O-desmethylnaproxen by high-performance liquid chromatography on a Chiral-AGP column 577(1992)362
- Andersen, M. P., see Jørgensen, M. 577(1992)167
- Anderson, C. B., see Mangino, M. J. 577(1992)157
- Anderson, G. M., see Chakraborty, M. 571(1991)235
- Anderson, K., see Boppana, V. K. 593(1992)29
- Anderson, K. L., see Wang, L. H. 579(1992)259
- Anderson, L., Cummings, J. and Smyth, J. F.
Rapid and selective isolation of radiolabelled inositol phosphates from cancer cells using solid-phase extraction 574(1992)150
- Anderson, L. W., Parker, R. J., Collins, J. M., Ahlgren, J. D., Wilkinson, D. and Strong, J. M.
Gas chromatographic-mass spectrometric method for routine monitoring of 5-fluorouracil in plasma of patients receiving low-level protracted infusions 581(1992)195
- Anderson, R. E., see Chen, H. 578(1992)124
- Andersson, J. T.
Comments on the Note "Liquid chromatographic determination of planar aromatic sulphur compounds in crude oil", *J. Chromatogr.*, 475 (1989) 421 585(1991)376
- Andersson, J. T.
Comments on the Note "Liquid chromatographic determination of planar aromatic sulphur compounds in crude oil", *J. Chromatogr.*, 475 (1989) 421 585(1991)376
- Andersson, J. T., see Fedorak, P. M. 591(1992)362
- Andersson, P. E., Demirbükler, M. and Blomberg, L. G.
Quantitative hydrocarbon group analysis of gasoline and diesel fuel by supercritical fluid chromatography 595(1992)301
- Andersson, S., Thompson, R. A. and Allenmark, S. G.
Direct liquid chromatographic separation of enantiomers on immobilized protein stationary phases. IX. Influence of the cross-linking reagent on the retentive and enantioselective properties of chiral sorbents based on bovine serum albumin 591(1992)65
- Andert, K., see Brandl, G. 586(1991)249
- Andiné, P., see Orwar, O. 566(1991)39
- Andoh, H., see Ohhashi, T. 585(1991)153
- Andonian-Haftvan, J., see Abraham, M. H. 588(1991)361
- André, D., see Vérité, P. 578(1992)134
- André, G., see Chauret, N. 594(1992)179
- Andresen, A. T., Jacobsen, P. B. and Rasmussen, K. E.
Automated high-performance liquid chromatography of iopentol in human plasma and whole blood using on-line dialysis as sample preparation 575(1992)93
- Andresen, A. T., Krogh, M. and Rasmussen, K. E.
On-line dialysis and weak cation-exchange enrichment of dialysate. Automated high-performance liquid chromatography of pholcodine in human plasma and whole blood 582(1992)123
- Andreux, J. P., see Merino-Dugay, A. 579(1992)73
- Andriamboavonjy, E., Flaschel, E. and Renken, A.
Separation of functionalized dextrans by reversed-phase high-performance liquid chromatography 587(1991)288
- Andriollo, O., Lartigue-Mattei, C., Chabard, J. L., Bargnoux, H., Petit, J., Berger, J. A. and Pognat, J. F.
Measurement of trazodone in plasma and brain of rat by capillary gas chromatography with a nitrogen-selective detector 575(1992)301
- Andrzejczak, J., see Paczkowski, D. 573(1992)235
- Anelli, P. L., Tomba, C. and Uggeri, F.
Optical resolution of 2-chloro-3-phenylmethoxypropanoic acid after derivatization with (*S*)-2-octanol by high-performance liquid chromatography 589(1992)346
- Ang, S.-G. and Wong, V. W. T.
Chromatographic analysis of low-molecular-mass copper-binding ligands from the crab species *Scylla serrata* and *Portunus pelagicus* 599(1992)21
- Angelo, H. R., see Christensen, H. R. 574(1992)161
- Angerer, J. and Hörsch, B.
Determination of aromatic hydrocarbons and their metabolites in human blood and urine (Review) 580(1992)229
- Angiboust, J.-F., see Debal, V. 581(1992)93
- Anigbogu, V. C., Muñoz de la Peña, A., Ndou, T. T. and Warner, I. M.
Retention behavior of β -cyclodextrin complexes of anthracene and pyrene using reversed-phase liquid chromatography. The effects of *tert*-butanol and cyclopentanol 594(1992)37
- Anliker, S. L., Hamilton, M., Bopp, R. J. and Goldberg, M. J.
Sensitive method for the quantitation of nortriptyline and 10-hydroxynortriptyline in human plasma by capillary gas chromatography with electron-capture detection 573(1992)141
- Annaert, W. G., see Llona, I. 596(1992)51
- Annapoorani, K. S., see Ragupathi, G. 574(1992)267
- Annette, D., see Graham, W. D. 594(1992)187
- Antia, F. D. and Horváth, C.
Analysis of isotachic patterns in displacement chromatography 556(1991)119
- Anton, K., Bach, M. and Geiser, A.
Supercritical fluid chromatography in the routine stability control of antipruritic preparations 553(1991)71
- Antonelli, A., Conte, L. S. and Lercker, G.
Applications of capillary gas chromatography to the quality control of butter and related products 552(1991)273
- Antoniewicz, S. M., Cook, J. A. and Brown, R. R.
Determination of ciclesanine in human plasma by high-performance liquid chromatography using automated column switching 573(1992)93
- Antonozzi, I., see Carducci, C. 553(1991)149
- Aoki, H., see Okazaki, O. 563(1991)313
- Aoki, I., Okumura, M. and Yashiki, T.
High-performance liquid chromatographic determination of lansoprazole and its metabolites in human serum and urine 571(1991)283
- Aoki, I., see Yamaguchi, M. 575(1992)123

- Aoki, T., see Ueno, R. 573(1992)333
- Aoyagi, H., see Ono, S. 597(1992)293
- Aoyama, H., see Komatsu, S. 568(1991)460
- Aoyama, R. G., McErlane, K. M., Erber, H., Kitts, D. D. and Burt, H. M.
High-performance liquid chromatographic analysis of oxytetracycline in chinook salmon following administration of medicated feed 588(1991)181
- Apffel, A. and Perry, M. L.
Quantitation and linearity for particle-beam liquid chromatography-mass spectrometry 554(1991)103
- Appelt, G., see Hänsler, M. 563(1991)63
- Arai, I., see Shibasaki, H. 576(1992)47
- Arai, T., see Ueno, M. 597(1992)197
- Araki, M., see Yamasaki, N. 597(1992)207
- Araki, T., see Kimata, K. 558(1991)19
- Araki, T., see Kimata, K. 594(1992)87
- Araki, T., see Kimata, K. 595(1992)77
- Arand, M., see Oesch, F. 582(1992)232
- Arcelloni, C., see Fermo, I. 593(1992)171
- Archambault, J., see Chauret, N. 588(1991)281
- Archambault, J., see Chauret, N. 594(1992)179
- Arcuri, F., see Castagnetta, L. A. 572(1991)25
- Ardemagni, R., see Lombardi, F. 576(1992)129
- Ardid, D., see Coudore, F. 584(1992)249
- Arimoto, H., see Kamizono, A. 567(1991)113
- Arin, M. J., Diez, M. T. and Resines, J. A.
Rapid and simple method for the determination of urinary benzoic and phenylacetic acids and their glycine conjugates in ruminants by reversed-phase high-performance liquid chromatography 582(1992)13
- Ariyoshi, Y., see Ito, K. 598(1992)237
- Arm, H., see Brügger, R. 557(1991)163
- Arm, H., see Brügger, R. 592(1992)309
- Arm, H., see Straub, R. 585(1991)195
- Armand, J.-P., see Barilero, I. 575(1992)275
- Armando, I., Grossman, E., Hoffman, A. and Goldstein, D. S.
Method for measuring endogenous 3-O-methyl dopa in urine and plasma 568(1991)45
- Armstead, I. P. and Ling, J. R.
Chromatographic separation of mixed peptides from amino acids in biological digests with volatile buffers 586(1991)259
- Arnaud, J., Fortis, I., Blachier, S., Kia, D. and Favier, A.
Simultaneous determination of retinol, α -tocopherol and β -carotene in serum by isocratic high-performance liquid chromatography 572(1991)103
- Arnold, P. J., Guserle, R., Luckow, V., Hemmer, R. and Grote, H.
Liquid chromatography-mass spectrometry in metabolic research. I. Metabolites of benzbromarone in human plasma and urine 544(1991)267
- Aronson, A. L., see Tyczkowska, K. L. 594(1992)195
- Aronson, A. L., see Voyksner, R. D. 567(1991)389
- Arpino, P., see Bruchet, A. 562(1991)469
- Arrigoni, M., see Abbiati, G. 566(1991)445
- Arrigoni Martelli, E., see Mancinelli, A. 553(1991)81
- Arrigoni Martelli, E., see Zezza, F. 593(1992)99
- Arrio, B., see Hill, M. 589(1992)101
- Arrowood, S. and Hoyt, Jr., A. M.
Determination of cimetidine in pharmaceutical preparations by capillary zone electrophoresis 586(1991)177
- Arrowood, S., Hoyt, Jr., A. M. and Sepaniak, M. J.
Monitoring of benzylpenicillin decomposition in gastric contents by capillary zone electrophoresis 583(1992)105
- Arshady, R.
Beaded polymer supports and gels (Review). I. Manufacturing techniques 586(1991)181
- Arshady, R.
Beaded polymer supports and gels (Review). II. Physico-chemical criteria and functionalization 586(1991)199
- Arts, C. J. M., Van Baak, M. J. and Den Hartog, J. M. P.
Control system for detection of the illegal use of naturally occurring steroids in calves 564(1991)429
- Arvidsson, E., Jansson, S. O. and Schill, G.
Retention processes on α_1 -acid glycoprotein-bonded stationary phase 591(1992)55
- Arvidsson, T., Collijn, E., Tivert, A.-M. and Rosén, L.
Peak distortion in the column liquid chromatographic determination of omeprazole dissolved in borax buffer 586(1991)271
- Asakawa, N., see Miwa, T. 566(1991)163
- Asakawa, N., see Oda, Y. 572(1991)133
- Asano, T., see Miwa, H. 568(1991)25
- Ascalone, V., Flaminio, L., Guinebault, P., Thénot, J. P. and Morselli, P. L.
Determination of zolpidem, a new sleep-inducing agent, and its metabolites in biological fluids: pharmacokinetics, drug metabolism and overdosing investigations in humans 581(1992)237
- Asenjo, J. A., see Leser, E. W. 584(1992)43
- Ashmore, S. P., see Mason, J. L. 570(1991)191
- Aspeslet, L. J., see Torok-Both, G. A. 579(1992)99
- Asquith, T. N., see Neubecker, T. A. 574(1992)340
- Assadullahi, T. P., Dagli, E. and Warner, J. O.
High-performance liquid chromatography method for serum methotrexate levels in children with severe steroid-dependent asthma 565(1991)349
- Assandri, A., see Giachetti, C. 585(1991)111
- Asthana, O. P., see Paliwal, J. K. 572(1991)219
- Astier, A.
Simultaneous high-performance liquid chromatographic determination of urinary metabolites of benzene, nitrobenzene, toluene, xylene and styrene 573(1992)318
- Atamna, I. Z., Issaq, H. J., Muschik, G. M. and Janini, G. M.
Optimization of resolution in capillary zone electrophoresis: combined effect of applied voltage and buffer concentration 588(1991)315
- Athalye, A. M., Gibbs, S. J. and Lightfoot, E. N.
Predictability of chromatographic protein separations. Study of size-exclusion media with narrow particle size distributions 589(1992)71
- Attard Barbini, D., see Di Muccio, A. 552(1991)241
- Attard Barbini, D., see Di Muccio, A. 553(1991)305
- Au, J. L.-S., see Wientjes, M. G. 563(1991)400

- Aubert, M.-C., Lee, C. R., Krstulović, A. M., Lesellier, E., Pêchard, M.-R. and Tchaplá, A.
Separation of *trans/cis* α - and β -carotenes by supercritical fluid chromatography. I. Effects of temperature, pressure and organic modifiers on the retention of carotenes 557(1991)47
- Aucoin, D. P., see Hedeén, K. M. 572(1991)239
- Aucoin, D. P., see Tyczkowska, K. L. 576(1992)305
- Auriola, S., Martinsen, A., Oksman-Caldentey, K.-M. and Naaranlahti, T.
Analysis of tropane alkaloids with thermospray high-performance liquid chromatography-mass spectrometry 562(1991)737
- Auriola, S., Naaranlahti, T. and Lapinjoki, S. P.
Determination of *Catharanthus roseus* alkaloids by high-performance liquid chromatography-isotope dilution thermospray-mass spectrometry 554(1991)227
- Auriola, S., see Pennanen, S. 568(1991)125
- Auriola, S., see Raatikainen, O. 585(1991)247
- Auriola, S. O. K., Lepistö, A.-M., Naaranlahti, T. and Lapinjoki, S. P.
Determination of taxol by high-performance liquid chromatography-thermospray mass spectrometry 594(1992)153
- Ausili, A., see Di Muccio, A. 552(1991)241
- Ausili, A., see Di Muccio, A. 553(1991)305
- Awadé, A., see Gonzalès, T. 584(1992)101
- Awadzi, K., see Hoaksey, P. E. 566(1991)244
- Awasthi, S., Ahmad, F., Sharma, R. and Ahmad, H.
Reversed-phase chromatographic method for specific determination of glutathione in cultured malignant cells 584(1992)167
- Axelson, J. E., see Wright, M. R. 565(1991)225
- Axelson, J. E., see Yeleswaram, K. 565(1991)383
- Ayesh, R., see Zhang, A. Q. 584(1992)141
- Aymard, N., see Traiffort, E. 571(1991)231
- Ayotte, C., see Massé, R. 562(1991)323
- Azad, A. R. M., see Nachman, M. 597(1992)155
- Ba, B., Eckert, G. and Leube, J.
Use of dabsylation, column switching and chiral separation for the determination of a renin inhibitor in rat, marmoset and human plasma 572(1991)277
- Baarslag, P. C., see De Bokx, P. K. 594(1992)9
- Baastoe, M. B. and Lundanes, E.
Capillary supercritical fluid chromatography of primary aliphatic amines using carbon dioxide and nitrous oxide as mobile phases 558(1991)458
- Baba, Y., Matsuura, T., Wakamoto, K. and Tshako, M.
Comparison of high-performance liquid chromatography with capillary gel electrophoresis in single-base resolution of polynucleotides 558(1991)273
- Bach, M., see Anton, K. 553(1991)71
- Bacosi, A., see Pichini, S. 581(1992)143
- Baczynskyj, L.
Particle beam liquid chromatography-mass spectrometry on a double-focusing sector instrument 562(1991)13
- Badger, S. E., see Levison, P. R. 590(1992)49
- Badoud, R., Fay, L., Richli, U. and Hušek, P.
Gas chromatographic determination of N-carboxymethyl amino acids, the periodate oxidation products of Amadori compounds 552(1991)345
- Baessler, M., Boeden, H.-F., Koelsch, R. and Lasch, J.
Cellulose beads: a weak leaking affinity support 589(1992)93
- Baeyens, W., see Lansens, P. 586(1991)329
- Baeyens, W. R. G.
Fluorometric analysis in biomedical chemistry; trends and techniques including HPLC applications (edited by N. Ichinose, G. Schwedt, F. M. Schnepel and K. Adachi) (Book Review) 590(1992)373
- Baeyens, W. R. G., see Lin Ling, B. 553(1991)433
- Bagchi, D., see Shara, M. A. 576(1992)221
- Baglin, A., see Roux, A. 570(1991)453
- Bagnati, R., Paleologo Oriundi, M., Russo, V., Danese, M., Berti, F. and Fanelli, R.
Determination of zeranol and β -zeaxanthin in calf urine by immunoaffinity extraction and gas chromatography-mass spectrometry after repeated administration of zeranol 564(1991)493
- Baguley, B. C., see Kestell, P. 564(1991)315
- Bailey, A. J., see Sims, T. J. 582(1992)49
- Bailey, C. A., see Guzman, N. A. 598(1992)123
- Bailey, F. J., see Lehman, E. D. 574(1992)225
- Bailey, L. C., see Wilhelm, J. A. 578(1992)231
- Baillet, A., see Rakotomanga, S. 570(1991)277
- Baillet, A., see Taverna, M. 558(1991)105
- Bailon, P., see Nachman, M. 597(1992)155
- Baiocchi, C., see Silvestro, L. 563(1991)443
- Baiocchi, C., see Silvestro, L. 591(1992)225
- Baird, W. M., see Teffera, Y. 577(1992)69
- Baj, S. and Kulicki, Z.
High-performance liquid chromatographic determination of products of autoxidation of isopropylbenzene derivatives 588(1991)33
- Baker, G. B., see Paetsch, P. R. 573(1992)313
- Baker, G. B., see Torok-Both, G. A. 579(1992)99
- Bakri, D., see Jouve, J. 567(1991)331
- Balaña, R., see Alvarez-Bujidos, M. L. 578(1992)321
- Balcells, J., Guada, J. A., Peiró, J. M. and Parker, D. S.
Simultaneous determination of allantoin and oxypurines in biological fluids by high-performance liquid chromatography 575(1992)153
- Balconi, L., see Blo, G. 556(1991)249
- Baldo, C., see Rizzolo, A. 553(1991)187
- Baldwin, M. A., see Abdulrahman, S. 562(1991)713

- Baliková, M.
Selective system of identification and determination of antidepressants and neuroleptics in serum or plasma by solid-phase extraction followed by high-performance liquid chromatography with photodiode-array detection in analytical toxicology 581(1992)75
- Balmér, K., Lagerström, P.-O., Persson, B.-A. and Schill, G.
Reversed retention order and other stereoselective effects in the separation of amino alcohols on Chiralcel OD 592(1992)331
- Balmér, K., Persson, A., Lagerström, P.-O., Persson, B.-A. and Schill, G.
Liquid chromatographic separation of the enantiomers of metoprolol and its α -hydroxy metabolite on Chiralcel OD for determination in plasma and urine 553(1991)391
- Balmér, K., see Ahnoff, M. 592(1992)323
- Balogh, M. P. and Stacey, C. C.
Chromatographic developments in low flow delivery for liquid chromatography-mass spectrometry 562(1991)73
- Baltas, M., Benbakkar, M., Gorrichon, L. and Zedde, C.
Plant growth regulators G₁, G₂, G₃. Synthesis, extraction and determination of leaf content in *Eucalyptus grandis* 600(1992)323
- Bán, T., Papp, E. and Inczédy, J.
Reversed-phase high-performance liquid chromatography of anionic and ethoxylated non-ionic surfactants and pesticides in liquid pesticide formulations 593(1992)227
- Bánhegyi, G., Cserháti, T. and Valkó, K.
Retention of some ethylenediamine oligomers in reversed-phase chromatography 553(1991)407
- Bank, R. A., Eriksson, A. W. and Pals, G.
High-performance liquid chromatography: purification and chromatographic behaviour of molecular variants of pepsinogen A from human urine 571(1991)47
- Banke, N., Hansen, K. and Diers, I.
Detection of enzyme activity in fractions collected from free solution capillary electrophoresis of complex samples 559(1991)325
- Banner, C. D.
Sources of plant sterol contaminants encountered in low level steroid analysis 572(1991)307
- Banno, K.
Analytical studies on the chiral separation and simultaneous determination of pantothenic acid and hopantenic acid enantiomers in rat plasma by gas chromatography-mass fragmentography. Reply to Küsters *et al.* 576(1992)181
- Banno, K., Horimoto, S. and Mabuchi, M.
Assay of nicergoline and three metabolites in human plasma and urine by high-performance liquid chromatography-atmospheric pressure ionization mass spectrometry 568(1991)375
- Banno, K., Horimoto, S. and Matsuoka, M.
Analytical studies on the chiral separation and simultaneous determination of pantothenic acid and hopantenic acid enantiomers in rat plasma by gas chromatography-mass fragmentography 564(1991)1
- Banno, K., see Ishii, K. 564(1991)338
- Banno, K., see Matsuoka, M. 581(1992)65
- Bannwarth, B., see Leroy, P. 564(1991)258
- Bao, J., see Towns, J. K. 599(1992)227
- Barberi, M., Merlin, J. L. and Weber, B.
Sensitive determination of free and plasma protein-bound dipyrindamole by high-performance liquid chromatography 565(1991)511
- Barberi-Heyob, M., Merlin, J. L. and Weber, B.
Determination of 5-fluorouracil and its main metabolites in plasma by high-performance liquid chromatography 573(1992)247
- Barberi-Heyob, M., Merlin, J. L. and Weber, B.
Analysis of 5-fluorouracil in plasma and urine by high-performance liquid chromatography 581(1992)281
- Barbhayya, R. H., see Janiszewski, J. S. 577(1992)151
- Barbhayya, R. H., see Papp, E. A. 574(1992)93
- Barbier, B., see Jaussaud, P. 573(1992)136
- Barbier, F., see De Vos, M. 564(1991)296
- Barceló, D., Durand, G., Vreeken, R. J., De Jong, G. J., Lingeman, H. and Brinkman, U. A. T.
Evaluation of eluents in thermospray liquid chromatography-mass spectrometry for identification and determination of pesticides in environmental samples 553(1991)311
- Barceló, D., see Durand, G. 554(1991)233
- Barceló, D., see Durand, G. 562(1991)507
- Bárcena, J. A., see Pascual, P. 581(1992)49
- Barchielli, M., see Coppi, G. 563(1991)385
- Bard, M., see Goss, T. A. 588(1991)157
- Barefoot, A. C., see Reiser, R. W. 554(1991)91
- Barford, R. A., see Brewster, J. D. 598(1992)23
- Bargnoux, H., see Andriollo, O. 575(1992)301
- Bargossi, A. M., see Grossi, G. 593(1992)217
- Barham, C., see Liu, H. 576(1992)163
- Barilero, I., Gandia, D., Armand, J.-P., Mathieu-Boué, A., Ré, M., Gouyette, A. and Chabot, G. G.
Simultaneous determination of the camptothecin analogue CPT-11 and its active metabolite SN-38 by high-performance liquid chromatography: application to plasma pharmacokinetic studies in cancer patients 575(1992)275
- Barinaga, C. J., see Smith, R. D. 559(1991)197
- Baron, A., see Delage, E. 555(1991)125
- Baron, B., see Bories, C. 579(1992)143
- Barré, J., see Hervé, F. 577(1992)43
- Barrett, D., see Buck, C. F. 594(1992)225
- Barrett, D. A., Shaw, P. N. and Davis, S. S.
Determination of morphine and 6-acetylmorphine in plasma by high-performance liquid chromatography with fluorescence detection 566(1991)135
- Barry, E. F., see Ombaba, J. M. 598(1992)97
- Bartels, M. J. and Kastl, P. E.
Analysis of 3,5,6-trichloropyridinol in human urine using negative-ion chemical ionization gas chromatography-mass spectrometry 575(1992)69
- Bartha, Á., Ståhlberg, J. and Szokoli, F.
Extension of the electrostatic retention model of reversed-phase ion-pair high-performance liquid chromatography to include the effect of the eluent pH 552(1991)13

- Barthomeuf, C., Pourrat, H. and Regerat, F.
Densitometry for analysis of protein and peptide hydrolysates: application to luteinizing hormone releasing hormone 568(1991)451
- Bartle, K. D., Baulch, D. L., Clifford, A. A. and Coleby, S. E.
Magnitude of the diffusion coefficient anomaly in the critical region and its effect on supercritical fluid chromatography 557(1991)69
- Bartlett, K., see Pourfarzam, M. 570(1991)253
- Bartolucci, G. B., see Sturaro, A. 590(1992)223
- Bartunik, H. D., see Jacob, L. 587(1991)85
- Bartunik, L. J., see Jacob, L. 587(1991)85
- Baruffini, A., De Lorenzi, E., Gandini, C., Kitsos, M. and Massolini, G.
High-performance liquid chromatographic determination of α -tocopheryl nicotinate in cosmetic preparations 593(1992)95
- Basak, A., Yuan, X. W., Seidah, N. G., Chrétien, M. and Lazure, C.
Affinity purification of proteinases by a combination of immobilized peptidyl aldehyde and semicarbazone 581(1992)17
- Baselier, M. R. P., see Van Tellingen, O. 573(1992)328
- Bassi, F. A., see Castagnola, M. 572(1991)51
- Bast, A., see De Jong, J. 570(1991)209
- Bast, A., see De Jong, J. 574(1992)273
- Bastian, T. L., see Herring, V. L. 567(1991)221
- Bastide, P., see Sautou, V. 571(1991)298
- Batley, M., see Haynes, P. A. 581(1992)187
- Battaglia, F. C., see Loy, G. L. 562(1991)169
- Battino, M., see Grossi, G. 593(1992)217
- Baud, F. J., see Houzé, P. 574(1992)305
- Baudendistel, L. J., see Scott, M. D. 576(1992)149
- Bauer, P., see Royer-Morrot, M. J. 581(1992)297
- Bauer, T., see Sturm, G. 562(1991)351
- Bauer, W., see Böcker, R. H. 568(1991)363
- Baulch, D. L., see Bartle, K. D. 557(1991)69
- Baum, O., see Josić, D. 590(1992)59
- Baumann, J., see Yamamoto, H. 593(1992)313
- Baumes, R. L., see Voirin, S. G. 590(1992)313
- Baumes, R. L., see Voirin, S. G. 595(1992)269
- Baune, B., see Fernandez, C. 572(1991)195
- Baurain, R., see Van Tellingen, O. 553(1991)47
- Baurain, R., see Van Tellingen, O. 574(1992)293
- Bavazzano, P., see Busini, S. 577(1992)180
- Bavazzano, P., see Santoni, G. 581(1992)287
- Bayer, E., see Buszewski, B. 552(1991)415
- Bayer, E., see Unger, K. K. 556(1991)395
- Baylocq-Ferrier, D., see Rakotomanga, S. 570(1991)277
- Baylocq-Ferrier, D., see Taverna, M. 558(1991)105
- Bayne, S., Hoogerbrugge, C. M., Thomsen, J., Skriver, L., Van Buul-Offers, S. C. and Van den Brande, J. L.
Primary sequences of insulin-like growth factors 1 and 2 isolated from porcine plasma 562(1991)391
- Bayne, W. F., see Constanzer, M. L. 566(1991)127
- Bayonove, C. L., see Voirin, S. G. 590(1992)313
- Bayonove, C. L., see Voirin, S. G. 595(1992)269
- Beard, A., Naikwadi, K. and Karasek, F. W.
Comparison of extraction methods for polychlorinated dibenzo-*p*-dioxins and dibenzofurans in fly ash using gas chromatography-mass spectrometry 589(1992)265
- Beaumier, P., see Matassa, L. C. 573(1992)43
- Becchi, M., see Ventura, R. 562(1991)723
- Becher, M., see Saeed, K. 567(1991)185
- Beck, O., Boreus, L. O., Lafolie, P. and Jacobsson, G.
Chiral analysis of methadone in plasma by high-performance liquid chromatography 570(1991)198
- Beck, O., see Helander, A. 579(1992)340
- Becker, M., see Boeden, H.-F. 552(1991)389
- Becker, R. E., see Unni, L. K. 573(1992)99
- Becker, R. E., see Unni, L. K. 573(1992)275
- Beckers, J. L., see Ackermans, M. T. 585(1991)123
- Beckers, J. L., see Ackermans, M. T. 590(1992)341
- Beckers, J. L., see Ackermans, M. T. 595(1992)327
- Beckers, J. L., see Ackermans, M. T. 596(1992)101
- Becket, J. M., see Jones, H. A. 570(1991)361
- Becue, T. and Broquaire, M.
Confirmation of the structure of by-products in the synthesis of Modafinil by liquid chromatography-mass spectrometry 557(1991)489
- Beecher, G. R., see Khachik, F. 582(1992)153
- Beecken, V., see Jacob, L. 587(1991)85
- Beeley, J. A.
Clinical applications of electrophoresis of human salivary proteins (Review) 569(1991)261
- Beetens, J., see Weber, C. 562(1991)599
- Beetens, J., see Weber, C. 577(1992)1
- Begley, P., Corbin, R., Foulger, B. E. and Simmonds, P. G.
Photoemissive ionisation source for ion mobility detectors 588(1991)239
- Begutter, H., see Pachinger, A. 558(1991)369
- Beijnen, J. H., see Burger, D. M. 584(1992)239
- Beijnen, J. H., see Kaijser, G. P. 571(1991)121
- Beijnen, J. H., see Kaijser, G. P. 583(1992)175
- Beijnen, J. H., see Van Tellingen, O. 553(1991)47
- Beijnen, J. H., see Van Tellingen, O. 573(1992)328
- Beijnen, J. H., see Van Tellingen, O. 574(1992)293
- Beijnen, J. H., see Van Tellingen, O. 576(1992)158
- Beke, R., see De Vos, M. 564(1991)296
- Beker, D. and Radić, B.
Fast determination of ochratoxin A in serum by liquid chromatography: comparison with enzymic spectrofluorimetric method 570(1991)441
- Bekkers, M. H. J., see De Bokx, P. K. 598(1992)115
- Beksaç, S., see Kutlar, F. 567(1991)359
- Bélanger, P.-M., see Abolfathi, Z. 579(1992)366
- Belenkii, B. G., see Tennikova, T. B. 555(1991)97
- Belik, M. Y., see Galushko, S. V. 553(1991)143
- Belik, M. Y., see Galushko, S. V. 600(1992)79
- Belin, J.-M., see Serrano-Carreón, L. 584(1992)129
- Bell, C. J., see Bernert, Jr., J. T. 578(1992)1
- Bell, H., see Tallaksen, C. M. E. 564(1991)127
- Bell, R. G.
Preparative high-performance liquid chromatographic separation and isolation of bacitracin components and their relationship to microbiological activity 590(1992)163

- Bellato, P., see Mariot, R. 576(1992)143
- Belliardo, F., Bertolino, A., Brandolo, G. and Lucarelli, C.
Micro-liquid chromatography method for the determination of ciclopiroxolamine after pre-column derivatization in topical formulations 553(1991)41
- Belliardo, F., see Lucarelli, C. 573(1992)150
- Belloq, J., see Budzinski, H. 590(1992)297
- Bellon, B., see Rugani, N. 583(1992)246
- Bellotti, V., see Pianezzola, E. 574(1992)170
- Bellotto, N., Reiner, V., Verga, F., Canobbio, G., Lietti, F., Magnani, P. and Lucarelli, C.
Determination of 2'-deoxy-5-iodouridine and its metabolite 5-iodouracil by high-performance liquid chromatography with ultraviolet absorbance detection in human serum 572(1991)327
- Belpaire, F. M., see Rosseel, M. T. 568(1991)239
- Belpaire, F. M., see Vermeulen, A. M. 567(1991)472
- Bemgård, A., Colmsjö, A. and Lundmark, B.-O.
Gas chromatographic analysis of high-molecular-weight polynuclear aromatic hydrocarbons. I. Molecular weight 328 595(1992)247
- Benassi, S., see Santoni, G. 581(1992)287
- Benatar, M. F., see Houzé, P. 574(1992)305
- Benavente, O., see Castillo, J. 555(1991)285
- Benbakkar, M., see Baltas, M. 600(1992)323
- Beneš, M. J., see Turková, J. 597(1992)19
- Benet, L. Z., see Han, C. 579(1992)237
- Benfenati, E., see Celardo, A. 568(1991)407
- Benlyas, M., see Moussaoui, M. 558(1991)71
- Benlyas, M., see Moussaoui, M. 591(1992)115
- Bennett, J. L., McMillan, J. and Dimitrov, N. V.
High-performance liquid chromatographic method for the analysis of Oltipraz in human serum and urine 573(1992)146
- Bennett, S. M., see Jajoo, H. K. 577(1992)299
- Benoit, E., Jaussaud, P., Besse, S., Videmann, B., Courtot, D., Delatour, P. and Bonnaire, Y.
Identification of a benzhydrylic metabolite of ketoprofen in horses by gas chromatography-mass spectrometry and high-performance liquid chromatography 583(1992)167
- Benoît, F., see Varin, F. 574(1992)57
- Benowitz, N. L., see Jacob, III, P. 583(1992)145
- Bensoussan, M., see Serrano-Carreón, L. 584(1992)129
- Bente III, P. F., see Huang, X. C. 600(1992)289
- Bentrop, D. and Engelhardt, H.
Chromatographic characterization of ion exchangers for high-performance liquid chromatography of proteins. I. Chromatographic determination of loading capacity for low- and high-molecular mass anions 556(1991)363
- Benziger, D. P., see Skee, D. M. 568(1991)494
- Benzo, M., see Venturelli, E. 582(1992)7
- Beran, M., Přikrylová, V., Sedmera, P., Novák, J., Zima, J., Blumauerová, M. and Todorov, T. K.
Isolation of erythromycin A N-oxide and pseudoerythromycin A hemiketal from fermentation broth of *Saccharopolyspora erythraea* by thin-layer and high-performance liquid chromatography 558(1991)265
- Berdecia-Rodriguez, J., see Hendrich, C. E. 577(1992)19
- Berek, D., see Macko, T. 592(1992)109
- Berezkin, V. G., Viktorova, E. N., Gavrichev, V. S., Rang, S. A., Kuningas, K. R. and Meister, A. E.
Gas chromatographic separation of unsaturated polar compounds on inorganic stationary phases containing silver nitrate 552(1991)313
- Berezkina, L. G., Borisova, S. I., Melnikova, S. V. and Souhodolova, V. I.
Reaction-gas chromatography for physico-chemical investigations on aggressive gases 552(1991)319
- Berg-Candolfi, M., Borlakoglu, J. T., Dulery, B., Jehl, F. and Haegele, K. D.
Assessment of the biotransformation of the cardiotoxic agent piroximone by high-performance liquid chromatography and gas chromatography-mass spectrometry 593(1992)1
- Bergamini, E., see Fierabracci, V. 570(1991)285
- Berghem, N., see Borén, K. 588(1991)139
- Bergens, A.
Reductive electrochemical detection in liquid chromatography with a zinc amalgam scrubber column 598(1992)195
- Berger, J. A., see Andriollo, O. 575(1992)301
- Berger, M., Cadet, J., Berube, R., Langlois, R. and Van Lier, J. E.
Reversed-phase high-performance liquid chromatography-thermospray mass spectrometry of radiation-induced decomposition products of thymine and thymidine 593(1992)133
- Berger, M. R., see Gebelein, M. 577(1992)146
- Berger, R., see Tröltzsch, C. 594(1992)111
- Berger, T. A. and Deye, J. F.
Correlation between column surface area and retention of polar solutes in packed-column supercritical fluid chromatography 594(1992)291
- Berger, T. A., see Blumberg, L. M. 596(1992)1
- Bergot, B. J. and Egan, W.
Separation of synthetic phosphorothioate oligodeoxynucleotides from their oxygenated (phosphodiester) defect species by strong-anion-exchange high-performance liquid chromatography 599(1992)35
- Bergqvist, Y., Eckerbom, S., Larsson, H. and Malekzadeh, M.
Reversed-phase liquid chromatographic method for the simultaneous determination of the antimalarial drugs sulfadoxine, pyrimethamine, mefloquine and its major carboxylic metabolite in plasma 571(1991)169
- Bergqvist, Y., see Mount, D. L. 564(1991)181
- Berlin, K. D., see Chen, C. L. 583(1992)274
- Berman, J., see Millington, D. S. 562(1991)47
- Berman, S. S., see Siu, K. W. M. 554(1991)27
- Berna, R. A. and Zacchei, A. G.
High-performance liquid chromatographic method for the determination of a novel leukotriene D₄ antagonist (MK-0571) in biological specimens 563(1991)458
- Bernal-Barragan, H., see Dehnhard, M. 566(1991)101
- Bernert, Jr., J. T., Bell, C. J., McGuffey, J. E. and Waymack, P. P.
Determination of "free" glycerol in human serum reference materials by isotope-dilution gas chromatography-mass spectrometry 578(1992)1

- Berrisford, R. G., see Clark, B. J. 553(1991)383
- Berry, M. J. and Davies, J.
Use of antibody fragments in immunoaffinity chromatography. Comparison of FV fragments, VH fragments and paralog peptides 597(1992)239
- Berry, M. J., Davies, J., Smith, C. G. and Smith, I.
Immobilization of Fv antibody fragments on porous silica and their utility in affinity chromatography 587(1991)161
- Berthod, A., Borgerding, M. F. and Hinze, W. L.
Investigation of the causes of reduced efficiency in micellar liquid chromatography 556(1991)263
- Berti, F., see Bagnati, R. 564(1991)493
- Bertin, P., see Payan, E. 566(1991)9
- Bertini, J., see Triolo, A. 568(1991)281
- Bertino, J. R., see Smith-Rogers, J. A. 566(1991)147
- Bertolasi, V., see Pietrogrande, M. C. 592(1992)65
- Bertolino, A., see Belliardo, F. 553(1991)41
- Bertram, D., see Boeden, H.-F. 552(1991)389
- Bertrand, M. J., see Carrier, A. 591(1992)129
- Bertrand, M. J., see Gagné, J.-P. 554(1991)47
- Bertrand, M. J., see Gagné, J.-P. 554(1991)61
- Bertrand, M. J., see Gagné, J.-P. 554(1991)293
- Bertrand, O., see Kong Sing, Y. L. 558(1991)43
- Bertrand, O., see Kong Sing, Y. L. 598(1992)181
- Berube, R., see Berger, M. 593(1992)133
- Beschke, K., see Greischel, A. 568(1991)246
- Besse, S., see Benoit, E. 583(1992)167
- Besson, C., see Gonzalés, T. 584(1992)101
- Bettendorff, L.
Application of high-performance liquid chromatography to the study of thiamine metabolism and in particular thiamine triphosphatase 566(1991)397
- Betto, P., Ciranni-Signoretti, E. and Di Fava, R.
Determination of cimetidine and related impurities in pharmaceutical formulations by high-performance liquid chromatography 586(1991)149
- Betto, P., Gabriele, R. and Galeffi, C.
Determination of the norlignan glucosides of Hypoxidaceae by high-performance liquid chromatography 594(1992)131
- Betto, P., Gianbenedetti, M., Ponti, F., Ferretti, R., Settini, G., Gargiulo, M. and Lorenzini, R.
Application of a high-performance liquid chromatography coulometric method for the estimation of mebendazole and its metabolites in human sera 563(1991)115
- Betto, P., Meneguz, A., Ricciarello, G. and Pichini, S.
Simultaneous high-performance liquid chromatographic analysis of buspirone and its metabolite 1-(2-pyrimidinyl)-piperazine in plasma using electrochemical detection 575(1992)117
- Betto, P., Ricciarello, G., Pichini, S., Dello Strologo, L. and Rizzoni, G.
High-performance liquid chromatography-electrochemical detection of 3-methylhistidine in human urine 584(1992)256
- Betto, P., see Zuccaro, P. 567(1991)485
- Betts, T. J.
Study of the response of three liquid crystals as stationary phases for the gas chromatography of some cyclic monoterpene volatile oil constituents with short retention times 587(1991)343
- Betts, T. J.
Comparison of four liquid crystal stationary phases used above and below their melting point temperatures for the gas chromatography of some volatile oil constituents 588(1991)231
- Betts, T. J.
Cholesteryl acetate as a stationary phase for the gas chromatography of some volatile oil constituents 600(1992)337
- Betz, W. R. and Lambiase, S. J.
Dynamic gas-solid chromatographic techniques for characterizing carbon molecular sieves 556(1991)433
- Bevan, D. R., see Ducharme, J. 573(1992)79
- Beynon, R. J., see Leyland, D. M. 581(1992)179
- Bézar, J. A., see Semporé, B. G. 557(1991)227
- Bézar, J. A., see Semporé, B. G. 596(1992)185
- Bhat, R., see Chari, G. 571(1991)263
- Bhatti, A. G., see Khuhawar, M. Y. 558(1991)187
- Bhatti, A. R., see C.-Mendoza, C. E. 580(1992)355
- Bhatti, M. M. and Foster, R. T.
Stereospecific high-performance liquid chromatographic assay of metoprolol 579(1992)361
- Bhatti, T., see C.-Mendoza, C. E. 580(1992)355
- Bhaumik, K.
K562 cells: a source for embryonic globin chains 571(1991)37
- Bi, H., Cooper, S. F. and Côté, M. G.
Determination and identification of amiloride in human urine by high-performance liquid chromatography and gas chromatography-mass spectrometry 582(1992)93
- Bi, H., see Massé, R. 562(1991)323
- Bialek, K., see Michalczyk, L. 596(1992)294
- Bianchi, A. P., Varney, M. S. and Phillips, J.
Analysis of industrial solvent mixtures in water using a miniature purge-and-trap device with thermal desorption and capillary gas chromatography-mass spectrometry 557(1991)429
- Bianchi, E., see Pessi, A. 557(1991)307
- Bianchi, G. P., see Grossi, G. 593(1992)217
- Bianchi-Bosisio, A., D'Agrosa, F., Gaboardi, F., Gianazza, E. and Righetti, P. G.
Sodium dodecyl sulphate electrophoresis of urinary proteins (Review) 569(1991)243
- Bianchi-Bosisio, A., see Righetti, P. G. 569(1991)197
- Biarez, O., see Tod, M. 575(1992)171
- Bidgood, A. M., see Musson, D. G. 565(1991)89
- Biersack, H., see Boege, F. 587(1991)3
- Biggs, W. B., see Jinno, K. 557(1991)459
- Biggs, W. R., see Jinno, K. 594(1992)105
- Billion-Rey, F., see Guilluy, R. 562(1991)341
- Biondi, P. A., Guidotti, L., Negri, A. and Secchi, C.
High-performance liquid chromatographic determination of D-amino acid oxidase activity 566(1991)377
- Birarelli, M., see Carducci, C. 553(1991)149

- Bird, R., see Evans, M. B. 552(1991)169
- Birkenmeir, G., see Goubran-Botros, H. 597(1992)357
- Birnbaum, S. and Nilsson, K. G. I.
Dehydrogenase-silica as a stationary phase for the separation of alcohols and ketones 587(1991)268
- Bishop, J. F., see Webster, L. K. 565(1991)482
- Bissbort, S., see Ubbink, J. B. 565(1991)441
- Bissbort, S. H., see Ubbink, J. B. 566(1991)369
- Bisse, E. and Wieland, H.
Coupling of *m*-aminophenylboronic acid to *s*-triazine-activated Sephacryl: use in the affinity chromatography of glycosylated hemoglobins 575(1992)223
- Bisset, N. G., see De, B. 587(1991)318
- Bitteur, S. M., see Voirin, S. G. 590(1992)313
- Björkman, S., see Eriksson, T. 582(1992)211
- Blaas, W., see Jiao, Y. 595(1992)364
- Blachier, S., see Arnaud, J. 572(1991)103
- Black, R. M., Clarke, R. J. and Read, R. W.
Analysis of 1,1'-sulphonylbis[2-(methylsulphonyl)ethane] and 1-methylsulphonyl-2-[2-(methylthio)ethylsulphonyl]ethane, metabolites of sulphur mustard, in urine using gas chromatography-mass spectrometry 558(1991)405
- Black, R. M. and Read, R. W.
Methods for the analysis of thiodiglycol sulphoxide, a metabolite of sulphur mustard, in urine using gas chromatography-mass spectrometry 558(1991)393
- Blackwell, J. A. and Carr, P. W.
Ion- and ligand-exchange chromatography of proteins using porous zirconium oxide supports in organic and inorganic Lewis base eluents 596(1992)27
- Blagbrough, I. S., Daykin, M. M., Doherty, M., Patrick, M. and Shaw, P. N.
High-performance liquid chromatographic determination of naproxen, ibuprofen and diclofenac in plasma and synovial fluid in man 578(1992)251
- Blair, I. A., see Edeki, T. 577(1992)190
- Blake, T. J. A., see Boppana, V. K. 593(1992)29
- Blanchard, J., see Ugwu, S. O. 584(1992)175
- Blanchot, G., see Roux, A. 570(1991)453
- Blanco, C. G., see Guillén, M. D. 591(1992)287
- Blanco, R., see Lin, S. 557(1991)369
- Blank, C. L., see Ikarashi, Y. 575(1992)29
- Blaschke, G., see Fieger, H. 575(1992)255
- Blaschke, G., see Hollenhorst, T. 585(1991)329
- Blaschke, G., see Radler, S. 567(1991)229
- Blaschke, G., see Westhoff, F. 578(1992)265
- Blay, P., see Pleasance, S. 558(1991)155
- Blažek, J., see Černý, J. 600(1992)243
- Bleha, M., see Tennikova, T. B. 555(1991)97
- Bleiberg, B., Steinberg, J. J., Katz, S. D., Wexler, J. and LeJemtel, T.
Determination of plasma lactic acid concentration and specific activity using high-performance liquid chromatography 568(1991)301
- Bleyzac, N., see Bouliou, R. 567(1991)480
- Bleyzac, N., see Bouliou, R. 571(1991)331
- Blo, G., Remelli, M., Pedrielli, F., Balconi, L., Sigon, F. and Dondi, F.
Peak-shape analysis and noise evaluation in suppressed ion chromatography for ultra-trace ion analysis 556(1991)249
- Blo, G., see Accorsi, C. A. 555(1991)65
- Bloemendal, M., see Özçimder, M. 570(1991)19
- Bloemhof, H., Van Dijk, K. N., De Graaf, S. S. N., Vendrig, D. E. M. M. and Uges, D. R. A.
Sensitive method for the determination of vincristine in human serum by high-performance liquid chromatography after on-line column-extraction 572(1991)171
- Blomberg, L. G., see Andersson, P. E. 595(1992)301
- Blomberg, L. G., see Demirbükler, M. 600(1992)358
- Blomqvist, A., see Lindberg, C. 554(1991)215
- Bloom, J. W.
Reversed-phase high-performance liquid chromatographic tryptic digest peptide map comparisons of monoclonal antibodies to human tumor necrosis factor 574(1992)219
- Blumauerová, M., see Beran, M. 558(1991)265
- Blumberg, L. M. and Berger, T. A.
Variance of a zone migrating in a non-uniform time-invariant linear medium 596(1992)1
- Blumer, J. L., see Allen, E. 578(1992)175
- Blumer, J. L., see Ogata, T. 575(1992)39
- Boaretto, A., see Pastore, P. 591(1992)219
- Boček, P.
Analytical biotechnology: Capillary electrophoresis and chromatography (edited by Cs. Horváth and J. N. Nikely) (Book Review) 585(1991)378
- Boček, P., see Klepárník, K. 569(1991)3
- Bochner, F., see Hayball, P. J. 570(1991)446
- Bockbrader, H. N., see Janiczek, N. 571(1991)179
- Böcker, R. H., Peter, R., Machbert, G. and Bauer, W.
Identification and determination of the two principal metabolites of minocycline in humans 568(1991)363
- Boddy, A. V. and Idle, J. R.
Combined thin-layer chromatography-photography-densitometry for the quantification of ifosfamide and its principal metabolites in urine, cerebrospinal fluid and plasma 575(1992)137
- Boeden, H.-F., Pommerening, K., Becker, M., Rupprich, C., Heltzauer, M., Loth, F., Müller, R. and Bertram, D.
Bead cellulose derivatives as supports for immobilization and chromatographic purification of proteins 552(1991)389
- Boeden, H.-F., see Baeseler, M. 589(1992)93
- Boege, F., Gieseler, F., Biersack, H. and Clark, M.
Use of anion-exchange chromatography and chromatofocusing to reveal the structural and functional heterogeneity of topoisomerase II in a HL-60 cell line resistant to multi-drug treatment 587(1991)3
- Bogoni, P., see Zonta, F. 594(1992)137
- Bogusz, M.
High-performance liquid chromatographic determination of morphine, morphine-3-glucuronide, morphine-6-glucuronide and codeine in biological samples using multi-wavelength forward optical detection: a reply 579(1992)189
- Bohm, B. A., see Koupai-Abyazani, M. R. 594(1992)117
- Böhmer, T., see Tallaksen, C. M. E. 564(1991)127
- Bohrer, A., see Turk, J. 575(1992)183
- Bohuon, G., see Delage, E. 555(1991)125

- Boison, J. O., Korsrud, G. O., MacNeil, J. D., Keng, L. and Papich, M.
Determination of penicillin G in bovine plasma by high-performance liquid chromatography after pre-column derivatization 576(1992)315
- Boje, K. M., see Miller, R. 578(1992)103
- Bojesen, G., see Justesen, U. 562(1991)59
- Bolelli, G. F., see Pellegatti, M. 573(1992)105
- Boltralik, J. J., see McCue, B. A. 565(1991)255
- Bonanni, L., see Menichini, E. 555(1991)211
- Bonati, M., see Celardo, A. 568(1991)407
- Bonekamp, A., see Lendrath, G. 588(1991)303
- Bonet Domingo, E., Medina Hernández, M. J., Ramis Ramos, G. and García Alvarez-Coque, M. C.
High-performance liquid chromatographic determination of diuretics in urine by micellar liquid chromatography 582(1992)189
- Boniface, P. J., Martin, I. C., Nolan, S. L. and Tan, S. T.
Development of a method for the determination of zopiclone in whole blood 584(1992)199
- Bonifaci, L., Frezzotti, D., Cavalca, G., Malaguti, E. and Ravanetti, G. P.
Analysis of ϵ -caprolactam and its cyclic oligomers by high-performance liquid chromatography 585(1991)333
- Bonn, G., see Seitz, U. 559(1991)499
- Bonn, G. K., see Haidacher, D. 591(1992)351
- Bonn, G. K., see Huber, C. G. 599(1992)113
- Bonnaire, Y., see Benoit, E. 583(1992)167
- Bonnaire, Y., see Jaussaud, P. 573(1992)136
- Bontempelli, G., see Tagliaro, F. 563(1991)299
- Boobis, A. R., see Jones, H. A. 570(1991)361
- Boobis, A. R., see Murray, S. 568(1991)341
- Boomsma, F., Alberts, G., Van der Hoorn, F. A. J., Man in 't Veld, A. J. and Schalekamp, M. A. D. H.
Simultaneous determination of free catecholamines and epinine and estimation of total epinine and dopamine in plasma and urine by high-performance liquid chromatography with fluorimetric detection 574(1992)109
- Boomsma, F., see Alberts, G. 583(1992)236
- Boomsma, F., see Van Der Hoorn, F. A. J. 563(1991)348
- Boos, K.-S., Lintelmann, J. and Kettrup, A.
Coupled-column high-performance liquid chromatographic method for the determination of 1-hydroxypyrene in urine of subjects exposed to polycyclic aromatic hydrocarbons 600(1992)189
- Bootsma, H. J., Van den Berg, C. W. and Van Dijk, H.
Rapid isolation of human complement component C9 to verify the specificity of a haemolytic C9 microassay 591(1992)187
- Bopp, R. J., see Anliker, S. L. 573(1992)141
- Boppana, V. K., Geschwindt, L., Cyronak, M. J. and Rhodes, G.
Determination of the enantiomers of fenoldopam in human plasma by reversed-phase high-performance liquid chromatography after chiral derivatization 592(1992)317
- Boppana, V. K., Simpson, R. C., Anderson, K., Miller-Stein, C., Blake, T. J. A., Hwang, B. Y.-H. and Rhodes, G. R.
High-performance liquid chromatographic determination of monohydroxy compounds by a combination of pre-column derivatization and post-column reaction detection 593(1992)29
- Borák, J., see Macka, M. 586(1991)291
- Borén, K., Larsson, M., Bergenhem, N. and Carlsson, U.
Rapid ion-exchange chromatography for preparative separation of proteins. IV. Application to bovine carbonic anhydrase III from skeletal muscle 588(1991)139
- Boreus, L. O., see Beck, O. 570(1991)198
- Borg, S., see Helander, A. 579(1992)340
- Borgerding, M. F., see Berthod, A. 556(1991)263
- Bories, C., Cardot, P. J. P., Abramowski, V., Poüs, C., Merino-Dugay, A., Baron, B. and Mougeot, G.
Elution mode of *Pneumocystis carinii* cysts in gravitational field-flow fractionation 579(1992)143
- Bories, C., see Merino, A. 572(1991)291
- Borisova, S. I., see Berezkina, L. G. 552(1991)319
- Borlakoglu, J. T., see Berg-Candolfi, M. 593(1992)1
- Borner, E., see Borner, K. 579(1992)285
- Borner, K., Borner, E. and Lode, H.
Determination of sparfloxacin in serum and urine by high-performance liquid chromatography 579(1992)285
- Börnsen, K. O., see Steiner, V. 586(1991)43
- Borrego, F., see Castillo, J. 555(1991)285
- Borrull, F., see Calull, M. 589(1992)151
- Borrull, F., see Calull, M. 590(1992)215
- Bortolomeazzi, R., see Moret, S. 591(1992)175
- Boschetti, E., see Santambien, P. 597(1992)315
- Bosman, J., see Witte, D. T. 553(1991)365
- Bosman, J., see Witte, D. T. 558(1991)333
- Bostrom, B. C., see Erdmann, G. R. 571(1991)149
- Böswart, J., Kostiuk, P., Vymlátíl, J., Schmidt, T., Pacáková, V. and Štulík, K.
High-performance liquid chromatographic determination of cholesteryl esters in the blood of obese children 571(1991)19
- Both, D. A. and Jemal, M.
Stereoisomeric purity determination of captopril by capillary gas chromatography 558(1991)257
- Both, D. A. and Jemal, M.
Determination of parts per million levels of trifluoroacetic acid in ceronapril bulk substance by headspace capillary gas chromatography 596(1992)85
- Both, D. A., Ribick, M. and Jemal, M.
Determination of methylboronic acid in teboroxime by capillary gas chromatography 585(1991)348
- Both-Miedema, R., see Jansen, E. H. J. M. 553(1991)123
- Botsoglou, N. A.
High-performance liquid chromatographic method for the determination of free gossypol in chicken liver 587(1991)333
- Bottaro, S., see Cracco, A. T. 574(1992)156
- Bottiglieri, T., see Hyland, K. 579(1992)55
- Boucherat, M., see Daumas, L. 570(1991)89
- Bouguerra, M. L., see Driss, M. R. 552(1991)213

- Bouguerra, M. L., see Sabbah, S. 552(1991)223
- Boulieu, R., Bleyzac, N. and Ferry, S.
High-performance liquid chromatographic determination of ganciclovir in plasma 567(1991)480
- Boulieu, R., Bleyzac, N. and Ferry, S.
Modified high-performance liquid chromatographic method for the determination of ganciclovir in plasma from patients with severe renal impairment 571(1991)331
- Bouquillon, A. I., Freeman, D. and Moulin, D. E.
Simultaneous solid-phase extraction and chromatographic analysis of morphine and hydromorphone in plasma by high-performance liquid chromatography with electrochemical detection 577(1992)354
- Bourell, J., see Frenz, J. 557(1991)289
- Bourguignon, B. and Massart, D. L.
Simultaneous optimization of several chromatographic performance goals using Derringer's desirability function 586(1991)11
- Bourguignon, B., Vankeerberghen, P. and Massart, D. L.
CRISEBOOK, a Hypermedia version of an expert system for the selection of optimization criteria in high-performance liquid chromatography 592(1992)51
- Boutagy, J. and Shenfield, G. M.
Simplified procedure for the determination of sotalol in plasma by high-performance liquid chromatography 565(1991)523
- Boutin, J. A., Ernoult, A.-P., Ferry, G., Genton, A. and Alpert, A. J.
Use of hydrophilic interaction chromatography for the study of tyrosine protein kinase specificity 583(1992)137
- Boutin-Muma, B., see Cohen, H. 595(1992)143
- Bouvier, O., see Cociglio, M. 572(1991)269
- Bouygues-De Ferran, A. M., Pham-Huy, C., Postaire, M. and Hamon, M.
Determination of trace amounts of fluoride in raw materials for pharmaceuticals by gas-liquid chromatography 585(1991)289
- Bowers, G. D., Highton, D. M., Manchec, G. R., Oxford, J. and Saynor, D. A.
Thermospray liquid chromatography-mass spectrometry for the characterisation of sulphate ester conjugates 554(1991)175
- Bradshaw, J. S., see Johnson, D. F. 594(1992)283
- Braggio, S., see Pellegatti, M. 573(1992)105
- Brambilla, P., see Magni, F. 573(1992)127
- Branch, S. K., see Warner, C. D. 562(1991)625
- Brandl, G., Kastner, F., Mannschreck, A., Nölting, B., Andert, K. and Wetzl, R.
Chiroptical detection during liquid chromatography. III. Non-stop acquisition of circular dichroism spectra during liquid chromatography 586(1991)249
- Brandolo, G., see Belliaro, F. 553(1991)41
- Branham, K. E., see Meng, Q. C. 579(1992)63
- Braquet, P., see Watanabe, M. 597(1992)309
- Brazier, J. L., see Desage, M. 571(1991)250
- Brazier, J. L., see Guilluy, R. 562(1991)341
- Brazier, M., see Kamel, S. 574(1992)255
- Brazzell, R. K., see McCue, B. A. 565(1991)255
- Brearley, G. M., see Scott, D. A. 573(1992)309
- Breckenridge, W. C., see Kuksis, A. 564(1991)11
- Breckenridge, W. C., see Kuksis, A. 579(1992)13
- Breda, M., Pianezzola, E. and Strolin Benedetti, M.
Determination of 3'-deamino-3'-[2(S)-methoxy-4-morpholinyl]doxorubicin, a new morpholinyl anthracycline, in plasma by high-performance liquid chromatography with fluorescence detection 578(1992)309
- Brega, A., Villa, P., Quadrini, G., Quadri, A. and Lucarelli, C.
High-performance liquid chromatographic determination of acetone in blood and urine in the clinical diagnostic laboratory 553(1991)249
- Breimer, D. D., see Roosemalen, M. C. M. 565(1991)516
- Breimer, D. D., see Venizelos, V. 573(1992)259
- Breitenbach, M., see Rizzi, A. M. 582(1992)35
- Bremner, J. B., see Richardson, D. E. 595(1992)155
- Brennan, J. M., see LaCreta, F. P. 571(1991)271
- Brennan, S. O., see Haynes, P. A. 581(1992)187
- Brennan, T. M., see Huang, X. C. 600(1992)289
- Brenneisen, R., see Helmlin, H.-J. 593(1992)87
- Brenneisen, R., see Mathys, K. 593(1992)79
- Bresson, M., see Rop, P. P. 573(1992)87
- Brewer, Jr., H. B., see Shibusawa, Y. 596(1992)118
- Brewster, J. D., Lightfield, A. R. and Barford, R. A.
Evaluation of restricted access media for high-performance liquid chromatographic analysis of sulfonamide antibiotic residues in bovine serum 598(1992)23
- Brewster, J. D. and Piotrowski, E. G.
Rapid electroanalytical clean-up of biological samples for high-performance liquid chromatography 585(1991)213
- Brier, G., see Kovach, P. M. 567(1991)129
- Brilis, G. M., see Brumley, W. C. 558(1991)223
- Brindle, I. D., see Marvin, C. H. 555(1991)147
- Bringmann, G., Hesselmann, C. and Feineis, D.
Endogenous alkaloids in man. XI. Analysis of glyoxylate-derived 1,3-thiazolidines by ion-pair-assisted reversed-phase chromatography 595(1992)351
- Brinkman, U. A. T., see Barceló, D. 553(1991)311
- Brinkman, U. A. T., see Debets, A. J. J. 600(1992)163
- Brinkman, U. A. T., see Farjam, A. 589(1992)141
- Brinkman, U. A. T., see Goosens, E. C. 552(1991)489
- Brinkman, U. A. T., see Kwakman, P. J. M. 553(1991)345
- Brinkman, U. A. T., see Somsen, G. W. 552(1991)635
- Brinkman, U. A. T., see Steijger, O. M. 557(1991)13
- Brinkman, U. A. T., see Van de Nesse, R. J. 552(1991)613
- Brinkman, U. A. T., see Van der Horst, F. A. L. 567(1991)161
- Briskman, V. A., see Gelfi, C. 598(1992)277
- Brisson, A. M., see Girault, J. 564(1991)43
- Brisson, J., see Hopkins, N. K. 577(1992)87
- Briza, P., see Rizzi, A. M. 582(1992)35
- Břízová, K., Králová, B., Demnerová, K. and Vinš, I.
Isolation and characterization of α -glucosidase from *Aspergillus niger* 593(1992)125
- Broccoli, G., see Dal Bo, L. 573(1992)158
- Brockmüller, J., see Staffeldt, B. 571(1991)133

- Brocks, D. R., Pasutto, F. M. and Jamali, F.
Analytical and semi-preparative high-performance liquid chromatographic separation and assay of hydroxychloroquine enantiomers 581(1992)83
- Brocks, D. R., see Liang, W. T. C. 577(1992)317
- Brocks, D. R., see Wright, M. R. 583(1992)259
- Bronz, I. and Olsen, I.
Intra-injector methylation of free fatty acids from aerobically and anaerobically cultured *Actinobacillus actinomycetemcomitans* and *Haemophilus aphrophilus* 576(1992)328
- Bronz, I. and Olsen, I.
Intra-injector formation of methyl esters from phenoxy acid pesticides 598(1992)309
- Bronner, W. E. and Xu, A. S.
Gas chromatographic-mass spectrometric methods of analysis for detection of 11-nor- Δ^9 -tetrahydrocannabinol-9-carboxylic acid in biological matrices (Review) 580(1992)63
- Brooks, M. A., see Tsai, E. W. 596(1992)217
- Brookshier, L. A., see Hoyer, G. L. 572(1991)159
- Broquaire, M., see Becue, T. 557(1991)489
- Broquet, C., see Watanabe, M. 597(1992)309
- Brørs, O., see Holthe, M. R. 579(1992)376
- Brossard, S., Lafosse, M. and Dreux, M.
Comparison of ethoxylated alcohols and polyethylene glycols by high-performance liquid chromatography and supercritical fluid chromatography using evaporative light-scattering detection 591(1992)149
- Brostedt, P., see Silberring, J. 554(1991)83
- Brown, A. R., James, C. D., Welch, R. M. and Harrelson, J. C.
Stereoselective high-performance liquid chromatographic assay with fluorometric detection for the isomers of mivacurium in human plasma 578(1992)302
- Brown, J., see Nijjar, M. S. 568(1991)393
- Brown, L. L., see Donahue, E. P. 571(1991)29
- Brown, L. W., Hundt, H. K. L. and Swart, K. J.
Automated high-performance liquid chromatographic method for the determination of mianserin in plasma using electrochemical detection 582(1992)268
- Brown, M., see Robertson, A. M. 588(1991)15
- Brown, M. A., see Kim, I. S. 589(1991)177
- Brown, N. D., Phillips, L. R., Leader, H. and Chiang, P. K.
Isolation and identification of β -hydroxyethylapropen: a urinary metabolite of aprocphen in rats 563(1991)466
- Brown, P. R., see Amari, J. V. 590(1992)153
- Brown, P. R., see Roman, M. 592(1992)3
- Brown, P. R., see Weston, A. 593(1992)289
- Brown, R. R., see Antoniewicz, S. M. 573(1992)93
- Brown, Jr., F. and Heath, P. C.
Separation and purification of azetidiny methyl sulfonates using preparative high-performance liquid chromatography 595(1992)355
- Brown-Thomas, J. M., Christensen, R. G., Rieger, R., Malone, W. and May, W. E.
Determination of glycyrrhetic acid in human plasma by high-performance liquid chromatography 568(1991)232
- Browne, S. P., Tebbett, I. R., Moore, C. M., Dusick, A., Covert, R. and Yee, G. T.
Analysis of meconium for cocaine in neonates 575(1992)158
- Browner, M. F., see Luong, C. B. H. 584(1992)77
- Brownlee, M., see Steinberg, J. J. 574(1992)41
- Brownlee, R. G., see Schwartz, H. E. 559(1991)267
- Brownrigg, C. M., see Brumley, W. C. 558(1991)223
- Brownsill, R., Wallace, D., Taylor, A. and Campbell, B.
Study of human urinary metabolism of fenfluramine using gas chromatography-mass spectrometry 562(1991)267
- Bruchet, A., Legrand, M. F., Arpino, P. and Dilettato, D.
Recent methods for the determination of volatile and non-volatile organic compounds in natural and purified drinking water 562(1991)469
- Brückner, H. and Gah, C.
High-performance liquid chromatographic separation of DL-amino acids derivatized with chiral variants of Sanger's reagent 555(1991)81
- Brügger, R. and Arm, H.
Investigations of the influence of silanol groups on the separation of enantiomers by liquid and supercritical fluid chromatography 592(1992)309
- Brügger, R., Krähenbühl, P., Marti, A., Straub, R. and Arm, H.
Chiral π -donor stationary phases with (*R*)-*N*-pivaloylnaphthylethylamide groups for direct enantiomer separation by gas, liquid and supercritical fluid chromatography 557(1991)163
- Bruin, G. J. M., Stegeman, G., Van Asten, A. C., Xu, X., Kraak, J. C. and Poppe, H.
Optimization and evaluation of the performance of arrangements for UV detection in high-resolution separations using fused-silica capillaries 559(1991)163
- Bruins, A. P.
Liquid chromatography-mass spectrometry with ionspray and electrospray interfaces in pharmaceutical and biomedical research 554(1991)39
- Brumley, W. C., Brownrigg, C. M. and Brilis, G. M.
Characterization of nitrogen-containing aromatic compounds in soil and sediment by capillary gas chromatography-mass spectrometry after fractionation 558(1991)223
- Brune, K., see Geisslinger, G. 568(1991)165
- Brune, K., see Geisslinger, G. 573(1992)163
- Brune, K., see Schmidt, N. 583(1992)195
- Brunmark, P., Persson, P. and Skarping, G.
Determination of 4,4'-methylenedianiline in hydrolysed human urine by micro liquid chromatography with ultraviolet detection 579(1992)350
- Brunmark, P., Skarping, G. and Schütz, A.
Determination of methylmercury in human blood using capillary gas chromatography and selected-ion monitoring 573(1992)35
- Brunner, L. A. and Luders, R. C.
Determination of a potential anxiolytic drug (CGS 20625) in human plasma by high-performance liquid chromatography 568(1991)487

- Bruno, A., see Pietta, P. 553(1991)223
 Bruno, A., see Pietta, P. 593(1992)165
 Bruno, G., see Lucarelli, C. 573(1992)150
 Bruno, G., see Mancinelli, A. 553(1991)81
 Bruno, R., see Vergniol, J. C. 582(1992)273
 Bruns, A., Waldhoff, H., Wilsch-Irrgang, A. and Winkle, W.
 Automated high-performance liquid chromatographic and
 size-exclusion chromatographic sample preparation by
 means of a robotic workstation 592(1992)249
 Buchanan, J. S., see Gurley, L. R. 559(1991)411
 Buchberger, W., Haddad, P. R. and Alexander, P. W.
 Separation of metal complexes of
 ethylenediaminetetraacetic acid in environmental water
 samples by ion chromatography with UV and
 potentiometric detection 558(1991)181
 Bucheli, F., see Wyss, R. 576(1992)111
 Bucheli, F., see Wyss, R. 593(1992)55
 Buck, C. F., Mayewski, P. A., Spencer, M. J., Whitlow, S.,
 Twickler, M. S. and Barrett, D.
 Determination of major ions in snow and ice cores by ion
 chromatography 594(1992)225
 Buckel, W., see Michel, C. 587(1991)93
 Buckley, D. J., see Burse, V. W. 566(1991)117
 Budzinski, H., Garrigues, P. and Bellocq, J.
 Gas chromatographic retention behaviour of
 dibenzothiophene derivatives on a smectic liquid
 crystalline polysiloxane stationary phase 590(1992)297
 Budzinski, H., see Sellier, N. 557(1991)451
 Büeler, M. R., see Wiederkehr, F. 566(1991)77
 Buinevicius, Z., see Sarin, M. 563(1991)129
 Bullock, J., see Snider, J. 599(1992)141
 Bult, A., see Kaijser, G. P. 571(1991)121
 Bult, A., see Kaijser, G. P. 583(1992)175
 Bureiko, A. S., see Marinichev, A. N. 600(1992)251
 Burford, M. D., see Langenfeld, J. J. 594(1992)297
 Burford, M. D., see Smith, R. M. 600(1992)175
 Burger, B. V., Le Roux, M., Munro, Z. M. and Wilken, M. E.
 Production and use of capillary traps for headspace gas
 chromatography of airborne volatile organic compounds
 552(1991)137
 Burger, D. M., Rosing, H., Van Gijn, R., Meenhorst, P. L.,
 Van Tellingen, O. and Beijnen, J. H.
 Determination of stavudine, a new antiretroviral agent, in
 human plasma by reversed-phase high-performance liquid
 chromatography with ultraviolet detection 584(1992)239
 Burghouts, J. T. M., see Van Tellingen, O. 573(1992)328
 Burgi, D. S., see Chien, R.-L. 559(1991)141
 Burgi, D. S., see Chien, R.-L. 559(1991)153
 Burgi, D. S., see Salomon, K. 559(1991)69
 Burke, C. J., see Lehman, E. D. 574(1992)225
 Burke, J. A., see Pirkle, W. H. 598(1992)159
 Burke, J. F., see Warner, C. D. 562(1991)625
 Burke, III, J. A., see Pirkle, W. H. 557(1991)173
 Burke, III, J. A., see Pirkle, W. H. 598(1992)1
 Burkow, I. C. and Henderson, R. J.
 Isolation and quantification of polymers from autoxidized
 fish oils by high-performance size-exclusion
 chromatography with an evaporative mass detector
 552(1991)501
 Burns, K. I., see Elchuk, S. 558(1991)197
 Burri, B. J., Kutnink, M. A. and Neidlinger, T. R.
 Assay of human transthyretin-bound holo-retinol-binding
 protein with reversed-phase high-performance liquid
 chromatography 567(1991)369
 Burse, V. W., Korver, M. P., McClure, P. C., Holler, J. S.,
 Fast, D. M., Head, S. L., Miller, D. T., Buckley, D. J.,
 Nassif, J. and Timperi, R. J.
 Problems associated with interferences in the analysis of
 serum for polychlorinated biphenyls 566(1991)117
 Burt, H. M., see Aoyama, R. G. 588(1991)181
 Burton, F. W., see Gadde, R. R. 593(1992)41
 Burton, S. C., Haggarty, N. W. and Harding, D. R. K.
 Efficient substitution of 1,1'-carbonyldiimidazole activated
 cellulose and Sepharose matrices with amino acyl spacer
 arms 587(1991)271
 Burton, S. D., Hutchins, J. E., Fredericksen, T. L., Ricks, C.
 and Tyczkowski, J. K.
 High-performance liquid chromatographic method for the
 determination of spectinomycin in turkey plasma
 571(1991)209
 Busby, S., see Downham, M. 584(1992)59
 Busch, M. P., see Schwartz, H. E. 559(1991)267
 Buser, H.-R., see Tóth, M. 598(1992)303
 Bushway, R. J., Hurst, H. L., Kugabalasooriar, J. and
 Perkins, L. B.
 Determination of carbendazim in blueberries by reversed-
 phase high-performance liquid chromatography
 587(1991)321
 Busini, S., Bavazzano, P. and Occhiato, E.
 High-performance liquid chromatographic determination
 of 3,5-dimethylhippuric acid in the occupational exposure
 to trimethylbenzenes 577(1992)180
 Busson, R., see Naidong, W. 586(1991)67
 Buszewski, B., Schmid, J., Albert, K. and Bayer, E.
 Chemically bonded phases for the reversed-phase high-
 performance liquid chromatographic separation of basic
 substances 552(1991)415
 Bütikofer, P., see Kuypers, F. A. 562(1991)191
 Butts, E. T., see Levison, P. R. 590(1992)49
 Bystrický, S., see Strašák, M. 585(1991)337
- C**
- Cabras, P., Melis, M., Spanedda, L. and Tuberoso, C.
 High-performance liquid chromatographic determination
 of dinitroaniline herbicides in soil and water
 585(1991)164

- Caccia, S., Confalonieri, S., Guiso, G., Celeste, S. and Marini, P. P.
Liquid chromatography of the potential memory-enhancing agent CL 275,838 and its main metabolites, using a post-column photochemical reactor and fluorimetric detection 581(1992)109
- Caccia, S., see Guiso, G. 572(1991)346
- Caccia, S., see Sarati, S. 563(1991)323
- Caccialanza, G., see Kitsos, M. 553(1991)1
- Cachet, T., Lannoo, P., Paesen, J., Janssen, G. and Hoogmartens, J.
Determination of erythromycin ethylsuccinate by liquid chromatography 600(1992)99
- Cadet, J., see Berger, M. 593(1992)133
- Caffaro, L. E., see Paetsch, P. R. 573(1992)313
- Caixach, J., see Guardiola, A. 562(1991)481
- Cajigas, A., see Steinberg, J. J. 574(1992)41
- Calabró, M., see Castagnetta, L. A. 572(1991)25
- Caldwell, K. D., see Li, J. 555(1991)260
- Caldwell, W. M., see Griest, W. H. 600(1992)273
- Caliceti, P., see Švagrová, I. 563(1991)95
- Callebat, I., see Maisonneuve, A. 583(1992)155
- Calull, M., López, E., Marcé, R. M., Olucha, J. C. and Borrull, F.
Optimization of an ion-exchange high-performance liquid chromatographic method for the determination of carboxylic acids, sugars, glycerol and ethanol in wines 589(1992)151
- Calull, M., Marcé, R. M. and Borrull, F.
Determination of carboxylic acids, sugars, glycerol and ethanol in wine and grape must by ion-exchange high-performance liquid chromatography with refractive index detection 590(1992)215
- Camagna, M., see Tarditi, L. 599(1992)13
- Camamo, M., see Haken, J. K. 595(1992)283
- Cammack, J., Oke, A. and Adams, R. N.
Simultaneous high-performance liquid chromatographic determination of ascorbic acid and dehydroascorbic acid in biological samples 565(1991)529
- Camoin, L., see Dandeu, J.-P. 599(1992)105
- Camoni, I., see Di Muccio, A. 553(1991)305
- Campbell, B., see Brownsill, R. 562(1991)267
- Campbell, J. A., see Harvey, S. D. 587(1991)300
- Campestrini, J., see Rouan, M. C. 573(1992)59
- Campestrini, J., see Rouan, M. C. 573(1992)65
- Campestrini, J., see Rouan, M. C. 577(1992)387
- Campins-Falcó, P., see Herráez-Hernández, R. 582(1992)181
- Campos-Barreda, F., see Pou-Clave, L. 563(1991)211
- Canafax, D. M., see Erdmann, G. R. 571(1991)149
- Cann-Moisán, C., Caroff, J. and Girin, E.
Standardization in the determination of red blood cell polyamines by high-performance liquid chromatography 574(1992)134
- Cannell, G. R., Mortimer, R. H., Maguire, D. J. and Addison, R. S.
Liquid chromatographic analysis of prednisolone, prednisone and their 20-reduced metabolites in perfusion media 563(1991)341
- Cannell, G. R., Williams, J. P., Yap, A. S. and Mortimer, R. H.
Selective liquid chromatographic assay for propylthiouracil in plasma 564(1991)310
- Canobbio, G., see Belotto, N. 572(1991)327
- Cantafora, A. and Masella, R.
Improved determination of individual molecular species of phosphatidylcholine in biological samples by high-performance liquid chromatography with internal standards 593(1992)139
- Cao, X.-L.
Determination of specific retention volumes at 20°C for hydrocarbons on microporous carbons 586(1991)161
- Cao, X. L., Colenutt, B. A. and Sing, K. S. W.
Study of microporous carbons by gas chromatographic determination of heats of physisorption 555(1991)183
- Caprioli, R. M., see Suter, M. J.-F. 553(1991)101
- Cardace, G., see Mancinelli, A. 553(1991)81
- Cardot, P. J. P., Gerota, J. and Martin, M.
Separation of living red blood cells by gravitational field-flow fractionation 568(1991)93
- Cardot, P. J. P., see Bories, C. 579(1992)143
- Cardot, P. J. P., see Merino, A. 572(1991)291
- Cardot, P. J. P., see Merino-Dugay, A. 579(1992)73
- Carducci, C., Moretti, F., Birarelli, M. and Antonozzi, I.
Rapid and sensitive high-performance liquid chromatographic method for the analysis of tryptophan, tyrosine and phenylalanine in biological samples 553(1991)149
- Carelse, O., Chetsanga, C. J. and Mubumbila, M. V.
Purification of common bean chloroplast DNA by DEAE cellulose column chromatography 558(1991)318
- Carenzi, A., see Longo, A. 562(1991)639
- Carleton, B. C., see Erdmann, G. R. 571(1991)149
- Carli, G., see Tagliaro, F. 563(1991)299
- Carlqvist, J., see Lanbeck-Vallén, K. 567(1991)121
- Carlqvist, J., see Lanbeck-Vallén, K. 578(1992)71
- Carlsson, U., see Borén, K. 588(1991)139
- Carlsten, J., see Lönnroth, P. 568(1991)419
- Carlucci, G. and Palumbo, G.
Analytical procedure for the determination of rifloxacin, a new pyridobenzothiazine, in human serum and urine by high-performance liquid chromatography 564(1991)346
- Caroff, J., see Cann-Moisán, C. 574(1992)134
- Caron, J., see Trivier, J. M. 579(1992)269
- Carpené, G., Vettoretti, A., Pedini, F., Rocco, S., Mantero, F. and Opocher, G.
Hypertensive congenital adrenal enzymatic defects detected by high-performance liquid chromatography of corticosteroids 553(1991)201
- Carpenter, A. and Purdy, W. C.
Preparation of heparin-glyceryl controlled-pore glass affinity media for the separation of α - and β -lipoproteins 573(1992)132
- Carpita, N. C., see Gibeau, D. M. 587(1991)284
- Carr, D., see Yang, Y.-B. 590(1992)35
- Carr, M. R., see Reville, A. T. 589(1992)281
- Carr, P. W., see Blackwell, J. A. 596(1992)27
- Carr, P. W., see Lucy, C. A. 556(1991)159

- Carr, P. W., see Park, J. H. 586(1991)1
Carr, P. W., see Schafer, W. A. 587(1991)137
Carr, P. W., see Schafer, W. A. 587(1991)149
Carr, R. A., Foster, R. T., Freitag, D. and Pasutto, F. M.
Stereospecific high-performance liquid chromatographic determination of tocinamide 566(1991)155
Carrai, P., see Colombini, M. P. 592(1992)255
Carrea, G., Pasta, P., Colonna, S. and Gaggero, N.
High-performance liquid chromatographic separation of chiral metallocenic ketones and alcohols 600(1992)320
Carrier, A., Gagné, J.-P. and Bertrand, M. J.
Limits and effects of precolumn addition of thioglycerol in liquid chromatographic-fast atom bombardment mass spectrometric systems 591(1992)129
Carrier, A., see Gagné, J.-P. 554(1991)47
Carrier, A., see Gagné, J.-P. 554(1991)61
Carrier, J., see Chauret, N. 588(1991)281
Carruba, G., see Castagnetta, L. A. 572(1991)25
Carter, S. R., Duke, C. C., Cutler, D. J. and Holder, G. M.
Sensitive stereospecific assay of warfarin in plasma: reversed-phase high-performance liquid chromatographic separation using diastereoisomeric esters of (-)-(1*S*,2*R*,4*R*)-endo-1,4,5,6,7,7-hexachlorobicyclo[2.2.1]hept-5-ene-2-carboxylic acid 574(1992)77
Cartoni, G. P., Castellani, L., Goretti, G., Russo, M. V. and Zacchei, P.
Gas-liquid microcapillary columns precoated with graphitized carbon black 552(1991)197
Caruel, H., Phemius, P., Rigal, L. and Gaset, A.
Chromatographic separation of polyols by ligand exchange. Effects of the ion-exchange resin cross-linking and size 594(1992)125
Caruel, H., Rigal, L. and Gaset, A.
Carbohydrate separation by ligand-exchange liquid chromatography. Correlation between the formation of sugar-cation complexes and the elution order 558(1991)89
Caruso, U., Fowler, B., Erceg, M. and Romano, C.
Determination of very-long-chain fatty acids in plasma by a simplified gas chromatographic-mass spectrometric procedure 562(1991)147
Casais Laiño, C., see Lansens, P. 586(1991)329
Cashman, J. R., see Yang, Z. C. 596(1992)79
Caslavská, J., Gebauer, P. and Thormann, W.
Purification of ovalbumin and lysozyme from a commercial product by recycling isotachopheresis 585(1991)145
Caslavská, J. and Thormann, W.
Continuous separation and purification of proteins by recycling isotachopheresis 594(1992)361
Caslavská, J., see Thormann, W. 589(1992)321
Caslavská, J., see Thormann, W. 593(1992)275
Caspary, W. F., see Stein, J. 576(1992)53
Cassiano, L., see Castagnola, M. 572(1991)51
Cassidy, R. M., see Elchuk, S. 558(1991)197
Cassier, P., see Cherton, J.-C. 566(1991)511
Castagnetta, L. A., Granata, O. M., Lo Casto, M., Calabró, M., Arcuri, F. and Carruba, G.
Simple approach to measure metabolic pathways of steroids in living cells 572(1991)25
Castagnola, M., Cassiano, L., Rabino, R., Rossetti, D. V. and Bassi, F. A.
Peptide mapping through the coupling of capillary electrophoresis and high-performance liquid chromatography: map prediction of the tryptic digest of myoglobin 572(1991)51
Castagnoli, Jr., N., see Igarashi, K. 579(1992)277
Castellani, L., see Cartoni, G. P. 552(1991)197
Castello, G. and D'Amato, G.
Determination of sulphur and nitrogen gases by gas chromatography on polystyrene porous polymer columns 585(1991)93
Castello, G., Vezzani, S. and Gerbino, T. C.
Gas chromatographic separation and automatic identification of complex mixtures of organic solvents in industrial wastes 585(1991)273
Castiglioni, C., see Abbiati, G. 566(1991)445
Castiglioni, C., see Fossati, T. 577(1992)382
Castillo, J., Benavente, O., Borrego, F. and Del Rio, J. A.
Analysis of commercial hesperidin methyl chalcone by high-performance liquid chromatography 555(1991)285
Castledine, J. B., Fell, A. F., Modin, R. and Sellberg, B.
Assessment of chromatographic peak purity by means of multi-wavelength detection and correlation-based algorithms 592(1992)27
Catalano, T., see Kersten, B. S. 588(1991)187
Caturla, M. C. and Cusido, E.
Solid-phase extraction for the high-performance liquid chromatographic determination of indomethacin, suxibuzone, phenylbutazone and oxyphenbutazone in plasma, avoiding degradation of compounds 581(1992)101
Caturla, M. C., Cusido, E. and Westerlund, D.
High-performance liquid chromatography method for the determination of aminoglycosides based on automated pre-column derivatization with *o*-phthalaldehyde 593(1992)69
Caude, M., see Villermet, A. 557(1991)85
Caudill, D., see Rodriguez, P. A. 563(1991)271
Cavalca, G., see Bonifaci, L. 585(1991)333
Cavalleri, A., see Venturelli, E. 582(1992)7
Cavallini, P., see Dalla Libera, L. 566(1991)427
Cavenaghi, L., see Riva, E. 553(1991)35
Cazaussus, A., see Sellier, N. 557(1991)451
Cebrian Perez, J. A., see Ortin, A. 558(1991)357
Cecchini, D. J., see Al-Deen, A. N. 600(1992)229
Cecon, L., see Lo Coco, F. 590(1992)235
Ceconi, O., Ruggieri, S. and Mugnai, G.
Use of N-acetylpsychosine as internal standard for quantitative high-performance liquid chromatographic analysis of glycosphingolipids 555(1991)267
Cech, D., see Jäschke, A. 585(1991)57
Cech, D., see Jäschke, A. 596(1992)165
Cecil, T. L. and Rutan, S. C.
Fluorescence detection in liquid chromatography with an intensified diode-array detector 556(1991)495
Čelap, M. B., see Janjić, T. J. 596(1992)91
Čelap, M. B., see Tešić, Ž. L. 585(1991)359
Čelap, M. B., see Vučković, G. 585(1991)181

- Celardo, A., Dell'Elba, G., Frassanito, R., Benfenati, E. and Bonati, M.
Simultaneous determination of isbufylline and its major metabolites in rabbit blood and urine by reversed-phase high-performance liquid chromatography 568(1991)407
- Celeste, S., see Caccia, S. 581(1992)109
- Celma, C. and Giralt, E.
Determination of the enantiomeric purity of synthetic peptides by gas chromatography-mass spectrometry 562(1991)447
- Ceriani, G., see Dal Bo, L. 573(1992)158
- Černý, J., Šebor, G. and Blažek, J.
Use of basic alumina in fractionation of fossil fuels 600(1992)243
- Cert, A., see Ruiz-Gutierrez, V. 575(1992)1
- Cessna, A. J. and Grover, R.
Determination of the herbicide diclofop in human urine 600(1992)327
- Ceva, P., see Pietta, P. 553(1991)223
- Chabanet, C. and Yvon, M.
Prediction of peptide retention time in reversed-phase high-performance liquid chromatography 599(1992)211
- Chabard, J. L., see Andriollo, O. 575(1992)301
- Chabot, G. G., see Barilero, I. 575(1992)275
- Chabraoui, L., Mathian, B., Patricot, M. C. and Revol, A.
Specific assay for unconjugated dehydroepiandrosterone in human plasma by capillary gas chromatography with electron-capture detection 567(1991)299
- Chai, T. J., see Liu, C. Y. 555(1991)291
- Chaiken, I.
Interactions and uses of antisense peptides in affinity technology (Review) 597(1992)29
- Chakraborty, M., Lahiri, P., Anderson, G. M. and Chatterjee, D.
Use of high-performance liquid chromatography for assay of glutamic acid decarboxylase. Its limitation in use for *post-mortem* brain 571(1991)235
- Chakravarthy, B. K., see Mangalan, S. 567(1991)498
- Chalom, J., see Tod, M. 594(1992)386
- Chalykh, A. E., Kolomiets, L. N., Larionov, O. G. and Vinogradova, N. I.
Investigation of solid surfaces by high-performance liquid chromatography 592(1992)121
- Chambaz, D. and Haerdi, W.
Determination of divalent trace metals in natural waters by preconcentration on N,N,N',N'-tetra(2-aminoethyl)ethylenediamine-silica followed by on-line ion chromatography 600(1992)203
- Chamberlin, B. A., see Schultz, G. A. 590(1992)329
- Chambers, L. M., see Warner, C. D. 562(1991)625
- Chamkasem, N., Hill, K. D. and Sewell, G. W.
High-performance liquid chromatographic column-switching technique for the determination of intermediates of anaerobic degradation of toluene in ground water microcosm 587(1991)185
- Champluvier, B., see Woker, R. 584(1992)85
- Champney, T. H., Hanneman, W. H. and Nichols, M. A.
 γ -Aminobutyric acid, catecholamine and indoleamine determinations from the same brain region by high-performance liquid chromatography with electrochemical detection 579(1992)334
- Chan, E. C. S., Siboo, I. R. and Siboo, R.
Purification of phospholipase C by hydrophobic interaction affinity chromatography 568(1991)85
- Chan, K., Lau, O. W. and Wong, Y. C.
Determination of pethidine and its major metabolites in human urine by gas chromatography 565(1991)247
- Chan, K., see Walubo, A. 567(1991)261
- Chan, K. Y., George, R. C., Chen, T.-M. and Okerholm, R. A.
Direct enantiomeric separation of terfenadine and its major acid metabolite by high-performance liquid chromatography, and the lack of stereoselective terfenadine enantiomer biotransformation in man 571(1991)291
- Chan, P. F., see Law, B. 552(1991)429
- Chandra Sekharan, P., see Ragupathi, G. 574(1992)267
- Chanev, C., see Gavrilo, T. B. 552(1991)179
- Chang, F. C., see Iida, T. 558(1991)451
- Chang, F.-M., see Cheng, F.-C. 582(1992)19
- Chang, J.-P., see Pirkle, W. H. 598(1992)1
- Chang, S. Y., Moore, T. A., Devaud, L. L., Taylor, L. C. E. and Hollingsworth, E. B.
Analysis of rat brain microdialysate by gas chromatography-high-resolution selected-ion monitoring mass spectrometry 562(1991)111
- Chang, T., see Janiczek, N. 571(1991)179
- Chang, X., see Geng, X. 599(1992)185
- Chari, G.
High-performance liquid chromatographic determination of morphine, morphine-3-glucuronide, morphine-6-glucuronide and codeine in biological samples using multi-wavelength forward optical detection. Reply to Bogusz 579(1992)191
- Chari, G., Gulati, A., Bhat, R. and Tebbett, I. R.
High-performance liquid chromatographic determination of morphine, morphine-3-glucuronide, morphine-6-glucuronide and codeine in biological samples using multi-wavelength forward optical detection 571(1991)263
- Charles, B. G., see Taylor, P. J. 581(1992)152
- Chase, H. A. and Draeger, N. M.
Affinity purification of proteins using expanded beds 597(1992)129
- Chase, H. A., see McCreath, G. E. 597(1992)189
- Chatterjee, D., see Chakraborty, M. 571(1991)235
- Chauret, N., Carrier, J., Mancini, M., Neufeld, R., Weber, M. and Archambault, J.
Gas chromatographic-mass spectrometric analysis of ginkgolides produced by *Ginkgo biloba* cell culture 588(1991)281
- Chauret, N., Côté, J., Archambault, J. and André, G.
High-performance gel permeation chromatographic analysis of immunoglobulin M produced by hybridoma cell culture 594(1992)179

- Chawla, B. and Di Sanzo, F.
Determination of sulfur components in light petroleum streams by high-resolution gas chromatography with chemiluminescence detection 589(1992)271
- Cecchini, L., see Rodriguez, O. M. 555(1991)221
- Cheli, P. J., see Phillips, D. J. 599(1992)239
- Chelminska-Bertilsson, M. and Allenmark, S.
Liquid chromatographic monitoring of pseudochoolinesterase activity: comparison of methods 575(1992)237
- Chemke, J., see Zamir, I. 567(1991)319
- Chen, C.-F., see Tsai, T.-H. 567(1991)405
- Chen, C.-F., see Tsai, T.-H. 576(1992)170
- Chen, C.-F., see Tsai, T.-H. 598(1992)143
- Chen, C. L., Lesseley, B. A., Clarke, C. R., Roder, J. D., Sangiah, S., Berlin, K. D., Garrison, G. L., Scherlag, B. J., Lazzara, R. and Patterson, E.
High-performance liquid chromatographic determination of BRB-I-28, a novel antiarrhythmic agent, in dog plasma and urine 583(1992)274
- Chen, D. Y., Swerdlow, H. P., Harke, H. R., Zhang, J. Z. and Dovichi, N. J.
Low-cost, high-sensitivity laser-induced fluorescence detection for DNA sequencing by capillary gel electrophoresis 559(1991)237
- Chen, F.-M., Liu, C.-Z., Gaffar, S. A. and Epstein, A. L.
Chromatographic methods for large-scale preparation of immunoglobulin G_{2a} monoclonal antibodies Lym-1 and TNT-1 F(ab')₂ fragments 563(1991)243
- Chen, F.-T. A.
Rapid protein analysis by capillary electrophoresis 559(1991)445
- Chen, H. and Anderson, R. E.
Quantitation of phenacyl esters of retinal fatty acids by high-performance liquid chromatography 578(1992)124
- Chen, I.-W., Lin, J. H. and Dorley, J.
Quantitation of a new cholecystokinin and gastrin receptor antagonist (L-365,260) in dog and rat plasma by high-performance liquid chromatography 576(1992)363
- Chen, J., see Fritz, J. S. 555(1991)329
- Chen, J.W., Cohen, A.S. and Karger, B.L.
Identification of DNA molecules by pre-column hybridization using capillary electrophoresis 559(1991)295
- Chen, M. J., see Liu, C. Y. 555(1991)291
- Chen, M.-L., see Huang, H.-S. 564(1991)195
- Chen, S., Ocampo, A. P. and Kucera, P. J.
Liquid chromatographic method for the determination of calcium cyanamide using pre-column derivatization 558(1991)141
- Chen, S., see Mariot, R. 576(1992)143
- Chen, S.-H., Wu, S.-M. and Wu, H.-L.
Stereochemical analysis of betamethasone and dexamethasone by derivatization and high-performance liquid chromatography 595(1992)203
- Chen, T. K., Erhard, K. F., Last, T., Eggleston, D. S. and Ho, M. Y. K.
Direct high-performance liquid chromatographic separation of enantiomeric peptidoleukotriene antagonists 596(1992)123
- Chen, T.-M., see Chan, K. Y. 571(1991)291
- Chen, Y., Potter, J. M. and Ravenscroft, P. J.
High-performance liquid chromatographic method for the simultaneous determination of monoethylglycinexylidide and lignocaine 574(1992)361
- Chen, Y.-H., see Oshima, T. 585(1991)255
- Cheng, C. and Huang, L.-H.
Application of ligand-exchange chromatography to the assay of L-alanine from DL-aspartic acid by *Pseudomonas dacunhae* 555(1991)272
- Cheng, F.-C., Yang, L.-L., Chang, F.-M., Chia, L.-G. and Kuo, J.-S.
Simultaneous measurement of serotonin, catecholamines and their metabolites in cat and human plasma by *in vitro* microdialysis-microbore high-performance liquid chromatography with amperometric detection 582(1992)19
- Cheng, J., see Wu, C. 594(1992)243
- Cheng, K. N., Elsom, L. F. and Hawkins, D. R.
Identification of metabolites of halofantrine, a new candidate anti-malarial drug, by gas chromatography-mass spectrometry 581(1992)203
- Cherton, J.-C., Chevassus-Clement, M.-T., Lange, C., Convert, O., Fritz, R. and Leroux, P.
Behaviour studies of the fungicide cymoxanil in two strains of the fungus *Botrytis cinerea* and in haemolymph of locust and lobster. I. *In situ* monitoring by internal surface reversed-phase high-performance liquid chromatography 566(1991)499
- Cherton, J.-C., Lange, C., Mulheim, C., Pais, M., Cassier, P. and Vey, A.
Direct *in vitro* and *in vivo* monitoring of destruxins metabolism in insects using internal surface reversed-phase high-performance liquid chromatography. I. Behaviour of E destruxin in locusts 566(1991)511
- Chesson, S. M., see Hsu, R. S. 572(1991)352
- Chetsanga, C. J., see Carelse, O. 558(1991)318
- Cheung, S. W., Tang, S. W. and Remington, G.
Simultaneous quantitation of loxapine, amoxapine and their 7- and 8-hydroxy metabolites in plasma by high-performance liquid chromatography 564(1991)213
- Chevalier, A., see Dorbon, M. 557(1991)155
- Chevanne, F., see Le Verge, R. 574(1992)283
- Chevassus-Clement, M.-T., see Cherton, J.-C. 566(1991)499
- Chia, L.-G., see Cheng, F.-C. 582(1992)19
- Chiandetti, L., see Cracco, A. T. 574(1992)156
- Chiang, P. K., see Brown, N. D. 563(1991)466
- Chiappinelli, L., see Pessi, A. 557(1991)307
- Chiari, M., Etti, C. and Righetti, P. G.
Capillary zone electrophoresis analysis of acrylamido buffers for isoelectric focusing in immobilized pH gradients 559(1991)119
- Chiari, M., Giacomini, M., Micheletti, C. and Righetti, P. G.
Synthesis of a new acrylamido buffer (acryloylhistamine) for isoelectric focusing in immobilized pH gradients and its analysis by capillary zone electrophoresis 558(1991)285
- Chiari, M., Micheletti, C., Righetti, P. G. and Poli, G.
Polyacrylamide gel polymerization under non-oxidizing conditions, as monitored by capillary zone electrophoresis 598(1992)287

- Chiba, K., see Kobayashi, K. 579(1992)299
- Chiba, M., see Marvin, C. H. 555(1991)147
- Chiba, R. and Ishii, Y.
Simultaneous determination of yohimbine hydrochloride, strychnine nitrate and methyltestosterone by ion-pair high-performance liquid chromatography 588(1991)344
- Chien, R.-L. and Burgi, D. S.
Field amplified sample injection in high-performance capillary electrophoresis 559(1991)141
- Chien, R.-L. and Burgi, D. S.
Field-amplified polarity-switching sample injection in high-performance capillary electrophoresis 559(1991)153
- Chiou, R. H.-Y. and Lo, M.-W.
High-performance liquid chromatographic determination of angiotensin II receptor antagonists in human plasma and urine. I. DuP 532 (L-694,492) 581(1992)165
- Chiou, R. H.-Y. and Lo, M.-W.
Determination of remoxipride in human plasma and urine by reversed-phase ion-pair high-performance liquid chromatography 581(1992)300
- Chipman, G. R. and Cruickshank, K. A.
Characterization of N-acyl-D-biotinols by particle-beam liquid chromatography-mass spectrometry. An alternative to probe mass spectrometry for thermally labile samples 554(1991)141
- Chiu, S.-H. L., see Alvaro, R. F. 578(1992)327
- Chloupek, R. C., Hancock, W. S. and Snyder, L. R.
Computer simulation as a tool for the rapid optimization of the high-performance liquid chromatographic separation of a tryptic digest of human growth hormone 594(1992)65
- Chmel, K. and Gajdůšková, V.
New Czechoslovak sorbent, Ekosorb 552(1991)449
- Chmelík, J. and Thormann, W.
Isoelectric focusing field-flow fractionation. III. Investigation of the influence of different experimental parameters on focusing of cytochrome *c* in the trapezoidal cross-section channel 600(1992)297
- Chmelík, J. and Thormann, W.
Isoelectric focusing field-flow fractionation. IV. Investigations on protein separations in the trapezoidal cross-section channel 600(1992)305
- Chmelík, J., see Pazourek, J. 593(1992)357
- Chmelík, J., see Thormann, W. 589(1992)321
- Chmelík, J., see Urbánková, E. 583(1992)27
- Cholerton, S., Idle, M. E., Vas, A., Gonzalez, F. J. and Idle, J. R.
Comparison of a novel thin-layer chromatographic-fluorescence detection method with a spectrofluorometric method for the determination of 7-hydroxycoumarin in human urine 575(1992)325
- Chollet, D. and Künstner, P.
Fast systematic approach for the determination of drugs in biological fluids by fully automated high-performance liquid chromatography with on-line solid-phase extraction and automated cartridge exchange. Application to cefaracetam in human urine 577(1992)335
- Chollet, D. and Salanon, M.
Determination of Zy 17617B in plasma by solid-phase extraction and liquid chromatography with automated pre-column exchange 593(1992)73
- Choma, I., Dawidowicz, A. L. and Lodkowski, R.
High-performance liquid chromatography of benzodiazepines using sorbents with thermally immobilized Carbowax 20M 600(1992)109
- Choma, J., see Witkiewicz, Z. 556(1991)441
- Chopineau, J., see Sautou, V. 571(1991)298
- Chovan, J. P., Zane, P. A. and Greenberg, G. E.
Automated-extraction, high-performance liquid chromatographic method and pharmacokinetics in rats of a highly A₂-selective adenosine agonist, CGS 21680 578(1992)77
- Chow, M. S. S., see Fan, C.-D. 570(1991)217
- Chrambach, A., see Rampino, N. J. 596(1992)141
- Chrétien, J. R., see Righezza, M. 556(1991)169
- Chrétien, J. R., see Soják, L. 557(1991)241
- Chrétien, M., see Basak, A. 581(1992)17
- Chrisley, B. M., McNair, H. M. and Driskell, J. A.
Separation and quantification of the B₆ vitamers in plasma and 4-pyridoxic acid in urine of adolescent girls by reversed-phase high-performance liquid chromatography 563(1991)369
- Christensen, H. R., Angelo, H. R. and Skajaa, K.
Determination of isradipine and its pyridine metabolite in serum by capillary column gas chromatography with nitrogen-selective detection 574(1992)161
- Christensen, J. M., see Hansen, Å. M. 578(1992)85
- Christensen, J. V.
Determination of (+)-5-(2,3-dihydrobenzofuran-7-yl)-3-methyl-8-nitro-2,3,4,5-tetrahydro-1H-3-benzazepin-7-ol (NNC 01-0687), a novel dopamine D-1 receptor antagonist, in plasma by solid-phase extraction and high-performance liquid chromatography 577(1992)366
- Christensen, R. G. and Malone, W.
Determination of Oltipraz in serum by high-performance liquid chromatography with optical absorbance and mass spectrometric detection 584(1992)207
- Christensen, R. G., see Brown-Thomas, J. M. 568(1991)232
- Christie, W. W., see Hoving, E. B. 565(1991)103
- Chu, S., see Gustavson, L. E. 574(1992)313
- Chu, S.-Y., Sennello, L. T. and Sonders, R. C.
Simultaneous determination of clarithromycin and 14(*R*)-hydroxyclearithromycin in plasma and urine using high-performance liquid chromatography with electrochemical detection 571(1991)199
- Chua, S. C., see Lee, B. L. 581(1992)41
- Chuchalin, A. G., see Gorchakov, V. D. 563(1991)166
- Churáček, J., Pechová, H., Horna, A., Kotrla, R. and Ventura, K.
Chromatographic trace analysis of some organic compounds in the environment using derivatization-sorption concentration techniques. I. Gas chromatographic analysis of acrylates in air 557(1991)523
- Churchill, F. C., see Mount, D. L. 564(1991)181
- Cigánek, M., Dressler, M. and Teplý, J.
Gas chromatographic properties of immobilized poly(ethylene glycol) stationary phases 588(1991)225
- Ciranni-Signoretti, E., see Betto, P. 586(1991)149
- Cisowski, W., see Makuch, B. 594(1992)145

- Cissik, J. H., see Schwertner, H. A. 578(1992)297
- Cividalli, G., see Zamir, I. 565(1991)424
- Claeys, M., see Qimin, L. 562(1991)435
- Clapot, P., see Daumas, L. 570(1991)89
- Clare, R. A., Huang, S., Doig, M. V. and Gibson, G. G.
Gas chromatographic-mass spectrometric
characterisation of some novel hydroxyeicosatetraenoic
acids formed on incubation of arachidonic acid with
microsomes from induced rat livers 562(1991)237
- Clare, R. A., see Doig, M. V. 554(1991)181
- Clark, B. J., Hamdi, A., Berrisford, R. G., Sabanathan, S. and
Mearns, A. J.
Reversed-phase and chiral high-performance liquid
chromatographic assay of bupivacaine and its
enantiomers in clinical samples after continuous
extraplural infusion 553(1991)383
- Clark, D., see Tharakan, J. 595(1992)103
- Clark, J. C., see Jones, H. A. 570(1991)361
- Clark, L. A., Wichmann, J. K., Kucharczyk, N. and Sofia, R.
D.
Determination of the anticonvulsant felbamate in beagle
dog plasma by high-performance liquid chromatography
573(1992)113
- Clark, M., see Boege, F. 587(1991)3
- Clarke, C. R., see Chen, C. L. 583(1992)274
- Clarke, R. J., see Black, R. M. 558(1991)405
- Clarke, S. J., see Saska, M. 590(1992)147
- Claus, R., see Dehnhard, M. 566(1991)101
- Clawson, D. C., see Hoyer, G. L. 572(1991)159
- Clifford, A. A., see Bartle, K. D. 557(1991)69
- Cline Love, L. J., see Tomasella, F. P. 587(1991)325
- Clow, A., see Halket, J. M. 562(1991)279
- C.-Mendoza, C. E., Bhatti, T. and Bhatti, A. R.
Electrophoretic analysis of snake venoms 580(1992)355
- Coca, J., see Vega, A. 586(1991)303
- Cochet, S., see Kong Sing, Y. L. 598(1992)181
- Cociglio, M., Alric, R. and Bouvier, O.
Performance analysis of a reversed-phase liquid
chromatographic assay of lamotrigine in plasma using
solvent-demixing extraction 572(1991)269
- Cody, J. T.
Determination of methamphetamine enantiomer ratios in
urine by gas chromatography-mass spectrometry
(Review) 580(1992)77
- Coene, J., Ghijs, M., Van den Eeckhout, E., Van den Bossche,
W. and Sandra, P.
Evaluation of different packings for high-performance
liquid chromatographic analysis of alkyl
lysophospholipids 553(1991)285
- Coenegracht, P. M. J., see Wieling, J. 594(1992)45
- Coenen, A. J. J. M., Henckens, L. H. G., Mengerink, Y., Van
der Wal, S., Quaeflieg, P. J. L. M., Koole, L. H. and
Meijer, E. M.
Optimization of the separation of the Rp and Sp
diastereomers of phosphate-methylated DNA and RNA
dinucleotides 596(1992)59
- Coenen, A. J. J. M., Kerkhoff, M. J. G., Heringa, R. M. and
Van der Wal, S.
Comparison of several methods for the determination of
trace amounts of polar aliphatic monocarboxylic acids by
high-performance liquid chromatography 593(1992)243
- Cohen, A. I., see Whigan, D. B. 568(1991)385
- Cohen, A. S., see Chen, J. W. 559(1991)295
- Cohen, H. and Boutin-Muma, B.
Fluorescence detection of trichothecene mycotoxins as
coumarin-3-carbonyl chloride derivatives by high-
performance liquid chromatography 595(1992)143
- Cohen, J. D., see Michalczuk, L. 596(1992)294
- Cohen, K. A., see Palladino, D. E. H. 599(1992)3
- Cohen, S., see Iohan, F. 564(1991)27
- Colagrande, F., see Jackman, G. P. 566(1991)234
- Cole, R. O., Sepaniak, M. J., Hinze, W. L., Gorse, J. and
Oldiges, K.
Bile salt surfactants in micellar electrokinetic capillary
chromatography. Application to hydrophobic molecule
separations 557(1991)113
- Cole, S. M., Macrae, P. V., Merson, J. R., Pullen, F. S. and
Rance, D. J.
Microbore liquid chromatography coupled to a flow fast
atom bombardment probe for the on-line detection of the
Tyr-Pro cleavage of a nonapeptide by recombinant HIV-1
protease 562(1991)67
- Coleby, S. E., see Bartle, K. D. 557(1991)69
- Colenutt, B. A., see Cao, X. L. 555(1991)183
- Colin, H., Hilaireau, P. and Martin, M.
Flip-flop elution concept in preparative liquid
chromatography 557(1991)137
- Collier, G. R., see Uhe, A. M. 564(1991)81
- Collier, P. D., see Van Bergen, C. A. 589(1992)109
- Collijn, E., see Arvidsson, T. 586(1991)271
- Collins, C. H., see De A. Leone, P. 553(1991)399
- Collins, J. M., see Anderson, L. W. 581(1992)195
- Collister, J. W., see Lichtfouse, E. 585(1991)177
- Colmsjö, A., see Bemgård, A. 595(1992)247
- Colombini, M. P., Carrai, P., Fuoco, R. and Abete, C.
Rapid and sensitive determination of urinary 2,5-
hexanedione by reversed-phase high-performance liquid
chromatography 592(1992)255
- Colonna, S., see Carrea, G. 600(1992)320
- Colzani, V., see Lombardi, F. 576(1992)129
- Colzi, A., see Santoni, G. 581(1992)287
- Compton, B. J.
Electrophoretic mobility modeling of proteins in free zone
capillary electrophoresis and its application to
monoclonal antibody microheterogeneity analysis
559(1991)357
- Conder, J. R., see Leaver, G. 590(1992)101
- Condoret, J. S., see Graber, M. 584(1992)115
- Cone, E. J. and Darwin, W. D.
Rapid assay of cocaine, opiates and metabolites by gas
chromatography-mass spectrometry (Review)
580(1992)43
- Cone, E. J. and Deyl, Z.
Foreword 580(1992)1
- Confalonieri, S., see Caccia, S. 581(1992)109
- Confalonieri, S., see Guiso, G. 572(1991)346
- Constant, H., see Duthel, J. M. 579(1992)85
- Constanzer, M. L., Matuszewski, B. K. and Bayne, W. F.
High-performance liquid chromatographic method for the
determination of finasteride in human plasma at
therapeutic doses 566(1991)127
- Constanzer, M. L., see Herold, M. L. 581(1992)227

- Conte, L. S., see Antonelli, A. 552(1991)273
- Conti, P., see Hamoir, T. 589(1992)31
- Convert, O., see Cherton, J.-C. 566(1991)499
- Cook, J. A., see Antoniewicz, S. M. 573(1992)93
- Cook, J. A., see Skee, D. M. 568(1991)494
- Cooke, N., see Guttman, A. 559(1991)285
- Cooke, N., see Guttman, A. 593(1991)297
- Cooks, R. G., see Kaiser, Jr., R. E. 562(1991)3
- Cooles, G. P., O'Brien, A. P. and Watt, J. J.
Quality control procedure for the gas chromatographic determination of light hydrocarbons in petroleum liquid 588(1991)259
- Cooper, D. A., see Lurie, I. S. 598(1992)59
- Cooper, S. F., see Bi, H. 582(1992)93
- Coppet, L., see Maisonneuve, A. 583(1992)155
- Coppi, G. and Barchielli, M.
Simple high-performance liquid chromatographic method for the determination of PGT/IA, a new immunostimulating drug, in biological fluids 563(1991)385
- Coquart, V. and Hennion, M.-C.
Interference removal in the organic trace-level analysis of aqueous environmental samples by on-line liquid chromatographic preconcentration techniques with two precolumns 553(1991)329
- Coquart, V. and Hennion, M.-C.
Determination of chlorotriazines in aqueous environmental samples at the ng/l level using preconcentration with a cation exchanger and on-line high-performance liquid chromatography 585(1991)67
- Coquart, V. and Hennion, M.-C.
Trace-level determination of polar phenolic compounds in aqueous samples by high-performance liquid chromatography and on-line preconcentration on porous graphitic carbon 600(1992)195
- Coquet, A., Veuthey, J.-L. and Haerd, W.
Selective post-column fluorogenic reaction with benzamidine for trace level detection of reducing saccharides in liquid chromatography 553(1991)255
- Corbin, R., see Begley, P. 588(1991)239
- Cordonnier, P., see Moreau, J. 576(1992)103
- Cordonnier, P., see Palette, C. 563(1991)103
- Corradetti, E., see Menichini, E. 555(1991)211
- Corthout, J., see Qimin, L. 562(1991)435
- Corvetta, A., Della Bitta, R., Luchetti, M. M. and Pomponio, G.
5-Methylcytosine content of DNA in blood, synovial mononuclear cells and synovial tissue from patients affected by autoimmune rheumatic diseases 566(1991)481
- Costi, P., see Zecca, L. 567(1991)425
- Côté, J., see Chauret, N. 594(1992)179
- Côté, M. G., see Bi, H. 582(1992)93
- Cottet-Emard, J. M., see Pagliari, R. 563(1991)23
- Couderc, F., see Le Quééré, J. L. 562(1991)659
- Coudore, F., Ardid, D., Eschalier, A., Lavarenne, J. and Fialip, J.
High-performance liquid chromatographic determination of amitriptyline and its main metabolites using a silica column with reversed-phase eluent. Application in mice 584(1992)249
- Courcoux, P., Serot, T., Larre, C. and Popineau, Y.
Characterization and identification of wheat cultivars by multi-dimensional analysis of reversed-phase high-performance liquid chromatograms 596(1992)225
- Courthaudon, L. O., see Fujinari, E. M. 592(1992)209
- Courtheyn, D., Desaever, C. and Verhe, R.
High-performance liquid chromatographic determination of clenbuterol and cimaterol using post-column derivatization 564(1991)537
- Courtheyn, D., see Vermoesen, A. 564(1991)385
- Courtot, D., see Benoit, E. 583(1992)167
- Courtot, D., see Jaussaud, P. 573(1992)136
- Coustille, J. L., see Latxague, L. 586(1991)166
- Cousty-Berlin, D., see Vicat, P. 575(1992)315
- Coutant, J. E., Westmark, P. A., Nardella, P. A., Walter, S. M. and Okerholm, R. A.
Determination of terfenadine and terfenadine acid metabolite in plasma using solid-phase extraction and high-performance liquid chromatography with fluorescence detection 570(1991)139
- Coutts, R. T., see Alessi-Severini, S. 582(1992)173
- Coutts, R. T., see Hussain, M. D. 582(1992)203
- Coutts, R. T., see Paetsch, P. R. 573(1992)313
- Coutts, R. T., see Torok-Both, G. A. 579(1992)99
- Coventry, P. A., see Hoaksey, P. E. 566(1991)244
- Covert, R., see Browne, S. P. 575(1992)158
- Cox, G. B.
Influence of operating parameters on the preparative gradient elution chromatography of insulins 599(1992)195
- Cox, G. B. and Snyder, L. R.
Displacement effects in preparative gradient high-performance liquid chromatographic separations 590(1992)17
- Cox, R., see Kuykendall, J. R. 585(1991)322
- Cox, S. K., Wilke, A. V. and Frazier, D.
Determination of adriamycin in plasma and tissue biopsies 564(1991)322
- Cox, S. R., see Schmith, V. D. 568(1991)253
- Cozzone, P. J., see Masson, S. 563(1991)231
- Cozzone, P. J., see Rugani, N. 583(1992)246
- Cracco, A. T., Dall'Amico, R., Ferrari, V., Bottaro, S., Zacchello, G., Chiandetti, L. and Zacchello, F.
Determination of 8-methoxypsoralen in plasma by gas chromatography-mass spectrometry using selected-ion monitoring 574(1992)156
- Craft, N. E., see Epler, K. S. 595(1992)89
- Craft, N. E., Wise, S. A. and Soares, Jr., J. H.
Optimization of an isocratic high-performance liquid chromatographic separation of carotenoids 589(1992)171
- Cragoe, Jr., E. J., see Alliegro, M. A. 582(1992)217
- Cragoe, Jr., E. J., see Holthe, M. R. 579(1992)376
- Cramer, H., see Rissler, K. 564(1991)67
- Cramer, S. M., see Gerstner, J. A. 596(1992)173
- Cramers, C. A.
Unified separation science (by J. C. Giddings) (Book Review) 585(1991)380
- Cramers, C.A., see Houben, R.J. 554(1991)351
- Cramers, C. A., see Janssen, H.-G. 552(1991)527

- Cravedi, J. P. and Delous, G.
Use of an immobilized enzyme reactor for the analysis of residues of 17 α -methyltestosterone in trout by high-performance liquid chromatography 564(1991)461
- Creaser, C. S., Koupai-Abyazani, M. R. and Stephenson, G. R.
Origin and control of multi-peak formation in the analysis of trimethylsilyl derivatives of flavanone aglycones by capillary column gas chromatography 586(1991)323
- Cremer, J. and Henning, K.
Application of reversed-phase medium-pressure liquid chromatography to the isolation, separation and amino acid analysis of two closely related peptide toxins of the cyanobacterium *Microcystis aeruginosa* strain PCC 7806 587(1991)71
- Cremer, J., Meyer, H. and Henning, K.
Separation of methylated and non-methylated cyanoginosin-LR homologues of the cyanobacterium *Microcystis aeruginosa* strain PCC 7806 by reversed-phase medium-pressure liquid chromatography 558(1991)430
- Crétier, G., El Khabchi, M. and Rocca, J. L.
Preparative liquid chromatography. II. Existence of optimum injection conditions for overloaded gradient elution separations 596(1992)15
- Crétier, G., Macherel, L. and Rocca, J. L.
Preparative liquid chromatography. I. Influence of column efficiency on optimum injection conditions under isocratic elution 590(1992)175
- Crimmins, D. L. and Thoma, R. S.
Chromatographic analysis of tropomyosins from rabbit skeletal, chicken gizzard and earthworm muscle 599(1992)51
- Crinis, N. A., see Webster, L. K. 565(1991)482
- Crocq, A., see Fournion, J. M. 556(1991)287
- Cromie, D. D. O., see Van Bergen, C. A. 589(1992)109
- Crommelin, D. J. A., see Grit, M. 585(1991)239
- Crooks, B., see Gaskell, R. M. 553(1991)357
- Cruickshank, K. A., see Chipman, G. R. 554(1991)141
- Cserháti, T.
Dependence of the silanophilic effect on the concentration of preadsorbed salts and on the chemical structure of peptides in reversed-phase thin-layer chromatography 553(1991)467
- Cserháti, T.
Support matrix effects in the reversed-phase thin-layer chromatography of some peptides 600(1992)149
- Cserháti, T. and Magyar, K.
Lipophilicity determination of some monoamine oxidase inhibitors: the effect of methanol and ammonium chloride 575(1992)57
- Cserháti, T., see Bánhegyi, G. 553(1991)407
- Cserháti, T., see Forgács, E. 592(1992)75
- Cserháti, T., see Forgács, E. 600(1992)43
- Cubría, J. C., see Alvarez-Bujidos, M. L. 578(1992)321
- Cubría, J. C., see Negro, A. 576(1992)135
- Cummings, J., see Anderson, L. 574(1992)150
- Cunico, B., see Dollinger, G. 592(1992)215
- Cunico, R. L., Gruhn, V., Kresin, L., Nitecki, D. E. and Wiktorowicz, J. E.
Characterization of polyethylene glycol modified proteins using charge-reversed capillary electrophoresis 559(1991)467
- Cupo, J. F.
Electrophoretic analysis of nuclear matrix proteins and the potential clinical applications (Review) 569(1991)389
- Curtis, M. A., see Pullen, R. H. 573(1992)49
- Cusido, E., see Caturla, M. C. 581(1992)101
- Cusido, E., see Caturla, M. C. 593(1992)69
- Cutler, D. J., see Carter, S. R. 574(1992)77
- Cutler, D. J., see McLachlan, A. J. 570(1991)119
- Cypess, A. M., see Lloyd, D. K. 568(1991)117
- Cyronak, M. J., see Boppana, V. K. 592(1992)317
- Cysewski, P., see Lemque, R. 599(1992)255
- Czech, J., see Leyssens, L. 564(1991)515
- Czok, M., see Katti, A. M. 556(1991)205
- Czok, M., see Merino-Dugay, A. 579(1992)73

D

- Dąbrowski, R., see Mazur, J. 600(1992)123
- Da Col, R., see Silvestro, L. 591(1992)225
- Daenens, P., see Debrabandere, L. 564(1991)557
- Daeseleire, E., De Guesquière, A. and Van Peteghem, C.
Derivatization and gas chromatographic-mass spectrometric detection of anabolic steroid residues isolated from edible muscle tissues 562(1991)673
- Daeseleire, E., De Guesquière, A. and Van Peteghem, C.
Combined high-performance liquid chromatography and radioimmunoassay for the screening of 19-nortestosterone and methyltestosterone residues in meat samples 564(1991)445
- Daeseleire, E., De Guesquière, A. and Van Peteghem, C.
Detection of the illegal use of ethinylestradiol in cattle urine by gas chromatography-mass spectrometry 564(1991)469
- Dagli, E., see Assadullahi, T. P. 565(1991)349
- D'Agostino, P. A. and Provost, L. R.
Determination of chemical warfare agents, their hydrolysis products and related compounds in soil 589(1992)287
- D'Agostino, P. A. and Provost, L. R.
Mass spectrometric identification of products formed during degradation of ethyl dimethylphosphoramidocyanidate (tabun) 598(1992)89
- D'Agostino, P. A. and Provost, L. R.
Capillary column gas chromatography-ammonia and deuterated ammonia chemical ionization mass spectrometry of sulfur vesicants 600(1992)267

- D'Agrosa, F., see Bianchi-Bosisio, A. 569(1991)243
- DaGue, B. B., see Suter, M. J.-F. 553(1991)101
- Dahlke, S., see Gassmann, G. 598(1992)313
- Dahms, T. E., see Scott, M. D. 576(1992)149
- Daitch, C. E., see Khachik, F. 582(1992)153
- Dakshinamurti, K., see Sharma, S. K. 578(1992)45
- Dal Bo, L., Ceriani, G. and Broccoli, G.
Determination of vincamine in human plasma by high-performance liquid chromatography with ultraviolet detection 573(1992)158
- Dale, A. D. and Evans, M. B.
Gas-liquid chromatography in qualitative analysis. XIX. The use of antioxidants to delay the oxidation of polyoxyethylene glycol stationary phases 552(1991)161
- Dalet-Fumeron, V., Guinec, N. and Pagano, M.
High-performance liquid chromatographic method for the simultaneous purification of cathepsins B, H and L from human liver 568(1991)55
- Dalla Libera, L., Fasolo, M., Cavallini, P., Ratti, E. and Gaviraghi, G.
Use of anion-exchange high-performance liquid chromatography for the study of smooth muscle myosin light-chain kinase and its catalytic domain 566(1991)427
- Dall'Amico, R., see Cracco, A. T. 574(1992)156
- Damaj, M. I., Trouvin, J. H., Lambrey, B. and Jacquot, C.
Determination of a new 2-amino-2-oxazoline (COR 3224) in plasma and brain tissue of the rat by high-performance liquid chromatography with electrochemical detection 563(1991)476
- Damani, L. A., see Epemolu, R. O. 573(1992)178
- Damani, L. A., see On, N. T. 574(1992)352
- Damani, L. A., see Stevenson, P. J. 563(1991)419
- Damani, L. A., see Usansky, J. I. 563(1991)283
- D'Amato, G., see Castello, G. 585(1991)93
- Damodaran, C., see Ragupathi, G. 574(1992)267
- Dandekar, K. A., see Janiszewski, J. S. 567(1991)274
- Dandekar, K. A., see Janiszewski, J. S. 577(1992)151
- Dandeu, J.-P., Rabillon, J., Lux, M., David, B., Guillaume, J.-L. and Camoin, L.
Isolation of *Der p1*, the *Dermatophagoides pteronyssinus* major mite allergen, from a crude mite culture extract, purification by ion-chromatography, and comparison between the material obtained and a cDNA-coded *Der p1* 599(1992)105
- Dandeu, J.-P., see Krstulović, A. M. 584(1992)1
- Danese, M., see Bagnati, R. 564(1991)493
- Danhof, M., see Heeremans, C. E. M. 554(1991)205
- Däppen, R. and Molnar, I.
Application of the gradient elution technique. Demonstration with a special test mixture and the DryLab G/plus method development software 592(1992)133
- Däppen, R., see Katti, A. 590(1992)127
- D'Aranno, V., Mancinelli, A. and Manzini, S.
Determination of the tricyclic compound adosupine and its three metabolites in plasma and brain of rat using high-performance liquid chromatography 574(1992)319
- Darbellay, C., see Slacanin, I. 557(1991)391
- Darwin, W. D., see Cone, E. J. 580(1992)43
- Dasnois, C., see Vandebroek, M. 564(1991)405
- Datta, A., see Wagh, J. S. 587(1991)280
- Daudon, F., see Dufay, C. 563(1991)224
- Daumas, L., Sabot, J.-F., Vermeulen, E., Clapot, P., Allegre, F., Pinatel, H., Boucherat, M. and Francois, B.
Determination of debrisoquine and metabolites in human urine by gas chromatography-mass spectrometry 570(1991)89
- David, B., see Dandeu, J.-P. 599(1992)105
- David, C., see Fournion, J. M. 556(1991)287
- Davidson, V. P. and Traiger, G. J.
Determination of *p*-methylthiobenzamide and *p*-methylthiobenzamide-S-oxide from rat plasma using solid-phase extraction and high-performance liquid chromatography 567(1991)213
- Davies, J., see Berry, M. J. 587(1991)161
- Davies, J., see Berry, M. J. 597(1992)239
- Davies, J. H., see Poon, G. K. 576(1992)235
- Davies, N. W., see Jack, D. S. 584(1992)189
- Davies, N. W., see Neill, G. P. 565(1991)207
- Davis, J. M., see Oros, F. J. 591(1992)1
- Davis, S. S., see Barrett, D. A. 566(1991)135
- Dawes, J., see Hodson, B. A. 565(1991)416
- Dawidowicz, A. L., see Choma, I. 600(1992)109
- Dawodu, O. F. and Meisen, A.
Identification of products resulting from carbonyl sulphide-induced degradation of diethanolamine 587(1991)237
- Dawra, R. K., see Sharma, O. P. 587(1991)351
- Day, B. W., Naylor, S., Gan, L.-S., Sahali, Y., Nguyen, T. T., Skipper, P. L., Wishnok, J. S. and Tannenbaum, S. R.
Gas chromatographic-mass spectrometric analysis of diols and tetrols from reactions of polycyclic aromatic hydrocarbon epoxides with hemoglobin 562(1991)563
- Daykin, M. M., see Blagbrough, I. S. 578(1992)251
- De, B. and Bisset, N. G.
Separation of *Strychnos nux-vomica* alkaloids by high-performance liquid chromatography 587(1991)318
- De A. Leone, P. and Collins, C. H.
Isocratic reversed-phase high-performance liquid chromatographic separation of simple perhalogenated compounds 553(1991)399
- Debal, V., Morjani, H., Millot, J.-M., Angiboust, J.-F., Gourdiere, B. and Manfait, M.
Determination of vinorelbine (Navelbine) in tumour cells by high-performance liquid chromatography 581(1992)93
- De Bersaques, J. P., see Meyer, E. 570(1991)149
- De Bertrand, N., see Durand, G. 554(1991)233
- De Bertrand, N., see Durand, G. 562(1991)507
- De Besi, P., see Gelfi, C. 598(1992)277
- Debets, A. J. J., Hupe, K.-P., Kok, W. T. and Brinkman, U. A. T.
Electrodialytic sample treatment coupled on-line with column liquid chromatography for the determination of basic and acidic compounds in environmental samples 600(1992)163
- De Boer, J. H., see Witte, D. T. 558(1991)333
- De Boer, T., see Witte, D. T. 553(1991)365

- De Bokx, P. K., Baarslag, P. C. and Urbach, H. P.
Modelling of displacement chromatography using non-ideal isotherms 594(1992)9
- De Bokx, P. K., Gillissen, E. E. A., Van de Weijer, P., Bekkers, M. H. J., Van Bommel, C. H. M. and Janssen, H.-G.
Fluorescence detector cell for use in an integrated electrically driven separation system 598(1992)115
- Debordes, L., see Maisonneuve, A. 583(1992)155
- De Brabander, H. F., see Vermoesen, A. 564(1991)385
- Debrabandere, L., Van Boven, M. and Daenens, P.
High-performance liquid chromatography with electrochemical detection of buprenorphine and its major metabolite in urine 564(1991)557
- De Bruijn, A. A. G. M., see Dirven, H. A. A. M. 564(1991)266
- De Bruijn, E. A., see Pattyn, G. 564(1991)352
- Debruyckere, G. and Van Peteghem, C.
Detection of 19-nortestosterone and its urinary metabolites in miniature pigs by gas chromatography-mass spectrometry 564(1991)393
- De Castro, A., see Montaña, A. 594(1992)259
- De Clerck, F., see Weber, C. 562(1991)599
- De Clerck, F., see Weber, C. 577(1992)1
- Deegan, R. J., see He, H. B. 574(1992)213
- Deffo, P., see Piovetti, L. 588(1991)99
- De Fluiter, P., see Jansen, E. H. J. M. 580(1992)325
- Degen, P., see Flesch, G. 568(1991)261
- Degen, P. H. and Schneider, W.
Rapid and sensitive determination of low concentrations of nicotine in plasma by gas chromatography with nitrogen-specific detection 563(1991)193
- Degen, P. H. and Schneider, W.
Determination of 4-hydroxyandrostenedione in plasma and urine by extractive alkylation and electron-capture gas chromatography 565(1991)67
- Degen, P. H., see Flesch, G. 581(1992)147
- De Goede, P. N. F. C., see Van der Horst, A. 574(1992)166
- De Graaf, G., see Dos Ramos, F. J. 558(1991)125
- De Graaf, S. S. N., see Bloemhof, H. 572(1991)171
- De Graeve, J., see Vandenbroeck, M. 564(1991)405
- De Guesquière, A., see Daeseleire, E. 562(1991)673
- De Guesquière, A., see Daeseleire, E. 564(1991)445
- De Guesquière, A., see Daeseleire, E. 564(1991)469
- Dehnhard, M., Bernal-Barragan, H. and Claus, R.
Rapid and accurate high-performance liquid chromatographic method for the determination of 3-methylindole (skatole) in faeces of various species 566(1991)101
- Deinzer, M., see Svoboda, M. 562(1991)403
- De Jong, A. P. J. M., see Somsen, G. W. 552(1991)635
- De Jong, D., see Goosens, E. C. 552(1991)489
- De Jong, G. J., see Barceló, D. 553(1991)311
- De Jong, G. J., see Goosens, E. C. 552(1991)489
- De Jong, G. J., see Kwakman, P. J. M. 553(1991)345
- De Jong, G. J., see Steijger, O. M. 557(1991)13
- De Jong, G. J., see Zech, K. 600(1992)161
- De Jong, J., Guérand, W. S., Schoofs, P. R., Bast, A. and Vander Vijgh, W. J. F.
Simple and sensitive quantification of anthracyclines in mouse atrial tissue using high-performance liquid chromatography and fluorescence detection 570(1991)209
- De Jong, J., Munniksmas, C. N., Guérand, W. S., Bast, A. and Van der Vijgh, W. J. F.
Fully automated determination of a new anthracycline N-*l*-leucyldoxorubicin and six metabolites in plasma by high-performance liquid chromatography with on-line sample handling 574(1992)273
- De Jong, J., Van Polanen, A. and Driessen, J. J. M.
Determination of ethylenediaminetetraacetic acid and its salts in canned mushrooms by reversed-phase ion-pair liquid chromatography 553(1991)243
- De Klippel, N., see Sarre, S. 575(1992)207
- De Kok, T. M. C. M., Levels, P. J., Van Faassen, A., Hazen, M., Ten Hoor, F. and Kleinjans, J. C. S.
Chromatographic methods for the determination of toxicants in faeces (Review) 580(1992)135
- De Kraker, J., see Kaijser, G. P. 571(1991)121
- De Kraker, J., see Kaijser, G. P. 583(1992)175
- Delabays, N., see Slacanian, I. 557(1991)391
- De La Fournière, L., see Rugani, N. 583(1992)246
- Delage, E., Bohuon, G., Baron, A. and Drilleau, J.-F.
High-performance liquid chromatography of the phenolic compounds in the juice of some French cider apple varieties 555(1991)125
- Delahaut, P., see Vandenbroeck, M. 564(1991)405
- De Las Rivas, J., Milicua, J. C. G. and Gomez, R.
Determination of carotenoid pigments in several tree leaves by reversed-phase high-performance liquid chromatography 585(1991)168
- Delatour, P., see Benoit, E. 583(1992)167
- De la Vigne, U., Jänchen, D. E. and Weber, W. H.
Application of high-performance thin-layer chromatography and automated multiple development for the identification and determination of pesticides in water 553(1991)489
- Del Cont Bernard, A., Royer Morrot, M. J., Zhiri, A., Rambourg, M. and Royer, R. J.
Automated determination of disopyramide and N-monoalkyldisopyramide in plasma by reversed-phase liquid chromatography with a column switching system 574(1992)365
- De Leenheer, A. P., see Meyer, E. 570(1991)149
- Delépine, B. and Sanders, P.
Determination of chloramphenicol in muscle using a particle beam interface for combining liquid chromatography with negative-ion chemical ionization mass spectrometry 582(1992)113
- Deleu, D., see Sarre, S. 575(1992)207
- Delgado, C., Sancho, P., Mendieta, J. and Luque, J.
Ligand-receptor interactions in affinity cell partitioning. Studies with transferrin covalently linked to monomethoxypoly(ethylene glycol) and rat reticulocytes 594(1992)97
- Delgado, C., see Tejedor, M. C. 589(1992)127

- Delgado, M., see Liu, H. 576(1992)163
- Delhotal Landes, B., Miscoria, G. and Flouvat, B.
Determination of lansoprazole and its metabolites in plasma by high-performance liquid chromatography using a loop column 577(1992)117
- De Ligny, C. L., see Gelsema, W. J. 594(1992)217
- Della Bitta, R., see Corvetta, A. 566(1991)481
- Dellacherie, E., see Ubrich, N. 584(1992)17
- Dell'Elba, G., see Celardo, A. 568(1991)407
- Dello Strologo, L., see Betto, P. 584(1992)256
- Delmas, M., see Snini, A. 590(1992)369
- Del Nero, S., Di Somma, M. and Vigevani, A.
High-performance liquid chromatographic analysis of FCE 24304 (6-methylenandrosta-1,4-diene-3,17-dione) and FCE 24928 (4-aminoandrosta-1,4,6-triene-3,17-dione), two new aromatase inhibitors 593(1991)25
- De Lorenzi, E., see Baruffini, A. 593(1992)95
- De Lorenzi, E., see Kitsos, M. 553(1991)1
- Delorenzo, O., see Qimin, L. 562(1991)435
- Delous, G., see Cravedi, J. P. 564(1991)461
- Del Rio, J. A., see Castillo, J. 555(1991)285
- DeLuca, P. P., see Lee, K. C. 555(1991)73
- Delucas, L. J., see Meng, Q. C. 579(1992)63
- De Man, H. C. J., see Mengerink, Y. 552(1991)593
- Demby, K. B., White, T. C. and Ghanayem, B. I.
Determination of methacrylonitrile in serum using gas chromatography and flame ionization detection 579(1992)153
- De Merulis, G., see Di Muccio, A. 552(1991)241
- Demilo, A. B., see Warthen, Jr., J. D. 590(1992)133
- Demina, N., see Polanuer, B. 594(1992)173
- Demirbüker, M. and Blomberg, L. G.
Permanganate-impregnated packed capillary columns for group separation of triacylglycerols using supercritical media as mobile phases 600(1992)358
- Demirbüker, M., see Andersson, P. E. 595(1992)301
- Demiroglou, A., see Jennissen, H. P. 597(1992)93
- Demnerová, K., see Břizová, K. 593(1992)125
- De Moel, J. J. M., see Van de Nesse, R. J. 552(1991)613
- Demont, N., see Le Quéré, J. L. 562(1991)659
- DeMonte, L. B., see Tarditi, L. 599(1992)13
- Demorest, D. and Dubrow, R.
Factors influencing the resolution and quantitation of oligonucleotides separated by capillary electrophoresis on a gel-filled capillary 559(1991)43
- Dempsey, C. A., Lavicky, J. and Dunn, A. J.
Apparent inter-channel interference in dual-electrode electrochemical detection 596(1992)110
- Denaro, M., see Milana, M. R. 552(1991)205
- De Neve, W., see Van Belle, S. J.-P. 578(1992)223
- Deng, C., see Liu, S. 598(1992)298
- Den Hartog, J. M. P., see Arts, C. J. M. 564(1991)429
- Dennhardt, R., see Zimmermann, J. 567(1991)240
- Dennis, M. F., see Stratford, M. R. L. 582(1992)145
- Dennis, M. J., see Gulaid, A. A. 566(1991)228
- Dent, M. R., see Price, G. J. 585(1991)83
- De Potter, W. P., see Llona, I. 596(1992)51
- Deratani, A., see Thuaud, N. 555(1991)53
- De Reynies, A., see Houzé, P. 574(1992)305
- Dernovaya, L. A., Eltekov, Y. A., Hradil, J. and Svec, F.
Estimation of cellulose polarity by gas chromatography 552(1991)365
- De Ruiter, G. A., Smid, P., Schols, H. A., Van Boom, J. H. and Rombouts, F. M.
Detection of fungal carbohydrate antigens by high-performance immunoaffinity chromatography using a protein A column with covalently linked immunoglobulin G 584(1992)69
- Desaever, C., see Courtheyn, D. 564(1991)537
- Desage, M., Rousseau-Tsangaris, M., Lecompte, D. and Brazier, J. L.
Quantitation of trihexyphenidyl from plasma using a mass-selective detector and electron-impact ionization 571(1991)250
- Desai, D. M. and Gal, J.
Reversed-phase high-performance liquid chromatographic separation of the stereoisomers of labetalol via derivatization with chiral and non-chiral isothiocyanate reagents 579(1992)165
- De Sandro, V., see Ducrottoy, G. 566(1991)415
- Desbène, P.-L., Abderrezag, A. and Desmazières, B.
Analytical study of heavy crude oil fractions by coupling of the transalkylation reaction with supercritical fluid chromatography 593(1992)321
- Desbène, P. L., Essayegh, M., Desmazières, B. and Villeneuve, F.
Analysis of biomass pyrolysis oils by a combination of various liquid chromatographic techniques and gas chromatography-mass spectrometry 553(1991)211
- Desbène, P.-L., see Oliveros, L. 589(1992)53
- Desideri, P. G., see Rodriguez, O. M. 555(1991)221
- Desiderio, C., see Sakodinskaya, I. K. 596(1992)95
- Desiderio, D. M., see Harajiri, S. 575(1992)213
- Desiderio, D. M., see Kusmierz, J. J. 574(1992)189
- Desiderio, D. M., see Lovelace, J. L. 562(1991)573
- Desmangles, R. J. P., Flipo, D., Fournier, M. and Mateescu, M. A.
Fast separation of macrophages by retention on cross-linked amylose and release by enzymatic amylolysis of the chromatographic material 584(1992)121
- Desmazières, B., see Desbène, P. L. 553(1991)211
- Desmazières, B., see Desbène, P.-L. 593(1992)321
- Desmazières, B., see Oliveros, L. 589(1992)53
- Desmet, G., see Kamel, S. 574(1992)255
- De Smet, M., see Hamoir, T. 589(1992)31
- De Smet, M., see Van Belle, S. J.-P. 576(1992)351
- De Smet, M., see Van Belle, S. J.-P. 578(1992)223
- Desmoulin, F., see Masson, S. 563(1991)231
- Deterding, L. J., Moseley, M. A., Tomer, K. B. and Jorgenson, J. W.
Nanoscale separations combined with tandem mass spectrometry 554(1991)73
- Deterding, L. J., Parker, C. E., Perkins, J. R., Moseley, M. A., Jorgenson, J. W. and Tomer, K. B.
Nanoscale separations. Capillary liquid chromatography-mass spectrometry and capillary zone electrophoresis-mass spectrometry for the determination of peptides and proteins 554(1991)329

- Deutsch, J., Hegedus, L., Greig, N. H., Rapoport, S. I. and Soncrant, T. T.
Electron-impact and chemical ionization detection of nicotine and cotinine by gas chromatography-mass spectrometry in rat plasma and brain 579(1992)93
- DeVane, C. L., see Miller, R. L. 570(1991)412
- Devaud, L. L., see Chang, S. Y. 562(1991)111
- De Vecchi, E., see Fermo, I. 593(1992)171
- Devilbiss, E. D., see Warthen, Jr., J. D. 590(1992)133
- De Vos, M., Verdievel, H., Schoonjans, R., Beke, R., De Weerd, G. A. and Barbier, F.
High-performance liquid chromatographic assay for the determination of 5-aminosalicylic acid and acetyl-5-aminosalicylic acid concentrations in endoscopic intestinal biopsy in humans 564(1991)296
- De Vries, A. J. and Nooter, K.
Quantification of mitoxantrone in bone marrow by high-performance liquid chromatography with electrochemical detection 563(1991)435
- De Vries, J. X. and Kymber, K. A.
Thermospray and particle beam liquid chromatographic-mass spectrometric analysis of coumarin anticoagulants 562(1991)31
- De Vries, M. and Odink, J.
Simultaneous measurement of serotonin and 5-hydroxyindoleacetic acid in rat brain using a liquid chromatographic method with electrochemical detection 564(1991)250
- Dewaele, C., see Lin Ling, B. 553(1991)433
- Dewaele, C., see Maes, P. C. 558(1991)343
- De Weerd, G. A., see De Vos, M. 564(1991)296
- Deye, J. F., see Berger, T. A. 594(1992)291
- Deyl, Z.
Advances in electrophoresis, Vol. 3 (edited by A. Chrambach, M. J. Dunn and B. J. Radola) (Book Review) 565(1991)533
- Deyl, Z.
The use of HPLC in receptor biochemistry (edited by A. R. Kerlavage) (Book Review) 566(1991)266
- Deyl, Z.
High-performance liquid chromatography in biotechnology (edited by W. S. Hancock) (Book Review) 568(1991)501
- Deyl, Z. and Struzinsky, R.
Capillary zone electrophoresis: its applicability and potential in biochemical analysis (Review) 569(1991)63
- Deyl, Z., see Cone, E. J. 580(1992)1
- Deyl, Z., see Hanis, T. 553(1991)93
- Deyl, Z., see Hodny, Z. 578(1992)53
- Deyl, Z., see Righetti, P. G. 569(1991)1
- Dezan, C., see Rugani, N. 583(1992)246
- De Zeeuw, R. A.
Recent developments in therapeutic drug monitoring and clinical toxicology (edited by I. Sunshine) (Book Review) 584(1992)280
- DeZeeuw, R.A., see Witte, D.T. 553(1991)365
- De Zeeuw, R. A., see Witte, D. T. 558(1991)333
- Dhermy, D., see Kong Sing, Y. L. 558(1991)43
- Dhermy, D., see Kong Sing, Y. L. 598(1992)181
- Di, Z. S., see Kishino, S. 582(1992)246
- Diallo, B., Vanhaelen-Fastré, R. and Vanhaelen, M.
Direct coupling of high-speed counter-current chromatography to thin-layer chromatography. Application to the separation of asiaticoside and madecassoside from *Centella asiatica* 558(1991)446
- Dias, M. M., see Rodrigues, A. E. 590(1992)93
- Di Carlo, S., see Pichini, S. 581(1992)143
- Dickson, P. H., see Shara, M. A. 576(1992)221
- Di Domenico, A., see Menichini, E. 555(1991)211
- Diekmann, N., see Lohmann, A. 564(1991)283
- Diekmann, N., see Lohmann, A. 564(1991)289
- Diers, I., see Banke, N. 559(1991)325
- Dietrich, R. F., see Reiser, R. W. 554(1991)91
- Diez, M. T., see Arin, M. J. 582(1992)13
- Diez-Cascón, A., see Pruñonosa, J. 574(1992)127
- Diez-Cascón, A., see Pruñonosa, J. 581(1992)219
- Di Fava, R., see Betto, P. 586(1991)149
- Di Giovannandrea, R., see Severini, G. 563(1991)147
- DiLeo, E. M., see Hsu, R. S. 572(1991)352
- Dilettato, D., see Bruchet, A. 562(1991)469
- Dilks, Jr., C. H., see Kirkland, J. J. 593(1992)339
- Dillon, J. G. and Hughes, M. K.
Determination of cholesterol and cortisone absorption in polyurethane. I. Methodology using size-exclusion chromatography and dual detection 572(1991)41
- Dimitrov, N. V., see Bennett, J. L. 573(1992)146
- Dimov, N. and Moskovkina, M.
New aspects of quantitative structure-retention relationships in chromatography 552(1991)59
- Di Muccio, A., Ausili, A., Dommarco, R., Attard Barbini, D., Santilio, A., Vergori, F., De Merulis, G. and Sernicola, L.
Solid-matrix partition for separation of organochlorine pesticide residues from fatty materials 552(1991)241
- Di Muccio, A., Camoni, I., Vergori, L., Dommarco, R., Attard Barbini, D., Vergori, F., Ausili, A. and Santilio, A.
Screening for coumatetralyl in soft drinks by solid-matrix extraction and high-performance liquid chromatography with diode-array detection 553(1991)305
- Ding, S. F., see Peng, C. T. 586(1991)85
- Dingler, E., see Lohmann, A. 567(1991)506
- DiNunzio, J. E. and Gadde, R. R.
Development of stability indicating assay methods for the determination of hydroxamic acids in topical formulations 594(1992)209
- Dion, D. M., see Phillips, D. J. 599(1992)239
- Diop, A., see Pastore, P. 591(1992)219
- Diquet, B., see Thomare, P. 583(1992)217
- Diquet, B., see Van Der Meersch-Mougeot, V. 567(1991)441
- Dirven, H. A. A. M., De Bruijn, A. A. G. M., Sessink, P. J. M. and Jongeneelen, F. J.
Determination of the cytochrome P-450 IV marker, ω -hydroxylauric acid, by high-performance liquid chromatography and fluorimetric detection 564(1991)266
- Di Sanzo, F., see Chawla, B. 589(1992)271
- Di Somma, M., see Del Nero, S. 593(1991)25
- Di Toro, M., see Longo, A. 562(1991)639
- Dittberner, K., see Zobel, H. 553(1991)441

- Dixit, V. and Dixit, V. M.
Solid-phase extraction of 11-nor- Δ -9-tetrahydrocannabinol-9-carboxylic acid from human urine with gas chromatographic-mass spectrometric confirmation 567(1991)81
- Dixit, V., Nguyen, H. and Dixit, V. M.
Solid-phase extraction of fluoxetine and norfluoxetine from serum with gas chromatography-electron-capture detection 563(1991)379
- Dixit, V. M., see Dixit, V. 563(1991)379
- Dixit, V. M., see Dixit, V. 567(1991)81
- Dizdar, N., Henriksson, A. and Kågedal, B.
Determination of L-3,4-dihydroxyphenylalanine in biological fluids and tissues 565(1991)1
- Djordjevic, N. M., see Liu, G. 592(1992)239
- Djordjevic, N. M., see Liu, G. 598(1992)153
- Do, L. and Raulin, F.
Gas chromatography of Titan's atmosphere. III. Analysis of low-molecular-weight hydrocarbons and nitriles with a CP-Sil-5 CB WCOT capillary column 591(1992)297
- Dobbs, N. A. and Twelves, C. J.
Measurement of epidoxorubicin and its metabolites by high-performance liquid chromatography using an advanced automated sample processor 572(1991)211
- Döe De Maindreville, M., see Le Verge, R. 574(1992)283
- Doelemeyer, A., see Pollak, V. A. 596(1992)241
- Doerge, D. R. and Yee, A. B. K.
Liquid chromatographic determination of ethylenethiourea using pulsed amperometric detection 586(1991)158
- Dognini, G. L., see Abbiati, G. 566(1991)445
- Doherty, M., see Blagbrough, I. S. 578(1992)251
- Doherty, R. M., see Abraham, M. H. 587(1991)213
- Doherty, R. M., see Abraham, M. H. 587(1991)229
- Doig, M. V. and Clare, R. A.
Use of thermospray liquid chromatography-mass spectrometry to aid in the identification of urinary metabolites of a novel antiepileptic drug, Lamotrigine 554(1991)181
- Doig, M. V., see Clare, R. A. 562(1991)237
- Dolan, J. W., see Lewis, J. A. 592(1992)183
- Dolan, J. W., see Lewis, J. A. 592(1992)197
- Doležalová, M.
On-line solid-phase extraction and high-performance liquid chromatographic determination of nortriptyline and amitriptyline in serum 579(1992)291
- Dollinger, G., Cunico, B., Kunitani, M., Johnson, D. and Jones, R.
Practical on-line determination of biopolymer molecular weights by high-performance liquid chromatography with classical light-scattering detection 592(1992)215
- Dollinger, G., see Kunitani, M. 588(1991)125
- Dolphin, P. J., see Zhou, G. 568(1991)69
- Domar, U., see Kirchnerberger, J. 574(1992)237
- Dominguez, A., see Guillén, M. D. 591(1992)287
- Dommarco, R., see Di Muccio, A. 552(1991)241
- Dommarco, R., see Di Muccio, A. 553(1991)305
- Donahue, E. P., Brown, L. L., Flakoll, P. J. and Abumrad, N. N.
Rapid measurement of leucine-specific activity in biological fluids by ion-exchange chromatography and post-column ninhydrin detection 571(1991)29
- Donati, F., see Ducharme, J. 573(1992)79
- Dondi, F., see Blo, G. 556(1991)249
- Dondi, F., see Pietrogrande, M. C. 592(1992)65
- Donnelly, J. R. and Sovocool, G. W.
Gas chromatographic elution patterns of chlorinated dioxins versus column polarity 594(1992)269
- Doorn, L., Jansen, E. H. J. M. and Van Leeuwen, F. X. R.
Comparison of high-performance liquid chromatographic detection methods for thyronine and tyrosine residues in toxicological studies of the thyroid 553(1991)135
- Doorn, L., see Jansen, E. H. J. M. 553(1991)123
- Doorn, L., see Jansen, E. H. J. M. 566(1991)471
- Doorn, L., see Reinerink, E. J. M. 553(1991)233
- Doornbos, D. A., see Wieling, J. 594(1992)45
- Doornbos, D. A., see Witte, D. T. 558(1991)333
- Dorbon, M., Lamaison, S. and Chevalier, A.
Simulated distillation of distillates on capillary columns: influence of the polarity of the stationary phase 557(1991)155
- Doretta, L., see Sturaro, A. 590(1992)223
- Dorizzi, R., Michelot, D., Tagliaro, F. and Ghielmi, S.
Methods for chromatographic determination of amanitins and related toxins in biological samples (Review) 580(1992)279
- Dorizzi, R., see Tagliaro, F. 563(1991)11
- Dorizzi, R., see Tagliaro, F. 566(1991)333
- Dorley, J., see Chen, I.-W. 576(1992)363
- Dosi, L., see Italia, A. 553(1991)15
- Dos Ramos, F. J., Noronha da Silveira, I. and De Graaf, G.
Column liquid chromatographic determination of carbadox and olaquinox in feeds 558(1991)125
- Dostert, P., see Allievi, C. 568(1991)271
- Dostert, P., see Niwa, T. 578(1992)109
- Dovich, N. J., see Chen, D. Y. 559(1991)237
- Dowle, C. J., see Robertson, A. M. 588(1991)15
- Downham, M., Busby, S., Jefferis, R. and Lyddiatt, A.
Immunoaffinity chromatography in biorecovery: an application of recombinant DNA technology to generic adsorption processes 584(1992)59
- Dowsett, M., see Poon, G. K. 565(1991)75
- Draeger, N. M., see Chase, H. A. 597(1992)129
- Drawert, F., see Nitz, S. 557(1991)187
- Drehmel, D. C., see Lindgren, E. R. 585(1991)353
- Drenth, B. F. H., see Witte, D. T. 553(1991)365
- Dressler, M., see Cigánek, M. 588(1991)225
- Dreux, M., see Brossard, S. 591(1992)149
- Dreyer, U., Melzer, H. and Möckel, H. J.
Increasing data precision in reversed-phase liquid chromatography leads to new information level 592(1992)13
- Driessen, C., see Leyssens, L. 564(1991)515
- Driessen, J. J. M., see De Jong, J. 553(1991)243
- Drilleau, J.-F., see Delage, E. 555(1991)125
- Driscoll, K. E., see Neubecker, T. A. 574(1992)340

- Driskell, J. A., Mc. Chrisley, B., Reynolds, L. K. and Moak, S. W.
Plasma B₆ vitamer and plasma and urinary 4-pyridoxic acid concentrations of middle-aged obese black women 568(1991)333
- Driskell, J. A., see Chrisley, B. M. 563(1991)369
- Driss, M. R., Sabbah, S. and Bouguerra, M. L.
Chromatographie en phase gazeuse haute résolution-spectrométrie de masse des congénères de biphényles polychlorés dans quelques échantillons d'origine biologique 552(1991)213
- Drogue, S., Rolet, M.-C., Thiébaud, D. and Rosset, R.
Separation of pristinamycins by high-speed counter-current chromatography. I. Selection of solvent system and preliminary preparative studies 593(1992)363
- Drohan, W., see Tharakan, J. 595(1992)103
- Drusany, I., see Prošek, M. 553(1991)477
- Du, P., see Massé, R. 562(1991)323
- Dua, V. K., Sarin, R. and Sharma, V. P.
Determination of sulfalene in plasma, red blood cells and whole blood by high-performance liquid chromatography 563(1991)333
- Duafala, M. E., see Smith-Rogers, J. A. 566(1991)147
- Dubiel, W., see Ziegler, M. 563(1991)172
- Dubrow, R., see Demorest, D. 559(1991)43
- Dubs, M.-C., Altschuh, D. and Van Regenmortel, M. H. V.
Mapping of viral epitopes with conformationally specific monoclonal antibodies using biosensor technology 597(1992)391
- Duch, D. S., see Mullin, R. J. 555(1991)254
- Ducharme, J., Varin, F., Bevan, D. R., Donati, F. and Théorêt, Y.
High-performance liquid chromatography-electrochemical detection of vecuronium and its metabolites in human plasma 573(1992)79
- Duchateau, A. L. L., Hillemans, M. G., Schepers, C. H. M., Ketelaar, P. E. F., Hermes, H. F. M. and Kamphuis, J.
Determination of biocatalyst consumption in an aminopeptidase process using automated sample preparation and high-performance liquid chromatography 566(1991)493
- Duchateau, A. L. L., Munsters, B. H. M., Kwakkenbos, G. T. C. and Van Leuken, R. G. J.
Selection of buffers and of an ion-pairing agent for thermospray liquid chromatographic-mass spectrometric analysis of ionic compounds 552(1991)605
- Duché, J.-C., see Hervé, F. 577(1992)43
- Ducros, V. and Favier, A.
Gas chromatographic-mass spectrometric method for the determination of selenium in biological samples 583(1992)35
- Ducrotot, G., Richert, L., De Sandro, V., Lurier, D. and Pacaud, E.
Determination of ¹²⁵I-labelled thyroxine glucuronide by reversed-phase high-performance liquid chromatography using on-line radiochemical detection to determine UDPglucuronosyltransferase activity 566(1991)415
- Dudley, R. L., see Sonnet, P. E. 586(1991)255
- Dufay, C., Lecron, J. C., Daudon, F. and Gombert, J.
Rapid and simple procedure for the determination of salsolinol in urine using high-performance liquid chromatography with electrochemical detection 563(1991)224
- Dufour, A., see Julien-Larose, C. 562(1991)39
- Dugal, R., see Massé, R. 562(1991)323
- Dugan, M. E. R., Sauer, W. C., Lien, K. A. and Fenton, T. W.
Ion-pair high-performance liquid chromatography of diaminopimelic acid in hydrolysates of physiological samples 582(1992)242
- Duke, C. C., see Carter, S. R. 574(1992)77
- Ďulák, K., see Petro, M. 595(1992)337
- Dulery, B., see Berg-Candolfi, M. 593(1992)1
- Dulery, B. D., Schoun, J. and Haeghele, K. D.
Determination of an irreversible inhibitor of monoamine oxidase B (MDL 72974A) in human plasma and urine by gas chromatography-positive-ion chemical ionization mass spectrometry 571(1991)241
- Dülffer, T., see Puhlmann, A. 581(1992)129
- Dumasia, M. C. and Houghton, E.
Screening and confirmatory analysis of β -agonists, β -antagonists and their metabolites in horse urine by capillary gas chromatography-mass spectrometry 564(1991)503
- Duncan, G. F., see Franc, J. E. 570(1991)129
- Dunlap, R. B., see Terry, Jr., A. V. 585(1991)101
- Dunn, A. J., see Dempsey, C. A. 596(1992)110
- Dunn, D. L., see Sternitzke, K. D. 589(1992)159
- Dunn, L. C., see Talmadge, K. W. 590(1992)83
- Dunn, M. J. and Hahn, D. A.
Aqueous high-performance size-exclusion chromatographic assay for high-molecular-weight impurities in ceftiofur sodium 595(1992)185
- Dunnett, M. and Harris, R. C.
Determination of carnosine and other biogenic imidazoles in equine plasma by isocratic reversed-phase ion-pair high-performance liquid chromatography 579(1992)45
- Durand, G., De Bertrand, N. and Barceló, D.
Applications of thermospray liquid chromatography-mass spectrometry in photochemical studies of pesticides in water 554(1991)233
- Durand, G., De Bertrand, N. and Barceló, D.
Mobile phase variations in thermospray liquid chromatography-mass spectrometry of pesticides 562(1991)507
- Durand, G., see Barceló, D. 553(1991)311
- Durhan, E., see Lukasewycz, M. 580(1992)215
- Dürsch, I., see Meyer, H. H. D. 564(1991)551
- Dusick, A., see Browne, S. P. 575(1992)158
- Duthel, J. M., Constant, H., Vallon, J. J., Rochet, T. and Miachon, S.
Quantitation by gas chromatography with selected-ion monitoring mass spectrometry of "natural" diazepam, N-desmethyldiazepam and oxazepam in normal human serum 579(1992)85

- Duthel, J. M. and Vallon, J. J.
Indolic urinary melanogens: separation and identification by gas chromatography with selected-ion monitoring mass spectrometry of 5-hydroxy-6-methoxyindole-2-carboxylic and 5-methoxy-6-hydroxyindole-2-carboxylic acids 570(1991)166
- Duthu, G. S., see Shepard, R. M. 565(1991)321
- Dutoit, J.-C.
Gas chromatographic retention behaviour of some solutes on structurally similar polar and non-polar stationary phases 555(1991)191
- Dutta, P. K., Shanley, M. and O'Donovan, G. A.
Monitoring the accumulation of nucleoside triphosphates by high-performance liquid chromatography 599(1992)137
- Dutton, M. V., Rastall, R. A. and Evans, C. S.
Improved high-performance liquid chromatographic separation for the analysis of oxalate in fungal culture media 587(1991)297
- Dyas, A. M., Robinson, M. L. and Fell, A. F.
Direct separation of nadolol enantiomers on a Pirkle-type chiral stationary phase 586(1991)351
- Dybczyński, R.
Role of ion-exchange and extraction chromatography in neutron activation analysis (Review) 600(1992)17
- Dyer, K. D., see Alliegro, M. A. 582(1992)217
- Dyr, J. E. and Suttner, J.
Affinity high-performance liquid chromatography on heparin-Glc-Spheron 563(1991)124
- Dyr, J. E., see Suttner, J. 563(1991)184
- Dzido, T. H., see Waksmundzka-Hajnos, M. 600(1992)51
- E**
- Ebenstein, D. B., see Quinn Baumann, P. 573(1992)11
- Ebermann, R., see Pichorner, H. 594(1992)400
- Ebinger, G., see Sarre, S. 575(1992)207
- Ebinger, G., see Sarre, S. 582(1992)29
- Eckardt, C. B., Keely, B. J. and Maxwell, J. R.
Identification of chlorophyll transformation products in a lake sediment by combined liquid chromatography-mass spectrometry 557(1991)271
- Eckelman, W. C., see Rosenspire, K. C. 574(1992)119
- Eckerbom, S., see Bergqvist, Y. 571(1991)169
- Eckert, G., see Ba, B. 572(1991)277
- Eckert-Tilotta, S. E., Hawthorne, S. B. and Miller, D. J.
Comparison of commercially available atomic emission and chemiluminescence detectors for sulfur-selective gas chromatographic detection 591(1992)313
- Eddy, C. L., see Rodriguez, P. A. 563(1991)271
- Edeki, T., Robin, D. W., Prakash, C., Blair, I. A. and Wood, A. J. J.
Sensitive assay for triazolam in plasma following low oral doses 577(1992)190
- Eder, C., see Küsters, E. 576(1992)179
- Eder, K., Reichlmayr-Lais, A. M. and Kirchgessner, M.
Gas chromatographic analysis of fatty acid methyl esters: avoiding discrimination by programmed temperature vaporizing injection 588(1991)265
- Eder, K., Reichlmayr-Lais, A. M. and Kirchgessner, M.
Simultaneous determination of amounts of major phospholipid classes and their fatty acid composition in erythrocyte membranes using high-performance liquid chromatography and gas chromatography 598(1992)33
- Edfort, M. J., see Kvalo, L. T. 565(1991)391
- Edlund, P. O. and Johansen, K.
Clean-up of plasma extracts by gel permeation chromatography during analysis of isosorbide nitrates by capillary gas chromatography 553(1991)21
- Edmonds, C. G., see Smith, R. D. 559(1991)197
- Edwards, G., see Hoaksey, P. E. 566(1991)244
- Edwards, G., see Thomas, C. G. 583(1992)131
- Edwards, R. H. T., see Leyland, D. M. 581(1992)179
- Effland, R. C., see Hsu, R. S. 572(1991)352
- Efthymiopoulos, C., see Lave, T. 572(1991)203
- Egan, D. A. and O'Kennedy, R.
Rapid and sensitive determination of coumarin and 7-hydroxycoumarin and its glucuronide conjugate in urine and plasma by high-performance liquid chromatography 582(1992)137
- Egan, W., see Bergot, B. J. 599(1992)35
- Egeler, E., see Jacob, K. 572(1991)317
- Eggers, C., see Liu, H. 576(1992)163
- Eggleston, D. S., see Chen, T. K. 596(1992)123
- Eguchi, M., see Johnson, D. F. 594(1992)283
- Ehwald, R., see Jäschke, A. 585(1991)57
- Ehwald, R., see Jäschke, A. 596(1992)165
- Einarsson, S., see Orwar, O. 566(1991)39
- Eisenbeiss, A., Reuke, S. and Türck, M.
Determination of diacetoneketogulonic acid in water samples by high-performance thin-layer chromatography 589(1992)390
- Eisman, M. S., see Wilson, T. C. M. 582(1992)195
- Eisner, E., see Pachinger, A. 558(1991)369
- Eiteman, M. A. and Gainer, J. L.
Predicting partition coefficients in polyethylene glycol-potassium phosphate aqueous two-phase systems 586(1991)341
- Eitoku, F., see Watanabe, K. 566(1991)109
- Ekberg, H., see Eriksson, T. 582(1992)211
- El-Kak, A. and Vijayalakshmi, M. A.
Study of the separation of mouse monoclonal antibodies by pseudobioaffinity chromatography using matrix-linked histidine and histamine 570(1991)29
- El-Shahawi, M. S., see Farag, A. B. 552(1991)371
- Elchuk, S., Burns, K. I., Cassidy, R. M. and Lucy, C. A.
Reversed-phase separation of transition metals, lanthanides and actinides by elution with mandelic acid 558(1991)197

- El Fallah, M. Z. and Martin, M.
Quantitative determination limit in chromatography: computer-based simulations 557(1991)23
- El Khabchi, M., see Crétier, G. 596(1992)15
- El'kin, Y. N., see Voinov, V. G. 586(1991)360
- Elliott, W. H., see Ferreira, H. E. C. S. 562(1991)697
- Elliott, W. H., see Shoemaker, J. D. 562(1991)125
- Ellis, P. P., see Riegel, M. R. 568(1991)467
- Elmroth, I., Larsson, L., Westerdahl, G. and Odham, G.
Determination of muramic acid by high-performance liquid chromatography-plasma spray mass spectrometry 598(1992)43
- Eloranta, T. O., see Hyvönen, T. 574(1992)17
- El Rassi, Z., see Nashabeh, W. 559(1991)367
- El Rassi, Z., see Nashabeh, W. 596(1992)251
- El Rassi, Z., see Nashabeh, W. 600(1992)279
- Elsom, L. F., see Cheng, K. N. 581(1992)203
- El Tayar, N., Tsai, R.-S., Vallat, P., Altomare, C. and Testa, B.
Measurement of partition coefficients by various centrifugal partition chromatographic techniques. A comparative evaluation 556(1991)181
- Eltekov, Y. A., see Dernovaya, L. A. 552(1991)365
- Elzinga, H., see Hessels, J. 563(1991)1
- Emmerling, M., see Glässgen, W. E. 598(1992)81
- Emmrich, F., see Guse, A. H. 593(1992)157
- Endert, E., see Takkenberg, B. 565(1991)430
- Endo, J., see Tabata, M. 597(1992)435
- Endo, T., see Kobata, A. 597(1992)111
- Endoh, Y. S., Yoshimura, H., Sasaki, N., Ishihara, Y., Sasaki, H., Nakamura, S., Inoue, Y. and Nishikawa, M.
High-performance liquid chromatographic determination of pamaquine, primaquine and carboxy primaquine in calf plasma using electrochemical detection 579(1992)123
- Engelhardt, H., see Bentrop, D. 556(1991)363
- Englert, G., see Khachik, F. 582(1992)153
- Enjalbert, F., Gallion, C., Jehl, F., Monteil, H. and Faulstich, H.
Simultaneous assay for amatoxins and phallotoxins in *Amanita phalloides* Fr. by high-performance liquid chromatography 598(1992)227
- Ensing, K., see Witte, D. T. 553(1991)365
- Epemolu, R. O., Singh, S., Hider, R. C. and Damani, L. A.
High-performance liquid chromatographic determination of 1,2-diethyl-3-hydroxypyridin-4-one and its 2-(1-hydroxyethyl) metabolite in rat blood 573(1992)178
- Epler, K. S., Sander, L. C., Ziegler, R. G., Wise, S. A. and Craft, N. E.
Evaluation of reversed-phase liquid chromatographic columns for recovery and selectivity of selected carotenoids 595(1992)89
- Epstein, A. L., see Chen, F.-M. 563(1991)243
- Erber, H., see Aoyama, R. G. 588(1991)181
- Erceg, M., see Caruso, U. 562(1991)147
- Erdin, R., see Timm, U. 593(1992)63
- Erdmann, G. R., Steury, J. C., Carleton, B. C., Stafford, R. J., Bostrom, B. C. and Canafax, D. M.
Reversed-phase high-performance liquid chromatographic approach to determine total lymphocyte concentrations of 6-thioguanine, methylmercaptopyrine and methylthioguanine in humans 571(1991)149
- Erhard, K. F., see Chen, T. K. 596(1992)123
- Eriksson, A. W., see Bank, R. A. 571(1991)47
- Eriksson, B.-M. and Wikström, M.
Determination of vanilmandelic acid in urine by coupled-column liquid chromatography combining affinity to boronate and separation by anion exchange 567(1991)1
- Eriksson, B.-M. and Wikström, M.
Determination of catecholamines in urine by liquid chromatography and electrochemical detection after on-line sample purification on immobilized boronic acid 593(1992)185
- Eriksson, T., Björkman, S., Fyge, Å. and Ekberg, H.
Determination of thalidomide in plasma and blood by high-performance liquid chromatography: avoiding hydrolytic degradation 582(1992)211
- Erlandsson, P., see Katti, A. 590(1992)127
- Erlandsson, P., see Marle, I. 586(1991)233
- Erler, U. and Heublein, G.
Synthesis and characterization of stationary phases on the basis of silicas modified with epoxidized polybutadienes. IV. Chromatographic experiments on a new amino-functionalized stationary phase for high-performance liquid chromatography 588(1991)340
- Erni, F.
Foreword 592(1992)1
- Erni, F., see Liu, G. 592(1992)239
- Erni, F., see Liu, G. 598(1992)153
- Erni, F., see Yamamoto, H. 593(1992)313
- Ernould, A.-P., see Boutin, J. A. 583(1992)137
- Escarzaga, M., see Watts, M. T. 577(1992)289
- Eschalié, A., see Coudore, F. 584(1992)249
- Escott, R. E. A., McDowell, P. G. and Porter, N. P.
Use of non-volatile ion-pairing agents for liquid chromatographic-mass spectrometric analyses with a moving-belt interface 554(1991)281
- Escott, R. E. A. and Mortimer, N.
Analysis of polyethylene glycol and derivatives by high-performance liquid chromatography using elevated temperatures and low-wavelength ultraviolet detection, and supercritical fluid chromatography 553(1991)423
- Essayegh, M., see Desbène, P. L. 553(1991)211
- Essink, A. W. G., Lohuis, C. P. G. G., Klein Elhorst, J. T. and Rutten, W. J.
Selective and quantitative isolation and determination of apomorphine in human plasma 570(1991)419
- Esteban, A., Graells, M., Satorre, J. and Pérez-Mateo, M.
Determination of paracetamol and its four major metabolites in mouse plasma by reversed-phase ion-pair high-performance liquid chromatography 573(1992)121
- Estiva, E. C., see Peterson, J. C. 564(1991)205
- Eto, S., Noda, H. and Noda, A.
High-performance liquid chromatographic method for direct separation of 5-(*p*-hydroxyphenyl)-5-phenylhydantoin enantiomers using a chiral tris(4-methylbenzoate) column 568(1991)157
- Eto, S., Noda, H. and Noda, A.
Chiral separation of barbiturates and hydantoins by reversed-phase high-performance liquid chromatography using a 25 or 50 mm short ODS cartridge column via β -cyclodextrin inclusion complexes 579(1992)253

- Etoh, T., Miyazaki, M., Harada, K., Nakayama, M. and Sugii, A.
Rapid analysis of human serum albumin by high-performance liquid chromatography 578(1992)292
- Ettori, C., see Chiari, M. 559(1991)119
- Ettre, L. S., see Meyer, V. R. 600(1992)3
- Evans, C. S., see Dutton, M. V. 587(1991)297
- Evans, L., see Mitchell, E. P. 581(1992)31
- Evans, M. B., Bird, R. and O'Sullivan, H.
Gas chromatography in qualitative analysis. XX. The deactivation of diatomaceous supports by batyl alcohol 552(1991)169
- Evans, M. B., see Dale, A. D. 552(1991)161
- Evans, P. D., see Macfarlane, R. G. 562(1991)585
- Evans, R. R. and Relling, M. V.
Automated high-performance liquid chromatographic assay for the determination of 7-ethoxycoumarin and umbelliferone 578(1992)141
- Everaerts, F. M., see Ackermans, M. T. 585(1991)123
- Everaerts, F. M., see Ackermans, M. T. 590(1992)341
- Everaerts, F. M., see Ackermans, M. T. 595(1992)327
- Everaerts, F. M., see Ackermans, M. T. 596(1992)101
- Evershed, R. P., Prescott, M. C. and Goad, L. J.
Deuteration as an aid to the high-temperature gas chromatography-mass spectrometry of steryl fatty acyl esters 590(1992)305
- Evershed, R. P., see Leyland, D. M. 581(1992)179
- Ezra, I., see Massil, S. E. 589(1992)375
- F**
- Fahimi, A., see Snini, A. 590(1992)369
- Falick, A. M., see Shackleton, C. H. L. 562(1991)175
- Fan, C.-D., Zhao, H. and Chow, M. S. S.
Simple and rapid high-performance liquid chromatographic assay for esmolol 570(1991)217
- Fan, D.-P., see Wang, Q.-S. 587(1991)359
- Fan, T. Y., see Sternitzke, K. D. 589(1992)159
- Fanali, S., see Pessi, A. 557(1991)307
- Fanali, S., see Quaglia, M. G. 593(1992)259
- Fanali, S., see Sakodinskaya, I. K. 596(1992)95
- Fanelli, R., see Bagnati, R. 564(1991)493
- Fang, F. W., Aguilar, M. I. and Hearn, M. T. W.
High-performance liquid chromatography of amino acids, peptides and proteins. CXX. Evaluation of bandwidth behaviour of proteins chromatographed on tentacle-type anion exchangers 599(1992)163
- Fang, S., see Lebl, M. 586(1991)145
- Fang, X.-H., see Zeng, Z.-R. 589(1992)368
- Fantazzini, C., see Marcato, B. 553(1991)415
- Farag, A. B. and El-Shahawi, M. S.
Removal of organic pollutants from aqueous solution. V. Comparative study of the extraction, recovery and chromatographic separation of some organic insecticides using unloaded polyurethane foam columns 552(1991)371
- Farina, M., see Rossini, M. L. 593(1992)47
- Farinotti, R., see Fernandez, C. 572(1991)195
- Farinotti, R., see Tod, M. 594(1992)386
- Farjam, A., Van de Merbel, N. C., Nieman, A. A., Lingeman, H. and Brinkman, U. A. T.
Determination of aflatoxin M1 using a dialysis-based immunoaffinity sample pretreatment system coupled on-line to liquid chromatography. Reusable immunoaffinity columns 589(1992)141
- Farkaš, P., see Soják, L. 557(1991)241
- Farmen, R. H., see Franc, J. E. 570(1991)129
- Fasolo, M., see Dalla Libera, L. 566(1991)427
- Fassina, G.
Oriented immobilization of peptide ligands on solid supports 591(1992)99
- Fast, D. M., see Burse, V. W. 566(1991)117
- Fathi, M., Tsacopoulos, M., Raverdino, V. and Porthault, M.
Separation of nucleotides in homogenates of octopus retina by ion-pair reversed-phase liquid chromatography and identification by mass spectrometry 563(1991)356
- Fatmi, A. A., see Pullen, R. H. 574(1992)101
- Fauler, J., see Tsikas, D. 574(1992)181
- Faulstich, H., see Enjalbert, F. 598(1992)227
- Fauntleroy, L. P., see Thedford, R. 582(1992)253
- Favarato, M., Mizzen, C. A. and McLachlan, D. R.
Resolution of serum aluminum-binding proteins by size-exclusion chromatography: identification of a new carrier of aluminum in human serum 576(1992)271
- Favier, A., see Arnaud, J. 572(1991)103
- Favier, A., see Ducros, V. 583(1992)35
- Favier, A., see Richard, M. J. 577(1992)9
- Fay, L., see Badoud, R. 552(1991)345
- Fábián, V., see Morvai, M. 600(1992)87
- Fábián, V., Morvai, M., Pintér-Szakács, M. and Molnár-Perl, I.
Standardization of cation-exchange clean-up prior to gas chromatography of amino acids 553(1991)87
- Fadeev, A. Y., Lisichkin, G. V., Runov, V. K. and Staroverov, S. M.
Diffusion of sorbed pyrene in the bonded layer of reversed-phase silicas. Effect of alkyl chain length and pore diameter 558(1991)31
- Fadeev, A. Y., Mingalyov, P. G., Staroverov, S. M., Lunina, E. V., Lisichkin, G. V., Gaida, A. V. and Monastyrsky, V. A.
Silica sorbents with one- and two-site attached bacitracin in affinity chromatography 596(1992)114
- Fägerstam, L. G., Frostell-Karlsson, Å., Karlsson, R., Persson, B. and Rönnberg, I.
Biospecific interaction analysis using surface plasmon resonance detection applied to kinetic, binding site and concentration analysis 597(1992)397

- Fayer, B. E., Huselton, C. A., Garland, W. A. and Liberato, D. J.
Quantification of acitretin in human plasma by microbore liquid chromatography-negative chemical ionization mass spectrometry 568(1991)135
- Feather, T., see Kuksis, A. 564(1991)11
- Feather, T., see Kuksis, A. 579(1992)13
- Fedorak, P. M. and Andersson, J. T.
Decomposition of two methylbenzothiophene sulphoxides in a commercial gas chromatography injection port liner 591(1992)362
- Feigl, E. O., see Herrmann, S. C. 574(1992)247
- Feijlbrief, M., see Welling, G. W. 599(1992)157
- Feinberg, H., see Sarin, M. 563(1991)129
- Feineis, D., see Bringmann, G. 595(1992)351
- Feldman, B. J., Mogadeddi, H. and Osterloh, J. D.
Determination of Pb²⁺ in water by isotope dilution gas chromatography-mass spectrometry of tetraethyllead formed by reaction with sodium tetraethylborate 594(1992)275
- Feliciani, R., see Milana, M. R. 552(1991)205
- Felinger, A. and Guiochon, G.
Optimization of the experimental conditions and the column design parameters in overloaded elution chromatography 591(1992)31
- Fell, A. F., see Castledine, J. B. 592(1992)27
- Fell, A. F., see Dyas, A. M. 586(1991)351
- Fell, A. F., see Zimina, T. M. 593(1992)233
- Feng, N., Minder, E. I., Grampp, T. and Vonderschmitt, D. J.
Identification and quantification of ergotamine in human plasma by gas chromatography-mass spectrometry 575(1992)289
- Fennessey, P. V., see Loy, G. L. 562(1991)169
- Fenske, M. and Schönheiter, H.
Thin-layer chromatography on silica-coated aluminium sheet as an adjunct to radioimmunoassay of steroids 563(1991)178
- Fenton, T. W., see Dugan, M. E. R. 582(1992)242
- Fermo, I., Arcelloni, C., De Vecchi, E., Vigano', S. and Paroni, R.
High-performance liquid chromatographic method with fluorescence detection for the determination of total homocyst(e)ine in plasma 593(1992)171
- Fernandez, C., Baune, B., Gimenez, F., Thuillier, A. and Farinotti, R.
Determination of zopiclone enantiomers in plasma by liquid chromatography using a chiral cellulose carbamate column 572(1991)195
- Fernández Cruz, E., see Ramis, I. 557(1991)507
- Fernández-Sánchez, E., Fernández-Torres, A., García-Domínguez, J. A. and Salvador-Moya, M. D.
Gas chromatographic comparative study of Superox 20M immobilized in different ways 556(1991)485
- Fernández-Torres, A., see Fernández-Sánchez, E. 556(1991)485
- Ferraina, R. A., see Shepard, R. M. 565(1991)321
- Ferrari, V., see Cracco, A. T. 574(1992)156
- Ferrario, P., see Zecca, L. 567(1991)425
- Ferre, S., see Parra, P. 570(1991)185
- Ferreira, H. E. C. S. and Elliott, W. H.
Pre-column derivatization of free bile acids for high-performance liquid chromatographic and gas chromatographic-mass spectrometric analysis 562(1991)697
- Ferretti, R., see Betto, P. 563(1991)115
- Ferris, R., Missbichler, A., Hartter, E., Mann-Buxbaum, E., Schalkhammer, T., Urban, G. and Pittner, F.
Construction and use of two α -human atrial natriuretic peptide-fragment affinity chromatography columns in the isolation of C- and N-terminal epitope-specific antibodies for use in a prototype α -hANP biosensor 577(1992)251
- Ferro, R., see Silvestro, L. 563(1991)443
- Ferry, G., see Boutin, J. A. 583(1992)137
- Ferry, S., see Bouliou, R. 567(1991)480
- Ferry, S., see Bouliou, R. 571(1991)331
- Feste, A. S.
Reversed-phase chromatography of phenylthiocarbamyl amino acid derivatives of physiological amino acids: an evaluation and a comparison with analysis by ion-exchange chromatography 574(1992)23
- Fettinger, J. C., see Manz, A. 593(1992)253
- Fetzer, J. C., see Jinno, K. 557(1991)459
- Fetzer, J. C., see Jinno, K. 594(1992)105
- Fey, M. L., see Rodriguez, P. A. 563(1991)271
- Fialip, J., see Coudore, F. 584(1992)249
- Fieger, H. and Blaschke, G.
Direct determination of the enantiomeric ratio of verapamil, its major metabolite norverapamil and gallopamil in plasma by chiral high-performance liquid chromatography 575(1992)255
- Fielden, P. R., see Packham, A. J. 552(1991)575
- Fielding, R. M., see Wang, L. H. 579(1992)259
- Fierabracci, V., Masiello, P., Novelli, M. and Bergamini, E.
Application of amino acid analysis by high-performance liquid chromatography with phenyl isothiocyanate derivatization to the rapid determination of free amino acids in biological samples 570(1991)285
- Filipek, M., see Paczkowski, D. 573(1992)235
- Filippich, L. J., see Zhu, J. 577(1992)77
- Finegan, B. A., see Hussain, M. D. 582(1992)203
- Finlay, D. C., see Mills, M. H. 564(1991)93
- Fiorella, P. L., see Grossi, G. 593(1992)217
- Fiorelli, G., see Pittalis, S. 573(1992)29
- Fiorini, F., see Garibotto, G. 572(1991)11
- Firby, P., see Matassa, L. C. 573(1992)43
- Firmin, J. L., see Price, N. P. J. 598(1992)51
- Firth, J., see Poon, G. K. 565(1991)75
- Fischer, C.-H.
Trace analysis of phthalocyanine pigments by high-performance liquid chromatography 592(1992)261
- Fisher, D. H., see Li, W. 588(1991)273
- Flakoll, P. J., see Donahue, E. P. 571(1991)29
- Flament, I., see Wünsche, L. 552(1991)539
- Flaminio, L., see Ascalone, V. 581(1992)237
- Flaschel, E., see Andriamboavonjy, E. 587(1991)288
- Fleminger, G., Neufeld, T., Star-Weinstock, M., Litvak, M. and Solomon, B.
Calcium-modulated conformational affinity chromatography. Application to the purification of calmodulin and S100 proteins 597(1992)263

- Fleminger, G., see Solomon, B. 597(1992)257
- Flesch, G., Francotte, E., Hell, F. and Degen, P. H.
Determination of the *R*-(-) and *S*-(+) enantiomers of the monohydroxylated metabolite of oxcarbazepine in human plasma by enantioselective high-performance liquid chromatography 581(1992)147
- Flesch, G., Tominaga, N. and Degen, P.
Improved determination of the bisphosphonate pamidronate disodium in plasma and urine by pre-column derivatization with fluorecamine, high-performance liquid chromatography and fluorescence detection 568(1991)261
- Fletcher, R. J., see Luong, C. B. H. 584(1992)77
- Flipo, D., see Desmangles, R. J. P. 584(1992)121
- Florance, J. R., Konteatis, Z. D., Macielag, M. J., Lessor, R. A. and Galdes, A.
Capillary zone electrophoresis studies of motilin peptides. Effects of charge, hydrophobicity, secondary structure and length 559(1991)391
- Flouvat, B., see Delhotal Landes, B. 577(1992)117
- Flouvat, B., see Roux, A. 570(1991)453
- Flynn, J. W. and O'Brien, J. E.
High-performance liquid chromatographic method for the simultaneous measurement of remacemide (a novel anticonvulsant and cerebroprotectant) and an active metabolite in human plasma 583(1992)91
- Fogiel, A. J., see Reiser, R. W. 554(1991)91
- Foglia, J. P., Sorisio, D. and Perel, J. M.
Determination of imipramine, desipramine and their hydroxy metabolites by reversed-phase chromatography with ultraviolet and coulometric detection 572(1991)247
- Folestad, S., see Orwar, O. 566(1991)39
- Foley, J. P., see Jeansonne, M. S. 594(1992)1
- Foltz, R. L., see Nelson, C. C. 580(1992)97
- Font, G., see Moltó, J. C. 555(1991)137
- Fontanarava, E., see Masson, S. 563(1991)231
- Forgács, E. and Cserhádi, T.
High-performance liquid chromatographic retention behaviour of ring-substituted aniline derivatives on a porous graphitized carbon column 600(1992)43
- Forgács, E., Cserhádi, T. and Valkó, K.
Retention behaviour of some ring-substituted phenol derivatives on a porous graphitized carbon column 592(1992)75
- Forges, N., see Tellez, S. 581(1992)257
- Forman, L. J., see Liu, H. 576(1992)163
- Forni, E., Polesello, A., Montefiori, D. and Maestrelli, A.
High-performance liquid chromatographic analysis of the pigments of blood-red prickly pear (*Opuntia ficus indica*) 593(1992)177
- Forrow, S., see Preece, N. E. 573(1992)227
- Fortis, I., see Arnaud, J. 572(1991)103
- Fossati, T., Parisi, S., Abbiati, G. and Castiglioni, C.
Determination of picotamide in human plasma and urine by high-performance liquid chromatography 577(1992)382
- Fossati, T., see Abbiati, G. 566(1991)445
- Foster, R. T., see Bhatti, M. M. 579(1992)361
- Foster, R. T., see Carr, R. A. 566(1991)155
- Fougnion, J. M., David, C., Crocq, A. and Genty, C.
Optimisation pentaparamétrique de la résolution en chromatographie phase gazeuse par la technique du diagramme à fenêtres. Cas du couplage de deux colonnes 556(1991)287
- Foulger, B. E., see Begley, P. 588(1991)239
- Fournier, M., see Desmangles, R. J. P. 584(1992)121
- Fourtillan, J. B., see Girault, J. 564(1991)43
- Fowler, B., see Caruso, U. 562(1991)147
- Fowler, J. S., see MacGregor, R. R. 590(1992)354
- Fowler, W. K., Stewart, D. C., Weinberg, D. S. and Sarver, E. W.
Gas chromatographic determination of the lewisite hydrolysate, 2-chlorovinylarsonous acid, after derivatization with 1,2-ethanedithiol 558(1991)235
- Fraisse, D., see Ventura, R. 562(1991)723
- Franc, J. E., Duncan, G. F., Farmen, R. H. and Pittman, K. A.
High-performance liquid chromatographic method for the determination of nefazodone and its metabolites in human plasma using laboratory robotics 570(1991)129
- Franceschetti, F., see Pellegatti, M. 573(1992)105
- Franchi, D., see Tagliaro, F. 580(1992)161
- Francis, H. W., see Jack, D. S. 584(1992)189
- Francois, B., see Daumas, L. 570(1991)89
- Francotte, E. and Junker-Buchheit, A.
Preparative chromatographic separation of enantiomers (Review) 576(1992)1
- Francotte, E. and Wolf, R. M.
Chromatographic resolution on methylbenzoylcellulose beads. Modulation of the chiral recognition by variation of the position of the methyl group on the aromatic ring 595(1992)63
- Francotte, E., see Flesch, G. 581(1992)147
- Frank, D., see Allen, E. 578(1992)175
- Frantsuzova, N. A., see Klyashchitsky, B. A. 577(1992)267
- Frassanito, R., see Celardo, A. 568(1991)407
- Frazier, D., see Cox, S. K. 564(1991)322
- Fredericksen, T. L., see Burton, S. D. 571(1991)209
- Fredrickson, H. L., see Klöppel, K.-D. 562(1991)369
- Freeman, D., see Bouquillon, A. I. 577(1992)354
- Freeman, K. H., see Lichtfouse, E. 585(1991)177
- Frei, R. W., see Özçimder, M. 570(1991)19
- Freitag, D., see Carr, R. A. 566(1991)155
- Frenz, J., Quan, C. P., Hancock, W. S. and Bourell, J.
Characterization of a tryptic digest by high-performance displacement chromatography and mass spectrometry 557(1991)289
- Frezzotti, D., see Bonifaci, L. 585(1991)333
- Friebe, S., Krauss, G. J. and Nitsche, H.
High-performance liquid chromatographic separation of *cis-trans* isomers of proline-containing peptides. I. Separation on cyclodextrin-bonded silica 598(1992)139
- Fried, K., see Lloyd, D. K. 578(1992)283
- Friedmann, E. I., see Matsumoto, G. I. 598(1992)267
- Friedrich, A., see Hagedorn, H.-W. 577(1992)195
- Friedrich, G., see Rissler, K. 565(1991)375
- Fritz, J. S. and Chen, J.
Ion exchange in analytical chemistry (by H. E. Walton and R. D. Rocklin) (Book Review) 555(1991)329

- Fritz, J.S., see Sun, J.J. 590(1992)197
 Fritz, R., see Cherton, J.-C. 566(1991)499
 Froger, M., see Teh, L.-C. 582(1992)65
 Frölich, J. C., see Tsikas, D. 574(1992)181
 Frostell-Karlsson, Å., see Fägerstam, L. G. 597(1992)397
 Frutiger, S., see Toussi, A. 582(1992)87
 Frydman, A., see Vergniol, J. C. 582(1992)273
 Fu, C., see Wang, Z. 589(1992)349
 Fuchs, D., see Werner, E. R. 570(1991)43
 Fuchs, M. J. and Keim, V.
 Separation of rat pancreatic secretory proteins by cation-exchange fast protein liquid chromatography 576(1992)287
 Fujima, H., see Haginaka, J. 592(1992)301
 Fujimaki, Y., Sudo, K. and Hakusui, H.
 Simultaneous determination of nefiracetam and its metabolites by high-performance liquid chromatography 575(1992)261
 Fujimoto, K., see Okamoto, J. 583(1992)53
 Fujimura, Y., see Hida, M. 585(1991)345
 Fujinari, E. M. and Courthaudon, L. O.
 Nitrogen-specific liquid chromatography detector based on chemiluminescence. Application to the analysis of ammonium nitrogen in waste water 592(1992)209
 Fujioka, T., Inoue, F., Sumita, S., Kuriyama, M., Hanawa, T. and Katagi, T.
 Purification and reconstitution of μ -opioid receptors in liposome 597(1992)429
 Fujita, Y., see Watanabe, H. 568(1991)445
 Fukamachi, M., see Kawaguchi, S. 567(1991)11
 Fukuda, H., Yamatodani, A., Imamura, I., Maeyama, K., Watanabe, T. and Wada, H.
 High-performance liquid chromatographic determination of histamine N-methyltransferase activity 567(1991)459
 Fukuda, M., see Koizumi, K. 585(1991)233
 Fukuda, M., see Koizumi, K. 595(1992)340
 Fukunaga, T., see Kane, M. 569(1991)297
 Fukuoka, H., see Takahara, E. 576(1992)174
 Fukushima, S., Akane, A., Matsubara, K., Shiono, H., Morishita, H. and Nakada, F.
 Simultaneous determination of testosterone and androstadienone (sex attractant) in human plasma by gas chromatography-mass spectrometry with high-resolution selected-ion monitoring 565(1991)35
 Fukuyama, T., see Nishi, H. 553(1991)503
 Fuller, R. W., see Murray, S. 567(1991)289
 Funaki, T., see Roosemalen, M. C. M. 565(1991)516
 Funakubo, H., see Yoshioka, M. 566(1991)361
 Fung, H.-L., see Tzeng, T.-B. 571(1991)109
 Funkenbusch, E. F., see Schafer, W. A. 587(1991)137
 Fuoco, R., see Colombini, M. P. 592(1992)255
 Furst, W. and Hallström, S.
 Simultaneous determination of myocardial nucleotides, nucleosides, purine bases and creatine phosphate by ion-pair high-performance liquid chromatography 578(1992)39
 Furtek, C. I. and Lo, M.-W.
 Simultaneous determination of a novel angiotensin II receptor blocking agent, losartan, and its metabolite in human plasma and urine by high-performance liquid chromatography 573(1992)295
 Furusaki, S., see Kim, M. 585(1991)45
 Furusaki, S., see Kim, M. 586(1991)27
 Furuta, T., Katayama, M., Shibasaki, H. and Kasuya, Y.
 Simultaneous determination of stable isotopically labelled L-histidine and urocanic acid in human plasma by stable isotope dilution mass spectrometry 576(1992)213
 Furuta, T., see Shibasaki, H. 576(1992)47
 Furuta, T., see Shibasaki, H. 579(1992)193
 Furuta, T., Takahashi, H. and Kasuya, Y.
 Reversible reaction via a carbanion intermediate in the elimination of ammonia from L-histidine catalysed by histidine ammonia-lyase 562(1991)363
 Fuss, L., see Massil, S. E. 589(1992)375
 Futani, S., see Yao, K. 581(1992)11
 Fyge, Å., see Eriksson, T. 582(1992)211
- ## G
- Gaboardi, F., see Bianchi-Bosisio, A. 569(1991)243
 Gabriele, R., see Betto, P. 594(1992)131
 Gabrielsson, J., see Hellenäs, K.-E. 573(1992)69
 Gabrielsson, M., Hoffmann, K.-J. and Regårdh, C. G.
 Determination of four carboxylic acid metabolites of felodipine in plasma by high-performance liquid chromatography 573(1992)265
 Gadde, R. R. and Burton, F. W.
 Simple reversed-phase high-performance liquid chromatographic method for 13-*cis*-retinoic acid in serum 593(1992)41
 Gadde, R. R., see DiNunzio, J. E. 594(1992)209
 Gaede, F.-P., Kirchner, M., Steinhilber, D. and Roth, H. J.
 Determination of picomole amounts of lipoxins C₄, D₄ and E₄ by high-performance liquid chromatography with electrochemical detection 585(1991)53
 Gaffar, S. A., see Chen, F.-M. 563(1991)243
 Gaggero, N., see Carrea, G. 600(1992)320
 Gagné, J.-P., Carrier, A. and Bertrand, M. J.
 Source of band broadening in liquid chromatographic-fast atom bombardment mass spectrometric systems with precolumn addition of viscous matrix to the mobile phase 554(1991)47
 Gagné, J.-P., Carrier, A. and Bertrand, M. J.
 Effect of the addition of viscous matrices to the mobile phase on chromatographic performance in liquid chromatography-fast atom bombardment mass spectrometry 554(1991)61
 Gagné, J.-P., Roussis, S. G. and Bertrand, M. J.
 Simple direct liquid introduction system usable as an interface for liquid chromatography-mass spectrometry on quadrupole and magnetic-sector mass spectrometers 554(1991)293
 Gagné, J.-P., see Carrier, A. 591(1992)129

- Gah, C., see Brückner, H. 555(1991)81
- Gaida, A. V., see Fadeev, A. Y. 596(1992)114
- Gaillard, Y., Gay-Montchamp, J. P. and Ollagnier, M.
Gas chromatographic determination of meprobamate in serum or plasma after solid-phase extraction 577(1992)171
- Gainer, J. L., see Eiteman, M. A. 586(1991)341
- Gajdůšková, V., see Chmel, K. 552(1991)449
- Gal, J., see Desai, D. M. 579(1992)165
- Galantino, M., see Rusconi, L. 589(1992)362
- Galdes, A., see Florance, J. R. 559(1991)391
- Gale, R. J., see Ghowsi, K. 559(1991)95
- Galeffi, C., see Betto, P. 594(1992)131
- Galensa, R., see Krause, M. 588(1991)41
- Galensa, R., see Mögele, R. 591(1992)165
- Galimberti, C., see Longo, A. 562(1991)639
- Galli Kienle, M., see Magni, F. 573(1992)127
- Gallion, C., see Enjalbert, F. 598(1992)227
- Gallion, C., see Molina, F. 563(1991)205
- Gallo, J. M., see Hassan, E. E. 582(1992)225
- Gallo, J. M., see Qian, M. 578(1992)316
- Gallois, A., see Morin, P. 586(1991)171
- Galushko, S. V.
Calculation of retention and selectivity in reversed-phase liquid chromatography 552(1991)91
- Galushko, S. V., Belik, M. Y., Solodenko, V. A., Kasheva, T. N. and Kukhar, V. P.
Normal- and reversed-phase high-performance liquid chromatography of some phosphonodipeptides 553(1991)143
- Galushko, S. V., Belik, M. Y., Solodenko, V. A., Kasheva, T. N. and Kukhar, V. P.
Ion-exchange high-performance liquid chromatography of diastereoisomers of some phosphonodipeptides 600(1992)79
- Galushko, S. V., Shishkina, I. P., Gerus, I. I. and Kolycheva, M. T.
High-performance ligand-exchange liquid chromatography of fluoro derivatives of alanine 600(1992)83
- Galushko, S. V., Shishkina, I. P. and Soloshonok, V. A.
High-performance ligand-exchange chromatography of some amino acids containing two chiral centres 592(1992)345
- Games, D. E., see Scalia, S. 574(1992)197
- Gan, L.-S., see Day, B. W. 562(1991)563
- Gandhi, T. P., see Mangalan, S. 567(1991)498
- Gandia, D., see Barilero, I. 575(1992)275
- Gandini, C., see Baruffini, A. 593(1992)95
- Gandini, C., see Kitsos, M. 553(1991)1
- Gandul-Rojas, B., see Mínguez-Mosquera, M. I. 585(1991)259
- Ganhao, M. F., Hattingh, J., Hurwitz, M. L. and Pitts, N. I.
Evaluation of a simple plasma catecholamine extraction procedure prior to high-performance liquid chromatography and electrochemical detection 564(1991)55
- Gantier, J.-C., see Merino, A. 572(1991)291
- Gao, W., see Wu, C. 594(1992)243
- García Alvarez-Coque, M. C., see Bonet Domingo, E. 582(1992)189
- García-Baudin, J. M., see Sanchez-Brunete, C. 562(1991)525
- García-Domínguez, J. A., see Fernández-Sánchez, E. 556(1991)485
- Gardana, C., see Pietta, P. 558(1991)296
- Gardner, G. J., see Siu, K. W. M. 554(1991)27
- Gardrat, C., see Latxague, L. 586(1991)166
- Garg, V. and Jusko, W. J.
Simultaneous analysis of prednisone, prednisolone and their major hydroxylated metabolites in urine by high-performance liquid chromatography 567(1991)39
- Gargiulo, M., see Betto, P. 563(1991)115
- Garibotto, G., Ancarani, P., Russo, R., Sala, M. R., Fiorini, F. and Paoletti, E.
Reversed-phase high-performance liquid chromatographic analysis of branched-chain keto acid hydrazone derivatives: optimization of techniques and application to branched-chain keto acid balance studies across the forearm 572(1991)11
- Garland, W. A., see Fayer, B. E. 568(1991)135
- Garrido-Fernández, J., see Mínguez-Mosquera, M. I. 585(1991)259
- Garrigues, P., see Budzinski, H. 590(1992)297
- Garrison, G. L., see Chen, C. L. 583(1992)274
- Garver, F. A., see Kutlar, F. 567(1991)359
- Gaš, B., see Poláček, M. 596(1992)265
- Gaset, A., see Caruel, H. 558(1991)89
- Gaset, A., see Caruel, H. 594(1992)125
- Gaset, A., see Snini, A. 590(1992)369
- Gasiewicz, T. A., see Hahn, M. E. 563(1991)363
- Gaskell, R. M. and Crooks, B.
Practical strategy for the analytical separation of enantiomers by high-performance liquid chromatography 553(1991)357
- Gaspar, G.
High-speed gas chromatography. Theoretical and practical aspects 556(1991)331
- Gaspar, P., see Vandenberg, M. 564(1991)405
- Gaspari, F., Mainardi, L., Ruggenti, P. and Remuzzi, G.
High-performance liquid chromatographic determination of iothalamate acid in human plasma and urine 570(1991)435
- Gaspari, F., Taiocchi, L. and Pochobradsky, M. G.
Determination of brodimoprim and its hydroxy metabolite in human plasma, blood and urine by high-performance liquid chromatography 577(1992)123
- Gasparič, J. and Skutil, J.
Comparative study of chromatography on thin layers impregnated with organic stationary phases. Chromatographic separation of nitrophenols 558(1991)415
- Gasser, H., Strohmaier, W., Schlag, G., Schmid, E. R. and Allmaier, G.
Characterization of phosphatidylcholines in rabbit lung lavage fluid by positive and negative ion fast-atom bombardment mass spectrometry 562(1991)257
- Gassmann, G. and Dahlke, S.
Flame-photometric detection of nitrous oxide in addition to phosphine 598(1992)313

- Gaviraghi, G., see Dalla Libera, L. 566(1991)427
- Gavrichiev, V. S., see Berezkin, V. G. 552(1991)313
- Gavrilloff, C., see Leroy, P. 564(1991)258
- Gavrilova, T. B., Nikitin, Y. S., Vlasenko, E. V., Topalova, I., Petsev, N. and Chaney, C.
Gas chromatographic investigation of silica supports modified in a high-frequency low-temperature plasma 552(1991)179
- Gawdzik, B.
Retention of basic drugs on porous polymers in high-performance liquid chromatography 600(1992)115
- Gay-Montchamp, J. P., see Gaillard, Y. 577(1992)171
- Gaydou, E., see Morin, P. 586(1991)171
- Gazda, K., see Makuch, B. 594(1992)145
- Gazzaniga, P. P., see Pulcinelli, F. M. 575(1992)51
- Ge, H. and Wallace, G. G.
High-performance liquid chromatography on polypyrrole-modified silica 588(1991)25
- Gebauer, P. and Thormann, W.
Isotachopheresis of proteins in uncoated open-tubular fused-silica capillaries with a simple approach for column conditioning 558(1991)423
- Gebauer, P., see Caslavská, J. 585(1991)145
- Gebauer, P., see Thormann, W. 593(1992)275
- Gebelein, M., Merdes, G. and Berger, M. R.
Nucleotide preparation from cells and determination of nucleotides by ion-pair high-performance liquid chromatography 577(1992)146
- Geerts, F., see Van Belle, S. J.-P. 576(1992)351
- Gegout, P., see Payan, E. 566(1991)9
- Geher, K., see Kuksis, A. 564(1991)11
- Geher, K., see Kuksis, A. 579(1992)13
- Gehrke, C. W., see Mitchell, E. P. 581(1992)31
- Geiser, A., see Anton, K. 553(1991)71
- Geisslinger, G., Menzel-Soglowek, S., Kamp, H.-D. and Brune, K.
Stereoselective high-performance liquid chromatographic determination of the enantiomers of ketamine and norketamine in plasma 568(1991)165
- Geisslinger, G., Menzel-Soglowek, S., Schuster, O. and Brune, K.
Stereoselective high-performance liquid chromatographic determination of flurbiprofen in human plasma 573(1992)163
- Geisslinger, G., see Schmidt, N. 583(1992)195
- Gelfi, C., De Besi, P., Alloni, A., Righetti, P. G., Lyubimova, T. and Briskman, V. A.
Kinetics of acrylamide photopolymerization as investigated by capillary zone electrophoresis 598(1992)277
- Gélinas, H., see Massé, R. 562(1991)323
- Gelpi, E., see Abián, J. 554(1991)155
- Gelpi, E., see Abián, J. 562(1991)153
- Gelpi, E., see Ramis, I. 557(1991)507
- Gelpi, E., see Ramis, I. 575(1992)143
- Gelsema, W. J., De Ligny, C. L., Huigen, Y. M. and Schuring, J.
Determination of the mean ionic charge of the components of three ^{99m}Tc bone scanning agents by gel chromatography with two eluents of different electrolyte composition 594(1992)217
- Geng, X. and Chang, X.
High-performance hydrophobic interaction chromatography as a tool for protein refolding 599(1992)185
- Genton, A., see Boutin, J. A. 583(1992)137
- Genty, C., see Fournion, J. M. 556(1991)287
- Geodakyan, S. V., Voskoboinikova, I. V., Kolesnik, J. A., Tjukavkina, N. A., Litvinenko, V. I. and Glyzin, V. I.
High-performance liquid chromatographic method for the determination of mangiferin, likvirin and dihydroquercetin in rat plasma and urine 577(1992)371
- Geodakyan, S. V., see Tjukavkina, N. A. 571(1991)312
- George, R. C., see Chan, K. Y. 571(1991)291
- Gerber, G., see Grune, T. 553(1991)193
- Gerber, G., see Grune, T. 563(1991)53
- Gerber, G., see Werner, A. 553(1991)205
- Gerber, G., see Ziegler, M. 563(1991)172
- Gerbino, T. C., see Castello, G. 585(1991)273
- Gerig, T. M., see Jones, L. A. 595(1992)221
- Gerota, J., see Cardot, P. J. P. 568(1991)93
- Gerstner, J. A., Hamilton, R. and Cramer, S. M.
Membrane chromatographic systems for high-throughput protein separations 596(1992)173
- Gerus, I. I., see Galushko, S. V. 600(1992)83
- Geschier, C., see Regnault, V. 584(1992)93
- Geschwindt, L., see Boppana, V. K. 592(1992)317
- Getek, T., see Talaska, G. 580(1992)293
- Getek, T. A., Vestal, M. L. and Alexander, T. G.
Analysis of gentamicin sulfate by high-performance liquid chromatography combined with thermospray mass spectrometry 554(1991)191
- Ghabrial, H., see Martin, G. 574(1992)205
- Ghanayem, B. I., see Demby, K. B. 579(1992)153
- Ghatineh, S., see Preece, N. E. 573(1992)227
- Ghielmi, S., see Dorizzi, R. 580(1992)279
- Ghielmi, S., see Tagliaro, F. 566(1991)333
- Ghielmi, S., see Tagliaro, F. 580(1992)161
- Ghijis, M., see Coene, J. 553(1991)285
- Ghodbane, S., Oweimreen, G. A. and Martire, D. E.
Thermodynamics of solution of non-mesomorphic solutes at infinite dilution in the smectic-A, nematic and isotropic phases of *p-n*-octyl-*p'*-cyanobiphenyl. A gas-liquid chromatographic study 556(1991)317
- Ghowsi, K. and Gale, R. J.
Field effect electroosmosis 559(1991)95
- Giachetti, C., Assandri, A. and Zanolo, G.
Gas chromatographic-mass spectrometric determination of ethyl carbamate as the xanthylamide derivative in Italian aqua vitae (grappa) samples 585(1991)111
- Giacomini, M., see Chiari, M. 558(1991)285
- Gianazza, E. and Righetti, P. G.
Biomedical relevance of two-dimensional protein mapping (Review) 569(1991)43
- Gianazza, E., see Bianchi-Bosisio, A. 569(1991)243
- Gianazza, E., see Righetti, P. G. 569(1991)197
- Gianbenedetti, M., see Betto, P. 563(1991)115
- Giannini, R., see Zezza, F. 593(1992)99
- Giard-Pasquier, C. and Rickenbacher, U.
Purification of canine prolactin and growth hormone by fast protein liquid chromatography 553(1991)117

- Gibbs, S. J., see Athalye, A. M. 589(1992)71
- Gibeaut, D. M. and Carpita, N. C.
Clean-up procedure for partially methylated alditol acetate derivatives of polysaccharides 587(1991)284
- Gibson, G. G., see Clare, R. A. 562(1991)237
- Giese, R. W., see Al-Deen, A. N. 600(1992)229
- Giese, R. W., see Li, W. 588(1991)273
- Gieseler, F., see Boege, F. 587(1991)3
- Gilbert, M., see Abolfathi, Z. 579(1992)366
- Gill, R., see Smith, R. M. 592(1992)85
- Gillissen, E. E. A., see De Bokx, P. K. 598(1992)115
- Gilpin, R. K. and Wu, L.
Use of the reordering/resolution of alkyl-modified silica to characterize the microscopic heterogeneity of silica via liquid chromatography 556(1991)415
- Gimenez, F., see Fernandez, C. 572(1991)195
- Giralt, E., see Celma, C. 562(1991)447
- Girault, J., Istin, B., Malgouyat, J. M., Brisson, A. M. and Fourtillan, J. B.
Simultaneous determination of beclomethasone, beclomethasone monopropionate and beclomethasone dipropionate in biological fluids using a particle beam interface for combining liquid chromatography with negative-ion chemical ionization mass spectrometry 564(1991)43
- Girin, E., see Cann-Moisán, C. 574(1992)134
- Girón, J., see Alaiz, M. 591(1992)181
- Girot, P., see Santambien, P. 597(1992)315
- Giuliano, G., see Santoni, G. 581(1992)287
- Glaser, B. M., see Alliegro, M. A. 582(1992)217
- Glässgen, W. E., Hofmann, R., Emmerling, M., Neumann, G. D. and Seitz, H. U.
Structure elucidation of saccharides in anthocyanins and flavonols by means of methylation analysis and gas chromatography 598(1992)81
- Glatz, J. F. C., see Van Nieuwenhoven, F. A. 570(1991)173
- Glavač, D.
 R_M values of some colchicines and colchicineamides determined by reversed-phase thin-layer chromatography 591(1992)367
- Gleispach, H., see Leis, H. J. 578(1992)116
- Glennon, J. J., see Jones, L. A. 595(1992)209
- Glennon, J. J., see Jones, L. A. 595(1992)221
- Glód, B. K., Haddad, P. R. and Alexander, P. W.
Potentiometric detection of carboxylic acids and inorganic anions in ion-exclusion chromatography using camphorsulphonic acid as eluent 589(1992)209
- Glód, B. K., Haddad, P. R. and Alexander, P. W.
Ion-exclusion chromatography using mobile phases containing β -cyclodextrin 595(1992)149
- Glód, B. K., see Alexander, P. W. 589(1992)201
- Glover, V., see Halket, J. M. 562(1991)279
- Główniak, K.
Comparison of selectivity of silica and Florisil in the separation of natural pyranocoumarins 552(1991)453
- Glyzin, V. I., see Geodakyan, S. V. 577(1992)371
- Glyzin, V. I., see Tjukavkina, N. A. 571(1991)312
- Goad, L. J., see Evershed, R. P. 590(1992)305
- Godbillon, J., see Rouan, M. C. 573(1992)59
- Godbillon, J., see Rouan, M. C. 573(1992)65
- Godbillon, J., see Rouan, M. C. 577(1992)387
- Godbillon, J., see Sioufi, A. 565(1991)401
- Godbillon, J., see Sioufi, A. 571(1991)87
- Goewie, C. E., see Hogendoorn, E. A. 552(1991)113
- Golc-Wondra, A., see Prošek, M. 553(1991)477
- Goldberg, J. I., see Stoley, B. D. 567(1991)49
- Goldberg, M. J., see Anliker, S. L. 573(1992)141
- Goldstein, D. S., see Armando, J. 568(1991)45
- Golovchenko, N. P., Kataeva, I. A. and Akimenko, V. K.
Analysis of pH-dependent protein interactions with gel filtration medium 591(1992)121
- Golovnya, R. V., Zhuravleva, I. L., Yakush, E. V. and Schenderjuk, V. V.
Retentions of alkyloxazoles and alkylthiazoles and their prediction based on non-linear additivity concepts in gas chromatography 552(1991)1
- Gombert, J., see Dufay, C. 563(1991)224
- Gómez, G., see Ramis, I. 557(1991)507
- Gomez, L. E. and Lehmann F., P. A.
Plasma determination of the novel anticonvulsant D,L-3-hydroxy-3-ethyl-3-phenylpropionamide and preliminary pharmacokinetic studies in the rat 575(1992)306
- Gomez, R., see De Las Rivas, J. 585(1991)168
- Gomez-Belinchon, J. I., see Rosell, A. 562(1991)493
- Gong, S., see Wu, C. 594(1992)243
- González, T., Awadé, A., Besson, C. and Robert-Baudouy, J.
Purification and characterization of recombinant pyrrolidone carboxyl peptidase of *Bacillus subtilis* 584(1992)101
- González, C., see Rodríguez, P. 562(1991)547
- Gonzalez, F. J., see Cholerton, S. 575(1992)325
- González, M. J., see Tabera, J. 557(1991)481
- González, M. J., see Vidal-Madjar, C. 584(1992)11
- Goodwin, B. L., see Halket, J. M. 562(1991)279
- Gooijer, C.
Luminescence techniques in chemical and biochemical analysis (edited by W. R. G. Bayens, P. De Keukeleire and K. Korkides) (Book Review) 588(1991)367
- Gooijer, C., see Schreurs, M. 552(1991)625
- Gooijer, C., see Somsen, G. W. 552(1991)635
- Gooijer, C., see Van de Nesse, R. J. 552(1991)613
- Goosens, E. C., De Jong, D., Van den Berg, J. H. M., De Jong, G. J. and Brinkman, U. A. T.
Reversed-phase liquid chromatography coupled on-line with capillary gas chromatography. I. Introduction of large volumes of aqueous mixtures through an on-column interface 552(1991)489
- Gorchakov, V. D., Sakodinskii, K. I., Lebedin, Y. S. and Chuchalin, A. G.
Immunoabsorbents for clinical use: *ex vivo* immunoglobulin E removal in allergy 563(1991)166
- Goren, M. P.
Determination of urinary 2- and 3-dechloroethylated metabolites of ifosfamide by high-performance liquid chromatography 570(1991)351
- Goretti, G., see Cartoni, G. P. 552(1991)197
- Gori, G., see Sturaro, A. 590(1992)223
- Görög, S., see Herényi, B. 592(1992)297

- Gorrichon, L., see Baltas, M. 600(1992)323
- Gorrod, J. W., see Kajbaf, M. 575(1992)75
- Gorse, J., see Cole, R. O. 557(1991)113
- Goss, T. A., Bard, M. and Jarrett, H. W.
High-performance affinity chromatography of messenger RNA 588(1991)157
- Gosselet, M. and Sebillé, B.
Effects of β -cyclodextrin in the mobile phase on the retention and indirect detection of non-electrolytes in reversed-phase liquid chromatography. I. Study of aliphatic alcohols 552(1991)563
- Gosselet, M., see Agnus, B. 552(1991)583
- Goto, A., see Akimoto, K. 588(1991)165
- Goto, H., see Ohmori, S. 566(1991)1
- Goto, J., Hasegawa, K., Nambara, T. and Iida, T.
Studies on steroids. CCLIV. Gas chromatographic-mass spectrometric determination of 4- and 6-hydroxylated bile acids in human urine with negative ion chemical ionization detection 574(1992)1
- Goto, J., Saisho, Y. and Nambara, T.
Studies on steroids. CCLII. Separation and characterization of 3-oxobile acids in serum by high-performance liquid chromatography with fluorescence detection 567(1991)343
- Goto, J., Teraya, Y., Nambara, T. and Iida, T.
Studies on steroids. CCLIII. Capillary gas chromatographic behaviour of diethylhydrogensilyl-diethylsilylene derivatives of stereoisomeric bile acids 585(1991)281
- Goto, J., see Iida, T. 558(1991)451
- Goto, N., Kamata, T. and Ikegami, K.
Trace analysis of quinapril and its active metabolite, quinaprilat, in human plasma and urine by gas chromatography-negative-ion chemical ionization mass spectrometry 578(1992)195
- Goto, N., Sato, T., Shigetoshi, M. and Ikegami, K.
Determination of dioxopiperazine metabolites of quinapril in biological fluids by gas chromatography-mass spectrometry 578(1992)203
- Goto, T., see Tsuyama, Y. 596(1992)181
- Gotsick, J. T. and Schmidt, Jr., D. E.
Utility of small-particle silica in preparative chromatography 590(1992)77
- Gottschall, K., see Abou-Rebyeh, H. 566(1991)351
- Gottschall, K., see Oberdorfer, F. 552(1991)483
- Goubran-Botros, H., Nanak, E., Abdul Nour, J., Birkenmeir, G. and Vijayalakshmi, M. A.
Immobilized metal ion affinity electrophoresis. A study with several model proteins containing histidine 597(1992)357
- Gouesclou, L., see Pruñonosa, J. 574(1992)127
- Gouesclou, L., see Pruñonosa, J. 581(1992)219
- Gould, I. M., see Taylor, R. B. 563(1991)451
- Gourdiér, B., see Debal, V. 581(1992)93
- Gouyette, A., see Barilero, I. 575(1992)275
- Gower, D. B., see Mallet, A. I. 562(1991)647
- Graber, M. and Condoret, J. S.
Preparative anion-exchange chromatography of soybean trypsin inhibitor: the alternative of column-overload methods 584(1992)115
- Graells, M., see Esteban, A. 573(1992)121
- Graham, W. D. and Annette, D.
Determination of ascorbic and dehydroascorbic acid in potatoes (*Solanum tuberosum*) and strawberries using ion-exclusion chromatography 594(1992)187
- Grajek, H., Neffe, S. and Witkiewicz, Z.
Chromatographic determination of the physico-chemical parameters of adsorption on activated carbon fibres 600(1992)67
- Grajek, H., see Witkiewicz, Z. 556(1991)441
- Gramiccioni, L., see Milana, M. R. 552(1991)205
- Gramm, H.-J., see Zimmermann, J. 567(1991)240
- Grampp, T., see Feng, N. 575(1992)289
- Granata, O. M., see Castagnetta, L. A. 572(1991)25
- Granley, K., see Singh, A. K. 568(1991)351
- Granneman, G. R. and Varga, L. L.
High-performance liquid chromatographic procedures for the determination of temafloxacin in biological matrices 568(1991)197
- Grases, J. M., see Rodríguez, P. 562(1991)547
- Grate, J. W., see Abraham, M. H. 588(1991)361
- Grateful, T. M. and Lightfoot, Jr., E. N.
Finite difference modelling of continuous-flow electrophoresis 594(1992)341
- Gray, D. O., see Price, N. P. J. 598(1992)51
- Greaves, J., see Hale, R. C. 580(1992)257
- Green, B. N., see Shackleton, C. H. L. 562(1991)175
- Green, M. D., see Millington, D. S. 562(1991)47
- Greenberg, G. E., see Chovan, J. P. 578(1992)77
- Greenberg, J., see Phillips, M. 564(1991)242
- Greenhagen, R. D., see O'Shea, T. J. 593(1992)305
- Greenshaw, A. J., see Paetsch, P. R. 573(1992)313
- Greig, L. G., see Haynes, P. A. 588(1991)107
- Greig, N. H., see Deutsch, J. 579(1992)93
- Greischel, A., Beschke, K., Rapp, H. and Roth, W.
Quantitation of the new hypoglycaemic agent AG-EE 388 ZW in human plasma by automated high-performance liquid chromatography with electrochemical detection 568(1991)246
- Griest, W. H., Ramsey, R. S., Ho, C.-H. and Caldwell, W. M.
Supercritical fluid extraction of chemical warfare agent simulants from soil 600(1992)273
- Griggs, L. J., see Poon, G. K. 572(1991)143
- Grill, C. M., see Painuly, P. 590(1992)139
- Grimaldi, F., see Rop, P. P. 573(1992)87
- Grimalt, J. O., see Rosell, A. 562(1991)493
- Grimmelt, B., see Nijjar, M. S. 568(1991)393
- Grit, M., Crommelin, D. J. A. and Lang, J.
Determination of phosphatidylcholine, phosphatidylglycerol and their lyso forms from liposome dispersions by high-performance liquid chromatography using high-sensitivity refractive index detection 585(1991)239
- Groen, C. P., see Van Zijtveld, J. 600(1992)211
- Groppi, A., see Poletini, A. 564(1991)529
- Gross, G. A. and Grüter, A.
Quantitation of mutagenic/carcinogenic heterocyclic aromatic amines in food products 592(1992)271

- Grossi, G., Bargossi, A. M., Fiorella, P. L., Piazzini, S., Battino, M. and Bianchi, G. P.
Improved high-performance liquid chromatographic method for the determination of coenzyme Q₁₀ in plasma 593(1992)217
- Grossman, E., see Armando, I. 568(1991)45
- Grossman, P. D. and Soane, D. S.
Capillary electrophoresis of DNA in entangled polymer solutions 559(1991)257
- Grote, H., see Arnold, P. J. 544(1991)267
- Grover, P. K., see Paliwal, J. K. 572(1991)219
- Grover, R., see Cessna, A. J. 600(1992)327
- Gruhn, V., see Cunico, R. L. 559(1991)467
- Grummt, I., see Heilgental, G. 587(1991)25
- Grunau, J. A. and Swiader, J. M.
Chromatography of 99 amino acids and other ninhydrin-reactive compounds in the Pickering lithium gradient system 594(1992)165
- Grune, T., Siems, W., Gerber, G., Tikhonov, Y. V., Pimenov, A. M. and Toguzov, R. T.
Changes of nucleotide patterns in liver, muscle and blood during the growth of Ehrlich ascites cells: application of the reversed-phase and ion-pair reversed-phase high-performance liquid chromatography with radial compression column 563(1991)53
- Grune, T., Siems, W. G., Gerber, G. and Uhlig, R.
Determination of the ultraviolet absorbance and radioactivity of purine compounds separated by high-performance liquid chromatography. Application to metabolic flux rate analysis 553(1991)193
- Grüner, B., Plzák, Z. and Vinš, I.
Chromatographic behaviour of the *cis*-[B₁₂H₁₂]²⁻ derivatives on hydroxyethylmethacrylate gels 588(1991)201
- Grüner, B., Plzák, Z. and Vinš, I.
Purity assay of sodium mercaptododecaborate by high-performance liquid chromatography 595(1992)169
- Grunow, M. and Schöpp, W.
Purification of pro- and eukaryotic superoxide dismutases by charge-controlled hydrophobic chromatography 590(1992)247
- Grupeli, M., see Tejedor, M. C. 589(1992)127
- Grushka, E.
Effect of hydrostatic flow on the efficiency in capillary electrophoresis 559(1991)81
- Grushka, E., see Zamir, I. 565(1991)424
- Grushka, E., see Zamir, I. 567(1991)319
- Grüter, A., see Gross, G. A. 592(1992)271
- Grygiel, J., see Havard, J. 584(1992)270
- Guada, J. A., see Balcells, J. 575(1992)153
- Guan, Y. and Zhou, L.
Live retention database for identification in multi-step temperature-programmed capillary gas chromatography 552(1991)187
- Guardiola, A., Ventura, F., Matia, L., Caixach, J. and Rivera, J.
Gas chromatographic-mass spectrometric characterization of volatile organic compounds in Barcelona tap water 562(1991)481
- Guattari, B.
High-performance liquid chromatographic determination, with ultraviolet detection, of S-adenosyl-L-methionine and S-adenosyl-L-homocysteine in rat tissues and simultaneously of normetanephrine and metanephrine for phenylethanolamine-N-methyltransferase or catechol-O-methyltransferase activities 567(1991)254
- Guentert, T. W., see Stebler, T. 564(1991)330
- Guérand, W. S., see De Jong, J. 570(1991)209
- Guérand, W. S., see De Jong, J. 574(1992)273
- Guernet, M., see Merino-Dugay, A. 579(1992)73
- Guevremont, R., see Siu, K. W. M. 554(1991)27
- Guidotti, L., see Biondi, P. A. 566(1991)377
- Guieu, D., see Jaussaud, P. 573(1992)136
- Guillaume, J.-L., see Dandeu, J.-P. 599(1992)105
- Guillén, M. D., Iglesias, M. J., Dominguez, A. and Blanco, C. G.
Polynuclear aromatic hydrocarbon retention indices on SE-54 stationary phase of the volatile components of a coal tar pitch. Relationships between chromatographic retention and thermal reactivity 591(1992)287
- Guilluy, R., Billion-Rey, F. and Brazier, J. L.
On-line measurements of ¹³C enrichments in rat breath. Non-invasive method for *in vivo* study of drug enzymatic induction 562(1991)341
- Guinebault, P., see Ascalone, V. 581(1992)237
- Guinec, N., see Dalet-Fumeron, V. 568(1991)55
- Guiochon, G.
Foreword 590(1992)1
- Guiochon, G., see Felinger, A. 591(1992)31
- Guiochon, G., see Jacobson, S. C. 590(1992)119
- Guiochon, G., see Jacobson, S. C. 600(1992)37
- Guiochon, G., see Jandera, P. 588(1991)1
- Guiochon, G., see Katti, A. M. 556(1991)205
- Guiochon, G., see Roles, J. 589(1992)223
- Guiochon, G., see Roles, J. 591(1992)233
- Guiochon, G., see Roles, J. 591(1992)245
- Guiochon, G., see Roles, J. 591(1992)267
- Guiochon, G., see Yang, Y.-B. 590(1992)35
- Guiochon, G., see Zhu, J. 552(1991)71
- Guiraud, P., see Richard, M. J. 577(1992)9
- Guiso, G., Confalonieri, S. and Caccia, S.
High-performance liquid chromatographic method for the determination of a potential memory-enhancing compound (CL 275,838) and its desbenzyl metabolite in rat plasma or serum 572(1991)346
- Guiso, G., see Caccia, S. 581(1992)109
- Guiso, G., see Sarati, S. 563(1991)323
- Guix, T., see Parra, P. 570(1991)185
- Gulaid, A. A., Jahn, G. A., Maslen, C. and Dennis, M. J.
Simultaneous determination of trimipramine and its major metabolites by high-performance liquid chromatography 566(1991)228
- Gulati, A.
High-performance liquid chromatographic determination of morphine, morphine-3-glucuronide, morphine-6-glucuronide and codeine in biological samples using multi-wavelength forward optical detection. Reply to Zweipfenning 578(1992)335

- Gulati, A., see Chari, G. 571(1991)263
- Gumbhir, K. and Mason, W. D.
High-performance liquid chromatographic method for the determination of tris(hydroxymethyl)aminomethane (tromethamine) in human plasma 583(1992)99
- Gumbleton, M., see Han, C. 579(1992)237
- Gumprecht, D. L.
Comparison of commercially available thin-layer chromatography plates with mixtures of dyes, analgesics and phenols 595(1992)368
- Gunata, Z. Y., see Voirin, S. G. 590(1992)313
- Guo, R., Lim, C. K. and Peters, T. J.
High-performance liquid chromatographic assays for protoporphyrinogen oxidase and ferrochelatase in human leucocytes 566(1991)383
- Gupta, R. C., see Paliwal, J. K. 572(1991)219
- Gupta, R. N.
Analysis for drugs and metabolites including anti-infective agents (edited by E. Reid and I. D. Wilson) (Book Review) 565(1991)534
- Gupta, R. N.
Liquid chromatography in biomedical analysis (edited by T. Hanai) (Book Review) 571(1991)334
- Gupta, R. N.
Drug level monitoring: antidepressants (Review) 576(1992)183
- Gupta, R. N. and Whelton, C.
Determination of plasma homovanillic acid by liquid chromatography with electrochemical detection 582(1992)236
- Gurgey, A., see Kutlar, F. 567(1991)359
- Gurley, L. R., Buchanan, J. S., London, J. E., Stavert, D. M. and Lehnert, B. E.
High-performance capillary electrophoresis of proteins from the fluid lining of the lungs of rats exposed to perfluoroisobutylene 559(1991)411
- Gurley, L. R., London, J. E. and Valdez, J. G.
High-performance capillary electrophoresis of histones 559(1991)431
- Guse, A. H. and Emmrich, F.
Determination of inositol polyphosphates from human T-lymphocyte cell lines by anion-exchange high-performance liquid chromatography and post-column derivatization 593(1992)157
- Guserle, R., see Arnold, P. J. 544(1991)267
- Gustafson, A., see Wang, W.-Q. 581(1992)139
- Gustavson, L. E. and Chu, S.
High-performance liquid chromatographic procedure for the determination of tiagabine concentrations in human plasma using electrochemical detection 574(1992)313
- Gutiérrez, M. C., see Linares, P. 558(1991)147
- Gutnikov, G. and Streng, J. R.
Rapid high-performance liquid chromatographic determination of fatty acid profiles of lipids by conversion to their hydroxamic acids 587(1991)292
- Guttman, A. and Cooke, N.
Effect of temperature on the separation of DNA restriction fragments in capillary gel electrophoresis 559(1991)285
- Guttman, A., Nelson, R. J. and Cooke, N.
Prediction of migration behavior of oligonucleotides in capillary gel electrophoresis 593(1991)297
- Guyer, G., see Shaikh, B. 571(1991)189
- Guzman, N. A., Ali, H., Moschera, J., Iqbal, K. and Malick, A. W.
Assessment of capillary electrophoresis in pharmaceutical applications. Analysis and quantification of a recombinant cytokine in an injectable dosage form 559(1991)307
- Guzman, N. A., Moschera, J., Bailey, C. A., Iqbal, K. and Malick, A. W.
Assay of protein drug substances present in solution mixtures by fluorescamine derivatization and capillary electrophoresis 598(1992)123

H

- Haagsma, N., see Moretti, V. M. 583(1992)77
- Haagsma, N., see Van de Water, C. 566(1991)173
- Haanappel, V. A. C., Stevens, T. W., Hovestad, A., Skolnik, V. and Visser, R.
Gas chromatographic-mass spectrometric analysis of tar compounds formed during pyrolysis of rice husks 562(1991)531
- Haasnoot, W., see Ploum, M. E. 564(1991)413
- Häberlein, H., see Sturm, G. 562(1991)351
- Häberlein, I.
Separation of the complete thioredoxin pattern of soybean leaves (*Glycine max*) by high-performance anion-exchange chromatography on Mono Q 587(1991)109
- Hachey, D. L., see Roy, D. 567(1991)309
- Haddad, P. R., see Alexander, P. W. 589(1992)201
- Haddad, P. R., see Buchberger, W. 558(1991)181
- Haddad, P. R., see Glód, B. K. 589(1992)209
- Haddad, P. R., see Glód, B. K. 595(1992)149
- Haddad, P. R., see Mills, M. H. 564(1991)93
- Hadjimichael, J., see Rossomando, E. F. 583(1992)11
- Hadjimichael, J., see Rossomando, E. F. 583(1992)19
- Hadjimichael, J., see Spiegel, L. B. 573(1992)23
- Haegele, K. D., see Berg-Candolfi, M. 593(1992)1
- Haegele, K. D., see Dulery, B. D. 571(1991)241
- Haerdi, W., see Chambaz, D. 600(1992)203
- Haerdi, W., see Coquet, A. 553(1991)255
- Haerdi, W., see Maeder, G. 593(1992)9
- Hage, D. S., see Loun, B. 579(1992)225
- Hage, D. S., see Noctor, T. A. G. 577(1992)305
- Hagedorn, H.-W., Schulz, R. and Friedrich, A.
Detection of methandienone (methandrostenolone) and metabolites in horse urine by gas chromatography-mass spectrometry 577(1992)195

- Hagel, L.
Peak capacity of columns for size-exclusion chromatography 591(1992)47
- Haggarty, N. W., see Burton, S. C. 587(1991)271
- Haginaka, J. and Seyama, C.
Determination of enantiomers of 1-benzyl-4-[(5,6-dimethoxy-1-indanon)-2-yl]methylpiperidine hydrochloride (E2020), a centrally acting acetylcholine esterase inhibitor, in plasma by liquid chromatography with fluorometric detection 577(1992)95
- Haginaka, J., Seyama, C., Yasuda, H., Fujima, H. and Wada, H.
Retention and enantioselectivity of racemic solutes on a modified ovomucoid-bonded column. I. Cross-linking with glutaraldehyde 592(1992)301
- Haginaka, J., Seyama, C., Yasuda, H. and Takahashi, K.
Investigation of enantioselectivity and enantiomeric elution order of propranolol and its ester derivatives on an ovomucoid-bonded column 598(1992)67
- Haginaka, J. and Wakai, J.
Preparation and characterization of mixed functional phase silica materials using phenyl-, butyl- or octylchlorosilane as a silylating agent 596(1992)151
- Haginaka, J., see Kimata, K. 558(1991)19
- Haginaka, J., see Miwa, T. 566(1991)163
- Hagiwara, H., Nagasawa, T., Lodhi, K. M., Kozuka, M., Ito, T. and Hirose, S.
Affinity chromatographic purification of bovine lung endothelin receptor using biotinylated endothelin and avidin-agarose 597(1992)331
- Hague, G. R., see Mulholland, F. 589(1992)380
- Hahn, A., see Sander, S. 558(1991)115
- Hahn, D. A., see Dunn, M. J. 595(1992)185
- Hahn, G., see Spahn-Langguth, H. 584(1992)229
- Hahn, M. E. and Gasiewicz, T. A.
Determination of individual porphyrins in rodent urine using high-performance liquid chromatography following clean-up by anion-exchange chromatography 563(1991)363
- Hahn, M.-K., see Tan, H. S. I. 568(1991)145
- Haidacher, D., Bonn, G. K., Scherz, H., Nitsch, E. and Wutka, R.
Thin-layer electrophoresis of hydroxyethyl starches on a modified silica gel support 591(1992)351
- Håkanson, H., see Nilsson, M. 597(1992)383
- Haken, J. K.
Stationary phases in gas chromatography (by H. Rotzsche) (Book Review) 600(1992)378
- Haken, J.K. and Camamo, M.
Degradation of polysulphones by alkali fusion 595(1992)283
- Haken, J. K. and Iddamalgoda, P.
Reactor for prechromatographic fusion reactions 600(1992)352
- Hakusui, H., see Fujimaki, Y. 575(1992)261
- Hakusui, H., see Nomura, H. 572(1991)333
- Hakusui, H., see Okazaki, O. 563(1991)313
- Haldna, Ü.
Carbonate eluents in ion chromatography. Fractions of resin capacity linked to anions present in the eluent 589(1992)197
- Hale, R. C. and Greaves, J.
Methods for the analysis of persistent chlorinated hydrocarbons in tissues (Review) 580(1992)257
- Halket, J. M., Watkins, P. J., Przyborowska, A., Goodwin, B. L., Clow, A., Glover, V. and Sandler, M.
Isatin (indole-2,3-dione) in urine and tissues. Detection and determination by gas chromatography-mass spectrometry 562(1991)279
- Hall, C. D., see Marvin, C. H. 555(1991)147
- Hallström, S., see Fürst, W. 578(1992)39
- Halmekoski, J., see Salo, M. 592(1992)127
- Halvorsen, K., Ader, J. C., Rico, I. and Lattes, A.
Analyse par chromatographie en phase gazeuse de la réaction de macrolactonisation de Mukaiyama 557(1991)501
- Hamaguchi, K., see Ide, H. 596(1992)203
- Hamann, H.-J., see Kunath, A. 588(1991)352
- Hamdi, A., see Clark, B. J. 553(1991)383
- Hamelijnck, M. A. F., see Stevenson, P. J. 563(1991)419
- Hamers, A. R. M., see Ploum, M. E. 564(1991)413
- Hamilton, M., see Anliker, S. L. 573(1992)141
- Hamilton, R., see Gerstner, J. A. 596(1992)173
- Hammers, W. E., see Lebeau, B. 596(1992)285
- Hammes, W. and Weyhenmeyer, R.
New assay method for the determination of vinpocetine in human plasma by gas chromatography-mass spectrometry without transesterification caused by solvents: a reply. 567(1991)504
- Hammond, P. M., see Scott, D. A. 573(1992)309
- Hamoir, T., De Smet, M., Piryns, H., Conti, P., Vanden Driessche, N., Massart, D. L., Maris, F., Hindriks, H. and Schoenmakers, P. J.
Feasibility study for the construction of an integrated expert system in high-performance liquid chromatography 589(1992)31
- Hamon, M., see Bouygues-De Ferran, A. M. 585(1991)289
- Han, C., Gumbleton, M., Lau, D. T. W. and Benet, L. Z.
Improved gas chromatographic assay for the simultaneous determination of nitroglycerin and its mono- and dinitrate metabolites 579(1992)237
- Han, S. S., see Lee, H. S. 574(1992)175
- Hanai, K., see Kawai, S. 585(1991)318
- Hanawa, T., see Fujioka, T. 597(1992)429
- Hancock, W. S., see Chloupek, R. C. 594(1992)65
- Hancock, W. S., see Frenz, J. 557(1991)289
- Hang Huynh, N., see Heldin, E. 592(1992)339
- Hanis, T., Deyl, Z., Struzinsky, R. and Miksik, I.
Separation of elastin cross-links as phenylisothiocyanate derivatives 553(1991)93
- Hann, J. T., see Perez-Souto, N. 593(1992)209
- Hannant, M. E., see Unni, L. K. 573(1992)99
- Hanneman, W. H., see Champney, T. H. 579(1992)334
- Hansen, Å. M., Poulsen, O. M., Christensen, J. M. and Hansen, S. H.
Determination of 2-naphthylamine in urine by a novel reversed-phase high-performance liquid chromatography method 578(1992)85
- Hansen, G. R., see Köppel, C. 575(1992)87
- Hansen, K., see Banke, N. 559(1991)325

- Hansen, S. H. and Tjørnelund, J.
High-performance liquid chromatography on dynamically modified silica. IX. Modification of silica with 3-(N,N-dimethylpalmitylammonium) propanesulphonate for reversed-phase chromatography 556(1991)353
- Hansen, S. H., see Andersen, J. V. 577(1992)325
- Hansen, S. H., see Andersen, J. V. 577(1992)362
- Hansen, S. H., see Hansen, Å. M. 578(1992)85
- Hansen, S. H., see Jørgensen, M. 577(1992)167
- Hansen, S. H., see Tjørnelund, J. 570(1991)109
- Hansen, S. H., see Tjørnelund, J. 570(1991)224
- Hänsler, M., Appelt, G. and Rogos, R.
Improved detection of human pancreatic proteins separated by two-dimensional isoelectric focusing/sodium dodecyl sulphate gel electrophoresis using a combined staining procedure 563(1991)63
- Hanson, M., Unger, K. K., Mant, C. T. and Hodges, R. S.
Polymer-coated reversed-phase packings with controlled hydrophobic properties. I. Effect on the selectivity of protein separations 599(1992)65
- Hanson, M., Unger, K. K., Mant, C. T. and Hodges, R. S.
Polymer-coated reversed-phase packings with controlled hydrophobic properties. II. Effect on the selectivity of peptide separations 599(1992)77
- Hansson, L., see Marle, I. 586(1991)233
- Harada, K., see Etoh, T. 578(1992)292
- Harada, M., see Hiraoka, B. Y. 563(1991)142
- Haraguchi, H., see Hu, W. 557(1991)441
- Harajiri, S., Wood, G. and Desiderio, D. M.
Analysis of proenkephalin A, proopiomelanocortin and protachykinin neuropeptides in human lumbar cerebrospinal fluid by reversed-phase high-performance liquid chromatography, radioimmunoassay and enzymolysis 575(1992)213
- Harding, D. R. K., see Burton, S. C. 587(1991)271
- Harke, H. R., see Chen, D. Y. 559(1991)237
- Harkrader, R., see Reinhart, P. 570(1991)425
- Härmälä, P., Vuorela, H. and Rahko, E.-L.
Retention behaviour of closely related coumarins in thin-layer chromatographic preassays for high-performance liquid chromatography according to the "PRISMA" model 593(1992)329
- Harmatha, J., see Piš, J. 596(1992)271
- Harrelson, J. C., see Brown, A. R. 578(1992)302
- Harrington, S. J., Varro, R. and Li, T. M.
High-performance capillary electrophoresis as a fast in-process control method for enzyme-labelled monoclonal antibody conjugates 559(1991)385
- Harris, R. C., see Dunnett, M. 579(1992)45
- Harris, R. M., see Peterson, J. C. 564(1991)205
- Harrison, D. J., see Manz, A. 593(1992)253
- Harrison, K., see Yang, Y.-B. 590(1992)35
- Harrison, M. E., see Abdulrahman, S. 562(1991)713
- Hartter, E., see Ferris, R. 577(1992)251
- Härtter, S. and Hiemke, C.
Column switching and high-performance liquid chromatography in the analysis of amitriptyline, nortriptyline and hydroxylated metabolites in human plasma or serum 578(1992)273
- Hartwick, R. A., see Wang, T. 589(1992)307
- Hartwick, R. A., see Wang, T. 594(1992)325
- Hartwick, R. A., see Wu, N. 582(1992)77
- Harvey, D. J.
Identification by gas chromatography and mass spectrometry of lipids from the rat Harderian gland 565(1991)27
- Harvey, D. J., Samara, E. and Mechoulam, R.
Urinary metabolites of cannabidiol in dog, rat and man and their identification by gas chromatography-mass spectrometry 562(1991)299
- Harvey, S. D., Campbell, J. A., Kelsey, R. G. and Vance, N. C.
Separation of taxol from related taxanes in *Taxus brevifolia* extracts by isocratic elution reversed-phase microcolumn high-performance liquid chromatography 587(1991)300
- Hase, M., see Nawata, M. 597(1992)415
- Hasegawa, K., see Goto, J. 574(1992)1
- Hashem, A. and Lubczyk, B.
Determination of isoxsuprine in equine plasma by high-performance liquid chromatography with electrochemical detection 563(1991)216
- Hashimoto, A., Nishikawa, T., Oka, T., Takahashi, K. and Hayashi, T.
Determination of free amino acid enantiomers in rat brain and serum by high-performance liquid chromatography after derivatization with N-tert-butylloxycarbonyl-L-cysteine and o-phthaldialdehyde 582(1992)41
- Hassan, E. E. and Gallo, J. M.
High-performance liquid chromatographic analysis of the anticancer drug oxantrazole in rat whole blood and tissues 582(1992)225
- Hasui, M., see Takamatsu, T. 584(1992)261
- Hatada, K., Kimura, M., Ono, I. and Ozaki, M.
Determination of a new calcium antagonist and its main metabolite in plasma by thermospray liquid chromatography-mass spectrometry 583(1992)116
- Hatano, H., see Imoto, T. 557(1991)383
- Hathout, Y., see Serrano-Carreón, L. 584(1992)129
- Hattingh, J., see Ganhaio, M. F. 564(1991)55
- Hattori, H., Yamamoto, S., Iwata, M., Takashima, E., Yamada, T. and Suzuki, O.
Sensitive determination of phenothiazines in body fluids by gas chromatography with surface ionization detection 579(1992)247
- Hattori, H., Yamamoto, S., Iwata, M., Takashima, E., Yamada, T. and Suzuki, O.
Determination of diphenylmethane antihistaminic drugs and their analogues in body fluids by gas chromatography with surface ionization detection 581(1992)213
- Hattori, H., Yamamoto, S., Yamada, T. and Suzuki, O.
Determination of local anaesthetics in body fluids by gas chromatography with surface ionization detection 564(1991)278
- Haupt, K., see Wu, X. 584(1992)35
- Hausen, A., see Werner, E. R. 570(1991)43

- Havard, J., Grygiel, J. and Sampson, D.
Determination by high-performance liquid chromatography of hydroxyurea in human plasma 584(1992)270
- Hawkes, W. C. and Kutnink, M. A.
High-performance liquid chromatographic determination of selenocysteine with the fluorescent reagent, N-(iodoacetylaminoethyl)-5-naphthylamine-1-sulfonic acid 576(1992)263
- Hawkins, D. R., see Cheng, K. N. 581(1992)203
- Hawthorne, S. B., see Eckert-Tilotta, S. E. 591(1992)313
- Hawthorne, S. B., see Langenfeld, J. J. 594(1992)297
- Hayakawa, K., see Yamamoto, A. 585(1991)315
- Hayashi, A., see Matsubara, T. 562(1991)119
- Hayashi, N., see Noma, J. 568(1991)35
- Hayashi, T., see Hashimoto, A. 582(1992)41
- Hayashi, Y., see Ishibashi, M. 562(1991)613
- Hayashi, Y., see Matsuda, R. 585(1991)187
- Hayatsu, H.
Cellulose bearing covalently linked copper phthalocyanine trisulphonate as an adsorbent selective for polycyclic compounds and its use in studies of environmental mutagens and carcinogens (Review) 597(1992)37
- Hayball, P. J., Nation, R. L., Bochner, F. and Le Leu, R. K.
Enantiospecific analysis of ketoprofen in plasma by high-performance liquid chromatography 570(1991)446
- Haymore, B. L., see Luong, C. B. H. 584(1992)77
- Haynes, P. A., Batley, M., Peach, R. J., Brennan, S. O. and Redmond, J. W.
Characterisation of oligosaccharides from a glycoprotein variant of human serum albumin (albumin Casebrook) using high-performance anion-exchange chromatography and nuclear magnetic resonance spectroscopy 581(1992)187
- Haynes, P. A., Sheumack, D., Greig, L. G., Kibby, J. and Redmond, J. W.
Applications of automated amino acid analysis using 9-fluorenylmethyl chloroformate 588(1991)107
- Hayward Vermaak, W. J., see Ubbink, J. B. 565(1991)441
- Hazen, M., see De Kok, T. M. C. M. 580(1992)135
- He, H. B., Deegan, R. J., Wood, M. and Wood, A. J. J.
Optimization of high-performance liquid chromatographic assay for catecholamines.
Determination of optimal mobile phase composition and elimination of species-dependent differences in extraction recovery of 3,4-dihydroxybenzylamine 574(1992)213
- He, L.-Y., see Oshima, T. 585(1991)255
- Head, S. L., see Burse, V. W. 566(1991)117
- Hearn, M. T. W., see Fang, F. W. 599(1992)163
- Hearn, M. T. W., see Johnston, A. 557(1991)335
- Hearn, M. T. W., see Purcell, A. W. 593(1992)103
- Hearn, M. T. W., see Zachariou, M. 599(1992)171
- Heath, P. C., see Brown, Jr., F. 595(1992)355
- Heckenberg, A. L., see Weston, A. 593(1992)289
- Hecker, M., Kaw, S. and Walsh, D. T.
Determination of endothelin-converting enzyme activity by high-performance liquid chromatography-on-line radioactive flow monitoring 574(1992)344
- Hedeen, K. M., Tyczkowska, K., Aucoin, D. P. and Norton, R. M.
Rapid high-performance liquid chromatographic method for the determination of propranolol levels in canine and feline plasma 572(1991)239
- Hedner, T., see Nyström, L. 574(1992)137
- Heebner, E. T., see Holland, M. L. 567(1991)433
- Heeremans, C. E. M., Stijnen, A. M., Van der Hoeven, R. A. M., Niessen, W. M. A., Danhof, M. and Van der Greef, J.
Liquid chromatography-thermospray tandem mass spectrometry for identification of a heptabarbital metabolite and sample work-up artefacts 554(1991)205
- Heeremans, C. E. M., see Poon, G. K. 576(1992)235
- Hegedus, L., see Deutsch, J. 579(1992)93
- Heilenthal, G. and Grummt, J.
Isolation of multiple protein factors involved in ribosomal DNA transcription 587(1991)25
- Hein, S. J., Piepmeier, E. H. and Thomas, L. C.
Laser-excited fluorescence detection of gas-phase chromatography eluates 557(1991)39
- Heinzen, V. E. F. and Yunes, R. A.
Relationship between gas chromatographic retention indices and molecular connectivity indices of chlorinated pesticides and structurally related compounds 598(1992)243
- Hekstra, D., see Van der Mast, C. A. 564(1991)115
- Helander, A., Beck, O. and Borg, S.
Determination of urinary 5-hydroxytryptophol by high-performance liquid chromatography with electrochemical detection 579(1992)340
- Heldin, E., Hang Huynh, N. and Pettersson, C.
(2*S*,3*S*)-Dicyclohexyl tartrate as mobile phase additive for the determination of the enantiomeric purity of (*S*)-atropine in tablets 592(1992)339
- Heldin, E., Huynh, N. H. and Pettersson, C.
(2*R*,3*R*)-Dicyclohexyl tartrate as a chiral mobile phase additive 585(1991)35
- Hell, F., see Flesch, G. 581(1992)147
- Hellenäs, K.-E., Nyman, A., Slanina, P., Löf, L. and Gabrielsson, J.
Determination of potato glycoalkaloids and their aglycone in blood serum by high-performance liquid chromatography. Application to pharmacokinetic studies in humans 573(1992)69
- Hellendoorn, L., see Schreurs, M. 552(1991)625
- Helmer, J. C., see Salomon, K. 559(1991)69
- Helmlin, H.-J. and Brenneisen, R.
Determination of psychotropic phenylalkylamine derivatives in biological matrices by high-performance liquid chromatography with photodiode-array detection 593(1992)87
- Helms, M., see Zobel, H. 553(1991)441
- Helusová, Š., see Turková, J. 597(1992)19
- Hemmer, R., see Arnold, P. J. 544(1991)267
- Hench, B. D., see Ruckmick, S. C. 565(1991)277
- Henckens, L. H. G., see Coenen, A. J. J. M. 596(1992)59
- Henderson, R. J., see Burkow, I. C. 552(1991)501

- Hendrich, C. E., Berdecia-Rodriguez, J., Wiedmeier, V. T. and Porterfield, S. P.
Method for the quantitation of iodothyronines in body tissues and fluids using high-performance liquid chromatography 577(1992)19
- Hendry, I. R., see Kearns, G. L. 574(1992)356
- Hendry, I. R., see Wells, T. G. 570(1991)235
- Henion, J. D., see Johansson, I. M. 554(1991)311
- Henion, J. D., see Johansson, I. M. 559(1991)515
- Henion, J. D., see McLaughlin, L. G. 591(1992)195
- Henion, J. D., see Rule, G. S. 582(1992)103
- Henke, D., see Liebmann, B. 572(1991)181
- Henke, W., see Ziegler, M. 563(1991)172
- Hennel, B., see Jacob, K. 572(1991)317
- Henning, K., see Cremer, J. 558(1991)430
- Henning, K., see Cremer, J. 587(1991)71
- Hennion, M.-C., see Coquart, V. 553(1991)329
- Hennion, M.-C., see Coquart, V. 585(1991)67
- Hennion, M.-C., see Coquart, V. 600(1992)195
- Henriksson, A., see Dizdar, N. 565(1991)1
- Henry, R., see Le Quéré, J. L. 562(1991)659
- Heo, G. S., see Lee, K.-J. 559(1991)317
- Heo, G. S., see Lee, K.-J. 577(1992)135
- Herault, J., see Jouve, J. 567(1991)331
- Herault, J., see Jouve, J. 574(1992)9
- Herbretau, B., see Morin-Allory, L. 590(1992)203
- Hercelin, B., see Leroy, P. 564(1991)258
- Herényi, B. and Görög, S.
Chiral high-performance liquid chromatographic separations on an α_1 -acid glycoprotein column. II. Separation of the diastereomeric and enantiomeric analogues of vinpocetine (Cavinton) 592(1992)297
- Heringa, R. M., see Coenen, A. J. J. M. 593(1992)243
- Hermansson, J., see Wiese, B. 567(1991)175
- Hermes, H. F. M., see Duchateau, A. L. L. 566(1991)493
- Hernandez, A., see Alton, K. B. 579(1992)307
- Hernandez, L., see Tellez, S. 581(1992)257
- Hernández, L. M., see Tabera, J. 557(1991)481
- Hernández, L. M., see Vidal-Madjar, C. 584(1992)11
- Herold, D. A., see Aggarwal, S. K. 576(1992)297
- Herold, M. L., Constanzer, M. L. and Matuszewski, B. K.
Determination of picogram levels of heptylphysostigmine in human plasma using high-performance liquid chromatography with fluorescence detection 581(1992)227
- Heron, S. and Tchaplá, A.
Description of retention mechanism by solvophobic theory. Influence of organic modifiers on the retention behaviour of homologous series in reversed-phase liquid chromatography 556(1991)219
- Herráez-Hernández, R., Campins-Falcó, P. and Sevillano-Cabeza, A.
Determination of acetazolamide in human urine samples by reversed-phase high-performance liquid chromatography in the presence of xanthines 582(1992)181
- Herregóts, P., see Sarre, S. 575(1992)207
- Herring, V. L., Bastian, T. L. and Lalonde, R. L.
Solid-phase extraction and direct high-performance liquid chromatographic determination of metoprolol enantiomers in plasma 567(1991)221
- Herrmann, S. C. and Feigl, E. O.
Subtraction method for the high-performance liquid chromatographic measurement of plasma adenosine 574(1992)247
- Hervé, F., Duché, J.-C., Barré, J., Millot, M.-C. and Tillement, J.-P.
pH titration curves of the desialylated human α_1 -acid glycoprotein variants by combined isoelectrofocusing-electrophoresis: utilization in the development of a fractionation method for the protein variants by chromatography on immobilized metal affinity adsorbent 577(1992)43
- Herweck, U., Zimmerman, H. and Reichling, J.
Suitable chiral packing material for the high-performance liquid chromatographic separation of derivatives of 1'-hydroxyeugenol 600(1992)312
- Herzog, R., see Zech, K. 553(1991)55
- Hesselmann, C., see Bringmann, G. 595(1992)351
- Hessels, J., Kingma, A. W., Sturkenboom, M. C. J. M., Elzinga, H., Van den Berg, G. A. and Muskiet, F. A. J.
Gas chromatographic determination of N-acetylisoputrescine- γ -lactam, a unique catabolite of N¹-acetylspermidine 563(1991)1
- Hessing, M., Van Schijndel, H. B., Van Grunsven, W. M. J., Wolf, H. and Middeldorp, J. M.
Purification and quantification of recombinant Epstein-Barr viral glycoproteins gp350/220 from Chinese hamster ovary cells 599(1992)267
- Hesso, A., see Rosenberg, C. 552(1991)265
- Hesso, A., see Shapi, M. M. 562(1991)681
- Hetmanski, M. T. and Scudamore, K. A.
Detection of zearalenone in cereal extracts using high-performance liquid chromatography with post-column derivatization 588(1991)47
- Heublein, G., see Erler, U. 588(1991)340
- Heykants, J., see Woestenborghs, R. 583(1992)223
- Hida, M., Mitsui, T. and Fujimura, Y.
Determination of succinic acid by pyrolysis-gas chromatography in the presence of sodium sulfite and iron powder 585(1991)345
- Hider, R. C., see Epemolu, R. O. 573(1992)178
- Hiemke, C., see Härtter, S. 578(1992)273
- Hiemke, C., see Süß, S. 565(1991)363
- Hiemke, C., see Weigmann, H. 583(1992)209
- Hiemstra, Y., see Welling, G. W. 599(1992)157
- Highsmith, F., see Tharakan, J. 595(1992)103
- Highsmith, F. A., see Orthner, C. L. 558(1991)55
- Higton, D. M., see Bowers, G. D. 554(1991)175
- Higuchi, S., see Usui, T. 584(1992)213
- Hilaireau, P., see Colin, H. 557(1991)137
- Hill, K. D., see Chamkasem, N. 587(1991)185
- Hill, M. and Arrio, B.
Activation of matrices by 4,6-diphenylthieno[3,4-*d*]-1,3-dioxol-2-one-5,5-dioxide. High-performance liquid affinity chromatographic separations 589(1992)101

- Hillemans, M. G., see Duchateau, A. L. L. 566(1991)493
- Hillert, A., see Süß, S. 565(1991)363
- Himmel, H. M., Sadony, V. and Ravens, U.
Quantitation of hypoxanthine in plasma from patients with ischemic heart disease: adaption of a high-performance liquid chromatographic method 568(1991)105
- Hindriks, H., see Hamoir, T. 589(1992)31
- Hinze, W. L., see Berthod, A. 556(1991)263
- Hinze, W. L., see Cole, R. O. 557(1991)113
- Hirabayashi, J. and Kasai, K.
Arginine-tail method, an affinity tag procedure utilizing anhydrotrypsin agarose 597(1992)181
- Hirabayashi, N., see Ikegawa, S. 577(1992)229
- Hirachi, N., see Kawaguchi, S. 567(1991)11
- Hirahara, K., see Ohsawa, K. 597(1992)285
- Hirai, E., see Okamoto, J. 583(1992)45
- Hirai, E., see Okamoto, J. 583(1992)53
- Hirai, S.-I., see Yamaguchi, M. 575(1992)123
- Hirai, Y., see Tomokuni, K. 567(1991)65
- Hiramatsu, K., see Okuda, T. 567(1991)141
- Hirano, K., see Nakanishi, A. 591(1992)159
- Hiraoka, B. Y. and Harada, M.
Use of L-prolyl-L-leucylglycinamide (MIF-1) for the high-performance liquid chromatographic determination of proline iminopeptidase activity in rat liver 563(1991)142
- Hirasawa, N., see Watanabe, M. 597(1992)309
- Hirata, H., see Okutani, R. 577(1992)25
- Hirata, H., see Sakaki, K. 585(1991)117
- Hirata, N., Tamura, Y., Kasai, M., Yanagihara, Y. and Noguchi, K.
New, stable polyamine-bonded polymer gel column 592(1992)93
- Hirayama, F., see Oshima, T. 585(1991)255
- Hirayama, S. and Maruyama, M.
Determination of a small amount of niacin in foodstuffs by high-performance liquid chromatography 588(1991)171
- Hirokawa, T., Omori, A., Yokota, Y., Hu, J.-Y. and Kiso, Y.
Isotachophoretic separation of minor components from a matrix component in the case of strong electrolytes 585(1991)297
- Hirokawa, T., Sugino, R. and Kiso, Y.
Electrophoretic mobility and dissociation constants of tripeptides evaluated by isotachopheresis 585(1991)364
- Hirokawa, T., Taka, T., Yokota, Y. and Kiso, Y.
Dilution effect of carbonate ion on isotachophoretic zones observed with scanning UV photometric detection 555(1991)247
- Hirokawa, T., Yokota, Y., Takaba, H., Nakahara, K. and Kiso, Y.
Isotachophoretic analyser with options for operational electrolyte selection and repeated analysis 555(1991)321
- Hirokawa, T., see Hu, J.-Y. 589(1992)339
- Hirokawa, T., see Hu, J.-Y. 594(1992)371
- Hirokawa, T., see Polášek, M. 596(1992)265
- Hirose, S., see Hagiwara, H. 597(1992)331
- Hirota, T., see Ohmori, S. 566(1991)1
- Hirth, W., see Rosenspire, K. C. 574(1992)119
- Hizukuri, S., see Koizumi, K. 585(1991)233
- Hjelm, M., see Lynes, G. W. 562(1991)213
- Hjertén, S., see Liao, J.-L. 586(1991)21
- Hlavačková, E. and Reya, R.
Comparison of colorimetric and chromatographic determination of *tert*-butylpyrocatechol 595(1992)137
- Ho, C.-H., see Griest, W. H. 600(1992)273
- Ho, M. Y. K., see Chen, T. K. 596(1992)123
- Ho, S.-T., Wang, J.-J., Ho, W. and Hu, O. Y.-P.
Determination of buprenorphine by high-performance liquid chromatography with fluorescence detection: application to human and rabbit pharmacokinetic studies 570(1991)339
- Ho, W., see Ho, S.-T. 570(1991)339
- Hoaksey, P. E., Awadzi, K., Ward, S. A., Coventry, P. A., Orme, M. L.E. and Edwards, G.
Rapid and sensitive method for the determination of albandazole and albandazole sulphoxide in biological fluids 566(1991)244
- Hobo, T., see Watabe, K. 590(1992)289
- Hochstrasser, D. F., see Toussi, A. 582(1992)87
- Hochuli, E., see Loetscher, P. 595(1992)113
- Hodgdon, H. L., see Phillips, D. J. 599(1992)239
- Hodges, R. S., see Hanson, M. 599(1992)65
- Hodges, R. S., see Hanson, M. 599(1992)77
- Hodges, R. S., see Zhu, B.-Y. 594(1992)75
- Hodny, Z., Struzinsky, R. and Deyl, Z.
Silver staining of collagen type I after sodium dodecylsulphate polyacrylamide gel electrophoresis: effect of Maillard reaction 578(1992)53
- Hodný, Z., see Syrový, I. 569(1991)175
- Hodr, R., see Knobloch, E. 566(1991)89
- Hodson, B. A., Pepper, D. S. and Dawes, J.
Novel affinity chromatographic system for the single-step purification of glycosaminoglycans from complex systems using volatile buffers 565(1991)416
- Hoffman, A., see Armando, I. 568(1991)45
- Hoffman, D. A., Wallace, S. M. and Verbeeck, R. K.
Simple method for the determination of inorganic sulfate in human serum and urine using single-column ion chromatography 565(1991)447
- Hoffman, J. L.
Ion chromatographic analysis of the purity and synthesis of sulfonium and selenium ions 588(1991)211
- Hoffman, N. E., see Welch, K. J. 591(1992)75
- Hoffmann, K.-J., see Gabrielsson, M. 573(1992)265
- Hofmann, R., see Glässgen, W. E. 598(1992)81
- Höft, E., see Kunath, A. 588(1991)352
- Hogeland, K., see Svoboda, M. 562(1991)403
- Hogendoorn, E. A., Hoogerbrugge, R., Goewie, C. E., Van Zoonen, P. and Schoenmakers, P. J.
Development of a rational optimisation procedure for the automated sample clean-up with column switching in pesticide residue analysis 552(1991)113
- Holada, K., see Suttner, J. 563(1991)184
- Holder, G. M., see Carter, S. R. 574(1992)77
- Holland, K. T., see Mallet, A. I. 562(1991)647

- Holland, M. L. and Heebner, E. T.
High-performance liquid chromatographic assay with ultraviolet detection for the determination of etoperidone and two active metabolites, 5-(1-hydroxyethyl)etoperidone and 1-(3-chlorophenyl)piperazine, in plasma 567(1991)433
- Hollenhorst, T. and Blaschke, G.
Direct separation of the enantiomers of propafenone, diprafenone and their major metabolites by high-performance liquid chromatography on modified cellulose and amylose chiral stationary phases 585(1991)329
- Holler, J. S., see Burse, V. W. 566(1991)117
- Höller, M., see Weber, C. 562(1991)599
- Hollingsworth, E. B., see Chang, S. Y. 562(1991)111
- Holthe, M. R., Brørs, O., Cragoe, Jr., E. J., Veel, T. and Ræder, M.
Determination of active 5-(N,N-hexamethylene)amiloride in pig plasma 579(1992)376
- Holthuis, J. J. M., see Steijger, O. M. 557(1991)13
- Holthuis, J. J. M., see Van der Horst, F. A. L. 567(1991)161
- Holtzhauser, M., see Boeden, H.-F. 552(1991)389
- Homandberg, G. A., see Kang, K. 567(1991)57
- Homma, M., see Kanai, M. 567(1991)415
- Honda, A., Shoda, J., Tanaka, N., Matsuzaki, Y., Osuga, T., Shigematsu, N., Tohma, M. and Miyazaki, H.
Simultaneous assay of the activities of two key enzymes in cholesterol metabolism by gas chromatography-mass spectrometry 565(1991)53
- Honda, S., Taga, A., Kakehi, K., Koda, S. and Okamoto, Y.
Determination of cefixime and its metabolites by high-performance capillary electrophoresis 590(1992)364
- Honda, S., Taga, A., Suzuki, K., Suzuki, S. and Kakehi, K.
Determination of the association constant of monovalent mode protein-sugar interaction by capillary zone electrophoresis 597(1992)377
- Honda, S., Yamamoto, K., Suzuki, S., Ueda, M. and Kakehi, K.
High-performance capillary zone electrophoresis of carbohydrates in the presence of alkaline earth metal ions 588(1991)327
- Hoogerbrugge, C. M., see Bayne, S. 562(1991)391
- Hoogerbrugge, R., see Hogendoorn, E. A. 552(1991)113
- Hoogland, G. J. M., see Van de Nesse, R. J. 552(1991)613
- Hoogland, H., see Ackermans, M. T. 596(1992)101
- Hoogmartens, J., see Cachet, T. 600(1992)99
- Hoogmartens, J., see Naidong, W. 586(1991)61
- Hoogmartens, J., see Naidong, W. 586(1991)67
- Hopkins, N. K., Wagner, C. M., Brisson, J. and Addison, T. E.
Validation of the simultaneous determination of methylprednisolone and methylprednisolone acetate in human plasma by high-performance liquid chromatography 577(1992)87
- Hoppel, C. L., see Krahenbuhl, S. 573(1992)3
- Horie, M., see Saito, K. 595(1992)163
- Horimoto, S., see Banno, K. 564(1991)1
- Horimoto, S., see Banno, K. 568(1991)375
- Horimoto, S., see Matsuoka, M. 581(1992)65
- Hormazabal, V. and Rogstad, A.
Simultaneous determination of sulphadiazine and trimethoprim in plasma and tissues of cultured fish for residual and pharmacokinetic studies 583(1992)201
- Horna, A., see Churáček, J. 557(1991)523
- Horowitz, J. D., see Sallustio, B. C. 576(1992)321
- Hörsch, B., see Angerer, J. 580(1992)229
- Horváth, C., see Antia, F. D. 556(1991)119
- Horváth, C., see Hung-Jye Lin, F. 589(1992)185
- Horváth, G., see Tomori, É. 578(1992)91
- Hoschützky, H., see Zimmerhackl, L. B. 587(1991)81
- Hosoya, K., see Kimata, K. 558(1991)19
- Hosoya, K., see Kimata, K. 595(1992)77
- Hostettmann, K., see Slacanian, I. 557(1991)391
- Hothi, H. S., see Sud, D. 596(1992)281
- Houben, R. J., Leclercq, P. A. and Cramers, C. A.
Ionization mechanisms in capillary supercritical fluid chromatography-chemical ionization mass spectrometry 554(1991)351
- Houbová, H., see Stehlik, P. 565(1991)477
- Houghton, E., see Dumasia, M. C. 564(1991)503
- House, R. M., see Palladino, D. E. H. 599(1992)3
- Houzé, P., De Reynies, A., Baud, F. J., Benatar, M. F. and Pays, M.
Simultaneous determination of chloroquine and its three metabolites in human plasma, whole blood and urine by ion-pair high-performance liquid chromatography 574(1992)305
- Hovestad, A., see Haanappel, V. A. C. 562(1991)531
- Hoving, E. B., Muskiet, F. A. J. and Christie, W. W.
Separation of cholesterol esters by silver ion chromatography using high-performance liquid chromatography or solid-phase extraction columns packed with a bonded sulphonic acid phase 565(1991)103
- Howell, J. A., see Leaver, G. 590(1992)101
- Hoyer, G. L.
Pirmenol determination by high-performance liquid chromatography 565(1991)497
- Hoyer, G. L., Clawson, D. C., Brookshier, L. A., Nolan, P. E. and Marcus, F. I.
High-performance liquid chromatographic method for the quantitation of quinidine and selected quinidine metabolites 572(1991)159
- Hoyt, Jr., A. M., see Arrowood, S. 583(1992)105
- Hoyt, Jr., A. M., see Arrowood, S. 586(1991)177
- Hrabák, P., see Tvřická, E. 563(1991)188
- Hradil, J., see Dernovaya, L. A. 552(1991)365
- Hruby, V. J., see Lebl, M. 586(1991)145
- Hsieh, C.-Y. and Huang, J.
Two-dimensional high-performance liquid chromatographic method to assay *p*-hydroxyphenylphenylhydantoin enantiomers in biological fluids and stereoselectivity of enzyme induction in phenytoin metabolism 575(1992)109
- Hsieh, J. Y.-K., see Kline, W. F. 578(1992)31
- Hsieh, J. Y.-K., see Robinett, R. S. R. 570(1991)157
- Hsu, M.-C. and Huang, W. F.
Collaborative study of the determination of cloxacillin by column liquid chromatography 600(1992)333

- Hsu, R. S., DiLeo, E. M., Chesson, S. M., Klein, J. T. and Effland, R. C.
Determination of HP 749, a potential therapeutic agent for Alzheimer's disease, in plasma by high-performance liquid chromatography 572(1991)352
- Hu, J.-Y., Hirokawa, T., Nishiyama, F. and Kiso, Y.
Study of isotachophoretic separation behaviour of metal cations by means of particle-induced X-ray emission. II. Separation efficiency of lanthanoid model mixture 589(1992)339
- Hu, J.-Y., Hirokawa, T., Nishiyama, F., Kiso, Y., Ito, K. and Shoto, E.
Study of isotachophoretic separation behaviour of metal cations by means of particle-induced X-ray emission. III. Analysis of a crude rare earth chloride from monazite 594(1992)371
- Hu, J.-Y., see Hirokawa, T. 585(1991)297
- Hu, O. Y.-P., see Ho, S.-T. 570(1991)339
- Hu, W., Haraguchi, H. and Takeuchi, T.
Analysis of nucleotides by high-performance liquid chromatography with phosphorus-selective detection 557(1991)441
- Hua, R. L., see Peng, C. T. 589(1992)231
- Huang, B. L., see Kiang, C.-H. 567(1991)195
- Huang, C.-Y., see Wen, K.-C. 593(1992)191
- Huang, E. C., see Johansson, I. M. 554(1991)311
- Huang, H.-S., Wu, J.-R. and Chen, M.-L.
Reversed-phase high-performance liquid chromatography of amphoteric β -lactam antibiotics: effects of columns, ion-pairing reagents and mobile phase pH on their retention times 564(1991)195
- Huang, J., see Hsieh, C.-Y. 575(1992)109
- Huang, L.-H., see Cheng, C. 555(1991)272
- Huang, S., see Clare, R. A. 562(1991)237
- Huang, S.-K., see Xu, D.-K. 567(1991)451
- Huang, W. F., see Hsu, M.-C. 600(1992)333
- Huang, X. C., Stuart, S. G., Bente III, P. F. and Brennan, T. M.
Capillary gel electrophoresis of single-stranded DNA fragments with UV detection 600(1992)289
- Huang, Y.-S., see Redden, P. R. 567(1991)21
- Huang, Z.-F., see Zeng, Z.-R. 589(1992)368
- Huber, C. G., Oefner, P. J. and Bönn, G. K.
High-performance liquid chromatographic separation of detritylated oligonucleotides on highly cross-linked poly(styrene-divinylbenzene) particles 599(1992)113
- Huber, O., see Toussi, A. 582(1992)87
- Huber, R. and Schildnitz, U.
Sensitive determination of the phosphodiesterase III/IV inhibitor zardaverine in human serum by direct sample injection, automated precolumn clean-up and high-performance liquid chromatography 553(1991)65
- Hubert, P., see Traifort, E. 571(1991)231
- Hubert, P., see Ubrich, N. 584(1992)17
- Hudson, M. K., see Mofidi, M. 589(1992)241
- Huf, F. A., see Ploum, M. E. 564(1991)413
- Hughes, G. J., see Toussi, A. 582(1992)87
- Hughes, M. K., see Dillon, J. G. 572(1991)41
- Huh, H., Staba, E. J. and Singh, J.
Supercritical fluid chromatographic analysis of polyprenols in *Ginkgo biloba* L. 600(1992)364
- Hühnerfuss, H. and Kallenborn, R.
Chromatographic separation of marine organic pollutants (Review) 580(1992)191
- Huie, C. W., see Aiken, J. H. 584(1992)181
- Huie, C. W., see Aiken, J. H. 588(1991)295
- Huie, C. W., see Wu, N. 582(1992)77
- Huigen, Y. M., see Gelsema, W. J. 594(1992)217
- Huisman, T. H. J., see Kutlar, F. 567(1991)359
- Hujer, R., see Knobloch, E. 566(1991)89
- Hulak, I., see Santambien, P. 597(1992)315
- Humbert, P., see Makki, S. 563(1991)407
- Hundt, H. K. L., see Brown, L. W. 582(1992)268
- Hung, C. T., see Wanwimolruk, S. 572(1991)227
- Hung, M.-F., see Kastrissios, H. 577(1992)103
- Hung-Jye Lin, F. and Horváth, C.
Measurement of stability constants for complexes of divalent cations with tartaric acid by ion chromatography 589(1992)185
- Hunt, P., see Vicat, P. 575(1992)315
- Hüntemann, S. B., see Seitz, J. 587(1991)55
- Hupe, K.-P., see Debets, A. J. J. 600(1992)163
- Hupe, K.-P., see Zech, K. 600(1992)161
- Hurni, W.M. and Miller, W.J.
Analysis of a vaccine purification process by capillary electrophoresis 559(1991)337
- Hurst, H. L., see Bushway, R. J. 587(1991)321
- Hurwitz, M. L., see Ganhaio, M. F. 564(1991)55
- Husain, S., Nageswara Sarma, P., Swamy, N. S., Alvi, S. N. and Nageswara Rao, R.
High-performance liquid chromatographic separation and determination of small amounts of 2-naphthaleneacetic acid in 1-naphthaleneacetic acid 558(1991)435
- Husain, S., Narsimha, R., Alvi, S. N. and Rao, R. N.
Studies on photoisomerization of 4,4'-diaminostilbene-2,2'-disulphonic acid for quality assurance by high-performance liquid chromatography 596(1992)127
- Husain, S., Narsimha, R., Alvi, S. N. and Rao, R. N.
Monitoring the effluents of the trichloroacetic acid process by high-performance liquid chromatography 600(1992)316
- Hušek, P.
Rapid derivatization and gas chromatographic determination of amino acids 552(1991)289
- Hušek, P., see Badoud, R. 552(1991)345
- Huselton, C. A., see Fayer, B. E. 568(1991)135
- Hussain, M. D., Tam, Y. K., Finegan, B. A. and Coutts, R. T.
Simple and sensitive high-performance liquid chromatographic method for the determination of diltiazem and six of its metabolites in human plasma 582(1992)203
- Hutchins, J. E., see Burton, S. D. 571(1991)209
- Huupponen, R., Ojala-Karlsson, P., Rouru, J. and Koulu, M.
Determination of metformin in plasma by high-performance liquid chromatography 583(1992)270

- Huwe, J. K.
Artifact formation during gas chromatographic-mass spectrometric analysis of a methylsulfinyl-containing metabolite 575(1992)287
- Huynh, N. H., see Heldin, E. 585(1991)35
- Hwang, B. Y.-H., see Boppana, V. K. 593(1992)29
- Hyland, K. and Bottiglieri, T.
Measurement of total plasma and cerebrospinal fluid homocysteine by fluorescence following high-performance liquid chromatography and precolumn derivatization with *o*-phthaldialdehyde 579(1992)55
- Hyvönen, T., Keinänen, T. A., Khomutov, A. R., Khomutov, R. M. and Eloranta, T. O.
Monitoring of the uptake and metabolism of aminoxy analogues of polyamines in cultured cells by high-performance liquid chromatography 574(1992)17
- I**
- Iannetti, P., see Pichini, S. 581(1992)143
- Ichiba, M., see Tomokuni, K. 567(1991)65
- Ichimiya, H., see Sakurai, T. 571(1991)1
- Iddamalgoda, P., see Haken, J. K. 600(1992)352
- Ide, H., Hamaguchi, K., Kobata, S., Murakami, A., Kimura, Y., Makino, K., Kamada, M., Miyamoto, S., Nagaya, T., Kamogawa, K. and Izumi, Y.
Purification of serine hydroxymethyltransferase from *Bacillus stearothermophilus* with ion-exchange high-performance liquid chromatography 596(1992)203
- Ide, T., see Murata, M. 579(1992)329
- Idle, J. R., see Boddy, A. V. 575(1992)137
- Idle, J. R., see Cholerton, S. 575(1992)325
- Idle, M. E., see Cholerton, S. 575(1992)325
- Iga, T., see Yamamura, Y. 567(1991)151
- Igarashi, K. and Castagnoli, Jr., N.
Determination of the pyridinium metabolite derived from haloperidol in brain tissue, plasma and urine by high-performance liquid chromatography with fluorescence detection 579(1992)277
- Iglesias, M. J., see Guillén, M. D. 591(1992)287
- Iida, T., Tamaru, T., Chang, F. C., Goto, J. and Nambara, T.
Capillary gas chromatographic analysis of oxo, oxo-hydroxy and unsaturated bile acid glycine conjugates 558(1991)451
- Iida, T., see Goto, J. 574(1992)1
- Iida, T., see Goto, J. 585(1991)281
- Iida, Y., Kinouchi, Y., Takeichi, Y., Imai, T. and Otagiri, M.
Simultaneous determination of a new dihydropyridine calcium antagonist (MPC-1304) and its metabolite in dog plasma by high-performance liquid chromatography with electrochemical detection 571(1991)277
- Ijichi, S., see Akasaka, K. 596(1992)197
- Ikarashi, Y., Iwatsuki, H., Blank, C. L. and Maruyama, Y.
Glassy carbon pre-column for direct determination of acetylcholine and choline in biological samples using liquid chromatography with electrochemical detection 575(1992)29
- Ikarashi, Y. and Maruyama, Y.
Applicability of high-performance liquid chromatography-continuous-flow fast atom bombardment mass spectrometry for simultaneous quantitation of multiple neurochemicals 587(1991)306
- Ikeda, M. and Kusaka, T.
Liquid chromatography-mass spectrometry of hydroxy and non-hydroxy fatty acids as amide derivatives 575(1992)197
- Ikeda, M., see Ohhashi, T. 585(1991)153
- Ikegami, K., see Goto, N. 578(1992)195
- Ikegami, K., see Goto, N. 578(1992)203
- Ikegami, Y., see Sano, K. 578(1992)63
- Ikegawa, S., Hirabayashi, N., Yoshimura, T., Tohma, M., Maeda, M. and Tsuji, A.
Determination of conjugated bile acids in human urine by high-performance liquid chromatography with chemiluminescence detection 577(1992)229
- Ikemoto, S., see Omi, T. 570(1991)399
- Ikewaki, K., see Shibusawa, Y. 596(1992)118
- Imai, K., see Toyo'oka, T. 588(1991)61
- Imai, T., see Iida, Y. 571(1991)277
- Imamura, I., see Fukuda, H. 567(1991)459
- Imanari, T., Toyoda, H., Yamanashi, S., Shinomiya, K., Koshiishi, I. and Oguma, T.
Study of the measurement of chondroitin sulphates in rabbit plasma and serum 574(1992)142
- Imanari, T., see Akiyama, H. 579(1992)203
- Imanari, T., see Toyoda, H. 565(1991)141
- Imbimbo, B. P., see Vree, T. B. 573(1992)168
- Imoto, T., Yamamoto, F. M., Miyasaka, A. and Hatano, H.
High-performance liquid chromatography-atmospheric pressure ionization mass spectrometry of gymnemic acids 557(1991)383
- Inczédy, J., see Bán, T. 593(1992)227
- Ingwersen, S. H.
Column liquid chromatographic assay of the dopamine reuptake inhibitor vanoxerine (GBR 12909) in human serum 571(1991)305
- Inotsume, N., see Kamizono, A. 567(1991)113
- Inoue, F., see Fujioka, T. 597(1992)429
- Inoue, H., see Watanabe, K. 566(1991)109
- Inoue, K., see Watanabe, H. 568(1991)445
- Inoue, Y., see Endoh, Y. S. 579(1992)123
- Inoue, Y., see Matsumoto, M. 562(1991)139
- Iohan, F., Vincze, I., Monder, C. and Cohen, S.
High-performance liquid chromatographic determination of cortolic and cortolonic acids as pyrenyl ester derivatives 564(1991)27
- Ip, D. P., see Tsai, E. W. 596(1992)217
- Iqbal, K., see Guzman, N. A. 559(1991)307
- Iqbal, K., see Guzman, N. A. 598(1992)123
- Iqbal, K., see Saska, M. 590(1992)147

- Iredale, J. and Wainer, I. W.
Determination of hydroxychloroquine and its major metabolites in plasma using sequential achiral-chiral high-performance liquid chromatography 573(1992)253
- Irth, H., see Reinhoud, N. J. 574(1992)327
- Irth, H., see Venizelos, V. 573(1992)259
- Isaksson, R., see Marle, I. 586(1991)233
- Iseki, K., see Kishino, S. 582(1992)246
- Ishibashi, E., see Wakui, Y. 575(1992)131
- Ishibashi, M., Watanabe, K., Ohyama, Y., Mizugaki, M., Hayashi, Y. and Takasaki, W.
Novel derivatization and immunoextraction to improve microanalysis of 11-dehydrothromboxane B₂ in human urine 562(1991)613
- Ishibashi, M., see Toyo'oka, T. 588(1991)61
- Ishida, J., Sonezaki, S. and Yamaguchi, M.
4,5-Diaminophthalhydrazide as a highly sensitive chemiluminescence derivatization reagent for α -dicarbonyl compounds in high-performance liquid chromatography 598(1992)203
- Ishigaki, I., see Kim, M. 585(1991)45
- Ishigaki, I., see Kim, M. 586(1991)27
- Ishihara, Y., see Endoh, Y. S. 579(1992)123
- Ishii, K., Banno, K., Miyamoto, T. and Kakimoto, T.
Determination of diltiazem hydrochloride enantiomers in dog plasma using chiral stationary-phase liquid chromatography 564(1991)338
- Ishii, S.
Preface 597(1992)1
- Ishii, Y., see Chiba, R. 588(1991)344
- Ishikawa, F., see Yamane, M. 579(1992)25
- Ishizaki, T., see Kobayashi, K. 579(1992)299
- Iso, S., see Uehara, N. 586(1991)356
- Isobe, T., Uchida, K., Taoka, M., Shinkai, F., Manabe, T. and Okuyama, T.
Automated two-dimensional liquid chromatographic system for mapping proteins in highly complex mixtures 588(1991)115
- Isosaki, K., Seno, N., Matsumoto, I., Koyama, T. and Moriguchi, S.
Immobilization of protein ligands with methyl vinyl ether-maleic anhydride copolymer 597(1992)123
- Issaq, H. J., see Atamna, I. Z. 588(1991)315
- Istin, B., see Girault, J. 564(1991)43
- Italia, A., Dosi, L. and Schiavi, M.
Improved high-performance liquid chromatographic resolution of the geometric isomers of 6-hydroxy-4-(1-hydroxy-1-methylethyl)-1-cyclohexene-1-ethanol and by-products with β -cyclodextrin 553(1991)15
- Ito, K., Ariyoshi, Y. and Sunahara, H.
Determination of inorganic anions in salt solutions by ion chromatography using C₁₈ reversed-phase columns coated with cetyltrimethylammonium 598(1992)237
- Ito, K., see Hu, J.-Y. 594(1992)371
- Ito, K., see Kawai, S. 585(1991)318
- Ito, N., see Abe, I. 565(1991)183
- Ito, S., see Ota, A. 593(1992)37
- Ito, T., see Hagiwara, H. 597(1992)331
- Ito, Y., see Shibusawa, Y. 596(1992)118
- Itoh, H., see Kawasaki, S. 595(1992)193
- Itoh, H., see Nimura, N. 585(1991)207
- Itoh, K., see Jinno, K. 594(1992)105
- Itoh, Y., see Okutani, R. 577(1992)25
- Itokawa, H., see Oshima, T. 585(1991)255
- Ittarat, I., Webster, H. K. and Yuthavong, Y.
High-performance liquid chromatographic determination of dihydroorotate dehydrogenase of *Plasmodium falciparum* and effects of antimalarials on enzyme activity 582(1992)57
- Ivanov, A., see Kostadinova, S. 568(1991)315
- Ivanova, N., see Lekova, K. 552(1991)353
- Iványi, T., see Van de Heijning, B. J. M. 565(1991)159
- Iwase, H.
Ultramicrodetermination of cyanocobalamin in elemental diet by solid-phase extraction and high-performance liquid chromatography with visible detection 590(1992)359
- Iwata, M., see Hattori, H. 579(1992)247
- Iwata, M., see Hattori, H. 581(1992)213
- Iwata, Y., see Kanai, M. 567(1991)415
- Iwatani, K., see Nakagawa, Y. 562(1991)99
- Iwatsuki, H., see Ikarashi, Y. 575(1992)29
- Izumi, Y., see Ide, H. 596(1992)203

J

- Jack, D. S., Rumble, R. H., Davies, N. W. and Francis, H. W.
Enantiospecific gas chromatographic-mass spectrometric procedure for the determination of ketoprofen and ibuprofen in synovial fluid and plasma: application to protein binding studies 584(1992)189
- Jackman, G. P., Colagrande, F. and Louis, W. J.
Validation of a solid-phase extraction high-performance liquid chromatographic assay for doxazosin 566(1991)234
- Jackowski, G., see Shi, Q. 572(1991)59
- Jackson, C. M., see Winzor, D. J. 597(1992)57
- Jackson, J., see Shaikh, B. 571(1991)189
- Jackson, P. E. and Jones, W. R.
Membrane-based sample preparation device for the pretreatment of acidic samples prior to cation analysis by ion chromatography 586(1991)283
- Jacob, I., see Royer-Morrot, M. J. 581(1992)297
- Jacob, K., Egeler, E., Hennel, B., Neumeier, D. and Lupp, P.
Application of ion-pair high-performance liquid chromatography to detection of the atypical coproporphyrin isomers II and IV in human faeces 572(1991)317

- Jacob, L., Beecken, V., Bartunik, L. J., Rose, M. and Bartunik, H. D.
Purification and crystallization of yeast hexokinase isoenzymes. Characterization of different forms by chromatofocusing 587(1991)85
- Jacob, III, P., Shulgin, A. T., Yu, L. and Benowitz, N. L.
Determination of the nicotine metabolite *trans*-3'-hydroxycotinine in urine of smokers using gas chromatography with nitrogen-selective detection or selected ion monitoring 583(1992)145
- Jacobs, A., see Leyssens, L. 564(1991)515
- Jacobsen, P. B., see Andresen, A. T. 575(1992)93
- Jacobson, S. C. and Guiochon, G.
Experimental study of the production rate of pure enantiomers from racemic mixtures 590(1992)119
- Jacobson, S. C. and Guiochon, G.
Contribution of ionically immobilized bovine serum albumin to the retention of enantiomers 600(1992)37
- Jacobsson, G., see Beck, O. 570(1991)198
- Jacquot, C., see Damaj, M. I. 563(1991)476
- Jagodzinski, J. J., Marshall, G. T., Poulsen, B. J., Raza, G. and Rolls, W. A.
Chromatography on Poly-RP and its cyano and diol derivatives using both polar and non-polar solvent systems 591(1992)89
- Jahanshahi, M., see Kajbaf, M. 575(1992)75
- Jahn, G. A., see Gulaid, A. A. 566(1991)228
- Jajoo, H. K., Bennett, S. M. and Kornhauser, D. M.
Thermospray liquid chromatographic-mass spectrometric analysis of anti-AIDS nucleosides: quantification of 2',3'-dideoxycytidine in plasma samples 577(1992)299
- Jakober, B., see Liebich, H. M. 572(1991)1
- Jakobs, C., see Zee, T. 574(1992)335
- Jamali, F., see Alessi-Severini, S. 582(1992)173
- Jamali, F., see Brocks, D. R. 581(1992)83
- Jamali, F., see Liang, W. T. C. 577(1992)317
- Jamali, F., see Palylyk, E. L. 568(1991)187
- Jamali, F., see Wright, M. R. 583(1992)259
- James, C. D., see Brown, A. R. 578(1992)302
- James, G. T., see Peloquin, C. A. 563(1991)472
- Janák, J., see Soják, L. 557(1991)241
- Janas, J., see Voelkel, A. 555(1991)205
- Jänchen, D. E., see De la Vigne, U. 553(1991)489
- Jandera, P. and Guiochon, G.
Effect of the sample solvent on band profiles in preparative liquid chromatography using non-aqueous reversed-phase high-performance liquid chromatography 588(1991)1
- Jandera, P. and Rozkošná, J.
Method for characterization of selectivity in reversed-phase liquid chromatography. V. Calibration of the retention scale for chromatographic systems with low concentrations of organic solvents in the mobile phase 556(1991)145
- Jandik, P., see Weston, A. 593(1992)289
- Jane, F., see Parra, P. 570(1991)185
- Janero, D. R., Yarwood, C. and Thakkar, J. K.
Application of solid-phase extraction on anion-exchange cartridges to quantify 5'-nucleotidase activity 573(1992)207
- Jang, M. D., see Park, J. H. 586(1991)1
- Jang, M. D., see Park, J. H. 595(1992)45
- Jang, Y., see Singh, A. K. 568(1991)351
- Janiczek, N., Bockbrader, H. N., Chang, T., Amidon, G. L. and Smith, D. E.
Stereoselective high-performance liquid chromatographic assay for pirlmenol enantiomers in dog plasma 571(1991)179
- Janini, G. M., see Atamna, I. Z. 588(1991)315
- Janiszewski, J. S., Mulvana, D. E., Kaul, S., Dandekar, K. A. and Barbhaiya, R. H.
High-performance liquid chromatographic determination of 2',3'-didehydro-3'-deoxythymidine, a new anti-human immunodeficiency virus agent, in human plasma and urine 577(1992)151
- Janiszewski, J. S., Shukla, U. A., Kaul, S., Dandekar, K. A. and Pittman, K. A.
High-performance liquid chromatographic assay of α -(4-fluorophenyl)-4-(5-fluoro-2-pyrimidinyl)-1-piperazine butanol (BMV 14802), a potential antipsychotic drug, in monkey and rat plasma 567(1991)274
- Janjić, T. J., Milojković, D. M., Vučković, G. N. and Čelap, M. B.
Thin-layer chromatography on polyacrylonitrile. IV. Investigation of the separation mechanisms for tris-(alkylxanthato)cobalt(III) complexes 596(1992)91
- Janjić, T. J., see Tešić, Ž. L. 585(1991)359
- Jansen, E. H. J. M. and De Fluiter, P.
Detection of the enzymatic activity of cytochrome P-450 enzymes by high-performance liquid chromatography (Review) 580(1992)325
- Jansen, E. H. J. M., Doorn, L. and Van Leeuwen, F. X. R.
Determination of proteolytic hydrolysis of thyroglobulin 566(1991)471
- Jansen, E. H. J. M. and Van den Berg, R. H.
High-performance liquid chromatographic investigation of product formation in the horseradish peroxidase-enhanced chemiluminescence of luminol with different enhancers 566(1991)461
- Jansen, E. H. J. M., Van den Berg, R. H., Both-Miedema, R. and Doorn, L.
Advantages and limitations of pre-column derivatization of amino acids with dansyl chloride 553(1991)123
- Jansen, E. H. J. M., Van Ginkel, L. A., Van den Berg, R. H. and Stephany, R. W.
High-performance liquid chromatographic separation and detection methods for anabolic compounds (Review) 580(1992)111
- Jansen, E. H. J. M., see Doorn, L. 553(1991)135
- Jansen, E. H. J. M., see Reinerink, E. J. M. 553(1991)233
- Jansen, E. H. J. M., see Van Look, L. J. 564(1991)451
- Jansen, R. W., see Molema, G. 579(1992)107
- Janssen, G., see Cachet, T. 600(1992)99
- Janssen, H.-G., Schoenmakers, P. J. and Cramers, C. A.
Effects of modifiers in packed and open-tubular supercritical fluid chromatography 552(1991)527
- Janssen, H.-G., see De Bokx, P. K. 598(1992)115
- Jansson, S. O., see Arvidsson, E. 591(1992)55
- Jäppinen, P., see Rosenberg, C. 552(1991)265

- Jarman, M., see Poon, G. K. 565(1991)75
 Jarman, M., see Poon, G. K. 572(1991)143
 Jarman, M., see Poon, G. K. 576(1992)235
 Jarrett, H. W., see Goss, T. A. 588(1991)157
 Jarrett, H. W., see Massom, L. R. 600(1992)221
 Jäschke, A., Cech, D. and Ehwald, R.
 Chromatographic fractionation of nucleic acids using microcapsules made from plant cells 585(1991)57
 Jäschke, A., Cech, D. and Ehwald, R.
 Inclusion and fractionated release of nucleic acids using microcapsules made from plant cells 596(1992)165
 Jauhainen, M., see Zhou, G. 568(1991)69
 Jaulmes, A., see Lemque, R. 599(1992)255
 Jaulmes, A., see Vidal-Madjar, C. 584(1992)11
 Jaussaud, P., Guieu, D., Courtot, D., Barbier, B. and Bonnaire, Y.
 Identification of a tolfenamic acid metabolite in the horse by gas chromatography-mass spectrometry 573(1992)136
 Jaussaud, P., see Benoit, E. 583(1992)167
 Jeansonne, M. S. and Foley, J. P.
 Improved equations for the calculation of chromatographic figures of merit for ideal and skewed chromatographic peaks 594(1992)1
 Jefferies, T. M., see Patel, H. B. 555(1991)21
 Jefferis, R., see Downham, M. 584(1992)59
 Jegorov, A., Trnka, T. and Stuchlík, J.
 High-performance liquid chromatographic detection of enantiomeric amino alcohols after derivatization with α -phthalaldehyde and various thiosugars 558(1991)311
 Jehl, F., see Berg-Candolfi, M. 593(1992)1
 Jehl, F., see Enjalbert, F. 598(1992)227
 Jehl, F., see Molina, F. 563(1991)205
 Jelinek, I., Snopek, J. and Smolková-Keulemansová, E.
 Influence of counter-ion inclusion complexation on the quality of cyclodextrin-supported separations in isotachopheresis 557(1991)215
 Jelinek, I., see Snopek, J. 559(1991)215
 Jelínková, M., see Kuchař, M. 557(1991)399
 Jellum, E., Thorsrud, A. K. and Time, E.
 Capillary electrophoresis for diagnosis and studies of human disease, particularly metabolic disorders 559(1991)455
 Jemal, M., see Both, D. A. 558(1991)257
 Jemal, M., see Both, D. A. 585(1991)348
 Jemal, M., see Both, D. A. 596(1992)85
 Jen, S. C. D. and Pinto, N. G.
 Theory of optimization of ideal displacement chromatography of binary mixtures 590(1992)3
 Jeng, C.-Y. and Langer, S. H.
 Hydroquinone oxidation kinetics in adsorptive liquid chromatographic beds 556(1991)383
 Jeng, C.-Y. and Langer, S. H.
 Reaction kinetics and kinetic processes in modern liquid chromatographic reactors (Review) 589(1992)1
 Jenke, D. R.
 Determination of solute-polymer interaction constants via chromatography with polymer capillaries 594(1992)29
 Jennissen, H. P. and Demiroglou, A.
 Base-atom recognition in protein adsorption to alkyl agaroses 597(1992)93
 Jessner, G., see Pichorner, H. 594(1992)400
 Jezorek, J. R., see Ji, D. 590(1992)189
 Ji, D. and Jezorek, J. R.
 Chromatographic effects of residual amino groups on liquid chromatographic stationary phases derived from aminopropyl-silica gel 590(1992)189
 Jiao, Y., Blaas, W., Rühl, C. and Weber, R.
 Identification of ochratoxin A in food samples by chemical derivatization and gas chromatography-mass spectrometry 595(1992)364
 Jiménez, B., see Tabera, J. 557(1991)481
 Jiménez, B., see Vidal-Madjar, C. 584(1992)11
 Jiménez, R., see Reuvers, T. 564(1991)477
 Jiménez, R. M., see Rodríguez, F. J. 578(1992)146
 Jinno, K., Saito, Y., Malhan née Chopra, R., Pesek, J. J., Fetzer, J. C. and Biggs, W. B.
 Retention behaviour of polycyclic aromatic hydrocarbons on a liquid-crystal bonded phase in reversed-phase liquid chromatography 557(1991)459
 Jinno, K., Yamamoto, K., Ueda, T., Nagashima, H., Itoh, K., Fetzer, J. C. and Biggs, W. R.
 Liquid chromatographic separation of all-carbon molecules C₆₀ and C₇₀ with multi-legged phenyl group bonded silica phases 594(1992)105
 Jinno, K., see Okamoto, M. 556(1991)407
 Jockisch, W., see Matucha, M. 588(1991)251
 Johansen, K., see Edlund, P. O. 553(1991)21
 Johansen, M., Tønnesen, F. and Rasmussen, K. E.
 Column-switching high-performance liquid chromatographic detection of pholcodine and its metabolites in urine with fluorescence and electrochemical detection 573(1992)283
 Johansson, I. M., Huang, E. C., Henion, J. D. and Zweigenbaum, J.
 Capillary electrophoresis-atmospheric pressure ionization mass spectrometry for the characterization of peptides. Instrumental considerations for mass spectrometric detection 554(1991)311
 Johansson, I. M., Pavelka, R. and Henion, J. D.
 Determination of small drug molecules by capillary electrophoresis-atmospheric pressure ionization mass spectrometry 559(1991)515
 Johansson, K. and Olin, Å.
 Determination of selenium by capillary gas chromatography after high-temperature derivatization with 1,2-diamino-3,5-dibromobenzene 598(1992)105
 Johnson, D., see Dollinger, G. 592(1992)215
 Johnson, D., see Kunitani, M. 588(1991)125
 Johnson, D. F., Bradshaw, J. S., Eguchi, M., Rossiter, B. E., Lee, M. L., Petersson, P. and Markides, K. E.
 Synthesis of (1*R*-*trans*)-N,N'-1,2-cyclohexylenebisbenzamide oligodimethylsiloxane copolymers for use as chiral stationary phases for capillary supercritical fluid chromatography 594(1992)283
 Johnson, H. D., see Katti, P. S. 566(1991)29
 Johnson, L., see Lönnroth, P. 568(1991)419

- Johnson, V. A., see Kearns, G. L. 574(1992)356
- Johnson, W. R., see Reiser, R. W. 554(1991)91
- Johnston, A. and Hearn, M. T. W.
High-performance liquid chromatography of amino acids, peptides and proteins. CXIV. Protein interactions with porous coulombic sorbents: comparison of experimental findings with predictions of several adsorption models 557(1991)335
- Johnston, G. A. R., see Yap, B. K. 573(1992)183
- Joly, J. M. and Williams, W. M.
High-performance liquid chromatographic analysis of 5-ethylpyrimidines and 5-methylpyrimidines in plasma 563(1991)392
- Jones, A. T. and Roberts, N. B.
Peptide maps of five human pepsin isoenzymes and other aspartic proteinases 599(1992)179
- Jones, H. A., Rhodes, C. G., Law, M. P., Becket, J. M., Clark, J. C., Boobis, A. R. and Taylor, G. W.
Rapid analysis for metabolites of ¹⁴C-labelled drugs: fate of [¹⁴C]-S-4-(*tert*-butylamino-2-hydroxypropoxy) benzimidazol-2-one in the dog 570(1991)361
- Jones, L. A., Glennon, J. J. and Reiss, W. H.
Critical capillary column examination of the relationship between separation number and height equivalent to a theoretical plate and their dependence on temperature 595(1992)209
- Jones, L. A., Reiss, W. H., Glennon, J. J. and Gerig, T. M.
Van Deemter-type relationship for determining the optimum initial flow-rate and optimum pressure programming rate in temperature/pressure-programmed capillary column gas chromatography utilizing separation numbers 595(1992)221
- Jones, M. D., see Robinson, P. R. 564(1991)147
- Jones, R., see Dollinger, G. 592(1992)215
- Jones, W. R., see Jackson, P. E. 586(1991)283
- Jones, W. R., see Weston, A. 593(1992)289
- Jongen, M. J. M., see Leenheers, L. H. 578(1992)189
- Jongeneelen, F. J., see Dirven, H. A. A. M. 564(1991)266
- Jongepier, U., see Michaelis, H. C. 571(1991)257
- Jonkman, J. H. G., see Wieling, J. 594(1992)45
- Jönsson, B. and Skarping, G.
Method for the biological monitoring of hexahydrophthalic anhydride by the determination of hexahydrophthalic acid in urine using gas chromatography and selected-ion monitoring 572(1991)117
- Jönsson, B., Welinder, H. and Skarping, G.
Determination of hexahydrophthalic anhydride in air using gas chromatography 558(1991)247
- Jönsson, J. Å.
Non-chromatographic continuous separation techniques (by M. Valcárcel and M. D. Luque de Castro) (Book Review) 587(1991)369
- Jönsson, J. Å., see Karlsson, L. 557(1991)99
- Jönsson, J. Å., see Lindegård, B. 573(1992)191
- Jordan, L. M. and Russo-Marie, F.
Purification and partial characterization of phospholipase A₂ isoforms from human placenta 597(1992)299
- Jørgensen, M., Andersen, M. P. and Hansen, S. H.
Simultaneous determination of nitroglycerin and its dinitrate metabolites by capillary gas chromatography with electron-capture detection 577(1992)167
- Jorgenson, J. W., see Deterding, L. J. 554(1991)73
- Jorgenson, J. W., see Deterding, L. J. 554(1991)329
- Josić, D., Reusch, J., Löster, K., Baum, O. and Reutter, W.
High-performance membrane chromatography of serum and plasma membrane proteins 590(1992)59
- Josić, D., see Abou-Rebyeh, H. 566(1991)341
- Josić, D., see Abou-Rebyeh, H. 566(1991)351
- Joubert, J. R., see Le Roux, A. M. 581(1992)306
- Jouve, J., Bakri, D., Herault, J. and Muh, J. P.
Determination of urinary vanillic acid and plasma dihydroxyphenylalanine as markers of non-secreting neuroblastoma by high-performance liquid chromatography with electrochemical detection 567(1991)331
- Jouve, J., Herault, J., Tournade, H. and Muh, J. P.
Determination by high-performance liquid chromatography with electrochemical detection of free and conjugated N-acetyldopamine excretion in urine of children with neuroblastoma and nephroblastoma 574(1992)9
- Jouzeau, J. Y., see Payan, E. 566(1991)9
- Joyce, J. G., see Lehman, E. D. 574(1992)225
- Jozefowicz, M., see Letourneur, D. 589(1992)87
- Juergens, U. H.
Analysis of sulthiame in serum by narrow-bore high-performance liquid chromatography. Comparison of direct sample injection with pre-column switching and extrelut extraction 553(1991)7
- Julien-Larose, C., Voirin, P., Mas-Chamberlin, C. and Dufour, A.
Use of particle beam liquid chromatography-electron impact mass spectrometry for structure elucidation of oxodipine and three of its metabolites 562(1991)39
- Jung, K.-H., see Youn, D. Y. 591(1992)19
- Jung, L., see Lave, T. 572(1991)203
- Jung, M., Schmalzing, D. and Schurig, V.
Theoretical approach to the gas chromatographic separation of enantiomers on dissolved cyclodextrin derivatives 552(1991)43
- Junker-Buchheit, A., see Francotte, E. 576(1992)1
- Juranić, N., see Vučković, G. 585(1991)181
- Jurisson, S., see Rosenspire, K. C. 574(1992)119
- Jursík, T., Stránský, K. and Ubik, K.
Trapping system for trace organic volatiles 586(1991)315
- Jusko, W. J., see Garg, V. 567(1991)39
- Just, U., see Pasch, H. 589(1992)295
- Justesen, U. and Bojesen, G.
Analysis of some eicosanoids by continuous-flow fast atom bombardment mass spectrometry 562(1991)59
- Juvet, Jr., R. S., see Lundeen, J. T. 556(1991)305

K

- Kaczvinsky, Jr., J. R. and Read, S. A.
Development and use of a stability-indicating high-performance liquid chromatographic assay for Meldrum's acid 575(1992)177
- Kadaba, P. K., see Stevenson, P. J. 563(1991)419
- Kage, S., Nagata, T. and Kudo, K.
Determination of polysulphides in blood by gas chromatography and gas chromatography-mass spectrometry 564(1991)163
- Kågedal, B., see Dizdar, N. 565(1991)1
- Kahie, Y. D., see Pietrogrande, M. C. 592(1992)65
- Kaijser, G. P., Beijnen, J. H., Bult, A., Wiese, G., De Kraker, J., Keizer, H. J. and Underberg, W. J. M.
Gas chromatographic determination of 2- and 3-dechloroethylifosfamide in plasma and urine 583(1992)175
- Kaijser, G. P., Beijnen, J. H., Bult, A., Wiese, G., De Kraker, J. and Underberg, W. J. M.
Gas chromatographic determination of ifosfamide in microvolumes of urine and plasma 571(1991)121
- Kaiser, Jr., R. E., Williams, J. D., Lammert, S. A., Cooks, R. G. and Zakett, D.
Thermospray liquid chromatography-mass spectrometry with a quadrupole ion trap mass spectrometer 562(1991)3
- Kajbaf, M., Jahanshahi, M., Pattichis, K., Gorrod, J. W. and Naylor, S.
Rapid and efficient purification of cimetropium bromide and mifentidine drug metabolite mixtures derived from microsomal incubates for analysis by mass spectrometry 575(1992)75
- Kajbaf, M., Sepai, O., Lamb, J. H. and Naylor, S.
Identification of metabolites of 4,4'-diaminodiphenylmethane (methylene dianiline) using liquid chromatographic and mass spectrometric techniques 583(1992)63
- Kaji, H., see Terazawa, K. 565(1991)453
- Kajii, E., see Omi, T. 570(1991)399
- Kajiwara, H.
Application of high-performance capillary electrophoresis to the analysis of conformation and interaction of metal-binding proteins 559(1991)345
- Takehi, K., see Honda, S. 588(1991)327
- Takehi, K., see Honda, S. 590(1992)364
- Takehi, K., see Honda, S. 597(1992)377
- Kakemoto, M.
Simultaneous determination of sorbic acid, dehydroacetic acid and benzoic acid by gas chromatography-mass spectrometry 594(1992)253
- Kakimoto, T., see Ishii, K. 564(1991)338
- Kakinoki, S., see Kishino, S. 582(1992)246
- Kakinuma, H., see Yokoyama, Y. 566(1991)19
- Kakinuma, H., see Yokoyama, Y. 583(1992)1
- Kalasinaky, K. S., see Kuhlman, Jr., J. J. 578(1992)207
- Kaleja, M., see Oberdorfer, F. 552(1991)483
- Kallenborn, R., see Hühnerfuss, H. 580(1992)191
- Kamada, M., see Ide, H. 596(1992)203
- Kamahori, M., Taki, M., Watanabe, Y. and Miura, J.
Analysis of plasma catecholamines by high-performance liquid chromatography with fluorescence detection: simple sample preparation for pre-column fluorescence derivatization 567(1991)351
- Kamata, K., Motohashi, N., Meyer, R. and Yamamoto, Y.
Determination of 7-methylbenz[c]acridines by capillary gas chromatography with electron-capture detection 596(1992)233
- Kamata, T., see Goto, N. 578(1992)195
- Kamberov, E., see Kostadinova, S. 568(1991)315
- Kamel, S., Brazier, M., Desmet, G., Picard, C., Mennecier, I. and Seberty, J.
High-performance liquid chromatographic determination of 3-hydroxypyridinium derivatives as new markers of bone resorption 574(1992)255
- Kamesaki, T., see Omi, T. 570(1991)399
- Kamiński, M.
Simple test for determination of the degree of distortion of the liquid-phase flow profile in columns for preparative liquid chromatography 589(1992)61
- Kamizono, A., Inotsume, N., Miyamoto, K., Ueda, K., Miyakawa, T., Arimoto, H. and Nakano, M.
Determination of sultopride and tiapride in serum by gas chromatography using a surface ionisation detector 567(1991)113
- Kammaing, D. A., see Kwakman, P. J. M. 553(1991)345
- Kamogawa, K., see Ide, H. 596(1992)203
- Kamp, H.-D., see Geisslinger, G. 568(1991)165
- Kamphuis, J., see Duchateau, A. L. L. 566(1991)493
- Kanai, M., Kouno, Y., Homma, M., Yamada, K., Oka, K., Noguchi, M., Abe, M., Sakakura, K. and Iwata, Y.
Simple and accurate determination of methylpyrazines in biofluids using high-performance liquid chromatography 567(1991)415
- Kanaji, Y., see Matsushita, S. 586(1991)363
- Kanda, H., see Watabe, K. 590(1992)289
- Kane, M., Fukunaga, T., Yamamoto, Y., Yamada, M. and Tatsuno, Y.
Electrophoretic phenotyping of erythrocyte enzymes (Review) 569(1991)297
- Kane, R. E., see Shansky, R. E. 589(1992)165
- Kaneta, T., see Tanaka, S. 587(1991)364
- Kang, K., Schwarz, D., Sky-Peck, H., Kuettner, K. E. and Homandberg, G. A.
Novel method of purification of human leukocytic elastase using adsorption on a high-performance liquid chromatography gel permeation column 567(1991)57
- Kaniansky, D., see Madajová, V. 589(1992)329
- Kantarjian, H. M., see Saunders, P. P. 577(1992)37
- Kapás, M., see Pudleiner, P. 553(1991)29
- Kapil, R. P., Padovani, P. K., King, S.-Y. P. and Lam, G. N.
Nanogram level quantitation of oxycodone in human plasma by capillary gas chromatography using nitrogen-phosphorus selective detection 577(1992)283

- Kaplan, B., Pras, M. and Ravid, M.
Isolation and purification of amyloid protein A by sodium dodecyl sulphate polyacrylamide gel electrophoresis and reversed-phase high-performance liquid chromatography 573(1992)17
- Karaiskakis, G., see Koliadima, A. 595(1992)237
- Karasek, F. W., see Beard, A. 589(1992)265
- Karege, F. and Rudolph, W.
High-performance liquid chromatographic determination of phenylacetic acid in human plasma extracted with ethyl acetate 570(1991)376
- Karger, B. L., see Chen, J. W. 559(1991)295
- Karger, B. L., see Lin, S. 557(1991)369
- Karl, V., Schmarr, H.-G. and Mosandl, A.
Simultaneous stereoanalysis of 2-alkyl-branched acids, esters and alcohols using a selectivity-adjusted column system in multi-dimensional gas chromatography 587(1991)347
- Karlsen, J., see Tallaksen, C. M. E. 564(1991)127
- Karlsson, L., Mathiasson, L., Åkesson, J. and Jönsson, J. Å.
Quantitative aspects of the determination of compounds with widely varying polarity using capillary supercritical fluid chromatography 557(1991)99
- Karlsson, R., see Fägerstam, L. G. 597(1992)397
- Kårnsnäs, P. and Lindblom, T.
Characterization of hydrophobic interaction and hydrophobic interaction chromatography media by multivariate analysis 599(1992)131
- Kasahara, T., see Okutani, R. 577(1992)25
- Kasai, K.
Trypsin and affinity chromatography (Review) 597(1992)3
- Kasai, K., see Hirabayashi, J. 597(1992)181
- Kasai, K.-I., see Abe, I. 565(1991)183
- Kasai, M., see Hirata, N. 592(1992)93
- Kasheva, T. N., see Galushko, S. V. 553(1991)143
- Kasheva, T. N., see Galushko, S. V. 600(1992)79
- Kashiwagi, S., see Nakamura, K. 597(1992)351
- Kašička, V. and Prusík, Z.
Application of capillary isotachopheresis in peptide analysis (Review) 569(1991)123
- Kastl, P. E., see Bartels, M. J. 575(1992)69
- Kastner, F., see Brandl, G. 586(1991)249
- Kastner, M. and Neubert, D.
High-performance metal chelate affinity chromatography of cytochromes P-450 using Chelating Superose 587(1991)43
- Kastner, M. and Neubert, D.
Isolation of cytochrome P-450 components from marmoset liver microsomes by high-performance liquid chromatography 587(1991)117
- Kastrissios, H., Hung, M.-F. and Triggs, E. J.
High-performance liquid chromatographic method for the quantitation of bupivacaine, 2,6-pipecoloxylidide and 4'-hydroxybupivacaine in plasma and urine 577(1992)103
- Kasuya, Y., see Furuta, T. 562(1991)363
- Kasuya, Y., see Furuta, T. 576(1992)213
- Kasuya, Y., see Shibasaki, H. 576(1992)47
- Kasuya, Y., see Shibasaki, H. 579(1992)193
- Kataeva, I. A., see Golovchenko, N. P. 591(1992)121
- Katagi, T., see Fujioka, T. 597(1992)429
- Kataoka, H., Ohru, S., Miyamoto, Y. and Makita, M.
Determination of urinary β -phenylethylamine as its N-benzenesulphonamide derivative by gas chromatography with flame photometric detection 578(1992)120
- Katayama, M., Masuda, Y. and Taniguchi, H.
Determination of alcohols by high-performance liquid chromatography after pre-column derivatization with 2-(4-carboxyphenyl)-5,6-dimethylbenzimidazole 585(1991)219
- Katayama, M., see Furuta, T. 576(1992)213
- Kato, Y. and Numajiri, Y.
Chloride attachment negative-ion mass spectra of sugars by combined liquid chromatography and atmospheric pressure chemical ionization mass spectrometry 562(1991)81
- Kato, H., see Takahara, E. 576(1992)174
- Kato, I., see Ohara, K. 586(1991)35
- Kato, T., see Ono, S. 597(1992)293
- Kato, Y., see Kobayashi, K. 579(1992)299
- Kato, Y., see Nozaki, O. 570(1991)1
- Katoh, T., see Ohba, S. 568(1991)325
- Katsanos, N. A. and Vassilakos, C.
Theoretical analysis for measurement of building pollution parameters by gas chromatography 557(1991)469
- Katsanos, N. A., see Koliadima, A. 595(1992)237
- Katti, A., Erlandsson, P. and Däppen, R.
Application of preparative liquid chromatography to the isolation of enantiomers of a benzodiazepinone derivative 590(1992)127
- Katti, A. M.
Experimental study on the effect of the sample size on the band profile for a binary mixture showing no competitive interaction 556(1991)195
- Katti, A. M., Czok, M. and Guiochon, G.
Prediction of single and binary profiles in overloaded elution chromatography using various semi-ideal models 556(1991)205
- Katti, A. M., see Katti, P. S. 566(1991)29
- Katti, A. M., see Zhu, J. 552(1991)71
- Katti, P. S., Katti, A. M. and Johnson, H. D.
Determination of heat-exposure effects on the concentration of catecholamines in bovine plasma and milk 566(1991)29
- Katz, R. N., see Roboz, J. 575(1992)281
- Katz, S. D., see Bleiberg, B. 568(1991)301
- Kaufmann, P., see Olsson, N. U. 600(1992)257
- Kaufmann, S. H. E., see Schoel, B. 587(1991)19
- Kaul, S., see Janiszewski, J. S. 567(1991)274
- Kaul, S., see Janiszewski, J. S. 577(1992)151
- Kavanagh, P. E., see Shalliker, R. A. 558(1991)440
- Kaw, S., see Hecker, M. 574(1992)344
- Kawagoe, J. L., see Pollack, G. M. 570(1991)406
- Kawaguchi, S., Hirachi, N. and Fukamachi, M.
New analyser for the determination of urinary vanillylmandelic acid, homovanillic acid and creatinine 567(1991)11

- Kawahara, J., see Otsuka, K. 559(1991)209
 Kawahara, S., see Nakano, M. 564(1991)235
 Kawahara, Y., see Nakanishi, A. 591(1992)159
 Kawahara, Y., see Shioya, H. 568(1991)309
 Kawai, H., see Shinkai, H. 570(1991)203
 Kawai, S., Hanai, K., Ito, K., Kitahara, S. and Kuwae, A.
 High-performance liquid chromatographic separation of
p-hydroxyphenylpyruvic acid 585(1991)318
 Kawai, S., see Tomita, M. 566(1991)239
 Kawai, S., see Tomita, M. 571(1991)324
 Kawai, T., see Okutani, R. 577(1992)25
 Kawakami, J., see Yamamura, Y. 567(1991)151
 Kawakami, Y., Ohga, T., Shimamoto, C., Satoh, N. and
 Ohmori, S.
 Determination of free N-acetylamino acids in biological
 samples and N-terminal acetylamino acids of proteins
 576(1992)63
 Kawamoto, M., see Morita, I. 576(1992)334
 Kawanishi, E., see Makino, Y. 567(1991)248
 Kawasaki, S., Ueda, H., Itoh, H. and Tadano, J.
 Screening of organophosphorus pesticides using liquid
 chromatography-atmospheric pressure chemical
 ionization mass spectrometry 595(1992)193
 Kawasaki, T., see Nakanishi, A. 591(1992)159
 Kawase, M., see Ohmori, S. 574(1992)35
 Kawashima, Y., see Ota, A. 593(1992)37
 Kayano, M., see Takamatsu, T. 584(1992)261
 Kazama, M., see Abe, I. 565(1991)183
 Kazlauskas, R., see Lisi, A. M. 563(1991)257
 Kazlauskas, R., see Lisi, A. M. 581(1992)57
 Kazlauskas, R., see Yap, B. K. 573(1992)183
 Kažoka, H. A., see Shatz, V. D. 552(1991)23
 Kearns, G. L., Johnson, V. A., Hendry, I. R. and Wells, T. G.
 Microanalytical high-performance liquid chromatographic
 assay for cefpirome in human milk and urine
 574(1992)356
 Kearns, G. L., see Wells, T. G. 570(1991)235
 Keely, B. J., see Eckardt, C. B. 557(1991)271
 Keim, V., see Fuchs, M. J. 576(1992)287
 Keinänen, T. A., see Hyvönen, T. 574(1992)17
 Keizer, H. J., see Kaijser, G. P. 583(1992)175
 Keller, U., see Küry, D. 572(1991)302
 Keller, U., see Wünsche, L. 552(1991)539
 Kelly, J., see Pleasance, S. 591(1992)325
 Kelly, M. T., see Sagar, K. A. 577(1992)109
 Kelsey, R. G., see Harvey, S. D. 587(1991)300
 Kema, I. P., see Muskiet, F. A. J. 579(1992)184
 Kemper, K., see Oberdorfer, F. 552(1991)483
 Keng, L., see Boison, J. O. 576(1992)315
 Kennler, E. and Schwer, C.
 Peak dispersion and separation efficiency in high-
 performance zone electrophoresis with gel-filled
 capillaries 595(1992)313
 Kenney, M. E., see Zuk, M. M. 568(1991)437
 Keppler, C., see Seitz, J. 587(1991)55
 Kerkhoff, M. J. G., see Coenen, A. J. J. M. 593(1992)243
 Kermici, M., see Krien, P. M. 576(1992)255
 Kermici, M., see Vingler, P. 571(1991)73
 Kerner, J., see Zezza, F. 593(1992)99
 Kersten, B. S., Catalano, T. and Rozenman, Y.
 Ion-pairing high-performance liquid chromatographic
 method for the determination of 5-aminosalicylic acid
 and related impurities in bulk chemical 588(1991)187
 Kerzel, P. and Mersmann, A.
 Adsorption behaviour of lipase from *Staphylococcus*
carnosus on a hydrophobic adsorbent 584(1992)109
 Kestell, P., McKeage, M. J. and Baguley, B. C.
 Determination of xanthenone-4-acetic acid in mouse
 plasma by high-performance liquid chromatography
 564(1991)315
 Ketelaar, P. E. F., see Duchateau, A. L. L. 566(1991)493
 Kettrup, A., see Boos, K.-S. 600(1992)189
 Kever, J. J., see Zimina, T. M. 593(1992)233
 Kežić, S. and Monster, A. C.
 Determination of 2,5-hexanedione in urine and serum by
 gas chromatography after derivatization with O-
 (pentafluorobenzyl)hydroxylamine and solid-phase
 extraction 563(1991)199
 Khachik, F., Englert, G., Daitch, C. E., Beecher, G. R.,
 Tonucci, L. H. and Lusby, W. R.
 Isolation and structural elucidation of the geometrical
 isomers of lutein and zeaxanthin in extracts from human
 plasma 582(1992)153
 Khaledi, M. G., see Smith, S. C. 559(1991)57
 Khaledi, M. G., see Strasters, J. K. 586(1991)221
 Khatter, N., see Ngo, T. T. 597(1992)101
 Khetarpal, V. K., Storbeck, L. S., Wells, D. and Meacham,
 Jr., R. H.
 Method for separation and determination of lactone and
 hydroxy acid forms of a new HMG CoA reductase
 inhibitor (RG 12561) in plasma 567(1991)491
 Khomutov, A. R., see Hyvönen, T. 574(1992)17
 Khomutov, R. M., see Hyvönen, T. 574(1992)17
 Khoo, S. B., see Yik, Y. F. 585(1991)139
 Khoo, S. B., see Yik, Y. F. 589(1992)333
 Khuhawar, M. Y. and Bhatti, A. G.
 Thermoanalytical and chromatographic studies of copper
 (II), nickel(II) and oxovanadium(IV) complexes of
 tetradentate Schiff bases derived from β -diketones and
 2,3-diaminopentane 558(1991)187
 Khuhawar, M. Y., see Sarafraz-Yazdi, A. 594(1992)395
 Khvostova, A. I., see Tjukavkina, N. A. 571(1991)312
 Kia, D., see Arnaud, J. 572(1991)103
 Kiang, C.-H. and Lee, C.
 High-performance liquid chromatographic determination
 of the 1,4-diketo and 1,4-diketo-2,3-dihydroxy
 metabolites of lonapalene in rat urine 565(1991)339
 Kiang, C.-H., Nolan, T., Huang, B. L. and Lee, C. P.-P.
 Determination of femtomole/milliliter concentrations of
 enprostil acid in human plasma using high-performance
 liquid chromatography-laser-induced fluorescence
 detection 567(1991)195
 Kibby, J., see Haynes, P. A. 588(1991)107
 Kibe, T., see Ohba, S. 568(1991)325
 Kidouchi, K., see Ohba, S. 568(1991)325
 Kiefer, C. R., see Kutlar, F. 567(1991)359
 Kiełbasiński, P., see Kudzin, Z. H. 588(1991)307
 Kietzmann, D., see Michaelis, H. C. 571(1991)257

- Kihira, Y. and Aiba, S.
Artificial immunoglobulin G-binding protein mimetic to staphylococcal protein A. Its production and application to affinity purification of immunoglobulin G 597(1992)277
- Kikuchi, K., see Shinkai, H. 570(1991)203
- Kikuta, C., see Mascher, H. 583(1992)122
- Kiltz, H.-H., see Ozyhar, A. 587(1991)11
- Kim, H.-J. and Richardson, M.
Determination of 5-hydroxymethylfurfural by ion-exclusion chromatography with UV detection 593(1992)153
- Kim, H. J., see Rozman, P. 574(1992)146
- Kim, I. S., Sasinos, F. I., Rishi, D. K., Stephens, R. D. and Brown, M. A.
Determination of aromatic sulfonic acids in aqueous environmental samples by anion-exchange chromatography coupled to particle beam mass spectrometry and UV spectrophotometry 589(1991)177
- Kim, M., Saito, K., Furusaki, S., Sato, T., Sugo, T. and Ishigaki, I.
Adsorption and elution of bovine γ -globulin using an affinity membrane containing hydrophobic amino acids as ligands 585(1991)45
- Kim, M., Saito, K., Furusaki, S., Sugo, T. and Ishigaki, I.
Protein adsorption capacity of a porous phenylalanine-containing membrane based on a polyethylene matrix 586(1991)27
- Kim, N. J., see Lee, K.-J. 577(1992)135
- Kim, S.-T., see Strasters, J. K. 586(1991)221
- Kim, T.-J., Park, J.-S. and Shin, H.-S.
Identification of new urinary metabolites of trimeprazine in rats by gas chromatography-mass spectrometry 575(1992)295
- Kimata, K., Hosoya, K., Tanaka, N., Araki, T. and Patterson, Jr., D. G.
Preparation of nitrophenylethylsilylated silica gel and its chromatographic properties in the separation of polychlorinated dibenzo-*p*-dioxins 595(1992)77
- Kimata, K., Hosoya, K., Tanaka, N., Araki, T., Tsuboi, R. and Haginaka, J.
Effect of stationary phase structure on retention and selectivity of restricted-access reversed-phase packing materials 558(1991)19
- Kimata, K., Tanaka, N. and Araki, T.
Suppression of the effect of metal impurities in alkylsilylated silica packing materials 594(1992)87
- Kimura, M., see Hatada, K. 583(1992)116
- Kimura, M., see Ohsawa, K. 597(1992)285
- Kimura, M., see Yamasaki, N. 597(1992)207
- Kimura, S., see Kitagawa, T. 567(1991)267
- Kimura, Y., see Ide, H. 596(1992)203
- Kinahan, I. M. and Smyth, M. R.
High-performance liquid chromatographic determination of PEG 600 in human urine 565(1991)297
- King, C. R., see Maguire, M. H. 575(1992)243
- King, D. N., see Patel, H. B. 555(1991)21
- King, S. J., see Meng, Q. C. 579(1992)63
- King, S.-Y. P., see Kapil, R. P. 577(1992)283
- Kingma, A. W., see Hessels, J. 563(1991)1
- Kinoshita, T., see Nimura, N. 585(1991)207
- Kinoshita, T., see Yokoyama, T. 568(1991)212
- Kinosita, I., see Sano, K. 578(1992)63
- Kinouchi, Y., see Iida, Y. 571(1991)277
- Kint, A. H., see Meyer, E. 570(1991)149
- Kinter, M., Sullivan, S., Roberts, R. J. and Spitz, D.
Trace quantitation of 4-hydroxy-2-nonenal in biological samples as its oxime-bis-*tert*-butyldimethylsilyl derivative using 3-hydroxynonanal as an internal standard 578(1992)9
- Kinter, M., see Aggarwal, S. K. 576(1992)297
- Kintz, P.
Gas chromatographic analysis of nicotine and cotinine in hair 580(1992)347
- Kintz, P., Tracqui, A. and Mangin, P.
Determination of nalbuphine using high-performance liquid chromatography coupled to photodiode-array detection and gas chromatography coupled to mass spectrometry 579(1992)172
- Kinuta, M., see Yao, K. 581(1992)11
- Kira, R., see Ôi, N. 592(1992)291
- Kirchberger, J., Domar, U., Kopperschläger, G. and Stigbrand, T.
Interactions of human alkaline phosphatase isoenzymes with triazine dyes using affinity partitioning, affinity chromatography and difference spectroscopy 574(1992)237
- Kirchberger, J., Kopperschläger, G. and Vijayalakshmi, M. A.
Dye-ligand affinity partitioning of lactate dehydrogenase isoenzymes 557(1991)325
- Kirchessner, D. A., see Lindgren, E. R. 585(1991)353
- Kirchessner, M., see Eder, K. 588(1991)265
- Kirchessner, M., see Eder, K. 598(1992)33
- Kirchner, M., see Gaede, F.-P. 585(1991)53
- Kirkland, J. J., Dilks, Jr., C. H., Rementer, S. W. and Yau, W. W.
Asymmetric-channel flow field-flow fractionation with exponential force-field programming 593(1992)339
- Kirkland, J. J., see Yau, W. W. 556(1991)111
- Kishino, S., Di, Z. S., Sugawara, M., Iseki, K., Miyazaki, K., Kakinoki, S., Nomura, A. and Kitabatake, A.
Rapid and simple method for the determination of α_1 -acid glycoprotein in serum by column liquid chromatography 582(1992)246
- Kiso, Y., see Hirokawa, T. 555(1991)247
- Kiso, Y., see Hirokawa, T. 555(1991)321
- Kiso, Y., see Hirokawa, T. 585(1991)297
- Kiso, Y., see Hirokawa, T. 585(1991)364
- Kiso, Y., see Hu, J.-Y. 589(1992)339
- Kiso, Y., see Hu, J.-Y. 594(1992)371
- Kiss, F., see Macka, M. 586(1991)291
- Kissinger, P. T.
Electroanalysis of biologically important compounds (by J. P. Hart) (Book Review) 568(1991)270
- Kissinger, P.T., see Linhares, M.C. 578(1992)157
- Kitabatake, A., see Kishino, S. 582(1992)246
- Kitagawa, T., see Okamoto, J. 583(1992)45
- Kitagawa, T., see Okamoto, J. 583(1992)53

- Kitagawa, T., Tsuboi, I., Kimura, S. and Sasamoto, Y.
Rapid method for preparing a β -glucan-specific sensitive fraction from *Limulus (Tachypleus tridentatus)* amoebocyte lysate 567(1991)267
- Kitahara, H., see Ōi, N. 592(1992)291
- Kitahara, S., see Kawai, S. 585(1991)318
- Kitamura, F., see Ueda, T. 592(1992)229
- Kitamura, F., see Ueda, T. 593(1992)265
- Kitaoka, Y., Kobayashi, M. and Ujeno, Y.
Studies of the electrophoretic behaviour of *p*-dihydroxyborylphenylalanine and related compounds by the three-spot method 585(1991)369
- Kitsos, M., Gandini, C., Massolini, G., De Lorenzi, E. and Caccialanza, G.
High-performance liquid chromatography post-column derivatization with fluorescence detection to study the influence of ambroxol on dipalmitoylphosphatidylcholine levels in rabbit eustachian tube washings 553(1991)1
- Kitsos, M., see Baruffini, A. 593(1992)95
- Kitts, D. D., see Aoyama, R. G. 588(1991)181
- Klaassen, C. D., see Rozman, P. 574(1992)146
- Klein, J. T., see Hsu, R. S. 572(1991)352
- Klein, R. F. X., see Lurie, I. S. 598(1992)59
- Kleine, A. H., see Van Nieuwenhoven, F. A. 570(1991)173
- Klein Elhorst, J. T., see Essink, A. W. G. 570(1991)419
- Kleinjans, J. C. S., see De Kok, T. M. C. M. 580(1992)135
- Kleinmann, I., see Šmídl, P. 598(1992)15
- Klepárník, K. and Boček, P.
Theoretical background for clinical and biomedical applications of electromigration techniques (Review) 569(1991)3
- Klette, K. L., see Kuhlman, Jr., J. J. 578(1992)207
- Klimeš, J., Klimešová, V. and Waissner, K.
Chromatographic behaviour of dipyritylsulphides. Relationship between log *k'* values and structure by reversed-phase high-performance liquid chromatography 595(1992)334
- Klimeš, J., Sochor, J., Zahradníček, M. and Sedláček, J.
Simultaneous high-performance liquid chromatographic determination of salicylates in whole blood, plasma and isolated erythrocytes 584(1992)221
- Klimešová, V., see Klimeš, J. 595(1992)334
- Kline, W. F. and Matuszewski, B. K.
Improved determination of the bisphosphonate alendronate in human plasma and urine by automated precolumn derivatization and high-performance liquid chromatography with fluorescence and electrochemical detection 583(1992)183
- Kline, W. F., Matuszewski, B. K. and Hsieh, J. Y.-K.
Determination of a cyclic heptapeptide, a novel fibrinogen receptor antagonist, in human plasma by high-performance liquid chromatography with automated pre-column derivatization, column switching and fluorescence detection 578(1992)31
- Klöppel, K.-D. and Fredrickson, H. L.
Fast atom bombardment mass spectrometry as a rapid means of screening mixtures of ether-linked polar lipids from extremely halophilic archaeobacteria for the presence of novel chemical structures. 562(1991)369
- Klunk, L. J., see Russell, J. W. 572(1991)321
- Klus, H., see Pachinger, A. 558(1991)369
- Klyashchitsky, B. A., Mitina, V. K., Frantsuzova, N. A., Krasnopolsky, Y. M. and Shvets, V. I.
Use of biospecific interactions of collagen, fibronectin and their fragments in affinity chromatography 577(1992)267
- Knadler, M. P.
Quantification of a decapeptide anticoagulant in rat and monkey plasma by high-performance liquid chromatography 579(1992)371
- Knauz, D., see Morvai, M. 552(1991)337
- Kneer, J., see Rissler, K. 565(1991)375
- Knobloch, E., Mandys, F., Hodr, R., Hujer, R. and Mader, R.
Study of the mechanism of the photoisomerization and photooxidation of bilirubin using a model for the phototherapy of hyperbilirubinemia 566(1991)89
- Knupp, C. A., see Papp, E. A. 574(1992)93
- Kobata, A. and Endo, T.
Immobilized lectin columns: useful tools for the fractionation and structural analysis of oligosaccharides (Review) 597(1992)111
- Kobata, S., see Ide, H. 596(1992)203
- Kobayashi, K., Chiba, K., Sohn, D.-R., Kato, Y. and Ishizaki, T.
Simultaneous determination of omeprazole and its metabolites in plasma and urine by reversed-phase high-performance liquid chromatography with an alkaline-resistant polymer-coated C_{18} column 579(1992)299
- Kobayashi, M., see Kitaoka, Y. 585(1991)369
- Kobayashi, M., see Ohba, S. 568(1991)325
- Kobold, U., see Puhlmann, A. 581(1992)129
- Koda, S., see Honda, S. 590(1992)364
- Kodama, H., see Sugahara, K. 565(1991)408
- Kodama, H., see Sugahara, K. 579(1992)318
- Kodama, H., see Watanabe, H. 568(1991)445
- Kodo, N., see Millington, D. S. 562(1991)47
- Koekkoek-van den Herik, I., see Van de Heijning, B. J. M. 565(1991)159
- Koelsch, R., see Baeseler, M. 589(1992)93
- Koffel, J. C., see Lave, T. 572(1991)203
- Kohata, K., Watanabe, M. and Yamanaka, K.
Highly sensitive determination of photosynthetic pigments in marine *in situ* samples by high-performance liquid chromatography 558(1991)131
- Kohno, T. and Kuwata, K.
Preconcentration technique for introducing gaseous or volatile compounds into a capillary gas chromatographic column 587(1991)338
- Kohriyama, K., see Okuda, T. 567(1991)141
- Koizumi, K., Fukuda, M. and Hizukuri, S.
Estimation of the distributions of chain length of amylopectins by high-performance liquid chromatography with pulsed amperometric detection 585(1991)233
- Koizumi, K., Kubota, Y., Ozaki, H., Shigenobu, K., Fukuda, M. and Tanimoto, T.
Analyses of isomeric mono-O-methyl-D-glucoses, D-glucobioses and D-glucose monophosphates by high-performance anion-exchange chromatography with pulsed amperometric detection 595(1992)340

- Koizumi, Y., see Yasukawa, K. 597(1992)271
- Kojima, K., Ogawa, H. K., Seno, N. and Matsumoto, I.
Affinity purification and affinity characterization of carbohydrate-binding proteins in bovine kidney 597(1992)323
- Kojima, K., see Ueno, M. 597(1992)197
- Kok, W. T., see Debets, A. J. J. 600(1992)163
- Kokkonen, P. S., Niessen, W. M. A., Tjaden, U. R. and Van der Greef, J.
Bioanalysis of erythromycin 2'-ethylsuccinate in plasma using phase-system switching continuous-flow fast atom bombardment liquid chromatography-mass spectrometry 565(1991)265
- Kolesnik, J. A., see Geodakyan, S. V. 577(1992)371
- Kolesnik, J. A., see Tjukavkina, N. A. 571(1991)312
- Koliadima, A., Karaiskakis, G., Katsanos, N. A. and Roth, M.
Activity coefficients in binary liquid mixtures measured by reversed-flow gas chromatography 595(1992)237
- Kolker, H. J., see Van 't Klooster, G. A. E. 571(1991)157
- Kolker, H. J., see Van 't Klooster, G. A. E. 579(1992)158
- Kolker, H. J., see Van 't Klooster, G. A. E. 579(1992)355
- Kollie, T. O. and Poole, C. F.
Influence of solute size and the non-polar interaction term on the selection of test solutes for the classification of stationary phase selectivity in gas chromatography 556(1991)457
- Kollmannsberger, H., see Nitz, S. 557(1991)187
- Kolomiets, L. N., see Chalykh, A. E. 592(1992)121
- Kolycheva, M. T., see Galushko, S. V. 600(1992)83
- Komatsu, S., Murata, S., Aoyama, H., Zenki, T., Ozawa, N., Tateishi, M. and Vrbanc, J. J.
Micro-quantitative determination of ciprofloxacin in plasma by gas chromatography-mass spectrometry coupled with an antibody extraction 568(1991)460
- Kominami, G., see Murai, Y. 597(1992)201
- Komine, Y., see Shimada, K. 565(1991)111
- Komulainen, H., see Pennanen, S. 568(1991)125
- Konami, Y., Yamamoto, K. and Osawa, T.
Purification and characterization of carbohydrate-binding peptides from *Lotus tetragonolobus* and *Ulex europaeus* seed lectins using affinity chromatography 597(1992)213
- Konami, Y., see Yamamoto, K. 597(1992)221
- Kondo, A., see Ohara, K. 586(1991)35
- Kondo, J., see Nakanishi, A. 591(1992)159
- Kondo, M., see Shimohigashi, Y. 597(1992)425
- Kondoh, Y., see Ohmori, S. 566(1991)1
- Kong Sing, Y. L., Algiman, E., Kroviarski, Y., Massot, C., Dhermy, D. and Bertrand, O.
Modification of reversed-phase columns with dyed surfactants. Preparation of mechanically resistant efficient immobilized dyes for protein purification 558(1991)43
- Kong Sing, Y. L., Kroviarski, Y., Cochet, S., Dhermy, D. and Bertrand, O.
High-performance hydrophobic interaction chromatography of proteins on reversed-phase supports coated with non-ionic surfactants of polyoxyethylene type. Purification of a fungal aspartic proteinase 598(1992)181
- Konishi, M., see Murai, Y. 597(1992)201
- Kononenko, V. L. and Shimkus, J. K.
Non-equilibrium integral Doppler anemometric analysis of particle mixtures in a channel flow using an intrinsic hydrodynamic focusing force biased by another force 553(1991)517
- Kononenko, V. L. and Shimkus, J. K.
Use of integral Doppler anemometry in field-flow fractionation 600(1992)139
- Konteatis, Z. D., see Florance, J. R. 559(1991)391
- Koole, L. H., see Coenen, A. J. J. M. 596(1992)59
- Koopman-Kimenai, P. M., see Verwey-van Wissen, C. P. W. G. M. 570(1991)309
- Kootstra, P. R., see Somsen, G. W. 552(1991)635
- Köppel, C. and Tenczer, J.
Mass spectral characterization of urinary metabolites of D,L-kavain 562(1991)207
- Köppel, C., Tenczer, J. and Peixoto-Menezes, K. M.
Formation of formaldehyde adducts from various drugs by use of methanol in a toxicological screening procedure with gas chromatography-mass spectrometry 563(1991)73
- Köppel, C. and Wagemann, A.
Plasma level monitoring of D,L-verapamil and three of its metabolites by reversed-phase high-performance liquid chromatography 570(1991)229
- Köppel, C., Wagemann, A., Hansen, G. R. and Müller, C.
Monitoring of ajmaline in plasma with high-performance liquid chromatography 575(1992)87
- Kopperschläger, G., see Kirchberger, J. 557(1991)325
- Kopperschläger, G., see Kirchberger, J. 574(1992)237
- Körber, F., see Abou-Rebyeh, H. 566(1991)341
- Körber, F., see Abou-Rebyeh, H. 566(1991)351
- Kornhauser, D. M., see Jajoo, H. K. 577(1992)299
- Korori, S. A. A., see Pichorner, H. 594(1992)400
- Korsrud, G. O., see Boisson, J. O. 576(1992)315
- Koruna, I., see Ryska, M. 562(1991)289
- Korver, M. P., see Burse, V. W. 566(1991)117
- Koscielny, M. L., see Levison, P. R. 590(1992)49
- Kosh, J. W., see Terry, Jr., A. V. 585(1991)101
- Koshiishi, I., see Imanari, T. 574(1992)142
- Kostadinova, S., Ivanov, A. and Kamberov, E.
Purification and some properties of phospholipase C from *Achromobacter xylosoxidans* 568(1991)315
- Kostenko, V. G., see Nemirovskii, V. D. 594(1992)23
- Kostiainen, R.
Identification of trichothecenes by thermospray, plasmaspay and dynamic fast-atom bombardment liquid chromatography-mass spectrometry 562(1991)555
- Kostiuk, P., see Böswart, J. 571(1991)19
- Kotagal, U. R., see Tan, H. S. I. 568(1991)145
- Kotaki, H., see Yamamura, Y. 567(1991)151
- Kotrla, R., see Churáček, J. 557(1991)523
- Köttgen, E., see Righetti, P. G. 569(1991)197
- Kotyński, A., see Kudzin, Z. H. 588(1991)307
- Koulu, M., see Huupponen, R. 583(1992)270
- Kouno, Y., see Kanai, M. 567(1991)415
- Koupai-Abyazani, M. R., McCallum, J. and Bohm, B. A.
Identification of the constituent flavanoid units in sainfoin proanthocyanidins by reversed-phase high-performance liquid chromatography 594(1992)117

- Koupai-Abyazani, M. R., see Creaser, C. S. 586(1991)323
- Kouwatli, H., see Tod, M. 594(1992)386
- Kovach, P. M., Lantz, R. J. and Brier, G.
High-performance liquid chromatographic determination of loracarbef, a potential metabolite, cefaclor and cephalixin in human plasma, serum and urine 567(1991)129
- Kovács-Hadady, K.
Study of the retention behaviour of barbiturates by overpressured layer chromatography using silica gel bonded with tricaprylmethylammonium chloride 589(1992)301
- Kovács-Hadady, K. and Szilágyi, J.
Separation of minoxidil and its intermediates by overpressured layer chromatography using a stationary phase bonded with tricaprylmethylammonium chloride 553(1991)459
- Koyama, M., see Ohsawa, K. 597(1992)285
- Koyama, T., see Isosaki, K. 597(1992)123
- Koza, T., see Wurst, M. 593(1992)201
- Kozuka, M., see Hagiwara, H. 597(1992)331
- Kraak, J. C., see Bruin, G. J. M. 559(1991)163
- Kraft, R., see Lohmann, A. 564(1991)283
- Krähenbühl, P., see Brügger, R. 557(1991)163
- Krahenbühl, S., Minkler, P. E. and Hoppel, C. L.
Derivatization of isolated endogenous butyrobetaine with 4'-bromophenacyl trifluoromethanesulfonate followed by high-performance liquid chromatography 573(1992)3
- Králová, B., see Břizová, K. 593(1992)125
- Krämer, B. K., see Mayer, F. 572(1991)339
- Kramer, P. A., see Weaver, M. L. 581(1992)293
- Krasnopolsky, Y. M., see Klyashchitsky, B. A. 577(1992)267
- Kraus, E., see Kuchař, M. 557(1991)399
- Kraus, G., Schulz, H.-U. and Lohmann, A.
Determination of apovincaminic acid in serum by means of high-performance liquid chromatography 573(1992)323
- Kraus, L., see Lendrath, G. 588(1991)303
- Krause, M. and Galensa, R.
High-performance liquid chromatography of diastereomeric flavanone glycosides in *Citrus* on a β -cyclodextrin-bonded stationary phase (Cyclobond I) 588(1991)41
- Krause, W. and Kühne, G.
High-performance liquid chromatographic procedure for specificity testing of radioimmunoassays: rolipram 573(1992)303
- Krauss, G. J., see Friebe, S. 598(1992)139
- Kreimer-Birnbaum, M., see Zuk, M. M. 568(1991)437
- Kresin, L., see Cunico, R. L. 559(1991)467
- Kresin, L., see Kunitani, M. 588(1991)125
- Krien, P., see Vingler, P. 571(1991)73
- Krien, P. M., Margou, V. and Kermici, M.
Electrochemical determination of femtomole amounts of free reduced and oxidized glutathione. Application to human hair follicles 576(1992)255
- Krishna, D. R., see Venkataramana, D. 575(1992)167
- Kristjánsson, F.
Sensitive determination of buspirone in serum by solid-phase extraction and two-dimensional high-performance liquid chromatography 566(1991)250
- Križanovič, K., see Petro, M. 595(1992)337
- Kroboth, P. D., see Schmith, V. D. 568(1991)253
- Krogh, M., see Andresen, A. T. 582(1992)123
- Kroner, K. H., see Langlotz, P. 591(1992)107
- Kroviarski, Y., see Kong Sing, Y. L. 558(1991)43
- Kroviarski, Y., see Kong Sing, Y. L. 598(1992)181
- Krstulović, A. M.
Chromatography and modification of nucleosides, Part A (edited by C. W. Gehrke and K. C. T. Kuo) (Book Review) 563(1991)228
- Krstulović, A. M. and Dandeu, J.-P.
Preface 584(1992)1
- Krstulovic, A. M., Tagliaro, F. and Marigo, M.
Foreword 566(1991)273
- Krstulović, A. M., see Aubert, M.-C. 557(1991)47
- Krstulović, A.M., see Lesellier, E. 557(1991)59
- Krüger, H., see Pasch, H. 589(1992)295
- Krull, I. R.
Advances in chromatography, Vol. 32 (edited by J. C. Giddings, E. Grushka and P. R. Brown) (Book Review) 600(1992)376
- Krull, I. S., see Mhatre, R. M. 591(1992)139
- Kubo, M., Sasabe, H. and Shimizu, T.
Highly sensitive method for the determination of 5-fluorouracil in biological samples in the presence of 2'-deoxy-5-fluorouridine by gas chromatography-mass spectrometry 564(1991)137
- Kubota, Y., see Koizumi, K. 595(1992)340
- Kucera, P. J., see Chen, S. 558(1991)141
- Kuchař, M., Kraus, E. and Jelínková, M.
Influence of mobile phase composition on evaluation of lipophilicity by partition chromatography 557(1991)399
- Kucharczyk, N., see Clark, L. A. 573(1992)113
- Kudo, K., see Kage, S. 564(1991)163
- Kudo, M., see Ohkubo, T. 573(1992)289
- Kudzin, Z. H., Kotyński, A. and Kiełbasiński, P.
Application of the iodine-azide reagent for selective detection of thiophosphoryl compounds in thin-layer chromatography 588(1991)307
- Kuettner, K. E., see Kang, K. 567(1991)57
- Kugabalasooriar, J., see Bushway, R. J. 587(1991)321
- Kuhara, T., see Matsumoto, M. 562(1991)139
- Kühlkamp, V., see Mayer, F. 572(1991)339
- Kuhlman, Jr., J. J., Levine, B., Klette, K. L., Maglulio, Jr., J., Kalasinsky, K. S. and Smith, M. L.
Measurement of azacyclonol in urine and serum of humans following terfenadine (Seldane) administration using gas chromatography-mass spectrometry 578(1992)207
- Kühne, G., see Krause, W. 573(1992)303
- Kuijpers, A., see Van Tellingen, O. 573(1992)328
- Kukhar, V. P., see Galushko, S. V. 553(1991)143
- Kukhar, V. P., see Galushko, S. V. 600(1992)79
- Kuksis, A., Marai, L. and Myher, J. J.
Reversed-phase liquid chromatography-mass spectrometry of complex mixtures of natural triacylglycerols with chloride-attachment negative chemical ionization 588(1991)73

- Kuksis, A., Myher, J. J., Geher, K., Breckenridge, W. C., Feather, T., McGuire, V. and Little, J. A.
Gas chromatographic profiles of plasma total lipids as indicators of dietary history. Correlation with fat intake based on 24-h dietary recall 564(1991)11
- Kuksis, A., Myher, J. J., Geher, K., Breckenridge, W. C., Feather, T., McGuire, V. and Little, J. A.
Gas chromatographic profiles of plasma total lipids as indicators of dietary history. Correlation with carbohydrate and alcohol intake based on 24-h dietary recall 579(1992)13
- Kula, M.-R., see Woker, R. 584(1992)85
- Kulemeier, J., see Stein, J. 576(1992)53
- Kulicki, Z., see Baj, S. 588(1991)33
- Kulik, W., see Van Rhijn, J. A. 595(1992)289
- Kumashiro, S., see Watanabe, A. 583(1992)241
- Kumps, A., see Mardens, Y. 577(1992)341
- Kunath, A., Höft, E., Hamann, H.-J. and Wagner, J.
Enantiomeric separation of racemic hydroperoxides and related alcohols 588(1991)352
- Kuningas, K. R., see Berezkin, V. G. 552(1991)313
- Kunitani, M., Dollinger, G., Johnson, D. and Kresin, L.
On-line characterization of polyethylene glycol-modified proteins 588(1991)125
- Kunitani, M., see Dollinger, G. 592(1992)215
- Kunkel, M. E., Seo, A. and Thomas, R. L.
Determination of profiles of non-collagenous proteins from rat bones by sodium dodecyl sulfate high-performance liquid chromatography 589(1992)135
- Kunsmann, C. M., see Kunsmann, G. W. 568(1991)427
- Kunsmann, G. W., Manno, J. E., Przekop, M. A., Manno, B. R., Llorens, K. A. and Kunsmann, C. M.
Determination of temazepam and temazepam glucuronide by reversed-phase high-performance liquid chromatography 568(1991)427
- Künstner, P., see Chollet, D. 577(1992)335
- Kuo, J.-S., see Cheng, F.-C. 582(1992)19
- Kuo, K., see Mitchell, E. P. 581(1992)31
- Kuriyama, M., see Fujioka, T. 597(1992)429
- Kuroda, N., see Nakashima, K. 584(1992)275
- Kurono, M., Yoshida, K. and Naruto, S.
Determination of AJ-3941, a possible agent for the treatment of cerebrovascular disorders, in plasma and brain by means of high-performance liquid chromatography with fluorescence detection 578(1992)152
- Kurosawa-Ohsawa, K., see Ohsawa, K. 597(1992)285
- Kurumaji, A., see Watanabe, A. 583(1992)241
- Küry, D. and Keller, U.
Trimethylsilyl-O-methyloxime derivatives for the measurement of [6,6-²H₂]-D-glucose-enriched plasma samples by gas chromatography-mass spectrometry 572(1991)302
- Kusaka, T., see Ikeda, M. 575(1992)197
- Kusmierz, J. J. and Desiderio, D. M.
Characterization of an aminopeptidase in cerebrospinal fluid. Structure elucidation of enzyme hydrolysis products of synthetic methionine-enkephalin by reversed-phase high-performance liquid chromatography and mass spectrometry 574(1992)189
- Kusmierz, J. J., see Lovelace, J. L. 562(1991)573
- Küstners, E., Spöndlin, C. and Eder, C.
Analytical studies on the chiral separation and simultaneous determination of pantothenic acid and hopantenic acid enantiomers in rat plasma by gas chromatography-mass fragmentography: a reply 576(1992)179
- Kutlar, F., Moscoso, H., Kiefer, C. R., Garver, F. A., Beksaç, S., Önderoglu, L., Gurgey, A., Altay, Ç. and Huisman, T. H. J.
Quantities of adult, fetal and embryonic globin chains in the blood of eighteen- to twenty-week-old human fetuses 567(1991)359
- Kutnink, M. A., see Burri, B. J. 567(1991)369
- Kutnink, M. A., see Hawkes, W. C. 576(1992)263
- Kuwae, A., see Kawai, S. 585(1991)318
- Kuwana, T., see Ueda, T. 593(1992)265
- Kuwata, K., see Kohno, T. 587(1991)338
- Kuykendall, J. R. and Cox, R.
Rapid and quantitative separation of nicotinamide and its N¹-methylated metabolite by Dowex AG50-X4 chromatography 585(1991)322
- Kuypers, F. A., Bütikofer, P. and Shackleton, C. H. L.
Application of liquid chromatography-thermospray mass spectrometry in the analysis of glycerophospholipid molecular species 562(1991)191
- Kuznetsova, T. A., see Voinov, V. G. 586(1991)360
- Kvalo, L. T., Wilhelm, J. A., Edfort, M. J. and Venturella, V. S.
Determination of [³H]briofentanil, a potent narcotic analgesic, from rat serum by high-performance liquid chromatography with on-line radioactive detection 565(1991)391
- Květina, J., see Nobilis, M. 581(1992)251
- Kwakkenbos, G. T. C., see Duchateau, A. L. L. 552(1991)605
- Kwakman, P. J. M., Kamminga, D. A., Brinkman, U. A. T. and De Jong, G. J.
Sensitive liquid chromatographic determination of alkyl-, nitro- and chlorophenols by precolumn derivatization with dansyl chloride, postcolumn photolysis and peroxyoxalate chemiluminescence detection 553(1991)345
- Kymer, K. A., see De Vries, J. X. 562(1991)31
- Kyoyama, K., see Okuda, T. 567(1991)141
- Kysilka, R., see Wurst, M. 593(1992)201

L

- Labdon, J. E., see Pramanik, B. N. 562(1991)377
- LaCreta, F. P., Brennan, J. M., Tinsley, P. W. and O'Dwyer, P. J.
High-performance liquid chromatographic determination of ethacrynic acid in human plasma 571(1991)271
- LaCreta, F. P., see Tinsley, P. W. 564(1991)303
- La Croix, R., see Pianezzola, E. 574(1992)170
- Ladisch, C. M., see Yang, Y. 598(1992)169
- Ladisch, M. R., see Yang, Y. 598(1992)169
- Lafolie, P., see Beck, O. 570(1991)198
- Lafont, O., see Vérité, P. 578(1992)134
- Lafosse, M., see Brossard, S. 591(1992)149
- Laganà, A. and Marino, A.
General and selective isolation procedure for high-performance liquid chromatographic determination of anabolic steroids in tissues 588(1991)89
- Lagendijk, J., Ubbink, J. B. and Vermaak, W. J. H.
Quantification of erythrocyte S-adenosyl-L-methionine levels and its application in enzyme studies 576(1992)95
- Lagerström, P.-O., see Balmér, K. 553(1991)391
- Lagerström, P.-O., see Balmér, K. 592(1992)331
- Lagerwerf, A. J., Vanlinthout, L. E. H. and Vree, T. B.
Rapid determination of succinylcholine in human plasma by high-performance liquid chromatography with fluorescence detection 570(1991)390
- Lagomarsino, R. J. and Latner, N.
Wide dynamic range electron-capture detection using the electron-capture detector pulse frequency signal 595(1992)359
- La Grone, J., see Miller, R. 578(1992)103
- Lagu, A. L., see Strege, M. A. 555(1991)109
- Lahiri, P., see Chakraborty, M. 571(1991)235
- Lalonde, R. L., see Herring, V. L. 567(1991)221
- Lam, G. N., see Kapil, R. P. 577(1992)283
- Lamaison, S., see Dorbon, M. 557(1991)155
- Lamb, J. H., see Kajbaf, M. 583(1992)63
- Lambert, W. E., see Meyer, E. 570(1991)149
- Lambertsen, G., see Lie, Ø. 565(1991)119
- Lambiase, S. J., see Betz, W. R. 556(1991)433
- Lambrey, B., see Damaj, M. I. 563(1991)476
- Lammert, S. A., see Kaiser, Jr., R. E. 562(1991)3
- La Morgia, R., see Santoni, G. 581(1992)287
- Lanbeck-Vallén, K. and Carlqvist, J.
Determination of cefcanel in plasma and urine by high-performance liquid chromatography using coupled columns, after administration of the new cephalosporin prodrug cefcanel daloxate hydrochloride 578(1992)71
- Lanbeck-Vallén, K., Carlqvist, J. and Nordgren, T.
Determination of ampicillin in biological fluids by coupled-column liquid chromatography and post-column derivatization 567(1991)121
- Lane, L., see Levison, P. R. 590(1992)49
- Lang, J., see Grit, M. 585(1991)239
- Lang, V.
Polychlorinated biphenyls in the environment (Review) 595(1992)1
- Lange, C., see Cherton, J.-C. 566(1991)499
- Lange, C., see Cherton, J.-C. 566(1991)511
- Langenfeld, J. J., Burford, M. D., Hawthorne, S. B. and Miller, D. J.
Effects of collection solvent parameters and extraction cell geometry on supercritical fluid extraction efficiencies 594(1992)297
- Langer, S. H., see Jeng, C.-Y. 556(1991)383
- Langer, S. H., see Jeng, C.-Y. 589(1992)1
- Langguth, P., see Spahn-Langguth, H. 584(1992)229
- Langley, G. J., see Preece, N. E. 573(1992)227
- Langlois, R., see Berger, M. 593(1992)133
- Langlois, S., see Wood, S. 569(1991)421
- Langlotz, P. and Kroner, K. H.
Surface-modified membranes as a matrix for protein purification 591(1992)107
- Langvardt, P. W. and Nolan, R. J.
Determination of styrene-7,8-oxide in whole rat blood by gas chromatography-mass spectrometry 567(1991)93
- Lanin, S. N. and Nikitin, Y. S.
Retention data for five ketotrichothecenes in reversed-phase high-performance liquid chromatography with different eluent systems 558(1991)81
- Lannoo, P., see Cachet, T. 600(1992)99
- Lansens, P., Casais Laiño, C., Meuleman, C. and Baeyens, W.
Evaluation of gas chromatographic columns for the determination of methylmercury in aqueous head space extracts from biological samples 586(1991)329
- Lantz, R. J., see Kovach, P. M. 567(1991)129
- Lapicque, F., see Payan, E. 566(1991)9
- Lapinjoki, S. P., see Auriola, S. 554(1991)227
- Lapinjoki, S. P., see Auriola, S. O. K. 594(1992)153
- Larionov, O. G., Petrenko, V. V. and Platonova, N. P.
Determination of sorbent polarity and selectivity by linear regression of retention indices 552(1991)31
- Larionov, O. G., see Chalykh, A. E. 592(1992)121
- La Rosa, C., see Lucarelli, C. 573(1992)150
- Larre, C., see Courcoux, P. 596(1992)225
- Larsson, H., see Bergqvist, Y. 571(1991)169
- Larsson, L., see Elmroth, I. 598(1992)43
- Larsson, L., see Magnusson, P. 576(1992)79
- Larsson, M., see Borén, K. 588(1991)139
- Lartigue-Mattei, C., see Andriollo, O. 575(1992)301
- Lasch, J., see Baeseler, M. 589(1992)93
- Last, T., see Chen, T. K. 596(1992)123
- Latner, N., see Lagomarsino, R. J. 595(1992)359
- Lattes, A., see Halvorsen, K. 557(1991)501
- Latxague, L., Gardrat, C., Coustille, J. L., Viaud, M. C. and Rollin, P.
Identification of enzymatic degradation products from synthesized glucobrassicin by gas chromatography-mass spectrometry 586(1991)166

- Lau, C. E.
Determination of cocaethylene, cocaine and their metabolites in rat serum microsomes by high-performance liquid chromatography, and its application to pharmacokinetic studies in rodents 582(1992)167
- Lau, D. T. W., see Han, C. 579(1992)237
- Lau, O. W., see Chan, K. 565(1991)247
- Lau-Cam, C. A., see Roos, R. W. 555(1991)278
- Laufen, H., see Riedel, K.-D. 576(1992)358
- Lavagnini, I., see Pastore, P. 591(1992)219
- Lavarenne, J., see Coudore, F. 584(1992)249
- Lave, T., Efthymiopoulos, C., Koffel, J. C. and Jung, L.
Determination of tertatolol enantiomers in biological fluids by high-performance liquid chromatography 572(1991)203
- Lavicky, J., see Dempsey, C. A. 596(1992)110
- Law, B. and Chan, P. F.
Possible role for stationary phase metal interactions in the chromatography of hydroxyamines on silica 552(1991)429
- Law, M. P., see Jones, H. A. 570(1991)361
- Lázaro, F., see Linares, P. 558(1991)147
- Lazure, C., see Basak, A. 581(1992)17
- Lazzara, R., see Chen, C. L. 583(1992)274
- Leader, H., see Brown, N. D. 563(1991)466
- Leaver, G., Howell, J. A. and Conder, J. R.
Adsorption kinetics of albumin on a cross-linked cellulose chromatographic ion exchanger 590(1992)101
- Leavitt, R. K., see Matassa, L. C. 573(1992)43
- Lebeau, B. and Hammers, W. E.
Estimation of tobacco blend compositions using closed-loop stripping analysis and stepwise multiple linear regression and partial least-squares techniques 596(1992)285
- Lebedin, Y. S., see Gorchakov, V. D. 563(1991)166
- Lebl, M., Fang, S. and Hruby, V. J.
High-performance liquid chromatography of peptides at reduced temperatures: separation of isomers 586(1991)145
- Le Blanc, J. C. Y., see Siu, K. W. M. 554(1991)27
- LeBoeuf, E., see Zimmerman, C. L. 563(1991)83
- Lebon, M., see Sellier, N. 557(1991)451
- Lebrón-Aguilar, R., see Quintanilla-López, J. E. 591(1992)303
- Lecaillon, J. B., see Rouan, M. C. 573(1992)59
- Lecaillon, J. B., see Rouan, M. C. 573(1992)65
- Lecaillon, J. B., see Rouan, M. C. 577(1992)387
- Le Clanche, V., see Rouan, M. C. 573(1992)65
- Leclercq, P. A., see Houben, R. J. 554(1991)351
- Leclercq, P. A., see Pacáková, V. 555(1991)229
- Lecompte, D., see Desage, M. 571(1991)250
- Le Corre, P., see Le Verge, R. 574(1992)283
- Lecron, J. C., see Dufay, C. 563(1991)224
- Ledda, S., see Manca, L. 563(1991)158
- Le Duigou, F., see Rouan, M. C. 573(1992)59
- Lee, B. L., Chua, S. C., Ong, H. Y. and Ong, C. N.
High-performance liquid chromatographic method for routine determination of vitamins A and E and β -carotene in plasma 581(1992)41
- Lee, B. L., Ong, H. Y. and Ong, C. N.
Determination of thiamine and its phosphate esters by gradient-elution high-performance liquid chromatography 567(1991)71
- Lee, C., see Kiang, C.-H. 565(1991)339
- Lee, C. P.-P., see Kiang, C.-H. 567(1991)195
- Lee, C. R., see Aubert, M.-C. 557(1991)47
- Lee, C. R., see Lesellier, E. 557(1991)59
- Lee, C. S., Wu, C.-T., Lopes, T. and Patel, B.
Analysis of separation efficiency in capillary electrophoresis with direct control of electroosmosis by using an external electric field 559(1991)133
- Lee, H.-B. and Peart, T. E.
Supercritical carbon dioxide extraction of resin and fatty acids from sediments at pulp mill sites 594(1992)309
- Lee, H. K., see Ng, C. L. 598(1992)133
- Lee, H. K., see Ong, C. P. 559(1991)529
- Lee, H. K., see Ong, C. P. 559(1991)537
- Lee, H. K., see Ong, C. P. 588(1991)335
- Lee, H. K., see Yeo, S. K. 585(1991)133
- Lee, H. K., see Yeo, S. K. 594(1992)335
- Lee, H. K., see Yik, Y. F. 585(1991)139
- Lee, H. K., see Yik, Y. F. 589(1992)333
- Lee, H. S., Shin, H. C., Han, S. S. and Roh, J. K.
High-performance liquid chromatographic determination of rifapentine in serum using column switching 574(1992)175
- Lee, J. E., see Park, J. H. 586(1991)1
- Lee, K. C. and DeLuca, P. P.
Simultaneous determination of prostaglandins E₁, A₁ and B₁ by reversed-phase high-performance liquid chromatography for the kinetic studies of prostaglandin E₁ in solution 555(1991)73
- Lee, K.-J. and Heo, G. S.
Free solution capillary electrophoresis of proteins using untreated fused-silica capillaries 559(1991)317
- Lee, K.-J., Heo, G. S., Kim, N. J. and Moon, D.-C.
Separation of theophylline and its analogues by micellar electrokinetic chromatography: application to the determination of theophylline in human plasma 577(1992)135
- Lee, M. G., see Park, K. H. 572(1991)259
- Lee, M. H., see Park, K. H. 572(1991)259
- Lee, M. L., see Johnson, D. F. 594(1992)283
- Lee, S., see Ono, S. 597(1992)293
- Lee, T., Yeung, E. S. and Sharma, M.
Micellar electrokinetic capillary chromatographic separation and laser-induced fluorescence detection of 2'-deoxynucleoside 5'-monophosphates of normal and modified bases 565(1991)197
- Lee, T. T. and Yeung, E. S.
High-sensitivity laser-induced fluorescence detection of native proteins in capillary electrophoresis 595(1992)319
- Lee Chin, J. R.
Determination of the catecholamines and serotonin, their precursors tyrosine and tryptophan, and their main metabolites in rat brain using reversed-phase high-performance liquid chromatography with fluorimetric and oxidative amperometric detection in series 578(1992)17

- Leenheers, L. H., Van Breugel, D. C., Ravensberg, J. C., Meuling, W. J. A. and Jongen, M. J. M.
Determination of 2-isopropoxyphenol in urine by capillary gas chromatography and mass-selective detection 578(1992)189
- Lefebvre, R. A., see Rosseel, M. T. 565(1991)504
- Legaz, M. E., see Vicente, C. 553(1991)271
- Legendre, J.-Y., see Tod, M. 594(1992)386
- Legrand, M. F., see Bruchet, A. 562(1991)469
- Lehman, E. D., Schaefer, T. F., Przysiecki, C. T., Joyce, J. G., Bailey, F. J., Schulman, C. A., Burke, C. J., Ramjit, H. G. and Miller, W. J.
Large-scale purification and characterization of recombinant tick anticoagulant peptide 574(1992)225
- Lehman-McKeeman, L. D., see Rodriguez, P. A. 563(1991)271
- Lehmann F., P. A., see Gomez, L. E. 575(1992)306
- Lehnert, B. E., see Gurley, L. R. 559(1991)411
- Leibman, E., see Solomon, B. 597(1992)257
- Leis, H. J., Ozand, P. T., Al Odaib, A. and Gleispach, H.
Quantitative measurement of N-acetyl-L-aspartic acid in urine by gas chromatography with negative-ion chemical ionization mass spectrometry 578(1992)116
- LeJemtel, T., see Bleiberg, B. 568(1991)301
- Lekova, K. and Ivanova, N.
Gas chromatographic analysis of some glycol ether analytes on a specially packed column 552(1991)353
- Le Leu, R. K., see Hayball, P. J. 570(1991)446
- Lelievre, G., see Thuaud, N. 555(1991)53
- Lembcke, B., see Stein, J. 576(1992)53
- Lemque, R., Jaulmes, A., Sébille, B., Vidal-Madjar, C. and Cysewski, P.
Study of the adsorption of self-associating proteins on an anion exchanger. Application to the chromatography of β -lactoglobulin B 599(1992)255
- Lemque, R., Vidal-Madjar, C., Racine, M., Piquion, J. and Sébille, B.
Anion-exchange chromatographic properties of α -lactalbumin eluted from quaternized polyvinylimidazole. Study of the role of the polymer coating 553(1991)165
- Lendrath, G., Bonekamp, A. and Kraus, L.
Quantitative planar chromatography of phospholipids with different fatty acid compositions 588(1991)303
- Lenhoff, A. M., see Shankar, A. 556(1991)235
- Lennard, L. and Singleton, H. J.
High-performance liquid chromatographic assay of the methyl and nucleotide metabolites of 6-mercaptopurine: quantitation of red blood cell 6-thioguanine nucleotide, 6-thioinosinic acid and 6-methylmercaptopurine metabolites in a single sample 583(1992)83
- Leoncini, R., see Terzuoli, L. 586(1991)153
- Leonhardt, B. A., see Warthen, Jr., J. D. 590(1992)133
- Lepistö, A.-M., see Auriola, S. O. K. 594(1992)153
- Lepojević, Ž., see Vrbaški, L. 558(1991)328
- Lepri, L., see Rodriguez, O. M. 555(1991)221
- Le Quéré, J. L., Sébédio, J. L., Henry, R., Couderc, F., Demont, N. and Promé, J. C.
Gas chromatography-mass spectrometry and gas chromatography-tandem mass spectrometry of cyclic fatty acid monomers isolated from heated fats 562(1991)659
- Lercker, G., see Antonelli, A. 552(1991)273
- Lercker, G., see Moret, S. 591(1992)175
- Le Roux, A. M., Wium, C. A., Joubert, J. R. and Van Jaarsveld, P. P.
Evaluation of a high-performance thin-layer chromatographic technique for the determination of salbutamol serum levels in clinical trials 581(1992)306
- Le Roux, M., see Burger, B. V. 552(1991)137
- Leroux, P., see Cherton, J.-C. 566(1991)499
- Leroy, P., Nicolas, A., Gavrilloff, C., Matt, M., Netter, P., Bannwarth, B., Herculim, B. and Mazza, M.
Determination of 2-mercaptopropionylglycine and its metabolite, 2-mercaptopropionic acid, in plasma by ion-pair reversed-phase high-performance liquid chromatography with post-column derivatization 564(1991)258
- Lesellier, E., Tchaplal, A., Péchard, M.-R., Lee, C. R. and Krstulović, A. M.
Separation of *trans/cis* α - and β -carotenes by supercritical fluid chromatography. II. Effect of the type of octadecyl-bonded stationary phase on retention and selectivity of carotenes 557(1991)59
- Lesellier, E., see Aubert, M.-C. 557(1991)47
- Leser, E. W. and Asenjo, J. A.
Rational design of purification processes for recombinant proteins 584(1992)43
- Lesseley, B. A., see Chen, C. L. 583(1992)274
- Lessor, R. A., see Florance, J. R. 559(1991)391
- Letarte, M., see Tarditi, L. 599(1992)13
- Letavernier, J.-F., see Voirin, A. 553(1991)155
- Letourneur, D., Migonney, V., Muller, D. and Jozefowicz, M.
Phosphorylated polystyrene resins in high-performance ion-exchange chromatography 589(1992)87
- Letter, W. S.
Preparative isolation of vitamin D₂ from previtamin D₂ by recycle high-performance liquid chromatography 590(1992)169
- Leube, J., see Ba, B. 572(1991)277
- Leuk, B., see Zimmerhackl, L. B. 587(1991)81
- Leung, P. Y. and Tsao, C. S.
Preparation of an optimum mobile phase for the simultaneous determination of neurochemicals in mouse brain tissues by high-performance liquid chromatography with electrochemical detection 576(1992)245
- Levels, P. J., see De Kok, T. M. C. M. 580(1992)135
- Le Verge, R., Le Corre, P., Chevanne, F., Döe De Maingreville, M., Royer, D. and Levy, J.
Determination of yohimbine and its two hydroxylated metabolites in humans by high-performance liquid chromatography and mass spectral analysis 574(1992)283
- Levesque, J., see Rey, J.-P. 587(1991)314
- Levesque, J., see Rey, J.-P. 596(1992)276

- Levin, S. and Abu-Lafi, S.
Adsorption isotherms of phenylalanine in a chromatographic column measured simultaneously by system peaks analysis and frontal analysis 556(1991)277
- Levine, B., see Kuhlman, Jr., J. J. 578(1992)207
- Levine, L., see Watanabe, M. 597(1992)309
- Levison, P. R., Badger, S. E., Toome, D. W., Koscielny, M. L., Lane, L. and Butts, E. T.
Economic considerations important in the scale-up of an ovalbumin separation from hen egg-white on the anion-exchange cellulose DE92 590(1992)49
- Levitsky, S., see Sarin, M. 563(1991)129
- Levy, J., see Le Verge, R. 574(1992)283
- Lewis, I. A. S., see Poon, G. K. 572(1991)143
- Lewis, J. A., Dolan, J. W., Snyder, L. R. and Molnar, I.
Computer simulation for the prediction of separation as a function of pH for reversed-phase high-performance liquid chromatography. II. Resolution as a function of simultaneous change in pH and solvent strength 592(1992)197
- Lewis, J. A., Lommen, D. C., Raddatz, W. D., Dolan, J. W., Snyder, L. R. and Molnar, I.
Computer simulation for the prediction of separation as a function of pH for reversed-phase high-performance liquid chromatography. I. Accuracy of a theory-based model 592(1992)183
- Lewis, R. C., see Rossi, D. T. 566(1991)257
- Leyland, D. M., Evershed, R. P., Edwards, R. H. T. and Beynon, R. J.
Application of gas chromatography-mass spectrometry with selected ion monitoring to the urinalysis of 4-pyridoxic acid 581(1992)179
- Leyssens, L., Driessen, C., Jacobs, A., Czech, J. and Raus, J.
Determination of β_2 -receptor agonists in bovine urine and liver by gas chromatography-tandem mass spectrometry 564(1991)515
- Lhermitte, M., see Trivier, J. M. 579(1992)269
- Li, J. and Caldwell, K. D.
Improved accuracy in the determination of field-flow fractionation elution volumes 555(1991)260
- Li, J., see Park, J. H. 586(1991)1
- Li, K.
Determination of sialic acids in human serum by reversed-phase liquid chromatography with fluorimetric detection 579(1992)209
- Li, S. F. Y., see Ng, C. L. 598(1992)133
- Li, S. F. Y., see Ong, C. P. 559(1991)529
- Li, S. F. Y., see Ong, C. P. 559(1991)537
- Li, S. F. Y., see Ong, C. P. 588(1991)335
- Li, S. F. Y., see Yeo, S. K. 585(1991)133
- Li, S. F. Y., see Yeo, S. K. 594(1992)335
- Li, S. F. Y., see Yik, Y. F. 585(1991)139
- Li, S. F. Y., see Yik, Y. F. 589(1992)333
- Li, T. M., see Harrington, S. J. 559(1991)385
- Li, W., Sotiriou-Leventis, C., Abdel-Baky, S., Fisher, D. H. and Giese, R. W.
Superoxide chemical transformation of diolepoxide polyaromatic hydrocarbon DNA adducts. Determination of benzo[a]pyrene-*r*-7,*t*-8,9,*c*-10-tetrahydrotretol by gas chromatography 588(1991)273
- Liang, W. T. C., Brocks, D. R. and Jamali, F.
Stereospecific high-performance liquid chromatographic assay of pirprofen enantiomers in rat plasma and urine 577(1992)317
- Liao, J.-L., Zhang, R. and Hjertén, S.
Continuous beds for standard and micro high-performance liquid chromatography 586(1991)21
- Liapis, A. I. and McCoy, M. A.
Theory of perfusion chromatography 599(1992)87
- Liau, L. S., see Ong, C.-N. 576(1992)346
- Liautard, J.-P.
Behavior of large nucleic acids in reversed-phase high-performance liquid chromatography 584(1992)135
- Liberato, D. J., see Fayer, B. E. 568(1991)135
- Libersa, C., see Trivier, J. M. 579(1992)269
- Lichtfouse, E., Freeman, K. H., Collister, J. W. and Merritt, D. A.
Enhanced resolution of organic compounds from sediments by isotopic gas chromatography-combustion-mass spectrometry 585(1991)177
- Lichý, A., Macek, J. and Adam, M.
Determination of hydroxylysylpyridinoline in tissues and urine by high-performance liquid chromatography 563(1991)153
- Lie, Ø. and Lambertsen, G.
Fatty acid composition of glycerophospholipids in seven tissues of cod (*Gadus morhua*), determined by combined high-performance liquid chromatography and gas chromatography 565(1991)119
- Liebich, H. M.
HPLC in clinical chemistry (by I. N. Papadoyannis) (Book Review) 574(1992)179
- Liebich, H. M., Wirth, C. and Jakober, B.
Analysis of polyunsaturated fatty acids in blood serum after fish oil administration 572(1991)1
- Liebich, H. M., see Mayer, F. 572(1991)339
- Liebmann, B., Henke, D., Spahn-Langguth, H. and Mutschler, E.
Determination of the quaternary compound ciclotropium in human biological material after hydrolysis and derivatization with the fluorophor flunoxapfen chloride 572(1991)181
- Liehr, J. G., see Roy, D. 567(1991)309
- Lien, K. A., see Dugan, M. E. R. 582(1992)242
- Liesiené, J., see Maruška, A. 596(1992)157
- Lietti, F., see Belotto, N. 572(1991)327
- Lightfield, A. R., see Brewster, J. D. 598(1992)23
- Lightfoot, E. N., see Athalye, A. M. 589(1992)71
- Lightfoot, Jr., E. N., see Grateful, T. M. 594(1992)341
- Lim, C. K., see Guo, R. 566(1991)383
- Lim, C. K., see Luo, J. 566(1991)409
- Limon, A., see Parra, P. 570(1991)185
- Lin, J. H., see Chen, I.-W. 576(1992)363
- Lin, S., Blanco, R. and Karger, B. L.
Adsorption-desorption isotherm hysteresis of β -lactoglobulin A with a weakly hydrophobic surface 557(1991)369
- Lin, S.-N., see Suter, M. J.-F. 553(1991)101
- Lin, S.-X., see Yang, F. 582(1992)71

- Linares, P., Gutiérrez, M. C., Lázaro, F., Luque de Castro, M. D. and Valcárcel, M.
Determination of benzocaine, dextromethorphan and cetylpyridinium ion by high-performance liquid chromatography with UV detection 558(1991)147
- Linares, P., Luque de Castro, M. D. and Valcárcel, M.
Determination of polyphosphates in intermediate materials for detergent manufacture by ion high-performance liquid chromatography with post-column derivatization 585(1991)267
- Lindberg, C., Paulson, J. and Blomqvist, A.
Evaluation of an automated thermospray liquid chromatography-mass spectrometry system for quantitative use in bioanalytical chemistry 554(1991)215
- Lindberg, C., see Paulson, J. 554(1991)149
- Lindblom, T., see Kårnsnäs, P. 599(1992)131
- Lindgård, B., Jönsson, J. Å. and Mathiasson, L.
Liquid membrane work-up of blood plasma samples applied to gas chromatographic determination of aliphatic amines 573(1992)191
- Linder, D., see Michel, C. 587(1991)93
- Lindgren, E. R., Pershing, D. W., Kirchgessner, D. A. and Drehmel, D. C.
Near-continuous measurement of hydrogen sulfide and carbonyl sulfide by an automatic gas chromatograph 585(1991)353
- Lindman, Y., see Ahnoff, M. 592(1992)323
- Lindner, W.
Chiral separations by liquid chromatography (edited by S. Ahuja) (Book Review) 595(1992)375
- Lindner, W., Uray, G. and Steiner, U.
(S,S)-Diphenylethylethanediamine derivatives as chiral selectors. II. Gasparrini-type bound chiral stationary phase with high enantioselectivity for naphthylamides 553(1991)373
- Ling, J. R., see Armstead, I. P. 586(1991)259
- Lingeman, H., see Barceló, D. 553(1991)311
- Lingeman, H., see Farjam, A. 589(1992)141
- Lingeman, H., see Özçimder, M. 570(1991)19
- Linhares, M. C. and Kissinger, P. T.
In vivo sampling using loop microdialysis probes coupled to a liquid chromatograph 578(1992)157
- Linhart, I., Vosmanská, M., Mitera, J. and Šmejkal, J.
Biotransformation of diethenylbenzenes. IV. A simple high-performance liquid chromatographic method for separation of urinary metabolites of 1,3-diethenylbenzene on analytical and semi-preparative scales 578(1992)215
- Lin Ling, B., Dewaele, C. and Baeyens, W. R. G.
Micro liquid chromatography with fluorescence detection of thiols and disulphides 553(1991)433
- Lintelmann, J., see Boos, K.-S. 600(1992)189
- Lipkowski, J., see Witkiewicz, Z. 600(1992)1
- Lisi, A. M., Kazlauskas, R. and Trout, G. J.
Diuretic screening in human urine by gas chromatography-mass spectrometry: use of a macroreticular acrylic copolymer for the efficient removal of the coextracted phase-transfer reagent after derivatization by direct extractive alkylation 581(1992)57
- Lisi, A. M., Trout, G. J. and Kazlauskas, R.
Screening for diuretics in human urine by gas chromatography-mass spectrometry with derivatization by direct extractive alkylation 563(1991)257
- Lisi, D. D., Stuart, J. D. and Snyder, L. R.
Computer simulation of gradient elution separation. Accuracy of predictions for non-linear gradients 555(1991)1
- Lisichkin, G. V., see Fadeev, A. Y. 558(1991)31
- Lisichkin, G. V., see Fadeev, A. Y. 596(1992)114
- Little, J. A., see Kuksis, A. 564(1991)11
- Little, J. A., see Kuksis, A. 579(1992)13
- Little, R. J., see Tsuji, K. 594(1992)317
- Littlejohn, D., see Robertson, A. M. 588(1991)15
- Litvak, M., see Fleminger, G. 597(1992)263
- Litvinenko, V. I., see Geodakyan, S. V. 577(1992)371
- Liu, C. Y., Chen, M. J. and Chai, T. J.
N-(Hydroxymethyl)thioamide resin as stationary phase in ion-exchange chromatography for metal ion separation 555(1991)291
- Liu, C.-Z., see Chen, F.-M. 563(1991)243
- Liu, D.-M.
Gas chromatographic separation of diastereomeric amino acid derivatives on chiral stationary phases. Application to the determination of enantiomeric composition in (S)-(+)-2-butanol 589(1992)249
- Liu, F.-S., see Wen, K.-C. 593(1992)191
- Liu, G., Djordjevic, N. M. and Erni, F.
High-temperature open-tubular capillary column liquid chromatography 592(1992)239
- Liu, G., Djordjevic, N. M. and Erni, F.
Reversed- and normal-phase separations by high-temperature open-tubular column liquid chromatography 598(1992)153
- Liu, H., Forman, L. J., Montoya, J., Eggers, C., Barham, C. and Delgado, M.
Determination of valproic acid by high-performance liquid chromatography with photodiode-array and fluorescence detection 576(1992)163
- Liu, H. and Wehmeyer, K. R.
Solid-phase extraction with supercritical fluid elution as a sample preparation technique for the ultratrace analysis of flavone in blood plasma 577(1992)61
- Liu, H. and Worthen, H. G.
Measurement of free amino acid levels in ultrafiltrates of blood plasma by high-performance liquid chromatography with automatic pre-column derivatization 579(1992)215
- Liu, J., Shirota, O. and Novotny, M.
Separation of fluorescent oligosaccharide derivatives by microcolumn techniques based on electrophoresis and liquid chromatography 559(1991)223
- Liu, S., Zhao, M. and Deng, C.
Separation and determination of trace amounts of vanadium(V), chromium(III) and iron(III) with 2-(2-thienylazo)-5-diethylaminophenol chelates by high-performance liquid chromatography 598(1992)298
- Liu, W., see Zhou, W. 589(1992)358
- Liu, X., see Lou, X. 586(1991)139

- Liu, X.-Q., see Xu, D.-K. 567(1991)451
- Liu, Y., see Lou, X. 552(1991)153
- Liu, Y.-M. and Sheu, S.-J.
Determination of ephedrine alkaloids by capillary electrophoresis 600(1992)370
- Llona, I., Annaert, W. G. and De Potter, W. P.
Simultaneous purification of the neuroproteins synapsin I and synaptophysin 596(1992)51
- Llorens, K. A., see Kunsman, G. W. 568(1991)427
- Lloyd, D. K., Cypess, A. M. and Wainer, I. W.
Determination of cytosine- β -D-arabinoside in plasma using capillary electrophoresis 568(1991)117
- Lloyd, D. K., Fried, K. and Wainer, I. W.
Determination of N-acetylator phenotype using caffeine as a probe compound: a comparison of high-performance liquid chromatography and capillary electrophoresis methods 578(1992)283
- Lo, M.-W., see Chiou, R. H.-Y. 581(1992)165
- Lo, M.-W., see Chiou, R. H.-Y. 581(1992)300
- Lo, M.-W., see Furtak, C. I. 573(1992)295
- Lo Casto, M., see Castagnetta, L. A. 572(1991)25
- Lo Coco, F., Cecon, L., Valentini, C. and Novelli, V.
High-performance liquid chromatographic determination of 2-furaldehyde in spirits 590(1992)235
- Lode, H., see Borner, K. 579(1992)285
- Lodhi, K. M., see Hagiwara, H. 597(1992)331
- Lodkowski, R., see Choma, I. 600(1992)109
- Lodkowski, R., see Supryniewicz, Z. 552(1991)463
- Loetscher, P., Mottlau, L. and Hochuli, E.
Immobilization of monoclonal antibodies for affinity chromatography using a chelating peptide 595(1992)113
- Loew, S., see Pichorner, H. 594(1992)400
- Löfgren, L. and Petersson, G.
Photoionization assessment of C₃-C₅ alkadienes and alkenes in urban air 591(1992)358
- Löfman, O., see Magnusson, P. 576(1992)79
- Lohmann, A. and Diekmann, N.
Determination of fendiline in human plasma by means of capillary gas chromatography and nitrogen-phosphorus selective detection 564(1991)289
- Lohmann, A., Diekmann, N., Walters, R. and Kraft, R.
Determination of bencyclane in human plasma by means of capillary gas chromatography and nitrogen-phosphorus selective detection 564(1991)283
- Lohmann, A. and Dingler, E.
New assay method for the determination of vinpocetine in human plasma by gas chromatography-mass spectrometry without transesterification caused by solvents 567(1991)506
- Lohmann, A., see Kraus, G. 573(1992)323
- Lohuis, C. P. G. G., see Essink, A. W. G. 570(1991)419
- Lombardi, F., Ardemagni, R., Colzani, V. and Visconti, M.
High-performance liquid chromatographic determination of rifloxacin and its main active metabolite in biological fluids 576(1992)129
- Lommen, D. C., see Lewis, J. A. 592(1992)183
- Lo Moro, A., see Raspi, G. 593(1992)119
- London, J. E., see Gurley, L. R. 559(1991)411
- London, J. E., see Gurley, L. R. 559(1991)431
- Long-Zhu, J., see Peng, S. 572(1991)73
- Longdon, K., see Witherow, L. 553(1991)497
- Longo, A., Di Toro, M., Galimberti, C. and Carenzi, A.
Determination of N-acetylcysteine in human plasma by gas chromatography-mass spectrometry 562(1991)639
- Lonkar, S. T., see Patil, S. F. 600(1992)344
- Lönnroth, P., Carlsten, J., Johnson, L. and Smith, U.
Measurements by microdialysis of free tissue concentrations of propranolol 568(1991)419
- Lödf, L., see Hellenäs, K.-E. 573(1992)69
- Lopes, J. C., see Rodrigues, A. E. 590(1992)93
- Lopes, T., see Lee, C. S. 559(1991)133
- Lopes Marques, R. M. and Schoenmakers, P. J.
Modelling retention in reversed-phase liquid chromatography as a function of pH and solvent composition 592(1992)157
- López, E., see Calull, M. 589(1992)151
- López-Barea, J., see Alhama, J. 586(1991)51
- López-Barea, J., see Alhama, J. 589(1992)121
- López-Pérez, M. J., see Ortin, A. 558(1991)357
- Lorber, B., see Meng, Q. C. 579(1992)63
- Lorenzini, R., see Betto, P. 563(1991)115
- Lork, K. D., see Unger, K. K. 556(1991)395
- Lorreyne, W., see Woestenborghs, R. 583(1992)223
- Löster, K., see Josić, D. 590(1992)59
- Loth, F., see Boeden, H.-F. 552(1991)389
- Lottspeich, F., see Ohhashi, T. 585(1991)153
- Lou, X., Liu, X., Zhang, S. and Zhou, L.
Enantiomeric separation of α -phenylethylamine and its substituted isomers by gas chromatography 586(1991)139
- Lou, X., Liu, Y. and Zhou, L.
Investigation of cross-linking chiral stationary phases within capillary columns 552(1991)153
- Louis, W. J., see Jackman, G. P. 566(1991)234
- Loun, B. and Hage, D. S.
Characterization of thyroxine-albumin binding using high-performance affinity chromatography. I. Interactions at the warfarin and indole sites of albumin 579(1992)225
- Loureiro, J. M., see Rodrigues, A. E. 590(1992)93
- Louter, A. J. H., see Özçimder, M. 570(1991)19
- Lovelace, J. L., Kusmier, J. J. and Desiderio, D. M.
Analysis of methionine enkephalin in human pituitary by multi-dimensional reversed-phase high-performance liquid chromatography, radioreceptor assay, radioimmunoassay, fast atom bombardment mass spectrometry, and mass spectrometry-mass spectrometry 562(1991)573
- Low, S. P., see Ong, C. P. 559(1991)529
- Lowe, C. R., see McCreath, G. E. 597(1992)189
- Lowes, S., Rose, M. E., Mills, G. A. and Pollitt, R. J.
Identification of urinary acylcarnitines using gas chromatography-mass spectrometry: preliminary clinical applications 577(1992)205
- Loy, G. L., Quick, Jr., A. N., Battaglia, F. C., Meschia, G. and Fennessey, P. V.
Measurement of leucine and α -ketoisocaproic acid fluxes in the fetal/placental unit 562(1991)169
- Lu, X., see Wu, C. 594(1992)243
- Lu, Y., see Min, S. 581(1992)272

- Lu, Z. P., see Rodrigues, A. E. 590(1992)93
- Lubczyk, B., see Hashem, A. 563(1991)216
- Lubli, G., see Tagliaro, F. 580(1992)161
- Lucarelli, C., Pelloso, R., Bruno, G., La Rosa, C. and Belliardo, F.
Evaluation of 2-(2-thiophenecarboxy)benzoic acid and related active metabolites in biological samples 573(1992)150
- Lucarelli, C., see Belliardo, F. 553(1991)41
- Lucarelli, C., see Belotto, N. 572(1991)327
- Lucarelli, C., see Brega, A. 553(1991)249
- Lucas, R. A., see Van Bergen, C. A. 589(1992)109
- Luchetti, M. M., see Corvetta, A. 566(1991)481
- Luckow, V., see Arnold, P. J. 544(1991)267
- Lucy, C. A. and Carr, P. W.
Analysis of the separability of plate height into overload and intrinsic contributions using the kinetic model of non-linear chromatography 556(1991)159
- Lucy, C. A., see Elchuk, S. 558(1991)197
- Luders, R. C., see Brunner, L. A. 568(1991)487
- Lüdi, H., see Manz, A. 593(1992)253
- Ludwig, R. C.
Application of a pellicular anion-exchange resin to the separation of inorganic and organic anions by single-column anion exchange 592(1992)101
- Lukasewycz, M. and Durhan, E.
Strategies for the identification of non-polar toxicants in aqueous environmental samples using toxicity-based fractionation and gas chromatography-mass spectrometry (Review) 580(1992)215
- Lundanes, E., see Baastoe, M. B. 558(1991)458
- Lundeen, J. T. and Juvet, Jr., R. S.
Quantitative resolution of severely overlapping gas chromatographic peaks. Isothermal and temperature-programmed operation 556(1991)305
- Lundmark, B.-O., see Bengård, A. 595(1992)247
- Lunina, E. V., see Fadeev, A. Y. 596(1992)114
- Lunte, C. E., see O'Shea, T. J. 593(1992)305
- Lunte, S. M., see O'Shea, T. J. 593(1992)305
- Luo, J. and Lim, C. K.
Random decarboxylation of uroporphyrinogen III by human hepatic uroporphyrinogen decarboxylase 566(1991)409
- Luo, T., see Xie, M. 555(1991)33
- Luong, C. B. H., Browner, M. F., Fletterick, R. J. and Haymore, B. L.
Purification of glycogen phosphorylase isozymes by metal-affinity chromatography 584(1992)77
- Luppa, P., see Jacob, K. 572(1991)317
- Luque, J., see Delgado, C. 594(1992)97
- Luque, J., see Tejedor, M. C. 589(1992)127
- Luque de Castro, M. D. and Valcárcel, M.
New approaches to coupling flow-injection analysis and high-performance liquid chromatography 600(1992)183
- Luque de Castro, M. D., see Linares, P. 558(1991)147
- Luque de Castro, M. D., see Linares, P. 585(1991)267
- Lurie, I. S., Cooper, D. A. and Klein, R. F. X.
High-performance liquid chromatographic analysis of benzodiazepines using diode array, electrochemical and thermospray mass spectrometric detection 598(1992)59
- Lurier, D., see Ducrottoy, G. 566(1991)415
- Lusby, W. R., see Khachik, F. 582(1992)153
- Lusby, W. R., see Warthen, Jr., J. D. 590(1992)133
- Luther, M. A., see Neidhardt, E. A. 590(1992)255
- Lux, M., see Dandeu, J.-P. 599(1992)105
- Lyddiatt, A., see Downham, M. 584(1992)59
- Lyddiatt, A., see Mohan, S. B. 584(1992)23
- Lynch, R. J., see Perez-Souto, N. 593(1992)209
- Lynes, G. W. and Hjelm, M.
Phenylalanine in whole blood and plasma — a candidate reference method 562(1991)213
- Lyttle, D. S., see Peterson, J. C. 564(1991)205
- Lyubimova, T., see Gelfi, C. 598(1992)277

M

- Mabuchi, M., see Banno, K. 568(1991)375
- Mabuchi, M., see Matsuoka, M. 581(1992)65
- Máca, B., see Pospíšil, J. 574(1992)71
- Macek, J., see Lichý, A. 563(1991)153
- Macek, K.
CRC handbook of chromatography: drugs, Vols. III-VI (edited by R.N. Gupta) (Book Review) 564(1991)355
- Macek, K.
Handbook of thin-layer chromatography (edited by J. Sherma and B. Fried) (Book Review) 588(1991)365
- Macek, K., see Nobilis, M. 581(1992)251
- MacFarlane, N., see O'Dowd, A. 577(1992)347
- Macfarlane, R. G., Midgley, J. M., Watson, D. G. and Evans, P. D.
Biogenic amines: their occurrence, biosynthesis and metabolism in the locust, *Schistocerca gregaria*, by gas chromatography-negative-ion chemical ionisation mass spectrometry 562(1991)585
- MacGregor, R. R., Fowler, J. S. and Wolf, A. P.
Determination of the enantiomeric composition of samples of cocaine by normal-phase high-performance liquid chromatography with UV detection 590(1992)354
- Machbert, G., see Böcker, R. H. 568(1991)363
- Macherel, L., see Crétier, G. 590(1992)175
- Machulskis, G. E., see Wilson, T. C. M. 582(1992)195
- Macielag, M. J., see Florance, J. R. 559(1991)391
- Mack, G.
Improved high-performance liquid chromatographic determination of ciprofloxacin and its metabolites in human specimens 582(1992)263
- Macka, M., Borák, J. and Kiss, F.
Separation of some platinum(II) complexes by ionic strength gradient on a solvent-generated ion-exchange sorbent 586(1991)291

- Macko, T. and Berek, D.
Experimental evaluation of sorbent bed homogeneity in high-performance liquid chromatographic columns 592(1992)109
- MacNeil, J. D., see Boison, J. O. 576(1992)315
- Macrae, P. V., see Cole, S. M. 562(1991)67
- Mada, A., see Akiyama, H. 579(1992)203
- Madajová, V., Turcelová, E. and Kaniansky, D.
Influence of poly(vinylpyrrolidone) on isotachophoretic separations of inorganic anions in aqueous electrolyte systems 589(1992)329
- Maddock, J., see Robinson, P. R. 564(1991)147
- Mader, R., see Knobloch, E. 566(1991)89
- Madhu, C., see Rozman, P. 574(1992)146
- Madsen, R., see Mitchell, E. P. 581(1992)31
- Madurawe, R. D., see Orthner, C. L. 558(1991)55
- Maeba, R., see Werner, A. 553(1991)205
- Maeda, M., see Ikegawa, S. 577(1992)229
- Maeder, G., Pelletier, M. and Haerdi, W.
Determination of amphetamines by high-performance liquid chromatography with ultraviolet detection. On-line pre-column derivatization with 9-fluorenylmethyl chloroformate and preconcentration 593(1992)9
- Maes, P. C., Nagels, L. J., Dewaele, C. and Alderweireldt, F. C.
Characterization of cellulase-based enzyme reactors for the high-performance liquid chromatographic determination of β -D-glucan oligosaccharides 558(1991)343
- Maestrelli, A., see Forni, E. 593(1992)177
- Maeyama, K., see Fukuda, H. 567(1991)459
- Maftouh, M., Ponthus, C., Tuong, A. and Picard, C.
Liquid secondary ion mass spectrometry applied to structural confirmation of enzymically prepared C-terminal-truncated derivatives of recombinant hirudin 562(1991)421
- Maggio, A., see Milana, M. R. 552(1991)205
- Maghuin-Rogister, G., see Vandebroek, M. 564(1991)405
- Maglulio, Jr., J., see Kuhlman, Jr., J. J. 578(1992)207
- Magnani, P., see Belotto, N. 572(1991)327
- Magni, F., Monti, L., Brambilla, P., Poma, R., Pozza, G. and Galli Kienle, M.
Determination of plasma [6,6-²H₂]glucose enrichment by a simple and accurate gas chromatographic-mass spectrometric method 573(1992)127
- Magnusson, O., Nilsson, L. B. and Westerlund, D.
Determination of dopamine and two of its acidic metabolites by high-performance liquid chromatography with electrochemical detection: a retrospective article (Chromatography Classic) 582(1992)1
- Magnusson, P., Löfman, O. and Larsson, L.
Determination of alkaline phosphatase isoenzymes in serum by high-performance liquid chromatography with post-column reaction detection 576(1992)79
- Maguire, D. J., see Cannell, G. R. 563(1991)341
- Maguire, M. H., Szabo, I., Slegel, P. and King, C. R.
Determination of concentrations of adenosine and other purines in human term placenta by reversed-phase high-performance liquid chromatography with photodiode-array detection: evidence for pathways of purine metabolism in the placenta 575(1992)243
- Magyar, K., see Cserhádi, T. 575(1992)57
- Mahuzier, G., see Tod, M. 594(1992)386
- Mainardi, L., see Gaspari, F. 570(1991)435
- Maisonneuve, A., Callebat, I., Debordes, L. and Coppet, L.
Specific and sensitive quantitation of 2,2'-dichlorodiethyl sulphide (sulphur mustard) in water, plasma and blood: application to toxicokinetic study in the rat after intravenous intoxication 583(1992)155
- Makino, K., see Ide, H. 596(1992)203
- Makino, M., see Moriyama, T. 571(1991)61
- Makino, Y. and Kawanishi, E.
High-performance liquid chromatographic separation of human apotransferrin and monoferric and diferric transferrins 567(1991)248
- Makino, Y. and Nishimura, S.
High-performance liquid chromatographic separation of human apolactoferrin and monoferric and diferric lactoferrins 579(1992)346
- Makita, M., see Kataoka, H. 578(1992)120
- Makita, T., see Sato, J. 575(1992)63
- Makki, S., Treffel, P., Humbert, P. and Agache, P.
High-performance liquid chromatographic determination of citropten and bergapten in suction blister fluid after solar product application in humans 563(1991)407
- Makuch, B., Gazda, K. and Cisowski, W.
Isolation of aristolochic acids HA₂I and HA₂II by preparative liquid chromatography 594(1992)145
- Malaguti, E., see Bonifaci, L. 585(1991)333
- Malaguti Aliberti, L., see Severini, G. 563(1991)147
- Malavasi, F., see Tarditi, L. 599(1992)13
- Malekzadeh, M., see Bergqvist, Y. 571(1991)169
- Malgouyat, J. M., see Girault, J. 564(1991)43
- Malhan née Chopra, R., see Jinno, K. 557(1991)459
- Malick, A. W., see Guzman, N. A. 559(1991)307
- Malick, A. W., see Guzman, N. A. 598(1992)123
- Mallee, L., see Scouten, W. H. 597(1992)231
- Mallet, A. I., Holland, K. T., Rennie, P. J., Watkins, W. J. and Gower, D. B.
Applications of gas chromatography-mass spectrometry in the study of androgen and odorous 16-androstene metabolism by human axillary bacteria 562(1991)647
- Mallett, C. D., see Page, K. M. 575(1992)319
- Mallia, A. K., see Nevens, J. R. 597(1992)247
- Malone, W., see Brown-Thomas, J. M. 568(1991)232
- Malone, W., see Christensen, R. G. 584(1992)207
- Maltby, D., see Peng, C. T. 586(1991)113
- Maltby, D., see Peng, C. T. 589(1992)231
- Mal'tsev, I. I., see Voinov, V. G. 586(1991)360
- Mamet, R., see Schoenfeld, N. 570(1991)51
- Manabe, T., see Isobe, T. 588(1991)115
- Manca, L., Masala, B., Ledda, S. and Naitana, S.
Separation of caprine globin chains by reversed-phase high-performance liquid chromatography: evidence for the presence of a silent β^B -globin allele in Sardinian sheep 563(1991)158
- Manchee, G. R., see Bowers, G. D. 554(1991)175
- Mancinelli, A., Bruno, G., Cardace, G., Morabito, E., Marzo, A. and Arrigoni Martelli, E.
High-performance liquid chromatographic evaluation of Med 15 and its metabolites Med 5 and tolmetin in rat plasma 553(1991)81

- Mancinelli, A., Pace, S., Marzo, A., Martelli, E. A. and Passetti, G.
Determination of pentoxifylline and its metabolites in human plasma by high-performance liquid chromatography with solid-phase extraction 575(1992)101
- Mancinelli, A., see D'Aranno, V. 574(1992)319
- Mancini, M., see Chauret, N. 588(1991)281
- Mandys, F., see Knobloch, E. 566(1991)89
- Manera, E., see Pietta, P. 553(1991)223
- Mañes, J., see Moltó, J. C. 555(1991)137
- Maness, N. O., Miranda, E. T. and Mort, A. J.
Recovery of sugar derivatives from 2-aminopyridine labeling mixtures for high-performance liquid chromatography using UV or fluorescence detection 587(1991)177
- Manfait, M., see Debal, V. 581(1992)93
- Mangalan, S., Patel, R. B., Gandhi, T. P. and Chakravathy, B. K.
Detection and determination of free and plasma protein-bound astemizole by thin-layer chromatography: a useful technique for bioavailability studies 567(1991)498
- Mangin, P., see Kintz, P. 579(1992)172
- Mangino, M. J., Zografakis, J., Murphy, M. K. and Anderson, C. B.
Improved and simplified tissue extraction method for quantitating long-chain acyl-coenzyme A thioesters with picomolar detection using high-performance liquid chromatography 577(1992)157
- Mangino, M. J., see Turk, J. 575(1992)183
- Mangoni, P., see Sioufi, A. 565(1991)401
- Man in 't Veld, A. J., see Alberts, G. 583(1992)236
- Man in 't Veld, A. J., see Boomsma, F. 574(1992)109
- Man in 't Veld, A. J., see Van Der Hoorn, F. A. J. 563(1991)348
- Mann-Buxbaum, E., see Ferris, R. 577(1992)251
- Manninen, A., see Pennanen, S. 568(1991)125
- Manning, M. A., see Tan, H. S. I. 568(1991)145
- Manno, B. R., see Kunsman, G. W. 568(1991)427
- Manno, J. E., see Kunsman, G. W. 568(1991)427
- Mannscheck, A., see Brandl, G. 586(1991)249
- Mannucci, C., see Triolo, A. 567(1991)105
- Mannucci, C., see Triolo, A. 568(1991)281
- Mant, C. T., see Hanson, M. 599(1992)65
- Mant, C. T., see Hanson, M. 599(1992)77
- Mant, C. T., see Zhu, B.-Y. 594(1992)75
- Mantero, F., see Carpenè, G. 553(1991)201
- Manz, A., Harrison, D. J., Verpoorte, E. M. J., Fettingler, J. C., Paulus, A., Lüdi, H. and Widmer, H. M.
Planar chips technology for miniaturization and integration of separation techniques into monitoring systems. Capillary electrophoresis on a chip 593(1992)253
- Manzari, A., see Venturelli, E. 582(1992)7
- Manzini, S., see D'Aranno, V. 574(1992)319
- Marai, L., see Kuksis, A. 588(1991)73
- Marasas, W. F. O., see Shephard, G. S. 566(1991)195
- Marcato, B., Fantazzini, C. and Sevinci, F.
Determination of polymeric hindered amine light stabilizers in polyolefins by high-performance liquid chromatography 553(1991)415
- Marcé, R. M., see Calull, M. 589(1992)151
- Marcé, R. M., see Calull, M. 590(1992)215
- Marcus, F. I., see Hoyer, G. L. 572(1991)159
- Mardens, Y., Kumps, A., Planchon, C. and Wurth, C.
Comparison of two extraction procedures for urinary organic acids prior to gas chromatography-mass spectrometry 577(1992)341
- Mareš, P., Řezanka, T. and Novák, M.
Analysis of human blood plasma triacylglycerols using capillary gas chromatography, silver ion thin-layer chromatographic fractionation and desorption chemical ionization mass spectrometry 568(1991)1
- Mareš, P., see Tvrzická, E. 563(1991)188
- Margou, V., see Krien, P. M. 576(1992)255
- Mariani, M., see Tarditi, L. 599(1992)13
- Marigo, M., see Krstulovic, A. M. 566(1991)273
- Marigo, M., see Tagliaro, F. 563(1991)11
- Marigo, M., see Tagliaro, F. 563(1991)299
- Marigo, M., see Tagliaro, F. 566(1991)333
- Marigo, M., see Tagliaro, F. 580(1992)161
- Marinello, E., see Terzuoli, L. 586(1991)153
- Marini, P. P., see Caccia, S. 581(1992)109
- Marinichev, A. N., Vitenberg, A. G. and Bureiko, A. S.
Efficiency of gas extraction in headspace analysis 600(1992)251
- Marino, A., see Laganà, A. 588(1991)89
- Mariot, R., Zangirolami, L., Bellato, P., Chen, S., Pieraccini, G. and Moneti, G.
Determination of the coumarin derivative cloricromene acid in rabbit plasma and platelets 576(1992)143
- Maris, F., see Hamoir, T. 589(1992)31
- Markides, K. E., see Johnson, D. F. 594(1992)283
- Markman, M., see Smith-Rogers, J. A. 566(1991)147
- Marle, I., Erlandsson, P., Hansson, L., Isaksson, R., Pettersson, C. and Pettersson, G.
Separation of enantiomers using cellulase (CBH I) silica as a chiral stationary phase 586(1991)233
- Marmont, F., see Silvestro, L. 563(1991)443
- Marrero, D., see Russell, J. W. 572(1991)321
- Maršálek, B., see Zahradníčková, H. 555(1991)239
- Marshall, G. T., see Jagodzinski, J. J. 591(1992)89
- Marshall, T., Williams, J. and Williams, K. M.
Electrophoresis of serum isoenzymes and proteins following acute myocardial infarction (Review) 569(1991)323
- Marston, A., see Slacanin, I. 557(1991)391
- Martelli, E. A., see Mancinelli, A. 575(1992)101
- Martens, H. J. M., see Van der Horst, A. 574(1992)166
- Marti, A., see Brügger, R. 557(1991)163
- Martin, G., Ghabrial, H., Morgan, D. J., Smallwood, R. A. and Sewell, R. B.
High-performance liquid chromatographic method to resolve and determine lipopolysaccharide sub-groups of *Escherichia coli* endotoxin in isolated perfused rat liver perfusate 574(1992)205
- Martin, I. C., see Boniface, P. J. 584(1992)199
- Martin, J. and White, I. N. H.
Fluorimetric determination of oxidised and reduced glutathione in cells and tissues by high-performance liquid chromatography following derivatization with dansyl chloride 568(1991)219

- Martin, M., see Cardot, P. J. P. 568(1991)93
 Martin, M., see Colin, H. 557(1991)137
 Martin, M., see El Fallah, M. Z. 557(1991)23
 Martin-Villacorta, J., see Méndez, R. 579(1992)115
 Martinez-Lara, E., see Pascual, P. 581(1992)49
 Martinez di Montemuros, F., see Pittalis, S. 573(1992)29
 Martinsen, A., see Auriola, S. 562(1991)737
 Martire, D. E., see Ghodbane, S. 556(1991)317
 Martire, D. E., see Riester, R. L. 588(1991)289
 Marubini, E., see Venturelli, E. 582(1992)7
 Maruška, A., Šerys, A., Liesienė, J., Urbonavičienė, J. and Žygas, A.
 Evaluation of morphological structure of packings by gel permeation chromatography 596(1992)157
 Maruta, T., see Oshima, T. 585(1991)255
 Maruyama, M. and Nagayoshi, T.
 Reversed-phase liquid chromatographic behaviour of alkylamines with amperometric detection and its application to trace analysis 594(1992)159
 Maruyama, M., see Hirayama, S. 588(1991)171
 Maruyama, W., see Niwa, T. 578(1992)109
 Maruyama, Y., see Ikarashi, Y. 575(1992)29
 Maruyama, Y., see Ikarashi, Y. 587(1991)306
 Marvin, C. A., see Sarre, S. 582(1992)29
 Marvin, C. H., Brindle, I. D., Hall, C. D. and Chiba, M.
 Rapid on-line precolumn high-performance liquid chromatographic method for the determination of benomyl, carbendazim and aldicarb species in drinking water 555(1991)147
 Maryanoff, C. A., see Shah, R. D. 588(1991)348
 Marzo, A., see Mancinelli, A. 553(1991)81
 Marzo, A., see Mancinelli, A. 575(1992)101
 Mas-Chamberlin, C., see Julien-Larose, C. 562(1991)39
 Masala, B., see Manca, L. 563(1991)158
 Mascher, H., Kikuta, C., Metz, R. and Vergin, H.
 New, high-sensitivity high-performance liquid chromatographic method for the determination of acyclovir in human plasma, using fluorometric detection 583(1992)122
 Masella, R., see Cantafora, A. 593(1992)139
 Mashige, F. and Ohkubo, A.
 Determination of proteins in urine by high-performance liquid chromatography with spectrophotometric detection using a pyrogallol red-molybdate complex 565(1991)173
 Masiello, P., see Fierabracci, V. 570(1991)285
 Maslen, C., see Gulaid, A. A. 566(1991)228
 Mason, J. L., Ashmore, S. P. and Aitkenhead, A. R.
 Simple method for the determination of morphine and its active glucuronide metabolite in human plasma by high-performance liquid chromatography with electrochemical detection 570(1991)191
 Mason, S. R., Ward, L. C. and Reilly, P. E. B.
 Fluorimetric detection of serum corticosterone using high-performance liquid chromatography 581(1992)267
 Mason, W. D., see Gumbhir, K. 583(1992)99
 Masotti, P., see Zonta, F. 594(1992)137
 Massart, D. L., see Bourguignon, B. 586(1991)11
 Massart, D. L., see Bourguignon, B. 592(1992)51
 Massart, D. L., see Hamoir, T. 589(1992)31
 Massart, D. L., see Van Belle, S. J.-P. 576(1992)351
 Massart, D. L., see Van Belle, S. J.-P. 578(1992)223
 Massart, D. L., see Vandebosch, C. 592(1992)37
 Massé, R., Bi, H., Ayotte, C., Du, P., Gélinas, H. and Dugal, R.
 Studies on anabolic steroids. V. Sequential reduction of methandienone and structurally related steroid A-ring substituents in humans: gas chromatographic-mass spectrometric study of the corresponding urinary metabolites 562(1991)323
 Massil, S. E., Ezra, I. and Fuss, L.
 Direct gas chromatographic analysis of O,O-dimethyl- and O,O-diethyldithiophosphoric acids and identification of their by-products 589(1992)375
 Massolini, G., see Baruffini, A. 593(1992)95
 Massolini, G., see Kitsos, M. 553(1991)1
 Massom, L. R. and Jarrett, H. W.
 High-performance affinity chromatography of DNA. II. Porosity effects 600(1992)221
 Masson, S., Sciaky, M., Desmoulin, F., Fontanarava, E. and Cozzone, P. J.
 Simple cation-exchange high-performance liquid chromatography optimized to the measurement of metabolites in the effluents from perfused rat livers using refractive index and ultraviolet detectors 563(1991)231
 Massot, C., see Kong Sing, Y. L. 558(1991)43
 Masuda, M., see Oshima, T. 585(1991)255
 Masuda, Y., see Katayama, M. 585(1991)219
 Masuoka, N., see Yao, K. 581(1992)11
 Matassa, L. C., Woodard, D., Leavitt, R. K., Firby, P. and Beaumier, P.
 Solid-phase extraction techniques for the determination of glycopyrrolate from equine urine by liquid chromatography-tandem mass spectrometry and gas chromatography-mass spectrometry 573(1992)43
 Mateescu, M. A., see Desmangles, R. J. P. 584(1992)121
 Mateos, J. L., see Vicente, C. 553(1991)271
 Mathian, B., see Chabraoui, L. 567(1991)299
 Mathiasson, L., see Karlsson, L. 557(1991)99
 Mathiasson, L., see Lindegård, B. 573(1992)191
 Mathieu-Boué, A., see Barilero, I. 575(1992)275
 Mathys, K. and Brenneisen, R.
 Determination of (*S*)-(-)-cathinone and its metabolites (*R,S*)-(-)-norephedrine and (*R,R*)-(-)-norpseudoephedrine in urine by high-performance liquid chromatography with photodiode-array detection 593(1992)79
 Matia, L., see Guardiola, A. 562(1991)481
 Matsubara, K., see Fukushima, S. 565(1991)35
 Matsubara, T. and Hayashi, A.
 Fragmentation pathways of O-trimethylsilyl ethers of dihydroxy long-chain bases analysed by linked-scan mass spectrometry 562(1991)119
 Matsubara, T., see Okamoto, J. 583(1992)45
 Matsubara, T., see Okamoto, J. 583(1992)53
 Matsuda, R., Hayashi, Y., Suzuki, T. and Saito, Y.
 Simplex optimization of liquid chromatography with the function of mutual information as a criterion 585(1991)187

- Matsui, H., see Ohhira, S. 566(1991)207
- Matsui, T., see Yoshioka, M. 566(1991)361
- Matsuishi, T., see Ohsawa, K. 597(1992)285
- Matsumoto, G. I., Friedmann, E. I., Watanuki, K. and Ocampo-Friedmann, R.
Novel long-chain *anteiso*-alkanes and *anteiso*-alkanoic acids in Antarctic rocks colonized by living and fossil cryptoendolithic microorganisms 598(1992)267
- Matsumoto, H., see Shimohigashi, Y. 597(1992)425
- Matsumoto, I., see Isosaki, K. 597(1992)123
- Matsumoto, I., see Kojima, K. 597(1992)323
- Matsumoto, I., see Matsumoto, M. 562(1991)139
- Matsumoto, I., see Ueno, M. 597(1992)197
- Matsumoto, M., Kuhara, T., Inoue, Y., Shinka, T. and Matsumoto, I.
Abnormal fatty acid metabolism in patients in hopantenate therapy during clinical episodes 562(1991)139
- Matsumoto, M., see Takeba, K. 596(1992)67
- Matsumoto, U., Nakayama, H., Shibusawa, Y. and Niimura, T.
Separation of human serum lipoproteins into three major classes by hydroxyapatite chromatography 566(1991)67
- Matsunaga, A., see Yamamoto, A. 585(1991)315
- Matsuo, S., see Ueno, H. 566(1991)57
- Matsuoka, M., see Banno, K. 564(1991)1
- Matsuoka, M., Horimoto, S., Mabuchi, M. and Banno, K.
Determination of three metabolites of a new angiotensin-converting enzyme inhibitor, imidapril, in plasma and urine by gas chromatography-mass spectrometry using multiple ion detection 581(1992)65
- Matsushima, E. and Sato, T.
Determination of bisaramil and its metabolite in plasma using high-performance liquid chromatography with electrochemical detection 573(1992)339
- Matsushita, S., Sugita, M., Motooka, I. and Kanaji, Y.
Determination of calcium ion in the presence of phosphate anion and collagen by capillary-type isotachopheresis 586(1991)363
- Matsushita, Y., see Watanabe, M. 597(1992)309
- Matsuura, T., see Baba, Y. 558(1991)273
- Matsuzaki, Y., see Honda, A. 565(1991)53
- Matsuzaki, Y., see Wakui, Y. 575(1992)131
- Matt, M., see Leroy, P. 564(1991)258
- Mattiasson, B., see Nilsson, M. 597(1992)383
- Matucha, M., Jockisch, W., Verner, P. and Anders, G.
Isotope effect in gas-liquid chromatography of labelled compounds 588(1991)251
- Matulík, F., see Urbánková, E. 583(1992)27
- Matuszewski, B., see Woolf, E. 577(1992)129
- Matuszewski, B. K., see Constanzer, M. L. 566(1991)127
- Matuszewski, B. K., see Herold, M. L. 581(1992)227
- Matuszewski, B. K., see Kline, W. F. 578(1992)31
- Matuszewski, B. K., see Kline, W. F. 583(1992)183
- Matz, S. G.
Determination of Cyasorb UV 1084 and its degradation products in low-density polyethylene film by high-performance liquid chromatography 587(1991)205
- Maurer, H. H.
Systematic toxicological analysis of drugs and their metabolites by gas chromatography-mass spectrometry (Review) 580(1992)3
- Mauri, P., see Pietta, P. 553(1991)223
- Mauri, P., see Pietta, P. 558(1991)296
- Mauri, P., see Pietta, P. 593(1992)165
- Mauri, P. L., Pietta, P. G. and Pace, M.
Automated high-performance liquid chromatographic assay of enzymatic activities 566(1991)327
- Maurich, V., Zacchigna, M. and Pitotti, A.
p-Nitrophenyllaurate: a substrate for the high-performance liquid chromatographic determination of lipase activity 566(1991)453
- Maxwell, J. R., see Eckardt, C. B. 557(1991)271
- May, W. E., see Brown-Thomas, J. M. 568(1991)232
- Mayer, F., Krämer, B. K., Röss, K. M., Kühlkamp, V., Liebich, H. M., Risler, T. and Seipel, L.
Simplified, rapid and inexpensive extraction procedure for a high-performance liquid chromatographic method for determination of disopyramide and its main metabolite mono-N-dealkylated disopyramide in serum 572(1991)339
- Mayewski, P. A., see Buck, C. F. 594(1992)225
- Maylin, G., see Steffenrud, S. 577(1992)221
- Mazur, J., Witkiewicz, Z. and Dąbrowski, R.
Gas chromatographic separation of *cis-trans* isomers of alkylcyclohexylbenzenes on a capillary column with a liquid crystalline stationary phase 600(1992)123
- Mazza, M., see Leroy, P. 564(1991)258
- Mazzanti, L., see Menichini, E. 555(1991)211
- Mberu, E. K., Muhia, D. K. and Watkins, W. M.
Measurement of halofantrine and its major metabolite desbutylhalofantrine in plasma and blood by high-performance liquid chromatography: a new methodology 581(1992)156
- Mberu, E. K., Ward, S. A., Winstanley, P. A. and Watkins, W. M.
Measurement of quinine in filter paper-absorbed blood by high-performance liquid chromatography 570(1991)180
- Mc Chrisley, B., see Driskell, J. A. 568(1991)333
- McCague, R., see Poon, G. K. 572(1991)143
- McCague, R., see Poon, G. K. 576(1992)235
- McCague, R., see Van Bakergem, E. 598(1992)189
- McCallister, J. D., see Patrick, K. S. 583(1992)254
- McCallum, J., see Koupai-Abyazani, M. R. 594(1992)117
- McCarthy, E., see Peloquin, C. A. 563(1991)472
- McClain, C. J., see Schmidt, J. 568(1991)207
- McCloskey, J. A., see Takeda, N. 562(1991)225
- McClure, P. C., see Burse, V. W. 566(1991)117
- McCoy, M. A., see Liapis, A. I. 599(1992)87
- McCreath, G. E., Chase, H. A., Purvis, D. R. and Lowe, C. R.
Novel affinity separations based on perfluorocarbon emulsions. Use of a perfluorocarbon affinity emulsion for the purification of human serum albumin from blood plasma in a fluidised bed 597(1992)189
- McCue, B. A., Park, Y. H., Brazzell, R. K. and Boltralik, J. J.
Capillary gas chromatographic-electron-capture assay for the aldose reductase inhibitor imirestat in lens and plasma 565(1991)255

- McDonnell, T. and Pawliszyn, J.
Capillary isotachopheresis with concentration-gradient detection. An application to the separation of synthetic peptides 559(1991)489
- McDowell, P. G., see Escott, R. E. A. 554(1991)281
- McErlane, K. M., see Aoyama, R. G. 588(1991)181
- McGee, J. P., Palin, K. J., Shaw, P. N. and Potter, C.
High-performance liquid chromatographic analysis of trilostane and ketotrilostane in rat plasma 567(1991)282
- McGhee, C. N. J., see Watson, D. G. 571(1991)101
- McGhie, T. K.
Analysis of serum methylmalonic acid for the determination of cobalt deficiency in cattle 566(1991)215
- McGuffey, J. E., see Bernert, Jr., J. T. 578(1992)1
- McGuire, V., see Kuksis, A. 564(1991)11
- McGuire, V., see Kuksis, A. 579(1992)13
- McKeage, M. J., see Kestell, P. 564(1991)315
- McKenna, K. F., see Torok-Both, G. A. 579(1992)99
- McLachlan, A. J., Tett, S. E. and Cutler, D. J.
High-performance liquid chromatographic separation of the enantiomers of hydroxychloroquine and its major metabolites in biological fluids using an α_1 -acid glycoprotein stationary phase 570(1991)119
- McLachlan, D. R., see Favarato, M. 576(1992)271
- McLaughlin, L. G. and Henion, J. D.
Determination of aminoglycoside antibiotics by reversed-phase ion-pair high-performance liquid chromatography coupled with pulsed amperometry and ion spray mass spectrometry 591(1992)195
- McLean, S., see Neill, G. P. 565(1991)207
- McLennan, E. A., see Uhe, A. M. 564(1991)81
- McMillan, J., see Bennett, J. L. 573(1992)146
- McMorland, G. H., see Wright, M. R. 565(1991)225
- McNair, H. M., see Chrisley, B. M. 563(1991)369
- Meacham, Jr., R. H., see Khetarpal, V. K. 567(1991)491
- Mearns, A. J., see Clark, B. J. 553(1991)383
- Measures, G., see Perez-Souto, N. 593(1992)209
- Mechoulam, R., see Harvey, D. J. 562(1991)299
- Medica, A., see Santoni, G. 581(1992)287
- Medina, M. B. and Schwartz, D. P.
Thin-layer chromatographic detection of zeranol and estradiol in fortified plasma and tissue extracts with Fast Corinth V 581(1992)119
- Medina Hernández, M. J., see Bonet Domingo, E. 582(1992)189
- Medzihradzsky, D., see Urbányi, Z. 578(1992)130
- Meenhorst, P. L., see Burger, D. M. 584(1992)239
- Meguro, H., see Akasaka, K. 596(1992)197
- Meier, P. and Thormann, W.
Determination of thiopental in human serum and plasma by high-performance capillary electrophoresis-micellar electrokinetic chromatography 559(1991)505
- Meijer, D. K. F., see Molema, G. 579(1992)107
- Meijer, E. M., see Coenen, A. J. J. M. 596(1992)59
- Meisen, A., see Dawodu, O. F. 587(1991)237
- Meister, A. E., see Berezkin, V. G. 552(1991)313
- Melander, A., see Rydberg, T. 564(1991)223
- Melenevskaya, E. Y., see Zimina, T. M. 593(1992)233
- Melis, M., see Cabras, P. 585(1991)164
- Melnikova, S. V., see Berezkina, L. G. 552(1991)319
- Melzer, H., see Dreyer, U. 592(1992)13
- Melzi d'Eril, G. V. and Rizzo, V.
Determination of free DOPA and 3-O-methyl-DOPA in human plasma by high-performance liquid chromatography with electrochemical detection 553(1991)265
- Ménager, S., see Vérité, P. 578(1992)134
- Méndez, R., Negro, A. and Martín-Villacorta, J.
High-performance liquid chromatographic methods for the determination of the penems SCH 29482 and FCE 22101 in human serum and urine 579(1992)115
- Méndez, R., see Negro, A. 576(1992)135
- Mendieta, J., see Delgado, C. 594(1992)97
- Meneguz, A., see Betto, P. 575(1992)117
- Meng, Q. C., King, S. J., Branham, K. E., Delucas, L. J., Lorber, B. and Oparil, S.
Preparative isolation of angiotensin-converting enzyme from human lung 579(1992)63
- Mengerink, Y., De Man, H. C. J. and Van der Wal, S.
Use of an evaporative light scattering detector in reversed-phase high-performance liquid chromatography of oligomeric surfactants 552(1991)593
- Mengerink, Y., see Coenen, A. J. J. M. 596(1992)59
- Menichini, E., Di Domenico, A., Bonanni, L., Corradetti, E., Mazzanti, L. and Zucchetti, G.
Reliability assessment of a gas chromatographic method for polycyclic aromatic hydrocarbons in olive oil 555(1991)211
- Mennecier, I., see Kamel, S. 574(1992)255
- Mensink, C. K., see Wieling, J. 594(1992)45
- Menzel-Soglowek, S., see Geisslinger, G. 568(1991)165
- Menzel-Soglowek, S., see Geisslinger, G. 573(1992)163
- Meo, J., see Richard, M. J. 577(1992)9
- Merati, R., see Riva, E. 553(1991)35
- Merdes, G., see Gebelein, M. 577(1992)146
- Merino, A., Bories, C., Gantier, J.-C. and Cardot, P. J. P.
Isolation of microfilariae from blood by gravitational field-flow fractionation 572(1991)291
- Merino-Dugay, A., Cardot, P. J. P., Czok, M., Guernet, M. and Andreux, J. P.
Monitoring of an experimental red blood cell pathology with gravitational field-flow fractionation 579(1992)73
- Merino-Dugay, A., see Bories, C. 579(1992)143
- Merlin, J. L., see Barberi, M. 565(1991)511
- Merlin, J. L., see Barberi-Heyob, M. 573(1992)247
- Merlin, J. L., see Barberi-Heyob, M. 581(1992)281
- Merritt, D. A., see Lichtfouse, E. 585(1991)177
- Mersmann, A., see Kerzel, P. 584(1992)109
- Merson, J. R., see Cole, S. M. 562(1991)67
- Mertens, K., Rogiers, V., Sonck, W. and Vercruyssen, A.
Measurement of reduced and oxidized glutathione in cultures of adult rat hepatocytes 565(1991)149
- Meschia, G., see Loy, G. L. 562(1991)169
- Mészáros, S., see Tomori, É. 578(1992)91
- Metcalfe, T., see Ueda, T. 593(1992)265
- Metz, R., see Mascher, H. 583(1992)122
- Metzler, M., see Spranger, B. 564(1991)485
- Meuleman, C., see Lansens, P. 586(1991)329

- Meuling, W. J. A., see Leenheers, L. H. 578(1992)189
- Meuwly, P., see Saugy, M. 564(1991)567
- Meyer, E., Lambert, W. E., De Leenheer, A. P., De Bersaques, J. P. and Kint, A. H.
Improved quantitation of 13-*cis*- and all-*trans*-acitretin in human plasma by normal-phase high-performance liquid chromatography 570(1991)149
- Meyer, H., see Cremer, J. 558(1991)430
- Meyer, H. H. D., Rinke, L. and Dürsch, I.
Residue screening for the β -agonists clenbuterol, salbutamol and cimaterol in urine using enzyme immunoassay and high-performance liquid chromatography 564(1991)551
- Meyer, R., see Kamata, K. 596(1992)233
- Meyer, V. R.
Peak purity, yield and throughput in preparative liquid chromatography with self-displacement 592(1992)17
- Meyer, V. R. and Ettre, L. S.
Early evolution of chromatography: the activities of Charles Dhéré 600(1992)3
- Mhatre, R. M. and Krull, I. S.
Interfacing gradient elution ion-exchange chromatography and low-angle laser light-scattering photometry for analysis of proteins 591(1992)139
- Miachon, S., see Duthel, J. M. 579(1992)85
- Michaelis, H. C., Kietzmann, D., Neurath, H., Jongepier, U. and Schilling, B.
Sensitive determination of piritramide in human plasma by gas chromatography 571(1991)257
- Michalczyk, L., Bialek, K. and Cohen, J. D.
Rapid determination of free tryptophan in plant samples by gas chromatography-selected ion monitoring mass spectrometry 596(1992)294
- Michalik, M., see Tröltzsch, C. 594(1992)111
- Michalski, W. P.
Resolution of three forms of superoxide dismutase by immobilised metal affinity chromatography 576(1992)340
- Michel, C., Buckel, W. and Linder, D.
Purification of the coenzyme B₁₂-containing 2-methyleneglutarate mutase from *Clostridium barkeri* by high-performance liquid chromatography 587(1991)93
- Micheletti, C., see Chiari, M. 558(1991)285
- Micheletti, C., see Chiari, M. 598(1992)287
- Michelot, D., see Dorizzi, R. 580(1992)279
- Michotte, Y., see Sarre, S. 575(1992)207
- Michotte, Y., see Sarre, S. 582(1992)29
- Middeldorp, J. M., see Hessing, M. 599(1992)267
- Midgley, J. M., see Macfarlane, R. G. 562(1991)585
- Midgley, J. M., see Watson, D. G. 571(1991)101
- Mielniczuk, Z., see Paczkowski, D. 573(1992)235
- Migonney, V., see Letourneur, D. 589(1992)87
- Mihara, R., see Sano, K. 578(1992)63
- Mikhailov, V. V., see Voinov, V. G. 586(1991)360
- Miksik, I., see Hanis, T. 553(1991)93
- Milana, M. R., Maggio, A., Denaro, M., Feliciani, R. and Gramiccioni, L.
Modern approach to the quantitative determination of volatiles in solid samples. Multiple headspace extraction gas chromatography for the determination of cyclohexanone residues in soil 552(1991)205
- Miles, C. J.
Determination of National Survey of Pesticides analytes in groundwater by liquid chromatography with postcolumn reaction detection 592(1992)283
- Milicua, J. C. G., see De Las Rivas, J. 585(1991)168
- Miljević, D., see Vučković, G. 585(1991)181
- Miller, D. J., see Eckert-Tilotta, S. E. 591(1992)313
- Miller, D. J., see Langenfeld, J. J. 594(1992)297
- Miller, D. J., see O'Dowd, A. 577(1992)347
- Miller, D. T., see Burse, V. W. 566(1991)117
- Miller, M., see Pesek, J. J. 556(1991)373
- Miller, N. S., see Neubecker, T. A. 574(1992)340
- Miller, R., La Grone, J., Skolnick, P. and Boje, K. M.
High-performance liquid chromatographic assay for 1-aminocyclopropanecarboxylic acid from plasma and brain 578(1992)103
- Miller, R. L. and DeVane, C. L.
Determination of cocaine, benzoylecgonine and ecgonine methyl ester in plasma by reversed-phase high-performance liquid chromatography 570(1991)412
- Miller, W. J., see Hurni, W. M. 559(1991)337
- Miller, W. J., see Lehman, E. D. 574(1992)225
- Miller-Stein, C., see Boppana, V. K. 593(1992)29
- Milligan, P. A.
Determination of piroxicam and its major metabolites in the plasma, urine and bile of humans by high-performance liquid chromatography 576(1992)121
- Millington, D. S., Norwood, D. L., Kodo, N., Moore, R., Green, M. D. and Berman, J.
Biomedical applications of high-performance liquid chromatography-mass spectrometry with continuous-flow fast atom bombardment 562(1991)47
- Millot, J.-M., see Debal, V. 581(1992)93
- Millot, M.-C., see Hervé, F. 577(1992)43
- Mills, G. A., see Lowes, S. 577(1992)205
- Mills, M. H., Finlay, D. C. and Haddad, P. R.
Determination of melatonin and monoamines in rat pineal using reversed-phase ion-interaction chromatography with fluorescence detection 564(1991)93
- Mills, T. A. and Phillips, C. S. G.
Study of salt hydrates by gas-solid chromatography 557(1991)495
- Milne, R. W., Nation, R. L., Reynolds, G. D., Somogyi, A. A. and Van Crugten, J. T.
High-performance liquid chromatographic determination of morphine and its 3- and 6-glucuronide metabolites: improvements to the method and application to stability studies 565(1991)457
- Milojković, D. M., see Janjić, T. J. 596(1992)91
- Milon, H.
Identification of poly(ethylene terephthalate) cyclic oligomers by liquid chromatography-mass spectrometry 554(1991)305
- Mimura, Y., Nakamura, K. and Takeo, K.
Analysis of the interaction between an $\alpha(1\rightarrow6)$ dextran-specific mouse hybridoma antibody and dextran B512 by affinity electrophoresis 597(1992)345
- Min, S., Yisheng, Y. and Lu, Y.
Determination of urinary 3-methylhistidine by high-performance liquid chromatography with *o*-phthalaldehyde precolumn derivatization 581(1992)272

- Minárik, M., see Sitkey, V. 585(1991)373
- Minder, E. I., see Feng, N. 575(1992)289
- Mingalyov, P. G., see Fadeev, A. Y. 596(1992)114
- Minger, A., see Thormann, W. 593(1992)275
- Mínguez-Mosquera, M. I., Gandul-Rojas, B., Montaña-Asquerino, A. and Garrido-Fernández, J.
Determination of chlorophylls and carotenoids by high-performance liquid chromatography during olive lactic fermentation 585(1991)259
- Minguillón, C., see Oliveros, L. 589(1992)53
- Minkler, P. E., see Krahenbuhl, S. 573(1992)3
- Minobe, S., see Nawata, M. 597(1992)415
- Minouchi, T., see Shibata, N. 566(1991)187
- Miranda, E. T., see Maness, N. O. 587(1991)177
- Misawa, S., see Tanaka, E. 584(1992)267
- Míscoria, G., see Delhotal Landes, B. 577(1992)117
- Mishra, U., see Singh, A. K. 568(1991)351
- Misra, S., see Sharma, S. D. 594(1992)379
- Missbichler, A., see Ferris, R. 577(1992)251
- Mitamura, K., see Shimada, K. 565(1991)111
- Mitchell, E. P., Evans, L., Schultz, P., Madsen, R., Yarbrow, J. W., Gehrke, C. W. and Kuo, K.
Modified nucleosides in human serum 581(1992)31
- Mitchell, R., see Ueda, T. 592(1992)229
- Mitchell, R., see Ueda, T. 593(1992)265
- Mitchell, S. C., see Zhang, A. Q. 584(1992)141
- Mitera, J., see Linhart, I. 578(1992)215
- Mitina, V. K., see Klyashchitsky, B. A. 577(1992)267
- Mitsui, T., see Hida, M. 585(1991)345
- Miura, J., see Kamahori, M. 567(1991)351
- Miwa, H., Yamamoto, M. and Asano, T.
High-performance liquid chromatographic analysis of fatty acid compositions of platelet phospholipids as their 2-nitrophenylhydrazides 568(1991)25
- Miwa, T., Sakashita, S., Ozawa, H., Haginaka, J., Asakawa, N. and Miyake, Y.
Application of an ovomucoid-conjugated polymer column for the enantiospecific determination of chlorprenaline concentrations in plasma 566(1991)163
- Miyabara, Y., Tabata, M., Suzuki, J. and Suzuki, S.
Separation and sensitive determination of i-urobilin and 1-stercobilin by high-performance liquid chromatography with fluorimetric detection 574(1992)261
- Miyajima, A., see Svoboda, M. 562(1991)403
- Miyakawa, T., see Kamizono, A. 567(1991)113
- Miyake, Y., see Miwa, T. 566(1991)163
- Miyamoto, K., see Kamizono, A. 567(1991)113
- Miyamoto, S., see Ide, H. 596(1992)203
- Miyamoto, T., see Ishii, K. 564(1991)338
- Miyamoto, Y., see Kataoka, H. 578(1992)120
- Miyasaka, A., see Imoto, T. 557(1991)383
- Miyazaki, H., see Honda, A. 565(1991)53
- Miyazaki, H., see Sakurai, T. 571(1991)1
- Miyazaki, K., see Kishino, S. 582(1992)246
- Miyazaki, M., see Etoh, T. 578(1992)292
- Miyazaki, M., see Yamamoto, A. 585(1991)315
- Mizugaki, M., see Ishibashi, M. 562(1991)613
- Mizukami, E., see Yamamoto, A. 585(1991)315
- Mizutani, T., see Oshima, T. 585(1991)255
- Mizzen, C. A., see Favarrato, M. 576(1992)271
- Moak, S. W., see Driskell, J. A. 568(1991)333
- Moats, W. A.
High-performance liquid chromatographic determination of penicillin G, penicillin V and cloxacillin in beef and pork tissues 593(1992)15
- Moberg, C., Norin, T., Rákos, L. and Sjödin, K.
Ligand-exchange chromatography of alkenes on stationary phases containing palladium(II) complexes. Enantiomeric separation of *trans*-1,2-divinylcyclohexane 585(1991)309
- Möckel, H. J., see Dreyer, U. 592(1992)13
- Modin, R., see Castledine, J. B. 592(1992)27
- Modrek, B., see Ahmed, F. 599(1992)25
- Moeller, M. R.
Drug detection in hair by chromatographic procedures (Review) 580(1992)125
- Mofidi, M., Hudson, M. K. and Underhill, K.
Flame infrared emission detection of hydrogen 589(1992)241
- Mogadeddi, H., see Feldman, B. J. 594(1992)275
- Mögele, R., Pabel, B. and Galensa, R.
Determination of organic acids, amino acids and saccharides by high-performance liquid chromatography and a postcolumn enzyme reactor with amperometric detection 591(1992)165
- Mohan, S. B. and Lyddiatt, A.
Ligand location and biochemical productivity of silica-based immunoaffinity adsorbents 584(1992)23
- Möhrke, W., see Spahn-Languth, H. 584(1992)229
- Mokashi, A. A., see Wagh, J. S. 587(1991)280
- Molema, G., Jansen, R. W., Visser, J. and Meijer, D. K. F.
Simultaneous analysis of azidothymidine and its mono-, di- and triphosphate derivatives in biological fluids, tissue and cultured cells by a rapid high-performance liquid chromatographic method 579(1992)107
- Molina, F., Jehl, F., Gallion, C., Penner, F. and Monteil, H.
Determination of the third generation oral cephalosporin cefpodoxime in biological fluids by high-speed high-performance liquid chromatography 563(1991)205
- Molnar, I., see Däppen, R. 592(1992)133
- Molnar, I., see Lewis, J. A. 592(1992)183
- Molnar, I., see Lewis, J. A. 592(1992)197
- Molnár-Perl, I., see Fábíán, V. 553(1991)87
- Molnár-Perl, I., see Morvai, M. 552(1991)337
- Molnár-Perl, I., see Morvai, M. 600(1992)87
- Molteni, S., see Thormann, W. 589(1992)321
- Molteni, S., see Thormann, W. 593(1992)275
- Moltó, J. C., Picó, Y., Font, G. and Mañes, J.
Determination of triazines and organophosphorus pesticides in water samples using solid-phase extraction 555(1991)137
- Monakhova, N. I., see Nemirovskii, V. D. 594(1992)23
- Monastyrsky, V. A., see Fadeev, A. Y. 596(1992)114
- Moncrieff, J.
Determination of theophylline in serum and saliva in the presence of caffeine and its metabolites 568(1991)177

- Moncrieff, J.
Extractionless determination of diclofenac sodium in serum using reversed-phase high-performance liquid chromatography with fluorimetric detection 577(1992)185
- Moncrieff, J.
Determination of cetirizine in serum using reversed-phase high-performance liquid chromatography with ultraviolet spectrophotometric detection 583(1992)128
- Monde, K. and Takasugi, M.
Studies on stress metabolites. XVI. High-performance liquid chromatographic analysis of cruciferous phytoalexins using complex ternary mobile phase gradients 598(1992)147
- Monder, C., see Iohan, F. 564(1991)27
- Moneti, G., see Mariot, R. 576(1992)143
- Monger, L. S. and Olliff, C. J.
Measurement of β -endorphin in human plasma by high-performance liquid chromatography with electrochemical detection: validation of a method employing the simultaneous purification and concentration of β -endorphin 577(1992)239
- Monger, L. S. and Olliff, C. J.
Reversed-phase chromatographic behaviour of β -endorphin: evidence of conformational change 595(1992)125
- Monsaert, C., see Van Belle, S. J.-P. 576(1992)351
- Monsaert, C., see Van Belle, S. J.-P. 578(1992)223
- Monster, A. C., see Kežić, S. 563(1991)199
- Montagna, M., see Poletini, A. 564(1991)529
- Montaño, A., De Castro, A., Rejano, L. and Sánchez, A.-H.
Analysis of zapatera olives by gas and high-performance liquid chromatography 594(1992)259
- Montaño-Asquerino, A., see Mínguez-Mosquera, M. I. 585(1991)259
- Montay, G., see Vergniol, J. C. 582(1992)273
- Montefiori, D., see Forni, E. 593(1992)177
- Monteil, H., see Enjalbert, F. 598(1992)227
- Monteil, H., see Molina, F. 563(1991)205
- Montero, E., see Ruiz-Gutierrez, V. 581(1992)171
- Monti, L., see Magni, F. 573(1992)127
- Montoya, J., see Liu, H. 576(1992)163
- Moon, D.-C., see Lee, K.-J. 577(1992)135
- Moore, C. M., see Browne, S. P. 575(1992)158
- Moore, R., see Millington, D. S. 562(1991)47
- Moore, T. A., see Chang, S. Y. 562(1991)111
- Moore, W. T., see Suter, M. J.-F. 553(1991)101
- Morabito, E., see Mancinelli, A. 553(1991)81
- Morcol, T., see Orthner, C. L. 558(1991)55
- Moreau, J., Palette, C., Cordonnier, P., Naline, E., Advenier, C. and Pays, M.
Separation and identification of the 4-hydroxyantipyrine sulphoconjugate 576(1992)103
- Morehead, H., see Talmadge, K. W. 590(1992)83
- Morehead, H. W., Talmadge, K. W., O'Shannessy, D. J. and Siebert, C. J.
Optimization of oxidation of glycoproteins: an assay for predicting coupling to hydrazide chromatographic supports 587(1991)171
- Moret, S., Bortolomeazzi, R. and Lercker, G.
Improvement of extraction procedure for biogenic amines in foods and their high-performance liquid chromatographic determination 591(1992)175
- Moretti, F., see Carducci, C. 553(1991)149
- Moretti, V. M., Van de Water, C. and Haagsma, N.
Automated high-performance liquid chromatographic determination of chloramphenicol in milk and swine muscle tissue using on-line immunoaffinity sample clean-up 583(1992)77
- Morgan, D. J., see Martin, G. 574(1992)205
- Mori, H., see Nagasawa, M. 577(1992)275
- Mori, S., see Mukoyama, Y. 588(1991)195
- Moriguchi, S., see Isosaki, K. 597(1992)123
- Morin, P., Gallois, A., Richard, H. and Gaydou, E.
Fast separation of polymethoxylated flavones by carbon dioxide supercritical fluid chromatography 586(1991)171
- Morin-Allory, L. and Herbreteau, B.
High-performance liquid chromatography and supercritical fluid chromatography of monosaccharides and polyols using light-scattering detection. Chemometric studies of the retentions 590(1992)203
- Morishita, H., see Fukushima, S. 565(1991)35
- Morita, I., Kawamoto, M. and Yoshida, H.
Difference in the concentration of tryptophan metabolites between maternal and umbilical foetal blood 576(1992)334
- Moritani, C., see Ohhashi, T. 585(1991)153
- Moritz, R. L. and Simpson, R. J.
Application of capillary reversed-phase high-performance liquid chromatography to high-sensitivity protein sequence analysis 599(1992)119
- Moriyama, H., see Nozaki, O. 570(1991)1
- Moriyama, T., Tozawa, T., Yamashita, H., Onodera, S., Nobuoka, M. and Makino, M.
Separation and characterization of sialic acid-containing salivary-type amylase from patients' sera with immunoglobulin A-type myeloma 571(1991)61
- Morjani, H., see Debal, V. 581(1992)93
- Morris, R. G., see Sallustio, B. C. 576(1992)321
- Morrissey, M. A. and Widmer, H. M.
Ion-mobility spectrometry as a detection method for packed-column supercritical fluid chromatography 552(1991)551
- Morselli, P. L., see Ascalone, V. 581(1992)237
- Mort, A. J., see Maness, N. O. 587(1991)177
- Mortimer, N., see Escott, R. E. A. 553(1991)423
- Mortimer, R. H., see Cannell, G. R. 563(1991)341
- Mortimer, R. H., see Cannell, G. R. 564(1991)310
- Morvai, M., Fábán, V. and Molnár-Perl, I.
Buffer and pH dependence of the retention of phenylthiocarbamylamino acids in reversed-phase high-performance liquid chromatography 600(1992)87
- Morvai, M., Molnár-Perl, I. and Knausz, D.
Simultaneous gas-liquid chromatographic determination of sugars and organic acids as trimethylsilyl derivatives in vegetables and strawberries 552(1991)337
- Morvai, M., see Fábán, V. 553(1991)87
- Mosandl, A., see Karl, V. 587(1991)347

- Moschera, J., see Guzman, N. A. 559(1991)307
Moschera, J., see Guzman, N. A. 598(1992)123
Moscoso, H., see Kutlar, F. 567(1991)359
Moseley, M. A., see Deterding, L. J. 554(1991)73
Moseley, M. A., see Deterding, L. J. 554(1991)329
Moskovkina, M., see Dimov, N. 552(1991)59
Motohashi, M., see Yamashita, K. 570(1991)329
Motohashi, M., see Yamashita, K. 577(1992)174
Motohashi, N., see Kamata, K. 596(1992)233
Motoki, K., see Toyoda, H. 565(1991)141
Motooka, I., see Matsushita, S. 586(1991)363
Mottlau, L., see Loetscher, P. 595(1992)113
Mougeot, G., see Bories, C. 579(1992)143
Moulin, D. E., see Bouquillon, A. I. 577(1992)354
Mouloungui, Z., see Snini, A. 590(1992)369
Mount, D. L., Churchill, F. C. and Bergqvist, Y.
Determination of mefloquine in blood, filter paper-
absorbed blood and urine by 9-fluorenylmethyl
chloroformate derivatization followed by liquid
chromatography with fluorescence detection
564(1991)181
Mounts, T. L., see Abidi, S. L. 598(1992)209
Moussaoui, M., Benlyas, M. and Wahl, P.
Diffusion of proteins in the chromatographic gel AcA-34
558(1991)71
Moussaoui, M., Benlyas, M. and Wahl, P.
Diffusion of proteins in Sepharose Cl-B gels
591(1992)115
Mubumbila, M. V., see Carelse, O. 558(1991)318
Much, H., see Pasch, H. 589(1992)295
Muda, M., see Salomoni, M. 572(1991)312
Mue, S., see Watanabe, M. 597(1992)309
Mugnai, G., see Cecconi, O. 555(1991)267
Muh, J. P.
Sensitivity and specificity of urinary N-acetyldopamine as
a marker for neuroblastomas: comparison with
traditional urinary catecholamine metabolites. Reply to
Muskiet and Kema 579(1992)187
Muh, J. P., see Jouve, J. 567(1991)331
Muh, J. P., see Jouve, J. 574(1992)9
Muhia, D. K., see Mberu, E. K. 581(1992)156
Muñoz Blanco, M. T., see Ortin, A. 558(1991)357
Mukai, Y., see Yoshioka, M. 566(1991)361
Mukaida, N., see Okutani, R. 577(1992)25
Mukoyama, Y., Shimizu, N., Sakata, T.-I. and Mori, S.
Elution behaviour of polyamic acid and polyamide-imide
in size-exclusion chromatography 588(1991)195
Mulder, G. J., see Venizelos, V. 573(1992)259
Mulders, T. M. T., see Venizelos, V. 573(1992)259
Mulheim, C., see Cherton, J.-C. 566(1991)511
Mulholland, F. and Hague, G. R.
Monitoring 4-methoxy-2,3,6-trimethylbenzenesulphonyl
deprotection of arginine-containing synthetic peptides
using capillary zone electrophoresis 589(1992)380
Müller, C., see Köppel, C. 575(1992)87
Muller, D., see Letourneur, D. 589(1992)87
Müller, H.-M., see Schwadorf, K. 595(1992)259
Muller, N., see Payan, E. 566(1991)9
Müller, R., see Boeden, H.-F. 552(1991)389
Müller-Werner, B., see Thurl, S. 568(1991)291
Mullin, R. J. and Duch, D. S.
Preparation of a chiral matrix. Resolution of (6*R,S*)-N⁵-
formyltetrahydrofolate 555(1991)254
Mullins, M. A., see Shepard, R. M. 565(1991)321
Mulvana, D. E., see Janiszewski, J. S. 577(1992)151
Munafa, A., see Saugy, M. 564(1991)567
Munniksmä, C. N., see De Jong, J. 574(1992)273
Muñoz de la Peña, A., see Anigbogu, V. C. 594(1992)37
Munro, P. D., see Winzor, D. J. 597(1992)57
Munro, Z. M., see Burger, B. V. 552(1991)137
Munsters, B. H. M., see Duchateau, A. L. L. 552(1991)605
Murachi, T., see Tabata, M. 597(1992)435
Murai, Y., Ohno, M., Konishi, M. and Kominami, G.
Selective clean-up method using an immunoaffinity
column following radioimmunoassay of prostaglandin F_{2α}
in biological fluids 597(1992)201
Murakami, A., see Ide, H. 596(1992)203
Muramatsu, I., see Tamaki, M. 574(1992)65
Muramatsu, S., see Watanabe, M. 597(1992)309
Murano, M., see Noguchi, H. 573(1992)336
Murata, M. and Ide, T.
Determination of cholesterol in sub-nanomolar quantities
in biological fluids by high-performance liquid
chromatography 579(1992)329
Murata, S., see Komatsu, S. 568(1991)460
Murphy, G. M., see Stubbs, A. P. 570(1991)293
Murphy, M. K., see Mangino, M. J. 577(1992)157
Murray, S. and Boobis, A. R.
Combined assay for phenacetin and paracetamol in
plasma using capillary column gas chromatography-
negative-ion mass spectrometry 568(1991)341
Murray, S., Wellings, R., Taylor, I. K., Fuller, R. W. and
Taylor, G. W.
Gas chromatographic-mass spectrometric assay to
measure urinary N¹-methylhistamine excretion in man
567(1991)289
Murray, Y. L., see Thedford, R. 582(1992)253
Muschik, G. M., see Atamna, I. Z. 588(1991)315
Muskiet, F. A. J. and Kema, I. P.
Sensitivity and specificity of urinary N-acetyldopamine as
a marker for neuroblastomas: comparison with
traditional urinary catecholamine metabolites
579(1992)184
Muskiet, F. A. J., see Hessels, J. 563(1991)1
Muskiet, F. A. J., see Hoving, E. B. 565(1991)103
Mussini, E., see Salomoni, M. 572(1991)312
Musson, D. G., Bidgood, A. M. and Olejnik, O.
Assay methodology for prednisolone, prednisolone
acetate and prednisolone sodium phosphate in rabbit
aqueous humor and ocular physiological solutions
565(1991)89
Mutschler, E., see Liebmann, B. 572(1991)181
Mutschler, E., see Spahn-Langguth, H. 584(1992)229
Mutter, M., see Steiner, V. 586(1991)43
Myers, C., see Allen, E. 578(1992)175
Myers, C., see Ogata, T. 575(1992)39
Myher, J. J., see Kuksis, A. 564(1991)11
Myher, J. J., see Kuksis, A. 579(1992)13
Myher, J. J., see Kuksis, A. 588(1991)73

N

- Naaranlahti, T., see Auriola, S. 554(1991)227
 Naaranlahti, T., see Auriola, S. 562(1991)737
 Naaranlahti, T., see Auriola, S. O. K. 594(1992)153
 Nachman, M.
 Kinetic aspects of membrane-based immunoaffinity chromatography 597(1992)167
 Nachman, M., Azad, A. R. M. and Bailon, P.
 Membrane-based receptor affinity chromatography 597(1992)155
 Nagabhushan, T. L., see Pramanik, B. N. 562(1991)377
 Nagae, N., see Nimura, N. 585(1991)207
 Nagae, N., see Takeuchi, T. 595(1992)121
 Nagai, K., see Sakihama, N. 597(1992)147
 Naganuma, H., see Nakanishi, A. 591(1992)159
 Nagasawa, M., Sasabe, H., Shimizu, T. and Mori, H.
 Use of a carrier for quantitation of a new dihydropyridine calcium antagonist (OPC-13340) in human plasma by highly sensitive gas chromatography with negative-ion chemical ionization mass spectrometry 577(1992)275
 Nagasawa, T., Sakai, T. and Onodera, R.
 Simple and sensitive determination of plasma N⁺-methylhistidine by high-performance liquid chromatography using pre-column derivative formation with *o*-phthalaldehyde-2-mercaptoethanol 566(1991)223
 Nagasawa, T., see Hagiwara, H. 597(1992)331
 Nagasawa, T., see Sakai, T. 576(1992)155
 Nagashima, H., see Jinno, K. 594(1992)105
 Nagata, O., see Takahara, E. 576(1992)174
 Nagata, T. and Saeki, M.
 Determination of thiamphenicol residues in chicken muscles by column liquid chromatography 565(1991)471
 Nagata, T., see Kage, S. 564(1991)163
 Nagata, Y., Yamamoto, K. and Shimojo, T.
 Determination of D- and L-amino acids in mouse kidney by high-performance liquid chromatography 575(1992)147
 Nagatomo, A., Sakai, K., Nose, T., Shimohigashi, Y. and Ohno, M.
 Occurrence of an allosteric transition in the modification of papain with L-1-acetyl-2,3-dihydropyrrolo[2,3-*b*]-indole-2-carboxamide 597(1992)411
 Nagatsu, T.
 Application of high-performance liquid chromatography to the study of biogenic amine-related enzymes (Review) 566(1991)287
 Nagatsu, T., see Niwa, T. 578(1992)109
 Nagaya, T., see Ide, H. 596(1992)203
 Nagayoshi, T., see Maruyama, M. 594(1992)159
 Nagels, L. J., see Maes, P. C. 558(1991)343
 Nageswara Rao, R., see Husain, S. 558(1991)435
 Nageswara Sarma, P., see Husain, S. 558(1991)435
 Naggi, A., see Silvestro, L. 591(1992)225
 Naidong, W., Verresen, K., Busson, R., Roets, E. and Hoogmartens, J.
 Isolation of doxycycline, 6-epidoxycycline and 2-acetyl-2-decarboxamidometacycline from commercial metacycline by preparative column liquid chromatography on silica gel 586(1991)67
 Naidong, W., Verresen, K., Roets, E. and Hoogmartens, J.
 Determination of metacycline and related substances by column liquid chromatography on poly(styrene-divinylbenzene) 586(1991)61
 Naikwadi, K., see Beard, A. 589(1992)265
 Naitana, S., see Manca, L. 563(1991)158
 Nakada, F., see Fukushima, S. 565(1991)35
 Nakagawa, G., see Yasui, T. 596(1992)73
 Nakagawa, Y. and Iwatani, K.
 Scanning thin-layer chromatography-liquid secondary ion mass spectrometry and its application for investigation of drug metabolites 562(1991)99
 Nakagomi, K., see Saito, K. 595(1992)163
 Nakahara, D., see Niwa, T. 578(1992)109
 Nakahara, K., see Hirokawa, T. 555(1991)321
 Nakajima, M., see Wakabayashi, H. 573(1992)154
 Nakajima, M., see Yamato, S. 590(1992)241
 Nakamoto, A., see Ueda, T. 592(1992)229
 Nakamoto, A., see Ueda, T. 593(1992)265
 Nakamura, K., Kashiwagi, S. and Takeo, K.
 Characterization of the interaction between human plasma fibronectin and collagen by means of affinity electrophoresis 597(1992)351
 Nakamura, K., see Mimura, Y. 597(1992)345
 Nakamura, K., see Takeo, K. 597(1992)365
 Nakamura, S., see Endoh, Y. S. 579(1992)123
 Nakamura, S., see Watanabe, K. 566(1991)109
 Nakanishi, A., Naganuma, H., Kondo, J., Watanabe, K., Hirano, K., Kawasaki, T. and Kawahara, Y.
 3-Bromomethyl-7-methoxy-1,4-benzoxazin-2-one as a highly sensitive fluorescence derivatization reagent for carboxylic acids in high-performance liquid chromatography 591(1992)159
 Nakano, M. and Kawahara, S.
 High-performance liquid chromatographic method for simultaneous determination of enantiomers of 5-dimethylsulphamoyl-6,7-dichloro-2,3-dihydrobenzofuran-2-carboxylic acid and its N-monodemethyl metabolite in monkey plasma and urine after chiral derivatization 564(1991)235
 Nakano, M., see Kamizono, A. 567(1991)113
 Nakano, S., see Noguchi, H. 583(1992)266
 Nakashima, K., Okamoto, M., Yoshida, K., Kuroda, N., Akiyama, S. and Yamaguchi, M.
 Preparation of a fluorescent derivative of benzoyllecgonine, and preliminary studies of its application to the analysis of urine 584(1992)275
 Nakashima, K., see Toyo'oka, T. 588(1991)61
 Nakata, H.
 Affinity chromatography in purification of A₁ adenosine receptors 597(1992)335
 Nakata, K., see Ohmori, S. 566(1991)1

- Nakata, K., see Ohmori, S. 574(1992)35
- Nakayama, F., see Sakurai, T. 571(1991)1
- Nakayama, H., see Matsumoto, U. 566(1991)67
- Nakayama, M., see Etoh, T. 578(1992)292
- Nakazawa, H., see Okamoto, M. 588(1991)177
- Nakazawa, H., see Saito, K. 595(1992)163
- Nakazawa, H., see Takeba, K. 596(1992)67
- Naline, E., see Moreau, J. 576(1992)103
- Naline, E., see Palette, C. 563(1991)103
- Nambara, T.
Advances in steroid analysis (edited by S. Görög) (Book Review) 594(1992)403
- Nambara, T., see Goto, J. 567(1991)343
- Nambara, T., see Goto, J. 574(1992)1
- Nambara, T., see Goto, J. 585(1991)281
- Nambara, T., see Iida, T. 558(1991)451
- Nanak, E., see Goubbran-Botros, H. 597(1992)357
- Naoi, M., see Niwa, T. 578(1992)109
- Narang, P. K., see Rossi, D. T. 566(1991)257
- Nardella, P. A., see Coutant, J. E. 570(1991)139
- Nardi, A., see Pessi, A. 557(1991)307
- Nardi, A., see Quaglia, M. G. 593(1992)259
- Nardi, A., see Sakodinskaya, I. K. 596(1992)95
- Narsimha, R., see Husain, S. 596(1992)127
- Narsimha, R., see Husain, S. 600(1992)316
- Naruse, Y., see Uda, T. 586(1991)131
- Naruto, S., see Kurono, M. 578(1992)152
- Naruto, S., see Noguchi, H. 573(1992)336
- Naruto, S., see Noguchi, H. 583(1992)266
- Nashabeh, W. and El Rassi, Z.
Capillary zone electrophoresis of proteins with hydrophilic fused-silica capillaries 559(1991)367
- Nashabeh, W. and El Rassi, Z.
Enzymophoresis of nucleic acids by tandem capillary enzyme reactor–capillary zone electrophoresis 596(1992)251
- Nashabeh, W. and El Rassi, Z.
Capillary zone electrophoresis of linear and branched oligosaccharides 600(1992)279
- Nassif, J., see Burse, V. W. 566(1991)117
- Nation, R. L., see Hayball, P. J. 570(1991)446
- Nation, R. L., see Milne, R. W. 565(1991)457
- Navarro, J. L., see Alaiz, M. 591(1992)181
- Navarro-González, R., Negrón-Mendoza, A. and Albarrán, G.
Analysis of keto acids as their methyl esters of 2,4-dinitrophenylhydrazone derivatives by gas chromatography and gas chromatography–mass spectrometry 587(1991)247
- Navvab, M., see Talmadge, K. W. 590(1992)83
- Nawata, M., Minobe, S., Hase, M., Watanabe, T., Sato, T. and Tosa, T.
Specific assay for endotoxin using immobilized histidine, *Limulus* amoebocyte lysate and a chromogenic substrate 597(1992)415
- Nayak, V. S.
New solid adsorbents for the separation of lower hydrocarbons and permanent gases. II. Ammonium molybdophosphate 556(1991)425
- Naylor, S., see Day, B. W. 562(1991)563
- Naylor, S., see Kajbaf, M. 575(1992)75
- Naylor, S., see Kajbaf, M. 583(1992)63
- Nazarova, V. I., Shcherbakova, K. D. and Shcherbakova, O. A.
Chromatographic properties of graphitized thermal carbon black modified with a monolayer of liquid crystal 600(1992)59
- Ndou, T. T., see Anigbogu, V. C. 594(1992)37
- Neff, W. E., Zeitoun, M. A. M. and Weisleder, D.
Resolution of lipolysis mixtures from soybean oil by a solid-phase extraction procedure 589(1992)353
- Neffe, S., see Grajek, H. 600(1992)67
- Negawa, M. and Shoji, F.
Optical resolution by simulated moving-bed adsorption technology 590(1992)113
- Negri, A., see Biondi, P. A. 566(1991)377
- Negro, A., Alvarez-Bujidos, M. L., Ortiz, A. I., Cubría, J. C., Méndez, R. and Ordóñez, D.
Reversed-phase ion-pair high-performance liquid chromatographic determination of triclabendazole metabolites in serum and urine 576(1992)135
- Negro, A., see Alvarez-Bujidos, M. L. 578(1992)321
- Negro, A., see Méndez, R. 579(1992)115
- Negrón-Mendoza, A., see Navarro-González, R. 587(1991)247
- Neidhardt, E. A., Luther, M. A. and Recny, M. A.
Rapid, two-step purification process for the preparation of pyrogen-free murine immunoglobulin G₁ monoclonal antibodies 590(1992)255
- Neidlinger, T. R., see Burri, B. J. 567(1991)369
- Neill, G. P., Davies, N. W. and McLean, S.
Automated screening procedure using gas chromatography–mass spectrometry for identification of drugs after their extraction from biological samples 565(1991)207
- Nelson, C. C. and Foltz, R. L.
Chromatographic and mass spectrometric methods for determination of lysergic acid diethylamide (LSD) and metabolites in body fluids (Review) 580(1992)97
- Nelson, R. J., see Guttman, A. 593(1991)297
- Nelson, T. J.
Deconvolution method for accurate determination of overlapping peak areas in chromatograms 587(1991)129
- Nemirovskii, V. D., Monakhova, N. I. and Kostenko, V. G.
High-performance liquid chromatographic study of the correlation between the physico-chemical parameters of furan and benzene aldehydes and the rate of their metabolism by yeast 594(1992)23
- Netter, P., see Leroy, P. 564(1991)258
- Netter, P., see Payan, E. 566(1991)9
- Neubecker, T. A., Miller, N. S., Asquith, T. N. and Driscoll, K. E.
Measurement of histamine in rat bronchoalveolar lavage fluid by high-performance cation-exchange chromatography coupled with post-column derivatization to *o*-phthalaldehyde 574(1992)340
- Neubert, D., see Kastner, M. 587(1991)43
- Neubert, D., see Kastner, M. 587(1991)117
- Neufeld, R., see Chauret, N. 588(1991)281

- Neufeld, T., see Fleming, G. 597(1992)263
- Neumann, G. D., see Glässgen, W. E. 598(1992)81
- Neumeier, D., see Jacob, K. 572(1991)317
- Neurath, H., see Michaelis, H. C. 571(1991)257
- Nevens, J. R., Mallia, A. K., Wendt, M. W. and Smith, P. K.
Affinity chromatographic purification of immunoglobulin M antibodies utilizing immobilized mannan binding protein 597(1992)247
- Neville, C., see Snider, J. 599(1992)141
- Ng, C. L., Lee, H. K. and Li, S. F. Y.
Systematic optimization of capillary electrophoretic separation of sulphonamides 598(1992)133
- Ng, C. L., see Ong, C. P. 559(1991)537
- Ng, C. L., see Ong, C. P. 588(1991)335
- Ng, J., see Zhu, J. 577(1992)77
- Ngo, T. T. and Khatter, N.
Avid AL, a synthetic ligand affinity gel mimicking immobilized bacterial antibody receptor for purification of immunoglobulin G 597(1992)101
- Nguyen, B. L., see Patrick, K. S. 583(1992)254
- Nguyen, H., see Dixit, V. 563(1991)379
- Nguyen, P. T., see Salamoun, J. 596(1992)43
- Nguyen, T. T., see Day, B. W. 562(1991)563
- Nichols, M. A., see Champney, T. H. 579(1992)334
- Nicolas, A., see Leroy, P. 564(1991)258
- Nicolas, P., see Tod, M. 575(1992)171
- Nicolotti, A., see Pellegrino, S. 592(1992)279
- Nielen, M. W. F.
Quantitative aspects of indirect UV detection in capillary zone electrophoresis 588(1991)321
- Nielsen, E., see Vidal, P. 573(1992)201
- Nieman, A. A., see Farjam, A. 589(1992)141
- Niessen, W. M. A.
Quantitative chemical analysis (by D. C. Harris) (Book Review) 587(1991)368
- Niessen, W. M. A., Tjaden, U. R. and Van der Greef, J.
Strategies in developing interfaces for coupling liquid chromatography and mass spectrometry 554(1991)3
- Niessen, W. M. A., see Van Bakergem, E. 598(1992)189
- Niessen, W. M. A., see Heeremans, C. E. M. 554(1991)205
- Niessen, W. M. A., see Kokkonen, P. S. 565(1991)265
- Niessen, W. M. A., see Poon, G. K. 576(1992)235
- Niessen, W. M. A., see Tinke, A. P. 554(1991)119
- Niessen, W. M. A., see Verheij, E. R. 554(1991)339
- Nigo, Y., see Tomita, M. 571(1991)324
- Niimura, T., see Matsumoto, U. 566(1991)67
- Nijjar, M. S., Grimmelt, B. and Brown, J.
Purification of domoic acid from toxic blue mussels (*Mytilus edulis*) and phytoplankton 568(1991)393
- Nikitin, D. I., see Řezanka, T. 558(1991)215
- Nikitin, Y. S., see Gavrilo, T. B. 552(1991)179
- Nikitin, Y. S., see Lanin, S. N. 558(1991)81
- Nilsson, K. G. I., see Birnbaum, S. 587(1991)268
- Nilsson, L. B.
Optimization of a peak compression system for a remoxipride metabolite (FLA797) and its application to bioanalysis 591(1992)207
- Nilsson, L. B., see Magnusson, O. 582(1992)1
- Nilsson, L. B., see Pettersson, K.-J. 581(1992)161
- Nilsson, M., Håkanson, H. and Mattiasson, B.
Process monitoring by flow-injection immunoassay. Evaluation of a sequential competitive binding assay 597(1992)383
- Nimura, N., Itoh, H., Kinoshita, T., Nagae, N. and Nomura, M.
Fast protein separation by reversed-phase high-performance liquid chromatography on octadecylsilyl-bonded non-porous silica gel. Effect of particle size of column packing on column efficiency 585(1991)207
- Nishi, H., Fukuyama, T. and Terabe, S.
Chiral separation by cyclodextrin-modified micellar electrokinetic chromatography 553(1991)503
- Nishida, M., see Okuda, T. 567(1991)141
- Nishihara, K., see Ohmori, S. 574(1992)35
- Nishikawa, M., see Endoh, Y. S. 579(1992)123
- Nishikawa, T., see Hashimoto, A. 582(1992)41
- Nishimura, S., see Makino, Y. 579(1992)346
- Nishioka, R., Umeda, I., Ōi, N., Tabata, S. and Uno, K.
Determination of felodipine and its metabolites in plasma using capillary gas chromatography with electron-capture detection and their identification by gas chromatography-mass spectrometry 565(1991)237
- Nishiyama, F., see Hu, J.-Y. 589(1992)339
- Nishiyama, F., see Hu, J.-Y. 594(1992)371
- Nitecki, D. E., see Cunico, R. L. 559(1991)467
- Nitsch, E., see Haidacher, D. 591(1992)351
- Nitsche, H., see Friebe, S. 598(1992)139
- Nitta, K. and Watanabe, A.
Determination of calcium ions tightly bound to proteins 585(1991)173
- Nityanand, S., see Paliwal, J. K. 572(1991)219
- Nitz, S., Kollmannsberger, H., Weinreich, B. and Drawert, F.
Enantiomeric distribution and ¹³C/¹²C isotope ratio determination of γ -lactones: appropriate methods for the differentiation between natural and non-natural flavours? 557(1991)187
- Niwa, T., Maruyama, W., Nakahara, D., Takeda, N., Yoshizumi, H., Tatematsu, A., Takahashi, A., Dostert, P., Naoi, M. and Nagatsu, T.
Endogenous synthesis of N-methylsalsolinol, an analogue of 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine, in rat brain during *in vivo* microdialysis with salsolinol, as demonstrated by gas chromatography-mass spectrometry 578(1992)109
- Nobilis, M., Pastera, J., Svoboda, D., Května, J. and Macek, K.
High-performance liquid chromatographic determination of ambroxol in human plasma 581(1992)251
- Nobuhara, K., see Okamoto, M. 556(1991)407
- Nobuoka, M., see Moriyama, T. 571(1991)61
- Noctor, T. A. G., Wainer, I. W. and Hage, D. S.
Allosteric and competitive displacement of drugs from human serum albumin by octanoic acid, as revealed by high-performance liquid affinity chromatography, on a human serum albumin-based stationary phase 577(1992)305
- Noda, A., see Eto, S. 568(1991)157
- Noda, A., see Eto, S. 579(1992)253

- Noda, H., see Eto, S. 568(1991)157
 Noda, H., see Eto, S. 579(1992)253
 Noguchi, H., Tomita, N., Naruto, S. and Nakano, S.
 Determination of gliclazide in serum by high-performance liquid chromatography using solid-phase extraction 583(1992)266
 Noguchi, H., Yoshida, K., Murano, M. and Naruto, S.
 Determination of nadolol in serum by high-performance liquid chromatography with fluorimetric detection 573(1992)336
 Noguchi, K., see Abe, I. 565(1991)183
 Noguchi, K., see Hirata, N. 592(1992)93
 Noguchi, K., see Yasukawa, K. 597(1992)271
 Noguchi, M., see Kanai, M. 567(1991)415
 Nolan, P. E., see Hoyer, G. L. 572(1991)159
 Nolan, R. J., see Langvardt, P. W. 567(1991)93
 Nolan, S. L., see Boniface, P. J. 584(1992)199
 Nolan, T., see Kiang, C.-H. 567(1991)195
 Nölting, B., see Brandl, G. 586(1991)249
 Noma, J., Hayashi, N. and Sekiba, K.
 Automated direct high-performance liquid chromatographic assay for estretol, estriol, cortisone and cortisol in serum and amniotic fluid 568(1991)35
 Nomura, A., see Kishino, S. 582(1992)246
 Nomura, H., Teshima, E. and Hakusui, H.
 Simple isocratic high-performance liquid chromatographic method for measurement of iodixanol in human plasma 572(1991)333
 Nomura, M., see Nimura, N. 585(1991)207
 Noordhoek, J., see Van 't Klooster, G. A. E. 579(1992)158
 Noordhoek, J., see Van 't Klooster, G. A. E. 579(1992)355
 Nooter, K., see De Vries, A. J. 563(1991)435
 Nooyen, W. J., see Van Tellingen, O. 553(1991)47
 Nooyen, W. J., see Van Tellingen, O. 573(1992)328
 Nooyen, W. J., see Van Tellingen, O. 574(1992)293
 Nooyen, W. J., see Van Tellingen, O. 576(1992)158
 Nordgren, T., see Lanbeck-Vallén, K. 567(1991)121
 Nordholm, L.
 Determination of the neuroprotective agent 6-nitro-7-sulphamoylbenzof[quinoxaline-2,3-dione in plasma using high-performance liquid chromatography 568(1991)481
 Norin, T., see Moberg, C. 585(1991)309
 Noronha da Silveira, I., see Dos Ramos, F. J. 558(1991)125
 Norris, R., see Taylor, P. J. 581(1992)152
 Norton, R. M., see Hedeén, K. M. 572(1991)239
 Norwood, D. L., see Millington, D. S. 562(1991)47
 Nose, N., see Saito, K. 595(1992)163
 Nose, T., see Nagatomo, A. 597(1992)411
 Nose, Y., see Ohmori, S. 566(1991)1
 Nouws, J. F. M., see Vree, T. B. 579(1992)131
 Novák, J., see Beran, M. 558(1991)265
 Novák, M., see Mareš, P. 568(1991)1
 Nováková, N., see Urbánková, E. 583(1992)27
 Novaković, J., see Tvrzická, E. 563(1991)188
 Novelli, M., see Fierabracci, V. 570(1991)285
 Novelli, V., see Lo Cocco, F. 590(1992)235
 Novomeský, P., see Strašák, M. 585(1991)337
 Novotný, L., see Reichelová, V. 588(1991)147
 Novotny, M., see Liu, J. 559(1991)223
 Novotny, M., see Snopek, J. 559(1991)215
 Novotny, M. V., see Soini, H. 559(1991)547
 Nowotnik, D. P., see Rosenspire, K. C. 574(1992)119
 Nozaki, O., Ohata, T., Ohba, Y., Moriyama, H. and Kato, Y.
 Determination of serum cortisol by reversed-phase liquid chromatography using precolumn sulphuric acid-ethanol fluorescence derivatization and column switching 570(1991)1
 Numajiri, Y., see Kato, Y. 562(1991)81
 Nunn, A. D., see Rosenspire, K. C. 574(1992)119
 Nyberg, F., see Silberring, J. 554(1991)83
 Nyberg, F., see Silberring, J. 562(1991)459
 Nyman, A., see Hellenäs, K.-E. 573(1992)69
 Nyström, L., Salén, G. and Hedner, T.
 Improved method for the determination of 5,6-dihydroxytryptamine and 5,7-dihydroxytryptamine in tissue using high-performance liquid chromatography with electrochemical detection 574(1992)137
 Nyssönen, K. and Salonen, J. T.
 Comparison of gel permeation chromatography, density gradient ultracentrifugation and precipitation methods for quantitation of very-low-, low- and high-density lipoprotein cholesterol 570(1991)382
- ## O
- Obach, R., see Pruñonosa, J. 574(1992)127
 Obach, R., see Pruñonosa, J. 581(1992)219
 Oberdorfer, F., Kemper, K., Kaleja, M., Reusch, J. and Gottschall, K.
 Carbohydrate analysis with ion chromatography using Eurokat stationary phases. Preparative separation of monosaccharides and their fluorinated derivatives 552(1991)483
 Obrezkov, O. N., Shpigun, O. A., Zolotov, Y. A. and Shlyamin, V. I.
 Ion chromatographic determination of anions, especially sulphur-containing anions, with conductimetric and kinetic detection 558(1991)209
 O'Brien, A. P., see Cooles, G. P. 588(1991)259
 O'Brien, J. E., see Flynn, J. W. 583(1992)91
 Ocampo, A. P., see Chen, S. 558(1991)141
 Ocampo-Friedmann, R., see Matsumoto, G. I. 598(1992)267
 Occhiato, E., see Busini, S. 577(1992)180
 Oda, Y., Asakawa, N., Abe, S., Yoshida, Y. and Sato, I.
 Avidin protein-conjugated column for direct injection analysis of drug enantiomers in plasma by high-performance liquid chromatography 572(1991)133
 O'Dea, K., see Uhe, A. M. 564(1991)81
 Odham, G., see Elmroth, I. 598(1992)43
 Odink, J., see De Vries, M. 564(1991)250
 Odle, J., see Van Kempen, T. A. T. G. 584(1992)157

- Odom, J. D., see Terry, Jr., A. V. 585(1991)101
- O'Donovan, G. A., see Dutta, P. K. 599(1992)137
- O'Dowd, A., O'Dowd, J. J., O'Dowd, J. J. M., MacFarlane, N., Abe, H. and Miller, D. J.
Analysis of novel imidazoles from isolated perfused rabbit heart by two high-performance liquid chromatographic methods 577(1992)347
- O'Dowd, J. J., see O'Dowd, A. 577(1992)347
- O'Dowd, J. J. M., see O'Dowd, A. 577(1992)347
- O'Dwyer, P. J., see LaCreta, F. P. 571(1991)271
- O'Dwyer, P. J., see Tinsley, P. W. 564(1991)303
- Oe, T., see Shimada, K. 558(1991)306
- Oefner, P., see Seitz, U. 559(1991)499
- Oefner, P. J., see Huber, C. G. 599(1992)113
- Oesch, F., Wagner, H., Platt, K. L. and Arand, M.
Improved sample preparation for the testosterone hydroxylation assay using disposable extraction columns 582(1992)232
- Offermanns, J., see Thurl, S. 568(1991)291
- Ogata, T., Myers, C. and Blumer, J. L.
Determination of platelet monoamine oxidase activity by high-performance liquid chromatography with electrochemical detection 575(1992)39
- Ogawa, H., see Ohmori, S. 566(1991)1
- Ogawa, H. K., see Kojima, K. 597(1992)323
- Ogawa, H. K., see Ueno, M. 597(1992)197
- O'Grady, B. V., see Richardson, D. E. 595(1992)155
- Oguma, T., see Imanari, T. 574(1992)142
- O'Hara, G., see Pleasance, S. 558(1991)155
- Ohara, K., Sano, M., Kondo, A. and Kato, I.
Two-dimensional mapping by high-performance liquid chromatography of pyridylamino oligosaccharides from various glycosphingolipids 586(1991)35
- Ohata, T., see Nozaki, O. 570(1991)1
- Ohba, S., Kidouchi, K., Katoh, T., Kibe, T., Kobayashi, M. and Wada, Y.
Automated determination of orotic acid, uracil and pseudouridine in urine by high-performance liquid chromatography with column switching 568(1991)325
- Ohba, Y., see Nozaki, O. 570(1991)1
- Ohga, T., see Kawakami, Y. 576(1992)63
- Ohhashi, T., Moritani, C., Andoh, H., Satoh, S., Ohmori, S., Lottspeich, F. and Ikeda, M.
Preparative high-yield electroelution of proteins after separation by sodium dodecyl sulphate-polyacrylamide gel electrophoresis and its application to analysis of amino acid sequences and to raise antibodies 585(1991)153
- Ohhira, S. and Matsui, H.
Comparison of sulphur-mode and tin-mode flame photometric detectors for the gas chromatographic determination of organotin compounds 566(1991)207
- Ohkawa, T., see Takamatsu, T. 584(1992)261
- Ohkubo, A., see Mashige, F. 565(1991)173
- Ohkubo, T., Kudo, M. and Sugawara, K.
Determination of ofloxacin in human serum by high-performance liquid chromatography with column switching 573(1992)289
- Ohkubo, T., see Seto, H. 570(1991)301
- Ohlson, S., see Wikström, M. 597(1992)83
- Ohmori, H., see Sakihama, N. 597(1992)147
- Ohmori, S., Nakata, K., Nishihara, K., Yamamoto, S., Kawase, M. and Tsuboi, S.
Sensitive determination of cystathionine and assays for cystathionine β - and γ -lyase, as well as cystathionine β -synthase, using high-performance liquid chromatography 574(1992)35
- Ohmori, S., Nose, Y., Ogawa, H., Tsuyama, K., Hirota, T., Goto, H., Yano, Y., Kondoh, Y., Nakata, K. and Tsuboi, S.
Fluorimetric and high-performance liquid chromatographic determination of D-lactate in biological samples 566(1991)1
- Ohmori, S., see Kawakami, Y. 576(1992)63
- Ohmori, S., see Ohhashi, T. 585(1991)153
- Ohmori, S., see Otsuka, M. 577(1992)215
- Ohno, M., see Murai, Y. 597(1992)201
- Ohno, M., see Nagatomo, A. 597(1992)411
- Ohno, M., see Ono, S. 597(1992)293
- Ohno, M., see Shimohigashi, Y. 597(1992)425
- Ohru, H., see Akasaka, K. 596(1992)197
- Ohru, S., see Kataoka, H. 578(1992)120
- Ohsawa, K., Kimura, M., Kurosawa-Ohsawa, K., Takahashi, M., Koyama, M., Abiko, Y., Hirahara, K., Matsuishi, T. and Tanaka, S.
Purification of sufficiently γ -carboxylated recombinant protein C and its derivatives. Calcium-dependent affinity shift in immunoaffinity and ion-exchange chromatography 597(1992)285
- Ohta, J., see Sugahara, K. 579(1992)318
- Ohta, J., see Yao, K. 581(1992)11
- Ohtaka, H., see Tanaka, S. 587(1991)364
- Ohuchi, K., see Watanabe, M. 597(1992)309
- Ohya, K., see Akimoto, K. 588(1991)165
- Ohyama, Y., see Ishibashi, M. 562(1991)613
- Öi, N., Kitahara, H. and Kira, R.
Direct separation of enantiomers by high-performance liquid chromatography on a new chiral ligand-exchange phase 592(1992)291
- Öi, N., see Nishioka, R. 565(1991)237
- Ojala-Karlsson, P. and Scheinin, M.
Improved high-performance liquid chromatographic method for the routine determination of unconjugated 3-methoxy-4-hydroxyphenylethylene glycol in human plasma using solid-phase extraction and electrochemical detection 565(1991)131
- Ojala-Karlsson, P., see Huupponen, R. 583(1992)270
- Oka, K., see Kanai, M. 567(1991)415
- Oka, T., see Hashimoto, A. 582(1992)41
- Okada, T.
Secondary equilibrium size-exclusion chromatography of ions with polymeric mobile phase additives 586(1991)277
- Okamoto, J., Hirai, E., Kitagawa, T. and Matsubara, T.
Capillary gas chromatographic method for the determination of the thromboxane A_2 receptor antagonist S-1452 and its metabolites in human plasma 583(1992)45
- Okamoto, J., Hirai, E., Suzuki, Y., Fujimoto, K., Kitagawa, T. and Matsubara, T.
Capillary gas chromatographic method for the determination of the thromboxane A_2 receptor antagonist S-1452 and its metabolites in human urine 583(1992)53

- Okamoto, M.
Preparation and evaluation of octadecyl- or phenylpropyl-treated porous glass for the high-performance liquid chromatographic analysis of bufadienolides in *Bufo venenum* 552(1991)381
- Okamoto, M. and Nakazawa, H.
Reversal of elution order during direct enantiomeric separation of pyriproxyfen on a cellulose-based chiral stationary phase 588(1991)177
- Okamoto, M., Nobuhara, K. and Jinno, K.
Effect of silanol groups on heat-treated silicas by calcination and retention behaviour in high-performance liquid chromatography 556(1991)407
- Okamoto, M., see Nakashima, K. 584(1992)275
- Okamoto, Y., see Honda, S. 590(1992)364
- Okazaki, O., Aoki, H. and Hokusui, H.
High-performance liquid chromatographic determination of (S)-(-)-ofloxacin and its metabolites in serum and urine using a solid-phase clean-up 563(1991)313
- Oke, A., see Cammack, J. 565(1991)529
- O'Kennedy, R., see Egan, D. A. 582(1992)137
- Okerholm, R. A., see Chan, K. Y. 571(1991)291
- Okerholm, R. A., see Coutant, J. E. 570(1991)139
- Oksman-Caldentey, K.-M., see Auriola, S. 562(1991)737
- Okuda, T., Nishida, M., Sameshima, I., Kyoyama, K., Hiramatsu, K., Takehara, Y. and Kohriyama, K.
Determination of atropine in biological specimens by high-performance liquid chromatography 567(1991)141
- Okumura, M., see Aoki, I. 571(1991)283
- Okuno, K., see Uda, T. 586(1991)131
- Okutani, R., Itoh, Y., Hirata, H., Kasahara, T., Mukaida, N. and Kawai, T.
Simple and high-yield purification of urine protein 1 using immunoaffinity chromatography: evidence for the identity of urine protein 1 and human Clara cell 10-kilodalton protein 577(1992)25
- Okuyama, T., see Isobe, T. 588(1991)115
- Okuyama, T., see Tomita, M. 566(1991)239
- Okuyama, T., see Tomita, M. 571(1991)324
- Oldiges, K., see Cole, R. O. 557(1991)113
- Olejnik, O., see Musson, D. G. 565(1991)89
- Olin, Å., see Johansson, K. 598(1992)105
- Oliveros, L., Minguillón, C., Desmazières, B. and Desbène, P.-L.
Silica-bonded chiral stationary phases with structurally simple π -donor chiral selectors for high-performance liquid chromatography 589(1992)53
- Oliw, E. H.
Enantioselective separation of some polyunsaturated epoxy fatty acids by high-performance liquid chromatography on a cellulose phenylcarbamate (Chiralcel OC) stationary phase 583(1992)231
- Ollagnier, M., see Gaillard, Y. 577(1992)171
- Olley, P. M., see Shi, Q. 572(1991)59
- Olliff, C. J., see Monger, L. S. 577(1992)239
- Olliff, C. J., see Monger, L. S. 595(1992)125
- Olsen, B. A., Stafford, J. D. and Reed, D. E.
Determination of dirithromycin purity and related substances by high-performance liquid chromatography 594(1992)203
- Olsen, I., see Brondz, I. 576(1992)328
- Olsen, I., see Brondz, I. 598(1992)309
- Olsson, A., see Pettersson, K.-J. 563(1991)414
- Olsson, N. U. and Kaufmann, P.
Optimized method for the determination of 1,2-diacyl-*sn*-glycero-3-phosphocholine and 1,2-diacyl-*sn*-glycero-3-phosphoethanolamine molecular species by enzymatic hydrolysis and gas chromatography 600(1992)257
- Olucha, J. C., see Calull, M. 589(1992)151
- Ombaba, J. M. and Barry, E. F.
Determination of organotin species by capillary gas chromatography with alternating current plasma emission detection 598(1992)97
- Omi, T., Kamesaki, T., Kajii, E. and Ikemoto, S.
Method for detecting the lectin activity of *Momordica charantia* transferred from micro two-dimensional electrophoretic gel on to nitrocellulose 570(1991)399
- Omori, A., see Hirokawa, T. 585(1991)297
- On, N. T., Damani, L. A. and Rance, D. J.
Quantitation of 6-(pyridin-3-yl)quinolin-2(1H)-one, a cardiac stimulant, and its N-oxide metabolite in dog plasma by high-performance liquid chromatography 574(1992)352
- Önderoglu, L., see Kutlar, F. 567(1991)359
- Ong, C.-N., Liao, L. S. and Ong, H. Y.
Separation of globins using free zone capillary electrophoresis 576(1992)346
- Ong, C. N., see Lee, B. L. 567(1991)71
- Ong, C. N., see Lee, B. L. 581(1992)41
- Ong, C. P., Ng, C. L., Lee, H. K. and Li, S. F. Y.
Separation of Dns-amino acids and vitamins by micellar electrokinetic chromatography 559(1991)537
- Ong, C. P., Ng, C. L., Lee, H. K. and Li, S. F. Y.
Determination of antihistamines in pharmaceuticals by capillary electrophoresis 588(1991)335
- Ong, C. P., Pang, S. F., Low, S. P., Lee, H. K. and Li, S. F. Y.
Migration behaviour of catechols and catecholamines in capillary electrophoresis 559(1991)529
- Ong, C. P., see Yik, Y. F. 589(1992)333
- Ong, H. Y., see Lee, B. L. 567(1991)71
- Ong, H. Y., see Lee, B. L. 581(1992)41
- Ong, H. Y., see Ong, C.-N. 576(1992)346
- Ono, I., see Hatada, K. 583(1992)116
- Ono, S., Kato, T., Lee, S., Aoyagi, H. and Ohno, M.
Interaction of amphipathic model lipopeptides with phospholipid bilayers 597(1992)293
- Ono, T., see Shibata, N. 566(1991)187
- Onodera, M., see Shibata, K. 570(1991)13
- Onodera, R., see Nagasawa, T. 566(1991)223
- Onodera, S.
Characterization and determination of organic compounds in the mutagenic XAD-2 extracts of drinking water 557(1991)413
- Onodera, S., see Moriyama, T. 571(1991)61
- Oparil, S., see Meng, Q. C. 579(1992)63
- Opocher, G., see Carpenè, G. 553(1991)201
- Ordóñez, D., see Alvarez-Bujidos, M. L. 578(1992)321
- Ordóñez, D., see Negro, A. 576(1992)135

- Ordunez, C., see Talmadge, K. W. 590(1992)83
- Orme, M. L.E., see Hoaksey, P. E. 566(1991)244
- Oros, F. J. and Davis, J. M.
Comparison of statistical theories of spot overlap in two-dimensional separations and verification of means for estimating the number of zones 591(1992)1
- O'Rourke, B. D., see Quinn Baumann, P. 573(1992)11
- Orthner, C. L., Highsmith, F. A., Tharakan, J., Madurawe, R. D., Morcol, T. and Velandar, W. H.
Comparison of the performance of immunosorbents prepared by site-directed or random coupling of monoclonal antibodies 558(1991)55
- Ortin, A., López-Pérez, M. J., Muiño Blanco, M. T. and Cebrian Perez, J. A.
Extraction of xanthine oxidase from milk by counter-current distribution in an aqueous two-phase system 558(1991)357
- Ortiz, A., see Alvarez-Bujidos, M. L. 578(1992)321
- Ortiz, A. I., see Negro, A. 576(1992)135
- Örvell, C., see Welling, G. W. 599(1992)157
- Orwar, O., Folestad, S., Einarsson, S., Andiné, P. and Sandberg, M.
Automated determination of neuroactive acidic sulphur-containing amino acids and γ -glutamyl peptides using liquid chromatography with fluorescence and electrochemical detection 566(1991)39
- Osawa, T., see Konami, Y. 597(1992)213
- Osawa, T., see Yamamoto, K. 597(1992)221
- O'Shannessy, D. J., see Morehead, H. W. 587(1991)171
- O'Shea, T. J., Greenhagen, R. D., Lunte, S. M., Lunte, C. E., Smyth, M. R., Radzik, D. M. and Watanabe, N.
Capillary electrophoresis with electrochemical detection employing an on-column Nafion joint 593(1992)305
- Oshima, T., Hirayama, F., Masuda, M., Maruta, T., Sagara, K., Mizutani, T., He, L.-Y., Tong, Y.-Y., Chen, Y.-H. and Itokawa, H.
Determination of sennoside A and B in the pharmaceutical preparation Otsuji-to using ion-pair high-performance liquid chromatography with column switching 585(1991)255
- Osman, S., see Sonnet, P. E. 586(1991)255
- Osselton, M. D., see Smith, R. M. 592(1992)85
- Osterloh, J. D., see Feldman, B. J. 594(1992)275
- Österlund, B. R.
Foreword 599(1992)1
- Ostrovský, I., see Soják, L. 557(1991)241
- Osuga, T., see Honda, A. 565(1991)53
- O'Sullivan, H., see Evans, M. B. 552(1991)169
- Ota, A., Ito, S. and Kawashima, Y.
Direct high-performance liquid chromatographic resolution of a novel benzothiazine Ca^{2+} antagonist and related compounds 593(1992)37
- Otagiri, M., see Iida, Y. 571(1991)277
- Ōtake, N., see Shinkai, H. 570(1991)203
- Otsuka, K., Kawahara, J., Tatekawa, K. and Terabe, S.
Chiral separations by micellar electrokinetic chromatography with sodium N-dodecanoyl-L-valinate 559(1991)209
- Otsuka, M. and Ohmori, S.
Simple and sensitive determination of diacetyl and acetoin in biological samples and alcoholic drinks by gas chromatography with electron-capture detection 577(1992)215
- Ottaviani, G., see Pichini, S. 577(1992)358
- Otvos, Jr., L., Urge, L. and Thurin, J.
Influence of different N- and O-linked carbohydrates on the retention times of synthetic peptides in reversed-phase high-performance liquid chromatography 599(1992)43
- Overmyer, S. K., see Rossi, D. T. 566(1991)257
- Oweimreen, G. A., see Ghodbane, S. 556(1991)317
- Oxford, J., see Bowers, G. D. 554(1991)175
- Oyama, T., see Wakui, Y. 575(1992)131
- Ozaki, H., see Koizumi, K. 595(1992)340
- Ozaki, M., see Hatada, K. 583(1992)116
- Ozand, P. T., see Leis, H. J. 578(1992)116
- Ozawa, H., see Miwa, T. 566(1991)163
- Ozawa, N., see Komatsu, S. 568(1991)460
- Özçimder, M., Louter, A. J. H., Lingeman, H., Voogt, W. H., Frei, R. W. and Bloemendal, M.
Determination of oxidized, reduced and protein-bound glutathione in eye lenses by high-performance liquid chromatography and electrochemical detection 570(1991)19
- Ožyhar, A. and Kiltz, H.-H.
High-resolution gel filtration of the ecdysteroid receptor-DNA complex —an alternative to the electrophoretic mobility shift assay 587(1991)11

P

- Pabel, B., see Mögele, R. 591(1992)165
- Pacáková, V. and Leclercq, P. A.
Gas chromatography-mass spectrometry and high-performance liquid chromatographic analyses of thermal degradation products of common plastics 555(1991)229
- Pacáková, V., see Böswart, J. 571(1991)19
- Pacáková, V., see Štulík, K. 552(1991)439
- Pacáková, V., see Švagrová, I. 563(1991)95
- Pacaud, E., see Ducrotoy, G. 566(1991)415
- Pace, M., see Mauri, P. L. 566(1991)327
- Pace, S., see Mancinelli, A. 575(1992)101
- Pachinger, A., Eisner, E., Begutter, H. and Klus, H.
Determination of a metabolite of the herbicide pyridate in drinking and groundwater using high-performance liquid chromatography with amperometric detection 558(1991)369
- Pacifici, R., see Pichini, S. 568(1991)267
- Pacifici, R., see Pichini, S. 577(1992)358

- Pacifici, R., see Pichini, S. 581(1992)143
 Pacifici, R., see Zuccaro, P. 567(1991)485
 Packham, A. J. and Fielden, P. R.
 Column switching for the high-performance liquid chromatographic analysis of polynuclear aromatic hydrocarbons in petroleum products 552(1991)575
 Paczkowski, D., Filipek, M., Mielniczuk, Z., Andrzejczak, J., Popławska, W. and Sitkiewicz, D.
 Simultaneous determination of mexiletine and four hydroxylated metabolites in human serum by high-performance liquid chromatography and its application to pharmacokinetic studies 573(1992)235
 Padfield, A. D., see Woollard, P. M. 562(1991)249
 Padovani, P. K., see Kapil, R. P. 577(1992)283
 Paesen, J., see Cachet, T. 600(1992)99
 Paetsch, P. R., Baker, G. B., Caffaro, L. E., Greenshaw, A. J., Rauw, G. A. and Coutts, R. T.
 Electron-capture gas chromatographic procedure for simultaneous determination of amphetamine and N-methylamphetamine 573(1992)313
 Pagani, R., see Terzuoli, L. 586(1991)153
 Pagano, M., see Dalet-Fumeron, V. 568(1991)55
 Page, K. M., Mallett, C. D. and Page, M. R.
 Determination of the novel hydroxymethyl glutaryl coenzyme A reductase inhibitor (RP 61969) and its dihydroxy acid hydrolysis product in human plasma by reversed-phase high-performance liquid chromatography 575(1992)319
 Page, M. R., see Page, K. M. 575(1992)319
 Pagès, M., see Abián, J. 554(1991)155
 Pagliari, R., Cottet-Emard, J. M. and Peyrin, L.
 Determination of free and conjugated normetanephrine and metanephrine in human plasma by high-performance liquid chromatography with electrochemical detection 563(1991)23
 Pagliuca, G., see Piretti, M. V. 585(1991)342
 Painuly, P. and Grill, C. M.
 Purification of erucic acid by preparative high-performance liquid chromatography and crystallization 590(1992)139
 Païs, M., see Cherton, J.-C. 566(1991)511
 Paisse, O., see Ventura, R. 562(1991)723
 Paleologo Oriundi, M., see Bagnati, R. 564(1991)493
 Palermo, K. M., see Pullen, R. H. 573(1992)49
 Palette, C., Cordonnier, P., Naline, E., Advenier, C. and Pays, M.
 High-performance liquid chromatographic method for the determination of the three main oxidative and 3-carboxylic antipyrine metabolites in human urine 563(1991)103
 Palette, C., see Moreau, J. 576(1992)103
 Palin, K. J., see McGee, J. P. 567(1991)282
 Paliwal, J. K., Gupta, R. C., Grover, P. K., Asthana, O. P. and Nityanand, S.
 Simultaneous determination of centbutindole and its hydroxy metabolite in serum by high-performance liquid chromatography 572(1991)219
 Palladino, D. E. H., House, R. M. and Cohen, K. A.
 Measurement of amino acid compositions of glycoprotein systems by gas-phase hydrolysis and reversed-phase high-performance liquid chromatography 599(1992)3
 Pals, G., see Bank, R. A. 571(1991)47
 Palumbo, G., see Carlucci, G. 564(1991)346
 Palylyk, E. L. and Jamali, F.
 Simultaneous determination of ketoprofen enantiomers and probenecid in plasma and urine by high-performance liquid chromatography 568(1991)187
 Pang, K. S., see Zimmerman, C. L. 563(1991)83
 Pang, S. F., see Ong, C. P. 559(1991)529
 Pannu, B. S., see Sud, D. 596(1992)281
 Paoletti, E., see Garibotto, G. 572(1991)11
 Pappis, M., see Swart, K. J. 593(1992)21
 Papich, M., see Boison, J. O. 576(1992)315
 Papp, E., see Bán, T. 593(1992)227
 Papp, E. A., Knupp, C. A. and Barbhuiya, R. H.
 High-performance liquid chromatographic assays for the quantification of amikacin in human plasma and urine 574(1992)93
 Paquet, N., see Toussi, A. 582(1992)87
 Parisi, A., see Tarditi, L. 599(1992)13
 Parisi, S., see Fossati, T. 577(1992)382
 Park, G., see Ally, A. 575(1992)19
 Park, J. H., Jang, M. D. and Shin, M. J.
 Solvatochromic hydrogen bond donor acidity of cyclodextrins and reversed-phase liquid chromatographic retention of small molecules on a β -cyclodextrin-bonded silica stationary phase 595(1992)45
 Park, J. H., Lee, J. E., Jang, M. D., Li, J.- and Carr, P. W.
 UNIFAC model as a heuristic guide for estimating retention in chromatography 586(1991)1
 Park, J.-S., see Kim, T.-J. 575(1992)295
 Park, K. H., Lee, M. H. and Lee, M. G.
 Simultaneous determination of haloperidol and its metabolite, reduced haloperidol, in plasma, blood, urine and tissue homogenates by high-performance liquid chromatography 572(1991)259
 Park, Y. H., see McCue, B. A. 565(1991)255
 Parker, C. E., see Deterding, L. J. 554(1991)329
 Parker, D. S., see Balcells, J. 575(1992)153
 Parker, R. J., see Anderson, L. W. 581(1992)195
 Parkin, J. E.
 Assay for thiomersal (thimerosal) with adaptation to the quantitation of total ethylmercury available in degraded samples 587(1991)329
 Paroni, R., see Fermo, I. 593(1992)171
 Parra, P., Limon, A., Ferre, S., Guix, T. and Jane, F.
 High-performance liquid chromatographic separation of caffeine, theophylline, theobromine and paraxanthine in rat brain and serum. 570(1991)185
 Parson, K. A., see Schafer, W. A. 587(1991)137
 Parviainen, M. T., see Pikkarainen, S. A. 577(1992)163
 Parvoli, G., see Sturaro, A. 590(1992)223
 Pascale, M. R., see Zezza, F. 593(1992)99
 Pasch, H., Krüger, H., Much, H. and Just, U.
 Analysis of benzyloxy-terminated poly(1,3,6-trioxocane)s by supercritical fluid chromatography 589(1992)295
 Pascual, P., Martinez-Lara, E., Bárcena, J. A. and Toribio, F.
 Direct assay of glutathione peroxidase activity using high-performance capillary electrophoresis 581(1992)49
 Pascual-Mostaza, C., see Pou-Clave, L. 563(1991)211

- Paseiro Losada, P., Paz Abuín, S., Vázquez Odériz, L., Simal Lozano, J. and Simal Gándara, J.
Quality control of cured epoxy resins. Determination of residual free monomers (*m*-xylylenediamine and bisphenol A diglycidyl ether) in the finished product 585(1991)75
- Passetti, G., see Mancinelli, A. 575(1992)101
- Pasta, P., see Carrea, G. 600(1992)320
- Pastera, J., see Nobilis, M. 581(1992)251
- Pastore, P., Boaretto, A., Lavagnini, I. and Diop, A.
UV spectrophotometric detection in the ion chromatographic analysis of alkali and alkaline earth metal and ammonium cations 591(1992)219
- Pasutto, F. M., see Alessi-Severini, S. 582(1992)173
- Pasutto, F. M., see Brocks, D. R. 581(1992)83
- Pasutto, F. M., see Carr, R. A. 566(1991)155
- Patel, B., see Lee, C. S. 559(1991)133
- Patel, H. B., King, D. N. and Jefferies, T. M.
Application of solute and mobile phase partition coefficient to describe solute retention in reversed-phase high-performance liquid chromatography 555(1991)21
- Patel, R. B., see Mangalan, S. 567(1991)498
- Pátfalusi, M., see Tomori, É. 578(1992)91
- Patil, S. F. and Lonkar, S. T.
Thermal desorption-gas chromatography for the determination of benzene, aniline, nitrobenzene and chlorobenzene in workplace air 600(1992)344
- Patin, E., see Rivat, C. 576(1992)71
- Patrick, J. E., see Alton, K. B. 579(1992)307
- Patrick, K. S., Nguyen, B. L. and McCallister, J. D.
Gas chromatographic-mass spectrometric determination of plasma selegiline using a deuterated internal standard 583(1992)254
- Patricot, M. C., see Chabraoui, L. 567(1991)299
- Patsalos, P. N., see Semba, J. 565(1991)357
- Patterson, E., see Chen, C. L. 583(1992)274
- Patterson, S. D.
Use of solvent selectivity optimization procedures for high-performance liquid chromatographic method development 592(1992)43
- Patterson, W. R., see Schwertner, H. A. 578(1992)297
- Patterson, Jr., D. G., see Kimata, K. 595(1992)77
- Patthy, M.
Gradient elution with shorter equilibration times in reversed-phase ion-pair chromatography 592(1992)143
- Pattichis, K., see Kajbaf, M. 575(1992)75
- Patrick, M., see Blagbrough, I. S. 578(1992)251
- Pattyn, G., Van Oosterom, A. T., De Bruijn, E. A. and Tjaden, U. R.
Determination of the new anticancer agent KW 2149, 7-N-[2-((2-(γ -L-glutamylamino)ethyl)dithio)ethyl]mitomycin C, an analogue of mitomycin C 564(1991)352
- Patzelová, V., see Pospíšil, J. 574(1992)71
- Paulson, J. and Lindberg, C.
Increasing thermospray response for cortisol by derivatization 554(1991)149
- Paulson, J., see Lindberg, C. 554(1991)215
- Paulus, A., see Manz, A. 593(1992)253
- Paulussen, R. J. A., see Ploum, M. E. 564(1991)413
- Pavelka, R., see Johansson, I. M. 559(1991)515
- Pawliszyn, J. and Wu, J.
Moving boundary capillary electrophoresis with concentration gradient detection 559(1991)111
- Pawliszyn, J., see McDonnell, T. 559(1991)489
- Payan, E., Jouzeau, J. Y., Lapique, F., Muller, N., Payan, J. P., Gegout, P., Bertin, P. and Netter, P.
Assay of synovial fluid hyaluronic acid using high-performance liquid chromatography of hyaluronidase digests 566(1991)9
- Payan, J. P., see Payan, E. 566(1991)9
- Pays, M., see Houzé, P. 574(1992)305
- Pays, M., see Moreau, J. 576(1992)103
- Pays, M., see Palettc, C. 563(1991)103
- Paz Abuín, S., see Paseiro Losada, P. 585(1991)75
- Pazourek, J. and Chmelík, J.
Simple solution of velocity profiles of laminar flows in channels of various cross-sections used in field-flow fractionation 593(1992)357
- Peach, R. J., see Haynes, P. A. 581(1992)187
- Pearson, S. J. and Reynolds, G. P.
Determination of 3-hydroxykynurenine in human brain and plasma by high-performance liquid chromatography with electrochemical detection. Increased concentrations in hepatic encephalopathy 565(1991)436
- Peart, T. E., see Lee, H.-B. 594(1992)309
- Péchar, M.-R., see Aubert, M.-C. 557(1991)47
- Péchar, M.-R., see Lesellier, E. 557(1991)59
- Pechová, H., see Churáček, J. 557(1991)523
- Pedini, F., see Carpenè, G. 553(1991)201
- Pedrielli, F., see Blo, G. 556(1991)249
- Pedrosa, M. M., see Vicente, C. 553(1991)271
- Peeters, A., see Vree, T. B. 573(1992)168
- Peiffer, G., see Piovetti, L. 588(1991)99
- Peiró, J. M., see Balcells, J. 575(1992)153
- Peixoto-Menezes, K. M., see Köppel, C. 563(1991)73
- Pellegatti, M., Braggio, S., Sartori, S., Franceschetti, F. and Bolelli, G. F.
Validation of a high-performance liquid chromatographic-radioimmunoassay method for the determination of lacidipine in plasma 573(1992)105
- Pellegrino, S., Petrarulo, M., Testa, E. and Nicolotti, A.
Rapid high-performance liquid chromatographic determination of urinary N-(1-methylethyl)-N'-phenyl-1,4-benzenediamine in workers exposed to aromatic amines 592(1992)279
- Pellegrino, S., see Petrarulo, M. 579(1992)324
- Pellerin, F., see Rakotomanga, S. 570(1991)277
- Pelletier, M., see Maeder, G. 593(1992)9
- Pelloso, R., see Lucarelli, C. 573(1992)150
- Peloquin, C. A., James, G. T. and McCarthy, E.
Improved high-performance liquid chromatographic assay for the determination of ethionamide in serum 563(1991)472
- Pelzer, G., see Vandenbroeck, M. 564(1991)405
- Peng, C. T., Hua, R. L. and Maltby, D.
Prediction of retention indexes. IV. Chain branching in alkylbenzene isomers with C₁₀₋₁₃ alkyl chains identified in a scintillator solvent 589(1992)231

- Peng, C. T., Yang, Z. C. and Ding, S. F.
Prediction of retention indexes. II. Structure-retention index relationship on polar columns 586(1991)85
- Peng, C. T., Yang, Z. C. and Maltby, D.
Prediction of retention indexes. III. Silylated derivatives of polar compounds 586(1991)113
- Peng, S., Xiao-Quan, S., Yan, Z., Long-Zhu, J. and Wei-Bing, X.
Determination of dietary cadmium-induced metallothioneins in rabbit kidneys and cadmium in metallothioneins by anion-exchange high-performance liquid chromatography coupled with graphite furnace atomic absorption spectrometry 572(1991)73
- Pennanen, S., Auriola, S., Manninen, A. and Komulainen, H.
Identification of the main metabolites of 2-ethylhexanoic acid in rat urine using gas chromatography-mass spectrometry 568(1991)125
- Penner, F., see Molina, F. 563(1991)205
- Penther, J. M.
Analysis of alanine and aspartate aminotransferase isoforms in mustard (*Sinapis alba* L.) cotyledons 587(1991)101
- Pepper, D. S., see Hodson, B. A. 565(1991)416
- Perel, J. M., see Foglia, J. P. 572(1991)247
- Pérez, S., see Sánchez-Brunete, C. 552(1991)235
- Pérez-Mateo, M., see Esteban, A. 573(1992)121
- Perez-Souto, N., Lynch, R. J., Measures, G. and Hann, J. T.
Use of high-performance liquid chromatographic peak deconvolution and peak labelling to identify antiparasitic components in plant extracts 593(1992)209
- Perico, A., see Santoni, G. 581(1992)287
- Perico, A., see Triolo, A. 567(1991)105
- Perico, A., see Triolo, A. 568(1991)281
- Perkins, J. R., see Deterding, L. J. 554(1991)329
- Perkins, L. B., see Bushway, R. J. 587(1991)321
- Perlík, F., see Pospíšil, J. 574(1992)71
- Permanyer, J., see Rodríguez, P. 562(1991)547
- Perogordo, E., see Reuvers, T. 564(1991)477
- Perry, M. L., see Apffel, A. 554(1991)103
- Pershing, D. W., see Lindgren, E. R. 585(1991)353
- Persson, A., see Balmér, K. 553(1991)391
- Persson, B., see Fägerstam, L. G. 597(1992)397
- Persson, B.-A., see Balmér, K. 553(1991)391
- Persson, B.-A., see Balmér, K. 592(1992)331
- Persson, P., see Brunmark, P. 579(1992)350
- Pesek, J. J., Vidensek, M. A. and Miller, M.
Synthesis of chemically bonded liquid crystals for high-performance liquid chromatography. New phases via the organochlorosilane pathway 556(1991)373
- Pesek, J. J., see Jinno, K. 557(1991)459
- Pesonen, K.
Variation of hydrophobicity of human urinary epidermal growth factor 568(1991)226
- Pessi, A., Bianchi, E., Chiappinelli, L., Nardi, A. and Fanali, S.
Application of capillary zone electrophoresis to the characterization of multiple antigen peptides 557(1991)307
- Pestellini, V., see Triolo, A. 568(1991)281
- Peter, R., see Böcker, R. H. 568(1991)363
- Peters, T. J., see Guo, R. 566(1991)383
- Peterson, J. C., Estiva, E. C., Lyttle, D. S. and Harris, R. M.
High-performance liquid chromatographic determination of 4,4'-methylenedianiline in human urine 564(1991)205
- Petersson, G., see Löfgren, L. 591(1992)358
- Petersson, G., see Strömvall, A.-M. 589(1992)385
- Petersson, P., see Johnson, D. F. 594(1992)283
- Petit, J., see Andriollo, O. 575(1992)301
- Petitjean, O., see Tod, M. 575(1992)171
- Petrarulo, M., Pellegrino, S. and Testa, E.
High-performance liquid chromatographic microassay for methyl ethyl ketone in urine as the 2,4-dinitrophenyl-hydrazone derivative 579(1992)324
- Petrarulo, M., see Pellegrino, S. 592(1992)279
- Petrauskas, A. A. and Švedas, V. K.
Hydrophobicity of β -lactam antibiotics. Explanation and prediction of their behaviour in various partitioning solvent systems and reversed-phase chromatography 585(1991)3
- Petrenko, V. V., see Larionov, O. G. 552(1991)31
- Petro, M., Križanovič, K. and Ďulák, K.
High-performance liquid chromatography of isopropylphenols 595(1992)337
- Petrukhin, O. M., see Timerbaev, A. R. 555(1991)163
- Petsev, N., see Gavrilova, T. B. 552(1991)179
- Pettersson, C., see Heldin, E. 585(1991)35
- Pettersson, C., see Heldin, E. 592(1992)339
- Pettersson, C., see Marle, I. 586(1991)233
- Pettersson, G., see Marle, I. 586(1991)233
- Pettersson, K.-J. and Nilsson, L. B.
Determination of dextropropoxyphene and norpropoxyphene in plasma by high-performance liquid chromatography 581(1992)161
- Pettersson, K.-J. and Olsson, A.
Liquid chromatographic determination of the enantiomers of ibuprofen in plasma using a chiral AGP column 563(1991)414
- Pettigrew, A., see Allen, E. 578(1992)175
- Pévet, P., see Raynaud, F. 564(1991)103
- Peyrin, L., see Pagliari, R. 563(1991)23
- Pfau, W. and Phillips, D. H.
Improved reversed-phase high-performance liquid chromatographic separation of ^{32}P -labelled nucleoside 3',5'-bisphosphate adducts of polycyclic aromatic hydrocarbons 570(1991)65
- Pfeffer, P. E., see Sonnet, P. E. 586(1991)255
- Pfeffer, W. D. and Yeung, E. S.
Electroosmotically driven electrochromatography of anions having similar electrophoretic mobilities by ion pairing 557(1991)125
- Pfister, M., see Straub, R. 585(1991)195
- Pfleiderer, B., see Unger, K. K. 556(1991)395
- Pfleiderer, W., see Werner, E. R. 570(1991)43
- Pham-Huy, C., see Bouygues-De Ferran, A. M. 585(1991)289
- Phemius, P., see Caruel, H. 594(1992)125
- Phillips, C. S. G., see Mills, T. A. 557(1991)495
- Phillips, D. H., see Pfau, W. 570(1991)65

- Phillips, D. J., Cheli, P. J., Dion, D. M., Hodgdon, H. L., Pomfret, A. M. and San Souci, B. R.
Schemes for efficient protein purification on a family of polymeric ion exchangers in glass columns 599(1992)239
- Phillips, J., see Bianchi, A. P. 557(1991)429
- Phillips, L. R., see Brown, N. D. 563(1991)466
- Phillips, M. and Greenberg, J.
Method for the collection and analysis of volatile compounds in the breath 564(1991)242
- Phillipson, J. D., see Abdulrahman, S. 562(1991)713
- Pianezzola, E., Bellotti, V., La Croix, R. and Strolin Benedetti, M.
Determination of cabergoline in plasma and urine by high-performance liquid chromatography with electrochemical detection 574(1992)170
- Pianezzola, E., see Breda, M. 578(1992)309
- Piazzini, S., see Grossi, G. 593(1992)217
- Picard, C., see Kamel, S. 574(1992)255
- Picard, C., see Maftouh, M. 562(1991)421
- Pichini, S., Altieri, I., Bacosi, A., Di Carlo, S., Zuccaro, P., Iannetti, P. and Pacifici, R.
High-performance liquid chromatographic-mass spectrometric assay of busulfan in serum and cerebrospinal fluid 581(1992)143
- Pichini, S., Altieri, I., Pacifici, R., Rosa, M., Ottaviani, G. and Zuccaro, P.
Simultaneous determination of cotinine and *trans*-3'-hydroxycotinine in human serum by high-performance liquid chromatography 577(1992)358
- Pichini, S., Altieri, I., Pacifici, R., Rosa, M. and Zuccaro, P.
Elimination of caffeine interference in high-performance liquid chromatographic determination of cotinine in human plasma 568(1991)267
- Pichini, S., see Betto, P. 575(1992)117
- Pichini, S., see Betto, P. 584(1992)256
- Pichini, S., see Zuccaro, P. 567(1991)485
- Pichorner, H., Loew, S., Korori, S. A. A., Jessner, G. and Ebermann, R.
Fast and sensitive staining technique for catalase in polyacrylamide gel 594(1992)400
- Picó, Y., see Moltó, J. C. 555(1991)137
- Piepmeyer, E. H., see Hein, S. J. 557(1991)39
- Pieraccini, G., see Mariot, R. 576(1992)143
- Pieters, L. A. C., see Qimin, L. 562(1991)435
- Pietrogrande, M. C., Reschiglian, P., Dondi, F., Kahie, Y. D. and Bertolasi, V.
Correlations between high-performance liquid chromatographic retention, X-ray structural and ¹³C NMR spectroscopic data of flavonoid compounds 592(1992)65
- Pietrzyk, D. J., see Yuan, D. 557(1991)315
- Pietta, P., Bruno, A., Mauri, P. and Rava, A.
Separation of flavonol-2-O-glycosides from *Calendula officinalis* and *Sambucus nigra* by high-performance liquid and micellar electrokinetic capillary chromatography 593(1992)165
- Pietta, P., Gardana, C., Mauri, P. and Zecca, L.
High-performance liquid chromatographic analysis of flavonol glycosides of *Solidago virgaurea* 558(1991)296
- Pietta, P., Mauri, P., Bruno, A., Rava, A., Manera, E. and Ceva, P.
Identification of flavonoids from *Ginkgo biloba* L., *Anthemis nobilis* L. and *Equisetum arvense* L. by high-performance liquid chromatography with diode-array UV detection 553(1991)223
- Pietta, P. G., see Mauri, P. L. 566(1991)327
- Pikkarainen, S. A. and Parviainen, M. T.
Determination of retinyl palmitate and total vitamin A content in liver and liver-based ready-to-eat foods 577(1992)163
- Pilorz, K., see Supryniewicz, Z. 552(1991)463
- Pimenov, A. M., see Grune, T. 563(1991)53
- Pimenov, A. M., see Ziegler, M. 563(1991)172
- Pinatel, H., see Dumas, L. 570(1991)89
- Pini, L. A., see Zuccaro, P. 567(1991)485
- Pintér-Szakács, M., see Fábrián, V. 553(1991)87
- Pinto, N. G., see Jen, S. C. D. 590(1992)3
- Piotrowski, E. G., see Brewster, J. D. 585(1991)213
- Piovetto, L., Deffo, P., Valls, R. and Peiffer, G.
Determination of sterols and diterpenoids from brown algae (Cystoseiraceae) 588(1991)99
- Piquion, J., see Lemque, R. 553(1991)165
- Piretti, M. V. and Pagliuca, G.
Regeneration of column activity after the gas chromatographic separation of membrane lipids on thermostable SE-52 phase 585(1991)342
- Pirkle, W. H.
Unusual effect of temperature on the retention of enantiomers on a chiral column 558(1991)1
- Pirkle, W. H. and Burke, J. A.
Separation of the enantiomers of the 3,5-dinitrobenzamide derivatives of α -amino phosphonates on four chiral stationary phases 598(1992)159
- Pirkle, W. H. and Burke, III, J. A.
Chiral stationary phase designed for β -blockers 557(1991)173
- Pirkle, W. H., Chang, J.-P. and Burke, III, J. A.
Contribution of specific hydrophobic interactions to chiral recognition 598(1992)1
- Pirkle, W. H. and Welch, C. J.
Effect of superfluous remote polar functionality on chiral recognition 589(1992)45
- Piryns, H., see Hamoir, T. 589(1992)31
- Piš, J. and Harmatha, J.
Phenylboronic acid as a versatile derivatization agent for chromatography of ecdysteroids 596(1992)271
- Pisaříková, A., see Tvřická, E. 563(1991)188
- Pitotti, A., see Maurich, V. 566(1991)453
- Pittalis, S., Martinez di Montemuros, F., Tavazzi, D. and Fiorelli, G.
Rapid isolation of glucose-6-phosphate dehydrogenase from human erythrocytes by combined affinity and anion-exchange chromatography for biochemical characterization of variants 573(1992)29
- Pittman, K. A., see Franc, J. E. 570(1991)129
- Pittman, K. A., see Janiszewski, J. S. 567(1991)274
- Pittner, F., see Ferris, R. 577(1992)251
- Pitts, N. I., see Ganhao, M. F. 564(1991)55

- Pivarnik, P. E., see Amari, J. V. 590(1992)153
- Planchon, C., see Mardens, Y. 577(1992)341
- Plank, C., see Rizzi, A. M. 557(1991)199
- Plantinga, W. J., Van Toorn, W. G. and Van der Stégen, G. H. D.
Determination of 3-chloropropane-1,2-diol in liquid hydrolysed vegetable proteins by capillary gas chromatography with flame ionization detection 555(1991)311
- Platonova, N. P., see Larionov, O. G. 552(1991)31
- Platt, D., see Rieck, W. 578(1992)259
- Platt, K. L., see Oesch, F. 582(1992)232
- Plaum, T., see Sturm, G. 562(1991)351
- Pleasance, S., Blay, P., Quilliam, M. A. and O'Hara, G.
Determination of sulfonamides by liquid chromatography, ultraviolet diode array detection and ion-spray tandem mass spectrometry with application to cultured salmon flesh 558(1991)155
- Pleasance, S., Thibault, P. and Kelly, J.
Comparison of liquid-junction and coaxial interfaces for capillary electrophoresis-mass spectrometry with application to compounds of concern to the aquaculture industry 591(1992)325
- Plicka, J., see Šmídl, P. 598(1992)15
- Ploum, M. E., Haasnoot, W., Paulussen, R. J. A., Van Bruchem, G. D., Hamers, A. R. M., Schilt, R. and Huf, F. A.
Test strip enzyme immunoassays and the fast screening of nortestosterone and clenbuterol residues in urine samples at the parts per billion level 564(1991)413
- Plzák, Z., see Grüner, B. 588(1991)201
- Plzák, Z., see Grüner, B. 595(1992)169
- Pochobradsky, M. G., see Gaspari, F. 577(1992)123
- Podrazký, V., see Štovičková, J. 575(1992)229
- Pognat, J. F., see Andriollo, O. 575(1992)301
- Poláková, L., see Ryska, M. 562(1991)289
- Polanuer, B., Sholin, A., Demina, N. and Rumiantseva, N.
Determination of glutamine, glutamic acid and pyroglutamic acids using high-performance liquid chromatography on dynamically modified silica 594(1992)173
- Polanuer, B. M.
Direct aqueous injection gas chromatography on a potassium fluoride crystal hydrate-containing sorbent. Determination of volatile organic solvents in the fermentation broth of *Clostridium* strains 596(1992)138
- Poláček, M., Gaš, B., Hirokawa, T. and Vacík, J.
Determination of limiting ionic mobilities and dissociation constants of some local anaesthetics 596(1992)265
- Polesello, A., see Forni, E. 593(1992)177
- Polesello, A., see Rizzolo, A. 553(1991)187
- Polettini, A., Ricossa, M. C., Groppi, A. and Montagna, M.
Determination of clenbuterol in urine as its cyclic boronate derivative by gas chromatography-mass spectrometry 564(1991)529
- Poli, G., see Chiari, M. 598(1992)287
- Polišenská, M., see Zahradníčková, H. 555(1991)239
- Pollack, G. M. and Kawagoe, J. L.
Determination of methanol in whole blood by capillary gas chromatography with direct on-column injection 570(1991)406
- Pollak, V. A., Doelemeyer, A., Winkler, W. and Schulze-Clewing, J.
Important design features of a system for the densitometric analysis of two-dimensional flat-bed separations 596(1992)241
- Pollitt, R. J., see Lowes, S. 577(1992)205
- Polman, C. H., see Van der Horst, A. 574(1992)166
- Polo-Diez, L. M., see Quintanilla-López, J. E. 591(1992)303
- Poma, R., see Magni, F. 573(1992)127
- Pomerantz, S. C., see Takeda, N. 562(1991)225
- Pomfret, A. M., see Phillips, D. J. 599(1992)239
- Pommerening, K., see Boeden, H.-F. 552(1991)389
- Pommery, J., see Trivier, J. M. 579(1992)269
- Pommier, F., see Sioufi, A. 571(1991)87
- Pomponio, G., see Corvetta, A. 566(1991)481
- Ponthus, C., see Maftouh, M. 562(1991)421
- Ponti, F., see Betto, P. 563(1991)115
- Poole, C. F., see Kollie, T. O. 556(1991)457
- Poon, G. K., Jarman, M., McCague, R., Davies, J. H., Heeremans, C. E. M., Van der Hoeven, R. A. M., Niessen, W. M. A. and Van der Greef, J.
Identification of 4-hydroxyandrost-4-ene-3,17-dione metabolites in prostatic cancer patients by liquid chromatography-mass spectrometry 576(1992)235
- Poon, G. K., Jarman, M., Rowlands, M. G., Dowsett, M. and Firth, J.
Determination of 4-hydroxyandrost-4-ene-3,17-dione metabolism in breast cancer patients using high-performance liquid chromatography-mass spectrometry 565(1991)75
- Poon, G. K., McCague, R., Griggs, L. J., Jarman, M. and Lewis, I. A. S.
Characterisation of metabolites of 3-ethyl-3-(4-pyridyl)-piperidine-2,6-dione, a potential breast cancer drug 572(1991)143
- Poon, G. K., see Van Bakergem, E. 598(1992)189
- Popineau, Y., see Courcoux, P. 596(1992)225
- Popławska, W., see Paczkowski, D. 573(1992)235
- Popp, M., see Seitz, U. 559(1991)499
- Poppe, H.
New approach for calculating ideal chromatograms from arbitrary composite distribution isotherms 556(1991)95
- Poppe, H., see Bruin, G. J. M. 559(1991)163
- Porter, N. P., see Escott, R. E. A. 554(1991)281
- Porterfield, S. P., see Hendrich, C. E. 577(1992)19
- Porthault, M., see Fathi, M. 563(1991)356
- Pospíšil, J., Patzelová, V., Perlík, F. and Máca, B.
Determination of ethyl biscoumacetate and its metabolite 7-hydroxyethyl biscoumacetate in human serum by high-performance liquid chromatography and mass spectrometry 574(1992)71
- Post, M. H., see Van der Horst, F. A. L. 567(1991)161
- Postaire, M., see Bouygues-De Ferran, A. M. 585(1991)289
- Potschka, M.
Interfacial effects in size-exclusion chromatography of latex 587(1991)276

- Potter, C., see McGee, J. P. 567(1991)282
- Potter, J. M., see Chen, Y. 574(1992)361
- Pou-Clave, L., Campos-Barreda, F. and Pascual-Mostaza, C.
Determination of ciprofloxacin in human serum by liquid chromatography 563(1991)211
- Poulou, M., see Tod, M. 594(1992)386
- Poulsen, B. J., see Jagodzinski, J. J. 591(1992)89
- Poulsen, O. M., see Hansen, Å. M. 578(1992)85
- Pourfarzam, M. and Bartlett, K.
Synthesis, characterisation and high-performance liquid chromatography of C₆-C₁₆ dicarboxyl-*mono-coenzyme A* and *-mono-carnitine esters* 570(1991)253
- Pourrat, H., see Barthomeuf, C. 568(1991)451
- Poüs, C., see Bories, C. 579(1992)143
- Pousset, J.-L., see Rey, J.-P. 587(1991)314
- Pousset, J.-L., see Rey, J.-P. 596(1992)276
- Pouwelse, A. V., see Van Zijtveld, J. 600(1992)211
- Povel, F. R., see Tuinstra, L. G. M. T. 552(1991)259
- Pozza, G., see Magni, F. 573(1992)127
- Prabhasankar, P., see Ragupathi, G. 574(1992)267
- Prakash, C., see Edeki, T. 577(1992)190
- Pramanik, B. N., Tsarbopoulos, A., Labdon, J. E., Trotta, P. P. and Nagabhushan, T. L.
Structural analysis of biologically active peptides and recombinant proteins and their modified counterparts by mass spectrometry 562(1991)377
- Prapamontol, T. and Stevenson, D.
Rapid method for the determination of organochlorine pesticides in milk 552(1991)249
- Pras, M., see Kaplan, B. 573(1992)17
- Preece, N. E., Forrow, S., Ghatineh, S., Langley, G. J. and Timbrell, J. A.
Determination of hydrazine in biofluids by capillary gas chromatography with nitrogen-sensitive or mass spectrometric detection 573(1992)227
- Prescott, M. C., see Evershed, R. P. 590(1992)305
- Preston, H. D., see Van Bergen, C. A. 589(1992)109
- Price, C. P., see Scott, D. A. 573(1992)309
- Price, G. J.
Prediction of retention in gas-liquid chromatography using the UNIFAC group contribution method. III. Recent developments in UNIFAC 586(1991)297
- Price, G. J. and Dent, M. R.
Prediction of retention in gas-liquid chromatography using the UNIFAC group contribution method. II. Polymer stationary phases 585(1991)83
- Price, N. P. J., Firmin, J. L. and Gray, D. O.
Screening for amines by dansylation and automated high-performance liquid chromatography 598(1992)51
- Přikrylová, V., see Beran, M. 558(1991)265
- Pristach, J., see Suttner, J. 563(1991)184
- Promé, J. C., see Le Quééré, J. L. 562(1991)659
- Prošek, M., Drusany, I. and Golc-Wondra, A.
Quantitative two-dimensional thin-layer chromatography 553(1991)477
- Provost, L. R., see D'Agostino, P. A. 589(1992)287
- Provost, L. R., see D'Agostino, P. A. 598(1992)89
- Provost, L. R., see D'Agostino, P. A. 600(1992)267
- Pruñonosa, J., Obach, R., Diez-Cascón, A. and Gouesclou, L.
Determination of cicletanine enantiomers in plasma by high-performance capillary electrophoresis 574(1992)127
- Pruñonosa, J., Obach, R., Diez-Cascón, A. and Gouesclou, L.
Comparison of high-performance liquid chromatography and high-performance capillary electrophoresis for the determination of cicletanine in plasma 581(1992)219
- Prusík, Z., see Kašička, V. 569(1991)123
- Prusiková, M., see Taimr, L. 587(1991)355
- Przekop, M. A., see Kunsman, G. W. 568(1991)427
- Przyborowska, A., see Halket, J. M. 562(1991)279
- Przybylski, M., see Svoboda, M. 562(1991)403
- Przysiecki, C. T., see Lehman, E. D. 574(1992)225
- Pudleiner, P., Kapás, M. and Vereczkey, L.
High-performance liquid chromatographic method for the determination of RGH-5702 in plasma samples 553(1991)29
- Puhlmann, A., Dülffer, T. and Kobold, U.
Multidimensional high-performance liquid chromatography on Pinkerton ISRP and RP18 columns: direct serum injection to quantify creatinine 581(1992)129
- Pulcinelli, F. M., Gazzaniga, P. P. and Salganicoff, L.
Use of Zn-pyrophosphatase in the high-performance liquid chromatographic analysis of cell extracts containing ³²P-labelled inositol phosphates 575(1992)51
- Pullen, F. S., see Cole, S. M. 562(1991)67
- Pullen, R. H. and Fatmi, A. A.
Determination of fluvoxamine in human plasma by high-performance liquid chromatography with fluorescence detection 574(1992)101
- Pullen, R. H., Palermo, K. M. and Curtis, M. A.
Determination of an antipsychotic agent (ICI 204,636) and its 7-hydroxy metabolite in human plasma by high-performance liquid chromatography and gas chromatography-mass spectrometry 573(1992)49
- Purcell, A. W., Aguilar, M. I. and Hearn, M. T. W.
High-performance liquid chromatography of amino acids, peptides and proteins. CXV. Thermodynamic behaviour of peptides in reversed-phase chromatography 593(1992)103
- Purdy, W. C., see Carpenter, A. 573(1992)132
- Purdy, W. C., see Tadey, T. 583(1992)111
- Purnell, H.
A practical guide to the care, maintenance and troubleshooting of capillary gas chromatographic systems. (by D. Rood) (Book Review) 585(1991)379
- Purnell, J. H. and Wattan, M. H.
Theoretical and experimental comparison of serially linked and mixed-packing gas-liquid chromatography columns 555(1991)173
- Purvis, D. R., see McCreath, G. E. 597(1992)189
- Püttmann, W.
Thermodesorption-gas chromatography-mass spectrometric analysis of biological materials for potential molecular precursors of the constituents of the crude oils 552(1991)325

Q

- Qian, M. and Gallo, J. M.
High-performance liquid chromatographic determination of the calcium channel blocker nimodipine in monkey plasma 578(1992)316
- Qimin, L., Van den Heuvel, H., Delorenzo, O., Corthout, J., Pieters, L. A. C., Vlietinck, A. J. and Claeys, M.
Mass spectral characterization of C-glycosidic flavonoids isolated from a medicinal plant (*Passiflora incarnata*) 562(1991)435
- Quadri, A., see Brega, A. 553(1991)249
- Quadrini, G., see Brega, A. 553(1991)249
- Quaedflieg, P. J. L. M., see Coenen, A. J. J. M. 596(1992)59
- Quaglia, M. G., Fanali, S., Nardi, A., Rossi, C. and Ricci, M.
Separation of leucino-statins by capillary zone electrophoresis 593(1992)259
- Quan, C. P., see Frenz, J. 557(1991)289
- Quick, Jr., A. N., see Loy, G. L. 562(1991)169
- Quicke, J., see Rop, P. P. 573(1992)87
- Quilliam, M. A., see Pleasance, S. 558(1991)155
- Quinn Baumann, P., Ebenstein, D. B., O'Rourke, B. D. and Sreekumaran Nair, K.
High-performance liquid chromatographic technique for non-derivatized leucine purification: evidence for carbon isotope fractionation 573(1992)11
- Quintanilla-López, J. E., Lebrón-Aguilar, R. and Polo-Díez, L. M.
Comparative study of clean-up and fractionation methods for the determination of organochlorine pesticides in lipids by gas chromatography 591(1992)303

R

- Raatikainen, O.
High-performance liquid chromatography of heptaene polyenes: assay of heptaene produced by *Streptomyces griseoviridis* 588(1991)356
- Raatikainen, O., Auriola, S. and Tuomisto, J.
Identification of aromatic moieties and mycosamine in antifungal heptaenes with high-performance liquid chromatography, high-performance liquid chromatography-mass spectrometry and gas chromatography-mass spectrometry 585(1991)247
- Rabillon, J., see Dandeu, J.-P. 599(1992)105
- Rabino, R., see Castagnola, M. 572(1991)51
- Racine, M., see Lemque, R. 553(1991)165
- Raddatz, W. D., see Lewis, J. A. 592(1992)183
- Rader, D. J., see Shibusawa, Y. 596(1992)118
- Radić, B., see Beker, D. 570(1991)441
- Radler, S. and Blaschke, G.
Enantioselective determination of dimethindene in urine after oral administration of racemic dimethindene 567(1991)229
- Radzik, D. M., see O'Shea, T. J. 593(1992)305
- Ræder, M., see Holthe, M. R. 579(1992)376
- Ragupathi, G., Prabhasankar, P., Chandra Sekharan, P., Annapoorani, K. S. and Damodaran, C.
Novel solid-state fluorodensitometric method for the determination of haptens in protein-hapten conjugates. Demonstration with a toxic glycoside of *Cleistanthus collinus* 574(1992)267
- Rahal, S., see Tazerouti, A. 596(1992)132
- Rahko, E.-L., see Härmälä, P. 593(1992)329
- Raina, A. K.
Chromatography and isolation of insect hormones and pheromones (edited by A. R. McCaffery and I. D. Wilson) (Book Review) 600(1992)373
- Rákos, L., see Moberg, C. 585(1991)309
- Rakotomanga, S., Baillet, A., Pellerin, F. and Baylocq-Ferrier, D.
Simultaneous determination of gluconolactone, galactonolactone and galactitol in urine by reversed-phase liquid chromatography: application to galactosemia 570(1991)277
- Ramanadham, S., see Turk, J. 575(1992)183
- Rambourg, M., see Del Cont Bernard, A. 574(1992)365
- Rambourg, M., see Royer-Morrot, M. J. 581(1992)297
- Ramis, I., Roselló-Catafau, J. and Gelpí, E.
In vivo transformation of arachidonic acid into 12-hydroxy-5,8,10,14-eicosatetraenoic acid by human nasal mucosa 575(1992)143
- Ramis, I., Roselló-Catafau, J., Gómez, G., Zabay, J. M., Fernández Cruz, E. and Gelpí, E.
Cyclooxygenase and lipooxygenase arachidonic acid metabolism by monocytes from human immune deficiency virus-infected drug users 557(1991)507
- Ramis Ramos, G., see Bonet Domingo, E. 582(1992)189
- Ramjit, H. G., see Lehman, E. D. 574(1992)225
- Ramos Carrillo, A., see Rotllán Pascual, P. 563(1991)37
- Rampino, N. J. and Chrambach, A.
DNA electrophoresis in uncross-linked polyacrylamide solution, studied by epifluorescence microscopy 596(1992)141
- Ramsey, R. S., see Griest, W. H. 600(1992)273
- Ramus, T. L., see Thomas, L. C. 586(1991)309
- Ramzan, I.
High-performance liquid chromatographic assay for laudanoline in biological fluids and tissue for neurotoxic studies in rats 565(1991)465
- Rance, D. J., see Cole, S. M. 562(1991)67
- Rance, D. J., see On, N. T. 574(1992)352
- Rang, S. A., see Berezkin, V. G. 552(1991)313
- Rao, R. N., see Husain, S. 596(1992)127

- Rao, R. N., see Husain, S. 600(1992)316
- Rapoport, S. I., see Deutsch, J. 579(1992)93
- Rapp, H., see Greischel, A. 568(1991)246
- Rappuoli, P., see Scapol, L. 600(1992)235
- Rasmussen, K. E., see Agasøster, T. 564(1991)171
- Rasmussen, K. E., see Agasøster, T. 570(1991)99
- Rasmussen, K. E., see Andresen, A. T. 575(1992)93
- Rasmussen, K. E., see Andresen, A. T. 582(1992)123
- Rasmussen, K. E., see Johansen, M. 573(1992)283
- Raspi, G., Lo Moro, A. and Spinetti, M.
Direct determination of kallikrein by high-performance liquid chromatography 593(1992)119
- Rastall, R. A., see Dutton, M. V. 587(1991)297
- Ratna, S., see Zimmerman, C. L. 563(1991)83
- Ratnaraj, N., see Semba, J. 565(1991)357
- Ratti, E., see Dalla Libera, L. 566(1991)427
- Raulin, F., see Do, L. 591(1992)297
- Raus, J., see Leyssens, L. 564(1991)515
- Rauw, G. A., see Paetsch, P. R. 573(1992)313
- Rava, A., see Pietta, P. 553(1991)223
- Rava, A., see Pietta, P. 593(1992)165
- Ravanetti, G. P., see Bonifaci, L. 585(1991)333
- Ravens, U., see Himmel, H. M. 568(1991)105
- Ravensberg, J. C., see Leenheers, L. H. 578(1992)189
- Ravenscroft, P. J., see Chen, Y. 574(1992)361
- Ravenscroft, P. J., see Taylor, P. J. 581(1992)152
- Raverdino, V.
On-line derivatization of eluted substances in dynamic high-performance liquid chromatography-mass spectrometry through the particle-beam interface 554(1991)125
- Raverdino, V., see Fathi, M. 563(1991)356
- Ravid, M., see Kaplan, B. 573(1992)17
- Ravis, W. R., see Shaikh, B. 571(1991)189
- Raviv, O., see Solomon, B. 597(1992)257
- Raynaud, F. and Pévet, P.
Determination of 5-methoxyindoles in pineal gland and plasma samples by high-performance liquid chromatography with electrochemical detection 564(1991)103
- Raza, G., see Jagodzinski, J. J. 591(1992)89
- Ré, M., see Barilero, I. 575(1992)275
- Read, R. W., see Black, R. M. 558(1991)393
- Read, R. W., see Black, R. M. 558(1991)405
- Read, S. A., see Kaczvinsky, Jr., J. R. 575(1992)177
- Rechnitz, G. A., see Tanaka, S. 597(1992)443
- Recny, M. A., see Neidhardt, E. A. 590(1992)255
- Redden, P. R. and Huang, Y.-S.
Automated separation and quantitation of lipid fractions by high-performance liquid chromatography and mass detection 567(1991)21
- Redmond, J. W., see Haynes, P. A. 581(1992)187
- Redmond, J. W., see Haynes, P. A. 588(1991)107
- Reed, D. E., see Olsen, B. A. 594(1992)203
- Rees, L. W., see Robinson, P. R. 564(1991)147
- Reeuwijk, H. J. E. M., Tjaden, U. R. and Van der Greef, J.
Simultaneous determination of furosemide and amiloride in plasma using high-performance liquid chromatography with fluorescence detection 575(1992)269
- Regårdh, C. G., see Gabrielsson, M. 573(1992)265
- Regerat, F., see Barthomeuf, C. 568(1991)451
- Regnault, V., Rivat, C., Vallar, L., Geschier, C. and Stoltz, J. F.
Purification of biologically active human plasma transthyretin by dye-affinity chromatography: studies on dye leakage and possibility of heat treatment for virus inactivation 584(1992)93
- Regnault, V., Rivat, C., Vallar, L. and Stoltz, J.-F.
Dye-affinity purification of transthyretin from an unexploited by-product of human plasma chromatographic fractionation 576(1992)87
- Regnault, V., see Ubrich, N. 584(1992)17
- Regnier, F. E., see Towns, J. K. 599(1992)227
- Rehner, G., see Sander, S. 558(1991)115
- Reibnegger, G., see Werner, E. R. 570(1991)43
- Reichelová, V., Novotný, L. and Zima, D.
High-performance liquid chromatographic study of some biologically active analogues of arabinosylcytosine 588(1991)147
- Reichling, J., see Herweck, U. 600(1992)312
- Reichlmayr-Lais, A. M., see Eder, K. 588(1991)265
- Reichlmayr-Lais, A. M., see Eder, K. 598(1992)33
- Reid, R. G., see Taylor, R. B. 563(1991)451
- Reijenga, J. C.
Data compression in computerized signal processing for isocratic chromatography 585(1991)160
- Reijenga, J. C.
GCSIM: a gas-liquid chromatography simulator for educational purposes 588(1991)217
- Reilly, P. E. B., see Mason, S. R. 581(1992)267
- Reimer, G. J. and Suarez, A.
Development of a screening method for five sulfonamides in salmon muscle tissue using thin-layer chromatography 555(1991)315
- Reinacher, M., see Schuler, G. 587(1991)61
- Reiner, V., see Belotto, N. 572(1991)327
- Reinerink, E. J. M., Doorn, L., Jansen, E. H. J. M. and Van Iersel, A. A. J.
Measurement of enzyme activities of cytochrome P-450 isoenzymes by high-performance liquid chromatographic analysis of products 553(1991)233
- Reinhart, P., Harkrader, R., Wylie, R., Yewey, G. and Van Horne, K. C.
Sanguinarine levels in biological samples by high-performance liquid chromatography 570(1991)425
- Reinhoud, N. J., Tjaden, U. R., Irth, H. and Van der Greef, J.
Bioanalysis of some anthracyclines in human plasma by capillary electrophoresis with laser-induced fluorescence detection 574(1992)327
- Reiser, R. W., Barefoot, A. C., Dietrich, R. F., Fogiel, A. J., Johnson, W. R. and Scott, M. T.
Application of microcolumn liquid chromatography-continuous-flow fast atom bombardment mass spectrometry in environmental studies of sulfonylurea herbicides 554(1991)91
- Reiss, W. H., see Jones, L. A. 595(1992)209
- Reiss, W. H., see Jones, L. A. 595(1992)221
- Rejano, L., see Montaña, A. 594(1992)259

- Relling, M. V., see Evans, R. R. 578(1992)141
- Remelli, M., see Blo, G. 556(1991)249
- Rementer, S. W., see Kirkland, J. J. 593(1992)339
- Remien, J., see Šalamoun, J. 596(1992)43
- Remington, G., see Cheung, S. W. 564(1991)213
- Rommel, R. P., see Sinz, M. W. 571(1991)217
- Remuzzi, G., see Gaspari, F. 570(1991)435
- Ren, Z., see Xie, M. 555(1991)33
- Renken, A., see Andriamboavonjy, E. 587(1991)288
- Rennie, P. J., see Mallet, A. I. 562(1991)647
- Reschiglian, P., see Pietrogrande, M. C. 592(1992)65
- Resines, J. A., see Arin, M. J. 582(1992)13
- Ress, K. M., see Mayer, F. 572(1991)339
- Reuke, S., see Eisenbeiss, A. 589(1992)390
- Reusch, J., see Abou-Rebyeh, H. 566(1991)341
- Reusch, J., see Josić, D. 590(1992)59
- Reusch, J., see Oberdorfer, F. 552(1991)483
- Reutter, W., see Josić, D. 590(1992)59
- Reuvers, T., Perogordo, E. and Jiménez, R.
Rapid screening method for the determination of diethylstilbestrol in edible animal tissue by column liquid chromatography with electrochemical detection 564(1991)477
- Revill, A. T., Carr, M. R. and Rowland, S. J.
Use of oxidative degradation followed by capillary gas chromatography-mass spectrometry and multi-dimensional scaling analysis to fingerprint unresolved complex mixtures of hydrocarbons 589(1992)281
- Revol, A., see Chabraoui, L. 567(1991)299
- Rey, J.-P., Levesque, J., Pousset, J.-L. and Roblot, F.
Analytical and quantitative studies of californin and protopin in aerial part extracts of *Eschscholtzia californica* Cham. with high-performance liquid chromatography 587(1991)314
- Rey, J.-P., Levesque, J., Pousset, J.-L. and Roblot, F.
Analytical studies of isorhoeadine and rhoegenine in petal extracts of *Papaver rhoeas* L. using high-performance liquid chromatography 596(1992)276
- Reya, R., see Hlavačková, E. 595(1992)137
- Reynolds, G. D., see Milne, R. W. 565(1991)457
- Reynolds, G. P., see Pearson, S. J. 565(1991)436
- Reynolds, L. K., see Driskell, J. A. 568(1991)333
- Řezanka, T.
Practical manual on lipid analysis. A series of monographs. I. Fatty acids (by J. G. Alvarez and J. C. Touchstone) (Book Review) 575(1992)182
- Řezanka, T.
Analysis of sterol esters from alga and yeast by high-performance liquid chromatography and capillary gas chromatography-mass spectrometry with chemical ionization 598(1992)219
- Řezanka, T., Zlatkin, I. V., Viden, I., Slabova, O. I. and Nikitin, D. I.
Capillary gas chromatography-mass spectrometry of unusual and very long-chain fatty acids from soil oligotrophic bacteria 558(1991)215
- Řezanka, T., see Mareš, P. 568(1991)1
- Rhodes, C. G., see Jones, H. A. 570(1991)361
- Rhodes, G., see Boppana, V. K. 592(1992)317
- Rhodes, G. R., see Boppana, V. K. 593(1992)29
- Ribick, M., see Both, D. A. 585(1991)348
- Ricci, M., see Quaglia, M. G. 593(1992)259
- Ricciarelo, G., see Betto, P. 575(1992)117
- Ricciarelo, G., see Betto, P. 584(1992)256
- Ricciarelo, G., see Zuccaro, P. 567(1991)485
- Richard, H., see Morin, P. 586(1991)171
- Richard, J., see Sioufi, A. 565(1991)401
- Richard, M. J., Guiraud, P., Meo, J. and Favier, A.
High-performance liquid chromatographic separation of malondialdehyde-thiobarbituric acid adduct in biological materials (plasma and human cells) using a commercially available reagent 577(1992)9
- Richardson, D. E., Bremner, J. B. and O'Grady, B. V.
Quantitative analysis of total resin acids by high-performance liquid chromatography of their coumarin ester derivatives 595(1992)155
- Richardson, M., see Kim, H.-J. 593(1992)153
- Richardson, R. A., see Sutton, B. M. 581(1992)277
- Richert, L., see Ducrottoy, G. 566(1991)415
- Richli, U., see Badoud, R. 552(1991)345
- Rickenbacher, U., see Giard-Pasquier, C. 553(1991)117
- Ricks, C., see Burton, S. D. 571(1991)209
- Rico, I., see Halvorsen, K. 557(1991)501
- Ricossa, M. C., see Poletini, A. 564(1991)529
- Rieck, W. and Platt, D.
High-performance liquid chromatographic method for the determination of alprazolam in plasma using the column-switching technique 578(1992)259
- Riedel, K.-D., Wildfeuer, A., Laufen, H. and Zimmermann, T.
Equivalence of a high-performance liquid chromatographic assay and a bioassay of azithromycin in human serum samples 576(1992)358
- Riegel, M. R. and Ellis, P. P.
Determination of trifluorothymidine in the eye using high-performance liquid chromatography 568(1991)467
- Rieger, R., see Brown-Thomas, J. M. 568(1991)232
- Riekkola, M.-L., see Sirén, H. 590(1992)263
- Riester, R. L., Yan, C. and Martire, D. E.
Determination of the hold-up time and column outlet density for capillary supercritical fluid chromatography 588(1991)289
- Rigal, L., see Caruel, H. 558(1991)89
- Rigal, L., see Caruel, H. 594(1992)125
- Riggs, K. W., see Wright, M. R. 565(1991)225
- Righetti, P. G.
Advances in electrophoresis, Vol. 4 (edited by A. Chrambach, M. J. Dunn and B. J. Radola) (Book Review) 600(1992)375
- Righetti, P. G. and Deyl, Z.
Preface 569(1991)1
- Righetti, P. G., Gianazza, E., Bianchi-Bosisio, A., Sinha, P. and Köttgen, E.
Isoelectric focusing in immobilized pH gradients: applications in clinical chemistry and forensic analysis (Review) 569(1991)197
- Righetti, P. G., see Bianchi-Bosisio, A. 569(1991)243
- Righetti, P. G., see Chiari, M. 558(1991)285
- Righetti, P. G., see Chiari, M. 598(1992)287

- Righetti, P. G., see Chiari, M. 559(1991)119
- Righetti, P. G., see Gelfi, C. 598(1992)277
- Righetti, P. G., see Gianazza, E. 569(1991)43
- Righezza, M. and Chrétien, J. R.
Factor analysis and experimental design in high-performance liquid chromatography. XI. Factor analysis maps and chromatographic information 556(1991)169
- Rihter, B. D., see Zuk, M. M. 568(1991)437
- Rijks, J., see Rutten, G. 590(1992)271
- Rimmer, D. A. and Rose, M. E.
Some novel homochiral derivatizing agents for the gas chromatographic analysis of enantiomeric secondary alcohols 598(1992)251
- Rinke, L., see Meyer, H. H. D. 564(1991)551
- Rios, J. J., see Ruiz-Gutierrez, V. 575(1992)1
- Rishi, D. K., see Kim, I. S. 589(1991)177
- Risler, T., see Mayer, F. 572(1991)339
- Rissler, K. and Cramer, H.
Marked improvement of a substance P radioimmunoassay by reduction of ¹²⁵I-labelled [Tyr⁸]-substance P prepared by the chloramine-T method with mercaptoethanol and subsequent purification by reversed-phase liquid chromatography 564(1991)67
- Rissler, K., Friedrich, G. and Kneer, J.
Determination of the retinoid (*E*)-1,2,3,4-tetrahydro-1,1,4,4-tetramethyl-6-(1-methyl-2-phenylethenyl)-naphthalene and its phenolic metabolite in human plasma by reversed-phase high-performance liquid chromatography 565(1991)375
- Riva, E., Merati, R. and Cavenaghi, L.
High-performance liquid chromatographic determination of rifapentine and its metabolite in human plasma by direct injection into a shielded hydrophobic phase column 553(1991)35
- Rivat, C., Sertillanges, P., Patin, E. and Stoltz, J.-F.
Single-step method for purification of human transferrin from a by-product of chromatographic fractionation of plasma 576(1992)71
- Rivat, C., see Regnault, V. 576(1992)87
- Rivat, C., see Regnault, V. 584(1992)93
- Rivat, C., see Ubrich, N. 584(1992)17
- Rivera, J., see Guardiola, A. 562(1991)481
- Rivier, L., see Saugy, M. 564(1991)567
- Rizzi, A. M., Briza, P. and Breitenbach, M.
Determination of D-alanine and D-glutamic acid in biological samples by coupled-column chromatography using β -cyclodextrin as mobile phase additive 582(1992)35
- Rizzi, A. M. and Plank, C.
Coupled column chromatography in chiral separations: systems employing β -cyclodextrin phases for chiral separation 557(1991)199
- Rizzo, V., see Melzi d'Eril, G. V. 553(1991)265
- Rizzolo, A., Baldo, C. and Polesello, A.
Application of high-performance liquid chromatography to the analysis of niacin and biotin in Italian almond cultivars 553(1991)187
- Rizzoni, G., see Betto, P. 584(1992)256
- Robben, A. J. P. M., see Visser, S. 599(1992)205
- Robert-Baudouy, J., see Gonzalès, T. 584(1992)101
- Roberts, M. F., see Abdulrahman, S. 562(1991)713
- Roberts, M. S., see Shen, J. 565(1991)309
- Roberts, N. B., see Jones, A. T. 599(1992)179
- Roberts, R. J., see Kinter, M. 578(1992)9
- Robertson, A. M., Littlejohn, D., Brown, M. and Dowle, C. J.
Assessment of a thermospray interface for the coupling of high-performance liquid chromatography and Fourier transform infrared spectrometry 588(1991)15
- Robin, D. W., see Edeki, T. 577(1992)190
- Robinett, R. S. R. and Hsieh, J. Y.-K.
Stereoselective determination of *R*(-)- and *S*(+)-MK-571, a leukotriene D₄ antagonist, in human plasma by chiral high-performance liquid chromatography 570(1991)157
- Robinson, M. L., see Dyas, A. M. 586(1991)351
- Robinson, P. R., Jones, M. D., Maddock, J. and Rees, L. W.
Simultaneous determination of clobopride and a major metabolite N-desbenzylclobopride in plasma by capillary gas chromatography-negative-ion chemical ionization mass spectrometry 564(1991)147
- Roblot, F., see Rey, J.-P. 587(1991)314
- Roblot, F., see Rey, J.-P. 596(1992)276
- Roboz, J. and Katz, R. N.
Diagnosis of disseminated candidiasis based on serum D/L-arabinitol ratios using negative chemical ionization mass spectrometry 575(1992)281
- Rocca, J. L., see Crétier, G. 590(1992)175
- Rocca, J. L., see Crétier, G. 596(1992)15
- Rocco, S., see Carpenè, G. 553(1991)201
- Rochet, T., see Duthel, J. M. 579(1992)85
- Rodchenkov, G. M., Vedenin, A. N., Uralets, V. P. and Semenov, V. A.
Characterization of prednisone, prednisolone and their metabolites by gas chromatography-mass spectrometry 565(1991)45
- Roder, J. D., see Chen, C. L. 583(1992)274
- Rodgers, M. A. J., see Zuk, M. M. 568(1991)437
- Rodrigues, A. E., Lopes, J. C., Lu, Z. P., Loureiro, J. M. and Dias, M. M.
Importance of intraparticle convection in the performance of chromatographic processes 590(1992)93
- Rodríguez, F. J., Jiménez, R. M. and Alonso, R. M.
Separation and determination of aminohalogenbenzophenones by high-performance liquid chromatography with electrochemical detection 578(1992)146
- Rodríguez, O. M., Desideri, P. G., Lepri, L. and Checchini, L.
Simultaneous separation and determination of hydrocarbons and organochlorine compounds by using a two-step microcolumn 555(1991)221
- Rodríguez, P., Permanyer, J., Grases, J. M. and González, C.
Confirmation method for the identification and determination of some organophosphorus and organochlorine pesticides in cocoa beans by gas chromatography-mass spectrometry 562(1991)547

- Rodríguez, P. A., Takigiku, R., Lehman-McKeeman, L. D., Fey, M. L., Eddy, C. L. and Caudill, D.
Design of a gas chromatograph with parallel radioactivity and mass spectrometric detection. Application to the identification of the major metabolite of *d*-limonene associated with α 2u-globulin 563(1991)271
- Rodríguez, R., see Zhu, M. 559(1991)479
- Rodríguez-Gonzalo, E., Sánchez-Martín, M. J. and Sánchez-Camazano, M.
Determination of acephate by liquid chromatography in the presence of aqueous soil extracts 585(1991)324
- Rodríguez Del Castillo, A., see Rotllán Pascual, P. 563(1991)37
- Roemen, T. H. M. and Van der Vusse, G. J.
Assessment of fatty acids in cardiac tissue as 9-anthryldiazomethane esters by high-performance liquid chromatography 570(1991)243
- Roets, E., see Naidong, W. 586(1991)61
- Roets, E., see Naidong, W. 586(1991)67
- Rogiers, V., see Mertens, K. 565(1991)149
- Rogos, R., see Hänslér, M. 563(1991)63
- Rogstad, A., see Hormazabal, V. 583(1992)201
- Roh, J. H., see Talaska, G. 580(1992)293
- Roh, J. K., see Lee, H. S. 574(1992)175
- Roldán, J. M., see Alhama, J. 589(1992)121
- Roles, J. and Guiochon, G.
Prediction of the elution profiles of high-concentration bands in gas chromatography 589(1992)223
- Roles, J. and Guiochon, G.
Experimental determination of adsorption isotherm data for the study of the surface energy distribution of various solid surfaces by inverse gas-solid chromatography 591(1992)233
- Roles, J. and Guiochon, G.
Precision and accuracy of the gas-solid adsorption isotherms derived by the elution-by-characteristic-points method 591(1992)245
- Roles, J. and Guiochon, G.
Validity of the model used to relate the energy distribution and the adsorption isotherm 591(1992)267
- Rolet, M.-C., see Drogue, S. 593(1992)363
- Rolle, P., see Abbati, G. 566(1991)445
- Rollin, P., see Latxague, L. 586(1991)166
- Rolls, W. A., see Jagodzinski, J. J. 591(1992)89
- Roman, M. and Brown, P. R.
Separation techniques for biotechnology in the 1990s 592(1992)3
- Roman, M. C. and Siergiej, R. W.
Accuracy of computer-simulated gas chromatographic separations based on a linear elution strength model 589(1992)215
- Romano, C., see Caruso, U. 562(1991)147
- Rombouts, F. M., see De Ruiter, G. A. 584(1992)69
- Róna, K. and Szabó, I.
Determination of mephenytoin stereoselective oxidative metabolism in urine by chiral liquid chromatography employing β -cyclodextrin as a mobile phase additive 573(1992)173
- Rongen, R., see Scouten, W. H. 597(1992)231
- Rönnerberg, I., see Fägerstam, L. G. 597(1992)397
- Roos, A. H., see Tuinstra, L. G. M. T. 552(1991)259
- Roos, P. H.
Immobilized metal affinity chromatography as a means of fractionating microsomal cytochrome P-450 isozymes 587(1991)33
- Roos, R. W. and Lau-Cam, C. A.
Utility of dansyl derivatization to the high-performance liquid chromatographic analysis of 2-phenylethylamine drugs 555(1991)278
- Roosemalen, M. C. M., Soons, P. A., Funaki, T. and Breimer, D. D.
High-performance liquid chromatographic determination of the polar metabolites of nifedipine in plasma, blood and urine 565(1991)516
- Roots, I., see Staffeldt, B. 571(1991)133
- Rop, P. P., Grimaldi, F., Bresson, M., Spinazzola, J., Quicke, J. and Viala, A.
Determination of dextromoramide in plasma and whole blood using high-performance liquid chromatography with ultraviolet absorbance detection 573(1992)87
- Rosa, M., see Pichini, S. 568(1991)267
- Rosa, M., see Pichini, S. 577(1992)358
- Rose, M., see Jacob, L. 587(1991)85
- Rose, M. E., see Lowes, S. 577(1992)205
- Rose, M. E., see Rimmer, D. A. 598(1992)251
- Rosegay, A., see Alvaro, R. F. 578(1992)327
- Rosell, A., Gomez-Belinchon, J. I. and Grimalt, J. O.
Gas chromatographic-mass spectrometric analysis of urban-related aquatic and airborne volatile organic compounds. Study of the extracts obtained by water closed-loop stripping and air adsorption with charcoal and polyurethane foam 562(1991)493
- Roselló-Catafau, J., see Ramis, I. 557(1991)507
- Roselló-Catafau, J., see Ramis, I. 575(1992)143
- Rosén, L., see Arvidsson, T. 586(1991)271
- Rosenberg, C., Aalto, T., Tornaues, J., Hesso, A., Jäppinen, P. and Vainio, H.
Identification by capillary gas chromatography-mass spectrometry of volatile organohalogen compounds formed during bleaching of kraft pulp 552(1991)265
- Rosenspire, K. C., Hirth, W., Jurisson, S., Nowotnik, D. P., Eckelman, W. C. and Nunn, A. D.
Direct analysis of whole blood by internal surface reversed-phase chromatography: an examination of the binding and metabolism of technetium dioxime complexes 574(1992)119
- Rosing, H., see Burger, D. M. 584(1992)239
- Rosseel, M. T. and Lefebvre, R. A.
Determination of cetirizine in human urine by high-performance liquid chromatography 565(1991)504
- Rosseel, M. T., Vermeulen, A. M. and Belpaire, F. M.
Reversed-phase high-performance liquid chromatographic analysis of atenolol enantiomers in plasma after chiral derivatization with (+)-1-(9-fluorenyl)ethyl chloroformate 568(1991)239
- Rosseel, M. T., see Vermeulen, A. M. 567(1991)472
- Rosset, R., see Drogue, S. 593(1992)363
- Rosset, R., see Villermet, A. 557(1991)85

- Rossetti, D. V., see Castagnola, M. 572(1991)51
- Rossi, C., see Quaglia, M. G. 593(1992)259
- Rossi, D. T., Overmyer, S. K., Lewis, R. C. and Narang, P. K.
Quantitation of a novel antiemetic (ADR-851) in plasma and urine by reversed-phase high-performance liquid chromatography with fluorescence detection 566(1991)257
- Rossini, M. L. and Farina, M.
Stability studies with a high-performance liquid chromatographic method for the determination of a new anthracycline analogue, 3'-deamino-3'-[2-(*S*)-methoxy-4-morpholino]doxorubicin (FCE 23762), in the final drug formulation 593(1992)47
- Rossiter, B. E., see Johnson, D. F. 594(1992)283
- Rossomando, E. F.
High-performance liquid chromatography in enzymatic analysis. Opening lecture 566(1991)275
- Rossomando, E. F., White, L. B. and Hadjimichael, J.
Immunomagnetic separation of tumor necrosis factor α . II. *In situ* procedure for the human gingival space 583(1992)19
- Rossomando, E. F., White, L. B., Hadjimichael, J. and Shafer, D.
Immunomagnetic separation of tumor necrosis factor α . I. Batch procedure for human temporomandibular fluid 583(1992)11
- Rossomando, E. F., see Spiegel, L. B. 573(1992)23
- Roth, H. J., see Gaede, F.-P. 585(1991)53
- Roth, M., see Koliadima, A. 595(1992)237
- Roth, W., see Greischel, A. 568(1991)246
- Rotllán Pascual, P., Ramos Carrillo, A. and Rodríguez Del Castillo, A.
Reversed-phase high-performance liquid chromatography of dinucleoside polyphosphates 563(1991)37
- Rotzsche, H.
Gas chromatographic analysis of fatty acid salts 552(1991)281
- Rouan, M. C., Campestrini, J., Lecaillon, J. B. and Godbillon, J.
Rapid determination of propyphenazone in plasma by high-performance liquid chromatography 577(1992)387
- Rouan, M. C., Campestrini, J., Le Clanche, V., Lecaillon, J. B. and Godbillon, J.
Automated microanalysis of carbamazepine and its epoxide and *trans*-diol metabolites in plasma by column liquid chromatography 573(1992)65
- Rouan, M. C., Le Duigou, F., Campestrini, J., Lecaillon, J. B. and Godbillon, J.
Fast liquid chromatography for the determination of drugs in plasma and combination with liquid-solid extraction in a fully automated system 573(1992)59
- Rouleau, J. R., see Abolfathi, Z. 579(1992)366
- Rouru, J., see Huupponen, R. 583(1992)270
- Rousseau-Tsangaris, M., see Desage, M. 571(1991)250
- Roussin, A., see Tellez, S. 581(1992)257
- Roussis, S. G., see Gagné, J.-P. 554(1991)293
- Roux, A., Blanchot, G., Baglin, A. and Flouvat, B.
Liquid chromatographic analysis of propranolol enantiomers in human blood using precolumn derivatization with (+)-1-(9-fluorenyl)ethyl chloroformate 570(1991)453
- Roux-de Balmann, H. and Sanchez, V.
Continuous-flow electrophoresis: a separation criterion applied to the separation of model proteins 594(1992)351
- Rowland, S. J., see Revill, A. T. 589(1992)281
- Rowlands, M. G., see Poon, G. K. 565(1991)75
- Roy, D., Hachey, D. L. and Liehr, J. G.
Determination of estradiol 2- and 4-hydroxylase activities by gas chromatography with electron-capture detection 567(1991)309
- Royer, D., see Le Verge, R. 574(1992)283
- Royer, R. J., see Del Cont Bernard, A. 574(1992)365
- Royer, R. J., see Royer-Morrot, M. J. 581(1992)297
- Royer-Morrot, M. J., Rambourg, M., Jacob, I., Bauer, P. and Royer, R. J.
Determination of zopiclone in plasma using column liquid chromatography with ultraviolet detection 581(1992)297
- Royer Morrot, M. J., see Del Cont Bernard, A. 574(1992)365
- Rozenman, Y., see Kersten, B. S. 588(1991)187
- Rozkošná, J., see Jandera, P. 556(1991)145
- Rozman, P., Kim, H. J., Madhu, C. and Klaassen, C. D.
Tissue sulfate determination by ion chromatography 574(1992)146
- Ruane, R. J., see Witherow, L. 553(1991)497
- Ruban, V. F.
Rapid determination of phenylalanine in plasma by capillary high-performance liquid chromatography with electrochemical detection 563(1991)134
- Rubino, F. M. and Zecca, L.
Application of triple quadrupole tandem mass spectrometry to the analysis of pyridine-containing derivatives of long-chain acids and alcohols 579(1992)1
- Ruckmick, S. C. and Hench, B. D.
Direct analysis of the dopamine agonist (-)-2-(*N*-propyl-*N*-2-thienylethylamino)-5-hydroxytetralin hydrochloride in plasma by high-performance liquid chromatography using two-dimensional column switching 565(1991)277
- Rudolph, W., see Karege, F. 570(1991)376
- Rugani, N., Dezan, C., De La Fournière, L., Cozzone, P. J., Bellon, B. and Sarda, L.
Separation and characterization of the precursor and activated forms of porcine and human pancreatic colipase by reversed-phase liquid chromatography 583(1992)246
- Ruggenti, P., see Gaspari, F. 570(1991)435
- Ruggieri, S., see Cecconi, O. 555(1991)267
- Rühl, C., see Jiao, Y. 595(1992)364
- Ruiz-Gutierrez, V., Cert, A. and Rios, J. J.
Determination of phospholipid fatty acid and triacylglycerol composition of rat caecal mucosa 575(1992)1
- Ruiz-Gutierrez, V., Montero, E. and Villar, J.
Determination of fatty acid and triacylglycerol composition of human adipose tissue 581(1992)171
- Rule, G. S. and Henion, J. D.
Determination of drugs from urine by on-line immunoaffinity chromatography-high-performance liquid chromatography-mass spectrometry 582(1992)103
- Rumble, R. H., see Jack, D. S. 584(1992)189

- Rumiantseva, N., see Polanuer, B. 594(1992)173
- Runov, V. K., see Fadeev, A. Y. 558(1991)31
- Rupprich, C., see Boeden, H.-F. 552(1991)389
- Rurak, D. W., see Wright, M. R. 565(1991)225
- Rurak, D. W., see Yeleswaram, K. 565(1991)383
- Rusconi, L. and Galantino, M.
Solvent system for the counter-current purification of hydrophobic bombesin analogues 589(1992)362
- Russell, I. M., see Shalliker, R. A. 558(1991)440
- Russell, J. W., Marrero, D., Whiterock, V. J., Klunk, L. J. and Starrett, J. E.
Determination of 9-[2-phosphonylmethoxy]ethyl]adenine in rat urine by high-performance liquid chromatography with fluorescence detection 572(1991)321
- Russo, M. V., see Cartoni, G. P. 552(1991)197
- Russo, R., see Garibotto, G. 572(1991)11
- Russo, V., see Bagnati, R. 564(1991)493
- Russo-Marie, F., see Jordan, L. M. 597(1992)299
- Rutan, S. C., see Cecil, T. L. 556(1991)495
- Rutten, G. and Rijks, J.
Activity testing and surface characterization of pretreated fused-silica capillaries for gas chromatography. A new modification of existing intermediate column tests 590(1992)271
- Rutten, W. J., see Essink, A. W. G. 570(1991)419
- Rydberg, T., Wählin-Boll, E. and Melander, A.
Determination of glibenclamide and its two major metabolites in human serum and urine by column liquid chromatography 564(1991)223
- Rykert, U. M., see Schwende, F. J. 565(1991)488
- Ryll, T. and Wagner, R.
Improved ion-pair high-performance liquid chromatographic method for the quantification of a wide variety of nucleotides and sugar-nucleotides in animal cells 570(1991)77
- Ryska, M., Koruna, I., Poláková, L. and Svátek, E.
Mass spectrometric structure determination of metabolites of 3-[2-(N,N-dimethylaminomethyl)phenylthio]phenol 562(1991)289
- Sabbah, S., see Driss, M. R. 552(1991)213
- Sabot, J.-F., see Daumas, L. 570(1991)89
- Sadlej-Sosnowska, N. and Śledzińska, I.
Validation of chromatographic retention models in reversed-phase high-performance liquid chromatography by fitting experimental data to the relevant equations 595(1992)53
- Sadony, V., see Himmel, H. M. 568(1991)105
- Saeed, K. and Becher, M.
On-line solid-phase extraction of piroxicam prior to its determination by high-performance liquid chromatography 567(1991)185
- Saeki, M., see Nagata, T. 565(1991)471
- Sagar, K. A., Kelly, M. T. and Smyth, M. R.
Analysis of terbutaline in human plasma by high-performance liquid chromatography with electrochemical detection using a micro-electrochemical flow cell 577(1992)109
- Sagara, K., see Oshima, T. 585(1991)255
- Sahali, Y., see Day, B. W. 562(1991)563
- Saini, G., see Silvestro, L. 563(1991)443
- Saisho, Y., see Goto, J. 567(1991)343
- Saito, K., Horie, M., Nose, N., Nakagomi, K. and Nakazawa, H.
High-performance liquid chromatography of histamine and 1-methylhistamine with on-column fluorescence derivatization 595(1992)163
- Saito, K., see Kim, M. 585(1991)45
- Saito, K., see Kim, M. 586(1991)27
- Saito, Y., see Jinno, K. 557(1991)459
- Saito, Y., see Matsuda, R. 585(1991)187
- Saitoh, K., see Shibata, Y. 598(1992)73
- Sakai, K., see Nagatomo, A. 597(1992)411
- Sakai, T. and Nagasawa, T.
Simple, rapid and sensitive determination of plasma taurine by high-performance liquid chromatography using pre-column derivative formation with fluorescamine 576(1992)155
- Sakai, T., see Nagasawa, T. 566(1991)223
- Sakaki, K. and Hirata, H.
Separation of secondary alcohol enantiomers using supercritical fluid chromatography 585(1991)117
- Sakakura, K., see Kanai, M. 567(1991)415
- Sakamoto, H., see Shimohigashi, Y. 597(1992)425
- Sakashita, S., see Miwa, T. 566(1991)163
- Sakata, T.-I., see Mukoyama, Y. 588(1991)195
- Sakihama, N., Nagai, K., Ohmori, H., Tomizawa, H., Tsujita, M. and Shin, M.
Immobilized ferredoxins for affinity chromatography of ferredoxin-dependent enzymes 597(1992)147
- Sakodinskaya, I. K., Desiderio, C., Nardi, A. and Fanali, S.
Micellar electrokinetic chromatographic study of hydroquinone and some of its ethers. Determination of hydroquinone in skin-toning cream 596(1992)95
- Sakodynskii, K. I., see Gorchakov, V. D. 563(1991)166
- Sakurai, T., Ichimiya, H., Miyazaki, H. and Nakayama, F.
Profiling of eicosanoids in inflamed gall bladder wall by gas chromatography with selected-ion monitoring 571(1991)1
- Sá, M. M. and Sereno, A. M.
Effect of column material on sorption isotherms obtained by inverse gas chromatography 600(1992)341
- Saar, J., see Yinon, J. 586(1991)73
- Sabanathan, S., see Clark, B. J. 553(1991)383
- Sabbah, S. and Bouguerra, M. L.
Analyse par chromatographie en phase gazeuse haute résolution-spectrométrie de masse des mélanges techniques de Phénochlor 552(1991)223

S

- Sala, M. R., see Garibotto, G. 572(1991)11
- Salamoun, J., Nguyen, P. T. and Remien, J.
Cation-exchange liquid chromatography of choline and acetylcholine on free shielded silanols of silica-based reversed-phase stationary phases 596(1992)43
- Salanon, M., see Chollet, D. 593(1992)73
- Salén, G., see Nyström, L. 574(1992)137
- Salganicoff, L., see Pulcinelli, F. M. 575(1992)51
- Sallustio, B. C., Morris, R. G. and Horowitz, J. D.
High-performance liquid chromatographic determination of sotalol in plasma. I. Application to the disposition of sotalol enantiomers in humans 576(1992)321
- Salm, P., see Taylor, P. J. 581(1992)152
- Salmon, J. A., see Woollard, P. M. 562(1991)249
- Salo, M., Vuorela, H. and Halmekoski, J.
Effect of temperature and mobile phase on the retention of retinoates in reversed-phase liquid chromatography 592(1992)127
- Salomon, K., Burgi, D. S. and Helmer, J. C.
Evaluation of fundamental properties of a silica capillary used for capillary electrophoresis 559(1991)69
- Salomoni, M., Muda, M., Zuccato, E. and Mussini, E.
High-performance liquid chromatographic determination of desmosine and isodesmosine after phenylisothiocyanate derivatization 572(1991)312
- Salonen, J. T., see Nyysönen, K. 570(1991)382
- Salto, T., see Sanchez-Brunete, C. 562(1991)525
- Salvador-Moya, M. D., see Fernández-Sánchez, E. 556(1991)485
- Samara, E., see Harvey, D. J. 562(1991)299
- Sameshima, I., see Okuda, T. 567(1991)141
- Sampson, D., see Havard, J. 584(1992)270
- Sánchez, A.-H., see Montaña, A. 594(1992)259
- Sanchez, V., see Roux-de Balmann, H. 594(1992)351
- Sánchez-Brunete, C., Pérez, S. and Tadeo, J. L.
Determination of phenoxy ester herbicides by gas and high-performance liquid chromatography 552(1991)235
- Sanchez-Brunete, C., Salto, T., Garcia-Baudin, J. M. and Tadeo, J. L.
Analysis of triallate residues in cereals and soil by gas chromatography with ion-trap detection 562(1991)525
- Sánchez-Camazano, M., see Rodríguez-Gonzalo, E. 585(1991)324
- Sánchez-Martín, M. J., see Rodríguez-Gonzalo, E. 585(1991)324
- Sancho, P., see Delgado, C. 594(1992)97
- Sandberg, M., see Orwar, O. 566(1991)39
- Sander, L. C., see Epler, K. S. 595(1992)89
- Sander, S., Hahn, A., Stein, J. and Rehner, G.
Comparative studies on the high-performance liquid chromatographic determination of thiamine and its phosphate esters with chloroethylthiamine as an internal standard using pre- and post-column derivatization procedures 558(1991)115
- Sanders, C., see Vermoesen, A. 564(1991)385
- Sanders, P., see Delépine, B. 582(1992)113
- Sandler, M., see Halket, J. M. 562(1991)279
- Sandra, P., see Coene, J. 553(1991)285
- Sangiah, S., see Chen, C. L. 583(1992)274
- Sano, A., Takimoto, N. and Takitani, S.
High-performance liquid chromatographic determination of cyanide in human red blood cells by pre-column fluorescence derivatization 582(1992)131
- Sano, K., Kinoshita, I., Mihara, R., Ikegami, Y. and Uesugi, T.
High-performance liquid chromatographic determination of sulphobromophthalein and its conjugates 578(1992)63
- Sano, M., see Ohara, K. 586(1991)35
- Sano, M., see Yokouchi, Y. 555(1991)297
- Sano, Y., see Yamamoto, S. 597(1992)173
- San Souci, B. R., see Phillips, D. J. 599(1992)239
- Santa, T., see Yamamura, Y. 567(1991)151
- Santambien, P., Girot, P., Hulak, I. and Boschetti, E.
Immunoenzymatic assay of dyes currently used in affinity chromatography for protein purification 597(1992)315
- Santilio, A., see Di Muccio, A. 552(1991)241
- Santilio, A., see Di Muccio, A. 553(1991)305
- Santoni, G., Bavazzano, P., Perico, A., Colzi, A., Benassi, S., Medica, A., La Morgia, R. and Giuliano, G.
High-performance liquid chromatographic determination of N-methylformamide and N-methyl-N-(hydroxymethyl) formamide in human urine 581(1992)287
- Sapis, J.-C., see Voirin, S. G. 595(1992)269
- Sarafraz-Yazdi, A., Khuahwar, M. Y. and Uden, P. C.
Gas chromatography of chloride and bromide as phenylboronic acid/mercuric nitrate derivatives with microwave induced plasma atomic emission detection 594(1992)395
- Sarati, S., Guiso, G., Spinelli, R. and Caccia, S.
Determination of piribedil and its basic metabolites in plasma by high-performance liquid chromatography 563(1991)323
- Sarda, L., see Rugani, N. 583(1992)246
- Saria, A.
Identification of α - and β -species of calcitonin gene-related peptide in the rat amygdala after separation with capillary zone electrophoresis 573(1992)219
- Sarin, M., Buinevicius, Z., Levitsky, S. and Feinberg, H.
Isocratic high-performance liquid chromatographic analysis of myocardial creatine phosphate and adenine nucleotides 563(1991)129
- Sarin, R., see Dua, V. K. 563(1991)333
- Sarre, S., Michotte, Y., Herregodts, P., Deleu, D., De Klippel, N. and Ebinger, G.
High-performance liquid chromatography with electrochemical detection for the determination of levodopa, catecholamines and their metabolites in rat brain dialysates 575(1992)207
- Sarre, S., Michotte, Y., Marvin, C. A. and Ebinger, G.
Microbore liquid chromatography with dual electrochemical detection for the determination of serotonin and 5-hydroxyindoleacetic acid in rat brain dialysates 582(1992)29
- Sartori, S., see Pellegatti, M. 573(1992)105
- Sarver, E. W., see Fowler, W. K. 558(1991)235
- Sasabe, H., see Kubo, M. 564(1991)137
- Sasabe, H., see Nagasawa, M. 577(1992)275
- Sasaki, H., see Endoh, Y. S. 579(1992)123
- Sasaki, H., see Wakui, Y. 575(1992)131
- Sasaki, N., see Endoh, Y. S. 579(1992)123

- Sasamoto, Y., see Kitagawa, T. 567(1991)267
- Sasinos, F. I., see Kim, I. S. 589(1991)177
- Saska, M., Clarke, S. J., Wu, M. D. and Iqbal, K.
Glucose-fructose equilibria on Dowex Monosphere 99
CA resin under overloaded conditions 590(1992)147
- Sasunkevich, V. A., see Voinov, V. G. 586(1991)360
- Sato, H., see Yokoyama, Y. 555(1991)155
- Sato, H., see Yokoyama, Y. 566(1991)19
- Sato, H., see Yokoyama, Y. 583(1992)1
- Sato, J., Watanabe, K., Takahama, S. and Makita, T.
Quantification of 5-fluoro-2'-deoxyuridine in plasma by
gas chromatography and negative-ion chemical ionization
mass spectrometry 575(1992)63
- Sato, K., see Watabe, K. 590(1992)289
- Sato, T., see Goto, N. 578(1992)203
- Sato, T., see Kim, M. 585(1991)45
- Sato, T., see Matsushima, E. 573(1992)339
- Sato, T., see Nawata, M. 597(1992)415
- Sato, T., see Oda, Y. 572(1991)133
- Satoh, N., see Kawakami, Y. 576(1992)63
- Satoh, S., see Ohhashi, T. 585(1991)153
- Satorre, J., see Esteban, A. 573(1992)121
- Sattari, S., see Wright, M. R. 583(1992)259
- Sauer, W. C., see Dugan, M. E. R. 582(1992)242
- Saugy, M., Meuwly, P., Munafo, A. and Rivier, L.
Rapid high-performance liquid chromatographic
determination with fluorescence detection of furosemide
in human body fluids and its confirmation by gas
chromatography-mass spectrometry 564(1991)567
- Saunders, P. P., Alvarez, E. and Kantarjian, H. M.
Determination of nicotinamide-adenine dinucleotide and
thiazole-4-carboxamide-adenine dinucleotide in human
leukocytes by reversed-phase high-performance liquid
chromatography 577(1992)37
- Sautou, V., Chopineau, J., Terrisse, M. P. and Bastide, P.
Solid-phase extraction of midazolam and two of its
metabolites from plasma for high-performance liquid
chromatographic analysis 571(1991)298
- Sawada, Y., see Yamamura, Y. 567(1991)151
- Sawatzki, G., see Thurl, S. 568(1991)291
- Saynor, D. A., see Bowers, G. D. 554(1991)175
- Scalia, S. and Games, D. E.
Analysis of conjugated bile acids by packed-column
supercritical fluid chromatography 574(1992)197
- Scapol, L., Rappuoli, P. and Viscomi, G. C.
Purification of recombinant human interferon- β by
immobilized antisense peptides 600(1992)235
- Schad, H., Schäfer, F., Weber, L. and Seidel, H. J.
Determination of benzene metabolites in urine of mice by
solid-phase extraction and high-performance liquid
chromatography 593(1992)147
- Schaefer, T. F., see Lehman, E. D. 574(1992)225
- Schäfer, F., see Schad, H. 593(1992)147
- Schafer, W. A. and Carr, P. W.
Chromatographic characterization of a phosphate-
modified zirconia support for bio-chromatographic
applications 587(1991)149
- Schafer, W. A., Carr, P. W., Funkenbusch, E. F. and Parson,
K. A.
Physical and chemical characterization of a porous
phosphate-modified zirconia substrate 587(1991)137
- Schalekamp, M. A. D. H., see Alberts, G. 583(1992)236
- Schalekamp, M. A. D. H., see Boomsma, F. 574(1992)109
- Schalekamp, M. A. D. H., see Van Der Hoorn, F. A. J.
563(1991)348
- Schalkhammer, T., see Ferris, R. 577(1992)251
- Schär, M., see Steiner, V. 586(1991)43
- Scheinin, M., see Ojala-Karlsson, P. 565(1991)131
- Schenderjuk, V. V., see Golovnya, R. V. 552(1991)1
- Schepers, C. H. M., see Duchateau, A. L. L. 566(1991)493
- Scherlag, B. J., see Chen, C. L. 583(1992)274
- Scherz, H., see Haidacher, D. 591(1992)351
- Schiavi, M., see Italia, A. 553(1991)15
- Schiavon, G., see Tagliaro, F. 563(1991)11
- Schiavon, G., see Tagliaro, F. 563(1991)299
- Schildnitz, U., see Huber, R. 553(1991)65
- Schill, G., see Arvidsson, E. 591(1992)55
- Schill, G., see Balmér, K. 553(1991)391
- Schill, G., see Balmér, K. 592(1992)331
- Schilling, B., see Michaelis, H. C. 571(1991)257
- Schilt, R., see Ploum, M. E. 564(1991)413
- Schlag, G., see Gasser, H. 562(1991)257
- Schmalzing, D., see Jung, M. 552(1991)43
- Schmarr, H.-G., see Karl, V. 587(1991)347
- Schmid, E. R., see Gasser, H. 562(1991)257
- Schmid, J., see Buszewski, B. 552(1991)415
- Schmidt, J. and McClain, C. J.
Separation of *o*-phthalaldehyde-mercaptoethanol
derivatives of amino acids from blood plasma on
reversed-phase Nova-Pak C₁₈ cartridges 568(1991)207
- Schmidt, N., Brune, K. and Geisslinger, G.
Stereoselective determination of the enantiomers of
methadone in plasma using high-performance liquid
chromatography 583(1992)195
- Schmidt, T., see Bösward, J. 571(1991)19
- Schmidt, Jr., D. E., see Gotsick, J. T. 590(1992)77
- Schmith, V. D., Cox, S. R., Zemaitis, M. A. and Kroboth, P.
D.
New high-performance liquid chromatographic method
for the determination of alprazolam and its metabolites in
serum: instability of 4-hydroxyalprazolam 568(1991)253
- Schmitter, J.-M.
Performances and limits of plasma desorption mass
spectrometry in the primary structure determination of
proteins 557(1991)359
- Schneider, R. E., see Takigiku, R. 559(1991)247
- Schneider, W., see Degen, P. H. 563(1991)193
- Schneider, W., see Degen, P. H. 565(1991)67
- Schoel, B. and Kaufmann, S. H. E.
Hydrophobic interaction chromatography for the
purification of a mycobacterial heat shock protein of
relative molecular mass 60 000 587(1991)19
- Schoenfeld, N. and Mamet, R.
High-performance liquid chromatographic detection of
pitfalls in porphobilinogen deaminase determination
570(1991)51
- Schoenmakers, P. J., see Hamoir, T. 589(1992)31
- Schoenmakers, P. J., see Hogendoorn, E. A. 552(1991)113
- Schoenmakers, P. J., see Janssen, H.-G. 552(1991)527
- Schoenmakers, P. J., see Lopes Marques, R. M. 592(1992)157

- Schöllhammer, G., see Süß, S. 565(1991)363
- Schols, H. A., see De Ruiter, G. A. 584(1992)69
- Schönheiter, H., see Fenske, M. 563(1991)178
- Schoofs, P. R., see De Jong, J. 570(1991)209
- Schoonjans, R., see De Vos, M. 564(1991)296
- Schöpp, W., see Grunow, M. 590(1992)247
- Schornagel, J. H., see Van Tellingen, O. 576(1992)158
- Schoun, J., see Dulery, B. D. 571(1991)241
- Schreck, C. E., see Warthen, Jr., J. D. 590(1992)133
- Schreurs, J., see Svoboda, M. 562(1991)403
- Schreurs, M., Hellendoorn, L., Gooijer, C. and Velthorst, N. H.
Time-resolved luminescence detection of derivatized thiols in column liquid chromatography 552(1991)625
- Schrijver, J., see Van Vliet, T. 553(1991)179
- Schröder, H. F.
Polar, hydrophilic compounds in drinking water produced from surface water. Determination by liquid chromatography-mass spectrometry 554(1991)251
- Schubert-Rehberg, K., see Abou-Rebyeh, H. 566(1991)341
- Schubert-Rehberg, K., see Abou-Rebyeh, H. 566(1991)351
- Schuler, G. and Reinacher, M.
Development and optimization of a single-step procedure using protein A affinity chromatography to isolate murine IgG₁ monoclonal antibodies from hybridoma supernatants 587(1991)61
- Schulman, C. A., see Lehman, E. D. 574(1992)225
- Schultz, G. A., Chamberlin, B. A., Sweeley, C. C., Watson, J. T. and Allison, J.
Complex mixture analysis based on gas chromatography-mass spectrometry with time array detection using a beam deflection time-of-flight mass spectrometer 590(1992)329
- Schultz, P., see Mitchell, E. P. 581(1992)31
- Schulz, H.-U., see Kraus, G. 573(1992)323
- Schulz, R., see Hagedorn, H.-W. 577(1992)195
- Schulze-Clewing, J., see Pollak, V. A. 596(1992)241
- Schurig, V., see Jung, M. 552(1991)43
- Schuring, J., see Gelsema, W. J. 594(1992)217
- Schuster, O., see Geisslinger, G. 573(1992)163
- Schütz, A., see Brunmark, P. 573(1992)35
- Schwadorf, K. and Müller, H.-M.
Determination of α - and β -zearealenol and zearealenone in cereals by gas chromatography with ion-trap detection 595(1992)259
- Schwartz, D., see Sonnet, P. E. 586(1991)255
- Schwartz, D. P., see Medina, M. B. 581(1992)119
- Schwartz, H. E., Ulfelder, K., Sunzeri, F. J., Busch, M. P. and Brownlee, R. G.
Analysis of DNA restriction fragments and polymerase chain reaction products towards detection of the AIDS (HIV-1) virus in blood 559(1991)267
- Schwarz, D., see Kang, K. 567(1991)57
- Schwende, F. J. and Rykert, U. M.
Determination of alantamol hydrobromide, a novel antipsychotic agent, in human blood plasma and urine by high-performance liquid chromatography with fluorescence detection and solid-phase extraction 565(1991)488
- Schwer, C., see Kenndler, E. 595(1992)313
- Schwertner, H. A., Patterson, W. R., Cissik, J. H. and Wilson, K. W.
New extraction procedure and high-performance liquid chromatographic method for analyzing polyethylene glycol-400 in urine 578(1992)297
- Sciaky, M., see Masson, S. 563(1991)231
- Scott, D. A., Hammond, P. M., Brearley, G. M. and Price, C. P.
Identification by high-performance liquid chromatography of tyrosine ammonia-lyase activity in purified fractions of *Phaseolus vulgaris* phenylalanine ammonia-lyase 573(1992)309
- Scott, M. D., Baudendistel, L. J. and Dahms, T. E.
Rapid separation of creatine, phosphocreatine and adenosine metabolites by ion-pair reversed-phase high-performance liquid chromatography in plasma and cardiac tissue 576(1992)149
- Scott, M. T., see Reiser, R. W. 554(1991)91
- Scouten, W. H., Adler, C. M., Rongen, R. and Mallee, L.
Evaluation of the mode of binding of immunoglobulin to activated agarose 597(1992)231
- Scudamore, K. A., see Hetmanski, M. T. 588(1991)47
- Seay, S. S., see Tyczkowska, K. L. 576(1992)305
- Sébidio, J. L., see Le Quéré, J. L. 562(1991)659
- Sebert, J., see Kamel, S. 574(1992)255
- Sebille, B., see Agnus, B. 552(1991)583
- Sebille, B., see Gosselet, M. 552(1991)563
- Sébille, B., see Lemque, R. 553(1991)165
- Sébille, B., see Lemque, R. 599(1992)255
- Sebille, B., see Thuaud, N. 555(1991)53
- Sébille, B., see Vidal-Madjar, C. 584(1992)11
- Sébille, B., see Vivarat-Perrin, M.-P. 584(1992)3
- Sebille, B., see Voirin, A. 553(1991)155
- Šebor, G., see Černý, J. 600(1992)243
- Secchi, C., see Biondi, P. A. 566(1991)377
- Secreto, G., see Venturelli, E. 582(1992)7
- Sedláček, J., see Klimeš, J. 584(1992)221
- Sedmera, P., see Beran, M. 558(1991)265
- Seelen, I. G. J. A., see Ackermans, M. T. 590(1992)341
- Segura, J., see Ventura, R. 562(1991)723
- Sehgal, R. K., see Amari, J. V. 590(1992)153
- Seidah, N. G., see Basak, A. 581(1992)17
- Seidel, H. J., see Schad, H. 593(1992)147
- Seiler, W., see Süß, S. 565(1991)363
- Seipel, L., see Mayer, F. 572(1991)339
- Seitz, H. U., see Glässgen, W. E. 598(1992)81
- Seitz, J., Keppler, C. and Hüntemann, S. B.
Purification and characterization of transglutaminases from the genital tract of the male rat 587(1991)55
- Seitz, U., Bonn, G., Oefner, P. and Popp, M.
Isotachophoretic analysis of flavonoids and phenolcarboxylic acids of relevance to phytopharmaceutical industry 559(1991)499
- Sekiba, K., see Noma, J. 568(1991)35
- Sellberg, B., see Castledine, J. B. 592(1992)27
- Sellier, N., Cazaussus, A., Budzinski, H. and Lebon, M.
Structure determination of sesquiterpenes in Chinese vetiver oil by gas chromatography-tandem mass spectrometry 557(1991)451

- Semba, J., Ratnaraj, N. and Patsalos, P. N.
Simple and rapid micro-analytical procedures for the estimation of milacemide and its metabolite glycynamide in rat plasma and cerebrospinal fluid by high-performance liquid chromatography 565(1991)357
- Semba, J., see Watanabe, A. 583(1992)241
- Semenov, S. N.
Integral Doppler anemometry in porous membranes for the analysis of liquid mixtures and examination of membrane properties 600(1992)129
- Semenov, S. N.
Transverse particle redistribution in a flat channel for SPLITT or integral Doppler anemometry 600(1992)133
- Semenov, V. A., see Rodchenkov, G. M. 565(1991)45
- Semporé, B. G. and Bézard, J. A.
Enantiomer separation by chiral-phase liquid chromatography of urethane derivatives of natural diacylglycerols previously fractionated by reversed-phase liquid chromatography 557(1991)227
- Semporé, B. G. and Bézard, J. A.
Separation of monoacylglycerols by reversed-phase high-performance liquid chromatography 596(1992)185
- Sennello, L. T., see Chu, S.-Y. 571(1991)199
- Seno, N., see Isosaki, K. 597(1992)123
- Seno, N., see Kojima, K. 597(1992)323
- Seno, N., see Ueno, M. 597(1992)197
- Seo, A., see Kunkel, M. E. 589(1992)135
- Sepai, O., see Kajbaf, M. 583(1992)63
- Sepaniak, M. J., see Arrowood, S. 583(1992)105
- Sepaniak, M. J., see Cole, R. O. 557(1991)113
- Serenó, A. M., see Sá, M. M. 600(1992)341
- Sernicola, L., see Di Muccio, A. 552(1991)241
- Serot, T., see Courcoux, P. 596(1992)225
- Serrano-Carreón, L., Hathout, Y., Bensoussan, M. and Belin, J.-M.
Quantitative separation of *Trichoderma* lipid classes on a bonded phase column 584(1992)129
- Sertillanges, P., see Rivat, C. 576(1992)71
- Šěrys, A., see Maruška, A. 596(1992)157
- Sessink, P. J. M., see Dirven, H. A. A. M. 564(1991)266
- Seto, H. and Ohkubo, T.
Determination of a guanosine-malonaldehyde adduct in urine by high-performance liquid chromatography with a thiobarbituric acid reaction detector 570(1991)301
- Settimj, G., see Betto, P. 563(1991)115
- Severini, G., Malaguti Aliberti, L. and Di Giovannandrea, R.
Diagnostic aspects of alkaline phosphatase: separation of isoenzymes in normal and pathological human serum by high-performance liquid chromatography 563(1991)147
- Sevillano-Cabeza, A., see Herráez-Hernández, R. 582(1992)181
- Sevini, F., see Marcato, B. 553(1991)415
- Sewell, G. W., see Chamkasem, N. 587(1991)185
- Sewell, R. B., see Martin, G. 574(1992)205
- Seyama, C., see Haginaka, J. 577(1992)95
- Seyama, C., see Haginaka, J. 592(1992)301
- Seyama, C., see Haginaka, J. 598(1992)67
- Shackleton, C. H. L., Falick, A. M., Green, B. N. and Witkowska, H. E.
Electrospray mass spectrometry in the clinical diagnosis of variant hemoglobins 562(1991)175
- Shackleton, C. H. L., see Kuypers, F. A. 562(1991)191
- Shafer, D., see Rossomando, E. F. 583(1992)11
- Shafiq-Ur-Rehman,
Rapid isocratic method for the separation and quantification of major phospholipid classes by high-performance liquid chromatography 567(1991)29
- Shah, R. D. and Maryanoff, C. A.
High-performance liquid chromatographic analysis of diastereomers and enantiomers of pyrroloisoquinoline antidepressants 588(1991)348
- Shaikh, B., Jackson, J., Guyer, G. and Ravis, W. R.
Determination of neomycin in plasma and urine by high-performance liquid chromatography. Application to a preliminary pharmacokinetic study 571(1991)189
- Shalliker, R. A., Kavanagh, P. E. and Russell, I. M.
Reversed-phase separations of high-molecular-weight polystyrenes in a dichloromethane-acetonitrile solvent system 558(1991)440
- Shankar, A. and Lenhoff, A. M.
Dispersion in round tubes and its implications for extra-column dispersion 556(1991)235
- Shanley, M., see Dutta, P. K. 599(1992)137
- Shansky, R. E. and Kane, R. E.
Separation of soy lecithin using gel permeation chromatography 589(1992)165
- Shantha, N. C. and Ackman, R. G.
Behaviour of a common phthalate plasticizer (dioctyl phthalate) during the alkali- and/or acid-catalysed steps in an AOCs method for the preparation of methyl esters 587(1991)263
- Shapi, M. M. and Hesso, A.
Gas chromatographic-mass spectrometric analysis of some potential toxicants amongst volatile compounds emitted during large-scale thermal degradation of poly (acrylonitrile-butadiene-styrene) plastic 562(1991)681
- Shara, M. A., Dickson, P. H., Bagchi, D. and Stohs, S. J.
Excretion of formaldehyde, malondialdehyde, acetaldehyde and acetone in the urine of rats in response to 2,3,7,8-tetrachlorodibenzo-*p*-dioxin, paraquat, endrin and carbon tetrachloride 576(1992)221
- Sharma, M., see Lee, T. 565(1991)197
- Sharma, O. P. and Dawra, R. K.
Thin-layer chromatographic separations of lantadenes, the pentacyclic triterpenoids from lantana (*Lantana camara*) plant 587(1991)351
- Sharma, R., see Awasthi, S. 584(1992)167
- Sharma, S. D. and Misra, S.
Quantitative separation of Cr³⁺ from Mo⁶⁺, W⁶⁺, Hg₂²⁺, Cu²⁺ and Pb²⁺. Chromatographic behaviour of 51 cations on papers impregnated with Sn(IV)-based inorganic ion exchangers in complex-forming acid systems 594(1992)379
- Sharma, S. K. and Dakshinamurti, K.
Determination of vitamin B₆ vitamers and pyridoxic acid in biological samples 578(1992)45
- Sharma, V. P., see Dua, V. K. 563(1991)333
- Shatz, V. D. and Kažoka, H. A.
Adsorption and partition mode in high-performance liquid chromatography of highly polar solutes on silica 552(1991)23

- Shaw, P. N., see Barrett, D. A. 566(1991)135
- Shaw, P. N., see Blagbrough, I. S. 578(1992)251
- Shaw, P. N., see McGee, J. P. 567(1991)282
- Shcherbakova, K. D., see Nazarova, V. I. 600(1992)59
- Shcherbakova, O. A., see Nazarova, V. I. 600(1992)59
- Shearer, G., see Tarbin, J. A. 577(1992)376
- Shearer, G., see Tarbin, J. A. 579(1992)177
- Shen, G., Weirong, T. and Shixiang, W.
Simple high-performance liquid chromatographic method for the determination of captopril in biological fluids 582(1992)258
- Shen, J., Wanwimolruk, S. and Roberts, M. S.
Novel direct high-performance liquid chromatographic method for determination of salicylate glucuronide conjugates in human urine 565(1991)309
- Shenfield, G. M., see Boutagy, J. 565(1991)523
- Shepard, R. M., Duthu, G. S., Ferraina, R. A. and Mullins, M. A.
High-performance liquid chromatographic assay with electrochemical detection for azithromycin in serum and tissues 565(1991)321
- Shepard, T. A., see Wilhelm, J. A. 578(1992)231
- Shephard, G. S., Thiel, P. G. and Sydenham, E. W.
Determination of fumonisin B₁ in plasma and urine by high-performance liquid chromatography 574(1992)299
- Shephard, G. S., Thiel, P. G., Sydenham, E. W., Vleggaar, R. and Marasas, W. F. O.
Reversed-phase high-performance liquid chromatography of tenuazonic acid and related tetramic acids 566(1991)195
- Sheu, S.-J., see Liu, Y.-M. 600(1992)370
- Sheumack, D., see Haynes, P. A. 588(1991)107
- Shi, Q., Olley, P. M. and Jackowski, G.
Comparison of ion-exchange chromatography, isoelectric precipitation and reversed-phase high-performance liquid chromatography for the separation of individual cardiac myosin light chains 572(1991)59
- Shibasaki, H., Arai, I., Furuta, T. and Kasuya, Y.
Simultaneous determination of cortisol and cortisone in human plasma by stable-isotope dilution mass spectrometry 576(1992)47
- Shibasaki, H., Furuta, T. and Kasuya, Y.
Stable isotope dilution mass spectrometry for the simultaneous determination of cortisol, cortisone, prednisolone and prednisone in plasma 579(1992)193
- Shibasaki, H., see Furuta, T. 576(1992)213
- Shibata, K. and Onodera, M.
High-performance liquid chromatographic determination of 3-hydroxykynurenine with fluorimetric detection; comparison of preovulatory phase and postovulatory phase urinary excretion 570(1991)13
- Shibata, N., Akabane, M., Minouchi, T., Ono, T. and Shimakawa, H.
Fluorimetric determination of mexiletine in serum by high-performance liquid chromatography using pre-column derivatization with fluorecamine 566(1991)187
- Shibata, Y., Saitoh, K. and Suzuki, N.
Control of the retention selectivity of rare earth octaethylporphyrins in reversed-phase high-performance liquid chromatography using amines as mobile phase additives 598(1992)73
- Shibukawa, A. and Wainer, I. W.
Simultaneous direct determination of the enantiomers of verapamil and norverapamil in plasma using a derivatized amylose high-performance liquid chromatographic chiral stationary phase 574(1992)85
- Shibusawa, Y., Ito, Y., Ikewaki, K., Rader, D. J. and Brewer, Jr., H. B.
Counter-current chromatography of lipoproteins with a polymer phase system using the cross-axis synchronous coil planet centrifuge 596(1992)118
- Shibusawa, Y., see Matsumoto, U. 566(1991)67
- Shida, N., see Yasukawa, K. 597(1992)271
- Shidawara, S., see Akiyama, H. 579(1992)203
- Shigematsu, N., see Honda, A. 565(1991)53
- Shigenobu, K., see Koizumi, K. 595(1992)340
- Shigetoshi, M., see Goto, N. 578(1992)203
- Shijo, Y., see Uehara, N. 586(1991)356
- Shima, K., see Yasukawa, K. 597(1992)271
- Shimada, K., Komine, Y. and Mitamura, K.
High-performance liquid chromatographic separation of bile acid pyrenacyl esters with cyclodextrin-containing mobile phase 565(1991)111
- Shimada, K., Oe, T. and Suzuki, M.
Effect of derivatization of steroids on their retention behaviour in inclusion chromatography using cyclodextrin as a mobile phase additive 558(1991)306
- Shimada, K., see Wakabayashi, H. 573(1992)154
- Shimada, K., see Yamato, S. 590(1992)241
- Shimakawa, H., see Shibata, N. 566(1991)187
- Shimamoto, C., see Kawakami, Y. 576(1992)63
- Shimasaki, H., see Werner, A. 553(1991)205
- Shimizu, N., see Mukoyama, Y. 588(1991)195
- Shimizu, T., see Kubo, M. 564(1991)137
- Shimizu, T., see Nagasawa, M. 577(1992)275
- Shimkus, J. K., see Kononenko, V. L. 553(1991)517
- Shimkus, J. K., see Kononenko, V. L. 600(1992)139
- Shimohigashi, Y., Takada, K., Sakamoto, H., Matsumoto, H., Yasunaga, T., Kondo, M. and Ohno, M.
Discriminative affinity labelling of opioid receptors by enkephalin and morphiceptin analogues containing 3-nitro-2-pyridinesulphenyl-activated thiol residues 597(1992)425
- Shimohigashi, Y., see Nagatomo, A. 597(1992)411
- Shimojo, M., see Shioya, H. 568(1991)309
- Shimojo, T., see Nagata, Y. 575(1992)147
- Shin, H. C., see Lee, H. S. 574(1992)175
- Shin, H.-S., see Kim, T.-J. 575(1992)295
- Shin, M., see Sakihama, N. 597(1992)147
- Shin, M. J., see Park, J. H. 595(1992)45
- Shinka, T., see Matsumoto, M. 562(1991)139
- Shinkai, F., see Isobe, T. 588(1991)115
- Shinkai, H., Takahashi, H., Kikuchi, K., Kawai, H. and Otake, N.
Determination of the new morpholino anthracycline MX2-HCl and its metabolites in biological samples by high-performance liquid chromatography 570(1991)203
- Shinomiya, K., see Imanari, T. 574(1992)142
- Shinomiya, K., see Toyoda, H. 565(1991)141

- Shintani, H.
Solid-phase extraction and high-performance liquid chromatographic analysis of a toxic compound from γ -irradiated polyurethane 600(1992)93
- Shiono, H., see Fukushima, S. 565(1991)35
- Shioya, H., Shimojo, M. and Kawahara, Y.
Determination in plasma of angiotensin-converting enzyme inhibitor by inhibitor-binding assay 568(1991)309
- Shirota, O., see Liu, J. 559(1991)223
- Shishkina, I. P., see Galushko, S. V. 592(1992)345
- Shishkina, I. P., see Galushko, S. V. 600(1992)83
- Shixiang, W., see Shen, G. 582(1992)258
- Shlyamin, V. I., see Obrezkov, O. N. 558(1991)209
- Shoda, J., see Honda, A. 565(1991)53
- Shoemaker, J. D. and Elliott, W. H.
Automated screening of urine samples for carbohydrates, organic and amino acids after treatment with urease 562(1991)125
- Shoji, F., see Negawa, M. 590(1992)113
- Sholin, A., see Polanuer, B. 594(1992)173
- Shoto, E., see Hu, J.-Y. 594(1992)371
- Shpigun, O. A., see Obrezkov, O. N. 558(1991)209
- Shuely, W. J., see Abraham, M. H. 587(1991)213
- Shuely, W. J., see Abraham, M. H. 587(1991)229
- Shukla, U. A., see Janiszewski, J. S. 567(1991)274
- Shulgin, A. T., see Jacob, III, P. 583(1992)145
- Shvets, V. I., see Klyashchitsky, B. A. 577(1992)267
- Siboo, I. R., see Chan, E. C. S. 568(1991)85
- Siboo, R., see Chan, E. C. S. 568(1991)85
- Siebert, C. J., see Morehead, H. W. 587(1991)171
- Siebert, C. J., see Talmadge, K. W. 590(1992)83
- Siems, W., see Grune, T. 563(1991)53
- Siems, W. G., see Grune, T. 553(1991)193
- Siergiej, R. W., see Roman, M. C. 589(1992)215
- Sigon, F., see Blo, G. 556(1991)249
- Silberring, J., Brostedt, P., Thörnwall, M. and Nyberg, F.
Approach to studying proteinase specificity by continuous-flow fast atom bombardment mass spectrometry and high-performance liquid chromatography combined with photodiode-array ultraviolet detection 554(1991)83
- Silberring, J. and Nyberg, F.
Analysis of tyrosine- and methionine-containing neuropeptides by fast atom bombardment mass spectrometry 562(1991)459
- Silks III, L. A., see Terry, Jr., A. V. 585(1991)101
- Silvestro, L., Viano, I., Baiocchi, C., Saini, G., Marmont, F. and Ferro, R.
Quantitation of melphalan in plasma of patients by reversed-phase high-performance liquid chromatography with electrochemical detection 563(1991)443
- Silvestro, L., Viano, I., Naggi, A., Torri, G., Da Col, R. and Baiocchi, C.
High-performance liquid chromatographic-mass spectrometric analysis of oligosaccharides from enzymatic digestion of glycosaminoglycans. Application to human samples 591(1992)225
- Simal Gándara, J., see Paseiro Losada, P. 585(1991)75
- Simal Lozano, J., see Paseiro Losada, P. 585(1991)75
- Simmonds, P. G., see Begley, P. 588(1991)239
- Simpson, C. F., see Akapo, S. O. 557(1991)515
- Simpson, R. C., see Boppana, V. K. 593(1992)29
- Simpson, R. J., see Moritz, R. L. 599(1992)119
- Sims, T. J. and Bailey, A. J.
Quantitative analysis of collagen and elastin cross-links using a single-column system 582(1992)49
- Sing, K. S. W., see Cao, X. L. 555(1991)183
- Singh, A. K.
Quantitative analysis of inositol lipids and inositol phosphates in synaptosomes and microvessels by column chromatography: comparison of the mass analysis and the radiolabelling methods 581(1992)1
- Singh, A. K., Jang, Y., Mishra, U. and Granley, K.
Simultaneous analysis of flunixin, naproxen, ethacrynic acid, indomethacin, phenylbutazone, mefenamic acid and thiosalicylic acid in plasma and urine by high-performance liquid chromatography and gas chromatography-mass spectrometry 568(1991)351
- Singh, J., see Huh, H. 600(1992)364
- Singh, S., see Epemolu, R. O. 573(1992)178
- Singleton, H. J., see Lennard, L. 583(1992)83
- Sinha, P., see Righetti, P. G. 569(1991)197
- Sinkkonen, S.
Isolation of chlorinated dibenzothiophenes by high-performance thin-layer chromatography 553(1991)453
- Sinz, M. W. and Rimmel, R. P.
Analysis of lamotrigine and lamotrigine 2-N-glucuronide in guinea pig blood and urine by reserved-phase ion-pairing liquid chromatography 571(1991)217
- Siouffi, A.-M.
Some aspects of optimization in planar chromatography (Review) 556(1991)81
- Sioufi, A., Pommier, F. and Godbillon, J.
Determination of diclofenac in plasma and urine by capillary gas chromatography-mass spectrometry with possible simultaneous determination of deuterium-labelled diclofenac 571(1991)87
- Sioufi, A., Richard, J., Mangoni, P. and Godbillon, J.
Determination of diclofenac in plasma using a fully automated analytical system combining liquid-solid extraction with liquid chromatography 565(1991)401
- Sips, J. H. M., see Van Tellingen, O. 576(1992)158
- Sirén, H. and Riekkola, M.-L.
Determination of metal ions by on-line complexation and ion-pair chromatography 590(1992)263
- Sissons, D. J., see Van Bergen, C. A. 589(1992)109
- Sitkey, V., Travěncová, I. and Minárik, M.
Isolation and purification of bacterial proteinases by means of autofocusing 585(1991)373
- Sitkiewicz, D., see Paczkowski, D. 573(1992)235
- Siu, K. W. M., Guevremont, R., Le Blanc, J. C. Y., Gardner, G. J. and Berman, S. S.
Electrospray interfacing for the coupling of ion-exchange and ion-pairing chromatography to mass spectrometry 554(1991)27
- Sjödín, K., see Moberg, C. 585(1991)309
- Skajaa, K., see Christensen, H. R. 574(1992)161

- Skarping, G., see Brunmark, P. 573(1992)35
 Skarping, G., see Brunmark, P. 579(1992)350
 Skarping, G., see Jönsson, B. 558(1991)247
 Skarping, G., see Jönsson, B. 572(1991)117
 Skee, D. M., Cook, J. A. and Benziger, D. P.
 Determination of a novel steroidal androgen receptor antagonist (Win 49596) in human plasma using solid-phase extraction and high-performance liquid chromatography with ultraviolet detection 568(1991)494
 Skinner, M. H., see Wong, Y. W. J. 571(1991)318
 Skipper, P. L., see Day, B. W. 562(1991)563
 Skolnick, P., see Miller, R. 578(1992)103
 Skolnik, V., see Haanappel, V. A. C. 562(1991)531
 Skriver, L., see Bayne, S. 562(1991)391
 Skultéty, L. and Toman, R.
 Improved procedure for the drying and storage of polyacrylamide slab gels 582(1992)249
 Skutil, J., see Gasparič, J. 558(1991)415
 Sky-Peck, H., see Kang, K. 567(1991)57
 Slabova, O. I., see Rezanaka, T. 558(1991)215
 Slacanin, I., Marston, A., Hostettmann, K., Delabays, N. and Darbellay, C.
 Isolation and determination of flavonol glycosides from *Epilobium* species 557(1991)391
 Slangen, C. J., see Visser, S. 599(1992)205
 Slanina, P., see Hellenäs, K.-E. 573(1992)69
 Šleboda, M., see Wodecki, Z. J. 558(1991)302
 Śledzińska, I., see Sadlej-Sosnowska, N. 595(1992)53
 Slegel, P., see Maguire, M. H. 575(1992)243
 Ślégel, P., see Valkó, K. 592(1992)59
 Sloyer, B. D. and Goldberg, J. I.
 Determination of γ -glutamyl conjugates of monoamines by means of high-performance liquid chromatography with electrochemical detection and application to gastropod tissues 567(1991)49
 Smallwood, R. A., see Martin, G. 574(1992)205
 Šmejkal, J., see Linhart, I. 578(1992)215
 Smid, P., see De Ruiter, G. A. 584(1992)69
 Šmidl, P., Plicka, J. and Kleinmann, I.
 Separation of HEMA modified for immobilized metal ion affinity chromatographic separation of proteins 598(1992)15
 Smit, L. A., see Van 't Klooster, G. A. E. 571(1991)157
 Smith, C. G., see Berry, M. J. 587(1991)161
 Smith, D. A., see Walker, D. K. 568(1991)475
 Smith, D. E., see Janiczek, N. 571(1991)179
 Smith, D. L., see Teffera, Y. 577(1992)69
 Smith, I., see Berry, M. J. 587(1991)161
 Smith, I. D. and Waters, D. G.
 Determination of topanol antioxidants in methacrylates using capillary gas-liquid chromatography 596(1992)290
 Smith, M. L., see Kuhlman, Jr., J. J. 578(1992)207
 Smith, P. C., see Wang, L. H. 579(1992)259
 Smith, P. K., see Nevens, J. R. 597(1992)247
 Smith, R. D., Udseth, H. R., Barinaga, C. J. and Edmonds, C. G.
 Instrumentation for high-performance capillary electrophoresis-mass spectrometry 559(1991)197
 Smith, R. L., see Zhang, A. Q. 584(1992)141
 Smith, R. M. and Burford, M. D.
 Optimization of supercritical fluid extraction of volatile constituents from a model plant matrix 600(1992)175
 Smith, R. M., Westlake, J. P., Gill, R. and Osselton, M. D.
 Retention reproducibility of basic drugs in high-performance liquid chromatography on a silica column with a methanol-high-pH buffer eluent. Changes in selectivity with the age of the stationary phase 592(1992)85
 Smith, S. C., Strasters, J. K. and Khaledi, M. G.
 Influence of operating parameters on reproducibility in capillary electrophoresis 559(1991)57
 Smith, U., see Lönnroth, P. 568(1991)419
 Smith-Rogers, J. A., Tong, W. P., Duafala, M. E., Markman, M. and Bertino, J. R.
 High-performance liquid chromatographic method for the simultaneous measurement of floxuridine and fluorouracil in human body fluids 566(1991)147
 Smolková-Keulemansová, E., see Jelínek, I. 557(1991)215
 Smolkova-Keulemansova, E., see Snopek, J. 559(1991)215
 Smuts, C. M. and Tichelaar, H. Y.
 Simple thin-layer chromatographic purification procedure for the determination of cholesterol ester fatty acid compositions 564(1991)272
 Smyth, J. F., see Anderson, L. 574(1992)150
 Smyth, M. R., see Kinahan, I. M. 565(1991)297
 Smyth, M. R., see O'Shea, T. J. 593(1992)305
 Smyth, M. R., see Sagar, K. A. 577(1992)109
 Snider, J., Neville, C., Yuan, L.-C. and Bullock, J.
 Characterization of the heterogeneity of polyethylene glycol-modified superoxide dismutase by chromatographic and electrophoretic techniques 599(1992)141
 Snini, A., Fahimi, A., Mouloungui, Z., Delmas, M. and Gaset, A.
 Separation and preparative isolation of phenolic dialdehydes by on-line overpressured layer chromatography 590(1992)369
 Snopek, J., Soini, H., Novotny, M., Smolkova-Keulemansova, E. and Jelinek, I.
 Selected applications of cyclodextrin selectors in capillary electrophoresis 559(1991)215
 Snopek, J., see Jelínek, I. 557(1991)215
 Snyder, L. R., see Chloupek, R. C. 594(1992)65
 Snyder, L. R., see Cox, G. B. 590(1992)17
 Snyder, L. R., see Lewis, J. A. 592(1992)183
 Snyder, L. R., see Lewis, J. A. 592(1992)197
 Snyder, L. R., see Lisi, D. D. 555(1991)1
 Soane, D. S., see Grossman, P. D. 559(1991)257
 Soares, Jr., J. H., see Craft, N. E. 589(1992)171
 Sochor, J., see Klimeš, J. 584(1992)221
 Søreberg, H., see Vinther, A. 559(1991)3
 Søreberg, H., see Vinther, A. 559(1991)27
 Søreberg, H., see Vinther, A. 589(1992)315
 Sofia, R. D., see Clark, L. A. 573(1992)113
 Sohn, D.-R., see Kobayashi, K. 579(1992)299
 Soini, H., Tsuda, T. and Novotny, M. V.
 Electrochromatographic solid-phase extraction for determination of cimetidine in serum by micellar electrokinetic capillary chromatography 559(1991)547

- Soini, H., see Snopek, J. 559(1991)215
- Soják, L., Farkaš, P., Ostrovský, I., Janák, J. and Chrétien, J. R.
Capillary gas chromatography of C₅-C₁₃ branched alkynes on squalane and liquid crystal stationary phases 557(1991)241
- Solodenko, V. A., see Galushko, S. V. 553(1991)143
- Solodenko, V. A., see Galushko, S. V. 600(1992)79
- Solomon, B., Raviv, O., Leibman, E. and Fleminger, G.
Affinity purification of antibodies using immobilized FB domain of protein A 597(1992)257
- Solomon, B., see Fleminger, G. 597(1992)263
- Soloshonok, V. A., see Galushko, S. V. 592(1992)345
- Somogyi, A. A., see Milne, R. W. 565(1991)457
- Somsen, G. W., Van de Nesse, R. J., Gooijer, C., Brinkman, U. A. T., Velthorst, N. H., Visser, T., Kootstra, P. R. and De Jong, A. P. J. M.
Spray jet assembly interface for the coupling of reversed-phase narrow-bore liquid chromatography and Fourier transform infrared spectrometry 552(1991)635
- Sonck, W., see Mertens, K. 565(1991)149
- Soncrant, T. T., see Deutsch, J. 579(1992)93
- Sonders, R. C., see Chu, S.-Y. 571(1991)199
- Sonezaki, S., see Ishida, J. 598(1992)203
- Sonnet, P. E., Dudley, R. L., Osman, S., Pfeffer, P. E. and Schwartz, D.
Configuration analysis of unsaturated hydroxy fatty acids 586(1991)255
- Soons, P. A., see Roosemalen, M. C. M. 565(1991)516
- Sorisio, D., see Foglia, J. P. 572(1991)247
- Sotiriou-Leventis, C., see Li, W. 588(1991)273
- Souhodolova, V. I., see Berezkina, L. G. 552(1991)319
- Soumillion, J. P., see Tazerouti, A. 596(1992)132
- Southwell, I.
Eucalyptus oils: use, chemistry, distillation and marketing (edited by D. J. Boland, J. J. Brophy and A. P. N. House) (Book Review) 598(1992)316
- Sovocool, G. W., see Donnelly, J. R. 594(1992)269
- Spaethe, S. M., see VanAlstyne, E. L. 579(1992)37
- Spahn-Langguth, H., Hahn, G., Mutschler, E., Möhrke, W. and Langguth, P.
Enantiospecific high-performance liquid chromatographic assay with fluorescence detection for the monoamine oxidase inhibitor tranlylcypromine and its applicability in pharmacokinetic studies 584(1992)229
- Spahn-Langguth, H., see Liebmann, B. 572(1991)181
- Spanedda, L., see Cabras, P. 585(1991)164
- Spencer, M. J., see Buck, C. F. 594(1992)225
- Spiegel, L. B., Hadjimichael, J. and Rossomando, E. F.
Assay of sialyltransferase activity by reversed-phase ion-pair high-performance liquid chromatography 573(1992)23
- Spinazzola, J., see Rop, P. P. 573(1992)87
- Spinelli, R., see Sarati, S. 563(1991)323
- Spinetti, M., see Raspi, G. 593(1992)119
- Spitz, D., see Kinter, M. 578(1992)9
- Spöndlin, C., see Küsters, E. 576(1992)179
- Spranger, B. and Metzler, M.
Disposition of 17 β -trenbolone in humans 564(1991)485
- Spurway, T. D., see Witherow, L. 553(1991)497
- Sreekumaran Nair, K., see Quinn Baumann, P. 573(1992)11
- Staba, E. J., see Huh, H. 600(1992)364
- Stacey, C. C., see Balogh, M. P. 562(1991)73
- Staffeldt, B., Brockmöller, J. and Roots, I.
Determination of S-carboxymethyl-L-cysteine and some of its metabolites in urine and serum by high-performance liquid chromatography using fluorescent pre-column labelling 571(1991)133
- Stafford, J. D., see Olsen, B. A. 594(1992)203
- Stafford, R. J., see Erdmann, G. R. 571(1991)149
- Ståhlberg, J., see Bartha, Á. 552(1991)13
- Stalker, D. J., see Sturm, G. 562(1991)351
- Stancher, B., see Zonta, F. 594(1992)137
- Star-Weinstock, M., see Fleminger, G. 597(1992)263
- Stárek, J.
Theory of linear non-equilibrium chromatography with beds of a finite length 591(1992)273
- Staroverov, S. M., see Fadeev, A. Y. 558(1991)31
- Staroverov, S. M., see Fadeev, A. Y. 596(1992)114
- Starrett, J. E., see Russell, J. W. 572(1991)321
- Stavert, D. M., see Gurley, L. R. 559(1991)411
- Stebler, T. and Guentert, T. W.
Determination of diazepam and nordazepam in milk and plasma in the presence of oxazepam and temazepam 564(1991)330
- Steed, J. W., see Abraham, M. H. 588(1991)361
- Steel, C. C.
Radio-detection high-performance liquid chromatographic enzyme assay for inhibitors of fungal sterol Δ^14 -reductase 566(1991)435
- Steffenrud, S. and Maylin, G.
Thermospray liquid chromatography-mass spectrometry of corticosteroids 577(1992)221
- Stegehuis, D. S., Tjaden, U. R. and Van der Greef, J.
Analyte focusing in capillary electrophoresis using on-line isotachopheresis 591(1992)341
- Stegeman, G., see Bruin, G. J. M. 559(1991)163
- Stehlík, P. and Houbová, H.
Determination of amidepin in human plasma by reversed-phase high-performance liquid chromatography with fluorescence detection 565(1991)477
- Steijger, O. M., De Jong, G. J., Holthuis, J. J. M. and Brinkman, U. A. T.
On-line electrochemical reagent generation for liquid chromatography with luminol-based chemiluminescence detection 557(1991)13
- Stein, J., Kulemeier, J., Lembcke, B. and Caspary, W. F.
Simple and rapid method for determination of short-chain fatty acids in biological materials by high-performance liquid chromatography with ultraviolet detection 576(1992)53
- Stein, J., see Sander, S. 558(1991)115
- Steinberg, J. J., Cajigas, A. and Brownlee, M.
Enzymatic shot-gun 5'-phosphorylation and 3'-sister phosphate exchange: a two-dimensional thin-layer chromatographic technique to measure DNA deoxynucleotide modification 574(1992)41
- Steinberg, J. J., see Bleiberg, B. 568(1991)301

- Steiner, U., see Lindner, W. 553(1991)373
- Steiner, V., Schär, M., Börnsen, K. O. and Mutter, M.
Retention behaviour of a template-assembled synthetic protein and its amphiphilic building blocks on reversed-phase columns 586(1991)43
- Steinhilber, D., see Gaede, F.-P. 585(1991)53
- Stellaard, F., see Zee, T. 574(1992)335
- Stephany, R. W., see Jansen, E. H. J. M. 580(1992)111
- Stephens, R. D., see Kim, I. S. 589(1991)177
- Stephenson, G. R., see Creaser, C. S. 586(1991)323
- Sternieri, E., see Zuccaro, P. 567(1991)485
- Sternitzke, K. D., Fan, T. Y. and Dunn, D. L.
High-performance liquid chromatographic determination of pilocarpine hydrochloride and its degradation products using a β -cyclodextrin column 589(1992)159
- Steury, J. C., see Erdmann, G. R. 571(1991)149
- Stevens, T. W., see Haanappel, V. A. C. 562(1991)531
- Stevenson, D., see Prapamontol, T. 552(1991)249
- Stevenson, K., see Zhou, G. 568(1991)69
- Stevenson, P. J., Hamelijck, M. A. F., Kadaba, P. K. and Damani, L. A.
Triazolines. XXIII. High-performance liquid chromatographic assay in rat blood for a novel triazoline anticonvulsant (ADD17014) 563(1991)419
- Stewart, D. C., see Fowler, W. K. 558(1991)235
- Stigbrand, T., see Kirchberger, J. 574(1992)237
- Stijnen, A. M., see Heeremans, C. E. M. 554(1991)205
- Stoev, G.
Application and chiral recognition of heptakis (2,6-di-O-methyl-3-O-trifluoroacetyl)- β -cyclodextrin as a stationary phase for the gas chromatographic separation of enantiomers 589(1992)257
- Stohs, S. J., see Shara, M. A. 576(1992)221
- Stokes, K. H., see Webster, L. K. 565(1991)482
- Stokes, M., see Thedford, R. 582(1992)253
- Stoltz, J.-F., see Regnault, V. 576(1992)87
- Stoltz, J. F., see Regnault, V. 584(1992)93
- Stoltz, J.-F., see Rivat, C. 576(1992)71
- Stopher, D. A., see Walker, D. K. 568(1991)475
- Storbeck, L. S., see Khetarpal, V. K. 567(1991)491
- Storme, G. A., see Van Belle, S. J.-P. 576(1992)351
- Storme, G. A., see Van Belle, S. J.-P. 578(1992)223
- Storr-Hansen, E.
Comparative analysis of thirty polychlorinated biphenyl congeners on two capillary columns of different polarity with non-linear multi-level calibration 558(1991)375
- Stoskopf, M. K., see Tyczkowska, K. L. 576(1992)305
- Štovičková, J., Podrazký, V. and Trnavský, K.
Affinity chromatography of serine proteinases from the bovine intervertebral disc on BPTI-Separon HEMA 1000 575(1992)229
- Stránský, K., see Jursík, T. 586(1991)315
- Strašák, M., Novomeský, P. and Bystrický, S.
Rapid optical resolution of electrically neutral cobalt(III) chelate complexes by gel permeation chromatography 585(1991)337
- Strasters, J. K., Kim, S.-T. and Khaledi, M. G.
Multiparameter optimizations in micellar liquid chromatography using the iterative regression optimization strategy 586(1991)221
- Strasters, J. K., see Smith, S. C. 559(1991)57
- Stratford, M. R. L. and Dennis, M. F.
High-performance liquid chromatographic determination of nicotinamide and its metabolites in human and murine plasma and urine 582(1992)145
- Straub, R., Pfister, M. and Arm, H.
New chiral covalently bonded, π -donor stationary phases for high-performance liquid chromatography, based on derivatives of optically active 1-(1-naphthyl)ethylamine. Part II 585(1991)195
- Straub, R., see Brügger, R. 557(1991)163
- Strege, M. A. and Lagu, A. L.
Anion-exchange chromatography of DNA restriction fragments 555(1991)109
- Streng, J. R., see Gutnikov, G. 587(1991)292
- Strohmaier, W., see Gasser, H. 562(1991)257
- Strolin Benedetti, M., see Allievi, C. 568(1991)271
- Strolin Benedetti, M., see Breda, M. 578(1992)309
- Strolin Benedetti, M., see Pianezzola, E. 574(1992)170
- Strömvall, A.-M. and Petersson, G.
Protection of terpenes against oxidative and acid decomposition on adsorbent cartridges 589(1992)385
- Strong, J. M., see Anderson, L. W. 581(1992)195
- Struzinsky, R., see Deyl, Z. 569(1991)63
- Struzinsky, R., see Hanis, T. 553(1991)93
- Struzinsky, R., see Hodny, Z. 578(1992)53
- Stuart, J. D., see Lisi, D. D. 555(1991)1
- Stuart, S. G., see Huang, X. C. 600(1992)289
- Stubbs, A. P., Murphy, G. M. and Wilkinson, M. L.
Isocratic high-performance liquid chromatographic measurement of optimal 5α -steroid reductase activity in Hep-G2 cells 570(1991)293
- Stuchlík, J., see Jegorov, A. 558(1991)311
- Štulík, K., Pacáková, V. and Wang, H.
High-performance liquid chromatography of amino acids and dipeptides on new ion exchangers of the HEMA series 552(1991)439
- Štulík, K., see Böswart, J. 571(1991)19
- Štulík, K., see Švagrová, I. 563(1991)95
- Stump, W. T., see Turk, J. 575(1992)183
- Sturaro, A., Parvoli, G., Zanchetta, S., Doretto, L., Gori, G. and Bartolucci, G. B.
Mass spectrometric and gas and high-performance liquid chromatographic behaviour of an impurity in 2,5-hexanedione 590(1992)223
- Sturkenboom, M. C. J. M., see Hessels, J. 563(1991)1
- Sturm, G., Häberlein, H., Bauer, T., Plaum, T. and Stalker, D. J.
Mass spectrometric and high-performance liquid chromatographic studies of medroxyprogesterone acetate metabolites in human plasma 562(1991)351
- Suarez, A., see Reimer, G. J. 555(1991)315
- Sud, D., Hothi, H. S. and Pannu, B. S.
Role of metal ions in the ligand-exchange separation of amino acids 596(1992)281
- Sudo, K., see Fujimaki, Y. 575(1992)261
- Sudoh, Y., see Yakabe, Y. 558(1991)323
- Sugahara, K. and Kodama, H.
Liquid chromatography-mass spectrometry for simultaneous analyses of iminodipeptides containing an N-terminal or a C-terminal proline 565(1991)408

- Sugahara, K., Ohta, J., Takemura, M. and Kodama, H.
Determination of cystathionine and perhydro-1,4-thiazepine-3,5-dicarboxylic acid in the urine of a patient with cystathioninuria using column liquid chromatography-mass spectrometry 579(1992)318
- Sugahara, K., see Watanabe, H. 568(1991)445
- Sugawara, K., see Ohkubo, T. 573(1992)289
- Sugawara, M., see Kishino, S. 582(1992)246
- Sugii, A., see Etoh, T. 578(1992)292
- Sugimoto, H., see Watanabe, K. 566(1991)109
- Sugino, R., see Hirokawa, T. 585(1991)364
- Sugita, M., see Matsushita, S. 586(1991)363
- Sugo, T., see Kim, M. 585(1991)45
- Sugo, T., see Kim, M. 586(1991)27
- Sullivan, S., see Kinter, M. 578(1992)9
- Sumita, S., see Fujioka, T. 597(1992)429
- Sun, J. J. and Fritz, J. S.
Chemically modified resins for solid-phase extraction 590(1992)197
- Sunahara, H., see Ito, K. 598(1992)237
- Sunzeri, F. J., see Schwartz, H. E. 559(1991)267
- Suprynowicz, Z., Pilorz, K. and Lodkowski, R.
Evaluation of the application of liquid-phase titration to the examination of the adsorption activity of silica-based high-performance liquid chromatographic packings 552(1991)463
- Surber, B. W., see Thomas, S. B. 586(1991)265
- Surtel, D. A. M., see Van Nieuwenhoven, F. A. 570(1991)173
- Süss, S., Seiler, W., Hiemke, C., Schöllnhammer, G., Wetzel, H. and Hillert, A.
Determination of benperidol and its reduced metabolite in human plasma by high-performance liquid chromatography and electrochemical detection 565(1991)363
- Šušterčić, M., see Alajbeg, A. 557(1991)255
- Suter, M. J.-F., DaGue, B. B., Moore, W. T., Lin, S.-N. and Caprioli, R. M.
Recent advances in liquid chromatography-mass spectrometry and capillary zone electrophoresis-mass spectrometry for protein analysis 553(1991)101
- Suttner, J., Dyr, J. E., Pristach, J. and Holada, K.
Versatile modular microelectrophoresis system and the simultaneous preparation of large numbers of polyacrylamide gels 563(1991)184
- Suttner, J., see Dyr, J. E. 563(1991)124
- Sutton, B. M. and Richardson, R. A.
Assay of timolol in human plasma using gas chromatography with electron-capture detection 581(1992)277
- Suzuki, J., see Miyabara, Y. 574(1992)261
- Suzuki, K., see Honda, S. 597(1992)377
- Suzuki, K., see Watanabe, K. 566(1991)109
- Suzuki, M., see Shimada, K. 558(1991)306
- Suzuki, N., see Shibata, Y. 598(1992)73
- Suzuki, O., see Hattori, H. 564(1991)278
- Suzuki, O., see Hattori, H. 579(1992)247
- Suzuki, O., see Hattori, H. 581(1992)213
- Suzuki, S., see Honda, S. 588(1991)327
- Suzuki, S., see Honda, S. 597(1992)377
- Suzuki, S., see Miyabara, Y. 574(1992)261
- Suzuki, T., see Matsuda, R. 585(1991)187
- Suzuki, T., see Uda, T. 586(1991)131
- Suzuki, Y.
Electrophoresis of mutant proteins in inherited diseases (Review) 569(1991)407
- Suzuki, Y., see Okamoto, J. 583(1992)53
- Suzuno, R., see Takeo, K. 597(1992)365
- Švagrová, I., Štulík, K., Pacáková, V., Caliceti, P. and Veronese, F. M.
Determination of narciclasine in serum by reversed-phase high-performance liquid chromatography: comparison of amperometric, ultraviolet photometric and fluorescence detection 563(1991)95
- Svátek, E., see Ryska, M. 562(1991)289
- Svec, F., see Dernovaya, L. A. 552(1991)365
- Švec, F., see Tennikova, T. B. 555(1991)97
- Švedas, V. K., see Petrauskas, A. A. 585(1991)3
- Švel-Cerovečki, S., see Alajbeg, A. 557(1991)255
- Svoboda, D., see Nobilis, M. 581(1992)251
- Svoboda, M., Przybylski, M., Schreurs, J., Miyajima, A., Hogeland, K. and Deinzer, M.
Mass spectrometric determination of glycosylation sites and oligosaccharide composition of insect-expressed mouse interleukin-3 562(1991)403
- Swamy, N. S., see Husain, S. 558(1991)435
- Swart, K. J. and Paggis, M.
Automated high-performance liquid chromatographic method for the determination of rifampicin in plasma 593(1992)21
- Swart, K. J., see Brown, L. W. 582(1992)268
- Sweeley, C. C., see Schultz, G. A. 590(1992)329
- Swerdlow, H. P., see Chen, D. Y. 559(1991)237
- Swiader, J. M., see Grunau, J. A. 594(1992)165
- Sydenham, E. W., see Shephard, G. S. 566(1991)195
- Sydenham, E. W., see Shephard, G. S. 574(1992)299
- Syrový, I. and Hodný, Z.
Staining and quantification of proteins separated by polyacrylamide gel electrophoresis (Review) 569(1991)175
- Szabo, I., see Maguire, M. H. 575(1992)243
- Szabó, I., see Róna, K. 573(1992)173
- Szilágyi, J., see Kovács-Hadady, K. 553(1991)459
- Szokoli, F., see Bartha, Á. 552(1991)13

T

- Tabata, M., Murachi, T., Endo, J. and Totani, M.
Use of various types of column reactors for flow-injection analysis 597(1992)435
- Tabata, M., see Miyabara, Y. 574(1992)261

- Tabata, S., see Nishioka, R. 565(1991)237
- Tabata, T., see Yamaguchi, M. 575(1992)123
- Tabera, J., Jiménez, B., Hernández, L. M. and González, M. J.
Optimization of the gas chromatographic analysis of a standard mixture of polychlorodibenzo-*p*-dioxins and polychlorodibenzofurans 557(1991)481
- Tadano, J., see Kawasaki, S. 595(1992)193
- Tadeo, J. L., see Sánchez-Brunete, C. 552(1991)235
- Tadeo, J. L., see Sanchez-Brunete, C. 562(1991)525
- Tadey, T. and Purdy, W. C.
Characterization of plasma apolipoproteins by capillary electrophoresis 583(1992)111
- Taga, A., see Honda, S. 590(1992)364
- Taga, A., see Honda, S. 597(1992)377
- Taga, M., see Tanaka, S. 587(1991)364
- Tagliaro, F., Dorizzi, R., Ghielmi, S. and Marigo, M.
Direct injection high-performance liquid chromatographic method with electrochemical detection for the determination of ethanol and methanol in plasma using an alcohol oxidase reactor 566(1991)333
- Tagliaro, F., Lubli, G., Ghielmi, S., Franchi, D. and Marigo, M.
Chromatographic methods for blood alcohol determination (Review) 580(1992)161
- Tagliaro, F., Schiavon, G., Bontempelli, G., Carli, G. and Marigo, M.
Improved high-performance liquid chromatographic determination with amperometric detection of α -amanitin in human plasma based on its voltammetric study 563(1991)299
- Tagliaro, F., Schiavon, G., Dorizzi, R. and Marigo, M.
Development of post-column enzymic reactors with immobilized alcohol oxidase for use in the high-performance liquid chromatographic assay of alcohols with electrochemical detection 563(1991)11
- Tagliaro, F., see Dorizzi, R. 580(1992)279
- Tagliaro, F., see Krstulovic, A. M. 566(1991)273
- Taimr, L. and Prusíková, M.
Chromatographic behaviour of the antidegradant ethoxyquin and its transformation products 587(1991)355
- Taiocchi, L., see Gaspari, F. 577(1992)123
- Taka, T., see Hirokawa, T. 555(1991)247
- Takaba, H., see Hirokawa, T. 555(1991)321
- Takada, K., see Shimohigashi, Y. 597(1992)425
- Takagi, T., see Takahara, E. 576(1992)174
- Takahagi, H., see Watanabe, M. 597(1992)309
- Takahama, S., see Sato, J. 575(1992)63
- Takahara, E., Fukuoka, H., Takagi, T., Nagata, O. and Kato, H.
Simultaneous determination of a new gastrointestinal prokinetic agent (HSR-803) and its metabolites in human serum and urine by high-performance liquid chromatography using automated column-switching 576(1992)174
- Takahashi, A., see Niwa, T. 578(1992)109
- Takahashi, H., see Furuta, T. 562(1991)363
- Takahashi, H., see Shinkai, H. 570(1991)203
- Takahashi, K., see Haginaka, J. 598(1992)67
- Takahashi, K., see Hashimoto, A. 582(1992)41
- Takahashi, M., see Ohsawa, K. 597(1992)285
- Takahata, Y., see Yakabe, Y. 558(1991)323
- Takamatsu, T., Yamazaki, K., Kayano, M., Takenaka, F., Hasui, M. and Ohkawa, T.
Determination of eperisone in human plasma by gas chromatography-mass spectrometry 584(1992)261
- Takasaki, W., see Ishibashi, M. 562(1991)613
- Takashima, E., see Hattori, H. 579(1992)247
- Takashima, E., see Hattori, H. 581(1992)213
- Takasugi, M., see Monde, K. 598(1992)147
- Takatori, T., see Terazawa, K. 565(1991)453
- Takeba, K., Matsumoto, M. and Nakazawa, H.
Determination of nitroxynil in cow milk by reversed-phase high-performance liquid chromatography with dual-electrode coulometric detection 596(1992)67
- Takeda, N. and Akiyama, Y.
Pre-column derivatization of sulfa drugs with fluorescamine and high-performance liquid chromatographic determination at their residual levels in meat and meat products 558(1991)175
- Takeda, N., Pomerantz, S. C. and McCloskey, J. A.
Detection of ribose-methylated nucleotides in enzymatic hydrolysates of RNA by thermospray liquid chromatography-mass spectrometry 562(1991)225
- Takeda, N., see Niwa, T. 578(1992)109
- Takeda, S., see Wakui, Y. 575(1992)131
- Takeda, Y., see Toyo'oka, T. 588(1991)61
- Takehara, Y., see Okuda, T. 567(1991)141
- Takeichi, Y., see Iida, Y. 571(1991)277
- Takemura, M., see Sugahara, K. 579(1992)318
- Takenaka, F., see Takamatsu, T. 584(1992)261
- Takeo, K., Nakamura, K. and Suzuno, R.
Separation of monoclonal antibodies from antihapten antisera by two-dimensional affinity electrophoresis 597(1992)365
- Takeo, K., see Mimura, Y. 597(1992)345
- Takeo, K., see Nakamura, K. 597(1992)351
- Taketa, K.
Clinical applications of lectin affinity electrophoresis (Review) 569(1991)229
- Takeuchi, T. and Nagae, N.
Stationary phase effect on enantioselectivity of dansyl phenylalanine in microcolumn liquid chromatography with γ -cyclodextrin as mobile phase additive 595(1992)121
- Takeuchi, T., see Hu, W. 557(1991)441
- Takeuchi, T., see Tanaka, S. 597(1992)443
- Taki, M., see Kamahori, M. 567(1991)351
- Takigiku, R. and Schneider, R. E.
Reproducibility and quantitation of separation for ribonucleoside triphosphates and deoxyribonucleoside triphosphates by capillary zone electrophoresis 559(1991)247
- Takigiku, R., see Rodriguez, P. A. 563(1991)271
- Takimoto, N., see Sano, A. 582(1992)131
- Takitani, S., see Sano, A. 582(1992)131

- Takkenberg, B., Endert, E., Van Ingen, H. E. and Ackermans, M.
Improved method for the determination of serotonin in plasma by high-performance liquid chromatography using on-line sample pre-treatment 565(1991)430
- Talaska, G., Roh, J. H. and Getek, T.
³²P-Postlabelling and mass spectrometric methods for analysis of bulky, polyaromatic carcinogen-DNA adducts in humans (Review) 580(1992)293
- Tallaksen, C. M. E., Bøhmer, T., Bell, H. and Karlsen, J.
Concomitant determination of thiamine and its phosphate esters in human blood and serum by high-performance liquid chromatography 564(1991)127
- Talmadge, K. W., Dunn, L. C., Abouelezz, M., Morehead, H., Navvab, M., Ordunez, C., Tisch, T. L. and Siebert, C. J.
Characterization of synthetic macroporous packing materials in low-pressure cartridges and columns 590(1992)83
- Talmadge, K. W., see Morehead, H. W. 587(1991)171
- Tam, Y. K., see Hussain, M. D. 582(1992)203
- Tamaki, M., Akabori, S. and Muramatsu, I.
Studies on cyclic peptides related to graminin by reversed-phase high-performance liquid chromatography 574(1992)65
- Tamaru, T., see Iida, T. 558(1991)451
- Tamura, Y., see Hirata, N. 592(1992)93
- Tan, H. G. H., see Tan, H. S. I. 568(1991)145
- Tan, H. S. I., Manning, M. A., Hahn, M.-K., Tan, H. G. H. and Kotagal, U. R.
Determination of benzyl alcohol and its metabolite in plasma by reversed-phase high-performance liquid chromatography 568(1991)145
- Tan, S. T., see Boniface, P. J. 584(1992)199
- Tanaka, E.
Simultaneous determination of caffeine and its primary demethylated metabolites in human plasma by high-performance liquid chromatography 575(1992)311
- Tanaka, E. and Misawa, S.
Improved method for the determination of trimethadione and its demethylated metabolite, dimethadione, in human serum by gas chromatography 584(1992)267
- Tanaka, N., see Honda, A. 565(1991)53
- Tanaka, N., see Kimata, K. 558(1991)19
- Tanaka, N., see Kimata, K. 594(1992)87
- Tanaka, N., see Kimata, K. 595(1992)77
- Tanaka, S., Kaneta, T., Taga, M., Yoshida, H. and Ohtaka, H.
Capillary tube isotachopheric separation of catecholamines using cyclodextrin in the leading electrolyte 587(1991)364
- Tanaka, S., Takeuchi, T. and Rechnitz, G. A.
Non-isotopic receptor assay for benzodiazepines using a biotin-labeled ligand and biotin-immobilized microtiter plate 597(1992)443
- Tanaka, S., see Ohsawa, K. 597(1992)285
- Tang, S. W., see Cheung, S. W. 564(1991)213
- Taniguchi, H., see Katayama, M. 585(1991)219
- Tanikawa, M., see Toyoda, H. 565(1991)141
- Tanimoto, T., see Koizumi, K. 595(1992)340
- Tannenbaum, S. R., see Day, B. W. 562(1991)563
- Tanner, J. K., see Woodcock, E. A. 581(1992)134
- Tanzer, J. M., see Weaver, M. L. 581(1992)293
- Taoka, M., see Isobe, T. 588(1991)115
- Tapiero, C., see Voirin, S. G. 590(1992)313
- Tarbin, J. A. and Shearer, G.
High-performance liquid chromatographic determination of imidocarb in cattle kidney with cation-exchange clean-up 577(1992)376
- Tarbin, J. A. and Shearer, G.
Improved high-performance liquid chromatographic procedure for the determination of lasalocid in chicken tissues and egg using polymeric and porous graphitic carbon columns 579(1992)177
- Tarditi, L., Camagna, M., Parisi, A., Vassarotto, C., DeMonte, L. B., Letarte, M., Malavasi, F. and Mariani, M.
Selective high-performance liquid chromatographic purification of bispecific monoclonal antibodies 599(1992)13
- Tateishi, M., see Komatsu, S. 568(1991)460
- Tatekawa, K., see Otsuka, K. 559(1991)209
- Tatematsu, A., see Niwa, T. 578(1992)109
- Tatsuno, Y., see Kane, M. 569(1991)297
- Tavazzi, D., see Pittalis, S. 573(1992)29
- Taverna, M., Baillet, A., Werner, R. and Bayloq-Ferrier, D.
Liquid chromatographic method for the determination of the carbohydrate moiety of glycoproteins. Application to α_1 -acid glycoprotein and tissue plasminogen activator 558(1991)105
- Tayeb, N., see Traiffort, E. 571(1991)231
- Taylor, A., see Brownsill, R. 562(1991)267
- Taylor, G. W., see Jones, H. A. 570(1991)361
- Taylor, G. W., see Murray, S. 567(1991)289
- Taylor, I. K., see Murray, S. 567(1991)289
- Taylor, L. C. E., see Chang, S. Y. 562(1991)111
- Taylor, P. J., Charles, B. G., Norris, R., Salm, P. and Ravenscroft, P. J.
Measurement of dothiepin and its major metabolites in plasma by high-performance liquid chromatography 581(1992)152
- Taylor, R. B., Reid, R. G. and Gould, I. M.
Determination of teicoplanin in plasma using microbore high-performance liquid chromatography and injection-generated gradients 563(1991)451
- Taylor, S. M., see Wright, M. R. 565(1991)225
- Tazerouti, A., Rahal, S. and Soumillion, J. P.
Simultaneous gas chromatographic analysis of heptyl chloride-heptanesulphonyl chloride isomeric mixtures 596(1992)132
- Tchapla, A., see Aubert, M.-C. 557(1991)47
- Tchapla, A., see Heron, S. 556(1991)219
- Tchapla, A., see Lesellier, E. 557(1991)59
- Tebbett, I. R., see Browne, S. P. 575(1992)158
- Tebbett, I. R., see Chari, G. 571(1991)263
- Teeuwesen, J., see Van der Horst, F. A. L. 567(1991)161
- Teffera, Y., Baird, W. M. and Smith, D. L.
Detection of benzo[a]pyrene sulfate and glucuronide conjugates in cell culture medium by directly coupled microbore high-performance liquid chromatography-fast atom bombardment mass spectrometry 577(1992)69

- Tegtmeier, F., see Weber, C. 562(1991)599
- Tegtmeier, F., see Weber, C. 577(1992)1
- Teh, L.-C. and Froger, M.
Evaluation of the chromatographic procedure for the preparation of a high-purity C1-esterase inhibitor concentrate from cryosupernatant plasma 582(1992)65
- Tejedor, M. C., Delgado, C., Grupeli, M. and Luque, J.
Affinity partitioning of erythrocytic phosphofructokinase in aqueous two-phase systems containing poly(ethylene glycol)-bound Cibacron Blue. Influence of pH, ionic strength and substrates/ effectors 589(1992)127
- Tellez, S., Forges, N., Roussin, A. and Hernandez, L.
Coupling of microdialysis with capillary electrophoresis: a new approach to the study of drug transfer between two compartments of the body in freely moving rats 581(1992)257
- Ten Bokkel Huinink, W. W., see Van Tellingen, O. 553(1991)47
- Ten Bokkel Huinink, W. W., see Van Tellingen, O. 574(1992)293
- Tenczer, J., see Köppel, C. 562(1991)207
- Tenczer, J., see Köppel, C. 563(1991)73
- Ten Hoor, F., see De Kok, T. M. C. M. 580(1992)135
- Tennikova, T. B., Bleha, M., Švec, F., Almazova, T. V. and Belenkii, B. G.
High-performance membrane chromatography of proteins, a novel method of protein separation 555(1991)97
- Teplý, J., see Cigánek, M. 588(1991)225
- Terabe, S., see Nishi, H. 553(1991)503
- Terabe, S., see Otsuka, K. 559(1991)209
- Teraoka, T., see Yao, K. 581(1992)11
- Teraya, Y., see Goto, J. 585(1991)281
- Terazawa, K., Kaji, H., Akabane, H. and Takatori, T.
Determination of dimethyl sulphide in blood and adipose tissue by headspace gas analysis 565(1991)453
- Terrisse, M. P., see Sautou, V. 571(1991)298
- Terry, Jr., A. V., Silks III, L. A., Dunlap, R. B., Odom, J. D. and Kosh, J. W.
Synthesis of novel selenium-containing choline and acetylcholine analogues and their quantitation using a pyrolysis-gas chromatography-mass spectrometry assay 585(1991)101
- Terzian, J. A., see Aiken, J. H. 584(1992)181
- Terzuoli, L., Leoncini, R., Vannoni, D., Marinello, E. and Pagani, R.
High-performance liquid chromatography of thiazolidinic compounds obtained by condensation of pyridoxal 5'-phosphate or pyridoxal with amino thiols (L- or D-cysteine, cysteamine, L-cysteine ethyl ester) 586(1991)153
- Teshima, E., see Nomura, H. 572(1991)333
- Tešić, Ž. L., Janjić, T. J. and Čelap, M. B.
Effect of the substituents of β -diketonato ligands on R_f values of tris(chelate) transition metal complexes obtained by normal- and reversed-phase thin-layer chromatography on unmodified silica gel 585(1991)359
- Testa, B., see El Tayar, N. 556(1991)181
- Testa, E., see Pellegrino, S. 592(1992)279
- Testa, E., see Petrarulo, M. 579(1992)324
- Tett, S. E., see McLachlan, A. J. 570(1991)119
- Thakkar, J. K., see Janero, D. R. 573(1992)207
- Tharakan, J., Highsmith, F., Clark, D. and Drohan, W.
Physical and biochemical characterization of five commercial resins for immunoaffinity purification of factor IX 595(1992)103
- Tharakan, J., see Orthner, C. L. 558(1991)55
- Theford, R., Fautleroy, L. P., Murray, Y. L., Stokes, M. and Thornton, C. L.
High-performance liquid chromatographic monitoring of metabolic products resulting from the treatment of mouse hepatoma cells with N⁶-cycloalkylated nucleosides 582(1992)253
- Thénot, J. P., see Ascalone, V. 581(1992)237
- Théorêt, Y. and Varin, F.
Simple, rapid and selective method using high-performance liquid chromatography for the determination of bretylium in plasma 575(1992)162
- Théorêt, Y., see Ducharme, J. 573(1992)79
- Théorêt, Y., see Varin, F. 574(1992)57
- Thibault, P., see Pleasance, S. 591(1992)325
- Thiébaud, D., see Drogue, S. 593(1992)363
- Thiébaud, D., see Villermet, A. 557(1991)85
- Thiel, P. G., see Shephard, G. S. 566(1991)195
- Thiel, P. G., see Shephard, G. S. 574(1992)299
- Thoma, R. S., see Crimmins, D. L. 599(1992)51
- Thomare, P., Wang, K., Van Der Meersch-Mougeot, V. and Diquet, B.
Sensitive micromethod for column liquid chromatographic determination of fluoxetine and norfluoxetine in human plasma 583(1992)217
- Thomas, C. G., Ward, S. A. and Edwards, G.
Selective determination, in plasma, of artemether and its major metabolite, dihydroartemisinin, by high-performance liquid chromatography with ultraviolet detection 583(1992)131
- Thomas, L. C. and Ramus, T. L.
Measurements using gas chromatography with coelution and dual-isotope atomic emission detection 586(1991)309
- Thomas, L. C. and Weichmann, W.
Quantitative measurements via co-elution and dual-isotope detection by gas chromatography-mass spectrometry 587(1991)255
- Thomas, L. C., see Hein, S. J. 557(1991)39
- Thomas, R. L., see Kunkel, M. E. 589(1992)135
- Thomas, S. B. and Surber, B. W.
Preparative separation and analysis of the enantiomers of [³H]Abbott-69992, an HIV anti-infective nucleoside, by ligand-exchange high-performance liquid chromatography 586(1991)265
- Thompson, R. A., see Andersson, S. 591(1992)65
- Thompson, S., see Allen, E. 578(1992)175
- Thomsen, J., see Bayne, S. 562(1991)391
- Thormann, W., Caslavská, J., Molteni, S. and Chmelík, J.
Capillary isoelectric focusing with electroosmotic zone displacement and on-column multichannel detection 589(1992)321

- Thormann, W., Minger, A., Molteni, S., Caslavská, J. and Gebauer, P.
Determination of substituted purines in body fluids by micellar electrokinetic capillary chromatography with direct sample injection 593(1992)275
- Thormann, W., see Caslavská, J. 585(1991)145
- Thormann, W., see Caslavská, J. 594(1992)361
- Thormann, W., see Chmelik, J. 600(1992)297
- Thormann, W., see Chmelik, J. 600(1992)305
- Thormann, W., see Gebauer, P. 558(1991)423
- Thormann, W., see Meier, P. 559(1991)505
- Thornton, C. L., see Thedford, R. 582(1992)253
- Thörnwall, M., see Silberring, J. 554(1991)83
- Thorsrud, A. K., see Jellum, E. 559(1991)455
- Thuaud, N., Sebillé, B., Deratani, A. and Lelievre, G.
Retention behavior and chiral recognition of β -cyclodextrin-derivative polymer adsorbed on silica for warfarin, structurally related compounds and Dns-amino acids 555(1991)53
- Thuillier, A., see Fernandez, C. 572(1991)195
- Thurin, J., see Otvos, Jr., L. 599(1992)43
- Thurl, S., Offermanns, J., Müller-Werner, B. and Sawatzki, G.
Determination of neutral oligosaccharide fractions from human milk by gel permeation chromatography 568(1991)291
- Tichá, M., see Turková, J. 597(1992)19
- Tichelaar, H. Y., see Smuts, C. M. 564(1991)272
- Tikhonov, Y. V., see Grune, T. 563(1991)53
- Tikhonov, Y. V., see Ziegler, M. 563(1991)172
- Tillement, J.-P., see Hervé, F. 577(1992)43
- Timbrell, J. A., see Preece, N. E. 573(1992)227
- Time, E., see Jellum, E. 559(1991)455
- Timerbaev, A. R., Tsoi, I. G. and Petrukhin, O. M.
Correlation analysis in liquid chromatography of metal chelates. II. One-dimensional retention-mobile phase composition (physicochemical property) models in reversed-phase liquid chromatography 555(1991)163
- Timm, U. and Erdin, R.
Determination of the catechol-O-methyltransferase inhibitor Ro 40-7592 in human plasma by high-performance liquid chromatography with coulometric detection 593(1992)63
- Timperi, R. J., see Burse, V. W. 566(1991)117
- Tinke, A. P., Van der Hoeven, R. A. M., Niessen, W. M. A., Tjaden, U. R. and Van der Greef, J.
Some aspects of peak broadening in particle-beam liquid chromatography-mass spectrometry 554(1991)119
- Tinsley, P. W., O'Dwyer, P. J. and LaCreta, F. P.
Chromatographic analysis of methylmercaptopyrimidine riboside in human plasma and urine 564(1991)303
- Tinsley, P. W., see LaCreta, F. P. 571(1991)271
- Tisch, T. L., see Talmadge, K. W. 590(1992)83
- Tissot, J.-D., see Toussi, A. 582(1992)87
- Tivert, A.-M., see Arvidsson, T. 586(1991)271
- Tjaden, U. R., see Kokkonen, P. S. 565(1991)265
- Tjaden, U. R., see Niessen, W. M. A. 554(1991)3
- Tjaden, U. R., see Pattyn, G. 564(1991)352
- Tjaden, U. R., see Reeuwijk, H. J. E. M. 575(1992)269
- Tjaden, U. R., see Reinhoud, N. J. 574(1992)327
- Tjaden, U. R., see Stegehuis, D. S. 591(1992)341
- Tjaden, U. R., see Tinke, A. P. 554(1991)119
- Tjaden, U. R., see Van Bakergem, E. 598(1992)189
- Tjaden, U. R., see Venizelos, V. 573(1992)259
- Tjaden, U. R., see Verheij, E. R. 554(1991)339
- Tjørnelund, J. and Hansen, S. H.
High-performance liquid chromatographic assay of 5-aminosalicylic acid (5-ASA) and its metabolites N- β -D-glucopyranosyl-5-ASA, N-acetyl-5-ASA, N-formyl-5-ASA and N-butyryl-5-ASA in biological fluids 570(1991)109
- Tjørnelund, J. and Hansen, S. H.
Stability of 5-aminosalicylic acid and its metabolites in plasma at -20°C . Formation of N- β -D-glucopyranosyl-5-aminosalicylic acid 570(1991)224
- Tjørnelund, J., see Hansen, S. H. 556(1991)353
- Tjukavkina, N. A., Geodakyan, S. V., Khvostova, A. I., Kolesnik, J. A. and Glyzin, V. I.
Simple high-performance liquid chromatographic method for the determination of a new phytochemical drug, fellavine, and its metabolites in human and rat plasma and urine 571(1991)312
- Tjukavkina, N. A., see Geodakyan, S. V. 577(1992)371
- Tod, M., Biarez, O., Nicolas, P. and Petitjean, O.
Sensitive determination of josamycin and rokitamycin in plasma by high-performance liquid chromatography with fluorescence detection 575(1992)171
- Tod, M., Legendre, J.-Y., Chalom, J., Kouwatli, H., Poulou, M., Farinotti, R. and Mahuzier, G.
Primary and secondary amine derivatization with luminarins 1 and 2: separation by liquid chromatography with peroxyoxalate chemiluminescence detection 594(1992)386
- Todorić, A., see Alajbeg, A. 557(1991)255
- Todorov, T. K., see Beran, M. 558(1991)265
- Toguzov, R. T., see Grune, T. 563(1991)53
- Toguzov, R. T., see Ziegler, M. 563(1991)172
- Tohda, K., see Watanabe, K. 566(1991)109
- Tohma, M., see Honda, A. 565(1991)53
- Tohma, M., see Ikegawa, S. 577(1992)229
- Toida, T., see Akiyama, H. 579(1992)203
- Toman, R., see Škultěty, Ľ. 582(1992)249
- Tomasella, F. P., Zuting, P. and Cline Love, L. J.
Determination of sun-screen agents in cosmetic products by micellar liquid chromatography 587(1991)325
- Tomassen, M. J. H., see Ackermans, M. T. 596(1992)101
- Tomba, C., see Anelli, P. L. 589(1992)346
- Tomer, K. B., see Deterding, L. J. 554(1991)73
- Tomer, K. B., see Deterding, L. J. 554(1991)329
- Tominaga, N., see Flesch, G. 568(1991)261
- Tomita, M., Okuyama, T., Nigo, Y., Uno, B. and Kawai, S.
Determination of glyphosate and its metabolite, (aminomethyl)phosphonic acid, in serum using capillary electrophoresis 571(1991)324
- Tomita, M., Okuyama, T., Watanabe, S., Uno, B. and Kawai, S.
High-performance liquid chromatographic determination of glyphosate and (aminomethyl)phosphonic acid in human serum after conversion into *p*-toluenesulphonyl derivatives 566(1991)239

- Tomita, N., see Noguchi, H. 583(1992)266
- Tomizawa, H., see Sakihama, N. 597(1992)147
- Tomokuni, K., Hirai, Y. and Ichiba, M.
Fluorimetric determination of hepatic δ -aminolevulinic acid synthase activity by high-performance liquid chromatography 567(1991)65
- Tomori, É., Horváth, G., Pátfalusi, M., Mészáros, S. and Vereczkey, L.
Pharmacokinetic and metabolism studies on girisopam by chromatographic and spectrometric methods in humans 578(1992)91
- Tomsová, Z.
Thin-layer chromatography of urinary 17-oxosteroids using dansylhydrazine as a prelabelling reagent 570(1991)396
- Tong, W. P., see Smith-Rogers, J. A. 566(1991)147
- Tong, Y.-Y., see Oshima, T. 585(1991)255
- Tønnesen, F., see Johansen, M. 573(1992)283
- Tonucci, L. H., see Khachik, F. 582(1992)153
- Toome, D. W., see Levison, P. R. 590(1992)49
- Topalova, I., see Gavrilova, T. B. 552(1991)179
- Toribio, F., see Alhama, J. 586(1991)51
- Toribio, F., see Alhama, J. 589(1992)121
- Toribio, F., see Pascual, P. 581(1992)49
- Torlop, B., see Wodecki, Z. J. 558(1991)302
- Tornaues, J., see Rosenberg, C. 552(1991)265
- Torok-Both, G. A., Baker, G. B., Coutts, R. T., McKenna, K. F. and Aspeslet, L. J.
Simultaneous determination of fluoxetine and norfluoxetine enantiomers in biological samples by gas chromatography with electron-capture detection 579(1992)99
- Torri, G., see Silvestro, L. 591(1992)225
- Toru, M., see Watanabe, A. 583(1992)241
- Tosa, T., see Nawata, M. 597(1992)415
- Totani, M., see Tabata, M. 597(1992)435
- Tóth, M. and Buser, H.-R.
Simple method for collecting volatile compounds from single insects and other point sources for gas chromatographic analysis 598(1992)303
- Touchstone, J. C., see Alvarez, J. G. 577(1992)142
- Tournade, H., see Jouve, J. 574(1992)9
- Toussi, A., Paquet, N., Huber, O., Frutiger, S., Tissot, J.-D., Hughes, G. J. and Hochstrasser, D. F.
Polypeptide marker and disease patterns found while mapping proteins in ascitis 582(1992)87
- Towns, J. K., Bao, J. and Regnier, F. E.
Synthesis and evaluation of epoxy polymer coatings for the analysis of proteins by capillary zone electrophoresis 599(1992)227
- Toyoda, H., Motoki, K., Tanikawa, M., Shinomiya, K., Akiyama, H. and Imanari, T.
Determination of human urinary hyaluronic acid, chondroitin sulphate and dermatan sulphate as their unsaturated disaccharides by high-performance liquid chromatography 565(1991)141
- Toyoda, H., see Akiyama, H. 579(1992)203
- Toyoda, H., see Imanari, T. 574(1992)142
- Toyo'oka, T., Ishibashi, M., Takeda, Y., Nakashima, K., Akiyama, S., Uzu, S. and Imai, K.
Precolumn fluorescence tagging reagent for carboxylic acids in high-performance liquid chromatography: 4-substituted-7-aminoalkylamino-2,1,3-benzoxadiazoles 588(1991)61
- Tozawa, T.
Electrophoretic analysis of enzyme-linked immunoglobulins and their clinical significance (Review) 569(1991)347
- Tozawa, T., see Moriyama, T. 571(1991)61
- Traag, W. A., see Van Rhijn, J. A. 595(1992)289
- Tracqui, A., see Kintz, P. 579(1992)172
- Traiffort, E., Hubert, P., Tayeb, N. and Aymard, N.
Rapid determination of 5-hydroxytryptamine in whole blood by liquid chromatography with fluorimetric detection 571(1991)231
- Traiger, G. J., see Davidson, V. P. 567(1991)213
- Trathnigg, B.
Determination of chemical composition of polymers by size-exclusion chromatography with coupled density and refractive index detection. III. Polyethylene oxide and polytetrahydrofuran 552(1991)507
- Travěncová, I., see Sitkey, V. 585(1991)373
- Treffel, P., see Makki, S. 563(1991)407
- Triggs, E. J., see Kastrissios, H. 577(1992)103
- Triolo, A., Bertini, J., Mannucci, C., Perico, A. and Pestellini, V.
Analysis of platelet-activating factor by gas chromatography-mass spectrometry: low-energy electron impact of the corresponding 3-acetyl-2-*tert*-butyldimethylsilyl derivative 568(1991)281
- Triolo, A., Mannucci, C. and Perico, A.
Gas chromatographic-mass spectrometric determination of two new antimycotic agents, 1-[(5-chloro-2-benzofuranyl)(2-chlorophenyl)methyl]-1*H*-imidazole and 1-[(5-bromo-2-benzofuranyl)phenylmethyl]-1*H*-imidazole, in rabbit plasma following topical administration: a preliminary comparison with bifonazole 567(1991)105
- Trivier, J. M., Pommery, J., Libersa, C., Caron, J. and Lhermitte, M.
High-performance liquid chromatographic assay for amiodarone N-deethylation in microsomes of rat liver 579(1992)269
- Trnavský, K., see Štovičková, J. 575(1992)229
- Trnka, T., see Jegorov, A. 558(1991)311
- Tröltzsch, C., Berger, R. and Michalik, M.
Liquid chromatographic isolation and structural elucidation of methoxymethylene dimethyl dodecatrienes catalytically synthesized from methanol and three moles of isoprene 594(1992)111
- Trotta, P. P., see Pramanik, B. N. 562(1991)377
- Trout, G. J., see Lisi, A. M. 563(1991)257
- Trout, G. J., see Lisi, A. M. 581(1992)57
- Trouvin, J. H., see Damaj, M. I. 563(1991)476
- Tsacopoulos, M., see Fathi, M. 563(1991)356
- Tsai, E. W., Ip, D. P. and Brooks, M. A.
Determination of alendronate in pharmaceutical dosage formulations by ion chromatography with conductivity detection 596(1992)217

- Tsai, R.-S., see El Tayar, N. 556(1991)181
- Tsai, T.-H. and Chen, C.-F.
High-performance liquid chromatographic determination of 18 α -glycyrrhetic acid and 18 β -glycyrrhetic acid in rat plasma: application to pharmacokinetic study 567(1991)405
- Tsai, T.-H. and Chen, C.-F.
Determination of glycyrrhizin in rabbit plasma by high-performance liquid chromatography with photodiode-array ultraviolet detection and its pharmacokinetics application 576(1992)170
- Tsai, T.-H. and Chen, C.-F.
Identification and determination of honokiol and magnolol from *Magnolia officinalis* by high-performance liquid chromatography with photodiode-array UV detection 598(1992)143
- Tsao, C. S., see Leung, P. Y. 576(1992)245
- Tsarbopoulos, A., see Pramanik, B. N. 562(1991)377
- Tsikakos, D., Fauler, J. and Frölich, J. C.
Enzymic preparation of dioxygen-18 labelled leukotriene E₄ and its use in quantitative gas chromatography-mass spectrometry 574(1992)181
- Tsoi, I. G., see Timerbaev, A. R. 555(1991)163
- Tsuboi, I., see Kitagawa, T. 567(1991)267
- Tsuboi, R., see Kimata, K. 558(1991)19
- Tsuboi, S., see Ohmori, S. 566(1991)1
- Tsuboi, S., see Ohmori, S. 574(1992)35
- Tsuchiya, M., see Yokoyama, Y. 566(1991)19
- Tsuchiya, M., see Yokoyama, Y. 583(1992)1
- Tsuda, T. and Zare, R. N.
Split injector for capillary zone electrophoresis 559(1991)103
- Tsuda, T., see Soini, H. 559(1991)547
- Tsuhako, M., see Baba, Y. 558(1991)273
- Tsuji, A., see Ikegawa, S. 577(1992)229
- Tsuji, K. and Little, R. J.
Charge-reversed, polymer-coated capillary column for the analysis of a recombinant chimeric glycoprotein 594(1992)317
- Tsujita, M., see Sakihama, N. 597(1992)147
- Tsuyama, K., see Ohmori, S. 566(1991)1
- Tsuyama, Y., Uchida, T. and Goto, T.
Analysis of underivatized C₁₂-C₁₈ fatty acids by reversed-phase ion-pair high-performance liquid chromatography with conductivity detection 596(1992)181
- Tu, T. M., see Varin, F. 574(1992)57
- Tuberoso, C., see Cabras, P. 585(1991)164
- Tucker, D. J., see Uhe, A. M. 564(1991)81
- Tuinstra, L. G. M. T., Povel, F. R. and Roos, A. H.
Multi-matrix-multi-pesticide method for agricultural products 552(1991)259
- Tuinstra, L. G. M. T., see Van Rhijn, J. A. 552(1991)517
- Tuinstra, L. G. M. T., see Van Rhijn, J. A. 592(1992)265
- Tuinstra, L. G. M. T., see Van Rhijn, J. A. 595(1992)289
- Tuomisto, J., see Raatikainen, O. 585(1991)247
- Tuong, A., see Maftouh, M. 562(1991)421
- Turcelová, E., see Madajová, V. 589(1992)329
- Türck, M., see Eisenbeiss, A. 589(1992)390
- Turcotte, J. G., see Amari, J. V. 590(1992)153
- Turgeon, J., see Abolfathi, Z. 579(1992)366
- Turk, J., Bohrer, A., Stump, W. T., Ramanadham, S. and Mangino, M. J.
Quantification of distinct molecular species of the 2-lyso metabolite of platelet-activating factor by gas chromatography-negative-ion chemical ionization mass spectrometry 575(1992)183
- Turková, J., Vohník, S., Helusová, Š., Beneš, M. J. and Tichá, M.
Galactosylation as a tool for the stabilization and immobilization of proteins (Review) 597(1992)19
- Tvrzická, E., Mareš, P., Písařiková, A., Novaković, J. and Hrabák, P.
Simplified gas chromatographic method for the simultaneous determination of phytosterols and cholesterol 563(1991)188
- Twelves, C. J., see Dobbs, N. A. 572(1991)211
- Twickler, M. S., see Buck, C. F. 594(1992)225
- Tyczkowska, K., see Hedeon, K. M. 572(1991)239
- Tyczkowska, K. L., Seay, S. S., Stoskopf, M. K. and Aucoin, D. P.
Determination of ceftazidime in dolphin serum by liquid chromatography with ultraviolet-visible detection and confirmation by thermospray liquid chromatography-mass spectrometry 576(1992)305
- Tyczkowska, K. L., Voyksner, R. D. and Aronson, A. L.
Solvent degradation of cloxacillin *in vitro*. Tentative identification of degradation products using thermospray liquid chromatography-mass spectrometry 594(1992)195
- Tyczkowska, K. L., see Voyksner, R. D. 567(1991)389
- Tyczkowski, J. K., see Burton, S. D. 571(1991)209
- Tzeng, T.-B. and Fung, H.-L.
Gas chromatographic analysis of isomeric organic mononitrates in plasma 571(1991)109

U

- Ubbink, J. B., Hayward Vermaak, W. J. and Bissbort, S.
Rapid high-performance liquid chromatographic assay for total homocysteine levels in human serum 565(1991)441
- Ubbink, J. B., Vermaak, W. J. H. and Bissbort, S. H.
High-performance liquid chromatographic assay of human lymphocyte kynureninase activity levels 566(1991)369
- Ubbink, J. B., see Legendijk, J. 576(1992)95
- Ubik, K., see Jursik, T. 586(1991)315

- Ubrich, N., Hubert, P., Regnault, V., Dellacherie, E. and Rivat, C.
Compared stability of Sepharose-based immunoabsorbents prepared by various activation methods 584(1992)17
- Ubuka, T., see Yao, K. 581(1992)11
- Uchida, K., see Isobe, T. 588(1991)115
- Uchida, T., see Tsuyama, Y. 596(1992)181
- Uchida, T., see Yasukawa, K. 597(1992)271
- Uchino, K., see Yamamura, Y. 567(1991)151
- Uda, T., Okuno, K., Suzuki, T. and Naruse, Y.
Gas chromatography for measurement of hydrogen isotopes at tritium processing 586(1991)131
- Udagawa, A., see Yoshioka, M. 566(1991)361
- Uden, P. C., see Sarafraz-Yazdi, A. 594(1992)395
- Udseth, H. R., see Smith, R. D. 559(1991)197
- Uebis, V.
Determination of a new blood glucose-lowering agent in biological fluids by capillary gas chromatography using electron-capture detection 567(1991)465
- Ueda, H., see Kawasaki, S. 595(1992)193
- Ueda, K., see Kamizono, A. 567(1991)113
- Ueda, M., see Honda, S. 588(1991)327
- Ueda, T., Mitchell, R., Kitamura, F., Metcalf, T., Kuwana, T. and Nakamoto, A.
Separation of naphthalene-2,3-dicarboxaldehyde-labeled amino acids by high-performance capillary electrophoresis with laser-induced fluorescence detection 593(1992)265
- Ueda, T., Mitchell, R., Kitamura, F. and Nakamoto, A.
Constant-potential amperometric detection of carbohydrates at metal electrodes in high-performance anion-exchange chromatography 592(1992)229
- Ueda, T., see Jinno, K. 594(1992)105
- Uehara, N., Iso, S. and Shijo, Y.
Retention behavior of salicylidene-glycinate-aluminum(III) in reversed-phase high-performance liquid chromatography 586(1991)356
- Ueno, H. and Matsuo, S.
High-performance liquid chromatography followed by radioimmunoassay for the determination of a luteinizing hormone-releasing hormone analogue, leuprorelin, and its metabolite 566(1991)57
- Ueno, K. and Umeda, T.
Electrochemical and chromatographic properties of selected hydrazine and hydrazide derivatives of carbonyl compounds 585(1991)225
- Ueno, M., Arai, T., Kojima, K., Ogawa, H. K., Matsumoto, I. and Seno, N.
Improved affinity chromatographic purification of D-mannose-N-acetyl-D-glucosamine-specific lectin from the bark of *Sophora japonica* eliminating the loss by sugar specific self-aggregation 597(1992)197
- Ueno, R., Uno, K. and Aoki, T.
Determination of oxytetracycline in blood serum by high-performance liquid chromatography with direct injection 573(1992)333
- Uesugi, T., see Sano, K. 578(1992)63
- Ueta, N., see Werner, A. 553(1991)205
- Uges, D. R. A., see Bloemhof, H. 572(1991)171
- Uggeri, F., see Anelli, P. L. 589(1992)346
- Ugwu, S. O. and Blanchard, J.
High-performance liquid chromatographic assay for the α -melanotropin[4,10] fragment analogue (Melanotan-II) in rat plasma 584(1992)175
- Uhe, A. M., Collier, G. R., McLennan, E. A., Tucker, D. J. and O'Dea, K.
Quantitation of tryptophan and other plasma amino acids by automated pre-column o-phthalaldehyde derivatization high-performance liquid chromatography: improved sample preparation 564(1991)81
- Uhlig, R., see Grune, T. 553(1991)193
- Ujeno, Y., see Kitaoka, Y. 585(1991)369
- Ulfelder, K., see Schwartz, H. E. 559(1991)267
- Umeda, I., see Nishioka, R. 565(1991)237
- Umeda, T., see Ueno, K. 585(1991)225
- Underberg, W. J. M., see Kaijser, G. P. 571(1991)121
- Underberg, W. J. M., see Kaijser, G. P. 583(1992)175
- Underhill, K., see Mofidi, M. 589(1992)241
- Unger, K. K., Lork, K. D., Pfeleiderer, B., Albert, K. and Bayer, E.
Impact of acidic/hydrothermal treatment on pore structural and chromatographic properties of porous silicas. I. The conventional approach 556(1991)395
- Unger, K. K., see Hanson, M. 599(1992)65
- Unger, K. K., see Hanson, M. 599(1992)77
- Unni, L. K. and Becker, R. E.
Determination of heptylphysostigmine in plasma by high-performance liquid chromatography with electrochemical detection 573(1992)275
- Unni, L. K., Hannant, M. E. and Becker, R. E.
High-performance liquid chromatographic method using ultraviolet detection for measuring metrifonate and dichlorvos levels in human plasma 573(1992)99
- Uno, B., see Tomita, M. 566(1991)239
- Uno, B., see Tomita, M. 571(1991)324
- Uno, K., see Nishioka, R. 565(1991)237
- Uno, K., see Ueno, R. 573(1992)333
- Unteregger, G.
Utility of protein electrophoretic analysis in the characterization of malignant tissues (Review) 569(1991)367
- Uralets, V. P., see Rodchenkov, G. M. 565(1991)45
- Uray, G., see Lindner, W. 553(1991)373
- Urbach, H. P., see De Bokx, P. K. 594(1992)9
- Urban, G., see Ferris, R. 577(1992)251
- Urbánková, E., Vacek, A., Nováková, N., Matulík, F. and Chmelík, J.
Investigation of red blood cell fractionation by gravitational field-flow fractionation 583(1992)27
- Urbányi, Z. and Medzihradzky, D.
Rapid method to isolate serum amyloid P component from human plasma. Characterization of the isolated protein 578(1992)130
- Urbonavičienė, J., see Maruška, A. 596(1992)157
- Urge, L., see Otvos, Jr., L. 599(1992)43
- Usansky, J. I. and Damani, L. A.
Assay of metyrapone, metyrapol and the isomeric mono-N-oxides of metyrapone in biological fluids by high-performance liquid chromatography 563(1991)283

- Usui, T., Watanabe, T. and Higuchi, S.
Determination of a new bisphosphonate, YM175, in plasma, urine and bone by high-performance liquid chromatography with electrochemical detection 584(1992)213
- Uzu, S., see Toyo'oka, T. 588(1991)61
- V**
- Vacek, A., see Urbánková, E. 583(1992)27
- Vacík, J., see Poláček, M. 596(1992)265
- Vainio, H., see Rosenberg, C. 552(1991)265
- Valcárcel, M., see Linares, P. 558(1991)147
- Valcárcel, M., see Linares, P. 585(1991)267
- Valcárcel, M., see Luque de Castro, M. D. 600(1992)183
- Valdez, J. G., see Gurley, L. R. 559(1991)431
- Valentin, P.
Is chromatography a separation process? The zonoid answer (Review) 556(1991)25
- Valentini, C., see Lo Coco, F. 590(1992)235
- Valkó, K. and Slégel, P.
Chromatographic separation and molecular modelling of triazines with respect to their inhibition of the growth of L1210/R71 cells 592(1992)59
- Valkó, K., see Bánhegyi, G. 553(1991)407
- Valkó, K., see Forgács, E. 592(1992)75
- Vallar, L., see Regnault, V. 576(1992)87
- Vallar, L., see Regnault, V. 584(1992)93
- Vallat, P., see El Tayar, N. 556(1991)181
- Vallon, J. J., see Duthel, J. M. 570(1991)166
- Vallon, J. J., see Duthel, J. M. 579(1992)85
- Valls, R., see Piovetti, L. 588(1991)99
- VanAlstyne, E. L. and Spaethe, S. M.
Rapid method for automated on-line extraction and fractionation of plasma leukotrienes and 12-hydroxy-5,8,10,14-eicosatetraenoic acids by reversed-phase high-performance liquid chromatography 579(1992)37
- Van Asten, A. C., see Bruin, G. J. M. 559(1991)163
- Van Baak, M. J., see Arts, C. J. M. 564(1991)429
- Van Bakergem, E., Van der Hoeven, R. A. M., Niessen, W. M. A., Tjaden, U. R., Van der Greef, J., Poon, G. K. and McCague, R.
On-line continuous-flow dialysis thermospray tandem mass spectrometry for quantitative screening of drugs in plasma: roglitimide 598(1992)189
- Van Belle, S. J.-P., De Smet, M., De Neve, W., Monsaert, C., Storme, G. A. and Massart, D. L.
Determination of vinca alkaloids in mouse tissues by high-performance liquid chromatography 578(1992)223
- Van Belle, S. J.-P., De Smet, M., Monsaert, C., Geerts, F., Storme, G. A. and Massart, D. L.
High-performance liquid chromatographic determination of navelbine in MO₄ mouse fibrosarcoma cells and biological fluids 576(1992)351
- Van Bergen, C. A., Collier, P. D., Cromie, D. D. O., Lucas, R. A., Preston, H. D. and Sissons, D. J.
Determination of chloropropanols in protein hydrolysates 589(1992)109
- Van Bommel, C. H. M., see De Bokx, P. K. 598(1992)115
- Van Boom, J. H., see De Ruiter, G. A. 584(1992)69
- Van Boven, M., see Debrabandere, L. 564(1991)557
- Van Breugel, D. C., see Leenheers, L. H. 578(1992)189
- Van Bruchem, G. D., see Ploum, M. E. 564(1991)413
- Van Buul-Offers, S. C., see Bayne, S. 562(1991)391
- Vance, N. C., see Harvey, S. D. 587(1991)300
- Van Crugten, J. T., see Milne, R. W. 565(1991)457
- Van de Heijning, B. J. M., Koekkoek-van den Herik, I., Iványi, T. and Van Wimersma Greidanus, T. B.
Solid-phase extraction of plasma vasopressin: evaluation, validation and application 565(1991)159
- Van de Merbel, N. C., see Farjam, A. 589(1992)141
- Van den Berg, C. W., see Bootsma, H. J. 591(1992)187
- Van den Berg, G. A., see Hessels, J. 563(1991)1
- Van den Berg, J. H. M., see Goosens, E. C. 552(1991)489
- Van den Berg, R. H., see Jansen, E. H. J. M. 553(1991)123
- Van den Berg, R. H., see Jansen, E. H. J. M. 566(1991)461
- Van den Berg, R. H., see Jansen, E. H. J. M. 580(1992)111
- Van den Berg, R. H., see Van Look, L. J. 564(1991)451
- Van den Biggelaar-Martea, M., see Vree, T. B. 573(1992)168
- Van den Biggelaar-Martea, M., see Vree, T. B. 578(1992)239
- Vandenbosch, C., Vannecke, C. and Massart, D. L.
Optimization of the separation of (6R)- and (6S)-leucovorin and evaluation of the robustness of the optimum 592(1992)37
- Van den Bossche, W., see Coene, J. 553(1991)285
- Van den Brande, J. L., see Bayne, S. 562(1991)391
- Vandenbroeck, M., Van Vyncht, G., Gaspar, P., Dasnois, C., Delahaut, P., Pelzer, G., De Graeve, J. and Maghuin-Rogister, G.
Identification and characterization of 19-nortestosterone in urine of meat-producing animals 564(1991)405
- Vanden Driessche, N., see Hamoir, T. 589(1992)31
- Van den Eeckhout, E., see Coene, J. 553(1991)285
- Van de Nesse, R. J., Hoogland, G. J. M., De Moel, J. J. M., Gooijer, C., Brinkman, U. A. T. and Velthorst, N. H.
On-line coupling of liquid chromatography to thin-layer chromatography for the identification of polycyclic aromatic hydrocarbons in marine sediment by fluorescence excitation and emission spectroscopy 552(1991)613
- Van de Nesse, R. J., see Somsen, G. W. 552(1991)635
- Van den Heuvel, H., see Qimin, L. 562(1991)435
- Van der Greef, J., see Heeremans, C. E. M. 554(1991)205
- Van der Greef, J., see Kokkonen, P. S. 565(1991)265
- Van der Greef, J., see Niessen, W. M. A. 554(1991)3
- Van der Greef, J., see Poon, G. K. 576(1992)235
- Van der Greef, J., see Reeuwijk, H. J. E. M. 575(1992)269
- Van der Greef, J., see Reinhoud, N. J. 574(1992)327

- Van der Greef, J., see Stegehuis, D. S. 591(1992)341
 Van der Greef, J., see Tinke, A. P. 554(1991)119
 Van der Greef, J., see Van Bakergem, E. 598(1992)189
 Van der Greef, J., see Venizelos, V. 573(1992)259
 Van der Greef, J., see Verheij, E. R. 554(1991)339
 Van der Hoeven, R. A. M., see Heeremans, C. E. M. 554(1991)205
 Van der Hoeven, R. A. M., see Poon, G. K. 576(1992)235
 Van der Hoeven, R. A. M., see Tinke, A. P. 554(1991)119
 Van der Hoeven, R. A. M., see Van Bakergem, E. 598(1992)189
 Van der Hoorn, F. A. J., see Boomsma, F. 574(1992)109
 Van Der Hoorn, F. A. J., Boomsma, F., Man in 't Veld, A. J. and Schalekamp, M. A. D. H.
 Improved measurement of urinary catecholamines by liquid-liquid extraction, derivatization and high-performance liquid chromatography with fluorimetric detection 563(1991)348
 Van der Horst, A., De Goede, P. N. F. C., Van Diemen, H. A. M., Polman, C. H. and Martens, H. J. M.
 Determination of 4-aminopyridine in serum by solid-phase extraction and high-performance liquid chromatography 574(1992)166
 Van der Horst, F. A. L., Van Opstal, M. A. J., Teeuwssen, J., Post, M. H., Holthuis, J. J. M. and Brinkman, U. A. T.
 Comparative study on the determination of the anti-neoplastic drug teniposide in plasma using micellar liquid chromatography and surfactant-mediated plasma clean-up 567(1991)161
 Van der Mast, C. A., Hekstra, D. and Voorma, H. O.
 Separation of translationally active mRNAs by reversed-phase ion-pair high-performance liquid chromatography 564(1991)115
 Van Der Meersch-Mougeot, V. and Diquet, B.
 Sensitive one-step extraction procedure for column liquid chromatographic determination of fluvoxamine in human and rat plasma 567(1991)441
 Van Der Meersch-Mougeot, V., see Thomare, P. 583(1992)217
 Van der Stegen, G. H. D., see Plantinga, W. J. 555(1991)311
 Van der Vijgh, W. J. F., see De Jong, J. 570(1991)209
 Van der Vijgh, W. J. F., see De Jong, J. 574(1992)273
 Van der Vusse, G. J., see Roemen, T. H. M. 570(1991)243
 Van der Vusse, G. J., see Van Nieuwenhoven, F. A. 570(1991)173
 Van der Wal, S., see Coenen, A. J. J. M. 593(1992)243
 Van der Wal, S., see Coenen, A. J. J. M. 596(1992)59
 Van der Wal, S., see Mengerink, Y. 552(1991)593
 Van der Weyde, M. P., see Wright, M. R. 565(1991)225
 Van der Woude, H. R., see Van Tellingen, O. 553(1991)47
 Van der Woude, H. R., see Van Tellingen, O. 574(1992)293
 Van de Water, C. and Haagsma, N.
 Analysis of chloramphenicol residues in swine tissues and milk: comparative study using different screening and quantitative methods 566(1991)173
 Van de Water, C., see Moretti, V. M. 583(1992)77
 Van de Weijer, P., see De Bokx, P. K. 598(1992)115
 Van Diemen, H. A. M., see Van der Horst, A. 574(1992)166
 Van Dijk, H., see Bootsma, H. J. 591(1992)187
 Van Dijk, K. N., see Bloemhof, H. 572(1991)171
 Van Ede, J., see Welling, G. W. 599(1992)157
 Van Ewijk-Beneken Kolmer, E. W. J., see Vree, T. B. 579(1992)131
 Van Faassen, A., see De Kok, T. M. C. M. 580(1992)135
 Van Gijn, R., see Burger, D. M. 584(1992)239
 Van Ginkel, L. A.
 Review. Immunoaffinity chromatography, its applicability and limitations in multi-residue analysis of anabolizing and doping agents 564(1991)363
 Van Ginkel, L. A., see Jansen, E. H. J. M. 580(1992)111
 Van Grunsven, W. M. J., see Hessing, M. 599(1992)267
 Vanhaelen, M., see Diallo, B. 558(1991)446
 Vanhaelen-Fastré, R., see Diallo, B. 558(1991)446
 Van Horne, K. C., see Reinhart, P. 570(1991)425
 Van Iersel, A. A. J., see Reinerink, E. J. M. 553(1991)233
 Van Ingen, H. E., see Takkenberg, B. 565(1991)430
 Van Jaarsveld, P. P., see Le Roux, A. M. 581(1992)306
 Vankeerberghen, P., see Bourguignon, B. 592(1992)51
 Van Kempen, T. A. T. G. and Odle, J.
 Quantification of carnitine esters by high-performance liquid chromatography. Effect of feeding medium-chain triglycerides on the plasma carnitine ester profile 584(1992)157
 Van Leeuwen, F. X. R., see Doorn, L. 553(1991)135
 Van Leeuwen, F. X. R., see Jansen, E. H. J. M. 566(1991)471
 Van Leuken, R. G. J., see Duchateau, A. L. L. 552(1991)605
 Van Lier, J. E., see Berger, M. 593(1992)133
 Vanlinthout, L. E. H., see Lagerwerf, A. J. 570(1991)390
 Van Look, L. J., Jansen, E. H. J. M., Van den Berg, R. H., Zomer, G., Vanoosthuize, K. E. and Van Peteghem, C.
 Development of a competitive enzyme immunoassay for 17 α -19-nortestosterone 564(1991)451
 Van Miert, A. S. J. P. A. M., see Van 't Klooster, G. A. E. 579(1992)158
 Van Miert, A. S. J. P. A. M., see Van 't Klooster, G. A. E. 579(1992)355
 Vannoeke, C., see Vandenbosch, C. 592(1992)37
 Van Nieuwenhoven, F. A., Vork, M. M., Surtel, D. A. M., Kleine, A. H., Van der Vusse, G. J. and Glatz, J. F. C.
 High-yield two-step chromatographic procedure for purification of fatty acid-binding protein from human heart 570(1991)173
 Vannoni, D., see Terzuoli, L. 586(1991)153
 Van Oosterom, A. T., see Pattyn, G. 564(1991)352
 Vanoosthuize, K. E., see Van Look, L. J. 564(1991)451
 Van Opstal, M. A. J., see Van der Horst, F. A. L. 567(1991)161
 Van Peteghem, C.
 Preface 564(1991)361
 Van Peteghem, C., see Daeseleire, E. 562(1991)673
 Van Peteghem, C., see Daeseleire, E. 564(1991)445
 Van Peteghem, C., see Daeseleire, E. 564(1991)469
 Van Peteghem, C., see Debruyckere, G. 564(1991)393
 Van Peteghem, C., see Van Look, L. J. 564(1991)451
 Van Polanen, A., see De Jong, J. 553(1991)243
 Van Regenmortel, M. H. V., see Dubs, M.-C. 597(1992)391

- Van Rhijn, J. A., Traag, W. A., Kulik, W. and Tuinstra, L. G. M. T.
Automated clean-up procedure for the gas chromatographic-high-resolution mass spectrometric determination of polychlorinated dibenzo-*p*-dioxins and dibenzofurans in milk 595(1992)289
- Van Rhijn, J. A. and Tuinstra, L. G. M. T.
Miniaturisation of size-exclusion chromatography as a powerful clean-up tool in residue analysis 552(1991)517
- Van Rhijn, J. A., Viveen, J. and Tuinstra, L. G. M. T.
Automated determination of aflatoxin B₁ in cattle feed by two-column solid-phase extraction with on-line high-performance liquid chromatography 592(1992)265
- Van Rompaey, F., see Woestenborghs, R. 583(1992)223
- Van Schaik, F., see Van Vliet, T. 553(1991)179
- Van Schijndel, H. B., see Hessing, M. 599(1992)267
- Van Schoonhoven, J., see Van Vliet, T. 553(1991)179
- Van Seeventer, P. B., see Van 't Klooster, G. A. E. 571(1991)157
- Van Tellingen, O., Beijnen, J. H., Baurain, R., Ten Bokkel Huinink, W. W., Van der Woude, H. R. and Nooyen, W. J.
High-performance liquid chromatographic determination of vinblastine, 4-O-deacetylvinblastine and the potential metabolite 4-O-deacetylvinblastine-3-oic acid in biological fluids 553(1991)47
- Van Tellingen, O., Beijnen, J. H., Baurain, R., Ten Bokkel Huinink, W. W., Van der Woude, H. R. and Nooyen, W. J.
Biodetermination of N-(deacetyl-O-4-vinblastoyl-23)-L-tryptophan, a metabolite of vintriptol, by high-performance liquid chromatography with fluorescence detection 574(1992)293
- Van Tellingen, O., Kuijpers, A., Beijnen, J. H., Baselier, M. R. P., Burghouts, J. T. M. and Nooyen, W. J.
Bio-analysis of vinorelbine by high-performance liquid chromatography with fluorescence detection 573(1992)328
- Van Tellingen, O., Sips, J. H. M., Beijnen, J. H., Schornagel, J. H. and Nooyen, W. J.
Bioanalysis of the investigational anti-tumour drug 5,10-dideaza-5,6,7,8-tetrahydrofolic acid by high-performance liquid chromatography with ultraviolet detection 576(1992)158
- Van Tellingen, O., see Burger, D. M. 584(1992)239
- Van 't Klooster, G. A. E., Kolker, H. J., Woutersen-van Nijnanten, F. M. A., Noordhoek, J. and Van Miert, A. S. J. P. A. M.
Determination of trimethoprim and its oxidative metabolites in cell culture media and microsomal incubation mixtures by high-performance liquid chromatography 579(1992)355
- Van 't Klooster, G. A. E., Van Seeventer, P. B., Kolker, H. J., Smit, L. A. and Witkamp, R. F.
High-performance liquid chromatographic method for the routine determination of sulphadimidine, its hydroxy metabolites and N4-acetylsulphadimidine in body fluids and cell culture media 571(1991)157
- Van 't Klooster, G. A. E., Woutersen-van Nijnanten, F. M. A., Kolker, H. J., Noordhoek, J. and Van Miert, A. S. J. P. A. M.
Improved high-performance liquid chromatographic method for the determination of ethylmorphine and its metabolites in microsomal incubations and cell culture media 579(1992)158
- Van Toorn, W. G., see Plantinga, W. J. 555(1991)311
- Van Vliet, T., Van Schaik, F., Van Schoonhoven, J. and Schrijver, J.
Determination of several retinoids, carotenoids and E vitamers by high-performance liquid chromatography. Application to plasma and tissues of rats fed a diet rich in either β -carotene or canthaxanthin 553(1991)179
- Van Vyncht, G., see Vandebroeck, M. 564(1991)405
- Van Wimersma Greidanus, T. B., see Van de Heijning, B. J. M. 565(1991)159
- Van Zijtveld, J., Pouwelse, A. V. and Groen, C. P.
Application of an internal surface reversed-phase column for the automated determination of flucycloxon residues 600(1992)211
- Van Zoonen, P., see Hogendoorn, E. A. 552(1991)113
- Varga, L. L., see Granneman, G. R. 568(1991)197
- Varin, F., Tu, T. M., Benoît, F., Villeneuve, J.-P. and Théorêt, Y.
High-performance liquid chromatographic determination of spironolactone and its metabolites in human biological fluids after solid-phase extraction 574(1992)57
- Varin, F., see Ducharme, J. 573(1992)79
- Varin, F., see Théorêt, Y. 575(1992)162
- Varney, M. S., see Bianchi, A. P. 557(1991)429
- Varro, R., see Harrington, S. J. 559(1991)385
- Vas, A., see Cholerton, S. 575(1992)325
- Vassarotto, C., see Tarditi, L. 599(1992)13
- Vassilakos, C., see Katsanos, N. A. 557(1991)469
- Vázquez Odériz, L., see Paseiro Losada, P. 585(1991)75
- Vedenin, A. N., see Rodchenkov, G. M. 565(1991)45
- Veel, T., see Holthe, M. R. 579(1992)376
- Vega, A. and Coca, J.
Activity coefficients at infinite dilution determined by gas-liquid chromatography. Organic solvents in Apiezon L 586(1991)303
- Velander, W. H., see Orthner, C. L. 558(1991)55
- Velayudhan, A., see Yang, Y. 598(1992)169
- Velthorst, N. H., see Schreurs, M. 552(1991)625
- Velthorst, N. H., see Somsen, G. W. 552(1991)635
- Velthorst, N. H., see Van de Nesse, R. J. 552(1991)613
- Vendrig, D. E. M. M., see Bloemhof, H. 572(1991)171
- Venizelos, V., Irth, H., Tjaden, U. R., Van der Greef, J., Breimer, D. D., Mulders, T. M. T. and Mulder, G. J.
Enantioselective determination of *R*- and *S*-(α -bromoisovaleryl)urea in plasma using high-performance liquid chromatography after solid-phase extraction 573(1992)259
- Venkataramana, D. and Krishna, D. R.
High-performance liquid chromatographic determination of usnic acid in plasma 575(1992)167
- Ventura, F., see Guardiola, A. 562(1991)481
- Ventura, K., see Churáček, J. 557(1991)523

- Ventura, R., Fraisse, D., Becchi, M., Paisse, O. and Segura, J.
Approach to the analysis of diuretics and masking agents
by high-performance liquid chromatography-mass
spectrometry in doping control 562(1991)723
- Venturella, V. S., see Kvalo, L. T. 565(1991)391
- Venturella, V. S., see Wilhelm, J. A. 578(1992)231
- Venturelli, E., Manzari, A., Cavalleri, A., Benzo, M., Secretò,
G. and Marubini, E.
Urinary testosterone measurement by gas
chromatography after solid-phase extraction and high-
performance liquid chromatography 582(1992)7
- Verbeeck, R. K., see Hoffman, D. A. 565(1991)447
- Vercammen, J., see Vermoesen, A. 564(1991)385
- Vercruyse, A., see Mertens, K. 565(1991)149
- Verdievel, H., see De Vos, M. 564(1991)296
- Vereczkey, L., see Pudleiner, P. 553(1991)29
- Vereczkey, L., see Tomori, É. 578(1992)91
- Verga, F., see Belotto, N. 572(1991)327
- Vergin, H., see Mascher, H. 583(1992)122
- Vergniol, J. C., Bruno, R., Montay, G. and Frydman, A.
Determination of Taxotere in human plasma by a semi-
automated high-performance liquid chromatographic
method 582(1992)273
- Vergori, F., see Di Muccio, A. 552(1991)241
- Vergori, F., see Di Muccio, A. 553(1991)305
- Vergori, L., see Di Muccio, A. 553(1991)305
- Verhe, R., see Courtheyn, D. 564(1991)537
- Verheij, E. R., Tjaden, U. R., Niessen, W. M. A. and Van der
Greef, J.
Pseudo-electrochromatography-mass spectrometry: a new
alternative 554(1991)339
- Vérité, P., André, D., Ménager, S. and Lafont, O.
Gas chromatographic-mass spectrometric method for
analysis of a pyrimido-*s*-triazine and some of its
metabolites in dog urine 578(1992)134
- Vermaak, W. J. H., see Lagendijk, J. 576(1992)95
- Vermaak, W. J. H., see Ubbink, J. B. 566(1991)369
- Vermeulen, A. M., Rosseel, M. T. and Belpaire, F. M.
High-performance liquid chromatographic method for the
simultaneous determination of *R*-(-)- and *S*-(+)-
hexobarbital in rat plasma 567(1991)472
- Vermeulen, A. M., see Rosseel, M. T. 568(1991)239
- Vermeulen, E., see Daumas, L. 570(1991)89
- Vermoesen, A., Vercammen, J., Sanders, C., Courtheyn, D.
and De Brabander, H. F.
Comparison of two mass-selective detectors with special
reference to the analysis of residues of anabolics
564(1991)385
- Verner, P., see Matucha, M. 588(1991)251
- Veronese, F. M., see Švagrová, I. 563(1991)95
- Verpoorte, E. M. J., see Manz, A. 593(1992)253
- Verresen, K., see Naidong, W. 586(1991)61
- Verresen, K., see Naidong, W. 586(1991)67
- Verwey-van Wissen, C. P. W. G. M., Koopman-Kimenai, P.
M. and Vree, T. B.
Direct determination of codeine, norcodeine, morphine
and normorphine with their corresponding *O*-glucuronide
conjugates by high-performance liquid chromatography
with electrochemical detection 570(1991)309
- Verwey-van Wissen, C. P. W. G. M., see Vree, T. B.
578(1992)239
- Vestal, M. L., see Getek, T. A. 554(1991)191
- Vettoretti, A., see Carpenè, G. 553(1991)201
- Veuthey, J.-L., see Coquet, A. 553(1991)255
- Vey, A., see Cherton, J.-C. 566(1991)511
- Vezzani, S., see Castello, G. 585(1991)273
- Viala, A., see Rop, P. P. 573(1992)87
- Viano, I., see Silvestro, L. 563(1991)443
- Viano, I., see Silvestro, L. 591(1992)225
- Viaud, M. C., see Latxague, L. 586(1991)166
- Vicat, P., Cousty-Berlin, D. and Hunt, P.
Determination of the enantiomers of a novel 20,21-
dinoreburnamenine derivative in rat plasma and brain by
high-performance liquid chromatography using a chiral
stationary phase 575(1992)315
- Vicente, C., Mateos, J. L., Pedrosa, M. M. and Legaz, M. E.
High-performance liquid chromatographic determination
of sugars and polyols in extracts of lichens and sugarcane
juice 553(1991)271
- Vidal, P., Nielsen, E. and Welinder, B. S.
Effect of glycation on the heterogeneity of human serum
albumin analysed by reversed-phase high-performance
liquid chromatography in a solvent containing formic
acid 573(1992)201
- Vidal-Madjar, C., Jaulmes, A., Sébille, B., González, M. J.,
Jiménez, B. and Hernández, L. M.
Study of protein-ligand binding effects by direct
chromatographic on-column injection 584(1992)11
- Vidal-Madjar, C., see Lemque, R. 553(1991)165
- Vidal-Madjar, C., see Lemque, R. 599(1992)255
- Vidal-Madjar, C., see Vivarat-Perrin, M.-P. 584(1992)3
- Videmann, B., see Benoit, E. 583(1992)167
- Viden, I., see Řezanka, T. 558(1991)215
- Vidensek, M. A., see Pesek, J. J. 556(1991)373
- Vigano, S., see Fermo, I. 593(1992)171
- Vigevani, A., see Del Nero, S. 593(1991)25
- Vijayalakshmi, M. A., see El-Kak, A. 570(1991)29
- Vijayalakshmi, M. A., see Goubbran-Botros, H. 597(1992)357
- Vijayalakshmi, M. A., see Kirchnerberger, J. 557(1991)325
- Vijayalakshmi, M. A., see Wu, X. 584(1992)35
- Viktorova, E. N., see Berezkin, V. G. 552(1991)313
- Villa, P., see Brega, A. 553(1991)249
- Villar, J., see Ruiz-Gutierrez, V. 581(1992)171
- Villeneuve, F., see Desbène, P. L. 553(1991)211
- Villeneuve, J.-P., see Varin, F. 574(1992)57
- Villermet, A., Thiébaud, D., Caude, M. and Rosset, R.
Packed column supercritical fluid chromatography with
carbon dioxide-polar modifiers. Influence of carbon
dioxide density on retention 557(1991)85
- Vincze, I., see Iohan, F. 564(1991)27
- Vingler, P., Kermici, M. and Krien, P.
Direct quantitative digital autoradiography-thin-layer
chromatography of 3 α ,3 β - and 5 α -reduced and 17 β -
dehydrogenated androgens derived from testosterone
metabolism 571(1991)73
- Vinogradova, N. I., see Chalykh, A. E. 592(1992)121
- Vinš, I., see Břízová, K. 593(1992)125
- Vinš, I., see Grüner, B. 588(1991)201

- Vinš, I., see Grüner, B. 595(1992)169
- Vinther, A. and Sørensen, H.
Mathematical model describing dispersion in free solution capillary electrophoresis under stacking conditions 559(1991)3
- Vinther, A. and Sørensen, H.
Temperature elevations of the sample zone in free solution capillary electrophoresis under stacking conditions 559(1991)27
- Vinther, A. and Sørensen, H.
Radial pH distribution during capillary electrophoresis with electroosmotic flow. Analysis with high ionic strength buffers 589(1992)315
- Vioque, E., see Alaiz, M. 591(1992)181
- Viscomi, G. C., see Scapol, L. 600(1992)235
- Visconti, M., see Lombardi, F. 576(1992)129
- Visser, J., see Molema, G. 579(1992)107
- Visser, R., see Haanappel, V. A. C. 562(1991)531
- Visser, S., Slangen, C. J. and Robben, A. J. P. M.
Determination of molecular mass distributions of whey protein hydrolysates by high-performance size-exclusion chromatography 599(1992)205
- Visser, T., see Somsen, G. W. 552(1991)635
- Vitenberg, A. G.
Methods of equilibrium concentration for the gas chromatographic determination of trace volatiles (Review) 556(1991)1
- Vitenberg, A. G., see Marinichev, A. N. 600(1992)251
- Vivarat-Perrin, M.-P., Sébille, B. and Vidal-Madjar, C.
Study of affinity supports based on reactive polymers immobilized on silica: affinity constant determination from isocratic zonal elution 584(1992)3
- Viveen, J., see Van Rhijn, J. A. 592(1992)265
- Vlasenko, E. V., see Gavrilova, T. B. 552(1991)179
- Vlegaar, R., see Shephard, G. S. 566(1991)195
- Vlietinck, A. J., see Qimin, L. 562(1991)435
- Voelkel, A. and Janas, J.
Attempts to use Laffort's solubility factors as polarity parameters for organic compounds in inverse gas chromatography 555(1991)205
- Vohnik, S., see Turková, J. 597(1992)19
- Voinov, V. G., El'kin, Y. N., Kuznetsova, T. A., Mal'tsev, I. I., Mikhailov, V. V. and Sasunkevich, V. A.
Use of mass spectrometry for the detection and identification of bromine-containing diphenyl ethers 586(1991)360
- Voirin, A., Letavernier, J.-F. and Sebille, B.
Improvement of extraction and concentration of milk peptides with solid-phase cartridges for analysis by high-performance liquid chromatography 553(1991)155
- Voirin, P., see Julien-Larose, C. 562(1991)39
- Voirin, S. G., Baumes, R. L., Gunata, Z. Y., Bitteur, S. M., Bayonove, C. L. and Tapiero, C.
Analytical methods for monoterpene glycosides in grape and wine. I. XAD-2 extraction and gas chromatographic-mass spectrometric determination of synthetic glycosides 590(1992)313
- Voirin, S. G., Baumes, R. L., Sapis, J.-C. and Bayonove, C. L.
Analytical methods for monoterpene glycosides in grape and wine. II. Qualitative and quantitative determination of monoterpene glycosides in grape 595(1992)269
- Volin, P.
Determination of urinary normetanephrine, metanephrine and 3-methoxytyramine by high-performance liquid chromatography with electrochemical detection: comparison between automated column-switching and manual dual-column sample purification methods 578(1992)165
- Volin, P.
Simultaneous determination of serum cortisol and cortisone by reversed-phase liquid chromatography with ultraviolet detection 584(1992)147
- Vonderschmitt, D. J., see Feng, N. 575(1992)289
- Vonderschmitt, D. J., see Wiederkehr, F. 566(1991)77
- Voogt, W. H., see Özçimder, M. 570(1991)19
- Voorma, H. O., see Van der Mast, C. A. 564(1991)115
- Vork, M. M., see Van Nieuwenhoven, F. A. 570(1991)173
- Voskoboinikova, I. V., see Geodakyan, S. V. 577(1992)371
- Vosmanská, M., see Linhart, I. 578(1992)215
- Voyksner, R. D., Tyczkowska, K. L. and Aronson, A. L.
Development of analytical methods for some penicillins in bovine milk by ion-paired chromatography and confirmation by thermospray mass spectrometry 567(1991)389
- Voyksner, R. D., see Tyczkowska, K. L. 594(1992)195
- Vrbnac, J. J., see Komatsu, S. 568(1991)460
- Vrbaški, L. and Lepojević, Ž.
Rapid identification of microbial starch degradation products from a complex nutrient medium by a thin-layer chromatographic method 558(1991)328
- Vree, T. B., Van den Biggelaar-Martea, M., Peeters, A. and Imbimbo, B. P.
High-performance liquid chromatography and preliminary pharmacokinetics of rifloxacin and its metabolites, N-desmethylrifloxacin and rifloxacin sulfoxide, in urine of Rhesus monkey *Macaca mulatta* 573(1992)168
- Vree, T. B., Van den Biggelaar-Martea, M. and Verwey-van Wissen, C. P. W. G. M.
Determination of naproxen and its metabolite O-desmethylnaproxen with their acyl glucuronides in human plasma and urine by means of direct gradient high-performance liquid chromatography 578(1992)239
- Vree, T. B., Van Ewijk-Beneken Kolmer, E. W. J. and Nouws, J. F. M.
Direct-gradient high-performance liquid chromatographic analysis and preliminary pharmacokinetics of flumequine and flumequine acyl glucuronide in humans: effect of probenecid 579(1992)131
- Vree, T. B., see Lagerwerf, A. J. 570(1991)390
- Vree, T. B., see Verwey-van Wissen, C. P. W. G. M. 570(1991)309
- Vreeken, R. J., see Barceló, D. 553(1991)311
- Vučković, G., Juranić, N., Miljević, D. and Čelap, M. B.
Simultaneous effect of geometrical isomerism and chelate ring size of tris(aminocarboxylato)cobalt(III) complexes on their behaviour in thin-layer chromatography on silica gel and alumina 585(1991)181
- Vučković, G. N., see Janjić, T. J. 596(1992)91

Vuik, J.

Rapid determination of abamectin in lettuce and cucumber by high-performance liquid chromatography 553(1991)299

Vuorela, H., see Härmälä, P. 593(1992)329

Vuorela, H., see Salo, M. 592(1992)127

Vymlátíl, J., see Böswart, J. 571(1991)19

W

- Wachter, H., see Werner, E. R. 570(1991)43
- Wada, H., see Fukuda, H. 567(1991)459
- Wada, H., see Haginaka, J. 592(1992)301
- Wada, H., see Yasui, T. 596(1992)73
- Wada, Y., see Ohba, S. 568(1991)325
- Wagemann, A., see Köppel, C. 570(1991)229
- Wagemann, A., see Köppel, C. 575(1992)87
- Wagh, J. S., Mokashi, A. A. and Datta, A.
A high-performance liquid chromatographic method for the monitoring and quantification of the synthesis of *p*-hydroxyphenylacetamide 587(1991)280
- Wagner, C. M., see Hopkins, N. K. 577(1992)87
- Wagner, H., see Oesch, F. 582(1992)232
- Wagner, J., see Kunath, A. 588(1991)352
- Wagner, R., see Ryll, T. 570(1991)77
- Wahl, P., see Moussaoui, M. 558(1991)71
- Wahl, P., see Moussaoui, M. 591(1992)115
- Wählin-Boll, E., see Rydberg, T. 564(1991)223
- Wainer, I. W., see Iredale, J. 573(1992)253
- Wainer, I. W., see Lloyd, D. K. 568(1991)117
- Wainer, I. W., see Lloyd, D. K. 578(1992)283
- Wainer, I. W., see Noctor, T. A. G. 577(1992)305
- Wainer, I. W., see Shibukawa, A. 574(1992)85
- Waisser, K., see Klimeš, J. 595(1992)334
- Wakabayashi, H., Nakajima, M., Yamato, S. and Shimada, K.
Determination of idebenone in rat serum and brain by high-performance liquid chromatography using platinum catalyst reduction and electrochemical detection 573(1992)154
- Wakabayashi, H., see Yamato, S. 590(1992)241
- Wakai, J., see Haginaka, J. 596(1992)151
- Wakamoto, K., see Baba, Y. 558(1991)273
- Wakmundzka-Hajnos, M., Wawrzynowicz, T. and Dzido, T. H.
Comparison of adsorption properties of Florisil and silica in high-performance liquid chromatography. I. Retention behaviour of monofunctional model solutes 600(1992)51
- Wakui, Y., Yanagisawa, E., Ishibashi, E., Matsuzaki, Y., Takeda, S., Sasaki, H., Aburada, M. and Oyama, T.
Determination of baicalin and baicalein in rat plasma by high-performance liquid chromatography with electrochemical detection 575(1992)131
- Waldhoff, H., see Bruns, A. 592(1992)249
- Walker, D. K., Smith, D. A. and Stopher, D. A.
Liquid-liquid extraction and high-performance liquid chromatography for the determination of a novel antidysrhythmic agent (UK-68,798) in human urine 568(1991)475
- Walker, L. V., Walsh, J. R. and Webber, J. J.
High-performance liquid chromatography of sulphonamides extracted from bovine and porcine muscle by solid-phase dispersion 595(1992)179
- Walker, L. V., see Walsh, J. R. 596(1992)211
- Wallace, D., see Brownsill, R. 562(1991)267
- Wallace, G. G., see Ge, H. 588(1991)25
- Wallace, S. M., see Hoffman, D. A. 565(1991)447
- Walsh, D. T., see Hecker, M. 574(1992)344
- Walsh, J. R., Walker, L. V. and Webber, J. J.
Determination of tetracyclines in bovine and porcine muscle by high-performance liquid chromatography using solid-phase extraction 596(1992)211
- Walsh, J. R., see Walker, L. V. 595(1992)179
- Walter, S. M., see Coutant, J. E. 570(1991)139
- Walters, R., see Lohmann, A. 564(1991)283
- Walters, R. R., see Wu, D. 598(1992)7
- Walubo, A., Chan, K. and Wong, C. L.
Simultaneous assay for isoniazid and hydrazine metabolite in plasma and cerebrospinal fluid in the rabbit 567(1991)261
- Wang, H., see Štulík, K. 552(1991)439
- Wang, J.-J., see Ho, S.-T. 570(1991)339
- Wang, J.-Y., see Yang, F. 582(1992)71
- Wang, K., see Thomare, P. 583(1992)217
- Wang, L. H., Smith, P. C., Anderson, K. L. and Fielding, R. M.
High-performance liquid chromatographic analysis of amphotericin B in plasma, blood, urine and tissues for pharmacokinetic and tissue distribution studies 579(1992)259
- Wang, Q.-S. and Fan, D.-P.
Optimization of separation of rare earths in high-performance thin-layer chromatography 587(1991)359
- Wang, T. and Hartwick, R. A.
Binary buffers for indirect absorption detection in capillary zone electrophoresis 589(1992)307
- Wang, T. and Hartwick, R. A.
Capillary modification and evaluation using streaming potential and frontal chromatography for protein analysis in capillary electrophoresis 594(1992)325
- Wang, T., see Wu, N. 582(1992)77
- Wang, W.-Q. and Gustafson, A.
One-dimensional thin-layer chromatographic separation of phospholipids and lysophospholipids from tissue lipid extracts 581(1992)139
- Wang, Y.-T., see Zeng, Z.-R. 589(1992)368
- Wang, Z., Xu, H. and Fu, C.
Sensitive fluorescence detection of some nitrosamines by precolumn derivatization with dansyl chloride and high-performance liquid chromatography 589(1992)349

- Wanwimolruk, S., Zoest, A. R., Wanwimolruk, S. Z. and Hung, C. T.
Sensitive high-performance liquid chromatographic determination of famotidine in plasma. Application to pharmacokinetic study 572(1991)227
- Wanwimolruk, S., see Shen, J. 565(1991)309
- Wanwimolruk, S. Z., see Wanwimolruk, S. 572(1991)227
- Ward, L. C., see Mason, S. R. 581(1992)267
- Ward, S. A., see Hoaksey, P. E. 566(1991)244
- Ward, S. A., see Mberu, E. K. 570(1991)180
- Ward, S. A., see Thomas, C. G. 583(1992)131
- Warner, C. D., Branch, S. K., Burke, J. F., Chambers, L. M. and Young, V. R.
Microderivatization of 4-[¹⁸O]hydroxyproline and quantitation with a benchtop mass spectrometer 562(1991)625
- Warner, I. M., see Anigbogu, V. C. 594(1992)37
- Warner, J. O., see Assadullahi, T. P. 565(1991)349
- Warthen, Jr., J. D., Demilo, A. B., Leonhardt, B. A., Lusby, W. R., Devilbiss, E. D. and Schreck, C. E.
Separation of *cis* and *trans* isomers from a mosquito repellent, ClC-4, via semi-preparative high-performance liquid chromatography and the repellent effect of each 590(1992)133
- Watabe, K., Kanda, H., Sato, K. and Hobo, T.
High-temperature continuous counter-current gas-liquid chromatography 590(1992)289
- Watanabe, A., Semba, J., Kurumaji, A., Kumashiro, S. and Toru, M.
Measurement of glutamate, aspartate and glycine and its potential precursors in human brain using high-performance liquid chromatography by pre-column derivatization with diethylaminoazobenzene sulphonyl chloride 583(1992)241
- Watanabe, A., see Nitta, K. 585(1991)173
- Watanabe, H., Sugahara, K., Inoue, K., Fujita, Y. and Kodama, H.
Liquid chromatographic-mass spectrometric analysis for screening of patients with cystinuria, and identification of cystine stone 568(1991)445
- Watanabe, K., Tohda, K., Sugimoto, H., Eitoku, F., Inoue, H., Suzuki, K. and Nakamura, S.
Ion-sensitive field effect transistor as a monovalent cation detector for ion chromatography and its application to the measurement of Na⁺ and K⁺ concentrations in serum 566(1991)109
- Watanabe, K., see Akasaka, K. 596(1992)197
- Watanabe, K., see Ishibashi, M. 562(1991)613
- Watanabe, K., see Nakanishi, A. 591(1992)159
- Watanabe, K., see Sato, J. 575(1992)63
- Watanabe, M., Watanabe, T., Hirasawa, N., Mue, S., Muramatsu, S., Matsushita, Y., Takahagi, H., Braquet, P., Broquet, C., Levine, L. and Ohuchi, K.
Preparation of immunoaffinity mini-columns for the analysis of platelet activating factor (PAF) in biological samples 597(1992)309
- Watanabe, M., see Kohata, K. 558(1991)131
- Watanabe, N., see O'Shea, T. J. 593(1992)305
- Watanabe, S., see Tomita, M. 566(1991)239
- Watanabe, T., see Fukuda, H. 567(1991)459
- Watanabe, T., see Nawata, M. 597(1992)415
- Watanabe, T., see Usui, T. 584(1992)213
- Watanabe, T., see Watanabe, M. 597(1992)309
- Watanabe, Y., see Kamahori, M. 567(1991)351
- Watanuki, K., see Matsumoto, G. I. 598(1992)267
- Waters, D. G., see Smith, I. D. 596(1992)290
- Watkins, P. J., see Halket, J. M. 562(1991)279
- Watkins, W. J., see Mallet, A. I. 562(1991)647
- Watkins, W. M., see Mberu, E. K. 570(1991)180
- Watkins, W. M., see Mberu, E. K. 581(1992)156
- Watson, D. G., Midgley, J. M. and McGhee, C. N. J.
Determination of (-)-bunolol and its metabolite, dihydro-(-)-bunolol, in human aqueous humour by gas chromatography-negative ion chemical ionisation mass spectrometry 571(1991)101
- Watson, D. G., see Macfarlane, R. G. 562(1991)585
- Watson, E. and Yao, F.
Fast protein monitoring in fermentation broth using non-porous micropellicular reversed-phase columns 594(1992)392
- Watson, J. T., see Schultz, G. A. 590(1992)329
- Watt, J. J., see Cooles, G. P. 588(1991)259
- Wattan, M. H., see Purnell, J. H. 555(1991)173
- Watts, M. T., Escarzaga, M. and Williams, C. H.
Gas chromatographic headspace analysis of sevoflurane in blood 577(1992)289
- Wawrzynowicz, T., see Waksmundzka-Hajnos, M. 600(1992)51
- Waymack, P. P., see Bernert, Jr., J. T. 578(1992)1
- Weaver, M. L., Tanzer, J. M. and Kramer, P. A.
High-performance liquid chromatographic determination of pilocarpine in plasma 581(1992)293
- Webber, J. J., see Walker, L. V. 595(1992)179
- Webber, J. J., see Walsh, J. R. 596(1992)211
- Weber, B., see Barberi, M. 565(1991)511
- Weber, B., see Barberi-Heyob, M. 573(1992)247
- Weber, B., see Barberi-Heyob, M. 581(1992)281
- Weber, C., Beetens, J., De Clerck, F. and Tegtmeier, F.
Gas chromatographic-mass spectrometric determination of 11-dehydrothromboxane B₂ in human urine 577(1992)1
- Weber, C., Höller, M., Beetens, J., De Clerck, F. and Tegtmeier, F.
Determination of 6-keto-PGF_{1α}, 2,3-dinor-6-keto-PGF_{1α}, thromboxane B₂, 2,3-dinor-thromboxane B₂, PGE₂, PGD₂ and PGF_{2α} in human urine by gas chromatography-negative ion chemical ionization mass spectrometry 562(1991)599
- Weber, L.
Gas chromatographic determination of urinary phenol conjugates after acid hydrolysis/extractive acetylation 574(1992)349
- Weber, L., see Schad, H. 593(1992)147
- Weber, M., see Chauret, N. 588(1991)281
- Weber, R., see Jiao, Y. 595(1992)364
- Weber, W. H., see De la Vigne, U. 553(1991)489
- Webster, H. K., see Ittarat, I. 582(1992)57

- Webster, L. K., Crinis, N. A., Stokes, K. H. and Bishop, J. F.
High-performance liquid chromatographic method for the determination of toremifene and its major human metabolites 565(1991)482
- Wehmeyer, K. R., see Liu, H. 577(1992)61
- Wehr, T., see Zhu, M. 559(1991)479
- Wei, H.-C. and Wu, C.-L.
Use of a Sep-Pak cartridge for preparative collection in gas chromatography 555(1991)302
- Wei, J.-L., see Wei, J.-Q. 552(1991)103
- Wei, J.-Q., Wei, J.-L. and Zhou, X.-T.
Computer-aided optimization of the experimental conditions for the isocratic reversed-phase high-performance liquid chromatographic separation of hormonal steroids 552(1991)103
- Wei-Bing, X., see Peng, S. 572(1991)73
- Weichmann, W., see Thomas, L. C. 587(1991)255
- Weigmann, H. and Hiemke, C.
Determination of clozapine and its major metabolites in human serum using automated solid-phase extraction and subsequent isocratic high-performance liquid chromatography with ultraviolet detection 583(1992)209
- Weinberg, D. S., see Fowler, W. K. 558(1991)235
- Weinreich, B., see Nitz, S. 557(1991)187
- Weirong, T., see Shen, G. 582(1992)258
- Weisleder, D., see Neff, W. E. 589(1992)353
- Welch, C. J., see Pirkle, W. H. 589(1992)45
- Welch, K. J. and Hoffman, N. E.
Physico-chemical properties of electron-acceptor stationary phases in liquid chromatography 591(1992)75
- Welch, R. M., see Brown, A. R. 578(1992)302
- Welham, K. J., see Abdulrahman, S. 562(1991)713
- Welinder, B. S., see Vidal, P. 573(1992)201
- Welinder, H., see Jönsson, B. 558(1991)247
- Welling, G. W., Hiemstra, Y., Feijlbrief, M., Örvell, C., Van Ede, J. and Welling-Wester, S.
Comparison of detergents for extraction and ion-exchange high-performance liquid chromatography of Sendai virus membrane proteins 599(1992)157
- Welling-Wester, S., see Welling, G. W. 599(1992)157
- Wellings, R., see Murray, S. 567(1991)289
- Wells, D., see Khetarpal, V. K. 567(1991)491
- Wells, T. G., Hendry, I. R. and Kearns, G. L.
Measurement of bumetanide in plasma and urine by high-performance liquid chromatography and application to bumetanide disposition 570(1991)235
- Wells, T. G., see Kearns, G. L. 574(1992)356
- Wels, G., see Werner, E. R. 570(1991)43
- Wen, K.-C., Huang, C.-Y. and Liu, F.-S.
Determination of cinnamic acid and paeoniflorin in traditional chinese medicinal preparations by high-performance liquid chromatography 593(1992)191
- Wendt, M. W., see Nevens, J. R. 597(1992)247
- Werner, A., Maeba, R., Shimasaki, H., Ueta, N. and Gerber, G.
Purine and pyrimidine compounds in murine peritoneal macrophages cultured *in vitro* 553(1991)205
- Werner, E. R., Werner-Felmayer, G., Fuchs, D., Hausen, A., Reibnegger, G., Wels, G., Yim, J. J., Pfeleiderer, W. and Wachter, H.
6-Pyruvoyl tetrahydropterin synthase assay in extracts of cultured human cells using high-performance liquid chromatography with fluorescence detection of biopterin. 570(1991)43
- Werner, R., see Taverna, M. 558(1991)105
- Werner-Felmayer, G., see Werner, E. R. 570(1991)43
- Westerdahl, G., see Elmroth, I. 598(1992)43
- Westerlund, D., see Caturla, M. C. 593(1992)69
- Westerlund, D., see Magnusson, O. 582(1992)1
- Westhoff, F. and Blaschke, G.
High-performance liquid chromatographic determination of the stereoselective biotransformation of the chiral drug praziquantel 578(1992)265
- Westlake, J. P., see Smith, R. M. 592(1992)85
- Westmark, P. A., see Coutant, J. E. 570(1991)139
- Weston, A., Brown, P. R., Jandik, P., Jones, W. R. and Heckenberg, A. L.
Factors affecting the separation of inorganic metal cations by capillary electrophoresis 593(1992)289
- Wetzel, H., see Süß, S. 565(1991)363
- Wetzel, R., see Brandl, G. 586(1991)249
- Wexler, J., see Bleiberg, B. 568(1991)301
- Weyhenmeyer, R., see Hammes, W. 567(1991)504
- Whelton, C., see Gupta, R. N. 582(1992)236
- Whigan, D. B. and Cohen, A. I.
High-performance liquid chromatographic determination of 1- β -D-arabinofuranosyl-*E*-5-(2-bromovinyl)uracil and its metabolite (*E*)-5-(2-bromovinyl)uracil in serum 568(1991)385
- White, I. N. H., see Martin, J. 568(1991)219
- White, L. B., see Rossomando, E. F. 583(1992)11
- White, L. B., see Rossomando, E. F. 583(1992)19
- White, T. C., see Demby, K. B. 579(1992)153
- Whiterock, V. J., see Russell, J. W. 572(1991)321
- Whiting, G. S., see Abraham, M. H. 587(1991)213
- Whiting, G. S., see Abraham, M. H. 587(1991)229
- Whiting, G. S., see Abraham, M. H. 588(1991)361
- Whiting, G. S., see Abraham, M. H. 594(1992)229
- Whitlow, S., see Buck, C. F. 594(1992)225
- Wičar, S.
Split injection into a capillary column at very low split ratios 557(1991)1
- Wichmann, J. K., see Clark, L. A. 573(1992)113
- Widmer, H. M., see Manz, A. 593(1992)253
- Widmer, H. M., see Morrissey, M. A. 552(1991)551
- Wiederkehr, F.
Analysis of cerebrospinal fluid proteins by electrophoresis (Review) 569(1991)281
- Wiederkehr, F., Büeler, M. R. and Vonderschmitt, D. J.
Chromatographic and electrophoretic studies of circulating immune complexes in plasma 566(1991)77
- Wiedmeier, V. T., see Hendrich, C. E. 577(1992)19
- Wieland, H., see Bisse, E. 575(1992)223

- Wieling, J., Coenegracht, P. M. J., Mensink, C. K., Jonkman, J. H. G. and Doornbos, D. A.
Selection of robust combinations of extraction liquid composition and internal standard. Monte Carlo simulation of improvement of assay methods with liquid-liquid extraction prior to high-performance liquid chromatography 594(1992)45
- Wientjes, M. G. and Au, J. L.-S.
High-performance liquid chromatographic analysis of 2',3'-dideoxynosine in biological samples 563(1991)400
- Wiese, B. and Hermansson, J.
Bioanalysis of diclofenac as its fluorescent carbazole acetic acid derivative by a post-column photoderivatization high-performance liquid chromatographic method 567(1991)175
- Wiese, G., see Kaijser, G. P. 571(1991)121
- Wiese, G., see Kaijser, G. P. 583(1992)175
- Wikström, M. and Ohlson, S.
Computer simulation of weak affinity chromatography 597(1992)83
- Wikström, M., see Eriksson, B.-M. 567(1991)1
- Wikström, M., see Eriksson, B.-M. 593(1992)185
- Wiktorowicz, J. E., see Cunico, R. L. 559(1991)467
- Wildfeuer, A., see Riedel, K.-D. 576(1992)358
- Wilhelm, J. A., Bailey, L. C., Shepard, T. A. and Venturella, V. S.
Simultaneous determination of phenolphthalein and phenolphthalein glucuronide from dog serum, urine and bile by high-performance liquid chromatography 578(1992)231
- Wilhelm, J. A., see Kvalo, L. T. 565(1991)391
- Wilke, A. V., see Cox, S. K. 564(1991)322
- Wilken, M. E., see Burger, B. V. 552(1991)137
- Wilkinson, D., see Anderson, L. W. 581(1992)195
- Wilkinson, M. L., see Stubbs, A. P. 570(1991)293
- Williams, C. H., see Watts, M. T. 577(1992)289
- Williams, J., see Marshall, T. 569(1991)323
- Williams, J. D., see Kaiser, Jr., R. E. 562(1991)3
- Williams, J. P., see Cannell, G. R. 564(1991)310
- Williams, K. M., see Marshall, T. 569(1991)323
- Williams, W. M., see Joly, J. M. 563(1991)392
- Wilsch-Irrgang, A., see Bruns, A. 592(1992)249
- Wilson, G. S., see Wimalasena, R. L. 572(1991)85
- Wilson, I. D., see Witherow, L. 553(1991)497
- Wilson, K. W., see Schwertner, H. A. 578(1992)297
- Wilson, T. C. M., Eisman, M. S. and Machulskis, G. E.
Quantitation of the novel anticonvulsant remacemide in rat and dog plasma and urine: application of the plasma methodology to measure the plasma protein binding of remacemide 582(1992)195
- Wimalasena, R. L. and Wilson, G. S.
Factors affecting the specific activity of immobilized antibodies and their biologically active fragments 572(1991)85
- Winkle, W., see Bruns, A. 592(1992)249
- Winkler, W., see Pollak, V. A. 596(1992)241
- Winstanley, P. A., see Mberu, E. K. 570(1991)180
- Winzor, D. J.
Recent developments in quantitative affinity chromatography (Review) 597(1992)67
- Winzor, D. J., Munro, P. D. and Jackson, C. M.
Study of high-affinity interactions by quantitative affinity chromatography. Analytical expressions in terms of total ligand concentration 597(1992)57
- Wirth, C., see Liebich, H. M. 572(1991)1
- Wise, S. A., see Craft, N. E. 589(1992)171
- Wise, S. A., see Epler, K. S. 595(1992)89
- Wishnok, J. S., see Day, B. W. 562(1991)563
- Witherow, L., Spurway, T. D., Ruane, R. J., Wilson, I. D. and Longdon, K.
Problems and solutions in chiral thin-layer chromatography: a two-phase "Pirkle" modified amino-bonded plate 553(1991)497
- Witkamp, R. F., see Van 't Klooster, G. A. E. 571(1991)157
- Witkiewicz, Z., Grajek, H. and Choma, J.
Chromatographic determination of the physico-chemical parameters of adsorption on active carbons 556(1991)441
- Witkiewicz, Z. and Lipkowski, J.
Foreword 600(1992)1
- Witkiewicz, Z., see Grajek, H. 600(1992)67
- Witkiewicz, Z., see Mazur, J. 600(1992)123
- Witkowska, H. E., see Shackleton, C. H. L. 562(1991)175
- Witte, D. T., Bosman, J., De Boer, T., Drenth, B. F. H., Ensing, K. and De Zeeuw, R. A.
Influence of chemical structure of tricyclic tertiary dimethylamines on chiral separation by reversed-phase high-performance liquid chromatography after derivatization with (-)-menthylchloroformate 553(1991)365
- Witte, D. T., Bosman, J., De Zeeuw, R. A., De Boer, J. H. and Doornbos, D. A.
Optimization of the direct enantiomeric separation by high-performance liquid chromatography of the D-2 dopamine agonist N-0437 using response surface methodology and the multi-criteria decision-making method 558(1991)333
- Wium, C. A., see Le Roux, A. M. 581(1992)306
- Wodecki, Z. J., Torlop, B. and Ślebioda, M.
Chromatographic determination of citric acid for monitoring the mould process 558(1991)302
- Woestenborghs, R., Lorreyne, W., Van Rompaey, F. and Heykants, J.
Determination of risperidone and 9-hydroxyrisperidone in plasma, urine and animal tissues by high-performance liquid chromatography 583(1992)223
- Woker, R., Champluvier, B. and Kula, M.-R.
Purification of S-oxynitrilase from *Sorghum bicolor* by immobilized metal ion affinity chromatography on different carrier materials 584(1992)85
- Wolf, A. P., see MacGregor, R. R. 590(1992)354
- Wolf, H., see Hessing, M. 599(1992)267
- Wolf, R. M., see Francotte, E. 595(1992)63
- Wong, C. L., see Walubo, A. 567(1991)261
- Wong, V. W. T., see Ang, S.-G. 599(1992)21
- Wong, Y. C., see Chan, K. 565(1991)247
- Wong, Y. W. J. and Skinner, M. H.
Rapid method for the determination of ketanserin in rat serum by high-performance liquid chromatography with fluorimetric detection 571(1991)318

- Wood, A. J. J., see Edeki, T. 577(1992)190
- Wood, A. J. J., see He, H. B. 574(1992)213
- Wood, G., see Harajiri, S. 575(1992)213
- Wood, M., see He, H. B. 574(1992)213
- Wood, S. and Langlois, S.
DNA typing in hereditary disease (Review) 569(1991)421
- Woodard, D., see Matassa, L. C. 573(1992)43
- Woodcock, E. A. and Tanner, J. K.
Preparation of samples for high-performance liquid chromatography of inositol phosphates 581(1992)134
- Woolf, E. and Matuszewski, B.
High-performance liquid chromatographic procedure for the determination of a non-nucleoside HIV-1 reverse transcriptase inhibitor in human plasma 577(1992)129
- Woollard, P. M., Salmon, J. A. and Padfield, A. D.
Use of high-performance liquid chromatography–thermospray mass spectrometry and gas chromatography–electron-impact mass spectrometry in the identification of the metabolites of α -methylacetoxyhydroxamic acids, potential anti-asthmatic agents 562(1991)249
- Worthen, H. G., see Liu, H. 579(1992)215
- Woutersen-van Nijnanten, F. M. A., see Van 't Klooster, G. A. E. 579(1992)158
- Woutersen-van Nijnanten, F. M. A., see Van 't Klooster, G. A. E. 579(1992)355
- Wright, B. E.
Construction of kinetic models to understand metabolism *in vivo* 566(1991)309
- Wright, M. R., Axelson, J. E., Abbott, F. S., Riggs, K. W., Van der Weyde, M. P., Taylor, S. M., McMorland, G. H. and Rurak, D. W.
Determination of ritodrine in biological fluids of the pregnant sheep by fused-silica capillary gas chromatography using electron-capture detection 565(1991)225
- Wright, M. R., Sattari, S., Brocks, D. R. and Jamali, F.
Improved high-performance liquid chromatographic assay method for the enantiomers of ibuprofen 583(1992)259
- Wroński, M.
Determination of volatile thiols by gas chromatography using separation as tributyltin mercaptides 555(1991)306
- Wu, C., Cheng, J., Gao, W., Zeng, Z., Lu, X. and Gong, S.
Preparation and capillary column gas chromatographic characterization of aza-crown ether polysiloxane stationary phases 594(1992)243
- Wu, C.-L., see Wei, H.-C. 555(1991)302
- Wu, C.-T., see Lee, C. S. 559(1991)133
- Wu, C.-Y., see Zeng, Z.-R. 589(1992)368
- Wu, D. and Walters, R. R.
Effects of stationary phase ligand density on high-performance ion-exchange chromatography of proteins 598(1992)7
- Wu, H.-L., see Chen, S.-H. 595(1992)203
- Wu, J., see Pawliszyn, J. 559(1991)111
- Wu, J.-R., see Huang, H.-S. 564(1991)195
- Wu, L., see Gilpin, R. K. 556(1991)415
- Wu, M. D., see Saska, M. 590(1992)147
- Wu, N., Wang, T., Hartwick, R. A. and Huie, C. W.
Separation of serum bilirubin species by micellar electrokinetic chromatography with direct sample injection 582(1992)77
- Wu, S.-M., see Chen, S.-H. 595(1992)203
- Wu, X., Haupt, K. and Vijayalakshmi, M. A.
Separation of immunoglobulin G by high-performance pseudo-bioaffinity chromatography with immobilized histidine. I. Preliminary report on the influence of the silica support and the coupling mode 584(1992)35
- Wünsche, L., Keller, U. and Flament, I.
Combination of supercritical fluid chromatography with thin-layer chromatography on a semi-preparative scale 552(1991)539
- Wurst, M., Kysilka, R. and Koza, T.
Analysis and isolation of indole alkaloids of fungi by high-performance liquid chromatography 593(1992)201
- Wurth, C., see Mardens, Y. 577(1992)341
- Wutka, R., see Haidacher, D. 591(1992)351
- Wylie, R., see Reinhart, P. 570(1991)425
- Wyss, R. and Bucheli, F.
Quantitative analysis of retinoids in biological fluids by high-performance liquid chromatography using column switching. III. Determination of the arotinoid sumarotene and its Z-isomer in human and animal plasma 576(1992)111
- Wyss, R. and Bucheli, F.
Use of direct injection precolumn techniques for the high-performance liquid chromatographic determination of the retinoids acitretin and 13-*cis*-acitretin in plasma 593(1992)55

X

- Xiao-Quan, S., see Peng, S. 572(1991)73
- Xie, M., Zhou, C., Ren, Z. and Luo, T.
Experimental model for predicting solute retention time as a function of mobile phase composition in ternary reversed-phase high-performance liquid chromatography 555(1991)33
- Xu, A. S., see Bronner, W. E. 580(1992)63
- Xu, D.-K., Zhou, J.-H., Yuan, Y.-S., Liu, X.-Q. and Huang, S.-K.
High-performance liquid chromatographic assay for amiloride in plasma and urine 567(1991)451
- Xu, H., see Wang, Z. 589(1992)349
- Xu, X., see Bruin, G. J. M. 559(1991)163

Y

- Yakabe, Y., Sudoh, Y. and Takahata, Y.
Immobilization method for polyethylene glycol using a cross-linking co-agent 558(1991)323
- Yakush, E. V., see Golovnya, R. V. 552(1991)1
- Yamada, K., see Kanai, M. 567(1991)415
- Yamada, M., see Kane, M. 569(1991)297
- Yamada, T., see Hattori, H. 564(1991)278
- Yamada, T., see Hattori, H. 579(1992)247
- Yamada, T., see Hattori, H. 581(1992)213
- Yamaguchi, M., Yamashita, K., Aoki, I., Tabata, T., Hirai, S.-I. and Yashiki, T.
Determination of manidipine enantiomers in human serum using chiral chromatography and column-switching liquid chromatography 575(1992)123
- Yamaguchi, M., see Ishida, J. 598(1992)203
- Yamaguchi, M., see Nakashima, K. 584(1992)275
- Yamaguchi, O., see Yamasaki, N. 597(1992)207
- Yamamoto, A., Matsunaga, A., Mizukami, E., Hayakawa, K. and Miyazaki, M.
Ligand-exchange ion chromatographic determination of malic acid enantiomers in apple juice with photometric detection 585(1991)315
- Yamamoto, F. M., see Imoto, T. 557(1991)383
- Yamamoto, H., Baumann, J. and Erni, F.
Electrokinetic reversed-phase chromatography with packed capillaries 593(1992)313
- Yamamoto, K., Konami, Y. and Osawa, T.
Determination of the carbohydrate-binding site of *Bauhinia purpurea* lectin by affinity chromatography 597(1992)221
- Yamamoto, K., see Honda, S. 588(1991)327
- Yamamoto, K., see Jinno, K. 594(1992)105
- Yamamoto, K., see Konami, Y. 597(1992)213
- Yamamoto, K., see Nagata, Y. 575(1992)147
- Yamamoto, M., see Miwa, H. 568(1991)25
- Yamamoto, S. and Sano, Y.
Short-cut method for predicting the productivity of affinity chromatography 597(1992)173
- Yamamoto, S., see Hattori, H. 564(1991)278
- Yamamoto, S., see Hattori, H. 579(1992)247
- Yamamoto, S., see Hattori, H. 581(1992)213
- Yamamoto, S., see Ohmori, S. 574(1992)35
- Yamamoto, Y., see Kamata, K. 596(1992)233
- Yamamoto, Y., see Kane, M. 569(1991)297
- Yamamura, Y., Kawakami, J., Santa, T., Kotaki, H., Uchino, K., Sawada, Y. and Iga, T.
Selective high-performance liquid chromatographic method for the determination of glycyrrhizin and glycyrrhetic acid-3-O-glucuronide in biological fluids: application of ion-pair extraction and fluorescence labelling agent 567(1991)151
- Yamanaka, K., see Kohata, K. 558(1991)131
- Yamanashi, S., see Imanari, T. 574(1992)142
- Yamane, M. and Abe, A.
High-performance liquid chromatography-thermospray mass spectrometry of prostaglandin and thromboxane acetyl derivatives 568(1991)11
- Yamane, M. and Abe, A.
High-performance liquid chromatography-thermospray mass spectrometry of hydroxy-polyunsaturated fatty acid acetyl derivatives 575(1992)7
- Yamane, M., Abe, A., Yamane, S. and Ishikawa, F.
High-performance liquid chromatography-thermospray mass spectrometry of hydroperoxy polyunsaturated fatty acid acetyl derivatives 579(1992)25
- Yamane, S., see Yamane, M. 579(1992)25
- Yamasaki, N., Kimura, M., Yamaguchi, O. and Araki, M.
Isolation and characterization of isolectins from *Erythrina variegata* seeds 597(1992)207
- Yamashita, H., see Moriyama, T. 571(1991)61
- Yamashita, K., Motohashi, M. and Yashiki, T.
Column-switching techniques for high-performance liquid chromatography of ibuprofen and mefenamic acid in human serum with short-wavelength ultraviolet detection 570(1991)329
- Yamashita, K., Motohashi, M. and Yashiki, T.
Automated high-performance liquid chromatographic method for the simultaneous determination of cefotiam and Δ^3 -cefotiam in human plasma using column switching 577(1992)174
- Yamashita, K., see Yamaguchi, M. 575(1992)123
- Yamato, S., Nakajima, M., Wakabayashi, H. and Shimada, K.
Specific detection of acetyl-coenzyme A by reversed-phase ion-pair high-performance liquid chromatography with an immobilized enzyme reactor 590(1992)241
- Yamato, S., see Wakabayashi, H. 573(1992)154
- Yamatodani, A., see Fukuda, H. 567(1991)459
- Yamazaki, K., see Takamatsu, T. 584(1992)261
- Yan, C., see Riester, R. L. 588(1991)289
- Yan, Z., see Peng, S. 572(1991)73
- Yanagihara, Y., see Hirata, N. 592(1992)93
- Yanagisawa, E., see Wakui, Y. 575(1992)131
- Yang, F., Zhu, D.-W., Wang, J.-Y. and Lin, S.-X.
Rapid purification yielding highly active 17β -hydroxysteroid dehydrogenase: application of hydrophobic interaction and affinity fast protein liquid chromatography 582(1992)71
- Yang, L.-L., see Cheng, F.-C. 582(1992)19
- Yang, S. S.
Determination of maleic hydrazide in tobacco by micellar liquid chromatography 595(1992)346
- Yang, Y., Velayudhan, A., Ladisch, C. M. and Ladisch, M. R.
Protein chromatography using a continuous stationary phase 598(1992)169
- Yang, Y.-B., Harrison, K., Carr, D. and Guiochon, G.
Factors affecting the separation and loading capacity of proteins in preparative gradient elution high-performance liquid chromatography 590(1992)35
- Yang, Z. C. and Cashman, J. R.
Structure-retention index relationships for derivatized monosaccharides on non-polar gas chromatography columns 596(1992)79

- Yang, Z. C., see Peng, C. T. 586(1991)85
- Yang, Z. C., see Peng, C. T. 586(1991)113
- Yano, Y., see Ohmori, S. 566(1991)1
- Yao, F., see Watson, E. 594(1992)392
- Yao, K., Ubuka, T., Masuoka, N., Kinuta, M., Ohta, J., Teraoka, T. and Futani, S.
Assay of sialidase activity using ion-exchange chromatography and acidic ninhydrin reaction 581(1992)11
- Yap, A. S., see Cannell, G. R. 564(1991)310
- Yap, B. K., Johnston, G. A. R. and Kazlauskas, R.
Routine screening and quantitation of urinary corticosteroids using bench-top gas chromatography-mass-selective detection 573(1992)183
- Yarbro, J. W., see Mitchell, E. P. 581(1992)31
- Yarwood, C., see Janero, D. R. 573(1992)207
- Yashiki, T., see Aoki, I. 571(1991)283
- Yashiki, T., see Yamaguchi, M. 575(1992)123
- Yashiki, T., see Yamashita, K. 570(1991)329
- Yashiki, T., see Yamashita, K. 577(1992)174
- Yasuda, H., see Haginaka, J. 592(1992)301
- Yasuda, H., see Haginaka, J. 598(1992)67
- Yasui, T., Yuchi, A., Wada, H. and Nakagawa, G.
Reversed-phase high-performance liquid chromatography of several metal-8-quinolinethiol complexes 596(1992)73
- Yasukawa, K., Abe, F., Shida, N., Koizumi, Y., Uchida, T., Noguchi, K. and Shima, K.
High-performance affinity chromatography system for the rapid, efficient assay of glycated albumin 597(1992)271
- Yasunaga, T., see Shimohigashi, Y. 597(1992)425
- Yau, W. W. and Kirkland, J. J.
Improved computer algorithm for characterizing skewed chromatographic band broadening. I. Method 556(1991)111
- Yau, W. W., see Kirkland, J. J. 593(1992)339
- Yee, A. B. K., see Doerge, D. R. 586(1991)158
- Yee, G. T., see Browne, S. P. 575(1992)158
- Yeleswaram, K., Axelson, J. E. and Rurak, D. W.
Sensitive microbore high-performance liquid chromatographic assay for labetalol in the biological fluids of pregnant sheep 565(1991)383
- Yeo, S. K., Lee, H. K. and Li, S. F. Y.
Separation of antibiotics by high-performance capillary electrophoresis with photodiode-array detection 585(1991)133
- Yeo, S. K., Lee, H. K. and Li, S. F. Y.
Separation of plant growth regulators by capillary electrophoresis 594(1992)335
- Yeung, E. S., see Lee, T. 565(1991)197
- Yeung, E. S., see Lee, T. T. 595(1992)319
- Yeung, E. S., see Pfeffer, W. D. 557(1991)125
- Yewey, G., see Reinhart, P. 570(1991)425
- Yik, Y. F., Lee, H. K., Li, S. F. Y. and Khoo, S. B.
Micellar electrokinetic capillary chromatography of vitamin B₆ with electrochemical detection 585(1991)139
- Yik, Y. F., Ong, C. P., Khoo, S. B., Lee, H. K. and Li, S. F. Y.
Separation of polycyclic aromatic hydrocarbons by micellar electrokinetic chromatography with cyclodextrins as modifiers 589(1992)333
- Yim, J. J., see Werner, E. R. 570(1991)43
- Yim, K. W.
Fractionation of the human recombinant tissue plasminogen activator (rtPA) glycoforms by high-performance capillary zone electrophoresis and capillary isoelectric focusing 559(1991)401
- Yinon, J. and Saar, J.
Analysis of dyes extracted from textile fibers by thermospray high-performance liquid chromatography-mass spectrometry 586(1991)73
- Yisheng, Y., see Min, S. 581(1992)272
- Yokota, Y., see Hirokawa, T. 555(1991)247
- Yokota, Y., see Hirokawa, T. 555(1991)321
- Yokota, Y., see Hirokawa, T. 585(1991)297
- Yokouchi, Y. and Sano, M.
Trace determination of volatile organic compounds in soil based on thermal vaporization followed by Tenax-GC trapping and capillary gas chromatography-mass spectrometry 555(1991)297
- Yokoyama, T. and Kinoshita, T.
High-performance liquid chromatographic determination of taurine in biological fluids by post-column fluorescence reaction with thiamine 568(1991)212
- Yokoyama, Y. and Sato, H.
Reversed-phase high-performance liquid chromatographic determination of linear alkylbenzenesulphonates in river water at ppb levels by precolumn concentration 555(1991)155
- Yokoyama, Y., Sato, H., Tsuchiya, M. and Kakinuma, H.
Simultaneous determination of urinary creatinine and aromatic amino acids by cation-exchange chromatography with ultraviolet detection 566(1991)19
- Yokoyama, Y., Tsuchiya, M., Sato, H. and Kakinuma, H.
Determination of creatinine and ultraviolet-absorbing amino acids and organic acids in urine by reversed-phase high-performance liquid chromatography 583(1992)1
- Yoshida, H., see Morita, I. 576(1992)334
- Yoshida, H., see Tanaka, S. 587(1991)364
- Yoshida, K., see Kurono, M. 578(1992)152
- Yoshida, K., see Nakashima, K. 584(1992)275
- Yoshida, K., see Noguchi, H. 573(1992)336
- Yoshida, T.
Determination of *p*-chloronitrobenzene in plasma by reversed-phase high-performance liquid chromatography 570(1991)321
- Yoshida, Y., see Oda, Y. 572(1991)133
- Yoshimura, H., see Endoh, Y. S. 579(1992)123
- Yoshimura, T., see Ikegawa, S. 577(1992)229
- Yoshioka, M., Mukai, Y., Matsui, T., Udagawa, A. and Funakubo, H.
Immobilization of ultra-thin layer of monoclonal antibody on glass surface 566(1991)361
- Yoshizumi, H., see Niwa, T. 578(1992)109
- Youn, D. Y., Yun, S. J. and Jung, K.-H.
Improved algorithm for resolution of overlapped asymmetric chromatographic peaks 591(1992)19
- Young, V. R., see Warner, C. D. 562(1991)625
- Yu, P. H., see Zuo, D. M. 567(1991)381
- Yuan, D. and Pietrzyk, D. J.
Separation and indirect detection of amino acids as acetylated derivatives 557(1991)315

- Yuan, L.-C., see Snider, J. 599(1992)141
 Yuan, X. W., see Basak, A. 581(1992)17
 Yuan, Y.-S., see Xu, D.-K. 567(1991)451
 Yu, L., see Jacob, III, P. 583(1992)145
 Yuchi, A., see Yasui, T. 596(1992)73
 Yun, S. J., see Youn, D. Y. 591(1992)19
 Yunes, R. A., see Heinzen, V. E. F. 598(1992)243
 Yuthavong, Y., see Ittarat, I. 582(1992)57
 Yvon, M., see Chabanet, C. 599(1992)211
- Z**
- Zabay, J. M., see Ramis, I. 557(1991)507
 Zacchei, A. G., see Berna, R. A. 563(1991)458
 Zacchei, P., see Cartoni, G. P. 552(1991)197
 Zacchello, F., see Cracco, A. T. 574(1992)156
 Zacchello, G., see Cracco, A. T. 574(1992)156
 Zacchigna, M., see Maurich, V. 566(1991)453
 Zachariou, M. and Hearn, M. T. W.
 High-performance liquid chromatography of amino acids, peptides and proteins. CXXI. 8-Hydroxyquinoline-metal chelate chromatographic support: an additional mode of selectivity in immobilized-metal affinity chromatography 599(1992)171
 Zahradníček, M., see Klimeš, J. 584(1992)221
 Zahradníčková, H., Maršálek, B. and Polišenská, M.
 High-performance thin-layer chromatographic and high-performance liquid chromatographic determination of abscisic acid produced by cyanobacteria 555(1991)239
 Zakett, D., see Kaiser, Jr., R. E. 562(1991)3
 Zamir, I.
 Derivatization of saturated long-chain fatty acids with phenacyl bromide in non-ionic micelles 586(1991)347
 Zamir, I., Grushka, E. and Chemke, J.
 Separation and determination of saturated very-long-chain free fatty acids in plasma of patients with adrenoleukodystrophy using solid-phase extraction and high-performance liquid chromatographic analysis 567(1991)319
 Zamir, I., Grushka, E. and Cividalli, G.
 High-performance liquid chromatographic analysis of free palmitic and stearic acids in cerebrospinal fluid 565(1991)424
 Zanchetta, S., see Sturaro, A. 590(1992)223
 Zane, P. A., see Chovan, J. P. 578(1992)77
 Zangirolami, L., see Mariot, R. 576(1992)143
 Zanolio, G., see Giachetti, C. 585(1991)111
 Zare, R. N., see Tsuda, T. 559(1991)103
 Zecca, L., Ferrario, P. and Costi, P.
 Determination of diclofenac and its metabolites in plasma and cerebrospinal fluid by high-performance liquid chromatography with electrochemical detection 567(1991)425
 Zecca, L., see Pietta, P. 558(1991)296
 Zecca, L., see Rubino, F. M. 579(1992)1
 Zech, K., De Jong, G. J. and Hupe, K.-P.
 Foreword 600(1992)161
 Zech, K. and Herzog, R.
 Two-dimensional high-performance liquid chromatography at low ng/ml levels of the anti-proliferative agent B859-35 in serum with automated sample clean-up, solid-phase trapping and ultraviolet detection 553(1991)55
 Zedde, C., see Baltas, M. 600(1992)323
 Zee, T., Stellaard, F. and Jakobs, C.
 Analysis of pipecolic acid in biological fluids using capillary gas chromatography with electron-capture detection and [²H₁₁]pipecolic acid as internal standard 574(1992)335
 Zeitoun, M. A. M., see Neff, W. E. 589(1992)353
 Zemaitis, M. A., see Schmith, V. D. 568(1991)253
 Zeng, Z., see Wu, C. 594(1992)243
 Zeng, Z.-R., Wu, C.-Y., Fang, X.-H., Huang, Z.-F. and Wang, Y.-T.
 Study of dihydroxy-substituted saturated urushiol crown ether as a stationary phase in capillary gas chromatography 589(1992)368
 Zenki, T., see Komatsu, S. 568(1991)460
 Zezza, F., Kerner, J., Pascale, M. R., Giannini, R. and Arrigoni Martelli, E.
 Rapid determination of amino acids by high-performance liquid chromatography: release of amino acids by perfused rat liver 593(1992)99
 Zhang, A. Q., Mitchell, S. C., Ayesh, R. and Smith, R. L.
 Determination of trimethylamine and related aliphatic amines in human urine by head-space gas chromatography 584(1992)141
 Zhang, J. Z., see Chen, D. Y. 559(1991)237
 Zhang, R., see Liao, J.-L. 586(1991)21
 Zhang, S., see Lou, X. 586(1991)139
 Zhao, H., see Fan, C.-D. 570(1991)217
 Zhao, M., see Liu, S. 598(1992)298
 Zhiri, A., see Del Cont Bernard, A. 574(1992)365
 Zhong, W. Z.
 Determination of 3-(5-cyclopropyl-1,2,4-oxadiazol-3-yl)-5-(1-methylethyl)imidazo[1,5-a]quinoxalin-4(5H)-one in serum by high-performance liquid chromatography 563(1991)427
 Zhou, C., see Xie, M. 555(1991)33
 Zhou, G., Jauhainen, M., Stevenson, K. and Dolphin, P. J.
 Human plasma lecithin:cholesterol acyltransferase. Preparation and use of immobilized *p*-aminophenyl-arsenoxide as a catalytic site-directed covalent ligand in enzyme purification 568(1991)69
 Zhou, J.-H., see Xu, D.-K. 567(1991)451
 Zhou, L., see Guan, Y. 552(1991)187
 Zhou, L., see Lou, X. 552(1991)153
 Zhou, L., see Lou, X. 586(1991)139
 Zhou, W., Liu, W. and An, D.
 Simultaneous ion chromatographic determination of chloride and calcium contained in Ringer's injection using a chelating agent as eluent 589(1992)358

- Zhou, X.-T., see Wei, J.-Q. 552(1991)103
- Zhu, B.-Y., Mant, C. T. and Hodges, R. S.
Mixed-mode hydrophilic and ionic interaction chromatography rivals reversed-phase liquid chromatography for the separation of peptides 594(1992)75
- Zhu, D.-W., see Yang, F. 582(1992)71
- Zhu, J., Katti, A. M. and Guiochon, G.
Comparison of various isotherm models for predicting competitive adsorption data 552(1991)71
- Zhu, J., Ng, J. and Filippich, L. J.
Determination of tannic acid and its phenolic metabolites in biological fluids by high-performance liquid chromatography 577(1992)77
- Zhu, M., Rodriguez, R. and Wehr, T.
Optimizing separation parameters in capillary isoelectric focusing 559(1991)479
- Zhuravleva, I. L., see Golovnya, R. V. 552(1991)1
- Ziegler, M., Dubiel, W., Henke, W., Gerber, G., Pimenov, A. M., Tikhonov, Y. V. and Toguzov, R. T.
Application of ion-pair high-performance liquid chromatography with radioisotope detection to *in vitro* studies of nucleoside metabolism in mitochondria 563(1991)172
- Ziegler, R. G., see Epler, K. S. 595(1992)89
- Zilli, M. A.
Simple and fast chromatographic method for the simultaneous determination of vanillylmandelic acid and homovanillic acid in human urine 570(1991)371
- Zima, D., see Reichelová, V. 588(1991)147
- Zima, J., see Beran, M. 558(1991)265
- Zimina, T. M., Kever, J. J., Melenevskaya, E. Y. and Fell, A. F.
Analysis of block copolymers by high-performance liquid chromatography under critical conditions 593(1992)233
- Zimmerhackl, L. B., Leuk, B. and Hoschützky, H.
The cytoskeletal protein villin as a parameter for early detection of tubular damage in the human kidney 587(1991)81
- Zimmerman, C. L., Ratna, S., LeBoeuf, E. and Pang, K. S.
High-performance liquid chromatographic method for the direct determination of 4-methylumbelliferone and its glucuronide and sulfate conjugates. Application to studies in the single-pass *in situ* perfused rat intestine-liver preparation 563(1991)83
- Zimmerman, H., see Herweck, U. 600(1992)312
- Zimmermann, J., Dennhardt, R. and Gramm, H.-J.
Measurement of plasma catecholamines by high-performance liquid chromatography with electrochemical detection in intensive care patients after dobutamine infusion 567(1991)240
- Zimmermann, T., see Riedel, K.-D. 576(1992)358
- Zlatkin, I. V., see Řezanka, T. 558(1991)215
- Zobel, H., Helms, M. and Dittberner, K.
Quantitative structure-type analysis of heterocompound-hydrocarbon mixtures. VI. Characterization of the nature of neutral heterocompounds in a product of brown coal refining 553(1991)441
- Zoest, A. R., see Wanwimolruk, S. 572(1991)227
- Zografakis, J., see Mangino, M. J. 577(1992)157
- Zolotov, Y. A., see Obrezkov, O. N. 558(1991)209
- Zomer, G., see Van Look, L. J. 564(1991)451
- Zonta, F., Stancher, B., Bogoni, P. and Masotti, P.
High-performance liquid chromatography of okadaic acid and free fatty acids in mussels 594(1992)137
- Zuccaro, P., Altieri, I., Betto, P., Pacifici, R., Ricciarello, G., Pini, L. A., Sternieri, E. and Pichini, S.
Determination of naltrexone and 6 β -naltrexol in plasma by high-performance liquid chromatography with coulometric detection 567(1991)485
- Zuccaro, P., see Pichini, S. 568(1991)267
- Zuccaro, P., see Pichini, S. 577(1992)358
- Zuccaro, P., see Pichini, S. 581(1992)143
- Zuccato, E., see Salomoni, M. 572(1991)312
- Zucchetti, G., see Menichini, E. 555(1991)211
- Zuk, M. M., Rihter, B. D., Kenney, M. E., Rodgers, M. A. J. and Kreimer-Birnbaum, M.
Photosensitizers for tumor therapy: determination of bis(di-isobutyl octadecylsiloxy)silicon 2,3-naphthalocyanine (isoBOSINC) in rat tissue and serum by high-performance liquid chromatography 568(1991)437
- Zuo, D. M. and Yu, P. H.
High-performance liquid chromatographic procedure for the simultaneous determination of aromatic L-amino acid decarboxylase activity towards 3,4-dihydroxyphenylalanine and 5-hydroxytryptophan 567(1991)381
- Zuting, P., see Tomasella, F. P. 587(1991)325
- Zweigenbaum, J., see Johansson, I. M. 554(1991)311
- Zweipfenning, P. G. M.
High-performance liquid chromatographic determination of morphine, morphine-3-glucuronide, morphine-6-glucuronide and codeine in biological samples using multi-wavelength forward optical detection: a reply 578(1992)333
- Žygys, A., see Maruška, A. 596(1992)157

SUBJECT INDEX

VOLUMES 551–600

A

Absorbance ratio

Assessment of chromatographic peak purity by means of multi-wavelength detection and correlation-based algorithms 592(1992)27

Acid–base separations

Analysis of biomass pyrolysis oils by a combination of various liquid chromatographic techniques and gas chromatography–mass spectrometry 553(1991)211

α_1 -Acid glycoprotein-bonded phase

Retention processes on α_1 -acid glycoprotein-bonded stationary phase 591(1992)55

Acidic/hydrothermal treatment

Impact of acidic/hydrothermal treatment on pore structural and chromatographic properties of porous silicas. I. The conventional approach 556(1991)395

Acidity

Determination of major ions in snow and ice cores by ion chromatography 594(1992)225

Actinobacillus actinomycetemcomitans

Intra-injector methylation of free fatty acids from aerobically and anaerobically cultured *Actinobacillus actinomycetemcomitans* and *Haemophilus aphrophilus* 576(1992)328

Activated carbon fibres

Chromatographic determination of the physico-chemical parameters of adsorption on activated carbon fibres 600(1992)67

Activated charcoal traps

Production and use of capillary traps for headspace gas chromatography of airborne volatile organic compounds 552(1991)137

Activation methods

Activation of matrices by 4,6-diphenylthieno[3,4-*d*]-1,3-dioxol-2-one-5,5-dioxide. High-performance liquid affinity chromatographic separations 589(1992)101

Active carbon

Chromatographic determination of the physico-chemical parameters of adsorption on active carbons 556(1991)441

Activity coefficients

Thermodynamics of solution of non-mesomorphic solutes at infinite dilution in the smectic-A, nematic and isotropic phases of *p*-*n*-octyl-*p*'-cyanobiphenyl. A gas–liquid chromatographic study 556(1991)317

UNIFAC model as a heuristic guide for estimating retention in chromatography 586(1991)1

Activity coefficients at infinite dilution determined by gas–liquid chromatography. Organic solvents in Apiezon L 586(1991)303

Activity coefficients in binary liquid mixtures measured by reversed-flow gas chromatography 595(1992)237

Activity testing

Activity testing and surface characterization of pretreated fused-silica capillaries for gas chromatography. A new modification of existing intermediate column tests 590(1992)271

Adsorbate–adsorbent interactions

Chromatographic determination of the physico-chemical parameters of adsorption on active carbons 556(1991)441

Adsorbents

see also Affinity adsorbents

New Czechoslovak sorbent, Ekosorb 552(1991)449

Study of microporous carbons by gas chromatographic determination of heats of physisorption 555(1991)183

New solid adsorbents for the separation of lower hydrocarbons and permanent gases. II. Ammonium molybdophosphate 556(1991)425

- Determination of hexahydrophthalic anhydride in air using gas chromatography 558(1991)247
- Protection of terpenes against oxidative and acid decomposition on adsorbent cartridges 589(1992)385
- Cellulose bearing covalently linked copper phthalocyanine trisulphonate as an adsorbent selective for polycyclic compounds and its use in studies of environmental mutagens and carcinogens (Review) 597(1992)37
- Theory of perfusion chromatography 599(1992)87
- Use of basic alumina in fractionation of fossil fuels 600(1992)243
- Adsorption**
- Adsorption and partition mode in high-performance liquid chromatography of highly polar solutes on silica 552(1991)23
- Gas chromatographic investigation of silica supports modified in a high-frequency low-temperature plasma 552(1991)179
- Evaluation of the application of liquid-phase titration to the examination of the adsorption activity of silica-based high-performance liquid chromatographic packings 552(1991)463
- High-performance liquid chromatography of amino acids, peptides and proteins. CXIV. Protein interactions with porous coulombic sorbents: comparison of experimental findings with predictions of several adsorption models 557(1991)335
- Theoretical analysis for measurement of building pollution parameters by gas chromatography 557(1991)469
- Immunoaffinity chromatography in biorecovery: an application of recombinant DNA technology to generic adsorption processes 584(1992)59
- Cation-exchange liquid chromatography of choline and acetylcholine on free shielded silanols of silica-based reversed-phase stationary phases 596(1992)43
- Theory of perfusion chromatography 599(1992)87
- Study of the adsorption of self-associating proteins on an anion exchanger. Application to the chromatography of β -lactoglobulin B 599(1992)255
- Comparison of adsorption properties of Florisil and silica in high-performance liquid chromatography. I. Retention behaviour of monofunctional model solutes 600(1992)51
- Chromatographic determination of the physico-chemical parameters of adsorption on activated carbon fibres 600(1992)67
- Adsorption capacity**
- Protein adsorption capacity of a porous phenylalanine-containing membrane based on a polyethylene matrix 586(1991)27
- Adsorption energy**
- Estimation of cellulose polarity by gas chromatography 552(1991)365
- Study of microporous carbons by gas chromatographic determination of heats of physisorption 555(1991)183
- Adsorption isotherms**
- Comparison of various isotherm models for predicting competitive adsorption data 552(1991)71
- Evaluation of the application of liquid-phase titration to the examination of the adsorption activity of silica-based high-performance liquid chromatographic packings 552(1991)463
- New approach for calculating ideal chromatograms from arbitrary composite distribution isotherms 556(1991)95
- Analysis of isotachic patterns in displacement chromatography 556(1991)119
- Prediction of single and binary profiles in overloaded elution chromatography using various semi-ideal models 556(1991)205
- Adsorption isotherms of phenylalanine in a chromatographic column measured simultaneously by system peaks analysis and frontal analysis 556(1991)277
- High-performance liquid chromatography on dynamically modified silica. IX. Modification of silica with 3-(N,N-dimethylpalmitylammonium) propanesulphonate for reversed-phase chromatography 556(1991)353
- Dynamic gas-solid chromatographic techniques for characterizing carbon molecular sieves 556(1991)433
- Chromatographic determination of the physico-chemical parameters of adsorption on active carbons 556(1991)441
- Adsorption-desorption isotherm hysteresis of β -lactoglobulin A with a weakly hydrophobic surface 557(1991)369
- Adsorption and elution of bovine γ -globulin using an affinity membrane containing hydrophobic amino acids as ligands 585(1991)45
- Prediction of the elution profiles of high-concentration bands in gas chromatography 589(1992)223
- Theory of optimization of ideal displacement chromatography of binary mixtures 590(1992)3
- Displacement effects in preparative gradient high-performance liquid chromatographic separations 590(1992)17
- Optical resolution by simulated moving-bed adsorption technology 590(1992)113
- Glucose-fructose equilibria on Dowex Monosphere 99 CA resin under overloaded conditions 590(1992)147

- Optimization of the experimental conditions and the column design parameters in overloaded elution chromatography 591(1992)31
- Experimental determination of adsorption isotherm data for the study of the surface energy distribution of various solid surfaces by inverse gas-solid chromatography 591(1992)233
- Precision and accuracy of the gas-solid adsorption isotherms derived by the elution-by-characteristic-points method 591(1992)245
- Validity of the model used to relate the energy distribution and the adsorption isotherm 591(1992)267
- Investigation of solid surfaces by high-performance liquid chromatography 592(1992)121
- Modelling of displacement chromatography using non-ideal isotherms 594(1992)9
- Ion- and ligand-exchange chromatography of proteins using porous zirconium oxide supports in organic and inorganic Lewis base eluents 596(1992)27
- Contribution of ionically immobilized bovine serum albumin to the retention of enantiomers 600(1992)37
- Chromatographic determination of the physico-chemical parameters of adsorption on activated carbon fibres 600(1992)67
- Effect of column material on sorption isotherms obtained by inverse gas chromatography 600(1992)341
- Adsorption kinetics**
- Adsorption kinetics of albumin on a cross-linked cellulose chromatographic ion exchanger 590(1992)101
- Computer simulation of weak affinity chromatography 597(1992)83
- Membrane-based receptor affinity chromatography 597(1992)155
- Kinetic aspects of membrane-based immunoaffinity chromatography 597(1992)167
- Adsorption measurements**
- Capillary modification and evaluation using streaming potential and frontal chromatography for protein analysis in capillary electrophoresis 594(1992)325
- Adsorption traps**
- Production and use of capillary traps for headspace gas chromatography of airborne volatile organic compounds 552(1991)137
- Affinity adsorbents**
- Bead cellulose derivatives as supports for immobilization and chromatographic purification of proteins 552(1991)389
- Modification of reversed-phase columns with dyed surfactants. Preparation of mechanically resistant efficient immobilized dyes for protein purification 558(1991)43
- Comparison of the performance of immunosorbents prepared by site-directed or random coupling of monoclonal antibodies 558(1991)55
- Adsorption and elution of bovine γ -globulin using an affinity membrane containing hydrophobic amino acids as ligands 585(1991)45
- Protein adsorption capacity of a porous phenylalanine-containing membrane based on a polyethylene matrix 586(1991)27
- High-performance affinity chromatography of NADP⁺ dehydrogenases from cell-free extracts using a nucleotide analogue as general ligand 586(1991)51
- Immobilization of Fv antibody fragments on porous silica and their utility in affinity chromatography 587(1991)161
- Optimization of oxidation of glycoproteins: an assay for predicting coupling to hydrazide chromatographic supports 587(1991)171
- Efficient substitution of 1,1'-carbonyldiimidazole activated cellulose and Sepharose matrices with amino acyl spacer arms 587(1991)271
- High-performance affinity chromatography of messenger RNA 588(1991)157
- Cellulose beads: a weak leaking affinity support 589(1992)93
- Characterization of synthetic macroporous packing materials in low-pressure cartridges and columns 590(1992)83
- Oriented immobilization of peptide ligands on solid supports 591(1992)99
- Determination of catecholamines in urine by liquid chromatography and electrochemical detection after on-line sample purification on immobilized boronic acid 593(1992)185
- Immobilization of monoclonal antibodies for affinity chromatography using a chelating peptide 595(1992)113
- Simultaneous purification of the neuroproteins synapsin I and synaptophysin 596(1992)51
- Silica sorbents with one- and two-site attached bacitracin in affinity chromatography 596(1992)114
- Trypsin and affinity chromatography (Review) 597(1992)3
- Galactosylation as a tool for the stabilization and immobilization of proteins (Review) 597(1992)19

- Study of high-affinity interactions by quantitative affinity chromatography. Analytical expressions in terms of total ligand concentration 597(1992)57
- Recent developments in quantitative affinity chromatography (Review) 597(1992)67
- Immobilized lectin columns: useful tools for the fractionation and structural analysis of oligosaccharides (Review) 597(1992)111
- Immobilization of protein ligands with methyl vinyl ether-maleic anhydride copolymer 597(1992)123
- Affinity purification of proteins using expanded beds 597(1992)129
- Immobilized ferredoxins for affinity chromatography of ferredoxin-dependent enzymes 597(1992)147
- Membrane-based receptor affinity chromatography 597(1992)155
- Kinetic aspects of membrane-based immunoaffinity chromatography 597(1992)167
- Novel affinity separations based on perfluorocarbon emulsions. Use of a perfluorocarbon affinity emulsion for the purification of human serum albumin from blood plasma in a fluidised bed 597(1992)189
- Improved affinity chromatographic purification of D-mannose-N-acetyl-D-glucosamine-specific lectin from the bark of *Sophora japonica* eliminating the loss by sugar specific self-aggregation 597(1992)197
- Evaluation of the mode of binding of immunoglobulin to activated agarose 597(1992)231
- Use of antibody fragments in immunoaffinity chromatography. Comparison of FV fragments, VH fragments and paralog peptides 597(1992)239
- Artificial immunoglobulin G-binding protein mimetic to staphylococcal protein A. Its production and application to affinity purification of immunoglobulin G 597(1992)277
- Immunoenzymatic assay of dyes currently used in affinity chromatography for protein purification 597(1992)315
- Affinity chromatography in purification of A₁ adenosine receptors 597(1992)335
- Specific assay for endotoxin using immobilized histidine, *Limulus* amoebocyte lysate and a chromogenic substrate 597(1992)415
- Purification and reconstitution of μ -opioid receptors in liposome 597(1992)429
- Purification of recombinant human interferon- β by immobilized antisense peptides 600(1992)235
- Affinity chromatography**
see also Special issue index
- Affinity electrophoresis**
Analysis of the interaction between an $\alpha(1\rightarrow6)$ dextran-specific mouse hybridoma antibody and dextran B512 by affinity electrophoresis 597(1992)345
- Characterization of the interaction between human plasma fibronectin and collagen by means of affinity electrophoresis 597(1992)351
- Immobilized metal ion affinity electrophoresis. A study with several model proteins containing histidine 597(1992)357
- Separation of monoclonal antibodies from antihapten antisera by two-dimensional affinity electrophoresis 597(1992)365
- Affinity gels**
Avid AL, a synthetic ligand affinity gel mimicking immobilized bacterial antibody receptor for purification of immunoglobulin G 597(1992)101
- Affinity labelling**
Discriminative affinity labelling of opioid receptors by enkephalin and morphiceptin analogues containing 3-nitro-2-pyridinesulphenyl-activated thiol residues 597(1992)425
- Affinity ligands**
Computer simulation of weak affinity chromatography 597(1992)83
- Affinity membranes**
Surface-modified membranes as a matrix for protein purification 591(1992)107
- Affinity partitioning**
Affinity partitioning of erythrocytic phosphofructokinase in aqueous two-phase systems containing poly(ethylene glycol)-bound Cibacron Blue. Influence of pH, ionic strength and substrates/ effectors 589(1992)127
- Ligand-receptor interactions in affinity cell partitioning. Studies with transferrin covalently linked to monomethoxypoly(ethylene glycol) and rat reticulocytes 594(1992)97
- Affinity purification**
Affinity purification of antibodies using immobilized FB domain of protein A 597(1992)257
- Affinity shift, calcium-dependent**
Purification of sufficiently γ -carboxylated recombinant protein C and its derivatives. Calcium-dependent affinity shift in immunoaffinity and ion-exchange chromatography 597(1992)285

Affinity supports

Study of affinity supports based on reactive polymers immobilized on silica: affinity constant determination from isocratic zonal elution 584(1992)3

Affinity tag procedures

Arginine-tail method, an affinity tag procedure utilizing anhydrotrypsin agarose 597(1992)181

Agarose

Diffusion of proteins in Sepharose Cl-B gels 591(1992)115

Evaluation of the mode of binding of immunoglobulin to activated agarose 597(1992)231

Agglutinins

Immobilized lectin columns: useful tools for the fractionation and structural analysis of oligosaccharides (Review) 597(1992)111

Alcohol dehydrogenase-silica

Dehydrogenase-silica as a stationary phase for the separation of alcohols and ketones 587(1991)268

Alkaline fusion

Degradation of polysulphones by alkali fusion 595(1992)283

Reactor for prechromatographic fusion reactions 600(1992)352

Alkylsilylated silica

Suppression of the effect of metal impurities in alkylsilylated silica packing materials 594(1992)87

Allosteric transition

Occurrence of an allosteric transition in the modification of papain with L-1-acetyl-2,3-dihydropyrrolo[2,3-*b*]-indole-2-carboxamide 597(1992)411

Alternating current plasma emission detector

Determination of organotin species by capillary gas chromatography with alternating current plasma emission detection 598(1992)97

Amanita phalloides

Simultaneous assay for amatoxins and phallotoxins in *Amanita phalloides* Fr. by high-performance liquid chromatography 598(1992)227

Amino groups

Chromatographic effects of residual amino groups on liquid chromatographic stationary phases derived from aminopropyl-silica gel 590(1992)189

Aminopropyl-silica gel

Chromatographic effects of residual amino groups on liquid chromatographic stationary phases derived from aminopropyl-silica gel 590(1992)189

Ammonium molybdophosphate

New solid adsorbents for the separation of lower hydrocarbons and permanent gases. II. Ammonium molybdophosphate 556(1991)425

Amphiphilic copolymer resins

Beaded polymer supports and gels (Review). I. Manufacturing techniques 586(1991)181

Amplifier

High-speed gas chromatography. Theoretical and practical aspects 556(1991)331

Amylose

Fast separation of macrophages by retention on cross-linked amylose and release by enzymatic amyolysis of the chromatographic material 584(1992)121

Analyte focusing

Analyte focusing in capillary electrophoresis using on-line isotachopheresis 591(1992)341

Anhydrotrypsin agarose

Arginine-tail method, an affinity tag procedure utilizing anhydrotrypsin agarose 597(1992)181

Anion exchangers

see also Ion exchangers

Anion-exchange chromatography of DNA restriction fragments 555(1991)109

High-performance liquid chromatography of amino acids, peptides and proteins. CXIV. Protein interactions with porous coulombic sorbents: comparison of experimental findings with predictions of several adsorption models 557(1991)335

Determination of aromatic sulfonic acids in aqueous environmental samples by anion-exchange chromatography coupled to particle beam mass spectrometry and UV spectrophotometry 589(1991)177

Application of a pellicular anion-exchange resin to the separation of inorganic and organic anions by single-column anion exchange 592(1992)101

Antarctic rocks

Novel long-chain *anteiso*-alkanes and *anteiso*-alkanoic acids in Antarctic rocks colonized by living and fossil cryptoendolithic microorganisms 598(1992)267

Antigen-antibody interaction

Kinetic aspects of membrane-based immunoaffinity chromatography 597(1992)167

Antipruritic agents

Supercritical fluid chromatography in the routine stability control of antipruritic preparations 553(1991)71

Antisense peptide recognition

Interactions and uses of antisense peptides in affinity technology (Review) 597(1992)29

Apple juice

High-performance liquid chromatography of the phenolic compounds in the juice of some French cider apple varieties 555(1991)125

Applied voltage

Optimization of resolution in capillary zone electrophoresis: combined effect of applied voltage and buffer concentration 588(1991)315

Aqueous two-phase systems

Extraction of xanthine oxidase from milk by counter-current distribution in an aqueous two-phase system 558(1991)357

Arginine-tail method

Arginine-tail method, an affinity tag procedure utilizing anhydrotropin agarose 597(1992)181

Artemisia absinthium

Use of high-performance liquid chromatographic peak deconvolution and peak labelling to identify antiparasitic components in plant extracts 593(1992)209

Aspergillus oryzae

Detection of enzyme activity in fractions collected from free solution capillary electrophoresis of complex samples 559(1991)325

Association constants

Determination of the association constant of monovalent mode protein-sugar interaction by capillary zone electrophoresis 597(1992)377

Asymmetric channels

Asymmetric-channel flow field-flow fractionation with exponential force-field programming 593(1992)339

Atmospheric pressure chemical ionization

Screening of organophosphorus pesticides using liquid chromatography-atmospheric pressure chemical ionization mass spectrometry 595(1992)193

Atmospheric pressure ionization

Capillary electrophoresis-atmospheric pressure ionization mass spectrometry for the characterization of peptides. Instrumental considerations for mass spectrometric detection 554(1991)311

Determination of small drug molecules by capillary electrophoresis-atmospheric pressure ionization mass spectrometry 559(1991)515

Atomic emission spectroscopy

Measurements using gas chromatography with coelution and dual-isotope atomic emission detection 586(1991)309

Comparison of commercially available atomic emission and chemiluminescence detectors for sulfur-selective gas chromatographic detection 591(1992)313

Gas chromatography of chloride and bromide as phenylboronic acid/mercuric nitrate derivatives with microwave induced plasma atomic emission detection 594(1992)395

Autofocusing

Isolation and purification of bacterial proteinases by means of autofocusing 585(1991)373

Automated sample clean-up

Development of a rational optimisation procedure for the automated sample clean-up with column switching in pesticide residue analysis 552(1991)113

Automation

Near-continuous measurement of hydrogen sulfide and carbonyl sulfide by an automatic gas chromatograph 585(1991)353

Development and optimization of a single-step procedure using protein A affinity chromatography to isolate murine IgG₁ monoclonal antibodies from hybridoma supernatants 587(1991)61

Automated two-dimensional liquid chromatographic system for mapping proteins in highly complex mixtures 588(1991)115

Automated high-performance liquid chromatographic and size-exclusion chromatographic sample preparation by means of a robotic workstation 592(1992)249

Automated determination of aflatoxin B₁ in cattle feed by two-column solid-phase extraction with on-line high-performance liquid chromatography 592(1992)265

Automated high-performance liquid chromatographic method for the determination of rifampicin in plasma 593(1992)21

Determination of catecholamines in urine by liquid chromatography and electrochemical detection after on-line sample purification on immobilized boronic acid 593(1992)185

Comparison of several methods for the determination of trace amounts of polar aliphatic monocarboxylic acids by high-performance liquid chromatography 593(1992)243

Screening for amines by dansylation and automated high-performance liquid chromatography 598(1992)51

Coupled-column high-performance liquid chromatographic method for the determination of 1-hydroxypyrene in urine of subjects exposed to polycyclic aromatic hydrocarbons 600(1992)189

Application of an internal surface reversed-phase column for the automated determination of flucycloxon residues 600(1992)211

Avid AL

Avid AL, a synthetic ligand affinity gel mimicking immobilized bacterial antibody receptor for purification of immunoglobulin G 597(1992)101

Avidin-biotin technique

Non-isotopic receptor assay for benzodiazepines using a biotin-labeled ligand and biotin-immobilized microtiter plate 597(1992)443

Avidin column

Avidin protein-conjugated column for direct injection analysis of drug enantiomers in plasma by high-performance liquid chromatography 572(1991)133

Axial diffusion

Finite difference modelling of continuous-flow electrophoresis 594(1992)341

Axial dispersion

Theory of perfusion chromatography 599(1992)87

Azoxydiphenetole

Comparison of four liquid crystal stationary phases used above and below their melting point temperatures for the gas chromatography of some volatile oil constituents 588(1991)231

B***Bacillus stearothermophilus***

Purification of serine hydroxymethyltransferase from *Bacillus stearothermophilus* with ion-exchange high-performance liquid chromatography 596(1992)203

Bacillus subtilis

Purification and characterization of recombinant pyrrolidone carboxyl peptidase of *Bacillus subtilis* 584(1992)101

Balance, electronic

Improved accuracy in the determination of field-flow fractionation elution volumes 555(1991)260

Band ...

see also Peak ...

Band broadening

Source of band broadening in liquid chromatographic-fast atom bombardment mass spectrometric systems with precolumn addition of viscous matrix to the mobile phase 554(1991)47

Improved computer algorithm for characterizing skewed chromatographic band broadening. I. Method 556(1991)111

High-performance liquid chromatography of amino acids, peptides and proteins. CXX. Evaluation of bandwidth behaviour of proteins chromatographed on tentacle-type anion exchangers 599(1992)163

Band profiles

see also Peak shape

Experimental study on the effect of the sample size on the band profile for a binary mixture showing no competitive interaction 556(1991)195

Prediction of single and binary profiles in overloaded elution chromatography using various semi-ideal models 556(1991)205

Effect of the sample solvent on band profiles in preparative liquid chromatography using non-aqueous reversed-phase high-performance liquid chromatography 588(1991)1

Prediction of the elution profiles of high-concentration bands in gas chromatography 589(1992)223

Experimental study of the production rate of pure enantiomers from racemic mixtures 590(1992)119

Optimization of the experimental conditions and the column design parameters in overloaded elution chromatography 591(1992)31

Validity of the model used to relate the energy distribution and the adsorption isotherm 591(1992)267

Band splitting

Effect of the sample solvent on band profiles in preparative liquid chromatography using non-aqueous reversed-phase high-performance liquid chromatography 588(1991)1

Bandwidth

High-performance liquid chromatography of amino acids, peptides and proteins. CXX. Evaluation of bandwidth behaviour of proteins chromatographed on tentacle-type anion exchangers 599(1992)163

Base-atom recognition

Base-atom recognition in protein adsorption to alkyl agaroses 597(1992)93

Base pairing

High-performance affinity chromatography of DNA. II. Porosity effects 600(1992)221

Basic alumina

Use of basic alumina in fractionation of fossil fuels 600(1992)243

Batch separation

Immunomagnetic separation of tumor necrosis factor α . I. Batch procedure for human temporomandibular fluid 583(1992)11

Bauhinia purpurea

Determination of the carbohydrate-binding site of *Bauhinia purpurea* lectin by affinity chromatography 597(1992)221

Beaded polymer supports

Beaded polymer supports and gels (Review). I. Manufacturing techniques 586(1991)181

Beaded polymer supports and gels (Review). II. Physico-chemical criteria and functionalization 586(1991)199

Bead size

Beaded polymer supports and gels (Review). II. Physico-chemical criteria and functionalization 586(1991)199

Bidirectional column test

Simple test for determination of the degree of distortion of the liquid-phase flow profile in columns for preparative liquid chromatography 589(1992)61

Binary buffers

Binary buffers for indirect absorption detection in capillary zone electrophoresis 589(1992)307

Binary solvents

Comparison of selectivity of silica and Florisil in the separation of natural pyranocoumarins 552(1991)453

Binding constants

Study of high-affinity interactions by quantitative affinity chromatography. Analytical expressions in terms of total ligand concentration 597(1992)57

Binding site analysis

Biospecific interaction analysis using surface plasmon resonance detection applied to kinetic, binding site and concentration analysis 597(1992)397

Biological recognition

see also Social issue index

Biosensors

Mapping of viral epitopes with conformationally specific monoclonal antibodies using biosensor technology 597(1992)391

Biospecific interaction analysis using surface plasmon resonance detection applied to kinetic, binding site and concentration analysis 597(1992)397

Biospecific adsorbents

Galactosylation as a tool for the stabilization and immobilization of proteins (Review) 597(1992)19

Biospecific interaction analysis

Biospecific interaction analysis using surface plasmon resonance detection applied to kinetic, binding site and concentration analysis 597(1992)397

Biospecific ligands

Protein adsorption capacity of a porous phenylalanine-containing membrane based on a polyethylene matrix 586(1991)27

Biospecific recognition

Computer simulation of weak affinity chromatography 597(1992)83

Biotinylation

Affinity chromatographic purification of bovine lung endothelin receptor using biotinylated endothelin and avidin-agarose 597(1992)331

Biotransformation

Assessment of the biotransformation of the cardiotoxic agent piroximone by high-performance liquid chromatography and gas chromatography-mass spectrometry 593(1992)1

Bis(methoxybenzylideneanilbitoluidine)

Comparison of four liquid crystal stationary phases used above and below their melting point temperatures for the gas chromatography of some volatile oil constituents 588(1991)231

Bis(methoxybenzylideneanilchloroaniline)

Comparison of four liquid crystal stationary phases used above and below their melting point temperatures for the gas chromatography of some volatile oil constituents 588(1991)231

Blue cotton

Cellulose bearing covalently linked copper phthalocyanine trisulphonate as an adsorbent selective for polycyclic compounds and its use in studies of environmental mutagens and carcinogens (Review) 597(1992)37

Boiling point

Simulated distillation of distillates on capillary columns: influence of the polarity of the stationary phase 557(1991)155

Book reviews

Ion exchange in analytical chemistry (by H. E. Walton and R. D. Rocklin) 555(1991)329

Chromatography and modification of nucleosides, Part A (edited by C. W. Gehrke and K. C. T. Kuo) 563(1991)228

CRC handbook of chromatography: drugs, Vols. III-VI (edited by R.N. Gupta) 564(1991)355

Advances in electrophoresis, Vol. 3 (edited by A. Chrambach, M. J. Dunn and B. J. Radola) 565(1991)533

Analysis for drugs and metabolites including anti-infective agents (edited by E. Reid and I. D. Wilson) 565(1991)534

The use of HPLC in receptor biochemistry (edited by A. R. Kerlavage) 566(1991)266

Electroanalysis of biologically important compounds (by J. P. Hart) 568(1991)270

- High-performance liquid chromatography in biotechnology (edited by W. S. Hancock) 568(1991)501
- Liquid chromatography in biomedical analysis (edited by T. Hanai) 571(1991)334
- HPLC in clinical chemistry (by I. N. Papadoyannis) 574(1992)179
- Practical manual on lipid analysis. A series of monographs. I. Fatty acids (by J. G. Alvarez and J. C. Touchstone) 575(1992)182
- Recent developments in therapeutic drug monitoring and clinical toxicology (edited by I. Sunshine) 584(1992)280
- Analytical biotechnology: Capillary electrophoresis and chromatography (edited by Cs. Horváth and J. N. Nikely) 585(1991)378
- A practical guide to the care, maintenance and troubleshooting of capillary gas chromatographic systems (by D. Rood) 585(1991)379
- Unified separation science (by J. C. Giddings) 585(1991)380
- Quantitative chemical analysis (by D. C. Harris) 587(1991)368
- Non-chromatographic continuous separation techniques (by M. Valcárcel and M. D. Luque de Castro) 587(1991)369
- Handbook of thin-layer chromatography (edited by J. Sherma and B. Fried) 588(1991)365
- Luminescence techniques in chemical and biochemical analysis (edited by W. R. G. Bayens, P. De Keukeleire and K. Korkides) 588(1991)367
- Fluorometric analysis in biomedical chemistry; trends and techniques including HPLC applications (edited by N. Ichinose, G. Schwedt, F. M. Schnepel and K. Adachi) 590(1992)373
- Advances in steroid analysis (edited by S. Görög) 594(1992)403
- Chiral separations by liquid chromatography (edited by S. Ahuja) 595(1992)375
- Eucalyptus oils: use, chemistry, distillation and marketing (edited by D. J. Boland, J. J. Brophy and A. P. N. House) 598(1992)316
- Chromatography and isolation of insect hormones and pheromones (edited by A. R. McCaffery and I. D. Wilson) 600(1992)373
- Advances in electrophoresis, Vol. 4 (edited by A. Chrambach, M. J. Dunn and B. J. Radola) 600(1992)375
- Advances in chromatography, Vol. 32 (edited by J. C. Giddings, E. Grushka and P. R. Brown) 600(1992)376
- Stationary phases in gas chromatography (by H. Rotzsche) 600(1992)378
- Boundary effects**
Theory of linear non-equilibrium chromatography with beds of a finite length 591(1992)273
- Bovine serum albumin**
Study of protein-ligand binding effects by direct chromatographic on-column injection 584(1992)11
- Breakthrough curves**
Adsorption and elution of bovine γ -globulin using an affinity membrane containing hydrophobic amino acids as ligands 585(1991)45
- Adsorption kinetics of albumin on a cross-linked cellulose chromatographic ion exchanger 590(1992)101
- Short-cut method for predicting the productivity of affinity chromatography 597(1992)173
- Theory of perfusion chromatography 599(1992)87
- Brown coals**
Quantitative structure-type analysis of heterocompound-hydrocarbon mixtures. VI. Characterization of the nature of neutral heterocompounds in a product of brown coal refining 553(1991)441
- Brush type stationary phases**
New chiral covalently bonded, π -donor stationary phases for high-performance liquid chromatography, based on derivatives of optically active 1-(1-naphthyl)ethylamine. Part II 585(1991)195
- Bubblers**
Determination of hexahydrophthalic anhydride in air using gas chromatography 558(1991)247
- Buffer concentration**
Optimization of resolution in capillary zone electrophoresis: combined effect of applied voltage and buffer concentration 588(1991)315
- Buffers**
Chromatographic separation of mixed peptides from amino acids in biological digests with volatile buffers 586(1991)259
- Binary buffers for indirect absorption detection in capillary zone electrophoresis 589(1992)307
- Butteroil**
Reversed-phase liquid chromatography-mass spectrometry of complex mixtures of natural triacylglycerols with chloride-attachment negative chemical ionization 588(1991)73

C

Calcination

Effect of silanol groups on heat-treated silicas by calcination and retention behaviour in high-performance liquid chromatography 556(1991)407

Capacity factors

High-performance liquid chromatography of amino acids and dipeptides on new ion exchangers of the HEMA series 552(1991)439

High-performance liquid chromatographic study of some biologically active analogues of arabinosylcytosine 588(1991)147

Chromatographic behaviour of the *cis*-[B₁₂H₁₂]²⁻ derivatives on hydroxyethylmethacrylate gels 588(1991)201

Capillary action

Studies of the electrophoretic behaviour of *p*-dihydroxyborylphenylalanine and related compounds by the three-spot method 585(1991)369

Capillary columns

Recent advances in liquid chromatography–mass spectrometry and capillary zone electrophoresis–mass spectrometry for protein analysis 553(1991)101

Application of microcolumn liquid chromatography–continuous-flow fast atom bombardment mass spectrometry in environmental studies of sulfonylurea herbicides 554(1991)91

Activity testing and surface characterization of pretreated fused-silica capillaries for gas chromatography. A new modification of existing intermediate column tests 590(1992)271

Experimental determination of adsorption isotherm data for the study of the surface energy distribution of various solid surfaces by inverse gas–solid chromatography 591(1992)233

High-temperature open-tubular capillary column liquid chromatography 592(1992)239

Charge-reversed, polymer-coated capillary column for the analysis of a recombinant chimeric glycoprotein 594(1992)317

Capillary modification and evaluation using streaming potential and frontal chromatography for protein analysis in capillary electrophoresis 594(1992)325

Critical capillary column examination of the relationship between separation number and height equivalent to a theoretical plate and their dependence on temperature 595(1992)209

Gas chromatographic analysis of high-molecular-weight polynuclear aromatic hydrocarbons. I. Molecular weight 328 595(1992)247

Reversed- and normal-phase separations by high-temperature open-tubular column liquid chromatography 598(1992)153

Application of capillary reversed-phase high-performance liquid chromatography to high-sensitivity protein sequence analysis 599(1992)119

Capillary columns, packed

Effects of modifiers in packed and open-tubular supercritical fluid chromatography 552(1991)527

Capillary electrophoresis

see also Special issue index

Capillary electrophoresis

see also Micellar electrokinetic chromatography

Recent advances in liquid chromatography–mass spectrometry and capillary zone electrophoresis–mass spectrometry for protein analysis 553(1991)101

Nanoscale separations combined with tandem mass spectrometry 554(1991)73

Capillary electrophoresis–atmospheric pressure ionization mass spectrometry for the characterization of peptides. Instrumental considerations for mass spectrometric detection 554(1991)311

Nanoscale separations. Capillary liquid chromatography–mass spectrometry and capillary zone electrophoresis–mass spectrometry for the determination of peptides and proteins 554(1991)329

Bile salt surfactants in micellar electrokinetic capillary chromatography. Application to hydrophobic molecule separations 557(1991)113

Electroosmotically driven electrochromatography of anions having similar electrophoretic mobilities by ion pairing 557(1991)125

Application of capillary zone electrophoresis to the characterization of multiple antigen peptides 557(1991)307

Comparison of high-performance liquid chromatography with capillary gel electrophoresis in single-base resolution of polynucleotides 558(1991)273

Synthesis of a new acrylamido buffer (acryloylhistamine) for isoelectric focusing in immobilized pH gradients and its analysis by capillary zone electrophoresis 558(1991)285

Mathematical model describing dispersion in free solution capillary electrophoresis under stacking conditions 559(1991)3

- Temperature elevations of the sample zone in free solution capillary electrophoresis under stacking conditions 559(1991)27
- Factors influencing the resolution and quantitation of oligonucleotides separated by capillary electrophoresis on a gel-filled capillary 559(1991)43
- Influence of operating parameters on reproducibility in capillary electrophoresis 559(1991)57
- Evaluation of fundamental properties of a silica capillary used for capillary electrophoresis 559(1991)69
- Effect of hydrostatic flow on the efficiency in capillary electrophoresis 559(1991)81
- Field effect electroosmosis 559(1991)95
- Split injector for capillary zone electrophoresis 559(1991)103
- Moving boundary capillary electrophoresis with concentration gradient detection 559(1991)111
- Capillary zone electrophoresis analysis of acrylamido buffers for isoelectric focusing in immobilized pH gradients 559(1991)119
- Analysis of separation efficiency in capillary electrophoresis with direct control of electroosmosis by using an external electric field 559(1991)133
- Field amplified sample injection in high-performance capillary electrophoresis 559(1991)141
- Field-amplified polarity-switching sample injection in high-performance capillary electrophoresis 559(1991)153
- Optimization and evaluation of the performance of arrangements for UV detection in high-resolution separations using fused-silica capillaries 559(1991)163
- Instrumentation for high-performance capillary electrophoresis-mass spectrometry 559(1991)197
- Chiral separations by micellar electrokinetic chromatography with sodium N-dodecanoyl-L-valinate 559(1991)209
- Selected applications of cyclodextrin selectors in capillary electrophoresis 559(1991)215
- Separation of fluorescent oligosaccharide derivatives by microcolumn techniques based on electrophoresis and liquid chromatography 559(1991)223
- Low-cost, high-sensitivity laser-induced fluorescence detection for DNA sequencing by capillary gel electrophoresis 559(1991)237
- Reproducibility and quantitation of separation for ribonucleoside triphosphates and deoxyribonucleoside triphosphates by capillary zone electrophoresis 559(1991)247
- Capillary electrophoresis of DNA in entangled polymer solutions 559(1991)257
- Analysis of DNA restriction fragments and polymerase chain reaction products towards detection of the AIDS (HIV-1) virus in blood 559(1991)267
- Effect of temperature on the separation of DNA restriction fragments in capillary gel electrophoresis 559(1991)285
- Identification of DNA molecules by pre-column hybridization using capillary electrophoresis 559(1991)295
- Assessment of capillary electrophoresis in pharmaceutical applications. Analysis and quantification of a recombinant cytokine in an injectable dosage form 559(1991)307
- Free solution capillary electrophoresis of proteins using untreated fused-silica capillaries 559(1991)317
- Detection of enzyme activity in fractions collected from free solution capillary electrophoresis of complex samples 559(1991)325
- Analysis of a vaccine purification process by capillary electrophoresis 559(1991)337
- Application of high-performance capillary electrophoresis to the analysis of conformation and interaction of metal-binding proteins 559(1991)345
- Electrophoretic mobility modeling of proteins in free zone capillary electrophoresis and its application to monoclonal antibody microheterogeneity analysis 559(1991)357
- Capillary zone electrophoresis of proteins with hydrophilic fused-silica capillaries 559(1991)367
- High-performance capillary electrophoresis as a fast in-process control method for enzyme-labelled monoclonal antibody conjugates 559(1991)385
- Capillary zone electrophoresis studies of motilin peptides. Effects of charge, hydrophobicity, secondary structure and length 559(1991)391
- Fractionation of the human recombinant tissue plasminogen activator (rtPA) glycoforms by high-performance capillary zone electrophoresis and capillary isoelectric focusing 559(1991)401
- High-performance capillary electrophoresis of proteins from the fluid lining of the lungs of rats exposed to perfluoroisobutylene 559(1991)411

- High-performance capillary electrophoresis of histones 559(1991)431
- Rapid protein analysis by capillary electrophoresis 559(1991)445
- Capillary electrophoresis for diagnosis and studies of human disease, particularly metabolic disorders 559(1991)455
- Characterization of polyethylene glycol modified proteins using charge-reversed capillary electrophoresis 559(1991)467
- Determination of thiopental in human serum and plasma by high-performance capillary electrophoresis–micellar electrokinetic chromatography 559(1991)505
- Determination of small drug molecules by capillary electrophoresis–atmospheric pressure ionization mass spectrometry 559(1991)515
- Migration behaviour of catechols and catecholamines in capillary electrophoresis 559(1991)529
- Separation of Dns-amino acids and vitamins by micellar electrokinetic chromatography 559(1991)537
- Electrochromatographic solid-phase extraction for determination of cimetidine in serum by micellar electrokinetic capillary chromatography 559(1991)547
- Micellar electrokinetic capillary chromatographic separation and laser-induced fluorescence detection of 2'-deoxynucleoside 5'-monophosphates of normal and modified bases 565(1991)197
- Capillary zone electrophoresis: its applicability and potential in biochemical analysis (Review) 569(1991)63
- Separation of globins using free zone capillary electrophoresis 576(1992)346
- Separation of antibiotics by high-performance capillary electrophoresis with photodiode-array detection 585(1991)133
- Determination of cimetidine in pharmaceutical preparations by capillary zone electrophoresis 586(1991)177
- Optimization of resolution in capillary zone electrophoresis: combined effect of applied voltage and buffer concentration 588(1991)315
- Quantitative aspects of indirect UV detection in capillary zone electrophoresis 588(1991)321
- High-performance capillary zone electrophoresis of carbohydrates in the presence of alkaline earth metal ions 588(1991)327
- Determination of antihistamines in pharmaceuticals by capillary electrophoresis 588(1991)335
- Binary buffers for indirect absorption detection in capillary zone electrophoresis 589(1992)307
- Radial pH distribution during capillary electrophoresis with electroosmotic flow. Analysis with high ionic strength buffers 589(1992)315
- Capillary isoelectric focusing with electroosmotic zone displacement and on-column multichannel detection 589(1992)321
- Monitoring 4-methoxy-2,3,6-trimethylbenzenesulphonyl deprotection of arginine-containing synthetic peptides using capillary zone electrophoresis 589(1992)380
- Comparison of isotachopheresis, capillary zone electrophoresis and high-performance liquid chromatography for the determination of salbutamol, terbutaline sulphate and fenoterol hydrobromide in pharmaceutical dosage forms 590(1992)341
- Determination of cefixime and its metabolites by high-performance capillary electrophoresis 590(1992)364
- Comparison of liquid-junction and coaxial interfaces for capillary electrophoresis–mass spectrometry with application to compounds of concern to the aquaculture industry 591(1992)325
- Analyte focusing in capillary electrophoresis using on-line isotachopheresis 591(1992)341
- Planar chips technology for miniaturization and integration of separation techniques into monitoring systems. Capillary electrophoresis on a chip 593(1992)253
- Separation of leucinostatin by capillary zone electrophoresis 593(1992)259
- Separation of naphthalene-2,3-dicarboxaldehyde-labeled amino acids by high-performance capillary electrophoresis with laser-induced fluorescence detection 593(1992)265
- Factors affecting the separation of inorganic metal cations by capillary electrophoresis 593(1992)289
- Prediction of migration behavior of oligonucleotides in capillary gel electrophoresis 593(1991)297
- Capillary electrophoresis with electrochemical detection employing an on-column Nafion joint 593(1992)305
- Charge-reversed, polymer-coated capillary column for the analysis of a recombinant chimeric glycoprotein 594(1992)317
- Capillary modification and evaluation using streaming potential and frontal chromatography for protein analysis in capillary electrophoresis 594(1992)325

Separation of plant growth regulators by capillary electrophoresis 594(1992)335

Peak dispersion and separation efficiency in high-performance zone electrophoresis with gel-filled capillaries 595(1992)313

High-sensitivity laser-induced fluorescence detection of native proteins in capillary electrophoresis 595(1992)319

Determination of sulphonamides in pork meat extracts by capillary zone electrophoresis 596(1992)101

DNA electrophoresis in uncross-linked polyacrylamide solution, studied by epifluorescence microscopy 596(1992)141

Enzymophoresis of nucleic acids by tandem capillary enzyme reactor-capillary zone electrophoresis 596(1992)251

Determination of the association constant of monovalent mode protein-sugar interaction by capillary zone electrophoresis 597(1992)377

Assay of protein drug substances present in solution mixtures by fluorescamine derivatization and capillary electrophoresis 598(1992)123

Systematic optimization of capillary electrophoretic separation of sulphonamides 598(1992)133

Kinetics of acrylamide photopolymerization as investigated by capillary zone electrophoresis 598(1992)277

Polyacrylamide gel polymerization under non-oxidizing conditions, as monitored by capillary zone electrophoresis 598(1992)287

Characterization of the heterogeneity of polyethylene glycol-modified superoxide dismutase by chromatographic and electrophoretic techniques 599(1992)141

Synthesis and evaluation of epoxy polymer coatings for the analysis of proteins by capillary zone electrophoresis 599(1992)227

Capillary zone electrophoresis of linear and branched oligosaccharides 600(1992)279

Capillary gel electrophoresis of single-stranded DNA fragments with UV detection 600(1992)289

Determination of ephedrine alkaloids by capillary electrophoresis 600(1992)370

Carbon adsorbents

Chromatographic determination of the physico-chemical parameters of adsorption on activated carbon fibres 600(1992)67

Carbon isotope ratio

Enantiomeric distribution and $^{13}\text{C}/^{12}\text{C}$ isotope ratio determination of γ -lactones: appropriate methods for the differentiation between natural and non-natural flavours? 557(1991)187

Carbon molecular sieves

Dynamic gas-solid chromatographic techniques for characterizing carbon molecular sieves 556(1991)433

Carbowax 20M

High-performance liquid chromatography of benzodiazepines using sorbents with thermally immobilized Carbowax 20M 600(1992)109

Carrier effects

Quantitation and linearity for particle-beam liquid chromatography-mass spectrometry 554(1991)103

Carrier gases

Gas chromatographic separation of unsaturated polar compounds on inorganic stationary phases containing silver nitrate 552(1991)313

Catalysis

Liquid chromatographic isolation and structural elucidation of methoxymethylene dimethyl dodecatrienes catalytically synthesized from methanol and three moles of isoprene 594(1992)111

Cation-exchange behaviour

Cation-exchange liquid chromatography of choline and acetylcholine on free shielded silanols of silica-based reversed-phase stationary phases 596(1992)43

Cation exchangers

see also Ion exchangers

Standardization of cation-exchange clean-up prior to gas chromatography of amino acids 553(1991)87

Recovery of sugar derivatives from 2-aminopyridine labeling mixtures for high-performance liquid chromatography using UV or fluorescence detection 587(1991)177

Quantitative separation of Cr^{3+} from Mo^{6+} , W^{6+} , Hg_2^{2+} , Cu^{2+} and Pb^{2+} . Chromatographic behaviour of 51 cations on papers impregnated with Sn(IV)-based inorganic ion exchangers in complex-forming acid systems 594(1992)379

Effects of stationary phase ligand density on high-performance ion-exchange chromatography of proteins 598(1992)7

Cell cultures

High-performance gel permeation chromatographic analysis of immunoglobulin M produced by hybridoma cell culture 594(1992)179

Cell separations

Separation techniques for biotechnology in the 1990s 592(1992)3

Cellulase-silica

Separation of enantiomers using cellulase (CBH I) silica as a chiral stationary phase 586(1991)233

Cellulose, 1,1'-carbonyldiimidazole substituted

Efficient substitution of 1,1'-carbonyldiimidazole activated cellulose and Sepharose matrices with amino acyl spacer arms 587(1991)271

Cellulose, cross-linked

Adsorption kinetics of albumin on a cross-linked cellulose chromatographic ion exchanger 590(1992)101

Cellulose acetate fibres

Analysis of dyes extracted from textile fibers by thermospray high-performance liquid chromatography-mass spectrometry 586(1991)73

Cellulose beads

Cellulose beads: a weak leaking affinity support 589(1992)93

Cellulose-copper phthalocyanine trisulphonate

Cellulose bearing covalently linked copper phthalocyanine trisulphonate as an adsorbent selective for polycyclic compounds and its use in studies of environmental mutagens and carcinogens (Review) 597(1992)37

Cellulose tris(3,5-dimethylphenylcarbamate)

Direct high-performance liquid chromatographic separation of enantiomeric peptidoleukotriene antagonists 596(1992)123

¹³C enrichment

On-line measurements of ¹³C enrichments in rat breath. Non-invasive method for *in vivo* study of drug enzymatic induction 562(1991)341

Centella asiatica

Direct coupling of high-speed counter-current chromatography to thin-layer chromatography. Application to the separation of asiaticoside and madecassoside from *Centella asiatica* 558(1991)446

Centrifugal partition chromatography

Measurement of partition coefficients by various centrifugal partition chromatographic techniques. A comparative evaluation 556(1991)181

Ceramic particles

Experimental determination of adsorption isotherm data for the study of the surface energy distribution of various solid surfaces by inverse gas-solid chromatography 591(1992)233

Cereals

Determination of α - and β -zearealenol and zearealenone in cereals by gas chromatography with ion-trap detection 595(1992)259

Chain length distribution

Estimation of the distributions of chain length of amylopectins by high-performance liquid chromatography with pulsed amperometric detection 585(1991)233

Channel flow

Non-equilibrium integral Doppler anemometric analysis of particle mixtures in a channel flow using an intrinsic hydrodynamic focusing force biased by another force 553(1991)517

Charge discrimination enhanced injection

Field-amplified polarity-switching sample injection in high-performance capillary electrophoresis 559(1991)153

Charge-reversed capillaries

Characterization of polyethylene glycol modified proteins using charge-reversed capillary electrophoresis 559(1991)467

Cheese

Improvement of extraction procedure for biogenic amines in foods and their high-performance liquid chromatographic determination 591(1992)175

Chelate ring size

Simultaneous effect of geometrical isomerism and chelate ring size of tris(aminocarboxylato)cobalt(III) complexes on their behaviour in thin-layer chromatography on silica gel and alumina 585(1991)181

Chelates

Thermoanalytical and chromatographic studies of copper(II), nickel(II) and oxovanadium(IV) complexes of tetradentate Schiff bases derived from β -diketones and 2,3-diaminopentane 558(1991)187

Chelating ligands

Effect of the substituents of β -diketonato ligands on R_f values of tris(chelate) transition metal complexes obtained by normal- and reversed-phase thin-layer chromatography on unmodified silica gel 585(1991)359

Separon HEMA modified for immobilized metal ion affinity chromatographic separation of proteins 598(1992)15

Chelating silicas

Determination of divalent trace metals in natural waters by preconcentration on N,N,N',N'-tetra(2-aminoethyl)ethylenediamine-silica followed by on-line ion chromatography 600(1992)203

Chelating Superose

High-performance metal chelate affinity chromatography of cytochromes P-450 using Chelating Superose 587(1991)43

Chemical ionization mass fragmentography

Simultaneous determination of sorbic acid, dehydroacetic acid and benzoic acid by gas chromatography-mass spectrometry 594(1992)253

Chemical reactors

Reaction kinetics and kinetic processes in modern liquid chromatographic reactors (Review) 589(1992)1

Chemical shift

Correlations between high-performance liquid chromatographic retention, X-ray structural and ^{13}C NMR spectroscopic data of flavonoid compounds 592(1992)65

Chemical waste

Gas chromatographic separation and automatic identification of complex mixtures of organic solvents in industrial wastes 585(1991)273

Chemiluminescence detector

Time-resolved luminescence detection of derivatized thiols in column liquid chromatography 552(1991)625

Sensitive liquid chromatographic determination of alkyl-, nitro- and chlorophenols by precolumn derivatization with dansyl chloride, postcolumn photolysis and peroxyoxalate chemiluminescence detection 553(1991)345

On-line electrochemical reagent generation for liquid chromatography with luminol-based chemiluminescence detection 557(1991)13

Determination of sulfur components in light petroleum streams by high-resolution gas chromatography with chemiluminescence detection 589(1992)271

Nitrogen-specific liquid chromatography detector based on chemiluminescence. Application to the analysis of ammonium nitrogen in waste water 592(1992)209

Primary and secondary amine derivatization with luminarins 1 and 2: separation by liquid chromatography with peroxyoxalate chemiluminescence detection 594(1992)386

4,5-Diaminophthalhydrazide as a highly sensitive chemiluminescence derivatization reagent for α -dicarbonyl compounds in high-performance liquid chromatography 598(1992)203

Chemometrics

High-performance liquid chromatography and supercritical fluid chromatography of monosaccharides and polyols using light-scattering detection. Chemometric studies of the retentions 590(1992)203

Chinook salmon

High-performance liquid chromatographic analysis of oxytetracycline in chinook salmon following administration of medicated feed 588(1991)181

Chiral additives

Practical strategy for the analytical separation of enantiomers by high-performance liquid chromatography 553(1991)357

Chiral centres

High-performance ligand-exchange chromatography of some amino acids containing two chiral centres 592(1992)345

Chirality

Chiroptical detection during liquid chromatography. III. Non-stop acquisition of circular dichroism spectra during liquid chromatography 586(1991)249

Chiral mobile phase additive

(2*R*,3*R*)-Dicyclohexyl tartrate as a chiral mobile phase additive 585(1991)35

Chiral reagents

High-performance liquid chromatographic separation of DL-amino acids derivatized with chiral variants of Sanger's reagent 555(1991)81

Chiral recognition

Chiral separation by cyclodextrin-modified micellar electrokinetic chromatography 553(1991)503

Unusual effect of temperature on the retention of enantiomers on a chiral column 558(1991)1

Effect of superfluous remote polar functionality on chiral recognition 589(1992)45

Application and chiral recognition of heptakis (2,6-di-O-methyl-3-O-trifluoroacetyl)- β -cyclodextrin as a stationary phase for the gas chromatographic separation of enantiomers 589(1992)257

Retention and enantioselectivity of racemic solutes on a modified ovomucoid-bonded column. I. Cross-linking with glutaraldehyde 592(1992)301

Direct high-performance liquid chromatographic resolution of a novel benzothiazine Ca^{2+} antagonist and related compounds 593(1992)37

Synthesis of (1*R*-*trans*)-N,N'-1,2-cyclohexylenebisbenzamideo-ligodimethylsiloxane copolymers for use as chiral stationary phases for capillary supercritical fluid chromatography 594(1992)283

Contribution of specifiable hydrophobic interactions to chiral recognition 598(1992)1

Separation of the enantiomers of the 3,5-dinitrobenzamide derivatives of α -amino phosphonates on four chiral stationary phases 598(1992)159

Chiral selectors

Practical strategy for the analytical separation of enantiomers by high-performance liquid chromatography 553(1991)357

(*S,S*)-Diphenylethylethanediamine derivatives as chiral selectors. II. Gasparri-type bound chiral stationary phase with high enantioselectivity for naphthylamides 553(1991)373

- Problems and solutions in chiral thin-layer chromatography: a two-phase "Pirkle" modified amino-bonded plate 553(1991)497
- (2*R*,3*R*)-Dicyclohexyl tartrate as a chiral mobile phase additive 585(1991)35
- New chiral covalently bonded, π -donor stationary phases for high-performance liquid chromatography, based on derivatives of optically active 1-(1-naphthyl)ethylamine. Part II 585(1991)195
- Direct separation of nadolol enantiomers on a Pirkle-type chiral stationary phase 586(1991)351
- Silica-bonded chiral stationary phases with structurally simple π -donor chiral selectors for high-performance liquid chromatography 589(1992)53
- Investigations of the influence of silanol groups on the separation of enantiomers by liquid and supercritical fluid chromatography 592(1992)309
- (2*S*,3*S*)-Dicyclohexyl tartrate as mobile phase additive for the determination of the enantiomeric purity of (*S*)-atropine in tablets 592(1992)339
- Chiral sites**
Reversed retention order and other stereoselective effects in the separation of amino alcohols on Chiralcel OD 592(1992)331
- Chiral stationary phases, GC**
Investigation of cross-linking chiral stationary phases within capillary columns 552(1991)153
- Chiral π -donor stationary phases with (*R*)-*N*-pivaloylnaphthylethylamide groups for direct enantiomer separation by gas, liquid and supercritical fluid chromatography 557(1991)163
- Stereoisomeric purity determination of captopril by capillary gas chromatography 558(1991)257
- Application and chiral recognition of heptakis (2,6-di-*O*-methyl-3-*O*-trifluoroacetyl)- β -cyclodextrin as a stationary phase for the gas chromatographic separation of enantiomers 589(1992)257
- Chiral stationary phases, LC**
Practical strategy for the analytical separation of enantiomers by high-performance liquid chromatography 553(1991)357
- (*S*,*S*)-Diphenylethylethanediamine derivatives as chiral selectors. II. Gasparrini-type bound chiral stationary phase with high enantioselectivity for naphthylamides 553(1991)373
- Liquid chromatographic separation of the enantiomers of metoprolol and its α -hydroxy metabolite on Chiralcel OD for determination in plasma and urine 553(1991)391
- Retention behavior and chiral recognition of β -cyclodextrin-derivative polymer adsorbed on silica for warfarin, structurally related compounds and Dns-amino acids 555(1991)53
- Preparation of a chiral matrix. Resolution of (*6R,S*)-*N*⁵-formyltetrahydrofolate 555(1991)254
- Chiral π -donor stationary phases with (*R*)-*N*-pivaloylnaphthylethylamide groups for direct enantiomer separation by gas, liquid and supercritical fluid chromatography 557(1991)163
- Chiral stationary phase designed for β -blockers 557(1991)173
- Unusual effect of temperature on the retention of enantiomers on a chiral column 558(1991)1
- Preparative chromatographic separation of enantiomers (Review) 576(1992)1
- New chiral covalently bonded, π -donor stationary phases for high-performance liquid chromatography, based on derivatives of optically active 1-(1-naphthyl)ethylamine. Part II 585(1991)195
- Separation of enantiomers using cellulase (CBH I) silica as a chiral stationary phase 586(1991)233
- Preparative separation and analysis of the enantiomers of [³H]Abbott-69992, an HIV anti-infective nucleoside, by ligand-exchange high-performance liquid chromatography 586(1991)265
- Direct separation of nadolol enantiomers on a Pirkle-type chiral stationary phase 586(1991)351
- Effect of superfluous remote polar functionality on chiral recognition 589(1992)45
- Silica-bonded chiral stationary phases with structurally simple π -donor chiral selectors for high-performance liquid chromatography 589(1992)53
- Optical resolution by simulated moving-bed adsorption technology 590(1992)113
- Direct liquid chromatographic separation of enantiomers on immobilized protein stationary phases. IX. Influence of the cross-linking reagent on the retentive and enantioselective properties of chiral sorbents based on bovine serum albumin 591(1992)65
- Direct separation of enantiomers by high-performance liquid chromatography on a new chiral ligand-exchange phase 592(1992)291
- Chiral high-performance liquid chromatographic separations on an α_1 -acid glycoprotein column. II. Separation of the diastereomeric and enantiomeric analogues of vinpocetine (Cavinton) 592(1992)297

Retention and enantioselectivity of racemic solutes on a modified ovomucoid-bonded column. I. Cross-linking with glutaraldehyde 592(1992)301

Investigations of the influence of silanol groups on the separation of enantiomers by liquid and supercritical fluid chromatography 592(1992)309

High-performance ligand-exchange chromatography of some amino acids containing two chiral centres 592(1992)345

Synthesis of (1*R*-*trans*)-*N,N'*-1,2-cyclohexylenebisbenzamideoligodimethylsiloxane copolymers for use as chiral stationary phases for capillary supercritical fluid chromatography 594(1992)283

Chromatographic resolution on methylbenzoylcellulose beads. Modulation of the chiral recognition by variation of the position of the methyl group on the aromatic ring 595(1992)63

Direct high-performance liquid chromatographic separation of enantiomeric peptidoleukotriene antagonists 596(1992)123

Contribution of specifiable hydrophobic interactions to chiral recognition 598(1992)1

Investigation of enantioselectivity and enantiomeric elution order of propranolol and its ester derivatives on an ovomucoid-bonded column 598(1992)67

Separation of the enantiomers of the 3,5-dinitrobenzamide derivatives of α -amino phosphonates on four chiral stationary phases 598(1992)159

Contribution of ionically immobilized bovine serum albumin to the retention of enantiomers 600(1992)37

Suitable chiral packing material for the high-performance liquid chromatographic separation of derivatives of 1'-hydroxyeugenol 600(1992)312

Chiral stationary phases, SFC

Chiral π -donor stationary phases with (*R*)-*N*-pivaloylnaphthylethylamide groups for direct enantiomer separation by gas, liquid and supercritical fluid chromatography 557(1991)163

Investigations of the influence of silanol groups on the separation of enantiomers by liquid and supercritical fluid chromatography 592(1992)309

Chloramine-T method

Marked improvement of a substance P radioimmunoassay by reduction of ^{125}I -labelled [Tyr⁸]-substance P prepared by the chloramine-T method with mercaptoethanol and subsequent purification by reversed-phase liquid chromatography 564(1991)67

Chlorella kessleri

Analysis of sterol esters from alga and yeast by high-performance liquid chromatography and capillary gas chromatography-mass spectrometry with chemical ionization 598(1992)219

Chloride attachment

Chloride attachment negative-ion mass spectra of sugars by combined liquid chromatography and atmospheric pressure chemical ionization mass spectrometry 562(1991)81

Chromatobiosis

Immunomagnetic separation of tumor necrosis factor α . II. *In situ* procedure for the human gingival space 583(1992)19

Chromatofocusing

Use of anion-exchange chromatography and chromatofocusing to reveal the structural and functional heterogeneity of topoisomerase II in a HL-60 cell line resistant to multi-drug treatment 587(1991)3

Purification and crystallization of yeast hexokinase isoenzymes. Characterization of different forms by chromatofocusing 587(1991)85

Chromatographic figures of merit

Improved equations for the calculation of chromatographic figures of merit for ideal and skewed chromatographic peaks 594(1992)1

Chromatographic history

Early evolution of chromatography: the activities of Charles Dhéré 600(1992)3

Chromatographic parameters

Effect of the addition of viscous matrices to the mobile phase on chromatographic performance in liquid chromatography-fast atom bombardment mass spectrometry 554(1991)61

Limits and effects of precolumn addition of thioglycerol in liquid chromatographic-fast atom bombardment mass spectrometric systems 591(1992)129

Retention behaviour of closely related coumarins in thin-layer chromatographic preassays for high-performance liquid chromatography according to the "PRISMA" model 593(1992)329

Chromatographic reactor

Reaction kinetics and kinetic processes in modern liquid chromatographic reactors (Review) 589(1992)1

Chromatographic theory

Theoretical approach to the gas chromatographic separation of enantiomers on dissolved cyclodextrin derivatives 552(1991)43

Comparison of various isotherm models for predicting competitive adsorption data 552(1991)71

Is chromatography a separation process? The zonoid answer (Review) 556(1991)25

New approach for calculating ideal chromatograms from arbitrary composite distribution isotherms 556(1991)95

Analysis of the separability of plate height into overload and intrinsic contributions using the kinetic model of non-linear chromatography 556(1991)159

Determination of the hold-up time and column outlet density for capillary supercritical fluid chromatography 588(1991)289

Prediction of the elution profiles of high-concentration bands in gas chromatography 589(1992)223

Experimental determination of adsorption isotherm data for the study of the surface energy distribution of various solid surfaces by inverse gas-solid chromatography 591(1992)233

Precision and accuracy of the gas-solid adsorption isotherms derived by the elution-by-characteristic-points method 591(1992)245

Validity of the model used to relate the energy distribution and the adsorption isotherm 591(1992)267

Theory of linear non-equilibrium chromatography with beds of a finite length 591(1992)273

Modelling of displacement chromatography using non-ideal isotherms 594(1992)9

Validation of chromatographic retention models in reversed-phase high-performance liquid chromatography by fitting experimental data to the relevant equations 595(1992)53

Critical capillary column examination of the relationship between separation number and height equivalent to a theoretical plate and their dependence on temperature 595(1992)209

Van Deemter-type relationship for determining the optimum initial flow-rate and optimum pressure programming rate in temperature/pressure-programmed capillary column gas chromatography utilizing separation numbers 595(1992)221

Variance of a zone migrating in a non-uniform time-invariant linear medium 596(1992)1

Theory of perfusion chromatography 599(1992)87

High-performance liquid chromatography of amino acids, peptides and proteins. CXX. Evaluation of bandwidth behaviour of proteins chromatographed on tentacle-type anion exchangers 599(1992)163

Chromatographic titration

Evaluation of the application of liquid-phase titration to the examination of the adsorption activity of silica-based high-performance liquid chromatographic packings 552(1991)463

Chromatography

see also Special issue index

Circular dichroism spectroscopy

Capillary zone electrophoresis studies of motilin peptides. Effects of charge, hydrophobicity, secondary structure and length 559(1991)391

Chiroptical detection during liquid chromatography. III. Non-stop acquisition of circular dichroism spectra during liquid chromatography 586(1991)249

Citrus oil

Fast separation of polymethoxylated flavones by carbon dioxide supercritical fluid chromatography 586(1991)171

Classification

Characterization and identification of wheat cultivars by multi-dimensional analysis of reversed-phase high-performance liquid chromatograms 596(1992)225

Clean-up methods

Solid-matrix partition for separation of organochlorine pesticide residues from fatty materials 552(1991)241

Rapid method for the determination of organochlorine pesticides in milk 552(1991)249

Multi-matrix-multi-pesticide method for agricultural products 552(1991)259

Removal of organic pollutants from aqueous solution. V. Comparative study of the extraction, recovery and chromatographic separation of some organic insecticides using unloaded polyurethane foam columns 552(1991)371

New Czechoslovak sorbent, Ekosorb 552(1991)449

Miniaturisation of size-exclusion chromatography as a powerful clean-up tool in residue analysis 552(1991)517

Column switching for the high-performance liquid chromatographic analysis of polynuclear aromatic hydrocarbons in petroleum products 552(1991)575

Analysis of sulthiame in serum by narrow-bore high-performance liquid chromatography. Comparison of direct sample injection with pre-column switching and extrelut extraction 553(1991)7

Clean-up of plasma extracts by gel permeation chromatography during analysis of isosorbide nitrates by capillary gas chromatography 553(1991)21

Sensitive determination of the phosphodiesterase III/IV inhibitor zardaverine in human serum by direct sample injection, automated precolumn clean-up and high-performance liquid chromatography 553(1991)65

- Standardization of cation-exchange clean-up prior to gas chromatography of amino acids 553(1991)87
- Improvement of extraction and concentration of milk peptides with solid-phase cartridges for analysis by high-performance liquid chromatography 553(1991)155
- Reversed-phase and chiral high-performance liquid chromatographic assay of bupivacaine and its enantiomers in clinical samples after continuous extraplural infusion 553(1991)383
- Liquid chromatography–thermospray tandem mass spectrometry for identification of a heptabarbital metabolite and sample work-up artefacts 554(1991)205
- Simultaneous separation and determination of hydrocarbons and organochlorine compounds by using a two-step microcolumn 555(1991)221
- Comparative studies on the high-performance liquid chromatographic determination of thiamine and its phosphate esters with chloroethylthiamine as an internal standard using pre- and post-column derivatization procedures 558(1991)115
- Characterization of nitrogen-containing aromatic compounds in soil and sediment by capillary gas chromatography–mass spectrometry after fractionation 558(1991)223
- Chromatographic determination of citric acid for monitoring the mould process 558(1991)302
- Methods for the analysis of thiodiglycol sulphoxide, a metabolite of sulphur mustard, in urine using gas chromatography–mass spectrometry 558(1991)393
- Analysis of 1,1'-sulphonylbis[2-(methylsulphonyl)ethane] and 1-methylsulphonyl-2-[2-(methylthio)ethylsulphonyl]ethane, metabolites of sulphur mustard, in urine using gas chromatography–mass spectrometry 558(1991)405
- Rapid electro-dialytic clean-up of biological samples for high-performance liquid chromatography 585(1991)213
- Determination of polyphosphates in intermediate materials for detergent manufacture by ion high-performance liquid chromatography with post-column derivatization 585(1991)267
- Clean-up procedure for partially methylated alditol acetate derivatives of polysaccharides 587(1991)284
- Detection of zearalenone in cereal extracts using high-performance liquid chromatography with post-column derivatization 588(1991)47
- Determination of a small amount of niacin in foodstuffs by high-performance liquid chromatography 588(1991)171
- High-performance liquid chromatographic–mass spectrometric analysis of oligosaccharides from enzymatic digestion of glycosaminoglycans. Application to human samples 591(1992)225
- Comparative study of clean-up and fractionation methods for the determination of organochlorine pesticides in lipids by gas chromatography 591(1992)303
- Rapid and sensitive determination of urinary 2,5-hexanedione by reversed-phase high-performance liquid chromatography 592(1992)255
- Rapid high-performance liquid chromatographic determination of urinary N-(1-methylethyl)-N'-phenyl-1,4-benzenediamine in workers exposed to aromatic amines 592(1992)279
- High-performance liquid chromatographic determination of penicillin G, penicillin V and cloxacillin in beef and pork tissues 593(1992)15
- Determination of Zy 17617B in plasma by solid-phase extraction and liquid chromatography with automated pre-column exchange 593(1992)73
- Improved determination of individual molecular species of phosphatidylcholine in biological samples by high-performance liquid chromatography with internal standards 593(1992)139
- Analysis and isolation of indole alkaloids of fungi by high-performance liquid chromatography 593(1992)201
- Improved high-performance liquid chromatographic method for the determination of coenzyme Q₁₀ in plasma 593(1992)217
- High-performance liquid chromatography of okadaic acid and free fatty acids in mussels 594(1992)137
- Effects of collection solvent parameters and extraction cell geometry on supercritical fluid extraction efficiencies 594(1992)297
- Automated clean-up procedure for the gas chromatographic–high-resolution mass spectrometric determination of polychlorinated dibenzo-*p*-dioxins and dibenzofurans in milk 595(1992)289
- Identification of ochratoxin A in food samples by chemical derivatization and gas chromatography–mass spectrometry 595(1992)364
- Selective clean-up method using an immunoaffinity column following radioimmunoassay of prostaglandin F_{2α} in biological fluids 597(1992)201
- Closed-loop stripping analysis**
Gas chromatographic–mass spectrometric characterization of volatile organic compounds in Barcelona tap water 562(1991)481

Estimation of tobacco blend compositions using closed-loop stripping analysis and stepwise multiple linear regression and partial least-squares techniques 596(1992)285

Clostridium acetobutlicum

Direct aqueous injection gas chromatography on a potassium fluoride crystal hydrate-containing sorbent. Determination of volatile organic solvents in the fermentation broth of *Clostridium* strains 596(1992)138

Clostridium barkeri

Purification of the coenzyme B₁₂-containing 2-methyleneglutarate mutase from *Clostridium barkeri* by high-performance liquid chromatography 587(1991)93

Cluster analysis

Use of oxidative degradation followed by capillary gas chromatography-mass spectrometry and multi-dimensional scaling analysis to fingerprint unresolved complex mixtures of hydrocarbons 589(1992)281

Cnidoscolus urens

Analysis of keto acids as their methyl esters of 2,4-dinitrophenylhydrazone derivatives by gas chromatography and gas chromatography-mass spectrometry 587(1991)247

Coal tar pitch

Polynuclear aromatic hydrocarbon retention indices on SE-54 stationary phase of the volatile components of a coal tar pitch. Relationships between chromatographic retention and thermal reactivity 591(1992)287

Coated columns

Synthesis and evaluation of epoxy polymer coatings for the analysis of proteins by capillary zone electrophoresis 599(1992)227

Coating layer thickness

Anion-exchange chromatographic properties of α -lactalbumin eluted from quaternized polyvinylimidazole. Study of the role of the polymer coating 553(1991)165

Coating mechanism

Application of a pellicular anion-exchange resin to the separation of inorganic and organic anions by single-column anion exchange 592(1992)101

Coating methods

Gas-liquid microcapillary columns precoated with graphitized carbon black 552(1991)197

Capillary zone electrophoresis of proteins with hydrophilic fused-silica capillaries 559(1991)367

Coating reagents

Charge-reversed, polymer-coated capillary column for the analysis of a recombinant chimeric glycoprotein 594(1992)317

Coatings

Synthesis and evaluation of epoxy polymer coatings for the analysis of proteins by capillary zone electrophoresis 599(1992)227

Coelution

Quantitative measurements via co-elution and dual-isotope detection by gas chromatography-mass spectrometry 587(1991)255

Coil planet centrifuge

Counter-current chromatography of lipoproteins with a polymer phase system using the cross-axis synchronous coil planet centrifuge 596(1992)118

Collaborative study

Collaborative study of the determination of cloxacillin by column liquid chromatography 600(1992)333

Collection solvent

Effects of collection solvent parameters and extraction cell geometry on supercritical fluid extraction efficiencies 594(1992)297

Column capacity

High-performance affinity chromatography of DNA. II. Porosity effects 600(1992)221

Column conditioning

Isotachopheresis of proteins in uncoated open-tubular fused-silica capillaries with a simple approach for column conditioning 558(1991)423

Column coupling

Optimisation pentamétrique de la résolution en chromatographie phase gazeuse par la technique du diagramme à fenêtres. Cas du couplage de deux colonnes 556(1991)287

Column efficiency

Investigation of the causes of reduced efficiency in micellar liquid chromatography 556(1991)263

High-speed gas chromatography. Theoretical and practical aspects 556(1991)331

Effect of hydrostatic flow on the efficiency in capillary electrophoresis 559(1991)81

Fast protein separation by reversed-phase high-performance liquid chromatography on octadecylsilyl-bonded non-porous silica gel. Effect of particle size of column packing on column efficiency 585(1991)207

Chromatographic characterization of a phosphate-modified zirconia support for bio-chromatographic applications 587(1991)149

Simple test for determination of the degree of distortion of the liquid-phase flow profile in columns for preparative liquid chromatography 589(1992)61

Preparative liquid chromatography. I. Influence of column efficiency on optimum injection conditions under isocratic elution 590(1992)175

Peak capacity of columns for size-exclusion chromatography 591(1992)47

High-temperature open-tubular capillary column liquid chromatography 592(1992)239

Electrokinetic reversed-phase chromatography with packed capillaries 593(1992)313

Charge-reversed, polymer-coated capillary column for the analysis of a recombinant chimeric glycoprotein 594(1992)317

Critical capillary column examination of the relationship between separation number and height equivalent to a theoretical plate and their dependence on temperature 595(1992)209

Preparative liquid chromatography. II. Existence of optimum injection conditions for overloaded gradient elution separations 596(1992)15

Column hold-up time

Determination of the hold-up time and column outlet density for capillary supercritical fluid chromatography 588(1991)289

Column length

Optimisation pentaparamétrique de la résolution en chromatographie phase gazeuse par la technique du diagramme à fenêtres. Cas du couplage de deux colonnes 556(1991)287

Factors affecting the separation and loading capacity of proteins in preparative gradient elution high-performance liquid chromatography 590(1992)35

Column liquid chromatography

see also Special issue index

Column material

Effect of column material on sorption isotherms obtained by inverse gas chromatography 600(1992)341

Column overloading

Effect of the sample solvent on band profiles in preparative liquid chromatography using non-aqueous reversed-phase high-performance liquid chromatography 588(1991)1

Column preparation

Experimental determination of adsorption isotherm data for the study of the surface energy distribution of various solid surfaces by inverse gas-solid chromatography 591(1992)233

Column reactors

Use of various types of column reactors for flow-injection analysis 597(1992)435

Column regeneration

Regeneration of column activity after the gas chromatographic separation of membrane lipids on thermostable SE-52 phase 585(1991)342

Columns, GC

Evaluation of gas chromatographic columns for the determination of methylmercury in aqueous head space extracts from biological samples 586(1991)329

Columns, LC

Experimental evaluation of sorbent bed homogeneity in high-performance liquid chromatographic columns 592(1992)109

Columns, permanganate-impregnated

Permanganate-impregnated packed capillary columns for group separation of triacylglycerols using supercritical media as mobile phases 600(1992)358

Column stability

New, stable polyamine-bonded polymer gel column 592(1992)93

Column switching

Development of a rational optimisation procedure for the automated sample clean-up with column switching in pesticide residue analysis 552(1991)113

Column switching for the high-performance liquid chromatographic analysis of polynuclear aromatic hydrocarbons in petroleum products 552(1991)575

Use of an evaporative light scattering detector in reversed-phase high-performance liquid chromatography of oligomeric surfactants 552(1991)593

Analysis of sulthiame in serum by narrow-bore high-performance liquid chromatography. Comparison of direct sample injection with pre-column switching and extrelut extraction 553(1991)7

Sensitive determination of the phosphodiesterase III/IV inhibitor zardaverine in human serum by direct sample injection, automated precolumn clean-up and high-performance liquid chromatography 553(1991)65

Reversed-phase and chiral high-performance liquid chromatographic assay of bupivacaine and its enantiomers in clinical samples after continuous extraplural infusion 553(1991)383

Rapid on-line precolumn high-performance liquid chromatographic method for the determination of benomyl, carbendazim and aldicarb species in drinking water 555(1991)147

Coupled column chromatography in chiral separations: systems employing β -cyclodextrin phases for chiral separation 557(1991)199

Determination of sennoside A and B in the pharmaceutical preparation Otsuji-to using ion-pair high-performance liquid chromatography with column switching 585(1991)255

High-performance liquid chromatographic column-switching technique for the determination of intermediates of anaerobic degradation of toluene in ground water microcosm 587(1991)185

Simultaneous stereoanalysis of 2-alkyl-branched acids, esters and alcohols using a selectivity-adjusted column system in multi-dimensional gas chromatography 587(1991)347

Determination of aflatoxin M1 using a dialysis-based immunoaffinity sample pretreatment system coupled on-line to liquid chromatography. Reusable immunoaffinity columns 589(1992)141

Determination of the catechol-O-methyltransferase inhibitor Ro 40-7592 in human plasma by high-performance liquid chromatography with coulometric detection 593(1992)63

Determination of catecholamines in urine by liquid chromatography and electrochemical detection after on-line sample purification on immobilized boronic acid 593(1992)185

Improved high-performance liquid chromatographic method for the determination of coenzyme Q₁₀ in plasma 593(1992)217

Coupled-column high-performance liquid chromatographic method for the determination of 1-hydroxypyrene in urine of subjects exposed to polycyclic aromatic hydrocarbons 600(1992)189

Application of an internal surface reversed-phase column for the automated determination of flucycloxuron residues 600(1992)211

Column temperature

Optimisation pentaparamétrique de la résolution en chromatographie phase gazeuse par la technique du diagramme à fenêtres. Cas du couplage de deux colonnes 556(1991)287

Effect of temperature on the separation of DNA restriction fragments in capillary gel electrophoresis 559(1991)285

High-temperature open-tubular capillary column liquid chromatography 592(1992)239

Reversed- and normal-phase separations by high-temperature open-tubular column liquid chromatography 598(1992)153

Column tests

Activity testing and surface characterization of pretreated fused-silica capillaries for gas chromatography. A new modification of existing intermediate column tests 590(1992)271

Column throughput

Theory of optimization of ideal displacement chromatography of binary mixtures 590(1992)3

Complexation, on-line

Determination of metal ions by on-line complexation and ion-pair chromatography 590(1992)263

Complexation constants

Measurement of stability constants for complexes of divalent cations with tartaric acid by ion chromatography 589(1992)185

Complex formation

Secondary equilibrium size-exclusion chromatography of ions with polymeric mobile phase additives 586(1991)277

Complex mixture analysis

Complex mixture analysis based on gas chromatography-mass spectrometry with time array detection using a beam deflection time-of-flight mass spectrometer 590(1992)329

Computer algorithm

Improved computer algorithm for characterizing skewed chromatographic band broadening. I. Method 556(1991)111

Computer-assisted chromatography

Computer-aided optimization of the experimental conditions for the isocratic reversed-phase high-performance liquid chromatographic separation of hormonal steroids 552(1991)103

Development of a rational optimisation procedure for the automated sample clean-up with column switching in pesticide residue analysis 552(1991)113

Computer simulation of gradient elution separation. Accuracy of predictions for non-linear gradients 555(1991)1

Optimisation pentaparamétrique de la résolution en chromatographie phase gazeuse par la technique du diagramme à fenêtres. Cas du couplage de deux colonnes 556(1991)287

Accuracy of computer-simulated gas chromatographic separations based on a linear elution strength model 589(1992)215

CRISEBOOK, a Hypermedia version of an expert system for the selection of optimization criteria in high-performance liquid chromatography 592(1992)51

Application of the gradient elution technique. Demonstration with a special test mixture and the DryLab G/plus method development software 592(1992)133

Computer simulation for the prediction of separation as a function of pH for reversed-phase high-performance liquid chromatography. I. Accuracy of a theory-based model 592(1992)183

Computer simulation for the prediction of separation as a function of pH for reversed-phase high-performance liquid chromatography. II. Resolution as a function of simultaneous change in pH and solvent strength 592(1992)197

Optimization of the separation of the Rp and Sp diastereomers of phosphate-methylated DNA and RNA dinucleotides 596(1992)59

Computer simulation of weak affinity chromatography 597(1992)83

Computer-assisted method development

Optimization of separation of rare earths in high-performance thin-layer chromatography 587(1991)359

Computer-controlled densitometry

Quantitative two-dimensional thin-layer chromatography 553(1991)477

Computer programs

Quantitative two-dimensional thin-layer chromatography 553(1991)477

Computer simulation

Computer simulation of gradient elution separation. Accuracy of predictions for non-linear gradients 555(1991)1

Quantitative determination limit in chromatography: computer-based simulations 557(1991)23

Isotachophoretic separation of minor components from a matrix component in the case of strong electrolytes 585(1991)297

GCSIM: a gas-liquid chromatography simulator for educational purposes 588(1991)217

Preparative liquid chromatography. I. Influence of column efficiency on optimum injection conditions under isocratic elution 590(1992)175

Computer simulation as a tool for the rapid optimization of the high-performance liquid chromatographic separation of a tryptic digest of human growth hormone 594(1992)65

Concentration gradient detector

Moving boundary capillary electrophoresis with concentration gradient detection 559(1991)111

Capillary isotachopheresis with concentration-gradient detection. An application to the separation of synthetic peptides 559(1991)489

Conductivity detector

Ion chromatographic determination of anions, especially sulphur-containing anions, with conductimetric and kinetic detection 558(1991)209

Configuration analysis

Configuration analysis of unsaturated hydroxy fatty acids 586(1991)255

Conformational changes

Reversed-phase chromatographic behaviour of β -endorphin: evidence of conformational change 595(1992)125

Conformational isomers

High-performance liquid chromatography of peptides at reduced temperatures: separation of isomers 586(1991)145

High-performance liquid chromatographic separation of *cis-trans* isomers of proline-containing peptides. I. Separation on cyclodextrin-bonded silica 598(1992)139

Connective indices

Effect of temperature and mobile phase on the retention of retinoates in reversed-phase liquid chromatography 592(1992)127

Connectivity model

Relationship between gas chromatographic retention indices and molecular connectivity indices of chlorinated pesticides and structurally related compounds 598(1992)243

Continuous beds

Continuous beds for standard and micro high-performance liquid chromatography 586(1991)21

Continuous-flow dialysis

On-line continuous-flow dialysis thermospray tandem mass spectrometry for quantitative screening of drugs in plasma: roglitamide 598(1992)189

Continuous-flow electrophoresis

Finite difference modelling of continuous-flow electrophoresis 594(1992)341

Continuous-flow electrophoresis: a separation criterion applied to the separation of model proteins 594(1992)351

Continuous stationary phase

Protein chromatography using a continuous stationary phase 598(1992)169

Convection

Protein chromatography using a continuous stationary phase 598(1992)169

Convective diffusion

Dispersion in round tubes and its implications for extra-column dispersion 556(1991)235

Conversion efficiency

Kinetics of acrylamide photopolymerization as investigated by capillary zone electrophoresis 598(1992)277

Coomassie Brilliant Blue staining

Improved detection of human pancreatic proteins separated by two-dimensional isoelectric focusing/sodium dodecyl sulphate gel electrophoresis using a combined staining procedure 563(1991)63

Copolymer phases

Synthesis of (1*R*-*trans*)-N,N'-1,2-cyclohexylenebisbenzamide-*ligodimethylsiloxane* copolymers for use as chiral stationary phases for capillary supercritical fluid chromatography 594(1992)283

Copper electrode

Mixed potential response at metallic electrodes in potentiometric detection for ion chromatography 589(1992)201

Core-shell grafts

Beaded polymer supports and gels (Review). I. Manufacturing techniques 586(1991)181

Correlation coefficient

Assessment of chromatographic peak purity by means of multi-wavelength detection and correlation-based algorithms 592(1992)27

Correspondence factor analysis

Factor analysis and experimental design in high-performance liquid chromatography. XI. Factor analysis maps and chromatographic information 556(1991)169

Counter-current chromatography

Direct coupling of high-speed counter-current chromatography to thin-layer chromatography. Application to the separation of asiaticoside and madecassoside from *Centella asiatica* 558(1991)446

Solvent system for the counter-current purification of hydrophobic bombesin analogues 589(1992)362

Separation of pristinamycins by high-speed counter-current chromatography. I. Selection of solvent system and preliminary preparative studies 593(1992)363

Counter-current chromatography of lipoproteins with a polymer phase system using the cross-axis synchronous coil planet centrifuge 596(1992)118

Counter-current distribution

Extraction of xanthine oxidase from milk by counter-current distribution in an aqueous two-phase system 558(1991)357

Counter-current gas-liquid chromatography

High-temperature continuous counter-current gas-liquid chromatography 590(1992)289

Counter-flow technique

Isotachophoretic analyser with options for operational electrolyte selection and repeated analysis 555(1991)321

Counter-ion inclusion complexation

Influence of counter-ion inclusion complexation on the quality of cyclodextrin-supported separations in isotachopheresis 557(1991)215

Coupled-column methods

Coupled-column high-performance liquid chromatographic method for the determination of 1-hydroxypyrene in urine of subjects exposed to polycyclic aromatic hydrocarbons 600(1992)189

Coupled columns

Theoretical and experimental comparison of serially linked and mixed-packing gas-liquid chromatography columns 555(1991)173

Coupled density detectors

Determination of chemical composition of polymers by size-exclusion chromatography with coupled density and refractive index detection. III. Polyethylene oxide and polytetrahydrofuran 552(1991)507

Coupling methods

Efficient substitution of 1,1'-carbonyldiimidazole activated cellulose and Sepharose matrices with amino acyl spacer arms 587(1991)271

Craig machine

Preparative liquid chromatography. II. Existence of optimum injection conditions for overloaded gradient elution separations 596(1992)15

CRAIGSIM

Preparative liquid chromatography. I. Influence of column efficiency on optimum injection conditions under isocratic elution 590(1992)175

Critical conditions

Analysis of block copolymers by high-performance liquid chromatography under critical conditions 593(1992)233

Cross-linked amylose

Fast separation of macrophages by retention on cross-linked amylose and release by enzymatic amyolysis of the chromatographic material 584(1992)121

Cross-linking co-agents

Immobilization method for polyethylene glycol using a cross-linking co-agent 558(1991)323

Cross-linking reagent

Direct liquid chromatographic separation of enantiomers on immobilized protein stationary phases. IX. Influence of the cross-linking reagent on the retentive and enantioselective properties of chiral sorbents based on bovine serum albumin 591(1992)65

Crude oil

Gas chromatographic retention behaviour of dibenzothiophene derivatives on a smectic liquid crystalline polysiloxane stationary phase 590(1992)297

Analytical study of heavy crude oil fractions by coupling of the transalkylation reaction with supercritical fluid chromatography 593(1992)321

Cryofocusing

Preconcentration technique for introducing gaseous or volatile compounds into a capillary gas chromatographic column 587(1991)338

Cryostat system

Preconcentration technique for introducing gaseous or volatile compounds into a capillary gas chromatographic column 587(1991)338

Cryptioendolithic microorganisms

Novel long-chain *anteiso*-alkanes and *anteiso*-alkanoic acids in Antarctic rocks colonized by living and fossil cryptioendolithic microorganisms 598(1992)267

Crystallization

Purification of erucic acid by preparative high-performance liquid chromatography and crystallization 590(1992)139

Cyanobacteria

High-performance thin-layer chromatographic and high-performance liquid chromatographic determination of abscisic acid produced by cyanobacteria 555(1991)239

Cycle time

Flip-flop elution concept in preparative liquid chromatography 557(1991)137

Cyclic voltammetry

Micellar electrokinetic capillary chromatography of vitamin B₆ with electrochemical detection 585(1991)139

Cyclodextrin-bonded silicas

High-performance liquid chromatographic separation of *cis-trans* isomers of proline-containing peptides. I. Separation on cyclodextrin-bonded silica 598(1992)139

Cystoseiraceae

Determination of sterols and diterpenoids from brown algae (Cystoseiraceae) 588(1991)99

D

Dansylation

Screening for amines by dansylation and automated high-performance liquid chromatography 598(1992)51

Darcy's law

Determination of the hold-up time and column outlet density for capillary supercritical fluid chromatography 588(1991)289

Data acquisition system

High-speed gas chromatography. Theoretical and practical aspects 556(1991)331

Data compression

Data compression in computerized signal processing for isocratic chromatography 585(1991)160

Data precision

Increasing data precision in reversed-phase liquid chromatography leads to new information level 592(1992)13

Dead volume

Theory of linear non-equilibrium chromatography with beds of a finite length 591(1992)273

Increasing data precision in reversed-phase liquid chromatography leads to new information level 592(1992)13

DEAE-cellulose

Purification of common bean chloroplast DNA by DEAE cellulose column chromatography 558(1991)318

Deconvolution methods

Quantitative resolution of severely overlapping gas chromatographic peaks. Isothermal and temperature-programmed operation 556(1991)305

Deconvolution method for accurate determination of overlapping peak areas in chromatograms 587(1991)129

Improved algorithm for resolution of overlapped asymmetric chromatographic peaks 591(1992)19

Use of high-performance liquid chromatographic peak deconvolution and peak labelling to identify antiparasitic components in plant extracts 593(1992)209

Denaturation

Retention behaviour of a template-assembled synthetic protein and its amphiphilic building blocks on reversed-phase columns 586(1991)43

Densitometric equipment

Important design features of a system for the densitometric analysis of two-dimensional flat-bed separations 596(1992)241

Densitometry

Important design features of a system for the densitometric analysis of two-dimensional flat-bed separations 596(1992)241

Density drop

Determination of the hold-up time and column outlet density for capillary supercritical fluid chromatography 588(1991)289

Density gradients

Packed column supercritical fluid chromatography with carbon dioxide-polar modifiers. Influence of carbon dioxide density on retention 557(1991)85

Deoxygenation

Reductive electrochemical detection in liquid chromatography with a zinc amalgam scrubber column 598(1992)195

Derivatization

Increasing thermospray response for cortisol by derivatization 554(1991)149

Derivatization, electrophoresis

Separation of fluorescent oligosaccharide derivatives by microcolumn techniques based on electrophoresis and liquid chromatography 559(1991)223

High-performance capillary zone electrophoresis of carbohydrates in the presence of alkaline earth metal ions 588(1991)327

Separation of naphthalene-2,3-dicarboxaldehyde-labeled amino acids by high-performance capillary electrophoresis with laser-induced fluorescence detection 593(1992)265

Assay of protein drug substances present in solution mixtures by fluorescence derivatization and capillary electrophoresis 598(1992)123

Derivatization, GC

Rapid derivatization and gas chromatographic determination of amino acids 552(1991)289

Gas chromatographic determination of N-carboxymethyl amino acids, the periodate oxidation products of Amadori compounds 552(1991)245

Chromatographic trace analysis of some organic compounds in the environment using derivatization-sorption concentration techniques. I. Gas chromatographic analysis of acrylates in air 557(1991)523

Gas chromatographic determination of the lewisite hydrolysate, 2-chlorovinylarsonous acid, after derivatization with 1,2-ethanedithiol 558(1991)235

Methods for the analysis of thiodiglycol sulphoxide, a metabolite of sulphur mustard, in urine using gas chromatography-mass spectrometry 558(1991)393

Analysis of 1,1'-sulphonylbis[2-(methylsulphinyl)ethane] and 1-methylsulphinyl-2-[2-(methylthio)ethylsulphonyl]ethane, metabolites of sulphur mustard, in urine using gas chromatography-mass spectrometry 558(1991)405

Capillary gas chromatographic analysis of oxo, oxo-hydroxy and unsaturated bile acid glycine conjugates 558(1991)451

Intra-injector methylation of free fatty acids from aerobically and anaerobically cultured *Actinobacillus actinomycetemcomitans* and *Haemophilus aphrophilus* 576(1992)328

Gas chromatographic-mass spectrometric determination of ethyl carbamate as the xanthylamide derivative in Italian aqua vitae (grappa) samples 585(1991)111

Studies on steroids. CCLIII. Capillary gas chromatographic behaviour of diethylhydrogensilyl-diethylsilylene derivatives of stereoisomeric bile acids 585(1991)281

Determination of trace amounts of fluoride in raw materials for pharmaceuticals by gas-liquid chromatography 585(1991)289

Origin and control of multi-peak formation in the analysis of trimethylsilyl derivatives of flavanone aglycones by capillary column gas chromatography 586(1991)323

Analysis of keto acids as their methyl esters of 2,4-dinitrophenylhydrazone derivatives by gas chromatography and gas chromatography-mass spectrometry 587(1991)247

Clean-up procedure for partially methylated alditol acetate derivatives of polysaccharides 587(1991)284

Gas chromatographic separation of diastereomeric amino acid derivatives on chiral stationary phases. Application to the determination of enantiomeric composition in (S)-(+)-2-butanol 589(1992)249

Analytical methods for monoterpene glycosides in grape and wine. I. XAD-2 extraction and gas chromatographic-mass spectrometric determination of synthetic glycosides 590(1992)313

Determination of Pb²⁺ in water by isotope dilution gas chromatography-mass spectrometry of tetraethyllead formed by reaction with sodium tetraethylborate 594(1992)275

Structure-retention index relationships for derivatized monosaccharides on non-polar gas chromatography columns 596(1992)79

Determination of 7-methylbenz[c]acridines by capillary gas chromatography with electron-capture detection 596(1992)233

Structure elucidation of saccharides in anthocyanins and flavonols by means of methylation analysis and gas chromatography 598(1992)81

Determination of selenium by capillary gas chromatography after high-temperature derivatization with 1,2-diamino-3,5-dibromobenzene 598(1992)105

Some novel homochiral derivatizing agents for the gas chromatographic analysis of enantiomeric secondary alcohols 598(1992)251

Intra-injector formation of methyl esters from phenoxy acid pesticides 598(1992)309

Derivatization, LC

Time-resolved luminescence detection of derivatized thiols in column liquid chromatography 552(1991)625

Separation of elastin cross-links as phenylisothiocyanate derivatives 553(1991)93

Selective post-column fluorogenic reaction with benzamidine for trace level detection of reducing saccharides in liquid chromatography 553(1991)255

Sensitive liquid chromatographic determination of alkyl-, nitro- and chlorophenols by precolumn derivatization with dansyl chloride, postcolumn photolysis and peroxyoxalate chemiluminescence detection 553(1991)345

Influence of chemical structure of tricyclic tertiary dimethylamines on chiral separation by reversed-phase high-performance liquid chromatography after derivatization with (-)-menthylchloroformate 553(1991)365

On-line derivatization of eluted substances in dynamic high-performance liquid chromatography-mass spectrometry through the particle-beam interface 554(1991)125

High-performance liquid chromatographic separation of DL-amino acids derivatized with chiral variants of Sanger's reagent 555(1991)81

Utility of dansyl derivatization to the high-performance liquid chromatographic analysis of 2-phenylethylamine drugs 555(1991)278

Enantiomer separation by chiral-phase liquid chromatography of urethane derivatives of natural diacylglycerols previously fractionated by reversed-phase liquid chromatography 557(1991)227

Diffusion of proteins in the chromatographic gel AcA-34 558(1991)71

Comparative studies on the high-performance liquid chromatographic determination of thiamine and its phosphate esters with chloroethylthiamine as an internal standard using pre- and post-column derivatization procedures 558(1991)115

Liquid chromatographic method for the determination of calcium cyanamide using pre-column derivatization 558(1991)141

Pre-column derivatization of sulfa drugs with fluorescamine and high-performance liquid chromatographic determination at their residual levels in meat and meat products 558(1991)175

Effect of derivatization of steroids on their retention behaviour in inclusion chromatography using cyclodextrin as a mobile phase additive 558(1991)306

High-performance liquid chromatographic detection of enantiomeric amino alcohols after derivatization with *o*-phthaldialdehyde and various thiosugars 558(1991)311

Separation of fluorescent oligosaccharide derivatives by microcolumn techniques based on electrophoresis and liquid chromatography 559(1991)223

Determination of alcohols by high-performance liquid chromatography after pre-column derivatization with 2-(4-carboxyphenyl)-5,6-dimethylbenzimidazole 585(1991)219

Electrochemical and chromatographic properties of selected hydrazine and hydrazide derivatives of carbonyl compounds 585(1991)225

Determination of polyphosphates in intermediate materials for detergent manufacture by ion high-performance liquid chromatography with post-column derivatization 585(1991)267

Derivatization of saturated long-chain fatty acids with phenacyl bromide in non-ionic micelles 586(1991)347

Recovery of sugar derivatives from 2-aminopyridine labeling mixtures for high-performance liquid chromatography using UV or fluorescence detection 587(1991)177

Rapid high-performance liquid chromatographic determination of fatty acid profiles of lipids by conversion to their hydroxamic acids 587(1991)292

Detection of zearalenone in cereal extracts using high-performance liquid chromatography with post-column derivatization 588(1991)47

Separation of neutral mono- and oligosaccharides derivatized with ethyl *p*-aminobenzoate by high-performance liquid chromatography on an amine-bonded vinyl alcohol copolymer column 588(1991)53

Precolumn fluorescence tagging reagent for carboxylic acids in high-performance liquid chromatography: 4-substituted-7-aminoalkylamino-2,1,3-benzoxadiazoles 588(1991)61

Applications of automated amino acid analysis using 9-fluorenylmethyl chloroformate 588(1991)107

Optical resolution of 2-chloro-3-phenylmethoxypropanoic acid after derivatization with (*S*)-2-octanol by high-performance liquid chromatography 589(1992)346

Sensitive fluorescence detection of some nitrosamines by precolumn derivatization with dansyl chloride and high-performance liquid chromatography 589(1992)349

High-performance liquid chromatographic determination of 2-furaldehyde in spirits 590(1992)235

3-Bromomethyl-7-methoxy-1,4-benzoxazin-2-one as a highly sensitive fluorescence derivatization reagent for carboxylic acids in high-performance liquid chromatography 591(1992)159

Amino acid analysis by high-performance liquid chromatography after derivatization with diethyl ethoxymethylenemalonate 591(1992)181

Determination of the enantiomers of fenoldopam in human plasma by reversed-phase high-performance liquid chromatography after chiral derivatization 592(1992)317

By-products in the derivatization of amines with the chiral reagent 2,3,4,6-tetra-O-acetyl- β -D-glucopyranosyl isothiocyanate and their elimination 592(1992)323

Determination of amphetamines by high-performance liquid chromatography with ultraviolet detection. On-line pre-column derivatization with 9-fluorenylmethyl chloroformate and preconcentration 593(1992)9

High-performance liquid chromatographic determination of monohydroxy compounds by a combination of pre-column derivatization and post-column reaction detection 593(1992)29

High-performance liquid chromatography method for the determination of aminoglycosides based on automated pre-column derivatization with *o*-phthalaldehyde 593(1992)69

High-performance liquid chromatographic method with fluorescence detection for the determination of total homocyst(e)ine in plasma 593(1992)171

Comparison of several methods for the determination of trace amounts of polar aliphatic monocarboxylic acids by high-performance liquid chromatography 593(1992)243

Primary and secondary amine derivatization with luminarins 1 and 2: separation by liquid chromatography with peroxyoxalate chemiluminescence detection 594(1992)386

Fluorescence detection of trichothecene mycotoxins as coumarin-3-carbonyl chloride derivatives by high-performance liquid chromatography 595(1992)143

Quantitative analysis of total resin acids by high-performance liquid chromatography of their coumarin ester derivatives 595(1992)155

High-performance liquid chromatography of histamine and 1-methylhistamine with on-column fluorescence derivatization 595(1992)163

High-performance liquid chromatography and post-column derivatization with diphenyl-1-pyrenylphosphine for fluorimetric determination of triacylglycerol hydroperoxides 596(1992)197

Phenylboronic acid as a versatile derivatization agent for chromatography of ecdysteroids 596(1992)271

Affinity chromatographic purification of bovine lung endothelin receptor using biotinylated endothelin and avidin-agarose 597(1992)331

Screening for amines by dansylation and automated high-performance liquid chromatography 598(1992)51

4,5-Diaminophthalhydrazide as a highly sensitive chemiluminescence derivatization reagent for α -dicarbonyl compounds in high-performance liquid chromatography 598(1992)203

Derivatization, SFC

Separation of secondary alcohol enantiomers using supercritical fluid chromatography 585(1991)117

Derivatization, TLC

Phenylboronic acid as a versatile derivatization agent for chromatography of ecdysteroids 596(1992)271

Dermatophagoides pteronyssinus

Isolation of *Der pI*, the *Dermatophagoides pteronyssinus* major mite allergen, from a crude mite culture extract, purification by ion-chromatography, and comparison between the material obtained and a cDNA-coded *Der pI* 599(1992)105

Derringer's desirability function

Simultaneous optimization of several chromatographic performance goals using Derringer's desirability function 586(1991)11

Desirability function

Simultaneous optimization of several chromatographic performance goals using Derringer's desirability function 586(1991)11

Desorption

Production and use of capillary traps for headspace gas chromatography of airborne volatile organic compounds 552(1991)137

Study of high-affinity interactions by quantitative affinity chromatography. Analytical expressions in terms of total ligand concentration 597(1992)57

Desorption isotherms

Adsorption-desorption isotherm hysteresis of β -lactoglobulin A with a weakly hydrophobic surface 557(1991)369

Detection sensitivity

Effects of β -cyclodextrin in the mobile phase on the retention and indirect detection of non-electrolytes in reversed-phase liquid chromatography. I. Study of aliphatic alcohols 552(1991)563

Detector cells

Fluorescence detector cell for use in an integrated electrically driven separation system 598(1992)115

Detectors, electrophoresis

Dilution effect of carbonate ion on isotachophoretic zones observed with scanning UV photometric detection 555(1991)247

Moving boundary capillary electrophoresis with concentration gradient detection 559(1991)111

Optimization and evaluation of the performance of arrangements for UV detection in high-resolution separations using fused-silica capillaries 559(1991)163

Low-cost, high-sensitivity laser-induced fluorescence detection for DNA sequencing by capillary gel electrophoresis 559(1991)237

Capillary isotachopheresis with concentration-gradient detection. An application to the separation of synthetic peptides 559(1991)489

Determination of small drug molecules by capillary electrophoresis-atmospheric pressure ionization mass spectrometry 559(1991)515

Micellar electrokinetic capillary chromatography of vitamin B₆ with electrochemical detection 585(1991)139

Quantitative aspects of indirect UV detection in capillary zone electrophoresis 588(1991)321

Binary buffers for indirect absorption detection in capillary zone electrophoresis 589(1992)307

Capillary isoelectric focusing with electroosmotic zone displacement and on-column multichannel detection 589(1992)321

Capillary electrophoresis with electrochemical detection employing an on-column Nafion joint 593(1992)305

High-sensitivity laser-induced fluorescence detection of native proteins in capillary electrophoresis 595(1992)319

Capillary gel electrophoresis of single-stranded DNA fragments with UV detection 600(1992)289

Detectors, GC

Multi-matrix-multi-pesticide method for agricultural products 552(1991)259

Simultaneous separation and determination of hydrocarbons and organochlorine compounds by using a two-step microcolumn 555(1991)221

High-speed gas chromatography. Theoretical and practical aspects 556(1991)331

Laser-excited fluorescence detection of gas-phase chromatography eluates 557(1991)39

Structure determination of sesquiterpenes in Chinese vetiver oil by gas chromatography-tandem mass spectrometry 557(1991)451

Sensitive determination of phenothiazines in body fluids by gas chromatography with surface ionization detection 579(1992)247

Measurements using gas chromatography with coelution and dual-isotope atomic emission detection 586(1991)309

Evaluation of gas chromatographic columns for the determination of methylmercury in aqueous head space extracts from biological samples 586(1991)329

Photoemissive ionisation source for ion mobility detectors 588(1991)239

Flame infrared emission detection of hydrogen 589(1992)241

Determination of sulfur components in light petroleum streams by high-resolution gas chromatography with chemiluminescence detection 589(1992)271

Comparison of commercially available atomic emission and chemiluminescence detectors for sulfur-selective gas chromatographic detection 591(1992)313

Gas chromatography of chloride and bromide as phenylboronic acid/mercuric nitrate derivatives with microwave induced plasma atomic emission detection 594(1992)395

Determination of α - and β -zeaxenol and zeaxenone in cereals by gas chromatography with ion-trap detection 595(1992)259

Wide dynamic range electron-capture detection using the electron-capture detector pulse frequency signal 595(1992)359

Determination of organotin species by capillary gas chromatography with alternating current plasma emission detection 598(1992)97

Flame-photometric detection of nitrous oxide in addition to phosphine 598(1992)313

Detectors, LC

Determination of chemical composition of polymers by size-exclusion chromatography with coupled density and refractive index detection. III. Polyethylene oxide and polytetrahydrofuran 552(1991)507

Use of an evaporative light scattering detector in reversed-phase high-performance liquid chromatography of oligomeric surfactants 552(1991)593

Time-resolved luminescence detection of derivatized thiols in column liquid chromatography 552(1991)625

Spray jet assembly interface for the coupling of reversed-phase narrow-bore liquid chromatography and Fourier transform infrared spectrometry 552(1991)635

Comparison of high-performance liquid chromatographic detection methods for thyronine and tyrosine residues in toxicological studies of the thyroid 553(1991)135

Fluorescence detection in liquid chromatography with an intensified diode-array detector 556(1991)495

On-line electrochemical reagent generation for liquid chromatography with luminol-based chemiluminescence detection 557(1991)13

Separation and indirect detection of amino acids as acetylated derivatives 557(1991)315

Analysis of nucleotides by high-performance liquid chromatography with phosphorus-selective detection 557(1991)441

Liquid chromatographic method for the determination of the carbohydrate moiety of glycoproteins. Application to α_1 -acid glycoprotein and tissue plasminogen activator 558(1991)105

Determination of benzocaine, dextromethorphan and cetylpyridinium ion by high-performance liquid chromatography with UV detection 558(1991)147

Ion chromatographic determination of anions, especially sulphur-containing anions, with conductimetric and kinetic detection 558(1991)209

Optimization and evaluation of the performance of arrangements for UV detection in high-resolution separations using fused-silica capillaries 559(1991)163

Determination of phosphatidylcholine, phosphatidylglycerol and their lyso forms from liposome dispersions by high-performance liquid chromatography using high-sensitivity refractive index detection 585(1991)239

Chiroptical detection during liquid chromatography. III. Non-stop acquisition of circular dichroism spectra during liquid chromatography 586(1991)249

Assessment of a thermospray interface for the coupling of high-performance liquid chromatography and Fourier transform infrared spectrometry 588(1991)15

Mixed potential response at metallic electrodes in potentiometric detection for ion chromatography 589(1992)201

Potentiometric detection of carboxylic acids and inorganic anions in ion-exclusion chromatography using camphorsulphonic acid as eluent 589(1992)209

High-performance liquid chromatography and supercritical fluid chromatography of monosaccharides and polyols using light-scattering detection. Chemometric studies of the retentions 590(1992)203

Interfacing gradient elution ion-exchange chromatography and low-angle laser light-scattering photometry for analysis of proteins 591(1992)139

Comparison of ethoxylated alcohols and polyethylene glycols by high-performance liquid chromatography and supercritical fluid chromatography using evaporative light-scattering detection 591(1992)149

UV spectrophotometric detection in the ion chromatographic analysis of alkali and alkaline earth metal and ammonium cations 591(1992)219

Nitrogen-specific liquid chromatography detector based on chemiluminescence. Application to the analysis of ammonium nitrogen in waste water 592(1992)209

Practical on-line determination of biopolymer molecular weights by high-performance liquid chromatography with classical light-scattering detection 592(1992)215

Constant-potential amperometric detection of carbohydrates at metal electrodes in high-performance anion-exchange chromatography 592(1992)229

Determination of National Survey of Pesticides analytes in groundwater by liquid chromatography with postcolumn reaction detection 592(1992)283

High-performance liquid chromatographic analysis of benzodiazepines using diode array, electrochemical and thermospray mass spectrometric detection 598(1992)59

Fluorescence detector cell for use in an integrated electrically driven separation system 598(1992)115

Detectors, TLC

Use of hematoporphyrin as a fluorescent stain for detection of lipids in high-performance thin-layer chromatography 588(1991)295

Determination limit

Quantitative determination limit in chromatography: computer-based simulations 557(1991)23

Deuteration

Deuteration as an aid to the high-temperature gas chromatography-mass spectrometry of steryl fatty acyl esters 590(1992)305

Dhéré, C.

Early evolution of chromatography: the activities of Charles Dhéré 600(1992)3

Dialysis

Improved affinity chromatographic purification of D-mannose-N-acetyl-D-glucosamine-specific lectin from the bark of *Sophora japonica* eliminating the loss by sugar specific self-aggregation 597(1992)197

Diastereomer separation

High-performance liquid chromatographic separation of DL-amino acids derivatized with chiral variants of Sanger's reagent 555(1991)81

High-performance liquid chromatography of diastereomeric flavanone glycosides in *Citrus* on a β -cyclodextrin-bonded stationary phase (Cyclobond I) 588(1991)41

Optimization of the separation of the Rp and Sp diastereomers of phosphate-methylated DNA and RNA dinucleotides 596(1992)59

Diatomaceous supports

Gas chromatography in qualitative analysis. XX. The deactivation of diatomaceous supports by batyl alcohol 552(1991)169

Dielectric constant

Correlation analysis in liquid chromatography of metal chelates. II. One-dimensional retention-mobile phase composition (physicochemical property) models in reversed-phase liquid chromatography 555(1991)163

Diesel fuel

Quantitative hydrocarbon group analysis of gasoline and diesel fuel by supercritical fluid chromatography 595(1992)301

Differential absorbance

Chiroptical detection during liquid chromatography. III. Non-stop acquisition of circular dichroism spectra during liquid chromatography 586(1991)249

Differential chromatography

Predictability of chromatographic protein separations. Study of size-exclusion media with narrow particle size distributions 589(1992)71

Differential migration

Immobilized metal ion affinity electrophoresis. A study with several model proteins containing histidine 597(1992)357

Differential thermal analysis

Thermoanalytical and chromatographic studies of copper(II), nickel(II) and oxovanadium(IV) complexes of tetradentate Schiff bases derived from β -diketones and 2,3-diaminopentane 558(1991)187

Diffusion

Diffusion of sorbed pyrene in the bonded layer of reversed-phase silicas. Effect of alkyl chain length and pore diameter 558(1991)31

Finite difference modelling of continuous-flow electrophoresis 594(1992)341

Diffusion coefficients

Source of band broadening in liquid chromatographic-fast atom bombardment mass spectrometric systems with precolumn addition of viscous matrix to the mobile phase 554(1991)47

Magnitude of the diffusion coefficient anomaly in the critical region and its effect on supercritical fluid chromatography 557(1991)69

Diffusion of proteins in the chromatographic gel AcA-34 558(1991)71

Diffusion of proteins in Sepharose Cl-B gels 591(1992)115

Peak dispersion and separation efficiency in high-performance zone electrophoresis with gel-filled capillaries 595(1992)313

Dihydroxy-saturated urushiol crown ether

Study of dihydroxy-substituted saturated urushiol crown ether as a stationary phase in capillary gas chromatography 589(1992)368

Diode-array detector

Identification of flavonoids from *Ginkgo biloba* L., *Anthemis nobilis* L. and *Equisetum arvense* L. by high-performance liquid chromatography with diode-array UV detection 553(1991)223

Fluorescence detection in liquid chromatography with an intensified diode-array detector 556(1991)495

Determination of benzocaine, dextromethorphan and cetylpyridinium ion by high-performance liquid chromatography with UV detection 558(1991)147

Determination of sulfonamides by liquid chromatography, ultraviolet diode array detection and ion-spray tandem mass spectrometry with application to cultured salmon flesh 558(1991)155

Separation of antibiotics by high-performance capillary electrophoresis with photodiode-array detection 585(1991)133

Determination of (S)-(-)-cathinone and its metabolites (R,S)-(-)-norephedrine and (R,R)-(-)-norpseudoephedrine in urine by high-performance liquid chromatography with photodiode-array detection 593(1992)79

Determination of psychotropic phenylalkylamine derivatives in biological matrices by high-performance liquid chromatography with photodiode-array detection 593(1992)87

High-performance liquid chromatographic analysis of benzodiazepines using diode array, electrochemical and thermospray mass spectrometric detection 598(1992)59

Dipolarity

Hydrogen bonding. XVI. A new solute solvation parameter, π^H_2 , from gas chromatographic data 587(1991)213

Hydrogen bonding. XVII. The characterisation of 24 gas-liquid chromatographic stationary phases studied by Poole and co-workers, including molten salts, and evaluation of solute-stationary phase interactions 587(1991)229

Dipole-dipole interactions

Preparation and capillary column gas chromatographic characterization of aza-crown ether polysiloxane stationary phases 594(1992)243

Direct liquid introduction

Strategies in developing interfaces for coupling liquid chromatography and mass spectrometry 554(1991)3

Simple direct liquid introduction system usable as an interface for liquid chromatography-mass spectrometry on quadrupole and magnetic-sector mass spectrometers 554(1991)293

Discrimination

Characterization and identification of wheat cultivars by multi-dimensional analysis of reversed-phase high-performance liquid chromatograms 596(1992)225

Dispersion

Dispersion in round tubes and its implications for extra-column dispersion 556(1991)235

Mathematical model describing dispersion in free solution capillary electrophoresis under stacking conditions 559(1991)3

Variance of a zone migrating in a non-uniform time-invariant linear medium 596(1992)1

Dispersion interaction

Validation of chromatographic retention models in reversed-phase high-performance liquid chromatography by fitting experimental data to the relevant equations 595(1992)53

Dispersion operators

Source of band broadening in liquid chromatographic-fast atom bombardment mass spectrometric systems with precolumn addition of viscous matrix to the mobile phase 554(1991)47

Displacement, mobility-selective

Binary buffers for indirect absorption detection in capillary zone electrophoresis 589(1992)307

Displacement chromatography

Analysis of isotachic patterns in displacement chromatography 556(1991)119

Characterization of a tryptic digest by high-performance displacement chromatography and mass spectrometry 557(1991)289

Theory of optimization of ideal displacement chromatography of binary mixtures 590(1992)3

Peak purity, yield and throughput in preparative liquid chromatography with self-displacement 592(1992)17

Modelling of displacement chromatography using non-ideal isotherms 594(1992)9

Displacement effects

Displacement effects in preparative gradient high-performance liquid chromatographic separations 590(1992)17

Experimental study of the production rate of pure enantiomers from racemic mixtures 590(1992)119

Influence of operating parameters on the preparative gradient elution chromatography of insulins 599(1992)195

Displacement electrophoresis

see Isotachophoresis

Displacement model

Validation of chromatographic retention models in reversed-phase high-performance liquid chromatography by fitting experimental data to the relevant equations 595(1992)53

Displacer salts

High-performance liquid chromatography of amino acids and dipeptides on new ion exchangers of the HEMA series 552(1991)439

High-performance liquid chromatography of amino acids, peptides and proteins. CXX. Evaluation of bandwidth behaviour of proteins chromatographed on tentacle-type anion exchangers 599(1992)163

Dissociation constants

Electrophoretic mobility and dissociation constants of tripeptides evaluated by isotachophoresis 585(1991)364

Determination of limiting ionic mobilities and dissociation constants of some local anaesthetics 596(1992)265

Distillation

Simulated distillation of distillates on capillary columns: influence of the polarity of the stationary phase 557(1991)155

Distribution coefficients

Analysis of block copolymers by high-performance liquid chromatography under critical conditions 593(1992)233

Distribution equilibrium

New approach for calculating ideal chromatograms from arbitrary composite distribution isotherms 556(1991)95

Domains

Affinity purification of antibodies using immobilized FB domain of protein A 597(1992)257

Drug monitoring

On-line continuous-flow dialysis thermospray tandem mass spectrometry for quantitative screening of drugs in plasma: rogletimide 598(1992)189

DryLab G

Computer simulation of gradient elution separation. Accuracy of predictions for non-linear gradients 555(1991)1

DryLab G/plus

Application of the gradient elution technique. Demonstration with a special test mixture and the DryLab G/plus method development software 592(1992)133

DryLab GC

Accuracy of computer-simulated gas chromatographic separations based on a linear elution strength model 589(1992)215

Drylab I/mp

Computer simulation for the prediction of separation as a function of pH for reversed-phase high-performance liquid chromatography. I. Accuracy of a theory-based model 592(1992)183

Computer simulation for the prediction of separation as a function of pH for reversed-phase high-performance liquid chromatography. II. Resolution as a function of simultaneous change in pH and solvent strength 592(1992)197

Dual-column systems

Comparative analysis of thirty polychlorinated biphenyl congeners on two capillary columns of different polarity with non-linear multi-level calibration 558(1991)375

Dual detection, GC

Photoionization assessment of C₃-C₅ alkadienes and alkenes in urban air 591(1992)358

Dual-electrode electrochemical detector

Apparent inter-channel interference in dual-electrode electrochemical detection 596(1992)110

Dual-isotope atomic emission detector

Measurements using gas chromatography with coelution and dual-isotope atomic emission detection 586(1991)309

Dual-isotope detection

Quantitative measurements via co-elution and dual-isotope detection by gas chromatography-mass spectrometry 587(1991)255

Dye labelling

Optimization of oxidation of glycoproteins: an assay for predicting coupling to hydrazide chromatographic supports 587(1991)171

Dye-ligand affinity partitioning

Dye-ligand affinity partitioning of lactate dehydrogenase isoenzymes 557(1991)325

Dynamically modified silica

Study of the retention behaviour of barbiturates by overpressured layer chromatography using silica gel bonded with tricaprilmethylammonium chloride 589(1992)301

Determination of glutamine, glutamic acid and pyroglutamic acids using high-performance liquid chromatography on dynamically modified silica 594(1992)173

Determination of inorganic anions in salt solutions by ion chromatography using C₁₈ reversed-phase columns coated with cetyltrimethylammonium 598(1992)237

Dynamic binding constants

Determination of solute-polymer interaction constants via chromatography with polymer capillaries 594(1992)29

Dynamic ion exchangers

Modification of reversed-phase columns with dyed surfactants. Preparation of mechanically resistant efficient immobilized dyes for protein purification 558(1991)43

Reversed-phase separation of transition metals, lanthanides and actinides by elution with mandelic acid 558(1991)197

Dynamic processes

Reaction kinetics and kinetic processes in modern liquid chromatographic reactors (Review) 589(1992)1

Dynamic protein binding capacities

Characterization of synthetic macroporous packing materials in low-pressure cartridges and columns 590(1992)83

Dynamic solute adsorption

Determination of solute-polymer interaction constants via chromatography with polymer capillaries 594(1992)29

E**Edgeworth-Cramér fitting**

Peak-shape analysis and noise evaluation in suppressed ion chromatography for ultra-trace ion analysis 556(1991)249

Effective mobility

Dilution effect of carbonate ion on isotachophoretic zones observed with scanning UV photometric detection 555(1991)247

Influence of poly(vinylpyrrolidone) on isotachophoretic separations of inorganic anions in aqueous electrolyte systems 589(1992)329

Determination of sulphonamides in pork meat extracts by capillary zone electrophoresis 596(1992)101

Efficiency

see Column efficiency

Eigenzones

Experimental evaluation of sorbent bed homogeneity in high-performance liquid chromatographic columns 592(1992)109

Einstein–Nernst equation

Peak dispersion and separation efficiency in high-performance zone electrophoresis with gel-filled capillaries 595(1992)313

Electrically driven separations

Fluorescence detector cell for use in an integrated electrically driven separation system 598(1992)115

Electrochemical detector

Determination of free DOPA and 3-O-methyl-DOPA in human plasma by high-performance liquid chromatography with electrochemical detection 553(1991)265

Micellar electrokinetic capillary chromatography of vitamin B₆ with electrochemical detection 585(1991)139

Electrochemical and chromatographic properties of selected hydrazine and hydrazide derivatives of carbonyl compounds 585(1991)225

Constant-potential amperometric detection of carbohydrates at metal electrodes in high-performance anion-exchange chromatography 592(1992)229

Capillary electrophoresis with electrochemical detection employing an on-column Nafion joint 593(1992)305

Apparent inter-channel interference in dual-electrode electrochemical detection 596(1992)110

High-performance liquid chromatographic analysis of benzodiazepines using diode array, electrochemical and thermospray mass spectrometric detection 598(1992)59

Electrochemical reagent generation

On-line electrochemical reagent generation for liquid chromatography with luminol-based chemiluminescence detection 557(1991)13

Electrochromatographic solid-phase extraction

Electrochromatographic solid-phase extraction for determination of cimetidine in serum by micellar electrokinetic capillary chromatography 559(1991)547

Electrochromatography

Electroosmotically driven electrochromatography of anions having similar electrophoretic mobilities by ion pairing 557(1991)125

Electrodialysis

Rapid electro-dialytic clean-up of biological samples for high-performance liquid chromatography 585(1991)213

Electrodialytic sample treatment coupled on-line with column liquid chromatography for the determination of basic and acidic compounds in environmental samples 600(1992)163

Electroelution

Preparative high-yield electroelution of proteins after separation by sodium dodecyl sulphate–polyacrylamide gel electrophoresis and its application to analysis of amino acid sequences and to raise antibodies 585(1991)153

Electroendosmotic elution

Electrokinetic reversed-phase chromatography with packed capillaries 593(1992)313

Electro-injection

Field amplified sample injection in high-performance capillary electrophoresis 559(1991)141

Field-amplified polarity-switching sample injection in high-performance capillary electrophoresis 559(1991)153

Electrokinetic chromatography

Electrokinetic reversed-phase chromatography with packed capillaries 593(1992)313

Electrolyte systems

Relationship between zone length and step height in isotachophoresis 595(1992)327

Electron-acceptor stationary phases

Physico-chemical properties of electron-acceptor stationary phases in liquid chromatography 591(1992)75

Electron-capture detector

Wide dynamic range electron-capture detection using the electron-capture detector pulse frequency signal 595(1992)359

Electronic scanning

Important design features of a system for the densitometric analysis of two-dimensional flat-bed separations 596(1992)241

Electron-impact mass spectrometry

Some aspects of peak broadening in particle-beam liquid chromatography–mass spectrometry 554(1991)119

Characterization of N-acyl-D-biotinols by particle-beam liquid chromatography–mass spectrometry. An alternative to probe mass spectrometry for thermally labile samples 554(1991)141

Characterization of nitrogen-containing aromatic compounds in soil and sediment by capillary gas chromatography–mass spectrometry after fractionation 558(1991)223

Simultaneous determination of sorbic acid, dehydroacetic acid and benzoic acid by gas chromatography–mass spectrometry 594(1992)253

Electroosmosis

Field effect electroosmosis 559(1991)95

Analysis of separation efficiency in capillary electrophoresis with direct control of electroosmosis by using an external electric field 559(1991)133

Finite difference modelling of continuous-flow electrophoresis 594(1992)341

Electroosmotic flow

Evaluation of fundamental properties of a silica capillary used for capillary electrophoresis 559(1991)69

Studies of the electrophoretic behaviour of *p*-dihydroxyborylphenylalanine and related compounds by the three-spot method 585(1991)369

Capillary tube isotachopheric separation of catecholamines using cyclodextrin in the leading electrolyte 587(1991)364

Radial pH distribution during capillary electrophoresis with electroosmotic flow. Analysis with high ionic strength buffers 589(1992)315

Capillary isoelectric focusing with electroosmotic zone displacement and on-column multichannel detection 589(1992)321

Finite difference modelling of continuous-flow electrophoresis 594(1992)341

Electroosmotic mobility

Continuous-flow electrophoresis: a separation criterion applied to the separation of model proteins 594(1992)351

Electrophoresis

see also Special issue index

Electrophoresis

see also Capillary electrophoresis

Thin-layer electrophoresis of hydroxyethyl starches on a modified silica gel support 591(1992)351

Finite difference modelling of continuous-flow electrophoresis 594(1992)341

Important design features of a system for the densitometric analysis of two-dimensional flat-bed separations 596(1992)241

Analysis of the interaction between an $\alpha(1 \rightarrow 6)$ dextran-specific mouse hybridoma antibody and dextran B512 by affinity electrophoresis 597(1992)345

Characterization of the interaction between human plasma fibronectin and collagen by means of affinity electrophoresis 597(1992)351

Separation of monoclonal antibodies from antihapten antisera by two-dimensional affinity electrophoresis 597(1992)365

Characterization of the heterogeneity of polyethylene glycol-modified superoxide dismutase by chromatographic and electrophoretic techniques 599(1992)141

Electrophoretic migration

Studies of the electrophoretic behaviour of *p*-dihydroxyborylphenylalanine and related compounds by the three-spot method 585(1991)369

Electrophoretic mobility

Influence of operating parameters on reproducibility in capillary electrophoresis 559(1991)57

Electrophoretic mobility modeling of proteins in free zone capillary electrophoresis and its application to monoclonal antibody microheterogeneity analysis 559(1991)357

Determination of some drugs by micellar electrokinetic capillary chromatography. The pseudo-effective mobility as parameter for screening 585(1991)123

Electrophoretic mobility and dissociation constants of tripeptides evaluated by isotachopheresis 585(1991)364

Quantitative aspects of indirect UV detection in capillary zone electrophoresis 588(1991)321

Continuous-flow electrophoresis: a separation criterion applied to the separation of model proteins 594(1992)351

Peak dispersion and separation efficiency in high-performance zone electrophoresis with gel-filled capillaries 595(1992)313

Electrospray ionization

Recent advances in liquid chromatography–mass spectrometry and capillary zone electrophoresis–mass spectrometry for protein analysis 553(1991)101

Electrospray interfacing for the coupling of ion-exchange and ion-pairing chromatography to mass spectrometry 554(1991)27

Liquid chromatography–mass spectrometry with ionspray and electrospray interfaces in pharmaceutical and biomedical research 554(1991)39

Nanoscale separations. Capillary liquid chromatography–mass spectrometry and capillary zone electrophoresis–mass spectrometry for the determination of peptides and proteins 554(1991)329

Electrostatic interactions

Carbohydrate analysis with ion chromatography using Eurokat stationary phases. Preparative separation of monosaccharides and their fluorinated derivatives 552(1991)483

Analysis of pH-dependent protein interactions with gel filtration medium 591(1992)121

Electrostatic retention model

Extension of the electrostatic retention model of reversed-phase ion-pair high-performance liquid chromatography to include the effect of the eluent pH 552(1991)13

Elemental analysis

Physical and chemical characterization of a porous phosphate-modified zirconia substrate 587(1991)137

Eluent composition

Reversed-phase liquid chromatography coupled on-line with capillary gas chromatography. I. Introduction of large volumes of aqueous mixtures through an on-column interface 552(1991)489

Eluotropic scale

Ion- and ligand-exchange chromatography of proteins using porous zirconium oxide supports in organic and inorganic Lewis base eluents 596(1992)27

Elution behaviour

Purification and crystallization of yeast hexokinase isoenzymes. Characterization of different forms by chromatofocusing 587(1991)85

Elution of characteristic points

Experimental determination of adsorption isotherm data for the study of the surface energy distribution of various solid surfaces by inverse gas-solid chromatography 591(1992)233

Precision and accuracy of the gas-solid adsorption isotherms derived by the elution-by-characteristic-points method 591(1992)245

Validity of the model used to relate the energy distribution and the adsorption isotherm 591(1992)267

Elution order reversal

Reversal of elution order during direct enantiomeric separation of pyriproxyfen on a cellulose-based chiral stationary phase 588(1991)177

Elution profiles

Prediction of the elution profiles of high-concentration bands in gas chromatography 589(1992)223

Glucose-fructose equilibria on Dowex Monosphere 99 CA resin under overloaded conditions 590(1992)147

Precision and accuracy of the gas-solid adsorption isotherms derived by the elution-by-characteristic-points method 591(1992)245

Elution volume

Improved accuracy in the determination of field-flow fractionation elution volumes 555(1991)260

Enantiomeric purity

Application of preparative liquid chromatography to the isolation of enantiomers of a benzodiazepinone derivative 590(1992)127

(2*S*,3*S*)-Dicyclohexyl tartrate as mobile phase additive for the determination of the enantiomeric purity of (*S*)-atropine in tablets 592(1992)339

Enantiomer separation

Theoretical approach to the gas chromatographic separation of enantiomers on dissolved cyclodextrin derivatives 552(1991)43

Investigation of cross-linking chiral stationary phases within capillary columns 552(1991)153

Practical strategy for the analytical separation of enantiomers by high-performance liquid chromatography 553(1991)357

Influence of chemical structure of tricyclic tertiary dimethylamines on chiral separation by reversed-phase high-performance liquid chromatography after derivatization with (–)-menthylchloroformate 553(1991)365

Reversed-phase and chiral high-performance liquid chromatographic assay of bupivacaine and its enantiomers in clinical samples after continuous extrapleural infusion 553(1991)383

Liquid chromatographic separation of the enantiomers of metoprolol and its α -hydroxy metabolite on Chiralcel OD for determination in plasma and urine 553(1991)391

Problems and solutions in chiral thin-layer chromatography: a two-phase "Pirkle" modified amino-bonded plate 553(1991)497

Chiral separation by cyclodextrin-modified micellar electrokinetic chromatography 553(1991)503

Retention behavior and chiral recognition of β -cyclodextrin-derivative polymer adsorbed on silica for warfarin, structurally related compounds and Dns-amino acids 555(1991)53

Chiral π -donor stationary phases with (*R*)-*N*-pivaloylnaphthylethylamide groups for direct enantiomer separation by gas, liquid and supercritical fluid chromatography 557(1991)163

Chiral stationary phase designed for β -blockers 557(1991)173

Enantiomeric distribution and $^{13}\text{C}/^{12}\text{C}$ isotope ratio determination of γ -lactones: appropriate methods for the differentiation between natural and non-natural flavours? 557(1991)187

Coupled column chromatography in chiral separations: systems employing β -cyclodextrin phases for chiral separation 557(1991)199

Enantiomer separation by chiral-phase liquid chromatography of urethane derivatives of natural diacylglycerols previously fractionated by reversed-phase liquid chromatography 557(1991)227

- Unusual effect of temperature on the retention of enantiomers on a chiral column 558(1991)1
- Stereoisomeric purity determination of captopril by capillary gas chromatography 558(1991)257
- High-performance liquid chromatographic detection of enantiomeric amino alcohols after derivatization with *o*-phthaldialdehyde and various thiosugars 558(1991)311
- Optimization of the direct enantiomeric separation by high-performance liquid chromatography of the D-2 dopamine agonist N-0437 using response surface methodology and the multi-criteria decision-making method 558(1991)333
- Chiral separations by micellar electrokinetic chromatography with sodium N-dodecanoyl-L-valinate 559(1991)209
- Selected applications of cyclodextrin selectors in capillary electrophoresis 559(1991)215
- Preparative chromatographic separation of enantiomers (Review) 576(1992)1
- (2*R*,3*R*)-Dicyclohexyl tartrate as a chiral mobile phase additive 585(1991)35
- Separation of secondary alcohol enantiomers using supercritical fluid chromatography 585(1991)117
- New chiral covalently bonded, π -donor stationary phases for high-performance liquid chromatography, based on derivatives of optically active 1-(1-naphthyl)ethylamine. Part II 585(1991)195
- Ligand-exchange chromatography of alkenes on stationary phases containing palladium(II) complexes. Enantiomeric separation of *trans*-1,2-divinylcyclohexane 585(1991)309
- Ligand-exchange ion chromatographic determination of malic acid enantiomers in apple juice with photometric detection 585(1991)315
- Direct separation of the enantiomers of propafenone, diprafenone and their major metabolites by high-performance liquid chromatography on modified cellulose and amylose chiral stationary phases 585(1991)329
- Rapid optical resolution of electrically neutral cobalt(III) chelate complexes by gel permeation chromatography 585(1991)337
- Enantiomeric separation of α -phenylethylamine and its substituted isomers by gas chromatography 586(1991)139
- Separation of enantiomers using cellulase (CBH I) silica as a chiral stationary phase 586(1991)233
- Chiroptical detection during liquid chromatography. III. Non-stop acquisition of circular dichroism spectra during liquid chromatography 586(1991)249
- Preparative separation and analysis of the enantiomers of [³H]Abbott-69992, an HIV anti-infective nucleoside, by ligand-exchange high-performance liquid chromatography 586(1991)265
- Direct separation of nadolol enantiomers on a Pirkle-type chiral stationary phase 586(1991)351
- Simultaneous stereoisomer analysis of 2-alkyl-branched acids, esters and alcohols using a selectivity-adjusted column system in multi-dimensional gas chromatography 587(1991)347
- Reversal of elution order during direct enantiomeric separation of pyriproxyfen on a cellulose-based chiral stationary phase 588(1991)177
- High-performance liquid chromatographic analysis of diastereomers and enantiomers of pyrrolisoquinoline antidepressants 588(1991)348
- Enantiomeric separation of racemic hydroperoxides and related alcohols 588(1991)352
- Silica-bonded chiral stationary phases with structurally simple π -donor chiral selectors for high-performance liquid chromatography 589(1992)53
- Gas chromatographic separation of diastereomeric amino acid derivatives on chiral stationary phases. Application to the determination of enantiomeric composition in (*S*)-(+)-2-butanol 589(1992)249
- Application and chiral recognition of heptakis (2,6-di-*O*-methyl-3-*O*-trifluoroacetyl)- β -cyclodextrin as a stationary phase for the gas chromatographic separation of enantiomers 589(1992)257
- Optical resolution by simulated moving-bed adsorption technology 590(1992)113
- Experimental study of the production rate of pure enantiomers from racemic mixtures 590(1992)119
- Application of preparative liquid chromatography to the isolation of enantiomers of a benzodiazepinone derivative 590(1992)127
- Determination of the enantiomeric composition of samples of cocaine by normal-phase high-performance liquid chromatography with UV detection 590(1992)354
- Retention processes on α_1 -acid glycoprotein-bonded stationary phase 591(1992)55

- Direct liquid chromatographic separation of enantiomers on immobilized protein stationary phases. IX. Influence of the cross-linking reagent on the retentive and enantioselective properties of chiral sorbents based on bovine serum albumin 591(1992)65
- Direct separation of enantiomers by high-performance liquid chromatography on a new chiral ligand-exchange phase 592(1992)291
- Chiral high-performance liquid chromatographic separations on an α_1 -acid glycoprotein column. II. Separation of the diastereomeric and enantiomeric analogues of vinpocetine (Cavinton) 592(1992)297
- Retention and enantioselectivity of racemic solutes on a modified ovomucoid-bonded column. I. Cross-linking with glutaraldehyde 592(1992)301
- Investigations of the influence of silanol groups on the separation of enantiomers by liquid and supercritical fluid chromatography 592(1992)309
- Determination of the enantiomers of fenoldopam in human plasma by reversed-phase high-performance liquid chromatography after chiral derivatization 592(1992)317
- Reversed retention order and other stereoselective effects in the separation of amino alcohols on Chiralcel OD 592(1992)331
- (2*S*,3*S*)-Dicyclohexyl tartrate as mobile phase additive for the determination of the enantiomeric purity of (*S*)-atropine in tablets 592(1992)339
- High-performance ligand-exchange chromatography of some amino acids containing two chiral centres 592(1992)345
- Direct high-performance liquid chromatographic resolution of a novel benzothiazine Ca^{2+} antagonist and related compounds 593(1992)37
- Chromatographic resolution on methylbenzoylcellulose beads. Modulation of the chiral recognition by variation of the position of the methyl group on the aromatic ring 595(1992)63
- Stationary phase effect on enantioselectivity of dansyl phenylalanine in microcolumn liquid chromatography with γ -cyclodextrin as mobile phase additive 595(1992)121
- Direct high-performance liquid chromatographic separation of enantiomeric peptidoleukotriene antagonists 596(1992)123
- Investigation of enantioselectivity and enantiomeric elution order of propranolol and its ester derivatives on an ovomucoid-bonded column 598(1992)67
- Separation of the enantiomers of the 3,5-dinitrobenzamide derivatives of α -amino phosphonates on four chiral stationary phases 598(1992)159
- Some novel homochiral derivatizing agents for the gas chromatographic analysis of enantiomeric secondary alcohols 598(1992)251
- Contribution of ionically immobilized bovine serum albumin to the retention of enantiomers 600(1992)37
- High-performance ligand-exchange liquid chromatography of fluoro derivatives of alanine 600(1992)83
- Suitable chiral packing material for the high-performance liquid chromatographic separation of derivatives of 1'-hydroxyeugenol 600(1992)312
- High-performance liquid chromatographic separation of chiral metallocenic ketones and alcohols 600(1992)320
- Enantioselectivity**
Investigation of cross-linking chiral stationary phases within capillary columns 552(1991)153
- (*S,S*)-Diphenylethylethanediamine derivatives as chiral selectors. II. Gasparrini-type bound chiral stationary phase with high enantioselectivity for naphthylamides 553(1991)373
- Chiral separation by cyclodextrin-modified micellar electrokinetic chromatography 553(1991)503
- (2*R*,3*R*)-Dicyclohexyl tartrate as a chiral mobile phase additive 585(1991)35
- New chiral covalently bonded, π -donor stationary phases for high-performance liquid chromatography, based on derivatives of optically active 1-(1-naphthyl)ethylamine. Part II 585(1991)195
- Direct liquid chromatographic separation of enantiomers on immobilized protein stationary phases. IX. Influence of the cross-linking reagent on the retentive and enantioselective properties of chiral sorbents based on bovine serum albumin 591(1992)65
- Retention and enantioselectivity of racemic solutes on a modified ovomucoid-bonded column. I. Cross-linking with glutaraldehyde 592(1992)301
- Investigations of the influence of silanol groups on the separation of enantiomers by liquid and supercritical fluid chromatography 592(1992)309
- Synthesis of (1*R*-*trans*)-N,N'-1,2-cyclohexylenebisbenzamideoligodimethylsiloxane copolymers for use as chiral stationary phases for capillary supercritical fluid chromatography 594(1992)283
- Separation of the enantiomers of the 3,5-dinitrobenzamide derivatives of α -amino phosphonates on four chiral stationary phases 598(1992)159

Energy distribution

Validity of the model used to relate the energy distribution and the adsorption isotherm 591(1992)267

Enthalpy changes

Effect of temperature and mobile phase on the retention of retinoates in reversed-phase liquid chromatography 592(1992)127

Entropy-enthalpy compensation

Physico-chemical properties of electron-acceptor stationary phases in liquid chromatography 591(1992)75

Environmental monitoring

Gas chromatographic elution patterns of chlorinated dioxins versus column polarity 594(1992)269

Enzymatic transfer

Separation of functionalized dextrans by reversed-phase high-performance liquid chromatography 587(1991)288

Enzyme activities

Measurement of enzyme activities of cytochrome P-450 isoenzymes by high-performance liquid chromatographic analysis of products 553(1991)233

Enzyme-antibody conjugates

High-performance capillary electrophoresis as a fast in-process control method for enzyme-labelled monoclonal antibody conjugates 559(1991)385

Enzyme-linked immunosorbent assay

Immobilization of ultra-thin layer of monoclonal antibody on glass surface 566(1991)361

Enzyme reactors

Characterization of cellulase-based enzyme reactors for the high-performance liquid chromatographic determination of β -D-glucan oligosaccharides 558(1991)343

Enzymophoresis of nucleic acids by tandem capillary enzyme reactor-capillary zone electrophoresis 596(1992)251

Epifluorescence microscopy

DNA electrophoresis in uncross-linked polyacrylamide solution, studied by epifluorescence microscopy 596(1992)141

Epilobium

Isolation and determination of flavonol glycosides from *Epilobium* species 557(1991)391

Epitope mapping

Mapping of viral epitopes with conformationally specific monoclonal antibodies using biosensor technology 597(1992)391

Epoxy resins, cured

Quality control of cured epoxy resins. Determination of residual free monomers (*m*-xylylenediamine and bisphenol A diglycidyl ether) in the finished product 585(1991)75

Epstein-Barr virus

Purification and quantification of recombinant Epstein-Barr viral glycoproteins gp350/220 from Chinese hamster ovary cells 599(1992)267

Equilibration times

Gradient elution with shorter equilibration times in reversed-phase ion-pair chromatography 592(1992)143

Equilibrium concentration

Methods of equilibrium concentration for the gas chromatographic determination of trace volatiles (Review) 556(1991)1

Equilibrium constants

Ion-exclusion chromatography using mobile phases containing β -cyclodextrin 595(1992)149

Equilibrium interaction constants

Determination of solute-polymer interaction constants via chromatography with polymer capillaries 594(1992)29

Equilibrium parameters

Predictability of chromatographic protein separations. Study of size-exclusion media with narrow particle size distributions 589(1992)71

Erythrina variegata

Isolation and characterization of isolectins from *Erythrina variegata* seeds 597(1992)207

Erythrocyte membranes

Simultaneous determination of amounts of major phospholipid classes and their fatty acid composition in erythrocyte membranes using high-performance liquid chromatography and gas chromatography 598(1992)33

Eschscholtzia californica

Analytical and quantitative studies of californin and protopin in aerial part extracts of *Eschscholtzia californica* Cham. with high-performance liquid chromatography 587(1991)314

Eucalyptus grandis

Plant growth regulators G₁, G₂, G₃. Synthesis, extraction and determination of leaf content in *Eucalyptus grandis* 600(1992)323

Euclidean distance

Use of oxidative degradation followed by capillary gas chromatography-mass spectrometry and multi-dimensional scaling analysis to fingerprint unresolved complex mixtures of hydrocarbons 589(1992)281

Eupergit C-melittin

Calcium-modulated conformational affinity chromatography. Application to the purification of calmodulin and S100 proteins 597(1992)263

Eupergit C 30N

Activation of matrices by 4,6-diphenylthieno[3,4-*d*]-1,3-dioxol-2-one-5,5-dioxide. High-performance liquid affinity chromatographic separations 589(1992)101

Evaporative light-scattering detector

High-performance liquid chromatography and supercritical fluid chromatography of monosaccharides and polyols using light-scattering detection. Chemometric studies of the retentions 590(1992)203

Comparison of ethoxylated alcohols and polyethylene glycols by high-performance liquid chromatography and supercritical fluid chromatography using evaporative light-scattering detection 591(1992)149

Exclusion limit

Bead cellulose derivatives as supports for immobilization and chromatographic purification of proteins 552(1991)389

Expanded beds

Affinity purification of proteins using expanded beds 597(1992)129

Expert systems

Feasibility study for the construction of an integrated expert system in high-performance liquid chromatography 589(1992)31

CRISEBOOK, a Hypermedia version of an expert system for the selection of optimization criteria in high-performance liquid chromatography 592(1992)51

Exponential force-field programming

Asymmetric-channel flow field-flow fractionation with exponential force-field programming 593(1992)339

Exponentially modified Gaussian peak model

Improved algorithm for resolution of overlapped asymmetric chromatographic peaks 591(1992)19

Extinction coefficients

Analysis of ϵ -caprolactam and its cyclic oligomers by high-performance liquid chromatography 585(1991)333

Extra-column dispersion

Dispersion in round tubes and its implications for extra-column dispersion 556(1991)235

Extraction cell

Effects of collection solvent parameters and extraction cell geometry on supercritical fluid extraction efficiencies 594(1992)297

Extraction eluents

Optimization of supercritical fluid extraction of volatile constituents from a model plant matrix 600(1992)175

Extraction methods

Determination of Cyasorb UV 1084 and its degradation products in low-density polyethylene film by high-performance liquid chromatography 587(1991)205

Comparison of extraction methods for polychlorinated dibenzo-*p*-dioxins and dibenzofurans in fly ash using gas chromatography-mass spectrometry 589(1992)265

Improvement of extraction procedure for biogenic amines in foods and their high-performance liquid chromatographic determination 591(1992)175

Quantitation of mutagenic/carcinogenic heterocyclic aromatic amines in food products 592(1992)271

Determination of cinnamic acid and paeoniflorin in traditional chinese medicinal preparations by high-performance liquid chromatography 593(1992)191

Analysis and isolation of indole alkaloids of fungi by high-performance liquid chromatography 593(1992)201

Improved high-performance liquid chromatographic method for the determination of coenzyme Q₁₀ in plasma 593(1992)217

Selection of robust combinations of extraction liquid composition and internal standard. Monte Carlo simulation of improvement of assay methods with liquid-liquid extraction prior to high-performance liquid chromatography 594(1992)45

Solvent degradation of cloxacillin *in vitro*. Tentative identification of degradation products using thermospray liquid chromatography-mass spectrometry 594(1992)195

Supercritical carbon dioxide extraction of resin and fatty acids from sediments at pulp mill sites 594(1992)309

Quantitative analysis of total resin acids by high-performance liquid chromatography of their coumarin ester derivatives 595(1992)155

Comparison of detergents for extraction and ion-exchange high-performance liquid chromatography of Sendai virus membrane proteins 599(1992)157

Purification of DNA-derived deoxynucleotides from leukocytes involving nuclease elution of an ion-exchange column 600(1992)229

F**Factor analysis**

Factor analysis and experimental design in high-performance liquid chromatography. XI. Factor analysis maps and chromatographic information 556(1991)169

Factorial design

Computer-aided optimization of the experimental conditions for the isocratic reversed-phase high-performance liquid chromatographic separation of hormonal steroids 552(1991)103

Retention processes on α_1 -acid glycoprotein-bonded stationary phase 591(1992)55

Fast atom bombardment

Source of band broadening in liquid chromatographic-fast atom bombardment mass spectrometric systems with precolumn addition of viscous matrix to the mobile phase 554(1991)47

Effect of the addition of viscous matrices to the mobile phase on chromatographic performance in liquid chromatography-fast atom bombardment mass spectrometry 554(1991)61

Limits and effects of precolumn addition of thioglycerol in liquid chromatographic-fast atom bombardment mass spectrometric systems 591(1992)129

Fast atom bombardment, continuous-flow

Nanoscale separations combined with tandem mass spectrometry 554(1991)73

Approach to studying proteinase specificity by continuous-flow fast atom bombardment mass spectrometry and high-performance liquid chromatography combined with photodiode-array ultraviolet detection 554(1991)83

Application of microcolumn liquid chromatography-continuous-flow fast atom bombardment mass spectrometry in environmental studies of sulfonylurea herbicides 554(1991)91

Nanoscale separations. Capillary liquid chromatography-mass spectrometry and capillary zone electrophoresis-mass spectrometry for the determination of peptides and proteins 554(1991)329

Pseudo-electrochromatography-mass spectrometry: a new alternative 554(1991)339

Applicability of high-performance liquid chromatography-continuous-flow fast atom bombardment mass spectrometry for simultaneous quantitation of multiple neurochemicals 587(1991)306

Fast atom bombardment mapping

Structural analysis of biologically active peptides and recombinant proteins and their modified counterparts by mass spectrometry 562(1991)377

Feedstocks, particulate-containing

Affinity purification of proteins using expanded beds 597(1992)129

Fermentation brines

Analysis of zapatera olives by gas and high-performance liquid chromatography 594(1992)259

Fermentation media

Determination of glutamine, glutamic acid and pyroglutamic acids using high-performance liquid chromatography on dynamically modified silica 594(1992)173

Fast protein monitoring in fermentation broth using non-porous micropellicular reversed-phase columns 594(1992)392

Ferredoxin affinity adsorbents

Immobilized ferredoxins for affinity chromatography of ferredoxin-dependent enzymes 597(1992)147

Field-amplified sample injection

Field amplified sample injection in high-performance capillary electrophoresis 559(1991)141

Field-amplified polarity-switching sample injection in high-performance capillary electrophoresis 559(1991)153

Field effect electroosmosis

Field effect electroosmosis 559(1991)95

Field-flow fractionation

Non-equilibrium integral Doppler anemometric analysis of particle mixtures in a channel flow using an intrinsic hydrodynamic focusing force biased by another force 553(1991)517

Improved accuracy in the determination of field-flow fractionation elution volumes 555(1991)260

Separation of living red blood cells by gravitational field-flow fractionation 568(1991)93

Investigation of red blood cell fractionation by gravitational field-flow fractionation 583(1992)27

Asymmetric-channel flow field-flow fractionation with exponential force-field programming 593(1992)339

Simple solution of velocity profiles of laminar flows in channels of various cross-sections used in field-flow fractionation 593(1992)357

Integral Doppler anemometry in porous membranes for the analysis of liquid mixtures and examination of membrane properties 600(1992)129

Transverse particle redistribution in a flat channel for SPLITT or integral Doppler anemometry 600(1992)133

Use of integral Doppler anemometry in field-flow fractionation 600(1992)139

Isoelectric focusing field-flow fractionation. III. Investigation of the influence of different experimental parameters on focusing of cytochrome *c* in the trapezoidal cross-section channel 600(1992)297

Isoelectric focusing field-flow fractionation. IV. Investigations on protein separations in the trapezoidal cross-section channel 600(1992)305

Film thickness

Production and use of capillary traps for headspace gas chromatography of airborne volatile organic compounds 552(1991)137

Fish oils

Isolation and quantification of polymers from autoxidized fish oils by high-performance size-exclusion chromatography with an evaporative mass detector 552(1991)501

Flame infrared emission detector

Flame infrared emission detection of hydrogen 589(1992)241

Flame photometric detector

Flame-photometric detection of nitrous oxide in addition to phosphine 598(1992)313

Flavour analysis

Enantiomeric distribution and $^{13}\text{C}/^{12}\text{C}$ isotope ratio determination of γ -lactones: appropriate methods for the differentiation between natural and non-natural flavours? 557(1991)187

Flip-flop elution

Flip-flop elution concept in preparative liquid chromatography 557(1991)137

Flory-Huggins interaction parameters

Activity coefficients at infinite dilution determined by gas-liquid chromatography. Organic solvents in Apiezon L 586(1991)303

Flow cells

Optimization and evaluation of the performance of arrangements for UV detection in high-resolution separations using fused-silica capillaries 559(1991)163

Flow-injection analysis

Use of various types of column reactors for flow-injection analysis 597(1992)435

New approaches to coupling flow-injection analysis and high-performance liquid chromatography 600(1992)183

Flow-injection immunoassay

Process monitoring by flow-injection immunoassay. Evaluation of a sequential competitive binding assay 597(1992)383

Flow-injection manifold

New approaches to coupling flow-injection analysis and high-performance liquid chromatography 600(1992)183

Flow-meter

Determination of the hold-up time and column outlet density for capillary supercritical fluid chromatography 588(1991)289

Flow profiles

Simple test for determination of the degree of distortion of the liquid-phase flow profile in columns for preparative liquid chromatography 589(1992)61

Van Deemter-type relationship for determining the optimum initial flow-rate and optimum pressure programming rate in temperature/pressure-programmed capillary column gas chromatography utilizing separation numbers 595(1992)221

Flow-rate

Improved accuracy in the determination of field-flow fractionation elution volumes 555(1991)260

Optimisation pentaparamétrique de la résolution en chromatographie phase gazeuse par la technique du diagramme à fenêtres. Cas du couplage de deux colonnes 556(1991)287

Chromatographic developments in low flow delivery for liquid chromatography-mass spectrometry 562(1991)73

Membrane chromatographic systems for high-throughput protein separations 596(1992)173

Fluorescence detector

Selective post-column fluorogenic reaction with benzamidine for trace level detection of reducing saccharides in liquid chromatography 553(1991)255

Micro liquid chromatography with fluorescence detection of thiols and disulphides 553(1991)433

Fluorescence detector cell for use in an integrated electrically driven separation system 598(1992)115

Fluorescence detector, GC

Laser-excited fluorescence detection of gas-phase chromatography eluates 557(1991)39

Fluorescence spectroscopy

On-line coupling of liquid chromatography to thin-layer chromatography for the identification of polycyclic aromatic hydrocarbons in marine sediment by fluorescence excitation and emission spectroscopy 552(1991)613

Fluorescent stains

Use of hematoporphyrin as a fluorescent stain for detection of lipids in high-performance thin-layer chromatography 588(1991)295

Fluorimetric detectors

Use of hematoporphyrin as a fluorescent stain for detection of lipids in high-performance thin-layer chromatography 588(1991)295

Flux

Adsorption and elution of bovine γ -globulin using an affinity membrane containing hydrophobic amino acids as ligands 585(1991)45

Fly ash

Comparison of extraction methods for polychlorinated dibenzo-*p*-dioxins and dibenzofurans in fly ash using gas chromatography-mass spectrometry 589(1992)265

Forensic science

see also Special issue index

Fourier transform

Deconvolution method for accurate determination of overlapping peak areas in chromatograms 587(1991)129

Fourier transform infrared reflection

Immobilization of ultra-thin layer of monoclonal antibody on glass surface 566(1991)361

Fractionation methods

Use of basic alumina in fractionation of fossil fuels 600(1992)243

Frontal analysis

Comparison of various isotherm models for predicting competitive adsorption data 552(1991)71

Adsorption isotherms of phenylalanine in a chromatographic column measured simultaneously by system peaks analysis and frontal analysis 556(1991)277

Theory of perfusion chromatography 599(1992)87

Frontal chromatography

Capillary modification and evaluation using streaming potential and frontal chromatography for protein analysis in capillary electrophoresis 594(1992)325

Fruit juices

Determination of 5-hydroxymethylfurfural by ion-exclusion chromatography with UV detection 593(1992)153

FUMI

Simplex optimization of liquid chromatography with the function of mutual information as a criterion 585(1991)187

Functional endgroups

Analysis of benzyloxy-terminated poly(1,3,6-trioxocane)s by supercritical fluid chromatography 589(1992)295

Functionality constant

Prediction of retention indexes. III. Silylated derivatives of polar compounds 586(1991)113

Fused-silica capillaries

Evaluation of fundamental properties of a silica capillary used for capillary electrophoresis 559(1991)69

Free solution capillary electrophoresis of proteins using untreated fused-silica capillaries 559(1991)317

Capillary zone electrophoresis of proteins with hydrophilic fused-silica capillaries 559(1991)367

G

Galactosylation

Galactosylation as a tool for the stabilization and immobilization of proteins (Review) 597(1992)19

Gas chromatograph, automatic

Near-continuous measurement of hydrogen sulfide and carbonyl sulfide by an automatic gas chromatograph 585(1991)353

Gas extraction

Efficiency of gas extraction in headspace analysis 600(1992)251

Gas-liquid chromatography simulator

GCSIM: a gas-liquid chromatography simulator for educational purposes 588(1991)217

Gasoline

Quantitative hydrocarbon group analysis of gasoline and diesel fuel by supercritical fluid chromatography 595(1992)301

Gas-phase compressibility

Prediction of the elution profiles of high-concentration bands in gas chromatography 589(1992)223

Gas-solid reactions

Theoretical analysis for measurement of building pollution parameters by gas chromatography 557(1991)469

Gaussian function

Deconvolution method for accurate determination of overlapping peak areas in chromatograms 587(1991)129

GCSIM

GCSIM: a gas-liquid chromatography simulator for educational purposes 588(1991)217

Gel electrophoresis

Theoretical background for clinical and biomedical applications of electromigration techniques (Review) 569(1991)3

Gel filtration

Analysis of pH-dependent protein interactions with gel filtration medium 591(1992)121

Geometrical isomerism

Simultaneous effect of geometrical isomerism and chelate ring size of tris(aminocarboxylato)cobalt(III) complexes on their behaviour in thin-layer chromatography on silica gel and alumina 585(1991)181

Geometric isomers

Improved high-performance liquid chromatographic resolution of the geometric isomers of 6-hydroxy-4-(1-hydroxy-1-methylethyl)-1-cyclohexene-1-ethanol and by-products with β -cyclodextrin 553(1991)15

High-performance liquid chromatography of peptides at reduced temperatures: separation of isomers 586(1991)145

Ginkgo biloba

Gas chromatographic-mass spectrometric analysis of ginkgolides produced by *Ginkgo biloba* cell culture 588(1991)281

Glycine max

Separation of the complete thioredoxin pattern of soybean leaves (*Glycine max*) by high-performance anion-exchange chromatography on Mono Q 587(1991)109

Gradient elution

Analysis of polyethylene glycol and derivatives by high-performance liquid chromatography using elevated temperatures and low-wavelength ultraviolet detection, and supercritical fluid chromatography 553(1991)423

Computer simulation of gradient elution separation. Accuracy of predictions for non-linear gradients 555(1991)1

Reversed-phase separations of high-molecular-weight polystyrenes in a dichloromethane-acetonitrile solvent system 558(1991)440

Separation of some platinum(II) complexes by ionic strength gradient on a solvent-generated ion-exchange sorbent 586(1991)291

Factors affecting the separation and loading capacity of proteins in preparative gradient elution high-performance liquid chromatography 590(1992)35

Interfacing gradient elution-ion-exchange chromatography and low-angle laser light-scattering photometry for analysis of proteins 591(1992)139

Application of the gradient elution technique. Demonstration with a special test mixture and the DryLab G/plus method development software 592(1992)133

Gradient elution with shorter equilibration times in reversed-phase ion-pair chromatography 592(1992)143

Chromatography of 99 amino acids and other ninhydrin-reactive compounds in the Pickering lithium gradient system 594(1992)165

Preparative liquid chromatography. II. Existence of optimum injection conditions for overloaded gradient elution separations 596(1992)15

Optimization of the separation of the Rp and Sp diastereomers of phosphate-methylated DNA and RNA dinucleotides 596(1992)59

Studies on stress metabolites. XVI. High-performance liquid chromatographic analysis of cruciferous phytoalexins using complex ternary mobile phase gradients 598(1992)147

Application of capillary reversed-phase high-performance liquid chromatography to high-sensitivity protein sequence analysis 599(1992)119

Influence of operating parameters on the preparative gradient elution chromatography of insulins 599(1992)195

Gradient parameters

Displacement effects in preparative gradient high-performance liquid chromatographic separations 590(1992)17

Gradient slope

Influence of operating parameters on the preparative gradient elution chromatography of insulins 599(1992)195

Grapes

Analytical methods for monoterpene glycosides in grape and wine. I. XAD-2 extraction and gas chromatographic-mass spectrometric determination of synthetic glycosides 590(1992)313

Graphitized carbon

Retention behaviour of some ring-substituted phenol derivatives on a porous graphitized carbon column 592(1992)75

Graphitized carbon black

Gas-liquid microcapillary columns precoated with graphitized carbon black 552(1991)197

Chromatographic properties of graphitized thermal carbon black modified with a monolayer of liquid crystal 600(1992)59

Grappa

Gas chromatographic-mass spectrometric determination of ethyl carbamate as the xanthylamide derivative in Italian aqua vitae (grappa) samples 585(1991)111

Gravitational field-flow fractionation

Investigation of red blood cell fractionation by gravitational field-flow fractionation 583(1992)27

Group-type separations

Quantitative structure-type analysis of heterocompound-hydrocarbon mixtures. VI. Characterization of the nature of neutral heterocompounds in a product of brown coal refining 553(1991)441

Group resolution parameter

Physico-chemical properties of electron-acceptor stationary phases in liquid chromatography 591(1992)75

Group retention contribution

Correlations between high-performance liquid chromatographic retention, X-ray structural and ^{13}C NMR spectroscopic data of flavonoid compounds 592(1992)65

Group retention factor

Prediction of retention indexes. II. Structure-retention index relationship on polar columns 586(1991)85

Prediction of retention indexes. III. Silylated derivatives of polar compounds 586(1991)113

H***Haemophilus aphrophilus***

Intra-injector methylation of free fatty acids from aerobically and anaerobically cultured *Actinobacillus actinomycetemcomitans* and *Haemophilus aphrophilus* 576(1992)328

Hair follicles

Electrochemical determination of femtomole amounts of free reduced and oxidized glutathione. Application to human hair follicles 576(1992)255

Headspace analysis

Production and use of capillary traps for headspace gas chromatography of airborne volatile organic compounds 552(1991)137

Modern approach to the quantitative determination of volatiles in solid samples. Multiple headspace extraction gas chromatography for the determination of cyclohexanone residues in soil 552(1991)205

Determination of volatile thiols by gas chromatography using separation as tributyltin mercaptides 555(1991)306

Methods of equilibrium concentration for the gas chromatographic determination of trace volatiles (Review) 556(1991)1

Analysis of industrial solvent mixtures in water using a miniature purge-and-trap device with thermal desorption and capillary gas chromatography-mass spectrometry 557(1991)429

Evaluation of gas chromatographic columns for the determination of methylmercury in aqueous head space extracts from biological samples 586(1991)329

Determination of parts per million levels of trifluoroacetic acid in ceronapril bulk substance by headspace capillary gas chromatography 596(1992)85

Efficiency of gas extraction in headspace analysis 600(1992)251

Heat of adsorption

Estimation of cellulose polarity by gas chromatography 552(1991)365

Study of microporous carbons by gas chromatographic determination of heats of physisorption 555(1991)183

Heat transfer coefficients

Temperature elevations of the sample zone in free solution capillary electrophoresis under stacking conditions 559(1991)27

Height equivalent to a theoretical plate

Importance of intraparticle convection in the performance of chromatographic processes 590(1992)93

Helium ionization detector

Determination of sulphur and nitrogen gases by gas chromatography on polystyrene porous polymer columns 585(1991)93

 α -Helix

Retention behaviour of a template-assembled synthetic protein and its amphiphilic building blocks on reversed-phase columns 586(1991)43

Heparin-glyceryl controlled-pore glass

Preparation of heparin-glyceryl controlled-pore glass affinity media for the separation of α - and β -lipoproteins 573(1992)132

High-performance liquid chromatography, coupling to FIA

New approaches to coupling flow-injection analysis and high-performance liquid chromatography 600(1992)183

High-speed gas chromatography

High-speed gas chromatography. Theoretical and practical aspects 556(1991)331

High-temperature counter-current GC

High-temperature continuous counter-current gas-liquid chromatography 590(1992)289

Hold-up time

Determination of the hold-up time and column outlet density for capillary supercritical fluid chromatography 588(1991)289

Hollow-fibre membranes

Adsorption and elution of bovine γ -globulin using an affinity membrane containing hydrophobic amino acids as ligands 585(1991)45

Protein adsorption capacity of a porous phenylalanine-containing membrane based on a polyethylene matrix 586(1991)27

Membrane-based receptor affinity chromatography 597(1992)155

Kinetic aspects of membrane-based immunoaffinity chromatography 597(1992)167

Hollow-fibre sample preparation device

Membrane-based sample preparation device for the pretreatment of acidic samples prior to cation analysis by ion chromatography 586(1991)283

Homologous series

Description of retention mechanism by solvophobic theory. Influence of organic modifiers on the retention behaviour of homologous series in reversed-phase liquid chromatography 556(1991)219

Determination of the hold-up time and column outlet density for capillary supercritical fluid chromatography 588(1991)289

Thin-layer chromatography on polyacrylonitrile. IV. Investigation of the separation mechanisms for tris-(alkylxanthato)cobalt(III) complexes 596(1992)91

Honey

Determination of 5-hydroxymethylfurfural by ion-exclusion chromatography with UV detection 593(1992)153

Human immunodeficiency virus

Cyclooxygenase and lipoxigenase arachidonic acid metabolism by monocytes from human immune deficiency virus-infected drug users 557(1991)507

Analysis of DNA restriction fragments and polymerase chain reaction products towards detection of the AIDS (HIV-1) virus in blood 559(1991)267

Hybrid hybridoma

Selective high-performance liquid chromatographic purification of bispecific monoclonal antibodies 599(1992)13

Hybridization

Identification of DNA molecules by pre-column hybridization using capillary electrophoresis 559(1991)295

Hybridoma supernatant

Development and optimization of a single-step procedure using protein A affinity chromatography to isolate murine IgG₁ monoclonal antibodies from hybridoma supernatants 587(1991)61

Hydrodynamic effect

Diffusion of proteins in the chromatographic gel AcA-34 558(1991)71

Hydrodynamic focusing

Non-equilibrium integral Doppler anemometric analysis of particle mixtures in a channel flow using an intrinsic hydrodynamic focusing force biased by another force 553(1991)517

Hydrodynamic voltammograms

Determination of picomole amounts of lipoxins C₄, D₄ and E₄ by high-performance liquid chromatography with electrochemical detection 585(1991)53

Constant-potential amperometric detection of carbohydrates at metal electrodes in high-performance anion-exchange chromatography 592(1992)229

Hydrogen bond donor acidity

Solvatochromic hydrogen bond donor acidity of cyclodextrins and reversed-phase liquid chromatographic retention of small molecules on a β -cyclodextrin-bonded silica stationary phase 595(1992)45

Hydrogen bonding

Carbohydrate analysis with ion chromatography using Eurokat stationary phases. Preparative separation of monosaccharides and their fluorinated derivatives 552(1991)483

Hydrogen bonding. XVI. A new solute solvation parameter, π^H_2 , from gas chromatographic data 587(1991)213

Hydrogen bonding. XVII. The characterisation of 24 gas-liquid chromatographic stationary phases studied by Poole and co-workers, including molten salts, and evaluation of solute-stationary phase interactions 587(1991)229

Hydrogen bonding. XIX. The characterisation of two poly(methylphenylsiloxane)s 588(1991)361

Correlations between high-performance liquid chromatographic retention, X-ray structural and ¹³C NMR spectroscopic data of flavonoid compounds 592(1992)65

Hydrogen bonding. XXI. Solvation parameters for alkylaromatic hydrocarbons from gas-liquid chromatographic data 594(1992)229

Preparation and capillary column gas chromatographic characterization of aza-crown ether polysiloxane stationary phases 594(1992)243

Hydrolyzed vegetable protein

Determination of 3-chloropropane-1,2-diol in liquid hydrolysed vegetable proteins by capillary gas chromatography with flame ionization detection 555(1991)311

Hydrophilic interactions

Effect of stationary phase structure on retention and selectivity of restricted-access reversed-phase packing materials 558(1991)19

Mixed-mode hydrophilic and ionic interaction chromatography rivals reversed-phase liquid chromatography for the separation of peptides 594(1992)75

Hydrophilicity

Mixed-mode hydrophilic and ionic interaction chromatography rivals reversed-phase liquid chromatography for the separation of peptides 594(1992)75

Hydrophobic interactions

Effect of stationary phase structure on retention and selectivity of restricted-access reversed-phase packing materials 558(1991)19

Purification and crystallization of yeast hexokinase isoenzymes. Characterization of different forms by chromatofocusing 587(1991)85

Study of the retention behaviour of barbiturates by overpressured layer chromatography using silica gel bonded with tricaprilmethylammonium chloride 589(1992)301

Analysis of pH-dependent protein interactions with gel filtration medium 591(1992)121

Thin-layer chromatography on polyacrylonitrile. IV. Investigation of the separation mechanisms for tris-(alkylxanthato)cobalt(III) complexes 596(1992)91

Contribution of specific hydrophobic interactions to chiral recognition 598(1992)1

High-performance liquid chromatographic separation of *cis-trans* isomers of proline-containing peptides. I. Separation on cyclodextrin-bonded silica 598(1992)139

Characterization of hydrophobic interaction and hydrophobic interaction chromatography media by multivariate analysis 599(1992)131

Hydrophobicity

see also Lipophilicity

Calculation of retention and selectivity in reversed-phase liquid chromatography 552(1991)91

Correlation analysis in liquid chromatography of metal chelates. II. One-dimensional retention-mobile phase composition (physicochemical property) models in reversed-phase liquid chromatography 555(1991)163

Hydrophobicity of β -lactam antibiotics. Explanation and prediction of their behaviour in various partitioning solvent systems and reversed-phase chromatography 585(1991)3

Predicting partition coefficients in polyethylene glycol-potassium phosphate aqueous two-phase systems 586(1991)341

Hydrophobic interaction chromatography for the purification of a mycobacterial heat shock protein of relative molecular mass 60 000 587(1991)19

High-performance liquid chromatographic study of some biologically active analogues of arabinosylcytosine 588(1991)147

Chromatographic separation and molecular modelling of triazines with respect to their inhibition of the growth of L1210/R71 cells 592(1992)59

Polymer-coated reversed-phase packings with controlled hydrophobic properties. I. Effect on the selectivity of protein separations 599(1992)65

Characterization of hydrophobic interaction and hydrophobic interaction chromatography media by multivariate analysis 599(1992)131

Hydrostatic flow

Effect of hydrostatic flow on the efficiency in capillary electrophoresis 559(1991)81

Hydrostatic pressure elution

Electrokinetic reversed-phase chromatography with packed capillaries 593(1992)313

N-(Hydroxymethyl)thioamide resin

N-(Hydroxymethyl)thioamide resin as stationary phase in ion-exchange chromatography for metal ion separation 555(1991)291

8-Hydroxyquinoline metal chelate support

High-performance liquid chromatography of amino acids, peptides and proteins. CXXI. 8-Hydroxyquinoline-metal chelate chromatographic support: an additional mode of selectivity in immobilized-metal affinity chromatography 599(1992)171

Hypermedia

CRISEBOOK, a Hypermedia version of an expert system for the selection of optimization criteria in high-performance liquid chromatography 592(1992)51

I**Ideal chromatograms**

New approach for calculating ideal chromatograms from arbitrary composite distribution isotherms 556(1991)95

Image processing

Important design features of a system for the densitometric analysis of two-dimensional flat-bed separations 596(1992)241

Immobilization

Spray jet assembly interface for the coupling of reversed-phase narrow-bore liquid chromatography and Fourier transform infrared spectrometry 552(1991)635

Gas chromatographic comparative study of Superox 20M immobilized in different ways 556(1991)485

Immobilization method for polyethylene glycol using a cross-linking co-agent 558(1991)323

Immobilization of Fv antibody fragments on porous silica and their utility in affinity chromatography 587(1991)161

Gas chromatographic properties of immobilized poly(ethylene glycol) stationary phases 588(1991)225

Oriented immobilization of peptide ligands on solid supports 591(1992)99

Immobilization of monoclonal antibodies for affinity chromatography using a chelating peptide 595(1992)113

Silica sorbents with one- and two-site attached bacitracin in affinity chromatography 596(1992)114

Immobilization of protein ligands with methyl vinyl ether-maleic anhydride copolymer 597(1992)123

Evaluation of the mode of binding of immunoglobulin to activated agarose 597(1992)231

Affinity purification of antibodies using immobilized FB domain of protein A 597(1992)257

Immobilized antibodies

Comparison of the performance of immunosorbents prepared by site-directed or random coupling of monoclonal antibodies 558(1991)55

Use of antibody fragments in immunoaffinity chromatography. Comparison of FV fragments, VH fragments and paralog peptides 597(1992)239

Immobilized antisense peptides

Purification of recombinant human interferon- β by immobilized antisense peptides 600(1992)235

Immobilized dyes

Modification of reversed-phase columns with dyed surfactants. Preparation of mechanically resistant efficient immobilized dyes for protein purification 558(1991)43

Immobilized enzyme reactors

Characterization of cellulase-based enzyme reactors for the high-performance liquid chromatographic determination of β -D-glucan oligosaccharides 558(1991)343

Specific detection of acetyl-coenzyme A by reversed-phase ion-pair high-performance liquid chromatography with an immobilized enzyme reactor 590(1992)241

Determination of organic acids, amino acids and saccharides by high-performance liquid chromatography and a postcolumn enzyme reactor with amperometric detection 591(1992)165

Enzymophoresis of nucleic acids by tandem capillary enzyme reactor-capillary zone electrophoresis 596(1992)251

Use of various types of column reactors for flow-injection analysis 597(1992)435

Immobilized lectins

Immobilized lectin columns: useful tools for the fractionation and structural analysis of oligosaccharides (Review) 597(1992)111

Immobilized ligands

Bead cellulose derivatives as supports for immobilization and chromatographic purification of proteins 552(1991)389

Trypsin and affinity chromatography (Review) 597(1992)3

Study of high-affinity interactions by quantitative affinity chromatography. Analytical expressions in terms of total ligand concentration 597(1992)57

Affinity chromatography in purification of A₁ adenosine receptors 597(1992)335

Biospecific interaction analysis using surface plasmon resonance detection applied to kinetic, binding site and concentration analysis 597(1992)397

Immobilized metal affinity chromatography

Immobilized metal affinity chromatography as a means of fractionating microsomal cytochrome P-450 isozymes 587(1991)33

High-performance liquid chromatography of amino acids, peptides and proteins. CXXI. 8-Hydroxyquinoline-metal chelate chromatographic support: an additional mode of selectivity in immobilized-metal affinity chromatography 599(1992)171

Immobilized metal ions

Immobilization of monoclonal antibodies for affinity chromatography using a chelating peptide 595(1992)113

Immobilized metal ion affinity electrophoresis. A study with several model proteins containing histidine 597(1992)357

Separon HEMA modified for immobilized metal ion affinity chromatographic separation of proteins 598(1992)15

Immobilized pH gradients

Synthesis of a new acrylamido buffer (acryloylhistamine) for isoelectric focusing in immobilized pH gradients and its analysis by capillary zone electrophoresis 558(1991)285

Isoelectric focusing in immobilized pH gradients: applications in clinical chemistry and forensic analysis (Review) 569(1991)197

Immobilized proteins

Direct liquid chromatographic separation of enantiomers on immobilized protein stationary phases. IX. Influence of the cross-linking reagent on the retentive and enantioselective properties of chiral sorbents based on bovine serum albumin 591(1992)65

Galactosylation as a tool for the stabilization and immobilization of proteins (Review) 597(1992)19

Immobilized receptors

Membrane-based receptor affinity chromatography 597(1992)155

Immunoabsorbents

Compared stability of Sepharose-based immunoabsorbents prepared by various activation methods 584(1992)17

Immunoaffinity

Kinetic aspects of membrane-based immunoaffinity chromatography 597(1992)167

Immunoaffinity adsorbents

Immobilization of Fv antibody fragments on porous silica and their utility in affinity chromatography 587(1991)161

Purification and determination of glutamine synthetase by high-performance immunoaffinity chromatography 589(1992)121

Characterization of synthetic macroporous packing materials in low-pressure cartridges and columns 590(1992)83

Immobilization of monoclonal antibodies for affinity chromatography using a chelating peptide 595(1992)113

Selective clean-up method using an immunoaffinity column following radioimmunoassay of prostaglandin F_{2α} in biological fluids 597(1992)201

Preparation of immunoaffinity mini-columns for the analysis of platelet activating factor (PAF) in biological samples 597(1992)309

Immunoaffinity interactions

Comparison of the performance of immunosorbents prepared by site-directed or random coupling of monoclonal antibodies 558(1991)55

Immunoaffinity precolumn

Determination of aflatoxin M1 using a dialysis-based immunoaffinity sample pretreatment system coupled on-line to liquid chromatography. Reusable immunoaffinity columns 589(1992)141

Immunoaffinity resins

Physical and biochemical characterization of five commercial resins for immunoaffinity purification of factor IX 595(1992)103

Immunoconjugates

Immobilization of Fv antibody fragments on porous silica and their utility in affinity chromatography 587(1991)161

Immunoconjugates

Comparison of the performance of immunosorbents prepared by site-directed or random coupling of monoclonal antibodies 558(1991)55

Impregnated layers

Comparative study of chromatography on thin layers impregnated with organic stationary phases. Chromatographic separation of nitrophenols 558(1991)415

Impregnated paper

Quantitative separation of Cr³⁺ from Mo⁶⁺, W⁶⁺, Hg₂²⁺, Cu²⁺ and Pb²⁺. Chromatographic behaviour of 51 cations on papers impregnated with Sn(IV)-based inorganic ion exchangers in complex-forming acid systems 594(1992)379

Impregnated silica gel layers

Separation of minoxidil and its intermediates by overpressured layer chromatography using a stationary phase bonded with tricaprilmethylammonium chloride 553(1991)459

Inclusion complexes

Effects of β-cyclodextrin in the mobile phase on the retention and indirect detection of non-electrolytes in reversed-phase liquid chromatography. II. Steroids 552(1991)583

Improved high-performance liquid chromatographic resolution of the geometric isomers of 6-hydroxy-4-(1-hydroxy-1-methylethyl)-1-cyclohexene-1-ethanol and by-products with β -cyclodextrin 553(1991)15

Capillary tube isotachopheric separation of catecholamines using cyclodextrin in the leading electrolyte 587(1991)364

Indirect detection

Effects of β -cyclodextrin in the mobile phase on the retention and indirect detection of non-electrolytes in reversed-phase liquid chromatography. I. Study of aliphatic alcohols 552(1991)563

Effects of β -cyclodextrin in the mobile phase on the retention and indirect detection of non-electrolytes in reversed-phase liquid chromatography. II. Steroids 552(1991)583

Separation and indirect detection of amino acids as acetylated derivatives 557(1991)315

Quantitative aspects of indirect UV detection in capillary zone electrophoresis 588(1991)321

Binary buffers for indirect absorption detection in capillary zone electrophoresis 589(1992)307

Infinite dilution

Activity coefficients at infinite dilution determined by gas-liquid chromatography. Organic solvents in Apiezon L 586(1991)303

Infrared spectrometry, Fourier transform

Spray jet assembly interface for the coupling of reversed-phase narrow-bore liquid chromatography and Fourier transform infrared spectrometry 552(1991)635

Assessment of a thermospray interface for the coupling of high-performance liquid chromatography and Fourier transform infrared spectrometry 588(1991)15

Injection concentration

Preparative liquid chromatography. II. Existence of optimum injection conditions for overloaded gradient elution separations 596(1992)15

Injection conditions

Preparative liquid chromatography. I. Influence of column efficiency on optimum injection conditions under isocratic elution 590(1992)175

Injection methods

Analysis of sulthiame in serum by narrow-bore high-performance liquid chromatography. Comparison of direct sample injection with pre-column switching and extrelut extraction 553(1991)7

Split injection into a capillary column at very low split ratios 557(1991)1

Field amplified sample injection in high-performance capillary electrophoresis 559(1991)141

Field-amplified polarity-switching sample injection in high-performance capillary electrophoresis 559(1991)153

Origin and control of multi-peak formation in the analysis of trimethylsilyl derivatives of flavanone aglycones by capillary column gas chromatography 586(1991)323

Gas chromatographic analysis of fatty acid methyl esters: avoiding discrimination by programmed temperature vaporizing injection 588(1991)265

Comparison of liquid-junction and coaxial interfaces for capillary electrophoresis-mass spectrometry with application to compounds of concern to the aquaculture industry 591(1992)325

Use of direct injection precolumn techniques for the high-performance liquid chromatographic determination of the retinoids acitretin and 13-*cis*-acitretin in plasma 593(1992)55

Isoelectric focusing field-flow fractionation. III. Investigation of the influence of different experimental parameters on focusing of cytochrome *c* in the trapezoidal cross-section channel 600(1992)297

Injection port liner

Decomposition of two methylbenzothiophene sulphoxides in a commercial gas chromatography injection port liner 591(1992)362

Injectors

Split injector for capillary zone electrophoresis 559(1991)103

Integral Doppler anemometry

Non-equilibrium integral Doppler anemometric analysis of particle mixtures in a channel flow using an intrinsic hydrodynamic focusing force biased by another force 553(1991)517

Integral Doppler anemometry in porous membranes for the analysis of liquid mixtures and examination of membrane properties 600(1992)129

Transverse particle redistribution in a flat channel for SPLITT or integral Doppler anemometry 600(1992)133

Use of integral Doppler anemometry in field-flow fractionation 600(1992)139

Integration programs

Quantitative two-dimensional thin-layer chromatography 553(1991)477

Interactive surface contact area

High-performance liquid chromatography of amino acids, peptides and proteins. CXV. Thermodynamic behaviour of peptides in reversed-phase chromatography 593(1992)103

Interconversion rate

Origin and control of multi-peak formation in the analysis of trimethylsilyl derivatives of flavanone aglycones by capillary column gas chromatography 586(1991)323

Interface configuration

Comparison of liquid-junction and coaxial interfaces for capillary electrophoresis-mass spectrometry with application to compounds of concern to the aquaculture industry 591(1992)325

Interfaces, electrophoresis-MS

Instrumentation for high-performance capillary electrophoresis-mass spectrometry 559(1991)197

Comparison of liquid-junction and coaxial interfaces for capillary electrophoresis-mass spectrometry with application to compounds of concern to the aquaculture industry 591(1992)325

Interfaces, LC-IR

Spray jet assembly interface for the coupling of reversed-phase narrow-bore liquid chromatography and Fourier transform infrared spectrometry 552(1991)635

Assessment of a thermospray interface for the coupling of high-performance liquid chromatography and Fourier transform infrared spectrometry 588(1991)15

Interfaces, LC-MS

Strategies in developing interfaces for coupling liquid chromatography and mass spectrometry 554(1991)3

Electrospray interfacing for the coupling of ion-exchange and ion-pairing chromatography to mass spectrometry 554(1991)27

Liquid chromatography-mass spectrometry with ionspray and electrospray interfaces in pharmaceutical and biomedical research 554(1991)39

Quantitation and linearity for particle-beam liquid chromatography-mass spectrometry 554(1991)103

Some aspects of peak broadening in particle-beam liquid chromatography-mass spectrometry 554(1991)119

Use of non-volatile ion-pairing agents for liquid chromatographic-mass spectrometric analyses with a moving-belt interface 554(1991)281

Simple direct liquid introduction system usable as an interface for liquid chromatography-mass spectrometry on quadrupole and magnetic-sector mass spectrometers 554(1991)293

Nanoscale separations. Capillary liquid chromatography-mass spectrometry and capillary zone electrophoresis-mass spectrometry for the determination of peptides and proteins 554(1991)329

Pseudo-electrochromatography-mass spectrometry: a new alternative 554(1991)339

Interfaces, SFC-MS

Ionization mechanisms in capillary supercritical fluid chromatography-chemical ionization mass spectrometry 554(1991)351

Interfacial effects

Interfacial effects in size-exclusion chromatography of latex 587(1991)276

Internal standards

Use of N-acetylpsychosine as internal standard for quantitative high-performance liquid chromatographic analysis of glycosphingolipids 555(1991)267

Selection of robust combinations of extraction liquid composition and internal standard. Monte Carlo simulation of improvement of assay methods with liquid-liquid extraction prior to high-performance liquid chromatography 594(1992)45

Internal-surface reversed-phase packing

Evaluation of restricted access media for high-performance liquid chromatographic analysis of sulfonamide antibiotic residues in bovine serum 598(1992)23

Application of an internal surface reversed-phase column for the automated determination of flucyclozuron residues 600(1992)211

Intraparticle convection

Importance of intraparticle convection in the performance of chromatographic processes 590(1992)93

Inverse gas chromatography

Attempts to use Laffort's solubility factors as polarity parameters for organic compounds in inverse gas chromatography 555(1991)205

Inverse gas-solid chromatography

Experimental determination of adsorption isotherm data for the study of the surface energy distribution of various solid surfaces by inverse gas-solid chromatography 591(1992)233

Precision and accuracy of the gas-solid adsorption isotherms derived by the elution-by-characteristic-points method 591(1992)245

In vivo metabolism

Construction of kinetic models to understand metabolism *in vivo* 566(1991)309

Ion-exchange capacity

Carbonate eluents in ion chromatography. Fractions of resin capacity linked to anions present in the eluent 589(1992)197

Ion exchangers

Bead cellulose derivatives as supports for immobilization and chromatographic purification of proteins 552(1991)389

High-performance liquid chromatography of amino acids and dipeptides on new ion exchangers of the HEMA series 552(1991)439

Anion-exchange chromatographic properties of α -lactalbumin eluted from quaternized polyvinylimidazole. Study of the role of the polymer coating 553(1991)165

Chromatographic characterization of ion exchangers for high-performance liquid chromatography of proteins. I. Chromatographic determination of loading capacity for low- and high-molecular mass anions 556(1991)363

Phosphorylated polystyrene resins in high-performance ion-exchange chromatography 589(1992)87

Economic considerations important in the scale-up of an ovalbumin separation from hen egg-white on the anion-exchange cellulose DE92 590(1992)49

Schemes for efficient protein purification on a family of polymeric ion exchangers in glass columns 599(1992)239

Ion-exchange sites

Effects of stationary phase ligand density on high-performance ion-exchange chromatography of proteins 598(1992)7

Ion exclusion

Determination of ascorbic and dehydroascorbic acid in potatoes (*Solanum tuberosum*) and strawberries using ion-exclusion chromatography 594(1992)187

Ionic charge

Determination of the mean ionic charge of the components of three ^{99m}Tc bone scanning agents by gel chromatography with two eluents of different electrolyte composition 594(1992)217

Ionic interactions

Mixed-mode hydrophilic and ionic interaction chromatography rivals reversed-phase liquid chromatography for the separation of peptides 594(1992)75

Ionic mobility

Influence of poly(vinylpyrrolidone) on isotachophoretic separations of inorganic anions in aqueous electrolyte systems 589(1992)329

Peak dispersion and separation efficiency in high-performance zone electrophoresis with gel-filled capillaries 595(1992)313

Ionic strength

Affinity partitioning of erythrocytic phosphofructokinase in aqueous two-phase systems containing poly(ethylene glycol)-bound Cibacron Blue. Influence of pH, ionic strength and substrates/effectors 589(1992)127

Radial pH distribution during capillary electrophoresis with electroosmotic flow. Analysis with high ionic strength buffers 589(1992)315

Analysis of pH-dependent protein interactions with gel filtration medium 591(1992)121

Ionic strength gradient

Separation of some platinum(II) complexes by ionic strength gradient on a solvent-generated ion-exchange sorbent 586(1991)291

Ion interaction

Reversed-phase liquid chromatographic behaviour of alkylamines with amperometric detection and its application to trace analysis 594(1992)159

Ionization conditions

Simple direct liquid introduction system usable as an interface for liquid chromatography-mass spectrometry on quadrupole and magnetic-sector mass spectrometers 554(1991)293

Ionization efficiencies

Quantitative measurements via co-elution and dual-isotope detection by gas chromatography-mass spectrometry 587(1991)255

Ionization mechanisms

Ionization mechanisms in capillary supercritical fluid chromatography-chemical ionization mass spectrometry 554(1991)351

Ionization methods

Evaluation of eluents in thermospray liquid chromatography-mass spectrometry for identification and determination of pesticides in environmental samples 553(1991)311

Strategies in developing interfaces for coupling liquid chromatography and mass spectrometry 554(1991)3

Liquid chromatography-mass spectrometry with ionspray and electrospray interfaces in pharmaceutical and biomedical research 554(1991)39

Polar, hydrophilic compounds in drinking water produced from surface water. Determination by liquid chromatography-mass spectrometry 554(1991)251

Determination of taxol by high-performance liquid chromatography-thermospray mass spectrometry 594(1992)153

Simultaneous determination of sorbic acid, dehydroacetic acid and benzoic acid by gas chromatography-mass spectrometry 594(1992)253

Screening of organophosphorus pesticides using liquid chromatography-atmospheric pressure chemical ionization mass spectrometry 595(1992)193

Ion mobility detector

Photoemissive ionisation source for ion mobility detectors 588(1991)239

Ion-mobility spectrometry

Ion-mobility spectrometry as a detection method for packed-column supercritical fluid chromatography 552(1991)551

Ion-pair extraction

Sensitive liquid chromatographic determination of alkyl-, nitro- and chlorophenols by precolumn derivatization with dansyl chloride, postcolumn photolysis and peroxyoxalate chemiluminescence detection 553(1991)345

Ion-pairing agents

Extension of the electrostatic retention model of reversed-phase ion-pair high-performance liquid chromatography to include the effect of the eluent pH 552(1991)13

Selection of buffers and of an ion-pairing agent for thermospray liquid chromatographic-mass spectrometric analysis of ionic compounds 552(1991)605

Use of non-volatile ion-pairing agents for liquid chromatographic-mass spectrometric analyses with a moving-belt interface 554(1991)281

Gradient elution with shorter equilibration times in reversed-phase ion-pair chromatography 592(1992)143

Ion radius, hydrated

Dependence of the silanophilic effect on the concentration of preadsorbed salts and on the chemical structure of peptides in reversed-phase thin-layer chromatography 553(1991)467

Ion-sensitive field effect transistor

Ion-sensitive field effect transistor as a monovalent cation detector for ion chromatography and its application to the measurement of Na⁺ and K⁺ concentrations in serum 566(1991)109

Ion spray interface

Liquid chromatography-mass spectrometry with ionspray and electrospray interfaces in pharmaceutical and biomedical research 554(1991)39

High-performance liquid chromatographic-mass spectrometric analysis of oligosaccharides from enzymatic digestion of glycosaminoglycans. Application to human samples 591(1992)225

Ion spray mass spectrometry

Determination of sulfonamides by liquid chromatography, ultraviolet diode array detection and ion-spray tandem mass spectrometry with application to cultured salmon flesh 558(1991)155

Determination of aminoglycoside antibiotics by reversed-phase ion-pair high-performance liquid chromatography coupled with pulsed amperometry and ion spray mass spectrometry 591(1992)195

Ion-trap detector

Determination of α - and β -zearalenol and zearalenone in cereals by gas chromatography with ion-trap detection 595(1992)259

Ion trap mass spectrometer

Multi-matrix-multi-pesticide method for agricultural products 552(1991)259

Thermospray liquid chromatography-mass spectrometry with a quadrupole ion trap mass spectrometer 562(1991)3

Isoelectric focusing

Synthesis of a new acrylamido buffer (acryloylhistamine) for isoelectric focusing in immobilized pH gradients and its analysis by capillary zone electrophoresis 558(1991)285

Capillary zone electrophoresis analysis of acrylamido buffers for isoelectric focusing in immobilized pH gradients 559(1991)119

Fractionation of the human recombinant tissue plasminogen activator (rtPA) glycoforms by high-performance capillary zone electrophoresis and capillary isoelectric focusing 559(1991)401

Optimizing separation parameters in capillary isoelectric focusing 559(1991)479

Theoretical background for clinical and biomedical applications of electromigration techniques (Review) 569(1991)3

Isoelectric focusing in immobilized pH gradients: applications in clinical chemistry and forensic analysis (Review) 569(1991)197

Isolation and purification of bacterial proteinases by means of autofocusing 585(1991)373

Capillary isoelectric focusing with electroosmotic zone displacement and on-column multichannel detection 589(1992)321

Characterization of the heterogeneity of polyethylene glycol-modified superoxide dismutase by chromatographic and electrophoretic techniques 599(1992)141

Isoelectric focusing field-flow fractionation. III. Investigation of the influence of different experimental parameters on focusing of cytochrome *c* in the trapezoidal cross-section channel 600(1992)297

Isoelectric focusing field-flow fractionation. IV. Investigations on protein separations in the trapezoidal cross-section channel 600(1992)305

Isotachic patterns

Analysis of isotachic patterns in displacement chromatography 556(1991)119

Isotachopheresis

Dilution effect of carbonate ion on isotachopheretic zones observed with scanning UV photometric detection 555(1991)247

Influence of counter-ion inclusion complexation on the quality of cyclodextrin-supported separations in isotachopheresis 557(1991)215

Isotachopheresis of proteins in uncoated open-tubular fused-silica capillaries with a simple approach for column conditioning 558(1991)423

Selected applications of cyclodextrin selectors in capillary electrophoresis 559(1991)215

Capillary isotachopheresis with concentration-gradient detection. An application to the separation of synthetic peptides 559(1991)489

Isotachopheretic analysis of flavonoids and phenolcarboxylic acids of relevance to phytopharmaceutical industry 559(1991)499

Theoretical background for clinical and biomedical applications of electromigration techniques (Review) 569(1991)3

Application of capillary isotachopheresis in peptide analysis (Review) 569(1991)123

Purification of ovalbumin and lysozyme from a commercial product by recycling isotachopheresis 585(1991)145

Isotachopheretic separation of minor components from a matrix component in the case of strong electrolytes 585(1991)297

Determination of calcium ion in the presence of phosphate anion and collagen by capillary-type isotachopheresis 586(1991)363

Capillary tube isotachopheretic separation of catecholamines using cyclodextrin in the leading electrolyte 587(1991)364

Influence of poly(vinylpyrrolidone) on isotachopheretic separations of inorganic anions in aqueous electrolyte systems 589(1992)329

Study of isotachopheretic separation behaviour of metal cations by means of particle-induced X-ray emission. II. Separation efficiency of lanthanoid model mixture 589(1992)339

Comparison of isotachopheresis, capillary zone electrophoresis and high-performance liquid chromatography for the determination of salbutamol, terbutaline sulphate and fenoterol hydrobromide in pharmaceutical dosage forms 590(1992)341

Analyte focusing in capillary electrophoresis using on-line isotachopheresis 591(1992)341

Continuous separation and purification of proteins by recycling isotachopheresis 594(1992)361

Study of isotachopheretic separation behaviour of metal cations by means of particle-induced X-ray emission. III. Analysis of a crude rare earth chloride from monazite 594(1992)371

Relationship between zone length and step height in isotachopheresis 595(1992)327

Determination of limiting ionic mobilities and dissociation constants of some local anaesthetics 596(1992)265

Isotachopheretic analyzer

Isotachopheretic analyser with options for operational electrolyte selection and repeated analysis 555(1991)321

Isotherm hysteresis

Adsorption-desorption isotherm hysteresis of β -lactoglobulin A with a weakly hydrophobic surface 557(1991)369

Isotope dilution gas chromatography

Determination of Pb^{2+} in water by isotope dilution gas chromatography-mass spectrometry of tetraethyllead formed by reaction with sodium tetraethylborate 594(1992)275

Isotope dilution mass spectrometry

Determination of *Catharanthus roseus* alkaloids by high-performance liquid chromatography-isotope dilution thermospray-mass spectrometry 554(1991)227

Phenylalanine in whole blood and plasma — a candidate reference method 562(1991)213

Isotope effects

Isotope effect in gas-liquid chromatography of labelled compounds 588(1991)251

Isotopic gas chromatography-combustion mass spectrometry

Enhanced resolution of organic compounds from sediments by isotopic gas chromatography-combustion-mass spectrometry 585(1991)177

Isotopic labelling

Quantitative measurements via co-elution and dual-isotope detection by gas chromatography-mass spectrometry 587(1991)255

Iterative regression optimization

Multiparameter optimizations in micellar liquid chromatography using the iterative regression optimization strategy 586(1991)221

K**KES**

Feasibility study for the construction of an integrated expert system in high-performance liquid chromatography 589(1992)31

Keto-enol tautomerism

High-performance liquid chromatographic separation of *p*-hydroxyphenylpyruvic acid 585(1991)318

Kinetic data

High-performance liquid chromatographic study of the correlation between the physico-chemical parameters of furan and benzene aldehydes and the rate of their metabolism by yeast 594(1992)23

Kinetic detection

Ion chromatographic determination of anions, especially sulphur-containing anions, with conductimetric and kinetic detection 558(1991)209

Kinetic measurements

Biospecific interaction analysis using surface plasmon resonance detection applied to kinetic, binding site and concentration analysis 597(1992)397

Kinetic model, non-linear chromatography

Analysis of the separability of plate height into overload and intrinsic contributions using the kinetic model of non-linear chromatography 556(1991)159

Kinetic models

Construction of kinetic models to understand metabolism *in vivo* 566(1991)309

Kinetics

Simultaneous determination of prostaglandins E₁, A₁ and B₁ by reversed-phase high-performance liquid chromatography for the kinetic studies of prostaglandin E₁ in solution 555(1991)73

Knox equation

Investigation of the causes of reduced efficiency in micellar liquid chromatography 556(1991)263

L**Laminar flow**

Non-equilibrium integral Doppler anemometric analysis of particle mixtures in a channel flow using an intrinsic hydrodynamic focusing force biased by another force 553(1991)517

Simple solution of velocity profiles of laminar flows in channels of various cross-sections used in field-flow fractionation 593(1992)357

Langmuir isotherms

see Adsorption isotherms

Lantana camara

Thin-layer chromatographic separations of lantadenes, the pentacyclic triterpenoids from lantana (*Lantana camara*) plant 587(1991)351

Large-pore supports

Importance of intraparticle convection in the performance of chromatographic processes 590(1992)93

Laser-excited fluorescence detector

Laser-excited fluorescence detection of gas-phase chromatography eluates 557(1991)39

Laser-induced fluorescence detector

Determination of sultopride and tiapride in serum by gas chromatography using a surface ionisation detector 567(1991)113

Separation of naphthalene-2,3-dicarboxaldehyde-labeled amino acids by high-performance capillary electrophoresis with laser-induced fluorescence detection 593(1992)265

High-sensitivity laser-induced fluorescence detection of native proteins in capillary electrophoresis 595(1992)319

Lateral diffusion

Diffusion of sorbed pyrene in the bonded layer of reversed-phase silicas. Effect of alkyl chain length and pore diameter 558(1991)31

Leading electrolyte

Influence of poly(vinylpyrrolidone) on isotachophoretic separations of inorganic anions in aqueous electrolyte systems 589(1992)329

Lectin affinity electrophoresis

Clinical applications of lectin affinity electrophoresis (Review) 569(1991)229

Leucocytes

Purification of DNA-derived deoxynucleotides from leukocytes involving nuclease elution of an ion-exchange column 600(1992)229

Lewis base eluents

Ion- and ligand-exchange chromatography of proteins using porous zirconium oxide supports in organic and inorganic Lewis base eluents 596(1992)27

Ligand attachment

Silica sorbents with one- and two-site attached bacitracin in affinity chromatography 596(1992)114

Ligand concentration

Study of high-affinity interactions by quantitative affinity chromatography. Analytical expressions in terms of total ligand concentration 597(1992)57

Ligand density

Protein adsorption capacity of a porous phenylalanine-containing membrane based on a polyethylene matrix 586(1991)27

Effects of stationary phase ligand density on high-performance ion-exchange chromatography of proteins 598(1992)7

Ligand exchange

Ligand-exchange ion chromatographic determination of malic acid enantiomers in apple juice with photometric detection 585(1991)315

Preparative separation and analysis of the enantiomers of [³H]Abbott-69992, an HIV anti-infective nucleoside, by ligand-exchange high-performance liquid chromatography 586(1991)265

High-performance ligand-exchange chromatography of some amino acids containing two chiral centres 592(1992)345

Chromatographic separation of polyols by ligand exchange. Effects of the ion-exchange resin cross-linking and size 594(1992)125

Ion- and ligand-exchange chromatography of proteins using porous zirconium oxide supports in organic and inorganic Lewis base eluents 596(1992)27

Role of metal ions in the ligand-exchange separation of amino acids 596(1992)281

Ligand-exchange phase

Direct separation of enantiomers by high-performance liquid chromatography on a new chiral ligand-exchange phase 592(1992)291

Ligand leakage

Cellulose beads: a weak leaking affinity support 589(1992)93

Ligands

Reversed-phase separation of transition metals, lanthanides and actinides by elution with mandelic acid 558(1991)197

High-performance affinity chromatography of NADP⁺ dehydrogenases from cell-free extracts using a nucleotide analogue as general ligand 586(1991)51

Recent developments in quantitative affinity chromatography (Review) 597(1992)67

Light-scattering detector

Use of an evaporative light scattering detector in reversed-phase high-performance liquid chromatography of oligomeric surfactants 552(1991)593

Liquid chromatographic method for the determination of the carbohydrate moiety of glycoproteins. Application to α_1 -acid glycoprotein and tissue plasminogen activator 558(1991)105

Practical on-line determination of biopolymer molecular weights by high-performance liquid chromatography with classical light-scattering detection 592(1992)215

Light stabilizers

Determination of Cyasorb UV 1084 and its degradation products in low-density polyethylene film by high-performance liquid chromatography 587(1991)205

Limiting ionic mobility

Determination of limiting ionic mobilities and dissociation constants of some local anaesthetics 596(1992)265

***Limulus* amoebocyte lysate test**

Specific assay for endotoxin using immobilized histidine, *Limulus* amoebocyte lysate and a chromogenic substrate 597(1992)415

Linear-elution strength model

Accuracy of computer-simulated gas chromatographic separations based on a linear elution strength model 589(1992)215

Linear dependence

Thin-layer chromatography on polyacrylonitrile. IV. Investigation of the separation mechanisms for tris-(alkylxanthato)cobalt(III) complexes 596(1992)91

Linear mobile phase velocity

Short-cut method for predicting the productivity of affinity chromatography 597(1992)173

Linear non-equilibrium chromatography

Theory of linear non-equilibrium chromatography with beds of a finite length 591(1992)273

Linear solvation energy relationship

Solvatochromic hydrogen bond donor acidity of cyclodextrins and reversed-phase liquid chromatographic retention of small molecules on a β -cyclodextrin-bonded silica stationary phase 595(1992)45

Linomat C

Direct coupling of high-speed counter-current chromatography to thin-layer chromatography. Application to the separation of asiaticoside and madecassoside from *Centella asiatica* 558(1991)446

Lipophilicity

see also Hydrophobicity

Influence of mobile phase composition on evaluation of lipophilicity by partition chromatography 557(1991)399

R_M values of some colchicines and colchicineamides determined by reversed-phase thin-layer chromatography 591(1992)367

Support matrix effects in the reversed-phase thin-layer chromatography of some peptides 600(1992)149

Liquid chromatographic reactor

Hydroquinone oxidation kinetics in adsorptive liquid chromatographic beds 556(1991)383

Liquid chromatography-gas chromatography

Reversed-phase liquid chromatography coupled on-line with capillary gas chromatography. I. Introduction of large volumes of aqueous mixtures through an on-column interface 552(1991)489

Liquid chromatography-mass spectrometry

see also Special issue index

Liquid chromatography-mass spectrometry

Selection of buffers and of an ion-pairing agent for thermospray liquid chromatographic-mass spectrometric analysis of ionic compounds 552(1991)605

Strategies in developing interfaces for coupling liquid chromatography and mass spectrometry 554(1991)3

Electrospray interfacing for the coupling of ion-exchange and ion-pairing chromatography to mass spectrometry 554(1991)27

Source of band broadening in liquid chromatographic-fast atom bombardment mass spectrometric systems with precolumn addition of viscous matrix to the mobile phase 554(1991)47

Nanoscale separations combined with tandem mass spectrometry 554(1991)73

On-line derivatization of eluted substances in dynamic high-performance liquid chromatography-mass spectrometry through the particle-beam interface 554(1991)125

Increasing thermospray response for cortisol by derivatization 554(1991)149

Use of methyl oxime derivatives to enhance structural information in thermospray high-performance liquid chromatography-mass spectrometry. Analysis of linoleic acid lipoxygenase metabolites in maize embryos 554(1991)155

Analysis of gentamicin sulfate by high-performance liquid chromatography combined with thermospray mass spectrometry 554(1991)191

Liquid chromatography-thermospray tandem mass spectrometry for identification of a heptabarbital metabolite and sample work-up artefacts 554(1991)205

Evaluation of an automated thermospray liquid chromatography-mass spectrometry system for quantitative use in bioanalytical chemistry 554(1991)215

Determination of *Catharanthus roseus* alkaloids by high-performance liquid chromatography-isotope dilution thermospray-mass spectrometry 554(1991)227

Applications of thermospray liquid chromatography-mass spectrometry in photochemical studies of pesticides in water 554(1991)233

Polar, hydrophilic compounds in drinking water produced from surface water. Determination by liquid chromatography-mass spectrometry 554(1991)251

Use of non-volatile ion-pairing agents for liquid chromatographic-mass spectrometric analyses with a moving-belt interface 554(1991)281

Identification of poly(ethylene terephthalate) cyclic oligomers by liquid chromatography-mass spectrometry 554(1991)305

Nanoscale separations. Capillary liquid chromatography-mass spectrometry and capillary zone electrophoresis-mass spectrometry for the determination of peptides and proteins 554(1991)329

Identification of chlorophyll transformation products in a lake sediment by combined liquid chromatography-mass spectrometry 557(1991)271

High-performance liquid chromatography-atmospheric pressure ionization mass spectrometry of gymnemic acids 557(1991)383

Confirmation of the structure of by-products in the synthesis of Modafinil by liquid chromatography-mass spectrometry 557(1991)489

Determination of sulfonamides by liquid chromatography, ultraviolet diode array detection and ion-spray tandem mass spectrometry with application to cultured salmon flesh 558(1991)155

Thermospray and particle beam liquid chromatographic-mass spectrometric analysis of coumarin anticoagulants 562(1991)31

Use of particle beam liquid chromatography–electron impact mass spectrometry for structure elucidation of oxodipine and three of its metabolites 562(1991)39

Analysis of tropane alkaloids with thermospray high-performance liquid chromatography–mass spectrometry 562(1991)737

Identification of aromatic moieties and mycosamine in antifungal heptaenes with high-performance liquid chromatography, high-performance liquid chromatography–mass spectrometry and gas chromatography–mass spectrometry 585(1991)247

Applicability of high-performance liquid chromatography–continuous-flow fast atom bombardment mass spectrometry for simultaneous quantitation of multiple neurochemicals 587(1991)306

Reversed-phase liquid chromatography–mass spectrometry of complex mixtures of natural triacylglycerols with chloride-attachment negative chemical ionization 588(1991)73

Determination of aromatic sulfonic acids in aqueous environmental samples by anion-exchange chromatography coupled to particle beam mass spectrometry and UV spectrophotometry 589(1991)177

Limits and effects of precolumn addition of thioglycerol in liquid chromatographic–fast atom bombardment mass spectrometric systems 591(1992)129

Determination of aminoglycoside antibiotics by reversed-phase ion-pair high-performance liquid chromatography coupled with pulsed amperometry and ion spray mass spectrometry 591(1992)195

High-performance liquid chromatographic–mass spectrometric analysis of oligosaccharides from enzymatic digestion of glycosaminoglycans. Application to human samples 591(1992)225

Reversed-phase high-performance liquid chromatography–thermospray mass spectrometry of radiation-induced decomposition products of thymine and thymidine 593(1992)133

Determination of muramic acid by high-performance liquid chromatography–plasma spray mass spectrometry 598(1992)43

Liquid chromatography–thin layer chromatography, on-line
On-line coupling of liquid chromatography to thin-layer chromatography for the identification of polycyclic aromatic hydrocarbons in marine sediment by fluorescence excitation and emission spectroscopy 552(1991)613

Liquid crystals

Thermodynamics of solution of non-mesomorphic solutes at infinite dilution in the smectic-A, nematic and isotropic phases of *p*-*n*-octyl-*p*'-cyanobiphenyl. A gas–liquid chromatographic study 556(1991)317

Synthesis of chemically bonded liquid crystals for high-performance liquid chromatography. New phases via the organochlorosilane pathway 556(1991)373

Capillary gas chromatography of C₅–C₁₃ branched alkynes on squalane and liquid crystal stationary phases 557(1991)241

Retention behaviour of polycyclic aromatic hydrocarbons on a liquid-crystal bonded phase in reversed-phase liquid chromatography 557(1991)459

Study of the response of three liquid crystals as stationary phases for the gas chromatography of some cyclic monoterpene volatile oil constituents with short retention times 587(1991)343

Comparison of four liquid crystal stationary phases used above and below their melting point temperatures for the gas chromatography of some volatile oil constituents 588(1991)231

Gas chromatographic retention behaviour of dibenzothiophene derivatives on a smectic liquid crystalline polysiloxane stationary phase 590(1992)297

Chromatographic properties of graphitized thermal carbon black modified with a monolayer of liquid crystal 600(1992)59

Gas chromatographic separation of *cis*–*trans* isomers of alkylcyclohexylbenzenes on a capillary column with a liquid crystalline stationary phase 600(1992)123

Liquid–liquid extraction

Selection of robust combinations of extraction liquid composition and internal standard. Monte Carlo simulation of improvement of assay methods with liquid–liquid extraction prior to high-performance liquid chromatography 594(1992)45

Liquid phases

Attempts to use Laffort's solubility factors as polarity parameters for organic compounds in inverse gas chromatography 555(1991)205

Loading capacity

Chromatographic characterization of ion exchangers for high-performance liquid chromatography of proteins. I. Chromatographic determination of loading capacity for low- and high-molecular mass anions 556(1991)363

Chromatographic characterization of a phosphate-modified zirconia support for bio-chromatographic applications 587(1991)149

Factors affecting the separation and loading capacity of proteins in preparative gradient elution high-performance liquid chromatography 590(1992)35

Lotus tetragonolobus

Purification and characterization of carbohydrate-binding peptides from *Lotus tetragonolobus* and *Ulex europeus* seed lectins using affinity chromatography 597(1992)213

Low-angle laser light scattering photometry

Interfacing gradient elution ion-exchange chromatography and low-angle laser light-scattering photometry for analysis of proteins 591(1992)139

Luminescence detector

Time-resolved luminescence detection of derivatized thiols in column liquid chromatography 552(1991)625

Lung fluid

High-performance capillary electrophoresis of proteins from the fluid lining of the lungs of rats exposed to perfluoroisobutylene 559(1991)411

M

Macrolactonization

Analyse par chromatographie en phase gazeuse de la réaction de macrolactonisation de Mukaiyama 557(1991)501

Macrophages

Fast separation of macrophages by retention on cross-linked amylose and release by enzymatic amylolysis of the chromatographic material 584(1992)121

Manufacturing processes

Beaded polymer supports and gels (Review). I. Manufacturing techniques 586(1991)181

Mass balance equations

Prediction of the elution profiles of high-concentration bands in gas chromatography 589(1992)223

Mass fragmentography

Simultaneous determination of sorbic acid, dehydroacetic acid and benzoic acid by gas chromatography-mass spectrometry 594(1992)253

Mass overload

Displacement effects in preparative gradient high-performance liquid chromatographic separations 590(1992)17

Peak purity, yield and throughput in preparative liquid chromatography with self-displacement 592(1992)17

Mass spectrometry

see also Special issue index

Mass spectrometry

Recent advances in liquid chromatography-mass spectrometry and capillary zone electrophoresis-mass spectrometry for protein analysis 553(1991)101

Simple direct liquid introduction system usable as an interface for liquid chromatography-mass spectrometry on quadrupole and magnetic-sector mass spectrometers 554(1991)293

Capillary electrophoresis-atmospheric pressure ionization mass spectrometry for the characterization of peptides. Instrumental considerations for mass spectrometric detection 554(1991)311

High-performance liquid chromatographic analysis of benzodiazepines using diode array, electrochemical and thermospray mass spectrometric detection 598(1992)59

Mass spectrometric identification of products formed during degradation of ethyl dimethylphosphoramidocyanidate (tabun) 598(1992)89

Mass spectrometry, beam deflection time-of-flight

Complex mixture analysis based on gas chromatography-mass spectrometry with time array detection using a beam deflection time-of-flight mass spectrometer 590(1992)329

Mass transfer

Anion-exchange chromatographic properties of α -lactalbumin eluted from quaternized polyvinylimidazole. Study of the role of the polymer coating 553(1991)165

Theory of perfusion chromatography 599(1992)87

Matrix activation

Activation of matrices by 4,6-diphenylthieno[3,4-*d*]-1,3-dioxol-2-one-5,5-dioxide. High-performance liquid affinity chromatographic separations 589(1992)101

Matrix architecture

Beaded polymer supports and gels (Review). II. Physico-chemical criteria and functionalization 586(1991)199

Matrix effects

Support matrix effects in the reversed-phase thin-layer chromatography of some peptides 600(1992)149

Matrix solid-phase dispersion

Development of a screening method for five sulfonamides in salmon muscle tissue using thin-layer chromatography 555(1991)315

McReynolds solutes

Attempts to use Laffort's solubility factors as polarity parameters for organic compounds in inverse gas chromatography 555(1991)205

Meat

Pre-column derivatization of sulfa drugs with fluorecamine and high-performance liquid chromatographic determination at their residual levels in meat and meat products 558(1991)175

Mellitin-Eupergit C

Calcium-modulated conformational affinity chromatography. Application to the purification of calmodulin and S100 proteins 597(1992)263

Membrane-based immunoaffinity system

Kinetic aspects of membrane-based immunoaffinity chromatography 597(1992)167

Membrane-based sample preparation device

Membrane-based sample preparation device for the pretreatment of acidic samples prior to cation analysis by ion chromatography 586(1991)283

Membrane chromatography

High-performance membrane chromatography of serum and plasma membrane proteins 590(1992)59

Membrane chromatographic systems for high-throughput protein separations 596(1992)173

Membranes

High-performance membrane chromatography of proteins, a novel method of protein separation 555(1991)97

Adsorption and elution of bovine γ -globulin using an affinity membrane containing hydrophobic amino acids as ligands 585(1991)45

Protein adsorption capacity of a porous phenylalanine-containing membrane based on a polyethylene matrix 586(1991)27

Surface-modified membranes as a matrix for protein purification 591(1992)107

Integral Doppler anemometry in porous membranes for the analysis of liquid mixtures and examination of membrane properties 600(1992)129

Menhaden oil

Reversed-phase liquid chromatography-mass spectrometry of complex mixtures of natural triacylglycerols with chloride-attachment negative chemical ionization 588(1991)73

Metabolic disorders

Capillary electrophoresis for diagnosis and studies of human disease, particularly metabolic disorders 559(1991)455

Metal electrodes

Constant-potential amperometric detection of carbohydrates at metal electrodes in high-performance anion-exchange chromatography 592(1992)229

Metal interactions

Possible role for stationary phase metal interactions in the chromatography of hydroxyamines on silica 552(1991)429

Metal-ligand complexes

Reversed-phase separation of transition metals, lanthanides and actinides by elution with mandelic acid 558(1991)197

Metal-silica interactions

Suppression of the effect of metal impurities in alkylsilylated silica packing materials 594(1992)87

Method development

Use of solvent selectivity optimization procedures for high-performance liquid chromatographic method development 592(1992)43

CRISEBOOK, a Hypermedia version of an expert system for the selection of optimization criteria in high-performance liquid chromatography 592(1992)51

Application of the gradient elution technique. Demonstration with a special test mixture and the DryLab G/plus method development software 592(1992)133

Computer simulation for the prediction of separation as a function of pH for reversed-phase high-performance liquid chromatography. II. Resolution as a function of simultaneous change in pH and solvent strength 592(1992)197

Methylbenzoyl cellulose

Chromatographic resolution on methylbenzoylcellulose beads. Modulation of the chiral recognition by variation of the position of the methyl group on the aromatic ring 595(1992)63

Micellar electrokinetic chromatography

see also Capillary electrophoresis

Chiral separation by cyclodextrin-modified micellar electrokinetic chromatography 553(1991)503

Bile salt surfactants in micellar electrokinetic capillary chromatography. Application to hydrophobic molecule separations 557(1991)113

Chiral separations by micellar electrokinetic chromatography with sodium N-dodecanoyl-L-valinate 559(1991)209

Determination of thiopental in human serum and plasma by high-performance capillary electrophoresis-micellar electrokinetic chromatography 559(1991)505

Separation of Dns-amino acids and vitamins by micellar electrokinetic chromatography 559(1991)537

Electrochromatographic solid-phase extraction for determination of cimetidine in serum by micellar electrokinetic capillary chromatography 559(1991)547

- Micellar electrokinetic capillary chromatographic separation and laser-induced fluorescence detection of 2'-deoxynucleoside 5'-monophosphates of normal and modified bases 565(1991)197
- Theoretical background for clinical and biomedical applications of electromigration techniques (Review) 569(1991)3
- Determination of some drugs by micellar electrokinetic capillary chromatography. The pseudo-effective mobility as parameter for screening 585(1991)123
- Micellar electrokinetic capillary chromatography of vitamin B₆ with electrochemical detection 585(1991)139
- Determination of antihistamines in pharmaceuticals by capillary electrophoresis 588(1991)335
- Separation of polycyclic aromatic hydrocarbons by micellar electrokinetic chromatography with cyclodextrins as modifiers 589(1992)333
- Separation of flavonol-2-O-glycosides from *Calendula officinalis* and *Sambucus nigra* by high-performance liquid and micellar electrokinetic capillary chromatography 593(1992)165
- Determination of substituted purines in body fluids by micellar electrokinetic capillary chromatography with direct sample injection 593(1992)275
- Micellar electrokinetic chromatographic study of hydroquinone and some of its ethers. Determination of hydroquinone in skin-toning cream 596(1992)95
- Micellar liquid chromatography**
Investigation of the causes of reduced efficiency in micellar liquid chromatography 556(1991)263
- Multiparameter optimizations in micellar liquid chromatography using the iterative regression optimization strategy 586(1991)221
- Micellar mobile phases**
Aqueous high-performance size-exclusion chromatographic assay for high-molecular-weight impurities in ceftiofur sodium 595(1992)185
- Micellar phase-transfer catalysis**
Derivatization of saturated long-chain fatty acids with phenacyl bromide in non-ionic micelles 586(1991)347
- Micelles**
Separation of soy lecithin using gel permeation chromatography 589(1992)165
- Microbial starch degradation**
Rapid identification of microbial starch degradation products from a complex nutrient medium by a thin-layer chromatographic method 558(1991)328
- Microcapillary columns**
Gas-liquid microcapillary columns precoated with graphitized carbon black 552(1991)197
- Microcapsules**
Chromatographic fractionation of nucleic acids using microcapsules made from plant cells 585(1991)57
- Microcystis aeruginosa***
Separation of methylated and non-methylated cyanoginosin-LR homologues of the cyanobacterium *Microcystis aeruginosa* strain PCC 7806 by reversed-phase medium-pressure liquid chromatography 558(1991)430
- Microdialysis probes**
In vivo sampling using loop microdialysis probes coupled to a liquid chromatograph 578(1992)157
- Microelectrodes**
Capillary electrophoresis with electrochemical detection employing an on-column Nafion joint 593(1992)305
- Microfiltration**
Protein adsorption capacity of a porous phenylalanine-containing membrane based on a polyethylene matrix 586(1991)27
- Micromachining**
Planar chips technology for miniaturization and integration of separation techniques into monitoring systems. Capillary electrophoresis on a chip 593(1992)253
- Micro-membrane suppressors**
Use of non-volatile ion-pairing agents for liquid chromatographic-mass spectrometric analyses with a moving-belt interface 554(1991)281
- Micropellicular supports**
Fast protein monitoring in fermentation broth using non-porous micropellicular reversed-phase columns 594(1992)392
- Microporous carbons**
Determination of specific retention volumes at 20°C for hydrocarbons on microporous carbons 586(1991)161
- Microwave-induced plasma detector**
Evaluation of gas chromatographic columns for the determination of methylmercury in aqueous head space extracts from biological samples 586(1991)329
- Gas chromatography of chloride and bromide as phenylboronic acid/mercuric nitrate derivatives with microwave induced plasma atomic emission detection 594(1992)395

Migration behaviour

Determination of antihistamines in pharmaceuticals by capillary electrophoresis 588(1991)335

Separation of polycyclic aromatic hydrocarbons by micellar electrokinetic chromatography with cyclodextrins as modifiers 589(1992)333

Prediction of migration behavior of oligonucleotides in capillary gel electrophoresis 593(1991)297

Separation of plant growth regulators by capillary electrophoresis 594(1992)335

Migration distance

Continuous-flow electrophoresis: a separation criterion applied to the separation of model proteins 594(1992)351

Migration reproducibility

Influence of operating parameters on reproducibility in capillary electrophoresis 559(1991)57

Migration time

Comparison of high-performance liquid chromatography with capillary gel electrophoresis in single-base resolution of polynucleotides 558(1991)273

Optimization of resolution in capillary zone electrophoresis: combined effect of applied voltage and buffer concentration 588(1991)315

Migration velocity

Dilution effect of carbonate ion on isotachophoretic zones observed with scanning UV photometric detection 555(1991)247

Milk

Automated clean-up procedure for the gas chromatographic-high-resolution mass spectrometric determination of polychlorinated dibenzo-*p*-dioxins and dibenzofurans in milk 595(1992)289

Miniaturization

Planar chips technology for miniaturization and integration of separation techniques into monitoring systems. Capillary electrophoresis on a chip 593(1992)253

Minor component analysis

Isotachophoretic separation of minor components from a matrix component in the case of strong electrolytes 585(1991)297

Mite

Isolation of *Der pI*, the *Dermatophagoides pteronyssinus* major mite allergen, from a crude mite culture extract, purification by ion-chromatography, and comparison between the material obtained and a cDNA-coded *Der pI* 599(1992)105

Mixed carrier system

Determination of antihistamines in pharmaceuticals by capillary electrophoresis 588(1991)335

Mixed functional phase silica

Preparation and characterization of mixed functional phase silica materials using phenyl-, butyl- or octylchlorosilane as a silylating agent 596(1992)151

Mixed potential response

Mixed potential response at metallic electrodes in potentiometric detection for ion chromatography 589(1992)201

Mobile phase additive

(2*S*,3*S*)-Dicyclohexyl tartrate as mobile phase additive for the determination of the enantiomeric purity of (*S*)-atropine in tablets 592(1992)339

Mobile phase composition

Adsorption and partition mode in high-performance liquid chromatography of highly polar solutes on silica 552(1991)23

Development of a rational optimisation procedure for the automated sample clean-up with column switching in pesticide residue analysis 552(1991)113

Effects of β -cyclodextrin in the mobile phase on the retention and indirect detection of non-electrolytes in reversed-phase liquid chromatography. I. Study of aliphatic alcohols 552(1991)563

Use of an evaporative light scattering detector in reversed-phase high-performance liquid chromatography of oligomeric surfactants 552(1991)593

Selection of buffers and of an ion-pairing agent for thermospray liquid chromatographic-mass spectrometric analysis of ionic compounds 552(1991)605

Improved high-performance liquid chromatographic resolution of the geometric isomers of 6-hydroxy-4-(1-hydroxy-1-methylethyl)-1-cyclohexene-1-ethanol and by-products with β -cyclodextrin 553(1991)15

Effect of the addition of viscous matrices to the mobile phase on chromatographic performance in liquid chromatography-fast atom bombardment mass spectrometry 554(1991)61

Quantitation and linearity for particle-beam liquid chromatography-mass spectrometry 554(1991)103

Analysis of gentamicin sulfate by high-performance liquid chromatography combined with thermospray mass spectrometry 554(1991)191

Ionization mechanisms in capillary supercritical fluid chromatography-chemical ionization mass spectrometry 554(1991)351

- Application of solute and mobile phase partition coefficient to describe solute retention in reversed-phase high-performance liquid chromatography 555(1991)21
- Experimental model for predicting solute retention time as a function of mobile phase composition in ternary reversed-phase high-performance liquid chromatography 555(1991)33
- Correlation analysis in liquid chromatography of metal chelates. II. One-dimensional retention-mobile phase composition (physicochemical property) models in reversed-phase liquid chromatography 555(1991)163
- Application of ligand-exchange chromatography to the assay of L-alanine from DL-aspartic acid by *Pseudomonas dacunhae* 555(1991)272
- Some aspects of optimization in planar chromatography (Review) 556(1991)81
- Method for characterization of selectivity in reversed-phase liquid chromatography. V. Calibration of the retention scale for chromatographic systems with low concentrations of organic solvents in the mobile phase 556(1991)145
- Description of retention mechanism by solvophobic theory. Influence of organic modifiers on the retention behaviour of homologous series in reversed-phase liquid chromatography 556(1991)219
- Separation of *trans/cis* α - and β -carotenes by supercritical fluid chromatography. I. Effects of temperature, pressure and organic modifiers on the retention of carotenes 557(1991)47
- Packed column supercritical fluid chromatography with carbon dioxide-polar modifiers. Influence of carbon dioxide density on retention 557(1991)85
- Separation and indirect detection of amino acids as acetylated derivatives 557(1991)315
- Influence of mobile phase composition on evaluation of lipophilicity by partition chromatography 557(1991)399
- Unusual effect of temperature on the retention of enantiomers on a chiral column 558(1991)1
- Retention data for five ketotrichothecenes in reversed-phase high-performance liquid chromatography with different eluent systems 558(1991)81
- Reversed-phase separation of transition metals, lanthanides and actinides by elution with mandelic acid 558(1991)197
- Optimization of the direct enantiomeric separation by high-performance liquid chromatography of the D-2 dopamine agonist N-0437 using response surface methodology and the multi-criteria decision-making method 558(1991)333
- Comparative study of chromatography on thin layers impregnated with organic stationary phases. Chromatographic separation of nitrophenols 558(1991)415
- Reversed-phase separations of high-molecular-weight polystyrenes in a dichloromethane-acetonitrile solvent system 558(1991)440
- Capillary supercritical fluid chromatography of primary aliphatic amines using carbon dioxide and nitrous oxide as mobile phases 558(1991)458
- Studies on anabolic steroids. V. Sequential reduction of methandienone and structurally related steroid A-ring substituents in humans: gas chromatographic-mass spectrometric study of the corresponding urinary metabolites 562(1991)323
- Mobile phase variations in thermospray liquid chromatography-mass spectrometry of pesticides 562(1991)507
- (2*R*,3*R*)-Dicyclohexyl tartrate as a chiral mobile phase additive 585(1991)35
- Peak distortion in the column liquid chromatographic determination of omeprazole dissolved in borax buffer 586(1991)271
- Secondary equilibrium size-exclusion chromatography of ions with polymeric mobile phase additives 586(1991)277
- Reversal of elution order during direct enantiomeric separation of pyriproxyfen on a cellulose-based chiral stationary phase 588(1991)177
- Elution behaviour of polyamic acid and polyamide-imide in size-exclusion chromatography 588(1991)195
- Chromatographic behaviour of the *closa*-[B₁₂H₁₂]²⁻ derivatives on hydroxyethylmethacrylate gels 588(1991)201
- Enantiomeric separation of racemic hydroperoxides and related alcohols 588(1991)352
- Optimization of an ion-exchange high-performance liquid chromatographic method for the determination of carboxylic acids, sugars, glycerol and ethanol in wines 589(1992)151
- Optimization of an isocratic high-performance liquid chromatographic separation of carotenoids 589(1992)171
- Carbonate eluents in ion chromatography. Fractions of resin capacity linked to anions present in the eluent 589(1992)197
- Potentiometric detection of carboxylic acids and inorganic anions in ion-exclusion chromatography using camphorsulphonic acid as eluent 589(1992)209

Displacement effects in preparative gradient high-performance liquid chromatographic separations 590(1992)17

Limits and effects of precolumn addition of thioglycerol in liquid chromatographic-fast atom bombardment mass spectrometric systems 591(1992)129

Optimization of the separation of (6*R*)- and (6*S*)-leucovorin and evaluation of the robustness of the optimum 592(1992)37

Use of solvent selectivity optimization procedures for high-performance liquid chromatographic method development 592(1992)43

Modelling retention in reversed-phase liquid chromatography as a function of pH and solvent composition 592(1992)157

Computer simulation for the prediction of separation as a function of pH for reversed-phase high-performance liquid chromatography. II. Resolution as a function of simultaneous change in pH and solvent strength 592(1992)197

Reversed retention order and other stereoselective effects in the separation of amino alcohols on Chiralcel OD 592(1992)331

Mixed-mode hydrophilic and ionic interaction chromatography rivals reversed-phase liquid chromatography for the separation of peptides 594(1992)75

Evaluation of reversed-phase liquid chromatographic columns for recovery and selectivity of selected carotenoids 595(1992)89

Stationary phase effect on enantioselectivity of dansyl phenylalanine in microcolumn liquid chromatography with γ -cyclodextrin as mobile phase additive 595(1992)121

Ion-exclusion chromatography using mobile phases containing β -cyclodextrin 595(1992)149

Ion- and ligand-exchange chromatography of proteins using porous zirconium oxide supports in organic and inorganic Lewis base eluents 596(1992)27

Buffer and pH dependence of the retention of phenylthiocarbamylamino acids in reversed-phase high-performance liquid chromatography 600(1992)87

Determination of erythromycin ethylsuccinate by liquid chromatography 600(1992)99

Mobility

see Electrophoretic mobility

Mobilization agents

Optimizing separation parameters in capillary isoelectric focusing 559(1991)479

Modifiers

Effects of modifiers in packed and open-tubular supercritical fluid chromatography 552(1991)527

Separation of polycyclic aromatic hydrocarbons by micellar electrokinetic chromatography with cyclodextrins as modifiers 589(1992)333

Separation of plant growth regulators by capillary electrophoresis 594(1992)335

Molar response factor

Characterization of cellulase-based enzyme reactors for the high-performance liquid chromatographic determination of β -D-glucan oligosaccharides 558(1991)343

Molecular connectivity indices

Relationship between gas chromatographic retention indices and molecular connectivity indices of chlorinated pesticides and structurally related compounds 598(1992)243

Molecular descriptors

Determination of sorbent polarity and selectivity by linear regression of retention indices 552(1991)31

Molecular mass

Clean-up of plasma extracts by gel permeation chromatography during analysis of isosorbide nitrates by capillary gas chromatography 553(1991)21

Molecular mass determination

Practical on-line determination of biopolymer molecular weights by high-performance liquid chromatography with classical light-scattering detection 592(1992)215

Molecular mass distribution

Determination of chemical composition of polymers by size-exclusion chromatography with coupled density and refractive index detection. III. Polyethylene oxide and polytetrahydrofuran 552(1991)507

Separation of soy lecithin using gel permeation chromatography 589(1992)165

Analysis of benzyloxy-terminated poly(1,3,6-trioxocane)s by supercritical fluid chromatography 589(1992)295

Analysis of block copolymers by high-performance liquid chromatography under critical conditions 593(1992)233

Determination of molecular mass distributions of whey protein hydrolysates by high-performance size-exclusion chromatography 599(1992)205

Molecular modelling

Chromatographic separation and molecular modelling of triazines with respect to their inhibition of the growth of L1210/R71 cells 592(1992)59

Molecular recognition

Trypsin and affinity chromatography (Review) 597(1992)3

Immobilized metal ion affinity electrophoresis. A study with several model proteins containing histidine 597(1992)357

Molecular shape

Gas chromatographic retention behaviour of dibenzothiophene derivatives on a smectic liquid crystalline polysiloxane stationary phase 590(1992)297

Molecular sieves

Dynamic gas-solid chromatographic techniques for characterizing carbon molecular sieves 556(1991)433

Molecular structure

Effect of temperature and mobile phase on the retention of retinoates in reversed-phase liquid chromatography 592(1992)127

Momordica charantia

Method for detecting the lectin activity of *Momordica charantia* transferred from micro two-dimensional electrophoretic gel on to nitrocellulose 570(1991)399

Monomeric phases

Evaluation of reversed-phase liquid chromatographic columns for recovery and selectivity of selected carotenoids 595(1992)89

Monte Carlo simulation

Selection of robust combinations of extraction liquid composition and internal standard. Monte Carlo simulation of improvement of assay methods with liquid-liquid extraction prior to high-performance liquid chromatography 594(1992)45

Mosquito repellent efficiency

Separation of *cis* and *trans* isomers from a mosquito repellent, CIC-4, via semi-preparative high-performance liquid chromatography and the repellent effect of each 590(1992)133

Moving-bed adsorption

Optical resolution by simulated moving-bed adsorption technology 590(1992)113

Moving belt interface

Use of non-volatile ion-pairing agents for liquid chromatographic-mass spectrometric analyses with a moving-belt interface 554(1991)281

Mukaiyama reaction

Analyse par chromatographie en phase gazeuse de la réaction de macrolactonisation de Mukaiyama 557(1991)501

Multi-component adsorption

Theory of perfusion chromatography 599(1992)87

Multi-component distribution function

Modelling of displacement chromatography using non-ideal isotherms 594(1992)9

Multi-criteria decision making

Optimization of the direct enantiomeric separation by high-performance liquid chromatography of the D-2 dopamine agonist N-0437 using response surface methodology and the multi-criteria decision-making method 558(1991)333

Multidimensional scaling analysis

Use of oxidative degradation followed by capillary gas chromatography-mass spectrometry and multi-dimensional scaling analysis to fingerprint unresolved complex mixtures of hydrocarbons 589(1992)281

Multi-legged phenyl bonded silicas

Liquid chromatographic separation of all-carbon molecules C₆₀ and C₇₀ with multi-legged phenyl group bonded silica phases 594(1992)105

Multi-residue determinations

Determination of National Survey of Pesticides analytes in groundwater by liquid chromatography with postcolumn reaction detection 592(1992)283

Multiple development

Application of high-performance thin-layer chromatography and automated multiple development for the identification and determination of pesticides in water 553(1991)489

Multiple peaks

Origin and control of multi-peak formation in the analysis of trimethylsilyl derivatives of flavanone aglycones by capillary column gas chromatography 586(1991)323

Multivariate analysis

Characterization of hydrophobic interaction and hydrophobic interaction chromatography media by multivariate analysis 599(1992)131

Multiwavelength detection

Assessment of chromatographic peak purity by means of multi-wavelength detection and correlation-based algorithms 592(1992)27

Murine peritoneal macrophages

Purine and pyrimidine compounds in murine peritoneal macrophages cultured *in vitro* 553(1991)205

Mussels

High-performance liquid chromatography of okadaic acid and free fatty acids in mussels 594(1992)137

Determination of organotin species by capillary gas chromatography with alternating current plasma emission detection 598(1992)97

Mustard cotyledons

Analysis of alanine and aspartate aminotransferase isoforms in mustard (*Sinapis alba* L.) cotyledons 587(1991)101

Mustard oil care

Role of metal ions in the ligand-exchange separation of amino acids 596(1992)281

N

Nafion joints

Capillary electrophoresis with electrochemical detection employing an on-column Nafion joint 593(1992)305

Neutron activation analysis

Role of ion-exchange and extraction chromatography in neutron activation analysis (Review) 600(1992)17

Noise evaluation

Peak-shape analysis and noise evaluation in suppressed ion chromatography for ultra-trace ion analysis 556(1991)249

Non-ideal isotherms

Modelling of displacement chromatography using non-ideal isotherms 594(1992)9

Non-linear behaviour

Unusual effect of temperature on the retention of enantiomers on a chiral column 558(1991)1

Glucose-fructose equilibria on Dowex Monosphere 99 CA resin under overloaded conditions 590(1992)147

Non-linear chromatography

Comparison of various isotherm models for predicting competitive adsorption data 552(1991)71

Precision and accuracy of the gas-solid adsorption isotherms derived by the elution-by-characteristic-points method 591(1992)245

Modelling of displacement chromatography using non-ideal isotherms 594(1992)9

Computer simulation of weak affinity chromatography 597(1992)83

Non-linearity

Quantitation and linearity for particle-beam liquid chromatography-mass spectrometry 554(1991)103

Non-linear multilevel calibration

Comparative analysis of thirty polychlorinated biphenyl congeners on two capillary columns of different polarity with non-linear multi-level calibration 558(1991)375

Non-mesomorphic solutes

Thermodynamics of solution of non-mesomorphic solutes at infinite dilution in the smectic-A, nematic and isotropic phases of *p*-*n*-octyl-*p*'-cyanobiphenyl. A gas-liquid chromatographic study 556(1991)317

Non-polar interactions

Influence of solute size and the non-polar interaction term on the selection of test solutes for the classification of stationary phase selectivity in gas chromatography 556(1991)457

Non-specific interaction

Biosep-SEC-S high-performance size-exclusion chromatographic columns for proteins and peptides 599(1992)25

Non-stop acquisition

Chiroptical detection during liquid chromatography. III. Non-stop acquisition of circular dichroism spectra during liquid chromatography 586(1991)249

Nuclear magnetic resonance spectrometry

Structural investigation of oligomeric *n*-octylsilyl reversed phases 557(1991)515

Correlations between high-performance liquid chromatographic retention, X-ray structural and ¹³C NMR spectroscopic data of flavonoid compounds 592(1992)65

Separation of synthetic phosphorothioate oligodeoxynucleotides from their oxygenated (phosphodiester) defect species by strong-anion-exchange high-performance liquid chromatography 599(1992)35

Nuclear magnetic resonance spectrometry, solid-state

Physical and chemical characterization of a porous phosphate-modified zirconia substrate 587(1991)137

Nuclear medicine

Determination of the mean ionic charge of the components of three ^{99m}Tc bone scanning agents by gel chromatography with two eluents of different electrolyte composition 594(1992)217

Purity assay of sodium mercaptododecaborate by high-performance liquid chromatography 595(1992)169

Nutritive media

Rapid identification of microbial starch degradation products from a complex nutrient medium by a thin-layer chromatographic method 558(1991)328



Octahedral complexes

Simultaneous effect of geometrical isomerism and chelate ring size of tris(aminocarboxylato)cobalt(III) complexes on their behaviour in thin-layer chromatography on silica gel and alumina 585(1991)181

n-Octylsilyl-silica

Structural investigation of oligomeric *n*-octylsilyl reversed phases 557(1991)515

Oil

Gas chromatography and gas chromatography-mass spectrometry study of hydrocarbons in Vlasta oil (Adriatic Basin) as the basis for geochemical interpretation 557(1991)255

Use of oxidative degradation followed by capillary gas chromatography-mass spectrometry and multi-dimensional scaling analysis to fingerprint unresolved complex mixtures of hydrocarbons 589(1992)281

Oligotrophic soil bacteria

Capillary gas chromatography-mass spectrometry of unusual and very long-chain fatty acids from soil oligotrophic bacteria 558(1991)215

Olive oil

Reliability assessment of a gas chromatographic method for polycyclic aromatic hydrocarbons in olive oil 555(1991)211

Olives

Determination of chlorophylls and carotenoids by high-performance liquid chromatography during olive lactic fermentation 585(1991)259

On-column derivatization

High-performance liquid chromatography of histamine and 1-methylhistamine with on-column fluorescence derivatization 595(1992)163

On-line sample processing

Coupled-column high-performance liquid chromatographic method for the determination of 1-hydroxypyrene in urine of subjects exposed to polycyclic aromatic hydrocarbons 600(1992)189

Open-tubular capillary columns

see also Capillary columns

High-temperature open-tubular capillary column liquid chromatography 592(1992)239

Reversed- and normal-phase separations by high-temperature open-tubular column liquid chromatography 598(1992)153

Operating parameters

Influence of operating parameters on the preparative gradient elution chromatography of insulins 599(1992)195

Optical resolution

Optical resolution by simulated moving-bed adsorption technology 590(1992)113

Optimization

Some aspects of optimization in planar chromatography (Review) 556(1991)81

Optimisation pentaparamétrique de la résolution en chromatographie phase gazeuse par la technique du diagramme à fenêtres. Cas du couplage de deux colonnes 556(1991)287

Optimization of the gas chromatographic analysis of a standard mixture of polychlorodibenzo-*p*-dioxins and polychlorodibenzofurans 557(1991)481

Optimization of the direct enantiomeric separation by high-performance liquid chromatography of the D-2 dopamine agonist N-0437 using response surface methodology and the multi-criteria decision-making method 558(1991)333

Simplex optimization of liquid chromatography with the function of mutual information as a criterion 585(1991)187

Simultaneous optimization of several chromatographic performance goals using Derringer's desirability function 586(1991)11

Multiparameter optimizations in micellar liquid chromatography using the iterative regression optimization strategy 586(1991)221

Optimization of separation of rare earths in high-performance thin-layer chromatography 587(1991)359

Feasibility study for the construction of an integrated expert system in high-performance liquid chromatography 589(1992)31

Optimization of an ion-exchange high-performance liquid chromatographic method for the determination of carboxylic acids, sugars, glycerol and ethanol in wines 589(1992)151

Accuracy of computer-simulated gas chromatographic separations based on a linear elution strength model 589(1992)215

Theory of optimization of ideal displacement chromatography of binary mixtures 590(1992)3

Optimization of the experimental conditions and the column design parameters in overloaded elution chromatography 591(1992)31

Optimization of the separation of (6*R*)- and (6*S*)-leucovorin and evaluation of the robustness of the optimum 592(1992)37

Use of solvent selectivity optimization procedures for high-performance liquid chromatographic method development 592(1992)43

Modelling retention in reversed-phase liquid chromatography as a function of pH and solvent composition 592(1992)157

Retention behaviour of closely related coumarins in thin-layer chromatographic preassays for high-performance liquid chromatography according to the "PRISMA" model 593(1992)329

Computer simulation as a tool for the rapid optimization of the high-performance liquid chromatographic separation of a tryptic digest of human growth hormone 594(1992)65

Optimization of the separation of the Rp and Sp diastereomers of phosphate-methylated DNA and RNA dinucleotides 596(1992)59

Systematic optimization of capillary electrophoretic separation of sulphonamides 598(1992)133

Optimization criteria

CRISEBOOK, a Hypermedia version of an expert system for the selection of optimization criteria in high-performance liquid chromatography 592(1992)51

Organic modifiers

Ion-mobility spectrometry as a detection method for packed-column supercritical fluid chromatography 552(1991)551

Description of retention mechanism by solvophobic theory. Influence of organic modifiers on the retention behaviour of homologous series in reversed-phase liquid chromatography 556(1991)219

Retention data for five ketotrichothecenes in reversed-phase high-performance liquid chromatography with different eluent systems 558(1991)81

Retention behavior of β -cyclodextrin complexes of anthracene and pyrene using reversed-phase liquid chromatography. The effects of *tert*.-butanol and cyclopentanol 594(1992)37

Organochlorosilane pathway

Synthesis of chemically bonded liquid crystals for high-performance liquid chromatography. New phases via the organochlorosilane pathway 556(1991)373

Ortho effect

Prediction of retention indexes. II. Structure-retention index relationship on polar columns 586(1991)85

Overlapping zones

Comparison of statistical theories of spot overlap in two-dimensional separations and verification of means for estimating the number of zones 591(1992)1

Overloaded band profiles

Precision and accuracy of the gas-solid adsorption isotherms derived by the elution-by-characteristic-points method 591(1992)245

Overloaded conditions

Glucose-fructose equilibria on Dowex Monosphere 99 CA resin under overloaded conditions 590(1992)147

Preparative liquid chromatography. II. Existence of optimum injection conditions for overloaded gradient elution separations 596(1992)15

Overloaded elution

Experimental study on the effect of the sample size on the band profile for a binary mixture showing no competitive interaction 556(1991)195

Overloaded elution chromatography

Prediction of single and binary profiles in overloaded elution chromatography using various semi-ideal models 556(1991)205

Optimization of the experimental conditions and the column design parameters in overloaded elution chromatography 591(1992)31

Overpressured layer chromatography

Separation and preparative isolation of phenolic dialdehydes by on-line overpressured layer chromatography 590(1992)369

Ovomucoid-conjugated polymer column

Application of an ovomucoid-conjugated polymer column for the enantiospecific determination of chlorprenaline concentrations in plasma 566(1991)163

Oxidative degradation

Use of oxidative degradation followed by capillary gas chromatography-mass spectrometry and multi-dimensional scaling analysis to fingerprint unresolved complex mixtures of hydrocarbons 589(1992)281

P**Packed capillary columns**

Electrokinetic reversed-phase chromatography with packed capillaries 593(1992)313

Packing density distribution

Experimental evaluation of sorbent bed homogeneity in high-performance liquid chromatographic columns 592(1992)109

Pairing ions

Determination of aminoglycoside antibiotics by reversed-phase ion-pair high-performance liquid chromatography coupled with pulsed amperometry and ion spray mass spectrometry 591(1992)195

Palaeoenvironmental data

Determination of major ions in snow and ice cores by ion chromatography 594(1992)225

Palladium catalysts

Liquid chromatographic isolation and structural elucidation of methoxymethylene dimethyl dodecatrienes catalytically synthesized from methanol and three moles of isoprene 594(1992)111

***Papaver rhoeas* L.**

Analytical studies of isorhoeadine and rhoeagenine in petal extracts of *Papaver rhoeas* L. using high-performance liquid chromatography 596(1992)276

Para effect

Prediction of retention indexes. II. Structure-retention index relationship on polar columns 586(1991)85

Particle beam mass spectrometry

Quantitation and linearity for particle-beam liquid chromatography-mass spectrometry 554(1991)103

Some aspects of peak broadening in particle-beam liquid chromatography-mass spectrometry 554(1991)119

On-line derivatization of eluted substances in dynamic high-performance liquid chromatography-mass spectrometry through the particle-beam interface 554(1991)125

Characterization of N-acyl-D-biotinols by particle-beam liquid chromatography-mass spectrometry. An alternative to probe mass spectrometry for thermally labile samples 554(1991)141

Particle beam liquid chromatography-mass spectrometry on a double-focusing sector instrument 562(1991)13

Determination of aromatic sulfonic acids in aqueous environmental samples by anion-exchange chromatography coupled to particle beam mass spectrometry and UV spectrophotometry 589(1991)177

Particle-induced X-ray emission

Study of isotachophoretic separation behaviour of metal cations by means of particle-induced X-ray emission. II. Separation efficiency of lanthanoid model mixture 589(1992)339

Study of isotachophoretic separation behaviour of metal cations by means of particle-induced X-ray emission. III. Analysis of a crude rare earth chloride from monazite 594(1992)371

Particle size

Fast protein separation by reversed-phase high-performance liquid chromatography on octadecylsilyl-bonded non-porous silica gel. Effect of particle size of column packing on column efficiency 585(1991)207

Utility of small-particle silica in preparative chromatography 590(1992)77

Optical resolution by simulated moving-bed adsorption technology 590(1992)113

Computer simulation of weak affinity chromatography 597(1992)83

Partition behaviour

Extraction of xanthine oxidase from milk by counter-current distribution in an aqueous two-phase system 558(1991)357

Partition coefficients

Application of solute and mobile phase partition coefficient to describe solute retention in reversed-phase high-performance liquid chromatography 555(1991)21

Measurement of partition coefficients by various centrifugal partition chromatographic techniques. A comparative evaluation 556(1991)181

Influence of mobile phase composition on evaluation of lipophilicity by partition chromatography 557(1991)399

Hydrophobicity of β -lactam antibiotics. Explanation and prediction of their behaviour in various partitioning solvent systems and reversed-phase chromatography 585(1991)3

Predicting partition coefficients in polyethylene glycol-potassium phosphate aqueous two-phase systems 586(1991)341

Hydrogen bonding. XVII. The characterisation of 24 gas-liquid chromatographic stationary phases studied by Poole and co-workers, including molten salts, and evaluation of solute-stationary phase interactions 587(1991)229

Hydrogen bonding. XIX. The characterisation of two poly (methylphenylsiloxane)s 588(1991)361

Determination of the mean ionic charge of the components of three ^{99m}Tc bone scanning agents by gel chromatography with two eluents of different electrolyte composition 594(1992)217

Partitioning

Comparative study of chromatography on thin layers impregnated with organic stationary phases. Chromatographic separation of nitrophenols 558(1991)415

Adsorption and partition mode in high-performance liquid chromatography of highly polar solutes on silica 552(1991)23

Ligand-receptor interactions in affinity cell partitioning. Studies with transferrin covalently linked to monomethoxypoly (ethylene glycol) and rat reticulocytes 594(1992)97

Partition ratio

Validation of chromatographic retention models in reversed-phase high-performance liquid chromatography by fitting experimental data to the relevant equations 595(1992)53

Path analysis

New approach for calculating ideal chromatograms from arbitrary composite distribution isotherms 556(1991)95

Peak ...

see also Band ...

Peak area

Deconvolution method for accurate determination of overlapping peak areas in chromatograms 587(1991)129

Peak asymmetry factors

Retention of basic drugs on porous polymers in high-performance liquid chromatography 600(1992)115

Peak broadening

Some aspects of peak broadening in particle-beam liquid chromatography-mass spectrometry 554(1991)119

Limits and effects of precolumn addition of thioglycerol in liquid chromatographic-fast atom bombardment mass spectrometric systems 591(1992)129

Peak capacity

Peak capacity of columns for size-exclusion chromatography 591(1992)47

Computer simulation of weak affinity chromatography 597(1992)83

Peak compression

Optimization of a peak compression system for a remoxipride metabolite (FLA797) and its application to bioanalysis 591(1992)207

Peak dispersion

Peak dispersion and separation efficiency in high-performance zone electrophoresis with gel-filled capillaries 595(1992)313

Peak distortion

Peak distortion in the column liquid chromatographic determination of omeprazole dissolved in borax buffer 586(1991)271

Peak identification

Determination of psychotropic phenylalkylamine derivatives in biological matrices by high-performance liquid chromatography with photodiode-array detection 593(1992)87

Peak overlap

Quantitative resolution of severely overlapping gas chromatographic peaks. Isothermal and temperature-programmed operation 556(1991)305

Improved algorithm for resolution of overlapped asymmetric chromatographic peaks 591(1992)19

Peak profiles

Improved algorithm for resolution of overlapped asymmetric chromatographic peaks 591(1992)19

Peak purity

Peak purity, yield and throughput in preparative liquid chromatography with self-displacement 592(1992)17

Assessment of chromatographic peak purity by means of multi-wavelength detection and correlation-based algorithms 592(1992)27

Determination of psychotropic phenylalkylamine derivatives in biological matrices by high-performance liquid chromatography with photodiode-array detection 593(1992)87

Peak separation

Deconvolution method for accurate determination of overlapping peak areas in chromatograms 587(1991)129

Peak shape

see also Band profiles

Gas chromatography in qualitative analysis. XX. The deactivation of diatomaceous supports by batyl alcohol 552(1991)169

Possible role for stationary phase metal interactions in the chromatography of hydroxyamines on silica 552(1991)429

Effects of modifiers in packed and open-tubular supercritical fluid chromatography 552(1991)527

Improved computer algorithm for characterizing skewed chromatographic band broadening. I. Method 556(1991)111

Peak-shape analysis and noise evaluation in suppressed ion chromatography for ultra-trace ion analysis 556(1991)249

Activity testing and surface characterization of pretreated fused-silica capillaries for gas chromatography. A new modification of existing intermediate column tests 590(1992)271

Improved equations for the calculation of chromatographic figures of merit for ideal and skewed chromatographic peaks 594(1992)1

Peak splitting

Peak distortion in the column liquid chromatographic determination of omeprazole dissolved in borax buffer 586(1991)271

Pellicular particles

Beaded polymer supports and gels (Review). I. Manufacturing techniques 586(1991)181

Pellicular resins

Application of a pellicular anion-exchange resin to the separation of inorganic and organic anions by single-column anion exchange 592(1992)101

Peptide-lipid interactions

Interaction of amphipathic model lipopeptides with phospholipid bilayers 597(1992)293

Peptide mapping

Improvement of extraction and concentration of milk peptides with solid-phase cartridges for analysis by high-performance liquid chromatography 553(1991)155

Application of capillary reversed-phase high-performance liquid chromatography to high-sensitivity protein sequence analysis 599(1992)119

Peptide maps of five human pepsin isoenzymes and other aspartic proteinases 599(1992)179

Perfluorocarbon emulsion

Novel affinity separations based on perfluorocarbon emulsions. Use of a perfluorocarbon affinity emulsion for the purification of human serum albumin from blood plasma in a fluidised bed 597(1992)189

Performance concept

High-speed gas chromatography. Theoretical and practical aspects 556(1991)331

Perfusion chromatography

Theory of perfusion chromatography 599(1992)87

Permanganate-impregnated capillary columns

Permanganate-impregnated packed capillary columns for group separation of triacylglycerols using supercritical media as mobile phases 600(1992)358

Peroxyoxalate chemiluminescence

Sensitive liquid chromatographic determination of alkyl-, nitro- and chlorophenols by precolumn derivatization with dansyl chloride, postcolumn photolysis and peroxyoxalate chemiluminescence detection 553(1991)345

Petroleum products

Column switching for the high-performance liquid chromatographic analysis of polynuclear aromatic hydrocarbons in petroleum products 552(1991)575

Pharmaceutical preparations

Collaborative study of the determination of cloxacillin by column liquid chromatography 600(1992)333

Phase equilibria

Prediction of retention in gas-liquid chromatography using the UNIFAC group contribution method. II. Polymer stationary phases 585(1991)83

Phase transitions

Study of salt hydrates by gas-solid chromatography 557(1991)495

pH effects

Affinity partitioning of erythrocytic phosphofructokinase in aqueous two-phase systems containing poly(ethylene glycol)-bound Cibacron Blue. Influence of pH, ionic strength and substrates/ effectors 589(1992)127

Buffer and pH dependence of the retention of phenylthiocarbamylamino acids in reversed-phase high-performance liquid chromatography 600(1992)87

Phenylazo-8-quinolinol silica gel

Chromatographic effects of residual amino groups on liquid chromatographic stationary phases derived from aminopropyl-silica gel 590(1992)189

pH gradients

Peak distortion in the column liquid chromatographic determination of omeprazole dissolved in borax buffer 586(1991)271

Radial pH distribution during capillary electrophoresis with electroosmotic flow. Analysis with high ionic strength buffers 589(1992)315

Isoelectric focusing field-flow fractionation. III. Investigation of the influence of different experimental parameters on focusing of cytochrome *c* in the trapezoidal cross-section channel 600(1992)297

Phosphorus-selective detection

Analysis of nucleotides by high-performance liquid chromatography with phosphorus-selective detection 557(1991)441

Photobleaching

Diffusion of proteins in the chromatographic gel AcA-34 558(1991)71

Photoemissive ionization source

Photoemissive ionisation source for ion mobility detectors
588(1991)239

Photoionization detector

Photoionization assessment of C₃-C₅ alkadienes and alkenes
in urban air 591(1992)358

Photoisomerization

Studies on photoisomerization of 4,4'-diaminostilbene-2,2'-
disulphonic acid for quality assurance by high-performance
liquid chromatography 596(1992)127

Photolysis

Determination of National Survey of Pesticides analytes in
groundwater by liquid chromatography with postcolumn
reaction detection 592(1992)283

Photopolymerization

Kinetics of acrylamide photopolymerization as investigated by
capillary zone electrophoresis 598(1992)277

Physicochemical criteria

Beaded polymer supports and gels (Review). II. Physico-
chemical criteria and functionalization 586(1991)199

Physicochemical parameters

High-performance liquid chromatographic study of the
correlation between the physico-chemical parameters of furan
and benzene aldehydes and the rate of their metabolism by
yeast 594(1992)23

Phytoplankton biomass

Highly sensitive determination of photosynthetic pigments in
marine *in situ* samples by high-performance liquid
chromatography 558(1991)131

 π -Acceptors

New chiral covalently bonded, π -donor stationary phases for
high-performance liquid chromatography, based on derivatives
of optically active 1-(1-naphthyl)ethylamine. Part II
585(1991)195

Silica-bonded chiral stationary phases with structurally simple
 π -donor chiral selectors for high-performance liquid
chromatography 589(1992)53

Pickering lithium gradient system

Chromatography of 99 amino acids and other ninhydrin-
reactive compounds in the Pickering lithium gradient system
594(1992)165

 π -Donors

Silica-bonded chiral stationary phases with structurally simple
 π -donor chiral selectors for high-performance liquid
chromatography 589(1992)53

Planar chips technology

Planar chips technology for miniaturization and integration of
separation techniques into monitoring systems. Capillary
electrophoresis on a chip 593(1992)253

Plasma desorption mass spectrometry

Performances and limits of plasma desorption mass
spectrometry in the primary structure determination of
proteins 557(1991)359

Structural analysis of biologically active peptides and
recombinant proteins and their modified counterparts by mass
spectrometry 562(1991)377

Plasmon resonance detector

Biospecific interaction analysis using surface plasmon
resonance detection applied to kinetic, binding site and
concentration analysis 597(1992)397

Plate height

Anion-exchange chromatographic properties of α -lactalbumin
eluted from quaternized polyvinylimidazole. Study of the role
of the polymer coating 553(1991)165

Analysis of the separability of plate height into overload and
intrinsic contributions using the kinetic model of non-linear
chromatography 556(1991)159

Effect of hydrostatic flow on the efficiency in capillary
electrophoresis 559(1991)81

Plate number

Comparison of high-performance liquid chromatography with
capillary gel electrophoresis in single-base resolution of
polynucleotides 558(1991)273

Computer simulation for the prediction of separation as a
function of pH for reversed-phase high-performance liquid
chromatography. II. Resolution as a function of simultaneous
change in pH and solvent strength 592(1992)197

***Pneumocystis carinii* cysts**

Elution mode of *Pneumocystis carinii* cysts in gravitational
field-flow fractionation 579(1992)143

Poiseuille flow

Non-equilibrium integral Doppler anemometric analysis of
particle mixtures in a channel flow using an intrinsic
hydrodynamic focusing force biased by another force
553(1991)517

Polarity

Attempts to use Laffort's solubility factors as polarity
parameters for organic compounds in inverse gas
chromatography 555(1991)205

Quantitative aspects of the determination of compounds with
widely varying polarity using capillary supercritical fluid
chromatography 557(1991)99

Simulated distillation of distillates on capillary columns:
influence of the polarity of the stationary phase 557(1991)155

Correlations between high-performance liquid chromatographic retention, X-ray structural and ^{13}C NMR spectroscopic data of flavonoid compounds 592(1992)65

Polarizability

Hydrogen bonding. XVI. A new solute solvation parameter, π^{H}_2 , from gas chromatographic data 587(1991)213

Pollution parameters

Theoretical analysis for measurement of building pollution parameters by gas chromatography 557(1991)469

Polyacrylamide gels

Versatile modular microelectrophoresis system and the simultaneous preparation of large numbers of polyacrylamide gels 563(1991)184

Polyacrylamide gel polymerization under non-oxidizing conditions, as monitored by capillary zone electrophoresis 598(1992)287

Polyacrylamide slab gels

Improved procedure for the drying and storage of polyacrylamide slab gels 582(1992)249

Polydispersity

On-line characterization of polyethylene glycol-modified proteins 588(1991)125

Predictability of chromatographic protein separations. Study of size-exclusion media with narrow particle size distributions 589(1992)71

Polyester fibres

Analysis of dyes extracted from textile fibers by thermospray high-performance liquid chromatography-mass spectrometry 586(1991)73

Polyethylene glycol phases

Gas chromatographic properties of immobilized poly(ethylene glycol) stationary phases 588(1991)225

Polyglycidyl methacrylate disks

High-performance membrane chromatography of serum and plasma membrane proteins 590(1992)59

Polymer backbone

Beaded polymer supports and gels (Review). II. Physico-chemical criteria and functionalization 586(1991)199

Polymer capillaries

Determination of solute-polymer interaction constants via chromatography with polymer capillaries 594(1992)29

Polymer-coated silica

Anion-exchange chromatographic properties of α -lactalbumin eluted from quaternized polyvinylimidazole. Study of the role of the polymer coating 553(1991)165

High-performance liquid chromatography on polypyrrole-modified silica 588(1991)25

Polymer-coated reversed-phase packings with controlled hydrophobic properties. I. Effect on the selectivity of protein separations 599(1992)65

Polymer coatings

High-performance liquid chromatography on polypyrrole-modified silica 588(1991)25

Polymeric phases

Evaluation of reversed-phase liquid chromatographic columns for recovery and selectivity of selected carotenoids 595(1992)89

Polymeric reagents

Secondary equilibrium size-exclusion chromatography of ions with polymeric mobile phase additives 586(1991)277

Polymerization

Analysis of benzyloxy-terminated poly(1,3,6-trioxocane)s by supercritical fluid chromatography 589(1992)295

Kinetics of acrylamide photopolymerization as investigated by capillary zone electrophoresis 598(1992)277

Polyacrylamide gel polymerization under non-oxidizing conditions, as monitored by capillary zone electrophoresis 598(1992)287

Polymer phase systems

Counter-current chromatography of lipoproteins with a polymer phase system using the cross-axis synchronous coil planet centrifuge 596(1992)118

Polymethacrylate-coated silica

Polymer-coated reversed-phase packings with controlled hydrophobic properties. II. Effect on the selectivity of peptide separations 599(1992)77

Polynomial equations

Relationship between gas chromatographic retention indices and molecular connectivity indices of chlorinated pesticides and structurally related compounds 598(1992)243

Polystyrene, phosphorylated

Phosphorylated polystyrene resins in high-performance ion-exchange chromatography 589(1992)87

Poly(styrene-divinylbenzene)

Chromatography on Poly-RP and its cyano and diol derivatives using both polar and non-polar solvent systems 591(1992)89

High-performance liquid chromatographic separation of detritylated oligonucleotides on highly cross-linked poly(styrene-divinylbenzene) particles 599(1992)113

Poly(styrene-divinylbenzene), chemically modified

Chemically modified resins for solid-phase extraction
590(1992)197

Pore fraction

Peak capacity of columns for size-exclusion chromatography
591(1992)47

Pore size

High-performance affinity chromatography of DNA. II.
Porosity effects 600(1992)221

Pore size distribution

Characterization of synthetic macroporous packing materials
in low-pressure cartridges and columns 590(1992)83

Porosity

Beaded polymer supports and gels (Review). II. Physico-
chemical criteria and functionalization 586(1991)199

Porosity effects

High-performance affinity chromatography of DNA. II.
Porosity effects 600(1992)221

Porous graphitic carbon

High-performance liquid chromatographic retention behaviour
of ring-substituted aniline derivatives on a porous graphitized
carbon column 600(1992)43

Trace-level determination of polar phenolic compounds in
aqueous samples by high-performance liquid chromatography
and on-line preconcentration on porous graphitic carbon
600(1992)195

Porous silica

Immobilization of Fv antibody fragments on porous silica and
their utility in affinity chromatography 587(1991)161

Portunus pelagicus

Chromatographic analysis of low-molecular-mass copper-
binding ligands from the crab species *Scylla serrata* and
Portunus pelagicus 599(1992)21

Positional isomers

Use of methyl oxime derivatives to enhance structural
information in thermospray high-performance liquid
chromatography-mass spectrometry. Analysis of linoleic acid
lipoxygenase metabolites in maize embryos 554(1991)155

Chromatographic behaviour of the *closo*-[B₁₂H₁₂]²⁻
derivatives on hydroxyethylmethacrylate gels 588(1991)201

Post-column extraction

Evaluation of eluents in thermospray liquid chromatography-
mass spectrometry for identification and determination of
pesticides in environmental samples 553(1991)311

Post-column photolysis

Sensitive liquid chromatographic determination of alkyl-,
nitro- and chlorophenols by precolumn derivatization with
dansyl chloride, postcolumn photolysis and peroxyoxalate
chemiluminescence detection 553(1991)345

Post-column reactors

High-performance liquid chromatography post-column
derivatization with fluorescence detection to study the
influence of ambroxol on dipalmitoylphosphatidylcholine
levels in rabbit eustachian tube washings 553(1991)1

Selective post-column fluorogenic reaction with benzamidine
for trace level detection of reducing saccharides in liquid
chromatography 553(1991)255

Analysis of nucleotides by high-performance liquid
chromatography with phosphorus-selective detection
557(1991)441

Comparative studies on the high-performance liquid
chromatographic determination of thiamine and its phosphate
esters with chloroethylthiamine as an internal standard using
pre- and post-column derivatization procedures 558(1991)115

Separation of metal complexes of ethylenediaminetetraacetic
acid in environmental water samples by ion chromatography
with UV and potentiometric detection 558(1991)181

Characterization of cellulase-based enzyme reactors for the
high-performance liquid chromatographic determination of
 β -D-glucan oligosaccharides 558(1991)343

Determination of polyphosphates in intermediate materials for
detergent manufacture by ion high-performance liquid
chromatography with post-column derivatization
585(1991)267

Detection of zearalenone in cereal extracts using high-
performance liquid chromatography with post-column
derivatization 588(1991)47

Determination of organic acids, amino acids and saccharides
by high-performance liquid chromatography and a postcolumn
enzyme reactor with amperometric detection 591(1992)165

Determination of National Survey of Pesticides analytes in
groundwater by liquid chromatography with postcolumn
reaction detection 592(1992)283

High-performance liquid chromatographic determination of
monohydroxy compounds by a combination of pre-column
derivatization and post-column reaction detection
593(1992)29

Determination of inositol polyphosphates from human T-
lymphocyte cell lines by anion-exchange high-performance
liquid chromatography and post-column derivatization
593(1992)157

High-performance liquid chromatography and post-column derivatization with diphenyl-1-pyrenylphosphine for fluorimetric determination of triacylglycerol hydroperoxides 596(1992)197

Determination of divalent trace metals in natural waters by preconcentration on N,N,N',N'-tetra(2-aminoethyl)ethylenediamine-silica followed by on-line ion chromatography 600(1992)203

³²P-Postlabelling

³²P-Postlabelling and mass spectrometric methods for analysis of bulky, polyaromatic carcinogen-DNA adducts in humans (Review) 580(1992)293

Potential barrier

Analysis of pH-dependent protein interactions with gel filtration medium 591(1992)121

Potentiometric detector

Mixed potential response at metallic electrodes in potentiometric detection for ion chromatography 589(1992)201

Potentiometric detection of carboxylic acids and inorganic anions in ion-exclusion chromatography using camphorsulphonic acid as eluent 589(1992)209

Preadsorbed salts

Dependence of the silanophilic effect on the concentration of preadsorbed salts and on the chemical structure of peptides in reversed-phase thin-layer chromatography 553(1991)467

Precolumn addition

Limits and effects of precolumn addition of thioglycerol in liquid chromatographic-fast atom bombardment mass spectrometric systems 591(1992)129

Precolumn derivatization

Time-resolved luminescence detection of derivatized thiols in column liquid chromatography 552(1991)625

Comparison of high-performance liquid chromatographic detection methods for thyronine and tyrosine residues in toxicological studies of the thyroid 553(1991)135

High-performance liquid chromatographic microassay for methyl ethyl ketone in urine as the 2,4-dinitrophenylhydrazone derivative 579(1992)324

High-performance liquid chromatography method for the determination of aminoglycosides based on automated precolumn derivatization with *o*-phthalaldehyde 593(1992)69

Precolumns

Determination of aflatoxin M1 using a dialysis-based immunoaffinity sample pretreatment system coupled on-line to liquid chromatography. Reusable immunoaffinity columns 589(1992)141

Activity testing and surface characterization of pretreated fused-silica capillaries for gas chromatography. A new modification of existing intermediate column tests 590(1992)271

Preconcentration methods

Interference removal in the organic trace-level analysis of aqueous environmental samples by on-line liquid chromatographic preconcentration techniques with two precolumns 553(1991)329

Rapid on-line precolumn high-performance liquid chromatographic method for the determination of benomyl, carbendazim and aldicarb species in drinking water 555(1991)147

Peak-shape analysis and noise evaluation in suppressed ion chromatography for ultra-trace ion analysis 556(1991)249

Determination of a metabolite of the herbicide pyridate in drinking and groundwater using high-performance liquid chromatography with amperometric detection 558(1991)369

Determination of chlorotriazines in aqueous environmental samples at the ng/l level using preconcentration with a cation exchanger and on-line high-performance liquid chromatography 585(1991)67

Preconcentration technique for introducing gaseous or volatile compounds into a capillary gas chromatographic column 587(1991)338

Determination of amphetamines by high-performance liquid chromatography with ultraviolet detection. On-line pre-column derivatization with 9-fluorenylmethyl chloroformate and preconcentration 593(1992)9

Cellulose bearing covalently linked copper phthalocyanine trisulphonate as an adsorbent selective for polycyclic compounds and its use in studies of environmental mutagens and carcinogens (Review) 597(1992)37

Determination of divalent trace metals in natural waters by preconcentration on N,N,N',N'-tetra(2-aminoethyl)ethylenediamine-silica followed by on-line ion chromatography 600(1992)203

Preparative chromatography

see also Special issue index

Preparative chromatography

Preparation of a chiral matrix. Resolution of (6*R*,*S*)-N⁵-formyltetrahydrofolate 555(1991)254

Use of a Sep-Pak cartridge for preparative collection in gas chromatography 555(1991)302

Analysis of the separability of plate height into overload and intrinsic contributions using the kinetic model of non-linear chromatography 556(1991)159

- Flip-flop elution concept in preparative liquid chromatography 557(1991)137
- Preparative chromatographic separation of enantiomers (Review) 576(1992)1
- Chromatographic fractionation of nucleic acids using microcapsules made from plant cells 585(1991)57
- Isolation of doxycycline, 6-epidoxycycline and 2-acetyl-2-decarboxamidometacycline from commercial metacycline by preparative column liquid chromatography on silica gel 586(1991)67
- Preparative separation and analysis of the enantiomers of [³H]Abbott-69992, an HIV anti-infective nucleoside, by ligand-exchange high-performance liquid chromatography 586(1991)265
- Separation of functionalized dextrans by reversed-phase high-performance liquid chromatography 587(1991)288
- Effect of the sample solvent on band profiles in preparative liquid chromatography using non-aqueous reversed-phase high-performance liquid chromatography 588(1991)1
- Simple test for determination of the degree of distortion of the liquid-phase flow profile in columns for preparative liquid chromatography 589(1992)61
- Theory of optimization of ideal displacement chromatography of binary mixtures 590(1992)3
- Displacement effects in preparative gradient high-performance liquid chromatographic separations 590(1992)17
- Factors affecting the separation and loading capacity of proteins in preparative gradient elution high-performance liquid chromatography 590(1992)35
- Economic considerations important in the scale-up of an ovalbumin separation from hen egg-white on the anion-exchange cellulose DE92 590(1992)49
- Utility of small-particle silica in preparative chromatography 590(1992)77
- Characterization of synthetic macroporous packing materials in low-pressure cartridges and columns 590(1992)83
- Adsorption kinetics of albumin on a cross-linked cellulose chromatographic ion exchanger 590(1992)101
- Experimental study of the production rate of pure enantiomers from racemic mixtures 590(1992)119
- Application of preparative liquid chromatography to the isolation of enantiomers of a benzodiazepinone derivative 590(1992)127
- Purification of erucic acid by preparative high-performance liquid chromatography and crystallization 590(1992)139
- Glucose-fructose equilibria on Dowex Monosphere 99 CA resin under overloaded conditions 590(1992)147
- Isolation of experimental anti-AIDS glycerophospholipids by micro-preparative reversed-phase high-performance liquid chromatography 590(1992)153
- Preparative high-performance liquid chromatographic separation and isolation of bacitracin components and their relationship to microbiological activity 590(1992)163
- Preparative isolation of vitamin D₂ from previtamin D₂ by recycle high-performance liquid chromatography 590(1992)169
- Preparative liquid chromatography. I. Influence of column efficiency on optimum injection conditions under isocratic elution 590(1992)175
- Rapid, two-step purification process for the preparation of pyrogen-free murine immunoglobulin G₁ monoclonal antibodies 590(1992)255
- Optimization of the experimental conditions and the column design parameters in overloaded elution chromatography 591(1992)31
- Liquid chromatographic isolation and structural elucidation of methoxymethylene dimethyl dodecatrienes catalytically synthesized from methanol and three moles of isoprene 594(1992)111
- Isolation of aristolochic acids HA_sI and HA_sII by preparative liquid chromatography 594(1992)145
- Separation and purification of azetidiny methyl sulfinates using preparative high-performance liquid chromatography 595(1992)355
- Preparative liquid chromatography. II. Existence of optimum injection conditions for overloaded gradient elution separations 596(1992)15
- Membrane chromatographic systems for high-throughput protein separations 596(1992)173
- Short-cut method for predicting the productivity of affinity chromatography 597(1992)173
- Influence of operating parameters on the preparative gradient elution chromatography of insulins 599(1992)195
- Preparative isotachopheresis**
Continuous separation and purification of proteins by recycling isotachopheresis 594(1992)361

Pressure profiles

Prediction of the elution profiles of high-concentration bands in gas chromatography 589(1992)223

Pressure programming

Van Deemter-type relationship for determining the optimum initial flow-rate and optimum pressure programming rate in temperature/pressure-programmed capillary column gas chromatography utilizing separation numbers 595(1992)221

Pressure variations

Experimental evaluation of sorbent bed homogeneity in high-performance liquid chromatographic columns 592(1992)109

Pre-therapeutic screening

Use of anion-exchange chromatography and chromatofocusing to reveal the structural and functional heterogeneity of topoisomerase II in a HL-60 cell line resistant to multi-drug treatment 587(1991)3

Principal component analysis

Dependence of the silanophilic effect on the concentration of preadsorbed salts and on the chemical structure of peptides in reversed-phase thin-layer chromatography 553(1991)467

Factor analysis and experimental design in high-performance liquid chromatography. XI. Factor analysis maps and chromatographic information 556(1991)169

High-performance liquid chromatography and supercritical fluid chromatography of monosaccharides and polyols using light-scattering detection. Chemometric studies of the retentions 590(1992)203

Retention behaviour of some ring-substituted phenol derivatives on a porous graphitized carbon column 592(1992)75

Characterization of hydrophobic interaction and hydrophobic interaction chromatography media by multivariate analysis 599(1992)131

High-performance liquid chromatographic retention behaviour of ring-substituted aniline derivatives on a porous graphitized carbon column 600(1992)43

Support matrix effects in the reversed-phase thin-layer chromatography of some peptides 600(1992)149

PRISMA model

Retention behaviour of closely related coumarins in thin-layer chromatographic preassays for high-performance liquid chromatography according to the "PRISMA" model 593(1992)329

Process effluents

Monitoring the effluents of the trichloroacetic acid process by high-performance liquid chromatography 600(1992)316

Process monitoring

Flame infrared emission detection of hydrogen 589(1992)241

Process-scale separation

Economic considerations important in the scale-up of an ovalbumin separation from hen egg-white on the anion-exchange cellulose DE92 590(1992)49

Production rate

Experimental study of the production rate of pure enantiomers from racemic mixtures 590(1992)119

Optimization of the experimental conditions and the column design parameters in overloaded elution chromatography 591(1992)31

Isolation of aristolochic acids HA₃J and HA₃II by preparative liquid chromatography 594(1992)145

Productivity

Short-cut method for predicting the productivity of affinity chromatography 597(1992)173

Profiling studies

Applications of automated amino acid analysis using 9-fluorenylmethyl chloroformate 588(1991)107

Programmed temperature vaporizing injection

Gas chromatographic analysis of fatty acid methyl esters: avoiding discrimination by programmed temperature vaporizing injection 588(1991)265

Protein A column

Development and optimization of a single-step procedure using protein A affinity chromatography to isolate murine IgG₁ monoclonal antibodies from hybridoma supernatants 587(1991)61

Protein conformation

Determination of calcium ions tightly bound to proteins 585(1991)173

Protein-conjugated columns

Avidin protein-conjugated column for direct injection analysis of drug enantiomers in plasma by high-performance liquid chromatography 572(1991)133

Protein-DNA interactions

High-resolution gel filtration of the ecdysteroid receptor-DNA complex—an alternative to the electrophoretic mobility shift assay 587(1991)11

Protein hydrolysate

Determination of chloropropanols in protein hydrolysates 589(1992)109

Protein-Pak HR

Schemes for efficient protein purification on a family of polymeric ion exchangers in glass columns 599(1992)239

Protein refolding

High-performance hydrophobic interaction chromatography as a tool for protein refolding 599(1992)185

Protein sequence analysis

Performances and limits of plasma desorption mass spectrometry in the primary structure determination of proteins 557(1991)359

Pseudo-effective mobility

Determination of some drugs by micellar electrokinetic capillary chromatography. The pseudo-effective mobility as parameter for screening 585(1991)123

Pseudo-electrochromatography

Pseudo-electrochromatography-mass spectrometry: a new alternative 554(1991)339

Pseudomonas dacunhae

Application of ligand-exchange chromatography to the assay of L-alanine from DL-aspartic acid by *Pseudomonas dacunhae* 555(1991)272

Pulsed amperometric detector

Liquid chromatographic determination of ethylenethiourea using pulsed amperometric detection 586(1991)158

Constant-potential amperometric detection of carbohydrates at metal electrodes in high-performance anion-exchange chromatography 592(1992)229

Purge-and-trap methods

Analysis of industrial solvent mixtures in water using a miniature purge-and-trap device with thermal desorption and capillary gas chromatography-mass spectrometry 557(1991)429

Purification methods

Improved affinity chromatographic purification of D-mannose-N-acetyl-D-glucosamine-specific lectin from the bark of *Sophora japonica* eliminating the loss by sugar specific self-aggregation 597(1992)197

Purity assays

Purity assay of sodium mercaptododecaborate by high-performance liquid chromatography 595(1992)169

Pyrolysis

Synthesis of novel selenium-containing choline and acetylcholine analogues and their quantitation using a pyrolysis-gas chromatography-mass spectrometry assay 585(1991)101

Determination of succinic acid by pyrolysis-gas chromatography in the presence of sodium sulfite and iron powder 585(1991)345

Pyrolysis oils

Analysis of biomass pyrolysis oils by a combination of various liquid chromatographic techniques and gas chromatography-mass spectrometry 553(1991)211

Q**Quadrupole ion trap mass spectrometer**

Thermospray liquid chromatography-mass spectrometry with a quadrupole ion trap mass spectrometer 562(1991)3

Quality control

Quality control procedure for the gas chromatographic determination of light hydrocarbons in petroleum liquid 588(1991)259

Quantitation

Factors influencing the resolution and quantitation of oligonucleotides separated by capillary electrophoresis on a gel-filled capillary 559(1991)43

Quantitation, SFC

Quantitative aspects of the determination of compounds with widely varying polarity using capillary supercritical fluid chromatography 557(1991)99

Quantitative determination limit

Quantitative determination limit in chromatography: computer-based simulations 557(1991)23

R**Radiation-induced grafting**

Protein adsorption capacity of a porous phenylalanine-containing membrane based on a polyethylene matrix 586(1991)27

Radiochemical detector

Use of thermospray liquid chromatography-mass spectrometry to aid in the identification of urinary metabolites of a novel antiepileptic drug, Lamotrigine 554(1991)181

Radiochemical separation

Role of ion-exchange and extraction chromatography in neutron-activation analysis (Review) 600(1992)17

Radioimmunoassay

Selective clean-up method using an immunoaffinity column following radioimmunoassay of prostaglandin F_{2α} in biological fluids 597(1992)201

Rapid ion-exchange chromatography

Rapid ion-exchange chromatography for preparative separation of proteins. IV. Application to bovine carbonic anhydrase III from skeletal muscle 588(1991)139

Rate constants

Characterization of cellulase-based enzyme reactors for the high-performance liquid chromatographic determination of β -D-glucan oligosaccharides 558(1991)343

Rat liver microsomes

Assessment of the biotransformation of the cardiotoxic agent piroximone by high-performance liquid chromatography and gas chromatography-mass spectrometry 593(1992)1

Rat reticulocytes

Ligand-receptor interactions in affinity cell partitioning. Studies with transferrin covalently linked to monomethoxypoly(ethylene glycol) and rat reticulocytes 594(1992)97

Reaction kinetics

Reaction-gas chromatography for physico-chemical investigations on aggressive gases 552(1991)319

Reaction kinetics and kinetic processes in modern liquid chromatographic reactors (Review) 589(1992)1

Reactors

Reactor for prechromatographic fusion reactions 600(1992)352

Recovery

Methods for the analysis of thiodiglycol sulphoxide, a metabolite of sulphur mustard, in urine using gas chromatography-mass spectrometry 558(1991)393

Experimental study of the production rate of pure enantiomers from racemic mixtures 590(1992)119

Selection of robust combinations of extraction liquid composition and internal standard. Monte Carlo simulation of improvement of assay methods with liquid-liquid extraction prior to high-performance liquid chromatography 594(1992)45

Recycle chromatography

Preparative isolation of vitamin D₂ from previtamin D₂ by recycle high-performance liquid chromatography 590(1992)169

Recycling isotachopheresis

Purification of ovalbumin and lysozyme from a commercial product by recycling isotachopheresis 585(1991)145

Continuous separation and purification of proteins by recycling isotachopheresis 594(1992)361

Recycling partition equilibrium studies

Study of high-affinity interactions by quantitative affinity chromatography. Analytical expressions in terms of total ligand concentration 597(1992)57

Reduced mobilities

Ion-mobility spectrometry as a detection method for packed-column supercritical fluid chromatography 552(1991)551

Reduced plate height

Electrokinetic reversed-phase chromatography with packed capillaries 593(1992)313

Reductive electrochemical detection

Reductive electrochemical detection in liquid chromatography with a zinc amalgam scrubber column 598(1992)195

Refolding

High-performance hydrophobic interaction chromatography as a tool for protein refolding 599(1992)185

Refractive index detector

Determination of chemical composition of polymers by size-exclusion chromatography with coupled density and refractive index detection. III. Polyethylene oxide and polytetrahydrofuran 552(1991)507

Determination of phosphatidylcholine, phosphatidylglycerol and their lyso forms from liposome dispersions by high-performance liquid chromatography using high-sensitivity refractive index detection 585(1991)239

Reimer Tiemann reaction

Separation and preparative isolation of phenolic dialdehydes by on-line overpressured layer chromatography 590(1992)369

Relative capacity factors

Retention reproducibility of basic drugs in high-performance liquid chromatography on a silica column with a methanol-high-pH buffer eluent. Changes in selectivity with the age of the stationary phase 592(1992)85

Remote polar groups

Effect of superfluous remote polar functionality on chiral recognition 589(1992)45

Reordering/resolution

Use of the reordering/resolution of alkyl-modified silica to characterize the microscopic heterogeneity of silica via liquid chromatography 556(1991)415

Resin counter-ion selection

Carbohydrate separation by ligand-exchange liquid chromatography. Correlation between the formation of sugar-complexes and the elution order 558(1991)89

Resolution

Optimisation pentaparamétrique de la résolution en chromatographie phase gazeuse par la technique du diagramme à fenêtres. Cas du couplage de deux colonnes 556(1991)287

Optimization of the direct enantiomeric separation by high-performance liquid chromatography of the D-2 dopamine agonist N-0437 using response surface methodology and the multi-criteria decision-making method 558(1991)333

Factors influencing the resolution and quantitation of oligonucleotides separated by capillary electrophoresis on a gel-filled capillary 559(1991)43

Optimization of resolution in capillary zone electrophoresis: combined effect of applied voltage and buffer concentration 588(1991)315

Peak capacity of columns for size-exclusion chromatography 591(1992)47

Computer simulation for the prediction of separation as a function of pH for reversed-phase high-performance liquid chromatography. I. Accuracy of a theory-based model 592(1992)183

Direct high-performance liquid chromatographic resolution of a novel benzothiazine Ca^{2+} antagonist and related compounds 593(1992)37

Charge-reversed, polymer-coated capillary column for the analysis of a recombinant chimeric glycoprotein 594(1992)317

Critical capillary column examination of the relationship between separation number and height equivalent to a theoretical plate and their dependence on temperature 595(1992)209

Resolution mapping

Application of the gradient elution technique. Demonstration with a special test mixture and the DryLab G/plus method development software 592(1992)133

Resonance electron-capture mass spectrometry

Use of mass spectrometry for the detection and identification of bromine-containing diphenyl ethers 586(1991)360

Response factors

Quantitative aspects of indirect UV detection in capillary zone electrophoresis 588(1991)321

Relationship between zone length and step height in isotachophoresis 595(1992)327

Response surface methodology

Optimization of the direct enantiomeric separation by high-performance liquid chromatography of the D-2 dopamine agonist N-0437 using response surface methodology and the multi-criteria decision-making method 558(1991)333

Response surface modelling

Optimization of an ion-exchange high-performance liquid chromatographic method for the determination of carboxylic acids, sugars, glycerol and ethanol in wines 589(1992)151

Restricted access media

Evaluation of restricted access media for high-performance liquid chromatographic analysis of sulfonamide antibiotic residues in bovine serum 598(1992)23

Retention

Effect of stationary phase structure on retention and selectivity of restricted-access reversed-phase packing materials 558(1991)19

Retention behaviour

Application of solute and mobile phase partition coefficient to describe solute retention in reversed-phase high-performance liquid chromatography 555(1991)21

Gas chromatographic retention behaviour of some solutes on structurally similar polar and non-polar stationary phases 555(1991)191

Capillary gas chromatography of C_5 - C_{13} branched alkynes on squalane and liquid crystal stationary phases 557(1991)241

Retention behaviour of a template-assembled synthetic protein and its amphiphilic building blocks on reversed-phase columns 586(1991)43

Study of the retention behaviour of barbiturates by overpressured layer chromatography using silica gel bonded with tricaprilmethylammonium chloride 589(1992)301

Correlations between high-performance liquid chromatographic retention, X-ray structural and ^{13}C NMR spectroscopic data of flavonoid compounds 592(1992)65

Retention behaviour of some ring-substituted phenol derivatives on a porous graphitized carbon column 592(1992)75

Modelling retention in reversed-phase liquid chromatography as a function of pH and solvent composition 592(1992)157

Reversed-phase high-performance liquid chromatography of anionic and ethoxylated non-ionic surfactants and pesticides in liquid pesticide formulations 593(1992)227

Retention behaviour of closely related coumarins in thin-layer chromatographic preassays for high-performance liquid chromatography according to the "PRISMA" model 593(1992)329

Retention behavior of β -cyclodextrin complexes of anthracene and pyrene using reversed-phase liquid chromatography. The effects of *tert.*-butanol and cyclopentanol 594(1992)37

Mixed-mode hydrophilic and ionic interaction chromatography rivals reversed-phase liquid chromatography for the separation of peptides 594(1992)75

Control of the retention selectivity of rare earth octaethylporphyrins in reversed-phase high-performance liquid chromatography using amines as mobile phase additives 598(1992)73

Comparison of adsorption properties of Florisil and silica in high-performance liquid chromatography. I. Retention behaviour of monofunctional model solutes 600(1992)51

Retention data

Increasing data precision in reversed-phase liquid chromatography leads to new information level 592(1992)13

Chromatographic separation and molecular modelling of triazines with respect to their inhibition of the growth of L1210/R71 cells 592(1992)59

Retention factor

Unusual effect of temperature on the retention of enantiomers on a chiral column 558(1991)1

Retention gaps

Reversed-phase liquid chromatography coupled on-line with capillary gas chromatography. I. Introduction of large volumes of aqueous mixtures through an on-column interface 552(1991)489

Retention indices

Retentions of alkyloxazoles and alkylthiazoles and their prediction based on non-linear additivity concepts in gas chromatography 552(1991)1

Determination of sorbent polarity and selectivity by linear regression of retention indices 552(1991)31

Live retention database for identification in multi-step temperature-programmed capillary gas chromatography 552(1991)187

Gas chromatographic retention behaviour of some solutes on structurally similar polar and non-polar stationary phases 555(1991)191

Attempts to use Laffort's solubility factors as polarity parameters for organic compounds in inverse gas chromatography 555(1991)205

Method for characterization of selectivity in reversed-phase liquid chromatography. V. Calibration of the retention scale for chromatographic systems with low concentrations of organic solvents in the mobile phase 556(1991)145

Gas chromatographic-mass spectrometric analysis of tar compounds formed during pyrolysis of rice husks 562(1991)531

Determination of sulphur and nitrogen gases by gas chromatography on polystyrene porous polymer columns 585(1991)93

Prediction of retention indexes. II. Structure-retention index relationship on polar columns 586(1991)85

Prediction of retention indexes. III. Silylated derivatives of polar compounds 586(1991)113

Prediction of retention indexes. IV. Chain branching in alkylbenzene isomers with C₁₀₋₁₃ alkyl chains identified in a scintillator solvent 589(1992)231

Gas chromatographic retention behaviour of dibenzothiophene derivatives on a smectic liquid crystalline polysiloxane stationary phase 590(1992)297

Polynuclear aromatic hydrocarbon retention indices on SE-54 stationary phase of the volatile components of a coal tar pitch. Relationships between chromatographic retention and thermal reactivity 591(1992)287

Gas chromatographic elution patterns of chlorinated dioxins *versus* column polarity 594(1992)269

Gas chromatographic analysis of high-molecular-weight polynuclear aromatic hydrocarbons. I. Molecular weight 328 595(1992)247

Structure-retention index relationships for derivatized monosaccharides on non-polar gas chromatography columns 596(1992)79

Relationship between gas chromatographic retention indices and molecular connectivity indices of chlorinated pesticides and structurally related compounds 598(1992)243

Retention of basic drugs on porous polymers in high-performance liquid chromatography 600(1992)115

Retention mechanisms

Adsorption and partition mode in high-performance liquid chromatography of highly polar solutes on silica 552(1991)23

Description of retention mechanism by solvophobic theory. Influence of organic modifiers on the retention behaviour of homologous series in reversed-phase liquid chromatography 556(1991)219

High-performance liquid chromatography on dynamically modified silica. IX. Modification of silica with 3-(N,N-dimethylpalmitylammonium) propanesulphonate for reversed-phase chromatography 556(1991)353

Retention data for five ketotrichothecenes in reversed-phase high-performance liquid chromatography with different eluent systems 558(1991)81

Chromatographic characterization of a phosphate-modified zirconia support for bio-chromatographic applications 587(1991)149

Retention processes on α_1 -acid glycoprotein-bonded stationary phase 591(1992)55

Retention modelling

Modelling retention in reversed-phase liquid chromatography as a function of pH and solvent composition 592(1992)157

Retention models

Extension of the electrostatic retention model of reversed-phase ion-pair high-performance liquid chromatography to include the effect of the eluent pH 552(1991)13

Secondary equilibrium size-exclusion chromatography of ions with polymeric mobile phase additives 586(1991)277

Measurement of stability constants for complexes of divalent cations with tartaric acid by ion chromatography 589(1992)185

Validation of chromatographic retention models in reversed-phase high-performance liquid chromatography by fitting experimental data to the relevant equations 595(1992)53

Effects of stationary phase ligand density on high-performance ion-exchange chromatography of proteins 598(1992)7

Retention order

Reversed retention order and other stereoselective effects in the separation of amino alcohols on Chiralcel OD 592(1992)331

Retention prediction

Retentions of alkyloxazoles and alkylthiazoles and their prediction based on non-linear additivity concepts in gas chromatography 552(1991)1

Extension of the electrostatic retention model of reversed-phase ion-pair high-performance liquid chromatography to include the effect of the eluent pH 552(1991)13

New aspects of quantitative structure-retention relationships in chromatography 552(1991)59

Calculation of retention and selectivity in reversed-phase liquid chromatography 552(1991)91

Development of a rational optimisation procedure for the automated sample clean-up with column switching in pesticide residue analysis 552(1991)113

Experimental model for predicting solute retention time as a function of mobile phase composition in ternary reversed-phase high-performance liquid chromatography 555(1991)33

Prediction of retention in gas-liquid chromatography using the UNIFAC group contribution method. II. Polymer stationary phases 585(1991)83

UNIFAC model as a heuristic guide for estimating retention in chromatography 586(1991)1

Prediction of retention indexes. II. Structure-retention index relationship on polar columns 586(1991)85

Prediction of retention in gas-liquid chromatography using the UNIFAC group contribution method. III. Recent developments in UNIFAC 586(1991)297

Accuracy of computer-simulated gas chromatographic separations based on a linear elution strength model 589(1992)215

Computer simulation for the prediction of separation as a function of pH for reversed-phase high-performance liquid chromatography. I. Accuracy of a theory-based model 592(1992)183

Prediction of peptide retention time in reversed-phase high-performance liquid chromatography 599(1992)211

Retention properties

Retention reproducibility of basic drugs in high-performance liquid chromatography on a silica column with a methanol-high-pH buffer eluent. Changes in selectivity with the age of the stationary phase 592(1992)85

Solvatochromic hydrogen bond donor acidity of cyclodextrins and reversed-phase liquid chromatographic retention of small molecules on a β -cyclodextrin-bonded silica stationary phase 595(1992)45

Retention scale calibration

Method for characterization of selectivity in reversed-phase liquid chromatography. V. Calibration of the retention scale for chromatographic systems with low concentrations of organic solvents in the mobile phase 556(1991)145

Retention-thermal reactivity relationship

Polynuclear aromatic hydrocarbon retention indices on SE-54 stationary phase of the volatile components of a coal tar pitch. Relationships between chromatographic retention and thermal reactivity 591(1992)287

Retention times

Live retention database for identification in multi-step temperature-programmed capillary gas chromatography 552(1991)187

Secondary equilibrium size-exclusion chromatography of ions with polymeric mobile phase additives 586(1991)277

Retention volumes

Estimation of cellulose polarity by gas chromatography 552(1991)365

Determination of specific retention volumes at 20°C for hydrocarbons on microporous carbons 586(1991)161

Retention volumes, specific

Thermodynamics of solution of non-mesomorphic solutes at infinite dilution in the smectic-A, nematic and isotropic phases of *p*-*n*-octyl-*p*'-cyanobiphenyl. A gas-liquid chromatographic study 556(1991)317

Reticulation

Chromatographic separation of polyols by ligand exchange. Effects of the ion-exchange resin cross-linking and size 594(1992)125

Reversed-flow gas chromatography

Theoretical analysis for measurement of building pollution parameters by gas chromatography 557(1991)469

Activity coefficients in binary liquid mixtures measured by reversed-flow gas chromatography 595(1992)237

Reviews

Methods of equilibrium concentration for the gas chromatographic determination of trace volatiles (Review) 556(1991)1

Is chromatography a separation process? The zonoid answer (Review) 556(1991)25

Some aspects of optimization in planar chromatography (Review) 556(1991)81

Application of high-performance liquid chromatography to the study of biogenic amine-related enzymes (Review) 566(1991)287

Theoretical background for clinical and biomedical applications of electromigration techniques (Review) 569(1991)3

Biomedical relevance of two-dimensional protein mapping (Review) 569(1991)43

Capillary zone electrophoresis: its applicability and potential in biochemical analysis (Review) 569(1991)63

Application of capillary isotachopheresis in peptide analysis (Review) 569(1991)123

Staining and quantification of proteins separated by polyacrylamide gel electrophoresis (Review) 569(1991)175

Isoelectric focusing in immobilized pH gradients: applications in clinical chemistry and forensic analysis (Review) 569(1991)197

Clinical applications of lectin affinity electrophoresis (Review) 569(1991)229

Sodium dodecyl sulphate electrophoresis of urinary proteins (Review) 569(1991)243

Clinical applications of electrophoresis of human salivary proteins (Review) 569(1991)261

Analysis of cerebrospinal fluid proteins by electrophoresis (Review) 569(1991)281

Electrophoretic phenotyping of erythrocyte enzymes (Review) 569(1991)297

Electrophoresis of serum isoenzymes and proteins following acute myocardial infarction (Review) 569(1991)323

Electrophoretic analysis of enzyme-linked immunoglobulins and their clinical significance (Review) 569(1991)347

Utility of protein electrophoretic analysis in the characterization of malignant tissues (Review) 569(1991)367

Electrophoretic analysis of nuclear matrix proteins and the potential clinical applications (Review) 569(1991)389

Electrophoresis of mutant proteins in inherited diseases (Review) 569(1991)407

DNA typing in hereditary disease (Review) 569(1991)421

Preparative chromatographic separation of enantiomers (Review) 576(1992)1

Drug level monitoring: antidepressants (Review) 576(1992)183

Systematic toxicological analysis of drugs and their metabolites by gas chromatography-mass spectrometry (Review) 580(1992)3

Rapid assay of cocaine, opiates and metabolites by gas chromatography-mass spectrometry (Review) 580(1992)43

Gas chromatographic-mass spectrometric methods of analysis for detection of 11-nor- Δ^9 -tetrahydrocannabinol-9-carboxylic acid in biological matrices (Review) 580(1992)63

Determination of methamphetamine enantiomer ratios in urine by gas chromatography-mass spectrometry (Review) 580(1992)77

Chromatographic and mass spectrometric methods for determination of lysergic acid diethylamide (LSD) and metabolites in body fluids (Review) 580(1992)97

High-performance liquid chromatographic separation and detection methods for anabolic compounds (Review) 580(1992)111

Drug detection in hair by chromatographic procedures (Review) 580(1992)125

Chromatographic methods for the determination of toxicants in faeces (Review) 580(1992)135

Chromatographic methods for blood alcohol determination (Review) 580(1992)161

Chromatographic separation of marine organic pollutants (Review) 580(1992)191

Strategies for the identification of non-polar toxicants in aqueous environmental samples using toxicity-based fractionation and gas chromatography-mass spectrometry (Review) 580(1992)215

Determination of aromatic hydrocarbons and their metabolites in human blood and urine (Review) 580(1992)229

Methods for the analysis of persistent chlorinated hydrocarbons in tissues (Review) 580(1992)257

Methods for chromatographic determination of amanitins and related toxins in biological samples (Review) 580(1992)279

³²P-Postlabelling and mass spectrometric methods for analysis of bulky, polyaromatic carcinogen-DNA adducts in humans (Review) 580(1992)293

Detection of the enzymatic activity of cytochrome P-450 enzymes by high-performance liquid chromatography (Review) 580(1992)325

Beaded polymer supports and gels (Review). I. Manufacturing techniques 586(1991)181

Beaded polymer supports and gels (Review). II. Physico-chemical criteria and functionalization 586(1991)199

Reaction kinetics and kinetic processes in modern liquid chromatographic reactors (Review) 589(1992)1

Polychlorinated biphenyls in the environment (Review) 595(1992)1

Trypsin and affinity chromatography (Review) 597(1992)3

Galactosylation as a tool for the stabilization and immobilization of proteins (Review) 597(1992)19

Interactions and uses of antisense peptides in affinity technology (Review) 597(1992)29

Cellulose bearing covalently linked copper phthalocyanine trisulphonate as an adsorbent selective for polycyclic compounds and its use in studies of environmental mutagens and carcinogens (Review) 597(1992)37

Recent developments in quantitative affinity chromatography (Review) 597(1992)67

Immobilized lectin columns: useful tools for the fractionation and structural analysis of oligosaccharides (Review) 597(1992)111

Early evolution of chromatography: the activities of Charles Dhéré 600(1992)3

Role of ion-exchange and extraction chromatography in neutron activation analysis (Review) 600(1992)17

Ringer's injection

Simultaneous ion chromatographic determination of chloride and calcium contained in Ringer's injection using a chelating agent as eluent 589(1992)358

Rinsing, capillaries

Influence of operating parameters on reproducibility in capillary electrophoresis 559(1991)57

River water

Reversed-phase high-performance liquid chromatographic determination of linear alkylbenzenesulphonates in river water at ppb levels by precolumn concentration 555(1991)155

Robotic workstation

Automated high-performance liquid chromatographic and size-exclusion chromatographic sample preparation by means of a robotic workstation 592(1992)249

S

Saccharomyces cerevisiae

Analysis of sterol esters from alga and yeast by high-performance liquid chromatography and capillary gas chromatography-mass spectrometry with chemical ionization 598(1992)219

Saccharopolyspora erythraea

Isolation of erythromycin A N-oxide and pseudoerythromycin A hemiketal from fermentation broth of *Saccharopolyspora erythraea* by thin-layer and high-performance liquid chromatography 558(1991)265

Salmon

Development of a screening method for five sulfonamides in salmon muscle tissue using thin-layer chromatography 555(1991)315

High-performance liquid chromatographic analysis of oxytetracycline in chinook salmon following administration of medicated feed 588(1991)181

Salt adsorption

Dependence of the silanophilic effect on the concentration of preadsorbed salts and on the chemical structure of peptides in reversed-phase thin-layer chromatography 553(1991)467

Sample degradation

Solvent degradation of cloxacillin *in vitro*. Tentative identification of degradation products using thermospray liquid chromatography-mass spectrometry 594(1992)195

Sample handling

see also Special issue index

Sample loading

Displacement effects in preparative gradient high-performance liquid chromatographic separations 590(1992)17

Sample preparation

Reversed-phase liquid chromatography coupled on-line with capillary gas chromatography. I. Introduction of large volumes of aqueous mixtures through an on-column interface 552(1991)489

Clean-up of plasma extracts by gel permeation chromatography during analysis of isosorbide nitrates by capillary gas chromatography 553(1991)21

Interference removal in the organic trace-level analysis of aqueous environmental samples by on-line liquid chromatographic preconcentration techniques with two precolumns 553(1991)329

Reversed-phase and chiral high-performance liquid chromatographic assay of bupivacaine and its enantiomers in clinical samples after continuous extraplural infusion 553(1991)383

Liquid chromatography-thermospray tandem mass spectrometry for identification of a heptabarbital metabolite and sample work-up artefacts 554(1991)205

Characterization of nitrogen-containing aromatic compounds in soil and sediment by capillary gas chromatography-mass spectrometry after fractionation 558(1991)223

Methods for the analysis of thiodiglycol sulphoxide, a metabolite of sulphur mustard, in urine using gas chromatography-mass spectrometry 558(1991)393

Determination of trace amounts of fluoride in raw materials for pharmaceuticals by gas-liquid chromatography 585(1991)289

Membrane-based sample preparation device for the pretreatment of acidic samples prior to cation analysis by ion chromatography 586(1991)283

High-performance liquid chromatographic column-switching technique for the determination of intermediates of anaerobic degradation of toluene in ground water microcosm 587(1991)185

Analyte focusing in capillary electrophoresis using on-line isotachopheresis 591(1992)341

Automated high-performance liquid chromatographic and size-exclusion chromatographic sample preparation by means of a robotic workstation 592(1992)249

Solvent degradation of cloxacillin *in vitro*. Tentative identification of degradation products using thermospray liquid chromatography-mass spectrometry 594(1992)195

Determination of major ions in snow and ice cores by ion chromatography 594(1992)225

Electrodialytic sample treatment coupled on-line with column liquid chromatography for the determination of basic and acidic compounds in environmental samples 600(1992)163

Sample size

Experimental study on the effect of the sample size on the band profile for a binary mixture showing no competitive interaction 556(1991)195

Sample solvent effect

Effect of the sample solvent on band profiles in preparative liquid chromatography using non-aqueous reversed-phase high-performance liquid chromatography 588(1991)1

Sample stacking

Mathematical model describing dispersion in free solution capillary electrophoresis under stacking conditions 559(1991)3

Temperature elevations of the sample zone in free solution capillary electrophoresis under stacking conditions 559(1991)27

Continuous separation and purification of proteins by recycling isotachopheresis 594(1992)361

Sample throughput

Flip-flop elution concept in preparative liquid chromatography 557(1991)137

Utility of small-particle silica in preparative chromatography 590(1992)77

Sample zone temperatures

Temperature elevations of the sample zone in free solution capillary electrophoresis under stacking conditions 559(1991)27

Sampling methods

High-speed gas chromatography. Theoretical and practical aspects 556(1991)331

Determination of hexahydrophthalic anhydride in air using gas chromatography 558(1991)247

Determination of major ions in snow and ice cores by ion chromatography 594(1992)225

Sanger's reagent

High-performance liquid chromatographic separation of DL-amino acids derivatized with chiral variants of Sanger's reagent 555(1991)81

Saturated fractional factorial design

Optimization of the separation of (6*R*)- and (6*S*)-leucovorin and evaluation of the robustness of the optimum 592(1992)37

Scaled relative retention indices

Gas chromatographic elution patterns of chlorinated dioxins versus column polarity 594(1992)269

Scintillator solvent

Prediction of retention indexes. IV. Chain branching in alkylbenzene isomers with C₁₀₋₁₃ alkyl chains identified in a scintillator solvent 589(1992)231

Scylla serrata

Chromatographic analysis of low-molecular-mass copper-binding ligands from the crab species *Scylla serrata* and *Portunus pelagicus* 599(1992)21

Secondary equilibria

Secondary equilibrium size-exclusion chromatography of ions with polymeric mobile phase additives 586(1991)277

Secondary structure

Retention behaviour of a template-assembled synthetic protein and its amphiphilic building blocks on reversed-phase columns 586(1991)43

Sediments

Characterization of nitrogen-containing aromatic compounds in soil and sediment by capillary gas chromatography-mass spectrometry after fractionation 558(1991)223

Selectivity

Method for characterization of selectivity in reversed-phase liquid chromatography. V. Calibration of the retention scale for chromatographic systems with low concentrations of organic solvents in the mobile phase 556(1991)145

Influence of solute size and the non-polar interaction term on the selection of test solutes for the classification of stationary phase selectivity in gas chromatography 556(1991)457

Effect of stationary phase structure on retention and selectivity of restricted-access reversed-phase packing materials 558(1991)19

Chromatographic characterization of a phosphate-modified zirconia support for bio-chromatographic applications 587(1991)149

Reaction kinetics and kinetic processes in modern liquid chromatographic reactors (Review) 589(1992)1

Modelling retention in reversed-phase liquid chromatography as a function of pH and solvent composition 592(1992)157

Separation of plant growth regulators by capillary electrophoresis 594(1992)335

Reversed- and normal-phase separations by high-temperature open-tubular column liquid chromatography 598(1992)153

Selectivity mapping

Characterization of hydrophobic interaction and hydrophobic interaction chromatography media by multivariate analysis 599(1992)131

Selectivity points

Retention behaviour of closely related coumarins in thin-layer chromatographic preassays for high-performance liquid chromatography according to the "PRISMA" model 593(1992)329

Selectivity tuning

Pseudo-electrochromatography-mass spectrometry: a new alternative 554(1991)339

Semi-permeable surface packings

Evaluation of restricted access media for high-performance liquid chromatographic analysis of sulfonamide antibiotic residues in bovine serum 598(1992)23

Sendai virus

Comparison of detergents for extraction and ion-exchange high-performance liquid chromatography of Sendai virus membrane proteins 599(1992)157

Sensor chips

Biospecific interaction analysis using surface plasmon resonance detection applied to kinetic, binding site and concentration analysis 597(1992)397

Separation efficiency

Analysis of separation efficiency in capillary electrophoresis with direct control of electroosmosis by using an external electric field 559(1991)133

Isotachophoretic separation of minor components from a matrix component in the case of strong electrolytes 585(1991)297

Separation factor

Theoretical approach to the gas chromatographic separation of enantiomers on dissolved cyclodextrin derivatives 552(1991)43

Separation number

Critical capillary column examination of the relationship between separation number and height equivalent to a theoretical plate and their dependence on temperature 595(1992)209

Van Deemter-type relationship for determining the optimum initial flow-rate and optimum pressure programming rate in temperature/pressure-programmed capillary column gas chromatography utilizing separation numbers 595(1992)221

Sepharose, 1,1'-carbonyldiimidazole substituted

Efficient substitution of 1,1'-carbonyldiimidazole activated cellulose and Sepharose matrices with amino acyl spacer arms 587(1991)271

Sepharose-based immunoadsorbents

Compared stability of Sepharose-based immunoadsorbents prepared by various activation methods 584(1992)17

Sepharose Cl-B

Diffusion of proteins in Sepharose Cl-B gels 591(1992)115

Sep-Pak cartridges

Use of a Sep-Pak cartridge for preparative collection in gas chromatography 555(1991)302

Sequencing

Low-cost, high-sensitivity laser-induced fluorescence detection for DNA sequencing by capillary gel electrophoresis 559(1991)237

Automated two-dimensional liquid chromatographic system for mapping proteins in highly complex mixtures 588(1991)115

Sequential competitive binding assay

Process monitoring by flow-injection immunoassay. Evaluation of a sequential competitive binding assay 597(1992)383

Shielded hydrophobic phase packings

Evaluation of restricted access media for high-performance liquid chromatographic analysis of sulfonamide antibiotic residues in bovine serum 598(1992)23

Shielded hydrophobic phases

High-performance liquid chromatographic determination of rifampentine and its metabolite in human plasma by direct injection into a shielded hydrophobic phase column 553(1991)35

Signal processing

Data compression in computerized signal processing for isocratic chromatography 585(1991)160

Silanol groups

Evaluation of the application of liquid-phase titration to the examination of the adsorption activity of silica-based high-performance liquid chromatographic packings 552(1991)463

Effect of silanol groups on heat-treated silicas by calcination and retention behaviour in high-performance liquid chromatography 556(1991)407

Use of the reordering/resolution of alkyl-modified silica to characterize the microscopic heterogeneity of silica via liquid chromatography 556(1991)415

Investigations of the influence of silanol groups on the separation of enantiomers by liquid and supercritical fluid chromatography 592(1992)309

Cation-exchange liquid chromatography of choline and acetylcholine on free shielded silanols of silica-based reversed-phase stationary phases 596(1992)43

Silanols

Retention behavior of salicylidene-glycinatoaluminium(III) in reversed-phase high-performance liquid chromatography 586(1991)356

Silanophilic effects

Dependence of the silanophilic effect on the concentration of preadsorbed salts and on the chemical structure of peptides in reversed-phase thin-layer chromatography 553(1991)467

Silica

Retention reproducibility of basic drugs in high-performance liquid chromatography on a silica column with a methanol-high-pH buffer eluent. Changes in selectivity with the age of the stationary phase 592(1992)85

Polymer-coated reversed-phase packings with controlled hydrophobic properties. I. Effect on the selectivity of protein separations 599(1992)65

Comparison of adsorption properties of Florisil and silica in high-performance liquid chromatography. I. Retention behaviour of monofunctional model solutes 600(1992)51

Silica gel

Use of the reordering/resolution of alkyl-modified silica to characterize the microscopic heterogeneity of silica via liquid chromatography 556(1991)415

Beaded polymer supports and gels (Review). II. Physico-chemical criteria and functionalization 586(1991)199

Silica gel, dynamically modified

High-performance liquid chromatography on dynamically modified silica. IX. Modification of silica with 3-(N,N-dimethylpalmitylammonium) propanesulphonate for reversed-phase chromatography 556(1991)353

Silica gel, heat-treated

Effect of silanol groups on heat-treated silicas by calcination and retention behaviour in high-performance liquid chromatography 556(1991)407

Silica gel, nitrophenylethylsilylated

Preparation of nitrophenylethylsilylated silica gel and its chromatographic properties in the separation of polychlorinated dibenzo-*p*-dioxins 595(1992)77

Silica gel, plasma modified

Gas chromatographic investigation of silica supports modified in a high-frequency low-temperature plasma 552(1991)179

Silica gel, polymethacrylate-coated

Polymer-coated reversed-phase packings with controlled hydrophobic properties. II. Effect on the selectivity of peptide separations 599(1992)77

Silica gel, porous

Impact of acidic/hydrothermal treatment on pore structural and chromatographic properties of porous silicas. I. The conventional approach 556(1991)395

Silver/silver chloride electrode

Mixed potential response at metallic electrodes in potentiometric detection for ion chromatography 589(1992)201

Silver staining

Improved detection of human pancreatic proteins separated by two-dimensional isoelectric focusing/sodium dodecyl sulphate gel electrophoresis using a combined staining procedure 563(1991)63

Silver staining of collagen type I after sodium dodecylsulphate polyacrylamide gel electrophoresis: effect of Maillard reaction 578(1992)53

Silylating agents

Preparation and characterization of mixed functional phase silica materials using phenyl-, butyl- or octylchlorosilane as a silylating agent 596(1992)151

Simplex method

Simplex optimization of liquid chromatography with the function of mutual information as a criterion 585(1991)187

Optimization of an ion-exchange high-performance liquid chromatographic method for the determination of carboxylic acids, sugars, glycerol and ethanol in wines 589(1992)151

Optimization of the separation of (6*R*)- and (6*S*)-leucovorin and evaluation of the robustness of the optimum 592(1992)37

Simulated moving bed process

Glucose-fructose equilibria on Dowex Monosphere 99 CA resin under overloaded conditions 590(1992)147

Size-exclusion mechanism

Biosep-SEC-S high-performance size-exclusion chromatographic columns for proteins and peptides 599(1992)25

Skin effect

Evaluation of morphological structure of packings by gel permeation chromatography 596(1992)157

Skin-toning cream

Micellar electrokinetic chromatographic study of hydroquinone and some of its ethers. Determination of hydroquinone in skin-toning cream 596(1992)95

Soil

Modern approach to the quantitative determination of volatiles in solid samples. Multiple headspace extraction gas chromatography for the determination of cyclohexanone residues in soil 552(1991)205

Characterization of nitrogen-containing aromatic compounds in soil and sediment by capillary gas chromatography-mass spectrometry after fractionation 558(1991)223

Soil extract

Determination of acephate by liquid chromatography in the presence of aqueous soil extracts 585(1991)324

Solid-phase dispersion

High-performance liquid chromatography of sulphonamides extracted from bovine and porcine muscle by solid-phase dispersion 595(1992)179

Solid-phase extraction

Solid-matrix partition for separation of organochlorine pesticide residues from fatty materials 552(1991)241

Removal of organic pollutants from aqueous solution. V. Comparative study of the extraction, recovery and chromatographic separation of some organic insecticides using unloaded polyurethane foam columns 552(1991)371

Improvement of extraction and concentration of milk peptides with solid-phase cartridges for analysis by high-performance liquid chromatography 553(1991)155

Screening for coumatetralyl in soft drinks by solid-matrix extraction and high-performance liquid chromatography with diode-array detection 553(1991)305

Electrochromatographic solid-phase extraction for determination of cimetidine in serum by micellar electrokinetic capillary chromatography 559(1991)547

On-line solid-phase extraction and high-performance liquid chromatographic determination of nortriptyline and amitriptyline in serum 579(1992)291

General and selective isolation procedure for high-performance liquid chromatographic determination of anabolic steroids in tissues 588(1991)89

Resolution of lipolysis mixtures from soybean oil by a solid-phase extraction procedure 589(1992)353

Determination of diacetoneketogulonic acid in water samples by high-performance thin-layer chromatography 589(1992)390

Chemically modified resins for solid-phase extraction 590(1992)197

Determination of carboxylic acids, sugars, glycerol and ethanol in wine and grape must by ion-exchange high-performance liquid chromatography with refractive index detection 590(1992)215

Quantitation of mutagenic/carcinogenic heterocyclic aromatic amines in food products 592(1992)271

Use of direct injection precolumn techniques for the high-performance liquid chromatographic determination of the retinoids acitretin and 13-*cis*-acitretin in plasma 593(1992)55

Determination of Zy 17617B in plasma by solid-phase extraction and liquid chromatography with automated precolumn exchange 593(1992)73

Determination of tetracyclines in bovine and porcine muscle by high-performance liquid chromatography using solid-phase extraction 596(1992)211

Solid-phase extraction and high-performance liquid chromatographic analysis of a toxic compound from γ -irradiated polyurethane 600(1992)93

Solid sorbents

Determination of hexahydrophthalic anhydride in air using gas chromatography 558(1991)247

Solid surfaces

Experimental determination of adsorption isotherm data for the study of the surface energy distribution of various solid surfaces by inverse gas-solid chromatography 591(1992)233

Validity of the model used to relate the energy distribution and the adsorption isotherm 591(1992)267

Investigation of solid surfaces by high-performance liquid chromatography 592(1992)121

Solubility factors

Attempts to use Laffort's solubility factors as polarity parameters for organic compounds in inverse gas chromatography 555(1991)205

Solubility parameters

Activity coefficients at infinite dilution determined by gas-liquid chromatography. Organic solvents in Apiezon L 586(1991)303

Solute affinity

High-performance liquid chromatography of amino acids, peptides and proteins. CXV. Thermodynamic behaviour of peptides in reversed-phase chromatography 593(1992)103

Solute-ligand complexes

Study of high-affinity interactions by quantitative affinity chromatography. Analytical expressions in terms of total ligand concentration 597(1992)57

Solute-ligand interactions

Recent developments in quantitative affinity chromatography (Review) 597(1992)67

Solute partition

Recent developments in quantitative affinity chromatography (Review) 597(1992)67

Solute-polymer interaction constants

Determination of solute-polymer interaction constants via chromatography with polymer capillaries 594(1992)29

Solute-stationary phase interactions

Calculation of retention and selectivity in reversed-phase liquid chromatography 552(1991)91

Reaction kinetics and kinetic processes in modern liquid chromatographic reactors (Review) 589(1992)1

Solvatochromic hydrogen bond donor acidity of cyclodextrins and reversed-phase liquid chromatographic retention of small molecules on a β -cyclodextrin-bonded silica stationary phase 595(1992)45

Biosep-SEC-S high-performance size-exclusion chromatographic columns for proteins and peptides 599(1992)25

Solute size

Influence of solute size and the non-polar interaction term on the selection of test solutes for the classification of stationary phase selectivity in gas chromatography 556(1991)457

Solvation

Hydrogen bonding. XVI. A new solute solvation parameter, π^H_2 , from gas chromatographic data 587(1991)213

Hydrogen bonding. XVII. The characterisation of 24 gas-liquid chromatographic stationary phases studied by Poole and co-workers, including molten salts, and evaluation of solute-stationary phase interactions 587(1991)229

Solvation equation

Hydrogen bonding. XIX. The characterisation of two poly(methylphenylsiloxane)s 588(1991)361

Solvation parameters

Hydrogen bonding. XXI. Solvation parameters for alkylaromatic hydrocarbons from gas-liquid chromatographic data 594(1992)229

Solvatochromic solvent polarity

Influence of mobile phase composition on evaluation of lipophilicity by partition chromatography 557(1991)399

Solvent diagrams

Retention behaviour of closely related coumarins in thin-layer chromatographic preassays for high-performance liquid chromatography according to the "PRISMA" model 593(1992)329

Solvent mixtures

Analysis of industrial solvent mixtures in water using a miniature purge-and-trap device with thermal desorption and capillary gas chromatography-mass spectrometry 557(1991)429

Solvent selectivity

Use of solvent selectivity optimization procedures for high-performance liquid chromatographic method development 592(1992)43

Solvent strength

Application of solute and mobile phase partition coefficient to describe solute retention in reversed-phase high-performance liquid chromatography 555(1991)21

Computer simulation for the prediction of separation as a function of pH for reversed-phase high-performance liquid chromatography. II. Resolution as a function of simultaneous change in pH and solvent strength 592(1992)197

Retention behaviour of closely related coumarins in thin-layer chromatographic preassays for high-performance liquid chromatography according to the "PRISMA" model 593(1992)329

Solvent systems

Quantitative two-dimensional thin-layer chromatography 553(1991)477

Effect of the substituents of β -diketonato ligands on R_f values of tris(chelate) transition metal complexes obtained by normal- and reversed-phase thin-layer chromatography on unmodified silica gel 585(1991)359

Purification of erucic acid by preparative high-performance liquid chromatography and crystallization 590(1992)139

Separation of pristinamycins by high-speed counter-current chromatography. I. Selection of solvent system and preliminary preparative studies 593(1992)363

Thin-layer chromatography on polyacrylonitrile. IV. Investigation of the separation mechanisms for tris-(alkylxanthato)cobalt(III) complexes 596(1992)91

Solvophobic interaction

Reversed-phase liquid chromatographic behaviour of alkylamines with amperometric detection and its application to trace analysis 594(1992)159

Solvophobic theory

Calculation of retention and selectivity in reversed-phase liquid chromatography 552(1991)91

Description of retention mechanism by solvophobic theory. Influence of organic modifiers on the retention behaviour of homologous series in reversed-phase liquid chromatography 556(1991)219

Sorbent bed homogeneity

Experimental evaluation of sorbent bed homogeneity in high-performance liquid chromatographic columns 592(1992)109

Sorbent distribution

Experimental evaluation of sorbent bed homogeneity in high-performance liquid chromatographic columns 592(1992)109

Sorbents

see also Adsorbents

Calculation of retention and selectivity in reversed-phase liquid chromatography 552(1991)91

Normal- and reversed-phase high-performance liquid chromatography of some phosphonodipeptides 553(1991)143

Separon HEMA modified for immobilized metal ion affinity chromatographic separation of proteins 598(1992)15

Sorption effect

Prediction of the elution profiles of high-concentration bands in gas chromatography 589(1992)223

Sorption energy

Retentions of alkyloxazoles and alkylthiazoles and their prediction based on non-linear additivity concepts in gas chromatography 552(1991)1

Sorption layers

Theory of linear non-equilibrium chromatography with beds of a finite length 591(1992)273

Sorption mechanism

Adsorption and partition mode in high-performance liquid chromatography of highly polar solutes on silica 552(1991)23

Southern blotting

Identification of DNA molecules by pre-column hybridization using capillary electrophoresis 559(1991)295

Soxhlet extraction

Supercritical carbon dioxide extraction of resin and fatty acids from sediments at pulp mill sites 594(1992)309

Spacer arm coupling

Efficient substitution of 1,1'-carbonyldiimidazole activated cellulose and Sepharose matrices with amino acyl spacer arms 587(1991)271

Spacer arms

Beaded polymer supports and gels (Review). II. Physico-chemical criteria and functionalization 586(1991)199

Separon HEMA modified for immobilized metal ion affinity chromatographic separation of proteins 598(1992)15

Specific retention volumes

Determination of specific retention volumes at 20°C for hydrocarbons on microporous carbons 586(1991)161

Split injection

Split injection into a capillary column at very low split ratios 557(1991)1

Split injector

Split injector for capillary zone electrophoresis 559(1991)103

SPLITT

Transverse particle redistribution in a flat channel for SPLITT or integral Doppler anemometry 600(1992)133

Spot counting procedures

Comparison of statistical theories of spot overlap in two-dimensional separations and verification of means for estimating the number of zones 591(1992)1

Spray jet assembly

Spray jet assembly interface for the coupling of reversed-phase narrow-bore liquid chromatography and Fourier transform infrared spectrometry 552(1991)635

Spray reagents

Application of the iodine-azide reagent for selective detection of thiophosphoryl compounds in thin-layer chromatography 588(1991)307

Stability constants

Measurement of stability constants for complexes of divalent cations with tartaric acid by ion chromatography 589(1992)185

Stability indicating assay

Development of stability indicating assay methods for the determination of hydroxamic acids in topical formulations 594(1992)209

Staining methods

Fast and sensitive staining technique for catalase in polyacrylamide gel 594(1992)400

Staphylococcus carnosus

Adsorption behaviour of lipase from *Staphylococcus carnosus* on a hydrophobic adsorbent 584(1992)109

Stationary phase additives

Gas-liquid chromatography in qualitative analysis. XIX. The use of antioxidants to delay the oxidation of polyoxyethylene glycol stationary phases 552(1991)161

Stationary phases, GC

Determination of sorbent polarity and selectivity by linear regression of retention indices 552(1991)31

Gas-liquid chromatography in qualitative analysis. XIX. The use of antioxidants to delay the oxidation of polyoxyethylene glycol stationary phases 552(1991)161

Gas chromatography in qualitative analysis. XX. The deactivation of diatomaceous supports by batyl alcohol 552(1991)169

Gas chromatographic separation of unsaturated polar compounds on inorganic stationary phases containing silver nitrate 552(1991)313

Gas chromatographic analysis of some glycol ether analytes on a specially packed column 552(1991)353

Theoretical and experimental comparison of serially linked and mixed-packing gas-liquid chromatography columns 555(1991)173

Gas chromatographic retention behaviour of some solutes on structurally similar polar and non-polar stationary phases 555(1991)191

Thermodynamics of solution of non-mesomorphic solutes at infinite dilution in the smectic-A, nematic and isotropic phases of *p*-*n*-octyl-*p*'-cyanobiphenyl. A gas-liquid chromatographic study 556(1991)317

New solid adsorbents for the separation of lower hydrocarbons and permanent gases. II. Ammonium molybdophosphate 556(1991)425

Dynamic gas-solid chromatographic techniques for characterizing carbon molecular sieves 556(1991)433

Influence of solute size and the non-polar interaction term on the selection of test solutes for the classification of stationary phase selectivity in gas chromatography 556(1991)457

Gas chromatographic comparative study of Superox 20M immobilized in different ways 556(1991)485

Capillary gas chromatography of C₅-C₁₃ branched alkynes on squalane and liquid crystal stationary phases 557(1991)241

Immobilization method for polyethylene glycol using a cross-linking co-agent 558(1991)323

Prediction of retention in gas-liquid chromatography using the UNIFAC group contribution method. II. Polymer stationary phases 585(1991)83

Evaluation of gas chromatographic columns for the determination of methylmercury in aqueous head space extracts from biological samples 586(1991)329

Hydrogen bonding. XVII. The characterisation of 24 gas-liquid chromatographic stationary phases studied by Poole and co-workers, including molten salts, and evaluation of solute-stationary phase interactions 587(1991)229

Study of the response of three liquid crystals as stationary phases for the gas chromatography of some cyclic monoterpene volatile oil constituents with short retention times 587(1991)343

Gas chromatographic properties of immobilized poly(ethylene glycol) stationary phases 588(1991)225

- Study of dihydroxy-substituted saturated urushiol crown ether as a stationary phase in capillary gas chromatography 589(1992)368
- Gas chromatographic retention behaviour of dibenzothiophene derivatives on a smectic liquid crystalline polysiloxane stationary phase 590(1992)297
- Preparation and capillary column gas chromatographic characterization of aza-crown ether polysiloxane stationary phases 594(1992)243
- Direct aqueous injection gas chromatography on a potassium fluoride crystal hydrate-containing sorbent. Determination of volatile organic solvents in the fermentation broth of *Clostridium* strains 596(1992)138
- Chromatographic properties of graphitized thermal carbon black modified with a monolayer of liquid crystal 600(1992)59
- Gas chromatographic separation of *cis-trans* isomers of alkylcyclohexylbenzenes on a capillary column with a liquid crystalline stationary phase 600(1992)123
- Cholesteryl acetate as a stationary phase for the gas chromatography of some volatile oil constituents 600(1992)337
- Stationary phases, LC**
- Estimation of cellulose polarity by gas chromatography 552(1991)365
- Chemically bonded phases for the reversed-phase high-performance liquid chromatographic separation of basic substances 552(1991)415
- Possible role for stationary phase metal interactions in the chromatography of hydroxyamines on silica 552(1991)429
- High-performance liquid chromatography of amino acids and dipeptides on new ion exchangers of the HEMA series 552(1991)439
- Evaluation of the application of liquid-phase titration to the examination of the adsorption activity of silica-based high-performance liquid chromatographic packings 552(1991)463
- High-performance liquid chromatographic determination of rifampine and its metabolite in human plasma by direct injection into a shielded hydrophobic phase column 553(1991)35
- Anion-exchange chromatographic properties of α -lactalbumin eluted from quaternized polyvinylimidazole. Study of the role of the polymer coating 553(1991)165
- Evaluation of different packings for high-performance liquid chromatographic analysis of alkyl lysophospholipids 553(1991)285
- Anion-exchange chromatography of DNA restriction fragments 555(1991)109
- N-(Hydroxymethyl)thioamide resin as stationary phase in ion-exchange chromatography for metal ion separation 555(1991)291
- High-performance liquid chromatography on dynamically modified silica. IX. Modification of silica with 3-(N,N-dimethylpalmitylammonium) propanesulphonate for reversed-phase chromatography 556(1991)353
- Synthesis of chemically bonded liquid crystals for high-performance liquid chromatography. New phases via the organochlorosilane pathway 556(1991)373
- Impact of acidic/hydrothermal treatment on pore structural and chromatographic properties of porous silicas. I. The conventional approach 556(1991)395
- Effect of silanol groups on heat-treated silicas by calcination and retention behaviour in high-performance liquid chromatography 556(1991)407
- Use of the reordering/resolution of alkyl-modified silica to characterize the microscopic heterogeneity of silica via liquid chromatography 556(1991)415
- High-performance liquid chromatography of amino acids, peptides and proteins. CXIV. Protein interactions with porous coulombic sorbents: comparison of experimental findings with predictions of several adsorption models 557(1991)335
- Retention behaviour of polycyclic aromatic hydrocarbons on a liquid-crystal bonded phase in reversed-phase liquid chromatography 557(1991)459
- Structural investigation of oligomeric *n*-octylsilyl reversed phases 557(1991)515
- Effect of stationary phase structure on retention and selectivity of restricted-access reversed-phase packing materials 558(1991)19
- Diffusion of sorbed pyrene in the bonded layer of reversed-phase silicas. Effect of alkyl chain length and pore diameter 558(1991)31
- Modification of reversed-phase columns with dyed surfactants. Preparation of mechanically resistant efficient immobilized dyes for protein purification 558(1991)43
- Chromatographic fractionation of nucleic acids using microcapsules made from plant cells 585(1991)57
- Fast protein separation by reversed-phase high-performance liquid chromatography on octadecylsilyl-bonded non-porous silica gel. Effect of particle size of column packing on column efficiency 585(1991)207

- Ligand-exchange chromatography of alkenes on stationary phases containing palladium(II) complexes. Enantiomeric separation of *trans*-1,2-divinylcyclohexane 585(1991)309
- Continuous beds for standard and micro high-performance liquid chromatography 586(1991)21
- High-performance affinity chromatography of NADP⁺ dehydrogenases from cell-free extracts using a nucleotide analogue as general ligand 586(1991)51
- Physical and chemical characterization of a porous phosphate-modified zirconia substrate 587(1991)137
- Chromatographic characterization of a phosphate-modified zirconia support for bio-chromatographic applications 587(1991)149
- Optimization of oxidation of glycoproteins: an assay for predicting coupling to hydrazide chromatographic supports 587(1991)171
- Dehydrogenase-silica as a stationary phase for the separation of alcohols and ketones 587(1991)268
- High-performance liquid chromatography on polypyrrole-modified silica 588(1991)25
- Comparison of four liquid crystal stationary phases used above and below their melting point temperatures for the gas chromatography of some volatile oil constituents 588(1991)231
- Synthesis and characterization of stationary phases on the basis of silicas modified with epoxidized polybutadienes. IV. Chromatographic experiments on a new amino-functionalized stationary phase for high-performance liquid chromatography 588(1991)340
- Predictability of chromatographic protein separations. Study of size-exclusion media with narrow particle size distributions 589(1992)71
- Phosphorylated polystyrene resins in high-performance ion-exchange chromatography 589(1992)87
- Cellulose beads: a weak leaking affinity support 589(1992)93
- Activation of matrices by 4,6-diphenylthieno[3,4-*d*]-1,3-dioxol-2-one-5,5-dioxide. High-performance liquid affinity chromatographic separations 589(1992)101
- Utility of small-particle silica in preparative chromatography 590(1992)77
- Importance of intraparticle convection in the performance of chromatographic processes 590(1992)93
- Chromatographic effects of residual amino groups on liquid chromatographic stationary phases derived from aminopropyl-silica gel 590(1992)189
- Retention processes on α_1 -acid glycoprotein-bonded stationary phase 591(1992)55
- Physico-chemical properties of electron-acceptor stationary phases in liquid chromatography 591(1992)75
- Chromatography on Poly-RP and its cyano and diol derivatives using both polar and non-polar solvent systems 591(1992)89
- Oriented immobilization of peptide ligands on solid supports 591(1992)99
- Retention behaviour of some ring-substituted phenol derivatives on a porous graphitized carbon column 592(1992)75
- New, stable polyamine-bonded polymer gel column 592(1992)93
- Application of a pellicular anion-exchange resin to the separation of inorganic and organic anions by single-column anion exchange 592(1992)101
- High-performance liquid chromatography of amino acids, peptides and proteins. CXV. Thermodynamic behaviour of peptides in reversed-phase chromatography 593(1992)103
- Mixed-mode hydrophilic and ionic interaction chromatography rivals reversed-phase liquid chromatography for the separation of peptides 594(1992)75
- Suppression of the effect of metal impurities in alkylsilylated silica packing materials 594(1992)87
- Liquid chromatographic separation of all-carbon molecules C₆₀ and C₇₀ with multi-legged phenyl group bonded silica phases 594(1992)105
- Fast protein monitoring in fermentation broth using non-porous micropellicular reversed-phase columns 594(1992)392
- Preparation of nitrophenylethylsilylated silica gel and its chromatographic properties in the separation of polychlorinated dibenzo-*p*-dioxins 595(1992)77
- Evaluation of reversed-phase liquid chromatographic columns for recovery and selectivity of selected carotenoids 595(1992)89
- Physical and biochemical characterization of five commercial resins for immunoaffinity purification of factor IX 595(1992)103
- Stationary phase effect on enantioselectivity of dansyl phenylalanine in microcolumn liquid chromatography with γ -cyclodextrin as mobile phase additive 595(1992)121

- Ion- and ligand-exchange chromatography of proteins using porous zirconium oxide supports in organic and inorganic Lewis base eluents 596(1992)27
- Cation-exchange liquid chromatography of choline and acetylcholine on free shielded silanols of silica-based reversed-phase stationary phases 596(1992)43
- Preparation and characterization of mixed functional phase silica materials using phenyl-, butyl- or octylchlorosilane as a silylating agent 596(1992)151
- Evaluation of morphological structure of packings by gel permeation chromatography 596(1992)157
- Immobilized lectin columns: useful tools for the fractionation and structural analysis of oligosaccharides (Review) 597(1992)111
- Effects of stationary phase ligand density on high-performance ion-exchange chromatography of proteins 598(1992)7
- Separon HEMA modified for immobilized metal ion affinity chromatographic separation of proteins 598(1992)15
- Evaluation of restricted access media for high-performance liquid chromatographic analysis of sulfonamide antibiotic residues in bovine serum 598(1992)23
- Protein chromatography using a continuous stationary phase 598(1992)169
- High-performance hydrophobic interaction chromatography of proteins on reversed-phase supports coated with non-ionic surfactants of polyoxyethylene type. Purification of a fungal aspartic proteinase 598(1992)181
- Biosep-SEC-S high-performance size-exclusion chromatographic columns for proteins and peptides 599(1992)25
- Polymer-coated reversed-phase packings with controlled hydrophobic properties. I. Effect on the selectivity of protein separations 599(1992)65
- Polymer-coated reversed-phase packings with controlled hydrophobic properties. II. Effect on the selectivity of peptide separations 599(1992)77
- High-performance liquid chromatographic separation of detritylated oligonucleotides on highly cross-linked poly(styrene-divinylbenzene) particles 599(1992)113
- High-performance liquid chromatography of amino acids, peptides and proteins. CXX. Evaluation of bandwidth behaviour of proteins chromatographed on tentacle-type anion exchangers 599(1992)163
- High-performance liquid chromatography of amino acids, peptides and proteins. CXXI. 8-Hydroxyquinoline-metal chelate chromatographic support: an additional mode of selectivity in immobilized-metal affinity chromatography 599(1992)171
- Schemes for efficient protein purification on a family of polymeric ion exchangers in glass columns 599(1992)239
- High-performance liquid chromatography of benzodiazepines using sorbents with thermally immobilized Carbowax 20M 600(1992)109
- Application of an internal surface reversed-phase column for the automated determination of flucycloxuron residues 600(1992)211
- High-performance affinity chromatography of DNA. II. Porosity effects 600(1992)221
- Stationary phases, SFC**
Separation of *trans/cis* α - and β -carotenes by supercritical fluid chromatography. II. Effect of the type of octadecyl-bonded* stationary phase on retention and selectivity of carotenes 557(1991)59
- Correlation between column surface area and retention of polar solutes in packed-column supercritical fluid chromatography 594(1992)291
- Stationary phases, TLC**
Comparison of commercially available thin-layer chromatography plates with mixtures of dyes, analgesics and phenols 595(1992)368
- Stationary phase stability**
Chromatographic characterization of a phosphate-modified zirconia support for bio-chromatographic applications 587(1991)149
- Statistical moments**
Theory of linear non-equilibrium chromatography with beds of a finite length 591(1992)273
- Statistical theories**
Comparison of statistical theories of spot overlap in two-dimensional separations and verification of means for estimating the number of zones 591(1992)1
- Step gradients**
Development of a rational optimisation procedure for the automated sample clean-up with column switching in pesticide residue analysis 552(1991)113
- Stereoselective effects**
Reversed retention order and other stereoselective effects in the separation of amino alcohols on Chiralcel OD 592(1992)331

Stereoselectivity

Enantiomeric separation of racemic hydroperoxides and related alcohols 588(1991)352

Steric disposition

Application and chiral recognition of heptakis (2,6-di-O-methyl-3-O-trifluoroacetyl)- β -cyclodextrin as a stationary phase for the gas chromatographic separation of enantiomers 589(1992)257

Stokes radius

Diffusion of proteins in the chromatographic gel AcA-34 558(1991)71

Streaming potential

Capillary modification and evaluation using streaming potential and frontal chromatography for protein analysis in capillary electrophoresis 594(1992)325

Structure-activity relationships

Hydrophobicity of β -lactam antibiotics. Explanation and prediction of their behaviour in various partitioning solvent systems and reversed-phase chromatography 585(1991)3

High-performance liquid chromatographic study of some biologically active analogues of arabinosylcytosine 588(1991)147

Chromatographic separation and molecular modelling of triazines with respect to their inhibition of the growth of L1210/R71 cells 592(1992)59

Direct high-performance liquid chromatographic resolution of a novel benzothiazine Ca^{2+} antagonist and related compounds 593(1992)37

Structure-retention relationships

Retentions of alkyloxazoles and alkylthiazoles and their prediction based on non-linear additivity concepts in gas chromatography 552(1991)1

Determination of sorbent polarity and selectivity by linear regression of retention indices 552(1991)31

New aspects of quantitative structure-retention relationships in chromatography 552(1991)59

Calculation of retention and selectivity in reversed-phase liquid chromatography 552(1991)91

Influence of mobile phase composition on evaluation of lipophilicity by partition chromatography 557(1991)399

Prediction of retention indexes. II. Structure-retention index relationship on polar columns 586(1991)85

Prediction of retention indexes. IV. Chain branching in alkylbenzene isomers with C_{10-13} alkyl chains identified in a scintillator solvent 589(1992)231

Gas chromatographic retention behaviour of dibenzothiophene derivatives on a smectic liquid crystalline polysiloxane stationary phase 590(1992)297

Correlations between high-performance liquid chromatographic retention, X-ray structural and ^{13}C NMR spectroscopic data of flavonoid compounds 592(1992)65

Effect of temperature and mobile phase on the retention of retinoates in reversed-phase liquid chromatography 592(1992)127

Gas chromatographic elution patterns of chlorinated dioxins versus column polarity 594(1992)269

Chromatographic behaviour of dipyritylsulphides. Relationship between $\log k'$ values and structure by reversed-phase high-performance liquid chromatography 595(1992)334

Structure-retention index relationships for derivatized monosaccharides on non-polar gas chromatography columns 596(1992)79

Strychnos nux-vomica

Separation of *Strychnos nux-vomica* alkaloids by high-performance liquid chromatography 587(1991)318

Sugar process juices

Determination of volatile and non-volatile organic acids in technical sugar solutions by ion-exclusion chromatography 555(1991)65

Sulphur chemiluminescence detector

Comparison of commercially available atomic emission and chemiluminescence detectors for sulfur-selective gas chromatographic detection 591(1992)313

Supercritical fluid chromatography

see also Special issue index

Supercritical fluid chromatography

Effects of modifiers in packed and open-tubular supercritical fluid chromatography 552(1991)527

Ion-mobility spectrometry as a detection method for packed-column supercritical fluid chromatography 552(1991)551

Supercritical fluid chromatography in the routine stability control of antipruritic preparations 553(1991)71

Analysis of polyethylene glycol and derivatives by high-performance liquid chromatography using elevated temperatures and low-wavelength ultraviolet detection, and supercritical fluid chromatography 553(1991)423

Ionization mechanisms in capillary supercritical fluid chromatography-chemical ionization mass spectrometry 554(1991)351

- Separation of *trans/cis* α - and β -carotenes by supercritical fluid chromatography. I. Effects of temperature, pressure and organic modifiers on the retention of carotenes 557(1991)47
- Separation of *trans/cis* α - and β -carotenes by supercritical fluid chromatography. II. Effect of the type of octadecyl-bonded stationary phase on retention and selectivity of carotenes 557(1991)59
- Magnitude of the diffusion coefficient anomaly in the critical region and its effect on supercritical fluid chromatography 557(1991)69
- Packed column supercritical fluid chromatography with carbon dioxide-polar modifiers. Influence of carbon dioxide density on retention 557(1991)85
- Quantitative aspects of the determination of compounds with widely varying polarity using capillary supercritical fluid chromatography 557(1991)99
- Chiral π -donor stationary phases with (*R*)-*N*-pivaloylnaphthylethylamide groups for direct enantiomer separation by gas, liquid and supercritical fluid chromatography 557(1991)163
- Capillary supercritical fluid chromatography of primary aliphatic amines using carbon dioxide and nitrous oxide as mobile phases 558(1991)458
- Separation of secondary alcohol enantiomers using supercritical fluid chromatography 585(1991)117
- Fast separation of polymethoxylated flavones by carbon dioxide supercritical fluid chromatography 586(1991)171
- Determination of the hold-up time and column outlet density for capillary supercritical fluid chromatography 588(1991)289
- Analysis of benzyloxy-terminated poly(1,3,6-trioxocane)s by supercritical fluid chromatography 589(1992)295
- High-performance liquid chromatography and supercritical fluid chromatography of monosaccharides and polyols using light-scattering detection. Chemometric studies of the retentions 590(1992)203
- Comparison of ethoxylated alcohols and polyethylene glycols by high-performance liquid chromatography and supercritical fluid chromatography using evaporative light-scattering detection 591(1992)149
- Investigations of the influence of silanol groups on the separation of enantiomers by liquid and supercritical fluid chromatography 592(1992)309
- Analytical study of heavy crude oil fractions by coupling of the transalkylation reaction with supercritical fluid chromatography 593(1992)321
- Synthesis of (1*R-trans*)-*N,N'*-1,2-cyclohexylenebisbenzamide-oligodimethylsiloxane copolymers for use as chiral stationary phases for capillary supercritical fluid chromatography 594(1992)283
- Correlation between column surface area and retention of polar solutes in packed-column supercritical fluid chromatography 594(1992)291
- Quantitative hydrocarbon group analysis of gasoline and diesel fuel by supercritical fluid chromatography 595(1992)301
- Supercritical fluid chromatographic analysis of polyprenols in *Ginkgo biloba* L 600(1992)364
- Supercritical fluid chromatography, semi preparative**
Combination of supercritical fluid chromatography with thin-layer chromatography on a semi-preparative scale 552(1991)539
- Supercritical fluid extraction**
Effects of collection solvent parameters and extraction cell geometry on supercritical fluid extraction efficiencies 594(1992)297
- Supercritical carbon dioxide extraction of resin and fatty acids from sediments at pulp mill sites 594(1992)309
- Optimization of supercritical fluid extraction of volatile constituents from a model plant matrix 600(1992)175
- Supercritical fluid extraction of chemical warfare agent simulants from soil 600(1992)273
- Superox 20M**
Gas chromatographic comparative study of Superox 20M immobilized in different ways 556(1991)485
- Superoxide chemical transformation**
Superoxide chemical transformation of diepoxide polyaromatic hydrocarbon DNA adducts. Determination of benzo[*a*]pyrene-*r*-7,*t*-8,9,*c*-10-tetrahydrotetrol by gas chromatography 588(1991)273
- Surface analysis**
Investigation of solid surfaces by high-performance liquid chromatography 592(1992)121
- Surface area**
Beaded polymer supports and gels (Review). II. Physico-chemical criteria and functionalization 586(1991)199
- Correlation between column surface area and retention of polar solutes in packed-column supercritical fluid chromatography 594(1992)291
- Surface characterization**
Activity testing and surface characterization of pretreated fused-silica capillaries for gas chromatography. A new modification of existing intermediate column tests 590(1992)271

Surface energies

Chromatographic separation and molecular modelling of triazines with respect to their inhibition of the growth of L1210/R71 cells 592(1992)59

Surface energy distribution

Experimental determination of adsorption isotherm data for the study of the surface energy distribution of various solid surfaces by inverse gas-solid chromatography 591(1992)233

Surface ionization detector

Determination of sultopride and tiapride in serum by gas chromatography using a surface ionisation detector 567(1991)113

Sensitive determination of phenothiazines in body fluids by gas chromatography with surface ionization detection 579(1992)247

Surface plasmon resonance

Biospecific interaction analysis using surface plasmon resonance detection applied to kinetic, binding site and concentration analysis 597(1992)397

Surface properties

Evaluation of fundamental properties of a silica capillary used for capillary electrophoresis 559(1991)69

Surface tension

Correlation analysis in liquid chromatography of metal chelates. II. One-dimensional retention-mobile phase composition (physicochemical property) models in reversed-phase liquid chromatography 555(1991)163

Surface water

Polar, hydrophilic compounds in drinking water produced from surface water. Determination by liquid chromatography-mass spectrometry 554(1991)251

Surfactants

Modification of reversed-phase columns with dyed surfactants. Preparation of mechanically resistant efficient immobilized dyes for protein purification 558(1991)43

Swelling behaviour

Beaded polymer supports and gels (Review). II. Physicochemical criteria and functionalization 586(1991)199

System peaks

Effects of β -cyclodextrin in the mobile phase on the retention and indirect detection of non-electrolytes in reversed-phase liquid chromatography. I. Study of aliphatic alcohols 552(1991)563

Adsorption isotherms of phenylalanine in a chromatographic column measured simultaneously by system peaks analysis and frontal analysis 556(1991)277

System selectivity

Enzymophoresis of nucleic acids by tandem capillary enzyme reactor-capillary zone electrophoresis 596(1992)251

T**Tailing reducers**

Gas chromatography in qualitative analysis. XX. The deactivation of diatomaceous supports by batyl alcohol 552(1991)169

Tandem mass spectrometry

Electrospray interfacing for the coupling of ion-exchange and ion-pairing chromatography to mass spectrometry 554(1991)27

Nanoscale separations combined with tandem mass spectrometry 554(1991)73

Liquid chromatography-thermospray tandem mass spectrometry for identification of a heptabarbital metabolite and sample work-up artefacts 554(1991)205

Polar, hydrophilic compounds in drinking water produced from surface water. Determination by liquid chromatography-mass spectrometry 554(1991)251

Structure determination of sesquiterpenes in Chinese vetiver oil by gas chromatography-tandem mass spectrometry 557(1991)451

Determination of sulfonamides by liquid chromatography, ultraviolet diode array detection and ion-spray tandem mass spectrometry with application to cultured salmon flesh 558(1991)155

Comparison of liquid-junction and coaxial interfaces for capillary electrophoresis-mass spectrometry with application to compounds of concern to the aquaculture industry 591(1992)325

On-line continuous-flow dialysis thermospray tandem mass spectrometry for quantitative screening of drugs in plasma: rogletimide 598(1992)189

Taxus brevifolia

Separation of taxol from related taxanes in *Taxus brevifolia* extracts by isocratic elution reversed-phase microcolumn high-performance liquid chromatography 587(1991)300

Temperature effects

Unusual effect of temperature on the retention of enantiomers on a chiral column 558(1991)1

Effect of temperature on the separation of DNA restriction fragments in capillary gel electrophoresis 559(1991)285

High-performance liquid chromatographic study of some biologically active analogues of arabinosylcytosine 588(1991)147

High-temperature open-tubular capillary column liquid chromatography 592(1992)239

Reversed- and normal-phase separations by high-temperature open-tubular column liquid chromatography 598(1992)153

Temperature gradients

Temperature elevations of the sample zone in free solution capillary electrophoresis under stacking conditions 559(1991)27

Temperature programming

Accuracy of computer-simulated gas chromatographic separations based on a linear elution strength model 589(1992)215

Van Deemter-type relationship for determining the optimum initial flow-rate and optimum pressure programming rate in temperature/pressure-programmed capillary column gas chromatography utilizing separation numbers 595(1992)221

Template assembled synthetic proteins

Retention behaviour of a template-assembled synthetic protein and its amphiphilic building blocks on reversed-phase columns 586(1991)43

Tenax GC trapping

Trace determination of volatile organic compounds in soil based on thermal vaporization followed by Tenax-GC trapping and capillary gas chromatography-mass spectrometry 555(1991)297

Tentacle-type anion exchangers

High-performance liquid chromatography of amino acids, peptides and proteins. CXX. Evaluation of bandwidth behaviour of proteins chromatographed on tentacle-type anion exchangers 599(1992)163

Terminating electrolyte

Dilution effect of carbonate ion on isotachophoretic zones observed with scanning UV photometric detection 555(1991)247

Ternary gradients

Studies on stress metabolites. XVI. High-performance liquid chromatographic analysis of cruciferous phytoalexins using complex ternary mobile phase gradients 598(1992)147

Ternary solvents

Comparison of selectivity of silica and Florisil in the separation of natural pyranocoumarins 552(1991)453

Test solutes

Influence of solute size and the non-polar interaction term on the selection of test solutes for the classification of stationary phase selectivity in gas chromatography 556(1991)457

Textile fibres

Analysis of dyes extracted from textile fibers by thermospray high-performance liquid chromatography-mass spectrometry 586(1991)73

Thermal conductivity detector

Determination of sulphur and nitrogen gases by gas chromatography on polystyrene porous polymer columns 585(1991)93

Thermal decomposition

Reaction-gas chromatography for physico-chemical investigations on aggressive gases 552(1991)319

Thermal desorption

Thermodesorption-gas chromatography-mass spectrometric analysis of biological materials for potential molecular precursors of the constituents of the crude oils 552(1991)325

Analysis of industrial solvent mixtures in water using a miniature purge-and-trap device with thermal desorption and capillary gas chromatography-mass spectrometry 557(1991)429

Thermal desorption-gas chromatography for the determination of benzene, aniline, nitrobenzene and chlorobenzene in workplace air 600(1992)344

Thermal vaporization

Trace determination of volatile organic compounds in soil based on thermal vaporization followed by Tenax-GC trapping and capillary gas chromatography-mass spectrometry 555(1991)297

Thermodynamic activity coefficients

Prediction of retention in gas-liquid chromatography using the UNIFAC group contribution method. II. Polymer stationary phases 585(1991)83

Thermodynamic behaviour

High-performance liquid chromatography of amino acids, peptides and proteins. CXV. Thermodynamic behaviour of peptides in reversed-phase chromatography 593(1992)103

Thermodynamic critical conditions

Analysis of block copolymers by high-performance liquid chromatography under critical conditions 593(1992)233

Thermodynamic parameters

Live retention database for identification in multi-step temperature-programmed capillary gas chromatography 552(1991)187

Effect of temperature and mobile phase on the retention of retinoates in reversed-phase liquid chromatography 592(1992)127

Thermodynamics of solution

Thermodynamics of solution of non-mesomorphic solutes at infinite dilution in the smectic-A, nematic and isotropic phases of *p*-*n*-octyl-*p*'-cyanobiphenyl. A gas-liquid chromatographic study 556(1991)317

Thermogravimetric analysis

Thermoanalytical and chromatographic studies of copper(II), nickel(II) and oxovanadium(IV) complexes of tetradentate Schiff bases derived from β -diketones and 2,3-diaminopentane 558(1991)187

Thermospray interface

Thermospray liquid chromatography-mass spectrometry for the characterisation of sulphate ester conjugates 554(1991)175

Thermospray ionization

Evaluation of eluents in thermospray liquid chromatography-mass spectrometry for identification and determination of pesticides in environmental samples 553(1991)311

Increasing thermospray response for cortisol by derivatization 554(1991)149

Use of methyl oxime derivatives to enhance structural information in thermospray high-performance liquid chromatography-mass spectrometry. Analysis of linoleic acid lipoxygenase metabolites in maize embryos 554(1991)155

Use of thermospray liquid chromatography-mass spectrometry to aid in the identification of urinary metabolites of a novel antiepileptic drug, Lamotrigine 554(1991)181

Analysis of gentamicin sulfate by high-performance liquid chromatography combined with thermospray mass spectrometry 554(1991)191

Liquid chromatography-thermospray tandem mass spectrometry for identification of a heptabarbital metabolite and sample work-up artefacts 554(1991)205

Evaluation of an automated thermospray liquid chromatography-mass spectrometry system for quantitative use in bioanalytical chemistry 554(1991)215

Determination of *Catharanthus roseus* alkaloids by high-performance liquid chromatography-isotope dilution thermospray-mass spectrometry 554(1991)227

Polar, hydrophilic compounds in drinking water produced from surface water. Determination by liquid chromatography-mass spectrometry 554(1991)251

Determination of taxol by high-performance liquid chromatography-thermospray mass spectrometry 594(1992)153

Thermospray mass spectrometry

Solvent degradation of cloxacillin *in vitro*. Tentative identification of degradation products using thermospray liquid chromatography-mass spectrometry 594(1992)195

Thin-layer chromatography, coupling to counter-current chromatography

Direct coupling of high-speed counter-current chromatography to thin-layer chromatography. Application to the separation of asiaticoside and madecassoside from *Centella asiatica* 558(1991)446

Thin-layer electrophoresis

Thin-layer electrophoresis of hydroxyethyl starches on a modified silica gel support 591(1992)351

Thin-layer plates

Comparison of commercially available thin-layer chromatography plates with mixtures of dyes, analgesics and phenols 595(1992)368

Throughput

Economic considerations important in the scale-up of an ovalbumin separation from hen egg-white on the anion-exchange cellulose DE92 590(1992)49

Adsorption kinetics of albumin on a cross-linked cellulose chromatographic ion exchanger 590(1992)101

Membrane chromatographic systems for high-throughput protein separations 596(1992)173

Time array detection

Complex mixture analysis based on gas chromatography-mass spectrometry with time array detection using a beam deflection time-of-flight mass spectrometer 590(1992)329

Time-resolved luminescence

Time-resolved luminescence detection of derivatized thiols in column liquid chromatography 552(1991)625

Titan's atmosphere

Gas chromatography of Titan's atmosphere. III. Analysis of low-molecular-weight hydrocarbons and nitriles with a CP-Sil-5 CB WCOT capillary column 591(1992)297

Tobacco

Determination of maleic hydrazide in tobacco by micellar liquid chromatography 595(1992)346

Estimation of tobacco blend compositions using closed-loop-stripping analysis and stepwise multiple linear regression and partial least-squares techniques 596(1992)285

Toria oil care

Role of metal ions in the ligand-exchange separation of amino acids 596(1992)281

Toxicity-based fractionation

Strategies for the identification of non-polar toxicants in aqueous environmental samples using toxicity-based fractionation and gas chromatography-mass spectrometry (Review) 580(1992)215

Toxicology

see also Special issue index

Trace analysis

Trace determination of volatile organic compounds in soil based on thermal vaporization followed by Tenax-GC trapping and capillary gas chromatography-mass spectrometry 555(1991)297

Peak-shape analysis and noise evaluation in suppressed ion chromatography for ultra-trace ion analysis 556(1991)249

Determination of a metabolite of the herbicide pyridate in drinking and groundwater using high-performance liquid chromatography with amperometric detection 558(1991)369

Determination of trace amounts of fluoride in raw materials for pharmaceuticals by gas-liquid chromatography 585(1991)289

Comparison of several methods for the determination of trace amounts of polar aliphatic monocarboxylic acids by high-performance liquid chromatography 593(1992)243

Reversed-phase liquid chromatographic behaviour of alkylamines with amperometric detection and its application to trace analysis 594(1992)159

Determination of major ions in snow and ice cores by ion chromatography 594(1992)225

Determination of inorganic anions in salt solutions by ion chromatography using C₁₈ reversed-phase columns coated with cetyltrimethylammonium 598(1992)237

Determination of divalent trace metals in natural waters by preconcentration on N,N,N',N'-tetra(2-aminoethyl)ethylenediamine-silica followed by on-line ion chromatography 600(1992)203

Trace enrichment

Rapid method for the determination of organochlorine pesticides in milk 552(1991)249

Interference removal in the organic trace-level analysis of aqueous environmental samples by on-line liquid chromatographic preconcentration techniques with two precolumns 553(1991)329

Strategies in developing interfaces for coupling liquid chromatography and mass spectrometry 554(1991)3

Rapid on-line precolumn high-performance liquid chromatographic method for the determination of benomyl, carbendazim and aldicarb species in drinking water 555(1991)147

Chromatographic trace analysis of some organic compounds in the environment using derivatization-sorption concentration techniques. I. Gas chromatographic analysis of acrylates in air 557(1991)523

Coupled-column high-performance liquid chromatographic method for the determination of I-hydroxypyrene in urine of subjects exposed to polycyclic aromatic hydrocarbons 600(1992)189

Trace-level determination of polar phenolic compounds in aqueous samples by high-performance liquid chromatography and on-line preconcentration on porous graphitic carbon 600(1992)195

Transalkylation reaction

Analytical study of heavy crude oil fractions by coupling of the transalkylation reaction with supercritical fluid chromatography 593(1992)321

Transport parameters

Predictability of chromatographic protein separations. Study of size-exclusion media with narrow particle size distributions 589(1992)71

Transverse particle distribution

Transverse particle redistribution in a flat channel for SPLIT or integral Doppler anemometry 600(1992)133

Trapezoidal cross-section channel

Isoelectric focusing field-flow fractionation. III. Investigation of the influence of different experimental parameters on focusing of cytochrome *c* in the trapezoidal cross-section channel 600(1992)297

Isoelectric focusing field-flow fractionation. IV. Investigations on protein separations in the trapezoidal cross-section channel 600(1992)305

Trapping methods

Simple method for collecting volatile compounds from single insects and other point sources for gas chromatographic analysis 598(1992)303

Trapping system

Trapping system for trace organic volatiles 586(1991)315

s-Triazine-activated Sephacryl

Coupling of *m*-aminophenylboronic acid to *s*-triazine-activated Sephacryl: use in the affinity chromatography of glycosylated hemoglobins 575(1992)223

Trichoderma

Quantitative separation of *Trichoderma* lipid classes on a bonded phase column 584(1992)129

Tryptic digests

Recent advances in liquid chromatography–mass spectrometry and capillary zone electrophoresis–mass spectrometry for protein analysis 553(1991)101

Peptide maps of five human pepsin isoenzymes and other aspartic proteinases 599(1992)179

Tryptic mapping

Characterization of a tryptic digest by high-performance displacement chromatography and mass spectrometry 557(1991)289

Two-dimensional electrophoresis

Important design features of a system for the densitometric analysis of two-dimensional flat-bed separations 596(1992)241

Two-dimensional liquid chromatography

Automated two-dimensional liquid chromatographic system for mapping proteins in highly complex mixtures 588(1991)115

Two-dimensional mapping

Two-dimensional mapping by high-performance liquid chromatography of pyridylamino oligosaccharides from various glycosphingolipids 586(1991)35

Two-dimensional separations

Comparison of statistical theories of spot overlap in two-dimensional separations and verification of means for estimating the number of zones 591(1992)1

Two-dimensional separations, semi preparative

Combination of supercritical fluid chromatography with thin-layer chromatography on a semi-preparative scale 552(1991)539

Two-phase systems

Ligand–receptor interactions in affinity cell partitioning. Studies with transferrin covalently linked to monomethoxypoly(ethylene glycol) and rat reticulocytes 594(1992)97

Tyrosine-proline cleavage

Microbore liquid chromatography coupled to a flow fast atom bombardment probe for the on-line detection of the Tyr-Pro cleavage of a nonapeptide by recombinant HIV-1 protease 562(1991)67

U***Ulex europaeus***

Purification and characterization of carbohydrate-binding peptides from *Lotus tetragonolobus* and *Ulex europaeus* seed lectins using affinity chromatography 597(1992)213

Ultrafiltration membranes

Chromatographic fractionation of nucleic acids using microcapsules made from plant cells 585(1991)57

Ultraviolet absorbance detector

Capillary gel electrophoresis of single-stranded DNA fragments with UV detection 600(1992)289

Ultraviolet spectrophotometry

UV spectrophotometric detection in the ion chromatographic analysis of alkali and alkaline earth metal and ammonium cations 591(1992)219

UNIFAC group contribution method

Prediction of retention in gas–liquid chromatography using the UNIFAC group contribution method. II. Polymer stationary phases 585(1991)83

Prediction of retention in gas–liquid chromatography using the UNIFAC group contribution method. III. Recent developments in UNIFAC 586(1991)297

UNIFAC model

UNIFAC model as a heuristic guide for estimating retention in chromatography 586(1991)1

Universal calibration

Interfacial effects in size-exclusion chromatography of latex 587(1991)276

On-line characterization of polyethylene glycol-modified proteins 588(1991)125

Universal sulphur chemiluminescence detector

Determination of sulfur components in light petroleum streams by high-resolution gas chromatography with chemiluminescence detection 589(1992)271

Unmodified silica

Adsorption and partition mode in high-performance liquid chromatography of highly polar solutes on silica 552(1991)23

Unresolved complex mixtures

Use of oxidative degradation followed by capillary gas chromatography–mass spectrometry and multi-dimensional scaling analysis to fingerprint unresolved complex mixtures of hydrocarbons 589(1992)281

Urban air

Photoionization assessment of C₃-C₅ alkadienes and alkenes in urban air 591(1992)358

V**Vaccines**

Analysis of a vaccine purification process by capillary electrophoresis 559(1991)337

Van Deemter equation, modified

Protein chromatography using a continuous stationary phase 598(1992)169

Van Deemter plots

Source of band broadening in liquid chromatographic-fast atom bombardment mass spectrometric systems with precolumn addition of viscous matrix to the mobile phase 554(1991)47

Utility of small-particle silica in preparative chromatography 590(1992)77

Limits and effects of precolumn addition of thioglycerol in liquid chromatographic-fast atom bombardment mass spectrometric systems 591(1992)129

Critical capillary column examination of the relationship between separation number and height equivalent to a theoretical plate and their dependence on temperature 595(1992)209

Van Deemter-type relationship for determining the optimum initial flow-rate and optimum pressure programming rate in temperature/pressure-programmed capillary column gas chromatography utilizing separation numbers 595(1992)221

Van 't Hoff plots

Unusual effect of temperature on the retention of enantiomers on a chiral column 558(1991)1

Variable coefficient parameters

Optimization of an ion-exchange high-performance liquid chromatographic method for the determination of carboxylic acids, sugars, glycerol and ethanol in wines 589(1992)151

Variable size simplex approach

Optimization of the separation of (6*R*)- and (6*S*)-leucovorin and evaluation of the robustness of the optimum 592(1992)37

Velocity profiles

Simple solution of velocity profiles of laminar flows in channels of various cross-sections used in field-flow fractionation 593(1992)357

Vesicles

Chromatographic fractionation of nucleic acids using microcapsules made from plant cells 585(1991)57

Vesicular packing

Inclusion and fractionated release of nucleic acids using microcapsules made from plant cells 596(1992)165

Vetiver oil

Structure determination of sesquiterpenes in Chinese vetiver oil by gas chromatography-tandem mass spectrometry 557(1991)451

Viscosity

Correlation analysis in liquid chromatography of metal chelates. II. One-dimensional retention-mobile phase composition (physicochemical property) models in reversed-phase liquid chromatography 555(1991)163

Viscous matrices

Effect of the addition of viscous matrices to the mobile phase on chromatographic performance in liquid chromatography-fast atom bombardment mass spectrometry 554(1991)61

Vlasta oil

Gas chromatography and gas chromatography-mass spectrometry study of hydrocarbons in Vlasta oil (Adriatic Basin) as the basis for geochemical interpretation 557(1991)255

Void volume

Separation of *trans/cis* α - and β -carotenes by supercritical fluid chromatography. II. Effect of the type of octadecyl-bonded stationary phase on retention and selectivity of carotenes 557(1991)59

Volatile oils

Cholesteryl acetate as a stationary phase for the gas chromatography of some volatile oil constituents 600(1992)337

Voltammetry

Capillary electrophoresis with electrochemical detection employing an on-column Nafion joint 593(1992)305

W**Water, mutagenic XAD-2 extracts**

Characterization and determination of organic compounds in the mutagenic XAD-2 extracts of drinking water 557(1991)413

Water analysis

Gas chromatographic-mass spectrometric characterization of volatile organic compounds in Barcelona tap water 562(1991)481

Water pollution

Characterization and determination of organic compounds in the mutagenic XAD-2 extracts of drinking water 557(1991)413

Western blotting

Purification of canine prolactin and growth hormone by fast protein liquid chromatography 553(1991)117

Wheat cultivars

Characterization and identification of wheat cultivars by multi-dimensional analysis of reversed-phase high-performance liquid chromatograms 596(1992)225

Whey protein hydrolysate

Determination of molecular mass distributions of whey protein hydrolysates by high-performance size-exclusion chromatography 599(1992)205

Window analysis

Theoretical and experimental comparison of serially linked and mixed-packing gas-liquid chromatography columns 555(1991)173

Window diagram method

Optimisation pentaparamétrique de la résolution en chromatographie phase gazeuse par la technique du diagramme à fenêtres. Cas du couplage de deux colonnes 556(1991)287

Wine

Optimization of an ion-exchange high-performance liquid chromatographic method for the determination of carboxylic acids, sugars, glycerol and ethanol in wines 589(1992)151

Analytical methods for monoterpene glycosides in grape and wine. I. XAD-2 extraction and gas chromatographic-mass spectrometric determination of synthetic glycosides 590(1992)313

X**X-Ray photoelectron spectroscopy**

Physical and chemical characterization of a porous phosphate-modified zirconia substrate 587(1991)137

Z**Zapatera olives**

Analysis of zapatera olives by gas and high-performance liquid chromatography 594(1992)259

Zinc amalgam scrubber column

Reductive electrochemical detection in liquid chromatography with a zinc amalgam scrubber column 598(1992)195

Zirconium oxide supports

Physical and chemical characterization of a porous phosphate-modified zirconia substrate 587(1991)137

Chromatographic characterization of a phosphate-modified zirconia support for bio-chromatographic applications 587(1991)149

Ion- and ligand-exchange chromatography of proteins using porous zirconium oxide supports in organic and inorganic Lewis base eluents 596(1992)27

Zone electrophoresis

Theoretical background for clinical and biomedical applications of electromigration techniques (Review) 569(1991)3

Zone electrophoretic sample treatment

Rapid electro-dialytic clean-up of biological samples for high-performance liquid chromatography 585(1991)213

Zone length

Relationship between zone length and step height in isotachopheresis 595(1992)327

Zone migration

Variance of a zone migrating in a non-uniform time-invariant linear medium 596(1992)1

Zone overlap

Comparison of statistical theories of spot overlap in two-dimensional separations and verification of means for estimating the number of zones 591(1992)1

Zonoid

Is chromatography a separation process? The zonoid answer (Review) 556(1991)25

COMPOUND INDEX

VOLUMES 551-600

A

- Abamectin**
Rapid determination of abamectin in lettuce and cucumber by high-performance liquid chromatography 553(1991)299
- Abscisic acid**
High-performance thin-layer chromatographic and high-performance liquid chromatographic determination of abscisic acid produced by cyanobacteria 555(1991)239
- Absynthin**
Use of high-performance liquid chromatographic peak deconvolution and peak labelling to identify antiparasitic components in plant extracts 593(1992)209
- Acenocoumarol**
Thermospray and particle beam liquid chromatographic-mass spectrometric analysis of coumarin anticoagulants 562(1991)31
- Acephate**
Determination of acephate by liquid chromatography in the presence of aqueous soil extracts 585(1991)324
- Acetaldehyde**
Excretion of formaldehyde, malondialdehyde, acetaldehyde and acetone in the urine of rats in response to 2,3,7,8-tetrachlorodibenzo-*p*-dioxin, paraquat, endrin and carbon tetrachloride 576(1992)221
- Analysis of zapatera olives by gas and high-performance liquid chromatography 594(1992)259
- Acetaminophen**
see Paracetamol
- Acetanilide**
Reversed-phase liquid chromatography coupled on-line with capillary gas chromatography. I. Introduction of large volumes of aqueous mixtures through an on-column interface 552(1991)489
- Acetazolamide**
Determination of acetazolamide in human urine samples by reversed-phase high-performance liquid chromatography in the presence of xanthines 582(1992)181
- Acetic acid**
Determination of volatile and non-volatile organic acids in technical sugar solutions by ion-exclusion chromatography 555(1991)65
- Acetoin**
Simple and sensitive determination of diacetyl and acetoin in biological samples and alcoholic drinks by gas chromatography with electron-capture detection 577(1992)215
- Acetone**
Estimation of cellulose polarity by gas chromatography 552(1991)365
- High-performance liquid chromatographic determination of acetone in blood and urine in the clinical diagnostic laboratory 553(1991)249
- Development and use of a stability-indicating high-performance liquid chromatographic assay for Meldrum's acid 575(1992)177
- Excretion of formaldehyde, malondialdehyde, acetaldehyde and acetone in the urine of rats in response to 2,3,7,8-tetrachlorodibenzo-*p*-dioxin, paraquat, endrin and carbon tetrachloride 576(1992)221
- Direct aqueous injection gas chromatography on a potassium fluoride crystal hydrate-containing sorbent. Determination of volatile organic solvents in the fermentation broth of *Clostridium* strains 596(1992)138
- Acetonitrile**
Estimation of cellulose polarity by gas chromatography 552(1991)365

Acetoxyscirpenols

Identification of trichothecenes by thermospray, plasmaspray and dynamic fast-atom bombardment liquid chromatography-mass spectrometry 562(1991)555

Acetylacetone

Photoemissive ionisation source for ion mobility detectors 588(1991)239

N-Acetylamino acids

Determination of free N-acetylamino acids in biological samples and N-terminal acetylamino acids of proteins 576(1992)63

Acetylamino-*n*-propylmorpholine

Polyacrylamide gel polymerization under non-oxidizing conditions, as monitored by capillary zone electrophoresis 598(1992)287

Acetyl-5-aminosalicylic acid

High-performance liquid chromatographic assay for the determination of 5-aminosalicylic acid and acetyl-5-aminosalicylic acid concentrations in endoscopic intestinal biopsy in humans 564(1991)296

High-performance liquid chromatographic assay of 5-aminosalicylic acid (5-ASA) and its metabolites N- β -D-glucopyranosyl-5-ASA, N-acetyl-5-ASA, N-formyl-5-ASA and N-butyryl-5-ASA in biological fluids 570(1991)109

N-Acetyl-L-aspartic acid

Quantitative measurement of N-acetyl-L-aspartic acid in urine by gas chromatography with negative-ion chemical ionization mass spectrometry 578(1992)116

Acetylcholine

Application of high-performance liquid chromatography to the study of biogenic amine-related enzymes (Review) 566(1991)287

Glassy carbon pre-column for direct determination of acetylcholine and choline in biological samples using liquid chromatography with electrochemical detection 575(1992)29

Cation-exchange liquid chromatography of choline and acetylcholine on free shielded silanols of silica-based reversed-phase stationary phases 596(1992)43

Acetylcholine esterase inhibitor

Determination of enantiomers of 1-benzyl-4-[(5,6-dimethoxy-1-indanon)-2-yl]methylpiperidine hydrochloride (E2020), a centrally acting acetylcholine esterase inhibitor, in plasma by liquid chromatography with fluorometric detection 577(1992)95

Acetyl-coenzyme A

Specific detection of acetyl-coenzyme A by reversed-phase ion-pair high-performance liquid chromatography with an immobilized enzyme reactor 590(1992)241

Acetyl-Cys-Asn- β , β -dimethylthioprolino-*p*-aminomethylphenyl-alanine-Gly-Asp-Cys-OH

Determination of a cyclic heptapeptide, a novel fibrinogen receptor antagonist, in human plasma by high-performance liquid chromatography with automated pre-column derivatization, column switching and fluorescence detection 578(1992)31

N-Acetylcysteine

Determination of N-acetylcysteine in human plasma by gas chromatography-mass spectrometry 562(1991)639

2-Acetyl-2-decarboxamidometacycline

Isolation of doxycycline, 6-epidoxycycline and 2-acetyl-2-decarboxamidometacycline from commercial metacycline by preparative column liquid chromatography on silica gel 586(1991)67

L-1-Acetyl-2,3-dihydropyrrolo[2,3-*b*]-indole-2-carboxamide

Occurrence of an allosteric transition in the modification of papain with L-1-acetyl-2,3-dihydropyrrolo[2,3-*b*]-indole-2-carboxamide 597(1992)411

N-Acetyldopamine

Determination by high-performance liquid chromatography with electrochemical detection of free and conjugated N-acetyldopamine excretion in urine of children with neuroblastoma and nephroblastoma 574(1992)9

Sensitivity and specificity of urinary N-acetyldopamine as a marker for neuroblastomas: comparison with traditional urinary catecholamine metabolites 579(1992)184

Sensitivity and specificity of urinary N-acetyldopamine as a marker for neuroblastomas: comparison with traditional urinary catecholamine metabolites. Reply to Muskiet and Kema 579(1992)187

N-Acetylisoptreanine- γ -lactam

Gas chromatographic determination of N-acetylisoptreanine- γ -lactam, a unique catabolite of N¹-acetylspermidine 563(1991)1

6-Acetylmorphine

Determination of morphine and 6-acetylmorphine in plasma by high-performance liquid chromatography with fluorescence detection 566(1991)135

Rapid assay of cocaine, opiates and metabolites by gas chromatography-mass spectrometry (Review) 580(1992)43

N-Acetyloligosaccharides

Capillary zone electrophoresis of linear and branched oligosaccharides 600(1992)279

N-Acetylpsychosine

Use of N-acetylpsychosine as internal standard for quantitative high-performance liquid chromatographic analysis of glycosphingolipids 555(1991)267

Acetylsalicylic acid

Simultaneous high-performance liquid chromatographic determination of salicylates in whole blood, plasma and isolated erythrocytes 584(1992)224

Acetylselenonium choline

Synthesis of novel selenium-containing choline and acetylcholine analogues and their quantitation using a pyrolysis-gas chromatography-mass spectrometry assay 585(1991)101

N4-Acetylsulphadimidine

High-performance liquid chromatographic method for the routine determination of sulphadimidine, its hydroxy metabolites and N4-acetylsulphadimidine in body fluids and cell culture media 571(1991)157

 α_1 -Acid glycoprotein

Liquid chromatographic method for the determination of the carbohydrate moiety of glycoproteins. Application to α_1 -acid glycoprotein and tissue plasminogen activator 558(1991)105

Rapid and simple method for the determination of α_1 -acid glycoprotein in serum by column liquid chromatography 582(1992)246

 α_1 -Acid glycoprotein, desialylated

pH titration curves of the desialylated human α_1 -acid glycoprotein variants by combined isoelectrofocusing-electrophoresis: utilization in the development of a fractionation method for the protein variants by chromatography on immobilized metal affinity adsorbent 577(1992)43

Acidic sulphur-containing amino acids

Automated determination of neuroactive acidic sulphur-containing amino acids and γ -glutamyl peptides using liquid chromatography with fluorescence and electrochemical detection 566(1991)39

Acid phosphatase

Electrophoretic phenotyping of erythrocyte enzymes (Review) 569(1991)297

Acids

Application of triple quadrupole tandem mass spectrometry to the analysis of pyridine-containing derivatives of long-chain acids and alcohols 579(1992)1

Prediction of retention indexes. III. Silylated derivatives of polar compounds 586(1991)113

Acisoga

Gas chromatographic determination of N-acetylispoutreanine- γ -lactam, a unique catabolite of N¹-acetylspermidine 563(1991)1

Acitretin

Quantification of acitretin in human plasma by microbore liquid chromatography-negative chemical ionization mass spectrometry 568(1991)135

Improved quantitation of 13-*cis*- and all-*trans*-acitretin in human plasma by normal-phase high-performance liquid chromatography 570(1991)149

Use of direct injection precolumn techniques for the high-performance liquid chromatographic determination of the retinoids acitretin and 13-*cis*-acitretin in plasma 593(1992)55

Acridine

Characterization of nitrogen-containing aromatic compounds in soil and sediment by capillary gas chromatography-mass spectrometry after fractionation 558(1991)223

Acrylamido buffers

Capillary zone electrophoresis analysis of acrylamido buffers for isoelectric focusing in immobilized pH gradients 559(1991)119

Acrylates

Chromatographic trace analysis of some organic compounds in the environment using derivatization-sorption concentration techniques. I. Gas chromatographic analysis of acrylates in air 557(1991)523

Acrylic acid

Continuous beds for standard and micro high-performance liquid chromatography 586(1991)21

Acryloyl histamine

Synthesis of a new acrylamido buffer (acryloylhistamine) for isoelectric focusing in immobilized pH gradients and its analysis by capillary zone electrophoresis 558(1991)285

Actinides

Reversed-phase separation of transition metals, lanthanides and actinides by elution with mandelic acid 558(1991)197

Activation agents

Precolumn fluorescence tagging reagent for carboxylic acids in high-performance liquid chromatography: 4-substituted-7-aminoalkylamino-2,1,3-benzoxadiazoles 588(1991)61

Acyclovir

New, high-sensitivity high-performance liquid chromatographic method for the determination of acyclovir in human plasma, using fluorometric detection 583(1992)122

Acylbiotinols

Characterization of N-acyl-D-biotinols by particle-beam liquid chromatography-mass spectrometry. An alternative to probe mass spectrometry for thermally labile samples 554(1991)141

Acylcarnitines

Biomedical applications of high-performance liquid chromatography-mass spectrometry with continuous-flow fast atom bombardment 562(1991)47

Identification of urinary acylcarnitines using gas chromatography-mass spectrometry: preliminary clinical applications 577(1992)205

Acyl-coenzyme A thioesters

Biomedical applications of high-performance liquid chromatography–mass spectrometry with continuous-flow fast atom bombardment 562(1991)47

Improved and simplified tissue extraction method for quantitating long-chain acyl-coenzyme A thioesters with picomolar detection using high-performance liquid chromatography 577(1992)157

ADD17014

Triazolines. XXIII. High-performance liquid chromatographic assay in rat blood for a novel triazoline anticonvulsant (ADD17014) 563(1991)419

Adenine

Extension of the electrostatic retention model of reversed-phase ion-pair high-performance liquid chromatography to include the effect of the eluent pH 552(1991)13

Determination of concentrations of adenosine and other purines in human term placenta by reversed-phase high-performance liquid chromatography with photodiode-array detection: evidence for pathways of purine metabolism in the placenta 575(1992)243

Adenine nucleotides

Isocratic high-performance liquid chromatographic analysis of myocardial creatine phosphate and adenine nucleotides 563(1991)129

Simple cation-exchange high-performance liquid chromatography optimized to the measurement of metabolites in the effluents from perfused rat livers using refractive index and ultraviolet detectors 563(1991)231

Adenosine

Subtraction method for the high-performance liquid chromatographic measurement of plasma adenosine 574(1992)247

Determination of concentrations of adenosine and other purines in human term placenta by reversed-phase high-performance liquid chromatography with photodiode-array detection: evidence for pathways of purine metabolism in the placenta 575(1992)243

Enzymophoresis of nucleic acids by tandem capillary enzyme reactor–capillary zone electrophoresis 596(1992)251

Adenosine, metabolites

Rapid separation of creatine, phosphocreatine and adenosine metabolites by ion-pair reversed-phase high-performance liquid chromatography in plasma and cardiac tissue 576(1992)149

A₁ Adenosine receptor

Affinity chromatography in purification of A₁ adenosine receptors 597(1992)335

Adenosine 5'-triphosphate

Use of hydrophilic interaction chromatography for the study of tyrosine protein kinase specificity 583(1992)137

S-Adenosyl-L-homocysteine

High-performance liquid chromatographic determination, with ultraviolet detection, of S-adenosyl-L-methionine and S-adenosyl-L-homocysteine in rat tissues and simultaneously of normetanephrine and metanephrine for phenylethanolamine-N-methyltransferase or catechol-O-methyltransferase activities 567(1991)254

S-Adenosyl-L-methionine

High-performance liquid chromatographic determination, with ultraviolet detection, of S-adenosyl-L-methionine and S-adenosyl-L-homocysteine in rat tissues and simultaneously of normetanephrine and metanephrine for phenylethanolamine-N-methyltransferase or catechol-O-methyltransferase activities 567(1991)254

Quantification of erythrocyte S-adenosyl-L-methionine levels and its application in enzyme studies 576(1992)95

Adosupine

Determination of the tricyclic compound adosupine and its three metabolites in plasma and brain of rat using high-performance liquid chromatography 574(1992)319

ADR-851

Quantitation of a novel antiemetic (ADR-851) in plasma and urine by reversed-phase high-performance liquid chromatography with fluorescence detection 566(1991)257

Adrenaline

see Epinephrine

Adriamycin

see Doxorubicin

Aflatoxins

Determination of aflatoxin M1 using a dialysis-based immunoaffinity sample pretreatment system coupled on-line to liquid chromatography. Reusable immunoaffinity columns 589(1992)141

Automated determination of aflatoxin B₁ in cattle feed by two-column solid-phase extraction with on-line high-performance liquid chromatography 592(1992)265

AG-EE 388 ZW

Quantitation of the new hypoglycaemic agent AG-EE 388 ZW in human plasma by automated high-performance liquid chromatography with electrochemical detection 568(1991)246

Aglycones

Measurement of epidoxorubicin and its metabolites by high-performance liquid chromatography using an advanced automated sample processor 572(1991)211

Determination of potato glycoalkaloids and their aglycone in blood serum by high-performance liquid chromatography. Application to pharmacokinetic studies in humans 573(1992)69

High-performance liquid chromatographic analysis of the pigments of blood-red prickly pear (*Opuntia ficus indica*) 593(1992)177

Analytical methods for monoterpene glycosides in grape and wine. II. Qualitative and quantitative determination of monoterpene glycosides in grape 595(1992)269

β -Agonists

Screening and confirmatory analysis of β -agonists, β -antagonists and their metabolites in horse urine by capillary gas chromatography-mass spectrometry 564(1991)503

Residue screening for the β -agonists clenbuterol, salbutamol and cimaterol in urine using enzyme immunoassay and high-performance liquid chromatography 564(1991)551

AJ-3941

Determination of AJ-3941, a possible agent for the treatment of cerebrovascular disorders, in plasma and brain by means of high-performance liquid chromatography with fluorescence detection 578(1992)152

Ajmalicine

Determination of *Catharanthus roseus* alkaloids by high-performance liquid chromatography-isotope dilution thermospray-mass spectrometry 554(1991)227

Ajmaline

Monitoring of ajmaline in plasma with high-performance liquid chromatography 575(1992)87

Ajmaline N-oxide

Monitoring of ajmaline in plasma with high-performance liquid chromatography 575(1992)87

Alanine

Application of ligand-exchange chromatography to the assay of L-alanine from DL-aspartic acid by *Pseudomonas dacunhae* 555(1991)272

Determination of D-alanine and D-glutamic acid in biological samples by coupled-column chromatography using β -cyclodextrin as mobile phase additive 582(1992)35

Alanine aminotransferase

Electrophoretic analysis of enzyme-linked immunoglobulins and their clinical significance (Review) 569(1991)347

Analysis of alanine and aspartate aminotransferase isoforms in mustard (*Sinapis alba* L.) cotyledons 587(1991)101

Albendazole

Rapid and sensitive method for the determination of albendazole and albendazole sulphoxide in biological fluids 566(1991)244

Albendazole sulphoxide

Rapid and sensitive method for the determination of albendazole and albendazole sulphoxide in biological fluids 566(1991)244

Direct liquid chromatographic separation of enantiomers on immobilized protein stationary phases. IX. Influence of the cross-linking reagent on the retentive and enantioselective properties of chiral sorbents based on bovine serum albumin 591(1992)65

Albumin

see also Human serum albumin

Chromatographic characterization of ion exchangers for high-performance liquid chromatography of proteins. I. Chromatographic determination of loading capacity for low- and high-molecular mass anions 556(1991)363

High-performance liquid chromatography of amino acids, peptides and proteins. CXIV. Protein interactions with porous coulombic sorbents: comparison of experimental findings with predictions of several adsorption models 557(1991)335

Adsorption kinetics of albumin on a cross-linked cellulose chromatographic ion exchanger 590(1992)101

High-sensitivity laser-induced fluorescence detection of native proteins in capillary electrophoresis 595(1992)319

Novel affinity separations based on perfluorocarbon emulsions. Use of a perfluorocarbon affinity emulsion for the purification of human serum albumin from blood plasma in a fluidised bed 597(1992)189

High-performance affinity chromatography system for the rapid, efficient assay of glycated albumin 597(1992)271

Assay of protein drug substances present in solution mixtures by fluorescamine derivatization and capillary electrophoresis 598(1992)123

Contribution of ionically immobilized bovine serum albumin to the retention of enantiomers 600(1992)37

Albumin Casebrook

Characterisation of oligosaccharides from a glycoprotein variant of human serum albumin (albumin Casebrook) using high-performance anion-exchange chromatography and nuclear magnetic resonance spectroscopy 581(1992)187

Alcohols

Gas chromatography in qualitative analysis. XX. The deactivation of diatomaceous supports by butyl alcohol 552(1991)169

Gas chromatographic separation of unsaturated polar compounds on inorganic stationary phases containing silver nitrate 552(1991)313

Gas chromatographic analysis of some glycol ether analytes on a specially packed column 552(1991)353

Effects of β -cyclodextrin in the mobile phase on the retention and indirect detection of non-electrolytes in reversed-phase liquid chromatography. I. Study of aliphatic alcohols 552(1991)563

Use of an evaporative light scattering detector in reversed-phase high-performance liquid chromatography of oligomeric surfactants 552(1991)593

Chromatographic determination of the physico-chemical parameters of adsorption on active carbons 556(1991)441

Application of triple quadrupole tandem mass spectrometry to the analysis of pyridine-containing derivatives of long-chain acids and alcohols 579(1992)1

Separation of secondary alcohol enantiomers using supercritical fluid chromatography 585(1991)117

Determination of alcohols by high-performance liquid chromatography after pre-column derivatization with 2-(4-carboxyphenyl)-5,6-dimethylbenzimidazole 585(1991)219

Prediction of retention indexes. III. Silylated derivatives of polar compounds 586(1991)113

Predicting partition coefficients in polyethylene glycol-potassium phosphate aqueous two-phase systems 586(1991)341

Dehydrogenase-silica as a stationary phase for the separation of alcohols and ketones 587(1991)268

Enantiomeric separation of racemic hydroperoxides and related alcohols 588(1991)352

Preparation and capillary column gas chromatographic characterization of aza-crown ether polysiloxane stationary phases 594(1992)243

Some novel homochiral derivatizing agents for the gas chromatographic analysis of enantiomeric secondary alcohols 598(1992)251

High-performance liquid chromatographic separation of chiral metallocenic ketones and alcohols 600(1992)320

Aldicarb

Rapid on-line precolumn high-performance liquid chromatographic method for the determination of benomyl, carbendazim and aldicarb species in drinking water 555(1991)147

Alditol acetates

Structure elucidation of saccharides in anthocyanins and flavonols by means of methylation analysis and gas chromatography 598(1992)81

Alditol acetates, methylated

Clean-up procedure for partially methylated alditol acetate derivatives of polysaccharides 587(1991)284

Alditols

Structure-retention index relationships for derivatized monosaccharides on non-polar gas chromatography columns 596(1992)79

Aldohexoses

see also Sugars

Carbohydrate analysis with ion chromatography using Eurokat stationary phases. Preparative separation of monosaccharides and their fluorinated derivatives 552(1991)483

Aldolase

Recent developments in quantitative affinity chromatography (Review) 597(1992)67

Alendronate

Improved determination of the bisphosphonate alendronate in human plasma and urine by automated precolumn derivatization and high-performance liquid chromatography with fluorescence and electrochemical detection 583(1992)183

Determination of alendronate in pharmaceutical dosage formulations by ion chromatography with conductivity detection 596(1992)217

Alentamol

Determination of alentamol hydrobromide, a novel antipsychotic agent, in human blood plasma and urine by high-performance liquid chromatography with fluorescence detection and solid-phase extraction 565(1991)488

Alkadienes

Photoionization assessment of C₃-C₅ alkadienes and alkenes in urban air 591(1992)358

Alkali metals

Secondary equilibrium size-exclusion chromatography of ions with polymeric mobile phase additives 586(1991)277

Membrane-based sample preparation device for the pretreatment of acidic samples prior to cation analysis by ion chromatography 586(1991)283

UV spectrophotometric detection in the ion chromatographic analysis of alkali and alkaline earth metal and ammonium cations 591(1992)219

Factors affecting the separation of inorganic metal cations by capillary electrophoresis 593(1992)289

Alkali metal salts

Modelling of displacement chromatography using non-ideal isotherms 594(1992)9

Alkaline earth metals

Membrane-based sample preparation device for the pretreatment of acidic samples prior to cation analysis by ion chromatography 586(1991)283

High-performance capillary zone electrophoresis of carbohydrates in the presence of alkaline earth metal ions 588(1991)327

UV spectrophotometric detection in the ion chromatographic analysis of alkali and alkaline earth metal and ammonium cations 591(1992)219

Factors affecting the separation of inorganic metal cations by capillary electrophoresis 593(1992)289

Alkaline phosphatase

High-performance capillary electrophoresis as a fast in-process control method for enzyme-labelled monoclonal antibody conjugates 559(1991)385

Diagnostic aspects of alkaline phosphatase: separation of isoenzymes in normal and pathological human serum by high-performance liquid chromatography 563(1991)147

Electrophoretic analysis of enzyme-linked immunoglobulins and their clinical significance (Review) 569(1991)347

Interactions of human alkaline phosphatase isoenzymes with triazine dyes using affinity partitioning, affinity chromatography and difference spectroscopy 574(1992)237

Alkaline phosphatase isoenzymes

Determination of alkaline phosphatase isoenzymes in serum by high-performance liquid chromatography with post-column reaction detection 576(1992)79

Alkaline protease

Detection of enzyme activity in fractions collected from free solution capillary electrophoresis of complex samples 559(1991)325

Alkaloids

Determination of *Catharanthus roseus* alkaloids by high-performance liquid chromatography-isotope dilution thermospray-mass spectrometry 554(1991)227

Pseudo-electrochromatography-mass spectrometry: a new alternative 554(1991)339

High-performance liquid chromatographic-mass spectrometric assay of high-value compounds for pharmaceutical use from plant cell tissue culture: *Cinchona* alkaloids 562(1991)713

Analysis of tropane alkaloids with thermospray high-performance liquid chromatography-mass spectrometry 562(1991)737

Determination of yohimbine and its two hydroxylated metabolites in humans by high-performance liquid chromatography and mass spectral analysis 574(1992)283

Separation of *Strychnos nux-vomica* alkaloids by high-performance liquid chromatography 587(1991)318

Analysis and isolation of indole alkaloids of fungi by high-performance liquid chromatography 593(1992)201

Analytical studies of isorhoadine and rhoegenine in petal extracts of *Papaver rhoeas* L. using high-performance liquid chromatography 596(1992)276

Alkanes

Live retention database for identification in multi-step temperature-programmed capillary gas chromatography 552(1991)187

Study of microporous carbons by gas chromatographic determination of heats of physisorption 555(1991)183

Thermodynamics of solution of non-mesomorphic solutes at infinite dilution in the smectic-A, nematic and isotropic phases of *p-n*-octyl-*p'*-cyanobiphenyl. A gas-liquid chromatographic study 556(1991)317

Enhanced resolution of organic compounds from sediments by isotopic gas chromatography-combustion-mass spectrometry 585(1991)177

Trapping system for trace organic volatiles 586(1991)315

Increasing data precision in reversed-phase liquid chromatography leads to new information level 592(1992)13

Analytical study of heavy crude oil fractions by coupling of the transalkylation reaction with supercritical fluid chromatography 593(1992)321

Critical capillary column examination of the relationship between separation number and height equivalent to a theoretical plate and their dependence on temperature 595(1992)209

Van Deemter-type relationship for determining the optimum initial flow-rate and optimum pressure programming rate in temperature/pressure-programmed capillary column gas chromatography utilizing separation numbers 595(1992)221

Anteiso-Alkanes

Novel long-chain anteiso-alkanes and anteiso-alkanoic acids in Antarctic rocks colonized by living and fossil cryptoendolithic microorganisms 598(1992)267

Alkanes, perdeuterated

Isotope effect in gas-liquid chromatography of labelled compounds 588(1991)251

Anteiso-Alkanoic acids

Novel long-chain *anteiso*-alkanes and *anteiso*-alkanoic acids in Antarctic rocks colonized by living and fossil cryptoendolithic microorganisms 598(1992)267

Alkanoic acids

see Carboxylic acids

2-Alkanones

Method for characterization of selectivity in reversed-phase liquid chromatography. V. Calibration of the retention scale for chromatographic systems with low concentrations of organic solvents in the mobile phase 556(1991)145

Alkenes

Photoionization assessment of C₃-C₅ alkadienes and alkenes in urban air 591(1992)358

3-Alkoxy carbonyl-2-pyrazolines

Method for characterization of selectivity in reversed-phase liquid chromatography. V. Calibration of the retention scale for chromatographic systems with low concentrations of organic solvents in the mobile phase 556(1991)145

Alkylamines

Reversed-phase liquid chromatographic behaviour of alkylamines with amperometric detection and its application to trace analysis 594(1992)159

Alkylaromatic hydrocarbons

Hydrogen bonding. XXI. Solvation parameters for alkylaromatic hydrocarbons from gas-liquid chromatographic data 594(1992)229

Alkyl aryl ketones

Retention of basic drugs on porous polymers in high-performance liquid chromatography 600(1992)115

Alkylbenzenes

Chemically bonded phases for the reversed-phase high-performance liquid chromatographic separation of basic substances 552(1991)415

Gas chromatographic-mass spectrometric characterization of volatile organic compounds in Barcelona tap water 562(1991)481

Prediction of retention indexes. IV. Chain branching in alkylbenzene isomers with C₁₀₋₁₃ alkyl chains identified in a scintillator solvent 589(1992)231

Analytical study of heavy crude oil fractions by coupling of the transalkylation reaction with supercritical fluid chromatography 593(1992)321

Alkylbenzene sulphonates

Reversed-phase high-performance liquid chromatography of anionic and ethoxylated non-ionic surfactants and pesticides in liquid pesticide formulations 593(1992)227

Alkylbenzenesulphonates, linear

Reversed-phase high-performance liquid chromatographic determination of linear alkylbenzenesulphonates in river water at ppb levels by precolumn concentration 555(1991)155

2-Alkyl-branched acids

Simultaneous stereoanalysis of 2-alkyl-branched acids, esters and alcohols using a selectivity-adjusted column system in multi-dimensional gas chromatography 587(1991)347

Alkylcyclohexylbenzenes

Gas chromatographic separation of *cis-trans* isomers of alkylcyclohexylbenzenes on a capillary column with a liquid crystalline stationary phase 600(1992)123

Alkyl lysophospholipids

Evaluation of different packings for high-performance liquid chromatographic analysis of alkyl lysophospholipids 553(1991)285

3-*n*-Alkyl-6-methyluracils

Method for characterization of selectivity in reversed-phase liquid chromatography. V. Calibration of the retention scale for chromatographic systems with low concentrations of organic solvents in the mobile phase 556(1991)145

Alkylloxazoles

Retentions of alkylloxazoles and alkylthiazoles and their prediction based on non-linear additivity concepts in gas chromatography 552(1991)1

Alkylphenols

Sensitive liquid chromatographic determination of alkyl-, nitro- and chlorophenols by precolumn derivatization with dansyl chloride, postcolumn photolysis and peroxyoxalate chemiluminescence detection 553(1991)345

Alkylsulphates

Quantitative aspects of indirect UV detection in capillary zone electrophoresis 588(1991)321

Alkylthiazoles

Retentions of alkylloxazoles and alkylthiazoles and their prediction based on non-linear additivity concepts in gas chromatography 552(1991)1

Alkynes

Capillary gas chromatography of C₅-C₁₃ branched alkynes on squalane and liquid crystal stationary phases 557(1991)241

Allantoin

Simultaneous determination of allantoin and oxypurines in biological fluids by high-performance liquid chromatography 575(1992)153

Allergens

Isolation of *Der pI*, the *Dermatophagoides pteronyssinus* major mite allergen, from a crude mite culture extract, purification by ion-chromatography, and comparison between the material obtained and a cDNA-coded *Der pI* 599(1992)105

Alprazolam

New high-performance liquid chromatographic method for the determination of alprazolam and its metabolites in serum: instability of 4-hydroxyalprazolam 568(1991)253

High-performance liquid chromatographic method for the determination of alprazolam in plasma using the column-switching technique 578(1992)259

Aluminium complexes

Retention behavior of salicylidene-glycinatoaluminium(III) in reversed-phase high-performance liquid chromatography 586(1991)356

Aluminum-binding proteins

Resolution of serum aluminum-binding proteins by size-exclusion chromatography: identification of a new carrier of aluminum in human serum 576(1992)271

Amanitins

Improved high-performance liquid chromatographic determination with amperometric detection of α -amanitin in human plasma based on its voltammetric study 563(1991)299

Methods for chromatographic determination of amanitins and related toxins in biological samples (Review) 580(1992)279

Simultaneous assay for amatoxins and phallotoxins in *Amanita phalloides* Fr. by high-performance liquid chromatography 598(1992)227

Amatoxins

Simultaneous assay for amatoxins and phallotoxins in *Amanita phalloides* Fr. by high-performance liquid chromatography 598(1992)227

Ambroxol

High-performance liquid chromatography post-column derivatization with fluorescence detection to study the influence of ambroxol on dipalmitoylphosphatidylcholine levels in rabbit eustachian tube washings 553(1991)1

High-performance liquid chromatographic determination of ambroxol in human plasma 581(1992)251

Amidepin

Determination of amidepin in human plasma by reversed-phase high-performance liquid chromatography with fluorescence detection 565(1991)477

Amides

Correlation between column surface area and retention of polar solutes in packed-column supercritical fluid chromatography 594(1992)291

Amikacin

High-performance liquid chromatographic assays for the quantification of amikacin in human plasma and urine 574(1992)93

High-performance liquid chromatography method for the determination of aminoglycosides based on automated pre-column derivatization with *o*-phthalaldehyde 593(1992)69

Amiloride

High-performance liquid chromatographic assay for amiloride in plasma and urine 567(1991)451

Simultaneous determination of furosemide and amiloride in plasma using high-performance liquid chromatography with fluorescence detection 575(1992)269

Determination and identification of amiloride in human urine by high-performance liquid chromatography and gas chromatography-mass spectrometry 582(1992)93

High-performance liquid chromatographic method for quantitating plasma levels of amiloride and its analogues 582(1992)217

Amine drugs

Selective system of identification and determination of antidepressants and neuroleptics in serum or plasma by solid-phase extraction followed by high-performance liquid chromatography with photodiode-array detection in analytical toxicology 581(1992)75

Amines

see also Catecholamines

Capillary supercritical fluid chromatography of primary aliphatic amines using carbon dioxide and nitrous oxide as mobile phases 558(1991)458

Biogenic amines: their occurrence, biosynthesis and metabolism in the locust, *Schistocerca gregaria*, by gas chromatography-negative-ion chemical ionisation mass spectrometry 562(1991)585

Liquid membrane work-up of blood plasma samples applied to gas chromatographic determination of aliphatic amines 573(1992)191

Prediction of retention indexes. III. Silylated derivatives of polar compounds 586(1991)113

Improvement of extraction procedure for biogenic amines in foods and their high-performance liquid chromatographic determination 591(1992)175

Retention reproducibility of basic drugs in high-performance liquid chromatography on a silica column with a methanol-high-pH buffer eluent. Changes in selectivity with the age of the stationary phase 592(1992)85

Gradient elution with shorter equilibration times in reversed-phase ion-pair chromatography 592(1992)143

Investigations of the influence of silanol groups on the separation of enantiomers by liquid and supercritical fluid chromatography 592(1992)309

By-products in the derivatization of amines with the chiral reagent 2,3,4,6-tetra-O-acetyl- β -D-glucoopyranosyl isothiocyanate and their elimination 592(1992)323

Correlation between column surface area and retention of polar solutes in packed-column supercritical fluid chromatography 594(1992)291

Primary and secondary amine derivatization with luminarins 1 and 2: separation by liquid chromatography with peroxyoxalate chemiluminescence detection 594(1992)386

Screening for amines by dansylation and automated high-performance liquid chromatography 598(1992)51

Control of the retention selectivity of rare earth octaethylporphyrins in reversed-phase high-performance liquid chromatography using amines as mobile phase additives 598(1992)73

***p*-Aminoacetophenone**

Identification of aromatic moieties and mycosamine in antifungal heptaenes with high-performance liquid chromatography, high-performance liquid chromatography-mass spectrometry and gas chromatography-mass spectrometry 585(1991)247

L-Amino acid decarboxylase A, aromatic

High-performance liquid chromatographic procedure for the simultaneous determination of aromatic L-amino acid decarboxylase activity towards 3,4-dihydroxyphenylalanine and 5-hydroxytryptophan 567(1991)381

D-Amino acid oxidase

High-performance liquid chromatographic determination of D-amino acid oxidase activity 566(1991)377

Amino acids

Investigation of cross-linking chiral stationary phases within capillary columns 552(1991)153

Rapid derivatization and gas chromatographic determination of amino acids 552(1991)289

High-performance liquid chromatography of amino acids and dipeptides on new ion exchangers of the HEMA series 552(1991)439

Standardization of cation-exchange clean-up prior to gas chromatography of amino acids 553(1991)87

Advantages and limitations of pre-column derivatization of amino acids with dabsyl chloride 553(1991)123

Rapid and sensitive high-performance liquid chromatographic method for the analysis of tryptophan, tyrosine and phenylalanine in biological samples 553(1991)149

High-performance liquid chromatographic separation of DL-amino acids derivatized with chiral variants of Sanger's reagent 555(1991)81

Application of ligand-exchange chromatography to the assay of L-alanine from DL-aspartic acid by *Pseudomonas dacunhae* 555(1991)272

Chiral separations by micellar electrokinetic chromatography with sodium N-dodecanoyl-L-valinate 559(1991)209

Automated screening of urine samples for carbohydrates, organic and amino acids after treatment with urease 562(1991)125

Measurement of leucine and α -ketoisocaproic acid fluxes in the fetal/placental unit 562(1991)169

Quantitation of tryptophan and other plasma amino acids by automated pre-column *o*-phthalaldehyde derivatization high-performance liquid chromatography: improved sample preparation 564(1991)81

Simultaneous determination of urinary creatinine and aromatic amino acids by cation-exchange chromatography with ultraviolet detection 566(1991)19

Automated determination of neuroactive acidic sulphur-containing amino acids and γ -glutamyl peptides using liquid chromatography with fluorescence and electrochemical detection 566(1991)39

Determination of proteolytic hydrolysis of thyroglobulin 566(1991)471

Separation of *o*-phthalaldehyde-mercaptoethanol derivatives of amino acids from blood plasma on reversed-phase Nova-Pak C₁₈ cartridges 568(1991)207

Liquid chromatographic-mass spectrometric analysis for screening of patients with cystinuria, and identification of cystine stone 568(1991)445

Densitometry for analysis of protein and peptide hydrolysates: application to luteinizing hormone releasing hormone 568(1991)451

Capillary zone electrophoresis: its applicability and potential in biochemical analysis (Review) 569(1991)63

Application of amino acid analysis by high-performance liquid chromatography with phenyl isothiocyanate derivatization to the rapid determination of free amino acids in biological samples 570(1991)285

- Reversed-phase chromatography of phenylthiocarbamyl amino acid derivatives of physiological amino acids: an evaluation and a comparison with analysis by ion-exchange chromatography 574(1992)23
- High-performance liquid chromatographic determination of 3-hydroxypyridinium derivatives as new markers of bone resorption 574(1992)255
- Determination of D- and L-amino acids in mouse kidney by high-performance liquid chromatography 575(1992)147
- Determination of free N-acetylamino acids in biological samples and N-terminal acetylamino acids of proteins 576(1992)63
- Measurement of free amino acid levels in ultrafiltrates of blood plasma by high-performance liquid chromatography with automatic pre-column derivatization 579(1992)215
- Determination of free amino acid enantiomers in rat brain and serum by high-performance liquid chromatography after derivatization with N-*tert.*-butyloxycarbonyl-L1-cysteine and *o*-phthalaldehyde 582(1992)31
- Determination of creatinine and ultraviolet-absorbing amino acids and organic acids in urine by reversed-phase high-performance liquid chromatography 583(1992)1
- Preparative high-yield electroelution of proteins after separation by sodium dodecyl sulphate-polyacrylamide gel electrophoresis and its application to analysis of amino acid sequences and to raise antibodies 585(1991)153
- Prediction of retention indexes. III. Silylated derivatives of polar compounds 586(1991)113
- Multiparameter optimizations in micellar liquid chromatography using the iterative regression optimization strategy 586(1991)221
- Chromatographic separation of mixed peptides from amino acids in biological digests with volatile buffers 586(1991)259
- Application of reversed-phase medium-pressure liquid chromatography to the isolation, separation and amino acid analysis of two closely related peptide toxins of the cyanobacterium *Microcystis aeruginosa* strain PCC 7806 587(1991)71
- Applications of automated amino acid analysis using 9-fluorenylmethyl chloroformate 588(1991)107
- Determination of organic acids, amino acids and saccharides by high-performance liquid chromatography and a postcolumn enzyme reactor with amperometric detection 591(1992)165
- Direct separation of enantiomers by high-performance liquid chromatography on a new chiral ligand-exchange phase 592(1992)291
- High-performance ligand-exchange chromatography of some amino acids containing two chiral centres 592(1992)345
- Rapid determination of amino acids by high-performance liquid chromatography: release of amino acids by perfused rat liver 593(1992)99
- High-performance liquid chromatographic method with fluorescence detection for the determination of total homocyst(e)ine in plasma 593(1992)171
- Capillary electrophoresis with electrochemical detection employing an on-column Nafion joint 593(1992)305
- Chromatography of 99 amino acids and other ninhydrin-reactive compounds in the Pickering lithium gradient system 594(1992)165
- Determination of glutamine, glutamic acid and pyroglutamic acids using high-performance liquid chromatography on dynamically modified silica 594(1992)173
- Role of metal ions in the ligand-exchange separation of amino acids 596(1992)281
- Assay of protein drug substances present in solution mixtures by fluorescamine derivatization and capillary electrophoresis 598(1992)123
- Reductive electrochemical detection in liquid chromatography with a zinc amalgam scrubber column 598(1992)195
- Application of capillary reversed-phase high-performance liquid chromatography to high-sensitivity protein sequence analysis 599(1992)119
- High-performance liquid chromatography of amino acids, peptides and proteins. CXX. Evaluation of bandwidth behaviour of proteins chromatographed on tentacle-type anion exchangers 599(1992)163
- Amino acids, N-acetylated**
Separation and indirect detection of amino acids as acetylated derivatives 557(1991)315
- Amino acids, N-[2,2-bis(ethoxycarbonyl)vinyl] derivatives**
Amino acid analysis by high-performance liquid chromatography after derivatization with diethyl ethoxymethylenemalonate 591(1992)181
- Amino acids, 2-butyl TFA esters**
Gas chromatographic separation of diastereomeric amino acid derivatives on chiral stationary phases. Application to the determination of enantiomeric composition in (*S*)-(+)-2-butanol 589(1992)249
- Amino acids, 1-cyano-2-substituted-benz[*f*]isoindole derivatives**
Separation of naphthalene-2,3-dicarboxaldehyde-labeled amino acids by high-performance capillary electrophoresis with laser-induced fluorescence detection 593(1992)265

Amino acids, Dns derivatives

Retention behavior and chiral recognition of β -cyclodextrin-derivative polymer adsorbed on silica for warfarin, structurally related compounds and Dns-amino acids 555(1991)53

Separation of Dns-amino acids and vitamins by micellar electrokinetic chromatography 559(1991)537

Amino acids, phenylthiocarbamyl derivatives

Measurement of amino acid compositions of glycoprotein systems by gas-phase hydrolysis and reversed-phase high-performance liquid chromatography 599(1992)3

Buffer and pH dependence of the retention of phenylthiocarbamylamino acids in reversed-phase high-performance liquid chromatography 600(1992)87

Amino alcohols

High-performance liquid chromatographic detection of enantiomeric amino alcohols after derivatization with *o*-phthalaldehyde and various thiosugars 558(1991)311

Prediction of retention indexes. III. Silylated derivatives of polar compounds 586(1991)113

Reversed retention order and other stereoselective effects in the separation of amino alcohols on Chiralcel OD 592(1992)331

4-Aminoandrost-1,4,6-triene-3,17-dione

High-performance liquid chromatographic analysis of FCE 24304 (6-methylenandrost-1,4-diene-3,17-dione) and FCE 24928 (4-aminoandrost-1,4,6-triene-3,17-dione), two new aromatase inhibitors 593(1991)25

***p*-Aminobenzamide**

Study of affinity supports based on reactive polymers immobilized on silica: affinity constant determination from isocratic zonal elution 584(1992)3

2-Amino-5-bromo-2'-fluorobenzophenone

Separation and determination of aminohalogenbenzophenones by high-performance liquid chromatography with electrochemical detection 578(1992)146

 γ -Aminobutyric acid

γ -Aminobutyric acid, catecholamine and indoleamine determinations from the same brain region by high-performance liquid chromatography with electrochemical detection 579(1992)334

Reductive electrochemical detection in liquid chromatography with a zinc amalgam scrubber column 598(1992)195

Aminocaproic acid

Efficient substitution of 1,1'-carbonyldiimidazole activated cellulose and Sepharose matrices with amino acyl spacer arms 587(1991)271

2-Amino-5-chlorobenzophenone

Separation and determination of aminohalogenbenzophenones by high-performance liquid chromatography with electrochemical detection 578(1992)146

4-Amino-7-chloroquinoline

Simultaneous determination of chloroquine and its three metabolites in human plasma, whole blood and urine by ion-pair high-performance liquid chromatography 574(1992)305

1-Aminocyclopropanecarboxylic acid

High-performance liquid chromatographic assay for 1-aminocyclopropanecarboxylic acid from plasma and brain 578(1992)103

1-Amino-1-deoxy-2-ketoses

Gas chromatographic determination of *N*-carboxymethyl amino acids, the periodate oxidation products of Amadori compounds 552(1991)345

2-Amino-2',5-dichlorobenzophenone

Separation and determination of aminohalogenbenzophenones by high-performance liquid chromatography with electrochemical detection 578(1992)146

2-Amino-1-(2,4-dichlorophenyl)ethanol

(2*R*,3*R*)-Dicyclohexyl tartrate as a chiral mobile phase additive 585(1991)35

Aminoglycosides

Determination of aminoglycoside antibiotics by reversed-phase ion-pair high-performance liquid chromatography coupled with pulsed amperometry and ion spray mass spectrometry 591(1992)195

High-performance liquid chromatography method for the determination of aminoglycosides based on automated pre-column derivatization with *o*-phthalaldehyde 593(1992)69

Aminohalogenbenzophenones

Separation and determination of aminohalogenbenzophenones by high-performance liquid chromatography with electrochemical detection 578(1992)146

4-Amino-1-hydroxybutane-1,1-bisphosphonic acid

Improved determination of the bisphosphonate alendronate in human plasma and urine by automated precolumn derivatization and high-performance liquid chromatography with fluorescence and electrochemical detection 583(1992)183

 δ -Aminolevulinic acid synthase

Fluorimetric determination of hepatic δ -aminolevulinic acid synthase activity by high-performance liquid chromatography 567(1991)65

(Aminomethyl)phosphonic acid

High-performance liquid chromatographic determination of glyphosate and (aminomethyl)phosphonic acid in human serum after conversion into *p*-toluenesulfonyl derivatives 566(1991)239

Determination of glyphosate and its metabolite, (aminomethyl) phosphonic acid, in serum using capillary electrophoresis 571(1991)324

2-Amino-2-oxazoline

Determination of a new 2-amino-2-oxazoline (COR 3224) in plasma and brain tissue of the rat by high-performance liquid chromatography with electrochemical detection 563(1991)476

Aminopeptidase

Determination of biocatalyst consumption in an aminopeptidase process using automated sample preparation and high-performance liquid chromatography 566(1991)493

Characterization of an aminopeptidase in cerebrospinal fluid. Structure elucidation of enzyme hydrolysis products of synthetic methionine-enkephalin by reversed-phase high-performance liquid chromatography and mass spectrometry 574(1992)189

Aminophenols

Extension of the electrostatic retention model of reversed-phase ion-pair high-performance liquid chromatography to include the effect of the eluent pH 552(1991)13

Correlation between column surface area and retention of polar solutes in packed-column supercritical fluid chromatography 594(1992)291

***p*-Aminophenylarsenoxide**

Human plasma lecithin:cholesterol acyltransferase. Preparation and use of immobilized *p*-aminophenylarsenoxide as a catalytic site-directed covalent ligand in enzyme purification 568(1991)69

***m*-Aminophenylboronic acid**

Coupling of *m*-aminophenylboronic acid to *s*-triazine-activated Sephacryl: use in the affinity chromatography of glycosylated hemoglobins 575(1992)223

α -Aminophosphonic acids

Separation of the enantiomers of the 3,5-dinitrobenzamide derivatives of α -amino phosphonates on four chiral stationary phases 598(1992)159

2-Aminopyridine

Recovery of sugar derivatives from 2-aminopyridine labeling mixtures for high-performance liquid chromatography using UV or fluorescence detection 587(1991)177

4-Aminopyridine

Determination of 4-aminopyridine in serum by solid-phase extraction and high-performance liquid chromatography 574(1992)166

5-Aminosalicylic acid

High-performance liquid chromatographic assay for the determination of 5-aminosalicylic acid and acetyl-5-aminosalicylic acid concentrations in endoscopic intestinal biopsy in humans 564(1991)296

High-performance liquid chromatographic assay of 5-aminosalicylic acid (5-ASA) and its metabolites *N*- β -D-glucopyranosyl-5-ASA, *N*-acetyl-5-ASA, *N*-formyl-5-ASA and *N*-butyryl-5-ASA in biological fluids 570(1991)109

Stability of 5-aminosalicylic acid and its metabolites in plasma at -20°C . Formation of *N*- β -D-glucopyranosyl-5-aminosalicylic acid 570(1991)224

Ion-pairing high-performance liquid chromatographic method for the determination of 5-aminosalicylic acid and related impurities in bulk chemical 588(1991)187

2-Amino-1-[spiro(cyclopropane-1,1'-inden)-3'-yl]ethanol (2*R*,3*R*)-Dicyclohexyl tartrate as a chiral mobile phase additive 585(1991)35

Aminothiazolyl cephalosporins

Microanalytical high-performance liquid chromatographic assay for cefpirome in human milk and urine 574(1992)356

Aminothiols

High-performance liquid chromatography of thiazolidinic compounds obtained by condensation of pyridoxal 5'-phosphate or pyridoxal with aminothiols (*L*- or *D*-cysteine, cysteamine, *L*-cysteine ethyl ester) 586(1991)153

Amiodarone

High-performance liquid chromatographic assay for amiodarone *N*-deethylation in microsomes of rat liver 579(1992)269

Amitriptyline

Column switching and high-performance liquid chromatography in the analysis of amitriptyline, nortriptyline and hydroxylated metabolites in human plasma or serum 578(1992)273

On-line solid-phase extraction and high-performance liquid chromatographic determination of nortriptyline and amitriptyline in serum 579(1992)291

High-performance liquid chromatographic determination of amitriptyline and its main metabolites using a silica column with reversed-phase eluent. Application in mice 584(1992)249

Simultaneous optimization of several chromatographic performance goals using Derringer's desirability function 586(1991)11

Amitriptyline *N*-oxide

High-performance liquid chromatographic determination of amitriptyline and its main metabolites using a silica column with reversed-phase eluent. Application in mice 584(1992)249

Ammonia

Reaction-gas chromatography for physico-chemical investigations on aggressive gases 552(1991)319

Ammonium ion

UV spectrophotometric detection in the ion chromatographic analysis of alkali and alkaline earth metal and ammonium cations 591(1992)219

Ammonium nitrogen compounds

Nitrogen-specific liquid chromatography detector based on chemiluminescence. Application to the analysis of ammonium nitrogen in waste water 592(1992)209

Ammonium 7-N-piperazino-2,1,3-benzoxadiazole-4-sulphonate
Precolumn fluorescence tagging reagent for carboxylic acids in high-performance liquid chromatography: 4-substituted-7-aminoalkylamino-2,1,3-benzoxadiazoles 588(1991)61

Ammonium salts

Selection of buffers and of an ion-pairing agent for thermospray liquid chromatographic-mass spectrometric analysis of ionic compounds 552(1991)605

Amoxapine

Simultaneous quantitation of loxapine, amoxapine and their 7- and 8-hydroxy metabolites in plasma by high-performance liquid chromatography 564(1991)213

Amoxicillin

Reversed-phase high-performance liquid chromatography of amphoteric β -lactam antibiotics: effects of columns, ion-pairing reagents and mobile phase pH on their retention times 564(1991)195

Development of analytical methods for some penicillins in bovine milk by ion-paired chromatography and confirmation by thermospray mass spectrometry 567(1991)389

Separation of antibiotics by high-performance capillary electrophoresis with photodiode-array detection 585(1991)133

Amphetamines

Electron-capture gas chromatographic procedure for simultaneous determination of amphetamine and N-methylamphetamine 573(1992)313

Determination of methamphetamine enantiomer ratios in urine by gas chromatography-mass spectrometry (Review) 580(1992)77

Determination of amphetamines by high-performance liquid chromatography with ultraviolet detection. On-line pre-column derivatization with 9-fluorenylmethyl chloroformate and preconcentration 593(1992)9

Amphotericin B

High-performance liquid chromatographic analysis of amphotericin B in plasma, blood, urine and tissues for pharmacokinetic and tissue distribution studies 579(1992)259

High-performance liquid chromatography of heptaene polyenes: assay of heptaene produced by *Streptomyces griseoviridis* 588(1991)356

Ampicillin

Reversed-phase high-performance liquid chromatography of amphoteric β -lactam antibiotics: effects of columns, ion-pairing reagents and mobile phase pH on their retention times 564(1991)195

Determination of ampicillin in biological fluids by coupled-column liquid chromatography and post-column derivatization 567(1991)121

Development of analytical methods for some penicillins in bovine milk by ion-paired chromatography and confirmation by thermospray mass spectrometry 567(1991)389

Separation of antibiotics by high-performance capillary electrophoresis with photodiode-array detection 585(1991)133

Amylase

Electrophoretic analysis of enzyme-linked immunoglobulins and their clinical significance (Review) 569(1991)347

Separation and characterization of sialic acid-containing salivary-type amylase from patients' sera with immunoglobulin A-type myeloma 571(1991)61

Amylocaine

Determination of limiting ionic mobilities and dissociation constants of some local anaesthetics 596(1992)265

Amyloid

Rapid method to isolate serum amyloid P component from human plasma. Characterization of the isolated protein 578(1992)130

Amyloid protein A

Isolation and purification of amyloid protein A by sodium dodecyl sulphate polyacrylamide gel electrophoresis and reversed-phase high-performance liquid chromatography 573(1992)17

Amylopectins

Estimation of the distributions of chain length of amylopectins by high-performance liquid chromatography with pulsed amperometric detection 585(1991)233

Anabolic agents

see also Special issue index

Anabolic agents

Studies on anabolic steroids. V. Sequential reduction of methandienone and structurally related steroid A-ring substituents in humans: gas chromatographic-mass spectrometric study of the corresponding urinary metabolites 562(1991)323

Derivatization and gas chromatographic-mass spectrometric detection of anabolic steroid residues isolated from edible muscle tissues 562(1991)673

Review. Immunoaffinity chromatography, its applicability and limitations in multi-residue analysis of anabolizing and doping agents 564(1991)363

High-performance liquid chromatographic separation and detection methods for anabolic compounds (Review) 580(1992)111

General and selective isolation procedure for high-performance liquid chromatographic determination of anabolic steroids in tissues 588(1991)89

Analgesics

Comparison of commercially available thin-layer chromatography plates with mixtures of dyes, analgesics and phenols 595(1992)368

Androgens

see also Steroids

Direct quantitative digital autoradiography–thin-layer chromatography of $3\alpha,3\beta$ - and 5α -reduced and 17β -dehydrogenated androgens derived from testosterone metabolism 571(1991)73

Androstadienone

Simultaneous determination of testosterone and androstadienone (sex attractant) in human plasma by gas chromatography–mass spectrometry with high-resolution selected-ion monitoring 565(1991)35

16-Androstenedione

Applications of gas chromatography–mass spectrometry in the study of androgen and odorous 16-androstene metabolism by human axillary bacteria 562(1991)647

Androsterone

On-line derivatization of eluted substances in dynamic high-performance liquid chromatography–mass spectrometry through the particle-beam interface 554(1991)125

Anethole

Comparison of four liquid crystal stationary phases used above and below their melting point temperatures for the gas chromatography of some volatile oil constituents 588(1991)231

Angiotensin III

Analyte focusing in capillary electrophoresis using on-line isotachopheresis 591(1992)341

Angiotensin-converting enzyme

Preparative isolation of angiotensin-converting enzyme from human lung 579(1992)63

Angiotensin-converting enzyme inhibitor

Determination in plasma of angiotensin-converting enzyme inhibitor by inhibitor-binding assay 568(1991)309

Angiotensin II receptor antagonist

High-performance liquid chromatographic determination of angiotensin II receptor antagonists in human plasma and urine. I. DuP 532 (L-694,492) 581(1992)165

Anilines

Chemically bonded phases for the reversed-phase high-performance liquid chromatographic separation of basic substances 552(1991)415

Chromatographic effects of residual amino groups on liquid chromatographic stationary phases derived from aminopropyl-silica gel 590(1992)189

Computer simulation for the prediction of separation as a function of pH for reversed-phase high-performance liquid chromatography. I. Accuracy of a theory-based model 592(1992)183

Computer simulation for the prediction of separation as a function of pH for reversed-phase high-performance liquid chromatography. II. Resolution as a function of simultaneous change in pH and solvent strength 592(1992)197

High-performance liquid chromatographic retention behaviour of ring-substituted aniline derivatives on a porous graphitized carbon column 600(1992)43

Thermal desorption–gas chromatography for the determination of benzene, aniline, nitrobenzene and chlorobenzene in workplace air 600(1992)344

Anions

Dilution effect of carbonate ion on isotachophoretic zones observed with scanning UV photometric detection 555(1991)247

Electroosmotically driven electrochromatography of anions having similar electrophoretic mobilities by ion pairing 557(1991)125

Ion chromatographic determination of anions, especially sulphur-containing anions, with conductimetric and kinetic detection 558(1991)209

Determination of major ions in snow and ice cores by ion chromatography 594(1992)225

Relationship between zone length and step height in isotachopheresis 595(1992)327

Anions, inorganic

see Inorganic anions

Anserine

Determination of carnosine and other biogenic imidazoles in equine plasma by isocratic reversed-phase ion-pair high-performance liquid chromatography 579(1992)45

Anthocyanins

Structure elucidation of saccharides in anthocyanins and flavonols by means of methylation analysis and gas chromatography 598(1992)81

Anthracene

Measurements using gas chromatography with coelution and dual-isotope atomic emission detection 586(1991)309

Quantitative measurements via co-elution and dual-isotope detection by gas chromatography-mass spectrometry 587(1991)255

Retention behavior of β -cyclodextrin complexes of anthracene and pyrene using reversed-phase liquid chromatography. The effects of *tert.*-butanol and cyclopentanol 594(1992)37

Anthracyclines

Simple and sensitive quantification of anthracyclines in mouse atrial tissue using high-performance liquid chromatography and fluorescence detection 570(1991)209

Stability studies with a high-performance liquid chromatographic method for the determination of a new anthracycline analogue, 3'-deamino-3'-[2-(*S*)-methoxy-4-morpholino]doxorubicin (FCE 23762), in the final drug formulation 593(1992)47

Anthraquinone

Quantitative measurements via co-elution and dual-isotope detection by gas chromatography-mass spectrometry 587(1991)255

Anthraquinone-1,8-disulphonic acid

Electrodialytic sample treatment coupled on-line with column liquid chromatography for the determination of basic and acidic compounds in environmental samples 600(1992)163

Antibiotics

Analysis of gentamicin sulfate by high-performance liquid chromatography combined with thermospray mass spectrometry 554(1991)191

Determination of sulfonamides by liquid chromatography, ultraviolet diode array detection and ion-spray tandem mass spectrometry with application to cultured salmon flesh 558(1991)155

Isolation of erythromycin A N-oxide and pseudoerythromycin A hemiketal from fermentation broth of *Saccharopolyspora erythraea* by thin-layer and high-performance liquid chromatography 558(1991)265

Equivalence of a high-performance liquid chromatographic assay and a bioassay of azithromycin in human serum samples 576(1992)358

Hydrophobicity of β -lactam antibiotics. Explanation and prediction of their behaviour in various partitioning solvent systems and reversed-phase chromatography 585(1991)3

Separation of antibiotics by high-performance capillary electrophoresis with photodiode-array detection 585(1991)133

Identification of aromatic moieties and mycosamine in antifungal heptaenes with high-performance liquid chromatography, high-performance liquid chromatography-mass spectrometry and gas chromatography-mass spectrometry 585(1991)247

Isolation of doxycycline, 6-epidoxycycline and 2-acetyl-2-decarboxamidometacycline from commercial metacycline by preparative column liquid chromatography on silica gel 586(1991)67

High-performance liquid chromatographic analysis of oxytetracycline in chinook salmon following administration of medicated feed 588(1991)181

High-performance liquid chromatography of heptaene polyenes: assay of heptaene produced by *Streptomyces griseoviridis* 588(1991)356

Preparative high-performance liquid chromatographic separation and isolation of bacitracin components and their relationship to microbiological activity 590(1992)163

Determination of aminoglycoside antibiotics by reversed-phase ion-pair high-performance liquid chromatography coupled with pulsed amperometry and ion spray mass spectrometry 591(1992)195

High-performance liquid chromatographic determination of penicillin G, penicillin V and cloxacillin in beef and pork tissues 593(1992)15

Automated high-performance liquid chromatographic method for the determination of rifampicin in plasma 593(1992)21

Separation of pristinamycins by high-speed counter-current chromatography. I. Selection of solvent system and preliminary preparative studies 593(1992)363

Solvent degradation of cloxacillin *in vitro*. Tentative identification of degradation products using thermospray liquid chromatography-mass spectrometry 594(1992)195

Determination of dirithromycin purity and related substances by high-performance liquid chromatography 594(1992)203

Aqueous high-performance size-exclusion chromatographic assay for high-molecular-weight impurities in ceftiofur sodium 595(1992)185

Determination of nitroxylin in cow milk by reversed-phase high-performance liquid chromatography with dual-electrode coulometric detection 596(1992)67

Determination of tetracyclines in bovine and porcine muscle by high-performance liquid chromatography using solid-phase extraction 596(1992)211

Evaluation of restricted access media for high-performance liquid chromatographic analysis of sulfonamide antibiotic residues in bovine serum 598(1992)23

Determination of erythromycin ethylsuccinate by liquid chromatography 600(1992)99

Collaborative study of the determination of cloxacillin by column liquid chromatography 600(1992)333

Antibodies

see also Monoclonal antibodies

Factors affecting the specific activity of immobilized antibodies and their biologically active fragments 572(1991)85

Preparative high-yield electroelution of proteins after separation by sodium dodecyl sulphate-polyacrylamide gel electrophoresis and its application to analysis of amino acid sequences and to raise antibodies 585(1991)153

Immobilization of Fv antibody fragments on porous silica and their utility in affinity chromatography 587(1991)161

Characterization of synthetic macroporous packing materials in low-pressure cartridges and columns 590(1992)83

Use of antibody fragments in immunoaffinity chromatography. Comparison of FV fragments, VH fragments and paralog peptides 597(1992)239

Affinity purification of antibodies using immobilized FB domain of protein A 597(1992)257

Anticonvulsants

High-performance liquid chromatographic procedure for the determination of tiagabine concentrations in human plasma using electrochemical detection 574(1992)313

Antidepressants

Drug level monitoring: antidepressants (Review) 576(1992)183

Selective system of identification and determination of antidepressants and neuroleptics in serum or plasma by solid-phase extraction followed by high-performance liquid chromatography with photodiode-array detection in analytical toxicology 581(1992)75

Antidextran antibodies

Analysis of the interaction between an $\alpha(1\rightarrow6)$ dextran-specific mouse hybridoma antibody and dextran B512 by affinity electrophoresis 597(1992)345

Antigens

Immobilization of Fv antibody fragments on porous silica and their utility in affinity chromatography 587(1991)161

Antihistamines

Determination of antihistamines in pharmaceuticals by capillary electrophoresis 588(1991)335

Antihistaminics

Determination of diphenylmethane antihistaminic drugs and their analogues in body fluids by gas chromatography with surface ionization detection 581(1992)213

Anti-infective agents

Preparative separation and analysis of the enantiomers of [³H]Abbott-69992, an HIV anti-infective nucleoside, by ligand-exchange high-performance liquid chromatography 586(1991)265

Anti-inflammatory agents

High-performance liquid chromatographic evaluation of Med 15 and its metabolites Med 5 and tolmetin in rat plasma 553(1991)81

Antioxidants

Gas-liquid chromatography in qualitative analysis. XIX. The use of antioxidants to delay the oxidation of polyoxyethylene glycol stationary phases 552(1991)161

Antipyrene

High-performance liquid chromatographic method for the determination of the three main oxidative and 3-carboxylic antipyrene metabolites in human urine 563(1991)103

Separation and identification of the 4-hydroxyantipyrene sulphoconjugate 576(1992)103

Antisense peptides

Interactions and uses of antisense peptides in affinity technology (Review) 597(1992)29

Anti-Tac-H

Membrane-based receptor affinity chromatography 597(1992)155

Antiviral drugs

Pseudo-electrochromatography-mass spectrometry: a new alternative 554(1991)339

Apolactoferrin

High-performance liquid chromatographic separation of human apolactoferrin and monoferric and diferric lactoferrins 579(1992)346

Apolipoprotein B

Compared stability of Sepharose-based immunoabsorbents prepared by various activation methods 584(1992)17

Apolipoproteins

Characterization of plasma apolipoproteins by capillary electrophoresis 583(1992)111

Apomorphine

Possible role for stationary phase metal interactions in the chromatography of hydroxyamines on silica 552(1991)429

Selective and quantitative isolation and determination of apomorphine in human plasma 570(1991)419

Apotransferrin

High-performance liquid chromatographic separation of human apotransferrin and monoferric and diferric transferrins 567(1991)248

Apovincaminic acid

Determination of apovincaminic acid in serum by means of high-performance liquid chromatography 573(1992)323

Aprophen

Isolation and identification of β -hydroxyethylaprophen: a urinary metabolite of aprophen in rats 563(1991)466

Arabinitol

Diagnosis of disseminated candidiasis based on serum D/L-arabinitol ratios using negative chemical ionization mass spectrometry 575(1992)281

1- β -D-Arabinofuranosyl-*E*-5-(2-bromovinyl)uracil

High-performance liquid chromatographic determination of 1- β -D-arabinofuranosyl-*E*-5-(2-bromovinyl)uracil and its metabolite (*E*)-5-(2-bromovinyl)uracil in serum 568(1991)385

Arabinose

High-performance liquid chromatographic determination of sugars and polyols in extracts of lichens and sugarcane juice 553(1991)271

Arabinosylcytosine

High-performance liquid chromatographic study of some biologically active analogues of arabinosylcytosine 588(1991)147

Arabitol

Chromatographic separation of polyols by ligand exchange. Effects of the ion-exchange resin cross-linking and size 594(1992)125

Arachidonic acid

Gas chromatographic-mass spectrometric characterisation of some novel hydroxyeicosatetraenoic acids formed on incubation of arachidonic acid with microsomes from induced rat livers 562(1991)237

Arachidonic acid metabolites

Cyclooxygenase and lipoxygenase arachidonic acid metabolism by monocytes from human immune deficiency virus-infected drug users 557(1991)507

(Arene)tricarbonylchromium ketones

High-performance liquid chromatographic separation of chiral metallocenic ketones and alcohols 600(1992)320

Arginine

Liquid chromatographic-mass spectrometric analysis for screening of patients with cystinuria, and identification of cystine stone 568(1991)445

Arginine-containing peptides

Monitoring 4-methoxy-2,3,6-trimethylbenzenesulphonyl deprotection of arginine-containing synthetic peptides using capillary zone electrophoresis 589(1992)380

Aristolochic acids

Isolation of aristolochic acids HA_sI and HA_sII by preparative liquid chromatography 594(1992)145

Aromatase

High-performance liquid chromatographic analysis of FCE 24304 (6-methylenandrosta-1,4-diene-3,17-dione) and FCE 24928 (4-aminoandrosta-1,4,6-triene-3,17-dione), two new aromatase inhibitors 593(1991)25

Aromatic amines

see also Amines

New chiral covalently bonded, π -donor stationary phases for high-performance liquid chromatography, based on derivatives of optically active 1-(1-naphthyl)ethylamine. Part II 585(1991)195

Aromatic amines

see also Amines

Quantitation of mutagenic/carcinogenic heterocyclic aromatic amines in food products 592(1992)271

Rapid high-performance liquid chromatographic determination of urinary N-(1-methylethyl)-N'-phenyl-1,4-benzenediamine in workers exposed to aromatic amines 592(1992)279

Comparison of adsorption properties of Florisil and silica in high-performance liquid chromatography. I. Retention behaviour of monofunctional model solutes 600(1992)51

Aromatic compounds

Preparation and capillary column gas chromatographic characterization of aza-crown ether polysiloxane stationary phases 594(1992)243

Aromatic hydrocarbons

see also Hydrocarbons

Gas chromatographic investigation of silica supports modified in a high-frequency low-temperature plasma 552(1991)179

Determination of aromatic hydrocarbons and their metabolites in human blood and urine (Review) 580(1992)229

Use of basic alumina in fractionation of fossil fuels 600(1992)243

Arsenobetaine

Electrospray interfacing for the coupling of ion-exchange and ion-pairing chromatography to mass spectrometry 554(1991)27

Arsenocholine

Electrospray interfacing for the coupling of ion-exchange and ion-pairing chromatography to mass spectrometry 554(1991)27

Artemether

Selective determination, in plasma, of artemether and its major metabolite, dihydroartemisinin, by high-performance liquid chromatography with ultraviolet detection 583(1992)131

Artemisin

Use of high-performance liquid chromatographic peak deconvolution and peak labelling to identify antiparasitic components in plant extracts 593(1992)209

Arylalkanoic acids

Influence of mobile phase composition on evaluation of lipophilicity by partition chromatography 557(1991)399

Ascorbic acid

see Vitamin C

Asiaticoside

Direct coupling of high-speed counter-current chromatography to thin-layer chromatography. Application to the separation of asiaticoside and madecassoside from *Centella asiatica* 558(1991)446

Aspartame

Selection of buffers and of an ion-pairing agent for thermospray liquid chromatographic-mass spectrometric analysis of ionic compounds 552(1991)605

Aspartate

Measurement of glutamate, aspartate and glycine and its potential precursors in human brain using high-performance liquid chromatography by pre-column derivatization with diethylaminoazobenzene sulphonyl chloride 583(1992)241

Aspartate aminotransferase

Electrophoresis of serum isoenzymes and proteins following acute myocardial infarction (Review) 569(1991)323

Electrophoretic analysis of enzyme-linked immunoglobulins and their clinical significance (Review) 569(1991)347

Analysis of alanine and aspartate aminotransferase isoforms in mustard (*Sinapis alba* L.) cotyledons 587(1991)101

Aspartic acid

Application of ligand-exchange chromatography to the assay of L-alanine from DL-aspartic acid by *Pseudomonas dacunhae* 555(1991)272

Aspartic proteinase

High-performance hydrophobic interaction chromatography of proteins on reversed-phase supports coated with non-ionic surfactants of polyoxyethylene type. Purification of a fungal aspartic proteinase 598(1992)181

Asphaltenes

Detection and determination of heavy crude oil fractions by coupling of the transalkylation reaction with supercritical fluid chromatography 593(1992)321

Astemizole

Detection and determination of free and plasma protein-bound astemizole by thin-layer chromatography: a useful technique for bioavailability studies 567(1991)498

Atenolol

Reversed-phase high-performance liquid chromatographic analysis of atenolol enantiomers in plasma after chiral derivatization with (+)-1-(9-fluorenyl)ethyl chloroformate 568(1991)239

Atrazine

Application of high-performance thin-layer chromatography and automated multiple development for the identification and determination of pesticides in water 553(1991)489

Determination of chlorotriazines in aqueous environmental samples at the ng/l level using preconcentration with a cation exchanger and on-line high-performance liquid chromatography 585(1991)67

Atriopeptidase

High-performance liquid chromatographic determination of N-[2(S)-(mercaptomethyl)-3-(2-methylphenyl)-1-oxopropyl]-L-methionine, the active plasma metabolite of a prodrug atriopeptidase inhibitor (SCH 42495), using a thiol selective (Au/Hg) amperometric detector 579(1992)307

Atropine

Determination of atropine in biological specimens by high-performance liquid chromatography 567(1991)141

Retention processes on α_1 -acid glycoprotein-bonded stationary phase 591(1992)55

(2S,3S)-Dicyclohexyl tartrate as mobile phase additive for the determination of the enantiomeric purity of (S)-atropine in tablets 592(1992)339

Aureofungin

Identification of aromatic moieties and mycosamine in antifungal heptaenes with high-performance liquid chromatography, high-performance liquid chromatography-mass spectrometry and gas chromatography-mass spectrometry 585(1991)247

High-performance liquid chromatography of heptaene polyenes: assay of heptaene produced by *Streptomyces griseoviridis* 588(1991)356

Avermectins

Rapid determination of abamectin in lettuce and cucumber by high-performance liquid chromatography 553(1991)299

Azaarenes

Characterization of nitrogen-containing aromatic compounds in soil and sediment by capillary gas chromatography-mass spectrometry after fractionation 558(1991)223

Azacyclonol

Measurement of azacyclonol in urine and serum of humans following terfenadine (Seldane) administration using gas chromatography-mass spectrometry 578(1992)207

Azetidinyl methyl sulphates

Separation and purification of azetidiny methyl sulfates using preparative high-performance liquid chromatography 595(1992)355

3'-Azido-3'-deoxythymidine monophosphate diglyceride

Isolation of experimental anti-AIDS glycerophospholipids by micro-preparative reversed-phase high-performance liquid chromatography 590(1992)153

Azidothymidine

Simultaneous analysis of azidothymidine and its mono-, di- and triphosphate derivatives in biological fluids, tissue and cultured cells by a rapid high-performance liquid chromatographic method 579(1992)107

Azidothymidine phosphates

Simultaneous analysis of azidothymidine and its mono-, di- and triphosphate derivatives in biological fluids, tissue and cultured cells by a rapid high-performance liquid chromatographic method 579(1992)107

Azithromycin

High-performance liquid chromatographic assay with electrochemical detection for azithromycin in serum and tissues 565(1991)321

Equivalence of a high-performance liquid chromatographic assay and a bioassay of azithromycin in human serum samples 576(1992)358

Azodiphenylmethane

Identification of metabolites of 4,4'-diaminodiphenylmethane (methylene dianiline) using liquid chromatographic and mass spectrometric techniques 583(1992)63

Azodrine

Removal of organic pollutants from aqueous solution. V. Comparative study of the extraction, recovery and chromatographic separation of some organic insecticides using unloaded polyurethane foam columns 552(1991)371

Azoxydiphenylmethane

Identification of metabolites of 4,4'-diaminodiphenylmethane (methylene dianiline) using liquid chromatographic and mass spectrometric techniques 583(1992)63

B**B859-35**

Two-dimensional high-performance liquid chromatography at low ng/ml levels of the anti-proliferative agent B859-35 in serum with automated sample clean-up, solid-phase trapping and ultraviolet detection 553(1991)55

Bacitracin

Preparative high-performance liquid chromatographic separation and isolation of bacitracin components and their relationship to microbiological activity 590(1992)163

Silica sorbents with one- and two-site attached bacitracin in affinity chromatography 596(1992)114

Baicalein

Determination of baicalin and baicalein in rat plasma by high-performance liquid chromatography with electrochemical detection 575(1992)131

Baicalin

Determination of baicalin and baicalein in rat plasma by high-performance liquid chromatography with electrochemical detection 575(1992)131

Bambuterol

Dehydrogenase-silica as a stationary phase for the separation of alcohols and ketones 587(1991)268

Barbital

Effect of stationary phase structure on retention and selectivity of restricted-access reversed-phase packing materials 558(1991)19

Barbiturates

Chiral separation by cyclodextrin-modified micellar electrokinetic chromatography 553(1991)503

Chiral separation of barbiturates and hydantoins by reversed-phase high-performance liquid chromatography using a 25 or 50 mm short ODS cartridge column via β -cyclodextrin inclusion complexes 579(1992)253

Study of the retention behaviour of barbiturates by overpressured layer chromatography using silica gel bonded with tricaprilmethylammonium chloride 589(1992)301

Basic compounds

Retention reproducibility of basic drugs in high-performance liquid chromatography on a silica column with a methanol-high-pH buffer eluent. Changes in selectivity with the age of the stationary phase 592(1992)85

Basic drugs

Retention of basic drugs on porous polymers in high-performance liquid chromatography 600(1992)115

Beclomethasone

Simultaneous determination of beclomethasone, beclomethasone monopropionate and beclomethasone dipropionate in biological fluids using a particle beam interface for combining liquid chromatography with negative-ion chemical ionization mass spectrometry 564(1991)43

Bencyclane

Determination of bencyclane in human plasma by means of capillary gas chromatography and nitrogen-phosphorus selective detection 564(1991)283

Benomyl

Rapid on-line precolumn high-performance liquid chromatographic method for the determination of benomyl, carbendazim and aldicarb species in drinking water 555(1991)147

Determination of carbendazim in blueberries by reversed-phase high-performance liquid chromatography 587(1991)321

Benperidol

Determination of benperidol and its reduced metabolite in human plasma by high-performance liquid chromatography and electrochemical detection 565(1991)363

Benzaldehyde

Determination of benzyl alcohol and its metabolite in plasma by reversed-phase high-performance liquid chromatography 568(1991)145

Benzbromarone

Liquid chromatography-mass spectrometry in metabolic research. I. Metabolites of benzbromarone in human plasma and urine 544(1991)267

Benzene

Study of microporous carbons by gas chromatographic determination of heats of physisorption 555(1991)183

Gas chromatographic separation and automatic identification of complex mixtures of organic solvents in industrial wastes 585(1991)273

Activity coefficients in binary liquid mixtures measured by reversed-flow gas chromatography 595(1992)237

Chromatographic determination of the physico-chemical parameters of adsorption on activated carbon fibres 600(1992)67

Thermal desorption-gas chromatography for the determination of benzene, aniline, nitrobenzene and chlorobenzene in workplace air 600(1992)344

Benzene aldehydes

High-performance liquid chromatographic study of the correlation between the physico-chemical parameters of furan and benzene aldehydes and the rate of their metabolism by yeast 594(1992)23

Benzene derivatives

Electrokinetic reversed-phase chromatography with packed capillaries 593(1992)313

Benzenes, halogenated

Relationship between gas chromatographic retention indices and molecular connectivity indices of chlorinated pesticides and structurally related compounds 598(1992)243

Benzocaine

Determination of benzocaine, dextromethorphan and cetylpyridinium ion by high-performance liquid chromatography with UV detection 558(1991)147

Determination of benzyl alcohol and its metabolite in plasma by reversed-phase high-performance liquid chromatography 568(1991)145

Benzodiazepines

New aspects of quantitative structure-retention relationships in chromatography 552(1991)59

Influence of mobile phase composition on evaluation of lipophilicity by partition chromatography 557(1991)399

Determination of small drug molecules by capillary electrophoresis-atmospheric pressure ionization mass spectrometry 559(1991)515

Non-isotopic receptor assay for benzodiazepines using a biotin-labeled ligand and biotin-immobilized microtiter plate 597(1992)443

High-performance liquid chromatographic analysis of benzodiazepines using diode array, electrochemical and thermospray mass spectrometric detection 598(1992)59

High-performance liquid chromatography of benzodiazepines using sorbents with thermally immobilized Carbowax 20M 600(1992)109

Benzodiazepinone

Application of preparative liquid chromatography to the isolation of enantiomers of a benzodiazepinone derivative 590(1992)127

Benzohydroxamine acids

Development of stability indicating assay methods for the determination of hydroxamic acids in topical formulations 594(1992)209

Benzoic acids

Determination of benzyl alcohol and its metabolite in plasma by reversed-phase high-performance liquid chromatography 568(1991)145

Rapid and simple method for the determination of urinary benzoic and phenylacetic acids and their glycine conjugates in ruminants by reversed-phase high-performance liquid chromatography 582(1992)13

High-performance liquid chromatographic column-switching technique for the determination of intermediates of anaerobic degradation of toluene in ground water microcosm 587(1991)185

Chromatographic effects of residual amino groups on liquid chromatographic stationary phases derived from aminopropyl-silica gel 590(1992)189

Computer simulation for the prediction of separation as a function of pH for reversed-phase high-performance liquid chromatography. II. Resolution as a function of simultaneous change in pH and solvent strength 592(1992)197

Simultaneous determination of sorbic acid, dehydroacetic acid and benzoic acid by gas chromatography-mass spectrometry 594(1992)253

Benzoic acids, substituted

Computer simulation for the prediction of separation as a function of pH for reversed-phase high-performance liquid chromatography. I. Accuracy of a theory-based model 592(1992)183

Benzo[ghi]perylene

Reliability assessment of a gas chromatographic method for polycyclic aromatic hydrocarbons in olive oil 555(1991)211

Benzophenone

Calculation of retention and selectivity in reversed-phase liquid chromatography 552(1991)91

Benzo[a]pyrene glucuronide

Detection of benzo[a]pyrene sulfate and glucuronide conjugates in cell culture medium by directly coupled microbore high-performance liquid chromatography-fast atom bombardment mass spectrometry 577(1992)69

Benzo[a]pyrene sulphate

Detection of benzo[a]pyrene sulfate and glucuronide conjugates in cell culture medium by directly coupled microbore high-performance liquid chromatography-fast atom bombardment mass spectrometry 577(1992)69

Benzo[a]pyrene-*r*-7,*t*-8,9,*c*-10-tetrahydrotretol

Superoxide chemical transformation of diolepoxide polyaromatic hydrocarbon DNA adducts. Determination of benzo[a]pyrene-*r*-7,*t*-8,9,*c*-10-tetrahydrotretol by gas chromatography 588(1991)273

Benzoquinone

Hydroquinone oxidation kinetics in adsorptive liquid chromatographic beds 556(1991)383

Photoemissive ionisation source for ion mobility detectors 588(1991)239

Benzothiazines

Direct high-performance liquid chromatographic resolution of a novel benzothiazine Ca^{2+} antagonist and related compounds 593(1992)37

Benzothiophene

Determination of sulfur components in light petroleum streams by high-resolution gas chromatography with chemiluminescence detection 589(1992)271

2,1,3-Benzoxadiazoles, 4,7-substituted

Precolumn fluorescence tagging reagent for carboxylic acids in high-performance liquid chromatography: 4-substituted-7-aminoalkylamino-2,1,3-benzoxadiazoles 588(1991)61

Benzoylacetone

Photoemissive ionisation source for ion mobility detectors 588(1991)239

N-Benzoylalanine

Contribution of ionically immobilized bovine serum albumin to the retention of enantiomers 600(1992)37

Benzoylcegonine

Determination of cocaine, benzoylcegonine and ecgonine methyl ester in plasma by reversed-phase high-performance liquid chromatography 570(1991)412

Rapid assay of cocaine, opiates and metabolites by gas chromatography-mass spectrometry (Review) 580(1992)43

^{32}P -Postlabelling and mass spectrometric methods for analysis of bulky, polyaromatic carcinogen-DNA adducts in humans (Review) 580(1992)293

Preparation of a fluorescent derivative of benzoylcegonine, and preliminary studies of its application to the analysis of urine 584(1992)275

3-Benzoyl-2-naphthaldehyde

Separation of fluorescent oligosaccharide derivatives by microcolumn techniques based on electrophoresis and liquid chromatography 559(1991)223

Benzoylphenylurea pesticides

Application of an internal surface reversed-phase column for the automated determination of flucycloxuron residues 600(1992)211

Benzyl alcohol

Determination of benzyl alcohol and its metabolite in plasma by reversed-phase high-performance liquid chromatography 568(1991)145

7-Benzyl-7-aza-3-thiabicyclo[3,3,1]nonane · HClO₄

High-performance liquid chromatographic determination of BRB-I-28, a novel antiarrhythmic agent, in dog plasma and urine 583(1992)274

1-Benzyl-4-[(5,6-dimethoxy-1-indanon)-2-yl]methylpiperidine

Determination of enantiomers of 1-benzyl-4-[(5,6-dimethoxy-1-indanon)-2-yl]methylpiperidine hydrochloride (E2020), a centrally acting acetylcholine esterase inhibitor, in plasma by liquid chromatography with fluorometric detection 577(1992)95

Bergapten

High-performance liquid chromatographic determination of citropten and bergapten in suction blister fluid after solar product application in humans 563(1991)407

Bestatin-sensitive aminopeptidase

Characterization of an aminopeptidase in cerebrospinal fluid. Structure elucidation of enzyme hydrolysis products of synthetic methionine-enkephalin by reversed-phase high-performance liquid chromatography and mass spectrometry 574(1992)189

Betacyanins

High-performance liquid chromatographic analysis of the pigments of blood-red prickly pear (*Opuntia ficus indica*) 593(1992)177

Betalaines

High-performance liquid chromatographic analysis of the pigments of blood-red prickly pear (*Opuntia ficus indica*) 593(1992)177

Betamethasone

Stereochemical analysis of betamethasone and dexamethasone by derivatization and high-performance liquid chromatography 595(1992)203

Betain

High-performance liquid chromatographic analysis of the pigments of blood-red prickly pear (*Opuntia ficus indica*) 593(1992)177

Biferric transferrin

High-performance liquid chromatographic separation of human apotransferrin and monoferric and diferric transferrins 567(1991)248

Bifonazole

Gas chromatographic-mass spectrometric determination of two new antimycotic agents, 1-[(5-chloro-2-benzofuranyl)(2-chlorophenyl)methyl]-1*H*-imidazole and 1-[(5-bromo-2-benzofuranyl)phenylmethyl]-1*H*-imidazole, in rabbit plasma following topical administration: a preliminary comparison with bifonazole 567(1991)105

Bile acid pyrenacyl esters

High-performance liquid chromatographic separation of bile acid pyrenacyl esters with cyclodextrin-containing mobile phase 565(1991)111

Bile acids

Capillary gas chromatographic analysis of oxo, oxo-hydroxy and unsaturated bile acid glycine conjugates 558(1991)451

Pre-column derivatization of free bile acids for high-performance liquid chromatographic and gas chromatographic-mass spectrometric analysis 562(1991)697

Studies on steroids. CCLIV. Gas chromatographic-mass spectrometric determination of 4- and 6-hydroxylated bile acids in human urine with negative ion chemical ionization detection 574(1992)1

Analysis of conjugated bile acids by packed-column supercritical fluid chromatography 574(1992)197

Determination of conjugated bile acids in human urine by high-performance liquid chromatography with chemiluminescence detection 577(1992)229

Studies on steroids. CCLIII. Capillary gas chromatographic behaviour of diethylhydrogensilyl-diethylsilylene derivatives of stereoisomeric bile acids 585(1991)281

Bile salts

Bile salt surfactants in micellar electrokinetic capillary chromatography. Application to hydrophobic molecule separations 557(1991)113

Bilirubin

Study of the mechanism of the photoisomerization and photooxidation of bilirubin using a model for the phototherapy of hyperbilirubinemia 566(1991)89

Separation of serum bilirubin species by micellar electrokinetic chromatography with direct sample injection 582(1992)77

Bilobetin

Identification of flavonoids from *Ginkgo biloba* L., *Anthemis nobilis* L. and *Equisetum arvense* L. by high-performance liquid chromatography with diode-array UV detection 553(1991)223

1,1'-Bi-2-naphthol

Problems and solutions in chiral thin-layer chromatography: a two-phase "Pirkle" modified amino-bonded plate 553(1991)497

Biotin

Application of high-performance liquid chromatography to the analysis of niacin and biotin in Italian almond cultivars 553(1991)187

Biphenyl

Reversed-phase liquid chromatography coupled on-line with capillary gas chromatography. I. Introduction of large volumes of aqueous mixtures through an on-column interface 552(1991)489

Biphenyls, halogenated

Relationship between gas chromatographic retention indices and molecular connectivity indices of chlorinated pesticides and structurally related compounds 598(1992)243

Bisaramil

Determination of bisaramil and its metabolite in plasma using high-performance liquid chromatography with electrochemical detection 573(1992)339

Bis(2-chloroethylthiol)ethyl ether

Capillary column gas chromatography–ammonia and deuterated ammonia chemical ionization mass spectrometry of sulfur vesicants 600(1992)267

Bisdesethylchloroquine

High-performance liquid chromatographic separation of the enantiomers of hydroxychloroquine and its major metabolites in biological fluids using an α_1 -acid glycoprotein stationary phase 570(1991)119

Determination of hydroxychloroquine and its major metabolites in plasma using sequential achiral–chiral high-performance liquid chromatography 573(1992)253

Simultaneous determination of chloroquine and its three metabolites in human plasma, whole blood and urine by ion-pair high-performance liquid chromatography 574(1992)305

N,O-Bis(diethylhydrogensilyl)trifluoroacetamide

Studies on steroids. CCLIII. Capillary gas chromatographic behaviour of diethylhydrogensilyl–diethylsilylene derivatives of stereoisomeric bile acids 585(1991)281

Bis(di-isobutyl octadecylsiloxy)silicon 2,3-naphthalocyanine

Photosensitizers for tumor therapy: determination of bis(di-isobutyl octadecylsiloxy)silicon 2,3-naphthalocyanine (isoBOSINC) in rat tissue and serum by high-performance liquid chromatography 568(1991)437

Bishomoallylic hydroxy fatty acids, methyl esters

Configuration analysis of unsaturated hydroxy fatty acids 586(1991)255

Bispecific monoclonal antibodies

Selective high-performance liquid chromatographic purification of bispecific monoclonal antibodies 599(1992)13

Bisphenol A diglycidyl ether

Quality control of cured epoxy resins. Determination of residual free monomers (*m*-xylylenediamine and bisphenol A diglycidyl ether) in the finished product 585(1991)75

Bisphosphorates

Determination of a new bisphosphonate, YM175, in plasma, urine and bone by high-performance liquid chromatography with electrochemical detection 584(1992)213

Bispidinol

Determination of bisaramil and its metabolite in plasma using high-performance liquid chromatography with electrochemical detection 573(1992)339

 β -Blockers

Chiral stationary phase designed for β -blockers 557(1991)173

Screening and confirmatory analysis of β -agonists, β -antagonists and their metabolites in horse urine by capillary gas chromatography–mass spectrometry 564(1991)503

Separation of enantiomers using cellulase (CBH I) silica as a chiral stationary phase 586(1991)233

Block polymers

Analysis of block copolymers by high-performance liquid chromatography under critical conditions 593(1992)233

BMY-14802

High-performance liquid chromatographic assay of α -(4-fluorophenyl)-4-(5-fluoro-2-pyrimidinyl)-1-piperazine butanol (BMY 14802), a potential antipsychotic drug, in monkey and rat plasma 567(1991)274

Bombesin

Solvent system for the counter-current purification of hydrophobic bombesin analogues 589(1992)362

High-performance liquid chromatography of amino acids, peptides and proteins. CXV. Thermodynamic behaviour of peptides in reversed-phase chromatography 593(1992)103

Bone proteins

Determination of profiles of non-collagenous proteins from rat bones by sodium dodecyl sulfate high-performance liquid chromatography 589(1992)135

Bone scanning agents

Determination of the mean ionic charge of the components of three ^{99m}Tc bone scanning agents by gel chromatography with two eluents of different electrolyte composition 594(1992)217

Borane cage compounds

Chromatographic behaviour of the *closo*-[B₁₂H₁₂]²⁻ derivatives on hydroxyethylmethacrylate gels 588(1991)201

Bornanes

Application and chiral recognition of heptakis (2,6-di-O-methyl-3-O-trifluoroacetyl)- β -cyclodextrin as a stationary phase for the gas chromatographic separation of enantiomers 589(1992)257

Bradykinin

Effect of the addition of viscous matrices to the mobile phase on chromatographic performance in liquid chromatography–fast atom bombardment mass spectrometry 554(1991)61

Branched-chain keto acids

Reversed-phase high-performance liquid chromatographic analysis of branched-chain keto acid hydrazone derivatives: optimization of techniques and application to branched-chain keto acid balance studies across the forearm 572(1991)11

Bretylum

Simple, rapid and selective method using high-performance liquid chromatography for the determination of bretylum in plasma 575(1992)162

Brifentanil

Determination of [³H]brifentanil, a potent narcotic analgesic, from rat serum by high-performance liquid chromatography with on-line radioactive detection 565(1991)391

Brij 76

Modification of reversed-phase columns with dyed surfactants. Preparation of mechanically resistant efficient immobilized dyes for protein purification 558(1991)43

Brodiprim

Determination of brodimoprim and its hydroxy metabolite in human plasma, blood and urine by high-performance liquid chromatography 577(1992)123

Bromide

Mixed potential response at metallic electrodes in potentiometric detection for ion chromatography 589(1992)201

Gas chromatography of chloride and bromide as phenylboronic acid/mercuric nitrate derivatives with microwave induced plasma atomic emission detection 594(1992)395

Determination of inorganic anions in salt solutions by ion chromatography using C₁₈ reversed-phase columns coated with cetyltrimethylammonium 598(1992)237

1-[(5-Bromo-2-benzofuranyl)phenylmethyl]-1H-imidazole

Gas chromatographic–mass spectrometric determination of two new antimycotic agents, 1-[(5-chloro-2-benzofuranyl)(2-chlorophenyl)methyl]-1H-imidazole and 1-[(5-bromo-2-benzofuranyl)phenylmethyl]-1H-imidazole, in rabbit plasma following topical administration: a preliminary comparison with bifonazole 567(1991)105

(α -Bromoisovaleryl)urea

Enantioselective determination of *R*- and *S*-(α -bromoisovaleryl)urea in plasma using high-performance liquid chromatography after solid-phase extraction 573(1992)259

3-Bromomethyl-7-methoxy-1,4-benzoxazine-2-one

3-Bromomethyl-7-methoxy-1,4-benzoxazin-2-one as a highly sensitive fluorescence derivatization reagent for carboxylic acids in high-performance liquid chromatography 591(1992)159

(*E*)-5-(2-Bromovinyl)uracil

High-performance liquid chromatographic determination of 1- β -D-arabinofuranosyl-*E*-5-(2-bromovinyl)uracil and its metabolite (*E*)-5-(2-bromovinyl)uracil in serum 568(1991)385

Brucine

Separation of *Strychnos nux-vomica* alkaloids by high-performance liquid chromatography 587(1991)318

Buckminsterfullerenes

Liquid chromatographic separation of all-carbon molecules C₆₀ and C₇₀ with multi-legged phenyl group bonded silica phases 594(1992)105

Budesonide

Evaluation of an automated thermospray liquid chromatography–mass spectrometry system for quantitative use in bioanalytical chemistry 554(1991)215

Bufadienolides

Preparation and evaluation of octadecyl- or phenylpropyl-treated porous glass for the high-performance liquid chromatographic analysis of bufadienolides in *Bufois venenum* 552(1991)381

Bufaralol

Chiral stationary phase designed for β -blockers 557(1991)173

Bumetanide

Measurement of bumetanide in plasma and urine by high-performance liquid chromatography and application to bumetanide disposition 570(1991)235

Bunolol

Determination of (–)-bunolol and its metabolite, dihydro-(–)-bunolol, in human aqueous humour by gas chromatography–negative ion chemical ionisation mass spectrometry 571(1991)101

Bupivacaine

Reversed-phase and chiral high-performance liquid chromatographic assay of bupivacaine and its enantiomers in clinical samples after continuous extrapleural infusion 553(1991)383

Determination of local anaesthetics in body fluids by gas chromatography with surface ionization detection 564(1991)278

High-performance liquid chromatographic method for the quantitation of bupivacaine, 2,6-pipecoloxylidide and 4'-hydroxybupivacaine in plasma and urine 577(1992)103

Determination of limiting ionic mobilities and dissociation constants of some local anaesthetics 596(1992)265

Buprenorphine

High-performance liquid chromatography with electrochemical detection of buprenorphine and its major metabolite in urine 564(1991)557

Determination of buprenorphine by high-performance liquid chromatography with fluorescence detection: application to human and rabbit pharmacokinetic studies 570(1991)339

Rapid assay of cocaine, opiates and metabolites by gas chromatography–mass spectrometry (Review) 580(1992)43

³²P-Postlabelling and mass spectrometric methods for analysis of bulky, polyaromatic carcinogen–DNA adducts in humans (Review) 580(1992)293

Buspirone

Sensitive determination of buspirone in serum by solid-phase extraction and two-dimensional high-performance liquid chromatography 566(1991)250

Simultaneous high-performance liquid chromatographic analysis of buspirone and its metabolite 1-(2-pyrimidinyl)-piperazine in plasma using electrochemical detection 575(1992)117

Busulfan

High-performance liquid chromatographic–mass spectrometric assay of busulfan in serum and cerebrospinal fluid 581(1992)143

Butanol

Direct aqueous injection gas chromatography on a potassium fluoride crystal hydrate-containing sorbent. Determination of volatile organic solvents in the fermentation broth of *Clostridium* strains 596(1992)138

2-Butanol

Gas chromatographic separation of diastereomeric amino acid derivatives on chiral stationary phases. Application to the determination of enantiomeric composition in (*S*)-(+)-2-butanol 589(1992)249

Analysis of zapatera olives by gas and high-performance liquid chromatography 594(1992)259

2-Butanone

Estimation of cellulose polarity by gas chromatography 552(1991)365

Butyl acrylate

Continuous beds for standard and micro high-performance liquid chromatography 586(1991)21

S-4-(*tert*-Butylamino-2-hydroxypropoxy)benzimidazol-2-one

Rapid analysis for metabolites of ¹¹C-labelled drugs: fate of [¹¹C]-S-4-(*tert*-butylamino-2-hydroxypropoxy)benzimidazol-2-one in the dog 570(1991)361

4-*tert*-Butyl pyrocatechol

Comparison of colorimetric and chromatographic determination of *tert*-butylpyrocatechol 595(1992)137

***n*-Butylthiol**

Chromatographic trace analysis of some organic compounds in the environment using derivatization–sorption concentration techniques. I. Gas chromatographic analysis of acrylates in air 557(1991)523

Butyric acid

Applications of capillary gas chromatography to the quality control of butter and related products 552(1991)273

Butyrobetaine

Derivatization of isolated endogenous butyrobetaine with 4'-bromophenacyl trifluoromethanesulfonate followed by high-performance liquid chromatography 573(1992)3

N-Butyryl-5-aminosalicylic acid

High-performance liquid chromatographic assay of 5-aminosalicylic acid (5-ASA) and its metabolites N-β-D-glucopyranosyl-5-ASA, N-acetyl-5-ASA, N-formyl-5-ASA and N-butyryl-5-ASA in biological fluids 570(1991)109

C

C₉

Rapid isolation of human complement component C9 to verify the specificity of a haemolytic C9 microassay 591(1992)187

Cabergoline

Determination of cabergoline in plasma and urine by high-performance liquid chromatography with electrochemical detection 574(1992)170

Cadaverine

Improvement of extraction procedure for biogenic amines in foods and their high-performance liquid chromatographic determination 591(1992)175

Cadmium

Separation of metal complexes of ethylenediaminetetraacetic acid in environmental water samples by ion chromatography with UV and potentiometric detection 558(1991)181

Determination of dietary cadmium-induced metallothioneins in rabbit kidneys and cadmium in metallothioneins by anion-exchange high-performance liquid chromatography coupled with graphite furnace atomic absorption spectrometry 572(1991)73

Determination of divalent trace metals in natural waters by preconcentration on N,N,N',N'-tetra(2-aminoethyl)ethylenediamine-silica followed by on-line ion chromatography 600(1992)203

Caffeine

Elimination of caffeine interference in high-performance liquid chromatographic determination of cotinine in human plasma 568(1991)267

High-performance liquid chromatographic separation of caffeine, theophylline, theobromine and paraxanthine in rat brain and serum 570(1991)185

Simultaneous determination of caffeine and its primary demethylated metabolites in human plasma by high-performance liquid chromatography 575(1992)311

Determination of N-acetylator phenotype using caffeine as a probe compound: a comparison of high-performance liquid chromatography and capillary electrophoresis methods 578(1992)283

Simplex optimization of liquid chromatography with the function of mutual information as a criterion 585(1991)187

Determination of substituted purines in body fluids by micellar electrokinetic capillary chromatography with direct sample injection 593(1992)275

Calcitonin gene-related peptide

Identification of α - and β -species of calcitonin gene-related peptide in the rat amygdala after separation with capillary zone electrophoresis 573(1992)219

Calcium

Determination of calcium ions tightly bound to proteins 585(1991)173

Determination of calcium ion in the presence of phosphate anion and collagen by capillary-type, isotachopheresis 586(1991)363

Simultaneous ion chromatographic determination of chloride and calcium contained in Ringer's injection using a chelating agent as eluent 589(1992)358

Calcium antagonists

Direct high-performance liquid chromatographic resolution of a novel benzothiazine Ca^{2+} antagonist and related compounds 593(1992)37

Calcium-binding proteins

Application of high-performance capillary electrophoresis to the analysis of conformation and interaction of metal-binding proteins 559(1991)345

Calcium cyanamide

Liquid chromatographic method for the determination of calcium cyanamide using pre-column derivatization 558(1991)141

Californin

Analytical and quantitative studies of californin and protopin in aerial part extracts of *Eschscholtzia californica* Cham. with high-performance liquid chromatography 587(1991)314

Calmodulin

Application of high-performance capillary electrophoresis to the analysis of conformation and interaction of metal-binding proteins 559(1991)345

Base-atom recognition in protein adsorption to alkyl agaroses 597(1992)93

Calcium-modulated conformational affinity chromatography. Application to the purification of calmodulin and S100 proteins 597(1992)263

Camphorsulphonic acid

Potentiometric detection of carboxylic acids and inorganic anions in ion-exclusion chromatography using camphorsulphonic acid as eluent 589(1992)209

Camptothecin

Simultaneous determination of the camptothecin analogue CPT-11 and its active metabolite SN-38 by high-performance liquid chromatography: application to plasma pharmacokinetic studies in cancer patients 575(1992)275

Selective and sensitive determination of lactone and hydroxy acid forms of camptothecin and two derivatives (CPT-11 and SN-38) by high-performance liquid chromatography with fluorescence detection 588(1991)165

Candididin

Identification of aromatic moieties and mycosamine in antifungal heptaenes with high-performance liquid chromatography, high-performance liquid chromatography-mass spectrometry and gas chromatography-mass spectrometry 585(1991)247

High-performance liquid chromatography of heptaene polyenes: assay of heptaene produced by *Streptomyces griseoviridis* 588(1991)356

Candimycin

Identification of aromatic moieties and mycosamine in antifungal heptaenes with high-performance liquid chromatography, high-performance liquid chromatography-mass spectrometry and gas chromatography-mass spectrometry 585(1991)247

High-performance liquid chromatography of heptaene polyenes: assay of heptaene produced by *Streptomyces griseoviridis* 588(1991)356

Cannabidiol

Urinary metabolites of cannabidiol in dog, rat and man and their identification by gas chromatography-mass spectrometry 562(1991)299

Canthaxanthin

Determination of several retinoids, carotenoids and E vitamers by high-performance liquid chromatography. Application to plasma and tissues of rats fed a diet rich in either β -carotene or canthaxanthin 553(1991)179

ϵ -Caprolactam

Analysis of ϵ -caprolactam and its cyclic oligomers by high-performance liquid chromatography 585(1991)333

Captopril

Stereoisomeric purity determination of captopril by capillary gas chromatography 558(1991)257

Simple high-performance liquid chromatographic method for the determination of captopril in biological fluids 582(1992)258

Carbadox

Column liquid chromatographic determination of carbadox and olaquinox in feeds 558(1991)125

Determination of some drugs by micellar electrokinetic capillary chromatography. The pseudo-effective mobility as parameter for screening 585(1991)123

Carbamates

Evaluation of eluents in thermospray liquid chromatography-mass spectrometry for identification and determination of pesticides in environmental samples 553(1991)311

High-performance liquid chromatographic determination of monohydroxy compounds by a combination of pre-column derivatization and post-column reaction detection 593(1992)29

Carbamazepine

Determination of thiopental in human serum and plasma by high-performance capillary electrophoresis-micellar electrokinetic chromatography 559(1991)505

Fast liquid chromatography for the determination of drugs in plasma and combination with liquid-solid extraction in a fully automated system 573(1992)59

Automated microanalysis of carbamazepine and its epoxide and *trans*-diol metabolites in plasma by column liquid chromatography 573(1992)65

Preparation and characterization of mixed functional phase silica materials using phenyl-, butyl- or octylchlorosilane as a silylating agent 596(1992)151

Carbendazim

Rapid on-line precolumn high-performance liquid chromatographic method for the determination of benomyl, carbendazim and aldicarb species in drinking water 555(1991)147

Determination of carbendazim in blueberries by reversed-phase high-performance liquid chromatography 587(1991)321

Carbohydrate antigens

Detection of fungal carbohydrate antigens by high-performance immunoaffinity chromatography using a protein A column with covalently linked immunoglobulin G 584(1992)69

Carbohydrate-binding peptides

Purification and characterization of carbohydrate-binding peptides from *Lotus tetragonolobus* and *Ulex europaeus* seed lectins using affinity chromatography 597(1992)213

Determination of the carbohydrate-binding site of *Bauhinia purpurea* lectin by affinity chromatography 597(1992)221

Carbohydrate-binding proteins

Affinity purification and affinity characterization of carbohydrate-binding proteins in bovine kidney 597(1992)323

Carbohydrates

see also Sugars and Saccharides

Simultaneous gas-liquid chromatographic determination of sugars and organic acids as trimethylsilyl derivatives in vegetables and strawberries 552(1991)337

High-performance liquid chromatographic determination of sugars and polyols in extracts of lichens and sugarcane juice 553(1991)271

Carbohydrate separation by ligand-exchange liquid chromatography. Correlation between the formation of sugar-cation complexes and the elution order 558(1991)89

Rapid identification of microbial starch degradation products from a complex nutrient medium by a thin-layer chromatographic method 558(1991)328

Characterization of cellulase-based enzyme reactors for the high-performance liquid chromatographic determination of β -D-glucan oligosaccharides 558(1991)343

Automated screening of urine samples for carbohydrates, organic and amino acids after treatment with urease 562(1991)125

Construction of kinetic models to understand metabolism *in vivo* 566(1991)309

Two-dimensional mapping by high-performance liquid chromatography of pyridylamino oligosaccharides from various glycosphingolipids 586(1991)35

Recovery of sugar derivatives from 2-aminopyridine labeling mixtures for high-performance liquid chromatography using UV or fluorescence detection 587(1991)177

High-performance capillary zone electrophoresis of carbohydrates in the presence of alkaline earth metal ions 588(1991)327

Synthesis and characterization of stationary phases on the basis of silicas modified with epoxidized polybutadienes. IV. Chromatographic experiments on a new amino-functionalized stationary phase for high-performance liquid chromatography 588(1991)340

Determination of carboxylic acids, sugars, glycerol and ethanol in wine and grape must by ion-exchange high-performance liquid chromatography with refractive index detection 590(1992)215

Constant-potential amperometric detection of carbohydrates at metal electrodes in high-performance anion-exchange chromatography 592(1992)229

Galactosylation as a tool for the stabilization and immobilization of proteins (Review) 597(1992)19

Analysis of the interaction between an $\alpha(1\rightarrow6)$ dextran-specific mouse hybridoma antibody and dextran B512 by affinity electrophoresis 597(1992)345

Influence of different N- and O-linked carbohydrates on the retention times of synthetic peptides in reversed-phase high-performance liquid chromatography 599(1992)43

Carbon

Liquid chromatographic separation of all-carbon molecules C_{60} and C_{70} with multi-legged phenyl group bonded silica phases 594(1992)105

Carbonate

Carbonate eluents in ion chromatography. Fractions of resin capacity linked to anions present in the eluent 589(1992)197

Carbon dioxide

Dilution effect of carbonate ion on isotachophoretic zones observed with scanning UV photometric detection 555(1991)247

Packed column supercritical fluid chromatography with carbon dioxide-polar modifiers. Influence of carbon dioxide density on retention 557(1991)85

Capillary supercritical fluid chromatography of primary aliphatic amines using carbon dioxide and nitrous oxide as mobile phases 558(1991)458

Carbonic anhydrase

High-performance liquid chromatography of amino acids, peptides and proteins. CXIV. Protein interactions with porous coulombic sorbents: comparison of experimental findings with predictions of several adsorption models 557(1991)335

Application of high-performance capillary electrophoresis to the analysis of conformation and interaction of metal-binding proteins 559(1991)345

Purification of carbonic anhydrase isoenzymes by high-performance affinity chromatography and hydrophobic interaction chromatography 566(1991)351

Rapid ion-exchange chromatography for preparative separation of proteins. IV. Application to bovine carbonic anhydrase III from skeletal muscle 588(1991)139

Carbonyl compounds

see also Ketones

Electrochemical and chromatographic properties of selected hydrazine and hydrazide derivatives of carbonyl compounds 585(1991)225

1,1'-Carbonyldiimidazole

Efficient substitution of 1,1'-carbonyldiimidazole activated cellulose and Sepharose matrices with amino acyl spacer arms 587(1991)271

Carbonyl sulphide

Near-continuous measurement of hydrogen sulfide and carbonyl sulfide by an automatic gas chromatograph 585(1991)353

Identification of products resulting from carbonyl sulphide-induced degradation of diethanolamine 587(1991)237

Carboplatin

Separation of some platinum(II) complexes by ionic strength gradient on a solvent-generated ion-exchange sorbent 586(1991)291

3-(4-Carboxybenzoyl)-2-quinolinecarboxaldehyde

Separation of fluorescent oligosaccharide derivatives by microcolumn techniques based on electrophoresis and liquid chromatography 559(1991)223

1-(4'-Carboxybutyl)-3,7-dimethylxanthine

Determination of pentoxifylline and its metabolites in human plasma by high-performance liquid chromatography with solid-phase extraction 575(1992)101

2-[p-(2-Carboxyethyl)phenylethylamine]-5'-N-ethylcarboxamido-adenosine

Automated-extraction, high-performance liquid chromatographic method and pharmacokinetics in rats of a highly A_2 -selective adenosine agonist, CGS 21680 578(1992)77

Carboxylic acids

see also Fatty acids and Organic acids

Use of an evaporative light scattering detector in reversed-phase high-performance liquid chromatography of oligomeric surfactants 552(1991)593

Use of non-volatile ion-pairing agents for liquid chromatographic-mass spectrometric analyses with a moving-belt interface 554(1991)281

Determination of volatile and non-volatile organic acids in technical sugar solutions by ion-exclusion chromatography 555(1991)65

Influence of counter-ion inclusion complexation on the quality of cyclodextrin-supported separations in isotachopheresis 557(1991)215

Assessment of a thermospray interface for the coupling of high-performance liquid chromatography and Fourier transform infrared spectrometry 588(1991)15

Pre-column fluorescence tagging reagent for carboxylic acids in high-performance liquid chromatography: 4-substituted-7-aminoalkylamino-2,1,3-benzoxadiazoles 588(1991)61

Optimization of an ion-exchange high-performance liquid chromatographic method for the determination of carboxylic acids, sugars, glycerol and ethanol in wines 589(1992)151

Potentiometric detection of carboxylic acids and inorganic anions in ion-exclusion chromatography using camphorsulphonic acid as eluent 589(1992)209

Determination of carboxylic acids, sugars, glycerol and ethanol in wine and grape must by ion-exchange high-performance liquid chromatography with refractive index detection 590(1992)215

3-Bromomethyl-7-methoxy-1,4-benzoxazin-2-one as a highly sensitive fluorescence derivatization reagent for carboxylic acids in high-performance liquid chromatography 591(1992)159

Comparison of several methods for the determination of trace amounts of polar aliphatic monocarboxylic acids by high-performance liquid chromatography 593(1992)243

Suppression of the effect of metal impurities in alkylsilylated silica packing materials 594(1992)87

Ion-exclusion chromatography using mobile phases containing β -cyclodextrin 595(1992)149

Quantitative analysis of total resin acids by high-performance liquid chromatography of their coumarin ester derivatives 595(1992)155

N-Carboxymethyl amino acids

Gas chromatographic determination of N-carboxymethyl amino acids, the periodate oxidation products of Amadori compounds 552(1991)345

S-Carboxymethyl-L-cysteine

Determination of S-carboxymethyl-L-cysteine and some of its metabolites in urine and serum by high-performance liquid chromatography using fluorescent pre-column labelling 571(1991)133

2-(4-Carboxyphenyl)-5,6-dimethylbenzimidazole

Determination of alcohols by high-performance liquid chromatography after pre-column derivatization with 2-(4-carboxyphenyl)-5,6-dimethylbenzimidazole 585(1991)219

Carboxyprimaquine

High-performance liquid chromatographic determination of pamaquine, primaquine and carboxy primaquine in calf plasma using electrochemical detection 579(1992)123

1-(3'-Carboxypropyl)-3,7-dimethylxanthine

Determination of pentoxifylline and its metabolites in human plasma by high-performance liquid chromatography with solid-phase extraction 575(1992)101

Carcinogen-DNA adducts

³²P-Postlabelling and mass spectrometric methods for analysis of bulky, polyaromatic carcinogen-DNA adducts in humans (Review) 580(1992)293

Carcinogens

Cellulose bearing covalently linked copper phthalocyanine trisulphonate as an adsorbent selective for polycyclic compounds and its use in studies of environmental mutagens and carcinogens (Review) 597(1992)37

Cardiotonic agents

Assessment of the biotransformation of the cardiotonic agent piroximone by high-performance liquid chromatography and gas chromatography-mass spectrometry 593(1992)1

Carnitine

Synthesis, characterisation and high-performance liquid chromatography of C₆-C₁₆ dicarboxyl-*mono*-coenzyme A and *mono*-carnitine esters 570(1991)253

L-Carnitine

Rapid determination of amino acids by high-performance liquid chromatography: release of amino acids by perfused rat liver 593(1992)99

Carnitine esters

Quantification of carnitine esters by high-performance liquid chromatography. Effect of feeding medium-chain triglycerides on the plasma carnitine ester profile 584(1992)157

Carnosine

Determination of carnosine and other biogenic imidazoles in equine plasma by isocratic reversed-phase ion-pair high-performance liquid chromatography 579(1992)45

 β -Carotene

Determination of several retinoids, carotenoids and E vitamins by high-performance liquid chromatography. Application to plasma and tissues of rats fed a diet rich in either β -carotene or canthaxanthin 553(1991)179

Simultaneous determination of retinol, α -tocopherol and β -carotene in serum by isocratic high-performance liquid chromatography 572(1991)103

High-performance liquid chromatographic method for routine determination of vitamins A and E and β -carotene in plasma 581(1992)41

Carotenes

Separation of *trans/cis* α - and β -carotenes by supercritical fluid chromatography. I. Effects of temperature, pressure and organic modifiers on the retention of carotenes 557(1991)47

Separation of *trans/cis* α - and β -carotenes by supercritical fluid chromatography. II. Effect of the type of octadecyl-bonded stationary phase on retention and selectivity of carotenes 557(1991)59

Optimization of an isocratic high-performance liquid chromatographic separation of carotenoids 589(1992)171

Carotenoids

Determination of several retinoids, carotenoids and E vitamins by high-performance liquid chromatography. Application to plasma and tissues of rats fed a diet rich in either β -carotene or canthaxanthin 553(1991)179

Determination of carotenoid pigments in several tree leaves by reversed-phase high-performance liquid chromatography 585(1991)168

Determination of chlorophylls and carotenoids by high-performance liquid chromatography during olive lactic fermentation 585(1991)259

Optimization of an isocratic high-performance liquid chromatographic separation of carotenoids 589(1992)171

Evaluation of reversed-phase liquid chromatographic columns for recovery and selectivity of selected carotenoids 595(1992)89

Carvone

Effects of collection solvent parameters and extraction cell geometry on supercritical fluid extraction efficiencies 594(1992)297

Optimization of supercritical fluid extraction of volatile constituents from a model plant matrix 600(1992)175

Caryophyllene

Comparison of four liquid crystal stationary phases used above and below their melting point temperatures for the gas chromatography of some volatile oil constituents 588(1991)231

Optimization of supercritical fluid extraction of volatile constituents from a model plant matrix 600(1992)175

Catalase

Fast and sensitive staining technique for catalase in polyacrylamide gel 594(1992)400

Catechin

High-performance liquid chromatography of the phenolic compounds in the juice of some French cider apple varieties 555(1991)125

Identification of the constituent flavanoid units in sainfoin proanthocyanidins by reversed-phase high-performance liquid chromatography 594(1992)117

Catechol

Determination of benzene metabolites in urine of mice by solid-phase extraction and high-performance liquid chromatography 593(1992)147

Catecholamines

Migration behaviour of catechols and catecholamines in capillary electrophoresis 559(1991)529

Analysis of rat brain microdialysate by gas chromatography-high-resolution selected-ion monitoring mass spectrometry 562(1991)111

Improved measurement of urinary catecholamines by liquid-liquid extraction, derivatization and high-performance liquid chromatography with fluorimetric detection 563(1991)348

Evaluation of a simple plasma catecholamine extraction procedure prior to high-performance liquid chromatography and electrochemical detection 564(1991)55

Determination of heat-exposure effects on the concentration of catecholamines in bovine plasma and milk 566(1991)29

Application of high-performance liquid chromatography to the study of biogenic amine-related enzymes (Review) 566(1991)287

Measurement of plasma catecholamines by high-performance liquid chromatography with electrochemical detection in intensive care patients after dobutamine infusion 567(1991)240

- High-performance liquid chromatographic determination, with ultraviolet detection, of S-adenosyl-L-methionine and S-adenosyl-L-homocysteine in rat tissues and simultaneously of normetanephrine and metanephrine for phenylethanolamine-N-methyltransferase or catechol-O-methyltransferase activities 567(1991)254
- Analysis of plasma catecholamines by high-performance liquid chromatography with fluorescence detection: simple sample preparation for pre-column fluorescence derivatization 567(1991)351
- Simultaneous determination of free catecholamines and epinine and estimation of total epinine and dopamine in plasma and urine by high-performance liquid chromatography with fluorimetric detection 574(1992)109
- Optimization of high-performance liquid chromatographic assay for catecholamines. Determination of optimal mobile phase composition and elimination of species-dependent differences in extraction recovery of 3,4-dihydroxybenzylamine 574(1992)213
- High-performance liquid chromatography with electrochemical detection for the determination of levodopa, catecholamines and their metabolites in rat brain dialysates 575(1992)207
- Preparation of an optimum mobile phase for the simultaneous determination of neurochemicals in mouse brain tissues by high-performance liquid chromatography with electrochemical detection 576(1992)245
- Determination of the catecholamines and serotonin, their precursors tyrosine and tryptophan, and their main metabolites in rat brain using reversed-phase high-performance liquid chromatography with fluorimetric and oxidative amperometric detection in series 578(1992)17
- Sensitivity and specificity of urinary N-acetyldopamine as a marker for neuroblastomas: comparison with traditional urinary catecholamine metabolites 579(1992)184
- Sensitivity and specificity of urinary N-acetyldopamine as a marker for neuroblastomas: comparison with traditional urinary catecholamine metabolites. Reply to Muskiet and Kema 579(1992)187
- γ -Aminobutyric acid, catecholamine and indoleamine determinations from the same brain region by high-performance liquid chromatography with electrochemical detection 579(1992)334
- Simultaneous measurement of serotonin, catecholamines and their metabolites in cat and human plasma by *in vitro* microdialysis-microbore high-performance liquid chromatography with amperometric detection 582(1992)19
- Simultaneous determination of catecholamines and dobutamine in human plasma and urine by high-performance liquid chromatography with fluorimetric detection 583(1992)236
- Capillary tube isotachophoretic separation of catecholamines using cyclodextrin in the leading electrolyte 587(1991)364
- Computer simulation for the prediction of separation as a function of pH for reversed-phase high-performance liquid chromatography. I. Accuracy of a theory-based model 592(1992)183
- Determination of catecholamines in urine by liquid chromatography and electrochemical detection after on-line sample purification on immobilized boronic acid 593(1992)185
- Apparent inter-channel interference in dual-electrode electrochemical detection 596(1992)110
- Catechol-O-methyltransferase**
High-performance liquid chromatographic determination, with ultraviolet detection, of S-adenosyl-L-methionine and S-adenosyl-L-homocysteine in rat tissues and simultaneously of normetanephrine and metanephrine for phenylethanolamine-N-methyltransferase or catechol-O-methyltransferase activities 567(1991)254
- Application of an organic solvent extraction to the determination of catechol-O-methyltransferase activity by high-performance liquid chromatography in human mononuclear cells 578(1992)175
- Catechols**
Migration behaviour of catechols and catecholamines in capillary electrophoresis 559(1991)529
- Catharanthine**
Determination of *Catharanthus roseus* alkaloids by high-performance liquid chromatography-isotope dilution thermospray-mass spectrometry 554(1991)227
- Cathepsins**
High-performance liquid chromatographic method for the simultaneous purification of cathepsins B, H and L from human liver 568(1991)55
- Cathinone**
Determination of (S)-(-)-cathinone and its metabolites (R,S)-(-)-norephedrine and (R,R)-(-)-norpseudoephedrine in urine by high-performance liquid chromatography with photodiode-array detection 593(1992)79
- Cations**
Carbohydrate separation by ligand-exchange liquid chromatography. Correlation between the formation of sugar-cation complexes and the elution order 558(1991)89

Determination of major ions in snow and ice cores by ion chromatography 594(1992)225

Relationship between zone length and step height in isotachopheresis 595(1992)327

Cations, divalent

Measurement of stability constants for complexes of divalent cations with tartaric acid by ion chromatography 589(1992)185

Cebaracetam

Fast systematic approach for the determination of drugs in biological fluids by fully automated high-performance liquid chromatography with on-line solid-phase extraction and automated cartridge exchange. Application to cebaracetam in human urine 577(1992)335

Cedrene

Effects of collection solvent parameters and extraction cell geometry on supercritical fluid extraction efficiencies 594(1992)297

Cedrol

Effects of collection solvent parameters and extraction cell geometry on supercritical fluid extraction efficiencies 594(1992)297

Cefaclor

Reversed-phase high-performance liquid chromatography of amphoteric β -lactam antibiotics: effects of columns, ion-pairing reagents and mobile phase pH on their retention times 564(1991)195

High-performance liquid chromatographic determination of loracarbef, a potential metabolite, cefaclor and cephalixin in human plasma, serum and urine 567(1991)129

Cefcanel

Determination of cefcanel in plasma and urine by high-performance liquid chromatography using coupled columns, after administration of the new cephalosporin prodrug cefcanel daloxate hydrochloride 578(1992)71

Cefixime

Determination of cefixime and its metabolites by high-performance capillary electrophoresis 590(1992)364

Cefotiam

Automated high-performance liquid chromatographic method for the simultaneous determination of cefotiam and Δ^3 -cefotiam in human plasma using column switching 577(1992)174

Cefpirome

Microanalytical high-performance liquid chromatographic assay for cefpirome in human milk and urine 574(1992)356

Cefpodoxime

Determination of the third generation oral cephalosporin cefpodoxime in biological fluids by high-speed high-performance liquid chromatography 563(1991)205

Ceftazidime

Determination of ceftazidime in dolphin serum by liquid chromatography with ultraviolet-visible detection and confirmation by thermospray liquid chromatography-mass spectrometry 576(1992)305

α -Cellulose

Optimization of supercritical fluid extraction of volatile constituents from a model plant matrix 600(1992)175

Centbutindole

Simultaneous determination of centbutindole and its hydroxy metabolite in serum by high-performance liquid chromatography 572(1991)219

Cephalexin

Reversed-phase high-performance liquid chromatography of amphoteric β -lactam antibiotics: effects of columns, ion-pairing reagents and mobile phase pH on their retention times 564(1991)195

High-performance liquid chromatographic determination of loracarbef, a potential metabolite, cefaclor and cephalixin in human plasma, serum and urine 567(1991)129

Cephalomannine

Separation of taxol from related taxanes in *Taxus brevifolia* extracts by isocratic elution reversed-phase microcolumn high-performance liquid chromatography 587(1991)300

Cephalosporins

Determination of the third generation oral cephalosporin cefpodoxime in biological fluids by high-speed high-performance liquid chromatography 563(1991)205

Microanalytical high-performance liquid chromatographic assay for cefpirome in human milk and urine 574(1992)356

Hydrophobicity of β -lactam antibiotics. Explanation and prediction of their behaviour in various partitioning solvent systems and reversed-phase chromatography 585(1991)3

Aqueous high-performance size-exclusion chromatographic assay for high-molecular-weight impurities in ceftiofur sodium 595(1992)185

Cephradine

Reversed-phase high-performance liquid chromatography of amphoteric β -lactam antibiotics: effects of columns, ion-pairing reagents and mobile phase pH on their retention times 564(1991)195

Ceronapril

Determination of parts per million levels of trifluoroacetic acid in ceronapril bulk substance by headspace capillary gas chromatography 596(1992)85

C1-esterase inhibitor

Evaluation of the chromatographic procedure for the preparation of a high-purity C1-esterase inhibitor concentrate from cryosupernatant plasma 582(1992)65

Cetiofur sodium

Aqueous high-performance size-exclusion chromatographic assay for high-molecular-weight impurities in cetiofur sodium 595(1992)185

Cetirizine

Determination of cetirizine in human urine by high-performance liquid chromatography 565(1991)504

Determination of cetirizine in serum using reversed-phase high-performance liquid chromatography with ultraviolet spectrophotometric detection 583(1992)128

Cetylpyridinium ion

Determination of benzocaine, dextromethorphan and cetylpyridinium ion by high-performance liquid chromatography with UV detection 558(1991)147

Cetyltrimethylammonium bromide

Determination of metal ions by on-line complexation and ion-pair chromatography 590(1992)263

CGS 20625

Determination of a potential anxiolytic drug (CGS 20625) in human plasma by high-performance liquid chromatography 568(1991)487

CGS 21680

Automated-extraction, high-performance liquid chromatographic method and pharmacokinetics in rats of a highly A₂-selective adenosine agonist, CGS 21680 578(1992)77

Chaconines

Determination of potato glycoalkaloids and their aglycone in blood serum by high-performance liquid chromatography. Application to pharmacokinetic studies in humans 573(1992)69

Chalcones

Origin and control of multi-peak formation in the analysis of trimethylsilyl derivatives of flavanone aglycones by capillary column gas chromatography 586(1991)323

Chelating compounds

Suppression of the effect of metal impurities in alkylsilylated silica packing materials 594(1992)87

Chemical warfare agents

Methods for the analysis of thiodiglycol sulphoxide, a metabolite of sulphur mustard, in urine using gas chromatography-mass spectrometry 558(1991)393

Determination of chemical warfare agents, their hydrolysis products and related compounds in soil 589(1992)287

Mass spectrometric identification of products formed during degradation of ethyl dimethylphosphoramidocyanidate (tabun) 598(1992)89

Capillary column gas chromatography-ammonia and deuterated ammonia chemical ionization mass spectrometry of sulfur vesicants 600(1992)267

Supercritical fluid extraction of chemical warfare agent simulants from soil 600(1992)273

Chimasorb 905

Determination of polymeric hindered amine light stabilizers in polyolefins by high-performance liquid chromatography 553(1991)415

Chimasorb 944

Determination of polymeric hindered amine light stabilizers in polyolefins by high-performance liquid chromatography 553(1991)415

Chinchocaine

Determination of limiting ionic mobilities and dissociation constants of some local anaesthetics 596(1992)265

Chloramphenicol

Selected applications of cyclodextrin selectors in capillary electrophoresis 559(1991)215

Analysis of chloramphenicol residues in swine tissues and milk: comparative study using different screening and quantitative methods 566(1991)173

Determination of chloramphenicol in muscle using a particle beam interface for combining liquid chromatography with negative-ion chemical ionization mass spectrometry 582(1992)113

Automated high-performance liquid chromatographic determination of chloramphenicol in milk and swine muscle tissue using on-line immunoaffinity sample clean-up 583(1992)77

Chloride

Peak-shape analysis and noise evaluation in suppressed ion chromatography for ultra-trace ion analysis 556(1991)249

Mixed potential response at metallic electrodes in potentiometric detection for ion chromatography 589(1992)201

Simultaneous ion chromatographic determination of chloride and calcium contained in Ringer's injection using a chelating agent as eluent 589(1992)358

Gas chromatography of chloride and bromide as phenylboronic acid/mercuric nitrate derivatives with microwave induced plasma atomic emission detection 594(1992)395

Chlorinated hydrocarbons

Simultaneous separation and determination of hydrocarbons and organochlorine compounds by using a two-step microcolumn 555(1991)221

Methods for the analysis of persistent chlorinated hydrocarbons in tissues (Review) 580(1992)257

Chlorobenzene

Estimation of cellulose polarity by gas chromatography 552(1991)365

High-temperature open-tubular capillary column liquid chromatography 592(1992)239

Reversed- and normal-phase separations by high-temperature open-tubular column liquid chromatography 598(1992)153

Thermal desorption-gas chromatography for the determination of benzene, aniline, nitrobenzene and chlorobenzene in workplace air 600(1992)344

1-[(5-Chloro-2-benzofuranyl)(2-chlorophenyl)methyl]-1*H*-imidazole

Gas chromatographic-mass spectrometric determination of two new antimycotic agents, 1-[(5-chloro-2-benzofuranyl)(2-chlorophenyl)methyl]-1*H*-imidazole and 1-[(5-bromo-2-benzofuranyl)phenylmethyl]-1*H*-imidazole, in rabbit plasma following topical administration: a preliminary comparison with bifonazole 567(1991)105

2-Chloroethyl (2-chloroethoxy)ethyl sulphide

Capillary column gas chromatography-ammonia and deuterated ammonia chemical ionization mass spectrometry of sulfur vesicants 600(1992)267

Chloroethylethylsulphide

Supercritical fluid extraction of chemical warfare agent simulants from soil 600(1992)273

Chlorogenic acids

High-performance liquid chromatography of the phenolic compounds in the juice of some French cider apple varieties 555(1991)125

***p*-Chloronitrobenzene**

Determination of *p*-chloronitrobenzene in plasma by reversed-phase high-performance liquid chromatography 570(1991)321

2-Chloro-4-nitrophenyl acetate

Carrier membrane as a stationary phase for affinity chromatography and kinetic studies of membrane-bound enzymes 566(1991)341

Chlorophenols

Calculation of retention and selectivity in reversed-phase liquid chromatography 552(1991)91

Sensitive liquid chromatographic determination of alkyl-, nitro- and chlorophenols by precolumn derivatization with dansyl chloride, postcolumn photolysis and peroxyoxalate chemiluminescence detection 553(1991)345

2-Chloro-3-phenylmethoxypropanoic acid

Optical resolution of 2-chloro-3-phenylmethoxypropanoic acid after derivatization with (*S*)-2-octanol by high-performance liquid chromatography 589(1992)346

5-(4-Chlorophenyl)-2-[(4-methylphenyl)sulphonyl]-4-pentynoic acid

Determination of a new blood glucose-lowering agent in biological fluids by capillary gas chromatography using electron-capture detection 567(1991)465

***m*-Chlorophenylpiperazine**

High-performance liquid chromatographic assay with ultraviolet detection for the determination of etoperidone and two active metabolites, 5-(1-hydroxyethyl)etoperidone and 1-(3-chlorophenyl)piperazine, in plasma 567(1991)433

High-performance liquid chromatographic method for the determination of nefazodone and its metabolites in human plasma using laboratory robotics 570(1991)129

Chlorophyllide *a*

Highly sensitive determination of photosynthetic pigments in marine *in situ* samples by high-performance liquid chromatography 558(1991)131

Chlorophylls

Thermodesorption-gas chromatography-mass spectrometric analysis of biological materials for potential molecular precursors of the constituents of the crude oils 552(1991)325

Identification of chlorophyll transformation products in a lake sediment by combined liquid chromatography-mass spectrometry 557(1991)271

Highly sensitive determination of photosynthetic pigments in marine *in situ* samples by high-performance liquid chromatography 558(1991)131

Determination of carotenoid pigments in several tree leaves by reversed-phase high-performance liquid chromatography 585(1991)168

Determination of chlorophylls and carotenoids by high-performance liquid chromatography during olive lactic fermentation 585(1991)259

Chloroplast DNA

Purification of common bean chloroplast DNA by DEAE cellulose column chromatography 558(1991)318

3-Chloropropane-1,2-diol

Determination of 3-chloropropane-1,2-diol in liquid hydrolysed vegetable proteins by capillary gas chromatography with flame ionization detection 555(1991)311

Chloropropanols

Determination of chloropropanols in protein hydrolysates 589(1992)109

Chloroquine

High-performance liquid chromatographic separation of the enantiomers of hydroxychloroquine and its major metabolites in biological fluids using an α_1 -acid glycoprotein stationary phase 570(1991)119

Simultaneous determination of chloroquine and its three metabolites in human plasma, whole blood and urine by ion-pair high-performance liquid chromatography 574(1992)305

Chlorotetracycline

Separation of antibiotics by high-performance capillary electrophoresis with photodiode-array detection 585(1991)133

Chlorotriazines

Evaluation of eluents in thermospray liquid chromatography-mass spectrometry for identification and determination of pesticides in environmental samples 553(1991)311

Interference removal in the organic trace-level analysis of aqueous environmental samples by on-line liquid chromatographic preconcentration techniques with two precolumns 553(1991)329

Determination of chlorotriazines in aqueous environmental samples at the ng/l level using preconcentration with a cation exchanger and on-line high-performance liquid chromatography 585(1991)67

2-Chlorovinylarsonous acid

Gas chromatographic determination of the lewisite hydrolysate, 2-chlorovinylarsonous acid, after derivatization with 1,2-ethanedithiol 558(1991)235

Chlorpheniramine

Retention and enantioselectivity of racemic solutes on a modified ovomucoid-bonded column. I. Cross-linking with glutaraldehyde 592(1992)301

Chlorprenaline

Application of an ovomucoid-conjugated polymer column for the enantiospecific determination of chlorprenaline concentrations in plasma 566(1991)163

Chlorthalidone

Coupled column chromatography in chiral separations: systems employing β -cyclodextrin phases for chiral separation 557(1991)199

Chlortoluron

Application of high-performance thin-layer chromatography and automated multiple development for the identification and determination of pesticides in water 553(1991)489

Cholanoic aids

Studies on steroids. CCLIV. Gas chromatographic-mass spectrometric determination of 4- and 6-hydroxylated bile acids in human urine with negative ion chemical ionization detection 574(1992)1

Cholesterol

Simplified gas chromatographic method for the simultaneous determination of phytosterols and cholesterol 563(1991)188

Simple thin-layer chromatographic purification procedure for the determination of cholesterol ester fatty acid compositions 564(1991)272

Automated separation and quantitation of lipid fractions by high-performance liquid chromatography and mass detection 567(1991)21

Comparison of gel permeation chromatography, density gradient ultracentrifugation and precipitation methods for quantitation of very-low-, low- and high-density lipoprotein cholesterol 570(1991)382

Determination of cholesterol and cortisone absorption in polyurethane. I. Methodology using size-exclusion chromatography and dual detection 572(1991)41

Sources of plant sterol contaminants encountered in low level steroid analysis 572(1991)307

Determination of cholesterol in sub-nanomolar quantities in biological fluids by high-performance liquid chromatography 579(1992)329

Effect of the sample solvent on band profiles in preparative liquid chromatography using non-aqueous reversed-phase high-performance liquid chromatography 588(1991)1

Use of hematoporphyrin as a fluorescent stain for detection of lipids in high-performance thin-layer chromatography 588(1991)295

Cholesterol esters

Simple thin-layer chromatographic purification procedure for the determination of cholesterol ester fatty acid compositions 564(1991)272

Separation of cholesterol esters by silver ion chromatography using high-performance liquid chromatography or solid-phase extraction columns packed with a bonded sulphonic acid phase 565(1991)103

Automated separation and quantitation of lipid fractions by high-performance liquid chromatography and mass detection 567(1991)21

High-performance liquid chromatographic determination of cholesteryl esters in the blood of obese children 571(1991)19

Cholesterol 7 α -hydroxylase

Simultaneous assay of the activities of two key enzymes in cholesterol metabolism by gas chromatography-mass spectrometry 565(1991)53

Cholesteryl acetate

Cholesteryl acetate as a stationary phase for the gas chromatography of some volatile oil constituents 600(1992)337

Cholesteryl esters

Use of hematoporphyrin as a fluorescent stain for detection of lipids in high-performance thin-layer chromatography 588(1991)295

Choline

Glassy carbon pre-column for direct determination of acetylcholine and choline in biological samples using liquid chromatography with electrochemical detection 575(1992)29

Cation-exchange liquid chromatography of choline and acetylcholine on free shielded silanols of silica-based reversed-phase stationary phases 596(1992)43

Choline analogues

Synthesis of novel selenium-containing choline and acetylcholine analogues and their quantitation using a pyrolysis-gas chromatography-mass spectrometry assay 585(1991)101

Chondroitin sulphate

Chromatographic characterization of ion exchangers for high-performance liquid chromatography of proteins. I. Chromatographic determination of loading capacity for low- and high-molecular mass anions 556(1991)363

Determination of human urinary hyaluronic acid, chondroitin sulphate and dermatan sulphate as their unsaturated disaccharides by high-performance liquid chromatography 565(1991)141

Study of the measurement of chondroitin sulphates in rabbit plasma and serum 574(1992)142

Chemiluminescence high-performance liquid chromatography for the determination of hyaluronic acid, chondroitin sulphate and dermatan sulphate 579(1992)203

Chromium

Effect of the substituents of β -diketonato ligands on R_f values of tris(chelate) transition metal complexes obtained by normal- and reversed-phase thin-layer chromatography on unmodified silica gel 585(1991)359

Separation and determination of trace amounts of vanadium (V), chromium(III) and iron(III) with 2-(2-thienylazo)-5-diethylaminophenol chelates by high-performance liquid chromatography 598(1992)298

Cicletanine

Determination of cicletanine in human plasma by high-performance liquid chromatography using automated column switching 573(1992)93

Determination of cicletanine enantiomers in plasma by high-performance capillary electrophoresis 574(1992)127

Comparison of high-performance liquid chromatography and high-performance capillary electrophoresis for the determination of cicletanine in plasma 581(1992)219

Ciclopiroxolamine

Micro-liquid chromatography method for the determination of ciclopiroxolamine after pre-column derivatization in topical formulations 553(1991)41

Ciclotropium

Determination of the quaternary compound ciclotropium in human biological material after hydrolysis and derivatization with the fluorophor flunoxaprofen chloride 572(1991)181

Cimaterol

High-performance liquid chromatographic determination of clenbuterol and cimaterol using post-column derivatization 564(1991)537

Residue screening for the β -agonists clenbuterol, salbutamol and cimaterol in urine using enzyme immunoassay and high-performance liquid chromatography 564(1991)551

Cimetidine

Electrochromatographic solid-phase extraction for determination of cimetidine in serum by micellar electrokinetic capillary chromatography 559(1991)547

Determination of cimetidine and related impurities in pharmaceutical formulations by high-performance liquid chromatography 586(1991)149

Determination of cimetidine in pharmaceutical preparations by capillary zone electrophoresis 586(1991)177

Cimetropium bromide

Rapid and efficient purification of cimetropium bromide and mifentidine drug metabolite mixtures derived from microsomal incubates for analysis by mass spectrometry 575(1992)75

Cinchona alkaloids

High-performance liquid chromatographic-mass spectrometric assay of high-value compounds for pharmaceutical use from plant cell tissue culture: *Cinchona* alkaloids 562(1991)713

Cinchonine

High-performance liquid chromatographic-mass spectrometric assay of high-value compounds for pharmaceutical use from plant cell tissue culture: *Cinchona* alkaloids 562(1991)713

Cinnamic acid

Determination of cinnamic acid and paeoniflorin in traditional chinese medicinal preparations by high-performance liquid chromatography 593(1992)191

Ciprofloxacin

Determination of ciprofloxacin in human serum by liquid chromatography 563(1991)211

Improved high-performance liquid chromatographic determination of ciprofloxacin and its metabolites in human specimens 582(1992)263

Ciprostene

Micro-quantitative determination of ciprostene in plasma by gas chromatography-mass spectrometry coupled with an antibody extraction 568(1991)460

Cisplatin

Separation of some platinum(II) complexes by ionic strength gradient on a solvent-generated ion-exchange sorbent 586(1991)291

Citric acids

Determination of volatile and non-volatile organic acids in technical sugar solutions by ion-exclusion chromatography 555(1991)65

Chromatographic determination of citric acid for monitoring the mould process 558(1991)302

Citropten

High-performance liquid chromatographic determination of citropten and bergapten in suction blister fluid after solar product application in humans 563(1991)407

CL 275,838

High-performance liquid chromatographic method for the determination of a potential memory-enhancing compound (CL 275,838) and its desbenzyl metabolite in rat plasma or serum 572(1991)346

Liquid chromatography of the potential memory-enhancing agent CL 275,838 and its main metabolites, using a post-column photochemical reactor and fluorimetric detection 581(1992)109

Clarithromycin

Simultaneous determination of clarithromycin and 14(*R*)-hydroxyclearithromycin in plasma and urine using high-performance liquid chromatography with electrochemical detection 571(1991)199

Clebopride

Simultaneous determination of clebopride and a major metabolite N-desbenzylclebopride in plasma by capillary gas chromatography-negative-ion chemical ionization mass spectrometry 564(1991)147

Clenbuterol

Test strip enzyme immunoassays and the fast screening of nortestosterone and clenbuterol residues in urine samples at the parts per billion level 564(1991)413

Determination of β_2 -receptor agonists in bovine urine and liver by gas chromatography-tandem mass spectrometry 564(1991)515

Determination of clenbuterol in urine as its cyclic boronate derivative by gas chromatography-mass spectrometry 564(1991)529

High-performance liquid chromatographic determination of clenbuterol and cimaterol using post-column derivatization 564(1991)537

Residue screening for the β -agonists clenbuterol, salbutamol and cimaterol in urine using enzyme immunoassay and high-performance liquid chromatography 564(1991)551

Cloricromene acid

Determination of the coumarin derivative cloricromene acid in rabbit plasma and platelets 576(1992)143

Cloxacillin

Development of analytical methods for some penicillins in bovine milk by ion-paired chromatography and confirmation by thermospray mass spectrometry 567(1991)389

High-performance liquid chromatographic determination of penicillin G, penicillin V and cloxacillin in beef and pork tissues 593(1992)15

Solvent degradation of cloxacillin *in vitro*. Tentative identification of degradation products using thermospray liquid chromatography-mass spectrometry 594(1992)195

Collaborative study of the determination of cloxacillin by column liquid chromatography 600(1992)333

Clozapine

Determination of clozapine and its major metabolites in human serum using automated solid-phase extraction and subsequent isocratic high-performance liquid chromatography with ultraviolet detection 583(1992)209

Clozapine N-oxide

Determination of clozapine and its major metabolites in human serum using automated solid-phase extraction and subsequent isocratic high-performance liquid chromatography with ultraviolet detection 583(1992)209

Coagulation factors

Comparison of the performance of immunosorbents prepared by site-directed or random coupling of monoclonal antibodies 558(1991)55

Physical and biochemical characterization of five commercial resins for immunoaffinity purification of factor IX 595(1992)103

Cobalt

Determination of cobalt in urine by gas chromatography–mass spectrometry employing nickel as an internal standard 576(1992)297

Effect of the substituents of β -diketonato ligands on R_F values of tris(chelate) transition metal complexes obtained by normal- and reversed-phase thin-layer chromatography on unmodified silica gel 585(1991)359

Determination of divalent trace metals in natural waters by preconcentration on N,N,N',N'-tetra(2-aminoethyl)ethylenediamine-silica followed by on-line ion chromatography 600(1992)203

Cobalt complexes

Rapid optical resolution of electrically neutral cobalt(III) chelate complexes by gel permeation chromatography 585(1991)337

Cocaethylene

Rapid assay of cocaine, opiates and metabolites by gas chromatography–mass spectrometry (Review) 580(1992)43

³²P-Postlabelling and mass spectrometric methods for analysis of bulky, polyaromatic carcinogen–DNA adducts in humans (Review) 580(1992)293

Determination of cocaethylene, cocaine and their metabolites in rat serum microsamples by high-performance liquid chromatography, and its application to pharmacokinetic studies in rodents 582(1992)167

Cocaine

Determination of cocaine, benzoylecgonine and ecgonine methyl ester in plasma by reversed-phase high-performance liquid chromatography 570(1991)412

Analysis of meconium for cocaine in neonates 575(1992)158

Rapid assay of cocaine, opiates and metabolites by gas chromatography–mass spectrometry (Review) 580(1992)43

Determination of cocaethylene, cocaine and their metabolites in rat serum microsamples by high-performance liquid chromatography, and its application to pharmacokinetic studies in rodents 582(1992)167

Determination of the enantiomeric composition of samples of cocaine by normal-phase high-performance liquid chromatography with UV detection 590(1992)354

Determination of limiting ionic mobilities and dissociation constants of some local anaesthetics 596(1992)265

Codeine

Direct determination of codeine, norcodeine, morphine and normorphine with their corresponding O-glucuronide conjugates by high-performance liquid chromatography with electrochemical detection 570(1991)309

High-performance liquid chromatographic determination of morphine, morphine-3-glucuronide, morphine-6-glucuronide and codeine in biological samples using multi-wavelength forward optical detection 571(1991)263

High-performance liquid chromatographic determination of morphine, morphine-3-glucuronide, morphine-6-glucuronide and codeine in biological samples using multi-wavelength forward optical detection: a reply 578(1992)333

High-performance liquid chromatographic determination of morphine, morphine-3-glucuronide, morphine-6-glucuronide and codeine in biological samples using multi-wavelength forward optical detection. Reply to Zweipfenning 578(1992)335

High-performance liquid chromatographic determination of morphine, morphine-3-glucuronide, morphine-6-glucuronide and codeine in biological samples using multi-wavelength forward optical detection: a reply 579(1992)189

High-performance liquid chromatographic determination of morphine, morphine-3-glucuronide, morphine-6-glucuronide and codeine in biological samples using multi-wavelength forward optical detection. Reply to Bogusz 579(1992)191

Rapid assay of cocaine, opiates and metabolites by gas chromatography–mass spectrometry (Review) 580(1992)43

Retention reproducibility of basic drugs in high-performance liquid chromatography on a silica column with a methanol–high-pH buffer eluent. Changes in selectivity with the age of the stationary phase 592(1992)85

Coenzyme A

Synthesis, characterisation and high-performance liquid chromatography of C₆-C₁₆ dicarboxylyl-mono-coenzyme A and -mono-carnitine esters 570(1991)253

Coenzyme B₁₂

Purification of the coenzyme B₁₂-containing 2-methyleneglutarate mutase from *Clostridium barkeri* by high-performance liquid chromatography 587(1991)93

Coenzyme Q

Improved high-performance liquid chromatographic method for the determination of coenzyme Q₁₀ in plasma 593(1992)217

Coenzymes

Application of a pellicular anion-exchange resin to the separation of inorganic and organic anions by single-column anion exchange 592(1992)101

Colchiceinamides

R_M values of some colchicines and colchiceinamides determined by reversed-phase thin-layer chromatography 591(1992)367

Colchicines

R_M values of some colchicines and colchiceinamides determined by reversed-phase thin-layer chromatography 591(1992)367

Colipase

Separation and characterization of the precursor and activated forms of porcine and human pancreatic colipase by reversed-phase liquid chromatography 583(1992)246

Collagen

Use of biospecific interactions of collagen, fibronectin and their fragments in affinity chromatography 577(1992)267

Silver staining of collagen type I after sodium dodecylsulphate polyacrylamide gel electrophoresis: effect of Maillard reaction 578(1992)53

Quantitative analysis of collagen and elastin cross-links using a single-column system 582(1992)49

Determination of calcium ion in the presence of phosphate anion and collagen by capillary-type isotachopheresis 586(1991)363

Characterization of the interaction between human plasma fibronectin and collagen by means of affinity electrophoresis 597(1992)351

Colloids

Asymmetric-channel flow field-flow fractionation with exponential force-field programming 593(1992)339

Colubrine

Separation of *Strychnos nux-vomica* alkaloids by high-performance liquid chromatography 587(1991)318

Complement

Rapid isolation of human complement component C9 to verify the specificity of a haemolytic C9 microassay 591(1992)187

Conalbumin

High-sensitivity laser-induced fluorescence detection of native proteins in capillary electrophoresis 595(1992)319

Copoly(styrene-acrylamide)s

Beaded polymer supports and gels (Review). II. Physico-chemical criteria and functionalization 586(1991)199

Copper

Separation of metal complexes of ethylenediaminetetraacetic acid in environmental water samples by ion chromatography with UV and potentiometric detection 558(1991)181

Thermoanalytical and chromatographic studies of copper(II), nickel(II) and oxovanadium(IV) complexes of tetradentate Schiff bases derived from β -diketones and 2,3-diaminopentane 558(1991)187

Determination of divalent trace metals in natural waters by preconcentration on N,N,N',N'-tetra(2-aminoethyl)ethylenediamine-silica followed by on-line ion chromatography 600(1992)203

Copper-binding proteins

Chromatographic analysis of low-molecular-mass copper-binding ligands from the crab species *Scylla serrata* and *Portunus pelagicus* 599(1992)21

Copper sulphate

Study of salt hydrates by gas-solid chromatography 557(1991)495

Coproporphyrin

Application of ion-pair high-performance liquid chromatography to detection of the atypical coproporphyrin isomers II and IV in human faeces 572(1991)317

COR 3224

Determination of a new 2-amino-2-oxazoline (COR 3224) in plasma and brain tissue of the rat by high-performance liquid chromatography with electrochemical detection 563(1991)476

Corticosteroids

Hypertensive congenital adrenal enzymatic defects detected by high-performance liquid chromatography of corticosteroids 553(1991)201

Evaluation of an automated thermospray liquid chromatography-mass spectrometry system for quantitative use in bioanalytical chemistry 554(1991)215

Characterization of prednisone, prednisolone and their metabolites by gas chromatography–mass spectrometry 565(1991)45

Routine screening and quantitation of urinary corticosteroids using bench-top gas chromatography–mass-selective detection 573(1992)183

Thermospray liquid chromatography–mass spectrometry of corticosteroids 577(1992)221

Stable isotope dilution mass spectrometry for the simultaneous determination of cortisol, cortisone, prednisolone and prednisone in plasma 579(1992)193

Corticosterone

Thin-layer chromatography on silica-coated aluminium sheet as an adjunct to radioimmunoassay of steroids 563(1991)178

Fluorimetric detection of serum corticosterone using high-performance liquid chromatography 581(1992)267

Cortisol

On-line derivatization of eluted substances in dynamic high-performance liquid chromatography–mass spectrometry through the particle-beam interface 554(1991)125

Increasing thermospray response for cortisol by derivatization 554(1991)149

Thin-layer chromatography on silica-coated aluminium sheet as an adjunct to radioimmunoassay of steroids 563(1991)178

Automated direct high-performance liquid chromatographic assay for estetrol, estriol, cortisone and cortisol in serum and amniotic fluid 568(1991)35

Determination of serum cortisol by reversed-phase liquid chromatography using precolumn sulphuric acid–ethanol fluorescence derivatization and column switching 570(1991)1

Simultaneous determination of cortisol and cortisone in human plasma by stable-isotope dilution mass spectrometry 576(1992)47

Stable isotope dilution mass spectrometry for the simultaneous determination of cortisol, cortisone, prednisolone and prednisone in plasma 579(1992)193

Simultaneous determination of serum cortisol and cortisone by reversed-phase liquid chromatography with ultraviolet detection 584(1992)147

Cortisone

On-line derivatization of eluted substances in dynamic high-performance liquid chromatography–mass spectrometry through the particle-beam interface 554(1991)125

Automated direct high-performance liquid chromatographic assay for estetrol, estriol, cortisone and cortisol in serum and amniotic fluid 568(1991)35

Determination of cholesterol and cortisone absorption in polyurethane. I. Methodology using size-exclusion chromatography and dual detection 572(1991)41

Simultaneous determination of cortisol and cortisone in human plasma by stable-isotope dilution mass spectrometry 576(1992)47

Thermospray liquid chromatography–mass spectrometry of corticosteroids 577(1992)221

Stable isotope dilution mass spectrometry for the simultaneous determination of cortisol, cortisone, prednisolone and prednisone in plasma 579(1992)193

Simultaneous determination of serum cortisol and cortisone by reversed-phase liquid chromatography with ultraviolet detection 584(1992)147

Corticoic acids

High-performance liquid chromatographic determination of corticoic and cortolonic acids as pyrenyl ester derivatives 564(1991)27

Cortolic acid

High-performance liquid chromatographic determination of cortolic and cortolonic acids as pyrenyl ester derivatives 564(1991)27

Cortolonic acid

High-performance liquid chromatographic determination of cortolic and cortolonic acids as pyrenyl ester derivatives 564(1991)27

Cotinine

Elimination of caffeine interference in high-performance liquid chromatographic determination of cotinine in human plasma 568(1991)267

Simultaneous determination of cotinine and *trans*-3'-hydroxycotinine in human serum by high-performance liquid chromatography 577(1992)358

Electron-impact and chemical ionization detection of nicotine and cotinine by gas chromatography–mass spectrometry in rat plasma and brain 579(1992)93

Gas chromatographic analysis of nicotine and cotinine in hair 580(1992)347

***p*-Coumaric acid**

High-performance liquid chromatography of the phenolic compounds in the juice of some French cider apple varieties 555(1991)125

Coumarins

Thermospray and particle beam liquid chromatographic-mass spectrometric analysis of coumarin anticoagulants 562(1991)31

Rapid and sensitive determination of coumarin and 7-hydroxycoumarin and its glucuronide conjugate in urine and plasma by high-performance liquid chromatography 582(1992)137

Retention behaviour of closely related coumarins in thin-layer chromatographic preassays for high-performance liquid chromatography according to the "PRISMA" model 593(1992)329

Coumatetralyl

Screening for coumatetralyl in soft drinks by solid-matrix extraction and high-performance liquid chromatography with diode-array detection 553(1991)305

CPT-11

Simultaneous determination of the camptothecin analogue CPT-11 and its active metabolite SN-38 by high-performance liquid chromatography: application to plasma pharmacokinetic studies in cancer patients 575(1992)275

Selective and sensitive determination of lactone and hydroxy acid forms of camptothecin and two derivatives (CPT-11 and SN-38) by high-performance liquid chromatography with fluorescence detection 588(1991)165

Creatine

Rapid determination of creatine, phosphocreatine, purine bases and nucleotides (ATP, ADP, AMP, GTP, GDP) in heart biopsies by gradient ion-pair reversed-phase liquid chromatography 575(1992)19

Rapid separation of creatine, phosphocreatine and adenosine metabolites by ion-pair reversed-phase high-performance liquid chromatography in plasma and cardiac tissue 576(1992)149

Creatine kinase

Electrophoresis of serum isoenzymes and proteins following acute myocardial infarction (Review) 569(1991)323

Electrophoretic analysis of enzyme-linked immunoglobulins and their clinical significance (Review) 569(1991)347

Creatine phosphate

Isocratic high-performance liquid chromatographic analysis of myocardial creatine phosphate and adenine nucleotides 563(1991)129

Simultaneous determination of myocardial nucleotides, nucleosides, purine bases and creatine phosphate by ion-pair high-performance liquid chromatography 578(1992)39

Creatinine

Simultaneous determination of urinary creatinine and aromatic amino acids by cation-exchange chromatography with ultraviolet detection 566(1991)19

New analyser for the determination of urinary vanillylmandelic acid, homovanillic acid and creatinine 567(1991)11

Multidimensional high-performance liquid chromatography on Pinkerton ISRP and RP18 columns: direct serum injection to quantify creatinine 581(1992)129

Determination of creatinine and ultraviolet-absorbing amino acids and organic acids in urine by reversed-phase high-performance liquid chromatography 583(1992)1

Use of various types of column reactors for flow-injection analysis 597(1992)435

Cresols

Calculation of retention and selectivity in reversed-phase liquid chromatography 552(1991)91

Gas chromatographic determination of urinary phenol conjugates after acid hydrolysis/extractive acetylation 574(1992)349

High-performance liquid chromatographic column-switching technique for the determination of intermediates of anaerobic degradation of toluene in ground water microcosm 587(1991)185

Crotamiton

Supercritical fluid chromatography in the routine stability control of antipruritic preparations 553(1991)71

Crown ethers

Preparation and capillary column gas chromatographic characterization of aza-crown ether polysiloxane stationary phases 594(1992)243

 β -Cryptoxanthin

Optimization of an isocratic high-performance liquid chromatographic separation of carotenoids 589(1992)171

Cumene

High-performance liquid chromatographic determination of products of autoxidation of isopropylbenzene derivatives 588(1991)33

Curdlan

Rapid method for preparing a β -glucan-specific sensitive fraction from *Limulus* (*Tachypleus tridentatus*) amoebocyte lysate 567(1991)267

Cyanide

High-performance liquid chromatographic determination of cyanide in human red blood cells by pre-column fluorescence derivatization 582(1992)131

Cyanoazaarenes

Characterization of nitrogen-containing aromatic compounds in soil and sediment by capillary gas chromatography-mass spectrometry after fractionation 558(1991)223

Cyanocobalamin

Ultramicrodetermination of cyanocobalamin in elemental diet by solid-phase extraction and high-performance liquid chromatography with visible detection 590(1992)359

Cyanoginosins

Separation of methylated and non-methylated cyanoginosin-LR homologues of the cyanobacterium *Microcystis aeruginosa* strain PCC 7806 by reversed-phase medium-pressure liquid chromatography 558(1991)430

Cyasorb UV 1084

Determination of Cyasorb UV 1084 and its degradation products in low-density polyethylene film by high-performance liquid chromatography 587(1991)205

Cyclic heptapeptides

Determination of a cyclic heptapeptide, a novel fibrinogen receptor antagonist, in human plasma by high-performance liquid chromatography with automated pre-column derivatization, column switching and fluorescence detection 578(1992)31

N⁶-Cycloalkylated nucleosides

High-performance liquid chromatographic monitoring of metabolic products resulting from the treatment of mouse hepatoma cells with N⁶-cycloalkylated nucleosides 582(1992)253

Cyclodextrins

Effects of β -cyclodextrin in the mobile phase on the retention and indirect detection of non-electrolytes in reversed-phase liquid chromatography. I. Study of aliphatic alcohols 552(1991)563

Effects of β -cyclodextrin in the mobile phase on the retention and indirect detection of non-electrolytes in reversed-phase liquid chromatography. II. Steroids 552(1991)583

Improved high-performance liquid chromatographic resolution of the geometric isomers of 6-hydroxy-4-(1-hydroxy-1-methylethyl)-1-cyclohexene-1-ethanol and by-products with β -cyclodextrin 553(1991)15

Chiral separation by cyclodextrin-modified micellar electrokinetic chromatography 553(1991)503

Retention behavior and chiral recognition of β -cyclodextrin-derivative polymer adsorbed on silica for warfarin, structurally related compounds and Dns-amino acids 555(1991)53

Coupled column chromatography in chiral separations: systems employing β -cyclodextrin phases for chiral separation 557(1991)199

Influence of counter-ion inclusion complexation on the quality of cyclodextrin-supported separations in isotachopheresis 557(1991)215

Effect of derivatization of steroids on their retention behaviour in inclusion chromatography using cyclodextrin as a mobile phase additive 558(1991)306

Selected applications of cyclodextrin selectors in capillary electrophoresis 559(1991)215

Separation of functionalized dextrans by reversed-phase high-performance liquid chromatography 587(1991)288

Separation of polycyclic aromatic hydrocarbons by micellar electrokinetic chromatography with cyclodextrins as modifiers 589(1992)333

Retention behavior of β -cyclodextrin complexes of anthracene and pyrene using reversed-phase liquid chromatography. The effects of *tert.*-butanol and cyclopentanol 594(1992)37

Solvatochromic hydrogen bond donor acidity of cyclodextrins and reversed-phase liquid chromatographic retention of small molecules on a β -cyclodextrin-bonded silica stationary phase 595(1992)45

Stationary phase effect on enantioselectivity of dansyl phenylalanine in microcolumn liquid chromatography with γ -cyclodextrin as mobile phase additive 595(1992)121

Ion-exclusion chromatography using mobile phases containing β -cyclodextrin 595(1992)149

Cyclohexane

Study of microporous carbons by gas chromatographic determination of heats of physisorption 555(1991)183

Cyclohexanecarboxylic acid

Analysis of zapatera olives by gas and high-performance liquid chromatography 594(1992)259

Cyclohexanone

Modern approach to the quantitative determination of volatiles in solid samples. Multiple headspace extraction gas chromatography for the determination of cyclohexanone residues in soil 552(1991)205

Peak purity, yield and throughput in preparative liquid chromatography with self-displacement 592(1992)17

Cyclohexylamine

Liquid membrane work-up of blood plasma samples applied to gas chromatographic determination of aliphatic amines 573(1992)191

Cyclopentanone

Peak purity, yield and throughput in preparative liquid chromatography with self-displacement 592(1992)17

Cyclophosphamide

Gas chromatographic determination of ifostamide in microvolumes of urine and plasma 571(1991)121

3-(5-Cyclopropyl-1,2,4-oxadiazol-3-yl)-5-(1-methylethyl)imidazo[1,5-a]quinoxalin-4(5H)-one

Determination of 3-(5-cyclopropyl-1,2,4-oxadiazol-3-yl)-5-(1-methylethyl)imidazo[1,5-a]quinoxalin-4(5H)-one in serum by high-performance liquid chromatography 563(1991)427

p-Cymene

High-performance liquid chromatographic determination of products of autoxidation of isopropylbenzene derivatives 588(1991)33

Cymoxanil

Behaviour studies of the fungicide cymoxanil in two strains of the fungus *Botrytis cinerea* and in haemolymph of locust and lobster. I. *In situ* monitoring by internal surface reversed-phase high-performance liquid chromatography 566(1991)499

Cypermethrin

Reversed-phase high-performance liquid chromatography of anionic and ethoxylated non-ionic surfactants and pesticides in liquid pesticide formulations 593(1992)227

Cystathionine

Sensitive determination of cystathionine and assays for cystathionine β - and γ -lyase, as well as cystathionine β -synthase, using high-performance liquid chromatography 574(1992)35

Determination of cystathionine and perhydro-1,4-thiazepine-3,5-dicarboxylic acid in the urine of a patient with cystathioninuria using column liquid chromatography-mass spectrometry 579(1992)318

Cystine

Liquid chromatographic-mass spectrometric analysis for screening of patients with cystinuria, and identification of cystine stone 568(1991)445

Cytidine

Enzymophoresis of nucleic acids by tandem capillary enzyme reactor-capillary zone electrophoresis 596(1992)251

Cytidine deaminase

High-performance liquid chromatographic study of some biologically active analogues of arabinosylcytosine 588(1991)147

Cytochrome c

Instrumentation for high-performance capillary electrophoresis-mass spectrometry 559(1991)197

Immobilized metal ion affinity electrophoresis. A study with several model proteins containing histidine 597(1992)357

High-performance liquid chromatography of amino acids, peptides and proteins. CXXI. 8-Hydroxyquinoline-metal chelate chromatographic support: an additional mode of selectivity in immobilized-metal affinity chromatography 599(1992)171

Isoelectric focusing field-flow fractionation. III. Investigation of the influence of different experimental parameters on focusing of cytochrome *c* in the trapezoidal cross-section channel 600(1992)297

Cytochrome P-450

Measurement of enzyme activities of cytochrome P-450 isoenzymes by high-performance liquid chromatographic analysis of products 553(1991)233

Detection of the enzymatic activity of cytochrome P-450 enzymes by high-performance liquid chromatography (Review) 580(1992)325

Immobilized metal affinity chromatography as a means of fractionating microsomal cytochrome P-450 isozymes 587(1991)33

High-performance metal chelate affinity chromatography of cytochromes P-450 using Chelating Superose 587(1991)43

Isolation of cytochrome P-450 components from marmoset liver microsomes by high-performance liquid chromatography 587(1991)117

Cytochrome P-450 IV

Determination of the cytochrome P-450 IV marker, ω -hydroxylauric acid, by high-performance liquid chromatography and fluorimetric detection 564(1991)266

Cytokines

Assessment of capillary electrophoresis in pharmaceutical applications. Analysis and quantification of a recombinant cytokine in an injectable dosage form 559(1991)307

Cytoprotective agents

High-performance liquid chromatographic method for the determination of RGH-5702 in plasma samples 553(1991)29

Cytosine- β -D-arabinoside

Determination of cytosine- β -D-arabinoside in plasma using capillary electrophoresis 568(1991)117

D**Dansylamides**

Influence of mobile phase composition on evaluation of lipophilicity by partition chromatography 557(1991)399

Dansyl chloride

Sensitive fluorescence detection of some nitrosamines by precolumn derivatization with dansyl chloride and high-performance liquid chromatography 589(1992)349

Dansyl leucine

Optimization of resolution in capillary zone electrophoresis: combined effect of applied voltage and buffer concentration 588(1991)315

Dansyl methionine

Optimization of resolution in capillary zone electrophoresis: combined effect of applied voltage and buffer concentration 588(1991)315

Dapsone

Determination of some drugs by micellar electrokinetic capillary chromatography. The pseudo-effective mobility as parameter for screening 585(1991)123

Daunorubicin

Determination of 3'-deamino-3'-[2(S)-methoxy-4-morpholinyl]doxorubicin, a new morpholinyl anthracycline, in plasma by high-performance liquid chromatography with fluorescence detection 578(1992)309

N-(Deacetyl-O-4-vinblastoyl)-L-tryptophan

Biodetermination of N-(deacetyl-O-4-vinblastoyl-23)-L-tryptophan, a metabolite of vintriptyl, by high-performance liquid chromatography with fluorescence detection 574(1992)293

(Deaminohydroxy)toremifene

High-performance liquid chromatographic method for the determination of toremifene and its major human metabolites 565(1991)482

3'-Deamino-3'-[2(S)-methoxy-4-morpholino]doxorubicin

Determination of 3'-deamino-3'-[2(S)-methoxy-4-morpholinyl]doxorubicin, a new morpholinyl anthracycline, in plasma by high-performance liquid chromatography with fluorescence detection 578(1992)309

Stability studies with a high-performance liquid chromatographic method for the determination of a new anthracycline analogue, 3'-deamino-3'-[2-(S)-methoxy-4-morpholino]doxorubicin (FCE 23762), in the final drug formulation 593(1992)47

Debrisoquine

Determination of debrisoquine and metabolites in human urine by gas chromatography-mass spectrometry 570(1991)89

Decahydronaphthalene

High-temperature continuous counter-current gas-liquid chromatography 590(1992)289

Decapeptides

Quantification of a decapeptide anticoagulant in rat and monkey plasma by high-performance liquid chromatography 579(1992)371

Dechloroethylfosfamide

Gas chromatographic determination of ifosfamide in microvolumes of urine and plasma 571(1991)121

Dechloroethylfosfamides

Gas chromatographic determination of 2- and 3-dechloroethylfosfamide in plasma and urine 583(1992)175

Dehydroacetic acid

Simultaneous determination of sorbic acid, dehydroacetic acid and benzoic acid by gas chromatography-mass spectrometry 594(1992)253

Dehydroascorbic acid

Simultaneous high-performance liquid chromatographic determination of ascorbic acid and dehydroascorbic acid in biological samples 565(1991)529

Determination of ascorbic and dehydroascorbic acid in potatoes (*Solanum tuberosum*) and strawberries using ion-exclusion chromatography 594(1992)187

Dehydrochloramphenicol

Determination of chloramphenicol in muscle using a particle beam interface for combining liquid chromatography with negative-ion chemical ionization mass spectrometry 582(1992)113

Dehydroepiandrosterone

Specific assay for unconjugated dehydroepiandrosterone in human plasma by capillary gas chromatography with electron-capture detection 567(1991)299

11-Dehydrothromboxane B₂

Novel derivatization and immunoextraction to improve microanalysis of 11-dehydrothromboxane B₂ in human urine 562(1991)613

Gas chromatographic-mass spectrometric determination of 11-dehydrothromboxane B₂ in human urine 577(1992)1

N-Demethylminocycline

Identification and determination of the two principal metabolites of minocycline in humans 568(1991)363

Deoxyaldohexoses

Carbohydrate analysis with ion chromatography using Eurokat stationary phases. Preparative separation of monosaccharides and their fluorinated derivatives 552(1991)483

Deoxycorticosterone

Thin-layer chromatography on silica-coated aluminium sheet as an adjunct to radioimmunoassay of steroids 563(1991)178

7-Deoxydoxorubicin aglycone

Fully automated determination of a new anthracycline N-/leucyldoxorubicin and six metabolites in plasma by high-performance liquid chromatography with on-line sample handling 574(1992)273

7-Deoxydoxorubicinol aglycone

Fully automated determination of a new anthracycline N-/leucyldoxorubicin and six metabolites in plasma by high-performance liquid chromatography with on-line sample handling 574(1992)273

7-Deoxydoxorubicinolon

Simple and sensitive quantification of anthracyclines in mouse atrial tissue using high-performance liquid chromatography and fluorescence detection 570(1991)209

7-Deoxydoxorubicinon

Simple and sensitive quantification of anthracyclines in mouse atrial tissue using high-performance liquid chromatography and fluorescence detection 570(1991)209

N-(2-Deoxy- β -D-erythro-pentafuranosyl)formamide

Reversed-phase high-performance liquid chromatography-thermospray mass spectrometry of radiation-induced decomposition products of thymine and thymidine 593(1992)133

Deoxyfluoroaldohexoses

Carbohydrate analysis with ion chromatography using Eurokat stationary phases. Preparative separation of monosaccharides and their fluorinated derivatives 552(1991)483

2'-Deoxy-5-fluorouridine

Highly sensitive method for the determination of 5-fluorouracil in biological samples in the presence of 2'-deoxy-5-fluorouridine by gas chromatography-mass spectrometry 564(1991)137

2'-Deoxy-5-iodouridine

Determination of 2'-deoxy-5-iodouridine and its metabolite 5-iodouracil by high-performance liquid chromatography with ultraviolet absorbance detection in human serum 572(1991)327

Deoxynivalenol

Identification of trichothecenes by thermospray, plasmaspray and dynamic fast-atom bombardment liquid chromatography-mass spectrometry 562(1991)555

2'-Deoxynucleoside 5'-monophosphates

Micellar electrokinetic capillary chromatographic separation and laser-induced fluorescence detection of 2'-deoxynucleoside 5'-monophosphates of normal and modified bases 565(1991)197

Deoxynucleotide phosphorothioates

Separation of synthetic phosphorothioate oligodeoxynucleotides from their oxygenated (phosphodiester) defect species by strong-anion-exchange high-performance liquid chromatography 599(1992)35

Deoxynucleotides

Purification of DNA-derived deoxynucleotides from leukocytes involving nuclease elution of an ion-exchange column. 600(1992)229

Deoxyribonucleoside triphosphates

Reproducibility and quantitation of separation for ribonucleoside triphosphates and deoxyribonucleoside triphosphates by capillary zone electrophoresis 559(1991)247

Deoxyuridine

Enzymatic shot-gun 5'-phosphorylation and 3'-sister phosphate exchange: a two-dimensional thin-layer chromatographic technique to measure DNA deoxynucleotide modification 574(1992)41

2'-Deoxyuridine 3-monophosphate

Enzymatic shot-gun 5'-phosphorylation and 3'-sister phosphate exchange: a two-dimensional thin-layer chromatographic technique to measure DNA deoxynucleotide modification 574(1992)41

Dermatan sulphate

Determination of human urinary hyaluronic acid, chondroitin sulphate and dermatan sulphate as their unsaturated disaccharides by high-performance liquid chromatography 565(1991)141

Chemiluminescence high-performance liquid chromatography for the determination of hyaluronic acid, chondroitin sulphate and dermatan sulphate 579(1992)203

High-performance liquid chromatographic-mass spectrometric analysis of oligosaccharides from enzymatic digestion of glycosaminoglycans. Application to human samples 591(1992)225

Dermorphine derivatives

Influence of mobile phase composition on evaluation of lipophilicity by partition chromatography 557(1991)399

25-Desacetyl rifapentine

High-performance liquid chromatographic determination of rifapentine and its metabolite in human plasma by direct injection into a shielded hydrophobic phase column 553(1991)35

Desacetylvinorelbine

Determination of vinorelbine (Navelbine) in tumour cells by high-performance liquid chromatography 581(1992)93

N-Desalkylbuprenorphine

High-performance liquid chromatography with electrochemical detection of buprenorphine and its major metabolite in urine 564(1991)557

N-Desalkylpropafenone

Direct separation of the enantiomers of propafenone, diprafenone and their major metabolites by high-performance liquid chromatography on modified cellulose and amylose chiral stationary phases 585(1991)329

N-Desbenzyleclopriide

Simultaneous determination of clevopride and a major metabolite N-desbenzyleclopriide in plasma by capillary gas chromatography-negative-ion chemical ionization mass spectrometry 564(1991)147

Desbutylhalofantrine

Measurement of halofantrine and its major metabolite desbutylhalofantrine in plasma and blood by high-performance liquid chromatography: a new methodology 581(1992)156

N-Desethylamidarone

High-performance liquid chromatographic assay for amidarone N-deethylation in microsomes of rat liver 579(1992)269

Desethylatrazine

Application of high-performance thin-layer chromatography and automated multiple development for the identification and determination of pesticides in water 553(1991)489

Desethylchloroquine

High-performance liquid chromatographic separation of the enantiomers of hydroxychloroquine and its major metabolites in biological fluids using an α_1 -acid glycoprotein stationary phase 570(1991)119

Determination of hydroxychloroquine and its major metabolites in plasma using sequential achiral-chiral high-performance liquid chromatography 573(1992)253

Simultaneous determination of chloroquine and its three metabolites in human plasma, whole blood and urine by ion-pair high-performance liquid chromatography 574(1992)305

Desethyleneciprofloxacin

Improved high-performance liquid chromatographic determination of ciprofloxacin and its metabolites in human specimens 582(1992)263

Desethylhydroxychloroquine

High-performance liquid chromatographic separation of the enantiomers of hydroxychloroquine and its major metabolites in biological fluids using an α_1 -acid glycoprotein stationary phase 570(1991)119

Determination of hydroxychloroquine and its major metabolites in plasma using sequential achiral-chiral high-performance liquid chromatography 573(1992)253

Desipramine

Determination of imipramine, desipramine and their hydroxy metabolites by reversed-phase chromatography with ultraviolet and coulometric detection 572(1991)247

Isotope effect in gas-liquid chromatography of labelled compounds 588(1991)251

N-Desmethylozapine

Determination of clozapine and its major metabolites in human serum using automated solid-phase extraction and subsequent isocratic high-performance liquid chromatography with ultraviolet detection 583(1992)209

N-Desmethyldiazepam

Quantitation by gas chromatography with selected-ion monitoring mass spectrometry of "natural" diazepam, N-desmethyldiazepam and oxazepam in normal human serum 579(1992)85

O-Desmethylnaproxen

Simultaneous quantitative determination of naproxen, its metabolite 6-O-desmethylnaproxen and their five conjugates in plasma and urine samples by high-performance liquid chromatography on dynamically modified silica 577(1992)325

Simultaneous determination of (*R*)- and (*S*)-naproxen and (*R*)- and (*S*)-6-O-desmethylnaproxen by high-performance liquid chromatography on a Chiral-AGP column 577(1992)362

Determination of naproxen and its metabolite O-desmethylnaproxen with their acyl glucuronides in human plasma and urine by means of direct gradient high-performance liquid chromatography 578(1992)239

Desmethylnortriptyline

High-performance liquid chromatographic determination of amitriptyline and its main metabolites using a silica column with reversed-phase eluent. Application in mice 584(1992)249

Desmethyl-(S)-(-)-ofloxacin

High-performance liquid chromatographic determination of (*S*)-(-)-ofloxacin and its metabolites in serum and urine using a solid-phase clean-up 563(1991)313

O-Desmethylquinidine

High-performance liquid chromatographic method for the quantitation of quinidine and selected quinidine metabolites 572(1991)159

N-Desmethylrifaxin

High-performance liquid chromatography and preliminary pharmacokinetics of rifloxacin and its metabolites, N-desmethylrifloxacin and rifloxacin sulfoxide, in urine of Rhesus monkey *Macaca mulatta* 573(1992)168

N-Desmethyltoremifene

High-performance liquid chromatographic method for the determination of toremifene and its major human metabolites 565(1991)482

N-Desmethylzopiclone

Development of a method for the determination of zopiclone in whole blood 584(1992)199

Desmosine

Separation of elastin cross-links as phenylisothiocyanate derivatives 553(1991)93

High-performance liquid chromatographic determination of desmosine and isodesmosine after phenylisothiocyanate derivatization 572(1991)312

Destruxins

Direct *in vitro* and *in vivo* monitoring of destruxins metabolism in insects using internal surface reversed-phase high-performance liquid chromatography. I. Behaviour of E destruxin in locusts 566(1991)511

Detergents

Comparison of detergents for extraction and ion-exchange high-performance liquid chromatography of Sendai virus membrane proteins 599(1992)157

Deuterium

Gas chromatography for measurement of hydrogen isotopes at tritium processing 586(1991)131

Dexamethasone

Stereochemical analysis of betamethasone and dexamethasone by derivatization and high-performance liquid chromatography 595(1992)203

Dextran B512

Analysis of the interaction between an $\alpha(1\rightarrow6)$ dextran-specific mouse hybridoma antibody and dextran B512 by affinity electrophoresis 597(1992)345

Dextrans

Biosep-SEC-S high-performance size-exclusion chromatographic columns for proteins and peptides 599(1992)25

Dextrins

Separation of functionalized dextrins by reversed-phase high-performance liquid chromatography 587(1991)288

Dextromethorphan

Liquid chromatography-mass spectrometry with ionspray and electrospray interfaces in pharmaceutical and biomedical research 554(1991)39

Determination of benzocaine, dextromethorphan and cetylpyridinium ion by high-performance liquid chromatography with UV detection 558(1991)147

Dextromoramide

Determination of dextromoramide in plasma and whole blood using high-performance liquid chromatography with ultraviolet absorbance detection 573(1992)87

Dextropropoxyphene

Determination of dextropropoxyphene and norpropoxyphene in plasma by high-performance liquid chromatography 581(1992)161

Dextrophan

Liquid chromatography-mass spectrometry with ionspray and electrospray interfaces in pharmaceutical and biomedical research 554(1991)39

Diacetoneketogulonic acid

Determination of diacetoneketogulonic acid in water samples by high-performance thin-layer chromatography 589(1992)390

Diacetyl

Simple and sensitive determination of diacetyl and acetoin in biological samples and alcoholic drinks by gas chromatography with electron-capture detection 577(1992)215

Diacylglycerols

Enantiomer separation by chiral-phase liquid chromatography of urethane derivatives of natural diacylglycerols previously fractionated by reversed-phase liquid chromatography 557(1991)227

Resolution of lipolysis mixtures from soybean oil by a solid-phase extraction procedure 589(1992)353

1,2-Diacyl-*sn*-3-glycerophosphocholine

Optimized method for the determination of 1,2-diacyl-*sn*-glycero-3-phosphocholine and 1,2-diacyl-*sn*-glycero-3-phosphoethanolamine molecular species by enzymatic hydrolysis and gas chromatography 600(1992)257

1,2-Diacyl-*sn*-3-glycerophosphoethanolamine

Optimized method for the determination of 1,2-diacyl-*sn*-glycero-3-phosphocholine and 1,2-diacyl-*sn*-glycero-3-phosphoethanolamine molecular species by enzymatic hydrolysis and gas chromatography 600(1992)257

Diadenosine polyphosphates

Reversed-phase high-performance liquid chromatography of dinucleoside polyphosphates 563(1991)37

Dialdehydes

Separation and preparative isolation of phenolic dialdehydes by on-line overpressured layer chromatography 590(1992)369

4,4'-Diaminodibenzyl-2,2'-disulphonic acid

Studies on photoisomerization of 4,4'-diaminostilbene-2,2'-disulphonic acid for quality assurance by high-performance liquid chromatography 596(1992)127

1,2-Diamino-3,5-dibromobenzene

Determination of selenium by capillary gas chromatography after high-temperature derivatization with 1,2-diamino-3,5-dibromobenzene 598(1992)105

4,4'-Diaminodiphenylmethane

Solid-phase extraction and high-performance liquid chromatographic analysis of a toxic compound from γ -irradiated polyurethane 600(1992)93

1,12-Diaminododecane

Simple direct liquid introduction system usable as an interface for liquid chromatography-mass spectrometry on quadrupole and magnetic-sector mass spectrometers 554(1991)293

4,4'-Diaminophenylmethane

Identification of metabolites of 4,4'-diaminodiphenylmethane (methylene dianiline) using liquid chromatographic and mass spectrometric techniques 583(1992)63

4,5-Diaminophthalhydrazide

4,5-Diaminophthalhydrazide as a highly sensitive chemiluminescence derivatization reagent for α -dicarbonyl compounds in high-performance liquid chromatography 598(1992)203

Diaminopimelic acid

Ion-pair high-performance liquid chromatography of diaminopimelic acid in hydrolysates of physiological samples 582(1992)242

4,4'-Diaminostilbene-2,2'-disulphonic acid

Studies on photoisomerization of 4,4'-diaminostilbene-2,2'-disulphonic acid for quality assurance by high-performance liquid chromatography 596(1992)127

cis-Diamminedichloroplatinum

see Cisplatin

2,5-Diaryltetrahydrofuran

Determination of enantiomeric concentrations of a 2,5-diaryltetrahydrofuran (L-668,750), a platelet-activating factor receptor antagonist, in rat plasma using a chiral α_1 -acid glycoprotein high-performance liquid chromatographic column 578(1992)327

Diazepam

Determination of diazepam and nordazepam in milk and plasma in the presence of oxazepam and temazepam 564(1991)330

Quantitation by gas chromatography with selected-ion monitoring mass spectrometry of "natural" diazepam, N-desmethyldiazepam and oxazepam in normal human serum 579(1992)85

Simultaneous optimization of several chromatographic performance goals using Derringer's desirability function 586(1991)11

Dibenzothiophenes

Comments on the Note "Liquid chromatographic determination of planar aromatic sulphur compounds in crude oil", *J. Chromatogr.*, 475 (1989) 421 585(1991)376

Determination of sulfur components in light petroleum streams by high-resolution gas chromatography with chemiluminescence detection 589(1992)271

Gas chromatographic retention behaviour of dibenzothiophene derivatives on a smectic liquid crystalline polysiloxane stationary phase 590(1992)297

Dibenzothiophenes, chlorinated

Isolation of chlorinated dibenzothiophenes by high-performance thin-layer chromatography 553(1991)453

Dibutyl phthalate

Effect of silanol groups on heat-treated silicas by calcination and retention behaviour in high-performance liquid chromatography 556(1991)407

Characterization and determination of organic compounds in the mutagenic XAD-2 extracts of drinking water 557(1991)413

 α -Dicarbonyl compounds

4,5-Diaminophthalhydrazide as a highly sensitive chemiluminescence derivatization reagent for α -dicarbonyl compounds in high-performance liquid chromatography 598(1992)203

Dicarboxyl-*mono*-carnitine esters

Synthesis, characterisation and high-performance liquid chromatography of C₆-C₁₆ dicarboxyl-*mono*-coenzyme A and -*mono*-carnitine esters 570(1991)253

Dicarboxyl-*mono*-coenzyme A esters

Synthesis, characterisation and high-performance liquid chromatography of C₆-C₁₆ dicarboxyl-*mono*-coenzyme A and -*mono*-carnitine esters 570(1991)253

2,2'-Dichloroethyl sulphide

Specific and sensitive quantitation of 2,2'-dichlorodiethyl sulphide (sulphur mustard) in water, plasma and blood: application to toxicokinetic study in the rat after intravenous intoxication 583(1992)155

Dichloromethane

Effect of the sample solvent on band profiles in preparative liquid chromatography using non-aqueous reversed-phase high-performance liquid chromatography 588(1991)1

Dichlorprop

Application of high-performance thin-layer chromatography and automated multiple development for the identification and determination of pesticides in water 553(1991)489

Dichlorvos

High-performance liquid chromatographic method using ultraviolet detection for measuring metrifonate and dichlorvos levels in human plasma 573(1992)99

Diclofenac

Determination of diclofenac in plasma using a fully automated analytical system combining liquid-solid extraction with liquid chromatography 565(1991)401

Bioanalysis of diclofenac as its fluorescent carbazole acetic acid derivative by a post-column photoderivatization high-performance liquid chromatographic method 567(1991)175

Determination of diclofenac and its metabolites in plasma and cerebrospinal fluid by high-performance liquid chromatography with electrochemical detection 567(1991)425

Determination of diclofenac in plasma and urine by capillary gas chromatography-mass spectrometry with possible simultaneous determination of deuterium-labelled diclofenac 571(1991)87

Extractionless determination of diclofenac sodium in serum using reversed-phase high-performance liquid chromatography with fluorimetric detection 577(1992)185

High-performance liquid chromatographic determination of naproxen, ibuprofen and diclofenac in plasma and synovial fluid in man 578(1992)251

Diclofop

Determination of the herbicide diclofop in human urine 600(1992)327

Dicoumarol

Thermospray and particle beam liquid chromatographic-mass spectrometric analysis of coumarin anticoagulants 562(1991)31

(2*R*,3*R*)-Dicyclohexyl tartrate

(2*R*,3*R*)-Dicyclohexyl tartrate as a chiral mobile phase additive 585(1991)35

5,10-Dideaza-5,6,7,8-tetrahydrofolic acid

Bioanalysis of the investigational anti-tumour drug 5,10-dideaza-5,6,7,8-tetrahydrofolic acid by high-performance liquid chromatography with ultraviolet detection 576(1992)158

2',3'-Didehydro-3'-deoxythymidine

High-performance liquid chromatographic determination of 2',3'-didehydro-3'-deoxythymidine, a new anti-human immunodeficiency virus agent, in human plasma and urine 577(1992)151

2',3'-Dideoxycytidine

Thermospray liquid chromatographic-mass spectrometric analysis of anti-AIDS nucleosides: quantification of 2',3'-dideoxycytidine in plasma samples 577(1992)299

2',3'-Dideoxycytidine monophosphate diglyceride

Isolation of experimental anti-AIDS glycerophospholipids by micro-preparative reversed-phase high-performance liquid chromatography 590(1992)153

2',3'-Dideoxyinosine

High-performance liquid chromatographic analysis of 2',3'-dideoxyinosine in biological samples 563(1991)400

Dieldrin

Solid-matrix partition for separation of organochlorine pesticide residues from fatty materials 552(1991)241

Diethanolamide

Identification of products resulting from carbonyl sulphide-induced degradation of diethanolamine 587(1991)237

Diethenylbenzenes

Biotransformation of diethenylbenzenes. IV. A simple high-performance liquid chromatographic method for separation of urinary metabolites of 1,3-diethenylbenzene on analytical and semi-preparative scales 578(1992)215

O,O-Diethyldithiophosphoric acid

Direct gas chromatographic analysis of O,O-dimethyl- and O,O-diethyldithiophosphoric acids and identification of their by-products 589(1992)375

Diethyl ether

Experimental determination of adsorption isotherm data for the study of the surface energy distribution of various solid surfaces by inverse gas-solid chromatography 591(1992)233

Precision and accuracy of the gas-solid adsorption isotherms derived by the elution-by-characteristic-points method 591(1992)245

Validity of the model used to relate the energy distribution and the adsorption isotherm 591(1992)267

Diethyl ethoxymethylenemalonate

Amino acid analysis by high-performance liquid chromatography after derivatization with diethyl ethoxymethylenemalonate 591(1992)181

Di(2-ethylhexyl) phthalate

Characterization and determination of organic compounds in the mutagenic XAD-2 extracts of drinking water 557(1991)413

Behaviour of a common phthalate plasticizer (dioctyl phthalate) during the alkali- and/or acid-catalysed steps in an AOCS method for the preparation of methyl esters 587(1991)263

1,2-Diethyl-3-hydroxypyridin-4-one

High-performance liquid chromatographic determination of 1,2-diethyl-3-hydroxypyridin-4-one and its 2-(1-hydroxyethyl) metabolite in rat blood 573(1992)178

Diethyl phthalate

High-temperature continuous counter-current gas-liquid chromatography 590(1992)289

Diethylstilbestrol

Rapid screening method for the determination of diethylstilbestrol in edible animal tissue by column liquid chromatography with electrochemical detection 564(1991)477

General and selective isolation procedure for high-performance liquid chromatographic determination of anabolic steroids in tissues 588(1991)89

3,3-Difluoroalanine

High-performance ligand-exchange liquid chromatography of fluoro derivatives of alanine 600(1992)83

Diguanosine polyphosphates

Reversed-phase high-performance liquid chromatography of dinucleoside polyphosphates 563(1991)37

Dihydroajmaline

Monitoring of ajmaline in plasma with high-performance liquid chromatography 575(1992)87

Dihydroanthracene

Quantitative measurements via co-elution and dual-isotope detection by gas chromatography-mass spectrometry 587(1991)255

Dihydroartemisinin

Selective determination, in plasma, of artemether and its major metabolite, dihydroartemisinin, by high-performance liquid chromatography with ultraviolet detection 583(1992)131

(+)-5-(2,3-Dihydrobenzofuran-7-yl)-3-methyl-8-nitro-2,3,4,5-tetrahydro-1H-3-benzazepin-7-ol

Determination of (+)-5-(2,3-dihydrobenzofuran-7-yl)-3-methyl-8-nitro-2,3,4,5-tetrahydro-1H-3-benzazepin-7-ol (NNC 01-0687), a novel dopamine D-1 receptor antagonist, in plasma by solid-phase extraction and high-performance liquid chromatography 577(1992)366

Dihydrobunolol

Determination of (-)-bunolol and its metabolite, dihydro(-)-bunolol, in human aqueous humour by gas chromatography-negative ion chemical ionisation mass spectrometry 571(1991)101

10,11-Dihydro-10-hydroxycarbamazepine

Determination of the *R*-(-) and *S*-(+) enantiomers of the monohydroxylated metabolite of oxcarbazepine in human plasma by enantioselective high-performance liquid chromatography 581(1992)147

[3*R*-(+)-2,3-Dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzodiazepine-3-yl]-N-(3-methylphenyl)urea

Quantitation of a new cholecystokinin and gastrin receptor antagonist (L-365,260) in dog and rat plasma by high-performance liquid chromatography 576(1992)363

Dihydroorotate dehydrogenase

High-performance liquid chromatographic determination of dihydroorotate dehydrogenase of *Plasmodium falciparum* and effects of antimalarials on enzyme activity 582(1992)57

Dihydropyridine calcium antagonist

Simultaneous determination of a new dihydropyridine calcium antagonist (MPC-1304) and its metabolite in dog plasma by high-performance liquid chromatography with electrochemical detection 571(1991)277

Dihydroquercetin

High-performance liquid chromatographic method for the determination of mangiferin, likviritin and dihydroquercetin in rat plasma and urine 577(1992)371

Dihydroquinidine

High-performance liquid chromatographic method for the quantitation of quinidine, and selected quinidine metabolites 572(1991)159

Dihydrostreptomycin

Determination of aminoglycoside antibiotics by reversed-phase ion-pair high-performance liquid chromatography coupled with pulsed amperometry and ion spray mass spectrometry 591(1992)195

5,6-Dihydrothymidine

Reversed-phase high-performance liquid chromatography-thermospray mass spectrometry of radiation-induced decomposition products of thymine and thymidine 593(1992)133

2,5-Dihydroxybenzohydrazide

Electrochemical and chromatographic properties of selected hydrazine and hydrazide derivatives of carbonyl compounds 585(1991)225

2,4-Dihydroxybenzoic acid

Extension of the electrostatic retention model of reversed-phase ion-pair high-performance liquid chromatography to include the effect of the eluent pH 552(1991)13

3,5-Dihydroxybenzoic acid

Effect of the addition of viscous matrices to the mobile phase on chromatographic performance in liquid chromatography-fast atom bombardment mass spectrometry 554(1991)61

3,4-Dihydroxybenzylamine

Optimization of high-performance liquid chromatographic assay for catecholamines. Determination of optimal mobile phase composition and elimination of species-dependent differences in extraction recovery of 3,4-dihydroxybenzylamine 574(1992)213

***p*-Dihydroxyborylphenylalanine**

Studies of the electrophoretic behaviour of *p*-dihydroxyborylphenylalanine and related compounds by the three-spot method 585(1991)369

Dihydroxycarbamazepine

Fast liquid chromatography for the determination of drugs in plasma and combination with liquid–solid extraction in a fully automated system 573(1992)59

Automated microanalysis of carbamazepine and its epoxide and *trans*-diol metabolites in plasma by column liquid chromatography 573(1992)65

***cis*-1,2-Dihydroxycyclohexyl methyl ketone**

Synthesis of (*1R-trans*)-N,N'-1,2-cyclohexylenebisbenzamide-oligodimethylsiloxane copolymers for use as chiral stationary phases for capillary supercritical fluid chromatography 594(1992)283

5,6-Dihydroxy-5,6-dihydrothymidine

Reversed-phase high-performance liquid chromatography–thermospray mass spectrometry of radiation-induced decomposition products of thymine and thymidine 593(1992)133

Dihydroxy long-chain bases

Fragmentation pathways of O-trimethylsilyl ethers of dihydroxy long-chain bases analysed by linked-scan mass spectrometry 562(1991)119

2,4-Dihydroxy-4-methyl-5-nitrobenzophenone

Determination of the catechol-O-methyltransferase inhibitor Ro 40-7592 in human plasma by high-performance liquid chromatography with coulometric detection 593(1992)63

3,4-Dihydroxyphenylacetic acid

Optimization of high-performance liquid chromatographic assay for catecholamines. Determination of optimal mobile phase composition and elimination of species-dependent differences in extraction recovery of 3,4-dihydroxybenzylamine 574(1992)213

Simultaneous measurement of serotonin, catecholamines and their metabolites in cat and human plasma by *in vitro* microdialysis–microbore high-performance liquid chromatography with amperometric detection 582(1992)19

3,4-Dihydroxyphenylalanine

Determination of free DOPA and 3-O-methyl-DOPA in human plasma by high-performance liquid chromatography with electrochemical detection 553(1991)265

Determination of L-3,4-dihydroxyphenylalanine in biological fluids and tissues 565(1991)1

Determination of urinary vanillic acid and plasma dihydroxyphenylalanine as markers of non-secreting neuroblastoma by high-performance liquid chromatography with electrochemical detection 567(1991)331

High-performance liquid chromatographic procedure for the simultaneous determination of aromatic L-amino acid decarboxylase activity towards 3,4-dihydroxyphenylalanine and 5-hydroxytryptophan 567(1991)381

Method for measuring endogenous 3-O-methyl-dopa in urine and plasma 568(1991)45

Optimization of high-performance liquid chromatographic assay for catecholamines. Determination of optimal mobile phase composition and elimination of species-dependent differences in extraction recovery of 3,4-dihydroxybenzylamine 574(1992)213

High-performance liquid chromatography with electrochemical detection for the determination of levodopa, catecholamines and their metabolites in rat brain dialysates 575(1992)207

4,5-Dihydroxy-3-(*p*-sulphophenylazo)-2,7-naphthalene-disulphonic acid

Isotachopheretic separation of minor components from a matrix component in the case of strong electrolytes 585(1991)297

Dihydroxytryptamines

Improved method for the determination of 5,6-dihydroxytryptamine and 5,7-dihydroxytryptamine in tissue using high-performance liquid chromatography with electrochemical detection 574(1992)137

3,3'-Diindolylmethane

Identification of enzymatic degradation products from synthesized glucobrassicin by gas chromatography–mass spectrometry 586(1991)166

Diiodofluorescein

Fluorescence detector cell for use in an integrated electrically driven separation system 598(1992)115

Diisopropylbenzenes

High-performance liquid chromatographic determination of products of autoxidation of isopropylbenzene derivatives 588(1991)33

Diisopropylfluorophosphate

Supercritical fluid extraction of chemical warfare agent simulants from soil 600(1992)273

Diisopropylmethylphosphonate

Supercritical fluid extraction of chemical warfare agent simulants from soil 600(1992)273

1,4-Diketo-2,3-dihydroxyionapalene

High-performance liquid chromatographic determination of the 1,4-diketo and 1,4-diketo-2,3-dihydroxy metabolites of ionapalene in rat urine 565(1991)339

1,4-Diketolonapalene

High-performance liquid chromatographic determination of the 1,4-diketo and 1,4-diketo-2,3-dihydroxy metabolites of ionapalene in rat urine 565(1991)339

Diltiazem

Determination of diltiazem hydrochloride enantiomers in dog plasma using chiral stationary-phase liquid chromatography 564(1991)338

Simple and sensitive high-performance liquid chromatographic method for the determination of diltiazem and six of its metabolites in human plasma 582(1992)203

Dimethadione

Improved method for the determination of trimethadione and its demethylated metabolite, dimethadione, in human serum by gas chromatography 584(1992)267

Dimethindene

Enantioselective determination of dimethindene in urine after oral administration of racemic dimethindene 567(1991)229

Dimethoate

Removal of organic pollutants from aqueous solution. V. Comparative study of the extraction, recovery and chromatographic separation of some organic insecticides using unloaded polyurethane foam columns 552(1991)371

Dimethylalkynes

Capillary gas chromatography of C₅-C₁₃ branched alkynes on squalane and liquid crystal stationary phases 557(1991)241

N-[4-[2-(Dimethylamino)ethoxy]benzyl]-3,4-dimethoxybenzamide hydrochloride

Simultaneous determination of a new gastrointestinal prokinetic agent (HSR-803) and its metabolites in human serum and urine by high-performance liquid chromatography using automated column-switching 576(1992)174

3-[2-(N,N-Dimethylaminomethyl)phenylthio]phenol

Mass spectrometric structure determination of metabolites of 3-[2-(N,N-dimethylaminomethyl)phenylthio]phenol 562(1991)289

N-(2-Dimethylaminopropyl)iminodibenzyl

Influence of chemical structure of tricyclic tertiary dimethylamines on chiral separation by reversed-phase high-performance liquid chromatography after derivatization with (-)-menthylchloroformate 553(1991)365

4-(N,N-Dimethylaminosulphonyl)-7-N-cadaverino-2,1,3-benzoxadiazole

Precolumn fluorescence tagging reagent for carboxylic acids in high-performance liquid chromatography: 4-substituted-7-aminoalkylamino-2,1,3-benzoxadiazoles 588(1991)61

4-(N,N-Dimethylaminosulphonyl)-7-N-piperazino-2,1,3-benzoxadiazole

Precolumn fluorescence tagging reagent for carboxylic acids in high-performance liquid chromatography: 4-substituted-7-aminoalkylamino-2,1,3-benzoxadiazoles 588(1991)61

Dimethylanilines

Chromatography on Poly-RP and its cyano and diol derivatives using both polar and non-polar solvent systems 591(1992)89

3-[2-(4,7-Dimethylbenzoxayolyl)methyl]amino-5-ethyl-6-methyl-1H-pyridin-2-one

High-performance liquid chromatographic procedure for the determination of a non-nucleoside HIV-1 reverse transcriptase inhibitor in human plasma 577(1992)129

2,2-Dimethylbutane

Study of microporous carbons by gas chromatographic determination of heats of physisorption 555(1991)183

N,N-Dimethylcyclohexylamine

Liquid membrane work-up of blood plasma samples applied to gas chromatographic determination of aliphatic amines 573(1992)191

O,O-Dimethyldithiophosphoric acid

Direct gas chromatographic analysis of O,O-dimethyl- and O,O-diethyldithiophosphoric acids and identification of their by-products 589(1992)375

N,N-Dimethylethylamine

Liquid membrane work-up of blood plasma samples applied to gas chromatographic determination of aliphatic amines 573(1992)191

N,N-Dimethylformamide

Elution behaviour of polyamic acid and polyamide-imide in size-exclusion chromatography 588(1991)195

3,5-Dimethylhippuric acid

High-performance liquid chromatographic determination of 3,5-dimethylhippuric acid in the occupational exposure to trimethylbenzenes 577(1992)180

3,7-Dimethyl-1-(5'-hydroxyethyl)xanthine

Determination of pentoxifylline and its metabolites in human plasma by high-performance liquid chromatography with solid-phase extraction 575(1992)101

Dimethylmethylphosphonate

Supercritical fluid extraction of chemical warfare agent simulants from soil 600(1992)273

3-(N,N-Dimethylpalmitylammonium)propanesulphonate

High-performance liquid chromatography on dynamically modified silica. IX. Modification of silica with 3-(N,N-dimethylpalmitylammonium) propanesulphonate for reversed-phase chromatography 556(1991)353

Dimethylphenols

Chromatography on Poly-RP and its cyano and diol derivatives using both polar and non-polar solvent systems 591(1992)89

N-[(2,6-Dimethylphenylcarbamoyl)methyl]succinamic acid

Simultaneous determination of nefiracetam and its metabolites by high-performance liquid chromatography 575(1992)261

Dimethyl phthalate

Effect of silanol groups on heat-treated silicas by calcination and retention behaviour in high-performance liquid chromatography 556(1991)407

High-temperature continuous counter-current gas-liquid chromatography 590(1992)289

Dimethylpyridines

Chromatography on Poly-RP and its cyano and diol derivatives using both polar and non-polar solvent systems 591(1992)89

5-Dimethylsulphamoyl-6,7-dichloro-2,3-dihydrobenzofuran-2-carboxylic acid

High-performance liquid chromatographic method for simultaneous determination of enantiomers of 5-dimethylsulphamoyl-6,7-dichloro-2,3-dihydrobenzofuran-2-carboxylic acid and its N-monodemethyl metabolite in monkey plasma and urine after chiral derivatization 564(1991)235

Dimethyl sulphide

Determination of dimethyl sulphide in blood and adipose tissue by headspace gas analysis 565(1991)453

Dimethyl sulphoxide

Determination of 7-methylbenz[*c*]acridines by capillary gas chromatography with electron-capture detection 596(1992)233

Dimetridazole

Determination of some drugs by micellar electrokinetic capillary chromatography. The pseudo-effective mobility as parameter for screening 585(1991)123

Dinitramine

High-performance liquid chromatographic determination of dinitroaniline herbicides in soil and water 585(1991)164

N-(3,5-Dinitrobenzoyl)leucine

Effect of superfluous remote polar functionality on chiral recognition 589(1992)45

2,4-Dinitrophenylhydrazine

Analysis of keto acids as their methyl esters of 2,4-dinitrophenylhydrazone derivatives by gas chromatography and gas chromatography-mass spectrometry 587(1991)247

2,4-Dinitrophenylhydrazones

High-performance liquid chromatographic determination of 2-furaldehyde in spirits 590(1992)235

4,4'-Dinitrostilbene-2,2'-disulphonic acid

Studies on photoisomerization of 4,4'-diaminostilbene-2,2'-disulphonic acid for quality assurance by high-performance liquid chromatography 596(1992)127

20,21-Dinoreburnamenine

Determination of the enantiomers of a novel 20,21-dinoreburnamenine derivative in rat plasma and brain by high-performance liquid chromatography using a chiral stationary phase 575(1992)315

Dinucleoside polyphosphates

Reversed-phase high-performance liquid chromatography of dinucleoside polyphosphates 563(1991)37

Diocaine

Determination of limiting ionic mobilities and dissociation constants of some local anaesthetics 596(1992)265

Diocetyl phthalate

Effect of silanol groups on heat-treated silicas by calcination and retention behaviour in high-performance liquid chromatography 556(1991)407

Behaviour of a common phthalate plasticizer (diocetyl phthalate) during the alkali- and/or acid-catalysed steps in an AOCS method for the preparation of methyl esters 587(1991)263

Diolepoxide polyaromatic hydrocarbon DNA adducts

Superoxide chemical transformation of diolepoxide polyaromatic hydrocarbon DNA adducts. Determination of benzo[*a*]pyrene-*r*-7,*t*-8,9,*c*-10-tetrahydroretrol by gas chromatography 588(1991)273

Diols

Gas chromatographic-mass spectrometric analysis of diols and tetrols from reactions of polycyclic aromatic hydrocarbon epoxides with hemoglobin 562(1991)563

Prediction of retention indexes. III. Silylated derivatives of polar compounds 586(1991)113

Synthesis of (1*R*-*trans*)-*N,N'*-1,2-cyclohexylenebisbenzamide-oligodimethylsiloxane copolymers for use as chiral stationary phases for capillary supercritical fluid chromatography 594(1992)283

Dioxins

Comparison of extraction methods for polychlorinated dibenzo-*p*-dioxins and dibenzofurans in fly ash using gas chromatography-mass spectrometry 589(1992)265

Gas chromatographic elution patterns of chlorinated dioxins versus column polarity 594(1992)269

Automated clean-up procedure for the gas chromatographic-high-resolution mass spectrometric determination of polychlorinated dibenzo-*p*-dioxins and dibenzofurans in milk 595(1992)289

Dipalmitoylphosphatidylcholine

High-performance liquid chromatography post-column derivatization with fluorescence detection to study the influence of ambroxol on dipalmitoylphosphatidylcholine levels in rabbit eustachian tube washings 553(1991)1

Dipeptides

High-performance liquid chromatography of amino acids and dipeptides on new ion exchangers of the HEMA series 552(1991)439

Diperodone

Determination of limiting ionic mobilities and dissociation constants of some local anaesthetics 596(1992)265

Diphenyl ethers, bromine-containing

Use of mass spectrometry for the detection and identification of bromine-containing diphenyl ethers 586(1991)360

4,4'-Diphenylmethanediisocyanate

Automated high-performance liquid chromatographic and size-exclusion chromatographic sample preparation by means of a robotic workstation 592(1992)249

4,6-Diphenylthieno[3,4-*d*]-1,3-dioxol-2-one-5,5-dioxide

Activation of matrices by 4,6-diphenylthieno[3,4-*d*]-1,3-dioxol-2-one-5,5-dioxide. High-performance liquid affinity chromatographic separations 589(1992)101

Dipipanone

Retention reproducibility of basic drugs in high-performance liquid chromatography on a silica column with a methanol-high-pH buffer eluent. Changes in selectivity with the age of the stationary phase 592(1992)85

Diprafenone

Direct separation of the enantiomers of propafenone, diprafenone and their major metabolites by high-performance liquid chromatography on modified cellulose and amylose chiral stationary phases 585(1991)329

Dipyridamole

Effect of stationary phase structure on retention and selectivity of restricted-access reversed-phase packing materials 558(1991)19

Sensitive determination of free and plasma protein-bound dipyridamole by high-performance liquid chromatography 565(1991)511

Dipyridylsulphides

Chromatographic behaviour of dipyridylsulphides. Relationship between log *k'* values and structure by reversed-phase high-performance liquid chromatography 595(1992)334

Diquat

Electrodialytic sample treatment coupled on-line with column liquid chromatography for the determination of basic and acidic compounds in environmental samples 600(1992)163

Diradylglycerobenzoates

Improved determination of individual molecular species of phosphatidylcholine in biological samples by high-performance liquid chromatography with internal standards 593(1992)139

Dirithromycin

Determination of dirithromycin purity and related substances by high-performance liquid chromatography 594(1992)203

Disodium dihydrogen(cycloheptylamino)methylenebisphosphate monohydrate

Determination of a new bisphosphonate, YM175, in plasma, urine and bone by high-performance liquid chromatography with electrochemical detection 584(1992)213

Disopyramide

Simplified, rapid and inexpensive extraction procedure for a high-performance liquid chromatographic method for determination of disopyramide and its main metabolite mono-N-dealkylated disopyramide in serum 572(1991)339

Automated determination of disopyramide and N-monodealkylidisopyramide in plasma by reversed-phase liquid chromatography with a column switching system 574(1992)365

Disulphides

Micro liquid chromatography with fluorescence detection of thiols and disulphides 553(1991)433

Diterpenes

Determination of sterols and diterpenoids from brown algae (Cystoseiraceae) 588(1991)99

Determination of taxol by high-performance liquid chromatography-thermospray mass spectrometry 594(1992)153

Diuretics

Approach to the analysis of diuretics and masking agents by high-performance liquid chromatography-mass spectrometry in doping control 562(1991)723

Screening for diuretics in human urine by gas chromatography-mass spectrometry with derivatisation by direct extractive alkylation 563(1991)257

Diuretic screening in human urine by gas chromatography–mass spectrometry: use of a macroreticular acrylic copolymer for the efficient removal of the coextracted phase-transfer reagent after derivatization by direct extractive alkylation 581(1992)57

High-performance liquid chromatographic determination of diuretics in urine by micellar liquid chromatography 582(1992)189

trans-1,2-Divinylcyclohexane

Ligand-exchange chromatography of alkenes on stationary phases containing palladium(II) complexes. Enantiomeric separation of *trans*-1,2-divinylcyclohexane 585(1991)309

DNA

Purification of common bean chloroplast DNA by DEAE cellulose column chromatography 558(1991)318

Low-cost, high-sensitivity laser-induced fluorescence detection for DNA sequencing by capillary gel electrophoresis 559(1991)237

Capillary electrophoresis of DNA in entangled polymer solutions 559(1991)257

Analysis of DNA restriction fragments and polymerase chain reaction products towards detection of the AIDS (HIV-1) virus in blood 559(1991)267

Effect of temperature on the separation of DNA restriction fragments in capillary gel electrophoresis 559(1991)285

Identification of DNA molecules by pre-column hybridization using capillary electrophoresis 559(1991)295

5-Methylcytosine content of DNA in blood, synovial mononuclear cells and synovial tissue from patients affected by autoimmune rheumatic diseases 566(1991)481

DNA typing in hereditary disease (Review) 569(1991)421

High-resolution gel filtration of the ecdysteroid receptor–DNA complex —an alternative to the electrophoretic mobility shift assay 587(1991)11

Isolation of multiple protein factors involved in ribosomal DNA transcription 587(1991)25

Separation techniques for biotechnology in the 1990s 592(1992)3

DNA electrophoresis in uncross-linked polyacrylamide solution, studied by epifluorescence microscopy 596(1992)141

Biospecific interaction analysis using surface plasmon resonance detection applied to kinetic, binding site and concentration analysis 597(1992)397

High-performance affinity chromatography of DNA. II. Porosity effects 600(1992)221

Purification of DNA-derived deoxynucleotides from leukocytes involving nuclease elution of an ion-exchange column 600(1992)229

Capillary gel electrophoresis of single-stranded DNA fragments with UV detection 600(1992)289

DNA adducts

Enzymatic shot-gun 5'-phosphorylation and 3'-sister phosphate exchange: a two-dimensional thin-layer chromatographic technique to measure DNA deoxynucleotide modification 574(1992)41

Superoxide chemical transformation of diepoxide polyaromatic hydrocarbon DNA adducts. Determination of benzo[*a*]pyrene-*r*-7,*t*-8,9,*c*-10-tetrahydrotetrol by gas chromatography 588(1991)273

DNA dinucleotides

Optimization of the separation of the Rp and Sp diastereomers of phosphate-methylated DNA and RNA dinucleotides 596(1992)59

DNA restriction fragments

Anion-exchange chromatography of DNA restriction fragments 555(1991)109

DNA topoisomerases

Use of anion-exchange chromatography and chromatofocusing to reveal the structural and functional heterogeneity of topoisomerase II in a HL-60 cell line resistant to multi-drug treatment 587(1991)3

Dobutamine

Simultaneous determination of catecholamines and dobutamine in human plasma and urine by high-performance liquid chromatography with fluorimetric detection 583(1992)236

closo-Dodecahydro-dodecaborate anion

Chromatographic behaviour of the *closo*-[B₁₂H₁₂]²⁻ derivatives on hydroxyethylmethacrylate gels 588(1991)201

Dodecane

Quantitative resolution of severely overlapping gas chromatographic peaks. Isothermal and temperature-programmed operation 556(1991)305

Domoic acid

Purification of domoic acid from toxic blue mussels (*Mytilus edulis*) and phytoplankton 568(1991)393

Dopa

see 3,4-Dihydroxyphenylalanine

Dopamine

Improved measurement of urinary catecholamines by liquid-liquid extraction, derivatization and high-performance liquid chromatography with fluorimetric detection 563(1991)348

Evaluation of a simple plasma catecholamine extraction procedure prior to high-performance liquid chromatography and electrochemical detection 564(1991)55

Determination of γ -glutamyl conjugates of monoamines by means of high-performance liquid chromatography with electrochemical detection and application to gastropod tissues 567(1991)49

Measurement of plasma catecholamines by high-performance liquid chromatography with electrochemical detection in intensive care patients after dobutamine infusion 567(1991)240

Analysis of plasma catecholamines by high-performance liquid chromatography with fluorescence detection: simple sample preparation for pre-column fluorescence derivatization 567(1991)351

Simultaneous determination of free catecholamines and epinine and estimation of total epinine and dopamine in plasma and urine by high-performance liquid chromatography with fluorimetric detection 574(1992)109

Optimization of high-performance liquid chromatographic assay for catecholamines. Determination of optimal mobile phase composition and elimination of species-dependent differences in extraction recovery of 3,4-dihydroxybenzylamine 574(1992)213

γ -Aminobutyric acid, catecholamine and indoleamine determinations from the same brain region by high-performance liquid chromatography with electrochemical detection 579(1992)334

Determination of dopamine and two of its acidic metabolites by high-performance liquid chromatography with electrochemical detection: a retrospective article (Chromatography Classic) 582(1992)1

Simultaneous measurement of serotonin, catecholamines and their metabolites in cat and human plasma by *in vitro* microdialysis-microbore high-performance liquid chromatography with amperometric detection 582(1992)19

Simultaneous determination of catecholamines and dobutamine in human plasma and urine by high-performance liquid chromatography with fluorimetric detection 583(1992)236

Determination of catecholamines in urine by liquid chromatography and electrochemical detection after on-line sample purification on immobilized boronic acid 593(1992)185

Dopamine agonists

Optimization of the direct enantiomeric separation by high-performance liquid chromatography of the D-2 dopamine agonist N-0437 using response surface methodology and the multi-criteria decision-making method 558(1991)333

Direct analysis of the dopamine agonist (-)-2-(N-propyl-N-2-thienylethylamino)-5-hydroxytetralin hydrochloride in plasma by high-performance liquid chromatography using two-dimensional column switching 565(1991)277

Doping agents

see also Special issue index

Doping agents

Review. Immunoaffinity chromatography, its applicability and limitations in multi-residue analysis of anabolizing and doping agents 564(1991)363

Dothiepin

Measurement of dothiepin and its major metabolites in plasma by high-performance liquid chromatography 581(1992)152

Dothiepin-S-oxide

Measurement of dothiepin and its major metabolites in plasma by high-performance liquid chromatography 581(1992)152

Doxazosin

Validation of a solid-phase extraction high-performance liquid chromatographic assay for doxazosin 566(1991)234

Doxorubicin

Determination of adriamycin in plasma and tissue biopsies 564(1991)322

Simple and sensitive quantification of anthracyclines in mouse atrial tissue using high-performance liquid chromatography and fluorescence detection 570(1991)209

Fully automated determination of a new anthracycline N-*l*-leucyldoxorubicin and six metabolites in plasma by high-performance liquid chromatography with on-line sample handling 574(1992)273

Bioanalysis of some anthracyclines in human plasma by capillary electrophoresis with laser-induced fluorescence detection 574(1992)327

Determination of 3'-deamino-3'-[2(*S*)-methoxy-4-morpholinyl]doxorubicin, a new morpholinyl anthracycline, in plasma by high-performance liquid chromatography with fluorescence detection 578(1992)309

Doxorubicinol

Simple and sensitive quantification of anthracyclines in mouse atrial tissue using high-performance liquid chromatography and fluorescence detection 570(1991)209

Fully automated determination of a new anthracycline N-*l*-leucyldoxorubicin and six metabolites in plasma by high-performance liquid chromatography with on-line sample handling 574(1992)273

Doxorubicinol aglycone

Fully automated determination of a new anthracycline N-*l*-leucyldoxorubicin and six metabolites in plasma by high-performance liquid chromatography with on-line sample handling 574(1992)273

Doxycycline

Determination of metacycline and related substances by column liquid chromatography on poly(styrene-divinylbenzene) 586(1991)61

Isolation of doxycycline, 6-epidoxycycline and 2-acetyl-2-decarboxamidometacycline from commercial metacycline by preparative column liquid chromatography on silica gel 586(1991)67

Drinking water

Recent methods for the determination of volatile and non-volatile organic compounds in natural and purified drinking water 562(1991)469

Drug metabolites

Scanning thin-layer chromatography-liquid secondary ion mass spectrometry and its application for investigation of drug metabolites 562(1991)99

Drugs

On-line measurements of ^{13}C enrichments in rat breath. Non-invasive method for *in vivo* study of drug enzymatic induction 562(1991)341

Formation of formaldehyde adducts from various drugs by use of methanol in a toxicological screening procedure with gas chromatography-mass spectrometry 563(1991)73

Automated screening procedure using gas chromatography-mass spectrometry for identification of drugs after their extraction from biological samples 565(1991)207

Allosteric and competitive displacement of drugs from human serum albumin by octanoic acid, as revealed by high-performance liquid affinity chromatography, on a human serum albumin-based stationary phase 577(1992)305

Fast systematic approach for the determination of drugs in biological fluids by fully automated high-performance liquid chromatography with on-line solid-phase extraction and automated cartridge exchange. Application to cebaracetam in human urine 577(1992)335

Systematic toxicological analysis of drugs and their metabolites by gas chromatography-mass spectrometry (Review) 580(1992)3

Drug detection in hair by chromatographic procedures (Review) 580(1992)125

Coupling of microdialysis with capillary electrophoresis: a new approach to the study of drug transfer between two compartments of the body in freely moving rats 581(1992)257

Determination of drugs from urine by on-line immunoaffinity chromatography-high-performance liquid chromatography-mass spectrometry 582(1992)103

DuP 532

High-performance liquid chromatographic determination of angiotensin II receptor antagonists in human plasma and urine. I. DuP 532 (L-694,492) 581(1992)165

Dyes

Analysis of dyes extracted from textile fibers by thermospray high-performance liquid chromatography-mass spectrometry 586(1991)73

Comparison of commercially available thin-layer chromatography plates with mixtures of dyes, analgesics and phenols 595(1992)368

Immunoenzymatic assay of dyes currently used in affinity chromatography for protein purification 597(1992)315

Fluorescence detector cell for use in an integrated electrically driven separation system 598(1992)115

Dynorphins

Approach to studying proteinase specificity by continuous-flow fast atom bombardment mass spectrometry and high-performance liquid chromatography combined with photodiode-array ultraviolet detection 554(1991)83

E**E2020**

Determination of enantiomers of 1-benzyl-4-[(5,6-dimethoxy-1-indanon)-2-yl]methylpiperidine hydrochloride (E2020), a centrally acting acetylcholine esterase inhibitor, in plasma by liquid chromatography with fluorometric detection 577(1992)95

Ecdysteroid receptor

High-resolution gel filtration of the ecdysteroid receptor-DNA complex—an alternative to the electrophoretic mobility shift assay 587(1991)11

Ecdysteroids

Phenylboronic acid as a versatile derivatization agent for chromatography of ecdysteroids 596(1992)271

Ecgonine methyl ester

Determination of cocaine, benzoylecgonine and ecgonine methyl ester in plasma by reversed-phase high-performance liquid chromatography 570(1991)412

Rapid assay of cocaine, opiates and metabolites by gas chromatography-mass spectrometry (Review) 580(1992)43

Echinonone

Optimization of an isocratic high-performance liquid chromatographic separation of carotenoids 589(1992)171

Evaluation of reversed-phase liquid chromatographic columns for recovery and selectivity of selected carotenoids 595(1992)89

Eicosanoids

Analysis of some eicosanoids by continuous-flow fast atom bombardment mass spectrometry 562(1991)59

Profiling of eicosanoids in inflamed gall bladder wall by gas chromatography with selected-ion monitoring 571(1991)1

In vivo transformation of arachidonic acid into 12-hydroxy-5,8,10,14-eicosatetraenoic acid by human nasal mucosa 575(1992)143

Elastase

Novel method of purification of human leukocytic elastase using adsorption on a high-performance liquid chromatography gel permeation column 567(1991)57

Elastin

Separation of elastin cross-links as phenylisothiocyanate derivatives 553(1991)93

Quantitative analysis of collagen and elastin cross-links using a single-column system 582(1992)49

Ellagic acid

Determination of tannic acid and its phenolic metabolites in biological fluids by high-performance liquid chromatography 577(1992)77

Endin

Solid-matrix partition for separation of organochlorine pesticide residues from fatty materials 552(1991)241

Endoglucanase

Analysis of pH-dependent protein interactions with gel filtration medium 591(1992)121

Endopeptidases

Approach to studying proteinase specificity by continuous-flow fast atom bombardment mass spectrometry and high-performance liquid chromatography combined with photodiode-array ultraviolet detection 554(1991)83

 β -Endorphin

Measurement of β -endorphin in human plasma by high-performance liquid chromatography with electrochemical detection: validation of a method employing the simultaneous purification and concentration of β -endorphin 577(1992)239

High-performance liquid chromatography of amino acids, peptides and proteins. CXV. Thermodynamic behaviour of peptides in reversed-phase chromatography 593(1992)103

Reversed-phase chromatographic behaviour of β -endorphin: evidence of conformational change 595(1992)125

Endothelin-converting enzyme

Determination of endothelin-converting enzyme activity by high-performance liquid chromatography-on-line radioactive flow monitoring 574(1992)344

Endothelin receptor

Affinity chromatographic purification of bovine lung endothelin receptor using biotinylated endothelin and avidin-agarose 597(1992)331

Endotoxins

High-performance liquid chromatographic method to resolve and determine lipopolysaccharide sub-groups of *Escherichia coli* endotoxin in isolated perfused rat liver perfusate 574(1992)205

Specific assay for endotoxin using immobilized histidine, *Limulus* amoebocyte lysate and a chromogenic substrate 597(1992)415

Enkephalins

Discriminative affinity labelling of opioid receptors by enkephalin and morphiceptin analogues containing 3-nitro-2-pyridinesulphenyl-activated thiol residues 597(1992)425

Enprostil acid

Determination of femtomole/milliliter concentrations of enprostil acid in human plasma using high-performance liquid chromatography-laser-induced fluorescence detection 567(1991)195

Enzyme-linked immunoglobulins

Electrophoretic analysis of enzyme-linked immunoglobulins and their clinical significance (Review) 569(1991)347

Enzymes

see also Special issue index

Enzymes

see also Proteins

Measurement of enzyme activities of cytochrome P-450 isoenzymes by high-performance liquid chromatographic analysis of products 553(1991)233

Approach to studying proteinase specificity by continuous-flow fast atom bombardment mass spectrometry and high-performance liquid chromatography combined with photodiode-array ultraviolet detection 554(1991)83

Use of methyl oxime derivatives to enhance structural information in thermospray high-performance liquid chromatography-mass spectrometry. Analysis of linoleic acid lipoxygenase metabolites in maize embryos 554(1991)155

Dye-ligand affinity partitioning of lactate dehydrogenase isoenzymes 557(1991)325

- Diffusion of proteins in the chromatographic gel AcA-34 558(1991)71
- Extraction of xanthine oxidase from milk by counter-current distribution in an aqueous two-phase system 558(1991)357
- Detection of enzyme activity in fractions collected from free solution capillary electrophoresis of complex samples 559(1991)325
- Reversible reaction via a carbanion intermediate in the elimination of ammonia from L-histidine catalysed by histidine ammonia-lyase 562(1991)363
- Use of L-prolyl-L-leucylglycinamide (MIF-1) for the high-performance liquid chromatographic determination of proline iminopeptidase activity in rat liver 563(1991)142
- Simultaneous assay of the activities of two key enzymes in cholesterol metabolism by gas chromatography-mass spectrometry 565(1991)53
- High-performance liquid chromatography in enzymatic analysis. Opening lecture 566(1991)275
- Application of high-performance liquid chromatography to the study of biogenic amine-related enzymes (Review) 566(1991)287
- Construction of kinetic models to understand metabolism *in vivo* 566(1991)309
- Automated high-performance liquid chromatographic assay of enzymatic activities 566(1991)327
- Carrier membrane as a stationary phase for affinity chromatography and kinetic studies of membrane-bound enzymes 566(1991)341
- Purification of carbonic anhydrase isoenzymes by high-performance affinity chromatography and hydrophobic interaction chromatography 566(1991)351
- High-performance liquid chromatographic assay of human lymphocyte kynureninase activity levels 566(1991)369
- High-performance liquid chromatographic determination of D-amino acid oxidase activity 566(1991)377
- High-performance liquid chromatographic assays for protoporphyrinogen oxidase and ferrochelatase in human leucocytes 566(1991)383
- Determination of ¹²⁵I-labelled thyroxine glucuronide by reversed-phase high-performance liquid chromatography using on-line radiochemical detection to determine UDPglucuronosyltransferase activity 566(1991)415
- Radio-detection high-performance liquid chromatographic enzyme assay for inhibitors of fungal sterol Δ^{14} -reductase 566(1991)435
- High-performance liquid chromatographic assay of L- α -glycerophosphorylcholine using a two-step enzymic conversion 566(1991)445
- p*-Nitrophenyllaurate: a substrate for the high-performance liquid chromatographic determination of lipase activity 566(1991)453
- High-performance liquid chromatographic investigation of product formation in the horseradish peroxidase-enhanced chemiluminescence of luminol with different enhancers 566(1991)461
- Determination of biocatalyst consumption in an aminopeptidase process using automated sample preparation and high-performance liquid chromatography 566(1991)493
- Novel method of purification of human leukocytic elastase using adsorption on a high-performance liquid chromatography gel permeation column 567(1991)57
- Fluorimetric determination of hepatic δ -aminolevulinic acid synthase activity by high-performance liquid chromatography 567(1991)65
- High-performance liquid chromatographic determination, with ultraviolet detection, of S-adenosyl-L-methionine and S-adenosyl-L-homocysteine in rat tissues and simultaneously of normetanephrine and metanephrine for phenylethanolamine-N-methyltransferase or catechol-O-methyltransferase activities 567(1991)254
- High-performance liquid chromatographic procedure for the simultaneous determination of aromatic L-amino acid decarboxylase activity towards 3,4-dihydroxyphenylalanine and 5-hydroxytryptophan 567(1991)381
- High-performance liquid chromatographic determination of histamine N-methyltransferase activity 567(1991)459
- Human plasma lecithin:cholesterol acyltransferase. Preparation and use of immobilized *p*-aminophenylarsenoxide as a catalytic site-directed covalent ligand in enzyme purification 568(1991)69
- Purification of phospholipase C by hydrophobic interaction affinity chromatography 568(1991)85
- Purification and some properties of phospholipase C from *Achromobacter xylosoxidans* 568(1991)315
- Electrophoretic phenotyping of erythrocyte enzymes (Review) 569(1991)297
- Electrophoresis of serum isoenzymes and proteins following acute myocardial infarction (Review) 569(1991)323
- Electrophoretic analysis of enzyme-linked immunoglobulins and their clinical significance (Review) 569(1991)347

- 6-Pyruvoyl tetrahydropterin synthase assay in extracts of cultured human cells using high-performance liquid chromatography with fluorescence detection of biopterin. 570(1991)43
- High-performance liquid chromatographic detection of pitfalls in porphobilinogen deaminase determination 570(1991)51
- Isocratic high-performance liquid chromatographic measurement of optimal 5 α -steroid reductase activity in Hep-G2 cells 570(1991)293
- Separation and characterization of sialic acid-containing salivary-type amylase from patients' sera with immunoglobulin A-type myeloma 571(1991)61
- Use of high-performance liquid chromatography for assay of glutamic acid decarboxylase. Its limitation in use for *post-mortem* brain 571(1991)235
- Assay of sialyltransferase activity by reversed-phase ion-pair high-performance liquid chromatography 573(1992)23
- Rapid isolation of glucose-6-phosphate dehydrogenase from human erythrocytes by combined affinity and anion-exchange chromatography for biochemical characterization of variants 573(1992)29
- Application of solid-phase extraction on anion-exchange cartridges to quantify 5'-nucleotidase activity 573(1992)207
- Identification by high-performance liquid chromatography of tyrosine ammonia-lyase activity in purified fractions of *Phaseolus vulgaris* phenylalanine ammonia-lyase 573(1992)309
- Characterization of an aminopeptidase in cerebrospinal fluid. Structure elucidation of enzyme hydrolysis products of synthetic methionine-enkephalin by reversed-phase high-performance liquid chromatography and mass spectrometry 574(1992)189
- Interactions of human alkaline phosphatase isoenzymes with triazine dyes using affinity partitioning, affinity chromatography and difference spectroscopy 574(1992)237
- Determination of endothelin-converting enzyme activity by high-performance liquid chromatography-on-line radioactive flow monitoring 574(1992)344
- Determination of platelet monoamine oxidase activity by high-performance liquid chromatography with electrochemical detection 575(1992)39
- Use of Zn-pyrophosphatase in the high-performance liquid chromatographic analysis of cell extracts containing ³²P-labelled inositol phosphates 575(1992)51
- Affinity chromatography of serine proteinases from the bovine intervertebral disc on BPTI-Separon HEMA 1000 575(1992)229
- Liquid chromatographic monitoring of pseudocholinesterase activity: comparison of methods 575(1992)237
- Determination of alkaline phosphatase isoenzymes in serum by high-performance liquid chromatography with post-column reaction detection 576(1992)79
- Separation of rat pancreatic secretory proteins by cation-exchange fast protein liquid chromatography 576(1992)287
- Resolution of three forms of superoxide dismutase by immobilised metal affinity chromatography 576(1992)340
- Application of an organic solvent extraction to the determination of catechol-O-methyltransferase activity by high-performance liquid chromatography in human mononuclear cells 578(1992)175
- Preparative isolation of angiotensin-converting enzyme from human lung 579(1992)63
- Detection of the enzymatic activity of cytochrome P-450 enzymes by high-performance liquid chromatography (Review) 580(1992)325
- Assay of sialidase activity using ion-exchange chromatography and acidic ninhydrin reaction 581(1992)11
- Affinity purification of proteinases by a combination of immobilized peptidyl aldehyde and semicarbazone 581(1992)17
- Direct assay of glutathione peroxidase activity using high-performance capillary electrophoresis 581(1992)49
- High-performance liquid chromatographic determination of dihydroorotate dehydrogenase of *Plasmodium falciparum* and effects of antimalarials on enzyme activity 582(1992)57
- Evaluation of the chromatographic procedure for the preparation of a high-purity CI-esterase inhibitor concentrate from cryosupernatant plasma 582(1992)65
- Rapid purification yielding highly active 17 β -hydroxysteroid dehydrogenase: application of hydrophobic interaction and affinity fast protein liquid chromatography 582(1992)71
- Use of hydrophilic interaction chromatography for the study of tyrosine protein kinase specificity 583(1992)137
- Purification of S-oxynitrilase from *Sorghum bicolor* by immobilized metal ion affinity chromatography on different carrier materials 584(1992)85
- Purification and characterization of recombinant pyrrolidone carboxyl peptidase of *Bacillus subtilis* 584(1992)101
- Adsorption behaviour of lipase from *Staphylococcus carnosus* on a hydrophobic adsorbent 584(1992)109

- Isolation and purification of bacterial proteinases by means of autofocusing 585(1991)373
- High-performance affinity chromatography of NADP⁺ dehydrogenases from cell-free extracts using a nucleotide analogue as general ligand 586(1991)51
- Purification and characterization of transglutaminases from the genital tract of the male rat 587(1991)55
- Purification and crystallization of yeast hexokinase isoenzymes. Characterization of different forms by chromatofocusing 587(1991)85
- Purification of the coenzyme B₁₂-containing 2-methyleneglutarate mutase from *Clostridium barkeri* by high-performance liquid chromatography 587(1991)93
- Analysis of alanine and aspartate aminotransferase isoforms in mustard (*Sinapis alba* L.) cotyledons 587(1991)101
- Isolation of cytochrome P-450 components from marmoset liver microsomes by high-performance liquid chromatography 587(1991)117
- Rapid ion-exchange chromatography for preparative separation of proteins. IV. Application to bovine carbonic anhydrase III from skeletal muscle 588(1991)139
- Affinity partitioning of erythrocytic phosphofructokinase in aqueous two-phase systems containing poly(ethylene glycol)-bound Cibacron Blue. Influence of pH, ionic strength and substrates/ effectors 589(1992)127
- Specific detection of acetyl-coenzyme A by reversed-phase ion-pair high-performance liquid chromatography with an immobilized enzyme reactor 590(1992)241
- Purification of pro- and eukaryotic superoxide dismutases by charge-controlled hydrophobic chromatography 590(1992)247
- Practical on-line determination of biopolymer molecular weights by high-performance liquid chromatography with classical light-scattering detection 592(1992)215
- Direct determination of kallikrein by high-performance liquid chromatography 593(1992)119
- Isolation and characterization of α -glucosidase from *Aspergillus niger* 593(1992)125
- Silica sorbents with one- and two-site attached bacitracin in affinity chromatography 596(1992)114
- Purification of serine hydroxymethyltransferase from *Bacillus stearothermophilus* with ion-exchange high-performance liquid chromatography 596(1992)203
- Trypsin and affinity chromatography (Review) 597(1992)3
- Recent developments in quantitative affinity chromatography (Review) 597(1992)67
- Affinity purification of proteins using expanded beds 597(1992)129
- Immobilized ferredoxins for affinity chromatography of ferredoxin-dependent enzymes 597(1992)147
- Purification and partial characterization of phospholipase A₂ isoforms from human placenta 597(1992)299
- Occurrence of an allosteric transition in the modification of papain with L-1-acetyl-2,3-dihydropyrrolo[2,3-*b*]-indole-2-carboxamide 597(1992)411
- Use of various types of column reactors for flow-injection analysis 597(1992)435
- High-performance hydrophobic interaction chromatography of proteins on reversed-phase supports coated with non-ionic surfactants of polyoxyethylene type. Purification of a fungal aspartic proteinase 598(1992)181
- Theory of perfusion chromatography 599(1992)87
- Peptide maps of five human pepsin isoenzymes and other aspartic proteinases 599(1992)179
- Purification of DNA-derived deoxynucleotides from leukocytes involving nuclease elution of an ion-exchange column 600(1992)229
- Ephedrine**
Determination of ephedrine alkaloids by capillary electrophoresis 600(1992)370
- Ephedrine alkaloids**
Determination of ephedrine alkaloids by capillary electrophoresis 600(1992)370
- Epicatechin**
High-performance liquid chromatography of the phenolic compounds in the juice of some French cider apple varieties 555(1991)125
- Epidermal growth factor**
Variation of hydrophobicity of human urinary epidermal growth factor 568(1991)226
- Epidoxorubicin**
Simple and sensitive quantification of anthracyclines in mouse atrial tissue using high-performance liquid chromatography and fluorescence detection 570(1991)209
- Measurement of epidoxorubicin and its metabolites by high-performance liquid chromatography using an advanced automated sample processor 572(1991)211

Epidoxorubicinol

Simple and sensitive quantification of anthracyclines in mouse atrial tissue using high-performance liquid chromatography and fluorescence detection 570(1991)209

Measurement of epidoxorubicin and its metabolites by high-performance liquid chromatography using an advanced automated sample processor 572(1991)211

6-Epidoxycycline

Determination of metacycline and related substances by column liquid chromatography on poly(styrene-divinylbenzene) 586(1991)61

Isolation of doxycycline, 6-epidoxycycline and 2-acetyl-2-decarboxamidometacycline from commercial metacycline by preparative column liquid chromatography on silica gel 586(1991)67

4-Epimetacycline

Determination of metacycline and related substances by column liquid chromatography on poly(styrene-divinylbenzene) 586(1991)61

Epinephrine

Improved measurement of urinary catecholamines by liquid-liquid extraction, derivatization and high-performance liquid chromatography with fluorimetric detection 563(1991)348

Evaluation of a simple plasma catecholamine extraction procedure prior to high-performance liquid chromatography and electrochemical detection 564(1991)55

Determination of heat-exposure effects on the concentration of catecholamines in bovine plasma and milk 566(1991)29

Measurement of plasma catecholamines by high-performance liquid chromatography with electrochemical detection in intensive care patients after dobutamine infusion 567(1991)240

Analysis of plasma catecholamines by high-performance liquid chromatography with fluorescence detection: simple sample preparation for pre-column fluorescence derivatization 567(1991)351

Simultaneous determination of free catecholamines and epinine and estimation of total epinine and dopamine in plasma and urine by high-performance liquid chromatography with fluorimetric detection 574(1992)109

Optimization of high-performance liquid chromatographic assay for catecholamines. Determination of optimal mobile phase composition and elimination of species-dependent differences in extraction recovery of 3,4-dihydroxybenzylamine 574(1992)213

Simultaneous measurement of serotonin, catecholamines and their metabolites in cat and human plasma by *in vitro* microdialysis-microbore high-performance liquid chromatography with amperometric detection 582(1992)19

Simultaneous determination of catecholamines and dobutamine in human plasma and urine by high-performance liquid chromatography with fluorimetric detection 583(1992)236

Determination of catecholamines in urine by liquid chromatography and electrochemical detection after on-line sample purification on immobilized boronic acid 593(1992)185

Epinine

Simultaneous determination of free catecholamines and epinine and estimation of total epinine and dopamine in plasma and urine by high-performance liquid chromatography with fluorimetric detection 574(1992)109

Epirubicin

Bioanalysis of some anthracyclines in human plasma by capillary electrophoresis with laser-induced fluorescence detection 574(1992)327

Epoxy carbamazepine

Fast liquid chromatography for the determination of drugs in plasma and combination with liquid-solid extraction in a fully automated system 573(1992)59

Automated microanalysis of carbamazepine and its epoxide and *trans*-diol metabolites in plasma by column liquid chromatography 573(1992)65

Epoxy fatty acids

Enantioselective separation of some polyunsaturated epoxy fatty acids by high-performance liquid chromatography on a cellulose phenylcarbamate (Chiralcel OC) stationary phase 583(1992)231

Ergotamine

Identification and quantification of ergotamine in human plasma by gas chromatography-mass spectrometry 575(1992)289

Eriodictyol

Origin and control of multi-peak formation in the analysis of trimethylsilyl derivatives of flavanone aglycones by capillary column gas chromatography 586(1991)323

Erucic acid

Purification of erucic acid by preparative high-performance liquid chromatography and crystallization 590(1992)139

Erythrocytes

Separation of living red blood cells by gravitational field-flow fractionation 568(1991)93

Monitoring of an experimental red blood cell pathology with gravitational field-flow fractionation 579(1992)73

Investigation of red blood cell fractionation by gravitational field-flow fractionation 583(1992)27

Erythromycin

Determination of dirithromycin purity and related substances by high-performance liquid chromatography 594(1992)203

Erythromycin A N-oxide

Isolation of erythromycin A N-oxide and pseudoerythromycin A hemiketal from fermentation broth of *Saccharopolyspora erythraea* by thin-layer and high-performance liquid chromatography 558(1991)265

Erythromycin 2'-ethylsuccinate

Bioanalysis of erythromycin 2'-ethylsuccinate in plasma using phase-system switching continuous-flow fast atom bombardment liquid chromatography-mass spectrometry 565(1991)265

Erythromycin ethylsuccinate

Determination of erythromycin ethylsuccinate by liquid chromatography 600(1992)99

Erythromycylamine

Determination of dirithromycin purity and related substances by high-performance liquid chromatography 594(1992)203

Eseroline

Determination of heptylphosphostigmine in plasma by high-performance liquid chromatography with electrochemical detection 573(1992)275

Esmolol

Simple and rapid high-performance liquid chromatographic assay for esmolol 570(1991)217

Esperisone

Determination of esperisone in human plasma by gas chromatography-mass spectrometry 584(1992)261

Essential oils

Structure determination of sesquiterpenes in Chinese vetiver oil by gas chromatography-tandem mass spectrometry 557(1991)451

Esterase D

Electrophoretic phenotyping of erythrocyte enzymes (Review) 569(1991)297

Estetrol

Automated direct high-performance liquid chromatographic assay for estetrol, estriol, cortisone and cortisol in serum and amniotic fluid 568(1991)35

Estradiol

Determination of estradiol 2- and 4-hydroxylase activities by gas chromatography with electron-capture detection 567(1991)309

Thin-layer chromatographic detection of zeranol and estradiol in fortified plasma and tissue extracts with Fast Corinth V 581(1992)119

General and selective isolation procedure for high-performance liquid chromatographic determination of anabolic steroids in tissues 588(1991)89

Estra-1,3,5(10)-trienes

Influence of mobile phase composition on evaluation of lipophilicity by partition chromatography 557(1991)399

Estriol

Automated direct high-performance liquid chromatographic assay for estetrol, estriol, cortisone and cortisol in serum and amniotic fluid 568(1991)35

Estrogens

Automated direct high-performance liquid chromatographic assay for estetrol, estriol, cortisone and cortisol in serum and amniotic fluid 568(1991)35

Ethacrynic acid

Simultaneous analysis of flunixin, naproxen, ethacrynic acid, indomethacin, phenylbutazone, mefenamic acid and thiosalicylic acid in plasma and urine by high-performance liquid chromatography and gas chromatography-mass spectrometry 568(1991)351

High-performance liquid chromatographic determination of ethacrynic acid in human plasma 571(1991)271

Ethafuralin

High-performance liquid chromatographic determination of dinitroaniline herbicides in soil and water 585(1991)164

Ethane

Preconcentration technique for introducing gaseous or volatile compounds into a capillary gas chromatographic column 587(1991)338

1,2-Ethanedithiol

Gas chromatographic determination of the lewisite hydrolysate, 2-chlorovinylarsonous acid, after derivatization with 1,2-ethanedithiol 558(1991)235

Ethanol

Estimation of cellulose polarity by gas chromatography 552(1991)365

Development of post-column enzymic reactors with immobilized alcohol oxidase for use in the high-performance liquid chromatographic assay of alcohols with electrochemical detection 563(1991)11

Direct injection high-performance liquid chromatographic method with electrochemical detection for the determination of ethanol and methanol in plasma using an alcohol oxidase reactor 566(1991)333

Chromatographic methods for blood alcohol determination (Review) 580(1992)161

Optimization of an ion-exchange high-performance liquid chromatographic method for the determination of carboxylic acids, sugars, glycerol and ethanol in wines 589(1992)151

Determination of carboxylic acids, sugars, glycerol and ethanol in wine and grape must by ion-exchange high-performance liquid chromatography with refractive index detection 590(1992)215

Analysis of zapatera olives by gas and high-performance liquid chromatography 594(1992)259

Direct aqueous injection gas chromatography on a potassium fluoride crystal hydrate-containing sorbent. Determination of volatile organic solvents in the fermentation broth of *Clostridium* strains 596(1992)138

Ethanol oxidation metabolites

Simple cation-exchange high-performance liquid chromatography optimized to the measurement of metabolites in the effluents from perfused rat livers using refractive index and ultraviolet detectors 563(1991)231

Ethenzamide

Simplex optimization of liquid chromatography with the function of mutual information as a criterion 585(1991)187

Ethinylestradiol

Comparison of two mass-selective detectors with special reference to the analysis of residues of anabolics 564(1991)385

Detection of the illegal use of ethinylestradiol in cattle urine by gas chromatography-mass spectrometry 564(1991)469

Ethionamide

Improved high-performance liquid chromatographic assay for the determination of ethionamide in serum 563(1991)472

7-Ethoxycoumarin

Automated high-performance liquid chromatographic assay for the determination of 7-ethoxycoumarin and umbelliferone 578(1992)141

Ethoxylated alcohols

Comparison of ethoxylated alcohols and polyethylene glycols by high-performance liquid chromatography and supercritical fluid chromatography using evaporative light-scattering detection 591(1992)149

Ethoxylated alkyl alcohols

Use of an evaporative light scattering detector in reversed-phase high-performance liquid chromatography of oligomeric surfactants 552(1991)593

2-Ethoxy-N-[α -(2-methyl-1-propyl)-2-piperidinobenzyl]-4-carbamoylmethylbenzoic acid

Quantitation of the new hypoglycaemic agent AG-EE 388 ZW in human plasma by automated high-performance liquid chromatography with electrochemical detection 568(1991)246

Ethoxyquin

Chromatographic behaviour of the antidegradant ethoxyquin and its transformation products 587(1991)355

6-Ethoxy-2,2,4-trimethyl-1,2-dihydroquinoline

Chromatographic behaviour of the antidegradant ethoxyquin and its transformation products 587(1991)355

Ethyl *p*-aminobenzoate

Separation of neutral mono- and oligosaccharides derivatized with ethyl *p*-aminobenzoate by high-performance liquid chromatography on an amine-bonded vinyl alcohol copolymer column 588(1991)53

Ethyl biscoumacetate

Determination of ethyl biscoumacetate and its metabolite 7-hydroxyethyl biscoumacetate in human serum by high-performance liquid chromatography and mass spectrometry 574(1992)71

Ethyl carbamate

Gas chromatographic-mass spectrometric determination of ethyl carbamate as the xanthylamide derivative in Italian aqua vitae (grappa) samples 585(1991)111

Ethylene

Preconcentration technique for introducing gaseous or volatile compounds into a capillary gas chromatographic column 587(1991)338

Ethylenediamine

Retention of some ethylenediamine oligomers in reversed-phase chromatography 553(1991)407

Measurement of stability constants for complexes of divalent cations with tartaric acid by ion chromatography 589(1992)185

Ethylenediaminetetraacetic acid

Determination of ethylenediaminetetraacetic acid and its salts in canned mushrooms by reversed-phase ion-pair liquid chromatography 553(1991)243

Separation of metal complexes of ethylenediaminetetraacetic acid in environmental water samples by ion chromatography with UV and potentiometric detection 558(1991)181

Simultaneous ion chromatographic determination of chloride and calcium contained in Ringer's injection using a chelating agent as eluent 589(1992)358

Ethylenethiourea

Liquid chromatographic determination of ethylenethiourea using pulsed amperometric detection 586(1991)158

2-Ethylhexanoic acid

Identification of the main metabolites of 2-ethylhexanoic acid in rat urine using gas chromatography–mass spectrometry 568(1991)125

2-Ethylhexyl-*p*-dimethylaminobenzoate

Determination of sun-screen agents in cosmetic products by micellar liquid chromatography 587(1991)325

2-Ethylhexyl-*p*-methoxycinnamate

Determination of sun-screen agents in cosmetic products by micellar liquid chromatography 587(1991)325

Ethylmercury

Evaluation of gas chromatographic columns for the determination of methylmercury in aqueous head space extracts from biological samples 586(1991)329

Assay for thiomersal (thimerosal) with adaptation to the quantitation of total ethylmercury available in degraded samples 587(1991)329

Ethyl 2-methylbutanoate

Simultaneous stereoanalysis of 2-alkyl-branched acids, esters and alcohols using a selectivity-adjusted column system in multi-dimensional gas chromatography 587(1991)347

Ethylmorphine

Improved high-performance liquid chromatographic method for the determination of ethylmorphine and its metabolites in microsomal incubations and cell culture media 579(1992)158

Ethyl-naphthalenes

Analytical study of heavy crude oil fractions by coupling of the transalkylation reaction with supercritical fluid chromatography 593(1992)321

Ethylphenethyl alcohol

Dehydrogenase-silica as a stationary phase for the separation of alcohols and ketones 587(1991)268

7-Ethyl-10-[4-(1-piperidino)-1-piperidino]carbonyloxycamptothecin

Simultaneous determination of the camptothecin analogue CPT-11 and its active metabolite SN-38 by high-performance liquid chromatography: application to plasma pharmacokinetic studies in cancer patients 575(1992)275

3-Ethyl-3-(4-pyridyl)piperidine-2,6-dione

Characterisation of metabolites of 3-ethyl-3-(4-pyridyl)-piperidine-2,6-dione, a potential breast cancer drug 572(1991)143

5-Ethylpyrimidines

High-performance liquid chromatographic analysis of 5-ethylpyrimidines and 5-methylpyrimidines in plasma 563(1991)392

Etoperidone

High-performance liquid chromatographic assay with ultraviolet detection for the determination of etoperidone and two active metabolites, 5-(1-hydroxyethyl)etoperidone and 1-(3-chlorophenyl)piperazine, in plasma 567(1991)433

 β -Eucaine

Determination of limiting ionic mobilities and dissociation constants of some local anaesthetics 596(1992)265

Eugenol

Effects of collection solvent parameters and extraction cell geometry on supercritical fluid extraction efficiencies 594(1992)297

Optimization of supercritical fluid extraction of volatile constituents from a model plant matrix 600(1992)175

F**Factor IX**

Physical and biochemical characterization of five commercial resins for immunoaffinity purification of factor IX 595(1992)103

Famotidine

Sensitive high-performance liquid chromatographic determination of famotidine in plasma. Application to pharmacokinetic study 572(1991)227

Fatty acid-binding protein

High-yield two-step chromatographic procedure for purification of fatty acid-binding protein from human heart 570(1991)173

Fatty acid methyl esters

Gas-liquid microcapillary columns precoated with graphitized carbon black 552(1991)197

Gas chromatographic analysis of fatty acid salts 552(1991)281

Analysis of human blood plasma triacylglycerols using capillary gas chromatography, silver ion thin-layer chromatographic fractionation and desorption chemical ionization mass spectrometry 568(1991)1

Gas chromatographic analysis of fatty acid methyl esters: avoiding discrimination by programmed temperature vaporizing injection 588(1991)265

Fatty acid monomers

Gas chromatography–mass spectrometry and gas chromatography–tandem mass spectrometry of cyclic fatty acid monomers isolated from heated fats 562(1991)659

Fatty acids

see also Carboxylic acids and Organic acids

Applications of capillary gas chromatography to the quality control of butter and related products 552(1991)273

Thermodesorption–gas chromatography–mass spectrometric analysis of biological materials for potential molecular precursors of the constituents of the crude oils 552(1991)325

Isolation and quantification of polymers from autoxidized fish oils by high-performance size-exclusion chromatography with an evaporative mass detector 552(1991)501

Some aspects of peak broadening in particle-beam liquid chromatography–mass spectrometry 554(1991)119

Capillary gas chromatography–mass spectrometry of unusual and very long-chain fatty acids from soil oligotrophic bacteria 558(1991)215

Abnormal fatty acid metabolism in patients in hopantenate therapy during clinical episodes 562(1991)139

Determination of very-long-chain fatty acids in plasma by a simplified gas chromatographic–mass spectrometric procedure 562(1991)147

Simple thin-layer chromatographic purification procedure for the determination of cholesterol ester fatty acid compositions 564(1991)272

Fatty acid composition of glycerophospholipids in seven tissues of cod (*Gadus morhua*), determined by combined high-performance liquid chromatography and gas chromatography 565(1991)119

High-performance liquid chromatographic analysis of free palmitic and stearic acids in cerebrospinal fluid 565(1991)424

Separation and determination of saturated very-long-chain free fatty acids in plasma of patients with adrenoleukodystrophy using solid-phase extraction and high-performance liquid chromatographic analysis 567(1991)319

High-performance liquid chromatographic analysis of fatty acid compositions of platelet phospholipids as their 2-nitrophenylhydrazides 568(1991)25

Assessment of fatty acids in cardiac tissue as 9-anthryldiazomethane esters by high-performance liquid chromatography 570(1991)243

Analysis of polyunsaturated fatty acids in blood serum after fish oil administration 572(1991)1

Determination of phospholipid fatty acid and triacylglycerol composition of rat caecal mucosa 575(1992)1

High-performance liquid chromatography–thermospray mass spectrometry of hydroxy-polyunsaturated fatty acid acetyl derivatives 575(1992)7

Liquid chromatography–mass spectrometry of hydroxy and non-hydroxy fatty acids as amide derivatives 575(1992)197

Simple and rapid method for determination of short-chain fatty acids in biological materials by high-performance liquid chromatography with ultraviolet detection 576(1992)53

Intra-injector methylation of free fatty acids from aerobically and anaerobically cultured *Actinobacillus actinomycetemcomitans* and *Haemophilus aphrophilus* 576(1992)328

Quantitation of phenacyl esters of retinal fatty acids by high-performance liquid chromatography 578(1992)124

Application of triple quadrupole tandem mass spectrometry to the analysis of pyridine-containing derivatives of long-chain acids and alcohols 579(1992)1

High-performance liquid chromatography–thermospray mass spectrometry of hydroperoxy polyunsaturated fatty acid acetyl derivatives 579(1992)25

Determination of fatty acid and triacylglycerol composition of human adipose tissue 581(1992)171

Derivatization of saturated long-chain fatty acids with phenacyl bromide in non-ionic micelles 586(1991)347

Rapid high-performance liquid chromatographic determination of fatty acid profiles of lipids by conversion to their hydroxamic acids 587(1991)292

Isotope effect in gas–liquid chromatography of labelled compounds 588(1991)251

Quantitative planar chromatography of phospholipids with different fatty acid compositions 588(1991)303

Resolution of lipolysis mixtures from soybean oil by a solid-phase extraction procedure 589(1992)353

Purification of erucic acid by preparative high-performance liquid chromatography and crystallization 590(1992)139

Reversed-phase high-performance liquid chromatography of anionic and ethoxylated non-ionic surfactants and pesticides in liquid pesticide formulations 593(1992)227

High-performance liquid chromatography of okadaic acid and free fatty acids in mussels 594(1992)137

Analysis of zapatera olives by gas and high-performance liquid chromatography 594(1992)259

Supercritical carbon dioxide extraction of resin and fatty acids from sediments at pulp mill sites 594(1992)309

Analysis of underivatized C₁₂–C₁₈ fatty acids by reversed-phase ion-pair high-performance liquid chromatography with conductivity detection 596(1992)181

Simultaneous determination of amounts of major phospholipid classes and their fatty acid composition in erythrocyte membranes using high-performance liquid chromatography and gas chromatography 598(1992)33

Fatty acid salts

Gas chromatographic analysis of fatty acid salts 552(1991)281

Fatty alcohols

Application of triple quadrupole tandem mass spectrometry to the analysis of pyridine-containing derivatives of long-chain acids and alcohols 579(1992)1

Fatty amines

Reversed-phase high-performance liquid chromatography of anionic and ethoxylated non-ionic surfactants and pesticides in liquid pesticide formulations 593(1992)227

FB Fragment

Affinity purification of antibodies using immobilized FB domain of protein A 597(1992)257

FCE 23762

Stability studies with a high-performance liquid chromatographic method for the determination of a new anthracycline analogue, 3'-deamino-3'-[2-(*S*)-methoxy-4-morpholino]doxorubicin (FCE 23762), in the final drug formulation 593(1992)47

Felbamate

Determination of the anticonvulsant felbamate in beagle dog plasma by high-performance liquid chromatography 573(1992)113

Fellavine

Simple high-performance liquid chromatographic method for the determination of a new phytochemical drug, fellavine, and its metabolites in human and rat plasma and urine 571(1991)312

Felodipine

Determination of felodipine and its metabolites in plasma using capillary gas chromatography with electron-capture detection and their identification by gas chromatography–mass spectrometry 565(1991)237

Determination of four carboxylic acid metabolites of felodipine in plasma by high-performance liquid chromatography 573(1992)265

Fenbendazole

Determination of some drugs by micellar electrokinetic capillary chromatography. The pseudo-effective mobility as parameter for screening 585(1991)123

Fendiline

Determination of fendiline in human plasma by means of capillary gas chromatography and nitrogen–phosphorus selective detection 564(1991)289

Fenfluramine

Study of human urinary metabolism of fenfluramine using gas chromatography–mass spectrometry 562(1991)267

Fenoldopam

Determination of the enantiomers of fenoldopam in human plasma by reversed-phase high-performance liquid chromatography after chiral derivatization 592(1992)317

Fenoterol

Determination of β_2 -receptor agonists in bovine urine and liver by gas chromatography–tandem mass spectrometry 564(1991)515

Fenoterol hydrobromide

Comparison of isotachopheresis, capillary zone electrophoresis and high-performance liquid chromatography for the determination of salbutamol, terbutaline sulphate and fenoterol hydrobromide in pharmaceutical dosage forms 590(1992)341

Ferredoxin

Immobilized ferredoxins for affinity chromatography of ferredoxin-dependent enzymes 597(1992)147

Ferritin

High-performance liquid chromatography of amino acids, peptides and proteins. CXIV. Protein interactions with porous coulombic sorbents: comparison of experimental findings with predictions of several adsorption models 557(1991)335

Ferrochelatase

High-performance liquid chromatographic assays for protoporphyrinogen oxidase and ferrochelatase in human leucocytes 566(1991)383

Fibrinogen

Base-atom recognition in protein adsorption to alkyl agaroses 597(1992)93

Fibronectin

Use of biospecific interactions of collagen, fibronectin and their fragments in affinity chromatography 577(1992)267

Characterization of the interaction between human plasma fibronectin and collagen by means of affinity electrophoresis 597(1992)351

Finasteride

High-performance liquid chromatographic method for the determination of finasteride in human plasma at therapeutic doses 566(1991)127

FLA797

Optimization of a peak compression system for a remoxipride metabolite (FLA797) and its application to bioanalysis 591(1992)207

Flavanoids

Identification of the constituent flavanoid units in sainfoin proanthocyanidins by reversed-phase high-performance liquid chromatography 594(1992)117

Flavan-3-ols

Identification of the constituent flavanoid units in sainfoin proanthocyanidins by reversed-phase high-performance liquid chromatography 594(1992)117

Flavanone aglycones

Origin and control of multi-peak formation in the analysis of trimethylsilyl derivatives of flavanone aglycones by capillary column gas chromatography 586(1991)323

Flavanone glycosides

High-performance liquid chromatography of diastereomeric flavanone glycosides in *Citrus* on a β -cyclodextrin-bonded stationary phase (Cyclobond I) 588(1991)41

Flavanones

Analysis of commercial hesperidin methyl chalcone by high-performance liquid chromatography 555(1991)285

Origin and control of multi-peak formation in the analysis of trimethylsilyl derivatives of flavanone aglycones by capillary column gas chromatography 586(1991)323

Flavone

Solid-phase extraction with supercritical fluid elution as a sample preparation technique for the ultratrace analysis of flavone in blood plasma 577(1992)61

Flavonoids

Identification of flavonoids from *Ginkgo biloba* L., *Anthemis nobilis* L. and *Equisetum arvense* L. by high-performance liquid chromatography with diode-array UV detection 553(1991)223

Quantitative two-dimensional thin-layer chromatography 553(1991)477

Isotachophoretic analysis of flavonoids and phenolcarboxylic acids of relevance to phytopharmaceutical industry 559(1991)499

Mass spectral characterization of C-glycosidic flavonoids isolated from a medicinal plant (*Passiflora incarnata*) 562(1991)435

High-performance liquid chromatography of diastereomeric flavanone glycosides in *Citrus* on a β -cyclodextrin-bonded stationary phase (Cyclobond I) 588(1991)41

Correlations between high-performance liquid chromatographic retention, X-ray structural and ^{13}C NMR spectroscopic data of flavonoid compounds 592(1992)65

Separation of flavonol-2-O-glycosides from *Calendula officinalis* and *Sambucus nigra* by high-performance liquid and micellar electrokinetic capillary chromatography 593(1992)165

Flavonol glycosides

Isolation and determination of flavonol glycosides from *Epilobium* species 557(1991)391

High-performance liquid chromatographic analysis of flavonol glycosides of *Solidago virgaurea* 558(1991)296

Separation of flavonol-2-O-glycosides from *Calendula officinalis* and *Sambucus nigra* by high-performance liquid and micellar electrokinetic capillary chromatography 593(1992)165

Flavonols

High-performance liquid chromatographic analysis of flavonol glycosides of *Solidago virgaurea* 558(1991)296

Structure elucidation of saccharides in anthocyanins and flavonols by means of methylation analysis and gas chromatography 598(1992)81

Floxuridine

High-performance liquid chromatographic method for the simultaneous measurement of floxuridine and fluorouracil in human body fluids 566(1991)147

Flucyclohexan

Application of an internal surface reversed-phase column for the automated determination of flucyclohexan residues 600(1992)211

Flufenamic acid

Simultaneous optimization of several chromatographic performance goals using Derringer's desirability function 586(1991)11

Flumequine

Automated analysis of oxolinic acid and flumequine in salmon whole blood and plasma using dialysis combined with trace enrichment as on-line sample preparation for high-performance liquid chromatography 564(1991)171

Direct-gradient high-performance liquid chromatographic analysis and preliminary pharmacokinetics of flumequine and flumequine acyl glucuronide in humans: effect of probenecid 579(1992)131

Flumequine acyl glucuronide

Direct-gradient high-performance liquid chromatographic analysis and preliminary pharmacokinetics of flumequine and flumequine acyl glucuronide in humans: effect of probenecid 579(1992)131

Flunixin

Simultaneous analysis of flunixin, naproxen, ethacrynic acid, indomethacin, phenylbutazone, mefenamic acid and thiosalicylic acid in plasma and urine by high-performance liquid chromatography and gas chromatography-mass spectrometry 568(1991)351

Fluorene

Laser-excited fluorescence detection of gas-phase chromatography eluates 557(1991)39

Fluorescamine

Pre-column derivatization of sulfa drugs with fluorescamine and high-performance liquid chromatographic determination at their residual levels in meat and meat products 558(1991)175

Assay of protein drug substances present in solution mixtures by fluorescamine derivatization and capillary electrophoresis 598(1992)123

Fluorescein

Fluorescence detector cell for use in an integrated electrically driven separation system 598(1992)115

Fluoride

Determination of trace amounts of fluoride in raw materials for pharmaceuticals by gas-liquid chromatography 585(1991)289

3-Fluoroalanine

High-performance ligand-exchange liquid chromatography of fluoro derivatives of alanine 600(1992)83

5-Fluoro-2'-deoxyuridine

High-performance liquid chromatographic method for the simultaneous measurement of floxuridine and fluorouracil in human body fluids 566(1991)147

Determination of 5-fluorouracil and its main metabolites in plasma by high-performance liquid chromatography 573(1992)247

Quantification of 5-fluoro-2'-deoxyuridine in plasma by gas chromatography and negative-ion chemical ionization mass spectrometry 575(1992)63

Fluorodeoxyuridine monophosphate

Determination of 5-fluorouracil and its main metabolites in plasma by high-performance liquid chromatography 573(1992)247

(±)-(E)-1-(3-Fluoro-6,11-dihydrodibenz[*b,e*]oxepin-11-yl)-4-(3-phenyl-2-propenyl)piperazine dimaleate

Determination of AJ-3941, a possible agent for the treatment of cerebrovascular disorders, in plasma and brain by means of high-performance liquid chromatography with fluorescence detection 578(1992)152

α-(4-Fluorophenyl)-4-(5-fluoro-2-pyrimidinyl)-1-piperazine butanol

High-performance liquid chromatographic assay of α-(4-fluorophenyl)-4-(5-fluoro-2-pyrimidinyl)-1-piperazine butanol (BMY 14802), a potential antipsychotic drug, in monkey and rat plasma 567(1991)274

5-Fluorouracil

Highly sensitive method for the determination of 5-fluorouracil in biological samples in the presence of 2'-deoxy-5-fluorouridine by gas chromatography-mass spectrometry 564(1991)137

High-performance liquid chromatographic method for the simultaneous measurement of floxuridine and fluorouracil in human body fluids 566(1991)147

Determination of 5-fluorouracil and its main metabolites in plasma by high-performance liquid chromatography 573(1992)247

Gas chromatographic-mass spectrometric method for routine monitoring of 5-fluorouracil in plasma of patients receiving low-level protracted infusions 581(1992)195

Analysis of 5-fluorouracil in plasma and urine by high-performance liquid chromatography 581(1992)281

Fluorouridine

Determination of 5-fluorouracil and its main metabolites in plasma by high-performance liquid chromatography 573(1992)247

Fluoxetine

Solid-phase extraction of fluoxetine and norfluoxetine from serum with gas chromatography-electron-capture detection 563(1991)379

Simultaneous determination of fluoxetine and norfluoxetine enantiomers in biological samples by gas chromatography with electron-capture detection 579(1992)99

Sensitive micromethod for column liquid chromatographic determination of fluoxetine and norfluoxetine in human plasma 583(1992)217

Flurazepam

Determination of small drug molecules by capillary electrophoresis-atmospheric pressure ionization mass spectrometry 559(1991)515

Flurbiprofen

Stereoselective high-performance liquid chromatographic determination of flurbiprofen in human plasma 573(1992)163

Retention and enantioselectivity of racemic solutes on a modified ovomucoid-bonded column. I. Cross-linking with glutaraldehyde 592(1992)301

Fluvoxamine

Sensitive one-step extraction procedure for column liquid chromatographic determination of fluvoxamine in human and rat plasma 567(1991)441

Determination of fluvoxamine in human plasma by high-performance liquid chromatography with fluorescence detection 574(1992)101

Food dyes

Electrospray interfacing for the coupling of ion-exchange and ion-pairing chromatography to mass spectrometry 554(1991)27

Formaldehyde

Excretion of formaldehyde, malondialdehyde, acetaldehyde and acetone in the urine of rats in response to 2,3,7,8-tetrachlorodibenzo-*p*-dioxin, paraquat, endrin and carbon tetrachloride 576(1992)221

Automated high-performance liquid chromatographic and size-exclusion chromatographic sample preparation by means of a robotic workstation 592(1992)249

Formaldehyde adducts

Formation of formaldehyde adducts from various drugs by use of methanol in a toxicological screening procedure with gas chromatography-mass spectrometry 563(1991)73

Formic acid

Determination of volatile and non-volatile organic acids in technical sugar solutions by ion-exclusion chromatography 555(1991)65

N-Formyl-5-aminosalicylic acid

High-performance liquid chromatographic assay of 5-aminosalicylic acid (5-ASA) and its metabolites N- β -D-glucopyranosyl-5-ASA, N-acetyl-5-ASA, N-formyl-5-ASA and N-butyryl-5-ASA in biological fluids 570(1991)109

(R,S)-N⁵-Formyltetrahydrofolate

Preparation of a chiral matrix. Resolution of (6*R*,*S*)-N⁵-formyltetrahydrofolate 555(1991)254

FRC-8653

Determination of a new calcium antagonist and its main metabolite in plasma by thermospray liquid chromatography-mass spectrometry 583(1992)116

Free fatty acids

High-performance liquid chromatographic analysis of free palmitic and stearic acids in cerebrospinal fluid 565(1991)424

Separation and determination of saturated very-long-chain free fatty acids in plasma of patients with adrenoleukodystrophy using solid-phase extraction and high-performance liquid chromatographic analysis 567(1991)319

Fructose

Glucose-fructose equilibria on Dowex Monosphere 99 CA resin under overloaded conditions 590(1992)147

Fuels

Use of basic alumina in fractionation of fossil fuels 600(1992)243

Fullerenes

Liquid chromatographic separation of all-carbon molecules C₆₀ and C₇₀ with multi-legged phenyl group bonded silica phases 594(1992)105

Fumonisin B₁

Determination of fumonisin B₁ in plasma and urine by high-performance liquid chromatography 574(1992)299

Fungicides

see Pesticides

2-Furaldehyde

High-performance liquid chromatographic determination of 2-furaldehyde in spirits 590(1992)235

Furaltadone

Determination of some drugs by micellar electrokinetic capillary chromatography. The pseudo-effective mobility as parameter for screening 585(1991)123

Furan aldehydes

High-performance liquid chromatographic study of the correlation between the physico-chemical parameters of furan and benzene aldehydes and the rate of their metabolism by yeast 594(1992)23

Furans

Comparison of extraction methods for polychlorinated dibenzo-*p*-dioxins and dibenzofurans in fly ash using gas chromatography-mass spectrometry 589(1992)265

Furosemide

Rapid high-performance liquid chromatographic determination with fluorescence detection of furosemide in human body fluids and its confirmation by gas chromatography-mass spectrometry 564(1991)567

Simultaneous determination of furosemide and amiloride in plasma using high-performance liquid chromatography with fluorescence detection 575(1992)269

Fusion protein

Comparison of detergents for extraction and ion-exchange high-performance liquid chromatography of Sendai virus membrane proteins 599(1992)157

Fv antibody fragments

Immobilization of Fv antibody fragments on porous silica and their utility in affinity chromatography 587(1991)161

G

Galactitol

High-performance liquid chromatographic determination of sugars and polyols in extracts of lichens and sugarcane juice 553(1991)271

Simultaneous determination of gluconolactone, galactonolactone and galactitol in urine by reversed-phase liquid chromatography: application to galactosemia 570(1991)277

Galactonolactone

Simultaneous determination of gluconolactone, galactonolactone and galactitol in urine by reversed-phase liquid chromatography: application to galactosemia 570(1991)277

Galactose

High-performance liquid chromatographic determination of sugars and polyols in extracts of lichens and sugarcane juice 553(1991)271

β -Galactosidase

Immunoaffinity chromatography in biorecovery: an application of recombinant DNA technology to generic adsorption processes 584(1992)59

Gallic acid

Determination of tannic acid and its phenolic metabolites in biological fluids by high-performance liquid chromatography 577(1992)77

Gallocatechin

Identification of the constituent flavanoid units in sainfoin proanthocyanidins by reversed-phase high-performance liquid chromatography 594(1992)117

Gallopamil

Direct determination of the enantiomeric ratio of verapamil, its major metabolite norverapamil and gallopamil in plasma by chiral high-performance liquid chromatography 575(1992)255

Ganciclovir

High-performance liquid chromatographic determination of ganciclovir in plasma 567(1991)480

Modified high-performance liquid chromatographic method for the determination of ganciclovir in plasma from patients with severe renal impairment 571(1991)331

GBR 12909

Column liquid chromatographic assay of the dopamine reuptake inhibitor vanoxerine (GBR 12909) in human serum 571(1991)305

Gentamicin sulphate

Analysis of gentamicin sulfate by high-performance liquid chromatography combined with thermospray mass spectrometry 554(1991)191

Gentamycin

High-performance liquid chromatography method for the determination of aminoglycosides based on automated pre-column derivatization with *o*-phthalaldehyde 593(1992)69

Gentisic acid

Novel direct high-performance liquid chromatographic method for determination of salicylate glucuronide conjugates in human urine 565(1991)309

Simultaneous high-performance liquid chromatographic determination of salicylates in whole blood, plasma and isolated erythrocytes 584(1992)221

Ginkgetin

Identification of flavonoids from *Ginkgo biloba* L., *Anthemis nobilis* L. and *Equisetum arvense* L. by high-performance liquid chromatography with diode-array UV detection 553(1991)223

Ginkgolides

Gas chromatographic-mass spectrometric analysis of ginkgolides produced by *Ginkgo biloba* cell culture 588(1991)281

Girosepam

Pharmacokinetic and metabolism studies on girosepam by chromatographic and spectrometric methods in humans 578(1992)91

Gliadins

Characterization and identification of wheat cultivars by multi-dimensional analysis of reversed-phase high-performance liquid chromatograms 596(1992)225

Glibenclamide

Determination of glibenclamide and its two major metabolites in human serum and urine by column liquid chromatography 564(1991)223

Gliclazide

Determination of gliclazide in serum by high-performance liquid chromatography using solid-phase extraction 583(1992)266

Globin chains

Quantities of adult, fetal and embryonic globin chains in the blood of eighteen- to twenty-week-old human fetuses 567(1991)359

K562 cells: a source for embryonic globin chains 571(1991)37

Globins

Separation of globins using free zone capillary electrophoresis 576(1992)346

α 2u-Globulin

Design of a gas chromatograph with parallel radioactivity and mass spectrometric detection. Application to the identification of the major metabolite of *d*-limonene associated with α 2u-globulin 563(1991)271

 γ -Globulins

Optimizing separation parameters in capillary isoelectric focusing 559(1991)479

Glucagon

High-performance liquid chromatography of amino acids, peptides and proteins. CXV. Thermodynamic behaviour of peptides in reversed-phase chromatography 593(1992)103

 β -D-Glucan oligosaccharides

Characterization of cellulase-based enzyme reactors for the high-performance liquid chromatographic determination of β -D-glucan oligosaccharides 558(1991)343

 β -Glucans

Rapid method for preparing a β -glucan-specific sensitive fraction from *Limulus (Tachypleus tridentatus)* amoebocyte lysate 567(1991)267

D-Glucobioses

Analyses of isomeric mono-O-methyl-D-glucoses, D-glucobioses and D-glucose monophosphates by high-performance anion-exchange chromatography with pulsed amperometric detection 595(1992)340

Glucobrassicin

Identification of enzymatic degradation products from synthesized glucobrassicin by gas chromatography-mass spectrometry 586(1991)166

Glucocorticoids

Automated direct high-performance liquid chromatographic assay for estretol, estriol, cortisone and cortisol in serum and amniotic fluid 568(1991)35

Gluconic acid

Determination of volatile and non-volatile organic acids in technical sugar solutions by ion-exclusion chromatography 555(1991)65

Gluconolactone

Simultaneous determination of gluconolactone, galactonolactone and galactitol in urine by reversed-phase liquid chromatography: application to galactosemia 570(1991)277

N- β -D-Glucopyranosyl-5-aminosalicylic acid

High-performance liquid chromatographic assay of 5-aminosalicylic acid (5-ASA) and its metabolites N- β -D-glucopyranosyl-5-ASA, N-acetyl-5-ASA, N-formyl-5-ASA and N-butyryl-5-ASA in biological fluids 570(1991)109

Stability of 5-aminosalicylic acid and its metabolites in plasma at -20°C . Formation of N- β -D-glucopyranosyl-5-aminosalicylic acid 570(1991)224

Glucosaccharides

Characterization of cellulase-based enzyme reactors for the high-performance liquid chromatographic determination of β -D-glucan oligosaccharides 558(1991)343

Glucose

Trimethylsilyl-O-methyloxime derivatives for the measurement of [6,6- $^2\text{H}_2$]-D-glucose-enriched plasma samples by gas chromatography-mass spectrometry 572(1991)302

Determination of plasma [6,6- $^2\text{H}_2$]glucose enrichment by a simple and accurate gas chromatographic-mass spectrometric method 573(1992)127

Glucose-fructose equilibria on Dowex Monosphere 99 CA resin under overloaded conditions 590(1992)147

D-Glucose monophosphates

Analyses of isomeric mono-O-methyl-D-glucoses, D-glucobioses and D-glucose monophosphates by high-performance anion-exchange chromatography with pulsed amperometric detection 595(1992)340

Glucose oxidase

Galactosylation as a tool for the stabilization and immobilization of proteins (Review). 597(1992)19

Glucose-6-phosphate dehydrogenase

Rapid isolation of glucose-6-phosphate dehydrogenase from human erythrocytes by combined affinity and anion-exchange chromatography for biochemical characterization of variants 573(1992)29

High-performance affinity chromatography of NADP⁺ dehydrogenases from cell-free extracts using a nucleotide analogue as general ligand 586(1991)51

 α -Glucosidase

Isolation and characterization of α -glucosidase from *Aspergillus niger* 593(1992)125

Glucosides

Determination of the norlignan glucosides of Hypoxidaceae by high-performance liquid chromatography 594(1992)131

Glucosinolates

Identification of enzymatic degradation products from synthesized glucobrassicin by gas chromatography-mass spectrometry 586(1991)166

Glutamate

Measurement of glutamate, aspartate and glycine and its potential precursors in human brain using high-performance liquid chromatography by pre-column derivatization with diethylaminoazobenzene sulphonyl chloride 583(1992)241

Glutamic acid

Determination of D-alanine and D-glutamic acid in biological samples by coupled-column chromatography using β -cyclodextrin as mobile phase additive 582(1992)35

Determination of glutamine, glutamic acid and pyroglutamic acids using high-performance liquid chromatography on dynamically modified silica 594(1992)173

Glutamic acid decarboxylase

Use of high-performance liquid chromatography for assay of glutamic acid decarboxylase. Its limitation in use for *post-mortem* brain 571(1991)235

Glutamine

Determination of glutamine, glutamic acid and pyroglutamic acids using high-performance liquid chromatography on dynamically modified silica 594(1992)173

Glutamine synthetase

Purification and determination of glutamine synthetase by high-performance immunoaffinity chromatography 589(1992)121

7-N-[2-((γ -L-Glutamylamino)ethyl)dithio]ethyl]mitomycin C
Determination of the new anticancer agent KW 2149, 7-N-[2-((γ -L-glutamylamino)ethyl)dithio]ethyl]mitomycin C, an analogue of mitomycin C 564(1991)352

 γ -Glutamyl dopamine

Determination of γ -glutamyl conjugates of monoamines by means of high-performance liquid chromatography with electrochemical detection and application to gastropod tissues 567(1991)49

 γ -Glutamyl-5-hydroxytryptamine

Determination of γ -glutamyl conjugates of monoamines by means of high-performance liquid chromatography with electrochemical detection and application to gastropod tissues 567(1991)49

 γ -Glutamyl monoamines

Determination of γ -glutamyl conjugates of monoamines by means of high-performance liquid chromatography with electrochemical detection and application to gastropod tissues 567(1991)49

 γ -Glutamyl peptides

Automated determination of neuroactive acidic sulphur-containing amino acids and γ -glutamyl peptides using liquid chromatography with fluorescence and electrochemical detection 566(1991)39

Glutaraldehyde

Automated high-performance liquid chromatographic and size-exclusion chromatographic sample preparation by means of a robotic workstation 592(1992)249

Glutathione

Time-resolved luminescence detection of derivatized thiols in column liquid chromatography 552(1991)625

Determination of oxidized, reduced and protein-bound glutathione in eye lenses by high-performance liquid chromatography and electrochemical detection 570(1991)19

Electrochemical determination of femtomole amounts of free reduced and oxidized glutathione. Application to human hair follicles 576(1992)255

Reversed-phase chromatographic method for specific determination of glutathione in cultured malignant cells 584(1992)167

Glutathione, oxidized

Measurement of reduced and oxidized glutathione in cultures of adult rat hepatocytes 565(1991)149

Fluorimetric determination of oxidised and reduced glutathione in cells and tissues by high-performance liquid chromatography following derivatization with dansyl chloride 568(1991)219

Glutathione, reduced

Measurement of reduced and oxidized glutathione in cultures of adult rat hepatocytes 565(1991)149

Fluorimetric determination of oxidised and reduced glutathione in cells and tissues by high-performance liquid chromatography following derivatization with dansyl chloride 568(1991)219

Glutathione peroxidase

Direct assay of glutathione peroxidase activity using high-performance capillary electrophoresis 581(1992)49

Glutathione reductase

High-performance affinity chromatography of NADP⁺ dehydrogenases from cell-free extracts using a nucleotide analogue as general ligand 586(1991)51

Glycated albumin

High-performance affinity chromatography system for the rapid, efficient assay of glycated albumin 597(1992)271

Glycerobenzoates

Application of liquid chromatography-thermospray mass spectrometry in the analysis of glycerophospholipid molecular species 562(1991)191

Glycerol

Determination of "free" glycerol in human serum reference materials by isotope-dilution gas chromatography-mass spectrometry 578(1992)1

Optimization of an ion-exchange high-performance liquid chromatographic method for the determination of carboxylic acids, sugars, glycerol and ethanol in wines 589(1992)151

Determination of carboxylic acids, sugars, glycerol and ethanol in wine and grape must by ion-exchange high-performance liquid chromatography with refractive index detection 590(1992)215

Glycerophospholipids

Application of liquid chromatography–thermospray mass spectrometry in the analysis of glycerophospholipid molecular species 562(1991)191

Fatty acid composition of glycerophospholipids in seven tissues of cod (*Gadus morhua*), determined by combined high-performance liquid chromatography and gas chromatography 565(1991)119

Isolation of experimental anti-AIDS glycerophospholipids by micro-preparative reversed-phase high-performance liquid chromatography 590(1992)153

Optimized method for the determination of 1,2-diacyl-*sn*-glycero-3-phosphocholine and 1,2-diacyl-*sn*-glycero-3-phosphoethanolamine molecular species by enzymatic hydrolysis and gas chromatography 600(1992)257

L- α -Glycerophosphorylcholine

High-performance liquid chromatographic assay of L- α -glycerophosphorylcholine using a two-step enzymic conversion 566(1991)445

Glyceryl dinitrate

Improved gas chromatographic assay for the simultaneous determination of nitroglycerin and its mono- and dinitrate metabolites 579(1992)237

Glyceryl mononitrate

Improved gas chromatographic assay for the simultaneous determination of nitroglycerin and its mono- and dinitrate metabolites 579(1992)237

Glyceryl nitrates

Simultaneous determination of nitroglycerin and its dinitrate metabolites by capillary gas chromatography with electron-capture detection 577(1992)167

Glycinamide

Simple and rapid micro-analytical procedures for the estimation of milacemide and its metabolite glycinamide in rat plasma and cerebrospinal fluid by high-performance liquid chromatography 565(1991)357

Glycine

Measurement of glutamate, aspartate and glycine and its potential precursors in human brain using high-performance liquid chromatography by pre-column derivatization with diethylaminoazobenzene sulphonyl chloride 583(1992)241

Glycoalkaloids

Determination of potato glycoalkaloids and their aglycone in blood serum by high-performance liquid chromatography. Application to pharmacokinetic studies in humans 573(1992)69

Glycogen phosphorylase

Purification of glycogen phosphorylase isozymes by metal-affinity chromatography 584(1992)77

Glycol ethers

Gas chromatographic analysis of some glycol ether analytes on a specially packed column 552(1991)353

Glycolic acid

Determination of volatile and non-volatile organic acids in technical sugar solutions by ion-exclusion chromatography 555(1991)65

Glycolysis metabolites

Simple cation-exchange high-performance liquid chromatography optimized to the measurement of metabolites in the effluents from perfused rat livers using refractive index and ultraviolet detectors 563(1991)231

Glycopeptides

Influence of different N- and O-linked carbohydrates on the retention times of synthetic peptides in reversed-phase high-performance liquid chromatography 599(1992)43

Glycoproteins

Liquid chromatographic method for the determination of the carbohydrate moiety of glycoproteins. Application to α_2 -acid glycoprotein and tissue plasminogen activator 558(1991)105

Fractionation of the human recombinant tissue plasminogen activator (rtPA) glycoforms by high-performance capillary zone electrophoresis and capillary isoelectric focusing 559(1991)401

Optimization of oxidation of glycoproteins: an assay for predicting coupling to hydrazide chromatographic supports 587(1991)171

Charge-reversed, polymer-coated capillary column for the analysis of a recombinant chimeric glycoprotein 594(1992)317

Galactosylation as a tool for the stabilization and immobilization of proteins (Review) 597(1992)19

Immobilized lectin columns: useful tools for the fractionation and structural analysis of oligosaccharides (Review) 597(1992)111

Measurement of amino acid compositions of glycoprotein systems by gas-phase hydrolysis and reversed-phase high-performance liquid chromatography 599(1992)3

Purification and quantification of recombinant Epstein–Barr viral glycoproteins gp350/220 from Chinese hamster ovary cells 599(1992)267

Purification of recombinant human interferon- β by immobilized antisense peptides 600(1992)235

Glycopyrrolate

Solid-phase extraction techniques for the determination of glycopyrrolate from equine urine by liquid chromatography–tandem mass spectrometry and gas chromatography–mass spectrometry 573(1992)43

Glycosaminoglycans

Novel affinity chromatographic system for the single-step purification of glycosaminoglycans from complex systems using volatile buffers 565(1991)416

Chemiluminescence high-performance liquid chromatography for the determination of hyaluronic acid, chondroitin sulphate and dermatan sulphate 579(1992)203

High-performance liquid chromatographic–mass spectrometric analysis of oligosaccharides from enzymatic digestion of glycosaminoglycans. Application to human samples 591(1992)225

Glycosides

Analytical methods for monoterpene glycosides in grape and wine. I. XAD-2 extraction and gas chromatographic–mass spectrometric determination of synthetic glycosides 590(1992)313

Analytical methods for monoterpene glycosides in grape and wine. II. Qualitative and quantitative determination of monoterpene glycosides in grape 595(1992)269

C-Glycosidic flavonoids

Mass spectral characterization of C-glycosidic flavonoids isolated from a medicinal plant (*Passiflora incarnata*) 562(1991)435

Glycosphingolipids

Use of N-acetylpsychosine as internal standard for quantitative high-performance liquid chromatographic analysis of glycosphingolipids 555(1991)267

Two-dimensional mapping by high-performance liquid chromatography of pyridylamino oligosaccharides from various glycosphingolipids 586(1991)35

Glycyrrhetic acid-3-O-glucuronide

Selective high-performance liquid chromatographic method for the determination of glycyrrhizin and glycyrrhetic acid-3-O-glucuronide in biological fluids: application of ion-pair extraction and fluorescence labelling agent 567(1991)151

Glycyrrhetic acid

High-performance liquid chromatographic determination of 18 α -glycyrrhetic acid and 18 β -glycyrrhetic acid in rat plasma: application to pharmacokinetic study 567(1991)405

Determination of glycyrrhetic acid in human plasma by high-performance liquid chromatography 568(1991)232

Glycyrrhizin

Selective high-performance liquid chromatographic method for the determination of glycyrrhizin and glycyrrhetic acid-3-O-glucuronide in biological fluids: application of ion-pair extraction and fluorescence labelling agent 567(1991)151

Determination of glycyrrhizin in rabbit plasma by high-performance liquid chromatography with photodiode-array ultraviolet detection and its pharmacokinetics application 576(1992)170

Glyoxal

Automated high-performance liquid chromatographic and size-exclusion chromatographic sample preparation by means of a robotic workstation 592(1992)249

Glyphosate

High-performance liquid chromatographic determination of glyphosate and (aminomethyl)phosphonic acid in human serum after conversion into *p*-toluenesulphonyl derivatives 566(1991)239

Determination of glyphosate and its metabolite, (aminomethyl)phosphonic acid, in serum using capillary electrophoresis 571(1991)324

Gossypol

High-performance liquid chromatographic method for the determination of free gossypol in chicken liver 587(1991)333

Granulocyte colony stimulating factor

Fast protein monitoring in fermentation broth using non-porous micropellicular reversed-phase columns 594(1992)392

Gratisin

Studies on cyclic peptides related to gratisin by reversed-phase high-performance liquid chromatography 574(1992)65

Growth hormones

Purification of canine prolactin and growth hormone by fast protein liquid chromatography 553(1991)117

Characterization of a tryptic digest by high-performance displacement chromatography and mass spectrometry 557(1991)289

Mathematical model describing dispersion in free solution capillary electrophoresis under stacking conditions 559(1991)3

Temperature elevations of the sample zone in free solution capillary electrophoresis under stacking conditions 559(1991)27

Radial pH distribution during capillary electrophoresis with electroosmotic flow. Analysis with high ionic strength buffers 589(1992)315

Computer simulation as a tool for the rapid optimization of the high-performance liquid chromatographic separation of a tryptic digest of human growth hormone 594(1992)65

Growth promoters

Column liquid chromatographic determination of carbadox and olaquinox in feeds 558(1991)125

G substances

Plant growth regulators G₁, G₂, G₃. Synthesis, extraction and determination of leaf content in *Eucalyptus grandis* 600(1992)323

Guaifenesin

Simplex optimization of liquid chromatography with the function of mutual information as a criterion 585(1991)187

Guanosine

Determination of a guanosine-malonaldehyde adduct in urine by high-performance liquid chromatography with a thiobarbituric acid reaction detector 570(1991)301

Determination of concentrations of adenosine and other purines in human term placenta by reversed-phase high-performance liquid chromatography with photodiode-array detection: evidence for pathways of purine metabolism in the placenta 575(1992)243

Gymnemic acids

High-performance liquid chromatography-atmospheric pressure ionization mass spectrometry of gymnemic acids 557(1991)383

H

Haemagglutinin-neuraminidase

Comparison of detergents for extraction and ion-exchange high-performance liquid chromatography of Sendai virus membrane proteins 599(1992)157

Haematoporphyrin

Use of hematoporphyrin as a fluorescent stain for detection of lipids in high-performance thin-layer chromatography 588(1991)295

Haemoglobins

Electrospray mass spectrometry in the clinical diagnosis of variant hemoglobins 562(1991)175

Gas chromatographic-mass spectrometric analysis of diols and tetrols from reactions of polycyclic aromatic hydrocarbon epoxides with hemoglobin 562(1991)563

Separation of caprine globin chains by reversed-phase high-performance liquid chromatography: evidence for the presence of a silent β^0 -globin allele in Sardinian sheep 563(1991)158

Quantities of adult, fetal and embryonic globin chains in the blood of eighteen- to twenty-week-old human fetuses 567(1991)359

K562 cells: a source for embryonic globin chains 571(1991)37

Coupling of *m*-aminophenylboronic acid to *s*-triazine-activated Sephacryl: use in the affinity chromatography of glycosylated hemoglobins 575(1992)223

Halofantrine

Measurement of halofantrine and its major metabolite desbutylhalofantrine in plasma and blood by high-performance liquid chromatography: a new methodology 581(1992)156

Halofantrine, metabolites

Identification of metabolites of halofantrine, a new candidate anti-malarial drug, by gas chromatography-mass spectrometry 581(1992)203

Halogenated compounds

Relationship between gas chromatographic retention indices and molecular connectivity indices of chlorinated pesticides and structurally related compounds 598(1992)243

Haloperidol

Simultaneous determination of haloperidol and its metabolite, reduced haloperidol, in plasma, blood, urine and tissue homogenates by high-performance liquid chromatography 572(1991)259

Haloperidol pyridinium metabolite

Determination of the pyridinium metabolite derived from haloperidol in brain tissue, plasma and urine by high-performance liquid chromatography with fluorescence detection 579(1992)277

Hamycin

Identification of aromatic moieties and mycosamine in antifungal heptaenes with high-performance liquid chromatography, high-performance liquid chromatography-mass spectrometry and gas chromatography-mass spectrometry 585(1991)247

High-performance liquid chromatography of heptaene polyenes: assay of heptaene produced by *Streptomyces griseoviridis* 588(1991)356

α -hANP-specific antibodies

Construction and use of two α -human atrial natriuretic peptide-fragment affinity chromatography columns in the isolation of C- and N-terminal epitope-specific antibodies for use in a prototype α -hANP biosensor 577(1992)251

Haptens

Novel solid-state fluorodensitometric method for the determination of haptens in protein-hapten conjugates. Demonstration with a toxic glycoside of *Cleistanthus collinus* 574(1992)267

Heat shock protein

Hydrophobic interaction chromatography for the purification of a mycobacterial heat shock protein of relative molecular mass 60 000 587(1991)19

Heparin

Affinity high-performance liquid chromatography on heparin-Glc-Spheron 563(1991)124

High-performance liquid chromatographic-mass spectrometric analysis of oligosaccharides from enzymatic digestion of glycosaminoglycans. Application to human samples 591(1992)225

Hepatitis B surface antigen

Analysis of a vaccine purification process by capillary electrophoresis 559(1991)337

Heptabarbital

Liquid chromatography-thermospray tandem mass spectrometry for identification of a heptabarbital metabolite and sample work-up artefacts 554(1991)205

Heptaene polyenes

High-performance liquid chromatography of heptaene polyenes: assay of heptaene produced by *Streptomyces griseoviridis* 588(1991)356

Heptaenes

Identification of aromatic moieties and mycosamine in antifungal heptaenes with high-performance liquid chromatography, high-performance liquid chromatography-mass spectrometry and gas chromatography-mass spectrometry 585(1991)247

Heptakis(2,3,6-tri-O-ethyl)- β -cyclodextrin

Simultaneous stereoanalysis of 2-alkyl-branched acids, esters and alcohols using a selectivity-adjusted column system in multi-dimensional gas chromatography 587(1991)347

Heptamethoxyflavone

Fast separation of polymethoxylated flavones by carbon dioxide supercritical fluid chromatography 586(1991)171

Heptanesulphonyl chlorides

Simultaneous gas chromatographic analysis of heptyl chloride-heptanesulphonyl chloride isomeric mixtures 596(1992)132

Heptyl chlorides

Simultaneous gas chromatographic analysis of heptyl chloride-heptanesulphonyl chloride isomeric mixtures 596(1992)132

Heptylphysostigmine

Determination of heptylphysostigmine in plasma by high-performance liquid chromatography with electrochemical detection 573(1992)275

Determination of picogram levels of heptylphysostigmine in human plasma using high-performance liquid chromatography with fluorescence detection 581(1992)227

Herbicides

see Pesticides

Heroin

Rapid assay of cocaine, opiates and metabolites by gas chromatography-mass spectrometry (Review) 580(1992)43

Hesperidin methyl chalcone

Analysis of commercial hesperidin methyl chalcone by high-performance liquid chromatography 555(1991)285

Hesperitin

Origin and control of multi-peak formation in the analysis of trimethylsilyl derivatives of flavanone aglycones by capillary column gas chromatography 586(1991)323

Hetacillin

Development of analytical methods for some penicillins in bovine milk by ion-paired chromatography and confirmation by thermospray mass spectrometry 567(1991)389

Hexachlorobenzene

New Czechoslovak sorbent, Ekosorb 552(1991)449

Ionization mechanisms in capillary supercritical fluid chromatography-chemical ionization mass spectrometry 554(1991)351

Hexadecane

Activity coefficients in binary liquid mixtures measured by reversed-flow gas chromatography 595(1992)237

Hexahydrophthalic acid

Method for the biological monitoring of hexahydrophthalic anhydride by the determination of hexahydrophthalic acid in urine using gas chromatography and selected-ion monitoring 572(1991)117

Hexahydrophthalic anhydride

Determination of hexahydrophthalic anhydride in air using gas chromatography 558(1991)247

Method for the biological monitoring of hexahydrophthalic anhydride by the determination of hexahydrophthalic acid in urine using gas chromatography and selected-ion monitoring 572(1991)117

5-(N,N-Hexamethylene)amiloride

Determination of active 5-(N,N-hexamethylene)amiloride in pig plasma 579(1992)376

Hexane

Activity coefficients in binary liquid mixtures measured by reversed-flow gas chromatography 595(1992)237

2,5-Hexanedione

Determination of 2,5-hexanedione in urine and serum by gas chromatography after derivatization with O-(pentafluorobenzyl)hydroxylamine and solid-phase extraction 563(1991)199

Mass spectrometric and gas and high-performance liquid chromatographic behaviour of an impurity in 2,5-hexanedione 590(1992)223

Rapid and sensitive determination of urinary 2,5-hexanedione by reversed-phase high-performance liquid chromatography 592(1992)255

Hexobarbital

Coupled column chromatography in chiral separations: systems employing β -cyclodextrin phases for chiral separation 557(1991)199

High-performance liquid chromatographic method for the simultaneous determination of *R*-(-)- and *S*-(+)-hexobarbital in rat plasma 567(1991)472

Retention and enantioselectivity of racemic solutes on a modified ovomucoid-bonded column. I. Cross-linking with glutaraldehyde 592(1992)301

Hexokinase

Purification and crystallization of yeast hexokinase isoenzymes. Characterization of different forms by chromatofocusing 587(1991)85

Hexoses

Carbohydrate separation by ligand-exchange liquid chromatography. Correlation between the formation of sugar-cation complexes and the elution order 558(1991)89

High-density lipoproteins

Separation of human serum lipoproteins into three major classes by hydroxyapatite chromatography 566(1991)67

Hippuric acid

Determination of benzyl alcohol and its metabolite in plasma by reversed-phase high-performance liquid chromatography 568(1991)145

Simultaneous high-performance liquid chromatographic determination of urinary metabolites of benzene, nitrobenzene, toluene, xylene and styrene 573(1992)318

Rapid and simple method for the determination of urinary benzoic and phenylacetic acids and their glycine conjugates in ruminants by reversed-phase high-performance liquid chromatography 582(1992)13

Hirudin

Liquid secondary ion mass spectrometry applied to structural confirmation of enzymically prepared C-terminal-truncated derivatives of recombinant hirudin 562(1991)421

Histamine

Measurement of histamine in rat bronchoalveolar lavage fluid by high-performance cation-exchange chromatography coupled with post-column derivatization to *o*-phthaldialdehyde 574(1992)340

Improvement of extraction procedure for biogenic amines in foods and their high-performance liquid chromatographic determination 591(1992)175

Primary and secondary amine derivatization with luminarins 1 and 2: separation by liquid chromatography with peroxyoxalate chemiluminescence detection 594(1992)386

High-performance liquid chromatography of histamine and 1-methylhistamine with on-column fluorescence derivatization 595(1992)163

Histamine N-methyltransferase

High-performance liquid chromatographic determination of histamine N-methyltransferase activity 567(1991)459

Histidine

Reversible reaction via a carbanion intermediate in the elimination of ammonia from L-histidine catalysed by histidine ammonia-lyase 562(1991)363

Simultaneous determination of stable isotopically labelled L-histidine and urocanic acid in human plasma by stable isotope dilution mass spectrometry 576(1992)213

Determination of carnosine and other biogenic imidazoles in equine plasma by isocratic reversed-phase ion-pair high-performance liquid chromatography 579(1992)45

Immobilized metal ion affinity electrophoresis. A study with several model proteins containing histidine 597(1992)357

Specific assay for endotoxin using immobilized histidine, *Limulus* amoebocyte lysate and a chromogenic substrate 597(1992)415

Histidine ammonia-lyase

Reversible reaction via a carbanion intermediate in the elimination of ammonia from L-histidine catalysed by histidine ammonia-lyase 562(1991)363

Histones

High-performance capillary electrophoresis of histones 559(1991)431

Homatropine

(2*R*,3*R*)-Dicyclohexyl tartrate as a chiral mobile phase additive 585(1991)35

Retention processes on α_1 -acid glycoprotein-bonded stationary phase 591(1992)55

(2*S*,3*S*)-Dicyclohexyl tartrate as mobile phase additive for the determination of the enantiomeric purity of (*S*)-atropine in tablets 592(1992)339

Homocysteine

Rapid high-performance liquid chromatographic assay for total homocysteine levels in human serum 565(1991)441

Measurement of total plasma and cerebrospinal fluid homocysteine by fluorescence following high-performance liquid chromatography and precolumn derivatization with *o*-phthalaldehyde 579(1992)55

Homocystine

High-performance liquid chromatographic method with fluorescence detection for the determination of total homocyst(e)ine in plasma 593(1992)171

Homovanillic acid

Analysis of rat brain microdialysate by gas chromatography–high-resolution selected-ion monitoring mass spectrometry 562(1991)111

New analyser for the determination of urinary vanillylmandelic acid, homovanillic acid and creatinine 567(1991)11

Simple and fast chromatographic method for the simultaneous determination of vanillylmandelic acid and homovanillic acid in human urine 570(1991)371

Improved method for the determination of 5,6-dihydroxytryptamine and 5,7-dihydroxytryptamine in tissue using high-performance liquid chromatography with electrochemical detection 574(1992)137

Simultaneous measurement of serotonin, catecholamines and their metabolites in cat and human plasma by *in vitro* microdialysis–microbore high-performance liquid chromatography with amperometric detection 582(1992)19

Determination of plasma homovanillic acid by liquid chromatography with electrochemical detection 582(1992)236

Honokiol

Identification and determination of honokiol and magnolol from *Magnolia officinalis* by high-performance liquid chromatography with photodiode-array UV detection 598(1992)143

Hopantenate

Abnormal fatty acid metabolism in patients in hopantenate therapy during clinical episodes 562(1991)139

Hopantenic acid

Analytical studies on the chiral separation and simultaneous determination of pantothenic acid and hopantenic acid enantiomers in rat plasma by gas chromatography–mass fragmentography 564(1991)1

Analytical studies on the chiral separation and simultaneous determination of pantothenic acid and hopantenic acid enantiomers in rat plasma by gas chromatography–mass fragmentography: a reply 576(1992)179

Analytical studies on the chiral separation and simultaneous determination of pantothenic acid and hopantenic acid enantiomers in rat plasma by gas chromatography–mass fragmentography. Reply to Küsters *et al.* 576(1992)181

Hormones

Computer-aided optimization of the experimental conditions for the isocratic reversed-phase high-performance liquid chromatographic separation of hormonal steroids 552(1991)103

Purification of canine prolactin and growth hormone by fast protein liquid chromatography 553(1991)117

Mathematical model describing dispersion in free solution capillary electrophoresis under stacking conditions 559(1991)3

Temperature elevations of the sample zone in free solution capillary electrophoresis under stacking conditions 559(1991)27

Radial pH distribution during capillary electrophoresis with electroosmotic flow. Analysis with high ionic strength buffers 589(1992)315

Validation of chromatographic retention models in reversed-phase high-performance liquid chromatography by fitting experimental data to the relevant equations 595(1992)53

Horseradish peroxidase

High-performance liquid chromatographic investigation of product formation in the horseradish peroxidase-enhanced chemiluminescence of luminol with different enhancers 566(1991)461

HSR-803

Simultaneous determination of a new gastrointestinal prokinetic agent (HSR-803) and its metabolites in human serum and urine by high-performance liquid chromatography using automated column-switching 576(1992)174

HT-2 toxin

Identification of trichothecenes by thermospray, plasmaspray and dynamic fast-atom bombardment liquid chromatography–mass spectrometry 562(1991)555

α -Human atrial natriuretic peptide

Construction and use of two α -human atrial natriuretic peptide-fragment affinity chromatography columns in the isolation of C- and N-terminal epitope-specific antibodies for use in a prototype α -hANP biosensor 577(1992)251

Human serum albumin

Effect of glycation on the heterogeneity of human serum albumin analysed by reversed-phase high-performance liquid chromatography in a solvent containing formic acid 573(1992)201

pH titration curves of the desialylated human α_1 -acid glycoprotein variants by combined isoelectrofocusing–electrophoresis: utilization in the development of a fractionation method for the protein variants by chromatography on immobilized metal affinity adsorbent 577(1992)43

Allosteric and competitive displacement of drugs from human serum albumin by octanoic acid, as revealed by high-performance liquid affinity chromatography, on a human serum albumin-based stationary phase 577(1992)305

Rapid analysis of human serum albumin by high-performance liquid chromatography 578(1992)292

Characterization of thyroxine–albumin binding using high-performance affinity chromatography. I. Interactions at the warfarin and indole sites of albumin 579(1992)225

Hyaluronic acid

Determination of human urinary hyaluronic acid, chondroitin sulphate and dermatan sulphate as their unsaturated disaccharides by high-performance liquid chromatography 565(1991)141

Assay of synovial fluid hyaluronic acid using high-performance liquid chromatography of hyaluronidase digests 566(1991)9

Chemiluminescence high-performance liquid chromatography for the determination of hyaluronic acid, chondroitin sulphate and dermatan sulphate 579(1992)203

Hydantoin

Chiral separation of barbiturates and hydantoins by reversed-phase high-performance liquid chromatography using a 25 or 50 mm short ODS cartridge column via β -cyclodextrin inclusion complexes 579(1992)253

Hydrazides

Electrochemical and chromatographic properties of selected hydrazine and hydrazide derivatives of carbonyl compounds 585(1991)225

Hydrazines

Simultaneous assay for isoniazid and hydrazine metabolite in plasma and cerebrospinal fluid in the rabbit 567(1991)261

Determination of hydrazine in biofluids by capillary gas chromatography with nitrogen-sensitive or mass spectrometric detection 573(1992)227

Electrochemical and chromatographic properties of selected hydrazine and hydrazide derivatives of carbonyl compounds 585(1991)225

Hydrocarbons

Gas chromatographic investigation of silica supports modified in a high-frequency low-temperature plasma 552(1991)179

Thermodesorption–gas chromatography–mass spectrometric analysis of biological materials for potential molecular precursors of the constituents of the crude oils 552(1991)325

Quantitative structure-type analysis of heterocompound–hydrocarbon mixtures. VI. Characterization of the nature of neutral heterocompounds in a product of brown coal refining 553(1991)441

Simultaneous separation and determination of hydrocarbons and organochlorine compounds by using a two-step microcolumn 555(1991)221

New solid adsorbents for the separation of lower hydrocarbons and permanent gases. II. Ammonium molybdophosphate 556(1991)425

Gas chromatography and gas chromatography–mass spectrometry study of hydrocarbons in Vlasta oil (Adriatic Basin) as the basis for geochemical interpretation 557(1991)255

Analysis of industrial solvent mixtures in water using a miniature purge-and-trap device with thermal desorption and capillary gas chromatography–mass spectrometry 557(1991)429

Gas chromatographic–mass spectrometric characterization of volatile organic compounds in Barcelona tap water 562(1991)481

Determination of aromatic hydrocarbons and their metabolites in human blood and urine (Review) 580(1992)229

Methods for the analysis of persistent chlorinated hydrocarbons in tissues (Review) 580(1992)257

Determination of specific retention volumes at 20°C for hydrocarbons on microporous carbons 586(1991)161

Quality control procedure for the gas chromatographic determination of light hydrocarbons in petroleum liquid 588(1991)259

Use of oxidative degradation followed by capillary gas chromatography–mass spectrometry and multi-dimensional scaling analysis to fingerprint unresolved complex mixtures of hydrocarbons 589(1992)281

Gas chromatography of Titan's atmosphere. III. Analysis of low-molecular-weight hydrocarbons and nitriles with a CP-Sil-5 CB WCOT capillary column 591(1992)297

Quantitative hydrocarbon group analysis of gasoline and diesel fuel by supercritical fluid chromatography 595(1992)301

Use of basic alumina in fractionation of fossil fuels 600(1992)243

Cholesteryl acetate as a stationary phase for the gas chromatography of some volatile oil constituents 600(1992)337

Hydrocortisone

Supercritical fluid chromatography in the routine stability control of antipruritic preparations 553(1991)71

Thermospray liquid chromatography-mass spectrometry of corticosteroids 577(1992)221

Hydrogen, organic

Flame infrared emission detection of hydrogen 589(1992)241

Hydrogencarbonate

Carbonate eluents in ion chromatography. Fractions of resin capacity linked to anions present in the eluent 589(1992)197

Hydrogen fluoride

Reaction-gas chromatography for physico-chemical investigations on aggressive gases 552(1991)319

Hydrogen isotopes

Gas chromatography for measurement of hydrogen isotopes at tritium processing 586(1991)131

Hydrogen peroxide

On-line electrochemical reagent generation for liquid chromatography with luminol-based chemiluminescence detection 557(1991)13

Hydrogen sulphide

Near-continuous measurement of hydrogen sulfide and carbonyl sulfide by an automatic gas chromatograph 585(1991)353

Preconcentration technique for introducing gaseous or volatile compounds into a capillary gas chromatographic column 587(1991)338

Hydromorphone

Simultaneous solid-phase extraction and chromatographic analysis of morphine and hydromorphone in plasma by high-performance liquid chromatography with electrochemical detection 577(1992)354

Hydroperoxides

Enantiomeric separation of racemic hydroperoxides and related alcohols 588(1991)352

High-performance liquid chromatography and post-column derivatization with diphenyl-1-pyrenylphosphine for fluorimetric determination of triacylglycerol hydroperoxides 596(1992)197

Hydroperoxy polyunsaturated fatty acids

High-performance liquid chromatography-thermospray mass spectrometry of hydroperoxy polyunsaturated fatty acid acetyl derivatives 579(1992)25

Hydroquinone

Hydroquinone oxidation kinetics in adsorptive liquid chromatographic beds 556(1991)383

Determination of benzene metabolites in urine of mice by solid-phase extraction and high-performance liquid chromatography 593(1992)147

Micellar electrokinetic chromatographic study of hydroquinone and some of its ethers. Determination of hydroquinone in skin-toning cream 596(1992)95

Hydroxamic acids

Rapid high-performance liquid chromatographic determination of fatty acid profiles of lipids by conversion to their hydroxamic acids 587(1991)292

Development of stability indicating assay methods for the determination of hydroxamic acids in topical formulations 594(1992)209

Hydroxide

Carbonate eluents in ion chromatography. Fractions of resin capacity linked to anions present in the eluent 589(1992)197

Hydroxylalprazolam

New high-performance liquid chromatographic method for the determination of alprazolam and its metabolites in serum: instability of 4-hydroxylalprazolam 568(1991)253

Hydroxyamines

Possible role for stationary phase metal interactions in the chromatography of hydroxyamines on silica 552(1991)429

1-Hydroxy-3-aminopropylidene-1,1-biphosphonate

Improved determination of the bisphosphonate pamidronate disodium in plasma and urine by pre-column derivatization with fluorecamine, high-performance liquid chromatography and fluorescence detection 568(1991)261

10-Hydroxyamitriptyline

Column switching and high-performance liquid chromatography in the analysis of amitriptyline, nortriptyline and hydroxylated metabolites in human plasma or serum 578(1992)273

High-performance liquid chromatographic determination of amitriptyline and its main metabolites using a silica column with reversed-phase eluent. Application in mice 584(1992)249

4-Hydroxyandrost-4-ene-3,17-dione

Determination of 4-hydroxyandrostenedione in plasma and urine by extractive alkylation and electron-capture gas chromatography 565(1991)67

Determination of 4-hydroxyandrost-4-ene-3,17-dione metabolism in breast cancer patients using high-performance liquid chromatography–mass spectrometry 565(1991)75

Identification of 4-hydroxyandrost-4-ene-3,17-dione metabolites in prostatic cancer patients by liquid chromatography–mass spectrometry 576(1992)235

3-Hydroxyanthranilic acid

High-performance liquid chromatographic assay of human lymphocyte kynureninase activity levels 566(1991)369

Difference in the concentration of tryptophan metabolites between maternal and umbilical foetal blood 576(1992)334

4-Hydroxyantipyrine, sulphoconjugate

Separation and identification of the 4-hydroxyantipyrine sulphoconjugate 576(1992)103

Hydroxy-1,3-benzenedicarboxaldehydes

Separation and preparative isolation of phenolic dialdehydes by on-line overpressured layer chromatography 590(1992)369

***p*-Hydroxybenzoic acid**

Simple direct liquid introduction system usable as an interface for liquid chromatography–mass spectrometry on quadrupole and magnetic-sector mass spectrometers 554(1991)293

Limits and effects of precolumn addition of thioglycerol in liquid chromatographic–fast atom bombardment mass spectrometric systems 591(1992)129

Hydroxy bile acids

Capillary gas chromatographic analysis of oxo, oxo-hydroxy and unsaturated bile acid glycine conjugates 558(1991)451

Hydroxybrodiprim

Determination of brodiprim and its hydroxy metabolite in human plasma, blood and urine by high-performance liquid chromatography 577(1992)123

4'-Hydroxybupivacaine

High-performance liquid chromatographic method for the quantitation of bupivacaine, 2,6-pipecoloxylidide and 4'-hydroxybupivacaine in plasma and urine 577(1992)103

Hydroxychloroquine

High-performance liquid chromatographic separation of the enantiomers of hydroxychloroquine and its major metabolites in biological fluids using an α_1 -acid glycoprotein stationary phase 570(1991)119

Determination of hydroxychloroquine and its major metabolites in plasma using sequential achiral–chiral high-performance liquid chromatography 573(1992)253

Analytical and semi-preparative high-performance liquid chromatographic separation and assay of hydroxychloroquine enantiomers 581(1992)83

14(*R*)-Hydroxyclearithromycin

Simultaneous determination of clarithromycin and 14(*R*)-hydroxyclearithromycin in plasma and urine using high-performance liquid chromatography with electrochemical detection 571(1991)199

***trans*-3'-Hydroxycotinine**

Simultaneous determination of cotinine and *trans*-3'-hydroxycotinine in human serum by high-performance liquid chromatography 577(1992)358

Determination of the nicotine metabolite *trans*-3'-hydroxycotinine in urine of smokers using gas chromatography with nitrogen-selective detection or selected ion monitoring 583(1992)145

7-Hydroxycoumarin

Comparison of a novel thin-layer chromatographic–fluorescence detection method with a spectrofluorometric method for the determination of 7-hydroxycoumarin in human urine 575(1992)325

Rapid and sensitive determination of coumarin and 7-hydroxycoumarin and its glucuronide conjugate in urine and plasma by high-performance liquid chromatography 582(1992)137

7-Hydroxycoumarin glucuronide

Rapid and sensitive determination of coumarin and 7-hydroxycoumarin and its glucuronide conjugate in urine and plasma by high-performance liquid chromatography 582(1992)137

Hydroxydebrisoquine

Determination of debrisoquine and metabolites in human urine by gas chromatography–mass spectrometry 570(1991)89

2-Hydroxydesipramine

Determination of imipramine, desipramine and their hydroxy metabolites by reversed-phase chromatography with ultraviolet and coulometric detection 572(1991)247

2-Hydroxydicarboxylic acids

Abnormal fatty acid metabolism in patients in hopantenate therapy during clinical episodes 562(1991)139

5-Hydroxy-5,6-dihydrothymidine

Reversed-phase high-performance liquid chromatography–thermospray mass spectrometry of radiation-induced decomposition products of thymine and thymidine 593(1992)133

5-Hydroxydiprafenone

Direct separation of the enantiomers of propafenone, diprafenone and their major metabolites by high-performance liquid chromatography on modified cellulose and amylose chiral stationary phases 585(1991)329

4-Hydroxydodecanedioic acids

Abnormal fatty acid metabolism in patients in hopantenate therapy during clinical episodes 562(1991)139

Hydroxyeicosatetraenoic acids

Gas chromatographic-mass spectrometric characterisation of some novel hydroxyeicosatetraenoic acids formed on incubation of arachidonic acid with microsomes from induced rat livers 562(1991)237

In vivo transformation of arachidonic acid into 12-hydroxy-5,8,10,14-eicosatetraenoic acid by human nasal mucosa 575(1992)143

Rapid method for automated on-line extraction and fractionation of plasma leukotrienes and 12-hydroxy-5,8,10,14-eicosatetraenoic acids by reversed-phase high-performance liquid chromatography 579(1992)37

Hydroxyestradiols

Determination of estradiol 2- and 4-hydroxylase activities by gas chromatography with electron-capture detection 567(1991)309

 β -Hydroxyethylapropfen

Isolation and identification of β -hydroxyethylapropfen: a urinary metabolite of apropfen in rats 563(1991)466

7-Hydroxyethyl biscoumacetate

Determination of ethyl biscoumacetate and its metabolite 7-hydroxyethyl biscoumacetate in human serum by high-performance liquid chromatography and mass spectrometry 574(1992)71

5R,6S,8R-6-(1-Hydroxyethyl)-2-carbamoyloxymethylpenem-3-carboxylic acid

High-performance liquid chromatographic methods for the determination of the penems SCH 29482 and FCE 22101 in human serum and urine 579(1992)115

5R,6R,8R-6-(1-Hydroxyethyl)-2-ethylthiopenem-3-carboxylic acid

High-performance liquid chromatographic methods for the determination of the penems SCH 29482 and FCE 22101 in human serum and urine 579(1992)115

5-(1-Hydroxyethyl)etoperidone

High-performance liquid chromatographic assay with ultraviolet detection for the determination of etoperidone and two active metabolites, 5-(1-hydroxyethyl)etoperidone and 1-(3-chlorophenyl)piperazine, in plasma 567(1991)433

D,L-3-Hydroxy-3-ethyl-3-phenylpropionamide

Plasma determination of the novel anticonvulsant D,L-3-hydroxy-3-ethyl-3-phenylpropionamide and preliminary pharmacokinetic studies in the rat 575(1992)306

Hydroxyethyl starch

Thin-layer electrophoresis of hydroxyethyl starches on a modified silica gel support 591(1992)351

1'-Hydroxyeugenol

Suitable chiral packing material for the high-performance liquid chromatographic separation of derivatives of 1'-hydroxyeugenol 600(1992)312

Hydroxy fatty acids

Liquid chromatography-mass spectrometry of hydroxy and non-hydroxy fatty acids as amide derivatives 575(1992)197

Configuration analysis of unsaturated hydroxy fatty acids 586(1991)255

Hydroxyglibenclamide

Determination of glibenclamide and its two major metabolites in human serum and urine by column liquid chromatography 564(1991)223

(6S,4R)-(-)-6-Hydroxy-4-(1-hydroxy-1-methylethyl)-1-cyclohexene-1-ethanol

Improved high-performance liquid chromatographic resolution of the geometric isomers of 6-hydroxy-4-(1-hydroxy-1-methylethyl)-1-cyclohexene-1-ethanol and by-products with β -cyclodextrin 553(1991)15

2-Hydroxyimipramine

Determination of imipramine, desipramine and their hydroxy metabolites by reversed-phase chromatography with ultraviolet and coulometric detection 572(1991)247

5-Hydroxyindoleacetic acid

Analysis of rat brain microdialysate by gas chromatography-high-resolution selected-ion monitoring mass spectrometry 562(1991)111

Simultaneous measurement of serotonin and 5-hydroxyindoleacetic acid in rat brain using a liquid chromatographic method with electrochemical detection 564(1991)250

Improved method for the determination of 5,6-dihydroxytryptamine and 5,7-dihydroxytryptamine in tissue using high-performance liquid chromatography with electrochemical detection 574(1992)137

Difference in the concentration of tryptophan metabolites between maternal and umbilical foetal blood 576(1992)334

γ -Aminobutyric acid, catecholamine and indoleamine determinations from the same brain region by high-performance liquid chromatography with electrochemical detection 579(1992)334

Simultaneous measurement of serotonin, catecholamines and their metabolites in cat and human plasma by *in vitro* microdialysis–microbore high-performance liquid chromatography with amperometric detection 582(1992)19

Microbore liquid chromatography with dual electrochemical detection for the determination of serotonin and 5-hydroxyindoleacetic acid in rat brain dialysates 582(1992)29

α -Hydroxyisobutyric acid

Study of isotachopheretic separation behaviour of metal cations by means of particle-induced X-ray emission. II. Separation efficiency of lanthanoid model mixture 589(1992)339

3-Hydroxykynurenine

Determination of 3-hydroxykynurenine in human brain and plasma by high-performance liquid chromatography with electrochemical detection. Increased concentrations in hepatic encephalopathy 565(1991)436

High-performance liquid chromatographic determination of 3-hydroxykynurenine with fluorimetric detection; comparison of preovulatory phase and postovulatory phase urinary excretion 570(1991)13

4 β -Hydroxylated bile acids

Studies on steroids. CCLIV. Gas chromatographic–mass spectrometric determination of 4- and 6-hydroxylated bile acids in human urine with negative ion chemical ionization detection 574(1992)1

Hydroxylauric acid

Determination of the cytochrome P-450 IV marker, ω -hydroxylauric acid, by high-performance liquid chromatography and fluorimetric detection 564(1991)266

Hydroxylysylpyridinoline

Determination of hydroxylysylpyridinoline in tissues and urine by high-performance liquid chromatography 563(1991)153

High-performance liquid chromatographic determination of 3-hydroxypyridinium derivatives as new markers of bone resorption 574(1992)255

5-Hydroxy-6-methoxyindole-2-carboxylic acid

Indolic urinary melanogens: separation and identification by gas chromatography with selected-ion monitoring mass spectrometry of 5-hydroxy-6-methoxyindole-2-carboxylic and 5-methoxy-6-hydroxyindole-2-carboxylic acids 570(1991)166

5-Hydroxymethylfurfural

Determination of 5-hydroxymethylfurfural by ion-exclusion chromatography with UV detection 593(1992)153

3-Hydroxy-3-methylglutaryl-coenzyme A

Simultaneous assay of the activities of two key enzymes in cholesterol metabolism by gas chromatography–mass spectrometry 565(1991)53

Hydroxymethylglutaryl coenzyme A reductase inhibitor

Method for separation and determination of lactone and hydroxy acid forms of a new HMG CoA reductase inhibitor (RG 12561) in plasma 567(1991)491

Determination of the novel hydroxymethyl glutaryl coenzyme A reductase inhibitor (RP 61969) and its dihydroxy acid hydrolysis product in human plasma by reversed-phase high-performance liquid chromatography 575(1992)319

Hydroxymethylmexiletine

Simultaneous determination of mexiletine and four hydroxylated metabolites in human serum by high-performance liquid chromatography and its application to pharmacokinetic studies 573(1992)235

***p*-Hydroxymexiletine**

Simultaneous determination of mexiletine and four hydroxylated metabolites in human serum by high-performance liquid chromatography and its application to pharmacokinetic studies 573(1992)235

Hydroxymidazolam

Solid-phase extraction of midazolam and two of its metabolites from plasma for high-performance liquid chromatographic analysis 571(1991)298

9-Hydroxymincycline

Identification and determination of the two principal metabolites of minocycline in humans 568(1991)363

6-Hydroxynaphthalene-2-sulphonic acid

Extension of the electrostatic retention model of reversed-phase ion-pair high-performance liquid chromatography to include the effect of the eluent pH 552(1991)13

Hydroxynefazodone

High-performance liquid chromatographic method for the determination of nefazodone and its metabolites in human plasma using laboratory robotics 570(1991)129

Hydroxynefracetam

Simultaneous determination of nefracetam and its metabolites by high-performance liquid chromatography 575(1992)261

4-Hydroxy-2-nonenal

Trace quantitation of 4-hydroxy-2-nonenal in biological samples as its oxime–bis-*tert*-butyldimethylsilyl derivative using 3-hydroxynonanal as an internal standard 578(1992)9

10-Hydroxynortriptyline

Sensitive method for the quantitation of nortriptyline and 10-hydroxynortriptyline in human plasma by capillary gas chromatography with electron-capture detection 573(1992)141

Column switching and high-performance liquid chromatography in the analysis of amitriptyline, nortriptyline and hydroxylated metabolites in human plasma or serum 578(1992)273

High-performance liquid chromatographic determination of amitriptyline and its main metabolites using a silica column with reversed-phase eluent. Application in mice 584(1992)249

Hydroxyomeprazole

Simultaneous determination of omeprazole and its metabolites in plasma and urine by reversed-phase high-performance liquid chromatography with an alkaline-resistant polymer-coated C₁₈ column 579(1992)299

***p*-Hydroxyphenylacetamide**

A high-performance liquid chromatographic method for the monitoring and quantification of the synthesis of *p*-hydroxyphenylacetamide 587(1991)280

***p*-Hydroxyphenylphenylhydantoin**

High-performance liquid chromatographic method for direct separation of 5-(*p*-hydroxyphenyl)-5-phenylhydantoin enantiomers using a chiral tris(4-methylbenzoate) column 568(1991)157

Two-dimensional high-performance liquid chromatographic method to assay *p*-hydroxyphenylphenylhydantoin enantiomers in biological fluids and stereoselectivity of enzyme induction in phenytoin metabolism 575(1992)109

Chiral separation of barbiturates and hydantoins by reversed-phase high-performance liquid chromatography using a 25 or 50 mm short ODS cartridge column via β -cyclodextrin inclusion complexes 579(1992)253

***p*-Hydroxyphenylpyruvic acid**

High-performance liquid chromatographic separation of *p*-hydroxyphenylpyruvic acid 585(1991)318

***p*-Hydroxypiribedil**

Determination of piribedil and its basic metabolites in plasma by high-performance liquid chromatography 563(1991)323

5-Hydroxypiroxicam

Determination of piroxicam and its major metabolites in the plasma, urine and bile of humans by high-performance liquid chromatography 576(1992)121

5-Hydroxypiroxicam glucuronide

Determination of piroxicam and its major metabolites in the plasma, urine and bile of humans by high-performance liquid chromatography 576(1992)121

Hydroxy-polyunsaturated fatty acids

High-performance liquid chromatography-thermospray mass spectrometry of hydroxy-polyunsaturated fatty acid acetyl derivatives 575(1992)7

Hydroxyprednisolones

Simultaneous analysis of prednisone, prednisolone and their major hydroxylated metabolites in urine by high-performance liquid chromatography 567(1991)39

20 β -Hydroxyprednisone

Simultaneous analysis of prednisone, prednisolone and their major hydroxylated metabolites in urine by high-performance liquid chromatography 567(1991)39

17-Hydroxyprogesterone

Thin-layer chromatography on silica-coated aluminium sheet as an adjunct to radioimmunoassay of steroids 563(1991)178

Hydroxyproline

Microderivatization of 4-[¹⁸O]hydroxyproline and quantitation with a benchtop mass spectrometer 562(1991)625

5-Hydroxypropafenone

Direct separation of the enantiomers of propafenone, diprafenone and their major metabolites by high-performance liquid chromatography on modified cellulose and amylose chiral stationary phases 585(1991)329

Hydroxypropylmethylcellulose

Isotachopheresis of proteins in uncoated open-tubular fused-silica capillaries with a simple approach for column conditioning 558(1991)423

Capillary isoelectric focusing with electroosmotic zone displacement and on-column multichannel detection 589(1992)321

1-Hydroxypyrene

Coupled-column high-performance liquid chromatographic method for the determination of 1-hydroxypyrene in urine of subjects exposed to polycyclic aromatic hydrocarbons 600(1992)189

3-Hydroxypyridinium derivatives

High-performance liquid chromatographic determination of 3-hydroxypyridinium derivatives as new markers of bone resorption 574(1992)255

3-Hydroxyquinidine

High-performance liquid chromatographic method for the quantitation of quinidine and selected quinidine metabolites 572(1991)159

9-Hydroxyrisperidone

Determination of risperidone and 9-hydroxyrisperidone in plasma, urine and animal tissues by high-performance liquid chromatography 583(1992)223

17 β -Hydroxysteroid dehydrogenase

Rapid purification yielding highly active 17 β -hydroxysteroid dehydrogenase: application of hydrophobic interaction and affinity fast protein liquid chromatography 582(1992)71

5-Hydroxysulphadimidine

High-performance liquid chromatographic method for the routine determination of sulphadimidine, its hydroxy metabolites and N4-acetylsulphadimidine in body fluids and cell culture media 571(1991)157

4-Hydroxytetradecanedioic acids

Abnormal fatty acid metabolism in patients in hopantenate therapy during clinical episodes 562(1991)139

Hydroxytrimethoprim

Determination of trimethoprim and its oxidative metabolites in cell culture media and microsomal incubation mixtures by high-performance liquid chromatography 579(1992)355

2-Hydroxytrimipramine

Simultaneous determination of trimipramine and its major metabolites by high-performance liquid chromatography 566(1991)228

5-Hydroxytryptamine

see Serotonin

5-Hydroxytryptophan

High-performance liquid chromatographic procedure for the simultaneous determination of aromatic L-amino acid decarboxylase activity towards 3,4-dihydroxyphenylalanine and 5-hydroxytryptophan 567(1991)381

5-Hydroxytryptophol

Determination of urinary 5-hydroxytryptophol by high-performance liquid chromatography with electrochemical detection 579(1992)340

Hydroxyurea

Determination by high-performance liquid chromatography of hydroxyurea in human plasma 584(1992)270

Hydroxyvecuronium

High-performance liquid chromatography-electrochemical detection of vecuronium and its metabolites in human plasma 573(1992)79

Hydroxyyohimbines

Determination of yohimbine and its two hydroxylated metabolites in humans by high-performance liquid chromatography and mass spectral analysis 574(1992)283

Hygromycin

Determination of aminoglycoside antibiotics by reversed-phase ion-pair high-performance liquid chromatography coupled with pulsed amperometry and ion spray mass spectrometry 591(1992)195

Hyoscyamine

Analysis of tropane alkaloids with thermospray high-performance liquid chromatography-mass spectrometry 562(1991)737

Hypoxanthine

Quantitation of hypoxanthine in plasma from patients with ischemic heart disease: adaption of a high-performance liquid chromatographic method 568(1991)105

Simultaneous determination of allantoin and oxypurines in biological fluids by high-performance liquid chromatography 575(1992)153

Determination of concentrations of adenosine and other purines in human term placenta by reversed-phase high-performance liquid chromatography with photodiode-array detection: evidence for pathways of purine metabolism in the placenta 575(1992)243

I**Ibuprofen**

(*S,S*)-Diphenylethylethanediamine derivatives as chiral selectors. II. Gasparrini-type bound chiral stationary phase with high enantioselectivity for naphthylamides 553(1991)373

Simple direct liquid introduction system usable as an interface for liquid chromatography-mass spectrometry on quadrupole and magnetic-sector mass spectrometers 554(1991)293

Liquid chromatographic determination of the enantiomers of ibuprofen in plasma using a chiral AGP column 563(1991)414

Column-switching techniques for high-performance liquid chromatography of ibuprofen and mefenamic acid in human serum with short-wavelength ultraviolet detection 570(1991)329

High-performance liquid chromatographic determination of naproxen, ibuprofen and diclofenac in plasma and synovial fluid in man 578(1992)251

Improved high-performance liquid chromatographic assay method for the enantiomers of ibuprofen 583(1992)259

Enantiospecific gas chromatographic-mass spectrometric procedure for the determination of ketoprofen and ibuprofen in synovial fluid and plasma: application to protein binding studies 584(1992)189

Retention and enantioselectivity of racemic solutes on a modified ovomucoid-bonded column. I. Cross-linking with glutaraldehyde 592(1992)301

Icajine

Separation of *Strychnos nux-vomica* alkaloids by high-performance liquid chromatography 587(1991)318

ICI 204,636

Determination of an antipsychotic agent (ICI 204,636) and its 7-hydroxy metabolite in human plasma by high-performance liquid chromatography and gas chromatography-mass spectrometry 573(1992)49

Idaverine

Reversed-phase liquid chromatography coupled on-line with capillary gas chromatography. I. Introduction of large volumes of aqueous mixtures through an on-column interface 552(1991)489

Idebenone

Determination of idebenone in rat serum and brain by high-performance liquid chromatography using platinum catalyst reduction and electrochemical detection 573(1992)154

Ifosfamide

Determination of urinary 2- and 3-dechloroethylated metabolites of ifosfamide by high-performance liquid chromatography 570(1991)351

Gas chromatographic determination of ifosfamide in microvolumes of urine and plasma 571(1991)121

Combined thin-layer chromatography–photography–densitometry for the quantification of ifosfamide and its principal metabolites in urine, cerebrospinal fluid and plasma 575(1992)137

Gas chromatographic determination of 2- and 3-dechloroethylifosfamide in plasma and urine 583(1992)175

Imidapril

Determination of three metabolites of a new angiotensin-converting enzyme inhibitor, imidapril, in plasma and urine by gas chromatography–mass spectrometry using multiple ion detection 581(1992)65

Imidazoleamines

Application of high-performance liquid chromatography to the study of biogenic amine-related enzymes (Review) 566(1991)287

Imidazoles

Analysis of novel imidazoles from isolated perfused rabbit heart by two high-performance liquid chromatographic methods 577(1992)347

Determination of carnosine and other biogenic imidazoles in equine plasma by isocratic reversed-phase ion-pair high-performance liquid chromatography 579(1992)45

Imidazolium iodides

Use of non-volatile ion-pairing agents for liquid chromatographic–mass spectrometric analyses with a moving-belt interface 554(1991)281

Imidocarb

High-performance liquid chromatographic determination of imidocarb in cattle kidney with cation-exchange clean-up 577(1992)376

Iminodipeptides

Liquid chromatography–mass spectrometry for simultaneous analyses of iminodipeptides containing an N-terminal or a C-terminal proline 565(1991)408

Imipramine

High-performance liquid chromatography on dynamically modified silica. IX. Modification of silica with 3-(N,N-dimethylpalmitylammonium) propanesulphonate for reversed-phase chromatography 556(1991)353

Effect of stationary phase structure on retention and selectivity of restricted-access reversed-phase packing materials 558(1991)19

Determination of imipramine, desipramine and their hydroxy metabolites by reversed-phase chromatography with ultraviolet and coulometric detection 572(1991)247

Retention reproducibility of basic drugs in high-performance liquid chromatography on a silica column with a methanol–high-pH buffer eluent. Changes in selectivity with the age of the stationary phase 592(1992)85

Imirestat

Capillary gas chromatographic–electron-capture assay for the aldose reductase inhibitor imirestat in lens and plasma 565(1991)255

Immobilines

Capillary zone electrophoresis analysis of acrylamido buffers for isoelectric focusing in immobilized pH gradients 559(1991)119

Immobilized antibodies

Factors affecting the specific activity of immobilized antibodies and their biologically active fragments 572(1991)85

Immune complexes

Chromatographic and electrophoretic studies of circulating immune complexes in plasma 566(1991)77

Immunoabsorbents

Immunoabsorbents for clinical use: *ex vivo* immunoglobulin E removal in allergy 563(1991)166

Immunoglobulins

Bead cellulose derivatives as supports for immobilization and chromatographic purification of proteins 552(1991)389

High-performance capillary electrophoresis as a fast in-process control method for enzyme-labelled monoclonal antibody conjugates 559(1991)385

Immunoabsorbents for clinical use: *ex vivo* immunoglobulin E removal in allergy 563(1991)166

Chromatographic methods for large-scale preparation of immunoglobulin G_{2a} monoclonal antibodies Lym-1 and TNT-1 F(ab')₂ fragments 563(1991)243

Electrophoretic analysis of enzyme-linked immunoglobulins and their clinical significance (Review) 569(1991)347

Study of the separation of mouse monoclonal antibodies by pseudobioaffinity chromatography using matrix-linked histidine and histamine 570(1991)29

Factors affecting the specific activity of immobilized antibodies and their biologically active fragments 572(1991)85

Ligand location and biochemical productivity of silica-based immunoaffinity adsorbents 584(1992)23

Separation of immunoglobulin G by high-performance pseudo-bioaffinity chromatography with immobilized histidine. I. Preliminary report on the influence of the silica support and the coupling mode 584(1992)35

Rapid, two-step purification process for the preparation of pyrogen-free murine immunoglobulin G₁ monoclonal antibodies 590(1992)255

Separation techniques for biotechnology in the 1990s 592(1992)3

High-performance gel permeation chromatographic analysis of immunoglobulin M produced by hybridoma cell culture 594(1992)179

Avid AL, a synthetic ligand affinity gel mimicking immobilized bacterial antibody receptor for purification of immunoglobulin G 597(1992)101

Evaluation of the mode of binding of immunoglobulin to activated agarose 597(1992)231

Affinity chromatographic purification of immunoglobulin M antibodies utilizing immobilized mannan binding protein 597(1992)247

Artificial immunoglobulin G-binding protein mimetic to staphylococcal protein A. Its production and application to affinity purification of immunoglobulin G 597(1992)277

Separation of monoclonal antibodies from antihapten antisera by two-dimensional affinity electrophoresis 597(1992)365

Indicaxanthin

High-performance liquid chromatographic analysis of the pigments of blood-red prickly pear (*Opuntia ficus indica*) 593(1992)177

Indole

Characterization of thyroxine-albumin binding using high-performance affinity chromatography. I. Interactions at the warfarin and indole sites of albumin 579(1992)225

Indoleacetic acid

Difference in the concentration of tryptophan metabolites between maternal and umbilical foetal blood 576(1992)334

3-Indoleacetonitrile

Identification of enzymatic degradation products from synthesized glucobrassicin by gas chromatography-mass spectrometry 586(1991)166

Indole alkaloids

Analysis and isolation of indole alkaloids of fungi by high-performance liquid chromatography 593(1992)201

Indoleamines

Application of high-performance liquid chromatography to the study of biogenic amine-related enzymes (Review) 566(1991)287

Indole compounds

Studies on stress metabolites. XVI. High-performance liquid chromatographic analysis of cruciferous phytoalexins using complex ternary mobile phase gradients 598(1992)147

Indole-2,3-dione

Isatin (indole-2,3-dione) in urine and tissues. Detection and determination by gas chromatography-mass spectrometry 562(1991)279

Indolelactic acid

Difference in the concentration of tryptophan metabolites between maternal and umbilical foetal blood 576(1992)334

3-Indolemethanol

Identification of enzymatic degradation products from synthesized glucobrassicin by gas chromatography-mass spectrometry 586(1991)166

Indolepropionic acid

Difference in the concentration of tryptophan metabolites between maternal and umbilical foetal blood 576(1992)334

Indomethacin

Simultaneous analysis of flunixin, naproxen, ethacrynic acid, indomethacin, phenylbutazone, mefenamic acid and thiosalicylic acid in plasma and urine by high-performance liquid chromatography and gas chromatography-mass spectrometry 568(1991)351

Solid-phase extraction for the high-performance liquid chromatographic determination of indomethacin, suxibuzone, phenylbutazone and oxyphenbutazone in plasma, avoiding degradation of compounds 581(1992)101

Inorganic acids

Ion chromatographic determination of anions, especially sulphur-containing anions, with conductimetric and kinetic detection 558(1991)209

Inorganic anions

Potentiometric detection of carboxylic acids and inorganic anions in ion-exclusion chromatography using camphorsulphonic acid as eluent 589(1992)209

Influence of poly(vinylpyrrolidone) on isotachophoretic separations of inorganic anions in aqueous electrolyte systems 589(1992)329

Application of a pellicular anion-exchange resin to the separation of inorganic and organic anions by single-column anion exchange 592(1992)101

Determination of inorganic anions in salt solutions by ion chromatography using C_{18} reversed-phase columns coated with cetyltrimethylammonium 598(1992)237

Inosine

Determination of concentrations of adenosine and other purines in human term placenta by reversed-phase high-performance liquid chromatography with photodiode-array detection: evidence for pathways of purine metabolism in the placenta 575(1992)243

Enzymophoresis of nucleic acids by tandem capillary enzyme reactor-capillary zone electrophoresis 596(1992)251

Inositol lipids

Quantitative analysis of inositol lipids and inositol phosphates in synaptosomes and microvessels by column chromatography: comparison of the mass analysis and the radiolabelling methods 581(1992)1

Inositol phosphates

Rapid and selective isolation of radiolabelled inositol phosphates from cancer cells using solid-phase extraction 574(1992)150

Quantitative analysis of inositol lipids and inositol phosphates in synaptosomes and microvessels by column chromatography: comparison of the mass analysis and the radiolabelling methods 581(1992)1

Preparation of samples for high-performance liquid chromatography of inositol phosphates 581(1992)134

Inositol polyphosphates

Determination of inositol polyphosphates from human T-lymphocyte cell lines by anion-exchange high-performance liquid chromatography and post-column derivatization 593(1992)157

Insect hormones

Simple method for collecting volatile compounds from single insects and other point sources for gas chromatographic analysis 598(1992)303

Insecticides

see Pesticides

Insulin

Instrumentation for high-performance capillary electrophoresis-mass spectrometry 559(1991)197

Influence of operating parameters on the preparative gradient elution chromatography of insulins 599(1992)195

Insulin-like growth factors

Primary sequences of insulin-like growth factors 1 and 2 isolated from porcine plasma 562(1991)391

Biospecific interaction analysis using surface plasmon resonance detection applied to kinetic, binding site and concentration analysis 597(1992)397

Interferon

Assessment of capillary electrophoresis in pharmaceutical applications. Analysis and quantification of a recombinant cytokine in an injectable dosage form 559(1991)307

Structural analysis of biologically active peptides and recombinant proteins and their modified counterparts by mass spectrometry 562(1991)377

Immobilization of monoclonal antibodies for affinity chromatography using a chelating peptide 595(1992)113

Kinetic aspects of membrane-based immunoaffinity chromatography 597(1992)167

Assay of protein drug substances present in solution mixtures by fluorescence derivatization and capillary electrophoresis 598(1992)123

High-performance hydrophobic interaction chromatography as a tool for protein refolding 599(1992)185

Purification of recombinant human interferon- β by immobilized antisense peptides 600(1992)235

Interleukin-2

On-line characterization of polyethylene glycol-modified proteins 588(1991)125

Membrane-based receptor affinity chromatography 597(1992)155

Kinetic aspects of membrane-based immunoaffinity chromatography 597(1992)167

Interleukin-2 receptor

Kinetic aspects of membrane-based immunoaffinity chromatography 597(1992)167

Interleukin-3

Mass spectrometric determination of glycosylation sites and oligosaccharide composition of insect-expressed mouse interleukin-3 562(1991)403

Iodate

Determination of inorganic anions in salt solutions by ion chromatography using C_{18} reversed-phase columns coated with cetyltrimethylammonium 598(1992)237

Iodide

Mixed potential response at metallic electrodes in potentiometric detection for ion chromatography 589(1992)201

Iodixanol

Simple isocratic high-performance liquid chromatographic method for measurement of iodixanol in human plasma 572(1991)333

N-(Iodoacetylaminoethyl)-5-naphthylamine-1-sulphonic acid

High-performance liquid chromatographic determination of selenocysteine with the fluorescent reagent, N-(iodoacetylaminoethyl)-5-naphthylamine-1-sulfonic acid 576(1992)263

Iodobenzoic acids

Selected applications of cyclodextrin selectors in capillary electrophoresis 559(1991)215

Iodothyronines

Method for the quantitation of iodothyronines in body tissues and fluids using high-performance liquid chromatography 577(1992)19

Iodotyrosines

Method for the quantitation of iodothyronines in body tissues and fluids using high-performance liquid chromatography 577(1992)19

5-Iodouracil

Determination of 2'-deoxy-5-iodouridine and its metabolite 5-iodouracil by high-performance liquid chromatography with ultraviolet absorbance detection in human serum 572(1991)327

Ions

Theoretical background for clinical and biomedical applications of electromigration techniques (Review) 569(1991)3

Iopentol

Automated high-performance liquid chromatography of iopentol in human plasma and whole blood using on-line dialysis as sample preparation 575(1992)93

Iothalamic acid

High-performance liquid chromatographic determination of iothalamic acid in human plasma and urine 570(1991)435

Iprodione

Development of a rational optimisation procedure for the automated sample clean-up with column switching in pesticide residue analysis 552(1991)113

Irganox 565

Assessment of a thermospray interface for the coupling of high-performance liquid chromatography and Fourier transform infrared spectrometry 588(1991)15

Irganox 1076

Ionization mechanisms in capillary supercritical fluid chromatography-chemical ionization mass spectrometry 554(1991)351

Iron

Separation of metal complexes of ethylenediaminetetraacetic acid in environmental water samples by ion chromatography with UV and potentiometric detection 558(1991)181

Separation and determination of trace amounts of vanadium (V), chromium(III) and iron(III) with 2-(2-thienylazo)-5-diethylaminophenol chelates by high-performance liquid chromatography 598(1992)298

Iron γ -oxide

Investigation of solid surfaces by high-performance liquid chromatography 592(1992)121

Iron(II)-1,10-phenanthroline

Separation and indirect detection of amino acids as acetylated derivatives 557(1991)315

Iron powder

Determination of succinic acid by pyrolysis-gas chromatography in the presence of sodium sulfite and iron powder 585(1991)345

Isatin

Isatin (indole-2,3-dione) in urine and tissues. Detection and determination by gas chromatography-mass spectrometry 562(1991)279

Isbufylline

Simultaneous determination of isbufylline and its major metabolites in rabbit blood and urine by reversed-phase high-performance liquid chromatography 568(1991)407

Isoajmaline

Monitoring of ajmaline in plasma with high-performance liquid chromatography 575(1992)87

Isobetanin

High-performance liquid chromatographic analysis of the pigments of blood-red prickly pear (*Opuntia ficus indica*) 593(1992)177

IsoBOSINC

Photosensitizers for tumor therapy: determination of bis(di-isobutyl octadecylsiloxy)silicon 2,3-naphthalocyanine (isoBOSINC) in rat tissue and serum by high-performance liquid chromatography 568(1991)437

Isodesmosine

Separation of elastin cross-links as phenylisothiocyanate derivatives 553(1991)93

High-performance liquid chromatographic determination of desmosine and isodesmosine after phenylisothiocyanate derivatization 572(1991)312

Isoenzymes

Purification of carbonic anhydrase isoenzymes by high-performance affinity chromatography and hydrophobic interaction chromatography 566(1991)351

Electrophoresis of serum isoenzymes and proteins following acute myocardial infarction (Review) 569(1991)323

Determination of alkaline phosphatase isoenzymes in serum by high-performance liquid chromatography with post-column reaction detection 576(1992)79

Purification of glycogen phosphorylase isozymes by metal-affinity chromatography 584(1992)77

Isoidide mononitrate

Gas chromatographic analysis of isomeric organic mononitrates in plasma 571(1991)109

Isolectins

Isolation and characterization of isolectins from *Erythrina variegata* seeds 597(1992)207

Isolongifolol

Use of a Sep-Pak cartridge for preparative collection in gas chromatography 555(1991)302

Isomannide mononitrate

Gas chromatographic analysis of isomeric organic mononitrates in plasma 571(1991)109

Isoniazid

Simultaneous assay for isoniazid and hydrazine metabolite in plasma and cerebrospinal fluid in the rabbit 567(1991)261

Isonicotinic acid methyl ester

Assessment of the biotransformation of the cardiotoxic agent piroximone by high-performance liquid chromatography and gas chromatography-mass spectrometry 593(1992)1

Isopentane

Study of microporous carbons by gas chromatographic determination of heats of physisorption 555(1991)183

Isopilcarpine

High-performance liquid chromatographic determination of pilocarpine hydrochloride and its degradation products using a β -cyclodextrin column 589(1992)159

Isoprene

Method for the collection and analysis of volatile compounds in the breath 564(1991)242

Liquid chromatographic isolation and structural elucidation of methoxymethylene dimethyl dodecatrienes catalytically synthesized from methanol and three moles of isoprene 594(1992)111

Isoprenoids

Novel long-chain *anteiso*-alkanes and *anteiso*-alkanoic acids in Antarctic rocks colonized by living and fossil cryptoendolithic microorganisms 598(1992)267

Isopropalin

High-performance liquid chromatographic determination of dinitroaniline herbicides in soil and water 585(1991)164

2-Isopropoxyphenol

Determination of 2-isopropoxyphenol in urine by capillary gas chromatography and mass-selective detection 578(1992)189

Isopropylbenzenes

High-performance liquid chromatographic determination of products of autoxidation of isopropylbenzene derivatives 588(1991)33

Isopropylphenols

High-performance liquid chromatography of isopropylphenols 595(1992)337

N-Isopropyltropinium

Determination of the quaternary compound ciclotropium in human biological material after hydrolysis and derivatization with the fluorophor flunoxapofen chloride 572(1991)181

Isorhamnetin-3-O-rutinoside

High-performance liquid chromatographic analysis of flavonol glycosides of *Solidago virgaurea* 558(1991)296

Isorhoeadine

Analytical studies of isorhoeadine and rheoagenine in petal extracts of *Papaver rhoeas* L. using high-performance liquid chromatography 596(1992)276

Isoricinoleic acid, methyl ester

Configuration analysis of unsaturated hydroxy fatty acids 586(1991)255

Isosorbide dinitrate

Clean-up of plasma extracts by gel permeation chromatography during analysis of isosorbide nitrates by capillary gas chromatography 553(1991)21

Isosorbide mononitrate

Gas chromatographic analysis of isomeric organic mononitrates in plasma 571(1991)109

Isotopically labelled compounds

Measurements using gas chromatography with coelution and dual-isotope atomic emission detection 586(1991)309

Isosuprine

Determination of isosuprine in equine plasma by high-performance liquid chromatography with electrochemical detection 563(1991)216

Isozymogens

High-performance liquid chromatography: purification and chromatographic behaviour of molecular variants of pepsinogen A from human urine 571(1991)47

Isradipine

Determination of isradipine and its pyridine metabolite in serum by capillary column gas chromatography with nitrogen-selective detection 574(1992)161

J**Josamycin**

Sensitive determination of josamycin and rokitamycin in plasma by high-performance liquid chromatography with fluorescence detection 575(1992)171

K**Kaempferol-3-O-robinobioside**

High-performance liquid chromatographic analysis of flavonol glycosides of *Solidago virgaurea* 558(1991)296

Kaempferol-3-O-rutinoside

High-performance liquid chromatographic analysis of flavonol glycosides of *Solidago virgaurea* 558(1991)296

Kallikrein

Direct determination of kallikrein by high-performance liquid chromatography 593(1992)119

Kanamycin

High-performance liquid chromatography method for the determination of aminoglycosides based on automated pre-column derivatization with *o*-phthalaldehyde 593(1992)69

Kappa light-chain

Immunoaffinity chromatography in biorecovery: an application of recombinant DNA technology to generic adsorption processes 584(1992)59

D,L-Kawain

Mass spectral characterization of urinary metabolites of D,L-kawain 562(1991)207

Ketamine

Stereoselective high-performance liquid chromatographic determination of the enantiomers of ketamine and norketamine in plasma 568(1991)165

Ketanserin

Rapid method for the determination of ketanserin in rat serum by high-performance liquid chromatography with fluorimetric detection 571(1991)318

Keto acids

Analysis of keto acids as their methyl esters of 2,4-dinitrophenylhydrazone derivatives by gas chromatography and gas chromatography-mass spectrometry 587(1991)247

2-Ketoalkanes

Computer simulation of gradient elution separation. Accuracy of predictions for non-linear gradients 555(1991)1

Ketodiols

Use of methyl oxime derivatives to enhance structural information in thermospray high-performance liquid chromatography-mass spectrometry. Analysis of linoleic acid lipoxygenase metabolites in maize embryos 554(1991)155

Ketogenesis metabolites

Simple cation-exchange high-performance liquid chromatography optimized to the measurement of metabolites in the effluents from perfused rat livers using refractive index and ultraviolet detectors 563(1991)231

Ketoisocaproic acid

Measurement of leucine and α -ketoisocaproic acid fluxes in the fetal/placental unit 562(1991)169

Reversed-phase high-performance liquid chromatographic analysis of branched-chain keto acid hydrazone derivatives: optimization of techniques and application to branched-chain keto acid balance studies across the forearm 572(1991)11

Ketoisovaleric acid

Reversed-phase high-performance liquid chromatographic analysis of branched-chain keto acid hydrazone derivatives: optimization of techniques and application to branched-chain keto acid balance studies across the forearm 572(1991)11

Ketols

Use of methyl oxime derivatives to enhance structural information in thermospray high-performance liquid chromatography-mass spectrometry. Analysis of linoleic acid lipoxygenase metabolites in maize embryos 554(1991)155

Ketomethylvaleric acid

Reversed-phase high-performance liquid chromatographic analysis of branched-chain keto acid hydrazone derivatives: optimization of techniques and application to branched-chain keto acid balance studies across the forearm 572(1991)11

Ketones

Quantitative structure-type analysis of heterocompound-hydrocarbon mixtures. VI. Characterization of the nature of neutral heterocompounds in a product of brown coal refining 553(1991)441

Dehydrogenase-silica as a stationary phase for the separation of alcohols and ketones 587(1991)268

Peak purity, yield and throughput in preparative liquid chromatography with self-displacement 592(1992)17

High-performance liquid chromatographic separation of chiral metallocenic ketones and alcohols 600(1992)320

Ketopelenolid-A

Use of high-performance liquid chromatographic peak deconvolution and peak labelling to identify antiparasitic components in plant extracts 593(1992)209

Ketoprofen

Simultaneous determination of ketoprofen enantiomers and probenecid in plasma and urine by high-performance liquid chromatography 568(1991)187

Enantiospecific analysis of ketoprofen in plasma by high-performance liquid chromatography 570(1991)446

Identification of a benzhydrolic metabolite of ketoprofen in horses by gas chromatography-mass spectrometry and high-performance liquid chromatography 583(1992)167

Enantiospecific gas chromatographic-mass spectrometric procedure for the determination of ketoprofen and ibuprofen in synovial fluid and plasma: application to protein binding studies 584(1992)189

Retention and enantioselectivity of racemic solutes on a modified ovomucoid-bonded column. I. Cross-linking with glutaraldehyde 592(1992)301

Ketoprofen, benzhydrolic metabolite

Identification of a benzhydrolic metabolite of ketoprofen in horses by gas chromatography-mass spectrometry and high-performance liquid chromatography 583(1992)167

Ketosteroids

Electrochemical and chromatographic properties of selected hydrazine and hydrazide derivatives of carbonyl compounds 585(1991)225

Ketotifen

Selected applications of cyclodextrin selectors in capillary electrophoresis 559(1991)215

Ketotrilostane

High-performance liquid chromatographic analysis of trilostane and ketotrilostane in rat plasma 567(1991)282

Khellacetone

Comparison of selectivity of silica and Florisil in the separation of natural pyranocoumarins 552(1991)453

Klenow polymerase

Characterization of synthetic macroporous packing materials in low-pressure cartridges and columns 590(1992)83

KW 2149

Determination of the new anticancer agent KW 2149, 7-N-[2-((2-(γ -L-glutamylamino)ethyl)dithio)ethyl]mitomycin C, an analogue of mitomycin C 564(1991)352

Kynureninase

High-performance liquid chromatographic assay of human lymphocyte kynureninase activity levels 566(1991)369

Kynurenine

Difference in the concentration of tryptophan metabolites between maternal and umbilical foetal blood 576(1992)334

L**L-365,260**

Quantitation of a new cholecystokinin and gastrin receptor antagonist (L-365,260) in dog and rat plasma by high-performance liquid chromatography 576(1992)363

L-668,750

Determination of enantiomeric concentrations of a 2,5-diaryltetrahydrofuran (L-668,750), a platelet-activating factor receptor antagonist, in rat plasma using a chiral α_1 -acid glycoprotein high-performance liquid chromatographic column 578(1992)327

L-694,492

High-performance liquid chromatographic determination of angiotensin II receptor antagonists in human plasma and urine. I. DuP 532 (L-694,492) 581(1992)165

Labetalol

Sensitive microbore high-performance liquid chromatographic assay for labetalol in the biological fluids of pregnant sheep 565(1991)383

Reversed-phase high-performance liquid chromatographic separation of the stereoisomers of labetalol via derivatization with chiral and non-chiral isothiocyanate reagents 579(1992)165

Lacidipine

Validation of a high-performance liquid chromatographic-radioimmunoassay method for the determination of lacidipine in plasma 573(1992)105

α -Lactalbumin

Anion-exchange chromatographic properties of α -lactalbumin eluted from quaternized polyvinylimidazole. Study of the role of the polymer coating 553(1991)165

β -Lactam antibiotics

see also Antibiotics

Reversed-phase high-performance liquid chromatography of amphoteric β -lactam antibiotics: effects of columns, ion-pairing reagents and mobile phase pH on their retention times 564(1991)195

Hydrophobicity of β -lactam antibiotics. Explanation and prediction of their behaviour in various partitioning solvent systems and reversed-phase chromatography 585(1991)3

Aqueous high-performance size-exclusion chromatographic assay for high-molecular-weight impurities in ceftiofur sodium 595(1992)185

D-Lactate

Fluorimetric and high-performance liquid chromatographic determination of D-lactate in biological samples 566(1991)1

Lactate dehydrogenase

Dye-ligand affinity partitioning of lactate dehydrogenase isoenzymes 557(1991)325

Electrophoresis of serum isoenzymes and proteins following acute myocardial infarction (Review) 569(1991)323

Electrophoretic analysis of enzyme-linked immunoglobulins and their clinical significance (Review) 569(1991)347

Study of high-affinity interactions by quantitative affinity chromatography. Analytical expressions in terms of total ligand concentration 597(1992)57

Lactic acid

Determination of volatile and non-volatile organic acids in technical sugar solutions by ion-exclusion chromatography 555(1991)65

Determination of plasma lactic acid concentration and specific activity using high-performance liquid chromatography 568(1991)301

Analysis of zapatera olives by gas and high-performance liquid chromatography 594(1992)259

Lactobionic acid

Determination of the association constant of monovalent mode protein-sugar interaction by capillary zone electrophoresis 597(1992)377

Lactoferrins

High-performance liquid chromatographic separation of human apolactoferrin and monoferric and diferric lactoferrins 579(1992)346

β -Lactoglobulin A

Adsorption-desorption isotherm hysteresis of β -lactoglobulin A with a weakly hydrophobic surface 557(1991)369

β -Lactoglobulin B

Study of the adsorption of self-associating proteins on an anion exchanger. Application to the chromatography of β -lactoglobulin B 599(1992)255

Lactones

Enantiomeric distribution and $^{13}\text{C}/^{12}\text{C}$ isotope ratio determination of γ -lactones: appropriate methods for the differentiation between natural and non-natural flavours? 557(1991)187

Analyse par chromatographie en phase gazeuse de la réaction de macrolactonisation de Mukaiyama 557(1991)501

Optimization of supercritical fluid extraction of volatile constituents from a model plant matrix 600(1992)175

Laminaran

Rapid method for preparing a β -glucan-specific sensitive fraction from *Limulus (Tachypleus tridentatus)* amebocyte lysate 567(1991)267

Lamotrigine

Use of thermospray liquid chromatography-mass spectrometry to aid in the identification of urinary metabolites of a novel antiepileptic drug, Lamotrigine 554(1991)181

Analysis of lamotrigine and lamotrigine 2-N-glucuronide in guinea pig blood and urine by reserved-phase ion-pairing liquid chromatography 571(1991)217

Performance analysis of a reversed-phase liquid chromatographic assay of lamotrigine in plasma using solvent-demixing extraction 572(1991)269

Lamotrigine 2-N-glucuronide

Analysis of lamotrigine and lamotrigine 2-N-glucuronide in guinea pig blood and urine by reserved-phase ion-pairing liquid chromatography 571(1991)217

Lannate

Removal of organic pollutants from aqueous solution. V. Comparative study of the extraction, recovery and chromatographic separation of some organic insecticides using unloaded polyurethane foam columns 552(1991)371

Lansoprazole

High-performance liquid chromatographic determination of lansoprazole and its metabolites in human serum and urine 571(1991)283

Determination of lansoprazole and its metabolites in plasma by high-performance liquid chromatography using a loop column 577(1992)117

Lansoprazole sulphide

Determination of lansoprazole and its metabolites in plasma by high-performance liquid chromatography using a loop column 577(1992)117

Lansoprazole sulphone

Determination of lansoprazole and its metabolites in plasma by high-performance liquid chromatography using a loop column 577(1992)117

Lantadenes

Thin-layer chromatographic separations of lantadenes, the pentacyclic triterpenoids from lantana (*Lantana camara*) plant 587(1991)351

Lanthanides

Reversed-phase separation of transition metals, lanthanides and actinides by elution with mandelic acid 558(1991)197

Factors affecting the separation of inorganic metal cations by capillary electrophoresis 593(1992)289

Study of isotachophoretic separation behaviour of metal cations by means of particle-induced X-ray emission. III. Analysis of a crude rare earth chloride from monazite 594(1992)371

Lanthanoids

Study of isotachophoretic separation behaviour of metal cations by means of particle-induced X-ray emission. II. Separation efficiency of lanthanoid model mixture 589(1992)339

Lasalocid

Improved high-performance liquid chromatographic procedure for the determination of lasalocid in chicken tissues and egg using polymeric and porous graphitic carbon columns 579(1992)177

Latex

Interfacial effects in size-exclusion chromatography of latex 587(1991)276

Laudanosine

High-performance liquid chromatographic assay for laudanosine in biological fluids and tissue for neurotoxic studies in rats 565(1991)465

Lauric acid

Determination of the cytochrome P-450 IV marker, ω -hydroxylauric acid, by high-performance liquid chromatography and fluorimetric detection 564(1991)266

LCAT

Human plasma lecithin:cholesterol acyltransferase. Preparation and use of immobilized *p*-aminophenylarsenoxide as a catalytic site-directed covalent ligand in enzyme purification 568(1991)69

Lead

Separation of metal complexes of ethylenediaminetetraacetic acid in environmental water samples by ion chromatography with UV and potentiometric detection 558(1991)181

Determination of Pb^{2+} in water by isotope dilution gas chromatography-mass spectrometry of tetraethyllead formed by reaction with sodium tetraethylborate 594(1992)275

Determination of divalent trace metals in natural waters by preconcentration on N,N,N',N'-tetra(2-aminoethyl)ethylenediamine-silica followed by on-line ion chromatography 600(1992)203

Lecithin

Use of hematoporphyrin as a fluorescent stain for detection of lipids in high-performance thin-layer chromatography 588(1991)295

Quantitative planar chromatography of phospholipids with different fatty acid compositions 588(1991)303

Separation of soy lecithin using gel permeation chromatography 589(1992)165

Lecithin:cholesterol acyltransferase

Human plasma lecithin:cholesterol acyltransferase. Preparation and use of immobilized *p*-aminophenylarsenoxide as a catalytic site-directed covalent ligand in enzyme purification 568(1991)69

Lectins

Bead cellulose derivatives as supports for immobilization and chromatographic purification of proteins 552(1991)389

Method for detecting the lectin activity of *Momordica charantia* transferred from micro two-dimensional electrophoretic gel on to nitrocellulose 570(1991)399

Arginine-tail method, an affinity tag procedure utilizing anhydrotrypsin agarose 597(1992)181

Improved affinity chromatographic purification of D-mannose-N-acetyl-D-glucosamine-specific lectin from the bark of *Sophora japonica* eliminating the loss by sugar specific self-aggregation 597(1992)197

Isolation and characterization of isolectins from *Erythrina variegata* seeds 597(1992)207

Purification and characterization of carbohydrate-binding peptides from *Lotus tetragonolobus* and *Ulex europaeus* seed lectins using affinity chromatography 597(1992)213

Determination of the carbohydrate-binding site of *Bauhinia purpurea* lectin by affinity chromatography 597(1992)221

Determination of the association constant of monovalent mode protein-sugar interaction by capillary zone electrophoresis 597(1992)377

Leucine

Measurement of leucine and α -ketoisocaproic acid fluxes in the fetal/placental unit 562(1991)169

Rapid measurement of leucine-specific activity in biological fluids by ion-exchange chromatography and post-column ninhydrin detection 571(1991)29

High-performance liquid chromatographic technique for non-derivatized leucine purification: evidence for carbon isotope fractionation 573(1992)11

Leucine-enkephalin

Limits and effects of precolumn addition of thioglycerol in liquid chromatographic-fast atom bombardment mass spectrometric systems 591(1992)129

Leucinocaine

Determination of limiting ionic mobilities and dissociation constants of some local anaesthetics 596(1992)265

Leucinostatins

Separation of leucinostatins by capillary zone electrophoresis 593(1992)259

Leucovorin

Optimization of the separation of (6*R*)- and (6*S*)-leucovorin and evaluation of the robustness of the optimum 592(1992)37

N-*l*-Leucyldoxorubicin

Fully automated determination of a new anthracycline N-*l*-leucyldoxorubicin and six metabolites in plasma by high-performance liquid chromatography with on-line sample handling 574(1992)273

N-*l*-Leucyldoxorubicinol

Fully automated determination of a new anthracycline N-*l*-leucyldoxorubicin and six metabolites in plasma by high-performance liquid chromatography with on-line sample handling 574(1992)273

Leukotriene antagonists

Direct high-performance liquid chromatographic separation of enantiomeric peptidoleukotriene antagonists 596(1992)123

Leukotriene D₄ antagonist

High-performance liquid chromatographic method for the determination of a novel leukotriene D₄ antagonist (MK-0571) in biological specimens 563(1991)458

Stereoselective determination of *R*(-)- and *S*(+)-MK-571, a leukotriene D₄ antagonist, in human plasma by chiral high-performance liquid chromatography 570(1991)157

Leukotrienes

Cyclooxygenase and lipoxygenase arachidonic acid metabolism by monocytes from human immune deficiency virus-infected drug users 557(1991)507

Enzymic preparation of dioxygen-18 labelled leukotriene E₄ and its use in quantitative gas chromatography-mass spectrometry 574(1992)181

Rapid method for automated on-line extraction and fractionation of plasma leukotrienes and 12-hydroxy-5,8,10,14-eicosatetraenoic acids by reversed-phase high-performance liquid chromatography 579(1992)37

Leuprorelin

High-performance liquid chromatography followed by radioimmunoassay for the determination of a luteinizing hormone-releasing hormone analogue, leuprorelin, and its metabolite 566(1991)57

Levodopa

see 3,4-Dihydroxyphenylalanine

Lewisite

Gas chromatographic determination of the lewisite hydrolysate, 2-chlorovinylarsonous acid, after derivatization with 1,2-ethanedithiol 558(1991)235

Lichenan

Rapid method for preparing a β -glucan-specific sensitive fraction from *Limulus* (*Tachypleus tridentatus*) amoebocyte lysate 567(1991)267

Lidocaine

Determination of local anaesthetics in body fluids by gas chromatography with surface ionization detection 564(1991)278

Determination of limiting ionic mobilities and dissociation constants of some local anaesthetics 596(1992)265

Lignocaine

High-performance liquid chromatographic method for the simultaneous determination of monoethylglycinexylidide and lignocaine 574(1992)361

Likviritin

High-performance liquid chromatographic method for the determination of mangiferin, likviritin and dihydroquercetin in rat plasma and urine 577(1992)371

Limonene

Design of a gas chromatograph with parallel radioactivity and mass spectrometric detection. Application to the identification of the major metabolite of *d*-limonene associated with α 2u-globulin 563(1991)271

Protection of terpenes against oxidative and acid decomposition on adsorbent cartridges 589(1992)385

Optimization of supercritical fluid extraction of volatile constituents from a model plant matrix 600(1992)175

Linalol

Cholesteryl acetate as a stationary phase for the gas chromatography of some volatile oil constituents 600(1992)337

Linoleic acid

Use of methyl oxime derivatives to enhance structural information in thermospray high-performance liquid chromatography-mass spectrometry. Analysis of linoleic acid lipoxygenase metabolites in maize embryos 554(1991)155

Linoleic acid lipoxygenase

Use of methyl oxime derivatives to enhance structural information in thermospray high-performance liquid chromatography-mass spectrometry. Analysis of linoleic acid lipoxygenase metabolites in maize embryos 554(1991)155

Lipases

Adsorption behaviour of lipase from *Staphylococcus carnosus* on a hydrophobic adsorbent 584(1992)109

p-Nitrophenyllaurate: a substrate for the high-performance liquid chromatographic determination of lipase activity 566(1991)453

Lipids

Enantiomer separation by chiral-phase liquid chromatography of urethane derivatives of natural diacylglycerols previously fractionated by reversed-phase liquid chromatography 557(1991)227

Gas chromatographic profiles of plasma total lipids as indicators of dietary history. Correlation with fat intake based on 24-h dietary recall 564(1991)11

Identification by gas chromatography and mass spectrometry of lipids from the rat Harderian gland 565(1991)27

Automated separation and quantitation of lipid fractions by high-performance liquid chromatography and mass detection 567(1991)21

Separation of acidic and neutral lipids by aminopropyl-bonded silica gel column chromatography 577(1992)142

Gas chromatographic profiles of plasma total lipids as indicators of dietary history. Correlation with carbohydrate and alcohol intake based on 24-h dietary recall 579(1992)13

Determination of cholesterol in sub-nanomolar quantities in biological fluids by high-performance liquid chromatography 579(1992)329

Quantitative separation of *Trichoderma* lipid classes on a bonded phase column 584(1992)129

Regeneration of column activity after the gas chromatographic separation of membrane lipids on thermostable SE-52 phase 585(1991)342

High-performance liquid chromatography of phosphatidic acids and related polar lipids 587(1991)193

Rapid high-performance liquid chromatographic determination of fatty acid profiles of lipids by conversion to their hydroxamic acids 587(1991)292

Use of hematoporphyrin as a fluorescent stain for detection of lipids in high-performance thin-layer chromatography 588(1991)295

Comparative study of clean-up and fractionation methods for the determination of organochlorine pesticides in lipids by gas chromatography 591(1992)303

Lipids, ether-linked polar

Fast atom bombardment mass spectrometry as a rapid means of screening mixtures of ether-linked polar lipids from extremely halophilic archaeobacteria for the presence of novel chemical structures 562(1991)369

Lipopeptides

Interaction of amphipathic model lipopeptides with phospholipid bilayers 597(1992)293

Lipopolysaccharides

High-performance liquid chromatographic method to resolve and determine lipopolysaccharide sub-groups of *Escherichia coli* endotoxin in isolated perfused rat liver perfusate 574(1992)205

Specific assay for endotoxin using immobilized histidine, *Limulus* amoebocyte lysate and a chromogenic substrate 597(1992)415

Lipoproteins

Separation of human serum lipoproteins into three major classes by hydroxyapatite chromatography 566(1991)67

Comparison of gel permeation chromatography, density gradient ultracentrifugation and precipitation methods for quantitation of very-low-, low- and high-density lipoprotein cholesterol 570(1991)382

Preparation of heparin-glyceryl controlled-pore glass affinity media for the separation of α - and β -lipoproteins 573(1992)132

Characterization of plasma apolipoproteins by capillary electrophoresis 583(1992)111

Counter-current chromatography of lipoproteins with a polymer phase system using the cross-axis synchronous coil planet centrifuge 596(1992)118

Liposomes

Determination of phosphatidylcholine, phosphatidylglycerol and their lyso forms from liposome dispersions by high-performance liquid chromatography using high-sensitivity refractive index detection 585(1991)239

Lipoxins

Determination of picomole amounts of lipoxins C₄, D₄ and E₄ by high-performance liquid chromatography with electrochemical detection 585(1991)53

Lometrexol

Bioanalysis of the investigational anti-tumour drug 5,10-dideaza-5,6,7,8-tetrahydrofolic acid by high-performance liquid chromatography with ultraviolet detection 576(1992)158

Lonapalene

High-performance liquid chromatographic determination of the 1,4-diketo and 1,4-diketo-2,3-dihydroxy metabolites of lonapalene in rat urine 565(1991)339

Long-chain bases

Fragmentation pathways of O-trimethylsilyl ethers of dihydroxy long-chain bases analysed by linked-scan mass spectrometry 562(1991)119

Longifolene

Use of a Sep-Pak cartridge for preparative collection in gas chromatography 555(1991)302

Lopirazepam

Direct liquid chromatographic separation of enantiomers on immobilized protein stationary phases. IX. Influence of the cross-linking reagent on the retentive and enantioselective properties of chiral sorbents based on bovine serum albumin 591(1992)65

Loracarbef

High-performance liquid chromatographic determination of loracarbef, a potential metabolite, cefaclor and cephalixin in human plasma, serum and urine 567(1991)129

Lormetazepam

(S,S)-Diphenylethylethanediamine derivatives as chiral selectors. II. Gasparrini-type bound chiral stationary phase with high enantioselectivity for naphthylamides 553(1991)373

Losartan

Simultaneous determination of a novel angiotensin II receptor blocking agent, losartan, and its metabolite in human plasma and urine by high-performance liquid chromatography 573(1992)295

Low-density lipoproteins

Separation of human serum lipoproteins into three major classes by hydroxyapatite chromatography 566(1991)67

Low-density polyethylene

Determination of Cyasorb UV 1084 and its degradation products in low-density polyethylene film by high-performance liquid chromatography 587(1991)205

Loxapine

Simultaneous quantitation of loxapine, amoxapine and their 7- and 8-hydroxy metabolites in plasma by high-performance liquid chromatography 564(1991)213

LSD

Chromatographic and mass spectrometric methods for determination of lysergic acid diethylamide (LSD) and metabolites in body fluids (Review) 580(1992)97

Luminarins

Primary and secondary amine derivatization with luminarins 1 and 2: separation by liquid chromatography with peroxyoxalate chemiluminescence detection 594(1992)386

Luminol

High-performance liquid chromatographic investigation of product formation in the horseradish peroxidase-enhanced chemiluminescence of luminol with different enhancers 566(1991)461

Lung fluid proteins

High-performance capillary electrophoresis of proteins from the fluid lining of the lungs of rats exposed to perfluoroisobutylene 559(1991)411

Lutein

Isolation and structural elucidation of the geometrical isomers of lutein and zeaxanthin in extracts from human plasma 582(1992)153

Determination of carotenoid pigments in several tree leaves by reversed-phase high-performance liquid chromatography 585(1991)168

Optimization of an isocratic high-performance liquid chromatographic separation of carotenoids 589(1992)171

Evaluation of reversed-phase liquid chromatographic columns for recovery and selectivity of selected carotenoids 595(1992)89

Luteinizing hormone releasing hormone

Densitometry for analysis of protein and peptide hydrolysates: application to luteinizing hormone releasing hormone 568(1991)451

Luxabendazole

Determination of luxabendazole in biological fluids by high-performance liquid chromatography 578(1992)321

Lycopene

Optimization of an isocratic high-performance liquid chromatographic separation of carotenoids 589(1992)171

Evaluation of reversed-phase liquid chromatographic columns for recovery and selectivity of selected carotenoids 595(1992)89

Lym-1 F(ab')₂ fragment

Chromatographic methods for large-scale preparation of immunoglobulin G_{2a} monoclonal antibodies Lym-1 and TNT-1 F(ab')₂ fragments 563(1991)243

Lysergic acid diethylamide

Chromatographic and mass spectrometric methods for determination of lysergic acid diethylamide (LSD) and metabolites in body fluids (Review) 580(1992)97

Lysine

Liquid chromatographic-mass spectrometric analysis for screening of patients with cystinuria, and identification of cystine stone 568(1991)445

Lysophospholipids

One-dimensional thin-layer chromatographic separation of phospholipids and lysophospholipids from tissue lipid extracts 581(1992)139

Lysozyme

Purification of ovalbumin and lysozyme from a commercial product by recycling isotachopheresis 585(1991)145

Immobilization of Fv antibody fragments on porous silica and their utility in affinity chromatography 587(1991)161

Use of antibody fragments in immunoaffinity chromatography. Comparison of FV fragments, VH fragments and paralog peptides 597(1992)239

2-Lyso-platelet activating factor

Quantification of distinct molecular species of the 2-lyso metabolite of platelet-activating factor by gas chromatography-negative-ion chemical ionization mass spectrometry 575(1992)183

Lysylpyridinoline

High-performance liquid chromatographic determination of 3-hydroxypyridinium derivatives as new markers of bone resorption 574(1992)255

M

Mabuterol

Determination of β_2 -receptor agonists in bovine urine and liver by gas chromatography-tandem mass spectrometry 564(1991)515

Macrolactones

Analyse par chromatographie en phase gazeuse de la réaction de macrolactonisation de Mukaiyama 557(1991)501

Macrolides

Sensitive determination of josamycin and rokitamycin in plasma by high-performance liquid chromatography with fluorescence detection 575(1992)171

Madecassoside

Direct coupling of high-speed counter-current chromatography to thin-layer chromatography. Application to the separation of asiaticoside and madecassoside from *Centella asiatica* 558(1991)446

Magnolol

Identification and determination of honokiol and magnolol from *Magnolia officinalis* by high-performance liquid chromatography with photodiode-array UV detection 598(1992)143

Maleic anhydride

Immobilization of protein ligands with methyl vinyl ether-maleic anhydride copolymer 597(1992)123

Maleic hydrazide

Determination of maleic hydrazide in tobacco by micellar liquid chromatography 595(1992)346

Malic acid

Determination of volatile and non-volatile organic acids in technical sugar solutions by ion-exclusion chromatography 555(1991)65

Ligand-exchange ion chromatographic determination of malic acid enantiomers in apple juice with photometric detection 585(1991)315

Determination of carboxylic acids, sugars, glycerol and ethanol in wine and grape must by ion-exchange high-performance liquid chromatography with refractive index detection 590(1992)215

Malonaldehyde

Determination of a guanosine-malonaldehyde adduct in urine by high-performance liquid chromatography with a thiobarbituric acid reaction detector 570(1991)301

Malondialdehyde

Excretion of formaldehyde, malondialdehyde, acetaldehyde and acetone in the urine of rats in response to 2,3,7,8-tetrachlorodibenzo-*p*-dioxin, paraquat, endrin and carbon tetrachloride 576(1992)221

High-performance liquid chromatographic separation of malondialdehyde-thiobarbituric acid adduct in biological materials (plasma and human cells) using a commercially available reagent 577(1992)9

Malonic acid

Development and use of a stability-indicating high-performance liquid chromatographic assay for Meldrum's acid 575(1992)177

Maltitol

Separation of functionalized dextrans by reversed-phase high-performance liquid chromatography 587(1991)288

Maltosaccharides

Estimation of the distributions of chain length of amylopectins by high-performance liquid chromatography with pulsed amperometric detection 585(1991)233

Maltose

Separation of functionalized dextrans by reversed-phase high-performance liquid chromatography 587(1991)288

Mandelic acid

Simultaneous high-performance liquid chromatographic determination of urinary metabolites of benzene, nitrobenzene, toluene, xylene and styrene 573(1992)318

Experimental study of the production rate of pure enantiomers from racemic mixtures 590(1992)119

Contribution of ionically immobilized bovine serum albumin to the retention of enantiomers 600(1992)37

Mangiferin

High-performance liquid chromatographic method for the determination of mangiferin, likviritin and dihydroquercetin in rat plasma and urine 577(1992)371

Manidipine

Determination of manidipine enantiomers in human serum using chiral chromatography and column-switching liquid chromatography 575(1992)123

Mannan-binding protein

Affinity chromatographic purification of immunoglobulin M antibodies utilizing immobilized mannan binding protein 597(1992)247

Mannitol

Chromatographic separation of polyols by ligand exchange. Effects of the ion-exchange resin cross-linking and size 594(1992)125

Marine organic pollutants

Chromatographic separation of marine organic pollutants (Review) 580(1992)191

Marine toxins

Comparison of liquid-junction and coaxial interfaces for capillary electrophoresis-mass spectrometry with application to compounds of concern to the aquaculture industry 591(1992)325

Masking agents

Approach to the analysis of diuretics and masking agents by high-performance liquid chromatography-mass spectrometry in doping control 562(1991)723

MDL 72974A

Determination of an irreversible inhibitor of monoamine oxidase B (MDL 72974A) in human plasma and urine by gas chromatography-positive-ion chemical ionization mass spectrometry 571(1991)241

Mebendazole

Application of a high-performance liquid chromatography coulometric method for the estimation of mebendazole and its metabolites in human sera 563(1991)115

Meconium

Analysis of meconium for cocaine in neonates 575(1992)158

Med 5

High-performance liquid chromatographic evaluation of Med 15 and its metabolites Med 5 and tolmetin in rat plasma 553(1991)81

Med 15

High-performance liquid chromatographic evaluation of Med 15 and its metabolites Med 5 and tolmetin in rat plasma 553(1991)81

Medroxyprogesterone acetate

Mass spectrometric and high-performance liquid chromatographic studies of medroxyprogesterone acetate metabolites in human plasma 562(1991)351

Mefenamic acid

Simultaneous analysis of flunixin, naproxen, ethacrynic acid, indomethacin, phenylbutazone, mefenamic acid and thiosalicylic acid in plasma and urine by high-performance liquid chromatography and gas chromatography-mass spectrometry 568(1991)351

Column-switching techniques for high-performance liquid chromatography of ibuprofen and mefenamic acid in human serum with short-wavelength ultraviolet detection 570(1991)329

Mefloquine

Determination of mefloquine in blood, filter paper-absorbed blood and urine by 9-fluorenylmethyl chloroformate derivatization followed by liquid chromatography with fluorescence detection 564(1991)181

Reversed-phase liquid chromatographic method for the simultaneous determination of the antimalarial drugs sulfadoxine, pyrimethamine, mefloquine and its major carboxylic metabolite in plasma 571(1991)169

Melanogens

Indolic urinary melanogens: separation and identification by gas chromatography with selected-ion monitoring mass spectrometry of 5-hydroxy-6-methoxyindole-2-carboxylic and 5-methoxy-6-hydroxyindole-2-carboxylic acids 570(1991)166

Melanotron-II

High-performance liquid chromatographic assay for the α -melanotropin[4,10] fragment analogue (Melanotan-II) in rat plasma 584(1992)175

 α -Melanotropin[4,10] fragment

High-performance liquid chromatographic assay for the α -melanotropin[4,10] fragment analogue (Melanotan-II) in rat plasma 584(1992)175

Melatonin

Determination of melatonin and monoamines in rat pineal using reversed-phase ion-interaction chromatography with fluorescence detection 564(1991)93

Determination of 5-methoxyindoles in pineal gland and plasma samples by high-performance liquid chromatography with electrochemical detection 564(1991)103

Meldrum's acid

Development and use of a stability-indicating high-performance liquid chromatographic assay for Meldrum's acid 575(1992)177

Melphalan

Quantitation of melphalan in plasma of patients by reversed-phase high-performance liquid chromatography with electrochemical detection 563(1991)443

Membrane proteins

see also Proteins

High-performance membrane chromatography of serum and plasma membrane proteins 590(1992)59

Comparison of detergents for extraction and ion-exchange high-performance liquid chromatography of Sendai virus membrane proteins 599(1992)157

Memory-enhancing agent

Liquid chromatography of the potential memory-enhancing agent CL 275,838 and its main metabolites, using a post-column photochemical reactor and fluorimetric detection 581(1992)109

Mephénytoin

Determination of mephénytoin stereoselective oxidative metabolism in urine by chiral liquid chromatography employing β -cyclodextrin as a mobile phase additive 573(1992)173

Synthesis of (1*R*-*trans*)-*N,N'*-1,2-cyclohexylenebisbenzamide-oligodimethylsiloxane copolymers for use as chiral stationary phases for capillary supercritical fluid chromatography 594(1992)283

Mepivacaine

Determination of local anaesthetics in body fluids by gas chromatography with surface ionization detection 564(1991)278

Meprobamate

Gas chromatographic determination of meprobamate in serum or plasma after solid-phase extraction 577(1992)171

Mercaptalbumin

Rapid analysis of human serum albumin by high-performance liquid chromatography 578(1992)292

***N*-[2(*S*)-(Mercaptomethyl)-3-(2-methylphenyl)-1-oxopropyl]-*L*-methionine**

High-performance liquid chromatographic determination of *N*-[2(*S*)-(mercaptomethyl)-3-(2-methylphenyl)-1-oxopropyl]-*L*-methionine, the active plasma metabolite of a prodrug atriopeptidase inhibitor (SCH 42495), using a thiol selective (Au/Hg) amperometric detector 579(1992)307

2-Mercaptopropionic acid

Determination of 2-mercaptopropionylglycine and its metabolite, 2-mercaptopropionic acid, in plasma by ion-pair reversed-phase high-performance liquid chromatography with post-column derivatization 564(1991)258

2-Mercaptopropionylglycine

Determination of 2-mercaptopropionylglycine and its metabolite, 2-mercaptopropionic acid, in plasma by ion-pair reversed-phase high-performance liquid chromatography with post-column derivatization 564(1991)258

6-Mercaptopurine

High-performance liquid chromatographic assay of the methyl and nucleotide metabolites of 6-mercaptopurine: quantitation of red blood cell 6-thioguanine nucleotide, 6-thioinosinic acid and 6-methylmercaptopurine metabolites in a single sample 583(1992)83

Mescaline

Determination of psychotropic phenylalkylamine derivatives in biological matrices by high-performance liquid chromatography with photodiode-array detection 593(1992)87

Mesityl oxide

Optimization of resolution in capillary zone electrophoresis: combined effect of applied voltage and buffer concentration 588(1991)315

Metabolites

Simple cation-exchange high-performance liquid chromatography optimized to the measurement of metabolites in the effluents from perfused rat livers using refractive index and ultraviolet detectors 563(1991)231

Metacycline

Determination of metacycline and related substances by column liquid chromatography on poly(styrene-divinylbenzene) 586(1991)61

Isolation of doxycycline, 6-epidoxycycline and 2-acetyl-2-decarboxamidometacycline from commercial metacycline by preparative column liquid chromatography on silica gel 586(1991)67

Metal chelates

Correlation analysis in liquid chromatography of metal chelates. II. One-dimensional retention-mobile phase composition (physicochemical property) models in reversed-phase liquid chromatography 555(1991)163

Metal complexes

Separation of metal complexes of ethylenediaminetetraacetic acid in environmental water samples by ion chromatography with UV and potentiometric detection 558(1991)181

Reversed-phase high-performance liquid chromatography of several metal-8-quinolinethiol complexes 596(1992)73

Metal ions

Measurement of stability constants for complexes of divalent cations with tartaric acid by ion chromatography 589(1992)185

Study of isotachophoretic separation behaviour of metal cations by means of particle-induced X-ray emission. II. Separation efficiency of lanthanoid model mixture 589(1992)339

Determination of metal ions by on-line complexation and ion-pair chromatography 590(1992)263

Factors affecting the separation of inorganic metal cations by capillary electrophoresis 593(1992)289

Study of isotachophoretic separation behaviour of metal cations by means of particle-induced X-ray emission. III. Analysis of a crude rare earth chloride from monazite 594(1992)371

Quantitative separation of Cr^{3+} from Mo^{6+} , W^{6+} , Hg_2^{2+} , Cu^{2+} and Pb^{2+} . Chromatographic behaviour of 51 cations on papers impregnated with Sn(IV)-based inorganic ion exchangers in complex-forming acid systems 594(1992)379

Metallothionein

Determination of dietary cadmium-induced metallothioneins in rabbit kidneys and cadmium in metallothioneins by anion-exchange high-performance liquid chromatography coupled with graphite furnace atomic absorption spectrometry 572(1991)73

Metals

N-(Hydroxymethyl)thioamide resin as stationary phase in ion-exchange chromatography for metal ion separation 555(1991)291

Suppression of the effect of metal impurities in alkylsilylated silica packing materials 594(1992)87

Metanephrine

Determination of free and conjugated normetanephrine and metanephrine in human plasma by high-performance liquid chromatography with electrochemical detection 563(1991)23

High-performance liquid chromatographic determination, with ultraviolet detection, of S-adenosyl-L-methionine and S-adenosyl-L-homocysteine in rat tissues and simultaneously of normetanephrine and metanephrine for phenylethanolamine-N-methyltransferase or catechol-O-methyltransferase activities 567(1991)254

Determination of urinary normetanephrine, metanephrine and 3-methoxytyramine by high-performance liquid chromatography with electrochemical detection: comparison between automated column-switching and manual dual-column sample purification methods 578(1992)165

Metformin

Determination of metformin in plasma by high-performance liquid chromatography 583(1992)270

Methacrylonitrile

Determination of methacrylonitrile in serum using gas chromatography and flame ionization detection 579(1992)153

Methadone

Chiral analysis of methadone in plasma by high-performance liquid chromatography 570(1991)198

Stereoselective determination of the enantiomers of methadone in plasma using high-performance liquid chromatography 583(1992)195

Methamphetamine

Determination of methamphetamine enantiomer ratios in urine by gas chromatography-mass spectrometry (Review) 580(1992)77

Methandienone

Studies on anabolic steroids. V. Sequential reduction of methandienone and structurally related steroid A-ring substituents in humans: gas chromatographic-mass spectrometric study of the corresponding urinary metabolites 562(1991)323

Detection of methandienone (methandrostenolone) and metabolites in horse urine by gas chromatography–mass spectrometry 577(1992)195

Methandrostenolone

Detection of methandienone (methandrostenolone) and metabolites in horse urine by gas chromatography–mass spectrometry 577(1992)195

1-(4-Methanesulphonamidophenoxy)-2-[N-(4-methanesulphonamidophenethyl)-N-methylamino]ethane

Liquid–liquid extraction and high-performance liquid chromatography for the determination of a novel antidysrhythmic agent (UK-68,798) in human urine 568(1991)475

Methanethiol

Preconcentration technique for introducing gaseous or volatile compounds into a capillary gas chromatographic column 587(1991)338

Methanol

Estimation of cellulose polarity by gas chromatography 552(1991)365

Bile salt surfactants in micellar electrokinetic capillary chromatography. Application to hydrophobic molecule separations 557(1991)113

Development of post-column enzymic reactors with immobilized alcohol oxidase for use in the high-performance liquid chromatographic assay of alcohols with electrochemical detection 563(1991)11

Direct injection high-performance liquid chromatographic method with electrochemical detection for the determination of ethanol and methanol in plasma using an alcohol oxidase reactor 566(1991)333

Determination of methanol in whole blood by capillary gas chromatography with direct on-column injection 570(1991)406

Analysis of zapatera olives by gas and high-performance liquid chromatography 594(1992)259

Methionine

Analysis of tyrosine- and methionine-containing neuropeptides by fast atom bombardment mass spectrometry 562(1991)459

Methionine-enkephalin

Effect of the addition of viscous matrices to the mobile phase on chromatographic performance in liquid chromatography–fast atom bombardment mass spectrometry 554(1991)61

Analysis of methionine enkephalin in human pituitary by multi-dimensional reversed-phase high-performance liquid chromatography, radioreceptor assay, radioimmunoassay, fast atom bombardment mass spectrometry, and mass spectrometry–mass spectrometry 562(1991)573

Characterization of an aminopeptidase in cerebrospinal fluid. Structure elucidation of enzyme hydrolysis products of synthetic methionine-enkephalin by reversed-phase high-performance liquid chromatography and mass spectrometry 574(1992)189

Limits and effects of precolumn addition of thioglycerol in liquid chromatographic–fast atom bombardment mass spectrometric systems 591(1992)129

Methocarbamol

High-performance liquid chromatographic analysis of methocarbamol enantiomers in biological fluids 582(1992)173

Methotrexate

High-performance liquid chromatography method for serum methotrexate levels in children with severe steroid-dependent asthma 565(1991)349

5-Methoxy-6-hydroxyindole-2-carboxylic acid

Indolic urinary melanogens: separation and identification by gas chromatography with selected-ion monitoring mass spectrometry of 5-hydroxy-6-methoxyindole-2-carboxylic and 5-methoxy-6-hydroxyindole-2-carboxylic acids 570(1991)166

3-Methoxy-4-hydroxyphenylalanine

Method for measuring endogenous 3-O-methyldopa in urine and plasma 568(1991)45

3-Methoxy-4-hydroxyphenylglycol

Analysis of rat brain microdialysate by gas chromatography–high-resolution selected-ion monitoring mass spectrometry 562(1991)111

Improved high-performance liquid chromatographic method for the routine determination of unconjugated 3-methoxy-4-hydroxyphenylethyleneglycol in human plasma using solid-phase extraction and electrochemical detection 565(1991)131

5-Methoxyindoleacetic acid

Determination of 5-methoxyindoles in pineal gland and plasma samples by high-performance liquid chromatography with electrochemical detection 564(1991)103

5-Methoxyindoles

Determination of 5-methoxyindoles in pineal gland and plasma samples by high-performance liquid chromatography with electrochemical detection 564(1991)103

Methoxymethylene dimethyl dodecatrienes

Liquid chromatographic isolation and structural elucidation of methoxymethylene dimethyl dodecatrienes catalytically synthesized from methanol and three moles of isoprene 594(1992)111

Methoxyphenylacetates

Selected applications of cyclodextrin selectors in capillary electrophoresis 559(1991)215

2-(4-Methoxyphenyl)-5,6,7,8,9,10-hexahydrocyclohepta[*b*]pyrazolo[3,4-*d*]pyridin-3(2*H*)-one

Determination of a potential anxiolytic drug (CGS 20625) in human plasma by high-performance liquid chromatography 568(1991)487

8-Methoxypsoralen

Determination of 8-methoxypsoralen in plasma by gas chromatography-mass spectrometry using selected-ion monitoring 574(1992)156

4-Methoxy-2,3,6-trimethylbenzenesulphonyl groups

Monitoring 4-methoxy-2,3,6-trimethylbenzenesulphonyl deprotection of arginine-containing synthetic peptides using capillary zone electrophoresis 589(1992)380

5-Methoxytryptamine

Determination of 5-methoxyindoles in pineal gland and plasma samples by high-performance liquid chromatography with electrochemical detection 564(1991)103

5-Methoxytryptophol

Determination of 5-methoxyindoles in pineal gland and plasma samples by high-performance liquid chromatography with electrochemical detection 564(1991)103

3-Methoxytyramine

Determination of urinary normetanephrine, metanephrine and 3-methoxytyramine by high-performance liquid chromatography with electrochemical detection: comparison between automated column-switching and manual dual-column sample purification methods 578(1992)165

3-Methoxytyrosine

Method for measuring endogenous 3-O-methyldopa in urine and plasma 568(1991)45

 α -Methylacetohydroxamic acids

Use of high-performance liquid chromatography-thermospray mass spectrometry and gas chromatography-electron-impact mass spectrometry in the identification of the metabolites of α -methylacetohydroxamic acids, potential anti-asthmatic agents 562(1991)249

Methylalkynes

Capillary gas chromatography of C₅-C₁₃ branched alkynes on squalane and liquid crystal stationary phases 557(1991)241

Methylamines

Determination of trimethylamine and related aliphatic amines in human urine by head-space gas chromatography 584(1992)141

N-Methyl-*p*-aminoacetophenone

Identification of aromatic moieties and mycosamine in antifungal heptaenes with high-performance liquid chromatography, high-performance liquid chromatography-mass spectrometry and gas chromatography-mass spectrometry 585(1991)247

N-Methylamphetamine

Electron-capture gas chromatographic procedure for simultaneous determination of amphetamine and N-methylamphetamine 573(1992)313

7-Methylbenz[*c*]acridines

Determination of 7-methylbenz[*c*]acridines by capillary gas chromatography with electron-capture detection 596(1992)233

Methylbenzothiophene sulphoxides

Decomposition of two methylbenzothiophene sulphoxides in a commercial gas chromatography injection port liner 591(1992)362

Methylboronic acid

Determination of methylboronic acid in teboroxime by capillary gas chromatography 585(1991)348

2-Methylbutanoic acid

Simultaneous stereoanalysis of 2-alkyl-branched acids, esters and alcohols using a selectivity-adjusted column system in multi-dimensional gas chromatography 587(1991)347

2-Methylbutan-1-ol

Simultaneous stereoanalysis of 2-alkyl-branched acids, esters and alcohols using a selectivity-adjusted column system in multi-dimensional gas chromatography 587(1991)347

(*S*)-(+)-2-Methylcarboxypropyl chloromethyl ether

Some novel homochiral derivatizing agents for the gas chromatographic analysis of enantiomeric secondary alcohols 598(1992)251

3-Methylcyclopent-2-ene

Mass spectrometric and gas and high-performance liquid chromatographic behaviour of an impurity in 2,5-hexanedione 590(1992)223

5-Methylcytosine

5-Methylcytosine content of DNA in blood, synovial mononuclear cells and synovial tissue from patients affected by autoimmune rheumatic diseases 566(1991)481

Methyl 3,5-diamino-6-chloropyrazinecarboxylate

Determination and identification of amiloride in human urine by high-performance liquid chromatography and gas chromatography-mass spectrometry 582(1992)93

3-O-Methyldopa

Determination of free DOPA and 3-O-methyl-DOPA in human plasma by high-performance liquid chromatography with electrochemical detection 553(1991)265

Method for measuring endogenous 3-O-methyldopa in urine and plasma 568(1991)45

6-Methylenandrosta-1,4-diene-3,17-dione

High-performance liquid chromatographic analysis of FCE 24304 (6-methylenandrosta-1,4-diene-3,17-dione) and FCE 24928 (4-aminoandrosta-1,4,6-triene-3,17-dione), two new aromatase inhibitors 593(1991)25

N,N-Methylenebisacrylamide

Continuous beds for standard and micro high-performance liquid chromatography 586(1991)21

4,4'-Methylenedianiline

High-performance liquid chromatographic determination of 4,4'-methylenedianiline in human urine 564(1991)205

Determination of 4,4'-methylenedianiline in hydrolysed human urine by micro liquid chromatography with ultraviolet detection 579(1992)350

Methylenedianiline

Identification of metabolites of 4,4'-diaminodiphenylmethane (methylene dianiline) using liquid chromatographic and mass spectrometric techniques 583(1992)63

Methylenedioxyamphetamine

Determination of psychotropic phenylalkylamine derivatives in biological matrices by high-performance liquid chromatography with photodiode-array detection 593(1992)87

Methylenedioxymethylamphetamine

Determination of psychotropic phenylalkylamine derivatives in biological matrices by high-performance liquid chromatography with photodiode-array detection 593(1992)87

2-Methyleneglutarate mutase

Purification of the coenzyme B₁₂-containing 2-methyleneglutarate mutase from *Clostridium barkeri* by high-performance liquid chromatography 587(1991)93

Methyl ethyl ketone

High-performance liquid chromatographic microassay for methyl ethyl ketone in urine as the 2,4-dinitrophenylhydrazone derivative 579(1992)324

N-(1-Methylethyl)-N'-phenyl-1,4-benzenediamine

Rapid high-performance liquid chromatographic determination of urinary N-(1-methylethyl)-N'-phenyl-1,4-benzenediamine in workers exposed to aromatic amines 592(1992)279

1-Methylfluorene

Laser-excited fluorescence detection of gas-phase chromatography eluates 557(1991)39

N-Methylformamide

High-performance liquid chromatographic determination of N-methylformamide and N-methyl-N-(hydroxymethyl)formamide in human urine 581(1992)287

4-O-Methylgallic acid

Determination of tannic acid and its phenolic metabolites in biological fluids by high-performance liquid chromatography 577(1992)77

Methylglycosides

Liquid chromatographic method for the determination of the carbohydrate moiety of glycoproteins. Application to α_1 -acid glycoprotein and tissue plasminogen activator 558(1991)105

3-Methylhippuric acid

Simultaneous high-performance liquid chromatographic determination of urinary metabolites of benzene, nitrobenzene, toluene, xylene and styrene 573(1992)318

N^r-Methylhistamine

Gas chromatographic-mass spectrometric assay to measure urinary N^r-methylhistamine excretion in man 567(1991)289

1-Methylhistamine

High-performance liquid chromatography of histamine and 1-methylhistamine with on-column fluorescence derivatization 595(1992)163

N^r-Methylhistidine

Simple and sensitive determination of plasma N^r-methylhistidine by high-performance liquid chromatography using pre-column derivative formation with *o*-phthalaldehyde-2-mercaptoethanol 566(1991)223

1-Methylhistidine

Determination of carnosine and other biogenic imidazoles in equine plasma by isocratic reversed-phase ion-pair high-performance liquid chromatography 579(1992)45

3-Methylhistidine

Determination of urinary 3-methylhistidine by high-performance liquid chromatography with *o*-phthalaldehyde precolumn derivatization 581(1992)272

High-performance liquid chromatography-electrochemical detection of 3-methylhistidine in human urine 584(1992)256

N-Methyl-N-(hydroxymethyl)formamide

High-performance liquid chromatographic determination of N-methylformamide and N-methyl-N-(hydroxymethyl)formamide in human urine 581(1992)287

4-Methylhydroxysulphadimidine

High-performance liquid chromatographic method for the routine determination of sulphadimidine, its hydroxy metabolites and N4-acetylsulphadimidine in body fluids and cell culture media 571(1991)157

3-Methylindole

Rapid and accurate high-performance liquid chromatographic method for the determination of 3-methylindole (skatole) in faeces of various species 566(1991)101

2-Methylitaconate Δ -isomerase

Purification of the coenzyme B₁₂-containing 2-methyleneglutarate mutase from *Clostridium barkeri* by high-performance liquid chromatography 587(1991)93

Methylmalonic acid

Analysis of serum methylmalonic acid for the determination of cobalt deficiency in cattle 566(1991)215

Methylmannoside

Study of high-affinity interactions by quantitative affinity chromatography. Analytical expressions in terms of total ligand concentration 597(1992)57

Methylmercaptapurine

Reversed-phase high-performance liquid chromatographic approach to determine total lymphocyte concentrations of 6-thioguanine, methylmercaptapurine and methylthioguanine in humans 571(1991)149

6-Methylmercaptapurine

High-performance liquid chromatographic assay of the methyl and nucleotide metabolites of 6-mercaptapurine: quantitation of red blood cell 6-thioguanine nucleotide, 6-thioinosinic acid and 6-methylmercaptapurine metabolites in a single sample 583(1992)83

Methylmercaptapurine riboside

Chromatographic analysis of methylmercaptapurine riboside in human plasma and urine 564(1991)303

Methylmercury

Determination of methylmercury in human blood using capillary gas chromatography and selected-ion monitoring 573(1992)35

Evaluation of gas chromatographic columns for the determination of methylmercury in aqueous head space extracts from biological samples 586(1991)329

Methyl methacrylate

Determination of topanol antioxidants in methacrylates using capillary gas-liquid chromatography 596(1992)290

Methyl 2-methylbutanoate

Simultaneous stereoanalysis of 2-alkyl-branched acids, esters and alcohols using a selectivity-adjusted column system in multi-dimensional gas chromatography 587(1991)347

1-Methyl-4-methylsulfinyltetrachlorobenzene

Artifact formation during gas chromatographic-mass spectrometric analysis of a methylsulfinyl-containing metabolite 575(1992)287

N-Methylmorpholine

Liquid membrane work-up of blood plasma samples applied to gas chromatographic determination of aliphatic amines 573(1992)191

Methylmyristate

Automated high-performance liquid chromatographic and size-exclusion chromatographic sample preparation by means of a robotic workstation 592(1992)249

2-Methylpentane

Study of microporous carbons by gas chromatographic determination of heats of physisorption 555(1991)183

Methylphenobarbital

Chiral separation of barbiturates and hydantoins by reversed-phase high-performance liquid chromatography using a 25 or 50 mm short ODS cartridge column via β -cyclodextrin inclusion complexes 579(1992)253

(\pm)-Methyl 3-phenyl-2(*E*)-propenyl 1,4-dihydro-2,6-dimethyl-4-(3-nitrophenyl)-3,5-pyridinedicarboxylate

Use of a carrier for quantitation of a new dihydropyridine calcium antagonist (OPC-13340) in human plasma by highly sensitive gas chromatography with negative-ion chemical ionization mass spectrometry 577(1992)275

1-Methyl-4-phenyl-1,2,3,6-tetrahydropyridine

Liquid chromatography-mass spectrometry with ionspray and electrospray interfaces in pharmaceutical and biomedical research 554(1991)39

Endogenous synthesis of N-methylsalsolinol, an analogue of 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine, in rat brain during *in vivo* microdialysis with salsolinol, as demonstrated by gas chromatography-mass spectrometry 578(1992)109

Methylprednisolone

Validation of the simultaneous determination of methylprednisolone and methylprednisolone acetate in human plasma by high-performance liquid chromatography 577(1992)87

Methylprednisolone acetate

Validation of the simultaneous determination of methylprednisolone and methylprednisolone acetate in human plasma by high-performance liquid chromatography 577(1992)87

Methylpyrazines

Simple and accurate determination of methylpyrazines in biofluids using high-performance liquid chromatography 567(1991)415

5-Methylpyrimidines

High-performance liquid chromatographic analysis of 5-ethylpyrimidines and 5-methylpyrimidines in plasma 563(1991)392

N-Methylsalsolinol

Endogenous synthesis of N-methylsalsolinol, an analogue of 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine, in rat brain during *in vivo* microdialysis with salsolinol, as demonstrated by gas chromatography-mass spectrometry 578(1992)109

1-Methylsulphinyl-2-[2-(methylthio)ethylsulphonyl]ethane

Analysis of 1,1'-sulphonylbis[2-(methylsulphinyl)ethane] and 1-methylsulphinyl-2-[2-(methylthio)ethylsulphonyl]ethane, metabolites of sulphur mustard, in urine using gas chromatography-mass spectrometry 558(1991)405

Methylsulphonium ions

Ion chromatographic analysis of the purity and synthesis of sulfonium and selenonium ions 588(1991)211

(5 α ,17 α)-1'-(Methylsulphonyl)-1'-H-pregn-20-yno[3,2-c]-pyrazol-17-ol

Determination of a novel steroidal androgen receptor antagonist (Win 49596) in human plasma using solid-phase extraction and high-performance liquid chromatography with ultraviolet detection 568(1991)494

Methyltestosterone

Comparison of two mass-selective detectors with special reference to the analysis of residues of anabolics 564(1991)385

Combined high-performance liquid chromatography and radioimmunoassay for the screening of 19-nortestosterone and methyltestosterone residues in meat samples 564(1991)445

Use of an immobilized enzyme reactor for the analysis of residues of 17 α -methyltestosterone in trout by high-performance liquid chromatography 564(1991)461

Simultaneous determination of yohimbine hydrochloride, strychnine nitrate and methyltestosterone by ion-pair high-performance liquid chromatography 588(1991)344

p-Methylthiobenzamide

Determination of p-methylthiobenzamide and p-methylthiobenzamide-S-oxide from rat plasma using solid-phase extraction and high-performance liquid chromatography 567(1991)213

p-Methylthiobenzamide-S-oxide

Determination of p-methylthiobenzamide and p-methylthiobenzamide-S-oxide from rat plasma using solid-phase extraction and high-performance liquid chromatography 567(1991)213

Methylthioguanine

Reversed-phase high-performance liquid chromatographic approach to determine total lymphocyte concentrations of 6-thioguanine, methylmercaptapurine and methylthioguanine in humans 571(1991)149

4-Methylumbelliferone

High-performance liquid chromatographic method for the direct determination of 4-methylumbelliferone and its glucuronide and sulfate conjugates. Application to studies in the single-pass *in situ* perfused rat intestine-liver preparation 563(1991)83

Methyl vinyl ether-maleic anhydride copolymer

Immobilization of protein ligands with methyl vinyl ether-maleic anhydride copolymer 597(1992)123

Metoprolol

Liquid chromatographic separation of the enantiomers of metoprolol and its α -hydroxy metabolite on Chiralcel OD for determination in plasma and urine 553(1991)391

Chiral stationary phase designed for β -blockers 557(1991)173

Solid-phase extraction and direct high-performance liquid chromatographic determination of metoprolol enantiomers in plasma 567(1991)221

Stereospecific high-performance liquid chromatographic assay of metoprolol 579(1992)361

Reversed retention order and other stereoselective effects in the separation of amino alcohols on Chiralcel OD 592(1992)331

Metoxychlor

Solid-matrix partition for separation of organochlorine pesticide residues from fatty materials 552(1991)241

Metrifonate

High-performance liquid chromatographic method using ultraviolet detection for measuring metrifonate and dichlorvos levels in human plasma 573(1992)99

Metyrapol

Assay of metyrapone, metyrapol and the isomeric mono-N-oxides of metyrapone in biological fluids by high-performance liquid chromatography 563(1991)283

Metyrapone

Assay of metyrapone, metyrapol and the isomeric mono-N-oxides of metyrapone in biological fluids by high-performance liquid chromatography 563(1991)283

Metyrapone N-oxides

Assay of metyrapone, metyrapol and the isomeric mono-N-oxides of metyrapone in biological fluids by high-performance liquid chromatography 563(1991)283

Mexiletine

Fluorimetric determination of mexiletine in serum by high-performance liquid chromatography using pre-column derivatization with fluorecamine 566(1991)187

Simultaneous determination of mexiletine and four hydroxylated metabolites in human serum by high-performance liquid chromatography and its application to pharmacokinetic studies 573(1992)235

Improved high-performance liquid chromatographic assay for the stereoselective determination of mexiletine in plasma 579(1992)366

Mianserin

Automated high-performance liquid chromatographic method for the determination of mianserin in plasma using electrochemical detection 582(1992)268

Microfilariae

Isolation of microfilariae from blood by gravitational field-flow fractionation 572(1991)291

Midazolam

Solid-phase extraction of midazolam and two of its metabolites from plasma for high-performance liquid chromatographic analysis 571(1991)298

MIF-1

Use of L-prolyl-L-leucylglycinamide (MIF-1) for the high-performance liquid chromatographic determination of proline iminopeptidase activity in rat liver 563(1991)142

Mifentidine

Rapid and efficient purification of cimetropium bromide and mifentidine drug metabolite mixtures derived from microsomal incubates for analysis by mass spectrometry 575(1992)75

Milacemide

Simple and rapid micro-analytical procedures for the estimation of milacemide and its metabolite glycinamide in rat plasma and cerebrospinal fluid by high-performance liquid chromatography 565(1991)357

Milk peptides

Improvement of extraction and concentration of milk peptides with solid-phase cartridges for analysis by high-performance liquid chromatography 553(1991)155

Minocycline

Identification and determination of the two principal metabolites of minocycline in humans 568(1991)363

Minoxidil

Separation of minoxidil and its intermediates by overpressured layer chromatography using a stationary phase bonded with tricaprilmethylammonium chloride 553(1991)459

Mitomycin C

Determination of the new anticancer agent KW 2149, 7-N-[2-((2-(γ -L-glutamylamino)ethyl)dithio)ethyl]mitomycin C, an analogue of mitomycin C 564(1991)352

Mitoxanthrone

Quantification of mitoxanthrone in bone marrow by high-performance liquid chromatography with electrochemical detection 563(1991)435

Mivacurium

Stereoselective high-performance liquid chromatographic assay with fluorometric detection for the isomers of mivacurium in human plasma 578(1992)302

MK-571

High-performance liquid chromatographic method for the determination of a novel leukotriene D₄ antagonist (MK-0571) in biological specimens 563(1991)458

Stereoselective determination of *R*(-)- and *S*(+)-MK-571, a leukotriene D₄ antagonist, in human plasma by chiral high-performance liquid chromatography 570(1991)157

Modafinil

Confirmation of the structure of by-products in the synthesis of Modafinil by liquid chromatography-mass spectrometry 557(1991)489

Molybdate

Determination of proteins in urine by high-performance liquid chromatography with spectrophotometric detection using a pyrogallol red-molybdate complex 565(1991)173

Monoacylglycerols

Resolution of lipolysis mixtures from soybean oil by a solid-phase extraction procedure 589(1992)353

Separation of monoacylglycerols by reversed-phase high-performance liquid chromatography 596(1992)185

Monoamine oxidase

Determination of platelet monoamine oxidase activity by high-performance liquid chromatography with electrochemical detection 575(1992)39

Monoamine oxidase B

Determination of an irreversible inhibitor of monoamine oxidase B (MDL 72974A) in human plasma and urine by gas chromatography-positive-ion chemical ionization mass spectrometry 571(1991)241

Monoamine oxidase inhibitors

Lipophilicity determination of some monoamine oxidase inhibitors: the effect of methanol and ammonium chloride 575(1992)57

Monoamines

Determination of melatonin and monoamines in rat pineal using reversed-phase ion-interaction chromatography with fluorescence detection 564(1991)93

Determination of γ -glutamyl conjugates of monoamines by means of high-performance liquid chromatography with electrochemical detection and application to gastropod tissues 567(1991)49

Monocarboxylic acids, aliphatic

Comparison of several methods for the determination of trace amounts of polar aliphatic monocarboxylic acids by high-performance liquid chromatography 593(1992)243

Monochloroacetic acid

Isotachophoretic separation of minor components from a matrix component in the case of strong electrolytes 585(1991)297

Monoclonal antibodies

see also Antibodies

Comparison of the performance of immunosorbents prepared by site-directed or random coupling of monoclonal antibodies 558(1991)55

Electrophoretic mobility modeling of proteins in free zone capillary electrophoresis and its application to monoclonal antibody microheterogeneity analysis 559(1991)357

Immobilization of ultra-thin layer of monoclonal antibody on glass surface 566(1991)361

Study of the separation of mouse monoclonal antibodies by pseudobioaffinity chromatography using matrix-linked histidine and histamine 570(1991)29

Reversed-phase high-performance liquid chromatographic tryptic digest peptide map comparisons of monoclonal antibodies to human tumor necrosis factor 574(1992)219

Development and optimization of a single-step procedure using protein A affinity chromatography to isolate murine IgG₁ monoclonal antibodies from hybridoma supernatants 587(1991)61

Immobilization of Fv antibody fragments on porous silica and their utility in affinity chromatography 587(1991)161

Rapid, two-step purification process for the preparation of pyrogen-free murine immunoglobulin G₁ monoclonal antibodies 590(1992)255

Surface-modified membranes as a matrix for protein purification 591(1992)107

High-performance gel permeation chromatographic analysis of immunoglobulin M produced by hybridoma cell culture 594(1992)179

Immobilization of monoclonal antibodies for affinity chromatography using a chelating peptide 595(1992)113

Computer simulation of weak affinity chromatography 597(1992)83

Membrane-based receptor affinity chromatography 597(1992)155

Use of antibody fragments in immunoaffinity chromatography. Comparison of FV fragments, VH fragments and paralog peptides 597(1992)239

Affinity chromatographic purification of immunoglobulin M antibodies utilizing immobilized mannan binding protein 597(1992)247

Analysis of the interaction between an $\alpha(1 \rightarrow 6)$ dextran-specific mouse hybridoma antibody and dextran B512 by affinity electrophoresis 597(1992)345

Separation of monoclonal antibodies from antihapten antisera by two-dimensional affinity electrophoresis 597(1992)365

Mapping of viral epitopes with conformationally specific monoclonal antibodies using biosensor technology 597(1992)391

Biospecific interaction analysis using surface plasmon resonance detection applied to kinetic, binding site and concentration analysis 597(1992)397

Selective high-performance liquid chromatographic purification of bispecific monoclonal antibodies 599(1992)13

Purification and quantification of recombinant Epstein-Barr viral glycoproteins gp350/220 from Chinese hamster ovary cells 599(1992)267

N-Monodealkyldisopyramide

Simplified, rapid and inexpensive extraction procedure for a high-performance liquid chromatographic method for determination of disopyramide and its main metabolite mono-N-dealkylated disopyramide in serum 572(1991)339

Automated determination of disopyramide and N-monodealkyldisopyramide in plasma by reversed-phase liquid chromatography with a column switching system 574(1992)365

Monodesmethyltrimipramine

Simultaneous determination of trimipramine and its major metabolites by high-performance liquid chromatography 566(1991)228

Monoethylglycinexylidide

High-performance liquid chromatographic method for the simultaneous determination of monoethylglycinexylidide and lignocaine 574(1992)361

Monoferric transferrin

High-performance liquid chromatographic separation of human apotransferrin and monoferric and diferric transferrins 567(1991)248

Monohydroxy compounds

High-performance liquid chromatographic determination of monohydroxy compounds by a combination of pre-column derivatization and post-column reaction detection 593(1992)29

Mono-O-methyl-D-glucose

Analyses of isomeric mono-O-methyl-D-glucoses, D-glucobioses and D-glucose monophosphates by high-performance anion-exchange chromatography with pulsed amperometric detection 595(1992)340

Mononitrates, organic

Gas chromatographic analysis of isomeric organic mononitrates in plasma 571(1991)109

Monosaccharides

Carbohydrate analysis with ion chromatography using Eurokat stationary phases. Preparative separation of monosaccharides and their fluorinated derivatives 552(1991)483

High-performance liquid chromatographic determination of sugars and polyols in extracts of lichens and sugarcane juice 553(1991)271

Separation of neutral mono- and oligosaccharides derivatized with ethyl *p*-aminobenzoate by high-performance liquid chromatography on an amine-bonded vinyl alcohol copolymer column 588(1991)53

Study of dihydroxy-substituted saturated urushiol crown ether as a stationary phase in capillary gas chromatography 589(1992)368

High-performance liquid chromatography and supercritical fluid chromatography of monosaccharides and polyols using light-scattering detection. Chemometric studies of the retentions 590(1992)203

New, stable polyamine-bonded polymer gel column 592(1992)93

Structure-retention index relationships for derivatized monosaccharides on non-polar gas chromatography columns 596(1992)79

Monoterpene glycosides

Analytical methods for monoterpene glycosides in grape and wine. I. XAD-2 extraction and gas chromatographic-mass spectrometric determination of synthetic glycosides 590(1992)313

Analytical methods for monoterpene glycosides in grape and wine. II. Qualitative and quantitative determination of monoterpene glycosides in grape 595(1992)269

Monoterpenes, cyclic

Study of the response of three liquid crystals as stationary phases for the gas chromatography of some cyclic monoterpene volatile oil constituents with short retention times 587(1991)343

Morphiceptin

Discriminative affinity labelling of opioid receptors by enkephalin and morphiceptin analogues containing 3-nitro-2-pyridinesulphenyl-activated thiol residues 597(1992)425

Morphine

High-performance liquid chromatographic determination of morphine and its 3- and 6-glucuronide metabolites: improvements to the method and application to stability studies 565(1991)457

Determination of morphine and 6-acetylmorphine in plasma by high-performance liquid chromatography with fluorescence detection 566(1991)135

Simple method for the determination of morphine and its active glucuronide metabolite in human plasma by high-performance liquid chromatography with electrochemical detection 570(1991)191

Direct determination of codeine, norcodeine, morphine and normorphine with their corresponding O-glucuronide conjugates by high-performance liquid chromatography with electrochemical detection 570(1991)309

High-performance liquid chromatographic determination of morphine, morphine-3-glucuronide, morphine-6-glucuronide and codeine in biological samples using multi-wavelength forward optical detection 571(1991)263

Column-switching high-performance liquid chromatographic detection of pholcodine and its metabolites in urine with fluorescence and electrochemical detection 573(1992)283

Simultaneous solid-phase extraction and chromatographic analysis of morphine and hydromorphone in plasma by high-performance liquid chromatography with electrochemical detection 577(1992)354

High-performance liquid chromatographic determination of morphine, morphine-3-glucuronide, morphine-6-glucuronide and codeine in biological samples using multi-wavelength forward optical detection: a reply 578(1992)333

High-performance liquid chromatographic determination of morphine, morphine-3-glucuronide, morphine-6-glucuronide and codeine in biological samples using multi-wavelength forward optical detection. Reply to Zweipfenning 578(1992)335

Improved high-performance liquid chromatographic method for the determination of ethylmorphine and its metabolites in microsomal incubations and cell culture media 579(1992)158

High-performance liquid chromatographic determination of morphine, morphine-3-glucuronide, morphine-6-glucuronide and codeine in biological samples using multi-wavelength forward optical detection: a reply 579(1992)189

High-performance liquid chromatographic determination of morphine, morphine-3-glucuronide, morphine-6-glucuronide and codeine in biological samples using multi-wavelength forward optical detection. Reply to Bogusz 579(1992)191

Rapid assay of cocaine, opiates and metabolites by gas chromatography-mass spectrometry (Review) 580(1992)43

Morphine glucuronides

High-performance liquid chromatographic determination of morphine and its 3- and 6-glucuronide metabolites: improvements to the method and application to stability studies 565(1991)457

Simple method for the determination of morphine and its active glucuronide metabolite in human plasma by high-performance liquid chromatography with electrochemical detection 570(1991)191

High-performance liquid chromatographic determination of morphine, morphine-3-glucuronide, morphine-6-glucuronide and codeine in biological samples using multi-wavelength forward optical detection 571(1991)263

High-performance liquid chromatographic determination of morphine, morphine-3-glucuronide, morphine-6-glucuronide and codeine in biological samples using multi-wavelength forward optical detection: a reply 578(1992)333

High-performance liquid chromatographic determination of morphine, morphine-3-glucuronide, morphine-6-glucuronide and codeine in biological samples using multi-wavelength forward optical detection. Reply to Zweipfenning 578(1992)335

High-performance liquid chromatographic determination of morphine, morphine-3-glucuronide, morphine-6-glucuronide and codeine in biological samples using multi-wavelength forward optical detection: a reply 579(1992)189

High-performance liquid chromatographic determination of morphine, morphine-3-glucuronide, morphine-6-glucuronide and codeine in biological samples using multi-wavelength forward optical detection. Reply to Bogusz 579(1992)191

Morpholino anthracycline

Determination of the new morpholino anthracycline MX2-HCl and its metabolites in biological samples by high-performance liquid chromatography 570(1991)203

3-O-Morpholinoethylmorphine

On-line dialysis and weak cation-exchange enrichment of dialysate. Automated high-performance liquid chromatography of pholcodine in human plasma and whole blood 582(1992)123

Motilin

Capillary zone electrophoresis studies of motilin peptides. Effects of charge, hydrophobicity, secondary structure and length 559(1991)391

MPC-1304

Simultaneous determination of a new dihydropyridine calcium antagonist (MPC-1304) and its metabolite in dog plasma by high-performance liquid chromatography with electrochemical detection 571(1991)277

Muconic acid

Determination of benzene metabolites in urine of mice by solid-phase extraction and high-performance liquid chromatography 593(1992)147

Multiple antigen peptides

Application of capillary zone electrophoresis to the characterization of multiple antigen peptides 557(1991)307

Multiple proteins factors

Isolation of multiple protein factors involved in ribosomal DNA transcription 587(1991)25

Muramic acid

Determination of muramic acid by high-performance liquid chromatography-plasma spray mass spectrometry 598(1992)43

Murine immunoglobulin G₁

Rapid, two-step purification process for the preparation of pyrogen-free murine immunoglobulin G₁ monoclonal antibodies 590(1992)255

Mustard

Determination of chemical warfare agents, their hydrolysis products and related compounds in soil 589(1992)287

Mutagens

Cellulose bearing covalently linked copper phthalocyanine trisulphonate as an adsorbent selective for polycyclic compounds and its use in studies of environmental mutagens and carcinogens (Review) 597(1992)37

MX2-HCl

Determination of the new morpholino anthracycline MX2-HCl and its metabolites in biological samples by high-performance liquid chromatography 570(1991)203

Mycosamine

Identification of aromatic moieties and mycosamine in antifungal heptaenes with high-performance liquid chromatography, high-performance liquid chromatography-mass spectrometry and gas chromatography-mass spectrometry 585(1991)247

Mycotoxins

Retention data for five ketotrichothecenes in reversed-phase high-performance liquid chromatography with different eluent systems 558(1991)81

Detection of zearalenone in cereal extracts using high-performance liquid chromatography with post-column derivatization 588(1991)47

Automated determination of aflatoxin B₁ in cattle feed by two-column solid-phase extraction with on-line high-performance liquid chromatography 592(1992)265

Fluorescence detection of trichothecene mycotoxins as coumarin-3-carbonyl chloride derivatives by high-performance liquid chromatography 595(1992)143

Determination of α - and β -zearealenol and zearalenone in cereals by gas chromatography with ion-trap detection 595(1992)259

Identification of ochratoxin A in food samples by chemical derivatization and gas chromatography-mass spectrometry 595(1992)364

Myoglobin

Electrospray interfacing for the coupling of ion-exchange and ion-pairing chromatography to mass spectrometry 554(1991)27

Instrumentation for high-performance capillary electrophoresis-mass spectrometry 559(1991)197

Peptide mapping through the coupling of capillary electrophoresis and high-performance liquid chromatography: map prediction of the tryptic digest of myoglobin 572(1991)51

Myosin light-chain kinase

Use of anion-exchange high-performance liquid chromatography for the study of smooth muscle myosin light-chain kinase and its catalytic domain 566(1991)427

Myosin light chains

Comparison of ion-exchange chromatography, isoelectric precipitation and reversed-phase high-performance liquid chromatography for the separation of individual cardiac myosin light chains 572(1991)59

Myrcene

Protection of terpenes against oxidative and acid decomposition on adsorbent cartridges 589(1992)385

N

β -NAD⁺

Determination of concentrations of adenosine and other purines in human term placenta by reversed-phase high-performance liquid chromatography with photodiode-array detection: evidence for pathways of purine metabolism in the placenta 575(1992)243

NAD

Determination of nicotinamide-adenine dinucleotide and thiazole-4-carboxamide-adenine dinucleotide in human leukocytes by reversed-phase high-performance liquid chromatography 577(1992)37

NADH

Study of high-affinity interactions by quantitative affinity chromatography. Analytical expressions in terms of total ligand concentration 597(1992)57

Nadolol

Possible role for stationary phase metal interactions in the chromatography of hydroxyamines on silica 552(1991)429

Determination of nadolol in serum by high-performance liquid chromatography with fluorimetric detection 573(1992)336

Direct separation of nadolol enantiomers on a Pirkle-type chiral stationary phase 586(1991)351

Nalbuphine

Determination of nalbuphine using high-performance liquid chromatography coupled to photodiode-array detection and gas chromatography coupled to mass spectrometry 579(1992)172

6 β -Naltrexol

Determination of naltrexone and 6 β -naltrexol in plasma by high-performance liquid chromatography with coulometric detection 567(1991)485

Naltrexone

Determination of naltrexone and 6 β -naltrexol in plasma by high-performance liquid chromatography with coulometric detection 567(1991)485

Naphthalene

Calculation of retention and selectivity in reversed-phase liquid chromatography 552(1991)91

Reversed-phase liquid chromatography coupled on-line with capillary gas chromatography. I. Introduction of large volumes of aqueous mixtures through an on-column interface 552(1991)489

Magnitude of the diffusion coefficient anomaly in the critical region and its effect on supercritical fluid chromatography 557(1991)69

1-Naphthaleneacetic acid

High-performance liquid chromatographic separation and determination of small amounts of 2-naphthaleneacetic acid in 1-naphthaleneacetic acid 558(1991)435

2-Naphthaleneacetic acid

High-performance liquid chromatographic separation and determination of small amounts of 2-naphthaleneacetic acid in 1-naphthaleneacetic acid 558(1991)435

Naphthalene-2,3-dicarboxaldehyde

Separation of naphthalene-2,3-dicarboxaldehyde-labeled amino acids by high-performance capillary electrophoresis with laser-induced fluorescence detection 593(1992)265

Naphthols

Chromatography on Poly-RP and its cyano and diol derivatives using both polar and non-polar solvent systems 591(1992)89

 α -Naphthylamide

Practical strategy for the analytical separation of enantiomers by high-performance liquid chromatography 553(1991)357

Naphthylamides

(*S,S*)-Diphenylethylethanediamine derivatives as chiral selectors. II. Gasparrini-type bound chiral stationary phase with high enantioselectivity for naphthylamides 553(1991)373

2-Naphthylamine

Determination of 2-naphthylamine in urine by a novel reversed-phase high-performance liquid chromatography method 578(1992)85

(*R*)-(–)-1-(1-Naphthyl)ethyl isocyanate

Separation of secondary alcohol enantiomers using supercritical fluid chromatography 585(1991)117

 α -Naphthylethylisocyanate

Configuration analysis of unsaturated hydroxy fatty acids 586(1991)255

Naphthylphosphates

Selected applications of cyclodextrin selectors in capillary electrophoresis 559(1991)215

Naproxen

Simultaneous analysis of flunixin, naproxen, ethacrynic acid, indomethacin, phenylbutazone, mefenamic acid and thiosalicylic acid in plasma and urine by high-performance liquid chromatography and gas chromatography-mass spectrometry 568(1991)351

Simultaneous quantitative determination of naproxen, its metabolite 6-O-desmethylnaproxen and their five conjugates in plasma and urine samples by high-performance liquid chromatography on dynamically modified silica 577(1992)325

Simultaneous determination of (*R*)- and (*S*)-naproxen and (*R*)- and (*S*)-6-O-desmethylnaproxen by high-performance liquid chromatography on a Chiral-AGP column 577(1992)362

Determination of naproxen and its metabolite O-desmethylnaproxen with their acyl glucuronides in human plasma and urine by means of direct gradient high-performance liquid chromatography 578(1992)239

High-performance liquid chromatographic determination of naproxen, ibuprofen and diclofenac in plasma and synovial fluid in man 578(1992)251

Narciclasine

Determination of narciclasine in serum by reversed-phase high-performance liquid chromatography: comparison of amperometric, ultraviolet photometric and fluorescence detection 563(1991)95

Naringin

High-performance liquid chromatography of diastereomeric flavanone glycosides in *Citrus* on a β -cyclodextrin-bonded stationary phase (Cyclobond I) 588(1991)41

Narirutin

High-performance liquid chromatography of diastereomeric flavanone glycosides in *Citrus* on a β -cyclodextrin-bonded stationary phase (Cyclobond I) 588(1991)41

Navelbine

High-performance liquid chromatographic determination of navelbine in MO₄ mouse fibrosarcoma cells and biological fluids 576(1992)351

Nefazodone

High-performance liquid chromatographic method for the determination of nefazodone and its metabolites in human plasma using laboratory robotics 570(1991)129

Nefiracetam

Simultaneous determination of nefiracetam and its metabolites by high-performance liquid chromatography 575(1992)261

 α -Neoendorphins

Approach to studying proteinase specificity by continuous-flow fast atom bombardment mass spectrometry and high-performance liquid chromatography combined with photodiode-array ultraviolet detection 554(1991)83

Neohesperidin

High-performance liquid chromatography of diastereomeric flavanone glycosides in *Citrus* on a β -cyclodextrin-bonded stationary phase (Cyclobond I) 588(1991)41

Neomycin

Determination of neomycin in plasma and urine by high-performance liquid chromatography. Application to a preliminary pharmacokinetic study 571(1991)189

Neurochemicals

Preparation of an optimum mobile phase for the simultaneous determination of neurochemicals in mouse brain tissues by high-performance liquid chromatography with electrochemical detection 576(1992)245

Applicability of high-performance liquid chromatography-continuous-flow fast atom bombardment mass spectrometry for simultaneous quantitation of multiple neurochemicals 587(1991)306

Neuroleptics

Selective system of identification and determination of antidepressants and neuroleptics in serum or plasma by solid-phase extraction followed by high-performance liquid chromatography with photodiode-array detection in analytical toxicology 581(1992)75

Neuropeptides

see also β -Endorphin and Enkephalins

Approach to studying proteinase specificity by continuous-flow fast atom bombardment mass spectrometry and high-performance liquid chromatography combined with photodiode-array ultraviolet detection 554(1991)83

Analysis of tyrosine- and methionine-containing neuropeptides by fast atom bombardment mass spectrometry 562(1991)459

Analysis of proenkephalin A, proopioidmelanocortin and protachykinin neuropeptides in human lumbar cerebrospinal fluid by reversed-phase high-performance liquid chromatography, radioimmunoassay and enzymolysis 575(1992)213

Neurophysin

Oriented immobilization of peptide ligands on solid supports 591(1992)99

Neuroproteins

Simultaneous purification of the neuroproteins synapsin I and synaptophysin 596(1992)51

Neurotransmitters

Cation-exchange liquid chromatography of choline and acetylcholine on free shielded silanols of silica-based reversed-phase stationary phases 596(1992)43

Neutrophil protease

Novel method of purification of human leukocytic elastase using adsorption on a high-performance liquid chromatography gel permeation column 567(1991)57

Niacin

Application of high-performance liquid chromatography to the analysis of niacin and biotin in Italian almond cultivars 553(1991)187

Determination of a small amount of niacin in foodstuffs by high-performance liquid chromatography 588(1991)171

Nicarbazin

Determination of some drugs by micellar electrokinetic capillary chromatography. The pseudo-effective mobility as parameter for screening 585(1991)123

Nicergoline

Assay of nicergoline and three metabolites in human plasma and urine by high-performance liquid chromatography-atmospheric pressure ionization mass spectrometry 568(1991)375

Nickel

Separation of metal complexes of ethylenediaminetetraacetic acid in environmental water samples by ion chromatography with UV and potentiometric detection 558(1991)181

Thermoanalytical and chromatographic studies of copper(II), nickel(II) and oxovanadium(IV) complexes of tetradentate Schiff bases derived from β -diketones and 2,3-diaminopentane 558(1991)187

Determination of cobalt in urine by gas chromatography-mass spectrometry employing nickel as an internal standard 576(1992)297

Determination of divalent trace metals in natural waters by preconcentration on N,N,N',N'-tetra(2-aminoethyl)ethylenediamine-silica followed by on-line ion chromatography 600(1992)203

Nicotinamide

High-performance liquid chromatographic determination of nicotinamide and its metabolites in human and murine plasma and urine 582(1992)145

Rapid and quantitative separation of nicotinamide and its N¹-methylated metabolite by Dowex AG50-X4 chromatography 585(1991)322

Nicotine

Rapid and sensitive determination of low concentrations of nicotine in plasma by gas chromatography with nitrogen-specific detection 563(1991)193

Elimination of caffeine interference in high-performance liquid chromatographic determination of cotinine in human plasma 568(1991)267

Electron-impact and chemical ionization detection of nicotine and cotinine by gas chromatography-mass spectrometry in rat plasma and brain 579(1992)93

Gas chromatographic analysis of nicotine and cotinine in hair 580(1992)347

Determination of the nicotine metabolite *trans*-3'-hydroxycotinine in urine of smokers using gas chromatography with nitrogen-selective detection or selected ion monitoring 583(1992)145

Nifedipine

Effect of stationary phase structure on retention and selectivity of restricted-access reversed-phase packing materials 558(1991)19

High-performance liquid chromatographic determination of the polar metabolites of nifedipine in plasma, blood and urine 565(1991)516

Nimodipine

High-performance liquid chromatographic determination of the calcium channel blocker nimodipine in monkey plasma 578(1992)316

Nitrate

Mixed potential response at metallic electrodes in potentiometric detection for ion chromatography 589(1992)201

Determination of inorganic anions in salt solutions by ion chromatography using C_{18} reversed-phase columns coated with cetyltrimethylammonium 598(1992)237

Nitric acid

Monitoring the effluents of the trichloroacetic acid process by high-performance liquid chromatography 600(1992)316

Nitriles

Gas chromatography of Titan's atmosphere. III. Analysis of low-molecular-weight hydrocarbons and nitriles with a CP-Sil-5 CB WCOT capillary column 591(1992)297

5-Nitrosalicylic acid

Ion-pairing high-performance liquid chromatographic method for the determination of 5-aminosalicylic acid and related impurities in bulk chemical 588(1991)187

Nitrite

Mixed potential response at metallic electrodes in potentiometric detection for ion chromatography 589(1992)201

Nitroarenes

Comparison of adsorption properties of Florisil and silica in high-performance liquid chromatography. I. Retention behaviour of monofunctional model solutes 600(1992)51

Nitrobenzene

Calculation of retention and selectivity in reversed-phase liquid chromatography 552(1991)91

Thermal desorption-gas chromatography for the determination of benzene, aniline, nitrobenzene and chlorobenzene in workplace air 600(1992)344

***p*-Nitrobenzoic acid**

Chromatographic effects of residual amino groups on liquid chromatographic stationary phases derived from aminopropyl-silica gel 590(1992)189

Nitro compounds

Study of dihydroxy-substituted saturated urushiol crown ether as a stationary phase in capillary gas chromatography 589(1992)368

Nitrogen

Chromatographic determination of the physico-chemical parameters of adsorption on activated carbon fibres 600(1992)67

Nitrogen-containing aromatic compounds

Characterization of nitrogen-containing aromatic compounds in soil and sediment by capillary gas chromatography-mass spectrometry after fractionation 558(1991)223

Nitrogen-containing compounds

Quantitative aspects of the determination of compounds with widely varying polarity using capillary supercritical fluid chromatography 557(1991)99

Nitrogen gases

Determination of sulphur and nitrogen gases by gas chromatography on polystyrene porous polymer columns 585(1991)93

Nitroglycerin

Improved gas chromatographic assay for the simultaneous determination of nitroglycerin and its mono- and dinitrate metabolites 579(1992)237

Nitromethane

Estimation of cellulose polarity by gas chromatography 552(1991)365

Nitrophenols

Effects of modifiers in packed and open-tubular supercritical fluid chromatography 552(1991)527

Sensitive liquid chromatographic determination of alkyl-, nitro- and chlorophenols by precolumn derivatization with dansyl chloride, postcolumn photolysis and peroxyoxalate chemiluminescence detection 553(1991)345

Comparative study of chromatography on thin layers impregnated with organic stationary phases. Chromatographic separation of nitrophenols 558(1991)415

Simultaneous high-performance liquid chromatographic determination of urinary metabolites of benzene, nitrobenzene, toluene, xylene and styrene 573(1992)318

4-Nitrophenyl acetate

Carrier membrane as a stationary phase for affinity chromatography and kinetic studies of membrane-bound enzymes 566(1991)341

Nitrophenylaminopropanediol

Determination of chloramphenicol in muscle using a particle beam interface for combining liquid chromatography with negative-ion chemical ionization mass spectrometry 582(1992)113

***p*-Nitrophenyllaurate**

p-Nitrophenyllaurate: a substrate for the high-performance liquid chromatographic determination of lipase activity 566(1991)453

p-Nitrophenylmannoside

Study of high-affinity interactions by quantitative affinity chromatography. Analytical expressions in terms of total ligand concentration 597(1992)57

3-(2-Nitrophenylmethyl)-2-thiazolidinone

High-performance liquid chromatographic method for the determination of RGH-5702 in plasma samples 553(1991)29

4-Nitro-7-N-piperazino-2,1,3-benzoxadiazole

Precolumn fluorescence tagging reagent for carboxylic acids in high-performance liquid chromatography: 4-substituted-7-aminoalkylamino-2,1,3-benzoxadiazoles 588(1991)61

Nitrosamines

Sensitive fluorescence detection of some nitrosamines by precolumn derivatization with dansyl chloride and high-performance liquid chromatography 589(1992)349

4-Nitroso-4'-aminodiphenylmethane

Identification of metabolites of 4,4'-diaminodiphenylmethane (methylene dianiline) using liquid chromatographic and mass spectrometric techniques 583(1992)63

Nitrosochloramphenicol

Determination of chloramphenicol in muscle using a particle beam interface for combining liquid chromatography with negative-ion chemical ionization mass spectrometry 582(1992)113

6-Nitro-7-sulphamoylbenzo[*l*]quinoxaline-2,3-dione

Determination of the neuroprotective agent 6-nitro-7-sulphamoylbenzo[*l*]quinoxaline-2,3-dione in plasma using high-performance liquid chromatography 568(1991)481

Nitrous oxide

Capillary supercritical fluid chromatography of primary aliphatic amines using carbon dioxide and nitrous oxide as mobile phases 558(1991)458

Flame-photometric detection of nitrous oxide in addition to phosphine 598(1992)313

Nitroxynil

Determination of nitroxynil in cow milk by reversed-phase high-performance liquid chromatography with dual-electrode coulometric detection 596(1992)67

Nivalenol

Retention data for five ketotrichothecenes in reversed-phase high-performance liquid chromatography with different eluent systems 558(1991)81

NNC 01-0687

Determination of (+)-5-(2,3-dihydrobenzofuran-7-yl)-3-methyl-8-nitro-2,3,4,5-tetrahydro-1H-3-benzazepin-7-ol (NNC 01-0687), a novel dopamine D-1 receptor antagonist, in plasma by solid-phase extraction and high-performance liquid chromatography 577(1992)366

Nobiletin

Fast separation of polymethoxylated flavones by carbon dioxide supercritical fluid chromatography 586(1991)171

Nonapeptide

Microbore liquid chromatography coupled to a flow fast atom bombardment probe for the on-line detection of the Tyr-Pro cleavage of a nonapeptide by recombinant HIV-1 protease 562(1991)67

Non-collagenous proteins

Determination of profiles of non-collagenous proteins from rat bones by sodium dodecyl sulfate high-performance liquid chromatography 589(1992)135

Nonmercaptalbumin

Rapid analysis of human serum albumin by high-performance liquid chromatography 578(1992)292

Non-nucleoside HIV-1 reverse transcriptase inhibitor

High-performance liquid chromatographic procedure for the determination of a non-nucleoside HIV-1 reverse transcriptase inhibitor in human plasma 577(1992)129

Non-polar toxicants

Strategies for the identification of non-polar toxicants in aqueous environmental samples using toxicity-based fractionation and gas chromatography-mass spectrometry (Review) 580(1992)215

Non-volatiles

Recent methods for the determination of volatile and non-volatile organic compounds in natural and purified drinking water 562(1991)469

Nonylphenols

Reversed-phase high-performance liquid chromatography of anionic and ethoxylated non-ionic surfactants and pesticides in liquid pesticide formulations 593(1992)227

Noradrenaline

see Norepinephrine

Norcocaethylene

Determination of cocaethylene, cocaine and their metabolites in rat serum microsamples by high-performance liquid chromatography, and its application to pharmacokinetic studies in rodents 582(1992)167

Norcocaine

Rapid assay of cocaine, opiates and metabolites by gas chromatography-mass spectrometry (Review) 580(1992)43

Determination of cocaethylene, cocaine and their metabolites in rat serum microsamples by high-performance liquid chromatography, and its application to pharmacokinetic studies in rodents 582(1992)167

Norcodeine

Direct determination of codeine, norcodeine, morphine and normorphine with their corresponding O-glucuronide conjugates by high-performance liquid chromatography with electrochemical detection 570(1991)309

Nordazepam

Determination of diazepam and nordazepam in milk and plasma in the presence of oxazepam and temazepam 564(1991)330

Norephedrine

Determination of (*S*)-(-)-cathinone and its metabolites (*R,S*)-(-)-norephedrine and (*R,R*)-(-)-norpseudoephedrine in urine by high-performance liquid chromatography with photodiode-array detection 593(1992)79

Norepinephrine

Improved measurement of urinary catecholamines by liquid-liquid extraction, derivatization and high-performance liquid chromatography with fluorimetric detection 563(1991)348

Evaluation of a simple plasma catecholamine extraction procedure prior to high-performance liquid chromatography and electrochemical detection 564(1991)55

Determination of heat-exposure effects on the concentration of catecholamines in bovine plasma and milk 566(1991)29

Measurement of plasma catecholamines by high-performance liquid chromatography with electrochemical detection in intensive care patients after dobutamine infusion 567(1991)240

Analysis of plasma catecholamines by high-performance liquid chromatography with fluorescence detection: simple sample preparation for pre-column fluorescence derivatization 567(1991)351

Simultaneous determination of free catecholamines and epinine and estimation of total epinine and dopamine in plasma and urine by high-performance liquid chromatography with fluorimetric detection 574(1992)109

Optimization of high-performance liquid chromatographic assay for catecholamines. Determination of optimal mobile phase composition and elimination of species-dependent differences in extraction recovery of 3,4-dihydroxybenzylamine 574(1992)213

γ -Aminobutyric acid, catecholamine and indoleamine determinations from the same brain region by high-performance liquid chromatography with electrochemical detection 579(1992)334

Simultaneous measurement of serotonin, catecholamines and their metabolites in cat and human plasma by *in vitro* microdialysis-microbore high-performance liquid chromatography with amperometric detection 582(1992)19

Simultaneous determination of catecholamines and dobutamine in human plasma and urine by high-performance liquid chromatography with fluorimetric detection 583(1992)236

Micellar electrokinetic capillary chromatography of vitamin B₆ with electrochemical detection 585(1991)139

Determination of catecholamines in urine by liquid chromatography and electrochemical detection after on-line sample purification on immobilized boronic acid 593(1992)185

Norethylmorphine

Improved high-performance liquid chromatographic method for the determination of ethylmorphine and its metabolites in microsomal incubations and cell culture media 579(1992)158

Norfluoxetine

Solid-phase extraction of fluoxetine and norfluoxetine from serum with gas chromatography-electron-capture detection 563(1991)379

Simultaneous determination of fluoxetine and norfluoxetine enantiomers in biological samples by gas chromatography with electron-capture detection 579(1992)99

Sensitive micromethod for column liquid chromatographic determination of fluoxetine and norfluoxetine in human plasma 583(1992)217

Norketamine

Stereoselective high-performance liquid chromatographic determination of the enantiomers of ketamine and norketamine in plasma 568(1991)165

Norlignan glucosides

Determination of the norlignan glucosides of Hypoxidaceae by high-performance liquid chromatography 594(1992)131

Normetanephrine

Determination of free and conjugated normetanephrine and metanephrine in human plasma by high-performance liquid chromatography with electrochemical detection 563(1991)23

High-performance liquid chromatographic determination, with ultraviolet detection, of S-adenosyl-L-methionine and S-adenosyl-L-homocysteine in rat tissues and simultaneously of normetanephrine and metanephrine for phenylethanolamine-N-methyltransferase or catechol-O-methyltransferase activities 567(1991)254

Determination of urinary normetanephrine, metanephrine and 3-methoxytyramine by high-performance liquid chromatography with electrochemical detection: comparison between automated column-switching and manual dual-column sample purification methods 578(1992)165

Normorphine

Direct determination of codeine, norcodeine, morphine and normorphine with their corresponding O-glucuronide conjugates by high-performance liquid chromatography with electrochemical detection 570(1991)309

Norpropoxyphene

Determination of dextropropoxyphene and norpropoxyphene in plasma by high-performance liquid chromatography 581(1992)161

Norpseudoephedrine

Determination of (*S*)-(-)-cathinone and its metabolites (*R,S*)-(-)-norephedrine and (*R,R*)-(-)-norpseudoephedrine in urine by high-performance liquid chromatography with photodiode-array detection 593(1992)79

Nortestosterone

Comparison of two mass-selective detectors with special reference to the analysis of residues of anabolics 564(1991)385

Detection of 19-nortestosterone and its urinary metabolites in miniature pigs by gas chromatography-mass spectrometry 564(1991)393

Identification and characterization of 19-nortestosterone in urine of meat-producing animals 564(1991)405

Test strip enzyme immunoassays and the fast screening of nortestosterone and clenbuterol residues in urine samples at the parts per billion level 564(1991)413

Combined high-performance liquid chromatography and radioimmunoassay for the screening of 19-nortestosterone and methyltestosterone residues in meat samples 564(1991)445

Development of a competitive enzyme immunoassay for 17 α -19-nortestosterone 564(1991)451

11-Nor- Δ -9-tetrahydrocannabinol-9-carboxylic acid

Solid-phase extraction of 11-nor- Δ -9-tetrahydrocannabinol-9-carboxylic acid from human urine with gas chromatographic-mass spectrometric confirmation 567(1991)81

Gas chromatographic-mass spectrometric methods of analysis for detection of 11-nor- Δ -9-tetrahydrocannabinol-9-carboxylic acid in biological matrices (Review) 580(1992)63

Northiaden

Measurement of dothiepin and its major metabolites in plasma by high-performance liquid chromatography 581(1992)152

Northiaden-S-oxide

Measurement of dothiepin and its major metabolites in plasma by high-performance liquid chromatography 581(1992)152

Nortriptyline

Sensitive method for the quantitation of nortriptyline and 10-hydroxynortriptyline in human plasma by capillary gas chromatography with electron-capture detection 573(1992)141

Column switching and high-performance liquid chromatography in the analysis of amitriptyline, nortriptyline and hydroxylated metabolites in human plasma or serum 578(1992)273

On-line solid-phase extraction and high-performance liquid chromatographic determination of nortriptyline and amitriptyline in serum 579(1992)291

High-performance liquid chromatographic determination of amitriptyline and its main metabolites using a silica column with reversed-phase eluent. Application in mice 584(1992)249

Retention reproducibility of basic drugs in high-performance liquid chromatography on a silica column with a methanol-high-pH buffer eluent. Changes in selectivity with the age of the stationary phase 592(1992)85

Norverapamil

Simultaneous direct determination of the enantiomers of verapamil and norverapamil in plasma using a derivatized amylose high-performance liquid chromatographic chiral stationary phase 574(1992)85

Direct determination of the enantiomeric ratio of verapamil, its major metabolite norverapamil and gallopamil in plasma by chiral high-performance liquid chromatography 575(1992)255

Noscapine

Simultaneous determination of yohimbine hydrochloride, strychnine nitrate and methyltestosterone by ion-pair high-performance liquid chromatography 588(1991)344

Novacine

Separation of *Strychnos nux-vomica* alkaloids by high-performance liquid chromatography 587(1991)318

Nuclear matrix proteins

Electrophoretic analysis of nuclear matrix proteins and the potential clinical applications (Review) 569(1991)389

Nuclease

Purification of DNA-derived deoxynucleotides from leukocytes involving nuclease elution of an ion-exchange column 600(1992)229

Nucleic acids

see also DNA and RNA

Capillary zone electrophoresis: its applicability and potential in biochemical analysis (Review) 569(1991)63

Behavior of large nucleic acids in reversed-phase high-performance liquid chromatography 584(1992)135

Chromatographic fractionation of nucleic acids using microcapsules made from plant cells 585(1991)57

Inclusion and fractionated release of nucleic acids using microcapsules made from plant cells 596(1992)165

Enzymophoresis of nucleic acids by tandem capillary enzyme reactor-capillary zone electrophoresis 596(1992)251

Nucleobases

Determination of the ultraviolet absorbance and radioactivity of purine compounds separated by high-performance liquid chromatography. Application to metabolic flux rate analysis 553(1991)193

Optimization and evaluation of the performance of arrangements for UV detection in high-resolution separations using fused-silica capillaries 559(1991)163

Changes of nucleotide patterns in liver, muscle and blood during the growth of Ehrlich ascites cells: application of the reversed-phase and ion-pair reversed-phase high-performance liquid chromatography with radial compression column 563(1991)53

Nucleoside 3',5'-bisphosphate adducts

Improved reversed-phase high-performance liquid chromatographic separation of ³²P-labelled nucleoside 3',5'-bisphosphate adducts of polycyclic aromatic hydrocarbons 570(1991)65

Nucleosides

Determination of the ultraviolet absorbance and radioactivity of purine compounds separated by high-performance liquid chromatography. Application to metabolic flux rate analysis 553(1991)193

Purine and pyrimidine compounds in murine peritoneal macrophages cultured *in vitro* 553(1991)205

Changes of nucleotide patterns in liver, muscle and blood during the growth of Ehrlich ascites cells: application of the reversed-phase and ion-pair reversed-phase high-performance liquid chromatography with radial compression column 563(1991)53

Application of ion-pair high-performance liquid chromatography with radioisotope detection to *in vitro* studies of nucleoside metabolism in mitochondria 563(1991)172

Subtraction method for the high-performance liquid chromatographic measurement of plasma adenosine 574(1992)247

Quantification of 5-fluoro-2'-deoxyuridine in plasma by gas chromatography and negative-ion chemical ionization mass spectrometry 575(1992)63

Simultaneous determination of myocardial nucleotides, nucleosides, purine bases and creatine phosphate by ion-pair high-performance liquid chromatography 578(1992)39

Modified nucleosides in human serum 581(1992)31

High-performance liquid chromatographic monitoring of metabolic products resulting from the treatment of mouse hepatoma cells with N⁶-cycloalkylated nucleosides 582(1992)253

Preparative separation and analysis of the enantiomers of [³H]Abbott-69992, an HIV anti-infective nucleoside, by ligand-exchange high-performance liquid chromatography 586(1991)265

Nucleoside triphosphates

Monitoring the accumulation of nucleoside triphosphates by high-performance liquid chromatography 599(1992)137

5'-Nucleotidase

Application of solid-phase extraction on anion-exchange cartridges to quantify 5'-nucleotidase activity 573(1992)207

Nucleotides

Determination of the ultraviolet absorbance and radioactivity of purine compounds separated by high-performance liquid chromatography. Application to metabolic flux rate analysis 553(1991)193

Purine and pyrimidine compounds in murine peritoneal macrophages cultured *in vitro* 553(1991)205

Pseudo-electrochromatography-mass spectrometry: a new alternative 554(1991)339

Analysis of nucleotides by high-performance liquid chromatography with phosphorus-selective detection 557(1991)441

Reproducibility and quantitation of separation for ribonucleoside triphosphates and deoxyribonucleoside triphosphates by capillary zone electrophoresis 559(1991)247

Detection of ribose-methylated nucleotides in enzymatic hydrolysates of RNA by thermospray liquid chromatography-mass spectrometry 562(1991)225

Changes of nucleotide patterns in liver, muscle and blood during the growth of Ehrlich ascites cells: application of the reversed-phase and ion-pair reversed-phase high-performance liquid chromatography with radial compression column 563(1991)53

Separation of nucleotides in homogenates of octopus retina by ion-pair reversed-phase liquid chromatography and identification by mass spectrometry 563(1991)356

Improved ion-pair high-performance liquid chromatographic method for the quantification of a wide variety of nucleotides and sugar-nucleotides in animal cells 570(1991)77

Rapid determination of creatine, phosphocreatine, purine bases and nucleotides (ATP, ADP, AMP, GTP, GDP) in heart biopsies by gradient ion-pair reversed-phase liquid chromatography 575(1992)19

Determination of nicotinamide-adenine dinucleotide and thiazole-4-carboxamide-adenine dinucleotide in human leukocytes by reversed-phase high-performance liquid chromatography 577(1992)37

Nucleotide preparation from cells and determination of nucleotides by ion-pair high-performance liquid chromatography 577(1992)146

Simultaneous determination of myocardial nucleotides, nucleosides, purine bases and creatine phosphate by ion-pair high-performance liquid chromatography 578(1992)39

High-performance affinity chromatography of NADP⁺ dehydrogenases from cell-free extracts using a nucleotide analogue as general ligand 586(1991)51

Application of a pellicular anion-exchange resin to the separation of inorganic and organic anions by single-column anion exchange 592(1992)101

Gradient elution with shorter equilibration times in reversed-phase ion-pair chromatography 592(1992)143

Peak dispersion and separation efficiency in high-performance zone electrophoresis with gel-filled capillaries 595(1992)313

Important design features of a system for the densitometric analysis of two-dimensional flat-bed separations 596(1992)241

Nystatin

Separation of antibiotics by high-performance capillary electrophoresis with photodiode-array detection 585(1991)133

O

Ochratoxin A

Fast determination of ochratoxin A in serum by liquid chromatography: comparison with enzymic spectrofluorimetric method 570(1991)441

Identification of ochratoxin A in food samples by chemical derivatization and gas chromatography-mass spectrometry 595(1992)364

Octaethylporphyrin complexes

Control of the retention selectivity of rare earth octaethylporphyrins in reversed-phase high-performance liquid chromatography using amines as mobile phase additives 598(1992)73

Octahydro-3H-2-benzopyran-3-one

Separation of *cis* and *trans* isomers from a mosquito repellent, CIC-4, via semi-preparative high-performance liquid chromatography and the repellent effect of each 590(1992)133

Octane

Activity coefficients in binary liquid mixtures measured by reversed-flow gas chromatography 595(1992)237

Octanoic acid

Allosteric and competitive displacement of drugs from human serum albumin by octanoic acid, as revealed by high-performance liquid affinity chromatography, on a human serum albumin-based stationary phase 577(1992)305

Octopamine

Extension of the electrostatic retention model of reversed-phase ion-pair high-performance liquid chromatography to include the effect of the eluent pH 552(1991)13

***p*-*n*-Octyl-*p*'-cyanobiphenyl**

Thermodynamics of solution of non-mesomorphic solutes at infinite dilution in the smectic-A, nematic and isotropic phases of *p*-*n*-octyl-*p*'-cyanobiphenyl. A gas-liquid chromatographic study 556(1991)317

Oestr ...

see Estr ...

Ofloxacin

High-performance liquid chromatographic determination of (*S*)-(-)-ofloxacin and its metabolites in serum and urine using a solid-phase clean-up 563(1991)313

Determination of ofloxacin in human serum by high-performance liquid chromatography with column switching 573(1992)289

(*S*)-(-)-Ofloxacin N-oxide

High-performance liquid chromatographic determination of (*S*)-(-)-ofloxacin and its metabolites in serum and urine using a solid-phase clean-up 563(1991)313

Okadaic acid

High-performance liquid chromatography of okadaic acid and free fatty acids in mussels 594(1992)137

Olaquinox

Column liquid chromatographic determination of carbadox and olaquinox in feeds 558(1991)125

Olefins

Gas chromatographic separation of unsaturated polar compounds on inorganic stationary phases containing silver nitrate 552(1991)313

Ligand-exchange chromatography of alkenes on stationary phases containing palladium(II) complexes. Enantiomeric separation of *trans*-1,2-divinylcyclohexane 585(1991)309

Oligodeoxyribonucleotides

Chromatographic fractionation of nucleic acids using microcapsules made from plant cells 585(1991)57

Oligomeric surfactants

Use of an evaporative light scattering detector in reversed-phase high-performance liquid chromatography of oligomeric surfactants 552(1991)593

Oligomers

Ion-mobility spectrometry as a detection method for packed-column supercritical fluid chromatography 552(1991)551

Retention of some ethylenediamine oligomers in reversed-phase chromatography 553(1991)407

Analysis of ϵ -caprolactam and its cyclic oligomers by high-performance liquid chromatography 585(1991)333

Analysis of benzyloxy-terminated poly(1,3,6-trioxane)s by supercritical fluid chromatography 589(1992)295

Oligonucleotides

Factors influencing the resolution and quantitation of oligonucleotides separated by capillary electrophoresis on a gel-filled capillary 559(1991)43

Optimization and evaluation of the performance of arrangements for UV detection in high-resolution separations using fused-silica capillaries 559(1991)163

Prediction of migration behavior of oligonucleotides in capillary gel electrophoresis 593(1991)297

Peak dispersion and separation efficiency in high-performance zone electrophoresis with gel-filled capillaries 595(1992)313

Enzymophoresis of nucleic acids by tandem capillary enzyme reactor-capillary zone electrophoresis 596(1992)251

High-performance liquid chromatographic separation of detritylated oligonucleotides on highly cross-linked poly(styrene-divinylbenzene) particles 599(1992)113

High-performance affinity chromatography of DNA. II. Porosity effects 600(1992)221

Oligosaccharides

Characterization of cellulase-based enzyme reactors for the high-performance liquid chromatographic determination of β -D-glucan oligosaccharides 558(1991)343

Separation of fluorescent oligosaccharide derivatives by microcolumn techniques based on electrophoresis and liquid chromatography 559(1991)223

Determination of neutral oligosaccharide fractions from human milk by gel permeation chromatography 568(1991)291

Characterisation of oligosaccharides from a glycoprotein variant of human serum albumin (albumin Casebrook) using high-performance anion-exchange chromatography and nuclear magnetic resonance spectroscopy 581(1992)187

Two-dimensional mapping by high-performance liquid chromatography of pyridylamino oligosaccharides from various glycosphingolipids 586(1991)35

Recovery of sugar derivatives from 2-aminopyridine labeling mixtures for high-performance liquid chromatography using UV or fluorescence detection 587(1991)177

Separation of neutral mono- and oligosaccharides derivatized with ethyl *p*-aminobenzoate by high-performance liquid chromatography on an amine-bonded vinyl alcohol copolymer column 588(1991)53

High-performance liquid chromatographic-mass spectrometric analysis of oligosaccharides from enzymatic digestion of glycosaminoglycans. Application to human samples 591(1992)225

New, stable polyamine-bonded polymer gel column 592(1992)93

Immobilized lectin columns: useful tools for the fractionation and structural analysis of oligosaccharides (Review) 597(1992)111

Capillary zone electrophoresis of linear and branched oligosaccharides 600(1992)279

Oltipraz

High-performance liquid chromatographic method for the analysis of Oltipraz in human serum and urine 573(1992)146

Determination of Oltipraz in serum by high-performance liquid chromatography with optical absorbance and mass spectrometric detection 584(1992)207

Omeprazole

(*S,S*)-Diphenylethylethanediamine derivatives as chiral selectors. II. Gasparrini-type bound chiral stationary phase with high enantioselectivity for naphthylamides 553(1991)373

Simultaneous determination of omeprazole and its metabolites in plasma and urine by reversed-phase high-performance liquid chromatography with an alkaline-resistant polymer-coated C_{18} column 579(1992)299

Peak distortion in the column liquid chromatographic determination of omeprazole dissolved in borax buffer 586(1991)271

Omeprazole sulphone

Simultaneous determination of omeprazole and its metabolites in plasma and urine by reversed-phase high-performance liquid chromatography with an alkaline-resistant polymer-coated C_{18} column 579(1992)299

Ondansetron

Thermospray liquid chromatography-mass spectrometry for the characterisation of sulphate ester conjugates 554(1991)175

OPC-13340

Use of a carrier for quantitation of a new dihydropyridine calcium antagonist (OPC-13340) in human plasma by highly sensitive gas chromatography with negative-ion chemical ionization mass spectrometry 577(1992)275

Opiates

Rapid assay of cocaine, opiates and metabolites by gas chromatography-mass spectrometry (Review) 580(1992)43

Opioid peptides

Approach to studying proteinase specificity by continuous-flow fast atom bombardment mass spectrometry and high-performance liquid chromatography combined with photodiode-array ultraviolet detection 554(1991)83

Opioid receptors

Discriminative affinity labelling of opioid receptors by enkephalin and morphiceptin analogues containing 3-nitro-2-pyridinesulphenyl-activated thiol residues 597(1992)425

Purification and reconstitution of μ -opioid receptors in liposome 597(1992)429

Orciprenaline

Determination of β_2 -receptor agonists in bovine urine and liver by gas chromatography-tandem mass spectrometry 564(1991)515

Organic acids

see also Carboxylic acids and Fatty acids

Simultaneous gas-liquid chromatographic determination of sugars and organic acids as trimethylsilyl derivatives in vegetables and strawberries 552(1991)337

Automated screening of urine samples for carbohydrates, organic and amino acids after treatment with urease 562(1991)125

Comparison of two extraction procedures for urinary organic acids prior to gas chromatography-mass spectrometry 577(1992)341

Determination of creatinine and ultraviolet-absorbing amino acids and organic acids in urine by reversed-phase high-performance liquid chromatography 583(1992)1

Complex mixture analysis based on gas chromatography-mass spectrometry with time array detection using a beam deflection time-of-flight mass spectrometer 590(1992)329

Determination of organic acids, amino acids and saccharides by high-performance liquid chromatography and a postcolumn enzyme reactor with amperometric detection 591(1992)165

Application of a pellicular anion-exchange resin to the separation of inorganic and organic anions by single-column anion exchange 592(1992)101

Correlation between column surface area and retention of polar solutes in packed-column supercritical fluid chromatography 594(1992)291

Organic acids, urinary

Comparison of two extraction procedures for urinary organic acids prior to gas chromatography-mass spectrometry 577(1992)341

Organic anions

Application of a pellicular anion-exchange resin to the separation of inorganic and organic anions by single-column anion exchange 592(1992)101

Organic compounds

Recent methods for the determination of volatile and non-volatile organic compounds in natural and purified drinking water 562(1991)469

Organic pollutants

Chromatographic separation of marine organic pollutants (Review) 580(1992)191

Organochlorine compounds

Solid-matrix partition for separation of organochlorine pesticide residues from fatty materials 552(1991)241

Rapid method for the determination of organochlorine pesticides in milk 552(1991)249

Identification by capillary gas chromatography-mass spectrometry of volatile organohalogen compounds formed during bleaching of kraft pulp 552(1991)265

Miniaturisation of size-exclusion chromatography as a powerful clean-up tool in residue analysis 552(1991)517

Confirmation method for the identification and determination of some organophosphorus and organochlorine pesticides in cocoa beans by gas chromatography-mass spectrometry 562(1991)547

Comparative study of clean-up and fractionation methods for the determination of organochlorine pesticides in lipids by gas chromatography 591(1992)303

Organohalogen compounds

Identification by capillary gas chromatography-mass spectrometry of volatile organohalogen compounds formed during bleaching of kraft pulp 552(1991)265

Organophosphorus compounds

Evaluation of eluents in thermospray liquid chromatography-mass spectrometry for identification and determination of pesticides in environmental samples 553(1991)311

Determination of triazines and organophosphorus pesticides in water samples using solid-phase extraction 555(1991)137

Confirmation method for the identification and determination of some organophosphorus and organochlorine pesticides in cocoa beans by gas chromatography–mass spectrometry 562(1991)547

Application of the iodine–azide reagent for selective detection of thiophosphoryl compounds in thin-layer chromatography 588(1991)307

Screening of organophosphorus pesticides using liquid chromatography–atmospheric pressure chemical ionization mass spectrometry 595(1992)193

Organotin compounds

Comparison of sulphur-mode and tin-mode flame photometric detectors for the gas chromatographic determination of organotin compounds 566(1991)207

Determination of organotin species by capillary gas chromatography with alternating current plasma emission detection 598(1992)97

Ornithine

Liquid chromatographic–mass spectrometric analysis for screening of patients with cystinuria, and identification of cystine stone 568(1991)445

Orotic acid

Automated determination of orotic acid, uracil and pseudouridine in urine by high-performance liquid chromatography with column switching 568(1991)325

Ovalbumin

Purification of ovalbumin and lysozyme from a commercial product by recycling isotachopheresis 585(1991)145

Economic considerations important in the scale-up of an ovalbumin separation from hen egg-white on the anion-exchange cellulose DE92 590(1992)49

Oxadiazon

Characterization and determination of organic compounds in the mutagenic XAD-2 extracts of drinking water 557(1991)413

Oxalate

Improved high-performance liquid chromatographic separation for the analysis of oxalate in fungal culture media 587(1991)297

Oxanthrazole

High-performance liquid chromatographic analysis of the anticancer drug oxanthrazole in rat whole blood and tissues 582(1992)225

Oxazepam

Determination of diazepam and nordazepam in milk and plasma in the presence of oxazepam and temazepam 564(1991)330

Quantitation by gas chromatography with selected-ion monitoring mass spectrometry of “natural” diazepam, N-desmethyldiazepam and oxazepam in normal human serum 579(1992)85

Direct liquid chromatographic separation of enantiomers on immobilized protein stationary phases. IX. Influence of the cross-linking reagent on the retentive and enantioselective properties of chiral sorbents based on bovine serum albumin 591(1992)65

Oxiracetam

High-performance liquid chromatographic determination of monohydroxy compounds by a combination of pre-column derivatization and post-column reaction detection 593(1992)29

Oxobile acids

Capillary gas chromatographic analysis of oxo, oxo-hydroxy and unsaturated bile acid glycine conjugates 558(1991)451

Studies on steroids. CCLJII. Separation and characterization of 3-oxobile acids in serum by high-performance liquid chromatography with fluorescence detection 567(1991)343

Oxociprofloxacin

Improved high-performance liquid chromatographic determination of ciprofloxacin and its metabolites in human specimens 582(1992)263

Oxodipine

Use of particle beam liquid chromatography–electron impact mass spectrometry for structure elucidation of oxodipine and three of its metabolites 562(1991)39

Oxolinic acid

Automated analysis of oxolinic acid and flumequine in salmon whole blood and plasma using dialysis combined with trace enrichment as on-line sample preparation for high-performance liquid chromatography 564(1991)171

17-Oxosteroids

Thin-layer chromatography of urinary 17-oxosteroids using dansylhydrazine as a prelabelling reagent 570(1991)396

Oxovanadium complexes

Thermoanalytical and chromatographic studies of copper(II), nickel(II) and oxovanadium(IV) complexes of tetradentate Schiff bases derived from β -diketones and 2,3-diaminopentane 558(1991)187

Oxprenolol

Chiral stationary phase designed for β -blockers 557(1991)173

Retention and enantioselectivity of racemic solutes on a modified ovomucoid-bonded column. I. Cross-linking with glutaraldehyde 592(1992)301

Oxybenzone

Determination of sun-screen agents in cosmetic products by micellar liquid chromatography 587(1991)325

Oxydone

Nanogram level quantitation of oxycodone in human plasma by capillary gas chromatography using nitrogen-phosphorus selective detection 577(1992)283

S-Oxynitrilase

Purification of S-oxynitrilase from *Sorghum bicolor* by immobilized metal ion affinity chromatography on different carrier materials 584(1992)85

Oxyphenbutazone

Solid-phase extraction for the high-performance liquid chromatographic determination of indomethacin, suxibuzone, phenylbutazone and oxyphenbutazone in plasma, avoiding degradation of compounds 581(1992)101

Oxypurines

Simultaneous determination of allantoin and oxypurines in biological fluids by high-performance liquid chromatography 575(1992)153

Oxytetracycline

Fully automated high-performance liquid chromatographic analysis of whole blood and plasma samples using on-line dialysis as sample preparation. Determination of oxytetracycline in bovine and salmon whole blood and plasma 570(1991)99

Determination of oxytetracycline in blood serum by high-performance liquid chromatography with direct injection 573(1992)333

Determination of metacycline and related substances by column liquid chromatography on poly(styrene-divinylbenzene) 586(1991)61

High-performance liquid chromatographic analysis of oxytetracycline in chinook salmon following administration of medicated feed 588(1991)181

Oxytocin

High-performance liquid chromatography of peptides at reduced temperatures: separation of isomers 586(1991)145

Interactions and uses of antisense peptides in affinity technology (Review) 597(1992)29

P**Pachyman**

Rapid method for preparing a β -glucan-specific sensitive fraction from *Limulus (Tachypleus tridentatus)* amoebocyte lysate 567(1991)267

Paeoniflorin

Determination of cinnamic acid and paeoniflorin in traditional chinese medicinal preparations by high-performance liquid chromatography 593(1992)191

Palmitic acid

Some aspects of peak broadening in particle-beam liquid chromatography-mass spectrometry 554(1991)119

High-performance liquid chromatographic analysis of free palmitic and stearic acids in cerebrospinal fluid 565(1991)424

Pamaquine

High-performance liquid chromatographic determination of pamaquine, primaquine and carboxy primaquine in calf plasma using electrochemical detection 579(1992)123

Pamidronate

Improved determination of the bisphosphonate pamidronate disodium in plasma and urine by pre-column derivatization with fluorescamine, high-performance liquid chromatography and fluorescence detection 568(1991)261

Pancreatic enzymes

Separation of rat pancreatic secretory proteins by cation-exchange fast protein liquid chromatography 576(1992)287

Pantothenic acid

Analytical studies on the chiral separation and simultaneous determination of pantothenic acid and hopantenic acid enantiomers in rat plasma by gas chromatography-mass fragmentography 564(1991)1

Analytical studies on the chiral separation and simultaneous determination of pantothenic acid and hopantenic acid enantiomers in rat plasma by gas chromatography-mass fragmentography: a reply 576(1992)179

Analytical studies on the chiral separation and simultaneous determination of pantothenic acid and hopantenic acid enantiomers in rat plasma by gas chromatography-mass fragmentography. Reply to Küsters *et al.* 576(1992)181

Papain

Occurrence of an allosteric transition in the modification of papain with L-1-acetyl-2,3-dihydropyrrolo[2,3-*b*]-indole-2-carboxamide 597(1992)411

Papaverine

Simultaneous optimization of several chromatographic performance goals using Derringer's desirability function 586(1991)11

Parabens

Determination of sun-screen agents in cosmetic products by micellar liquid chromatography 587(1991)325

Paracetamol

Combined assay for phenacetin and paracetamol in plasma using capillary column gas chromatography-negative-ion mass spectrometry 568(1991)341

Determination of paracetamol and its four major metabolites in mouse plasma by reversed-phase ion-pair high-performance liquid chromatography 573(1992)121

In vivo sampling using loop microdialysis probes coupled to a liquid chromatograph 578(1992)157

Simplex optimization of liquid chromatography with the function of mutual information as a criterion 585(1991)187

Paraquat

Electrodialytic sample treatment coupled on-line with column liquid chromatography for the determination of basic and acidic compounds in environmental samples 600(1992)163

Paraxanthine

High-performance liquid chromatographic separation of caffeine, theophylline, theobromine and paraxanthine in rat brain and serum 570(1991)185

Simultaneous determination of caffeine and its primary demethylated metabolites in human plasma by high-performance liquid chromatography 575(1992)311

Parvalbumin

Application of high-performance capillary electrophoresis to the analysis of conformation and interaction of metal-binding proteins 559(1991)345

PEG 600

High-performance liquid chromatographic determination of PEG 600 in human urine 565(1991)297

Pendimethalin

High-performance liquid chromatographic determination of dinitroaniline herbicides in soil and water 585(1991)164

Penems

High-performance liquid chromatographic methods for the determination of the penems SCH 29482 and FCE 22101 in human serum and urine 579(1992)115

Penicillins

Determination of penicillin G in bovine plasma by high-performance liquid chromatography after pre-column derivatization 576(1992)315

Development of analytical methods for some penicillins in bovine milk by ion-paired chromatography and confirmation by thermospray mass spectrometry 567(1991)389

Monitoring of benzylpenicillin decomposition in gastric contents by capillary zone electrophoresis 583(1992)105

Separation of antibiotics by high-performance capillary electrophoresis with photodiode-array detection 585(1991)133

Hydrophobicity of β -lactam antibiotics. Explanation and prediction of their behaviour in various partitioning solvent systems and reversed-phase chromatography 585(1991)3

High-performance liquid chromatographic determination of penicillin G, penicillin V and cloxacillin in beef and pork tissues 593(1992)15

Pentoses

Carbohydrate separation by ligand-exchange liquid chromatography. Correlation between the formation of sugar-cation complexes and the elution order 558(1991)89

Pentoxifylline

Determination of pentoxifylline and its metabolites in human plasma by high-performance liquid chromatography with solid-phase extraction 575(1992)101

Pepsin isoenzymes

Peptide maps of five human pepsin isoenzymes and other aspartic proteinases 599(1992)179

Pepsinogen A

High-performance liquid chromatography: purification and chromatographic behaviour of molecular variants of pepsinogen A from human urine 571(1991)47

Peptides

see also Special issue index

Peptides

Improvement of extraction and concentration of milk peptides with solid-phase cartridges for analysis by high-performance liquid chromatography 553(1991)155

Dependence of the silanophilic effect on the concentration of preadsorbed salts and on the chemical structure of peptides in reversed-phase thin-layer chromatography 553(1991)467

Nanoscale separations combined with tandem mass spectrometry 554(1991)73

Capillary electrophoresis-atmospheric pressure ionization mass spectrometry for the characterization of peptides. Instrumental considerations for mass spectrometric detection 554(1991)311

Nanoscale separations. Capillary liquid chromatography-mass spectrometry and capillary zone electrophoresis-mass spectrometry for the determination of peptides and proteins 554(1991)329

- Characterization of a tryptic digest by high-performance displacement chromatography and mass spectrometry 557(1991)289
- Application of capillary zone electrophoresis to the characterization of multiple antigen peptides 557(1991)307
- Performances and limits of plasma desorption mass spectrometry in the primary structure determination of proteins 557(1991)359
- Capillary zone electrophoresis studies of motilin peptides. Effects of charge, hydrophobicity, secondary structure and length 559(1991)391
- Capillary isotachopheresis with concentration-gradient detection. An application to the separation of synthetic peptides 559(1991)489
- Biomedical applications of high-performance liquid chromatography-mass spectrometry with continuous-flow fast atom bombardment 562(1991)47
- Structural analysis of biologically active peptides and recombinant proteins and their modified counterparts by mass spectrometry 562(1991)377
- Analysis of tyrosine- and methionine-containing neuropeptides by fast atom bombardment mass spectrometry 562(1991)459
- Automated determination of neuroactive acidic sulphur-containing amino acids and γ -glutamyl peptides using liquid chromatography with fluorescence and electrochemical detection 566(1991)39
- Densitometry for analysis of protein and peptide hydrolysates: application to luteinizing hormone releasing hormone 568(1991)451
- Capillary zone electrophoresis: its applicability and potential in biochemical analysis (Review) 569(1991)63
- Application of capillary isotachopheresis in peptide analysis (Review) 569(1991)123
- Identification of α - and β -species of calcitonin gene-related peptide in the rat amygdala after separation with capillary zone electrophoresis 573(1992)219
- Studies on cyclic peptides related to grastisin by reversed-phase high-performance liquid chromatography 574(1992)65
- Reversed-phase high-performance liquid chromatographic tryptic digest peptide map comparisons of monoclonal antibodies to human tumor necrosis factor 574(1992)219
- Large-scale purification and characterization of recombinant tick anticoagulant peptide 574(1992)225
- Determination of a cyclic heptapeptide, a novel fibrinogen receptor antagonist; in human plasma by high-performance liquid chromatography with automated pre-column derivatization, column switching and fluorescence detection 578(1992)31
- Electrophoretic mobility and dissociation constants of tripeptides evaluated by isotachopheresis 585(1991)364
- Retention behaviour of a template-assembled synthetic protein and its amphiphilic building blocks on reversed-phase columns 586(1991)43
- High-performance liquid chromatography of peptides at reduced temperatures: separation of isomers 586(1991)145
- Multiparameter optimizations in micellar liquid chromatography using the iterative regression optimization strategy 586(1991)221
- Chromatographic separation of mixed peptides from amino acids in biological digests with volatile buffers 586(1991)259
- Application of reversed-phase medium-pressure liquid chromatography to the isolation, separation and amino acid analysis of two closely related peptide toxins of the cyanobacterium *Microcystis aeruginosa* strain PCC 7806 587(1991)71
- Applications of automated amino acid analysis using 9-fluorenylmethyl chloroformate 588(1991)107
- Solvent system for the counter-current purification of hydrophobic bombesin analogues 589(1992)362
- Monitoring 4-methoxy-2,3,6-trimethylbenzenesulphonyl deprotection of arginine-containing synthetic peptides using capillary zone electrophoresis 589(1992)380
- Oriented immobilization of peptide ligands on solid supports 591(1992)99
- Analyte focusing in capillary electrophoresis using on-line isotachopheresis 591(1992)341
- Separation techniques for biotechnology in the 1990s 592(1992)3
- High-performance liquid chromatography of amino acids, peptides and proteins. CXV. Thermodynamic behaviour of peptides in reversed-phase chromatography 593(1992)103
- Separation of leucino-statins by capillary zone electrophoresis 593(1992)259
- Computer simulation as a tool for the rapid optimization of the high-performance liquid chromatographic separation of a tryptic digest of human growth hormone 594(1992)65

Mixed-mode hydrophilic and ionic interaction chromatography rivals reversed-phase liquid chromatography for the separation of peptides 594(1992)75

Trypsin and affinity chromatography (Review) 597(1992)3

Interactions and uses of antisense peptides in affinity technology (Review) 597(1992)29

Use of antibody fragments in immunoaffinity chromatography. Comparison of FV fragments, VH fragments and paralog peptides 597(1992)239

Assay of protein drug substances present in solution mixtures by fluorescamine derivatization and capillary electrophoresis 598(1992)123

High-performance liquid chromatographic separation of *cis-trans* isomers of proline-containing peptides. I. Separation on cyclodextrin-bonded silica 598(1992)139

Biosep-SEC-S high-performance size-exclusion chromatographic columns for proteins and peptides 599(1992)25

Influence of different N- and O-linked carbohydrates on the retention times of synthetic peptides in reversed-phase high-performance liquid chromatography 599(1992)43

Polymer-coated reversed-phase packings with controlled hydrophobic properties. II. Effect on the selectivity of peptide separations 599(1992)77

Isolation of *Der pI*, the *Dermatophagoides pteronyssinus* major mite allergen, from a crude mite culture extract, purification by ion-chromatography, and comparison between the material obtained and a cDNA-coded *Der pI* 599(1992)105

Application of capillary reversed-phase high-performance liquid chromatography to high-sensitivity protein sequence analysis 599(1992)119

High-performance liquid chromatography of amino acids, peptides and proteins. CXX. Evaluation of bandwidth behaviour of proteins chromatographed on tentacle-type anion exchangers 599(1992)163

Determination of molecular mass distributions of whey protein hydrolysates by high-performance size-exclusion chromatography 599(1992)205

Prediction of peptide retention time in reversed-phase high-performance liquid chromatography 599(1992)211

Support matrix effects in the reversed-phase thin-layer chromatography of some peptides 600(1992)149

Peptides, synthetic

Determination of the enantiomeric purity of synthetic peptides by gas chromatography-mass spectrometry 562(1991)447

Peptide toxins

Application of reversed-phase medium-pressure liquid chromatography to the isolation, separation and amino acid analysis of two closely related peptide toxins of the cyanobacterium *Microcystis aeruginosa* strain PCC 7806 587(1991)71

Perchloric acid

Reaction-gas chromatography for physico-chemical investigations on aggressive gases 552(1991)319

Perfluorocycloalkanes

Wide dynamic range electron-capture detection using the electron-capture detector pulse frequency signal 595(1992)359

Perfluoroisobutylene

High-performance capillary electrophoresis of proteins from the fluid lining of the lungs of rats exposed to perfluoroisobutylene 559(1991)411

Perhalogenated compounds

Isocratic reversed-phase high-performance liquid chromatographic separation of simple perhalogenated compounds 553(1991)399

Perhydro-1,4-thiazepine-3,5-dicarboxylic acid

Determination of cystathionine and perhydro-1,4-thiazepine-3,5-dicarboxylic acid in the urine of a patient with cystathioninuria using column liquid chromatography-mass spectrometry 579(1992)318

Periodic acid

Determination of 7-methylbenz[c]acridines by capillary gas chromatography with electron-capture detection 596(1992)233

Pesticides

Development of a rational optimisation procedure for the automated sample clean-up with column switching in pesticide residue analysis 552(1991)113

Determination of phenoxy ester herbicides by gas and high-performance liquid chromatography 552(1991)235

Solid-matrix partition for separation of organochlorine pesticide residues from fatty materials 552(1991)241

Rapid method for the determination of organochlorine pesticides in milk 552(1991)249

Multi-matrix-multi-pesticide method for agricultural products 552(1991)259

Removal of organic pollutants from aqueous solution. V. Comparative study of the extraction, recovery and chromatographic separation of some organic insecticides using unloaded polyurethane foam columns 552(1991)371

Miniaturisation of size-exclusion chromatography as a powerful clean-up tool in residue analysis 552(1991)517

- Evaluation of eluents in thermospray liquid chromatography–mass spectrometry for identification and determination of pesticides in environmental samples 553(1991)311
- Interference removal in the organic trace-level analysis of aqueous environmental samples by on-line liquid chromatographic preconcentration techniques with two precolumns 553(1991)329
- Application of high-performance thin-layer chromatography and automated multiple development for the identification and determination of pesticides in water 553(1991)489
- Application of microcolumn liquid chromatography–continuous-flow fast atom bombardment mass spectrometry in environmental studies of sulfonylurea herbicides 554(1991)91
- Applications of thermospray liquid chromatography–mass spectrometry in photochemical studies of pesticides in water 554(1991)233
- Determination of triazines and organophosphorus pesticides in water samples using solid-phase extraction 555(1991)137
- Rapid on-line precolumn high-performance liquid chromatographic method for the determination of benomyl, carbendazim and aldicarb species in drinking water 555(1991)147
- Simultaneous separation and determination of hydrocarbons and organochlorine compounds by using a two-step microcolumn 555(1991)221
- Characterization and determination of organic compounds in the mutagenic XAD-2 extracts of drinking water 557(1991)413
- Determination of a metabolite of the herbicide pyridate in drinking and groundwater using high-performance liquid chromatography with amperometric detection 558(1991)369
- Mobile phase variations in thermospray liquid chromatography–mass spectrometry of pesticides 562(1991)507
- Determination of chlorotriazines in aqueous environmental samples at the ng/l level using preconcentration with a cation exchanger and on-line high-performance liquid chromatography 585(1991)67
- High-performance liquid chromatographic determination of dinitroaniline herbicides in soil and water 585(1991)164
- Determination of acephate by liquid chromatography in the presence of aqueous soil extracts 585(1991)324
- Liquid chromatographic determination of ethylenethiourea using pulsed amperometric detection 586(1991)158
- Determination of carbendazim in blueberries by reversed-phase high-performance liquid chromatography 587(1991)321
- Comparative study of clean-up and fractionation methods for the determination of organochlorine pesticides in lipids by gas chromatography 591(1992)303
- Determination of National Survey of Pesticides analytes in groundwater by liquid chromatography with postcolumn reaction detection 592(1992)283
- Reversed-phase high-performance liquid chromatography of anionic and ethoxylated non-ionic surfactants and pesticides in liquid pesticide formulations 593(1992)227
- Screening of organophosphorus pesticides using liquid chromatography–atmospheric pressure chemical ionization mass spectrometry 595(1992)193
- Relationship between gas chromatographic retention indices and molecular connectivity indices of chlorinated pesticides and structurally related compounds 598(1992)243
- Intra-injector formation of methyl esters from phenoxy acid pesticides 598(1992)309
- Application of an internal surface reversed-phase column for the automated determination of flucycloxiuron residues 600(1992)211
- Determination of the herbicide diclofop in human urine 600(1992)327
- Pethidine**
Determination of pethidine and its major metabolites in human urine by gas chromatography 565(1991)247
- Petroleum**
Quality control procedure for the gas chromatographic determination of light hydrocarbons in petroleum liquid 588(1991)259
- Petroleum asphaltenes**
Analytical study of heavy crude oil fractions by coupling of the transalkylation reaction with supercritical fluid chromatography 593(1992)321
- Petroleum products**
Simulated distillation of distillates on capillary columns: influence of the polarity of the stationary phase 557(1991)155
- PGT/1A**
Simple high-performance liquid chromatographic method for the determination of PGT/1A, a new immunostimulating drug, in biological fluids 563(1991)385
- Phallacidin**
Simultaneous assay for amatoxins and phallotoxins in *Amanita phalloides* Fr. by high-performance liquid chromatography 598(1992)227

Phallisacin

Simultaneous assay for amatoxins and phallotoxins in *Amanita phalloides* Fr. by high-performance liquid chromatography 598(1992)227

Phallisin

Simultaneous assay for amatoxins and phallotoxins in *Amanita phalloides* Fr. by high-performance liquid chromatography 598(1992)227

Phalloidin

Simultaneous assay for amatoxins and phallotoxins in *Amanita phalloides* Fr. by high-performance liquid chromatography 598(1992)227

Phalloin

Simultaneous assay for amatoxins and phallotoxins in *Amanita phalloides* Fr. by high-performance liquid chromatography 598(1992)227

Phallotoxins

Methods for chromatographic determination of amanitins and related toxins in biological samples (Review) 580(1992)279

Phenacetin

Combined assay for phenacetin and paracetamol in plasma using capillary column gas chromatography–negative-ion mass spectrometry 568(1991)341

Phenaceturic acid

Rapid and simple method for the determination of urinary benzoic and phenylacetic acids and their glycine conjugates in ruminants by reversed-phase high-performance liquid chromatography 582(1992)13

Phenacyl bromide

Derivatization of saturated long-chain fatty acids with phenacyl bromide in non-ionic micelles 586(1991)347

Phenanthrene

Reversed-phase liquid chromatography coupled on-line with capillary gas chromatography. I. Introduction of large volumes of aqueous mixtures through an on-column interface 552(1991)489

Phenobarbital

Simultaneous optimization of several chromatographic performance goals using Derünger's desirability function 586(1991)11

Preparation and characterization of mixed functional phase silica materials using phenyl-, butyl- or octylchlorosilane as a silylating agent 596(1992)151

Phenochlor

Chromatographie en phase gazeuse haute résolution–spectrométrie de masse des congénères de biphényles polychlorés dans quelques échantillons d'origine biologique 552(1991)213

Analyse par chromatographie en phase gazeuse haute résolution–spectrométrie de masse des mélanges techniques de Phénochlor 552(1991)223

Phenolcarboxylic acids

Isotachophoretic analysis of flavonoids and phenolcarboxylic acids of relevance to phytopharmaceutical industry 559(1991)499

Phenol conjugates

Gas chromatographic determination of urinary phenol conjugates after acid hydrolysis/extractive acetylation 574(1992)349

Phenolethanolamine-N-methyltransferase

High-performance liquid chromatographic determination, with ultraviolet detection, of S-adenosyl-L-methionine and S-adenosyl-L-homocysteine in rat tissues and simultaneously of normetanephrine and metanephrine for phenylethanolamine-N-methyltransferase or catechol-O-methyltransferase activities 567(1991)254

Phenolic acids

Capillary electrophoresis with electrochemical detection employing an on-column Nafion joint 593(1992)305

Phenolic compounds

High-performance liquid chromatography of the phenolic compounds in the juice of some French cider apple varieties 555(1991)125

Assessment of a thermospray interface for the coupling of high-performance liquid chromatography and Fourier transform infrared spectrometry 588(1991)15

Trace-level determination of polar phenolic compounds in aqueous samples by high-performance liquid chromatography and on-line preconcentration on porous graphitic carbon 600(1992)195

Phenolphthalein

Simultaneous determination of phenolphthalein and phenolphthalein glucuronide from dog serum, urine and bile by high-performance liquid chromatography 578(1992)231

Phenolphthalein glucuronide

Simultaneous determination of phenolphthalein and phenolphthalein glucuronide from dog serum, urine and bile by high-performance liquid chromatography 578(1992)231

Phenols

Calculation of retention and selectivity in reversed-phase liquid chromatography 552(1991)91

Quantitative two-dimensional thin-layer chromatography 553(1991)477

Packed column supercritical fluid chromatography with carbon dioxide–polar modifiers. Influence of carbon dioxide density on retention 557(1991)85

Simultaneous high-performance liquid chromatographic determination of urinary metabolites of benzene, nitrobenzene, toluene, xylene and styrene 573(1992)318

High-performance liquid chromatographic column-switching technique for the determination of intermediates of anaerobic degradation of toluene in ground water microcosm 587(1991)185

Study of dihydroxy-substituted saturated urushiol crown ether as a stationary phase in capillary gas chromatography 589(1992)368

Chemically modified resins for solid-phase extraction 590(1992)197

Retention behaviour of some ring-substituted phenol derivatives on a porous graphitized carbon column 592(1992)75

Determination of benzene metabolites in urine of mice by solid-phase extraction and high-performance liquid chromatography 593(1992)147

Suppression of the effect of metal impurities in alkylsilylated silica packing materials 594(1992)87

Comparison of commercially available thin-layer chromatography plates with mixtures of dyes, analgesics and phenols 595(1992)368

Comparison of adsorption properties of Florisil and silica in high-performance liquid chromatography. I. Retention behaviour of monofunctional model solutes 600(1992)51

Phenothiazines

Sensitive determination of phenothiazines in body fluids by gas chromatography with surface ionization detection 579(1992)247

Phenoxy acids

Evaluation of eluents in thermospray liquid chromatography-mass spectrometry for identification and determination of pesticides in environmental samples 553(1991)311

Intra-injector formation of methyl esters from phenoxy acid pesticides 598(1992)309

Phenoxy esters

Determination of phenoxy ester herbicides by gas and high-performance liquid chromatography 552(1991)235

Phenprocoumon

Thermospray and particle beam liquid chromatographic-mass spectrometric analysis of coumarin anticoagulants 562(1991)31

Phenylacetic acid

High-performance liquid chromatographic determination of phenylacetic acid in human plasma extracted with ethyl acetate 570(1991)376

Rapid and simple method for the determination of urinary benzoic and phenylacetic acids and their glycine conjugates in ruminants by reversed-phase high-performance liquid chromatography 582(1992)13

Phenylalanine

Selection of buffers and of an ion-pairing agent for thermospray liquid chromatographic-mass spectrometric analysis of ionic compounds 552(1991)605

Rapid and sensitive high-performance liquid chromatographic method for the analysis of tryptophan, tyrosine and phenylalanine in biological samples 553(1991)149

Adsorption isotherms of phenylalanine in a chromatographic column measured simultaneously by system peaks analysis and frontal analysis 556(1991)277

Phenylalanine in whole blood and plasma — a candidate reference method 562(1991)213

Rapid determination of phenylalanine in plasma by capillary high-performance liquid chromatography with electrochemical detection 563(1991)134

Studies of the electrophoretic behaviour of *p*-dihydroxyborylphenylalanine and related compounds by the three-spot method 585(1991)369

Protein adsorption capacity of a porous phenylalanine-containing membrane based on a polyethylene matrix 586(1991)27

Stationary phase effect on enantioselectivity of dansyl phenylalanine in microcolumn liquid chromatography with γ -cyclodextrin as mobile phase additive 595(1992)121

Phenylalanine ammonia-lyase

Identification by high-performance liquid chromatography of tyrosine ammonia-lyase activity in purified fractions of *Phaseolus vulgaris* phenylalanine ammonia-lyase 573(1992)309

1-Phenylalkylamides

New chiral covalently bonded, π -donor stationary phases for high-performance liquid chromatography, based on derivatives of optically active 1-(1-naphthyl)ethylamine. Part II 585(1991)195

Phenylalkylamines

Investigations of the influence of silanol groups on the separation of enantiomers by liquid and supercritical fluid chromatography 592(1992)309

Determination of psychotropic phenylalkylamine derivatives in biological matrices by high-performance liquid chromatography with photodiode-array detection 593(1992)87

Phenylalkyl dinitrobenzoyl derivatives

New chiral covalently bonded, π -donor stationary phases for high-performance liquid chromatography, based on derivatives of optically active 1-(1-naphthyl)ethylamine. Part II 585(1991)195

Phenylalkylhydroxamic acids

Development of stability indicating assay methods for the determination of hydroxamic acids in topical formulations 594(1992)209

Phenylboronic acid

Phenylboronic acid as a versatile derivatization agent for chromatography of ecdysteroids 596(1992)271

1-Phenyl-1-butanol

Dehydrogenase-silica as a stationary phase for the separation of alcohols and ketones 587(1991)268

Phenylbutazone

Simultaneous analysis of flunixin, naproxen, ethacrynic acid, indomethacin, phenylbutazone, mefenamic acid and thiosalicylic acid in plasma and urine by high-performance liquid chromatography and gas chromatography-mass spectrometry 568(1991)351

Solid-phase extraction for the high-performance liquid chromatographic determination of indomethacin, suxibuzone, phenylbutazone and oxyphenbutazone in plasma, avoiding degradation of compounds 581(1992)101

***p*-Phenylenediamine**

Quantitation and linearity for particle-beam liquid chromatography-mass spectrometry 554(1991)103

Phenylephrine

Retention reproducibility of basic drugs in high-performance liquid chromatography on a silica column with a methanol-high-pH buffer eluent. Changes in selectivity with the age of the stationary phase 592(1992)85

2-Phenylethanol

Prediction of single and binary profiles in overloaded elution chromatography using various semi-ideal models 556(1991)205

Phenylethylamines

Utility of dansyl derivatization to the high-performance liquid chromatographic analysis of 2-phenylethylamine drugs 555(1991)278

Determination of urinary β -phenylethylamine as its N-benzenesulphonamide derivative by gas chromatography with flame photometric detection 578(1992)120

Enantiomeric separation of α -phenylethylamine and its substituted isomers by gas chromatography 586(1991)139

Improvement of extraction procedure for biogenic amines in foods and their high-performance liquid chromatographic determination 591(1992)175

Phenylglycine dinitrobenzoyl derivatives

New chiral covalently bonded, π -donor stationary phases for high-performance liquid chromatography, based on derivatives of optically active 1-(1-naphthyl)ethylamine. Part II 585(1991)195

Phenylglyoxylic acid

Simultaneous high-performance liquid chromatographic determination of urinary metabolites of benzene, nitrobenzene, toluene, xylene and styrene 573(1992)318

Phenylmercury

Evaluation of gas chromatographic columns for the determination of methylmercury in aqueous head space extracts from biological samples 586(1991)329

1-Phenyl-3-methyl-5-pyrazolone

High-performance capillary zone electrophoresis of carbohydrates in the presence of alkaline earth metal ions 588(1991)327

3-Phenylpropanol

Prediction of single and binary profiles in overloaded elution chromatography using various semi-ideal models 556(1991)205

Phenylpropanolamine

Effect of stationary phase structure on retention and selectivity of restricted-access reversed-phase packing materials 558(1991)19

Phenylserine

High-performance ligand-exchange chromatography of some amino acids containing two chiral centres 592(1992)345

Phenylureas

Evaluation of eluents in thermospray liquid chromatography-mass spectrometry for identification and determination of pesticides in environmental samples 553(1991)311

Interference removal in the organic trace-level analysis of aqueous environmental samples by on-line liquid chromatographic preconcentration techniques with two precolumns 553(1991)329

Phenytol

Two-dimensional high-performance liquid chromatographic method to assay *p*-hydroxyphenylphenylhydantoin enantiomers in biological fluids and stereoselectivity of enzyme induction in phenytol metabolism 575(1992)109

Preparation and characterization of mixed functional phase silica materials using phenyl-, butyl- or octylchlorosilane as a silylating agent 596(1992)151

Pheromones

Simple method for collecting volatile compounds from single insects and other point sources for gas chromatographic analysis 598(1992)303

Phloridzin

High-performance liquid chromatography of the phenolic compounds in the juice of some French cider apple varieties 555(1991)125

Phloroglucinol

Effect of the addition of viscous matrices to the mobile phase on chromatographic performance in liquid chromatography-fast atom bombardment mass spectrometry 554(1991)61

Trace-level determination of polar phenolic compounds in aqueous samples by high-performance liquid chromatography and on-line preconcentration on porous graphitic carbon 600(1992)195

Pholcodine

Column-switching high-performance liquid chromatographic detection of pholcodine and its metabolites in urine with fluorescence and electrochemical detection 573(1992)283

On-line dialysis and weak cation-exchange enrichment of dialysate. Automated high-performance liquid chromatography of pholcodine in human plasma and whole blood 582(1992)123

Phosphates

Gas chromatographic-mass spectrometric characterization of volatile organic compounds in Barcelona tap water 562(1991)481

Determination of calcium ion in the presence of phosphate anion and collagen by capillary-type isotachopheresis 586(1991)363

Phosphatidic acids

High-performance liquid chromatography of phosphatidic acids and related polar lipids 587(1991)193

Phosphatidylcholine

Characterization of phosphatidylcholines in rabbit lung lavage fluid by positive and negative ion fast-atom bombardment mass spectrometry 562(1991)257

Fatty acid composition of glycerophospholipids in seven tissues of cod (*Gadus morhua*), determined by combined high-performance liquid chromatography and gas chromatography 565(1991)119

Rapid isocratic method for the separation and quantification of major phospholipid classes by high-performance liquid chromatography 567(1991)29

Determination of phosphatidylcholine, phosphatidylglycerol and their lyso forms from liposome dispersions by high-performance liquid chromatography using high-sensitivity refractive index detection 585(1991)239

Gas chromatographic analysis of fatty acid methyl esters: avoiding discrimination by programmed temperature vaporizing injection 588(1991)265

Improved determination of individual molecular species of phosphatidylcholine in biological samples by high-performance liquid chromatography with internal standards 593(1992)139

High-performance liquid chromatographic separation of molecular species of neutral phospholipids 598(1992)209

Phosphatidylcholine ethanolamine

High-performance liquid chromatographic separation of molecular species of neutral phospholipids 598(1992)209

Phosphatidylethanolamine

Fatty acid composition of glycerophospholipids in seven tissues of cod (*Gadus morhua*), determined by combined high-performance liquid chromatography and gas chromatography 565(1991)119

Rapid isocratic method for the separation and quantification of major phospholipid classes by high-performance liquid chromatography 567(1991)29

Gas chromatographic analysis of fatty acid methyl esters: avoiding discrimination by programmed temperature vaporizing injection 588(1991)265

Phosphatidylglycerol

Determination of phosphatidylcholine, phosphatidylglycerol and their lyso forms from liposome dispersions by high-performance liquid chromatography using high-sensitivity refractive index detection 585(1991)239

Phosphatidylinositol

Fatty acid composition of glycerophospholipids in seven tissues of cod (*Gadus morhua*), determined by combined high-performance liquid chromatography and gas chromatography 565(1991)119

Phosphatidylserine

Fatty acid composition of glycerophospholipids in seven tissues of cod (*Gadus morhua*), determined by combined high-performance liquid chromatography and gas chromatography 565(1991)119

Rapid isocratic method for the separation and quantification of major phospholipid classes by high-performance liquid chromatography 567(1991)29

Phosphine

Flame-photometric detection of nitrous oxide in addition to phosphine 598(1992)313

Phosphocreatine

Rapid determination of creatine, phosphocreatine, purine bases and nucleotides (ATP, ADP, AMP, GTP, GDP) in heart biopsies by gradient ion-pair reversed-phase liquid chromatography 575(1992)19

Rapid separation of creatine, phosphocreatine and adenosine metabolites by ion-pair reversed-phase high-performance liquid chromatography in plasma and cardiac tissue 576(1992)149

Phosphofructokinase

Affinity partitioning of erythrocytic phosphofructokinase in aqueous two-phase systems containing poly(ethylene glycol)-bound Cibacron Blue. Influence of pH, ionic strength and substrates/effectors 589(1992)127

Affinity purification of proteins using expanded beds 597(1992)129

Phosphoglucomutase

Electrophoretic phenotyping of erythrocyte enzymes (Review) 569(1991)297

Phospholipase A₂

Purification and partial characterization of phospholipase A₂ isoforms from human placenta 597(1992)299

Phospholipase C

Purification of phospholipase C by hydrophobic interaction affinity chromatography 568(1991)85

Purification and some properties of phospholipase C from *Achromobacter xylooxidans* 568(1991)315

Phospholipids

Evaluation of different packings for high-performance liquid chromatographic analysis of alkyl lysophospholipids 553(1991)285

Application of liquid chromatography–thermospray mass spectrometry in the analysis of glycerophospholipid molecular species 562(1991)191

Automated separation and quantitation of lipid fractions by high-performance liquid chromatography and mass detection 567(1991)21

Rapid isocratic method for the separation and quantification of major phospholipid classes by high-performance liquid chromatography 567(1991)29

High-performance liquid chromatographic analysis of fatty acid compositions of platelet phospholipids as their 2-nitrophenylhydrazides 568(1991)25

Determination of phospholipid fatty acid and triacylglycerol composition of rat caecal mucosa 575(1992)1

One-dimensional thin-layer chromatographic separation of phospholipids and lysophospholipids from tissue lipid extracts 581(1992)139

Determination of phosphatidylcholine, phosphatidylglycerol and their lyso forms from liposome dispersions by high-performance liquid chromatography using high-sensitivity refractive index detection 585(1991)239

High-performance liquid chromatography of phosphatidic acids and related polar lipids 587(1991)193

Quantitative planar chromatography of phospholipids with different fatty acid compositions 588(1991)303

Isolation of experimental anti-AIDS glycerophospholipids by micro-preparative reversed-phase high-performance liquid chromatography 590(1992)153

Interaction of amphipathic model lipopeptides with phospholipid bilayers 597(1992)293

Simultaneous determination of amounts of major phospholipid classes and their fatty acid composition in erythrocyte membranes using high-performance liquid chromatography and gas chromatography 598(1992)33

High-performance liquid chromatographic separation of molecular species of neutral phospholipids 598(1992)209

Optimized method for the determination of 1,2-diacyl-*sn*-glycero-3-phosphocholine and 1,2-diacyl-*sn*-glycero-3-phosphoethanolamine molecular species by enzymatic hydrolysis and gas chromatography 600(1992)257

Phosphonodipeptides

Normal- and reversed-phase high-performance liquid chromatography of some phosphonodipeptides 553(1991)143

Ion-exchange high-performance liquid chromatography of diastereoisomers of some phosphonodipeptides 600(1992)79

9-[(2-Phosphonylmethoxy)ethyl]adenine

Determination of 9-[(2-phosphonylmethoxy)ethyl]adenine in rat urine by high-performance liquid chromatography with fluorescence detection 572(1991)321

Phosphorothioate oligodeoxynucleotides

Separation of synthetic phosphorothioate oligodeoxynucleotides from their oxygenated (phosphodiester) defect species by strong-anion-exchange high-performance liquid chromatography 599(1992)35

Phosphorylase b

Base-atom recognition in protein adsorption to alkyl agaroses 597(1992)93

Phosphotransacetylase

Specific detection of acetyl-coenzyme A by reversed-phase ion-pair high-performance liquid chromatography with an immobilized enzyme reactor 590(1992)241

Photosynthetic pigments

Highly sensitive determination of photosynthetic pigments in marine *in situ* samples by high-performance liquid chromatography 558(1991)131

Phthalates

Behaviour of a common phthalate plasticizer (dioctyl phthalate) during the alkali- and/or acid-catalysed steps in an AOCS method for the preparation of methyl esters 587(1991)263

***o*-Phthaldialdehyde**

High-performance liquid chromatographic detection of enantiomeric amino alcohols after derivatization with *o*-phthaldialdehyde and various thiosugars 558(1991)311

Phthalocyanine pigments

Trace analysis of phthalocyanine pigments by high-performance liquid chromatography 592(1992)261

Physostigmine

Determination of picogram levels of heptylphysostigmine in human plasma using high-performance liquid chromatography with fluorescence detection 581(1992)227

Phytadienes

Thermodesorption-gas chromatography-mass spectrometric analysis of biological materials for potential molecular precursors of the constituents of the crude oils 552(1991)325

Phytoalexins

Studies on stress metabolites. XVI. High-performance liquid chromatographic analysis of cruciferous phytoalexins using complex ternary mobile phase gradients 598(1992)147

Phytosterols

Simplified gas chromatographic method for the simultaneous determination of phytosterols and cholesterol 563(1991)188

Piazselenol

Determination of selenium by capillary gas chromatography after high-temperature derivatization with 1,2-diamino-3,5-dibromobenzene 598(1992)105

Picotamide

Determination of picotamide in human plasma and urine by high-performance liquid chromatography 577(1992)382

Picric acid

Isotachophoretic separation of minor components from a matrix component in the case of strong electrolytes 585(1991)297

Pigments

High-performance liquid chromatographic analysis of the pigments of blood-red prickly pear (*Opuntia ficus indica*) 593(1992)177

Pilocarpic acid

High-performance liquid chromatographic determination of pilocarpine hydrochloride and its degradation products using a β -cyclodextrin column 589(1992)159

Pilocarpine

High-performance liquid chromatographic determination of pilocarpine in plasma 581(1992)293

High-performance liquid chromatographic determination of pilocarpine hydrochloride and its degradation products using a β -cyclodextrin column 589(1992)159

Pindolol

Chiral stationary phase designed for β -blockers 557(1991)173

 α -Pinene

Use of a Sep-Pak cartridge for preparative collection in gas chromatography 555(1991)302

Effects of collection solvent parameters and extraction cell geometry on supercritical fluid extraction efficiencies 594(1992)297

Pinenes

Protection of terpenes against oxidative and acid decomposition on adsorbent cartridges 589(1992)385

Pipecolic acid

Analysis of pipecolic acid in biological fluids using capillary gas chromatography with electron-capture detection and [$^2\text{H}_1$]pipecolic acid as internal standard 574(1992)335

2,6-Pipecoloxylidide

High-performance liquid chromatographic method for the quantitation of bupivacaine, 2,6-pipecoloxylidide and 4'-hydroxybupivacaine in plasma and urine 577(1992)103

Piperidine derivative

Experimental study on the effect of the sample size on the band profile for a binary mixture showing no competitive interaction 556(1991)195

Piribedil

Determination of piribedil and its basic metabolites in plasma by high-performance liquid chromatography 563(1991)323

Piribedil, catechol metabolite

Determination of piribedil and its basic metabolites in plasma by high-performance liquid chromatography 563(1991)323

Piribedil N-oxide

Determination of piribedil and its basic metabolites in plasma by high-performance liquid chromatography 563(1991)323

Piritramide

Sensitive determination of piritramide in human plasma by gas chromatography 571(1991)257

Pirfenol

Pirfenol determination by high-performance liquid chromatography 565(1991)497

Stereoselective high-performance liquid chromatographic assay for pirfenol enantiomers in dog plasma 571(1991)179

Piroxicam

On-line solid-phase extraction of piroxicam prior to its determination by high-performance liquid chromatography 567(1991)185

Determination of piroxicam and its major metabolites in the plasma, urine and bile of humans by high-performance liquid chromatography 576(1992)121

Piroximone

Assessment of the biotransformation of the cardiotonic agent piroximone by high-performance liquid chromatography and gas chromatography-mass spectrometry 593(1992)1

Pirprofen

Stereospecific high-performance liquid chromatographic assay of pirprofen enantiomers in rat plasma and urine 577(1992)317

Plant growth regulators

Separation of plant growth regulators by capillary electrophoresis 594(1992)335

Plant hormones

High-performance thin-layer chromatographic and high-performance liquid chromatographic determination of abscisic acid produced by cyanobacteria 555(1991)239

Plant growth regulators G₁, G₂, G₃. Synthesis, extraction and determination of leaf content in *Eucalyptus grandis* 600(1992)323

Plasma total lipids

Gas chromatographic profiles of plasma total lipids as indicators of dietary history. Correlation with fat intake based on 24-h dietary recall 564(1991)11

Plasminogens

Immobilized urokinase column as part of a specific detection system for plasminogen species separated by high-performance affinity chromatography 565(1991)183

Plastics

Gas chromatography-mass spectrometry and high-performance liquid chromatographic analyses of thermal degradation products of common plastics 555(1991)229

Platelet-activating factor

Analysis of platelet-activating factor by gas chromatography-mass spectrometry: low-energy electron impact of the corresponding 3-acetyl-2-*tert.*-butyldimethylsilyl derivative 568(1991)281

Quantification of distinct molecular species of the 2-lyso metabolite of platelet-activating factor by gas chromatography-negative-ion chemical ionization mass spectrometry 575(1992)183

Preparation of immunoaffinity mini-columns for the analysis of platelet activating factor (PAF) in biological samples 597(1992)309

Platinum complexes

Separation of some platinum(II) complexes by ionic strength gradient on a solvent-generated ion-exchange sorbent 586(1991)291

Polar compounds

Adsorption and partition mode in high-performance liquid chromatography of highly polar solutes on silica 552(1991)23

Gas-liquid microcapillary columns precoated with graphitized carbon black 552(1991)197

Chemically bonded phases for the reversed-phase high-performance liquid chromatographic separation of basic substances 552(1991)415

Polar, hydrophilic compounds in drinking water produced from surface water. Determination by liquid chromatography-mass spectrometry 554(1991)251

Correlation between column surface area and retention of polar solutes in packed-column supercritical fluid chromatography 594(1992)291

Pollutants

Polar, hydrophilic compounds in drinking water produced from surface water. Determination by liquid chromatography-mass spectrometry 554(1991)251

Chromatographic separation of marine organic pollutants (Review) 580(1992)191

Poly(1,3,6-trioxocanes)

Analysis of benzyloxy-terminated poly(1,3,6-trioxocane)s by supercritical fluid chromatography 589(1992)295

Polyacrylamide, uncross-linked

DNA electrophoresis in uncross-linked polyacrylamide solution, studied by epifluorescence microscopy 596(1992)141

Polyacrylamides

Beaded polymer supports and gels (Review). II. Physico-chemical criteria and functionalization 586(1991)199

Polyadenylic acid

High-performance affinity chromatography of messenger RNA 588(1991)157

Polyamic acid

Elution behaviour of polyamic acid and polyamide-imide in size-exclusion chromatography 588(1991)195

Polyamide-imide

Elution behaviour of polyamic acid and polyamide-imide in size-exclusion chromatography 588(1991)195

Polyamines

Application of high-performance liquid chromatography to the study of biogenic amine-related enzymes (Review) 566(1991)287

Monitoring of the uptake and metabolism of aminoxy analogues of polyamines in cultured cells by high-performance liquid chromatography 574(1992)17

Standardization in the determination of red blood cell polyamines by high-performance liquid chromatography 574(1992)134

Polyaromatic hydrocarbons

Effects of collection solvent parameters and extraction cell geometry on supercritical fluid extraction efficiencies 594(1992)297

Use of basic alumina in fractionation of fossil fuels 600(1992)243

Polychlorinated biphenyls

Chromatographie en phase gazeuse haute résolution-spectrométrie de masse des congénères de biphényles polychlorés dans quelques échantillons d'origine biologique 552(1991)213

Analyse par chromatographie en phase gazeuse haute résolution-spectrométrie de masse des mélanges techniques de Phénochlor 552(1991)223

New Czechoslovak sorbent, Ekosorb 552(1991)449

Comparative analysis of thirty polychlorinated biphenyl congeners on two capillary columns of different polarity with non-linear multi-level calibration 558(1991)375

Problems associated with interferences in the analysis of serum for polychlorinated biphenyls 566(1991)117

Polychlorinated biphenyls in the environment (Review) 595(1992)1

Polychlorinated dibenzo-*p*-dioxins

Optimization of the gas chromatographic analysis of a standard mixture of polychlorodibenzo-*p*-dioxins and polychlorodibenzofurans 557(1991)481

Comparison of extraction methods for polychlorinated dibenzo-*p*-dioxins and dibenzofurans in fly ash using gas chromatography-mass spectrometry 589(1992)265

Preparation of nitrophenylethylsilylated silica gel and its chromatographic properties in the separation of polychlorinated dibenzo-*p*-dioxins 595(1992)77

Automated clean-up procedure for the gas chromatographic-high-resolution mass spectrometric determination of polychlorinated dibenzo-*p*-dioxins and dibenzofurans in milk 595(1992)289

Polychlorinated dibenzofurans

Optimization of the gas chromatographic analysis of a standard mixture of polychlorodibenzo-*p*-dioxins and polychlorodibenzofurans 557(1991)481

Comparison of extraction methods for polychlorinated dibenzo-*p*-dioxins and dibenzofurans in fly ash using gas chromatography-mass spectrometry 589(1992)265

Polyclonal antibodies

Purification and determination of glutamine synthetase by high-performance immunoaffinity chromatography 589(1992)121

Polycyclic aromatic hydrocarbons

see Polynuclear aromatic hydrocarbons

Polyesters

Reactor for prechromatographic fusion reactions 600(1992)352

Polyethylene

Gas chromatography-mass spectrometry and high-performance liquid chromatographic analyses of thermal degradation products of common plastics 555(1991)229

Determination of Cyasorb UV 1084 and its degradation products in low-density polyethylene film by high-performance liquid chromatography 587(1991)205

Polyethylene glycol

Analysis of polyethylene glycol and derivatives by high-performance liquid chromatography using elevated temperatures and low-wavelength ultraviolet detection, and supercritical fluid chromatography 553(1991)423

Immobilization method for polyethylene glycol using a cross-linking co-agent 558(1991)323

High-performance liquid chromatographic determination of PEG 600 in human urine 565(1991)297

New extraction procedure and high-performance liquid chromatographic method for analyzing polyethylene glycol-400 in urine 578(1992)297

Predicting partition coefficients in polyethylene glycol-potassium phosphate aqueous two-phase systems 586(1991)341

Comparison of ethoxylated alcohols and polyethylene glycols by high-performance liquid chromatography and supercritical fluid chromatography using evaporative light-scattering detection 591(1992)149

Ligand-receptor interactions in affinity cell partitioning. Studies with transferrin covalently linked to monomethoxypoly (ethylene glycol) and rat reticulocytes 594(1992)97

Characterization of the heterogeneity of polyethylene glycol-modified superoxide dismutase by chromatographic and electrophoretic techniques 599(1992)141

Polyethylene glycol-modified proteins

Characterization of polyethylene glycol modified proteins using charge-reversed capillary electrophoresis 559(1991)467

Polyethylene glycol monodecyl ether

Use of an evaporative light scattering detector in reversed-phase high-performance liquid chromatography of oligomeric surfactants 552(1991)593

Poly(ethylene terephthalate)

Identification of poly(ethylene terephthalate) cyclic oligomers by liquid chromatography-mass spectrometry 554(1991)305

Polyhydroxy aromatic compounds

Chemically modified resins for solid-phase extraction 590(1992)197

Polymerase chain reaction products

Analysis of DNA restriction fragments and polymerase chain reaction products towards detection of the AIDS (HIV-1) virus in blood 559(1991)267

Polymers

Isolation and quantification of polymers from autoxidized fish oils by high-performance size-exclusion chromatography with an evaporative mass detector 552(1991)501

Determination of chemical composition of polymers by size-exclusion chromatography with coupled density and refractive index detection. III. Polyethylene oxide and polytetrahydrofuran 552(1991)507

Analysis of block copolymers by high-performance liquid chromatography under critical conditions 593(1992)233

Asymmetric-channel flow field-flow fractionation with exponential force-field programming 593(1992)339

Determination of solute-polymer interaction constants via chromatography with polymer capillaries 594(1992)29

Comparison of colorimetric and chromatographic determination of *tert.*-butylpyrocatechol 595(1992)137

Polymethacrylates

Beaded polymer supports and gels (Review). II. Physico-chemical criteria and functionalization 586(1991)199

Polymethoxylated flavones

Fast separation of polymethoxylated flavones by carbon dioxide supercritical fluid chromatography 586(1991)171

Polynuclear aromatic hydrocarbons

Column switching for the high-performance liquid chromatographic analysis of polynuclear aromatic hydrocarbons in petroleum products 552(1991)575

On-line coupling of liquid chromatography to thin-layer chromatography for the identification of polycyclic aromatic hydrocarbons in marine sediment by fluorescence excitation and emission spectroscopy 552(1991)613

Spray jet assembly interface for the coupling of reversed-phase narrow-bore liquid chromatography and Fourier transform infrared spectrometry 552(1991)635

Reliability assessment of a gas chromatographic method for polycyclic aromatic hydrocarbons in olive oil 555(1991)211

Gas chromatography-mass spectrometry and high-performance liquid chromatographic analyses of thermal degradation products of common plastics 555(1991)229

Fluorescence detection in liquid chromatography with an intensified diode-array detector 556(1991)495

Laser-excited fluorescence detection of gas-phase chromatography eluates 557(1991)39

Packed column supercritical fluid chromatography with carbon dioxide-polar modifiers. Influence of carbon dioxide density on retention 557(1991)85

Characterization and determination of organic compounds in the mutagenic XAD-2 extracts of drinking water 557(1991)413

Retention behaviour of polycyclic aromatic hydrocarbons on a liquid-crystal bonded phase in reversed-phase liquid chromatography 557(1991)459

Characterization of nitrogen-containing aromatic compounds in soil and sediment by capillary gas chromatography-mass spectrometry after fractionation 558(1991)223

Gas chromatographic-mass spectrometric analysis of tar compounds formed during pyrolysis of rice husks 562(1991)531

Gas chromatographic-mass spectrometric analysis of diols and tetrols from reactions of polycyclic aromatic hydrocarbon epoxides with hemoglobin 562(1991)563

Improved reversed-phase high-performance liquid chromatographic separation of ³²P-labelled nucleoside 3',5'-bisphosphate adducts of polycyclic aromatic hydrocarbons 570(1991)65

Comments on the Note "Liquid chromatographic determination of planar aromatic sulphur compounds in crude oil", *J. Chromatogr.*, 475 (1989) 421 585(1991)376

Separation of polycyclic aromatic hydrocarbons by micellar electrokinetic chromatography with cyclodextrins as modifiers 589(1992)333

Polynuclear aromatic hydrocarbon retention indices on SE-54 stationary phase of the volatile components of a coal tar pitch. Relationships between chromatographic retention and thermal reactivity 591(1992)287

Gas chromatographic analysis of high-molecular-weight polynuclear aromatic hydrocarbons. I. Molecular weight 328 595(1992)247

Coupled-column high-performance liquid chromatographic method for the determination of 1-hydroxypyrene in urine of subjects exposed to polycyclic aromatic hydrocarbons 600(1992)189

Polynuclear aromatic sulphur heterocycles

Comments on the Note "Liquid chromatographic determination of planar aromatic sulphur compounds in crude oil", *J. Chromatogr.*, 475 (1989) 421 585(1991)376

Gas chromatographic retention behaviour of dibenzothiophene derivatives on a smectic liquid crystalline polysiloxane stationary phase 590(1992)297

Polynucleotides

see also Special issue index

Polynucleotides

Comparison of high-performance liquid chromatography with capillary gel electrophoresis in single-base resolution of polynucleotides 558(1991)273

High-performance affinity chromatography of messenger RNA 588(1991)157

Polyolefins

Determination of polymeric hindered amine light stabilizers in polyolefins by high-performance liquid chromatography 553(1991)415

Polyols

High-performance liquid chromatographic determination of sugars and polyols in extracts of lichens and sugarcane juice 553(1991)271

Chromatographic separation of polyols by ligand exchange. Effects of the ion-exchange resin cross-linking and size 594(1992)125

Polyoxyethylene glycols

Gas-liquid chromatography in qualitative analysis. XIX. The use of antioxidants to delay the oxidation of polyoxyethylene glycol stationary phases 552(1991)161

Polyoxyethylenes

High-performance hydrophobic interaction chromatography of proteins on reversed-phase supports coated with non-ionic surfactants of polyoxyethylene type. Purification of a fungal aspartic proteinase 598(1992)181

Polyphosphates

Determination of polyphosphates in intermediate materials for detergent manufacture by ion high-performance liquid chromatography with post-column derivatization 585(1991)267

Polyprenols

Supercritical fluid chromatographic analysis of polyprenols in *Ginkgo biloba* L. 600(1992)364

Polypropylene

Gas chromatography-mass spectrometry and high-performance liquid chromatographic analyses of thermal degradation products of common plastics 555(1991)229

Determination of solute-polymer interaction constants via chromatography with polymer capillaries 594(1992)29

Polysaccharides

Characterization of cellulase-based enzyme reactors for the high-performance liquid chromatographic determination of β -D-glucan oligosaccharides 558(1991)343

Beaded polymer supports and gels (Review). II. Physico-chemical criteria and functionalization 586(1991)199

Clean-up procedure for partially methylated alditol acetate derivatives of polysaccharides 587(1991)284

Polystyrene

Reversed-phase separations of high-molecular-weight polystyrenes in a dichloromethane-acetonitrile solvent system 558(1991)440

Beaded polymer supports and gels (Review). II. Physico-chemical criteria and functionalization 586(1991)199

Polysulphides

Determination of polysulphides in blood by gas chromatography and gas chromatography-mass spectrometry 564(1991)163

Polysulphones

Degradation of polysulphones by alkali fusion 595(1992)283

Polyunsaturated fatty acids

Fatty acid composition of glycerophospholipids in seven tissues of cod (*Gadus morhua*), determined by combined high-performance liquid chromatography and gas chromatography 565(1991)119

Quantitation of phenacyl esters of retinal fatty acids by high-performance liquid chromatography 578(1992)124

High-performance liquid chromatography–thermospray mass spectrometry of hydroperoxy polyunsaturated fatty acid acetyl derivatives 579(1992)25

Polyurethane

Solid-phase extraction and high-performance liquid chromatographic analysis of a toxic compound from γ -irradiated polyurethane 600(1992)93

Poly(vinyl chloride)

Determination of solute–polymer interaction constants via chromatography with polymer capillaries 594(1992)29

Porphobilinogen deaminase

High-performance liquid chromatographic detection of pitfalls in porphobilinogen deaminase determination 570(1991)51

Porphyrins

Determination of individual porphyrins in rodent urine using high-performance liquid chromatography following clean-up by anion-exchange chromatography 563(1991)363

Use of eluent containing surfactant for the liquid chromatographic analysis of porphyrins by direct serum injection 584(1992)181

Potassium

Ion-sensitive field effect transistor as a monovalent cation detector for ion chromatography and its application to the measurement of Na^+ and K^+ concentrations in serum 566(1991)109

Potassium phosphate

Predicting partition coefficients in polyethylene glycol–potassium phosphate aqueous two-phase systems 586(1991)341

Potassium superoxide

Superoxide chemical transformation of diolepoxide polyaromatic hydrocarbon DNA adducts. Determination of benzo[*a*]pyrene-*r*-7,*t*-8,9,*c*-10-tetrahydrotetrol by gas chromatography 588(1991)273

Praziquantel

High-performance liquid chromatographic determination of the stereoselective biotransformation of the chiral drug praziquantel 578(1992)265

Prednisolone

Liquid chromatographic analysis of prednisolone, prednisone and their 20-reduced metabolites in perfusion media 563(1991)341

Characterization of prednisone, prednisolone and their metabolites by gas chromatography–mass spectrometry 565(1991)45

Assay methodology for prednisolone, prednisolone acetate and prednisolone sodium phosphate in rabbit aqueous humor and ocular physiological solutions 565(1991)89

Simultaneous analysis of prednisone, prednisolone and their major hydroxylated metabolites in urine by high-performance liquid chromatography 567(1991)39

Thermospray liquid chromatography–mass spectrometry of corticosteroids 577(1992)221

Stable isotope dilution mass spectrometry for the simultaneous determination of cortisol, cortisone, prednisolone and prednisone in plasma 579(1992)193

Prednisolone acetate

Assay methodology for prednisolone, prednisolone acetate and prednisolone sodium phosphate in rabbit aqueous humor and ocular physiological solutions 565(1991)89

Prednisolone sodium phosphate

Assay methodology for prednisolone, prednisolone acetate and prednisolone sodium phosphate in rabbit aqueous humor and ocular physiological solutions 565(1991)89

Prednisone

Liquid chromatographic analysis of prednisolone, prednisone and their 20-reduced metabolites in perfusion media 563(1991)341

Characterization of prednisone, prednisolone and their metabolites by gas chromatography–mass spectrometry 565(1991)45

Simultaneous analysis of prednisone, prednisolone and their major hydroxylated metabolites in urine by high-performance liquid chromatography 567(1991)39

Thermospray liquid chromatography–mass spectrometry of corticosteroids 577(1992)221

Stable isotope dilution mass spectrometry for the simultaneous determination of cortisol, cortisone, prednisolone and prednisone in plasma 579(1992)193

Pregnanolone

Effects of β -cyclodextrin in the mobile phase on the retention and indirect detection of non-electrolytes in reversed-phase liquid chromatography. II. Steroids 552(1991)583

Primaquine

High-performance liquid chromatographic determination of pamaquine, primaquine and carboxy primaquine in calf plasma using electrochemical detection 579(1992)123

Pristinamycins

Separation of pristinamycins by high-speed counter-current chromatography. I. Selection of solvent system and preliminary preparative studies 593(1992)363

Proanthocyanidins

Identification of the constituent flavanoid units in sainfoin proanthocyanidins by reversed-phase high-performance liquid chromatography 594(1992)117

Probenecid

Simultaneous determination of ketoprofen enantiomers and probenecid in plasma and urine by high-performance liquid chromatography. 568(1991)187

Procaine

Determination of limiting ionic mobilities and dissociation constants of some local anaesthetics. 596(1992)265

Procion navy

Modification of reversed-phase columns with dyed surfactants. Preparation of mechanically resistant efficient immobilized dyes for protein purification. 558(1991)43

Procolipase

Separation and characterization of the precursor and activated forms of porcine and human pancreatic colipase by reversed-phase liquid chromatography. 583(1992)246

Procyanidins

High-performance liquid chromatography of the phenolic compounds in the juice of some French cider apple varieties. 555(1991)125

Procymidone

Development of a rational optimisation procedure for the automated sample clean-up with column switching in pesticide residue analysis. 552(1991)113

Proenkephalin A

Analysis of proenkephalin A, proopiomelanocortin and protachykinin neuropeptides in human lumbar cerebrospinal fluid by reversed-phase high-performance liquid chromatography, radioimmunoassay and enzymolysis. 575(1992)213

Progesterone

Effects of β -cyclodextrin in the mobile phase on the retention and indirect detection of non-electrolytes in reversed-phase liquid chromatography. II. Steroids. 552(1991)583

Thin-layer chromatography on silica-coated aluminium sheet as an adjunct to radioimmunoassay of steroids. 563(1991)178

General and selective isolation procedure for high-performance liquid chromatographic determination of anabolic steroids in tissues. 588(1991)89

Prolactin

Purification of canine prolactin and growth hormone by fast protein liquid chromatography. 553(1991)117

Proline

High-performance liquid chromatographic separation of *cis-trans* isomers of proline-containing peptides. I. Separation on cyclodextrin-bonded silica. 598(1992)139

Proline iminopeptidase

Use of L-prolyl-L-leucylglycinamide (MIF-1) for the high-performance liquid chromatographic determination of proline iminopeptidase activity in rat liver. 563(1991)142

Prolintane

Retention reproducibility of basic drugs in high-performance liquid chromatography on a silica column with a methanol-high-pH buffer eluent. Changes in selectivity with the age of the stationary phase. 592(1992)85

L-Prolyl-L-leucylglycinamide

Use of L-prolyl-L-leucylglycinamide (MIF-1) for the high-performance liquid chromatographic determination of proline iminopeptidase activity in rat liver. 563(1991)142

Promethazine

Influence of chemical structure of tricyclic tertiary dimethylamines on chiral separation by reversed-phase high-performance liquid chromatography after derivatization with (-)-menthylchloroformate. 553(1991)365

Pronethanol

Chiral stationary phase designed for β -blockers. 557(1991)173

Prooppiomelanocortin

Analysis of proenkephalin A, prooppiomelanocortin and protachykinin neuropeptides in human lumbar cerebrospinal fluid by reversed-phase high-performance liquid chromatography, radioimmunoassay and enzymolysis. 575(1992)213

Propafenone

Direct separation of the enantiomers of propafenone, diprafenone and their major metabolites by high-performance liquid chromatography on modified cellulose and amylose chiral stationary phases. 585(1991)329

Propane

Preconcentration technique for introducing gaseous or volatile compounds into a capillary gas chromatographic column. 587(1991)338

n-Propanol

Analysis of zapatera olives by gas and high-performance liquid chromatography. 594(1992)259

Propazine

Determination of chlorotriazines in aqueous environmental samples at the ng/l level using preconcentration with a cation exchanger and on-line high-performance liquid chromatography. 585(1991)67

Propolis

Quantitative two-dimensional thin-layer chromatography. 553(1991)477

Propranolol

Chiral stationary phase designed for β -blockers. 557(1991)173

Measurements by microdialysis of free tissue concentrations of propranolol 568(1991)419

Liquid chromatographic analysis of propranolol enantiomers in human blood using precolumn derivatization with (+)-1-(9-fluorenyl)ethyl chloroformate 570(1991)453

Rapid high-performance liquid chromatographic method for the determination of propranolol levels in canine and feline plasma 572(1991)239

Separation of enantiomers using cellulase (CBH I) silica as a chiral stationary phase 586(1991)233

Investigation of enantioselectivity and enantiomeric elution order of propranolol and its ester derivatives on an ovomucoid-bonded column 598(1992)67

***n*-Propyl carbamate esters**

High-performance liquid chromatographic determination of monohydroxy compounds by a combination of pre-column derivatization and post-column reaction detection 593(1992)29

N-(*n*-Propyl)-N-(4-pyridinyl)-1H-indol-1-amine hydrochloride
Determination of HP 749, a potential therapeutic agent for Alzheimer's disease, in plasma by high-performance liquid chromatography 572(1991)352

2-(*N*-Propyl-*N*-2-thienylethylamino)-5-hydroxytetralin
Direct analysis of the dopamine agonist (-)-2-(*N*-propyl-*N*-2-thienylethylamino)-5-hydroxytetralin hydrochloride in plasma by high-performance liquid chromatography using two-dimensional column switching 565(1991)277

Optimization of the direct enantiomeric separation by high-performance liquid chromatography of the D-2 dopamine agonist N-0437 using response surface methodology and the multi-criteria decision-making method 558(1991)333

Propylthiouracil

Selective liquid chromatographic assay for propylthiouracil in plasma 564(1991)310

Propyphenazone

Rapid determination of propyphenazone in plasma by high-performance liquid chromatography 577(1992)387

Prostaglandins

Simultaneous determination of prostaglandins E₁, A₁ and B₁ by reversed-phase high-performance liquid chromatography for the kinetic studies of prostaglandin E₁ in solution 555(1991)73

Cyclooxygenase and lipoygenase arachidonic acid metabolism by monocytes from human immune deficiency virus-infected drug users 557(1991)507

Analysis of some eicosanoids by continuous-flow fast atom bombardment mass spectrometry 562(1991)59

Determination of 6-keto-PGF_{1α}, 2,3-dinor-6-keto-PGF_{1α}, thromboxane B₂, 2,3-dinor-thromboxane B₂, PGE₂, PGD₂ and PGF_{2α} in human urine by gas chromatography-negative ion chemical ionization mass spectrometry 562(1991)599

High-performance liquid chromatography-thermospray mass spectrometry of prostaglandin and thromboxane acetyl derivatives 568(1991)11

Gas chromatographic-mass spectrometric determination of 11-dehydrothromboxane B₂ in human urine 577(1992)1

Selective clean-up method using an immunoaffinity column following radioimmunoassay of prostaglandin F_{2α} in biological fluids 597(1992)201

Protachykinin

Analysis of proenkephalin A, proopiomelanocortin and protachykinin neuropeptides in human lumbar cerebrospinal fluid by reversed-phase high-performance liquid chromatography, radioimmunoassay and enzymolysis 575(1992)213

Protein-ligand complexes

Study of protein-ligand binding effects by direct chromatographic on-column injection 584(1992)11

Proteinases

Affinity purification of proteinases by a combination of immobilized peptidyl aldehyde and semicarbazone 581(1992)17

Isolation and purification of bacterial proteinases by means of autofocusing 585(1991)373

Silica sorbents with one- and two-site attached bacitracin in affinity chromatography 596(1992)114

Peptide maps of five human pepsin isoenzymes and other aspartic proteinases 599(1992)179

Protein C

Comparison of the performance of immunosorbents prepared by site-directed or random coupling of monoclonal antibodies 558(1991)55

Purification of sufficiently γ-carboxylated recombinant protein C and its derivatives. Calcium-dependent affinity shift in immunoaffinity and ion-exchange chromatography 597(1992)285

Proteins

see also Special issue index

Proteins

Bead cellulose derivatives as supports for immobilization and chromatographic purification of proteins 552(1991)389

- Recent advances in liquid chromatography–mass spectrometry and capillary zone electrophoresis–mass spectrometry for protein analysis 553(1991)101
- Anion-exchange chromatographic properties of α -lactalbumin eluted from quaternized polyvinylimidazole. Study of the role of the polymer coating 553(1991)165
- Nanoscale separations. Capillary liquid chromatography–mass spectrometry and capillary zone electrophoresis–mass spectrometry for the determination of peptides and proteins 554(1991)329
- High-performance membrane chromatography of proteins, a novel method of protein separation 555(1991)97
- Dilution effect of carbonate ion on isotachophoretic zones observed with scanning UV photometric detection 555(1991)247
- High-performance liquid chromatography of amino acids, peptides and proteins. CXIV. Protein interactions with porous coulombic sorbents: comparison of experimental findings with predictions of several adsorption models 557(1991)335
- Performances and limits of plasma desorption mass spectrometry in the primary structure determination of proteins 557(1991)359
- Adsorption–desorption isotherm hysteresis of β -lactoglobulin A with a weakly hydrophobic surface 557(1991)369
- Effect of stationary phase structure on retention and selectivity of restricted-access reversed-phase packing materials 558(1991)19
- Modification of reversed-phase columns with dyed surfactants. Preparation of mechanically resistant efficient immobilized dyes for protein purification 558(1991)43
- Diffusion of proteins in the chromatographic gel AcA-34 558(1991)71
- Extraction of xanthine oxidase from milk by counter-current distribution in an aqueous two-phase system 558(1991)357
- Isotachophoresis of proteins in uncoated open-tubular fused-silica capillaries with a simple approach for column conditioning 558(1991)423
- Optimization and evaluation of the performance of arrangements for UV detection in high-resolution separations using fused-silica capillaries 559(1991)163
- Free solution capillary electrophoresis of proteins using untreated fused-silica capillaries 559(1991)317
- Electrophoretic mobility modeling of proteins in free zone capillary electrophoresis and its application to monoclonal antibody microheterogeneity analysis 559(1991)357
- Capillary zone electrophoresis of proteins with hydrophilic fused-silica capillaries 559(1991)367
- High-performance capillary electrophoresis of proteins from the fluid lining of the lungs of rats exposed to perfluoroisobutylene 559(1991)411
- Rapid protein analysis by capillary electrophoresis 559(1991)445
- Characterization of polyethylene glycol modified proteins using charge-reversed capillary electrophoresis 559(1991)467
- Optimizing separation parameters in capillary isoelectric focusing 559(1991)479
- Structural analysis of biologically active peptides and recombinant proteins and their modified counterparts by mass spectrometry 562(1991)377
- Primary sequences of insulin-like growth factors 1 and 2 isolated from porcine plasma 562(1991)391
- Improved detection of human pancreatic proteins separated by two-dimensional isoelectric focusing/sodium dodecyl sulphate gel electrophoresis using a combined staining procedure 563(1991)63
- Affinity high-performance liquid chromatography on heparin-Glc-Spheron 563(1991)124
- Versatile modular microelectrophoresis system and the simultaneous preparation of large numbers of polyacrylamide gels 563(1991)184
- Determination of proteins in urine by high-performance liquid chromatography with spectrophotometric detection using a pyrogallol red–molybdate complex 565(1991)173
- Densitometry for analysis of protein and peptide hydrolysates: application to luteinizing hormone releasing hormone 568(1991)451
- Biomedical relevance of two-dimensional protein mapping (Review) 569(1991)43
- Capillary zone electrophoresis: its applicability and potential in biochemical analysis (Review) 569(1991)63
- Staining and quantification of proteins separated by polyacrylamide gel electrophoresis (Review) 569(1991)175
- Sodium dodecyl sulphate electrophoresis of urinary proteins (Review) 569(1991)243
- Clinical applications of electrophoresis of human salivary proteins (Review) 569(1991)261
- Analysis of cerebrospinal fluid proteins by electrophoresis (Review) 569(1991)281

- Electrophoresis of serum isoenzymes and proteins following acute myocardial infarction (Review) 569(1991)323
- Utility of protein electrophoretic analysis in the characterization of malignant tissues (Review) 569(1991)367
- Electrophoretic analysis of nuclear matrix proteins and the potential clinical applications (Review) 569(1991)389
- Electrophoresis of mutant proteins in inherited diseases (Review) 569(1991)407
- High-yield two-step chromatographic procedure for purification of fatty acid-binding protein from human heart 570(1991)173
- Resolution of serum aluminum-binding proteins by size-exclusion chromatography: identification of a new carrier of aluminum in human serum 576(1992)271
- Separation of rat pancreatic secretory proteins by cation-exchange fast protein liquid chromatography 576(1992)287
- pH titration curves of the desialylated human α_1 -acid glycoprotein variants by combined isoelectrofocusing-electrophoresis: utilization in the development of a fractionation method for the protein variants by chromatography on immobilized metal affinity adsorbent 577(1992)43
- Rapid method to isolate serum amyloid P component from human plasma. Characterization of the isolated protein 578(1992)130
- Polypeptide marker and disease patterns found while mapping proteins in ascitis 582(1992)87
- Rational design of purification processes for recombinant proteins 584(1992)43
- Adsorption and elution of bovine γ -globulin using an affinity membrane containing hydrophobic amino acids as ligands 585(1991)45
- Purification of ovalbumin and lysozyme from a commercial product by recycling isotachopheresis 585(1991)145
- Preparative high-yield electroelution of proteins after separation by sodium dodecyl sulphate-polyacrylamide gel electrophoresis and its application to analysis of amino acid sequences and to raise antibodies 585(1991)153
- Determination of calcium ions tightly bound to proteins 585(1991)173
- Fast protein separation by reversed-phase high-performance liquid chromatography on octadecylsilyl-bonded non-porous silica gel. Effect of particle size of column packing on column efficiency 585(1991)207
- Continuous beds for standard and micro high-performance liquid chromatography 586(1991)21
- Protein adsorption capacity of a porous phenylalanine-containing membrane based on a polyethylene matrix 586(1991)27
- Retention behaviour of a template-assembled synthetic protein and its amphiphilic building blocks on reversed-phase columns 586(1991)43
- High-resolution gel filtration of the ecdysteroid receptor-DNA complex —an alternative to the electrophoretic mobility shift assay 587(1991)11
- Hydrophobic interaction chromatography for the purification of a mycobacterial heat shock protein of relative molecular mass 60 000 587(1991)19
- Isolation of multiple protein factors involved in ribosomal DNA transcription 587(1991)25
- The cytoskeletal protein villin as a parameter for early detection of tubular damage in the human kidney 587(1991)81
- Purification and crystallization of yeast hexokinase isoenzymes. Characterization of different forms by chromatofocusing 587(1991)85
- Chromatographic characterization of a phosphate-modified zirconia support for bio-chromatographic applications 587(1991)149
- Optimization of oxidation of glycoproteins: an assay for predicting coupling to hydrazide chromatographic supports 587(1991)171
- Applications of automated amino acid analysis using 9-fluorenylmethyl chloroformate 588(1991)107
- Automated two-dimensional liquid chromatographic system for mapping proteins in highly complex mixtures 588(1991)115
- Rapid ion-exchange chromatography for preparative separation of proteins. IV. Application to bovine carbonic anhydrase III from skeletal muscle 588(1991)139
- Predictability of chromatographic protein separations. Study of size-exclusion media with narrow particle size distributions 589(1992)71
- Phosphorylated polystyrene resins in high-performance ion-exchange chromatography 589(1992)87
- Determination of profiles of non-collagenous proteins from rat bones by sodium dodecyl sulfate high-performance liquid chromatography 589(1992)135

- Radial pH distribution during capillary electrophoresis with electroosmotic flow. Analysis with high ionic strength buffers 589(1992)315
- Capillary isoelectric focusing with electroosmotic zone displacement and on-column multichannel detection 589(1992)321
- Displacement effects in preparative gradient high-performance liquid chromatographic separations 590(1992)17
- Factors affecting the separation and loading capacity of proteins in preparative gradient elution high-performance liquid chromatography 590(1992)35
- Economic considerations important in the scale-up of an ovalbumin separation from hen egg-white on the anion-exchange cellulose DE92 590(1992)49
- High-performance membrane chromatography of serum and plasma membrane proteins 590(1992)59
- Peak capacity of columns for size-exclusion chromatography 591(1992)47
- Surface-modified membranes as a matrix for protein purification 591(1992)107
- Diffusion of proteins in Sepharose Cl-B gels 591(1992)115
- Analysis of pH-dependent protein interactions with gel filtration medium 591(1992)121
- Interfacing gradient elution ion-exchange chromatography and low-angle laser light-scattering photometry for analysis of proteins 591(1992)139
- Separation techniques for biotechnology in the 1990s 592(1992)3
- Practical on-line determination of biopolymer molecular weights by high-performance liquid chromatography with classical light-scattering detection 592(1992)215
- High-performance gel permeation chromatographic analysis of immunoglobulin M produced by hybridoma cell culture 594(1992)179
- Capillary modification and evaluation using streaming potential and frontal chromatography for protein analysis in capillary electrophoresis 594(1992)325
- Continuous-flow electrophoresis: a separation criterion applied to the separation of model proteins 594(1992)351
- Continuous separation and purification of proteins by recycling isotachopheresis 594(1992)361
- Fast protein monitoring in fermentation broth using non-porous micropellicular reversed-phase columns 594(1992)392
- Immobilization of monoclonal antibodies for affinity chromatography using a chelating peptide 595(1992)113
- Reversed-phase chromatographic behaviour of β -endorphin: evidence of conformational change 595(1992)125
- High-sensitivity laser-induced fluorescence detection of native proteins in capillary electrophoresis 595(1992)319
- Ion- and ligand-exchange chromatography of proteins using porous zirconium oxide supports in organic and inorganic Lewis base eluents 596(1992)27
- Simultaneous purification of the neuroproteins synapsin I and synaptophysin 596(1992)51
- Membrane chromatographic systems for high-throughput protein separations 596(1992)173
- Important design features of a system for the densitometric analysis of two-dimensional flat-bed separations 596(1992)241
- Base-atom recognition in protein adsorption to alkyl agaroses 597(1992)93
- Immobilization of protein ligands with methyl vinyl ether-maleic anhydride copolymer 597(1992)123
- Affinity purification of proteins using expanded beds 597(1992)129
- Short-cut method for predicting the productivity of affinity chromatography 597(1992)173
- Arginine-tail method, an affinity tag procedure utilizing anhydrotrypsin agarose 597(1992)181
- Novel affinity separations based on perfluorocarbon emulsions. Use of a perfluorocarbon affinity emulsion for the purification of human serum albumin from blood plasma in a fluidised bed 597(1992)189
- Affinity purification of antibodies using immobilized FB domain of protein A 597(1992)257
- Artificial immunoglobulin G-binding protein mimetic to staphylococcal protein A. Its production and application to affinity purification of immunoglobulin G 597(1992)277
- Immunoenzymatic assay of dyes currently used in affinity chromatography for protein purification 597(1992)315
- Affinity purification and affinity characterization of carbohydrate-binding proteins in bovine kidney 597(1992)323
- Affinity chromatographic purification of bovine lung endothelin receptor using biotinylated endothelin and avidin-agarose 597(1992)331

- Affinity chromatography in purification of A₁ adenosine receptors 597(1992)335
- Immobilized metal ion affinity electrophoresis. A study with several model proteins containing histidine 597(1992)357
- Determination of the association constant of monovalent mode protein-sugar interaction by capillary zone electrophoresis 597(1992)377
- Effects of stationary phase ligand density on high-performance ion-exchange chromatography of proteins 598(1992)7
- Separon HEMA modified for immobilized metal ion affinity chromatographic separation of proteins 598(1992)15
- Assay of protein drug substances present in solution mixtures by fluorescamine derivatization and capillary electrophoresis 598(1992)123
- Protein chromatography using a continuous stationary phase 598(1992)169
- Chromatographic analysis of low-molecular-mass copper-binding ligands from the crab species *Scylla serrata* and *Portunus pelagicus* 599(1992)21
- Biosep-SEC-S high-performance size-exclusion chromatographic columns for proteins and peptides 599(1992)25
- Chromatographic analysis of tropomyosins from rabbit skeletal, chicken gizzard and earthworm muscle 599(1992)51
- Polymer-coated reversed-phase packings with controlled hydrophobic properties. I. Effect on the selectivity of protein separations 599(1992)65
- Application of capillary reversed-phase high-performance liquid chromatography to high-sensitivity protein sequence analysis 599(1992)119
- Comparison of detergents for extraction and ion-exchange high-performance liquid chromatography of Sendai virus membrane proteins 599(1992)157
- High-performance liquid chromatography of amino acids, peptides and proteins. CXX. Evaluation of bandwidth behaviour of proteins chromatographed on tentacle-type anion exchangers 599(1992)163
- High-performance hydrophobic interaction chromatography as a tool for protein refolding 599(1992)185
- Synthesis and evaluation of epoxy polymer coatings for the analysis of proteins by capillary zone electrophoresis 599(1992)227
- Schemes for efficient protein purification on a family of polymeric ion exchangers in glass columns 599(1992)239
- Isoelectric focusing field-flow fractionation. IV. Investigations on protein separations in the trapezoidal cross-section channel 600(1992)305
- Proteins, polyethylene glycol modified**
On-line characterization of polyethylene glycol-modified proteins 588(1991)125
- Proteins, self-associating**
Study of the adsorption of self-associating proteins on an anion exchanger. Application to the chromatography of β -lactoglobulin B 599(1992)255
- Protocatechuic acid**
High-performance liquid chromatography of the phenolic compounds in the juice of some French cider apple varieties 555(1991)125
- Protopin**
Analytical and quantitative studies of californin and protopin in aerial part extracts of *Eschscholtzia californica* Cham. with high-performance liquid chromatography 587(1991)314
- Protoporphyrinogen oxidase**
High-performance liquid chromatographic assays for protoporphyrinogen oxidase and ferrochelatase in human leucocytes 566(1991)383
- Prunin**
High-performance liquid chromatography of diastereomeric flavanone glycosides in *Citrus* on a β -cyclodextrin-bonded stationary phase (Cyclobond I) 588(1991)41
- Pseudobrucine**
Separation of *Strychnos nux-vomica* alkaloids by high-performance liquid chromatography 587(1991)318
- Pseudocholesterase**
Liquid chromatographic monitoring of pseudocholesterase activity: comparison of methods 575(1992)237
- Pseudoerythromycin A hemiketal**
Isolation of erythromycin A N-oxide and pseudoerythromycin A hemiketal from fermentation broth of *Saccharopolyspora erythraea* by thin-layer and high-performance liquid chromatography 558(1991)265
- Pseudostrychnine**
Separation of *Strychnos nux-vomica* alkaloids by high-performance liquid chromatography 587(1991)318
- Pseudouridine**
Automated determination of orotic acid, uracil and pseudouridine in urine by high-performance liquid chromatography with column switching 568(1991)325
- Psicaine-neu**
Determination of limiting ionic mobilities and dissociation constants of some local anaesthetics 596(1992)265

Psilocin

Analysis and isolation of indole alkaloids of fungi by high-performance liquid chromatography 593(1992)201

Psilocybin

Analysis and isolation of indole alkaloids of fungi by high-performance liquid chromatography 593(1992)201

Pulegone

Use of a Sep-Pak cartridge for preparative collection in gas chromatography 555(1991)302

Purines

see also Nucleobases

Determination of the ultraviolet absorbance and radioactivity of purine compounds separated by high-performance liquid chromatography. Application to metabolic flux rate analysis 553(1991)193

Purine and pyrimidine compounds in murine peritoneal macrophages cultured *in vitro* 553(1991)205

Rapid determination of creatine, phosphocreatine, purine bases and nucleotides (ATP, ADP, AMP, GTP, GDP) in heart biopsies by gradient ion-pair reversed-phase liquid chromatography 575(1992)19

Simultaneous determination of myocardial nucleotides, nucleosides, purine bases and creatine phosphate by ion-pair high-performance liquid chromatography 578(1992)39

Determination of substituted purines in body fluids by micellar electrokinetic capillary chromatography with direct sample injection 593(1992)275

Puriproxyfen

Reversal of elution order during direct enantiomeric separation of puriproxyfen on a cellulose-based chiral stationary phase 588(1991)177

Putrescine

Monitoring of the uptake and metabolism of aminoxy analogues of polyamines in cultured cells by high-performance liquid chromatography 574(1992)17

Improvement of extraction procedure for biogenic amines in foods and their high-performance liquid chromatographic determination 591(1992)175

Pyranocoumarins

Comparison of selectivity of silica and Florisil in the separation of natural pyranocoumarins 552(1991)453

Pyrene

Diffusion of sorbed pyrene in the bonded layer of reversed-phase silicas. Effect of alkyl chain length and pore diameter 558(1991)31

Retention behavior of β -cyclodextrin complexes of anthracene and pyrene using reversed-phase liquid chromatography. The effects of *tert.*-butanol and cyclopentanol 594(1992)37

Pyridate

Determination of a metabolite of the herbicide pyridate in drinking and groundwater using high-performance liquid chromatography with amperometric detection 558(1991)369

Pyridinium iodides

Use of non-volatile ion-pairing agents for liquid chromatographic-mass spectrometric analyses with a moving-belt interface 554(1991)281

6-(Pyridin-3-yl)quinolin-2(1H)-one

Quantitation of 6-(pyridin-3-yl)quinolin-2(1H)-one, a cardiac stimulant, and its N-oxide metabolite in dog plasma by high-performance liquid chromatography 574(1992)352

Pyridobenzothiazines

Analytical procedure for the determination of rifloxacin, a new pyridobenzothiazine, in human serum and urine by high-performance liquid chromatography 564(1991)346

Pyridoxal

Determination of vitamin B₆ vitamers and pyridoxic acid in biological samples 578(1992)45

Micellar electrokinetic capillary chromatography of vitamin B₆ with electrochemical detection 585(1991)139

High-performance liquid chromatography of thiazolidinic compounds obtained by condensation of pyridoxal 5'-phosphate or pyridoxal with amino thiols (L- or D-cysteine, cysteamine, L-cysteine ethyl ester) 586(1991)153

Pyridoxal phosphate

Determination of vitamin B₆ vitamers and pyridoxic acid in biological samples 578(1992)45

High-performance liquid chromatography of thiazolidinic compounds obtained by condensation of pyridoxal 5'-phosphate or pyridoxal with amino thiols (L- or D-cysteine, cysteamine, L-cysteine ethyl ester) 586(1991)153

Pyridoxamine

Determination of vitamin B₆ vitamers and pyridoxic acid in biological samples 578(1992)45

Micellar electrokinetic capillary chromatography of vitamin B₆ with electrochemical detection 585(1991)139

Pyridoxamine phosphate

Determination of vitamin B₆ vitamers and pyridoxic acid in biological samples 578(1992)45

4-Pyridoxic acid

Separation and quantification of the B₆ vitamers in plasma and 4-pyridoxic acid in urine of adolescent girls by reversed-phase high-performance liquid chromatography 563(1991)369

Plasma B₆ vitamers and plasma and urinary 4-pyridoxic acid concentrations of middle-aged obese black women 568(1991)333

Determination of vitamin B₆ vitamers and pyridoxic acid in biological samples 578(1992)45

Application of gas chromatography–mass spectrometry with selected ion monitoring to the urinalysis of 4-pyridoxic acid 581(1992)179

Pyridoxine

Determination of vitamin B₆ vitamers and pyridoxic acid in biological samples 578(1992)45

Micellar electrokinetic capillary chromatography of vitamin B₆ with electrochemical detection 585(1991)139

Pyridylamino oligosaccharides

Two-dimensional mapping by high-performance liquid chromatography of pyridylamino oligosaccharides from various glycosphingolipids 586(1991)35

Pyrimethamine

Reversed-phase liquid chromatographic method for the simultaneous determination of the antimalarial drugs sulfadoxine, pyrimethamine, mefloquine and its major carboxylic metabolite in plasma 571(1991)169

Pyrimidines

Murine and pyrimidine compounds in murine peritoneal macrophages cultured *in vitro* 553(1991)205

1-(2-Pyrimidinyl)piperazine

Simultaneous high-performance liquid chromatographic analysis of buspirone and its metabolite 1-(2-pyrimidinyl)-piperazine in plasma using electrochemical detection 575(1992)117

Pyrimido-*s*-triazine

Gas chromatographic–mass spectrometric method for analysis of a pyrimido-*s*-triazine and some of its metabolites in dog urine 578(1992)134

Pyrocatechol

Trace-level determination of polar phenolic compounds in aqueous samples by high-performance liquid chromatography and on-line preconcentration on porous graphitic carbon 600(1992)195

Pyrogallol

Determination of tannic acid and its phenolic metabolites in biological fluids by high-performance liquid chromatography 577(1992)77

Pyrogallol red

Determination of proteins in urine by high-performance liquid chromatography with spectrophotometric detection using a pyrogallol red–molybdate complex 565(1991)173

Pyroglutamic acid

Determination of glutamine, glutamic acid and pyroglutamic acids using high-performance liquid chromatography on dynamically modified silica 594(1992)173

3-L-Pyroglutamyl-L-thiazolidine-4-carboxylic acid

Simple high-performance liquid chromatographic method for the determination of PGT/IA, a new immunostimulating drug, in biological fluids 563(1991)385

Pyrophosphatase

Use of Zn–pyrophosphatase in the high-performance liquid chromatographic analysis of cell extracts containing ³²P-labelled inositol phosphates 575(1992)51

Pyrrolidonecarboxylic acid

Determination of volatile and non-volatile organic acids in technical sugar solutions by ion-exclusion chromatography 555(1991)65

Pyrrolidone carboxyl peptidase

Purification and characterization of recombinant pyrrolidone carboxyl peptidase of *Bacillus subtilis* 584(1992)101

Pyrroloisoquinolines

High-performance liquid chromatographic analysis of diastereomers and enantiomers of pyrroloisoquinoline antidepressants 588(1991)348

6-Pyruvoyl tetrahydropterin synthase

6-Pyruvoyl tetrahydropterin synthase assay in extracts of cultured human cells using high-performance liquid chromatography with fluorescence detection of biopterin. 570(1991)43

Q

Quaternary ammonium compounds

Evaluation of eluents in thermospray liquid chromatography–mass spectrometry for identification and determination of pesticides in environmental samples 553(1991)311

Quinapril

Trace analysis of quinapril and its active metabolite, quinaprilat, in human plasma and urine by gas chromatography–negative-ion chemical ionization mass spectrometry 578(1992)195

Determination of dioxopiperazine metabolites of quinapril in biological fluids by gas chromatography–mass spectrometry 578(1992)203

Quinaprilat

Trace analysis of quinapril and its active metabolite, quinaprilat, in human plasma and urine by gas chromatography–negative-ion chemical ionization mass spectrometry 578(1992)195

Quinidine

High-performance liquid chromatographic–mass spectrometric assay of high-value compounds for pharmaceutical use from plant cell tissue culture: *Cinchona* alkaloids 562(1991)713

High-performance liquid chromatographic method for the quantitation of quinidine and selected quinidine metabolites 572(1991)159

Quinidine-N-oxide

High-performance liquid chromatographic method for the quantitation of quinidine and selected quinidine metabolites 572(1991)159

Quinine

Measurement of quinine in filter paper-absorbed blood by high-performance liquid chromatography 570(1991)180

Quinolines

Effects of modifiers in packed and open-tubular supercritical fluid chromatography 552(1991)527

Comparison of adsorption properties of Florisil and silica in high-performance liquid chromatography. I. Retention behaviour of monofunctional model solutes 600(1992)51

8-Quinolines-thiol complexes

Reversed-phase high-performance liquid chromatography of several metal–8-quinolines-thiol complexes 596(1992)73

Quinones

Spray jet assembly interface for the coupling of reversed-phase narrow-bore liquid chromatography and Fourier transform infrared spectrometry 552(1991)635

R**Rare earth elements**

Study of isotachophoretic separation behaviour of metal cations by means of particle-induced X-ray emission. III. Analysis of a crude rare earth chloride from monazite 594(1992)371

Rare earth octaethylporphyrins

Control of the retention selectivity of rare earth octaethylporphyrins in reversed-phase high-performance liquid chromatography using amines as mobile phase additives 598(1992)73

Rare earths

Optimization of separation of rare earths in high-performance thin-layer chromatography 587(1991)359

 β_2 -Receptor agonists

see also β -Agonists

Determination of β_2 -receptor agonists in bovine urine and liver by gas chromatography–tandem mass spectrometry 564(1991)515

Receptor proteins

Affinity chromatography in purification of A₁ adenosine receptors 597(1992)335

Recombinant proteins

Trypsin and affinity chromatography (Review) 597(1992)3

Recombivax HB

Analysis of a vaccine purification process by capillary electrophoresis 559(1991)337

Reduced haloperidol

Simultaneous determination of haloperidol and its metabolite, reduced haloperidol, in plasma, blood, urine and tissue homogenates by high-performance liquid chromatography 572(1991)259

Remacemide

Quantitation of the novel anticonvulsant remacemide in rat and dog plasma and urine: application of the plasma methodology to measure the plasma protein binding of remacemide 582(1992)195

High-performance liquid chromatographic method for the simultaneous measurement of remacemide (a novel, anticonvulsant and cerebroprotectant) and an active metabolite in human plasma 583(1992)91

Remoxipride

Determination of remoxipride in human plasma and urine by reversed-phase ion-pair high-performance liquid chromatography 581(1992)300

Optimization of a peak compression system for a remoxipride metabolite (FLA797) and its application to bioanalysis 591(1992)207

Renin inhibitor

Use of dabsylation, column switching and chiral separation for the determination of a renin inhibitor in rat, marmoset and human plasma 572(1991)277

Resin acids

Supercritical carbon dioxide extraction of resin and fatty acids from sediments at pulp mill sites 594(1992)309

Quantitative analysis of total resin acids by high-performance liquid chromatography of their coumarin ester derivatives 595(1992)155

Resorcinol

Trace-level determination of polar phenolic compounds in aqueous samples by high-performance liquid chromatography and on-line preconcentration on porous graphitic carbon 600(1992)195

Retinoates

Effect of temperature and mobile phase on the retention of retinoates in reversed-phase liquid chromatography 592(1992)127

13-*cis*-Retinoic acid

Simple reversed-phase high-performance liquid chromatographic method for 13-*cis*-retinoic acid in serum 593(1992)41

Retinoids

Determination of several retinoids, carotenoids and E vitamins by high-performance liquid chromatography. Application to plasma and tissues of rats fed a diet rich in either β -carotene or canthaxanthin 553(1991)179

Determination of the retinoid (*E*)-1,2,3,4-tetrahydro-1,1,4,4-tetramethyl-6-(1-methyl-2-phenylethenyl)naphthalene and its phenolic metabolite in human plasma by reversed-phase high-performance liquid chromatography 565(1991)375

Simultaneous determination of retinol, α -tocopherol and β -carotene in serum by isocratic high-performance liquid chromatography 572(1991)103

Quantitative analysis of retinoids in biological fluids by high-performance liquid chromatography using column switching. III. Determination of the retinoid sumatriptan and its *Z*-isomer in human and animal plasma 576(1992)111

Simple reversed-phase high-performance liquid chromatographic method for 13-*cis*-retinoic acid in serum 593(1992)41

Use of direct injection precolumn techniques for the high-performance liquid chromatographic determination of the retinoids acitretin and 13-*cis*-acitretin in plasma 593(1992)55

Retinol

Simultaneous determination of retinol, α -tocopherol and β -carotene in serum by isocratic high-performance liquid chromatography 572(1991)103

Retinol-binding protein

Assay of human transthyretin-bound holo-retinol-binding protein with reversed-phase high-performance liquid chromatography 567(1991)369

Retinyl palmitate

Determination of retinyl palmitate and total vitamin A content in liver and liver-based ready-to-eat foods 577(1992)163

RG 12561

Method for separation and determination of lactone and hydroxy acid forms of a new HMG CoA reductase inhibitor (RG 12561) in plasma 567(1991)491

RGH-5702

High-performance liquid chromatographic method for the determination of RGH-5702 in plasma samples 553(1991)29

Rhodamine B

Fluorescence detector cell for use in an integrated electrically driven separation system 598(1992)115

Rhoeagenine

Analytical studies of isorhoeagenine and rhoeagenine in petal extracts of *Papaver rhoeas* L. using high-performance liquid chromatography 596(1992)276

Ribonuclease

Immobilized metal ion affinity electrophoresis. A study with several model proteins containing histidine 597(1992)357

Ribonucleoside triphosphates

Reproducibility and quantitation of separation for ribonucleoside triphosphates and deoxyribonucleoside triphosphates by capillary zone electrophoresis 559(1991)247

Ribose-methylated nucleotides

Detection of ribose-methylated nucleotides in enzymatic hydrolysates of RNA by thermospray liquid chromatography-mass spectrometry 562(1991)225

Ricinoleic acid, methyl ester

Configuration analysis of unsaturated hydroxy fatty acids 586(1991)255

Rifampicin

Automated high-performance liquid chromatographic method for the determination of rifampicin in plasma 593(1992)21

Rifapentine

High-performance liquid chromatographic determination of rifapentine and its metabolite in human plasma by direct injection into a shielded hydrophobic phase column 553(1991)35

High-performance liquid chromatographic determination of rifapentine in serum using column switching 574(1992)175

Risperidone

Determination of risperidone and 9-hydroxyrisperidone in plasma, urine and animal tissues by high-performance liquid chromatography 583(1992)223

Ritodrine

Determination of ritodrine in biological fluids of the pregnant sheep by fused-silica capillary gas chromatography using electron-capture detection 565(1991)225

RNA

Purification of common bean chloroplast DNA by DEAE cellulose column chromatography 558(1991)318

mRNA

Separation of translationally active mRNAs by reversed-phase ion-pair high-performance liquid chromatography 564(1991)115

High-performance affinity chromatography of messenger RNA 588(1991)157

tRNA

Enzymophoresis of nucleic acids by tandem capillary enzyme reactor-capillary zone electrophoresis 596(1992)251

RNA dinucleotides

Optimization of the separation of the Rp and Sp diastereomers of phosphate-methylated DNA and RNA dinucleotides 596(1992)59

RNA polymerase

Isolation of multiple protein factors involved in ribosomal DNA transcription 587(1991)25

Ro 42-5892/001

Use of dabsylation, column switching and chiral separation for the determination of a renin inhibitor in rat, marmoset and human plasma 572(1991)277

Rogletimide

On-line continuous-flow dialysis thermospray tandem mass spectrometry for quantitative screening of drugs in plasma: rogetimide 598(1992)189

Rokitamycin

Sensitive determination of josamycin and rokitamycin in plasma by high-performance liquid chromatography with fluorescence detection 575(1992)171

Rolipram

High-performance liquid chromatographic procedure for specificity testing of radioimmunoassays: rolipram 573(1992)303

RP 61969

Determination of the novel hydroxymethyl glutaryl coenzyme A reductase inhibitor (RP 61969) and its dihydroxy acid hydrolysis product in human plasma by reversed-phase high-performance liquid chromatography 575(1992)319

Rubidium

Effect of the substituents of β -diketonato ligands on R_F values of tris(chelate) transition metal complexes obtained by normal- and reversed-phase thin-layer chromatography on unmodified silica gel 585(1991)359

Rufloxacin

Analytical procedure for the determination of rufloxacin, a new pyridobenzothiazine, in human serum and urine by high-performance liquid chromatography 564(1991)346

High-performance liquid chromatography and preliminary pharmacokinetics of rufloxacin and its metabolites, N-desmethylrufloxacin and rufloxacin sulfoxide, in urine of Rhesus monkey *Macaca mulatta* 573(1992)168

High-performance liquid chromatographic determination of rufloxacin and its main active metabolite in biological fluids 576(1992)129

Rufloxacin sulphoxide

High-performance liquid chromatography and preliminary pharmacokinetics of rufloxacin and its metabolites, N-desmethylrufloxacin and rufloxacin sulfoxide, in urine of Rhesus monkey *Macaca mulatta* 573(1992)168

Rutin

High-performance liquid chromatographic analysis of flavonol glycosides of *Solidago virgaurea* 558(1991)296

S

S-1452

Capillary gas chromatographic method for the determination of the thromboxane A_2 receptor antagonist S-1452 and its metabolites in human plasma 583(1992)45

Capillary gas chromatographic method for the determination of the thromboxane A_2 receptor antagonist S-1452 and its metabolites in human urine 583(1992)53

S100 proteins

Calcium-modulated conformational affinity chromatography. Application to the purification of calmodulin and S100 proteins 597(1992)263

Saccharides

Selective post-column fluorogenic reaction with benzamidine for trace level detection of reducing saccharides in liquid chromatography 553(1991)255

Assessment of a thermospray interface for the coupling of high-performance liquid chromatography and Fourier transform infrared spectrometry 588(1991)15

Determination of organic acids, amino acids and saccharides by high-performance liquid chromatography and a postcolumn enzyme reactor with amperometric detection 591(1992)165

New, stable polyamine-bonded polymer gel column 592(1992)93

Structure elucidation of saccharides in anthocyanins and flavonols by means of methylation analysis and gas chromatography 598(1992)81

Saccharose

Separation of functionalized dextrans by reversed-phase high-performance liquid chromatography 587(1991)288

Safrole

Comparison of four liquid crystal stationary phases used above and below their melting point temperatures for the gas chromatography of some volatile oil constituents 588(1991)231

Salbutamol

Possible role for stationary phase metal interactions in the chromatography of hydroxyamines on silica 552(1991)429

Thermospray liquid chromatography–mass spectrometry for the characterisation of sulphate ester conjugates 554(1991)175

Determination of β_2 -receptor agonists in bovine urine and liver by gas chromatography–tandem mass spectrometry 564(1991)515

Residue screening for the β -agonists clenbuterol, salbutamol and cimaterol in urine using enzyme immunoassay and high-performance liquid chromatography 564(1991)551

Evaluation of a high-performance thin-layer chromatographic technique for the determination of salbutamol serum levels in clinical trials 581(1992)306

Comparison of isotachopheresis, capillary zone electrophoresis and high-performance liquid chromatography for the determination of salbutamol, terbutaline sulphate and fenoterol hydrobromide in pharmaceutical dosage forms 590(1992)341

Salicyl acyl glucuronide

Novel direct high-performance liquid chromatographic method for determination of salicylate glucuronide conjugates in human urine 565(1991)309

Salicylamide

Simplex optimization of liquid chromatography with the function of mutual information as a criterion 585(1991)187

Salicylates

Simultaneous high-performance liquid chromatographic determination of salicylates in whole blood, plasma and isolated erythrocytes 584(1992)221

Salicylic acid

Novel direct high-performance liquid chromatographic method for determination of salicylate glucuronide conjugates in human urine 565(1991)309

Evaluation of 2-(2-thiophenecarboxy)benzoic acid and related active metabolites in biological samples 573(1992)150

Simultaneous high-performance liquid chromatographic determination of salicylates in whole blood, plasma and isolated erythrocytes 584(1992)221

Salicylideneglycinatoaluminium(III)

Retention behavior of salicylideneglycinatoaluminium(III) in reversed-phase high-performance liquid chromatography 586(1991)356

Salicyl phenolic glucuronide

Novel direct high-performance liquid chromatographic method for determination of salicylate glucuronide conjugates in human urine 565(1991)309

Salicyluric acid

Novel direct high-performance liquid chromatographic method for determination of salicylate glucuronide conjugates in human urine 565(1991)309

Simultaneous high-performance liquid chromatographic determination of salicylates in whole blood, plasma and isolated erythrocytes 584(1992)221

Salicyluric phenolic glucuronide

Novel direct high-performance liquid chromatographic method for determination of salicylate glucuronide conjugates in human urine 565(1991)309

Salsolinol

Rapid and simple procedure for the determination of salsolinol in urine using high-performance liquid chromatography with electrochemical detection 563(1991)224

Determination of free salsolinol concentrations in human urine using gas chromatography–mass spectrometry 568(1991)271

Endogenous synthesis of N-methylsalsolinol, an analogue of 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine, in rat brain during *in vivo* microdialysis with salsolinol, as demonstrated by gas chromatography–mass spectrometry 578(1992)109

Salt hydrates

Study of salt hydrates by gas–solid chromatography 557(1991)495

Samidin

Comparison of selectivity of silica and Florisil in the separation of natural pyranocoumarins 552(1991)453

Sandwicine

Monitoring of ajmaline in plasma with high-performance liquid chromatography 575(1992)87

Sanguinarine

Sanguinarine levels in biological samples by high-performance liquid chromatography 570(1991)425

Santonin

Use of high-performance liquid chromatographic peak deconvolution and peak labelling to identify antiparasitic components in plant extracts 593(1992)209

Effects of collection solvent parameters and extraction cell geometry on supercritical fluid extraction efficiencies 594(1992)297

Optimization of supercritical fluid extraction of volatile constituents from a model plant matrix 600(1992)175

Sarin

Determination of chemical warfare agents, their hydrolysis products and related compounds in soil 589(1992)287

Saxitoxin

Comparison of liquid-junction and coaxial interfaces for capillary electrophoresis-mass spectrometry with application to compounds of concern to the aquaculture industry 591(1992)325

Sciadopitysin

Identification of flavonoids from *Ginkgo biloba* L., *Anthemis nobilis* L. and *Equisetum arvense* L. by high-performance liquid chromatography with diode-array UV detection 553(1991)223

Scopolamine

Analysis of tropane alkaloids with thermospray high-performance liquid chromatography-mass spectrometry 562(1991)737

(2*S*,3*S*)-Dicyclohexyl tartrate as mobile phase additive for the determination of the enantiomeric purity of (*S*)-atropine in tablets 592(1992)339

Seldane

Measurement of azacyclonol in urine and serum of humans following terfenadine (Seldane) administration using gas chromatography-mass spectrometry 578(1992)207

Selegiline

Gas chromatographic-mass spectrometric determination of plasma selegiline using a deuterated internal standard 583(1992)254

Selenium

Gas chromatographic-mass spectrometric method for the determination of selenium in biological samples 583(1992)35

Determination of selenium by capillary gas chromatography after high-temperature derivatization with 1,2-diamino-3,5-dibromobenzene 598(1992)105

Selenocysteine

High-performance liquid chromatographic determination of selenocysteine with the fluorescent reagent, N-(iodoacetylaminoethyl)-5-naphthylamine-1-sulfonic acid 576(1992)263

Selenium choline

Synthesis of novel selenium-containing choline and acetylcholine analogues and their quantitation using a pyrolysis-gas chromatography-mass spectrometry assay 585(1991)101

Selenium ions

Ion chromatographic analysis of the purity and synthesis of sulfonium and selenonium ions 588(1991)211

Sennosides

Determination of sennoside A and B in the pharmaceutical preparation Otsuji-to using ion-pair high-performance liquid chromatography with column switching 585(1991)255

Serine hydroxymethyltransferase

Purification of serine hydroxymethyltransferase from *Bacillus stearothermophilus* with ion-exchange high-performance liquid chromatography 596(1992)203

Serine protease

Direct determination of kallikrein by high-performance liquid chromatography 593(1992)119

Serine proteinase

Affinity chromatography of serine proteinases from the bovine intervertebral disc on BPT1-Separon HEMA 1000 575(1992)229

Serotonin

Simultaneous measurement of serotonin and 5-hydroxyindoleacetic acid in rat brain using a liquid chromatographic method with electrochemical detection 564(1991)250

Improved method for the determination of serotonin in plasma by high-performance liquid chromatography using on-line sample pre-treatment 565(1991)430

Determination of γ -glutamyl conjugates of monoamines by means of high-performance liquid chromatography with electrochemical detection and application to gastropod tissues 567(1991)49

Rapid determination of 5-hydroxytryptamine in whole blood by liquid chromatography with fluorimetric detection 571(1991)231

Improved method for the determination of 5,6-dihydroxytryptamine and 5,7-dihydroxytryptamine in tissue using high-performance liquid chromatography with electrochemical detection 574(1992)137

Preparation of an optimum mobile phase for the simultaneous determination of neurochemicals in mouse brain tissues by high-performance liquid chromatography with electrochemical detection 576(1992)245

Difference in the concentration of tryptophan metabolites between maternal and umbilical foetal blood 576(1992)334

Determination of the catecholamines and serotonin, their precursors tyrosine and tryptophan, and their main metabolites in rat brain using reversed-phase high-performance liquid chromatography with fluorimetric and oxidative amperometric detection in series 578(1992)17

γ -Aminobutyric acid, catecholamine and indoleamine determinations from the same brain region by high-performance liquid chromatography with electrochemical detection 579(1992)334

Simultaneous measurement of serotonin, catecholamines and their metabolites in cat and human plasma by *in vitro* microdialysis-microbore high-performance liquid chromatography with amperometric detection 582(1992)19

Microbore liquid chromatography with dual electrochemical detection for the determination of serotonin and 5-hydroxyindoleacetic acid in rat brain dialysates 582(1992)29

Serum proteins
Rapid protein analysis by capillary electrophoresis 559(1991)445

Sesquimustard
Capillary column gas chromatography-ammonia and deuterated ammonia chemical ionization mass spectrometry of sulfur vesicants 600(1992)267

Sesquiterpene lactones
Use of high-performance liquid chromatographic peak deconvolution and peak labelling to identify antiparasitic components in plant extracts 593(1992)209

Sesquiterpenes
Structure determination of sesquiterpenes in Chinese vetiver oil by gas chromatography-tandem mass spectrometry 557(1991)451

Sevoflurane
Gas chromatographic headspace analysis of sevoflurane in blood 577(1992)289

Short-chain fatty acids

Simple and rapid method for determination of short-chain fatty acids in biological materials by high-performance liquid chromatography with ultraviolet detection 576(1992)53

Sialic acids

Determination of sialic acids in human serum by reversed-phase liquid chromatography with fluorimetric detection 579(1992)209

Sialidase

Assay of sialidase activity using ion-exchange chromatography and acidic ninhydrin reaction 581(1992)11

Sialyltransferase

Assay of sialyltransferase activity by reversed-phase ion-pair high-performance liquid chromatography 573(1992)23

Silylated derivatives

Prediction of retention indexes. III. Silylated derivatives of polar compounds 586(1991)113

Simazine

Determination of chlorotriazines in aqueous environmental samples at the ng/l level using preconcentration with a cation exchanger and on-line high-performance liquid chromatography 585(1991)67

Sinatophysin

Simultaneous purification of the neuroproteins synapsin I and synaptophysin 596(1992)51

Sinensetin

Fast separation of polymethoxylated flavones by carbon dioxide supercritical fluid chromatography 586(1991)171

Sitosterol

Sources of plant sterol contaminants encountered in low level steroid analysis 572(1991)307

Skatole

Rapid and accurate high-performance liquid chromatographic method for the determination of 3-methylindole (skatole) in faeces of various species 566(1991)101

SN-38

Simultaneous determination of the camptothecin analogue CPT-11 and its active metabolite SN-38 by high-performance liquid chromatography: application to plasma pharmacokinetic studies in cancer patients 575(1992)275

Selective and sensitive determination of lactone and hydroxy acid forms of camptothecin and two derivatives (CPT-11 and SN-38) by high-performance liquid chromatography with fluorescence detection 588(1991)165

Snake venoms

Electrophoretic analysis of snake venoms 580(1992)355

Sodium

Peak-shape analysis and noise evaluation in suppressed ion chromatography for ultra-trace ion analysis 556(1991)249

Ion-sensitive field effect transistor as a monovalent cation detector for ion chromatography and its application to the measurement of Na⁺ and K⁺ concentrations in serum 566(1991)109

Sodium cholate

Bile salt surfactants in micellar electrokinetic capillary chromatography. Application to hydrophobic molecule separations 557(1991)113

Sodium N-dodecanoyl-L-valinate

Chiral separations by micellar electrokinetic chromatography with sodium N-dodecanoyl-L-valinate 559(1991)209

Sodium dodecyl sulphate

Bile salt surfactants in micellar electrokinetic capillary chromatography. Application to hydrophobic molecule separations 557(1991)113

Sodium mercaptododecaborate

Purity assay of sodium mercaptododecaborate by high-performance liquid chromatography 595(1992)169

Sodium sulphite

Determination of succinic acid by pyrolysis-gas chromatography in the presence of sodium sulfite and iron powder 585(1991)345

Solanines

Determination of potato glycoalkaloids and their aglycone in blood serum by high-performance liquid chromatography. Application to pharmacokinetic studies in humans 573(1992)69

Solvents

Gas chromatographic separation and automatic identification of complex mixtures of organic solvents in industrial wastes 585(1991)273

Soman

Determination of chemical warfare agents, their hydrolysis products and related compounds in soil 589(1992)287

Sorbic acid

Simultaneous determination of sorbic acid, dehydroacetic acid and benzoic acid by gas chromatography-mass spectrometry 594(1992)253

Sorbitol

Simultaneous gas-liquid chromatographic determination of sugars and organic acids as trimethylsilyl derivatives in vegetables and strawberries 552(1991)337

Chromatographic separation of polyols by ligand exchange. Effects of the ion-exchange resin cross-linking and size 594(1992)125

Sotalol

Simplified procedure for the determination of sotalol in plasma by high-performance liquid chromatography 565(1991)523

High-performance liquid chromatographic determination of sotalol in plasma. I. Application to the disposition of sotalol enantiomers in humans 576(1992)321

Soybean trypsin inhibitor

Preparative anion-exchange chromatography of soybean trypsin inhibitor: the alternative of column-overload methods 584(1992)115

Sparfloxacin

Determination of sparfloxacin in serum and urine by high-performance liquid chromatography 579(1992)285

Spectinomycin

High-performance liquid chromatographic method for the determination of spectinomycin in turkey plasma 571(1991)209

Determination of aminoglycoside antibiotics by reversed-phase ion-pair high-performance liquid chromatography coupled with pulsed amperometry and ion spray mass spectrometry 591(1992)195

Spermidine

Monitoring of the uptake and metabolism of aminoxy analogues of polyamines in cultured cells by high-performance liquid chromatography 574(1992)17

Standardization in the determination of red blood cell polyamines by high-performance liquid chromatography 574(1992)134

Improvement of extraction procedure for biogenic amines in foods and their high-performance liquid chromatographic determination 591(1992)175

Spermine

Standardization in the determination of red blood cell polyamines by high-performance liquid chromatography 574(1992)134

Improvement of extraction procedure for biogenic amines in foods and their high-performance liquid chromatographic determination 591(1992)175

Sphingomyelin

Rapid isocratic method for the separation and quantification of major phospholipid classes by high-performance liquid chromatography 567(1991)29

Use of hematoporphyrin as a fluorescent stain for detection of lipids in high-performance thin-layer chromatography 588(1991)295

(+)-2,2'-Spiro[2H-chromene]

Chiroptical detection during liquid chromatography. III. Non-stop acquisition of circular dichroism spectra during liquid chromatography 586(1991)249

Spirolactams

Unusual effect of temperature on the retention of enantiomers on a chiral column 558(1991)1

Spirolactone

High-performance liquid chromatographic determination of spirinolactone and its metabolites in human biological fluids after solid-phase extraction 574(1992)57

Stanols

Identification of chlorophyll transformation products in a lake sediment by combined liquid chromatography-mass spectrometry 557(1991)271

Starches

Rapid identification of microbial starch degradation products from a complex nutrient medium by a thin-layer chromatographic method 558(1991)328

Thin-layer electrophoresis of hydroxyethyl starches on a modified silica gel support 591(1992)351

Stavudine

Determination of stavudine, a new antiretroviral agent, in human plasma by reversed-phase high-performance liquid chromatography with ultraviolet detection 584(1992)239

Stearic acid

High-performance liquid chromatographic analysis of free palmitic and stearic acids in cerebrospinal fluid 565(1991)424

Stercobilin

Separation and sensitive determination of i-urobilin and I-stercobilin by high-performance liquid chromatography with fluorimetric detection 574(1992)261

5 α -Steroid reductase

Isocratic high-performance liquid chromatographic measurement of optimal 5 α -steroid reductase activity in Hep-G2 cells 570(1991)293

Steroids

Computer-aided optimization of the experimental conditions for the isocratic reversed-phase high-performance liquid chromatographic separation of hormonal steroids 552(1991)103

Thermodesorption-gas chromatography-mass spectrometric analysis of biological materials for potential molecular precursors of the constituents of the crude oils 552(1991)325

Effects of β -cyclodextrin in the mobile phase on the retention and indirect detection of non-electrolytes in reversed-phase liquid chromatography. II. Steroids 552(1991)583

On-line derivatization of eluted substances in dynamic high-performance liquid chromatography-mass spectrometry through the particle-beam interface 554(1991)125

Effect of derivatization of steroids on their retention behaviour in inclusion chromatography using cyclodextrin as a mobile phase additive 558(1991)306

Particle beam liquid chromatography-mass spectrometry on a double-focusing sector instrument 562(1991)13

Studies on anabolic steroids. V. Sequential reduction of methandienone and structurally related steroid A-ring substituents in humans: gas chromatographic-mass spectrometric study of the corresponding urinary metabolites 562(1991)323

Mass spectrometric and high-performance liquid chromatographic studies of medroxyprogesterone acetate metabolites in human plasma 562(1991)351

Derivatization and gas chromatographic-mass spectrometric detection of anabolic steroid residues isolated from edible muscle tissues 562(1991)673

Thin-layer chromatography on silica-coated aluminium sheet as an adjunct to radioimmunoassay of steroids 563(1991)178

Simplified gas chromatographic method for the simultaneous determination of phytosterols and cholesterol 563(1991)188

Liquid chromatographic analysis of prednisolone, prednisone and their 20-reduced metabolites in perfusion media 563(1991)341

Review. Immunoaffinity chromatography, its applicability and limitations in multi-residue analysis of anabolizing and doping agents 564(1991)363

Comparison of two mass-selective detectors with special reference to the analysis of residues of anabolics 564(1991)385

Detection of 19-nortestosterone and its urinary metabolites in miniature pigs by gas chromatography-mass spectrometry 564(1991)393

Identification and characterization of 19-nortestosterone in urine of meat-producing animals 564(1991)405

Test strip enzyme immunoassays and the fast screening of nortestosterone and clenbuterol residues in urine samples at the parts per billion level 564(1991)413

Control system for detection of the illegal use of naturally occurring steroids in calves 564(1991)429

Combined high-performance liquid chromatography and radioimmunoassay for the screening of 19-nortestosterone and methyltestosterone residues in meat samples 564(1991)445

- Development of a competitive enzyme immunoassay for 17 α -19-nortestosterone 564(1991)451
- Use of an immobilized enzyme reactor for the analysis of residues of 17 α -methyltestosterone in trout by high-performance liquid chromatography 564(1991)461
- Detection of the illegal use of ethinylestradiol in cattle urine by gas chromatography-mass spectrometry 564(1991)469
- Rapid screening method for the determination of diethylstilbestrol in edible animal tissue by column liquid chromatography with electrochemical detection 564(1991)477
- Determination of zeranol and β -zearalanol in calf urine by immunoaffinity extraction and gas chromatography-mass spectrometry after repeated administration of zeranol 564(1991)493
- Determination of clenbuterol in urine as its cyclic boronate derivative by gas chromatography-mass spectrometry 564(1991)529
- High-performance liquid chromatographic determination of clenbuterol and cimaterol using post-column derivatization 564(1991)537
- Residue screening for the β -agonists clenbuterol, salbutamol and cimaterol in urine using enzyme immunoassay and high-performance liquid chromatography 564(1991)551
- Simultaneous determination of testosterone and androstadienone (sex attractant) in human plasma by gas chromatography-mass spectrometry with high-resolution selected-ion monitoring 565(1991)35
- Characterization of prednisone, prednisolone and their metabolites by gas chromatography-mass spectrometry 565(1991)45
- Determination of 4-hydroxyandrostenedione in plasma and urine by extractive alkylation and electron-capture gas chromatography 565(1991)67
- Determination of 4-hydroxyandrost-4-ene-3,17-dione metabolism in breast cancer patients using high-performance liquid chromatography-mass spectrometry 565(1991)75
- Assay methodology for prednisolone, prednisolone acetate and prednisolone sodium phosphate in rabbit aqueous humor and ocular physiological solutions 565(1991)89
- Simultaneous analysis of prednisone, prednisolone and their major hydroxylated metabolites in urine by high-performance liquid chromatography 567(1991)39
- Specific assay for unconjugated dehydroepiandrosterone in human plasma by capillary gas chromatography with electron-capture detection 567(1991)299
- Studies on steroids. CCLII. Separation and characterization of 3-oxobile acids in serum by high-performance liquid chromatography with fluorescence detection 567(1991)343
- Automated direct high-performance liquid chromatographic assay for estetrol, estriol, cortisone and cortisol in serum and amniotic fluid 568(1991)35
- Determination of serum cortisol by reversed-phase liquid chromatography using precolumn sulphuric acid-ethanol fluorescence derivatization and column switching 570(1991)1
- Direct quantitative digital autoradiography-thin-layer chromatography of 3 α ,3 β - and 5 α -reduced and 17 β -dehydrogenated androgens derived from testosterone metabolism 571(1991)73
- Simple approach to measure metabolic pathways of steroids in living cells 572(1991)25
- Sources of plant sterol contaminants encountered in low level steroid analysis 572(1991)307
- Routine screening and quantitation of urinary corticosteroids using bench-top gas chromatography-mass-selective detection 573(1992)183
- Simultaneous determination of cortisol and cortisone in human plasma by stable-isotope dilution mass spectrometry 576(1992)47
- Identification of 4-hydroxyandrost-4-ene-3,17-dione metabolites in prostatic cancer patients by liquid chromatography-mass spectrometry 576(1992)235
- Detection of methandienone (methandrostenolone) and metabolites in horse urine by gas chromatography-mass spectrometry 577(1992)195
- Thermospray liquid chromatography-mass spectrometry of corticosteroids 577(1992)221
- Stable isotope dilution mass spectrometry for the simultaneous determination of cortisol, cortisone, prednisolone and prednisone in plasma 579(1992)193
- High-performance liquid chromatographic separation and detection methods for anabolic compounds (Review) 580(1992)111
- Thin-layer chromatographic detection of zeranol and estradiol in fortified plasma and tissue extracts with Fast Corinth V 581(1992)119
- Fluorimetric detection of serum corticosterone using high-performance liquid chromatography 581(1992)267
- Urinary testosterone measurement by gas chromatography after solid-phase extraction and high-performance liquid chromatography 582(1992)7

Improved sample preparation for the testosterone hydroxylation assay using disposable extraction columns 582(1992)232

Simultaneous determination of serum cortisol and cortisone by reversed-phase liquid chromatography with ultraviolet detection 584(1992)147

General and selective isolation procedure for high-performance liquid chromatographic determination of anabolic steroids in tissues 588(1991)89

High-performance liquid chromatographic analysis of FCE 24304 (6-methylenandrosta-1,4-diene-3,17-dione) and FCE 24928 (4-aminoandrosta-1,4,6-triene-3,17-dione), two new aromatase inhibitors 593(1991)25

Validation of chromatographic retention models in reversed-phase high-performance liquid chromatography by fitting experimental data to the relevant equations 595(1992)53

Sterol esters

Analysis of sterol esters from alga and yeast by high-performance liquid chromatography and capillary gas chromatography-mass spectrometry with chemical ionization 598(1992)219

Sterol Δ^{14} -reductase

Radio-detection high-performance liquid chromatographic enzyme assay for inhibitors of fungal sterol Δ^{14} -reductase 566(1991)435

Sterols

Thermodesorption-gas chromatography-mass spectrometric analysis of biological materials for potential molecular precursors of the constituents of the crude oils 552(1991)325

Identification of chlorophyll transformation products in a lake sediment by combined liquid chromatography-mass spectrometry 557(1991)271

Sources of plant sterol contaminants encountered in low level steroid analysis 572(1991)307

Determination of sterols and diterpenoids from brown algae (Cystoseiraceae) 588(1991)99

Steryl fatty acyl esters

Deuteration as an aid to the high-temperature gas chromatography-mass spectrometry of steryl fatty acyl esters 590(1992)305

Stigmasterol

Sources of plant sterol contaminants encountered in low level steroid analysis 572(1991)307

Streptomycin

Determination of aminoglycoside antibiotics by reversed-phase ion-pair high-performance liquid chromatography coupled with pulsed amperometry and ion spray mass spectrometry 591(1992)195

Strychnine

Separation of *Strychnos nux-vomica* alkaloids by high-performance liquid chromatography 587(1991)318

Strychnine nitrate

Simultaneous determination of yohimbine hydrochloride, strychnine nitrate and methyltestosterone by ion-pair high-performance liquid chromatography 588(1991)344

Styrene

Comparison of colorimetric and chromatographic determination of *tert*.-butylpyrocatechol 595(1992)137

Styrene-butadiene block copolymer

Determination of solute-polymer interaction constants via chromatography with polymer capillaries 594(1992)29

Styrene-7,8-oxide

Determination of styrene-7,8-oxide in whole rat blood by gas chromatography-mass spectrometry 567(1991)93

Substance P

Marked improvement of a substance P radioimmunoassay by reduction of ^{125}I -labelled [Tyr⁸]-substance P prepared by the chloramine-T method with mercaptoethanol and subsequent purification by reversed-phase liquid chromatography 564(1991)67

Succinic acid

Determination of volatile and non-volatile organic acids in technical sugar solutions by ion-exclusion chromatography 555(1991)65

Determination of succinic acid by pyrolysis-gas chromatography in the presence of sodium sulfite and iron powder 585(1991)345

Succinylcholine

Rapid determination of succinylcholine in human plasma by high-performance liquid chromatography with fluorescence detection 570(1991)390

Succinylpurines

Capillary electrophoresis for diagnosis and studies of human disease, particularly metabolic disorders 559(1991)455

Sucrose

Effect of column material on sorption isotherms obtained by inverse gas chromatography 600(1992)341

Sugar-nucleotides

Improved ion-pair high-performance liquid chromatographic method for the quantification of a wide variety of nucleotides and sugar-nucleotides in animal cells 570(1991)77

Sugars

see also Carbohydrates and Saccharides

Chloride attachment negative-ion mass spectra of sugars by combined liquid chromatography and atmospheric pressure chemical ionization mass spectrometry 562(1991)81

Optimization of an ion-exchange high-performance liquid chromatographic method for the determination of carboxylic acids, sugars, glycerol and ethanol in wines 589(1992)151

Determination of the association constant of monovalent mode protein–sugar interaction by capillary zone electrophoresis 597(1992)377

Sulphadiazine

Development of a screening method for five sulfonamides in salmon muscle tissue using thin-layer chromatography 555(1991)315

Simultaneous determination of sulphadiazine and trimethoprim in plasma and tissues of cultured fish for residual and pharmacokinetic studies 583(1992)201

Determination of some drugs by micellar electrokinetic capillary chromatography. The pseudo-effective mobility as parameter for screening 585(1991)123

Sulphadimethoxine

Development of a screening method for five sulfonamides in salmon muscle tissue using thin-layer chromatography 555(1991)315

Determination of sulfonamides by liquid chromatography, ultraviolet diode array detection and ion-spray tandem mass spectrometry with application to cultured salmon flesh 558(1991)155

Sulphadimidine

High-performance liquid chromatographic method for the routine determination of sulphadimidine, its hydroxy metabolites and N4-acetylsulphadimidine in body fluids and cell culture media 571(1991)157

Determination of some drugs by micellar electrokinetic capillary chromatography. The pseudo-effective mobility as parameter for screening 585(1991)123

Sulphadoxine

Reversed-phase liquid chromatographic method for the simultaneous determination of the antimalarial drugs sulphadoxine, pyrimethamine, mefloquine and its major carboxylic metabolite in plasma 571(1991)169

Sulpha drugs

Pre-column derivatization of sulfa drugs with fluorescamine and high-performance liquid chromatographic determination at their residual levels in meat and meat products 558(1991)175

Rapid electro-dialytic clean-up of biological samples for high-performance liquid chromatography 585(1991)213

Sulphalene

Determination of sulfalene in plasma, red blood cells and whole blood by high-performance liquid chromatography 563(1991)333

Sulphamerazine

Development of a screening method for five sulfonamides in salmon muscle tissue using thin-layer chromatography 555(1991)315

Sulphamethazine

Development of a screening method for five sulfonamides in salmon muscle tissue using thin-layer chromatography 555(1991)315

Rapid electro-dialytic clean-up of biological samples for high-performance liquid chromatography 585(1991)213

Sulphapyridine

Development of a screening method for five sulfonamides in salmon muscle tissue using thin-layer chromatography 555(1991)315

Sulphate

Peak-shape analysis and noise evaluation in suppressed ion chromatography for ultra-trace ion analysis 556(1991)249

Tissue sulfate determination by ion chromatography 574(1992)146

Sulphate, inorganic

Simple method for the determination of inorganic sulfate in human serum and urine using single-column ion chromatography 565(1991)447

Sulphate ester conjugates

Thermospray liquid chromatography–mass spectrometry for the characterisation of sulphate ester conjugates 554(1991)175

Sulphobromophthalein

High-performance liquid chromatographic determination of sulphobromophthalein and its conjugates 578(1992)63

Sulphociprofloxacin

Improved high-performance liquid chromatographic determination of ciprofloxacin and its metabolites in human specimens 582(1992)263

Sulphonamides

Development of a screening method for five sulfonamides in salmon muscle tissue using thin-layer chromatography 555(1991)315

Determination of sulfonamides by liquid chromatography, ultraviolet diode array detection and ion-spray tandem mass spectrometry with application to cultured salmon flesh 558(1991)155

Determination of small drug molecules by capillary electrophoresis–atmospheric pressure ionization mass spectrometry 559(1991)515

Comparison of liquid-junction and coaxial interfaces for capillary electrophoresis–mass spectrometry with application to compounds of concern to the aquaculture industry 591(1992)325

Selection of robust combinations of extraction liquid composition and internal standard. Monte Carlo simulation of improvement of assay methods with liquid–liquid extraction prior to high-performance liquid chromatography 594(1992)45

High-performance liquid chromatography of sulphonamides extracted from bovine and porcine muscle by solid-phase dispersion 595(1992)179

Determination of sulphonamides in pork meat extracts by capillary zone electrophoresis 596(1992)101

Evaluation of restricted access media for high-performance liquid chromatographic analysis of sulfonamide antibiotic residues in bovine serum 598(1992)23

Systematic optimization of capillary electrophoretic separation of sulphonamides 598(1992)133

Sulphonic acids

Use of non-volatile ion-pairing agents for liquid chromatographic–mass spectrometric analyses with a moving-belt interface 554(1991)281

Determination of aromatic sulfonic acids in aqueous environmental samples by anion-exchange chromatography coupled to particle beam mass spectrometry and UV spectrophotometry 589(1991)177

Application of a pellicular anion-exchange resin to the separation of inorganic and organic anions by single-column anion exchange 592(1992)101

Electrodialytic sample treatment coupled on-line with column liquid chromatography for the determination of basic and acidic compounds in environmental samples 600(1992)163

Sulphonium ions

Ion chromatographic analysis of the purity and synthesis of sulfonium and selenonium ions 588(1991)211

1,1'-Sulphonylbis[2-(methylsulphinyl)ethane]

Analysis of 1,1'-sulphonylbis[2-(methylsulphinyl)ethane] and 1-methylsulphinyl-2-[2-(methylthio)ethylsulphonyl]ethane, metabolites of sulphur mustard, in urine using gas chromatography–mass spectrometry 558(1991)405

Sulphonylureas

Application of microcolumn liquid chromatography–continuous-flow fast atom bombardment mass spectrometry in environmental studies of sulphonylurea herbicides 554(1991)91

Sulphopeptides

Determination of picomole amounts of lipoxins C₄, D₄ and E₄ by high-performance liquid chromatography with electrochemical detection 585(1991)53

Sulphorhodamine B

Fluorescence detector cell for use in an integrated electrically driven separation system 598(1992)115

Sulphosuccinimidyl 2-(biotinamido)ethyl-1,3-dithiopropionate

Affinity chromatographic purification of bovine lung endothelin receptor using biotinylated endothelin and avidin–agarose 597(1992)331

Sulphur-containing compounds

Determination of sulfur components in light petroleum streams by high-resolution gas chromatography with chemiluminescence detection 589(1992)271

Comparison of commercially available atomic emission and chemiluminescence detectors for sulfur-selective gas chromatographic detection 591(1992)313

Sulphur gases

Determination of sulphur and nitrogen gases by gas chromatography on polystyrene porous polymer columns 585(1991)93

Sulphur heterocycles

Comments on the Note "Liquid chromatographic determination of planar aromatic sulphur compounds in crude oil", *J. Chromatogr.*, 475 (1989) 421 585(1991)376

Sulphur mustard

Methods for the analysis of thiodiglycol sulphoxide, a metabolite of sulphur mustard, in urine using gas chromatography–mass spectrometry 558(1991)393

Analysis of 1,1'-sulphonylbis[2-(methylsulphinyl)ethane] and 1-methylsulphinyl-2-[2-(methylthio)ethylsulphonyl]ethane, metabolites of sulphur mustard, in urine using gas chromatography–mass spectrometry 558(1991)405

Specific and sensitive quantitation of 2,2'-dichlorodiethyl sulphide (sulphur mustard) in water, plasma and blood: application to toxicokinetic study in the rat after intravenous intoxication 583(1992)155

Sulphur vesicants

Capillary column gas chromatography–ammonia and deuterated ammonia chemical ionization mass spectrometry of sulfur vesicants 600(1992)267

Sulthiame

Analysis of sulthiame in serum by narrow-bore high-performance liquid chromatography. Comparison of direct sample injection with pre-column switching and extrelut extraction 553(1991)7

Sultopride

Determination of sultopride and tiapride in serum by gas chromatography using a surface ionisation detector 567(1991)113

Sumarotene

Quantitative analysis of retinoids in biological fluids by high-performance liquid chromatography using column switching. III. Determination of the arotinoid sumarotene and its Z-isomer in human and animal plasma 576(1992)111

Sun-screen agents

Determination of sun-screen agents in cosmetic products by micellar liquid chromatography 587(1991)325

Superoxide dismutase

Resolution of three forms of superoxide dismutase by immobilised metal affinity chromatography 576(1992)340

Purification of pro- and eukaryotic superoxide dismutases by charge-controlled hydrophobic chromatography 590(1992)247

Characterization of the heterogeneity of polyethylene glycol-modified superoxide dismutase by chromatographic and electrophoretic techniques 599(1992)141

Surfactants

Use of an evaporative light scattering detector in reversed-phase high-performance liquid chromatography of oligomeric surfactants 552(1991)593

Reversed-phase high-performance liquid chromatographic determination of linear alkylbenzenesulphonates in river water at ppb levels by precolumn concentration 555(1991)155

Bile salt surfactants in micellar electrokinetic capillary chromatography. Application to hydrophobic molecule separations 557(1991)113

Quantitative aspects of indirect UV detection in capillary zone electrophoresis 588(1991)321

Reversed-phase high-performance liquid chromatography of anionic and ethoxylated non-ionic surfactants and pesticides in liquid pesticide formulations 593(1992)227

High-performance hydrophobic interaction chromatography of proteins on reversed-phase supports coated with non-ionic surfactants of polyoxyethylene type. Purification of a fungal aspartic proteinase 598(1992)181

Suxibuzone

Solid-phase extraction for the high-performance liquid chromatographic determination of indomethacin, suxibuzone, phenylbutazone and oxyphenbutazone in plasma, avoiding degradation of compounds 581(1992)101

Synapsin I

Simultaneous purification of the neuroproteins synapsin I and synaptophysin 596(1992)51

T**Tabersonine**

Determination of *Catharanthus roseus* alkaloids by high-performance liquid chromatography-isotope dilution thermospray-mass spectrometry 554(1991)227

Tabun

Mass spectrometric identification of products formed during degradation of ethyl dimethylphosphoramidocyanidate (tabun) 598(1992)89

Tangeretin

Fast separation of polymethoxylated flavones by carbon dioxide supercritical fluid chromatography 586(1991)171

Tannic acid

Determination of tannic acid and its phenolic metabolites in biological fluids by high-performance liquid chromatography 577(1992)77

Tar compounds

Gas chromatographic-mass spectrometric analysis of tar compounds formed during pyrolysis of rice husks 562(1991)531

Tartaric acid

Determination of volatile and non-volatile organic acids in technical sugar solutions by ion-exclusion chromatography 555(1991)65

Tartrate

Measurement of stability constants for complexes of divalent cations with tartaric acid by ion chromatography 589(1992)185

Taurine

Capillary electrophoresis for diagnosis and studies of human disease, particularly metabolic disorders 559(1991)455

High-performance liquid chromatographic determination of taurine in biological fluids by post-column fluorescence reaction with thiamine 568(1991)212

Simple, rapid and sensitive determination of plasma taurine by high-performance liquid chromatography using pre-column derivative formation with fluorescamine 576(1992)155

Taxol

Separation of taxol from related taxanes in *Taxus brevifolia* extracts by isocratic elution reversed-phase microcolumn high-performance liquid chromatography 587(1991)300

Determination of taxol by high-performance liquid chromatography–thermospray mass spectrometry 594(1992)153

Taxotere

Determination of Taxotere in human plasma by a semi-automated high-performance liquid chromatographic method 582(1992)273

Teboroxime

Determination of methylboronic acid in teboroxime by capillary gas chromatography 585(1991)348

Technetium complexes

Determination of the mean ionic charge of the components of three ^{99m}Tc bone scanning agents by gel chromatography with two eluents of different electrolyte composition 594(1992)217

Technetium dioxime complexes

Direct analysis of whole blood by internal surface reversed-phase chromatography: an examination of the binding and metabolism of technetium dioxime complexes 574(1992)119

Teicoplanin

Determination of teicoplanin in plasma using microbore high-performance liquid chromatography and injection-generated gradients 563(1991)451

Temafoxacin

High-performance liquid chromatographic procedures for the determination of temafoxacin in biological matrices 568(1991)197

Temazepam

Determination of diazepam and nordazepam in milk and plasma in the presence of oxazepam and temazepam 564(1991)330

Determination of temazepam and temazepam glucuronide by reversed-phase high-performance liquid chromatography 568(1991)427

Temazepam glucuronide

Determination of temazepam and temazepam glucuronide by reversed-phase high-performance liquid chromatography 568(1991)427

Teniposide

Comparative study on the determination of the anti-neoplastic drug teniposide in plasma using micellar liquid chromatography and surfactant-mediated plasma clean-up 567(1991)161

Tenuazonic acid

Reversed-phase high-performance liquid chromatography of tenuazonic acid and related tetramic acids 566(1991)195

Terbutaline

Determination of β_2 -receptor agonists in bovine urine and liver by gas chromatography–tandem mass spectrometry 564(1991)515

Analysis of terbutaline in human plasma by high-performance liquid chromatography with electrochemical detection using a micro-electrochemical flow cell 577(1992)109

Dehydrogenase-silica as a stationary phase for the separation of alcohols and ketones 587(1991)268

Terbutaline sulphate

Comparison of isotachopheresis, capillary zone electrophoresis and high-performance liquid chromatography for the determination of salbutamol, terbutaline sulphate and fenoterol hydrobromide in pharmaceutical dosage forms 590(1992)341

Terbutylazine

Determination of chlorotriazines in aqueous environmental samples at the ng/l level using preconcentration with a cation exchanger and on-line high-performance liquid chromatography 585(1991)67

Terfenadine

Determination of terfenadine and terfenadine acid metabolite in plasma using solid-phase extraction and high-performance liquid chromatography with fluorescence detection 570(1991)139

Direct enantiomeric separation of terfenadine and its major acid metabolite by high-performance liquid chromatography, and the lack of stereoselective terfenadine enantiomer biotransformation in man 571(1991)291

Measurement of azacyclonol in urine and serum of humans following terfenadine (Seldane) administration using gas chromatography–mass spectrometry 578(1992)207

Terfenadine acid metabolite

Determination of terfenadine and terfenadine acid metabolite in plasma using solid-phase extraction and high-performance liquid chromatography with fluorescence detection 570(1991)139

Terpene hydrocarbons

Comparison of four liquid crystal stationary phases used above and below their melting point temperatures for the gas chromatography of some volatile oil constituents 588(1991)231

Terpenes

Gas-liquid microcapillary columns precoated with graphitized carbon black 552(1991)197

Protection of terpenes against oxidative and acid decomposition on adsorbent cartridges 589(1992)385

Analytical methods for monoterpene glycosides in grape and wine. I. XAD-2 extraction and gas chromatographic-mass spectrometric determination of synthetic glycosides 590(1992)313

Determination of taxol by high-performance liquid chromatography-thermospray mass spectrometry 594(1992)153

Analytical methods for monoterpene glycosides in grape and wine. II. Qualitative and quantitative determination of monoterpene glycosides in grape 595(1992)269

Optimization of supercritical fluid extraction of volatile constituents from a model plant matrix 600(1992)175

Cholesteryl acetate as a stationary phase for the gas chromatography of some volatile oil constituents 600(1992)337

Tertatolol

Determination of tertatolol enantiomers in biological fluids by high-performance liquid chromatography 572(1991)203

Testosterone

On-line derivatization of eluted substances in dynamic high-performance liquid chromatography-mass spectrometry through the particle-beam interface 554(1991)125

Thin-layer chromatography on silica-coated aluminium sheet as an adjunct to radioimmunoassay of steroids 563(1991)178

Simultaneous determination of testosterone and androstadienone (sex attractant) in human plasma by gas chromatography-mass spectrometry with high-resolution selected-ion monitoring 565(1991)35

Direct quantitative digital autoradiography-thin-layer chromatography of 3 α ,3 β - and 5 α -reduced and 17 β -dehydrogenated androgens derived from testosterone metabolism 571(1991)73

Urinary testosterone measurement by gas chromatography after solid-phase extraction and high-performance liquid chromatography 582(1992)7

Improved sample preparation for the testosterone hydroxylation assay using disposable extraction columns 582(1992)232

General and selective isolation procedure for high-performance liquid chromatographic determination of anabolic steroids in tissues 588(1991)89

2,3,4,6-Tetra-O-acetyl- β -D-glucopyranosyl isothiocyanate
By-products in the derivatization of amines with the chiral reagent 2,3,4,6-tetra-O-acetyl- β -D-glucopyranosyl isothiocyanate and their elimination 592(1992)323

Tetrabutylammonium borate

Membrane-based sample preparation device for the pretreatment of acidic samples prior to cation analysis by ion chromatography 586(1991)283

Tetrabutylammonium hydroxide

Membrane-based sample preparation device for the pretreatment of acidic samples prior to cation analysis by ion chromatography 586(1991)283

Tetrabutyltin

Determination of organotin species by capillary gas chromatography with alternating current plasma emission detection 598(1992)97

Tetracaine

Determination of limiting ionic mobilities and dissociation constants of some local anaesthetics 596(1992)265

Tetrachlorothioanisole

Artifact formation during gas chromatographic-mass spectrometric analysis of a methylsulfinyl-containing metabolite 575(1992)287

Tetracyclines

Determination of tetracyclines in bovine and porcine muscle by high-performance liquid chromatography using solid-phase extraction 596(1992)211

Tetradecanedioic acid

Automated high-performance liquid chromatographic and size-exclusion chromatographic sample preparation by means of a robotic workstation 592(1992)249

Tetradecyltrimethylammonium bromide

Determination of metal ions by on-line complexation and ion-pair chromatography 590(1992)263

Tetrahydrofuran

Estimation of cellulose polarity by gas chromatography 552(1991)365

[2-(D-Tetrahydroisoquinoline carboxylic acid)]oxytocin

High-performance liquid chromatography of peptides at reduced temperatures: separation of isomers 586(1991)145

Tetrahydronaphthalene

High-temperature continuous counter-current gas-liquid chromatography 590(1992)289

(S)-(+)-Tetrahydro-5-oxo-2-furanmethyl chloromethyl ether

Some novel homochiral derivatizing agents for the gas chromatographic analysis of enantiomeric secondary alcohols 598(1992)251

(E)-1,2,3,4-Tetrahydro-1,1,4,4-tetramethyl-6-(1-methyl-2-phenylethenyl)naphthalene

Determination of the retinoid (*E*)-1,2,3,4-tetrahydro-1,1,4,4-tetramethyl-6-(1-methyl-2-phenylethenyl)naphthalene and its phenolic metabolite in human plasma by reversed-phase high-performance liquid chromatography 565(1991)375

Tetramethylarsonium

Electrospray interfacing for the coupling of ion-exchange and ion-pairing chromatography to mass spectrometry 554(1991)27

N,N,N',N'-Tetramethylethylenediamine

Optimizing separation parameters in capillary isoelectric focusing 559(1991)479

Tetramethylscutellarein

Fast separation of polymethoxylated flavones by carbon dioxide supercritical fluid chromatography 586(1991)171

Tetramic acids

Reversed-phase high-performance liquid chromatography of tenuazonic acid and related tetramic acids 566(1991)195

Tetrols

Gas chromatographic-mass spectrometric analysis of diols and tetrols from reactions of polycyclic aromatic hydrocarbon epoxides with hemoglobin 562(1991)563

Thalidomide

Determination of thalidomide in plasma and blood by high-performance liquid chromatography: avoiding hydrolytic degradation 582(1992)211

Theobromine

High-performance liquid chromatographic separation of caffeine, theophylline, theobromine and paraxanthine in rat brain and serum 570(1991)185

Simultaneous determination of caffeine and its primary demethylated metabolites in human plasma by high-performance liquid chromatography 575(1992)311

Theophylline

Effect of stationary phase structure on retention and selectivity of restricted-access reversed-phase packing materials 558(1991)19

Determination of theophylline in serum and saliva in the presence of caffeine and its metabolites 568(1991)177

High-performance liquid chromatographic separation of caffeine, theophylline, theobromine and paraxanthine in rat brain and serum 570(1991)185

Simultaneous determination of caffeine and its primary demethylated metabolites in human plasma by high-performance liquid chromatography 575(1992)311

Separation of theophylline and its analogues by micellar electrokinetic chromatography: application to the determination of theophylline in human plasma 577(1992)135

Determination of substituted purines in body fluids by micellar electrokinetic capillary chromatography with direct sample injection 593(1992)275

Thermolysin

Application of high-performance capillary electrophoresis to the analysis of conformation and interaction of metal-binding proteins 559(1991)345

Thiamine

Comparative studies on the high-performance liquid chromatographic determination of thiamine and its phosphate esters with chloroethylthiamine as an internal standard using pre- and post-column derivatization procedures 558(1991)115

Concomitant determination of thiamine and its phosphate esters in human blood and serum by high-performance liquid chromatography 564(1991)127

Application of high-performance liquid chromatography to the study of thiamine metabolism and in particular thiamine triphosphatase 566(1991)397

Determination of thiamine and its phosphate esters by gradient-elution high-performance liquid chromatography 567(1991)71

Thiamine phosphate esters

Comparative studies on the high-performance liquid chromatographic determination of thiamine and its phosphate esters with chloroethylthiamine as an internal standard using pre- and post-column derivatization procedures 558(1991)115

Concomitant determination of thiamine and its phosphate esters in human blood and serum by high-performance liquid chromatography 564(1991)127

Determination of thiamine and its phosphate esters by gradient-elution high-performance liquid chromatography 567(1991)71

Thiamphenicol

Determination of thiamphenicol residues in chicken muscles by column liquid chromatography 565(1991)471

Thiamylal

Determination of thiopental in human serum and plasma by high-performance capillary electrophoresis-micellar electrokinetic chromatography 559(1991)505

Thiazole-4-carboxamide-adenine dinucleotide

Determination of nicotinamide-adenine dinucleotide and thiazole-4-carboxamide-adenine dinucleotide in human leukocytes by reversed-phase high-performance liquid chromatography 577(1992)37

Thiazolidine-4-carboxylic acids

High-performance liquid chromatography of thiazolidinic compounds obtained by condensation of pyridoxal 5'-phosphate or pyridoxal with aminothiols (L- or D-cysteine, cysteamine, L-cysteine ethyl ester) 586(1991)153

1,3-Thiazolidines

Endogenous alkaloids in man. XI. Analysis of glyoxylate-derived 1,3-thiazolidines by ion-pair-assisted reversed-phase chromatography 595(1992)351

2-(2-Thienylazo)-5-diethylaminophenol chelates

Separation and determination of trace amounts of vanadium (V), chromium(III) and iron(III) with 2-(2-thienylazo)-5-diethylaminophenol chelates by high-performance liquid chromatography 598(1992)298

Thimerosal

Assay for thiomersal (thimerosal) with adaptation to the quantitation of total ethylmercury available in degraded samples 587(1991)329

Thiobarbituric acid

High-performance liquid chromatographic separation of malondialdehyde-thiobarbituric acid adduct in biological materials (plasma and human cells) using a commercially available reagent 577(1992)9

Thiodiglycol sulphoxide

Methods for the analysis of thiodiglycol sulphoxide, a metabolite of sulphur mustard, in urine using gas chromatography-mass spectrometry 558(1991)393

6-Thioguanine

Reversed-phase high-performance liquid chromatographic approach to determine total lymphocyte concentrations of 6-thioguanine, methylmercaptapurine and methylthioguanine in humans 571(1991)149

6-Thioguanine nucleotide

High-performance liquid chromatographic assay of the methyl and nucleotide metabolites of 6-mercaptopurine: quantitation of red blood cell 6-thioguanine nucleotide, 6-thioinosinic acid and 6-methylmercaptapurine metabolites in a single sample 583(1992)83

6-Thioinosinic acid

High-performance liquid chromatographic assay of the methyl and nucleotide metabolites of 6-mercaptopurine: quantitation of red blood cell 6-thioguanine nucleotide, 6-thioinosinic acid and 6-methylmercaptapurine metabolites in a single sample 583(1992)83

Thiols

Time-resolved luminescence detection of derivatized thiols in column liquid chromatography 552(1991)625

Micro liquid chromatography with fluorescence detection of thiols and disulphides 553(1991)433

Determination of volatile thiols by gas chromatography using separation as tributyltin mercaptides 555(1991)306

Capillary electrophoresis for diagnosis and studies of human disease, particularly metabolic disorders 559(1991)455

Thiopental

Determination of thiopental in human serum and plasma by high-performance capillary electrophoresis-micellar electrokinetic chromatography 559(1991)505

2-(2-Thiophenecarboxy)benzoic acid

Evaluation of 2-(2-thiophenecarboxy)benzoic acid and related active metabolites in biological samples 573(1992)150

2-Thiophenecarboxylic acid

Evaluation of 2-(2-thiophenecarboxy)benzoic acid and related active metabolites in biological samples 573(1992)150

Thiophenes

Quantitative structure-type analysis of heterocompound-hydrocarbon mixtures. VI. Characterization of the nature of neutral heterocompounds in a product of brown coal refining 553(1991)441

Determination of sulfur components in light petroleum streams by high-resolution gas chromatography with chemiluminescence detection 589(1992)271

Thiophosphoryl compounds

Application of the iodine-azide reagent for selective detection of thiophosphoryl compounds in thin-layer chromatography 588(1991)307

Thioredoxins

Separation of the complete thioredoxin pattern of soybean leaves (*Glycine max*) by high-performance anion-exchange chromatography on Mono Q 587(1991)109

Thioridazine

Selected applications of cyclodextrin selectors in capillary electrophoresis 559(1991)215

Thiosalicylic acid

Simultaneous analysis of flunixin, naproxen, ethacrynic acid, indomethacin, phenylbutazone, mefenamic acid and thiosalicylic acid in plasma and urine by high-performance liquid chromatography and gas chromatography–mass spectrometry 568(1991)351

Thiosugars

High-performance liquid chromatographic detection of enantiomeric amino alcohols after derivatization with *o*-phthalaldehyde and various thiosugars 558(1991)311

Thiosulphate

Mixed potential response at metallic electrodes in potentiometric detection for ion chromatography 589(1992)201

Threonine

Coupled column chromatography in chiral separations: systems employing β -cyclodextrin phases for chiral separation 557(1991)199

High-performance ligand-exchange chromatography of some amino acids containing two chiral centres 592(1992)345

Thrombin

Study of high-affinity interactions by quantitative affinity chromatography. Analytical expressions in terms of total ligand concentration 597(1992)57

Thromboxane A₂ receptor antagonist

Capillary gas chromatographic method for the determination of the thromboxane A₂ receptor antagonist S-1452 and its metabolites in human plasma 583(1992)45

Capillary gas chromatographic method for the determination of the thromboxane A₂ receptor antagonist S-1452 and its metabolites in human urine 583(1992)53

Thromboxanes

Cyclooxygenase and lipoxygenase arachidonic acid metabolism by monocytes from human immune deficiency virus-infected drug users 557(1991)507

Analysis of some eicosanoids by continuous-flow fast atom bombardment mass spectrometry 562(1991)59

Characteristic fragmentation of thromboxane B₂ in thermospray high-performance liquid chromatography–mass spectrometry 562(1991)153

Determination of 6-keto-PGF_{1 α} , 2,3-dinor-6-keto-PGF_{1 α} , thromboxane B₂, 2,3-dinor-thromboxane B₂, PGE₂, PGD₂ and PGF_{2 α} in human urine by gas chromatography–negative ion chemical ionization mass spectrometry 562(1991)599

High-performance liquid chromatography–thermospray mass spectrometry of prostaglandin and thromboxane acetyl derivatives 568(1991)11

Gas chromatographic–mass spectrometric determination of 11-dehydrothromboxane B₂ in human urine 577(1992)1

Thymidine

Reversed-phase high-performance liquid chromatography–thermospray mass spectrometry of radiation-induced decomposition products of thymine and thymidine 593(1992)133

Thymine

Reversed-phase high-performance liquid chromatography–thermospray mass spectrometry of radiation-induced decomposition products of thymine and thymidine 593(1992)133

Thymol

Comparison of four liquid crystal stationary phases used above and below their melting point temperatures for the gas chromatography of some volatile oil constituents 588(1991)231

Thyroglobulin

Determination of proteolytic hydrolysis of thyroglobulin 566(1991)471

Thyronine

Comparison of high-performance liquid chromatographic detection methods for thyronine and tyrosine residues in toxicological studies of the thyroid 553(1991)135

Thyroxine

Characterization of thyroxine–albumin binding using high-performance affinity chromatography. I. Interactions at the warfarin and indole sites of albumin 579(1992)225

Thyroxine glucuronide

Determination of ¹²⁵I-labelled thyroxine glucuronide by reversed-phase high-performance liquid chromatography using on-line radiochemical detection to determine UDPglucuronosyltransferase activity 566(1991)415

Tiagabine

High-performance liquid chromatographic procedure for the determination of tiagabine concentrations in human plasma using electrochemical detection 574(1992)313

Tiapride

Determination of sultopride and tiapride in serum by gas chromatography using a surface ionisation detector 567(1991)113

Tick anticoagulant peptide

Large-scale purification and characterization of recombinant tick anticoagulant peptide 574(1992)225

Timolol

Assay of timolol in human plasma using gas chromatography with electron-capture detection 581(1992)277

Tin

Comparison of sulphur-mode and tin-mode flame photometric detectors for the gas chromatographic determination of organotin compounds 566(1991)207

Tinuvin 770

Determination of polymeric hindered amine light stabilizers in polyolefins by high-performance liquid chromatography 553(1991)415

Ionization mechanisms in capillary supercritical fluid chromatography—chemical ionization mass spectrometry 554(1991)351

Tissue plasminogen activator

Fractionation of the human recombinant tissue plasminogen activator (rtPA) glycoforms by high-performance capillary zone electrophoresis and capillary isoelectric focusing 559(1991)401

TNT-1 F(ab')₂ fragment

Chromatographic methods for large-scale preparation of immunoglobulin G_{2a} monoclonal antibodies Lym-1 and TNT-1 F(ab')₂ fragments 563(1991)243

Tobramycin

High-performance liquid chromatography method for the determination of aminoglycosides based on automated pre-column derivatization with *o*-phthalaldehyde 593(1992)69

Tocainide

Stereospecific high-performance liquid chromatographic determination of tocainide 566(1991)155

Tocopherols

Thermodesorption—gas chromatography—mass spectrometric analysis of biological materials for potential molecular precursors of the constituents of the crude oils 552(1991)325

Combination of supercritical fluid chromatography with thin-layer chromatography on a semi-preparative scale 552(1991)539

 α -Tocopheryl nicotinate

High-performance liquid chromatographic determination of α -tocopheryl nicotinate in cosmetic preparations 593(1992)95

Tolfenamic acid metabolite

Identification of a tolfenamic acid metabolite in the horse by gas chromatography—mass spectrometry 573(1992)136

Tolmetin

High-performance liquid chromatographic evaluation of Med 15 and its metabolites Med 5 and tolmetin in rat plasma 553(1991)81

Tolperisone

Retention and enantioselectivity of racemic solutes on a modified ovomucoid-bonded column. I. Cross-linking with glutaraldehyde 592(1992)301

Toluene

Gas chromatographic separation and automatic identification of complex mixtures of organic solvents in industrial wastes 585(1991)273

High-performance liquid chromatographic column-switching technique for the determination of intermediates of anaerobic degradation of toluene in ground water microcosm 587(1991)185

Toluene-2,4-diisocyanate

Automated high-performance liquid chromatographic and size-exclusion chromatographic sample preparation by means of a robotic workstation 592(1992)249

Topanol antioxidants

Determination of topanol antioxidants in methacrylates using capillary gas—liquid chromatography 596(1992)290

Topoisomerase II

Use of anion-exchange chromatography and chromatofocusing to reveal the structural and functional heterogeneity of topoisomerase II in a HL-60 cell line resistant to multi-drug treatment 587(1991)3

Toremifene

High-performance liquid chromatographic method for the determination of toremifene and its major human metabolites 565(1991)482

Toxicants

Chromatographic methods for the determination of toxicants in faeces (Review) 580(1992)135

Strategies for the identification of non-polar toxicants in aqueous environmental samples using toxicity-based fractionation and gas chromatography—mass spectrometry (Review) 580(1992)215

Toxic substances

Gas chromatographic—mass spectrometric analysis of some potential toxicants amongst volatile compounds emitted during large-scale thermal degradation of poly(acrylonitrile—butadiene—styrene) plastic 562(1991)681

Toxins

Methods for chromatographic determination of amanitins and related toxins in biological samples (Review) 580(1992)279

Transcription factors

Isolation of multiple protein factors involved in ribosomal DNA transcription 587(1991)25

Transferrins

High-performance liquid chromatographic separation of human apotransferrin and monoferric and diferric transferrins 567(1991)248

Single-step method for purification of human transferrin from a by-product of chromatographic fractionation of plasma 576(1992)71

Ligand-receptor interactions in affinity cell partitioning. Studies with transferrin covalently linked to monomethoxypoly (ethylene glycol) and rat reticulocytes 594(1992)97

Transglutamases

Purification and characterization of transglutaminases from the genital tract of the male rat 587(1991)55

Transition metal complexes

Effect of the substituents of β -diketonato ligands on R_f values of tris(chelate) transition metal complexes obtained by normal- and reversed-phase thin-layer chromatography on unmodified silica gel 585(1991)359

Transition metals

Reversed-phase separation of transition metals, lanthanides and actinides by elution with mandelic acid 558(1991)197

Factors affecting the separation of inorganic metal cations by capillary electrophoresis 593(1992)289

Role of metal ions in the ligand-exchange separation of amino acids 596(1992)281

Transplatin

Separation of some platinum(II) complexes by ionic strength gradient on a solvent-generated ion-exchange sorbent 586(1991)291

Transthyretin

Assay of human transthyretin-bound holo-retinol-binding protein with reversed-phase high-performance liquid chromatography 567(1991)369

Dye-affinity purification of transthyretin from an unexploited by-product of human plasma chromatographic fractionation 576(1992)87

Purification of biologically active human plasma transthyretin by dye-affinity chromatography: studies on dye leakage and possibility of heat treatment for virus inactivation 584(1992)93

Tranlycypromine

Enantiospecific high-performance liquid chromatographic assay with fluorescence detection for the monoamine oxidase inhibitor tranlycypromine and its applicability in pharmacokinetic studies 584(1992)229

Trazodone

Measurement of trazodone in plasma and brain of rat by capillary gas chromatography with a nitrogen-selective detector 575(1992)301

17 β -Trenbolone

Disposition of 17 β -trenbolone in humans 564(1991)485

Trenbolone

General and selective isolation procedure for high-performance liquid chromatographic determination of anabolic steroids in tissues 588(1991)89

Triacylglycerol hydroperoxides

High-performance liquid chromatography and post-column derivatization with diphenyl-1-pyrenylphosphine for fluorimetric determination of triacylglycerol hydroperoxides 596(1992)197

Triacylglycerols

Applications of capillary gas chromatography to the quality control of butter and related products 552(1991)273

Isolation and quantification of polymers from autoxidized fish oils by high-performance size-exclusion chromatography with an evaporative mass detector 552(1991)501

Automated separation and quantitation of lipid fractions by high-performance liquid chromatography and mass detection 567(1991)21

Analysis of human blood plasma triacylglycerols using capillary gas chromatography, silver ion thin-layer chromatographic fractionation and desorption chemical ionization mass spectrometry 568(1991)1

Determination of phospholipid fatty acid and triacylglycerol composition of rat caecal mucosa 575(1992)1

Determination of fatty acid and triacylglycerol composition of human adipose tissue 581(1992)171

Reversed-phase liquid chromatography-mass spectrometry of complex mixtures of natural triacylglycerols with chloride-attachment negative chemical ionization 588(1991)73

Resolution of lipolysis mixtures from soybean oil by a solid-phase extraction procedure 589(1992)353

Permanganate-impregnated packed capillary columns for group separation of triacylglycerols using supercritical media as mobile phases 600(1992)358

Triallate

Analysis of triallate residues in cereals and soil by gas chromatography with ion-trap detection 562(1991)525

Triallyl isocyanurate

Immobilization method for polyethylene glycol using a cross-linking co-agent 558(1991)323

Triamterene

Simultaneous optimization of several chromatographic performance goals using Derringer's desirability function 586(1991)11

Triazines

Determination of triazines and organophosphorus pesticides in water samples using solid-phase extraction 555(1991)137

Chromatographic separation and molecular modelling of triazines with respect to their inhibition of the growth of L1210/R71 cells 592(1992)59

Triazolam

Sensitive assay for triazolam in plasma following low oral doses 577(1992)190

Triazolines

Triazolines. XXIII. High-performance liquid chromatographic assay in rat blood for a novel triazoline anticonvulsant (ADD17014) 563(1991)419

Tributyltin chloride

Determination of organotin species by capillary gas chromatography with alternating current plasma emission detection 598(1992)97

Tributyltin mercaptides

Determination of volatile thiols by gas chromatography using separation as tributyltin mercaptides 555(1991)306

Tricarboxylic acid cycle metabolites

Simple cation-exchange high-performance liquid chromatography optimized to the measurement of metabolites in the effluents from perfused rat livers using refractive index and ultraviolet detectors 563(1991)231

Trichloroacetic acid

Monitoring the effluents of the trichloroacetic acid process by high-performance liquid chromatography 600(1992)316

3,5,6-Trichloropyridinol

Analysis of 3,5,6-trichloropyridinol in human urine using negative-ion chemical ionization gas chromatography-mass spectrometry 575(1992)69

Trichomycin

Identification of aromatic moieties and mycosamine in antifungal heptaenes with high-performance liquid chromatography, high-performance liquid chromatography-mass spectrometry and gas chromatography-mass spectrometry 585(1991)247

High-performance liquid chromatography of heptaene polyenes: assay of heptaene produced by *Streptomyces griseoviridis* 588(1991)356

Trichothecenes

Retention data for five ketotrichothecenes in reversed-phase high-performance liquid chromatography with different eluent systems 558(1991)81

Identification of trichothecenes by thermospray, plasmaspray and dynamic fast-atom bombardment liquid chromatography-mass spectrometry 562(1991)555

Fluorescence detection of trichothecene mycotoxins as coumarin-3-carbonyl chloride derivatives by high-performance liquid chromatography 595(1992)143

Triclabendazole

Reversed-phase ion-pair high-performance liquid chromatographic determination of triclabendazole metabolites in serum and urine 576(1992)135

Trienantine

Applications of capillary gas chromatography to the quality control of butter and related products 552(1991)273

Triethylamine

Liquid membrane work-up of blood plasma samples applied to gas chromatographic determination of aliphatic amines 573(1992)191

Trifluoroacetic acid

Determination of parts per million levels of trifluoroacetic acid in ceronapril bulk substance by headspace capillary gas chromatography 596(1992)85

3,3,3-Trifluoroalanine

High-performance ligand-exchange liquid chromatography of fluoro derivatives of alanine 600(1992)83

2,2,2-Trifluoro-(9-anthryl)ethanol

Problems and solutions in chiral thin-layer chromatography: a two-phase "Pirkle" modified amino-bonded plate 553(1991)497

Trifluorothymidine

Determination of trifluorothymidine in the eye using high-performance liquid chromatography 568(1991)467

Trifluralin

High-performance liquid chromatographic determination of dinitroaniline herbicides in soil and water 585(1991)164

Triglycerides

see Triacylglycerols

Trihexyphenidyl

Quantitation of trihexyphenidyl from plasma using a mass-selective detector and electron-impact ionization 571(1991)250

Trihydroxy acids

Use of methyl oxime derivatives to enhance structural information in thermospray high-performance liquid chromatography-mass spectrometry. Analysis of linoleic acid lipoxigenase metabolites in maize embryos 554(1991)155

Trilinolein

High-performance liquid chromatography and post-column derivatization with diphenyl-1-pyrenylphosphine for fluorimetric determination of triacylglycerol hydroperoxides 596(1992)197

Trilostane

High-performance liquid chromatographic analysis of trilostane and ketotrilostane in rat plasma 567(1991)282

Trimecaine

Determination of limiting ionic mobilities and dissociation constants of some local anaesthetics 596(1992)265

Trimeprazine

Influence of chemical structure of tricyclic tertiary dimethylamines on chiral separation by reversed-phase high-performance liquid chromatography after derivatization with (–)-menthylchloroformate 553(1991)365

Identification of new urinary metabolites of trimeprazine in rats by gas chromatography–mass spectrometry 575(1992)295

Trimethadione

Improved method for the determination of trimethadione and its demethylated metabolite, dimethadione, in human serum by gas chromatography 584(1992)267

Trimethoprim

Determination of trimethoprim and its oxidative metabolites in cell culture media and microsomal incubation mixtures by high-performance liquid chromatography 579(1992)355

Simultaneous determination of sulphadiazine and trimethoprim in plasma and tissues of cultured fish for residual and pharmacokinetic studies 583(1992)201

Comparison of liquid-junction and coaxial interfaces for capillary electrophoresis–mass spectrometry with application to compounds of concern to the aquaculture industry 591(1992)325

Trimethoprim N-oxides

Determination of trimethoprim and its oxidative metabolites in cell culture media and microsomal incubation mixtures by high-performance liquid chromatography 579(1992)355

Trimethylamine

Determination of trimethylamine and related aliphatic amines in human urine by head-space gas chromatography 584(1992)141

Trimethylamine N-oxide

Determination of trimethylamine and related aliphatic amines in human urine by head-space gas chromatography 584(1992)141

Trimethylanilinium hydroxide

Intra-injector formation of methyl esters from phenoxy acid pesticides 598(1992)309

Trimethylbenzenes

High-performance liquid chromatographic determination of 3,5-dimethylhippuric acid in the occupational exposure to trimethylbenzenes 577(1992)180

Trimethylfluorosilane

Determination of trace amounts of fluoride in raw materials for pharmaceuticals by gas–liquid chromatography 585(1991)289

Trimipramine

Influence of chemical structure of tricyclic tertiary dimethylamines on chiral separation by reversed-phase high-performance liquid chromatography after derivatization with (–)-menthylchloroformate 553(1991)365

Simultaneous determination of trimipramine and its major metabolites by high-performance liquid chromatography 566(1991)228

Triolein

Use of hematoporphyrin as a fluorescent stain for detection of lipids in high-performance thin-layer chromatography 588(1991)295

Tripeptides

Electrophoretic mobility and dissociation constants of tripeptides evaluated by isotachopheresis 585(1991)364

Triphenylphosphine oxide

Experimental study on the effect of the sample size on the band profile for a binary mixture showing no competitive interaction 556(1991)195

Tris(alkylxanthato)cobalt(III) complexes

Thin-layer chromatography on polyacrylonitrile. IV. Investigation of the separation mechanisms for tris-(alkylxanthato)cobalt(III) complexes 596(1992)91

Tris(aminocarboxylato)cobalt(III) complexes

Simultaneous effect of geometrical isomerism and chelate ring size of tris(aminocarboxylato)cobalt(III) complexes on their behaviour in thin-layer chromatography on silica gel and alumina 585(1991)181

Tris(hydroxymethyl)aminomethane

High-performance liquid chromatographic method for the determination of tris(hydroxymethyl)aminomethane (tromethamine) in human plasma 583(1992)99

Triterpene glucuronides

High-performance liquid chromatography–atmospheric pressure ionization mass spectrometry of gymnemic acids 557(1991)383

Triterpenoids

Thin-layer chromatographic separations of lantadenes, the pentacyclic triterpenoids from lantana (*Lantana camara*) plant 587(1991)351

Tritium

Gas chromatography for measurement of hydrogen isotopes at tritium processing 586(1991)131

Triton X

Ion-mobility spectrometry as a detection method for packed-column supercritical fluid chromatography 552(1991)551

Tromethamine

High-performance liquid chromatographic method for the determination of tris(hydroxymethyl)aminomethane (tromethamine) in human plasma 583(1992)99

Tropacocaine

Determination of limiting ionic mobilities and dissociation constants of some local anaesthetics 596(1992)265

Tropane alkaloids

Analysis of tropane alkaloids with thermospray high-performance liquid chromatography-mass spectrometry 562(1991)737

Tropomyosins

Chromatographic analysis of tropomyosins from rabbit skeletal, chicken gizzard and earthworm muscle 599(1992)51

Trypsin

Automated high-performance liquid chromatographic assay of enzymatic activities 566(1991)327

Trypsin and affinity chromatography (Review) 597(1992)3

Tryptamine

Improvement of extraction procedure for biogenic amines in foods and their high-performance liquid chromatographic determination 591(1992)175

Tryptophan

Rapid and sensitive high-performance liquid chromatographic method for the analysis of tryptophan, tyrosine and phenylalanine in biological samples 553(1991)149

Quantitation of tryptophan and other plasma amino acids by automated pre-column *o*-phthaldialdehyde derivatization high-performance liquid chromatography: improved sample preparation 564(1991)81

Determination of the catecholamines and serotonin, their precursors tyrosine and tryptophan, and their main metabolites in rat brain using reversed-phase high-performance liquid chromatography with fluorimetric and oxidative amperometric detection in series 578(1992)17

High-sensitivity laser-induced fluorescence detection of native proteins in capillary electrophoresis 595(1992)319

Rapid determination of free tryptophan in plant samples by gas chromatography-selected ion monitoring mass spectrometry 596(1992)294

T-2 toxin

Identification of trichothecenes by thermospray, plasmaspray and dynamic fast-atom bombardment liquid chromatography-mass spectrometry 562(1991)555

Tulobuterol

Determination of β_2 -receptor agonists in bovine urine and liver by gas chromatography-tandem mass spectrometry 564(1991)515

Tumor necrosis factor α

Immunomagnetic separation of tumor necrosis factor α . I. Batch procedure for human temporomandibular fluid 583(1992)11

Immunomagnetic separation of tumor necrosis factor α . II. *In situ* procedure for the human gingival space 583(1992)19

TVX Q 5402

Determination of benperidol and its reduced metabolite in human plasma by high-performance liquid chromatography and electrochemical detection 565(1991)363

Tylosin tartrate

Separation of antibiotics by high-performance capillary electrophoresis with photodiode-array detection 585(1991)133

Tyramine

Improvement of extraction procedure for biogenic amines in foods and their high-performance liquid chromatographic determination 591(1992)175

Tyrosine

Comparison of high-performance liquid chromatographic detection methods for thyronine and tyrosine residues in toxicological studies of the thyroid 553(1991)135

Rapid and sensitive high-performance liquid chromatographic method for the analysis of tryptophan, tyrosine and phenylalanine in biological samples 553(1991)149

Analysis of tyrosine- and methionine-containing neuropeptides by fast atom bombardment mass spectrometry 562(1991)459

Determination of the catecholamines and serotonin, their precursors tyrosine and tryptophan, and their main metabolites in rat brain using reversed-phase high-performance liquid chromatography with fluorimetric and oxidative amperometric detection in series 578(1992)17

Tyrosine ammonia-lyase

Identification by high-performance liquid chromatography of tyrosine ammonia-lyase activity in purified fractions of *Phaseolus vulgaris* phenylalanine ammonia-lyase 573(1992)309

Tyrosine protein kinase

Use of hydrophilic interaction chromatography for the study of tyrosine protein kinase specificity 583(1992)137

[Tyr⁸]-substance P

Marked improvement of a substance P radioimmunoassay by reduction of ¹²⁵I-labelled [Tyr⁸]-substance P prepared by the chloramine-T method with mercaptoethanol and subsequent purification by reversed-phase liquid chromatography 564(1991)67

U

Ubiquinone

Improved high-performance liquid chromatographic method for the determination of coenzyme Q₁₀ in plasma 593(1992)217

UK-68,798

Liquid-liquid extraction and high-performance liquid chromatography for the determination of a novel antidysrhythmic agent (UK-68,798) in human urine 568(1991)475

Umbelliferone

Automated high-performance liquid chromatographic assay for the determination of 7-ethoxycoumarin and umbelliferone 578(1992)141

Unsaturated esters

Gas chromatographic separation of unsaturated polar compounds on inorganic stationary phases containing silver nitrate 552(1991)313

Unsaturated hydrocarbons

Gas chromatographic separation of unsaturated polar compounds on inorganic stationary phases containing silver nitrate 552(1991)313

Uracil

Automated determination of orotic acid, uracil and pseudouridine in urine by high-performance liquid chromatography with column switching 568(1991)325

Enzymatic shot-gun 5'-phosphorylation and 3'-sister phosphate exchange: a two-dimensional thin-layer chromatographic technique to measure DNA deoxynucleotide modification 574(1992)41

Urease

Automated screening of urine samples for carbohydrates, organic and amino acids after treatment with urease 562(1991)125

Uric acid

Simultaneous determination of allantoin and oxypurines in biological fluids by high-performance liquid chromatography 575(1992)153

Uridine 5'-diphosphateglucuronosyltransferase

Determination of ¹²⁵I-labelled thyroxine glucuronide by reversed-phase high-performance liquid chromatography using on-line radiochemical detection to determine UDPglucuronosyltransferase activity 566(1991)415

Urine protein 1

Simple and high-yield purification of urine protein 1 using immunoaffinity chromatography: evidence for the identity of urine protein 1 and human Clara cell 10-kilodalton protein 577(1992)25

Urobilin

Separation and sensitive determination of i-urobilin and 1-stercobilin by high-performance liquid chromatography with fluorimetric detection 574(1992)261

Urocanic acid

Simultaneous determination of stable isotopically labelled L-histidine and urocanic acid in human plasma by stable isotope dilution mass spectrometry 576(1992)213

Uroporphyrinogen III

Random decarboxylation of uroporphyrinogen III by human hepatic uroporphyrinogen decarboxylase 566(1991)409

Usnic acid

High-performance liquid chromatographic determination of usnic acid in plasma 575(1992)167

V

Val-Ser-Gln-Asn-Tyr-Pro-Ile-Val-Gln

Microbore liquid chromatography coupled to a flow fast atom bombardment probe for the on-line detection of the Tyr-Pro cleavage of a nonapeptide by recombinant HIV-1 protease 562(1991)67

Valproic acid

Determination of valproic acid by high-performance liquid chromatography with photodiode-array and fluorescence detection 576(1992)163

Vanadium

Separation and determination of trace amounts of vanadium (V), chromium(III) and iron(III) with 2-(2-thienylazo)-5-diethylaminophenol chelates by high-performance liquid chromatography 598(1992)298

Vanillic acid

Determination of urinary vanillic acid and plasma dihydroxyphenylalanine as markers of non-secreting neuroblastoma by high-performance liquid chromatography with electrochemical detection 567(1991)331

Vanillic acid

Effect of the addition of viscous matrices to the mobile phase on chromatographic performance in liquid chromatography–fast atom bombardment mass spectrometry 554(1991)61

Simple direct liquid introduction system usable as an interface for liquid chromatography–mass spectrometry on quadrupole and magnetic-sector mass spectrometers 554(1991)293

Vanillylmandelic acid

Determination of vanilmandelic acid in urine by coupled-column liquid chromatography combining affinity to boronate and separation by anion exchange 567(1991)1

New analyser for the determination of urinary vanillylmandelic acid, homovanillic acid and creatinine 567(1991)11

Simple and fast chromatographic method for the simultaneous determination of vanillylmandelic acid and homovanillic acid in human urine 570(1991)371

Vanoxerine

Column liquid chromatographic assay of the dopamine reuptake inhibitor vanoxerine (GBR 12909) in human serum 571(1991)305

Vasopressin

Solid-phase extraction of plasma vasopressin: evaluation, validation and application 565(1991)159

Interactions and uses of antisense peptides in affinity technology (Review) 597(1992)29

Vecuronium

High-performance liquid chromatography–electrochemical detection of vecuronium and its metabolites in human plasma 573(1992)79

Vecuronium bromide

Liquid chromatography–mass spectrometry with ionspray and electrospray interfaces in pharmaceutical and biomedical research 554(1991)39

Verapamil

Plasma level monitoring of D,L-verapamil and three of its metabolites by reversed-phase high-performance liquid chromatography 570(1991)229

Simultaneous direct determination of the enantiomers of verapamil and norverapamil in plasma using a derivatized amylose high-performance liquid chromatographic chiral stationary phase 574(1992)85

Direct determination of the enantiomeric ratio of verapamil, its major metabolite norverapamil and gallopamil in plasma by chiral high-performance liquid chromatography 575(1992)255

Veronal

Quantitative aspects of indirect UV detection in capillary zone electrophoresis 588(1991)321

Very-long-chain fatty acids

Determination of very-long-chain fatty acids in plasma by a simplified gas chromatographic–mass spectrometric procedure 562(1991)147

Very-long-chain free fatty acids

Separation and determination of saturated very-long-chain free fatty acids in plasma of patients with adrenoleukodystrophy using solid-phase extraction and high-performance liquid chromatographic analysis 567(1991)319

Very-low-density lipoproteins

Separation of human serum lipoproteins into three major classes by hydroxyapatite chromatography 566(1991)67

Villin

The cytoskeletal protein villin as a parameter for early detection of tubular damage in the human kidney 587(1991)81

Vinblastine

High-performance liquid chromatographic determination of vinblastine, 4-O-deacetylvinblastine and the potential metabolite 4-O-deacetylvinblastine-3-oic acid in biological fluids 553(1991)47

Determination of vinca alkaloids in mouse tissues by high-performance liquid chromatography 578(1992)223

Determination of vinorelbine (Navelbine) in tumour cells by high-performance liquid chromatography 581(1992)93

Vinca alkaloids

High-performance liquid chromatographic determination of vinblastine, 4-O-deacetylvinblastine and the potential metabolite 4-O-deacetylvinblastine-3-oic acid in biological fluids 553(1991)47

Biodetermination of N-(deacetyl-O-4-vinblastoyl-23)-L-tryptophan, a metabolite of vintriptol, by high-performance liquid chromatography with fluorescence detection 574(1992)293

Determination of vinca alkaloids in mouse tissues by high-performance liquid chromatography 578(1992)223

Vincamine

Determination of vincamine in human plasma by high-performance liquid chromatography with ultraviolet detection 573(1992)158

Vincristine

Sensitive method for the determination of vincristine in human serum by high-performance liquid chromatography after on-line column-extraction 572(1991)171

Vinorelbine

Bio-analysis of vinorelbine by high-performance liquid chromatography with fluorescence detection 573(1992)328

Determination of vinorelbine (Navelbine) in tumour cells by high-performance liquid chromatography 581(1992)93

Vinpocetine

New assay method for the determination of vinpocetine in human plasma by gas chromatography–mass spectrometry without transesterification caused by solvents: a reply. 567(1991)504

New assay method for the determination of vinpocetine in human plasma by gas chromatography–mass spectrometry without transesterification caused by solvents 567(1991)506

Chiral high-performance liquid chromatographic separations on an α_1 -acid glycoprotein column. II. Separation of the diastereomeric and enantiomeric analogues of vinpocetine (Cavinton) 592(1992)297

Vinriptol

Biodetermination of N-(deacetyl-O-4-vinblastoyl-23)-L-tryptophan, a metabolite of vinriptol, by high-performance liquid chromatography with fluorescence detection 574(1992)293

Vinyl acetate–vinyl chloride copolymer

Determination of solute–polymer interaction constants via chromatography with polymer capillaries 594(1992)29

B₆ Vitamers

Separation and quantification of the B₆ vitamers in plasma and 4-pyridoxic acid in urine of adolescent girls by reversed-phase high-performance liquid chromatography 563(1991)369

Plasma B₆ vitamer and plasma and urinary 4-pyridoxic acid concentrations of middle-aged obese black women 568(1991)333

Vitamin A

Assay of human transthyretin-bound holo-retinol-binding protein with reversed-phase high-performance liquid chromatography 567(1991)369

Determination of retinyl palmitate and total vitamin A content in liver and liver-based ready-to-eat foods 577(1992)163

High-performance liquid chromatographic method for routine determination of vitamins A and E and β -carotene in plasma 581(1992)41

Vitamin B₂

Separation of Dns-amino acids and vitamins by micellar electrokinetic chromatography 559(1991)537

Vitamin B₆

Separation of Dns-amino acids and vitamins by micellar electrokinetic chromatography 559(1991)537

Plasma B₆ vitamer and plasma and urinary 4-pyridoxic acid concentrations of middle-aged obese black women 568(1991)333

Vitamin B

High-performance liquid chromatographic method for routine determination of vitamins A and E and β -carotene in plasma 581(1992)41

Vitamin B₆

Application of gas chromatography–mass spectrometry with selected ion monitoring to the urinalysis of 4-pyridoxic acid 581(1992)179

Micellar electrokinetic capillary chromatography of vitamin B₆ with electrochemical detection 585(1991)139

Vitamin B₁₂

Ultramicrodetermination of cyanocobalamin in elemental diet by solid-phase extraction and high-performance liquid chromatography with visible detection 590(1992)359

Vitamin B₆ vitamers

Determination of vitamin B₆ vitamers and pyridoxic acid in biological samples 578(1992)45

Vitamin C

Simultaneous high-performance liquid chromatographic determination of ascorbic acid and dehydroascorbic acid in biological samples 565(1991)529

Determination of organic acids, amino acids and saccharides by high-performance liquid chromatography and a postcolumn enzyme reactor with amperometric detection 591(1992)165

Determination of ascorbic and dehydroascorbic acid in potatoes (*Solanum tuberosum*) and strawberries using ion-exclusion chromatography 594(1992)187

Vitamin D₂

Preparative isolation of vitamin D₂ from previtamin D₂ by recycle high-performance liquid chromatography 590(1992)169

Vitamin E

Determination of several retinoids, carotenoids and E vitamers by high-performance liquid chromatography. Application to plasma and tissues of rats fed a diet rich in either β -carotene or canthaxanthin 553(1991)179

Simultaneous determination of retinol, α -tocopherol and β -carotene in serum by isocratic high-performance liquid chromatography 572(1991)103

Volatile compounds

Production and use of capillary traps for headspace gas chromatography of airborne volatile organic compounds 552(1991)137

Trace determination of volatile organic compounds in soil based on thermal vaporization followed by Tenax-GC trapping and capillary gas chromatography-mass spectrometry 555(1991)297

Methods of equilibrium concentration for the gas chromatographic determination of trace volatiles (Review) 556(1991)1

Recent methods for the determination of volatile and non-volatile organic compounds in natural and purified drinking water 562(1991)469

Gas chromatographic-mass spectrometric characterization of volatile organic compounds in Barcelona tap water 562(1991)481

Gas chromatographic-mass spectrometric analysis of urban-related aquatic and airborne volatile organic compounds. Study of the extracts obtained by water closed-loop stripping and air adsorption with charcoal and polyurethane foam 562(1991)493

Gas chromatographic-mass spectrometric analysis of some potential toxicants amongst volatile compounds emitted during large-scale thermal degradation of poly(acrylonitrile-butadiene-styrene) plastic 562(1991)681

Method for the collection and analysis of volatile compounds in the breath 564(1991)242

Trapping system for trace organic volatiles 586(1991)315

Vomicine

Separation of *Strychnos nux-vomica* alkaloids by high-performance liquid chromatography 587(1991)318

W

Warfarin

Retention behavior and chiral recognition of β -cyclodextrin-derivative polymer adsorbed on silica for warfarin, structurally related compounds and Dns-amino acids 555(1991)53

Thermospray and particle beam liquid chromatographic-mass spectrometric analysis of coumarin anticoagulants 562(1991)31

Sensitive stereospecific assay of warfarin in plasma: reversed-phase high-performance liquid chromatographic separation using diastereoisomeric esters of (-)-(1*S*,2*R*,4*R*)-endo-1,4,5,6,7,7-hexachlorobicyclo[2.2.1]hept-5-ene-2-carboxylic acid 574(1992)77

Characterization of thyroxine-albumin binding using high-performance affinity chromatography. I. Interactions at the warfarin and indole sites of albumin 579(1992)225

Water

Recent methods for the determination of volatile and non-volatile organic compounds in natural and purified drinking water 562(1991)469

Win 49596

Determination of a novel steroidal androgen receptor antagonist (Win 49596) in human plasma using solid-phase extraction and high-performance liquid chromatography with ultraviolet detection 568(1991)494

X

Xanthenone-4-acetic acid

Determination of xanthenone-4-acetic acid in mouse plasma by high-performance liquid chromatography 564(1991)315

Xanthine

Simultaneous determination of allantoin and oxypurines in biological fluids by high-performance liquid chromatography 575(1992)153

Determination of concentrations of adenosine and other purines in human term placenta by reversed-phase high-performance liquid chromatography with photodiode-array detection: evidence for pathways of purine metabolism in the placenta 575(1992)243

Xanthine oxidase

Extraction of xanthine oxidase from milk by counter-current distribution in an aqueous two-phase system 558(1991)357

Xanthylamides

Gas chromatographic-mass spectrometric determination of ethyl carbamate as the xanthylamide derivative in Italian aqua vitae (grappa) samples 585(1991)111

***o*-Xylene**

Quantitative resolution of severely overlapping gas chromatographic peaks. Isothermal and temperature-programmed operation 556(1991)305

Xylenes

Gas chromatographic separation and automatic identification of complex mixtures of organic solvents in industrial wastes 585(1991)273

Chromatography on Poly-RP and its cyano and diol derivatives using both polar and non-polar solvent systems 591(1992)89

Xylenols

Gas chromatographic determination of urinary phenol conjugates after acid hydrolysis/extractive acetylation 574(1992)349

Xylitol

Chromatographic separation of polyols by ligand exchange. Effects of the ion-exchange resin cross-linking and size 594(1992)125

Xylose

High-performance liquid chromatographic determination of sugars and polyols in extracts of lichens and sugarcane juice 553(1991)271

***m*-Xylylenediamine**

Quality control of cured epoxy resins. Determination of residual free monomers (*m*-xylylenediamine and bisphenol A diglycidyl ether) in the finished product 585(1991)75

Y**Yohimbine**

Determination of *Catharanthus roseus* alkaloids by high-performance liquid chromatography–isotope dilution thermospray-mass spectrometry 554(1991)227

Determination of yohimbine and its two hydroxylated metabolites in humans by high-performance liquid chromatography and mass spectral analysis 574(1992)283

Yohimbine hydrochloride

Simultaneous determination of yohimbine hydrochloride, strychnine nitrate and methyltestosterone by ion-pair high-performance liquid chromatography 588(1991)344

Z**Zardaverine**

Sensitive determination of the phosphodiesterase III/IV inhibitor zardaverine in human serum by direct sample injection, automated precolumn clean-up and high-performance liquid chromatography 553(1991)65

 β -Zearalanol

Determination of zeranol and β -zearalanol in calf urine by immunoaffinity extraction and gas chromatography–mass spectrometry after repeated administration of zeranol 564(1991)493

Zearalenol

Determination of α - and β -zearalenol and zearalenone in cereals by gas chromatography with ion-trap detection 595(1992)259

Zearalenone

Detection of zearalenone in cereal extracts using high-performance liquid chromatography with post-column derivatization 588(1991)47

Determination of α - and β -zearalenol and zearalenone in cereals by gas chromatography with ion-trap detection 595(1992)259

Zeaxanthin

Isolation and structural elucidation of the geometrical isomers of lutein and zeaxanthin in extracts from human plasma 582(1992)153

Optimization of an isocratic high-performance liquid chromatographic separation of carotenoids 589(1992)171

Evaluation of reversed-phase liquid chromatographic columns for recovery and selectivity of selected carotenoids 595(1992)89

Zeranol

Determination of zeranol and β -zearalanol in calf urine by immunoaffinity extraction and gas chromatography–mass spectrometry after repeated administration of zeranol 564(1991)493

Thin-layer chromatographic detection of zeranol and estradiol in fortified plasma and tissue extracts with Fast Corinth V 581(1992)119

General and selective isolation procedure for high-performance liquid chromatographic determination of anabolic steroids in tissues 588(1991)89

Zinc

Use of Zn-pyrophosphatase in the high-performance liquid chromatographic analysis of cell extracts containing ³²P-labelled inositol phosphates 575(1992)51

Determination of divalent trace metals in natural waters by preconcentration on N,N,N',N'-tetra(2-aminoethyl)ethylenediamine-silica followed by on-line ion chromatography 600(1992)203

Zinc-binding proteins

Application of high-performance capillary electrophoresis to the analysis of conformation and interaction of metal-binding proteins 559(1991)345

Zolpidem

Determination of zolpidem, a new sleep-inducing agent, and its metabolites in biological fluids: pharmacokinetics, drug metabolism and overdosing investigations in humans 581(1992)237

Zopiclone

Determination of zopiclone enantiomers in plasma by liquid chromatography using a chiral cellulose carbamate column 572(1991)195

Determination of zopiclone in plasma using column liquid chromatography with ultraviolet detection 581(1992)297

Development of a method for the determination of zopiclone in whole blood 584(1992)199

Zopiclone N-oxide

Development of a method for the determination of zopiclone in whole blood 584(1992)199

Zy 17617B

Determination of Zy 17617B in plasma by solid-phase extraction and liquid chromatography with automated pre-column exchange 593(1992)73

SPECIAL ISSUE INDEX

VOLUMES 551–600

18th International symposium on chromatography, Amsterdam, September 23–28, 1990. Volumes 552 and 553 (1991)

7th International symposium on liquid chromatography–mass spectrometry. Supercritical fluid chromatography–mass spectrometry, capillary electrophoresis–mass spectrometry and tandem mass spectrometry, Montreux (Switzerland), October 31–November 2, 1990. Volume 554 (1991)

Honour volumes on the occasion of the 60th birthday of Georges A. Guiochon. Volumes 556 and 557 (1991)

3rd International symposium on high performance capillary electrophoresis, San Diego, CA (USA), February 3–6, 1991. Volume 559 (1991)

2nd International symposium on applied mass spectrometry in the health sciences, Barcelona (Spain), April 17–20, 1990. Volume 562 (1991)

2nd International symposium on the analysis of anabolizing and doping agents in biosamples, Ghent (Belgium), May 15–18, 1990. Volume 564/2 (1991)

1st International symposium on applications of HPLC in enzyme chemistry, Verona (Italy), September 18–21, 1990. Volume 566/2 (1991)

Biomedical applications of electrophoretic techniques. Volume 569 (1991)

Toxicological and forensic applications of chromatography. Volume 580 (1992)

Bio-chromatography and molecular biology. 4th European meeting of the groupe Français de bio-chromatographie, La Grande Motte (France), May 12–14, 1992. Volume 584/1 (1992)

6th Fast protein liquid chromatography seminar, Titisee (Germany), February 27–March 1, 1991. Volume 587/1 (1991)

8th International symposium on preparative chromatography, Arlington, VA (USA), May 13–15, 1991. Volume 590/1 (1992)

15th International symposium on column liquid chromatography, Basel (Switzerland), June 3–7, 1991. Volumes 592 and 593 (1992)

9th International symposium on affinity chromatography and biological recognition, Yokohama (Japan), September 24–28, 1991. Volume 597 (1992)

11th International symposium on high-performance liquid chromatography of proteins, peptides and polynucleotides, Washington, DC (USA), October 20–23, 1991. Volume 599 (1992)

8th Danube symposium on chromatography, Warsaw (Poland), September 2–6, 1991. Volume 600/1 (1992)

5th Symposium on handling of environmental and biological samples in chromatography, Baden-Baden (Germany), September 26–27, 1991. Volume 600/2 (1992)

PUBLICATION SCHEDULE FOR THE 1993 SUBSCRIPTION

Journal of Chromatography and Journal of Chromatography, Biomedical Applications

MONTH	O 1992	N 1992	D 1992	J	F	M	A	
Journal of Chromatography	623/1 623/2 624/1+2	625/1 625/2	626/1 626/2 627/1+2	628/1 628/2 629/1 629/2	630/1+2 631/1+2 632/1+2 633/1+2	634/1 634/2	635/1 635/2 636/1 636/2	°
Cumulative Indexes, Vols. 601-650								
Bibliography Section						649/1		
Biomedical Applications				612/1	612/2	613/1	613/2 614/1	

° The publication schedule for further issues will be published later.

INFORMATION FOR AUTHORS

(Detailed *Instructions to Authors* were published in Vol. 609, pp. 437-443. A free reprint can be obtained by application to the publisher, Elsevier Science Publishers B.V., P.O. Box 330, 1000 AH Amsterdam, The Netherlands.)

Types of Contributions. The following types of papers are published in the *Journal of Chromatography* and the section on *Biomedical Applications*: Regular research papers (Full-length papers), Review articles, Short Communications and Discussions. Short Communications are usually descriptions of short investigations, or they can report minor technical improvements of previously published procedures; they reflect the same quality of research as Full-length papers, but should preferably not exceed five printed pages. Discussions (one or two pages) should explain, amplify, correct or otherwise comment substantively upon an article recently published in the journal. For Review articles, see inside front cover under Submission of Papers.

Submission. Every paper must be accompanied by a letter from the senior author, stating that he/she is submitting the paper for publication in the *Journal of Chromatography*.

Manuscripts. Manuscripts should be typed in **double spacing** on consecutively numbered pages of uniform size. The manuscript should be preceded by a sheet of manuscript paper carrying the title of the paper and the name and full postal address of the person to whom the proofs are to be sent. As a rule, papers should be divided into sections, headed by a caption (*e.g.*, Abstract, Introduction, Experimental, Results, Discussion, etc.). All illustrations, photographs, tables, etc., should be on separate sheets.

Abstract. All articles should have an abstract of 50-100 words which clearly and briefly indicates what is new, different and significant. No references should be given.

Introduction. Every paper must have a concise introduction mentioning what has been done before on the topic described, and stating clearly what is new in the paper now submitted.

Illustrations. The figures should be submitted in a form suitable for reproduction, drawn in Indian ink on drawing or tracing paper. Each illustration should have a legend, all the *legends* being typed (with double spacing) together on a *separate sheet*. If structures are given in the text, the original drawings should be supplied. Coloured illustrations are reproduced at the author's expense, the cost being determined by the number of pages and by the number of colours needed. The written permission of the author and publisher must be obtained for the use of any figure already published. Its source must be indicated in the legend.

References. References should be numbered in the order in which they are cited in the text, and listed in numerical sequence on a separate sheet at the end of the article. Please check a recent issue for the layout of the reference list. Abbreviations for the titles of journals should follow the system used by *Chemical Abstracts*. Articles not yet published should be given as "in press" (journal should be specified), "submitted for publication" (journal should be specified), "in preparation" or "personal communication".

Dispatch. Before sending the manuscript to the Editor please check that the envelope contains four copies of the paper complete with references, legends and figures. One of the sets of figures must be the originals suitable for direct reproduction. Please also ensure that permission to publish has been obtained from your institute.

Proofs. One set of proofs will be sent to the author to be carefully checked for printer's errors. Corrections must be restricted to instances in which the proof is at variance with the manuscript. "Extra corrections" will be inserted at the author's expense.

Reprints. Fifty reprints will be supplied free of charge. Additional reprints can be ordered by the authors. An order form containing price quotations will be sent to the authors together with the proofs of their article.

Advertisements. The Editors of the journal accept no responsibility for the contents of the advertisements. Advertisement rates are available on request. Advertising orders and enquiries can be sent to the Advertising Manager, Elsevier Science Publishers B.V., Advertising Department, P.O. Box 211, 1000 AE Amsterdam, Netherlands; courier shipments to: Van de Sande Bakhuizenstraat 4, 1061 AG Amsterdam, Netherlands; Tel. (+31-20) 515 3220/515 3222, Telefax (+31-20) 6833 041, Telex 16479 els vi nl. UK: T. G. Scott & Son Ltd., Tim Blake, Portland House, 21 Narborough Road, Cosby, Leics. LE9 5TA, UK; Tel. (+44-533) 753 333, Telefax (+44-533) 750 522. USA and Canada: Weston Media Associates, Daniel S. Lipner, P.O. Box 1110, Greens Farms, CT 06436-1110, USA; Tel. (+1-203) 261 2500, Telefax (+1-203) 261 0101.

Announcing..

International Ion Chromatography Symposium 1993

**September 12-15, 1993
Hyatt Regency Inner Harbor
Baltimore, Maryland USA**

PROGRAM CHAIRMAN:

Richard M. Cassidy
Chemistry Department
University of Saskatchewan
Saskatoon, SK Canada S7N 0W0

Telephone: 306/966-4668
Fax: 306/966-4730

SESSION TOPICS

- Separation Selectivity and Column Technology
- Developments in Separation Methodology
- Advances in Detection
- Special Sample Treatment Procedures
- Novel Applications
- Carbohydrate Separations
- Process Monitoring and Control
- Separation of Metal Ions
- Pharmaceutical Applications
- Environmental Applications
- Ion Analysis in the Electrical Generating Industry
- Interactions at the Capillary/Wall Interface and Ion Separations
- Standard Methods and Data Processing

Poster Abstracts will be accepted until **June 1, 1993.**

For program details and registration information, write or call:

Century International, Inc.
P.O. Box 493 • 25 Lee Road
Medfield, MA 02052 USA
508/359-8777 • 508/359-8778 (FAX)