

VOL. 649*NO. 1 MARCH 19, 1993

Bibliography Section

JOURNAL OF

CHROMATOGRAPHY

INCLUDING ELECTROPHORESIS AND OTHER SEPARATION METHODS

EDITORS

U. A. Th. Brinkman (Amsterdam)
 R. W. Giese (Boston, MA)
 J. K. Haken (Kensington, N.S.W.)
 K. Macek (Prague)
 L. R. Snyder (Orinda, CA)

EDITORS, SYMPOSIUM VOLUMES,

E. Heftmann (Orinda, CA), Z. Deyl (Prague)

EDITORIAL BOARD

D. W. Armstrong (Rolla, MO)
 W. A. Aue (Halifax)
 P. Boček (Brno)
 A. A. Boulton (Saskatoon)
 P. W. Carr (Minneapolis, MN)
 N. H. C. Cooke (San Ramon, CA)
 V. A. Davankov (Moscow)
 Z. Deyl (Prague)
 S. Dilli (Kensington, N.S.W.)
 H. Engelhardt (Saarbrücken)
 F. Erni (Basle)
 M. B. Evans (Hatfield)
 J. L. Glajch (N. Billerica, MA)
 G. A. Guiochon (Knoxville, TN)
 P. R. Haddad (Hobart, Tasmania)
 I. M. Hais (Hradec Králové)
 W. S. Hancock (San Francisco, CA)
 S. Hjertén (Uppsala)
 S. Honda (Higashi-Osaka)
 Cs. Horváth (New Haven, CT)
 J. F. K. Huber (Vienna)
 K.-P. Hupe (Waldbronn)
 T. W. Hutchens (Houston, TX)
 J. Janák (Brno)
 P. Jandera (Pardubice)
 B. L. Karger (Boston, MA)
 J. J. Kirkland (Newport, DE)
 E. sz. Kováts (Lausanne)
 A. J. P. Martin (Cambridge)
 L. W. McLaughlin (Chestnut Hill, MA)
 E. D. Morgan (Keele)
 J. D. Pearson (Kalamazoo, MI)
 H. Poppe (Amsterdam)
 F. E. Regnier (West Lafayette, IN)
 P. G. Righetti (Milan)
 P. Schoenmakers (Eindhoven)
 R. Schwarzenbach (Dübendorf)
 R. E. Shoup (West Lafayette, IN)
 R. P. Singhal (Wichita, KS)
 A. M. Scliffi (Marseille)
 D. J. Trygum (Boston, MA)
 N. Tanaka (Kyoto)
 S. Terabe (Hyogo)
 K. K. Unger (Mainz)
 R. Verpoorte (Leiden)
 Gy. Vigh (College Station, TX)
 J. T. Watson (East Lansing, MI)
 B. D. Westergaard (Uppsala)

EDITORS, BIBLIOGRAPHY SECTION

Z. Deyl (Prague), J. Janák (Brno), V. Schwarz (Prague)

ELSEVIER

blio.
 1.649
 1
 193

JOURNAL OF CHROMATOGRAPHY

INCLUDING ELECTROPHORESIS AND OTHER SEPARATION METHODS

Scope. The *Journal of Chromatography* publishes papers on all aspects of chromatography, electrophoresis and related methods. Contributions consist mainly of research papers dealing with chromatographic theory, instrumental developments and their applications. The section *Biomedical Applications*, which is under separate editorship, deals with the following aspects: developments in and applications of chromatographic and electrophoretic techniques related to clinical diagnosis or alterations during medical treatment; screening and profiling of body fluids or tissues related to the analysis of active substances and to metabolic disorders; drug level monitoring and pharmacokinetic studies; clinical toxicology; forensic medicine; veterinary medicine, occupational medicine; results from basic medical research with direct consequences in clinical practice. In *Symposium volumes*, which are under separate editorship, proceedings of symposia on chromatography, electrophoresis and related methods are published.

Submission of Papers. The preferred medium of submission is on disk with accompanying manuscript (see *Electronic manuscripts* in the Instructions to Authors, which can be obtained from the publisher, Elsevier Science Publishers B.V., P.O. Box 330, 1000 AH Amsterdam, Netherlands). Manuscripts (in English; four copies are required) should be submitted to: Editorial Office of *Journal of Chromatography*, P.O. Box 681, 1000 AR Amsterdam, Netherlands, Telefax (+31-20) 5862 304, or to: The Editor of *Journal of Chromatography, Biomedical Applications*, P.O. Box 681, 1000 AR Amsterdam, Netherlands. Review articles are invited or proposed in writing to the Editors who welcome suggestions for subjects. An outline of the proposed review should first be forwarded to the Editors for preliminary discussion prior to preparation. Submission of an article is understood to imply that the article is original and unpublished and is not being considered for publication elsewhere. For copyright regulations, see below.

Publication. The *Journal of Chromatography* (incl. *Biomedical Applications*) has 40 volumes in 1993. The subscription prices for 1993 are:

J. Chromatogr. (incl. *Cum. Indexes, Vols. 601-650*) + *Biomed. Appl.* (Vols. 612-651):
Dfl. 8520.00 plus Dfl. 1320.00 (p.p.h.) (total ca. US\$ 5622.75)

J. Chromatogr. (incl. *Cum. Indexes, Vols. 601-650*) only (Vols. 623-651):
Dfl. 7047.00 plus Dfl. 957.00 (p.p.h.) (total ca. US\$ 4573.75)

Biomed. Appl. only (Vols. 612-622):

Dfl. 2783.00 plus Dfl. 363.00 (p.p.h.) (total ca. US\$ 1797.75).

Subscription Orders. The Dutch guilder price is definitive. The US\$ price is subject to exchange-rate fluctuations and is given as a guide. Subscriptions are accepted on a prepaid basis only, unless different terms have been previously agreed upon. Subscriptions orders can be entered only by calendar year (Jan. Dec.) and should be sent to Elsevier Science Publishers, Journal Department, P.O. Box 211, 1000 AE Amsterdam, Netherlands, Tel. (+31-20) 5803 642, Telefax (+31-20) 5803 598, or to your usual subscription agent. Postage and handling charges include surface delivery except to the following countries where air delivery via SAL (Surface Air Lift) mail is ensured: Argentina, Australia, Brazil, Canada, China, Hong Kong, India, Israel, Japan, Malaysia, Mexico, New Zealand, Pakistan, Singapore, South Africa, South Korea, Taiwan, Thailand, USA. For Japan air delivery (SAL) requires 25% additional charge of the normal postage and handling charge. For all other countries airmail rates are available upon request. Claims for missing issues must be made within three months of our publication (mailing) date, otherwise such claims cannot be honoured free of charge. Back volumes of the *Journal of Chromatography* (Vols. 1-611) are available at Dfl. 230.00 (plus postage). Customers in the USA and Canada wishing information on this and other Elsevier journals, please contact Journal Information Center, Elsevier Science Publishing Co. Inc., 655 Avenue of the Americas, New York, NY 10010, USA, Tel. (+1-212) 633 3750, Telefax (+1-212) 633 3764.

Abstracts/Contents Lists published in Analytical Abstracts, Biochemical Abstracts, Biological Abstracts, Chemical Abstracts, Chemical Titles, Chromatography Abstracts, Clinical Chemistry Lookout, Current Awareness in Biological Sciences (CABS), Current Contents/Life Sciences, Current Contents/Physical, Chemical & Earth Sciences, Deep-Sea Research/Part B: Oceanographic Literature Review, Excerpta Medica, Index Medicus, Mass Spectrometry Bulletin, PASCAL-CNRS, Pharmaceutical Abstracts, Referativnyi Zhurnal, Research Alert, Science Citation Index and Trends in Biotechnology.

US Mailing Notice. *Journal of Chromatography* (ISSN 0021-9673) is published weekly (total 52 issues) by Elsevier Science Publishers (Sara Burgerhartstraat 25, P.O. Box 211, 1000 AE Amsterdam, Netherlands). Annual subscription price in the USA US\$ 4573.75 (subject to change), including air speed delivery. Application to mail at second class postage rate is pending at Jamaica, NY 11431. **USA POSTMASTERS:** Send address changes to *Journal of Chromatography*, Publications Expediting, Inc., 200 Meacham Avenue, Elmont, NY 11003. Airfreight and mailing in the USA by Publication Expediting.

© 1993 ELSEVIER SCIENCE PUBLISHERS B.V. All rights reserved.

0021-9673/93/06.00

No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, without the prior written permission of the publisher, Elsevier Science Publishers B.V., Copyright and Permissions Department, P.O. Box 521, 1000 AM Amsterdam, Netherlands

Upon acceptance of an article by the journal, the author(s) will be asked to transfer copyright of the article to the publisher. The transfer will ensure the widest possible dissemination of information.

Special regulations for readers in the USA. This journal has been registered with the Copyright Clearance Center, Inc. Consent is given for copying of articles for personal or internal use, or for the personal use of specific clients. This consent is given on the condition that the copier pays through the Center the per-copy fee stated in the code on the first page of each article for copying beyond that permitted by Sections 107 or 108 of the US Copyright Law. The appropriate fee should be forwarded with a copy of the first page of the article to the Copyright Clearance Center, Inc., 27 Congress Street, Salem, MA 01970, USA. If no code appears in an article, the author has not given broad consent to copy and permission to copy must be obtained directly from the author. All articles published prior to 1980 may be copied for a per-copy fee of US\$ 2.25, also payable through the Center. This consent does not extend to other kinds of copying, such as for general distribution, resale, advertising and promotion purposes, or for creating new collective works. Special written permission must be obtained from the publisher for such copying.

No responsibility is assumed by the Publisher for any injury and/or damage to persons or property as a matter of products liability, negligence or otherwise, or from any use or operation of any methods, products, instructions or ideas contained in the materials herein. Because of rapid advances in the medical sciences, the Publisher recommends that independent verification of diagnoses and drug dosages should be made.

Although all advertising material is expected to conform to ethical (medical) standards, inclusion in this publication does not constitute a guarantee or endorsement of the quality or value of such product or of the claims made of it by its manufacturer.

This issue is printed on acid-free paper.

Printed in the Netherlands

CONTENTS

Bibliography Section

Liquid Column Chromatography	B1
Gas Chromatography	B65
Planar Chromatography	B97
Electrophoresis	B112

JOURNAL of CHROMATOGRAPHY

INCLUDING ELECTROPHORESIS AND OTHER SEPARATION METHODS

EDITORS

U. A. Th. BRINKMAN (Amsterdam), R. W. GIESE (Boston, MA), J. K. HAKEN (Kensington, N.S.W.), K. MACEK (Prague),
L. R. SNYDER (Orinda, CA)

EDITORS, SYMPOSIUM VOLUMES

E. HEFTMANN (Orinda, CA), Z. DEYL (Prague)

EDITORIAL BOARD

D. W. Armstrong (Rolla, MO), W. A. Aue (Halifax), P. Boček (Brno), A. A. Boulton (Saskatoon), P. W. Carr (Minneapolis, MN),
N. H. C. Cooke (San Ramon, CA), V. A. Davankov (Moscow), Z. Deyl (Prague), S. Dilli (Kensington, N.S.W.), F. Erni (Basle), M.
B. Evans (Hatfield), J. L. Glajch (N. Billerica, MA), G. A. Guiochon (Knoxville, TN), P. R. Haddad (Hobart, Tasmania), I. M. Hais
(Hradec Králové), W. S. Hancock (San Francisco, CA), S. Hjertén (Uppsala), S. Honda (Higashi-Osaka), Cs. Horváth (New
Haven, CT), J. F. K. Huber (Vienna), K.-P. Hupe (Waldbronn), T. W. Hutchens (Houston, TX), J. Janák (Brno), P. Jandera
(Pardubice), B. L. Karger (Boston, MA), J. J. Kirkland (Newport, DE), E. sz. Kováts (Lausanne), A. J. P. Martin (Cambridge), L.
W. McLaughlin (Chestnut Hill, MA), E. D. Morgan (Keele), J. D. Pearson (Kalamazoo, MI), H. Poppe (Amsterdam), F. E. Regnier
(West Lafayette, IN), P. G. Righetti (Milan), P. Schoenmakers (Eindhoven), R. Schwarzenbach (Dübendorf), R. E. Shoup (West
Lafayette, IN), R. P. Singhal (Wichita, KS), A. M. Siouffi (Marseille), D. J. Strydom (Boston, MA), N. Tanaka (Kyoto), S. Terabe
(Hyogo), K. K. Unger (Mainz), R. Verpoorte (Leiden), Gy. Vigh (College Station, TX), J. T. Watson (East Lansing, MI), B. D.
Westerlund (Uppsala)

EDITORS, BIBLIOGRAPHY SECTION

Z. Deyl (Prague), J. Janák (Brno), V. Schwarz (Prague)



ELSEVIER
AMSTERDAM — LONDON — NEW YORK — TOKYO

J. Chromatogr., Vol. 649 (1993)

The *Bibliography Section 1993* will be published in two volumes (649 and 650) of two issues each. Combined indexes to both volumes will appear in the last issue of the year, Vol. 650, No. 2.

© 1993 ELSEVIER SCIENCE PUBLISHERS B.V. All rights reserved.

0021-9673/93/\$06.00

No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, without the prior written permission of the publisher, Elsevier Science Publishers B.V., Copyright and Permissions Department, P.O. Box 521, 1000 AM Amsterdam, Netherlands.

Upon acceptance of an article by the journal, the author(s) will be asked to transfer copyright of the article to the publisher. The transfer will ensure the widest possible dissemination of information.

Special regulations for readers in the USA. This journal has been registered with the Copyright Clearance Center, Inc. Consent is given for copying of articles for personal or internal use, or for the personal use of specific clients. This consent is given on the condition that the copier pays through the Center the per-copy fee stated in the code on the first page of each article for copying beyond that permitted by Sections 107 or 108 of the US Copyright Law. The appropriate fee should be forwarded with a copy of the first page of the article to the Copyright Clearance Center, Inc., 27 Congress Street, Salem, MA 01970, USA. If no code appears in an article, the author has not given broad consent to copy and permission to copy must be obtained directly from the author. All articles published prior to 1980 may be copied for a per-copy fee of US\$ 2.25, also payable through the Center. This consent does not extend to other kinds of copying, such as for general distribution, resale, advertising and promotion purposes, or for creating new collective works. Special written permission must be obtained from the publisher for such copying.

No responsibility is assumed by the Publisher for any injury and/or damage to persons or property as a matter of products liability, negligence or otherwise, or from any use or operation of any methods, products, instructions or ideas contained in the materials herein. Because of rapid advances in the medical sciences, the Publisher recommends that independent verification of diagnoses and drug dosages should be made.

Although all advertising material is expected to conform to ethical (medical) standards, inclusion in this publication does not constitute a guarantee or endorsement of the quality or value of such product or of the claims made of it by its manufacturer.

This issue is printed on acid-free paper.

Printed in the Netherlands

BIBLIOGRAPHY SECTION

SUPPLEMENT TO THE
JOURNAL OF CHROMATOGRAPHY
1993

EDITORS:

Z. DEYL (Prague)

J. JANÁK (Brno)

V. SCHWARZ (Prague)

Bibliography Section

Liquid Column Chromatography

1. REVIEWS AND BOOKS

- 1 Acedo Sanchez, J.: (Analyzers in process control. II). *Ing. Quim (Madrid)*, 22 (1990) 191-206; C.A., 117 (1992) 114203h - a review with no refs.
- 2 Afeyan, N.B., Fulton, S.P. and Regnier, F.E.: Perfusion chromatography: recent developments and applications. In: Kelly, J.W. and Baldwin, T.O. (Editors), *Appl. Enzyme Biotechnol.*, [Proc. Tex. A&M Univ., IUCCP Symp.], 9th 1991, Plenum, New York, 1991, pp. 221-231; C.A., 117 (1992) 166865v - a review with 12 refs.
- 3 Antia, F.D. and Horvath, C.: Gradient elution chromatography. *NATO ASI Ser., Ser. E*, 204 (1991) 115-136; C.A., 117 (1992) 56429y - a review with 51 refs.
- 4 Bekassy, S. and Szepeszy, L.: (A new unit operation: high-performance liquid chromatography. Part II. Industrial applications of the elution technique). *Magy. Kem. Lapja*, 47 (1992) 185-191; C.A., 117 (1992) 51464w - a review with 18 refs.
- 5 Carta, G.: Simultaneous reaction and chromatography. *NATO ASI Ser., Ser. E*, 204 (1991) 429-442; C.A., 117 (1992) 72209g - a review with 36 refs.
- 6 Carta, G. and Byers, C.H.: Separations by continuous annular chromatography. *NATO ASI Ser., Ser. E*, 204(Chromatogr. Membr. Processes Biotechnol.) (1991) 101-114; C.A., 117 (1992) 85998p - a review with 37 refs.
- 7 Costa, C.A.V.: Modeling equilibrium and kinetics in chromatographic processes. *NATO ASI Ser., Ser. E*, 204 (1991) 3-24; C.A., 117 (1992) 97981r - a review with 63 refs.
- 8 Dolan, J.W.: Solvent recycling. *LC-GC*, 10 (1992) 426-427; C.A., 117 (1992) 142563u - a review with 2 refs.
- 9 Dubin, P.L.: Problems in aqueous size exclusion chromatography. *Adv. Chromatogr. (N.Y.)*, 31 (1992) 119-151; C.A., 117 (1992) 97950e - a review with 95 refs.
- 10 Felix, G.: Survey of packing in donor-acceptor complex chromatography. *Chromatogr. Sci.*, 57 (1992) 33-95; C.A., 117 (1992) 77127p - a review with 596 refs.
- 11 Haddad, P.R. and Jackson, P.E.: *Ion Chromatography: Principles and Applications*. *Chromatogr. Libr.*, Vol. 46, Elsevier, Amsterdam, 1990, 798 p..
- 12 Japan Society for Analytical Chemistry: (*Ion Chromatography Data Book*). Kagaku Shinbunsha, Tokyo, 1991, 548 p; C.A., 117 (1992) 61862p.
- 13 Kissinger, P.T.: Microdialysis and liquid chromatography. *Tech. Behav. Neural Sci.*, 7(Microdial. Neurosci.) (1991) 103-115; C.A., 117 (1992) 65989h.
- 14 Kuznetsov, P.V.: (Current trends in the development of classic and unconventional affinity chromatography (a review)). *Khim.-Farm. Zh.*, 26 (1992) 78-84; C.A., 117 (1992) 103252x - a review with 133 refs.
- 15 Majors, R.E. and Hardy, D.: Sample preparation for biochromatography: an overview. *LC-GC*, 10 (1992) 356-362; C.A., 117 (1992) 22572z - a review with 6 refs.
- 16 Marko-Varga, G.A.: Column liquid chromatography in combination with immobilized enzymes and electrochemical detection and its applications in some industrial processes. *Electroanalysis (N.Y.)*, 4 (1992) 403-427; C.A., 117 (1992) 86027h - a review with 206 refs.
- 17 Nicoud, R.M. and Perrut, M.: Operating modes, scale-up and optimization of chromatographic processes. *NATO ASI Ser., Ser. E*, 204(Chromatogr. Membr. Processes Biotechnol.) (1991) 381-413; C.A., 117 (1992) 69222a - a review with 39 refs.
- 18 Nondek, L.: Complexation in chromatography. *Chromatogr. Sci.*, 57 (1992) 1-31; C.A., 117 (1992) 56423s - a review with 126 refs.
- 19 Rodrigues, A.E., Dias, M.M. and Lopes, J.C.B.: Theory of linear and nonlinear chromatography. *NATO ASI Ser., Ser. E*, 204 (1991) 25-52; C.A., 117 (1992) 97982s - a review with 66 refs.
- 20 Scott, R.P.W.: Modern liquid chromatography. *Chem. Soc. Rev.*, 21 (1992) 137-145; C.A., 117 (1992) 56433v - a review with no refs.
- 21 Teremachi, S.: (Chromatography). *Kobunshi*, 40 (1991) 644-647; C.A., 117 (1992) 131788v - a review with 35 refs.
- 22 Tunlid, A. and Larsson, L. (Editors): *Chromatography and Mass Spectrometry in Microbiology. Second International Symposium on the Interface Between Analytical Chemistry and Microbiology, June 10-13, 1991, Lund University*. *J. Microbiol. Methods*, Elsevier, Amsterdam, 1992, 103 pp.; C.A., 117 (1992) 167147z.
- 23 Wu, S.L.: Rapid methods in applied microbiology: an update. *LC-GC*, 10 (1992) 430-434; C.A., 117 (1992) 86018f - a review with 20 refs.
- 24 Wyatt, P.J.: Combined differential light scattering with various liquid chromatography separation techniques. *Spec. Publ.-R. Soc. Chem.*, 99(Laser Light Scattering Biochem.) (1992) 35-58; C.A., 117 (1992) 86004y - a review with 22 refs.

For additional information see C.A.:
117 (1992) 167168g.

See also 36, 50, 52, 67, 75, 76, 83, 92, 93, 96, 101, 108, 109, 127, 129, 130, 137, 141, 146, 147, 162, 164, 165, 166, 167, 172, 173, 177, 180, 181, 182, 183, 185, 192, 193, 195, 196, 197, 198, 201, 208, 209, 210, 272, 277, 307, 327, 328, 335, 355, 362, 407, 455, 465, 466, 467, 469, 477, 478, 481, 482, 487, 492, 495, 502, 504, 519, 536, 538, 540, 552, 610, 611, 621, 646, 683, 685, 688, 690, 692, 698, 699, 706, 707, 721, 726, 797, 876, 1102, 1103, 1106, 1122, 1170, 1183, 1195, 1214, 1215, 1216, 1259, 1261, 1306, 1308, 1314, 1333, 1334, 1340, 1357, 1437, 1444, 1504, 1505, 1510, 1512, 1524, 1526, 1527, 1528, 1529, 1530, 1531, 1532, 1533, 1545, 1549, 1571, 1583, 1597.

2. FUNDAMENTALS, THEORY AND GENERAL

2a. General

- 25 Blumberg, L.M.: Outline of a theory of focusing in linear chromatography. *Anal. Chem.*, 64 (1992) 2459-2460.
- 26 Bonate, P.L.: Concepts in calibration theory, part III: weighted least-squares regression. *LC-GC*, 10 (1992) 448-450; *C.A.*, 117 (1992) 61697p.
- 27 Cases, J.: HPCPS: a new dimension for HPLC. *Am. Lab. (Shelton)*, 24 (1992) 66; *C.A.*, 116 (1992) 237956t.
- 28 Ceulemans, J.: Role of intermolecular interactions in chromatographic separations. In: Huyskens, P.L., Luck, W.A.P. and Zeegers-Huyskens, T. (Editors), *Intermol. Forces*, Springer, Berlin, 1991, pp. 459-483; *C.A.*, 117 (1992) 178846c.
- 29 Dolan, J.W.: LC problem guide. *LC-GC*, 10 (1992) 508-515; *C.A.*, 117 (1992) 142574y.
- 30 Fleming, P.: Column dead-time determination methods based on the isothermal retention-time behaviour of homologous series in chromatography. *Analyst (London)*, 117 (1992) 1553-1557.
- 31 Gaunitz, U.: Methods for qualitative and quantitative analysis using liquid chromatographic methods. *Ger. (East) DD 294,347 (Cl. G01N30/64)*, 26 Sep. 1991, Appl. 340,459, 08 May 1990; 15 p.; *C.A.*, 117 (1992) 103228u.
- 32 Hallstroem, A., Carlsson, A. and Staahle, L.: An efficient method to develop HPLC-assays. In: Rollema, H., Westerink, B.H.C. and Drijfhout, W.J. (Editors), *Monit. Mol. Neurosci., Proc. Int. Conf. In Vivo Methods, 5th 1991*, Univ. Cent. Pharm., Groningen, 1991, pp. 134-135; *C.A.*, 117 (1992) 146432y.
- 33 Jiao, Q.C. and Chen, Y.Z.: (Computer-aided prediction of optimal composition of multicomponent mobile phase in high performance liquid chromatography). *Yaouxue Xuebao*, 27 (1992) 213-219; *C.A.*, 117 (1992) 22615r.
- 34 Karjalainen, E.J. and Karjalainen, U.P.: Simultaneous analysis of multiple chromatographic runs and samples with alternating regression. *Chemom. Intell. Lab. Syst.*, 14 (1992) 423-427; *C.A.*, 117 (1992) 61951s.
- 35 Kim, S.U., Berninger, J.A., Yu, Q. and Wang, N.H.L.: Peak compression in stepwise pH elution with flow reversal in ion exchange chromatography. *Ind. Eng. Chem. Res.*, 31 (1992) 1717-1730; *C.A.*, 117 (1992) 49184m.
- 36 Kissinger, P.T.: The development of liquid chromatography/electrochemistry from a historical perspective. *Electroanalysis (N.Y.)*, 4 (1992) 359-366; *C.A.*, 117 (1992) 123725g - a review with 29 refs.
- 37 Lee, H.K. and Hoffman, N.E.: Injection solvent effect on peak height in ion exchange HPLC. *J. Chromatogr. Sci.*, 30 (1992) 415-421.
- 38 Manabe, K., Odashima, K. and Koga, K.: Chromatographic separation of aromatic guest compounds by a gel-immobilized water-soluble cyclophane. *Chem. Pharm. Bull.*, 40 (1992) 580-584.
- 39 Marengo, E., Gennaro, M.C. and Abrigo, C.: Experimental design and partial least squares for optimization of reversed-phase ion-interaction liquid chromatographic separation of nitrite, nitrate, and phenylenediamine isomers. *Anal. Chem.*, 64 (1992) 1885-1893.
- 40 Meyer, V.R.: Michael Tswett's columns: facts and speculations. *Chromatographia*, 34 (1992) 342-346.
- 41 Michigami, Y. and Yamamoto, Y.: Behaviour of the system peak in ion chromatography with indirect ultraviolet absorption detection and trimellitate as eluent. *J. Chromatogr.*, 623 (1992) 148-152.
- 42 Pownall, M.: Science and Leviathan: Fisheries research and whales. *Int. Lab.*, 22, No. 9 (1992) P4-P6.

For additional information see *C.A.*:
117 (1992) 103276h.

See also 8, 9, 11, 19, 20, 1086, 1144.

2b. Thermodynamics and theoretical relationships

- 43 Alvarez-Zapeda, A., Barman, B.N. and Martire, D.E.: Thermodynamic study of the marked differences between acetonitrile/water and methanol/water mobile-phase systems in reversed-phase liquid chromatography. *Anal. Chem.*, 64 (1992) 1978-1984.
- 44 Berthod, A. and de Carvalho, M.: Oil-in-water microemulsions as mobile phases in liquid chromatography. *Anal. Chem.*, 64 (1992) 2267-2272.
- 45 Chongying, L.: Silanol group effect in C₁₈ bonded phases in HPLC. *Chromatographia*, 34 (1992) 182-184.
- 46 El'tekova, N.A.: (Thermodynamics of adsorption and retention of aromatic compounds on hydrocarbon and silica sorbents). *Zh. Fiz. Khim.*, 66 (1992) 552-555; *C.A.*, 117 (1992) 138450n.
- 47 Felinga, A., Pasti, L. and Dondi, F.: Fourier analysis of multi-component chromatograms. Recognition of retention patterns. *Anal. Chem.*, 64 (1992) 2164-2174.
- 48 Felinger, A. and Guiochon, G.: Optimization of the experimental conditions and the column design parameters in displacement chromatography. *J. Chromatogr.*, 609 (1992) 35-47.
- 49 Gamzatov, A.Z., Kuznetsov, V.A., Ageev, A.N. and Yashin, Ya.I.: (Estimation of the energy of intermolecular interactions for comparison of the data of liquid adsorption chromatography and spectroscopy). *Zh. Fiz. Khim.*, 66 (1992) 723-729; *C.A.*, 117 (1992) 138444p.
- 50 Gertz, C.: (Optimization methods in HPLC. Testing and dimensioning of solvent transport system). *LaborPraxis*, 16 (1992) 496-505; *C.A.*, 117 (1992) 82650h - a review with 7 refs.

- 51 Grizodub, A.I., Levin, M.G., Asmolova, N.N. and Georgievaskii, V.P.: (Application of Shats' equation for the prediction of retention data in reversed- and normal-phase liquid chromatography). *Zh. Anal. Khim.*, 47 (1992) 514-521; C.A., 117 (1992) 130776c.
- 52 Katti, A.M., Diack, M., El Fallah, M.Z., Golshan-Shirazi, S., Jacobson, S.C., Seidel-Morgenstern, A. and Guiochon, G.: Prediction of high concentration band profiles in liquid chromatography. *Acc. Chem. Res.*, 25 (1992) 366-374; C.A., 117 (1992) 77136r - a review with 27 refs.
- 53 Kord, A.S. and Khaledi, M.G.: Controlling solvent strength and selectivity in micellar liquid chromatography: role of organic modifiers and micelles. *Anal. Chem.*, 64 (1992) 1894-1900.
- 54 Ma, Z. and Guiochon, G.: Shock layer thickness in the case of wide bands of single components and binary mixtures in non-linear liquid chromatography. *J. Chromatogr.*, 609 (1992) 19-33.
- 55 Mechkovskii, S.A., Ovsyanko, L.M., Akinchits, E.A., Abugoffa, A.A. and Muravskaya, N.V.: (Solvent effect in the formation of an analytical signal on surface-layer sorbents). *Izv. Vyssh. Uchebn. Zaved., Khim. Khim. Tekhnol.*, 35 (1992) 32-34; C.A., 117 (1992) 142442d.
- 56 Ugrazov, V.V.: (The codiffusion of mixture components in a capillary column). *Zh. Fiz. Khim.*, 66 (1992) 1049-1053; C.A., 117 (1992) 178927e.
- 57 Urbach, H.P.: Comparison of two hyperbolic systems in chromatography. *Proc. R. Soc. London, Ser. A*, 437 (1992) 403-428; C.A., 117 (1992) 119105n.
- 58 Wei, W., Wu, N.S. and Jiang, X.H.: New approach for area determination of an overlapped pair of chromatographic peaks. *J. Chromatogr.*, 623 (1992) 366-370.
- 59 Wu, N.S. and Wei, W.: Empirical equation for relative non-overlap. *Chromatographia*, 34 (1992) 450-452.
- 60 Zou, H.F., Zhang, Y.K., Hong, M.F. and Lu, P.C.: Combined effects of organic modifier concentration and column temperature on retention in reversed-phase ion-pair liquid chromatography. *Chromatographia*, 34 (1992) 14-18.
- 64 Berthod, A. and Dalaine, V.: (Use of the liquid nature of the stationary phase in countercurrent chromatography for octanol-water partition coefficient determination). *Analisis*, 20 (1992) 325-332; C.A., 117 (1992) 179258z.
- 65 Dubin, P.L., Edwards, S.L., Kaplan, J.I., Mehta, M.S., Tomalia, D. and Xia, J.: Carboxylated starburst dendrimers as calibration standards for aqueous size exclusion chromatography. *Anal. Chem.*, 64 (1992) 2344-2347.
- 66 Juergens, U.: (Narrow-, mini- or microbore-HPLC in routine analysis)? *GIT Spez. Chromatogr.*, 12 (1992) 27-29; C.A., 117 (1992) 82646m - a review with 2 refs.
- 67 Ma, Y.H., Lin, Y.S. and Fleming, H.L.: Chromatographic determinations of adsorption and diffusion for liquids in porous solids by HPLC. In: Mersmann, A.B. and Scholl, S.E. (Editors), *Int. Conf. Fundam. Adsorpt., 3rd 1989*, AICHE, New York, 1991, pp. 517-527; C.A., 117 (1992) 158087c - a review with 16 refs.
- 68 Savitskii, A.A. and Skorokhodov, I.I.: (Equation relating molecular weights of a polymer under study and a standard for exclusion chromatography derived using the nonlinear logarithmic dependence of intrinsic viscosity on molecular weight). *Vysokomol. Soedin., Ser. A*, 34 (1992) 143-148; C.A., 117 (1992) 172414p.
- 69 Wintermantel, M., Schmidt, M., Becker, A., Dorn, R., Kuehn, A. and Loesch, R.: (Structure determination of macromolecules with the gel-permeation chromatography). *Nachr. Chem. Tech. Lab.*, 40 (1992) 331-338; C.A., 117 (1992) 151675j.
- 70 Yamagami, C. and Takao, N.: Hydrophobicity parameters determined by reversed-phase liquid chromatography. VI. Prediction of log P of typical heteroaromatic compounds. *Chem. Express*, 7 (1992) 385-388; C.A., 117 (1992) 25867k.

For additional information see C.A.:

117 (1992) 77407e, 91182j, 133838k, 151749m.

See also 212, 232, 236, 240, 349, 352, 367, 383, 590, 1309, 1312, 1313, 1314, 1316, 1320, 1325, 1326, 1478, 1553.

For additional information see C.A.:

117 (1992) 56704j, 77305v.

See also 7, 25, 63, 155, 207, 217, 282, 637.

2c. Relationship between structure and chromatographic behaviour

- 61 Hirayama, C., Ihara, H., Nagaoka, S. and Syono, T.: Selective retention for structural isomers in aqueous liquid chromatography using poly(γ -methyl L-glutamate) spherical particles. *Chem. Lett.*, (1992) 971-974; C.A., 117 (1992) 103279m.
- 62 Liang, X., Zhang, Y., Huang, H., Zou, H. and Lu, P.: Overall design of recommendation of column system for HPLC. *Chin. Sci. Bull.*, 36 (1991) 1801-1805; C.A., 117 (1992) 39511b.

See also 26, 44, 447, 513, 1401.

2d. Measurement of physico-chemical and related values

- 63 Bailey, R. and Cassidy, R.M.: Mass transfer in the mobile and stationary phases in micellar liquid chromatography. *Anal. Chem.*, 64 (1992) 2277-2282.

3. GENERAL TECHNIQUES

3a. Apparatus and accessories

- 71 Chervet, J.P.: Micro flow processor. *Eur. Pat. Appl. EP 495,255* (Cl. G01N30/34), 22 Jul. 1992, Appl. 91/200,080, 17 Jan. 1991; 12 p.; C.A., 117 (1992) 142628u.
- 72 Day, W.R. and Balogh, M.P.: (LC/mass spectrometer pumping systems: one manufacturer's approach). *Chem. Aust.*, 59 (1992) 256-258; C.A., 117 (1992) 61940n.
- 73 Eray, B. and Unger, K.K.: (Micro-HPLC with packed 320 μ m fused silica capillary column. Part 2: system components and system optimization). *GIT Spez. Chromatogr.*, 12 (1992) 12-18; C.A., 117 (1992) 61943r.
- 74 Fathi, A.H.: Development and application of an on-line process size exclusion chromatograph. *Process Control Qual.*, 2 (1992) 335-346; C.A., 117 (1992) 133377j.
- 75 Figna, A.: (Phase-degassing methods in HPLC). *Lab. 2000* 5 (1991) 52-54; C.A., 117 (1992) 163137y - a review with no refs.

- 76 Goepel, W. and Oehme, M.: Chemical sensors and instrumentation in analytical chemistry. In: Goepel, W., Hesse, J. and Zemel, J.N. (Editors), *Sensors*, 3, VCH, Weinheim, 1992, pp. 819-846; C.A., 117 (1992) 142361b - a review with 36 refs.
- 77 Head, M. and Smith, K.: Flexible instrument control. *Am. Lab. (Shelton)*, 24 (1992) 50D-50H; C.A., 116 (1992) 227307t.
- 78 Kronwald, K.: Multi-component chromatograph column connection. *Ger. Offen. DE 4,014,605 (Cl. B01D15/08)*, 14 Nov. 1991, Appl. 07 May 1990; 8 p.; C.A., 117 (1992) 103309w.
- 79 Nylchreest, L.C. and Hail, M.E.: Multilayer flow electrospray ion source using improved sheath liquid. *U.S. US 5,122,670 (Cl. 250-423R; H01J49/10)*, 16 Jun. 1992, Appl. 702,235, 17 May 1991; 5 p.; C.A., 117 (1992) 162484x.
- 80 Righazza, M. and Siouffi, A.M.: Optimization of the design of a thin-layer cell with a single electrode for electrochemical detection in liquid chromatography. *Analisis*, 20 (1992) 333-340; C.A., 117 (1992) 163004c.
- 81 Simonson, L. and Nelson, K.: The adsorption of amitriptyline hydrochloride in HPLC injector loops. *LC-GC*, 10 (1992) 533-535; C.A., 117 (1992) 157606j.
- For additional information see C.A.:
117 (1992) 39408y, 39413w, 39416z, 39425b, 39575a, 56562m, 61879z, 61883w, 61891x, 82642g, 103321u, 107781s, 142626s, 184049m.
- See also 1, 84, 100, 256, 1522.
- 3b. Detectors and detection reagents**
- 82 Ando, T., Yamamoto, T., Kobayashi, N. and Munekata, E.: Synthesis of a highly luminescent terbium chelate and its application to actin. *Biochim. Biophys. Acta*, 1102 (1992) 186-194.
- 83 Budzikiewicz, H.: Selected reviews on mass-spectrometric topics. XLIV. Continuous flow FAB mass spectrometry. *Mass Spectrom. Rev.*, 11 (1992) 153; C.A., 117 (1992) 123730e - a review with 5 refs.
- 84 Claes, P., Dunford, M., Kenney, A. and Vardy, P.: An online dynamic light scattering instrument for macromolecular characterization. *Spec. Publ. - R. Soc. Chem.*, 99(Laser Light Scattering Biochem.) (1992) 66-76; C.A., 117 (1992) 65756e.
- 85 Dobrowolski, J.C., Strzemecki, G.J. and Jamroz, M.H.: Theory of the intensian method. A simple way to eliminate baseline interference with spectral band intensity. *Chemom. Intell. Lab. Syst.*, 15 (1992) 39-50; C.A., 117 (1992) 82476f.
- 86 Evgen'ev, M.I. and Levinson, F.S.: (Derivatives of dinitrobenzofurazan as selective reagents for organic analysis). *Khim. Geterotsikl. Soedin.*, (1991) 1565; C.A., 117 (1992) 123753q.
- 87 Garcia de Marina, A.: (New range of detection in HPLC: light-scattering detector. Theoretical principles desing advantages and limitation). *Tec. Lab.*, 14 (1992) 220-223; C.A., 117 (1992) 142599k.
- 88 Halasz, J. and Horvai, G.: Indirect amperometric detection of reducible analytes using a reductor column. *Electroanalysis (N.Y.)*, 4 (1992) 469-472; C.A., 117 (1992) 82696c.
- 89 Hjerten, S. and Srichaiyo, T.: Multipoint detection method for electrophoresis and chromatography in capillaries. *U.S. US 5,114,551 (Cl. 204-180.1; B01D57/02)*, 19 May 1992, Appl. 769,073, 30 Sep. 1991; 11 pp.; C.A., 117 (1992) 44039h.
- 90 Ikemoto, N., Lo, L.C. and Naganishi, K.: (Detection of compounds with hydroxyl and amino groups in subpicomole amounts using 2-naphthoylimidazole as a fluorescence label). *Angew. Chem.*, 104 (1992) 918-919; C.A., 117 (1992) 82664r.
- 91 Johnson, D.C. and LaCourse, W.R.: Pulsed electrochemical detection at noble metal electrodes in liquid chromatography. *Electroanalysis (N.Y.)*, 4 (1992) 367-380; C.A., 117 (1992) 123726h.
- 92 Jumel, K., Browne, P. and Kennedy, J.F.: The use of low angle laser light scattering with gel permeation chromatography for the molecular weight determination of biomolecules. *Spec. Publ. - R. Soc. Chem.*, 99(Laser Light Scattering Biochem.) (1992) 23-34; C.A., 117 (1992) 65728x - a review with 19 refs.
- 93 Lickl, E.: (Electrochemical measuring methods). *Oesterr. Chem. Z.*, 93 (1992) 20-21; C.A., 117 (1992) 61666c - a review with 5 refs.
- 94 Long, J.T. and Weber, S.G.: Voltammetry in static and flowing solutions with a large-amplitude sine wave potential. *Electroanalysis (N.Y.)*, 4 (1992) 429-437; C.A., 117 (1992) 123527u.
- 95 Loub, L., Opekar, F., Pacakova, V. and Stulik, K.: A solid polymer electrolyte amperometric detector for FIA and HPLC with mobile phases of low conductivity. *Electroanalysis (N.Y.)*, 4 (1992) 447-451; C.A., 117 (1992) 123738p.
- 96 Mannschreck, A.: Chiroptical detection during liquid chromatography: Part 4. Applications to stereanalysis and stereodynamics. *Chirality*, 4 (1992) 163-169; C.A., 117 (1992) 142564v - a review with 30 refs.
- 97 Mattusch, J., Knauer, H., Noack, J., Ehrhardt, H. and Findeisen, B.: Electrochemical detector for liquid chromatographs of flow-injection analyzers. *Ger. (East) DD 297,250 (Cl. G01N27/02)*, 02 Jan. 1992, Appl. 343,604, 23 Aug. 1990; 6 p.; C.A., 117 (1992) 103201e.
- 98 Msimanga, H.Z. and Sturrock, P.E.: The number of components in noisy data from a swept-potential electrochemical detector. *Electroanalysis (N.Y.)*, 4 (1992) 507-513; C.A., 117 (1992) 123776z.
- 99 Niederlander, H.A.G., Engelaer, F.W., Gooijer, C. and Velthorst, N.H.: HPLC Chemiluminescence detection based on photochemically on-line generated dioxetanes. In: Stanley, P.E. and Kricka, L.J. (Editors), *Biolumin. Chemilumin. Proc. Int. Symp.*, 6th 1990, Wiley, Chichester, 1991, pp. 227-230; C.A., 117 (1992) 142598j.
- 100 Obst, D. and Thietz, P.: Method and apparatus for chromatographic detection. *Ger. (East) DD 294,182 (Cl. B01D15/08)*, 26 Sep. 1991, Appl. 340,529, 10 May 1990; 6 p.; C.A., 117 (1992) 61869w.
- 101 Ozcimder, M. and Demirci, A.: Trends in on-line electrochemical generation of oxidizing and reducing agents for high pressure liquid chromatography. *Kim. Sanayi*, 32 (1992) 86-91; C.A., 117 (1992) 120315n - a review with 17 refs.
- 102 Sato, M. and Yamada, T.: Electrochemiluminiscent detection for HPLC and FIA. In: Stanley, P.E. and Kricka, L.J. (Editors), *Biolumin. Chemilumin. Proc. Int. Symp.*, 6th 1990, Wiley, Chichester, 1991, pp. 231-234; C.A., 117 (1992) 82651j.
- 103 Sorrels, J.W. and Dewald, H.D.: A reticulated vitreous carbon spectroelectrochemical detector for flow injection analysis and liquid chromatography. *Electroanalysis (N.Y.)*, 4 (1992) 487-493; C.A., 117 (1992) 123740h.

- 104 Stojanovic, R.S., Bond, A.M. and Butler, E.C.V.: A comparative study of the cylindrical wire, thin-layer, and wall-jet detector cells for the determination of inorganic arsenic by ion exclusion chromatography with constant and pulsed amperometric detection. *Electroanalysis (N.Y.)*, 4 (1992) 453-461; *C.A.*, 117 (1992) 118038f.
- 105 Szulc, M.E. and Krull, I.S.: Mixed-bed polymeric *o*-nitrobenzophenone reagents for the on-line derivatization of amines in high performance liquid chromatography. *Biomed. Chromatogr.*, 6 (1992) 269-277.
- 106 Tong, W.G.: High-resolution, sensitive nonlinear laser detection method for chemical analysis based on degenerate four-wave mixing. *Proc. SPIE-Int. Soc. Opt. Eng.*, 1637 (1992) 25-32; *C.A.*, 117 (1992) 142444f.
- 107 Tong, W.G.: Sensitive analytical detection method based on nonlinear laser wave mixing inside small capillary flow detector cells. *Proc. Int. Conf. Lasers 1991*, 14th, 1992, pp. 816-820; *C.A.*, 117 (1992) 183821v.
- 108 Uden, P.C. (Editor): *Element-Specific Chromatographic Detection by Atomic Emission Spectroscopy. Developed from a Symposium at the 199th National Meeting of the American Chemical Society, Boston, April 22-27, 1990. ACS Symp. Ser.*, Vol. 479, Washington, 1992, 350 p.; *C.A.*, 116 (1992) 142979p.
- 109 Worsfold, P.J.: Analytical applications of luminescence detection in liquid chromatography and flow injection analysis. In: Stanley, P.E. and Kricka, L.J. (Editors), *Biolumin, Chemilumin. Proc. Int. Symp.*, 6th 1990, Wiley, Chichester, 1991, pp. 371-378; *C.A.*, 117 (1992) 123484c - a review with 10 refs.
- 110 Wu, J., Lu, J. and Zhang, Z.: (Luminol chemiluminescence reaction in liquid chromatography). *Huaxue Shiji*, 14 (1992) 95-98; *C.A.*, 117 (1992) 82426q - a review with 42 refs.
- 111 Zhang, H.-M., Zhou, F.-X. and Krull, I.S.: Comparison of 9-fluorenylmethoxycarbonyl and 9-fluoreneacetyl-tagged silica-based derivatization reagents in high-performance liquid chromatography. *J. Pharm. Biomed. Anal.*, 10 (1992) 577-586.
- 112 Zhou, J. and Wang, E.: Electrocatalytic oxidation and flow detection of hydrazine compounds in liquid chromatography at a vitamin B12 adsorbed glassy carbon electrode. *Electroanalysis (N.Y.)*, 4 (1992) 473-479; *C.A.*, 117 (1992) 123739q.
- 113 Zhou, J. and Wang, W.: Sensitive amperometric detection of glucose by reversed phase liquid chromatography at a Prussian Blue chemically modified electrode of novel construction. *J. Electroanal. Chem.*, 331 (1992) 1029-1043; *C.A.*, 117 (1992) 184009y.
- For additional information see *C.A.*:
117 (1992) 22566a, 22731a, 61974b, 61989k.
- See also 24, 170, 242, 249, 325, 359, 431, 516, 562, 566, 578, 628, 638, 644, 685, 1077, 1217, 1600.
- 3c. *Sorbents and columns, packing procedures*
- 114 Burns, M.A.: Positive focusing in a magnetically stabilized fluidized bed. *U.S. US 5,084,184* (Cl. 210-656; B01D15/08), 28 Jan. 1992, Appl. 667,622, 11 Mar. 1991; 9 p.; *C.A.*, 117 (1992) 29743q.
- 115 Cabrera, K., Lubda, D. and Jung, G.: Alkaline-stable stationary phases for HPLC. Part 1. Aluspher aluminium. *Kontakte (Darmstadt)*, 1 (1992) 12-15; *C.A.*, 117 (1992) 61944s.
- 116 Dawidowicz, A.L.: Properties on controlled porosity glasses (CPG) with chemistry bonded diethylamino groups obtained using various modifiers in the preparation procedure. *Pol. J. Chem.*, 66 (1992) 687-696; *C.A.*, 117 (1992) 158323b.
- 117 Eray, B. and Unger, K.K.: (Micro-HPLC with packed 320 μ m fused silica capillary columns). *GIT Spez. Chromatogr.*, 12 (1992) 5-11; *C.A.*, 117 (1992) 61942q.
- 118 Felix, G., Thienpont, A. and Dentraygues, P.: Synthesis of acceptor bonded phases for donor-acceptor liquid chromatography. *Chromatographia*, 34 (1992) 177-181.
- 119 Fujimoto, C., Suzuki, M. and Jinno, K.: Planarity selectivity in supercritical and liquid chromatography using octadecylsilane-modified silicas with carbon dioxide. *Chromatographia*, 34 (1992) 121-124.
- 120 Funkenbusch, E.F., Carr, P.W., Hanggi, D.A. and Weber, T.P.: Carbon-clad zirconium oxide particles, their manufacture, and uses. *U.S. US 5,108,597* (Cl. 210-198.2; B01D15/08), 28 Apr. 1992, Appl. 497,594, 22 Mar. 1990; 25 p.; *C.A.*, 117 (1992) 174503r.
- 121 Hagel, L.: Pore structure of chromatography media. Comments. *Bioseparation*, 2 (1991) 1-2; *C.A.*, 117 (1992) 39533k.
- 122 Honda, F., Honda, H. and Koishi, M.: (Application of composite particles to column packing for HPLC (high performance liquid chromatography)). *Pharm. Tech. Jpn.*, 8 (1992) 599-605; *C.A.*, 117 (1992) 98306m.
- 123 Honda, F., Honda, H. and Koishi, M.: Application of non-porous silica ultramicrospheres to high-performance liquid chromatographic column packings. *J. Chromatogr.*, 609 (1992) 49-59.
- 124 Huhn, G., Mueller, H. and Eppert, G.: Syntheses of polymer-coated cation-exchangers on the basis of silica gels produced from ester silicates. *React. Polym.*, 17 (1992) 109-113; *C.A.*, 117 (1992) 28256w.
- 125 Inogamov, U.K., Kasymov, S.K., Karas, A.V., Skvortsov, V.T., Salikhov, S.I. and Gurvich, A.E.: Preparation of cellulose sorbent for affinity chromatography using epichlorohydrin and sodium borohydride. *U.S.S.R. SU 1,669,918* (Cl. C08B15/06), 15 Aug. 1991, Appl. 4,689,702, 25 May 1989; *C.A.*, 117 (1992) 51128q.
- 126 Jinno, K., Yamamoto, K., Kuwamoto, T., Nagashima, H., Ueda, T., Tajima, S., Kimura, E., Itoh, K., Fetzter, J.C. and Biggs, W.R.: Multidentate phenyl-bonded phases in high-performance liquid chromatography. *Chromatographia*, 34 (1992) 381-385.
- 127 Kimata, K., Hosoya, K. and Tanaka, N.: (Recent advances in HPLC packing materials). *Bunseki*, (1992) 450-457; *C.A.*, 117 (1992) 119026n - a review with 83 refs.
- 128 King, J.-K. and Pinto, N.G.: Short fibrous supports for preparative chromatographic separations of biomolecules. *J. Chromatogr.*, 609 (1992) 61-68.
- 129 McNair, H.: Chromatography columns: questions and answers. *LC-GC*, 10 (1992) 416-424; *C.A.*, 117 (1992) 142562t - a review with 6 refs.
- 130 Pesek, J.J. and Williamsen, E.J.: The potential uses of bonded liquid crystal materials in high-performance liquid chromatography and supercritical-fluid chromatography. *TrAC*, 11 (1992) 259-266 - a review with 26 refs.
- 131 Pirkle, W.H., Welch, C.J., Burke, J.A. and Lamm, B.: Target-directed desing of chiral stationary phases. *Anal. Proc. (London)*, 29 (1992) 225-226; *C.A.*, 117 (1992) 142573x.

- 132 Reinholdsson, P., Hargital, T., Isaksson, R., Nikitidis, A. and Andersson, C.: Poly(trimethylolpropane trimethacrylate) particles with acid chloride functionality. A convenient entry into support materials for catalysis and chromatography. *React. Polym.*, 17 (1992) 175-186; C.A., 117 (1992) 71157b.
- 133 Rongen, R.M.L., Adler, C.M. and Scouten, W.H.: S-Zephyr, a new high performance ion exchange chromatography column matrix. *Bioseparation*, 2 (1991) 297-308; C.A., 117 (1992) 65781j.
- 134 Shiao, M.S. and Lin, L.J.: Method for evaluating the performance of reverse-phase high-performance liquid chromatographic columns. U.S. US 5,057,224 (Cl. 210-635; B01D15/08), 15 Oct. 1991, Appl. 507,455, 11 Apr. 1990; 12 p.; C.A., 117 (1992) 61990d.
- 135 Takayanagi, H., Honda, G. and Ito, T.: Novel packing materials for chromatographic separation of proteins. In: Abe, M., Kataoka, T. and Suzuki, T. (Editors), *New Dev. Ion Exch. Proc. Int. Conf. Ion Exch.*, Kodansha, Tokyo, 1991, pp. 579-582; C.A., 117 (1992) 12719r.
- 136 Takeuchi, T. and Miwa, T.: Microcolumn liquid chromatography with silica gel dynamically modified with a bile acid derivative. *Chromatographia*, 34 (1992) 386-390.
- 137 Varshney, K.G.: Applications of synthetic inorganic ion exchangers in environmental analysis. In: Abe, M., Kataoka, T. and Suzuki, T. (Editors), *New Dev. Ion Exch., Proc. Int. Conf. Ion Exch.*, Kodansha, Tokyo, 1991, pp. 413-420; C.A., 117 (1992) 123500e - a review with 21 refs.
- 138 Vonk, N., Verstraeten, W.P. and Marinissen, J.W.: Miniaturized columns for the routine HPLC lab: high speed and minibore performance. *J. Chromatogr. Sci.*, 30 (1992) 296-300.
- 139 Waksmundzka-Hajnos, M.: Comparison of adsorption properties of Florisil and silica in high-performance liquid chromatography. II. Retention behaviour of bi- and trifunctional model solutes. *J. Chromatogr.*, 623 (1992) 15-21.

For additional information see C.A.:

117 (1992) 22634w, 22878d, 22903h, 28304k, 39582a, 51737n, 62004x, 72622m, 77148w, 82717k, 93001y, 93003a, 103085v, 103230p, 103325y, 113097q, 114492b, 114496f, 142588f, 146438e, 148780j, 153114z, 156270h, 158311w, 163125t, 166921k, 172298d.

See also 10, 40, 45, 62, 73, 181, 184, 185, 211, 351, 603, 686, 691, 704, 906, 985, 1133, 1161, 1401, 1578, 1603, 1620.

3d. Quantitative analysis

- 140 Astakhov, A.V., Shadrin, A.M. and Glazyrin, E.M.: Method of quantitative chromatographic analysis measuring the amount of the sample using an ionization detector. U.S.S.R. SU 1,693,538 (Cl. G01N30/78), 23 Nov. 1991, Appl. 4,717,275, 11 Jul. 1989; C.A., 117 (1992) 103332y.
- 141 Ivanovic, D. and Medenica, M.: (High performance liquid chromatography in quantitative analysis). *Arh. Farm.*, 41 (1991) 243-253; C.A., 117 (1992) 39499d - a review with 4 refs.

See also 282, 697, 1294.

3e. Preparative scale chromatography

- 142 Gareil, P.: Comparative experimental evaluation of the simple non-linear models available for rapid prediction of the sample size in preparative liquid chromatography. *Prep. Chromatogr.*, 1 (1991) 195-206; C.A., 117 (1992) 77335e.
- 143 Peskin, A.P. and Rudge, S.R.: Optimization of large-scale chromatography for biotechnological applications. *Appl. Biochem. Biotechnol.*, 34-35 (1992) 49-59; C.A., 117 (1992) 46589m.

See also 17, 128, 234, 268, 315, 352, 531, 626, 695, 970.

3f. Programmed temperature, pressure, vapors, gradients

- 144 Dolan, J.W.: The case of the mystery gradient peak. *LC-GC*, 10 (1992) 294-296; C.A., 117 (1992) 39534m.
- 145 Gazda, K.: Effect of sample volume on band width of a substance in gradient elution chromatography. *Chem. Anal. (Warsaw)*, 36 (1991) 507-523; C.A., 117 (1992) 82517v.
- 146 Gertz, C.: (Optimization methods in HPLC. Optimization of elution power, pH value, buffer and ion-pair concentration, and temperature of isocratic and gradient systems). *LaborPraxis*, 16 (1992) 642-653; C.A., 117 (1992) 123731f - a review with 30 refs.

See also 3, 66, 1405.

4. SPECIAL TECHNIQUES

4a. Automation

- 147 Brinkman, U.A.T. and Grossi, G.: (Automation of specimen handling in HPLC). *Lab. 2000*, 5 (1991) 62-72; C.A., 117 (1992) 103247z - a review with 19 refs.
- 148 Verillon, F., Qian, F. and Rasquin, P.: The use of on-line dialysis with HPLC for the automated preparation and analysis of complex samples. *Anal. Sci.*, 7 (1991) 1511-1517; C.A., 117 (1992) 142584b.
- 149 Vogel, W. and Steinwand, M.: Advanced sample processing in HPLC. *Int. Lab.*, 22, No. 8 (1992) P16-P27.

See also 74, 701, 1380, 1492.

4b. Computerization and modelling

- 150 Chunfeng, Z., Nina, M., Xianfeng, S., Hong, W., Peihong, Z. and Minggao, X.: An approximate experimental model for predicting solute retention times in RPHPLC with quaternary solvent systems. *J. High Resolut. Chromatogr.*, 15 (1992) 555-561.
- 151 Colomina, O.: (Current trends in the design of systems for treatment of analytical data). *Quim. Ind. (Madrid)*, 38 (1992) 119-120; C.A., 117 (1992) 162972e.
- 152 Danielsson, R. and Malmquist, G.: Multi-dimensional simplex interpolation, An approach to local models for prediction. *Chemom. Intell. Lab. Syst.*, 14 (1992) 115-128; C.A., 117 (1992) 119008h.

- 153 Dolan, J.W., Lewis, J.A., Raddatz, W.D. and Snyder, L.R.: Multi-parameter computer simulation for HPLC method development. *Am. Lab. (Shelton)*, 24 (1992) 40D, 40F-40J, 40L; C.A., 116 (1992) 268307m.
- 154 Dong, M.W., Vandemark, F.L. and Passalacqua, P.V.: Validating HPLC methods using an intelligent system controller. *Am. Lab. (Shelton)*, 24 (1992) 36-45; C.A., 117 (1992) 56085h.
- 155 Gerritsen, M.J.P., Tanis, H., Vandeginste, B.G.M. and Kateman, G.: Generalized rank annihilation factor analysis, iterative target transformation factor analysis, and residual bilinearization for the quantitative analysis of data from liquid chromatography with photodiode array detection. *Anal. Chem.*, 64 (1992) 2042-2056.
- 156 Green, A.: Benefits of client/server computing for analytical laboratories. *Int. Lab.*, 22, No. 10 (1992) 1-26.
- 157 Jaufmann, L.: (Good laboratory practice: validation in the chromatographic laboratory). *CLB Chem. Labor Biotech.*, 43 (1992) 426-427; C.A., 117 (1992) 184002r.
- 158 Kaljurand, M.: (Computer chromatography of nonstationary substance flows). *Zh. Anal. Khim.*, 47 (1992) 279-293; C.A., 117 (1992) 163159g.
- 159 Nau, V.A. and White, L.B.: A microrobotics approach to LC sample preparation. *Am. Lab. (Shelton)*, 24 (1992) 64-72; C.A., 116 (1992) 247591j.
- 160 Nelson, D.C., Dauciuinas, V., Lysakowski, R. and Duff, M.: The Standard for Chromatography Data Communication from the Analytical Instruments Association. *Int. Lab.*, 22, No. 10 (1992) 32-38.
- 161 Ouchi, G.I.: Custom reporting using a spreadsheet program. *LC-GC*, 10 (1992) 436-440; C.A., 117 (1992) 82455y.
- 162 Ouchi, G.I.: Using chromatographic data in a data base. *LC-GC*, 10 (1992) 304-308; C.A., 117 (1992) 39498c - a review with no refs.
- 163 Sato, H. and Watanabe, H.: Computer simulation of ion-exchange chromatography with bulkproperty detectors. *React. Polym.*, 17 (1992) 1-8; C.A., 117 (1992) 39287h.
- 164 Schmitz, F.: (Experiences on installation of a computer-supported operation of ion chromatography (IC) and result reporting by on-line coupling with laboratory data management). *Gewaesserschutz, Wasser, Abwasser*, 124 (1991) 122-141; C.A., 117 (1992) 142339a - a review with 4 refs.
- 165 Stienen, G.: (Requirements of hardware and software for rationalizing and increasing efficiency of chromatographic systems). *Gewaesserschutz, Wasser, Abwasser*, 124 (1991) 149-158; C.A., 117 (1992) 142559x - a review with no refs.
- 167 Arpino, P.J.: Combined liquid chromatography/mass spectrometry (LC/MS): a review. *NATO ASI Ser., Ser. C*, 353(Mass Spectrom. Biol. Sci.: Tutorial) (1992) 253-267; C.A., 117 (1992) 166859w - a review with 110 refs.
- 168 Hsu, F.F.: Water as a useful calibrant for thermospray mass spectrometry. *Biol. Mass Spectrom.*, 21 (1992) 363-364; C.A., 117 (1992) 103254z.
- 169 Iwahashi, H., Parker, C.E., Mason, R.P. and Tomer, K.B.: Combined liquid chromatography/electron paramagnetic resonance spectrometry/electrospray mass spectrometry for radical identification. *Anal. Chem.*, 64 (1992) 2244-2252.
- 170 Lago, P. and Corti, M.: Dynamic laser light scattering system for chromatographic analysis. *Conf. Proc.-Ital. Phys. Soc.*, 29 (1991) 351-354; C.A., 117 (1992) 162995q.
- 171 Lawrence, D.L.: The application of normal phase liquid chromatography/mass spectrometry by using coaxial continuous flow fast atom bombardment. *J. Am. Soc. Mass Spectrom.*, 3 (1992) 575-581; C.A., 117 (1992) 163142w.
- 172 Schmarr, H.G.: (Online coupling of HPLC with gas chromatography (GC)). *LaborPraxis*, 16 (1992) 716-721; C.A., 117 (1992) 183996f - a review with 8 refs.
- 173 Schwedt, G. and Meyer, A.: (Coupling techniques in chromatography). *LaborPraxis*, 16 (1992) 712-714; C.A., 117 (1992) 183995e - a review with 16 refs.
- 174 Spraul, M., Hofmann, M., Dvortsak, P., Nicholson, J.K. and Wilson, I.D.: Liquid chromatography coupled with high-field proton NMR for profiling human urine for endogenous compounds and drug metabolites. *J. Pharm. Biomed. Anal.*, 10 (1992) 601-605.
- 175 Vreeken, R.J., van Dongen, W.D., Chijsen, R.T., de Jong, G.J., Longeman, H., Brinkman, U.A.T., van Leuken, R.G.J., Kwakkenbos, G.T.C. and Deelder, R.S.: Coupling of column liquid chromatography and thermospray mass spectrometry via phase-system switching with an ion-exchange trapping column. *Biol. Mass Spectrom.*, 21 (1992) 305-314; C.A., 117 (1992) 39519k.
- See also 36, 72, 79, 108, 219, 220, 238, 241, 257, 266, 308, 322, 417, 450, 454, 460, 465, 467, 492, 500, 504, 677, 703, 789, 1131, 1180, 1226, 1263, 1274, 1305, 1344, 1429, 1433, 1444, 1473, 1482, 1520, 1530, 1544, 1590, 1604.

4d. Affinity chromatography (advances)

For additional information see C.A.:
117 (1992) 103095y.

See also 7, 33, 39, 47, 77, 142, 207, 253, 1600.

4c. Combination with other physico-chemical techniques (MS, IR etc.)

166 Arpino, P.: Combined liquid chromatography-mass spectrometry. Part III. Applications of thermospray. *Mass Spectrom. Rev.*, 11 (1992) 3-40; C.A., 117 (1992) 142561s - a review with 295 refs.

- 176 Cassidy, S.A., Janis, L.J. and Regnier, F.E.: Kinetic chromatographic sequential addition immunoassays using protein A affinity chromatography. *Anal. Chem.*, 64 (1992) 1973-1977.
- 177 Gasinska, E. and Maslowska, J.: (Affinity chromatography on special polymeric supports and its use in separation and purification of bioactive substances). *Wlokan Chem.*, 17 (1991) 15-27; C.A., 117 (1992) 166823e - a review with 10 refs.
- 178 Gilpin, R.K., Gao, H. and Jaroniec, M.: Liquid chromatographic studies of silica-immobilized bovine serum albumin under normal-phase conditions: separation of phenolic solutes using ternary mixtures of hydrogen chloride-diethyl ether-hexane as eluents. *J. Liq. Chromatogr.*, 15 (1992) 2503-2518.
- 179 Jack, G.W.: Immunoaffinity chromatography. *Methods Mol. Biol. (Totowa)*, 11(Pract. Protein Chromatogr.) (1992) 125-133; C.A., 117 (1992) 167053r.

- 180 Liapis, A.I.: Biospecific adsorption mechanisms and processes of affinity chromatography. In: Mersmann, A.B. and Scholl, S.E. (Editors), *Int. Conf. Fundam. Adsorpt., 3rd 1989*, AIChE, New York, 1991, pp. 25-61; C.A., 117 (1992) 127466w - a review with 66 refs.
- 181 Lowe, C.R.: Affinity chromatography: design of biospecific chromatographic materials. *NATO ASI Ser., Ser. E*, 204 (1991) 353-361; C.A., 117 (1992) 76279c - a review with 24 refs.
- 182 Lowe, C.R.: Affinity chromatography: principles and applications. *NATO ASI Ser., Ser. E*, 204 (1991) 297-308; C.A., 117 (1992) 56432u - a review with 30 refs.
- 183 McCoy, B.J.: Theory of affinity chromatography. *NATO ASI Ser., Ser. E*, 204 (1991) 297-308; C.A., 117 (1992) 56432u - a review with 30 refs.
- 184 Mohan, S.B. and Lyddiatt, A.: Silica-based solid phases for affinity chromatography: effect of pore size and ligand location upon biochemical productivity. *Biotechnol. Bioeng.*, 40 (1992) 549-563; C.A., 117 (1992) 166935t.
- 185 Mueller-Schulte, D., Schaefer-Bilger, U. and Brunner, H.: (New media for affinity chromatography on the basis of grafted polymers). *GIT Fachz. Lab.*, 36 (1992) 315-322; C.A., 117 (1992) 39507e - a review with 21 refs.
- 186 Ngo, T.T.: Facile activation of Trisacryl gels with 2-fluoro-1-methylpyridinium salt (FMP): applications in affinity chromatography and enzyme immobilization. In: Gebelein, C.G., Cheng, T.C. and Yang, V.C.-M. (Editors), *Cosmet. Pharm. Appl. Polym., [Proc. Am. Chem. Soc. Symp. Polym. Cosmet. Pharm. Appl.]*, 1990, Plenum, New York, 1991, pp. 385-398; C.A., 117 (1992) 107732b.
- 187 Ohlson, S.: Exploiting weak affinities. *Methods Mol. Biol. (Totowa)*, 11 (Pract. Protein Chromatogr.) (1992) 197-208; C.A., 117 (1992) 166910f.
- 188 Pepper, D.S.: Some alternative coupling chemistries for affinity chromatography. *Methods Mol. Biol. (Totowa)*, 11 (Pract. Protein Chromatogr.) (1992) 173-196; C.A., 117 (1992) 167113k.
- 189 Petropoulos, J.H., Liapis, A.I., Petrou, J.K., Kanellopoulos, N.K. and Kollopoulos, N.P.: Restricted diffusion of biomolecules in affinity adsorbents. In: Mersmann, A.B. and Scholl, S.E. (Editors), *Int. Conf. Fundam. Adsorpt., 3rd 1989*, AIChE, New York, 1991, pp. 665-678; C.A., 117 (1992) 127513j.
- 190 Rhee, J. and Andersen, N.H.: Preparation of imidazolyl acids tethered to agarose gel and assessment of the degree of coupling for affinity chromatography. *Bull. Korean Chem. Soc.*, 13 (1992) 208-210; C.A., 117 (1992) 146194x.
- 191 Scopes, R.K.: Biospecific affinity. *Methods Mol. Biol. (Totowa)*, 11 (Pract. Protein Chromatogr.) (1992) 209-221; C.A., 117 (1992) 166911g.
- 192 Shi, J. and Du, H.: (Application of immobilized enzyme reactors in high performance liquid chromatography). *Sepu*, 10 (1992) 204-206; C.A., 117 (1992) 127473w - a review with 27 refs.
- 193 Suwanrumpha, S., McLean, M.A., Fink, S.W., Wilder, C., Stachowiak, K., Dyckes, D.F. and Freas, R.B.: Determination of biomolecules by using liquid chromatography and thermospray mass spectrometry: from penicillins to peptides. *NATO ASI Ser., Ser. C*, 353 (Mass Spectrom. Biol. Sci.: Tutorial) (1992) 281-301; C.A., 117 (1992) 166860q - a review with 89 refs.
- 194 West, I. and Goldring, O.: Lectin affinity chromatography. *Methods Mol. Biol. (Totowa)*, 11 (Pract. Protein Chromatogr.) (1992) 81-89; C.A., 117 (1992) 166909n.
- 195 Xiong, K.: (Pseudobiospecific ligand affinity chromatography). *Shengwu Huaxue Yu Shengwu Wuli Jinzhan*, 19 (1992) 21-26; C.A., 117 (1992) 86034h - a review with 20 refs.
- 196 Yip, T.T. and Hutchens, T.W.: Immobilized metal ion affinity chromatography. *Methods Mol. Biol. (Totowa)*, 11 (Pract. Protein Chromatogr.) (1992) 17-31; C.A., 117 (1992) 166846q - a review with 29 refs.

For additional information see C.A.:

117 (1992) 22621q, 22648d, 107790u, 127800a, 129859a.

See also 14, 121, 125, 201, 274, 307, 385, 391, 402, 609, 683, 698, 702, 714, 723, 745, 746, 752, 759, 761, 763, 783, 815, 830, 869, 878, 923, 945, 988, 1005, 1008, 1070, 1102, 1333, 1472.

4e. Functional analysis

See 136.

4f. Trace analysis and pre-separation techniques

197 Fardy, J.J.: Preconcentration of trace elements by ion exchangers. In: Alfassi, Z.B. and Wai, C.M. (Editors), *Preconc. Tech. Trace Elem.*, CRC, Boca Raton, 1992, pp. 181-210; C.A., 117 (1992) 123493e - a review with 224 refs.

198 Haddad, P.R. and Jackson, P.E.: Chromatographic preconcentration methods for inorganic species. In: Alfassi, Z.B. and Wai, C.M. (Editors), *Preconc. Tech. Trace Elem.*, CRC, Boca Raton, 1992, pp. 243-274; C.A., 117 (1992) 103022x - a review with 91 refs.

199 Law, B. and Weir, S.: Fundamental studies in reversed-phase liquid-solid extraction of basic drugs; III: sample matrix effects. *J. Pharm. Biomed. Anal.*, 10 (1992) 487-493.

200 Pinto, C.G., Pavón, J.L.P. and Cordero, B.M.: Cloud point preconcentration and high-performance liquid chromatographic analysis with electrochemical detection. *Anal. Chem.*, 64 (1992) 2334-2338.

201 Van Ginkel, L.A., Stephany, R.W., van Rossum, H.J. and Zoontjes, P.W.: Perspectives in residue analysis; the use of immobilized antibodies in (multi) residue analysis. *TrAC*, 11 (1992) 294-298 - a review with 19 refs.

See also 90, 260, 1322, 1323, 1584, 1596.

4g. Enantiomers, separation

202 Altria, K.D., Goodall, D.M. and Rogan, M.M.: Chiral separation of β -amino alcohols by capillary electrophoresis using cyclodextrins as buffer additives. I. Effect of varying operating parameters. *Chromatographia*, 34 (1992) 19-24.

203 Casarini, D., Lunazzi, L., Pasquali, F., Gasparrini, F. and Villani, C.: Conformational studies by dynamic NMR. 47. Conformation, stereodynamics, and chiral separation of the rotational enantiomers of hindered naphthyl ketones. *J. Am. Chem. Soc.*, 114 (1992) 6521-6527; C.A., 117 (1992) 69299f.

204 Hargrave, C.P. and Waters, D.G.: High-performance liquid chromatographic determination of the enantioselectivity of a novel Reformatsky reaction. *Anal. Proc. (London)*, 29 (1992) 242-244; C.A., 117 (1992) 130876k.

- 205 Mano, N., Oda, Y., Asakawa, N., Yoshida, Y., Sato, T. and Miwa, T.: Development of a flavoprotein column for chiral separation by high-performance liquid chromatography. *J. Chromatogr.*, 623 (1992) 221-228.
- 206 Meeson, S.R.: The separation of enantiomers by high-performance liquid chromatography. Avail. *Univ. Microfilms Int.*, Order No. BRDX93606, 1990, 269 p.; *C.A.*, 117 (1992) 142597h.
- 207 Meyer, V.R.: Use of chromatographic peak-heights ratios for quantitative analysis: application to the separation of enantiomers. *J. Chromatogr.*, 623 (1992) 371-374.
- 208 Shibata, T.: Optical resolution by liquid chromatography with a chiral stationary phase. *Organomet. News*, (1991) 45-52; *C.A.*, 117 (1992) 123727j - a review with 23 refs.
- 209 Snopek, J., Jelínek, I. and Smolková-Keulemansová, E.: Chiral separation by analytical electromigration methods. *J. Chromatogr.*, 609 (1992) 1-17 - a review with 97 refs.
- 210 Srinivas, N. and Igwemezie, L.N.: Chiral separation by high-performance liquid chromatography. I. Review on direct separation of enantiomers as diastereomeric derivatives using ultraviolet, fluorescence and electrochemical detection. *Biomed. Chromatogr.*, 6 (1992) 163-167; *C.A.*, 117 (1992) 137741w - a review with many refs.
- 211 Straub, R. and Arm, H.: Thermal treatment of supports for chiral stationary phases and its influence on enantioselectivity. *J. Chromatogr.*, 623 (1992) 23-31.

For additional information see *C.A.*:
117 (1992) 69730q.

See also 96, 131, 243, 245, 288, 290, 291, 294, 403, 404, 409, 414, 460, 484, 512, 517, 541, 544, 573, 612, 985, 1114, 1122, 1128, 1133, 1135, 1151, 1291, 1327, 1329, 1330, 1331, 1339, 1351, 1355, 1357, 1361, 1362, 1363, 1374, 1389, 1401, 1409, 1415, 1431, 1435, 1439, 1440, 1479, 1486, 1548, 1549.

4h. Other special techniques

- 212 Fitzgerald, J.J., Nebo, C.O. and Fuerstenau, M.C.: Gel filtration chromatography of supernatants from aluminium silicate hydrogels of silicon/aluminium ratios from 3.33/1 to 13.32/1. *J. Colloid Interface Sci.*, 151 (1992) 317-332; *C.A.*, 117 (1992) 77391v.
- 213 Grob, K. and Tönz, B.: On-line solvent evaporator for coupled LC systems. *J. High Resolut. Chromatogr.*, 15 (1992) 594-600.
- 214 Hendriks, P.J.M., Claessens, H.A., Noij, T.H.M., Everaerts, F.M. and Cramers, C.A.: On-line isotachopheresis. A selective sample pretreatment prior to column liquid chromatographic analysis. *Chromatographia*, 33 (1992) 539-545.
- 215 Pettit, H.O. and Justice, J.B., Jr.: Procedures for microdialysis with small bore HPLC. *Tech. Behav. Neural Sci.*, 7(Microdial. Neurosci.) (1991) 117-153; *C.A.*, 117 (1992) 65990b.

See also 6, 13, 16, 27, 53, 63, 373, 620, 705, 1471, 1567.

5. HYDROCARBONS AND HALOGEN DERIVATIVES

5a. Aliphatic hydrocarbons

- 216 Sharer, J.E., Duescher, R.J. and Elfarra, A.A.: Species and tissue differences in the microsomal oxidation of 1,3-butadiene and the glutathione conjugation of butadiene monoxide in mice and rats. Possible role in 1,3-butadiene-induced toxicity. *Drug Metab. Disp.*, 20 (1992) 658-664.

See also 520.

5b. Cyclic hydrocarbons

- 217 Barman, B.N. and Martire, D.E.: Factors influencing retention and resolution of substituted alkylbenzenes in reversed-phase liquid chromatography. *Chromatographia*, 34 (1992) 347-356.
- 218 Bhyrappa, P., Penicaud, A., Kawamoto, M. and Reed, C.A.: Improved chromatographic separation and purification of C₆₀ and C₇₀ fullerenes. *J. Chem. Soc., Chem. Commun.*, (1992) 936-937; *C.A.*, 117 (1992) 102813u.
- 219 Diack, M., Hettich, R.L., Compton, R.N. and Guiochon, G.: Contribution to the isolation and characterization of buckminsterfullerenes. *Anal. Chem.*, 64 (1992) 2143-2148.
- 220 Jinno, K., Uemura, T., Nagashima, H. and Itoh, K.: An LC-MS approach to the on-line identification of fullerenes. *J. High Resolut. Chromatogr.*, 15 (1992) 627-628.
- 221 Lamprecht, G. and Huber, J.F.K.: Direct determination of benzo[a]pyrene in oil distillates by on-line two-dimensional HPLC with column switching. *Chromatographia*, 34 (1992) 376-380.
- 222 Rinderle, S.J., Black, S.D., Sharma, P.K. and Witiak, D.T.: Comparative metabolism *in vitro* of a novel carcinogenic polycyclic aromatic hydrocarbon, 1,2,3,4-tetrahydro-7,12-dimethylbenz[a]anthracene, and its two regioisomeric B-ring fluoro analogues. *Cancer Res.*, 52 (1992) 3035-3042.
- 223 Swaminathan, S. and Reznikoff, C.A.: Metabolism and nucleic acid binding of N-hydroxy-4-acetylaminobiphenyl and N-acetoxy-4-acetylaminobiphenyl by cultured human uroepithelial cells. *Cancer Res.*, 52 (1992) 3286-3294.
- 224 Tamakawa, K., Seki, T. and Tsunoda, A.: HPLC detection of polycyclic aromatic hydrocarbons. *Food Sci. Technol.*, 52(Food Anal. HPLC) (1992) 697-711; *C.A.*, 117 (1992) 110246b.
- 225 Tatarovicova, V. and Bekarek, V.: Determination of various aromatic hydrocarbons in slurries, soil, silage, and impregnation oils by high-pressure liquid chromatography and infrared spectroscopy. *Acta Univ. Palacki. Olomuc., Fac. Rerum Nat.*, 102 (1991) 67-75; *C.A.*, 117 (1992) 123793c.
- 226 Velasco Oetega, J., Laborda Grima, R. and Balasch Parisi, S.: (Determination of benzo(a)pyrene by on-line multidimensional liquid chromatography). *Tec. Lab.*, 14 (1992) 202-207; *C.A.*, 117 (1992) 123792b.
- 227 Welch, C.J. and Pirkle, W.H.: Progress in the design of selectors for buckminsterfullerene. *J. Chromatogr.*, 609 (1992) 89-101.

See also 44, 46, 118, 126, 130, 1085, 1135, 1504.

5c. Halogen derivatives

- 228 Bamhart, E.R., Banhart, E.R., Kimata, K., Hosoya, K., Terabe, S., Alexander, L.R. and Patterson, D.G., Jr.: HPLC elution order of polychlorinated biphenyl congeners with octadecyl-, pyrenylethyl-, and nitrated naphthylethyl-silylated silica gel. *Organohalogen Compd.*, 2(EPRI PCB-Semin., Anal., AQ-QC, Brominated Compd., Short-Chain Aliphatics) (1990) 87-90; C.A., 117 (1992) 165350m.
- 229 Nilsson, N.L. and Colmsjö, A.L.: Retention characteristics of chlorinated polycyclic aromatic hydrocarbons in normal phase HPLC. II. Chloro-substituted PAHs. *Chromatographia*, 34 (1992) 115-120.
- 230 Siren, H., Hyvönen, H., Saarinen, M., Rovio, S. and Riekkola, M.-L.: Comparison of HPLC and solid phase clean-up methods for identification of PCBs in cod-liver oil by HRGC/MS-SIM technique. *Chromatographia*, 34 (1992) 421-430.
- 231 Wells, D.E., Maier, E.A. and Griepink, B.: Development in the analysis for chlorobiphenyls in environmental matrices for certification purposes. *Int. J. Environ. Anal. Chem.*, 46 (1992) 265-275; C.A., 117 (1992) 82667u.

See also 279, 1268.

5d. Complex hydrocarbon mixtures (incl. analysis of tars, bitumens and mineral oils)

- 232 Acevedo, S., Escobar, G., Gutierrez, L.B. and D'Aquino, J.: Synthesis and isolation of octylated asphaltene standards for calibration of GPC columns and determination of asphaltene molecular weight. *Fuel*, 71 (1992) 1077-1079; C.A., 117 (1992) 174728t.
- 233 Branthaver, J.F., Catalfomo, M.W. and Petersen, J.C.: Ion exchange chromatography separation of SHRP asphalts. *Fuel Sci. Technol. Int.*, 10 (1992) 855-885; C.A., 117 (1992) 96246t.
- 234 Branthaver, J.F., Duvall, J.J., Petersen, J.C., Plancher, H. and Robertson, R.E.: Preparative size-exclusion chromatography separation of SHRP asphalts: correlation with viscoelastic properties. *Fuel Sci. Technol. Int.*, 10 (1992) 1003-1032; C.A., 117 (1992) 96251r.
- 235 Cosgrove, M.: New analytical techniques for monitoring chemical changes in rolling oils which affect steel surface cleanliness. *Commun. Eur. Communities, (Rep.) EUR*, EUR 14113 (1992) 117-124; C.A., 117 (1992) 134113g.
- 236 Georgiev, I. and Minkova, V.: (Study of the shale oil heavy fraction using gel chromatography in combination with the other methods). *Zh. Prikl. Khim. (S.-Peterburg)*, 64 (1991) 2551-2556; C.A., 117 (1992) 72835h.
- 237 Jennigs, P.W., Pribanic, J.A.S., Mendes, T.M. and Smith, J.A.: High performance gel permeation chromatography in the characterization of self-assemblies in asphalt. I. *Fuel Sci. Technol. Int.*, 10 (1992) 809-823; C.A., 117 (1992) 96244r.
- 238 Kelly, G.W., Bartle, K.D., Clifford, A.A. and Robinson, R.E.: Application of coupled LC-GC to the analysis of the polar fraction of diesel particulate matter. *J. High Resolut. Chromatogr.*, 15 (1992) 526-530.
- 239 Ma, S., Yuan, H., Yang, M. and Lu, W.: Hydrocarbon group-type analysis of heavy oils by HPLC. In: Hou, X. (Editor), *Proc. Int. Conf. Pet. Refin. Petrochem. Process. 2*, Int. Acad. Publ., Beijing, 1991, pp. 834-839; C.A., 117 (1992) 174702e.

- 240 Nerlich, B., Zobel, H. and Helms, M.: (Hydrocarbon mixtures from liquid-phase hydrogenation of brown coals. Gel permeation chromatography of heterocompound-containing hydrocarbon mixtures). *Erdoel Kohle, Erdgas, Petrochem.*, 45 (1992) 305-307; C.A., 117 (1992) 154283r.
- 241 Qian, K. and Hsu, C.S.: Molecular transformation in hydrotreating processes studied by on-line liquid chromatography/mass spectrometry. *Anal. Chem.*, 64 (1992) 2327-2333.

For additional information see C.A.:

117 (1992) 154129v.

See also 221, 1142.

6. ALCOHOLS

- 242 Barisci, J.N. and Wallace, G.G.: Photoelectrochemical detection of alcohols. *Electroanalysis (N.Y.)*, 4 (1992) 439-445; C.A., 117 (1992) 82695b.
- 243 Caccamese, S. and Maccagnano, M.G.: Liquid chromatographic separation of chiral diarylcarbinols. *Chirality*, 4 (1992) 170-173; C.A., 117 (1992) 163156d.
- 244 Caputi, A., Jr., Christensen, E., Biedenweg, N. and Miller, S.: Liquid chromatographic method for determination of glycerol in wine and grape juice. *J. AOAC Int.*, 75 (1992) 379-383; C.A., 117 (1992) 46854u.
- 245 Hosoya, K., Tanaka, N., Araki, T., Kimata, K., Tsuboi, R., Takabe, K. and Yoda, H.: Molecular recognition of acyclic stereoisomer using a simple high performance liquid chromatography. II. *J. Liq. Chromatogr.*, 15 (1992) 2445-2465.
- 246 Sakakihara, Y. and Kamoshita, S.: Partition of free dolichol in human urine. *Biochem. Cell Biol.*, 70 (1992) 518-521.
- 247 Sánchez Munoz, G., Garcia Romero, E., Pérez Coello, M.S. and Martín Alvarez, P.J.: Determination of glycerol in wines by high-performance liquid chromatography: comparison with enzymatic method. *J. Chromatogr.*, 623 (1992) 43-47.
- 248 Swiezewska, E., Chojnacki, T., Jankowski, W.J., Singh, A.K. and Olsson, J.: The occurrence of long chain polyprenols in leaves of plants of *Rosaceae* family and their isolation by time-extended liquid chromatography. *Biochem. Cell Biol.*, 70 (1992) 448-454.
- 249 Yoshida, T., Moriyama, Y. and Taniguchi, H.: 6-Methoxy-2-methylsulphonylquinoline-4-carbonyl chloride as a fluorescence derivatization reagent for alcohols in high-performance liquid chromatography. *Anal. Sci.*, 8 (1992) 355-359; C.A., 117 (1992) 103300m.

For additional information see C.A.:

117 (1992) 184026b.

See also 216, 324, 420, 1153, 1157, 1158, 1159, 1512.

7. PHENOLS

- 250 Arfelli, G., Chiavari, G., Castellari, M. and Amati, A.: (Effect of vinification procedures on the polyphenol content of wines obtained from different grape cultivars. Note I. Determination of polyphenols in musts and wines by HPLC). *Vignevini*, 19 (1992) 53-58; C.A., 117 (1992) 130079c.
- 251 Goncharov, V.V., Goryunova, V.V. and Tul'chinskii, V.M.: (Isocratic groupwise separation of phenols by HPLC). *Zh. Prikl. Khim. (S.-Peterburg)*, 64 (1991) 2673-2678; C.A., 117 (1992) 96968e.
- 252 Marko-Varga, G. and Barceló, D.: Liquid chromatographic retention and separation of phenols and related aromatic compounds on reversed phase columns. *Chromatographia*, 34 (1992) 146-154.
- 255 Maslowska, J. and Starzynski, S.: (Separation of phenol derivatives by high performance liquid chromatography). *Chem. Anal. (Warsaw)*, 36 (1991) 219-223; C.A., 117 (1992) 39530g.
- 256 Melcher, R.G., Bakke, D.W. and Hughes, G.H.: On-line membrane/liquid chromatographic analyzer for pentachlorophenol and other trace phenols in wastewater. *Anal. Chem.*, 64 (1992) 2258-2262.
- 257 Mellon, F.A.: Liquid chromatography/mass spectrometry and fast atom bombardment mass spectrometry of polyphenols. In: Galletti, G.C. (Editor), *Prod. Util. Lignocellul. [Proc. Workshop]*, 1990, Elsevier, London, 1991, pp. 151-162; C.A., 117 (1992) 48131e.
- 258 Montedoro, G., Servili, M., Baldioli, M. and Miniati, E.: Simple and hydrolyzable phenolic compounds in virgin olive oil. 2. Initial characterization of the hydrolyzable fraction. *J. Agric. Food Chem.*, 40 (1992) 1577-1580.
- 259 Seitz, L.M.: Identification of 5-(2-oxoalkyl)resorcinols and 5-(2-oxoalkenyl)resorcinols in wheat and rye grains. *J. Agric. Food Chem.*, 40 (1992) 1541-1546.
- 260 Wen, Y.H., Wu, S.S., Lin, S.J. and Wu, H.L.: Trace analysis for phenol by derivatization and high performance liquid chromatography. *Zhonghua Yaoxue Zazhi*, 44 (1992) 221-227; C.A., 117 (1992) 184038g.

See also 178, 263, 427, 1294, 1324, 1526, 1531, 1546, 1547.

8. SUBSTANCES CONTAINING HETEROCYCLIC OXYGEN

8a. Flavonoids

- 261 Andriollo, N., Guarini, A. and Cassani, G.: Isolation and characterization of pseudobactin B: a pseudobactin-type siderophore from *Pseudomonas* species strain PD 30. *J. Agric. Food Chem.*, 40 (1992) 1245-1248.

- 262 Davies, A.J. and Mazza, G.: Separation and characterization of anthocyanins of *Monarda fistulosa* by high-performance liquid chromatography. *J. Agric. Food Chem.*, 40 (1992) 1341-1345.
- 263 De Simon, B.F., Perez-Illarbe, J., Hernandez, T., Gomez-Cordoves, C. and Estrella, I.: Importance of phenolic compounds for the characterization of fruit juices. *J. Agric. Food Chem.*, 40 (1992) 1531-1535.
- 264 Gueldner, R.C., Snook, M.E., Widstrom, N.W. and Wiseman, B.R.: TLC screen for maysin, chlorogenic acid and other possible resistance factors to the fall armyworm and the corn earworm in *Zea mays*. *J. Agric. Food Chem.*, 40 (1992) 1211-1213.
- 265 Hertog, M.G.L., Hollman, P.C.H. and Venema, D.P.: Optimization of a quantitative HPLC determination of potentially anticarcinogenic flavonoids in vegetables and fruits. *J. Agric. Food Chem.*, 40 (1992) 1591-1598.
- 266 Iida, J. and Murata, T.: Application of thermospray liquid chromatography/mass spectrometry to analysis of glycosides. *Anal. Sci.*, 7(Suppl., Proc. Int. Congr. Anal. Sci., 1991, Pt. 1) (1991) 963-966; C.A., 117 (1992) 43967x.
- 267 Mizuno, M., Tsuchida, H., Kozukue, N. and Mizuno, S.: Rapid quantitative analysis and distribution of free quercetin in vegetables and fruits. *Nippon Shokuhin Kogyo Gakkaishi*, 39 (1992) 88-92; C.A., 117 (1992) 149598t.
- 268 Proliac, A., Chaboud, A. and Raynaud, J.: Isolement et identification de trois C-glycosylflavones dans les tiges feuillées d'*Arum dracunculus*. *Pharmazie*, 47 (1992) 646-647.
- 269 Smit, G., Puvanesarajah, V., Carlson, R.W., Barbour, W.M. and Stacey, G.: *Bradyrhizobium japonicum* nodD₁ can be specifically induced by soybean flavonoids that do not induce the nodYABCSUIJ operon. *J. Biol. Chem.*, 267 (1992) 310-318.
- 270 Tomas-Barberan, F.A., Gil, M.I., Ferreres, F., Garcia-Viguera, C. and Tomas-Lorente, F.: (Detection of honey in Jijona nougat by HPLC of flavonoids). *Aliment., Equipos Tecnol.*, 11 (1992) 93-95; C.A., 117 (1992) 130038p.

See also 1517, 1518, 1556.

8b. Aflatoxins and other mycotoxins

- 271 Abbas, H.K., Mirocha, C.J., Shier, W. and Gunther, R.: Bioassay, extraction, and purification procedure for wortmannin, the hemorrhagic factor produced by *Fusarium oxysporum* N17-B grown on rice. *J. Assoc. Off. Anal. Chem.*, 75 (1992) 474-480.
- 272 Adensam, L. and Lebedova, M.: (Determination of aflatoxins in peanuts). *Prum. Potravin*, 43 (1992) 277-279; C.A., 117 (1992) 130017f - a review with 4 refs.
- 273 Ansari, A.A. and Shrivastava, A.K.: High performance liquid chromatographic analysis of norsolorinic acid produced by *Aspergillus parasiticus* mutant. *Proc. Indian Natl. Sci. Acad., Part B*, 57 (1991) 403-405; C.A., 116 (1992) 190401r.
- 274 Candlish, A.A.G., Stimson, W.H. and Smith, J.E.: Immunological methods for aflatoxins: collaborative study of immunoaffinity column chromatography method. In: Morgan, M.R.A., Smith, C.J. and Williams, P.A. (Editors), *Food Saf. Qual. Assur.: Appl. Immunoassay Syst., Proc., 1st 1991*, Elsevier, London, 1992, 103-110; C.A., 117 (1992) 88868g.

- 275 Cole, R.O., Holland, R.D. and Sepaniak, M.J.: Factors influencing performance in the rapid separation of aflatoxins by micellar electrokinetic capillary chromatography. *Talanta*, 39 (1992) 1139-1147; C.A., 117 (1992) 165379c.
- 276 Gilbert, J., Boenke, A. and Wagstaffe, P.J.: Deoxynivalenol in wheat and maize flour reference materials. 1. An intercomparison of methods. *Food Addit. Contam.*, 9 (1992) 71-81; C.A., 117 (1992) 130033h.
- 277 Holcomb, M., Wilson, D.M., Trucksess, M.W. and Thompson, H.C., Jr.: Determination of aflatoxins in food products by chromatography. *J. Chromatogr.*, 624 (1992) 341-352 - a review with 97 refs.
- 278 Macho, M.L., Munoz, P. and Eguileor, I.: Aflatoxin M₁ analysis of milk and dairy products of total diet in the Basque country. In: Morgan, M.R.A., Smith, C.J. and Williams, P.A. (Editors), *Food Saf. Qual. Assur.: Appl. Immunoassay Syst., Proc., 1st 1991*, Elsevier, London, 1992, pp. 115-118; C.A., 117 (1992) 110234w.
- 279 Magnarini, C., Cossignani, L. and Damiani, P.: (Aflatoxins and chlorinated solvents in edible oils). *Lab. 2000*, 5 (1991) 138-140; C.A., 117 (1992) 110273h.
- 280 Mueller, M. and Lepom, P.: (Analysis of *Alternaria* mycotoxins in laboratory cultures). *Zentralbl. Mikrobiol.*, 147 (1992) 197-206; C.A., 117 (1992) 107524k.
- 281 Nesheim, S., Stack, M.E., Trucksess, M.W., Eppley, R.M. and Krogh, P.: Rapid solvent-efficient method for liquid chromatographic determination of ochratoxin A in corn, barley, and kidney: collaborative study. *J. AOAC Int.*, 75 (1992) 481-487; C.A., 117 (1992) 46857x.
- 282 Rotunno, T.: Combined Kalman filtering and sheepest sescent minimization for quantitative analysis of unresolved misaligned chromatographic peaks. The resolution of alternariol and altenuisol mycotoxins in mixtures by HPLC. *Chromatographia*, 34 (1992) 56-62.
- 283 Rubin, D. and Laposata, M.: Cellular interactions between *n*-6 and *n*-3 fatty acids: a mass analysis of fatty acid elongation/desaturation, distribution among complex lipids, and conversion to eicosanoids. *J. Lipid Res.*, 33 (1992) 1431-1440.
- 284 Sachse, J.: Identification and determination of mellein in cultures of the fungus *Septoria nodorum* (Berk.) by thin-layer and high-performance liquid chromatography. *J. Chromatogr.*, 609 (1992) 349-353.
- 285 Siemens, K. and Zawistowski, J.: Determination of *Penicillium roqueforti* toxin by reversed-phase high-performance liquid chromatography. *J. Chromatogr.*, 609 (1992) 205-211.
- 286 Sydenham, E.W., Shephard, G.S. and Thiel, P.G.: Liquid chromatographic determination of fumonisins B₁, B₂, and B₃ in foods and feeds. *J. AOAC Int.*, 75 (1992) 313-318; C.A., 117 (1992) 88904r.
- 287 Werner, G.: (Determination of feed aflatoxins following selective clean-up on an immunoaffinity column). *Agribiol. Res.*, 44 (1991) 289-298; C.A., 117 (1992) 24847s.
- 8c. *Other compounds with heterocyclic oxygen (incl. tannins)*
- 288 Evans, J.M., Smith, R.J. and Stemp, G.: Separation of the enantiomers of some potassium channel activators using an α -acid glycoprotein column. *J. Chromatogr.*, 623 (1992) 163-167.
- 289 Guerra-Hernández, E., García-Villanova, B. and Montilla-Gómez, J.: Determination of hydroxymethylfurfural in baby cereals by high performance liquid chromatography. *J. Liq. Chromatogr.*, 15 (1992) 2551-2559.
- 290 Hargitai, T., Reinholdsson, P. and Sandstroem, J.: Chromatographic enantiomer separation, circular dichroism spectrum, and absolute configuration of the 7-methoxycoumarin syn head-to-tail dimer. *Acta Chem. Scand.*, 45 (1991) 1076-1079; C.A., 117 (1992) 25810m.
- 291 Okamoto, M. and Nakazawa, H.: Direct chromatographic separation of the enantiomers of methyl jasmonate and its derivatives. *Biosci., Biotechnol., Biochem.*, 56 (1992) 1172-1173; C.A., 117 (1992) 166916n.
- 292 Trumble, J.T., Millar, J.G., Ott, D.E. and Carson, W.C.: Seasonal patterns and pesticidal effects on the phototoxic linear furanocoumarins in celery, *Apium graveolens* L. *J. Agric. Food Chem.*, 40 (1992) 1501-1506.
- 293 Yamagami, C. and Takao, N.: Hydrophobicity parameters determined by reversed-phase liquid chromatography. V. Relationship between the capacity factor and the octanol-water partition coefficient for simple heteroaromatic compounds and their ester derivatives. *Chem. Pharm. Bull.*, 40 (1992) 925-929.
- For additional information see C.A.:
117 (1992) 46850q, 149584k.
- See also 86, 262, 1292, 1305, 1306, 1481, 1509, 1532.
9. OXO COMPOUNDS, ETHERS, EPOXIDES AND QUINONES
- 294 Blee, E. and Schuber, F.: Enantioselectivity of the hydrolysis of linoleic acid monoepoxides catalyzed by soybean fatty acid epoxide hydrolase. *Biochem. Biophys. Res. Commun.*, 187 (1992) 171-177.
- 295 Chiavari, G., Torsi, G. and Asmundsdottir, A.M.: Different methods for HPLC analysis for aldehydes in aqueous solutions. *Ann. Chim. (Rome)*, 82 (1992) 349-356; C.A., 117 (1992) 163177m.
- 296 Horti, A., Glibin, E. and Nesterov, V.: Retention behaviour of crown ethers and actinomycin D in reversed-phase HPLC. *Chromatographia*, 34 (1992) 155-158.
- 297 Ladde, A. and Mueller, C.: (Isolation of ubiquinones from extracts of waste biomass via column chromatography). *Chem. Tech. (Leipzig)*, 44 (1992) 175-177; C.A., 117 (1992) 174004d.
- 298 Popa, I.M. and Savin, A.: Analysis of phenyl-glycidyl-ether by high-pressure liquid chromatography. *Bul. Inst. Politeh. Iasi, Sect. 2: Chim. Ing. Chim.*, 36 (1990) 31-34; C.A., 117 (1992) 163162c.
- 299 Roeper, K. and Zahn, H.: (Quantitative determination of glutaric aldehyde among formaldehyde in collagen samples. Part 2. Analysis of collagen samples treated with glutaraldehyde via reaction with 2,4-dinitrophenyl hydrazine (DNPH)). *Leder*, 43 (1992) 117-133; C.A., 117 (1992) 153130b.
- 300 Tsai, T.H., Chen, C.M. and Chen, C.F.: Disposition of asarone after intravenous administration to rabbits assessed using HPLC. *J. Pharm. Pharmacol.*, 44 (1992) 620-622; C.A., 117 (1992) 103469y.

- 301 Tsai, Y., Aoki, T., Maruta, H., Abe, H., Sakagami, H., Hatano, T., Okuda, T. and Tanuma, S.: Mouse mammary tumor virus gene expression is suppressed by oligomeric ellagitannins, novel inhibitors of poly(ADP-ribose) glycohydrolase. *J. Biol. Chem.*, 267 (1992) 14436-14442.
- 302 Zhang, D., Fan, C., Yu, W., Yu, B., Zhang, H. and Wang, Y.: (Study of ubiquinone-10 content in the myocardium of patients with Keshan disease and its effect on protection of the myocardium by high performance liquid chromatography). *Sepu*, 10 (1992) 115-117; C.A., 117 (1992) 65799w.
- 303 Zonta, F., Bogoni, P. and Masoti, P.: (Determination of carbonyl compounds in beverages and spirits by HPLC). *Ind. Bevande*, 21 (1992) 273-276; C.A., 117 (1992) 130085b.
- For additional information see C.A.:
117 (1992) 33803b.
- See also 267, 289, 584, 893, 1136, 1149, 1542, 1543.
10. CARBOHYDRATES
- 10a. Mono and oligosaccharides. Structural studies
- 304 Altmann, F.: Determination of amino sugars and amino acids in glycoconjugates using precolumn derivatization with *o*-phthalaldehyde. *Anal. Biochem.*, 204 (1992) 215-219.
- 305 Bugner, E. and Feinberg, M.: Determination of mono- and disaccharides in foods by interlaboratory study: quantitation of bias components for liquid chromatography. *J. AOAC Int.*, 75 (1992) 443-464; C.A., 117 (1992) 68598r.
- 306 Cheng, C., Chen, J.-T. and Huang, H.-R.: High-performance liquid chromatography of carbohydrates in enzymic hydrolysate of waste paper. *Chromatographia*, 34 (1992) 534-536.
- 307 Clonis, Y.A.: High-performance affinity chromatography for protein separation and purification. *Methods Mol. Biol. (Totowa)*, 11(Pract. Protein Chromatogr.) (1992) 105-123; C.A., 117 (1992) 146392k - a review with 15 refs.
- 308 Conboy, J.J. and Henion, J.: High-performance anion-exchange chromatography coupled with mass spectrometry for the determination of carbohydrates. *Biol. Mass Spectrom.*, 21 (1992) 397-407; C.A., 117 (1992) 127763r.
- 309 Damm, J.B.L., Overkluft, G.T., Vermeulen, B.W.M., Fluitsma, C.F. and van Dedem, G.W.K.: Separation of natural and synthetic heparin fragments by high-performance capillary electrophoresis. *J. Chromatogr.*, 608 (1992) 297-309.
- 310 Davies, M., Smith, K.D., Harbin, A.-M. and Hounsell, E.F.: High-performance liquid chromatography of oligosaccharide alditols and glycopeptides on a graphitized carbon column. *J. Chromatogr.*, 609 (1992) 125-131.
- 311 Del Nozal, M.J., Bernal, J.L., Gomez, F.J., Antolin, A. and Toribio, L.: Post-column derivatization of carbohydrates with ethanolamine-boric acid prior to their detection by high-performance liquid chromatography. *J. Chromatogr.*, 607 (1992) 191-198.
- 312 Grönberg, G., Lipniunas, P., Lundgren, T., Lindh, F. and Nilsson, B.: Structural analysis of five new monosialylated oligosaccharides from human milk. *Arch. Biochem. Biophys.*, 296 (1992) 597-610.
- 313 Gu, J., Meng, X., Tang, H. and Chen, H.: (High-performance liquid chromatographic determination of N-acetylglucosaminyltransferase III during induced hepatocarcinoma in rat with N-nitrosodiethylamine). *Shengwu Huaxue Yu Shengwu Wuli Xuebao*, 24 (1992) 48-54; C.A., 117 (1992) 165335k.
- 314 Haynes, P.A., Batley, M., Peach, R.J., Brennan, S.O. and Redmond, J.W.: Characterisation of oligosaccharides from a glycoprotein variant of human serum albumin (albumin Casebrook) using high-performance anion-exchange chromatography and nuclear magnetic resonance spectroscopy. *J. Chromatogr.*, 581 (1992) 187-193.
- 315 Heikkilä, H., Hyöky, G., Niittymäki, P., Viljava, T. and Myohanen, T.: Production of glucose and fructose from sucrose. *PCT Int. Appl. WO 92 07,097 (Cl. C13K3/00)*, 30 Apr. 1992, FI Appl. 90/5,069, 15 Oct. 1990; 24 p.; C.A., 117 (1992) 29136u.
- 316 Iwahara, S., Maeyama, T., Mishima, T., Jikibara, T., Takegawa, K. and Iwamoto, H.: Studies on the uronic acid-containing glycoproteins of *Fusarium* sp. M7-1: IV. Isolation and identification of four novel oligosaccharide units derived from the acidic polysaccharide chain. *J. Biochem. (Tokyo)*, 112 (1992) 355-359.
- 317 Jiang, M., Passaniti, A., Penno, M.B. and Hart, G.W.: Adrenal carcinoma tumor progression and penultimate cell surface oligosaccharides. *Cancer Res.*, 52 (1992) 2222-2227.
- 318 Klein, A., Carnoy, C., Wieruszkeski, J.-M., Strecker, G., Strang, A.-M., van Halbeek, H., Roussel, P. and Lamblin, G.: The broad diversity of neutral and sialylated oligosaccharides derived from human salivary mucins. *Biochemistry*, 31 (1992) 6152-6165.
- 319 Kokoh, K.B., Leger, J.M., Beden, B., Huser, H. and Lamy, C.: "On line" chromatographic analysis of the products resulting from the electrocatalytic oxidation of D-glucose on pure and adatoms modified platinum and gold electrodes. *Electrochim. Acta*, 37 (1992) 1909-1918; C.A., 117 (1992) 159525f.
- 320 Matsuhashi, S. and Inoue, S. and Hatanaka, C.: Simultaneous measurement of the galacturonate and neutral sugar contents of pectic substances by an enzymic-HPLC method. *Biosci., Biotechnol., Biochem.*, 56 (1992) 1053-1057; C.A., 117 (1992) 169688a.
- 321 Muzquiz, M., Rey, C., Cuadrado, C. and Fenwick, G.R.: Effect of gemination on the oligosaccharide content of lupin species. *J. Chromatogr.*, 607 (1992) 349-352.
- 322 Niessen, W.M.A., van der Hoeven, R.A.M., van der Greef, J., Schols, H.A., Lucas-Lokhorst, G., Voragen, A.G.J. and Bruggink, C.: High-performance anion-exchange chromatography/thermospray mass spectrometry in the analysis of oligosaccharides. *Rapid Commun. Mass Spectrom.*, 6 (1992) 474-478; C.A., 117 (1992) 86318d.
- 323 Pascale, M.C., Malagolini, N., Serafini-Cessi, F., Migliaccio, G., Leone, A. and Bonatti, S.: Biosynthesis and oligosaccharide structure of human CD8 glycoprotein expressed in a rat epithelial cell line. *J. Biol. Chem.*, 267 (1992) 9940-9947.
- 324 Rabinowitz, I.: Chromatographic method of purifying cyclitols. U.S. US 5,096,594 (Cl. 210-656; B01D15/08), 17 Mar. 1992, Appl. 634,096, 20 Dec. 1990; 5 p.; C.A., 117 (1992) 82711d.
- 325 Rakotomanga, S., Baillet, A., Pellerin, F. and Baylocq-Ferrier, D.: Liquid chromatographic analysis of monosaccharides with phenylisocyanate derivatization. *J. Pharm. Biomed. Anal.*, 10 (1992) 587-591.
- 326 Schomburg, G., Deege, A., Hinrichs, H., Hübinger, E. and Husmann, H.: Preparation, purification, and analysis of alkylated cyclodextrins. *J. High Resolut. Chromatogr.*, 15 (1992) 597-584.

- 327 Scott, F.W.: HPLC determination of carbohydrates in foods. *Food Sci. Technol.*, 52(Food Anal. HPLC) (1992) 259-274; *C.A.*, 117 (1992) 110174b - a review with 59 refs.
- 328 Solbrig-Lebuhn, H.: (Carbohydrate analysis. Anion-exchange chromatography with pulsed electrochemical detection (HPAEPAD)). *LaborPraxis*, 16 (1992) 786-789; *C.A.*, 117 (1992) 173689u - a review with 2 refs.
- 329 Spiro, M.J. and Spiro, R.G.: Monosaccharide determination of glycoconjugates by reverse-phase high-performance liquid chromatography of their phenylthiocarbonyl derivatives. *Anal. Biochem.*, 204 (1992) 152-157.
- 330 Strecker, G., Wieruszkeski, J.-M., Michalski, J.-C. and Montreuil, J.: ¹H and ¹³C-NMR spectroscopy of 2-oxo-3-deoxy-D-glycero-D-galactononulosonic acid-containing oligosaccharide-alditols bearing Lewis X, Lewis Y and 1-Lewis Y determinants isolated from the jelly coat of *Pleurodeles waltl* eggs. *Biochem. J.*, 287 (1992) 905-909.
- 331 Tanimoto, T., Tsujikawa, J. and Koizumi, K.: Preparation of three positional isomers of diglucosyl-cyclomaltohexaose. *Chem. Pharm. Bull.*, 40 (1992) 1125-1129.
- 332 Tomiya, N., Suzuki, T., Awaya, J., Mizuno, K., Matsubara, A., Nakano, K. and Kurono, M.: Determination of monosaccharides and sugar alcohols in tissues from diabetic rats by high-performance liquid chromatography with pulsed amperometric detection. *Anal. Biochem.*, 206 (1992) 98-104.
- 333 Urashima, T., Messer, M. and Bubbs, W.A.: Biosynthesis of marsupial milk oligosaccharides II: characterization of a β⁶-N-acetylglucosaminyltransferase in lactating mammary glands of the tammar wallaby, *Macropus eugenii*. *Biochim. Biophys. Acta*, 1117 (1992) 223-231.
- 334 Van de Merbel, N.C., Kool, I.M., Longeman, H., Brinkman, U.A.T., Kolhorn, A. and de Rijke, L.C.: On-line monitoring of fermentation processes by ultrafiltration and column liquid chromatography. *Chromatographia*, 33 (1992) 525-532.
- 335 Wang, K.: (Some progresses in structural research of carbohydrate). *Shengwu Huaxue Yu Shengwu Wuli Jinzhan*, 18 (1991) 405-408; *C.A.*, 117 (1992) 65702j - a review with 11 refs.
- 336 Watson, E., Bhide, A., Kenney, W.C. and Lin, F.-K.: High-performance anion-exchange chromatography of asparagine-linked oligosaccharides. *Anal. Biochem.*, 205 (1992) 90-95.
- 337 White, D.R., Jr. and Cancalon, P.F.: Detection of beet sugar adulteration of orange juice by liquid chromatography/pulsed amperometric detection with column switching. *J. AOAC Int.*, 75 (1992) 584-587; *C.A.*, 117 (1992) 46864x.
- 338 Yamada, S., Yoshida, K., Sugiura, M. and Sugahara, K.: One- and two-dimensional ¹H-NMR characterization of two series of sulfated disaccharides prepared from chondroitin sulfate and heparan sulfate/heparin by bacterial eliminase digestion. *J. Biochem. (Tokyo)*, 112 (1992) 440-447.
- For additional information see *C.A.*:
117 (1992) 88836v, 127802c.
- See also 90, 113, 247, 339, 344, 353, 371, 380, 381, 382, 393, 396, 987, 1000, 1002, 1037, 1153, 1178, 1525.
- 10b. *Polysaccharides, mucopolysaccharides, lipopolysaccharides*
- 339 Barry, F.P., Gaw, J.U., Young, C.N. and Neame, P.J.: Hyaluronan-binding region of aggrecan from pig laryngeal cartilage. Amino acid sequence, analysis of N-linked oligosaccharides and location of the keratan sulphate. *Biochem. J.*, 286 (1992) 761-769.
- 340 Berth, G.: Methodical aspects of characterization of alginate and pectate by light scattering and viscometry coupled with GPC. *Carbohydr. Polym.*, 19 (1992) 1-9; *C.A.*, 117 (1992) 13325n.
- 341 Calatroni, A., Vinci, R. and Ferlazzo, A.M.: Characteristics of the interactions between acid glycosaminoglycans and proteins in normal human plasma as revealed by the behaviour of the protein-polysaccharide complexes in ultrafiltration and chromatographic procedures. *Clin. Chim. Acta*, 206 (1992) 167-180.
- 342 Cooper, S., Pera, M.F., Bennett, W. and Finch, J.T.: A novel keratan sulphate proteoglycan from a human embryonal carcinoma cell line. *Biochem. J.*, 286 (1992) 959-966.
- 343 De Briel, D., Couderc, F., Riegel, P., Jehl, F. and Minck, R.: High-performance liquid chromatography of corynomycolic acids as a tool in identification of *Corynebacterium* species and related organisms. *J. Clin. Microbiol.*, 30 (1992) 1407-1417; *C.A.*, 117 (1992) 43756c.
- 344 Dohi, T., Nishikawa, A., Ishizuka, I., Totani, M., Yamaguchi, K., Nakagawa, K., Saitoh, O., Ohshiba, S. and Oshima, M.: Substrate specificity and distribution of UDP-GalNAc:sialylparagloboside N-acetylgalactosaminyltransferase in the human stomach. *Biochem. J.*, 288 (1992) 161-165.
- 345 Flannery, C., Stanescu, V., Mörgelein, M., Boynton, R., Gordy, J. and Sandy, J.: Variability in the G3 domain content of bovine aggrecan from cartilage extracts and chondrocyte cultures. *Arch. Biochem. Biophys.*, 297 (1992) 52-60.
- 346 Galis, Z.S., Alavi, M.Z. and Moore, S.: Sulfated proteoglycans of rabbit aorta: selective extraction and alternative method for glycosaminoglycan. *Anal. Biochem.*, 204 (1992) 390-397.
- 347 Hermentin, P. and Seidat, J.: Microscale analysis of N-acetylneuraminic acid. *GBF Monogr.*, 15(Protein Glycosylation: Cell, Biotechnol. Anal. Aspects) (1991) 185-188; *C.A.*, 117 (1992) 166906j.
- 348 Hu, W.-L. and Regoeczi, E.: Hepatic heparan sulphate proteoglycan and the recycling of transferrin. *Biochem. Cell Biol.*, 70 (1992) 535-538.
- 349 Huber, A., Schurz, J., Praznik, W. and Hinterstoisser, B.: (Size-exclusion chromatography/small-angle laser light scattering: application to polysaccharides). *Papier (Darmstadt)*, 45 (1991) 735-741; *C.A.*, 117 (1992) 28994k.
- 350 Ishihara, M., Kiefer, M.C., Barr, P.J., Guo, Y. and Swiedler, S.J.: Selection of COS cell mutants defective in the biosynthesis of heparan sulfate proteoglycan. *Anal. Biochem.*, 206 (1992) 400-407.
- 351 Kragten, E.A., Kamerling, J.P. and Vliegenterth, F.G.: Composition analysis of carboxymethylcellulose by high-pH anion-exchange chromatography with pulsed amperometric detection. *J. Chromatogr.*, 623 (1992) 49-53.
- 352 Kravtchenko, T.P., Berth, G., Voragen, A.G.J. and Pilnik, W.: Studies on the intermolecular distribution of industrial pectins by means of preparative size exclusion chromatography. *Carbohydr. Polym.*, 18 (1992) 253-263; *C.A.*, 117 (1992) 113915e.

- 353 Lindholt, K. and Lindahl, U.: Biosynthesis of heparin. The D-glucuronosyl- and N-acetyl-D-glucosaminyltransferase reactions and their relation to polymer modification. *Biochem. J.*, 287 (1992) 21-29.
- 354 Magnus, J.H., Stenstad, T., Husby, G. and Kolset, S.O.: Isolation and partial characterization of heparan sulphate proteoglycans from human hepatic amyloid. *Biochem. J.*, 288 (1992) 255-231.
- 355 Matsuda, F.: (Application of chromatographic separation technique to starch sweetener). *Gekkan, Fudo Kemikaru*, 8 (1992) 80-85; C.A., 117 (1992) 169654m - a review with 1 ref.
- 356 McBain, J.A. and Mueller, G.C.: Release of cell surface proteoglycans from differentiating colon cells proceeds by cleavage of lipophilic anchor peptides. *Biochem. J.*, 287 (1992) 131-140.
- 357 Nakanishi, H., Oguri, K., Yoshida, K., Itano, N., Takenaga, K., Kazama, T., Yoshida, A. and Okayama, M.: Structural differences between heparan sulphates of proteoglycan involved in the formation of basement membranes *in vivo* by Lewis-lung-carcinoma-derived cloned cells with different metastatic potentials. *Biochem. J.*, 288 (1992) 215-224.
- 358 Ng, C.K., Handley, C.J., Preston, B.N. and Robinson, H.C.: The extracellular processing and catabolism of hyaluronan in cultured adult articular cartilage explants. *Arch. Biochem. Biophys.*, 298 (1992) 70-79.
- 359 Rollings, J.E.: Use of online laser light scattering coupled to chromatographic separations for the determination of molecular weight, branching, size and shape distributions of polysaccharides. *Spec. Publ.-R. Soc. Chem.*, 99(Laser Light Scattering Biochem.) (1992) 275-293; C.A., 117 (1992) 86005z.
- 360 Schmidtchen, A. and Fransson, L.-A.: Analysis of glycosaminoglycan chains from different proteoglycan populations in human embryonic skin fibroblasts. *Eur. J. Biochem.*, 208 (1992) 537-546.
- 361 Seibel, M.J., Macaulay, W., Jelsma, R., Saed-Nejad, F. and Ratcliffe, A.: Antigenic properties of keratan sulfate: influence of antigen structure, monoclonal antibodies, and antibody valency. *Arch. Biochem. Biophys.*, 296 (1992) 410-418.
- 362 Shioda, K.: Chromatographic separation process [starch hydrolysis products]. In: Schenck, F.W. and Hebeda, R.E. (Editors), *Starch Hydrolysis Prod.*, VCH, New York, 1992, pp. 555-572; C.A., 117 (1992) 169642f - a review with 19 refs.
- 363 Tai, G.-H., Morris, H.G., Brown, G.M., Huckerby, T.N. and Niedzynski, I.A.: A sub-population of keratan sulphates derived from bovine articular cartilage is capped with $\alpha(2-6)$ -linked N-acetylneuraminic acid residues. Affinity chromatography using immobilized *Sambucus nigra* lectin and characterization using ^1H NMR spectroscopy. *Biochem. J.*, 286 (1992) 231-234.
- 364 Takeuchi, Y., Yanagishita, M. and Hascall, V.C.: Effects of MgCl_2 on the release and recycling of heparan sulfate proteoglycans in a rat parathyroid cell line. *Arch. Biochem. Biophys.*, 298 (1992) 371-379.
- 365 Theander, O.: Chemical characterization of some potential lignocellulosic raw materials in the swedish Agro-Fiber project. In: Galletti, G.C. (Editor), *Prod. Util. Lignocellul.*, 1990, Elsevier, London, 1991, pp. 49-61; C.A., 117 (1992) 152963p.
- 366 Vardanis, A.: Particulate glycogen of mammalian liver: specificity in binding phosphorylase and glycogen synthase. *Biochem. Cell Biol.*, 70 (1992) 523-527.
- 367 Zhou, R.: (GPC-LALLS in the measurement of molecular weight and its distribution of dextran). *Zhongguo Yaoxue Zazhi*, 27 (1992) 159-162; C.A., 117 (1992) 76579a.

For additional information see C.A.:
117 (1992) 113928m.

See also 328, 1478.

10c. Glycoproteins and their constituents

- 368 Aoki, T., Takasaki, T., Miura, H. and Watabe, H.: Development of radioimmunoassay of intact laminin and its clinical evaluation as a tumor marker. *Chem. Pharm. Bull.*, 40 (1992) 981-985.
- 369 Bhattacharya, L. and Brewer, C.F.: Formation of homogeneous carbohydrate-lectin cross-linked precipitates from mixtures of D-galactose/N-acetyl-D-galactosamine-specific lectins and multiantennary galactosyl carbohydrates. *Eur. J. Biochem.*, 208 (1992) 179-185.
- 370 Bihoreau, N., Fontaine-Aupart, M.-P., Lehegarat, A., Desmadril, M. and Yon, J.M.: First determination of the secondary structure of purified factor VIII light chain. *Biochem. J.*, 288 (1992) 35-40.
- 371 Björk, I., Ylänjärvi, K., Olson, S.T., Hermentin, P., Conradt, H.S. and Zettlmeissl, G.: Decreased affinity of recombinant anti-thrombin for heparin due to increased glycosylation. *Biochem. J.*, 286 (1992) 793-800.
- 372 Cantor, A.B. and Kornfeld, S.: A method for ^3H mannose labeling of Asn-linked oligosaccharides on recombinant glycoproteins synthesized in *Xenopus* oocytes. *Anal. Biochem.*, 205 (1992) 220-226.
- 373 Chen, T.-F., Yu, H. and Barofsky, F.: Centrifugal size-exclusion chromatographic method for rapid desalting and filtering of carbohydrate samples prior to fast atom bombardment mass spectrometry. *Anal. Chem.*, 64 (1992) 2014-2019.
- 374 De Jonge, B.L.M., Chang, Y., Gage, D. and Tomasz, A.: Peptidoglycan composition in heterogenous Tn551 mutants of a methicillin-resistant *Staphylococcus aureus* strain. *J. Biol. Chem.*, 267 (1992) 11255-11259.
- 375 De Jonge, B.L.M., Chang, Y., Gage, D. and Tomasz, A.: Peptidoglycan composition of a highly methicillin-resistant *Staphylococcus aureus* strain. The role of penicillin binding protein 2A. *J. Biol. Chem.*, 267 (1992) 11248-11254.
- 376 Do, S.I. and Cummings, R.D.: The hamster transferrin receptor contains Ser/Thr-linked oligosaccharides: use of a lectin-resistant CHO cell line to identify glycoproteins containing these linkages. *J. Biochem. Biophys. Methods*, 24 (1992) 153-165; C.A., 117 (1992) 67163h.
- 377 Fu, D. and van Halbeek, H.: N-Glycosylation site mapping of human serotransferrin by serial lectin affinity chromatography, fast atom bombardment-mass spectrometry, and ^1H nuclear magnetic resonance spectroscopy. *Anal. Biochem.*, 206 (1992) 53-63.
- 378 Gonzales, J., Takao, T., Hori, H., Besada, V., Rodriguez, R., Pardon, G. and Shimonishi, Y.: A method for determination of N-glycosylation sites in glycoproteins by collision-induced dissociation analysis in fast atom bombardment mass spectrometry: identification of the positions of carbohydrate-linked asparagine in recombinant α -amylase by treatment with peptide-N-glycosidase F in ^{18}O -labeled water. *Anal. Biochem.*, 205 (1992) 151-158.

- 379 Hageman, J.H. and Kuehn, G.D.: Boronic acid matrixes for the affinity purification of glycoproteins and enzymes. *Methods Mol. Biol. (Totowa)*, 11(Pract. Protein Chromatogr.) (1992) 45-71; *C.A.*, 117 (1992) 146449J.
- 380 Hard, K., van Zadelhoff, G., Moonen, P., Kamerling, J.P. and Vliegthart, J.F.G.: The Asn-linked carbohydrate chains of human Tamm-Horsfall glycoprotein of one male. Novel sulfated and novel N-acetylgalactosamine-containing N-linked carbohydrate chains. *Eur. J. Biochem.*, 209 (1992) 895-915.
- 381 Hayase, T., Rice, K.G., Dziegielewska, K.M., Kuhlenschmidt, M., Reilly, T. and Lee, Y.C.: Comparison of N-glycosides of fetuins from different species and human α_2 -HS-glycoprotein. *Biochemistry*, 31 (1992) 4915-4921.
- 382 Hermentin, P., Witzel, R., Doenges, R., Bauer, R., Haupt, H., Patel, T., Parekh, R.B. and Brazel, D.: The mapping of high-pH anion-exchange chromatography with pulsed amperometric detection and capillary electrophoresis of the carbohydrate moieties of human plasma α_1 -acid glycoprotein. *Anal. Biochem.*, 206 (1992) 419-429.
- 383 Kato, A., Kameyama, K. and Takagi, T.: Molecular weight determination and compositional analysis of dextran-protein conjugates using low-angle laser light scattering technique combined with high-performance gel chromatography. *Biochim. Biophys. Acta*, 1159 (1992) 22-28.
- 384 Kim, D.H., Kanno, C. and Mizokami, Y.: Purification and characterization of major glycoproteins, PAS-6 and PAS-7, from bovine milk fat globule membrane. *Biochim. Biophys. Acta*, 1122 (1992) 203-211.
- 385 Konami, Y., Yamamoto, K., Osawa, T. and Irimura, T.: The primary structure of the *Cytisus scoparius* seed lectin and a carbohydrate-binding peptide. *J. Biochem. (Tokyo)*, 112 (1992) 366-375.
- 386 Muldbjerg, M., Schousboe, I. and Halkier, T.: Purification of bovine histidine-rich glycoprotein. *Thromb. Res.*, 65 (1992) 815-819; *C.A.*, 117 (1992) 44025a.
- 387 Nakada, H., Inoue, M., Numata, Y., Tanaka, N., Funakoshi, I., Fukui, S. and Yamashita, I.: Cancer-associated glycoproteins defined by a monoclonal antibody, MLS 128, recognized the Th antigen. *Biochem. Biophys. Res. Commun.*, 187 (1992) 217-224.
- 388 Nishimura, H., Takao, T., Hase, S., Shimonishi, Y. and Iwanaga, S.: Human factor IX has a tetrasaccharide O-glycosidically linked to serine 61 through the fucose residue. *J. Biol. Chem.*, 267 (1992) 17520-17525.
- 389 Noguchi, S. and Nakano, M.: Structure of the acidic N-linked carbohydrate chains of the 55-kDa glycoprotein family (PZP3) from porcine zona pellucida. *Eur. J. Biochem.*, 209 (1992) 883-894.
- 390 Nuck, R., Orthen, B. and Reutter, W.: Occurrence of α 1-2-fucosylation in membrane glycoproteins of Morris hepatoma 7777 but not in liver. Aberrant type of fucosylation in a malignant tissue. *Eur. J. Biochem.*, 208 (1992) 669-676.
- 391 Paul, E., Hurtubise, Y. and LeBel, D.: Purification and characterization of the apical plasma membrane of the rat pancreatic acinar cell. *J. Membr. Biol.*, 127 (1992) 129-137; *C.A.*, 117 (1992) 65791n.
- 392 Reddy, M.S., Bobek, L.A., Haraszthy, G.G., Biesbrock, A.R. and Levine, M.J.: Structural features of the low-molecular-mass human salivary mucin. *Biochem. J.*, 287 (1992) 639-643.
- 393 Rice, K.G., Takahashi, N., Namiki, Y., Tran, A.D., Lisi, P.J. and Lee, Y.C.: Quantitative mapping of the N-linked sialyloligosaccharides of recombinant erythropoietin: combination of direct high-performance anion-exchange chromatography and 2-aminopyridine derivatization. *Anal. Biochem.*, 206 (1992) 278-287.
- 394 Ruffet, E., Paquet, N., Frutiger, S., Hughes, G.J. and Jaton, J.-C.: Structural and electron-microscopic studies of jacalin from jackfruit (*Artocarpus integrifolia*) show that this lectin is a 65 kDa tetramer. *Biochem. J.*, 286 (1992) 131-134.
- 395 Shirota, O., Rice, D. and Novotny, M.: High-sensitivity analysis of sialyl-oligosaccharide glycosylation sites in glycoproteins by miniaturized tryptic digestion and microcolumn liquid chromatography. *Anal. Biochem.*, 205 (1992) 189-192.
- 396 Suzuki, S., Kakehi, K. and Honda, S.: Two-dimensional mapping of N-glycosidically linked asialo-oligosaccharides from glycoproteins as reductively pyridylaminated derivatives using dual separation modes of high-performance capillary electrophoresis. *Anal. Biochem.*, 205 (1992) 227-236.
- 397 Townsend, R.R.: Assessment of protein glycosylation using high-pH anion-exchange chromatography with pulsed electrochemical detection. *GBF Monogr.*, 15(Protein Glycosylation: Cell, Biotechnol. Anal. Aspects) (1991) 147-160; *C.A.*, 117 (1992) 166905h.
- 398 Tulsiani, D.R.P. and Touster, O.: Evidence that swainsonine pretreatment of rats leads to the formation of autophagic vacuoles and endosomes with decreased capacity to mature to, or fuse with, active lysosomes. *Arch. Biochem. Biophys.*, 296 (1992) 556-561.
- 399 Warburton, M.J., Kimbell, R., Ferns, S.A., Hayman, A.R., Perusinghe, N. and Monaghan, P.: Characterisation of chondroitin/dermatan sulphate proteoglycans synthesised by rat mammary myoepithelial and fibroblastic cell lines. *Biochim. Biophys. Acta*, 1117 (1992) 291-300.
- 400 Wilson, M.R. and Easterbrook-Smith, S.B.: Clusterin binds by a multivalent mechanism to the Fc and Fab regions of IgG. *Biochim. Biophys. Acta*, 1159 (1992) 319-326.
- 401 Yanagishita, M.: Metabolic labeling of glycosylphosphatidylinositol-anchor of heparan sulfate proteoglycans in rat ovarian granulosa cells. *J. Biol. Chem.*, 267 (1992) 9499-9504.
- 402 Yip, T.T. and Hutchens, T.W.: Metal ion affinity adsorption of a zinc(II)-transport protein present in maternal plasma during lactation: structural characterization and identification as histidine-rich glycoprotein. *Protein Expression Purif.*, 2 (1991) 355-362; *C.A.*, 117 (1992) 22608r.

See also 90, 310, 665, 827, 1042.

11. ORGANIC ACIDS AND LIPIDS

11a. Organic acids and simple esters

- 403 Allenmark, S. and Andersson, S.: Some mechanistic aspects on chiral discrimination of organic acids by immobilized bovine serum albumin (BSA). *Chirality*, 4 (1992) 24-29; *C.A.*, 117 (1992) 107515h.

- 404 Allenmark, S. and Ohlsson, A.: Enantioselectivity of lipase-catalyzed hydrolysis of some 2-chloroethyl 2-arylpropanoates studied by chiral reversed-phase liquid chromatography. *Chirality*, 4 (1992) 98-102; C.A., 117 (1992) 86086b.
- 405 Antoine, C., Murphy, R.C., Henson, P.M. and MacIouf, J.: Time-dependent utilization of platelet arachidonic acid by the neutrophil in formation of 5-lipoxygenase products in platelet-neutrophil co-incubations. *Biochim. Biophys. Acta*, 1128 (1992) 139-146.
- 406 Bhuiyan, A.K.M.J., Jackson, S., Turnbull, D.M., Aynsley-Green, A., Leonard, J.V. and Bartlett, K.: The measurement of carnitine and acyl-carnitines: application to the investigation of patients with suspected inherited disorders of mitochondrial fatty acid oxidation. *Clin. Chim. Acta*, 207 (1992) 185-204.
- 407 Blanco Gomis, D.: HPLC analysis of organic acids. *Food Sci. Technol.*, 52(Food Anal. HPLC) (1992) 371-385; C.A., 117 (1992) 110176d - a review with 59 refs.
- 408 Brenes-Balbuena, M., Garcia-Garcia, P. and Garrido-Fernandez, A.: Phenolic compounds related to the black color formed during the processing of ripe olives. *J. Agric. Food Chem.*, 40 (1992) 1192-1196.
- 409 Duprat, F. and Coyard, V.: Partial separation of lactic acid enantiomers by HPLC using (S)-(+)-mandelic acid as chiral mobile-phase additive. *Chromatographia*, 34 (1992) 31-35.
- 410 Frimmel, F.H., Gremm, T. and Huber, S.: Liquid chromatographic characterization of refractory organic acids. *Sci. Total Environ.*, 117-118 (1992) 197-206; C.A., 117 (1992) 55504g.
- 411 Furuta, T. and Kihara, K.: Relationship between the effects on bactericidal activity of selected disinfectants and the hydrophobic characters of dibasic acid diesters. *Chem. Pharm. Bull.*, 40 (1992) 1309-1312.
- 412 Gerard, H.C., Osman, S.F., Fett, W.F. and Moreau, R.A.: Separation, identification and quantification of monomers from cutin polymers by high performance liquid chromatography and evaporative light scattering detection. *Phytochem. Anal.*, 3 (1992) 139-144; C.A., 117 (1992) 146444d.
- 413 Glasgow, W.C., Afshari, C.A., Barrett, J.C. and Eling, T.E.: Modulation of the epidermal growth factor mitogenic response by metabolites of linoleic and arachidonic acid in Syrian hamster embryo fibroblasts. Differential effects in tumor suppressor gene (+) and (-) phenotypes. *J. Biol. Chem.*, 267 (1992) 10771-10779.
- 414 Goodall, D.M., Wu, Z. and Lisseter, S.G.: Examination of fruit juices and ciders for adulteration by analysis of (-)-malic acid using high-performance liquid chromatography with dual ultraviolet/polarimetric detection. *Anal. Proc. (London)*, 29 (1992) 238-240; C.A., 117 (1992) 110270e.
- 415 Habib, A. and MacIouf, J.: Comparison of leukotriene A₄ metabolism into leukotriene C₄ by human platelets and endothelial cells. *Arch. Biochem. Biophys.*, 298 (1992) 544-552.
- 416 Henderson, W.R., Jr., Rashed, M., Yong, E.C., Fritsche, T.R. and Chiang, G.K.: *Toxoplasma gondii* stimulates the release of 13- and 9-hydroxyoctadecadienoic acids by human platelets. *Biochemistry*, 31 (1992) 5356-5362.
- 417 Hogge, L.R., Abrams, G.D., Abrams, S.R., Thibault, P. and Pleasance, S.: Characterization of abscisic acid and metabolites by combined liquid chromatography-mass spectrometry with ion-spray and plasma-spray ionization techniques. *J. Chromatogr.*, 623 (1992) 255-263.
- 418 Hvattum, E., Skrede, S., Bremer, J. and Solbakken, M.: The metabolism of tetradecylthiopropionic acid, a 4-thia stearic acid, in the rat. *In vivo* and *in vitro* studies. *Biochem. J.*, 286 (1992) 879-887.
- 419 Isaka, Y., Yamamoto, S., Fujiwara, Y., Shin, S., Ochi, S., Kamada, T. and Ueda, N.: (A new analytical method for simultaneous measurement of iothalamate and iohexol). *Rinsho Byori*, 40 (1992) 703-707; C.A., 117 (1992) 146447g.
- 420 Ishinaga, M., Yamauchi, T., Egusa, K. and Kuroda, K.: Changes of dolichol and dolichyl fatty acyl esters during the germination and development of soybean. *Biochem. Cell Biol.*, 70 (1992) 466-469.
- 421 Jareonkitmongkol, S., Kawashima, H., Shimizu, S. and Yamada, H.: Production of 5,8,11-*cis*-eicosatrienoic acid by a Δ 12-desaturase-defective mutant of *Martirella alpina* 1S-4. *J. Am. Oil Chem. Soc.*, 69 (1992) 939-944.
- 422 Konrad, R.J., Jolly, Y.C., Major, C. and Wolf, B.A.: Carbachol stimulation of phospholipase A₂ and insulin secretion in pancreatic islets. *Biochem. J.*, 287 (1992) 283-290.
- 423 Krausse, R. and Ullmann, U.: A modified procedure for the identification of anaerobic bacteria by high performance liquid chromatography - quantitative analysis of short-chain fatty acids. *Zentralbl. Bakteriol.*, 276 (1991) 1-8; C.A., 117 (1992) 127525q.
- 424 Lu, D.S., Feng, W.Y., Ling, D.H. and Hua, W.Z.: Determination of hydroxy acids as their copper(II) complexes by reversed-phase liquid chromatography with UV detection. *J. Chromatogr.*, 623 (1992) 55-62.
- 425 Marquez-Ruiz, G., Perez-Camino, M.C. and Dobarganes, M.C.: Digestibility of fatty acid monomers, dimers and polymers in the rat. *J. Am. Oil Chem. Soc.*, 69 (1992) 930-934.
- 426 Mohring, H. and Spittler, G.: Produkte der Dimerisierung ungesättigter Fettsäuren VIII: Über die Zusammensetzung der Fraktion der "Intermediates" bei der Fettsäuredimerisierung. *Fat Sci. Technol.*, 94 (1992) 241-245.
- 427 Montedoro, G., Servili, M., Baldioli, M. and Miniati, E.: Simple and hydrolyzable phenolic compounds in virgin olive oil. 1. Their extraction, separation and quantitative and semiquantitative evaluation by HPLC. *J. Agric. Food Chem.*, 40 (1992) 1571-1576.
- 428 Nikolova-Damyanova, B., Herslof, B.G. and Christie, W.W.: Silver ion high-performance liquid chromatography of derivatives of isomeric fatty acids. *J. Chromatogr.*, 609 (1992) 133-140.
- 429 Nilsson, A., Hjelte, L. and Strandvik, B.: Incorporation of dietary [¹⁴C]arachidonic acid and [³H]eicosapentaenoic acid into tissue lipids during absorption of a fish oil emulsion. *J. Lipid Res.*, 33 (1992) 1295-1305.
- 430 Onyeneho, S.N. and Hettiarachchy, N.S.: Antioxidant activity of durum wheat bran. *J. Agric. Food Chem.*, 40 (1992) 1496-1500.
- 431 Petrarulo, M., Marangella, M., Cosseddu, D. and Linari, F.: High-performance liquid chromatographic assay for L-glyceric acid in body fluids. Application in primary hyperoxaluria type 2. *Clin. Chim. Acta*, 211 (1992) 143-153.
- 432 Plakas, S.M., Khoo, L. and Barron, M.G.: 2,4-Dichlorophenoxyacetic acid disposition after oral administration in channel catfish. *J. Agric. Food Chem.*, 40 (1992) 1236-1239.
- 433 Rouseff, R.L., Seetharaman, K., Naim, M., Nagy, S. and Zehavi, U.: Improved HPLC determination of hydrocinnamic acids in orange juice using solvents containing THF. *J. Agric. Food Chem.*, 40 (1992) 1139-1143.

- 434 Schwenk, U., Morita, E., Engel, R. and Schröder, J.-M.: Identification of 5-oxo-15-hydroxy-6,8,11,13-eicosatetraenoic acid as a novel and potent human eosinophil chemotactic eicosanoid. *J. Biol. Chem.*, 267 (1992) 12482-12488.
- 435 Seki, T., Orita, Y., Yamamoto, S., Ueda, N., Yanagihara, Y. and Noguchi, K.: (A method for the simultaneous determination of iothalamic acid and creatinine clearance by coupled-column liquid chromatography). *Rinsho Byori*, 40 (1992) 697-702; *C.A.*, 117 (1992) 146446f.
- 436 Stanek, R., Gain, R.E., Glover, D.D. and Larsen, B.: High performance ion exclusion chromatographic characterization of the vaginal organic acids in women with bacterial vaginosis. *Biomed. Chromatogr.*, 6 (1992) 231-235.
- 437 Stijntjes, G.J., te Koppele, J.M. and Vermeulen, N.P.E.: High-performance liquid chromatography-fluorescence assay of pyruvic acid to determine cysteine conjugate β -lyase activity: application of S-1,2-dichlorovinyl-L-cysteine and S-2-benzothiazoyl-L-cysteine. *Anal. Biochem.*, 206 (1992) 334-343.

For additional information see C.A.:

117 (1992) 24706v, 46827m, 61958z, 86099h, 110261c, 127538w, 146715t, 149578m.

See also 37, 214, 258, 263, 294, 319, 445, 459, 462, 470, 488, 491, 544, 1133, 1193, 1525, 1526, 1535, 1543, 1603.

11b. Prostaglandins

- 438 Arai, H., Nagano, Y., Narumiya, S. and Kita, T.: Decreased arachidonate metabolism in mouse peritoneal macrophages after foam cell transformation with oxidized low-density lipoproteins. *J. Biochem. (Tokyo)*, 112 (1992) 482-487.
- 439 Dray, F., Vulliez-le Normand, B., Deroussent, A., Briquet, I., Gabelle, M.-M., Nakamura, S., Wahl, L.M., Gouyette, A. and Salahuddin, Z.S.: Active metabolism of arachidonic acid by Kaposi sarcoma cells cultured from lung biopsies (KS-3); identification by HPLC and MS/MS of the predominant metabolite secreted as the 11,12-epoxy-eicosatrienoic acid. *Biochim. Biophys. Acta*, 1180 (1992) 83-90.
- 440 Hotter, G., Gómez, G., Ramis, I., Bioque, G., Roselló-Catafau, J. and Gelpi, E.: Solid-phase extraction of prostanoids using an automatic sample preparation system. *J. Chromatogr.*, 607 (1992) 239-243.
- 441 Nicoll-Griffith, D. and Zamboni, R.: N-Acetyl [³H] leukotriene D₄: a stable internal standard for automated BIO-FAST HPLC extraction and separation of LTE₄ from human urine. *Prostaglandins*, 43 (1992) 523-532; *C.A.*, 117 (1992) 104369c.
- 442 Ohara, A., Sugiyama, S., Hoshino, H., Hamajima, E., Goto, H., Tsukamoto, Y. and Ozawa, T.: Reduction of adverse effects of indometacin by anti-allergic drugs in rat stomachs. *Arzneim.-Forsch.*, 42 (1992) 1115-1118.
- 443 Van der Donk, E.M.M., Vervaart, J.M.A., Verhagen, J., Veldink, G.A. and Vliegenthart, J.F.G.: 12-Lipoxygenase from rat basophilic leukemia cells, an oxygenase with Leukotriene A₄-synthase activity. *Biochim. Biophys. Acta*, 1128 (1992) 14-25.

See also 405.

11c. Lipids and their constituents

- 444 Abidi, S.L. and Mounts, T.L.: Separation of molecular species of phosphatidylserine by reverse-phase ion-pair HPLC. *J. Liq. Chromatogr.*, 15 (1992) 2487-2502.
- 445 Achenbach, H., Heffer-Bübl, U., Waibel, R., Constenla, M.A. and Hahn, H.: Gaschromatographische Analyse des Samenfettes der mittelamerikanischen *Fevillea cordifolia* (Cucurbitaceae). *Fat Sci. Technol.*, 94 (1992) 294-297.
- 446 Booker, M.L., LaMorte, W.W., Ahrendt, S.A., Lillemo, K.D. and Pitt, H.A.: Distribution of phosphatidylcholine molecular species between mixed micelles and phospholipid-cholesterol vesicles in human gallbladder bile: dependence on acyl chain length and unsaturation. *J. Lipid Res.*, 33 (1992) 1485-1492.
- 447 Bouteiller, J.C. and Maurin, R.: (Relation between triacylglycerol structure and HPLC retention data). *Rev. Fr. Corps Gras*, 38 (1991) 297-304; *C.A.*, 117 (1992) 43740t.
- 448 Bussolino, F., Sironi, M., Bocchietto, E. and Mantovani, A.: Synthesis of platelet-activating factor by polymorphonuclear neutrophils stimulated with interleukin-8. *J. Biol. Chem.*, 267 (1992) 14598-14603.
- 449 Calari, H., Howard, S. and Bittman, R.: Synthesis and antineoplastic properties of an ether glycerophosphocholine, an analog of ET-18-OCH₃-GPC. *Biochem. Biophys. Res. Commun.*, 187 (1992) 603-608.
- 450 Chen, S., Menon, G. and Traldi, P.: Identification of aminophospholipid stereoisomers by positive-ion fast atom bombardment combined with collisional activation mass-analyzed ion kinetic energy analysis and high-performance liquid chromatography. *Org. Mass Spectrom.*, 27 (1992) 215-218; *C.A.*, 117 (1992) 43969z.
- 451 Cronholm, T., Viestam-Rains, M. and Sjövall, J.: Decreased content of arachidonoyl species of phosphatidylinositol phosphates in pancreas of rats fed on an ethanol-containing diet. *Biochem. J.*, 287 (1992) 925-928.
- 452 Davey, M.W. and Lambein, F.: Quantitative derivatization and high-performance liquid chromatographic analysis of cyanobacterial heterocyst-type glycolipids. *Anal. Biochem.*, 206 (1992) 323-327.
- 453 Davey, M.W. and Lambein, F.: Semipreparative isolation of individual cyanobacterial heterocyst-type glycolipids by reverse-phase high-performance liquid chromatography. *Anal. Biochem.*, 206 (1992) 226-230.
- 454 Demirbueker, M., Blomberg, L.G., Urban, O.N., Bergqvist, M., Hersloef, B.G., Alvarado, J.F.: Characterization of triacylglycerols in the seeds of *Aquilegia vulgaris* by chromatographic and mass spectrometric methods. *Lipids*, 27 (1992) 436-441; *C.A.*, 117 (1992) 127511g.
- 455 Evershed, R.P. and Goad, L.J.: Chromatographic and mass-spectrometric methods for the analysis of the steryl esters of plants. *Stud. Nat. Prod. Chem.*, 9(Struct. Chem., Pt. B) (1991) 447-486; *C.A.*, 117 (1992) 22584e - a review with 126 refs.
- 456 Ford, D.A., Rosenbloom, K.B. and Gross, R.W.: The primary determinant of rabbit myocardial ethanolamine phosphotransferase substrate selectivity is the covalent nature of the sn-1 aliphatic group of diacyl glycerol acceptors. *J. Biol. Chem.*, 267 (1992) 11222-11228.

- 457 Hierro, M.T.G., Tomás, M.C., Fernández-Martín, F. and Santa-María, G.: Determination of the triglyceride composition of avocado oil by high-performance liquid chromatography using a light-scattering detector. *J. Chromatogr.*, 607 (1992) 329-338.
- 458 Hirose, S., Prince, G.M., Sevlever, D., Ravi, L., Rosenberry, T.L., Ueda, E. and Medof, M.E.: Characterization of putative glycoinositol phospholipid anchor precursors in mammalian cells. Localization of phosphoethanolamine. *J. Biol. Chem.*, 267 (1992) 16968-16974.
- 459 Hopia, A.I., Piironen, V.I., Koivistoinen, P.E. and Hyvönen, L.E.T.: Analysis of lipid classes by solid-phase extraction and high-performance size exclusion chromatography. *J. Am. Oil Chem. Soc.*, 69 (1992) 772-776.
- 460 Itabashi, Y., Marai, L. and Kuksis, A.: Identification of natural diacylglycerols as the 3,5-dinitrophenylurethanes by chiral phase liquid chromatography with mass spectrometry. *Lipids*, 26 (1991) 951-956; *C.A.*, 117 (1992) 22800x.
- 461 Kamal-Eldin, A., Yousif, G., Iskander, G.M. and Appelquist, L.-A.: Seed lipids of *Sesamum indicum* L. and related wild species in Sudan I: Fatty acids and triacylglycerols. *Fat Sci. Technol.*, 94 (1992) 254-259.
- 462 Kasurinen, J.: A novel fluorescent fatty acid, 5-methyl-BD3-3-dodecanoic acid, is a potential probe in lipid transport studies by incorporating selectively to lipid classes of BHK cells. *Biochem. Biophys. Res. Commun.*, 187 (1992) 1594-1601.
- 463 Kondo, Y., Miyazawa, T. and Mizutani, J.: Detection and time-course analysis of phospholipid hydroperoxide in soybean seedlings after treatment with fungal elicitor, by chemiluminescence-HPLC assay. *Biochim. Biophys. Acta*, 1127 (1992) 227-232.
- 464 Kropp, J., Ambrose, K.R., Knapp, F.F., Jr. and Nissen, H.P.: Incorporation of radiiodinated IPPA and BMIPP fatty acid analogs into complex lipids from isolated rat hearts. *Nucl. Med. Biol.*, 19 (1992) 283-288; *C.A.*, 117 (1992) 22630s.
- 465 Kuksis, A., Marai, L., Myher, J.J., Itabashi, Y. and Pind, S.: Applications of GC/MS, LC/MS and FAB/MS to determination of molecular species of glycerolipids. In: Perkins, E.G. (Editor), *Anal. Fats, Oils Lipoproteins*, AOCS, Champaign, 1991, pp. 464-467; *C.A.*, 117 (1992) 166826h - a review with 41 refs.
- 466 Kuksis, A., Marai, L., Myher, J.J., Itabashi, Y. and Pind, S.: Qualitative and quantitative analysis of molecular species of glycerolipids by HPLC. In: Perkins, E.G. (Editor), *Anal. Fats, Oils Lipoproteins*, AOCS, Champaign, 1991, pp. 214-232; *C.A.*, 117 (1992) 146374f - a review with 58 refs.
- 467 Liem, A.K.D., Baumann, R.A., de Jong, A.P.J.M., van der Velde, E.G. and van Zoonen, P.: Analysis of organic micropollutants in the lipid fraction of foodstuffs. *J. Chromatogr.*, 624 (1992) 317-339 - a review with 172 refs.
- 468 Macphee, C.H., Carter, A.N., Ruiz-Larrea, F., Ward, J.G., Young, R.C. and Downes, C.P.: The stereoselective recognition of substrates by phosphoinositide kinases. Studies using synthetic stereoisomers of dipalmitoyl phosphatidylinositol. *J. Biol. Chem.*, 267 (1992) 11137-11143.
- 469 Marini, D.: HPLC of lipids. *Food Sci. Technol.*, 52(Food Anal. HPLC) (1992) 169-240; *C.A.*, 117 (1992) 110172z - a review with 86 refs.
- 470 Marquez-Ruiz, G., Perez-Camino, M.C. and Dobarganes, M.C.: *In vitro* action of pancreatic lipase on complex glycerides from thermally oxidized oils. *Fat Sci. Technol.*, 94 (1992) 307-312.
- 471 Melton, S.L.: Analysis of soybean lecithins and beef phospholipids by HPLC with an evaporative light scattering detector. *J. Am. Oil Chem. Soc.*, 69 (1992) 784-788.
- 472 Mezin, I.A., Menzeleeva, G.K., Menzeleev, R.F., Pinchuk, A.N., Krasnopol'skii, Y.M. and Shvets, V.I.: (Complex preparative isolation of lipids from bovine brain). *Biol. Membr.*, 9 (1992) 301-307; *C.A.*, 117 (1992) 127677r.
- 473 Morikawa, S., Inubushi, T., Kitoh, K., Kido, C. and Nozaki, M.: Chemical assessment of phospholipid and phosphoenergetic metabolites in regenerating rat liver measured by *in vivo* and *in vitro* ^{31}P -NMR. *Biochim. Biophys. Acta*, 1117 (1992) 251-257.
- 474 Nozawa, M., Iwamoto, T., Tokoro, T. and Eto, Y.: Novel procedure for measuring psychosine derivatives by a HPLC method. *J. Neurochem.*, 59 (1992) 607-609; *C.A.*, 117 (1992) 107543r.
- 475 Olsson, N.U., Kaufmann, P. and Kroon, C.-G.: HPLC separation of molecular species of intact sphingomyelin, utilizing multivariate design and optimization. *Chromatographia*, 34 (1992) 529-533.
- 476 Ousley, A.H. and Morell, P.: Individual molecular species of phosphatidylcholine and phosphatidylethanolamine in myelin turn over at different rates. *J. Biol. Chem.*, 267 (1992) 10362-10369.
- 477 Purdon, M.P.: Application of HPLC to lipid separation and analysis: general qualitative and quantitative analysis. In: Perkins, E.G. (Editor), *Anal. Fats, Oils Lipoproteins*, AOCS, Champaign, 1991, pp. 193-209; *C.A.*, 117 (1992) 146373e - a review with 60 refs.
- 478 Purdon, M.P.: Application of HPLC to lipid separation and analysis: sample preparation. In: Perkins, E.G. (Editor), *Anal. Fats, Oils Lipoproteins*, AOCS, Champaign, 1991, pp. 145-165; *C.A.*, 117 (1992) 146371c - a review with 108 refs.
- 479 Salivaras, E. and McCurdy, A.R.: Detection of olive oil adulteration with canola oil from triacylglycerol analysis by reversed-phase high-performance liquid chromatography. *J. Am. Oil Chem. Soc.*, 69 (1992) 935-938.
- 480 Scarselli, L., Giampaoli, F., Amancipato, L. and Aloisi, U.: (The content of trilinolein in the triglyceride fraction of olive oils as an index of their genuineness. HPLC study with refractive index detection of olive oils marketed and produced in Lucca Province). *Boll. Chim. Ig., Parte Sci.*, 42 (1991) 681-690; *C.A.*, 117 (1992) 68606s.
- 481 Shukla, V.K.S.: Application of HPLC to lipid separation and analysis: lipid derivatives. In: Perkins, E.G. (Editor), *Anal. Fats, Oils Lipoproteins*, AOCS, Champaign, 1991, pp. 233-235; *C.A.*, 117 (1992) 169629g - a review with 48 refs.
- 482 Shukla, V.K.S.: Application of HPLC to lipid separation and analysis: neutral lipids. In: Perkins, E.G. (Editor), *Anal. Fats, Oils Lipoproteins*, AOCS, Champaign, 1991, pp. 210-213; *C.A.*, 117 (1992) 169628f - a review with 13 refs.
- 483 Singh, A.K.: Quantitative analysis of inositol lipids and inositol phosphates in synaptosomes and microvessels by column chromatography: comparison of the mass analysis and the radiolabelling methods. *J. Chromatogr.*, 581 (1992) 1-10.
- 484 Suzuki, T., Ota, T. and Takagi, T.: Temperature effect on enantiomeric separation of diacylglycerol derivatives by HPLC on various chiral columns. *J. Chromatogr. Sci.*, 30 (1992) 315-318.

- 485 Tomai, Y., Kojima, H., Saito, S., Takayama-Abe, K. and Horichi, H.: Characteristic distribution of glycolipids in gadoid fish nerve tissues and its bearing on phylogeny. *J. Lipid Res.*, 33 (1992) 1351-1359.
- 486 Van Wijk, G.M.T., Hostetler, K.Y. and van den Bosch, H.: Antiviral nucleoside diphosphate diglycerides: improved synthesis and facilitated purification. *J. Lipid Res.*, 33 (1992) 1211-1219.
- 487 Van der Meeren, P., Vanderdeelen, J. and Baert, L.: Phospholipid analysis by HPLC. *Food Sci. Technol.*, 52(Food Anal. HPLC) (1992) 241-258; *C.A.*, 117 (1992) 110173a - a review with 35 refs.
- 488 Wieland, E., Diedrich, F., Niedmann, P.D. and Seidel, D.: Measurement of free fatty acid, phospholipid, triglyceride, and cholesterol ester hydroperoxides by a CL-HPLC method. In: Stanley, P.E. and Kricka, L.J. (Editors), *Biolumin. Chemilumin. Proc. Int. Symp., 6th 1990*, Wiley, Chichester, 1991, pp. 235-238; *C.A.*, 117 (1992) 127510f.
- 489 Yamasaki, K., Kikuoka, M., Saeki, T., Kokusenyu, Y. and Matsuo, M.: (Determination of lecithin in gastric juice by HPLC with electrochemical detection using enzyme reactions). *Bunseki Kagaku*, 41 (1992) 275-279; *C.A.*, 117 (1992) 107525m.
- 490 Yamashita, G., Tazuma, S. and Kajiyama, G.: Effects of organic anions on biliary lipid secretion in rats. Importance of association with biliary lipid structures. *Biochem. J.*, 286 (1992) 193-196.
- 491 Yates, R.A. and Caldwell, J.C.: Adsorptive capacity of active filter aids for used cooking oil. *J. Am. Oil Chem. Soc.*, 69 (1992) 894-897.
- For additional information see *C.A.*:
117 (1992) 49145z.
- See also 235, 519, 523, 572, 961, 1158, 1159.
- 11d. *Lipoproteins and their constituents*
- 492 Alaupovic, P. and Koren, E.: Immunoaffinity chromatography of plasma lipoprotein particles. In: Perkins, E.G. (Editor), *Anal. Fats, Oils Lipoproteins*, AOCS, Champaign, 1991, pp. 599-622; *C.A.*, 117 (1992) 166830e - a review with 97 refs.
- 493 Burkey, B.F., Stuart, W.D. and Harmony, J.A.K.: Hepatic apolipoprotein J is secreted as a lipoprotein. *J. Lipid Res.*, 33 (1992) 1517-1526.
- 494 De Rijke, Y.B., Jürgens, G., Hessels, E.M.A., Hermann, A. and van Berkel, T.J.C.: *In vivo* fate and scavenger receptor recognition of oxidized lipoprotein [a] isoforms in rats. *J. Lipid Res.*, 33 (1992) 1315-1325.
- 495 Edelstein, C.: Conventional and high performance liquid chromatography as applied to human plasma apolipoproteins. In: Perkins, E.G. (Editor), *Anal. Fats, Oils Lipoproteins*, AOCS, Champaign, 1991, pp. 589-598; *C.A.*, 117 (1992) 146378k - a review with 22 refs.
- 496 Gerdes, L.U., Gerdes, C., Klausen, I.C. and Faergeman, O.: Generation of analytic plasma lipoprotein profiles using two prepacked Superose 6B columns. *Clin. Chim. Acta*, 205 (1992) 1-9.
- 497 Mezdoor, H., Nomura, S., Yamamura, T. and Yamamoto, A.: Concentration and distribution of apolipoproteins A-I and E in normolipidemic, WHHL and diet-induced hyperlipidemic rabbit sera. *Biochim. Biophys. Acta*, 1127 (1992) 116-123.
- 498 Mowri, H.-O., Patsch, W., Smith, L.C., Gotto, A.M., Jr. and Patsch, J.R.: Different reactivities of high density lipoprotein₂ subfractions with hepatic lipase. *J. Lipid Res.*, 33 (1992) 1269-1279.
- 499 Rumsey, S.C., Galeano, N.F., Arad, Y. and Deckelbaum, R.J.: Cryopreservation with sucrose maintains normal physical and biological properties of human plasma low density lipoproteins. *J. Lipid Res.*, 33 (1992) 1551-1561.
- See also 666.
12. ORGANIC PEROXIDES
- 500 Chamulitrat, W., Iwashashi, H., Kelman, D.J. and Mason, R.P.: Evidence against the 1:2:2:1 quartet DMPO spectrum as the radical adduct of the lipid alkoxyl radical. *Arch. Biochem. Biophys.*, 296 (1992) 645-649.
- See also 463, 488.
13. STEROIDS
- 13a. *General techniques*
- 501 Gardner, R.S., Walker, M. and Edwardson, P.A.D.: Development of sensitive HPLC assays for the analysis of steroids *in vitro* and *in vivo* studies. In: Sunshine, I. (Editor), *Recent Dev. Ther. Drug Monit. Clin. Toxicol.*, Dekker, New York, 1992, pp. 503-511; *C.A.*, 117 (1992) 97407h.
- 502 Jansen, E.H.J.M., van Ginkel, L.A., van den Berg, R.H. and Stephany, R.W.: High-performance liquid chromatographic separation and detection methods for anabolic compounds. *J. Chromatogr.*, 580 (1992) 111-124 - a review with 58 refs.
- 503 Miscicka, M., Sadlej-Sosnowska, N. and Wilczynska-Wojtulewicz, I.: (Determination of the components of mixtures containing hydrocortisone by high-performance liquid chromatography). *Acta Pol. Pharm.*, 47 (1990) 25-28; *C.A.*, 117 (1992) 137783m.
- 504 Van Ginkel, L.A., Jansen, E.H.J., Stephany, R.W., Zoontjes, P.W., Schwillens, P.L.W.J., van Rossum, H.J. and Visser, T.: Liquid chromatographic purification and detection of anabolic compounds. *J. Chromatogr.*, 624 (1992) 389-401 - a review with 29 refs.
- 505 Venkatesh, B., Tan, C.H., Kime, D.E. and Lam, T.J.: Steroid metabolism in teleost gonads: purification and identification of metabolites by high-performance liquid chromatography. *Steroids*, 57 (1992) 276-281; *C.A.*, 117 (1992) 43771d.
- 506 Wong, T., Shackleton, C.H.L., Covery, T.R. and Ellis, G.: Identification of the steroids in neonatal plasma that interfere with 17 α -hydroxyprogesterone radioimmunoassays. *Clin. Chem. (Winston-Salem)*, 38 (1992) 1830-1837.
- See also 201, 876, 1328.

13b. *Pregnane and androstane derivatives*

- 507 Cavina, G., Valvo, L., Gallinella, B., Porra, R. and Savella, A.L.: The identification of related substances in 9α -fluoroprednisolone-21 acetate by means of high-performance liquid chromatography with diode array detector and mass spectrometry. *J. Pharm. Biomed. Anal.*, 10 (1992) 437-446.
- 508 Gemzik, B., Green, J. and Parkinson, A.: Hydroxylation of 5α -androstane- $3\beta,17\beta$ -diol by rat prostate microsomes: effects of antibodies and chemical inhibitors of cytochrome P450 enzymes. *Arch. Biochem. Biophys.*, 296 (1992) 355-365.
- 509 Jayyosi, Z., Cooper, K.O. and Thomas, P.E.: Brain cytochrome P450 and testosterone metabolism by rat brain subcellular fractions: presence of cytochrome P450 3a immunoreactive protein in rat brain mitochondria. *Arch. Biochem. Biophys.*, 298 (1992) 265-270.
- 510 Mason, S.R., Ward, L.C. and Reilly, P.E.B.: Fluorimetric detection of serum corticosterone using high-performance liquid chromatography. *J. Chromatogr.*, 581 (1992) 267-271.
- 511 McBride, J.H., Ettenger, R.B., Rodgerson, D.O., Park, S.S. and Reyes, A.F.: Monitoring of plasma glucocorticosteroids as an indicator of noncompliance in pediatric renal transplant recipients. In: Sunshine, I. (Editor), *Recent Dev. Ther. Drug Monit. Clin. Toxicol.*, Dekker, New York, 1992, pp. 147-152; *C.A.*, 117 (1992) 63115w.
- 512 Noroski, J., Mayo, D. and Kirschbaum, J.J.: Liquid chromatographic resolution of the isomers of tipredane and phenylthio-proline using urea-solubilized β -cyclodextrin in the mobile phase. *J. Pharm. Biomed. Anal.*, 10 (1992) 447-455.
- 513 Parini, C., Bacigalupo, M.A., Colombi, S. and Saita, R.: Reversed phase TLC and HPLC of hemisuccinyl derivatives of 11α -hydroxyprogesterone. *J. Planar Chromatogr.*, 5 (1992) 251-254.
- 514 Payne, D.W., Packman, J.N. and Adashi, E.Y.: Follicle-stimulating hormone inhibits granulosa cell 5α -reductase activity. Possible role of 5α -reductase as a steroidogenic pubertal switch. *J. Biol. Chem.*, 267 (1992) 13348-13355.
- 515 Szecsi, M., Toth, I. and Faredin, I.: (Separation of radioactive androgenic steroids and precursors by high pressure liquid chromatography). *Izotoptech. Diagn.*, 34 (1991) 111-114; *C.A.*, 117 (1992) 83533j.
- 516 Yoshitake, T., Ishida, J., Sonezaki, S. and Yamaguchi, M.: High performance liquid chromatographic determination of $3\alpha,5\beta$ -tetrahydroaldosterone and cortisol in human urine with fluorescence detection. *Biomed. Chromatogr.*, 6 (1992) 217-221.

See also 1350.

13c. *Estrogens*

- 517 Oda, T., Sakakibara, Y., Sato, Y., Hanzawa, H. and Hata, T.: Effects of (+)-, (-) and (\pm)-indenestrols A and B on microtubule polymerization. *Chem. Pharm. Bull.*, 40 (1992) 588-592.
- 518 Su, S.Y., Shiu, G.K., Simmons, J., Viswanathan, C.T. and Skelly, J.P.: High performance liquid chromatographic analysis of six conjugated and unconjugated estrogens in serum. *Biomed. Chromatogr.*, 6 (1992) 265-268.

See also 502.

13d. *Sterols*

- 519 Fenton, M.: Chromatographic separation of cholesterol in foods. *J. Chromatogr.*, 624 (1992) 369-388 - a review with 117 refs.
- 520 Grob, K., Artho, A. and Mariani, C.: Determination of raffination of edible oils and fats by olefinic degradation products of sterols and squalene, using coupled LC-GC. *Fat Sci. Technol.*, 94 (1992) 394-400.
- 521 Jain, U.K., Higuchi, W.I., Liu, C.L., Lee, P.H. and Mazer, N.A.: Cholesterol (thermodynamic) activity determinations in bile salt-lecithin-cholesterol systems and cholesterol-rich liquid crystalline mesophase formation. *Pharm. Res.*, 9 (1992) 792-799; *C.A.*, 117 (1992) 40165y.
- 522 Lund, E., Diczfalusy, U. and Björkhem, I.: On the mechanism of oxidation of cholesterol at C-7 in a lipoxygenase system. *J. Biol. Chem.*, 267 (1992) 12462-12467.
- 523 Subbaiah, P.V., Liu, M., Bolan, P.J. and Paltauf, F.: Altered positional specificity of human plasma lecithin-cholesterol acyltransferase in the presence of sn-2 arachidonoyl phosphatidyl cholines. Mechanism of formation of saturated cholesteryl esters. *Biochim. Biophys. Acta*, 1128 (1992) 83-92.

See also 459, 525, 876.

13e. *Bile acids and alcohols*

- 524 Chen, N., Zhang, Y. and Lu, P.: (Recommendation of analytical method for bile acids in expert system for liquid chromatography and peak identification by using interaction index in reversed-phase high performance liquid chromatography). *Sepu*, 10 (1992) 177-179; *C.A.*, 117 (1992) 65795a.
- 525 Fuchs, M., Scheibner, J., Hörmann, E., Tauber, G. and Stange, E.F.: Enterohepatic circulation in hamsters with an extracorporeal bile duct. *J. Lipid Res.*, 33 (1992) 1383-1392.
- 526 Roda, A., Cerre, C., Simoni, P., Polimeni, C., Vaccari, C. and Pistillo, A.: Determination of free and amidated bile acids by high-performance liquid chromatography with evaporative light-scattering mass detection. *J. Lipid Res.*, 33 (1992) 1393-1402.
- 527 Zhang, Y.K., Chen, N., Zao, R.H., Li, T., Wang, Z.Y. and Lu, P.C.: The retention behaviour of conjugated bile acids in reversed phase high performance liquid chromatography. *Biomed. Chromatogr.*, 6 (1992) 255-257.

For additional information see *C.A.*:
117 (1992) 43744x, 56089n.

13f. *Ecdysones and other insect steroid hormones*

- 528 Zhang, M., Stout, M.J. and Kubo, I.: Isolation of ecdysteroids from *Vitex strickeri* using RLCC and recycling HPLC. *Phytochemistry*, 31 (1992, Pub. 1991) 247-250; *C.A.*, 117 (1992) 64699b.

13g. *Other sterols*

- 529 Chen, C. and Wu, R.: Analysis of intermediates of Ru486 by HPLC. *Zhongguo Yaoke Daxue Xuebao*, 23 (1992) 51-54; *C.A.*, 117 (1992) 157743b.

For additional information see *C.A.*:
117 (1992) 76600a.

14. STEROID GLYCOSIDES AND SAPONINS

- 530 Da, S., Tang, X., Da, H. and Yang, J.: (Determination of diosgenin in *Dioscorea zingiberensis* by using reversed-phase high performance liquid chromatography). *Sepu*, 10 (1992) 98-100; C.A., 117 (1992) 65798v.
- 531 Ferth, R. and Kopp, B.: Cardenolide aus *Ornithogalum umbellatum* L. *Pharmazie*, 47 (1992) 626-629.

See also 1516, 1517.

15. TERPENES AND OTHER VOLATILE AROMATIC COMPOUNDS

15a. Terpenes

- 532 Komae, H. and Hayashi, N.: Gel permeation chromatography of mono- and sesquiterpenes. In: Bhattacharyya, S.C., Sen, N. and Sethi, K.L. (Editors), *Proc.-Int. Congr. Essent. Oils, Fragrances Flavours*, 11th, Oxford and IBH, New Delhi, 1989, pp. 57-62; C.A., 117 (1992) 123770t.
- 533 Moreau, R.A., Preisig, C.L., Osman, S.F.: A rapid quantitative method for the analysis of sesquiterpene phytoalexins by high performance liquid chromatography. *Phytochem. Anal.*, 3 (1992) 125-128; C.A., 117 (1992) 146443c.
- 534 Striedner, J., Czygan, F.C. and Braunegg, G.: Contributions to the biotechnological production of sweeteners from *Stevia rebaudiana* Berton. I. A method for the serial analysis of diterpene glycosides by HPLC. *Acta Biotechnol.*, 11 (1991) 495-499; C.A., 117 (1992) 24841k.

See also 134, 1520.

15b. Essential oils

- 535 Verschuere, M., Sandra, P. and David, F.: Fractionation by SFE and microcolumn analysis of the essential oil and the bitter principles of hops. *J. Chromatogr. Sci.*, 30 (1992) 388-391.

For additional information see C.A.:
117 (1992) 118191a.

15c. Bitter substances

- 536 De Keukeleire, D., Vindevogel, J., Szűcs, R. and Sandra, P.: The history and analytical chemistry of beer bitter acids. *TrAC*, 11 (1992) 275-280 - a review with 34 refs.

See also 535, 1531.

16. NITRO AND NITROSO COMPOUNDS

- 537 Muir, A.D. and Soroka, J.J.: HPLC-ion chromatographic analysis of nitrate in plant tissues and its application to sweetclover (*Melilotus* spp.) seedlings. *J. Agric. Food Chem.*, 40 (1992) 1602-1605.

- 538 Sen, N.P.: Determination of N-nitroso compounds in foods and beverages using HPLC. *Food Sci. Technol.*, 52(Food Anal. HPLC) (1992) 673-696; C.A., 117 (1992) 110183d - a review with 95 refs.

For additional information see C.A.:
117 (1992) 42027d.

See also 86, 88, 1364.

17. AMINES, AMIDES AND RELATED NITROGEN COMPOUNDS

17a. Amines and polyamines

- 539 Bazylak, G.: Helical nickel(II) complexes as mobile phase additives in the reversed-phase high-performance liquid chromatographic separation of alkylamine enantiomers. *Analyst (London)*, 117 (1992) 1429-1433.
- 540 Debruyckere, G. and van Peteghem, C.: HPLC analysis of organic bases. *Food Sci. Technol.*, 52(Food Anal. HPLC) (1992) 643-671; C.A., 117 (1992) 110182c - a review with 149 refs.
- 541 Duchateau, A.L.L., Knuts, H., Boesten, J.M.M. and Guns, J.J.: Enantioseparation of amino compounds by derivatization with o-phthalaldehyde and D-3-mercapto-2-methylpropionic acid. *J. Chromatogr.*, 623 (1992) 232-245.
- 542 Flentge, F., Venema, K., Koch, T. and Korf, J.: An enzyme-reactor for electrochemical monitoring of choline and acetylcholine: applications in high-performance liquid chromatography, brain tissue, microdialysis and cerebrospinal fluid. *Anal. Biochem.*, 204 (1992) 305-310.
- 543 Gazzani, G., Stoppini, G., Gandini, C. and Bettero, A.: Reversed-phase high-performance liquid chromatographic and derivative UV spectrophotometric determination of α -phenylethylamine in phosphomycin. *J. Chromatogr.*, 609 (1992) 391-394.
- 544 Karlsson, A. and Pettersson, C.: Separation of enantiomeric amines and acids using chiral ion-pair chromatography on porous graphitic carbon. *Chirality*, 4 (1992) 323-332; C.A., 117 (1992) 166915m.
- 545 Lehtonen, P., Saarinen, M., Vesanto, M. and Riekkola, M.L.: Determination of wine amines by HPLC using automated precolumn derivatization with o-phthalaldehyde and fluorescence detection. *Z. Lebensm.-Unters. Forsch.*, 194 (1992) 434-437; C.A., 117 (1992) 149587p.
- 546 Maruyama, M.: Determination of trace amounts of 3,4-dichloroaniline by high-performance liquid chromatography with amperometric detection and its application to pesticide residue analysis. *Fresenius J. Anal. Chem.*, 343 (1992) 890-892.
- 547 McCrossen, S.D., Giles, R.G., Oxley, P.W. and McArdle, J.V.: High-performance liquid chromatography procedure for the determination of purity of di-N-n-propylamine. *J. Chromatogr.*, 623 (1992) 229-235.
- 548 Noda, N., Tsunefuka, S., Tanaka, R. and Miyahara, K.: Isolation of two 1-O-alkyl-sn-glycero-3-phosphocholines from the earthworm, *Pheretima asiatica*. *Chem. Pharm. Bull.*, 40 (1992) 1349-1351.
- 549 Oefner, P.J., Wongyai, S. and Bonn, G.: High-performance liquid chromatographic determination of free polyamines in human seminal plasma. *Clin. Chim. Acta*, 205 (1992) 11-18.

- 550 Plessi, M., Ferioli, V., Gamberini, G. and Monzani, A.: High-performance liquid chromatographic assay of bioactive amines in Parmigiano-Reggiano cheese. *Atti Soc. Nat. Mat. Modena*, 121 (1990) 1-9; *C.A.*, 117 (1992) 130023e.
- 551 Schwarzenbach, R. and Gubler, D.: Detection of heterocyclic aromatic amines in food flavours. *J. Chromatogr.*, 624 (1992) 491-495.
- 552 Smith, M.A.: Chromatographic methods for the identification and quantitation of polyamines. In: Slocum, R.D. and Flores, H.E. (Editors), *Biochem. Physiol. Polyamines Plants*, CRC, Boca Raton, 1991, pp. 229-242; *C.A.*, 117 (1992) 86023d - a review with 51 refs.
- 553 Sparks, T.C. and Geng, C.: Analysis of the biogenic amines in the central nervous system of the tobacco hornworm by high-performance liquid chromatography with 16-sensor electrochemical detection. *Anal. Biochem.*, 205 (1992) 319-325.
- 554 Yoshida, T., Hirata, M., Tabuchi, T., Miyajima, K. and Andoh, K.: (Amounts of urinary metabolites of *p*-chloroanilines and their half lives in a patient with acute poisoning). *Sangyo Igaku*, 34 (1992) 126-130; *C.A.*, 117 (1992) 105771q.
- For additional information see *C.A.*:
117 (1992) 127540r.
- See also 39, 105, 111, 112, 202, 406, 567, 568, 1133, 1134, 1245, 1374.
- 17b. *Catecholamines and their metabolites*
- 555 Kaplan, P., Tejral, M. and Nentwich, I.: The determination of urinary metanephrine and normetanephrine by high performance liquid chromatography with ultraviolet absorbance detection. *J. Liq. Chromatogr.*, 15 (1992) 2561-2579.
- 556 Lagerkvist, S.: Sample splitting provides a fast and selective method for determining brain dialysate dopamine and its metabolites. In: Rollema, H., Westerink, B.H.C. and Drijfhout, W.J. (Editors), *Monit. Mol. Neurosci., Proc. Int. Conf. In Vivo Methods, 5th 1991*, Univ. Cent. Pharm., Groningen, 1991, pp. 136-138; *C.A.*, 117 (1992) 163985y.
- 557 Maeno, Y.: Experimental studies on the vital reaction in wounded skin - simultaneous observation of biogenic amines in incised skin wounds. *Nagoya Med. J.*, 36 (1991) 23-41; *C.A.*, 117 (1992) 42072q.
- 558 Pagliari, R., Peyrin, L., Cottet-Emard, J.M. and Westerink, B.H.C.: *In vivo* monitoring of noradrenaline release from the frontal cortex of free moving rats using chronic microdialysis and HPLC detection. In: Rollema, H., Westerink, B.H.C. and Drijfhout, W.J. (Editors), *Monit. Mol. Neurosci., Proc. Int. Conf. In Vivo Methods, 5th 1991*, Univ. Cent. Pharm., Groningen, 1991, pp. 142-145; *C.A.*, 117 (1992) 63119a.
- 559 Parkin, H.M. and Macdonald, I.A.: Measurement of plasma noradrenaline and dihydroxyphenylglycol (DHPG) in plasma from healthy volunteers and patients with chest pain. In: Rollema, H., Westerink, B.H.C. and Drijfhout, W.J. (Editors), *Monit. Mol. Neurosci., Proc. Int. Conf. In Vivo Methods, 5th 1991*, Univ. Cent. Pharm., Groningen, 1991, pp. 171-172; *C.A.*, 117 (1992) 63120u.
- 560 Saito, H., Murai, S., Abe, E., Masuda, Y. and Itoh, T.: Rapid and simultaneous assay of monoamine neurotransmitters and their metabolites in discrete brain areas of mice by HPLC with coulometric detection. *Pharmacol., Biochem. Behav.*, 42 (1992) 351-356; *C.A.*, 117 (1992) 40626z.
- 561 Spicer, C.H., Forster, C.D. and Macdonald, I.A.: The assay of noradrenaline, adrenaline and isoprenaline in biological fluids. In: Rollema, H., Westerink, B.H.C. and Drijfhout, W.J. (Editors), *Monit. Mol. Neurosci., Proc. Int. Conf. In Vivo Methods, 5th 1991*, Univ. Cent. Pharm., Groningen, 1991, pp. 176-177; *C.A.*, 117 (1992) 63122w.
- 562 Villanueva Camanas, R.M., Sanchis Mollo, J.M., Simo Alfonso, E.F. and Ramis Ramos, G.: Thermal lens spectrometric detection of catecholamines after oxidation to aminochromes. *Anal. Lett.*, 25 (1992) 1425-1445.
- For additional information see *C.A.*:
117 (1992) 63103r, 63779r, 124679g, 124689k, 124769m.
- See also 32.
- 17c. *Urea and guanidine derivatives*
- See 1149.
- 17d. *Other amine derivatives and amides (excl. peptides)*
- 563 Aspley, S., Kulkarni, R.S., Fone, K.C.F., Bennett, G.W. and Marsden, C.A.: ACh measurement by HPLC-ED: the use of second IMER for the analysis of *in vitro* ACh release from striatal and hippocampal tissue slices. In: Rollema, H., Westerink, B.H.C. and Drijfhout, W.J. (Editors), *Monit. Mol. Neurosci., Proc. Int. Conf. In Vivo Methods, 5th 1991*, Univ. Cent. Pharm., Groningen, 1991, pp. 297-299; *C.A.*, 117 (1992) 63126a.
- 564 Baker, D.G., Don, H.F. and Brown, J.K.: Direct measurement of acetylcholine release in guinea pig tracheas. *Am. J. Physiol.*, 263 (1992) L142-L147; *C.A.*, 117 (1992) 143625c.
- 565 Ghanayem, B.I., Sanchez, I.M. and Burka, L.T.: Effects of dose, strain, and dosing vehicle on methacrylonitrile disposition in rats and identification of a novel-exhaled metabolite. *Drug Metab. Disp.*, 20 (1992) 643-652.
- 566 Jackson, P.E. and Kahler, B.: The determination of hydrazine in sulfuric acid by precolumn derivatization liquid chromatography. *J. High Resolut. Chromatogr.*, 15 (1992) 620-621.
- 567 Lagana, A., Mario, A., Fago, G. and Micheli, A.: Determination of free sphingosine in biological system by high performance liquid chromatography. *Ann. Chim. (Rome)*, 81 (1991) 721-734; *C.A.*, 117 (1992) 22616s.
- 568 McMaster, C.R. and Choy, P.C.: The determination of tissue ethanalamine levels by reverse-phase high-performance liquid chromatography. *Lipids*, 27 (1992) 560-563; *C.A.*, 117 (1992) 86116m.
- 569 Oomah, B.D., Mazza, G. and Kenaschuk, E.O.: Cyanogenic compounds in flaxseed. *J. Agric. Food Chem.*, 40 (1992) 1346-1348.
- 570 Ricny, J., Tucek, S., and Vins, I.: Sensitive method for HPLC determination of acetylcholine, choline and their analog using fluorometric detection. *J. Neurosci. Methods*, 41 (1992) 11-17; *C.A.*, 117 (1992) 40603q.

- 571 Santoni, G., Bavazzano, P., Perico, A., Colzi, A., Benassi, S., Medica, A., la Morgia, R. and Giuliano, G.: High-performance liquid chromatographic determination of N-methylformamide and N-methyl-N-(hydroxymethyl)-formamide in human urine. *J. Chromatogr.*, 581 (1992) 287-292.
- For additional information see C.A.:
117 (1992) 39584c.
- See also 473, 554, 893, 1318, 1331, 1513.
18. AMINO ACIDS AND PEPTIDES; CHEMICAL STRUCTURE OF PROTEINS
- 18a. *Amino acids and their derivatives*
- 572 Araki, S., Abe, S., Yamada, S., Satake, M., Fujiwara, N., Kon, K. and Ando, S.: Characterization of two novel pyruvylated glycosphingolipids containing 2'-aminoethylphosphoryl(+6)-galactose from the nervous system of *Aplysia kurodai*. *J. Biochem. (Tokyo)*, 112 (1992) 461-469.
- 573 Bruckner, H. and Strecker, B.: Chiral monochloro-s-triazines as derivatizing reagents for resolving DL-amino acids by HPLC. *Chromatographia*, 33 (1992) 586-587.
- 574 Brueckner, H., Langer, M. and Godel, H.: LC-determination of D- and L-amino acids in peptide hydrolyzates by pre-column derivatization with a chiral reagent. In: Angeletti, R.H. (Editor), *Tech. Protein Chem. 3; [Pap. Annu. Symp. Protein Soc.]*, 5th 1991, Academic, San Diego, 1992, 243-247; C.A., 117 (1992) 65788s.
- 575 Cahnmann, H.J., Goncalves, E., Ito, Y., Fales, H.M. and Sokolowski, E.A.: Synthesis and properties of N-bromoacetyl-L-tyroxine. *Anal. Biochem.*, 204 (1992) 344-350.
- 576 Canevari, L., Vieira, R., Aldegunde, M. and Dagani, F.: High-performance liquid chromatographic separation with electrochemical detection of amino acids focusing on neurochemical application. *Anal. Biochem.*, 205 (1992) 137-142.
- 577 Chorbanov, B., Iliev, I., Borresen, T. and Adler-Nissen, J.: Gel chromatography of protein hydrolyzate with high degree of hydrolysis. *Dokl. Bulg. Akad. Nauk*, 44 (1991) 45-47; C.A., 117 (1992) 65789t.
- 578 Ci, Y.-X., Tie, J.-K., Wang, Q.-W. and Chang, W.-B.: Flow injection and liquid chromatographic postcolumn detection of amino acids by mimetic peroxidase-catalysed chemiluminescence reaction. *Anal. Chim. Acta*, 269 (1992) 109-114.
- 579 Clancy, R.M. and Abramson, S.B.: Novel synthesis of S-nitrosoglutathione and degradation by human neutrophils. *Anal. Biochem.*, 204 (1992) 365-371.
- 580 Csapo, J. and Henics, Z.: Quantitative determination of bacterial protein from the diamino pimelic acid and D-alanine content of rumen liquor and intestines. *Acta Agron. Hung.*, 40 (1991) 159-173; C.A., 117 (1992) 86084z.
- 581 Dou, L. and Krull, I.S.: Identification of photochemical products of amino acids, peptides, and proteins in on-line, postcolumn photolytic derivatization detection by HPLC-electrochemistry. *Electroanalysis (N.Y.)*, 4 (1992) 381-391; C.A., 117 (1992) 107738h.
- 582 El-Sayed Metwally, M. and Belal, F.F.: Application of high-performance liquid chromatography in the kinetic study of α -methyl-dopa. *J. Chromatogr.*, 623 (1992) 75-82.
- 583 Epstein, W.W., Lever, D., Leining, L.M., Bruenger, E. and Rilling, H.C.: Quantitation of prenylcysteines by a selective cleavage reaction. *Proc. Natl. Acad. Sci. U.S.A.*, 88 (1991) 9668-9670; C.A., 117 (1992) 43730q.
- 584 Giron, J., Alaiz, M. and Vioque, E.: High-performance liquid chromatographic determination of N- ϵ -(2-propenyl)lysine in biological samples after derivatization with diethylethoxymethylene-malonate. *Anal. Biochem.*, 206 (1992) 155-160.
- 585 Henle, T., Walter, H., Krause, I. and Klostermeyer, H.: Efficient determination of individual Maillard compounds in heat-treated milk products by amino acid analysis. *Int. Dairy J.*, 1 (1992) 125-135; C.A., 117 (1992) 24857v.
- 586 Hirano, H., Komatsu, S., Takakura, H., Sakiyama, F. and Tsunawasa, S.: Deblocking and subsequent microsequence analysis of N^F-blocked proteins electroblotted onto PVDF membranes. *J. Biochem. (Tokyo)*, 111 (1992) 754-757.
- 587 Horz, K.H. and Reichling, J.: (Ion-pair chromatography of amino acids). *Bioforum*, 15 (1992) 155-159; C.A., 117 (1992) 86104f.
- 588 Huber, W.: (Determination of aminopolycarboxylic acids by ion pair chromatography). *Z. Wasser Abwasser Forsch.*, 25 (1992) 6-8; C.A., 117 (1992) 55503f.
- 589 Ikarashi, Y. and Maruyama, Y.: Identification of O-methyl-dopa in the ventricular fluid of patients with Parkinson's disease. *Biol. Mass Spectrom.*, 21 (1992) 380-384; C.A., 117 (1992) 127762q.
- 590 Kaur, P., Mundhara, G.L., Kar, H.S. and Tiwari, J.S.: Sorption-desorption behavior of the amino acids, aspartic acid and glutamic acid on chemically pretreated alumina in relation to chromatography. In: Mersmann, A.B. and Scholl, S.E. (Editors), *Int. Conf. Fundam. Adsorpt.*, 3rd 1989, AIChE, New York, 1991, pp. 379-394; C.A., 117 (1992) 119108r.
- 591 Landry, J., Delhaye, S. and Jones, D.G.: Determination of tryptophan in feedstuffs: comparison of two methods of hydrolysis prior to HPLC analysis. *J. Sci. Food Agric.*, 58 (1992) 439-441; C.A., 117 (1992) 68581e.
- 592 Maskos, Z., Rush, J.D. and Koppenol, W.H.: The hydroxylation of tryptophan. *Arch. Biochem. Biophys.*, 296 (1992) 514-520.
- 593 Min, S., Yisheng, Y. and Lu, Y.: Determination of urinary 3-methylhistidine by high-performance liquid chromatography with o-phthalaldehyde precolumn derivatization. *J. Chromatogr.*, 581 (1992) 272-276.
- 594 Miyata, T.: (Pancreatic function diagnosis (PFD) with p-aminosalicylic acid (PAS) load test of exocrine pancreatic function test. Study on the measurement of PABA and PAS in urine using high performance liquid chromatography). *Ochanomizu Igaku Zasshi*, 39 (1991) 113-122; C.A., 117 (1992) 127541s.
- 595 Molnár-Perl, I. and Morvai, M.: Cystine and cysteine analysis as the same phenylthiocarbonyl derivatives by HPLC in protein hydrolyzates. *Chromatographia*, 34 (1992) 132-136.
- 596 Nakamura, F., Yamazaki, K. and Suyama, K.: Isolation and structural characterization of a new crosslinking amino acid, cyclopentenosine, from the acid hydrolysate of elastin. *Biochem. Biophys. Res. Commun.*, 186 (1992) 1533-1538.
- 597 Nakazawa, M. and Manabe, K.: The direct hydrolysis of proteins containing tryptophan on polyvinylidene difluoride membranes by mercaptoethanesulfonic acid in the vapor phase. *Anal. Biochem.*, 206 (1992) 105-108.

- 598 Nokihara, K., Morita, N. and Kuriki, T.: Applications of an automated apparatus for two-dimensional electrophoresis, Model TEP-1, for microsequence analyses of proteins. *Electrophoresis (Weinheim)*, 13 (1992) 701-707.
- 599 Nomoto, H., Yasukawa, K. and Inoue, Y.: Separation of neutral glycoasparagines according to their content of cis diol groups. *Biosci., Biotechnol., Biochem.*, 56 (1992) 1090-1095; C.A., 117 (1992) 146454g.
- 600 Oates, M.D.: Amino acid analysis by microcolumn liquid chromatography. Avail. *Univ. Microfilms Int.*, Order No. DA9106130, 1990, 199 p.; C.A., 117 (1992) 184033b.
- 601 Orwar, O., Folestad, S. and Sandberg, M.: Determination of attomole quantities of neuroactive acidic sulfur-containing amino acids and γ -glutamyl peptides using argon ion laser-induced fluorescence detection and microcolumn liquid chromatography. In: Rollema, H., Westerink, B.H.C. and Drijfhout, W.J. (Editors), *Monit. Mol. Neurosci., Proc. Int. Conf. In Vivo Methods, 5th 1991*, Univ. Cent. Pharm., Groningen, 1991, pp. 125-127; C.A., 117 (1992) 127512h.
- 602 Perwnim, T., Miles, C. and Seifert, J.: A new method of L-kynurenine precolumn derivatization with naphthalene-2,3-dicarboxaldehyde and cyanide for high-performance liquid chromatography (HPLC). *Fresenius J. Anal. Chem.*, 343 (1992) 907-908.
- 603 Pochapsky, T.C. and Gopen, Q.: A chromatographic approach to the determination of relative free energies of interaction between hydrophobic and amphiphilic amino acid side chains. *Protein Sci.*, 1 (1992) 786-795; C.A., 117 (1992) 127524p.
- 604 Puchala, R., Piór, H., Kulasek, G.W. and Shelford, J.A.: Determination of diaminopimelic acid in biological materials using high-performance liquid chromatography. *J. Chromatogr.*, 623 (1992) 63-67.
- 605 Rajpurohit, R., Paik, W.K. and Kim, S.: Enzymatic methylation of heterogeneous nuclear ribonucleoprotein in isolated liver nuclei. *Biochim. Biophys. Acta*, 1122 (1992) 183-188.
- 606 Shah, M.A., Bergethon, P.R., Boak, A.M., Gallop, P.M. and Kagan, H.M.: Oxidation of peptidyl lysine by copper complexes of pyrroloquinoline quinone and other quinones. A model for oxidative pathochemistry. *Biochim. Biophys. Acta*, 1159 (1992) 311-318.
- 607 Slight, S.H., Prabhakaram, M., Shin, D.B., Feather, M.S. and Ortwerth, B.J.: The extent of N^ε-(carboxymethyl)lysine formation in lens proteins and polylysine by the autoxidation products of ascorbic acid. *Biochim. Biophys. Acta*, 1117 (1992) 199-206.
- 608 Turck, C.W.: Identification of phosphotyrosine residues in peptides by high performance liquid chromatography on-line derivative spectroscopy. *Pept. Res.*, 5 (1992) 156-160; C.A., 117 (1992) 107742e.
- 609 Vijayalakshmi, M.A.: Histidine ligand affinity chromatography. *Methods Mol. Biol. (Totowa)*, 11(Pract. Protein Chromatogr.) (1992) 33-44; C.A., 117 (1992) 166907k.
- 610 White, J.A. and Hart, R.J.: HPLC analysis of amino acids. *Food Sci. Technol.*, 52(Food Anal. HPLC) (1992) 75-115; C.A., 117 (1992) 110169d - a review with 142 refs.
- 611 White, J.A. and Hart, R.J.: Derivatization methods for liquid chromatographic separation of amino acids. *Food Sci. Technol.*, 52(Food Anal. HPLC) (1992) 53-74; C.A., 117 (1992) 110168c - a review with 74 refs.
- 612 Zukowski, J., Pawlowska, M. and Armstrong, D.W.: Efficient enantioselective separation and determination of trace impurities in secondary amino acids (i.e., imino acids). *J. Chromatogr.*, 623 (1992) 33-41.

For additional information see C.A.:

117 (1992) 22638a, 33792x, 33793y, 43741u, 66060k, 86342g, 169704c.

See also 149, 216, 384, 512, 541, 679, 682, 909, 1130, 1140, 1525, 1532, 1537.

18b. Peptides, peptidic and proteinous hormones, growth factors

- 613 Aguilar, M.I., Wilce, M.C.J., Round, A.J. and Hearn, M.T.W.: Surface mapping of peptides by computer graphics and HPLC. In: Smith, J.A. and Rivier, J.E. (Editors), *Pept.: Chem. Biol., Proc. Am. Pept. Symp., 12th 1991*, ESCOM, Leiden, 1992, pp. 580-582; C.A., 117 (1992) 127737k.
- 614 Amano, T., Okita, Y., Okinaga, T., Matsui, T., Nishiyama, I. and Hoshi, M.: ECG jelly component responsible for histone degradation and aerosome reaction in the starfish, *Asterina pectinifera*. *Biochem. Biophys. Res. Commun.*, 187 (1992) 274-278.
- 615 Bogerd, J., Li, K.W., Janssen-Dommerholt, C. and Goos, H.: Two gonadotropin-releasing hormones from african catfish (*Clarias gariepinus*). *Biochem. Biophys. Res. Commun.*, 187 (1992) 127-134.
- 616 Bulet, P., Cociancich, S., Reuland, M., Sauber, F., Bischoff, R., Hegy, G., van Dorsselaer, A., Hetru, C. and Hoffmann, J.A.: A novel insect defensin mediates the inducible antibacterial activity in larvae of the dragonfly *Aeschna cyanea* (Bleoptera, Odonata). *Eur. J. Biochem.*, 209 (1992) 977-984.
- 617 Buttner, K., Blondelle, S.E., Ostresh, J.M. and Houghten, R.A.: Perturbation of peptide conformations induced in anisotropic environments. *Biopolymers*, 32 (1992) 575-583; C.A., 117 (1992) 86270g.
- 618 Conlon, J.M., O'Harte, F., Smith, D.D., Tonon, M.-C. and Vaudry, H.: Isolation and primary structure of urotensin II from the brain of a tetrapod, the frog *Rana ridibunda*. *Biochem. Biophys. Res. Commun.*, 188 (1992) 578-583.
- 619 Crimmins, D.L.: Comparative analysis of synthetic peptides by free-solution capillary electrophoresis (FSCE) and strong-cation exchange (SCX)-HPLC. In: Angeletti, R.H. (Editor), *Tech. Protein Chem. 3, [Pap. Annu. Symp. Protein Soc.]*, 5th 1991, Academic, San Diego, 1992, pp. 171-181; C.A., 117 (1992) 65787r.
- 620 Davis, M.T. and Lee, T.D.: Analysis of peptide mixtures by capillary high performance liquid chromatography: A practical guide to small-scale separations. *Protein Sci.*, 1 (1992) 935-944; C.A., 117 (1992) 167118r.
- 621 Dorizzi, R., Michelot, D., Tagliaro, F. and Ghielmi, S.: Methods for chromatographic determination of amanitins and related toxins in biological samples. *J. Chromatogr.*, 580 (1992) 279-291 - a review with 55 refs.
- 622 Duvick, J.P., Rood, T., Rao, A.G. and Marshak, D.R.: Purification and characterization of a novel antimicrobial peptide from maize (*Zea mays* L.) kernels. *J. Biol. Chem.*, 267 (1992) 18814-18820.

- 623 Emly, J.F., Ratcliffe, W.A., Green, E., Bowden, S.J., Heath, D.A., Blight, A., Hughes, S. and Ratcliffe, J.G.: Immunochemical characterisation of parathyroid hormone-related protein from tumour and non-tumour cells. *Biochim. Biophys. Acta*, 1180 (1992) 58-64.
- 624 Fassina, G., Consonni, R., Zetta, L. and Cassani, G.: Design of hydrophatically complementary peptides for big endothelin affinity purification. *Int. J. Pept. Protein Res.*, 39 (1992) 540-548; C.A., 117 (1992) 164002n.
- 625 Fialaire, A., Postaire, E., Prognon, P., Pradier, F. and Pradeau, D.: Thermal decomposition of reduced glutathione in solution for organ preservation. *J. Pharm. Biomed. Anal.*, 10 (1992) 457-460.
- 626 Fulton, S., Meys, M., Protentis, J., Afeyan, N.B., Carlton, J. and Haycock, J.: Preparative peptide purification by cation-exchange and reversed-phase perfusion chromatography. *Bio-Techniques*, 12 (1992) 742-744; C.A., 117 (1992) 127509n.
- 627 Greenwald, J.E., Needleman, P., Siegel, N., Tetens, E., Biel, B. and Ritter, D.: Processing of atriopeptin prohormone by non-myocytic atrial cells. *Biochem. Biophys. Res. Commun.*, 188 (1992) 644-654.
- 628 Hasegawa, Y., Jette, D.C., Miyamoto, A., Kawasaki, H. and Yuki, H.: Chemiluminescent determination of amino-terminal residues of peptides after derivatization of anilinothiazolinone from Edman degradation with N-(4-aminoheptyl)-N-ethylisoluminol. *Anal. Sci.*, 7 (1992) 945-948; C.A., 117 (1992) 142602f.
- 629 Heath, W.F., Belagaje, R.M., Brooke, G.S., Chance, R.E., Hoffmann, J.A., Long, H.B., Reams, S.G., Roundtree, C., Shaw, W.N., Sliker, L.J. et al.: (A-C-B) Human proinsulin, a novel insulin agonist and intermediate in the synthesis of biosynthetic human insulin. *J. Biol. Chem.*, 267 (1992) 419-425.
- 630 Hermann, K., Rittweger, R., Phillips, M.I. and Ring, J.: Presence of angiotensin peptides in human urine. *Clin. Chem. (Winston-Salem)*, 38 (1992) 1768-1772.
- 631 Hirayama, M., Toyota, K., Yamaguchi, G., Hidaka, H. and Naito, H.: HPLC analysis of commercial casein phosphopeptides (CPP). *Biosci., Biotechnol., Biochem.*, 56 (1992) 1126-1127; C.A., 117 (1992) 169689b.
- 632 Ichiki, Y., Kitamura, K., Kangawa, K., Kawamoto, M., Matsuo, H. and Eto, T.: Organ distribution and characterization of porcine peptides (VIP, CGRP and PHI) that increase cAMP in rat platelets. *Biochem. Biophys. Res. Commun.*, 187 (1992) 1587-1593.
- 633 Kálmán, F., Cserhádi, T., Valkó, K. and Neubert, K.: Relationships between the reversed-phase liquid chromatographic retention characteristics and physico-chemical parameters of some β -casomorphin peptides. *Anal. Chim. Acta*, 268 (1992) 247-254.
- 634 Kohama, Y., Okabe, M., Tsujikawa, K., Oka, H., Teramoto, T., Kayamori, Y., Itoh, M. and Mimura, T.: Preparation of biologically active and site specifically radioiodinated recombinant human insulin-like growth factor-I. *Chem. Pharm. Bull.*, 40 (1992) 808-810.
- 635 Krull, I.S. and Dou, L.: Determination of peptides and proteins using on-line, post-column photolytic derivatization in liquid chromatography/electrochemistry (LC-hv-EC). *Curr. Sep.*, 11 (1992) 7-11; C.A., 117 (1992) 166904g.
- 636 Lakhani, S., Khanna, N.C. and Tewari, K.K.: Two distinct transcriptional activities of pea (*Pisum sativum*) chloroplasts share immunochemically related functional polypeptides. *Biochem. J.*, 286 (1992) 833-841.
- 637 Lundell, N. and Markides, K.: Two-dimensional liquid chromatography of peptides: an optimization strategy. *Chromatographia*, 34 (1992) 369-375.
- 638 Luo, P. and Baldwin, R.P.: LCEC of underivatized polypeptides and proteins at copper electrodes. *Electroanalysis (N.Y.)*, 4 (1992) 393-401; C.A., 117 (1992) 127729j.
- 639 Lupu, R., Wellstein, A., Sheridan, J., Ennis, B.W., Zugmaier, G., Katz, D., Lippman, M.E. and Dickson, R.B.: Purification and characterization of a novel growth factor from human breast cancer cells. *Biochemistry*, 31 (1992) 7330-7340.
- 640 Miyata, A., Jiang, L., Minamino, N. and Arimura, A.: Identification of calcitonin gene related polypeptide in ovine hypothalamic extract. *Biochem. Biophys. Res. Commun.*, 187 (1992) 1474-1479.
- 641 Nakamura, T., Asashima, M., Eto, Y., Takio, K., Uchiyama, H., Moriya, N., Ariizumi, T., Yashiro, T., Sugino, K., Titani, K. and Sugino, H.: Isolation and characterization of native activin B. *J. Biol. Chem.*, 267 (1992) 16385-16389.
- 642 Nanami, M., Zaitzu, K. and Ohkura, Y.: Preparation of S-acetylthioglycolyl insulins based on separation by anion-exchange high-performance liquid chromatography. *Chem. Pharm. Bull.*, 40 (1992) 957-960.
- 643 Ng, N.F.L. and Hew, C.L.: Structure of an antifreeze polypeptide from the sea raven. Disulfide bonds and similarity to lectin-binding proteins. *J. Biol. Chem.*, 267 (1992) 16069-16075.
- 644 Noguchi, K. and Higuchi, S.: Determination of glutathione isopropyl ester in rat, dog and human blood by high-performance liquid chromatography with fluorescence detection. *J. Pharm. Biomed. Anal.*, 10 (1992) 515-520.
- 645 Page, M.D., Edwards, C.A., Brown, R., Weeks, I., Woodhead, J.S. and Scanlon, M.F.: Immunochemiluminometric assays (ICMA) specific for growth hormone releasing hormone 1044 NH₂ and 1-40 OH. *Clin. Chim. Acta*, 210 (1992) 167-178.
- 646 Polo, M.C., Gonzalez de Llano, D. and Ramos, M.: Derivatization and liquid chromatographic separation of peptides. *Food Sci. Technol.*, 52 (Food Anal. HPLC) (1992) 117-140; C.A., 117 (1992) 110170x - a review with 128 refs.
- 647 Rao, U.S., Bauzon, D.D. and Scarborough, G.A.: Cytoplasmic location of amino acids 359-440 of the *Neurospora crassa* plasma membrane H⁺-ATPase. *Biochim. Biophys. Acta*, 1108 (1992) 153-158.
- 648 Rose, K., Regamey, P.-O., Anderegg, R., Wells, T.N.C. and Proudfoot, A.E.I.: Human interleukin-5 expressed in *Escherichia coli* has N-terminal modifications. *Biochem. J.*, 286 (1992) 825-828.
- 649 Sasaoki, K., Hiroshima, T., Kusumoto, S. and Nishi, K.: Deamidation at asparagine-88 in recombinant human interleukin 2. *Chem. Pharm. Bull.*, 40 (1992) 976-980.
- 650 Schnölzer, M., Jones, A., Alewood, P.F. and Kent, S.B.H.: Ion-spray tandem mass spectrometry in peptide synthesis: structural characterization of minor by-products in the synthesis of ACP(65-74). *Anal. Biochem.*, 204 (1992) 335-343.
- 651 Schweitz, H., Vigne, P., Moinier, D., Frelin, C. and Lazdunski, M.: A new member of the natriuretic peptide family is present in the venom of the green mamba (*Dendroaspis angusticeps*). *J. Biol. Chem.*, 267 (1992) 13928-13932.

- 652 Shiomi, K., Nakazato, M., Miyazato, M., Kangawa, K., Matsuo, H. and Matsukura, S.: Establishment of hypersensitive radioimmunoassay for islet amyloid polypeptide using antiserum specific for its N-terminal region. *Biochem. Biophys. Res. Commun.*, 186 (1992) 1065-1073.
- 653 Simon, D., Romestand, B., Huang, H., Badouaille, G., Fehrentz, J.-A., Pau, B., Marchand, J. and Corvol, P.: Direct, simplified, and sensitive assay of angiotensin II in plasma extracts performed with a high-affinity monoclonal antibody. *Clin. Chem. (Winston-Salem)*, 38 (1992) 1963-1967.
- 654 Soifer, N.E., Dee, K.E., Insogna, K.L., Burtis, W.J., Matovcik, L.M., Wu, T.L., Milstone, L.M., Broadus, A.E., Philbrick, W.M. and Stewart, A.F.: Parathyroid hormone-related protein. Evidence for secretion of a novel mid-region fragment by three different cell types. *J. Biol. Chem.*, 267 (1992) 18236-18243.
- 655 Stevenson, C.L., Williams, T.D., Anderegg, R.J. and Borchardt, R.T.: Identification and quantitation of tetrapeptide deamidation products by mass spectrometry. *J. Pharm. Biomed. Anal.*, 10 (1992) 567-575.
- 656 Suzuki, M., Itoh, T., Suzuki, T., Koga, N., Kato, K., Saito, T. and Mitsui, Y.: Identification of the hepatocyte mitogen in bovine spleen as heparin-binding growth factors. *Biochem. Biophys. Res. Commun.*, 186 (1992) 1192-1200.
- 657 Toomik, R., Ekman, P. and Engström, L.: A potential pitfall in protein kinase assay: phosphocellulose paper as an unreliable adsorbent of produced phosphopeptides. *Anal. Biochem.*, 204 (1992) 311-314.
- 658 Vallenweider, F., Irminger, J.-C., Gross, D.J., Villa-Komaroff, L. and Halban, P.A.: Processing of proinsulin by transfected hepatoma (FAO) cells. *J. Biol. Chem.*, 267 (1992) 14629-14636.
- 659 Wang, Y., Badgery-Parker, T., Lovas, S., Chartrel, N., Vaudry, H., Burcher, E. and Conlon, J.M.: Primary structure and receptor-binding properties of a neurokinin related peptide from frog gut. *Biochem. J.*, 287 (1992) 827-832.
- 660 Weitzmann, M.N. and Savage, N.: Nuclear internalisation and DNA binding activities of interleukin-1, interleukin-1 receptor and interleukin-1/receptor complexes. *Biochem. Biophys. Res. Commun.*, 187 (1992) 1166-1171.
- 661 Yasufuku, K., Ohashi, H., Katsuta-Enomoto, Y., Fukuroda, T., Noguchi, K. and Yano, M.: High-yield production of recombinant endothelin-1. *J. Biochem. (Tokyo)*, 112 (1992) 360-365.
- 662 Yasunaga, F., Kawano, A., Morikawa, N. and Takeyama, M.: Enzyme immunoassay of angiotensin II in human plasma. *Chem. Pharm. Bull.*, 40 (1992) 792-794.
- 663 Zhu, B.Y., Mant, C.T. and Hodges, R.S.: Mixed-mode hydrophilic and ionic interaction chromatography rivals reversed-phase chromatography for separation of peptides. In: Smith, J.A. and River, J.E. (Editors), *Pept.: Chem. Biol., Proc. Am. Pept. Symp., 12th 1991*, ESCOM, Leiden, 1992, pp. 546-548; *C.A.*, 117 (1992) 142589g.
- For additional information see *C.A.*:
117 (1992) 22619v, 22620p, 143623a.
- See also 339, 581, 597, 601, 682, 712, 772, 812, 902, 1018, 1472, 1480, 1536, 1560.
- 18c. *Elucidation of structure of proteins and enzymes*
- 664 Bongers, J., Lambros, T., Ahmad, M. and Heimer, E.P.: Kinetics of dipeptidyl peptidase IV proteolysis of growth hormone-releasing factor and analogs. *Biochim. Biophys. Acta*, 1122 (1992) 147-153.
- 665 Buck, F., Luth, C., Strupat, K. and Bretting, H.: Comparative investigations on the amino-acid sequences of different isolectins from the sponge *Axinella polypoides* (Schmidt). *Biochim. Biophys. Acta*, 1159 (1992) 1-8.
- 666 Choi-Miura, N.H., Takahashi, Y., Nakano, Y., Tobe, T. and Tomita, M.: Identification of the disulfide bonds in human plasma protein SP-40,40 (apolipoprotein-J). *J. Biochem. (Tokyo)*, 112 (1992) 557-561.
- 667 Edel, C.M., Hartog, A.F. and Berden, J.A.: Inhibition of mitochondrial F₁-ATPase activity by binding of (2-azido)-ADP to a slowly exchangeable non-catalytic nucleotide binding site. *Biochim. Biophys. Acta*, 1101 (1992) 329-338.
- 668 Hasegawa, Y., Ueda, Y., Watanabe, M. and Morita, F.: Studies on amino acid sequences of two isoforms of 17-kDa essential light chain of smooth muscle myosin from porcine aorta media. *J. Biochem. (Tokyo)*, 111 (1992) 798-803.
- 669 Hori, K. and Morita, F.: Actin-actin contact: inhibition of actin-polymerization by subdomain 4 peptide fragments. *J. Biochem. (Tokyo)*, 112 (1992) 401-408.
- 670 Hu, P.-S., Persson, B., Höög, J.-O., Jörnvall, H., Hartog, A.F., Berden, J.A., Holmberg, E. and Rydstrom, J.: Energy-linked transhydrogenase. Characterization of a nucleotide-binding sequence in nicotinamide nucleotide transhydrogenase from beef heart. *Biochim. Biophys. Acta*, 1102 (1992) 19-29.
- 671 Lendaro, E., Ippoliti, R., Brancaccio, A., Bellelli, A., Vallone, B., Ivaldi, G., Sciaratta, G.V., Castello, C., Tomova, S., Brunori, M. and Amiconi, G.: Hemoglobin Dallas ($\alpha 97$ (G4)Asn \rightarrow Lys): functional characterization of a high oxygen affinity natural mutant. *Biochim. Biophys. Acta*, 1180 (1992) 15-20.
- 672 Lin, S.-R. and Chang, C.-C.: Chemical modification of arginine residues in α -bungarotoxin. *Biochim. Biophys. Acta*, 1159 (1992) 255-261.
- 673 Moritz, R.L., Ward, L.D. and Simpson, R.J.: High-sensitivity peptide mapping utilizing reversed-phase microbore and microcolumn liquid chromatography. In: Angeletti, R.H. (Editor), *Tech. Protein Chem. 3, [Pap. Annu. Symp. Protein Soc.]*, 5th 1991, Academic, San Diego, 1992, pp. 97-106; *C.A.*, 117 (1992) 66002t.
- 674 Ortwerth, B.J., Slight, S.H., Prabhakaram, M., Sun, Y. and Smith, J.B.: Site-specific glycation of lens crystallins by ascorbic acid. *Biochim. Biophys. Acta*, 1117 (1992) 207-215.
- 675 Rua, J., Robertson, A.G.S. and Nimmo, H.G.: Identification of the histidine residue in *Escherichia coli* isocitrate lyase that reacts with diethylpyrocarbonate. *Biochim. Biophys. Acta*, 1122 (1992) 212-218.
- 676 Tao, H.P., Adalsteins, A.E. and Kasarda, D.D.: Intermolecular disulfide bonds link specific high-molecular-weight glutenin subunits in wheat endosperm. *Biochim. Biophys. Acta*, 1159 (1992) 13-21.
- 677 Van Breemen, R.B. and Davis, R.G.: Rates of peptide proteolysis measured using liquid chromatography and continuous-flow fast atom bombardment mass spectrometry. *Anal. Chem.*, 64 (1992) 2233-2237.

- 678 Wajcman, H., Kister, J., Marden, M., Lahary, A., Monconduit, M. and Galacteros, F.: Hemoglobin Rouen (α -140 (HC2) Tyr \rightarrow His): alteration of the α -chain C-terminal region and moderate increase in oxygen affinity. *Biochim. Biophys. Acta*, 1180 (1992) 53-57.
- 679 Watanabe, M., Hasegawa, Y., Katoh, T. and Morita, F.: Amino acid sequence of the 20-kDa regulatory light chain of porcine aorta media smooth muscle myosin. *J. Biochem. (Tokyo)*, 112 (1992) 431-432.
- See also 385, 608, 647, 736, 744, 770, 776, 780, 792, 798, 799, 801, 803, 812, 823, 884, 896, 928, 933, 934, 936, 948, 950, 959, 964, 990, 991, 1018, 1027, 1040, 1044, 1045, 1051, 1056, 1063, 1065.
19. PROTEINS
- 19a. General techniques
- 680 Afeyan, N.B., Gordon, N.F., Fulton, S.P. and Regnier, F.E.: Perfusion chromatography: a novel tool for protein purification and analysis. In: Angeletti, R.H. (Editor), *Tech. Protein Chem. 3, [Pap. Annu. Symp. Protein Soc.]*, 5th 1991, Academic, San Diego, 1992, pp. 135-149; C.A., 117 (1992) 65786q.
- 681 Battey, N.H. and Venis, M.A.: Calcium-dependent hydrophobic interaction chromatography. *Methods Mol. Biol. (Totowa)*, 11(Pract. Protein Chromatogr.) (1992) 73-80; C.A., 117 (1992) 166908m.
- 682 Boyd, V.L., Bozzini, M.L., Zon, G., Noble, R.L. and Mattaliano, R.J.: Sequencing of peptides and proteins from the carboxy terminus. *Anal. Biochem.*, 206 (1992) 344-352.
- 683 Burton, S.J.: Dye-ligand affinity chromatography. *Methods Mol. Biol. (Totowa)*, 11(Pract. Protein Chromatogr.) (1992) 91-103; C.A., 117 (1992) 146391j - a review with 13 refs.
- 684 Chervet, J.P. and Ursem, M.: (The role of micro-HPLC in the analysis of proteins). *GIT Spec. Chromatogr.*, 12 (1992) 38-44; C.A., 117 (1992) 61930j.
- 685 Cox, J.A. and Przyjazny, A.: Electrochemistry of biopolymers. *TrAC*, 11 (1992) 298-302 - a review with 40 refs.
- 686 Donovan, J., Rabel, F. and Zahran, J.: Protein separations on tentacle ion exchangers. Part II: scale-up and loading study. *Am. Biotechnol. Lab.*, 10 (1992) 38-40; C.A., 117 (1992) 166903f.
- 687 Harris, D.A.: Size-exclusion high-performance liquid chromatography of proteins. *Methods Mol. Biol. (Totowa)*, 11(Pract. Protein Chromatogr.) (1992) 223-236; C.A., 117 (1992) 166912h.
- 688 Hochuli, E.: Purification of histidine fusion proteins by metal chelate chromatography. In: Christiansen, C., Munck, L. and Villadsen, J. (Editors), *Proc. - Eur. Congr. Biotechnol.*, 5th 1990, Munksgaard, Copenhagen, 1990, pp. 1005-1009; C.A., 117 (1992) 146359e - a review with 15 refs.
- 689 Hoffman, R.D. and Lane, M.D.: Iodophenylarsine oxide and arsenical affinity chromatography: new probes for dithiol proteins. Application to tubulins and to components of the insulin receptor-glucose transporter signal transduction pathway. *J. Biol. Chem.*, 267 (1992) 14005-14011.
- 690 Hutchens, T.W.: Thiophilic adsorption chromatography. *Methods Mol. Biol. (Totowa)*, 11(Pract. Protein Chromatogr.) (1992) 1-15; C.A., 117 (1992) 168937u - a review with 22 refs.
- 691 Inoue, S. and Ohtaki, N.: Influence of chromatographic conditions on binding capacities of hydroxyapatite for globular proteins. *Anal. Sci.*, 8 (1992) 361-365; C.A., 117 (1992) 65792p.
- 692 Kenney, A. and Fowell, S. (Editors): *Practical Protein Chromatography. Methods Mol. Biol.*, Humana Press, Totowa, 1992, 327 p.; C.A., 117 (1992) 146688m.
- 693 Kenney, A.C.: Ion-exchange chromatography of proteins. *Methods Mol. Biol. (Totowa)*, 11(Pract. Protein Chromatogr.) (1992) 249-258; C.A., 117 (1992) 166914k.
- 694 Kuboi, R., Mori, Y., Yamada, Y., Komasa, I. and Luisi, P.L.: Separation of biopolymers with reverse micellar bulk liquid membranes. *Process Metall.*, 7B(Solvent Extr. 1990, Pt. B) (1992) 1863-1868; C.A., 117 (1992) 146569y.
- 695 Levison, P.R., Badger, S.E., Toome, D.W., Carcary, D. and Butts, E.T.: Use of anion-exchange cellulose for process-scale protein separation. In: Abe, M., Kataoka, T. and Suzuki, T. (Editors), *New Dev. Ion Exch. Proc. Int. Conf. Ion Exch.*, Kodansha, Tokyo, 1991, pp. 597-600; C.A., 117 (1992) 110023b.
- 696 Li, C.M. and Hutchens, T.W.: Chromatofocusing. *Methods Mol. Biol. (Totowa)*, 11(Pract. Protein Chromatogr.) (1992) 237-248; C.A., 117 (1992) 166913j.
- 697 Mhatre, R.M. and Krull, I.S.: Fast protein separation and characterization by hydrophobic interaction chromatography (HIC) and low angle laser light scattering photometry (LALLS). *Chromatographia*, 34 (1992) 357-362.
- 698 Porath, J.: Immobilized metal ion affinity chromatography. *Protein Expression Purif.*, 3 (1992) 263-281; C.A., 117 (1992) 146398s - a review with 77 refs.
- 699 Regnier, F.E., DeStefano, J.J., Hearn, M.T.W., Janson, J.C., Osterlund, B.R. and Unger, K.K. (Editors): *11th International Symposium on High-Performance Liquid Chromatography of Proteins, Peptides and Polynucleotides*, Washington, October 20-23, 1991. *J. Chromatogr.*, Vol. 599, No. 1-2, Elsevier, Amsterdam, 1992, 274 p.; C.A., 117 (1992) 66041e.
- 700 Renken, E., Kaaerd, A. and Kasche, V.: Fluorescence of proteins adsorbed to chromatographic supports: application in studies on adsorption-desorption kinetics. In: Mersmann, A.B. and Scholl, S.E. (Editors), *Int. Conf. Fundam. Adsorpt.*, 3rd 1989, AIChE, New York, 1991, pp. 715-722; C.A., 117 (1992) 127514k.
- 701 Stamm, W.W. and Kula, M.R.: Automated chromatography of proteins with product specific control based on online monitoring of enzymic activity. In: Christiansen, C., Munck, L. and Villadsen, J. (Editors), *Proc.-Eur. Congr. Biotechnol.*, 5th 1990, Munksgaard, Copenhagen, 1990, pp. 1032-1035; C.A., 117 (1992) 145803h.
- 702 Stewart, D.J., Purvis, D.R., Pitts, J.M. and Lowe, C.R.: Development of an enzyme-linked immunosorbent assay for C.I. Reactive Blue 2 and its application to a comparison of the stability and performance of a perfluorocarbon support with other immobilised C.I. Reactive Blue 2 affinity adsorbents. *J. Chromatogr.*, 623 (1992) 1-14.
- 703 Swiderek, K.M., Chen, S., Feistner, G.J., Shively, J.E. and Lee, T.D.: Applications of liquid chromatography-electrospray mass spectrometry (LC-ES/MS). In: Angeletti, R.H. (Editor), *Tech. Protein Chem. 3, [Pap. Annu. Symp. Protein Soc.]*, 5th 1991, Academic, San Diego, 1992, pp. 457-465; C.A., 117 (1992) 66008z.

- 704 Szepeszy, L. and Rippel, G.: Comparison and evaluation of HIC columns of different hydrophobicity. *Chromatographia*, 34 (1992) 391-397.
- 705 Takahashi, Y. and Goto, S.: Continuous separation and concentration of proteins using an annular chromatograph. *J. Chem. Eng. Jpn.*, 25 (1992) 403-407; *C.A.*, 117 (1992) 166938w.
- 706 Tudor, R.: (Thiophilic adsorption chromatography - efficient method of protein fractionation). *Stud. Cercet. Biochim.*, 35 (1992) 51-63; *C.A.*, 117 (1992) 166862s - a review with 27 refs.
- 707 Vervaeck, J. and Huyghebaert, A.: HPLC of food proteins. *Food Sci. Technol.*, 52(Food Anal. HPLC) (1992) 141-167; *C.A.*, 117 (1992) 110171y - a review with 62 refs.
- 708 Williamson, K.C., Duffy, P.E. and Kaslow, D.C.: Immunoaffinity chromatography using electroelution. *Anal. Biochem.*, 206 (1992) 359-362.

For additional information see C.A.:
117 (1992) 146710n.

See also 128, 135, 181, 194, 383, 397, 581, 635, 638, 718, 1108.

19b. *Proteins of cells, viruses and subcellular particles*

- 709 Bertrand, C., Dumoulin, R., Divry, P., Mathieu, M. and Vianey-Saban, C.: Purification of electron transfer flavoprotein from pig liver mitochondria and its application to the diagnosis of deficiencies of acyl-CoA dehydrogenases in human fibroblasts. *Clin. Chim. Acta*, 210 (1992) 75-91.
- 710 Caldentey, J. and Bamford, D.H.: The lytic enzyme of the *Pseudomonas* phage $\phi 6$. Purification and biochemical characterization. *Biochim. Biophys. Acta*, 1159 (1992) 44-50.
- 711 Furukawa, T., Uchiumi, T., Tokunaga, R. and Taketani, S.: Ribosomal protein P2, a novel iron-binding protein. *Arch. Biochem. Biophys.*, 298 (1992) 182-186.
- 712 Kareem, B.N., Rokkones, E., Hogset, A., Holmgren, E. and Gautvik, K.M.: A method for the evaluation of the efficiency of signal sequences for secretion and correct N-terminal processing of human parathyroid hormone produced in *Escherichia coli*. *Anal. Biochem.*, 204 (1992) 26-33.
- 713 Oldfield, S. and Proun, C.G.: Purification, phosphorylation and control of the guanine-nucleotide-exchange factor from rabbit reticulocyte lysates. *Eur. J. Biochem.*, 208 (1992) 73-81.
- 714 Rose, S., Hensley, P., O'Shannessy, D.J., Culp, J., Debouck, C. and Chaiken, I.: Characterization of HIV-1 p24 self-association using analytical affinity chromatography. *Proteins: Struct., Funct., Genet.*, 13 (1992) 112-119; *C.A.*, 117 (1992) 145570e.
- 715 Sabourin, C.L.K. and Ley, R.D.: Purification and characterization of fibroblast growth factors from the opossum, *Monodelphis domestica*, brain. *Neurosci. Lett.*, 140 (1992) 148-152; *C.A.*, 117 (1992) 83543n.
- 716 Schuette, H., Papamichael, N., Paulsen, J. and Hustedt, H.: Integration of hydrophobic adsorption into protein purification schemes. In: Mersmann, A.B. and Scholl, S.E. (Editors), *Int. Conf. Fundam. Adsorpt.*, 3rd 1989, AIChE, New York, 1991, pp. 795-803; *C.A.*, 117 (1992) 127732e.
- 717 Tremblay, L., Gingras, D., Boivin, D. and Béliveau, R.: Tyrosine protein kinase activity in renal brush-border membranes. *Biochim. Biophys. Acta*, 1108 (1992) 183-189.

- 718 Yildiz, E., Grüber, G., Hörger, S., Zimmermann, H., Echner, H., Stoeva, S. and Voelter, W.: Application of capillary electrophoresis in peptide research. *Electrophoresis (Weinheim)*, 13 (1992) 683-686.
- 719 Zoon, K.C., Miller, D., Bekisz, J., zur Nedden, D., Enterline, J.C., Nguyen, N.Y. and Hu, R.: Purification and characterization of multiple components of human lymphoblastoid interferon- α . *J. Biol. Chem.*, 267 (1992) 15210-15216.

For additional information see C.A.:
117 (1992) 22653b, 22768t.

See also 860.

19c. *Proteins synthesized by genetic manipulation, monoclonal antibodies*

- 720 Baksh, S., Burns, K., Busaan, J. and Michalak, M.: Expression and purification of recombinant and native calreticulin. *Protein Expression Purif.*, 3 (1992) 322-331; *C.A.*, 117 (1992) 166922m.
- 721 Costa, C.A. and Cabral, J.S. (Editors): *Chromatographic and Membrane Processes in Biotechnology. (Proceedings of the NATO Advanced Study Institute on Chromatography and Membrane processes in Biotechnology, S. Mique - Acores, Portugal, July 15-27, 1990.) NATO ASI Ser., Ser. E: Applied Sciences, 1991, 204, Kluwer, Dordrecht, 1991, 473 pp.; C.A.*, 117 (1992) 86325d.
- 722 Gordon, M., Chokshi, H. and Alpers, D.H.: *In vitro* expression and secretion of functional mammalian intrinsic factor using recombinant baculovirus. *Biochim. Biophys. Acta*, 1132 (1992) 276-283.
- 723 Nachman, M., Azad, A.R.M. and Bailon, P.: Efficient recovery of recombinant proteins using membrane-based immunoaffinity chromatography (MIC). *Biotechnol. Bioeng.*, 40 (1992) 564-571; *C.A.*, 117 (1992) 166936u.
- 724 Ragin, R.C., Donahoe, P.K., Kenneally, M.K. and Ahmad, M.F.: Human Muellierian inhibiting substance: enhanced purification imparts biochemical stability and restores antiproliferative effects. *Protein Expression Purif.*, 3 (1992) 236-245; *C.A.*, 117 (1992) 83546r.
- 725 Stofko-Hahn, R., Carr, D.W. and Scott, J.D.: A single step purification for recombinant proteins: characterization of a microtubule associated protein (MAP2) fragment which associates with the type II cAMP-dependent protein kinase. *FEBS Lett.*, 302 (1992) 274-278; *C.A.*, 117 (1992) 63892x.

For additional information see C.A.:
117 (1992) 85742a, 107522h.

See also 661, 814, 845, 953.

19d. *Microbial and plant proteins*

- 726 Bietz, J.A. and Simpson, D.G.: Electrophoresis and chromatography of wheat proteins: available methods, and procedures for statistical evaluation of the data. *J. Chromatogr.*, 624 (1992) 53-80 - a review with 192 refs.
- 727 Borgmeyer, J.R., Smith, C.E. and Huynh, Q.K.: Isolation and characterization of a 25 kDa antifungal protein from flax seeds. *Biochem. Biophys. Res. Commun.*, 187 (1992) 480-487.

- 728 Chambers, S.J., Bacon, J.R. and Lambert, N.: The quantitative analysis of seed proteins from peas using high performance liquid chromatography. *Phytochem. Anal.*, 3 (1992) 49-54; C.A., 117 (1992) 43757d.
- 729 Coso, O.A., Diaz Anel, A., Martinetto, H., Muschietti, J.P., Kazanietz, M., Fraidenraich, D., Torres, H.N. and Flawia, M.M.: Characterization of a G_i-protein from *Trypanosoma cruzi* epimastigote membranes. *Biochem. J.*, 287 (1992) 443-446.
- 730 Cover, T.L. and Blaser, M.J.: Purification and characterization of the vacuolating toxin from *Helicobacter pylori*. *J. Biol. Chem.*, 267 (1992) 10570-10575.
- 731 Ellis, C.M., Ford, R.C. and Holzenburg, A.: Detergent sensitivity of the tonoplast H⁺-ATPase and its purification from *Beta vulgaris*. *Biochim. Biophys. Acta*, 1136 (1992) 319-326.
- 732 Fernandes, P.A., Keen, J.N., Findlay, J.B.C. and Moradas-Ferreira, P.: A protein homologous to glyceraldehyde-3-phosphate dehydrogenase is induced in the cell wall of a flocculent *Kluyveromyces marxianus*. *Biochim. Biophys. Acta*, 1159 (1992) 67-73.
- 733 Huguenev, P., Römer, S., Kuntz, M. and Camara, B.: Characterization and molecular cloning of a flavoprotein catalyzing the synthesis of phytofluene and ζ -carotene in *Capsicum* chromoplasts. *Eur. J. Biochem.*, 209 (1992) 399-407.
- 734 Komisarenko, S.V., Bobrovnyk, S.P., Lyashchenko, K.P., Nartykov, A.V., Petrova, Yu.L. and Gorbenko, O.V.: (Chromatographic analysis and immunobiologic properties of tuberculin). *Ukr. Biokhim. Zh.*, 63 (1991) 21-26; C.A., 117 (1992) 169113j.
- 735 Liu, H., Yan, Y.- and Tong, J.: (An improved gel chromatographic technique for the isolation and purification of cholera enterotoxin). *Sepu*, 10 (1992) 179-180, 166; C.A., 117 (1992) 64351g.
- 736 McFarlan, S.C., Terrell, C.A. and Hogenkamp, H.P.C.: The purification, characterization, and primary structure of a small redox protein from *Methanobacterium thermoautotrophicum*, an archaeobacterium. *J. Biol. Chem.*, 267 (1992) 10561-10569.
- 737 Mu, J. and Kao, T.-h.: Expression of two S-ribonucleases of *Petunia inflata* using baculovirus expression system. *Biochem. Biophys. Res. Commun.*, 187 (1992) 299-304.
- 738 Sjöström, H., Friis, S.U., Norén, O. and Anthonson, D.: Purification and characterisation of antigenic gliadins in coeliac disease. *Clin. Chim. Acta*, 207 (1992) 227-237.
- 739 Sutton, K.H., Hay, R.L. and Mout, C.H.: The effect of kernel weight on the assessment of baking performance of wheats by RP-HPLC of glutenin subunits from single grains. *J. Cereal Sci.*, 15 (1992) 253-265; C.A., 117 (1992) 130036m.
- 740 Terras, F.R.G., Schoofe, H.M.E., De Bolle, M.F.C., van Leuven, F., Rees, S.B., Vanderleyden, J., Cammue, B.P.A. and Broekaert, W.F.: Analysis of two novel classes of plant antifungal proteins from radish (*Raphanus sativus* L.) seeds. *J. Biol. Chem.*, 267 (1992) 15301-15309.
- 741 Ullah, A.H.J., Neucere, J.N. and Cleveland, T.E.: Surface proteins of two aflatoxin-producing isolates of *Aspergillus flavus* and *Aspergillus parasiticus* mycelia. 2. HPLC mapping by gel-permeation, ion-exchange and reverse-phase chromatography. *J. Agric. Food Chem.*, 40 (1992) 1613-1616.
- 742 Van de Meent, E.J., Kobayashi, M., Erkelens, C., van Veelen, P.A., Otte, S.C.M., Inoue, K., Watanabe, T. and Ames, J.: The nature of the primary electron acceptor in green sulfur bacteria. *Biochim. Biophys. Acta*, 1102 (1992) 371-378.
- 743 Waters, E.J., Wallace, W. and Williams, P.J.: Identification of heat-unstable wine proteins and their resistance to peptidases. *J. Agric. Food Chem.*, 40 (1992) 1514-1519.
- 744 Zarins, Z. and Chrastil, J.: Separation and purification of rice oryzenin subunits by anion-exchange and gel-permeation chromatography. *J. Agric. Food Chem.*, 40 (1992) 1599-1601.
- 745 Zawistowska, U., Zawistowski, J. and Friesen, A.D.: Application of immobilized metal affinity chromatography for large-scale purification of endogenous α -amylase inhibitor from barley kernels. *Biotechnol. Appl. Biochem.*, 15 (1992) 160-170; C.A., 117 (1992) 24941t.

See also 676, 712, 788, 1068.

19e. *Proteins of blood, serum and blood cells*

- 746 Biedermann, K., Sabater, M., Soerensen, J., Fiedler, H. and Emborg, C.: Quantitative binding studies of a monoclonal antibody to immobilized protein-A. *Bioseparation*, 2 (1991) 309-314; C.A., 117 (1992) 107521g.
- 747 Favarato, M., Mizzen, C.A., Sutherland, M.K., Krishnan, B., Kruck, T.P.A. and McLachlan, D.R.C.: Aluminium-binding serum proteins: desferrioxamine alters serum aluminium speciation. *Clin. Chim. Acta*, 207 (1992) 41-55.
- 748 Fried, R., Wiederkehr, F. and Vonderschmitt, D.J.: Immune complexes in blood: a new strategy for the analysis of their antigen portion. *Electrophoresis (Weinheim)*, 13 (1992) 718-719.
- 749 Gaffney, D.K. and Sieber, F.: The role of serum and serum components in the merocyanine 540-sensitized photoinactivation of K562 leukemia cells. *Biochim. Biophys. Acta*, 1117 (1992) 321-325.
- 750 Groopman, J.D., Skipper, P.L., Donahue, P.R., Trudel, L.J., Wildschutte, M., Kadlubar, F.F. and Tannenbaum, S.R.: Monoclonal antibodies and rabbit antisera recognizing 4-aminobiphenyl-DNA adducts and application to immunoaffinity chromatography. *Carcinogenesis (London)*, 13 (1992) 917-922; C.A., 117 (1992) 125975n.
- 751 Harada, K., Uda, S., Nakayama, M. and Sugii, A.: High performance liquid chromatography of ^{99m}Tc labeled human serum albumin using and N-methylpyridinium polymer column. *Chem. Pharm. Bull.*, 40 (1992) 697-700.
- 752 Kelleher, P.C., Toftager-Larsen, K. and Walters, C.J.P.: Comparison of concanavalin A-affinity electrophoresis and concanavalin A-affinity chromatography in the analysis of amniotic fluid alpha₁-fetoprotein microheterogeneity. In: Breborowicz, J. and Mackiewicz, A. (Editors), *Affinity Electrophor.*, CRC, Boca Raton, 1992, pp. 89-95; C.A., 117 (1992) 167107m.
- 753 Koskinen, L. and Sillanaukee, P.: Determination of fetal hemoglobin by cation exchange liquid chromatography. *Clin. Chim. Acta*, 206 (1992) 215-223.
- 754 Koyama, T., Koike, Y., Sugahara, Y. and Aoki, N.: Purification of human α_2 -plasmin inhibitor by a one-step immunoaffinity chromatography procedure. *Thromb. Res.*, 66 (1992) 451-454; C.A., 117 (1992) 85662f.
- 755 Lepesheva, G.I. and Martsev, S.P.: (Modification of the rabbit IgG hinge disulfide and study of its interaction with the polyvalent antigen ferritin, protein A and anti-IgG). *Biokhimiya (Moscow)*, 57 (1992) 1089-1099.

- 756 Nakatani, S., Kato, Y., Nasu, M., Fujita, S. and Katayama, Y.: (Determination of serum glycosylated albumin by high-performance affinity chromatography). *Bunseki Kagaku*, 41 (1992) 387-391; C.A., 117 (1992) 146457k.
- 757 Neuenschwander, P.F. and Jesty, J.: Thrombin-activated and factor Xa-activated human factor VIII: differences in cofactor activity and decay rate. *Arch. Biochem. Biophys.*, 296 (1992) 426-434.
- 758 Polten, B.M., Jimenez, C. and Schliesser, T.: (Comparative study of a chromatographic (protein A column) and a nonchromatographic procedure (octanoic acid-ammonium sulfate precipitation) for concentration of immunoglobulin in rabbit antiserum). *J. Vet. Med., Ser. B*, 38 (1991) 714-718; C.A., 117 (1992) 24359j.
- 759 Rutishauser, H. and Reichmuth, A.: The influence of gravity on the weighing result. *Int. Lab.*, 22, No. 9 (1992) P14-P18.
- 760 Sakata, S., Maekawa, H., Ogawa, T., Sarui, H., Matsuda, M., Endo, J., Wada, S. and Nagai, K.: A case of Hashimoto's thyroiditis with antibodies to thyroid hormone analogues. *Clin. Chim. Acta*, 207 (1992) 129-136.
- 761 Samaraweera, P., Porath, J. and Law, J.H.: Separation of insect hemolymph proteins by cascade-mode multi-affinity chromatography. *Arch. Insect Biochem. Physiol.*, 20 (1992) 243-251; C.A., 117 (1992) 166927s.
- 762 Takami, M., Kasuya, I. and Tsunoo, H.: Polymerized albumin receptor on rat liver cells. *J. Biochem. (Tokyo)*, 111 (1992) 714-721.
- 763 Yamada, S., Kamiya, Y., Eto, N., Yamada, K., Murakami, H. and Majima, T.: Effective purification of monoclonal antibodies by fast flow affinity chromatography. In: Sasaki, R. and Ikura, K. (Editors), *Anim. Cell Cult. Prod. Biol., Proc. Annu. Meet. Jpn. Assoc. Anim. Cell Technol.*, 3rd 1990, Kluwer, Dordrecht, 1991, pp. 353-357; C.A., 117 (1992) 129443k.
- For additional information see C.A.:
117 (1992) 24679p, 24697t, 46544t, 66050g, 86340c, 88628d, 126946r, 127533r, 129864y, 146442b.
- See also 652, 846, 1235, 1563.
- 19f. *Structural and muscle proteins*
- 764 Arai, M., Yada, T., Suzuki, S. and Kimata, K.: Isolation and characterization of type IX collagen-proteoglycan from the Swarm rat chondrosarcoma. *Biochim. Biophys. Acta*, 1117 (1992) 60-70.
- 765 Bouillon, R., Vanderschueren, D., van Herck, E., Nielsen, H.K., Bex, M., Heyns, W. and van Baelen, H.: Homologous radioimmunoassay of human osteocalcin. *Clin. Chem. (Winston-Salem)*, 38 (1992) 2055-2060.
- 766 Brook, J.L., Branford-White, C. and Whish, W.J.: Calmodulin from the *Eucestoda* and its use as a probe for calmodulin binding proteins. *Biochem. Soc. Trans.*, 20 (1992) 300S; C.A., 117 (1992) 22653b.
- 767 Katakami, Y., Katakami, N., Janmey, P.A., Hartwig, J.H. and Stossel, T.P.: Isolation of the phosphatidylinositol 4-monophosphate dissociable high-affinity profilin-actin complex. *Biochim. Biophys. Acta*, 1122 (1992) 123-135.
- 768 Mani, R.S. and Kay, C.M.: Purification and characterization of a novel 12,000-Da calcium-binding protein from smooth muscle. *Arch. Biochem. Biophys.*, 296 (1992) 442-449.
- 769 Ngai, S. and Hodges, R.S.: Biologically important interactions between synthetic peptides of the N-terminal region of troponin I and troponin C. *J. Biol. Chem.*, 267 (1992) 15715-15720.
- 770 Sobotka-Briner, C. and Chelsky, D.: COOH-Terminal methylation of lamin B and inhibition of methylation by farnesylated peptides corresponding to lamin B and other CAAX motif proteins. *J. Biol. Chem.*, 267 (1992) 12116-12122.
- 771 Zhou, H.-Y., Ohnuma, Y., Takita, H., Fujisawa, R., Mizuno, M. and Kuboki, Y.: Effects of a bone lysine-rich 18 kDa protein on osteoblast-like MC3T3-E1 cells. *Biochem. Biophys. Res. Commun.*, 186 (1992) 1288-1293.
- For additional information see C.A.:
117 (1992) 86112g.
- See also 668, 669, 679, 787, 834.
- 19g. *Protamines, histones and other nuclear proteins*
- 772 Balhorn, R., Corzett, M. and Mazrimas, J.A.: Formation of intra-protamine disulfides *in vitro*. *Arch. Biochem. Biophys.*, 296 (1992) 384-393.
- 773 Baubichon-Cortay, H., Mallet, L., Denoroy, L. and Roux, B.: Histone H1a subtype presents structural differences compared to other histone H1 subtypes. Evidence for a specific motif in the C-terminal domain. *Biochim. Biophys. Acta*, 1122 (1992) 167-177.
- 774 Lassle, M., Richter, A. and Knippers, R.: Comparison of replicative and non-replicative chromatin assembly pathways in HeLa cell extracts. *Biochim. Biophys. Acta*, 1132 (1992) 1-10.
- 775 Lindner, H., Helliger, W., Dirschlmaier, A., Talasz, H., Wurm, M., Sarg, B., Jaquemar, M. and Puschendorf, B.: Separation of phosphorylated histone H1 variants by high-performance capillary electrophoresis. *J. Chromatogr.*, 608 (1992) 211-216.
- 776 Waterborg, J.H.: Identification of five sites of acetylation in alfalfa histone H4. *Biochemistry*, 31 (1992) 6211-6219.
- See also 614.
- 19h. *Chromoproteins and metalloproteins*
- 777 Balvers, W.G., Boersma, M.G., Veeger, C. and Rietjens, I.M.C.M.: Differential cumene hydroperoxide sensitivity of cytochrome P-450 enzymes IA1 and IB1 determined by their way of membrane incorporation. *Biochim. Biophys. Acta*, 1117 (1992) 179-187.
- 778 Greaves, J., Rainer, J.S. and Mangum, C.P.: Size-exclusion high performance liquid chromatography of the dodecameric and hexameric forms of hemocyanin from *Callinectes sapidus*. *Mar. Biol. (Berlin)*, 113 (1992) 33-36; C.A., 117 (1992) 65794r.
- 779 Isa, M., Cumps, J., Fossoul, C. and Atassi, G.: A new method for the purification of cytochrome P-450 from human liver microsomes. *Biomed. Chromatogr.*, 6 (1992) 248-250.
- 780 Ishioka, N., Iyori, N., Noji, J. and Kurioka, S.: Detection of abnormal haemoglobin by capillary electrophoresis and structural identification. *Biomed. Chromatogr.*, 6 (1992) 224-226.
- 781 Jouanneau, Y., Dupont, C., Meyer, C. and Gaillard, J.: Expression in *Escherichia coli* and characterization of a recombinant 7Fe ferredoxin of *Rhodobacter capsulatus*. *Biochem. J.*, 286 (1992) 269-273.

- 782 Kabzinski, A.K.M. and Paryjczak, T.: (Covalent chromatographic separation of apothionein as example of separation of polythiolate polypeptides). *Chem. Anal. (Warsaw)*, 36 (1991) 699-708; C.A., 117 (1992) 127547y.
- 783 Kovaleva, G.G. and Gorodetskii, V.K.: (Production and use of affinity sorbent containing phenylboric acid for separation and quantitative estimation of glycosylated hemoglobin). *Vopr. Med. Khim.*, 38 (1992) 53-54; C.A., 117 (1992) 127765t.
- 784 Lee, P.D.K., Sherman, L.D., O'Day, M.R., Rognerud, C.L. and Ou, C.N.: Comparisons of home blood glucose testing and glycosylated protein measurements. *Diabetes Res. Clin. Pract.*, 16 (1992) 53-62; C.A., 117 (1992) 65973y.
- 785 Maiavacca, R., Tedeschi, S., Mosca, A., Calmi, S., de Leonardi, P., Todaro, C., Zuliani, G., Buscaglia, M. and Cantu-Rajoldi, A.: Nonradioactive quantification of low concentrations of hemoglobin A by HPLC for midtrimester prenatal diagnosis of β -thalassemia. *Clin. Chem. (Winston-Salem)*, 38 (1992) 1906-1908.
- 786 Micallef, S., Couillard, Y., Campbell, P.G.C. and Tessier, A.: An evaluation of the HPLC-gel chromatographic method for analyzing metallothioneins in aquatic organisms. *Talanta*, 39 (1992) 1073-1079; C.A., 117 (1992) 165377a.
- 787 Picello, E., Damiani, E. and Margreth, A.: Low-affinity Ca^{2+} -binding sites versus Zn^{2+} -binding sites in histidine-rich Ca^{2+} -binding protein of skeletal muscle sarcoplasmic reticulum. *Biochem. Biophys. Res. Commun.*, 186 (1992) 659-667.
- 788 Swanson, R.V., Zhou, J., Leary, J.A., Williams, T., de Lorimier, R., Bryant, D.A. and Glazer, A.N.: Characterization of phycocyanin produced by *cpcE* and *cpcF* mutants and identification of an intergenic suppressor of the defect in bilin attachment. *J. Biol. Chem.*, 267 (1992) 16146-16154.
- 789 Takatera, K. and Watanabe, T.: Application of high-performance liquid chromatography/inductively coupled plasma mass spectrometry to the speciation of cadmium-binding metallothionein-like protein in a cyanobacterium. *Anal. Sci.*, 8 (1992) 469-474; C.A., 117 (1992) 146635s.
- 790 Van der Dijks, F.P.L., van den Berg, G.A., Schermer, J.G., Muskiet, F.D., Landman, H. and Muskiet, F.A.J.: Screening cord blood for hemoglobinopathies and thalassemia by HPLC. *Clin. Chem. (Winston-Salem)*, 38 (1992) 1864-1869.
- 791 Vasseur, C., Blouquit, Y., Kister, J., Prome, D., Kavanaugh, J.S., Rogers, P.H., Guillemin, C., Arnone, A., Galacteros, F., Poyart, C. et al.: Hemoglobin Thionville. An α -chain variant with a substitution of a glutamate for valine at NA-1 and having an acetylated methionine NH_2 terminus. *J. Biol. Chem.*, 267 (1992) 12682-12691.
- For additional information see C.A.:
117 (1992) 43750w, 127803d, 146714s.
- See also 377, 671, 678, 711, 747, 753, 768, 868.
- 19i. *Proteins of glands, gland products, various zymogens (incl. milk proteins)*
- 792 Huang, T.-F., Peng, H.-C., Peng, I.-S., Teng, C.-M. and Ouyang, C.: An antiplatelet peptide, gabonin, from *Bitis gabonica* snake venom. *Arch. Biochem. Biophys.*, 298 (1992) 13-20.
- 793 Kawaminami, M., Tanaka, K., Asawa, T., Osugi, S., Kawachi, H., Kurusu, S. and Hashimoto, I.: Association of annexin V with prolactin in the rat anterior pituitary gland. *Biochem. Biophys. Res. Commun.*, 186 (1992) 894-899.
- 794 Law, A.J.R., Papoff, C.M., Dalgleish, D.G. and Campus, R.L.: Quantitative fractionation of ovine casein by cation-exchange FPLC. *Milchwissenschaft*, 47 (1992) 279-282; C.A., 117 (1992) 149625z.
- 795 Nishi, N., Kagawa, Y., Miyataka, H., Oya, H. and Wada, F.: An anti-probasin monoclonal antibody recognizes and novel 40-kDa protein localized in rat liver and a specific region of kidney urinary tubule. *Biochim. Biophys. Acta*, 1117 (1992) 47-54.
- 796 Shetty, S., Hegde, S.N. and Bharathi, L.: Purification of a growth factor from pigeon milk. *Biochim. Biophys. Acta*, 1117 (1992) 193-198.
- 797 Strange, E.D., Malin, E.L., van Hekken, D.L. and Basch, J.J.: Chromatographic and electrophoretic methods used for analysis of milk proteins. *J. Chromatogr.*, 624 (1992) 81-102 - a review with 204 refs.
- 798 Waita, J.H., Jensen, R.A. and Morse, D.E.: Cement precursor proteins of the reef-building polychaete *Phragmatopoma californica* (fewkes). *Biochemistry*, 31 (1992) 5733-5738.
- 19j. *Proteins of brain, cerebrospinal fluid and eye*
- 799 Hasegawa, M., Morishima-Kawashima, M., Takio, K., Suzuki, M., Titani, K. and Ihara, Y.: Protein sequence and mass spectrometric analyses of tau in the Alzheimer's disease brain. *J. Biol. Chem.*, 267 (1992) 17047-17054.
- 800 Izykenova, G.A. and Taranova, N.P.: (Dalargin-binding proteins of brain synaptic membranes: isolation and properties in comparison with opiate receptors). *Biokhimiya (Moscow)*, 57 (1992) 663-670.
- 801 Neubert, T.A., Johnson, R.S., Hurley, J.B. and Walsh, K.A.: The rod transducin α subunit amino terminus is heterogeneously fatty acylated. *J. Biol. Chem.*, 267 (1992) 18274-18277.
- 802 Rao, V.P. and Zigler, J.S., Jr.: Purification and characterization of ζ -crystallin/quinone reductase from guinea pig liver. *Biochim. Biophys. Acta*, 1117 (1992) 315-320.
- 803 Roquemore, E.P., Dell, A., Morris, H.R., Panico, M., Reason, A.J., Savoy, L.-A., Wistow, G.J., Zigler, J.S., Jr., Earles, B.J. and Hart, G.W.: Vertebrate lens α -crystallins are modified by O-linked N-acetylglucosamine. *J. Biol. Chem.*, 267 (1992) 555-563.
- 804 Saito, Y., Tsubuki, S., Ito, H., Ohmi-Imajo, S. and Kawashima, S.: Possible involvement of clathrin in neurogenesis induced by a protease inhibitor (benzyloxycarbonyl-Leu-Leu-aldehyde) in PC12 cells. *J. Biochem. (Tokyo)*, 112 (1992) 448-455.
- 805 Santini, S.A., Mordente, A., Meucci, E., Miggiano, G.A.D. and Martorana, G.E.: Conformational stability of bovine α -crystallin. Evidence for a destabilizing effect of ascorbate. *Biochem. J.*, 287 (1992) 107-112.
- 806 Togashi, K., Kameya, T., Kurosawa, T., Hasegawa, N. and Kawakami, M.: Concentrations and molecular forms of C-type natriuretic peptide in brain and cerebrospinal fluid. *Clin. Chem. (Winston-Salem)*, 38 (1992) 2136-2139.
- 807 Torriglia, A. and Blanquet, P.R.: Purification of an active receptor for acidic and basic fibroblast growth factor from bovine retina. *Biochim. Biophys. Acta*, 1137 (1992) 215-224.

- 808 Yoneda, Y., and Ogita, K.: Attempts to isolate the NMDA receptor complex from rat brain. *Mol. Neuropharmacol.*, 2 (1992) 65-70; C.A., 117 (1992) 124695j.
- 809 Zarina, S., Abbasi, A. and Zaidi, Z.H.: Primary structure of β -crystallin from human lens. *Biochem. J.*, 287 (1992) 375-381.
- See also 674, 715, 835.
- 19k. *Proteins of neoplastic tissue and transformed cells*
- 810 Castronovo, V., Luyten, F., van den Brule, F. and Sobel, M.E.: Identification of a 14-kDa laminin binding protein (HLBP14) in human melanoma cells that is identical to the 14-kDa galactoside binding lectin. *Arch. Biochem. Biophys.*, 297 (1992) 132-138.
- 811 Sage, E.H.: Secretion of SPARC by endothelial cells transformed by polyoma middle T oncogene inhibits the growth of normal endothelial cells *in vitro*. *Biochem. Cell Biol.*, 70 (1992) 579-592.
- 19l. *Specific binding and receptor proteins*
- 812 Alexander, D.R., Brown, M.H., Tutt, A.L., Crumpton, M.J. and Shivnan, E.: CD3 and CD2 antigen-mediated CD3 γ -chain phosphorylation in permeabilized human T cells. Regulation by cytosolic phosphatases. *Biochem. J.*, 288 (1992) 69-77.
- 813 Bahr, B.A. and Parsons, S.M.: Purification of the vesamicol receptor. *Biochemistry*, 31 (1992) 5763-5769.
- 814 Bautsch, W., Kretschmar, T., Stühmer, T., Kola, A., Emde, M., Köhl, J., Klos, A. and Bitter-Suermann, D.: A recombinant hybrid anaphylatoxin with dual C3a/C5a activity. *Biochem. J.*, 288 (1992) 261-266.
- 815 Blanks, R. and McLaughlin, L.W.: Oligodeoxynucleotides for affinity chromatography. In: Eckstein, F. (Editor), *Oligonucleotides Analogues*, IRL, Oxford, 1991, pp. 2451-254; C.A., 117 (1992) 43761a.
- 816 Dalton, A., Hornby, D.P., Langston, S.P. and Blackburn, G.M.: Characterization and purification of a novel dATP-binding protein in eukaryotes. *Biochem. J.*, 287 (1992) 871-879.
- 817 Dean, B., McAdam, A.J., Sundram, S., Pavay, G., Harrison, L.C. and Copolov, D.L.: Identification of a dopamine-binding protein on the membrane of the human platelet. *Biochem. J.*, 287 (1992) 45-50.
- 818 Ferreri, K. and Menan, K.M.: Characterization and isolation of a high-density-lipoprotein-binding protein from bovine corpus luteum plasma membrane. *Biochem. J.*, 287 (1992) 841-848.
- 819 Goulet, F., Moore, K.G. and Sartorelli, A.C.: Glycosylation of annexin I and annexin II. *Biochem. Biophys. Res. Commun.*, 188 (1992) 554-558.
- 820 Graham, D., Esnaud, H. and Langer, S.Z.: Partial purification and characterization of the sodium-ion-coupled 5-hydroxytryptamine transporter of rat cerebral cortex. *Biochem. J.*, 286 (1992) 801-805.
- 821 Hagenbuechle, O. and Wellauer, P.K.: A rapid method for the isolation of DNA-binding proteins from purified nuclei of tissues and cells in culture. *Nucleic Acids Res.*, 20 (1992) 3555-3559; C.A., 117 (1992) 166919r.
- 822 Hauguel-De Mouzon, S., Louizeau, M. and Girard, J.: Functional alterations of type I insulin-like growth factor receptor in placenta of diabetic rats. *Biochem. J.*, 288 (1992) 273-279.
- 823 Higashi, H., Omori, A. and Yamagata, T.: Calmodulin, a ganglioside-binding protein. Binding of gangliosides to calmodulin in the presence of calcium. *J. Biol. Chem.*, 267 (1992) 9831-9838.
- 824 Hokland, B., Mendez, A.J. and Oram, J.F.: Cellular localization and characterization of proteins that bind high density lipoprotein. *J. Lipid Res.*, 33 (1992) 1335-1342.
- 825 Inano, K., Ishida, T., Omata, S. and Horigome, T.: *In vitro* formation of estrogen receptor-heat shock protein 90 complexes. *J. Biochem. (Tokyo)*, 112 (1992) 535-540.
- 826 Inomata, Y., Kawaguchi, H., Hiramoto, M., Wada, T. and Handa, H.: Direct purification of multiple ATF/E4TF3 polypeptides from HeLa cell crude nuclear extracts using DNA affinity latex particles. *Anal. Biochem.*, 206 (1992) 109-114.
- 827 Kalsi, G., Das, H.R., Babu, C.R. and Das, R.H.: Isolation and characterization of a lectin from peanut roots. *Biochim. Biophys. Acta*, 1117 (1992) 114-119.
- 828 Kayyem, J.F., Roman, J.M., von Boxberg, Y., Schwarz, U. and Dreyer, W.J.: A method for the generation of monoclonal antibodies against rare cell-surface molecules. *Eur. J. Biochem.*, 208 (1992) 1-8.
- 829 Koyama, H., Prael, J.M., Uhland, A., Nanjo, M., Inaba, M., Nishizawa, Y., Morii, H., Nishii, Y. and DeLuca, H.F.: A new, highly sensitive assay for 1,25-dihydroxyvitamin D not requiring high-performance liquid chromatography: application of monoclonal antibody against vitamin D receptor to radioreceptor assay. *Anal. Biochem.*, 205 (1992) 213-219.
- 830 Larson, C.J. and Verdine, G.L.: A high-capacity column for affinity purification of sequence-specific DNA-binding proteins. *Nucleic Acids Res.*, 20 (1992) 3525; C.A., 117 (1992) 146450c.
- 831 Lemansky, P. and Herzog, V.: Endocytosis of thyroglobulin is not mediated by mannose-6-phosphate receptors in thyrocytes. Evidence for low-affinity-binding sites operating in the uptake of thyroglobulin. *Eur. J. Biochem.*, 209 (1992) 111-119.
- 832 Li, M. and Ishibashi, T.: *An in vitro* reversible interconversion of rat liver fatty acid binding protein having different isoelectric points by virtue of the fatty acid content. *Arch. Biochem. Biophys.*, 298 (1992) 254-258.
- 833 Mahon, M.J. and Gasiewicz, T.A.: Chelatable metal ions are not required for aryl hydrocarbon receptor transformation to a DNA binding form: phenanthrolines are possible competitive antagonists of 2,3,7,8-tetrachlorodibenzo-p-dioxin. *Arch. Biochem. Biophys.*, 297 (1992) 1-8.
- 834 Murayama, T. and Ogawa, Y.: Purification and characterization of two ryanodine-binding protein isoforms from sarcoplasmic reticulum of bullfrog skeletal muscle. *J. Biochem. (Tokyo)*, 112 (1992) 514-522.
- 835 Nakata, Y., Takamatsu, H., Kuroyanagi, N., Nishio, H., Segawa, T., Akizawa, T., Hirai, Y. and Akiyama, M.: Isolation of substance P binding protein from rat brain. *Jpn. J. Pharmacol.*, 59 (1992) 313-319; C.A., 117 (1992) 143632c.
- 836 Nalecz, K.A., Nalecz, M.J. and Azzi, A.: Isolation of tocopherol-binding proteins from the cytosol of smooth muscle A7r5 cells. *Eur. J. Biochem.*, 209 (1992) 37-42.
- 837 Ogert, R.A., Brown, J.E., Singh, B.R., Shriver-Lake, L.C. and Liggler, F.S.: Detection of *Clostridium botulinum* toxin A using a fiber optic-based biosensor. *Anal. Biochem.*, 205 (1992) 306-312.

- 838 Peryt, B., Maurel, P. and Lesca, P.: Characterization of the 4 S polycyclic aromatic hydrocarbon-binding protein in human liver and cells. *Arch. Biochem. Biophys.*, 298 (1992) 420-430.
- 839 Pilloud-Dagher, M.-C., Jouan, A. and Vignais, P.V.: Purification and properties of a functional 47-kilodalton cytosolic factor required for NADPH-oxidase activation in bovine neutrophils. *Biochem. Biophys. Res. Commun.*, 186 (1992) 731-738.
- 840 Platt, F.M., Karlsson, G.B. and Jacob, G.S.: Modulation of cell-surface transferrin receptor by the imino sugar N-butyldeoxynojirimycin. *Eur. J. Biochem.*, 208 (1992) 187-193.
- 841 Ploug, M., Eriksen, J., Plesner, T., Hansen, N.E. and Dano, K.: A soluble form of the glycolipid-anchored receptor for urokinase-type plasminogen activator is secreted from peripheral blood leukocytes from patients with paroxysmal nocturnal hemoglobinuria. *Eur. J. Biochem.*, 208 (1992) 397-404.
- 842 Richardson, A. and Taylor, C.: Purification of the inositol 1,4,5-trisphosphate receptor by heparin affinity chromatography using decavandate. *Biochem. Soc. Trans.*, 20 (1992) 135S; C.A., 117 (1992) 22650y.
- 843 Rutberg, S.E. and Ronai, Z.: A simple and efficient method for the purification of specific DNA binding proteins. *Nucleic Acids Res.*, 20 (1992) 1815; C.A., 117 (1992) 65880r.
- 844 Santostefano, M., Piskorska-Pliszczynska, J., Morrison, V. and Safe, S.: Effects of ligand structure on the *in vitro* transformation of the rat cytosolic aryl hydrocarbon receptor. *Arch. Biochem. Biophys.*, 297 (1992) 73-79.
- 845 Stambuk, R.A. and Moon, R.T.: Purification and characterization of recombinant *Xenopus* poly(A)-binding proteins expressed in a baculovirus system. *Biochem. J.*, 287 (1992) 761-766.
- 846 Stojkovski, S., Goumakos, W. and Sarkar, B.: Iron(III)-binding polypeptide in human cord and adult serum: isolation, purification and partial characterization. *Biochim. Biophys. Acta*, 1137 (1992) 155-161.
- 847 Suzuki, T., Nagata, K.-i., Matsuura, Y., Okano, Y. and Nozawa, Y.: Purification and characterization of a low M_r GTP-binding protein, ram p25, expressed by baculovirus expression system. *Biochim. Biophys. Acta*, 1159 (1992) 162-168.
- 848 Takikawa, H., Arai, S. and Yamanaka, M.: Lithocholate binding by 4 and 4' proteins in bovine small intestine. *Arch. Biochem. Biophys.*, 298 (1992) 486-491.
- 849 Verma, R.S. and Antony, A.C.: Immunoreactive folate-binding proteins from human saliva. Isolation and comparison of two distinct species. *Biochem. J.*, 286 (1992) 707-715.
- 850 Wakamiya, N., Okuno, Y., Sasao, F., Ueda, S., Yoshimatsu, K., Naiki, M. and Kurimura, T.: Isolation and characterization of conglutinin as an influenza A virus inhibitor. *Biochem. Biophys. Res. Commun.*, 187 (1992) 1270-1278.
- 851 Wille, M., Just, I., Wegner, A. and Aktories, K.: ADP-Ribosylation of gelsolin-actin complexes by clostridial toxins. *J. Biol. Chem.*, 267 (1992) 50-55.
- For additional information see C.A.:
117 (1992) 127529u, 146585a.
- See also 643, 660, 708, 711, 765, 768, 787, 807, 808, 810, 976.
- 19m. *Urinary proteins*
- See 993.
- 19n. *Other proteins (incl. proteinous inhibitors of enzymic activity)*
- 852 Baert, J.-L., Britel, M., Sautiere, P. and Malecha, J.: Ovohemerythrin, a major 14-kDa yolk protein distinct from vitellogenin in leech. *Eur. J. Biochem.*, 209 (1992) 563-569.
- 853 Hoffmann, R., Jung, S. and Hofer, H.W.: Association of a heat-stable inhibitor protein with cyclic-3',5'-AMP-dependent protein kinase from the nematode *Ascaris suum*: purification and characterization of the inhibitor. *Arch. Biochem. Biophys.*, 297 (1992) 296-303.
- 854 Ishikawa, M., Kubo, T. and Natori, S.: Purification and characterization of a dipterin homologue from *Sarcophaga peregrina* (flesh fly). *Biochem. J.*, 287 (1992) 573-578.
- 855 Kaplan, B. and Pras, M.: Separation of amyloid proteins by size-exclusion chromatography in aqueous organic mobile phase. *J. Liq. Chromatogr.*, 15 (1992) 2467-2486.
- 856 Moore, K.G., Goulet, F. and Sartorelli, A.C.: Purification of annexin I and annexin II from human placental membranes by high-performance liquid chromatography. *Protein Expression Purif.*, 3 (1992) 1-7; C.A., 117 (1992) 22614q.
- 857 Tuong, A., Maftouh, M., Ponthus, C., Whitechurch, O., Roitsch, C. and Picard, C.: Characterization of the deamidated forms of recombinant hirudin. *Biochemistry*, 31 (1992) 8201-8299.
- 858 Van Iwaarden, J.F., Shimizu, H., van Golde, P.H.M., Voelker, D.R. and van Golde, L.M.G.: Rat surfactant protein D enhances the production of oxygen radicals by rat alveolar macrophages. *Biochem. J.*, 286 (1992) 5-8.
- 859 Wagner, S.L., Siegel, R.S., Vedvick, T.S., Raschke, W.C. and van Nostrand, W.E.: High level expression, purification, and characterization of the Kunitz-type protease inhibitor domain of pro-tease nexin-2/amyloid β -protein precursor. *Biochem. Biophys. Res. Commun.*, 186 (1992) 1138-1145.
- 860 Wang, C., Lin, J.-M. and Lazarides, E.: Methylations of 70000-Da heat shock proteins in 3T3 cells: alterations by arsenite treatment, by different stages of growth and by virus transformation. *Arch. Biochem. Biophys.*, 297 (1992) 169-175.
- See also 745, 767.
20. ENZYMES AND ENZYME ACTIVITY ESTIMATION
- See 379, 701.
- 20a. *Oxidoreductases*
- 861 Adachi, H., Konishi, K., Adachi, I. and Horikoshi, I.: Purification and properties of reduced nicotinamide adenine dinucleotide dehydrogenase from *Photobacterium phosphoreum*. *Chem. Pharm. Bull.*, 40 (1992) 427-431.
- 862 Adachi, T., Kodera, T., Ohta, H., Hayashi, K. and Hirano, K.: The heparin binding site of human extracellular-superoxide dismutase. *Arch. Biochem. Biophys.*, 297 (1992) 155-161.
- 863 Axén, E., Bergman, T. and Wikvall, K.: Purification and characterization of a vitamin D₃ 25-hydroxylase from pig liver microsomes. *Biochem. J.*, 287 (1992) 725-731.
- 864 Braun, H.-P. and Schmitz, U.K.: Affinity purification of cytochrome c reductase from potato mitochondria. *Eur. J. Biochem.*, 208 (1992) 761-767.

- 865 Champluvier, B. and Kula, M.R.: Dye-ligand membranes as selective adsorbents for rapid purification of enzymes: a case study. *Biotechnol. Bioeng.*, 40 (1992) 33-40; C.A., 117 (1992) 126998j.
- 866 Costa, H.S., Santos, H., Turner, D.L. and Xavier, A.V.: Involvement of a labile axial histidine in coupling electron and proton transfer in *Methylophilus metholotropus* cytochrome c". *Eur. J. Biochem.*, 208 (1992) 427-433.
- 867 De Jong, E., van Berkel, W.J.H., van der Zwan, R.P. and de Bont, J.A.M.: Purification and characterization of vanillylalcohol oxidase from *Penicillium simplicissimum*. A novel aromatic alcohol oxidase containing covalently bound FAD. *Eur. J. Biochem.*, 208 (1992) 651-657.
- 868 Dionysius, D.A., Herse, J.B. and Grieve, P.A.: Extraction of lactoperoxidase and lactoferrin from whey using batch ion exchange techniques. *Aust. J. Dairy Technol.*, 46 (1991) 72-76; C.A., 117 (1992) 24969h.
- 869 Felsch, J., Kotake, S. and Copeland, R.A.: Purification of cytochrome c oxidase by lysine-affinity chromatography. *Protein Expression Purif.*, 3 (1992) 36-40; C.A., 117 (1992) 22234r.
- 870 Halle, F. and Meyer, J.-M.: Ferrisiderophore reductases of *Pseudomonas*. Purification, properties and cellular location of the *Pseudomonas aeruginosa* ferripyoverdine reductase. *Eur. J. Biochem.*, 209 (1992) 613-620.
- 871 Hartmann, C. and de Montellano, P.R.O.: Baculovirus expression and characterization of catalytically active horseradish peroxidase. *Arch. Biochem. Biophys.*, 297 (1992) 61-72.
- 872 Hatchikian, E.C., Forget, N., Fernandez, V.M., Williams, R. and Cammack, R.: Further characterization of the [Fe]-hydrogenase from *Desulfovibrio desulfuricans* ATCC 7757. *Eur. J. Biochem.*, 209 (1992) 357-365.
- 873 Helander, A. and Gogoll, A.: Mechanisms for plasma-mediated activation of human blood cell aldehyde dehydrogenase. *Biochim. Biophys. Acta*, 1136 (1992) 259-264.
- 874 Hirôse, J., Kano, H., Kidani, Y., Iwamoto, H. and Hiromi, K.: Zinc deficient bovine erythrocyte superoxide dismutase has low specific activity. *Chem. Pharm. Bull.*, 40 (1992) 506-508.
- 875 Itoh, N., Morinaga, N. and Nomura, A.: A variety of catalases and bromoperoxidases in genus *Pseudomonas* and their characterization. *Biochim. Biophys. Acta*, 1122 (1992) 189-195.
- 876 Jansen, E.H.J.M. and de Fluiter, P.: Detection of the enzymatic activity of cytochrome P-450 enzymes by high-performance liquid chromatography. *J. Chromatogr.*, 580 (1992) 325-346 - a review with 97 refs.
- 877 Johnson, T.M., Pease, E.A., Li, J.K.-K. and Tien, M.: Production and characterization of recombinant lignin peroxidase isozyme H2 from *Phanerochaete chrysosporium* using recombinant baculovirus. *Arch. Biochem. Biophys.*, 296 (1992) 660-666.
- 878 Kunapuli, S.P. and Vaidyanathan, C.S.: New indole oxygenase from the leaves of *Tecoma stans* L. Part I: Affinity purification and properties. *J. Indian Inst. Sci.*, 71 (1991) 503-513; C.A., 117 (1992) 166303k.
- 879 Lamikanra, O., Kirby, S.D. and Musingo, M.N.: Muscadine grape polyphenoloxidase: partial purification by high pressure liquid chromatography and some properties. *J. Food Sci.*, 57 (1992) 686-689; C.A., 117 (1992) 43263q.
- 880 Lu, Z., Zhang, R. and Diasio, R.B.: Purification and characterization of dihydropyrimidine dehydrogenase from human liver. *J. Biol. Chem.*, 267 (1992) 17102-17109.
- 881 Matsushita, K., Takahashi, K., Takahashi, M., Ameyama, M. and Adachi, O.: Methanol and ethanol oxidase respiratory chains of the methylotrophic acetic acid bacterium, *Acetobacter methanolicus*. *J. Biochem. (Tokyo)*, 111 (1992) 739-747.
- 882 Mizoguchi, T., Nanjo, H., Umemura, T., Nishinaka, T., Iwata, C., Imanishi, T., Tanaka, T., Terada, T. and Nishihara, T.: A novel dihydrodiol dehydrogenase in bovine liver cytosol: purification and characterization of multiple forms of dihydrodiol dehydrogenase. *J. Biochem. (Tokyo)*, 112 (1992) 523-529.
- 883 Morel, F., Surla, A. and Vignais, P.V.: Purification of human placenta diamine oxidase. *Biochem. Biophys. Res. Commun.*, 187 (1992) 178-186.
- 884 Negri, A., Cecilian, F., Tadeschi, G., Simonic, T. and Ronchi, S.: The primary structure of the flavoprotein D-aspartate oxidase from beef kidney. *J. Biol. Chem.*, 267 (1992) 11865-11871.
- 885 Pascual, P., Martinez-Lara, E., Bárcena, J.A., López-Barea, J. and Toribio, F.: Direct assay of glutathione peroxidase activity using high-performance capillary electrophoresis. *J. Chromatogr.*, 581 (1992) 49-56.
- 886 Perez-Vicente, R., Alamillo, J.M., Cárdenas, J. and Pineda, M.: Purification and substrate inactivation of xanthine dehydrogenase from *Chlamydomonas reinhardtii*. *Biochim. Biophys. Acta*, 1117 (1992) 159-166.
- 887 Pierik, A.J., Hagen, W.R., Redeker, J.S., Wolbert, R.B.G., Boersma, M., Verhagen, M.F.J.M., Grande, H.J., Veeger, C., Mutsaers, P.H.A. et al.: Redox properties of the iron-sulfur clusters in activated Fe-hydrogenase from *Desulfovibrio vulgaris* (Hildenborough). *Eur. J. Biochem.*, 209 (1992) 63-72.
- 888 Rabinovitch-Chable, H., Cook-Moreau, J., Breton, J.-C. and Rigaud, M.: Purification of lipoygenase and hydroperoxide dehydrase in flaxseeds: interaction between these enzymatic activities. *Biochem. Biophys. Res. Commun.*, 188 (1992) 858-864.
- 889 Sandström, J., Carlsson, L., Marklund, S.L. and Edlund, T.: The heparin-binding domain of extracellular superoxide dismutase C and formation of variants with reduced heparin affinity. *J. Biol. Chem.*, 267 (1992) 18205-18209.
- 890 Shaw, J.P. and Harayama, S.: Purification and characterization of the NADH:acceptor reductase component of xylene monooxygenase encoded by the TOL plasmid pWWO of *Pseudomonas putida* mt-2. *Eur. J. Biochem.*, 209 (1992) 51-61.
- 891 Suzuki, Y., Higo, K., Hagiwara, K., Ohtsubo, K., Kobayashi, A. and Nozu, Y.: Production and use of monoclonal antibodies against rice embryo lipoygenase-3. *Biosci. Biotechnol., Biochem.*, 56 (1992) 678-679; C.A., 117 (1992) 106901u.
- 892 Taylor, K.L., Uhlinger, D.J. and Kinkade, J.M., Jr.: Expression of recombinant myeloperoxidase using a baculovirus expression system. *Biochem. Biophys. Res. Commun.*, 187 (1992) 1572-1578.
- 893 Zhang, J., Blusztajn, J.K. and Zeisel, S.H.: Measurement of the formation of betaine aldehyde and betaine in rat liver mitochondria by a high pressure liquid chromatography-radioenzymatic assay. *Biochim. Biophys. Acta*, 1117 (1992) 333-339.
- 894 Zirngibl, C., van Dongen, W., Schwörer, B., von Büнау, R., Richter, M., Klein, A. and Thauer, R.K.: H₂-Forming methylenetetrahydromethanopterin dehydrogenase, a novel type of hydrogenase without iron-sulfur clusters in methanogenic archaea. *Eur. J. Biochem.*, 208 (1992) 511-520.

For additional information see C.A.:
117 (1992) 43173k, 106911x.

See also 732.

20b. *Transferases (excl. E.C. 2.7.-.-)*

- 895 Andriampandry, C., Massarelli, R. and Kanfer, J.N.: Properties of a partially purified phosphodimethylethanolamine methyltransferase from rat brain cytosol. *Biochem. J.*, 288 (1992) 267-272.
- 896 Arai, K., Yoshida, K.-T., Komoda, T., Kobayashi, N. and Sakagishi, Y.: Differences in the enzymatic nature and the sugar-chain structure of γ -glutamyl transferase between normal and carcinomatous human kidney and prostate. *Clin. Chim. Acta*, 210 (1992) 35-46.
- 897 Bell, W., Klaassen, P., Ohnacker, M., Boller, T., Herweijer, M., Schoppink, P., van der Zee, P. and Wiemken, A.: Characterization of the 56-kDa subunit of yeast trehalose-6-phosphate synthase and cloning of its gene reveal its identity with the product of ClF1, a regulator of carbon catabolite inactivation. *Eur. J. Biochem.*, 209 (1992) 951-959.
- 898 Bergamini, C.M. and Signorini, M.: Purification of testicular transglutaminase by hydrophobic chromatography on phenyl-Sepharose. *Biochem. Int.*, 27 (1992) 557-565; *C.A.*, 117 (1992) 166403t.
- 899 Bessoule, J.-J., Creach, A., Lessire, R. and Cassagne, C.: Evaluation of the amount of acyl-CoA elongases in leek (*Allium porrum* L) leaves. *Biochim. Biophys. Acta*, 1117 (1992) 78-82.
- 900 Bosshart, H. and Berger, E.G.: Biosynthesis and intracellular transport of α -2,6-sialyltransferase in rat hepatoma cells. *Eur. J. Biochem.*, 208 (1992) 341-349.
- 901 Dakour, J., Zopf, D. and Lundblad, A.: Assay of α 1,3N-acetyl-D-galactosaminyl transferase by affinity chromatography. *Anal. Biochem.*, 204 (1992) 210-214.
- 902 De Biase, D., Maras, B., Bossa, F., Barra, D. and John, R.A.: Protein structure of pig liver 4-aminobutyrate aminotransferase and comparison with a cDNA-deduced sequence. *Eur. J. Biochem.*, 208 (1992) 351-357.
- 903 Gonzalez, J.C., Banerjee, R.V., Huang, S., Sumner, J.S. and Matthews, R.G.: Comparison of cobalamin-independent and cobalamin-dependent methionine synthases from *Escherichia coli*: two solutions to the same chemical problem. *Biochemistry*, 31 (1992) 6045-6056.
- 904 Holmes, E.H.: Presence of an essential lysine residue in a GDP-fucose protected site of the α 1 \rightarrow 3fucosyltransferase from human small cell lung carcinoma NCI-H69 cells. *Arch. Biochem. Biophys.*, 296 (1992) 562-568.
- 905 Hussey, A.J. and Hayes, J.D.: Characterization of a human class-Theta glutathione S-transferase with activity towards 1-methynaphthyl sulphate. *Biochem. J.*, 286 (1992) 929-935.
- 906 Ide, H., Hamaguchi, K., Kobata, S., Murakami, A., Makino, K. and Kamada, M.: Purification of serine hydroxymethyltransferase with a new ion exchange column. *Kontakte (Darmstadt)*, No. 2 (1992) 61-64.
- 907 Kamat, U., Garg, R. and Sharma, C.B.: Purification to homogeneity and characterization of a 1,3- β -glucan (callose) synthase) from germinating *Arachis hypogaea* cotyledons. *Arch. Biochem. Biophys.*, 298 (1992) 731-739.
- 908 Kamio, K., Gasa, S. and Makita, A.: Galactosylceramide containing ω -aminofatty acids: preparation, characterization, and sulfo-transferase acceptor. *J. Lipid Res.*, 33 (1992) 1227-1232.
- 909 Katusz, R.M., Bono, B. and Colman, R.F.: Identification of Tyr¹¹⁵ labeled by S-(4-bromo-2,3-dioxobutyl)glutathione in the hydrophobic substrate binding site of glutathione S-transferase, isoenzyme 3-3. *Arch. Biochem. Biophys.*, 298 (1992) 667-677.
- 910 Kelley, M. and Vessey, D.A.: Structural comparison between the mitochondrial aralkyl-CoA and arylacetyl-CoA N-acyltransferases. *Biochem. J.*, 288 (1992) 315-317.
- 911 Kim, D.-H., Kim, H.-S. and Kobashi, K.: Purification and characterization of novel sulfotransferase obtained from *Klebsiella* K-36, an intestinal bacterium of rat. *J. Biochem. (Tokyo)*, 112 (1992) 456-460.
- 912 Martinez-Lara, E., Pascual, P., Toribio, F., López-Barea, J. and Bárcena, J.A.: Rapid method for the determination of glutathione transferase isoenzymes in crude extracts. *J. Chromatogr.*, 609 (1992) 141-146.
- 913 Nishikawa, A., Ihara, Y., Hatakeyama, M., Kangawa, K. and Taniguchi, N.: Purification, cDNA cloning, and expression of UDP-N-acetylglucosamine: β -D-mannoside β -1,4N-acetylglucosaminyltransferase III from rat kidney. *J. Biol. Chem.*, 267 (1992) 18199-18204.
- 914 Petrarulo, M., Pellegrino, S., Marangella, M., Cosseddu, D. and Linari, F.: High-performance liquid chromatographic microassay for L-alanine:glyoxylate aminotransferase activity in human liver. *Clin. Chim. Acta*, 208 (1992) 183-192.
- 915 Ploux, O., Soularue, P., Marquet, A., Gloeckler, R. and Lemoine, Y.: Investigation of the first step of biotin biosynthesis in *Bacillus sphaericus*. Purification and characterization of the pimeloyl-CoA synthase, and uptake of pimelate. *Biochem. J.*, 287 (1992) 685-690.
- 916 Rocchigiani, M., Micheli, V., Duley, J.A. and Simmonds, H.A.: Determination of nicotinamide phosphoribosyltransferase activity in human erythrocytes: high-performance liquid chromatography-linked method. *Anal. Biochem.*, 205 (1992) 334-336.
- 917 Schmid, D., Juassi, R. and Christen, P.: Precursor of mitochondrial aspartate aminotransferase synthesized in *Escherichia coli* is complexed with heat-shock protein DnaK. *Eur. J. Biochem.*, 208 (1992) 699-704.
- 918 Shenoy, B.C., Xie, Y., Park, V.L., Kumar, G.K., Beegen, H., Wood, H.G. and Samols, D.: The importance of methionine residues for the catalysis of the biotin enzyme, transcarboxylase. Analysis by site-directed mutagenesis. *J. Biol. Chem.*, 267 (1992) 18407-18412.
- 919 Sindelar, P. and Valtersson, C.: A dolichol acyltransferase present in rat and human postheparin plasma. *Biochem. Cell Biol.*, 70 (1992) 470-474.
- 920 Tabata, S. and Dohi, Y.: An Assay for oligo-(1 \rightarrow 4) \rightarrow (1 \rightarrow 4)-glucantransferase activity in the glycogen debranching enzyme system by usig HPLC with a pulsed amperometric detector. *Carbohydr. Res.*, 23 (1992) 179-183; *C.A.*, 117 (1992) 65227h.
- 921 Vannier, M.P., Thoirion, B., Morvan, C. and Demarty, M.: Localization of methyltransferase activities throughout the endomembrane system of flax (*Linum usitatissimum* L.) hypocotyls. *Biochem. J.*, 286 (1992) 863-868.
- 922 Wang, Y., Abernethy, J.L., Eckhardt, A.E. and Hill, R.L.: Purification and characterization of a UDP-GalNAc:polypeptide N-acetyl-galactosaminyltransferase specific for glycosylation of threonine residues. *J. Biol. Chem.*, 267 (1992) 12709-12716.

- 20c. *Transferases transferring phosphorus containing groups (E.C. 2.7.-)*
- 923 Appukuttan, P.S. and Basu, D.: Human brain creatine kinase binding to immobilized cibacron blue F3GA: characterization and use in purification of the enzyme. *Indian J. Biochem. Biophys.*, 29 (1992) 346-349; *C.A.*, 117 (1992) 145922w.
- 924 Basta, P., Strickland, M.B., Holmes, W., Loomis, C.R., Ballas, L.M. and Burns, D.J.: Sequence and expression of human protein kinase C- ϵ . *Biochim. Biophys. Acta*, 1132 (1992) 154-160.
- 925 Bidwai, A.P., Hanna, D.E. and Glover, C.V.C.: Purification and characterization of casein kinase II (CKII) from $\delta cka1 \delta cka2$ *Saccharomyces cerevisiae* rescued by *Drosophila* CKII subunits. The free catalytic subunit of casein kinase II is not toxic *in vivo*. *J. Biol. Chem.*, 267 (1992) 18790-18796.
- 926 Burrows, J.A. and Goward, C.R.: Purification and properties of DNA polymerase from *Bacillus caldotenax*. *Biochem. J.*, 287 (1992) 971-977.
- 927 Champion, S., Maisonhaute, C., Kim, M.H. and Best-Belpomme, M.: Characterization of the reverse transcriptase of 1731, a *Drosophila melanogaster* retrotransposon. *Eur. J. Biochem.*, 209 (1992) 523-531.
- 928 Chavan, A.J., Nemoto, Y., Narumiya, S., Kozaki, S. and Haley, B.E.: NAD⁺ Binding site of *Clostridium botulinum* C3 ADP-ribosyltransferase. Identification of peptide in the adenine ring binding domain using 2-azido NAD⁺. *J. Biol. Chem.*, 267 (1992) 14866-14870.
- 929 Dickens, M., Chin, J.E., Roth, R.A., Ellis, L., Denton, R.M. and Tavare, J.M.: Characterization of insulin-stimulated protein serine/threonine kinases in CHO cells expressing human insulin receptors with point and deletion mutations. *Biochem. J.*, 287 (1992) 201-209.
- 930 Emanuelli, M., Natalini, P., Raffaelli, N., Ruggieri, S., Vita, A. and Magni, G.: NAD Biosynthesis in human placenta: purification and characterization of homogeneous NMM adenylyltransferase. *Arch. Biochem. Biophys.*, 298 (1992) 29-34.
- 931 Epand, R.M., Stafford, A.R. and Lester, D.S.: Lipid vesicles which can bind to protein kinase C and activate the enzyme in the presence of EGTA. *Eur. J. Biochem.*, 208 (1992) 327-332.
- 932 Gold, M.R., Sanghera, J.S., Stewart, J. and Pelech, S.L.: Selective activation of p42 mitogen-activated protein (MAP) kinase in murine B lymphoma cell lines by membrane immunoglobulin cross-linking. Evidence for protein kinase C-independent and -dependent mechanisms of activation. *Biochem. J.*, 287 (1992) 269-276.
- 933 Green, P.C., Latshaw, S.P., Lador, U.S. and Kemp, R.G.: Identification of critical lysyl residues in the pyrophosphate-dependent phosphofructo-1-kinase of *Propionibacterium freudenreichii*. *Biochemistry*, 31 (1992) 4815-4821.
- 934 Hughes, K., Ramakrishna, S., Benjamin, W.B. and Woodgett, J.R.: Identification of multifunctional ATP-citrate lyase kinase as the α -isoform of glycogen synthase kinase-3. *Biochem. J.*, 288 (1992) 309-314.
- 935 Inoue, H., Yoshioka, T. and Hotta, Y.: Partial purification and characterization of membrane-associated diacylglycerol kinase of *Drosophila* heads. *Biochim. Biophys. Acta*, 1122 (1992) 219-224.
- 936 Just, I., Mohr, C., Schallehn, G., Menard, L., Didsbury, J.R., Vandekerckhove, J., van Damme, J. and Aktories, K.: Purification and characterization of an ADP-ribosyltransferase produced by *Clostridium limosum*. *J. Biol. Chem.*, 267 (1992) 10274-10280.
- 937 Ketcham, C.M. and Kornfeld, S.: Purification of UDP-N-acetylglucosamine:glycoprotein N-acetylglucosamine-1-phosphotransferase from *Acanthamoeba castellanii* and identification of a subunit of the enzyme. *J. Biol. Chem.*, 267 (1992) 11645-11653.
- 938 Lew, J., Beaudette, K., Litwin, C.M.E. and Wang, J.H.: Purification and characterization of a novel proline-directed protein kinase from bovine brain. *J. Biol. Chem.*, 267 (1992) 13383-13390.
- 939 Logsdon, N., Lee, C.G.L. and Harper, J.W.: Selective 5' modification of T7 RNA polymerase transcripts. *Anal. Biochem.*, 205 (1992) 36-41.
- 940 Lydon, N.B., Gay, B., Mett, H., Murray, B., Liebertanz, J., Gotzwiller, A., Piwnica-Worms, H., Roberts, T.M. and McGlynn, E.: Purification and biochemical characterization of non-myristoylated recombinant pp60^{c-src} kinase. *Biochem. J.*, 287 (1992) 985-993.
- 941 Mackintosh, R.W., Davies, S.P., Clarke, P.R., Weekes, J., Gillespie, J.G., Gibb, B.J. and Hardie, D.G.: Evidence for a protein kinase cascade in higher plants. 3-Hydroxy-3-methylglutaryl-CoA reductase kinase. *Eur. J. Biochem.*, 209 (1992) 923-931.
- 942 Matter, W.F., Brown, R.F. and Vlahos, C.J.: The inhibition of phosphatidylinositol 3-kinase by quercetin and analogs. *Biochem. Biophys. Res. Commun.*, 186 (1992) 624-631.
- 943 Mitchell, L.L.W. and Cooperman, B.S.: Active site studies of human immunodeficiency virus reverse transcriptase. *Biochemistry*, 31 (1992) 7707-7713.
- 944 Mitev, V.I., Angelov, I.V. and Marekov, L.N.: Identification and purification of a novel serine/threonine messenger-independent growth-related protein kinase from lactating goat mammary gland. *Biochim. Biophys. Acta*, 1117 (1992) 90-96.
- 945 Mori, S., Yokota, T., Shigeta, S., Sato, M., Sasaki, M., Nagai, Y., Onami, T. and Sayama, S.: (Purification of varicella zoster virus thymidine kinase by thymidine agarose gel affinity chromatography). *Fukushima Igaku Zasshi*, 41 (1991) 341-347; *C.A.*, 117 (1992) 145906u.
- 946 Nakanishi, H. and Exton, J.H.: Purification and characterization of the ζ isoform of protein kinase C from bovine kidney. *J. Biol. Chem.*, 267 (1992) 16347-16354.
- 947 Okazaki, H., Tani, C., Ando, M., Ishii, K., Ishibashi, S., Nishimura, Y. and Kato, K.: Possible involvement of cathepsin L in processing of rat liver hexokinase to eliminate mitochondria-binding ability. *J. Biochem. (Tokyo)*, 112 (1992) 409-413.
- 948 Palczewski, K., Buczylo, J., van Hooser, P., Carr, A.A., Huddleston, M.J. and Crabb, J.W.: Identification of the autophosphorylation sites in rhodopsin kinase. *J. Biol. Chem.*, 267 (1992) 18991-18998.
- 949 Pontremoli, S., Viotti, P.L., Michetti, M., Salamino, F., Sparatore, B. and Melloni, E.: Modulation of inhibitory efficiency of rat skeletal muscle calpastatin by phosphorylation. *Biochem. Biophys. Res. Commun.*, 187 (1992) 751-759.
- 950 Salvucci, M.E., Chavan, A.J. and Haley, B.E.: Identification of peptides from the adenine binding domains of ATP and AMP in adenylylase kinase: isolation of photoaffinity-labeled peptides by metal chelate chromatography. *Biochemistry*, 31 (1992) 4479-4487.

- 951 Seger, R., Ahn, N.G., Posada, J., Munar, E.S., Jensen, A.M., Cooper, J.A., Cobb, M.H. and Krebs, E.G.: Purification and characterization of mitogen-activated protein kinase activator(s) from epidermal growth factor-stimulated A431 cells. *J. Biol. Chem.*, 267 (1992) 14373-14381.
- 952 Sharova, N.P., Eliseeva, E.D. and Mikhailov, V.S.: (Changes in activity of two forms of DNA polymerase α during embryogenesis of the teleost fish *Misgurnus fossilis* L.) *Biokhimiya (Moscow)*, 57 (1992) 937-944.
- 953 Stamford, N.P.J., Lilley, P.E. and Dixon, N.E.: Enriched sources of *Escherichia coli* replication proteins. The dnaG primase is a zinc metalloprotein. *Biochim. Biophys. Acta*, 1132 (1992) 17-25.
- See also 681, 737.
- 20d. *Hydrolases, acting on ester bonds (E.C. 3.1.-.-)*
- 954 Allenmark, S. and Ohlsson, A.: Studies of the heterogeneity of a *Candida cylindracea* (rugosa) lipase: monitoring of esterolytic activity and enantioselectivity by chiral liquid chromatography *Biocatalysis*, 6 (1992) 211-221; *C.A.*, 117 (1992) 166352a.
- 955 Chang, T.-C., Huang, S.-M., Huang, T.-M. and Chang, G.-G.: Human placental alkaline phosphatase. An improved purification procedure and kinetic studies. *Eur. J. Biochem.*, 209 (1992) 241-247.
- 956 Chihara-Siomi, M., Yoshikawa, K., Oshima-Hirayama, N., Yamamoto, K., Sogabe, Y., Nakatani, T., Nishioka, T. and Oda, J.: Purification, molecular cloning and expression of lipase from *Pseudomonas aeruginosa* *Arch. Biochem. Biophys.*, 296 (1992) 505-513.
- 957 Faber, A. and Aviram, I.: Human neutrophil cytosolic phospholipase C: partial characterization. *Biochim. Biophys. Acta*, 1128 (1992) 8-13.
- 958 Fonda, M.L.: Purification and characterization of vitamin B₆-phosphate phosphatase from erythrocytes. *J. Biol. Chem.*, 267 (1992) 15978-15983.
- 959 Friedrich, G., Klumpp, S., Russell, C.B., Hinrichsen, R.D., Keller, R. and Schultz, J.E.: Purification, characterization and structure of protein phosphatase 1 from the cilia of *Paramecium tetraurelia* *Eur. J. Biochem.*, 209 (1992) 43-49.
- 960 Fujimoto, S., Murakami, K., Hosoda, T., Yamamoto, Y., Watanabe, K., Morinaka, Y. and Ohara, A.: Characterization of cationic acid phosphatase isozyme from rat liver mitochondria. *Chem. Pharm. Bull.*, 40 (1992) 1231-1235.
- 961 Gadella, B.M., Colenbrander, B., van Golde, L.M.G. and Lopes-Cardozo, M.: Characterization of three arylsulfatases in semen: seminolipid sulfohydrolase activity is present in seminal plasma. *Biochim. Biophys. Acta*, 1128 (1992) 155-162.
- 962 Gite, S.U. and Shankar, V.: Affinity purification of ribonuclease T1 on 5'-AMP-Sepharose. *Biotechnol. Tech.*, 6 (1992) 245-248; *C.A.*, 117 (1992) 127031a.
- 963 Hiraga, A., Munakata, H., Hata, K., Suzuki, Y. and Tsuike, S.: Purification and characterization of a rat liver protein-tyrosine phosphatase with sequence similarity to src-homology region 2. *Eur. J. Biochem.*, 209 (1992) 195-206.
- 964 Hori, N., Iwai, S., Inoue, H. and Ohtsuka, E.: Photoaffinity labeling of T4 endonuclease V with a substrate containing a phenyldiazirine derivative. *J. Biol. Chem.*, 267 (1992) 15591-15594.
- 965 Jayanthi, L.D., Balasubramanian, N. and Balasubramanian, A.S.: Cholinesterases exhibiting aryl acylamidase activity in human amniotic fluid. *Clin. Chim. Acta*, 205 (1992) 157-155.
- 966 Jobbagy, Z., Izsvak, Z. and Duda, E.: Positive co-operative interaction between the subunits of CeqI restriction endonuclease. *Biochem. J.*, 286 (1992) 85-88.
- 967 Ketterman, A.J., Jayawardena, K.G.I. and Hemingway, J.: Purification and characterization of a carboxylesterase involved in insecticide resistance from the mosquito *Culex quinquefasciatus* *Biochem. J.*, 287 (1992) 355-360.
- 968 Kingston-Smith, A.H., Major, I., Parry, M.A.J. and Keys, A.J.: Purification and properties of a phosphatase in French bean (*Phaseolus vulgaris* L.) leaves that hydrolyses 2'-carboxy-D-arabinitol 1-phosphate. *Biochem. J.*, 287 (1992) 821-825.
- 969 Kobayashi, T., Honke, K., Jin, T., Gasa, S., Miyazaki, T. and Makita, A.: Components and proteolytic processing sites of arylsulfatase B from human placenta. *Biochim. Biophys. Acta*, 1159 (1992) 243-247.
- 970 Lessinger, J.M., Tavidou, A., Arzoglou, P. and Ferard, G.: Interest of using a purified, stable and commutable preparation of human pancreatic lipase in indirect assays. *Anal. Lett.*, 25 (1992) 1453-1468.
- 971 Mitsui, T., Inagaki, M. and Ikebe, M.: Purification and characterization of smooth muscle myosin-associated phosphatase from chicken gizzards. *J. Biol. Chem.*, 267 (1992) 16727-16735.
- 972 Murakami, M., Kudo, I., Suwa, Y. and Inoue, K.: Release of 14-kDa group-II phospholipase A₂ from activated mast cells and its possible involvement in the regulation of the degranulation process. *Eur. J. Biochem.*, 209 (1992) 257-265.
- 973 Nakamura, A., Koide, Y., Miyazaki, H., Kitamura, A., Masaki, H., Beppu, T. and Uozumi, T.: Gene cloning and characterization of a novel extracellular ribonuclease of *Bacillus subtilis* *Eur. J. Biochem.*, 209 (1992) 121-127.
- 974 Park, D., Jhon, D., Kriz, R., Knopf, J. and Rhee, S.G.: Cloning, sequencing, expression, and G_q-independent activation of phospholipase C- β 2. *J. Biol. Chem.*, 267 (1992) 16048-16055.
- 975 Peterson, J., Fujimoto, W.Y. and Brunzell, J.D.: Human lipoprotein lipase: relationship of activity, heparin affinity, and conformation as studied with monoclonal antibodies. *J. Lipid Res.*, 33 (1992) 1165-1170.
- 976 Polya, G.M. and Wettenhall, R.E.H.: Rapid purification and N-terminal sequencing of a potato tuber cyclic nucleotide binding phosphatase. *Biochim. Biophys. Acta*, 1159 (1992) 179-184.
- 977 Poretz, R.D., Yang, R.S., Canas, B., Lackland, H., Stein, S. and Manowitz, P.: The structural basis for the electrophoretic isoforms of normal and variant human platelet arylsulphatase A. *Biochem. J.*, 287 (1992) 979-983.
- 978 Shimizu, S., Kataoka, M., Shimizu, K., Hirakata, M., Sakamoto, K. and Yamada, H.: Purification and characterization of a novel lactonohydrolase, catalyzing the hydrolysis of aldonate lactones and aromatic lactones, from *Fusarium oxysporum* *Eur. J. Biochem.*, 209 (1992) 383-390.
- 979 Shnyreva, M.G., Tsuprun, V.L., Stel'mashchuk, V.Ya. and Egorov, S.N.: (Study of the quaternary structure of extracellular reversible acid phosphatase from the yeast *Saccharomyces cerevisiae*). *Biokhimiya (Moscow)*, 57 (1992) 1100-1108.
- 980 Spaargaren, M., Wissink, S., Defize, L.H.K., de Laat, S.W. and Boonstra, J.: Characterization and identification of an epidermal-growth-factor-activated phospholipase A₂. *Biochem. J.*, 287 (1992) 37-43.

- 981 Speth, M. and Schulze, H.-U.: The purification of a detergent-soluble glucose-6-phosphatase from rat liver. *Eur. J. Biochem.*, 208 (1992) 643-650.
- 982 Tavidou, A., Avranas, A. and Arzoglou, P.: A mathematical approach to lipolysis based on the interrelationship of physicochemical and biochemical data. *Biochem. Biophys. Res. Commun.*, 186 (1992) 746-752.
- 983 Wang, Y.-M., Lu, P.-J., Ho, C.-L. and Tsai, I.-H.: Characterization and molecular cloning of neurotoxic phospholipases A₂ from Taiwan viper (*Vipera russelli formosensis*). *Eur. J. Biochem.*, 209 (1992) 635-641.
- 984 Winder, S.J., Pato, M.D. and Walsh, M.P.: Purification and characterization of calponin phosphatase from smooth muscle. Effect of dephosphorylation on calponin function. *Biochem. J.*, 286 (1992) 197-203.
- 985 Yang, S.K., Hsieh, Y.-Y., Chang, W.-C. and Huang, J.-D.: Enantioselectivity of microsomal and cytosolic esterases in rat intestinal mucosa. *Drug Metab. Disp.*, 20 (1992) 719-725.
- For additional information see C.A.:
117 (1992) 43558q.
- See also 498.
- 20e. *Hydrolases, acting on glycosyl compounds (E.C. 3.2.-.-)*
- 986 Ajandouz, E.H., Abe, J.-i., Svensson, B. and Marchis-Mouren, G.: Barley malt- α -amylase. Purification, action pattern, and sub-site mapping of isozyme 1 and two members of the isozyme 2 subfamily using *p*-nitrophenylated maltooligosaccharide substrates. *Biochim. Biophys. Acta*, 1159 (1992) 193-202.
- 987 Al Daher, S., de Gasperi, R., Daniel, P., Hirani, S., Warren, C. and Winchester, B.: Substrate specificity of human liver neutral α -mannosidase. *Biochem. J.*, 286 (1992) 47-53.
- 988 Coelho, L.C.B.B., Hutchinson, W., Koshy, A., Robinson, D.S. and Stirling, J.L.: Affinity chromatography of β -N-acetylhexosaminidases on columns with mixed charge and affinity functions. *Appl. Biochem. Biotechnol.*, 32 (1992) 171-185; C.A., 117 (1992) 166307q.
- 989 De Gasperi, R., Al Daher, S., Winchester, B.G. and Warren, C.D.: Substrate specificity of the bovine and feline neutral α -mannosidases. *Biochem. J.*, 286 (1992) 55-63.
- 990 Inoue, M., Yamada, H., Yasukochi, T., Kuroki, R., Miki, T., Horiuchi, T. and Imoto, T.: Multiple role of hydrophobicity of tryptophan-108 in chicken lysozyme: structural stability, saccharide binding ability, and abnormal pK_a of glutamic acid-35. *Biochemistry*, 31 (1992) 5545-5553.
- 991 Kanaya, E. and Kikuchi, M.: Folding of human lysozyme *in vivo* by the formation of an alternative disulfide bond. *J. Biol. Chem.*, 267 (1992) 15111-15115.
- 992 Mullis, K.G. and Ketcham, C.M.: The synthesis of substrates and two assays for the detection of N-acetylglucosamine-1-phosphodiester α -N-acetylglucosaminidase (uncovering enzyme). *Anal. Biochem.*, 205 (1992) 200-207.
- 993 Omichi, K. and Hase, S.: Detection of human urinary α -amylase encoded by the AMY2B gene using a fluorogenic substrate, FG5P. *J. Biochem. (Tokyo)*, 112 (1992) 303-305.
- 994 Rapp, P.: Formation, separation and characterization of three β -1,3-glucanases from *Sclerotium glaucanicum* *Biochim. Biophys. Acta*, 1117 (1992) 7-14.
- 995 Skudlarek, M.D., Tulsiani, D.R.P. and Orgebin-Crist, M.-C.: Rat epididymal luminal fluid acid β -D-galactosidase optimally hydrolyses glycoprotein substrate at neutral pH. *Biochem. J.*, 286 (1992) 907-914.
- 996 Takagaki, K., Nakamura, T., Takeda, Y., Daidouji, K. and Endo, M.: A new endo- β -galactosidase acting on the Gal β 1-3Gal linkage of the proteoglycan linkage region. *J. Biol. Chem.*, 267 (1992) 18558-18563.
- 997 Tassi, C., Beccari, T., Casini, A. and Orlicchio, A.: β -N-Acetylhexosaminidase in the urine, kidney and serum of bromobenzene-intoxicated mice. *Clin. Chim. Acta*, 206 (1992) 231-239.
- 998 Wang, J., Xu, P. and Fincher, G.B.: Purification, characterization and gene structure of (1 \rightarrow 3)- β -glucanase isoenzyme GIII from barley (*Hordeum vulgare*). *Eur. J. Biochem.*, 209 (1992) 103-109.
- 999 Watanabe, T., Sato, T., Yoshioka, S., Koshijima, T. and Kuwahara, M.: Purification and properties of *Aspergillus niger* β -glucosidase. *Eur. J. Biochem.*, 209 (1992) 651-659.
- 1000 Yao, K., Ubuka, T., Masuoka, N., Kinuta, M., Ohta, J., Teraoka, T. and Futani, S.: Assay of sialidase activity using ion-exchange chromatography and acidic ninhydrin reaction. *J. Chromatogr.*, 581 (1992) 11-15.
- 1001 Zeng, Y.-c., Li, Y.-T., Gu, Y.-j. and Zhang, S.-Z.: Purification and characterization of a strictly specific β -D-fucosidase from *Aspergillus phoenicis* *Arch. Biochem. Biophys.*, 298 (1992) 226-230.
- For additional information see C.A.:
117 (1992) 107310n.
- See also 313.
- 20f. *Other hydrolases*
- 1002 Aeed, P.A., Guido, D.M., Mathews, W.R. and Elhammer, A.P.: Characterization of the oligosaccharide structures on recombinant human prorenin expressed in Chinese hamster ovary cells. *Biochemistry*, 31 (1992) 6951-6961.
- 1003 Baiyles, E.M. and Hutton, J.C.: Kinetic analysis of the type-1 proinsulin endopeptidase by a monoclonal antibody-based immunoabsorbent assay. *Biochem. J.*, 286 (1992) 223-229.
- 1004 Bar-Zvi, D., Bar, I., Yoshida, M. and Shavit, N.: Covalent binding of 3'-O-(4-benzoyl)benzoyl adenosine 5'-triphosphate (BzATP) to the isolated α and β subunit and the $\alpha_3\beta_3$ core complex of TF₁. Covalent binding of BzATP prevents association of α and β subunits and induces dissociation of the $\alpha_3\beta_3$ core complex. *J. Biol. Chem.*, 267 (1992) 11029-11033.
- 1005 Basak, A., Yuan, X.W., Seidah, N.G., Chrétien, M. and Lazure, C.: Affinity purification of proteinases by a combination of immobilized peptidyl aldehyde and semicarbazone. *J. Chromatogr.*, 581 (1992) 17-29.
- 1006 Bawab, W., Querido, E., Crine, P. and DesGroseillers, L.: Identification and characterization of aminopeptidases from *Aplysia californica* *Biochem. J.*, 286 (1992) 967-975.
- 1007 Boketsky, Yu.R., Skoblov, Yu.S., Khodova, O.M. and Rabinovich, P.M.: (Purification and properties of GTP-cyclohydrolase of *Bacillus subtilis*). *Biokhimiya (Moscow)*, 57 (1992) 1021-1030.

- 1008 Bos, R., Berger, L. and Nieuwenhuizen, W.: One-step purification of tissue-type plasminogen activator using affinity chromatography with a special monoclonal antibody under mild conditions. *Biochim. Biophys. Acta*, 1117 (1992) 188-192.
- 1009 Buttle, D.J. and Saklatvala, J.: Lysosomal cysteine endopeptidases mediate interleukin 1-stimulated cartilage proteoglycan degradation. *Biochem. J.*, 287 (1992) 657-661.
- 1010 Capobianco, J.O., Lerner, C.G. and Goldman, R.C.: Application of a fluorogenic substrate in the assay of proteolytic activity and in the discovery of a potent inhibitor of *Candida albicans* aspartic proteinase. *Anal. Biochem.*, 204 (1992) 96-102.
- 1011 Chiou, S.-H., Hung, C.-C. and Huang, K.-F.: Characterization of a protease with α - and β -fibrinogenase activity from the western diamondback rattlesnake, *Crotalus atrox*. *Biochem. Biophys. Res. Commun.*, 187 (1992) 389-396.
- 1012 Citarella, F., Aiuti, A., La Porta, C., Russo, G., Pietropaolo, C., Rinaldi, M. and Fantoni, A.: Control of human coagulation by recombinant serine proteases. Blood clotting is activated by recombinant factor XII deleted of five regulatory domains. *Eur. J. Biochem.*, 208 (1992) 23-30.
- 1013 Dahlmann, B., Kuehn, L., Grziwa, A., Zwickl, P. and Baumeister, W.: Biochemical properties of the proteasome from *Thermoplasma acidophilum*. *Eur. J. Biochem.*, 208 (1992) 789-797.
- 1014 Dahms, P. and Mentlein, R.: Purification of the main somatostatin-degrading proteases from rat and pig brains, their action on other neuropeptides, and their identification as endopeptidases 24.15 and 24.16. *Eur. J. Biochem.*, 208 (1992) 145-154.
- 1015 Dando, P.M. and Barrett, A.J.: Quantification of peptide aldehyde ligands immobilized for the affinity chromatography of endopeptidases. *Anal. Biochem.*, 204 (1992) 328-331.
- 1016 Dick, L.R., Moomaw, C.R., Pramanik, B.C., DeMartino, G.N. and Slaughter, C.A.: Identification and localization of a cysteinyl residue critical for the trypsin-like catalytic activity of the proteasome. *Biochemistry*, 31 (1992) 7347-7355.
- 1017 Ilian, M.A. and Forsberg, N.E.: Gene expression of calpains and their specific endogenous inhibitor, calpastatin, in skeletal muscle of fed and fasted rabbits. *Biochem. J.*, 287 (1992) 163-171.
- 1018 Khropycheva, R.P., Marinchenko, G.V. and Fedosimov, V.A.: (Prolactin hydrolysis by mitochondrial serine proteinase from mammary gland secretory cells). *Biokhimiya (Moscow)*, 57 (1992) 945-955.
- 1019 Kobayashi, R., Nakahima, K., Sugano, K., Nishikawa, M. and Akiyama, S.: Gas phase derivatization of ammonia with 4-fluoro-7-nitrobenzo-2-oxa-1,3-diazole and its application to urease assay. *Chem. Pharm. Bull.*, 40 (1992) 1327-1328.
- 1020 Kole, H.K., Smith, D.R. and Lenard, J.: Characterization and partial purification of an insulinase from *Neurospora crassa*. *Arch. Biochem. Biophys.*, 297 (1992) 199-204.
- 1021 Kronheim, S.R., Mumma, A., Greenstreet, T., Glackin, P.J., van Ness, K., March, C.J. and Black, R.A.: Purification of interleukin-1 β converting enzyme, the protease that cleaves the interleukin-1 β precursor. *Arch. Biochem. Biophys.*, 296 (1992) 698-703.
- 1022 Lindahl, P., Nycander, M., Ylännejärvi, K., Pol, E. and Björk, I.: Characterization by rapid-kinetic and equilibrium methods of the interaction between N-terminally truncated forms of chicken cystatin and the cysteine proteinases papain and actinidin. *Biochem. J.*, 286 (1992) 165-171.
- 1023 Makarchikov, A.F. and Chernikevich, I.P.: Purification and characterization of thiamine triphosphatase from bovine brain. *Biochim. Biophys. Acta*, 1117 (1992) 326-332.
- 1024 Mäkinen, K.K., Mäkinen, P.-L. and Syed, S.A.: Purification and substrate specificity of an endopeptidase from the human oral spirochete *Treponema denticola* ATCC 35405, active on furylacryloyl-Leu-Gly-Pro-Ala and bradykinin. *J. Biol. Chem.*, 267 (1992) 14285-14293.
- 1025 Meulenberg, R., Pronk, J.T., Frank, J., Hezeu, W., Bos, P. and Kuenen, J.G.: Purification and partial characterization of a thermostable trithionate hydrolase from the acidophilic sulphur oxidizer *Thiobacillus acidophilus*. *Eur. J. Biochem.*, 209 (1992) 367-374.
- 1026 Miller, D.A., Sayad, K.U., Kulathila, R., Beaudry, G.A., Merkler, D.J. and Bertelsen, A.H.: Characterization of a bifunctional peptidylglycine enzyme expressed in Chinese hamster ovary cells. *Arch. Biochem. Biophys.*, 298 (1992) 380-388.
- 1027 Miyata, T., Aruga, R., Umeyama, H., Bezeaud, A., Guillin, M.-C. and Iwanaga, S.: Prothrombin Salakta: substitution of glutamic acid-466 by alanine reduces the fibrinogen clotting activity and the esterase activity. *Biochemistry*, 31 (1992) 7457-7462.
- 1028 Nakamura-Imajo, N., Satomura, S., Matsuura, S. and Murakami, K.: Renin assay using a fluorogenic substrate and high performance liquid chromatography. *Clin. Chim. Acta*, 211 (1992) 47-57.
- 1029 Ohshita, T., Nikawa, T., Towatari, T. and Katunuma, N.: Effects of selective inhibition of cathepsin B and general inhibition of cysteine proteinases on lysosomal proteolysis in rat liver *in vivo* and *in vitro*. *Eur. J. Biochem.*, 209 (1992) 223-231.
- 1030 Paik, S.R. and Lewis, R.V.: Purification and characterization of a putative proenkephalin cleaving enzyme. *Arch. Biochem. Biophys.*, 296 (1992) 617-629.
- 1031 Retzios, A.D. and Markland, F.S., Jr.: Purification, characterization, and fibrinogen cleavage sites of three fibrinolytic enzymes from the venom of *Crotalus basiliscus basiliscus*. *Biochemistry*, 31 (1992) 4547-4557.
- 1032 Rhyu, M.R., Nishimura, T., Kato, Y., Okitani, A. and Kato, H.: Purification and properties of aminopeptidase H from chicken skeletal muscle. *Eur. J. Biochem.*, 208 (1992) 53-59.
- 1033 Richter, O.-M.H. and Schäfer, G.: Purification and enzymic characterization of the cytoplasmic pyrophosphatase from the thermoacidophilic archaeobacterium *Thermoplasma acidophilum*. *Eur. J. Biochem.*, 209 (1992) 343-349.
- 1034 Saito, H., Kurata, S. and Natori, S.: Purification and characterization of a hemocyte proteinase of *Sarcophaga*, possibly participating in elimination of foreign substances. *Eur. J. Biochem.*, 209 (1992) 939-944.
- 1035 Santana, J.M., Grellier, P., Rodier, M.-H., Schrevel, J. and Teixeira, A.: Purification and characterization of a new 120 kDa alkaline proteinase of *Trypanosoma cruzi*. *Biochem. Biophys. Res. Commun.*, 187 (1992) 1466-1473.
- 1036 Saxena, M., Singhal, S.S., Awasthi, S., Singh, S.V., Labelle, E.F., Zimniak, P. and Awasthi, Y.C.: Dinitrophenyl S-glutathione ATPase, purified from human muscle catalyzes ATP hydrolysis in the presence of leukotrienes. *Arch. Biochem. Biophys.*, 298 (1992) 231-237.
- 1037 Schönherr, E., Jones, G.A. and Slakey, L.L.: Gastric and salivary mucins inhibit angiotensin-converting enzyme. Inhibition is partly due to oligosaccharides. *Biochem. J.*, 286 (1992) 425-433.

- 1038 Shimizu, S., Imanaka, T., Takano, T. and Ohkuma, S.: Induction and characterization of two types of ATPase on rat liver peroxisomes. *J. Biochem. (Tokyo)*, 112 (1992) 376-384.
- 1039 Shimizu, S., Ogawa, J., Chung, M.C.-M. and Yamada, H.: Purification and characterization of a novel enzyme, arylalkyl acylamidase, from *Pseudomonas putida* Sc2. *Eur. J. Biochem.*, 209 (1992) 375-382.
- 1040 Stefanova, H.I., East, J.M., Gore, M.G. and Lee, A.G.: Labeling the (Ca²⁺-Mg²⁺)-ATPase of sarcoplasmic reticulum with 4-(bromomethyl)-6,7-dimethoxycoumarin: detection of conformational changes. *Biochemistry*, 31 (1992) 6023-6031.
- 1041 Tanty, A.M. and Brelle, J.L.: Determination of enzymes in rennets. Simplification of the official method. *Lait*, 72 (1992) 103-107; *C.A.*, 117 (1992) 68564b.
- 1042 Treuheit, M.J., Ateai, A., Kirley, T.L. and Wallick, E.T.: Purification of the glycosylated beta subunit of (Na,K)-ATPase by lectin affinity chromatography. *Chromatographia*, 33 (1992) 521-524.
- 1043 Zakharov, S.D. and Kuz'mina, V.P.: (Subunit compositions of ATP-synthase from the thermophilic bacterium *Thermus thermophilus*). *Biokhimiya (Moscow)*, 57 (1992) 777-786.
- 1044 Zhuo, S., Garrod, S., Miller, P. and Allison, W.S.: Irradiation of the bovine mitochondrial F₁-ATPase previously inactivated with 5'-p-fluorosulfonylbenzoyl-8-azido-[³H]adenosine cross-links His-β427 to Tyr-β345 within the same β subunit. *J. Biol. Chem.*, 267 (1992) 12916-12927.
- For additional information see *C.A.*:
117 (1992) 146716u.
- See also 647, 731.
- 20g. Lyases
- 1045 Aoki, T., Tanaka, T. and Watabe, H.: Purification and characterization of γ-enolase from various mammals. *Chem. Pharm. Bull.*, 40 (1992) 1236-1239.
- 1046 Baur, B., Dietz, K.-J. and Winter, K.: Regulatory protein phosphorylation of phosphoenolpyruvate carboxylase in the facultative crassulacean-acid-metabolism plant *Mesembryanthemum crystallinum* L. *Eur. J. Biochem.*, 209 (1992) 95-101.
- 1047 Bogaards, J.J.P., van Ommen, B. and van Bladeren, P.J.: Purification and characterization of eight glutathione S-transferase isoenzymes of hamster. Comparison of subunit composition of enzymes from liver, kidney, testis, pancreas and trachea. *Biochem. J.*, 286 (1992) 383-388.
- 1048 Cook, L., Nagi, M.N., Suneja, S.K., Hand, A.R. and Cinti, D.L.: Evidence that β-hydroxyacyl-CoA dehydrase purified from rat liver microsomes is of peroxisomal origin. *Biochem. J.*, 287 (1992) 91-100.
- 1049 Deans, N.L., Allison, R.D. and Purich, D.L.: Steady-state kinetic mechanism of bovine brain tubulin:tyrosine ligase. *Biochem. J.*, 286 (1992) 243-251.
- 1050 Khan, A.S., van Driessche, E., Kanarek, L. and Beeckmans, S.: The purification of physicochemical characterization of maize (*Zea mays* L.) isocitrate lyase. *Arch. Biochem. Biophys.*, 297 (1992) 9-18.
- 1051 Ko, Y.H., Cremona, C.R. and McFadden, B.A.: Vanadate-dependent photomodification of serine 319 and 321 in the active site of isocitrate lyase from *Escherichia coli* *J. Biol. Chem.*, 267 (1992) 91-95.
- 1052 Kohen, A., Jacob, A. and Naasov, T.: Mechanistic studies of 3-deoxy-D-manno-2-octulosonate-8-phosphate synthase from *Escherichia coli* *Eur. J. Biochem.*, 208 (1992) 443-449.
- 1053 Lill, U., Lefrank, S., Henschen, A. and Eggerer, H.: Conversion, by limited proteolysis, of an archaeobacterial citrate synthase into essentially a citryl-CoA hydrolase. *Eur. J. Biochem.*, 208 (1992) 459-466.
- 1054 Nicholson, D.W., Klemba, M.W., Rasper, D.M., Metters, K.M., Zamboni, R.J. and Ford-Hutchinson, A.W.: Purification of human leukotriene C₄ synthase from dimethylsulfoxide-differentiated U937 cells. *Eur. J. Biochem.*, 209 (1992) 725-734.
- 1055 Poulsen, C., van der Heijden, R. and Verpoorte, R.: Assay of isochorismate synthase from plant cell cultures by high-performance liquid chromatography. *Phytochemistry*, 30 (1991) 2873-2876; *C.A.*, 116 (1992) 54106w.
- 1056 Shantz, L.M., Stanley, B.A., Secrist, J.A. and Pegg, A.E.: Purification of human S-adenosylmethionine decarboxylase expressed in *Escherichia coli* and use of this protein to investigate the mechanism of inhibition by the irreversible inhibitors, 5'-deoxy-5'-[(3-hydrazinopropyl)methylamino]adenosine and 5'-{[(Z)-4-amino-2-butenyl]methylamino}-5'-deoxyadenosine. *Biochemistry*, 31 (1992) 6848-6855.
- 1057 Staffas, L., Mankowitz, L., Söderström, M., Blanck, A., Porsch-Hällström, I., Sundberg, C., Mannervik, B., Olin, B., Rydström, J. et al.: Further characterization of hormonal regulation of glutathione transferase in rat liver and adrenal glands. Sex differences and demonstration that growth hormone regulates the hepatic levels. *Biochem. J.*, 286 (1992) 65-72.
- 1058 Wang, P., Matthews, D.E. and van Etten, H.D.: Purification and characterization of cyanide hydratase from the phytopathogenic fungus *Gloeocercospora sorghi* *Arch. Biochem. Biophys.*, 298 (1992) 569-575.
- See also 437, 675.
- 20h. Isomerases
- 1059 Abe, I., Bai, M., Xiao, X.-Y. and Prestwich, G.D.: Affinity labeling of vertebrate oxidosqualene cyclases with a tritiated suicide substrate. *Biochem. Biophys. Res. Commun.*, 187 (1992) 32-38.
- 1060 Hornbogen, T., Pieper, R., Hoffmann, K., Kleinkauf, H. and Zocher, R.: Two new cyclophilins from *Fusarium sambucinum* and *Aspergillus niger*: resistance of cyclophilin/cyclosporin A complexes against proteolysis. *Biochem. Biophys. Res. Commun.*, 187 (1992) 791-796.
- 1061 Siegal, G., Turchi, J.J., Jessee, C.B., Myers, T.W. and Bambara, R.A.: A novel DNA helicase from calf thymus. *J. Biol. Chem.*, 267 (1992) 13629-13635.
- 20i. Ligases
- 1062 Aoki, H., Yaworsky, P.J., Patel, S.D., Margolin-Brzezinski, F., Park, K.S. and Ganoza, M.C.: The asparaginyl-tRNA synthetase gene encodes one of the complementing factors for thermosensitive translation in the *Escherichia coli* mutant strain, N4316. *Eur. J. Biochem.*, 209 (1992) 511-521.
- 1063 Bianchi, C., Perego, R. and Delmonte, U.: Polypeptide composition of the 8S form of prolyl-tRNA synthetase from rat liver. *Biochem. Biophys. Res. Commun.*, 187 (1992) 1071-1076.

- 1064 Bora, P.S., Wu, X., Spilburg, C.A. and Lange, L.G.: Purification and characterization of fatty acid ethyl ester synthase-II from human myocardium. *J. Biol. Chem.*, 267 (1992) 13217-13221.
- 1065 Li, S. and Cronan, J.E., Jr.: The genes encoding the two carboxyltransferase subunits of *Escherichia coli* acetyl-CoA carboxylase. *J. Biol. Chem.*, 267 (1992) 16841-16847.
- 1066 Pajares, M.A., Durán, C., Corrales, F., Pliego, M.M. and Mato, J.M.: Modulation of rat liver S-adenosylmethionine synthetase activity by glutathione. *J. Biol. Chem.*, 267 (1992) 17598-17605.
- 1067 Pollack, S.J., Freeman, S., Pompliano, D.L. and Knowles, J.R.: Cloning, overexpression and mechanistic studies of carboxyphosphoenolpyruvate mutase from *Streptomyces hygroscopicus*. *Eur. J. Biochem.*, 209 (1992) 735-743.
- 1068 Shen, Z., Fice, D. and Byers, D.M.: Preparation of fatty-acylated derivatives of acyl carrier protein using *Vibrio harveyi* acyl-ACP synthetase. *Anal. Biochem.*, 204 (1992) 34-39.
- 1069 Ting, S., Bogner, P. and Dignam, J.D.: Isolation of prolyl-tRNA synthetase as a free form and as a form associated with glutamyl-tRNA synthetase. *J. Biol. Chem.*, 267 (1992) 17701-17709.
- 20j. *Complex mixtures and incompletely identified enzymes*
- 1070 Andersen, N.H. and Rhee, J.: An approach to isolation of thromboxane synthase (TX-SYN) by ligand tethered affinity techniques. *Bull. Korean Chem. Soc.*, 13 (1992) 119-122; *C.A.*, 117 (1992) 145829w.
- 1071 Heller, M., Bieri, S. and Brodbeck, U.: A novel form of glycosylphosphatidylinositol-anchor converting activity with a specificity of a phospholipase D in mammalian liver membranes. *Biochim. Biophys. Acta*, 1109 (1992) 109-116.
- See also 190.
21. PURINES, PYRIMIDINES, NUCLEIC ACIDS AND THEIR CONSTITUENTS
- 21a. *Purines, pyrimidines, nucleosides, nucleotides*
- 1072 Angelastro, J.M. and Purich, D.L.: Adenine and guanine nucleotide content of Triton-extracted cytoskeletal fractions on non-muscle cells. *Anal. Biochem.*, 204 (1992) 47-52.
- 1073 Berube, R., Lemaire, D.G.E. and Ruzsicska, B.P.: Thermospray high-performance liquid chromatographic mass spectral analyses of the photoproducts of dTpdT and dTpdU. *Biol. Mass Spectrom.*, 21 (1992) 259-266; *C.A.*, 117 (1992) 107336a.
- 1074 Chang, C., Doong, S., Zhou, J.H., Beach, J.W., Jeong, L.S., Chu, C.K., Tsai, C. and Cheng, Y.: Deoxycytidine deaminase-resistant stereoisomer is the active form of (14S)-2',3'-dideoxy-3'-thiacytidine in the inhibition of hepatitis B virus replication. *J. Biol. Chem.*, 267 (1992) 13938-13942.
- 1075 Chrzanowska, M., Szczepaniak, W. and Sloderbach, A.: Determination of 6-mercaptopurine in the presence of 6-thiouric acid in urine by ion exchange chromatography with a spectrophotometric detector. *Chem. Anal. (Warsaw)*, 36 (1991) 685-690; *C.A.*, 117 (1992) 144900a.
- 1076 Di Stefano, A., Pizzichini, M., Leoncini, R., Vannoni, D., Pagani, R. and Marinello, E.: Quantitative separation of uric acid and allantoin from rat liver tissue. *Biochim. Biophys. Acta*, 1117 (1992) 1-6.
- 1077 Fujise, H., Higashijima, M., Sakamoto, Y., Sonoki, S., Hisamatsu, S., Ogawa, E. and Takahashi, R.: HPLC determination of adenine compounds as fluorescent derivatives in canine red blood cells. *J. Vet. Med. Sci.*, 54 (1992) 379-381; *C.A.*, 117 (1992) 22640v.
- 1078 Kabulov, B.D., Shapirov, A.K., Ishanukulova, M.K. and Muminov, Ya.V.: (HPLC analysis of uracil, adenosin, and phosphaden). *Khim.-Farm. Zh.*, 26 (1992) 86-87; *C.A.*, 117 (1992) 142608n.
- 1079 Klein, S. and Oesch, F.: Assay for O⁶-alkylguanine-DNA-alkyltransferase using oligonucleotides containing O⁶-methylguanine in a BamHI recognition site as substrate. *Anal. Biochem.*, 205 (1992) 294-299.
- 1080 Martens, R.: A comparison of soil adenine nucleotide measurements by HPLC and enzymic analysis. *Soil Biol. Biochem.*, 24 (1992) 639-645; *C.A.*, 117 (1992) 150016h.
- 1081 Mitchell, E.P., Evans, L., Schultz, P., Madsen, R., Yarbrow, J.W., Gehrke, C.W. and Kuo, K.: Modified nucleosides in human serum. *J. Chromatogr.*, 581 (1992) 31-40.
- 1082 Nagaoka, H., Nohta, H., Saito, M. and Ohkura, Y.: High-performance liquid chromatographic determination of ribonucleosides and 2'-deoxyribonucleosides based on precolumn fluorescence derivatization of the sugar moieties. *Anal. Sci.*, 8 (1992) 345-349; *C.A.*, 117 (1992) 82693z.
- 1083 Nishio, H., Ono, A., Matsuda, A. and Ueda, T.: Nucleosides and nucleotides 111. Thermal stability of oligodeoxyribonucleotide duplexes containing N⁶-hydroxyadenine in substitution for adenine. *Chem. Pharm. Bull.*, 40 (1992) 1355-1357.
- 1084 Park, E.M., Shigenaga, M.K., Degan, P., Korn, T.S., Kitzler, J.W., Wehr, C.M., Kolachana, P. and Ames, B.N.: Assay of excised oxidative DNA lesions: isolation of 8-oxoguanine and its nucleoside derivatives from biological fluids with a monoclonal antibody column. *Proc. Natl. Acad. Sci. U.S.A.*, 89 (1992) 3375-3379; *C.A.*, 117 (1992) 42034d.
- 1085 Peltonen, K., Canella, K. and Dipple, A.: High-performance liquid chromatographic separation of purine deoxyribonucleoside monophosphate benzo[a]pyrene adducts. *J. Chromatogr.*, 623 (1992) 247-254.
- 1086 Pompon, A., Lefebvre, I. and Imbach, J.L.: "On-line internal surface reversed-phase cleaning": the direct HPLC analysis of crude biological samples. Application to the kinetics of degradation of oligonucleotides in cell culture medium. *Biochem. Pharmacol.*, 43 (1992) 1769-1775; *C.A.*, 117 (1992) 22637z.
- 1087 Porta, R., Moltrasio, D., Pescador, R., Lanzarotti, E., Mantovani, M. and Prino, G.: High-performance liquid chromatography determination of polydeoxyribonucleotides in plasma: its application to the determination of defibrotide's pharmacokinetics in the rabbit. *Anal. Biochem.*, 204 (1992) 143-146.
- 1088 Resines, J.A., Arín, M.J. and Díez, M.T.: Determination of creatinine and purine derivatives in ruminants' urine by reversed-phase high-performance liquid chromatography. *J. Chromatogr.*, 607 (1992) 199-202.
- 1089 Sant, M.E., Lyons, S.D., Phillips, L. and Christopherson, R.I.: Antifolates induce inhibition of amido phosphoribosyltransferase in leukemia cells. *J. Biol. Chem.*, 267 (1992) 11038-11045.
- 1090 Shewach, D.S.: Quantitation of deoxyribonucleoside 5'-triphosphates by a sequential boronate and anion-exchange high-pressure liquid chromatographic procedure. *Anal. Biochem.*, 206 (1992) 178-182.

- 1091 Singhall, R.P., Hughbanks, D. and Xian, J.: Separation of dideoxyribonucleosides in trace amounts by automated liquid chromatography and capillary electrophoresis. *J. Chromatogr.*, 609 (1992) 147-161.
- 1092 Takenaka, S., Dohtsu, K. and Takagi, M.: Development of a high-performance liquid chromatographic gel carrying intercalator-like benzoates for analysis of oligonucleotides. *Anal. Sci.*, 8 (1992) 151-156; *C.A.*, 117 (1992) 107516j.
- 1093 Tuck, M.T.: Partial purification of a 6-methyladenine mRNA methyltransferase which modifies internal adenine residues. *Biochem. J.*, 288 (1992) 233-240.
- 1094 Van Schepdael, A., van Aerschot, A., Herdewijn, P., Roets, E. and Hoogmartens, J.: Liquid chromatographic separation of the anomers and isomers of 2'-deoxyuridine. *Chromatographia*, 33 (1992) 571-574.
- 1095 Van Schepdael, A., van Aerschot, A., Herdewijn, P., Roets, E. and Hoogmartens, J.: Liquid chromatographic separation of the anomers and isomers of 2'-deoxyuridine. *Chromatographia*, 33 (1992) 571-574.
- 1096 Vtorushina, I.A., Gorbunov, Yu.A., Zarytova, V.F., Komarova, N.I., Lebedev, A.V., Lohkov, S.G., Manaenko, A.V., Sinyakov, A.N. and Shishkina, I.G.: (Diastereomers of nonionic oligonucleotide analogs. VI. Separation of diastereomers of ethyl phosphotriesters of octanucleotide d(GCCAAACA) by means of high performance complementary (affinity) chromatography). *Bioorg. Khim.*, 18 (1992) 92-99; *C.A.*, 117 (1992) 22611m.
- 1097 Xu, Y.-Z. and Swann, P.F.: Chromatographic separation of oligonucleotides with identical length: application to purification of oligomers containing a modified base. *Anal. Biochem.*, 204 (1992) 185-189.
- 1098 Zhou, H., You, G., Chen, X., Liu, S., Wang, B.: (Determination of hypoxanthine and adenine with monoamine oxidase). *Fenxi Huaxue*, 20 (1992) 413-415; *C.A.*, 117 (1992) 43742v.

For additional information see C.A.:

117 (1992) 22643y, 66056p, 107462p, 127544v.

See also 473, 486, 540, 816, 1155, 1448, 1468, 1477, 1483.

21b. Nucleic acids, RNA

- 1099 Bochkariov, D.E. and Kogon, A.A.: Application of 3-[3-(3-trifluoromethyl) diazirin-3-yl]phenyl]-2,3-dihydroxypropionic acid, carbene-generating, cleavable cross-linking reagent for photoaffinity labeling. *Anal. Biochem.*, 204 (1992) 90-95.
- 1100 Holmes, W.M., Andraos-Selim, C., Roberts, I. and Wahab, S.Z.: Structural requirements for tRNA methylation. Action of *Escherichia coli* tRNA(guanosine-1)methyltransferase on tRNA₁^{Leu} structural variants. *J. Biol. Chem.*, 267 (1992) 13440-13445.
- 1101 Jahn, D.: Expression of the *Chlamydomonas reinhardtii* chloroplast tRNA^{Glu} gene in a homologous *in vitro* transcription system is independent of upstream promoter elements. *Arch. Biochem. Biophys.*, 298 (1992) 505-513.
- 1102 Sproat, B.S. and Lamond, A.I.: 2'-O-Methyloligoribonucleotides: synthesis and applications. In: Eckstein, F. (Editor), *Oligonucleotides Analogues*, IRL, Oxford, 1991, pp. 49-86; *C.A.*, 117 (1992) 49017j - a review with 18 refs.

21c. Nucleic acids, DNA

- 1103 Hirabayashi, J. and Kasai, K.: (New intelligent separation method for large DNA molecules. Slalom chromatography). *Tanpakushitsu Kakusan Koso*, 37 (1992) 2173-2182; *C.A.*, 117 (1992) 166869z - a review with 22 refs.
- 1104 Kapp, U. and Langowski, J.: Preparation of DNA topoisomers by RP-18 high-performance liquid chromatography. *Anal. Biochem.*, 206 (1992) 293-299.
- 1105 Lai, J.S. and Herr, W.: Ethidium bromide provides a simple tool for identifying genuine DNA-independent protein associations. *Proc. Natl. Acad. Sci. U.S.A.*, 89 (1992) 6958-6962; *C.A.*, 117 (1992) 167091b.
- 1106 Talaska, G., Roh, J.H. and Getek, R.: ³²P-Postlabelling and mass spectrometric methods for analysis of bulky, polyaromatic carcinogen-DNA adducts in human. *J. Chromatogr.*, 580 (1992) 293-323 - a review with 144 refs.
- 1107 Térouanne, B., Balaguer, P., Boussioux, A.M. and Nicolas, J.C.: Quantitative and qualitative analysis of amplified DNA sequences by a competitive hybridization assay. *Anal. Biochem.*, 205 (1992) 193-199.
- 1108 Willis, T.C. and Goldberg, B.: A model system for DNA removal from protein solutions. *Am. Biotechnol. Lab.*, 10 (1992) 24; *C.A.*, 117 (1992) 166902e.

For additional information see C.A.:

117 (1992) 171971f.

See also 1566.

21f. Complex mixtures of nucleic acids and their fragments

See 1102.

22. ALKALOIDS

- 1109 Bruckert, E., Jacob, N., Lemaire, L., Truffert, J., Percheron, F. and de Gennes, J.L.: Relationship between smoking status and serum lipids in a hyperlipidemic population and analysis of possible confounding factors. *Clin. Chem. (Winston-Salem)*, 38 (1992) 1698-1705.
- 1110 Campíns-Falcó, P., Bosch-Reig, F., Herráez-Hernández, R. and Sevilano-Cabeza, A.: Determination of theophylline and paraxanthine in urine samples by liquid chromatography using the H-point standard additions method. *Anal. Chim. Acta*, 268 (1992) 73-80.
- 1111 Debal, V., Morjani, H., Millot, J.-M., Angiboust, J.-F., Gourdiere, B. and Manfait, M.: Determination of vinorelbine (Navelbine) in tumour cells by high-performance liquid chromatography. *J. Chromatogr.*, 581 (1992) 93-99.
- 1112 Dostal, J., Taborska, E. and Slavik, J.: Preparative column chromatography of quaternary benzophenanthridine alkaloids of *Dicranostigma lactucoides* *Fitoterapia*, 63 (1992) 67-69; *C.A.*, 117 (1992) 86800e.
- 1113 Drewes, F.E., van Staden, J. and Drewes, S.E.: Benzoylation and high performance liquid chromatographic detection of solasodine. *Phytochem. Anal.*, 3 (1992) 85-87; *C.A.*, 117 (1992) 22649e.

- 1114 Fitos, I., Visy, J., Simonyi, M. and Hermansson, J.: Chiral high-performance liquid chromatographic separations of vinca alkaloid analogues on α_1 -acid glycoprotein and human serum albumin columns. *J. Chromatogr.*, 609 (1992) 163-171.
- 1115 Golkiewicz, W., Kuczynski, J. and Jusiak, L.: (Comparison of separation selectivity of some alkaloids in the systems: aqueous methanolic mobile phase and unmodified or chemically modified silica gel). *Chem. Anal. (Warsaw)*, 36 (1991) 209-213; C.A., 117 (1992) 39529p.
- 1116 Larsen, A.K., Jr. and Tebbett, I.R.: Direct analysis for cocaine in urine by high-performance liquid chromatography (HPLC) using a column-switching technique. *J. Forensic Sci.*, 37 (1992) 636-639; C.A., 117 (1992) 144940p.
- 1117 Manceau, F., Fliniaux, M.A., Jacquin-Dubreuil, A.: A high performance liquid chromatographic procedure for the analysis of tobacco alkaloids. Application to the evaluation of tobacco alkaloids in plants and cell suspension cultures. *Phytochem. Anal.*, 3 (1992) 65-68; C.A., 117 (1992) 43758e.
- 1118 Moore, C., Browne, S., Tebbett, I. and Negrusz, A.: Determination of cocaine and its metabolites in brain tissue using high-flow solid-phase extraction columns and high-performance liquid chromatography. *Forensic Sci. Int.*, 53 (1992) 215-219; C.A., 117 (1992) 84764x.
- 1119 Nishikawa, M., Nakajima, K., Igarashi, K., Kasuya, F., Fukui, M. and Tsuchihashi, H.: (Determination of morphine 3-glucuronide in human urine by LC/APCI-MS). *Jpn. J. Toxicol. Environ. Health*, 38 (1992) 121-126; C.A., 117 (1992) 42076u.
- 1120 Petersen, D., Schumann, G., Oellerich, M., Klauke, R. and Wittner, N.: HPLC reference method for theophylline: calibration by use of standart addition. In: Sunshine, I. (Editor), *Recent Dev. Ther. Drug Monit. Clin. Toxicol.*, Dekker, New York, 1992, pp. 277-284; C.A., 117 (1992) 76602c.
- 1121 Puopolo, P.R., Chamberlin, P. and Flood, J.G.: Detection and confirmation of cocaine and cocaethylene in serum emergency toxicology specimens. *Clin. Chem. (Winston-Salem)*, 38 (1992) 1838-1842.
- 1122 Salvadori, P., Pini, D., Rosini, C., Bertucci, C. and Uccello-Barretta, G.: Chiral discriminations with Cinchona alkaloids. *Chirality*, 4 (1992) 42-49; C.A., 117 (1992) 56080c - a review with 27 refs.
- 1123 Yu, C., Hong, Y., Zhang, H. and Xu, X.: (DL-Camphor-10-sulfonic acid as an ion-pair reagent for determination of berberine in plasma by reversed-phase high-performance liquid chromatography). *Sepu*, 10 (1992) 167-168; C.A., 117 (1992) 62174c.

For additional information see C.A.:

117 (1992) 105805d, 107461n, 110041f, 110265g.

See also 1235, 1388, 1410, 1515, 1526.

23. OTHER SUBSTANCES CONTAINING HETEROCYCLIC NITROGEN

23a. Porphyrins and other pyrroles

- 1124 Sliviok, J., Kus, P. and Kozera, A.: Chromatographic characteristics of porphyrine ether glycosides. *Chromatographia*, 34 (1992) 431-432.

For additional information see C.A.:

117 (1992) 125978r.

See also 293.

23b. Bile pigments

- 1125 Ihara, H., Nakamura, H., Aoki, Y., Aoki, T. and Yoshida, M.: *In vitro* effects of light on serum bilirubin subfractions measured by high-performance liquid chromatography: comparison with four routine methods. *Clin. Chem. (Winston-Salem)*, 38 (1992) 2124-2129.

23c. Indole derivatives and plant hormones (gibberelins)

- 1126 Akhmedzhanova, S.B., Balabolkin, I.I., Vasil'eva, E.M. and Eller, K.I.: (Whole blood histamine and serotonin measurements by high-performance liquid chromatography). *Lab. Delo*, (1991) 10-13; C.A., 117 (1992) 83537p.
- 1127 Anderson, G.M., Hall, L.M., Yang, J.X. and Cohen, D.J.: Platelet dense granule release reaction monitored by high-performance liquid chromatography-fluorometric determination of endogenous serotonin. *Anal. Biochem.*, 206 (1992) 64-67.
- 1128 Caccamese, S., Benincori, T. and Principato, G.: Direct liquid chromatographic separation of the enantiomers of a 1,3-dihydro-phenylindol-2-one and of a 2,3-dihydro-2-oxo-benzofuran derivative. *J. Chromatogr.*, 623 (1992) 159-162.
- 1129 Hansen-Moller, J.: Determination of indolic compounds in pig back fat by solid-phase extraction and gradient high-performance liquid chromatography with special emphasis on the boar taint compound skatole. *J. Chromatogr.*, 624 (1992) 479-490.
- 1130 Liao, W. and Lu, X.: (Determination of 5-hydroxytryptamine, free and bound tryptophan in blood by high performance liquid chromatography). *Sepu*, 10 (1992) 118-119; C.A., 117 (1992) 65800q.
- 1131 Oestin, A., Moritz, T. and Sandberg, G.: Liquid chromatography/mass spectrometry of conjugates and oxidative metabolites of indole-3-acetic acid. *Biol. Mass Spectrom.*, 21 (1992) 292-298; C.A., 117 (1992) 65984c.
- 1132 Vieira, R., Míguez, J., Lema, M. and Aldegunde, M.: Pineal and plasma melatonin as determined by high-performance liquid chromatography with electrochemical detection. *Anal. Biochem.*, 205 (1992) 300-305.

For additional information see C.A.:

117 (1992) 88882g.

See also 293, 547, 592.

23d. Pyridine derivatives

See 596.

23e. Other N-heterocyclic compounds

- 1133 Banks, M.R., Cadogan, J.I.G., Dawson, I.M., Gosney, I., Grant, K.J., Gaur, S., Hodgson, P.K.G. and Stevenson, D.E.: Chromatographic resolution of racemic amines, carboxylic acids, and alcohols by a new homochiral reagent derived from endo-borneol. *Chromatographia*, 34 (1992) 48-50.

- 1134 Endo, Y., Kikuchi, T., Nakamura, M. and Shinoda, H.: Determination of histamine and polyamines in calcified tissues of mice: contribution of mast cells and histidine decarboxylase to the amount of histamine in the bone. *Calcif. Hisashi*, 51 (1992) 67-71; C.A., 117 (1992) 164000k.
- 1135 Mann, G., Titelbach-Helmrich, V., Lippmann, T. and Müller, K.: Separation of stereoisomers of tetramethylcalix[4]arenoctol and derivatives by high-performance liquid chromatography. *Chromatographia*, 34 (1992) 453-456.
- 1136 McLellan, A.C., Phillips, S.A. and Thornalley, P.J.: The assay of methylglyoxal in biological systems by derivatization with 1,2-diamino-4,5-dimethoxybenzene. *Anal. Biochem.*, 206 (1992) 17-23.
- 1137 Özden, S., Özden, T., Attila, I., Küçükislamoglu, M. and Okatan, A.: Isolation and identification via high-performance liquid chromatography and thin-layer chromatography of benzoxazolinone precursors from *Consolida orientalis* flowers. *J. Chromatogr.*, 609 (1992) 402-406.
- 1138 Puhlmann, A., Dülffer, T. and Kobold, U.: Multidimensional high-performance liquid chromatography on Pinkerton ISRP and RP18 columns: direct serum injection to quantify creatinine. *J. Chromatogr.*, 581 (1992) 129-133.
- 1139 Serrano-Carreón, L., Baudron, S., Bensoussan, M. and Belin, J.M.: A simple protocol for extraction and detection of pyrazines produced by fungi in solid state fermentation. *Biotechnol. Tech.*, 6 (1992) 19-22; C.A., 117 (1992) 46812d.
- 1140 Shibata, K. and Onodera, M.: Simultaneous high-performance liquid chromatographic measurement of xanthurenic acid and 3-hydroxyanthranilic acid in urine. *Biosci., Biotechnol., Biochem.*, 56 (1992) 974; C.A., 117 (1992) 86114j.
- 1141 Wiseman, R.W., Moerland, T.S., Chase, P.B., Stuppard, R. and Kushmerick, M.J.: High-performance liquid chromatographic assays for free and phosphorylated derivatives of the creatine analogues β -guanidopropionic acid and 1-carboxy-methyl-2-iminoimidazolidine (cyclocreatine). *Anal. Biochem.*, 204 (1992) 383-389.
- See also 86, 1088, 1126, 1359, 1528.
24. ORGANIC SULPHUR COMPOUNDS (INCL. GLUCOSINOLATES)
- 1142 Bundt, J., Herbel, W. and Steinhart, H.: Determination of polycyclic aromatic sulfur heterocycles (PASH) in diesel fuel by high performance liquid chromatography and photodiode-array detection. *J. High Resolut. Chromatogr.*, 15 (1992) 682-685.
- 1143 Ciucu, A. and Baldwin, R.P.: Determination of 2-thiothiazolidine-4-carboxylic acid in urine by liquid chromatography with electrochemical detection. *Electroanalysis (N.Y.)*, 4 (1992) 515-519; C.A., 117 (1992) 84735p.
- 1144 Koubek, E. and Stewart, A.E.: The analysis of sulfur in coal: an ion chromatography experiment. *J. Chem. Educ.*, 69 (1992) A146, A148; C.A., 117 (1992) 47501p.
- 1145 Quinsac, A., Ribaillier, D., Rollin, P. and Dreux, M.: Analysis of 5-vinyl-1,3-oxazolidine-2-thione by liquid chromatography. *J. AOAC Int.*, 75 (1992) 529-536; C.A., 117 (1992) 42063n.
- 1146 Sakurai, A., Horibe, H. and Okumura, Y.: Improved isolation method of urothion. *Rep. Fac. Sci., Shizuoka Univ.*, 26 (1992) 57-63; C.A., 117 (1992) 43755b.
- 1147 Smith, J.D., Ross, D., Graham, A.B. and Skellern, G.G.: Isolation and characterization of the S-glucuronide of 4-nitrothiophenol formed by microsomal glucuronyl transferase. *J. Pharm. Biomed. Anal.*, 10 (1992) 461-463.
- 1148 Zhang, Y., Cho, C.-G., Posner, G.H. and Talalay, P.: Spectroscopic quantitation of organic isothiocyanates by cyclocondensation with vicinal dithiols. *Anal. Biochem.*, 205 (1992) 100-107.
- 1149 Zheng, J. and Hanzlik, R.P.: Bromo(monohydroxy)phenyl mercapturic acids. A new class of mercapturic acids from bromobenzene-treated rats. *Drug Metab. Disp.*, 20 (1992) 688-694.
- 1150 Zhou, C., Bahr, A. and Schwedt, G.: The separation and determination of xanthates using high performance liquid chromatography. *Trans. Nonferrous Met. Soc. China*, 1 (1991) 89-95; C.A., 117 (1992) 183826a.
- See also 60, 418, 1478, 1520, 1605.
25. ORGANIC PHOSPHORUS COMPOUNDS (INCL. SUGAR PHOSPHATES)
- 1151 Fuchs, G., Doguet, L., Barth, D. and Perrut, M.: Enantiomer fractionation of phosphine oxides by preparative subcritical fluid chromatography. *J. Chromatogr.*, 623 (1992) 329-336.
- 1152 Kaszkin, M., Seidler, L., Kast, R. and Kinzel, V.: Epidermal-growth-factor-induced production of phosphatidylalcohols by HeLa cells and A431 cells through activation of phospholipase D. *Biochem. J.*, 287 (1992) 51-57.
- 1153 Kean, E.L.: Studies on the activation by dolichol-P-mannose of the biosynthesis of GlcNAc-P-P-dolichol and the topography of the GlcNAc-transferases concerned with the synthesis of GlcNAc-P-P-dolichol and (GlcNAc)₂-P-P-dolichol: a review. *Biochem. Cell Biol.*, 70 (1992) 413-421.
- 1154 Macphee, C.H.: Granulocyte/macrophage colony-stimulating factor affects myo-inositol metabolism in a novel manner. Implications for its primary action on human neutrophils. *Biochem. J.*, 286 (1992) 535-540.
- 1155 Min, Q., Zhang, Z. and Ruan, J.: (Simultaneous determination of creatine phosphate and adenine nucleotides in muscles by reversed-phase high performance liquid chromatography with automatic wavelength adjustment). *Sepu*, 10 (1992) 237-239; C.A., 117 (1992) 146452e.
- 1156 Shchipanova, I.N., Kharatyan, E.F., Ogrel', O.D., Sibel'dina, L.A. and Ostrovsky, D.N.: (On the nature of a new macroergic compound synthesized in bacteria in response to oxidative stress). *Biokhimiya (Moscow)*, 57 (1992) 862-872.
- 1157 Wong, N.S., Barker, C.J., Morris, A.J., Craxton, A., Kirk, C.J. and Michell, R.H.: The inositol phosphates in WRK1 rat mammary tumour cells. *Biochem. J.*, 286 (1992) 459-468.
- 1158 Woodcock, E.A. and Tanner, J.K.: Preparation of samples for high-performance liquid chromatography of inositol phosphates. *J. Chromatogr.*, 581 (1992) 134-138.
- 1159 Yatomi, Y., Ozaki, Y. and Kume, S.: Synthesis of phosphatidylinositol 3,4-bisphosphate but not phosphatidylinositol 3,4,5-trisphosphate is closely correlated with protein-tyrosine phosphorylation in thrombin-activated human platelets. *Biochem. Biophys. Res. Commun.*, 186 (1992) 1480-1486.
- For additional information see C.A.:
117 (1992) 169693y.

See also 136, 444, 451, 473, 483, 486, 548, 608, 657, 1052, 1067, 1141.

26. ORGANOMETALLIC AND RELATED COMPOUNDS

26a. Organometallic compounds

- 1160 Honore Hansen, S., Larsen, E.H., Pritzl, G. and Cornett, C.: Separation of seven arsenic compounds by high-performance liquid chromatography with on-line detection by hydrogen-argon flame atomic absorption spectrometry and inductively coupled plasma mass spectrometry. *J. Anal. At. Spectrom.*, 7 (1992) 629-634; C.A., 117 (1992) 123796f.
- 1161 Morin, P., Amran, M.B., Lakkis, M.D. and Leroy, M.J.F.: Ion-pair reversed-phase liquid chromatography of arsenic species on polymeric styrene-divinylbenzene packed columns with an alkaline aqueous mobile phase. *Chromatographia*, 33 (1992) 581-585.
- 1162 Uden, P.C. and Zeng, Y.: A study of diastereoisomerism of oxovanadium Schiff base chelates by gas chromatography-atomic emission spectroscopy, and high performance liquid chromatography. *Chromatographia*, 34 (1992) 269-275.

See also 1293, 1469, 1549.

26c. Coordination compounds

- 1163 Dolgonosov, A.M.: Simultaneous determination of transition metals and anions by ion chromatography. *React. Polym.*, 17 (1992) 95-99; C.A., 117 (1992) 39397u.
- 1164 Kim, I.W., Kang, S.H., Lim, J.H., Park, K.S. and Lee, W.: (A study on the elution behavior and separation of metal- α -diketone imine chelates by reversed-phase liquid chromatography). *Punsok Kwahak*, 2 (1989) 61-75; C.A., 117 (1992) 183879v.

See also 45, 1579.

27. VITAMINS AND VARIOUS ANIMAL GROWTH FACTORS (NON-PEPTIDIC)

- 1165 Aksnes, L.: A simplified high-performance liquid chromatographic method for determination of vitamin D₃, 25-hydroxyvitamin D₂ and 25-hydroxyvitamin D₃ in human serum. *Scand. J. Clin. Lab. Invest.*, 52 (1992) 177-182; C.A., 117 (1992) 127526r.
- 1166 Andersson, I. and Oeste, R.: Loss of ascorbic acid, folic acid and vitamin B₁₂, and changes in oxygen content of UHT milk. I. Introduction and methods. *Milchwissenschaft*, 47 (1992) 223-224; C.A., 117 (1992) 88839y.
- 1167 Bates, C.J.: Use of homocysteine to stabilise ascorbic acid, or to reduce dehydroascorbic acid, during HPLC separation of large volumes of tissue extracts. *Clin. Chim. Acta*, 205 (1992) 249-252.
- 1168 Bognar, A.: (Determination of vitamin D in food by HPLC. Results of collaborative studies of the working group "Vitamin Analysis" according to paragraph 35 LMBG (German Food Act)). *Z. Lebensm.-Unters. Forsch.*, 194 (1992) 469-475; C.A., 117 (1992) 130073w.

- 1169 Booth, C.K., Nixon, P.F. and Winzor, D.J.: Gel chromatographic evaluation of the binding constant for the interaction of thiamin diphosphate with magnesium ion. *J. Chromatogr.*, 609 (1992) 83-87.
- 1170 Bramley, P.M.: Analysis of carotenoids by high performance liquid chromatography and diode-array detection. *Phytochem. Anal.*, 3 (1992) 97-104; C.A., 117 (1992) 146363b - a review with 40 refs.
- 1171 Carcelain, G., David, F., Lepage, S., Bonnefont-Rousselot, D., Delattre, J., Legrand, A., Peynet, J. and Troupelet S.: Simple method for quantifying α -tocopherol in low-density + very-low-density lipoproteins and in high-density lipoproteins. *Clin. Chem. (Winston-Salem)*, 38 (1992) 1792-1795.
- 1172 Chase, G.W., Jr., Landen, W.O., Eitenmiller, R.R. and Soliman, A.-G.M.: Liquid chromatographic determination of thiamine, riboflavin, and pyridoxine in infant formula. *J. Assoc. Off. Anal. Chem.*, 75 (1992) 561-565.
- 1173 Chase, G.W., Jr., Landen, W.O., Jr., Eitenmiller, R.R. and Soliman, A.-G.M.: Liquid chromatographic determination of thiamine, riboflavin, and pyridoxine in infant formula. *J. AOAC Int.*, 75 (1992) 561-565; C.A., 117 (1992) 46861u.
- 1174 Chen, B.H. and Chen, Y.Y.: Determination of carotenoids and chlorophylls in water convolvulus (*Ipomoea aquatica*) by liquid chromatography. *Food Chem.*, 45 (1992) 129-134; C.A., 117 (1992) 110282k.
- 1175 Delgado Zamarreno, M.M., Sánchez Pérez, A., Gómez Pérez, C. and Hernández Méndez, J.: High-performance liquid chromatography with electrochemical detection for the simultaneous determination of vitamin A, D₃ and E in milk. *J. Chromatogr.*, 623 (1992) 69-74.
- 1176 Gratzfeld-Huesgen, A., Schuster, R. and Haecker, W.: Analysis of selected vitamins by HPLC and electrochemical detection. *Tec. Lab.*, 14 (1992) 277-283; C.A., 117 (1992) 178385b.
- 1177 Hart, J.P., Norman, M.D. and Lacey, C.J.: Voltametric behaviour of vitamins D₂ and D₃ at glassy carbon electrode and their determination in pharmaceutical products by using liquid chromatography with amperometric detection. *Analyst (London)*, 117 (1992) 1441-1445.
- 1178 Hoffman, K.M., Brown, P. and Maugle, P.D.: Comparison of three HPLC methods for the analysis of vitamins of ascorbic acid in shrimp tissue. *J. Liq. Chromatogr.*, 15 (1992) 2581-2610.
- 1179 Iwase, H.: Determination of folic acid in an elemental diet by high-performance liquid chromatography with UV detection. *J. Chromatogr.*, 609 (1992) 399-401.
- 1180 Khachik, F., Beecher, G.R., Goli, M.B., Lusby, W.R. and Smith, J.C., Jr.: Separation and identification of carotenoids and their oxidation products in the extracts of human plasma. *Anal. Chem.*, 64 (1992) 2111-2122.
- 1181 Klein, J., Fernandes, D., Pastuszak, A. and Koren, G.: Therapeutic drug monitoring of retinoids. *Ther. Drug Monit.*, 14 (1992) 197-202; C.A., 117 (1992) 103448r.
- 1182 Lambert, D., Adjalla, C., Felden, F., Benhayoun, S., Nicolas, J.P. and Guéant, J.L.: Identification of vitamin B₁₂ and analogues by high-performance capillary electrophoresis and comparison with high-performance liquid chromatography. *J. Chromatogr.*, 608 (1992) 311-315.
- 1183 Lambert, W.E. and de Leenheer, A.: Quantitative determination of water-soluble vitamins using HPLC. *Food Sci. Technol.*, 52(Food Anal. HPLC) (1992) 341-369; C.A., 117 (1992) 110175c - a review with 107 refs.

- 1184 Lambert, W.E., Vanneste, L. and de Leenheer, A.P.: Enzymatic sample hydrolysis and HPLC in a study of phylloquinone concentration in human milk. *Clin. Chem. (Winston-Salem)*, 38 (1992) 1743-1748.
- 1185 Lee, B.L., Chua, S.C., Ong, H.Y. and Ong, C.N.: High-performance liquid chromatographic method for routine determination of vitamins A and E and β -carotene in plasma. *J. Chromatogr.*, 581 (1992) 41-47.
- 1186 Leyland, D.M., Evershed, R.P., Edwards, R.H.T. and Beynon, R.J.: Application of gas chromatography-mass spectrometry with selected ion monitoring to the urinalysis of 4-pyridoxic acid. *J. Chromatogr.*, 581 (1992) 179-185.
- 1187 Masuda, S., Okano, T. and Kobayashi, T.: A method for the simultaneous determination of vitamins D₂, D₃ and their metabolites in plasma and its application to plasma samples obtained from normal subjects and patients. *Food Chem.*, 45 (1992) 215-225; C.A., 117 (1992) 146456j.
- 1188 McPartlin, J., Courtney, G., McNulty, H., Weir, D. and Scott, J.: The quantitative analysis of endogenous folate catabolites in human urine. *Anal. Biochem.*, 206 (1992) 256-261.
- 1189 Muindi, J.R.F., Frankel, S.R., Husefson, C., DeGrazia, F., Garland, W.A., Young, C.W. and Warrell, R.P., Jr.: Clinical pharmacology of oral all-trans retinoic acid in patients with acute promyelocytic leukemia. *Cancer Res.*, 52 (1992) 2138-2142.
- 1190 Nierenberg, D.W. and Nann, S.L.: A method for determining concentrations of retinol, tocopherol, and five carotenoids in human plasma and tissue samples. *Am. J. Clin. Nutr.*, 56 (1992) 417-426; C.A., 117 (1992) 146451d.
- 1191 Nisperos-Carriedo, M.O., Buslig, B.S. and Shaw, P.E.: Simultaneous detection of dehydroascorbic, ascorbic and some organic acids in fruits and vegetables by HPLC. *J. Agric. Food Chem.*, 40 (1992) 1127-1130.
- 1192 Olmedilla, B., Granado, F., Blanco, I. and Rojas-Hidalgo, E.: Determination of nine carotenoids, retinol, retinyl palmitate and α -tocopherol in control human serum using two internal standards. *Food Chem.*, 45 (1992) 205-213; C.A., 117 (1992) 146455h.
- 1193 Pourfarzam, M. and Bartlett, K.: Intermediates of peroxisomal β -oxidation of [U-¹⁴C]hexadecanedioneate. A study of the acyl-CoA esters which accumulate during peroxisomal β -oxidation of [U-¹⁴C]hexadecanedionyl-mono-CoA. *Eur. J. Biochem.*, 208 (1992) 301-307.
- 1194 Rippin, J.J.: Analysis for fully oxidized neopterin in serum by high-performance liquid chromatography. *Clin. Chem. (Winston-Salem)*, 38 (1992) 1722-1724.
- 1195 Rizzolo, A. and Polesello, S.: Chromatographic determination of vitamins in foods. *J. Chromatogr.*, 624 (1992) 103-152 - a review with 286 refs.
- 1196 Rodewald, J.M., Moran, J.W., Donoho, A.L. and Coleman, M.R.: Determination of monensin in raw material, premix, and animal feeds by liquid chromatography with correlation to microbiological assay. *J. AOAC Int.*, 75 (1992) 272-279; C.A., 117 (1992) 88901.
- 1197 Ruotolo, G., Zhang, H., Bentsianov, V. and Lee, N.-A.: Protocol for the study of the metabolism of retinyl esters in plasma lipoproteins during postprandial lipemia. *J. Lipid Res.*, 33 (1992) 1541-1549.
- 1198 Russell, L.F. and Vanderslice, J.T.: Nondegradative extraction and simultaneous quantitation of riboflavin, flavin mononucleotide, and flavin adenine dinucleotide in foods by HPLC. *Food Chem.*, 43 (1992) 151-162; C.A., 117 (1992) 24862t.
- 1199 Saupe, J. and Shearer, M.J.: (Method for the determination of vitamin K₁ (phylloquinone) and possibilities for measuring vitamin K₂ (menaquinone)). *Klin. Labor.*, 38 (1992) 47-50; C.A., 117 (1992) 65780h.
- 1200 Seiffert, B., Swaczyna, H. and Schaefer, I.: (Simultaneous determination of L-ascorbic and D-isoascorbic acids in wine and beer by HPLC). *Dtsch. Lebensm.-Rundsch.*, 88 (1992) 38-40; C.A., 117 (1992) 169666s.
- 1201 Shimizu, M., Gao, Y., Aso, T., Nakatsu, K. and Yamada, S.: Fluorometric assay of 25-hydroxyvitamin D₃ and 24R,25-dihydroxyvitamin D₃ in plasma. *Anal. Biochem.*, 204 (1992) 258-264.
- 1202 Shimoda, M.: Simultaneous determination of tetrahydrofolate and N⁵-methyltetrahydrofolate in pig plasma by high-performance liquid chromatography with electrochemical detection. *J. Vet. Med. Sci.*, 54 (1992) 249-253; C.A., 117 (1992) 22639b.
- 1203 Sierra, C., Pastor, M.C. and de Ramón, M.: Liquid chromatography determination of α -tocopherol in erythrocytes. *Clin. Chim. Acta*, 208 (1992) 119-126.
- 1204 Skurikhin, V.N.: (Determination of vitamins A and E in biological samples by reversed-phase micro-column high-performance liquid chromatography). *Bull. Vses. Nauchno-Issled. Inst. Fiziol., Biokhim. Pitan. S-kh. Zhivotn.*, (1991) 79-81; C.A., 117 (1992) 146453f.
- 1205 Sliva, M.G., Green, A.E., Sanders, J.K., Euber, J.R. and Saucerman, J.R.: Reversed-phase liquid chromatographic determination of vitamin D in infant formulas and enteral nutritional. *J. AOAC Int.*, 75 (1992) 566-571; C.A., 117 (1992) 46862v.
- 1206 Sugimoto, T., Ogiwara, S., Teradaira, R., Fujita, K. and Nagatsu, T.: Determination of oncopterin isolated from urine of cancer patients as N²-(β -aminopropyl)biopterin. *Biog. Amines*, 9 (1992) 77-82; C.A., 117 (1992) 146448h.
- 1207 Tanusi, S., Matsushita, J. and Yamashita, K.: (Quantitative analysis by high-performance liquid chromatography of total vitamin in food after reduction with dithiothreitol). *Kenkyu Kiyo - Tokushima Bunri Daigaku*, 43 (1992) 221-227; C.A., 117 (1992) 130067x.
- 1208 Wouters, W., van Dun, J., Dillen, A., Coene, M.-C., Cools, W. and de Coster, R.: Effects of liarozole, a new antitumoral compound, on retinoic acid-induced inhibition of cell growth and on retinoic acid metabolism in MCF-7 human breast cancer cells. *Cancer Res.*, 52 (1992) 2841-2846.
- 1209 Ye, W., Liu, L. and Li, H.: (Determination of the fat-soluble vitamins and β -carotene in milk by high performance liquid chromatography). *Sepu*, 10 (1992) 240-241; C.A., 117 (1992) 130067d.

For additional information see C.A.:

117 (1992) 22756m, 33816h, 86341f, 88881f, 110271f, 127801t.

See also 406, 1308.

28. ANTIBIOTICS

- 1210 Aldana, I., Fos, D., Gonzales Penas, E., Gazzaniga, A., Gianesello, V., Ceppi Monti, N., Figini, P.G., Zato, M.A., Bruseghini, L. and Esteras, A.: Ocular pharmacokinetics of thiamphenicol in rabbits. *Arzneim.-Forsch.*, 42 (1992) 1236-1239.
- 1211 Backes, B.J. and Rychnovsky, S.D.: A reverse-phase HPLC assay for measuring the interaction of polyene macrolide antifungal agents with sterols. *Anal. Biochem.*, 205 (1992) 96-99.
- 1212 Bannister, B. and Zapotocky, B.A.: Protorubradirin, an antibiotic containing a C-nitroso-sugar fragment, is the true secondary metabolite produced by *Streptomyces achromogenes* var. *rubradiris*. Rubradirin, described earlier, is its photooxidation product. *J. Antibiot.*, 45 (1992) 1313-1324.
- 1213 Bano, M.C., Braco, L. and Abad, C.: A semi-empirical approach for the simulation of circular dichroism spectra of gramicidin A in a model membrane. *Biophys. J.*, 63 (1992) 70-77; *C.A.*, 117 (1992) 126785n.
- 1214 Barker, S.A. and Walker, C.C.: Chromatographic methods for tetracycline analysis in foods. *J. Chromatogr.*, 624 (1992) 195-209 - a review with 81 refs.
- 1215 Bobbitt, D.R. and Ng, K.W.: Chromatographic analysis of antibiotic materials in food. *J. Chromatogr.*, 624 (1992) 153-170 - a review with 53 refs.
- 1216 Boison, J.O.: Chromatographic methods of analysis for penicillins in food-animal tissues and their significance in regulatory programs for residue reduction and avoidance. *J. Chromatogr.*, 624 (1992) 171-194 - a review with 127 refs.
- 1217 Bryan, P.D., Hawkins, K.R., Stewart, J.T. and Capomacchia, A.C.: Analysis of chlortetracycline by high performance liquid chromatography with postcolumn alkaline-induced fluorescence detection. *Biomed. Chromatogr.*, 6 (1992) 305-310.
- 1218 Fomichova, E.V., Fedorova, G.B., Potapova, N.P. and Katrukha, G.S.: Isolation and identification of new anthracycline antibiotics, rubomycins F and H. *J. Antibiot.*, 45 (1992) 1185-1186.
- 1219 Georgopoulos, A., Czejka, M.J., Jaeger, W., Dieterich, H.A. and Fell, J.J.: (HPLC analysis of teicoplanin in plasma: a comparison with a microbiological method). *Fortschr. Antimikrob. Antineoplast. Chemother.*, 10-12 (Teicoplanin I) (1991) 165-170; *C.A.*, 117 (1992) 141f.
- 1220 Hakoda, S., Tsubotani, S., Iwasa, T., Suzuki, M., Kondo, M. and Harada, S.: Production and biological activities of a new antifungal antibiotic, TAN-950A. *J. Antibiot.*, 45 (1992) 854-860.
- 1221 Hedtmann, U., Fehlhaber, H.-W., Sukatsch, D.A., Weber, M., Hoffmann, D. and Kraemer, H.P.: The new cytotoxic antibiotic cytorhodin X, an unusual anthracyclinone-9 α -glycoside. *J. Antibiot.*, 45 (1992) 1373-1375.
- 1222 Hoff, H., Drautz, H., Fiedler, H.-P., Zähler, H., Schultz, J.E., Keller-Schierlein, W., Philipps, S., Ritzau, M. and Zeeck, A.: Metabolic products of microorganisms. 261. Obscurolides, a novel class of phosphodiesterase inhibitors from *Streptomyces*. I. Production, isolation, structural elucidation and biological activity of obscurolides A₁ to A₄. *J. Antibiot.*, 45 (1992) 1096-1108.
- 1223 Hofstead, S.J., Matson, J.A., Malacko, A.R. and Marquardt, H.: Kedarcidin, a new chromoprotein antitumor antibiotic. II. Isolation, purification and physico-chemical properties. *J. Antibiot.*, 45 (1992) 1250-1254.
- 1224 Hsu, M.-C., Chang, Y.-W. and Lee, Y.-T.: Column liquid chromatography and microbiological assay compared for determination of cefadroxil preparations. *J. Chromatogr.*, 609 (1992) 181-186.
- 1225 Hsu, M.C. and Hsu, P.W.: High-performance liquid chromatographic method for potency determination of amoxicillin in commercial preparations and for stability studies. *Antimicrob. Agents Chemother.*, 36 (1992) 1276-1279; *C.A.*, 117 (1992) 56099r.
- 1226 Ikai, Y., Oka, H., Hayakawa, J., Harada, K.-i. and Suzuki, M.: Structural characterization of bacitracin components by frit-fast atom bombardment (FAB) liquid chromatography/mass spectrometry (LC/MS). *J. Antibiot.*, 45 (1992) 1325-1334.
- 1227 Itoh, J. and Miyadoh, S.: SF2457, A new antibiotic related to ampicillin. *J. Antibiot.*, 45 (1992) 846-853.
- 1228 Iwaki, K., Okumura, N. and Yamazaki, M.: Determination of tetracycline antibiotics by reversed-phase high-performance liquid chromatography with fluorescence detection. *J. Chromatogr.*, 623 (1992) 153-158.
- 1229 Kádár-Pauncz, J., Podányi, B. and Horváth, H.: Isolation and structure elucidation of new antibiotics related to angolamycin. *J. Antibiot.*, 45 (1992) 1231-1238.
- 1230 Kawada, S.-Z., Yamashita, Y., Uosaki, Y., Gomi, K., Iwasaki, T., Takiguchi, T. and Nakano, H.: UCT4B, a new antitumor antibiotic with topoisomerase II mediated DNA cleavage activity, from *Streptomyces* sp. *J. Antibiot.*, 45 (1992) 1182-1184.
- 1231 Keukens, H.J., Aerts, M.M.L., Traag, W.A., Nouws, J.F.M., de Ruig, W.G., Beek, W.M.J. and den Hartog, J.M.P.: Analytical strategy for the regulatory control of residues of chloramphenicol in meat: preliminary studies in milk. *J. AOAC Int.*, 75 (1992) 245-256; *C.A.*, 117 (1992) 88897r.
- 1232 Komagata, D., Sawa, T., Muraoka, Y., Imada, C., Okami, Y. and Takeuchi, T.: TA-3037A, a new inhibitor of glutathione S-transferase, produced by actinomycetes. I. Production, isolation, physico-chemical properties and biological activities. *J. Antibiot.*, 45 (1992) 1117-1121.
- 1233 Krugers Dagneaux, P.G.L.C., Loohuis, C.P.G.G., Klein Elhorst, J.T. and van der Veer, T.S.: Liquid chromatographic estimation of tranlycypromine in human plasma. *Pharm. Weekbl., Sci. Ed.*, 14 (1992) 46-49; *C.A.*, 117 (1992) 39691k.
- 1234 Lai, F. and Sheehan, T.: Enhancement of detection sensitivity and cleanup selectivity for tobramycin through pre-column derivatization. *J. Chromatogr.*, 609 (1992) 173-179.
- 1235 Li, Y., Kobayashi, H., Hashimoto, Y. and Iwasaki, S.: Binding selectivity of rhizoxin, phomopsis A, vinblastine, and ansamitocin P-3 to fungal tubulins: differential interactions of these antimetabolic agents with brain and fungal tubulins. *Biochem. Biophys. Res. Commun.*, 187 (1992) 722-729.
- 1236 Matsumoto, T. and Komiya, I.: (Assay method of ME1207, a novel oral cephem antibiotic, in biological body fluids. II. High-performance liquid chromatographic assay method). *Chemotherapy (Tokyo)*, 40(Suppl. 2) (1992) 101-104; *C.A.*, 117 (1992) 19783p.
- 1237 Mehta, A.C., Hart-Davies, S. and Kay, E.A.: High-performance liquid chromatographic determination of ciprofloxacin in plasma. *J. Clin. Pharm. Ther.*, 17 (1992) 117-120; *C.A.*, 117 (1992) 19778r.

- 1238 Miyata, S., Ohhata, N., Murai, H., Masui, Y., Ezaki, M., Takase, S., Nishikawa, M., Kiyoto, S., Okuhara, M. and Kohsaka, M.: WS009A and B, new endothelin receptor antagonists isolated from *Streptomyces* sp. No. 89009. I. Taxonomy, fermentation, isolation, physico-chemical properties and biological activities. *J. Antibiot.*, 45 (1992) 1029-1040.
- 1239 Moats, W.A.: Determination of cloxacillin and penicillin V in milk using an automated liquid chromatography cleanup. *J. AOAC Int.*, 75 (1992) 257-260; C.A., 117 (1992) 88898s.
- 1240 Morishita, A., Murofushi, S., Ishizawa, K., Mutoh, N. and Yaginuma, S.: Isolation and characterization of sporeamicin C. *J. Antibiot.*, 45 (1992) 1011-1015.
- 1241 Naganuma, S., Sakai, K., Hasumi, K. and Endo, A.: Acaterin, a novel inhibitor of acyl-CoA:cholesterol acyltransferase produced by *Pseudomans* sp. A92. *J. Antibiot.*, 45 (1992) 1216-1221.
- 1242 Naidong, W., Thurani, J., Vermeulen, K., Roets, E. and Hoogmartens, J.: Quantitative analysis of minocycline by liquid chromatography on poly(styrene-divinylbenzene). *J. Liq. Chromatogr.*, 15 (1992) 2529-2549.
- 1243 Naidong, W., Vermeulen, K., Quintens, I., Roets, E. and Hoogmartens, J.: Evaluation of analytical methods. Analysis of minocycline by liquid chromatography. *Chromatographia*, 33 (1992) 560-566.
- 1244 Nakagawa, Y., Itai, S., Yoshida, T. and Nagai, T.: Physicochemical properties and stability in the acidic solution of a new macrolide antibiotic, clarithromycin, in comparison with erythromycin. *Chem. Pharm. Bull.*, 40 (1992) 725-728.
- 1245 Neely, F.L.: Determination of tylosin and related macrolides in fermentation broth by gradient elution ion-pair liquid chromatography. *Chromatographia*, 34 (1992) 51-55.
- 1246 Neely, F.L.: Determination of tylosin and related macrolides in fermentation broth by gradient elution ion-pair liquid chromatography. *Chromatographia*, 34 (1992) 51-55.
- 1247 Nelis, H.J., Vandenbranden, J., Verhaeghe, B., de Kruif, A., Mattheeuws, D. and de Leenheer, A.P.: Liquid chromatographic determination of ampicillin in bovine and dog plasma by using a tandem solid-phase extraction method. *Antimicrob. Agents Chemother.*, 36 (1992) 1606-1610; C.A., 117 (1992) 123930v.
- 1248 Ogawara, H., Higashi, K., Manita, S., Tanaka, K., Shimizu, Y. and Shufang, L.: An inhibitor for inositol-specific phospholipase C from *Actinomadura* sp. *J. Antibiot.*, 45 (1992) 1365-1366.
- 1249 Ohkuma, H., Naruse, N., Nishiyama, Y., Tsuno, T., Hoshino, Y., Sawada, Y., Konishi, M. and Oki, T.: Sultricin, a new antifungal and antitumor antibiotic from *Streptomyces roseiscleroticus*. Production, isolation, structure and biological activity. *J. Antibiot.*, 45 (1992) 1239-1249.
- 1250 Ohta, T., Hashimoto, E. and Hasegawa, M.: Cloning and analysis of a gene (*SMS13*) encoding sannamycin B-glycyltransferase from *Streptomyces sannanensis* and its distribution among actinomycetes. *J. Antibiot.*, 45 (1992) 1167-1175.
- 1251 Shan-Yang Lin, Wen-Hsiu Wu and Wing-Yiu Lui: *In vitro* release, pharmacokinetic and tissue distribution studies of doxorubicin hydrochloride (Adriamycin HCL®) encapsulated in lipidized w/o emulsions and w/o/w multiple emulsions. *Pharmazie*, 47 (1992) 439-443.
- 1252 Shoji, J., Hino, H., Katayama, T., Matsumoto, K., Tanimoto, T., Hattori, T., Higashiyama, I., Miwa, H., Motokawa, K. and Yoshida, T.: Isolation and characterization of new peptide antibiotics, plusbacins A₁-A₄ and B₁-B₄. *J. Antibiot.*, 45 (1992) 817-823.
- 1253 Shoji, J., Hino, H., Katayama, T., Nakagawa, J., Ikenishi, Y., Iwatani, K. and Yoshida, T.: Structures of new peptide antibiotics, plusbacins A₁-A₄ and B₁-B₄. *J. Antibiot.*, 45 (1992) 824-831.
- 1254 Tawara, S., Matsumoto, S., Matsumoto, Y., Kamimura, T. and Goto, S.: Structure-binding relationship and binding sites of cephalosporins in human serum albumin. *J. Antibiot.*, 45 (1992) 1346-1357.
- 1255 Toki, S., Ando, K., Kawamoto, I., Sano, H., Yoshida, M. and Matsuda, Y.: ES-242-2, -3, -4, -5, -6, -7, and -8, novel bioxanthracenes produced by *Verticillium* sp., which act on the N-methyl-D-aspartate receptor. *J. Antibiot.*, 45 (1992) 1047-1054.
- 1256 Tomoda, H., Nishida, H., Huang, X.-H., Masuma, R., Kim, Y.K. and Omura, S.: New cyclodepsipeptides, enniatins D, E. and F produced by *Fusarium* sp. FO-1305. *J. Antibiot.*, 45 (1992) 1207-1215.
- 1257 Tsukamoto, N., Fujii, I., Ebizuka, Y. and Sankawa, U.: Cloning of aklavinone biosynthesis genes from *Streptomyces galilaeus*. *J. Antibiot.*, 45 (1992) 1286-1294.
- 1258 Yoshimoto, A., Johdo, O., Fujii, S., Kubo, K., Nishida, H., Okamoto, R. and Takeuchi, T.: Anthracycline metabolites from baumycin-producing *Streptomyces* sp. D788. I. Isolation of antibiotic-blocked mutants and their characterization. *J. Antibiot.*, 45 (1992) 1255-1272.

For additional information see C.A.:

117 (1992) 19761e, 46681k, 46721y, 76593a, 149604s, 169582m, 178400c, 178404g.

See also 296, 1457, 1461, 1499, 1560.

29. INSECTICIDES, PESTICIDES AND OTHER AGROCHEMICALS

29a. General techniques

- 1259 Bushway, R.J.: Analysis of pesticide residues in food by HPLC. *Food Sci. Technol.*, 52(Food Anal. HPLC) (1992) 507-577; C.A., 117 (1992) 110180a - a review with 242 refs.
- 1260 Committee for analytical methods: Report of two co-operative trials of gel permeation chromatographic method for the isolation of pesticide residues from oils and fats. *Analyst (London)*, 117 (1992) 1451-1455.
- 1261 Fodor-Csorba, K.: Chromatographic methods for the determination of pesticides in foods. *J. Chromatogr.*, 624 (1992) 353-367 - a review with 62 refs.
- 1262 Heikes, D.L. and Craun, J.C.: Rapid multiresidue procedure for the determination of pesticides in anhydrous lanolin and lanolin-containing pharmaceutical preparations utilizing gel permeation chromatography cleanup with gas chromatographic and mass spectrometric techniques. *J. Agric. Food Chem.*, 40 (1992) 1586-1590; C.A., 117 (1992) 137764f.
- 1263 Hogendoorn, E.A., Verschraagen, C., Brinkman, U.A.T. and van Zoonen, P.: Coupled column liquid chromatography for the trace determination of polar pesticides in water using direct large-volume injection: method development strategy applied to methyl isothiocyanate. *Anal. Chim. Acta*, 268 (1992) 205-215.

- 1264 Kozlova, E.Yu. and Kobrina, V.N.: (Determination of pesticide mixtures used in vegetable cultivation). *Sib. Khim. Zh.*, (1992) 29-36; C.A., 117 (1992) 68571b.
- 1265 Reupert, R., Brausen, G. and Ploeger, E.: (Analytical method for the determination of pesticides by micro-HPLC). *GIT Spez. Chromatogr.*, 12 (1992) 30-36; C.A., 117 (1992) 118036d.
- 1266 Tonogai, Y., Tsumura, Y., Nakamura, Y. and Ito, Y.: (Studies on pesticides for post-harvest application. III. Analysis of post-harvest application pesticides in citrus fruits). *Shokuhin Eiseigaku Zasshi*, 33 (1992) 23-30; C.A., 117 (1992) 130034j.

For additional information see C.A.:
117 (1992) 46832k.

See also 1549, 1551.

29b. Chlorinated insecticides

- 1267 Singh, D.K. and Agarwal, H.C.: Chemical release and nature of soil-bound DDT residues. *J. Agric. Food Chem.*, 40 (1992) 1713-1716.
- 1268 Wells, D.E. and Echarri, I.: Determination of individual chlorobiphenyls (CBs), including non-ortho, and mono-ortho chloro substituted CBs in marine mammals from Scottish waters. *Int. J. Environ. Anal. Chem.*, 47 (1992) 75-97; C.A., 117 (1992) 84738s.

See also 546, 1278, 1279.

29c. Phosphorus insecticides

- 1269 Hanks, A.R.: Liquid chromatographic method for determination of methamidophos in technical products and pesticide formulations: CIPAC collaborative study. *J. Assoc. Off. Anal. Chem.*, 75 (1992) 698-700.
- 1270 Kwakman, P.J.M., Vreuls, J.J., Brinkman, U.A.T. and Ghijsen, R.T.: Determination of organophosphorus pesticides in aqueous samples by on-line membrane disk extraction and capillary gas chromatography. *Chromatographia*, 34 (1992) 41-47.
- 1271 Tieffova, P. and Matousek, M.: (A method for routine determination of ethephon residues in plant material). *Agrochimia (Bratislava)*, 31 (1991) 251-254; C.A., 116 (1992) 146022u.

29d. Carbamates

- 1272 De Kok, A., Hiemstra, M. and Brinkman, U.A.T.: Low ng/l-level determination of twenty N-methylcarbamate pesticides and twelve of their polar metabolites in surface water via off-line solid-phase extraction and high-performance liquid chromatography with post-column reaction and fluorescence detection. *J. Chromatogr.*, 623 (1992) 265-276.
- 1273 De la Colina, C., Pena-Heras, A. and Sánchez-Rasero, F.: Carbosulfan in technical concentrates and formulated products. Liquid chromatographic determination with photodiode-array detection. *J. Chromatogr.*, 607 (1992) 203-206.
- 1274 Pleasance, S., Anacleto, J.F., Bailey, M.R. and North, D.H.: An evaluation of atmospheric pressure ionization techniques for the analysis of N-methyl carbamate pesticides by liquid chromatography mass spectrometry. *J. Am. Soc. Mass Spectrom.*, 3 (1992) 378-397; C.A., 117 (1992) 64706b.

See also 1288.

29e. Herbicides

- 1275 Bagheri, H., Vreuls, J.J., Ghijsen, R.T. and Brinkman, U.A.T.: Determination of triazine herbicides in surface and drinking waters by off-line combination of liquid chromatography and gas chromatography-mass spectrometry. *Chromatographia*, 34 (1992) 5-13.
- 1276 Bagheri, H., Vreuls, J.J., Ghijsen, R.T. and Brinkman, U.A.T.: Determination of triazine herbicides in surface and drinking waters by off-line combination of liquid chromatography and gas chromatography-mass spectrometry. *Chromatographia*, 34 (1992) 5-13.
- 1277 Durand, G., Bouvot, V. and Barceló, D.: Determination of trace levels of herbicides in estuarine waters by gas and liquid chromatographic techniques. *J. Chromatogr.*, 607 (1992) 319-327.
- 1278 Hopper, M.L., McMahon, B., Griffitt, K.R., Cline, K., Fleming-Jones, M.E. and Kendall, D.C.: Analysis of fatty and nonfat foods for chlorophenoxy alkyl acids and pentachlorophenol. *J. Assoc. Off. Anal. Chem.*, 75 (1992) 707-713.
- 1279 Klöppel, H., Haider, J., Hoffmann, C. and Lüttecke, B.: Simultaneous determination of the herbicides isoproturon, dichlorprop and bifenoxy in soil using RP-HPLC. *Fresenius J. Anal. Chem.*, 344 (1992) 42-46.
- 1280 Ludwig, P., Gunkel, W. and Huehnerfuss, H.: Chromatographic separation of the enantiomers of marine pollutants. Part 5: enantioselective degradation of phenoxy-carboxylic acid herbicides by marine microorganisms. *Chemosphere*, 24 (1992) 1423-1429; C.A., 117 (1992) 64530q.
- 1281 Macomber, C., Bushway, R.J., Perkins, L.B., Baker, D., Fan, T.S. and Ferguson, B.S.: Determination of the ethanesulfonate metabolite of alachlor in water by high-performance liquid chromatography. *J. Agric. Food Chem.*, 40 (1992) 1450-1452.
- 1282 Miles, C.J. and Leong, G.: Glyphosate herbicide analysis by liquid chromatography with postcolumn reaction detection: column comparison and application notes. *LC-GC*, 10 (1992) 452-458; C.A., 117 (1992) 68593k.
- 1283 Suzuki, T. and Watanabe, S.: Liquid chromatographic screening method for fluorescent derivatives of chlorophenoxy acid herbicides in water. *J. Assoc. Off. Anal. Chem.*, 75 (1992) 720-724.
- 1284 Wigfield, Y.Y. and Lanouette, M.: A modified clean-up for the determination of glyphosate and its metabolite residues in lentils using high pressure liquid chromatography and post-column fluorogenic labelling. *Pestic. Sci.*, 33 (1991) 491-498; C.A., 116 (1992) 82331j.
- 1285 Zanco, M., Pfister, G. and Kettrup, A.: A new HPLC method for simultaneous determination of fluzifop-butyl and fluzifop in soil samples. *Fresenius J. Anal. Chem.*, 344 (1992) 39-41.

See also 214, 1271.

29f. Fungicides

- 1286 Koppen, B.: Liquid chromatographic determination of 2-hydroxy-3-aminophenazine and 2,3-diaminophenazine as impurities in pesticide formulations containing benomyl or carbendazim. *J. Chromatogr.*, 623 (1992) 168-173.

- 1287 Noguchi, A.: (Determination of thiabendazole in feed by high performance liquid chromatography). *Shiryō Kenkyū Hokoku (Tokyo Hishiryō Kensasho)*, 17 (1992) 42-49; C.A., 117 (1992) 130055s.
- 1288 Singh, R.P., Marvin, C.H., Brindle, I.D., Hall, C.D. and Chiba, M.: Stabilization of methyl [1-(butylcarbamoyl)-1H-benzimidazol-2-yl] carbamate (benomyl) in hydrochloric acid solutions. *J. Agric. Food Chem.*, 40 (1992) 1303-1306.
- 1289 Tonogai, Y., Tsumura, Y., Nakamura, Y., Matsuki, H. and Ito, Y.: (Studies on simultaneous determination of 12 kinds of fungicides containing unstable captan, captafol, etc. in preparation of test solution). *Jpn. J. Toxicol. Environ. Health*, 38 (1992) 270-281; C.A., 117 (1992) 130043m.
- 29g. *Other types of pesticides and various agrochemicals*
- 1290 Katagi, T.: Photodegradation of 3-phenoxybenzoic acid in water and on solid surfaces. *J. Agric. Food Chem.*, 40 (1992) 1269-1274.
- 1291 Okamoto, M. and Nakazawa, H.: Direct enantiomeric resolution of pyriproxyfen (Sumilarv) on a cellulose-based chiral stationary phase. A unique reversal of elution order on changing the nature of the mobile phase modifier. *Anal. Sci.*, 7(Suppl., Proc. Int. Congr. Anal. Sci., 1991, Pt. 1) (1991) 174-150; C.A., 117 (1992) 64700v.
- 1292 Rieger, S.: (Liquid chromatography of pyrethroids). *Acta Univ. Palacki. Olomuc., Fac. Rerum Nat.*, 102(Chem. 30) (1991) 25-34; C.A., 117 (1992) 106205v.
- 1293 Stáb, J.A., Rozing, M.J.M., van Hattum, B., Cofino, W.P. and Brinkman, U.A.T.: Normal-phase high-performance liquid chromatography with UV irradiation, morin complexation and fluorescence detection for the determination of organotin pesticides. *J. Chromatogr.*, 609 (1992) 195-203.
- 1294 Tomaselli, M. and Cozzolino, A.: (Pesticides in the tanning industry. Part II. Quantitative determination). *Cuoio, Pelli, Mater. Concianti*, 67 (1991) 221-235; C.A., 117 (1992) 29154y.
- For additional information see C.A.:
117 (1992) 106282t, 157263v.
30. SYNTHETIC AND NATURAL DYES
- 30a. *Synthetic dyes*
- 1295 Allen, J.L., Meinertz, J.R. and Gofus, J.E.: Determination of malachite green and its leuco form in water. *J. Assoc. Off. Anal. Chem.*, 75 (1992) 646-649.
- 1296 Gruenwald, M., Burtscher, E. and Bobleter, O.: (Determination of dyeing parameters by liquid chromatography). *Textilveredlung*, 27 (1992) 45-50; C.A., 117 (1992) 71542y.
- 1297 Heller, D.N.: Determination of leucogentian violet in chicken fat by liquid chromatography with UV detection. *J. Assoc. Off. Anal. Chem.*, 75 (1992) 650-654.
- 1298 Husain, S., Narsimha, R., Alvi, S.N. and Rao, R.N.: Rapid determination of 4-amino-4'-nitrostilbene-2,2'-disulfonic acid by HPLC. *J. High Resolut. Chromatogr.*, 15 (1992) 562-563.
- 1299 Kabasawa, Y., Tanimura, T., Nakazawa, H. and Shinomiya, K.: Application of counter alternative current chromatography to purification of food mono-azo dyes. *Anal. Sci.*, 8 (1992) 351-353; C.A., 117 (1992) 68603p.
- 1300 Legaz, M.E., Pedrosa, M.M., Mateos, J.L., Caffaro, S.V. and Vicente, C.: High-performance liquid chromatographic determination of acridine orange in nucleic acids isolated from dye-treated *Himantormia lugubris* thalli. *J. Chromatogr.*, 607 (1992) 245-251.
- 1301 Puch, E., Boada, E., Estrany, F. and Ros, J.: (Development of a method of separation of water-soluble fluorescent brighteners by HPLC). *Rev. Quim. Text.*, 102 (1991) 85-98; C.A., 117 (1992) 173375p.
- 1302 Roybal, J.E., Munns, R.K., Holland, D.C., Burkepille, R.G. and Hurlbut, J.A.: Liquid chromatographic determination of gentian violet in poultry feed. *J. AOAC Int.*, 75 (1992) 433-437; C.A., 117 (1992) 46856w.
- For additional information see C.A.:
117 (1992) 61947v.
- 30b. *Chloroplast and other natural pigments*
- 1303 Almela, L., Fernández-López, J.A. and López-Roca, J.M.: High-performance liquid chromatography-diode-array detection of photosynthetic pigments. *J. Chromatogr.*, 607 (1992) 215-219.
- 1304 DeCoster, B., Christensen, R.L., Gebhard, R., Lugtenburg, J., Farhoosh, R. and Frank, H.A.: Low-lying electronic states of carotenoids. *Biochim. Biophys. Acta*, 107-114.
- 1305 Glaessgen, W.E., Seitz, H.U. and Metzger, J.W.: High-performance liquid chromatography/electrospray mass spectrometry and tandem mass spectrometry of anthocyanins from plant tissues and cell cultures of *Daucus carota* L. *Biol. Mass Spectrom.*, 21 (1992) 271-277; C.A., 117 (1992) 65983b.
- 1306 Lee, H.S. and Hong, V.: Chromatographic analysis of anthocyanins. *J. Chromatogr.*, 624 (1992) 221-234 - a review with 126 refs.
- 1307 Li, S. and Inoue, H.: Preparation and determination of manganese(III) chlorophylls by reversed-phase high performance liquid chromatography. *Chromatographia*, 33 (1992) 567-570.
- 1308 O'Neil, C.A. and Schwartz, S.J.: Chromatographic analysis of *cis/trans* carotenoid isomers. *J. Chromatogr.*, 624 (1992) 235-252 - a review with 126 refs.
- 1309 Piccolo, A., Rausa, R. and Celano, G.: Characteristics of molecular size fractions of humic substances derived from oxidized coal. *Chemosphere*, 24 (1992) 1381-1387; C.A., 117 (1992) 30214z.
- 1310 Taylor, S.J. and McDowell, I.J.: Determination of the curcuminoid pigments in turmeric (*Curcuma domestica* Val) by reversed-phase high-performance liquid chromatography. *Chromatographia*, 34 (1992) 73-77.
- 1311 Yamashita, H. and Inoue, H.: Determination of zinc(II) chlorophylls and their derivatives by high performance liquid chromatography with fluorometric detection. *Anal. Sci.*, 7(Suppl., Proc. Int. Congr. Anal. Sci., 1991, Pt. 2) (1991) 1371-1374; C.A., 117 (1992) 65776m.
- For additional information see C.A.:
117 (1992) 46839t, 166819h.

See also 40, 410, 742, 1170, 1174, 1180, 1181, 1190, 1192, 1209, 1532, 1540.

31. PLASTICS AND THEIR INTERMEDIATES

- 1312 Babenko, E.V., Galimzyanov, R.M., Muratova, G.Ya. and Yusupova, M.K.: (New approach to the development of optimal formulations for polymeric composites). *Izv. Vyssh. Uchebn. Zaved., Khim. Khim. Tekhnol.*, 35 (1992) 78-82; C.A., 117 (1992) 112647g.
- 1313 Chubarova, E.V., Sudareva, N.N. and Nesterov, V.V.: (Degradation of high-molecular-weight polystyrene on macroporous glass membranes in size-exclusion chromatography). *Vysokomol. Soedin., Ser. A*, 34 (1992) 151-156; C.A., 117 (1992) 151750e.
- 1314 Cotts, P.M.: Characterization and solution properties of polyamic acid and ester precursors of polyimides. *J. Appl. Polym. Sci.: Appl. Polym. Symp.*, 51 (1992) 101-116; C.A., 117 (1992) 172342p - a review with 21 refs.
- 1315 Glockner, G. and Wolf, D.: Gradient high performance liquid chromatography of block copolymers of styrene and *t*-butyl methacrylate. II. Association phenomena in mixtures of block copolymer with polystyrene. *Chromatographia*, 34 (1992) 363-368.
- 1316 Khorshih, F., Lin, S., Jensen, A. and Kwoh, D.: Characterization of copolyester of poly(ethylene terephthalate) and poly(oxyethylene terephthalate) by proton NMR and gel permeation chromatography. *Polym. Bull. (Berlin)*, 28 (1992) 451-458; C.A., 117 (1992) 91179p.
- 1317 Kretschmer, K. and Helbig, W.: Differentiation of commercial polyester fibers using two-dimensional high-performance liquid chromatography (HPLC) and multivariate pattern recognition techniques. *J. Forensic Sci.*, 37 (1992) 727-737; C.A., 117 (1992) 126006j.
- 1318 Mourey, T.H., Guistina, R.A., Bryan, T.G., Thornbury, R.F., Blevis, R., Perry, R.J. and Turner, S.R.: N-Methylation of wholly aromatic polyamides for size-exclusion chromatography. *J. Appl. Polym. Sci.*, 45 (1992) 1983-1994; C.A., 117 (1992) 112231s.
- 1319 Naken, J.K. and Koopetngarm, S.: An approach to the analysis of crosslinked epoxy systems. *Chromatographia*, 34 (1992) 276-280.
- 1320 Opresnik, M., Sebenik, A., Zigon, M. and Osredkar, U.: The course of crosslinking of a brominated epoxy resin with dicyandiamine studied by GPC. *Vestn. Slov. Kem. Drus.*, 39 (1992) 33-46; C.A., 117 (1992) 152059y.
- 1321 Pasch, H., Brinkmann, C., Much, H. and Just, U.: Chromatographic investigations of macromolecules in the "critical range" of liquid chromatography. II. Two-dimensional separations of poly(ethylene oxide-block-propylene oxide). *J. Chromatogr.*, 623 (1992) 315-322.
- 1322 Rathmann, D.: Trace analysis of inorganic contamination of silicon water surface by ion chromatography. *Proc.-Electrochem. Soc.*, 92-12 (1992) 338-343; C.A., 117 (1992) 39337z.
- 1323 Rathmann, D. and Fabry, L.: Determination of inorganic trace contamination on thermoplastic carrier surface by ion chromatography. *Electrochem. Soc.*, 91-12 (1992) 344-349; C.A., 117 (1992) 39338a.
- 1324 Simal Gándara, J., Paz Abuín, S., López Mahía, P., Paseiro Losada, P. and Simal Lozano, J.: Identification of RP-HPLC peaks of bisphenol F and of bisphenol F diglycidyl ether and its hydrolysis products by thermospray mass spectrometry and gas chromatography/mass spectrometry. *Chromatographia*, 34 (1992) 67-72.
- 1325 Sreenivasan, K.: Characterization of poly(urethane-g-methyl methacrylate) by GPC. *Acta Polym.*, 43 (1992) 188-189; C.A., 117 (1992) 172416r.
- 1326 Vilenchik, L. and Ayotte, R.: New developments in liquid-chromatographic analysis of branched polymers. *J. Appl. Polym. Sci.: Appl. Polym. Symp.*, 51 (1992) 73-90; C.A., 117 (1992) 173173w.

For additional information see C.A.:

117 (1992) 27750r, 28597h, 49688d, 49689e, 91168j, 112396z, 131876x, 163193p, 163194q, 172411k.

See also 21, 68, 69, 74, 351.

32. DRUG ANALYSIS

32a. Drug analysis, general techniques

- 1327 Ahuja, S.: Chromatographic solutions to pharmaceutical analytical problems. *Chromatographia*, 34 (1992) 411-416.
- 1328 Bernal, J.L., del Nozal, M.J., Buj, G.A.G. and Juárez, J.M.: Use of methyl and ethyl acetate as organic modifiers in reversed-phase high-performance liquid chromatography. Application to impurity control in bulk drug steroids. *J. Chromatogr.*, 607 (1992) 175-181.
- 1329 Cooper, A.D., Jefferies, T.M. and Gaskell, R.M.: Resolution of racemic pharmaceuticals by high-performance liquid chromatography using β -cyclodextrin in the mobile phase. *Anal. Proc. (London)*, 29 (1992) 258-259; C.A., 117 (1992) 76609k.
- 1330 Cope, M.J.: Aspects of chiral high-performance liquid chromatography in pharmaceutical analysis: scientific and regulatory considerations. *Anal. Proc. (London)*, 29 (1992) 180-183; C.A., 117 (1992) 56087k.
- 1331 Guebitz, G., Pierer, B. and Wendelin, W.: Resolution of the enantiomers of drugs containing amino alcohol structure after derivatization with bromoacetic acid. *Chirality*, 4 (1992) 333-337; C.A., 117 (1992) 137780h.
- 1332 Izumoto, T., Aioi, A., Uenoyama, S., Kuriyama, K. and Azuma, M.: Relationship between the transference of a drug from a transdermal patch and the physicochemical properties. *Chem. Pharm. Bull.*, 40 (1992) 456-458.
- 1333 Katz, S.E. and Siewierski, M.: Drug residue analysis using immunoaffinity chromatography. *J. Chromatogr.*, 624 (1992) 403-409 - a review with 22 refs.
- 1334 Shepherd, M.J. and Carter, J.M.: Immunoaffinity chromatography for veterinary drug residue analysis. In: Morgan, M.R.A., Smith, C.J. and Williams, P.A. (Editors), *Food Saf. Qual. Assur.: Appl. Immunoassay Syst., Proc.*, 1st 1991, Elsevier, London, 1992, pp. 191-198; C.A., 117 (1992) 110163x - a review with 22 refs.

- 1335 Soltes, L.: (Prechromatographic preparation of biological samples by extraction in a liquid-solid sorbent system for determination of basic drugs). *Chem. Listy*, 86 (1992) 299-301; *C.A.*, 117 (1992) 142772m.
- 1336 Stefansson, M., Lu, B. and Westerlund, D.: Separation of glucuronides from urine by coupled-column separation using underivatized silica as precolumn. *J. Chromatogr. Sci.*, 30 (1992) 324-328.
- 1337 Sweeny, D.J. and Nellans, H.N.: Enantiomeric activation of glucuronidation in dog hepatic microsomes. *J. Biol. Chem.*, 267 (1992) 13171-13174.
- 1338 Vervoort, R.J.M., Maris, F.A. and Hindriks, H.: Comparison of high-performance liquid chromatographic methods for the analysis of basic drugs. *J. Chromatogr.*, 623 (1992) 207-22.
- 1339 Whelpton, R. and Buckley, D.G.: Comparison of Spherisorb Chiral 1 and Chiral 2 high-performance liquid chromatographic columns for the resolution of some drug enantiomers. *Anal. Proc. (London)*, 29 (1992) 249-251; *C.A.*, 117 (1992) 118588k.
- 1340 Wongyai, S.: (Analysis of drugs in biological fluids by direct injection into ISRP columns). *Warasan Phesatchasat*, 17 (1990) 36-44; *C.A.*, 117 (1992) 163239h - a review with 15 refs.
- 1341 Zhou, F.-X., Krull, I.S. and Feibush, B.: 9-Fluoreneacetyl-tagged, solid-phase reagent for derivatization in direct plasma injection. *J. Chromatogr.*, 609 (1992) 103-112.
- 1348 Leo, G.C. and Wu, W.-N.: The utility of one-dimensional homonuclear Hartmann-Hahn spectroscopy (1D HOHAHA) for identifying the suprofen glucuronides fraction from an HPLC separation. *J. Pharm. Biomed. Anal.*, 10 (1992) 607-613.
- 1349 Rummyantsev, D.O.: (Determination of indomethacin and orthophen in blood serum and dosage forms by HPLC). *Khim.-Farm. Zh.*, 26 (1992) 87-88; *C.A.*, 117 (1992) 76616k.
- 1350 Saari, H., Tulamo, R.M., Kontinen, Y.T. and Sorsa, T.: Methylprednisolone acetate induced release of cartilage proteoglycans: determination by high performance liquid chromatography. *Ann. Rheum. Dis.*, 51 (1992) 214-219; *C.A.*, 117 (1992) 40696x.
- 1351 Thomas, S.B., Surber, B.W. and Fitzgerald, M.: Preparative separation and analysis of the enantiomers of [¹⁴C]Zileuton, a 5-lipoxygenase inhibitor. *J. Chromatogr.*, 623 (1992) 390-394.
- 1352 Wang, J. and Moore, D.E.: A study of the photodegradation of benzydamine in pharmaceutical formulations using HPLC with diode array detection. *J. Pharm. Biomed. Anal.*, 10 (1992) 535-540.
- 1353 Wu, S.-M., Chen, S.-H. and Wu, H.-L.: Determination of beta-methasone and dexamethasone by derivatization and liquid chromatography. *Anal. Chim. Acta*, 268 (1992) 255-260.

For additional information see *C.A.*:
117 (1992) 19755f, 137768k.

See also 174, 1379, 1410, 1500.

For additional information see *C.A.*:

117 (1992) 33776v, 33862v, 163276t.

See also 32, 88, 154, 199, 205, 210, 1262, 1317, 1510, 1610.

32b. Antirheumatics and antiinflammatory drugs

- 1342 Caturla, M.C. and Cusido, E.: Solid-phase extraction for the high-performance liquid chromatographic determination of indomethacin, suxibuzone, phenylbutazone and oxyphenbutazone in plasma, avoiding degradation of compounds. *J. Chromatogr.*, 581 (1992) 101-107.
- 1343 Gonzales Tavares, L., Perez de la Cruz, M.J., Sanz Saiz, P., Camacho, M.A. and Martin, J.L.: High-pressure liquid chromatographic determination of the new nonsteroidal anti-inflammatory agent butibufen. *Arzneim.-Forsch.*, 42 (1992) 818-820; *C.A.*, 117 (1992) 33860t.
- 1344 Goodall, D.M., Riley, M.J., Wu, Z. and Wilson, I.D.: Determination of the ratio of R- and S-ibuprofen in purine using a solid-phase extraction procedure followed by high-performance liquid chromatography with dual polarimetric and ultraviolet absorbance detection. *Anal. Proc. (London)*, 29 (1992) 253-255; *C.A.*, 117 (1992) 62202k.
- 1345 Hochhaus, G., Hochhaus, R., Werber, G., Derendorf, H. and Möllmann, H.: A selective HPLC/RIA for dexamethasone and its prodrug dexamethasone-21-sulphobenzoate sodium in biological fluids. *Biomed. Chromatogr.*, 6 (1992) 283-286.
- 1346 Jagota, N.K. and Stewart, J.T.: Determination of mefenamic acid in human serum using solid-phase extraction and high-performance liquid chromatography-electrochemical detection. *LC-GC*, 10 (1992) 688-691; *C.A.*, 117 (1992) 163289z.
- 1347 Kim, H. and Chi, S.C.: Quantitation of flurbiprofen in isopropyl myristate by high performance liquid chromatography. *Yakche Hakhoechi*, 22 (1992) 63-68; *C.A.*, 117 (1992) 118594j.
- 32c. *Autonomic and cardiovascular drugs*
- 1354 Chabenat, C. and Boucly, P.: Determination of naphazoline in rat plasma using column liquid chromatography with ultraviolet detection. *Biomed. Chromatogr.*, 6 (1992) 241-243.
- 1355 Ching, C.B., Lim, B.G., Lee, E.J.D. and Ng, S.C.: Chromatographic resolution of the chiral isomers of several β -blockers over cellulose tris(3,5-dimethylphenylcarbamate) chiral stationary phase. *Chirality*, 4 (1992) 174-177; *C.A.*, 117 (1992) 97420g.
- 1356 Chiou, R.H.-Y. and Lo, M.-W.: High-performance liquid chromatographic determination of angiotensin II receptor antagonists in human plasma and urine. I. DuP 532 (L-694,492). *J. Chromatogr.*, 581 (1992) 165-170.
- 1357 Dyas, A.M.: The chiral chromatographic separation of β -adrenoceptor blocking drugs. *J. Pharm. Biomed. Anal.*, 10 (1992) 383-404 - a review with 152 refs.
- 1358 Erram, S. and Tipnis, H.P.: Simultaneous estimation of nadolol and hydrochlorothiazide in tablets by RP-HPLC. *Indian Drugs*, 29 (1992) 484-488; *C.A.*, 117 (1992) 157758k.
- 1359 Flores-Murrieta, F.J., Castaneda-Hernandez, G. and Hong, E.: Pharmacokinetics and antihypertensive effect of oral pelanserin in renal hypertensive dogs. *Arzneim.-Forsch.*, 42 (1992) 1105-1108.
- 1360 Flume, L., Mattioli, A., Busi, C., Corzani, S. and di Stefano, G.: High-pressure liquid chromatographic method for determining adenine arabinoside monophosphate coupled to lactosaminated albumin in plasma. *Pharm. Acta Helv.*, 66 (1991) 230-232; *C.A.*, 116 (1992) 86v.

- 1361 Gamberini, G., Ferioli, V., Rusticelli, C. and Vezzani, F.: Separation of chiral compounds with α -acid glycoprotein as selector. *Atti Soc. Nat. Mat. Modena*, 121 (1990) 111-120; *C.A.*, 117 (1992) 62166b.
- 1362 Goto, J., Shao, G., Ito, M., Kuriki, T. and Nambara, T.: High-performance liquid chromatographic determination of penbutolol enantiomers in plasma with fluorescence detection. *Anal. Sci.*, 7 (1991) 723-726; *C.A.*, 116 (1992) 66q.
- 1363 He, L. and Stewart, J.T.: A high performance liquid chromatographic method for the determination of albuterol enantiomers in human serum using solid phase extraction and chemical derivatization. *Biomed. Chromatogr.*, 6 (1992) 291-294.
- 1364 Jin, Z., Shen, X., Chen, E. and Xu, X.: (Determination of glycerol trinitrate in plasma by high-performance liquid chromatography-thermal energy analysis). *Sepu*, 10 (1992) 158-159; *C.A.*, 117 (1992) 62173b.
- 1365 Junnarkar, G. and Stavchansky, S.: Thermal degradation of famotidine in aqueous solution stability indicating assay. *Anal. Lett.*, 25 (1992) 1907-1913.
- 1366 Kintz, P., Lohner, S., Tracqui, A. and Mangin, P.: Rapid high-performance liquid chromatographic screening for calcium channel-blocking drugs in human plasma samples. *Toxicol. Methods*, 2 (1992) 42-45; *C.A.*, 117 (1992) 82804m.
- 1367 Miller, R.B. and Arnestoy, C.: A liquid chromatographic method for the determination of hydrochlorothiazide in human plasma. *J. Pharm. Biomed. Anal.*, 10 (1992) 541-545.
- 1368 Prunonosa, J., Obach, R., Diez-Cascón, A. and Gouesclou, L.: Comparison of high-performance liquid chromatography and high-performance capillary electrophoresis for the determination of cicletanine in plasma. *J. Chromatogr.*, 581 (1992) 219-226.
- 1369 Shi, D., Chen, G. and Zhuo, H.: (Simultaneous determination of propafenone and ethmozine concentrations in human plasma by high performance liquid chromatography). *Sepu*, 9 (1991) 372-374; *C.A.*, 116 (1992) 50770k.
- 1370 Smith, D.E., Guillard, S. and Rodriguez, C.A.: Effect of angiotensin II-induced changes in perfusion flow rate on chlorothiazide transport in the isolated perfused rat kidney. *J. Pharmacokin. Biopharm.*, 20 (1992) 195-207.
- 1371 Van Ginkel, L.A., Stephany, R.W. and van Rossum, H.J.: Development and validation of a multiresidue method for β -agonists in biological samples and animal feed. *J. Assoc. Off. Anal. Chem.*, 75 (1992) 554-560.
- 1372 Van Pelt, J., Verheesen, P.E., van Oosterhout, A.G.M., Beukeveld, G.J.J. and Brombacher, P.J.: Interference of dipyridamole in the analysis of porphyrins by HPLC. *Ann. Clin. Biochem.*, 29 (1992) 347-348; *C.A.*, 117 (1992) 39834j.
- 1373 Wall, G.M., Kenny, J.C., Fan, T.Y., Schafer, C., Ready, M.A., Baker, J.K., Ritter, P. and Scott, B.S.: Analysis of 3- and 4-monopivaloyl epinephrine, degradation products in dipivefrin hydrochloride drug substance and ophthalmic formulations. *J. Pharm. Biomed. Anal.*, 10 (1992) 465-471.
- 32d. *Central nervous system drugs*
- 1374 Altria, K.D., Goodall, D.M. and Roban, M.M.: Chiral separation of β -amino alcohols by capillary electrophoresis using cyclodextrins as buffer additives. I. Effect of varying operating parameters. *Chromatographia*, 34 (1992) 19-24.
- 1375 Arrizabalaga, P., Kamatari, A. and Landry, J.C.: (Determination of the composition of a fraudulent drug by coupled chromatographic methods (GC/MS, GC-FTIR, HPLC-DAD). The case of the Chinese pills, Chufong Toukuwan). *Chimia*, 46 (1992) 161-163; *C.A.*, 117 (1992) 137745a.
- 1376 Ascalone, V., Flaminio, L., Guinebault, P., Thénot, J.P. and Morselli, P.L.: Determination of zolpidem, a new sleep-inducing agent, and its metabolites in biological fluids: pharmacokinetics, drug metabolism and overdosing investigations in humans. *J. Chromatogr.*, 581 (1992) 237-250.
- 1377 Balkóvá, M.: Selective system of identification and determination of antidepressants and neuroleptics in serum or plasma by solid-phase extraction followed by high-performance liquid chromatography with photodiode-array detection in analytical toxicology. *J. Chromatogr.*, 581 (1992) 75-81.
- 1378 Bargo, E.S.: Liquid chromatographic determination of flurazepam hydrochloride in bulk drug and dosage forms: collaborative study. *J. AOAC Int.*, 75 (1992) 240-244; *C.A.*, 117 (1992) 76617m.
- 1379 Bousquet, E., Tirendi, S., Bonina, F.P., Montenegro, L., Bianchi, A.: Bioavailability of two formulations of acetylsalicylic acid gums. *Pharmazie*, 47 (1992) 607-609.
- 1380 Brunner, L.A. and Powell, M.L.: An automated method for the determination of a new potential antiepileptic agent (CGP 33101) in human plasma using high performance liquid chromatography. *Biomed. Chromatogr.*, 6 (1992) 278-282.
- 1381 Brunner, L.A., Stelling, M.S. and Luders, R.C.: Automated simultaneous determination of carbamazepine and its metabolite, carbamazepine-10,11-epoxide, in human plasma using high-performance liquid chromatography. *Lab. Rob. Autom.*, 4 (1992) 9-14; *C.A.*, 117 (1992) 82803k.
- 1382 Bulut, P. and Nigdeliöglu, M.: (Determination of lidocaine and chlohexidine in catheter lubricant gels by high-pressure liquid chromatography). *FABAD Farm. Bilimler Der.*, 17 (1992) 11-16; *C.A.*, 117 (1992) 33832k.
- 1383 Caccia, S., Confalonieri, S., Guiso, G., Celeste, S. and Marini, P.P.: Liquid chromatography of the potential memory-enhancing agent LC 275,838 and its main metabolites, using a post-column photochemical reactor and fluorimetric detection. *J. Chromatogr.*, 581 (1992) 109-117.
- 1384 Casas, M., Berrueta, L.A., Gallo, B. and Vicente, F.: Solid phase extraction conditions for the selective isolation of drugs from biological fluids predicted using liquid chromatography. *Chromatographia*, 34 (1992) 79-82.
- 1385 Chankvetadze, B.G., Okuyava, V.M., Chityashvili, Z.D. and Mikutidze, A.S.: Metabolic and pharmacokinetic study of the anti-epileptic drug pufemide. In: Sunshine, I. (Editor), *Recent Dev. Ther. Drug Monit. Clin. Toxicol.*, Dekker, New York, 1992, pp. 451-458; *C.A.*, 117 (1992) 82797m.
- 1386 Chiou, R.H.-Y. and Lo, M.W.: Determination of remoxipride in human plasma and urine by reversed-phase ion-pair high-performance liquid chromatography. *J. Chromatogr.*, 581 (1992) 300-305.

For additional information see *C.A.*:
117 (1992) 82793g, 137571r, 137772g.

See also 553, 1120, 1514.

- 1387 Diaz-Garcia, J.M., Evans, A.M. and Rowland, M.: Application of axial dispersion model of hepatic drug elimination to the kinetics of diazepam in the isolated perfused rat liver. *J. Pharmacokin. Biopharm.*, 20 (1992) 171-193.
- 1388 Doležalová, M.: Ion-pair high-performance liquid chromatographic determination of morphine and pseudomorphine in injections. *J. Pharm. Biomed. Anal.*, 10 (1992) 507-514.
- 1389 Flesch, G., Francotte, E., Hell, F. and Degen, P.H.: Determination of the R(-) and S(+) enantiomers of the monohydroxylated metabolite of oxcarbazepine in human plasma by enantioselective high-performance liquid chromatography. *J. Chromatogr.*, 581 (1992) 147-151.
- 1390 Fraser, A.D. and MacNeil, W.: HPLC analysis of gabapentin: a new anticonvulsant drug. In: Sunshine, I. (Editor), *Recent Dev. Ther. Drug Monit. Clin. Toxicol.*, Dekker, New York, 1992, pp. 313-320; *C.A.*, 117 (1992) 62183e.
- 1391 Fritze, J., Becker, T., Ziegler, V., Laux, G., Bieck, P., Sofic, E. and Riederer, P.: Brofaromine (CGP 11 305 A): estimation of plasma concentrations by a biologic technique as compared to liquid chromatography. *J. Neural Transm., Suppl.*, 32(Amine Oxidases Their Impact Neurobiol.) (1990) 197-201; *C.A.*, 117 (1992) 142768q.
- 1392 Gan, B.K., Baugh, D., Liu, R.H. and Walia, A.S.: Simultaneous analysis of amphetamine, methamphetamine, and 3,4-methylenedioxyamphetamine (MDMA) in urine samples by solid-phase extraction, derivatization, and gas chromatography/mass spectrometry. *J. Forensic Sci.*, 36 (1991) 1331-1341; *C.A.*, 117 (1992) 103456s.
- 1393 Göber, B., Franke, P. and Lisowski, H.: Zur Stabilität von Oxetacain und zur Struktur seiner Abbauprodukte. *Pharmazie*, 47 (1992) 584-587.
- 1394 Grosa, G., Caputo, O., Ceruti, M. and Biglino, G.: Metabolism of 7-(1,3-thiazolidin-2-yl-methyl)theophylline by rat liver microsomes. Evidence for a monooxygenase-dependent step in 1,3-thiazolidine ring cleavage. *Drug Metab. Disp.*, 20 (1992) 742-746.
- 1395 Hartmann, B., Schmieder, G., Tetzloff, W. and Töberich, H.: Bioäquivalenz und Bioverfügbarkeit nach Einmalgabe von löslichen Ranitidin-Tabletten. *Arzneim.-Forsch.*, 42 (1992) 1020-1022.
- 1396 Hartter, S., Wetzel, H. and Hiemke, C.: Automated determination of fluvoxamine in plasma by column-switching high-performance liquid chromatography. *Clin. Chem. (Winston-Salem)*, 38 (1992) 2082-2086.
- 1397 Herold, M.L., Constanzer, M.L. and Matuszewski, B.K.: Determination of picogram levels of heptylphosphostigmine in human plasma using high-performance liquid chromatography with fluorescence detection. *J. Chromatogr.*, 581 (1992) 227-236.
- 1398 Hewala, I.I.: GLC and HPLC determination of diazepam and its degradation products in pharmaceutical formulations. *Anal. Lett.*, 25 (1992) 1877-1905.
- 1399 Ivanovic, D., Herrenkhecht, C., Guernet-Nivaud, E. and Guernet, M.: High-performance liquid chromatography and UV-derivative spectrophotometric determination of acetylsalicylic acid and chlormezanone. *Anal. Lett.*, 25 (1992) 1693-1703.
- 1400 Kaliszan, R., Kaliszan, A., Noctor, T.A.G., Purcell, W.P. and Wainer, I.W.: Mechanism of retention of benzodiazepines in affinity, reversed-phase and adsorption high-performance liquid chromatography in view of quantitative structure-retention relationships. *J. Chromatogr.*, 609 (1992) 69-81.
- 1401 Kaliszan, R., Noctor, T.A.G. and Wainer, I.W.: Quantitative structure-enantioselective retention relationships for the chromatography of 1,4-benzodiazepines on a human serum albumin based HPLC chiral stationary phase: an approach to the computational prediction of retention and enantioselectivity. *Chromatographia*, 33 (1992) 546-550.
- 1402 Karlsson, M. and Wikberg, T.: Liquid chromatographic determination of a new catechol-O-methyltransferase inhibitor, entacapone, and its Z-isomer in human plasma and urine. *J. Pharm. Biomed. Anal.*, 10 (1992) 593-600.
- 1403 Kikura, R., Ishigami, A. and Nakahara, Y.: (Studies on comparison of metabolites in urine between deprenyl and methamphetamine. III. Enantiomeric composition analysis of metabolites in mouse urine by HPLC with GITC chiral reagent). *Jpn. J. Toxicol. Environ. Health*, 38 (1992) 136-141; *C.A.*, 117 (1992) 62157z.
- 1404 Krause, W. and Mechelke, B.: Placental transfer of the anxiolytic β -carboline abecarnil in rabbit. *Arzneim.-Forsch.*, 42 (1992) 1079-1083.
- 1405 Luhmann, I., Szatmary, S.C. and Grünert, I.: Determination of pipamperone in human plasma by high performance liquid chromatography. *Arzneim.-Forsch.*, 42 (1992) 1069-1072.
- 1406 Mazhar, M. and Binder, S.R.: Analysis of benzodiazepines and tricyclic antidepressants in whole blood using a common solid-phase clean-up and a common mobile phase. In: Sunshine, I. (Editor), *Recent Dev. Ther. Drug Monit. Clin. Toxicol.*, Dekker, New York, 1992, pp. 325-335; *C.A.*, 117 (1992) 62184f.
- 1407 McNulty, M.J., Deal, D.L., Nelson, F.R., Weller, S., Chandrasurin, P., Shockcor, J. and Findlay, J.W.A.: Disposition of acrivastine in the male beagle dogs. *Drug Metab. Disp.*, 20 (1992) 679-687.
- 1408 Misztal, G.: (Clozapine determination in blood serum by high-performance liquid chromatography). *Acta Pol. Pharm.*, 47 (1990) 7-9; *C.A.*, 117 (1992) 142779u.
- 1409 Nishikata, M., Nakai, A., Fushida, H., Miyake, K., Arita, T., Kitagawa, S., Kunitomo, M., Iseke, K. and Miyazaki, K.: Method for optical resolution of racemic homochlorcyclizine and comparison of optical isomers in antihistamine activity and pharmacokinetics. *Chem. Pharm. Bull.*, 40 (1992) 1341-1342.
- 1410 Patsi, J., Malkki, L. and Tammielto, S.: Simultaneous determination of caffeine, propylphenazone and salicylamide by Kalman filter spectrophotometry and high-performance liquid chromatography. *Acta Pharm. Nord.*, 4 (1992) 69-72; *C.A.*, 117 (1992) 97428r.
- 1411 Pettersson, K.-J. and Nilsson, L.B.: Determination of dextropropoxyphene and norpropoxyphene in plasma by high-performance liquid chromatography. *J. Chromatogr.*, 581 (1992) 161-164.
- 1412 Rose, M.D. and Shearer, G.: Determination of tranquilisers and carazolol residues in animal tissue using high-performance liquid chromatography with electrochemical detection. *J. Chromatogr.*, 624 (1992) 471-477.
- 1413 Royer-Morrot, M.J., Rambourg, M., Jacob, I., Bauer, P. and Royer, R.J.: Determination of zopiclone in plasma using column liquid chromatography with ultraviolet detection. *J. Chromatogr.*, 581 (1992) 297-299.
- 1414 Sagar, K.A., Hua, C., Kelly, M.T. and Smyth, M.R.: Analysis for salbutamol in human plasma by high-performance liquid chromatography with electrochemical detection using a micro-electrochemical flow cell. *Electroanalysis (N. Y.)*, 4 (1992) 481-486; *C.A.*, 117 (1992) 62197n.

- 1415 Sakurai, E., Yamasaki, S., Iizuka, Y., Hikichi, N. and Niwa, H.: The optical resolution of racemic chlorpheniramine and its stereoselective pharmacokinetics in rat plasma. *J. Pharm. Pharmacol.*, 44 (1992) 44-47; C.A., 116 (1992) 120320q.
- 1416 Sarkar, M.A., March, C. and Karnes, H.T.: Solid phase extraction and simultaneous high performance liquid chromatographic determination of antipyrine and its major metabolites in urine. *Biomed. Chromatogr.*, 6 (1992) 300-304.
- 1417 Srivastava, R.C., Nagappa, A.N., Raju, D.B. and Das, A.K.: Liquid membrane phenomena in the multiple actions of psychotropic drugs. *Indian J. Chem., Sect. A: Inorg., Bio-inorg., Phys., Theor. Anal. Chem.*, 31A (1992) 533-538; C.A., 117 (1992) 123907t.
- 1418 St. Louis, P., Rybczynski, J., Bissonnette, B. and Hartley, E.J.: Determination of bupivacaine in plasma by high-performance liquid chromatography. Levels after scalp infiltration in children. *Clin. Biochem.*, 24 (1991) 463-467; C.A., 116 (1992) 120290e.
- 1419 Suckow, R.F., Zhang, M.F. and Cooper, T.B.: Sensitive and selective liquid-chromatographic assay of fluoxetine and norfluoxetine in plasma with fluorescence detection after precolumn derivatization. *Clin. Chem. (Winston-Salem)*, 38 (1992) 1756-1761.
- 1420 Swart, P.J. and de Zeeuw, R.A.: Extensive gastrointestinal metabolic conversion limits the oral bioavailability of the dopamine D2 agonist N-0923 in freely moving rats. *Pharmazie*, 47 (1992) 613-615.
- 1421 Szumilo, H., Przyborowska, M., Przyborowski, L. and Misztal, G.: (Study of certain dibenzazepine and dibenzocycloheptadiene derivatives by thin-layer and high-performance liquid chromatography). *Acta Pol. Pharm.*, 47 (1990) 11-15; C.A., 117 (1992) 142778t.
- 1422 Tamisier-Karolak, L., Delhotal-Landes, B., Jolliet-Riant, P., Milliez, J., Jannet, D., Barre, J. and Flouvat, B.: Plasma assay of salbutamol by means of high-performance liquid chromatography with amperometric determination using a loop column for injection of plasma extracts. Application to the evaluation of subcutaneous administration of salbutamol. *Ther. Drug Monit.*, 14 (1992) 243-248; C.A., 117 (1992) 82821q.
- 1423 Tatsuno, M., Nishikawa, M., Katagi, M., Tsuchihashi, H., Igarashi, K., Kasuya, F. and Fukui, M.: (Determination of oxazepam glucuronide in human urine by liquid chromatography/ACPI-mass spectrometry). *Jpn. J. Toxicol. Environ. Health*, 38 (1992) 162-167; C.A., 117 (1992) 64337g.
- 1424 Taylor, P.J., Charles, B.G., Norris, R., Salm, P. and Ravenscroft, P.J.: Measurement of dothiepin and its major metabolites in plasma by high-performance liquid chromatography. *J. Chromatogr.*, 581 (1992) 152-155.
- 1425 Tebbett, I.R., Negrusz, A., Larsen, A.K., Jr. and Jensen, R.C.: A column switching technique for the determination of lidocaine and its metabolites in horse urine. *Chromatographia*, 34 (1992) 143-145.
- 1426 Thoma, K. and Schlütermann, B.: Beziehungen zwischen Herstellungsparametern und pharmazeutisch-technologischen Anforderungen an biodegradierbare Mikropartikeln. 6. Mitteilung: Zur Freisetzung von Cinchocain und Bupivacain aus biodegradierbaren Partikeln *in vitro Pharmazie*, 47 (1992) 436-439.
- 1427 Üstün, M., Sungur, S. and Ersoy, L.: Comparison of HPLC and derivative spectrophotometric method for the determination of paracetamol (acetaminophen) and acetylsalicylic acid in tablets. *Pharmazie*, 47 (1992) 558-559.
- 1428 Verjee, Z. and Giesbrecht, E.: An HPLC method for the quantitation of propafenone and its metabolites in pediatric samples. In: Sunshine, I. (Editor), *Recent Dev. Ther. Drug Monit. Clin. Toxicol.*, Dekker, New York, 1992, pp. 383-388; C.A., 117 (1992) 82796k.
- 1429 Verweij, A.M.A., Lipman, P.J.L. and Zweipfenning, P.G.M.: Quantitative liquid chromatography, thermospray/tandem mass spectrometry (LC/TSP/MS/MS) analysis of some thermolabile benzodiazepines in whole-blood. *Forensic Sci. Int.*, 54 (1992) 67-74; C.A., 117 (1992) 126014k.
- 1430 Walters-Thompson, K.M. and Mason, W.D.: Method for the determination of diphenhydramine in rabbit whole blood by high-performance liquid chromatography (HPLC) with ultraviolet (UV) detection in conjunction with gas chromatography (GC) with mass selective detection (MSD). *Pharm. Res.*, 9 (1992) 929-932; C.A., 117 (1992) 103447q.
- 1431 Weems, H. and Zamani, K.: Resolution of terfenadine enantiomers by β -cyclodextrin chiral stationary phase high-performance liquid chromatography. *Chirality*, 4 (1992) 268-272; C.A., 117 (1992) 178394d.
- 1432 William, H., Safi, H. and Westkaemper, R.B.: Initial studies on the N-glucosylation of phenobarbital by mouse liver microsomes using a radiochemical high-performance liquid chromatographic (HPLC) method. *Pharm. Res.*, 9 (1992) 613-616; C.A., 117 (1992) 39682h.
- 1433 Yamaguchi, J., Kokatsu, J. and Suwa, T.: Thermospray liquid chromatography/tandem mass spectrometry of thermally labile xenobiotic conjugates. *Biol. Mass Spectrom.*, 21 (1992) 285-291; C.A., 117 (1992) 62159b.
- 1434 Ye, L., Li, G., Yu, H., Ye, B. and Xue, H.: (Determination of lidocain by reversed-phase high performance liquid chromatography in healthy human fluids and fat tissue). *Sepu*, 10 (1992) 225-226; C.A., 117 (1992) 142770j.
- 1435 Zamani, K., Conner, D.P., Weems, H.B., Yang, S.K. and Cautilena, L.R., Jr.: Enantiomeric analysis of terfenadine in rat plasma by HPLC. *Chirality*, 3 (1991) 467-470; C.A., 116 (1992) 165681z.
- 1436 Zarparkar, S.S., Bhounsule, N.J. and Halkar, U.P.: High-performance liquid chromatographic determination of ketotifen hydroxifen fumarate in pharmaceuticals. *Indian Drugs*, 29 (1992) 365-366; C.A., 117 (1992) 33846t.

For additional information see C.A.:

117 (1992) 39692m, 55789d, 76587b, 76605f, 82786g, 84780z, 84781a, 118587j, 143614y.

See also 105, 202, 1233, 1327, 1341, 1349, 1497.

32e. Chemotherapeutics (exc. cytostatics and antibiotics)

- 1437 Agarwal, V.K.: High-performance liquid chromatographic methods for the determination of sulfonamides in tissue, milk and eggs. *J. Chromatogr.*, 624 (1992) 411-423 - a review with 45 refs.
- 1438 Arnér, E.S.J., Valentin, A. and Eriksson, S.: Thymidine and 3'-azido-3'-deoxythymidine metabolism in human peripheral blood lymphocytes and monocyte-derived macrophages. A study of both anabolic and catabolic pathways. *J. Biol. Chem.*, 267 (1992) 10968-10975.

- 1439 Blessington, B., Beiraghi, A., Lo, T.W., Drake, A. and Jonas, G.: Chiral HPLC-CD studies of the antituberculosis drug (+)-ethambutol. *Chirality*, 4 (1992) 227-229; C.A., 117 (1992) 118586h.
- 1440 Brocks, D.R., Pasutto, F.M. and Jamali, F.: Analytical and semi-preparative high-performance liquid chromatographic separation and assay of hydroxychloroquine enantiomers. *J. Chromatogr.*, 581 (1992) 83-92.
- 1441 Coppi, G. and Silingardi, S.: HPLC method for pharmacokinetic studies on ciclopiroxolamine in rabbits after intravenous and intravaginal administrations. *Farmaco*, 47(Suppl.) (1992) 779-786; C.A., 117 (1992) 142782q.
- 1442 Gavlick, W.K.: High-performance liquid chromatographic analysis of chlorhexidine and *p*-chloroaniline using a specialty column and photodiode-array detector. *J. Chromatogr.*, 623 (1992) 375-380.
- 1443 Gruszecka-Kowalik, E., Haugwitz, R.D. and Zalkow, L.H.: Quinobene, a new potent anti-HIV agent. *Biochem. Biophys. Res. Commun.*, 187 (1992) 1409-1417.
- 1444 Guggisberg, D., Mooser, A.E. and Koch, H.: Methods for the determination of sulphonamides in meat. *J. Chromatogr.*, 624 (1992) 425-437 - a review with 89 refs.
- 1445 Jarboe, H.H. and Kleinow, K.M.: Matrix solid-phase dispersion isolation and liquid chromatographic determination of oxolinic acid in channel catfish (*Ictalurus punctatus*) muscle tissue. *J. AOAC Int.*, 75 (1992) 428-432; C.A., 117 (1992) 46855v.
- 1446 Kawaguchi, S.: (Determination of sulfaquinoxaline in premix at trace level by high performance liquid chromatography). *Shiryō Kenkyū Hokoku (Tokyo Hishiryō Kensasho)*, 17 (1992) 52-57; C.A., 117 (1992) 130056t.
- 1447 Kijak, P.J.: Liquid chromatographic determination of ivermectin in bovine milk: collaborative study. *J. Assoc. Off. Anal. Chem.*, 75 (1992) 747-750.
- 1448 Kitade, Y., Suzuki, A., Hirota, K., Maki, Y., Nakane, H., Ono, K., Baba, M. and Shigeta, S.: Synthesis and anti-human immunodeficiency virus type 1 (HIV-1) activity of 3-substituted derivatives of 3'-azido-3'-deoxythymidine (AZT) and inhibition of HIV-1 reverse transcriptase by their 5'-triphosphates. *Chem. Pharm. Bull.*, 40 (1992) 920-924.
- 1449 Leroy, P., Gavrilloff, C., Nicolas, A., Archimbault, P. and Ambroggi, G.: Comparative assay of amoxicillin by high-performance liquid chromatography and microbiological methods for pharmacokinetic studies in calves. *Int. J. Pharm.*, 82 (1992) 157-164; C.A., 117 (1992) 82819v.
- 1450 Markus, J. and Sherma, J.: Regulatory analytical methods. Method I. Liquid chromatography/fluorescence determination of ivermectin in animal tissue and plasma. *J. Assoc. Off. Anal. Chem.*, 75 (1992) 757-767.
- 1451 Markus, J. and Sherma, J.: Regulatory analytical methods. Method II. Liquid chromatography/fluorescence confirmatory assay of ivermectin in cattle, sheep, and swine liver tissues. *J. Assoc. Off. Anal. Chem.*, 75 (1992) 767-771.
- 1452 Mberu, E.K., Muhia, D.K. and Watkins, W.M.: Measurement of halofantrine and its major metabolite desbutylhalofantrine in plasma and blood by high-performance liquid chromatography: a new methodology. *J. Chromatogr.*, 581 (1992) 156-160.
- 1453 Naesens, L., Balzarini, J. and de Clercq, E.: Pharmacokinetics in mice of the anti-retrovirus agent 9-(2-phosphonylmethoxyethyl)adenine. *Drug Metab. Disp.*, 20 (1992) 747-752.
- 1454 Okamura, T., Hirano, S., Sugiura, W. and Sakai, T.: Simultaneous determination of sulfamethoxazole and its metabolite N⁴-acetyl-sulfamethoxazole in human urine using solid-phase extraction technique. *Chem. Express*, 7 (1992) 373-376; C.A., 117 (1992) 19782n.
- 1455 Parks, O.W. and Kubena, L.F.: Liquid chromatographic determination of incurred nitrofurazone residues in chicken tissues. *J. AOAC Int.*, 75 (1992) 261-262; C.A., 117 (1992) 88899t.
- 1456 Pingarron Carrazon, J.M., Dominguez Recio, A. and Polo Diez, L.M.: Electroanalytical study of sulphamerazine at a glassy-carbon electrode and its determination in pharmaceutical preparations by HPLC with amperometric detection. *Talanta*, 39 (1992) 631-635; C.A., 117 (1992) 56111p.
- 1457 Rao, B.E., Raghuvver, S. and Srivastava, C.M.R.: HPLC determination of rifampicin, pyrazinamide and isoniazid in single and combined dosage forms. *Indian Drugs*, 29 (1992) 412-415; C.A., 117 (1992) 76611e.
- 1458 Reising, K.P.: Rapid analysis for ivermectin residue in liver and muscle tissue by liquid chromatography. *J. Assoc. Off. Anal. Chem.*, 75 (1992) 751-753.
- 1459 Schenck, F.J., Barker, S.A. and Long, A.R.: Matrix solid-phase dispersion extraction and liquid chromatographic determination of ivermectin in bovine liver tissue. *J. Assoc. Off. Anal. Chem.*, 75 (1992) 655-658.
- 1460 Schenck, F.J., Barker, S.A. and Long, A.R.: Matrix solid-phase dispersion extraction and liquid chromatographic determination of nicarbazin in chicken tissue. *J. Assoc. Off. Anal. Chem.*, 75 (1992) 659-662.
- 1461 Shah, Y., Khanna, S., Jindal, K.C. and Dighe, V.S.: Determination of rifampicin and isoniazid in pharmaceutical formulation by HPLC. *Drug Dev. Ind. Pharm.*, 18 (1992) 1589-1596; C.A., 117 (1992) 97434q.
- 1462 Shimomura, M. and Ishikuro, E.: (Determination of halofuginone calcium polystyrene-sulfonate in mixed feed at trace level by high performance liquid chromatography). *Shiryō Kenkyū Hokoku (Tokyo Hishiryō Kensasho)*, 17 (1992) 60-69; C.A., 117 (1992) 130057u.
- 1463 Siegmund, W., Zschiesche, M., Franke, G. and Wilke, A.: Bioavailability of metronidazole formulations (Vagimid®). *Pharmazie*, 47 (1992) 522-525.
- 1464 Tollomelli, B., Laffi, R., Rossato, S. and Colpo, G.: Liquid chromatographic assay of dimetridazole, ipronidazole, and ronidazole in feeds and premixes. *J. Assoc. Off. Anal. Chem.*, 75 (1992) 663-666.
- 1465 Valenta, A., Lexer, A. and Spiegl, P.: Analysis of clotrinazole in ointments by high-performance liquid chromatography. *Pharmazie*, 47 (1992) 641-642.
- 1466 Vinet, B., Comtois, R., Gervais, A. and Lemieux, C.: Clinical usefulness of high-pressure liquid chromatographic determination of serum pentamidine in AIDS patients. *Clin. Biochem.*, 25 (1992) 93-97; C.A., 117 (1992) 142775q.
- 1467 Wang, Y., Jiang, Y. and Fan, S.: (Determination of sulfonamide hydrophobicity by micellar liquid chromatography). *Sepu*, 10 (1992) 229-232; C.A., 117 (1992) 142771k.

For additional information see C.A.:

117 (1992) 39680f, 39734b, 46834n, 62165a, 68601m, 76358c, 76585z.

See also 411, 1287, 1382, 1524.

32f. *Cytostatics*

- 1468 Barberi-Heyob, M., Merlin, J.L. and Weber, B.: Analysis of 5-fluorouracil in plasma and urine by high-performance liquid chromatography. *J. Chromatogr.*, 581 (1992) 281-286.
- 1469 Berges, F. and Holler, E.: Ring-substituted diaqua(1,2-diphenylethylenediamine)platinum(II) sulfate reacts with DNA through a dissociable complex. *Eur. J. Biochem.*, 208 (1992) 573-579.
- 1470 Bissett, D., Graham, M.A., Setanoians, A., Chadwick, G.A., Wilson, P., Koier, I., Henrar, R., Schwartzmann, G., Cassidy, J., Kaye, S.B. and Kerr, D.J.: Phase I and pharmacokinetic study of rhizoxin. *Cancer Res.*, 52 (1992) 2894-2898.
- 1471 Li, Y.M., Jiang, S.X. and Chen, L.R.: Direct injection analysis of hexamethylene bisacetamide in biological fluids by an HPLC column switching technique with a micellar mobile phase. *Chromatographia*, 34 (1992) 63-66.
- 1472 Phillips, T.M.: Measurement of recombinant interferon levels by high performance immunoaffinity chromatography in body fluids of cancer patients on interferon therapy. *Biomed. Chromatogr.*, 6 (1992) 287-290.
- 1473 Pichini, S., Altieri, I., Bacosi, A., di Carlo, S., Zuccaro, P., Iannetti, P. and Pacifici, R.: High-performance liquid chromatographic-mass spectrometric assay of busulfan in serum and cerebrospinal fluid. *J. Chromatogr.*, 581 (1992) 143-146.
- 1474 Reid, J.M., Mathiesen, D.A., Benson, L.M., Kuffel, M.J. and Ames, M.M.: Murine pharmacokinetics and metabolism of penclomedine [3,5-dichloro-2,4-dimethoxy-6-(trichloromethyl)pyridine, NSC 338720]. *Cancer Res.*, 52 (1992) 2830-2834.
- 1475 Richeimer, S.L., Tinnermeier, D.M. and Timmons, D.W.: High-performance liquid chromatographic assay of taxol. *Anal. Chem.*, 64 (1992) 2323-2326.
- 1476 Swaffar, D.S., Harker, W.G., Pomerantz, S.C., Lim, H.-K. and Yost, G.S.: Isolation, purification, and characterization of two new chemical decomposition products of methylazoxypocarbazine. *Drug Metab. Disp.*, 20 (1992) 632-642.
- 1477 Zhu, Y., Gao, Y., Zheng, Y. and Qu, J.: (RP-HPLC determination of 1-hexylcarbamoyl-5-fluorouracil and its pharmacokinetic study in rats serum). *Yaowu Fenxi Zazhi*, 12 (1992) 142-145; *C.A.*, 117 (1992) 62172a.

For additional information see *C.A.*:
117 (1992) 76581v.

See also 265, 1251, 1520.

32g. *Other drug categories*

- 1478 Alban, S., Kraus, J. and Franz, G.: Synthesis of laminarin sulfates with anticoagulant activity. *Arzneim.-Forsch.*, 42 (1992) 1005-1008.
- 1479 Anderson, N.H., Johnston, D. and Vojvodic, P.R.: The enantiomeric resolution of ciprofibrate and related compounds by HPLC using chiral stationary phases. *J. Pharm. Biomed. Anal.*, 10 (1992) 501-505.
- 1480 Aubry, A.-F., Caude, M. and Rosset, R.: Separation and identification of dipeptides in a hydrolyzed brain extract. *Chromatographia*, 33 (1992) 533-538.
- 1481 Bakuridze, A.D., Tsgareishvili, N.T., Dargaeva, T.D., Patudin, A.V. and Berashvili, T.D.: (Chromatospectrophotometric determination of γ -pyrone substances contents in the above-ground parts of *Gentiana lutea* L.) *Rastit. Resur.*, 27 (1991) 115-119; *C.A.*, 117 (1992) 56095m.
- 1482 Christians, U., Braun, F., Schmidt, M., Kosian, N., Schiebel, H.-M., Ernst, L., Winkler, M., Kruse, C., Linck, A. and Sewing, K.-F.: Specific and sensitive measurement of FK506 and its metabolites in blood and urine of liver-graft recipients. *Clin. Chem. (Winston-Salem)*, 38 (1992) 2025-2032.
- 1483 Del Nozal, M.J., Bernal, J.L., Pampliega, A., Pastor, J.C. and Lopez, M.I.: Determination of 5-fluorouracil in vitreous gel and liquid by high-performance liquid chromatography. *J. Chromatogr.*, 607 (1992) 183-189.
- 1484 Di Pietra, A.M., Bonazzi, D. and Cavrini, V.: Analysis of chlorpropamide, tolbutamide and their related sulphonamide impurities by liquid chromatography (HPLC). *Farmaco*, 47 (1992) 787-797; *C.A.*, 117 (1992) 178401d.
- 1485 El-Yazigi, A. and Raines, D.A.: Simultaneous microdetermination of rifampin, deacetyl-rifampin, isoniazid, and acetylisoniazid in plasma by liquid chromatography with dual electrochemical and spectrophotometric detection. *Pharm. Res.*, 9 (1992) 812-816; *C.A.*, 117 (1992) 19793s.
- 1486 Feng, Z., Gollamudi, R., Han, G., Tang, Y. and Armstrong, D.W.: Chiral separation of nipecotic acid amides. *J. Chromatogr.*, 609 (1992) 187-193.
- 1487 Frontini, R. and Mielck, J.B.: Determination and quantitation of bendroflumethiazide and its degradation products using HPLC. *J. Liq. Chromatogr.*, 15 (1992) 2519-2528.
- 1488 Gilbert, J.D., Olah, T.V., Barrish, A. and Greber, T.F.: Determination of L-654,066, a new 5 α -reductase inhibitor in plasma by liquid chromatography/atmospheric pressure chemical ionization mass spectrometry. *Biol. Mass Spectrom.*, 21 (1992) 341-346; *C.A.*, 117 (1992) 82802j.
- 1489 Guo, Y., Chen, L., Lai, Y. and Hu, Z.: (Application of HPLC in the preparation of radioiodine labeled dopamine receptor imaging agents). *Tongweisu*, 3 (1990) 222-227, 221; *C.A.*, 117 (1992) 166694p.
- 1490 Herpol-Borremans, M. and Masse, M.O.: Amount of octyl triazone in cosmetic sunscreen products. *Int. J. Cosmet. Sci.*, 14 (1992) 113-119; *C.A.*, 117 (1992) 178077c.
- 1491 Imai, Y. and Kobayashi, S.: A simple method for the quantification of famotidine in human plasma and urine by paired-ion high performance liquid chromatography. *Biomed. Chromatogr.*, 6 (1992) 222-223.
- 1492 Lloyd, T.L., Perschy, T.D., Gooding, A.E. and Tomlinson, J.J.: Robotic solid phase extraction and high performance liquid chromatographic analysis of ranitidine in serum or plasma. *Biomed. Chromatogr.*, 6 (1992) 311-316.
- 1493 McMeekin, J.R., Fowles, S.E., Winton, C.F. and Pierce, D.M.: Rapid high-performance liquid chromatographic method for the analysis for the active antitherpes agent, penciclovir and its precursor, BRL 42359, in human plasma and urine following administration of the oral prodrug, famciclovir. *Anal. Proc. (London)*, 29 (1992) 178-180; *C.A.*, 117 (1992) 62170y.
- 1494 Nayak, V.G., Bhate, V.R., Dhupal, S.N., Nerurkar, V.R., Purandare, S.M., Dikshit, P.M. and Gaitonde, C.D.: Estimation of ipratropium bromide from aerosols by reversed-phase liquid chromatography. *Drug. Dev. Ind. Pharm.*, 18 (1992) 1681-1689; *C.A.*, 117 (1992) 157765k.

- 1495 Nobilis, M., Pastera, J., Svoboda, D., Květina, J. and Macek, K.: High-performance liquid chromatographic determination of ambroxol in human plasma. *J. Chromatogr.*, 581 (1992) 251-255.
- 1496 Noggle, F.T., Jr., Clark, C.R. and DeRuiter, J.: Liquid chromatographic and spectral analysis of the stereoisomers of dimethylaminorex. *J. Assoc. Off. Anal. Chem.*, 75 (1992) 423-427.
- 1497 Raju, N.R., Srilakshmi, P. and Krishna, U.M.: Simultaneous determination of chlorpheniramine maleate, ephedrine hydrochloride and codeine phosphate in cough syrups by reversed-phase high-performance liquid chromatography. *Indian Drugs*, 29 (1992) 408-411; *C.A.*, 117 (1992) 76610d.
- 1498 Reynolds, D.L., Eichmeier, L.S. and Giesing, D.H.: Determination of MDL 201,012 at femtomole/millilitre levels in human plasma by liquid chromatography with electrochemical detection. *Biomed. Chromatogr.*, 6 (1992) 295-299.
- 1499 Sattler, M., Guengerich, F.P., Yu, C.-H., Christians, U. and Sewing, K.-F.: Cytochrome P-450 3A enzymes are responsible for biotransformation of FK506 and rapamycin in man and rat. *Drug Metab. Disp.*, 20 (1992) 753-761.
- 1500 Scalia, S. and Massaccesi, M.: Simultaneous determination of bendazac and underivatized lysin in ophthalmic preparations by reversed-phase ion-pair high-performance liquid chromatography. *Int. J. Pharm.*, 82 (1992) 179-183; *C.A.*, 117 (1992) 97430k.
- 1501 Sips, J.H.M., van Tellingen, O., Beijnen, J.H., Underberg, W.J.M. and Nooijen, W.J.: (Determination of the diastereomers [6S]- and [6R]-folinate in plasma with on-line two-dimensional liquid chromatography.) *Pharm. Weekbl.*, 127 (1992) 356-361; *C.A.*, 117 (1992) 19760d.
- 1502 Weaver, M., Tanzer, J.M. and Kramer, P.A.: High-performance liquid chromatographic determination of pilocarpine in plasma. *J. Chromatogr.*, 581 (1992) 293-296.
- 1503 Weerasinghe, C.A., Lewis, D.O., Mathews, J.M., Jeffcoat, A.R., Troxler, P.M. and Wang, R.Y.: Aquatic photodegradation of albendazole and its major metabolites. 1. Photolysis rate and half-life for reactions in a tube. *J. Agric. Food Chem.*, 40 (1992) 1413-1418.
- For additional information see *C.A.*:
117 (1992) 62146v, 103429k, 157730v, 178387d.
- See also 419, 435, 1337, 1347, 1536, 1541, 1557, 1558.
- 32h. *Toxicological and forensic applications*
- 1504 Angerer, J. and Hörsch, B.: Determination of aromatic hydrocarbons and their metabolites in human blood and urine. *J. Chromatogr.*, 580 (1992) 229-255 - a review with 150 refs.
- 1505 De Kok, T.M.C.M., Levels, P.J., van Faassen, A., Hazen, M., ten Hoor, F. and Kleinjans, J.C.S.: Chromatographic methods for the determination of toxicants in faeces. *J. Chromatogr.*, 580 (1992) 135-159 - a review with 209 refs.
- 1506 Felscher, D.: (Application of high-pressure liquid chromatography with diode-array detection in emergency medicine). *Beitr. Gerichtl. Med.*, 49 (1991) 381-386; *C.A.*, 117 (1992) 165386c.
- 1507 Felscher, D.: (Application of high-pressure liquid chromatography with diode-array detection in forensic drug identification). *Beitr. Gerichtl. Med.*, 49 (1991) 375-379; *C.A.*, 117 (1992) 144912f.
- 1508 Koves, E.M. and Wells, J.: Evaluation of a photodiode array/HPLC-based system for the detection and quantitation of basic drugs in *postmortem* blood. *J. Forensic Sci.*, 37 (1992) 42-60; *C.A.*, 117 (1992) 144928r.
- 1509 Lehmann, T. and Brenneisen, R.: A new chromatographic method for the isolation of (-)- Δ^9 -(trans)-tetrahydrocannabinolic acid A. *Phytochem. Anal.*, 3 (1992) 88-90; *C.A.*, 117 (1992) 21568r.
- 1510 Moeller, M.R.: Drug detection in hair by chromatographic procedures. *J. Chromatogr.*, 580 (1992) 125-134 - a review with 60 refs.
- 1511 Nelson, C.C. and Foltz, R.L.: Chromatographic and mass spectrometric methods for determination of lysergic acid diethylamide (LSD) and metabolites in body fluids. *J. Chromatogr.*, 580 (1992) 97-109 - a review with 41 refs.
- 1512 Tagliaro, F., Lubli, G., Ghielmi, S., Franchi, D. and Marigo, M.: Chromatographic methods for blood alcohol determination. *J. Chromatogr.*, 580 (1992) 161-190 - a review with 208 refs.
- See also 252, 502, 554, 557, 571, 594, 621, 735, 876, 1085, 1088, 1106, 1116, 1118, 1119, 1121, 1143, 1145, 1268, 1423, 1425, 1515, 1530, 1549, 1570, 1573.
- 32i. *Plant extracts*
- 1513 Drost-Karbowska, K., Szauffer-Hajdrych, M. and Ozimek-Matti, J.: (Determination of capsaicinoids in capsicum fruits after chromatographic separation). *Acta Pol. Pharm.*, 47 (1990) 41-44; *C.A.*, 117 (1992) 137762d.
- 1514 Hashimoto, K., Yasuda, T. and Ohsawa, K.: Determination of synephrine from Chinese medicinal drugs originating from *Citrus* species by ion-pair high-performance liquid chromatography. *J. Chromatogr.*, 623 (1992) 386-389.
- 1515 Helmlin, H.-J., Bourquin, D. and Brenneisen, R.: Determination of phenylethylamines in hallucinogenic cactus species by high-performance liquid chromatography with photodiode-array detection. *J. Chromatogr.*, 623 (1992) 381-385.
- 1516 Maekawa, B. and Morimoto, K.: Development of analytical method for selective detection of cardenolides by high-performance liquid chromatography. *Biosci., Biotechnol., Biochem.*, 56 (1992) 967; *C.A.*, 117 (1992) 82816s.
- 1517 Nishioka, Y., Kyotani, S., Miyamura, M. and Kusunose, M.: Influence of time of administration of a Shosaiko-to extract granule on blood concentration of its active constituents. *Chem. Pharm. Bull.*, 40 (1992) 1335-1337.
- 1518 Sanz, M.J., Terencio, M.C. and Paya, M.: Isolation and hypotensive activity of a polymeric procyanidin fraction from *Pistacia lentiscus* L. *Pharmazie*, 47 (1992) 466-467.
- 1519 Sun, W.J., Sha, Z.F. and Gao, H.: Determination of arctiin and arctigenin in fructus arctii by reversed-phase HPLC. *Yaoxue Xuebao*, 27 (1992) 549-551; *C.A.*, 117 (1992) 157739e.
- 1520 Weinberg, D.S., Manier, M.L., Richardson, M.D., Haibach, F.G. and Rogers, T.S.: Identification and quantification of anticarcinogens in garlic extract and licorice root extract powder. *J. High Resolut. Chromatogr.*, 15 (1992) 641-654.
- 1521 Zarate, R., Sukrasno and Yeoman, M.M.: Application of two rapid techniques of column chromatography to separate the pungent principles of ginger, *Zingiber officinale* Roscoe. *J. Chromatogr.*, 609 (1992) 407-413.

For additional information see C.A.:

117 (1992) 33779y, 33783v, 33788a, 33789b.

See also 268, 300, 531, 1137, 1375, 1509, 1559.

33. CLINICO-CHEMICAL APPLICATIONS

33a. General papers and reviews

1522 Yamada, F., Ito, M. and Iida, H.: Automated liquid chromatographs for clinical analyses. *Jpn. Kokai Tokkyo Koho* JP 04 50,651 [92 50,651] (Cl. GO1N30/86), 19 Feb. 1992, Appl. 90/152,742, 13 Jun. 1990; 4 pp.; C.A., 117 (1992) 44041c.

33b. Complex mixtures and profiling (single compounds by cross-reference only)

1523 Takai, N., Nagata, Y., Kanazawa, H. and Matsushima, Y.: Anion-exchange chromatography of ultraviolet-absorbing compounds in body fluids. In: Abe, M., Kataoka, T. and Suzuki, T. (Editors), *New Dev. Ion Exch. Proc. Int. Sci.*, Kodansha, Tokyo, 1991, pp. 283-286; C.A., 117 (1992) 146439f.

See also 174, 341, 406, 431, 436, 442, 446, 496, 498, 506, 516, 549, 555, 556, 589, 630, 645, 653, 662, 747, 753, 760, 780, 785, 790, 806, 841, 896, 914, 965, 975, 997, 1028, 1081, 1109, 1125, 1138, 1165, 1171, 1185, 1187, 1190, 1192, 1194, 1199, 1384, 1581.

34. FOOD ANALYSIS

34a. General papers and reviews

1524 Botsoglou, N.A. and Kufidis, D.C.: Determination of antimicrobial residues in edible animal products by HPLC. *Food Sci. Technol.*, 52(Food Anal. HPLC) (1992) 457-505; C.A., 117 (1992) 110179g - a review with 211 refs.

1525 Elmroth, I., Sundin, P., Valeur, A., Larsson, L. and Odham, G.: Evaluation of chromatographic methods for the detection of bacterial contamination in biotechnical processes. *J. Microbiol. Methods*, 15 (1992) 215-228; C.A., 117 (1992) 129891e.

1526 Finger, A., Kuhr, S. and Engelhardt, U.H.: Chromatography of tea constituents. *J. Chromatogr.*, 624 (1992) 293-315 - a review with 219 refs.

1527 Flak, W. and Pilsbacher, L.: HPLC determination of additives and preservatives. *Food Sci. Technol.*, 52(Food Anal. HPLC) (1992) 421-456; C.A., 117 (1992) 110178f - a review with 173 refs.

1528 Knize, M.G., Felton, J.S. and Gross, G.A.: Chromatographic methods for the analysis of heterocyclic amine food mutagens/carcinogens. *J. Chromatogr.*, 624 (1992) 253-265 - a review with 28 refs.

1529 Lichon, M.J.: Sample preparation for chromatographic analysis of food. *J. Chromatogr.*, 624 (1992) 3-9 - a review with 49 refs.

1530 Luckas, B.: Phycotoxins in seafood - toxicological and chromatographic aspects. *J. Chromatogr.*, 624 (1992) 439-456 - a review with 101 refs.

1531 McMurrough, I. and Byrne, J.R.: HPLC analysis of bittering substances, phenolic compounds, and various compounds of alcoholic beverages. *Food Sci. Technol.*, 52(Food Anal. HPLC) (1992) 579-641; C.A., 117 (1992) 110181b - a review with 169 refs.

1532 Porretta, S.: Chromatographic analysis of Maillard reaction products. *J. Chromatogr.*, 624 (1992) 211-219 - a review with 77 refs.

1533 Siouffi, A.M.: HPLC. *Food Sci. Technol.*, 52(Food Anal. HPLC) (1992) 1-52; C.A., 117 (1992) 110167b - a review with 123 refs.

See also 277, 327, 407, 467, 469, 481, 482, 487, 504, 519, 536, 538, 540, 610, 611, 646, 707, 726, 797, 1183, 1195, 1214, 1215, 1216, 1261, 1306, 1308, 1333, 1437, 1444, 1571, 1597.

34b. Complex mixtures (single compounds by cross-reference only)

1534 Busturia, M.A.: (Determination of ingredients and additives in nonalcoholic beverages by isotachopheresis and high-pressure liquid chromatography). *Fluess. Obst.*, 59 (1992) 123-128; C.A., 117 (1992) 46836q.

1535 Calull, M., Marcé, R.M., Sánchez, G. and Borrull, F.: Determination of additives in wine by high-performance liquid chromatography. *J. Chromatogr.*, 607 (1992) 339-347.

1536 Hannisdal, A.: Analysis of acesulfame K, saccharin, and preservatives in beverages and jams by HPLC. *Z. Lebensm.-Unters. Forsch.*, 194 (1992) 517-519; C.A., 117 (1992) 169698d.

1537 Henle, T., Walter, H., Krause, I. and Klostermeyer, H.: Efficient determination of individual Maillard compounds in heat-treated milk products by amino acid analysis. *Int. Dairy J.*, 1 (1991) 125-135; C.A., 117 (1992) 149597s.

1538 Mareck-Engelke, U., Galensa, R. and Herrmann, K.: (Identification of exotic fruits in fruit products by HPLC). *Fluess. Obst.*, 58 (1991) 659-663; C.A., 117 (1992) 46819m.

1539 Okayama, A., Ohbayashi, H., Umoto, F., Tanaka, T., Tamase, K., Aoki, Y., Sasaki, M., Nakazawa, H., Suzuki, S. and Fujita, M.: (Elution behavior of soluble dietary fiber with various HPLC columns). *Nara-ken EiseiKenkyusho Nenpo*, 25 (1990; Pub. 1991) 135-138; C.A., 117 (1992) 110264f.

1540 Sujata, V., Ravishankar, G.A. and Venkataraman, L.V.: Methods for the analysis of the saffron metabolites crocin, crocetin, picrocrocin and safranal for the determination of the quality of the spice using thin-layer chromatography, high-performance liquid chromatography and gas chromatography. *J. Chromatogr.*, 624 (1992) 497-502.

1541 Yano, M., Shiba, S., Yokoyama, Y., Tagawa, Y., Masui, T., Ozawa, T., Warabi, Y., Saga, J., Hyodo, N. et al.: (Determination of saccharin in foods by high performance liquid chromatography). *Jpn. J. Toxicol. Environ. Health*, 38 (1992) 196-201; C.A., 117 (1992) 110225u.

For additional information see C.A.:

117 (1992) 46835p, 68623v, 130251c, 169860a.

See also 224, 244, 247, 250, 263, 265, 267, 270, 272, 274, 276, 278., 279, 281, 286, 287, 289, 303, 305, 320, 337, 408, 414, 427, 433, 457, 480, 534, 545, 550, 551, 585, 591, 631, 739, 745, 794, 868, 1041, 1129, 1139, 1166, 1168, 1173, 1175, 1184, 1191, 1196, 1198, 1200, 1205, 1207, 1209, 1231, 1239, 1259, 1264, 1266, 1278, 1289, 1299, 1302, 1371, 1412, 1445, 1446, 1447, 1455, 1462, 1464, 1556, 1558, 1601, 1606.

34c. *Organoleptically important compounds (flavors, odors, volatiles)*

1542 Borejsza-Wysocki, W., Goers, S.K., McArdle, R.N. and Hrazdina, G.: (*p*-Hydroxyphenyl)butan-2-one levels in raspberries determined by chromatographic and organoleptic methods. *J. Agric. Food Chem.*, 40 (1992) 1176-1177.

1543 Rouseff, R.L., Dettweiler, G.R., Swaine, R.M., Naim, M. and Zehavi, U.: Solid-phase extraction and HPLC determination of 4-vinyl guaiacol and its precursor, ferulic acid, in orange juice. *J. Chromatogr. Sci.*, 30 (1992) 383-387.

See also 551, 1526, 1540.

35. ENVIRONMENTAL ANALYSIS

35a. *General papers and reviews*

1544 Behymer, T.D., Bellar, T.A., Ho, J.S. and Budde, W.L.: Development of analytical methods for non-volatiles using liquid chromatography/particle beam/mass spectrometry. *Organohalogen. Compd.*, 2 (1990) 99-102; C.A., 117 (1992) 184039h.

1545 Rueedi, P.: (HPLC in environmental analysis). *Schweiz. Lab.-Z.*, 49 (1992) 119-122; C.A., 117 (1992) 96375j - a review with many refs.

For additional information see C.A.:
117 (1992) 54827c.

See also 137, 231, 467, 1505.

35c. *Water pollution (complex mixtures; single compounds by cross-reference only)*

1546 Butler, E.C.V. and Dal Pont, G.: Liquid chromatography-electrochemistry procedure for the determination of chlorophenolic compounds in pulp mill effluents and receiving waters. *J. Chromatogr.*, 609 (1992) 113-123.

1547 Clark, L.B., Rosen, R.T., Hartman, T.G., Louis, J.B., Suffet, I.H., Lippincott, R.L. and Rosen, J.D.: Determination of alkylphenol ethoxylates and their acetic acid derivatives in drinking water by particle beam liquid chromatography/mass spectrometry. *Int. J. Environ. Anal. Chem.*, 47 (1992) 167-180; C.A., 117 (1992) 96964a.

1548 Huehnerfuss, H.: (Chromatographic separation of the enantiomers of organic pollutants - a new access to process studies in the marine ecosystem). *GIT Fachz. Lab.*, 36 (1992) 489-497; C.A., 117 (1992) 76099u.

1549 Hühnerfuss, H. and Kallenborn, R.: Chromatographic separation of marine organic pollutants. *J. Chromatogr.*, 580 (1992) 191-214 - a review with 138 refs.

1550 Schroeder, H.F.: (Refractory substances in municipal waste waters). *Korresp. Abwasser*, 39 (1992) 387-394; C.A., 117 (1992) 55478b.

1551 Slobodnik, J., Brouwer, E.R., Geerdink, R.B., Mulder, W.H., Lingeman, H. and Brinkman, U.A.T.: Fully automated on-line liquid chromatographic separation system for polar pollutants in various types of water. *Anal. Chim. Acta*, 268 (1992) 55-65.

For additional information see C.A.:
117 (1992) 55205k, 96817e.

See also 253, 254, 256, 410, 588, 1263, 1275, 1276, 1277, 1280, 1283, 1293, 1295.

35d. *Soil pollution (complex mixtures; single compounds by cross-reference only)*

See 410, 1080, 1279, 1285, 1595.

36. SOME TECHNICAL PRODUCTS AND COMPLEX MIXTURES

See 80.

36a. *Surfactants*

1552 Lafosse, M., Marinier, P., Joseph, B. and Dreux, M.: Study of amphiphilic behaviour of alkylglycoside surfactants using reversed-phase liquid chromatography. *J. Chromatogr.*, 623 (1992) 277-287. 708, light scat. det.

1553 Mannhardt, K. and Novosad, J.J.: Modeling adsorption of surfactant mixtures using the surface excess formalism. In: Mersmann, A.B. and Scholl, S.E. (Editors), *Int. Conf. Fundam. Adsorpt.*, 3rd 1989, AIChE, New York, 1991, pp. 543-552; C.A., 117 (1992) 114777y.

1554 Okada, T.: Automated analyses of non-ionic surfactants in terms of hydrophobic and polyoxyethylated chain lengths. *J. Chromatogr.*, 609 (1992) 213-218.

1555 Zeman, I. and Paulovic, M.: (Chromatographic separation of surfactants. Analysis of propylene oxide derivatives of nonylphenol by liquid chromatography). *Petrochemia*, 30 (1990) 141-149; C.A., 117 (1992) 51306w.

36b. *Antioxidants and preservatives*

1556 Duh, P.D., Yeh, D.B. and Yen, G.C.: Extraction and identification of an antioxidative component from peanut hulls. *J. Am. Oil Chem. Soc.*, 69 (1992) 814-818.

1557 Ferioli, V., Vezzalini, F., Rustichelli, C. and Gamberini, G.: Determination of bronidox and bronopol in cosmetic products by reverse phase high-performance liquid chromatography. *Farmaco*, 47 (1992) 833-839; C.A., 117 (1992) 157366f.

1558 Irache, J.M., Vega, F.A. and Ezpeleta, I.: Antioxidants in some pharmaceuticals cosmetics and food from the European market. *Pharm. Acta Helv.*, 67 (1992) 152-155; C.A., 117 (1992) 97416k.

1559 Jitoe, A., Masuda, T., Tengah, I.G.P., Suprpta, D.N., Gara, I.W. and Nakatani, N.: Antioxidant activity of tropical ginger extracts and analysis of the contained curcuminoids. *J. Agric. Food Chem.*, 40 (1992) 1337-1340.

See also 430, 1527.

36c. *Complex mixtures, technical products and unidentified compounds*

- 1560 Aoyagi, T., Hatsu, M., Kojima, F., Hayashi, C., Hamada, M. and Takeuchi, T.: Benarthin: a new inhibitor of pyroglutamyl peptidase. I. Taxonomy, fermentation, isolation and biological activities. *J. Antibiot.*, 45 (1992) 1079-1083.
- 1561 Díaz, M., Moliner, R. and Ibarra, J.V.: Isolation of the phenolic fraction of coal pyrolysis tars by ion-exchange chromatography. *J. Chromatogr.*, 607 (1992) 353-360.
- 1562 Ito, K., Yamamoto, S., Takahashi, J. and Yasuda, Y.: Purification of early pregnancy factor from cattle sera. *Anim. Sci. Technol.*, 63 (1992) 394-397; *C.A.*, 117 (1992) 86109m.
- 1563 Maeda, Y.: (Establishment of an assay method for the advanced-stage Maillard product and changes in its level in the body). *Kobe Daigaku Ighakubu Kiyo*, 52 (1991) 245-252; *C.A.*, 117 (1992) 146529k.
- 1564 Mura, P., Piriou, P.Y., Lochon, D. and Reiss, D.: Rapid high-performance liquid chromatographic assay of chlorophacinone in human serum. *J. Anal. Toxicol.*, 16 (1992) 179-181; *C.A.*, 117 (1992) 42055m.
- 1565 Tokumura, A., Yotsumoto, T., Hoshikawa, T., Tanaka, T. and Tsukatani, H.: Quantitative analysis of platelet-activating factor in rat brain. *Life Sci.*, 51 (1992) 303-308; *C.A.*, 117 (1992) 67267v.

For additional information see *C.A.*:

117 (1992) 110226v, 114803d, 114809k.

See also 148, 235, 237, 365, 1309, 1322, 1323, 1537, 1577, 1587, 1602.

37. *CELLS, CELLULAR PARTICLES AND SUPRAMOLECULAR STRUCTURES*

- 1566 Allfrey, V.G. and Chen, T.A.: Nucleosomes of transcriptionally active chromatin: isolation of template-active nucleosomes by affinity chromatography. *Methods Cell Biol.*, 35(Funct. Organ. Nucleus: Lab. Guide) (1991) 315-335; *C.A.*, 117 (1992) 65785p.
- 1567 Shimeld, S.M. and Sharpe, P.T.: Cell-surface changes induced by ectopic expression of the murine homeobox gene Hox-3.3. *Biochim. Biophys. Acta*, 1136 (1992) 253-258.
- 1568 Walter, H., Raymond, F.D. and Fisher, D.: Erythrocyte partitioning in dextran-poly(ethylene glycol) aqueous phase systems. Events in phase and cell separation. *J. Chromatogr.*, 609 (1992) 219-227.

For additional information see *C.A.*:

117 (1992) 129838t, 146712q.

See also 391.

38. *INORGANIC COMPOUNDS*

38a. *Cations*

- 1569 Akiba, K., Ishibashi, Y. and Nakamura, S.: Application of liquid-liquid extraction of lanthanoid elements to centrifugal partition chromatography. *Process Metall.*, 7B (1992) 1677-1682; *C.A.*, 117 (1992) 142432a.
- 1570 Al-Rashdan, A., Vela, N.P., Caruso, J.A. and Heitkemper, D.T.: Lead speciation by gradient high-performance liquid chromatography with inductively coupled plasma mass spectrometric detection. *J. Anal. At. Spectrom.*, 7 (1992) 551-555; *C.A.*, 117 (1992) 21541b.
- 1571 Danielson, N.D. and Sherman, J.H.: Determination of anions and cations in food by HPLC. *Food Sci. Technol.*, 52(Food Anal. HPLC) (1992) 713-730; *C.A.*, 117 (1992) 110184e - a review with 72 refs.
- 1572 Du, A.: Radiochemical neutron activation analysis for cesium and rubidium in geological samples using zeolite (clinoptilolite) chromatography and sodium tetraphenylborate precipitation. *J. Radioanal. Nucl. Chem.*, 159 (1992) 293-299. *C.A.*, 117 (1992) 163056w.
- 1573 Elchuk, S., Lucy, C.A. and Burns, K.I.: High-resolution determination of ¹⁴⁷Pm in urine using dynamic ion-exchange chromatography. *Anal. Chem.*, 64 (1992) 2339-2343.
- 1574 Fadeeva, V.I., Tikhomirova, T.I., Kudryavtsev, G.V. and Yuferova, I.B.: (Sorption preconcentration, separation and determination of vanadium(V), molybdenum(VI) and tungsten(VI) using chemically modified silica gel). *Zh. Anal. Khim.*, 47 (1992) 466-477; *C.A.*, 117 (1992) 142475s.
- 1575 Gomensoro, S.A.: (Separation of mixtures of mono- and divalent cations by ion chromatography with electronic suppression). *Tech. Lab.*, 14 (1992) 208-211; *C.A.*, 117 (1992) 142433b.
- 1576 Hu, W., Takeuchi, T. and Haraguchi, H.: Ion chromatography of alkali metal ions using micellar bile salt coated stationary phase. *Anal. Sci.*, 8 (1992) 507-509; *C.A.*, 117 (1992) 163020e.
- 1577 Janos, P. and Broul, M.: (Determination of lanthanum, cerium, praseodymium and neodymium in polishing powders using the method of ion interaction chromatography in a reverse system). *Chem. Listy*, 86 (1992) 139-142; *C.A.*, 117 (1992) 136323z.
- 1578 Jones, P., Challenger, O.J., Hill, J.S. and Barnett, N.W.: Advances in chelating exchange ion chromatography for the determination of trace metals using dye-coated columns. *Analyst (London)*, 117 (1992) 1447-1480.
- 1579 Kuhnawar, M.Y. and Soomro, A.I.: Liquid and gas chromatographic procedures for the simultaneous determination of copper, nickel and palladium using tetradentate Schiff bases as complexing agents. *Anal. Chim. Acta*, 268 (1992) 49-53.
- 1580 Khukawar, M.Y. and Soomro, A.I.: High performance liquid chromatographic determination of copper(II) and nickel(II) by using solvent extraction and bis(acetylpyvalylmethane)ethylenediimine as complexing reagent. *Talanta*, 39 (1992) 609-612; *C.A.*, 117 (1992) 39389t.
- 1581 Nagaosa, Y. and Bond, A.M.: Determination of copper, iron, and aluminium in serum by HPLC with electrochemical and spectrophotometric detection. *J. High Resolut. Chromatogr.*, 15 (1992) 622-625.

- 1582 Nagaosa, Y., Kimata, M. and Miura, J.: (Determination of copper(II) as the diethyldithiophosphate complex by HPLC with an electrochemical detector). *Bunseki Kagaku*, 41 (1992) 371-374; C.A., 117 (1992) 146458m.
- 1583 Navratil, J.D., Murphy, A. and Sun, D.: Mixed solvent extraction-annular chromatographic systems for f-element separation and purification. *Process Metall.*, 7B (1992) 1151-1156; C.A., 117 (1992) 134762t - a review with 18 refs.
- 1584 Nguyen Van Suc, Desai, H.B., Parthasarathy, R. and Gangadharan, S.: Rare earth impurities in high-purity lanthanum oxide determined by neutron activation analysis. *J. Radioanal. Nucl. Chem.*, 164 (1992) 321-325; C.A., 117 (1992) 61766k.
- 1585 Oi, T., Ogino, H., Kawada, K., Hosoe, M. and Kakihana, H.: Ion exchange isotope separation factors of alkali and alkaline earth metals. In: Abe, M., Kataoka, T. and Suzuki, T. (Editors), *New Dev. Ion Exch. Proc. Int. Conf. Ion Exch.*, Kodansha, Tokyo, 1991, pp. 299-304; C.A., 117 (1992) 120242m.
- 1586 Park, K.S., Kim, N.B., Woo, H.J., Lee, K.Y., Hong, W. and Chun, S.K.: Determination of rare earth elements in steels by radiochemical neutron activation analysis. *J. Radioanal. Nucl. Chem.*, 160 (1992) 529-538; C.A., 117 (1992) 183909e.
- 1587 Ramakumar, K.L., Saxena, M.K., Raman, V.A., Sant, V.L., Kavimandan, V.D., Datta, B.P. and Jain, H.C.: Determination of zirconium in uranium-zirconium-aluminum and plutonium-zirconium-aluminum alloys by isotope dilution thermal ionization mass spectrometry. *J. Radioanal. Nucl. Chem.*, 158 (1992) 13-21; C.A., 117 (1992) 61761e.
- 1588 Richter, P., Fernandez-Romero, J.M., Luque de Castro, M.D. and Valcarcel, M.: Integrated FIA/HPLC method for preconcentration and determination of transition metal ions. *Chromatographia*, 34 (1992) 445-449.
- 1589 Ryan, E. and Meaney, M.: Determination of trace levels of copper(II), aluminium(III) and iron(III) by reversed-phase high-performance liquid chromatography using a novel on-line sample preconcentration technique. *Analyst (London)*, 117 (1992) 1435-1439.
- 1590 Shum, C.C.K., Pang, H. and Houk, R.S.: Speciation of mercury and lead compounds by microbore column liquid chromatography - inductively coupled plasma mass spectrometry with direct injection nebulization. *Anal. Chem.*, 64 (1992) 2444-2450.
- 1591 Spivakov, B.Ya., Maryutina, T.A. and Zolotov, Yu.A.: Multistage liquid-liquid extraction separation of inorganic ions using a planet centrifuge. *Process Metall.*, 7A (1992) 451-456; C.A., 117 (1992) 142436e.
- 1592 Usuda, S., Abe, H., Tachimori, S., Takeishi, H. and Murayama, W.: Application of centrifugal partition chromatography to separation of actinides and lanthanides in TBP extraction system. *Process Metall.*, 7A (1992) 717-722; C.A., 117 (1992) 120209f.
- 1593 Valle, A.I., González, M.J. and Marina, M.L.: Separation and quantitation of some metal ions by reversed-phase high-performance liquid chromatography using *in situ* complexation with (\pm -*trans*-1,2-diamine)cyclohexane-N,N',N',N'-tetraacetic acid. *J. Chromatogr.*, 607 (1992) 207-213.
- 1594 Verma, S.P.: Usefulness of liquid chromatography for determination of rare-earth elements in rocks and minerals. *Lanthanide Actinide Res.*, 3 (1991) 237-257; C.A., 117 (1992) 82528z.
- 1595 Whitten, M.G., Ritchie, G.S.P. and Willett, I.R.: Forms of soluble aluminum in acidic topsoils estimated by ion chromatography and 8-hydroxyquinoline and their correlation with growth of subterranean clover. *J. Soil Sci.*, 43 (1992) 283-293; C.A., 117 (1992) 150027n.
- 1596 Williams, T.P.: Determination of trace metals by ion chromatography with chemiluminescence detection. Avail. *Univ. Microfilms Int.*, Order No. BRDX93530, 1990, 241 p.; C.A., 117 (1992) 183843d.

For additional information see C.A.:

117 (1992) 39372g, 39444g, 61785r, 75873e, 82544b, 82614z, 103152q, 103167y, 144879s, 183791k, 183851e.

See also 55, 104, 124, 1163, 1607, 1608, 1609, 1611, 1613, 1614.

38b. Anions

- 1597 Fernández Pereira, C.F.: Application of ion chromatography to the determination of inorganic anions in foodstuffs. *J. Chromatogr.*, 624 (1992) 457-570 - a review with 92 refs.
- 1598 Groh, T. and Bächmann, K.: Indirect detection of anions with potassium hexacyanoferrate(II) and -(III) in the sub-pg range. *J. Liq. Chromatogr.*, 15 (1992) 2611-2622.
- 1599 Kaljurand, M., Urbas, E. and Haldna, U.: Correlation ion chromatography with indirect UV detection. *Chromatographia*, 34 (1992) 417-420.
- 1600 Kubota, M., Takai, N. and Okutani, T.: High-performance liquid chromatography of inorganic anions: use of multiwavelength detector as analytical method. In: Abe, M., Kataoka, T. and Suzuki, T. (Editors), *New Dev. Ion Exch. Proc. Int. Conf. Ion Exch.*, Kodansha, Tokyo, 1991, pp. 305-310; C.A., 117 (1992) 142409y.
- 1601 Murcia, M.A., Vera, A. and Garcia-Carmona, F.: Nitrate level in vegetables by ion chromatography. *Biochem. Soc. Trans.*, 20 (1992) 371S; C.A., 117 (1992) 149617y.
- 1602 Revankar, S.N.: Ion-chromatography in the determination of sulphur in steels. *Chem. Environ. Res.*, 1 (1992) 59-62; C.A., 117 (1992) 163836d.
- 1603 Saari-Nordhaus, R. and Anderson, J.M., Jr.: Ion chromatographic separation of inorganic anions and carboxylic acids on a mixed-mode stationary phase. *Anal. Chem.*, 64 (1992) 2283-2287.
- 1604 Salov, V.V., Yoshinaga, J., Shibata, Y. and Morita, M.: Determination of inorganic halogen species by liquid chromatography with inductively coupled argon plasma mass spectrometry. *Anal. Chem.*, 64 (1992) 2425-2428.
- 1605 Stuedel, R. and Münchow, V.: Sulphur compounds. CLIX. Determination of dithionite ($S_2O_4^{2-}$) and hydroxymethanesulphinate ($HOCH_2SO_2^-$; Rongalite) by ion-pair chromatography. *J. Chromatogr.*, 623 (1992) 174-177.
- 1606 Taguchi, N., Hirata, K., Amakawa, E., Ohnishi, K., Nishijima, M., Sakai, S. and Hamada, A.: (Determination of total bromine in a foods by ion chromatography). *Jpn. J. Toxicol. Environ. Health*, 38 (1992) 289-294; C.A., 117 (1992) 130044n.

For additional information see C.A.:

117 (1992) 39484v, 46830h, 61794t, 77341d, 114628a.

See also 39, 214, 1163, 1263, 1571, 1591, 1612.

38d. Volatile inorganic compounds

For additional information see C.A.:
117 (1992) 82712e.

39. RADIOACTIVE AND OTHER ISOTOPE COMPOUNDS

- 1607 Araujo, B.F., Matsuda, H.T., Carvalho, E.I. and Araujo, I.C.: Plutonium removal by ion exchange chromatography. *J. Radioanal. Nucl. Chem.*, 165 (1992) 209-217; C.A., 117 (1992) 120222e.
- 1608 Gorski, B., Gleisberg, B. and Nam, N.T.P.: Separation of the actinides, protactinium, thorium, uranium, neptunium and TPE including mendelevium by extraction methods. *Process Metall.*, 7A (1992) 711-716; C.A., 117 (1992) 120244p.
- 1609 Iwata, R., Kawamura, M., Ido, T. and Kimura, S.: Chromatographic purification of no-carrier-added magnesium-28 biological studies. *J. Radioanal. Nucl. Chem.*, 159 (1992) 233-237; C.A., 117 (1992) 139437a.
- 1610 Khachadoorian, S.K., Shafiq, Y.F. and Al-Janabi, M.A.A.: Comparative study of different chromatographic methods for quality control of radiopharmaceuticals labeled with metastable indium-113. *J. Radioanal. Nucl. Chem.*, 159 (1992) 3-8; C.A., 117 (1992) 76619p.
- 1611 Kimura, T. and Akatsu, J.: Applications of extraction chromatography in the bifunctional extractant-nitric acid system for separation of actinides in nuclear fuel cycle. *Process Metall.*, 7A (1992) 723-728; C.A., 117 (1992) 120210z.
- 1612 Lamb, J.D., Nordmeyer, F.R., Smith, R.G., van Orden, A., Allison, R. and Lash, R.P.: Separation of and analysis for radioiodine in nuclear reprocessing solutions of high ionic strength. *J. Radioanal. Nucl. Chem.*, 159 (1992) 285-292; C.A., 117 (1992) 163055v.
- 1613 Lopez, M.B.E., Serrano, G.J. and Fucugauchi, L.A.: (Separation of uranium from some fission products by ion-exchange chromatography). *Rev. Soc. Quim. Mex.*, 36 (1992) 34-39; C.A., 117 (1992) 78382e.
- 1614 McCabe, W.J., Ditchburn, R.G. and Whitehead, N.E.: Method for separation of polonium, thorium, protactinium and uranium in high yields from various matrixes. *J. Radioanal. Nucl. Chem.*, 159 (1992) 267-279; C.A., 117 (1992) 163016h.
- 1615 Oi, T., Takeda, T., Hosoe, M. and Kakahana, H.: Basic data of ion-exchange separation of boron isotopes. In: Abe, M., Kataoka, T. and Suzuki, T. (Editors), *New Dev. Ion Exch. Proc. Int. Conf. Ion Exch.*, Kodansha, Tokyo, 1991, pp. 293-298; C.A., 117 (1992) 99364r.
- 1616 Snyder, T.S. and Ferrari, H.M.: Uranium isotope separation by continuous anion exchange chromatography). *Brit. UK Pat. Appl. GB 2,250,451* (Cl. B01D15/08), 10 Jun. 1992, US Appl. 620,807, 03 Dec. 1990; 27 p.; C.A., 117 (1992) 120308n.
- 1617 Snyder, T.S. and Lee, E.D.: Zirconium and hafnium production in zero liquid discharge process. *U.S. US 5,112,493* (Cl. 210-656; B01D15/08), 12 May 1992, Appl. 624,782, 10 Dec. 1990; 10 p.; C.A., 117 (1992) 78503v.
- 1618 Thomas, L.C.: Quantitative comparisons using liquid chromatography with dual label radioactivity measurements for evaluation of isotope and impurity effects. *Talanta*, 39 (1992) 599-607; C.A., 117 (1992) 43891t.

For additional information see C.A.:
117 (1992) 33808g, 139436z.

See also 515, 1572, 1573, 1585, 1592.

Gas Chromatography

1. REVIEWS AND BOOKS

- 1 Berezkin, V.G. and Koshevnik, M.A.: (Scientometric analysis of modern chromatography.) *Zh. Anal. Khim.*, 47 (1992) 80-89.
- 2 Braun, T. and Zsindely, S.: The comparative standing of individual instrumental analytical techniques. *TrAC*, 11 (1992) 267-269.
- 3 Grob, K.: *On-line Coupled LC-GC*. Hüthig Verlag, Heidelberg, 1991, xxvii + 462 p.
- 4 McNally, M.E.P. and Bright, F.V.: Fundamental studies and applications of supercritical fluids. A review. *ACS Symp. Ser.*, 488 (1992) 1-15; *C.A.*, 117 (1992) 39200z - a review with 113 refs.
- 5 Rood, D.: *A Practical Guide to the Care, Maintenance and Trouble-shooting of Capillary Gas Chromatographic Systems*. Hüthig Verlag, Heidelberg, 1991, 191 p.
- 6 Schomburg, G.: Progress and problems with miniaturized analytical separation methods: GC, SFC, LC, and CE. *J. High Resolut. Chromatogr.*, 15 (1992) 493-494.

See also 42, 91, 98, 112, 138, 158, 163, 173, 187, 192, 204, 206, 209, 223, 231, 249, 289, 327, 329, 334, 359, 446, 482, 537, 540, 553, 570, 603, 624, 625, 634, 636, 642, 657, 680, 717, 721, 803.

2. FUNDAMENTALS, THEORY AND GENERAL

2a. General

- 7 Blumberg, L.M.: Outline of a theory of focusing in linear chromatography. *Anal. Chem.*, 64 (1992) 2459-2460.
- 8 Fleming, P.: Column dead-time determination methods based on the isothermal retention-time behaviour of a homologous series in chromatography. *Analyst (London)*, 117 (1992) 1553-1557.
- 9 Hinshaw, J.V.: A compendium of GC terms and techniques. *LC-GC Int.*, 5, No. 10 (1992) 22-26.
- 10 Li, S., He, D., He, Q. and Chen, X.: (A method of fitting and quick resolution of skewed and overlapped chromatographic peaks.) *Sepu*, 10 (1992) 251-254.
- 11 Moskvina, L.N. and Katykhin, G.S.: (Chromatography: methodological and nomenclature aspects.) *Zh. Anal. Khim.*, 47 (1992) 1508-1515.
- 12 Oszczapowicz, J.: Similarity models in gas chromatography. *J. Chim. Phys. Phys.-Chim. Biol.*, 89 (1992) 1603-1608; *C.A.*, 117 (1992) 138405b.
- 13 Wu, N.S. and Wei, W.: Empirical equation for relative non-overlap. *Chromatographia*, 34 (1992) 450-452.

2b. Thermodynamics and theoretical relationships

- 14 Poole, C.F., Kollie, T.O. and Poole, S.K.: Recent advances in solvation models for stationary phase characterization and the prediction of retention in gas chromatography. *Chromatographia*, 34 (1992) 281-302.
- 15 Reddy, K.S. and Kováts, E.sz.: Pair-wise interactions by gas chromatography. Part II: solvation free enthalpies of some hydrogen bonding solutes with non-associated primary alcohol groups. *Chromatographia*, 34 (1992) 249-252.
- 16 Reddy, K.S., Dutoit, J.-Cl. and Kováts, E.sz.: Pair-wise interaction by gas chromatography. I. Interaction free enthalpies of solutes with nonassociated primary alcohol groups. *J. Chromatogr.*, 609 (1992) 229-259.
- 17 Voelkel, A.: Inverse gas chromatography: the use of Laffort solubility factors and topological indices in structure-polarity relationships. *J. Chromatogr.*, 623 (1992) 83-91.

See also 207, 218, 220, 221.

2c. Relationship between structure and chromatographic behaviour

- 18 Chrétien, J.R., Righezza, M., Hassani, A. and Meklati, B.Y.: Factor analysis and experimental design in high performance liquid chromatography. XIII. Correspondence factor analysis of the heterogeneity of the behaviour of congeneric solutes series studied on homogeneous OV phases in chromatography. *J. Chromatogr.*, 609 (1992) 261-267.
- 19 Franke, J.P., Wijsbeek, J. and de Zeeuw, R.A.: Interlaboratory reproducibility of retention indexes in capillary gas chromatography. *J. Forensic Sci.*, 35 (1990) 813-820; *C.A.*, 117 (1992) 126009n.
- 20 Gornerova, T. and Singliar, M.: (Effect of the molecular weight of the stationary phase in gas chromatography on retention indexes.) *Petrochemia*, 30 (1990) 84-93; *C.A.*, 117 (1992) 89779r.
- 21 Zenkevich, I.G. and Kuznetsova, L.M.: (Utilization of physico-chemical constants for organic substances in their identification by chromatography - mass spectrometry.) *Zh. Anal. Khim.*, 47 (1992) 982-993.

See also 14, 16, 17, 248, 268, 298, 356, 458, 475, 606.

2d. Measurement of physico-chemical and related values

- 22 Calder, A.G., Anderson, S.E., Grant, I., McNurlan, M.A. and Gailick, P.J.: The determination of low d₅-phenylalanine enrichment (0.002-0.09 atom percent excess), after conversion to phenylethylamine, in relation to protein turnover studies by gas chromatography/electron ionization mass spectrometry. *Rapid Commun. Mass Spectrom.*, 6 (1992) 421-424; *C.A.*, 117 (1992) 127661f.

- 23 Cheng, R.J., Hsu, C.L. and Tsai, Y.F.: Constructing a complete temperature-composition diagram for a binary mixture: a new experiment that uses quantitative gas chromatography. *J. Chem. Educ.*, 69 (1992) 581-582.
- 24 Comor, J.J., Lausevic, Z.V., Kopecni, M.M. and Vukov, A.J.: Alkane adsorption on silicon carbide surface. *J. Colloid Interface Sci.*, 151 (1992) 498-504; *C.A.*, 117 (1992) 77282k.
- 25 Economopoulos, N., Athanassopoulos, N., Katsanos, N.A., Karaiskakis, G., Agathonos, P. and Vassilakos, C.: A plant kinetic study of alcoholic fermentation using reversed-flow gas chromatography. *Sep. Sci. Technol.*, 27 (1992) 2055-2070.
- 26 Gayrilova, T.B., Nikitin, Yu. S., Rudnitskaya, T.A. and Finkel'shtein, A.I.: (Investigation of adsorption properties of cyamelluric acid by gas chromatographic method.) *Zh. Fiz. Khim.*, 66 (1992) 1080-1088.
- 27 Jagiello, J., Bandosz, T.J. and Schwarz, J.A.: Inverse gas chromatographic study of activated carbons: the effect of controlled oxidation on microstructure and surface chemical functionality. *J. Colloid Interface Sci.*, 151 (1992) 433-445; *C.A.*, 117 (1992) 98174s.
- 28 Janák, K., Hägglund, I., Blomberg, L.G., Bergård, A.K. and Colmsjö, A.L.: Universal set-up for measurement of diffusion coefficients in supercritical carbon dioxide with flame ionization detection. *J. Chromatogr.*, 625 (1992) 311-321.
- 29 Khudyakov, V.L.: (Determination of thermodynamic characteristics of benzene and ethanol by means of precision gas chromatography.) *Zh. Fiz. Khim.*, 66 (1992) 785-786.
- 30 Kolb, B., Welter, C. and Bichler, C.: Determination of partition coefficients by automatic equilibrium headspace gas chromatography by vapor phase calibration. *Chromatographia*, 34 (1992) 235-240.
- 31 Kuo, H.-W., Wang, J.-D. and Lin, J.-M.: Determination of vapor pressure for 4,4'-bipyridine by gas saturation and gas chromatographic methods. *Chemosphere*, 24 (1992) 1679-1686.
- 32 Labows, J.H.: Surfactant solubilization behaviour via headspace analysis. *J. Am. Oil Chem. Soc.*, 69 (1992) 34-38.
- 33 Liao, K.: (Determination of the hydrogenated polyisoprene-benzene Flory-Huggins interaction parameter χ by inverse gas chromatography (IGC).) *Sepu*, 10 (1992) 304-305.
- 34 Liu, H., Wang, W. and Lu, H.: (A new model for the gas chromatographic measurements of vapor-liquid equilibrium of dilute solutions). *Beijing Huagong Xueyuan Xuebao*, 18, No. 1 (1991) 1-12; *C.A.*, 116 (1992) 159924q.
- 35 Matsuyama, Y., Yasuoka, T., Shimada, H. and Mitsuzawa, S.: (Measurement of hydrolysis rates of ethyl acetate by gas chromatography.) *Kagaku to Kyoiku*, 40 (1992) 387-389; *C.A.*, 117 (1992) 89570r.
- 36 Melsheimer, J. and Ziegler, D.: Ethane transformation on HZSM-5 studied by combined UV-visible spectroscopy and on-line gas chromatography. *J. Chem. Soc., Faraday Trans.*, 88 (1992) 2101-2108; *C.A.*, 117 (1992) 170738y.
- 37 Murko-Jezovsek, M., Mozetic, M. and Nagy, L.: AES and GC analyses of contact surfaces and protective gases in model reed switches. *Vacuum*; 43 (1992) 631-633; *C.A.*, 117 (1992) 122246q.
- 38 Mňuk, P., Smolková-Keulemansová, E. and Feltl, L.: Gas chromatographic study of the inclusion properties of dibenzo-24-crown-8 ether. *J. Chromatogr.*, 623 (1992) 297-304.
- 39 Oteo, J.L., Rubio, J., Tartaj, J., Fernandez, J.F. and Moure, C.: (Surface characterization of superconductive ceramic powders.) *Bol. Soc. Esp. Ceram. Vidrio*, 30 (1991) 458-461; *C.A.*, 117 (1992) 116861v.
- 40 Panzer, U. and Schreiber, H.P.: On the evaluation of surface interactions by inverse gas chromatography. *Macromolecules*, 25 (1992) 3633-3637; *C.A.*, 117 (1992) 27667u.
- 41 Roles, J.L.: An inverse gas-solid chromatographic study of the adsorption heterogeneity of aluminum oxide powders used in ceramic processing. *Avail. Univ. Microfilm Int., Order No. DA9133766*, 1991, 182 pp.; *C.A.*, 117 (1992) 156075y.
- 42 Rutan, S.C.: Discovering chemistry with chemometrics. *Chemometrics Intel. Lab. Systems*, 15 (1992) 137-141.
- 43 Schiller, M. and Gmehling, J.: Measurement of activity coefficients at infinite dilution using gas-liquid chromatography. 4. Results for alkylene glycol dialkyl ethers as stationary phases. *J. Chem. Eng. Data*, 37 (1992) 503-508; *C.A.*, 117 (1992) 158559h.
- 44 Shekurov, V.N. and Bereznoi, A.N.: (Method for chromatographic determination of coefficients of interdiffusion of binary systems in a gas phase.) *Teor. Osn. Khim. Tekhnol.*, 26 (1992) 354-363; *C.A.*, 117 (1992) 174171f.
- 45 Smith, J. and Patterson, D.: Application of gas-liquid chromatography to the study of gas-bitumen interactions. *Can. J. Chem. Eng.*, 70 (1992) 818-821; *C.A.*, 117 (1992) 134181c.
- 46 Suarez, J.J., Bueno, J.L., Medina, I. and Dizy, J.: (Applications of supercritical chromatography: determination of molecular diffusivity.) *Afinidad*, 49 (1992) 101-113; *C.A.*, 117 (1992) 33883c.
- 47 Tse, G., Orbey, H. and Sandler, S.I.: Infinite dilution activity coefficients and Henry's law coefficients of some priority water pollutants determined by a relative gas chromatographic method. *Environ. Sci. Technol.*, 26 (1992) 2017-2022; *C.A.*, 117 (1992) 137137d.
- 48 Vasil'ev, A.V. and Aleksandrov, E.N.: (Physico-chemical application of open tubular column chromatography.) *Usp. Khim.*, 61 (1992) 689-710.
- 49 Wu, L. and He, H.: Determination of absorption efficiencies of carbon dioxide absorbents by gas chromatography. *Talanta*, 39 (1992) 1081-1087.
- 50 Yu, Z.-R., Rizvi, S.S.H. and Zollweg, J.A.: Phase equilibria of oleic acid, methyl oleate, and anhydrous milk fat in supercritical carbon dioxide. *J. Supercrit. Fluids*, 5 (1992) 114-122.
- See also 15, 196, 213, 550.

3. GENERAL TECHNIQUES

3a. Apparatus and accessories

- 51 Bansagi, G., Langer, K. and Gyori, F.: (Flame ionization measuring instrument supplied from a hydrogen generator, for measuring C_nH_m [hydrocarbons].) *Hung. Töljes*, HU 58,914 (Cl. G01N27/62), 30 Mar. 1992, Appl. 89/6,815, 29 Dec. 1989; 8 pp.; *C.A.*, 117 (1992) 103319z.

- 52 Beche, M. and Roussel, P.: (Method and apparatus for injecting a sample into a column for gas-phase chromatographic separation.) *Fr. Demande* FR 2,664,385 (Cl. G01N30/24), 10 Jan. 1992, Appl. 90/8,913, 09 Jul. 1990; 23 pp.; C.A., 117 (1992) 39595g.
- 53 Behm, F. and Streit, P.: Capillary-tube method and apparatus for enriching samples for analysis in gas chromatography. *Eur. Pat. Appl.* EP 493,314 (Cl. G01N1/28), 01 Jul. 1992, CH Appl. 90/4,065, 20 Dec. 1990; 6 pp.; C.A., 117 (1992) 163130r.
- 54 Bolkunov, V.G., Kan, T.P., Korol'kov, A.V., Okhotnikov, B.P., Semenova, L.A. and Skornyakov, E.P.: (Evaporator arrangement for introducing samples into a gas chromatograph.) *U.S.S.R. SU* 1,702,301 (Cl. G01N30/18), 30 Dec. 1991, Appl. 4,769,829, 15 Dec. 1989; C.A., 117 (1992) 103318y.
- 55 Burford, M.D., Hawthorne, S.B., Miller, D.J. and Braggins, T.: Comparison of methods to prevent restrictor plugging during off-line supercritical extraction. *J. Chromatogr.*, 609 (1992) 321-332.
- 56 Cecchetti, W., Polloni, R., Bergamasco, G., Seraglia, R., Catinella, S., Cecchinato, F. and Traldi, P.: A new pyrolyzer for laser pyrolysis-gas chromatography/mass spectrometry experiments. *J. Anal. Appl. Pyrolysis*, 23 (1992) 165-173; C.A., 117 (1992) 132674s.
- 57 Cox, J., Kilminster, K., Dilworth, J. and Hosmer, G.: Use of pure nitrogen, air, and hydrogen generators with GC systems. *Am. Lab. (Fairfield)*, 24, No. 13 (1992) 34-35.
- 58 Gerstel, E.: (Evaporation-tube method and device for gas-chromatographic separation.) *Ger. Offen.* DE 4,011,350 (Cl. G01N30/12), 10 Oct. 1991, Appl. 07 Apr. 1990; 7 pp.; C.A., 117 (1992) 103329c.
- 59 Gerstel, E.: Rinsing-flow process and device for gas-chromatographic separation. *Eur. Pat. Appl.* EP 451,566 (Cl. G01N30/16), 16 Oct. 1991, DE Appl. 4,011,350, 07 Apr. 1990; 11 pp.; C.A., 117 (1992) 103330w.
- 60 Grob, K., Biedermann, M., Bernath, K., Neukom, H.-P. and Galli, M.: Call for fused silica tubing furnishing tight press-fit connections. *J. High Resolut. Chromatogr.*, 15 (1992) 613-614.
- 61 Grob, K., Brem, S. and Fröhlich, D.: Splitless injection of up to hundreds of microliters of liquid samples in capillary GC: part 1, concept. *J. High Resolut. Chromatogr.*, 15 (1992) 659-664.
- 62 Khatskevich, E.A.: (System for sampling and introduction of pure gases used for calibration and testing of chromatographs.) *Vysokochist. Veshchestva*, No. 3 (1992) 157-160; C.A., 117 (1992) 183792m.
- 63 Manura, J.J. and Hartman, T.G.: Applications of a short-path thermal desorption GC accessory. *Am. Lab. (Shelton)*, 24, No. 8 (1992) 46-52.
- 64 Matsui, M.: (Gas chromatograph for determination of formaldehyde and formic acid.) *Jpn. Kokai Tokkyo Koho* JP 04,120,462 [92,120,462] (Cl. G01N30/88), 21 Apr. 1992, Appl. 90/240,354, 11 Sep. 1990; 3 pp.; C.A., 117 (1992) 103320t.
- 65 Matsui, M.: (Gas chromatograph for determination of hydrogen and inorganic gases.) *Jpn. Kokai Tokkyo Koho* JP 04 66,868 [92 66,868] (Cl. G01N30/78), 03 Mar. 1992, Appl. 90/178,929, 06 Jul. 1990; 3pp.; C.A., 117 (1992) 39435e.
- 66 Newkirk, M.S. and Human, D.M.: Direct filter injection system for gas chromatograph. *U.S. US* 5,109,710 (Cl. 73-863.11; G01N30/20), 05 May 1992, Appl. 541,816, 21 Jun. 1990; 7 pp.; C.A., 117 (1992) 103323w.
- 67 Ono, Y. and Ito, M.: (Supercritical chromatographic separators.) *Jpn. Kokai Tokkyo Koho* JP 04 94,702 [92 94,702] (Cl. B01D11/00), 26 Mar. 1992, Appl. 90/214,050, 13 Aug. 1990; 5 pp.; C.A., 117 (1992) 51433k.
- 68 Phillips, J.B. and Liu, Z.: Two-dimensional gas-chromatograph technique and apparatus. *U.S. US* 5,135,549 (Cl. 55-67; B01D15/08), 04 Aug. 1992, Appl. 647,790, 30 Jan. 1991; 26 pp.; C.A., 117 (1992) 142627t.
- 69 Poshemanskii, V.M., Orgutsov, V.I. and Skornyakov, E.P.: (Analytical unit of a gas chromatograph.) *U.S.S.R. SU* 1,681,233 (Cl. G01N30/02), 30 Sep. 1991, Appl. 4,746,122, 03 Oct. 1989; C.A., 117 (1992) 184050e.
- 70 Pozhidaeva, K.A., Pzhidaev, V.M. and Korolev, A.A.: (A method for direct sample injection into a capillary column for gas chromatographic analysis of high-boiling substances.) *Zh. Anal. Khim.*, 47 (1992) 1439-1441.
- 71 Qian, J., Polymeropoulos, C.E. and Ulisse, R.: Liquid jet evolution from a gas chromatographic injector. *J. Chromatogr.*, 609 (1992) 269-276.
- 72 Reddy, K.T., Cernansky, N.P. and Cohen, R.S.: GC/on column injection technique to detect dodecyl hydroperoxides and their decomposition products. *Anal. Chem.*, 64 (1992) 2273-2276.
- 73 Rittenhouse, D. and Tait, T.: A new technology for more accurate gas flow rate measurements. *Am. Lab. (Fairfield)*, 24, No. 14 (1992) 32P, 32R, 32S.
- 74 Schlegelmilch, F.: (HSS-2-messsystem zur vollautomatischen headspace-GC.) *LaborPraxis*, 16 (1992) 862-865.
- 75 Seefeld, F. and Kebeilmann, L.: (Arrangement for metering liquid samples from pressurized stream in gas chromatograph.) *Ger. (East) DD* 295,715 (G01N30/04), 07 Nov. 1991, Appl. 302,316, 04 May 1987; 4 pp.; C.A., 117 (1992) 103314u.
- 76 Silvis, P.H., Walsh, J.W. and Shelow, D.M.: Sleeve units for inlet splitters of capillary gas chromatographs. *U.S. US* 5,119,669 (Cl. 73-23.41; G01N30/38), 09 Jun. 1992, Appl. 560,103, 31 Jul. 1990; 7 pp.; C.A., 117 (1992) 82710c.
- 77 Sivchenko, V.Ya., Mal'tsev, E.K., Bychkov, V.G., Kritskii, V.E., Ovcharenko, N.P. and Koval'chuk, V.I.: (Automatic portable gas chromatograph.) *U.S.S.R. SU* 1,702,300 (Cl. G01N30/10), 30 Dec. 1991, Appl. 4,643, 260, 26 Dec. 1988; C.A., 117 (1992) 103317x.
- 78 Song, G.: (Factors affecting the quantitative results in a capillary column split sampling.) *Tianranqi Huagong*, 16, No. 3 (1991) 54-57; C.A., 117 (1992) 184023y.
- 79 Stan'kov, I.N., Tarasov, S.N., Beresnev, A.N. and Morozova, O.T.: (Some features of direct liquid sample injection into a capillary column.) *Zh. Anal. Khim.*, 47 (1992) 522-529.
- 80 Staniewski, J. and Rijks, J.A.: Solvent elimination rate in temperature-programmed injections of large sample volumes in capillary gas chromatography. *J. Chromatogr.*, 623 (1992) 105-113.
- 81 Staniewski, J., Janssen, H.-G., Cramers, C.A. and Rijks, J.A.: Programmed-temperature injector for large-volume sample introduction in capillary gas chromatography and for liquid chromatography-gas chromatography interfacing. *J. Microcolumn Separ.*, 4 (1992) 331-338.
- 82 Vejrosta, J., Ansorgova, A. and Mikesova, M.: (Method for preparation of a sintered glass restrictor for chromatography with a supercritical mobile phase.) *Czech. CS* 267,115 (Cl. G01N30/00), 27 Jul. 1990, Appl. 87/8,276, 18 Nov. 1987; 4 pp.; C.A., 117 (1992) 163106n.

- 83 Wehner, W. and Koehler, T.: (A simple desorption device for gas chromatographic aroma analysis using headspace technique.) *Gartenbauwissenschaft*, 57 (1992) 126-129; *C.A.*, 117 (1992) 169700y.
- 84 Yashin, Ya.I.: (Analytical abilities of chromatographic instruments by Dzerzhinsk's OKBA NPO "Khimavtomatika" for environmental control.) *Zh. Anal. Khim.*, 47 (1992) 1516-1524.
- 85 Zelikman, A.M. and Revel'skii, I.A.: (Device for introducing samples into a gas chromatograph.) *U.S.S.R. SU* 1,681,235 (Cl. G01N30/24), 30 Sep. 1991, Appl. 4,697,652, 31 May 1989; *C.A.*, 117 (1992) 61980a.
- 86 Zelikman, A.M., Merkulov, V.D., Yakushin, V.L., Revel'skii, I.A. and Leenson, F.G.: (Gas chromatograph.) *U.S.S.R. SU* 1,673,951 (Cl. G01N30/02), 30 Aug. 1991, Appl. 4,715,188, 07 Jul. 1989; *C.A.*, 117 (1992) 39571w.

See also 164, 165, 168, 201, 212, 215, 266, 486, 701.

3b. Detectors and detection reagents

- 87 Blanco, C.G., Canga, J.S., Domínguez, A., Iglesias, M.J. and Guillén, M.D.: Flame ionization detection relative response factors of some polycyclic aromatic compounds. Determination of the main components of the coal tar pitch volatile fraction. *J. Chromatogr.*, 607 (1992) 295-302.
- 88 Bortolotto, G., Desages, A. and Romagnoli, J.: Optimal tuning of discrete PID controllers using Lyapunov's direct method robustness analysis. *Chem. Eng. Commun.*, 109 (1991) 1-18; *C.A.*, 116 (1992) 062630c.
- 89 Farzane, N.G., Ilyasov, L.V. and Zakharova, L.V.: (Method for calibration of a detector in gas-chromatographic thermal-conductometric analysis.) *U.S.S.R. SU* 1,679,367 (Cl. G01N30/78), 23 Sep. 1991, Appl. 4,483,409, 15 Sep. 1988; *C.A.*, 117 (1992) 61979g.
- 90 Grimsrud, E.P.: Gas-phase ion chemistry within the electron capture detector. *Mass Spectrom. Rev.*, 10 (1991) 457-489; *C.A.*, 117 (1992) 039505c.
- 91 Hill, H.H. and McMinn, D.: *Detectors for Capillary Chromatography*. Wiley, London, 1992, 464 p.
- 92 Klein, K.J., Larson, P.A. and Breckenridge, J.A.: Optimization of GC detector parameters using electronic pressure control. *J. High Resolut. Chromatogr.*, 15 (1992) 615-619.
- 93 Lechner-Fish, T.J., Cordill, L.D., Cooper, G.W. and Combs, G.L.: Applications and performance of a reduced volume TC detector in process GC. *Am. Lab. (Fairfield)*, 24, No. 11 (1992) 28J, 28L, 28N, 28P, 28R, 28T.
- 94 Park, Y.J., Yoo, H.S. and Lee, N.S.: Atomic emission detector for gas chromatography using cylindrical microwave cavity. *Bull. Korean Chem. Soc.*, 13 (1992) 349-351; *C.A.*, 117 (1992) 163153a.
- 95 Pedersen-Bjergaard, S. and Greibrokk, T.: Evaluation of a radio frequency plasma for oxygen-selective detection in capillary gas chromatography. *J. High Resolut. Chromatogr.*, 15 (1992) 677-681.
- 96 Pedersen-Bjergaard, S., Asp, T.N. and Greibrokk, T.: Determination of sulfur- and chlorine-containing compounds using capillary gas chromatography and atomic emission detection. *Anal. Chim. Acta*, 265 (1992) 87-92.
- 97 Shearer, R.L.: Development of flameless sulfur chemiluminescence detection: application to gas chromatography. *Anal. Chem.*, 64 (1992) 2192-2196.
- 98 Stanley, P.E.: A survey of more than 90 commercially available luminometers and imaging devices for low-light measurements of chemiluminescence and bioluminescence, including instruments for manual, automatic and specialized operation, for HPLC, LC, GLC and microtiter plates. Part 1: descriptions. *J. Biolumin. Chemilumin.*, 7 (1992) 77-108; *C.A.*, 117 (1992) 65714q - a review with 15 refs.
- 99 Sun, X.Y., Millier, B. and Aue, W.A.: Flame photometric detection of some transition metals. I. Calibrations and spectra. *Can. J. Chem.*, 70 (1992) 1129-1142; *C.A.*, 117 (1992) 163077d.
- 100 Vasin, S.V., Wentworth, W.E., Stearns, S.D. and Meyer, C.J.: Pulsed discharge emission detector - application to analytical spectroscopy of permanent gases. *Chromatographia*, 34 (1992) 226-234.
- 101 Wentworth, W.E., Vasin, S.V., Stearns, S.D. and Meyer, C.J.: Pulsed discharge helium ionization detector. *Chromatographia*, 34 (1992) 219-225.
- 102 Ye, F., Lin, C., Luo, C., Pei, Z. and Gao, Z.: (The quantitative performance of a new argon discharge detector.) *Sepu*, 10 (1992) 290-292.
- 103 Yu, A., Xu, W., Wang, F., Wang, X., Liu, Y. and Jin, Q.: A study on microwave induced plasma ionization detector for gas chromatograph. *Chem. Res. Chin. Univ.*, 7 (1991) 243-244; *C.A.*, 117 (1992) 142417z.
- 104 Zhang, Y., Bush, K.W. and Bush, M.A.: Evaluation of an improved burner design for a flame infrared emission (FIRE) gas chromatography detector. *Appl. Spectrosc.*, 46 (1992) 930-939.
- 105 Zheng, Y., Seeley, J.A., Dowling, T.M., Uden, P.C. and Khuahar, M.Y.: Element-selective atomic emission detection for capillary GC of metal-containing compounds. *J. High Resolut. Chromatogr.*, 15 (1992) 669-676.

See also 149, 202, 240, 256, 709, 799.

3c. Sorbents and columns, packing procedures

- 106 Berezkin, V.G. and Viktorova, E.N.: (Utilization of inorganic salts for gas chromatographic separation of unsaturated compounds.) *Zh. Anal. Khim.*, 47 (1992) 835-839.
- 107 Berezkin, V.G., Smiryayeva, V.E. and Popova, T.P.: (Open-tubular columns with a few layers of different immobilized stationary phases.) *Zh. Anal. Khim.*, 47 (1992) 825-831.
- 108 Betts, T.J.: Use of three molecularly toroid phases for the gas chromatography of some volatile oil constituents, and comparison with liquid crystal phases. *J. Chromatogr.*, 606 (1992) 281-284.
- 109 Castello, G. and D'Amato, G.: Fast evaluation of the polarity of gas chromatographic columns using the difference in apparent carbon number of linear alkanes and alcohols with the same retention. *J. Chromatogr.*, 623 (1992) 289-295.
- 110 Davudov, V.Ya., Mandrugina, A.A., Morozova, Yu.G., Roshchina, T.M. and Filatova, G.N.: (Adsorption and gas chromatographic properties of Silochrome with bonded aminopropyl groups.) *Zh. Fiz. Khim.*, 66 (1992) 736-743.

- 111 Egorova, K.V., Pakhomova, V.I., Petrova, E.I. and Vigdergauz, M.S.: (Fluorocarbons as low-temperature stationary phases.) *Izv. Vyssh. Uchebn. Zaved., Khim. Khim. Tekhnol.*, 34, No. 8 (1991) 63-68; C.A., 117 (1992) 39523g.
- 112 Ettre, L.S.: Open-tubular columns: past, present and future. *Chromatographia*, 34 (1992) 513-528; - a review with 125 refs.
- 113 Fu, R., Zhang, A., Ge, J., Deng, J., Jin, Y., Zhu, J., Gu, J., Huang, Z. and Zhang, B.: (The investigation of poly(crown ether)siloxane as a gas chromatographic stationary phase.) *Sepu*, 10 (1992) 198-203.
- 114 Funkenbusch, E.F., Carr, P.W., Hanggi, D.A. and Weber, T.P.: Carbon-clad zirconium oxide particles, their manufacture, and uses. U.S. US 5,108,597 (Cl. 210-198.2; B01D15/18), 28 Apr. 1992, Appl. 497,594, 22 Mar. 1990; 25 pp.; C.A., 117 (1992) 174503r.
- 115 Gromova, G.E., Lazaris, A.Ya., Terman, L.M. and Yul'chevskaya, S.D.: (Sorbents for gas chromatography with immobilized stationary phases.) *Zh. Anal. Khim.*, 47 (1992) 1276-1282.
- 116 Gu, X., Du, Y. and Cao, S.: (Gas chromatographic properties of propylammonium nitrate and isopropylammonium nitrate.) *Hebei Daxue Xuebao, Ziran Kexueban*, 11, No. 3 (1991) 29-33; C.A., 117 (1992) 82672s.
- 117 Guha, O.K. and Roy, J.: Nonpolar adsorbent for gas chromatography from anthracite. *Fuel Sci. Technol.*, 10 (1991) 83-84; C.A., 117 (1992) 142582z.
- 118 Hua, X., Zhang, W., Yang, X. and Li, G.: (Syntheses and properties of high-temperature liquid crystals as stationary phase in gas chromatography. IV. Phenylene bis(4,4'-alkylbiphenylcarboxylate)). *Huadong Huagong Xueyuan Xuebao*, 17 (1991) 700-709; C.A., 117 (1992) 184021w.
- 119 Kang, X.: (Development of wide-bore thick-film glass capillary column crosslinked with diethylene glycol succinate and its application.) *Fenxi Huaxue*, 20 (1992) 538-540; C.A., 117 (1992) 68589p.
- 120 Kang, X. and Zhou, Y.: (Radiation crosslinking of diethylene glycol-succinic acid copolymer glass capillary column with cobalt-60 gamma ray.) *Fenxi Huaxue*, 20 (1992) 39-41; C.A., 117 (1992) 49837b.
- 121 Karabanov, N.T.: (Binary systems in gas chromatography.) *Vysokochist. Veshchestva*, No. 3 (1992) 7-33; C.A., 117 (1992) 183994d - a review with 63 refs.
- 122 Kowalski, W.J.: Free radical crosslinking of the gas chromatographic stationary phase containing europium chelates. *Chromatographia*, 34 (1992) 266-268.
- 123 Kraus, A., Schumann, U., Kraus, G., Kohne, B. and Praefcke, K.: Investigation of a scyllitol-derived (saturated) disc-shaped liquid crystal stationary phase in capillary gas chromatography. *J. Chromatogr.*, 609 (1992) 277-281.
- 124 Leclercq, P.A.: Optimum performance of capillary GC columns as a function of column diameter and film thickness under various operating conditions: Part II: computer program extensions for *n*-alkane solutes and polydimethylsiloxane phases. *J. High Resolut. Chromatogr.*, 15 (1992) 531-534.
- 125 Long, Y., Kang, X. and Huang, T.: A novel derivative of β -cyclodextrin as chiral stationary phase in capillary gas chromatography. *Chin. Chem. Lett.*, 3 (1992) 457-458; C.A., 117 (1992) 184024z.
- 126 Mayer, S., Schmalzing, D., Jung, M. and Schleimer, M.: A chiral test mixture for permethylated β -cyclodextrin-polysiloxane gas-liquid chromatography phases: the Schurig test mixture. *LC-GC*, 10 (1992) 782-785.
- 127 Mdivnishvili, O.M., Uridiya, L.Ya. and Makharadze, L.V.: (Method of preparation of supports based on diatomite for gas-liquid chromatography.) U.S.S.R. SU 1,696,998 (Cl. G01N30/48), 07 Dec. 1991, Appl. 4,697,014, 25 May 1989; C.A., 117 (1992) 103231q.
- 128 Mikhalkin, A.P.: (Stationary phase for gas chromatography.) U.S.S.R. SU 1,003,645 (Cl. G01N30/48), 15 Nov. 1991, Appl. 2,959,157, 18 Jul. 1980; C.A., 117 (1992) 39583b.
- 129 Okana, H., Tokunaga, T., Suzuki, N., Ito, R. and Nagai, H.: (Quartz glass capillary column.) *Jpn. Kokai Tokkyo Koho* JP 04 13,966 [92 13,966] (Cl. G01N30/60), 17 Jan. 1992, Appl. 90/117,987, 07 May 1990; 8 pp.; C.A., 117 (1992) 61883w.
- 130 Platonova, N.P., Ulogov, V.O., Gradil, I. and Shvets, F.: (Gas chromatography on methacrylic polymers modified by cobalt cations.) *Zh. Fiz. Khim.*, 66 (1992) 1043-1048.
- 131 Platonova, N.P., Ulogov, V.O., Hradil, I. and Cvec, F.: (Gas chromatography on methacrylate polymers modified by cobalt cations.) *Zh. Fiz. Khim.*, 66 (1992) 1043-1048; C.A., 117 (1992) 178926d.
- 132 Sandra, P.: Miniaturization of chromatographic columns for the analysis of flavours and fragrances. *Analisis*, 20 (1992) M27-M30.
- 133 Shen, Y., Li, G. and Qin, Y.: (Syntheses and properties of poly(methylsiloxane)-type liquid crystals (PMMS) as gas chromatographic stationary phase. II. PMMSs with 1,4-benzenediol dibenzoate-type side chains.) *Huadong Huagong Xueyuan Xuebao*, 17 (1991) 694-700; C.A., 117 (1992) 172231b.
- 134 Shen, Y., Li, G., Qin, Y. and Zhang, W.: (Syntheses and properties of poly(methylsiloxane)-type liquid crystals (PMMS) as gas chromatographic stationary phase. I. PMMSs with biphenyl benzoate-type side chains.) *Huadong Huagong Xueyuan Xuebao*, 17 (1992) 687-694; C.A., 117 (1992) 172238j.
- 135 Sun, J.J. and Fritz, J.S.: Use of mercuric resins for selective abstraction of mercaptans in capillary gas chromatography. *J. High Resolut. Chromatogr.*, 15 (1992) 605-608.
- 136 Takeichi, T., Shimura, S., Toriyama, H., Takayama, Y. and Morikawa, M.: Carbamate derivatives of β -cyclodextrin as chiral stationary phases for metal capillary gas chromatography. *Chem. Lett.*, (1992) 1069-1072; C.A., 117 (1992) 112015z.
- 137 Volkov, S.A., Zel'venskii, V.Yu. and Smol'yaninov, G.A.: (Influence of thermal modification and properties of solid support on the characteristics of gas chromatographic adsorbents based on a poly(ethylene glycol).) *Zh. Anal. Khim.*, 47 (1992) 1442-1447.
- 138 Volpe, N. and Siouffi, A.-M.: Some questions for chromatographers. *Chromatographia*, 34 (1992) 213-218.
- 139 Zanozina, I.I., Akopova, O.B., Grigor'ov, V.V. and Diskina, D.E.: (Liquid-crystalline phase for chromatographic analysis of diarylalkanes.) *Khim. Tekhnol. Top. Masol*, 10 (1991) 34; C.A., 116 (1992) 024248b.
- 140 Zhang, G., Liu, J. and Gao, Y.: (Flexible glass capillary column modified by silicon film with strongly polar chromatography stationary phase.) *Fenxi Huaxue*, 20 (1992) 660-662; C.A., 117 (1992) 98219k.

- 141 Zheng, Z., Wang, C., Wu, C., Huang Z. and Wang, Y.: (Preparation and separation mechanism of glass capillary column with dicarboxyl crown ether stationary phase.) *Gaodeng Xuexiao Huaxue Xuebao*, 13 (1992) 752-755; C.A., 117 (1992) 142586d.
- 142 Zibarev, P.V., Ryzhova, G.L. and Salamatin, B.V.: (Method of producing a polymeric sorbent for gas chromatography by polymerizing acrylonitrile.) U.S.S.R. SU 1,236,739 (Cl. C08F257/02), 30 Dec. 1991, Appl. 3,575,433, 12 Jan. 1983; C.A., 117 (1992) 123813j.
- 143 Zuo, Y., Liu, S., Li, L. and Wang, Y.: (New ester-containing porous polymeric beads as gas chromatographic stationary phase.) *Gaodeng Xuexiao Huaxue Xuebao*, 13 (1992) 65-67; C.A., 117 (1992) 123765v.

See also 26, 43, 184, 191, 207, 208, 210, 216, 222, 227, 230, 238, 253, 254, 293, 356, 442, 681, 798.

3d. Quantitative analysis

- 144 Astakhov, A.V., Shadrin, A.M. and Glazyrin, E.M.: (Method of quantitative chromatographic analysis measuring the amount of the sample using an ionization detector.) U.S.S.R. SU 1,693,538 (Cl. G01N30/78), 23 Nov. 1991, Appl. 4,717,275, 11 Jul. 1989; C.A., 117 (1992) 103332y.
- 145 Kou, D., Yun, X., Li, Y. and Lu, X.: (Addition quantitative method in gas chromatography.) *Fenxi Huaxue*, 20 (1992) 597-599; C.A., 117 (1992) 103293m.
- 146 Li, S., He, D., He, Q. and Chen, X.: (Experimental study on linear rules between retention time and parameters of chromatographic peak.) *Sepu*, 10 (1992) 192-195.
- 147 MacTaggart, D.L. and Farwell, S.O.: Analytical use of linear regression. Part I: regression procedures for calibration and quantitation. *J. Assoc. Off. Anal. Chem.*, 75 (1992) 594-608.
- 148 MacTaggart, D.L. and Farwell, S.O.: Analytical use of linear regression. Part II: statistical error in both variables. *J. Assoc. Off. Anal. Chem.*, 75 (1992) 608-614.
- 149 Morvai, M., Pályka, I. and Molnár-Perl, I.: Flame ionization detector response factors using the effective carbon number concept in the quantitative analysis of esters. *J. Chromatogr. Sci.*, 30 (1992) 448-452.
- 150 Wei, W., Wu, N.S. and Jiang, X.H.: Short communication. New approach for area determination of an overlapped pair of chromatographic peaks. *J. Chromatogr.*, 623 (1992) 366-370.

See also 154, 259, 388, 510.

3e. Preparative scale chromatography

See 473.

3f. Programmed temperature, pressure, vapors, gradients

See 80.

4. SPECIAL TECHNIQUES

4a. Automation

- 151 Valente, A.L.P. and Augusto, F.: Gas chromatography: micro-computer program for calculation of the van Deemter equation. *Lab. Microcomput.*, 11 (1992) 21-29; C.A., 117 (1992) 34269u.
- 152 Wheeler, S. and Schubert, A.: Benefits of a chromatography carrier gas cylinder management program. *Am. Lab. (Fairfield)*, 24, No. 13 (1992) 32.
- 153 Xu, G., Li, H., Di, L., Liu, W., Zhang, Y. and Lu, P.: (The software for the recommendation of column system in expert system on gas chromatography V. The chromatogram base for verifying developed chromatographic methods.) *Sepu*, 10 (1992) 257-260.

See also 74.

4b. Computerization and modelling

- 154 Koch, C.A.: A smart autosampler for gas chromatography-mass spectrometry evolves. *LC-GC Int.*, 10 (1992) 709.
- 155 Mason, J.P., Kirk, I., Windsor, C.G., Tipler, A., Spragg, R.A. and Rendle, M.: A novel algorithm for chromatogram matching in qualitative analysis. *J. High Resolut. Chromatogr.*, 15 (1992) 539-547.
- 156 Nelson, D.C., Daucinas, V., Lysakowski, R. and Duff, M.: The standard for chromatography data communication from the analytical instruments association. *Am. Lab. (Fairfield)*, 24, No. 14 (1992) 52-57.
- 157 Yang, M.J., Pawliszyn, B. and Pawliszyn, J.: Optimization of calculation parameters and experimental conditions for multiplex gas chromatography using computer simulation. *J. Chromatogr. Sci.*, 30 (1992) 306-314.

See also 124.

4c. Combination with other physico-chemical techniques (MS, IR etc.)

- 158 Arpino, P.J.: Combined supercritical fluid chromatography/mass spectrometry (SFC/MS). *NATO ASI Ser., Ser. C*, 353 (1992) 269-280; C.A., 117 (1992) 142377m - a review with 55 refs.
- 159 Chung, R.J. and Aldridge, J.C.: On-line thermogravimetry/gas chromatography/mass spectrometry. *Anal. Instr.*, 20 (1992) 123-135.
- 160 Colby, B.N.: Spectral deconvolution for overlapping GC/MS components. *J. Am. Soc. Mass Spectrom.*, 3 (1992) 558-562; C.A., 117 (1992) 184030y.
- 161 Cole-Clarke, P. and Powell, J.: GC-FTIR of environmental compounds. *Chem. Aust.*, 59 (1992) 259-260; C.A., 117 (1992) 82657r.
- 162 Rubino, F.M. and Zecca, L.: Application of triple quadrupole tandem mass spectrometry to the analysis of pyridine-containing derivatives of long-chain acids and alcohols. *J. Chromatogr.*, 579 (1992) 1-12.
- 163 Sandra, P. and David, F.: Hyphenated techniques in capillary gas chromatography. *Chim. Oggi*, 10 (1992) 17-25; C.A., 117 (1992) 39500x - a review with 8 refs.

- 164 Sinha, M.P. and Gutnikov, G.: Role of a microbore capillary column in the development of a portable gas chromatograph-mass spectrometer. *J. Microcolumn Separ.*, 4 (1992) 405-410.
- 165 Snyder, A.P., Harden, C.S., Brittain, A.H., Kim, M.-G., Arnold, N.S. and Meuzelaar, H.L.C.: Portable, hand-held gas chromatography-ion mobility spectrometer. *Am. Lab. (Fairfield)*, 24, No. 15 (1992) 32B, 32D, 32E, 32F, 32H.
- 166 Torrent, A.: Les couplages en spectrométrie de masse. *Analisis*, 20 (1992) M66-M70.
- See also 2, 96, 179, 246, 258, 375, 419, 479, 484, 494, 680, 715, 750.
- 4f. *Trace analysis and pre-separation techniques*
- 167 Arthur, C.L., Potter, D.W., Buchholz, K.D., Motlagh, S. and Pawliszyn, J.: Solid-phase microextraction for the direct analysis of water: theory and practice. *LC-GC Int.*, 5, No. 10 (1992) 8-14.
- 168 Ashraf-Khorassani, M., Houck, R.K. and Levy, J.M.: Cryogenically cooled adsorbent trap for off-line supercritical fluid extraction. *J. Chromatogr. Sci.*, 30 (1992) 361-366.
- 169 Furton, K.G. and Rein, J.: Relative effect of experimental variables. Analytical supercritical fluid extraction efficiencies and correlations with chromatographic data. *ACS Symp. Ser.*, 488 (1992) 237-250; *C.A.*, 117 (1992) 61948w.
- 170 Ishihara, K. and Honma, N.: A novel device for the headspace trapping of volatiles: studies with a model solution. *J. Food Compos. Anal.*, 5 (1992) 139-145; *C.A.*, 117 (1992) 130032g.
- 171 Kapila, S., Nam, K.S., Liu, M.H., Puri, R.K. and Yanders, A.F.: Promises and pitfalls of supercritical fluid extraction in polychlorinated compound analyses. *Chemosphere*, 25 (1992) 11-16.
- 172 Lebo, J.A., Zajicek, J.L., Huckins, J.N., Petty, J.D. and Peterman, P.H.: Use of semipermeable membrane devices for in situ monitoring of polycyclic aromatic hydrocarbons in aquatic environments. *Chemosphere*, 25 (1992) 697-718.
- 173 Manninen, P. and Riekkola, M.L.: (Supercritical fluid extraction: a new method for sample preparation.) *Kem.-Kemi*, 19 (1992) 417-421; *C.A.*, 117 (1992) 142569a - a review with 62 refs.
- 174 Martin, F., Verdejo, T. and González-Vila, F.J.: Extraction of bituminous material from fossil organic matter using liquid carbon dioxide under liquid-vapour equilibrium conditions. *J. Chromatogr.*, 607 (1992) 377-379.
- 175 Mulcahey, L.J. and Tayler, L.T.: Collection efficiency of solid surface and sorbent traps in supercritical fluid extraction with modified carbon dioxide. *Anal. Chem.*, 64 (1992) 2352-2358.
- 176 Potter, D.W. and Pawliszyn, J.: Detection of substituted benzenes in water at the pg/ml level using solid-phase microextraction and gas chromatography-ion trap mass spectrometry. *J. Chromatogr.*, 625 (1992) 247-255.
- 177 Sunesson, A.-L., Nilsson, C.-A., Andersson, B. and Carlson, R.: Thermal desorption cold trap-injection in high-resolution gas chromatography: multivariate optimization of experimental conditions. *J. Chromatogr.*, 623 (1992) 93-103.
- 178 Terada, M., Yamamoto, T., Kuroiwa, Y. and Wakasugi, C.: Gel permeation chromatographic clean-up prior to determination of barbiturates in blood extract by gas chromatography. *Hochudoku*, 10, No. 1 (1992) 31-37; *C.A.*, 117 (1992) 84781a.
- 179 Vo, A.: Use of a centrifugal evaporator to reduce emissions of solvents used in EPA method 3510 prior to GC-MS Analysis. *J. High Resolut. Chromatogr.*, 15 (1992) 552-555.
- 180 Vreuls, J.J., Ghijsen, R.T., de Jong, G.J. and Brinkman, U.A.Th.: Drying step for introduction of water-free desorption solvent into a gas chromatograph after on-line liquid chromatographic trace enrichment of aqueous samples. *J. Chromatogr.*, 625 (1992) 237-245.
- 181 Wallace, J.C., Krieger, M.S. and Hites, R.A.: Reduction of contamination levels in on-line supercritical fluid extraction systems. *Anal. Chem.*, 64 (1992) 2655-2656.
- 182 Wang, H., Yu, B. and Jiang, W.: (Analysis of volatile organic pollutants with Curie point desorption gas chromatographic method. Part (1). Feasibility of using adsorption wire sampling.) *Huanjing Wuran Yu Fangzhi*, 14, No. 1 (1992) 35-37; *C.A.*, 117 (1992) 54862k.
- 183 Wenclawiak, B., Hoffmann, A. and Bogusz, M.: Supercritical fluid extraction of biological materials. *Fresenius J. Anal. Chem.*, 343 (1992) 116-117.
- See also 53, 135, 251, 354, 497, 502, 512, 521, 531, 558, 710, 716, 724, 726, 800.
- 4g. *Enantiomers, separation*
- 184 Bicchi, C., Artuffo, G., D'Amato, A., Galli, A. and Galli, M.: Cyclodextrin derivatives in the GC separation of racemic mixtures of volatile compounds: part IV. *J. High Resolut. Chromatogr.*, 15 (1992) 655-658.
- 185 Bornscheuer, U., Schapöhler, S., Scheper, T., Schügerl, K. and König, W.A.: Application of enantioselective capillary gas chromatography in lipase-catalysed transesterification reactions in organic media. *J. Chromatogr.*, 606 (1992) 288-290.
- 186 Buser, H.R., Mueller, M.D. and Rappe, C.: Enantioselective determination of chlordane components using chiral high-resolution gas chromatography-mass spectrometry with application to environmental samples. *Environ. Sci. Technol.*, 26 (1992) 1533-1540; *C.A.*, 117 (1992) 84717j.
- 187 Davin, L.B., Umezawa, T. and Lewis, N.G.: Evantioselective separations in phytochemistry. *Recent Adv. Phytochem.*, 25 (1991) 75-112; *C.A.*, 116 (1992) 170047b.
- 188 De Vries, N.K., Coussens, B. and Meier, R.J.: The separation of enantiomers on modified cyclodextrin columns: measurements and molecular modeling. *J. High Resolut. Chromatogr.*, 15 (1992) 499-504.
- 189 Fellous, R. and Derbesy, M.: Chromatographie gazeuse chirale et authenticité des arômes. *Analisis*, 20 (1992) M31-M33.
- 190 Koepke, T. and Mosandl, A.: (Stereoisomeric flavor compounds. Part 54. 8-Mercapto-p-menthan-3-one - optically pure stereoisomers and chiro-specific analysis.) *Z. Lebensm.-Unters. Forsch.*, 194 (1992) 372-376; *C.A.*, 117 (1992) 90516j.
- 191 Li, W.: Enantiomeric separation by capillary gas chromatography using derivatized cyclodextrin stationary phases. Avail. *Univ. Microfilms Int.*, Order No. DA9133826, 1991, 148pp.; *C.A.*, 117 (1992) 142594e.
- 192 Ojala, T.M., Hase, A. and Komppa, V.: (Supercritical chiral chromatography. A resolution technique for the analytical laboratory and production.) *Kem. - Kemi*, 19 (1992) 422-424; *C.A.*, 117 (1992) 142570u.

- 193 Pietruszka, J., Hochmuth, D.H., Gehrcke, B., Icheln, D., Runge, T. and Koenig, W.A.: Gas chromatographic enantioseparation of allenenes. *Tetrahedron: Asymmetry*, 3 (1992) 661-670; *C.A.*, 117 (1992) 69479q.
- 194 Schubert, V. and Mosandl, A.: (Stereoisomeric flavor compounds. Part 56. (Z)-4-Hepton-2-ol and its esters. Structure and properties of the enantiomers.) *Z. Lebensm.-Unters. Forsch.*, 194 (1992) 552-555; *C.A.*, 117 (1992) 168424z.
- See also 125, 126, 136, 217, 225, 316, 331, 393, 396, 438, 442, 494, 505, 570, 576, 599, 617, 681.
- 4h. *Other special techniques*
- 195 Burioko, A., Kaliurand, M., Marinichev, A.N. and Stolyarov, B.: Headspace analysis of rapid reaction systems by stroboscopic sampling. *Eesti Tead. Akad. Toim., Keem.*, 40 (1991) 157-165; *C.A.*, 116 (1992) 020495g.
- 196 Niotis, A. and Katsanos, N.A.: Hydrodenitrogenation catalysis studied by reversed-flow gas chromatography. *Chromatographia*, 34 (1992) 398-410.
- 197 Peremitin, I.V., Peremitina, V.T. and Ogloblina, I.P.: (Gas-chromatography method for analysis of a mixture of organic substances.) *U.S.S.R. SU 1,679,360 (Cl. G01N30/02)*, 23 Sep. 1991, *Appl.* 4,667,152, 19 Jan. 1989; *C.A.*, 117 (1992) 61995j.
- See also 357.
- 4i. *Supercritical fluid chromatography*
- 198 Ashraf, S., Bartle, K.D., Clifford, A.A. and Moulder, R.: The analysis of lubricating oil additives by supercritical fluid chromatography with packed and open tubular capillary columns. *J. High Resolut. Chromatogr.*, 15 (1992) 535-538.
- 199 Barbour, F.A. and Branthaver, J.F.: Analysis of SHRP asphalt nonpolar fractions by supercritical fluid chromatography. *Fuel Sci. Technol. Int.*, 10 (1992) 979-1001; *C.A.*, 117 (1992) 96250q.
- 200 Camel, V., Thiebaut, D. and Caude, M.: Supercritical fluid chromatography and extraction. *Analisis*, 20, No. 8 (1992) M18-M21.
- 201 Campbell, R.M., Cortes, H.J. and Green, L.S.: Large-volume injection in capillary supercritical fluid chromatography. *Anal. Chem.*, 64 (1992) 2852-2857.
- 202 Carey, J.M. and Caruso, J.A.: Plasma spectrometric detection for supercritical fluid chromatography. *TrAC*, 11 (1992) 287-293.
- 203 Edder, P., Haerdi, W., Veuthey, J.L. and Staub, C.: Quantitative capillary supercritical fluid chromatography and supercritical fluid extraction of basic drugs of abuse. *Chimia*, 46 (1992) 141-148.
- 204 Foley, J.P. and Crow, J.A.: Systematic multiparameter strategies for optimizing supercritical fluid chromatographic separations. *ACS Symp. Ser.*, 488 (1992) 304-335; *C.A.*, 117 (1992) 39506d - a review with 59 refs.
- 205 Fujimoto, C., Suzuki, M. and Jinno, K.: Planarity selectivity in supercritical fluid and liquid chromatography using octadecylsilane-modified silicas with carbon dioxide. *Chromatographia*, 34 (1992) 121-124.
- 206 Hedrick, J.L., Mulcahey, L.J. and Taylor, L.T.: Supercritical fluid extraction. *Mikrochim. Acta*, 108 (1992) 115-132 - a review with 39 refs.
- 207 Hütz, A. and Klesper, E.: Efficiency in supercritical fluid chromatography as a function of linear velocity, pressure/density, temperature and diffusion coefficient employing *n*-pentane as the eluent. *J. Chromatogr.*, 607 (1992) 79-89.
- 208 Janák, K., Demirbüker, M., Häggglund, I. and Blomberg, L.G.: Modifications of poly(methyl-3-propylthio)siloxane to give stationary phases for open tubular supercritical fluid chromatography. *Chromatographia*, 34 (1992) 335-341.
- 209 Jinno, K. (Editor): *Hyphenated Techniques in Supercritical Fluid Chromatography and Extraction*. Toyohashi University of Technology, Toyohashi, 1992, x + 334p.
- 210 Karlsson, L., Buttlar, T. and Mathiasson, L.: Retention and peak shape of acidic substances in capillary supercritical fluid chromatography with binary and ternary mobile phases. *J. Microcolumn Separ.*, 4 (1992) 423-430.
- 211 Kornfeld, R.: Reliable analytical determinations using supercritical fluid chromatography. *Am. Lab. (Fairfield)*, 24, No. 14 (1992) 25-32.
- 212 Koski, I.J., Markides, K.E., Richter, B.E. and Lee, M.L.: Microliter sample introduction for open tubular column supercritical fluid chromatography using a packed capillary for solute focusing. *Anal. Chem.*, 64 (1992) 1669-1675.
- 213 Langenfeld, J.J., Hawthorne, S.B., Miller, D.J. and Tehrani, J.: Method for determining the density of pure and modified supercritical fluids. *Anal. Chem.*, 64 (1992) 2263-2266.
- 214 Levy, J.M., Storozynsky, E. and Ashraf-Khorassani, M.: Use of modifiers in supercritical fluid extraction. In: Braight, F.V. and McNally, M.E.P. (Editors), *ACS Symp. Ser. 488, 1992*, ACS, Washington, 1992, 336-361; *C.A.*, 117 (1992) 184020v.
- 215 Pavlenko, I.V., Romanov, F.I. and Ryaboi, I.V.: (Supercritical fluid chromatography.) *U.S.S.R. SU 1,681,236 (Cl. G01N30/60)*, 30 Sep. 1991, *Appl.* 4,752,510, 24 Oct. 1989; *C.A.*, 117 (1992) 82706f.
- 216 Pesek, J.J. and Williamsen, E.J.: The potential uses of bonded liquid crystal materials in high-performance liquid chromatography and supercritical-fluid chromatography. *TrAC*, 11 (1992) 259-266.
- 217 Petersson, P., Lundell, N. and Markides, K.E.: Chiral separations in supercritical fluid chromatography: a multivariate optimization method. *J. Chromatogr.*, 623 (1992) 129-137.
- 218 Poe, D.P.: Model for retention and efficiency in open-tubular supercritical fluid chromatography. *J. Chromatogr.*, 625 (1992) 299-309.
- 219 Pyo, D. and Hwang, H.: Analysis of insecticides and fungicides employing supercritical carbon dioxide and modifiers. *Bull. Korean Chem. Soc.*, 13 (1992) 110-112; *C.A.*, 117 (1992) 21824w.
- 220 Roth, M.: Statistical thermodynamic treatment of high-pressure phase equilibria in supercritical fluid chromatography. *J. Phys. Chem.*, 96 (1992) 8548-8552.
- 221 Roth, M.: Statistical thermodynamic treatment of high-pressure phase equilibria in supercritical fluid chromatography. 2. Parameterization and testing of the model. *J. Phys. Chem.*, 96 (1992) 8552-8556.
- 222 Tong, D., Bartle, K.D., Clifford, A.A. and Robinson, R.E.: Unified gas and supercritical fluid chromatography on 50 μm i.d. columns. *J. High Resolut. Chromatogr.*, 15 (1992) 505-509.

- 223 Vérrillon, F., Heems, D., Pichon, B., Coleman, K. and Robert, J.-C.: Supercritical fluid chromatography with independent programming of mobile-phase pressure, composition, and flow rate. *Am. Lab. (Fairfield)*, 24, No. 9 (1992) 45-53 - a review with 17 refs.
- 224 Yan, C. and Martire, D.E.: Molecular theory of chromatography for blocklike solutes in isotropic stationary phases and its application to supercritical fluid chromatographic retention of PAHs. *J. Phys. Chem.*, 96 (1992) 7510-7517.
- See also 4, 46, 50, 55, 67, 82, 138, 158, 173, 175, 181, 183, 192, 246, 264, 278, 288, 374, 375, 426, 473, 551, 555, 557, 576, 596, 714, 731, 745, 780.
5. HYDROCARBONS AND HALOGEN DERIVATIVES
- 5a. *Aliphatic hydrocarbons*
- 225 Engewald, W., Reinhardt, R., Steinborn, A., Rohloff, J. and Herzschuh, R.: Gas chromatographic behaviour of isomeric tricyclo [10.4.0.0.4.9]hexadecanes. *Chromatographia*, 34 (1992) 253-258.
- 226 Ko, J., Nguyen, J. and Burleson, J.: Determination and confirmation of *cis*-9-tricosene in technical materials by capillary gas chromatography and gas chromatography/mass spectrometry. *J. Assoc. Off. Anal. Chem.*, 75 (1992) 878-882.
- 227 Soják, L., Ostrovsky, I., Kubinec, R., Kraus, G. and Kraus, A.: Separation and identification of all isomeric *n*-nonadecenes by capillary gas chromatography on a mesogenic stationary phase with Fourier transform infrared and mass spectrometric detection. *J. Chromatogr.*, 609 (1992) 283-288.
- 228 Tanaka, T.: (Simultaneous analysis of hydrocarbons and chlorinated hydrocarbons in air by using dual capillary columns.) *Kankyo Kagaku*, 2 (1992) 187-193; *C.A.*, 117 (1992) 117335p.
- 229 Östermark, U. and Petersson, G.: Assessment of hydrocarbons in vapours of conventional and alkylate-based petrol. *Chemosphere*, 25 (1992) 763-768.
- See also 24, 665, 777, 814, 815.
- 5b. *Cyclic hydrocarbons*
- 230 Andrews, A.R.J., Wu, Z. and Zlatkis, A.: The GC separation of C₂-naphthalene isomers using cyclodextrins and crown ethers. *Chromatographia*, 34 (1992) 163-165.
- 231 Angerer, J. and Hörsch, B.: Determination of aromatic hydrocarbons and their metabolites in human blood and urine. *J. Chromatogr.*, 580 (1992) 229-255 - a review with 150 refs.
- 232 Arey, J., Atkinson, R. and Zielinska, B.: GC/MS analysis of polycyclic aromatic hydrocarbons (PAH) and nitro-PAH in ambient air samples using high resolution GC/MS. In: Jennings, W.G. and Nikelly, J.G. (Editors), *Capillary Chromatography*, Huethig, Heidelberg, 1991, pp. 95-107; *C.A.*, 116 (1992) 026895].
- 233 Canton, L. and Grimalt, J.O.: Gas chromatographic-mass spectrometric characterization of polycyclic aromatic hydrocarbon mixtures in polluted coastal sediments. *J. Chromatogr.*, 607 (1992) 279-286.
- 234 Deshpande, L.S. and Sarin, R.: Development of gas-liquid technique for estimation of airborne polynuclear aromatic hydrocarbons hazardous to human health. *Asian J. Chem. Rev.*, 2 (1991) 136-149; *C.A.*, 117 (1992) 136644e.
- 235 Hartgers, W.A., Damste, J.S.S. and Leeuw, J.W.: Identification of C₂-C₄ alkylated benzenes in flash pyrolysates of kerogens, coals and asphaltenes. *J. Chromatogr.*, 606 (1992) 211-220.
- 236 Nazarova, V.I., Petrova, R.S. and Shcherbakova, K.D.: (Chromatostuctural analysis of fenalene molecule and products its hydrogenation.) *Zh. Fiz. Khim.*, 66 (1992) 325-328.
- 237 Oda, J. and Ichikawa, S.: (Analysis of polycyclic aromatic hydrocarbons in airborne particulates by gas chromatography/mass spectrometry and identifications of unknown mass spectra.) *Kankyo Kagaku*, 2 (1992) 402-403; *C.A.*, 117 (1992) 117336q.
- 238 Wang, H., Gen, T., Fu, R., Zhu, J., Wei, X., Li, H. and Liu, J.: Gas chromatographic separation of polycyclic aromatic hydrocarbons on a mesomorphic copolysiloxane stationary phase. *J. Chromatogr.*, 609 (1992) 414-418.
- See also 87, 139, 172, 224, 426, 559.
- 5c. *Halogen derivatives*
- 239 Badawy, M.I.: Evaluation of head space technique for the determination of trihalomethanes in water. *Bull. Environ. Contam. Toxicol.*, 48 (1992) 630-634; *C.A.*, 116 (1992) 158456b.
- 240 Bognar, J.A., Knighton, W.B. and Grimsrud, E.P.: Selective detection of brominated hydrocarbons by the photodetachment-modulated electron capture detector. *Anal. Chem.*, 64 (1992) 2451-2455.
- 241 Braby, M.F.: Sorption of methallyl chloride on wheat. *J. Stored Prod. Res.*, 28 (1992) 183-186; *C.A.*, 117 (1992) 165838b.
- 242 Brown, R.S., Pettit, K., Price, D. and Jones, P.W.: Thermal desorption gas chromatography: a quick screening technique for polychlorinated biphenyls. *Organohalogen Compd.*, 2 (1990) 117-120; *C.A.*, 117 (1992) 177704t.
- 243 Che, Z. and Gao, M.: (Gas chromatography/mass spectrometry (GC/MS) determination of impurities in the used chlorohydrocarbons.) *Sepu*, 10 (1992) 302-303.
- 244 Döring, J., Damberg, M., Gamradt, A. and Oehme, M.: Screening method based on the determination of perchlorinated aromatics for surface soil contaminated by copper slag containing high levels of polychlorinated dibenzofurans and dibenzo-*p*-dioxins. *Chemosphere*, 25 (1992) 755-762.
- 245 Frank, W., Frank, H. and Englert, R.: Automated analysis of C₁- and C₂-halocarbons in ambient air. *Organohalogen Compd.*, 2 (1990) 401-404; *C.A.*, 117 (1992) 177227q.
- 246 Fuoco, R. and Griffiths, P.R.: Analysis of polychlorobiphenyls (PCBs) in contaminated soil samples by supercritical fluid extraction/supercritical fluid chromatography coupled with FT-IR spectrometry. *Ann. Chim. (Rome)*, 82 (1992) 235-245; *C.A.*, 117 (1992) 150025k.
- 247 Gehring, D.G., Barsotti, D.J. and Gibbon, H.E.: Chlorofluorocarbons alternatives analysis, Part II: the determination of HCFC-141b purity by gas chromatography. *J. Chromatogr. Sci.*, 30 (1992) 301-305.

- 248 Gerbino, T.C., Castello, G. and D'Amato, G.: Gas chromatographic identification of halogenated hydrocarbons by using the correlation between their retention and boiling points. *J. Chromatogr.*, 609 (1992) 289-296.
- 249 Hale, R.C. and Greaves, J.: Methods for the analysis of persistent chlorinated hydrocarbons in tissues. *J. Chromatogr.*, 580 (1992) 257-278 - a review with 142 refs.
- 250 Hasegawa, A.: (Determination of dichlorodifluoromethane in air by using adsorbent collection/solvent extraction/gas chromatography.) *Kanagawa-ken Kankyo Kagaku Senta Kenkyu Hokoku*, 14 (1992) 12-15; *C.A.*, 117 (1992) 75397c.
- 251 Jiménez, B., Tabera, J., Hernández, L.M. and González, M.J.: Simplex optimization of the analysis of polychlorinated biphenyls. Application to the resolution of a complex mixture of congeners of interest on a single gas chromatographic column. *J. Chromatogr.*, 607 (1992) 271-278.
- 252 Kannan, N., Schulz-Bull, D.E., Petrick, G. and Duinker, J.C.: High resolution PCB analysis of Kanechlor, Phenoclor and Sovol mixtures using multidimensional gas chromatography. *Intern. J. Environ. Anal. Chem.*, 47 (1992) 201-215.
- 253 Larsen, B., Bwadt, S. and Facchetti, S.: Separation of toxic congeners from PCB mixtures on two series coupled narrow-bore columns (50 m SIL-8 and 25 m HT-5). *Intern. J. Environ. Anal. Chem.*, 47 (1992) 147-166.
- 254 Larsen, B., Bwadt, S. and Tilio, R.: Congener specific analysis of 140 chlorobiphenyls in technical mixtures on five narrow-bore GC columns. *Intern. J. Environ. Anal. Chem.*, 47 (1992) 47-68.
- 255 Larsen, B., Pelusio, F., Skej, H. and Paya-Perez, A.: Bioavailability of polychlorinated biphenyl congeners in the soil to earthworm (*L. rubellus*) system. *Intern. J. Environ. Anal. Chem.*, 46 (1992) 149-162.
- 256 Ohno, H. and Aoyama, T.: Simultaneous determination of volatile chlorinated hydrocarbons by dual detection using a semi-wide bore capillary column. *Eisei Kagaku*, 37 (1991) 387-394; *C.A.*, 116 (1992) 066760t.
- 257 Petrick, G., Schulz-Bull, D.E. and Duinker, J.C.: (Analyses for polychlorinated biphenyls and organic contaminants in drinking water.) *Z. Wasser- Abwasser-Forsch.*, 25 (1992) 115-121; *C.A.*, 117 (1992) 157274z.
- 258 Petrik, J. and Kocan, A.: (Isomer-specific analysis of technical mixtures of polychlorinated biphenyls by HRGC/MS.) *Chem. Listy*, 86 (1992) 694-698.
- 259 Pocklington, W.D.: Determination of tetrachloroethylene - results of a collaborative study and the standardized method. *J. Am. Oil Chem. Soc.*, 69 (1992) 789-793.
- 260 Thomas, V.N. and Ramstad, T.: A dynamic headspace GC method for the determination of vinyl chloride monomer in stored solutions of cefmetazole sodium in PVC bags. *Acta Pharm. Nord.*, 4 (1992) 97-104; *C.A.*, 117 (1992) 97429s.
- 261 Ueta, T., Nakamura, H. and Akiyama, K.: (Determination of three non-ortho-chlorine substituted PCBs in fish and shellfish by GC/MS selected ion monitoring method.) *Kenkyu Nenpo - Tokyo-toritsu Eisei Kenkyusho*, 42 (1991) 95-101; *C.A.*, 117 (1992) 149594p.
- 262 Welling, L., Paukku, R. and Mäntykoski, K.: PCB in recycled paper products. *Chemosphere*, 25 (1992) 293-295.
- 263 Wells, D.E. and Echarrí, I.: Determination of individual chlorobiphenyls (CBs), including non-ortho, and mono-ortho chlorosubstituted CBs in marine mammals from Scottish waters. *Intern. J. Environ. Anal. Chem.*, 47 (1992) 75-97.
- See also 167, 228, 300, 542, 558, 662, 675, 697, 698, 722, 724, 726, 732, 736, 743, 751, 756, 791.
- 5d. *Complex hydrocarbon mixtures (incl. analysis of tars, bitumens and mineral oils)*
- 264 Ashraf-Khorassani, M., Levy, J.M. and Dolata, L.A.: SFC determination of saturates and aromatics in petroleum streams. *Am. Lab. (Shelton)*, 24, No. 8 (1992) 29-35.
- 265 Li, Y.: (Improved chromatographic method for separating C₁-C₈ fractions.) *Shiyu Huagong*, 21 (1992) 112-116; *C.A.*, 117 (1992) 30105q.
- 266 Yasui, S.: (Method for determination of hydrocarbons and a gas chromatograph-flame ionization detector apparatus for this method.) *Jpn. Kokai Tokkyo Koho* JP 04,110,768 [92,110,768] (Cl. G01N30/88), 13 Apr. 1992, Appl. 90/231,409, 31 Aug. 1990; 3 pp.; *C.A.*, 117 (1992) 103341a.
- See also 697, 707, 750.
6. ALCOHOLS
- 267 Alugupalli, S., Mielniczuk, Z. and Larsson, L.: Gas chromatography-mass spectrometry methods for analysis of secondary alcohols present in the *Mycobacterium avium* complex. *J. Microbiol. Methods*, 15 (1992) 229-240; *C.A.*, 117 (1992) 127715b.
- 268 Boneva, S. and Kotov, S.: Capillary gas chromatography of some telomeric fluoroalkanols. *Chromatographia*, 34 (1992) 475-478.
- 269 Goto, K. and Ikemoto, S.: Analysis of ethyl alcohol concentration in blood using gas chromatograph equipped with blood gas sampler. *Nippon Hoigaku Zasshi*, 45 (1991) 393-399; *C.A.*, 117 (1992) 64338h.
- 270 Yu, S., Yu, Y., Jin, S. and Yang, J.: (Determination of the sex pheromone of *Procreas venosatus* with capillary GC/MS method.) *Sepu*, 9 (1991) 185-187.
- See also 16, 184, 185, 642, 658.
7. PHENOLS
- 271 Belova, Z.A. and Borzhunas, A.L.: (Gas chromatographic determination of phenol in atmospheric air.) *Khim. Prom.*, No. 4 (1992) 213-214.
- 272 Cruz, I. and Wells, D.E.: Determination of pentachlorophenol by exhaustive methylation and capillary gas chromatography in sewage sludge, contaminated water and suspended particulates. *Intern. J. Environ. Anal. Chem.*, 48 (1992) 101-113.
- 273 Kluyuev, N.A., Iilnikov, V.G., Brodskii, E.S. and Bocharov, B.V.: Chlorophenol determination in water samples using combined gas chromatography and mass spectrometry. *Organohalogen Compd.*, 2 (1990) 173-176; *C.A.*, 117 (1992) 177951w.
- 274 Korenman, Ya.I. and Fokin, V.N.: (Gas-chromatographic determination of phenol and cresols in natural waters and treated wastewaters.) *Khim. Tekhnol. Vody*, 14 (1992) 344-347; *C.A.*, 117 (1992) 157279e.

- 275 Kostkina, M.I., Vitenberg, A.G., Novikaite, N.V. and Strukova, T.P.: (High-temperature headspace analysis of trace volatile phenols in aqueous solutions.) *Zh. Anal. Khim.*, 47 (1992) 1665-1670.
- 276 Nick, K. and Schöler, H.F.: Gas-chromatographic determination of nitrophenols after derivatisation with diazomethane. *Fresenius J. Anal. Chem.* 343 (1992) 304-307.
- 277 Nolte, J. and Mayer, H.: (Determination of nitrophenols by capillary gas chromatography.) *GIT Fachz. Lab.*, 36 (1992) 339-340,342-344; *C.A.*, 117 (1992) 39569b.
- 278 Ong, C.P., Lee, H.K. and Li, S.F.Y.: Supercritical fluid chromatographic separation of 11 priority phenols using chlorodifluoromethane as modifier. *J. Chromatogr. Sci.*, 30 (1992) 319-323.
- 279 Salehuddin, S.M., Mottaleb, M.A. and Khan, A.H.: Determination of phenol in aqueous medium by gas-liquid chromatography. *J. Bangladesh Acad. Sci.*, 16 (1992) 69-74; *C.A.*, 117 (1992) 55497g.
- 280 Zatula, A.I. and Kivshik, L.S.: (Determination of phenol in natural waters and wastewater at the level of the maximum permissible concentration.) *Gidrokhim. Mater.*, 110 (1991) 123-129; *C.A.*, 117 (1992) 76105t.
- See also 171, 712.
8. SUBSTANCES CONTAINING HETEROCYCLIC OXYGEN
- 8a. Flavonoids
- 281 Creaser, C.S., Koupai-Abyazani, M.R. and Stephenson, G.R.: Gas chromatographic-mass spectrometric characterization of flavanones in citrus and grape juices. *Analyst (London)*, 117 (1992) 1105-1109.
- 282 English, S., Greenaway, W. and Whatley, F.R.: Analysis of phenolics in the bud exudates of *Populus deltoides*, *P. fremontii*, *P. sargentii* and *P. wislizenii* by GC-MS. *Phytochemistry*, 31 (1992) 1255-1260; *C.A.*, 117 (1992) 86715f.
- 283 Greenaway, W., English, S. and Whatley, F.R.: Relationships of *Populus acuminata* and *Populus generosa* with their parental species examined by gas chromatography - mass spectrometry of bud exudates. *Can. J. Bot.*, 70 (1992) 212-221; *C.A.*, 117 (1992) 44546w.
- 284 Greenaway, W., English, S., May, J. and Whatley, F.R.: Analysis of phenolics of bud exudates of *Populus cathayana* and *Populus szechuanica* by GC-MS. *Z. Naturforsch., C: Biosci.*, 47 (1992) 308-312; *C.A.*, 117 (1992) 44578h.
- 285 Greenaway, W., English, S., May, J. and Whatley, F.R.: Analysis of phenolics of bud exudates of *Populus koreana*, *Populus maximowiczii* and *Populus suaveolens* by GC-MS. *Z. Naturforsch., C: Biosci.*, 47 (1992) 313-317; *C.A.*, 117 (1992) 44579j.
- 286 Greenaway, W., May, J., Scaysbrook, T. and Whatley, F.R.: Analysis of phenolics of bud exudates of *Populus simonii* and *Populus yunnanensis* by GC-MS. *Z. Naturforsch., C: Biosci.*, 47 (1992) 477-480; *C.A.*, 117 (1992) 86804j.
- 287 Koupai-Abyazani, M.R., Creaser, C.S. and Stephenson, G.R.: Separation and identification of flavone, flavonol, isoflavone and flavanone aglycones by capillary column gas chromatography. *Phytochem. Anal.*, 3 (1992) 80-84; *C.A.*, 117 (1992) 43760z.
- 288 Suto, K., Masuda, M., Maruta, T., Sagara, K. and Mizutani, T.: (Application of supercritical fluid extraction and chromatography to determination of paeoniflorin and albiflorin in *Paeoniae Radix*.) *Shoyakugaku Zasshi*, 46 (1992) 9-13; *C.A.*, 117 (1992) 157732x.
- 8b. Aflatoxins and other mycotoxins
- 289 Dorizzi, R., Michelot, D., Tagliaro, F. and Ghielmi, S.: Methods for chromatographic determination of amanitins and related toxins in biological samples. *J. Chromatogr.*, 580 (1992) 279-291 - a review with 55 refs.
- 290 Spande, T.F., Garraffo, H.M., Daly, J.W., Tokuyama, T. and Shimada, A.: Identification of histriocotoxins by GC-MS and GC-FTIR: photo- and chemical-artefacts and revised carbon-13NMR assignments. *Tetrahedron*, 48 (1992) 1823-1826; *C.A.*, 117 (1992) 48961a.
- 8c. Other compounds with heterocyclic oxygen (incl. tannins)
- 291 Alexandrou, N., Miao, Z., Colquhoun, M., Pawliszyn, J. and Jenison, C.: Supercritical fluid extraction and cleanup with capillary GC-ion trap mass spectrometry for determination of polychlorinated dibenzo-p-dioxins and dibenzofurans in environmental samples. *J. Chromatogr. Sci.*, 30 (1992) 351-357.
- 292 Amemiya, T., Suzuki, S. and Nakamura, H.: (Determination of 1,4-dioxane in shampoos by headspace-capillary gas chromatography.) *Kenkyu Nenpo - Tokyo-toritsu Eisei Kenkyusho*, 42 (1991) 54-57; *C.A.*, 117 (1992) 137389n.
- 293 Bacher, R. and Ballschmitter, K.: Separation of polychlorodibenzo-p-dioxins and -dibenzofurans on the new polar stationary phase DB DIOXIN. *Chromatographia*, 34 (1992) 137-142.
- 294 Brodskii, E.S., linikov, V.G., Klyuev, N.A., Prokof'ev, A.K., Murenets, N.V. and Bocharov, B.V.: Tetrachlorbenzo-p-dioxins determination in surface and sewage water using GC, GC-LRMS and GC-HRMS. *Organohalogen Compd.*, 2 (1990) 115-116; *C.A.*, 117 (1992) 177950v.
- 295 Brodskii, E.S., Klyuev, N.A., Zhilnikov, V.C., Murenets, N.V., Bochorov, B.V. and Rusinov, G.L.: (Determination of tetrachlorinated dibenzo-p-dioxins, dibenzofurans and related compounds in commercial herbicide 2,4-D.) *Zh. Anal. Khim.*, 47 (1992) 1497-1503.
- 296 David, F., Sandra, P., Hoffmann, A. and Gerstel, J.: A novel CGC-CGC-MS configuration for the analysis of polychlorodibenzodioxins and polychlorodibenzofurans. *Chromatographia*, 34 (1992) 259-265.
- 297 Elli, G.: (Determination of dioxin by high-resolution gas chromatography/mass spectrometry (HRGC/MS).) *Lab. 2000*, 4, No. 7 (1990) 76-79; *C.A.*, 117 (1992) 163173g.
- 298 Golovnya, R.V., Misharina, T.A. and Beletskiy, I.V.: Influence of methyl, formyl and acetyl groups on retention of substituted furans and thiophenes in capillary GC. *Chromatographia*, 34 (1992) 497-501.
- 299 Huang, L.Q., Paiva, A., Tong, H., Monson, S.J. and Gross, M.L.: Application of gas chromatography/high-resolution mass spectrometry to the determination of trace monobromopolychlorodibenzo-p-dioxins in environmental samples. *J. Am. Soc. Mass Spectrom.*, 3 (1992) 248-259; *C.A.*, 117 (1992) 61967b.

- 300 Jia, C., Qiu, Y. and Yu, W.: (Multidimensional gas chromatography for the separation of TCDDs and PCBs.) *Huanjing Huaxue*, 11, No. 2 (1992) 52-57; *C.A.*, 117 (1992) 184022x.
- 301 Matsuzaki, Y., Ishibashi, E., Koguchi, S., Wakui, Y., Takeda, S., Aburada, M. and Oyama, T.: (Determination of gomisin A (TJN-101) and its metabolite in rat serum by gas chromatography-mass spectrometry. *Yakugaku Zasshi*, 111 (1991) 617-620; *C.A.*, 117 (1992) 19751b.
- 302 Pettit, K., Brown, R.S. and Jones, P.W.: The DB5 capillary column: the search for sensitivity. *Organohalogen Compd.*, 2 (1990) 203-206; *C.A.*, 117 (1992) 177269e.
- 303 Sovocool, G.W., Donnelly, J.R. and Munslow, W.D.: Gas chromatographic retention index modeling for environmental monitoring of halogenated dioxins. *Organohalogen Compd.*, 2 (1990) 361-364; *C.A.*, 117 (1992) 177271z.
- 304 Tanaka, K., Hara, S. and Makino, S.: (Development of new program for dioxin analysis using mass spectrometry.) *Kankyo Kagaku*, 2 (1992) 428-429; *C.A.*, 117 (1992) 177692n.
- 305 Telliard, W.A., McCarty, H.B. and King, J.R.: US EPA Method 1613. High resolution gas chromatography/high resolution mass spectrometry determination of tetra- through octa-chlorinated dibenzo-*p*-dioxins and dibenzofurans by isotope dilution. *Organohalogen Compd.*, 2 (1990) 303-306; *C.A.*, 117 (1992) 177707w.
- 306 Tong, H., Gross, M., Gibling, D., Monson, S., Huang, L., Moore, C., Moncur, J. and Ryan, P.: HRGC/HRMS with mass profile monitoring for the analysis of dioxin and related compounds. *Chemosphere*, 25 (1992) 21-24.
- 307 Yoshioka, K., Matuda, M. and Wakimoto, T.: (Study of analytical procedure of dibenzo-*p*-dioxin and dibenzofuran.) *Kankyo Kagaku*, 2 (1992) 356-357; *C.A.*, 117 (1992) 137208c.
- See also 244, 283, 284, 285, 358, 619, 651, 765, 791.
9. OXO COMPOUNDS, ETHERS, EPOXIDES AND QUINONES
- 308 Arias-Perez, R., Lorenzo-Ferreira, R.A. and Alvarez-Devesa, A.: Determination of formaldehyde in particleboards using head-space gas chromatography. *Wood Sci. Technol.*, 26 (1992) 219-226; *C.A.*, 117 (1992) 113852g.
- 309 Cardeal, Z.L., Hamon, M. and Pradeau, D.: (Standardized method for the determination of ethylene oxide.) *Anal. Chim. Acta*, 264 (1992) 365-368.
- 310 Ebeler, S.E., Hinrichs, S.H., Clifford, A.J. and Shibamoto, T.: Analysis of reactive carbonyls in the expired air transgenic mice. *Anal. Biochem.*, 205 (1992) 183-186.
- 311 Lyapkin, A.A., Plashina, E.L. and Churakova, N.S.: (Method for determination of formaldehyde using gas chromatography with flame-ionization detector.) *U.S.S.R. SU 1,709,208* (Cl. G01N31/00), 30 Jan. 1992, Appl. 4,786,876, 30 Jan. 1990; *C.A.*, 117 (1992) 103340z.
- 312 Minagawa, T., Suzuki, S. and Suzuki, T.: Analysis of benzophenones by gas chromatography/Fourier transform-infrared spectrometry. *Forensic Sci. Int.*, 51 (1991) 179-188; *C.A.*, 117 (1992) 42068t.
- 313 Vohra, K. and Gaind, V.S.: Gas chromatographic determination of 2,5-hexanedione in urine as an indicator of exposure to n-hexane. *J. Anal. Toxicol.*, 16 (1992) 176-178; *C.A.*, 117 (1992) 42054k.
- 314 Yamamoto, H.: Establishment of a simple and reliable method of capillary gas chromatography for simultaneous determination of chloral hydrate and its metabolites. *Mie Med. J.*, 41 (1991) 173-180; *C.A.*, 117 (1992) 19789v.
- 315 Yasuhara, A., Kataoka, M. and Shibamoto, T.: (Analysis of aldehydes in gasoline-car exhaust.) *Kankyo Kagaku*, 2 (1992) 448-449; *C.A.*, 117 (1992) 117337r.
- 316 Zeeck, E., Hardege, J.D., Willig, A., Krebber, R. and König, W.A.: Preparative separation of enantiomeric polychaete sex pheromones. *Naturwissenschaften*, 79 (1992) 182-183.
- See also 64, 422, 425, 671, 679.
10. CARBOHYDRATES
- 10a. Mono and oligosaccharides. Structural studies
- 317 Cherniak, R., Morris, L.C. and Turner, S.H.: Glucuronoxylomanan of *Cryptococcus neoformans* serotype D: structural analysis by gas-liquid chromatography-mass spectrometry and by ¹³C-nuclear magnetic resonance spectroscopy. *Carbohydr. Res.*, 223 (1992) 263-269; *C.A.*, 116 (1992) 147689d.
- 318 Fuelster, R.G.: Quantitation of sugars in street drug samples. *J. Forensic Sci.*, 37 (1992) 77-81; *C.A.*, 117 (1992) 165393c.
- 319 García-Raso, A., Páez, M.I., Martínez-Castro, I., Sanz, J. and Calvo, M.M.: Gas chromatographic retention of carbohydrate trimethylsilyl ethers. *J. Chromatogr.*, 607 (1992) 221-225.
- 320 Guo, Z.K., Lee, W.N.P., Katz, J. and Bergner, A.E.: Quantitation of positional isomers of deuterium-labeled glucose by gas chromatography/mass spectrometry. *Anal. Biochem.*, 204 (1992) 273-282.
- 321 Harris, P.J., Sadek, M., Brownlee, R.T.C., Blakeney, A.B., Webster, J. and Stone, B.A.: The detection and quantification of apiose by capillary gas chromatography of its alditol acetates. *Carbohydr. Res.*, 227 (1992) 365-370; *C.A.*, 117 (1992) 107518m.
- 322 Morvai, M. and Molnár-Perl, I.: Simultaneous gas chromatographic quantitation of sugars and acids in citrus fruits, pears, bananas, grapes, apples and tomatoes. *Chromatographia*, 34 (1992) 502-504.
- 323 Osman, S.F. and O'Connor, J.: Gas chromatography-mass spectrometry method for the determination of the reducing end of oligo- and polysaccharides. *J. Chromatogr.*, 606 (1992) 285-287.
- 324 Ukai, S., Yoshida, I., Honda, A., Nagai, K. and Kiho, T.: The distribution of carboxymethyl groups in O-(carboxymethyl)ated (1,4-darw.3)- β -D-glucans and (1,4-darw.3)- α -D-glucans. *Carbohydr. Res.*, 224 (1992) 201-208; *C.A.*, 116 (1992) 174597d.
- See also 760.

10b. *Polysaccharides, mucopolysaccharides, lipopolysaccharides*

- 325 Mischnick, P. and Koenig, W.A.: (Determination of the substituent distribution in modified polysaccharides by chemical degradation and gas chromatography/mass spectrometry.) *Papier (Darmstadt)*, 45 (1991) 757-764; *C.A.*, 117 (1992) 28995m.
- 326 Schomburg, G., Deege, A., Hinrichs, H., Hübinger, E. and Husmann, H.: Preparation, purification, and analysis of alkylated cyclodextrins. *J. High Resolut. Chromatogr.*, 15 (1992) 579-584.

See also 323, 353.

10c. *Glycoproteins and their constituents*

- 327 Kuksis, A., Marai, L., Myher, J.J., Itabashi, Y. and Pind, S.: Application of GC/MS, LC/MS and FAB/MS to determination of molecular species of glycerolipids. In: Perkins, E.G. (Editor), *Analysis of Fats, Oil Lipoproteins*, AOCS, Champaign, Ill., 1991, pp. 464-465; *C.A.*, 117 (1992) 166826h - a review with 41 refs.
- 328 Teixidor, P. and Grimalt J.O.: Gas chromatographic determination of isoprenoid alkylglycerol diethers in archaeobacterial cultures and environmental samples. *J. Chromatogr.*, 607 (1992) 253-259.

11. ORGANIC ACIDS AND LIPIDS

11a. *Organic acids and simple esters*

- 329 Ackman, R.G.: Application of gas-liquid chromatography to lipid separation and analysis: qualitative and quantitative analysis. In: Perkins, E.G. (Editor), *Analysis of Fats, Oils Lipoproteins*, AOCS, Champaign, Ill., 1991, pp. 270-300; *C.A.*, 117 (1992) 169630a.
- 330 Asilbekova, D.T., Gusakova, S.D. and Glushenkova, A.I.: (New hydroxy fatty acids from *Acanthopanax sessiliflorus*.) *Khim. Prir. Soedin.*, No. 5 (1991) 616-619; *C.A.*, 117 (1992) 150789n.
- 331 Barbeni, M., Allegrone, G., Cisero, M. and Guarda, P.A.: GC and NMR enantiodiscrimination of 2-methyl substituted aliphatic acids via deastereomeric esterification with (R)-pantolactone. *Flavour Fragrance J.*, 7 (1992) 163-167; *C.A.*, 117 (1992) 149575h.
- 332 Benard, G., Antoine, J.A., Ouattara, B. and Labie, C.: (Fatty acids of fatty livers and some animal and vegetable fats. Detection of the addition of exogenous fats to fatty liver products.) *Microbiol., Aliments, Nutr.*, 10 (1992) 29-34; *C.A.*, 117 (1992) 169696b.
- 333 Blanchemian, A., Grizeau, D. and Guary, J.C.: (Quantitative analysis of eicosapentaenoic acid from marine microalgae: advantages and disadvantages of some methods.) *Oceanis*, 18 (1992) 169-176; *C.A.*, 117 (1992) 127531p.
- 334 Bronner, W.E.: Gas chromatographic-mass spectrometric methods of analysis for detection of 11-nor- Δ^9 -tetrahydrocannabinol-9-carboxylic acid in biological matrices. *J. Chromatogr.*, 580 (1992) 63-75 - a review with 112 refs.
- 335 Cheng, G., Cao, S., Zhang, G., Liu, L., Wang, D. and Que, H.: (Detection of fatty acids in import palm oil by GC/MS.) *Fenxi Ceshi Tongbao*, 10 (1991) 43-45; *C.A.*, 116 (1991) 104723t.
- 336 Clemens, M. and Schöler, H.F.: Determination of halogenated acetic acids and 2,2-dichloropropionic acid in water samples. *Fresenius J. Anal. Chem.*, 344 (1992) 47-49.
- 337 Eder, K., Reichlmayr-Lais, A.M. and Kirchgessner, M.: Studies on the methanolysis of small amounts of purified phospholipids for gas chromatographic analysis of fatty acid methyl esters. *J. Chromatogr.*, 607 (1992) 55-67.
- 338 Frank, H. and Scholl, H.: Determination of TCA in spruce needles. *Organohalogen Compd.*, 2 (1990) 397-399; *C.A.*, 117 (1992) 165356t.
- 339 Gaetani, E., Laureri, C.F. and Vitto, M.: Ion trap detector - capillary gas chromatography of valproic acid and its mono-unsaturated metabolites in serum using methyl ester derivatives. *J. Pharm. Biomed. Anal.*, 10 (1992) 193-197; *C.A.*, 117 (1992) 103454q.
- 340 Higuchi, O.: (Analysis of fatty acids in human synovial fluid by gas chromatography chemical ionization mass spectrometry.) *Kurume Igakkai Zasshi*, 55 (1992) 301-311; *C.A.*, 117 (1992) 129264c.
- 341 Hill, E. and Murphy, R.C.: Quantitation of 20-hydroxy-5,8,11,14-eicosatetraenoic acid (20-HETE) produced by human polymorphonuclear leukocytes using electron capture ionization gas chromatography/mass spectrometry. *Biol. Mass Spectrom.*, 21 (1992) 249-253; *C.A.*, 117 (1992) 65778p.
- 342 Hillman, K., Williams, A.G. and Lloyd, D.: The influence of head-space gas composition on the production of VFA by rumen ciliate protozoa. *Lett. Appl. Microbiol.*, 13 (1991) 11-15; *C.A.*, 116 (1992) 102369h.
- 343 Jin, P.: (The precision and accuracy of gas chromatography in analysis of ethyl caproate in baijiu.) *Niangjiu*, No. 2 (1992) 8-12; *C.A.*, 117 (1992) 169675u.
- 344 Kageyama, K., Onoyama, Y., Nakajima, T., Tumura, M., Nishida, T., Kogawa, H. and Tanabe, K.: Rapid and simple gas chromatographic measurement of lactic acid in red blood cells, plasma, and tumor cells after hyperthermia. *Life Sci.*, 51 (1992) 407-410; *C.A.*, 117 (1992) 107536r.
- 345 Kang, X.: (Analysis of fatty acids in vegetable oils by cross-linked DEGS glass capillary column with large-diameter and thick film.) *Sepu*, 10 (1992) 300-302.
- 346 Kawase, M., Sonomoto, K. and Tanaka, A.: Inspection of an acyl-enzyme intermediate in a lipase reaction by gas chromatography-mass spectrometry and modeling of the reaction mechanism. *Biocatalysis*, 6 (1992) 43-50; *C.A.*, 117 (1992) 107138n.
- 347 Kozma-Kovacs, E., Halasz, A., Hajos, G., Sass, A. and Boross, F.: (Determination of lipoic acid in brewers' yeast by thin-layer and capillary gas chromatographic methods.) *Élelmiszervizsgálati Közl.*, 38 (1992) 18-26; *C.A.*, 117 (1992) 24856u.
- 348 Li, X., Chen, G., Gao, S. and Guan, Z.: (Rapid analysis of fatty acids in cerebrospinal fluid by gas chromatography and its clinical significance.) *Sepu*, 10 (1992) 248-249.
- 349 Li, X., la Motte, C.E., Stewart, C.R., Cloud, N.P., Wear-Bagnall, S. and Jiang, C.Z.: Determination of IAA and ABA in the same plant sample by a widely applicable method using GC-MS with selected ion monitoring. *J. Plant Growth Regul.*, 11 (1992) 55-65; *C.A.*, 117 (1992) 22631t.
- 350 Lie Ken Jie, M.S.F. and Choi, Y.C.: Gas chromatography-mass spectrometry of the picolinyl ester derivatives of deuterated acetylenic fatty acids. *J. Chromatogr.*, 625 (1992) 271-276.

- 351 Matter, L.: Determination of the animal of origin of dairy products and raw and cooked meats by GC analysis of the fatty acid methyl esters (FAME) obtained by transesterification. *J. High Resolut. Chromatogr.*, 15 (1992) 514-516.
- 352 Mattulat, A. and Baltes, W.: Determination of lipoic acid in meat of commercial quality. *Z. Lebensm.-Unters. Forsch.*, 194 (1992) 326-329; C.A., 117 (1992) 68608u.
- 353 Mielniczuk, Z., Alugupalli, S., Mielniczuk, E. and Larsson, L.: Gas chromatography-mass spectrometry of lipopolysaccharide 3-hydroxy fatty acids: comparison of pentafluorobenzoyl and trimethylsilyl methyl ester derivatives. *J. Chromatogr.*, 623 (1992) 115-122.
- 354 Murooka, M., Ozawa, H. and Matsubara, H.: (Examinations of the measurement of atmospheric low-molecular fatty acids by using beads coated with alkaline compound.) *Ehime-ken Kogai Gijutsu Senta Shoho*, 12 (1992) 79-84; C.A., 117 (1992) 177231m.
- 355 Nie, H., Huang, Z. and Pen, S.: (Rapid gas chromatographic method for determination of benzoic acid and sorbic acid in foods.) *Sepu*, 10 (1992) 244-246.
- 356 Promyshlennikova, E.P., Kirichenko, V.E., Pashkevich, K.I., Grigor'eva, D.N. and Golovnya, R.V.: (Influence of the stationary phase nature on the retention of polyfluorinated carboxylic acid esters.) *Zh. Anal. Khim.*, 47 (1992) 1070-1075.
- 357 Rodrigo, M.T. and Mendioroz, S.: Pulse-response chromatographic method as a technique for determination the selectivity of polyunsaturated acids in hydrogenation. *Chromatographia*, 34 (1992) 479-482.
- 358 Samukawa, K., Inoue, Y., Tatsumi, M., Tanaka, R. and Komai, Y.: (Chemical characterization of humic acids in soils with different fertilization histories by Curie-point pyrolysis-gas chromatography-mass spectrometry.) *Nippon Dojo Hiriyogaku Zasshi*, 63 (1992) 129-138; C.A., 117 (1992) 25375y.
- 359 Shimizu, A. and Nakanishi, T.: (Quantitative analysis of urinary organic acid.) *Rinsho Byori*, 40 (1992) 743-750; C.A., 117 (1992) 107467u - a review with 5 refs.
- 360 Smith, N.B. and Brooks, K.E.: Use of Toxi-Lab thin-layer chromatography in plasma very-long-chain fatty acid analysis. *Clin. Chim. Acta*, 206 (1992) 241-243.
- 361 Sollinger, S., Levsen, K. and Emmrich, M.: Determination of trace amounts of carboxylic acids in ambient air by capillary gas chromatography-mass spectrometry. *J. Chromatogr.*, 609 (1992) 297-304.
- 362 Ulberth, F. and Henninger, M.: One-step extraction/methylation method for determining the fatty acid composition of processed foods. *J. Am. Oil Chem. Soc.*, 69 (1992) 174-177.
- 363 Ulberth, F. and Henninger, M.: Simplified method for the determination of trans monoenes in edible fats by TLC-GLC. *J. Am. Oil Chem. Soc.*, 69 (1992) 829-831.
- 364 Varvaro, L.: (Identification of plant-pathogenic bacteria through cellular fatty acid analysis.) *Lab. 2000*, 6, No. 3 (1992) 61-64; C.A., 117 (1992) 127528t.
- 365 Vohra, K. and Gaiand, V.S.: Gas chromatographic determination of airborne organic acids via their anilides. *Analyst (London)*, 117 (1992) 1567-1570.
- 366 Wesén, C., Mu, H., Kvernheim, A.L. and Larsson, P.: Identification of chlorinated fatty acids in fish lipids by partitioning studies and by gas chromatography with Hall electrolytic conductivity detection. *J. Chromatogr.*, 625 (1992) 257-269.
- 367 Wolfe, M.F. and Seiber, J.N.: A method for the trace analysis of naptalam (N-1-naphthylphthalamic acid) in water. *Arch. Environ. Contam. Toxicol.*, 23 (1992) 137-141; C.A., 117 (1992) 76082h.
- 368 Wu, C., Pan, G. and Zeng, Z.: (Identification of mycobacteriums by gas chromatography (GC) with a crown ether capillary column.) *Sepu*, 10 (1992) 260-262.
- 369 Xu, Z., Li, Y., Li, X., Yu, C., Li, G. and Yang, N.: (Study of composition and content of fatty acids in the retina's lipids of rabbits by gas chromatography.) *Sepu*, 10 (1992) 283-285.
- 370 Yan, C.T. and Jen, J.F.: Determination of volatile fatty acids in landfill leachates by gas chromatography with distillation pretreatment. *Anal. Chim. Acta*, 264 (1992) 259-264.
- 371 Zheng, J., Lin, L. and Liu, P.: (Gas chromatographic determination of propyl p-hydroxybenzoate in foods.) *Shipin Kexue (Beijing)*, 144 (1991) 48-49; C.A., 117 (1992) 68576g.
- See also 64, 149, 162, 184, 322, 377, 420, 650, 651, 655, 660, 666, 668, 777, 781, 796.
- 11b. *Prostaglandins*
- 372 Ferretti, A., Flanagan, V.P. and Maida, E.J.: GC/MS/MS quantification of 11-dehydrothromboxane B₂ in human urine. *Prostaglandins, Leukotrienes Essent. Fatty Acids*, 46 (1992) 271-275; C.A., 117 (1992) 143637h.
- 11c. *Lipids and their constituents*
- 373 Ferreira, F. and Moyna, P.: Simple TLC-GC method for semi-quantitative separation of lipid classes in coastal sediments. *J. Liq. Chromatogr.*, 15 (1992) 1655-1663; C.A., 117 (1992) 84732k.
- 374 Ikawa, N., Furuta, S., Fukuzato, R., Imanishi, N., Kawana, H., Mitsuiki, M. and Suzuki, H.: Separation of polyunsaturated fatty acid from vegetable oil by supercritical fluid extraction and chromatography. *Process Metall.*, 7B (1992) 1707-1712; C.A., 117 (1992) 130175f.
- 375 Pinkston, J.D. and Bowling, D.J.: Use of SFC/MS for the confirmation of peak identities in the SFC/FID chromatogram of olestra. *J. Microcolumn Separ.*, 4 (1992) 295-303.
- 376 Plank, C. and Lorbeer, E.: Quality control of vegetable oil methyl esters used as diesel fuel substitutes: quantitative determination of mono-, di-, and triglycerides by capillary GC. *J. High Resolut. Chromatogr.*, 15 (1992) 609-612.
- 377 Ruiz-Gutierrez, V., Montero, E. and Villar, J.: Determination of fatty acid and triacylglycerol composition of human adipose tissue. *J. Chromatogr.*, 581 (1992) 171-178.
- See also 657.
12. **ORGANIC PEROXIDES**
- 378 Baechmann, K., Hauptmann, J., Polzer, J. and Schuetz, P.: Determination of organic alkyl- and 1-hydroxy hydroperoxides with chemiluminescence, fluorescence and GC/MS. *Fresenius J. Anal. Chem.*, 342 (1992) 809-812.
- See also 72.

13. STEROIDS

13b. Pregnane and androstane derivatives

- 379 Bi, H., Du, P. and Masse, R.: Studies on anabolic steroids-8. GC/MS characterization of unusual seco acidic metabolites of oxymetholone in human urine. *J. Steroid Biochem. Mol. Biol.*, 42 (1992) 229-242; *C.A.*, 117 (1992) 020715z.
- 380 Bohm, D., Goebel, R., Lippert, A., Tesch, D. and Lusky, K.: (Analysis of xylazine residues by gas chromatography/mass spectrometry.) *Bioforum*, 15 (1992) 212-214; *C.A.*, 117 (1992) 130070t.
- 381 De Boer, D., de Jong, E.G., Maes, R.A.A. and van Rossum, J.M.: The methyl-5 α -dihydrotestosterones mesterolone and drostanolone; gas chromatographic/mass spectrometric characterization of the urinary metabolites. *J. Steroid Biochem. Mol. Biol.*, 42 (1992) 411-419; *C.A.*, 117 (1992) 63110r.
- 382 Liu, C.S., Zhang, J. and Zhou, T.H.: (Investigation on the metabolism of dehydrochloromethyltestosterone in human.) *Yaoxue Xuebao*, 26 (1991) 682-687; *C.A.*, 117 (1992) 42067s.
- 383 Risk, M.S., Zakhari, N.A., Walsh, M.I., Toubar, S.S., Books, C.J.W. and Anderson, R.: Gas chromatography and mass spectrometry of dimethylethylsilyl ether derivatives of norethisterone metabolites in plasma. *Acta Pharm. Nord.*, 3 (1991) 205-210; *C.A.*, 116 (1992) 166411y.
- 384 Shibasaki, H., Furuta, T. and Kasuya, Y.: Stable isotope dilution mass spectrometry for the simultaneous determination of cortisol, cortisone, prednisolone and prednisone in plasma. *J. Chromatogr.*, 579 (1992) 193-202.
- 385 Trafford, D.J.H., Coldwell, R.D. and Makin, H.L.J.: Gas chromatography-mass spectrometry in the investigation of on-column dehydration of steroid hormones during gas-liquid chromatography. *J. Pharm. Biomed. Anal.*, 9 (1991) 1095-1105; *C.A.*, 117 (1992) 40608v.
- 386 Venturelli, E., Manzari, A., Cavalleri, A., Benzo, M., Secreto, G. and Marubini, E.: Urinary testosterone measurement by gas chromatography after solid-phase extraction and high-performance liquid chromatography. *J. Chromatogr.*, 582 (1992) 7-12.

See also 647, 648, 654.

13d. Sterols

- 387 Bernert, J.T., Jr., Akins, J.R., Cooper, G.R., Poulouse, A.K., Myers, G.L. and Sampson, E.J.: Factors influencing the accuracy of the national reference system total cholesterol reference method. *Clin. Chem. (Winston-Salem)*, 37 (1991) 2053-2061.
- 388 Lognay, G., Severin, M., Boenke, A. and Wagstaffe, P.J.: Edible fats and oils reference materials for sterols analysis with particular attention to cholesterol. Part I. Investigation of some analytical aspects by experienced laboratories. *Analyst (London)*, 117 (1992) 1093-1097.
- 389 Pioch, D., Lozano, P., Frater, C. and Graille, J.: A quick method for cholesterol titration in complex media. *Fat Sci. Technol.*, 94 (1992) 268-272; *C.A.*, 117 (1992) 149605t.

- 390 Stewart, G., Gosselin, C. and Pandian, S.: Selected ion monitoring of *tert*-butyldimethylsilyl cholesterol ethers for determination of total cholesterol content in foods. *Food Chem.*, 44 (1992) 377-380; *C.A.*, 117 (1992) 46866z.
- 391 Zhou, J. and Fu, C.: (Rapid determination of cholesterol in egg yolks by short capillary column gas chromatography.) *Sepu*, 10 (1992) 250.

See also 657.

15. TERPENES AND OTHER VOLATILE AROMATIC COMPOUNDS

15a. Terpenes

- 392 Boland, W., Feng, Z., Donath, J. and Gäbler, A.: Are acyclic C₁₁ and C₁₆ homoterpenes plant volatiles indicating herbivory? *Naturwissenschaften*, 79 (1992) 368-371.
- 393 Kreis, P., Juchelka, D., Motz, C. and Mosandl, A.: (Chiral components of essential oils. Part 9. Stereodifferentiation of borneol, isoborneol, and bornyl acetate.) *Dtsch. Apoth. Ztg.*, 131 (1991) 1984-1987; *C.A.*, 117 (1992) 137391g.
- 394 Lu, M., Wang, Y., Jing, S. and Chen, Z.: (Determination of trace isoprene and monoterpenes in the atmosphere by gas chromatography with the photoionization detector.) *Huanjing Huaxue*, 11, No. 2 (1992) 21-25; *C.A.*, 117 (1992) 117339t.
- 395 Ohta, S., Yamamitsu, T. and Suga, T.: An efficient method for following the enzymic reactions involved in camphor biosynthesis in *Cinnamomum camphora* by use of GC-mass spectroscopy and regiospecifically deuteriated substrate. *J. Labelled Compd. Radiopharm.*, 31 (1992) 397-404; *C.A.*, 117 (1992) 43991a.
- 396 Ravid, U., Putievsky, E., Katzir, I. and Ikan, R.: Chiral GC analysis of enantiomerically pure fenchone in essential oils. *Flavour Fragrance J.*, 7 (1992) 169-172; *C.A.*, 117 (1992) 137396n.
- 397 Valterová, I., Vrkok, J., Lindström, M. and Norin, T.: On the natural occurrence of (-)-3-carene, a component of termite defense secretions. *Naturwissenschaften*, 79 (1992) 416-417.
- 398 Wang, Q., Sun, F., Zhang, L., Ni, C., Zhang, Z. and Zhu, Y.: (Sampling and determination of camphor in workplace atmosphere and ambient air by gas chromatography.) *Zhongguo Huanjing Kexue*, 11 (1991) 232-234; *C.A.*, 117 (1992) 96400p.
- 399 Zhou, Y., He, G., Yang, M., Song, S., Zhu, J. and Zhu, C.: (Quantitative determination of the total tanshinone and borneol in "compound Danshen oral liquid".) *Zhongcaoyao*, 23 (1992) 292-293; *C.A.*, 117 (1992) 137760b.

See also 108, 184, 412, 689.

15b. Essential oils

- 400 Bicchi, C., Frattini, C., Pellegrino, G., Rubiolo, P., Ravertino, V. and Tsoupras, G.: Determination of sulphurated compounds in *Tagetes patula* cv. *nana* essential oil by gas chromatography with mass spectrometric, Fourier transform infrared and atomic emission spectrometric detection. *J. Chromatogr.*, 609 (1992) 305-313.

- 401 Chang, K.W., Moon, C.S., Lee, H.D., Lee, C.J. and Lee, U.C.: Determination of the effective components in the various parts of *Luffa cylindrica* (L.) Roemer and development to remove the flesh from its fruit: (I). Determination of chemical components in the various parts of *Luffa cylindrica* (L.) Roemer by GC and GC/MS. *Han'guk Nonghwa Hakhoechi*, 34 (1991) 366-372; C.A., 117 (1992) 23296z.
- 402 Chang, X., Shi, Y., Ma, B. and He, L.: (Chemical compositions of volatile oil from *Actinidia* leaves.) *Baiqiu'en Yike Daxue Xuebao*, 17 (1991) 569-570; C.A., 117 (1992) 157493v.
- 403 Dai, L., Yang, L., Guo, Y. and Peng, Q.: (Analysis of fragrance volatile chemical constituents of *Zhangzhou Narcissus*.) *Sepu*, 10 (1992) 280-283.
- 404 Jirovetz, L., Jaeger, W., Koch, H.P. and Remberg, G.: Investigation of the volatile constituents of the essential oil of Egyptian garlic by GC-MS and GC-FTIR. *Z. Lebensm.-Unters. Forsch.*, 194 (1992) 363-365; C.A., 117 (1992) 89146g.
- 405 Jung, D.S.: (Comparison on volatile constituents of *Houttuynia cordata* and *Saururus chinensis* by GC and GC-MS method.) *Nonmunjip - Chepu Taehakkyo, Chayon Kwahakpyon*, 33 (1991) 105-111; C.A., 117 (1992) 76304g.
- 406 Ma, X., Yu, X., Zheng, Z. and Mao, J.: (Investigation of volatile composition in frankincense and myrrh using analytical supercritical fluid extraction technique.) *Yaowu Fenxi Zazhi*, 12 (1992) 83-86; C.A., 117 (1992) 56093j.
- 407 Ohta, H., Nogata, Y., Yoza, K.-I. and Maeda, M.: Glass capillary gas chromatographic identification of volatile components recovered from orange essence by continuous liquid extraction. *J. Chromatogr.*, 607 (1992) 154-158.
- 408 Pan, J., Xu, Z., Ji, L., Zhao, Z. and Tang, X.: (Constituents of essential oils from leaves, stems, and fruits of *Perilla frutescens* (L.) Britt.) *Zhongguo Zhongyao Zazhi*, 17 (1992) 164-165; C.A., 117 (1992) 33787z.
- 409 Pang, N., Yan, Z. and Neu, Z.: (GC-MS analysis of essential oils in needles of *Pinus* in northeastern China.) *Zhongguo Zhongyao Zazhi*, 17 (1992) 166-168; C.A., 117 (1992) 66595p.
- 410 Pino, J.A.: Headspace sampling methods for the volatile components of garlic (*Allium sativum*). *J. Sci. Food Agric.*, 59 (1992) 131-133; C.A., 117 (1992) 110292p.
- 411 Sankar, K.U. and Narasimhan, S.: Quality evaluation of supercritical carbon dioxide extracted essential oils of pepper by GC finger printing and sensory analysis. *PAFAI J.*, 13, No.4 (1991) 35-38; C.A., 117 (1992) 24940s.
- 412 Veijanen, A., Kolehmainen, E., Kauppinen, R., Lahtipera, M. and Paasivirta, J.: Methods for the identification of tainting terpenoids and other compounds from algae. *Water Sci. Technol.*, 25 (1992) 165-170; C.A., 116 (1992) 262167u.
- 413 Vernin, G. and Lageot, C.: Couplage CG/SM pour l'analyse des arômes et des huiles essentielles. *Analisis*, 20 (1992) M34-M39.
- 414 Vernin, G., Metzger, J., Mondon, J.P. and Pieribattesti, J.C.: GC/MS analysis of the leaf oil of *Cryptomeria japonica* D. Don from Reunion Island. *J. Essent. Oil Res.*, 3 (1991) 197-207; C.A., 116 (1992) 080417t.
- 415 Widmer, W.W. and Collins, K.P.: Analysis of essential oils from selected cultivars of *Pelargonium quercifolium* by GC/MS. *J. Essent. Oil Res.*, 3 (1991) 331-340; C.A., 116 (1992) 027779m.
- 416 Ye, C., Feng, Y. and Teng, J.: (Analysis of β -eudesmol, magnolol, and honokiol in *Magnolia officinalis* by PTGC method.) *Yaowu Fenxi Zazhi*, 12 (1992) 159-162; C.A., 117 (1992) 76583x.
- 417 Ylipahkala, T.M. and Jalonen, J.E.: Isolation of very volatile compounds from the leaves of *Ledum palustre* using the purge and trap technique. *Chromatographia*, 34 (1992) 159-162.
- 418 Zhu, G.P.: (Comparison of chemical constituents of essential oils from *Elsholtzia splendens* and cultivated *Mosla chinensis* by GC-MS analysis.) *Yaowu Xuebao*, 27 (1992) 287-293; C.A., 117 (1992) 23321d.
- 419 Zupanc, M., Prosek, M. and Dusan, M.: Combined CI and EI mass spectra in the analysis of essential oils. *J. High Resolut. Chromatogr.*, 15 (1992) 510-513.

See also 155, 691.

15c. Bitter substances

- 420 Liu, Z., Liang, Z. and Wang, D.: (Method for quantitative determination of components in maleic modified rosin by gas chromatography.) *Linchan Huaxue Yu Gongye*, 11 (1991) 279-287; C.A., 117 (1992) 173632v.

16. NITRO AND NITROSO COMPOUNDS

- 421 Chen, S.: (Gas chromatographic determination of trace nitrobenzene in resin.) *Sepu*, 10 (1992) 247-248.
- 422 Galceran, M.T. and Moyano, E.: Effect of solvent polarity on the determination of oxo- and nitro-polycyclic aromatic hydrocarbons using capillary gas chromatography with splitless injection. *J. Chromatogr.*, 607 (1992) 287-294.
- 423 Han, C., Gumbleton, M., Lau, D.T.W. and Benet, L.Z.: Improved gas chromatographic assay for the simultaneous determination of nitroglycerin and its mono- and dinitrate metabolites. *J. Chromatogr.*, 579 (1992) 237-245.
- 424 Hastenteufel, S., Schmidt, M. and Wagner, R.: (Rüstungslastlaster.) *LaborPraxis*, 16 (1992) 888-894.
- 425 Lu, Y., Gu, Z. and Wang, A.: (Analysis of nitrobenzaldehyde with GC.) *Huaxue Shijie*, 33, No. 2 (1992) 84-85; C.A., 117 (1992) 123760q.
- 426 Paschke, T., Hawthorne, S.B., Miller, D.J. and Wenclawiak, B.: Supercritical fluid extraction of nitrated polycyclic aromatic hydrocarbons and polycyclic aromatic hydrocarbons from diesel exhaust particulate matter. *J. Chromatogr.*, 609 (1992) 333-340.
- 427 Shi, M., Tang, M. and Hu, Z.: (Gas chromatographic analysis of trace dinitrotoluene in water.) *Shanghai Huanjing Kexue*, 10, No. 9 (1991) 26-27; C.A., 117 (1992) 76065e.
- 428 Vlad, F.: (Gas chromatographic analysis of products from the mononitration of toluene and ethylbenzene.) *Rev. Chim. (Bucharest)*, 43 (1992) 162-166; C.A., 117 (1992) 123786c.
- 429 Yamashita, T., Yasuda, Y., Haraguchi, K., Sueta, S. and Kido, K.: (Determination of nitrobenzenes in ambient air.) *Taiki Osen Gakkaishi*, 27, No. 1 (1992) 65-72; C.A., 116 (1992) 261407k.

See also 232, 276, 791.

17. AMINES, AMIDES AND RELATED NITROGEN COMPOUNDS

17a. Amines and polyamines

- 430 Arimoto, H. and Fujii, T.: (Selective high-sensitivity analysis of quarternary ammonium salts.) *Jpn. Kokai Tokkyo Koho JP 03,267,755 [91,267,755](Cl. G01N30/06)*, 28 Nov. 1991, Appl. 90/67,455, 16 Mar. 1990; 4 pp.; C.A., 117 (1992) 39447k.
- 431 Jedrzejczak, K. and Gajnd, V.S.: A new polymeric reagent for the pentafluorobenzoylation of aliphatic amines quantified by gas chromatography. *Fresenius J. Anal. Chem.*, 344 (1992) 133-134.
- 432 Kataoka, H., Ohru, S., Miyamoto, Y. and Makita, M.: Determination of low molecular weight aliphatic primary amines in urine as their benzenesulphonyl derivatives by gas chromatography with flame photometric detection. *Biomed. Chromatogr.*, 6 (1992) 251-254.
- 433 Pieraccini, G., Luceri, F. and Moneti, G.: New gas-chromatography/mass-spectrometric method for the quantitative analysis of primary aromatic amines in main- and side-stream cigaret smoke. *Rapid Commun. Mass Spectrom.*, 6 (1992) 406-409; C.A., 117 (1992) 42557b.

17d. Other amine derivatives and amides (excl. peptides)

- 434 Okamoto, Y., Ohashi, N. and Sasano, H.: (Determination of trace N,N-dimethylformamide in environmental sample by gas chromatography-mass spectrometry.) *Kankyo Kagaku*, 2 (1992) 45-51; C.A., 117 (1992) 55467x.

See also 626.

18. AMINO ACIDS AND PEPTIDES; CHEMICAL STRUCTURE OF PROTEINS

18a. Amino acids and their derivatives

- 435 Bertacco, G., Boschelle, O. and Lercker, G.: Gas chromatographic determination of free amino acids in cheese. *Milch-wissenschaft*, 47 (1992) 348-350; C.A., 117 (1992) 169674t.
- 436 Carter, S.M.B., Midgley, J.M., Watson, D.G. and Logan, R.W.: Measurement of urinary medium chain acyl glycines by gas chromatography-negative ion chemical ionization mass spectrometry. *J. Pharm. Biomed. Anal.*, 9 (1991) 969-975; C.A., 117 (1992) 22834m.
- 437 Hennet, R.J.C., Holm, N.G. and Engel, M.H.: Abiotic synthesis of amino acids under hydrothermal conditions and the origin of life: a perpetual phenomenon? *Naturwissenschaften*, 79 (1992) 361-365.
- 438 Leal, W.S., Matsuyama, S., Kuwahara, Y., Wakamura, S. and Hasegawa, M.: An amino acid derivative as the sex pheromone of a scarab beetle. *Naturwissenschaften*, 79 (1992) 184-185.
- 439 Lou, X., Liu, Y., Wang, Q., Zhu, D. and Zhou, L.: (Determination of amino acids with enantiomer labelling by gas chromatography.) *Sepu*, 10 (1992) 195-199.
- 440 Toernqvist, M., Magnusson, A.L., Farmer, P.B., Tang, Y.-S., Jeffrey, A.M., Wazneh, L., Beuling, B.D.T., van der Waal, H. and van Sittert, N.J.: Ring test for low levels of N-(2-hydroxyethyl)valine in human hemoglobin. *Anal. Biochem.*, 203 (1992) 357-360.

- 441 Woo, K.L.: Gas-chromatographic determination of methylthiohydantoin amino acid as N(O)-butyldimethylsilyl derivatives in amino acid sequencing with methylisothiocyanate. *Han'guk Nonghwa Hakhoechi*, 35 (1992) 132-138; C.A., 117 (1992) 151304n.
- 442 Wu, L., Liu, G., Mi, A. and Jiang, Y.: (Gas chromatographic separation of enantiomers of α -amino acids and α -amino methylpyridines on chiral stationary phases.) *Sepu*, 10 (1992) 285-287.

See also 765.

18b. Peptides, peptidic and proteinous hormones, growth factors

- 443 Burinsky, D.J., Dunphy, R., Oylar, A.R., Shaw, C.J. and Cotter, M.L.: Characterization of a synthetic peptide impurity by fast-atom bombardment-tandem mass spectrometry and gas chromatography-mass spectrometry. *J. Pharm. Sci.*, 81 (1992) 597-600; C.A., 117 (1992) 118474v.

20. ENZYMES AND ENZYME ACTIVITY ESTIMATION

20d. Hydrolases, acting on ester bonds (E.C. 3.1.-.-)

See 346.

20g. Lyases

- 444 Pyzhik, T.N., Gerashchenko, D.Y. and Ostrovskii, Y.M.: (Gas chromatographic determination of hepatic ethanolamine phosphate phospho-lyase.) *Vestsi Akad. Navuk BSSR, Ser. Biyax, Navuk*, No. 6 (1991) 45-47; C.A., 117 (1992) 65218f.

21. PURINES, PYRIMIDINES, NUCLEIC ACIDS AND THEIR CONSTITUENTS

21a. Purines, pyrimidines, nucleosides, nucleotides

- 445 Anderson, L.W., Parker, R.J., Collins, J.M., Ahlgren, J.D., Wilkinson, D. and Strong, J.M.: Gas chromatographic-mass spectrometric method for routine monitoring of 5-fluorouracil in plasma of patients receiving low-level protracted infusions. *J. Chromatogr.*, 581 (1992) 195-201.
- 446 Halliwell, B. and Dizdaroglu, M.: The measurement of oxidative damage to DNA by HPLC and GC/MS techniques. *Free Radical Res. Commun.*, 16, No. 2 (1992) 75-87; C.A., 117 (1992) 84696b - a review with 54 refs.
- 447 Hammargen, W.M., Luffer, D.R., Schram, K.H., Reimer, M.L.J., Nakano, K., Yasaka, T. and Moorman, A.R.: The identification of 5'-deoxy-5'-methylthioguanosine in human urine by gas chromatography/mass spectrometry. *Nucleosides Nucleotides*, 11 (1992) 1275-1292; C.A., 117 (1992) 67644r.
- 448 Uchida, J., Umeno, Y. and Takeda, S.: (Significance of measuring 5-fluorouracil incorporated into RNA of tumor tissue as a parameter for the antitumor activity of 5-fluorouracil and its analogs.) *Gan to Kagaku Ryoho*, 19 (1992) 677-683; C.A., 117 (1992) 103440g.

22. ALKALOIDS

- 449 Deutsch, J., Hegedus, L., Greig, N.H., Rapoport, S.I. and Timothy, T.S.: Electron-impact and chemical ionization detection of nicotine and continine by gas chromatography-mass spectrometry in rat plasma and brain. *J. Chromatogr.*, 579 (1992) 93-98.
- 450 Ensing, J.G., Racamy, C. and de Zeeuw, R.A.: A rapid gas chromatographic method for the fingerprinting of illicit cocaine samples. *J. Forensic Sci.*, 37 (1992) 446-459; *C.A.*, 117 (1992) 126041s.
- 451 Fang, H.J., Cheng, K.D. and Xu, Y.Q.: (Analysis of Belladonna alkaloids by capillary GC and GC-MSD.) *Yaoxue Xuebao*, 27 (1992) 220-226; *C.A.*, 117 (1992) 56092h.
- 452 Henderson, G.L., Harkey, M.R., Zhou, C. and Jones, R.T.: Cocaine and metabolite concentrations in the hair of South American coca chewers. *J. Anal. Toxicol.*, 16 (1992) 199-201; *C.A.*, 117 (1992) 42079x.
- 453 Kintz, P.: Gas chromatographic analysis of nicotine and cotinine in hair. *J. Chromatogr.*, 580 (1992) 347-353.
- 454 Liu, R., Li, M., Jing, X., Sun, W., Feng, S. and Gao, L.: (Determination of morphine in the urine of heroin addicts.) *Zhongguo Yaoxue Zazhi*, 27, No. 2 (1992) 84-86; *C.A.*, 117 (1992) 105799e.
- 455 Martinsen, A., Hiltunen, K. and Huhtikangas, A.: Nitrogen-phosphorus detector potential in tropane alkaloid gas chromatography. *Phytochem. Anal.*, 3 (1992) 69-72; *C.A.*, 117 (1992) 43759f.
- 456 Yang, X., Qiao, W., Dong, S. and Wang, P.: (The study of determining method of nicotine in urine after administration of nicotine patch.) *Beijing Yike Daxue Xuebao*, 23 (1991) 403-405; *C.A.*, 117 (1992) 125968n.

See also 623, 624, 640, 705.

23. OTHER SUBSTANCES CONTAINING HETEROCYCLIC NITROGEN

23c. Indole derivatives and plant hormones (gibberelins)

- 457 Schneider, G., Sembdner, G., Jensen, E., Bernhard, U. and Wagenbreth, D.: GC-MS identification of native gibberellin-O-glucosides in pea seeds. *J. Plant Growth Regul.*, 11 (1992) 15-18; *C.A.*, 117 (1992) 23304a.

See also 349.

23d. Pyridine derivatives

See 31.

23e. Other N-heterocyclic compounds

- 458 Golovnya, R.V., Zhuravleva, I.L. and Sal'kova, M.A.: (Prediction of retention indices for alkylimidazoles under conditions for capillary gas chromatography.) *Zh. Anal. Khim.*, 47 (1992) 1269-1275.

24. ORGANIC SULPHUR COMPOUNDS (INCL. GLUCOSINOLATES)

- 459 Adeeva, V.G., Bobyleva, M.S., Kullkov, N.S., Kharchenko, V.G. and Yudovich, L.M.: Gas chromatography-mass spectrometry of the stereoisomers of heterocyclic compounds. Part 1. Perhydrothioxanthenes. *J. Chem. Soc., Perkin Trans. 2*, (1992) 965-969; *C.A.*, 117 (1992) 89648x.
- 460 Andersson, J.T.: Polycyclic aromatic sulfur heterocycles. I. Use of hydrogen peroxide oxidation for the group separation of polycyclic aromatic hydrocarbons and their sulfur analogs. *Intern. J. Environ. Anal. Chem.*, 48 (1992) 1-15.
- 461 Artacho Martin-Lagos, R., Olea Serrano, M.F. and Ruiz-Lopez, M.D.: Comparative study by gas chromatography-mass spectrometry of methods for the extraction of sulfur compounds in *Allium cepa* L. *Food Chem.*, 44 (1992) 305-308; *C.A.*, 117 (1992) 46842p.
- 462 Breeden, D.C. and Juvik, J.A.: An extraction method for the determination of dimethyl sulfide in cooked corn. *J. Food Compos. Anal.*, 5 (1992) 134-138; *C.A.*, 117 (1992) 130031f.
- 463 Buser, H.R.: Identification and sources of dioxin-like compounds: I. Polychlorodibenzothiophenes and polychlorothianthrenes, the sulfur-analogues of the polychlorodibenzofurans and polychlorodibenzodioxins. *Chemosphere*, 25 (1992) 45-48.
- 464 Christensen, K.R. and Reineccius, G.A.: Gas chromatographic analysis of volatile sulfur compounds from heated milk using static headspace sampling. *J. Dairy Sci.*, 75 (1992) 2098-2104; *C.A.*, 117 (1992) 169690v.
- 465 Imanaka, M., Kadota, M., Ogawa, N., Kumashiro, K. and Mori, T.: (Determination of 4- and 5-ring polycyclic aromatic sulfur heterocycles in tea leaves.) *Nippon Eiyo, Shokuryo Gakkaishi*, 45 (1992) 61-70; *C.A.*, 117 (1992) 24858w.
- 466 Liu, Y.: (Determination of organic sulfur compounds in wastewater.) *Zhongguo Huanjing Kexue*, 11 (1991) 384-387; *C.A.*, 116 (1992) 221189j.
- 467 Przyborski, G. and Bandion, F.: (Detection of methyl isothiocyanate in wine.) *Mitt. Klosterneuburg*, 42 (1992) 39-41; *C.A.*, 117 (1992) 149556c.
- 468 Ridgeway, R.G.J., Thornton, D.C. and Bandy, A.R.: Determination of trace aqueous dimethylsulfoxide concentrations by isotope dilution gas chromatography/mass spectrometry: application to rain and sea water. *J. Atmos. Chem.*, 14 (1992) 53-60; *C.A.*, 117 (1992) 33282f.
- 469 Sinkkonen, S., Kolehmainen, E. and Koistinen, J.: Substituted dibenzothiophenes I: synthesis, chromatography, mass spectrometry and structure elucidation by ¹H NMR spectroscopy. *Intern. J. Environ. Anal. Chem.*, 47 (1992) 7-20.
- 470 Sinkkonen, S., Paasivirta, J., Koistinen, J., Lahtipera, M. and Lammi, R.: Polychlorinated dibenzothiophenes in bleached pulp mill effluents. *Chemosphere*, 24 (1992) 1755-1763.
- 471 Slater, G.P.: Allyl thiocyanate - natural product or artifact in crucifer extracts? *Chromatographia*, 34 (1992) 461-467.
- 472 Tanzer, D. and Heumann, K.G.: Gas chromatographic trace-level determination of volatile organic sulfides and selenides and of methyl iodide in Atlantic surface waters. *Intern. J. Environ. Anal. Chem.*, 48 (1992) 17-31.

See also 97, 135, 190, 298, 400, 404, 618, 645, 748, 787, 810.

25. ORGANIC PHOSPHORUS COMPOUNDS (INCL. SUGAR PHOSPHATES)
- 473 Fuchs, G., Doguet, L., Barth, D. and Perrut, M.: Enantiomer fractionation of phosphine oxides by preparative subcritical fluid chromatography. *J. Chromatogr.*, 623 (1992) 329-336.
- 474 Hu, Y. and Yang, X.: (Rapid determination of DDVP in human plasma by gas chromatography.) *Sepu*, 10 (1992) 287-288.
- 475 Willis, T.W.: The effects of structure on retention in the gas chromatographic analysis of alkyl phosphates. *LC-GC*, 10 (1992) 770-773.
- 476 Wils, E.R.J., Hulst, A.G., Verwiël, P.E.J., van Krimpen, S.H. and Niederhauser, A.: Identification of an octyl methylphosphonofluoridate mixture in chemical industry samples. *Fresenius J. Anal. Chem.*, 343 (1992) 297-303.
- See also 635.
26. ORGANOMETALLIC AND RELATED COMPOUNDS
- 26a. *Organometallic compounds*
- 477 Baeyens, W.: Speciation of mercury in different compartments of the environment. *TrAC*, 11 (1992) 245-254 - a review with 56 refs.
- 478 Chakraborti, D.: Gas chromatography-atomic absorption spectrometry (GC-AAS) for specific determination of organolead compounds in the environment. *Chem. Environ. Res.*, 1 (1992) 3-12; C.A., 117 (1992) 136617y.
- 479 Chau, Y.K., Zhang, S. and Maguire, R.J.: Determination of butyltin species in sewage and sludge by gas chromatography-atomic absorption spectrometry. *Analyst (London)*, 117 (1992) 1162-1164.
- 480 David, F. and Sandra, P.: Analysis of organotin compounds using the HP 5921A atomic emission detector. *Analisis*, 20, No. 8 (1992) M53-M54.
- 481 Gomez-Ariza, J.L., Morales, E. and Ruiz-Benitez, M.: GC FPD method for the simultaneous speciation of butyltin and phenyltin compounds in waters. *Appl. Organomet. Chem.*, 6 (1992) 279-286; C.A., 117 (1992) 33284h.
- 482 Hill, S.J.: Analytical aspects of water quality. Advances in coupled techniques for speciation studies. *Anal. Proc.*, 29 (1992) 399-404 - a brief review with 7 refs.
- 483 Jiang, G., Ni, Z., Zhang, L., Li, A., Han, H. and Shan, X.: Determination of alkyl selenides by gas chromatography-electrothermal atomic absorption spectrometry. *J. Anal. At. Spectrom.*, 7 (1992) 447-450; C.A., 117 (1992) 39555u.
- 484 Kim, A., Hill, S., Ebdon, L. and Rowland, S.: Determination of organometallic compounds by capillary gas chromatography-inductively coupled plasma mass spectrometry. *J. High Resolut. Chromatogr.*, 15 (1992) 665-668.
- 485 Liou, D.C., Lou, Y.C. and Whang, C.W.: Determination of trace butyltin compounds in sea water by gas chromatography-atomic absorption spectrometry. *J. Chin. Chem. Soc. (Taipei)*, 39 (1992) 217-222; C.A., 117 (1992) 55512h.
- 486 Mori, H.: (Gas chromatograph prefiler for analysis column of alkylmercury and its use.) *Jpn. Kokai Tokkyo Koho JP 04,160,360 [92,160,360]* (Cl. G01N30/14), 03 Jun. 1992, Appl. 90/286,641, 23 Oct. 1990; 5 pp.; C.A., 117 (1992) 123811g.
- 487 Shiraiishi, H., Higashivama, T., Hashimoto, S. and Otsuki, A.: (Identification of metabolites of organotin compounds in mussel by GC/MS.) *Shitsuryo Bunseki*, 40 (1992) 137-146; C.A., 117 (1992) 125977q.
- See also 99, 105, 803.
- 26b. *Boranes, silanes and related non-metallic compounds*
- 488 Aggarwal, S.K., Kinter, M. and Herold, D.A.: Determination of selenium in urine by isotope dilution gas chromatography-mass spectrometry using 4-nitro-o-phenylenediamine, 3,5-dibromo-o-phenylenediamine, and 4-trifluoromethyl-o-phenylenediamine as derivatizing reagents. *Anal. Biochem.*, 202 (1992) 367-374.
- 489 Dai, K., Li, Y. and Shi, W.: (Gas-liquid chromatographic determination of trace arsenic after derivatization with 2,3-dimercaptopropanol.) *Sepu*, 10 (1992) 218-220.
- 490 Kaise, T. and Fukui, S.: The chemical form and acute toxicity of arsenic compounds in marine organisms. *Appl. Organomet. Chem.*, 6 (1992) 155-160; C.A., 117 (1992) 21536d.
- 491 Yamamoto, M., Tanaka, S. and Hashimoto, Y.: Selective determination of trace arsenic and antimony species in natural waters by gas chromatography with a photoionization detector. *Appl. Organomet. Chem.*, 6 (1992) 351-356; C.A., 117 (1992) 96982e.
- See also 472.
- 26c. *Coordination compounds*
- 492 Kito, A.: (Thermogravimetric analyses and gas-liquid chromatographic behavior of the copper (II) chelates of 1-(fluorophenyl)-4,4-dimethyl-1,3-pentanediones.) *Osaka Kogyo Gijutsu Shikensho Kiho*, 42 (1991) 136-141; C.A., 117 (1992) 61491s.
- 493 Staniewski, J.: (Gas chromatographic analysis of some hydroxyoxime extractants of metals.) *Chem. Anal. (Warsaw)*, 36 (1991) 325-333; C.A., 117 (1992) 39528n.
- 494 Uden, P.C. and Zeng, Y.: A study of diastereoisomerism of oxovanadium Schiff base chelates by gas chromatography-atomic emission spectroscopy, and high performance liquid chromatography. *Chromatographia*, 34 (1992) 269-275.
- See also 99, 105, 803.
27. VITAMINS AND VARIOUS ANIMAL GROWTH FACTORS (NON-PEPTIDIC)
- 495 Jiang, Z.: (Determination of fat-soluble vitamin E in sea buckthorn oil and variation in vitamin E content in refined oil.) *Linchuan Huaxue Yu Gongye*, 11 (1991) 241-246; C.A., 117 (1992) 89057d.
- 496 Leyland, D.M., Evershed, R.P., Edwards, R.H.T. and Beynon, R.J.: Application of gas chromatography-mass spectrometry with selected ion monitoring to the urinalysis of 4-pyridoxic acid. *J. Chromatogr.*, 581 (1992) 179-185.

See also 657.

29. INSECTICIDES, PESTICIDES AND OTHER AGROCHEMICALS

29a. General techniques

- 497 Bernal, J.L., del Nozal, M.J. and Jiménez, J.J.: Some observations on clean-up procedures using sulphuric acid and Florisil. *J. Chromatogr.*, 607 (1992) 303-309.
- 498 Gerhards, P.: (Bestimmung von Pestiziden mit GC und GC/MS.) *LaborPraxis*, 16 (1992) 1014-1018.
- 499 Germann, P., Bezegh, A. and Clerc, J.T.: Optimization of a GC separation of pesticides using a central composite design. *Pharm. Acta Helv.*, 67 (1992) 113-115; *C.A.*, 117 (1992) 106221x.
- 500 Gremm, T.J. and Frimmel, F.H.: Determination of organotin compounds in aqueous samples by means of HPGC-AED. *Water Res.*, 26 (1992) 1163-1169; *C.A.*, 117 (1992) 137230d.
- 501 Kenmotsu, K., Kobara, K., Ogino, Y. and Mori, T.: (Analysis of pesticides in effluent of golf links.) *Kankyo Kagaku*, 2 (1992) 412-413; *C.A.*, 117 (1992) 157252r.

See also 19, 676, 700, 714, 720.

29b. Chlorinated insecticides

- 502 Bernal, J.L., del Nozal, M.J. and Jiménez, J.J.: Use of a high-pressure Soxhlet extractor for the determination of organochlorine residues by gas chromatography. *Chromatographia*, 34 (1992) 468-474.
- 503 Galaktionova, M.A. and Mevinnaya, L.V.: (Rapid gas-chromatographic method for determination of organochlorine insecticides in water.) *Khim. Tekhnol. Vody*, 13 (1991) 987-991; *C.A.*, 116 (1992) 091010f.
- 504 Huehnerfuss, H., Faller, J., Koenig, W.A. and Ludwig, P.: Gas chromatographic separation of the enantiomers of marine pollutants. 4. Fate of hexachlorocyclohexane isomers in the Baltic and North Sea. *Environ. Sci. Technol.*, 26 (1992) 2127-2133; *C.A.*, 117 (1992) 177866x.
- 505 Ludwig, P., Huehnerfuss, H., Koenig, W.A. and Gunkel, W.: Gas chromatographic separation of the enantiomers of marine pollutants. Part 3. Enantioselective degradation of α -hexachlorocyclohexane and γ -hexachlorocyclohexane by marine microorganisms. *Mar. Chem.*, 38 (1992) 13-23; *C.A.*, 117 (1992) 118009x.
- 506 Mochizuki, E. and Kondo, M.: (Determination of chlorinated pesticides by capillary gas chromatography.) *Yamanashi-ken Eisei Kogai Kenkyusho Nenpo*, 34 (1990) 11-15; *C.A.*, 117 (1992) 64702x.
- 507 Nwankwoala, A.U. and Osibanjo, O.: Baseline levels of selected organochlorine pesticide residues in surface waters in Ibadan (Nigeria) by electron capture gas chromatography. *Sci. Total Environ.*, 119 (1992) 179-190; *C.A.*, 117 (1992) 55464u.
- 508 Poleze, L., Ribeiro, M.L., de Toledo, F., Regina, C., de Souza, C. and Joaquim, T.: (Rapid procedure for determination of heptachlor, aldrin and mirex in sugarcane rum.) *Ecletica Quim.*, 16 (1991) 55-62; *C.A.*, 117 (1992) 169691w.
- 509 Robbat, A., Jr., Liu, C. and Liu, T.-Y.: Field detection of organochlorine pesticides by thermal desorption gas chromatography-mass spectrometry. *J. Chromatogr.*, 625 (1992) 277-288.

- 510 Rymen, T., Clark, S., Boenke, A., Wagstaffe, P.J. and Lindsey, A.S.: Reference materials for PCB analysis: Production and certification of ten polychlorinated biphenyls in an iso-octane reference solution. *Fresenius J. Anal. Chem.*, 343 (1992) 553-560.
- 511 Schmidt, L.J. and Hesselberg, R.J.: A mass spectroscopic method for analysis of AHH-inducing and other polychlorinated biphenyl congeners and selected pesticides in fish. *Arch. Environ. Contam. Toxicol.*, 23 (1992) 37-44; *C.A.*, 117 (1992) 42064p.
- 512 Tan, G.H.: Comparison of solvent extraction and solid-phase extraction for the determination of organochlorine pesticide residues in water. *Analyst (London)*, 117 (1992) 1129-1132.

See also 171, 186, 242, 246, 257, 261, 675, 723, 730.

29c. Phosphorus insecticides

- 513 Allender, W.J. and Keegan, J.: Chromatographic analysis of *cis*- and *trans*- mevinphos in poisoned wildlife. *J. Chromatogr.*, 609 (1992) 315-320.
- 514 Allender, W.J., Major, E.J. and Cresswell, G.C.: Measurement of acephate and its principal metabolite in leaves by gas-liquid chromatography. *Pestic. Sci.*, 35 (1992) 91-94; *C.A.*, 117 (1992) 42578j.
- 515 Durand, G., Alonso, R. and Barcelo, D.: (Extraction and analysis of organophosphorus and triazine pesticides in water and soils.) *Quim. Anal. (Barcelona)*, 10 (1991) 157-168; *C.A.*, 117 (1992) 85102s.
- 516 Hasegawa, T., Namiki, A. and Itoh, K.: (Dichlorvos determination in air by gas chromatography.) *Nippon Kankyo Eisei Senta Shoho*, 18 (1991) 37-39; *C.A.*, 117 (1992) 54842d.
- 517 Hayakawa, T., Yoshimoto, M. and Nakamura, Y.: (Simultaneous determination of organophosphorus pesticides (chlorpyrifos-methyl and pyrimiphos-methyl) in feed by capillary column gas chromatography.) *Shiroyo Kenkyu Hokoku (Tokyo Hishiryu Kensasho)*, 17 (1992) 12-22; *C.A.*, 117 (1992) 149585m.
- 518 Heikes, D.L. and Craun, J.C.: Rapid multiresidue procedure for the determination of pesticides in anhydrous lanolin and lanolin-containing pharmaceutical preparations utilizing gel permeation chromatography cleanup with gas chromatographic and mass spectrometric techniques. *J. Agric. Food Chem.*, 40 (1992) 1586-1590.
- 519 Kojima, F. and Ishida, Y.: (Simultaneous determination of organophosphorus pesticides (diazinon, fenitrothion, malathion, MPP, parathion, PAP, EPN and DDVP) in feed by capillary column gas chromatography.) *Shiroyo Kenkyu Hokoku (Tokyo Hishiryu Kensasho)*, 17 (1992) 26-37; *C.A.*, 117 (1992) 130054r.
- 520 Morchio, G., de Andreis, R. and Verga, G.R.: (Organophosphorus residues in vegetable oils, with special reference to olive oil.) *Riv. Ital. Sostanze Grasse*, 69 (1992) 147-157.
- 521 Steinwandter, H.: Contribution to the ethyl acetate application in residue analysis. II. Micro on-line method for extracting organophosphorus pesticides from fruits, vegetables and feedstuffs. *Fresenius J. Anal. Chem.*, 343 (1992) 887-889.
- 522 Tieffova, P. and Matousek, M.: (A method for routine determination of ethephon residues in plant material.) *Agrochimia (Bratislava)*, 31 (1991) 251-254; *C.A.*, 116 (1992) 146022u.

- 523 Tsutsui, T., Ohta, M. and Adachi, T.: Cleanup procedure for determination of pesticide residues in vegetables by charcoal-Florisisil column. *Kyoto-fu Eisei Kogai Kenkyusho Nenpo*, 36 (1991) 35-42; C.A., 117 (1992) 46832k.

See also 219, 635, 730.

29d. Carbamates

- 524 Czege, G.: (Determination of urethane in fruit brandy by capillary gas chromatography with a nitrogen-selective detector.) *Szeszipar*, 39 (1991) 113-115; C.A., 117 (1992) 24845q.
- 525 Sancho, M.T., Muniategui, S., Huidobro, J.F. and Simal, J.: (Determination of fluvalinate residues in honey by GC/ECD.) *Rev. Agroquim. Technol. Aliment.*, 31 (1991) 417-422; C.A., 116 (1992) 172534g.

29e. Herbicides

- 526 An, Q. and Qian, W.: (A procedure for determination of trifluralin residues in soil by capillary GC.) *Turang (Nanjing)*, 24, No. 1 (1992) 48-50; C.A., 117 (1992) 165836z.
- 527 Firor, R.L.: (Evaluation of soil extracts containing 2,4,5-trichlorophenoxyacetic acid with an electron-capture detector.) *Qium. Ind. (Madrid)*, 38 (1992) 113-117; C.A., 117 (1992) 145236a.
- 528 Kawata, K.: (Simple GC and GC/MS determination of herbicides in air using quartz fiber filter and activated carbon fiber filter for sampling.) *Kankyo Kagaku*, 2 (1992) 181-186; C.A., 117 (1992) 106214x.
- 529 Kumazawa, T., Sato, K., Seno, H. and Suzuki, O.: Rapid isolation with Sep-Pak C₁₈ cartridges and capillary gas chromatography of triazine herbicides in human body fluids. *Forensic Sci. Int.*, 54 (1992) 159-166; C.A., 117 (1992) 144949y.
- 530 Stankova, O. and Smickova, J.: Determination of s-triazines in waters. *Acta Univ. Palacki. Olomuc., Fac. Rerum Nat.*, 102 (1991) 109-119; C.A., 117 (1992) 157250p.
- 531 Steinwandter, H.: Contribution to the ethyl acetate application in residue analysis. I. Micro method for extracting atrazine residues from soil samples. *Fresenius J. Anal. Chem.*, 343 (1992) 604-606.

See also 180, 515, 713.

29f. Fungicides

- 532 Allmendinger, H.: A method for determining residues of the fungicides Follicur and Bayfidan in plant material and soil by gas chromatography. *Pflanzenschutz-Nachr. Bayer*, 44 (1991) 5-66; C.A., 117 (1992) 21826y.
- 533 Butter, W. and Walker, G.: The determination of wood preserving agents on the surface of wood by gas chromatography and gas chromatography-mass spectrometry. *J. Anal. Chem.*, 343 (1992) 144.
- 534 Miyahara, M., Suzuki, T. and Saito, Y.: Capillary gas chromatographic determination of captafol in vegetables, fruits, and grains. *J. Agric. Food Chem.*, 40 (1992) 1150-1153.

See also 219, 241, 662.

29g. Other types of pesticides and various agrochemicals

- 535 Wells, C., Bertsch, W. and Perich, M.: Isolation of volatiles with insecticidal properties from the genus *Tagetes* (marigold). *Chromatographia*, 34 (1992) 241-248.

31. PLASTICS AND THEIR INTERMEDIATES

- 536 Ahmadi, A., Jalayer, P. and Mahabadi, P.: (Pyrolysis gas chromatography of polymers.) *Iran. J. Polym. Sci. Technol.*, 4 (1992) 272-278; C.A., 117 (1992) 131814a.
- 537 Barth, H.G. and Mays, J.W.: *Modern Methods of Polymer Characterization*. Wiley, New York, 1991, 547 p.
- 538 Becerra, M.R., Fernandez-Sanchez, E., Fernandez-Torres, A., Garcia-Dominguez, J.A. and Santiuste, J.M.: Thermodynamic characterization by inverse gas chromatography of a 50 % methyl, 50 % trifluoropropyl polysiloxane. *Macromolecules*, 25 (1992) 4665-4670; C.A., 117 (1992) 112364n.
- 539 Chehimi, M.M., Pigois-Landureau, E. and Delamar, M.: (Characterization of conducting polymers by inverse gas chromatography. Part 1. Despersive and acid-base properties of conducting polypyrrole and its related insulating form.) *J. Chim. Phys. Phys.-Chim. Biol.*, 89 (1992) 1173-1178; C.A., 117 (1992) 80968u.
- 540 Chehimi, M.M., Pigois-Landureau, E., Delamar, M., Watts, J.F., Jenkins, S.N. and Gibson, E.M.: The use of x-ray photoelectron spectroscopy, Fourier transform infra-red spectroscopy and inverse gas chromatography in the study of acid-base interactions of polymers, solvents and metal oxides. *Bull. Soc. Chim. Fr.*, 129 (1992) 137-144; C.A., 117 (1992) 49518y - a review with 46 refs.
- 541 Etxeberria, A., Alfageme, J., Uriarte, C. and Iruin, J.J.: Inverse gas chromatography in the characterization of polymeric materials. *J. Chromatogr.*, 607 (1992) 227-237.
- 542 Gangrade, N. and Price, J.C.: Simple gas chromatographic headspace analysis of residual organic solvent in microspheres. *J. Pharm. Sci.*, 81 (1992) 201-202; C.A., 117 (1992) 76608j.
- 543 Haken, J.K. and Koopetngarm, S.: An approach to the analysis of crosslinked epoxy systems. *Chromatographia*, 34 (1992) 276-280.
- 544 Ji, Y. and Liu, W.: (Identification of imide-containing aromatic polyether-polyketone and polyether-polysulfone by pyrolysis gas chromatography-mass spectrometry.) *Fenxi Huaxue*, 20 (1992) 28-31; C.A., 117 (1992) 70636v.
- 545 Kawarai, N. and Yamamoto, H.: (Discrimination of familiar plastics.) *Kagaku to Kyoiku*, 40 (1992) 330-331; C.A., 117 (1992) 69191q.
- 546 Mao, Y. and Wang, H.: (Inverse gas chromatographic determination of the solubility parameter of hydroxyl-terminated polybutadiene.) *Fenxi Ceshi Tongbao*, 10, No. 3 (1991) 46-50; C.A., 116 (1992) 060516w.
- 547 Mestdagh, H., Rolando, C., Sablier, M. and Rioux, J.-P.: Characterization of ketone resins by pyrolysis/gas chromatography/mass spectrometry. *Anal. Chem.*, 64 (1992) 2221-2226.

- 548 Mueller, T.W. and Fischer, S.A.: Determination of organic chloride impurities in polyaminoamide-epichlorohydrin resins by gas chromatography. *Tappi J.*, 75 (1992) 159-162; *C.A.*, 117 (1992) 91364v.
- 549 Nita, M., Negoescu, E., Chiriac, G. and Botea, M.: (Use of gas chromatography for the determination of molecular mass of Romanian polyethylene glycol.) *Rev. Chim. (Bucharest)*, 42 (1991) 336-339; *C.A.*, 117 (1992) 91167h.
- 550 Ozdemir, E., Acikses, A. and Coskun, M.: Determination of polystyrene-hydrocarbon interaction parameters and solubility parameter using inverse gas chromatography. *Macromol. Rep.*, A29 (1992) 63-70; *C.A.*, 117 (1992) 49650k.
- 551 Pasch, H., Brinkmann, C., Much, H. and Just, U.: Chromatographic investigations of macromolecules in the "critical range" of liquid chromatography. II. Two-dimensional separations of poly(ethylene oxide-block-propylene oxide). *J. Chromatogr.*, 623 (1992) 315-322.
- 552 Ryabikova, V.M., Zigel, A.N., Pirozhnaya, L.N. and Popova, G.S.: (Determination of the composition of bicomponent copolymers of tetrafluoroethylene with hexafluoropropylene, vinylidene fluoride and perfluoromethyl vinyl ether by pyrolytic gas chromatography method.) *Vysokomol. Soedin., Ser. B*, 34, No. 3 (1992) 54-59; *C.A.*, 117 (1992) 131849r.
- 553 Sato, S.: (Analysis of polymer components by various derivatization by Py-GC.) *Idemitsu Gihō*, 35 (1992) 211-216; *C.A.*, 117 (1992) 172336q.
- 554 Shchors, T.V., Tolchinskii, Yu.I., Kabachii, Yu.A., Gerner, E.Yu. and Val'kovskii, D.G.: (Equipment and technique for the study of thermochemical transformations of polymers and low-molecular-weight compounds using gas-chromatographic analysis of pyrolysis products.) *Vysokomol. Soedin., Ser. A*, 34, No. 4 (1992) 146-151; *C.A.*, 117 (1992) 151569c.
- 555 Silver, A.H. and Kalinoski, H.T.: Comparison of high-temperature gas chromatography and carbon dioxide supercritical fluid chromatography for the analysis of alcohol ethoxylates. *J. Am. Oil Chem. Soc.*, 69 (1992) 599-608.
- 556 Usami, T., Keitoku, F., Ohtami, H. and Tsuge, S.: Characterization of poly(2,6-dimethyl-1,4-phenylene ether) by high resolution pyrolysis-gas chromatography (HR-Pyg.c.) *Polymer*, 33 (1992) 3024-3030; *C.A.*, 117 (1992) 131878z.
- 557 Ute, K., Miyatake, N., Asada, T. and Hatada, K.: Stereoregular oligomers of methyl methacrylate. 6. Isolation of isotactic and syndiotactic methyl methacrylate oligomers from 19-mer to 29-mer by preparative supercritical fluid chromatography and their thermal analysis. *Polym. Bull. (Berlin)*, 28 (1992) 561-568; *C.A.*, 117 (1992) 112402y.
- 558 Wright, D.W., Mahler, K.O. and Davis, J.: The application of multidimensional GC techniques to the analysis of sub-PPB levels of vinyl chloride monomer in polyvinylchloride. *J. Chromatogr. Sci.*, 30 (1992) 291-295.
- 559 Xu, Z. and Zhang, L.: (Pyrolysis-gas chromatography of polynaphthalene.) *Sepu*, 10 (1992) 293-294.
- 560 Yang, X.: (Study on the effects of ACR on the heat stability and intermiscibility of PVC by pyrolysis gas chromatography (PGC).) *Sepu*, 10 (1992) 220-222.
- 561 Zaikin, V.G., Mardanov, R.G., Yakovlev, V.A. and Plate, N.A.: Composition and microstructure of butadiene-isoprene copolymers from pyrolysis-gas chromatography/mass spectrometric data. *J. Anal. Appl. Pyrolysis*, 23 (1992) 33-42.
- See also 33, 375, 776.
32. DRUG ANALYSIS
- 32a. *Drug analysis, general techniques*
- 562 Berridge, M.S., Cassidy, E.H. and Kozirowski, J.: Simple automatic quality control for radiolabeled gases for PET. *Appl. Radiat. Isot.*, 43 (1992) 1055-1056; *C.A.*, 117 (1992) 127539x.
- 563 Billot, P. and Pitard, B.: Taguchi design experiments for optimizing the gas chromatographic analysis of residual solvents in bulk pharmaceuticals. *J. Chromatogr.*, 623 (1992) 305-313.
- 564 Lavine, B.K.: Environmental applications of pattern recognition techniques. *Chemometrics Intel. Lab. Systems*, 15 (1992) 219-230.
- 565 Ramstad, T. and Walker, J.S.: Investigation of musty odour in pharmaceutical products by dynamic headspace gas chromatography. *Analyst (London)*, 117 (1992) 1361-1366.
- 566 Svinarov, D.A.: Unified liquid chromatographic and gas chromatographic assays for therapeutic drug monitoring and toxicology. In: Sunshine, I. (Editor), *Recent Dev. Ther. Drug Monit. Clin. Toxicol.*, Dekker, New York, 1992, 691-702 p.; *C.A.*, 117 (1992) 62194j.
- See also 260, 773.
- 32b. *Antirheumatics and antiinflammatory drugs*
- 567 Moreau, J., Palette, C., Cordonnier, P., Naline, E., Advenier, C. and Pays, M.: Separation and identification of the 4-hydroxyantipyrine sulfoconjugate. *J. Chromatogr.*, 576 (1992) 103-109; *C.A.*, 117 (1992) 019756g.
- 568 Newkirk, M.S.: Application of methods for determining total organic contribution to diesel particulates. *J. Eng. Gas Turbines Power*, 114 (1992) 578-589; *C.A.*, 117 (1992) 156694z.
- 32c. *Autonomic and cardiovascular drugs*
- 569 Bi, H., Cooper, S.F. and Côté, M.G.: Determination and identification of amiloride in human urine by high-performance liquid chromatography and gas chromatography-mass spectrometry. *J. Chromatogr.*, 582 (1992) 93-101.
- 570 Cody, J.T.: Determination of methamphetamine enantiomer ratios in urine by gas chromatography-mass spectrometry. *J. Chromatogr.*, 580 (1992) 77-95 - a review with 56 refs.
- 571 DePace, A., Verebey, K. and ElSohly, M.: Capillary gas-liquid chromatography separation of phenethylamines in amphetamine-positive urine samples. *J. Forensic Sci.*, 35 (1990) 1431-1435; *C.A.*, 117 (1992) 84788h.
- 572 Ho, J.W. and Moody, D.E.: Gas chromatography/mass spectrometry assays for the determination of debrisoquine and sparteine metabolites in microsomal fractions of rat liver. *Anal. Biochem.*, 203 (1992) 348-351.
- 573 Masayuki, M., Horimoto, S., Mabuchi, M. and Banno, K.: Determination of three metabolites of a new angiotensin converting enzyme inhibitor, imidapril, in plasma and urine by gas chromatography-mass spectrometry using multiple ion detection. *J. Chromatogr.*, 581 (1992) 65-73.

- 574 Quaglio, M.P., Bellini, A.M. and Minozzi, L.: Simultaneous determination of propranolol or metoprolol in the presence of benzodiazepines in the plasma by gas chromatography. *Farmaco*, 47 (1992) 799-809; C.A., 117 (1992) 142783r.
- 575 Rudolph, M., Janssen, W. and Strassner, M.: Determination of moxonidine (BDF 5895) in plasma by gas chromatography-negative ion chemical ionization mass spectrometry. *J. Pharm. Biomed. Anal.*, 10 (1992) 323-328; C.A., 117 (1992) 142793u.
- 576 Siret, L., Bargmann, N., Tambute, A. and Caude, M.: Direct enantiomeric separation of β -blockers on ChyRoSine-A by supercritical chromatography: supercritical carbon dioxide as transient in situ derivatizing agent. *Chirality*, 4 (1992) 252-262; C.A., 117 (1992) 178393c.
- 577 Sutton, B.M. and Richardson, R.A.: Short communication. Assay of timolol in human plasma using gas chromatography with electron-capture detection. *J. Chromatogr.*, 581 (1992) 277-280.
- 578 Tsai, F.Y., Lui, L.F. and Chang, B.: Analysis of diuretic doping agents by HPLC screening and GC-MSD confirmation. *J. Pharm. Biomed. Anal.*, 9 (1991) 1069-1076; C.A., 117 (1992) 21565n.
- 579 Wu, A.H.B., Wong, S.S., Johnson, K.G., Ballatore, A. and Seifert, W.E.J.: The conversion of ephedrine to methamphetamine and methamphetamine-like compounds during and prior to gas chromatographic /mass spectrometric analysis of CB and HFB derivatives. *Biol. Mass Spectrom.*, 21 (1992) 278-284; C.A., 117 (1992) 42077v.
- See also 423.
- 32d. *Central nervous system drugs*
- 580 Abdel-Moety, E.M., El-Tarras, M.F., El-Zeany, B.E.A. and Kelan, K.O.: First derivative spectrophotometric and gas-liquid chromatographic determination of caffeine in foods and pharmaceuticals. III. Simultaneous assay of caffeine and some antihistaminics. *Bull. Fac. Pharm.*, 28, No. 2 (1990) 1-5; C.A., 117 (1992) 33801z.
- 581 Baugh, L.D., Liu, R.H. and Walia, A.S.: Simultaneous gas chromatography/mass spectrometry assay of methadone and 2-ethylidene-1,5-dimethyl-3,3-diphenylpyrrolidine (EDDP) in urine. *J. Forensic Sci.*, 36 (1991) 548-555; C.A., 117 (1992) 105843q.
- 582 Beischlag, T.V. and Inaba, T.: Determination of nonderivatized *p*-hydroxylated metabolites of diazepam in biological fluids with a GC megabore column system. *J. Anal. Toxicol.*, 16 (1992) 236-239; C.A., 117 (1992) 103468x.
- 583 Berberic, D. and Uhrich, M.: (Analysis of phencyclidine (PCP) and codeine/morphine in urine by gas chromatography/ion-trap mass spectrometry.) *Lab. 2000*, 5, No. 7 (1991) 98-103; C.A., 117 (1992) 84748v.
- 584 Byrd, G.D., Paule, R.C., Sander, L.C., Sniegoski, L.T. and White, V.E.: Determination of 3-quinuclidinyl benzilate (QNB) and its major metabolites in urine by isotope dilution gas chromatography/mass spectrometry. *J. Anal. Toxicol.*, 16 (1992) 182-187; C.A., 117 (1992) 144876x.
- 585 Caturla, M.C., Salomo, J., Soler, M., Martinez, L. and Roser, R.: Validation of GC method to measure a new analgesic (E-4018) in biological samples. Comparison between HPLC and GC method. *J. Pharm. Biomed. Anal.*, 9 (1991) 1181-1184; C.A., 117 (1992) 19771h.
- 586 Choo, H.Y.P., Choi, J., Choi, M.J., Song, E.Y. and Park, J.: Quantitation of barbiturates in urine by GC/MS and its comparison to fluorescence polarization immunoassay. *Korean J. Toxicol.*, 7 (1991) 29-35; C.A., 117 (1992) 125981m.
- 587 Cyr, T.D., Lawrence, R.C. and Lovering, E.G.: Gas chromatographic methods for doxepin isomers, related compounds, and organic volatile impurities in raw materials and doxepin isomers in capsules. *J. Assoc. Off. Anal. Chem.*, 75 (1992) 804-809.
- 588 Duthel, J.M., Constant, H., Vallon, J.J., Rochet, T. and Miachon, S.: Quantitation by gas chromatography with selected-ion monitoring mass spectrometry of "natural" diazepam, N-desmethyldiazepam and oxazepam in normal human serum. *J. Chromatogr.*, 579 (1992) 85-91.
- 589 Geary, R. and Stavchansky, S.: Gas-liquid chromatographic determination of almitrine in rat intestinal fluid using flame ionization detection. *Anal. Lett.*, 25 (1992) 1843-1850.
- 590 Gjerde, H., Dahlin, E. and Christophersen, A.S.: Simultaneous determination of common benzodiazepines in blood using capillary gas chromatography. *J. Pharm. Biomed. Anal.*, 10 (1992) 317-322; C.A., 117 (1992) 144903d.
- 591 Hattori, H., Hoshino, N., Seno, H., Suzuki, O. and Yamada, T.: Capillary gas chromatography of underivatized barbiturates in human blood with a nitrogen-phosphorus detector and with splitless injection. *Hochudoku*, 10, No. 1 (1992) 16-22; C.A., 117 (1992) 84779f.
- 592 Hattori, H., Yamamoto, S., Iwata, M., Takashima, E., Yamada, T. and Suzuki, O.: Determination of diphenylmethane antihistaminic drugs and their analogues in body fluids by gas chromatography with surface ionization detection. *J. Chromatogr.*, 581 (1992) 213-218.
- 593 Hattori, H., Yamamoto, S., Iwata, M., Takashima, E., Yamada, T. and Suzuki, O.: Sensitive determination of phenothiazines in body fluids by gas chromatography with surface ionization detection. *J. Chromatogr.*, 579 (1992) 247-252.
- 594 Hayes, M.J., Khemani, L., Leal, M. and Powell, M.L.: Quantitative determination of a new anticonvulsant (CGS 18416A) in human plasma using capillary gas chromatography/mass spectrometry. *Biomed. Chromatogr.*, 6 (1992) 236-240.
- 595 Hewala, I.I.: GLC and HPLC determination of diazepam and its degradation products in pharmaceutical formulations. *Anal. Lett.*, 25 (1992) 1877-1905.
- 596 Jagota, N.K. and Stewart, J.T.: Analysis of diazepam and chlor-diazepoxide and their related compounds using supercritical fluid chromatography. *J. Liquid Chromatogr.*, 15 (1992) 2429-2443.
- 597 Kintz, P., Tracqui, A. and Mangin, P.: Determination of nalbuphine using high-performance liquid chromatography coupled to photodiode-array detection and gas chromatography coupled to mass spectrometry. *J. Chromatogr.*, 579 (1992) 172-176.
- 598 Kokatsu, J., Yomoda, R. and Suwa, T.: Selected ion monitoring for the determination of bromovaleerylurea in human plasma. *Chem. Pharm. Bull.*, 40 (1992) 1517-1519; C.A., 117 (1992) 142786u.
- 599 Kristensen, K. and Angelo, H.R.: Stereospecific gas chromatographic method for determination of methadone in serum. *Chirality*, 4 (1992) 263-267; C.A., 117 (1992) 105808g.

- 600 Kumazawa, T., Seno, H. and Suzuki, O.: Rapid isolation with Sep-Pak C₁₈ cartridges and wide-bore capillary gas chromatography of bromisovalum. *J. Anal. Toxicol.*, 16 (1992) 163-165; C.A., 117 (1992) 21571m.
- 601 Leal, M., Hayes, M.J. and Powell, M.L.: Quantitative determination of CGS 18102A, a new anxiolytic, in human plasma using capillary gas chromatography/mass spectrometry. *Biomed. Chromatogr.*, 6 (1992) 244-247.
- 602 Markovic, S. and Kusec, Z.: Determination of butalbital, caffeine and propyphenazone in pharmaceutical preparations by gas chromatography. *Pharmazie*, 46 (1991) 886; C.A., 117 (1992) 118597n.
- 603 Nelson, C.C. and Foltz, R.L.: Chromatographic and mass spectrometric methods for determination of lysergic acid diethylamide (LSD) and metabolites in body fluids. *J. Chromatogr.*, 580 (1992) 97-109; - a review with 41 refs.
- 604 Park, J., Choi, J., Choi, M.J., Song, E.Y. and Choo, H.Y.P.: Screening of benzodiazepines in urine by the immunoassay and quantitation by GC-NPD method. *Korean J. Toxicol.*, 7 (1991) 21-27; C.A., 117 (1992) 84774a.
- 605 Pokrajac, M., Miljkovic, B., Spiridonovic, D. and Varagic, V.M.: An improved gas chromatographic determination of valproic acid and valpromide in plasma. *Pharm. Acta Helv.*, 67 (1992) 237-240; C.A., 117 (1992) 163296z.
- 606 Roca, R.M.S., March, F.J.G., Fos, G.M.A., Domenech, R.G., Gimenez, F.P. and Alvarez, J.G.: Molecular topology and chromatographic retention parameters for benzodiazepines. *J. Chromatogr.*, 607 (1992) 91-95.
- 607 Singh, A.K., Granley, K., Misra, U., Naeem, K., White, T. and Jiang, Y.: Screening and confirmation of drugs in urine: interference of hordenine with the immunoassays and thin layer chromatography methods. *Forensic Sci. Int.*, 54 (1992) 9-22; C.A., 117 (1992) 126011g.
- 608 Soine, W.H., Duncan, W., Lambert, R., Middleberg, R., Finley, H. and O'Neil, D.J.: Differentiation of side chain isomers of ring-substituted amphetamines using gas chromatography/infrared/mass spectrometry (GC/IR/MS). *J. Forensic Sci.*, 37 (1992) 513-527; C.A., 117 (1992) 144937t.
- 609 Stephenson, J. and Tillmanns, U.: (Drug analysis with gas chromatography/mass spectrometry (GC/MS). Part 6. Barbiturates: identification and quantification with GC/MS.) *GIT Fachz. Lab.*, 36, No. 2 (1992) 130-139; C.A., 117 (1992) 105798d.
- 610 Torok-Both, G.A., Baker, G.B., Coutts, R.T., McKenna, K.F. and Launa, J.A.: Simultaneous determination of fluoxetine and norfluoxetine enantiomers in biological samples by gas chromatography with electron-capture detection. *J. Chromatogr.*, 579 (1992) 99-106.
- 611 Trinh, V.D. and Vernay, A.: A safer methylation procedure with boron trifluoride-methanol reagent for gas chromatographic analysis of ritalinic acid in urine. *J. Pharm. Biomed. Anal.*, 10 (1992) 187-191; C.A., 117 (1992) 82824t.
- 612 Vandenbergh, H. and Macdonald, J.C.: Simultaneous determination of clobazam and its metabolite desmethylclobazam in serum by gas chromatography with electron-capture detection. In: Sunshine, I. (Editor), *Recent Dev. Ther. Drug Monit. Clin. Toxicol.*, Dekker, New York, 1992, pp. 469-474; C.A., 117 (1992) 62190e.
- 613 Walters-Thompson, K.M. and Mason, W.D.: Method for the determination of diphenhydramine in rabbit whole blood by high-performance liquid chromatography (HPLC) with ultraviolet (UV) detection in conjunction with gas chromatography (GC) with mass selective detection (MSD). *Pharm. Res.*, 9 (1992) 929-932; C.A., 117 (1992) 103447q.
- 614 Wong, Y.C., Lau, O.W., Chan, K., Aun, C., Lowe, D.M. and Houghton, I.T.: Improved gas chromatographic method for the quantitation of plasma pethidine and norpethidine: application in determining pharmacokinetic parameters in Chinese patients after intramuscular administration of pethidine. *J. Biopharm. Sci.*, 1 (1990) 267-276; C.A., 117 (1992) 82808r.
- See also 178, 453, 622, 628, 638.
- 32e. *Chemotherapeutics (exc. cytostatics and antibiotics)*
- 615 Cheng, K.N., Elsom, L.F. and Hawkins, D.R.: Identification of metabolites of halofantrine, a new candidate anti-malarial drug, by gas chromatography-mass spectrometry. *J. Chromatogr.*, 581 (1992) 203-211.
- 616 Patterson, T.A. and Kosh, J.W.: Simultaneous quantitation of arecoline, acetylcholine, and choline in tissue using gas chromatography/electron impact mass spectrometry. *Biol. Mass Spectrom.*, 21 (1992) 299-304; C.A., 117 (1992) 39693n.
- 617 Schenkel, E., Duez, P. and Hanocq, M.: Stereoselective determination of cucurbitine in *Cucurbita* spp. seeds by gas chromatography and gas chromatography-mass spectrometry. *J. Chromatogr.*, 625 (1992) 289-298.
- 32f. *Cytostatics*
- 618 Weinberg, D.S., Manier, M.L., Richardson, M.D., Haibach, F.G. and Rogers, T.S.: Identification and quantification of anticarcinogens in garlic extract and licorice root extract powder. *J. High Resolut. Chromatogr.*, 15 (1992) 641-654.
- 32g. *Other drug categories*
- 619 Bankova, V., Christov, R., Stoev, G. and Popov, S.: Determination of phenolics from propolis by capillary gas chromatography. *J. Chromatogr.*, 607 (1992) 150-153.
- 620 Hagedorn, H.W. and Schulz, R.: Detection of diuretics in horse urine by GC/MS. *J. Anal. Toxicol.*, 16 (1992) 194-198; C.A., 117 (1992) 42078w.
- 621 Meyer, E., de Leenheer, A.P. and Sandra, P.: Aspects of the capillary GC analysis of all-trans and 13-cis-acitretin. *J. High Resolut. Chromatogr.*, 15 (1992) 637-640.
- 32h. *Toxicological and forensic applications*
- 622 Arrizabalaga, P., Kamatari, A. and Landry, J.C.: (Determination of the composition of a fraudulent drug by coupled chromatographic methods (GC/MS, GC-FTIR, HPLC-DAD). The case of the Chinese pills, Chufong Toukuwan.) *Chimia*, 46 (1992) 161-163; C.A., 117 (1992) 137745a.
- 623 Casale, J.F. and Waggoner, R.W., Jr.: A chromatographic impurity signature profile analysis for cocaine using capillary gas chromatography. *J. Forensic Sci.*, 36 (1991) 1312-1330; C.A., 117 (1992) 105827n.

- 624 Cone, E.J. and Darwin, W.D.: Rapid assay of cocaine, opiates and metabolites by gas chromatography-mass spectrometry. *J. Chromatogr.*, 580 (1992) 43-61 - a review with 117 refs.
- 625 De Kok, T.M.C.M., Levels, P.J., van Faassen, A., Hazen, M., ten Hoor, F. and Kleinjans, J.C.S.: Chromatographic methods for the determination of toxicants in faeces. *J. Chromatogr.*, 580 (1992) 135-159 - a review with 209 refs.
- 626 Demby, K.B., White, T.C. and Ghanayem, B.I.: Determination of methacrylonitrile in serum using gas chromatography and flame ionization detection. *J. Chromatogr.*, 579 (1992) 153-157.
- 627 Erdmann, F., Schuetz, H., Brose, C. and Rochholz, G.: (An optimized screening system for 170 pesticides.) *Beitr. Gerichtl. Med.*, 49 (1991) 121-126; *C.A.*, 117 (1992) 165282r.
- 628 Gan, B.K., Baugh, D., Liu, R.H. and Walia, A.S.: Simultaneous analysis of amphetamine, methamphetamine, and 3,4-methylenedioxymethamphetamine (MDMA) in urine samples by solid-phase extraction, derivatization, and gas chromatography/mass spectrometry. *J. Forensic Sci.*, 36 (1991) 1331-1341; *C.A.*, 117 (1992) 103456s.
- 629 Joern, W.A.: GC/MS assay of the marijuana carboxy metabolite: urine interference with the dimethyl derivative. *J. Anal. Toxicol.*, 16 (1992) 207; *C.A.*, 117 (1992) 021572n.
- 630 Johnson, K., Uhrich, M. and Tillmanns, U.: (Drug analysis with gas chromatography/mass spectrometry (GC/MS). Part 5. Identification and quantification of benzoylecgonine.) *GIT Fachz. Lab.*, 36, No. 1 (1992) 23-25; *C.A.*, 117 (1992) 84759z.
- 631 Kelly, P. and Jones, R.T.: Metabolism of tetrahydrocannabinol in frequent and infrequent marijuana users. *J. Anal. Toxicol.*, 16 (1992) 228-235; *C.A.*, 117 (1992) 104153c.
- 632 Kintz, P. and Mangin, P.: Analysis of opiates in human hair with FPIA, EMIT, and GC/MS. *Adli Tip Derg.*, 7 (1991) 129-132; *C.A.*, 117 (1992) 165413j.
- 633 Kintz, P., Ludes, B. and Mangin, P.: Detection of drugs in human hair using Abbott ADx, with confirmation by gas chromatography/mass spectrometry (GC/MS). *J. Forensic Sci.*, 37 (1992) 328-331; *C.A.*, 117 (1992) 126031p.
- 634 Maurer, H.H.: Systematic toxicological analysis of drugs and their metabolites by gas chromatography-mass spectrometry. *J. Chromatogr.*, 580 (1992) 3-41 - a review with 103 refs.
- 635 Miki, A., Tsuchihashi, H., Ueda, K. and Yamashita, M.: (Extractive derivatization of aqueous drugs using polymeric phase transfer catalysts. I. Sensitive analyses of dialkyl phosphates as pentafluorobenzyl derivatives.) *Jpn. J. Toxicol. Environ. Health*, 38 (1992) 168-175; *C.A.*, 117 (1992) 84766z.
- 636 Moeller, M.R.: Drug detection in hair by chromatographic procedures. *J. Chromatogr.*, 580 (1992) 125-134 - a review with 60 refs.
- 637 Moeller, M.R., Doerr, G. and Warth, S.: Simultaneous quantitation of delta-9-tetrahydrocannabinol (THC) and 11-nor-9-carboxy-delta-9-tetrahydrocannabinol (THC-COOH) in serum by GC/MS using deuterated internal standards and its application to a smoking study and forensic cases. *J. Forensic Sci.*, 37 (1992) 969-983; *C.A.*, 117 (1992) 165404g.
- 638 Nakahara, Y., Takahashi, K., Shimamine, M. and Takeda, Y.: Hair analysis for drug abuse: I. Determination of methamphetamine and amphetamine in hair by stable isotope dilution gas chromatography/mass spectrometry method. *J. Forensic Sci.*, 36 (1991) 70-78; *C.A.*, 117 (1992) 126018q.
- 639 Przekop, M.A., Manno, J.E., Kunsman, G.W., Cockerham, K.R. and Manno, B.R.: Evaluation of the Abbott ADx amphetamine/methamphetamine II abused drug assay: comparison to TDx, EMIT, and GC/MS methods. *J. Anal. Toxicol.*, 15 (1991) 323-326; *C.A.*, 116 (1992) 100708u.
- 640 Reuschel, S.A. and Smith, F.P.: Benzoylecgonine (cocaine metabolite) detection in hair samples of jail detainees using radioimmunoassay (RIA) and gas chromatography/mass spectrometry (GC/MS). *J. Forensic Sci.*, 36 (1991) 1179-1185; *C.A.*, 117 (1992) 144915j.
- 641 Streete, P.J., Ruprah, M., Ramsey, J.D. and Flanagan, R.J.: Detection and identification of volatile substances by headspace capillary gas chromatography to acid the diagnosis of acute poisoning. *Analyst (London)*, 117 (1992) 1111-1127.
- 642 Tagliaro, F., Lubli, G., Ghielmi, S., Franchi, D. and Marigo, M.: Chromatographic methods for blood alcohol determination. *J. Chromatogr.*, 580 (1992) 161-190 - a review with 28 refs.
- 643 Wu, A.H.B., Onigbinde, T.A., Wong, S.S. and Johnson, K.G.: Evaluation of full-scanning GC/ion trap MS analysis of NIDA drugs-of-abuse urine testing in urine. *J. Anal. Toxicol.*, 16 (1992) 202-206; *C.A.*, 117 (1992) 42080r.
- See also 203, 231, 269, 312, 334, 450, 452, 454, 571.
- 32i. *Plant extracts*
- 644 Jirovetz, L., Koch, H.P., Jaeger, W. and Remberg, G.: Investigations of German onion oil by GC-FID, GC-MS and GC-FTIR. *Pharmazie*, 47 (1992) 455-456; *C.A.*, 117 (1992) 157495x.
- 645 Yan, X., Wang, Z. and Barlow, P.: Quantitative estimation of garlic oil content in garlic oil-based health products. *Food Chem.*, 45 (1992) 135-139; *C.A.*, 117 (1992) 110283m.
- 646 Zhang, L., Zhang, Z., Sheng, L. and An, D.: (The determination of triptolide in *Tripterygium wilfordii* by capillary gas chromatography.) *Zhongguo Yaoke Daxue Xuebao*, 23 (1992) 158-160; *C.A.*, 117 (1992) 167708h.
- See also 622.
33. CLINICO-CHEMICAL APPLICATIONS
- 33b. *Complex mixtures and profiling (single compounds by cross-reference only)*
- 647 Holtan, D.W., Houghton, E., Silver, M., Fowden, A.L., Ousey, J. and Rosedale, P.D.: Plasma progestagens in the mare, fetus and newborn foal. *J. Reprod. Fertil., Suppl.*, 44 (1991) 517-528; *C.A.*, 117 (1992) 40619z.
- 648 Houghton, E., Holtan, D., Grainger, L., Voller, B.E., Rosedale, P.D. and Ousey, J.C.: Plasma progestagen concentrations in the normal and dysmature newborn foal. *J. Reprod. Fertil., Suppl.*, 44 (1991) 609-617; *C.A.*, 117 (1992) 40742j.
- 649 Kuksis, A., Myher, J.J., Geher, K., Breckenridge, W.C., Feather, T., McGuire, V. and Little, J.A.: Gas chromatographic profiles of plasma total lipids as indicators of dietary history. Correlation with carbohydrate and alcohol intake based on 24-h dietary recall. *J. Chromatogr.*, 579 (1992) 13-24.

- 650 Lehmann, W.D., Stephan, M. and Fuerstenberger, G.: Profiling assay for lipoxygenase products of linoleic and arachidonic acid by gas chromatography-mass spectrometry. *Anal. Biochem.*, 204 (1992) 158-170.
- 651 Liebich, H.M., Reichenmiller, A.-M., Wahl, H.-G., Tetschner, B., Risler, T., Eggstein, M. and Quellhorst, E.: GC-MS separation and quantitation of 3-carboxy-4-methyl-5-propyl-2-furanpropionic acid and hippuric acid from plasma of patients with renal failure: application to blood purification methods. *J. High Resolut. Chromatogr.*, 15 (1992) 601-604.
- 652 Lisi, A.M., Kazlauskas, R. and Trout, G.J.: Diuretic screening in human urine by gas chromatography-mass spectrometry: use of a macroreticular acrylic copolymer for the efficient removal of the coextracted phase-transfer reagent after derivatization by direct extractive alkylation. *J. Chromatogr.*, 581 (1992) 57-63.
- 653 Michitsuji, H., Ohara, A., Fukuda, M., Nakayama, K., Yamaguchi, K. and Fujiki, Y.: (Determination of acetone, methanol, and methyl ethyl ketone in urine using headspace gas chromatography (HS-GC).) *Sangyo Igaku*, 34 (1992) 243-252; *C.A.*, 117 (1992) 125979s.
- 654 Wudy, S.A., Wachter, U.A., Homoki, J., Teller, W.M. and Shackleton, C.H.L.: Androgen metabolism assessment by routine gas chromatography/mass spectrometry profiling of plasma steroids: part 1. Unconjugated steroids. *Steroids*, 57 (1992) 319-324; *C.A.*, 117 (1992) 83550n.
- 655 Yu, Z., Guo, Q. and Chang, L.: (Retention indices of 105 clinically important acidic metabolites on SE-54 and OV-210 capillary columns.) *Sepu*, 10 (1992) 278-280.
- See also 344, 348, 377, 436, 641.
34. FOOD ANALYSIS
- 34b. *Complex mixtures (single compounds by cross-reference only)*
- 656 Ammon, J., Mildau, G., Ruge, W. and Delincee, H.: (Detection of the irradiation of chicken meat by the gas chromatographic determination of fat radiolysis products.) *Dtsch. Lebensm.-Rundsch.*, 88, No. 2 (1992) 35-37; *C.A.*, 117 (1992) 169665r.
- 657 Artz, W.E.: Analysis of lipids by supercritical fluid chromatography. In: Perkins, E.G. (Editor), *Analysis of Fats, Oils Lipoproteins*, AOCS, Champaign, Ill., pp. 35-37; *C.A.*, 117 (1992) 169631b - a review with with 101 refs.
- 658 Caboni, M.F., Conte, L.S. and Lercker, G.: (Accidental contamination of food: gas chromatographic determination of mono- and diethylene glycols.) *Boll. Chim. Ig. Parte Sci.*, 42 (1991) 675-680; *C.A.*, 117 (1992) 68605r.
- 659 Cacho, J., Melendez, J. and Ferreira, V.: Development of a method for analyzing volatiles from foodstuff matrices, including microextraction by demixture. Application to the analysis of grapes. *Mikrochim. Acta*, 108 (1992) 61-72.
- 660 Castellon Arnau, A.: (Derivation and assessment of a general equation for evaluating the quantity of the material retained on the column during gas chromatography. Analysis of abused edible oils.) *Grasas Aceites*, 42 (1991) 404-413.
- 661 Chun, H.N. and Kim, Z.U.: (Evaluation of vegetable oil rancidity by headspace gas chromatographic analysis.) *Han'guk Non-gwhwa Hakhoechi*, 35 (1992) 36-41; *C.A.*, 117 (1992) 46992n.
- 662 Daft, J.L.: Headspace method for rapid determination of methyl bromide in assorted nut samples. *J. Assoc. Off. Anal. Chem.*, 75 (1992) 701-706.
- 663 Elmore, J.S. and Nursten, H.E.: A comparison of the headspace extracts of instant and ground coffee. In: Bessiere, Y. and Thomas, A.F. (Editors), *Flavour Science and Technology*, Wiley, Chichester, 1990, pp. 203-204; *C.A.*, 116 (1992) 005534a.
- 664 Furuta, M., Dohmaru, T., Katayama, T., Toratani, H. and Takeda, A.: Detection of irradiated frozen meat and poultry using carbon monoxide gas as a probe. *J. Agric. Food Chem.*, 40 (1992) 1099-1100.
- 665 Gao, H., Sun W. and Sha, Z.: (Determination of solvent remained in oil of seabuckthorn (*Hippophae rhamnoides*) by static head-space gas chromatography.) *Zhongcaoyao*, 23 (1992) 243,248; *C.A.*, 117 (1992) 137748d.
- 666 Gardner, D.R., Sanders, R.A., Henry, D.E., Tallmadge, D.H. and Wharton, H.W.: Characterization of used frying oils. Part 1: isolation and identification of compound classes. *J. Am. Oil Chem. Soc.*, 69 (1992) 499-508.
- 667 Guerzoni, M.E., Gardini, F., Sinigaglia, M. and Guglielmi, G.: HS-GC determination of lactic acid bacteria viable cells based on malolactic activity. *Cerevisia Biotechnol.*, 17 (1992) 19-23; *C.A.*, 117 (1992) 68575f.
- 668 Jirovetz, L., Jaeger, W., Remberg, G., Espinosa-Gonzalez, J., Morales, R., Woidich, A. and Nikiforov, A.: Analysis of the volatiles in the seed oil of *Hibiscus sabdariffa* (Malvaceae) by means of GC-MS and GC-FTIR. *J. Agric. Food Chem.*, 40 (1992) 1186-1187.
- 669 Kihara, K.: (An evaluation method for shoyu quality.) *Nippon Shoyu Kenkyusho Zasshi*, 18, No. 2 (1992) 66-69; *C.A.*, 117 (1992) 024851p.
- 670 Mattheis, J.P., Buchanan, D.A. and Fellman, J.K.: Identification of headspace volatile compounds from 'Bing' sweet cherry fruit. *Phytochemistry*, 31 (1992) 775-777; *C.A.*, 116 (1992) 192887c.
- 671 Park, P.S.W. and Goins, R.E.: Determination of volatile lipid oxidation products by dynamic headspace-capillary gas chromatographic analysis with application to milk-based nutritional products. *J. Agric. Food Chem.*, 40 (1992) 1581-1585.
- 672 Pino, J.: Headspace methods for volatile components of cocoa butter. *Nahrung*, 36 (1992) 175-180; *C.A.*, 117 (1992) 68586k.
- 673 Samukawa, K., Inoue, A. and Nagano, T.: (Characterization of rices by Curie-point pyrolysis-gas chromatography-mass spectrometry.) *Chem. Express*, 7 (1992) 449-452; *C.A.*, 117 (1992) 68602n.
- 674 Sjoberg, A.M., Tuominen, J.P., Kiutamo, T. and Luukkonen, S.M.: Evaluation of a gas chromatographic method for detection of irradiation of chicken and a chicken meat product. *J. Sci. Food Agric.*, 59 (1992) 65-75; *C.A.*, 117 (1992) 110291n.
- 675 Steinwandter, H.: Research in environmental pollution. VI. Identification of non-o,o'-Cl and mono-o,o'-Cl substituted PCB congeners in Hessian milk. *Fresenius J. Anal. Chem.*, 343 (1992) 378-381.
- 676 Yao, J., Wang, Z., Jiao, S., Zheng, Y., Zhao, J., Gao, X. and Liang, T.T.: A complete set of procedure for determination of multiresidue of pesticides in vegetables. *J. Environ. Sci. (China)*, 4 (1992) 106-116; *C.A.*, 117 (1992) 130053q.
- 677 Zimmer, M., Otteneder, H. and Bierl, I.: (Methyl isocyanate determination in wine.) *Weinwirtsch.*, No. 3 (1992) 10-11; *C.A.*, 117 (1992) 169670p.

- See also 241, 261, 281, 322, 332, 335, 345, 347, 351, 352, 355, 363, 366, 371, 374, 388, 390, 435, 462, 464, 465, 467, 487, 508, 511, 519, 520, 521, 522, 523, 524, 525, 532, 809, 812.
- 34c. *Organoleptically important compounds (flavors, odors, volatiles)*
- 678 Barker, R.L., Irwin, A.J. and Murray, C.R.: The relationship between fermentation variables and flavor volatiles by direct gas chromatographic injection of beer. *Tech. Q. - Master Brew. Assoc. Am.*, 29 (1992) 11-17; *C.A.*, 117 (1992) 88848a.
- 679 Braunsdorf, R., Hener, U. and Mosandl, A.: (Analytical differentiation between natural, fermentatively produced, and synthetic (nature-identical) aroma compounds. Part 2. GC-C-IRMS analysis of important flavor aldehydes - principles and applications.) *Z. Lebensm.-Unters. Forsch.*, 194 (1992) 426-430; *C.A.*, 117 (1992) 149586n.
- 680 Compton, S.V. and Stout, P.: Application of headspace GC-FTIR. Analysis of flavor oils. *Am. Lab. (Fairfield)*, 24, No. 9 (1992) 38-44 - a review with 10 refs.
- 681 Dietrich, A., Maas, B., Messer, W., Bruche, G., Karl, V., Kauszinger, A. and Mosandl, A.: Stereoisomeric flavor compounds, part LVIII: The use of heptakis(2,3-di-O-methyl-6-O-tert-butylidimethylsilyl)- β -cyclodextrin as a chiral stationary phase in flavor analysis. *J. High Resolut. Chromatogr.*, 15 (1992) 590-593.
- 682 Fischer, J. and Mueller, K.: (Studies of the volatile aromatics in potato. I. Isolation and enrichment of volatile aromatics using the dynamic headspace technique.) *Potato Res.*, 34 (1991) 159-167; *C.A.*, 116 (1992) 057794e.
- 683 Golovnya, R.V., Grigor'eva, D.N., Semina, L.A. and Misharina, T.A.: (Gas chromatographic and organoleptic estimation of the quality of meat flavours.) *Zh. Anal. Khim.*, 47 (1992) 844-849.
- 684 Lee, S.R., Macku, C. and Shibamoto, T.: Isolation and identification of headspace volatiles formed in heated butter. *J. Agric. Food Chem.*, 39 (1991) 1972-1975.
- 685 Leino, M.E.: Effect of freezing, freeze-drying, and air-drying on odor of chive characterized by headspace gas chromatography and sensory analyses. *J. Agric. Food Chem.*, 40 (1992) 1379-1384.
- 686 Misharina, T.A., Artamonova, M.P., Zhuravskaya, N.K. and Golovnya, R.V.: (Identification of volatile components of a model system with meat aroma.) *Zh. Anal. Khim.*, 47 (1992) 850-857.
- 687 Nagai, Y.: (Improved analysis of flavor compounds in alcoholic beverages by headspace gas chromatography.) *Nippon Shokuhin Kogyo Gakkaishi*, 39 (1992) 264-270; *C.A.*, 117 (1992) 169695a.
- 688 Narasimhan, S., Rajalakshmi, D. and Chand, N.: Quality of powdered black pepper (*Piper nigrum* L.) during storage. II. Principal components analyses of GC and sensory profiles. *J. Food Qual.*, 15 (1992) 67-83; *C.A.*, 117 (1992) 88995w.
- 689 Nitz, S., Spraul, M.H. and Drawert, F.: Gas chromatographic and sensory analysis of dill flowers (*Anethum graveolens*). *Chem., Mikrobiol., Technol. Lebensm.*, 13 (1992) 183-186; *C.A.*, 117 (1992) 24937w.
- 690 Náf, R.: La preparation d'extraits, une étape cruciale dans l'analyse des arômes. *Analisis*, 20 (1992) M23-M24.
- 691 Sheen, L.Y., Lin, S.Y. and Tsai, S.J.: (Odor assessments for volatile compounds of garlic and ginger essential oils by sniffing gas chromatography.) *Zhongguo Nongye Huaxue Huizhi*, 30, No. 1 (1992) 14-24; *C.A.*, 117 (1992) 130135t.
- 692 Um, K.W., Milton, E., Clarke, A.D. and Chao, R.R.: Concentration and identification of volatile compounds from heated beef fat using supercritical carbon dioxide extraction-gas liquid chromatography/mass spectrometry. *J. Agric. Food Chem.*, 40 (1992) 1641-1646.
- 693 Umamo, K.: (Computerized identification of volatile constituents using bench-top gas chromatograph/mass spectrometer.) *Fragrance J.*, 20 (1992) 27-32; *C.A.*, 117 (1992) 69205x.
- 694 Vernin, G., Metzger, J., Boniface, C., Murello, M.H., Siouffi, A., Larice, J.L. and Parkanyi, C.: Kinetics and thermal degradation of the fructose-methionine Amadori intermediates. GC-MS/SPECMA data bank identification of volatile aroma compounds. *Carbohydr. Res.*, 230 (1992) 15-29; *C.A.*, 117 (1992) 171975k.
- See also 83, 132, 189, 194, 413, 419, 471.
35. ENVIRONMENTAL ANALYSIS
- 35a. *General papers and reviews*
- 695 Brauch, H.J.: (Gas chromatography/mass spectrometry - a reliable and sensitive method for water analysis.) *Tech. Mess.*, 59 (1992) 154-159; *C.A.*, 117 (1992) 55217d.
- 696 Kenmotsu, K., Takano, H., Kobara, K. and Ogino, Y.: (Practice of determination of trace quantities of chemicals in environment by GC/MS.) *Shigen Kankyo Taisaku*, 28 (1992) 124-135; *C.A.*, 116 (1992) 206721h.
- 697 Kessels, H., Hoogerwerf, W. and Lips, J.: The determination of volatile organic compounds from EPA method 524.2 using purge and trap capillary gas chromatography, ECD and FID detection. *Analisis*, 20, No. 8 (1992) M55-M60.
- 698 Wells, D.E., Maier, E.A. and Griepink, B.: Development in the analysis of chlorobiphenyls in environmental matrices for certification purposes. *Intern. J. Environ. Anal. Chem.*, 46 (1992) 265-275.
- See also 477.
- 35b. *Air pollution (complex mixtures; single compounds by cross-reference only)*
- 699 Aceves, M. and Grimalt, J.O.: Gas chromatographic screening of organic compounds in urban aerosols. Selectivity effects in semi-polar columns. *J. Chromatogr.*, 607 (1992) 261-270.
- 700 Fujimori, K., Oki, N., Nakano, T., Tsuji, M. and Okuno, T.: (Analysis of pesticides in air by collection method using ODS cartridge.) *Kankyo Kagaku*, 1 (1991) 575-581; *C.A.*, 116 (1992) 180016e.
- 701 Goelen, E., Lambrechts, M., Geyskens, F. and Rymen, T.: Development and performance characteristics of a capillary dosage unit with in situ weight sensor for the preparation of known amounts of gaseous VOC's in air. *Intern. J. Environ. Anal. Chem.*, 47 (1992) 217-225.
- 702 Heavner, D.L., Ogden, M.W. and Nelson, P.R.: Multisorbent thermal desorption/gas chromatography/mass selective detection method for the determination of target volatile organic compounds in indoor air. *Environ. Sci. Technol.*, 26 (1992) 1737-1746; *C.A.*, 117 (1992) 96388r.

- 703 Helmig, D., Bauer, A., Mueller, J. and Klein, W.: Analysis of particulate organics in a forest atmosphere by thermodesorption GC/MS. Reply to comments. *Atmos. Environ.*, 26A (1992) 518-519; C.A., 116 (1992) 065954x.
- 704 Mueller, H. and Gellert, U.: (Air analysis in trace ranges with enrichment technique by the routine gas chromatograph RGC 202.) *Umwelt-Technol. Aktuell*, 3 (1992) 102-103; C.A., 117 (1992) 156557g.
- 705 Ogden, M.W.: Equivalency of gas chromatographic conditions in determination of nicotine in environmental tobacco smoke: minicolaborative study. *J. Assoc. Off. Anal. Chem.*, 75 (1992) 729-733.
- 706 Pfeil, R.M. and Mumma, R.O.: Air sampling of volatiles from *Agaricus bisporus* in a mushroom facility and from mushroom compost. *HortScience*, 27 (1992) 416-419; C.A., 117 (1992) 21914a.
- 707 Rhead, M.M. and Trier, C.J.: Fuel residues and organic combustion products in diesel exhaust emissions: sources, sampling and analysis. *TrAC*, 11 (1992) 255-259.
- 708 Simoneit, B.R.T.: Analysis of particulate organics in a forest atmosphere by thermodesorption GC/MS. Comments. *Atmos. Environ.*, 26A (1992) 517-518; C.A., 116 (1992) 090211k.
- 709 Yamashita, S., Ozawa, R., Yamaguchi, K., Hanai, Y., Katou, T. and Wylie, P.L.: Analysis of volatile organic compounds in air by gas chromatography with thermal desorption cold-trap injection and atomic emission and mass selective detection. *J. High Resolut. Chromatogr.*, 15 (1992) 549-551.
- See also 228, 232, 234, 237, 250, 271, 315, 354, 361, 365, 394, 398, 422, 426, 429, 516, 766, 804.
- 35c. *Water pollution (complex mixtures; single compounds by cross-reference only)*
- 710 Arthur, C.L., Potter, D.W., Buchholz, K.D., Motlagh, S. and Pawliszyn, J.: Solid-phase microextraction for the direct analysis of water: theory and practice. *LC-GC Int.*, 10 (1992) 656-661.
- 711 Brauch, H.-J.: The analysis of polar organic compounds in water. *Fresenius J. Anal. Chem.*, 343 (1992) 6-7.
- 712 Chen, Z.: (Determination of organic matters in dyeing wastewater by GC/MS.) *Shanghai Huanjing Kexue*, 10, No. 5 (1991) 27-31; C.A., 116 (1992) 113205c.
- 713 Durand, G., Bouvot, V. and Barceló, D.: Determination of trace levels of herbicides in estuarine waters by gas and liquid chromatographic techniques. *J. Chromatogr.*, 607 (1992) 319-327.
- 714 Ezzell, J.L. and Bruce, E.R.: Supercritical fluid extraction of pesticides and phthalate esters following solid phase extraction from water. *J. Microcolumn Separ.*, 4 (1992) 319-323.
- 715 Gurka, D.F., Pyle, S.M. and Titus, R.: Environmental analysis by direct aqueous injection. *Anal. Chem.*, 64 (1992) 1749-1754.
- 716 Hrivnak, J. and Ilavsky, J.: (Microextraction and capillary gas chromatography in water analysis for organic contaminants.) *Vodohospod. Cas.*, 39 (1991) 319-323; C.A., 117 (1992) 75811h - a review with 18 refs.
- 717 Hühnerfuss, H. and Kallenborn, R.: Chromatographic separation of marine organic pollutants. *J. Chromatogr.*, 580 (1992) 191-214 - a review with 138 refs.
- 718 Koehle, H. and Haberer, K.: (Dual channel evaluation system for the quality assurance in high-resolution gas chromatography.) *Vom Wasser*, 78 (1992) 305-312; C.A., 117 (1992) 157257w.
- 719 Lautamo, R.M.A., Decker, D., Jennings, W. and Mehran, M.F.: The "priority pollutant" analysis of water. *Chromatographia*, 34 (1992) 331-334.
- 720 Legrand, M.F., Costentin, E. and Bruchet, A.: Occurrence of 38 pesticides in various French surface and ground waters. *Environ. Technol.*, 12 (1991) 985-996; C.A., 116 (1992) 091005h.
- 721 Lukaszewicz, M. and Durhan, E.: Strategies for the identification of non-polar toxicants in aqueous environmental samples using toxicity-based fractionation and gas chromatography-mass spectrometry. *J. Chromatogr.*, 580 (1992) 215-228; - a review with 37 refs.
- 722 Matz, G. and Kesners, P.: Membrane probe for on-line measurement of organic compounds in water using a mobile GC/MS system. *Analisis*, 20, No. 3 (1992) M53-M54.
- 723 Melero, M.P.O., Alonso, J.S., Infante, A.P., Navascués, M.C.M., Guajardo, N.C., Cortabitarte, M.S.M. and Narvi6n, J.L.O.: Identification by gas chromatography-electron-capture detection and gas chromatography-mass spectrometry of the ozonation products in wastewater from dicofol and tetradifon manufacturing. *J. Chromatogr.*, 607 (1992) 311-318.
- 724 Molto, J.C., Pico, Y., Manes, J. and Font, G.: Analysis of polychlorinated biphenyls in aqueous samples using C₁₈ glass column extraction. *J. Assoc. Off. Anal. Chem.*, 75 (1992) 714-719.
- 725 Philippaerts, J., Vanhoof, C. and Vansant, E.F.: Wastewater analysis by purge and trap capillary GC-FTIR spectrometry. *Talanta*, 39 (1992) 681-685.
- 726 Pratt, K.F. and Pawliszyn, J.: Water monitoring system based on gas extraction with a single hollow fiber membrane and gas chromatographic cryotrapping. *Anal. Chem.*, 64 (1992) 2107-2110.
- 727 Takahashi, Y., Nakagawa, Z., Yosida, N., Sasano, H. and Morita, M.: (Research of monitoring organic compounds in river.) *Kankyo Kagaku*, 2 (1992) 438-439; C.A., 117 (1992) 117952n.
- 728 Tillmanns, U.: (Noxious matter trace analysis by combined gas chromatography and mass spectrometry.) *Umwelt-Technol. Aktuell*, 3 (1992) 97-101; C.A., 117 (1992) 157114x.
- 729 Woolfenden, E. and Barberio, J.: Purge-and-trap analysis of volatile organic compounds (VOCs) in water. *Inter. Lab.*, 22, No.9 (1992) 25-30.
- 730 Yamaguchi, Y., Fukushima, M., Fujita, T. and Yamada, A.: (Distribution and seasonal variation of pesticides in the Yodo River Basin.) *Kankyo Kagaku*, 1 (1991) 56-63; C.A., 116 (1992) 066731j.
- See also 47, 167, 172, 176, 239, 257, 272, 273, 274, 275, 279, 280, 294, 305, 336, 367, 412, 427, 466, 468, 472, 481, 485, 491, 500, 501, 503, 505, 507, 512, 515, 530, 695, 790, 810.
- 35d. *Soil pollution (complex mixtures; single compounds by cross-reference only)*
- 731 Emery, A.P., Chester, S.N. and MacCrehan, W.A.: Recovery of diesel fuel from clays by supercritical fluid extraction-gas chromatography. *J. Chromatogr.*, 606 (1992) 221-228.
- 732 Gerbino, T.C., Nadotti, S. and Castello, P.: Extraction and gas chromatographic determination of chlorinated solvents in contaminated soil. *J. Chromatogr.*, 623 (1992) 123-127.

- 733 Hewitt, A.D., Miyares, P.H., Leggett, D.C. and Jenkins, T.F.: Comparison of analytical methods for determination of volatile organic compounds in soils. *Environ. Sci. Technol.*, 26 (1992) 1932-1938; C.A., 117 (1992) 150008g.
- 734 Hou, J., Feng, J. and Ye, L.: (Tracing the contamination of soil by nonpolar compounds with adsorption wires.) *Fenxi Huaxue*, 20 (1992) 701-703; C.A., 117 (1992) 130411e.
- 735 Parr, J.L., Walters, G. and Hoffman, M.: Sampling and analysis of soils for gasoline range organics. *Hydrocarbon Contam. Soils Groundwater*, 1 (1992) 105-132; C.A., 117 (1992) 89423v.
- 736 Pavlostathis, S.G. and Mathavan, G.N.: Application of headspace analysis for the determination of volatile organic compounds in contaminated soils. *Eviron. Technol.*, 13 (1992) 23-33; C.A., 116 (1992) 268359e.
- 737 Samukawa, K.: Chemical characterization of soil organic matter by Curie-point pyrolysis-gas chromatography. *Chem. Express*, 7 (1992) 445-448; C.A., 117 (1992) 69038v.
- 738 Samukawa, K., Matsumoto, S. and Komai, Y.: (Characterization of effect of earthworm habitation on soil organic matter by Curie-point pyrolysis-gas chromatography-mass spectrometry.) *Nippon Dojo Hiriyogaku Zasshi*, 63 (1992) 139-145; C.A., 117 (1992) 47352r.
- See also 233, 244, 255, 373, 515, 531, 532.
36. SOME TECHNICAL PRODUCTS AND COMPLEX MIXTURES
- 36a. *Surfactants*
- See 32.
- 36b. *Antioxidants and preservatives*
- 739 De Koning, A.J. and van der Merve, G.: Determination of ethoxyquin and two of its oxidation products in fish meal by gas chromatography. *Analyst (London)*, 117 (1992) 1571-1576.
- 740 Goekeler, U.K.: Rapid and simple determination of oxygenated additives in gasoline. *Am. Lab. (Fairfield)*, 24, No. 11 (1992) 28D, 28F, 28H.
- See also 198, 714.
- 36c. *Complex mixtures, technical products and unidentified compounds*
- 741 Andreeva, G.C., Grigor'eva, I.A. and Trusov, S.R.: (Gas-liquid chromatographic analysis of 2,4-dichlorotoluene oxidation products.) *Khim.-Farm. Zh.*, 26, No. 2 (1992) 84-86; C.A., 117 (1992) 103305s.
- 742 Arakelyan, V.G., Dar'yan, L.A. and Lokhanin, A.K.: (Chromatographic method for diagnostic studies of high-voltage pulse capacitors during their fabrication.) *Elektrichestvo*, No. 1 (1992) 54-57; C.A., 117 (1992) 122611e.
- 743 Avgul, T.V., Churilin, V.S. and Rivina, Z.M.: (Gas chromatographic determination of organic impurities of the synthesis products of methane and ethane chloroderivatives on modified Silochrom S-80.) *Zh. Anal. Khim.*, 47 (1992) 398-402.
- 744 Bridle, K.A.: A new method for the analysis of plant tissues by GC-MS, with emphasis on floral induction in *Pharbitis nil* Choisy. Avail. *Univ. Microfilm Int.*, Order No. DA9135165, 1991, 193 pp.; C.A., 117 (1992) 146647x.
- 745 Brossard, S., Lafosse, M. and Dreux, M.: Analysis of synthetic mixtures of waxes by supercritical fluid chromatography with packed columns using evaporative light-scattering detection. *J. Chromatogr.*, 623 (1992) 323-328.
- 746 Chang, H.C.K., Bartle, K.D., Markides, K.E. and Lee, M.L.: Structural comparison of low-molecular-weight extractable compounds in different rank coals using capillary column gas chromatography. In: Meuzelaar, K.E. (Editor), *Adv. Coal Spectrosc.*, Plenum, New York, 1992, pp. 141-164; C.A., 117 (1992) 52059e.
- 747 Chu, J., Zhang, Z., Pu, F., Shi, Y., Perineau, F., Gaset, A. and Delmas, M.: Analysis of headspace of root of *Levisticum officinale* *Zhiwu Xuebao*, 33 (1991) 486-488; C.A., 116 (1992) 102647x.
- 748 Damsté, J.S.S., de las Heras, F.X.C. and de Leeuw, J.W.: Molecular analysis of sulphur-rich brown coals by flash pyrolysis-gas chromatography-mass spectrometry. The type III-S kerogen. *J. Chromatogr.*, 607 (1992) 361-376.
- 749 Dmitriev, V., Zubov, M.S., Baulov, V.I. and Toguzov, B.M.: (Improvement of gas-chromatographic determination of the composition of gas evolved from a battery.) *Zh. Prikl. Khim. (S.-Peterburg)*, 65 (1992) 210-212.
- 750 Dzidic, I., Petersen, H.A., Wadsworth, P.A. and Hart, H.V.: Townsend discharge nitric oxide chemical ionization gas chromatography/mass spectrometry for hydrocarbon analysis of the middle distillates. *Anal. Chem.*, 64 (1992) 2227-2232.
- 751 Falandysz, J., Yamashita, N., Tanabe, S. and Tatsukawa, R.: Composition of PCB isomers and congeners in technical chlorofen formulation produced in Poland. *Intern. J. Environ. Anal. Chem.*, 47 (1992) 129-136.
- 752 Fiebig, W., Mueller, H. and Preiss, H.: (Gas-chromatographic method for determination of lower hydrocarbons and permanent gases involving activated carbon.) *Ger. (East) DD 296,363 (Cl. G01N30/48)*, 28 Nov. 1991, Appl. 342,407, 02 Jul. 1990; 5 pp.; C.A., 117 (1992) 103331x.
- 753 Filippini, C., Sonnleitner, B. and Fiechter, A.: On-line monitoring of fast bioprocesses with flow injection analysis and gas chromatography. *GBF Monogr.*, 17 (1992) 531-534; C.A., 117 (1992) 68404z.
- 754 Garcia, C., Hernandez, T., Costa, F., Ceccanti, B. and Calcinaï, M.: A chemical-structure study of organic wastes and their humic acids during composting by means of pyrolysis-gas chromatography. *Sci. Total Environ.*, 119 (1992) 157-168; C.A., 117 (1992) 55179e.
- 755 González-Vila, F.J., Verdejo, T. and Martin, F.: Characterization of wastes from olive and sugarbeet processing industries and effects of their application upon the organic fraction of agricultural soils. *Intern. J. Environ. Anal. Chem.*, 46 (1992) 213-222.
- 756 Ivanov, V. and Sandell, E.: Characterization of polychlorinated biphenyl isomers in Sovol and trichlorodiphenyl formulations by high-resolution gas chromatography with electron capture detection and high-resolution gas chromatography-mass spectrometry techniques. *Environ. Sci. Technol.*, 26 (1992) 2012-2017; C.A., 117 (1992) 134109k.

- 757 Jordan, E.D., Hsieh, T.C.Y. and Fischer, N.H.: Volatile compounds from leaves of *Ceratiola ericoides* by dynamic headspace sampling. *Phytochemistry*, 31 (1992) 1203-1208; C.A., 117 (1992) 023299c.
- 758 Kawai, T., Kato, E. and Yamamoto, T.: (Gas chromatography method for determination of hexafluoroacetone in 2,2-bisarylhexafluoropropane.) *Jpn. Kokai Tokkyo Koho* JP 04 62,469 [92 62,469] (Cl. G01N30/06), 27 Feb. 1992, Appl. 90/172,229, 29 Jun. 1990; 4 pp.; C.A., 117 (1992) 62002v.
- 759 Kelly, G.W., Bartle, K.D., Clifford, A.A. and Robinson, R.E.: Application of coupled LC-GC to the analysis of the polar fraction of diesel particulate matter. *J. High Resolut. Chromatogr.*, 15 (1992) 526-530.
- 760 Kelly, J. and Helleur, R.: Quantitative analysis of the major saccharides in sulfite-treated wood pulps by pyrolysis-gas chromatography: the effect of metal ions. *J. Anal. Appl. Pyrolysis*, 23 (1992) 153-163; C.A., 117 (1992) 133165p.
- 761 Kokoh, K.B., Leger, J.M., Beden, B. and Lamy, C.: 'On line chromatographic analysis of the products resulting from the electrocatalytic oxidation of D-glucose on platinum, gold and adatoms modified platinum electrodes. Part I. Acid and neutral media. *Electrochim. Acta*, 37 (1992) 1333-1342; C.A., 117 (1992) 78622h.
- 762 Kolla, P.: Trace analysis of explosives from complex mixtures with sample pretreatment and selective detection. *J. Forensic Sci.*, 36 (1991) 1342-1359; C.A., 117 (1992) 105828p.
- 763 Kumano, Y.: (Gas chromatograph method for determination of paint odor components.) *Jpn. Kokai Tokkyo Koho* JP 04,140,659 [92,140,659] (Cl. G01N30/16), 14 May 1992, Appl. 90/260,388, 01 Oct. 1990; 4 pp.; C.A., 117 (1992) 103342b.
- 764 Kuntun, A., Kifili, H. and Hyong, C.K.: Gas chromatographic analysis of fragrances in palm-based white soaps. *J. Am. Oil Chem. Soc.*, 69 (1992) 614-620.
- 765 Lapolla, A., Gerhardinger, C., Baldo, L., Fedele, D., Favretto, D., Seraglia, R. and Traldi, P.: Pyrolysis/gas chromatography/mass spectrometry in the analysis of glycated poly-L-lysine. *Org. Mass Spektrom.*, 27 (1992) 183-187; C.A., 117 (1992) 22805c.
- 766 Larikov, V.V., Shurov, Yu.A., Golgel', O.V. and Novoselova, E.A.: (Gas chromatographic determination of some toxic substances in air and technological gases of cable production.) *Zh. Anal. Khim.*, 47 (1992) 1492-1496.
- 767 Leal, W.S., Hasegawa, M. and Sawada, M.: Identification of *Anomala schonfeldi* sex pheromone by high-resolution GC-behaviour bioassay. *Naturwissenschaften*, 79 (1992) 518-519.
- 768 Leonard, R.G., Brennan, M., Quigley, K. and Heatley, D.: Analysis of anaerobic sealants and adhesives. *Anal. Proc.*, 29 (1992) 393-398.
- 769 Lin, J., Hu, B. and Chen, W.: (Disillation and GC-MS-DS determination of cashew nutshell liquid.) *Linchen Huaxue Yu Gongye*, 11 (1991) 33-39; C.A., 116 (1992) 154153c.
- 770 Linkiewicz, M. and Morawski, I.: (Gas chromatographic determination of compounds in the synthesis of 2-ethylhexanoic acid.) *Chem. Anal. (Warsaw)*, 36 (1991) 625-628.
- 771 Malievskaya, L.G., Tsyru'nik, N.I., Kravchenko, O.B. and Pivovarova, S.N.: (Gas-chromatographic method for analysis of organic solutions containing water and acidic components.) *U.S.S.R. SU* 1,672,351 (Cl. G01N30/06), 23 Aug. 1991, Appl. 4,364,636, 13 Jan. 1988; C.A., 117 (1992) 61994h.
- 772 Morgan, S.L., Fox, A., Rogers, J.C. and Watt, B.E.: Identification of chemical markers for microbial differentiation and detection by gas chromatography-mass spectrometry. In: Nelson, W.H. (Editor), *Modern Technology for Rapid Microbiological Analysis*, VCH, New York, 1991, pp. 1-18; C.A., 117 (1992) 166958c.
- 773 Nagai, Y.: (Improved sensitivity of headspace gas chromatography and its application to residual solvents analysis of medicines.) *Bunseki Kagaku*, 41 (1992) 491-495.
- 774 Nikiforov, A., Buchbauer, G., Jirovets, L., Remberg, B. and Remberg, G.: Headspace constituents of Vetiver oil. *Z. Naturforsch., B: Chem. Sci.*, 47 (1992) 439-440; C.A., 116 (1992) 221309y.
- 775 Nowicki, J.: Analysis of fire debris samples by gas chromatography/mass spectrometry (GC-MS): case studies. *J. Forensic Sci.*, 36 (1991) 1536-1550; C.A., 117 (1992) 105831j.
- 776 Pleshkova, A.P., Uspenskaya, M.N. and Volkovich, S.V.: (Identification of organic substances as impurities in bisphenol A by combined chromatography - mass spectrometry.) *Zh. Anal. Khim.*, 47 (1992) 1066-1069.
- 777 Saiz-Jimenez, C.: Applications of pyrolysis-gas chromatography/mass spectrometry to the study of humic substances: evidence of aliphatic biopolymers in sedimentary and terrestrial humic acids. *Sci. Total Environ.*, 117-118 (1992) 13-25; C.A., 117 (1992) 72930k.
- 778 Samukawa, K.: (Effect of pyrofil temperature on soil humic acid analysis by Curie-point pyrolysis-gas chromatography/mass spectrometry.) *Chem. Express*, 7 (1992) 545-548; C.A., 117 (1992) 110704z.
- 779 Samukawa, K. and Itano, M.: (Chemical characterization of cyanobacteria by Curie-point pyrolysis-gas chromatography-mass spectrometry.) *Nippon Dojo Hiriyogaku Zasshi*, 63 (1992) 281-288; C.A., 117 (1992) 107849v.
- 780 Simonsick, W.J., Jr. and Litty, L.L.: Analytical applications of supercritical fluid extraction-chromatography in the coatings industry. In: Bright, F.V. and McNally, M.E.P. (Editors), *ACS Symp. Series 448*, ACS, Washington, pp. 288-303; C.A., 117 (1992) 152793h.
- 781 Smith, I.D. and Waters, D.G.: Rapid determination of impurities in 11-bromoundecyl methacrylate by capillary GC. *J. High Resolut. Chromatogr.*, 15 (1992) 696-697.
- 782 Sommermeyer, K., Cech, F., Hidebrand, U., Pfitzer, E. and Baumbach, C.: (Chromatographic studies on the polydispersity of hydroxyethyl starch.) *Starch/Staerke*, 44 (1992) 215-218; C.A., 117 (1992) 51188j.
- 783 Standen, G., Boucher, R.J., Eglinton, G., Hansen, G., Eglinton, T.I. and Larter, S.R.: Differentiation of German Tertiary brown coal lithotypes ('amorphous' and 'woody' keragens) using ruthenium tetroxide oxidation and pyrolysis-G.C.-M.S. *Fuel*, 71 (1992) 31-36; C.A., 116 (1992) 063226n.
- 784 Sugimoto, Y., Tunnukij, K., Miki, Y. and Yamadayo, S.: Characterization of compound-class fractions of vacuum gas oil by using capillary GC and GCMS. *Sekiyu Gakkaischi*, 35 (1992) 339-348; C.A., 117 (1992) 93382y.
- 785 Terekhova, G.P., Panina, L.I. and Sakodinskii, K.I.: (Gas chromatographic method for analysis of water-formaldehyde-methanol mixtures.) *U.S.S.R. SU* 1,693,535 (Cl. G01N30/48), 23 Nov. 1991, Appl. 4,703,525, 03 Apr. 1989; C.A., 117 (1992) 103327a.

- 786 Tranthim-Fryer, D.J.: The application of a simple and inexpensive modified carbon wire adsorption/solvent extraction technique to the analysis of accelerants and volatile organic compounds in arson debris. *J. Forensic Sci.*, 35 (1990) 271-280; C.A., 117 (1992) 126033r.
- 787 Ueda, H., Yamamoto, S., Yamada, K. and Wada, Y.: (Measurement of LP gas odorants by gas chromatography.) *Idemitsu Giho*, 34 (1991) 573-577; C.A., 116 (1992) 063115a.
- 788 Van Dam, J. and Daenens, P.: Evaluation of the thermal decomposition apparatus used for toxicity testing of hydraulic fluids. *J. Anal. Appl. Pyrolysis*, 23 (1992) 43-56; C.A., 117 (1992) 039521e.
- 789 Wang, S., Zhang, Z., Cai, L. and Ren, Y.: (Analysis of off-gas scrubbing effluent from manufacturing of phthalic anhydride by oxidation of *o*-xylene.) *Huaxue Yu Nianhe*, 4 (1991) 221-223; C.A., 117 (1992) 014089g.
- 790 Wang, Y. and Crosby, D.G.: (Determination of Quinclorac residues in water using bonded-phase silica extraction and chromatography.) *Huanjing Huaxue*, 11 (1992) 29-34; C.A., 116 (1992) 241591n.
- 791 Wienecke, J., Kruse, H. and Wassermann, O.: Organic compounds in the waste gasification and combustion process. *Chemosphere*, 25 (1992) 437-447.
- 792 Xue, L., Wang, G. and Wang, G.: (Chromatographic analysis of formaldehyde-water-benzene-metaformaldehyde liquid-liquid equilibrium system.) *Fenxi Huaxue*, 20 (1992) 740; C.A., 117 (1992) 123769z.
- 793 Zhang, K., Yu, Z., Wu, W. and Li, G.: (Mass spectrometric study of cholesteric liquid crystals. II. Identification of liquid crystal mixtures by GC-MS.) *Huadong Huagong Xueyuan Xuebao*, 17 (1991) 790-794; C.A., 117 (1992) 123761r.
- 794 Zhang, Y., Qu, L. and Wang, S.: (Determination of residual solvents in spice oleoresins by headspace gas chromatography.) *Sepu*, 10 (1992) 222-224.
- 795 Zunic, W.M., Ding, S., Durig, J.R. and Cohen, A.D.: Pyrolysis GC/FID/FT-IR, pyrolysis GC/MS, and FT-IR studies of pore water extracts derived from peat samples differing in depositional environments. *J. Mol. Struct.*, 267 (1992) 371-376; C.A., 116 (1992) 238632w.
- See also 72, 87, 159, 199, 211, 226, 229, 235, 243, 247, 252, 258, 260, 262, 264, 292, 295, 376, 401, 421, 424, 433, 476, 479, 482, 518, 565, 595, 659, 678, 699, 740.
37. CELLS, CELLULAR PARTICLES AND SUPRAMOLECULAR STRUCTURES
- 796 Dworzanski, J.P. and Meuzelaar, H.L.C.: Rapid detection and identification of microorganisms via pyrolytic on-line derivatization-gas chromatography-mass spectrometry. In: Nelson, W.H. (Editor), *Mod. Tech. Rapid Microbiol. Anal.*, VCH, New York, 1991, pp. 19-42; C.A., 117 (1992) 146649z.
- See also 368, 779.
38. INORGANIC COMPOUNDS
- 38b. Anions
- 797 Sun, M., Yin, W., Ou, W. and Wang, L.: (Assay of fluorine in water by gas chromatography with derivatization.) *Sepu*, 10 (1992) 288-290.
- See also 801.
- 38c. Permanent and rare gases
- 798 Andrews, A.R.J., Wu, Z. and Zlatkis, A.: The separation of hydrogen and deuterium homologues by inclusion gas chromatography. *Chromatographia*, 34 (1992) 457-460.
- 799 Shen, Z., Briens, C.L., Kwauk, M. and Bergougnou, M.A.: Improved thermal conductivity detector technique for helium measurements in a fluidized bed. *Powder Technol.*, 69 (1992) 249-253; C.A., 117 (1992) 029542y.
- 800 Yoshida, H., Nishina, A. and Hirano, M.: (Study on analysis of ultratrace impurities in gases. 2. Analysis of impurities in special gases, by quadrupole GC/MS.) *Koatsu Gasu*, 29 (1992) 140-147; C.A., 116 (1992) 206837a.
- See also 65, 100, 102, 664, 803.
- 38d. Volatile inorganic compounds
- 801 Dai, K., Li, Y. and Shi, W.: (Preconcentration and determination of dimethylarsonic acid by cation exchange and derivatization gas chromatography.) *Huanjing Huaxue*, 11 (1992) 63-68; C.A., 117 (1992) 142607m.
- 802 Ding, T., Liu, B., Zhang, H., Liu, Y., Gau, Z. and Li, Z.: Determination of trace amounts of water in high-purity silane, arsine and phosphine by gas chromatography. *Analyst (London)*, 117 (1992) 1577-1579.
- 803 Drugov, Yu.S.: (Gas chromatography of inorganic substances.) *Zh. Anal. Khim.*, 47 (1992) 246-270.
- 804 Haunold, W., Georgii, H.-W. and Ockelmann, G.: Gas chromatographic analysis of atmospheric sulfur dioxide and reduced sulfur compounds. *LC-GC Int.*, 5, No.10 (1992) 28-35.
- 805 Ishigaki, H.: (Gas chromatograph method for determination of sulfur dioxide in sulfur tetrafluoride.) *Jpn. Kokai Tokkyo Koho JP 04,157,362 [92,157,362]* (Cl. G01N30/88), 29 May 1992, Appl. 90/280,802, 19 Oct. 1990; 4 pp.; C.A., 117 (1992) 103246y.
- 806 Lin, B., Wang, S., Li, X. and Hu, L.: (Gas chromatographic determination of trace of water in pure carbon dioxide.) *Huaxue Shijie*, 33 (1992) 78-81; C.A., 117 (1992) 82555f.
- 807 McCardell, A. and Fuhrmann, J.J.: Determination of persulfate-oxidizable carbon by gas chromatography. *Soil Biol. Biochem.*, 24 (1992) 615-616; C.A., 117 (1992) 127527s.
- 808 Nakamura, R. and Yoshikawa, S.: (Gas-chromatography method and apparatus for determining a low-concentration impurity in fluorine or chlorine gas.) *Jpn. Kokai Tokkyo Koho JP 04 09,757 [92 09,757]* (Cl. G01N30/88), 14 Jan. 1992, Appl. 90/113,543, 27 Apr. 1990; 5 pp.; C.A., 117 (1992) 39424a.
- 809 Shan, S.: (Gas chromatographic determination of iodine content in kelp.) *Shipin Kexue (Beijing)*, 136 (1991) 46-48; C.A., 116 (1992) 104575w.

- 810 Shooter, D., de Mora, S.J., Grout, A., Wylie, D.J. and Zhi-Yun, H.: The chromatographic analysis of reduced sulfur gases in Antarctic waters following pre-concentration onto Tenax. *Intern. J. Environ. Anal. Chem.*, 47 (1992) 239-249.
- 811 Sitaula, B.K., Luo, J. and Bakken, L.R.: Rapid analysis of climate gases by wide bore capillary gas chromatography. *J. Environ. Qual.*, 21 (1992) 493-496; *C.A.*, 117 (1992) 136652f.
- 812 Sung, Y.J., Hotchkiss, J.H., Austic, R.E. and Dietert, R.R.: Direct measurement of nitric oxide in headspace gas produced by a chicken macrophage cell line in a closed culture system. *Biochem. Biophys. Res. Commun.*, 184 (1992) 36-42; *C.A.*, 117 (1992) 024373j.
- 813 Zou, Z.: (Gas chromatography for trace-moisture determination in pyromellitic acid dianhydride.) *Nanjing Huagong Xueyuan Xuebao*, 13, No. 4 (1991) 81-84; *C.A.*, 116 (1992) 187279b.

See also 667, 800.

39. RADIOACTIVE AND OTHER ISOTOPE COMPOUNDS

- 814 Sugimoto, A., Hong, X. and Wada, E.: Rapid and simple measurement of carbon isotope ratio of bubble methane using GC/C/IRMS. *Shitsuryo Bunseki*, 39 (1991) 261-266; *C.A.*, 116 (1992) 200014h.
- 815 Youn, D.Y., Hong, K.B. and Jung, K.H.: Direct measurement of the inverse secondary isotope effects of Rh(I)-C₂H₄ and Rh(I)-C₂H₄D utilizing gas chromatography. *J. Chromatogr.*, 607 (1992) 69-77.

See also 22, 320, 350, 562, 798.

Planar Chromatography

1. REVIEWS AND BOOKS

- 1 Ackman, R.G.: Application of thin-layer chromatography to lipid separation: detection methods. *Anal. Fats, Oils Lipoproteins*, (1991) 97-121; C.A., 117 (1992) 146369h - a review with 43 refs.
- 2 Ojanpera, I.: Toxicological drug screening by thin-layer chromatography. *Trends Anal. Chem.*, 11 (1992) 222-229; C.A., 117 (1992) 144841g - a review with 55 refs.

See also 20, 26, 42, 115, 146, 211, 214, 215, 217, 238, 239, 253, 261, 262, 277.

2. FUNDAMENTALS, THEORY AND GENERAL

2a. General

- 3 Palmer, K. and Rabel, F.: Planar chromatography background contaminants: some causes and curves. *Am. Lab. (Shelton)*, 24 (1992) 28BB, 28DD; C.A., 117 (1992) 61696n.
- 4 Poole, S.K. and Poole, C.F.: The influence of the solvent entry position on resolution in unidimensional multiple development thin layer chromatography. *J. Planar Chromatogr.*, 5 (1992) 221-228.

2b. Thermodynamics and theoretical relationships

- 5 Kowalska, T. and Witkowska-Kita, B.: On the mixed mechanism of solute retention in RPTLC with alkyl-bonded stationary phases. *J. Planar Chromatogr.*, 4 (1991) 446-450.
- 6 Uhegbu, C.E., Kearns, M.L., Knotts, K.D., Julian, L.A., Klein, M.L. and Nurok, D.: Relationship between ΔR_f and a solvent strength parameter. *J. Planar Chromatogr.*, 5 (1992) 259-262.

See also 161, 208, 256, 304.

2c. Relationship between structure and chromatographic behaviour

- 7 Glen, R.C., Rose, V.S., Lindon, J.C., Ruane, R.J., Wilson, I.D. and Nicholson, J.K.: Quantitative structure - chromatography relationships: prediction of TLC behavior using theoretically derived molecular properties. *J. Planar Chromatogr.*, 4 (1991) 432-438.
- 8 Palamareva, M.D. and Kozekov, I.D.: Chromatographic behaviour of diastereoisomers XI. Steric effects and solvent selectivity effects in retentions on silica of esters of maleic and fumaric acids. *J. Chromatogr.*, 606 (1992) 113-119 - 20 compounds, 21 solvent systems.
- 9 Pogány, G., Cserhádi, T., Oláh, J. and Valkó, K.: Relationships between the chromatographic retention data and the effects of nucleoside derivatives in highly metastatic 3LL cells. *J. Pharm. Biomed. Anal.*, 10 (1992) 495-500.

See also 30, 117, 141, 159, 160, 260, 275, 279, 299.

2d. Measurement of physico-chemical and related values

See 254.

3. GENERAL TECHNIQUES

- 10 Owada, K.: (Factors affecting rate of flow (R_f) values on thin-layer chromatography). *Gijutsu Joho - Shizuoka-ken Eisei Kan-kyo Senta*, 10 (1992) 4-7; C.A., 113 (1992) 127542t.

3a. Apparatus and accessories

See 24, 182.

3b. Detectors and detection reagents

- 11 Ardic, Z., Yuksel, U., Ilkme, B. and Yuksel, M.: Two new chromatographic properties of sulfonated lignin. *J. Fac. Sci., Ege Univ., Ser. A*, 14, No. 1 (1991) 36-39; C.A., 117 (1992) 152990v.
- 12 Evgen'ev, M.I. and Levinson, F.S.: Derivatives of dinitrobenzofuran as selective reagents for organic analysis. *Khim. Geterotsikl. Soedin.*, (1991) 1565; C.A., 117 (1992) 123753q.
- 13 Kartnig, T. and Göbel, I.: Determination of hypericin and pseudohypericin by thin-layer chromatography-densitometry. *J. Chromatogr.*, 609 (1992) 423-426.
- 14 Riemer, F. and Dalrup, T.: Das Forrest-Reagens als Detektionsmittel für die Dünnschichtchromatographie. *Pharmazie*, 47 (1992) 559.
- 15 Wu, N. and Huie, C.W.: Synchronization of timing in chemiluminescence thin-layer chromatographic system by coupling pneumatic nebulization with optical fiber-based detection. *Anal. Chem.*, 64 (1992) 2465-2468.

See also 1, 21, 22, 29, 33, 111, 144, 163.

3c. Sorbents and columns, packing procedures

- 16 Cynamon, M.H., Klemens, S.P. and Swenson, C.E.: TLC G-65 in combination with other agents in the therapy of *Mycobacterium avium* injection in beige mice. *J. Antimicrob. Chemother.*, 29 (1992) 693-699; C.A., 117 (1992) 76387m.
- 17 Litvinova, L.S. and Kurenbin, O.I.: (Taking into account sorbent characteristics in an equation for the rate of motion of an eluent in thin-layer chromatography). *Zh. Fiz. Khim.*, 66 (1992) 730-735; C.A., 117 (1992) 138445q.
- 18 Mack, M. and Kinkel, J.: Sorbents for thin-layer chromatographic separations, their preparation, and their use. *Ger. Offen DE 4,005,868 (Cl. C08F292/00)*, 29 Aug. 1991, Appl. 1991, Feb. 1990; 8 pp.; C.A., 117 (1992) 123812h.
- 19 Petrovic, S.M. and Acanski, M.: Modification of the silica surface with the vapors of polar solvents. *J. Planar Chromatogr.*, 4 (1991) 439-441.

- 20 Shantha, N.C.: Thin-layer chromatography-flame ionization detection latroscan system. *J. Chromatogr.*, 624 (1992) 21-35 - a review with 60 refs.
- See also 3, 27, 35, 37, 65, 117, 137, 161, 164, 169, 201, 249, 302, 304.
- 3d. *Quantitative analysis*
- 21 Pollak, V.A., Doelemeyer, A., Winkler, W. and Schulze-Clewing, J.: Important design features of a system for the densitometric analysis of two-dimensional flat-bed separations. *J. Chromatogr.*, 596 (1992) 241-249.
- 22 Regnault, C., Delvordre, P., Bonnier, H. and Postaire, E.: Development of a specific device for densitometry of thin-layer chromatographic sheets in planar chromatography. *J. Chromatogr.*, 607 (1992) 159-162.
- See also 13, 144.
- 3e. *Preparative scale chromatography*
- See 31, 39, 46, 111, 155, 243.
- 3f. *Programmed temperature, pressure, vapors, gradients*
- See 24, 46.
- 3g. *High performance procedures*
- See 3, 68, 190, 252.
4. SPECIAL TECHNIQUES
- 4b. *Computerization and modelling*
- 23 Markowski, W. and Soczewinski, E.: Computer-aided optimization of gradient multiple-development thin-layer chromatography. I. Two-stage development. *J. Chromatogr.*, 623 (1992) 139-147.
- 24 Matysik, G., Markowski, W., Soczewinski, E. and Polak, B.: Computer-aided optimization of stepwise gradient profiles in thin-layer chromatography. *Chromatographia*, 34 (1992) 303-307.
- 25 Qin-Sun, W., Zhuang-Ping, Z. and Chun-Xia, W.: Computer-assisted optimization of the separation of a sample of unknown composition by HPTLC. *J. Planar Chromatogr.*, 4 (1991) 442-445.
- See also 2, 48.
- 4c. *Combination with other physico-chemical techniques (MS, IR etc.)*
- 26 Busch, K.L.: Mass spectrometric detection for thin-layer chromatographic separations. *TrAC*, 11 (1992) 314-324 - a review with 45 refs.
- 27 Danielson, N.D., Katon, J.E., Bouffard, S.P. and Zhu, Z.: Zirconium oxide stationary phase for thin-layer chromatography with diffuse reflectance Fourier transform infrared detection. *Anal. Chem.*, 64 (1992) 2183-2186.
- 28 Tajima, T., Wada, K. and Ichimura, K.: Zone transfer technique for diffuse reflectance Fourier transform infrared spectrometry of analytes separated by thin-layer chromatography. *Vib. Spectrosc.*, 3 (1992) 211-216; *C.A.*, 117 (1992) 61929.
- See also 44, 56, 62, 173, 238, 248, 266.
- 4e. *Functional analysis*
- See 123.
- 4g. *Enantiomers, separation*
- See 171, 172.
- 4h. *Other special techniques*
- 29 Stojanovic, R.S., Bond, A.M. and Butler, E.C.V.: A comparative study of the cylindrical wire, thin-layer, and wall-jet detector cells for the determination of inorganic arsenic by ion exclusion chromatography with constant and pulsed amperometric detection. *Electroanalysis (N.Y.)*, 4 (1992) 453-461; *C.A.*, 117 (1992) 118038f.
- See also 169, 249.
5. HYDROCARBONS AND HALOGEN DERIVATIVES
- 5b. *Cyclic hydrocarbons*
- 30 Qureshi, P.M. and Varshney, M.: The possible role of charge transfer in the chromatographic behavior of model carcinogens. *J. Planar Chromatogr.*, 4 (1991) 497-498.
- See also 288, 289, 296.
- 5d. *Complex hydrocarbon mixtures (incl. analysis of tars, bitumens and mineral oils)*
- 31 Lian, H., Lee, C.Z.H., Wang, Y.Y. and Yen, T.F.: Characterization of asphalt with the preparative Chromatotron. *J. Planar Chromatogr.*, 5 (1992) 263-266.
6. ALCOHOLS
- 32 Kupferberg, A., Tomassoni, M.L. and Mersel, M.: Separation and purification of dolichol and dolichyl phosphate by anion-exchange paper chromatography: application to cultured cells. *Anal. Biochem.*, 206 (1992) 236-240.
- 33 Pyka, A. and Sliwiok, J.: Application of thymol blue and bromothymol blue to the detection of the alcohols in TLC. *J. Planar Chromatogr.*, 5 (1992) 282-283.

- 34 Swiezewska, E., Chojnacki, T., Jankowski, W.J., Singh, A.K. and Olsson, J.: The occurrence of long chain polyprenols in leaves of plants of *Rosaceae* family and their isolation by time-extended liquid chromatography. *Biochem. Cell Biol.*, 70 (1992) 448-454.

See also 51.

7. PHENOLS

- 35 Mohammad, A., Ajmal, M. and Anwar, S.: Retention behavior of phenols on silica gel layers impregnated with sodium nitrite. *J. Planar Chromatogr.*, 5 (1992) 286-289.
- 36 Montedoro, G., Servili, M., Baldioli, M. and Miniati, E.: Simple and hydrolyzable phenolic compounds in virgin olive oil. 2. Initial characterization of the hydrolyzable fraction. *J. Agric. Food Chem.*, 40 (1992) 1577-1580.
- 37 Petrovic, M., Kastelan-Macan, M. and Horvat, A.J.M.: Thin-layer chromatographic behaviour of substituted phenolic compounds on silica gel layers impregnated with Al(III) and Cu(II). *J. Chromatogr.*, 607 (1992) 163-167.

See also 11, 289.

8. SUBSTANCES CONTAINING HETEROCYCLIC OXYGEN

8a. Flavonoids

- 38 Gueldner, R.C., Snook, M.E., Widstrom, N.W. and Wiseman, B.R.: TLC screen for maysin, chlorogenic acid and other possible resistance factors to the fall armyworm and the corn earworm in *Zea mays*. *J. Agric. Food Chem.*, 40 (1992) 1211-1213.
- 39 Proliac, A., Chaboud, A. and Raynaud, J.: Isolement et identification de trois C-glycosylflavones dans les tiges feuillées d'*Arum dracuncululus*. *Pharmazie*, 47 (1992) 646-647.
- 40 Yu, S., Liu, X., Yin, S. and Wang, H.: (Determination of baicalin content in Xiaochaihutang oral liquid). *Zhongcaoyao*, 23 (1992) 239-240; *C.A.*, 117 (1992) 137747c.

See also 290, 293, 300.

8b. Aflatoxins and other mycotoxins

- 41 Drusany, I., Kravanja, R. and Prosek, M.: Quantitative two-dimensional thin layer chromatography of aflatoxins. *J. Planar Chromatogr.*, 4 (1991) 490-492.
- 42 Holcomb, M., Wilson, D.M., Trucksess, M.W. and Thompson, H.C., Jr.: Determination of aflatoxins in food products by chromatography. *J. Chromatogr.*, 624 (1992) 341-352 - a review with 97 refs.
- 43 Mellon, J.E.: Inhibition of aflatoxin production in *Aspergillus flavus* by cotton ovule extracts. *J. Am. Oil Chem. Soc.*, 69 (1992) 945-947.
- 44 Mueller, M. and Lepom, P.: (Analysis of *Alternaria* mycotoxins in laboratory cultures). *Zentralbl. Microbiol.*, 147 (1992) 197-206; *C.A.*, 117 (1992) 107524h.

- 45 Sachse, J.: Identification and determination of mellein in cultures of the fungus *Septoria nodorum* (Berk.) by thin-layer and high-performance liquid chromatography. *J. Chromatogr.*, 609 (1992) 349-353.

See also 143.

8c. Other compounds with heterocyclic oxygen (incl. tannins)

- 46 Cisowski, W., Palka-Gudyka, E., Krauze-Baranowska, M. and Króllicki, Z.A.: Preparative TLC of coumarins with gradient elution. *J. Planar Chromatogr.*, 4 (1991) 471-474.
- 47 Davies, A.J. and Mazza, G.: Separation and characterization of anthocyanins of *Monarda fistulosa* by high-performance liquid chromatography. *J. Agric. Food Chem.*, 40 (1992) 1341-1345.
- 48 Härmälä, P.: Retardation behavior and prediction of R_f values in TLC using the 'PRISMA' model. I. Regular part of the model. *J. Planar Chromatogr.*, 4 (1991) 460-466.
- 49 Wolski, T., Glowniak, T. and Doraczynska-Szopa, A.: (Tetrachloroethylene as a component of binary and ternary eluents for separation of coumarins by thin-layer chromatography). *Chem. Anal. (Warsaw)*, 36 (1991) 299-304; *C.A.*, 117 (1992) 39527m.

9. OXO COMPOUNDS, ETHERS, EPOXIDES AND QUINONES

- 50 Blee, E. and Schuber, F.: Enantioselectivity of the hydrolysis of linoleic acid monoepoxides catalyzed by soybean fatty acid epoxide hydrolase. *Biochem. Biophys. Res. Commun.*, 187 (1992) 171-177.
- 51 Piwowarczyk, K. and Kryszczyńska, H.: (Determination of products of the reduction of benzylideneacetone in chloride galvanizing baths agitated and not agitated with air). *Powloki Ochr.*, 18, No. 4-6 (1990) 16-27; *C.A.*, 117 (1992) 61969d.
- 52 Weng, X.C. and Gordon, M.H.: Antioxidant activity of quinones extracted from tanshen (*Salvia miltiorrhiza* Bunge). *J. Agric. Food Chem.*, 40 (1992) 1331-1336.

See also 5, 288.

10. CARBOHYDRATES

10a. Mono and oligosaccharides. Structural studies

- 53 Akiyama, H., Shidawara, S., Mada, A., Toyoda, H., Toida, T. and Imanari, T.: Chemiluminescence high-performance liquid chromatography for the determination of hyaluronic acid, chondroitin sulphate and dermatan sulphate. *J. Chromatogr.*, 579 (1992) 203-207.
- 54 Araki, S., Abe, S., Yamada, S., Satake, M., Fujiwara, N., Kon, K. and Ando, S.: Characterization of two novel pyruvylated glycosphingolipids containing 2'-aminoethylphosphoryl(→6)-galactose from the nervous system of *Aplysia kurodai*. *J. Biochem. (Tokyo)*, 112 (1992) 461-469.
- 55 Cantor, A.B. and Kornfeld, S.: A method for [3 H]mannose labeling of Asn-linked oligosaccharides on recombinant glycoproteins synthesized in *Xenopus* oocytes. *Anal. Biochem.*, 205 (1992) 220-226.

- 56 El-Kheir, A.A., Metwally, M.M.F. and Ayad, M.M.: Spectrophotometric methods in conjunction with thin layer chromatography applied for determination of sugars content in Egyptian honey and glucose syrup. *Egypt. J. Food Sci.*, 19 (1991) 41-51; C.A., 117 (1992) 130039q.
- 57 Nakada, H., Inoue, M., Numata, Y., Tanaka, N., Funakoshi, I., Fukui, S. and Yamashima, I.: Cancer-associated glycoproteins defined by a monoclonal antibody, MLS 128, recognized the Th antigen. *Biochem. Biophys. Res. Commun.*, 187 (1992) 217-224.
- 58 Skudlarek, M.D., Tulsiani, D.R.P. and Orgebin-Crist, M.-C.: Rat epididymal luminal fluid acid β -D-galactosidase optimally hydrolyses glycoprotein substrate at neutral pH. *Biochem. J.*, 286 (1992) 907-914.
- 59 Takeya, A., Hosomi, O., Shimoda, N. and Yazawa, S.: Biosynthesis of the blood group P antigen-like GalNAc β -3Gal β 1-4GlcNAc/Glc structure: a novel N-acetylgalactosaminyltransferase in human blood plasma. *J. Biochem. (Tokyo)*, 112 (1992) 389-395.
- 60 Urashima, T., Messer, M. and Bubb, W.A.: Biosynthesis of marsupial milk oligosaccharides II: characterization of a β 6-N-acetylglucosaminyltransferase in lactating mammary glands of the tammar wallaby, *Macropus eugenii* *Biochim. Biophys. Acta*, 1117 (1992) 223-231.
- 61 Wang, J., Xu, P. and Fincher, G.B.: Purification, characterization and gene structure of (1 \rightarrow 3)- β -glucanase isoenzyme GIII from barley (*Hordeum vulgare*). *Eur. J. Biochem.*, 209 (1992) 103-109.
- See also 109, 186.
- 10b. *Polysaccharides, mucopolysaccharides, lipopolysaccharides*
- 62 Pekic, B. and Lepojevic, Z.: (Determination of inulin by thin-layer chromatography on kieselguhr G). *Zb. Rad. - Technol. Fak. Novom Sadu*, (1990) 95-100; C.A., 117 (1992) 166901d.
11. ORGANIC ACIDS AND LIPIDS
- 11a. *Organic acids and simple esters*
- 63 Banerjee, P., Dawson, G. and Dasgupta, A.: Enrichment of saturated fatty acid containing phospholipids in sheep brain serotonin receptor preparations: use of microwave irradiation for rapid transesterification of phospholipids. *Biochim. Biophys. Acta*, 1110 (1992) 65-74.
- 64 Möhring, H. and Spittler, G.: Produkte der Dimerisierung ungesättigter Fettsäuren VIII: Über die Zusammensetzung der Fraktion der "Intermediates" bei der Fettsäuredimerisierung. *Fat Sci. Technol.*, 94 (1992) 241-245.
- 65 Nikolova-Damyanova, B., Herslof, B.G. and Christie, W.W.: Silver ion high-performance liquid chromatography of derivatives of isomeric fatty acids. *J. Chromatogr.*, 609 (1992) 133-140.
- 66 Qian, Z., Yan, W., He, Y. and Lu, G.: TLC-densitometric determination of benzoic acid and miconazole nitrate in paint. *Zhongguo Yiyuan Yaoxue Zazhi*, 11 (1991) 435-437; C.A., 117 (1992) 50797v.
- 67 Smith, N.B. and Brooks, K.E.: Use of Toxi-Lab thin layer chromatography in plasma very long chain fatty acid analysis. *Clin. Chim. Acta*, 206 (1992) 241-243.
- 68 Tibi, A., Steinmetz, A.C., Goury, V., Decool, A. and Darbord, J.C.: Mycolic acids and nocardomycolate analysis by overpressured thin layer chromatography. *J. Microbiol. Methods*, 15 (1992) 199-206; C.A., 117 (1992) 166899j.
- 69 Ulberth, F. and Henninger, M.: Simplified method for the determination of *trans* monoenes in edible fats by TLC-GLC. *J. Am. Oil Chem. Soc.*, 69 (1992) 829-831.
- 70 Winterbourn, C.C., van den Berg, J.J.M., Roitman, E. and Kuypers, F.A.: Chlorohydrin formation from unsaturated fatty acids reacted with hypochlorous acid. *Arch. Biochem. Biophys.*, 296 (1992) 547-555.
- See also 7, 8, 36, 37, 38, 50, 95, 99, 105, 289, 290.
- 11c. *Lipids and their constituents*
- 71 Anderson, V.C. and Thompson, D.H.: Photoinduced morphological changes in plasmalogen liposomes using visible light. *ACS Symp. Ser.*, 493 (Macromol. Assem. Polym. Syst.) (1992) 154-170; C.A., 117 (1992) 76410p.
- 72 Ardail, D., Lerne, F. and Louisot, P.: Phospholipid import into mitochondria: possible regulation mediated through lipid polymorphism. *Biochem. Biophys. Res. Commun.*, 186 (1992) 1384-1390.
- 73 Arduini, A., Tyurin, V., Tyuruna, Y., Arrigoni-Martelli, E., Molajoni, F., Dottori, S. and Federici, G.: Acyl-trafficking in membrane phospholipid fatty acid turnover: the transfer of fatty acid from the acyl-L-carnitine pool to membrane phospholipids in intact human erythrocytes. *Biochem. Biophys. Res. Commun.*, 187 (1992) 353-358.
- 74 Atsum, S., Nosaka, C., Iinuma, H. and Umezawa, K.: Inhibition of glucocerebrosidase and induction of neural abnormality by cyclophellitol in mice. *Arch. Biochem. Biophys.*, 297 (1992) 362-367.
- 75 Bai, H., Orlando, J. and Seyfried, T.N.: Altered ganglioside composition in virally transformed rat embryo fibroblasts. *Biochim. Biophys. Acta*, 1136 (1992) 23-27.
- 76 Bütikofer, P., Zollinger, M. and Brodbeck, U.: Alkylacyl glycerophosphoinositol in human and bovine erythrocytes. Molecular species composition and comparison with glycosylinositolphospholipid anchors of erythrocyte acetylcholinesterases. *Eur. J. Biochem.*, 208 (1992) 677-683.
- 77 Byrne, C.D., Wang, T.W.M. and Hales, C.N.: Control of Hep G2-cell triacylglycerol and apolipoprotein B synthesis and secretion of polyunsaturated non-esterified fatty acids and insulin. *Biochem. J.*, 288 (1992) 101-107.
- 78 Chabot, M.C., McPhail, L.C., Wykle, R.L., Kennerly, D.A. and McCall, C.E.: Comparison of diglyceride production from choline-containing phosphoglycerides in human neutrophils stimulated with N-formylmethionylleucylphenylalanine, ionophore A23187 or phorbol 12-myristate 13-acetate. *Biochem. J.*, 286 (1992) 693-699.
- 79 Chen, H., Born, E., Mathur, S.N., Johlin, F.C. and Field, F.J.: Sphingomyelin content of intestinal cell membranes regulates cholesterol absorption. Evidence for pancreatic and intestinal cell sphingomyelinase activity. *Biochem. J.*, 286 (1992) 771-777.

- 80 Chen, J.-W., Shi, K., Zhang, L. and Huang, F.: Effect of glycolipids on the phase behavior and dynamic properties of phospholipid liposomes. *Biochem. Biophys. Res. Commun.*, 186 (1992) 1294-1298.
- 81 Cronholm, T., Viestam-Rains, M. and Sjövall, J.: Decreased content of arachidonoyl species of phosphatidylinositol phosphates in pancreas of rats fed on an ethanol-containing diet. *Biochem. J.*, 287 (1992) 925-928.
- 82 Czarny, M., Sabala, P., Ucieklak, A., Kaczmarek, L. and Baranska, J.: Inhibition of phosphatidylserine synthesis by glutamate, acetylcholine, thapsigargin and ionophore A23187 in glioma C6 cells. *Biochem. Biophys. Res. Commun.*, 186 (1992) 1582-1587.
- 83 Dahiya, R., Boyle, B., Goldberg, B.C., Yoon, W.-H., Konety, B., Chen, K., Yen, T.-S.B., Blumenfeld, W. and Narayan, P.: Metastasis-associated alterations in phospholipids and fatty acids of human prostatic adenocarcinoma cell lines. *Biochem. Cell Biol.*, 70 (1992) 548-554.
- 84 Davey, M.W. and Lambein, F.: Quantitative derivatization and high-performance liquid chromatographic analysis of cyanobacterial heterocyst-type glycolipids. *Anal. Biochem.*, 206 (1992) 323-327.
- 85 Davey, M.W. and Lambein, F.: Semipreparative isolation of individual cyanobacterial heterocyst-type glycolipids by reverse-phase high-performance liquid chromatography. *Anal. Biochem.*, 206 (1992) 226-230.
- 86 Deeg, M.A., Murray, N.R. and Rosenberry, T.L.: Identification of glycoinositol phospholipids in rat liver by reductive radiomethylation of amines but not in H4IIE hepatoma cells or isolated hepatocytes by biosynthetic labeling with glucosamine. *J. Biol. Chem.*, 267 (1992) 18581-18588.
- 87 Fukami, K. and Takenawa, T.: Phosphatidic acid that accumulates in platelet-derived growth factor-stimulated Balb/c 3T3 cells is a potential mitogenic signal. *J. Biol. Chem.*, 267 (1992) 10988-10993.
- 88 Gadella, B.M., Colenbrander, B., van Golde, L.M.G. and Lopes-Cardozo, M.: Characterization of three arylsulfatases in semen: seminolipid sulfohydrolase activity is present in seminal plasma. *Biochim. Biophys. Acta*, 1128 (1992) 155-162.
- 89 Goldin, E., Roff, C.F., Miller, S.P.F., Rodriguez-Lafrasse, C., Vanier, M.T., Brady, R.O. and Pentchev, P.G.: Type C Niemann-Pick disease: a murine model of the lysosomal cholesterol lipidosis accumulates sphingosine and sphinganine in liver. *Biochim. Biophys. Acta*, 1127 (1992) 303-311.
- 90 Hattori, M.-a. and Horiuchi, R.: Enhancement of ganglioside GM3 synthesis in okadaic-acid-treated granulosa cells. *Biochim. Biophys. Acta*, 1137 (1992) 101-106.
- 91 Heller, M., Bieri, S. and Brodbeck, U.: A novel form of glycosylphosphatidylinositol-anchor converting activity with a specificity of a phospholipase D in mammalian liver membranes. *Biochim. Biophys. Acta*, 1109 (1992) 109-116.
- 92 Hirabayashi, Y., Nakao, T., Irie, F., Whittaker, V.P., Kon, K. and Ando, S.: Structural characterization of a novel cholinergic neuron-specific ganglioside in bovine brain. *J. Biol. Chem.*, 267 (1992) 12973-12978.
- 93 Holgersson, J., Bäcker, A.E., Breimer, M.E., Gustavsson, M.L., Jovall, P.-A., Karlsson, H., Pimlott, W. and Samuelsson, B.E.: The blood group B type-4 heptaglycosylceramide is a minor blood group B structure in human B kidneys in contrast to the corresponding A type-4 compound in A kidneys. Structural and *in vitro* biosynthetic studies. *Biochim. Biophys. Acta*, 1180 (1992) 33-43.
- 94 Hvattum, E., Skrede, S., Bremer, J. and Solbakken, M.: The metabolism of tetradecylthiopropionic acid, a 4-thia stearic acid, in the rat. *In vivo* and *in vitro* studies. *Biochem. J.*, 286 (1992) 879-887.
- 95 Johnson, S.B. and Brown, R.E.: Simplified derivatization for determining sphingolipid fatty acyl composition by gas chromatography-mass spectrometry. *J. Chromatogr.*, 605 (1992) 281-286.
- 96 Kamio, K., Gasa, S. and Makita, A.: Galactosylceramide containing ω -aminofatty acids: preparation, characterization, and sulfotransferase acceptor. *J. Lipid Res.*, 33 (1992) 1227-1232.
- 97 Kasurinen, J.: A novel fluorescent fatty acid, 5-methyl-BDY-3-dodecanoic acid, is a potential probe in lipid transport studies by incorporating selectively to lipid classes of BHK cells. *Biochem. Biophys. Res. Commun.*, 187 (1992) 1594-1601.
- 98 Kaszkin, M., Seidler, L., Kast, R. and Kinzel, V.: Epidermal-growth-factor-induced production of phosphatidylalcohols by HeLa cells and A431 cells through activation of phospholipase D. *Biochem. J.*, 287 (1992) 51-57.
- 99 Konrad, R.J., Jolly, Y.C., Major, C. and Wolf, B.A.: Carbachol stimulation of phospholipase A₂ and insulin excretion in pancreatic islets. *Biochem. J.*, 287 (1992) 283-290.
- 100 Kotani, M., Ozawa, H., Kawashima, I., Ando, S. and Tai, T.: Generation of one set of monoclonal antibodies specific for a pathway ganglio-series gangliosides. *Biochim. Biophys. Acta*, 1117 (1992) 97-103.
- 101 Kubo, H., Jiang, G.J., Irie, A., Morita, M., Matsubara, T. and Hoshi, M.: A novel ceramide trihexoside from the eggs of the sea urchin, *Hemicentrotus pulcherrimus*. *J. Biochem. (Tokyo)*, 111 (1992) 726-731.
- 102 Kuroki, Y., Gasa, S., Ogasawara, Y., Shiratori, M., Makita, A. and Akino, T.: Binding specificity of lung surfactant protein SP-D for glucosylceramide. *Biochem. Biophys. Res. Commun.*, 187 (1992) 963-969.
- 103 Laszlo, A., Papa, F. and David, H.L.: Thin-layer chromatography systems for the identification of *Mycobacterium tuberculosis*, *M. bovis* BCG, *M. kansasii*, *M. gastri* and *M. marinum* Egypt. *J. Biochem.*, 143 (1992) 519-524; C.A., 117 (1992) 187511s.
- 104 Lavery, S.B., Nudelman, E.D. and Hakomori, S.: Novel modification of glycosphingolipids by long-chain cyclic acetals: isolation and characterization of plasmalocerebroside from human brain. *Biochemistry*, 31 (1992) 5335-5340.
- 105 Li, M. and Ishibashi, T.: An *in vitro* reversible interconversion of rat liver fatty acid binding protein having different isoelectric points by virtue of the fatty acid content. *Arch. Biochem. Biophys.*, 298 (1992) 254-258.
- 106 Lin, P., Fung, W.-J.C. and Giffillan, A.M.: Phosphatidylcholine-specific phospholipase D-derived 1,2-diacylglycerol does not initiate protein kinase C activation in the RBL 2H3 mast-cell line. *Biochem. J.*, 287 (1992) 325-331.

- 107 Ly, H.L., Mortimer, B.-C., Baker, E. and Redgrave, T.G.: Clearance from plasma of lymph chylomicrons and chylomicron remnants labelled with ^{125}I -tyramine-cellobiose. *Biochem. J.*, 286 (1992) 937-943.
- 108 Matter, W.F., Brown, R.F. and Vlahos, C.J.: The inhibition of phosphatidylinositol 3-kinase by quercetin and analogs. *Biochem. Biophys. Res. Commun.*, 186 (1992) 624-631.
- 109 Milne, K.G., Ferguson, M.A.J. and Masterson, W.J.: Inhibition of the GlcNAc transferase of the glycosylphosphatidylinositol anchor biosynthesis in African trypanosomes. *Eur. J. Biochem.*, 208 (1992) 309-314.
- 110 Murakami, M., Kudo, I., Suwa, Y. and Inoue, K.: Release of 14-kDa group-II phospholipase A_2 from activated mast cells and its possible involvement in the regulation of the degranulation process. *Eur. J. Biochem.*, 209 (1992) 257-265.
- 111 Mürthing, J. and Unland, F.: Detection of gangliosides with the fluorochrome NBD dihexadecylamine and its application for preparative high performance thin layer chromatography. *Biomed. Chromatogr.*, 6 (1992) 227-230.
- 112 Nilsson, A., Hjelte, L. and Strandvik, B.: Incorporation of dietary [^{14}C]arachidonic acid and [^3H]eicosapentaenoic acid into tissue lipids during absorption of a fish oil emulsion. *J. Lipid Res.*, 33 (1992) 1295-1305.
- 113 Nohara, K., Suzuki, M., Inagaki, F., Sano, T. and Kaya, K.: A novel disialoganglioside in rat spleen lymphocytes. *J. Biol. Chem.*, 267 (1992) 14982-14986.
- 114 Nudelman, E.D., Levery, S.B., Igarashi, Y. and Hakomori, S.: Plasmalopsychosine, a novel plasmal (fatty aldehyde) conjugate of psychosine with cyclic acetal linkage. Isolation and characterization from human brain white matter. *J. Biol. Chem.*, 267 (1992) 11007-11016.
- 115 Olsson, N.U.: Advances in planar chromatography for the separation of food lipids. *J. Chromatogr.*, 624 (1992) 11-19 - a review with 40 refs.
- 116 Palmer, F.B.S.C., Byers, D.M., Spence, M.W. and Cook, H.W.: Calcium-independent effects of TMB-8. Modification of phospholipid metabolism in neuroblastoma cells by inhibition of choline uptake. *Biochem. J.*, 286 (1992) 505-512.
- 117 Parrish, C.C., Bodenne, G. and Gentien, P.: Separation of polyunsaturated and saturated lipids from marine phytoplankton on silica gel-coated Chromarods. *J. Chromatogr.*, 607 (1992) 97-104.
- 118 Paul, P., Bordmann, A., Rosenfelder, G. and Towbin, H.: Simultaneous determination of sugar incorporation into glycosphingolipids and glycoproteins. *Anal. Biochem.*, 204 (1992) 265-272.
- 119 Previti, M., Dotta, F., Pontieri, G.M., Di Mario, U. and Lenti, L.: Determination of gangliosides by high-performance liquid chromatography with photodiode-array detection. *J. Chromatogr.*, 605 (1992) 221-225.
- 120 Ren, S., Scarsdale, J.N., Ariga, T., Zhang, Y., Klein, R.A., Hartmann, R., Kushi, Y., Egge, H. and Yu, R.K.: O-Acetylated, gangliosides in bovine buttermilk. Characterization of 7-O-acetyl, and 7,9-di-O-acetyl G_{D3} . *J. Biol. Chem.*, 267 (1992) 12632-12638.
- 121 Rubin, D. and Laposata, M.: Cellular interactions between n -6 and n -3 fatty acids: a mass analysis of fatty acid elongation/desaturation, distribution among complex lipids, and conversion to eicosanoids. *J. Lipid Res.*, 33 (1992) 1431-1440.
- 122 Schuurmans Stekhoven, F.M.A.H., Tesser, G.I., Ramsteyn, G., Swarts, H.G.P. and de Pont, J.J.H.H.M.: Binding of ethylenediamine to phosphatidylserine is inhibitory to Na^+/K^+ -ATPase. *Biochim. Biophys. Acta*, 1109 (1992) 17-32.
- 123 Serizawa, S., Osawa, K., Togashi, K., Yamamoto, A., Ito, M., Hamanaka, S. and Otsuka, F.: Relationship between cholesterol sulfate and intercellular cohesion of the stratum corneum: demonstration using a push-pull meter and an improved high-performance thin-layer chromatographic separation system of all major stratum corneum lipids. *J. Invest. Dermatol.*, 99 (1992) 232-236; *C.A.*, 117 (1992) 128865n.
- 124 Sjöberg, E.R., Manzi, A.E., Khoo, K., Dell, A. and Varki, A.: Structural and immunological characterization of O-acetylated G_{D2} . Evidence that G_{D2} is an acceptor for ganglioside O-acetyltransferase in human melanoma cells. *J. Biol. Chem.*, 267 (1992) 16200-16211.
- 125 Slomiany, A., Grzelinska, E., Grabska, M., Yamaki, K.-I., Tamura, S., Kasinathan, C. and Slomiany, B.L.: Intracellular processes associated with glycoprotein transport and processing. *Arch. Biochem. Biophys.*, 298 (1992) 167-175.
- 126 Sonnino, S., Acquotti, D., Kirschner, G., Aguaglianza, A., Zecca, L., Rubino, F. and Tettamanti, G.: Preparation of lyso-GM1 ($^{13}\text{Neu}5\text{AcGgOse}_4$ -long chain bases) by a one-pot reaction. *J. Lipid Res.*, 33 (1992) 1221-1226.
- 127 Sorisky, A., King, W.G. and Rittenhouse, S.E.: Accumulation of $\text{PtdIns}(3,4)\text{P}_2$ and $\text{PtdIns}(3,4,5)\text{P}_3$ in thrombin-stimulated platelets. Different sensitivities of Ca^{2+} or functional integrin. *Biochem. J.*, 286 (1992) 581-584.
- 128 Tadano-Aritomi, K., Kasama, T., Handa, S. and Ishizuka, I.: Isolation and structural characterization of a mono-sulfated isoglobotetraosylceramide, the first sulfoglycosphingolipid of the isogloboseries, from rat kidney. *Eur. J. Biochem.*, 209 (1992) 305-313.
- 129 Taki, T., Rokukawa, C., Kasama, T., Kon, K., Ando, S., Abe, T. and Handa, S.: Human meconium gangliosides. Characterization of a novel I-type ganglioside with the $\text{NeuAc}\alpha 2\text{-6Gal}$ structure. *J. Biol. Chem.*, 267 (1992) 11811-11817.
- 130 Tamai, Y., Kojima, H., Saito, S., Takayama-Abe, K. and Horichi, H.: Characteristic distribution of glycolipids in gadoid fish nerve tissues and its bearing on phylogeny. *J. Lipid Res.*, 33 (1992) 1351-1359.
- 131 Terabayashi, T., Tsuda, M. and Kawanishi, Y.: The characteristic negative Cotton effect of ganglioside lactones observed by circular dichroism spectrometry. *Anal. Biochem.*, 204 (1992) 15-21.
- 132 Thorn, J.J., Levery, S.B., Salyan, M.E.K., Stroud, M.R., Cedergren, B., Nilsson, B., Hakamori, S. and Clausen, H.: Structural characterization of x_2 glycosphingolipid, its extended form, and its sialosyl derivatives: accumulation associated with the rare blood group p phenotype. *Biochemistry*, 31 (1992) 6509-6517.
- 133 Tomavo, S., Dubremetz, J.-F. and Schwarz, R.T.: A family of glycolipids from *Toxoplasma gondii*. Identification of candidate glycolipid precursor(s) for *Toxoplasma gondii* glycosylphosphatidylinositol membrane anchors. *J. Biol. Chem.*, 267 (1992) 11721-11728.
- 134 Triggiani, M., Fonteh, A.N. and Chilton, F.H.: Factors that influence the proportions of platelet-activating factor and 1-acyl-2-acetyl-sn-glycero-3-phosphocholine synthesized by the mast cell. *Biochem. J.*, 286 (1992) 497-503.

- 135 Van Veldhoven, P.P., Matthews, T.J., Bolognesi, D.P. and Bell, R.M.: Changes in bioactive lipids, alkylacylglycerol and ceramide, occur in HIV-infected cells. *Biochem. Biophys. Res. Commun.*, 187 (1992) 209-216.
- 136 Van Wijk, G.M.T., Hostetler, K.Y. and van den Bosch, H.: Antiviral nucleoside diphosphate diglycerides: improved synthesis and facilitated purification. *J. Lipid Res.*, 33 (1992) 1211-1219.
- 137 Wang, Wen-Qi and Gustafson, A.: One-dimensional thin-layer chromatographic separation of phospholipids and lysophospholipids from tissue lipid extracts. *J. Chromatogr.*, 581 (1992) 139-142.
- 138 Yamashita, G., Tazuma, S. and Kajiyama, G.: Effects of organic anions on biliary lipid secretion in rats. Importance of association with biliary lipid structures. *Biochem. J.*, 286 (1992) 193-196.
- 139 Yohe, H.C., Macala, L.J., Giordano, G. and McMurray, W.J.: GM_{1b} and GM_{1b}-GalNAc: major gangliosides of murine-derived macrophage-like WEHI-3 cells. *Biochim. Biophys. Acta*, 1109 (1992) 210-217.
- See also 1, 54, 63, 146, 152, 298.
13. STEROIDS
- 13a. General techniques
- See 6, 19, 259.
- 13b. Pregnane and androstane derivatives
- 140 Guo, W.: (Studies on a by-product with higher melting point in manufacture of epiandrosterone). *Zhongguo Yiyao Gongye Zazhi*, 22 (1991) 548-551; *C.A.*, 117 (1992) 118307t.
- 141 Parini, C., Bacigalupo, M.A., Colombi, S. and Saita, R.: Reversed phase TLC and HPLC of hemisuccinyl derivatives of 11 α -hydroxyprogesterone. *J. Planar Chromatogr.*, 5 (1992) 251-254.
- 142 Tang, G., Chen, Q. and He, B.: (Determination of triamcinolone acetonide in membrane by thin-layer chromatography). *Sepu*, 10 (1992) 186-187; *C.A.*, 117 (1992) 76600a.
- See also 259.
- 13c. Estrogens
- 143 Medina, M.B. and Schwartz, D.P.: Thin-layer chromatographic detection of zeranol and estradiol in fortified plasma and tissue extracts with Fast Corin V. *J. Chromatogr.*, 581 (1992) 119-128.
- 144 Windhorst, G. and de Kleijn, J.P.: Some technical recommendations for improving the performance of quantitative thin layer chromatography. *J. Planar Chromatogr.*, 5 (1992) 229-233.
- 13d. Sterols
- 145 Björkhem, I., Nyberg, B. and Einarsson, K.: 7 α -Hydroxylation of 27-hydroxycholesterol in human liver microsomes. *Biochim. Biophys. Acta*, 1128 (1992) 73-76.
- 146 Fenton, M.: Chromatographic separation of cholesterol in foods. *J. Chromatogr.*, 624 (1992) 369-388 - a review with 117 refs.
- 147 Genedella, R.J., Linton, L.L. and Moore, C.P.: Cholesterylene, a new recognized tissue lipid, found at high levels in the cornea. *Biochem. Biophys. Res. Commun.*, 186 (1992) 1647-1655.
- 148 Keesler, G.A., Casey, W.M. and Parks, L.W.: Stimulation by heme of steryl ester synthase and aerobic sterol exclusion in the yeast *Saccharomyces cerevisiae*. *Arch. Biochem. Biophys.*, 296 (1992) 474-481.
- 149 Maechler, P., Wollheim, C.B., Bentzen, C.L. and Niesor, E.: Role of the intestinal acyl-CoA:cholesterol acyltransferase activity in the hyperresponse of diabetic rats to dietary cholesterol. *J. Lipid Res.*, 33 (1992) 1475-1484.
- 150 Ruecker, G., Amer, M.M., Khattab, F.I. and El-Bayoumi, A.: Titrimetric determination of lynestrenol and norethynodrel with N-bromosuccinimide, and identification of the reaction products. *Egypt. J. Chem.*, 32 (1989, Publ. 1991) 207-217; *C.A.*, 117 (1992) 157750b.
- 151 Shefer, S., Nguyen, L.B., Salen, G., Ness, G.C., Chowdhary, I.R., Lerner, S., Batta, A.K. and Tint, G.S.: Differing effects of cholesterol and taurocholate on steady state hepatic HMG-CoA reductase and cholesterol 7 α -hydroxylase activities and mRNA levels in the rat. *J. Lipid Res.*, 33 (1992) 1193-1200.
- 152 Subbaiah, P.V., Liu, M., Bolan, P.J. and Paltauf, F.: Altered positional specificity of human plasma lecithin-cholesterol acyltransferase in the presence of sn-2 arachidonoyl phosphatidylcholines. Mechanism of formation of saturated cholesteryl esters. *Biochim. Biophys. Acta*, 1128 (1992) 83-92.
- See also 83.
- 13e. Bile acids and alcohols
- 153 Hashimoto, F., Sugimoto, C. and Hayashi, H.: Effects of aminotriazole treatment on biosyntheses of primary bile acids *in vivo*. *Chem. Pharm. Bull.*, 40 (1992) 795-798.
- 13g. Other steroids
- See 142.
14. STEROID GLYCOSIDES AND SAPONINS
- 154 Choi, K.J., Ko, S.R., Jeon, B.S. and Kim, M.W.: (Extraction yields on ginseng saponins in the extracts of crude drug preparations with various concentrations of ethanol). *Koryo Insam Hakhoechi*, 156 (1991) 192-196; *C.A.*, 117 (1992) 55756r.
- 155 Ferth, R. and Kopp, B.: Cardenolide aus *Ornithogalum umbellatum* L. *Pharmazie*, 47 (1992) 626-629.
- 156 Ko, S.R., Kim, S.C. and Choi, K.J.: (Extract yields and saponin contents of red ginseng extracts prepared with various concentrations of ethanol). *Saengyak Hakhoechi*, 23, No. 1 (1992) 24-28; *C.A.*, 117 (1992) 137496v.
- 157 Vampa, G., Benvenuti, S. and Rossi, T.: Determination of glycyrrhizin and 18 α - and 18 β -glycyrrhetic acid in rat plasma by high-performance thin-layer chromatography. *Farmaco*, 47 (1992) 825-832; *C.A.*, 117 (1992) 142784s.
- See also 24.

15. TERPENES AND OTHER VOLATILE AROMATIC COMPOUNDS

15b. *Essential oils*

- 158 Shin, S.H., Sim, Y., Kim, Y.S., Chi, H.J. and Lee, E.B.: (Studies on essential oils of *Lycopus lucidus* Turcz). *Saengyak Hak-hoechi*, 23, No. 1 (1992) 29-33; C.A., 117 (1992) 137497w.

16. NITRO AND NITROSO COMPOUNDS

See 287.

17. AMINES, AMIDES AND RELATED NITROGEN COMPOUNDS

17a. *Amines and polyamines*

- 159 Bazylak, G.: Retention behavior and molecular structure parameters of diethanolamine isomers in some reversed phase HPTLC systems. *J. Planar Chromatogr.*, 5 (1992) 275-279.
- 160 Bazylak, G.: Solute - solvent interactions of aliphatic aminoalcohols in underivatized silica - binary eluent HPTLC systems. *J. Planar Chromatogr.*, 5 (1992) 239-245.
- 161 Janjic, T.J., Milojkovic, D.M., Brocovic, A., Arbutina, Z.J. and Celap, M.B.: Thin-layer chromatography on polyacrylonitrile. V. Consideration of separation mechanisms of some aliphatic and aromatic amines. *J. Chromatogr.*, 609 (1992) 419-422.
- 162 Milhiet, P.-E., Dennin, F., Giocondi, M.-C., Le Grimellec, C., Garbay-Jaureguiberry, C., Boucheix, C. and Roques, B.P.: Detection of neutral endopeptidase-24.11/CD10 by flow cytometry and photomicroscopy using a new fluorescent inhibitor. *Anal. Biochem.*, 205 (1992) 57-64.
- 163 Ojanperä, I., Wähälä, K. and Hase, T.A.: Characterisation of amines by fast black K salt in thin-layer chromatography. *Analyst (London)*, 117 (1992) 1559-1565.
- 164 Singh, D.K. and Mishra, A.: Thin layer chromatography of aromatic amines on silica gel G impregnated with ammonium cerium(IV) nitrate. *J. Planar Chromatogr.*, 5 (1992) 284-286.
- 165 Voskoboynikov, V.V. and Gerasimenko, V.A.: (Method for determination of arylmono(di)isocyanates by thin-layer chromatography on silica gel). *U.S.S.R. SU 1,698,754* (Cl. G01N30/96), 15 Dec. 1991, Appl. 4,669,503, 28 Mar. 1989; C.A., 117 (1992) 61997m.

See also 12, 37, 297.

17c. *Urea and guanidine derivatives*

See 162.

17d. *Other amine derivatives and amides (excl. peptides)*

- 166 Boomgaarden, M., Heinrich, G. and Henklein, P.: Ein einfacher Zugang zu N-geschützten Aminosäureamiden. *Pharmazie*, 47 (1992) 710-711.

- 167 Perisic-Janjic, N.U., Petrovic, S.M., Djakovic, T., Nikolic, A.D. and Petrovic, S.D.: Retention behavior of some amides in reversed- and normal-phase thin-layer chromatography. *Chromatographia*, 34 (1992) 537-539.

See also 162.

18. AMINO ACIDS AND PEPTIDES; CHEMICAL STRUCTURE OF PROTEINS

18a. *Amino acids and their derivatives*

- 168 Caldentey, J. and Bamford, D.H.: The lytic enzyme of the *Pseudomonas* phage $\phi 6$. Purification and biochemical characterization. *Biochim. Biophys. Acta*, 1159 (1992) 44-50.
- 169 Jentsch, J.: The use of a new silica gel separation system in thin-layer chromatography. *Amino Acids*, 1 (1992) 279-281; C.A., 117 (1992) 86108k.
- 170 Kalinich, J.F. and McClain, D.E.: An *in vitro* method for radio-labeling proteins with ^{35}S . *Anal. Biochem.*, 205 (1992) 208-212.
- 171 Lepri, L., Coas, V. and Desideri, P.G.: Reversed phase planar chromatography of optically active fluorenylmethoxycarbonyl amino acids with bovine serum albumin in the mobile phase. *J. Planar Chromatogr.*, 5 (1992) 294-296.
- 172 Lepri, L., Coas, V., Desideri, P.G. and Zocchi, A.: Reversed phase planar chromatography of enantiomeric tryptophans with bovine serum albumin in the mobile phase. *J. Planar Chromatogr.*, 5 (1992) 234-238.
- 173 Lin, Y., Liang, D., Zhang, Y., Jin, S. and Huang, X.: Thin-layer chromatography-spectrophotometry for qualitative and quantitative analysis of bee-collected pollen. *Beijing Daxue Xuebao, Ziran Kexueban*, 117 (1991) 680-685; C.A., 117 (1992) 107523j.
- 174 Nakamura, F., Yamazaki, K. and Suyama, K.: Isolation and structural characterization of a new crosslinking amino acid, cyclopentenosine, from the acid hydrolysate of elastin. *Biochem. Biophys. Res. Commun.*, 186 (1992) 1533-1538.
- 175 Okada, S. and Ishimitsu, S.: (Studies on the application of amino acid chromatography for the purity test for amino acids in JP XI (1986)). *Iyakuin Kenkyu*, 23 (1992) 179-184; C.A., 117 (1992) 33793y.

See also 6, 166.

18b. *Peptides, peptidic and proteinous hormones, growth factors*

- 176 Dorizzi, R., Michelot, D., Tagliaro, F. and Ghielmi, S.: Methods for chromatographic determination of amanitins and related toxins in biological samples. *J. Chromatogr.*, 580 (1992) 279-291.
- 177 Hughes, K., Ramakrishna, S., Benjamin, W.B. and Woodgett, J.R.: Identification of multifunctional ATP-citrate lyase kinase as the α -isoform of glycogen synthase kinase-3. *Biochem. J.*, 288 (1992) 309-314.
- 178 Kuba, M., Ohmori, H. and Kumon, A.: Characterization of N^{ω} -phosphoarginine hydrolase from rat liver. *Eur. J. Biochem.*, 208 (1992) 747-752.
- 179 Nakao, M., Mizutani, T., Bhakta, A., Ribarac-Stepic, N. and Moudgil, V.K.: Phosphorylation of chicken oviduct progesterone receptor by cAMP-dependent protein kinase. *Arch. Biochem. Biophys.*, 298 (1992) 340-348.

180 Wang, C., Lin, J.-M. and Lazarides, E.: Methylations of 70000-Da heat shock proteins in 3T3 cells: alterations by arsenite treatment, by different stages of growth and by virus transformation. *Arch. Biochem. Biophys.*, 297 (1992) 169-175.

See also 22, 221.

19. PROTEINS

19a. General techniques

See 181.

19l. Specific binding and receptor proteins

181 Birnbaum, S., Uden, C., Magnusson, C.G.M. and Nilsson, S.: Latex-based thin-layer immunoaffinity chromatography for quantitation of protein analytes. *Anal. Biochem.*, 206 (1992) 168-171.

20. ENZYMES AND ENZYME ACTIVITY ESTIMATION

20d. Hydrolases, acting on ester bonds (E.C. 3.1.-.-)

182 Griban, G.A. and Minyaev, M.V.: (Estimation of cholinesterase activity using microthin-layer chromatography). *Vopr. Med. Khim.*, 38 (1992) 59-61; C.A., 117 (1992) 186997.

21. PURINES, PYRIMIDINES, NUCLEIC ACIDS AND THEIR CONSTITUENTS

21a. Purines, pyrimidines, nucleosides, nucleotides

183 Belikov, V.G., Lovtseva, E.A. and Saushkina, A.S.: (Optimization of conditions for adenosine chromatography by the experiment design method). *Farmatsiya (Moscow)*, 41 (1992) 31-34; C.A., 117 (1992) 166918q.

184 Dalton, A., Hornby, D.P., Langston, S.P. and Blackburn, G.M.: Characterization and purification of a novel dATP-binding protein in eukaryotes. *Biochem. J.*, 287 (1992) 871-879.

185 Ji, C. and Marnett, L.J.: Oxygen radical-dependent epoxidation of (7S,8S)-dihydroxy-7,8-dihydrobenzo[a]pyrene in mouse skin *in vivo*. Stimulation by phorbol esters and inhibition by anti-inflammatory steroids. *J. Biol. Chem.*, 267 (1992) 17842-17848.

186 Mullis, K.G. and Ketcham, C.M.: The synthesis of substrates and two assays for the detection of N-acetylglucosamine-1-phosphodiester α -N-acetylglucosaminidase (uncovering enzyme). *Anal. Biochem.*, 205 (1992) 200-207.

187 Nakamura, A., Koide, Y., Miyazaki, H., Kitamura, A., Masaki, H., Beppu, T. and Uozumi, T.: Gene cloning and characterization of a novel extracellular ribonuclease of *Bacillus subtilis*. *Eur. J. Biochem.*, 209 (1992) 121-127.

188 Nakamura, H., Natsumeda, Y., Nagai, M., Shiotani, T. and Weber, G.: Direct assay method for guanosine 5'-monophosphate reductase activity. *Anal. Biochem.*, 206 (1992) 115-118.

189 Talaska, G., Roh, J.H. and Getek, T.: ³²P-Postlabelling and mass spectrometric methods for analysis of bulky, polyaromatic carcinogen-DNA adducts in humans. *J. Chromatogr.*, 580 (1992) 293-323.

See also 9, 109, 136.

22. ALKALOIDS

190 Agbaba, D., Zivanov-Stakic, D. and Vukicevic, N.: Dissolution assay of theophylline, diprophylline and proxiphylline from a sustained-release dosage form by high-performance thin-layer chromatography. *Biomed. Chromatogr.*, 6 (1992) 141-142; C.A., 117 (1992) 33574c.

191 Beikin, S.G., Ruskikh, S.L. and Gaponenko, Ya.S.: (Identifying reaction of gindarine). *Farm. Zh. (Kiev)*, No. 3 (1992) 74-76; C.A., 117 (1992) 118578g.

192 Ensing, J.G. and de Zeeuw, R.A.: Detection, isolation, and identification of truxillines in illicit cocaine by means of thin-layer chromatography and mass spectrometry. *J. Forensic Sci.*, 36 (1991) 1299-1311; C.A., 117 (1992) 144919p.

193 Moeller, R.M.: Drug detection in hair by chromatographic procedures. *J. Chromatogr.*, 580 (1992) 125-134.

194 Monforte-Gonzalez, M., Ayora-Talavera, T., Maldonado-Mendoza, I.E. and Loyola-Vargas, V.M.: Quantitative analysis of serpentine and ajmalicine in plant tissues of *Catharanthus roseus* and hyoscyamine and scopolamine in root tissues of *Datura stramonium* by thin layer chromatography-densitometry. *Phytochem. Anal.*, 3 (1992) 117-121; C.A., 117 (1992) 157734z.

195 Nelson, C.C. and Foltz, R.L.: Chromatographic and mass spectrometric methods for determination of lysergic acid diethylamide (LSD) and metabolites in body fluids. *J. Chromatogr.*, 580 (1992) 97-109.

196 Nikolova-Damyanova, B., Tam, M.N. and Pyuskyulev, B.: Quantitative TLC of indole alkaloids. *J. Planar Chromatogr.*, 5 (1992) 271-274.

197 Obreshkova, D. and Naidenova, E.: (Quantitative determination of vincamine in Vincapan tablets). *Probl. Farmakol. Farm.*, 4 (1990) 68-71; C.A., 117 (1992) 76578z.

198 Singh, A.K., Granley, K., Misra, U., Naeem, K., White, T. and Jiang, Y.: Screening and confirmation of drugs in urine: interference of hordenine with the immunoassays and thin layer chromatography methods. *Forensic Sci. Int.*, 54 (1992) 9-22; C.A., 117 (1992) 126011q.

199 Wang, W., Wang, Y., Zhang, T. and Su, Y.: (Quantitative analysis of ephedrine in Tangxuanligi oral liquid by TLC densitometry). *Zhongcaoyao*, 23 (1992) 245-246; C.A., 117 (1992) 137749e.

200 Zhang, A., Qi, L., Liang, L. and Wang, X.: (Determination of quinidine sulfate concentration in plasma by thin-layer chromatographic and dual wavelength densitometry). *Xi'an Yike Daxue Xuebao*, 12 (1991) 84-86; C.A., 117 (1992) 103458u.

See also 294.

23. OTHER SUBSTANCES CONTAINING HETEROCYCLIC NITROGEN

23a. Porphyrins and other pyrroles

- 201 Saitoh, K., Kashiwa, R., Tada, R., Kiyohara, C. and Suzuki, N.: High-performance thin-layer chromatographic mobility of porphyrins and their copper complexes on different adsorbents. *J. Chromatogr.*, 609 (1992) 341-347.
- 202 Sliwiok, J., Kus, P. and Kozera, A.: Chromatographic characteristics of porphyrine ether glycosides. *Chromatographia*, 34 (1992) 431-432.

23d. Pyridine derivatives

- 203 Baranowska, I. and Kubik, T.: Comparison of the chromatographic behavior of diazines and their derivatives on silica gel thin layers. *J. Planar Chromatogr.*, 4 (1991) 467-470.

See also 174.

23e. Other N-heterocyclic compounds

- 204 Özden, S., Özden, T., Attila, I., Küçükislamoglu, M. and Okatan, A.: Isolation and identification via high-performance liquid chromatography and thin-layer chromatography of benzoxazolinone precursors from *Consolida orientalis* flowers. *J. Chromatogr.*, 609 (1992) 402-406.
- 205 Shimada, K. and Mizuguchi, T.: Sensitive and stable Cookson-type reagent for derivatization of conjugated dienes for high-performance liquid chromatography with fluorescence detection. *J. Chromatogr.*, 606 (1992) 133-135.

See also 203, 281.

24. ORGANIC SULPHUR COMPOUNDS (INCL. GLUCOSINOLATES)

- 206 Xie, B., Zhou, C., Song, R. and Zhang, R.: (Detection of sodium cyclamate in soft drinks). *Zhongguo Tiaoweipin*, (1992) 24-25; C.A., 117 (1992) 149577s.
- 207 Xun, L. and Xu, J.: (Study on the photochemical-fluorometric method. IV. Determination of sulfadiazine). *Fenxi Huaxue*, 20 (1992) 499-503; C.A., 117 (1992) 33850q.

See also 259, 278.

25. ORGANIC PHOSPHORUS COMPOUNDS (INCL. SUGAR PHOSPHATES)

See 32, 177, 179.

26. ORGANOMETALLIC AND RELATED COMPOUNDS

26a. Organometallic compounds

- 208 Vuckovic, G., Miljevic, D., Janjic, T.J. and Celap, M.B.: Salting-out thin-layer chromatography of transition metal complexes. I. Investigation of the behaviour of mixed aminocarboxylato cobalt(III) complexes on silica gel. *J. Chromatogr.*, 609 (1992) 427-431.

27. VITAMINS AND VARIOUS ANIMAL GROWTH FACTORS (NON-PEPTIDIC)

- 209 Chavan, J.D. and Khatri, J.M.: Simultaneous determination of retinol and α -tocopherol in human plasma by HPTLC. *J. Planar Chromatogr.*, 5 (1992) 280-282.
- 210 Igo'nikova, O.P., Akhmedkhodzhaeva, N.M. and Svechnikova, A.N.: (Chromatographic analysis of drug mixtures containing folic acid). *Farm. Zh. (Kiev)*, No. 3 (1992) 92-93; C.A., 117 (1992) 118604n.
- 211 Rizzolo, A. and Polesello, S.: Chromatographic determination of vitamins in foods. *J. Chromatogr.*, 624 (1992) 103-152 - a review with 286 refs.

See also 252.

28. ANTIBIOTICS

- 212 Aoyama, Y., Katayama, T., Yamamoto, M., Tanaka, H. and Kon, K.: A new antitumor antibiotic product, demethylchartreusin. Isolation and biological activities. *J. Antibiot.*, 45 (1992) 875-878.
- 213 Bannister, B. and Zapotocky, B.A.: Protobruidiridin, an antibiotic containing a C-nitro-sugar fragment, is the true secondary metabolite produced by *Streptomyces achromogenes* var. *rubradiridis*. Rubradiridin, described earlier, is its photooxidation product. *J. Antibiot.*, 45 (1992) 1313-1324.
- 214 Barker, S.A. and Walker, C.C.: Chromatographic methods for tetracycline analysis in foods. *J. Chromatogr.*, 624 (1992) 195-209 - a review with 81 refs.
- 215 Bobbitt, D.R. and Ng, K.W.: Chromatographic analysis of antibiotic materials in food. *J. Chromatogr.*, 624 (1992) 153-170 - a review with 53 refs.
- 216 Boeck, L.V.D., Berry, D.M., Mertz, F.P. and Wetzel, R.W.: A10255, a complex of novel growth-promoting thiopeptide antibiotics produced by a strain of *Streptomyces gardneri*. Taxonomy and fermentation studies. *J. Antibiot.*, 45 (1992) 1222-1230.
- 217 Boison, J.O.: Chromatographic methods of analysis for penicillins in food-animal tissues and their significance in regulatory programs for residue reduction and avoidance. *J. Chromatogr.*, 624 (1992) 171-194 - a review with 127 refs.
- 218 Dirlam, J.P., Bordner, J., Cullen, W.P., Jefferson, M.T. and Press-eau-Linabury, L.: The structure of CP-96,797, a polyether antibiotic related to K-41A and produced by *Streptomyces* sp. *J. Antibiot.*, 45 (1992) 1187-1189.

- 219 Fomicheva, E.V., Fedorova, G.B., Potapova, N.P. and Katrukha, G.S.: Isolation and identification of new anthracycline antibiotics, rubomycins F and H. *J. Antibiot.*, 45 (1992) 1185-1186.
- 220 Grabley, S., Hammann, P., Hütter, K., Kirsch, R., Kluge, H. and Thiericke, R.: Secondary metabolites by chemical screening. 20. Decaerestrictines, a new family of inhibitors of cholesterol biosynthesis from *Penicillium*: III. Decaerestrictines E to M. *J. Antibiot.*, 45 (1992) 1176-1181.
- 221 Hatsu, M., Tuda, M., Muraoka, Y., Aoyagi, T. and Takeuchi, T.: Benarthin: a new inhibitor of pyroglutamyl peptidase. II. Synthesis and structure-activity relationship. *J. Antibiot.*, 45 (1992) 1088-1095.
- 222 Hedtmann, U., Fehlhaber, H.-W., Sukatsch, D.A., Weber, M., Hoffmann, D. and Kraemer, H.P.: The new cytotoxic antibiotic cytorhodin X, an unusual anthracyclinone-9 α -glycoside. *J. Antibiot.*, 45 (1992) 1373-1375.
- 223 Hoff, H., Drautz, H., Fiedler, H.-P., Zähler, H., Schultz, J.E., Keller-Schierlein, W., Philipps, S., Ritzau, M. and Zeeck, A.: Metabolic products of microorganisms. 261. Obscurolidides, a novel class of phosphodiesterase inhibitors from *Streptomyces*. I. Production, isolation, structural elucidation and biological activity of obscurolidides A₁ to A₄. *J. Antibiot.*, 45 (1992) 1096-1108.
- 224 Itoh, J. and Miyadoh, S.: SF2457, A new antibiotic related to amicitin. *J. Antibiot.*, 45 (1992) 846-853.
- 225 Kádár-Pauncz, J., Podányi, B. and Horváth, G.: Isolation and structure elucidation of new antibiotics related to angolamycin. *J. Antibiot.*, 45 (1992) 1231-1238.
- 226 Kovacs-Haddady, K.: OPLC study of the retention behavior of several tetracyclines on thin layers impregnated with tricaprilmethylammonium chloride. *J. Planar Chromatogr.*, 4 (1991) 456-459.
- 227 Morishita, A., Murofushi, S., Ishizawa, K., Mutoh, N. and Yaginuma, S.: Isolation and characterization of sporeamicin C. *J. Antibiot.*, 45 (1992) 1011-1015.
- 228 Novakova, E.: (Detection of antibiotics in biological material in toxicological analysis. II. Detection of tetracyclines in biological material). *Cesk. Farm.*, 41 (1992) 105-107; *C.A.*, 117 (1992) 142785t.
- 229 Ogawara, H., Higashi, K., Manita, S., Tanaka, K., Shimizu, Y. and Shufang, L.: An inhibitor for inositol-specific phospholipase C from *Actinoadura* sp. *J. Antibiot.*, 45 (1992) 1365-1366.
- 230 Ohkuma, H., Naruse, N., Nishiyama, Y., Tsuno, T., Hoshino, Y., Sawada, Y., Konishi, M. and Oki, T.: Sultricin, a new antifungal and antitumor antibiotic from *Streptomyces roseiscleroticus*. Production, isolation, structure and biological activity. *J. Antibiot.*, 45 (1992) 1239-1249.
- 231 Parry, R.J. and Yang, N.: Isolation and characterization of furanomycin nonproducing *Streptomyces threomyceticus* mutants. *J. Antibiot.*, 45 (1992) 1161-1166.
- 232 Proksa, B., Uhrin, D., Adamcová, J. and Fuska, J.: Vermixocins A and B, two novel metabolites from *Penicillium vermiculatum* *J. Antibiot.*, 45 (1992) 1268-1272.
- 233 Shoji, J., Hino, H., Katayama, T., Nakagawa, J., Ikenishi, Y., Iwatani, K. and Yoshida, T.: Structures of new peptide antibiotics, plusbacins A₁~A₄ and B₁~B₄. *J. Antibiot.*, 45 (1992) 824-831.
- 234 Tabata, Y., Imoto, M. and Umezawa, K.: Biosynthetic preparation of radioactively labeled erbstatin. *J. Antibiot.*, 45 (1992) 1382-1384.
- 235 Tolstikov, V.V., Preobrazhenskaya, M.N., Balzarini, J. and de Clercq, E.: Chemical modification of antibiotic streptonigrin: synthesis and properties of 2'-decarboxy-2'-aminostreptonigrin (streptonigrone-2'-imine). *J. Antibiot.*, 45 (1992) 1002-1004.
- 236 Ueda, Y. and Vinet, V.: Synthesis and *in vitro* activity of novel quaternary ammonium carbapenems: 2-pyridiniopropyl and 1-pyridinioethyl carbapenems. *J. Antibiot.*, 45 (1992) 940-953.
- 237 Yoshimoto, A., Johdo, O., Fujii, S., Kubo, K., Nishida, H., Okamoto, R. and Takeuchi, T.: Anthracycline metabolites from baumycin-producing *Streptomyces* sp. D788. I. Isolation of antibiotic-blocked mutants and their characterization. *J. Antibiot.*, 45 (1992) 1255-1267.
29. INSECTICIDES, PESTICIDES AND OTHER AGROCHEMICALS
- 29a. *General techniques*
- 238 Erdmann, F., Schuetz, H., Brose, C. and Rochholz, G.: An optimized screening system for 170 pesticides. *Beitr. Gerichtl. Med.*, (1991) 121-126; *C.A.*, 117 (1992) 165282r - a review with 3 refs.
- 239 Fodor-Csorba, K.: Chromatographic methods for the determination of pesticides in foods. *J. Chromatogr.*, 624 (1992) 353-367 - a review with 62 refs.
- 240 Jork, H., Keller, G. and Kocher, U.: Application of AMD to the determination of crop-protection agents in drinking water. Part II: Limitations. *J. Planar Chromatogr.*, 5 (1992) 246-250.
- 29d. *Carbamates*
- 241 Rathore, H.S., Khan, H.A. and Sharma, R.: Thin layer chromatographic separation, detection, and quantitation of carbaryl in water. *J. Planar Chromatogr.*, 4 (1991) 494-496.
- 29e. *Herbicides*
- 242 Rathore, H.S. and Begum, T.: Chromatographic behavior of some carboxylic acid herbicides and plant growth regulators on impregnated papers. *J. Planar Chromatogr.*, 4 (1991) 451-455.
- 29g. *Other types of pesticides and various agrochemicals*
- 243 Gbewonyo, W.S.K. and Candy, D.J.: Chromatographic isolation of insecticidal amides from Piper guineense root. *J. Chromatogr.*, 607 (1992) 105-111.
- 244 Gevorgyan, A.A. and Bunyatyan, Yu.A.: (Determination of resin by thin-layer chromatography in air, water, beet tops, and root-crops). *Gig. Sanit.*, (1992) 61; *C.A.*, 117 (1992) 186562x.
- 245 Katagi, T.: Photodegradation of 3-phenoxybenzoic acid in water and on solid surfaces. *J. Agric. Food Chem.*, 40 (1992) 1269-1274.

30. SYNTHETIC AND NATURAL DYES

- 246 Politzer, I.R., Crago, K.T., Amos, K., Mitchell, K. and Hollin, T.: Thin-layer chromatography of laser dyes and dye analogs with cyclodextrine in the mobile phase. *Talanta*, 39 (1992) 953-958; C.A., 117 (1992) 132896r.

30a. Synthetic dyes

- 247 Bide, M.J. and Choi, H.: The thin layer chromatography of vat dyes. *J. Soc. Dyers Colour.*, 108 (1992) 133-138; C.A., 117 (1992) 92236s.
- 248 Fanali, S. and Schudel, M.: Some separation of black and red water-soluble fiber-tip pen inks by capillary zone electrophoresis and thin-layer chromatography. *J. Forensic Sci.*, 36 (1991) 1192-1197; C.A., 117 (1992) 144917m.
- 249 Tralic-Kulenovic, V., Mrsa, R. and Fischer-Jakic, L.: Densitometric examination of dye chromatograms. *Melliand Textilber.*, 73 (1992) E236, 512-514; C.A., 117 (1992) 132806m.

30b. Chloroplast and other natural pigments

- 250 Iborra, J.L., Castellar, M.R., Canovas, M. and Manjon, A.: TLC preparation purification of picrocrocine, HTCC and crocine from saffron. *J. Food Sci.*, 57 (1992) 714-716, 731; C.A., 117 (1992) 88999a.
- 251 Li, Z., Wu, J. and Tang, J.: (Densitometric determination of lycopene in tomato paste with thin-layer chromatography). *Sepu*, 10 (1992) 184-185; C.A., 117 (1992) 88883h.
- 252 Mercadante, A.Z. and Rodriguez-Arroyave, D.B.: Screening of carotenoids: comparison of thin-layer chromatography with high-efficiency thin-layer chromatography with multiple development. *Cienc. Technol. Aliment.*, 11 (1992) 200-209; C.A., 117 (1992) 190388n.
- 253 Watanabe, T. and Kobayashi, M.: (Separation of photosynthetic pigments). *Gendai Kagaku*, (1992) 384-390; C.A., 117 (1992) 166819h - a review with 3 refs.

See also 295.

31. PLASTICS AND THEIR INTERMEDIATES

- 254 Belay, M.T. and Poole, C.F.: Determination of molecular weight and oligomer distribution of poly(ethylene glycols) by thin layer chromatography and automated multiple development. *J. Planar Chromatogr.*, 4 (1991) 424-431.
- 255 Belen'kii, B.G., Gankina, E.S., Zgonnik, V.N., Malkhova, I.L. and Melenevskaya, E.Yu.: (Thin-layer chromatography of block copolymers: mechanism of separation, determination of polydispersity). *Vysokomol. Soedin., Ser. A*, 33 (1991) 2487-2493; C.A., 117 (1992) 131879a.
- 256 Belenkii, B.G., Gankina, E.S., Zgonnik, V.N., Malchova, I.I. and Melenevskaya, E.U.: Critical thin-layer chromatography as a novel method for the determination of the molecular weight and compositional inhomogeneity of block copolymers. *J. Chromatogr.*, 609 (1992) 355-362.

- 257 Petrovic, S.M. and Kolarov, L.A.: (Preparation and application of the aminoplast resin in thin layer chromatography). *Zb. Matice Srp. Priir. Nauke*, 78 (1990) 83-90; C.A., 117 (1992) 16945t.

32. DRUG ANALYSIS

32a. Drug analysis, general techniques

- 258 Ahuja, S.: Chromatographic solutions to pharmaceutical analytical problems. *Chromatographia*, 34 (1992) 411-416.
- 259 Ban, Y., Sun, Y. and Wang, Y.: (Optimization of the thin-layer chromatography solvent system for sulfa drugs). *Sepu*, 10 (1992) 103-105; C.A., 117 (1992) 118585g.
- 260 De Zeeuw, R.A., Franke, J.P., Dik, E., Ten, Dolle, D.W. and Kam, B.L.: Impact of tropical conditions on thin layer chromatography in analytical toxicology: high temperatures and moderate humidities. *J. Forensic Sci.*, 37 (1992) 984-990; C.A., 117 (1992) 165405h.
- 261 Gough, T.A. (Editor): *The Analysis of Drugs of Abuse*. Separation Science Series, Wiley, Chichester & New York, 1991, XVIII + 628 p.
- 262 Wilson, I.D.: Recent advances in thin-layer chromatography: application to drug metabolism studies. *J. Biopharm. Sci.*, 2 (1991) 173-189; C.A., 117 (1992) 82755w - a review with 34 refs.

See also 193, 310.

32b. Antirheumatics and antiinflammatory drugs

- 263 Ali, A. and Sharma, S.N.: Preparation and evaluation of solid dispersions of ibuprofen. *Indian J. Pharm. Sci.*, 53 (1991) 233-236; C.A., 117 (1992) 76345w.
- 264 Hickman, R.J.S.: Urinary excretion of paracetamol: a thin-layer chromatography exercise for nursing students. *J. Chem. Educ.*, 69 (1992) 510-511; C.A., 117 (1992) 89569x.
- 265 Kersten, B.S., Catalano, T. and Lucarelli, M.: TLC analysis of 5-aminosalicylic acid bulk chemical. *J. Planar Chromatogr.*, 4 (1991) 483-484.
- 266 Martin, P., Morden, W., Wall, P. and Wilson, I.: TLC combined with tandem mass spectrometry: application to the analysis of antipyrine and its metabolites in extracts of human urine. *J. Planar Chromatogr.*, 5 (1992) 255-258.
- 267 Sane, R.T., Tirodkar, V.B., Desai, A.J., Patel, M.K. and Kulkarni, U.D.: Application of different analytical methods for the determination of ketorolac tromethamine. *Indian Drugs*, 29 (1992) 489-493; C.A., 117 (1992) 157759m.

32c. Autonomic and cardiovascular drugs

- 268 Ling, Q. and Guo, Z.: (HPTLC determination of meriletine in plasma). *Yaowu Fenxi Zazhi*, 12 (1992) 211-213; C.A., 117 (1992) 184183a.
- 269 Marciniak, B., Kujawa, E. and Ogródowczyk, M.: Evaluation of nifedipine preparations by chromatographic-spectrophotometric methods. *Pharmazie*, 47 (1992) 502-504.

- 270 Turgeon, J., Pare, J.R.J., Lalonde, M., Grach-Belanger, O. and Belanger, P.M.: Isolation and structural characterization by spectroscopic methods of two glucuronide metabolites of mexiletine after N-oxidation and deamination. *Drug Metab. Disp.*, 20 (1992) 762-769.
- 271 Zivanov-Stakic, D., Agbaba, D., Vladimirov, S. and Ciric, L.J.: Spectrophotometric determination of diltiazem in dosage forms. *Farmaco*, 47 (1992) 393-396; *C.A.*, 117 (1992) 33833m.
- See also 220.
- 32d. Central nervous system drugs**
- 272 Cai, X., Chen, R. and Yi, D.: (Structure elucidation of impurities in crude carbamazepine). *Zhongguo Yiyao Gongye Zazhi*, 23 (1992) 78-80; *C.A.*, 117 (1992) 55789d.
- 273 Göber, B., Franke, P. and Lisowski, H.: Zur Stabilität von Oxetacain und zur Struktur seiner Abbauprodukte. *Pharmazie*, 47 (1992) 584-587.
- 274 Przyborowski, L., Zawisza, P. and Hopkala, H.: (Thin-layer chromatographic detection of glutethimide and methyprylone in blood). *Acta Pol. Pharm.*, 47 (1990) 21-24; *C.A.*, 117 (1992) 142777s.
- 275 Roca, R.M.S., March, F.J.G., Fos, G.M.A., Domenech, R.G., Gimenez, F.P. and Alvarez, J.G.: Molecular topology and chromatographic retention parameters for benzodiazepines. *J. Chromatogr.*, 607 (1992) 91-95.
- See also 14, 207, 258, 267.
- 32e. Chemotherapeutics (exc. cytostatics and antibiotics)**
- 276 Gruszecka-Kowalik, E., Haugwitz, R.D. and Zalkow, L.H.: Quinobene, a new potent anti-HIV agent. *Biochem. Biophys. Res. Commun.*, 187 (1992) 1409-1417.
- 277 Guggisberg, D., Mooser, A.E. and Koch, H.: Methods for the determination of sulphonamides in meat. *J. Chromatogr.*, 624 (1992) 425-437 - a review with 89 refs.
- 278 Hermann, T., Grzeskowiak, E. and Gruszka, I.: (Identification of some sulfadiazamide impurities and its determination). *Farm. Pol.*, 48 (1992) 189-195; *C.A.*, 117 (1992) 33834n.
- 279 Petrovic, S.M., Acanski, M., Perisic-Janjic, N.U. and Vlaovic, D.: Separation and retention behavior of some newly synthesized benzimidazoles in normal and reversed phase thin layer chromatography. *J. Planar Chromatogr.*, 4 (1991) 475-479.
- 280 Salomies, H.: Determination of tinidazole in dosage forms by quantitative HPTLC. *J. Planar Chromatogr.*, 5 (1992) 291-293.
- See also 157, 200, 207.
- 32f. Cytostatics**
- See 9, 197.
- 32g. Other drug categories**
- 281 De Weerd, G. and van der Meer, Y.G.: (Formulation and shelf-life of a co-trimoxazole intravenous infusion). *Ziekenhuisfarmacie*, 8, No. 1 (1992) 1-8; *C.A.*, 117 (1992) 55803d.
- 282 Hotho, M.R., Brers, O., Cragoe, E.J., Jr., Veel, T. and Raeder, M.: Determination of active 5-(N,N-hexamethylene)amiloride in pig plasma. *J. Chromatogr.*, 579 (1992) 376-381.
- 283 Kerc, J., Srcic, S., Urleb, U., Kanalec, A., Kofler, B. and Smid-Korbar, J.: Compatibility study between acetylcysteine and some commonly used tablet excipients. *J. Pharm. Pharmacol.*, 44 (1992) 515-518; *C.A.*, 117 (1992) 137538k.
- 284 Mangalan, S., Patel, R.B. and Chakravarthy, B.K.: Detection and determination of omeprazole by HPTLC: a reliable method for the estimation of plasma levels. *J. Planar Chromatogr.*, 4 (1991) 492-493.
- 285 Obreshkova, D. and Naidenova, E.: (Qualitative and quantitative analysis of Bellonal tablets). *Probl. Farmakol. Farm.*, 4, No. 7 (1990) 65-68; *C.A.*, 117 (1992) 157728a.
- 286 Obreshkova, L., Naidenova, E., Pavlova, I. and Kmetska, G.: (Qualitative and quantitative analysis of Tribestan film tablets). *Probl. Farmakol. Farm.*, 4, No. 7 (1990) 71-73; *C.A.*, 117 (1992) 118573b.
- 287 Shahjahan, M. and Enever, R.P.: A stability-indicating assay for nitrofurazone by paper chromatography. *Int. J. Pharm.*, 82 (1992) 215-221; *C.A.*, 117 (1992) 97431m.
- See also 16.
- 32h. Toxicological and forensic applications**
- 288 Kok, T.M.C.M., Levels, P.J., van Faassen, A., Hazen, M., ten Hoor, F. and Kleinjans, J.C.S.: Chromatographic methods for the determination of toxicants in faeces. *J. Chromatogr.*, 580 (1992) 135-159.
- 289 Linhart, I., Vosmanská, M., Mitera, J. and Šmejkal, J.: Biotransformation of diethenylbenzenes. IV. A simple high-performance liquid chromatographic method for separation of urinary metabolites of 1,3-diethenylbenzene on analytical and semi-preparative scales. *J. Chromatogr.*, 578 (1992) 215-221.
- See also 2, 176, 189, 195, 198, 260, 261.
- 32i. Plant extracts**
- 290 Kitanov, G.M.: Phenolic acids and flavonoids from *Stellaria media* (L.) Vill. (*Caryophyllaceae*). *Pharmazie*, 47 (1992) 470-471.
- 291 Mahran, G.H., El-Hosary, G., Seida, A.A. and Amin, W.: A study of the volatile oil content of *Ambrosia maritima* L. cultivated in Egypt. *Bull. Fac. Pharm. (Cairo Univ.)*, 28 (1990) 53-55; *C.A.*, 117 (1992) 76303f.
- 292 Obreshkova, L., Naidenová, E. and Pavlova, I.: (Qualitative and quantitative analysis of some drugs of plant origin in Neutracid tablets). *Probl. Farmakol. Farm.*, 4, No. 7 (1990) 62-65; *C.A.*, 117 (1992) 137744z.
- 293 Sanz, M.J., Terencio, M.C. and Paya, M.: Isolation and hypotensive activity of a polymeric procyanidin fraction from *Pistacia lentiscus* L. *Pharmazie*, 47 (1992) 466-467.
- 294 Zhu, D., Jiang, Y., Chen, T. and Gao, S.: (Research on chemical components and primary pharmaceutical actions of cultured bulb of *Fritillaria unibracteata*). *Zhongguo Yaokexue Xuebao*, 23 (1992) 118-121; *C.A.*, 117 (1992) 137498x.
- See also 13, 34, 39, 52, 155, 156, 194, 204, 243, 250.

33. CLINICO-CHEMICAL APPLICATIONS

33b. *Complex mixtures and profiling (single compounds by cross-reference only)*

See 67, 198, 200, 228.

34. FOOD ANALYSIS

34a. *General papers and reviews*

See 20, 42, 115, 146, 211, 214, 215, 217, 239, 277.

34b. *Complex mixtures (single compounds by cross-reference only)*

295 Sujata, V., Ravishankar, G.A. and Venkataraman, L.V.: Methods for the analysis of the saffron metabolites crocin, crocetin, picrocrocin and safranal for the determination of the quality of the spice using thin-layer chromatography, high-performance liquid chromatography and gas chromatography. *J. Chromatogr.*, 624 (1992) 497-502.

See also 56, 69, 206, 250, 251, 298, 300, 301.

34c. *Organoleptically important compounds (flavors, odors, volatiles)*

See 295.

35. ENVIRONMENTAL ANALYSIS

35b. *Air pollution (complex mixtures; single compounds by cross-reference only)*

See 244.

35c. *Water pollution (complex mixtures; single compounds by cross-reference only)*

296 Hühnerfuss, H. and Kallenborn, R.: Chromatographic separation of marine organic pollutants. *J. Chromatogr.*, 580 (1992) 191-214.

297 Stradomskaya, A.G., Tarasova, K.P. and Gavrilenko, N.I.: (Extraction-chromatographic method for determination of aniline in natural waters). *Gidrokhim. Mater.*, 110 (1991) 117-122; *C.A.*, 117 (1992) 55502e.

See also 240, 241, 244.

36. SOME TECHNICAL PRODUCTS AND COMPLEX MIXTURES

36a. *Surfactants*

298 Johansson, D. and Bergenstahl, B.: A influence of food emulsifiers on fat and sugar dispersions in oils. I. Adsorption, sedimentation. *J. Am. Oil Chem. Soc.*, 69 (1992) 705-717.

299 Szögyi, M. and Cserhati, T.: Determination of the lipophilicity of some non-ionic surfactants: the effect of organic modifiers. *J. Planar Chromatogr.*, 5 (1992) 267-270.

36b. *Antioxidants and preservatives*

300 Duh, P.D., Yeh, D.B. and Yen, G.C.: Extraction and identification of an antioxidative component from peanut hulls. *J. Am. Oil Chem. Soc.*, 69 (1992) 814-818.

301 Komaitis, M.E.: Thin-layer chromatographic detection of some common antioxidants. *Grasas Aceites (Seville)*, 42 (1991) 420-421; *C.A.*, 117 (1992) 88846y.

See also 52.

36c. *Complex mixtures, technical products and unidentified compounds*

See 66, 165, 221, 248, 257.

38. INORGANIC COMPOUNDS

38a. *Cations*

302 Panesar, K.S., Singh, O.V. and Tandon, S.N.: Reversed phase TLC behavior of 3d metal ions in high molecular weight amine-succinic acid systems. *J. Planar Chromatogr.*, 5 (1992) 289-291.

303 Ray, R.K. and Kauffman, G.B.: Chromatographic studies of metal complexes. Part-III. Adsorption, dissociation, and intermediate complex formation of some square planar biguanide complexes of copper(II), nickel(II) and palladium(II) on silica gel. *J. Indian Chem. Soc.*, 68 (1991) 291-293; *C.A.*, 117 (1992) 34313d.

304 Sharma, S.D. and Misra, S.: Effect of solvent composition and pH and R_f values of metal ions on titanium tungstate-impregnated papers in aqueous nitric acid, acetone-nitric acid and butanol-nitric acid systems. *J. Chromatogr.*, 606 (1992) 121-128 - 50 cations in 31 solvent systems.

305 Shimizu, T., Igarashi, R., Hayashi, Y. and Jindo, S.: TLC of inorganic ions on *p*-aminobenzyl cellulose in both sulfuric acid and acid - ammonium sulfate media. *J. Planar Chromatogr.*, 4 (1991) 487-489.

306 Zuanon, N.J.: (Separation and identification of cations by thin layer chromatography). *Eclética Quim.*, 16 (1991) 9-13; *C.A.*, 117 (1992) 163017j.

See also 208, 311.

38b. *Anions*

307 Mohammad, A. and Tiwari, S.: Effect of heavy metals on the chromatographic separation of periodate from iodate, bromate, iodide, molybdate, and ferrocyanide. *J. Planar Chromatogr.*, 4 (1991) 485-487.

See also 248.

39. RADIOACTIVE AND OTHER ISOTOPE COMPOUNDS

- 308 Billingham, M.W., Abrams, D.N. and Dupont, J.: A comparison of radiopharmaceutical labeling efficiency of chromatographic generator vs. MEK extraction [^{99m}Tc] pertechnetate. *Appl. Radiat. Isot.*, 43 (1992) 1045-1049; *C.A.*, 117 (1992) 137649x.
- 309 Ding, H.J. and Yeh, S.J.: Stability of dilute aqueous solutions of technetium-99-HMPAO diastereoisomers. *Appl. Radiat. Isot.*, 43 (1992) 1013-1017; *C.A.*, 117 (1992) 137648w.
- 310 Khachadorian, S.K., Shafiq, Y.F. and Al-Janabi, M.A.A.: Comparative study of different chromatographic methods for quality control of radiopharmaceuticals labeled with metastable indium-113. *J. Radioanal. Nucl. Chem.*, 159 (1992) 3-8; *C.A.*, 117 (1992) 76619p.
- 311 Koehler, B. and Taylor, D.M.: A rapid radiochromatographic test for metal-ligand complex formation. *Appl. Radiat. Isot.*, 43 (1992) 935-936; *C.A.*, 117 (1992) 127391t.
- 312 Nakayama, M., Saigo, H., Kai, E., Koda, A., Ozeki, H., Harada, K., Sugii, A., Tomiguchi, S., Kojima, A. et al.: Hydroxamamide as a chelating moiety for the preparation of technetium-99m radiopharmaceuticals (I). *Nucl. Med. Commun.*, 13 (1992) 445-449; *C.A.*, 117 (1992) 157607k.

See also 189.

Electrophoresis

1. REVIEWS AND BOOKS

- 1 Anonymous: (*The Whole of Electrophoresis*). Ishiyaku Publishers, Tokyo, 1992, 254 p.; C.A., 117 (1992) 98403r.
- 2 Burge, G.L.: Capillary electrophoresis what's in it for chemists? *Chem. Aust.*, 59 (1992) 253-255; C.A., 117 (1992) 61925m - a review with 8 refs.
- 3 Cai, S. and Yao, Z.: (Principles, applications and advances of electrophoresis. Part I.) *Sepu*, 10 (1992) 207-210; C.A., 117 (1992) 127474x - a review without refs.
- 4 Chebotarev, V.Yu. and Rekes, A.N.: (Classical and pulsed gel electrophoresis of biopolymers: theory and applications). *Biofizika*, 37 (1992) 243-289; C.A., 117 (1992) 158091z - a review with 85 refs.
- 5 Chrambach, A., Dunn, M.J. and Radola, B.J. (Editors): *Advances in Electrophoresis*. Vol. 5, VCH, Weinheim, 1992, 414 p.; C.A., 117 (1992) 107774s.

See also 62, 66, 85, 87, 88, 92, 99, 100, 103, 155, 159, 217, 221, 255, 317, 370, 374, 388, 595, 596, 635, 646, 649, 705, 706, 716, 721, 744, 748.

2. FUNDAMENTALS, THEORY AND GENERAL

2a. General

- 6 Allison, S.A. and Nambi, P.: Transport of charged macromolecules in an electric field by a numerical method. *Macromolecules*, 25 (1992) 3971-3978; C.A., 117 (1992) 49658u.
- 7 Debets, A.j.j., ElArabi, H., Hupe, K.P., Brinkman, A.T., Hendriks, P.J.M., Claessens, H.A., Everaerts, F.M. and Noij, T.H.M.: Electrophoretic sample pre-treatment techniques coupled on-line with column liquid chromatography. *Analyst (London)*, 117 (1992) 1355-1359.
- 8 Dhar, H.P., Nath, S. and Lewis, D.H.: On the effect of magnetic field on electrophoresis. *Tex. J. Sci.*, 43 (1991) 334-336; C.A., 117 (1992) 167016f.
- 9 Férard, G.: Quantities and units for electrophoresis in the clinical laboratory. *Clin. Chim. Acta*, 205 (1992) S17-S23.
- 10 Kōst, H.-P.: Evaluation of isoelectric focusing runs as specific resistance/volthour plots. *Electrophoresis (Weinheim)*, 13 (1992) 660-661.
- 11 Slater, G.W.: Anomalous electrophoresis, self-trapping and "freezing" of partially charged polyelectrolytes. *J. Phys. II*, 2 (1992) 1149-1158; C.A., 117 (1992) 56674z.

See also 2.

2b. Thermodynamics and theoretical relationships

- 12 Delinger, S.L. and Davis, J.M.: Influence of analyte plug width on plate number in capillary electrophoresis. *Anal. Chem.*, 64 (1992) 1947-1959.
- 13 Dubrovčáková, E., Gas, B., Vacik, J. and Smolková-Keulemansová, E.: Electromigration in systems with additives in background electrolytes. I. Addition of the neutral complexing agent. *J. Chromatogr.*, 623 (1992) 337-344.
- 14 Poppe, H.: Overloading and interaction phenomena in electrophoretic separations. *Anal. Chem.*, 64 (1992) 1908-1919.

See also 75, 82, 89, 625.

2d. Measurement of physico-chemical and related values

- 15 Sugrue, S.: Predicting and controlling colloid particle behavior using multiangle electrophoretic mobility and particle size measurements. *Adv. Filtr. Sep. Technol.*, 4 (1991) 76-80; C.A., 117 (1992) 72458n.
- 16 Sugrue, S.: Predicting and controlling colloid suspension stability using electrophoretic mobility and particle size measurements. *Am. Lab. (Shelton)*, 24 (1992) 64-71; C.A., 117 (1992) 98264w.
- 17 Van Leeuwen, J.M.J. and Kooiman, A.: The drift velocity in the Rubinstein-Duke model for electrophoresis. *Physica A (Amsterdam)*, 184 (1992) 79-97; C.A., 117 (1992) 172424s.

See also 94.

3. GENERAL TECHNIQUES

3a. Apparatus and accessories

- 19 Brenner, N. and Palmieri, R.H.: On-column pre-concentration of samples in capillary electrophoresis. *Eur. Pat. Appl. EP 500,211* (Cl. G01N27/447), 26 Aug. 1992, US Appl. 649,385, 01 Feb. 1991; 5 p.; C.A., 117 (1992) 163191m.
- 20 Carson, W.W., Fuchs, M. and Cheng, Y.F.: Method and apparatus for capillary electrophoresis fraction collection on a membrane. *Eur. Pat. Appl. EP 477,541* (Cl. G01N27/447), 01 Apr. 1992, US Appl. 575,111, 30 Aug. 1990; 10 p.; C.A., 117 (1992) 39574z.
- 21 Chervet, J.P.: Micro flow processor. *Eur. Pat. Appl. 495,255* (Cl. G01N30/34), 22 Jul. 1992, Appl. 91/2000,080, 17 Jan. 1991; 12 p.; C.A., 117 (1992) 142628u.
- 22 Chien, R.L. and Burgi, D.S.: System and method for improving sample concentration in capillary electrophoresis. *U.S. US 5,116,471* (Cl. 204-180.1; G01N27/26), 26 May 1992, Appl. 771,575, 04 Oct. 1991; 13 p.; C.A., 117 (1992) 61986g.

- 23 Harrison, D.J., Manz, A., Fan, Z., Lüdi, H. and Widmer, H.M.: Capillary electrophoresis and sample injection systems integrated on a planar glass chip. *Anal. Chem.*, 64 (1992) 1926-1932.
- 24 Isola, M.C., Franzoni, P.E. and Franzoni, L.: (Simple power supply for polyacrylamide electrophoresis and blotting of proteins). *An. Asoc. Quim. Argent.*, 79 (1991) 85-89; C.A., 117 (1992) 86069y.
- 25 Jorgenson, J.W. and Monning, C.A.: Electrophoresis method and device for high speed separation of complex molecules. *Eur. Pat. Appl. EP 475,760* (Cl. G01N27/447), 18 Mar. 1992, US Appl. 581,683, 13 Sep. 1990; 15 p.; C.A., 117 (1992) 61983d.
- 26 Kerr, R. and Jung, L.: Continuous positive charge capillary columns used in capillary electrophoresis. *PCT Int. Appl. WO 92 05,433* (Cl. G01N27/26), 02 Apr. 1992, FR Appl. 90/11,720, 19 Sep. 1990; 26 p.; C.A., 117 (1992) 123805h.
- 27 Kozulic, B. and Heimgartner, U.: Apparatus and method for submerged gel electrophoresis. *PCT Int. Appl. WO 92 08,969* (Cl. G01N27/26), 29 May 1992, GB Appl. 90/24,428, 09 Nov. 1990; 31 p.; C.A., 117 (1992) 123809n.
- 28 Lauer, H.H., Ooms, J.A., Hansen, M.C., Logtenberg, F.J. and Loth, K.A.: Method and apparatus for introduction of a volume of at least one fluid in a tube, suitable for capillary electrophoresis systems. *Eur. Pat. Appl. EP 475,533* (Cl. G01N27/26), 18 Mar. 1992, EP Appl. 90/202,412, 11 Sep. 1990; 13 p.; C.A., 117 (1992) 61884x.
- 29 Matsuoka, H., Hoshino, H., Shiwa, S., Matsuzawa, J., Yamaguchi, M., Suzuki, K. and Uchida, T.: Electrophoretic display panel. *Jpn. Kokai Tokkyo Koho JP 04 86,785* (92 86,785) (Cl. G09F9/37), 19 Mar. 1992, Appl. 90/202,663, 31 Jul. 1990; 5 p.; C.A., 117 (1992) 101203h.
- 30 Morrison, I.D., Hsieh, B.R. and Taylor, J.H.: Liquid electrophoretic developer containing curable liquid vehicle. *Eur. Pat. Appl. EP 455,343* (Cl. G03G9/12), 06 Nov. 1991, US Appl. 501,585, 30 Mar. 1990; 21 p.; C.A., 117 (1992) 100927k.
- 31 Papp, A.: Photoelectric electrophoresis controller. *U.S. US 5,120,419* (Cl. 204-299R; G01N27/26), 09 Jun. 1992, Appl. 541,626, 21 Jun. 1990; 6 p.; C.A., 117 (1992) 61991e.
- 32 Schmidt, J.L. and Cheh, H.Y.: Free-flow electrophoresis method and apparatus. *U.S. US 5,122,246* (Cl. 204-180.1; B01D57/02), 16 Jun. 1992, Appl. 721,303, 26 Jun. 1991; 8 pp.; C.A., 117 (1992) 86349q.
- 33 Schmidt, J.L. and Cheh, H.Y.: Membrane electrophoresis method and apparatus. *Eur. Pat. Appl. EP 491,528* (Cl. B01D57/02), 24 Jun. 1992, US Appl. 628,595, 17 Dec. 1990; 8 pp.; C.A., 117 (1992) 107784v.
- 34 Sylvester, K.V.: Vertically oriented gel electrophoresis apparatus. *U.S. US 5,112,470* (Cl. 204-299R; G01N27/28), 12 May 1992, Appl. 624,074, 07 Dec. 1990; 9 p.; C.A., 117 (1992) 39576b.
- 35 Watson, J.S.: Method and apparatus for continuous electrophoresis without flow perturbations. *U.S. US 5,082,541* (Cl. 204-180.1; B01D57/02), 21 Jan. 1992, Appl. 551,387, 10 Jul. 1990; 11 p.; C.A., 117 (1992) 29741n.
- 36 Woolley, C.L.: De-activated capillary columns for use in electrophoresis and micellar electrokinetic chromatography. *Eur. Pat. Appl. EP 494,544* (Cl. G01N27/447), 15 Jul. 1992, US Appl. 638,934, 09 Jan. 1991; 10 p.; C.A., 117 (1992) 184054j.
- 37 Yamaguchi, M., Hoshino, H., Shiwa, S., Matsuoka, H., Matsuzawa, J., Suzuki, K. and Uchida, T.: (Electrophoretic displaying liquid and display device therewith). *Jpn. Kokai, Tokkyo, Koho JP 04 86,784* (92 86,784) (Cl. G09F9/37) 19 Mar. 1992, Appl. 90/202,662, 31 Jul. 1990; 6 p.; C.A., 117 (1992) 121521p.
- 38 Yang, S., Li, Y., Hu, R. and Tian, Z.: (A new capillary isotachophoretic analyzer without pumps and valves II. Open-style sample introduction device). *Fenxi Ceshi Tongbao*, 11 (1992) 30-33; C.A., 117 (1992) 103058p.
- 39 Zhuo, X., Yang, S., Mu, J. and Tian, Z.: (A microcomputer-controlled capillary isotachophoretic system). *Sepu*, 10 (1992) 171-172; C.A., 117 (1992) 33299s.

See also 67, 76, 77, 83, 89, 94, 97, 98.

3b. Detectors and detection procedures

- 40 Bruin, G.J.M., van Asten, A.C., Xu, X. and Poppe, H.: Theoretical and experimental aspects of indirect detection in capillary electrophoresis. *J. Chromatogr.*, 608 (1992) 97-107.
- 41 Grossman, P.D., Hino, T. and Soane, D.S.: Dynamic light-scattering studies of hydroxyethyl cellulose solutions used as sieving media for electrophoretic separations. *J. Chromatogr.*, 608 (1992) 79-83.
- 42 Hjerten, S. and Srichaiyo, T.: Multipoint detection method for electrophoresis and chromatography in capillaries. *U.S. US 5,114,551* (Cl. 204-180.1; B01D57/02), 19 May 1992, Appl. 769,073, 30 Sep. 1991; 11 pp.; C.A., 117 (1992) 44039h.
- 43 Huang, X., Zare, R.N., Ewing, A.G. and Sloss, S.E.: End-column electrical and electrochemical detector for capillary zone electrophoresis. *Eur. Pat. Appl. EP 475,713* (Cl. G01N27/447), 18 Mar. 1992, US Appl. 580,259, 10 Sep. 1990; 9 p.; C.A., 117 (1992) 61982c.
- 44 Kovarik, A., Ondrkalova, M., Prachar, J. and Stofko, J.: Rapid and sensitive colloidal silver staining on cellulose acetate membranes. *Clin. Chim. Acta*, 208 (1992) 137-139.
- 45 Langley, K.H.: Developments of electrophoretic laser light scattering and some biochemical applications. *Spec. Publ. - R. Soc. Chem.*, 99(Laser Light Scattering Biochem.) (1992) 151-160; C.A., 117 (1992) 65757f.
- 46 Nielsen, M.W.F.: Indirect time-resolved luminescence detection in capillary zone electrophoresis. *J. Chromatogr.*, 608 (1992) 85-92.
- 47 Sackmann, E., Stelzle, M. and Weissmueller, G.: Biosensors based on capacitance measurements and 2D-microelectrophoresis on lipid membranes. *GBF Monogr.*, 17(Biosens.: Fundam., Technol. Appl.) (1992) 133-143; C.A., 117 (1992) 86046p.
- 48 Thompson, M.A., Ford, J.C. and Long, G.R.: On-line detection in slab gel electrophoresis using thermal lens spectrophotometry. *Appl. Spectrosc.*, 46 (1992) 945-947; C.A., 117 (1992) 86171a.
- 49 Tong, W.G.: High-resolution, sensitive nonlinear laser detection method for chemical analysis based on degenerate four-wave mixing. *Proc. SPIE-Int. Soc. Opt. Eng.*, 1637 (1992) 25-32; C.A., 117 (1992) 142444f.
- 50 Toulas, C. and Hernandez, L.: Applications of a laser-induced fluorescence detector for capillary electrophoresis to measure attomolar and zeptomolar amounts of compounds. *LC-GC*, 10 (1992) 471-476; C.A., 117 (1992) 103288p.

- 51 Xu, M., Wu, H., Xu, L. and Huang, X.: (A simple sensitive silver stain method following electrophoresis for detecting Bence Jones proteins in unconcentrated urine). *Tongji Yike Daxue Xuebao*, 20 (1991) 45-47; C.A., 117 (1992) 22730z.
- 52 Yeung, E.S., Wang, P., Li, W. and Giese, R.W.: Laser fluorescence detector for capillary electrophoresis. *J. Chromatogr.*, 608 (1992) 73-77.
- See also 99, 111, 115, 118, 161, 167, 207, 209, 216, 217, 360, 393, 471, 525, 556, 599, 626.
- 3c. *Stabilization media for electrophoresis*
- 53 Ceron, C.R., Rodrigues dos Santos, J. and Bicudo de Campos, H.E.: The use of gelatin to dry cellophane wound slab gels in an embroidering hoop. *Rev. Bras. Genet.*, 15 (1992) 201-203; C.A., 117 (1992) 22740c.
- 54 Mori, Y. and Mikami, M.: Micromolecule electrophoresis membranes. *Jpn. Kokai Tokkyo Koho* JP 03,296,657 (91,296,657) (Cl. G01N27/447), 27 Dec. 1991, Appl. 90/99,171, 13 Apr. 1990; 11 p.; C.A., 117 (1992) 71234z.
- 55 Rapp, R.L., Kowalchuk, W.K., Davis, K.L., Todd, E.A., Liu, K.-L. and Morris, M.D.: Acrylamide polymerization kinetics in gel electrophoresis capillaries. A Raman microprobe study. *Anal. Chem.*, 64 (1992) 2434-2437.
- 56 Righetti, P.G., Caglio, S., Saracchi, M. and Quaroni, S.: Laterally aggregated polyacrylamide gels for electrophoresis. *Electrophoresis (Weinheim)*, 13 (1992) 587-595.
- 57 Stellwagen, J. and Stellwagen, N.C.: The effect of gel structure on matrix orientation. *Electrophoresis (Weinheim)*, 13 (1992) 595-600.
- 58 Stellwagen, N.C.: Agarose gel pore radii are not dependent on the casting buffer. *Electrophoresis (Weinheim)*, 13 (1992) 601-603.
- 59 Tanaka, N., Tanigawa, T., Hosoya, K., Kimata, K., Araki, T. and Terabe, S.: Starburst dendrimers as carriers in electrokinetic chromatography. *Chem. Lett.*, (1992) 959-962; C.A., 117 (1992) 103278k.
- 60 Yoshida, T., Kawase, T. and Hiratsuka, N.: Microporous membranes for electrophoresis of serum proteins. *Jpn. Kokai Tokkyo Koho* JP 04 76,450 [92 76,450] (Cl. GO1N27/447), 11 Mar. 1992, Appl. 90/190,316, 18 Jul. 1990; 6 pp.; C.A., 117 (1992) 22888g.
- See also 562, 717.
- 3e. *Preparative scale electrophoresis*
- 61 Hartig, R., Hausmann, M., Schmitt, J., Herrmann, D.B.J., Riedmiller, M. and Cremer, C.: Preparative continuous separation of biological particles by means of free-flow magnetophoresis in a free-flow electrophoresis chamber. *Electrophoresis (Weinheim)*, 13 (1992) 674-676.
- 62 Hattori, H., Kaneda, T. and Ohtsuka, K.: (Preparative 2-dimensional electrophoresis. Slab NEPHGE/SDS-PAGE). *Seikagaku*, 64 (1992) 416-420; C.A., 117 (1992) 107456q - a review with 17 refs.
- 63 Righetti, P.G., Faupel, M. and Wensch, E.: Preparative electrophoresis with and without immobilized pH gradients. *Adv. Electrophor.*, 5 (1992) 159-200; C.A., 117 (1992) 127461r - a review with 89 refs.
- See also 463, 465, 509.
- 3f. *Programmed voltage and buffer gradients*
- See 612.
4. SPECIAL TECHNIQUES
- 4a. *Automation*
- 64 Lai, E., Birren, B.W., Stubblebine, W.H. and Bers, G.E.: Automatic control system for improvement of electrophoretic separation of macromolecules. *Ger. Offen. DE 4,132,694* (Cl. B01D57/02), 16 Apr. 1992, US Appl. 591,589, 02 Oct. 1990; 35 pp.; C.A., 117 (1992) 44058p.
- See also 194.
- 4b. *Computerization and modelling*
- 65 Onana, A.: Two dimensional finite element simulation of electrochemical separation process. *Hung. J. Ind. Chem.*, 20 (1992) 103-108; C.A., 117 (1992) 29436s.
- See also 39, 605, 651.
- 4c. *Combination with other physicochemical techniques, (MS, IR etc.)*
- See 73, 91, 178, 183, 187, 298.
- 4d. *Affinity electrophoresis*
- 66 Breborowicz, J. and Mackiewicz, A. (Editors): *Affinity Electrophoresis: Principles and Application* RCR, Boca Raton, 1992, 279 p.; C.A., 117 (1992) 127776x.
- See also 201, 294, 304, 561, 562.
- 4e. *Capillary zone electrophoresis and electrokinetic chromatography*
- 67 Abdel-Baky, S. and Giese, R.W.: Capillary electrophoresis washing technique. *J. Chromatogr.*, 608 (1992) 159.
- 68 Albin, M., Chen, S.-M., Louie, A., Pairaud, C., Colburn, J. and Wiktorowicz, J.: The use of capillary electrophoresis in a micro-preparative mode: methods and applications. *Anal. Biochem.*, 206 (1992) 382-388.
- 69 Altria, K.D., Goodall, D.M. and Rogan, M.M.: Chiral separation of β -amino alcohols by capillary electrophoresis using cyclodextrins as buffer additives. I. Effect of varying operating parameters. *Chromatographia*, 34 (1992) 19-24.
- 70 Belder, D. and Schomburg, G.: Modification of silica surfaces for CZE by adsorption of non-ionic hydrophilic polymers or use of radial electric field. *J. High Resolut. Chromatogr.*, 15 (1992) 686-693.
- 71 Cai, J. and El Rassi, Z.: Micellar electrokinetic capillary chromatography of neutral solutes with micelles of adjustable surface charge density. *J. Chromatogr.*, 608 (1992) 31-45.

- 72 Camilleri, P. and Okafu, G.N.: Halofantrine incorporates in sodium dodecyl sulphate (SDS) micelles: application in fluorescence and capillary electrophoresis study. *J. Chem. Soc., Chem. Commun.*, (1992) 530-532; *C.A.*, 117 (1992) 97296w.
- 73 Castoro, J.A., Chiu, R.W., Monnig, C.A. and Wilkins, C.L.: Matrix-assisted laser desorption/ionization of capillary electrophoresis effluents by Fourier transform mass spectrometry. *J. Am. Chem. Soc.*, 114 (1992) 7571-7572; *C.A.*, 117 (1992) 127753n.
- 74 Chang, H.-T. and Yeung, E.S.: Optimization of selectivity in capillary zone electrophoresis via dynamic pH gradient and dynamic flow gradient. *J. Chromatogr.*, 608 (1992) 65-72.
- 75 Chen, Y. and Zhu, A.: Flat capillary zone electrophoresis. *Sci. China, Ser. B.*, 35 (1992) 649-658; *C.A.*, 117 (1992) 158297w.
- 76 Cheng, Y.-F., Fuchs, M., Andrews, D. and Carson, W.: Membrane fraction collection for capillary electrophoresis. *J. Chromatogr.*, 608 (1992) 109-116.
- 77 Debets, A.J.J., Mazereeuw, M., Voogt, W.H., van Iperen, D.J., Lingeman, H., Hupe, K.-P. and Brinkman, U.A.T.: Switching valve with internal micro precolumn for on-line sample enrichment in capillary zone electrophoresis. *J. Chromatogr.*, 608 (1992) 151-158.
- 78 Fu, X., Lu, J. and Zhu, A.: (Quantitation by micellar electrokinetic capillary chromatography). *Sepu*, 10 (1992) 234-235; *C.A.*, 117 (1992) 184037f.
- 79 Gebauer, P., Thormann, W. and Boček, P.: Theoretical description of the zone electrophoretic separation of minor compounds in the presence of bulk amounts of a sample component with high mobility and like charge. *J. Chromatogr.*, 608 (1992) 47-57.
- 80 Hermentin, P., Witzel, R., Doenges, R., Bauer, R., Haupt, H., Patel, T., Parekh, R.B. and Brazel, D.: The mapping of high-pH anion-exchange chromatography with pulsed amperometric detection and capillary electrophoresis of the carbohydrate moieties of human plasma α_1 -acid glycoprotein. *Anal. Biochem.*, 206 (1992) 419-429.
- 81 Kaneta, T., Tanaka, S., Taga, M. and Yoshida, H.: Effect of addition of glucose on micellar electrokinetic capillary chromatography with sodium dodecyl sulphate. *J. Chromatogr.*, 609 (1992) 369-374.
- 82 Kenndler, E. and Friedl, W.: Adjustment of resolution and analysis time in capillary zone electrophoresis by varying the pH of the buffer. *J. Chromatogr.*, 608 (1992) 161-170.
- 83 Kobayashi, S.: Capillary electrophoresis apparatus for separating proteins and nucleic acids. *Jpn. Kokai Tokkyo Koho JP 04 84,752* [92 84,752] (Cl. GO1N27/447), 18 Mar. 1992, Appl. 90/199,861, 27 Jul. 1990; 5 pp.; *C.A.*, 117 (1992) 66045j.
- 84 Kobayashi, S.: Device for sample introduction into capillary electrophoresis apparatus for separation-determination of proteins and other substances. *Jpn. Kokai Tokkyo Koho JP 04,160,356* [92,160,356] (Cl. GO1N27/447), 03 Jun. 1992, Appl. 90/287,665, 24 Oct. 1990; 3 pp.; *C.A.*, 117 (1992) 146727y.
- 85 Kuhn, R. and Hoffstetter-Kuhn, S.: Chiral separation by capillary electrophoresis. *Chromatographia*, 34 (1992) 505-512 - a review with 49 refs.
- 86 Landers, J.P., Oda, R.P., Madden, B., Sismelich, T.P. and Spelsberg, T.C.: Reproducibility of sample separation using liquid or forced air convection thermostatted high performance capillary electrophoresis systems. *J. High Resolut. Chromatogr.*, 15 (1992) 517-525.
- 87 Okuyama, T. and Izumi, T.: (Capillary electrophoresis). *Gendai Kagaku*, 257 (1992) 36-42; *C.A.*, 117 (1992) 183992b - a review with no refs.
- 88 Rush, R.S.: High performance capillary electrophoresis of proteins and peptides: a minireview. In: Kelly, J.W. and Baldwin, T.O. (Editors), *Appl. Enzyme Biotechnol.*, [Proc. Tex. A&M Univ., IUCCP Symp.], 9th 1991, Plenum, New York, 1991, pp. 233-250; *C.A.*, 117 (1992) 166866w - a review with 78 refs.
- 89 Schützner, W. and Kenndler, E.: Electrophoresis in synthetic organic polymer capillaries: variation of electroosmotic velocity and zeta potential with pH and solvent composition. *Anal. Chem.*, 64 (1992) 1991-1995.
- 90 Smith, J.T. and El Rassi, Z.: Capillary zone electrophoresis of biological substances with surface-modified fused silica capillaries with switchable electroosmotic flow. *J. High Resolut. Chromatogr.*, 15 (1992) 573-578.
- 91 Smith, R.D., Barinaga, C.J. and Udseth, H.R.: Capillary zone electrophoresis/mass spectrometry an alternative to liquid chromatography/mass spectrometry? *Lab. 2000*; 5 (1991) 38-42; *C.A.*, 117 (1992) 103271c.
- 92 Terabe, S. and Matsubara, N.: (Capillary electrophoresis). *Kagaku Kogyo*, 43 (1992) 144-154; *C.A.*, 117 (1992) 103248a - a review with 19 refs.
- 93 Terabe, S., Matsubara, N., Ishihama, Y. and Okada, Y.: Microemulsion electrokinetic chromatography: comparison with micellar electrokinetic chromatography. *J. Chromatogr.*, 608 (1992) 23-29.
- 94 Towns, J.K. and Regnier, F.E.: Impact of polycation adsorption on efficiency and electroosmotically driven transport in capillary electrophoresis. *Anal. Chem.*, 64 (1992) 2473-2478.
- 95 Tribet, C., Gaboriaud, R. and Gareil, P.: Analogy between micelles and polymers of ionic surfactants. A capillary isotachophoretic study of small ionic aggregates in water-organic solutions. *J. Chromatogr.*, 608 (1992) 131-141.
- 96 Vespalec, R., Gebauer, P. and Bocek, P.: Identification of peaks in capillary zone electrophoresis based on actual mobilities. *Electrophoresis (Weinheim)*, 13 (1992) 677-682.
- 97 Whang, C.-W. and Chen, I.-C.: Cellulose acetate-coated porous polymer joint for capillary zone electrophoresis. *Anal. Chem.*, 64 (1992) 2461-2464.
- 98 Wu, C.-T., Lee, C.S. and Miller, C.J.: Ionized air for applying radial potential gradient in capillary electrophoresis. *Anal. Chem.*, 64 (1992) 2310-2311.
- 99 Yik, Y.F. and Li, S.F.Y.: Capillary electrophoresis with electrochemical detection. *TrAC*, 11 (1992) 325-332 - a review with 65 refs.
- 100 Zeniba, T.: (Bioscience and capillary electrophoresis (CE). 1.) *Cell Sci.*, 8 (1992) 141-144; *C.A.*, 117 (1992) 65712n - a review with 1 ref.

See also 2, 12, 19, 22, 23, 26, 28, 36, 39, 40, 41, 42, 43, 46, 50, 52, 59, 104, 105, 106, 107, 108, 109, 110, 111, 113, 115, 125, 129, 133, 134, 135, 158, 159, 160, 161, 162, 163, 164, 166, 167, 169, 170, 174, 176, 177, 178, 181, 182, 183, 185, 196, 197, 201, 203, 205, 208, 213, 215, 219, 220, 221, 245, 351, 359, 364, 447, 448, 464, 551, 561, 562, 563, 565, 566, 595, 611, 612, 630, 652, 669, 675, 699, 700, 701, 702, 703, 704, 707, 709, 711, 712, 713, 714, 717, 718, 719, 720, 721, 722, 723, 724, 725, 726, 728, 729, 730, 731, 732, 733, 734, 735, 737, 738, 739, 741, 750, 752, 753, 754, 755.

4f. Isotachopheresis

- 101 Burgstaller, W. and Schimer, F.: Isotachopheretic analysis of lipophilic cation tetraphenylphosphoniumbromide. *Fresenius J. Anal. Chem.*, 343 (1992) 893-895.
- 102 Hendriks, P.J.M., Claessens, H.A., Noij, T.H.M., Everaerts, F.M. and Cramers, C.A.: On-line isotachopheresis. A selective sample pretreatment prior to column liquid chromatographic analysis. *Chromatographia*, 33 (1992) 539-545.
- 103 Yoshida, H., Tanaka, S., Hiramata, Y. and Kan, M.: Separation and determination of trace elements by isotachopheresis. In: Alfassi, Z.B. and Wai, C.M. (Editors), *Preconc. Tech. Trace Elem.*, CRC, Boca Raton, 1992, pp. 417-426; C.A., 117 (1992) 123496h - a review with 26 refs.

See also 38, 136, 203, 533, 710, 721, 740.

4g. Enantiomers, separation

- 104 Cruzado, I.D. and Vigh, G.: Chiral separations by capillary electrophoresis using cyclodextrin-containing gels. *J. Chromatogr.*, 608 (1992) 421-425.
- 105 Schutzner, W. and Fanali, S.: Enantiomers resolution in capillary zone electrophoresis by using cyclodextrins. *Electrophoresis (Weinheim)*, 13 (1992) 687-690.
- 106 Wren, S.A.C. and Rowe, R.C.: Theoretical aspects of chiral separation in capillary electrophoresis. II. The role of organic solvent. *J. Chromatogr.*, 609 (1992) 363-367.

See also 69, 70, 85, 160, 164, 723, 725, 728.

4h. Two dimensional electrophoresis

See 62, 131, 146, 187, 189, 194, 198, 199, 214, 216, 231, 233, 242, 263, 268, 293, 316, 333, 337, 338, 342, 346, 366, 367, 446, 509, 553, 643, 698.

4i. Other special techniques

See 35, 47, 85, 107, 177, 331, 678, 712.

6. ALCOHOLS

See 703.

8. SUBSTANCES CONTAINING HETEROCYCLIC OXYGEN

8a. Flavonoids

- 107 Ng, C.L., Ong, C.P., Lee, H.K. and Li, S.F.Y.: Systematic optimization of micellar electrokinetic chromatographic separation of flavonoids. *Chromatographia*, 34 (1992) 166-172.

8c. Other compounds with heterocyclic oxygen (incl. tannins)

- 108 Corradini, D. and Corradini, C.: Separation and determination of 5-hydroxymethyl-2-furaldehyde and 2-furaldehyde in fruit juices by micellar electrokinetic capillary chromatography with direct sample injection. *J. Chromatogr.*, 624 (1992) 503-509.

See also 716.

10. CARBOHYDRATES

10a. Mono and oligosaccharides. Structural studies

- 109 Damm, J.B.L., Overklist, G.T., Vermeulen, B.W.M., Fluitsma, C.F. and van Dedem, G.W.K.: Separation of natural and synthetic heparin fragments by high-performance capillary electrophoresis. *J. Chromatogr.*, 608 (1992) 297-309.
- 110 Honda, S., Ueno, T. and Kakehi, K.: High-performance capillary electrophoresis of unsaturated oligosaccharides derived from glycosaminoglycans by digestion with chondroitinase ABC as 1-phenyl-3-methyl-5-pyrazolone derivatives. *J. Chromatogr.*, 608 (1992) 289-295.
- 111 Oefner, P.J., Vorndran, A.E., Grill, E., Huber, C. and Bonn, G.K.: Capillary zone electrophoretic analysis of carbohydrates by direct and indirect UV detection. *Chromatographia*, 34 (1992) 308-316.
- 112 Stack, R.J. and Sullivan, M.T.: Electrophoretic resolution and fluorescence detection of N-linked glycoprotein oligosaccharides after reductive amination with 8-aminonaphthalene-1,3,6-trisulfonic acid. *Glycobiology*, 2 (1992) 85-92; C.A., 117 (1992) 65881s.
- 113 Vorndran, A.E., Grill, E., Huber, C., Defner, P.J. and Bonn, G.K.: Capillary zone electrophoresis of aldoses, ketoses and uronic acids derivatized with ethyl *p*-aminobenzoate. *Chromatographia*, 34 (1992) 109-114.

See also 80, 90, 115, 129.

10b. Polysaccharides, mucopolysaccharides, lipopolysaccharides

- 114 Hiraoka, A., Harada, N., Uehara, T., Sekiguchi, M. and Maeda, M.: Capillary-isotachopheretic analyses of algal acidic polysaccharides and their application to a survey of heparinoid active sulfated polysaccharides in chlorophyta. *Chem. Pharm. Bull.*, 40 (1992) 783-785.
- 115 Kakehi, K.: Combination of capillary electrophoresis and laser-induced fluorescence can detect ten thousand carbohydrate molecules. *Trends Glycosci. Glycotechnol.*, 4 (1992) 106-107; C.A., 117 (1992) 22817h.

- 116 Kikuchi, A., Satoh, S. and Fujii, T.: Analysis of plant pectic substances by electrophoresis on a cellulose acetate membrane. *Biosci., Biotechnol., Biochem.*, 56 (1992) 1144-1145; C.A., 117 (1992) 149602q.
- 117 Sakane, T. and Yokota, A.: (Rapid method for lipopolysaccharide analysis with deoxycholate-polyacrylamide gel electrophoresis and PAS stain). *Seikagaku*, 64 (1992) 533-536; C.A., 117 (1992) 127648g.
- 118 Uchisawa, H., Takatani, Y., Hanamatsu, N., Matsue, H., Narumi, F., Kono, T. and Okubo, O.: Colorants from Cephalopoda as stains for detecting acidic polysaccharides and nucleic acids by cellulose acetate membrane electrophoresis. *Jpn. Kokai Tokkyo Koho JP 04 43,954 [92 43,954] (Cl. GO1N27/447)*, 13 Feb. 1992, Appl. 90/153,161, 11 Jun. 1990; 3 pp.; C.A., 117 (1992) 22906m.
- 10c. *Glycoproteins and their constituents*
- 119 Chen, F.T.A. and Sternberg, J.C.: Analysis of glycoprotein samples utilizing capillary zone electrophoresis. *U.S. US 5,120,413 (Cl.204-180.1; GO1N27/26)*, 09 Jun. 1992, Appl. 708,272, 31 May 1991; 13 pp.; C.A., 117 (1992) 127810d.
- 120 Gentile, F., Palumbo, G. and Salvatore, G.: The origin of the electrophoretic doublet of thyroglobulin. *Biochem. Biophys. Res. Commun.*, 186 (1992) 1185-1191.
- 121 Gianazza, E., Frattini, R., Michelagnoli, S., Cassinotti, M. and Sirtori, C.R.: Isoelectric focusing and immunoblotting of the platelet membrane glycoprotein complex IIb/IIIa following urea solubilization. *Electrophoresis (Weinheim)*, 13 (1992) 781-784.
- 122 Gonzales, J., Takao, T., Hori, H., Besada, V., Rodriguez, R., Pardon, G. and Shimonishi, Y.: A method for determination of N-glycosylation sites in glycoproteins by collision-induced dissociation analysis in fast atom bombardment mass spectrometry: identification of the positions of carbohydrate-linked asparagine in recombinant α -amylase by treatment with peptide-N-glycosidase F in ^{18}O -labeled water. *Anal. Biochem.*, 205 (1992) 151-158.
- 123 Kim, D.H., Kanno, C. and Mizokami, Y.: Purification and characterization of major glycoproteins, PAS-6 and PAS-7, from bovine milk fat globule membrane. *Biochim. Biophys. Acta*, 1122 (1992) 203-211.
- 124 Krueger, R.C., Jr., Hennig, A.K. and Schwartz, N.B.: Two immunologically and developmentally distinct chondroitin sulfate proteoglycans in embryonic chick brain. *J. Biol. Chem.*, 267 (1992) 12149-12161.
- 125 Landers, J.P., Oda, R.P., Madden, B.J. and Spelsberg, T.C.: High-performance capillary electrophoresis of glycoproteins: the use of modifiers of electroosmotic flow for analysis of microheterogeneity. *Anal. Biochem.*, 205 (1992) 115-124.
- 126 Loomis, C.A., Kolega, J., Manabe, M. and Sun, T.: Characterization of a keratinocyte-specific extracellular epitope of desmoglein. Implications for desmoglein heterogeneity and function. *J. Biol. Chem.*, 267 (1992) 16676-16684.
- 127 Mayinger, P. and Gawaz, M.: Photoaffinity labeling of integrin $\alpha\text{IIb}\beta\text{3}$ (glycoprotein IIb-IIIa) on intact platelets with 8-azido-[γ - ^{32}P]ATP. *Biochim. Biophys. Acta*, 1137 (1992) 77-81.
- 128 Reddy, M.S., Bobek, L.A., Haraszthy, G.G., Biesbrock, A.R. and Levine, M.J.: Structural features of the low-molecular-mass human salivary mucin. *Biochem. J.*, 287 (1992) 639-643.
- 129 Suzuki, S., Kakehi, K. and Honda, S.: Two-dimensional mapping of N-glycosidically linked asialo-oligosaccharides from glycoproteins as reductively pyridylaminated derivatives using dual separation modes of high-performance capillary electrophoresis. *Anal. Biochem.*, 205 (1992) 227-236.
- 130 Warburton, M.J., Kimbell, R., Ferns, S.A., Haynam, A.R., Perusinghe, N. and Monaghan, P.: Characterisation of chondroitin/dermatan sulphate proteoglycans synthesised by rat mammary myoepithelial and fibroblastic cell lines. *Biochim. Biophys. Acta*, 1117 (1992) 291-300.
- 131 Weiss, W., Postel, W. and Görg, A.: Application of sequential extraction procedures and glycoprotein blotting for the characterization of the 2-D polypeptide patterns of barley seed proteins. *Electrophoresis (Weinheim)*, 13 (1992) 770-773.
- 132 Wilson, M.R. and Easterbrook-Smith, S.B.: Clusterin binds by a multivalent mechanism to the Fc and Fab regions of IgG. *Biochim. Biophys. Acta*, 1159 (1992) 319-326.
- See also 80, 112, 237, 289, 309, 310, 367, 400, 412, 419, 455, 517.
11. ORGANIC ACIDS AND LIPIDS
- 11a. *Organic acids and simple esters*
- 133 Bjerregaard, C., Michaelsen, S. and Sorensen, H.: Determination of phenolic carboxylic acids by micellar electrokinetic capillary chromatography and evaluation of factors affecting the method. *J. Chromatogr.*, 608 (1992) 403-411.
- 134 Fu, X., Lu, J., Lu, Y. and Xu, H.: (Application of micellar electrokinetic capillary chromatographic analysis in the synthesis of mandelic acid). *Gaodeng Xuexiao Huaxue Xuebao*, 13 (1992) 45-47; C.A., 117 (1992) 103304r.
- 135 Tienstra, P.A., van Riel, J.A.M., Mingorance, M.D. and Olliman, C.: Assessment of the capabilities of capillary zone electrophoresis for the determination of hippuric and orotic acid in whey. *J. Chromatogr.*, 608 (1992) 357-361.
- 136 Zabranska, J., Pokorna, D. and Dohanyos, M.: Isotachophoretic determination of byproducts of anaerobic metabolism. *Sb. Vys. Sk. Chem.-Technol. Praze, F: Technol. Vody Prostredi*, F 28 (1991) 149-158; C.A., 117 (1992) 177949b.
- See also 102, 737.
- 11c. *Lipids and their constituents*
- 137 Stahl, N., Baldwin, M.A., Hecker, R., Pan, K., Burlingame, A.L. and Prusiner, S.B.: Glycosylinositol phospholipid anchors of the scrapie and cellular prion proteins contain sialic acid. *Biochemistry*, 31 (1992) 5043-5053.
- See also 117, 136, 285.

11d. Lipoproteins and their constituents

- 138 Bard, J.-M., Delattre-Lestavel, S., Clavey, V., Pont, P., Derudas, B., Parra, H.-J. and Fruchart, J.-C.: Isolation and characterization of two sub-species of Lp(a), one containing apo E and one free of apo E. *Biochim. Biophys. Acta*, 1127 (1992) 124-130.
- 139 Burkey, B.F., Stuart, W.D. and Harmony, J.A.K.: Hepatic apolipoprotein J is secreted as a lipoprotein. *J. Lipid Res.*, 33 (1992) 1517-1526.
- 140 De Rijke, Y.B., Jürgens, G., Hessels, E.M.A., Hermann, A. and van Berkel, T.J.C.: *In vivo* fate and scavenger receptor recognition of oxidized lipoprotein [a] isoforms in rats. *J. Lipid Res.*, 33 (1992) 1315-1325.
- 141 Dobiasova, M., Stribrna, J., Pritchard, P.H. and Frohlich, J.J.: Cholesterol esterification rate in plasma depleted of very low and low density lipoproteins is controlled by the proportion of HDL₂ and HDL₃ subclasses: study in hypertensive and normal middle-aged and septuagenarian men. *J. Lipid Res.*, 33 (1992) 1411-1418.
- 142 Fantappie, S., Corsini, A., Sidoli, A., Uboldi, P., Granata, A., Zanelli, T., Rossi, P., Marcovina, S., Fumagalli, R. and Catapano, A.L.: Monoclonal antibodies to human low density lipoprotein identity distinct areas on apolipoprotein B-100 relevant to the low density lipoprotein-receptor interaction. *J. Lipid Res.*, 33 (1992) 1111-1121.
- 143 Farrer, M., Game, F.L., Adams, P.C., Laker, M.F. and Alberti, K.G.M.M.: A simple, sensitive technique for classification of apolipoprotein(a) isoforms by sodium dodecyl sulphate-polyacrylamide gel electrophoresis. *Clin. Chim. Acta*, 207 (1992) 215-225.
- 144 Mamo, J.C.L., Hirano, T., Sainsbury, A., Fitzgerald, A.K. and Redgrave, T.G.: Hypertriglyceridemia is exacerbated by slow lipolysis of triacylglycerol-rich lipoproteins in fed but not fasted streptozotocin diabetic rats. *Biochim. Biophys. Acta*, 1128 (1992) 132-138.
- 145 Mezdour, H., Nomura, S., Yamamura, T. and Yamamoto, A.: Concentration and distribution of apolipoproteins A-I and E in normolipidemic, WHHL and diet-induced hyperlipidemic rabbit sera. *Biochim. Biophys. Acta*, 1127 (1992) 116-123.
- 146 Moriyama, K., Sasaki, J., Matsunaga, A., Arakawa, F., Takada, Y., Araki, K., Kaneko, S. and Arakawa, K.: Apolipoprotein E1 lys-146→Glu with type III hyperlipoproteinemia. *Biochim. Biophys. Acta*, 1128 (1992) 58-64.
- 147 Mowri, H.-O., Patsch, W., Smith, L.C., Gotto, A.M., Jr. and Patsch, J.R.: Different reactivities of high density lipoprotein subfractions with hepatic lipase. *J. Lipid Res.*, 33 (1992) 1269-1279.
- 148 Oppl, J.J.: Discontinuous and nonsequential polymeric gel systems for separation of macromolecules. *PCT Int. Appl. WO 92 10,277* (Cl. BO1D61/42), 25 Jun. 1992, US Appl. 627,054, 13 Dec. 1990; 46 pp.; *C.A.*, 117 (1992) 127817m.
- 149 Peterson, J., Fujimoto, W.Y. and Brunzell, J.D.: Human lipoprotein lipase: relationship of activity, heparin affinity, and conformation as studied with monoclonal antibodies. *J. Lipid Res.*, 33 (1992) 1165-1170.
- 150 Plonné, D., Winkler, L., Franke, H. and Dargel, R.: The visceral yolk sac - an important site of synthesis and secretion of apolipoprotein B containing lipoproteins in the foeto-placental unit of the rat. *Biochim. Biophys. Acta*, 1127 (1992) 174-185.
- 151 Rainwater, D.L., Kammerer, C.M., Cheng, M.L., Sparks, M.L. and VandeBerg, J.L.: Distribution of specific apolipoproteins determined by immunoblotting of baboon lipoproteins resolved by polyacrylamide gradient gel electrophoresis. *Biochem. Genet.*, 30 (1992) 143-158; *C.A.*, 117 (1992) 43994d.
- 152 Rumsey, S.C., Galeano, N.F., Arad, Y. and Deckelbaum, R.J.: Cryopreservation with sucrose maintains normal physical and biological properties of human plasma low density lipoproteins. *J. Lipid Res.*, 33 (1992) 1551-1561.
- 153 Ruzicka, V., März, W., Russ, A. and Gross, W.: Apolipoprotein B (Arg³⁵⁰⁰→Gln) allele specific polymerase chain reaction: large-scale screening of pooled blood samples. *J. Lipid Res.*, 33 (1992) 1563-1567.
- 154 Salmon, S., Santus, R., Mazière, J.C., Aubailly, M. and Haigle, J.: Modified apolipoprotein pattern after irradiation of human high-density lipoproteins by ultraviolet B. *Biochim. Biophys. Acta*, 1128 (1992) 167-173. 422, immmblot.
- 155 Shore, V.G.: Nondenaturing electrophoresis of lipoproteins in agarose and polyacrylamide gradient gels. In: Perkins, E.G. (Editor), *Anal. Fats, Oils Lipoproteins*, AOCS, Champaign, 1991, pp. 573-588; *C.A.*, 117 (1992) 166829m - a review with 4 refs.
- 156 Thomas, C.E.: The influence of medium components on Cu²⁺-dependent oxidation of low-density lipoproteins and its sensitivity to superoxide dismutase. *Biochim. Biophys. Acta*, 1128 (1992) 50-57.
- 157 Vilella, E., Joven, J., Bargalló, T., Turner, P.R. and Masana, L.: Interaction of oxidized low density lipoproteins with both apo B,E and scavenger receptors. A model for its production *in vitro* *Clin. Chim. Acta*, 210 (1992) 93-108.

See also 454.

15. TERPENES AND OTHER VOLATILE AROMATIC COMPOUNDS

15b. Essential oils

- 158 Verschuere, M., Sandra, P. and David, F.: Fractionation by SFE and microcolumn analysis of the essential oil and the bitter principles of hops. *J. Chromatogr. Sci.*, 30 (1992) 388-391.

15c. Bitter substances

- 159 De Keukeleire, D., Vindevogel, J., Szücs, R. and Sandra, P.: The history and analytical chemistry of beer bitter acids. *TRAC*, 11 (1992) 275-280 - a review with 34 refs.

See also 158.

17. AMINES, AMIDES AND RELATED NITROGEN COMPOUNDS

17a. Amines and polyamines

- 160 Höhne, E., Krauss, G.-J. and Gübitz, G.: Capillary zone electrophoresis of the enantiomers of aminoalcohols based on host-guest complexation with a chiral crown ether. *J. High Resolut. Chromatogr.*, 15 (1992) 698-700.

- 161 Ma, Y., Zhang, R. and Cooper, C.L.: Indirect photometric detection of polyamines in biological samples separated by high-performance capillary electrophoresis. *J. Chromatogr.*, 608 (1992) 93-96.
- See also 69, 728.
- 17d. *Other amine derivatives and amides (excl. peptides)*
- 162 Lukkari, P., Jumppanen, J., Holma, T., Sirén, H., Jinno, K., Elo, H. and Riekkola, M.-L.: Effect of the buffer solution on the elution order and separation of bis(amidinohydrazones) by micellar electrokinetic capillary chromatography. *J. Chromatogr.*, 608 (1992) 317-323.
- 163 Weiss, C.S., Hazlett, J.S., Datta, M.H. and Danzer, M.H.: Determination of quaternary ammonium compounds by capillary electrophoresis using direct and indirect UV detection. *J. Chromatogr.*, 608 (1992) 325-332.
- 164 Yamashoji, Y., Ariga, T., Asano, S. and Tanaka, M.: Chiral recognition and enantiomeric separation of alanine β -naphthylamide by cyclodextrins. *Anal. Chim. Acta*, 268 (1992) 39-47.
- See also 717, 718, 732.
18. AMINO ACIDS AND PEPTIDES; CHEMICAL STRUCTURE OF PROTEINS
- 18a. *Amino acids and their derivatives*
- 165 Lippmann, C., Lindschau, C. and Erdmann, V.A.: Thin-layer electrophoresis with PhastSystem facilitates analysis of phosphoamino acids from proteins bound to Immobilon. *Electrophoresis (Weinheim)*, 13 (1992) 666-668.
- 166 O'Shea, T.J., Weber, P.L., Bammel, B.P., Lunte, C.E., Lunte, S.M. and Smyth, M.R.: Monitoring excitatory amino acid release *in vivo* by microdialysis with capillary electrophoresis-electrochemistry. *J. Chromatogr.*, 608 (1992) 189-195.
- 167 Zhao, J.-Y., Chen, D.-Y. and Dovichi, N.J.: Low-cost laser-induced fluorescence detector for micellar capillary zone electrophoresis. Detection at the zeptomol level of tetramethylrhodamine thiocarbonyl amino acid derivatives. *J. Chromatogr.*, 608 (1992) 117-120.
- See also 40, 135, 169, 205, 211, 485, 498.
- 18b. *Peptides, peptidic and proteinous hormones, growth factors*
- 168 Amarowicz, R. and Kmita-Glazewska, H.: HPTLC and electrochromatography of rapeseed peptides. *Nahrung*, 36 (1992) 87-89; *C.A.*, 117 (1992) 65777n.
- 169 Bergman, T. and Joernvall, H.: Capillary electrophoresis for preparation of peptides and direct determination of amino acids. In: Angeletti, R.H. (Editor), *Tech. Protein Chem. 3, [Pap. Annu. Symp. Protein Soc.]*, 5th 1991, Academic, San Diego, 1992, pp. 129-134; *C.A.*, 117 (1992) 65893x.
- 170 Crimmins, D.L.: Comparative analysis of synthetic peptides by free-solution capillary electrophoresis (FSCE) and strong-cation exchange (SCX)-HPLC. In: Angeletti, R. (Editor), *Tech. Protein Chem. 3, [Pap. Annu. Symp. Protein Soc.]*, 5th 1991, Academic, San Diego, 1992, pp. 171-181; *C.A.*, 117 (1992) 65787r.
- 171 Domeyne, A., Pinset, C., Montarras, D., Garandel, V., Rosenfeld, R.G. and Barenton, B.: Preferential binding to insulin-like growth factor-II (IGF-II) to a putative $\alpha 2\beta 2$ IGF-II receptor type in C2 myoblasts. *Eur. J. Biochem.*, 208 (1992) 273-279.
- 172 Emly, J.F., Ratcliffe, W.A., Green, E., Bowden, S.J., Heath, D.A., Blight, A., Hughes, S. and Ratcliffe, J.G.: Immunochemical characterisation of parathyroid hormone-related protein from tumour and non-tumour cells. *Biochim. Biophys. Acta*, 1180 (1992) 58-64.
- 173 Heegaard, N.H.H. and Robey, F.A.: Use of capillary zone electrophoresis to evaluate the binding of anionic carbohydrates to synthetic peptides derived from human serum amyloid P component. *Anal. Chem.*, 64 (1992) 2479-2482.
- 174 Kašička, V., Prusik, Z. and Pospíšek, J.: Conversion of capillary zone electrophoresis to free-flow zone electrophoresis using a simple model of their correlation. Application to synthetic enkephalin-type peptide analysis and preparation. *J. Chromatogr.*, 608 (1992) 13-22.
- 175 Lakhani, S., Khanna, N.C. and Tewari, K.K.: Two distinct transcriptional activities of pea (*Pisum sativum*) chloroplasts share immunochemically related functional polypeptides. *Biochem. J.*, 286 (1992) 833-841.
- 176 Lauer, H.H.: Capillary electrophoresis of peptides in free solution. In: Rollema, H., Westerink, B.H.C. and Drijfhout, W.J. (Editors), *Monit. Mol. Neurosci., Proc. Int. Conf. In Vivo Methods*, 5th 1991, Univ. Cent. Pharm., Groningen, 1991, p. 121; *C.A.*, 117 (1992) 107649e.
- 177 Matsubara, N. and Terabe, S.: Separation of closely related peptides by capillary electrophoresis with a nonionic surfactant. *Chromatographia*, 34 (1992) 493-496.
- 178 Moseley, M.A., Jorgenson, J.W., Shabanowitz, J., Hunt, D.F. and Tomer, K.B.: Optimization of capillary zone electrophoresis/electrospray ionization parameters for the mass spectrometry and tandem mass spectrometry analysis of peptides. *J. Am. Soc. Mass Spectrom.*, 3 (1992) 289-300; *C.A.*, 117 (1992) 103302p.
- 179 Nakao, M., Mizutani, T., Bhakta, A., Ribarac-Stepic, N. and Moudgil, V.K.: Phosphorylation of chicken oviduct progesterone receptor by cAMP-dependent protein kinase. *Arch. Biochem. Biophys.*, 298 (1992) 340-348.
- 180 Sasaoki, K., Hiroshima, T., Kusumoto, S. and Nishi, K.: Deamidation at asparagine-88 in recombinant human interleukin 2. *Chem. Pharm. Bull.*, 40 (1992) 976-980.
- 181 Schwer, C. and Lottspeich, F.: Analytical and micropreparative separation of peptides by capillary zone electrophoresis using discontinuous buffer systems. *J. Chromatogr.*, 623 (1992) 345-355.
- 182 Soucheleau, J. and Denoroy, L.: Determination of vasoactive intestinal peptide in rat brain by high-performance capillary electrophoresis. *J. Chromatogr.*, 608 (1992) 181-188.
- 183 Tsuji, K., Baczynskyj, L. and Bronson, G.E.: Capillary electrophoresis - electrospray mass spectrometry for the analysis of recombinant bovine and porcine somatotropins. *Anal. Chem.*, 64 (1992) 1864-1870.
- 184 Young, P.M., Astephen, N.E. and Wheat, T.E.: Effects of pH and buffer composition on peptide separations by high performance liquid chromatography and capillary electrophoresis. *LC-GC*, 10 (1992) 26-32; *C.A.*, 117 (1992) 22795z.

- 185 Zhao, J.Y., Waldron, K.C., Miller, J., Zhang, J.Z., Harke, H. and Dovichi, N.J.: Attachment of a single fluorescent label to peptides for determination by capillary zone electrophoresis. *J. Chromatogr.*, 608 (1992) 239-242.
- See also 68, 191, 396, 486, 542, 707, 737.
- 18c. *Elucidation of structure of proteins and enzymes*
- 186 Aebersold, R., Patterson, S.D. and Hess, D.: Strategies for the isolation of peptides from low-abundance proteins for internal sequence analysis. In: Angeletti, R.H. (Editor), *Tech. Protein Chem. 3, [Pap. Annu. Symp. Protein Soc.]*, 5th 1991, Academic, San Diego, 1992, pp. 87-96; C.A., 117 (1992) 65890u.
- 187 Hall, S.C., Schindler, P.A., Masiarz, F.R. and Burlingame, A.L.: Tandem mass spectrometric sequencing of proteins isolated from 1- and 2-dimensional polyacrylamide gel electrophoresis. In: Angeletti, R.H. (Editor), *Tech. Protein Chem. 3, [Pap. Annu. Symp. Protein Soc.]*, 5th 1991, Academic, San Diego, 1992, pp. 525-532; C.A., 117 (1992) 65846j.
- 188 Hori, K. and Morita, F.: Actin-actin contact: inhibition of actin-polymerization by subdomain 4 peptide fragments. *J. Biochem. (Tokyo)*, 112 (1992) 401-408.
- 189 Hughes, G.J., Frutiger, S., Paquet, N., Ravier, F., Pasquali, C., Sanchez, J.-C., James, R., Tissot, J.-D., Bjellqvist, B. and Hochstrasser, D.F.: Plasma protein map: an update by microsequencing. *Electrophoresis (Weinheim)*, 13 (1992) 707-714.
- 190 Isoe, T., Naito, M., Shirai, A., Hirai, R. and Tsuruo, T.: Inhibition of different steps of the ubiquitin system by cisplatin and aciarubicin. *Biochim. Biophys. Acta*, 1117 (1992) 131-135.
- 191 Khropycheva, R.P., Marinchenko, G.V. and Fedosimov, V.A.: (Prolactin hydrolysis by mitochondrial serine proteinase from mammary gland secretory cells). *Biokhimiya (Moscow)*, 57 (1992) 945-955.
- 192 Kittelberger, R., Neale, T.J., Francky, K.T., Greenhill, N. and Gibson, G.J.: Cleavage of type VIII collagen by human neutrophil elastase. *Biochim. Biophys. Acta*, 1139 (1992) 295-299.
- 193 Moritz, R.L., Ward, L.D. and Simpson, R.J.: High-sensitivity peptide mapping utilizing reversed-phase microbore and microcolumn liquid chromatography. In: Angeletti, R.H. (Editor), *Tech. Protein Chem. 3, [Pap. Annu. Symp. Protein Soc.]*, 5th 1991, Academic, San Diego, 1992, pp. 97-106; C.A., 117 (1992) 66002t.
- 194 Nokihara, K., Morita, N. and Kuriki, T.: Applications of an automated apparatus for two-dimensional electrophoresis, Model TEP-1, for microsequence analyses of proteins. *Electrophoresis (Weinheim)*, 13 (1992) 701-707.
- 195 Seto-Young, D., Monk, B.C. and Perlin, D.S.: Assessing hydrophobic regions of the plasma membrane H⁺-ATPase from *Saccharomyces cerevisiae*. *Biochim. Biophys. Acta*, 1102 (1992) 213-219.
- 196 Smith, A., Kenny, J.W. and Ohms, J.I.: Micropreparative separation of tryptic digests by capillary electrophoresis and characterization by protein sequencing. In: Angeletti, R.H. (Editor), *Tech. Protein Chem. 3, [Pap. Annu. Symp. Protein Soc.]*, 5th 1991, Academic, San Diego, 1992, pp. 113-120; C.A., 117 (1992) 65891v.
- 197 Van de Goor, A.A.A.M., Wanders, B.J. and Henzel, W.J.: High-sensitivity peptide mapping with CE using laser-induced fluorescence detection. *Am. Biotechnol. Lab.*, 10 (1992) 16; C.A., 117 (1992) 167014d.
- See also 234, 274, 277, 278, 313, 318, 332, 343, 344, 347, 348, 359, 376, 380, 382, 396, 422, 444, 474, 477, 549, 550, 554.
19. PROTEINS
- 19a. *General techniques*
- 198 Baker, C.S., Corbett, J.M., May, A.J., Yacoub, M.H. and Dunn, M.J.: A human myocardial two-dimensional electrophoresis database: protein characterisation by microsequencing and immunoblotting. *Electrophoresis (Weinheim)*, 13 (1992) 723-726.
- 199 Burggraf, D., Andersson, K., Eckerskorn, C. and Lottspeich, F.: Towards a two-dimensional database of common human proteins. *Electrophoresis (Weinheim)*, 13 (1992) 729-732.
- 200 Chen, F.-T.: High-resolution protein analysis by automated capillary electrophoresis. *Clin. Chem. (Winston-Salem)*, 38 (1992) 1651-1653.
- 201 Chu, Y.H., Avila, L.Z., Biebuyck, H.A. and Whitesides, G.M.: Use of affinity capillary electrophoresis to measure binding constants of ligands to proteins. *J. Med. Chem.*, 35 (1992) 2915-2917; C.A., 117 (1992) 65899d.
- 202 Clark, W.M.: Electrophoresis-enhanced extractive separation. *CHEMTECH*, 22 (1992) 425-429; C.A., 117 (1992) 127642a.
- 203 Foret, F., Szoko, E. and Karger, B.L.: On-column transient and coupled column isotachophoretic preconcentration of protein samples in capillary zone electrophoresis. *J. Chromatogr.*, 608 (1992) 3-12.
- 204 Golias, T.L.: Immunofixation electrophoresis control system. U.S. US 5,137,614 (Cl. 204-199R; GO1N27/26), 11 Aug. 1992, Appl. 231,032, 11 Aug. 1988; 6 pp.; C.A., 117 (1992) 146729a.
- 205 Guzman, N.A., Moschera, J., Iqbal, K. and Malick, A.W.: Effect of buffer constituents on the determination of therapeutic proteins by capillary electrophoresis. *J. Chromatogr.*, 608 (1992) 197-204.
- 206 Heukeshoven, J. and Dernick, R.: Native horizontal ultrathin polyacrylamide gel electrophoresis of proteins under basic and acidic conditions. *Electrophoresis (Weinheim)*, 13 (1992) 654-659.
- 207 Kang, S., Zou, Z. and Qiao, X.: (The qualitative and quantitative analyses of protein with a novel fluorescence agent (EPQS) by fluorescent scanning on solid substrate). *Huaxi Yike Daxue Xuebao*, 22 (1991) 262-265; C.A., 117 (1992) 22664f.
- 208 Kraak, J.C., Busch, S. and Poppe, H.: Study of protein-drug binding using capillary zone electrophoresis. *J. Chromatogr.*, 608 (1992) 257-264.
- 209 Meier, T., Arni, S., Malarkannan, S., Poincelet, M. and Hoessli, D.: Immunodetection of biotinylated lymphocyte-surface proteins by enhanced chemiluminescence: a nonradioactive method for cell-surface protein analysis. *Anal. Biochem.*, 204 (1992) 220-226.

- 210 Mohan, S.B.: Determination of purity and yield. *Methods Mol. Biol. (Totowa)*, 11(Pract. Protein Chromatogr.) (1992) 307-323; C.A., 117 (1992) 167114m.
- 211 Nakazawa, M. and Manabe, K.: The direct hydrolysis of proteins containing tryptophan on polyvinylidene difluoride membranes by mercaptoethanesulfonic acid in the vapor phase. *Anal. Biochem.*, 206 (1992) 105-108.
- 212 Nelson, T.J.: Rapid isoelectric focusing of proteins in hydrolytically stable capillaries. *J. Chromatogr.*, 623 (1992) 357-365.
- 213 Pande, P.G., Nellore, R.V. and Bhagat, H.R.: Optimization and validation of analytical conditions for bovine serum albumin using capillary electrophoresis. *Anal. Biochem.*, 204 (1992) 103-106.
- 214 Praxmayer, C., Murach, K.-F., Baumgartner, B., Aberger, F., Schlegel, E. and Illmensee, K.: Protein synthesis in murine organs during postimplantation development detected by two-dimensional gel electrophoresis. *Electrophoresis (Weinheim)*, 13 (1992) 720-722.
- 215 Rohde, M.F., Stoney, K.S. and Wiktorowicz, J.E.: Estimation of protein isoelectric points by capillary electrophoresis using MICRO-COAT. In: Angeletti, R.H. (Editor), *Tech. Protein Chem. 3, [Pap. Annu. Symp. Protein Soc.]*, 5th 1991, Academic, San Diego, 1992, pp. 121-127; C.A., 117 (1992) 65892w.
- 216 Sanchez, J.-C., Ravier, F., Pasquali, C., Frutiger, S., Paquet, N., Bjellqvist, B., Hochstrasser, D.F. and Hughes, G.J.: Improving the detection of proteins after transfer to polyvinylidene difluoride membranes. *Electrophoresis (Weinheim)*, 13 (1992) 715-717.
- 217 Takagi, T.: (Electrophoretic light scattering. A new apparatus applicable to proteins of cells). *Seibutsu Butsuri Kagaku*, 36 (1992) 21-31; C.A., 117 (1992) 86030d - a review with 22 refs.
- 218 Thompson, D. and Larson, G.: Western blots using stained protein gels. *BioTechniques*, 12 (1992) 656-658; C.A., 117 (1992) 43928k.
- 219 Wu, D. and Regnier, F.E.: Sodium dodecyl sulfate-capillary gel electrophoresis of proteins using non-cross-linked polyacrylamide. *J. Chromatogr.*, 608 (1992) 349-356.
- 220 Wu, J. and Pawliszyn, J.: Application of capillary isoelectric focusing with universal concentration gradient detector to the analysis of protein samples. *J. Chromatogr.*, 608 (1992) 121-130.
- 221 Wu, S.L.: Rapid HPLC and capillary electrophoresis: a perspective of protein analysis from analytical biotechnology. *LC-GC*, 10 (1992) 430-434; C.A., 117 (1992) 86018f - a review with 20 refs.
- See also 3, 10, 24, 44, 68, 73, 83, 84, 88, 90, 165, 245.
- 19b. *Proteins of cells, viruses and subcellular particles*
- 222 Bertrand, C., Dumoulin, R., Divry, P., Matheiu, M. and Vianey-Saban, C.: Purification of electron transfer flavoprotein from pig liver mitochondria and its application to the diagnosis of deficiencies of acyl-CoA dehydrogenases in human fibroblasts. *Clin. Chim. Acta*, 210 (1992) 75-91.
- 223 Bybee, A. and Thomas, N.S.B.: The synthesis of p58^{cyclin A} and the phosphorylation of p34^{cdc2} are inhibited in human lymphoid cells arrested in G₁ by α -interferon. *Biochim. Biophys. Acta*, 1137 (1992) 73-76.
- 224 Caldentey, J. and Bamford, D.H.: The lytic enzyme of the *Pseudomonas* phage $\phi 6$. Purification and biochemical characterization. *Biochim. Biophys. Acta*, 1159 (1992) 44-50.
- 225 Celerier, J., Cibert, C., Barré-Sinoussi, F., Ammar, A. and Jamin, C.: Purification and identification of a CD25 expression inhibitory protein from cell-free supernatants of chronically HIV-infected promonocytic cells. *Biochim. Biophys. Acta*, 1139 (1992) 300-306.
- 226 Fisher, B., Kraft, P., Hahn, G.M. and Anderson, R.L.: Thermotolerance in the absence of induced heat shock proteins in a murine lymphoma. *Cancer Res.*, 52 (1992) 2954-2861.
- 227 Fisher, R.J., Koizumi, S., Kondoh, A., Mariano, J.M., Mavrothalassitis, G., Bhat, N.K. and Papas, T.S.: Human ETS1 oncoprotein. Purification, isoforms, -SH modification, and DNA sequence-specific binding. *J. Biol. Chem.*, 267 (1992) 17957-17965.
- 228 Gold, J.C. and Phillips, M.C.: Effects of membrane lipids and -proteins and cytoskeletal proteins on the kinetics of cholesterol exchange between high density lipoprotein and human red blood cells, ghosts and microvesicles. *Biochim. Biophys. Acta*, 1111 (1992) 103-110.
- 229 Kareem, B.N., Rokkones, E., Hogset, A., Holmgren, E. and Gautvik, K.M.: A method for the evaluation of the efficiency of signal sequences for secretion and correct N-terminal processing of human parathyroid hormone produced in *Escherichia coli*. *Anal. Biochem.*, 204 (1992) 26-33.
- 230 Kaufmann, S.H.: Expression of nuclear envelope lamins A and C in human myeloid leukemias. *Cancer Res.*, 52 (1992) 2847-2853.
- 231 Kovarova, H., Stulik, J., Macela, A., Lefkovits, I. and Skrabkova, Z.: Using two-dimensional gel electrophoresis to study immune response against intracellular bacterial infection. *Electrophoresis (Weinheim)*, 13 (1992) 741-742.
- 232 Kramer, W., Nicol, S.-B., Girbig, F., Gutjahr, U., Kowaleski, S. and Fasold, H.: Characterization and chemical modification of the Na⁺-dependent bile-acid transport system in brush-border membrane vesicles from rabbit ileum. *Biochim. Biophys. Acta*, 1111 (1992) 93-102.
- 233 Lavergne, J.-P., Reboud, A.-M., Sontag, B., Guillot, D. and Reboud, J.-P.: Binding of GDP to a ribosomal protein after elongation factor-2 dependent GTP hydrolysis. *Biochim. Biophys. Acta*, 1132 (1992) 284-289.
- 234 Lee, S., Wulffkuhle, J.D. and Otto, J.J.: Vinculin binding site mapped on talin with an anti-idiotypic antibody. *J. Biol. Chem.*, 267 (1992) 16355-16358.
- 235 Ochi, K.: Polyacrylamide gel electrophoresis analysis of ribosomal protein: a new approach for actinomycete taxonomy. *Gene*, 115 (1992) 261-265; C.A., 117 (1992) 107653b.
- 236 Ohtsuki, T., Suzu, S., Nagata, N. and Motoyoshi, K.: A human osteoblastic cell line, MG-63, produces two molecular types of macrophage-colony-stimulating factor. *Biochim. Biophys. Acta*, 1136 (1992) 297-301.
- 237 Okazaki, I., Himeno, M., Ezaki, J., Ishikawa, T. and Kato, K.: Purification and characterization of an 85 kDa sialoglycoprotein in rat liver lysosomal membranes. *J. Biochem. (Tokyo)*, 111 (1992) 763-769.
- 238 Oldfield, S. and Proud, C.G.: Purification, phosphorylation and control of the guanine-nucleotide-exchange factor from rabbit reticulocyte lysates. *Eur. J. Biochem.*, 208 (1992) 73-81.

- 239 Park, S., Liu, X., Pawson, T. and Jove, R.: Activated Src tyrosine kinase phosphorylates Tyr-457 of bovine GTPase-activating protein (GAP) *in vitro* and the corresponding residue of rat GAP *in vivo*. *J. Biol. Chem.*, 267 (1992) 17194-17200.
- 240 Slobodskaya, O.R. and Gorbalenya, A.E.: (Linker-inserted mutagenesis of the poliovirus 2A protease and its substrate simultaneously expressed in *Escherichia coli* cells). *Biokhimiya (Moscow)*, 57 (1992) 1057-1064.
- 241 Subramaniam, V.N., bin Mohd. Yusoff, A.R., Wong, S.H., Lim, G.B., Chew, M. and Hong, W.: Biochemical fractionation and characterization of proteins from Golgi-enriched membranes. *J. Biol. Chem.*, 267 (1992) 12016-12021.
- 242 Toku, S. and Tanaka, T.: The primary structure of chicken ribosomal protein L37a. *Biochim. Biophys. Acta*, 1132 (1992) 88-90.
- 243 Tremblay, L., Gingras, D., Boivin, D. and Béliveau, R.: Tyrosine protein kinase activity in renal brush-border membranes. *Biochim. Biophys. Acta*, 1108 (1992) 183-189.
- 244 Winters, D.K. and Cederbaum, A.I.: Time course characterization of the induction of cytochrome P-450 2E1 by pyrazole and 4-methylpyrazole. *Biochim. Biophys. Acta*, 1117 (1992) 15-24.
- 245 Yıldız, E., Grübler, G., Hörger, S., Zimmermann, H., Echner, H., Stoeva, S. and Voelter, W.: Application of capillary electrophoresis in peptide research. *Electrophoresis (Weinheim)*, 13 (1992) 683-686.
- See also 190, 209, 217, 287, 296, 303, 333, 335, 337, 346, 352, 389, 419, 436, 444, 460, 603.
- 19c. *Proteins synthesized by genetic manipulation, monoclonal antibodies*
- 246 Cheng, L., Hirst, K. and Piper, P.W.: Authentic temperature-regulation of a heat shock gene inserted into yeast on a high copy number vector. Influences of overexpression of HSP90 protein on high temperature growth and thermotolerance. *Biochim. Biophys. Acta*, 1132 (1992) 26-34.
- 247 Fujita, S., Maeda, K. and Maeda, Y.: Expression in *Escherichia coli* and a functional study of a β -troponin T 25 kDa fragment of rabbit skeletal muscle. *J. Biochem. (Tokyo)*, 112 (1992) 306-308.
- 248 Gordon, M., Chokshi, H. and Alpers, D.H.: *In vitro* expression and secretion of functional mammalian intrinsic factor using recombinant baculovirus. *Biochim. Biophys. Acta*, 1132 (1992) 276-283.
- 249 Kruse, E. and Kloppstech, K.: Integration of early light-inducible proteins into isolated thylakoid membranes. *Eur. J. Biochem.*, 208 (1992) 195-202.
- 250 Kurata, S.-i. and Matsumoto, M.: Expression of heme oxygenase and its RNA in mouse liver after injection of heme and splenectomy. *Biochim. Biophys. Acta*, 1132 (1992) 255-258.
- 251 Li, X.-K., Lijnen, H.R., Nelles, L., Hu, M.-H. and Collen, D.: Biochemical properties of recombinant mutants of nonglycosylated single chain urokinase-type plasminogen activator. *Biochim. Biophys. Acta*, 1159 (1992) 37-43.
- 252 Macheroux, P., Mulrooney, S.B., Williams, C.H., Jr. and Massey, V.: Direct expression of active spinach glycolate oxidase in *Escherichia coli*. *Biochim. Biophys. Acta*, 1132 (1992) 11-16.
- 253 White, S., Miller, K., Hopkins, C. and Trowbridge, I.S.: Monoclonal antibodies against defined epitopes of the human transferrin receptor cytoplasmic tail. *Biochim. Biophys. Acta*, 1136 (1992) 28-34.
- See also 311, 499, 695, 696.
- 19d. *Microbial and plant proteins*
- 254 Beers, E.P., Moreno, T.N. and Callis, J.: Subcellular localization of ubiquitin and ubiquitinated proteins in *Arabidopsis thaliana*. *J. Biol. Chem.*, 267 (1992) 15432-15439.
- 255 Bietz, J.A. and Simpson, D.G.: Electrophoresis and chromatography of wheat proteins: available methods, and procedures for statistical evaluation of the data. *J. Chromatogr.*, 624 (1992) 53-80 - a review with 192 refs.
- 256 Black, J.A. and Waggamon, K.A.: Wheat gliadin: digital imaging and database construction using a 4-band reference system of agarose isoelectric focusing patterns. *Electrophoresis (Weinheim)*, 13 (1992) 800-802.
- 257 Browning, K.S., Webster, C., Roberts, J.K.M. and Ravel, J.M.: Identification of an isozyme form of protein synthesis initiation factor 4F in plants. *J. Biol. Chem.*, 267 (1992) 10096-10100.
- 258 Caswell, R., Webster, C. and Busby, S.: Studies on the binding of the *Escherichia coli* MelR transcription activator protein to operator sequences at the MelAB promoter. *Biochem. J.*, 287 (1992) 501-508.
- 259 Caswell, R., Williams, J., Lyddiatt, A. and Busby, S.: Overexpression, purification and characterization of the *Escherichia coli* MelR transcription activator protein. *Biochem. J.*, 287 (1992) 493-499.
- 260 Ellis, C.M., Ford, R.C. and Holzenburg, A.: Detergent sensitivity of the tonoplast H⁺-ATPase and its purification from *Beta vulgaris*. *Biochim. Biophys. Acta*, 1136 (1992) 319-326.
- 261 Furst, V., Smith, I. and Holton, J.: Computer analysis of sulfur-35-labeled bacterial proteins. *J. Microbiol. Methods*, 15 (1992) 327-335; *C.A.*, 117 (1992) 127757s.
- 262 Höhne-Zell, B., Knoll, G., Riedel-Gras, U., Hofer, W. and Plattner, H.: A cortical phosphoprotein ('PP63') sensitive to exocytosis triggering in *Paramecium* cells. Immunolocalization and quenched-flow correlation of time course of dephosphorylation with membrane fusion. *Biochem. J.*, 286 (1992) 843-849.
- 263 Klier, H. and Lottspeich, F.: Detection of the hypusine-containing protein (HP=elF-5A) in crude yeast extracts by two-dimensional Western blots. *Electrophoresis (Weinheim)*, 13 (1992) 732-735.
- 264 Krasilnikov, O.V., Muratkhodjaev, J.N. and Zitzer, A.O.: The mode of action of *Vibrio cholerae* cytotoxin. The influences on both erythrocytes and planar lipid bilayers. *Biochim. Biophys. Acta*, 1111 (1992) 7-16.
- 265 Metakovskii, E.V. and Novoselskaya, A.Yu.: Gliadin allele identification in common wheat. I. Methodological aspects of the analysis of gliadin patterns by one-dimensional polyacrylamide gel electrophoresis. *J. Genet. Breed.*, 45 (1991) 317-323; *C.A.*, 117 (1992) 167019j.
- 266 Neucere, J.N.: Electrophoretic analysis of cationic proteins extracted from aflatoxin-resistant/susceptible varieties of corn. *J. Agric. Food Chem.*, 40 (1992) 1422-1424.

- 267 Nikolaev, I.V., Vavilova, E.A. and Vinetsky, Yu.P.: (Molecular structure of the lac-phenotype in the filamentous fungus *Penicillium canescens*: the existence of secreted and two cytosolic β -galactosidases). *Biokhimiya (Moscow)*, 57 (1992) 873-879.
- 268 Posch, A., van den Berg, B.M., Postel, W. and Görg, A.: Genetic variability of pepper (*Capsicum annum* L.) seed proteins studied by 2-D electrophoresis with immobilized pH gradients. *Electrophoresis (Weinheim)*, 13 (1992) 774-777.
- 269 Raimo, G., Masullo, M., Parente, A., Dello Russo, A. and Bocchini, V.: Molecular, functional and structural properties of an archaeobacterial elongation factor 2. *Biochim. Biophys. Acta*, 1132 (1992) 127-132.
- 270 Rajapakse, D.P., Imai, T. and Ishige, T.: Analysis of potato microtuber proteins by sodium dodecylsulfate polyacrylamide gel electrophoresis. *Potato Res.*, 34 (1991) 285-293; *C.A.*, 117 (1992) 24843n.
- 271 Rastogi, N., Labrousse, V. and Barreau, C.: A rapid microbead method for breaking pathogenic mycobacteria: application in SDS-PAGE and Western blot analysis. *Curr. Microbiol.*, 24 (1992) 311-317; *C.A.*, 117 (1992) 107748m.
- 272 Ruban, A.V. and Horton, P.: Mechanism of Δ pH-dependent dissipation of absorbed excitation energy by photosynthetic membranes. I. Spectroscopic analysis of isolated light-harvesting complexes. *Biochim. Biophys. Acta*, 1102 (1992) 30-38.
- 273 Rücker, W. and Maier, R.: Stress effects on hormonal growth factors in tobacco tissues indicated by special changes in the isoelectric peroxidase patterns. *Electrophoresis (Weinheim)*, 13 (1992) 798-799.
- 274 Segura-Nieto, M., Vazquez-Sanchez, N., Rubio-Velazquez, H., Olguin-Martinez, L.E., Rodriguez-Nester, C.E. and Herrera-Estrella, L.: Characterization of amaranth (*Amaranthus hypochondriacus* L.) seed proteins. *J. Agric. Food Chem.*, 40 (1992) 1553-1558.
- 275 Shokraii, E.H. and Esen, A.: Characterization of disulfide-linked forms of seed proteins in peanuts (*Arachis hypogaea* L.) *J. Agric. Food Chem.*, 40 (1992) 1491-1495.
- 276 Sjöström, H., Friis, S.U., Norén, O. and Anthonson, D.: Purification and characterisation of antigenic gliadins in coeliac disease. *Clin. Chim. Acta*, 207 (1992) 227-237.
- 277 Takada, S., Sakakibara, H. and Kamiya, R.: Three-headed outer arm dynein from *Chlamydomonas* that can functionally combine with outer-arm-missing axonemes. *J. Biochem. (Tokyo)*, 111 (1992) 758-762.
- 278 Tao, H.P., Adalsteins, A.E. and Kasarda, D.D.: Intermolecular disulfide bonds link specific high-molecular-weight glutenin subunits in wheat endosperm. *Biochim. Biophys. Acta*, 1159 (1992) 13-21.
- 279 Tokishita, S.-i., Kojima, A. and Mizuno, T.: Transmembrane signal transduction and osmoregulation in *Escherichia coli*: functional importance of the transmembrane regions of membrane-located protein kinase, EnvZ. *J. Biochem. (Tokyo)*, 111 (1992) 707-713.
- 280 Tokunaga, M., Kawamura, A. and Kohno, K.: Purification and characterization of BiP/Kar2 protein from *Saccharomyces cerevisiae*. *J. Biol. Chem.*, 267 (1992) 17553-17559.
- 281 Van der Staay, G.W.M., Brouwer, A., Baard, R.L., van Mourik, F. and Matthijs, H.C.P.: Separation of Photosystem I and II from the oxylchlorobacterium (prochlorophyte) *Prochlorothrix hollandica* and association of chlorophyll b binding antennae with Photosystem II. *Biochim. Biophys. Acta*, 1102 (1992) 230-228.
- 282 Venugopal, M.G., Wolfersberger, M.G. and Wallace, B.A.: Effects of pH on conformational properties related to the toxicity of *Bacillus thuringiensis* δ -endotoxin. *Biochim. Biophys. Acta*, 1159 (1992) 185-192.
- 283 Waters, E.J., Wallace, W. and Williams, P.J.: Identification of heat-unstable wine proteins and their resistance to peptidases. *J. Agric. Food Chem.*, 40 (1992) 1514-1519.
- 284 Weiss, W., Postel, W. and Görg, A.: Qualitative and quantitative changes in barley seed protein patterns during the malting process analyzed by sodium dodecyl sulfate-polyacrylamide gel electrophoresis with respect to malting quality. *Electrophoresis (Weinheim)*, 13 (1992) 787-797.
- 285 Zucker, B., Rehak, E. and Knop, C.: (Characterization of *Bordetella bronchiseptica* strains by polypeptide and lipopolysaccharide patterns analyses in SDS-PAGE). *Monatsh. Veterinaarmed.*, 47 (1992) 85-88; *C.A.*, 117 (1992) 167245e.

See also 131, 195, 229, 455, 715.

19e. *Proteins of blood, serum and blood cells*

- 286 Adrian, G.S., Herbert, D.C., Robinson, L.A.K., Walter, C.A., Buchanan, J.M., Adrian, E.K., Weaker, F.J., Eddy, C.A., Yang, F. and Bowman, B.H.: Expression of a human chimeric transferrin gene in senescent transgenic mice reflects the decrease of transferrin levels in aging humans. *Biochim. Biophys. Acta*, 1132 (1992) 168-176.
- 287 Aliev, D.A., Suleimanov, S.Yu., Guseinova, I.M., Asadov, A.A. and Ismailov, M.A.: (The effects of specific translation inhibitors on the polypeptide composition and spectral characteristics of wheat chloroplasts). *Biokhimiya (Moscow)*, 57 (1992) 679-686.
- 288 Anand, B. and Chou, I.: Microtubules and microtubule-associated proteins in resting and mitogenically activated normal human peripheral blood T cells. *J. Biol. Chem.*, 267 (1992) 10716-10722.
- 289 Back, S.A. and Alhadeff, J.A.: Differential isoform profiles of α_2 -macroglobulin from plasma of patients with chronic-progressive or relapsing-remitting multiple sclerosis. *Clin. Chim. Acta*, 211 (1992) 27-36.
- 290 Baillod, P., Affolter, B., Kurt, G.H. and Pflugshaupt, R.: Multimeric analysis of von Willebrand factor by vertical sodium dodecyl sulfate agarose gel electrophoresis, vacuum blotting technology, and sensitive visualization by alkaline phosphatase anti-alkaline phosphatase complex. *Thromb. Res.*, 66 (1992) 745-755; *C.A.*, 117 (1992) 127649h.
- 291 Barbi, G., Keil, L.B., Gaito, A.D. and DeBari, V.A.: Spectrotype distributions of circulating IgG from patients with systemic lupus erythematosus. *Clin. Chem. (Winston-Salem)*, 38 (1992) 1988-1994.
- 292 Bilinski, H., Horvath, L. and Trbojevic-Cepe, M.: Precipitation and characterization of an aluminosilicate from $AlCl_3-Na_2SiO_3-HCl$ in serum, of interest for Alzheimer disease. *Clin. Chem. (Winston-Salem)*, 38 (1992) 2019-2024.
- 293 Bini, L., Magi, B., Cellesi, C., Rossolini, A. and Pallini, V.: Two-dimensional electrophoresis analysis of human serum proteins during the acute-phase response. *Electrophoresis (Weinheim)*, 13 (1992) 743-746.

- 294 Breborowicz, J. and Gryska, K.: Determination of microheterogeneity of human α_1 -fetoprotein by affinity electrophoresis and by batch technique. In: Breborowicz, J. and Mackiewicz, A. (Editors), *Affinity Electrophor.*, CRC, Boca Raton, 1992, pp. 71-79; C.A., 117 (1992) 167106k.
- 295 Cameron, B.M., Merrill, C.R., Creed, G.J. and Vanderputten, D.: Electrophoretic and immunochemical methods for the diagnosis of chronic lower back and cervical pain. *PCT Int. Appl. WO 92 10,760* (Cl. GO1N33/68), 25 Jun. 1992, US Appl. 620,104, 30 Nov. 1990, 66 pp.; C.A., 117 (1992) 127812f.
- 296 Das, A.K., Kundu, M., Chakrabarti, P. and Basu, J.: Fatty acylation of a 55 kDa membrane protein of human erythrocytes. *Biochim. Biophys. Acta*, 1108 (1992) 128-132.
- 297 Deuticke, B., Lütke-meier, P. and Poser, B.: Tellurite-induced damage of the erythrocyte membrane. Manifestations and mechanisms. *Biochim. Biophys. Acta*, 1109 (1992) 97-107.
- 298 Eckerskorn, C., Strupat, K., Karas, M., Hillenkamp, F. and Lottspeich, F.: Mass spectrometric analysis of blotted proteins after gel electrophoretic separation by matrix-assisted laser desorption/ionization. *Electrophoresis (Weinheim)*, 13 (1992) 664-665.
- 299 Ferrari, P., Torielli, L., Salardi, S., Rizzo, A. and Bianchi, G.: Na⁺/K⁺/Cl⁻ cotransport in resealed ghosts from erythrocytes of the Milan hypertensive rats. *Biochim. Biophys. Acta*, 1111 (1992) 111-119.
- 300 Fried, R., Wiederkehr, F. and Vonderschmitt, D.J.: Immune complexes in blood: a new strategy for the analysis of their antigen portion. *Electrophoresis (Weinheim)*, 13 (1992) 718-719.
- 301 Fujii, T., Mimori, T., Hama, N., Suwa, A., Akizuki, M., Tojo, T. and Homma, M.: Characterization of autoantibodies that recognize U4/U6 small ribonucleoprotein particles in serum from a patient with primary Sjögren's syndrome. *J. Biol. Chem.*, 267 (1992) 16412-16416.
- 302 Grinenko, T.V., Skomorovskaya, E.V., Kudinov, S.A. and Zolotareva, K.N.: (Some peculiarities of plasminogen GLu- and Lys-forms interaction with native and partially hydrolyzed fibrin). *Biokhimiya (Moscow)*, 57 (1992) 728-738.
- 303 Itoh, K., Hara, T., Yamada, F. and Shibata, N.: Diphosphorylation of platelet myosin *ex vivo* in the initial phase of activation by thrombin. *Biochim. Biophys. Acta*, 1136 (1992) 52-56.
- 304 Kelleher, P.C., Toftager-Larsen, K. and Walters, C.J.P.: Comparison of concanavalin A-affinity electrophoresis and concanavalin A-affinity chromatography in the analysis of amniotic fluid α_1 -fetoprotein microheterogeneity. In: Breborowicz, J. and Mackiewicz, A. (Editors), *Affinity Electrophor.*, CRC, Boca Raton, 1992, pp. 89-95; C.A., 117 (1992) 167107m.
- 305 Khatler, N., Matson, R.S. and Ngo, T.: Utilization of a synthetic affinity ligand for immunoglobulin purification. *Int. Lab.*, 22, No. 9 (1992) P14-P18.
- 306 Knippel, E., Hein, J., Nazir, S., Schuett, W., Marlow, B., Fairhurst, D., Goetz, P., Bauer, I., Bittorf, T. et al.: Adsorption of serum components onto polystyrene particles. A diagnostic method for cystic fibrosis gene carriers? *Wiss. Z. Univ. Rostock, Naturwiss. Reihe*, 39 (1990) 76-102; C.A., 117 (1992) 169042k.
- 307 Kondo, N. and Kondo, J.: Identification of novel blood proteins specific for mammalian hibernation. *J. Biol. Chem.*, 267 (1992) 473-478.
- 308 Lepesheva, G.I. and Martsev, S.P.: (Modification of the rabbit IgG hinge disulfide and study of its interaction with the polyvalent antigen ferritin, protein A, and anti-IgG). *Biokhimiya (Moscow)*, 57 (1992) 1089-1099.
- 309 Lundy, F.T. and Wisdom, G.B.: The determination of asialoglycoforms of serum glycoproteins by lectin blotting with *Ricinus communis* agglutinin. *Clin. Chim. Acta*, 205 (1992) 187-195.
- 310 Mittermüller, J. and Weidinger, S.: Genetic study of orosomucoid by isoelectric focusing and immunoprinting in patients with carcinoma. *Electrophoresis (Weinheim)*, 13 (1992) 785-786.
- 311 Miura, Y., Tokunaga, F., Miyata, T., Moriyasu, M., Yoshikawa, K. and Iwanaga, S.: Preparation and properties of monoclonal antibodies against lipopolysaccharide-sensitive serine protease zymogen, factor C, from horseshoe crab (*Tachyplesus tridentatus*) hemocytes. *J. Biochem. (Tokyo)*, 112 (1992) 476-481.
- 312 Miyakoshi, H., Sugimoto, M. and Honda, H.: Method and reagents for detecting HTLV-I virus antibody in blood by electrophoresis-immunoassay. *Jpn. Kokai Tokkyo Koho JP 04 86,558* [92 86,558] (Cl. GO1N33/561), 19 Mar. 1992, Appl. 90/201,860, 30 Jul. 1990; 5 pp.; C.A., 117 (1992) 24680g.
- 313 Osawa, M., Sabbatini, A.R.M., Erukhimov, J., Werner, P.A.M. and Galbraith, R.M.: Binding of monoclonal antibody E12 to Ge globulin (vitamin D-binding protein) is inhibited by actin. *Biochim. Biophys. Acta*, 1117 (1992) 271-278.
- 314 Pala, A., di Ruzza, A., Rossetto, G., Moro, M., Strom, R., Padula, F. and Cosmi, E.V.: Distribution of SP₁, immunoreactivity among different plasma proteins: real molecular heterogeneity or adsorption of SP₁-beta to other plasma proteins? *Clin. Chim. Acta*, 207 (1992) 87-97.
- 315 Peach, R.J., Fellowes, A.P., Brennan, S.O. and George, P.M.: Albumin Rugby Park: a truncated albumin variant caused by a G→C splice-site mutation in intron 13. *Biochim. Biophys. Acta*, 1180 (1992) 107-110.
- 316 Petersen, A., Becker, W.-M. and Schlaak, M.: Examination of microheterogeneity in grass pollen allergens. *Electrophoresis (Weinheim)*, 13 (1992) 736-739.
- 317 Putnam, F.W.: (Electrophoretic and structural characterization of genetic variants of human serum albumin). *Seibutsu Butsuri Kagaku*, 36 (1992) 3-19; C.A., 117 (1992) 86029k - a review with 43 refs.
- 318 Rizzo, P., Tinello, C., Punturieri, A. and Taniuchi, H.: A study of hydrogen exchange of monoclonal antibodies: specificity of the antigen-binding induced conformational stabilization. *Biochim. Biophys. Acta*, 1159 (1992) 169-178.
- 319 Schreiber, W.E., Chiang, E. and Tse, S.S.L.: Electrophoresis underestimates the concentration of polyclonal immunoglobulins in serum. *Am. J. Clin. Pathol.*, 97 (1992) 610-613; C.A., 117 (1992) 109592e.
- 320 Sengeløv, H., Nielsen, M.H. and Borregaard, N.: Separation of human neutrophil plasma membrane from intracellular vesicles containing alkaline phosphatase and NADPH oxidase activity by free flow electrophoresis. *J. Biol. Chem.*, 267 (1992) 14912-14917.
- 321 Serdyuk, O.A. and Vasilov, R.G.: Analysis of human major histocompatibility complex antigens by electrophoretic methods with subsequent immunoblotting. *Biomed. Sci. (London)*, 1 (1990) 384-390; C.A., 117 (1992) 109588h.
- 322 Shakoory, A.R. and Aziz, F.: Effect of freezing (-30°C) on the electrophoretic pattern and estimations of blood serum proteins of freshwater fish, *Labeo rohita*. *Pak. J. Zool.*, 23 (1991) 369-372; C.A., 117 (1992) 43871m.
- 323 Sierra, F., Coeytaux, S., Juillerat, M., Ruffieux, C., Gauldie, J. and Guigoz, Y.: Serum T-kininogen levels increase two to four months before death. *J. Biol. Chem.*, 267 (1992) 10665-10669.

- 324 Speicher, D.W., Weglarz, L. and DeSilva, T.M.: Properties of human red cell spectrin heterodimer (side-to-side) assembly and identification of an essential nucleation site. *J. Biol. Chem.*, 267 (1992) 14775-14782.
- 325 Takami, M., Kasuya, I. and Tsunoo, H.: Polymerized albumin receptor on rat liver cells. *J. Biochem. (Tokyo)*, 111 (1992) 714-721.
- 326 Takanashi, M., Sakurai, T. and Tsuchiya, S.: Identification of the carboxymethyllysine residue in the advanced stage of glycosylated human serum albumin. *Chem. Pharm. Bull.*, 40 (1992) 705-708.
- 327 Taketa, K.: Affinity electrophoresis of alpha₁-fetoprotein in the diagnosis and management of cancer patients. In: Breborowicz, J. and Mackiewicz, A. (Editors), *Affinity Electrophoresis*, CRC, Boca Raton, 1992, pp. 199-210; C.A., 117 (1992) 167012b.
- 328 Torres, J.-M., Darracq, N. and Uriel, J.: Membrane proteins from lymphoblastoid cells showing cross-affinity for α -fetoprotein and albumin. Isolation and characterization. *Biochim. Biophys. Acta*, 1159 (1992) 60-66.
- 329 Tulp, A. and Verwoerd, D.: Analysis of members of the Ig-gene superfamily by thermal gradient polyacrylamide gel electrophoresis. *Electrophoresis (Weinheim)*, 13 (1992) 662-664.
- 330 Weidinger, S., Poller, W., Faber, J.-P. and Schwarzfischer, F.: Improved phenotyping of alpha₁-antichymotrypsin (ACT) by isoelectric focusing and immunoprinting: first demonstration of a deficient protein variant in the ACTA system. *Electrophoresis (Weinheim)*, 13 (1992) 778-780.
- 331 Zachariah, K., Kumar, P., Srivastava, D.K. and Rao, K.M.: Quantitation of low levels of IgM by electroimmunoassay. *Anal. Biochem.*, 204 (1992) 405-406.
- See also 51, 60, 121, 228, 253, 413, 419, 431, 444, 544.
- 19f. *Structural and muscle proteins*
- 332 Arai, M., Yada, T., Suzuki, S. and Kimata, K.: Isolation and characterization of type IX collagen-proteoglycan from the Swarm rat chondrosarcoma. *Biochim. Biophys. Acta*, 1117 (1992) 60-70.
- 333 Che, C., Müller, U. and Werdan, K.: Regulation of protein biosynthesis in neonatal rat cardiomyocytes by adrenoceptor-stimulation: investigations with high resolution two-dimensional polyacrylamide gel electrophoresis. *Electrophoresis (Weinheim)*, 13 (1992) 755-756.
- 334 Clark, C.J., Milligan, G., McLellan, A.R. and Connell, J.M.C.: Guanine nucleotide regulatory protein levels and function in spontaneously hypertensive rat vascular smooth-muscle cells. *Biochim. Biophys. Acta*, 1136 (1992) 290-296.
- 335 Endo, T.: SV 40 Large T inhibits myogenic differentiation partially through inducing c-jun. *J. Biochem. (Tokyo)*; 112 (1992) 321-329.
- 336 Hasegawa, Y. and Morita, F.: Role of 17-kDa essential light chain isoforms of aorta smooth muscle myosin. *J. Biochem. (Tokyo)*, 111 (1992) 804-809.
- 337 He, C., Müller, U., Oberthür, W. and Werdan, K.: Application of high resolution two-dimensional polyacrylamide gel electrophoresis of polypeptides from cultured neonatal rat cardiomyocytes: regulation of protein synthesis by catecholamines. *Electrophoresis (Weinheim)*, 13 (1992) 748-754.
- 338 Jungblut, P., Otto, A., Regitz, V., Fleck, E. and Wittmann-Liebold, B.: Identification of human myocard proteins separated by two-dimensional electrophoresis. *Electrophoresis (Weinheim)*, 13 (1992) 739-741.
- 339 Katakami, Y., Katakami, N., Janmey, P.A., Hartwig, J.H. and Stossel, T.P.: Isolation of the phosphatidylinositol 4-monophosphate dissociable high-affinity profilin-actin complex. *Biochim. Biophys. Acta*, 1122 (1992) 123-135.
- 340 Marinkovich, M.P., Lunstrum, G.P. and Burgeson, R.E.: The anchoring filament protein kalinin is synthesized and secreted as a high molecular weight precursor. *J. Biol. Chem.*, 267 (1992) 17900-17906.
- 341 Namba, Y., Ito, M., Zu, Y., Shigesada, K. and Maruyama, K.: Human T cell-L-plastin bundles actin filaments in a calcium-dependent manner. *J. Biochem. (Tokyo)*, 112 (1992) 503-507.
- 342 O'Brien, G.A., Corbett, J.M., Dunn, M.J., Cumming, D.V.E., May, A.J. and Yacoub, M.H.: Electrophoretic analysis of electrically trained skeletal muscle. *Electrophoresis (Weinheim)*, 13 (1992) 726-728.
- 343 Rehbein, H.: Fish species identification by peptide mapping of the myosin heavy chain. *Electrophoresis (Weinheim)*, 13 (1992) 805-806.
- 344 Rotman, E.I., de Jongh, K.S., Florio, V., Lai, Y. and Catterall, W.A.: Specific phosphorylation of a COOH-terminal site on the full-length form of the α 1 subunit of the skeletal muscle calcium channel by cAMP-dependent protein kinase. *J. Biol. Chem.*, 267 (1992) 16100-16105.
- 345 Takahashi, K., Hattori, A., Tatsumi, R. and Takai, K.: Calcium-induced splitting of connectin filaments into β -connectin and a 1,200-kDa subfragment. *J. Biochem. (Tokyo)*, 111 (1992) 778-782.
- 346 Weiss, H.-D., Betz, E. and Karsch, K.R.: Comparison of the cellular protein composition of aortic smooth muscle cells *in vivo* and *in vitro* by two-dimensional gel electrophoresis. *Electrophoresis (Weinheim)*, 13 (1992) 757-758.
- 347 Yoshida, M., Suzuki, A., Shimizu, T. and Ozawa, E.: Proteinase-sensitive sites on isolated rabbit dystrophin. *J. Biochem. (Tokyo)*, 112 (1992) 433-439.
- See also 188, 192, 198, 247, 302, 303, 361, 403, 409, 422, 446.
- 19g. *Protamines, histones and other nuclear proteins*
- 348 Baubichon-Cortay, H., Mallet, L., Denoroy, L. and Roux, B.: Histone H1a subtype presents structural differences compared to other histone H1 subtypes. Evidence for a specific motif in the C-terminal domain. *Biochim. Biophys. Acta*, 1122 (1992) 167-177.
- 349 Hozak, P., Roussel, P. and Hernandez-Verdun, D.: Procedures for specific detection of silver-stained nucleolar proteins on Western blots. *J. Histochem. Cytochem.*, 40 (1992) 1089-1096; C.A., 117 (1992) 107561v.
- 350 Hughes, E.N., Engelsberg, B.N. and Billings, P.C.: Purification of nuclear proteins that bind to cisplatin-damaged DNA. Identity with high mobility group proteins 1 and 2. *J. Biol. Chem.*, 267 (1992) 13520-13527.
- 351 Lindner, H., Helliger, W., Dirschlmaier, A., Talasz, H., Wurm, M., Sarg, B., Jaquemar, M. and Puschenodorf, B.: Separation of phosphorylated histone H1 variants by high-performance capillary electrophoresis. *J. Chromatogr.*, 608 (1992) 211-216.

- 352 Panin, L.E., Polyakov, L.M., Kuz'menko, A.P., Poteryaeva, O.N. and Skryl', G.Sh.: (Detection of apoprotein A-1 immunoreactivity in rat hepatocyte nuclear chromatin). *Biokhimiya (Moscow)*, 57 (1992) 826-832.
- 353 Rajpurohit, R., Paik, W.K. and Kim, S.: Enzymatic methylation of heterogeneous nuclear ribonucleoprotein in isolated liver nuclei. *Biochim. Biophys. Acta*, 1122 (1992) 183-188.
- 354 Thomassin, H., Ménard, L., Hengartner, C., Kirkland, J.B. and Poirier, G.G.: Poly(ADP-ribosylation) of chromatin in an *in vitro* poly(ADP-ribose)-turnover system. *Biochim. Biophys. Acta*, 1137 (1992) 171-181.
- 355 Vlasova, T.I., Demidenko, Z.N., Kirnos, M.D. and Vanyushin, B.F.: (Replicative cytosine DNA-methyltransferase from wheat seedling chromatin formed in the presence of 5-azacytidine). *Biokhimiya (Moscow)*, 57 (1992) 1049-1056.
- 356 Wang, R.: (Polyacrylamide slab gel electrophoresis of human brain glioma nonhistone chromatin protein (NHCP)). *Shengwu Huaxue Zazhi*, 7 (1991) 79-82; *C.A.*, 117 (1992) 43856k.
- See also 698.
- 19h. *Chromoproteins and metalloproteins*
- 357 Chevigne, R. and Janssen, J.: Method for separating HB_{A1c} from other hemoglobins by electrophoresis on chondroitin sulfate-contg. agarose gel. *Eur. Pat. Appl. EP 495,347* (Cl. GO1N27/447), 22 Jul. 1992, BE Appl. 91/27, 15 Jan. 1991; 7 pp.; *C.A.*, 117 (1992) 127813g.
- 358 Gladilov, V.V., Tulupov, G.V. and Zakkvatova, N.D.: Method for determination of hemoglobin heterogeneity. *U.S.S.R. SU 1,716,450* (Cl. GO1N33/72), 28 Feb. 1992, Appl. 4,630,647, 02 Jan. 1989; *C.A.*, 117 (1992) 127811e.
- 359 Ishioka, N., Iyori, N., Noji, J. and Kurioka, S.: Detection of abnormal haemoglobin by capillary electrophoresis and structural identification. *Biomed. Chromatogr.*, 6 (1992) 224-226.
- 360 Kalakutsky, K.L., Shvetsov, A.A., Bursakov, S.A., Letarov, A.V., Zabolotny, A.I. and L'vov, N.P.: (A highly sensitive method for detecting molybdenum-containing proteins). *Biokhimiya (Moscow)*, 57 (1992) 911-918.
- 361 Picello, E., Damiani, E. and Margreth, A.: Low-affinity Ca²⁺-binding sites versus Zn²⁺-binding sites in histidine-rich Ca²⁺-binding protein of skeletal muscle sarcoplasmic reticulum. *Biochem. Biophys. Res. Commun.*, 186 (1992) 659-667.
- 362 Vergeres, G. and Waskell, L.: Expression of cytochrome b₅ in yeast and characterization of mutants of the membrane-anchoring domain. *J. Biol. Chem.*, 267 (1992) 12583-12591.
- 363 Zaret, D.L., Morrison, N., Gulbranson, R. and Keren, D.F.: Immunofixation to quantify β_2 -transferrin in cerebrospinal fluid to detect leakage of cerebrospinal fluid from skull injury. *Clin. Chem. (Winston-Salem)*, 38 (1992) 1909-1912.
- 364 Zhu, M., Rodriguez, R., Wehr, T. and Siebert, C.: Capillary electrophoresis of hemoglobins and globin chains. *J. Chromatogr.*, 608 (1992) 225-237.
- See also 426, 696.
- 19i. *Proteins of glands, gland products, various zymogens (incl. milk proteins)*
- 365 Detrich, H.W., III, Fitzgerald, T.J., Dinsmore, J.H. and Marchese-Ragona, S.P.: Brain and egg tubulins from antarctic fishes are functionally and structurally distinct. *J. Biol. Chem.*, 267 (1992) 18766-18775.
- 366 Fenner, G.P., Johnson, L.A., Hruschka, W.R. and Bolt, D.J.: Two-dimensional electrophoresis and densitometric analysis of solubilized bovine sperm plasma membrane proteins detected by silver staining and radioiodination. *Arch. Androl.*, 29 (1992) 21-32; *C.A.*, 117 (1992) 167022e.
- 367 Häsler, M., Arendt, T. and Lange, K.: Effects of bile salt exposure on pancreatic duct barrier function and protein release determined by two-dimensional electrophoresis and lectin-affinoblotting. *Electrophoresis (Weinheim)*, 13 (1992) 747-748.
- 368 Kawaminami, M., Tanaka, K., Asawa, T., Osugi, S., Kawauchi, H., Kurusu, S. and Hashimoto, I.: Association of annexin V with prolactin in the rat anterior pituitary gland. *Biochem. Biophys. Res. Commun.*, 186 (1992) 894-899.
- 369 Mayer, W. and Hörtnner, H.: Discontinuous electrophoresis of β -caseins for the determination of bovine caseins in milk and dairy products. *Electrophoresis (Weinheim)*, 13 (1992) 803-804.
- 370 Mehrens, H.A. and Mueller, G.: (Research methods for milk proteins. Electrophoretic methods). *Lebensmittelchem., Lebensmittelqual.*, 4 (Milchproteine) (1991) 116-124; *C.A.*, 117 (1992) 146348a - a review with 34 refs.
- 371 Mendoza, C.E.C., Bhatti, T. and Bhatti, A.R.: Electrophoretic analysis of snake venoms. *J. Chromatogr.*, 580 (1992) 355-363.
- 372 Nishi, N., Kagawa, Y., Miyakawa, H., Oya, H. and Wada, F.: An anti-probasin monoclonal antibody recognizes a novel 40-kDa protein localized in rat liver and a specific region of kidney urinary tubule. *Biochim. Biophys. Acta*, 1117 (1992) 47-54.
- 373 Nishiura, T., Ishibashi, K. and Abe, K.: Rapid isolation by two electrophoretic steps of a cystatin induced in submandibular saliva of isoproterenol-treated rats. *Shika Kiso Igakkai Zasshi*, 33 (1991) 585-591; *C.A.*, 117 (1992) 43317k.
- 374 Strange, E.D., Malin, E.L., van Hekken, D.L. and Basch, J.J.: Chromatographic and electrophoretic methods used for analysis of milk proteins. *J. Chromatogr.*, 624 (1992) 81-102 - a review with 204 refs.
- 375 Tan, N.H. and Ponnudurai, G.: A comparative study on the electrophoretic patterns of snake venoms. *Comp. Biochem. Physiol., B: Comp. Biochem.*, 102B (1992) 103-109; *C.A.*, 117 (1992) 21696f.
- See also 352, 560, 736.
- 19j. *Proteins of brain, cerebrospinal fluid and eye*
- 376 Hu, R., Watanabe, M. and Bennett, V.: Characterization of human brain cDNA encoding the general isoform of β -spectrin. *J. Biol. Chem.*, 267 (1992) 18715-18722.
- 377 Ilincheta de Boschero, M.G. and Giusto, N.M.: Phosphatidic acid and polyphosphoinositide metabolism in rod outer segments. Differential role of soluble and peripheral proteins. *Biochim. Biophys. Acta*, 1127 (1992) 105-115.

- 378 Lindner, R. and Ungewickell, E.: Clathrin-associated proteins of bovine brain coated vesicles. An analysis of their number and assembly-promoting activity. *J. Biol. Chem.*, 267 (1992) 16567-16573.
- 379 Mashige, F., Shimizu, T., Iijima, S. and Ohkubo, A.: Analysis for cerebrospinal fluid proteins by sodium dodecyl sulfate-polyacrylamide gel electrophoresis. *Clin. Chim. (Winston-Salem)*, 38 (1992) 2008-2012.
- 380 Murphy, B.J. and Catterall, W.A.: Phosphorylation of purified rat brain Na⁺ channel reconstituted into phospholipid vesicles by protein kinase C. *J. Biol. Chem.*, 267 (1992) 16129-16134.
- 381 Nakamura, K., Okuya, Y., Katahira, M., Yoshida, S., Wada, S. and Okuno, M.: Analysis of tubulin isoforms by two-dimensional gel electrophoresis using SDS-polyacrylamide gel electrophoresis in the first dimension. *J. Biochem. Biophys. Methods*, 24 (1992) 195-203; *C.A.*, 117 (1992) 65910a.
- 382 Ortwerth, B.J., Slight, S.H., Prabhakaram, M., Sun, Y. and Smith, J.B.: Site-specific glycation of lens crystallins by ascorbic acid. *Biochim. Biophys. Acta*, 1117 (1992) 207-215.
- 383 Rao, P.V. and Zigler, J.S., Jr.: Purification and characterization of ζ -crystallin/quinone reductase from guinea pig liver. *Biochim. Biophys. Acta*, 1117 (1992) 315-320.
- 384 Roquemore, E.P., Dell, A., Morris, H.R., Panico, M., Reason, A.J., Savoy, L.-A., Wistow, G.J., Zigler, J.S., Jr., Earles, B.J. and Hart, G.W.: Vertebrate lens α -crystallins are modified by O-linked N-acetylglucosamine. *J. Biol. Chem.*, 267 (1992) 555-563.
- 385 Saito, Y., Tsubuki, S., Ito, H., Ohmi-Imajo, S. and Kawashima, S.: Possible involvement of clathrin in neuriteogenesis induced by a protease inhibitor (benzyloxycarbonyl-Leu-Leu-Leu-aldehyde) in PC12 cells. *J. Biochem. (Tokyo)*, 112 (1992) 448-455.
- 386 Torriglia, A. and Blanquet, P.R.: Purification of an active receptor for acidic and basic fibroblast growth factor from bovine retina. *Biochim. Biophys. Acta*, 1137 (1992) 215-224.
- 387 Van Boekel, M.A.M. and Hoenders, H.J.: *In vivo* glycation of bovine lens crystallins. *Biochim. Biophys. Acta*, 1159 (1992) 99-102.
- 388 Wiederkehr, F.: Electrophoresis of cerebrospinal fluid proteins. *Adv. Electrophor.*, 5 (1992) 241-283; *C.A.*, 117 (1992) 127462s - a review with 170 refs.
- See also 365, 411, 434, 587.
- 19k. *Proteins of neoplastic tissue and transformed cells*
- 389 Baker, M.S., Liang, X.-M. and Doe, W.F.: Occupancy of the cancer cell urokinase receptor (uPAR): effects of acid elution and exogenous uPA on cell surface urokinase (uPA). *Biochim. Biophys. Acta*, 1117 (1992) 143-152.
- 390 Henslee, J.G., Vachula, E.J., Paxton, R.J., Matias, M.S., Shively, J.E., Rittenhouse, H.G. and Tomita, J.T.: Identification and characterization of new antigenic fragments related to carcinoembryonic antigen in adult feces. *Cancer Res.*, 52 (1992) 4175-4182.
- 391 Koterov, A.N. and Risnik, V.V.: (SSB Proteins binding to DNA in Ehrlich ascites tumour cell chromatin). *Biokhimiya (Moscow)*, 57 (1992) 1083-1088.
- 392 Koyama, Y., Hayashi, T., Fujii, N. and Yoshida, T.: Recombinant mouse tumor necrosis factor expressed in mammalian cells: effect of glycosylation on cytotoxic activity. *Biochim. Biophys. Acta*, 1132 (1992) 188-194.
- 393 Krishnan, V., Narasimhan, T. and Safe, S.: Development of gel staining techniques for detecting the secretion of procathepsin D (52-kDa protein) in MCF-7 human breast cancer cells. *Anal. Biochem.*, 204 (1992) 137-142.
- 394 Sage, E.H.: Secretion of SPARC by endothelial cells transformed by polyoma middle T oncogene inhibits the growth of normal endothelial cells *in vitro*. *Biochem. Cell Biol.*, 70 (1992) 579-592.
- 395 Ueki, N., Nakazato, M., Ohkawa, T., Ikeda, T., Amuro, Y., Hada, T. and Higashino, K.: Excessive production of transforming growth-factor β 1 can play an important role in the development of tumorigenesis by its action for angiogenesis: validity of neutralizing antibodies to block tumor growth. *Biochim. Biophys. Acta*, 1137 (1992) 189-196.
- See also 172, 190, 230.
- 19l. *Specific binding and receptor proteins*
- 396 Alexander, D.R., Brown, M.H., Tutt, A.L., Crumpton, M.J. and Shivnan, E.: CD3 and CD2 antigen-mediated CD3 γ -chain phosphorylation in permeabilized human T cells. Regulation by cytosolic phosphatases. *Biochem. J.*, 288 (1992) 69-77.
- 397 Alexandru, D. and Iftode, C.: Purification of albumin binding proteins by successive polyacrylamide gel electrophoresis. *Rev. Roum. Biochim.*, 29 (1992) 3-5; *C.A.*, 117 (1992) 167009f.
- 398 Banerjee, P., Dawson, G. and Dasgupta, A.: Enrichment of saturated fatty acid containing phospholipids in sheep brain serotonin receptor preparations: use of microwave irradiation for rapid transesterification of phospholipids. *Biochim. Biophys. Acta*, 1110 (1992) 65-74.
- 399 Berger, L.C. and Seils, B.H.: Modification of mRNA-associated proteins during changes in growth conditions. *Biochem. Cell Biol.*, 70 (1992) 528-534.
- 400 Buck, F., Luth, C., Strupat, K. and Bretting, H.: Comparative investigations on the amino-acid sequences of different isolectins from the sponge *Axinella polypoides* (Schmidt). *Biochim. Biophys. Acta*, 1159 (1992) 1-8.
- 401 Dalton, A., Hornby, D.P., Langston, S.P. and Blackburn, G.M.: Characterization and purification of a novel dATP-binding protein in eukaryotes. *Biochem. J.*, 287 (1992) 871-879.
- 402 Dusanter-Fourt, I., Casadevall, N., Lacombe, C., Muller, O., Bilal, C., Fischer, S. and Mayeux, P.: Erythropoietin induces the tyrosine phosphorylation of its own receptor in human erythropoietin-responsive cells. *J. Biol. Chem.*, 267 (1992) 10670-10675.
- 403 Fukasawa, M., Arai, H. and Inoue, K.: Establishment of anti-human cholesteryl ester transfer protein monoclonal antibodies and radioimmunoassaying of the level of cholesteryl ester transfer protein in human plasma. *J. Biochem. (Tokyo)*, 111 (1992) 696-698.
- 404 Grässer, F.A., Göttel, S., Haiss, P., Boldyreff, B., Issinger, O.G. and Mueller-Lantzsch, N.: Phosphorylation of the Epstein-Barr virus nuclear antigen 2. *Biochem. Biophys. Res. Commun.*, 186 (1992) 1694-1701.
- 405 Hamaguchi, Y., Yamamoto, Y., Iwanari, H., Maruyama, S., Furukawa, T., Matsunami, N. and Honjo, T.: Biochemical and immunological characterization of the DNA binding protein (RBP-J_c) to mouse J_c recombination signal sequence. *J. Biochem. (Tokyo)*, 112 (1992) 314-320.

- 406 Hanasaki, K. and Arita, H.: Purification and characterization of a high-affinity binding protein for pancreatic-type phospholipase A₂. *Biochim. Biophys. Acta*, 1127 (1992) 233-241.
- 407 Hokland, B., Mendez, A.J. and Oram, J.F.: Cellular localization and characterization of proteins that bind high density lipoprotein. *J. Lipid Res.*, 33 (1992) 1355-1342.
- 408 Hotchin, N.A. and Watt, F.M.: Transcriptional and post-translational regulation of β_1 integrin expression during keratinocyte terminal differentiation. *J. Biol. Chem.*, 267 (1992) 14852-14858.
- 409 Imagawa, T., Nakai, J., Takeshima, H., Nakasaki, Y. and Shigekawa, M.: Expression of Ca²⁺ release channel activity from cardiac ryanodine receptor cDNA in Chinese hamster ovary cells. *J. Biochem. (Tokyo)*, 112 (1992) 508-513.
- 410 Inomata, Y., Kawaguchi, H., Hiramoto, M., Wada, T. and Handa, H.: Direct purification of multiple ATF/E4TF3 polypeptides from HeLa cell crude nuclear extracts using DNA affinity latex particles. *Anal. Biochem.*, 206 (1992) 109-114.
- 411 Izykova, G.A. and Taranova, N.P.: (Dalargin-binding proteins of brain synaptic membranes: isolation and properties in comparison with opiate receptors). *Biokhimiya (Moscow)*, 57 (1992) 663-670.
- 412 Kalsi, G., Das, H.R., Babu, C.R. and Das, R.H.: Isolation and characterization of a lectin from peanut roots. *Biochim. Biophys. Acta*, 1117 (1992) 114-119.
- 413 Kitteringham, N.R., Kenna, J.G., McLean, C., Clarke, J.B. and Park, B.K.: Conjugation of dinitrofluorobenzene to plasma proteins *in vivo* in the rat. *Drug Metab. Disp.*, 20 (1992) 625-631.
- 414 Kreienkamp, H.-J., Utkin, Y.N., Weise, C., Machold, J., Zsetlin, V.I. and Hucho, F.: Investigation of ligand-binding sites of the acetylcholine receptor using photoactivatable derivatives of neurotoxin II from *Naja naja oxiana*. *Biochemistry*, 31 (1992) 8239-8244.
- 415 MacKay, K., Danielpour, D., Miller, D., Border, W.A. and Robbins, A.R.: The 260-kDa transforming growth factor (TGF)- β binding protein in rat glomeruli is a complex comprised of 170- and 85-kDa TGF- β binding proteins. *J. Biol. Chem.*, 267 (1992) 11449-11454.
- 416 Mahon, M.J. and Gasiewicz, T.A.: Chelatable metal ions are not required for aryl hydrocarbon receptor transformation to a DNA binding form: phenanthrolines are possible competitive antagonists of 2,3,7,8-tetrachlorodibenzo-*p*-dioxin. *Arch. Biochem. Biophys.*, 297 (1992) 1-8.
- 417 Mangroo, D. and Gerber, G.E.: Photoaffinity labeling of fatty acid-binding proteins involved in long chain fatty acid transport in *Escherichia coli*. *J. Biol. Chem.*, 267 (1992) 17095-17101.
- 418 Mezqueldi, M., Fattoum, A., Derancourt, J. and Kassab, R.: Mapping of the functional domains in the amino-terminal region of calponin. *J. Biol. Chem.*, 267 (1992) 15943-15951.
- 419 Miyazaki, K., Hamano, T. and Hayashi, M.: Heat and autoclave resistance of cell-spreading activity of vitronectin. *Biochim. Biophys. Acta*, 1159 (1992) 215-222.
- 420 Modderman, P.W., Admiraal, L.G., Sonnenberg, A. von dem Borne, A.E.G.K.: Glycoproteins V and Ib-IX form a noncovalent complex in the platelet membrane. *J. Biol. Chem.*, 267 (1992) 364-369.
- 421 Moss, S.J., Doherty, C.A. and Haganir, R.L.: Identification of the cAMP-dependent protein kinase and protein kinase C phosphorylation sites within the major intracellular domains of the β_1 , γ_2S , and γ_2L subunits of the γ -aminobutyric acid type A receptor. *J. Biol. Chem.*, 267 (1992) 14470-14476.
- 422 Muryama, T. and Ogawa, Y.: Purification and characterization of two ryanodine-binding protein isoforms from sarcoplasmic reticulum of bullfrog skeletal muscle. *J. Biochem. (Tokyo)*, 112 (1992) 514-522.
- 423 Park, C.W. and Walker, M.D.: Subunit structure of cell-specific E box-binding proteins analyzed by quantitation of electrophoretic mobility shift. *J. Biol. Chem.*, 267 (1992) 15642-15649.
- 424 Ploug, M., Eriksen, J., Plesner, T., Hansen, N.E. and Dano, K.: A soluble form of the glycolipid-anchored receptor for urokinase-type plasminogen activator is secreted from peripheral blood leukocytes from patients with paroxysmal nocturnal hemoglobinuria. *Eur. J. Biochem.*, 208 (1992) 397-404.
- 425 Rabishaw, J.D., Kalman, V.K. and Proulx, K.L.: Production, processing, and partial purification of functional G protein $\beta\gamma$ subunits in baculovirus-infected insect cells. *Biochem. J.*, 286 (1992) 677-680.
- 426 Reed, J.C., Tanaka, S., Cuddy, M., Cho, D., Smith, J., Kallen, R., Saragovi, H.U. and Torigoe, T.: A strategy for generating monoclonal antibodies against recombinant baculovirus-produced proteins: application to the Bcl-2 oncoprotein. *Anal. Biochem.*, 205 (1992) 70-76.
- 427 Regazzi, R., Vallar, L., Ullrich, S., Ravazzola, M., Kikuchi, A., Takai, Y. and Wollheim, C.B.: Characterization of small-molecular-mass guanine-nucleotide-binding regulatory proteins in insulin-secreting cells and PC12 cells. *Eur. J. Biochem.*, 208 (1992) 729-737.
- 428 Schwartz, H.L., Strait, K.A., Ling, N.C. and Oppenheimer, J.H.: Quantitation of rat tissue thyroid hormone binding receptor isoforms by immunoprecipitation of nuclear triiodothyronine binding capacity. *J. Biol. Chem.*, 267 (1992) 11794-11799.
- 429 Siegenthaler, G., Tomatis, I., Chatellard-Gruaz, D., Jacconi, S., Eriksson, U. and Saurat, J.H.: Expression of CRABP-I and -II in human epidermal cells. Alteration of relative protein amounts is linked to the state of differentiation. *Biochem. J.*, 287 (1992) 383-389.
- 430 Slominski, A. and Pruski, D.: L-Dopa binding sites in rodent melanoma cells. *Biochim. Biophys. Acta*, 1139 (1992) 324-328.
- 431 Stojkovski, S., Goumakos, W. and Sarkar, B.: Iron(III)-binding polypeptide in human cord and adult serum: isolation, purification and partial characterization. *Biochim. Biophys. Acta*, 1137 (1992) 155-161.
- 432 Suzuki, T., Nagata, K.-i., Matsuura, Y., Okano, Y. and Nozawa, Y.: Purification and characterization of a low *M_r* GTP-binding protein, ram p25, expressed by baculovirus expression system. *Biochim. Biophys. Acta*, 1159 (1992) 162-168.
- 433 Takasuka, T., Adachi, M., Miyamoto, C., Furuichi, Y. and Watanabe, T.: Characterization of endothelin receptors ET_A and ET_B expressed in COS cells. *J. Biochem. (Tokyo)*, 112 (1992) 396-400.
- 434 Takasuka, T., Akiyama, N., Horii, I., Furuichi, Y. and Watanabe, T.: Different stability of ligand-receptor complex formed with two endothelin receptor species, ET_A and ET_B. *J. Biochem. (Tokyo)*, 111 (1992) 748-753.

- 435 Thoenes, G., Soudais, C., le Deist, F., Griscelli, C., Fischer, A. and Lisowska-Grospierre, B.: Structural analysis of low TCR-CD3 complex expression in T cells of an immunodeficient patient. *J. Biol. Chem.*, 267 (1992) 487-493.
- 436 Verheyden, K., Fransen, M., van Veldhoven, P.P. and Mannaerts, G.P.: Presence of small GTP-binding proteins in the peroxisomal membrane. *Biochim. Biophys. Acta*, 1109 (1992) 48-54.
- 437 Weinstein, S.P. and Haber, R.S.: Differential regulation of glucose transporter isoforms by thyroid hormone in rat heart. *Biochim. Biophys. Acta*, 1136 (1992) 302-308.
- 438 Wenisch, E., Righetti, P.G. and Weber, W.: Purification to single isoforms of a secreted epidermal growth factor receptor in a multicompartiment electrolyzer with isoelectric membranes. *Electrophoresis (Weinheim)*, 13 (1992) 668-673.
- 439 Xanthopoulos, K.G., Cannon, P.D., Robinson, G.S., Mirkovitch, J. and Darnell, J.E., Jr.: Transiently and stably introduced CCAAT/enhancer-binding-protein genes are constitutively expressed in cultured cells. *Eur. J. Biochem.*, 208 (1992) 501-509.
- 440 Yamakita, Y., Yamashiro, S. and Matsumura, F.: Characterization of mitotically phosphorylated caldesmon. *J. Biol. Chem.*, 267 (1992) 12022-12029.
- 441 Yip, C.C. and Jack, E.: Insulin receptors are bivalent as demonstrated by photoaffinity labeling. *J. Biol. Chem.*, 267 (1992) 13131-13134.
- 442 Zantema, A., Verlaan-De Vries, M., Maasdam, D., Bol, S. and van der Eb, A.: Heat shock protein 27 and α B-crystallin can form a complex, which dissociates by heat shock. *J. Biol. Chem.*, 267 (1992) 12936-12941.
- See also 142, 313, 361, 386, 391, 454, 458, 584, 603.
- 19m. *Urinary proteins*
- 443 Dakshinamurty, K.V., Melzer, H., Gladziwa, U., Kierdorf, H. and Mann, H.: (Protein gel electrophoresis of urinary proteins: urological results). *Nieren-Hochdruckkrankh.*, 21 (1992) 143-147; *C.A.*, 117 (1992) 65886x.
- 444 Zimmerman, U.-J., Speicher, D.W. and Fisher, A.B.: Secretagogue-induced proteolysis of lung spectrin in alveolar epithelial type II cells. *Biochim. Biophys. Acta*, 1137 (1992) 127-134.
- 19n. *Other proteins (incl. proteinous inhibitors of enzymic activity)*
- 445 Kaplan, B. and Pras, M.: Separation of amyloid proteins by size-exclusion chromatography in aqueous organic mobile phase. *J. Liq. Chromatogr.*, 15 (1992) 2467-2486.
- 446 Volodina, T.V., Nikitina, Z.K., Shishkin, S.S., Kurov, B.A., Rebrov, L.B. and Kozel'tsev, V.I.: (A comparative study of the protein composition of the human aorta tunica media). *Biokhimiya (Moscow)*, 57 (1992) 919-926.
- See also 249, 339.
20. ENZYMES AND ENZYME ACTIVITY ESTIMATION
- 447 Bao, J. and Regnier, F.E.: Ultramicro enzyme assays in a capillary electrophoretic system. *J. Chromatogr.*, 608 (1992) 217-224.
- 448 Chen, S.-M. and Wiktorowicz, J.E.: Isoelectric focusing by free solution capillary electrophoresis. *Anal. Biochem.*, 206 (1992) 84-90.
- See also 211.
- 20a. *Oxidoreductases*
- 449 Abadeh, S., Killackey, J., Benboubetra, M. and Harrison, R.: Purification and partial characterization of xanthine oxidase from human milk. *Biochim. Biophys. Acta*, 1117 (1992) 25-32.
- 450 Andresen, B.S., Bross, P., Jensen, U.B., Holme, E., Kolvraa, S., Gregersen, N. and Bolund, L.: Expression of wild-type and mutant medium-chain acyl-CoA dehydrogenase (MCAD) cDNA in eucaryotic cells. *Biochim. Biophys. Acta*, 1180 (1992) 65-72.
- 451 Arata, H., Shimizu, M. and Takamiya, K.-i.: Purification and properties of trimethylamine N-oxide reductase from aerobic photosynthetic bacterium *Roseobacter denitrificans*. *J. Biochem. (Tokyo)*, 112 (1992) 470-475.
- 452 Basaglia, F., Marchetti, M.G. and Cucchi, C.: The effects of phenylhydrazine and cobalt chloride on the electrophoretic and isoelectric focusing behavior of some enzymes in *Clarias gariepinus* (Clariidae, Teleostei). *Comp. Biochem. Physiol., B: Comp. Biochem.*, 102B (1992) 285-292; *C.A.*, 117 (1992) 42233t.
- 453 Chen, J.S., Preston, J.F., Wei, C., Hooshar, P., Gleeson, R.A. and Marshall, M.R.: Structural comparison of crustacean, potato and mushroom polyphenol oxidases. *J. Agric. Food Chem.*, 40 (1992) 1326-1330.
- 454 Cuthbert, J.A. and Lipsky, P.E.: Differential regulation of the expression of 3-hydroxy-3-methylglutaryl coenzyme A reductase, synthase, and low density lipoprotein receptor genes. *J. Lipid Res.*, 33 (1992) 1157-1163.
- 455 Fernandes, P.A., Keen, J.N., Findlay, J.B.C. and Moradas-Ferreira, P.: A protein homologous to glyceraldehyde-3-phosphate dehydrogenase is induced in the cell wall of a flocculent *Kluyveromyces marxianus*. *Biochim. Biophys. Acta*, 1159 (1992) 67-73.
- 456 Grebenshchikova, O.G., Alekseeva, A.E., Potapkina, T.A. and Prozorovsky, V.N.: (Study of structure-function peculiarities of lactate dehydrogenase isoenzymes using antipeptide antibodies against porcine lactate dehydrogenase isoform M₄). *Biokhimiya (Moscow)*, 57 (1992) 758-766.
- 457 Itoh, N., Morinaga, N. and Nomura, A.: A variety of catalases and bromoperoxidases in genus *Pseudomonas* and their characterization. *Biochim. Biophys. Acta*, 1122 (1992) 189-195.
- 458 Jackson, S.M., Wood, L.C., Lauer, S., Taylor, J.M., Cooper, A.D., Elias, P.M. and Feingold, K.R.: Effect of cutaneous permeability barrier disruption on HMG-CoA reductase, LDL receptor, and apolipoprotein E mRNA levels in the epidermis of hairless mice. *J. Lipid Res.*, 33 (1992) 1307-1314.
- 459 Johnson, T.M., Pease, E.A., Li, J.K.-K. and Tien, M.: Production and characterization of recombinant lignin peroxidase isozyme H2 from *Phanerochaete chrysosporium* using recombinant baculovirus. *Arch. Biochem. Biophys.*, 296 (1992) 660-666.
- 460 Matsushita, K., Takahashi, K., Takahashi, M., Ameyama, M. and Adachi, O.: Methanol and ethanol oxidase respiratory chains of the methylotrophic acetic acid bacterium, *Acetobacter methanolicus*. *J. Biochem. (Tokyo)*, 111 (1992) 739-747.

- 461 Mizoguchi, T., Nanjo, H., Umemura, T., Nishinaka, T., Iwata, C., Imanishi, T., Tanaka, T., Terada, T. and Nishihara, T.: A novel dihydrodiol dehydrogenase in bovine liver cytosol: purification and characterization of multiple forms of dihydrodiol dehydrogenase. *J. Biochem. (Tokyo)*, 112 (1992) 523-529.
- 462 Morel, F., Surla, A. and Vignais, P.V.: Purification of human placenta diamine oxidase. *Biochem. Biophys. Res. Commun.*, 187 (1992) 178-186.
- 463 Ohshima, T. and Ishida, M.: A large-scale preparative electrophoretic method for the purification of pyridine nucleotide-linked dehydrogenases. *Protein Expression Purif.*, 3 (1992) 121-125; C.A., 117 (1992) 126983a.
- 464 Pascual, P., Martínez-Lara, E., Bárcena, J.A., López-Barea, J. and Toribio, F.: Direct assay of glutathione peroxidase activity using high-performance capillary electrophoresis. *J. Chromatogr.*, 581 (1992) 49-56.
- 465 Pérez-Vicente, R., Alamillo, J.M., Cárdenas, J. and Pineda, M.: Purification and substrate inactivation of xanthine dehydrogenase from *Chlamydomonas reinhardtii*. *Biochim. Biophys. Acta*, 1117 (1992) 159-166.
- 466 Ryzlak, M.T., Ambroziak, W. and Schaffner, C.P.: Human prostatic aldehyde dehydrogenase of healthy controls and diseased prostates. *Biochim. Biophys. Acta*, 1139 (1992) 287-294.
- 467 Sharonov, B.P. and Churilova, I.V.: (Oxidative modification and inactivation of superoxide dismutase by hypochlorite). *Biochimiya (Moscow)*, 57 (1992) 719-727.
- 468 Shefer, S., Nguyen, L.B., Salen, G., Ness, G.C., Chowdhary, I.R., Lerner, S., Batta, A.K. and Tint, G.S.: Differing effects of cholesterol and taurocholate on steady state hepatic HMG-CoA reductase and cholesterol 7 α -hydroxylase activities and mRNA levels in the rat. *J. Lipid Res.*, 33 (1992) 1193-1200.
- 469 Taylor, K.L., Uhlinger, D.J. and Kinkade, J.M., Jr.: Expression of recombinant myeloperoxidase using a baculovirus expression system. *Biochem. Biophys. Res. Commun.*, 187 (1992) 1572-1578.
- 470 Yamada, J., Sakuma, M. and Suga, T.: Induction of peroxisomal β -oxidation enzymes by dehydroepiandrosterone and its sulfate in primary cultures of rat hepatocytes. *Biochim. Biophys. Acta*, 1137 (1992) 231-236.
- 471 Yang, T. and Mei, S.: (Comparison and study of positive and negative stains for SOD activity). *Shengwu Huaxue Yu Shengwu Wuli Jinzhan*, 18 (1991) 468-470; C.A., 117 (1992) 65217e.
- 472 Zhang, J., Blusztajn, J.K. and Zeisel, S.H.: Measurement of the formation of betaine aldehyde and betaine in rat liver mitochondria by a high pressure liquid chromatography-radioenzymatic assay. *Biochim. Biophys. Acta*, 1117 (1992) 333-339.
- 473 Zimmermann, P. and Kadenbach, B.: Modified structure and kinetics of cytochrome-c oxidase in fibroblasts from patients with Leigh syndrome. *Biochim. Biophys. Acta*, 1180 (1992) 99-106.
- See also 252, 577.
- 20b. *Transferases (excl. E.C. 2.7.-.-)*
- 474 Arai, K., Yoshida, K.-I., Komoda, T., Kobayashi, N. and Sakagishi, Y.: Differences in the enzymatic nature and the sugar-chain structure of γ -glutamyl transferase between normal and carcinomatous human kidney and prostate. *Clin. Chim. Acta*, 210 (1992) 35-46.
- 475 Bosshart, H. and Berger, E.G.: Biosynthesis and intracellular transport of α -2,6-sialyltransferase in rat hepatoma cells. *Eur. J. Biochem.*, 208 (1992) 341-349.
- 476 De la Rosa, J., LeGros, H.L., Jr., Geller, A.M. and Kotb, M.: Changes in the relative amount of subunits of methionine adenosyltransferase in human lymphocytes upon stimulation with a polyclonal T cell mitogen. *J. Biol. Chem.*, 267 (1992) 10699-10704.
- 477 Evjen, G. and Huseby, N.-E.: Characterization of the carbohydrate moiety of human γ -glutamyltransferases using lectin-blotting and glycosidase treatment. *Clin. Chim. Acta*, 209 (1992) 27-34.
- 478 Lee, K.B., Desai, U.R., Palcic, M.M., Hindsgaul, O. and Linhardt, R.J.: An electrophoresis-based assay for glycosyltransferase activity. *Anal. Biochem.*, 205 (1992) 108-114.
- 479 Lundqvist, G., Yücel-Lindberg, T. and Morgenstern, R.: The oligomeric structure of rat liver microsomal glutathione transferase studied by chemical cross-linking. *Biochim. Biophys. Acta*, 1159 (1992) 103-108.
- 480 Machida, Y., Fujii, K., Morimoto, S., Nomoto, T. and Motoyama, M.: (Study on the γ -GTP isozyme. 1. Basic study on a simple method). *Igaku Kansa*, 41 (1992) 880-884; C.A., 117 (1992) 85662z.
- 481 Rammesmayr, G. and Praznik, W.: Fast and sensitive simultaneous staining method of Q-enzyme, α -amylase, R-enzyme, phosphorylase and soluble starch synthase separated by starch-polyacrylamide gel electrophoresis. *J. Chromatogr.*, 623 (1992) 399-402.
- 482 Yokota, H. and Yuasa, A.: Activation and stabilization of UDP-glucuronyltransferase by lysophosphatidylcholine. *J. Biochem. (Tokyo)*, 112 (1992) 309-313.
- 20c. *Transferases transferring phosphorus containing groups (E.C. 2.7.-.-)*
- 483 Basta, P., Strickland, M.B., Holmes, W., Loomis, C.R., Ballas, L.M. and Burns, D.J.: Sequence and expression of human protein kinase C- ϵ . *Biochim. Biophys. Acta*, 1132 (1992) 154-160.
- 484 Borner, C., Guadagno, S.N., Fabbro, D. and Weinstein, I.B.: Expression of four protein kinase C isoforms in rat fibroblasts. Distinct subcellular distribution and regulation by calcium and phorbol esters. *J. Biol. Chem.*, 267 (1992) 12892-12899.
- 485 Gärtner, T., Kuhnelt, H., Raab, G., Raab, M., Strebhardt, K. and Rubsamen-Waigmann, H.: A strong protein-tyrosine kinase activity is associated with a baculovirus-expressed chicken tk1 gene. *Eur. J. Biochem.*, 208 (1992) 91-100.
- 486 Hughes, K., Ramakrishna, S., Benjamin, W.B. and Woodgett, J.R.: Identification of multifunctional ATP-citrate lyase kinase as the α -isoform of glycogen synthase kinase-3. *Biochem. J.*, 288 (1992) 309-314.
- 487 Hulkower, K.I., Sagi-Eisenberg, R., Traub, L.M., Georgescu, H.I. and Evans, C.H.: Synovial protein kinase C and its apparent insensitivity to interleukin-1. *Eur. J. Biochem.*, 209 (1992) 81-88.
- 488 Inoue, H., Hoshioka, T. and Hotta, Y.: Partial purification and characterization of membrane-associated diacylglycerol kinase of *Drosophila* heads. *Biochim. Biophys. Acta*, 1122 (1992) 219-224.

- 489 James, R.I., Menaya, J., Hudson, K., Devalia, V., Ryves, J., Evans, F.J., Thomas, S. and Clemens, M.J.: Role of protein kinase C in induction of gene expression and inhibition of cell proliferation by interferon α . *Eur. J. Biochem.*, 209 (1992) 813-822.
- 490 Kanemitsu, F. and Okigaki, T.: Cretine kinase MB isoforms for early diagnosis and monitoring of acute myocardial infarction. *Clin. Chim. Acta*, 206 (1992) 191-199.
- 491 Lee, S.A., Karaszkiwicz, J.W. and Anderson, W.B.: Elevated level of nuclear protein kinase C in multidrug-resistant MCF-7 human breast carcinoma cells. *Cancer Res.*, 52 (1992) 3750-3759.
- 492 Litchfield, D.W., Lüscher, B., Lozeman, F.J., Eisenman, R.N. and Krebs, E.G.: Phosphorylation of casein kinase II by p34^{cdc2} *in vitro* and at mitosis. *J. Biol. Chem.*, 267 (1992) 13943-13951.
- 493 Lydon, N.B., Gay, B., Mett, H., Murray, B., Liebentanz, J., Gutzwiller, A., Piwnica-Worms, H., Roberts, T.M. and McGlynn, E.: Purification and biochemical characterization of non-myristoylated recombinant pp60^{c-src} kinase. *Biochem. J.*, 287 (1992) 985-993.
- 494 Meckling-Gill, K.A., Yee, S.-P., Schrader, J.W. and Pawson, T.: A retrovirus encoding the v-fps protein-tyrosine kinase induces factor-independent growth and tumorigenicity in FDC-P1 cells. *Biochim. Biophys. Acta*, 1137 (1992) 65-72.
- 495 Mitev, V.I., Angelov, I.V. and Marekov, L.N.: Identification and purification of a novel serine/threonine messenger-independent growth-related protein kinase from lactating goat mammary gland. *Biochim. Biophys. Acta*, 1117 (1992) 90-96.
- 496 Okazaki, H., Tani, C., Ando, M., Ishii, K., Ishibashi, S., Nishimura, Y. and Kato, K.: Possible involvement of cathepsin L in processing of rat liver hexokinase to eliminate mitochondria-binding ability. *J. Biochem. (Tokyo)*, 112 (1992) 409-413.
- 497 Phillips, J.L., Haggren, W., Thomas, W.J., Ishida-Jones, T. and Adey, W.R.: Magnetic field-induced changes in specific gene transcription. *Biochim. Biophys. Acta*, 1132 (1992) 140-144.
- 498 Stahls, A.: The sulfhydryl reagent N-ethylmaleimide induces hyperphosphorylation on tyrosine residues in the Jurkat T-cell line. *Biochem. Biophys. Res. Commun.*, 187 (1992) 73-78.
- 499 Stamford, N.P., Lilley, P.E. and Dixon, N.E.: Enriched sources of *Escherichia coli* replication proteins. The dnaG primase is a zinc metalloprotein. *Biochim. Biophys. Acta*, 1132 (1992) 17-25.
- 500 Thompson, N.E., Hager, D.A. and Burgess, R.R.: Isolation and characterization of a polyol-responsive monoclonal antibody useful for gentle purification of *Escherichia coli* RNA polymerase. *Biochemistry*, 31 (1992) 7003-7008.
- See also 448.
- 20d. *Hydrolases, acting on ester bonds (E.C. 3.1.-.-)*
- 501 Chamberlain, B.R., Buttery, J.E. and Pannall, P.R.: A simple electrophoretic method for separating elevated liver and bone alkaline phosphatase isoenzymes in plasma after neuraminidase treatment. *Clin. Chim. Acta*, 208 (1992) 219-224.
- 502 Charton, E., Davies, C. and Macrae, A.R.: Use of specific polyclonal antibodies to detect heterogeneous lipases from *Geotrichum candidum*. *Biochim. Biophys. Acta*, 1127 (1992) 191-198.
- 503 Di Marco, S., Märki, F., Hofstetter, H., Schmitz, A., van Oostrum, J. and Grütter, M.G.: Purification, analysis, and enzymatic activity of recombinant human synovial fluid phospholipase A₂ and N-terminal variants. *J. Biochem. (Tokyo)*, 112 (1992) 350-354.
- 504 Fujimoto, S., Murakami, K., Hosoda, T., Yamamoto, Y., Watanabe, K., Morinaka, Y. and Ohara, A.: Characterization of cationic acid phosphatase isozyme from rat liver mitochondria. *Chem. Pharm. Bull.*, 40 (1992) 1231-1235.
- 505 Jayanthi, L.D., Balasubramanian, N. and Balasubramanian, A.S.: Cholinesterases exhibiting aryl acylamidase activity in human maniotic fluid. *Clin. Chim. Acta*, 205 (1992) 157-166.
- 506 Kadner, S.S., Katz, J. and Finlay, T.H.: Esterase-1: developmental expression in the mouse and distribution of related proteins in other species. *Arch. Biochem. Biophys.*, 296 (1992) 435-441.
- 507 Ketterman, A.J., Jayawardena, K.G.I. and Hemingway, J.: Purification and characterization of a carboxylesterase involved in insecticide resistance from the mosquito *Culex quinquefasciatus*. *Biochem. J.*, 287 (1992) 355-360.
- 508 Kobayashi, T., Honke, K., Jin, T., Gasa, S., Miyazaki, T. and Makita, A.: Components and proteolytic processing sites of arylsulfatase B from human placenta. *Biochim. Biophys. Acta*, 1159 (1992) 243-247.
- 509 Lessinger, J.M., Tavridou, A., Arzoglou, P. and Ferard, G.: Interest of using a purified, stable and commutable preparation of human pancreatic lipase in indirect assays. *Anal. Lett.*, 25 (1992) 1453-1468.
- 510 Masterson, C., Boehmer, P.E., McDonald, F., Chaudhuri, S., Hickson, I.D. and Emmerson, P.T.: Reconstitution of the activities of the RecBCD holoenzyme of *Escherichia coli* from the purified subunits. *J. Biol. Chem.*, 267 (1992) 13564-13572.
- 511 Masuno, H., Blanchette-Mackie, E.J., Schultz, C.J., Spaeth, A.E., Scow, R.O. and Okuda, H.: Retention of glucose by N-linked oligosaccharide chains impedes expression of lipoprotein lipase activity: effect of castanospermine. *J. Lipid Res.*, 33 (1992) 1343-1349.
- 512 Nishikawa, Y., Kanda, T., Yoshihara, H., Fukumoto, K. and Uematsu, I.: Cellulose acetate electrophoretic determination of bone alkaline phosphatase activity in healthy subjects and diabetic patients with and without osteopenia. *Clin. Chim. Acta*, 210 (1992) 13-22.
- 513 Onica, D., Torssander, J. and Waldenlind, L.: Recurrent transient hyperphosphatasemia of infancy in an adult. *Clin. Chem. (Winston-Salem)*, 38 (1992) 1913-1915.
- 514 Panteghini, M.: Electrophoretic fractionation of pancreatic lipase. *Clin. Chem. (Winston-Salem)*, 38 (1992) 1712-1716.
- 515 Poretz, R.D., Yang, R.S., Canas, B., Lackland, H., Stein, S. and Manowitz, P.: The structural basis for the electrophoretic isoforms of normal and variant human platelet arylsulphatase A. *Biochem. J.*, 287 (1992) 979-983.
- 516 Saeki, S., Ochi, M., Horiuchi, T., Sakagishi, Y., Fujimori-Arai, Y., Koyama, I., Komoda, T. and Takeuchi, N.: Intestinal type alkaline phosphatase hyperphosphatasemia associated with liver cirrhosis. *Clin. Chim. Acta*, 210 (1992) 63-73.
- 517 Sakayama, K., Masuno, H., Okumura, H., Shibata, T. and Okuda, H.: Glycosylation of lipoprotein lipase in human subcutaneous and omental adipose tissues. *Biochim. Biophys. Acta*, 1127 (1992) 153-156.

- 518 Shnyreva, M.G., Tsuprun, V.L., Stel'mashchuk, V.Ya. and Egorov, S.N.: (Study of the quaternary structure of extracellular repressible acid phosphatase from the yeast *Saccharomyces cerevisiae*). *Biokhimiya (Moscow)*, 57 (1992) 1100-1108.
- 519 Tamura, H.-o., Tameishi, K., Yamagata, H., Udaka, S., Kobayashi, T., Tomita, M. and Ikezawa, H.: Mass production of sphingomyelinase of *Bacillus cereus* by a protein-hyperproducing strain, *Bacillus brevis* 47, and its purification. *J. Biochem. (Tokyo)*, 112 (1992) 488-491.
- 520 Yasuda, T., Nadano, D., Tenjo, E., Takeshita, H. and Kishi, K.: The zymogram method for detection of ribonucleases after isoelectric focusing: analysis of multiple forms of human, bovine, and microbial enzymes. *Anal. Biochem.*, 106 (1992) 172-177.
- See also 147.
- 20e. *Hydrolases, acting on glycosyl compounds (E.C. 3.2.-.-)*
- 521 Ajandouz, E.H., Abe, J.-i., Svensson, B. and Marchis-Mouren, G.: Barley malt- α -amylase. Purification, action pattern, and sub-site mapping of isozyme 1 and two members of the isozyme 2 subfamily using *p*-nitrophenylated maltooligosaccharide substrates. *Biochim. Biophys. Acta*, 1159 (1992) 193-202.
- 522 Chambert, R., Rain-Guion, M.-C. and Petit-Glatron, M.-F.: Read-through of the *Bacillus subtilis* stop codon produces an extended enzyme displaying a higher polymerase activity. *Biochim. Biophys. Acta*, 1132 (1992) 145-153.
- 523 Holleran, W.M., Takagi, Y., Imokawa, G., Jackson, S., Lee, J.M. and Elias, P.M.: β -Glucocerebrosidase activity in murine epidermis: characterization and localization in relation to differentiation. *J. Lipid Res.*, 33 (1992) 1201-1209.
- 524 Isaksson, A., Hultberg, B. and Jonung, T.: Rat plasma clearance rate and organ distribution of β -hexosaminidase isoenzymes from human serum. *Clin. Chem. (Winston-Salem)*, 38 (1992) 1893-1898.
- 525 Mathew, R. and Rao, K.K.: Activity staining of endoglucanases in polyacrylamide gels. *Anal. Biochem.*, 206 (1992) 50-52.
- 526 Ohta, H., Ono, M., Sekiya, C. and Namiki, M.: Serum immunoreactive β -glucuronidase determined by an enzyme-linked immunosorbent assay in patients with hepatic diseases. *Clin. Chim. Acta*, 208 (1992) 9-21.
- 527 Raab, B.: Characterization of endopolygalacturonase (EC 3.2.1.15) from *Aspergillus niger* as glycoprotein by electrophoretic methods and lectin affino-blotting. *Electrophoresis (Weinheim)*, 13 (1992) 807-808.
- 528 Rapp, P.: Formation, separation and characterization of three β -1,3-glucanases from *Sclerotium gluconicum*. *Biochim. Biophys. Acta*, 1117 (1992) 7-14.
- 529 Wang, J., Xu, P. and Fincher, G.B.: Purification, characterization and gene structure of (1 \rightarrow 3)- β -glucanase isoenzyme GIII from barley (*Hordeum vulgare*). *Eur. J. Biochem.*, 209 (1992) 103-109.
- See also 250, 481.
- 20f. *Other hydrolases*
- 530 Basak, A., Yuan, X.W., Seidah, N.G., Chrétien, M. and Lazure, C.: Affinity purification of proteinases by a combination of immobilized peptidyl aldehyde and semicarbazone. *J. Chromatogr.*, 581 (1992) 17-29.
- 531 Bock, P.E.: Active-site-selective labeling of blood coagulation proteinases with fluorescence probes by the use of thioester peptide chloromethyl ketones. II. Properties of thrombin derivatives as receptors of prothrombin fragment 2 binding and specificity of the labeling approach for other proteinases. *J. Biol. Chem.*, 267 (1992) 14974-14981.
- 532 Boretzky, Yu.R., Skoblov, Yu.S., Khodova, O.M. and Rabinovich, P.M.: (Purification and properties of GTP-cyclohydrolase of *Bacillus subtilis*). *Biokhimiya (Moscow)*, 57 (1992) 1021-1030.
- 533 Carchon, H.A., Jansen, E. and Eggermont, E.: Determination of enterokinase activity by measuring the disappearance of trypsinogen. *Clin. Chim. Acta*, 207 (1992) 175-183.
- 534 Chiou, S.-H., Hung, C.-C. and Huang, K.-F.: Characterization of a protease with α - and β -fibrinogenase activity from the western diamondback rattlesnake, *Crotalus atrox*. *Biochem. Biophys. Res. Commun.*, 187 (1992) 389-396.
- 535 Edell, C.M., Hartog, A.F. and Berden, J.A.: Inhibition of mitochondrial F_1 -ATPase activity by binding of (2-azido)-ADP to a slowly exchangeable non-catalytic nucleotide binding site. *Biochim. Biophys. Acta*, 1101 (1992) 329-338.
- 536 Hamasur, B., Guerrieri, F., Zanotti, F. and Glaser, E.: F_0F_1 -ATP synthase of potato tuber mitochondria. Structural and functional characterization by resolution and reconstitution studies. *Biochim. Biophys. Acta*, 1101 (1992) 339-344.
- 537 Ilian, M.A. and Forsberg, N.E.: Gene expression of calpains and their specific endogenous inhibitor, calpastatin, in skeletal muscle of fed and fasted rabbits. *Biochem. J.*, 287 (1992) 163-171.
- 538 Ishiye, M. and Niwa, M.: Nucleotide sequence and expression in *Escherichia coli* of the Cephalosporin acylase gene of a *Pseudomonas* strain. *Biochim. Biophys. Acta*, 1132 (1992) 233-239.
- 539 Jackson, M.C. and Lenney, J.F.: Homocarnosinosis patients and great apes have a serum protein that cross-reacts with human serum carnosinase. *Clin. Chim. Acta*, 205 (1992) 109-116.
- 540 Kageyama, T., Ichinose, M., Tsukada, S., Miki, K., Kurokawa, K., Koizumi, O., Tanji, M., Yakabe, E., Athauda, S.B.P. and Takahashi, K.: Gastric procathepsin E and progastricsin from guinea pig. Purification, molecular cloning of cDNAs, and characterization of enzymatic properties, with special reference to procathepsin E. *J. Biol. Chem.*, 267 (1992) 16450-16459.
- 541 Kronheim, S.R., Mumma, A., Greenstreet, T., Glackin, P.J., van Ness, K., March, C.J. and Black, R.A.: Purification of interleukin-1 β converting enzyme, the protease that cleaves the interleukin-1 β precursor. *Arch. Biochem. Biophys.*, 296 (1992) 698-703.
- 542 Kuba, M., Ohmori, H. and Kumon, A.: Characterization of N ϵ -phosphoarginine hydrolase from rat liver. *Eur. J. Biochem.*, 208 (1992) 747-752.
- 543 Kyouden, T., Himeno, M., Ishikawa, T., Ohsumi, Y. and Kato, K.: Purification and characterization of dipeptidyl peptidase IV in rat liver lysosomal membranes. *J. Biochem. (Tokyo)*, 111 (1992) 770-777.

- 544 Ledgerwood, E.C., George, P.M., Bathurst, I.C. and Brennan, S.O.: The predicted proteinase furin is not the hepatic proalbumin convertase. *Biochim. Biophys. Acta*, 1159 (1992) 9-12.
- 545 Obata, K., Iwata, K., Okada, Y., Kohrin, Y., Ohuchi, E., Yoshida, S., Shinmei, M. and Hayakawa, T.: A one-step sandwich enzyme immunoassay for human matrix metalloproteinase 3 (stromelysin-1) using monoclonal antibodies. *Clin. Chim. Acta*, 211 (1992) 59-72.
- 546 Retzius, A.D. and Markland, F.S., Jr.: Purification, characterization, and fibrinogen cleavage sites of three fibrinolytic enzymes from the venom of *Crotalus basiliscus basiliscus*. *Biochemistry*, 31 (1992) 4547-4557.
- 547 Rowan, A.D., Mason, P., Mach, L. and Mort, J.S.: Rat procathepsin B. Proteolytic processing to the mature form *in vitro*. *J. Biol. Chem.*, 267 (1992) 15993-15999.
- 548 Southgate, K.M., Davies, M., Booth, R.F.G. and Newby, A.C.: Involvement of extracellular-matrix-degrading metalloproteinases in rabbit aortic smooth-muscle cell proliferation. *Biochem. J.*, 288 (1992) 93-99.
- 549 Tozawa, K., Odaka, M., Date, T. and Yoshida, M.: Molecular dissection of the β subunit of F₁-ATPase into peptide fragments. *J. Biol. Chem.*, 267 (1992) 16484-16490.
- 550 Treuheit, M.J., Ataei, A., Kirley, T.L. and Wallick, E.T.: Purification of the glycosylated beta subunit of (Na,K)-ATPase by lectin affinity chromatography. *Chromatographia*, 33 (1992) 521-524.
- 551 Vinther, A., Petersen, J. and Soeborg, H.: Capillary electrophoretic determination of the protease Savinase in cultivation broth. *J. Chromatogr.*, 608 (1992) 205-210.
- 552 Wnuk, W. and Loizeau, E.: Separation and characterization of human pepsinogens and pepsins by high-resolution discontinuous electrophoresis. *Adv. Exp. Med. Biol.*, 306 (Struct. Funct. Aspartic Proteinases) (1991) 91-94; C.A., 117 (1992) 106895v.
- 553 Zakharov, S.D. and Kuz'mina, V.P.: (Subunit composition of ATP-synthase from the thermophilic bacterium *Thermus thermophilus*). *Biokhimiya (Moscow)*, 57 (1992) 777-786.
- See also 251, 260.
- 20g. *Lyases*
- 554 Aoki, T., Tanaka, T. and Watabe, H.: Purification and characterization of γ -enolase from various mammals. *Chem. Pharm. Bull.*, 40 (1992) 1236-1239.
- 555 Baur, B., Dietz, K.-J. and Winter, K.: Regulatory protein phosphorylation of phosphoenolpyruvate carboxylase in the facultative crassulacean-acid-metabolism plant *Mesembryanthemum crystallinum* L. *Eur. J. Biochem.*, 209 (1992) 95-101.
- 556 Lee, H.J., Chiou, S.H. and Chang, G.G.: Staining of argininosuccinate lyase activity in polyacrylamide gel. *J. Biochem. Biophys. Methods*, 24 (1992) 205-214; C.A., 117 (1992) 106859m.
- 557 Lee, K.N., Maxwell, M.D., Patterson, M., Jr., Birckbichler, P.J. and Conway, E.: Identification of transglutaminase substrates in HT20 colon cancer cells: use of 5-(biotinamido)pentylamine as a transglutaminase-specific probe. *Biochim. Biophys. Acta*, 1136 (1992) 12-16.
- 20h. *Isomerases*
- See 452, 622.
- 20j. *Complex mixtures and incompletely identified enzymes*
- 558 Kadono-Okuda, K., Imanishi, S. and Inoue, H.: Identification of cell lines of *Bombyx mori* and some lepidopteran insects by electrophoresis. *Nippon Sanshigaku Zasshi*, 61 (1992) 184-187; C.A., 117 (1992) 127635a.
- 559 Noerrung, B.: Characterization of Danish isolates of *Listeria monocytogenes* by multilocus enzyme electrophoresis. *Int. J. Food Microbiol.*, 15 (1992) 51-59; C.A., 117 (1992) 167250c.
- 560 Perkins, P.S. and Pandol, S.J.: Cholecystokinin-induced changes in polysome structure regulate protein synthesis in pancreas. *Biochim. Biophys. Acta*, 1136 (1992) 265-271.
21. PURINES, PYRIMIDINES, NUCLEIC ACIDS AND THEIR CONSTITUENTS
- 21a. *Purines, pyrimidines, nucleosides, nucleotides*
- 561 Akashi, M., Sawa, T., Baba, Y. and Tshuhako, M.: Specific separation of oligodeoxynucleotides by capillary affinity gel electrophoresis (CAGE) using poly(9-vinyladenine)-polyacrylamide conjugated gel. *J. High Resolut. Chromatogr.*, 15 (1992) 625-626.
- 562 Baba, Y., Tshuhako, M., Sawa, T., Akashi, M. and Yashima, E.: Specific base recognition of oligodeoxynucleotides by capillary affinity gel electrophoresis using polyacrylamide-poly(9-vinyladenine) conjugated gel. *Anal. Chem.*, 64 (1992) 1920-1925.
- 563 Guttman, A., Arai, A. and Magyar, K.: Influence of pH on the migration properties of oligonucleotides in capillary gel electrophoresis. *J. Chromatogr.*, 608 (1992) 175-179.
- 564 Klein, S. and Oesch, F.: Assay for O⁶-alkylguanine-DNA-alkyltransferase using oligonucleotides containing O⁶-methylguanine in a BamHI recognition site as substrate. *Anal. Biochem.*, 205 (1992) 294-299.
- 565 Li, W., Moussa, A. and Giese, R.W.: Capillary electrophoresis of fluorescein-ethylenediamine-5'-deoxynucleotides. *J. Chromatogr.*, 608 (1992) 171-174.
- 566 Singhal, R.P., Hughbanks, D. and Xian, J.: Separation of dideoxyribonucleosides in trace amounts by automated liquid chromatography and capillary electrophoresis. *J. Chromatogr.*, 609 (1992) 147-161.
- See also 90, 118, 135, 598, 669, 698.
- 21b. *Nucleic acids, RNA*
- 567 Adachi, H., Ishida, N. and Tsujimoto, M.: Primary structure of rat renal dipeptidase and expression of its mRNA in rat tissues and COS-1 cells. *Biochim. Biophys. Acta*, 1132 (1992) 311-314.
- 568 Adamou, J. and Bag, J.: Alteration of translation and stability of mRNA for the poly(A)-binding protein during myogenesis. *Eur. J. Biochem.*, 209 (1992) 803-812.
- 569 Boardman, M., Cross, G.S., Jones, E.A. and Woodland, H.R.: Regulation of expression of a *Xenopus borealis* embryonic/larval α 3 skeletal-actin gene. *Eur. J. Biochem.*, 208 (1992) 241-249.
- 570 Capranico, G., Tinelli, S., Austin, C.A., Fisher, M. and Zunino, F.: Different patterns of gene expression of topoisomerase II isoforms in differentiated tissues during murine development. *Biochim. Biophys. Acta*, 1132 (1992) 43-48.

- 571 Connelly, I. and Possmayer, F.: cDNA sequence and alternative mRNA splicing of surfactant-associated protein C (SP-C) in rabbit lung. *Biochim. Biophys. Acta*, 1127 (1992) 199-207.
- 572 Crespo, P., González, C., Albajar, M., Rodríguez, J.C., Ortiz, J.M. and León, J.: Downregulation of hepatic albumin mRNA in response to induced hypercholesterolemia in rabbits. *Biochim. Biophys. Acta*, 1128 (1992) 77-82.
- 573 Endo, T. and Goto, S.: Retinoblastoma gene product Rb accumulates during myogenic differentiation and is deinduced by the expression of SV40 large T antigen. *J. Biochem. (Tokyo)*, 112 (1992) 427-430.
- 574 Fujita, T., Shirasawa, T., Uchida, K. and Maruyama, N.: Isolation of cDNA clone encoding rat senescence marker protein-30 (SMP30) and its tissue distribution. *Biochim. Biophys. Acta*, 1132 (1992) 297-305.
- 575 Gonzalez-Espinosa, C. and García-Sáinz, J.A.: Angiotensin II and active phorbol esters induce proto-oncogene expression in isolated rat hepatocytes. *Biochim. Biophys. Acta*, 1136 (1992) 309-314.
- 576 Goto, H., Chen, K.-s., Prael, J.M. and DeLuca, H.F.: A single receptor identical with that from intestine/T47D cells mediates the action of 1,25-dihydroxyvitamin D-3 in HL-60 cells. *Biochim. Biophys. Acta*, 1132 (1992) 103-108.
- 577 Green, A.K., McDowall, I.L., Richardson, S.-C.B. and Fisher, M.J.: The effect of vanadate upon the expression of phenylalanine hydroxylase in streptozotocin-diabetic rat liver. *Biochim. Biophys. Acta*, 1180 (1992) 21-27.
- 578 Guerrin, M., Prats, H., Mazars, P. and Valette, A.: Antiproliferative effect of phorbol esters on MCF-7 human breast adenocarcinoma cells: relationship with enhanced expression of transforming growth-factor- β_1 . *Biochim. Biophys. Acta*, 1137 (1992) 116-120.
- 579 Higashimoto, Y., Simchock, J. and Liddle, R.A.: Molecular cloning of rat glucose-dependent insulinotropic peptide (GIP). *Biochim. Biophys. Acta*, 1132 (1992) 72-74.
- 580 Igloi, G.L.: Affinity electrophoretic detection of primary amino groups in nucleic acids: application to modified bases of tRNA and to aminoacylation. *Anal. Biochem.*, 206 (1992) 363-368.
- 581 Kremling, H., Reinhart, N., Schlösser, M. and Engel, W.: The bovine protamine 2 gene: evidence for alternative splicing. *Biochim. Biophys. Acta*, 1132 (1992) 133-139.
- 582 Langerak, A.W., Dirks, R.P.H. and Versnel, M.A.: Splicing of the platelet-derived-growth-factor A-chain mRNA in human malignant mesothelioma cell lines and regulation of its expression. *Eur. J. Biochem.*, 208 (1992) 589-596.
- 583 Liu, L. and Pohajdak, B.: Cloning and sequencing of a human cDNA from cytolytic NK/T cells with homology to yeast SEC7. *Biochim. Biophys. Acta*, 1132 (1992) 75-78.
- 584 Okano, R., Mita, T. and Matsui, T.: Characterization of a novel promoter structure and its transcriptional regulation of the murine laminin B1 gene. *Biochim. Biophys. Acta*, 1132 (1992) 49-57.
- 585 Reddy, N.G., Hall, J.C., Kochins, J.G. and Perez, F.M.: Identification of novel RNA-binding proteins that interact in the coding region of protein D sense RNA *in vitro*. *Biochem. Biophys. Res. Commun.*, 187 (1992) 318-324.
- 586 Russ, A.P., Ruzicka, V., Maerz, W., Appelhans, H. and Gross, W.: Amplification and direct sequencing of a cDNA encoding human cytosolic 3-hydroxy-3-methylglutaryl-coenzyme A synthase. *Biochim. Biophys. Acta*, 1132 (1992) 329-331.
- 587 Sazuka, T., Tomooka, Y., Kathju, S., Ikawa, Y., Noda, M. and Kumar, S.: Identification of a developmentally regulated gene in the mouse central nervous system which encodes a novel proline rich protein. *Biochim. Biophys. Acta*, 1132 (1992) 240-248.
- 588 Vaughan, T.J., Pascall, J.C. and Brown, K.D.: Nucleotide sequence and tissue distribution of mouse transforming growth factor- α . *Biochim. Biophys. Acta*, 1132 (1992) 322-324.
- 589 Volm, M., Efferth, T., Mattern, J. and Pommerenke, E.W.: Resistance mechanisms in murine tumors with acquired multidrug resistance. *Arzneim.-Forsch.*, 42 (1992) 1163-1168.
- 590 Wundrack, I. and Dooley, S.: Nonradioactive ribonuclease protection analysis using digoxigenin labeling and chemiluminescent detection. *Electrophoresis (Weinheim)*, 13 (1992) 637-638.
- 591 Zhao, X., Araki, K., Miyazaki, J.-i. and Yamamura, K.-i.: Developmental and liver-specific expression directed by the serum amyloid P component promoter in transgenic mice. *J. Biochem. (Tokyo)*, 111 (1992) 736-738.
- 592 Zhao, Y., Kuntz, M.J., Harris, R.A. and Crabb, D.W.: Molecular cloning of the E1 β subunit of the rat branched chain α -ketoacid dehydrogenase. *Biochim. Biophys. Acta*, 1132 (1992) 207-210.

See also 83, 118, 244, 250, 437, 450, 454, 523, 597, 602, 613, 615, 618, 633, 684, 686.

21c. Nucleic acids, DNA

- 593 Andresen, B.S., Knudsen, I., Jensen, P.K.A., Rasmussen, K. and Gregersen, N.: Two novel nonradioactive polymerase chain reaction-based assays of dried blood spots, genomic DNA or whole cells for fast, reliable detection of Z nad S mutations in the α_1 -antitrypsin gene. *Clin. Chem. (Winston-Salem)*, 38 (1992) 2100-2107.
- 594 Aran, J.M. and Plagemann, P.G.W.: Nucleoside transport-deficient mutants of PK-15 pig kidney cell line. *Biochim. Biophys. Acta*, 1110 (1992) 51-58.
- 595 Baba, Y. and Tshako, M.: Gel-filled capillaries for nucleic acid separations in capillary electrophoresis. *TrAC*, 11 (1992) 280-287 - a review with 35 refs.
- 596 Bensaid, A., Young, J.R., Kaushal, A. and Tesle, A.J.: Pulsed-field gel electrophoresis and its application in the physical analysis of the bovine major histocompatibility complex. In: Schook, L.B., Lewin, H.A. and McLaren, D.G. (Editors), *Gene-Mapp. Tech. Appl.*, Dekker, New York, 1991, pp. 127-158; *C.A.*, 117 (1992) 88040f - a review with 32 refs.
- 597 Brown, C.A., McInnes, B., Dodelson de Kremer, R.D. and Mahuran, D.J.: Characterization of two HEXB gene mutations in Argentinean patients with Sandhoff disease. *Biochim. Biophys. Acta*, 1180 (1992) 91-98.
- 598 Bush, E. and Currie, R.A.: Determination of transcription factor isoelectric point by two-dimensional native isoelectric focusing and electrophoretic mobility shift analysis. *Anal. Biochem.*, 206 (1992) 189-194.
- 599 Cavatorta, F., Deriu, A., Malizia, F., Terech, P. and Middendorf, H.D.: Small-angle neutron scattering from DNA molecules during gel electrophoresis. *Physica B (Amsterdam)*, 180-181 (Pt. B) (1992) 770-772; *C.A.*, 117 (1992) 107633v.

- 600 Cazals-Hatem, D.L., Louie, D.C., Tanaka, S. and Reed, J.C.: Molecular cloning and DNA sequence analysis of cDNA encoding chicken homologue of the Bcl-2 oncoprotein. *Biochim. Biophys. Acta*, 1132 (1992) 109-113.
- 601 Court, D.S., Fedor, T., Gouldstone, M., Lincoln, P.J., Phillips, C.P., Tate, V., Thomson, J.A. and Watts, P.W.: Investigation of the between-gel and within-gel variation in fragment size determinations found when using single locus DNA probes. *Forensic Sci. Int.*, 53 (1992) 173-191; C.A., 117 (1992) 84761u.
- 602 Deffie, A.M., McPherson, J.P., Gupta, R.S., Hedley, D.W. and Goldenberg, G.J.: Multifactorial resistance to antineoplastic agents in drug-resistant P388 murine leukemia, Chinese hamster ovary, and human HeLa cells, with emphasis on the role of DNA topoisomerase II. *Biochem. Cell Biol.*, 70 (1992) 354-364.
- 603 Doye, V., le Gouvello, S., Dobransky, T., Chneiweiss, H., Beretta, L. and Sobel, A.: Expression of transfected stathmin cDNA reveals novel phosphorylated forms associated with developmental and functional cell regulation. *Biochem. J.*, 287 (1992) 549-554.
- 604 Duband-Goulet, I., Carot, V., Ulyanov, A.V., Douc-Rasy, S. and Prunell, A.: Chromatin reconstitution on small DNA rings. IV. DNA supercoiling and nucleosome sequence preference. *J. Mol. Biol.*, 224 (1992) 981-1001; C.A., 117 (1992) 85471m.
- 605 Duke, T.A.J. and Viovy, J.L.: Motion of megabase deoxyribonucleic acid during field-inversion gel electrophoresis: investigation by nonlocal Monte Carlo. *J. Chem. Phys.*, 96 (1992) 8552-8563; C.A., 117 (1992) 98308p.
- 606 Escarceller, M., Rodriguez-Frias, F., Jardi, R., San Segundo, B. and Eritja, R.: Detection of hepatitis B virus DNA in human serum samples: use of digoxigenin-labeled oligonucleotides as modified primers for the polymerase chain reaction. *Anal. Biochem.*, 206 (1992) 36-42.
- 607 Fawcett, J.S., Wheeler, D. and Chrambach, A.: Transverse agarose pore gradient gel electrophoresis of DNA. *J. Biochem. Biophys. Methods*, 24 (1992) 181-194; C.A., 117 (1992) 65909g.
- 608 Ferec, C., Audrezet, M.P., Mercier, B., Guillermit, H., Moulhier, P., Quere, I. and Verlingue, C.: Detection of over 98% cystic fibrosis mutations in a Celtic population. *Nat. Genet.*, 1 (1992) 188-191; C.A., 117 (1992) 84305y.
- 609 Gingras, F. and Labuda, D.: Efficiency of biotin end-labeling of DNA fragments is rapidly tested by gel retardation with streptavidin analog. *BioTechniques*, 11 (1991) 342-343; C.A., 117 (1992) 167013c.
- 610 Goyal, D.: A simplified method for screening and characterization of plasmid DNA in cyanobacteria. *J. Microbiol. Methods*, 15 (1992) 7-15; C.A., 117 (1992) 22739j.
- 611 Grossman, P.D.: Low-viscosity polymer solution for capillary electrophoresis of DNA or other biopolymers. U.S. 5,126,021 (Cl. 204-180.1; GO1N27/26), 30 Jun. 1992, Appl. 731,771, 17 Jul. 1991; 13 pp.; C.A., 117 (1992) 107799d.
- 612 Guttman, A., Wanders, B. and Cooke, N.: Enhanced separation of DNA restriction fragments by capillary gel electrophoresis using field strength gradients. *Anal. Chem.*, 64 (1992) 2348-2351.
- 613 Hashimoto, T., Suzuki, K., Mizuta, K. and Otaka, E.: Yeast ribosomal proteins: XIV. Complete nucleotide sequences of the two genes encoding *Saccharomyces cerevisiae* YL16. *Biochim. Biophys. Acta*, 1132 (1992) 195-198.
- 614 Heller, C. and Beck, S.: Field inversion gel electrophoresis in denaturing polyacrylamide gels. *Nucleic Acids Res.*, 20 (1992) 2447-2452; C.A., 117 (1992) 65901y.
- 615 Herget, T., Brooks, S.F., Broad, S. and Rozengurt, E.: Relationship between the major protein kinase C substrates acidic 80-kDa protein-kinase-C substrate (80K) and myristoylated alanine-rich C-kinase substrate (MARCKS). Members of a gene family or equivalent genes in different species. *Eur. J. Biochem.*, 209 (1992) 7-14.
- 616 Hirotsune, S., Hatada, I., Komatsubara, H., Nagai, H., Kuma, K., Kobayakawa, K., Kawara, T., Nakagawara, A., Fujii, K., Mukai, T. and Hayashizaki, Y.: New approach for detection of amplification in cancer DNA using restriction landmark genomic scanning. *Cancer Res.*, 52 (1992) 3642-3647.
- 617 Honeycutt, R.J., Sobral, B.W.S., McClelland, M. and Atherly, A.G.: Analysis of large DNA from soybean (*Glycine max* L. Merr.) by pulsed-field gel electrophoresis. *Plant J.*, 2 (1992) 133-135; C.A., 117 (1992) 125465w.
- 618 Huan, L.J., Xu, G., Forstner, G. and Forstner, J.: A serine, threonine and proline-rich region near the carboxyl-terminus of a rat intestinal mucin peptide. *Biochim. Biophys. Acta*, 1132 (1992) 79-82.
- 619 Imamura, Y., Iguchi-Arigo, S.M.M. and Ariga, H.: The upstream region of the mouse N-myc gene: identification of an enhancer element that functions preferentially in neuroblastoma IMR32 cells. *Biochim. Biophys. Acta*, 1132 (1992) 177-187.
- 620 Kapp, U. and Langowski, J.: Preparation of DNA topoisomers by RP-18 high-performance liquid chromatography. *Anal. Biochem.*, 206 (1992) 293-299.
- 621 Kato, M. and Shimizu, N.: Effect of the potential triplex DNA region on the *in vivo* expression of bacterial β -lactamase gene in superhelical recombinant plasmids. *J. Biochem. (Tokyo)*, 112 (1992) 492-494.
- 622 Kawada, S.-Z., Yamashita, Y., Uosaki, Y., Gomi, K., Iwasaki, T., Takiguchi, T. and Nakano, H.: UCT4B, a new antitumor antibiotic with topoisomerase II mediated DNA cleavage activity, from *Streptomyces* sp. *J. Antibiot.*, 45 (1992) 1182-1184.
- 623 Kotaka, T.: (New attempt to fractionation of macromolecular DNA). *Kagaku to Kogyo (Tokyo)*, 45 (1992) 938-940; C.A., 117 (1992) 41432v.
- 624 Lahiri, A. and Sen, S.: A semiempirical expression for the gel electrophoretic mobility of supercoiled DNA. *Biopolymers*, 32 (1992) 893-896; C.A., 117 (1992) 107651z.
- 625 Lahiri, A., Chaudhuri, S., Bhattacharyya, N.P.: Theoretical analysis of gel electrophoretic data for interaction of lysine rich histone with supercoiled DNA. *Biophys. Chem.*, 42 (1992) 223-228; C.A., 117 (1992) 22725b.
- 626 Lanan, M., Grossmann, D.W. and Morris, M.D.: Noninvasive imaging of nanogram quantities of DNA in agarose electrophoresis gels. *Anal. Chem.*, 64 (1992) 1967-1972.
- 627 Mantyla, A.L., Rossi, K.H., Vanhanen, S.A., Penttila, M.E., Suominen, P.L. and Nevalainen, K.M.H.: Electrophoretic karyotyping of wild-type and mutant *Trichoderma longibrachiatum* (reesei) strains. *Curr. Genet.*, 21 (1992) 471-477; C.A., 117 (1992) 84455x.
- 628 Matsumoto, M. and Doi, M.: Dynamics of DNA in gel electrophoresis. *AIP Conf. Proc.*, 256(Slow Dyn. Condens. Matter) (1992) 292-293; C.A., 117 (1992) 127645d.

- 629 Mäueler, W., Frank, G., Siedlaczek, I., Epplen, J.T. and Melmer, G.: PCR amplification products are of limited use for the study of DNA/protein interaction. *Electrophoresis (Weinheim)*, 13 (1992) 641-643.
- 630 Mayer, A., Sunzeri, F., Lee, T.H. and Busch, M.P.: Separation and detection of DNA polynucleotides using capillary electrophoresis: application to detection of polymerase chain reaction-amplified human immunodeficiency virus and HLA DNA. *Arch. Pathol. Lab. Med.*, 115 (1991) 1228-1234; *C.A.*, 117 (1992) 86199r.
- 631 Meersseman, G., Pennings, S. and Bradbury, E.M.: Mobile nucleosomes - a general behavior. *EMBO J.*, 11 (1992) 2951-2959; *C.A.*, 117 (1992) 165149c.
- 632 Merkulov, V.M. and Merkulova, T.I.: Nucleotide sequence of a fragment of the rat tryptophan oxygenase gene showing high affinity to glucocorticoid receptor *in vitro*. *Biochim. Biophys. Acta*, 1132 (1992) 100-102.
- 633 Michaelis, C. and Weeks, G.: Isolation and characterization of a cdc2 cDNA from *Dictyostelium discoideum*. *Biochim. Biophys. Acta*, 1132 (1992) 35-42.
- 634 Nobori, T., Reynolds, L., Orvis, L., Wu, D. and Carson, D.A.: A polymerase chain reaction-based method for isolation of gene-specific sequences from the interferon- α gene cluster. *Anal. Biochem.*, 205 (1992) 42-46.
- 635 Noolandi, J.: Theory of DNA gel electrophoresis. *Adv. Electrophor.*, 5 (1992) 1-57; *C.A.*, 117 (1992) 127459w - a review with 47 refs.
- 636 Pramatarova, A., Michaud, L., Gagnon, F. and Hamelin, C.: Electrophoretic transfer of DNA from gels to filters by using a homemade apparatus. *Genet. Anal.: Tech. Appl.*, 9 (1992) 26-30; *C.A.*, 117 (1992) 125483a.
- 637 Pulyaeva, H., Wheeler, D., Garner, M.M. and Chrambach, A.: Molecular sieving of lambda phage DNA in polyacrylamide solutions as a function of the molecular weight of the polymer. *Electrophoresis (Weinheim)*, 13 (1992) 608-614.
- 638 Riesner, D., Steger, G., Wiese, U., Wulfert, M., Heibey, M. and Henco, K.: Temperature-gradient gel electrophoresis for the detection of polymorphic DNA and for quantitative polymerase chain reaction. *Electrophoresis (Weinheim)*, 13 (1992) 632-636.
- 639 Sagawa, I. and Nagata, Y.: Analysis of chromosomal DNA of mushrooms in genus *Pleurotus* by pulse-field gel electrophoresis. *J. Gen. Appl. Microbiol.*, 38 (1992) 47-52; *C.A.*, 117 (1992) 86826t.
- 640 Sasaki, M., Konno, R., Nishio, M., Niwa, A., Yasumura, Y. and Enami, J.: A single-base-pair substitution abolishes D-aminooxidase activity in the mouse. *Biochim. Biophys. Acta*, 1139 (1992) 315-318.
- 641 Semenov, A.N. and Subbotin, A.V.: Mobility of a charged Rouse chain in gels under electric field. *Europhys. Lett.*, 19 (1992) 161-164; *C.A.*, 117 (1992) 107648d.
- 642 Serwer, P., Hayes, S.J., Moreno, E.T. and Park, C.Y.: A small (58-nm) attached sphere perturbs the sieving of 40-80-kilobase DNA in 0.2-2.5% agarose gel: analysis of bacteriophage T3 capsid-DNA complexes by use of pulsed field electrophoresis. *Biochemistry*, 31 (1992) 8397-8405.
- 643 Sheppard, R.D., Montagutelli, X., Xavier, J.W.C., Tsai, J.Y., Rose, A., Guenet, J.L., Cole, M.D. and Silver, L.M.: Two-dimensional gel analysis of complex DNA families: methodology and apparatus. *Mamm. Genome*, 1 (1991) 104-111; *C.A.*, 117 (1992) 63914f.
- 644 Steffensen, B., Magnuson, V.L., Potempa, C.L., Chen, D. and Klebe, R.J.: α_5 Integrin subunit expression changes during myogenesis. *Biochim. Biophys. Acta*, 1137 (1992) 95-100.
- 645 Tada, A., Ohashi, T., Yamagata, K., Shirasaki, Y., Okada, M., Fukushima, S. and Abe, H.: Detection of gene by electrophoresis. *Jpn. Kokai Tokkyo Koho JP 04 84,751 [92 84,751] (Cl. GO1N27/447)*, 18 Mar. 1992, Appl. 90/200,607, 27 Jul. 1990; 5 pp.; *C.A.*, 117 (1992) 41914d.
- 646 Talaska, G., Roh, J.H. and Getek, T.: ^{32}P -Postlabelling and mass spectrometric methods for analysis of bulky, polyaromatic carcinogen-DNA adducts in humans. *J. Chromatogr.*, 580 (1992) 293-323 - a review with 144 refs.
- 647 Tietz, D., Aldroubi, A., Pulyaeva, H., Guszczynski, T., Garner, M.M. and Chrambach, A.: Advances in DNA electrophoresis in polymer solutions. *Electrophoresis (Weinheim)*, 13 (1992) 614-616.
- 648 Vijayalaxmi, Tice, R.R. and Strauss, G.H.S.: Assessment of radiation-induced DNA damage in human blood lymphocytes using the single-cell gel electrophoresis technique. *Mutat. Res.*, 271 (1992) 243-252; *C.A.*, 117 (1992) 65641p.
- 649 Viovy, J.L., Duke, T. and Caron, F.: The physics of DNA electrophoresis. *Contemp. Phys.*, 33 (1992) 25-40; *C.A.*, 117 (1992) 127470t - a review with many refs.
- 650 Volkmuth, W.D. and Austin, R.H.: DNA electrophoresis in microlithographic arrays. *Nature (London)*, 358 (1992) 600-602; *C.A.*, 117 (1992) 127655g.
- 651 Wheeler, D., Orban, L., Garner, M.M. and Chrambach, A.: Computer-aided analysis of DNA curves on transverse gradient gels. *J. Biochem. Biophys. Methods*, 24 (1992) 171-180; *C.A.*, 117 (1992) 84307a.
- 652 Wheeler, D., Tietz, D. and Chrambach, A.: Information on DNA conformation derived from transverse pore gradient gel electrophoresis in conjunction with an advanced data analysis applied to capillary electrophoresis in polymer media. *Electrophoresis (Weinheim)*, 13 (1992) 604-608.
- 653 Whitaker, S.J. and McMillan, T.J.: Pulsed-field gel electrophoresis in the measurement of DNA double-strand break repair in xrs-6 and CHO cell lines: DNA degradation under some conditions interferes with the assessment of double-strand break rejoining. *Radiat. Res.*, 130 (1992) 389-392; *C.A.*, 117 (1992) 43584v.
- 654 White, H.W.: Rapid separation of DNA molecules by agarose gel electrophoresis: use of a new agarose matrix and a survey of running buffer effects. *BioTechniques*, 12 (1992) 574-579; *C.A.*, 117 (1992) 21222y.
- 655 Wilson, J.M., Grossman, M., Cabrera, J.A., Wu, C.H. and Wu, G.Y.: A novel mechanism for achieving transgene persistence *in vivo* after somatic gene transfer into hepatocytes. *J. Biol. Chem.*, 267 (1992) 11483-11489.
- 656 Yan, T., Wu, J., Sun, L. and Wei, K.: (Quantitation of single strand breaks in nonradioactive DNA by alkaline gel electrophoresis). *Shengwu Huaxue Yu Shengwu Wuli Jinzhan*, 18 (1991) 466-467; *C.A.*, 117 (1992) 43860g.
- 657 Yoo, H., Li, L., Sacks, P.G., Thompson, L.H., Becker, F.F. and Chan, J.Y.-H.: Alterations in expression and structure of the DNA repair gene XRCC1. *Biochem. Biophys. Res. Commun.*, 186 (1992) 900-910.

See also 56, 83, 146, 315, 571, 574, 581, 587, 589, 687, 694.

21d. Structural studies on RNA and RNA mapping

- 658 Fukumoto, F.: Difference in electrophoretic pattern of the rice dwarf virus genome segments in agarose and polyacrylamide gel electrophoresis. *Nippon Shokubutsu Byori Gakkaiho*, 58 (1992) 83-86; C.A., 117 (1992) 66144r.
- 659 Nakato, H., Izumi, S. and Tomino, S.: Structure and expression of gene coding for a pupal cuticle protein of *Bombyx mori*. *Biochim. Biophys. Acta*, 1132 (1992) 161-167.
- 660 Schoonmaker, D., Heimberger, T. and Birkhead, G.: Comparison of ribotyping and restriction enzyme analysis using pulsed-field gel electrophoresis for distinguishing *Legionella pneumophila* isolates obtained during a nosocomial outbreak. *J. Clin. Microbiol.*, 30 (1992) 1491-1498; C.A., 117 (1992) 44003s.
- 661 Shimba, S., Buckley, B., Reddy, R., Kiss, T. and Filipowicz, W.: Cap structure of U3 small nucleolar RNA in animal and plant cells is different. γ -Monomethyl phosphate cap structure in plant RNA. *J. Biol. Chem.*, 267 (1992) 13772-13777.
- 662 Shimeld, S.M. and Sharpe, P.T.: Cell-surface changes induced by ectopic expression of the murine homeobox gene Hox-3.3. *Biochim. Biophys. Acta*, 1136 (1992) 253-258.
- 663 Takasawa, S., Tohgo, A., Unno, M., Shiga, K., Yonekura, H. and Okamoto, H.: The primary structure of rat rig/ribosomal protein S15 gene. *Biochim. Biophys. Acta*, 1132 (1992) 228-230.
- 664 Tsuzuki, S., Miura, Y., Fushiki, T., Oomori, T., Satoh, T., Natori, Y. and Sugimoto, E.: Molecular cloning and characterization of genes encoding rat pancreatic cholecystokinin (CCK)-releasing peptide (monitor peptide) and pancreatic secretory trypsin inhibitor (PSTI). *Biochim. Biophys. Acta*, 1132 (1992) 199-202.
- See also 584, 587.
- 21e. Structural studies on DNA and DNA mapping
- 665 Ansoerge, W., Voss, H., Wiemann, S., Schwager, C., Sproat, B., Zimmermann, J., Stegemann, J., Erfle, H., Hewitt, N. and Rupp, T.: High-throughput automated DNA sequencing facility with fluorescent labels at the European Molecular Biology Laboratory. *Electrophoresis (Weinheim)*, 13 (1992) 616-619.
- 666 Bertenshaw, R., Sarne, D., Tornari, J., Weinberg, M. and Refetoff, S.: Sequencing of the variant thyroxine-binding globulin (TBG)-San Diego reveals two nucleotide substitutions. *Biochim. Biophys. Acta*, 1139 (1992) 307-310.
- 667 Bettink, S., Wullich, B., Christmann, A., Zwergel, T., Zang, K.-D. and Unteregger, G.: Genetic heterogeneity of prostatic carcinoma-derived cell lines as emphasized by DNA fingerprinting. *Electrophoresis (Weinheim)*, 13 (1992) 644-646.
- 668 Birkelund, S. and Stephens, R.S.: Construction of physical and genetic maps of *Chlamydia trachomatis* serovar L2 by pulsed-field gel electrophoresis. *J. Bacteriol.*, 174 (1992) 2742-2747; C.A., 117 (1992) 64055p.
- 669 Chiari, M., Nesi, M., Fazio, M. and Righetti, P.G.: Capillary electrophoresis of macromolecules in 'syrupy' solutions: facts and misfacts. *Electrophoresis (Weinheim)*, 13 (1992) 690-697.
- 670 Coffin, J.W., Condon, C., Compston, C.A., Potter, K.N., Lamontagne, L.R., Shafiq, J. and Kunimoto, D.Y.: Use of restriction fragment length polymorphisms resolved by pulsed-field gel electrophoresis for subspecies identification of mycobacteria in the *Mycobacterium avium* complex and for isolation of DNA probes. *J. Clin. Microbiol.*, 30 (1992) 1829-1836; C.A., 117 (1992) 66360h.
- 671 Drury, H.A., Politte, D.G., Ollinger, J.M., Green, P. and Thomas, L.J., Jr.: Maximum-likelihood estimation of restriction-fragment mobilities from 1-D electrophoretic agarose gels. *Proc. SPIE-Int. Soc. Opt. Eng.*, 1660(Biomed. Image Process. Three-Dimens. Microsc., Pt. 2) (1992) pp. 564-575; C.A., 117 (1992) 167018h.
- 672 Gomolka, M., Schwaiger, W., Weyers, E., Epplen, C., Buitkamp, J. and Epplen, J.T.: Towards a fingerprint method covering the immunological genome. *Electrophoresis (Weinheim)*, 13 (1992) 623-625.
- 673 Grossman, P.D., Menchen, S. and Hershey, D.: Quantitative analysis of DNA-sequencing electrophoresis. *Genet. Anal.: Tech. Appl.*, 9 (1992) 9-16; C.A., 117 (1992) 125482z.
- 674 Guo, Q., Lu, M. and Kallenbach, N.R.: Adenine affects the structure and stability of telomeric sequences. *J. Biol. Chem.*, 267 (1992) 15293-15300.
- 675 Harke, H.R., Bay, S., Zhang, J.Z., Rocheleau, M.J. and Dovichi, N.J.: Effect of total percent polyacrylamide in capillary gel electrophoresis for DNA sequencing of short fragments. A phenomenological model. *J. Chromatogr.*, 608 (1992) 143-150.
- 676 Herrmann, J.L., Bellenger, E., Perolat, P., Baranton, G. and Saint Girons, I.: Pulsed-field gel electrophoresis of NotI digests of leptospiral DNA: a new rapid method of serovar identification. *J. Clin. Microbiol.*, 30 (1992) 1696-1702; C.A., 117 (1992) 86204p.
- 677 Hoehm, H. and Karch, H.: DNA fingerprinting of *Escherichia coli* O157:H7 strains by pulsed-field gel electrophoresis. *J. Clin. Microbiol.*, 30 (1992) 2169-2172; C.A., 117 (1992) 107629y.
- 678 Huang, X.C., Quesada, M.A. and Mathies, R.A.: DNA sequencing using capillary array electrophoresis. *Anal. Chem.*, 64 (1992) 2149-2154.
- 679 Johnson, C.A., Forster, T.H., Winterford, C.M. and Allan, D.J.: Hydroxyurea induces apoptosis and regular DNA fragmentation in a Burkitt's lymphoma cell line. *Biochim. Biophys. Acta*, 1136 (1992) 1-4.
- 680 Kumaresan, K.R., Ramaswamy, M. and Yeung, A.T.: Structure of the DNA interstrand cross-link of 4,5',8-trimethylpsoralen. *Biochemistry*, 31 (1992) 6774-6783.
- 681 Liu, W.-M., Leeftang, E.P. and Schmid, C.W.: Unusual sequences of two old, inactive human Alu repeats. *Biochim. Biophys. Acta*, 1132 (1992) 306-308.
- 682 Matsumura, Y. and Tarin, D.: DNA fingerprinting survey of various human tumors and their metastases. *Cancer Res.*, 52 (1992) 2174-2179.
- 683 Melmer, G.: Simple repeats are not found more abundantly at G/C-rich regions. *Electrophoresis (Weinheim)*, 13 (1992) 639-640.
- 684 Nakamoto, M., Matsubara, S., Miyauchi, T., Obama, H., Ozawa, M. and Muramatsu, T.: A new family of heparin binding growth/differentiation factors: differential expression of the midkine (MK) and HB-GAM genes during mouse development. *J. Biochem. (Tokyo)*, 112 (1992) 346-349.

- 685 Ono, A., Ohdoi, C., Matsuda, A. and Ueda, T.: Nucleosides and nucleotides. 105. DNA bending in d(A)₄-d(T)₄ tracts containing 3-deazaadenine or 7-deazaadenine substituted for adenine. *Nucleosides Nucleotides*, 11 (1992) 227-235; *C.A.*, 117 (1992) 90693q.
- 686 Pierce, D.J., Jordan, E.M. and Breen, G.A.M.: Structural organization of a nuclear gene for the α -subunit of the bovine mitochondrial ATP synthase complex. *Biochim. Biophys. Acta*, 1132 (1992) 265-275.
- 687 Rodríguez, I.R., Gonzalez, P., Zigler, J.S., Jr. and Borrás, T.: A guinea-pig hereditary cataract contains a splice-site deletion in a crystallin gene. *Biochim. Biophys. Acta*, 1180 (1992) 44-52.
- 688 Römmling, U. and Tümmler, B.: A PacI/SwaI map of the *Pseudomonas aeruginosa* PAO chromosome. *Electrophoresis (Weinheim)*, 13 (1992) 649-651.
- 689 Römmling, U., Grothues, D., Heuer, T. and Tümmler, B.: Physical genome analysis of bacteria. *Electrophoresis (Weinheim)*, 13 (1992) 626-631.
- 690 Römmling, U., Grothues, D., Koopmann, U., Jahnke, B., Greipel, J. and Tümmler, B.: Pulsed-field gel electrophoresis analysis of a *Pseudomonas aeruginosa* pathovar. *Electrophoresis (Weinheim)*, 13 (1992) 646-648.
- 691 Schreiber, W.E. and Jamani, A.: Detection of a T/C polymorphism in the porphobilinogen deaminase gene by polymerase chain reaction amplification of specific alleles. *Clin. Chem. (Winston-Salem)*, 38 (1992) 2153-2155.
- 692 Turmel, C., Brassard, E. and Noolandi, J.: Manual sequencing using pulsed field. *Electrophoresis (Weinheim)*, 13 (1992) 620-622.
- 693 Winichagoon, P., Fucharoen, S., Wilairat, P., Chihara, K., Fukumaki, Y. and Wasi, P.: Identification of five rare mutations including a novel frameshift mutation causing β^0 -thalassemia/hemoglobin E disease. *Biochim. Biophys. Acta*, 1139 (1992) 280-286.
- 694 Wohl, T., Baur, M., Friedl, A.A. and Lottspeich, F.: Chromosomal localization of the HYP2-gene in *Saccharomyces cerevisiae* and use of pulsed-field gel electrophoresis for detection of irregular recombination events in gene disruption experiments. *Electrophoresis (Weinheim)*, 13 (1992) 651-653.
- 695 Yamamoto, K., Yee, C.C., Shirakawa, M. and Kyogoku, Y.: Characterization of the bacterially expressed *Drosophila engrailed* homeodomain. *J. Biochem. (Tokyo)*, 111 (1992) 793-797.
- 696 Yanase, T., Waterman, M.R., Zachmann, M., Winter, J.S.D., Simpson, E.R. and Kagimoto, M.: Molecular basis of apparent isolated 17,20-lyase deficiency: compound heterozygous mutations in the C-terminal region (Arg(496) \rightarrow Cys, Gln(461) \rightarrow Stop) actually cause combined 17 α -hydroxylase/17,20-lyase deficiency. *Biochim. Biophys. Acta*, 1139 (1992) 275-279.
- See also 315, 581, 587, 597, 612, 621, 629, 630, 632, 659.
- 21f. *Complex mixtures of nucleic acids and their fragments*
- 697 Garner, M.M.: Electrophoretic studies of DNA-protein interactions. *Adv. Electrophor.*, 5 (1992) 137-157; *C.A.*, 117 (1992) 127460q - a review with 52 refs.
- 698 Lässle, M., Richter, A. and Knippers, R.: Comparison of replicative and non-replicative chromatin assembly pathways in HeLa cell extracts. *Biochim. Biophys. Acta*, 1132 (1992) 1-10.
- See also 301, 625.
22. ALKALOIDS
- 699 Liu, Y.-M. and Sheu, S.-J.: Determination of quaternary alkaloids from *Coptidis rhizoma* by capillary electrophoresis. *J. Chromatogr.*, 623 (1992) 196-199.
- 700 Stuppner, H., Sturm, S. and Konwalinka, G.: Capillary electrophoretic analysis of oxindole alkaloids from *Uncaria tomentosa*. *J. Chromatogr.*, 609 (1992) 375-380.
- See also 737.
24. ORGANIC SULPHUR COMPOUNDS (INCL. GLUCOSINOLATES)
- 701 Desbène, P.L., Rony, C., Desmazieres, B. and Jacquier, J.C.: Analysis of alkylaromatic sulphonates by high-performance capillary electrophoresis. *J. Chromatogr.*, 608 (1992) 375-383.
- 702 Michaelsen, S., Moller, P. and Sorensen, H.: Factors influencing the separation and quantitation of intact glucosinolates and desulphoglucosinolates by micellar electrokinetic capillary chromatography. *J. Chromatogr.*, 608 (1992) 363-374.
25. ORGANIC PHOSPHORUS COMPOUNDS (INCL. SUGAR PHOSPHATES)
- 703 Henshall, A., Harrold, M.P. and Tso, J.M.Y.: Separation of inositol phosphates by capillary electrophoresis. *J. Chromatogr.*, 608 (1992) 413-419.
- See also 179, 485, 486, 498.
27. VITAMINS AND VARIOUS ANIMAL GROWTH FACTORS (NON-PEPTIDIC)
- 704 Lambert, D., Adjalla, C., Felden, F., Benhayoun, S., Nicolas, J.P. and Guéant, J.L.: Identification of vitamin B₁₂ and analogues by high-performance capillary electrophoresis and comparison with high-performance liquid chromatography. *J. Chromatogr.*, 608 (1992) 311-315.
- 705 Rizzolo, A. and Polesello, S.: Chromatographic determination of vitamins in foods. *J. Chromatogr.*, 624 (1992) 103-152 - a review with 286 refs.
28. ANTIBIOTICS
- 706 Boison, J.O.: Chromatographic methods of analysis for penicillins in food-animal tissues and their significance in regulatory programs for residue reduction and avoidance. *J. Chromatogr.*, 624 (1992) 171-194 - a review with 127 refs.
- 707 Carpenter, J.L., Camilleri, P., Dhanak, D. and Goodall, D.: A study of the binding of vancomycin to dipeptides using capillary electrophoresis. *J. Chem. Soc., Chem. Commun.*, (1992) 804-806; *C.A.*, 117 (1992) 70327v.

- 708 Hrdlicka, J.: (Identification of antibiotic residues in the tissues of slaughter animals by bioautographic detection electrophoresis). *Vet. Med. (Prague)*, 36 (1991) 695-701; C.A., 117 (1992) 88854z.

29. INSECTICIDES, PESTICIDES AND OTHER AGROCHEMICALS

29e. Herbicides

- 709 Desiderio, C. and Fanali, S.: Atrazine and simazine determination in river water samples by micellar electrokinetic capillary chromatography. *Electrophoresis (Weinheim)*, 13 (1992) 698-700.
- 710 Lucansky, D., Pukacova, A. and Dulak, K.: Isotachophoresis method for determination of ionogenic substances in manufacturing 5-amino-4-chloro-2-phenylpyridazin-3-one. *Czech CS 268,597 (Cl. G01N33/48)*, 31 Jul. 1990, Appl. 88/1,961, 25 Mar. 1988; 5 p.; C.A., 117 (1992) 82714g.
- 711 Wu, Q., Claessens, H.A. and Cramers, C.A.: The separation of herbicides by micellar electrophoretic capillary chromatography. *Chromatographia*, 34 (1992) 25-30.
- 712 Wui, Q., Claessens, H.A. and Cramers, C.A.: The separation of herbicides by micellar electrophoretic capillary chromatography. *Chromatographia*, 34 (1992) 25-30.

See also 102.

30. SYNTHETIC AND NATURAL DYES

30a. Synthetic dyes

- 713 Croft, S.N. and Lewis, D.M.: Analysis of reactive dyes and related derivatives using high-performance capillary electrophoresis. *Dyes Pigm.*, 18 (1992) 309-317; C.A., 117 (1992) 173369q.
- 714 Fanali, S. and Schudel, M.: Some separations of black and red water-soluble fiber-tip pen inks by capillary zone electrophoresis and thin-layer chromatography. *J. Forensic Sci.*, 36 (1991) 1192-1197; C.A., 117 (1992) 144917m.

30b. Chloroplast and other natural pigments

- 715 Kumagai, A. and Hirota, N.: Solubilization and isolation of chlorophyll-protein complexes from *Hizikia fusiformis*. *Nippon Suisan Gakkaishi*, 58 (1992) 1171-1178; C.A., 117 (1992) 107655d.
- 716 Lee, H.S. and Hong, V.: Chromatographic analysis of anthocyanins. *J. Chromatogr.*, 624 (1992) 221-234 - a review with 126 refs.

31. PLASTICS AND THEIR INTERMEDIATES

- 717 Gelfi, C., Alloni, A., de Besi, P. and Righetti, P.G.: Investigation of the properties of acrylamide bifunctional monomers (cross-linkers) by capillary zone electrophoresis. *J. Chromatogr.*, 608 (1992) 343-348.

- 718 Gelfi, C., de Besi, P., Alloni, A. and Righetti, P.G.: Investigation of the properties of novel acrylamido monomers by capillary zone electrophoresis. *J. Chromatogr.*, 608 (1992) 333-341.

See also 95.

32. DRUG ANALYSIS

32a. Drug analysis, general techniques

- 719 Evenson, M.A. and Wiktorowicz, J.E.: Automated capillary electrophoresis applied to therapeutic drug monitoring. *Clin. Chem. (Winston-Salem)*, 38 (1992) 1847-1852.
- 720 Gruebler, G., Mayer, H. and Voelter, W.: Capillary zone electrophoresis (CZE) for quality control of pharmaceuticals. Comparison of CZE with HPLC. *Bioforum*, 15 (1992) 50-51; C.A., 117 (1992) 178403f.
- 721 Klein, H. and Teichmann, R.: The application of capillary isotachophoresis in pharmaceutical analysis. *PZ Wiss.*, 5 (1992) 119-122; C.A., 117 (1992) 137743y - a review with 31 refs.
- 722 O'Shea, T.J., Telting-Diaz, M.W., Lunte, S.M., Lunte, C.E. and Smyth, M.R.: Capillary electrophoresis-electrochemistry of microdialysis samples for pharmacokinetic studies. *Electroanalysis (N.Y.)*, 4 (1992) 463-468; C.A., 117 (1992) 62196m.
- 723 Soini, H., Riekkola, M.-L. and Novotny, M.V.: Chiral separations of basic drugs and quantitation of bupivacaine enantiomers in serum by capillary electrophoresis with modified cyclodextrin buffers. *J. Chromatogr.*, 608 (1992) 265-274.
- 724 Tellez, S., Forges, N., Roussin, A. and Hernandez, L.: Coupling of microdialysis with capillary electrophoresis: a new approach to the study of drug transfer between two compartments of the body in freely moving rats. *J. Chromatogr.*, 581 (1992) 257-266.

32b. Antirheumatics and antiinflammatory drugs

- 725 D'Hulst, A. and Verbeke, N.: Chiral separation by capillary electrophoresis with oligosaccharides. *J. Chromatogr.*, 608 (1992) 275-287.

32c. Autonomic and cardiovascular drugs

- 726 Fanali, S., Cristalli, M., Nardi, A., Quaglia, M.G.: Capillary zone electrophoresis for drug analysis. Rapid determination of minoxidil in pharmaceutical formulations. *Farmaco*, 47 (1992) 711-722; C.A., 117 (1992) 157753e.
- 727 Prunonosa, J., Obach, R., Diez-Cascón, A. and Gouesclou, L.: Comparison of high-performance liquid chromatography and high-performance capillary electrophoresis for the determination of cicletanine in plasma. *J. Chromatogr.*, 581 (1992) 219-226.

See also 70, 723.

32d. Central nervous system drugs

- 728 Altria, K.D., Goodall, D.M. and Roban, M.M.: Chiral separation of β -amino alcohols by capillary electrophoresis using cyclodextrins as buffer additives. I. Effect of varying operating parameters. *Chromatographia*, 34 (1992) 19-24.

- 729 Lee, K.-J., Heo, G.S., Kim, N.J. and Moon, D.C.: Analysis of antiepileptic drugs in human plasma using micellar electrokinetic capillary chromatography. *J. Chromatogr.*, 608 (1992) 243-250.
- 730 Lloyd, D.K.: Quantitation of the stereoisomers of L-buthionine-(R,S)-sulfoximine in biological matrixes by capillary electrophoresis. *Anal. Proc. (London)*, 29 (1992) 169-170; *C.A.*, 117 (1992) 103433g.
- 731 Tellez, S., Forges, N., Roussin, A. and Hernandez, L.: Blood-brain barrier permeability to phenobarbital as studied by dual microdialysis coupled to capillary electrophoresis. In: Rollema, H., Westerink, B.H.C. and Drijfhout, W.J. (Editors), *Monit. Mol. Neurosci., Proc. Int. Conf. In Vivo Methods, 5th*, Univ. Cent. Pharm., Groningen, 1991, pp. 122-124; *C.A.*, 117 (1992) 82845a.
- See also 69, 723, 734.
- 32f. *Cytostatics*
- 732 Lukkari, P., Jumppanen, J., Jinno, K., Elo, H. and Riekkola, M.-L.: Determination of bis(amidinohydrazones) by micellar electrokinetic capillary chromatography. *J. Pharm. Biomed. Anal.*, 10 (1992) 561-565.
- 32g. *Other drug categories*
- 733 Shihabi, Z.K. and Constantinescu, M.S.: Iohexol in serum determined by capillary electrophoresis. *Clin. Chem. (Winston-Salem)*, 38 (1992) 2117-2120.
- See also 205, 737.
- 32h. *Toxicological and forensic applications*
- 734 Wernly, P. and Thormann, W.: Confirmation testing of 11-nor- Δ^9 -tetrahydrocannabinol-9-carboxylic acid in urine with micellar electrokinetic capillary chromatography. *J. Chromatogr.*, 608 (1992) 251-256.
- 735 Wernly, P. and Thormann, W.: Drug of abuse confirmation in human urine using stepwise solid-phase extraction and micellar electrokinetic capillary chromatography. *Anal. Chem.*, 64 (1992) 2155-2159.
- See also 208, 371, 646.
- 32i. *Plant extracts*
- See 699.
33. CLINICO-CHEMICAL APPLICATIONS
- 33a. *General papers and reviews*
- See 9, 719.
- 33b. *Complex mixtures and profiling (single compounds by cross-reference only)*
- See
- 141, 142, 143, 146, 147, 148, 149, 289, 291, 295, 306, 309, 310, 314, 327, 359, 424, 454, 474, 490, 501, 505, 512, 513, 514, 516, 524, 533, 539, 593, 691.
34. FOOD ANALYSIS
- 34a. *General papers and reviews*
- See 159, 255, 374, 705, 706, 716.
- 34b. *Complex mixtures (single compounds by cross-reference only)*
- 736 Amigo, L., Ramos, M., Calhau, L. and Barbosa, M.: Comparison of electrophoresis, isoelectric focusing, and immunodiffusion in determinations of cow's and goat's milk in Serra da Estrela cheeses. *Lait*, 72 (1992) 95-101; *C.A.*, 117 (1992) 68563a.
- 737 Pria, P.P., Ranghieri, M.C. and Schlabach, T.X.: (Multiscan capillary electrophoresis in the analysis of aspartame, caffeine and benzoic acid in beverages). *Lab. 2000*, 5 (1991) 60-61; *C.A.*, 117 (1992) 130048s.
- See also 108, 708, 751.
35. ENVIRONMENTAL ANALYSIS
- 35a. *General papers and reviews*
- 738 Takeda, S., Wakida, S., Yamane, M., Kawahara, A. and Higashi, K.: (Analysis of environmental pollutants by electrokinetic chromatography). *Kankyo Kagaku*, 2 (1992) 418-419; *C.A.*, 117 (1992) 137210x.
- 35c. *Water pollution (complex mixtures; single compounds by cross-reference only)*
- See 709.
36. SOME TECHNICAL PRODUCTS AND COMPLEX MIXTURES
- 739 Tong, W.G.: Sensitive analytical detection method based on nonlinear laser wave mixing inside small capillary low detector cells. *Proc. Int. Conf. Lasers 1991, 14th*, 1992, pp. 816-820; *C.A.*, 117 (1992) 183821v.
- 36a. *Surfactants*
- 740 Tribet, C., Gaboriaud, R. and Gareil, P.: Determination of C_8 - C_{20} saturated anionic and cationic surfactant mixtures by capillary isotachophoresis with conductivity detection. *J. Chromatogr.*, 609 (1992) 381-390.
- See also 163.

36c. Complex mixtures, technical products and unidentified compounds

- 741 Ewing, A.G.: Monitoring molecules in single neurons. In: Rollema, H., Westerink, B.H.C. and Drijfhout, W.J. (Editors), *Monit. Mol. Neurosci., Proc. Int. Conf. In Vivo Methods, 5th 1991*, Univ. Cent. Pharm., Groningen, 1991, pp. 3-5; C.A., 117 (1992) 127636b.

37. CELLS, CELLULAR PARTICLES AND SUPRAMOLECULAR STRUCTURES

- 742 Boxall, C. and Kelsall, G.H.: Photoelectrophoresis of colloidal semiconductors. Part III. Transient experiments on tin dioxide particles. *J. Electroanal. Chem.*, 328 (1992) 75-97; C.A., 117 (1992) 138452q.
- 743 Chen, S.B. and Keh, H.J.: Axisymmetric electrophoresis of multiple colloidal spheres. *J. Fluid Mech.*, 238 (1992) 251-276; C.A., 117 (1992) 138426j.
- 744 Costas, M.: Classification, identification and typing of bacteria by the analysis of their one-dimensional polyacrylamide gel electrophoretic protein patterns. *Adv. Electrophor.*, 5 (1992) 351-408; C.A., 117 (1992) 107452k - a review with 187 refs.
- 745 Dunstan, D.E. and Saville, D.A.: Electrophoretic mobility of colloidal alkane particles in electrolyte solutions. *J. Chem. Soc., Faraday Trans.*, 88 (1992) 2031-2033; C.A., 117 (1992) 138289s.
- 746 Hawker, D.T.L., Todd, P., Davis, R.H., Lawson, R.C. and Rudge, S.R.: Electrokinetic isolation of vesicles and ribosomes derived from *Serratia marcescens*. *Biotechnol. Prog.*, 8 (1992) 429-435; C.A., 117 (1992) 169476e.
- 747 Heaton, L.A.: Use of agarose gel electrophoresis to monitor conformational changes of some small, spherical plant viruses. *Phytopathology*, 82 (1992) 803-807; C.A., 117 (1992) 167015e.
- 748 Sasaki, H.: (Sieving of submicron particles by electrophoresis). *Kagaku Kogyo*, 43 (1992) 592-599; C.A., 117 (1992) 72225j - a review with 16 refs.
- 749 Sor, F.: Electrophoretic techniques for chromosome separation. *Methods Mol. Cell. Biol.*, 3 (1992) 65-70; C.A., 117 (1992) 105220j.

See also 61, 559.

38. INORGANIC COMPOUNDS

38a. Cations

- 750 Bächmann, K., Haumann, I. and Groh, T.: Simultaneous determination of inorganic cations and anions in capillary zone electrophoresis (CZE) with indirect fluorescence detection. *Fresenius J. Anal. Chem.*, 343 (1992) 901-902.
- 751 Singh, D., Sharma, G.P., Tyagi, B.P. and Dutt, P.: Determination of lead in chewing-gum by electrophoresis. *Sci. Phys. Sci.*, 4 (1992) 79-81; C.A., 117 (1992) 110248d.
- 752 Weston, A., Brown, P.R., Jandik, P., Heckenberg, A.L. and Jones, W.R.: Optimization of detection sensitivity in the analysis of inorganic cations by capillary ion electrophoresis using indirect photometric detection. *J. Chromatogr.*, 608 (1992) 395-402.

38b. Anions

- 753 Buchberger, W. and Haddad, P.R.: Effects of carrier electrolyte composition on separation selectivity in capillary zone electrophoresis of low-molecular-mass anions. *J. Chromatogr.*, 608 (1992) 59-64.
- 754 Carpio, R.A., Mariscal, R. and Welch, J.: Determination of boron and phosphorus in borophosphosilicate thin films on silicon substrates by capillary electrophoresis. *Anal. Chem.*, 64 (1992) 2123-2129.
- 755 Jones, W.R. and Jandik, P.: Various approaches to analysis of difficult sample matrices of anions using capillary ion electrophoresis. *J. Chromatogr.*, 608 (1992) 385-393.
- 756 Weber, I.: Electrophoretic method for separation of a calcium difluoridelithium monofluoride mixture on an electrical conductor. *Ger. (East)* DD 299,742 (Cl. C25D9/04), 07 May 1992, Appl. 168,341, 04 Jan. 1973; 2 p.; C.A., 117 (1992) 123715d.

See also 46, 102, 750.

PUBLICATION SCHEDULE FOR THE 1993 SUBSCRIPTION

Journal of Chromatography and Journal of Chromatography, Biomedical Applications

MONTH	O 1992	N 1992	D 1992	J	F	M	A
Journal of Chromatography	623/1 623/2 624/1 + 2	625/1 625/2	626/1 626/2 627/1 + 2	628/1 628/2 629/1 629/2	630/1 + 2 631/1 + 2 632/1 + 2 633/1 + 2	634/1 634/2	635/1 635/2 636/1 636/2 637/1
Cumulative Indexes, Vols. 601-650							601/1 + 2
Bibliography Section						649/1	
Biomedical Applications				612/1	612/2	613/1	613/2 614/1

The publication schedule for further issues will be published later.

INFORMATION FOR AUTHORS

(Detailed *Instructions to Authors* were published in Vol. 609, pp. 437-443. A free reprint can be obtained by application to the publisher, Elsevier Science Publishers B.V., P.O. Box 330, 1000 AH Amsterdam, The Netherlands.)

Types of Contributions. The following types of papers are published in the *Journal of Chromatography* and the section on *Biomedical Applications*: Regular research papers (Full-length papers), Review articles, Short Communications and Discussions. Short Communications are usually descriptions of short investigations, or they can report minor technical improvements of previously published procedures; they reflect the same quality of research as Full-length papers, but should preferably not exceed five printed pages. Discussions (one or two pages) should explain, amplify, correct or otherwise comment substantively upon an article recently published in the journal. For Review articles, see inside front cover under Submission of Papers.

Submission. Every paper must be accompanied by a letter from the senior author, stating that he/she is submitting the paper for publication in the *Journal of Chromatography*.

Manuscripts. Manuscripts should be typed in **double spacing** on consecutively numbered pages of uniform size. The manuscript should be preceded by a sheet of manuscript paper carrying the title of the paper and the name and full postal address of the person to whom the proofs are to be sent. As a rule, papers should be divided into sections, headed by a caption (*e.g.*, Abstract, Introduction, Experimental, Results, Discussion, etc.). All illustrations, photographs, tables, etc., should be on separate sheets.

Abstract. All articles should have an abstract of 50-100 words which clearly and briefly indicates what is new, different and significant. No references should be given.

Introduction. Every paper must have a concise introduction mentioning what has been done before on the topic described, and stating clearly what is new in the paper now submitted.

Illustrations. The figures should be submitted in a form suitable for reproduction, drawn in Indian ink on drawing or tracing paper. Each illustration should have a legend, all the *legends* being typed (with double spacing) together on a *separate sheet*. If structures are given in the text, the original drawings should be supplied. Coloured illustrations are reproduced at the author's expense, the cost being determined by the number of pages and by the number of colours needed. The written permission of the author and publisher must be obtained for the use of any figure already published. Its source must be indicated in the legend.

References. References should be numbered in the order in which they are cited in the text, and listed in numerical sequence on a separate sheet at the end of the article. Please check a recent issue for the layout of the reference list. Abbreviations for the titles of journals should follow the system used by *Chemical Abstracts*. Articles not yet published should be given as "in press" (journal should be specified), "submitted for publication" (journal should be specified), "in preparation" or "personal communication".

Dispatch. Before sending the manuscript to the Editor please check that the envelope contains four copies of the paper complete with references, legends and figures. One of the sets of figures must be the originals suitable for direct reproduction. Please also ensure that permission to publish has been obtained from your institute.

Proofs. One set of proofs will be sent to the author to be carefully checked for printer's errors. Corrections must be restricted to instances in which the proof is at variance with the manuscript. "Extra corrections" will be inserted at the author's expense.

Reprints. Fifty reprints will be supplied free of charge. Additional reprints can be ordered by the authors. An order form containing price quotations will be sent to the authors together with the proofs of their article.

Advertisements. The Editors of the journal accept no responsibility for the contents of the advertisements. Advertisement rates are available on request. Advertising orders and enquiries can be sent to the Advertising Manager, Elsevier Science Publishers B.V., Advertising Department, P.O. Box 211, 1000 AE Amsterdam, Netherlands; courier shipments to: Van de Sande Bakhuizenstraat 4, 1061 AG Amsterdam, Netherlands; Tel. (+31-20) 515 3220/515 3222, Telefax (+31-20) 6833 041, Telex 16479 els vi nl. UK: T. G. Scott & Son Ltd., Tim Blake, Portland House, 21 Narborough Road, Cosby, Leics. LE9 5TA, UK; Tel. (+44-533) 753 333, Telefax (+44-533) 750 522. USA and Canada: Weston Media Associates, Daniel S. Lipner, P.O. Box 1110, Greens Farms, CT 06436-1110, USA; Tel. (+1-203) 261 2500, Telefax (+1-203) 261 0101.

Capillary Electrophoresis

Principles, Practice and Applications

by S.F.Y. Li, National University of Singapore, Singapore

Journal of Chromatography Library Volume 52

Capillary Electrophoresis (CE) has had a very significant impact on the field of analytical chemistry in recent years as the technique is capable of very high resolution separations, requiring only small amounts of samples and reagents. Furthermore, it can be readily adapted to automatic sample handling and real time data processing. Many new methodologies based on CE have been reported. Rapid, reproducible separations of extremely small amounts of chemicals and biochemicals, including peptides, proteins, nucleotides, DNA, enantiomers, carbohydrates, vitamins, inorganic ions, pharmaceuticals and environmental pollutants have been demonstrated. A wide range of applications have been developed in greatly diverse fields, such as chemical, biotechnological, environmental and pharmaceutical analysis.

This book covers all aspects of CE, from the principles and technical aspects to the most important applications. It is intended to meet the growing need for a thorough and balanced treatment of CE. The book will serve as a comprehensive reference work and can also be used as a textbook for advanced undergraduate and graduate courses. Both the experienced analyst and the newcomer will find the text useful.

Contents:

1. Introduction. Historical Background. Overview of High Performance CE. Principles of Separations. Comparison with Other Separation Techniques.
2. Sample Injection Methods.

Introduction. Electrokinetic Injection. Hydrodynamic Injection. Electric Sample Splitter. Split Flow Syringe Injection System. Rotary Type Injector. Freeze Plug Injection. Sampling Device with Feeder. Microinjectors. Optical Gating. **3. Detection Techniques.** Introduction. UV-Visible Absorbance Detectors. Photodiode Array Detectors. Fluorescence Detectors. Laser-based Thermo-optical and Refractive Index Detectors. Indirect Detection. Conductivity Detection. Electrochemical Detection. Mass Spectrometric Detection. **4. Column Technology.** Uncoated Capillary Columns. Coated Columns. Gel-filled Columns. Packed Columns. Combining Packed and Open-Tubular Column. **5. Electrophoretic Media.** Electrophoretic Buffer Systems. Micellar Electrokinetic Capillary Chromatography. Inclusion Pseudophases. Metal-complexing Pseudophases. Other Types of Electrophoretic Media. **6. Special Systems and Methods.** Buffer Programming. Fraction Collection. Hyphenated Techniques. Field Effect Electroosmosis. Systematic Optimization of Separation. **7. Applications of CE.** Biomolecules. Pharmaceutical and Clinical Analysis. Inorganic Ions. Hydrocarbons. Foods and Drinks. Environmental Pollutants. Carbohydrates. Toxins. Polymers

and Particles. Natural Products. Fuel. Metal Chelates. Industrial Waste Water. Explosives. Miscellaneous Applications. **8. Recent Advances and Prospect for Growth.** Recent Reviews in CE. Advances in Injection Techniques. Novel Detection Techniques. Advances in Column Technology. Progress on Electrolyte Systems. New Systems and Methods. Additional Applications Based on CE. Future Trends. **References. Index.**

1992 xxvi + 586 pages
Price: US\$ 247.00 / Dfl. 395.00
ISBN 0-444-89433-0

"Everything seems to be there, any detection system you have ever dreamed of, any capillary coating, enough electrolyte systems to saturate your wits, and more..."

"...by far the most thorough and comprehensive book in the field yet to appear."

P.G. Righetti, Milan

ORDER INFORMATION

For USA and Canada
ELSEVIER SCIENCE PUBLISHERS
Judy Weislogel
P.O. Box 945
Madison Square Station, ...
New York, NY 10160-0757
Tel: (212) 989 5800
Fax: (212) 633 3880

In all other countries

ELSEVIER SCIENCE PUBLISHERS
P.O. Box 211
1000 AE Amsterdam
The Netherlands
Tel: (+31-20) 5803 753
Fax: (+31-20) 5803 705

US\$ prices are valid only for the USA & Canada and are subject to exchange rate fluctuations; in all other countries the Dutch guilder



ELSEVIER
SCIENCE PUBLISHERS