

JOURNAL OF

# CHROMATOGRAPHY A

INCLUDING ELECTROPHORESIS AND OTHER SEPARATION METHODS

## EDITORS

U.A.Th. Brinkman (Amsterdam)  
R.W. Giese (Boston, MA)  
J.K. Haken (Kensington, N.S.W.)  
L.R. Snyder (Orinda, CA)

EDITORS, SYMPOSIUM VOLUMES,  
E. Heftmann (Orinda, CA), Z. Deyl (Prague)

## EDITORIAL BOARD

D.W. Armstrong (Rolla, MO)  
W.A. Aue (Halifax)  
P. Boček (Brno)  
A.A. Boulton (Saskatoon)  
P.W. Carr (Minneapolis, MN)  
N.H.C. Cooke (San Ramon, CA)  
V.A. Davankov (Moscow)  
G.J. de Jong (Weesp)  
Z. Deyl (Prague)  
S. Dilli (Kensington, N.S.W.)  
Z. El Rassi (Stillwater, OK)  
H. Engelhardt (Saarbrücken)  
F. Erni (Basle)  
M.B. Evans (Hatfield)  
J.L. Glajch (N. Billerica, MA)  
G.A. Guiochon (Knoxville, TN)  
P.R. Haddad (Hobart, Tasmania)  
I.M. Hais (Hradec Králové)  
W.S. Hancock (Palo Alto, CA)  
S. Hjertén (Uppsala)  
S. Honda (Higashi-Osaka)  
Cs. Horváth (New Haven, CT)  
J.F.K. Huber (Vienna)  
K.-P. Hupe (Waldbronn)  
J. Janák (Brno)  
P. Jandera (Pardubice)  
B.L. Karger (Boston, MA)  
J.J. Kirkland (Newport, DE)  
E. sz. Kováts (Lausanne)  
K. Macek (Prague)  
A.J.P. Martin (Cambridge)  
L.W. McLaughlin (Chestnut Hill, MA)  
E.D. Morgan (Keele)  
J.D. Pearson (Kalamazoo, MI)  
H. Poppe (Amsterdam)  
F.E. Regnier (West Lafayette, IN)  
P.G. Righetti (Milan)  
P. Schoenmakers (Amsterdam)  
R. Schwarzenbach (Dübendorf)  
R.E. Shoup (West Lafayette, IN)  
R.P. Singhal (Wichita, KS)  
A.M. Siouffi (Marseille)  
D.J. Strydom (Boston, MA)  
N. Tanaka (Kyoto)  
S. Terabe (Hyogo)  
K.K. Unger (Mainz)  
R. Verpoorte (Leiden)  
Gy. Vigh (College Station, TX)  
J.T. Watson (East Lansing, MI)  
B.D. Westerlund (Uppsala)

## EDITORS, BIBLIOGRAPHY SECTION

Z. Deyl (Prague), J. Janák (Brno), V. Schwarz (Prague)

ELSEVIER

# JOURNAL OF CHROMATOGRAPHY A

INCLUDING ELECTROPHORESIS AND OTHER SEPARATION METHODS

**Scope.** The *Journal of Chromatography A* publishes papers on all aspects of **chromatography, electrophoresis** and related methods. Contributions consist mainly of research papers dealing with chromatographic theory, instrumental developments and their applications. In the *Symposium volumes*, which are under separate editorship, proceedings of symposia on chromatography, electrophoresis and related methods are published. *Journal of Chromatography B: Biomedical Applications*—This journal, which is under separate editorship, deals with the following aspects: developments in and applications of chromatographic and electrophoretic techniques related to clinical diagnosis or alterations during medical treatment; screening and profiling of body fluids or tissues related to the analysis of active substances and to metabolic disorders; drug level monitoring and pharmacokinetic studies; clinical toxicology; forensic medicine; veterinary medicine; occupational medicine; results from basic medical research with direct consequences in clinical practice.

**Submission of Papers.** The preferred medium of submission is on disk with accompanying manuscript (see *Electronic manuscripts* in the Instructions to Authors, which can be obtained from the publisher, Elsevier Science B.V., P.O. Box 330, 1000 AH Amsterdam, Netherlands). Manuscripts (in English; *four* copies are required) should be submitted to: Editorial Office of *Journal of Chromatography A*, P.O. Box 681, 1000 AR Amsterdam, Netherlands, Telefax (+31-20) 5862 304, or to: The Editor of *Journal of Chromatography B: Biomedical Applications*, P.O. Box 681, 1000 AR Amsterdam, Netherlands. Review articles are invited or proposed in writing to the Editors who welcome suggestions for subjects. An outline of the proposed review should first be forwarded to the Editors for preliminary discussion prior to preparation. Submission of an article is understood to imply that the article is original and unpublished and is not being considered for publication elsewhere. For copyright regulations, see below.

**Publication information.** *Journal of Chromatography A* (ISSN 0021-9673): for 1994 Vols. 652–682 are scheduled for publication. *Journal of Chromatography B: Biomedical Applications* (ISSN 0378-4347): for 1994 Vols. 652–662 are scheduled for publication. Subscription prices for *Journal of Chromatography A*, *Journal of Chromatography B: Biomedical Applications* or a combined subscription are available upon request from the publisher. Subscriptions are accepted on a prepaid basis only and are entered on a calendar year basis. Issues are sent by surface mail except to the following countries where air delivery via SAL is ensured: Argentina, Australia, Brazil, Canada, China, Hong Kong, India, Israel, Japan, Malaysia, Mexico, New Zealand, Pakistan, Singapore, South Africa, South Korea, Taiwan, Thailand, USA. For all other countries airmail rates are available upon request. Claims for missing issues must be made within six months of our publication (mailing) date. Please address all your requests regarding orders and subscription queries to: Elsevier Science B.V., Journal Department, P.O. Box 211, 1000 AE Amsterdam, Netherlands. Tel.: (+31-20) 5803 642; Fax: (+31-20) 5803 598. Customers in the USA and Canada wishing information on this and other Elsevier journals, please contact Journal Information Center, Elsevier Science Inc., 655 Avenue of the Americas, New York, NY 10010, USA, Tel. (+1-212) 633 3750, Telefax (+1-212) 633 3764.

**Abstracts/Contents Lists** published in Analytical Abstracts, Biochemical Abstracts, Biological Abstracts, Chemical Abstracts, Chemical Titles, Chromatography Abstracts, Current Awareness in Biological Sciences (CABS), Current Contents/Life Sciences, Current Contents/Physical, Chemical & Earth Sciences, Deep-Sea Research/Part B: Oceanographic Literature Review, Excerpta Medica, Index Medicus, Mass Spectrometry Bulletin, PASCAL-CNRS, Referativnyi Zhurnal, Research Alert and Science Citation Index.

**US Mailing Notice.** *Journal of Chromatography A* (ISSN 0021-9673) is published weekly (total 52 issues) by Elsevier Science B.V., (Sara Burgerhartstraat 25, P.O. Box 211, 1000 AE Amsterdam, Netherlands). Annual subscription price in the USA US\$ 4994.00 (US\$ price valid in North, Central and South America only) including air speed delivery. Second class postage paid at Jamaica, NY 11431. **USA POSTMASTERS:** Send address changes to *Journal of Chromatography A*, Publications Expediting, Inc., 200 Meacham Avenue, Elmont, NY 11003. Airfreight and mailing in the USA by Publications Expediting.

**See inside back cover** for Publication Schedule. Information for Authors and information on Advertisements.

© 1994 ELSEVIER SCIENCE B.V. All rights reserved.

0021-9673 94 \$07.00

No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, without the prior written permission of the publisher. Elsevier Science B.V. Copyright and Permissions Department, P.O. Box 521, 1000 AM Amsterdam, Netherlands.

Upon acceptance of an article by the journal, the author(s) will be asked to transfer copyright of the article to the publisher. The transfer will ensure the widest possible dissemination of information.

**Special regulations for readers in the USA:** This journal has been registered with the Copyright Clearance Center, Inc. Consent is given for copying of articles for personal or internal use, or for the personal use of specific clients. This consent is given on the condition that the copier pays through the Center the per-copy fee stated in the code on the first page of each article for copying beyond that permitted by Sections 107 or 108 of the US Copyright Law. The appropriate fee should be forwarded with a copy of the first page of the article to the Copyright Clearance Center, Inc., 27 Congress Street, Salem, MA 01970, USA. If no code appears in an article, the author has not given broad consent to copy and permission to copy must be obtained directly from the author. The fee indicated on the first page of an article in this issue will apply retroactively to all articles published in the journal, regardless of the year of publication. This consent does not extend to other kinds of copying, such as for general distribution, resale, advertising and promotion purposes, or for creating new collective works. Special written permission must be obtained from the publisher for such copying.

No responsibility is assumed by the Publisher for any injury and/or damage to persons or property as a matter of products liability, negligence or otherwise, or from any use or operation of any methods, products, instructions or ideas contained in the materials herein. Because of rapid advances in the medical sciences, the Publisher recommends that independent verification of diagnoses and drug dosages should be made.

Although all advertising material is expected to conform to ethical (medical) standards, inclusion in this publication does not constitute a guarantee or endorsement of the quality or value of such product or of the claims made of it by its manufacturer.

This issue is printed on acid-free paper.

Printed in the Netherlands

CONTENTS

*Bibliography Section*

Editorial . . . . .	VII
Liquid Column Chromatography . . . . .	B1
Gas Chromatography . . . . .	B58
Planar Chromatography . . . . .	B81
Electrophoresis . . . . .	B92



JOURNAL OF CHROMATOGRAPHY A  
VOL. 681 (1994)

ห้องสมุดมหาวิทยาลัยศรีปทุม

2537



# JOURNAL OF CHROMATOGRAPHY A

INCLUDING ELECTROPHORESIS AND OTHER SEPARATION METHODS

## EDITORS

U.A.Th. BRINKMAN (Amsterdam), R.W. GIESE (Boston, MA), J.K. HAKEN (Kensington, N.S.W.),  
L.R. SNYDER (Orinda, CA)

## EDITORS, SYMPOSIUM VOLUMES

E. HEFTMANN (Orinda, CA), Z. DEYL (Prague)

## EDITORIAL BOARD

D.W. Armstrong (Rolla, MO), W.A. Aue (Halifax), P. Boček (Brno), A.A. Boulton (Saskatoon), P.W. Carr (Minneapolis, MN), N.H.C. Cooke (San Ramon, CA), V.A. Davankov (Moscow), G.J. de Jong (Weesp), Z. Deyl (Prague), S. Dilli (Kensington, N.S.W.), Z. El Rassi (Stillwater, OK), H. Engelhardt (Saarbrücken), F. Erni (Basle), M.B. Evans (Hatfield), J.L. Glajch (N. Billerica, MA), G.A. Guiochon (Knoxville, TN), P.R. Haddad (Hobart, Tasmania), I.M. Hais (Hradec Králové), W.S. Hancock (Palo Alto, CA), S. Hjertén (Uppsala), S. Honda (Higashi-Osaka), Cs. Horváth (New Haven, CT), J.F.K. Huber (Vienna), K.-P. Hupe (Waldbronn), J. Janák (Brno), P. Jandera (Pardubice), B.L. Karger (Boston, MA), J.J. Kirkland (Newport, DE), E. sz. Kováts (Lausanne), K. Macek (Prague), A.J.P. Martin (Cambridge), L.W. McLaughlin (Chestnut Hill, MA), E.D. Morgan (Keele), J.D. Pearson (Kalamazoo, MI), H. Poppe (Amsterdam), F.E. Regnier (West Lafayette, IN), P.G. Righetti (Milan), P. Schoenmakers (Amsterdam), R. Schwarzenbach (Dübendorf), R.E. Shoup (West Lafayette, IN), R.P. Singhal (Wichita, KS), A.M. Siouffi (Marseille), D.J. Strydom (Boston, MA), N. Tanaka (Kyoto), S. Terabe (Hyogo), K.K. Unger (Mainz), R. Verpoorte (Leiden), Gy. Vigh (College Station, TX), J.T. Watson (East Lansing, MI), B.D. Westerlund (Uppsala)

## EDITORS, BIBLIOGRAPHY SECTION

Z. Deyl (Prague), J. Janák (Brno), V. Schwarz (Prague)



ELSEVIER  
AMSTERDAM — LONDON — NEW YORK — TOKYO

---

*J. Chromatogr. A*, Vol. 681 (1994)

The *Bibliography Section 1994* will be published in two volumes (681 and 682) of two issues each. Combined indexes to both volumes will appear in the last issue of the year, Vol. 682, No. 2.

© 1994 ELSEVIER SCIENCE B.V. All rights reserved.

0021-9673/94/\$07.00

No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, without the prior written permission of the publisher, Elsevier Science B.V., Copyright and Permissions Department, P.O. Box 521, 1000 AM Amsterdam, Netherlands.

Upon acceptance of an article by the journal, the author(s) will be asked to transfer copyright of the article to the publisher. The transfer will ensure the widest possible dissemination of information.

*Special regulations for readers in the USA* – This journal has been registered with the Copyright Clearance Center, Inc. Consent is given for copying of articles for personal or internal use, or for the personal use of specific clients. This consent is given on the condition that the copier pays through the Center the per-copy fee stated in the code on the first page of each article for copying beyond that permitted by Sections 107 or 108 of the US Copyright Law. The appropriate fee should be forwarded with a copy of the first page of the article to the Copyright Clearance Center, Inc., 27 Congress Street, Salem, MA 01970, USA. If no code appears in an article, the author has not given broad consent to copy and permission to copy must be obtained directly from the author. The fee indicated on the first page of an article in this issue will apply retroactively to all articles published in the journal, regardless of the year of publication. This consent does not extend to other kinds of copying, such as for general distribution, resale, advertising and promotion purposes, or for creating new collective works. Special written permission must be obtained from the publisher for such copying.

No responsibility is assumed by the Publisher for any injury and/or damage to persons or property as a matter of products liability, negligence or otherwise, or from any use or operation of any methods, products, instructions or ideas contained in the materials herein. Because of rapid advances in the medical sciences, the Publisher recommends that independent verification of diagnoses and drug dosages should be made.

Although all advertising material is expected to conform to ethical (medical) standards, inclusion in this publication does not constitute a guarantee or endorsement of the quality or value of such product or of the claims made of it by its manufacturer.

This issue is printed on acid-free paper.

Printed in the Netherlands



# BIBLIOGRAPHY SECTION

SUPPLEMENT TO THE  
JOURNAL OF CHROMATOGRAPHY A

1994

EDITORS:

Z. DEYL (Prague)

J. JANÁK (Brno)

V. SCHWARZ (Prague)



## Editorial

This year the Bibliography Section will be published in two volumes (N°681 and N°682). The rules of classification of individual entries will not differ from those used in the past year. There is a single change regarding fullerenes which do not fit anywhere into the classification scheme and we have decided to join them with section 5B. The Indexes for this year will be attached to the second volume of this year (N°682).

*Prague (Czech Republic)*  
*Brno (Czech Republic)*

Z. Deyl and V. Schwarz  
J. Janák





















## Bibliography Section

---

### Liquid Column Chromatography

1. REVIEWS AND BOOKS
- 1 Barker, P.E. and Ganetsos, G.: Continuous moving-column chromatographic systems. *Chromatogr. Sci.*, 61 (1993) 173-186; C.A., 119 (1993) 119875e - a review with 27 refs.
- 2 Betina, V.: Sampling sample preparation, extraction and clean-up. *J. Chromatogr. Libr.*, 54(Chromatography of Mycotoxins) (1993) 3-11; C.A., 119 (1993) 176868d.
- 3 Brazhnikov, V.V.: (Detectors for chromatography. Review). *Zavod. Lab.*, 59 (1993) 16-22; C.A., 119 (1993) 194730g - a review with 17 refs.
- 4 Bridges, S. and Barker, P.E.: Continuous cross-current chromatographic refiners. *Chromatogr. Sci.*, 61 (1993) 113-126; C.A., 119 (1993) 119874d - a review with 25 refs.
- 5 Candlish, A.A.G. and Stimson, W.H.: Emerging techniques: immunoaffinity chromatography. *J. Chromatogr. Libr.*, 54(Chromatography of Mycotoxins) (1993) 99-123; C.A., 119 (1993) 176872a.
- 6 Carr, P.W.: Solvatochromism, linear solvation energy relationships, and chromatography. *Microchem. J.*, 48 (1993) 4-28; C.A., 119 (1993) 189467x - a review with 54 refs.
- 7 Cho, J.J.: Principles and practice of high performance liquid chromatography. Applications in textiles. *Han'guk Somyu Konghakhoechi*, 29 (1992) 377-389; C.A., 119 (1993) 162232e - a review with 16 refs.
- 8 Conder, J.R.: Physicochemical measurement by chromatography: overview and solution thermodynamics. *NATO ASI Ser., Ser. C*, 383 (1992) 315-337; C.A., 119 (1993) 189639e - a review with many refs.
- 9 Gal, J.: Indirect methods for the chromatographic resolution of drug enantiomers. Synthesis and separation of diastereomeric derivatives. *Clin. Pharmacol.*, 18 (1993) 65-106; C.A., 119 (1993) 188683c - a review with 231 refs.
- 10 Ganetsos, G. and Barker, P.E. (Editors): *Preparative and Production Scale Chromatography. Chromatogr. Sci.*, Vol. 61, Dekker, New York, 1993, 786 p.; C.A., 119 (1993) 142081b.
- 11 Jinno, K.: Optimization of separation in high-performance liquid chromatography. *Bunseki*, (1993) 521-525; C.A., 119 (1993) 189038h - a review with 57 refs.
- 12 Kuronen, P.: Techniques of liquid column chromatography. *J. Chromatogr. Libr.*, 54(Chromatography of Mycotoxins) (1993) 36-77; C.A., 119 (1993) 176870y.
- 13 Miyabe, K. and Suzuki, M.: (Current status and related issues of studies on optimization strategies of operating conditions for reversed-phase liquid chromatography). *Seisan Kenkyu*, 45 (1993) 275-282; C.A., 119 (1993) 130433e - a review with 56 refs.
- 14 Nimura, N.: (HPLC on non-porous packing material). *Kuromatogurafi*, 13 (1992) 195-199; C.A., 119 (1993) 90327q - a review with 5 refs.
- 15 Nissen, H.P. and Kreysel, H.W.: Separation systems and instrumentation. In: Matissek, R. and Wittkowski, R. (Editors), *Perform. Liq. Chromatogr. Food Control Res.*, Technomic, Lancaster, 1993, pp. 63-83; C.A., 119 (1993) 155019n - a review with 20 refs.
- 16 Saito, H.: (Recent state of separation technology using liquid chromatography apparatus for production). *Kagaku Sochi*, 35 (1993) 63-68; C.A., 119 (1993) 120157d - a review with 10 refs.
- 17 Shaevich, A.B. and Muchnik, V.L.: (Computerized chromatographic identification of substances: method characteristics and data bases). *Zh. Fiz. Khim.*, 67 (1993) 827-832; C.A., 119 (1993) 194728n - a review with 27 refs.
- 18 Smith, W.T., Jr. and Patterson, J.M.: Analytical methods for alcohols, phenols, ethens and peroxides. In: Patai, S. (Editor), *Chem. Hydroxyl, Ether Peroxide Groups*, Wiley, Chichester, 1993, pp. 201-239; C.A., 119 (1993) 194721e - a review with 258 refs.
- 19 Stevenson, R.: Trickle down from the 150,000-L column. *Am. Lab. (Shelton)*, 25 (1993) 40ZZ-AAA; C.A., 119 (1993) 112449k - a review with no refs.
- 20 Timpa, J.D.: Size exclusion chromatography for characterization of cotton fiber. *Trends Polym. Sci.*, 1 (1993) 105-110; C.A., 119 (1993) 141043d - a review with 40 refs.
- 21 Vigdergauz, M.S. and Petrova, Ye.I.: (The alternation effect in regularities of chromatographic retention). *Usp. Khim.*, 61 (1992) 2172-2186; C.A., 119 (1993) 147096v - a review with 80 refs.
- 22 Yang, F.: (Basic relations of countercurrent chromatography). *Septu*, 11 (1993) 136-139; C.A., 119 (1993) 125716v - a review with 32 refs.
- 23 Yashin, Ya.I.: (Scientometric study of the state of the art and trends in chromatographic method and instrumentation). *Zh. Anal. Khim.*, 48 (1993) 415-426; C.A., 119 (1993) 151191q - a review with 5 refs.

See also 76, 133, 142, 156, 174, 196, 198, 226, 234, 236, 251, 256, 273, 276, 279, 297, 363, 364, 377, 422, 491, 510, 573, 578, 589, 593, 602, 606, 607, 876, 894, 904, 907, 910, 911, 955, 1003, 1081, 1082, 1087, 1105, 1111, 1151, 1152, 1154, 1159, 1160, 1163, 1165, 1293, 1359, 1360, 1361, 1362, 1363, 1364, 1365, 1368, 1369, 1373, 1374, 1393, 1418, 1424.

## 2. FUNDAMENTALS, THEORY AND GENERAL

### 2a. General

- 24 Baiocchi, C., Marengo, E., Saini, G., Roggero, M.A. and Giacomosa, D.: Reversed-phase high-performance liquid chromatography and chemometrics, a combined investigation tool for complex phytochemical problems. *J. Chromatogr.*, 644 (1993) 259-267.
- 25 Belfer, A.J. and Locke, D.C.: Calibration of instruments and chromatographs. *Eur. Pat. Appl. EP 554,604* (Cl. G01N30/62), 11 Aug. 1993, US Appl. 830,646, 04 Feb. 1992; 6 p.; C.A., 119 (1993) 130700q.
- 26 Chen, N., Liang, X., Zhang, Y. and Lu, P.: Chromatogram base in expert system for liquid chromatography. *Chin. Sci. Bull.*, 37 (1992) 1809-1812; C.A., 119 (1993) 90408s.
- 27 Dolan, J.W.: Avoiding the pitfalls of published methods. *LC-GC*, 11 (1993) 412-415; C.A., 119 (1993) 173257t.
- 28 Fornstedt, T. and Westerlund, D.: Effects on analyte peak performance by separated system peaks in ion-pair adsorption chromatography. *J. Chromatogr.*, 648 (1993) 315-324.
- 29 Herold, M.: SEC: Influence of salt concentration in the mobile phase. *Am. Lab. (Shelton)*, 25 (1993) 35-38; C.A., 119 (1993) 176866b.
- 30 Hu, W., Takeuchi, T. and Haraguchi, H.: Electrostatic ion chromatography. *Anal. Chem.*, 65 (1993) 2204-2208.
- 31 Kaufmann, M., Schwarz, T. and Bartholmes, P.: Continuous buffer exchange of column chromatographic eluates using a hollow-fibre membrane module. *J. Chromatogr.*, 639 (1993) 33-39.
- 32 Matsuda, R., Hayashi, Y., Suzuki, T. and Saito, Y.: Information theory of miniaturization advantages of column liquid chromatography for quantitative analysis. *Fresenius J. Anal. Chem.*, 347 (1993) 225-229.
- 33 Mulholland, M.: Expert systems and their implementation in chromatography. In: Bryce, C.F.A. (Editor), *Microcomput. Biochem.*, IRL, Oxford, 1992, pp. 243-266; C.A., 119 (1993) 155324h.
- 34 Potschka, M.: Mechanism of size-exclusion chromatography. I. Role of convection and obstructed diffusion in size-exclusion chromatography. *J. Chromatogr.*, 648 (1993) 41-69.
- 35 Schisla, D.K., Dink, H., Carr, P.W. and Cussler, E.L.: Polydisperse tube diameters compromise multiple open tubular chromatography. *AIChE J.*, 39 (1993) 946-953; C.A., 119 (1993) 108171w.
- 36 Schlüter, H. and Zidek, W.: Application of non-size-related separation effects to the purification of biologically active substances with a size-exclusion gel. *J. Chromatogr.*, 639 (1993) 17-22.
- 37 Stevenson, R.: Chromatography currents. Separation science in Japan: from the top down and the bottom up. *Int. Lab.*, 23, No. 9 (1993) P10-P15.
- 38 Touchstone, J.C.: History of chromatography. *J. Liq. Chromatogr.*, 16 (1993) 1647-1665.

For additional information see C.A.:

119 (1993) 173109w, 183509r, 194770v.

See also 6, 11, 13, 23, 44, 55, 68, 114, 132.

### 2b. Thermodynamics and theoretical relationships

- 39 Andreev, V.P. and Stefanovich, L.A.: Theory of field flow fractionation with the reversible adsorption on channel walls. *Chromatographia*, 37 (1993) 325-328.
- 40 Hatanaka, T. and Ishida, M.: The effect of adsorption isotherms and flow rates on separation performance in multicomponent continuous separation. *J. Chem. Eng. Jpn.*, 23 (1993) 242-246; C.A., 119 (1993) 98436v.
- 41 Janicka, M. and Rózylo, J.K.: The use of TLC in sandwich chambers for theoretical prediction of mixture separation in HPLC. *J. Planar Chromatogr.*, 6 (1993) 362-367.
- 42 Kazakevich, Y.V. and McNair, H.M.: Thermodynamic definition of HPLC dead volume. *J. Chromatogr. Sci.*, 31 (1993) 317-322.
- 43 King, J.W., Hopper, M.L., Luchtefeld, R.G., Taylor, S.L. and Orton, W.L.: Optimization of experimental conditions for the supercritical carbon dioxide extraction of pesticide residues from grain. *J. Assoc. Off. Anal. Chem.*, 76 (1993) 857-864.
- 44 Koo, Y.M. and Hong, J.: Scaling rules of nonlinear, coupled elution chromatography. In: Furusaki, S., Endo, I. and Matsuno, R. (Editors), *Biochem. Eng. 2001, Proc. Asia-Pac. Biochem. Eng. Conf.*, Springer, Tokyo, 1992, pp. 497-500; C.A., 119 (1993) 134674g.
- 45 Lebedev, Yu.Ya.: (Analysis of the deviation from equilibrium of interphase mass distributions for chromatographic bands). *Zh. Fiz. Chem.*, 67 (1993) 765-768; C.A., 119 (1993) 168322f.
- 46 Möckel, H.J. and Dreyer, U.: Second order retention effects in reversed-phase liquid chromatography. 1. Influence of solute size. *Chromatographia*, 37 (1993) 179-184.
- 47 Seidel-Morgenstern, A. and Guiochon, G.: Theoretical study of recycling in preparative chromatography. *AIChE J.*, 39 (1993) 809-819; C.A., 119 (1993) 104062p.
- 48 Shortt, D.W.: Differential molecular weight distributions in high performance size exclusion chromatography. *J. Liq. Chromatogr.*, 16 (1993) 3371-3391.
- 49 Tauler, R. and Barceló, D.: Multivariate curve resolution applied to liquid chromatography-diode array detection. *TrAC*, 12 (1993) 319-327.
- 50 Toft, J. and Kvalheim, O.M.: Resolution parameter for multidetection chromatography. *Anal. Chem.*, 65 (1993) 2270-2275.
- 51 Vigdergauz, M.S., Platonov, I.A., Lobachev, A.L. and Sushko, O.V.: (Retention times of nonsorbing materials in chromatography). *Zh. Fiz. Khim.*, 67 (1993) 857-860; C.A., 119 (1993) 189265e.
- 52 Yamamoto, A., Matsunaga, A., Mizukami, E., Hayakawa, K. and Miyazaki, M.: Adsorption isotherm of undissociated eluent acid and its relation to the retention of system peaks in non-suppressed ion chromatography. *J. Chromatogr.*, 644 (1993) 183-187.
- 53 Yashin, Ya. I.: (Retention rules for ion chromatography). *Zh. Fiz. Khim.*, 67 (1993) 769-772; C.A., 119 (1993) 168323g.

See also 34, 60, 113, 267, 497, 579, 1127.

## 2c. Relationship between structure and chromatographic behaviour

- 54 Cupid, B.C., Nicholson, J.K., Davis, P., Ruane, R.J., Wilson, I.D., Glen, R.C., Rose, V.S., Beddel, C.R. and Lindon, J.C.: Quantitative structure chromatography relationships in reversed-phase high performance liquid chromatography: prediction of retention behaviour using theoretically derived molecular properties. *Chromatographia*, 37 (1993) 241-249.
- 55 Hamoir, T., Massart, D.L., King, W., Kokot, S. and Douglas, K.: Prediction of initial chromatographic conditions in reversed-phase high-performance liquid chromatography. *J. Chromatogr. Sci.*, 31 (1993) 393-400.
- 56 Hanai, T., Hatano, H., Nimura, N. and Kinoshita, T.: QSRR in liquid chromatography aided by computational chemistry. *J. Liq. Chromatogr.*, 16 (1993) 1453-1462.
- 57 Law, B. and Stafford, L.E.: The use of ultraviolet spectra and chromatographic retention data as an aid to metabolite identification. *J. Pharm. Biomed. Anal.*, 11 (1993) 729-736.
- 58 Wyatt, P.J.: Mean square radius of molecules and secondary instrumental broadening. *J. Chromatogr.*, 648 (1993) 27-32.
- 59 Zou, H., Zhang, Y., Hong, M. and Lu, P.: Effect of Hammett's constant on the quantitative correlation of benzoic acid retention in RP-IPC with *n*-octanol/water partition coefficients. *J. Liq. Chromatogr.*, 16 (1993) 3433-3443.

See also 126, 454.

## 2d. Measurement of physico-chemical and related values

- 60 Nagy, D.J.: A Mark-Houwink equation for poly(vinyl alcohol) from SEC-viscometry. *J. Liq. Chromatogr.*, 16 (1993) 3041-3058.
- 61 Nam Tram Ho, Vo Van Huan, Buchmann, M., Sabra, F., Ruelle, P. and Kesselring, U.W.: Adsorption isotherms of water on a set of Fractosils and on Avicel PH 101 by frontal analysis chromatography. *Drug Dev. Ind. Pharm.*, 19 (1993) 1413-1445; *C.A.*, 119 (1993) 103190s.
- 62 Sanayei, R.A., Pang, S. and Rudin, A.: A new approach to establishing universal calibration curves for size-exclusion chromatography. *Polymer*, 34 (1993) 2320-2323; *C.A.*, 119 (1993) 96699j.
- 63 Vigdergauz, M.S. and Lange, P.K.: (Chromatography as a spectral method for investigations). *Zh. Fiz. Khim.*, 67 (1993) 815-816; *C.A.*, 119 (1993) 168276u.

See also 8, 48, 73, 167, 200, 221, 259, 287, 290, 294, 451, 1121, 1124, 1125, 1130, 1131, 1133, 1134, 1135, 1136, 1137, 1139, 1142, 1143, 1287, 1396.

## 3. GENERAL TECHNIQUES

## 3a. Apparatus and accessories

- 64 Andelman, M.D.: Flow-through capacitor for liquid chromatograph. *U.S. US 5,192,432* (Cl. 210-198.2; B01D15/08), 09 Mar. 1993, *US Appl.* 512,970, 23 Apr. 1990; 7 p.; *C.A.*, 119 (1993) 183502h.

- 65 Conway, W.D.: Multicolumn planet-centrifuge chromatograph. *U.S. US 5,217,608* (Cl. 210-198.2; B01D15/08), 08 Jun. 1993, *US Appl.* 223,024, 21 Jul. 1988; 17 p.; *C.A.*, 119 (1993) 108132j.
- 66 Maystre, F. and Bruno, A.: Liquid-chromatograph tube-system separation arrangement and optical method for analysis of samples. *Eur. Pat. Appl.* EP 547,282 (Cl. G01N30/74), 23 Jun. 1993, *Appl.* 91/810,991, 19 Dec. 1991; 11 p.; *C.A.*, 119 (1993) 130695s.
- 67 Sarmidi, M.R. and Barker, P.E.: Simultaneous biochemical reaction and separation in a rotating annular chromatograph. *Chem. Eng. Sci.*, 48 (1993) 2615-2623; *C.A.*, 119 (1993) 112605h.

For additional information see *C.A.*:

119 (1993) 173229k, 194662m, 194820m.

See also 1, 4, 25, 105, 169, 170, 194, 195, 197, 1042.

## 3b. Detectors and detection reagents

- 68 Boschetti, E., Duteil, X., Pouradier, N.C. and Moroux, Y.: Concerns and solutions for a proper decontamination of chromatographic packings. *Chim. Oggi*, 11 (1993) 29-35; *C.A.*, 119 (1993) 155149e.
- 69 Cameron, G., Jackson, P.E. and Gorenstein, M.V.: A new approach to peak purity assessment using photodiode array detection. *Chem. Aust.*, 60 (1993) 288-289; *C.A.*, 119 (1993) 108016z.
- 70 Fung, Y.M. and Saunders, P.: Liquid chromatography with electrochemical detection: the advantage of microcomposite electrodes. *Chem. Aust.*, 60 (1993) 278-280; *C.A.*, 119 (1993) 108174z.
- 71 Han, S.H., Lee, K.S., Cha, G.S., Liu, D. and Trojanowicz, M.: Potentiometric detection in ion chromatography using multi-ionophore membrane electrodes. *J. Chromatogr.*, 648 (1993) 283-288.
- 72 Idowu, O.R. and Adewuyi, G.O.: Sodium benzoxazole-2-sulfonate: a derivatisation reagent for the analysis of amines and amino acids by HPLC with fluorescence or UV detection. *J. Liq. Chromatogr.*, 16 (1993) 2501-2518.
- 73 Jeng, L. and Balke, S.T.: Evaluation of light-scattering detectors for size exclusion chromatography. II. Light-scattering equation selection. *J. Appl. Polym. Sci.*, 49 (1993) 1375-1385; *C.A.*, 119 (1993) 181837x.
- 74 Jeng, L., Balke, S.T., Mourey, T.H., Wheeler, L. and Romeo, P.: Evaluation of light-scattering detectors for size exclusion chromatography. *J. Appl. Polym. Sci.*, 49 (1993) 1359-1374; *C.A.*, 119 (1993) 181836w.
- 75 Jones, D., Brenton, A.G., Games, D.E., Brittain, A.H., Taylor, S., Kennedy, D. and Smith, P.: Ion mobility spectrometry as a detection technique for the separation sciences. *Rapid Commun. Mass Spectrom.*, 7 (1993) 561-566; *C.A.*, 119 (1993) 130666h.
- 76 Kientz, C.E. and Brinkman, U.A.T.: The use of gas chromatographic detectors in column liquid chromatography. *TrAC*, 12 (1993) 363-373 - a review with 52 refs.
- 77 Lago, P., Rovati, L., Cantu, L. and Corti, M.: A quasielastic light scattering detector for chromatographic analysis. *Rev. Sci. Instrum.*, 64 (1993) 1797-1802; *C.A.*, 119 (1993) 176852u.

- 78 Mason, P.B., Zhang, L., Carnahan, J.W. and Winans, R.E.: Helium microwave induced plasma atomic emission detection for liquid chromatography utilizing a moving band interface. *Anal. Chem.*, 65 (1993) 2596-2600.
- 79 Oliveros, L. and Minguillón, C.: Example of pitfalls in the UV detection used in the resolution of racemic compounds by liquid chromatography. *J. Chromatogr.*, 653 (1993) 144-147.
- 80 Rapkin, E.:  $\beta$ -Particle detection in HPLC by flow-through monitoring vs. liquid scintillation counting. *J. Liq. Chromatogr.*, 16 (1993) 1769-1781.
- 81 Rosenzweig, Z. and Yeung, E.S.: Laser-based double beam absorption detector for high-performance liquid chromatography. *J. Chromatogr.*, 645 (1993) 201-207.
- 82 Takeuchi, T. and Miwa, T.: Indirect conductometric detection of nonelectrolytes in microcolumn liquid chromatography. *Chromatographia*, 37 (1993) 281-283.
- 83 Toyo'oka, T., Chokshi, H.P., Givens, R.S., Carlson, R.G., Lunte, S.M. and Kuwana, T.: Fluorescence and chemiluminescence detection of oxazole-labelled amines and thiols. *Biomed. Chromatogr.*, 7 (1993) 208-216.
- 84 White, J.G., Soli, A.L. and Jorgenson, J.W.: Rapid scan microvoltammetric detection for liquid chromatography. *J. Liq. Chromatogr.*, 16 (1993) 1489-1496.
- For additional information see C.A.:  
119 (1993) 173321j, 183506n.
- See also 3, 49, 50, 139, 212, 227, 242, 244, 252, 274, 326, 342, 346, 351, 354, 367, 373, 398, 405, 426, 446, 448, 456, 464, 472, 524, 534, 741, 890, 923, 944, 961, 964, 971, 1000, 1048, 1055, 1062, 1098, 1160, 1415, 1420, 1435, 1438.
- 3c. *Sorbents and columns, packing procedures*
- 85 Alebic-Kolbah, T. and Wainer, I.W.: Application of an enzyme-based stationary phase to the determination of enzyme kinetic constants and types of inhibition. New high-performance liquid chromatographic approach utilizing an immobilized artificial membrane chromatographic support. *J. Chromatogr.*, 653 (1993) 122-129.
- 86 Alebic-Kolbah, T. and Wainer, I.W.: Enzyme-based high-performance liquid chromatography stationary phases as metabolic reactors. Immobilization of non-solubilized rat liver microsomes on an immobilized artificial membrane high-performance liquid chromatography support. *J. Chromatogr.*, 646 (1993) 289-295.
- 87 Bergmann-Leyder, N., Tambuté, A., Bégos, A. and Caude, M.: Synthesis and mechanistic study of novel  $\pi$ -basic chiral stationary phase derived from tyrosine. *Chromatographia*, 37 (1993) 433-443.
- 88 Buszewski, B., Kawka, S. and Wolski, T.: Polyamide PA-6 and silica gel as a mixed packing material for isolating dab syl amino acids using solid-phase extraction. *LC-GC*, 11 (1993) 366-369; C.A., 119 (1993) 134695q.
- 89 Chriswanto, H., Ge, H. and Wallace, G.G.: Polypyrrole-coated silica as a new stationary phase for liquid chromatography. *Chromatographia*, 37 (1993) 423-428.
- 90 Ding, X., He, X., Tu, G., Wang, P., Chu, A., Zhao, R., Liu, G. and Huang, X.: (Development of stationary phases based on controlled pore glass used for high-performance hydrophobic interaction chromatography). *Fenxi Huaxue*, 21 (1993) 410-415; C.A., 119 (1993) 134675h.
- 91 Farkas, G., Irgens, L.H., Quintero, G., Beeson, M.D., Al-Saeed, A. and Vigh, G.: Displacement chromatography on cyclodextrin silicas. IV. Separation of the enantiomers of ibuprofen. *J. Chromatogr.*, 645 (1993) 67-74.
- 92 Göhlin, K. and Larsson, M.: Study of polyorganosiloxanes (native and solvent swollen) for the preparation of narrow (5-15  $\mu$ m I.D.) and long (1-6 m) open tubular columns in reversed-phase liquid chromatography. *J. Chromatogr.*, 645 (1993) 41-56.
- 93 Grosse-Bley, M., Boemer, B., Grosser, R., Arlt, D. and Lange, W.: Preparation of optically active sulphur-containing amino acid derivatives and their polymers. *Ger. Offen.* DE 4,120,695 (Cl. C07C323/59), 24 Dec. 1992, Appl. 22 Jun. 1991; 14 p.; C.A., 119 (1993) 139964m.
- 94 Hagel, L.: Size-exclusion chromatography in an analytical perspective. *J. Chromatogr.*, 648 (1993) 19-25.
- 95 Hanson, M., Unger, K.K., Schmid, J., Albert, K. and Bayer, E.: Effect of the chain mobility of polymeric reversed-phase stationary phases on polypeptide retention. *Anal. Chem.*, 65 (1993) 2249-2253.
- 96 Hjertén, S., Mohammad, J. and Nakazato, K.: Improvement in flow properties and pH stability of compressed, continuous polymer beds for high-performance liquid chromatography. *J. Chromatogr.*, 646 (1993) 121-128.
- 97 Hosoya, K. and Frechet, J.M.J.: Influence of the seed polymer on the chromatographic properties of size monodisperse polymeric separation media prepared by a multi-step swelling and polymerization method. *J. Polym. Sci., Part A: Polym. Chem.*, 31 (1993) 2129-2141; C.A., 119 (1993) 96789p.
- 98 Hosoya, K., Kimata, K., Tanaka, N., Araki, T., Terashima, M. and Fréchet, J.M.J.: Influence of pore size and pore size distribution of polymer-based packing materials on chromatographic separation of carbon clusters. *J. Liq. Chromatogr.*, 16 (1993) 3059-3071.
- 99 Inoue, S. and Ohtaki, N.: Pyrophosphates as biocompatible packing materials for high-performance liquid chromatography. *J. Chromatogr.*, 645 (1993) 57-65.
- 100 Jinno, K., Ohta, H., Saito, Y., Uemura, T., Nagashima, H., Itoh, K., Chen, Y.-L., Luehr, G., Archer, J., Fetzer, J.C. and Biggs, W.R.: Dimethoxyphenylpropyl bonded silica phase for higher fullerenes separation by high-performance liquid chromatography. *J. Chromatogr.*, 648 (1993) 71-77.
- 101 Kaneko, S., Mitsuzawa, T., Ohmori, S., Nakamura, M., Nobuhara, K. and Masatani, M.: Silica-containing mixed-oxide gels prepared by a coprecipitation method as novel packing materials for liquid chromatography. *Chem. Lett.*, (1993) 1275-1278; C.A., 119 (1993) 173287c.
- 102 Kaster, J.A., de Oliveira, W., Glasser, W.G. and Velander, W.H.: Optimization of pressure-flow limits, strength, intraparticle transport and dynamic capacity by hydrogel solids content and bead size in cellulose immunosorbents. *J. Chromatogr.*, 648 (1993) 79-90.
- 103 Kibbey, C.E. and Meyerhoff, M.E.: Preparation and characterization of covalently bound tetraphenylporphyrin-silica gel stationary phases for reversed-phase and anion-exchange chromatography. *Anal. Chem.*, 65 (1993) 2189-2196.



- 104 Lee, H.G. and Jarrett, H.W.: Chromatographic stability of glucose-silica. *J. Chromatogr.*, 653 (1993) 130-134.
- 105 Majors, R.E.: New chromatography columns and accessories at the 1993 Pittsburgh Conference. Part II. *LC-GC*, 11 (1993) 258-269; *C.A.*, 119 (1993) 194738r.
- 106 Martin, P., Taberner, J., Fairbrother, A. and Wilson, I.D.: An investigation of the effects of carbon loading and endcapping on the solid-phase extraction of  $\beta$ -blockers onto  $C_{18}$  bonded silica gel. *J. Pharm. Biomed. Anal.*, 11 (1993) 671-677.
- 107 Matsui, J., Kato, T., Takeuchi, T., Suzuki, M., Yokoyama, K., Tamiya, E. and Karube, I.: Molecular recognition in continuous polymer rod prepared by a molecular imprinting technique. *Anal. Chem.*, 65 (1993) 2223-2224.
- 108 Mingalyov, P.G., Fadeev, A.Y., Staroverov, S.M., Lisichkin, G.V. and Lunina, E.V.: Nitroxide radicals in studies of the fine bonded layer structure of modified silicas. *J. Chromatogr.*, 646 (1993) 267-277.
- 109 Mueller-Schulte, D.: Synergistic-radiation grafting: a novel modification technique for the preparation of biomaterials. *Radiat. Phys. Chem.*, 42 (1993) 891-896; *C.A.*, 119 (1993) 188503u.
- 110 Nagaoka, S., Ihara, H. and Hirayama, C.: Preparation and characterization of spherical polymer packings from N,N-dialkylacrylamide for reversed-phase liquid chromatography. *Polym. J. (Tokyo)*, 25 (1993) 609-618; *C.A.*, 119 (1993) 173279b.
- 111 Nimura, N., Itoh, H. and Kinoshita, T.: (Binary-layered phase liquid chromatography). *Kuromatogurafi*, 13 (1992) 293-294; *C.A.*, 119 (1993) 85258p.
- 112 Porsch, B.: Epoxy- and diol-modified silica: optimization of surface bonding reaction. *J. Chromatogr.*, 653 (1993) 1-7.
- 113 Rodrigues, A.E., Lu, Z.P., Loureiro, J.M. and Carta, G.: Peak resolution in linear chromatography. Effects of intraparticle convection. *J. Chromatogr.*, 653 (1993) 189-198.
- 114 Salzman, A.T., Fulton, S., Gordon, N., Meys, M. and Carberry, R.: Development and execution of a biomolecule purification method. *Am. Biotechnol. Lab.*, 11 (1993) 40-44; *C.A.*, 119 (1993) 112583z.
- 115 Smith, R.E., Yourtee, D., Bean, T. and MacQuarrie, R.A.: Ion chromatography on a new moderate capacity anion exchange column. *J. Chromatogr. Sci.*, 31 (1993) 366-370.
- 116 Stalling, D.L., Guo, C.Y. and Saim, S.: Surface-linked  $C_{60}/70$ -polystyrene divinylbenzene beads as a new chromatographic material for enrichment of coplanar PCBs. *J. Chromatogr. Sci.*, 31 (1993) 265-278.
- 117 Su, W., Gregory, R.B. and Gilpin, R.K.: Liquid chromatographic studies of silica-immobilized HEW lysozyme. *J. Chromatogr. Sci.*, 31 (1993) 285-290.
- 118 Suhara, T., Kutsuna, H., Fukui, H. and Yamaguchi, M.: Modification of silica gel with diethylamino group by novel reaction and its characterization. *Colloid Polym. Sci.*, 271 (1993) 486-493; *C.A.*, 119 (1993) 134707v.
- 119 Takeuchi, T. and Miwa, T.: Determination of the specific surface area of silica gel by on-column titration with hydrochloric acid. *Anal. Chim. Acta*, 282 (1993) 565-570.
- 120 Vilenchik, L.Z., Asrar, J., Ayotte, R.C., Ternorutsky, L. and Hardiman, C.J.: Macromolecular porosimetry. *J. Chromatogr.*, 648 (1993) 9-17.
- 121 Wainer, I.W.: HPLC chiral stationary phases for the stereochemical resolution of enantiomeric compounds. The current state of the art. *Clin. Pharmacol.*, 18 (1993) 139-182; *C.A.*, 119 (1993) 188687g.
- 122 Wang, Q.C., Svec, F. and Fréchet, J.M.F.: Macroporous polymeric stationary-phase rod as continuous separation medium for reversed-phase chromatography. *Anal. Chem.*, 65 (1993) 2243-2248.
- 123 Wirth, H.-J. and Hearn, M.T.W.: High-performance liquid chromatography of amino acids, peptides and proteins. CXXX. Modified porous zirconia as sorbents in affinity chromatography. *J. Chromatogr.*, 646 (1993) 143-151.
- 124 Wirth, H.-J., Eriksson, K.-O., Holt, P., Aguilar, M. and Hearn, M.T.W.: High-performance liquid chromatography of amino acids, peptides and proteins. CXXIX. Ceramic-based particles as chemically stable chromatographic supports. *J. Chromatogr.*, 646 (1993) 129-141.
- 125 Yu, J. and El Rassi, Z.: Reversed-phase liquid chromatography of dansyl amino acids with microspherical octadecyl-silica and octadecyl-zirconia bonded stationary phases. *J. Liq. Chromatogr.*, 16 (1993) 2931-2959.
- 126 Zou, H., Zhang, Y., Hong, M. and Lu, P.: Effect of carbon content bonded on silica surface on retention values in reversed-phase liquid chromatography and ion-pair liquid chromatography. *J. Liq. Chromatogr.*, 16 (1993) 3445-3455.
- For additional information see *C.A.*:  
119 (1993) 96280r, 99057j, 108152r, 108159y, 118175q, 120473d, 135019j, 135020c, 145096w, 151410k, 155504s, 173298g, 173328s, 173334r, 173336t, 173337u, 173338v, 179163f, 181681s, 194683u, 194747t, 194814n.
- See also 14, 61, 182, 185, 188, 190, 201, 217, 262, 299, 471, 483, 500, 586, 595, 599, 806, 895, 896, 999, 1154, 1155, 1198, 1206, 1252, 1312.
- 3d. *Quantitative analysis*
- 127 Burrows, J.L.: Evaluation of calibration routines in pharmaceutical and bioanalytical separations, with reference to Monte Carlo simulations. *J. Pharm. Biomed. Anal.*, 11 (1993) 523-531.
- 128 Vigdergauz, M.S. and Krauze, I.M.: (Characteristics of quantitative chromatographic analysis). *Zh. Fiz. Khim.*, 67 (1993) 839-840; *C.A.*, 119 (1993) 194564f.
- 129 Vigdergauz, M.S. and Krauze, I.M.: (Development of a method for the quantitative interpretation of chromatograms using two internal standards). *Zh. Fiz. Khim.*, 67 (1993) 837-838; *C.A.*, 119 (1993) 194563e.
- 130 Wimley, W.C. and White, S.H.: Quantitation of electrostatic and hydrophobic membrane interactions by equilibrium dialysis and reverse-phase HPLC. *Anal. Biochem.*, 213 (1993) 213-217.
- For additional information see *C.A.*:  
119 (1993) 179506v.
- See also 49, 147, 330, 504, 1043, 1061, 1066.

## 3e. Preparative scale chromatography

- 131 Felinger, A. and Guiochon, G.: The change of pressure drop during large-scale chromatography of viscous samples. *Bio-technol. Prog.*, 9 (1993) 450-455; C.A., 119 (1993) 176862x.
- 132 Porsch, B.: Experimental consequences of differences between composition of the sample solvent and of the mobile phase in HPLC. *J. Liq. Chromatogr.*, 16 (1993) 3409-3421.
- 133 Samsonov, G.V.: (Preparative scale selective chromatography of biologically active substances (BAS) on heteronetwork, composite, and surface-laminated sorbents at low pressure). *Zh. Prikl. Khim. (S.-Peterburg)*, 65 (1992) 961-975; C.A., 119 (1993) 112523e - a review with 76 refs.
- 134 Viard, V. and Lameloise, M.L.: (Preparative chromatography: modeling on fixed bed and application to the performance of simulated mobile bed). *Recents Prog. Genie Procedes*, 5 (1991) 173-178; C.A., 119 (1993) 120171d.
- 135 Yamamoto, S. and Sano, Y.: Predicting the performance of production-scale chromatography. In: Furusaki, S., Endo, I. and Matsuno, R. (Editors), *Biochem. Eng. 2001, Proc. Asia-Pac. Biochem. Eng. Conf.*, Springer, Tokyo, 1992, pp. 548-550; C.A., 119 (1993) 115361m.

See also 10, 16, 47, 65, 190, 208, 209, 546, 589, 607, 757, 823.

## 3f. Programmed temperature, pressure, vapors, gradients

- 136 Moore, L.K. and Synovec, R.E.: Axial thermal gradient micro-bore liquid chromatography by flow programming. *Anal. Chem.*, 65 (1993) 2663-2670.

See also 1416.

## 4. SPECIAL TECHNIQUES

## 4a. Automation

- 137 Aloba, O.T., Bergman, D.E., Miller, R.T. and Daly, R.E.: Enhanced robotic dissolution system with concurrent off-line analysis. *J. Pharm. Biomed. Anal.*, 11 (1993) 861-869.
- 138 Davoli, E., Fanelli, R. and Bagnati, R.: Purification and analysis of drug residues in urine samples by on-line immunoaffinity chromatography/high-performance liquid chromatography/continuous-flow fast atom bombardment mass spectrometry. *Anal. Chem.*, 65 (1993) 2679-2685.
- 139 Jackson, C. and Yau, W.W.: Computer simulation study of size-exclusion chromatography with simultaneous viscometry and light scattering measurements. *J. Chromatogr.*, 645 (1993) 209-217.
- 140 McCoy, M.A., Liapis, A.I. and Unger, K.K.: Applications of mathematical modelling to the simulation of binary perfusion chromatography. *J. Chromatogr.*, 644 (1993) 1-9.
- 141 Mole, D., Mason, R.J. and McDowall, R.D.: The development of a strategy for the implementation of automation in a bioanalytical laboratory. *J. Pharm. Biomed. Anal.*, 11 (1993) 183-190.

See also 263, 470, 1055, 1230, 1379, 1383.

## 4b. Computerization and modelling

- 142 Ganetsos, G., Barker, P.E. and Akintoye, A.: Computer simulation of semicontinuous chromatographic refineries used in chemical and biochemical applications. *Chromatogr. Sci.*, 61 (1993) 769-782; C.A., 119 (1993) 98350n - a review with 22 refs.
- 143 Guillaume, Y. and Guinchard, C.: Optimizing mobile phase composition, its flow rate and column temperature in HPLC using an experimental design assisted with a simplex method. *J. Liq. Chromatogr.*, 16 (1993) 3457-3470.
- 144 Gureev, E.S., Nushtaeva, L.B. and Gulamov, L.M.: (Simulation of an extraction-chromatographic concentration process). *Radiokhimiya*, 35 (1993) 91-98; C.A., 119 (1993) 189252y.
- 145 Herold, E.G., Moll, W.T. and Grossman, S.J.: Operation of a centralized chromatography data system in a VAX-based, MS-DOS local area network environment. *Am. Lab. (Shelton)*, 25 (1993) 34H, 34J, 34L-34M; C.A., 119 (1993) 173061z.
- 146 Kaczmarek, K.: (Estimation of non-linear parameters of chromatographic column models). *Iz. Chem. Procesowa*, 13 (1992) 637-648; C.A., 119 (1993) 183595r.
- 147 Koskinen, M.O. and Koskinen, L.K.: Numerical integration of complex chromatograms using fitted Gaussian functions. *J. Liq. Chromatogr.*, 16 (1993) 3171-3184.
- 148 Kowalski, W.: (Mass transfer kinetics in liquid-liquid chromatography). *Inz. Chem. Procesowa*, 13 (1992) 609-619; C.A., 119 (1993) 183702y.
- 149 Kurbatova, S.V., Kolosova, E.A. and Vigdergauz, M.S.: (Operation of computerized chromatographic systems for the analysis of multicomponent mixtures). *Zh. Fiz. Khim.*, 67 (1993) 833-836; C.A., 119 (1993) 173284z.
- 150 Ouchi, G.I.: Creating control charts with a spreadsheet program. *LC-GC*, 11 (1993) 416-423; C.A., 119 (1993) 107998j.
- 151 Rigal, L., Phemius, P., Lattoque, S. and Monfort, J.P.: (Modeling of chromatographic separation in mobile bed of fructose-glucose mixture). *Recents Prog. Genie Procedes*, 5 (1991) 397-402; C.A., 119 (1993) 120174g.
- 152 Sebastian, M.J.: Extending the life of a chromatography data system by mixing computer platforms. *Chemom. Intell. Lab. Syst.*, 21 (1993) 51-64; C.A., 119 (1993) 130492y.
- 153 Seidel-Morgenstern, A. and Guiochon, G.: Modelling of the competitive isotherms and the chromatographic separation of two enantiomers. *Chem. Eng. Sci.*, 48 (1993) 2787-2797; C.A., 119 (1993) 167885e.
- 154 Smith, K.: Developments in the field of chromatography data handling. *Int. Lab.*, 23, No. 8 (1993) 20-22.

For additional information see C.A.:

119 (1993) 151193s, 161209r, 173096q, 183594q, 194754t.

See also 17, 48, 49, 56, 69, 105, 127, 134, 187, 680, 1091.

## 4c. Combination with other physico-chemical techniques (MS, IR etc.)

- 155 Banks, C.T.: The development and application of coupled HPLC-TLC for pharmaceutical analysis. *J. Pharm. Biomed. Anal.*, 11 (1993) 705-710.
- 156 Carey, J.M., Byrde, F.A. and Caruso, J.A.: Alternate methods of sample introduction for plasma mass spectrometry. *J. Chromatogr. Sci.*, 31 (1993) 330-344 - a review with 120 refs.

- 157 Chappell, C.G., Creaser, C.S. and Shepherd, M.J.: On-line high performance liquid chromatography - multidimensional gas chromatography and its application to the determination of stilbene hormones in corned beef. *J. High Resolut. Chromatogr.*, 16 (1993) 479-482.
- 158 Coulot, M., Domon, B., Grossenbacher, H., Guenat, C., Maerki, W., Mueller, D.R. and Richter, W.J.: LC-MS and MS/MS in the analysis of recombinant proteins. *J. Mol. Struct.*, 292 (1993) 89-103; *C.A.*, 119 (1993) 90710j.
- 159 Gagné, J.P., Carrier, A., Varfalvy, L. and Bertrand, M.J.: Evaluation of the performance of capillary liquid chromatography-fast atom bombardment mass spectrometry systems with precolumn addition of glycerol as a viscous matrix. *J. Chromatogr.*, 647 (1993) 13-20.
- 160 Gagné, J.P., Carrier, A., Varfalvy, L. and Bertrand, M.J.: Origin of the decrease in chromatographic resolution induced by the addition of viscous matrices in liquid chromatographic-fast atom bombardment mass spectrometric systems. *J. Chromatogr.*, 647 (1993) 21-29.
- 161 Genuit, W.J.L. and de Boer, J.: Applications of dual-beam thermospray liquid chromatography-mass spectrometry. *J. Chromatogr.*, 647 (1993) 73-78.
- 162 Hopfgartner, G., Bean, K., Henion, J. and Henry, R.: Ion spray mass spectrometric detection for liquid chromatography: a concentration- or a mass-flow-sensitive device? *J. Chromatogr.*, 647 (1993) 51-61.
- 163 Karlsson, K.-E.: Deuterium oxide as a reagent for the modification of mass spectra in electrospray microcolumn liquid chromatography-mass spectrometry. *J. Chromatogr.*, 647 (1993) 31-38.
- 164 Kienhuis, P.G.M.: Radiofrequency-only daughter scan mode to provide more spectral information in liquid chromatography-thermospray tandem mass spectrometry. *J. Chromatogr.*, 647 (1993) 39-50.
- 165 Krost, K.J.: Analysis of selected chemical groups by liquid chromatography/mass spectrometry. *Appl. Spectrosc.*, 47 (1993) 821-829; *C.A.*, 119 (1993) 108184c.
- 166 Mol, H.G.J., Janssen, H.-G.M., Cramers, C.A. and Brinkman, U.A.T.: Use of a temperature programmed injector with a packed liner for direct water analysis and on-line reversed phase LC-GC. *J. High Resolut. Chromatogr.*, 16 (1993) 459-463.
- 167 Mourey, T.H. and Balke, S.T.: A strategy for interpreting multidetector size-exclusion chromatography data. I. Development of a systematic approach. *ACS Symp. Ser.*, 521 (1993) 180-198; *C.A.*, 119 (1993) 161205m.
- 168 Müller, E. and Jork, H.: On-line coupling of HPLC, solid phase extraction, and TLC (HPLC-OSP-TLC). Assay of 4(5)-methylimidazole in caramel samples. *J. Planar Chromatogr.*, 6 (1993) 21-28.
- 169 Robins, R.H.: Probe-based electrospray adapter for thermospray-equipped quadrupole-based liquid chromatograph/mass spectrometer (LC/MS) systems. *U.S. Pat. Appl.* 5,235,186 (Cl. 250-288; B01D59/44), 10 Aug. 1993, Appl. 826,922, 24 Jan. 1992; 24 p.; *C.A.*, 119 (1993) 194818s.
- 170 Stillian, J.R., Barreto, V.M., Friedman, K.A., Rabin, S.B. and Toofan, M.: Ion-chromatograph system using electrochemical suppression and detector-effluent recycle. *Eur. Pat. Appl.* EP 555,962 (Cl. G01N30/96), 18 Aug. 1993, US Appl. 833,334, 10 Feb. 1992; 17 p.; *C.A.*, 119 (1993) 194813m.
- 171 Tinke, A.P., van der Hoeven, R.A.M., Niessen, W.M.A., Tjaden, U.R. and van der Greef, J.: New ionization strategies in particle-beam liquid chromatography-mass spectrometry based on the principle of surface ionization. *J. Chromatogr.*, 647 (1993) 63-72.
- For additional information see *C.A.*:  
119 (1993) 90720n, 90721p, 103462g, 112869x, 128049r.
- See also 66, 70, 75, 78, 138, 201, 230, 250, 255, 268, 275, 282, 323, 352, 393, 404, 424, 427, 434, 449, 451, 513, 522, 548, 563, 566, 567, 649, 755, 855, 858, 880, 881, 937, 966, 983, 984, 985, 1002, 1026, 1035, 1067, 1075, 1085, 1086, 1090, 1104, 1107, 1120, 1126, 1167, 1180, 1200, 1220, 1229, 1261, 1271, 1283, 1298, 1308, 1346, 1355, 1358, 1366, 1385, 1399, 1409.
- 4d. *Affinity chromatography (advances)*
- 172 Caron, M., Joubert-Caron, R., Cartier, J.R., Chadli, A. and Bladier, D.: Study of lectin-ganglioside interactions by high-performance liquid affinity chromatography. *J. Chromatogr.*, 646 (1993) 327-333.
- 173 Choe, T., Chung, I. and Kim, J.: Determination of elution conditions of immunoaffinity chromatography by ELISA (enzyme linked immunosorbent assay) method. In: Furusaki, S., Endo, I. and Matsuno, R. (Editors), *Biochem. Eng. 2001, Proc. Asia-Pac. Biochem. Eng. Conf.*, Springer, Tokyo, 1992, pp. 569-571; *C.A.*, 119 (1993) 115362n.
- 174 Jarrett, H.W.: Affinity chromatography with nucleic acid polymers. *J. Chromatogr.*, 618 (1993) 315-339 - a review with 88 refs.
- 175 Kabzinski, A.K.M. and Paryczak, T.: (Study of properties of immobilized enzymes by rapid immobilization on supports for affinity chromatography). *Przem. Chem.*, 72 (1993) 202-204; *C.A.*, 119 (1993) 112233k.
- 176 Katoh, S., Sohma, Y., Terashima, M. and Sada, E.: Immunorecognition of anti-peptide antibodies and application to separation of biomolecules. In: Furusaki, S., Endo, I. and Matsuno, R. (Editors), *Biochem. Eng. 2001, Proc. Asia-Pac. Biochem. Eng. Conf.*, Springer, Tokyo, 1992, pp. 509-512; *C.A.*, 119 (1993) 115356p.
- 177 Van Sommeren, A.P.G., Machielsens, P.A.G.M. and Gribnau, T.C.J.: Comparison of three activated agaroses for use in affinity chromatography: effects on coupling performance and ligand leakage. *J. Chromatogr.*, 639 (1993) 23-31.
- 178 Wheatley, J.B. and Schmidt, D.E., Jr.: Salt-induced immobilization of proteins on a high-performance liquid chromatographic epoxide affinity support. *J. Chromatogr.*, 644 (1993) 11-16.
- See also 5, 109, 123, 138, 267, 315, 526, 587, 598, 606, 608, 626, 733, 807, 817, 848.
- 4f. *Trace analysis and pre-separation techniques*
- 179 Kaneko, E., Hoshino, H. and Yotsuyanagi, T.: (Development of solvent substitution preconcentration method at the microliter level for ion-pair extraction-reversed-phase partition HPLC). *Biomed. Res. Trace Elem.*, 3 (1992) 207-208; *C.A.*, 119 (1993) 134854r.

- 180 Lau, S.-L. and Stenstrom, M.K.: Discussion on "Alternatives to methanol-water elution of solid-phase extraction columns for the fractionation of high log  $K_{ow}$  organic compounds in aqueous environmental samples" by Durhan et al., *J. Chromatogr.*, 629 (1993) 67-74. *J. Chromatogr.*, 646 (1993) 439-441.
- 181 Meyer, A., Kleiböhmer, W. and Cammann, K.: SFE of PAHs from soils with a high carbon content and analyte collection via combined liquid-solid trapping. *J. High Resolut. Chromatogr.*, 16 (1993) 491-494.

For additional information see C.A.:  
119 (1993) 155162d.

See also 106, 168, 205, 460, 470, 940, 1098, 1116, 1239, 1255, 1282, 1402, 1407, 1427, 1438.

#### 4g. Enantiomers, separation

- 182 Felix, G. and Zhang, T.: An amylopectin tris(phenylcarbamate) chiral stationary phase for high performance liquid chromatography. *J. High Resolut. Chromatogr.*, 16 (1993) 364-367.
- 183 Fujima, H., Kitagawa, H., Kitamura, H. and Wada, H.: (Characteristics of ovomucoid-bonded chiral separation column). *Kuro-matogurafi*, 13 (1992) 385-394; *C.A.*, 119 (1993) 112573w.
- 184 Hansson, L. and Isaksson, R.: Chromatographic determination of enantiomeric purity by achiral means. *J. Chromatogr.*, 653 (1993) 9-14.
- 185 Hu, W. and Haraguchi, H.: Enantiomeric separation with use of stationary phase coated with micellar bile salt for microcolumn liquid chromatography. *Bull. Chem. Soc. Jpn.*, 66 (1993) 1967-1970; *C.A.*, 119 (1993) 194767z.
- 186 Kurata, T. and Ando, S.: (Efficient chiral separation with a simulated moving-bed (SMB) process). *Res. Dev. Rev.-Mitsubishi Kasei Corp.*, 6 (1992) 102-105; *C.A.*, 119 (1993) 180099w.
- 187 Lipkowitz, K.B.: Atomistic molecular modeling in chiral chromatography. *Int. Lab.*, 23, No. 8 (1993) P8-P12.
- 188 Marle, I., Jönsson, S., Isaksson, R. and Pettersson, C.: Chiral stationary phases based on intact and fragmented cellobiohydrolase I immobilized on silica. *J. Chromatogr.*, 648 (1993) 333-347.
- 189 Oi, S., Shijo, M., Tanaka, H., Miyano, S. and Yamashita, J.: Chiral stationary phases consisting of axially dissymmetric 2'-substituted-1,1'-binaphthyl-2-carboxylic acids bonded to silica gel for high-performance liquid chromatographic separation of enantiomers. *J. Chromatogr.*, 645 (1993) 17-28.
- 190 Sinibaldi, M., Castellani, L., Federici, F. and Messina, A.: High-performance liquid displacement chromatography of enantiomers on a chiral poly-L-valyl-based stationary phase. *J. Liq. Chromatogr.*, 16 (1993) 2977-2992.
- 191 Wan, H., Wang, Y., Ou, Q. and Yu, W.: Improved enantiomeric separation with a 2,6-di-O-pentyl-3-O-trifluoroacetylated  $\beta$ -cyclodextrin and OV-7 mixed stationary phase chiral capillary column. *J. Chromatogr.*, 644 (1993) 202-207.

For additional information see C.A.:  
119 (1993) 112575y, 194764w.

See also 9, 87, 91, 93, 107, 121, 144, 153, 210, 237, 246, 328, 336, 349, 446, 453, 461, 476, 481, 486, 492, 496, 497, 500, 502, 516, 754, 875, 936, 958, 1022, 1041, 1147, 1155, 1163, 1174, 1176, 1177, 1181, 1182, 1203, 1206, 1207, 1208, 1214, 1218, 1223, 1225, 1230, 1236, 1242, 1254, 1262, 1266, 1276, 1311, 1324, 1333, 1334, 1335, 1336.

#### 4h. Other special techniques

- 192 Anderson, N.G. and Anderson, N.L.: Rotating processor for solid-phase reactions. *U.S. US 5,186,824* (Cl. 210-198.2; BO1D15/08), 16 Feb. 1993, Appl. 754,892, 04 Sep. 1991; 35 pp.; *C.A.*, 119 (1993) 90763d.
- 193 Barnes, N.: Developments in ion exchange chromatography. *Int. Lab.*, 23, No. 9 (1993) P4-P8.
- 194 Menet, J.-M. and Ito, Y.: Studies on a new cross-axis coil planet centrifuge for performing counter-current chromatography. II. Path and acceleration of coils and comparison with type J coil planet centrifuge. *J. Chromatogr.*, 644 (1993) 231-238.
- 195 Menet, J.-M., Shinomiya, K. and Ito, Y.: Studies on new cross-axis coil planet centrifuge for performing counter-current chromatography. III. Speculations on the hydrodynamic mechanism in stationary phase retention. *J. Chromatogr.*, 644 (1993) 239-252.
- 196 Okada, T.: (Field-flow fractionation using potential-controlled chromatography). *Kagaku (Kyoto)*, 48 (1993) 427; *C.A.*, 119 (1993) 130424c - a review with 5 refs.
- 197 Shinomiya, K., Menet, J.-M., Fales, H.M. and Ito, Y.: Studies on a new cross-axis coil planet centrifuge for performing counter-current chromatography. I. Design of the apparatus, retention of the stationary phase, and efficiency in the separation of proteins with polymer phase systems. *J. Chromatogr.*, 644 (1993) 215-229.
- 198 Wang, Z.: (Application of flash chromatography). *Zhongguo YiYao Gongye Zazhi*, 24 (1993) 92; *C.A.*, 119 (1993) 188682b - a review with 3 refs.

See also 22, 65, 67, 114, 220, 634, 1112, 1144, 1206, 1397, 1398, 1404.

## 5. HYDROCARBONS AND HALOGEN DERIVATIVES

### 5a. Aliphatic hydrocarbons

- 199 Akhlaq, M.S.: Rapid group-type analysis of crude oils using high-performance liquid chromatography and gas chromatography. *J. Chromatogr.*, 644 (1993) 253-258.
- 200 Hufton, J.R. and Danner, R.P.: Chromatographic study of alkanes in silicalite: transport properties. *AIChE J.*, 39 (1993) 962-974; *C.A.*, 119 (1993) 147042z.

See also 46.

5b. *Cyclic hydrocarbons, fullerenes*

- 201 Anacleto, J.F. and Quilliam, M.A.: Liquid chromatography/mass spectrometry investigation of the reversed-phase separation of fullerenes and their derivatives. *Anal. Chem.*, 65 (1993) 2236-2242.
- 202 Cabrera, K., Wieland, G. and Schäfer, M.: High-performance liquid chromatographic separation of fullerenes (C<sub>60</sub> and C<sub>70</sub>) using chemically bonded  $\gamma$ -cyclodextrin as stationary phase. *J. Chromatogr.*, 644 (1993) 396-399.
- 203 Chen, J., MacLeod, M.C., Zhao, R. and Geacintov, N.E.: Fluorescence HPLC methods for detecting benzo[a]pyrene-7,8-dihydrodiol 9,10-oxide-deoxyadenosine adducts in enzyme-digests of modified DNA: Improved sensitivity. *Carcinogenesis*, 14 (1993) 1049-1051; *C.A.*, 119 (1993) 110661z.
- 204 Chen, X. and Xiao, Y.: (Metabolism of the 1- and 3-nitrobenzo[a]pyrene mixture by rat liver microsomes). *Weisheng Dulixue Zazhi*, 6 (1992) 249-251; *C.A.*, 119 (1993) 110968e.
- 205 Dumont, J., Larocque-Lazure, F. and Iorio, C.: An alternative isolation procedure for the subsequent determination of benzo(a)pyrene in total particulate matter of cigarette smoke. *J. Chromatogr. Sci.*, 31 (1993) 371-374.
- 206 García, M.A., Vera, S., Bombin, M. and Marina, M.L.: Optimization of the separation selectivity of a group of benzene and naphthalene derivatives in micellar high-performance liquid chromatography using a C<sub>18</sub> column and alcohols as modifiers in the mobile phase. *J. Chromatogr.*, 646 (1993) 297-305.
- 207 Gügel, A. and Müllen, K.: Preparative separation of fullerene adducts by liquid chromatography on polystyrene gel. *Chromatographia*, 37 (1993) 387-391.
- 208 Herren, D., Thilgen, C., Calzaferri, G. and Diederich, F.: Preparative separation of higher fullerenes by high-performance liquid chromatography on a tetrachlorophthalimidopropyl-modified silica column. *J. Chromatogr.*, 644 (1993) 188-192.
- 209 Jinno, K., Uemura, T., Ohta, H., Nagashima, H. and Itoh, K.: Separation and identification of higher molecular weight fullerenes by high-performance liquid chromatography with monomeric and polymeric octadecylsilica bonded phases. *Anal. Chem.*, 65 (1993) 2650-2654.
- 210 Karlsson, A., Luthman, K., Pettersson, C. and Hacksell, U.: Enantioselective ion pair chromatography of phenolic 2-dipropylaminotetralin derivatives on achiral stationary phases: an experimental and theoretical study of chiral discrimination. *Acta Chem. Scand.*, 47 (1993) 469-481; *C.A.*, 119 (1993) 108163v.
- 211 Lehotay, J., Haimo, F., Brandšteterová, E. and Oktavec, D.: HPLC-DAD determination of benzo(a)pyrene in waste gas. *J. Liq. Chromatogr.*, 16 (1993) 3575-3584.
- 212 Marriott, P.J., Carpenter, P.D., Brady, P.H., McCormick, M.J., Griffiths, A.J., Hatvani, T.S.G. and Raddell, S.G.: Optimisation of fluorescence detection for polyaromatic hydrocarbon determination by using high performance liquid chromatography. *J. Liq. Chromatogr.*, 16 (1993) 3229-3247.
- 213 Millar, J.M., Greegan, K.M., Robbins, J.L., Robbins, W.K., Sherwood, R.D., Tindall, P.J. and Cox, D.M.: Chromatographic separations of fullerenes: discovery and characterization of C<sub>60</sub> mono-epoxide. *Synt. Met.*, 59 (1993) 317-331; *C.A.*, 119 (1993) 194241y.
- 214 Nondek, L., Kuzilek, M. and Krupicka, S.: LC Clean-up and GC/MS analysis of polycyclic aromatic hydrocarbons in river sediment. *Chromatographia*, 37 (1993) 381-386.

- 215 Rose, E., Martens, D., Lintelmann, J., Kicinski, H.G., Günther, W.J. and Kettrup, A.: Behaviour of polycyclic aromatic hydrocarbons and triazine herbicides in water and aquifer material of drinking water recharge plant. Part IV. PAHS in solid phase. *Fresenius J. Anal. Chem.*, 347 (1993) 44-51.
- 216 Wu, Y., Sun, Y., Gu, Z., Wang, Q., Zhou, X., Xiong, Y. and Jin, Z.: Analysis of C<sub>60</sub> and C<sub>70</sub> fullerenes by high-performance liquid chromatography. *J. Chromatogr.*, 648 (1993) 491-496.

For additional information see C.A.:

119 (1993) 163462s, 173108v, 194781z.

See also 89, 98, 100, 103, 181, 199, 222, 449, 1383.

5c. *Halogen derivatives*

- 217 Kimata, K., Hosoya, K., Araki, T., Tanaka, N., Barnhart, E.R., Alexander, L.R., Sirimanne, S., McClure, P.C., Crainger, J. and Patterson, D.G., Jr.: Electron-acceptor and electron-donor chromatographic stationary phases for the reversed-phase liquid chromatographic separation and isomer identification of polychlorinated dibenzo-p-dioxins. *Anal. Chem.*, 65 (1993) 2502-2509.
- 218 Tonelli, C., Turri, S., Gianotti, G. and Levi, M.: Chromatographic process for the separation of the non-, mono- and bifunctional species in perfluoro polyoxyalkylenes. *Eur. Pat. Appl. EP 538,827* (Cl. C08G65/30), 28 Apr. 1993, IT Appl. 91/Mi2774, 21 Oct. 1991; 10 p.; *C.A.*, 119 (1993) 181577f.

See also 116, 199.

5d. *Complex hydrocarbon mixtures (incl. analysis of tars, bitumens and mineral oils)*

- 219 Khan, Z.H., Marron, K. and Assi, S.H.: The development of less hazardous eluent mixtures for the liquid chromatographic separation of total polycyclic aromatic hydrocarbons in heavy hydrocarbon oil mixtures. *J. Liq. Chromatogr.*, 16 (1993) 1473-1487.
- 220 Menges, R.A., Spino, L.A. and Armstrong, D.W.: Separation and characterization of components of catalytic cracker feed using centrifugal partition chromatography. *Anal. Chem.*, 65 (1993) 2873-2877.
- 221 Sanchez, V. and Ceballo, C.D.: (Boiling-point distribution of heavy and extraheavy crude petroleum by gel-permeation chromatography). *Rev. Tec. INTEVEP*, 10 (1990) 145-150; *C.A.*, 119 (1993) 99363f.

For additional information see C.A.:

119 (1993) 184428a.

See also 110, 967.

## 6. ALCOHOLS

- 222 Lübke, M., le Quéré, J.-L. and Barron, D.: Retention behaviour of volatile compounds in normal-phase high-performance liquid chromatography on a diol column. *J. Chromatogr.*, 646 (1993) 307-316.

- 223 Meyer, T., Harms, D. and Gmehling, J.: Analysis of polyethylene glycols with respect to their oligomer distribution by high-performance liquid chromatography. *J. Chromatogr.*, 645 (1993) 135-139.
- 224 Nivinskene, O.Yu.: (Chromatographic determination of 2-butyne-1,4-diol and its transformation products formed during nickel electrodeposition). *Zashch. Met.*, 28 (1992) 1028-1032; *C.A.*, 119 (1993) 130688s.

For additional information see *C.A.*:  
119 (1993) 194790b.

See also 18, 82, 182, 189, 249, 1145, 1370.

## 7. PHENOLS

- 225 Barrero, A.F., Sánchez, J.F. and Rodríguez, I.: Use of alkenylresorcinols from *Ononis speciosa* as synthetic precursors of compounds with potential biological activity. *J. Natural Prod.*, 56 (1993) 1737-1746.
- 226 Dem'yanov, P.I.: (Chemical derivatization in chromatographic determination). *Zh. Anal. Khim.*, 47 (1992) 1942-1966; *C.A.*, 119 (1993) 108145r - a review with 187 refs.
- 227 Lee, B.L., Ong, H.Y., Shi, C.Y. and Ong, C.N.: Simultaneous determination of hydroquinone, catechol and phenol in urine using high-performance liquid chromatography with fluorimetric detection. *J. Chromatogr.*, 619 (1993) 259-266.
- 228 Park, J.H., Lee, Y.K., Cheong, N.Y. and Vang, M.D.: Reversed phase liquid chromatographic separation of some mono-substituted phenols with calix[6]arene-*p*-sulfonate-modified eluents. *Chromatographia*, 37 (1993) 221-223.
- 229 Webster, C., Smith, M., Wilson, P. and Cooke, M.: Determination of the total phenol content of soils by high speed liquid chromatography with electrochemical detection. *J. High Resolut. Chromatogr.*, 16 (1993) 549-551.
- 230 Wolfender, J.L. and Hostettmann, K.: Liquid chromatographic-UV detection and liquid chromatographic-thermospray mass spectrometric analysis of *Chironia* (Gentianaceae) species. *J. Chromatogr.*, 647 (1993) 191-202.

See also 18, 24, 41, 89, 222, 1358, 1368, 1390.

## 8. SUBSTANCES CONTAINING HETEROCYCLIC OXYGEN

### 8a. Flavonoids

- 231 Amic, D., Davidovic-Amic, D. and Trinajstić, N.: Application of topological indices to chromatographic data. Calculation of the retention indices of anthocyanins. *J. Chromatogr.*, 653 (1993) 115-121.
- 232 Tschiersch, K. and Hoelzl, J.: (Detection and analysis of acyl derivatives of apigenin 7- $\beta$ -D-glucoside). *PZ Wiss*, 5 (1992) 208-214; *C.A.*, 119 (1993) 125292d.

For additional information see *C.A.*:  
119 (1993) 146693a, 167852s.

See also 182, 1354.

### 8b. Aflatoxins and other mycotoxins

- 233 Breitholtz-Emanuelsson, A., Olsen, M., Oskarsson, A., Palminger, I. and Hult, K.: Ochratoxin A in cow's milk and human milk with corresponding human blood samples. *J. Assoc. Off. Anal. Chem.*, 76 (1993) 842-846.
- 234 Frisvad, J.C. and Thrane, U.: Liquid column chromatography of mycotoxins. *J. Chromatogr. Libr.*, Vol 54(Chromatography of Mycotoxins), Elsevier, Amsterdam, 1993, pp. 253-372; *C.A.*, 119 (1993) 132959e.
- 235 Garner, R.C., Whattam, M.M., Taylor, P.J.L. and Stow, M.W.: Analysis of United Kingdom purchased spices for aflatoxins using an immunoaffinity column clean-up procedure followed by high-performance liquid chromatographic analysis and post-column derivatisation with pyridinium bromide perbromide. *J. Chromatogr.*, 648 (1993) 485-490.
- 236 Scott, P.M.: Recent developments in methods of analysis for mycotoxins in foodstuffs. *TrAC*, 12 (1993) 373-381 - a review with 37 refs.

For additional information see *C.A.*:  
119 (1993) 115638g.

### 8c. Other compounds with heterocyclic oxygen (incl. tannins)

- 237 Geerlof, A., van Tol, J.B.A., Jongejan, J.A. and Duine, J.A.: Methods for the determination of the enantiomeric purity of the C<sub>3</sub>-synthons glycidol (2,3-epoxy-1-propanol) and solketal [2,2-dimethyl-4-(hydroxymethyl)-1,3-dioxolane]. *J. Chromatogr.*, 648 (1993) 119-129.
- 238 Roig, M.G., Bello, J.F., Kennedy, J.F., Rivera, Z.S. and Lloyd, L.L.: A reversed-phase HPLC method for measurement of 5-hydroxymethyl furfuraldehyde and furfuraldehyde in processed juices. *Bioseparation*, 3 (1992) 177-184; *C.A.*, 119 (1993) 115881f.
- 239 Speranza, G., Manitto, P., Cassara, P., Monti, D., de Castri, D. and Chiaiva, F.: Studies on aloe, 12. Furoaloesone, a new 5-methylchromone from Cape aloe. *J. Natural Prod.*, 56 (1993) 1089-1094.

See also 217, 222, 231, 1210, 1252.

## 9. OXO COMPOUNDS, ETHERS, EPOXIDES AND QUINONES

- 240 Attygalle, A.B., Xu, S.-C., Meinwald, J. and Eisner, T.: Defensive secretion of the millipede *Floridobolus penneri*. *J. Natural Prod.*, 56 (1993) 1700-1706.
- 241 Cais Fonseca, M.I. and Joglar Chitas, M.I.: (Formaldehyde in particleboard: Determination by high-pressure liquid chromatography (HPLC)). *Rev. Port. Farm.*, 42 (1992) 18-24; *C.A.*, 119 (1993) 183159b.
- 242 Cohen, H.P. and Tway, P.C.: High performance liquid chromatographic detection of residual formaldehyde in a hepatitis-A vaccine by use of hydralazine. *J. Liq. Chromatogr.*, 16 (1993) 1667-1684.

- 243 Coutrim, M.X., Nakamura, L.A. and Collins, C.H.: Quantification of 2,4-dinitrophenylhydrazones of low molecular mass aldehydes and ketones using HPLC. *Chromatographia*, 37 (1993) 185-190.
- 244 Fujita, M., Sano, M., Takeda, K. and Tomita, I.: Fluorescence detection of glutathione S conjugate with aldehyde by high-performance liquid chromatography with post-column derivatization. *Analyst (London)*, 118 (1993) 1289-1292.
- 245 Grömping, A.H.J., Karst, U. and Cammann, K.: Development of a method for simultaneous determinations of nitrogen oxides, aldehydes and ketones in air samples. *J. Chromatogr.*, 653 (1993) 341-347.
- 246 Miller, L. and Weyker, C.: Effects of compound structure and temperature on the resolution of enantiomers of cyclopentenones by liquid chromatography on derivatized cellulose chiral stationary phases. *J. Chromatogr.*, 653 (1993) 219-228.
- 247 Traoré, F., Farinotti, R. and Mahuzier, G.: Determination of malonaldehyde by coupled high-performance liquid chromatography-spectrofluorimetry after derivatization with luminarin 3. *J. Chromatogr.*, 648 (1993) 111-118.
- See also 24, 182, 222, 237, 238, 345, 442, 880, 938, 1112.
10. CARBOHYDRATES
- 10a. *Mono and oligosaccharides. Structural studies*
- 248 Adams, P., Zegeer, A., Bohnert, H.J. and Jensen, R.G.: Anion exchange separation and pulsed amperometric detection of inositols from flower petals. *Anal. Biochem.*, 214 (1993) 321-324.
- 249 Buttler, T.A., Johansson, K.A.J., Gorton, L.G.O. and Marko-Varga, G.A.: On-line fermentation process monitoring of carbohydrates and ethanol using tangential flow filtration and column liquid chromatography. *Anal. Chem.*, 65 (1993) 2628-2636.
- 250 Davies, M.J., Smith, K.D., Carruthers, R.A., Chai, W., Lawson, A.M. and Hounsell, E.F.: Use of a porous graphitised carbon column for the high-performance liquid chromatography of oligosaccharides, alditols and glycopeptides with subsequent mass spectrometry analysis. *J. Chromatogr.*, 646 (1993) 317-326.
- 251 Ganetsos, G. and Barker, P.E.: Semicontinuous countercurrent chromatographic refiners. *Chromatogr. Sci.*, 61 (1993) 233-255; C.A., 119 (1993) 98349u - a review with 26 refs.
- 252 Haginaka, J., Nishimura, Y. and Yasuda, H.: Determination of cyclodextrins in serum by reversed-phase chromatography with pulsed amperometric detection and a membrane reactor. *J. Pharm. Biomed. Anal.*, 11 (1993) 1023-1026.
- 253 Hicks, K.B. and Hotchkiss, A.T., Jr.: High performance liquid chromatography of cereal-derived carbohydrates. In: Alexander, R.J. and Zobel, H.F. (Editors), *Dev. Carbohydr. Chem.*, Am. Assoc. Cereal Chem., St. Paul, 1992, pp. 239-259; C.A., 119 (1993) 93772s.
- 254 Honda, S., Ueno, T., Ueda, M., Kakehi, K., Nishiura, S. and Tachibana, S.: (Micro analysis of glycosaminoglycan-derived oligosaccharides by HPLC). *Kuromatogurafi*, 13 (1992) 235-236; C.A., 119 (1993) 134673f.
- 255 Hugener, M., Tinke, A.P., Niessen, W.M.A., Tjaden, U.R. and van der Greef, J.: Pseudo-electrochromatography-negative-ion electrospray mass spectrometry of aromatic glucuronides and food colours. *J. Chromatogr.*, 647 (1993) 375-385.
- 256 Kakehi, K. and Honda, S.: Analysis of carbohydrates in glycoproteins by high-performance liquid chromatography and high-performance capillary electrophoresis. *Methods Mol. Biol. (Totowa)*, 14 (Glycoprotein Analysis in Biomedicine) (1993) 81-97; C.A., 119 (1993) 90318n - a review with 11 refs.
- 257 Karamanos, N.K.: High-performance liquid chromatographic determination of xylitol and hexosaminotols present in the reduced terminal of glycosaminoglycans. *J. Liq. Chromatogr.*, 16 (1993) 2639-2652.
- 258 Kiba, N., Goto, Y. and Furusawa, M.: Simultaneous determination of glucose and 1-deoxyglucose in serum by anion-exchange chromatography with an immobilized pyranose oxidase reactor. *J. Chromatogr.*, 620 (1993) 9-13.
- 259 Kobayashi, N. and Murakita, H.: (Simultaneous determinations of monosaccharides, oligosaccharides and molecular weight distribution of polysaccharide using column switching technique). *Kuromatogurafi*, 13 (1992) 251-252; C.A., 119 (1993) 108179e.
- 260 Kurosui, Y. and Sato, Y.: (Separation analysis of bichromophorically derivatized glycopyranosides). *Kuromatogurafi*, 13 (1992) 249-250; C.A., 119 (1993) 95948c.
- 261 Lamb, J.D., Myers, G.S. and Edge, N.: Ion chromatographic analysis of glucose, fructose, and sucrose concentrations in raw and processed vegetables. *J. Chromatogr. Sci.*, 31 (1993) 353-357.
- 262 Marioli, J.M., Luo, P.F. and Kuwana, T.: Nickel-chromium alloy electrode as a carbohydrate detector for liquid chromatography. *Anal. Chim. Acta*, 282 (1993) 571-580.
- 263 Marko-Varga, G., Dominguez, E., Persson, L.L. and Gorton, L.: Automated off-line sample clean-up of biotechnological samples by the use of robotics for the analysis of carbohydrates. *Lab. Rob. Autom.*, 5 (1993) 11-22; C.A., 119 (1993) 179216a.
- 264 Matsushashi, S., Nishikawa, N., Negishi, T. and Hatanaka, C.: Enzymic-HPLC determination of the amount and distribution of the galacturonan region in pectate molecules. *J. Liq. Chromatogr.*, 16 (1993) 3203-3215.
- 265 Meurens, M., Dupuy, N., Mirouze, F.D., Legrand, P. and Huvenne, J.P.: NIR and MID-IR analysis of isomerized glucose syrup. In: Hildrum, K.I. (Editor), *Near Infra-Red Spectrosc.*, Horwood, Chichester, 1992, pp. 47-50; C.A., 119 (1993) 179479p.
- 266 Mirmira, S.R., Schreiber, M.A. and Guessford, S.A.: High performance ion chromatographic analysis of glucose, isomaltose, and maltose in hydroxyethyl starch. *J. Liq. Chromatogr.*, 16 (1993) 2631-2638.
- 267 Munro, P.D., Winzor, D.J. and Cann, J.R.: Experimental and theoretical studies of rate constant evaluation by affinity chromatography. Determination of rate constants for the interaction of saccharides with concanavalin A. *J. Chromatogr.*, 646 (1993) 3-15.
- 268 Niessen, W.M.A., van der Hoeven, R.A.M., van der Greef, J., Schols, H.A., Voragen, A.G.J. and Bruggink, C.: Recent progress in high-performance anion-exchange chromatography-thermospray mass spectrometry of oligosaccharides. *J. Chromatogr.*, 647 (1993) 319-327.

- 269 Preller, A., Guixé, V., Torres, J.C., Hofer, H.W. and Ureta, T.: The separation and identification of picomole amounts of intermediates of glucose metabolism by high performance liquid chromatography on pellicular resins. *Biol. Res.*, 25 (1992) 73-78; *C.A.*, 119 (1993) 134696r.
- 270 Saito, K., Quearry, B.J., Saito, M., Nowak, T.S., Jr., Markey, S.P. and Heyes, M.P.: Kynurenine 3-hydroxylase in brain: species activity differences and effect of gerbil cerebral ischemia. *Arch. Biochem. Biophys.*, 307 (1993) 104-109.
- 271 Shibata, N., Kojima, C., Satoh, Y., Satoh, R., Suzuki, A., Kobayashi, H. and Suzuki, S.: Structural study of a cell-wall mannan of *Saccharomyces kluyveri* IFO 685 strain. Presence of a branched side chain and  $\beta$ -1,2-linkage. *Eur. J. Biochem.*, 217 (1993) 1-12.
- 272 Simms, P.J., Haines, R.M. and Hicks, K.B.: High-performance liquid chromatography of neutral oligosaccharides on a  $\beta$ -cyclodextrin bonded phase column. *J. Chromatogr.*, 648 (1993) 131-137.
- 273 Spies, T.: (HPLC of neutral carbohydrates). *Lebensm.-Biotechnol.*, 10 (1993) 82-84; *C.A.*, 119 (1993) 115600p - a review with 38 refs.
- 274 Takeuchi, T., Miwa, T., Hemmi, A. and Maeda, T.: Indirect conductimetric detection of cyclodextrins in liquid chromatography. *J. High Resolut. Chromatogr.*, 16 (1993) 419-421.
- 275 Tinke, A.P., van der Hoeven, R.A.M., Niessen, W.M.A., van der Greef, J., Vincken, J.-P. and Schols, H.A.: Electrospray mass spectrometry of neutral and acidic oligosaccharides: methylated cyclodextrins and identification of unknowns derived from fruit material. *J. Chromatogr.*, 647 (1993) 279-287.
- 276 Townsend, R.R.: Quantitative monosaccharide analysis of glycoproteins. High-performance liquid chromatography. *ACS Symp. Ser.*, 529(Chromatography in Biotechnology) (1993) 86-101; *C.A.*, 119 (1993) 176801b - a review with 36 refs.
- 277 Valiente, O., Fernandez, L.E., Perez, R.M. and Marquina, G.: Partial methanolysis of the agar-type sulfated galactan of the red seaweed *Laurencia gemmifera*. *Carbohydr. Res.*, 243 (1993) 191-197; *C.A.*, 119 (1993) 134684k.
- 278 Zhang, R., Zhang, Z., Liu, G., Hidaka, Y. and Shimonishi, Y.: High-performance liquid chromatographic determination of neutral and amino monosaccharides by ultraviolet and fluorescence detection of sugar 9-fluorenylmethoxycarbonyl hydrazones and 9-fluorenylmethoxycarbonyl amino sugars at picomole and subpicomole levels. *J. Chromatogr.*, 646 (1993) 45-52.
- For additional information see *C.A.*:  
119 (1993) 130682k, 134687p, 137713e, 154509k, 173281w, 176883e, 176885g.
- See also 1, 4, 117, 134, 142, 151, 285, 301, 312, 313, 314, 613, 629, 642, 768, 776, 816, 897, 974, 980, 1368.
- 10b. *Polysaccharides, mucopolysaccharides, lipopolysaccharides*
- 279 Barth, H.G. and Regnier, F.E.: High-performance gel-permeation chromatography of industrial gums: analysis of pectins and water-soluble celluloses. *Methods Carbohydr. Chem.*, 9 (1993) 105-114; *C.A.*, 119 (1993) 119876f - a review with 35 refs.
- 280 Birosel-Boettcher, N.: Measurement of starch in paper by high-pressure liquid chromatography. *Tappi J.*, 76 (1993) 207-208; *C.A.*, 119 (1993) 162691d.
- 281 Campos, A., Nunez, R., Koenig, C.S., Carey, D.J. and Brandan, E.: A lipid-anchored heparan sulfate proteoglycan is present in the surface of differentiated skeletal muscle cells. *Eur. J. Biochem.*, 216 (1993) 587-595.
- 282 Da Col, R., Silvestro, L., Naggi, A., Torri, G., Baiocchi, C., Moltrasio, D., Cedro, A. and Viano, I.: Characterization of the chemical structure of sulphated glycosaminoglycans after enzymatic digestion. Application of liquid chromatography-mass spectrometry with an atmospheric pressure interface. *J. Chromatogr.*, 647 (1993) 289-300.
- 283 Evanko, S.P. and Vogel, K.G.: Proteoglycan synthesis in fetal tendon is differentially regulated by cyclic compression *in vitro*. *Arch. Biochem. Biophys.*, 307 (1993) 153-164.
- 284 Gässler, N., Reissner, C., Janzen, N., Kähnert, H. and Kleesiek, K.: A high performance liquid chromatography method for the determination of glycosaminoglycans in human blood. *Eur. J. Clin. Chem. Clin. Biochem.*, 31 (1993) 503-511.
- 285 Hanisch, F.-G., Chai, W., Rosankiewicz, J.R., Lawson, A.M., Stoll, M.S. and Feizi, T.: Core-typing of O-linked glycans from human gastric mucins. Lack of evidence for the occurrence of the core sequence Gal1-6GalNAc. *Eur. J. Biochem.*, 217 (1993) 645-655.
- 286 Hayakawa, K., de Felice, C., Watanabe, T., Tanaka, T., Iinuma, K., Nihei, K., Higuchi, S., Ezoe, T., Hibi, I. and Kurosawa, K.: Determination of free N-acetylneuraminic acid in human body fluids by high-performance liquid chromatography with fluorimetric detection. *J. Chromatogr.*, 620 (1993) 25-31.
- 287 Hennessey, J.P., Jr., Bednar, B. and Manam, V.: Molecular size analysis of *Haemophilus influenzae* type B capsular polysaccharide. *J. Liq. Chromatogr.*, 16 (1993) 1715-1729.
- 288 Kamo, I., Kikuchi, A., Nonaka, I., Yamada, E. and Kondo, J.: Haemopoietic activity associated with biglycan like proteoglycan. *Biochem. Biophys. Res. Commun.*, 195 (1993) 1119-1126.
- 289 Kasinath, B.S.: Glomerular endothelial cell proteoglycans - regulation by TGF- $\beta$ 1. *Arch. Biochem. Biophys.*, 305 (1993) 370-377.
- 290 Komatsu, H., Yoshii, K., Ishimitsu, S., Okada, S. and Takahata, T.: Molecular mass determination of low-molecular-mass heparins. Application of wide collection angle measurements of light scattering using a high-performance gel permeation chromatographic system equipped with a low-angle laser light-scattering photometer. *J. Chromatogr.*, 644 (1993) 17-24.
- 291 Moody, S.F., Handman, E., McConville, M.J. and Bacic, A.: The structure of *Leishmania major* amastigote lipophosphoglycan. *J. Biol. Chem.*, 268 (1993) 18457-18466.
- 292 Pellerin, P. and Brillouet, J.M.: Study of red wine polysaccharides fractionated by ion-exchange chromatography. *Wein-Wiss.*, 47 (1992) 153-158; *C.A.*, 119 (1993) 138034w.
- 293 Reddy, G.P., Hayat, U., Bush, C.A. and Morris, J.G., Jr.: Capsular polysaccharide structure of a clinical isolate of *Vibrio vulnificus* strain BO 62316 determined by heteronuclear NMR spectroscopy and high-performance anion-exchange chromatography. *Anal. Biochem.*, 214 (1993) 106-115.



- 294 Rinaudo, M., Danhelka, J. and Milas, M.: A new approach to characterizing carboxymethylcelluloses by size-exclusion chromatography. *Carbohydr. Polym.*, 21 (1993) 1-5; C.A., 119 (1993) 183103d.
- 295 Sarazin, D., Francois, J., Roqueta, C. and Fleche, G.: Aqueous size-exclusion chromatography of hydroxyethyl amylopectin. *Carbohydr. Polym.*, 21 (1993) 29-38; C.A., 119 (1993) 183202k.
- 296 Silbert, J.E., Sugumaran, G. and Cogburn, J.N.: Sulphation of proteochondroitin and 4-methylumbelliferyl  $\beta$ -D-xyloside-chondroitin formed by mouse mastocytoma cells cultured in sulphate-deficient medium. *Biochem. J.*, 296 (1993) 119-126.
- 297 Sullivan, C.R., Corona, A. and Rollings, J.E.: Chromatographic technologies for macromolecular starch characterization. In: Alexander, R.J. and Zobel, H.F. (Editors), *Dev. Carbohydr. Chem.*, Am. Assoc. Cereal Chem., St. Paul, 1992, pp. 193-238; C.A., 119 (1993) 93771r - a review with 79 refs.
- 298 Thibert, L. and Lapiere, S.: Routine application of high-performance liquid chromatography for identification of mycobacteria. *J. Clin. Microbiol.*, 31 (1993) 1759-1763; C.A., 119 (1993) 134698t.
- 299 Tyler-Cross, R., Sobel, M., Marques, D., Soler, D.F. and Harris, R.B.: Heparin-von Willebrand factor binding as assessed by isothermal titration calorimetry and by affinity fractionation of heparins using synthetic peptides. *Arch. Biochem. Biophys.*, 306 (1993) 528-533.
- 300 Vijayagopal, P.: Enhanced synthesis and accumulation of proteoglycans in cholesterol-enriched arterial smooth muscle cells. *Biochem. J.*, 294 (1993) 603-611.
- 301 Yamada, T., Ogamo, A. and Nakagawa, Y.: A new chromatographic procedure for separation of typical  $\lambda$ -type molecular species from commercial  $\lambda$ -carrageenan. *Carbohydr. Polym.*, 20 (1993) 285-289; C.A., 119 (1993) 98367y.
- For additional information see C.A.:  
119 (1993) 155505t.
- See also 20, 257, 259, 313, 314, 317, 1396.
- 10c. *Glycoproteins and their constituents*
- 302 Bame, K.J.: Release of heparan sulfate glycosaminoglycans from proteoglycans in Chinese hamster ovary cells does not require proteolysis of the core protein. *J. Biol. Chem.*, 268 (1993) 19956-19964.
- 303 Burton-Wurster, N. and Lust, G.: Evidence for a glycosaminoglycan chain on a portion of articular cartilage fibronectins. *Arch. Biochem. Biophys.*, 306 (1993) 309-320.
- 304 Dam, T.K., Sarkar, M., Ghosal, J. and Choudhury, A.: Purification and partial characterization of a N-glycolylneuraminic acid specific lectin from the clam *Anadara granosa* (L.). *Biochem. Biophys. Res. Commun.*, 196 (1993) 422-429.
- 305 Eh, J., Seals, J.R., Murphy, C.I., van Halbeek, H. and Cummings, R.D.: Site-specific N-glycosylation and oligosaccharide structures of recombinant HIV-1 gp120 derived from a baculovirus expression system. *Biochemistry*, 32 (1993) 11087-11099.
- 306 Endo, T., Oda, O., Yamanaka, N., Maeda, K., Yoshida, M. and Kobata, A.: Alterations in the carbohydrate structures of an abnormal protein from sera of patients with rheumatoid arthritis. *Arch. Biochem. Biophys.*, 307 (1993) 119-125.
- 307 Kawasaki, N., Yokota, Y. and Kawasaki, T.: Differentiation of conglutination activity and sugar-binding activity of conglutinin after removal of NH<sub>2</sub>-terminal 54 amino acid residues by endogenous serine protease(s). *Arch. Biochem. Biophys.*, 305 (1993) 533-540.
- 308 Malette, B. and Bleau, G.: Biochemical characterization of hamster oviductin as a sulphated zona pellucida-binding glycoprotein. *Biochem. J.*, 295 (1993) 437-445.
- 309 Maynes, P.A., Gooley, A.A., Ferguson, M.A.J., Redmond, J.W. and Williams, K.L.: Post-translational modifications of the *Dicystoselium discoideum* glycoprotein PsA. Glycosylphosphatidylinositol membrane anchor and composition of O-linked oligosaccharides. *Eur. J. Biochem.*, 216 (1993) 729-737.
- 310 Mo, H., van Damme, E.J.M., Peumans, W.J. and Goldstein, I.J.: Purification and characterization of a mannose-specific lectin from shallot (*Allium ascalonicum*) bulbs. *Arch. Biochem. Biophys.*, 306 (1993) 431-438.
- 311 Nagpurkar, A., Hunt, D., Yang, C.-Y. and Mookerjee, S.: Degradation of rat C-reactive protein by macrophages. *Biochem. J.*, 295 (1993) 247-253.
- 312 Noguchi, S. and Nakano, M.: Structural characterization of the N-linked carbohydrate chains from mouse zona pellucida glycoproteins ZP2 and ZP3. *Biochim. Biophys. Acta*, 1158 (1993) 217-226.
- 313 Romero, P.A., Way, T. and Herscovics, A.: Lamp-1 does not acquire the large polylectosaminoglycans characteristic of F9 cells. *Biochem. J.*, 296 (1993) 253-257.
- 314 Saito, K., Komae, A., Kakuta, M., van Damme, E.J.M., Peumans, W.J., Goldstein, I.J. and Misaki, A.: The  $\alpha$ -mannosyl-binding lectin from leaves of the orchid twayblade (*Listera ovata*). Application to separation of  $\alpha$ -D-mannans from  $\alpha$ -D-glucans. *Eur. J. Biochem.*, 217 (1993) 677-681.
- 315 Schaumann, C., Oesch, F., Unger, K.K. and Wieser, R.J.: Analytical technique for studying the structure of glycoprotein N-glycans. *J. Chromatogr.*, 646 (1993) 227-234.
- 316 Schittny, J.C. and Schittny, C.M.: Role of the B1 short arm in laminin self-assembly. *Eur. J. Biochem.*, 216 (1993) 437-441.
- 317 Strang, A.-M., Allen, A.K., Holder, A.A. and van Halbeek, H.: The carbohydrate structures of *Trypanosoma brucei brucei* MITat 1.6 variant surface glycoprotein. A re-investigation of the C-terminal glycan. *Biochem. Biophys. Res. Commun.*, 196 (1993) 1430-1439.
- 318 Treuheit, M.J. and Halsall, H.B.: Restricted variation in the glycosylation of human  $\alpha_1$ -acid glycoprotein. *Chromatographia*, 37 (1993) 144-148.
- 319 Uhlin-Hansen, L. and Yanagishita, M.: Differential effect of brefeldin A on the biosynthesis of heparan sulfate and chondroitin/dermatan sulfate proteoglycans in rat ovarian granulosa cells in culture. *J. Biol. Chem.*, 268 (1993) 17370-17376.
- 320 Van Damme, E.J.M., Smeets, K., Torrekens, S., van Leuven, F. and Peumans, W.J.: The mannose-specific lectins from ransons (*Allium ursinum* L.) are encoded by three sets of genes. *Eur. J. Biochem.*, 217 (1993) 123-129.
- 321 Vargas-Albores, F., Hernández, J., Córdoba, F. and Zenteno, E.: Isolation of an immunosuppressive lectin from *Phaseolus vulgaris* L. cv Cacahuete using stroma. *Prepar. Biochem.*, 23 (1993) 473-483.
- 322 Wadsworth, J.D.F., Okuno, A. and Strong, P.N.: Assignment of laminin heavy chains using the lectin *Ricinus communis* agglutinin-1. *Biochem. J.*, 295 (1993) 537-541.

See also 250, 256, 643.

## 11. ORGANIC ACIDS AND LIPIDS

### 11a. Organic acids and simple esters

- 323 Alexander, J.N., IV and Quinn, C.J.: Organic acid analysis by ion chromatography-particle beam mass spectrometry. *J. Chromatogr.*, 647 (1993) 95-100.
- 324 Baer, A.N. and Green, F.A.: Fatty acid oxygenase activity of human hair roots. *J. Lipid Res.*, 34 (1993) 1505-1514.
- 325 Beyrich, T., Grünert, R. and Jira, T.: Vergleichende Untersuchungen zum katalytischen Einfluss von Cyclodextrinen auf Nitrophenylacetate und Nitrobenzoesäureester. *Pharmazie*, 48 (1993) 597-599.
- 326 Buckberry, L.D., Gallagher, S.D. and Shaw, P.N.: The development of a high performance liquid chromatographic assay method to determine pyruvate generated during the C-S lysis of cysteine conjugated xenobiotics. *J. Liq. Chromatogr.*, 16 (1993) 3073-3081.
- 327 Daret, D., Blin, P., Dorian, B., Rigaud, M. and Larrue, J.: Synthesis of monohydroxylated fatty acids from linoleic acid by rat aortic smooth muscle and tissues: influence on prostacyclin production. *J. Lipid Res.*, 34 (1993) 1473-1482.
- 328 Di Marzo, V., de Petrocellis, L., Gianfrani, C. and Cimino, G.: Biosynthesis, structure and biological activity of hydroxyeicosatetraenoic acids in *Hydra vulgaris*. *Biochem. J.*, 295 (1993) 23-29.
- 329 Dietz, E.A., Cortellucci, N.J. and Singley, K.F.: Determination of benzoic acid, chlorobenzoic acids and chlorendic acid in water. *J. Liq. Chromatogr.*, 16 (1993) 3331-3347.
- 330 Doadrio, A.L. and Clemente, B.: Microbore high-performance liquid chromatographic quantitative analysis of lysine acetyl salicylate and its degradation product salicylic acid. *An. R. Acad. Farm.*, 58 (1992) 259-268; *C.A.*, 119 (1993) 125335v.
- 331 Gravel, J., Falgout, J.P., Yergey, J., Trimble, L. and Riendeau, D.: Identification of 5-keto-(7E,9E,11Z,14Z)-eicosatetraenoic acid as a novel nonenzymatic rearrangement product of leukotriene A<sub>4</sub>. *Arch. Biochem. Biophys.*, 306 (1993) 469-475.
- 332 Harvey, B.A., Herrington, T.M. and Bee, R.: Transport studies of oil-soluble polymers. *ACS Symp. Ser.*, 520(Polymeric Delivery Systems) (1993) 220-231; *C.A.*, 119 (1993) 158543q.
- 333 Herrera, M.O., García, H.L., Mir, M.V. and Martínez, M.C.L.: Determination by high performance liquid chromatography of organic acids in Spanish rosé wines from the Alpujarra-Contraviesa region of Granada. *J. Liq. Chromatogr.*, 16 (1993) 3101-3112.
- 334 Husain, S., Narsimha, R., Alvi, S.N. and Rao, R.N.: Monitoring the levels of chloroacetic acids in the industrial environment by high performance liquid chromatography. *J. High Resolut. Chromatogr.*, 16 (1993) 381-383.
- 335 Ingalls, S.T., Kriaris, M.S., Xu, Y., DeWulf, D.W., Tserng, K.-Y. and Hoppel, C.L.: Method for isolation of non-esterified fatty acids and several other classes of plasma lipids by column chromatography on silica gel. *J. Chromatogr.*, 619 (1993) 9-19.
- 336 Kondo, J., Imaoka, T., Kawasaki, T., Nakanishi, A. and Kawahara, Y.: Fluorescence derivatization reagent for resolution of carboxylic acid enantiomers by high-performance liquid chromatography. *J. Chromatogr.*, 645 (1993) 75-81.
- 337 Lee, B.-L., New, A.-L., Kok, P.-W., Ong, H.-Y., Shi, C.-Y. and Ong, C.-N.: Urinary trans,trans-muconic acid determined by liquid chromatography: application in biological monitoring of benzene exposure. *Clin. Chem. (Washington)*, 39 (1993) 1788-1792.
- 338 Levi, V., Wehr, T., Talmadge, K. and Zhu, M.: Analysis of organic acids in wines by capillary electrophoresis and HPLC. *Int. Lab.*, 23, No. 8 (1993) P4-P7.
- 339 Levi, V., Wehr, T., Talmadge, K. and Zhu, M.: Analysis of organic acids in wines by capillary electrophoresis and HPLC. *Am. Lab. (Shelton)*, 25 (1993) 29-32; *C.A.*, 119 (1993) 115621w.
- 340 Lien, M.-H., Huang, B.-C. and Hsu, M.-C.: Determination of ascorbyl dipalmitate in cosmetic whitening powders by liquid chromatography. *J. Chromatogr.*, 645 (1993) 362-365.
- 341 Lladó, I., Palou, A. and Pons, A.: Combined enzymic and chromatographic techniques to determine specific radioactivity in free and triglyceride fatty acid plasma fractions. *J. Chromatogr.*, 619 (1993) 21-28.
- 342 Matsumoto, K., Takahashi, M., Takiyama, N., Misaki, H., Matsuo, N., Murano, S. and Yuki, H.: Enzyme reactor for urinary acylcarnitines assay by reversed-phase high-performance liquid chromatography. *Clin. Chim. Acta*, 216 (1993) 135-143.
- 343 McCaskill, D. and Croteau, R.: Procedures for the isolation and quantification of the intermediates of the mevalonic acid pathway. *Anal. Biochem.*, 215 (1993) 142-149.
- 344 Meuwly, P. and Métraux, J.-P.: *Ortho*-anisic acid as internal standard for the simultaneous quantitation of salicylic acid and its putative biosynthetic precursors in cucumber leaves. *Anal. Biochem.*, 214 (1993) 500-505.
- 345 Nagata, Y., Yamamoto, Y., Yamashita, S. and Niki, E.: Fluorometric analysis of free fatty acid hydroperoxides and their alcohols, and its application to blood plasma. *Int. Congr. Ser. - Excerpta Med.*, 998(Oxygen Radicals) (1992) 315-318; *C.A.*, 119 (1993) 112832e.
- 346 Nakashima, K., Taguchi, Y., Kuroda, N., Akiyama, S. and Duan, G.: 2-(4-Hydrazinocarbonylphenyl)-4,5-diphenylimidazole as a versatile fluorescent derivatization reagent for the high-performance liquid chromatographic analysis of free fatty acids. *J. Chromatogr.*, 619 (1993) 1-8.
- 347 Nikolova-Damyanova, B., Christie, W.W. and Herslöf, B.G.: High-performance liquid chromatography of fatty acid derivatives in the combined silver ion and reversed-phase modes. *J. Chromatogr.*, 653 (1993) 15-23.
- 348 Ohshima, T., Miyamoto, K.-I. and Sakai, R.: Simultaneous separation and sensitive measurement of free fatty acids in ancient pottery by high performance liquid chromatography. *J. Liq. Chromatogr.*, 16 (1993) 3217-3227.
- 349 Oliw, E.H., Hörnsten, L., Sprecher, H. and Hamberg, M.: Oxygenation of 5,8,11-eicosatrienoic acid by prostaglandin endoperoxide synthase and by cytochrome P450 monooxygenase: structure and mechanism of formation of major metabolites. *Arch. Biochem. Biophys.*, 305 (1993) 288-297.

- 350 Petrarulo, M., Cerelli, E., Marangella, M., Cosseddu, D., Vitale, C. and Linari, F.: (Quantitative determination of calcium oxalate in bone. Patients on regular dialysis treatment for primary hyperoxaluria and other nephropathies). *G. Ital. Chim. Clin.*, 17 (1992) 95-100; *C.A.*, 119 (1993) 134942t.
- 351 Poorthuis, B.J.H.M., Jille-Vlckova, T. and Onkenhout, W.: Determination of acylcarnitines in urine of patients with inborn errors of metabolism using high-performance liquid chromatography after derivatization with 4'-bromophenacylbromide. *Clin. Chim. Acta*, 216 (1993) 53-61.
- 352 Reynaud, D., Thickitt, C.P. and Pace-Asciak, C.R.: Facile preparation and structural determination of monohydroxy derivatives of docosahexaenoic acid (HDOHE) by  $\alpha$ -tocopherol-directed autoxidation. *Anal. Biochem.*, 214 (1993) 165-170.
- 353 Seki, T., Yamaji, K., Yanagihara, Y. and Noguchi, K.: Determination of vanillylmandelic acid and homovanillic acid in human urine by coupled column high-performance liquid chromatography with fluorimetric detection. *Int. Congr. Ser. - Excerpta Med.*, 991 (Progress in Clinical Biochemistry) (1992) 321-323; *C.A.*, 119 (1993) 112590z.
- 354 Stein, J., Milovic, V., Zeuzem, S. and Caspary, W.F.: Fluorometric high-performance liquid chromatography of free fatty acids using panacyl bromide. *J. Liq. Chromatogr.*, 16 (1993) 2915-2922.
- 355 Swadesh, J.K., Stewart, C.W., Jr. and Uden, P.C.: Comparison of liquid chromatographic methods for analysis of homologous alkyl esters of biphenyl-4,4'-dicarboxylic acid. *Analyst (London)*, 118 (1993) 1123-1125.
- 356 Villalobos C., A.E. and Calzada E., C.: (Determination of phthalates by HPLC). *Ing. Cienc. Quim.*, 14 (1992) 28-29; *C.A.*, 119 (1993) 140291w.
- For additional information see *C.A.*:  
119 (1993) 155154c, 158489b.
- See also 24, 59, 103, 115, 222, 358, 376, 485, 517, 559, 692, 749, 754, 992, 1349.
- 11b. *Prostaglandins*
- 357 Kråmer, H.J., Stevens, J. and Seeger, W.: Analysis of 2- and 3-series prostanoids by post-HPLC ELISA. *Anal. Biochem.*, 214 (1993) 535-543.
- 358 Lopez, S., Vila, L., Breviario, F. and de Castellarnau, C.: Interleukin-1 increases 15-hydroxyicosatetraenoic acid formation in cultured human endothelial cells. *Biochim. Biophys. Acta*, 1170 (1993) 17-24.
- 359 Tagari, P., Becker, A., Brideau, C., Frenette, R., Sadl, V., Thomas, E., Vickers, P. and Ford-Hutchinson, A.: Leukotriene generation and metabolism in dogs: inhibition of biosynthesis by MK-0591. *J. Pharmacol. Exp. Ther.*, 265 (1993) 416-425.
- 11c. *Lipids and their constituents*
- 360 Aitzetmüller, K.: HPLC von Triglyceriden ("Fingerprint-Methode"). *Gemeinschaftsarbeiten der DGF*, 136. Mitteilung: Deutsche Einheitsmethoden zur Untersuchung von Fetten, Fettprodukte, Tensiden und verwandten Stoffen, 103. Mitt.: Analyse von Fetten XXX. *Fat Sci. Technol.*, 95 (1993) 361-366.
- 361 Berggren, M.I., Gallegos, A., Dressler, L.A., Modest, E.J. and Powis, G.: Inhibition of the signalling enzyme phosphatidylinositol-3-kinase by antitumor ether lipid analogues. *Cancer Res.*, 53 (1993) 4297-4302.
- 362 Brühl, L., Schulte, E. and Thier, H.-P.: Fraktionierung der Triglyceride von Muttermilch durch HPLC an einer Silberionensäule und an RP-18-Material mit Lichtstreuendetektor. *Fat Sci. Technol.*, 95 (1993) 370-376.
- 363 Burkow, I.C. and Henderson, J.R.: Recent advances in lipid analysis by chromatography. In: Lambertsen, G. (Editor), *Proc.-Scand. Symp. Lipids, 16th*, Lipidforum, Bergen, 1991, pp. 60-69; *C.A.*, 119 (1993) 176804e - a review with 45 refs.
- 364 Camacho, M.L.: (HPLC determination of triglycerides in edible vegetable oils). *Aliment., Equipos Tecnol.*, 12 (1993) 45-49; *C.A.*, 119 (1993) 179456d - a review with 3 refs.
- 365 Conforti, F.D., Harris, C.H. and Rinehart, J.T.: High-performance liquid chromatographic analysis of wheat flour lipids using an evaporative light scattering detector. *J. Chromatogr.*, 645 (1993) 83-88.
- 366 Couto, A.S., de Lederkremer, R.M., Colli, W. and Alves, M.J.M.: The glycosylphosphatidylinositol anchor of the trypanostigote-specific Tc-85 glycoprotein from *Trypanosoma cruzi*. Metabolic-labeling and structural studies. *Eur. J. Biochem.*, 217 (1993) 597-602.
- 367 Davila, A.M., Marchal, R., Monin, N. and Vandecasteele, J.P.: Identification and determination of individual sophorolipids in fermentation products by gradient elution high-performance liquid chromatography with evaporative light-scattering detection. *J. Chromatogr.*, 648 (1993) 139-149.
- 368 Deffense, E.: (Novel HPLC procedure for the separation of monounsaturated triglyceride 1-2 and 1-3 positional isomers from vegetable oil. I. Description of the method). *Ref. Fr. Corps Gras*, 40 (1993) 33-39; *C.A.*, 119 (1993) 137700y.
- 369 Galella, G., Marangoni, F., Risé, P., Colombo, C., Galli, G. and Galli, C.: *n*-6 And *n*-3 fatty acid accumulation in thp-1 cell phospholipids. *Biochim. Biophys. Acta*, 1169 (1993) 280-290.
- 370 He, P., Hu, J. and Macher, B.A.: Glycosphingolipids of rabbit, sheep, and pig thymus. *Arch. Biochem. Biophys.*, 305 (1993) 350-361.
- 371 Hidari, K.I.-P.J., Irie, F., Suzuki, M., Kon, K., Ando, S. and Hirabayashi, Y.: A novel ganglioside with a free amino group in bovine brain. *Biochem. J.*, 296 (1993) 259-263.
- 372 Hierro, M.T.G., Najera, A.I. and Santa-Maria, G.: Analysis of triglycerides by reversed-phase HPLC with gradient elution using a light-scattering detector. *Rev. Esp. Cienc. Tecnol. Aliment.*, 32 (1992) 635-651; *C.A.*, 119 (1993) 137682u.
- 373 Hopia, A.I. and Ollilainen, V.-M.: Comparison of the evaporative light scattering detector (ELSD) and refractive index detector (RID) in lipid analysis. *J. Liq. Chromatogr.*, 16 (1993) 2469-2482.
- 374 Hopia, A.I., Lampi, A.M., Piironen, V.I., Hyvönen, L.E.T. and Koivistoinen, P.E.: Application of high-performance size-exclusion chromatography to study the autoxidation of unsaturated triacylglycerols. *J. Am. Oil Chem. Soc.*, 70 (1993) 779-784.
- 375 Hudiyono, S., Adenier, H. and Chaveron, H.: (Determination of the triglyceride composition of a hydrogenated palm oil fraction). *Rev. Fr. Corps Gra*, 40 (1993) 131-141; *C.A.*, 119 (1993) 115636e.

- 376 Kondo, Y., Kawai, Y., Hayashi, T., Ohnishi, M., Miyazawa, T., Itoh, S. and Mizutani, J.: Lipoygenase in soybean seedlings catalyzes the oxygenation of phospholipid and such activity changes after treatment with fungal elicitor. *Biochim. Biophys. Acta*, 1170 (1993) 301-306.
- 377 Leon, M.: (Determination of triglycerides in edible vegetable oils by high-performance liquid chromatography). *Tec. Lab.*, 15 (1993) 277-281; C.A., 119 (1993) 137667t - a review with 3 refs.
- 378 Li, R., Gage, D. and Ladisch, S.: Biosynthesis and shedding of murine lymphoma gangliosides. *Biochim. Biophys. Acta*, 1170 (1993) 283-290.
- 379 Liepkalns, V.A., Myher, J.J., Kuksis, A., Leli, U., Freysz, N. and Hauser, G.: Molecular species of glycerophospholipids and diacylglycerols of cultured SH-S454 human neuroblastoma cells. *Biochem. Cell Biol.*, 71 (1993) 141-149.
- 380 Lynch, S.M. and Frei, B.: Mechanism of copper- and iron-dependent oxidative modification of human low density lipoprotein. *J. Lipid Res.*, 34 (1993) 1745-1753.
- 381 Marai, L., Kuksis, A., Myher, J.J. and Itabashi, Y.: Liquid chromatography chloride attachment negative chemical ionization mass spectrometry of diacylglycerol dinitrophenylurethanes. *Biol. Mass Spectrom.*, 21 (1992) 541-547; C.A., 119 (1993) 112816c.
- 382 Moats, R.A., Lien, Y.-H.H., Filippi, D. and Ross, B.D.: Decrease in cerebral inositols in rats and humans. *Biochem. J.*, 295 (1993) 15-18.
- 383 Murphy, E.J., Stephens, R., Jurkowitz-Alexander, M. and Horrocks, L.A.: Acidic hydrolysis of plasmalogens followed by high-performance liquid chromatography. *Lipids*, 28 (1993) 565-568; C.A., 119 (1993) 112607k.
- 384 Nilsson, W.B. and Hudson, J.K.: Solubility of simple and mixed triacylglycerols in supercritical CO<sub>2</sub>. *J. Am. Oil Chem. Soc.*, 70 (1993) 749-754.
- 385 Norgauer, J., Dobos, G., Kownatzki, E., Dahinden, C., Burger, R., Kupper, R. and Gierschik, P.: Complement fragment C3a stimulates Ca<sup>2+</sup> influx in neutrophils via a pertussis-toxin-sensitive G protein. *Eur. J. Biochem.*, 217 (1993) 289-294.
- 386 Paradies, G., Ruggiero, F.M., Dinoi, P., Petrosillo, G. and Quagliariello, E.: Decreased cytochrome oxidase activity and changes in phospholipids in heart mitochondria from hypothyroid rats. *Arch. Biochem. Biophys.*, 307 (1993) 91-95.
- 387 Petersson, B., Podlaha, O. and Jirskog-Hed, B.: Triacylglycerol analysis of partially hydrogenated fats using high-performance liquid chromatography. *J. Chromatogr.*, 653 (1993) 25-35.
- 388 Ren, S., Ariga, T., Scarsdale, J.N., Zhang, Y., Slominski, A., Livingston, P.O., Ritter, G., Kushi, Y. and Yu, R.K.: Characterization of a hamster melanoma-associated ganglioside antigen as 7-O-acetylated disialoganglioside GD3. *J. Lipid Res.*, 34 (1993) 1565-1572.
- 389 Robinson, D.R., Xu, L.L., Knoell, C.T., Tateno, S. and Olesiak, W.: Modification of spleen phospholipid fatty acid composition by dietary fish oil and by n-fatty acid ethyl esters. *J. Lipid Res.*, 34 (1993) 1423-1434.
- 390 Salivaras, M. and McCurdy, A.R.: Adulteration of olive oil: detection through reversed-phase high performance liquid chromatography. *Dev. Food Sci.*, 32(Food Flavors, Ingredients and Composition) (1993) 279-300; C.A., 119 (1993) 179499v.
- 391 Sattler, W. and Stocker, R.: Greater selective uptake by Hep G2 cells of high-density lipoprotein cholesteryl ester hydroperoxides than of unoxidized cholesteryl esters. *Biochem. J.*, 294 (1993) 771-778.
- 392 Schneider, P., Rosat, J.-P., Ransijn, A., Ferguson, M.A.J. and McConville, M.J.: Characterization of glycoinositol phospholipids in the amastigote stage of the protozoan parasite *Leishmania major*. *Biochem. J.*, 295 (1993) 555-564.
- 393 Silvestro, L., da Col, R., Scappaticci, E., Libertucci, D., Biancone, L. and Camussi, G.: Development of a high-performance liquid chromatographic-mass spectrometric technique, with an ion-spray interface, for the determination of platelet-activating factor (PAF) and lyso-PAF in biological samples. *J. Chromatogr.*, 647 (1993) 261-269.
- 394 Tabernero, A., Bolanos, J.P. and Medina, J.M.: Lipogenesis from lactate in rat neurons and astrocytes in primary culture. *Biochem. J.*, 294 (1993) 635-638.
- 395 Tinsley, P.W. and Maerker, G.: Isolation and identification of palmitoylphosphocholinepropanediol from  $\gamma$ -irradiated dipalmitoylphosphatidylcholine. *J. Am. Oil Chem. Soc.*, 70 (1993) 815-816.

For additional information see C.A.:  
119 (1993) 90417u, 90419w, 158520e.

See also 115, 222, 341, 402, 754.

#### 11d. Lipoproteins and their constituents

- 396 Connelly, P.W., Maguire, G.F., Vezina, C., Hegele, R.A. and Little, J.A.: Identification of disulfide-linked apolipoprotein species in human lipoproteins. *J. Lipid Res.*, 34 (1993) 1717-1727.
- 397 Lehmann, R., Bhargava, A.S. and Günzel, P.: Serum lipoprotein pattern in rats, dogs and monkeys, including method comparison and influence of menstrual cycle in monkeys. *Eur. J. Clin. Chem. Clin. Biochem.*, 31 (1993) 633-637.
- 398 März, W., Siekmeier, R., Scharnagl, H., Seiffert, U.B. and Gross, W.: Fast lipoprotein chromatography: new method of analysis for plasma lipoproteins. *Clin. Chem. (Washington)*, 39 (1993) 2276-2281.
- 399 Razafindralambo, H., Paquot, M., Hbid, C., Jacques, P., Destain, J. and Thonart, P.: Purification of antifungal lipopeptides by reversed-phase high-performance liquid chromatography. *J. Chromatogr.*, 639 (1993) 81-85.
- 400 Tschantz, J.-C. and Sunahara, G.I.: Microaffinity chromatographic separation and characterization of lipoprotein fractions in rat and Mongolian gerbil serum. *Clin. Chem. (Washington)*, 39 (1993) 1861-1867.
- 401 Yang, C.-y., Gu, Z.-W., Valentinova, N., Pownall, H.J., Lee, B., Yang, M., Xie, Y.-h., Guyton, J.R., Vlasik, T.N., Fruchart, J.-C. and Gotto, A.M., Jr.: Human very low density lipoprotein structure: interaction of the C apolipoproteins with apolipoprotein B-100. *J. Lipid Res.*, 34 (1993) 1311-1321.

For additional information see C.A.:  
119 (1993) 176884f.

See also 706.

## 12. ORGANIC PEROXIDES

- 402 Aikens, J. and Dix, T.A.: Hydrodioxyl (perhydroxyl), peroxy and hydroxyl radical-initiated lipid peroxidation of large unilamellar vesicles (liposomes): comparative and mechanistic studies. *Arch. Biochem. Biophys.*, 305 (1993) 516-525.

See also 18, 345, 380, 391.

## 13. STEROIDS

## 13a. General techniques

- 403 Santos-Montes, A., Gasco-Lopez, A.I. and Izquierdo-Hornillos, R.: Optimization of the high-performance liquid chromatographic separation of a mixture of natural and synthetic corticosteroids. *J. Chromatogr.*, 620 (1993) 15-23.
- 404 Teffera, Y., Abramson, F.P., McLean, M. and Vestal, M.: Development of an isotope-selective high-performance liquid chromatography detector using chemical-reaction-interface mass spectrometry: application to deuterated cortisol metabolites in urine. *J. Chromatogr.*, 620 (1993) 89-96.

For additional information see C.A.:  
119 (1993) 174283k.

## 13b. Pregnane and androstane derivatives

- 405 Amin, M., Harrington, K. and von Wandruszka, R.: Determination of steroids in urine by sensitized terbium fluorescence. *Anal. Chem.*, 65 (1993) 2346-2351.
- 406 Ciranni Signoretti, E., Valvo, L., Savella, A.L. and Cavina, G.: Purity evaluation of 6 $\alpha$ -methylprednisolone acetate by HPLC. *J. Pharm. Biomed. Anal.*, 11 (1993) 587-593.
- 407 Ichikawa-Haraguchi, M., Sumizawa, T., Yoshimura, A., Furukawa, T., Hiramoto, S., Sugita, M. and Akiyama, S.-i.: Progesterone and its metabolites: the potent inhibitors of the transporting activity of P-glycoprotein in the adrenal gland. *Biochim. Biophys. Acta*, 1158 (1993) 201-208.
- 408 Laizure, S.C., Franklin, L.K., Kaiser, D.G., Williams, C.L., Stevens, R.C., Sanders, P.L. and Miller, M.: Disposition of tirilazad (U74006F), a 21-aminosteroid, in the plasma, heart, brain, and liver of the rat. *Drug Metab. Disp.*, 21 (1993) 951-954.
- 409 Li, Y.-M., Chen, L.-R. and Qu, Y.: Use of micellar mobile phases and an HPLC column switching system for direct injection determination of urinary free cortisol. *J. Liq. Chromatogr.*, 16 (1993) 2583-2594.
- 410 Samaan, G.J., Porquet, D., Demelier, J.F. and Biou, D.: Determination of cortisol and associated glucocorticoids in serum and urine by an automated liquid chromatographic assay. *Clin. Biochem.*, 26 (1993) 153-158; C.A., 119 (1993) 109150a.
- 411 Schmidt, R., Klaus, U. and Michl, K.: (A new method for easy and rapid determination of free cortisol in serum by HPLC). *Labor-Med.*, 16 (1993) 151-156; C.A., 119 (1993) 109160d.

For additional information see C.A.:  
119 (1993) 103487u.

See also 403, 404, 413.

## 13c. Estrogens

- 412 Fernandez, N., García, J.J., Diez, M.J., Terán, M.T. and Sierra, M.: Rapid high-performance liquid chromatographic assay of ethynloestradiol in rabbit plasma. *J. Chromatogr.*, 619 (1993) 143-147.
- 413 Ghosh, G., Roy, S.K. and Majumdar, M.K.: Separation and determination of ethinyl estradiol and norgestrel from oral contraceptive tablets by RP-HPLC. *Indian Drugs*, 30 (1993) 301-304; C.A., 119 (1993) 167857x.
- 414 Shimada, K., Kaji, H. and Kuwabara, K.: Determination of estril monoglucuronide derivatives in the Koenigs-Knorr reaction by high-performance liquid chromatography. *J. Chromatogr. Sci.*, 31 (1993) 363-365.

See also 157.

## 13d. Sterols

- 415 Chen, W.X., Li, P.Y., Wang, S., Dong, J. and Li, J.Z.: Serum cholesterol determined by liquid chromatography with 6-chlorostigmasterol as internal standard. *Clin. Chem. (Washington)*, 39 (1993) 1602-1607.
- 416 Hasan, S.Q. and Kushwaha, R.S.: Differences in 27-hydroxycholesterol concentrations in plasma and liver of baboons with high and low responses to dietary cholesterol and fat. *Biochim. Biophys. Acta*, 1182 (1993) 299-302.
- 417 Hwang, K.T. and Maerker, G.: Determination of 6-ketocholestanol in unirradiated and irradiated chicken meats. *J. Am. Oil Chem. Soc.*, 70 (1993) 789-792.
- 418 Kamido, H., Kuksis, A., Marai, L. and Myher, J.J.: Identification of core aldehydes among *in vitro* peroxidation products of cholesterol esters. *Lipids*, 28 (1993) 331-336; C.A., 119 (1993) 160628q.
- 419 McCluskey, S. and Devery, R.: Validation of chromatographic analysis of cholesterol oxides in dried foods. *Trends Food Sci. Technol.*, 4 (1993) 175-178; C.A., 119 (1993) 179428w.
- 420 Murui, T. and Wanaka K.: Measurement of sterylglucosides by high performance liquid chromatography with 1-anthroylnitrile derivatives. *Biosci. Biotechnol., Biochem.*, 57 (1993) 614-617; C.A., 119 (1993) 93830j.
- 421 Poorman, J.A., Buck, R.A., Smith, S.A., Overturf, M.L. and Loose-Mitchell, D.S.: Bile acid excretion and cholesterol 7 $\alpha$ -hydroxylase expression in hypercholesterolemia-resistant rabbits. *J. Lipid Res.*, 34 (1993) 1675-1685.
- 422 Smith, L.L.: Analysis of oxysterols by liquid chromatography. *J. Liq. Chromatogr.*, 16 (1993) 1731-1747 - a review with 88 refs.
- 423 Swaminathan, S., Wilson, W.K., Pinkerton, F.D., Gerst, N., Ramser, M. and Schroepfer, G.J., Jr.: Inhibitors of sterol synthesis. Chemical synthesis and properties of 3 $\beta$ -hydroxy-25,26,26,26,27,27,27-heptafluoro-5 $\alpha$ -cholest-8(14)-en-15-one and 25,26,26,26,27,27,27-heptafluorocholesterol and their effects on 3-hydroxy-3-methylglutaryl coenzyme A reductase activity in cultured mammalian cells. *J. Lipid Res.*, 34 (1993) 1805-1823.

See also 380, 391, 394.

13e. *Bile acids and alcohols*

- 424 Evans, J.E., Ghosh, A., Evans, B.A. and Natowicz, M.R.: Screening techniques for the detection of inborn errors of bile acid metabolism by direct injection and micro-high performance liquid chromatography-continuous flow-fast atom bombardment mass spectrometry. *Biol. Mass Spectrom.*, 22 (1993) 331-337; *C.A.*, 119 (1993) 177075e.
- 425 Niwa, T., Fujita, K., Goto, J. and Nambara, T.: Separation and characterization of ursodeoxycholate 7-N-acetylglucosaminides in human urine by high-performance liquid chromatography with fluorescence detection. *J. Liq. Chromatogr.*, 16 (1993) 2531-2544.
- 426 Roda, A., Gatti, R., Cavrini, V., Cerre, C. and Simoni, P.: HPLC study of the impurities present in different ursodeoxycholic acid preparations: comparative evaluation of four detectors. *J. Pharm. Biomed. Anal.*, 11 (1993) 751-760.

For additional information see C.A.:  
119 (1993) 160648w.

13f. *Ecdysones and other insect steroid hormones*

- 427 Evershed, R.P., Prescott, M.C., Kabbouh, M. and Rees, H.H.: High-performance liquid chromatography/mass spectrometry with thermospray ionization of free ecdysteroids. *Rapid Commun. Mass Spectrom.*, 7 (1993) 477-481; *C.A.*, 119 (1993) 134972c.

For additional information see C.A.:  
119 (1993) 112581x.

13g. *Other steroids*

- 428 Maier, M.S., Roccatagliata, A. and Seldes, A.M.: Two novel steroidal glycoside sulfates from the starfish *Cosmasterias lurida*. *J. Natural Prod.*, 56 (1993) 939-942.
- 429 Weiss, A.C., Jr., Elliger, C.A., Haddon, W.F. and Benson, M.: Insect inhibitory steroidal saccharide esters from *Physalis peruviana*. *J. Natural Prod.*, 56 (1993) 1365-1372.

For additional information see C.A.:  
119 (1993) 146716k.

## 14. STEROID GLYCOSIDES AND SAPONINS

- 430 Crespin, F., Ollivier, E., Elias, R., Maillard, C. and Balansard, G.: High-performance liquid chromatographic determination of the major saponin from *Opilia celtdifolia* Guill. Perr. *J. Chromatogr.*, 644 (1993) 404-406.
- 431 Iorizzi, M., Minale, L., Riccio, R. and Yasumoto, T.: Starfish saponins, part 51. Steroidal oligoglycosides from the starfish *Distolasterias nipon*. *J. Natural Prod.*, 56 (1993) 1786-1798.
- 432 Kim, B., Lee, M.Y., Cho, K.H., Park, J.H. and Park, M.K.: Analysis of ginseng saponins by HPLC with photoreduction fluorescence detection. *Arch. Pharmaceut. Res.*, 15 (1992) 328-332; *C.A.*, 119 (1993) 188698m.

- 433 Li, H.-Y., Koike, K., Ohmoto, T. and Ikeda, K.: Dianchinosides A and B, two new saponins from *Dianthus chinensis*. *J. Natural Prod.*, 56 (1993) 1065-1070.
- 434 Maillard, M.P. and Hostettmann, K.: Determination of saponins in crude plant extracts by liquid chromatography-thermospray mass spectrometry. *J. Chromatogr.*, 647 (1993) 137-146.

See also 1351, 1356.

## 15. TERPENES AND OTHER VOLATILE AROMATIC COMPOUNDS

15a. *Terpenes*

- 435 Achenbach, H., Waibel, R., Heftner-Bübl, U. and Constenla, M.A.: Constituents of *Fevillea cordifolia*: new corcucurbitacin and cucurbitacin glycosides. *J. Natural Prod.*, 56 (1993) 1506-1519.
- 436 Blay, G., Cardona, L., Garcia, B. and Pedro, J.R.: A short synthesis of (+)-colartin and (+)-arbusculin A from (-)-santonin. *J. Natural Prod.*, 56 (1993) 1723-1727.
- 437 Burnell, R.H., Coté, C. and Théberge, N.: Approaches to the synthesis of aromatic diterpenes oxygenated in the A ring. Synthesis of margocin. *J. Natural Prod.*, 56 (1993) 1459-1464.
- 438 Burnell, R.H., Dumont, N. and Théberge, N.: Synthesis of O-methyl nimbinone. *J. Natural Prod.*, 56 (1993) 1930-1936.
- 439 Cabral, J.A., McChesney, J.D. and Milhous, W.K.: A new anti-malarial quassinoid from *Simaba guianensis*. *J. Natural Prod.*, 56 (1993) 1954-1961.
- 440 Hufford, C.D., Jia, Y., Croom, E.M., Jr., Muhammed, I., Okunade, A.L., Clark, A.M. and Rogers, R.D.: Antimicrobial compounds from *Petalostemum purpureum*. *J. Natural Prod.*, 56 (1993) 1878-1889.
- 441 Inoue, H., Korenaga, T., Sagami, H., Koyama, T., Sugiyama, H. and Ogura, K.: Formation of farnesal and 3-hydroxy-2,3-dihydrofarnesal from farnesol by protoplasts of *Botryococcus braunii*. *Biochem. Biophys. Res. Commun.*, 196 (1993) 1401-1405.
- 442 Keller, R.K. and Vilsaint, F.: Regulation of isoprenoid metabolism in rat liver: near constant chain lengths of dolichyl phosphate and ubiquinone are maintained during greatly altered rates of cholesterologenesis. *Biochim. Biophys. Acta*, 1170 (1993) 204-210.
- 443 Lou, H., Li, X., Onda, M., Konda, Y., Machida, T., Toda, Y. and Harigaya, Y.: Further isolation of glycosides from *Cynanchum hancockianum*. *J. Natural Prod.*, 56 (1993) 1437-1443.
- 444 Ngassapa, O., Soejarto, D.D., Pezzuto, J.M., Farnsworth, N.R. and Che, C.-J.: Further lupane lactones from *Kokoona ochracea*. *J. Natural Prod.*, 56 (1993) 1676-1681.
- 445 Sundaram, K.M.S. and Curry, J.: High performance liquid chromatographic determination of azadirachtin in conifer and deciduous foliage, forest soils, leaf litter and stream water. *J. Liq. Chromatogr.*, 16 (1993) 3275-3290.
- 446 Zukowski, J. and Pawlowska, M.: Separation of monoterpene hydrocarbon enantiomers via complexation with  $\alpha$ -cyclodextrin in RP-HPLC. *J. High Resolut. Chromatogr.*, 16 (1993) 505-507.

See also 343, 975, 1318, 1358.

## 16. NITRO AND NITROSO COMPOUNDS

- 447 Baj, S., David, M. and Ambrozek, M.: High-performance liquid chromatographic determination of dynamites. *Chem. Anal. (Warsaw)*, 38 (1993) 323-328; *C.A.*, 119 (1993) 184097s.
- 448 Emmrich, M., Kaiser, M., Rüden, H. and Sollinger, S.: Determination of RDX, 2,4,6-trinitrotoluene and other nitroaromatic compounds by high-performance liquid chromatography with photodiode-array detection. *J. Chromatogr.*, 645 (1993) 89-94.
- 449 Hayes-Majstorovic, J., Guernet-Nivaud, E., Guernet, M. and Viel, C.: Mass spectrometry and liquid chromatography-mass spectrometry of  $\alpha$ -amino  $\beta$ -nitrostilbenes. *Anal. Lett.*, 26 (1993) 2239-2249.
- 450 Levsen, K., Musmann, P., Berger-Preiss, E., Preiss, A., Volmer, D. and Wuensch, G.: Analysis of nitroaromatic compounds and amines in ammunition wastewater and in aqueous samples from former ammunition plants and other military sites. *Acta Hydrochim Hydrobiol.*, 21 (1993) 153-166; *C.A.*, 119 (1993) 146199u.
- 458 Lin, Q.S., Su, W., Run, Q.Y. and Liu, Q.: Simultaneous determination of six monoamines in tissues and its application. *Int. Congr. Ser.-Excerpta Med.*, 991 (Progress in Clinical Biochemistry) (1992) 161-162; *C.A.*, 119 (1993) 86327x.
- 459 Lindahl, R., Levin, J.-O. and Andersson, K.: Determination of volatile amines in air by diffusive sampling, thiourea formation and high-performance liquid chromatography. *J. Chromatogr.*, 643 (1993) 35-41.
- 460 Pilipenko, A.T., Kurbansakhetov, Kh.K., Zul'figarov, O.S. and Orazmuradov, A.O.: (Ultrafiltration concentration of primary aromatic amines as diazo amino compounds and their determination by high-performance liquid chromatography in the presence of monohydric phenols). *Zavod. Lab.*, 59 (1993) 9-11; *C.A.*, 119 (1993) 194760s.
- 461 Uzunov, D. and Stoev, G.: Some aspects of the enantio-recognition of derivatized primary amines on a Pirkle-type chiral stationary phase utilizing tocainide and mexiletine as model compounds. *J. Chromatogr.*, 645 (1993) 233-239.

For additional information see *C.A.*:

119 (1993) 184091k.

See also 325, 505.

## 17. AMINES, AMIDES AND RELATED NITROGEN COMPOUNDS

## 17a. Amines and polyamines

- 451 Adams, P.N., Monkman, A.P. and Apperley, D.C.: Comparison of the molecular weights of polyaniline samples obtained from gel permeation chromatography and solid-state nitrogen-15 nuclear magnetic resonance spectroscopy. *Synth. Met.*, 55 (1993) 725-730; *C.A.*, 119 (1993) 161214p.
- 452 Coquart, V. and Hennion, M.-C.: Trace-level monitoring of chloroanilines in environmental waters using on-line trace-enrichment and liquid chromatography with UV and electrochemical detection. *Chromatographia*, 37 (1993) 392-398.
- 453 Fischer, L., Mueller, R., Mosbach, K. and Ekberg, B.: Method for separating enantiomers of aryloxypropranolamine derivatives. *PT Int. Appl. WO 93 09,075 (Cl. C07B57/00)*, 13 May 1993, SE Appl. 91/3,234, 04 Nov. 1991; 19 p.; *C.A.*, 119 (1993) 108196h.
- 454 Forgács, E.: Retention behavior of some ring-substituted aniline derivatives on polyethylene-coated and octadecylsilica columns. *J. Liq. Chromatogr.*, 16 (1993) 2483-2500.
- 455 Hiramatsu, K., Sugimoto, M., Kinoshita, K., Kamei, S. and Iwasaki, K.: Determination of free and acetyl polyamines excreted in urine of patients with urogenital cancer by the improved HPLC method. *Int. Congr. Ser.-Excerpta Med.*, 991 (Progress in Clinical Biochemistry) (1992) 957-959; *C.A.*, 119 (1993) 112591a.
- 456 Kamei, S., Hiramatsu, K., Sugimoto, M., Kinoshita, K. and Iwasaki, K.: Determination of polyamines by HPLC with an enzyme reactor. *Int. Congr. Ser.-Excerpta Med.*, 991 (Progress in Clinical Biochemistry) (1992) 117-121; *C.A.*, 119 (1993) 134678m.
- 457 Kotzabasis, K., Christakis-Hampsas, M.D. and Roubelakis-Angelakis, K.A.: A narrow-bore HPLC method for the identification and quantitation of free, conjugated, and bound polyamines. *Anal. Biochem.*, 214 (1993) 484-489.
- 462 Bergquist, J., Bergquist, S., Axelsson, R. and Ekman, R.: Demonstration of immunoglobulin G with affinity for dopamine in cerebrospinal fluid from psychotic patients. *Clin. Chim. Acta*, 217 (1993) 129-142.
- 463 Cheng, M.H., Lipsey, A.I., Lee, J. and Gamache, P.H.: Automated analysis of urinary VMA, HVA, and 5-HIAA by gradient HPLC using an array of eight coulometric electrochemical detectors. *Lab. Rob. Autom.*, 4 (1992) 297-303; *C.A.*, 119 (1993) 176890e.
- 464 Gamache, P.H., Kingery, M.L. and Acworth, I.N.: Urinary metanephrine and normetanephrine determined without extraction by using liquid chromatography and coulometric array detection. *Clin. Chem. (Washington)*, 39 (1993) 1825-1830.
- 465 Kiba, N., Itoi, M. and Furusawa, M.: Post-column immobilized tyrosinase reactor for determination of L-3,4-dihydroxyphenylalanine and L-tyrosine by high-performance liquid chromatography with fluorescence detection. *J. Chromatogr.*, 648 (1993) 481-484.
- 466 Kondo, N. and Hibi, K.: (Fully automatic urinary catecholamine analyzer). *Kuromatogurafi*, 13 (1992) 253-254; *C.A.*, 119 (1993) 112571u.
- 467 Li, J. and Shi, J.: (High-performance liquid chromatography with electrochemical detector for the determination of amniotic fluid catecholamine). *Fenxi Huaxue*, 21 (1993) 443-445; *C.A.*, 119 (1993) 86319w.
- 468 Ruban, V.F.: Determination of dopamine and its metabolites in microdialysates by capillary liquid chromatography with electrochemical detection. *J. Chromatogr.*, 619 (1993) 111-115.
- 469 Sakurai, E., Gunji, E., Iizuka, Y., Hikichi, N., Maeyama, K. and Watanabe, T.: *In vivo* microdialysis measurement of histamine in rat blood effects of compound 48/80 and histamine receptor antagonists. *J. Pharmacol. Toxicol. Methods*, 29 (1993) 105-109; *C.A.*, 119 (1993) 152226s.
- 470 Seiler, W. and Hiemke, C.: Determination of homovanillic acid in human plasma using HPLC with electrochemical detection and automated solid phase extraction. *J. Liq. Chromatogr.*, 16 (1993) 2813-2823.

See also 57, 72, 188, 189, 450, 1147, 1241.

## 17b. Catecholamines and their metabolites

- 471 Soga, T. and Inoue, Y.: Determination of catecholamines in urine and plasma by on-line sample pretreatment using an internal surface boronic acid gel. *J. Chromatogr.*, 620 (1993) 175-181.
- 472 Takahashi, M., Morita, M., Niwa, O. and Tabei, H.: Highly sensitive detection of catecholamines with interdigitated array microelectrodes in HPLC. *Sens. Actuators, B*, 13 (1993) 336-339; *C.A.*, 119 (1993) 152219s.
- 473 Zhang, H. and Stewart, J.T.: HPLC determination of norepinephrine bitartrate in 5% dextrose injection on underivatized silica with an aqueous-organic mobile phase. *J. Liq. Chromatogr.*, 16 (1993) 2861-2871.
- For additional information see *C.A.*:  
119 (1993) 112888c.
- See also 36, 70, 1204, 1233.
- 17d. *Other amine derivatives and amides (excl. peptides)*
- 474 Ikarashi, Y., Blank, C.L.R. and Maruyama, Y.: Binding characteristics of various neurochemicals to glassy carbon. *J. Chromatogr.*, 645 (1993) 219-231.
- 475 Kamimori, H., Hamashima, Y. and Konishi, M.: (Determination and urinary profiles of carnitine and acylcarnitines in human by high-performance liquid chromatography with fluorescence detection). *Kuromatogurafi*, 13 (1992) 267-268; *C.A.*, 119 (1993) 90415s.
- 476 Pirkle, W.H., Welch, C.J. and Zych, A.J.: Chromatographic investigation of the slowly interconverting atropisomers of hindered naphthamides. *J. Chromatogr.*, 648 (1993) 101-109.
- 477 Seto, Y. and Guengerich, F.P.: Liquid chromatographic determination of *para*-substituted N,N-dialkylaniline N-oxides. *J. Chromatogr.*, 619 (1993) 71-77.
- 478 Webb, L.E., Pavlina, T.M. and Hjelle, J.T.: A rapid high performance liquid chromatographic method for the analysis of choline in human plasma and peritoneal dialysis effluent: application in the assessment of choline loss in continuous ambulatory peritoneal dialysis. *Clin. Biochem.*, 26 (1993) 173-177; *C.A.*, 119 (1993) 155151z.
- For additional information see *C.A.*:  
119 (1993) 103443b, 117528v.
- See also 87, 222, 1425.
18. AMINO ACIDS AND PEPTIDES; CHEMICAL STRUCTURE OF PROTEINS
- 18a. *Amino acids and their derivatives*
- 479 Accinni, R., Belingheri, L., Giglioni, A., Micelli, G., Quaranta, M., Wei, J. and Lucarelli, L.: Quantitation of collagen/total protein ratio in biological samples by isocratic high performance liquid chromatography 4-hydroxyproline assay. *G. Ital. Chim. Clin.*, 17 (1992) 27-33; *C.A.*, 119 (1993) 112582y.
- 480 Andersson, A., Isaksson, A., Brattström, L. and Hultberg, B.: Monocysteine and other thiols determined in plasma by HPLC and thiol-specific postcolumn derivatization. *Clin. Chem. (Washington)*, 39 (1993) 1590-1597.
- 481 Armstrong, D.W., Zukowski, J., Ercal, N. and Gasper, M.: Stereochemistry of pipercolic acid found in the urine and plasma of subjects with peroxisomal deficiencies. *J. Pharm. Biomed. Anal.*, 11 (1993) 881-886.
- 482 Astephen, N. and Wheat, T.: An amino acid analysis method for assessing nutritional quality of infant formulas. *Am. Lab. (Shelton)*, 25 (1993) 24-27; *C.A.*, 119 (1993) 115626b.
- 483 Casillas, J.L., Addo-Yobo, F., Kenney, C.N., Aracil, J. and Martinez, M.: The use of modified divinylbenzene-polystyrene resins in the separation of fermentation products. A case study utilizing amino acids and a dipeptide. *J. Chem. Technol. Biotechnol.*, 55 (1992) 163-169; *C.A.*, 119 (1993) 96116s.
- 484 Chénais, B., Yapo, A., Lepoivre, M. and Tenu, J.-P.:  $\omega$ -Hydroxy-L-arginine, a reactional intermediate in nitric oxide biosynthesis, induced cytoxicity in human and murine tumor cells. *Biochem. Biophys. Res. Commun.*, 196 (1993) 1558-1565.
- 485 Darmaun, D., D'Amore, D. and Haymond, M.W.: Determination of glutamine and  $\alpha$ -ketoglutarate concentration and specific activity in plasma using high-performance liquid chromatography. *J. Chromatogr.*, 620 (1993) 33-38.
- 486 Dosena, A., Galaverna, G., Corradini, R. and Marchelli, R.: Two-dimensional high-performance liquid chromatographic system for the determination of enantiomeric excess in complex amino acid mixtures. Single amino acid analysis. *J. Chromatogr.*, 653 (1993) 229-234.
- 487 Endo, Y., Okayama, A., Endo, G., Horiguchi, S. and Nakazono, N.: (Improvement of urinary  $\delta$ -aminolevulinic acid determination by HPLC-fluorometry using pre-column derivatization). *Sangyo Igaku*, 35 (1993) 126-127; *C.A.*, 119 (1993) 88398v.
- 488 Fialaire, A., Postaire, E., Prognon, R. and Pradeau, D.: Bioavailability prediction of amino acids and peptides in nutritive mixtures by separation on non-polar stationary phases by high-performance liquid chromatography and direct UV detection at 210 nm. *J. Liq. Chromatogr.*, 16 (1993) 3003-3015.
- 489 Frey, J., Chamson, A. and Raby, N.: Separation of amino acids using ion-paired reversed-phase high-performance liquid chromatography with special reference to collagen hydrolysate. *Amino Acids*, 4 (1993) 45-51; *C.A.*, 119 (1993) 112603f.
- 490 Fujinari, E.M., Ribble, E. and Piserchio, M.V.: Nitrogen-specific liquid chromatography detection of amino acids, peptides, and proteins by HPLC-CLND. *Dev. Food Sci.*, 32(Food Flavors, Ingredients and Composition) (1993) 75-89; *C.A.*, 119 (1993) 179498u.
- 491 Hagan, R.L.: Determination of plasma homocysteine by HPLC with fluorescence detection: a survey of current methods. *J. Liq. Chromatogr.*, 16 (1993) 2701-2714 - a review with 36 refs.
- 492 Hancock, D.K. and Reeder, D.J.: Analysis and configuration assignments of the amino acids in a pyoverdine-type siderophore by reversed-phase high-performance liquid chromatography. *J. Chromatogr.*, 646 (1993) 335-343.
- 493 Hill, R.E., White, R.L. and Smith, K.C.: The identification of 5-hydroxy-L-norvaline in cultures of pyridoxine auxotrophs of *Escherichia coli*. *J. Natural Prod.*, 56 (1993) 1246-1254.
- 494 Hill, R.H., Jr., Caudill, S.P., Philen, R.M., Bailey, S.L., Flanders, W.D., Driskell, W.J., Kamb, M.L., Needham, L.L. and Sampson, E.J.: Contaminants in L-tryptophan associated with eosinophilia myalgia syndrome. *Arch. Environ. Contam. Toxicol.*, 25 (1993) 134-142; *C.A.*, 119 (1993) 125347a.



- 495 Hirano, H., Komatsu, S., Kajiwara, H., Takagi, Y. and Tsunawasa, S.: Microsequence analysis of the N-terminally blocked proteins immobilized on polyvinylidene difluoride membrane by Western blotting. *Electrophoresis (Weinheim)*, 14 (1993) 839-846.
- 496 Hyun, M.H., Ryoo, J.-J. and Lim, N.-E.: Optical resolution of racemic  $\alpha$ -amino acids on a dynamic chiral stationary phase by ligand exchange chromatography. *J. Liq. Chromatogr.*, 16 (1993) 3249-3261.
- 497 Iwaki, K., Yamazaki, M., Nimura, N., Kinoshita, T., Matsuda, K. and Hayashi, Y.: An information theory-based evaluation of stationary phases developed for optical resolution of amino acids. *Chromatographia*, 37 (1993) 156-158.
- 498 Jäger, W. and Koch, H.P.: Pharmacokinetics and organ distribution of liposome-encapsuled L-carnithine in rats. *Arzneim.-Forsch.*, 43 (1993) 974-977.
- 499 Jayatilleke, E. and Shaw, S.: A high-performance liquid chromatographic assay for reduced and oxidized glutathione in biological samples. *Anal. Biochem.*, 214 (1993) 452-457.
- 500 Lin, J.-Y. and Yang, M.-H.: High-performance liquid chromatographic separation of enantiomeric amino acids on bis[carbamoyl(alkyl)methylamino]-6-chloro-s-triazine-derived chiral stationary phases. *J. Chromatogr.*, 644 (1993) 277-283.
- 501 Mütsch-Eckner, M., Erdelmeier, C.A.J., Sticher, O. and Reuter, H.D.: A novel amino acid glycoside and three amino acids from *Allium sativum*. *J. Natural Prod.*, 56 (1993) 864-869.
- 502 Nagasawa, T., Ling, J.R. and Onodera, R.: Chiral high-performance liquid chromatographic separation of the three stereoisomers of 2,6-diaminopimelic acid without derivatisation. *J. Chromatogr.*, 653 (1993) 336-340.
- 503 Olken, N.M. and Marletta, M.A.: N<sup>G</sup>-Methyl-L-arginine functions as an alternate substrate and mechanism-based inhibitor of nitric oxide synthase. *Biochemistry*, 32 (1993) 9677-9685.
- 504 Parvy, P., Bardet, J., Rabier, D., Gasquet, M. and Kamoun, P.: Intra- and interlaboratory quality control for assay of amino acids in biological fluids: 14 years of the French experience. *Clin. Chem. (Washington)*, 39 (1993) 1831-1836.
- 505 Piotrovskij, V.K., Kállay, Z., Krejcy, K., Horecky, J., Trnovec, T. and Raberger, G.: N<sup>G</sup>-Nitro-L-arginine pharmacokinetics in rats after a single intravascular and oral dose: an appearance of secondary concentration time peaks. *Drug Metab. Disp.*, 21 (1993) 962-964.
- 506 Scislawski, P.W.D., Harris, I., Pickard, K., Brown, D.S. and Buchan, V.: Determination of L-methionine-dl-sulphoxide in tissue extracts. *J. Chromatogr.*, 619 (1993) 299-305.
- 507 Takahashi, M., Kushida, K., Kawana, K., Ishihara, C., Denda, M., Inoue, T. and Horiuchi, K.: Quantification of the cross-link pentosidine in serum from normal and uremic subjects. *Clin. Chem. (Washington)*, 39 (1993) 2162-2165.
- 508 Teerlink, T., Hennekes, M.W.T., van Leeuwen, P.A.M. and Houdijk, A.: Rapid-determination of glutamine in biological samples by high-performance liquid chromatography. *Clin. Chim. Acta*, 218 (1993) 159-168.
- 509 Tomokuni, K., Ichiba, M. and Hirai, Y.: Measurement of urinary  $\delta$ -aminolevulinic acid (ALA) by fluorometric HPLC and colorimetric methods. *Ind. Health*, 30 (1992) 119-128; *C.A.*, 119 (1993) 110646y.
- 510 Ueland, P.M., Refsum, H., Stabler, S.P., Malinow, M.R., Anderson, A. and Allen, R.H.: Total homocysteine in plasma or serum: methods and clinical applications. *Clin. Chem. (Washington)*, 39 (1993) 1764-1779 - a review with 129 refs.
- 511 Vaccher, C., Berthelot, P. and Debaert, M.: Direct separation of 4-amino-3-(4-chlorophenyl)butyric acid and analogues, GABA<sub>B</sub> ligands, using a chiral crown ether stationary phase. *J. Chromatogr.*, 645 (1993) 95-99.
- 512 Van Eijk, H.M.H., Rooyackers, D.R. and Deutz, N.E.P.: Rapid routine determination of amino acids in plasma by high-performance liquid chromatography with a 2-3  $\mu$ m Spherisorb ODS II column. *J. Chromatogr.*, 620 (1993) 143-148.
- 513 Van Leuken, R.G.J., Kwakkenbos, G.T.C. and Duchateau, A.L.L.: Volatile ion-pairing agents for liquid chromatographic-thermospray mass spectrometric determination of amino acids and amino acid amides. *J. Chromatogr.*, 647 (1993) 131-136.
- 514 Wang, F., Su, C., Hollfelder, K., Waddington, D. and Pan, Y.-C.E.: Electroblothing proteolytic products from native gel for direct N-terminal sequence analysis: An approach for studying protein-protein interaction. *Electrophoresis (Weinheim)*, 14 (1993) 847-851.
- 515 Wolf, C.-F., Hillmann, R., Keck, F.S., Wennauer, R. and Grünert, A.: Determination of thyroacetates in human sera by high performance liquid chromatography. *Eur. J. Clin. Chem. Clin. Biochem.*, 31 (1993) A57.
- 516 Yang, J. and Hage, D.S.: Characterization of the binding and chiral separation of D- and L-tryptophan on a high-performance immobilized human serum albumin column. *J. Chromatogr.*, 645 (1993) 241-250.
- 517 Yokoyama, Y., Tsuchiya, M., Sato, H. and Kakinuma, H.: Determination of urinary creatinine, amino acids and organic acids by reversed-phase high-performance liquid chromatography with ultraviolet detection. *Kuromatogurafi*, 13 (1992) 265-266; *C.A.*, 119 (1993) 90414r.
- 518 Zhou, F.-X., Krull, I.S. and Feibush, B.: Solid-phase derivatization of amino acids and peptides in high-performance liquid chromatography. *J. Chromatogr.*, 648 (1993) 357-365.
- 519 Zhu, T., Peng, Y., Lackland, H. and Stein, S.: Confirmation of oligonucleotide-peptide structure by amino acid analysis. *Anal. Biochem.*, 214 (1993) 585-587.
- For additional information see *C.A.*:  
119 (1993) 135021d, 160826c.
- See also 30, 72, 83, 87, 88, 93, 125, 189, 190, 309, 465, 525, 592, 601, 656, 685, 715, 802, 840, 896, 952, 971, 1276, 1414.
- 18b. *Peptides, peptidic and proteinous hormones, growth factors*
- 520 Agerberth, B., Boman, A., Andersson, M., Jörnvall, H., Mutt, V. and Boman, H.G.: Isolation of three antibacterial peptides from pig intestine: gastric inhibitory polypeptide(7-42), diazepam-binding inhibitor(32-86) and a novel factor, peptide 3910. *Eur. J. Biochem.*, 216 (1993) 623-629.
- 521 Aguilar, M.I., Mougos, S., Boublik, J., Rivier, J. and Hearn, M.T.W.: High-performance liquid chromatography of amino acids, peptides and proteins. CXXVIII. Effect of D-amino acid substitutions on the reversed-phase high-performance liquid chromatography retention behaviour of neuropeptide Y[18-36] analogues. *J. Chromatogr.*, 646 (1993) 53-65.

- 522 Alewood, P.F., Bailey, A.J., Brinkworth, R.I., Fairlie, D. and Jones, A.: Characterisation of TNF- $\alpha$ -related peptides by high-performance liquid chromatography-mass spectrometry and high-performance liquid chromatography-tandem mass spectrometry. *J. Chromatogr.*, 646 (1993) 185-191.
- 523 Büllesbach, E.E. and Schwabe, C.: Mouse relaxin: synthesis and biological activity of the first relaxin with an unusual cross-linking pattern. *Biochem. Biophys. Res. Commun.*, 196 (1993) 311-319.
- 524 Chen, L., Mazzeo, J., Krull, I.S. and Wu, S.-L.: Determination of peptide 520 in human plasma using post-column photolysis with electrochemical detection in liquid chromatography. *J. Pharm. Biomed. Anal.*, 11 (1993) 999-1007.
- 525 Conlon, J.M., Youson, J.H. and Mommsen, T.P.: Structure and biological activity of glucagon and glucagon-like peptide from a primitive bony fish, the bowfin (*Amia calva*). *Biochem. J.*, 295 (1993) 857-861.
- 526 Corona, G., Skaper, S.D. and Callegaro, L.: Purification of different biologically active forms of mouse  $\beta$ -nerve growth factor by immobilized metal ion affinity chromatography. In: Angeletti, R.H. (Editor), *Tech. Protein Chem. IV, [Pap. Protein Soc. Symp.] 6th, 1992*, Academic, San Diego, 1993, pp. 343-351; C.A., 119 (1993) 131743t.
- 527 Esposito, G., Fogolari, F., Viglino, P., Cattarinussi, S., de Magistris, M.T., Chiappinelli, L. and Pessi, A.: Conformational study of a short Pertussis toxin T cell epitope incorporated in a multiple antigen peptide template by CD and two-dimensional NMR. Analysis of the structural effects on the activity of synthetic immunogens. *Eur. J. Biochem.*, 217 (1993) 171-187.
- 528 Fukuda, T., Yun, A. and Imai, K.: A rapid and sensitive determination of salmon calcitonin in solutions containing bovine serum albumin or gelatin. *Biomed. Chromatogr.*, 7 (1993) 229-230.
- 529 Fuse, N., Hayashi, Y., Fukata, J., Tominaga, T., Ebisui, O., Satoh, Y., Isohara, T., Uno, I. and Imura, H.: Purification and characterization of new anti-adrenocorticotropin rabbit neutrophil peptides (defensin). *Eur. J. Biochem.*, 216 (1993) 653-659.
- 530 Goodearl, A.D., Davis, J.B., Mistry, K., Minghetti, L., Otsu, M., Waterfield, M.D. and Stroobant, P.: Purification of multiple forms of glial growth factor. *J. Biol. Chem.*, 268 (1993) 18095-18102.
- 531 Idei, M., Mezö, I., Vadász, Z., Horváth, A., Teplán, I. and Kéri, G.: Comparison of high-performance liquid chromatography and capillary electrophoresis in the analysis of somatostatin analogue peptides. *J. Chromatogr.*, 648 (1993) 251-256.
- 532 Igout, A., van Beeumen, J., Frankenne, F., Scippo, M.-L., Devereese, B. and Hennen, G.: Purification and biochemical characterization of recombinant human placental growth hormone produced in *Escherichia coli*. *Biochem. J.*, 295 (1993) 719-724.
- 533 Jones, O.T. and So, A.P.: Preparation and characterization of biotinylated analogs of the calcium channel blocker  $\omega$ -conotoxin. *Anal. Biochem.*, 214 (1993) 227-232.
- 534 Kai, M., Kojima, E., Ohkura, Y. and Iwasaki, M.: High-performance liquid chromatography on N-terminal tryptophan-containing peptides with precolumn fluorescence derivatization with glyoxal. *J. Chromatogr.*, 653 (1993) 235-240.
- 535 Kaneko, T., Wood, G., Crouch, W.L. and Desiderio, D.M.: Assay of preproenkephalin A processing enzymes in human lumbar cerebrospinal fluid. *Anal. Biochem.*, 215 (1993) 45-53.
- 536 Kieffer, B.L., Black, D. and Hirth, C.G.:  $^{32}$ P-Labeled opioid peptides with high affinity for the  $\delta$ -opioid receptor. *Anal. Biochem.*, 215 (1993) 1-8.
- 537 Kim, B.G. and Brooks, C.L.: Isolation and characterization of phosphorylated bovine prolactin. *Biochem. J.*, 296 (1993) 41-47.
- 538 Kirby, D.A., Miller, C.L. and Rivier, J.E.: Separation of neuropeptide Y diastereomers by high-performance liquid chromatography and capillary zone electrophoresis. *J. Chromatogr.*, 648 (1993) 257-265.
- 539 Lambros, T., Lema, P., III, Liu, W., Bongers, J., Felix, A.M. and Heimer, E.P.: Capillary electrophoresis of multiple antigenic-peptide (MAPS) of the malaria circumsporozoite protein epitopes. *J. Liq. Chromatogr.*, 16 (1993) 2039-2048.
- 540 Lebl, M.: Observation of a conformational effect in peptide molecule by reversed-phase high-performance liquid chromatography. *J. Chromatogr.*, 644 (1993) 285-287.
- 541 Ledwith, B.J., Cahill, M.K., Losse, L.S., Satiritz, S.M., Eydeloth, R.S., Dallob, A.L., Tanaka, W.K., Galloway, S.M. and Nichols, W.W.: Measurement of plasma angiotensin II: purification by cation-exchange chromatography. *Anal. Biochem.*, 213 (1993) 349-355.
- 542 Lerro, K.A., Orlando, R., Zhang, H., Usherwood, P.N.R. and Nakanishi, K.: Separation of the sticky peptides from membrane proteins by high-performance liquid chromatography in a normal-phase system. *Anal. Biochem.*, 215 (1993) 38-44.
- 543 Miller, B.T. and Kurosky, A.: Elevated intrinsic reactivity of seryl hydroxyl groups within the linear peptide triads His-Xaa-Ser or Ser-Xaa-His. *Biochem. Biophys. Res. Commun.*, 196 (1993) 461-467.
- 544 Miyagishima, T., Gasa, S., Honke, K., Sakai, M., Nishi, S., Yamamoto, M., Nishikawa, K., Miyazaki, T. and Makita, A.: Expression, purification and binding to the receptor of human insulin-like growth factor II. *Biochim. Biophys. Acta*, 1203 (1993) 155-161.
- 545 Moriyama, S., Duguay, S.J., Conlon, J.M., Duan, C., Dickhoff, W.W. and Plisetskaya, E.M.: Recombinant coho salmon insulin-like growth factor I. Expression in *Escherichia coli*, purification and characterization. *Eur. J. Biochem.*, 218 (1993) 205-211.
- 546 Narastoni, M., Scaranari, V., Romualdi, P., Donatini, A., Ferri, S. and Tomatis, R.: Studies on the anti-phospholipase A<sub>2</sub> and anti-inflammatory activities of a uteroglobin fragment and related peptides. *Arzneim.-Forsch.*, 43 (1993) 997-1000.
- 547 Neven, P., Iles, R.K., Howes, I., Sharma, K., Shepherd, J.H., Edwards, R., Collins, W.P. and Chard, T.: Substantial urinary concentrations of material resembling  $\beta$ -core fragment of chorionic gonadotropin  $\beta$ -subunit in mid-menstrual cycle. *Clin. Chem. (Washington)*, 39 (1993) 1857-1860.
- 548 Pittenauer, E., Capparros Rodriguez, M., dePedro, M.A., Allmaier, G. and Schmid, E.R.: HPLC and californium-252 plasma desorption-mass spectrometry of mucopeptides isolated from *E. coli*. *FEMS Symp.*, 65(Bacterial Growth and Lysis) (1993) 31-38; C.A., 119 (1993) 177072b.
- 549 Salzet, M., Bulet, P., van Dorsseleer, A. and Malecha, J.: Isolation, structural characterization and biological function of a lysine-conopressin in the central nervous system of the Pharyngobdellin leech *Erpobdella octoculata*. *Eur. J. Biochem.*, 217 (1993) 897-903.
- 550 Schoneich, C., Zhao, F., Wilson, G.S. and Borchardt, R.T.: Iron-thiolate induced oxidation of methionine to methionine sulfoxide in small model peptides. Intramolecular catalysis by histidine. *Biochim. Biophys. Acta*, 1158 (1993) 307-322.

- 551 Sereda, T.J., Mant, C.T., Quinn, A.M. and Hodges, R.S.: Effect of the  $\alpha$ -amino group on peptide retention behaviour in reversed-phase chromatography. Determination of the  $pK_a$  values of the  $\alpha$ -amino group of 19 different N-terminal amino acid residues. *J. Chromatogr.*, 646 (1993) 17-30.
- 552 Shiomi, K., Nakazato, M., Ihi, T., Kangawa, K., Matsuo, H. and Matsukura, S.: Establishment of radioimmunoassay for human neutrophil peptides and their increases in plasma and neutrophil in infection. *Biochem. Biophys. Res. Commun.*, 195 (1993) 1336-1344.
- 553 Sillard, R., Schulz-Knappe, P., Vogel, P., Raida, M., Bensch, K.W., Forssmann, W.-G. and Mutt, V.: A novel 77-residue peptide from porcine brain contains a leucine-zipper motif and is recognized by an antiserum to delta-sleep-inducing peptide. *Eur. J. Biochem.*, 216 (1993) 419-436.
- 554 Sugiyama, K. and Ogata, K.: High-performance liquid chromatographic determination of histatins in human saliva. *J. Chromatogr.*, 619 (1993) 306-309.
- 555 Teramoto, T., Kuwada, M., Niidome, T., Sawada, K., Nishizawa, Y. and Katayama, K.: A novel peptide from funnel web spider venom,  $\omega$ -Aga-TK, selectively blocks P-type calcium channels. *Biochem. Biophys. Res. Commun.*, 196 (1993) 134-140.
- 556 Tsuruoka, N., Yamshiro, K. and Tsujimoto, M.: Purification of soluble murine interleukin 5 (IL-5) receptor  $\alpha$  expressed in Chinese hamster ovary cells and its action as an IL-5 antagonist. *Arch. Biochem. Biophys.*, 307 (1993) 1133-1137.
- 557 Vandebroek, K., Fiten, P., Beuken, E., Martens, E., Janssen, A., van Damme, J., Opendakker, G. and Billiau, A.: Gene sequence, cDNA construction, expression in *Escherichia coli* and genetically approached purification of porcine interleukin-1 $\beta$ . *Eur. J. Biochem.*, 217 (1993) 45-52.
- 558 Varro, A. and Dockray, G.J.: Post-translational processing of progastrin: inhibition of cleavage, phosphorylation and sulphation by brefeldin A. *Biochem. J.*, 295 (1993) 813-819.
- 559 Westcott, J.Y., Ono, S., Oka, M., Jones, K. and Woelkel, N.F.: Inactivation of peptidoleukotrienes in the isolated perfused rat lung. *Arch. Biochem. Biophys.*, 306 (1993) 44-51.
- 560 Zamai, M. and Caiola, V.R.: Sequence-directed recognition peptides: inhibition of endothelin generation via a substrate-depletion mechanism. *Biochim. Biophys. Acta*, 1202 (1993) 337-340.
- 561 Zhang, J.-G., Reid, G.E., Moritz, R.L., Ward, L.D. and Simpson, R.J.: Specific covalent modification of the tryptophan residues in murine interleukin-6. Effect on biological activity and conformational stability. *Eur. J. Biochem.*, 217 (1993) 53-59.
- For additional information see C.A.:  
119 (1993) 103295e, 109149g, 111828w, 125303h, 152221m, 152223p, 188715q.
- See also 36, 83, 95, 130, 309, 311, 399, 480, 483, 488, 490, 492, 518, 519, 580, 586, 601, 607, 622, 625, 629, 637, 642, 645, 662, 708, 715, 777, 791.
- 18c. *Elucidation of structure of proteins and enzymes*
- 562 Fujiwara, Y., Oka, M., Hamamoto, T. and Sone, N.: Cytochrome c-551 of the thermophilic bacterium PS3, DNA sequence and analysis of the mature cytochrome. *Biochim. Biophys. Acta*, 1144 (1993) 213-219.
- 563 Grieve, P.A., Jones, A. and Alewood, P.F.: Analytical methods for differentiating minor sequence variations in related peptides. *J. Chromatogr.*, 646 (1993) 175-184.
- 564 Jones, A.T., Keen, J.N. and Roberts, N.B.: Human pepsin 3b peptide map sequence analysis, genotype and hydrophobic nature. *J. Chromatogr.*, 646 (1993) 207-212.
- 565 Koikeda, S., Ando, K., Kaji, H., Inoue, T., Murao, S., Takeuchi, K. and Samejima, T.: Molecular cloning of the gene for bilirubin oxidase from *Myrothecium verrucaria* and its expression in yeast. *J. Biol. Chem.*, 268 (1993) 18801-18809.
- 566 Legrand, R., Falconnet, J.-B., Prevost, D., Schoot, B. and Devaux, P.: Peptide mapping of recombinant human interferon- $\gamma$  by reversed-phase liquid chromatography with on-line identification by thermospray mass spectrometry and UV absorption spectrometry. *J. Chromatogr.*, 647 (1993) 3-12.
- 567 Mock, K., Hail, M., Mylchreest, I., Zhou, J., Johnson, K. and Jardine, I.: Rapid high-sensitivity peptide mapping by liquid chromatography-mass spectrometry. *J. Chromatogr.*, 646 (1993) 169-174.
- 568 Nirasawa, S., Liu, X., Nishino, T. and Kurihara, Y.: Disulfide bridge structure of the heat-stable sweet protein mabinlin II. *Biochim. Biophys. Acta*, 1202 (1993) 277-280.
- 569 Olah, T.V., Perrault, A.R., Cooperman, B.S., Montesano-Roditis, L., McWilliams, R. and Gritz, D.G.: Incorporation of dinitrophenyl derivatives of proteins S6, S13, S16 and S18 into the 30 S subunit of *Escherichia coli* ribosomes by total reconstitution. *J. Biol. Chem.*, 268 (1993) 18696-18700.
- 570 Paine, L.J., Perry, N., Popplewell, A.G., Gore, M.G. and Atkinson, T.: The identification of a lysine residue reactive to pyridoxal-5-phosphate in the glycerol dehydrogenase from the thermophile *Bacillus stearothermophilus*. *Biochim. Biophys. Acta*, 1202 (1993) 235-243.
- 571 Ramilo, C., Braus, G. and Evans, J.N.S.: A tyrosine residue is involved in the allosteric binding of tryptophan to yeast chorismate mutase. *Biochim. Biophys. Acta*, 1203 (1993) 71-76.
- 572 Salzer, U., Ahorn, H. and Prohaska, R.: Identification of the phosphorylation site on human erythrocyte band 7 integral membrane protein: implications for a monotopic protein structure. *Biochim. Biophys. Acta*, 1151 (1993) 149-152.
- 573 Tanabe, K.: (Fractionation of capillary HPLC peaks for protein sequencing). *Seikagaku*, 65 (1993) 474-477; *C.A.*, 119 (1993) 112512a - a review with 6 refs.
- 574 Wajcman, H., Kister, J., Promé, D., Galacteros, F. and Gilsanz, F.: Hb Villaverde [ $\beta$  89 (F5) Ser $\rightarrow$ Thr]: the structural modification of an intrasubunit contact is responsible for a high oxygen affinity. *Biochim. Biophys. Acta*, 1225 (1993) 89-94.
- 575 Wawrzynów, A. and Collins, J.H.: Chemical modification of the Ca<sup>2+</sup>-ATPase of rabbit skeletal muscle sarcoplasmic reticulum: identification of sites labeled with aryl isothiocyanates and thiol-directed conformational probes. *Biochim. Biophys. Acta*, 1203 (1993) 60-70.
- 576 Yonemoto, W., Garrod, S.M., Bell, S.M. and Taylor, S.S.: Identification of phosphorylation sites in the recombinant catalytic subunit of cAMP-dependent protein kinase. *J. Biol. Chem.*, 268 (1993) 18626-18632.
- 577 Zhang, Z. and Dixon, J.E.: Active site labeling of the *Yersinia* protein tyrosine phosphatase: the determination of the  $pK_a$  of the active site cysteine and the function of the conserved histidine 402. *Biochemistry*, 32 (1993) 9340-9345.

For additional information see C.A.:

119 (1993) 135023f.

See also 518, 594, 610, 612, 616, 625, 646, 647, 657, 677, 678, 681, 687, 709, 713, 725, 752, 801, 830, 852, 855, 857.

## 19. PROTEINS

### 19a. General techniques

- 578 Aguilar, M.I. and Hearn, M.T.W.: HPLC of biological macromolecules. *Chem. Aust.*, 60 (1993) 284-286; C.A., 119 (1993) 155020f - a review with 28 refs.
- 579 Belenkii, B.G., Podkladenko, A.M., Kurenbin, O.I., Mal'tsev, V.G., Nasledov, D.G. and Trushin, S.A.: Peculiarities of zone migration and band broadening in gradient reversed-phase high-performance liquid chromatography of proteins with respect to membrane chromatography. *J. Chromatogr.*, 645 (1993) 1-15.
- 580 Chin, C.C.Q. and Wold, F.: The use of tributylphosphine and 4-(aminosulfoxy)-7-fluoro-2,1,3-benzoxadiazol in the study of protein sulfhydryls and disulfides. *Anal. Biochem.*, 214 (1993) 128-134.
- 581 Corradini, D., Filippetti, R. and Corradini, C.: Salt mediated elution behavior of proteins on a silica-based stationary phase for size exclusion high performance liquid chromatography. *J. Liq. Chromatogr.*, 16 (1993) 3393-3408.
- 582 Cramer, S.M. and Brooks, C.A.: Ion-exchange displacement chromatography of proteins. Theoretical and experimental studies. *ACS Symp. Ser.*, 529(Chromatography in Biotechnology) (1993) 27-42; C.A., 119 (1993) 176867c.
- 583 Edwards, S.L. and Dubin, P.L.: pH Effects on non-ideal protein size-exclusion chromatography on Superose 6. *J. Chromatogr.*, 648 (1993) 3-7.
- 584 Ekman, P. and Jäger, O.: Quantitation of subnanomolar amounts of phosphate bound to seryl and threonyl residues in phosphoproteins using alkaline hydrolysis and malachite green. *Anal. Biochem.*, 214 (1993) 138-141.
- 585 Heath, T.G., Thanabal, V. and Ye, Q.Z.: Rapid, accurate and sensitive determination of molecular weight of nitrogen-15 and carbon-13 labeled recombinant proteins by electrospray ionization mass spectrometry. *Biotechnol. Tech.*, 7 (1993) 367-372; C.A., 119 (1993) 90726u.
- 586 Hjertén, S., Nakazato, K., Mohammad, J. and Eaker, D.: Reversed-phase chromatography of proteins and peptides on compressed continuous beds. *Chromatographia*, 37 (1993) 287-294.
- 587 Hu, S. and Do, D.D.: Histidine-ligand chromatography of proteins. Multiple modes of binding mechanism. *J. Chromatogr.*, 646 (1993) 31-35.
- 588 Jen, S.C.D. and Pinto, N.G.: Influence of displacer properties on displacement chromatography of proteins: a theoretical study. *React. Polym.*, 19 (1993) 145-161; C.A., 119 (1993) 90439c.
- 589 Jungbauer, A.: Preparative chromatography of biomolecules. *J. Chromatogr.*, 639 (1993) 3-16 - a review with 75 refs.
- 590 Kaltenbrunner, O., Tauer, C., Brunner, J. and Jungbauer, A.: Isoprotein analysis by ion-exchange chromatography using a linear pH gradient combined with a salt gradient. *J. Chromatogr.*, 639 (1993) 41-49.
- 591 Kamihira, M., Iijima, S. and Kobayashi, T.: Stabilities of antigen and antibody under elution conditions in immunoaffinity chromatography using monoclonal antibody. *Bioseparation*, 3 (1992) 185-188; C.A., 119 (1993) 134701p.
- 592 Kao, M.C.C. and Chung, M.C.M.: Pyridylethylation of cysteine residues in proteins. *Anal. Biochem.*, 215 (1993) 82-85.
- 593 Krull, I.S., Szulc, M.E. and Wu, S.L.: Principles of biopolymer detection in HPLC. LC-GC, 11 (1993) 350-355; C.A., 119 (1993) 176799g - a review with 7 refs.
- 594 Larmann, J.P., Jr., Lemmo, A.V., Moore, A.W., Jr. and Jorgenson, J.W.: Two-dimensional separations of peptides and proteins by comprehensive liquid chromatography-capillary electrophoresis. *Electrophoresis (Weinheim)*, 14 (1993) 439-447.
- 595 Lu, T., van Dyke, M. and Sawadogo, M.: Protein-protein interaction studies using immobilized oligohistidine fusion proteins. *Anal. Biochem.*, 213 (1993) 318-322.
- 596 Mao, Q.M., Prince, I.G. and Hearn, M.T.W.: High-performance liquid chromatography of amino acids, peptides and proteins. CXXXII. Optimisation of operating parameters for protein purification with chromatographic columns. *J. Chromatogr.*, 646 (1993) 81-89.
- 597 Mao, Q.M., Stockmann, R., Prince, I.G. and Hearn, M.T.W.: High-performance liquid chromatography of amino acids, peptides and proteins. CXXVI. Modelling of protein adsorption with non-porous and porous particles in a finite bath. *J. Chromatogr.*, 646 (1993) 67-80.
- 598 Mislovičová, D., Petro, M. and Berek, D.: Behaviour of polyhydroxyethyl methacrylate sorbent with dextran-filled macropores in dye-affinity chromatography of proteins. *J. Chromatogr.*, 646 (1993) 411-416.
- 599 O'Donoghue, G.V., Okarma, T.B. and Lee, Y.M.: An immune complex selective affinity column utilizing site-specific attachment of bovine conglutinin. *Anal. Biochem.*, 213 (1993) 310-317.
- 600 Ohshita, T.: Chromatographic analysis of lysosomal degradation of unlabeled native proteins *in vitro* by fluorescein-isothiocyanate labeling. *Anal. Biochem.*, 215 (1993) 17-23.
- 601 Reim, D.F. and Speicher, D.W.: High-sensitivity gas phase sequence analysis of proteins and peptides on PVDF membranes using short cycle tissues. *Anal. Biochem.*, 214 (1993) 87-95.
- 602 Sadana, A.: Inactivation of proteins and other biological macromolecules during chromatographic methods of bioseparation. *Bioseparation*, 3 (1992) 145-165; C.A., 119 (1993) 155074b - a review with many refs.
- 603 Shimoni, M., Reuveni, R. and Cais, M.: Non-dialysis method of rapid and facile sample preparation for the desalting and purification of enzymes and other proteins from plant extracts. *J. Chromatogr.*, 646 (1993) 99-105.
- 604 Strege, M.A. and Lagu, A.L.: Capillary electrophoresis as a tool for the analysis of protein folding. *J. Chromatogr.*, 652 (1993) 179-188.
- 605 Tennikova, T.B. and Svec, F.: High-performance membrane chromatography: highly efficient separation method for proteins in ion-exchange, hydrophobic interaction and reversed-phase modes. *J. Chromatogr.*, 646 (1993) 279-288.
- 606 Tomono, T.: (Protein separations by affinity chromatography). *Kagaku Kogyo*, 44 (1993) 430-437; C.A., 119 (1993) 155025m - a review with 64 refs.

- 607 Wisniewski, R.: Principles of the design and operational considerations of large scale high performance liquid chromatography (HPLC) systems for proteins and peptides purification. *Bioseparation*, 3 (1992) 77-143; C.A., 119 (1993) 155073a - a review with many refs.
- 608 Zachariou, M., Traverso, I. and Hearn, M.T.W.: High-performance liquid chromatography of amino acids, peptides and proteins. CXXXI. O-Phosphoserine as a new chelating ligand for use with hard Lewis metal ions in the immobilized-metal affinity chromatography of proteins. *J. Chromatogr.*, 646 (1993) 107-120.

For additional information see C.A.:  
119 (1993) 176823k.

See also 31, 96, 99, 109, 122, 123, 124, 158, 490, 755, 1166, 1364, 1431.

19b. *Proteins of cells, viruses and subcellular particles*

- 609 Ganoza, M.C., Aoki, H., Koziaradzki, I. and Schwartz, I.: Reconstruction of translation. Evidence for the involvement of the rescue protein in the association/dissociation of ribosomal subunits. *Eur. J. Biochem.*, 217 (1993) 839-847.
- 610 Gorman, J.J. and Shiell, B.J.: Isolation of carboxyl-termini and blocked amino-termini of viral proteins by high-performance cation-exchange chromatography. *J. Chromatogr.*, 646 (1993) 193-205.
- 611 Møller, J.V. and le Maire, M.: Detergent binding as a measure of hydrophobic surface area of integral membrane proteins. *J. Biol. Chem.*, 268 (1993) 18659-18672.
- 612 Nice, E., Layton, J., Fabri, L., Hellman, U., Engstrom, A., Persson, B. and Burgess, A.W.: Mapping of the antibody- and receptor-binding domains of granulocyte colony-stimulating factor using an optical biosensor. Comparison with enzyme-linked immunosorbent assay competition studies. *J. Chromatogr.*, 646 (1993) 159-168.
- 613 Scherzinger, E., Kruff, V. and Otto, S.: Purification of the large mobilization protein of plasmid RSF1010 and characterization of its site-specific DNA-cleaving/DNA-joining activity. *Eur. J. Biochem.*, 217 (1993) 929-938.
- 614 Watanabe, Y. and Takagi, T.: Characterization of further association of the trimeric membrane protein porin by low-angle laser-light scattering photometry coupled with high-performance gel chromatography. *J. Chromatogr.*, 653 (1993) 241-246.
- 615 Welling-Wester, S., Feijlbrief, M., Koedijk, D.G.A.M., Braaksma, M.A., Douma, B.R.K. and Welling, G.W.: Effect of different amounts of the non-ionic detergents C<sub>10</sub>E<sub>5</sub> and C<sub>12</sub>E<sub>5</sub> present in eluents for ion-exchange high-performance liquid chromatography of integral membrane proteins of Sendai virus. *J. Chromatogr.*, 646 (1993) 37-44.

See also 569, 594.

19c. *Proteins synthesized by genetic manipulation, monoclonal antibodies*

- 616 Facchetti, I., Valla, A. and Vigevani, A.: RP-HPLC peptide mapping methods for the analysis of recombinant human pro-urokinase. *J. Pharm. Biomed. Anal.*, 11 (1993) 737-744.

- 617 Yowell, G.G., Fazio, S.D. and Vivilecchia, R.V.: Analysis of a recombinant granulocyte macrophage colony stimulating factor dosage form by capillary electrophoresis, capillary isoelectric focusing and high-performance liquid chromatography. *J. Chromatogr.*, 652 (1993) 215-224.

See also 158, 621, 664, 680, 681, 903.

19d. *Microbial and plant proteins*

- 618 Coutos-Thevenot, P., Jouenne, T., Maes, O., Guerbette, F., Grosbois, M., le Caer, J.P., Boulay, M., Deloire, A., Kader, J.C. and Guern, J.: Four 9-kDa proteins excreted by somatic embryos of grapevine are isoforms of lipid-transfer proteins. *Eur. J. Biochem.*, 217 (1993) 885-889.
- 619 Ferreras, J.M., Barbieri, L., Girbés, T., Battelli, M.G., Rojo, M.A., Arias, F.J., Rocher, M.A., Soriano, F., Mendéz, E. and Stirpe, F.: Distribution and properties of major ribosome-inactivating proteins (28 S rRNA N-glycosidases) of the plant *Saponaria officinalis* L. (Caryophyllaceae). *Biochim. Biophys. Acta*, 1216 (1993) 31-42.
- 620 Ginnaga, A., Morokuma, K., Furukawa, S., Shigaki, T., Katsuki, N., Aihara, K., Oka, T., Sakoh, M., Tamura, C. and Imaizumi, A.: Further evaluation of the pertussis component vaccine produced by apoceruloplasmin affinity chromatography. *Dev. Biol. Stand.*, 73 (Pertussis: Evaluation and Research on Acellular Pertussis Vaccines) (1991) 233-241; C.A., 119 (1993) 136968e.
- 621 Morris, S.A., Revill, W.P., Staunton, J. and Leadlay, P.F.: Purification and separation of holo- and apo-forms of *Saccharopolyspora erythraea* acyl-carrier protein released from recombinant *Escherichia coli* by freezing and thawing. *Biochem. J.*, 294 (1993) 521-527.
- 622 Nishio, H., Kumagaye, K.Y., Kubo, S., Chen, Y.-n., Momiyama, A., Takahashi, T., Kimura, T. and Sakakibara, S.: Synthesis of  $\omega$ -agatoxin IV A and its related peptides. *Biochem. Biophys. Res. Commun.*, 196 (1993) 1447-1453.
- 623 Parente, A., de Luca, P., Bolognesi, A., Barbieri, L., Battelli, M.G., Abbondanza, A., Sande, M.J.W., Gigliano, G.S., Tazzari, P.L. and Stirpe, F.: Purification and partial characterization of single-chain ribosome-inactivating proteins from the seeds. *Biochim. Biophys. Acta*, 1216 (1993) 43-49.
- 624 Shone, C.C., Quinn, C.P., Wait, R., Hallis, B., Fooks, S.G. and Hambleton, P.: Proteolytic cleavage of synthetic fragments of vesicle-associated membrane protein, isoform-2 by botulinum type B neurotoxin. *Eur. J. Biochem.*, 217 (1993) 965-971.
- 625 Villalba, M., Batanero, E., López-Otín, C., Sánchez, L.M., Monsalve, R.I., González de la Pena, M.A., Lahoz, C. and Rodríguez, R.: The amino acid sequence of *Olea e i*, the major allergen from olive tree (*Olea europaea*) pollen. *Eur. J. Biochem.*, 216 (1993) 863-869.

See also 858.

19e. *Proteins of blood, serum and blood cells*

- 626 Basuyaux, B., Laroze, A. and Batard, T.: Pharmaceutical composition containing anti-allergenic antibodies from human plasma. *PCT Int. Appl. WO 93 11,160* (Cl. C07K15/06), 10 Jun. 1993, FR Appl. 91/14,954, 03 Dec. 1991; 26 p.; C.A., 119 (1993) 103310f.

- 627 Finch, J.W., Crouch, R.K., Knapp, D.R. and Schey, K.L.: Mass spectrometric identification of modifications to human serum albumin treated with hydrogen peroxide. *Arch. Biochem. Biophys.*, 305 (1993) 595-599.
- 628 Fleury, V., Loyau, S., Lijnen, H.R., Nieuwenhuizen, W. and Angles-Cano, E.: Molecular assembly of plasminogen and tissue-type plasminogen activator on an evolving fibrin surface. *Eur. J. Biochem.*, 216 (1993) 549-556.
- 629 Inoue, K. and Morita, T.: Identification of O-linked oligosaccharide chains in the activation peptides of blood coagulation factor X. The role of the carbohydrate moieties in the activation of factor X. *Eur. J. Biochem.*, 218 (1993) 153-163.
- 630 Ito, K., Baba, M., Yauda, Y., Sekiguchi, S. and Shimogori, Y.: Early pregnancy factor: Its characterization and purification from pregnant swine serum. *J. Reprod. Dev.*, 39 (1993) 129-138; C.A., 119 (1993) 134694p.
- 631 Jeppsson, J.-O., Kristensson, H. and Fimiani, C.: Carbohydrate-deficient transferrin quantified by HPLC to determine heavy consumption of alcohol. *Clin. Chem. (Washington)*, 39 (1993) 2115-2120.
- 632 Kowalska, M.A. and Tuszynski, G.P.: Interaction of thrombospondin with platelet glycoproteins GPI<sub>a</sub>-II<sub>a</sub> and GPII<sub>b</sub>-III<sub>a</sub>. *Biochem. J.*, 295 (1993) 725-730.
- 633 Leibl, H., Erber, W., Eibl, M.M. and Mannhalter, J.W.: Separation of polysaccharide-specific human immunoglobulin G subclasses using a Protein A Superose column with a pH gradient elution system. *J. Chromatogr.*, 639 (1993) 51-56.
- 634 Lütkemeyer, D., Bretschneider, M., Büntemeyer, H. and Lehmann, J.: Membrane chromatography for rapid purification of recombinant antithrombin III and monoclonal antibodies from cell culture supernatant. *J. Chromatogr.*, 639 (1993) 57-66.
- 635 Maruo, K., Akaike, T., Inada, Y., Ohkubo, I., Ono, T. and Maeda, H.: Effect of microbial and mite proteases on low and high molecular weight kininogens. Generation of kinin and inactivation of thiol protease inhibitory activity. *J. Biol. Chem.*, 268 (1993) 17711-17715.
- 636 Morley, S.J., Rau, M., Kay, J.E. and Pain, V.M.: Increased phosphorylation of eukaryotic initiation factor 4 $\alpha$  during early activation of T lymphocytes correlates with increased initiation factor 4F complex formation. *Eur. J. Biochem.*, 218 (1993) 39-48.
- 637 Nishihira, J., Koyama, Y., Sakai, M. and Nishi, S.: The fatty acid binding site of human  $\alpha$ -fetoprotein. *Biochem. Biophys. Res. Commun.*, 196 (1993) 1049-1057.
- 638 Pedroni, S., Lecomte, M.C., Gautero, H. and Dhermy, D.: Heterogeneous phosphorylation of erythrocyte spectrin  $\beta$ -chain in intact cells. *Biochem. J.*, 294 (1993) 841-846.
- 639 Rodriguez, W.V., Pritchard, P.H. and Hope, M.J.: The influence of size and composition on the cholesterol mobilizing properties of liposomes *in vivo*. *Biochim. Biophys. Acta*, 1153 (1993) 9-19.
- 640 Rosing, J., Bakker, H.M., Thomassen, M.C.L.G.D., Hemker, H.C. and Tans, G.: Characterization of two forms of human factor Va with different cofactor activities. *J. Biol. Chem.*, 268 (1993) 21130-21136.
- 641 Sugiyama, K., Ahorn, H., Maurer-Fogy, I. and Voss, T.: Expression of human interferon- $\alpha$ 2 in Sf9 cells. Characterization of O-linked glycosylation and protein heterogeneities. *Eur. J. Biochem.*, 217 (1993) 921-927.
- 642 Voss, T., Ergülen, E., Ahorn, H., Kubelka, V., Sugiyama, K., Maurer-Fogy, I. and Glössl, J.: Expression of human interferon  $\omega$ 1 in Sf9 cells. No evidence for complex-type N-linked glycosylation or sialylation. *Eur. J. Biochem.*, 217 (1993) 913-919.
- 643 Washizu, K., Kimura, S., Hiraiwa, H., Matsunaga, K., Kuwabara, M., Ariyoshi, Y., Kato, K. and Takeuchi, K.: Development and application of an enzyme immunoassay for tenascin. *Clin. Chim. Acta*, 219 (1993) 15-22.
- 644 Weykamp, C.W., Penders, T.J., Muskiet, F.A.J. and van der Slik, W.: Influence of hemoglobin variants and derivatives on glycohemoglobin determinations, as investigated by 102 laboratories using 16 methods. *Clin. Chem. (Washington)*, 39 (1993) 1717-1723.

For additional information see C.A.:

119 (1993) 133887s, 134693n, 136965b, 155506u.

See also 36, 590, 591, 720.

#### 19f. Structural and muscle proteins

- 645 Andreassen, L., Hojrup, P., Andersen, S.O. and Roepstorff, P.: Combined plasma-desorption mass spectrometry and Edman degradation applied to simultaneous sequence determination of isoforms of structural proteins from the cuticle of *Locusta migratoria*. *Eur. J. Biochem.*, 217 (1993) 267-273.
- 646 Resing, K.A., Johnson, R.S. and Walsh, K.A.: Characterization of protease processing sites during conversion of rat profilaggrin to filaggrin. *Biochemistry*, 32 (1993) 10036-10045.
- 647 Steinert, P.M., Marekov, L.N. and Parry, D.A.D.: Conservation of the structure of keratin intermediate filaments: molecular mechanism by which different keratin molecules integrate into pre-existing keratin intermediate filaments during differentiation. *Biochemistry*, 32 (1993) 10046-10056.
- 648 Vendeland, S.C., Beilstein, M.A., Chen, C.L., Jensen, O.N., Barofsky, E. and Whanger, P.D.: Purification and properties of selenoprotein W from rat muscle. *J. Biol. Chem.*, 268 (1993) 17103-17107.
- 649 Wu, C.: Simultaneous calibration of size exclusion chromatography and dynamic light scattering for the characterization of gelatine. *Macromolecules*, 26 (1993) 5423-5426; C.A., 119 (1993) 162789s.

See also 479, 580.

#### 19g. Protamines, histones and other nuclear proteins

- 650 Lindner, H., Wurm, M., Dirschlmaier, A., Sarg, B. and Helliger, W.: Application of high-performance capillary electrophoresis to the analysis of H1 histones. *Electrophoresis (Weinheim)*, 14 (1993) 480-485.
- 651 Zagariya, A., Khrapunov, S. and Zacharias, W.: Rapid method for the fractionation of nuclear proteins and their complexes by batch elution from hydroxyapatite. *J. Chromatogr.*, 648 (1993) 275-278.

See also 656.

19h. *Chromoproteins and metalloproteins*

- 652 Apostolova, M., Bontchev, P.R., Nachev, C. and Sirakova, I.: Apometallothionein in rat liver. *J. Chromatogr.*, 620 (1993) 191-197.
- 653 Caroli, S., la Torre, F., Petrucci, F., Senofonte, O. and Violante, N.: The chemical speciation of proteic iron using HPLC/ICP-AES. The case of ferritin. *Spectroscopy (Eugene)*, 8 (1993) 46-50; *C.A.*, 119 (1993) 112852m.
- 654 Chou, H.N. and Jeng, J.M.: High-performance liquid chromatography of biliprotein from *Bangia atropurpurea* 'Conchocelis'. *Hydrobiologia*, 260-261 (1993) 571-576; *C.A.*, 119 (1993) 176891f.
- 655 Cohen, M.P., Witt, J. and Wu, V.Y.: Purified hemoglobin preparations in the evaluation of HBA<sub>1c</sub> determination by ion exchange chromatography. *Ann. Clin. Biochem.*, 30 (1993) 265-271; *C.A.*, 119 (1993) 155168k.
- 656 Dallinger, R., Berger, B., Hunziker, P.E., Birchler, N., Hauer, C.R. and Kägi, J.H.R.: Purification and primary structure of snail metallothionein. Similarity of the N-terminal sequence with histones H4 and H2A. *Eur. J. Biochem.*, 216 (1993) 739-746.
- 657 Demuyneck, S., Li, K.W., van der Schors, R. and Dhainaut-Courtouis, N.: Amino acid sequence of the small cadmium. Binding protein (MPlI) from *Nereis diversicolor* (annelida, polychaeta). Evidence for a myohemerythrin structure. *Eur. J. Biochem.*, 217 (1993) 151-156.
- 658 Ganss, B. and Hoffmann, W.: Calcium binding to sialic acids and its effect on the conformation of ependymins. *Eur. J. Biochem.*, 217 (1993) 275-280.
- 659 Groten, J.P., Hissink, E. and van Bladeren, P.J.: Differences in chromatographic behavior of rat and pig metallothionein isoforms: A possible method of distinguishing between exogenous and endogenous metallothioneins. *IARC Sci. Publ.*, 118(Cadmium in the Human Environment) (1992) 219-224; *C.A.*, 119 (1993) 110793u.
- 660 Kimura, M., Jin, N. and Itokawa, Y.: (Determination of metallothionein in rat liver and kidney by high-performance liquid chromatography). *Biomed. Res. Trace Elem.*, 3 (1992) 139-140; *C.A.*, 119 (1993) 134690j.
- 661 Kutlar, F. and Huisman, T.H.J.: New ultra-micro high-performance liquid chromatographic method for determining the  $\gamma$  chain composition of hemoglobin F in normal adults. *J. Chromatogr.*, 620 (1993) 183-189.
- 662 Mackay, E.A., Overnell, J., Dunbar, B., Davidson, I., Hunziker, P.E., Kägi, J.H.R. and Fothergill, J.E.: Complete amino acid sequences of five dimeric and four monomeric forms of metallothionein from the edible mussel *Mytilus edulis*. *Eur. J. Biochem.*, 218 (1993) 183-194.
- 663 Manely, K., Ruangviriyachai, C. and Glennon, J.D.: Analysis of the microbial siderophore pseudobactin and its iron(III) complex by high-performance liquid chromatography with amperometric detection. *Anal. Proc.*, 30 (1993) 154-156; *C.A.*, 119 (1993) 134672e.
- 664 Martina, W.V., Martijn, E.G., van der Molen, M., Schermer, J.G. and Muskiet, F.A.J.:  $\beta$ -N-Terminal glycohemoglobins in subjects with common hemoglobinopathies: relation with fructosamine and mean erythrocyte age. *Clin. Chem. (Washington)*, 39 (1993) 2259-2265.
- 665 Richards, M.P. and Beattie, J.H.: Characterization of metallothionein isoforms. Comparison of capillary zone electrophoresis with reversed-phase high-performance liquid chromatography. *J. Chromatogr.*, 648 (1993) 459-468.
- 666 Roos, P.H.: Analysis of microsomal cytochrome P-450 patterns. Fast protein liquid chromatography with ion-exchange and immobilized metal affinity stationary phases. *ACS Symp. Ser.*, 529(Chromatography in Biotechnology) (1993) 112-131; *C.A.*, 119 (1993) 111840u.
- 667 Roush, D.J., Gill, D.S. and Willson, R.C.: Anion-exchange chromatographic behavior of recombinant rat cytochrome b<sub>5</sub>. Thermodynamic driving forces and temperature dependence of the stoichiometric displacement parameter Z. *J. Chromatogr.*, 653 (1993) 207-218.
- 668 Sonoda, Y., Endo, M., Ishida, K., Sato, Y., Fukusen, N. and Fukuhara, M.: Purification of a human cytochrome P-450 isozyme catalyzing lanosterol 14 $\alpha$ -demethylation. *Biochim. Biophys. Acta*, 1170 (1993) 92-97.
- 669 Steinebach, O.M. and Wolterbeek, H.T.: Determination of zinc-65, copper-64 and sulphur-35 labelled rat hepatoma tissue culture metallothioneins by high-performance liquid chromatography with on-line radioactivity detection. *J. Chromatogr.*, 619 (1993) 199-214.
- 670 Tamegai, H., Yamanaka, T. and Fukumori, Y.: Purification and properties of a 'cytochrome a<sub>1</sub>'-like hemoprotein from a magnetotactic bacterium, *Aquaspirillum magnetotacticum*. *Biochim. Biophys. Acta*, 1158 (1993) 237-243.

For additional information see *C.A.*:

119 (1993) 112606j, 112930k, 153509y.

See also 492, 644.

19i. *Proteins of glands, gland products, various zymogens (incl. milk proteins)*

- 671 De Jong, N., Visser, S. and Olieman, C.: Determination of milk proteins by capillary electrophoresis. *J. Chromatogr.*, 652 (1993) 207-213.
- 672 Garcia-Calvo, M., Leonard, R.J., Novick, J., Stevens, S.P., Schmalhofer, W., Kaczorowski, G.J. and Garcia, M.L.: Purification, characterization, and biosynthesis of margatoxin, a component of *Centruroides margaritatus* venom that selectively inhibits voltage-dependent potassium channels. *J. Biol. Chem.*, 268 (1993) 18866-18874.
- 673 Itagaki, T., Yoshida, M., Abe, S., Omichi, H. and Nishihira, Y.: Separation of human tear proteins with ceramic hydroxyapatite high-performance liquid chromatography. *J. Chromatogr.*, 620 (1993) 149-152.
- 674 Law, A.J.R. and Tziboula, A.: Fractionation of caprine  $\kappa$ -casein and examination of polymorphism by FPLC. *Milchwissenschaft*, 48 (1993) 68-71; *C.A.*, 119 (1993) 93815h.
- 675 Mori, E., Baba, T., Iwamatsu, A. and Mori, T.: Purification and characterization of a 38-kDa protein, sp38, with zona pellucida-binding property from porcine epididymal sperm. *Biochem. Biophys. Res. Commun.*, 196 (1993) 196-202.
- 676 Sharma, S.K., Hill, A.R. and Mittal, G.S.: An improved method to measure glycomacropptides (GMP) in renneted milk. *Milchwissenschaft*, 48 (1993) 71-73; *C.A.*, 119 (1993) 93816j.

- 677 Vita, C., Bontems, F., Bouet, F., Tauc, M., Poujeol, P., Vatanpour, H., Harvey, A.L., Menez, A. and Toma, F.: Synthesis of charybdotoxin and of two N-terminal truncated analogues. Structural and functional characteristics. *Eur. J. Biochem.*, 217 (1993) 157-169.
- 678 Yoo, S.H.: pH-Dependent binding of chromogranin B and secretory vesicle matrix proteins to the vesicle membrane. *Biochim. Biophys. Acta*, 1179 (1993) 239-246.

For additional information see C.A.:  
119 (1993) 112574x.

See also 533, 684, 695, 697, 705.

19j. *Proteins of brain, cerebrospinal fluid and eye*

- 679 Asano, T., Morishita, R., Matsuda, T., Fukada, Y., Yoshizawa, T. and Kato, K.: Purification of four forms of the  $\beta\gamma$  subunit complex of G proteins containing different  $\gamma$  subunits. *J. Biol. Chem.*, 268 (1993) 20512-20519.
- 680 Benedek, K.: Kinetics of recombinant human brain-derived neurotrophic factor unfolding under reversed-phase liquid chromatography conditions. *J. Chromatogr.*, 646 (1993) 91-98.
- 681 Fabri, L., Maruta, H., Muramatsu, H., Muramatsu, T., Simpson, R.J., Burgess, A.W. and Nice, E.C.: Structural characterisation of native and recombinant forms of the neurotrophic cytokine MK. *J. Chromatogr.*, 646 (1993) 213-225.
- 682 Gardela, J.E., Gorgone, G.A., Candela, L., Ghiso, J., Castano, E.M., Frangione, B. and Gorevic, P.D.: High-level expression and *in vitro* mutagenesis of a fibrillogenic 109-amino-acid C-terminal fragment of Alzheimer's disease amyloid precursor protein. *Biochem. J.*, 294 (1993) 667-674.
- 683 Huang, K.-P., Huang, F.L. and Chen, H.-C.: Characterization of a 7.5-kDa protein kinase C substrate (RC3 protein, neurogranin) from rat brain. *Arch. Biochem. Biophys.*, 305 (1993) 570-580.
- 684 Metz-Boutigue, M.-H., Garcia-Sablone, P., Hogue-Angeletti, R. and Aunis, D.: Intracellular and extracellular processing of chromogranin A. Determination of cleavage sites. *Eur. J. Biochem.*, 217 (1993) 247-257.
- 685 Nakajo, S., Tsukada, K., Omata, K., Nakamura, Y. and Nakaya, K.: A new brain-specific 14-kDa protein is a phosphoprotein. Its complete amino acid sequence and evidence for phosphorylation. *Eur. J. Biochem.*, 217 (1993) 1057-1063.
- 686 Tanaka, J., Ogawara, M., Ando, S., Shibata, M., Yatani, R., Kusagawa, M. and Inagaki, M.: Phosphorylation of a 62 kD porcine  $\alpha$ -internexin, a newly identified intermediate filament protein. *Biochem. Biophys. Res. Commun.*, 196 (1993) 115-123.
- 687 Van Haeringen, B., van den Bogaerde, M.R., Eden, D., van Grondelle, R. and Bloemendal, M.: Further characterization of structural and electric properties of non-spherical  $\alpha$ -crystallin. *Eur. J. Biochem.*, 217 (1993) 143-150.
- 688 Watkinson, A., Rogers, M. and Dockray, G.J.: Post-translational processing of chromogranin A: differential distribution of phosphorylated variants of pancreastatin and fragments 248-313 and 297-313 in bovine pancreas and ileum. *Biochem. J.*, 295 (1993) 649-654.
- 19k. *Proteins of neoplastic tissue and transformed cells*
- 689 Leith, I.R. and Russell, W.C.: Recognition of non-B DNA structures by cellular proteins. *Biochem. Biophys. Res. Commun.*, 196 (1993) 601-610.
- See also 637.
- 19l. *Specific binding and receptor proteins*
- 690 Aoki, T., Yamao, Y., Yonemasu, E., Kumasaki, Y. and Kako, Y.: Cross-linking of egg white riboflavin-binding protein by calcium phosphate. *Arch. Biochem. Biophys.*, 305 (1993) 242-246.
- 691 Backlund, P.S., Jr.: Carboxyl methylation of the low molecular weight GTP-binding protein G25K: regulation of carboxyl methylation by rhoGDI. *Biochem. Biophys. Res. Commun.*, 196 (1993) 534-542.
- 692 Banner, C.D., Göttlicher, M., Widmark, E., Sjövall, J., Rafter, J.J. and Gustafsson, J.-Å.: A systematic analytical chemistry/cell assay approach to isolate activators of orphan nuclear receptors from biological extracts: characterization of peroxisome proliferator-activated receptor activators in plasma. *J. Lipid Res.*, 34 (1993) 1583-1591.
- 693 Baxter, R.C., Suikkari, A.-M. and Martin, J.L.: Characterization of the binding defect in insulin-like growth factor binding protein-3 from pregnancy serum. *Biochem. J.*, 294 (1993) 847-852.
- 694 Bisgaier, C.L., Minton, L.L., Essenburg, A.D., White, A. and Homan, R.: Use of fluorescent cholesteryl ester microemulsions in cholesteryl ester transfer protein assay. *J. Lipid Res.*, 34 (1993) 1625-1634.
- 695 Desnoyers, L. and Manjunath, P.: Interaction of a novel class of phospholipid-binding proteins of bovine seminal fluid with different affinity matrices. *Arch. Biochem. Biophys.*, 305 (1993) 341-349.
- 696 Dutta-Roy, A.K., Leishman, D.J., Gordon, M.J., Campbell, F.M. and Duthie, G.G.: Identification of a low molecular mass (14.2 kDa)  $\alpha$ -tocopherol-binding protein in the cytosol of rat liver and heart. *Biochem. Biophys. Res. Commun.*, 196 (1993) 1108-1112.
- 697 Ehrlich, K.C.: Partial purification of a pea seed DNA-binding protein that specifically recognizes 5-methylcytosine. *Prepar. Biochem.*, 23 (1993) 423-438.
- 698 Ferron, L., Ferreiros, C.M., Criado, M.T. and Andrade, M.P.: Purification of the *Neisseria meningitidis* transferrin binding protein-2 (TBP2) to homogeneity using column chromatography. *FEMS Microbiol. Lett.*, 109 (1993) 159-165; C.A., 119 (1993) 90441x.
- 699 Gonzáles-Noriega, A., Ferrera, P., Coutino, R. and Michalak, C.: Isolation of an inhibitor for the independent mannose 6-phosphate receptor endocytosis of bovine  $\beta$ -glucuronidase by human fibroblasts. *Arch. Biochem. Biophys.*, 306 (1993) 331-336.
- 700 Grosbois, M., Guerbette, F., Jolliot, A., Quintin, F. and Kader, J.-C.: Control of maize lipid transfer protein activity by oxidoreducing conditions. *Biochim. Biophys. Acta*, 1170 (1993) 197-203.
- 701 Hakki, S., Robertson, D.C. and Waldman, S.A.: A 56 kDa binding protein for *Escherichia coli* heat-stable enterotoxin isolated from the cytoskeleton of rat intestinal membranes does not possess guanylate cyclase activity. *Biochim. Biophys. Acta*, 1152 (1993) 1-8.

See also 746.



- 702 Herr, F.M., Wardlaw, S.A., Kakkad, B., Albrecht, A., Quick, T.C. and Ong, D.E.: Intestinal vitamin A metabolism: coordinate distribution of enzymes and CRBP(II). *J. Lipid Res.*, 34 (1993) 1545-1554.
- 703 Jochen, A. and Hays, J.: Purification of the major substrate for palmitoylation in rat adipocytes: N-terminal homology with CD 36 and evidence for cell surface acylation. *J. Lipid Res.*, 34 (1993) 1783-1792.
- 704 Kramer, W., Girbig, F., Gutjahr, U., Kowalewski, S., Jouvenal, K., Müller, G., Tripier, D. and Wess, G.: Intestinal bile acid absorption. Na<sup>+</sup>-dependent bile acid transport activity in rabbit small intestine correlates with the coexpression of an integral 93-kDa and a peripheral 14-kDa bile acid-binding membrane protein along the duodenum-ileum axis. *J. Biol. Chem.*, 268 (1993) 18035-18046.
- 705 Kuiper, G.G.J.M., de Ruiter, P.E., Trapman, J., Jenster, G. and Brinkmann, A.O.: *In vitro* translation of androgen receptor cRNA results in an activated androgen receptor protein. *Biochem. J.*, 296 (1993) 161-167.
- 706 Kushwaha, R.S., Hasan, S.Q., McGill, H.C., Jr., Getz, G.S., Dunham, R.G. and Kanda, P.: Characterization of cholesteryl transfer protein inhibitor from plasma of baboons (*Papio* sp.) *J. Lipid Res.*, 34 (1993) 1285-1297.
- 707 Lee, Y.-S., Lee, H.-S., Park, M.-K., Hwang, E.-S., Park, E.-M., Kasai, H. and Chung, M.-H.: Identification of 8-hydroxyguanine glycosylase activity in mammalian tissues using 8-hydroxyguanine specific monoclonal antibody. *Biochem. Biophys. Res. Commun.*, 196 (1993) 1545-1551.
- 708 Moss, A.M. and Livingston, J.N.: Distinct  $\beta$ -subunits are present in hybrid insulin-like-growth-factor-1 receptors in the central nervous system. *Biochem. J.*, 294 (1993) 685-692.
- 709 Nielsen, S.U. and Spener, F.: Fatty acid-binding protein from rat heart is phosphorylated on Tyr<sup>19</sup> in response to insulin stimulation. *J. Lipid Res.*, 34 (1993) 1355-1366.
- 710 Ostergaard, J., Vergnolle, C., Schoentgen, F. and Kader, J.-C.: Acyl-binding/lipid-transfer proteins from rape seedlings, a novel category of proteins interacting with lipids. *Biochim. Biophys. Acta*, 1170 (1993) 109-117.
- 711 Pick, E., Gorzalczy, Y. and Engel, S.: Role of the rac1 p21-GDP-dissociation inhibitor for the heterodimer in the activation of the superoxide-forming NADPH oxidase of macrophages. *Eur. J. Biochem.*, 217 (1993) 441-455.
- 712 Ploug, M., Kjalke, M., Rønne, E., Weidle, U., Høyer-Hansen, G. and Danø, K.: Localization of the disulfide bonds in the NH<sub>2</sub>-terminal domain of the cellular receptor for human urokinase-type plasminogen activator. A domain structure belonging to a novel superfamily of glycolipid-anchored membrane proteins. *J. Biol. Chem.*, 268 (1993) 17539-17546.
- 713 Poletti, A., Conneely, O.M., McDonnell, D.P., Schrader, W.T., O'Malley, B.W. and Weigel, N.L.: Chicken progesterone receptor expressed in *Saccharomyces cerevisiae* is correctly phosphorylated at all four Ser-Pro phosphorylation sites. *Biochemistry*, 32 (1993) 9563-9569.
- 714 Schäfer, H. and Schmidt, W.E.: Characterization and purification of the solubilized pituitary adenylate-cyclase-activating polypeptide-1 receptor from porcine brain using a biotinylated ligand. *Eur. J. Biochem.*, 217 (1993) 823-830.
- 715 Schratzenholz, A., Godovac-Zimmermann, J., Schäfer, H.-J., Albuquerque, E.X. and Maelicke, A.: Photoaffinity labeling of *Torpedo* acetylcholine receptor by physostigmine. *Eur. J. Biochem.*, 216 (1993) 671-677.
- 716 Schultes, B.C., Reinsberg, J., Schlebusch, H., Oehr, P., Biersack, H.-J., Krebs, D. and Wagner, U.: Idiotypic cascades after injection of the monoclonal antibody OC125: a study in a mouse model. Induction of antibodies against OC125 and CA 125 after immunization with an anti-CA 125 (Mab OC125) monoclonal antibody by activation of the idiotypic network. *Eur. J. Clin. Chem. Clin. Biochem.*, 31 (1993) 427-432.
- 717 Srivastava, R.A.K., Bauchmann, D. and Schonfeld, G.: *In vivo* regulation of low-density lipoprotein receptors by estrogen differs at the post-transcriptional level in rat and mouse. *Eur. J. Biochem.*, 216 (1993) 527-538.
- 718 Staten, N.R., Byatt, J.C. and Krivi, G.G.: Ligand-specific dimerization of the extracellular domain of the bovine growth hormone receptor. *J. Biol. Chem.*, 268 (1993) 18467-18473.
- 719 Stuhlmüller, B., Kalden, J.R., Fahibusch, R., Hain, N. and Manger, B.: Immunological and functional properties of the acetylcholine receptor expressed on the human cell line TE671. *Eur. J. Clin. Chem. Clin. Biochem.*, 31 (1993) 657-665.
- 720 Tokumitsu, H., Masuda, E.S., Tsuboi, A., Arai, K.-i. and Arai, N.: Purification of the 120 kDa component of the human nuclear factor of activated T cells (NF-AT): reconstitution of binding activity to the cis-acting element of the GM-CSF and IL-2 promoter with AP-1. *Biochem. Biophys. Res. Commun.*, 196 (1993) 737-744.

For additional information see C.A.:  
119 (1993) 90412p, 112602e.

See also 172, 738.

#### 19m. Urinary proteins

721 Stapleton, A.M.F., Simpson, R.J. and Ryall, R.L.: Crystal matrix protein is related to human prothrombin. *Biochem. Biophys. Res. Commun.*, 195 (1993) 1199-1203.

#### 19n. Other proteins (incl. proteinous inhibitors of enzymic activity)

722 Chatrenet, B. and Chang, J.: The disulfide folding pathway of hirudin elucidated by stop/go folding experiments. *J. Biol. Chem.*, 268 (1993) 20988-20996.

723 Hermans, J.M. and Stone, S.R.: Interaction of activated protein C with serpins. *Biochem. J.*, 295 (1993) 239-245.

724 Teschauer, W.F., Mentele, R. and Sommerhoff, C.P.: Primary structure of a porcine leukocyte serpin. *Eur. J. Biochem.*, 217 (1993) 519-526.

725 Williamson, R.A., Smith, B.J., Angal, S. and Freedman, R.B.: Chemical modification of tissue inhibitor of metalloproteinases-1 and its inactivation by diethyl pyrocarbonate. *Biochim. Biophys. Acta*, 1203 (1993) 147-154.

See also 706.

## 20. ENZYMES AND ENZYME ACTIVITY ESTIMATION

See 603.

## 20a. Oxidoreductases

- 726 Abraham, Z.H.L., Lowe, D.J. and Smith, B.E.: Purification and characterization of the dissimilatory nitrite reductase from *Alcaligenes xylosoxidans* subsp. *xylosoxidans* (N.C.I.M.B. 11015): evidence for the presence of both type 1 and type 2 copper centres. *Biochem. J.*, 295 (1993) 587-593.
- 727 Adachi, H., Iida, S., Oguchi, S., Ohshima, H., Suzuki, H., Nagasaki, K., Kawasaki, H., Sugimura, T. and Esumi, H.: Molecular cloning of a cDNA encoding an inducible calmodulin-dependent nitric-oxide synthase from rat liver and its expression in COS 1 cells. *Eur. J. Biochem.*, 217 (1993) 37-43.
- 728 Ashida, A., Hatakeyama, K. and Kagamiyama, H.: cDNA Cloning, expression in *Escherichia coli* and purification of human 6-pyruvoyl-tetrahydropterin synthase. *Biochem. Biophys. Res. Commun.*, 195 (1993) 1386-1393.
- 729 Bonacker, L.G., Baudner, S., Mörschel, E., Böcher, R. and Thauer, R.K.: Properties of the two isoenzymes of methyl-coenzyme M reductase in *Methanobacterium thermoautotrophicum*. *Eur. J. Biochem.*, 217 (1993) 587-595.
- 730 Britsch, L., Dedio, J., Saedler, H. and Forkmann, G.: Molecular characterization of flavanone 3 $\beta$ -hydroxylases. Consensus sequence, comparison with related enzymes and the role of conserved histidine residues. *Eur. J. Biochem.*, 217 (1993) 745-754.
- 731 Chen, L., Liu, M.-Y., Legall, J., Fareleira, P., Santos, H. and Xavier, A.V.: Purification and characterization of an NADH-rubredoxin oxidoreductase involved in the utilization of oxygen by *Desulfovibrio gigas*. *Eur. J. Biochem.*, 216 (1993) 443-448.
- 732 Galaev, I.Yu. and Mattiasson, B.: Effect of synthetic polymers, poly(N-vinyl pyrrolidone) and poly(N-vinyl caprolactam), on elution of lactate dehydrogenase bound to Blue Sepharose. *J. Chromatogr.*, 648 (1993) 367-371.
- 733 Haupt, K. and Vijayalakshmi, M.A.: Interaction of catechol-2,3-dioxygenase of *Pseudomonas putida* with immobilized histidine and histamine. *J. Chromatogr.*, 644 (1993) 289-297.
- 734 Jimenez-Cervantes, C., Garcia-Borron, J., Valverde, P., Solano, F. and Lozano, J.A.: Tyrosinase isoenzyme in mammalian melanocytes. 1. Biochemical characterization of two melanosomal tyrosinases from B16 mouse melanoma. *Eur. J. Biochem.*, 217 (1993) 549-556.
- 735 Jones, A.M. and Hollocher, T.C.: Nitric oxide reductase of *Achromobacter cycloclastes*. *Biochim. Biophys. Acta*, 1144 (1993) 359-366.
- 736 Jouan, A., Pilloud-Dagher, M.-C., Fuchs, A. and Vignais, P.V.: A generally applicable ELISA for the detection and quantitation of the cytosolic factors of NADPH-oxidase activation in neutrophils. *Anal. Biochem.*, 214 (1993) 252-259.
- 737 Kalisz, H.M. and Schmid, R.D.: Separation of glucose oxidase isozymes from *Penicillium amagasakiense* by ion-exchange chromatography. *ACS Symp. Ser.*, 529(Chromatography in Biotechnology) (1993) 102-111; C.A., 119 (1993) 111903s.
- 738 Klinger, E., Sharabani, M. and Aviram, I.: The interaction of cytosolic components of neutrophil NADPH oxidase with phosphoinositides. *Biochem. J.*, 295 (1993) 565-570.
- 739 LoGrasso, P.V., Soltis, D.A. and Boettcher, B.R.: Overexpression, purification, and kinetic characterization of a carboxyl-terminal-truncated yeast squalene synthetase. *Arch. Biochem. Biophys.*, 307 (1993) 193-199.
- 740 Miranda, C.L., Collodi, P., Zhao, X., Barnes, D.W. and Buhler, D.R.: Regulation of cytochrome P450 expression in a novel liver cell line from Zebrafish (*Brachydanio rerio*). *Arch. Biochem. Biophys.*, 305 (1993) 320-327.
- 741 Nordling, M., Elmgren, M., Stahlberg, J., Pettersson, G. and Lindquist, S.-E.: A combined cellobiose oxidase/glucose oxidase biosensor for HPLC determination on-line of glucose and soluble cyclodextrines. *Anal. Biochem.*, 214 (1993) 389-396.
- 742 Ohashi, Y., Katayama, M., Hirata, K.-i., Suematsu, M., Kawashima, S. and Yokoyama, M.: Activation of nitric oxide synthase from cultured aortic endothelial cells by phospholipids. *Biochem. Biophys. Res. Commun.*, 195 (1993) 1314-1320.
- 743 Ohishi, N., Imaoka, S., Suzuki, T. and Funae, Y.: Characterization of two P-450 isozymes placed in the rat CYP2D subfamily. *Biochim. Biophys. Acta*, 1158 (1993) 227-236.
- 744 Ohmori, S., Horie, T., Guengerich, F.P., Kiuchi, M. and Kitada, M.: Purification and characterization of two forms of hepatic microsomal cytochrome P450 from untreated cynomolgus monkeys. *Arch. Biochem. Biophys.*, 305 (1993) 405-413.
- 745 Otto, A. and Birkenmeier, G.: Recognition and separation of isoenzymes by metal chelates. Immobilized metal ion affinity partitioning of lactate dehydrogenase isoenzymes. *J. Chromatogr.*, 644 (1993) 25-33.
- 746 Rajas, F. and Rousset, B.: A membrane-bound form of glutamate dehydrogenase possesses an ATP-dependent high-affinity microtubule-binding activity. *Biochem. J.*, 295 (1993) 447-455.
- 747 Reng, W., Riessland, R., Scheibe, R. and Jaenicke, R.: Cloning, site-specific mutagenesis, expression and characterization of full-length chloroplast NADP-malate dehydrogenase from *Pisum sativum*. *Eur. J. Biochem.*, 217 (1993) 189-197.
- 748 Rescigno, A., Sanjust, E., Curreli, N., Padiglia, A. and Floris, G.: Improved chromatographic purification of peroxidase and  $\beta$ -glucosidase from *Hordeum vulgare* seedlings. *Prepar. Biochem.*, 23 (1993) 485-492.
- 749 Romano, M., Chen, X.-S., Takahashi, Y., Yamamoto, S., Funk, C.D. and Serhan, C.N.: Lipoxin synthase activity of human platelet 12-lipoxygenase. *Biochem. J.*, 296 (1993) 127-133.
- 750 Sandhu, P., Baba, T. and Guengerich, F.P.: Expression of modified cytochrome P450 2C10 (2C9) in *Escherichia coli*, purification, and reconstitution of catalytic activity. *Arch. Biochem. Biophys.*, 306 (1993) 443-450.
- 751 Sandstrom, J., Karlsson, K., Edlund, T. and Marklund, S.L.: Heparin-affinity patterns and composition of extracellular superoxide dismutase in human plasma and tissues. *Biochem. J.*, 294 (1993) 853-857.
- 752 Sillard, R., Jörnvall, H. and Mutt, V.: Characterization of porcine intestinal cytochrome c oxidase subunit VIc, purified by affinity chromatography. *Biochem. Biophys. Res. Commun.*, 195 (1993) 746-750.
- 753 Slabaugh, M.B., Davis, R.E., Roseman, N.A. and Mathews, C.K.: Vaccinia virus ribonucleotide reductase expression and isolation of the recombinant large subunit. *J. Biol. Chem.*, 268 (1993) 17803-17810.

- 754 Takahashi, Y., Glasgow, W.C., Suzuki, H., Taketani, Y., Yamamoto, S., Anton, M., Kühn, H. and Brash, A.R.: Investigation of the oxygenation of phospholipids by the porcine leukocyte and human platelet arachidonate 12-lipoxygenases. *Eur. J. Biochem.*, 218 (1993) 165-171.
- 755 Vestling, M.M., Murphy, C.M., Keller, D.A., Fenselau, C., Dedinas, J., Ladd, D.L. and Olsen, M.A.: A strategy for characterization of polyethylene glycol-derivatized proteins. A mass-spectrometric analysis of the attachment sites of polyethylene glycol-derivatized superoxide dismutase. *Drug Metab. Disp.*, 21 (1993) 911-917.
- 756 Vianello, F., Di Paolo, M.L., Stevanato, R., Gasparini, R. and Rigo, A.: Purification and characterization of amine oxidase from soybean seedlings. *Arch. Biochem. Biophys.*, 307 (1993) 35-39.
- 757 Wimmerova, M., Glatz, Z., Janiczek, O. and Macholan, L.: Improved chromatographic purification of pea seedlings diamine oxidase. *Prep. Biochem.*, 23 (1993) 303-319; *C.A.*, 119 (1993) 134066k.
- 758 Yamamoto, T., Moriwaki, Y., Takahashi, S., Nasako, Y., Hiroishi, K. and Higashino, K.: Determination of erythrocyte aldehyde dehydrogenase activity by high-performance liquid chromatography. *Anal. Biochem.*, 215 (1993) 129-133.
- 759 Yong, H., Thomas, G.A. and Peticolas, W.L.: Metabolite-modulated complex formation between  $\alpha$ -glycerophosphate dehydrogenase and lactate dehydrogenase. *Biochemistry*, 32 (1993) 11124-1131.
- For additional information see *C.A.*:  
119 (1993) 155147c.
- See also 565.
- 20b. *Transferases (excl. E.C. 2.7.-.-)*
- 760 Bendiak, B., Ward, L.D. and Simpson, R.J.: Proteins of the Golgi apparatus. Purification to homogeneity, N-terminal sequence, and unusually large Stokes radius of the membrane-bound form of UDP-galactose:N-acetylglucosamine  $\beta$ 1-4-galactosyltransferase from rat liver. *Eur. J. Biochem.*, 216 (1993) 405-417.
- 761 Chen, A., Zhang, D. and Poulter, C.D.: (S)-Geranylgeranyl glyceryl phosphate synthase. Purification and characterization of the first pathway-specific enzyme in archaeobacterial membrane lipid biosynthesis. *J. Biol. Chem.*, 268 (1993) 21701-21705.
- 762 De Vries, T., Norberg, T., Lönn, H. and van den Eijnden, D.H.: The use of human milk fucosyltransferase in the synthesis of human-associated trimeric X determinants. *Eur. J. Biochem.*, 216 (1993) 769-777.
- 763 Derbel, M., Igarashi, T. and Satoh, T.: Differential induction of glutathione S-transferase subunits by phenobarbital, 3-methylcholanthrene and ethoxyquin in rat liver and kidney. *Biochim. Biophys. Acta*, 1158 (1993) 175-180.
- 764 Dieckmann-Schuppert, A., Bause, E. and Schwarz, R.T.: Studies on O-glycans of *Plasmodium falciparum*-infected human erythrocytes. Evidence for O-GlcNAc and O-GlcNAc-transferase in malaria parasites. *Eur. J. Biochem.*, 216 (1993) 779-788.
- 765 Habuchi, O., Matsui, Y., Kotoya, Y., Aoyama, Y., Yasuda, Y. and Noda, M.: Purification of chondroitin 6-sulfotransferase secreted from cultured chick embryo chondrocytes. *J. Biol. Chem.*, 268 (1993) 21968-21974.
- 766 Hagen, F.K., VanWuyckhuysse, B. and Tabak, L.A.: Purification, cloning, and expression of a bovine UDP-GalNAc:polypeptide N-acetyl-galactosaminyltransferase. *J. Biol. Chem.*, 268 (1993) 18960-18965.
- 767 Hussey, A.J. and Hayes, J.D.: Human Mu-class glutathione S-transferases present in liver, skeletal muscle and testicular tissue. *Biochim. Biophys. Acta*, 1203 (1993) 131-141.
- 768 Lee, Y.-C., Kurosawa, N., Hamamoto, T., Nakaoka, T. and Tsujii, S.: Molecular cloning and expression of Gal $\beta$ 1,3GalNAc $\alpha$ 2,3-sialyltransferase from mouse brain. *Eur. J. Biochem.*, 216 (1993) 377-385.
- 769 Liu, L.-F., Hong, J.-L., Tsai, S.-P., Hsieh, J.-C. and Tam, M.F.: Reversible modification of rat liver glutathione S-transferase 3-3 with 1-chloro-2,4-dinitrobenzene: specific labelling of Tyr-115. *Biochem. J.*, 296 (1993) 189-197.
- 770 Londesborough, J. and Vuorio, O.E.: Purification of trehalose synthase from baker's yeast. Its temperature-dependent activation by fructose 6-phosphate and inhibition by phosphate. *Eur. J. Biochem.*, 216 (1993) 841-848.
- 771 Mudgett, M.B. and Clarke, S.: Characterization of plant L-isospartyl methyltransferases that may be involved in seed survival: purification, cloning, and sequence analysis of the wheat germ enzyme. *Biochemistry*, 32 (1993) 11100-11111.
- 772 Petrarulo, M., Pellegrino, S., Marangella, M., Cosseddu, D. and Linari, F.: High-performance liquid chromatographic microassay for L-glutamate:glyoxylate aminotransferase activity in human liver. Application in primary hyperoxaluria type I. *Clin. Chim. Acta*, 218 (1993) 193-200.
- 773 Reed, D.W., Davin, L., Jain, J.C., Deluca, V., Nelson, L. and Underhill, E.W.: Purification and properties of UDP-glucose:thiohydroxamate glucosyltransferase from *Brassica napus* L. seedlings. *Arch. Biochem. Biophys.*, 305 (1993) 526-532.
- 774 Taha, T.S.M. and Deits, T.L.: Detection of glycineamide ribonucleotide by HPLC with pulsed amperometry: application to the assay for glutamine:5-phosphoribosyl-1-pyrophosphate amidotransferase (E.C. 2.4.2.14). *Anal. Biochem.*, 213 (1993) 323-328.
- 775 Tassin, J.P., Boutin, J.A., Ernould, A.P. and Atassi, G.: (Farnesyl transferase as a target for non-cytotoxic anticancer agents: First steps). *C.R. Seances Soc. Biol. Ses Fil.*, 185 (1991) 306-311; *C.A.*, 119 (1993) 108490z.
- 776 Vessels, J.M. and Radding, J.A.: Oligosaccharide mapping of fungal glucan synthase product by high-performance anion-exchange chromatography. *Anal. Biochem.*, 215 (1993) 150-155.
- 777 Vuorio, O.E., Kalkkinen, N. and Londesborough, J.: Cloning of the related genes encoding the 56-kDa and 123-kDa subunits of trehalose synthase from the yeast *Saccharomyces cerevisiae*. *Eur. J. Biochem.*, 216 (1993) 849-861.
- 778 Whitaker, D.P. and Goosey, M.W.: Purification and properties of the enzyme arylamine N-acetyltransferase from the housefly *Musca domestica*. *Biochem. J.*, 295 (1993) 149-154.
- 779 Wiedemann, M., Misawa, N. and Sandmann, G.: Purification and enzymatic characterization of the geranylgeranyl pyrophosphate synthase from *Erwinia uredovora* after expression in *Escherichia coli*. *Arch. Biochem. Biophys.*, 306 (1993) 152-157.

- 20c. *Transferases transferring phosphorus containing groups (E.C. 2.7.-.-)*
- 780 Bhikhabhai, R., Carlsson, T., Unge, T., Lövgren, S. and Strandberg, B.: Increased yield of homogeneous HIV-1 reverse transcriptase (p66/p51) using a slow purification approach. *J. Chromatogr.*, 639 (1993) 67-74.
- 781 Budde, R.J.A., Ramdas, L. and Ke, S.: Recombinant pp60<sup>c-src</sup> from baculovirus-infected insect cells: purification and characterization. *Prepar. Biochem.*, 23 (1993) 493-515.
- 782 Burrows, J.A. and Goward, C.R.: Preparation of DNA polymerase from *Bacillus caldotenax*. *J. Chromatogr.*, 639 (1993) 75-80.
- 783 Cutler, R.E., Jr., Maizels, E.T., Brooks, E.J., Mizuno, K., Ohno, S. and Hunzicker-Dunn, M.: Regulation of  $\delta$  protein kinase C during rat ovarian differentiation. *Biochim. Biophys. Acta*, 1179 (1993) 260-270.
- 784 Garcia-Paramio, P., Carmena, M.J., Gutierrez-Ocana, M.T., Recio, M.N. and Prieto, J.C.: Alteration of protein kinase C activity in diabetic rat prostate. *Biochem. Biophys. Res. Commun.*, 195 (1993) 166-172.
- 785 Komatsu, H. and Ikebe, M.: Affinity labelling of smooth-muscle myosin light-chain kinase with 5' [*p*-(fluorosulphonyl)benzoyl]adenosine. *Biochem. J.*, 296 (1993) 53-58.
- 786 Kopperschläger, G., Bär, J. and Stellwagen, E.: Limited proteolysis of yeast phosphofructokinase. Sequence location of cleavage sites created by the actions of different proteinases. *Eur. J. Biochem.*, 217 (1993) 527-533.
- 787 Maga, G., Verri, A., Bonizzi, L., Ponti, W., Poli, G., Garbesi, A., Niccola, D., Spadari, S. and Focher, F.: Lack of stereospecificity of suid pseudorabies virus thymidine kinase. *Biochem. J.*, 294 (1993) 381-385.
- 788 Mizoguchi, N., Eguchi, T., Sakura, N. and Ueda, K.: Erythrocyte galactokinase assay with high performance liquid chromatography. *Clin. Chim. Acta*, 216 (1993) 145-151.
- 789 Offermanns, S., Bombien, E. and Schultz, G.: Stimulation of tyrosine phosphorylation and mitogen-activated-protein (MAP) kinase activity in human SH-SY5Y neuroblastoma cells by carbachol. *Biochem. J.*, 294 (1993) 545-550.
- 790 Ohmichi, M., Zhu, G. and Saltiel, A.R.: Nerve growth factor activates calcium-insensitive protein kinase C- $\epsilon$  in PC-12 rat pheochromocytoma cells. *Biochem. J.*, 295 (1993) 767-772.
- 791 Sutherland, C., Alterio, J., Campbell, D.G., le Bourdellés, B., Mallet, J., Haavik, J. and Cohen, P.: Phosphorylation and activation of human tyrosine hydroxylase *in vitro* by mitogen-activated protein (MAP) kinase and MAP-kinase activated kinases 1 and 2. *Eur. J. Biochem.*, 217 (1993) 715-722.
- 792 Sutherland, C., Leighton, I.A. and Cohen, P.: Inactivation of glycogen synthase kinase-3 $\beta$  by phosphorylation: new kinase connections in insulin and growth-factor signalling. *Biochem. J.*, 296 (1993) 15-19.
- 793 Szumilo, T., Drake, R.R., York, J.L. and Elbein, A.D.: GDP-Mannose pyrophosphorylase. Purification to homogeneity, properties, and utilization to prepare photoaffinity analogs. *J. Biol. Chem.*, 268 (1993) 17943-17950.
- 794 Vandercammen, A. and van Schaftingen, E.: Species and tissue distribution of the regulatory protein of glucokinase. *Biochem. J.*, 294 (1993) 551-556.
- 795 Wang, L.-M., Kucera, G.L. and Capizzi, R.L.: Purification and characterization of deoxycytidine kinase from acute myeloid leukemia cell mitochondria. *Biochim. Biophys. Acta*, 1202 (1993) 309-316.
- 796 Wei, T. and Tao, M.: Human erythrocyte casein kinase II: characterization and phosphorylation of membrane cytoskeletal proteins. *Arch. Biochem. Biophys.*, 307 (1993) 206-216.
- 797 Wood, S.L., Emmison, N., Borthwick, A.C. and Yeaman, S.J.: The protein phosphatases responsible for dephosphorylation of hormone-sensitive lipase in isolated rat adipocytes. *Biochem. J.*, 295 (1993) 531-535.
- 798 Zimmerly, S., Drains, D., Sylvers, L.A. and Söll, D.: Identification of a 100-kDa protein associated with nuclear ribonuclease P activity in *Schizosaccharomyces pombe*. *Eur. J. Biochem.*, 217 (1993) 501-507.
- For additional information see C.A.:  
119 (1993) 134014s.
- See also 576.
- 20d. *Hydrolases, acting on ester bonds (E.C. 3.1.-.-)*
- 799 Belyakova, N.V., Kleiner, N.E., Kravetskaya, T.P., Legina, O.K., Naryzhny, S.N., Perrino, F.W., Shevelev, I.V. and Krutyakov, V.M.: Proof-reading 3'→5' exonucleases isolated from rat liver nuclein. *Eur. J. Biochem.*, 217 (1993) 493-500.
- 800 Bublitz, R., Armesto, J., Hoffmann-Blume, E., Schulze, M., Rhode, H., Horn, A., Aulwurm, S., Hannappel, E. and Fischer, W.: Heterogeneity of glycosylphosphatidylinositol-anchored alkaline phosphatase of calf intestine. *Eur. J. Biochem.*, 217 (1993) 199-207.
- 801 Chang, L.-S., Kuo, K.-W. and Chang, C.-C.: Identification of functional involvement of tryptophan residues in phospholipase A<sub>2</sub> from *Naja naja atra* (Taiwan cobra) snake venom. *Biochim. Biophys. Acta*, 1202 (1993) 216-220.
- 802 De Carvalho, M.S., McCormack, F.X. and Leslie, C.C.: The 85-kDa, arachidonic acid-specific phospholipase A<sub>2</sub> is expressed as an activated phosphoprotein in Sf9 cells. *Arch. Biochem. Biophys.*, 306 (1993) 534-540.
- 803 Dichek, H.L., Parrott, C., Ronan, R., Brunzell, J.D., Brewer, H.B., Jr. and Santamarina-Fojo, S.: Functional characterization of a chimeric lipase genetically engineered from human lipoprotein lipase and human hepatic lipase. *J. Lipid Res.*, 34 (1993) 1393-1401.
- 804 Gimble, F.S. and Thorner, J.: Purification and characterization of VDE, a site-specific endonuclease from the yeast *Saccharomyces cerevisiae*. *J. Biol. Chem.*, 268 (1993) 21844-21853.
- 805 Hattori, M., Arai, H. and Inoue, K.: Purification and characterization of bovine brain platelet-activating factor acetylhydrolase. *J. Biol. Chem.*, 268 (1993) 18748-18753.
- 806 Karunaratne, S.H.P.P., Jayawardena, K.G.I., Hemingway, J. and Ketterman, A.J.: Characterization of a B-type esterase involved in insecticide resistance from the mosquito *Culex quinquefasciatus*. *Biochem. J.*, 294 (1993) 575-579.

- 807 Komoda, T., Koyama, I., Fujimori-Arai, Y., Yamagishi, F., Sakagishi, Y. and Hirota, N.: Lectin affinity chromatography as a useful tool for determining the heterogeneity of alkaline phosphatases - intestinal-like enzymes. *Int. Congr. Ser. - Excerpta Med.*, 991 (Progress in Clinical Biochemistry) (1992) 345-349; *C.A.*, 119 (1993) 111826u.
- 808 Laboure, A.-M., Gagnon, J. and Lescure, A.-M.: Purification and characterization of a phytase (myo-inositol-hexakisphosphate phosphohydrolase) accumulated in maize (*Zea mays*) seedlings during germination. *Biochem. J.*, 295 (1993) 413-419.
- 809 Lee, C., Park, D.J., Lee, K., Kim, C.G. and Rhee, S.G.: Purification, molecular cloning, and sequencing of phospholipase C- $\beta$ 4. *J. Biol. Chem.*, 268 (1993) 21318-21327.
- 810 Lim, Y.-M. and Chung, M.C.M.: Isolation and characterization of pectin methylesterase from papaya. *Arch. Biochem. Biophys.*, 307 (1993) 15-20.
- 811 Mehra, R.S., Malhotra, V.P. and Rembhotkar, G.W.: Rapid purification of a restriction endonuclease from *Escherichia coli* RY13. *Biotechnol. Tech.*, 7 (1993) 411-414; *C.A.*, 119 (1993) 89673t.
- 812 Minami, T., Tojo, H., Shinomura, Y., Matsuzawa, Y. and Okamoto, M.: Purification and characterization of a phospholipase A<sub>2</sub> from human ileal mucosa. *Biochim. Biophys. Acta*, 1170 (1993) 125-130.
- 813 Mukherjee, J.J., Jay, F.T. and Choy, P.C.: Purification, characterization and modulation of a microsomal carboxylesterase in rat liver for the hydrolysis of acyl-CoA. *Biochem. J.*, 295 (1993) 81-86.
- 814 Robert, S., Domurado, D., Thomas, D. and Chopineau, J.: Fatty acid acylation of RNase A using reversed micelles as microreactors. *Biochem. Biophys. Res. Commun.*, 196 (1993) 447-454.
- 815 Sakamoto, W., Fujie, K., Nishihira, J. and Handa, H.: Effect of vitamin E on arachidonic acid-release in rat peritoneal macrophages. *Biochim. Biophys. Acta*, 1170 (1993) 296-300.
- 816 Sugo, T., Mas, E., Abouakil, N., Endo, T., Escribano, M.-J., Kobata, A. and Lombardo, D.: The structure of N-linked oligosaccharides of human pancreatic bile-salt-dependent lipase. *Eur. J. Biochem.*, 216 (1993) 799-805.
- 817 Thomas, T.C., Szekacz, A., Hammock, B.D., Wilson, B.W. and McNamee, M.G.: Affinity chromatography of neuropathy target esterase. *Chem.-Biol. Interact.*, 87 (1993) 347-360; *C.A.*, 119 (1993) 176415d.
- 818 Tokumoto, H., Croxtall, J.D., Choudhury, Q. and Flower, R.J.: Phospholipase A<sub>2</sub>-induced stimulation of A549 lung adenocarcinoma cell line proliferation. *Biochim. Biophys. Acta*, 1169 (1993) 236-242.
- 819 Tsujita, T. and Okuda, H.: Palmitoyl-coenzyme A hydrolyzing activity in rat kidney and its relationship to carboxylesterase. *J. Lipid Res.*, 34 (1993) 1773-1781.
- 820 Ueda, H., Kobayashi, T., Kishimoto, M., Tsutsumi, T., Watanabe, S. and Okuyama, H.: The presence of Ca<sup>2+</sup>-independent phospholipase A<sub>1</sub> highly specific for phosphatidylinositol in bovine brain. *Biochem. Biophys. Res. Commun.*, 195 (1993) 1272-1279.
- 821 Villela, E., Joven, J., Fernández, M., Villaro, S., Brunzell, J.D., Olivecrona, T. and Bengtsson-Olivecrona, G.: Lipoprotein lipase in human plasma is mainly inactive and associated with cholesterol-rich lipoproteins. *J. Lipid Res.*, 34 (1993) 1555-1564.
- 20e. *Hydrolases, acting on glycosyl compounds (E.C. 3.2.-.-)*
- 822 Afify, A.M., Stern, M., Guntenhöner, M. and Stern, R.: Purification and characterization of human serum hyaluronidase. *Arch. Biochem. Biophys.*, 305 (1993) 434-441.
- 823 Ajongwen, J.N., Akintoye, A., Barker, P.E., Ganetsos, G. and Shieh, M.T.: Large-scale purification of Leuconostoc mesenteroides NRRL B512F dextransucrase for use in the biosynthesis of dextran by batch and continuous chromatography. *Chem. Eng. J. (Lausanne)*, 51 (1993) B43-B50; *C.A.*, 119 (1993) 115380s.
- 824 Chen, Y.Q. and Wenger, D.A.: Galactocerebrosidase from human urine: purification and partial characterization. *Biochim. Biophys. Acta*, 1170 (1993) 53-61.
- 825 Evans, B.R., Margalit, R. and Woodward, J.: Attachment of pentaamineruthenium(III) to *Trichoderma reesei* cellobiohydrolase I increases its catalytic activity. *Biochem. Biophys. Res. Commun.*, 195 (1993) 497-503.
- 826 Fierobe, H.-P., Bagnara-Tardif, C., Gaudin, C., Guerlesquin, F., Sauve, P., Belaich, A. and Belaich, J.-P.: Purification and characterization of endoglucanase C from *Clostridium cellulolyticum*; catalytic comparison with endoglucanase A. *Eur. J. Biochem.*, 217 (1993) 557-565.
- 827 Forsyth, G.W., Romero, K.M., Alverson, J., VanderJagt, D.J. and Glew, R.H.: Variable expression of leukocyte cytosolic broad-specificity  $\beta$ -glucosidase activity. *Clin. Chim. Acta*, 216 (1993) 11-21.
- 828 Garcia-Campayo, V., McCrae, S.I., Zhang, J.-X., Flint, H.J. and Wood, T.M.: Mode of action, kinetic properties and physicochemical characterization of two different domains of a bifunctional (142- $\beta$ -D-xylanase from *Ruminococcus flavefaciens* expressed separately in *Escherichia coli*. *Biochem. J.*, 296 (1993) 235-243.
- 829 Mehran, H. and Storey, K.B.: Characterization of  $\alpha$ -glucosidase from rainbow trout liver. *Arch. Biochem. Biophys.*, 306 (1993) 188-194.
- See also 748, 849.
- 20f. *Other hydrolases*
- 830 Aggeler, R., Cai, S.X., Keana, J.F.W., Koike, T. and Capaldi, R.A.: The  $\gamma$  subunit of the *Escherichia coli* F<sub>1</sub>-ATPase can be cross-linked near the glycine-rich loop region of a  $\beta$  subunit when ADP + Mg<sup>2+</sup> occupies catalytic sites but not when ATP + Mg<sup>2+</sup> is bound. *J. Biol. Chem.*, 268 (1993) 20831-20837.
- 831 Blache, P., Kervran, A., Le-Nguyen, G., Dufour, M., Cohen-Solal, A., Duckworth, W. and Bataille, D.: Endopeptidase from rat liver membranes, which generates miniglucagon from glucagon. *J. Biol. Chem.*, 268 (1993) 21748-21753.
- 832 Crawford, C., Brown, N.R. and Willis, A.C.: Studies of the active site of m-calpain and the interaction with calpastatin. *Biochem. J.*, 296 (1993) 135-142.
- 833 Dando, P.M., Brown, M.A. and Barrett, A.J.: Human thimet oligopeptidase. *Biochem. J.*, 294 (1993) 451-457.
- 834 Freeman, S.-A., Peek, K., Prescott, M. and Daniel, R.: Characterization of a chelator-resistant proteinase from *Thermus* strain Rt4A2. *Biochem. J.*, 295 (1993) 463-469.
- 835 Hrycyna, C.A. and Clarke, S.: Purification and characterization of a novel metalloendopeptidase from *Saccharomyces cerevisiae*. *Biochemistry*, 32 (1993) 11293-11301.

- 836 Klein, J.R., Klein, U., Schad, M. and Plapp, R.: Cloning, DNA sequence analysis and partial characterization of pepN, a lysyl aminopeptidase from *Lactobacillus delbrückii* ssp. *lactis* DSM7290. *Eur. J. Biochem.*, 217 (1993) 105-114.
- 837 Kobayashi, M., Komeda, H., Nagasawa, T., Nishiyama, M., Hori-nouchi, S., Beppu, T., Yamada, H. and Shimizu, S.: Amidase coupled with low-molecular-mass nitrile hydratase from *Rhodococcus rhodochrous* U1. Sequencing and expression of the gene and purification and characterization of the gene product. *Eur. J. Biochem.*, 217 (1993) 327-336.
- 838 Kolkenbrock, H., Hecker-Kia, A., Orgel, D., Buchlow, G., Sörensen, H., Hauer, W. and Ulbrich, N.: A trypsin sensitive stromelysin isolated from rheumatoid synovial fluid is an activator for matrix metalloproteinases. *Eur. J. Clin. Chem. Clin. Biochem.*, 31 (1993) 625-631.
- 839 Lenfant, F., Petit, A., Labia, R., Maveyraud, L., Samama, J.-P. and Masson, J.-M.: Site-directed mutagenesis of  $\beta$ -lactamase TEM-1. Investigating the potential role of specific residues on the activity of *Pseudomonas*-specific enzyme. *Eur. J. Biochem.*, 217 (1993) 939-946.
- 840 Liau, D.F., Yin, N.X. and Ryan, S.F.: Isolation of human polymorphonuclear leukocyte elastase by chromatography on immobilized benzamidine. *Prepar. Biochem.*, 23 (1993) 439-447.
- 841 Lutter, R., Saraste, M., van Walraven, H.S., Runswick, M.J., Finel, M., Deatherage, J.F. and Walker, J.E.:  $F_1F_0$ -ATP synthase from bovine heart mitochondria: development of the purification of a monodisperse oligomycin-sensitive ATPase. *Biochem. J.*, 295 (1993) 799-806.
- 842 Mayer, U., Mann, K., Timpl, R. and Murphy, G.: Sites of nitrogen cleavage of proteases involved in tissue homeostasis and remodelling. *Eur. J. Biochem.*, 217 (1993) 877-884.
- 843 Miller, D.K., Ayala, J.M., Egger, L.A., Raju, S.M., Yamin, T., Ding, J.-F., Gaffney, E.P., Howard, A.D., Palyha, O.C., Rolando, A.M., Salley, J.P. et al.: Purification and characterization of active human interleukin-1 $\beta$ -converting enzyme from THP.1 monocytic cells. *J. Biol. Chem.*, 268 (1993) 18062-18069.
- 844 Nielsen, F.S. and Foltmann, B.: Activation of porcine pepsinogen A. The stability of two non-covalent activation intermediates at pH 8.5. *Eur. J. Biochem.*, 217 (1993) 137-142.
- 845 Page, A.E., Fuller, K., Chambers, T.J. and Warburton, M.J.: Purification and characterization of a tripeptidyl peptidase I from human osteoclastomas: evidence for its role in bone resorption. *Arch. Biochem. Biophys.*, 306 (1993) 354-359.
- 846 Page, M.G.P.: The kinetics of non-stoichiometric bursts of  $\beta$ -lactam hydrolysis catalysed by class C  $\beta$ -lactamases. *Biochem. J.*, 295 (1993) 295-304.
- 847 Perkins, S.J., Smith, K.F. and Sim, R.B.: Molecular modelling of the domain structure of factor I of human complement by X-ray and neutron solution scattering. *Biochem. J.*, 295 (1993) 101-108.
- 848 Peters, K., Fittkau, S., Steinert, A. and Ströhl, D.: Peptidyl methyl ketones as ligands in affinity chromatography of serine and cysteine proteinases. *J. Chromatogr.*, 648 (1993) 91-99.
- 849 Pshezhetsky, A.V. and Potier, M.: Stoichiometry of the human lysosomal carboxypeptidase- $\beta$ -galactosidase complex. *Biochem. Biophys. Res. Commun.*, 195 (1993) 354-362.
- 850 Sharma, K.K. and Ortwerth, B.J.: Bovine lens acylpeptide hydrolase. Purification and characterization of a tetrameric enzyme resistant to urea denaturation and proteolytic inactivation. *Eur. J. Biochem.*, 216 (1993) 631-637.
- 851 Su, Q. and Boschetti, A.: Partial purification and properties of enzymes involved in the processing of a chloroplast import protein from *Chlamydomonas reinhardtii*. *Eur. J. Biochem.*, 217 (1993) 1039-1047.
- 852 Tal, D.M., Goldshleger, R. and Karlish, S.J.D.: Separation of membrane-embedded tryptic peptides of Na,K-ATPase by size-exclusion chromatography. *J. Chromatogr.*, 646 (1993) 153-157.
- 853 Tsubuki, S., Kawasaki, H., Saito, Y., Miyashita, N., Inomata, M. and Kawashima, S.: Purification and characterization of a Z-Leu-Leu-MCA degrading protease expected to regulate neurite formation: a novel catalytic activity in proteasome. *Biochem. Biophys. Res. Commun.*, 196 (1993) 1195-1201.
- 854 Utaisincharoen, P., Mackessy, S.P., Miller, R.A. and Tu, A.T.: Complete primary structure and biochemical properties of gila-toxin, a serine protease with kallikrein-like and angiotensin-degrading activities. *J. Biol. Chem.*, 268 (1993) 21975-21983.
- 855 Van Dongen, W.D., Versluis, C., van Wassenaar, P.D., de Koster, C.G., Heerma, W. and Haverkamp, J.: Rapid analysis of enzymatic digests of a bacterial protease of the subtilisin type and a "bio-engineered" variant by high-performance liquid chromatography-frit fast atom bombardment mass spectrometry. *J. Chromatogr.*, 647 (1993) 301-309.
- 856 Williams, C.H., Yamamoto, T., Walsh, D.M. and Allsop, D.: Endopeptidase 3.4.24.11 converts N-1-(R,S)carboxy-3-phenylpropyl-Ala-Ala-Phe-p-carboxyanilide into a potent inhibitor of angiotensin-converting enzyme. *Biochem. J.*, 294 (1993) 681-684.
- 857 Yamagata, K., Daiho, T. and Kanazawa, T.: Labeling of lysine 492 with pyridoxal 5'-phosphate in the sarcoplasmic reticulum  $Ca^{2+}$ -ATPase. Lysine 492 residue is located outside the fluorescent 5-isothiocyanate-binding region in or near the ATP binding site. *J. Biol. Chem.*, 268 (1993) 20930-20936.

See also 175, 577, 728.

#### 20g. Lyases

- 858 Bourguignon, J., Vauclare, P., Merand, V., Forest, E., Neuburger, M. and Douce, R.: Glycine decarboxylase complex from higher plants. Molecular cloning, tissue distribution and mass spectrometry analyses of the T protein. *Eur. J. Biochem.*, 217 (1993) 377-386.
- 859 De Felice, C., Hayakawa, K., Watanabe, T. and Tanaka, T.: High-performance gel-permeation chromatographic analysis of protein aggregation. Application to bovine carbonic anhydrase. *J. Chromatogr.*, 645 (1993) 101-105.
- 860 Desai, U.R., Wang, H.-m. and Linhardt, R.J.: Substrate specificity of the heparin lyases from *Flavobacterium heparinum*. *Arch. Biochem. Biophys.*, 306 (1993) 461-468.
- 861 Hugo, E.R. and Byers, T.J.: S-Adenosyl-L-methionine decarboxylase of *Acanthamoeba castellanii* (Neff): purification and properties. *Biochem. J.*, 295 (1993) 203-209.
- 862 Maila, L.H., Siivari, K.M., Mäkelä, M.J., Jalonen, J.E., Latipää, P.M., Kunau, W.-H. and Hiltunen, J.K.: Enzymes converting D-3-hydroxyacetyl-CoA to trans-2-enoyl-CoA. Microsomal and peroxisomal isoenzymes in rat liver. *J. Biol. Chem.*, 268 (1993) 21578-21585.

- 863 Martinou, A., Kafetzopoulos, D. and Bouriotis, V.: Isolation of chitin deacetylase from *Mucor rouxii* by immunoaffinity chromatography. *J. Chromatogr.*, 644 (1993) 35-41.
- 864 Moore, J.D., Hawkins, A.R., Charles, I.G., Deka, R., Coggins, J.R., Cooper, A., Kelly, S.M. and Price, N.C.: Characterization of the type I dehydroquinase from *Salmonella typhi*. *Biochem. J.*, 295 (1993) 277-285.
- See also 837, 865.
- 20h. *Isomerases*
- 865 Roper, D.I. and Cooper, R.A.: Purification, nucleotide sequence and some properties of a bifunctional isomerase/decarboxylase from the homoprotocatechuate degradation pathway of *Escherichia coli* C. *Eur. J. Biochem.*, 217 (1993) 575-580.
- 866 Schaaf, P.M.M., Heide, L.E., Leistner, E.W., Tani, Y. and El-Olemy, M.M.: Immobilization of isochorismate hydroxymutase. Comparison of native versus immobilized enzyme. *J. Natural Prod.*, 56 (1993) 1304-1312.
- 867 Schaaf, P.M.M., Heide, L.E., Leistner, E.W., Tani, Y., Karas, M. and Deutzmann, R.: Properties of isochorismate hydroxymutase from flavobacterium K<sub>3-15</sub>. *J. Natural Prod.*, 56 (1993) 1294-1303.
- 20i. *Ligases*
- 868 Menguito, C.A., Papaconstantinou, J. and Weigel, P.H.: The histidyl-tRNA synthetase from *Streptococcus equisimilis*: overexpression in *Escherichia coli*, purification, and characterization. *Prepar. Biochem.*, 23 (1993) 449-472.
- 869 Um, H.-D. and Klein, C.: Evidence for allosteric regulation of succinyl-CoA synthetase. *Biochem. J.*, 295 (1993) 821-826.
- 20j. *Complex mixtures and incompletely identified enzymes*
- 870 Zomer, A.W.M., de Weerd, W.F.C., Langeveld, J. and van den Bosch, H.: Ether lipid synthesis: purification and identification of alkyl dihydroxyacetone phosphate synthase from guinea-pig liver. *Biochim. Biophys. Acta*, 1170 (1993) 189-196.
21. PURINES, PYRIMIDINES, NUCLEIC ACIDS AND THEIR CONSTITUENTS
- 21a. *Purines, pyrimidines, nucleosides, nucleotides*
- 871 Boudinot, F.D., Schinazi, R.F., Doshi, K.J., McClure, H.M. and Chu, C.K.: Pharmacokinetics and metabolism of 3'-azido-2',3'-dideoxy-5-methylcytidine in rhesus monkeys. *Drug Metab. Disp.*, 21 (1993) 855-860.
- 872 Chen, L.-S., Fujitaki, J. and Dixon, R.: An improved assay for adenosine in rat brain microdialysates using microbore high performance liquid chromatography. *J. Liq. Chromatogr.*, 16 (1993) 2791-2796.
- 873 Cihlář, T. and Rosenberg, I.: Efficient separation of natural ribonucleotides by low-pressure anion-exchange chromatography. *J. Chromatogr.*, 644 (1993) 299-305.
- 874 Dennis, D.M., Shryock, J. and Belardinelli, L.: Correlation of (-)N<sup>6</sup>-phenylisopropyladenosine blood levels with cardiac responses in the anesthetized guinea pig. *J. Pharmacol. Exp. Ther.*, 265 (1993) 543-550.
- 875 Di Marco, M.P., Evans, C.A., Dixit, D.M., Brown, W.L., Siddiqui, M.A., Tse, H.L.A., Jin, H., Nguyen-Ba, N. and Mansour, T.S.: High-performance liquid chromatographic determination of the isomeric purity of a series of dioxolane nucleoside analogues. *J. Chromatogr.*, 645 (1993) 107-114.
- 876 Grune, T. and Siems, W.G.: Reversed-phase high-performance liquid chromatography of purine compounds for investigation of biomedical problems: application to different tissues and body fluids. *J. Chromatogr.*, 618 (1993) 15-40 - a review with 107 refs.
- 877 Hamedani, M.P., Valkó, K., Qi, X., Welham, K.J. and Gibbons, W.A.: Two-dimensional high-performance liquid chromatographic method for assaying S-adenosyl-L-methionine and its related metabolites in tissues. *J. Chromatogr.*, 619 (1993) 191-198.
- 878 Hobbs, F.W. and Yarem, J.A.: Denaturing HPLC purification of tritylated oligonucleotides using tetraethylammonium hydroxide. *BioTechniques*, 14 (1993) 584-591; *C.A.*, 119 (1993) 155145a.
- 879 Kang, J.O., Gallagher, K.S. and Cohen, G.: Methylation of RNA purine bases by methyl radicals. *Arch. Biochem. Biophys.*, 306 (1993) 178-182.
- 880 Kostianen, R., Koivisto, P. and Peltonen, K.: Characterization of the products formed in the reaction between 1,3-butadiene monoxide and 2'-deoxyadenosine by liquid chromatography-continuous-flow fast atom bombardment mass spectrometry. *J. Chromatogr.*, 647 (1993) 91-94.
- 881 Lemièrre, F., Esmans, E.L., van Dongen, W., van den Eeckhout, E. and van Onckelen, H.: Evaluation of liquid chromatography-thermospray mass spectrometry in the determination of some phenylglycidyl ether-2'-deoxynucleoside adducts. *J. Chromatogr.*, 647 (1993) 211-218.
- 882 Mao, B., Li, B., Amin, S., Cosman, M. and Greacintov, N.E.: Opposite stereoselective resistance to digestion by phosphodiesterases I and II of benzo[a]pyrene diol epoxide-modified oligonucleotide adducts. *Biochemistry*, 32 (1993) 11785-11793.
- 883 Murakami, A., Tamura, Y., Ide, H. and Makino, K.: Optimization of the separation of oligodeoxyribonucleoside phosphoramidates and their characterization by circular dichroism spectroscopy. *J. Chromatogr.*, 648 (1993) 157-163.
- 884 Nakano, K., Nakao, T., Schram, K.H., Hammargren, W.M., McClure, T.D., Katz, M. and Petersen, E.: Urinary excretion of modified nucleosides as biological marker of RNA turnover in patients with cancer and AIDS. *Clin. Chim. Acta*, 218 (1993) 169-183.
- 885 Okamura-Ikeda, K., Ohmura, Y., Fujiwara, K. and Motokawa, Y.: Cloning and nucleotide sequence of the gcv operon encoding the *Escherichia coli* glycine-cleavage system. *Eur. J. Biochem.*, 216 (1993) 539-548.
- 886 Potaman, V.N., Chernov, I.P. and Demidov, V.V.: High-performance liquid chromatography of the photoproducts of nucleic acid components. III. Detection of the secondary structure differences in sequence isomeric self-complementary oligonucleotides. *J. Chromatogr.*, 648 (1993) 151-156.

- 887 Ravanat, J.-L., Douki, T., Incardona, M.-F. and Cadet, J.: HPLC separations of normal and modified nucleobases and nucleosides on an amino silica gel column. *J. Liq. Chromatogr.*, 16 (1993) 3185-3202.
- 888 Resines, J.A., Díez, M.T. and Arín, M.J.: Statistical evaluation of agreement between HPLC and colorimetric methods for analysis of allantoin in ruminants' urine. *J. Liq. Chromatogr.*, 16 (1993) 2853-2859.
- 889 Smolenski, R.T. and Yacoub, M.H.: Liquid chromatographic evaluation of purine production in the donor human heart during transplantation. *Biomed. Chromatogr.*, 7 (1993) 189-195.
- 890 Sonoki, S., Kadoike, Y., Kiyokawa, M. and Hisamatsu, S.: High-performance liquid chromatographic analysis of 2'-deoxynucleoside 5'-monophosphate using N-(dansyl)ethylenediamine as a fluorescent derivatizing reagent. *J. Liq. Chromatogr.*, 16 (1993) 2731-2739.
- 891 Teerlink, T., Hennekes, M., Bussemaker, J. and Goreneveld, J.: Simultaneous determination of creatine compounds and adenine nucleotides in myocardial tissue by high-performance liquid chromatography. *Anal. Biochem.*, 214 (1993) 278-283.
- 892 Topp, H., Duden, R. and Schöch, G.: 5,6-Dihydrouridine: a marker ribonucleoside for determining whole body degradation rates of transfer RNA in man and rats. *Clin. Chim. Acta*, 218 (1993) 73-82.
- 893 Warren, W.J. and Vella, G.: Analysis of synthetic oligodeoxyribonucleotides by capillary gel electrophoresis and anion-exchange HPLC. *BioTechniques*, 14 (1993) 598-606; *C.A.*, 119 (1993) 112827g.
- 894 Werner, A.: Reversed-phase and ion-pair separations of nucleotides, nucleosides and nucleobases: analysis of biological samples in health and disease. *J. Chromatogr.*, 618 (1993) 3-14 - a review with 90 refs.
- 895 Yildiz, S. and Pehlivan, E.: Study of parameters affecting the column elution and ligand-exchange kinetics by using cobalt(II)-loaded bDAEG-sporopollenin ligand exchange resin in chromatographic separations. In: Aydin, A. (Editor), *Kim. Kim. Muhendisligi Semp.*, 8th, Marmara Univ. Fac. Sci. Lett., Istanbul, 1992, pp. 145-150; *C.A.*, 119 (1993) 90424u.
- 896 Yildiz, S. and Pehlivan, E.: Synthesis of new natural ligand-exchanger "aDAEG-sporopollenin" [anti-diaminoethylglycol-sym-sporopollenin] and the investigation of ligand-exchanger kinetics of this resin (in the resin). In: Aydin, A. (Editor), *Kim. Kim. Muhendisligi Semp.*, 8th, Marmara Univ. Fac. Sci. Lett., Istanbul, 1992, pp. 169-174; *C.A.*, 119 (1993) 90422s.
- 897 Zocchi, E., Guida, L., Franco, L., Silvestro, L., Guerrini, M., Benatti, U. and de Flora, A.: Free ADP-ribose in human erythrocytes: pathways of intro-erythrocytic conversion and non-enzymic binding to membrane proteins. *Biochem. J.*, 295 (1993) 121-130.

For additional information see C.A.:  
119 (1993) 108280f, 155285w.

See also 203, 519, 525, 912, 915, 976, 977, 1212, 1344.

#### 21b. Nucleic acids, RNA

- 898 Johnson, D.F., Pfsay, K.S. and Innerarity, T.L.: The mechanism for apo-B mRNA editing is deamination. *Biochem. Biophys. Res. Commun.*, 195 (1993) 1204-1210.

- 899 Kanduc, D.: tRNA fractionation by HPLC. *Nucleic Acids Res.*, 21 (1993) 2778; *C.A.*, 119 (1993) 90440w.
- 900 Kanduc, D., Farber, E., Ghoshal, A. and Nagai, M.: Sequential alterations in tRNA population of 2-acetylaminofluorene-induced hepatocyte modules. *Biochem. Biophys. Res. Commun.*, 195 (1993) 1309-1313.
- 901 Sakamoto, K., Kawai, G., Niimi, T., Satoh, T., Sekine, M., Yamaizumi, Z., Nishimura, S., Miyazawa, T. and Yokoyama, S.: A modified uridine in the first position of the anticodon of a minor species of arginine tRNA, the argU gene product, from *Escherichia coli*. *Eur. J. Biochem.*, 216 (1993) 369-375.

See also 593, 879, 907.

#### 21c. Nucleic acids, DNA

- 902 Baba, Y., Tsuhako, M., Sawa, T., Akashi, M. and Yashima, E.: (Sequence-specific recognition of DNA by high-performance affinity chromatography and capillary affinity gel electrophoresis). *Kuromatogurafi*, 13 (1992) 333-334; *C.A.*, 119 (1993) 177061x.
- 903 Chene, J., Macon, K.J. and Volanakis, J.E.: cDNA cloning and characterization of the protein encoded by RD, a gene located in the class III region of the human major histocompatibility complex. *Biochem. J.*, 294 (1993) 589-593.
- 904 Frenkel, K. and Klein, C.B.: Methods used for analyses of "environmentally" damaged nucleic acids. *J. Chromatogr.*, 618 (1993) 289-314 - a review with 176 refs.
- 905 Jeltsch, A., Fritz, A., Alves, J., Wolfes, H. and Pingoud, A.: A fast and accurate enzyme-linked immunosorbent assay for the determination of the DNA cleavage activity of restriction endonucleases. *Anal. Biochem.*, 213 (1993) 234-240.
- 906 Joshi, A.K. and Smith, S.: Construction of a cDNA encoding the multifunctional animal fatty acid synthase and expression in *Spodoptera frugiperda* cells using baculovirus vectors. *Biochem. J.*, 296 (1993) 143-149.
- 907 Kasai, K.-i.: Size-dependent chromatographic separation of nucleic acids. *J. Chromatogr.*, 618 (1993) 203-221 - a review with 56 refs.
- 908 Liu, C.H.L., Liu, S. and Wang, E.: Expression of an EF-1 $\alpha$ -like rat cDNA, S1, in *E. coli* and production of a rabbit polyclonal antiserum to the recombinant protein. *Biochem. Biophys. Res. Commun.*, 195 (1993) 1371-1378.
- 909 Shastry, B.S.: Site-specific interaction of a partially purified human lens factor(s) with conserved sequences of the human gamma crystallin gene. *Arch. Biochem. Biophys.*, 306 (1993) 476-485.

For additional information see C.A.:  
119 (1993) 155178p.

See also 593, 689, 893.

#### 21e. Structural studies on DNA and DNA mapping

- 910 Cummings, J., French, R.C. and Smyth, J.F.: Application of high-performance liquid chromatography for recognition of covalent nucleic acid modification with anticancer drugs. *J. Chromatogr.*, 618 (1993) 251-276 - a review with 165 refs.



- 911 Hartley, J.A., Souhami, R.L. and Berardini, M.D.: Electrophoretic and chromatographic separation methods used to reveal inter-strand crosslinking of nucleic acids. *J. Chromatogr.*, 618 (1993) 277-288 - a review with 37 refs.
- 912 Ling, J., Fan, M., Wei, G., Liu, N., Qiao, W. and Yi, A.: (Detection of DNA G + C mol% in taxonomy of *Streptococcus mutans* by means of high performance liquid chromatograph). *Zhonghua Kouqiang Yixue Zazhi*, 27 (1992) 326-329; *C.A.*, 119 (1993) 134683j.
- 913 Posner, B., Lee, I., Itoh, T., Pyati, J., Graff, R., Thorton, G.B., la Polla, R. and Benkovic, S.J.: A revised strategy for cloning antibody gene fragments in bacteria. *Gene*, 128 (1993) 111-117; *C.A.*, 119 (1993) 152981j.
- 914 Saravanan, K., Venkatesan, P.V. and Nagarajan, B.: A modified flow injection analysis method to quantify DNA strand-breaks. *Med. Sci. Res.*, 21 (1993) 535-538; *C.A.*, 119 (1993) 110688p.
- 915 Sonoki, S., Hisamatsu, S. and Kiuchi, A.: High-performance liquid chromatographic determination of DNA base composition with fluorescence detection. *Nucleic Acids Res.*, 21 (1993) 2776; *C.A.*, 119 (1993) 134692m.
- 916 Zeillinger, R., Schneeberger, C., Speiser, P. and Kury, F.: Rapid quantitative analysis of differential PCR products by high-performance liquid chromatography. *BioTechniques*, 15 (1993) 89-95; *C.A.*, 119 (1993) 152928x.

See also 203.

## 22. ALKALOIDS

- 917 Ascah, T.L. and Fukunaga, N.: (New HPLC column provides improved of alkaloids and other substances of abuse). *Kuromatogurafi*, 14 (1993) 25R-31R; *C.A.*, 119 (1993) 108294p.
- 918 Berlinck, R.G.S., Braekman, J.C., Daloz, D., Bruno, I., Riccio, R., Ferri, S., Spampinato, S. and Speroni, E.: Polycyclic guanidine alkaloids from the marine sponge *Crambe crambe* and Ca<sup>++</sup> channel blocker activity of crambescidin 816. *J. Natural Prod.*, 56 (1993) 1007-1015.
- 919 Boyes-Korkis, J.M., Gurney, K.A., Penn, J., Mantle, P.G., Bilton, J.N. and Sheppard, R.N.: Anacine, a new benzodiazepine metabolite of *Penicillium aurantiogriseum* produced with other alkaloids in submerged fermentation. *J. Natural Prod.*, 56 (1993) 1707-1717.
- 920 Chaudhuri, P.K. and Thakur, R.S.: 1,2-Didemethylcolchicine: a new alkaloid from *Gloriosa superba*. *J. Natural Prod.*, 56 (1993) 1174-1176.
- 921 Fliieger, M., Sedmera, P., Havlicek, V., Cvak, L. and Stuchlik, J.: 10-Hydroxy-*cis*- and 10-hydroxy-*trans*-paspalic acid amide: new alkaloid from *Claviceps paspali*. *J. Natural Prod.*, 56 (1993) 810-814.
- 922 Fliniaux, M.-A., Manceau, F. and Jacquin-Dubreuil, A.: Simultaneous analysis of *l*-hyoscyamine, *l*-scopolamine and *dl*-tropic acid in plant material by reversed-phase high-performance liquid chromatography. *J. Chromatogr.*, 644 (1993) 193-197.
- 923 Fujinari, E.M.: Nitrogen-specific liquid chromatography detection by HPLC-CLND: a novel application for determining caffeine content in beverages. *Dev. Food Sci.*, 32(Food Flavors, Ingredients and Composition) (1993) 55-64; *C.A.*, 119 (1993) 158514f.
- 924 Girgis, E.H.: Ion-pair reversed-phase liquid chromatographic identification and quantitation of papaverine congeners. *J. Pharm. Sci.*, 82 (1993) 503-505.
- 925 Glass, R.L. and Johnson, E.L.: Comparison of high performance liquid chromatographic and gas chromatographic analyses of cocaine in coca leaves. *J. Liq. Chromatogr.*, 16 (1993) 3543-3555.
- 926 Jonker, H.H., Koops, A.J. and Hoogendoorn, J.C.: A rapid method for the quantification of steroidal glycoalkaloids by reversed-phase HPLC. *Potato Res.*, 35 (1992) 451-455; *C.A.*, 119 (1993) 115620v.
- 927 Klebovich, I., Arvela, P. and Pelkonen, O.: HPLC method for rapid determination of acetylator phenotype by measuring urinary caffeine metabolites. *J. Pharm. Biomed. Anal.*, 11 (1993) 1017-1021.
- 928 Kobayashi, J., Itagaki, F., Shigemori, H. and Sasaki, T.: Three new oonamide congeners from the Okinawan marine sponge *Theonella* sp. *J. Natural Prod.*, 56 (1993) 976-981.
- 929 Kraml, M.M. and DiCosmo, F.: A rapid high-performance liquid chromatographic method for the separation of the alkaloid precursor L-tyrosine and six tetrahydroisoquinoline alkaloids of *Papaver somniferum*. *Phytochem. Anal.*, 4 (1993) 103-104; *C.A.*, 119 (1993) 134680f.
- 930 Murphey, L.J. and Olsen, G.D.: A stereospecific microassay for the determination of morphine-6- $\beta$ -D-glucuronide and other active morphine metabolites in the neonatal guinea pig. *J. Liq. Chromatogr.*, 16 (1993) 2545-2561.
- 931 Paddle, B.M. and Dowling, M.H.: Simple high-performance liquid chromatographic method for assessing the deterioration of atropine-oxime mixtures employed as antidotes in the treatment of nerve agent poisoning. *J. Chromatogr.*, 648 (1993) 373-380.
- 932 Papadoyannis, I.N., Samanidou, V.F., Tsoukali, H. and Epivatianou, F.: Comparison of a RP-HPLC method with the therapeutic drug monitoring system TD<sub>x</sub> for the determination of theophylline in blood serum. *Anal. Lett.*, 26 (1993) 2127-2142.
- 933 Price, N.P.J., Firmin, J.L., Robins, R.J. and Gray, D.O.: High-performance liquid chromatography of the alkaloid perivine from *Catharanthus roseus* after derivatisation with dansyl chloride. *J. Chromatogr.*, 653 (1993) 161-166.
- 934 Reimer, G., Suarez, A. and Chui, Y.C.: A liquid chromatographic procedure for the analysis of yohimbine in equine serum and urine. *J. Anal. Toxicol.*, 17 (1993) 178-181; *C.A.*, 119 (1993) 108291k.
- 935 Seaton, M.J., Kyerematen, G.A. and Vesell, E.S.: Rates of excretion of cotinine, nicotine glucuronide, and 3-hydroxycotinine glucuronide in rat bile. *Drug Metab. Disp.*, 21 (1993) 927-932.
- 936 Stammel, W. and Thomas, H.: A simple and rapid method for the separation of the (R)- and (S)-enantiomers of the tetrahydroisoquinoline alkaloid salsolinol by high-performance liquid chromatography. *Anal. Lett.*, 26 (1993) 2513-2524.
- 937 Wada, K., Bando, H. and Kawahara, N.: Determination and quantitative analysis of *Aconitum* alkaloids in plants by liquid chromatography-atmospheric pressure chemical ionization mass spectrometry. *J. Chromatogr.*, 644 (1993) 43-48.

For additional information see *C.A.*:

119 (1993) 103498y, 125304j, 188692e, 188693f.

See also 84, 1232, 1258, 1265, 1338, 1345, 1347.

## 23. OTHER SUBSTANCES CONTAINING HETEROCYCLIC NITROGEN

23a. *Porphyrins and other pyrroles*

- 938 Flückiger, R., Paz, M., Mah, J., Bishop, A. and Gallop, P.M.: Characterization of the glycine-dependent redox-cycling activity in animal fluids and tissues using specific inhibitors and activators: evidence for presence of PQQ. *Biochem. Biophys. Res. Commun.*, 196 (1993) 61-68.
- 939 Jacob, K. and Doss, M.O.: Composition of urinary coproporphyrin isomers I-IV in human porphyrias. *Eur. J. Clin. Chem. Clin. Biochem.*, 31 (1993) 617-624.
- 940 Kennedy, S.W. and James, C.A.: Improved method to extract and concentrate porphyrins from liver tissue for analysis by high-performance liquid chromatography. *J. Chromatogr.*, 619 (1993) 127-132.
- 941 Lim, C.K., Guo, R., Luo, J.L. and Lawson, A.M.: Studies of porphyrin metabolism by HPLC and liquid secondary ion mass spectrometry. *Int. Congr. Ser. - Excerpta Med.*, 991 (Progress in Clinical Biochemistry) (1992) 103-107; C.A., 119 (1993) 112863r.

For additional information see C.A.:  
119 (1993) 194766y.

See also 1114.

23b. *Bile pigments*

- 942 Francoual, J., Myara, A., Benattar, C., Ravaud, A., Paumier, F., Magny, J.-F., Messaoui, C. and Trivin, F.: Investigation of total and conjugated bilirubin determination during the neonatal period. *Eur. J. Clin. Chem. Clin. Biochem.*, 31 (1993) 499-502.
- 943 Min, W. and Kim, M.: A sensitive screening test for urinary porphobilinogen using minicolumn chromatography. *Int. Congr. Ser. - Excerpta Med.*, 991 (Progress in Clinical Biochemistry) (1992) 325-327; C.A., 119 (1993) 134679n.

See also 1015.

23c. *Indole derivatives and plant hormones (gibberelins)*

- 944 Ishida, J., Iizuka, R. and Yamaguchi, M.: Serotonin and 5-hydroxyindole-3-acetic acid in human plasma and rat brain determined by liquid chromatography with post-column derivatization and fluorescence detection. *Clin. Chem. (Washington)*, 39 (1993) 2355-2356.
- 945 Peniston-Bird, J.F., Di, W.-L., Street, C.A., Kadva, A., Stalteri, M.A. and Silman, R.E.: HPLC assay of melatonin in plasma with fluorescence detection. *Clin. Chem. (Washington)*, 39 (1993) 2242-2247.
- 946 Rasmussen, T., Jensen, J., Anthoni, U., Christophersen, C. and Nielsen, P.H.: Structure and synthesis of bromoindoles from the marine sponge *Pseudosuberites hyalinus*. *J. Natural Prod.*, 56 (1993) 1553-1558.

See also 474, 995.

23d. *Pyridine derivatives*

- 947 Korte, W.D. and Shih, M.L.: Degradation of three related bis(pyridinium) aldoximes in aqueous solutions at high concentrations: examples of unexpectedly rapid amide group hydrolysis. *J. Pharm. Sci.*, 82 (1993) 782-786.

For additional information see C.A.:  
119 (1993) 153464e.

See also 222.

23e. *Other N-heterocyclic compounds*

- 948 Fujimoto, H., Ikeda, M., Yamamoto, K. and Yamazaki, M.: Structure of fischerin, a new toxic metabolite from an Ascomycete, *Neosartorya fischeri* var. *fischeri*. *J. Natural Prod.*, 56 (1993) 1268-1275.
- 949 Henle, T. and Klostermeyer, H.: Determination of protein-bound 2-amino-6-(2-formyl-1-pyrrolyl)-hexanoic acid (pyrraline) by ion-exchange chromatography with photodiode array detection. *Z. Lebensm.-Unters. Forsch.*, 196 (1993) 1-4; C.A., 119 (1993) 179524z.
- 950 Huskey, S.-E.W., Miller, R.R. and Chiu, S.-H.L.: N-Glucuronidation reactions. I. Tetrazole N-glucuronidation of selected angiotensin II receptor antagonists in hepatic microsomes from rats, dogs, monkeys, and humans. *Drug Metab. Disp.*, 21 (1993) 792-799.
- 951 Janeček, M., Quilliam, M.A. and Lawrence, J.F.: Analysis of paralytic shellfish poisoning toxins by automated pre-column oxidation and microcolumn liquid chromatography with fluorescence detection. *J. Chromatogr.*, 644 (1993) 321-331.
- 952 Kunte, H.J., Galinski, E.A. and Trueper, H.G.: A modified FMOC-method for the detection of amino acid-type osmolytes and tetrahydropyrimidines (ectoines). *J. Microbiol. Methods*, 17 (1993) 129-136; C.A., 119 (1993) 90426w.
- 953 Medina, J.H., de Stein, M.L., Wolfman, C., Wasowski, C., de Blas, A. and Paladini, A.C.: *In vivo* formation of benzodiazepine-like molecules in mammalian brain. *Biochem. Biophys. Res. Commun.*, 195 (1993) 1111-1118.
- 954 Moll, N., Chevrier, S. and Moll, M.: Determination and occurrence of 3,4-benzo[a]pyrene in smoked fish and meat products. *Dev. Food Sci.*, 32 (Food Flavors, Ingredients and Composition) (1993) 233-245; C.A., 119 (1993) 137728p.
- 955 Motohashi, N., Kamata, K. and Meyer, R.: Chromatographic techniques used to determine benz[c]acridines in environmental samples. *J. Chromatogr.*, 643 (1993) 1-10 - a review with 53 refs.
- 956 Musshoff, F., Daldrup, T. and Bonte, W.: Gas chromatographic-mass spectrometric screening procedure for the identification of formaldehyde-derived tetrahydroisoquinolines in human urine. *J. Chromatogr.*, 619 (1993) 235-242.
- 957 Renz, P., Endres, B., Kurz, B. and Marquart, J.: Biosynthesis of vitamin B12 in anaerobic bacteria. Transformation of 5-hydroxybenzimidazole and 5-hydroxy-6-methylbenzimidazole into 5,6-dimethylbenzimidazole in *Eubacterium limosum*. *Eur. J. Biochem.*, 217 (1993) 1117-1121.

- 958 Watt, A.P. and Rathbone, D.: Enantiomeric separation of substituted quinuclidinyl-pyrazines by HPLC using an acetylated  $\beta$ -cyclodextrin chiral stationary phase. *J. Liq. Chromatogr.*, 16 (1993) 3423-3431.

For additional information see C.A.:  
119 (1993) 103496w, 110656b.

See also 168, 222, 517, 938, 1251, 1252, 1317, 1320, 1343, 1423.

#### 24. ORGANIC SULPHUR COMPOUNDS (INCL. GLUCOSINOLATES)

- 959 Bolzoni, L., Sannino, A. and Bandini, M.: Determination of ethylenethiourea and propylenethiourea in tomato products and in fruit purees. *Food Chem.*, 47 (1993) 299-302; C.A., 119 (1993) 115656m.
- 960 Elfakir, C., Lafosse, M. and Dreux, M.: Retention of some glucosinolates in anion exchange chromatography. Part I: Qualitative study on a silica-trialkylammonium exchanger. *Chromatographia*, 37 (1993) 313-318.
- 961 Lee, D.W., Park, Y.H. and Choi, Y.W.: (A study of liquid chromatographic detection method for thiocarbamates by using photochemical reaction). *J. Korean Chem. Soc.*, 37 (1993) 453-461; C.A., 119 (1993) 102563d.
- 962 Liu, X. and Yun, Z.: High-performance liquid chromatographic determination of thiocyanate anion by derivatization with pentafluorobenzyl bromide. *J. Chromatogr.*, 653 (1993) 348-353.
- 963 Maderich, A.B. and Sugita, E.T.: Size-exclusion chromatographic determination of dextran sulfate in rat serum. *J. Chromatogr.*, 620 (1993) 137-142.
- 964 Mank, A.J.G., Molenaar, E.J., Lingeman, H., Gooijer, C., Brinkman, U.A.T. and Velthorst, N.H.: Visible diode laser induced fluorescence detection in liquid chromatography after pre-column derivatization of thiols. *Anal. Chem.*, 65 (1993) 2197-2203.
- 965 Müller, B. and Ruhnke, A.: Qualitätsprüfung von Knoblauchpulver. Vorschläge für eine Arzneibuchmonographie. *Dtsch. Apoth.-Ztg.*, 133 (1993) 37-47.
- 966 Niessen, W.M.A., McCarney, C.C., Moul, P.E.G., Tjaden, U.R. and van der Greef, J.: Liquid chromatography-mass spectrometry for the identification of minor components in benzothiazole derivatives. *J. Chromatogr.*, 647 (1993) 107-119.
- 967 Nishioka, M. and Tomich, R.S.: Isolation of aliphatic sulphur compounds in a crude oil by a nonreactive procedure. *Fuel*, 72 (1993) 1007-1010; C.A., 119 (1993) 142432s.
- 968 Ogasawara, Y., Ishii, K., Togawa, T. and Tanabe, S.: Determination of bound sulfur in serum by gas dialysis/high-performance liquid chromatography. *Anal. Biochem.*, 215 (1993) 73-81.
- 969 Raghuvveeran, C.D., Malhotra, R.C. and Dangi, R.S.: Reversed-phase high-performance liquid chromatography of sulphur mustard in water. *J. Liq. Chromatogr.*, 16 (1993) 1615-1624.
- 970 Simon, P. and Nicot, T.: Automated column-switching high-performance liquid chromatography for the determination of 2-thiothiazolidine-4-carboxylic acid in urine. *J. Chromatogr.*, 620 (1993) 47-53.

- 971 Vandenberg, P.J. and Johnson, D.C.: Pulsed electrochemical detection of cysteine, cystine, methionine, and glutathione at gold electrodes following their separation by liquid chromatography. *Anal. Chem.*, 65 (1993) 2713-2718.

- 972 Zou, H., Zhang, Y., Hong, M. and Lu, P.: Retention behaviour of aromatic sulphonic acids in reversed-phase ion-pair liquid chromatography with methanol and acetonitrile as organic modifiers. *J. Chromatogr.*, 644 (1993) 269-275.

For additional information see C.A.:  
119 (1993) 90722q, 112579c.

See also 83, 103, 222, 1138.

#### 25. ORGANIC PHOSPHORUS COMPOUNDS (INCL. SUGAR PHOSPHATES)

- 973 Fanjul, L.F., Marrero, I., González, J., Quintana, J., Santana, P., Estévez, F., Mato, J.M. and de Gálarreta, C.M.R.: Does oligosaccharide-phosphatidylinositol (glycosyl-phosphatidylinositol) hydrolysis mediate prolactin signal transduction in granulosa cells? *Eur. J. Biochem.*, 216 (1993) 747-755.
- 974 Kellner, M., Machicao, F., Berti, L., Sixt, B., Mushack, J., Seffer, E., Mosthaf, L., Ullrich, A. and Häring, H.U.: Inositol phospholipid oligosaccharides from rat fibroblasts and adipocytes stimulate 3-O-methylglucose transport. *Biochem. J.*, 295 (1993) 699-704.
- 975 Kennedy Keller, R. and Thompson, R.: Rapid synthesis of isoprenoid diphosphates and their isolation in one step using either thin layer or flash chromatography. *J. Chromatogr.*, 645 (1993) 161-167.
- 976 Lefrançois, L. and Gaudreau, P.: Basal and stimulated cyclic adenosine 3',5'-monophosphate production in rat adenopituitary homogenates: direct determination by high-performance liquid chromatography. *J. Chromatogr.*, 619 (1993) 116-120.
- 977 Rimerman, R.A., Prorok, G.D., Cordel, K.L., Shahwan, A.M. and Vaughan, W.P.: Improved high-performance liquid chromatographic analysis of intracellular deoxyribonucleoside triphosphate levels. *J. Chromatogr.*, 619 (1993) 29-35.
- 978 Rounds, M.A. and Nielsen, S.S.: Anion-exchange high-performance liquid chromatography with post-column detection for the analysis of phytic acid and other inositol phosphates. *J. Chromatogr.*, 653 (1993) 148-152.
- 979 Villa, S., Rossi, F., Biondi, P.A., Russo, V., Crimella, T., Fiorelli, G. and Zanella, A.: Determination of inositol hexaphosphate (IHP) in human IHP-loaded red blood cells by a simple high performance liquid chromatography method. *Adv. Exp. Med. Biol.*, 326 (1992) 41-49; C.A., 119 (1993) 103468p.
- 980 Wilson, I.B.H., Taylor, J.P., Webberley, M.C., Turner, N.J. and Flitsch, S.L.: A novel mono-branched lipid phosphate acts as a substrate for dolichyl phosphate mannose synthetase. *Biochem. J.*, 295 (1993) 195-201.
- 981 Woodcock, E.A., Anderson, K.E. and Land, S.L.: Lyophilization can generate artifacts in chromatographic profiles of inositol phosphates. *J. Chromatogr.* 619 (1993) 121-126.
- 982 Zhang, D. and Poulter, C.D.: Analysis and purification of phosphorylated isoprenoids by reversed-phase HPLC. *Anal. Biochem.*, 213 (1993) 356-361.

See also 442, 584, 802, 883.

## 26. ORGANOMETALLIC AND RELATED COMPOUNDS

## 26a. Organometallic compounds

- 983 Dauchy, X., Cottier, R., Batel, A., Jeannot, R., Borsier, M., Astruc, A. and Astruc, M.: Speciation of butyltin compounds by high-performance liquid chromatography with inductively coupled plasma mass spectrometry detection. *J. Chromatogr. Sci.*, 31 (1993) 416-421.

See also 156.

## 26b. Boranes, silanes and related non-metallic compounds

- 984 Rubio, R., Peralta, I., Albertí, J. and Rauret, G.: Arsenic species separation by IELC-ICP/OES: arsenocholine behavior. *J. Liq. Chromatogr.*, 16 (1993) 3531-3542.

## 26c. Coordination compounds

- 985 Careri, M., Mangia, A., Manini, P., Predieri, G., Raverdino, V., Tsoupras, G. and Sappa, E.: Particle beam liquid chromatography-mass spectrometry behaviour of polynuclear metal carbonyl compounds. *J. Chromatogr.*, 647 (1993) 79-90.
- 986 Khuhawar, M.Y. and Bhatti, A.G.: Chromatographic studies of oxovanadium(IV), palladium(II), copper(II) and nickel(II) chelates of some fluorinated tetradentate Schiff bases. *J. Chem. Soc. Pak.*, 14 (1992) 244-248; *C.A.*, 119 (1993) 107940j.
- 987 Otu, E., Robinson, C.W. and Byerley, J.J.: Inorganic modifiers in ion-pair chromatographic separation of dicyanoaurate (I). *Analyst (London)*, 118 (1993) 1277-1280.
- 988 Timerbaev, A.R., Semenova, O.P., Tsoi, I.G. and Petrukhin, O.M.: Correlation analysis in liquid chromatography of metal chelates. III. Multi-dimensional models in reversed-phase liquid chromatography. *J. Chromatogr.*, 648 (1993) 307-314.

## 27. VITAMINS AND VARIOUS ANIMAL GROWTH FACTORS (NON-PEPTIDIC)

- 989 Abdollahi, A., Rosenholtz, N.S. and Garwin, J.L.: Tocopherol micro-extraction method with application to quantitative analysis of lipophilic nutrients. *J. Food Sci.*, 58 (1993) 663-666; *C.A.*, 119 (1993) 158537r.
- 990 Baldet, P., Gerbling, H., Axiotis, S. and Douce, R.: Biotin biosynthesis in higher plant cells. Identification of intermediates. *Eur. J. Biochem.*, 217 (1993) 479-485.
- 991 Benlloch, R., Farré, R. and Frigola, A.: A quantitative estimate of ascorbic and isoascorbic acid by high performance liquid chromatography: application to citric juices. *J. Liq. Chromatogr.*, 16 (1993) 3113-3122.
- 992 Dieuaide, M., Couée, I., Pradet, A. and Raymond, P.: Effect of glucose starvation on the oxidation of fatty acids by maize root tip mitochondria and peroxisomes: evidence for mitochondrial fatty acid  $\beta$ -oxidation and acyl-CoA dehydrogenase activity in a higher plant. *Biochem. J.*, 296 (1993) 199-207.
- 993 Epler, K.S., Ziegler, R.G. and Craft, N.E.: Liquid chromatographic method for the determination of carotenoids, retinoids and tocopherols in human serum and in food. *J. Chromatogr.*, 619 (1993) 37-48.

- 994 Fraser, P.D., Albrecht, M. and Sandmann, G.: Development of high-performance liquid chromatographic systems for the separation of radiolabelled carotenes and precursors formed in specific enzymatic reactions. *J. Chromatogr.*, 645 (1993) 265-272.
- 995 Guerriero, A., D'Ambrosio, M., Peitra, F., Debitus, C. and Ribes, O.: Pteridines, sterols, and indole derivatives from the lithistid sponge *Corallistes undulatus* of the Coral Sea. *J. Natural Prod.*, 56 (1993) 1962-1970.
- 996 Honda, A., Nakashima, N., Shida, Y., Mori, Y., Nagata, A. and Ishizuka, S.: Modification of 1 $\alpha$ ,25-dihydroxyvitamin D<sub>3</sub> metabolism by introduction of 26,26,26,27,27,27-hexafluoro atoms in human promyelocytic leukaemia (HL-60) cells: isolation and identification of a novel bioactive metabolite, 26,26,26,27,27,27-hexafluoro-1 $\alpha$ ,23(S),25-trihydroxyvitamin D<sub>3</sub>. *Biochem. J.*, 295 (1993) 503-516.
- 997 Kimoto, E., Tanaka, H., Ohmoto, T. and Choami, M.: Analysis of the transformation products of dehydro-L-ascorbic acid by ion-pairing high-performance liquid chromatography. *Anal. Biochem.*, 214 (1993) 38-44.
- 998 Kozik, A. and Rapala-Kozik, M.: Ion-pair reversed-phase high-performance liquid chromatographic method for the separation of a set of unphosphorylated thiamine-related compounds. *J. Chromatogr.*, 648 (1993) 349-356.
- 999 Lesellier, E., Tchaplá, A. and Krstulovic, A.M.: Use of carotenoids in the characterization of octadecylsilane bonded columns and mechanism of retention of carotenoids on monomeric and polymeric stationary phases. *J. Chromatogr.*, 645 (1993) 29-39.
- 1000 Lin, L., Meltzer, N.M. and Honigberg, I.L.: Normal phase HPLC with EC detection: 1,25-dihydroxy vitamin D<sub>3</sub> and 25-hydroxy-16-ene-23-yne vitamin D<sub>3</sub>. *J. Liq. Chromatogr.*, 16 (1993) 3093-3100.
- 1001 Mascher, H.: Determination of total pyridoxal in human plasma following oral administration of vitamin B<sub>6</sub> by high-performance liquid chromatography with post-column derivatization. *J. Pharm. Sci.*, 82 (1993) 972-974.
- 1002 Micali, G., Lanuzza, F. and Currò, P.: Analysis of tocopherols in margarine by on-line HPLC-HRGC coupling. *J. High Resolut. Chromatogr.*, 16 (1993) 536-538.
- 1003 Mladek, C.: Vitamins. In: Matissek, R. and Wittkowski, R. (Editors), *High Perform. Liq. Chromatogr. Food Control Res.*, Technomic, Lancaster, 1993, pp. 183-233; *C.A.*, 119 (1993) 137620x - a review with 105 refs.
- 1004 Nakanishi, H., Suga, K. and Taniguchi, K.: (Determination of cyanocobalamin in tablets by high-performance liquid chromatography after pretreatment with gel filtration-solid phase extraction). *Iyakuin Kenkyu*, 24 (1993) 515-519; *C.A.*, 119 (1993) 146704e.
- 1005 Napolli, J.L.: Retinoic acid synthesis from  $\beta$ -carotene *in vitro*. *Methods Enzymol.*, 214(Carotenoids, Pt. B) (1993) 193-202; *C.A.*, 119 (1993) 134677k.
- 1006 Ohta, H., Maeda, M., Nogata, Y., Yoza, K.-I., Takeda, Y. and Osajima, Y.: A simple determination of thiamine in rice (*Oryza sativa* L.) by high-performance liquid chromatography with post-column derivatization. *J. Liq. Chromatogr.*, 16 (1993) 2617-2629.
- 1007 Olds, S.J., Vanderslice, J.T. and Brochetti, D.: Vitamin B<sub>6</sub> in raw and fried chicken by HPLC. *J. Food Sci.*, 58 (1993) 505-561; *C.A.*, 119 (1993) 158536q.

- 1008 Otlies, S. and Hisil, Y.: High pressure liquid chromatographic analysis of water-soluble vitamins in eggs. *Ital. J. Food Sci.*, 5 (1993) 69-75; C.A., 119 (1993) 158511c.
- 1009 Palfrey, S.M. and Labib, M.H.: A simple method for measuring neopterin in serum using HPLC. *Ann. Clin. Biochem.*, 30 (1993) 209-210; C.A., 119 (1993) 112578b.
- 1010 Pelzer, M., Northcott, S. and Hanson, G.: An improved method for the determination of nicotinic acid in human plasma by high-performance liquid chromatography. *J. Liq. Chromatogr.*, 16 (1993) 2563-2570.
- 1011 Rock, C.L. and Swendseid, M.E.: Plasma carotenoid levels in anorexia nervosa and in obese patients. *Methods Enzymol.*, 214(Carotenoids, Pt. B) (1993) 116-123; C.A., 119 (1993) 134676j.
- 1012 Rosebrough, S.F.: Plasma stability and pharmacokinetics of radiolabeled deferoxamine-biotin derivatives. *J. Pharmacol. Exp. Ther.*, 265 (1993) 408-415.
- 1013 Salyers, K.L., Cullum, M.E. and Zile, M.H.: Glucuronidation of all-trans-retinoic acid in liposomal membranes. *Biochim. Biophys. Acta*, 1152 (1993) 328-334.
- 1014 Santoro, M.I.R.M., Campos, P.M.B.G.M. and Hackmann, E.R.M.: Determining vitamins A, D<sub>3</sub> and E in creams: high performance liquid chromatography. *Cosmet. Toiletries*, 108 (1993) 71-74; C.A., 119 (1993) 124792e.
- 1015 Sarzanini, C., Mentasti, E., Vincenti, M., Nerva, M. and Gaido, F.: Determination of plasma tocopherols by high-performance liquid chromatography with coulometric detection. *J. Chromatogr.*, 620 (1993) 268-272.
- 1016 Satomura, Y., Kimura, M., Hiraike, H. and Itokawa, Y.: (A method for simultaneous determination of retinol and tocopherol homologues and its application to biological specimens). *B-tamin*, 67 (1993) 111-119; C.A., 119 (1993) 90425v.
- 1017 Scott, K.J. and Hart, D.J.: Further observations on problems associated with the analysis of carotenoids by HPLC-2: column temperature. *Food Chem.*, 47 (1993) 403-405; C.A., 119 (1993) 115668s.
- 1018 Shin, H.-C., Shimoda, M., Kokue, E. and Takahashi, Y.: Identification of 10-formyltetrahydrofolate, tetrahydrofolate and 5-methyltetrahydrofolate as major reduced folate derivatives in rat bile. *J. Chromatogr.*, 620 (1993) 39-46.
- 1019 Susin, S., Abián, J., Sánchez-Baeza, F., Peleato, M.L., Abadia, A., Galpi, E. and Abadia, J.: Riboflavin 3'- and 5'-sulfate, two novel flavins accumulating in the roots of iron-deficient sugar beet (*Beta vulgaris*). *J. Biol. Chem.*, 268 (1993) 20958-20965.
- 1020 Tachikawa, H., Yazawa, I. and Kitagawa, K.: (Separation characteristics of fat-soluble vitamins on amino-silica column, YMC-pack polyamine-II in normal phase). *Kuromatoguraji*, 13 (1992) 295-296; C.A., 119 (1993) 112572v.
- 1021 Tan, X., Meltzer, N. and Lindenbaum, S.: Determination of the kinetics of degradation of 13-cis-retinoic acid and all-trans-retinoic acid in solution. *J. Pharm. Biomed. Anal.*, 11 (1993) 817-822.
- 1022 Ueda, T., Ichikawa, H. and Igarashi, O.: Determination of  $\alpha$ -tocopherol stereoisomers in biological specimens using chiral phase high-performance liquid chromatography. *J. Nutr. Sci. Vitaminol.*, 39 (1993) 207-219; C.A., 119 (1993) 176882d.
- 1023 Vatassery, G.T., Smith, W.E. and Quach, H.T.: A liquid chromatographic method for the simultaneous determination of  $\alpha$ -tocopherol and tocopherolquinone in human red blood cells and other biological samples where tocopherol is easily oxidized during sample treatment. *Anal. Biochem.*, 214 (1993) 426-430.
- 1024 Verne-Mismer, J., Lamard, M. and Wagner, J.: Evaluation of deactivated reversed phases for the analysis of an N,N,N-trimethylethanaminium analogue of  $\alpha$ -tocopherol. Application to its purity control and determination in biological samples. *J. Chromatogr.*, 645 (1993) 251-258.
- 1025 White, S.: Determination of vitamins K1, K2 (MK-4) and K3 animal feeds by high-performance liquid chromatography. *Anal. Proc.*, 30 (1993) 266-267; C.A., 119 (1993) 158505d.
- 1026 Yeung, B., Vouros, P. and Satyanarayana Reddy, G.S.: Characterization of vitamin D<sub>3</sub> metabolites using continuous-flow fast atom bombardment tandem mass spectrometry and high-performance liquid chromatography. *J. Chromatogr.*, 645 (1993) 115-123.
- 1027 Zaman, Z., Fielden, P. and Frost, P.G.: Simultaneous determination of vitamins A and E and carotenoids in plasma by reversed-phase HPLC in elderly and younger subjects. *Clin. Chem. (Washington)*, 39 (1993) 2229-2234.

For additional information see C.A.:

119 (1993) 115651f, 137737r, 155156e, 194769b.

See also 462, 498, 702, 1116.

## 28. ANTIBIOTICS

- 1028 Aburaki, S., Okuyama, S., Hoshi, H., Kamachi, H., Nishio, M., Hasegawa, T., Masuyoshi, S., Iimura, S., Konishi, M. and Oki, T.: Synthesis and antifungal activity of pradimicin derivatives. Modifications on the aglycone part. *J. Antibiot.*, 46 (1993) 1447-1457.
- 1029 Aoyama, T., Zhao, W., Kojima, F., Muraoka, Y., Naganawa, H., Takeuchi, T. and Aoyagi, T.: Cysfluoretin, a new inhibitor of glutathione S-transferase, produced by *Streptomyces* sp. MI384-DF12. *J. Antibiot.*, 46 (1993) 1471-1474.
- 1030 Baertschi, S.W., Dorman, D.E., Occolowitz, J.L., Spangle, L.A., Collins, M.W., Wildfeuer, M.E. and Lorenz, L.J.: Isolation and structure elucidation of a novel product of the acidic degradation of cefaclor. *J. Pharm. Sci.*, 82 (1993) 622-626.
- 1031 Blanchflower, S.E., Banks, R.M., Everett, J.R. and Reading, C.: Further novel metabolites of the paraherquamide family. *J. Antibiot.*, 46 (1993) 1355-1363.
- 1032 Boger, D.L., Menezes, R.F. and Yang, W.: A simple method for the purification and isolation of bleomycin A<sub>2</sub>. *Bioorg. Med. Chem. Lett.*, 2 (1992) 959-962; C.A., 119 (1993) 125169u.
- 1033 Breinholt, J., Jensen, G.W., Nielsen, R.I., Olsen, C.E. and Frisvad, J.C.: Antifungal macrocyclic polyactones from *Penicillium verruculosum*. *J. Antibiot.*, 46 (1993) 1101-1108.
- 1034 Bryan, P.D. and Stewart, J.T.: Separation of tetracyclines by liquid chromatography with acidic mobile phases and polymeric columns. *J. Pharm. Biomed. Anal.*, 11 (1993) 971-976.

- 1035 Cannell, R.J.P., Dawson, M.J., Hale, R.S., Hall, R.M., Noble, D., Lynn, S. and Taylor, N.L.: The squalenylated, novel inhibitors of squalene synthase produced by a species of *Phoma*. IV. Preparation of fluorinated squalenylated by directed biosynthesis. *J. Antibiot.*, 46 (1993) 1381-1389.
- 1036 Charles, B. and Chulavatnatol, S.: Simple analysis of amoxicillin in plasma by high performance liquid chromatography with internal standardization and ultraviolet detection. *Biomed. Chromatogr.*, 7 (1993) 204-207.
- 1037 De Jong, J., Bast, A. and van der Vijgh, W.J.F.: Analysis of anthracycline antitumour drugs in tissues and body fluids using liquid chromatography. *TrAC*, 12 (1993) 422-428 - a review with 30 refs.
- 1038 Degroot, J.M., Wyhowski, de Bukanski, B. and Srebrnik, S.: Multiresidue analysis of tetracyclines in kidney by HPLC and photodiode array detection. *J. Liq. Chromatogr.*, 16 (1993) 3515-3529.
- 1039 Erlichman, C., Moore, M., Thiessen, J.J., Kerr, I.G., Walker, S., Goodman, P., Bjarnason, G., DeAngelis, C. and Bunting, P.: Phase I pharmacokinetic study of cyclosporin A combined with doxorubicin. *Cancer Res.*, 53 (1993) 4837-4842.
- 1040 Guzmán, M., Molpeceres, J., García, F., Aberturas, M.R. and Rodríguez, M.: Formation and characterization of cyclosporine-loaded nanoparticles. *J. Pharm. Sci.*, 82 (1993) 498-502.
- 1041 Hasinoff, B.B.: Enzymatic ring-opening reactions of the chiral cardioprotective agent (+)(S)-ICRF-187 and its (-)(R)-enantiomer ICRF-186 by dihydropyrimidine aminohydrolase. *Drug Metab. Disp.*, 21 (1993) 883-888.
- 1042 Hedge, V.R., Patel, M.G., Gullo, V.P., Horan, A.C., King, A.H., Gentile, F., Wagman, G.H., Puar, M.S. and Loebenberg, D.: A novel macrolactam-disaccharide antifungal antibiotic. Taxonomy, fermentation, isolation, physico-chemical properties, structure elucidation and biological activity. *J. Antibiot.*, 46 (1993) 1109-1115.
- 1043 Hendrix, C., Roets, E., Crommen, J., de Beer, J., Porqueras, E. van den Bossche, W. and Hoogmartens, J.: Collaborative study of the analysis of tetracycline by liquid chromatography on poly(styrene-divinylbenzene). *J. Liq. Chromatogr.*, 16 (1993) 3321-3329.
- 1044 Hendrix, C., Yongxin, Z., Pijcke, M., Roets, E. and Hoogmartens, J.: A comparative study of LC methods for analysis of cefradine. *J. Pharm. Biomed. Anal.*, 11 (1993) 595-599.
- 1045 Hori, Y., Abe, Y., Ezaki, M., Goto, T., Okuhara, M. and Kohsaka, M.: R1128 substances, novel non-steroidal estrogen-receptor antagonists produced by a *Streptomyces* I. Taxonomy, fermentation, isolation and biological properties. *J. Antibiot.*, 46 (1993) 1055-1062.
- 1046 Husek, A., Eibler, E. and Wiegrebe, W.: High-performance liquid chromatographic analysis of cyclosporine A in human skin. *Arch. Pharm. (Weinheim)*, 326 (1993) 365-368; *C.A.*, 119 (1993) 85312b.
- 1047 Isono, F., Sakaida, Y., Takahashi, S., Kinoshita, T., Nakamura, T. and Inukai, M.: Mureidomycins E and F, minor components of mureidomycins. *J. Antibiot.*, 46 (1993) 1203-1207.
- 1048 Iwaki, K., Okumura, N. and Yamazaki, M.: Rapid determination of tetracycline antibiotics in serum by reversed-phase high-performance liquid chromatography with fluorescence detection. *J. Chromatogr.*, 619 (1993) 319-323.
- 1049 Iwata-Reuyi, D., Basak, A., Silverman, L.S., Engle, C.A. and Townsend, C.A.: Synthesis and reaction of potential alternate substrates and mechanism-based inhibitors of clavamate synthase. *J. Natural Prod.*, 56 (1993) 1373-1396.
- 1050 Janeček, M., Quilliam, M.A., Bailey, M.R. and North, D.H.: Determination of erythromycin A by liquid chromatography and electrochemical detection, with application to salmon tissue. *J. Chromatogr.*, 619 (1993) 63-69.
- 1051 Johdo, O., Tone, H., Okamoto, R., Yoshimoto, A., Naganawa, H., Sawa, T. and Takeuchi, T.: Microbial glycosidation of some anthracycline antibiotics by an antibiotic-negative mutant of a aclarubicin producer. *J. Antibiot.*, 46 (1993) 1219-1231.
- 1052 Kamiyama, T., Itezone, Y., Umino, T., Satoh, T., Nakayama, N. and Yokose, K.: Tetrafrabricin, a novel fibrinogen receptor antagonist I. Structural elucidation. *J. Antibiot.*, 46 (1993) 1047-1054.
- 1053 Kamiyama, T., Umino, T., Fujisaki, N., Fujimori, K., Satoh, T., Yamashita, Y., Ohshima, S., Watanabe, J. and Yokose, K.: Tetrafrabricin, a novel fibrinogen receptor antagonist I. Taxonomy, fermentation, isolation, characterization and biological activities. *J. Antibiot.*, 46 (1993) 1039-1046.
- 1054 Kaniou, I.P., Zachariadis, G.A. and Stratis, J.A.: Separation and determination of five penicillins by reversed phase HPLC. *J. Liq. Chromatogr.*, 16 (1993) 2891-2897.
- 1055 Karnes, H.T., Wei, A.T. and Dimenna, G.P.: HPLC analysis of salinomycin in human plasma using pre-column oxidation and automated heart cut column switching. *J. Pharm. Biomed. Anal.*, 11 (1993) 823-827.
- 1056 Kumagai, K., Fukui, A., Tanaka, S., Ikemoto, M., Moriguchi, K. and Nabeshima, S.: PC-766B, a new macrolide antibiotic produced by *Nocardia brasiliensis*. II. Isolation, physico-chemical properties and structure elucidation. *J. Antibiot.*, 46 (1993) 1139-1144.
- 1057 Kuroda, K., Yoshida, M., Uosaki, Y., Ando, K., Kawamoto, I., Oishi, E., Onuma, H., Yamada, K. and Matsuda, Y.: AS-183, a novel inhibitor of acyl-CoA:cholesterol acyltransferase produced by *Scedosporium* sp. SPC-15549. *J. Antibiot.*, 46 (1993) 1196-1202.
- 1058 Laatsch, H., Kellner, M., Wolf, G., Lee, Y.-S., Hansske, F., Kohnetschny-Rapp, S., Pessara, U., Scheuer, W. and Stockinger, H.: Oligomycin F, a new immunosuppressive homologue of oligomycin A. *J. Antibiot.*, 46 (1993) 1334-1341.
- 1059 Lhoëst, G., Maton, N. and Verbeeck, R.K.: Isolation and identification of a novel isomerized epoxide metabolite of FK-506 from erythromycin-induced rabbit liver microsomes. *Drug Metab. Disp.*, 21 (1993) 850-854.
- 1060 Li, W.Y., Seah, S.K.L. and Koda, R.T.: Determination of mitomycin C in human aqueous humor and serum by high-performance liquid chromatography. *J. Chromatogr.*, 619 (1993) 148-153.
- 1061 Mathew, M., das Gupta, V. and Bethea, C.: Quantitation of cefaclor in pharmaceutical dosage forms using high performance liquid chromatography. *Drug. Dev. Ind. Pharm.*, 19 (1993) 1723-1729; *C.A.*, 119 (1993) 188730r.
- 1062 Nakashima, K., Kawaguchi, S., Akiyama, S. and Schulman, S.G.: High performance liquid chromatography of penicillins with penicillin-enhanced luminol chemiluminescence detection. *Biomed. Chromatogr.*, 7 (1993) 217-219.

- 1063 Namiki, Y., Kihara, N., Koda, S., Hane, K. and Yasuda, T.: Tautomeric phenomenon of a novel potent immunosuppressant (FK506) in solution. I. Isolation and structure determination of tautomeric compounds. *J. Antibiot.*, 46 (1993) 1149-1155.
- 1064 Oh, D.-M., Sinko, P.J. and Amidon, G.L.: Characterization of the oral absorption of some  $\beta$ -lactams: effect of the  $\alpha$ -amino side chain group. *J. Pharm. Sci.*, 82 (1993) 897-900.
- 1065 Olsen, B.A., Baertschi, S.W. and Riggan, R.M.: Multidimensional evaluation of impurity profiles for generic cephalexin and cefaclor antibiotics. *J. Chromatogr.*, 648 (1993) 165-173.
- 1066 Paesen, J., Calam, D.H., Miller, J.H.Mc.B., Raiola, G., Rozanski, A., Silver, B. and Hoogmartens, J.: Collaborative study of the analysis of erythromycin by liquid chromatography on wide-pore poly(styrene-divinylbenzene). *J. Liq. Chromatogr.*, 16 (1993) 1529-1544.
- 1067 Petersen, F., Zähler, H., Metzger, J.W., Freund, S. and Hummel, E.-P.: Germicidin, an autoregulative germination inhibitor of *Streptomyces viridochromogenes* NRRL B-1551. *J. Antibiot.*, 46 (1993) 1126-1138.
- 1068 Psomas, J.E. and losifidoy, E.G.: High performance liquid chromatographic analysis of residues of thiamphenicol in beef muscle. *J. Liq. Chromatogr.*, 16 (1993) 2653-2660.
- 1069 Rogstad, A. and Weng, B.: Extraction and analysis by high-performance liquid chromatography of antibiotics in a drug delivery system for farmed fish. *J. Pharm. Sci.*, 82 (1993) 518-520.
- 1070 Saito, A., Naito, S., Kobayashi, M., Tsuchiya, M., Toyama, T., Kaneko, S., Nanba, T. and Morisawa, Y.: Synthesis and antihelminthic activity of 13-alkoxymilbemycin derivatives. *J. Antibiot.*, 46 (1993) 1252-1264.
- 1071 Shafiee, A., Chen, T.S., Arison, B.S., Dumont, F.J., Colwell, L. and Kaplan, L.: Enzymatic synthesis and immunosuppressive activity of novel desmethylated immunomycins (ascomycins). *J. Antibiot.*, 46 (1993) 1397-1405.
- 1072 Shen, B., Nakayama, H. and Hutchinson, C.R.: Isolation and structural elucidation of tetracenomycin F2 and tetracenomycin F1: early intermediates in the biosynthesis of tetracenomycin C in *Streptomyces glaucescens*. *J. Natural Prod.*, 56 (1993) 1288-1293.
- 1073 Shimada, J., Hori, S., Oguma, T., Yoshikawa, T., Yamamoto, S., Nishikawa, T. and Yamada, H.: Effects of protein binding on the isomerization of ceftibuten. *J. Pharm. Sci.*, 82 (1993) 461-465.
- 1074 Shindo, K., Kamishohara, M., Odagawa, A., Matsuoka, M. and Kawai, H.: Vicenistatin, a novel 20-membered macrocyclic lactam antitumor antibiotic. *J. Antibiot.*, 46 (1993) 1076-1081.
- 1075 Straub, R.F. and Voyksner, R.D.: Determination of penicillin G, ampicillin, amoxicillin, cloxacillin and cephalixin by high-performance liquid chromatography-electrospray mass spectrometry. *J. Chromatogr.*, 647 (1993) 167-181.
- 1076 Takahashi, I., Maruta, R., Ando, K., Yoshida, M., Iwasaki, T., Kanazawa, J., Okabe, M. and Tamaoki, T.: UCA1064-B, a new antitumor antibiotic isolated from *Wallemia sebi*: production, isolation and structural determination. *J. Antibiot.*, 46 (1993) 1312-1314.
- 1077 Takahashi, S., Miyaoka, H., Tanaka, K., Enokita, R. and Okazaki, T.: Milbemycins  $\alpha_{11}$ ,  $\alpha_{12}$ ,  $\alpha_{13}$ ,  $\alpha_{14}$  and  $\alpha_{15}$ : a new family of milbemycins from *Streptomyces hygroscopicus* subsp. *aureolacrimosus*. Taxonomy, fermentation, isolation, structure elucidation and biological properties. *J. Antibiot.*, 46 (1993) 1364-1371.
- 1078 Tanaka, Y., Kanaya, I., Shiomi, K., Tanaka, H. and Omura, S.: Phthoxazolin, a specific inhibitor of cellulose biosynthesis from microbial origin. II. Isolation, physico-chemical properties, and structural elucidation. *J. Antibiot.*, 46 (1993) 1214-1218.
- 1079 Tokumura, T. and Horie, T.: Determination of a novel  $\beta$ -lactam antibiotic (E-1100) in rat plasma by high-performance liquid chromatography. *J. Chromatogr.*, 620 (1993) 153-157.
- 1080 Ueno, M., Yoshinaga, I., Amemiya, M., Someno, T., Inuma, H., Ishizuaka, M. and Takeuchi, T.: Delaminomycins, novel extracellular matrix receptor antagonists. V. Biosynthesis. *J. Antibiot.*, 46 (1993) 1390-1396.
- 1081 Ventura, N., Alvares de Souza, A. and Korolkovas, A.: (Determination of penicillins using colorimetric, mercurimetric, and chromatographic methods). *Rev. Port. Farm.*, 41 (1992) 15-22; *C.A.*, 119 (1993) 125283b - a review with 26 refs.
- 1082 Wallemacq, P.E. and Reding, R.: FK506 (tacrolimus), a novel immunosuppressant in organ transplantation: clinical, biomedical, and analytical aspects. *Clin. Chem. (Washington)*, 39 (1993) 2219-2228 - a review with 102 refs.
- 1083 White, C.R., Moats, W.A. and Kotula, K.L.: Comparative study of high performance liquid chromatographic methods for the determination of tetracycline antibiotics. *J. Liq. Chromatogr.*, 16 (1993) 2873-2890.
- 1084 Yamazaki, T., Tatee, T., Aoyama, T., Kojima, F., Takeuchi, T. and Aoyagi, T.: Bequinostatin C and D, new inhibitors of glutathione S-transferase, produced by *Streptomyces* sp. MI384-DF12. *J. Antibiot.*, 46 (1993) 1309-1311.
- 1085 Yoshikawa, Y., Ikai, K., Umeda, Y., Ogawa, A., Takesako, K. and Kato, I.: Isolation, structures, and antifungal activities of new aureobasidins. *J. Antibiot.*, 46 (1993) 1347-1354.

For additional information see C.A.:

119 (1993) 90421r, 93828q, 103481n, 103482p, 110692k, 115484d, 120194p, 130803a, 167581c, 167866z.

See also 130, 1306.

## 29. INSECTICIDES, PESTICIDES AND OTHER AGROCHEMICALS

- 1086 Bagheri, H., Brouwer, E.R., Ghijsen, R.T. and Brinkman, U.A.T.: On-line low-level screening of polar pesticides in drinking and surface waters by liquid chromatography-thermospray mass spectrometry. *J. Chromatogr.*, 647 (1993) 121-129.
- 1087 Barceló, D.: Environmental Protection Agency and other methods for the determination of priority pesticides and their transformation products in water. *J. Chromatogr.*, 643 (1993) 117-143 - a review with 103 refs.
- 1088 Chiron, S. and Barceló, D.: Determination of pesticides in drinking water by on-line solid-phase disk extraction followed by various liquid chromatographic systems. *J. Chromatogr.*, 645 (1993) 125-134.
- 1089 Hogendoorn, E.A., Brinkman, U.A.T. and van Zoonen, P.: Coupled-column reversed-phase liquid chromatography-UV analyser for the determination of polar pesticides in water. *J. Chromatogr.*, 644 (1993) 307-314.

- 1090 Volmer, D., Preiss, A., Levsen, K. and Wünsch, G.: Thermospray mass spectral studies of pesticides. Temperature and salt concentration effects on the ion abundances in thermospray mass spectra. *J. Chromatogr.*, 647 (1993) 235-259.

29a. *General techniques*

- 1091 Hogendoorn, E.A., Dijkman, E., Gort, S.M., Hoogerbrugge, R., van Zoonen, P. and Brinkman, U.A.T.: General strategy for multiresidue analysis of polar pesticides in ground water using coupled-column reversed-phase LC and step-gradient elution. *J. Chromatogr. Sci.*, 31 (1993) 433-439.

For additional information see C.A.:  
119 (1993) 158488a, 175460c.

See also 1382.

29b. *Chlorinated insecticides*

- 1092 Carabias Martínez, R., Rodríguez Gonzalo, E., Garay García, F. and Hernández Méndez, J.: Automated high-performance liquid chromatographic method for the determination of organophosphorus pesticides in waters with dual electrochemical (reductive-oxidative) detection. *J. Chromatogr.*, 644 (1993) 49-58.
- 1093 Newsome, W.H., Andrews, P., Conacher, H.B.S., Rao, R.C. and Chatt, A.: Total organochlorine content of fish from the Great Lakes. *J. Assoc. Off. Anal. Chem.*, 76 (1993) 703-706.

29c. *Phosphorus insecticides*

- 1094 Dhoot, J.S. and del Rosario, A.R.: Determination and confirmation of benomyl and carbendazim in water using high-performance liquid chromatography and diode array detection. *J. Chromatogr.*, 645 (1993) 178-181.

See also 116, 1381.

29d. *Carbamates*

- 1095 Ali, M.S., White, J.D., Bakowski, R.S., Stapleton, N.K., Williams, K.A., Johnson, R.C., Phillippo, E.T., Woods, R.W. and Ellis, R.L.: Extension of a liquid chromatographic method for N-methylcarbamate pesticides in cattle, swine, and poultry liver. *J. Assoc. Off. Anal. Chem.*, 76 (1993) 907-910.
- 1096 Bushway, R.J., Paradis, L.R., Perkins, L.B., Fan, T.S., Young, B.E.S. and Walsér, P.E.: Determination of 2-benzimidazolecarbamate in wine by competitive inhibition enzyme immunoassay. *J. Assoc. Off. Anal. Chem.*, 76 (1993) 851-856.
- 1097 Draper, W.M. and Wakeham, D.E.: Rate constants for metan-sodium cleavage and photodecomposition in water. *J. Agric. Food Chem.*, 41 (1993) 1129-1133.
- 1098 Howard, A.L., Braue, C. and Taylor, L.T.: Feasibility of thiocarbamate pesticide analysis in apples by supercritical fluid extraction and high-performance liquid chromatography. *J. Chromatogr. Sci.*, 31 (1993) 323-329.
- 1099 Simon, V.A., Pearson, K.S. and Taylor, A.: Determination of N-methylcarbamates and N-methylcarbamoyloximes in water by high-performance liquid chromatography with the use of fluorescence detection on a simple o-phthalaldehyde post-column reaction. *J. Chromatogr.*, 643 (1993) 317-320.

See also 43.

29e. *Herbicides*

- 1100 Bagheri, H., Slobodnik, J., Recasens, R.M.M., Ghijsen, R.T. and Brinkman, U.A.T.: Liquid chromatography - particle beam mass spectrometry for identification of unknown pollutants in water. *Chromatographia*, 37 (1993) 159-167.
- 1101 Balinova, A.: Solid-phase extraction followed by high-performance liquid chromatographic analysis for monitoring herbicides in drinking water. *J. Chromatogr.*, 643 (1993) 203-207.
- 1102 Corasaniti, M.T. and Nisticò, G.: Determination of paraquat in rat brain by high-performance liquid chromatography. *J. Chromatogr.*, 643 (1993) 419-425.
- 1103 Cserhádi, T. and Forgács, E.: Separation of some chlorophenoxyacetic acid congeners on a porous graphitized carbon column. *J. Chromatogr.*, 643 (1993) 331-336.
- 1104 Geerdink, R.B., Kienhuis, P.G.M. and Brinkman, U.A.T.: Fast screening method for eight phenoxyacid herbicides and bentazone in water. Optimization procedures for flow-injection analysis-thermospray tandem mass spectrometry. *J. Chromatogr.*, 647 (1993) 329-339.
- 1105 Tekel, J. and Kovačičová, J.: Chromatographic methods in the determination of herbicide residues in crops, food and environmental samples. *J. Chromatogr.*, 643 (1993) 291-303 - a review with 117 refs.
- 1106 Worobey, B.L.: Liquid chromatographic method for determination of diquat and paraquat herbicide in potatoes: collaborative study. *J. Assoc. Off. Anal. Chem.*, 76 (1993) 881-887.

See also 1374.

29f. *Fungicides*

- 1107 Barceló, D., Durand, G. and Vreeken, R.J.: Determination of quaternary amine pesticides by thermospray mass spectrometry. *J. Chromatogr.*, 647 (1993) 271-277.
- 1108 Ling, C.F., Melian, G.P., Jiminez-Conde, F. and Revilla, E.: High-performance liquid chromatographic analysis of carbofuran residues in tomatoes grown in hydroponics. *J. Chromatogr.*, 643 (1993) 351-355.

29g. *Other types of pesticides and various agrochemicals*

- 1109 Dardoize, F., Goasdoue, N., Goasdoue, C. and Couffignal, R.: High-performance liquid chromatography of chemical hybridizing agent in wheat. *J. Liq. Chromatogr.*, 16 (1993) 1517-1528.

30. SYNTHETIC AND NATURAL DYES

30a. *Synthetic dyes*

For additional information see C.A.:  
119 (1993) 137735p.

See also 255, 1369.



## 30b. Chloroplast and other natural pigments

- 1110 Ciavatta, C. and Govi, M.: Use of insoluble polyvinylpyrrolidone and isoelectric focusing in the study of humic substances in soils and organic wastes. *J. Chromatogr.*, 643 (1993) 261-270 - a review with 89 refs.
- 1111 De Leenheer, A.P., Lambert, W.E. and Meyer, E.: Chromatography of retinoids. *Clin. Dermatol.*, 5(Retinoids) (1993) 551-568; *C.A.*, 119 (1993) 90328r - a review with 59 refs.
- 1112 Hermans-Lokkerbol, A.C.J., van der Heijden, R. and Verpoorte, R.: Solvent system selection for separation of anthraquinones by means of centrifugal partition chromatography; application to an extract of a *Rubia tinctorum* hairy root culture. *J. Liq. Chromatogr.*, 16 (1993) 1433-1451.
- 1113 Inamura, I. and Uchida, K.: Gel chromatography of chlorophyllin (copper-sodium salt) in water and in aqueous poly(N-vinylpyrrolidone) solution. *Bull. Chem. Soc. Jpn.*, 66 (1993) 1828-1829; *C.A.*, 119 (1993) 147284e.
- 1114 Inoue, H., Imai, M., Naemura, T., Furuya, K. and Shizuri, Y.: Preparation and determination of zinc(II) chlorophylls by reversed-phase high-performance liquid chromatography. *J. Chromatogr.*, 645 (1993) 259-264.
- 1115 Ito, H., Tanaka, Y., Tsuji, H. and Tanaka, A.: Conversion of chlorophyll b to chlorophyll a by isolated cucumber etioplasts. *Arch. Biochem. Biophys.*, 306 (1993) 148-151.
- 1116 Marsili, R. and Callahan, D.: Comparison of a liquid solvent extraction technique and supercritical fluid extraction for the determination of  $\alpha$ - and  $\beta$ -carotene in vegetables. *J. Chromatogr. Sci.*, 31 (1993) 422-428.
- 1117 Parker, R.S.: Analysis of carotenoids in human plasma and tissues. *Methods Enzymol.*, 214(Carotenoids, Pt. B) (1993) 86-93; *C.A.*, 119 (1993) 112586c.
- 1118 Saitoh, K., Awaka, I. and Suzuki, N.: Separation of chlorophyll  $c_1$  and  $c_2$  by reversed-phase high-performance liquid chromatography. *J. Chromatogr.*, 653 (1993) 247-251.
- 1119 Schmitz, H.H., Poor, C.L., Gugger, E.T. and Erdman, J.W., Jr.: Analysis of carotenoids in human and animal tissues. *Methods Enzymol.*, 214(Carotenoids, Pt. B) (1993) 102-116; *C.A.*, 119 (1993) 112587d.
- See also 993, 1005, 1013, 1017, 1027, 1367.
31. PLASTICS AND THEIR INTERMEDIATES
- 1120 Auriola, S.O.K., Rönkkö, K.M. and Urtti, A.: Determination of polyethylene glycols by high-performance liquid chromatography-thermospray mass spectrometry. *J. Pharm. Biomed. Anal.*, 11 (1993) 1027-1032.
- 1121 Balke, S.T., Thitiratsakul, R., Lew, R., Cheung, P. and Mourey, T.H.: A strategy for interpreting multidetector size-exclusion chromatography data. II. Applications in plastic waste recovery. *ACS Symp. Ser.*, 521 (1993) 199-219; *C.A.*, 119 (1993) 182307m.
- 1122 Buerger, H.M., Mueller, H.M., Seebach, D., Boernsen, K.O., Schaer, M. and Widmer, H.M.: Matrix-assisted laser desorption and ionization as a mass spectrometric tool for the analysis of poly((R)-3-hydroxybutanoates). Comparison with gel permeation chromatography. *Macromolecules*, 26 (1993) 4783-4790; *C.A.*, 119 (1993) 118267w.
- 1123 DeGroot, A.W. and Hamre, W.J.: Characterization of high-molecular-mass polyethylenes by gel permeation chromatography-low-angle laser-light scattering. *J. Chromatogr.*, 648 (1993) 33-39.
- 1124 Goldwasser, J.M.: Absolute  $M_n$  determined by gel permeation chromatography-differential viscometry. *ACS Symp. Ser.*, 521 (1993) 243-251; *C.A.*, 119 (1993) 118266v.
- 1125 Gores, F. and Kitz, P.: Copolymer characterization using conventional size-exclusion chromatography and molar-mass-sensitive detectors. *ACS Symp. Ser.*, 521 (1993) 122-128; *C.A.*, 119 (1993) 161204k.
- 1126 Guarini, A., Guglielmetti, G. and Po, R.: Application of liquid chromatography-thermospray mass spectrometry to the analysis of polyester oligomers. *J. Chromatogr.*, 647 (1993) 311-318.
- 1127 Hunkeler, D., Macko, T. and Berek, D.: Critical conditions in the liquid chromatography of polymers. *ACS Symp. Ser.*, 521 (1993) 90-102; *C.A.*, 119 (1993) 161202h.
- 1128 Jenke, D.R.: Evaluation of various solvent systems for modeling solute sorption with plasticized polyvinyl chloride materials. *J. Pharm. Sci.*, 82 (1993) 617-621.
- 1129 Kuehn, A., Foerster, S., Loesch, R., Rommelfanger, M., Rosenauer, C. and Schmidt, M.: Size-exclusion chromatography of cationic and anionic polyelectrolytes in aqueous media. *Macromol. Chem., Rapid Commun.*, 14 (1993) 433-438; *C.A.*, 119 (1993) 96632g.
- 1130 Lehtinen, A. and Jakosuo-Jansson, H.: Molecular-weight determination of poly(4-methyl-1-pentene). *ACS Symp. Ser.*, 521 (1993) 277-291; *C.A.*, 119 (1993) 161206n.
- 1131 Mori, S. and Nishimura, Y.: Effects of addition of an electrolyte on size exclusion chromatography of polyamides using hexafluoro-2-propanol as mobile phase. *J. Liq. Chromatogr.*, 16 (1993) 3359-3370.
- 1132 Nikolov, R., Angelova, N. and Pishev, D.: (Chromatographic determination of oligomer content in aqueous solutions during caprolactam regeneration). *Khim. Volokna*, (1992) 57-59; *C.A.*, 119 (1993) 97809g.
- 1133 Pang, S. and Rudin, A.: Size-exclusion chromatographic assessment of long-chain branch frequency in polyethylenes. *ACS Symp. Ser.*, 521 (1993) 254-269; *C.A.*, 119 (1993) 161143q.
- 1134 Pang, S. and Rudin, A.: Size-exclusion chromatographic measurement of PVC molecular-weight distribution. *J. Appl. Polym. Sci.*, 49 (1993) 1189-1196; *C.A.*, 119 (1993) 161210j.
- 1135 Pasch, H.: Chromatographic investigation of macromolecules in the critical range of liquid chromatography. 5. Characterization of block copolymers of decyl and methyl methacrylate. *Macromol. Chem.*, 194 (1993) 2533-2541; *C.A.*, 119 (1993) 181696a.
- 1136 Sanayei, R.A., O'Driscoll, K.F. and Rudin, A.: Single-parameter universal calibration curve. *ACS Symp. Ser.*, 521 (1993) 103-112; *C.A.*, 119 (1993) 140157g.
- 1137 Shetty, S.S. and Garcia-Rubio, L.H.: Size-exclusion chromatography and end-group analysis of poly(methyl methacrylate). *ACS Symp. Ser.*, 521 (1993) 149-177; *C.A.*, 119 (1993) 140118v.

- 1138 Staal, W.J., Cools, P., van Herk, A.M. and German, A.L.: On-line extraction of polymers, oligomers, additives and monomers by multiple solvents on packed HPLC columns. *Chromatographia*, 37 (1993) 218-220.
- 1139 Szesztay, M. and Ghadir, M.: Gel permeation chromatography investigation of polyimide-siloxane copolymers: calibration procedure. *Angew. Macromol. Chem.*, 209 (1993) 111-117; *C.A.*, 119 (1993) 181691v.
- 1140 Trathnigg, B. and Yan, X.: Polymer analysis using size-exclusion chromatography with coupled density and refractive index detection. VI. Molecular mass dependence of preferential solvation of polyoxyethylenes in chloroform with different ethanol contents. *J. Chromatogr.*, 653 (1993) 199-206.
- 1141 Trathnigg, B., Thamer, D., Yan, X. and Kinugasa, S.: Analysis of polyethers by isocratic HPLC with universal detectors. I. Optimization of chromatographic conditions. *J. Liq. Chromatogr.*, 16 (1993) 2439-2452.
- 1142 Trathnigg, B., Thamer, D., Yan, X. and Kinugasa, S.: Analysis of polyethers by isocratic HPLC with universal detectors. II. Quantitative analysis and determination of molecular weight averages. *J. Liq. Chromatogr.*, 16 (1993) 2453-2467.
- 1143 Valic, S., Rakvin, B., Ranogajec, F., Vekslj, Z. and Grubisic-Galot, Z.: Study of irradiated polymers using double-modulation electron spin resonance and size exclusion chromatography. *Polymer*, 34 (1993) 2929-2933; *C.A.*, 119 (1993) 181763v.
- 1144 Van Asten, A.C., Venema, E., Kok, W.T. and Poppe, H.: Use of thermal field flow fractionation for the fractionation of polybutadiene in various organic solvents. *J. Chromatogr.*, 644 (1993) 83-94.
- 1145 Van Dam, J. and Daenens, P.: Molecular weight identification of polyethylene glycols in pharmaceutical preparations by gel permeation chromatography. *J. Pharm. Sci.*, 82 (1993) 938-941.
- 1146 Wu, C.S., Curry, J.F. and Senak, L.: Size-exclusion chromatography of cationic, nonionic, and anionic copolymers of vinylpyrrolidone. *ACS Symp. Ser.*, 521 (1993) 292-301; *C.A.*, 119 (1993) 140176n.
- For additional information see *C.A.*:  
119 (1993) 97756n, 118379j, 118380c, 181838y.
- See also 60, 82, 112, 167, 218, 1391.
32. DRUG ANALYSIS
- 32a. Drug analysis, general techniques
- 1147 Blom, Y. and Heldin, E.: Direct resolution of racemic drugs using cellulase silica as a chiral stationary phase. *J. Chromatogr.*, 653 (1993) 138-143.
- 1148 Bogusz, M., Franke, J.P., de Zeeuw, R.A. and Erkens, M.: An overview on standardisation of chromatographic methods for screening analysis in toxicology by means of retention indices and secondary standards. Part II. High-performance liquid chromatography. *Fresenius J. Anal. Chem.*, 347 (1993) 73-81.
- 1149 Campíns-Falcó, P., Herráez-Hernández, R. and Sevilla-Cabeza, A.: Column-switching techniques for high-performance liquid chromatography of drugs in biological samples. *J. Chromatogr.*, 619 (1993) 177-190.
- 1150 Chen, X.-H., Franke, J.-P., Wijsbeek, J. and de Zeeuw, R.A.: Reusability of Bond Elut Certify columns for the extraction of drugs from plasma. *J. Chromatogr.*, 619 (1993) 137-142.
- 1151 Covey, T. and Shushan, B.: The growing role of atmospheric pressure ionization LC/MS/MS in drug metabolism, and biopharmaceutical analysis. *Nippon Iyo Masu Supekutoru Gakkai Koenshu*, 16 (1991) 27-18; *C.A.*, 119 (1993) 85148c - a review with no refs.
- 1152 Guillemin, C.L.: Validation of chromatographic result on line or in laboratory. *Analisis*, 21 (1993) M38-M43; *C.A.*, 119 (1993) 151350r - a review with 6 refs.
- 1153 Gustavo Gonzalez, A. and Asuero, A.G.: Application of the extended Hildebrand solubility parameter treatment for optimizing reverse-phase high-performance liquid chromatography determination of pharmaceuticals. *Int. J. Pharm.*, 93 (1993) 183-188; *C.A.*, 119 (1993) 125291c.
- 1154 Haginaka, J.: (Multi-functional packing materials for high-performance liquid chromatography). *Yakugaku Kenkyu no Shinpo*, 9 (1993) 121-131; *C.A.*, 119 (1993) 108273f - a review with 35 refs.
- 1155 Haginaka, J., Murashima, T., Fujima, H. and Wada, H.: Direct injection assay of drug enantiomers in serum on ovomucoid-bonded silica materials by liquid chromatography. *J. Chromatogr.*, 620 (1993) 199-204.
- 1156 Kojima, T., Yashiki, M. and Miyazaki, T.: (Mass chromatographic detection of various drugs in biological materials). *Nippon Iyo Masu Supekutoru Gakkai Koenshu*, 16 (1991) 157-160; *C.A.*, 119 (1993) 110700m.
- 1157 Konishi, M., Okuno, K. and Hashimoto, H.: (Studies on interaction between cyclodextrins and drugs by high-performance liquid chromatography). *Kuromatogurafii*, 13 (1992) 273-274; *C.A.*, 119 (1993) 103098t.
- 1158 Kuchar, M. and Jelinkova, M.: Some problems of the chromatographic evaluation of lipophilicity. In: Kuchar, M. (Editor), *QSAR Des. Bioact. Compd.*, Prous, Barcelona, 1992, pp. 65-81; *C.A.*, 119 (1993) 155150y.
- 1159 Lu, M.: (Isolation of the dissociation compounds in RP-LC). *Yaowu Fenxi Zazhi*, 13 (1993) 206-210; *C.A.*, 119 (1993) 188684d - a review with 38 refs.
- 1160 Mahuzier, G., Prognon, P., Sargi, R., Kouwalti, H., Tod, M. and Farinotti, R.: (Luminescence and detection in liquid chromatography. Part I. Modification of the analytes environment). *Ann. Pharm. Fr.*, 51 (1993) 135-153; *C.A.*, 119 (1993) 146689d - a review with 63 refs.
- 1161 Park, M.K., Park, J.H., Cho, J.H., Kim, N.Y. and Kang, J.S.: Chemometric tool of chromatographic pattern recognition for the analysis of complex mixtures. *Arch. Pharmaceut. Res.*, 15 (1992) 376-378; *C.A.*, 119 (1993) 188689j.
- 1162 Pijnenburg, C.C., Teulings, C.J.A., Philipse, J.C.A., Barella, C.G.J., Conemans, J.M.H. and Duchateau, A.M.J.A.: (Separation of pharmaceuticals on end-capped octadecyl silyl HPLC columns: more than trial and error). *Ziekenhuisfarmacie*, 9 (1993) 24-28; *C.A.*, 119 (1993) 125287f.
- 1163 Santoro, M.I.R.M.: (Liquid chromatography with chiral stationary phases: New approach in drug regulatory analysis). *Rev. Farm. Bioquim. Univ. Sao Paulo*, 28 (1992) 1-29; *C.A.*, 119 (1993) 146684y - a review with 77 refs.

- 1164 Simonian, M.H., Dinh, S. and Fay, L.A.: Comparison of multi-component drug analysis by HPLC and UV/VIS spectroscopy. *Spectroscopy (Eugene)*, 8 (1993) 37-42; C.A., 119 (1993) 188716r.
- 1165 Steur, W.: A critical review of high-performance liquid chromatography, supercritical fluid chromatography and capillary zone electrophoresis in drug analysis. *Paperback APV*, 32 (1993) 95-105; C.A., 119 (1993) 146685z - a review with 18 refs.
- 1166 Sun, S.F. and Hsiao, C.L.: Hummel-Dreyer method in high-performance liquid chromatography for the determination of drug-protein binding parameters. *J. Chromatogr.*, 648 (1993) 325-331.
- 1167 Tiller, P.R.: Application of particle-beam mass spectrometry to drugs. An examination of the parameters affecting sensitivity. *J. Chromatogr.*, 647 (1993) 101-105.

For additional information see C.A.:  
119 (1993) 146690x.

See also 9, 28, 121, 127, 137, 155, 188, 198.

### 32b. Antirheumatics and antiinflammatory drugs

- 1168 Aggarwal, N.D., Goldman, D., Malick, A.W. and Sethachutkul, K.: Degradation profile and reversed-phase LC method development of the antiinflammatory drug, Ro 24-5913. *J. Pharm. Biomed. Anal.*, 11 (1993) 1037-1043.
- 1169 Avgerinos, A., Karidas, T. and Malamataris, S.: Extractionless high-performance liquid chromatographic method for the determination of diclofenac in human plasma and urine. *J. Chromatogr.*, 619 (1993) 324-329.
- 1170 Carretero, I., Morell, M. and Laserna, J.J.: Determination of dipyron metabolites in human plasma by micellar liquid chromatography. *J. Liq. Chromatogr.*, 16 (1993) 2767-2775.
- 1171 Högger, P. and Rohdewald, P.: Pharmacokinetics of bromfenac in healthy subjects after single oral administration of three different doses. *Arzneim.-Forsch.*, 43 (1993) 1114-1118.
- 1172 Imai, T., Fukuhara, A., Ueda, I. and Otagiri, M.: An evaluation of an anti-inflammatory-histamine H<sub>2</sub> antagonist drug complex on gastric erosions in the rat. *J. Pharmacol. Exp. Ther.*, 265 (1993) 328-333.
- 1173 Karidas, T., Avgerinos, A. and Malamataris, S.: Extractionless HPLC method for the determination of naproxen in human plasma and urine. *Anal. Lett.*, 26 (1993) 2341-2348.
- 1174 Lemko, C.H., Caillé, G. and Foster, R.T.: Stereospecific high-performance liquid chromatographic assay of ibuprofen: improved sensitivity and sample processing efficiency. *J. Chromatogr.*, 619 (1993) 330-335.
- 1175 Lopez-Bustamante, L.G., Troconiz, J.I.F. and Fos, D.: Tenoxicam: acute dose-dependent disposition studies in rats. *J. Pharm. Sci.*, 82 (1993) 851-853.
- 1176 Nicoll-Griffith, D., Scartozzi, M. and Chiem, N.: Automated derivatization and high-performance liquid chromatographic analysis of ibuprofen enantiomers. *J. Chromatogr.*, 653 (1993) 253-259.
- 1177 Paliwal, J.K., Smith, D.E., Cox, S.R., Berardi, R.R., Dunn-Kuchar, V.A. and Elta, G.H.: Stereoselective, competitive, and nonlinear plasma protein binding of ibuprofen enantiomers as determined *in vivo* in healthy subjects. *J. Pharmacokin. Biopharm.*, 21 (1993) 145-161.

- 1178 Schmitz, G., Lepper, H. and Estler, C.-J.: High-performance liquid chromatographic method for the routine determination of diclofenac and its hydroxy and methoxy metabolites from *in vitro* systems. *J. Chromatogr.*, 620 (1993) 158-163.
- 1179 Schulz, M. and Schmoldt, A.: Non-interaction of ranitidine with hydroxylation of diclofenac in rat and human liver microsomes. *Pharmazie*, 48 (1993) 625-626.
- 1180 Spraul, M., Hofmann, M., Wilson, I.D., Lenz, E., Nicholson, J.K. and Lindon, J.C.: Coupling of HPLC with <sup>19</sup>F- and <sup>1</sup>H-NMR spectroscopy to investigate the human urinary excretion of flurbiprofen metabolites. *J. Pharm. Biochem. Anal.*, 11 (1993) 1009-1015.
- 1181 Tsai, S.-W. and Wei, H.-J.: Self-normalized analysis of lipase-catalyzed conversion of naproxen enantiomers. *J. Liq. Chromatogr.*, 16 (1993) 2993-3001.
- 1182 Walker, J.S., Knihnicki, R.D., Seideman, P. and Day, R.O.: Pharmacokinetics of ibuprofen enantiomers in plasma and suction blister fluid in healthy volunteers. *J. Pharm. Sci.*, 82 (1993) 787-790.

For additional information see C.A.:

119 (1993) 103507a, 125316q, 125323q, 125333t.

See also 91, 406, 1222, 1265.

### 32c. Autonomic and cardiovascular drugs

- 1183 Ademola, J.I., Chow, C.A., Wester, R.C. and Maibach, H.I.: Metabolism of propranolol during percutaneous absorption in human skin. *J. Pharm. Sci.*, 82 (1993) 767-770.
- 1184 Agra, A.M. and Huang, S.-M.: High-performance liquid chromatographic determination of a potent A1 receptor antagonist (DMP 811) in plasma. *J. Chromatogr.*, 620 (1993) 205-209.
- 1185 Aluri, J.B. and Stavchansky, S.: Determination of guaifenesin in human plasma by liquid chromatography in the presence of pseudoephedrine. *J. Pharm. Biomed. Anal.*, 11 (1993) 803-808.
- 1186 Bachert, E.L. and Fung, H.-L.: High-performance liquid chromatographic method for stability and pharmacokinetic studies of nicorandil. *J. Chromatogr.*, 619 (1993) 336-341.
- 1187 Barbato, F., Cappello, B., Grumetto, L. and Morrica, P.: Analysis of calcium channel blocking dihydropyridines by high performance liquid chromatography. *Farmaco*, 48 (1993) 417-426; C.A., 119 (1993) 85261j.
- 1188 Barnes, A.R. and Nash, S.: HPLC determination of bendrofluzide in capsules. *Int. J. Pharm.*, 94 (1993) 231-234; C.A., 119 (1993) 188728w.
- 1189 Bauer, J.F., Krogh, S.K., Chang, Z.L. and Wong, C.F.: Determination of minor impurities in terazosin hydrochloride by high-performance liquid chromatography. *J. Chromatogr.*, 648 (1993) 175-181.
- 1190 Bonet-Domingo, E., Medina-Hernandez, M.J. and Garcia-Alvarez-Coque, M.C.: A micellar liquid chromatographic procedure for the determination of amiloride, bendroflumethiazide, chlorthalidone, spironolactone and triamterene in pharmaceuticals. *J. Pharm. Biomed. Anal.*, 11 (1993) 711-716.
- 1191 Bozkurt, A., Basci, N.E., Isimer, A. and Kayaalp, S.O.: Determination of debrisoquine and 4-hydroxydebrisoquine in urine by high-performance liquid chromatography with fluorescence detection after solid-phase extraction. *J. Pharm. Biomed. Anal.*, 11 (1993) 745-749.

- 1192 Campíns-Falcó, P., Herráez-Hernández, R. and Sevillano-Cabeza, A.: Simple and sensitive reversed-phase liquid chromatographic assay for analysis of chlorthalidone in urine. *J. Liq. Chromatogr.*, 16 (1993) 2571-2581.
- 1193 Carlucci, G., di Guiseppe, E. and Mazzeo, P.: Simultaneous determination of enalapril maleate and hydrochlorothiazide in tablets by derivative UV spectrophotometry and high-performance liquid chromatography. *Int. J. Pharm.*, 93 (1993) 245-248; C.A., 119 (1993) 125354a.
- 1194 Chen, C.L., Roder, J.D., Sangiah, S., Chen, H., Berlin, K.D., Garrison, G.L., Scherlag, B.J., Lazzara, R. and Patterson, E.: High-performance liquid chromatographic determination of SAZ-VII-22, a novel antiarrhythmic agent, in dog plasma and urine. *Anal. Sci.*, 9 (1993) 429-431; C.A., 119 (1993) 85283t.
- 1195 Costa, P., Bressole, F., Jarroux, E., Sarrazin, B., Mosser, J., Navratil, H. and Galtier, M.: Multiple-dose pharmacokinetics of moxislyte after oral administration to healthy volunteers. *J. Pharm. Sci.*, 82 (1993) 968-971.
- 1196 Costa, P., Bressolle, F., Sarrazin, B., Mosser, J. and Galtier, M.: Pharmacokinetics of moxislyte in healthy volunteers after intravenous and intracavernous administration. *J. Pharm. Sci.*, 82 (1993) 729-733.
- 1197 Criscuoli, M., Lippi, A., Mengozzi, G., Sardelli, G., Subissi, A. and Giachetti, A.: Pharmacokinetics and pharmacodynamics of idrapril in rats, dogs, and humans. *Drug Metab. Disp.*, 21 (1993) 835-840.
- 1198 Dawidowicz, A.L., Rauckyte, T. and Rogalski, J.: The preparation of sorbents for the analysis of human antithrombin III by means of high performance affinity chromatography. *Chromatographia*, 37 (1993) 168-172.
- 1199 DeMontigny, P. and Dave, J.: Analytical method development and preformulation stability studies of L-665,871 ( $\alpha$ -agonist) in swine feed. *J. Pharm. Biomed. Anal.*, 11 (1993) 947-954.
- 1200 Doerge, D.R., Bajic, S. and Lowes, S.: Analysis of clenbuterol in human plasma using liquid chromatography/atmospheric-pressure chemical-ionization mass spectrometry. *Rapid Commun. Mass Spectrom.*, 7 (1993) 462-464; C.A., 119 (1993) 110715v.
- 1201 Doroudian, A., Yeleswaram, K., Rurak, D.W., Abbott, F.S. and Axelson, J.E.: Sensitive high-performance liquid chromatographic method for direct separation of labetalol stereoisomers in biological fluids using and  $\alpha_1$ -acid glycoprotein stationary phase. *J. Chromatogr.*, 619 (1993) 79-86.
- 1202 Erram, S.V. and Tipnis, H.P.: Simple and rapid liquid chromatographic determination of eleven  $\beta$ -adrenoceptor antagonists in pharmaceuticals and plasma and correlation of HPLC retention times with their lipid solubilities. *Indian Drugs*, 30 (1993) 195-204; C.A., 110 (1993) 125336w.
- 1203 He, J., Shibukawa, A., Nakagawa, T., Wada, H., Fujima, H., Imai, E. and Gooh, Y.: (Direct injection analysis of atenolol enantiomers in plasma using an achiral/chiral coupled column system). *Kuromatogurafi*, 13 (1992) 313-317; C.A., 119 (1993) 108278m.
- 1204 Hussein, H., Mitrovic, V. and Schlepfer, M.: Rapid and sensitive assay of dobutamine in plasma by high-performance liquid chromatography and electrochemical detection. *J. Chromatogr.*, 620 (1993) 164-168.
- 1205 Indrayanto, G. and Handayani, R.: Quantitative determination of amroxol hydrochloride in tablets. *J. Pharm. Biomed. Anal.*, 11 (1993) 781-784.
- 1206 Josefsson, M., Carlsson, B. and Norlander, B.: Fast chromatographic separation of (-)-menthyl chloroformate derivatives of some chiral drugs, with specific reference to amlodipine, on porous graphitic carbon. *Chromatographia*, 37 (1993) 129-132.
- 1207 Kelly, M.J., Chambers, J. and MacNicol, A.D.: Simple and rapid method for the determination of the diastereomers of difenacoum in blood and liver using high-performance liquid chromatography with fluorescence detection. *J. Chromatogr.*, 620 (1993) 105-112.
- 1208 Kitani, M., Miyamoto, G. and Odomi, M.: Stereoselective high-performance liquid chromatographic assay for the determination of OPC-18790 enantiomers in human plasma and urine. *J. Chromatogr.*, 620 (1993) 97-104.
- 1209 Kountourellis, J.E., Markopoulou, C.K. and Stratis, J.A.: Quantitative high performance liquid chromatographic determination of bampine combined with tricyclic antidepressants and/or antipsychotics in pharmaceutical formulations. *Anal. Lett.*, 26 (1993) 2171-2182.
- 1210 Lamiable, D., Vistelle, R., Trenque, T., Fay, R., Millart, H. and Choisy, H.: Sensitive high-performance liquid chromatographic method for the determination of coumarin in plasma. *J. Chromatogr.*, 620 (1993) 273-277.
- 1211 Lee, Y.H., Shim, C.K., Lee, M.H. and Kim, S.K.: HPLC determination of diltiazem and deacetyldiltiazem in rat plasma. *Yakche Hakhoechi*, 22 (1992) 317-321; C.A., 119 (1993) 85272p.
- 1212 Mathôt, R.A.A., Appel, S., van Schaick, E.A., Soudijn, W., Ierman, A.P. and Danhof, M.: High-performance liquid chromatography of the adenosine A<sub>1</sub> agonist N<sup>6</sup>-cyclopentyladenosine and the A<sub>1</sub> antagonist 8-cyclopentyltheophylline and its application in a pharmacokinetic study in rats. *J. Chromatogr.*, 620 (1993) 113-120.
- 1213 Miller, J.H. McB., Robert, J.L. and Sorensen, A.M.: Reversed-phase ion-pair liquid chromatographic method for determining the impurities of furosemide. *J. Pharm. Biomed. Anal.*, 11 (1993) 257-261.
- 1214 Miller, L. and Bergeron, R.: Analytical and preparative resolution of enantiomers of verapamil and norverapamil using a cellulose-based chiral stationary phase in the reversed-phase mode. *J. Chromatogr.*, 648 (1993) 381-388.
- 1215 Modi, N.B., Veng-Pedersen, P., Grahm, D.J. and Dow, R.J.: Application of a system analysis approach to population pharmacokinetics and pharmacodynamics of nicardipine hydrochloride in healthy males. *J. Pharm. Sci.*, 82 (1993) 705-713.
- 1216 Neubeck, M., Becker, C., Henke, D., Rösch, W., Spahn-Languth, H. and Mutschler, E.: Pharmacokinetics of dilevalol and its conjugates in man. Assay method for plasma, urine and bile samples and preliminary pharmacokinetic studies. *Arzneim.-Forsch.*, 43 (1993) 953-957.
- 1217 Padval, M.V. and Bhargava, H.N.: Liquid chromatographic determination of indapamide in the presence of its degradation products. *J. Pharm. Biomed. Anal.*, 11 (1993) 1033-1036.
- 1218 Rawson, T.E., VanGorp, K.A., Yang, J. and Kogan, T.P.: Separation of 21-(R)- and 21-(S)-argatroban: solubility and activity of the individual diastereoisomers. *J. Pharm. Sci.*, 82 (1993) 672-673.
- 1219 Rosell, G., Camacho, A. and Parra, P.: Direct enantiomeric separation of *cis*-(±)diltiazem in plasma by high-performance liquid chromatography with ovomucoid column. *J. Chromatogr.*, 619 (1993) 87-92.

- 1220 Rossato, P., Scandola, M. and Grossi, P.: Investigation into lacidipine and related metabolites by high-performance liquid chromatography-mass spectrometry. *J. Chromatogr.*, 647 (1993) 155-166.
- 1221 Sahartova, O., Shatz, V. and Kalvins, I.: HPLC analysis of milrionate and its analogues in plasma. *J. Pharm. Biomed. Anal.*, 11 (1993) 1045-1047.
- 1222 Sane, R.T., Gangrade, M.G., Bapat, V.V., Surve, S.R. and Chonkar, N.L.: Simultaneous high performance liquid chromatographic determination of combined dosages, acebutol hydrochloride and hydrochlorothiazide xipamide and propranol hydrochloride methocarbamol and ibuprofen and trihexylphenidyl hydrochloride. *Indian Drugs*, 30 (1993) 205-219; *C.A.*, 119 (1993) 125337x.
- 1223 Suzuki, T., Horikiri, Y., Mizobe, M. and Noda, K.: Sensitive determination of bisoprolol enantiomers in plasma and urine by high-performance liquid chromatography using fluorescence detection, and application to preliminary study in humans. *J. Chromatogr.*, 619 (1993) 267-273.
- 1224 Tenjarla, S.N. and Chow, D.S.-L.: Stability indicating HPLC assay of propranolol for transdermal permeation study. *J. Liq. Chromatogr.*, 16 (1993) 2899-2913.
- See also 106, 473, 1057, 1179, 1265.
- 32d. *Central nervous system drugs*
- 1225 Aboul-Enein, H.Y., Serignese, V. and Bojarski, J.: Simple chiral liquid chromatographic enantioseparation of some racemic anti-epileptic drugs. *J. Liq. Chromatogr.*, 16 (1993) 2741-2749.
- 1226 Altria, K.D. and Conolly, P.C.: On-line solution stability determination of pharmaceuticals by capillary electrophoresis. *Chromatographia*, 37 (1993) 176-178.
- 1227 Aravagiri, M., Marder, S.R., van Putten, T. and Midha, K.K.: Determination of risperidone in plasma by high-performance liquid chromatography with electrochemical detection: application to therapeutic drug monitoring in schizophrenic patients. *J. Pharm. Sci.*, 82 (1993) 447-449.
- 1228 Ascalone, V., Guinebault, P. and Rouchouse, A.: Determination of mizolastine, a new antihistaminic drug, in human plasma by liquid-liquid extraction, solid-phase extraction and column-switching techniques in combination with high-performance liquid chromatography. *J. Chromatogr.*, 619 (1993) 275-284.
- 1229 Buggé, C.J.L., Tucker, M.D., Garcia, D.B., Kvalo, L.T. and Wilhelm, J.A.: Determination of mirfentanil hydrochloride (A-2508.HCl) in human plasma by liquid chromatography-mass spectrometry. *J. Pharm. Biomed. Anal.*, 11 (1993) 809-815.
- 1230 Butter, J.J., Kraak, J.C. and Poppe, H.: Determination of enantiomers of bupivacaine in serum using an on-line coupled three column liquid chromatographic system. *J. Pharm. Biomed. Anal.*, 11 (1993) 225-232.
- 1231 Chan, K. and Jones, R.D.M.: Simultaneous determination of flumazenil, midazolam and metabolites in human biological fluids by liquid chromatography. *J. Chromatogr.*, 619 (1993) 154-160.
- 1232 Chow, A.T., Meek, P.D. and Jusko, W.J.: High-pressure liquid chromatographic assay of theophylline in dog feces following oral administration of sustained-release products. *J. Pharm. Sci.*, 82 (1993) 956-958.
- 1233 Deleu, D., Sarre, S., Ebinger, G. and Michotte, Y.: Simultaneous monitoring of levodopa, dopamine and their metabolites in skeletal muscle and subcutaneous tissue in different pharmacological conditions using microdialysis. *J. Pharm. Biomed. Anal.*, 11 (1993) 577-585.
- 1234 El Maanni, A., Combourieu, I., Bonini, M. and Creppy, E.E.: Fluoxetine, an antidepressant, and norfluoxetine, its metabolite, determined by HPLC with a C<sub>8</sub> column and ultraviolet detection. *Clin. Chem. (Washington)*, 39 (1993) 1749-1750.
- 1235 Gillespie, T.A., Eckstein, J.A., Nardella, P. and Coutant, J.E.: Determination of dolasetron and its reduced metabolite in human plasma by GC-MS and LC. *J. Pharm. Biomed. Anal.*, 11 (1993) 955-962.
- 1236 Gopal, D., Grinberg, N., Dowling, T., Perpall, H., Bicker, G. and Tway, P.: Validation of a separation of diastereomers in the pharmaceutical industry. *J. Liq. Chromatogr.*, 16 (1993) 1749-1768.
- 1237 Graham, S.J., Beaulieu, N. and Lovering, E.G.: Sensitive high pressure liquid chromatographic assay method for formoterol fumarate. *J. Liq. Chromatogr.*, 16 (1993) 1497-1503.
- 1238 Griffin, R.J. and Burka, L.T.: Metabolism and elimination of oxazepam in B6C3F1 and Swiss-Webster mice. *Drug Metab. Disp.*, 21 (1993) 918-926.
- 1239 Gupta, R.N.: An improved solid phase extraction procedure for the determination of antidepressants in serum by column liquid chromatography. *J. Liq. Chromatogr.*, 16 (1993) 2751-2765.
- 1240 Haraguchi, T. and Korolkovas, A.: (Spectroscopic and chromatographic identification of buspirone). *Rev. Farm. Bioquim. Univ. Sao Paulo*, 27 (1991) 172-183; *C.A.*, 119 (1993) 125326t.
- 1241 Holland, D.C., Munns, R.K., Roybal, J.E., Hurlbut, J.A. and Long, A.R.: Simultaneous determination of xylazine and its major metabolite, 2,6-dimethylaniline, in bovine and swine kidney by liquid chromatography. *J. Assoc. Off. Anal. Chem.*, 76 (1993) 720-724.
- 1242 Hussein, Z., Mulford, D.J., Bopp, B.A. and Granneman, G.R.: Differences in the stereoselective pharmacokinetics of pazinaclole (DN-2327), a new anxiolytic, and its active metabolite after intravenous and oral single doses to dogs. *Drug Metab. Disp.*, 21 (1993) 805-810.
- 1243 Ibrahim, F.B.: Simultaneous determination and separation of several barbiturates and analgesic products by ion-pair high-performance liquid chromatography. *J. Liq. Chromatogr.*, 16 (1993) 2835-2851.
- 1244 Jayyosi, Z., Villoutrix, J., Ziegler, J.M., Batt, A.M., de Maack, F., Siest, G. and Thomas, P.E.: Identification of cytochrome P-450 isozymes involved in the hydroxylation of dantrolene by rat liver microsomes. *Drug Metab. Disp.*, 21 (1993) 939-945.
- 1245 Kamali, F.: Determination of plasma diazepam and desmethyl diazepam by solid-phase extraction and reversed-phase high-performance liquid chromatography. *J. Pharm. Biomed. Anal.*, 11 (1993) 625-627.
- 1246 Le Solleu, H., Demotes-Mainard, F., Vincon, G. and Bannwarth, B.: The determination of bromazepam in plasma by reversed-phase high-performance liquid chromatography. *J. Pharm. Biomed. Anal.*, 11 (1993) 771-775.
- 1247 Liu, L.L., Wang, Z., Feng, X.T. and Gao, S.: (Column switching HPLC method for determination of dextropran, an active metabolite of dextromethorphan, in plasma). *Yaoxue Xuebao*, 28 (1993) 374-378; *C.A.*, 119 (1993) 85290t.

- 1248 Martens, J. and Banditt, P.: Validation of the analysis of carbamazepine and its 10,11-epoxide metabolite by high-performance liquid chromatography from plasma: comparison with gas chromatography and the enzyme-multiplied immunoassay technique. *J. Chromatogr.*, 620 (1993) 169-173.
- 1249 Murillo, I., Costa, J. and Salvá, P.: Determination of bupivacaine in human plasma by HPLC. *J. Liq. Chromatogr.*, 16 (1993) 3509-3514.
- 1250 Nakasa, H., Komiya, M., Ohmori, S., Rikihisa, T. and Kitada, M.: Rat liver microsomal cytochrome P-450 responsible for reductive metabolism of zonisamide. *Drug Metab. Disp.*, 21 (1993) 777-781.
- 1251 Nicoll-Griffith, D.A., Chauret, N., Yergey, J.A., Trimble, L.A., Favreau, L., Zamboni, R., Grossman, S.J., Drey, J. and Herold, E.: Characterization of verlukast metabolites arising from an epoxide intermediate produced with hepatic microsomes from  $\beta$ -naphthoflavone-treated rodents (P-4501A1). *Drug Metab. Disp.*, 21 (1993) 861-867.
- 1252 Noctor, T.A.G., Diaz-Perez, M.J. and Wainer, I.W.: Use of a human serum albumin-based stationary phase for high-performance liquid chromatography as a tool for the rapid determination of drug-plasma protein binding. *J. Pharm. Sci.*, 82 (1993) 675-676.
- 1253 Ohkubo, T., Shimoyama, R. and Sugawara, K.: High performance liquid chromatographic determination of levopromazine in human breast milk and serum using solid phase extraction. *Biomed. Chromatogr.*, 7 (1993) 227-228.
- 1254 Okamoto, M., Ohgami, Y. and Nakazawa, H.: Direct stereochemical resolution of SM-11044, a novel anti-asthmatic drug, and its stereoisomers using a chiral immobilized protein stationary phase. *J. Chromatogr.*, 653 (1993) 261-265.
- 1255 Papadoyannis, I., Zotou, A., Samanidou, V., Theodoridis, G. and Zougrou, F.: Comparative study of different solid-phase extraction cartridges in the simultaneous RP-HPLC analysis of morphine and codeine in biological fluids. *J. Liq. Chromatogr.*, 16 (1993) 3017-3040.
- 1256 Parodi, B., Caviglioli, G., Bachi, A., Cafaggi, S. and Romussi, G.: Herbal mixtures with claimed slimming activity: determination by TLC and HPLC of illegally added drugs. *Pharmazie*, 48 (1993) 678-681.
- 1257 Piperaki, S., Parissi-Poulou, M. and Koupparis, M.: A separation study of tricyclic antidepressant drugs by HPLC with  $\beta$ -cyclodextrin bonded stationary phase. *J. Liq. Chromatogr.*, 16 (1993) 3487-3508.
- 1258 Rop, P.P., Grimaldi, F., Bresson, M., Fornaris, M. and Viala, A.: Liquid chromatographic analysis of cocaine, benzoylecgonine, local anaesthetic agents and some of their metabolites in biological fluids. *J. Liq. Chromatogr.*, 16 (1993) 2797-2811.
- 1259 Ryan, T.W.: Identification and quantification of tricyclic antidepressants by UV-photodiode array detection with multicomponent analysis. *J. Liq. Chromatogr.*, 16 (1993) 1545-1560.
- 1260 Scaglione, F., Scanni, A., Tomirotti, M., Dimaiuta, M., Ferrari, P. and Fraschini, F.: Pharmacokinetics and bioavailability of metoclopramide nasal spray versus metoclopramide intravenous in healthy volunteers and cancer patients. *Arzneim.-Forsch.*, 43 (1993) 986-988.
- 1261 Seay, S.S., Aucoin, D.P. and Tyczkowska, K.L.: Rapid high-performance liquid chromatographic method for the determination of ketamine and its metabolite dehydronorketamine in equine serum. *J. Chromatogr.*, 620 (1993) 281-287.
- 1262 Shimada, K., Mitamura, K., Morita, M. and Hirakata, K.: Separation of the diastereomers of baclofen by high performance liquid chromatography using cyclodextrin as a mobile phase additive. *J. Liq. Chromatogr.*, 16 (1993) 3311-3320.
- 1263 Song, H. and Abramson, F.: Drug metabolism studies using "intrinsic" and "extrinsic" labels. A demonstration using  $^{15}\text{N}$  vs. C1 in midazolam. *Drug Metab. Disp.*, 21 (1993) 868-873.
- 1264 Studenberg, S.D. and Brouwer, K.L.R.: Effect of phenobarbital and *p*-hydroxyphenobarbital glucuronide on acetaminophen metabolites in isolated rat hepatocytes: use of a kinetic model to examine the rates of formation and egress. *J. Pharmacokin. Biopharm.*, 21 (1993) 175-194.
- 1265 Sun, S.F. and Hsiao, C.L.: Qualitative study of the competition of drugs in binding to serum albumin. *Chromatographia*, 37 (1993) 329-335.
- 1266 Sztruhár, I., Ladányi, L., Zalavári-Dósa, L. and Beck, I.: Liquid chromatographic resolution of the enantiomers of psychotropic drug levomepromazine (methotrimeprazine) with  $\beta$ -cyclodextrin-bonded chiral stationary phase. *J. Pharm. Biomed. Anal.*, 11 (1993) 263-266.
- 1267 Tai, H.L., Osiewicz, R.J. and Bofinger, D.P.: A salt-free isocratic high-pressure liquid chromatographic method for the quantitation of benzodiazepines in serum. *Clin. Biochem.*, 26 (1993) 165-171; *C.A.*, 119 (1993) 108288q.
- 1268 Tracqui, A., Mikail, I., Kintz, P. and Mangin, P.: Nonfatal prolonged overdosage of pyrimethamine in an infant: measurement of plasma and urine levels using HPLC with diode-array detection. *J. Anal. Toxicol.*, 17 (1993) 248-250; *C.A.*, 119 (1993) 151551g.
- 1269 Vargay, Z., Horvath, G., Korponay, J., Kovacs, G., Kalman Mathe, I. and Banki, A.: (Survey of the analysis of Selegiline). *Acta Pharm. Hung.*, 62 (1992) 212-217; *C.A.*, 119 (1993) 103479t.
- 1270 Volpicelli, S.A., Centorrino, F., Puopolo, P.R., Kando, J., Frankenburg, F.R., Baldessarini, R.J. and Flood, J.G.: Determination of clozapine, norclozapine, and clozapine-N-oxide in serum by liquid chromatography. *Clin. Chem. (Washington)*, 39 (1993) 1656-1659.
- 1271 Woolf, T.F., Pool, W.F., Bjorge, S.M., Chang, T., Goel, O.P., Purchase II, C.F., Schoreder, M.C., Kunze, K.L. and Trager, W.F.: Bioactivation and irreversible binding of the cognition activator tacrine using human and rat liver microsomal preparations. Species difference. *Drug Metab. Disp.*, 21 (1993) 874-882.
- 1272 Yang, S.K.: Reversed-phase high-performance liquid chromatographic separation of 3-O-acyloxazepam and its hydrolysis and alcoholysis products. *J. Liq. Chromatogr.*, 16 (1993) 2605-2615.
- For additional information see *C.A.*:  
119 (1993) 103483q, 130812c, 153510s, 173475n, 175493r, 175497v.
- See also 143, 995, 1171, 1209, 1212, 1222, 1353.
- 32e. *Chemotherapeutics (exc. cytostatics and antibiotics)*
- 1273 Benchaoui, H.A. and McKellar, Q.A.: Determination of rafoxanide and closantel in ovine plasma by high performance liquid chromatography. *Biochem. Chromatogr.*, 7 (1993) 181-183.

- 1274 Bergqvist, Y., Al Kabbani, J., Pettersson, C. and Huynh, N.-H.: Enantioselective high-performance liquid chromatographic determination of (SR)- and (RS)-mefloquine in plasma using N-benzyloxycarbonyl-glycyl-L-proline as chiral counter ion. *J. Chromatogr.*, 620 (1993) 217-224.
- 1275 Breda, M., Pianezzola, E. and Benedetti, M.S.: Determination of exemestane, a new aromatase inhibitor, in plasma by high-performance liquid chromatography with ultraviolet detection. *J. Chromatogr.*, 620 (1993) 225-231.
- 1276 Brennan, J.M., O'Dwyer, P.J., Ozols, R.F. and LaCreta, F.P.: High-performance liquid chromatographic determination of the S- and R-diastereoisomers of L-buthionine (SR)-sulfoximine in human plasma and urine. *J. Chromatogr.*, 620 (1993) 121-128.
- 1277 Carlucci, G., Mazzeo, P. and Fantozzi, T.: Determination of ofloxacin in pharmaceutical forms by high-performance liquid chromatography and derivative UV-spectrophotometry. *Anal. Lett.*, 26 (1993) 2193-2201.
- 1278 Chu, P.-S., Wang, R.Y., Brandt, T.A. and Weerasinghe, C.A.: Determination of albendazole-2-aminosulfone in bovine milk using high-performance liquid chromatography with fluorometric detection. *J. Chromatogr.*, 620 (1993) 129-135.
- 1279 Cuiqing, C., Xingrong, L. and Rujin, W.: High-performance liquid chromatographic method for the determination of norfloxacin glutamate and glucuronate in solid and liquid dosage forms and its application to stability testing. *J. Pharm. Biomed. Anal.*, 11 (1993) 717-721.
- 1280 Dash, A.K.: A liquid chromatographic method for the determination of tolnaftate in pharmaceutical formulations. *J. Pharm. Biomed. Anal.*, 11 (1993) 847-853.
- 1281 Del Pilar da Silva, M., Procopio, J.R. and Hernández, L.: Evaluation of the capability of different chromatographic systems for the monitoring of thimerosal and its degradation products by high-performance liquid chromatography with amperometric detection. *J. Chromatogr.*, 653 (1993) 267-273.
- 1282 Fischer, J., Kelly, M.T., Smyth, M.R. and Jandera, P.: Determination of ivermectin in bovine plasma by column-switching LC using on-line solid-phase extraction and trace enrichment. *J. Pharm. Biomed. Anal.*, 11 (1993) 217-223.
- 1283 Horie, M., Saito, K., Nose, N., Tera, M. and Nakazawa, H.: Confirmation of residual oxolinic acid, nalidixic acid and piromidic acid in fish by thermospray liquid chromatography-mass spectrometry. *J. Liq. Chromatogr.*, 16 (1993) 1463-1472.
- 1284 Hormazabal, V., Steffenak, I. and Yndestad, M.: Simultaneous extraction and determination of sulfadiazine and trimethoprim in medicated fish feed by high-performance liquid chromatography. *J. Chromatogr.*, 648 (1993) 183-186.
- 1285 Ichikawa, N., Naora, K., Hayashibara, M. and Iwamoto, K.: High-performance liquid chromatographic determination of nalidixic acid in rat serum, brain and cerebrospinal fluid. *J. Pharm. Biomed. Anal.*, 11 (1993) 993-997.
- 1286 Kondo, F., Tsai, C.E., Hamada, E., Lin, S.Y. and Saitanu, K.: A continuous, simple and rapid method for the detection, extraction and identification of residual antibacterial agents in meat. *Microbios*, 73 (1993) 237-247; *C.A.*, 119 (1993) 137748v.
- 1287 Kristl, A., Vesnaver, G., Mrhar, A. and Kozjek, F.: Evaluation of partitioning and solubility data for some guanine derivatives in terms of mutual miscibility of octanol and water phase. *Pharmazie*, 48 (1993) 608-610.
- 1288 MacKichan, J.J. and Hink, W.F.: High-performance liquid chromatographic determination of CGA-184699 (lufenuron) in dog and cat blood. *J. Liq. Chromatogr.*, 16 (1993) 2595-2604.
- 1289 Morley, J.A. and Elrod, L., Jr.: Determination of fluoroquinolone antibacterials as N-acyl derivatives. *Chromatographia*, 37 (1993) 295-299.
- 1290 Rajaonarison, J.F., Lacarelle, B., Catalin, J., Durand, A. and Cano, J.P.: Effect of anticancer drugs on the glucuronidation of 3'-azido-3'-deoxythymidine in human liver microsomes. *Drug Metab. Disp.*, 21 (1993) 823-829.
- 1291 Saleh, M.I. and Loh, H.K.: Liquid chromatographic assay of pyronaridine in plasma and blood. *Anal. Chim. Acta*, 282 (1993) 559-564.
- 1292 Seifart, H.I., Kruger, P.B., Parkin, D.P., van Jaarsveld, P.P. and Donald, P.R.: Therapeutic monitoring of antituberculosis drugs by direct in-line extraction on a high-performance liquid chromatography system. *J. Chromatogr.*, 619 (1993) 285-290.
- 1293 Shaikh, B. and Moats, W.A.: Liquid chromatographic analysis of antibacterial drug residues in food products of animal origin. *J. Chromatogr.*, 643 (1993) 369-378 - a review with 97 refs.
- 1294 Shiga, N. and Matano, O.: High-performance liquid chromatographic method for the determination of oxolinic acid residues in crops. *J. Chromatogr.*, 643 (1993) 311-315.
- 1295 Szuna, A.J. and Blain, R.W.: Determination of a new antibacterial agent (Ro 23-9424) by multidimensional high-performance liquid chromatography with ultraviolet detection and direct plasma injection. *J. Chromatogr.*, 620 (1993) 211-216.
- 1296 Uetrecht, J.P., Shear, N.H. and Zahid, N.: N-Chlorination of sulfamethoxazole and dapsone by the myeloperoxidase system. *Drug Metab. Disp.*, 21 (1993) 830-834.
- 1297 Weber, J.D. and Smedley, M.D.: Liquid chromatographic method for determination of sulfamethazine residues in milk: collaborative study. *J. Assoc. Off. Anal. Chem.*, 76 (1993) 725-729.
- 1298 Zimmerman, C.L., Iyer, K.R., Faudskar II, A.L. and Rimmel, R.P.: Glucuronidation as a transient intermediate metabolic step in the elimination of (-)-carbovir. Identification of (-)-carbovir-5'-O-glucuronide in rat bile. *Drug Metab. Disp.*, 21 (1993) 902-910.

For additional information see C.A.:

119 (1993) 103518e, 115648k, 151520w, 151697j.

See also 1031, 1077, 1265, 1326, 1338.

### 32f. Cytostatics

- 1299 Bishop, S.M., Khoo, B.J., McRobert, A.J., Simpson, M.S.C., Phillips, D. and Beeby, A.: Characterisation of the photochemotherapeutic agent disulphonated aluminium phthalocyanine and its high-performance liquid chromatographic separated components. *J. Chromatogr.*, 646 (1993) 345-350.
- 1300 Blanz, J., Renner, U., Schmeer, K., Ehninger, G. and Zeller, K.-P.: Detection and identification of human urinary metabolites of biantrazole (CI-941). *Drug Metab. Disp.*, 21 (1993) 955-961.
- 1301 Borel, A.G. and Abbott, F.S.: Identification of carbamoylated thiol conjugates as metabolites of the antineoplastic 1-(2-chloroethyl)-3-cyclohexyl-1-nitrosourea, in rats and humans. *Drug Metab. Disp.*, 21 (1993) 889-901.

- 1302 Canal, P., Attal, M., Chatelut, E., Guichard, S., Hugué, F., Müller, C., Schlaifer, D., Laurent, G., Houin, G. and Bugat, R.: Plasma and cellular pharmacokinetics of mitoxantrone in high-dose chemotherapeutic regimen for refractory lymphomas. *Cancer Res.*, 53 (1993) 4850-4854.
- 1303 Delprino, L., Giacomotti, M., Dosio, F., Brusa, P., Ceruti, M., Grosa, G. and Cattel, L.: Toxin-target design for anticancer therapy. II: preparation and biological comparison of different chemically linked gelonin-antibody conjugates. *J. Pharm. Sci.*, 82 (1993) 699-704.
- 1304 Ehhardt, W.J., Sullivan, H.R., Wood, P.G., Woodland, J.M., Hamilton, M., Hamilton, C., Cornpropst, D., Grindey, G.B., Worzalla, J.F., Bewley, J.R. et al.: Pharmacokinetics of the anticancer agent sulofenur in mice, rats, monkeys and dogs. *J. Pharm. Sci.*, 82 (1993) 683-688.
- 1305 Elfarra, A.A. and Hwang, I.Y.: Targeting of 6-mercaptopurine to the kidneys. Metabolism and kidney-selectivity of S-(6-puriny)-L-cysteine analogs in rats. *Drug Metab. Disp.*, 21 (1993) 841-845.
- 1306 Embree, L., Gelmon, K.A., Lohr, A., Mayer, L.D., Coldman, A.J., Cullis, P.R., Palaitis, W., Pilkiewicz, F., Hudon, N.J., Heggie, J.R. and Goldie, J.H.: Chromatographic analysis and pharmacokinetics of liposome-encapsulated doxorubicin in non-small-cell lung cancer patients. *J. Pharm. Sci.*, 82 (1993) 627-634.
- 1307 Gonnet, F., Lemaire, D., Kozelka, J. and Chottard, J.-C.: Isolation of *cis*-[PtCl(NH<sub>3</sub>)<sub>2</sub>(H<sub>2</sub>O)](ClO<sub>4</sub>), the monohydrated form of the anti-tumour drug cisplatin, using cation-exchange high-performance liquid chromatography. *J. Chromatogr.*, 648 (1993) 279-282.
- 1308 Hadfield, J.A., McGown, A.T., Dawson, M.J., Thatcher, N. and Fox, B.W.: The suitability of carboplatin solutions for 14-day continuous infusion by ambulatory pump: an HPLC-dynamic FAB study. *J. Pharm. Biomed. Anal.*, 11 (1993) 723-727.
- 1309 Hüttmann, A., Mross, K. and Hossfeld, D.K.: Determination of the new podophyllotoxin derivative NK 611 in plasma by high-performance liquid chromatography with ultraviolet detection. *J. Chromatogr.*, 620 (1993) 233-238.
- 1310 Ishikawa, T. and Ali-Osman, F.: Glutathione-associated *cis*-diamminedichloroplatinum(II) metabolism and ATP-dependent efflux from leukemia cells. Molecular characterization of glutathione-platinum complex and its biological significance. *J. Biol. Chem.*, 268 (1993) 20116-20125.
- 1311 Kagel, J.R., Stella, V.J. and Riley, C.M.: A liquid chromatographic method for the determination of the enantiomeric purity of the anticancer drug, 9-amino-20(S)-camptothecin. *J. Pharm. Biomed. Anal.*, 11 (1993) 793-802.
- 1312 Ketchum, R.E.B. and Gibson, D.M.: Rapid isocratic reversed phase HPLC of taxanes on new columns developed specifically for taxol analysis. *J. Liq. Chromatogr.*, 16 (1993) 2519-2530.
- 1313 Malecki, F., Sułkowska, J. and Weiss-Gradzinska, W.: Reversed-phase high-performance liquid chromatographic determination of melphalan in pharmaceuticals. *Chem. Anal. (Warsaw)*, 38 (1993) 255-258; C.A., 119 (1993) 1887-22q.
- 1314 Rochard, E., Boutelet, H., Griesemann, E., Barthes, D. and Courtois, P.: Simultaneous high performance liquid chromatographic analysis of carboplatin in infusion fluids. *J. Liq. Chromatogr.*, 16 (1993) 1505-1516.
- 1315 Sakagami, H., Sakagami, T., Takeda, M., Iwaki, K. and Takeda, K.: Determination of sodium 5,6-benzylidene-L-ascorbate and related compounds by high-performance liquid chromatography. *J. Chromatogr.*, 653 (1993) 37-43.
- 1316 Sakai, A., Mori, N., Shuto, S. and Suzuki, T.: Deacylation-reacylation cycle: a possible absorption mechanism for the novel lymphotropic antitumor agent dipalmitoylphosphatidylfluoridine in rats. *J. Pharm. Sci.*, 82 (1993) 575-578.
- 1317 Walton, M.I. and Workman, P.: Pharmacokinetics and bioreductive metabolism of the novel benzotriazine di-N-oxide hypoxic cell cytotoxin tirapazamine (WIN 59075; SR 4233; NSC 130181) in mice. *J. Pharmacol. Exp. Ther.*, 265 (1993) 938-947.
- 1318 Wickremesinhe, E.R.M. and Arteca, R.N.: Methodology for the identification and purification of taxol and cephalomannine from *Taxus callus* cultures. *J. Liq. Chromatogr.*, 16 (1993) 3263-3274.

For additional information see C.A.:  
119 (1993) 130836p.

See also 1032, 1037, 1350.

### 32g. Other drug categories

- 1319 Andersson, T., Lagerstrom, P.-O., Miners, J.O., Veronese, M.E., Weidolf, L. and Birkett, D.J.: High-performance liquid chromatographic assay for human liver microsomal omeprazole metabolism. *J. Chromatogr.*, 619 (1993) 291-297.
- 1320 Balestrieri, F., Marini, D. and Melissano, M.: (HPLC determination of urocanic acid in sunscreen products). *Cosmet Toiletries Ed. Ital.*, 13 (1992) 34-38; C.A., 119 (1993) 146317f.
- 1321 Camacho-Torralba, P.L., Beeson, M.D., Vigh, G. and Thompson, D.H.: High-performance chiral displacement chromatographic separations in the normal-phase mode. II. Separation of the enantiomers of 1,2-O-dihexadecyl-*rac*-glycerol-3-O-(3,5-dinitrophenyl)carbamate using the Pirkle-type naphthylalanine silica stationary phase. *J. Chromatogr.*, 646 (1993) 259-266.
- 1322 Capacio, B.R., Byers, C.E., Jackson, T.K. and Matthews, R.L.: An HPLC method for the determination of granisetron in guinea pig plasma. *J. Anal. Toxicol.*, 17 (1993) 151-155; C.A., 119 (1993) 108290j.
- 1323 Chellquist, E.M. and Searle, R.: An LC method for measuring complex formation equilibria by competitive chelation. *J. Pharm. Biomed. Anal.*, 11 (1993) 985-992.
- 1324 De Vries, J.X. and Schmitz-Kummer, E.: Direct column liquid chromatographic enantiomer separation of the coumarin anticoagulants phenprocoumon, warfarin, acenocoumarol and metabolites on an  $\alpha_1$ -acid glycoprotein chiral stationary phase. *J. Chromatogr.*, 644 (1993) 315-320.
- 1325 De Vries, J.X., Voss, A., Kutschker, C. and Reiter, S.: Simultaneous determination of allopurinol and oxipurinol in human plasma and urine by high-performance liquid chromatography. *Arzneim.-Forsch.*, 43 (1993) 1072-1075.
- 1326 Den Hartigh, J., Langebroek, R. and Vermeij, P.: Ion-exchange liquid chromatographic analysis of bisphosphonates in pharmaceutical preparations. *J. Pharm. Biomed. Anal.*, 11 (1993) 977-983.
- 1327 Dixon, C.M., Saynor, D.A., Andrew, P.D., Oxford, J., Bradbury, A. and Tarbit, M.H.: Disposition of sumatriptan in laboratory animals and humans. *Drug Metab. Disp.*, 21 (1993) 761-769.
- 1328 Haldna, U. and Kojs, J.: Determination of sodium cyclamate and saccharin by ion chromatography. *Eesti Tead. Akad. Toim., Keem.*, 41 (1992) 69-72; C.A., 119 (1993) 137747u.



- 1329 Ishiwata, K., Ishii, S. and Senda, M.: HPLC using physiological saline for the quality control of radiopharmaceuticals used in PET studies. *Appl. Radiat. Isot.*, 44 (1993) 1119-1124; *C.A.*, 119 (1993) 154999v.
- 1330 Kim, B.K., Kim, Y.S., Park, S.B., Rhee, J.S., Jung, K.H. and Kim, K.N.: (Microanalysis of pancuronium bromide in urine and blood by HPLC). *Yakhak Hoechi*, 37 (1993) 30-35; *C.A.*, 119 (1993) 112595e.
- 1331 Konfino, M., Deltcheva, S. and Mindjova, K.: The isolation of lovastatin and its determination by densitometric TLC and by HPLC. *J. Planar Chromatogr.*, 6 (1993) 404-406.
- 1332 Kuo, B.-S., Poole, J.C., Hwang, K.-K. and Cheng, H.: Pharmacokinetics and metabolic interconversion of intravenous 4-amino-5-chloro-2-[(methylsulfinyl)ethoxy]-N-[2-(diethylamino)ethyl]benzamide and its sulfide and sulfone: metabolites in rats. *J. Pharm. Sci.*, 82 (1993) 694-696.
- 1333 Mascher, H., Kikuta, C. and Weyhenmeyer, R.: Diastereomeric separation of free and conjugated silibinin in plasma by reversed phase HPLC after specific extraction. *J. Liq. Chromatogr.*, 16 (1993) 2777-2789.
- 1334 Matuszewski, B.K. and Constanzer, M.L.: Indirect chiral separation and analyses in human biological fluids of the stereoisomers of a thienothiopyran-2-sulfonamide (TRUSOPT), a novel carbonic anhydrase inhibitor with two chiral centers in the molecule. *Chirality*, 4 (1992) 515-519; *C.A.*, 119 (1993) 85254j.
- 1335 Miyamoto, E., Demizu, Y., Murata, Y., Yamada, Y., Kawashima, S., Kontani, H. and Sakai, T.: High-performance liquid chromatographic preparation of oxybutynin enantiomers on a chiral stationary phase. *J. Chromatogr.*, 653 (1993) 135-137.
- 1336 Naidong, W. and Lee, J.W.: Development and validation of a high-performance liquid chromatographic method for the quantitation of warfarin enantiomers in human plasma. *J. Pharm. Biomed. Anal.*, 11 (1993) 785-792.
- 1337 Nebinger, P. and Koel, M.: Determination of acyclovir by ultrafiltration and high-performance liquid chromatography. *J. Chromatogr.*, 619 (1993) 342-344.
- 1338 Santoni, G., Tonsini, A., Gratterer, P., Mura, P., Furlanetto, S. and Pinzauti, S.: Determination of atropine sulphate and benzalkonium chloride in eye drops by HPLC. *Int. J. Pharm.*, 93 (1993) 239-243; *C.A.*, 119 (1993) 125353z.
- 1339 Teraoka, R., Otsuka, M. and Matsuda, Y.: Effects of temperature and relative humidity on the solid-state chemical stability of ranitidine hydrochloride. *J. Pharm. Sci.*, 62 (1993) 601-604.
- 1340 Uejima, Y., Kokubo, M., Oshida, J.-I., Kawabata, H., Kato, Y. and Fujii, K.: 5-Methyl-4H-3,1-benzoxazin-4-one derivatives: specific inhibitors of human leukocyte elastase. *J. Pharmacol. Exp. Ther.*, 265 (1993) 516-523.
- 1341 Van Rhijn, J.A., Traag, W.A. and Heskamp, H.H.: Confirmatory analysis of clenbuterol using two different derivatives simultaneously. *J. Chromatogr.*, 619 (1993) 243-249.
- 1342 Wagh, J.S., Mokashi, A.A. and Datta, A.: High-performance liquid chromatographic method for the monitoring of the synthesis of the precursor for tetramisole. *J. Chromatogr.*, 646 (1993) 428-433.
- See also 138, 426, 617, 620, 963, 1014, 1040, 1046, 1058, 1071, 1082, 1193, 1207, 1228, 1256, 1276, 1369, 1388.
- 32h. *Toxicological and forensic applications*
- 1343 Huang, J.L. and Morgan, D.J.: Simple direct injection high-performance liquid chromatographic method to determine quinidine in plasma. *J. Chromatogr.*, 620 (1993) 278-280.
- 1344 Schafer, F., Schad, H. and Weber, L.: Determination of phenylmercapturic acid in urine of benzene-exposed BDF-1 mice. *J. Chromatogr.*, 620 (1993) 239-242.
- 1345 Ubbink, J.B., Lagendijk, J. and Vermaak, W.H.H.: Simple high-performance liquid chromatographic method to verify the direct barbituric acid assay for urinary cotinine. *J. Chromatogr.*, 620 (1993) 254-259.
- 1346 Ventura, R., Nadal, T., Alcalde, P. and Segura, J.: Determination of mesocarb metabolites by high-performance liquid chromatography with UV detection and with mass spectrometry using a particle-beam interface. *J. Chromatogr.*, 647 (1993) 203-210.
- 1347 Zaromb, S., Alcaraz, J., Lawson, D. and Woo, C.S.: Detection of airborne cocaine and heroin by high-throughput liquid-absorption preconcentration and liquid chromatography-electrochemical detection. *J. Chromatogr.*, 643 (1993) 107-115.
- See also 204, 337, 487, 917, 931, 951, 970, 1200, 1246, 1255.
- 32i. *Plant extracts*
- 1348 Buszewski, B., Kawka, S., Suprynowicz, Z. and Wolski, T.: Simultaneous isolation of Rutin and Esculin from plant material and drugs using solid-phase extraction. *J. Pharm. Biomed. Anal.*, 11 (1993) 211-215.
- 1349 Feige, G.B., Lumbsch, H.T., Huneck, S. and Eltx, J.A.: Identification of lichen substances by a standardized high-performance liquid chromatographic method. *J. Chromatogr.*, 646 (1992) 417-427.
- 1350 Harris, R.K. and Haggerty, W.J.: Assays for potentially anticarcinogenic phytochemicals in flaxseed. *Cereal Foods World*, 38 (1993) 147-151; *C.A.*, 119 (1993) 137673s.
- 1351 Hasler, F., Krapf, R., Brenneisen, R., Bourquin, D. and Krähenbühl, S.: Determination of 18 $\beta$ -glycyrrhetic acid in biological fluids from humans and rats by solid-phase extraction and high-performance liquid chromatography. *J. Chromatogr.*, 620 (1993) 73-82.
- 1352 Kocsis, A., Szabo, L.F. and Podanyi, B.: New bis-iridoids from *Dipsacus laciniatus*. *J. Natural Prod.*, 56 (1993) 1486-1499.
- 1353 Kraemer, W. and Wiartalla, R.: (Determination of naphthodianthrones (total hypericin) in St. John's-wort (*Hypericum perforatum* L.)). *PZ Wiss.*, 5 (1992) 202-207; *C.A.*, 119 (1993) 103446e.
- 1354 Lamaison, J.L., Carnat, A., Petitjean-Freytet, C. and Carnat, A.P.: (Study on the pharmaceutical quality of batches of dried flowering tops of hawthorn collected in Auvergne (France)). *Plant. Med. Phytother.*, 25 (1991) 177-183; *C.A.*, 119 (1993) 167850q.
- 1355 Maillard, M.P., Wolfender, J.-L. and Hostettmann, K.: Use of liquid chromatography-thermospray mass spectrometry in phytochemical analysis of crude plant extracts. *J. Chromatogr.*, 647 (1993) 147-154.
- For additional information see *C.A.*:  
119 (1993) 86313q, 102972m, 103454f, 125331r, 125343w, 130802z, 137736q, 146312a, 146697e, 146717m, 151528e, 188719u.

- 1356 Miyamoto, F. and Saeki, M.: (Determination of glycyrrhizic acid in foods by high-performance liquid chromatography). *Chiba-ken Eisei Kenkyusho Kenkyu Hokoku*, 16 (1992) 23-28; *C.A.*, 119 (1993) 179531z.
- 1357 Shen, Y.-C. and Chen, C.-H.: New secoiridoid dilactones from *Fraxinus uhdei*. *J. Natural Prod.*, 56 (1993) 1905-1911.
- 1358 Wolfender, J.L., Maillard, M. and Hostettmann, K.: Liquid chromatographic-thermospray mass spectrometric analysis of crude plant extracts containing phenolic and terpene glycosides. *J. Chromatogr.*, 647 (1993) 183-190.
- 1359 Zheng, R. and Li, G.: (Application of HPLC in study of the Chinese materia medica). *Zhongcaoyao*, 24 (1993) 265-268; *C.A.*, 119 (1993) 167845s - a review with 42 refs.

For additional information see *C.A.*:

119 (1993) 103082h, 103448g, 125311j, 125314n, 167847u, 175469u, 188694g, 188736x.

See also 965, 1318, 1367.

### 33. CLINICO-CHEMICAL APPLICATIONS

#### 33a. General papers and reviews

See 510.

#### 33b. Complex mixtures and profiling (single compounds by cross-reference only)

See 227, 258, 286, 306, 318, 324, 342, 350, 351, 353, 354, 382, 396, 398, 409, 410, 411, 415, 425, 455, 464, 470, 480, 481, 485, 491, 504, 507, 508, 509, 515, 517, 547, 631, 644, 664, 772, 788, 821, 827, 838, 884, 889, 892, 939, 943, 944, 945, 956, 979, 993, 1009, 1011, 1023, 1344, 1395.

### 34. FOOD ANALYSIS

#### 34a. General papers and reviews

- 1360 Coors, U.: Components in foods with a high fat content. In: Matissek, R. and Wittkowski, R. (Editors), *High Perform. Liq. Chromatogr. Food Control Res.*, Technomic, Lancaster, 1993, pp. 151-182; *C.A.*, 119 (1993) 137619d - a review with 134 refs.
- 1361 Hitos Natera, Maria del Pilar.: (Use of HPLC in the investigation of some problems related to international trading in wines). *Tec. Lab.*, 15 (1993) 260-261; *C.A.*, 119 (1993) 179412m - a review with no refs.
- 1362 Matissek, R.: Modern high performance liquid chromatography-fundamentals and developments. In: Matissek, R. and Wittkowski, R. (Editors), *High Perform. Liq. Chromatogr. Food Control Res.*, Technomic, Lancaster, 1993, pp. 19-61; *C.A.*, 119 (1993) 137616a - a review with 73 refs.
- 1363 Matissek, R. and Wittkowski, R. (Editors): *High Performance Liquid Chromatography in Food Control and Research*. Technomic, Lancaster, 1993, 383 p.; *C.A.*, 119 (1993) 138047c.

1364 Mehrens, H.A. and Reimerdes, D.: Components in foods with a high protein content. In: Matissek, R. and Wittkowski, R. (Editors), *High Perform. Liq. Chromatogr. Food Control Res.*, Technomic, Lancaster, 1993, pp. 119-150; *C.A.*, 119 (1993) 137618c - a review with 161 refs.

1365 Roessner, B.: Ions and ionic components. In: Matissek, R. and Wittkowski, R. (Editors), *High Perform. Liq. Chromatogr. Food Control Res.*, Technomic, Lancaster, 1993, pp. 235-267; *C.A.*, 119 (1993) 137621y - a review with 68 refs.

See also 236, 1105.

#### 34b. Complex mixtures (single compounds by cross-reference only)

1366 Biedermann, M., Grob, K., Frohlich, D. and Meier, W.: Short description of an LC-(LC)-GC method for detecting irradiation of fatty foods. *Comm. Eur. Communities, [Rep.] EUR*, EUR 14315, 1993, pp. 323-331; *C.A.*, 119 (1993) 137703b.

1367 Castellar, M.R., Montijano, H., Manjón, A. and Iborra, J.L.: Preparative high-performance liquid chromatographic purification of saffron secondary metabolites. *J. Chromatogr.*, 648 (1993) 187-190.

1368 Galensa, R.: Components in foods with a high carbohydrate content. In: Matissek, R. and Wittkowski, R. (Editors), *High Perform. Liq. Chromatogr. Food Control Res.*, Technomic, Lancaster, 1993, pp. 87-117; *C.A.*, 119 (1993) 137617b - a review with 85 refs.

1369 Koebler, H.: Analysis of food additives. In: Matissek, R. and Wittkowski, R. (Editors), *High Perform. Liq. Chromatogr. Food Control Res.*, Technomic, Lancaster, 1993, pp. 271-296; *C.A.*, 119 (1993) 137692x.

1370 Paetzold, R., Fenz, R., Krause, M. and Galensa, R.: (Rapid detection of maize as malt substitute in beer). *Dtsch. Lebensm.-Rundsch.*, 88 (1992) 390-391; *C.A.*, 119 (1993) 158471q.

1371 Ramos, F. and Noronha da Silveira, M.I.: (Food additives: II. Robenidine in feeds). *Bol. Fac. Farm. Coimbra*, 15 (1991) 61-66; *C.A.*, 119 (1993) 93836r.

For additional information see *C.A.*:

119 (1993) 938854v.

See also 157, 168, 238, 255, 261, 265, 292, 332, 333, 338, 339, 364, 365, 368, 372, 375, 377, 390, 419, 420, 482, 674, 676, 923, 926, 954, 959, 989, 991, 993, 1002, 1003, 1006, 1007, 1008, 1017, 1025, 1038, 1068, 1098, 1106, 1116, 1199, 1200, 1283, 1286, 1297, 1328, 1356, 1424, 1432, 1433.

#### 34c. Organoleptically important compounds (flavors, odors, volatiles)

1372 Demian, B.A.: Trace analysis of vanillin in tobacco. *J. Liq. Chromatogr.*, 16 (1993) 3563-3574.

## 35. ENVIRONMENTAL ANALYSIS

## 35a. General papers and reviews

- 1373 Grob, R.L.: Environmental analyses utilizing gas and liquid chromatography. *J. Liq. Chromatogr.*, 16 (1993) 1783-1802 - a review with 20 refs.
- 1374 Singh, R.P. and Chiba, M.: Determination of benomyl and its degradation products by chromatographic methods in water, wettable powder formulations, and crops. *J. Chromatogr.*, 643 (1993) 249-260 - a review with 51 refs.

See also 116, 212, 1102, 1103, 1105, 1430.

## 35b. Air pollution (complex mixtures; single compounds by cross-reference only)

- 1375 Colli, M., Zabardini, L., Melzi d'Eril, G.V. and Marchetti, R.: Evaluation of a modified Marcali technique with high-performance liquid chromatography-ultraviolet detection for the determination of 2,4-toluene diisocyanate in air. *J. Chromatogr.*, 643 (1993) 51-54.
- 1376 Dolan, J.W.: Air Problems. *LC-GC*, 11 (1993) 270-273; *C.A.*, 119 (1993) 194749v.

See also 211, 245, 459, 955, 1347, 1438.

## 35c. Water pollution (complex mixtures; single compounds by cross-reference only)

- 1377 Bächmann, K., Haag, I., Prokop, T., Röder, A. and Wagner, P.: Chromatographic methods for the analysis of size-classified and individual raindrops. *J. Chromatogr.*, 643 (1993) 181-188.
- 1378 Maeda, K., Kobayashi, M., Nakamura, M., Nagasawa, K., Hashimoto, H. and Onodera, S.: Reactor water quality monitoring by on-line ion chromatography. *Water Chem. Nucl. React. Syst.*, 6 (1992) 214-218; *C.A.*, 119 (1993) 190357t.
- 1379 Sasurain, J. and Schwedt, G.: (Development of process ion chromatography for application in a waterworks). *Vom Wasser*, 80 (1993) 221-233; *C.A.*, 119 (1993) 124657q.
- 1380 Schröder, H.F.: Pollutants in drinking water and waste water. *J. Chromatogr.*, 643 (1993) 145-161.
- 1381 Suzuki, T., Yaguchi, K. and Kano, I.: Screening methods for asulam, oxine-copper and thiram in water by high-performance liquid chromatography after enrichment with a minicolumn. *J. Chromatogr.*, 643 (1993) 1713-179.
- 1382 Wotschokowsky, M.: (Analysis of pesticides in ground and drinking water). *GIT Fachz. Lab.*, 37 (1993) 404-408; *C.A.*, 119 (1993) 124686y.

For additional information see *C.A.*:

119 (1993) 107990a.

See also 329, 334, 450, 452, 1086, 1087, 1088, 1089, 1091, 1092, 1099, 1100, 1101, 1104, 1365, 1374, 1386, 1421, 1422, 1427, 1429.

35d. *Soil pollution (complex mixtures; single compounds by cross-reference only)*

- 1383 Gere, D.R., Knipe, C.R., Castelli, P., Hedrick, J., Randall Frank, L.G., Schulenberg-Schell, H., Schuster, R., Doherty, L., Orolin, J. and Lee, H.B.: Bridging the automation gap between sample preparation and analysis: an overview of SFE, GC, GC-MS, and HPLC applied to environmental samples. *J. Chromatogr. Sci.*, 31 (1993) 246-258.

See also 181, 212, 214, 229, 1110.

## 36. SOME TECHNICAL PRODUCTS AND COMPLEX MIXTURES

## 36a. Surfactants

- 1384 Marcomini, A., di Corcia, A., Samperi, R. and Capri, S.: Reversed-phase high-performance liquid chromatographic determination of linear alkylbenzene sulphonates, nonylphenol polyethoxylates and their carboxylic biotransformation products. *J. Chromatogr.*, 644 (1993) 59-71.
- 1385 Schröder, H.F.: Surfactants: non-biodegradable, significant pollutants in sewage treatment plant effluents. Separation, identification and quantification by liquid chromatography, flow-injection analysis-mass spectrometry and tandem mass spectrometry. *J. Chromatogr.*, 647 (1993) 219-234.
- 1386 Yokoyama, Y., Kondo, M. and Sato, H.: Determination of alkylbenzenesulphonates in environmental water by anion-exchange chromatography. *J. Chromatogr.*, 643 (1993) 169-172.

## 36b. Antioxidants and preservatives

- 1387 Barman, B.N.: LC-amperometric determination of *tert*-butylcatechol in *N*-methyl-2-pyrrolidone stream containing polar and non-polar impurities. *J. High Resolut. Chromatogr.*, 16 (1993) 404-406.
- 1388 Buergi, C. and Otz, T.: (Determination of preservatives in body shampoos and bubble baths). *Mitt. Geb. Lebensmittelunters Hyg.*, 83 (1992) 492-508; *C.A.*, 119 (1993) 102961g.
- 1389 Nielson, R.C.: Recent advances in polyolefin additive analysis. *J. Liq. Chromatogr.*, 16 (1993) 1625-1638.
- 1390 Page, B.D.: Liquid chromatographic method for the determination of nine phenolic antioxidants in butter oil: collaborative study. *J. Assoc. Off. Anal. Chem.*, 76 (1993) 765-779.
- 1391 Shamsi, S.A. and Danielson, N.D.: Ion chromatography of polyphosphates and polycarboxylates using a naphthalenetrisulfonate eluent with indirect photometric and conductivity detection. *J. Chromatogr.*, 653 (1993) 153-160.
- 1392 Thompson, M.J., Fell, A.F., Clark, B.J. and Robinson, M.L.: LC studies on the potential interaction of paraben preservatives with sorbitol and glycerol. *J. Pharm. Biomed. Anal.*, 11 (1993) 233-240.

See also 408, 1369.

36c. *Complex mixtures, technical products and unidentified compounds*

- 1393 Dahms, W., Schumacher, R. and Mayer, H.: (Analytics as a key to the safe process). *Metallberflaeche*, 47 (1993) 331-338; C.A., 119 (1993) 190617c - a review with 7 refs.
- 1394 Guthertz, L.S., Lim, S.D., Jang, Y. and Duffey, P.S.: Curvilinear-gradient high-performance liquid chromatography for identification of mycobacteria. *J. Clin. Microbiol.*, 31 (1993) 1876-1881; C.A., 119 (1993) 134699u.
- 1395 Schlüter, H., Kluth, B., Börjesson-Stoll, R., Nordhoff, E. and Zidek, W.: Partial purification and characterization of a circulating hypertensive factor in spontaneously hypertensive rats. *Eur. J. Biochem.*, 218 (1993) 67-73.
- 1396 Timpa, J.D.: Size-exclusion chromatography to characterize cotton fiber. *ACS Symp. Ser.*, 521 (1993) 302-313; C.A., 119 (1993) 182708t.

For additional information see C.A.:  
119 (1993) 114846d.

See also 7, 20, 149, 224, 241, 625, 1409, 1417.

37. CELLS, CELLULAR PARTICLES AND SUPRAMOLECULAR STRUCTURES

- 1397 Moon, M.H. and Giddings, J.C.: Size distribution of liposomes by flow field-flow fractionation. *J. Pharm. Biomed. Anal.*, 11 (1993) 911-920.

See also 498.

38. INORGANIC COMPOUNDS

38a. *Cations*

- 1398 Abe, H., Usuda, S., Takeishi, H. and Tachimori, S.: Separation of light lanthanoids by centrifugal partition chromatography in 30% TBP extraction system. *J. Liq. Chromatogr.*, 16 (1993) 2661-2672.
- 1399 Chou, C.L., Uthe, J.F. and Guy, R.D.: Determination of free and bound Cd, Zn, Cu, and Ag ions in lobster (*Homarus americanus*) digestive gland extracts by gel chromatography followed by atomic absorption spectrophotometry and polarography. *J. Assoc. Off. Anal. Chem.*, 76 (1993) 794-798.
- 1400 Frosini, M., Goreli, B., Matteini, M., Palmi, M., Valoti, M. and Sgaragli, G.P.: HPLC determination of inorganic cation levels in CSF and plasma of conscious rabbits. *J. Pharmacol. Toxicol. Methods*, 29 (1993) 99-104; C.A., 119 (1993) 155172g.
- 1401 Hall, K.E. and McCord, B.R.: The analysis of mono- and divalent cations present in explosive residues using ion chromatography with conductivity detection. *J. Forensic Sci.*, 38 (1993) 928-934; C.A., 119 (1993) 133023g.
- 1402 Hoshino, H., Kaneko, E. and Yotsuyanagi, T.: (A coupled scheme of HPLC and spectrophotometry as an approach to the highly sensitive and selective monitoring of trace elements). *Biomed. Res. Trace Elem.*, 3 (1992) 95-96; C.A., 119 (1993) 134689r.

- 1403 Jen, J.-F., Ou-Yang, G.-L., Chen, C.-S. and Yang, S.-M.: Simultaneous determination of chromium(III) and chromium(VI) with reversed-phase ion-pair high-performance liquid chromatography. *Analyst (London)*, 118 (1993) 1281-1284.
- 1404 Kitazume, E., Sato, N., Saito, Y. and Ito, Y.: Separation of heavy metals by high-speed countercurrent chromatography. *Anal. Chem.*, 65 (1993) 2225-2228.
- 1405 LeGras, C.A.A.: Simultaneous determination of anions and divalent cations using ion chromatography with ethylenediaminetetraacetic acid as eluent. *Analyst (London)*, 118 (1993) 1035-1041.
- 1406 Liu, Y. and Wang, Q.: Rapid and continuous ion chromatographic determination of trace heavy metal impurities in noble metals. *J. Chromatogr.*, 644 (1993) 73-82.
- 1407 Nam, N.P.T., Gleisberg, B. and Gorski, B.: Behavior of neptunium in the separation process of actinides from irradiated targets. *J. Radioanal. Nucl. Chem.*, 170 (1993) 359-366; C.A., 119 (1993) 130506f.
- 1408 Oguma, K., Sato, K. and Kuroda, R.: Microchemical determination of nine earth elements in silicate rocks by cation-exchange preconcentration-ion-interaction chromatography. *Chromatographia*, 37 (1993) 319-324.
- 1409 Okamoto, Y., Hahashi, K., Kawahara, Y. and Kumamaru, T.: Dual catio-selective membrane tube electro dialysis/ion chromatography for halide ions in strongly basic solutions and its application to the analysis of rock samples. *Bull. Sch. Soc. Jpn.*, 66 (1993) 1664-1668; C.A., 119 (1993) 130575c.
- 1410 Petrovic, M., Kaštelan-Macan, M., Turina, S. and Ivankovic, V.: Quantitative determination of Mg in Al-alloys by ion-exchange TLC. *J. Liq. Chromatogr.*, 16 (1993) 2673-2684.
- 1411 Riaz, M., Butt, S.B. and Ul-Haq, E.: Assay of uranium in the presence of various interfering ions using ion pair liquid chromatography. *J. Liq. Chromatogr.*, 16 (1993) 1589-1599.
- 1412 Saraswati, R. and Rao, T.H.: Ion-chromatographic separation and determination of some metal ions in stainless steels by pre-column chelation with 4-(2-thiazolylazo)resorcinol. *J. Liq. Chromatogr.*, 16 (1993) 1601-1614.
- 1413 Siebrands, T., Giersig, M., Mulvaney, P. and Fischer, G.H.: Steric exclusion chromatography of nanometer-sized gold particles. *Langmuir*, 9 (1993) 2297-2300; C.A., 119 (1993) 147272z.
- 1414 Subramanian, D.V.: Analysis of aluminum in amino acids by high performance liquid chromatography. *Can. Pat. Appl. CA 2,082, 899 (Cl. G01N30/02)*, 15 May 1993, US Appl. 792,124, 14 Nov. 1991; 13 p.; C.A., 119 (1993) 151416s.
- 1415 Walker, T.A.: The separation and indirect visible detection of rare earth metal cations on a dye-coated stationary phase. *J. Liq. Chromatogr.*, 16 (1993) 1573-1587.
- 1416 Watkins, R.T. and Le Roex, A.P.: The rare-earth element content of GSJ rock reference samples determined by gradient ion chromatography. *Geochem. J.*, 26 (1992) 241-249; C.A., 119 (1993) 194575k.
- 1417 Watkins, R.T. and Le Roex, A.P.: Rare earth element concentrations in USGS and CCRMP igneous rock standards determined by high performance ion chromatography. *Geostand. Newsl.*, 17 (1993) 105-111; C.A., 119 (1993) 130535q.
- 1418 Yi, Z., Zhuang, G. and Brown, P.R.: Analysis of metal ions by high performance liquid chromatography. *J. Liq. Chromatogr.*, 16 (1993) 3133-3170 - a review with 196 refs.

- 1419 Zhao, Y.: Composition of 2-(3,5-dibromo-2-pyridylazo)diethylaminophenol chelates by liquid chromatography. *Chromatographia*, 37 (1993) 284-286.

For additional information see C.A.:

119 (1993) 108061k, 112577a, 173142b, 194596t.

See also 71, 156, 653, 984, 1365, 1430.

### 38b. Anions

- 1420 Barnett, N.W., Jones, P. and Handley, H.W.: The use of zirconyl xylenol orange for the post-column spectroscopic detection of fluoride in ion chromatography. *Anal. Lett.*, 26 (1993) 2525-2529.
- 1421 Bignami, S., Davi, M.L., Milan, C., Moretti, M. and Navarra, F.: Evaluation of ion-exchange chromatography for nitrate determination in wastewaters. *J. Chromatogr.*, 644 (1993) 341-344.
- 1422 Davi, M.L., Bignami, S., Milan, C., Liboni, M. and Malfatto, M.G.: Determination of nitrate in surface waters by ion-exchange chromatography after oxidation of total organic nitrogen to nitrate. *J. Chromatogr.*, 644 (1993) 345-348.
- 1423 Debowski, J.K. and Gerber, F.A.: Determination of isocyanuric acid and its chlorinated derivatives in swimming pool waters by ion chromatography. *J. Chromatogr.*, 644 (1993) 400-403.
- 1424 Fung, Y.M.: (Determination of sulfites in food by HPLC with electrochemical detection). *Lebensm.-Biotechnol.*, 10 (1993) 88-89; C.A., 119 (1993) 115601q - a review with 14 refs.
- 1425 Gennaro, M.C., Abrigo, C., Marengo, E. and Liberatori, A.: A selective determination of azide by ion-interaction reversed-phase HPLC. *J. Liq. Chromatogr.*, 16 (1993) 2715-2730.
- 1426 Gorbunov, N. and Esposito, E.: Rapid and simple method for the determination of nitrite in synaptosomal superfusates by liquid chromatography with electrochemical detection. *J. Chromatogr.*, 619 (1993) 133-136.
- 1427 Kailu, L., Jin, L. and Siming, Y.: Ion chromatographic analysis of anions based on solid-phase extraction. *J. Liq. Chromatogr.*, 16 (1993) 3083-3092.
- 1428 Kura, G.: Anion-exchange selectivity of cyclic phosphate oligomers. *J. Chromatogr.*, 644 (1993) 198-201.
- 1429 Martin, M.W. and Giacomini, R.A.: Ultratrace anion analysis of high-purity water. A column comparison. *J. Chromatogr.*, 644 (1993) 333-340.
- 1430 Nguyen, V.D. and Rossbach, M.: Ion chromatographic investigation of brown algae (*Fucus vesiculosus*) of the German Environmental Specimen Bank. *J. Chromatogr.*, 643 (1993) 427-433.
- 1431 Saboori, A.M., Rose, N.R., Butscher, W.G. and Burek, C.L.: Modification of a nonincinerative method for determination of iodine in iodoproteins. *Anal. Biochem.*, 214 (1993) 335-338.

- 1432 Santillana, M.I., Ruiz, E., Nieto, M.T. and de Alba, M.: High performance ion chromatography determination of nitrite and nitrate in foodstuffs. *J. Liq. Chromatogr.*, 16 (1993) 1561-1571.
- 1433 Sertl, D. and Malone, W.: Liquid chromatographic method for determination of iodine in milk: collaborative study. *J. Assoc. Off. Anal. Chem.*, 76 (1993) 711-719.
- 1434 Stahl, R.: Ion chromatographic determination of chloride, chlorate, and perchlorate in sulfuric acid solutions. *Chromatographia*, 37 (1993) 300-302.
- 1435 Takeuchi, T., Miwa, T., Hemmi, A. and Maeda, T.: Microcolumns ion chromatography with conductometric detection. *Chromatographia*, 37 (1993) 173-175.

For additional information see C.A.:

119 (1993) 108066r, 108092w, 112576z, 151282v, 194714e.

See also 30, 43, 115, 962, 1391, 1405.

### 38d. Volatile inorganic compounds

- 1436 Hu, D., Feng, Y.P.: (HPLC detection of hydroxyl radicals in striatum extracellular fluid in rats subjected to reperfusion after cerebral ischemia and the action of vitamin E). *Yaoxue Xuebao*, 28 (1993) 337-341; C.A., 119 (1993) 155155d.
- 1437 Liu, Y. and Zhang, J.T.: (Direct measurement of oxy-radicals using HPLC-ECD). *Yaoxue Xuebao*, 28 (1993) 416-421; C.A., 119 (1993) 134697s.
- 1438 Rando, R.J., Poovey, H.G. and Chang, S.-N.: Collection and chemical derivatization of airborne phosgene with 1-(2-pyridyl)piperazine and determination by high performance liquid chromatography. *J. Liq. Chromatogr.*, 16 (1993) 3291-3209.
- 1439 Wu, W.S. and Gaid, V.: Determination of phosgene (carbonylchlorid) in air by high-performance liquid chromatography with a dual selective detection system. *Analyst (London)*, 118 (1993) 1285-1288.

For additional information see C.A.:

119 (1993) 110659e.

See also 245.

## 39. RADIOACTIVE AND OTHER ISOTOPE COMPOUNDS

See 80, 536.

## Gas Chromatography

### 1. REVIEWS AND BOOKS

- 1 Buffham, B.A., Mason, G. and Meacham, R.I.: Absolute gas chromatography. *Proc. R. Soc. London, Ser. A*, 440 (1993) 291-301; *C.A.*, 119 (1993) 15739x.
- 2 De Zeeuw, R.A., Franke, J.P., Machata, G., Moller, M.R., Muller, R.K., Graefe, A., Tiess, D., Pflieger, K. and von Mallinckrodt, M.G.: *Gas Chromatographic Retention Indices of Solvents and Other Volatile Substances for Use in Toxicological Analysis*. VCH Verlagsgesellschaft, Weinheim, 1992, 130 pp.
- 3 De Zeeuw, R.A., Franke, J.P., Maurer, H.H. and Pflieger, K.: *Gas Chromatographic Retention Indices of Toxicologically-Relevant Substances on Packed or Capillary Columns with Dimethylsilicone Stationary Phases*. VCH Verlagsgesellschaft, Weinheim, 1992, 407 pp.
- 4 Westwood, S.A.: *Supercritical Fluid Extraction and Its Use in Chromatographic Sample Preparation*. Blackie Academic and Professional, London, 1993, xii + 170 pp.
- 5 Yashin, Ya.I.: (Novel advances and trends of gas chromatography.) *Zh. Fiz. Khim.*, 67 (1993) 804-809 - a review with 22 refs.

See also 13, 62, 63, 66, 67, 69, 70, 71, 77, 79, 82, 84, 106, 119, 134, 140, 172, 175, 238, 240, 263, 267, 383, 472, 551.

### 2. FUNDAMENTALS, THEORY AND GENERAL

#### 2a. General

- 6 Davis, J.M.: Statistical theory of spot overlap for *n*-dimensional separations. *Anal. Chem.*, 65 (1993) 2014-2023.
- 7 Ettre, L.S.: The new IUPAC nomenclature for chromatography. *LC-GC Int.*, 6 (1993) 544-548.
- 8 Sun, Y., Huang, A., Zhang, R. and He, L.: Practical aspects in the utilization of the Sadtler Standard Gas Chromatography Retention Index Library. *J. Chromatogr.*, 648 (1993) 395-405.

See also 129.

#### 2b. Thermodynamics and theoretical relationships

- 9 Blumberg, L.M. and Berger, T.A.: Molecular basis of peak width in capillary gas chromatography under high column pressure drop. *Anal. Chem.*, 65 (1993) 2686-2689.
- 10 De Frutos, M., Sanz, J., Martínez-Castro, I. and Jiménez, M.I.: New parameters for the characterization of relationship between gas chromatographic retention and temperature. *Anal. Chem.*, 65 (1993) 2643-2649.
- 11 Larin, A.B.: (Dependence of peak profile on the column length in gas chromatography.) *Zh. Fiz. Khim.*, 67 (1993) 810-814.

- 12 Poole, C.F. and Kollie, T.O.: Interpretation of the influence of temperature on the solution properties of gas chromatographic phases using Abraham's solvation parameter model. *Anal. Chim. Acta*, 282 (1993) 1-17.
- 13 Roth, M.: (Thermodynamical measurements by column chromatography.) *Chem. Listy*, 87 (1993) 671-678 - a review with 139 refs.
- 14 Toft, J. and Kvalheim, O.M.: Resolution parameter for multidection chromatography. *Anal. Chem.*, 65 (1993) 2270-2275.
- 15 Vigdergauz, M.S. and Lange, P.K.: (Chromatography as a spectral method of investigation.) *Zh. Fiz. Khim.*, 67 (1993) 815-816.
- 16 Vigdergauz, M.S., Platonov, I.A., Lobachev, A.L. and Sushko, O.B.: (Retention time of non-sorbing substance in chromatography.) *Zh. Fiz. Khim.*, 67 (1993) 857-860.

#### 2c. Relationship between structure and chromatographic behaviour

- 17 Berezkin, V.G. and Korolev, A.A.: (Effect of nature of carrier gas on retention values of compounds in gas chromatography.) *Zavod. Lab.*, 59, No. 6 (1993) 15-16.
- 18 Grigor'eva, D.N., Vasil'ev, A.V. and Golovnya, R.V.: (Effect of temperature on the retention indices of homologous bifunctional compounds in capillary GC.) *Zh. Anal. Khim.*, 48 (1993) 1167-1174.
- 19 Jiang, X. and Zeng, X.: (The relationships between the structures and the retention indices of the homologues of alkyl pyridine and quinoline.) *Sepu*, 11 (1993) 154-156.
- 20 Li, L., Wu, C.-Y., Cai, L.-S. and Zeng, Z.-R.: Studies on the retention and thermodynamic properties of aromatic compounds on two types of crown ether polysiloxane stationary phase. *J. Chromatogr. A*, 653 (1993) 354-358.
- 21 Rohrschneider, L.: Prediction of retention data from stationary phase constants in gas chromatography. *Chromatographia*, 37 (1993) 250-258.
- 22 Zhang, W., Wu, Y. and Shi, J.: (A further study on the intersecting point of the curves of the logarithm of specific retention volume versus the reciprocal of temperature for homologues in gas-liquid chromatography.) *Sepu*, 11 (1993) 219-221.

See also 10, 27, 109, 112, 122, 180, 182, 184, 191, 198, 209, 219, 225, 351, 357, 363, 370, 436, 482, 539.

#### 2d. Measurement of physico-chemical and related values

- 23 Andronikashvili, T.G., Eprikashvili, L.G., Abulashvili, E.I., Leonidze, N.M. and Kordzakhia, T.N.: The investigation of chromatographic properties of Chromosorb-103 in the variant of surface-layer sorbent. *Izv. Akad. Nauk Gruz., Ser. Khim.*, 17 (1991) 219-223; *C.A.*, 118 (1993) 224597t.
- 24 Bhagat, S.D.: Inverse gas-chromatographic method for the estimation of the vapour pressure of less volatile compounds. *Frensenius J. Anal. Chem.*, 347 (1993) 365-367.

- 25 Hu, Q., Huang, D. and Wang, S.: A new method for predicting liquid-liquid equilibria from headspace analysis determinations over homogeneous liquid mixtures using the UNIQUAC model. *Fluid Phase Equilib.*, 84 (1993) 267-280; C.A., 118 (1993) 261718d.
- 26 Lee, H.L. and Luner, P.: Analysis of the adsorption of alkanes on high surface area cellulose by inverse gas chromatography. *J. Wood Chem. Technol.*, 13 (1993) 127-144; C.A., 118 (1993) 256810y.
- 27 Li, J., Zhang, Y. and Carr, P.W.: Development of a gas chromatographic scale of solute hydrogen bond acceptor basicity and characterization of some hydrogen bond donor phases by use of linear solvation energy relationships. *Anal. Chem.*, 65 (1993) 1969-1979.
- 28 Perlinger, J.A., Eisenreich, S.J. and Capel, P.D.: Application of headspace analysis to the study of sorption of hydrophobic organic chemicals to  $\alpha$ -alumina. *Environ. Sci. Technol.*, 27 (1993) 928-937.
- 29 Riedl, B. and Kamdem, P.D.: Estimation of the dispersive component of surface energy of polymer-grafted lignocellulosic fibers with inverse gas chromatography. *J. Adhes. Sci. Technol.*, 6 (1992) 1053-1067; C.A., 118 (1993) 215117j.
- 30 Riganakos, K.A., Demertzis, P.G. and Kontominas, M.G.: Effect of crystalline sucrose on the water sorption behavior of wheat flour as studied by inverse gas chromatography. *Food Sci. Technol. (London)*, 25 (1992) 389-394; C.A., 118 (1993) 211650e.
- 31 Robbins, G.A., Wang, S. and Stuart, J.D.: Using static headspace method to determine Henry's law constants. *Anal. Chem.*, 65 (1993) 3113-3118.
- 32 Voelkel, A.: Inverse gas chromatography. Solubility parameters of 1,3-bis- $[\omega$ -hexoxy-oligo(oxyethylene)]-propan-2-ols. *Acta Chromatogr. (Katowice)*, No. 1 (1992) 69-80.
- 33 Vorob'eva, L.M., Larionov, O.G. and Tchaliikh, A.E.: (Gas chromatographic study of adsorptive properties of  $\gamma$ -Fe<sub>2</sub>O<sub>3</sub> purified by dialysis.) *Zh. Fiz. Khim.*, 66 (1992) 3045-3051.
- 34 Wang, J.Y. and Charlet, G.: Thermodynamic study of ethyl cellulose by gas chromatography. *Macromolecules*, 26 (1993) 2413-2419; C.A., 118 (1993) 215133m.
- 35 Zheng, H., Du, S., Ma, F., Wang, J. and Lu, G.: (Study of the reduction mechanism of lanthanum nickel oxide (LaNiO<sub>3</sub>) by TPR-GC and XRD techniques.) *Wuji Huaxue Xuebao*, 7 (1991) 474-476; C.A., 118 (1993) 199211c.
- 38 Frolov, F.Ya. and Brazhnikov, V.V.: (Device for sampling and introducing vapor-phase samples into a gas chromatograph.) *U.S.S.R. SU 1,723,516* (Cl. G01N30/10), 3 Mar. 1992, Appl. 4,776,518, 30 Mar. 1990; C.A., 118 (1993) 204476z.
- 39 Grob, K. and Wagner, C.: Inserts providing complete sample evaporation above the column entrance in vaporizing GC injector. *J. High Resolut. Chromatogr.*, 16 (1993) 429-432.
- 40 Grob, K. and Wagner, C.: Procedure for testing inertness of inserts and insert packing material for GC injectors. *J. High Resolut. Chromatogr.*, 16 (1993) 464-468.
- 41 Gröngerg, L., Lövkvist, P. and Jönsson, J.Å.: An interface between a liquid flow system and a packed column gas chromatograph. *J. Chromatogr. Sci.*, 31 (1993) 384-386.
- 42 Hagman, G. and Roeraade, J.: Independently temperature programmed uncoated precolumn: a concept for improved sample introduction in capillary GC. *J. Microcolumn Sep.*, 5 (1993) 341-346.
- 43 Hagman, G. and Roeraade, J.: On-line liquid backflush of an uncoated precolumn for automated gas chromatographic analysis of complex mixtures. *J. Chromatogr. A*, 654 (1993) 287-298.
- 44 Hagman, G. and Roeraade, J.: On-line liquid backflush – a new technique for GC analysis of samples containing nonvolatile material. *J. High Resolut. Chromatogr.*, 16 (1993) 445-449.
- 45 Harter, D.A., Alcorn, W.B., Busche, S., Ellington, F.O., Kreuz, J.P., Brabston, M.A. and Ring, R.D.: On-line gas chromatograph for industrial process control. *Process Control Qual.*, 3 (1992) 145-151; C.A., 118 (1993) 194475g.
- 46 Ibáñez, C.: Construction of an all-glass/fused silica, zero dead volume multidimensional gas chromatographic system using press-fit connectors: preliminary results. *J. High Resolut. Chromatogr.*, 16 (1993) 552-554.
- 47 Klemp, M.A., Akard, M.L. and Sacks, R.D.: Cryofocusing inlet with reverse flow sample collection for gas chromatography. *Anal. Chem.*, 65 (1993) 2516-2521.
- 48 Lou, X., Janssen, H.-G. and Cramers, C.A.: Quantitative aspects of directly coupled supercritical fluid extraction–capillary gas chromatography with a conventional split/splitless injector as interface. *J. High Resolut. Chromatogr.*, 16 (1993) 425-428.
- 49 Louter, A.J.H., Brinkman, U.A.T. and Ghijzen, R.T.: Fully automated water analyzer based on on-line solid phase extraction–gas chromatography. *J. Microcolumn Sep.*, 5 (1993) 303-315.
- 50 Mol, H.G.J., Janssen, H.-G.M., Cramers, C.A. and Brinkman, U.A.T.: Use of a temperature programmed injector with a packed liner for direct water analysis and on-line reversed phase LC-GC. *J. High Resolut. Chromatogr.*, 16 (1993) 459-463.
- 51 Muto, H.: (Control apparatus for carrier-gas-flow rate in gas chromatographs.) *Jpn. Kokai Tokkyo Koho* JP 04,177,162 [92,177,162] (Cl. G01N30/32), 24 Jun. 1992, Appl. 90/304,046, 13 Nov. 1990; 7 pp.; C.A., 118 (1993) 224640b.
- 52 Nagayanagi, Y.: (Gas chromatograph.) *Jpn. Kokai Tokkyo Koho* JP 04,303,767 [92,303,767] (Cl. G01N30/54), 27 Oct. 1992, Appl. 91/92,857, 30 Mar. 1991; 3 pp.; C.A., 118 (1993) 246620x.
- 53 Onishi, A., Oguri, N. and Poongzag, K.: Development of a new injection system for Curie-point pyrolysis–gas chromatography. *J. Chromatogr. Sci.*, 31 (1993) 380-383.
- 54 Ridgeon, P.J. and Lillack, H.: Universelles Probenaufgabesystem für Kapillarsäulen. *LaborPraxis*, 17, No. 12 (1993) 22-27.

See also 420, 423, 528, 529.

### 3. GENERAL TECHNIQUES

#### 3a. Apparatus and accessories

- 36 Akard, M. and Sacks, R.: Column bifurcation with tunable selectivity for high-speed gas chromatography. *J. Chromatogr. Sci.*, 31 (1993) 297-304.
- 37 Blanchard, P., Shepson, P.B., Schiff, H.I. and Drummond, J.W.: Development of a gas chromatograph for trace level measurement of peroxyacetyl nitrate using chemical amplification. *Anal. Chem.*, 65 (1993) 2472-2477.

- 55 Señoráns, F.J., Tabera, J., Villén, J., Herraiz, M. and Reglero, G.: Variables affecting the introduction of large sample volumes in capillary gas chromatography using a programmed-temperature vaporizer. *J. Chromatogr.*, 648 (1993) 407-414.
- 56 Tajima, H., Shimamura, N. and Aoki, M.: (Development of large volume injection method for capillary GC.) *Bunseki Kagaku*, 42 (1993) 573-579.
- 57 Vreuls, J.J., de Jong, G.J., Ghijsen, R.T. and Brinkman, U.A.T.: On-line solid-phase extraction/thermal desorption for introduction of large volumes of aqueous samples into a capillary gas chromatograph: Part 2. *J. Microcolumn Sep.*, 5 (1993) 317-324.
- 58 Vtorov, B.G., Kalmanovskii, V.I., Rybalchenko, Yu.P., Savinov, I.M., Savinov, Yu.S. and Yashin, Ya.I.: (Gas chromatographic instrument of Dzerzhinsk construction bureau.) *Zh. Fiz. Khim.*, 67 (1993) 861-863.
- 59 Wu, Z. and Yu, W.: (Solvent venting procedure and its application to capillary gas chromatography.) *Sepeu*, 11 (1993) 91-93.
- See also 74, 83, 130, 133, 135, 164, 175, 503, 505, 517.
- 3b. *Detectors and detection reagents*
- 60 Adamiya, T.V., Bohdarenko, I.V., Budovich, V.L. and Shlyakhov, A.F.: (Photoionization detector with controlled selectivity for capillary gas chromatography.) *Zavod. Lab.*, 59, No. 5 (1993) 17-20.
- 61 Bernier, U.R. and Yost, R.A.: Vacuum operation of the flame-ionization detector for gas chromatography. *J. Chromatogr. Sci.*, 31 (1993) 358-362.
- 62 Bornhop, D.J. and Richter, B.E.: Detectors for capillary supercritical fluid chromatography. *Chem. Anal. (N. Y.)*, 121 (1992) 355-393; *C.A.*, 118 (1993) 265751g - a review with 156 refs.
- 63 Clement, R.E. and Reiner, E.J.: Mass-spectrometric detectors [for capillary chromatography]. *Chem. Anal. (N. Y.)*, 121 (1992) 327-353; *C.A.*, 118 (1993) 204399b - a review with 30 refs.
- 64 Conte, E.D. and Barry, E.F.: The performance of a dual alkali aerosol flame ionization detector for gas chromatography. *Anal. Lett.*, 26 (1993) 2711-2727.
- 65 D'Couto, S.: Electrolytic conductivity detector. *U.S. US* 5,194,814 (Cl. 324-446; G01N27/02), 16 Mar. 1993, Appl. 703,881, 22 May 1991; 11 pp.; *C.A.*, 118 (1993) 265835n.
- 66 Driscoll, J.N.: Far-UV ionization (photoionization) and absorbance detectors [for capillary chromatography]. *Chem. Anal. (N. Y.)*, 121 (1992) 51-81; *C.A.*, 118 (1993) 204393v - a review with 37 refs.
- 67 Fujii, T. and Arimoto, H.: The surface ionization detector [for capillary chromatography]. *Chem. Anal. (N. Y.)*, 121 (1992) 169-191; *C.A.*, 118 (1993) 204396y - a review with 24 refs.
- 68 Gelencsér, A., Szépvölgyi, J. and Hlavay, J.: Characterization of an element-specific detector for combined gas chromatography-atomic emission detection. *J. Chromatogr. A*, 654 (1993) 269-277.
- 69 Grimsrud, E.P.: The electron capture detector [for capillary gas chromatography]. *Chem. Anal. (N. Y.)*, 121 (1992) 83-107; *C.A.*, 118 (1993) 265750f - a review with 45 refs.
- 70 Hall, R.C.: The electrolytic conductivity (Hall) detector [for capillary chromatography]. *Chem. Anal. (N. Y.)*, 121 (1992) 109-137; *C.A.*, 118 (1993) 204394w.
- 71 Hutte, R.S. and Ray, J.D.: Sulfur-selective detectors [for capillary chromatography]. *Chem. Anal. (N. Y.)*, 121 (1992) 193-218; *C.A.*, 118 (1993) 204397z - a review with 84 refs.
- 72 Kageura, M., Fujiwara, Y., Hara, K., Hieda, Y. and Kashimura, S.: (A comparison of NPD with SID as a gas chromatography detector for drug analysis.) *Hochudoku*, 10 (1992) 144-145; *C.A.*, 118 (1993) 182639a.
- 73 Karpas, Z., Wang, Y.-F. and Eiceman, G.A.: Qualitative and quantitative response characteristics of a capillary gas chromatography/ion mobility spectrometer to halogenated compounds. *Anal. Chim. Acta*, 282 (1993) 19-31.
- 74 Khononzon, G. and Rutenberg, O.: Microsensors and actuators based on single-crystal silicon in miniature gas chromatographs and analyzers. *J. Micromech. Microeng.*, 2 (1992) 266-268; *C.A.*, 118 (1993) 265538t.
- 75 Kientz, C.E. and Brinkman, U.A.T.: The use of gas chromatographic detectors in column liquid chromatography. *TrAC*, 12 (1993) 363-373 - a review with 52 refs.
- 76 Lasa, J., Sliwka, I., Korus, A., Berski, M. and Cyganik, S.: (Photoionization detector for gas chromatograph.) *Pol. PL* 154,924 (Cl. G01N30/70), 31 Dec. 1991, Appl. 266,789, 10 Jul. 1987; 3 pp.; *C.A.*, 119 (1993) 19599z.
- 77 McMinin, D.G. and Hill, H.H.: The flame ionization detector [for capillary chromatography]. *Chem. Anal. (N. Y.)*, 121 (1992) 7-21; *C.A.*, 118 (1993) 204392u - a review with 41 refs.
- 78 Osato, I.: (Electron-capture detector used for chromatograph.) *Jpn. Kokai Tokkyo Koho* JP 04,303,759 [92,303,759] (Cl. G01N27/64), 27 Oct. 1992, Appl. 91/67,423, 30 Mar. 1991; 8 pp.; *C.A.*, 118 (1993) 246617b.
- 79 Patterson, P.L.: The nitrogen-phosphorus detector [for capillary chromatography]. *Chem. Anal. (N. Y.)*, 121 (1992) 139-168; *C.A.*, 118 (1993) 204395x - a review with 49 refs.
- 80 Pedersen-Bjergaard, S. and Greibrokk, T.: On-column bromine- and chlorine-selective detection for capillary gas chromatography using a radio frequency plasma. *Anal. Chem.*, 65 (1993) 1998-2002.
- 81 Purcell, J.E. and Dang, R.: Method of manufacturing glass beads for use in thermionic gas chromatographic detectors. *U.S. US* 5,198,009 (Cl. 65-59.22; C03C27/02), 3 Mar. 1993, Appl. 796,497, 22 Nov. 1991; 4 pp.; *C.A.*, 118 (1993) 265836p.
- 82 Ramsey, R.S. and Andrawes, F.F.: The modern helium ionization detector [for capillary chromatography]. *Chem. Anal. (N. Y.)*, 121 (1992) 23-49; *C.A.*, 118 (1993) 224398d - a review with 27 refs.
- 83 Staniewski, J., Janssen, H.-G.M., Rijks, J.A. and Cramers, C.A.: Introduction of large volumes of methylene chloride in capillary GC with electron capture detection. *J. Microcolumn Sep.*, 5 (1993) 429-432.
- 84 Uden, P.C.: Atomic plasma element-specific detection [for capillary chromatography]. *Chem. Anal. (N. Y.)*, 121 (1992) 219-250; *C.A.*, 118 (1993) 224579p - a review with 57 refs.
- 85 Yu, A., Wang, X. and Jin, Q.: (Studies on the response characteristics of microwave-induced plasma ionization detector.) *Gaodeng Xuexiao Huaxue Xuebao*, 13 (1992) 1533-1535; *C.A.*, 118 (1993) 224595r.
- See also 143, 167, 281, 365, 391, 462, 526.



## 3c. Sorbents and columns, packing procedures

- 86 Andronikashvili, T.G., Dzagania, M.A. and Eprikashvili, L.G.: (The effect of solvent nature on the character of coating and separating properties of liquid crystals in gas-liquid chromatography). *Soobshch. Akad. Nauk Gruz.*, 142 (1991) 321-324; C.A., 118 (1993) 220491f.
- 87 Berezkin, V.G., Shiryayeva, V.E. and Popova, T.P.: (About a possible realization of binary sorbents with immobilized stationary phases in gas chromatography.) *Zh. Fiz. Khim.*, 67 (1993) 841-842.
- 88 Betts, T.J.: Potential of three different  $\alpha$ -cyclodextrin modifications for the gas chromatographic evaluation of constituents of volatile oils. *J. Chromatogr. A*, 653 (1993) 167-172.
- 89 Bulanova, A.V., Vigdergauz, M.S. and Bashkinova, T.B.: (Development of chromatographic method at different pressures: Use of lower pressures.) *Zh. Fiz. Khim.*, 67 (1993) 854-856.
- 90 Correa, S.M.A., Barbosa, S.D.C.T., de Aquino, F.R. and Coelho, R.B.: (High-resolution capillary gas chromatography. 10. Reactivation and recovery of high-resolution capillary column packings used for analysis of wood tar components.) *Quim. Nova*, 15 (1992) 291-294; C.A., 118 (1993) 236127c.
- 91 Decker, D.R. and Spilkin, A.: Utilizing hydrogen carrier gas in volatiles analysis for maximum resolution in the shortest analysis time. *J. Chromatogr. Sci.*, 31 (1993) 305-309.
- 92 Dolgov, B.N. and Sidel'nikov, V.N.: (Evaluation of parameters of cryofocussing capillary for analysis of trapped concentrates.) *Zh. Fiz. Khim.*, 67 (1993) 817-820.
- 93 Fu, R., Ge, J., Deng, J., Zhu, J., Gu, J., Huang, Z. and Zhang, B.: (Behavior of capillary columns coated with poly(crown ether)siloxane.) *Fenxi Huaxue*, 21 (1993) 379-383; C.A., 119 (1993) 19557j.
- 94 Fu, R., Huang, C., Huang, Z. and Xu, W.: Preparation of benzo-18-crown-6 ether side-chain polysiloxane used as open tubular column gas chromatographic stationary phase. *J. Chromatogr. A*, 653 (1993) 173-177.
- 95 Fu, R., Jing, P., Gu, J., Huang, Z. and Chen, Y.: Side-chain liquid crystalline polysiloxane containing crown ether used as stationary phase for capillary gas chromatography. *Anal. Chem.*, 65 (1993) 2141-2144.
- 96 Ichikawa, H., Yokoyama, A., Hirata, K., Wada, H. and Kodera, K.: (Supports for gas chromatography from aggregates of fine carbon particles.) *Jpn. Kokai Tokkyo Koho* JP 05 02,012 [93 02,012] (Cl. G01N30/48), 8 Jan. 1993, Appl. 91/181,780, 26 Jun. 1991; 6 pp.; C.A., 118 (1993) 236901a.
- 97 Kawai, T.: (Method for formation of fine pores on the surface of a stationary phase for gas chromatography.) *Jpn. Kokai Tokkyo Koho* JP 04,303,766 [92,303,766] (Cl. G01N30/48), 27 Oct. 1992, Appl. 91/93,172, 30 Mar. 1991; 4 pp.; C.A., 118 (1993) 224649m.
- 98 Kou, D., Yun, X., Li, Y. and Lu, X.: (Quantitative analysis of trace organics by using gradient coated column.) *Sepu*, 11 (1993) 89-91.
- 99 Krestov, A.G., Blokhina, S.V., Galyametdinov, Yu.G., Ol'knovich, M.V. and Lokhanov, V.V.: (Sorbent based on a mesomorphic copper complex in the gas chromatography of organic compounds.) *Zh. Fiz. Khim.*, 67 (1993) 151-154.
- 100 Maas, B., Deitrich, A., Karl, V., Kaunzinger, A., Lehman, D., Köpke, T. and Mosandl, A.: *tert*-Butyldimethylsilyl-substituted cyclodextrin derivatives as versatile chiral stationary phase in capillary GC. *J. Microcolumn Sep.*, 5 (1993) 421-427.
- 101 Mazur, J., Dabrowski, R. and Witkiewicz, Z.: (Separation of mixture of organic compounds by gas chromatography.) *Pol. PL* 155,124 (Cl. G01N30/48), 3 Apr. 1992, Appl. 268,841, 17 Nov. 1987; 5 pp.; C.A., 119 (1993) 19608b.
- 102 Novikov, V.F.: (Organic derivatives of phosphorus and arsenic as stationary phases in gas chromatography.) *Zh. Fiz. Khim.*, 67 (1993) 848-853.
- 103 Okano, H., Tokunaga, T. and Ito, R.: (Metal capillary columns for gas chromatography.) *Jpn. Kokai Tokkyo Koho* JP 04,370,760 [92,370,760] (Cl. G01N30/60), 24 Dec. 1992, Appl. 91/147,521, 19 Jun. 1991; 4 pp.; C.A., 118 (1993) 194118t.
- 104 Onuchak, L.A., Vigdergauz, M.S. and Fedosova, E.V.: (Anisotropy of gas chromatographic sorption on silica fused capillary columns with nematic *p,p'*-azoxy-phenetol.) *Zh. Fiz. Khim.*, 67 (1993) 843-847.
- 105 Shalaeva, M.E., Zheivot, V.I., Fenelonov, V.B., Fridman, L.I. and Makakhov, V.V.: (Gas chromatographic studies of carbeneous adsorbents. Carbon fibrous material Actilen.) *Zh. Anal. Khim.*, 48 (1993) 1608-1614.
- 106 Vigdergauz, M.S.: (Use of liquid crystals in chromatography.) *Izv. Vyssh. Uchebn. Zaved., Khim. Khim. Tekhnol.*, 36, No. 1 (1993) 3-12; C.A., 118 (1993) 265762m - a review with 121 refs.
- 107 Vonk, N., de Zeeuw, J. and Buyten, J.: Ultimetal capillary columns for process GC analysis: A breakthrough. *Process Control Qual.*, 3 (1992) 137-144; C.A., 118 (1993) 204417f.
- 108 Wang, A. and Sun, Y.: (Qualitative analysis by using a serially connected binary gas chromatographic capillary columns.) *Sepu*, 11 (1993) 128-131.
- 109 Wasiaak, W.: Chemically bonded chelates as selective complexing sorbents for gas chromatography. II. Ketones, ethers and nitroalkanes. *J. Chromatogr. A*, 653 (1993) 63-69.
- 110 Wasiaak, W., Urbaniak, W., Obst, I. and Wawrzyniak, R.: Synthesis of silane with  $\beta$ -diketonate groups and their application in modification of silica gel for complexation gas chromatography. *Acta Chromatogr. (Katowice)*, No. 1 (1992) 56-68.
- 111 Wu, C.-Y., Cai, L.-S., Wang, Y.-J., Han, H.-M., Zeng, Z.-R., Yu, Z.-Y. and Yuan, H.: Preparation and gas chromatographic characterization of dibenzyl crown ether stationary phases. *Chromatographia*, 37 (1993) 374-380.
- 112 Xu, G., Zhang, Y. and Lu, P.: (Use of gas chromatography for sample requirement evaluation.) *Fenxi Huaxue*, 21 (1993) 276-281; C.A., 118 (1993) 265793x.
- 113 Xu, Y.: (Preparation of SCOT capillary columns with liquid crystal as stationary phase.) *Sepu*, 11 (1993) 217-218.

See also 12, 20, 21, 122, 131, 132, 161, 165, 168, 179, 233.

## 3d. Quantitative analysis

- 114 Dondi, F., Betti, A., Pasti, L., Pietrogrogrande, M.C. and Felinger, A.: Fourier analysis of multicomponent chromatograms. Application to experimental chromatograms. *Anal. Chem.*, 65 (1993) 2209-2222.
- 115 Rhoderick, G.C. and Miller, W.R.: Development of hydrocarbon gas standards. *J. Chromatogr. A*, 653 (1993) 71-81.

- 116 Soják, L., Farka, P., Kubinec, R. and Berezkin, V.G.: (The role of adsorption phenomena in capillary gas-liquid-solid chromatography.) *Zh. Anal. Khim.*, 48 (1993) 1604-1607.
- 117 Vigdergauz, M.S. and Krauze, I.M.: (Development of the method of quantitative interpretation of chromatograms by means of a double internal standard.) *Zh. Fiz. Khim.*, 67 (1993) 837-838.
- 118 Vigdergauz, M.S. and Krauze, I.M.: (Peculiarities of quantitative chromatographic analysis.) *Zh. Fiz. Khim.*, 67 (1993) 839-840.

See also 48, 73, 553.

### 3e. Preparative scale chromatography

- 119 Coca, J., Adrio, G., Jeng, C.Y. and Langer, S.H.: Gas and liquid chromatographic reactors. *Chromatogr. Sci.*, 61 (1993) 449-475; *C.A.*, 118 (1993) 215428m.
- 120 Schroll, W. and Nitz, S.: Preparative multidimensional gas chromatography with packed columns. *Chem., Mikrobiol., Technol. Lebensm.*, 14 (1992) 104-107; *C.A.*, 118 (1993) 187130n.

### 3f. Programmed temperature, pressure, vapors, gradients

- 121 Hawkes, S.J.: Viscosities of carrier gases at gas chromatograph temperatures and pressures. *Chromatographia*, 37 (1993) 399-401.
- 122 Messadi, D. and Ali-Mokhnache, S.: Calculation of retention indices and peak widths in temperature programmed gas-liquid chromatography. *Chromatographia*, 37 (1993) 264-270.

See also 173, 405.

## 4. SPECIAL TECHNIQUES

### 4a. Automation

- 123 Hurst, W.J. and Martin, R.A., Jr.: New analytical technologies and the challenge to automation. *Lab. Rob. Autom.*, 4 (1992) 193-196; *C.A.*, 119 (1993) 10945b.
- 124 Koskinen, M.O. and Koskinen, L.K.: Numerical integration of complex chromatograms using fitted gaussian functions. *J. Liq. Chromatogr.*, 16 (1993) 3171-3184.
- 125 Kurbatova, S.V., Kolosova, E.A. and Vigdergauz, M.S.: (Elaboration of computing-chromatographic system for the analysis of multicomponent mixtures.) *Zh. Fiz. Khim.*, 67 (1993) 833-836.
- 126 Shaeovich, A.B. and Mychnik, V.L.: (Computer-based chromatographic identification of compounds: Methodological peculiarities, algorithms, software, data bases.) *Zh. Fiz. Khim.*, 67 (1993) 827-832.
- 127 Smith, K.: Developments in the field of chromatography data handling. *Int. Lab.*, 23, No. 8 (1993) 20-22.
- 128 Vigdergauz, M.S., Kolosova, E.A. and Kurbatova, S.V.: (Automatic bases of chromatographic data: Structure and application.) *Zavod. Lab.*, 59, No. 6 (1993) 7-10.

See also 49, 107, 314, 501.

### 4b. Computerization and modelling

- 129 Dostál, M., Ventura, K. and Prudek, L.: (GC-CHAR: Programme for the calculation of gas-chromatographic characteristics.) *Chem. Listy*, 87 (1993) 814-818.
- 130 Gawlowski, J., Franaszczuk, K., Garnuszek, P., Niedzielski, J. and Bartulewicz, J.: Simple system for computer-aided chromatography. *Chem. Anal. (Warsaw)*, 37 (1992) 743-746.
- 131 Xu, G., Zhang, Y. and Lu, P.: (Software for recommendation of column systems in expert system on gas chromatography. III. Recommendation of stationary phases used for gas chromatographic analyses of organic compounds.) *Sepu*, 11 (1993) 65-68.
- 132 Xu, G., Zhang, Y. and Lu, P.: (The softwares for recommendation of column systems in expert system on gas chromatography: IV. Recommendation of stationary phases used for gas chromatographic analyses of inorganic gases and light hydrocarbons.) *Sepu*, 11 (1993) 131-135.

See also 125, 126.

### 4c. Combination with other physico-chemical techniques (MS, IR etc.)

- 133 Birch, E.: Solvent venting technique for gas chromatography with microwave-induced plasma atomic emission spectroscopy. *Anal. Chim. Acta*, 282 (1993) 451-458.
- 134 Carey, J.M., Byrde, F.A. and Caruso, J.A.: Alternate methods of sample introduction for plasma mass spectrometry. *J. Chromatogr. Sci.*, 31 (1993) 330-344 - a review with 120 refs.
- 135 Griffiths, P.R., Norton, K.L. and Bonanno, A.S.: SFC/FT-IR measurements involving elimination of the mobile phase. In: Jinno, K. (Editor), *Hyphenated Tech. Supercrit. Fluid Chromatogr. Extr.; J. Chromatogr. Libr.*, Vol. 53, Elsevier, Amsterdam, 1992, pp. 83-101; *C.A.*, 118 (1993) 204430e.
- 136 Jackson, P., Dent, G., Carter, D., Schofield, D.J., Chalmers, J.M., Visser, T. and Vredenburg, M.: Investigation of high sensitivity GC-FTIR as an analytical tool for structural identification. *J. High Resolut. Chromatogr.*, 16 (1993) 515-521.
- 137 Joern, W.A.: Ion ratio instability of a GC/MS system is related to fluctuations in room air flow. *J. Anal. Toxicol.*, 17 (1993) 122; *C.A.*, 118 (1993) 227686g.
- 138 Krock, K.A., Ragunathan, N. and Wilkins, C.L.: Parallel cryogenic trapping multidimensional gas chromatography with directly linked infrared and mass spectral detection. *J. Chromatogr.*, 645 (1993) 153-159.
- 139 McClellan, W.H., Buchanen, R.M., Arnold, N.S., Dworzanski, J.P. and Neuzelaar, H.L.C.: Thermogravimetry/gas chromatography/mass spectrometry and thermogravimetry/gas chromatography/Fourier transform infrared spectroscopy: Novel hyphenated methods in thermal analysis. *Anal. Chem.*, 65 (1993) 2819-2823.
- 140 Saito, M. and Yamauchi, Y.: Recycle supercritical fluid chromatography - on-line photodiode-array multiwavelength UV/VIS spectrometry/IR spectrometry/gas chromatography. In: Jinno, K. (Editor), *Hyphenated Tech. Supercrit. Fluid Chromatogr. Extr.; J. Chromatogr. Libr.*, Vol. 53, Elsevier, Amsterdam, 1992, pp. 129-149; *C.A.*, 118 (1993) 233109n.
- 141 Schwedt, G. and Meyer, A.: Kopplungstechniken in der Chromatographie. *LaborPraxis*, 17, No. 12 (1993) 12-21.

- 142 Tyrefors, L.N., Moulder, R.X. and Markides, K.E.: Interface for open tubular column supercritical fluid chromatography/atmospheric pressure chemical ionization mass spectrometry. *Anal. Chem.*, 65 (1993) 2835-2840.
- 143 Van Ysacker, P.G., Janssen, H.-G.M., Snijders, H.M.J., Leclercq, P.A., Cramers, C.A. and van Gruchten, H.J.M.: High-speed GC/MS using narrow-bore columns and ion trap detection. *J. Microcolumn Sep.*, 5 (1993) 413-419.
- 144 Xu, Y., Liu, J. and Gao, X.: (Gas chromatography combined with coulometry.) *Sepu*, 11 (1993) 95-97.
- See also 73, 176, 177, 189, 288, 289, 295, 300, 319, 341, 343, 354, 358, 365, 371, 375, 386, 391, 393, 440, 501, 512, 524.
- 4e. *Functional analysis*
- 145 Wei, J.-F. and Paulis, J.: Catalyzed acylation of phenols and aromatic amines with TFAC. *Microchem. J.*, 47 (1993) 2-9.
- 4f. *Trace analysis and preseparation techniques*
- 146 Bonin, M.A., Ashley, D.L., Cardinali, F.L., McCraw, J.M. and Patterson, D.G.J.: Importance of enhanced mass resolution in removing interferences when measuring volatile organic compounds in human blood by using purge-and-trap gas chromatography/mass spectrometry. *J. Am. Soc. Mass Spectrom.*, 3 (1992) 831-841; *C.A.*, 118 (1993) 208894g.
- 147 Cao, X.-L. and Hewitt, C.N.: Thermal desorption efficiencies for different adsorbate/adsorbent systems typically used in air monitoring programmes. *Chemosphere*, 27 (1993) 695-705.
- 148 Infante, R., Gutierrez, C. and Perez, C.: The use of liquid-solid extraction, capillary column GC/MS for the analysis of chlorinated acids and phenols in drinking water. *Water Sci. Technol.*, 26 (1992) 2583-2586; *C.A.*, 118 (1993) 131627n.
- 149 Jordan, L.: Automating a solid-phase extraction method. *LC-GC Int.*, 6 (1993) 594-599.
- 150 Kirsh, S.I., Kulagina, L.A. and Vigdergauz, M.S.: (Elution-displacement method for gas chromatographic determination of impurity components.) *Zh. Anal. Khim.*, 48 (1993) 1615-1620.
- 151 Kristiansen, N.K., Lundanes, E., Frshaug, M. and Becher, G.: Evaluation of the open-loop stripping technique used for the determination of volatile organic compounds in water. *Anal. Chim. Acta*, 280 (1993) 111-117.
- 152 Lucke, R.B., Campbell, J.A., Ross, G.A., Goheen, S.C. and Hoppe, E.W.: Closed-system, solid-phase extraction cleanup method for removal of normal paraffin hydrocarbon from samples prior to purge-and-trap volatile analysis. *Anal. Chem.*, 65 (1993) 2229-2235.
- 153 Matz, G. and Kesners, P.: Spray and trap method for water analysis by thermal desorption gas chromatography/mass spectrometry in field applications. *Anal. Chem.*, 65 (1993) 2366-2371.
- 154 Mitra, S. and Yun, C.: Continuous gas chromatographic monitoring of low concentration sample streams using an on-line microtrap. *J. Chromatogr.*, 648 (1993) 415-421.
- 155 Mol, H.G.J., Janssen, H.G. and Cramers, C.A.: Use of open-tubular trapping columns for on-line extraction—capillary gas chromatography of aqueous samples. *J. High Resolut. Chromatogr.*, 16 (1993) 413-418.
- 156 Morabito, P.L. and Melcher, R.G.: Automated membrane/gas chromatographic system for extraction and determination of trace organics in aqueous samples using large volume injection. *Process Control Qual.*, 3 (1992) 35-42; *C.A.*, 118 (1993) 194471c.
- 157 Takahata, J., Takahashi, Y. and Seki, T.: (Analytical method of pesticides used in golf courses. II. Analytical method by solid phase extraction and GC/MS.) *Sendai-shi Eisei Kenkyushoho*, 21 (1992) 262-264; *C.A.*, 118 (1993) 163046p.
- 158 Verweij, A.M.A. and Lipman, P.J.L.: Comparison of mass spectrometric techniques for the analysis of trace amount of 1-methylaminoanthraquinone, used as smoke dye in exploding money suitcases. *J. Chromatogr. A*, 653 (1993) 359-362.
- See also 37, 42, 92, 138, 186, 190, 217, 224, 244, 249, 255, 371, 512, 526, 558, 562.
- 4g. *Enantiomers, separation*
- 159 Hinshaw, J.V.: Separation of enantiomers by gas chromatography. *LC-GC*, 11 (1993) 644-648.
- 160 Hinshaw, J.V.: Separation of enantiomers by gas chromatography. *LC-GC Int.*, 6 (1993) 606-610.
- 161 Shitangkoon, A. and Vigh, G.: Gas chromatographic enantiomer separation using derivatized corn syrup polysaccharides as stationary phase. *J. High Resolut. Chromatogr.*, 16 (1993) 504-505.
- 162 Smith, I.D. and Simpson, C.F.: Investigation into the GC separation of enantiomers on a trifluoroacetylated cyclodextrin. Part II: Effect of derivatization on stereoselectivity towards alcohols. *J. High Resolut. Chromatogr.*, 16 (1993) 530-535.
- See also 100, 179, 180, 218, 272, 312, 317, 320, 334, 337, 350, 399.
- 4h. *Other special techniques*
- 163 Wang, T.S.: Adaptive multivariable PID control of a distillation column with unknown dead times. Avail. *Univ. Microfilms Int.*, Order No. DA9213859, 1991, 384 pp.; *C.A.*, 118 (1993) 150016d.
- See also 24, 53.
- 4i. *Supercritical fluid chromatography*
- 164 Burford, M.D., Hawthorne, S.B., Miller, D.J. and Macomber, J.: Construction of a robust stainless-steel clad fused-silica restrictor for use in supercritical fluid extraction. *J. Chromatogr.*, 648 (1993) 445-449.
- 165 Chester, T.L. and Innis, D.P.: Investigation of retention and selectivity in high-temperature, high-pressure, open-tubular supercritical fluid chromatography with CO<sub>2</sub> mobile phase. *J. Microcolumn Sep.*, 5 (1993) 441-449.
- 166 Gelencser, A. and de Leer, E.W.B.: (Testing the feasibility of using the ion-trap detector for monitoring environmental samples.) *Magy. Kem. Foly.*, 99 (1993) 127-130; *C.A.*, 118 (1993) 265807e.

- 167 Hill, H.H. and Atkinson, D.A.: General detection problems in SFC. In: Jinno, K. (Editor), *Hyphenated Tech. Supercrit. Fluid Chromatogr. Extr.*; *J. Chromatogr. Libr.*, Vol. 53, Elsevier, Amsterdam, 1992, pp. 1-8; *C.A.*, 118 (1993) 220264j.
- 168 Ibáñez, E., Alvarez, P.J.M., Reglero, G. and Herraiz, M.: Large particle micropacked columns in supercritical fluid chromatography. *J. Microcolumn Sep.*, 5 (1993) 371-381.
- 169 Jackson, P.: Advances of combined gas chromatography-mass spectrometry-matrix isolation Fourier transform infrared analysis. *Anal. Proc.*, 30 (1993) 394-395.
- 170 Karlsson, L., Järemo, M., Emilsson, M., Mathiasson, L. and Jönsson, J.Å.: Retention of coupled capillary column supercritical fluid chromatography with lipids as model compounds. *Chromatographia*, 37 (1993) 402-410.
- 171 Malik, A., Li, W. and Lee, M.L.: Preparation of long packed capillary columns using carbon dioxide slurries. *J. Microcolumn Sep.*, 5 (1993) 361-369.
- 172 Markides, K.E.: Instrumental analysis in bioanalysis, especially supercritical fluid chromatography. *Methodol. Surv. Biochem. Anal.*, 22 (1992) 279-289; *C.A.*, 118 (1993) 182537r - a review with 14 refs.
- 173 Pyo, D. and Huang, H.: Monitoring the mobile phase composition in supercritical fluid chromatography. *Anal. Chim. Acta*, 280 (1993) 103-109.
- 174 Raynie, D.E.: Warning concerning the use of nitrous oxide in supercritical fluid extractions. *Anal. Chem.*, 65 (1993) 3127-3128.
- 175 Riekkola, M.L., Manninen, P. and Hartonen, K.: SFE, SFE/GC and SFE/SFC: Instrumentation and applications. In: Jinno, K. (Editor), *Hyphenated Tech. Supercrit. Fluid Chromatogr. Extr.*; *J. Chromatogr. Libr.*, Vol. 53, Elsevier, Amsterdam, 1992, pp. 275-304; *C.A.*, 118 (1993) 229373b - a review with 42 refs.
- 176 Sadoun, F., Virelizier, H. and Arpino, P.J.: Packed-column supercritical fluid chromatography coupled with electrospray ionization mass spectrometry. *J. Chromatogr.*, 647 (1993) 351-359.
- 177 Takeuchi, M. and Saito, T.: Advances in semimicro packed column SFC and its hyphenation. In: Jinno, K. (Editor), *Hyphenated Tech. Supercrit. Fluid Chromatogr. Extr.*; *J. Chromatogr. Libr.*, Vol. 53, Elsevier, Amsterdam, 1992, pp. 47-63; *C.A.*, 118 (1993) 220265k.
- 178 Van Puyvelde, F., Chimowitz, E.H. and van Rompay, P.: Simulation and optimization in supercritical fluid chromatography. In: *Supercritical Fluid Engineering Science. ACS Symp. Ser.*, Vol. 514, ACS, Washington, 1993, pp. 281-293; *C.A.*, 118 (1993) 204409e.
- 179 Yi, G., Bradshaw, J.S., Rossiter, B.E., Reese, S.L., Petersson, P., Markides, K.E. and Lee, M.L.: Novel cyclodextrin-oligosiloxane copolymers for use as stationary phases to separate enantiomers in open tubular column supercritical fluid chromatography. *J. Org. Chem.*, 58 (1993) 2561-2565; *C.A.*, 118 (1993) 214069h.
- See also 4, 48, 62, 123, 134, 140, 142, 149, 181, 195, 242, 248, 253, 264, 280, 281, 301, 306, 380, 382, 393, 415, 422, 492, 527, 543.
5. HYDROCARBONS AND HALOGEN DERIVATIVES
- 5a. *Aliphatic hydrocarbons*
- 180 Boneva, S. and Balbolov, E.: Capillary gas chromatography of hydrogenation products of 1,5,9-cyclododecatriene stereoisomers. *Chromatographia*, 37 (1993) 277-280.
- 181 Bounoshita, M., Saito, M., Hibi, K., Jinno, K., Nagashima, H. and Itoh, K.: (Supercritical fluid extraction and chromatography of fullerenes.) *Kuromatogurafi*, 13 (1992) 357-358. *C.A.*, 118 (1993) 265571y.
- 182 Hu, Z. and Zhang, H.: Prediction of gas chromatographic retention indices of alkenes from the total solubility parameters. *J. Chromatogr. A*, 653 (1993) 275-282.
- 183 Moodie, I.M. and Labadarios, D.: Determination of pentane in human breath by gas chromatography. *J. High Resolut. Chromatogr.*, 16 (1993) 437-440.
- 184 Orav, A., Kuningas, K. and Rang, S.: A comparison of different retention index systems with unsaturated and aromatic hydrocarbons in capillary gas chromatography on PEG 20M. *Chromatographia*, 37 (1993) 411-414.
- 185 Pritts, W.A., Vieira, K.L. and Peters, D.G.: Quantitative determination of volatile products formed in electrolyses of organic compounds. *Anal. Chem.*, 65 (1993) 2145-2149.
- 186 Rohwer, E.R., Chen, G.-C. and Hassett, A.J.: Preparation and storage of ultra high purity hexane for solvent effect gas chromatographic analysis. *J. High Resolut. Chromatogr.*, 16 (1993) 561-564.
- 187 Tang, Y.Z., Tran, Q., Fellin, P., Cheng, W.K. and Drummond, I.: Determination of C<sub>1</sub>-C<sub>4</sub> hydrocarbons in air. *Anal. Chem.*, 65 (1993) 1932-1935.
- 188 Yu, Y.: (Improvement of gas chromatographic analysis of commercial butadiene.) *Hecheng Xiangjiao Gongye*, 15 (1992) 363; *C.A.*, 118 (1993) 193414z.
- See also 115, 132, 504, 508, 538.
- 5b. *Cyclic hydrocarbons, fullerenes*
- 189 Diehl, J.W., Finkbeiner, J.W. and DiSanzo, F.P.: Determination of benzene, toluene, ethylbenzene, and xylenes in gasolines by gas chromatography/deuterium isotope dilution Fourier transform infrared spectroscopy. *Anal. Chem.*, 65 (1993) 2493-2496.
- 190 Ehret-Henry, J., Ducruet, V., Luciani, A. and Feigenbaum, A.: Purge and trap gas chromatography determination of styrene and ethylbenzene. Performance in model medium. *Analisis*, 21 (1993) 277-280.
- 191 Heinzen, V.E.F. and Yunes, R.A.: Correlation between gas chromatographic retention indices of linear alkylbenzene isomers and molecular connectivity indices. *J. Chromatogr. A*, 654 (1993) 183-189.
- 192 Kubinec, R., Kurán, R., Ostrovsky, I. and Soják, L.: Determination of polycyclic aromatic hydrocarbons from bitumen concrete roads in drainage water by microextraction, large-volume sampling and gas chromatography-mass spectrometry with selected ion monitoring. *J. Chromatogr. A*, 653 (1993) 363-368.

- 193 Lancas, F.M. and Peralba, M.C.R.: CGC and GC/MS identification of aromatic compounds in bitumen and its correlation to geochemical properties. *Fuel Sci. Technol. Int.*, 11 (1993) 541-560; *C.A.*, 118 (1993) 194821s.
- 194 Lee, H.-B., Peart, T.E., Hong-You, R.L. and Gere, D.R.: Super-critical carbon dioxide extraction of polycyclic aromatic hydrocarbons from sediments. *J. Chromatogr. A*, 653 (1993) 83-91.
- 195 Levy, J.M., Dolata, L.A. and Ravey, R.M.: Considerations of SFE for GC/MS determination of polynuclear aromatic hydrocarbons in soils and sediments. *J. Chromatogr. Sci.*, 31 (1993) 349-352.
- 196 Ni, B. and Hu, X.: (Determination of blood benzene by gas chromatographic headspace analysis.) *Sepu*, 11 (1993) 121-122.
- 197 Nondek, L., Kuzílek, M. and Krupicka, Š.: LC clean-up and GC/MS analysis of polycyclic aromatic hydrocarbons in river sediment. *Chromatographia*, 37 (1993) 381-386.
- 198 Škrbic, B.D. and Cvejanov, J.Dj.: Unified retention indices of alkylbenzenes on OV-101 and SE-30. *Chromatographia*, 37 (1993) 215-217.
- 199 Tian, S.: (Analysis of the tower bottom oil of dimethylbenzene rectifying tower and C<sub>9</sub> aromatic hydrocarbon fraction by capillary gas chromatography.) *Sepu*, 11 (1993) 202-206.
- 200 Ye, F., Zhang, X., Di, L., Luo, C. and Zhang, Y.: (The optimization for separation of PNA in naphtha.) *Sepu*, 11 (1993) 156-158.
- See also 110, 147, 184, 470, 506, 519, 537.
- 5c. *Halogen derivatives*
- 201 Blum, W. and Aichholz, R.: High temperature stable CH<sub>3</sub>O-terminated poly(diphenyl/dimethyl)- and poly(diphenyl/3,3,3-trifluoropropylmethyl)siloxane copolymer stationary phases for capillary gas chromatography. *J. Microcolumn Sep.*, 5 (1993) 297-302.
- 202 Cammann, K. and Hübner, K.: False results in headspace-gas chromatographic analysis of trihalomethanes in swimming pool water due to elevated headspace temperatures. *J. Chromatogr.*, 648 (1993) 294-298.
- 203 Daft, J.L.: Methyl bromide determination in selected foods by headspace technique. *J. Assoc. Off. Anal. Chem.*, 76 (1993) 1083-1091.
- 204 Foerst, C., Simon, H. and Stieglitz, L.: Determination of chlorophenols and chlorobenzenes in leachate by headspace analysis. *Chemosphere*, 26 (1993) 1355-1365.
- 205 Galceran, M.T., Santos, F.J., Caixach, J. and Rivera, J.: PCBs and chlorinated pesticides in shellfish of a deltaic environment. *Chemosphere*, 27 (1993) 1183-1200.
- 206 Guo, Q.: (Determination of hydrocarbons in the air by gas chromatography with direct injection.) *Tianranqi Huagong*, 17 (1992) 50-53; *C.A.*, 119 (1993) 14297q.
- 207 Han, H. and Liu, C.: (Determination of vinyl chloride monomer in water by gas chromatography at sub- $\mu$ g/L level.) *Sepu*, 11 (1993) 171-172.
- 208 Himberg, K.K.: Coplanar polychlorinated biphenyls in some Finnish food commodities. *Chemosphere*, 27 (1993) 1235-1243.
- 209 Jalali-Heravi, M. and Garkani-Nejad, Z.: Prediction of gas chromatographic retention indices of some benzene derivatives. *J. Chromatogr.*, 648 (1993) 389-393.
- 210 Junk, S.A. and Meisch, H.-U.: Determination of chlorinated paraffins by GC-MS. *Fresenius J. Anal. Chem.*, 347 (1993) 361-364.
- 211 Kuwabara, K., Andoh, K., Nishimune, T., Sueki, K. and Fujita, M.: (Studies on analysis of volatile halogenated hydrocarbons (VHHs) in food. Trial construction of clean air chamber for the removal of VHHs in the air coming into food sample and the results of VHH measurement in food by means of duplicate portion study in 1991 and 1992.) *Osaka-furitsu Koshu Eisei Kenkyusho Kenkyu Hokoku, Shokuhin Eisei Hen*, 23 (1992) 29-40; *C.A.*, 118 (1993) 190223a.
- 212 Kvitka, L.A., Makhno, A.Ya., Smirnov, D.N. and Shevchuk, I.A.: (Method for determining organochlorine compounds in water by gas chromatography.) *U.S.S.R. SU 1,775,663 (Cl. G01N30/90)*, 15 Nov. 1992, Appl. 4,850,393, 28 Jun. 1990; *C.A.*, 119 (1993) 19627g.
- 213 Morosini, M., Schreitmüller, J., Reuter, U. and Ballschmiter, K.: Correlation between C-6/C-14 chlorinated hydrocarbons levels in the vegetation and in the boundary layer of the troposphere. *Environ. Sci. Technol.*, 27 (1993) 1517-1523.
- 214 Nakanishi, S. and Hino, T.: (Determination of volatile chlorinated hydrocarbons in soil using sampling vials for storage and extraction by means of water-hexane extraction method.) *Jpn. J. Toxicol. Environ. Health*, 39 (1993) 29-36; *C.A.*, 118 (1993) 211881f.
- 215 Nilsson, U.L. and Östman, C.E.: Chlorinated polycyclic aromatic hydrocarbons: methods of analysis and their occurrence in urban air. *Environ. Sci. Technol.*, 27 (1993) 1826-1831.
- 216 Ozawa, H., Murooka, M. and Matsubara, H.: (Determination of volatile halogenated hydrocarbons in water samples by purge and trap-capillary GC/MS.) *Zenkoku Kogaiken Kaishi*, 17 (1992) 189-193; *C.A.*, 119 (1993) 14959g.
- 217 Page, B.D. and Lacroix, G.: Application of solid-phase microextraction to the headspace gas chromatographic analysis of halogenated volatiles in selected foods. *J. Chromatogr.*, 648 (1993) 199-211.
- 218 Schurig, V. and Glausch, A.: Enantiomer separation of atropisomeric polychlorinated biphenyls (PCBs) by gas chromatography on Chirasil-Dex. *Naturwissenschaften*, 80 (1993) 468-469.
- 219 Spencer, J.L., Hendricks, J.P. and Kerr, D.: Least-squares analysis of gas chromatographic data for polychlorinated biphenyl mixtures. *J. Chromatogr. A*, 654 (1993) 143-149.
- 220 Sturrock, G.A., Simmonds, P.G., Nickless, G. and Zwiep, D.: Analysis of chlorofluorocarbon replacement compounds by capillary gas chromatography. *J. Chromatogr.*, 648 (1993) 423-431.
- 221 Sugiyama, H. and Abe, A.: (Gas chromatographic determination of volatile chlorinated hydrocarbons in soil by a technique of thermal desorption with water.) *Bunseki Kagaku*, 42 (1993) 531-536.
- 222 Vartiainen, T., Pukkala, E., Rienoja, T., Strandman, T. and Kakkonen, K.: Population exposure to tri- and tetrachloroethene and cancer risk: two cases of drinking water pollution. *Chemosphere*, 27 (1993) 1171-1181.
- 223 Williams, D.T., Kennedy, B. and LeBel, G.L.: Chlorinated naphthalenes in human adipose tissue from Ontario municipalities. *Chemosphere*, 27 (1993) 795-806.

- 224 Wittsiepe, J., Wallschläger, D., Selenka, F. and Jackwerth, E.: Gas-chromatographic determination of trace amounts of vinyl chloride and dichloromethane in water after purge-and-trap pre-concentration and conversion into their dibrominated derivatives. *Fresenius J. Anal. Chem.*, 346 (1993) 1028-1034.
- 225 Zenkevich, I.G., Konyukhova, S.V. and Maksimov, B.N.: (Dependence of gas chromatographic retention parameters of simple halogen containing compounds on their physico-chemical characteristics.) *Zh. Fiz. Khim.*, 67 (1993) 1474-1479.

See also 18, 83, 470, 505.

5d. *Complex hydrocarbon mixtures (incl. analysis of tars, bitumens and mineral oils)*

See 43, 152, 498, 525.

6. ALCOHOLS

- 226 Artyushina, G.G. and Stan'kov, I.N.: (Gas chromatographic separation and determination of tertiary hexyl alcohols having different substituents to the tertiary carbon atoms.) *Zh. Anal. Khim.*, 48 (1993) 1175-1179.
- 227 Gogineni, P.V., Crooks, P.A. and Murty, R.B.: Gas chromatographic determination of residual levels of *tert.*-butanol from lyophilized liposomal formulations. *J. Chromatogr.*, 620 (1993) 83-88.
- 228 Hallett, R.H., Gries, G., Gries, R., Borden, J.H., Czyzewska, E., Oehlschlager, A.C., Pierce, H.D., Jr., Angerilli, N.P.D. and Rauf, A.: Aggregation pheromones of two Asian palm weevils, *Rhyncophorus ferrugineus* and *R. vulneratus*. *Naturwissenschaften*, 80 (1993) 328-331.
- 229 Houzé, P., Chaussard, J., Harry, P. and Pays, M.: Simultaneous determination of ethylene glycol, propylene glycol, 1,3-butylene glycol and 2,3-butylene glycol in human serum and urine by wide-bore column gas chromatography. *J. Chromatogr.*, 619 (1993) 251-257.
- 230 Kern, F. and Bestmann, H.J.: Antennal electrophysiological responsiveness of the ponerine ant *Leptogenys diminuta* to trail and recruitment pheromones and its structure analogs. Pheromones, 94. *Naturwissenschaften*, 80 (1993) 424-427.
- 231 Molina, F., Bosque-Sendra, J.M., Pla, A. and Megias, L.: Determination of ethanol in biological samples by head-space gas chromatography. *J. Pharm. Biomed. Anal.*, 10 (1992) 1069-1071; *C.A.*, 118 (1993) 185230w.
- 232 Vitali, M., Leoni, V., Chiavarini, S. and Cremisini, C.: Determination of 2-ethyl-1-hexanol as contaminant in drinking water. *J. Assoc. Off. Anal. Chem.*, 76 (1993) 1133-1137.

See also 162, 254, 474.

7. PHENOLS

- 233 Andronikashvili, T.G., Gvelesiani, Z.A., Kiknadze, L.O. and Dzagania, M.A.: (Separation of cresol isomers by gas chromatography on a liquid-crystalline stationary phase.) *Izv. Akad. Nauk Gruz., Ser. Khim.*, 17 (1991) 247-252; *C.A.*, 118 (1993) 204434j.
- 234 English, S., Greenaway, W. and Whatley, F.R.: Analysis of phenolics of bud exudate of *Populus tristis* by GC/MS. *Z. Naturforsch., C: Biosci.*, 47 (1992) 512-515; *C.A.*, 118 (1993) 77033a.
- 235 Kvernheim, A.L., Martinsen, K., Carlberg, G.E., Berg, B.E., Fresvig, M. and Greibrokk, T.: Characterization of organohalogen matter in sediments from the receiving water of a bleach plant. *Chemosphere*, 27 (1993) 733-745.
- 236 Louch, J.R., LaFleur, L.E., Wilson, G., Bautz, D., Woodrow, D., Teitzel, H., Jones, J. and Mark, M.: Development and validation of analytical methods for the determination of phenolic compounds in pulp and wastewater treatment plant sludges. *Tappi J.*, 76, No. 3 (1993) 71-80; *C.A.*, 119 (1993) 14725c.
- 237 Nabivach, V.M. and Burjan, P.: (Pyrolysis of higher *o*-monoalkylphenols.) *Koks Khim.*, No. 8 (1991) 20-22; *C.A.*, 118 (1993) 215271e.

See also 145, 148, 259, 332, 443, 537.

8. SUBSTANCES CONTAINING HETEROCYCLIC OXYGEN

8b. *Aflatoxins and other mycotoxins*

- 238 Braun, T., Maczelka, H. and Zsindely, S.: Lags in the publication of research findings in analytical chemistry. *TrAC*, 12 (1993) 309-313.
- 239 Raza, S.K., Howell, S.A. and Mallet, A.I.: Identification of mycotoxins in keratomycosis-derived *Fusarium* isolates by gas chromatography-mass spectrometry. *J. Chromatogr.*, 620 (1993) 243-249.
- 240 Scott, P.M.: Recent developments in methods of analysis for mycotoxins in foodstuffs. *TrAC*, 12 (1993) 373-381 - a review with 37 refs.
- 241 Seidel, V., Lang, B., Fraissler, S., Lang, C., Schiller, K., Filek, G. and Lindner, W.: Analysis of trace levels of trichothecene mycotoxins in Austrian cereals by gas chromatography with electron capture detection. *Chromatographia*, 37 (1993) 191-201.
- 242 Young, J.C. and Games, D.E.: Analysis of *Fusarium* mycotoxins by supercritical fluid chromatography with ultraviolet or mass spectrometric detection. *J. Chromatogr. A*, 653 (1993) 374-379.
- 243 Zotou, A., Jefferies, T.M., Brough, P.A. and Gallagher, T.: Determination of anatoxin-a and homoanatoxin-a in blue-green algal extracts by high-performance liquid chromatography and gas chromatography-mass spectrometry. *Analyst (Cambridge)*, 118 (1993) 753-758.

## 8c. Other compounds with heterocyclic oxygen (incl. tannins)

- 244 Chang, R.R., Jarman, W.M. and Hennings, J.A.: Sample clean-up by solid-phase extraction for the ultratrace determination of polychlorinated dibenzo-*p*-dioxins and dibenzofurans in biological samples. *Anal. Chem.*, 65 (1993) 2420-2427.
- 245 Galletti, G.C., Ward, R.S., Pelter, A. and Goubet, D.: Generation of isobenzofuran derivatives by pyrolysis-gas chromatography/mass spectrometry. *J. Anal. Appl. Pyrolysis*, 24 (1992) 139-146.
- 246 King, T.L., Uthe, J.F. and Musial, C.J.: Rapid screening of fish tissue for polychlorinated dibenzo-*p*-dioxins and dibenzofurans. *Analyst (Cambridge)*, 119 (1993) 1269-1275.
- 247 Kronberg, L. and Franzén, R.: Determination of chlorinated furanones, hydroxyfuranones, and butenedioic acids in chlorine-treated water and in pulp bleaching liquor. *Environ. Sci. Technol.*, 27 (1993) 1811-1818.
- 248 Onuska, F.I., Terry, K.A. and Wilkinson, R.J.: The analysis of chlorinated dibenzofurans in municipal fly ash: supercritical fluid extraction vs Soxhlet. *J. High Resolut. Chromatogr.*, 16 (1993) 407-412.
- 249 Qiu, Y., Ou, Q. and Yu, W.: (SIM/GC/ITD determination of polychlorinated dibenzo-*p*-dioxins at parts per trillion level in soil.) *Fenxi Ceshi Tongbao*, 11, No. 2 (1992) 67-70; *C.A.*, 119 (1993) 14723a.
- 250 Thompson, T.S., Kolic, T.M., Townsend, J.A. and Mercer, R.S.: Determination of polychlorinated dibenzo-*p*-dioxins and dibenzofurans in tire fire runoff oil. *J. Chromatogr.*, 648 (1993) 213-219.
- 251 Zelinski, V., Lorenz, W. and Bahadir, M.: Brominated flame retardants and resulting PBDD/F in accidental fire residues from private residences. *Chemosphere*, 27 (1993) 1519-1528.

See also 276.

## 9. OXO COMPOUNDS, ETHERS, EPOXIDES AND QUINONES

- 252 Bakola-Christianopoulou, M.N., Papageorgiou, V.P. and Apazidou, K.K.: Gas chromatographic-mass spectrometric study of the reductive silylation of hydroxyquinones. *J. Chromatogr.*, 645 (1993) 293-301.
- 253 Kane, M., Dean, J.R., Hitchen, S.M., Tomlinson, W.R., Tranter, R.L. and Dowle, C.J.: Quantitative structure-extraction relationships: A model for supercritical fluid extraction. *Analyst (Cambridge)*, 118 (1993) 1261-1264.
- 254 Kissa, E.: Determination of acetylacetone and alcohols in titanium chelates. *Anal. Chim. Acta*, 281 (1993) 385-389.
- 255 Otson, R., Fellin, P., Tran, Q. and Stoyanoff, R.: Examination of sampling methods for assessment of personal exposures of airborne aldehydes. *Analyst (Cambridge)*, 118 (1993) 1253-1259.
- 256 Park, P.S.W.: Loss of volatile lipid oxidation products during thermal desorption in dynamic headspace-capillary gas chromatography. *J. Food Sci.*, 58 (1993) 220-222; *C.A.*, 118 (1993) 253484j.
- 257 Perkel', A.L., Freidin, B.G., Voronina, S.G. and Perkel', R.L.: (Determination of carbonyl-containing compounds in products of organic matter oxidation by gas chromatographic and photometric methods.) *Zh. Anal. Khim.*, 48 (1993) 1399-1406.

- 258 Sepai, O., Anderson, D., Street, B., Bird, I., Farmer, P.B. and Bailey, E.: Monitoring of exposure to styrene oxide by GC-MS analysis of phenylhydroxyethyl esters in hemoglobin. *Arch. Toxicol.*, 67 (1993) 28-33; *C.A.*, 119 (1993) 2535c.
- 259 Smith, T.J., Wearne, R.H. and Wallis, A.F.A.: Determination of chlorinated benzaldehydes and acetophenones in pulp bleaching effluents by gas chromatography. *J. Chromatogr.*, 648 (1993) 289-293.
- 260 Snell, R.P.: Gas chromatographic determination of cyclohexanone leached from hemodialysis tubing. *J. Assoc. Off. Anal. Chem.*, 76 (1993) 1127-1132.

See also 109, 158, 228.

## 10. CARBOHYDRATES

## 10a. Mono and oligosaccharides. Structural studies

- 261 Beylot, M., Previs, S.F., David, F. and Brunnengraber, H.: Determination of the <sup>13</sup>C-labelling pattern of glucose by gas chromatography-mass spectrometry. *Anal. Biochem.*, 212 (1993) 526-531.
- 262 Lin, Y.Y., Cheng, W.B. and Wright, C.E.: Regulation of glucose metabolism in cultured cells by mass isotopomer analysis. *Nippon Iyo Masu Supekutoru Gakkai Koenshu*, 17 (1992) 239-244; *C.A.*, 119 (1993) 4364v.
- 263 Seymour, F.R.: Identification and characterization of saccharides by GLC (gas-liquid chromatography) separation and MS analysis of their peracetylated aldononitrile (PAAN) and ketoxime (PAKO) derivatives. *Methods Carbohydr. Chem.*, 9 (1993) 59-85; *C.A.*, 118 (1993) 224581h - a review with 51 refs.
- 264 Xiang, Z.: (Separation of monosaccharides by supercritical fluid chromatography (SFC) with packed column.) *Sepu*, 11 (1993) 160-162.

See also 472.

## 10c. Glycoproteins and their constituents

- 265 Kawamura, Y., Hikidi, S., Maruyama, K., Uchiyama, S. and Saito, Y.: (Improvement of the direct determination method of linamarin in beans and bean paste products.) *Shokuhin Eiseigaku Zasshi*, 34 (1993) 74-79; *C.A.*, 118 (1993) 211510j.
- 266 Krotkiewski, H., Lisowska, E., Nilsson, G., Groenberg, G. and Nilsson, B.: An improved approach to the analysis of the structure of small oligosaccharides of glycoproteins: application to the O-linked oligosaccharides from human glycoprotein A. *Carbohydr. Res.*, 239 (1993) 35-50; *C.A.*, 119 (1993) 4371v.

## 11. ORGANIC ACIDS AND LIPIDS

## 11a. Organic acids and simple esters

- 267 Ackman, R.G.: Application of gas-liquid chromatography to lipid separation and analysis: qualitative and quantitative analysis. *Food Sci. Technol. (N. Y.)*, 53 (1992) 47-63; *C.A.*, 119 (1993) 7356e - a review with 57 refs.

- 268 Ackman, R.G.: Gas-liquid chromatography and the arachidonic acid content of Atlantic food fish. *Food Res. Int.*, 25 (1992) 453-456; C.A., 118 (1993) 253473e.
- 269 Aitzetmüller, K.: Capillary GLC fatty acid fingerprints of seed lipids – a tool in plant chemotaxonomy? *J. High Resolut. Chromatogr.*, 16 (1993) 488-490.
- 270 Akira, K., Nakamura, T., Shinohara, Y. and Baba, S.: Profiling of arachidonic acid metabolites in rabbit platelets by radio gas chromatography. *Lipids*, 28 (1993) 361-364.
- 271 Ballesteros, E., Gallero, M. and Valcárcel, M.: Automatic method for on-line preparation of fatty acid methyl esters from olive oil and other types of oil prior to their gas chromatographic determination. *Anal. Chim. Acta*, 282 (1993) 581-588.
- 272 Belfield, K.D., Hofmeister, T.S. and Seo, J.: Direct determination of enantiomeric excess of carboxylic esters by chiral capillary gas chromatography. *J. Chromatogr.*, 648 (1993) 497-500.
- 273 Garcia-Barceló, M., Luquin, M., Belda, F. and Ausina, V.: Gas chromatographic whole-cell fatty acid analysis as an aid for the identification of mixed mycobacterial cultures. *J. Chromatogr.*, 617 (1993) 299-303.
- 274 Hu, Z.: (Systematic analysis of various fatty acids in plants by gas chromatography (GC) and gas chromatography–mass spectrometry (GC-MS).) *Sepu*, 11 (1993) 197-199.
- 275 Li, X.: (Analysis of  $\alpha$ -isopropyl-*p*-chlorobenzylacetyl chloride by gas chromatography.) *Huaxue Shijie*, 33 (1992) 314-316; C.A., 118 (1993) 185712e.
- 276 Lisi, A.M., Kazlauskas, R. and Trout, G.J.: Gas chromatographic–mass spectrometric quantitation of urinary 11-nor- $\Delta^9$ -tetrahydrocannabinol-9-carboxylic acid after derivatization by direct extractive alkylation. *J. Chromatogr.*, 617 (1993) 265-270.
- 277 Lladó, I., Palou, A. and Pons, A.: Combined enzymic and chromatographic techniques to determine specific radioactivity in free and triglyceride fatty acid plasma fractions. *J. Chromatogr.*, 619 (1993) 21-28.
- 278 May, W.E. and Hume, D.J.: An automated gas-liquid chromatographic method of measuring free fatty acids in canola. *J. Am. Oil Chem. Soc.*, 70 (1993) 229-233.
- 279 Rudenko, S.D., Shcherbina, F.F., Voloshin, N.L., Zerzeva, I.M., Glavati, O.L. and Kravchuk, G.G.: (Gas-liquid chromatography of esters of alkylsalicylic and alkylhydroxyisophthalic acids.) *Neftepererab. Neftekhim. (Kiev)*, 43 (1992) 62-64; C.A., 119 (1993) 19584r.
- 280 Sakaki, K.: Supercritical fluid chromatographic separation of fatty acid methylesters on aminopropyl-bonded silica stationary phase. *J. Chromatogr.*, 648 (1993) 451-457.
- 281 Staby, A., Borch-Jensen, C., Møllerup, J. and Jensen, B.: Flame ionization detector responses to ethyl esters of sand eel (*Ammodytes lancea*) fish oil compared for different gas and supercritical fluid chromatographic systems. *J. Chromatogr.*, 648 (1993) 221-232.
- 282 Straczek, J., Felden, F., Douset, B., Gueant, J.L. and Belleville, F.: Quantification of methylmalonic acid in serum measured by capillary gas chromatography–mass spectrometry as *tert*-butyldimethylsilyl derivatives. *J. Chromatogr.*, 620 (1993) 1-7.
- 283 Stufka-Olczyk, J. and Pszonka, B.: Determination of fatty and resin acids in wastewater after biological treatment using capillary gas chromatography. *Chem. Anal. (Warsaw)*, 37 (1992) 307-312.
- 284 Sweetman, L.: Quantitative GCMS of urinary organic acids, plasma methylmalonic acid and plasma and urine propionic and isovaleric acids. *Nippon Iyo Masu Supekutoru Gakkai Koenshu*, 17 (1992) 49-53; C.A., 119 (1993) 4363u.
- 285 Wanke, T. and Eberle, S.H.: (Gas chromatographic determination of diethylenetriaminepentaacetic acid in surface water.) *Acta Hydrochim. Hydrobiol.*, 20 (1992) 192-196; C.A., 118 (1993) 260596a.
- 286 Yang, Y., Yang, Z. and Li, S.: (Determination of *ortho*-*tert*-butylcyclohexyl acetate by capillary gas chromatography.) *Sepu*, 11 (1993) 247-249.
- 287 Ye, X. and Zhu, G.: (Determination of fatty acids in suspended matter by capillary gas chromatography.) *Haiyang Yu Huzhao*, 23 (1992) 669-676; C.A., 118 (1993) 219316w.

See also 18, 100, 148, 335, 338, 464, 472.

#### 11b. Prostaglandins

- 288 Chiabrando, C., Rivoltella, L., Alberti, E., Bonollo, M., Djurup, R. and Fanelli, R.: Urinary excretion and origin of 11-dehydro-2,3-dinor-thromboxane B<sub>2</sub> in man. *Prostaglandins*, 45 (1993) 401-411.
- 289 Hizugaki, M., Nishikawa, M., Hishinuma, T., Ohyama, Y. and Ishibashi, M.: Analysis of the eicosapentanoic acid (EPA) metabolite  $\Delta^{17}$ -6-keto-PGF 1 $\alpha$  in human plasma by GC/SIM using [<sup>18</sup>O] $\Delta^{17}$ -6-keto-PGF 1 $\alpha$ . *Prostaglandins*, 44 (1992) 317-327.
- 290 Knott, I., Raes, M., Dieu, M., Lenoir, G., Burton, M. and Remacle, J.: Routine prostaglandin assay by GC-MS in multiwell tissue culture plates: Application to human synoviocytes and chondrocytes. *Anal. Biochem.*, 210 (1993) 360-365.

See also 448.

#### 11c. Lipids and their constituents

- 291 Khaled, M.Y., McNair, H.M. and Hanson, D.J.: High temperature chromatographic analysis of monoacylglycerols and diacylglycerols. *J. Chromatogr. Sci.*, 31 (1993) 375-379.
- 292 Lu, X.J., Myers, M.R. and Artz, W.E.: Supercritical fluid chromatographic analysis of the propoxylated glycerol esters of oleic acid. *J. Am. Oil Chem. Soc.*, 70 (1993) 355-360.

See also 170, 277.

#### 12. ORGANIC PEROXIDES

- 293 Perkel, A.L., Freidin, B.G. and Voronina, S.G.: (Elimination of the interferences from peroxide compounds in gas chromatographic determination of hydroxyl-carbonyl-containing and ester products of organic matter oxidation.) *Zh. Anal. Khim.*, 48 (1993) 1697-1705.
- 294 Polzer, J. and Bächmann, K.: Sensitive determination of alkyl hydroperoxides by high-resolution gas chromatography–mass spectrometry and high-resolution gas chromatography with flame ionization detection. *J. Chromatogr. A*, 653 (1993) 283-291.



- 295 Turnipseed, S.B., Allentoff, A.J. and Thompson, J.A.: Analysis of trimethylsilylperoxy derivatives of thermally labile hydroperoxides by gas chromatography-mass spectrometry. *Anal. Biochem.*, 213 (1993) 218-225.

See also 37.

### 13. STEROIDS

#### 13b. Pregnane and androstane derivatives

- 296 Zhang, J., Liu, C. and Zhou, T.: (Analysis of metabolites of the contraceptive norethisterone in urine by GC-MS.) *Gaodeng Xuexiao Huaxue Xuebao*, 14 (1993) 184-186; *C.A.*, 118 (1993) 205356x.

#### 13c. Estrogens

- 297 Chappell, C.G., Creaser, C.S. and Shepherd, M.J.: On-line high performance liquid chromatography-multidimensional gas chromatography and its application to the determination of stilbene hormones in corned beef. *J. High Resolut. Chromatogr.*, 16 (1993) 479-482.
- 298 Moretti, G., Citti, G., Cammarata, P. and Cozzani, R.: (Determination of synthetic estrogens in animal excretions using GC-MS.) *Riv. Soc. Ital. Sci. Aliment.*, 21 (1992) 135-140; *C.A.*, 118 (1993) 205345t.

See also 449.

#### 13d. Sterols

- 299 Lognag, G.C., Lacoste, F., Marlier, M., Mordret, F., Auge, C., Raoux, R., Wagstaffe, P.J., Boenke, A. and Severin, M.: The certification of the identity of individual sterols in three BCR oil and fat reference materials by gas chromatography-mass spectrometry (GC-MS). *Fat Sci. Technol.*, 95 (1993) 98-104.
- 300 Neese, R.A., Faix, D., Kletke, C., Wu, K., Wang, A.C., Shackleton, C.H.L. and Hellerstein, M.K.: Measurement of endogenous synthesis of plasma cholesterol in rats and humans using MIDA. *Am. J. Physiol.*, 264 (1993) E136-E147; *C.A.*, 118 (1993) 187298y.
- 301 Nomura, A., Yamada, J., Takatsu, A., Horimoto, Y. and Yarita, T.: Supercritical fluid chromatographic determination of cholesterol and cholesteryl esters in serum on ODS-silica gel column. *Anal. Chem.*, 65 (1993) 1994-1997.
- 302 Plank, C. and Lorbeer, E.: Analysis of free and esterified sterols in vegetable oil methyl esters by capillary GC. *J. High Resolut. Chromatogr.*, 16 (1993) 483-487.
- 303 Tamasawa, N. and Takebe, K.: Analytical examination of oxidized free and esterified 7-ketocholesterol and related oxysterols in human plasma incubated with copper. *Tohoku J. Exp. Med.*, 168 (1992) 37-45; *C.A.*, 118 (1993) 208708z.
- 304 Thompson, R.H. and Merola, G.V.: A simplified alternative to the AOAC official method for cholesterol in multicomponent foods. *J. Assoc. Off. Anal. Chem.*, 76 (1993) 1057-1068.
- 305 Yuan, H. and Zeng, Z.: (The determination of sterols in three species of *Spirulina* by using GLC/MS.) *Zhiwu Ziyuan Yu Huanjing*, 1, No. 4 (1992) 57-61; *C.A.*, 118 (1993) 187419p.

### 15. TERPENES AND OTHER VOLATILE AROMATIC COMPOUNDS

- 306 Raynor, M.W., Roberts, S.L. and Mulholland, D.A.: Capillary supercritical fluid chromatography of limonoids. *J. High Resolut. Chromatogr.*, 16 (1993) 469-472.

#### 15a. Terpenes

- 307 Hasler, A. and Meier, B.: Determination of terpenes in *Ginkgo biloba* L. by capillary gas chromatography. *Pharm. Pharmacol. Lett.*, 2 (1992) 187-190; *C.A.*, 118 (1993) 219941w.
- 308 Humphrey, A.M., Harris, J.R., Mansfield, C.A., Michalkowicz, D.M., Milchard, M., Moyler, D.A., Osbiston, A., Smith, S., Starr, B., Stevens, T.M. and Wilson, J.J.: Application of gas-liquid chromatography to the analysis of essential oils. Part XVI. Monographs for five essential oils. *Analyst (Cambridge)*, 118 (1993) 1089-1098.
- 309 Kanamori, H., Terauchi, M., Fuse, J. and Sakamoto, I.: (Studies on the evaluation of *Chamomillae* Flos. (Part 1). Simultaneous and quantitative analysis of fat-soluble compounds.) *Shoyakugaku Zasshi*, 46 (1992) 384-388; *C.A.*, 118 (1993) 261098b.
- 310 Kaniyas, G.D., Tsiatsa, E., Loukis, A. and Kilikoglou, V.: Determination and statistical analysis of trace element and active constituent concentrations in the medicinal plant *Eucalyptus camaldulensis* DehnH (*E. rostratus* Schlecht). *J. Radioanal. Nucl. Chem.*, 169 (1993) 483-491; *C.A.*, 118 (1993) 261140j.
- 311 Przyborski, H. and Bandion, F.: (Determination of  $\beta$ -asarone, pulegone, safrone, santonin, and thujone in spirits and wines.) *Mitt. Klosterneuburg*, 42 (1992) 171-178; *C.A.*, 119 (1993) 7419c.

See also 230, 312, 470.

#### 15b. Essential oils

- 312 Dugo, G., Stagno D'Alcontres, I., Donato, M.G. and Dugo, P.: On the genuineness of citrus essential oils. Part XXXVI. Detection of added reconstituted lemon oil in genuine cold-pressed lemon essential oil by high-resolution gas chromatography with chiral capillary columns. *J. Essent. Oil Res.*, 5 (1993) 21-26; *C.A.*, 118 (1993) 197759p.
- 313 Fang, H., Duan, H., Xu, Y., Zhou, T. and Lin, J.: (Studies on the chemical components of essential oil of *Elsholtzia blanda*.) *Sepu*, 11 (1993) 69-71.
- 314 Gavini, P., Angelini, E. and Fellous, R.: (Standardized chromatographic profiles. Automatic comparisons.) *Parfums, Cosmet., Aromes*, 101 (1991) 70-74; *C.A.*, 118 (1993) 240439y.
- 315 Guo, Y., Dai, L., Yang, L. and Ren, Q.: (Study on the stability of the quality of flower resource during the blossom period for the *Jasminum sambac* (*L.*) *aiton* in Fuzhou region (I) - analysis of the chemical components of essential oil.) *Sepu*, 11 (1993) 191-196.
- 316 Jing, Z., Wang, M. and Chen, S.: (Study on chemical constituents of essential oil of *Ledum palustre* L. var. *augustum* by gas chromatography/mass spectrometry (GC/MS).) *Sepu*, 11 (1993) 244-245.
- 317 Leach, D.N., Wyllie, S.G., Hall, J.G. and Kyrtatzis, I.: Enantiomeric composition of the principal components of the oil of *Melaleuca alternifolia* J. Agric. Food Chem., 41 (1993) 1627-1632.

- 318 Liu, J., Zhang, S. and Wei, L.: (Analysis and comparison of the volatile oil from *Daphne genkwa* Sied. et Zucc. before and after processing by GC-MS.) *Zhongguo Zhongyao Zazhi*, 18 (1993) 25-26; C.A., 118 (1993) 251446z.
- 319 Ma, Y., Hong, T., Chen, D. and Sun, S.: (The analysis of essential oil of *Krylovia limonifolia* (Less.) *schischk* by a gas chromatography-gas chromatography-mass spectroscopy (GC-GC-MS) system.) *Sepu*, 11 (1993) 148-150.
- 320 Maas, B., Dietrich, A. and Mosandl, A.: Enantioselective capillary gas chromatography-olfactometry in essential oil analysis. *Naturwissenschaften*, 80 (1993) 470-473.
- 321 Stashenko, E., Martinez, J.R., Macku, C. and Shibamoto, T.: HRGC and GC-MS analysis of essential oil from colombian Ylang-Ylang (*Cananga odorata* Hook Fil. et Thompson, *forma genuina*). *J. High Resolut. Chromatogr.*, 16 (1993) 441-444.

See also 88, 308, 480.

## 16. NITRO AND NITROSO COMPOUNDS

- 322 Pressmar, F., Neidlein, R., Strein, K. and Bartsch, W.: Pharmacokinetics of the organic nitrates *trans*-2-amino-2-methyl-N-(4-nitroxycyclohexylmethyl)-propionamide in dogs, and of 4-(2-nitroxyethyl)-piperidine in dogs and in monkeys. *Arzneim.-Forsch.*, 42 (1992) 1186-1191; C.A., 118 (1993) 246822q.
- 323 Slack, G.C., McNair, H.M., Hawthorne, S.B. and Miller, D.J.: Coupled solid phase extraction-supercritical fluid extraction-on-line gas chromatography of explosives from water. *J. High Resolut. Chromatogr.*, 16 (1993) 473-478.

See also 37, 109, 503, 506.

## 17. AMINES, AMIDES AND RELATED NITROGEN COMPOUNDS

### 17a. Amines and polyamines

- 324 Bartels, M.J., Timchalk, C. and Smith, F.A.: Gas chromatographic/tandem mass spectrometric identification and quantitation of metabolic 4-acetyltoluene-2,4-diamine from the F344 rat. *Biol. Mass Spectrom.*, 22 (1993) 194-200; C.A., 119 (1993) 2538f.
- 325 Guan, F., Liu, Y., Luo, Y., Liu, F., Hu, X., Li, Q. and Kang, Z.: (Identification of tetramine in intoxication samples by gas chromatography coupled with mass spectrometry and evaluation of artificial carbonic kidney for the removal of tetramine from poisoned persons.) *Fenxi Huaxue*, 21 (1993) 452-454; C.A., 118 (1993) 249401n.
- 326 Lundh, T. and Åkesson, B.: Gas chromatographic determination of primary and secondary low-molecular-mass aliphatic amines in urine using derivatization with isobutyl chloroformate. *J. Chromatogr.*, 617 (1993) 191-196.
- 327 Scheepers, P.T.J., Velders, D.D., Straetemans, M.M.E., Ouwkerk, J.C., van Vliet, L.A. and Bos, R.P.: Solid-phase extraction and gas chromatographic-mass spectrometric determination after hydrolysis of 2-aminofluorene haemoglobin adducts in blood of rats. *J. Chromatogr.*, 619 (1993) 215-221.

- 328 Straub, R.F., Voyksner, R.D. and Keever, J.T.: Determination of aromatic amines originating from azo dyes by hydrogen-palladium reduction combined with gas chromatography/mass spectrometry. *Anal. Chem.*, 65 (1993) 2131-2136.
- 329 Wu, L. and Cheng, Y.: (Gas chromatographic analytical determination of aniline in workshop air.) *Zhonghua Yufang Yixue Zazhi*, 26 (1992) 297-299; C.A., 119 (1993) 14294m.

See also 332.

### 17b. Catecholamines and their metabolites

- 330 Kema, I.P., Meiborg, G., Nagel, G.T., Stob, G.J. and Muskiet, F.A.J.: Isotope dilution ammonia chemical ionization mass fragmentographic analysis of urinary 3-O-methylated catecholamine metabolites. Rapid sample clean-up and extraction of lyophilized samples. *J. Chromatogr.*, 617 (1993) 181-189.

### 17d. Other amine derivatives and amides (excl. peptides)

- 331 Huang, Z. and Nie, H.: (Identification of brominated chlordimeform by gas chromatography and gas chromatography/mass spectroscopy.) *Sepu*, 11 (1993) 97-98.
- 332 Kumar, A. and Panwar, A.: Gas chromatographic separation of isomeric aminophenols, aniline, phenol, benzoquinone and azobenzene on an HP-1 capillary column. *Mikrochim. Acta*, 111 (1993) 177-182.
- 333 Raymer, J.H., Sparacino, C.M., Velez, G.R., Padilla, S., MacPhail, R.C. and Crofton, K.M.: Determination of acrylamide in rat serum and sciatic nerve by gas chromatography-electron-capture detection. *J. Chromatogr.*, 619 (1993) 223-234.
- 334 Taylor, W.G., Vedres, D.D. and Elder, J.L.: Capillary gas chromatographic separation of some diastereomeric amides from carbonyldiimidazole-mediated microgram-scale derivatizations of the acid moiety of permethrin insecticide. *J. Chromatogr.*, 645 (1993) 303-310.
- 335 Yang, D., Beylot, M., Agarwal, K.C., Soloviev, M.V. and Brunnergruber, H.: Assay of the human liver citric acid probe phenylacetylglutamine and of phenylacetate in plasma by gas chromatography-mass spectrometry. *Anal. Biochem.*, 212 (1993) 277-282.

See also 145, 478.

## 18. AMINO ACIDS AND PEPTIDES; CHEMICAL STRUCTURE OF PROTEINS

### 18a. Amino acids and their derivatives

- 336 Beylot, M., David, F. and Brunnengraber, H.: Determination of the <sup>13</sup>C-labelling pattern of glutamate by gas chromatography-mass spectrometry. *Anal. Biochem.*, 212 (1993) 532-536.
- 337 Domergue, N., Pugnieri, M. and Previero, A.: One-step conversion of amino acids into N-methyloxycarbonyl alkyl ester derivatives for chiral gas chromatography. *Anal. Biochem.*, 214 (1993) 420-425.
- 338 Fan, T.W.-M., Colmer, T.D., Lane, A.N. and Higashi, R.M.: Determination of metabolites by <sup>1</sup>H NMR and GC: Analysis for organic osmolytes in crude tissue extracts. *Anal. Biochem.*, 214 (1993) 260-271.

- 339 Nang, J., Cai, D. and Ouyang, Z.: (Determination of bound sulfur-containing amino acids by indirect gas chromatography.) *Sepu*, 11 (1993) 93-95.
- 340 Stämpfli, A.A., Ballèvre, O. and Fay, L.B.: Determination of taurine metabolism by measurement of <sup>15</sup>N-enriched taurine in cat urine by gas chromatography-mass spectrometry. *J. Chromatogr.*, 617 (1993) 197-203.
- See also 160.
21. PURINES, PYRIMIDINES, NUCLEIC ACIDS AND THEIR CONSTITUENTS
- 21a. *Purines, pyrimidines, nucleosides, nucleotides*
- 341 Chiu, C.S., Saha, M., Abushamas, A. and Giese, R.W.: Chemical transformation/derivation of O<sup>6</sup>-methyl- and O<sup>6</sup>-(hydroxyethyl)guanine for detection by GC-EC/MS. *Anal. Chem.*, 65 (1993) 3071-3875.
- 342 Prevost, V., Shuker, D.E.G., Friesen, M.D., Eberle, G., Rajewsky, M.F. and Bartsch, H.: Immunoaffinity purification and gas chromatography-mass spectrometric quantification of 3-alkyladenines in urine: Metabolism studies and basal excretion levels in man. *Carcinogenesis (London)*, 14 (1993) 199-204; *C.A.*, 118 (1993) 185580k.
- 343 Teixeira, A.J.R., Gommers-Ampt, J.H., van de Werken, G., Westra, J.G., Stavenuiter, J.F.C. and de Jong, A.P.J.M.: Method for the analysis of oxidized nucleosides by gas chromatography/mass spectrometry. *Anal. Biochem.*, 214 (1993) 474-483.
22. ALKALOIDS
- 344 Glass, R.L. and Johnson, E.L.: Comparison of high performance liquid chromatographic and gas chromatographic analyses of cocaine in coca leaves. *J. Liq. Chromatogr.*, 16 (1993) 3543-3555.
- 345 Jacob, P., III, Yu, L., Liang, G., Shulglin, A.T. and Benowitz, N.L.: Gas chromatographic-mass spectrometric method for determination of anabasine, anapipine and other tobacco alkaloids in urine of smokers and smokeless tobacco users. *J. Chromatogr.*, 619 (1993) 49-61.
- 346 Moore, J.M., Cooper, D.A., Kram, T.C. and Klein, R.F.X.: Sensitive detection of nicotine after its novel perfluoroacylation and analysis using capillary gas chromatography-electron-capture detection. *J. Chromatogr.*, 645 (1993) 273-281.
- 347 Pirozhkov, S.V., Watson, R.R. and Eskelson, C.D.: Gas chromatographic detection of cocaine and cocaethylene in hair of mice chronically injected with cocaine or cocaethylene and fed ethanol. *Forensic Sci. Int.*, 57 (1992) 99-107; *C.A.*, 118 (1993) 207050s.
- See also 459.
23. OTHER SUBSTANCES CONTAINING HETEROCYCLIC NITROGEN
- 23a. *Porphyryns and other pyrroles*
- 348 Xu, Z.: (Pyrolysis-gas chromatography of tetraphenylporphyrin and its metallic complexes.) *Sepu*, 11 (1993) 225-227.
- 23c. *Indole derivatives and plant hormones (gibberelins)*
- 349 Shen, Z., Zhou, Y. and Jiang, H.: (High pressure liquid chromatography and gas chromatography of gibberellic acid.) *Zhiwu Shenglixue Tongxun*, 28 (1992) 290-293; *C.A.*, 118 (1993) 208716a.
- 23d. *Pyridine derivatives*
- 350 Fischer, C., Schoenberger, F., Muck, W., Heuck, K. and Eichelbaum, M.: Simultaneous assessment of the intravenous and oral disposition of the enantiomers of racemic nimodipine by chiral stationary-phase high-performance liquid chromatography and gas chromatography/mass spectroscopy combined with a stable isotope technique. *J. Pharm. Sci.*, 82 (1993) 244-250; *C.A.*, 118 (1993) 224793d.
- 351 Nabivach, V.M.: (Correlation of structure and sorption of heterocyclic nitrogen compounds.) *Zh. Fiz. Khim.*, 67 (1993) 821-826.
- See also 19, 478.
- 23e. *Other N-heterocyclic compounds*
- 352 Musshoff, F., Daldrup, T. and Bonte, W.: Gas chromatographic-mass spectrometric screening procedure for the identification of formaldehyde-derived tetraisoquinolines in human urine. *J. Chromatogr.*, 619 (1993) 235-242.
- See also 243, 351, 432, 478.
24. ORGANIC SULPHUR COMPOUNDS (INCL. GLUCOSINOLATES)
- 353 Boccacci Mariani, M., Vinci, G. and Milana, M.R.: Quantitative determination of methyl isothiocyanate in wines by static headspace gas chromatography. *Ann. Chim. (Rome)*, 82 (1992) 589-586; *C.A.*, 118 (1993) 211529x.
- 354 D'Agostino, P.A. and Provost, L.R.: Capillary column electron impact and ammonia chemical ionization gas chromatographic-mass spectrometric and gas chromatographic-tandem mass spectrometric analysis of mustard hydrolysis products. *J. Chromatogr.*, 645 (1993) 283-292.
- 355 Goldstein, H., Rader, S. and Murakami, A.A.: Determination of 3-methyl-2-butene-1-thiol in beer. *J. Am. Soc. Brew. Chem.*, 51, No. 2 (1993) 70-74; *C.A.*, 119 (1993) 7452h.
- 356 Lavigne, V., Boidron, J.N. and Dubourdieu, D.: (Determination of low-molecular-weight volatile sulfur compounds in wine by gas chromatography with flame photometric detection.) *J. Int. Sci. Vigne Vin*, 27 (1993) 1-12; *C.A.*, 119 (1993) 7441d.
- 357 Miki, Y., Sugimoto, Y. and Yamadaya, S.: (Synthesis and GC-MS analysis of alkylthiophenes.) *Nippon Kagaku Kaishi*, No. 1 (1993) 79-85; *C.A.*, 118 (1993) 254661b.

- 358 Slater, G.P. and Manville, J.F.: Analysis of thiocyanates and isothiocyanates by ammonia chemical ionization gas chromatography—mass spectrometry and gas chromatography—Fourier transform infrared spectroscopy. *J. Chromatogr.*, 648 (1993) 433-443.
- 359 Terada, H., Nagai, Y. and Miyabe, M.: (Determination of methyl isothiocyanate in wine by using solid phase extraction/HPLC, GC, and GC/mass spectrometry.) *Shokuhin Eiseigaku Zasshi*, 33 (1992) 586-592; *C.A.*, 119 (1993) 7424a.
- 360 Uchiyama, S., Takeda, H., Kobayashi, A., Ito, S., Sakurai, H., Tada, Y., Aoki, G., Hosogai, T., Yamanaka, T. et al.: (Determination of methyl isothiocyanate in wine by GC and GC/mass spectrometry.) *Shokuhin Eiseigaku Zasshi*, 33 (1992) 603-608; *C.A.*, 119 (1993) 7423z.
- See also 339, 520, 559.
25. ORGANIC PHOSPHORUS COMPOUNDS (INCL. SUGAR PHOSPHATES)
- 361 Efer, J., Müller, S., Engewald, W., Knobloch, T. and Levsen, K.: Indirect GC determination of Ethepon in drinking water by a combination of reactive headspace sampling with adsorptive enrichment/thermal desorption. *Chromatographia*, 37 (1993) 361-364.
- 362 Gilyarov, V.A., Shcherbina, T.M., Laretina, A.P. and Kabachnik, M.I.: (GC-MS investigation of imide-amide rearrangement.) *Izv. Akad. Nauk, Ser. Khim.*, (1992) 2452-2455; *C.A.*, 118 (1993) 234150f.
- 363 Stan'kov, I.N., Beresnev, A.N., Agureev, V.G. and Lapin, S.N.: (Relation between the structure and characteristic retention of phosphoorganic compounds on stationary phase SE-30.) *Zh. Fiz. Khim.*, 66 (1992) 3321-3328.
- See also 543.
26. ORGANOMETALLIC AND RELATED COMPOUNDS
- 26a. Organometallic compounds
- 364 Diaz Ara, R.C.: (Gas-chromatographic determination of mercury after derivatization to diphenylmercury.) *Rev. Acad. Cienc. Exactas, Fis., Quim. Nat. Zaragoza*, 47 (1992) 191-199; *C.A.*, 118 (1993) 190220x.
- 365 Erteborg, H., Baxter, D.C. and Frech, W.: Speciation of mercury in natural waters by capillary gas chromatography with a microwave-induced plasma emission detector following preconcentration using a dithiocarbamate resin microcolumn installed in a closed flow injection system. *Analyst (Cambridge)*, 118 (1993) 1007-1013.
- 366 Forsyth, D.S., Weber, D. and Barlow, L.: The determination of organotin compounds in fruit juices using gas chromatography—atomic absorption spectrometry. *Appl. Organomet. Chem.*, 6 (1992) 579-585; *C.A.*, 118 (1993) 253467f.
- 367 Forsyth, D.S., Weber, D. and Dalgligh, K.: The determination of organotin compounds in edible oils by gas chromatography—atomic absorption spectrometry. *Talanta*, 40 (1993) 299-305.
- 368 Kawata, K., Minagawa, M., Fujieda, Y. and Yasuhara, A.: Sampling method of organotin compounds in air using a quartz-fibre filter and an activated carbon-fibre filter for gas chromatographic determination. *J. Chromatogr. A*, 653 (1993) 369-373.
- 369 Krylov, V.A., Krasotskii, S.G., Sarkisov, A.V. and Chernova, O.Yu.: (Analysis of high-purity diethyltellurium by capillary gas chromatography.) *Vysokochist. Veshchestva*, (1993) 133-141; *C.A.*, 118 (1993) 246592q.
- 370 Kupchik, E.J.: Structure—gas chromatographic retention time models of tetra-*n*-alkylsilanes and tetra-*n*-alkylgermanes using topological indexes. A correction. *J. Chromatogr.*, 645 (1993) 182-184.
- 371 Lobinski, R., Boutron, C.F., Candelone, J.-P., Hong, S., Szpunar-Lobinska, J. and Adams, F.C.: Speciation analysis of organolead compounds in Greenland snow at the femtogram-per-gram level by capillary gas chromatography/atomic emission spectrometry. *Anal. Chem.*, 65 (1993) 2510-2515.
- 372 Nagase, M. and Hasebe, K.: Determination of tributyltin and triphenyltin compounds in fish by gas chromatography with flame photometric detection. *Anal. Sci.*, 9 (1993) 517-522.
- 373 Ohnogi, H., Korenaga, T., Tanaka, M. and Shinoda, S.: (Determination of butyltin compounds by gas chromatography with flame photometric detection. Application to survey of water pollution by organic tin compounds.) *Kankyo Saigyo*, 14 (1992) 28-34; *C.A.*, 118 (1993) 197658e.
- 374 Quevauviller, P., Martin, F., Belin, C. and Donard, O.F.X.: Matrix effects in the determination of butyltin compounds in environmental samples by GC AA after hydride generation. *Appl. Organomet. Chem.*, 7 (1993) 149-157; *C.A.*, 118 (1993) 265822f.
- 375 Ståb, J.A., Cofino, W.P., van Hattum, B. and Brinkman, U.A.T.: Comparison of GC/MSD and GC/AED for the determination of organotin compounds in the environment. *Fresenius J. Anal. Chem.*, 347 (1993) 247-255.
- 376 Zhao, R., Sun, A. and Lü, Y.: (Determination of ferrocene solution by gas capillary chromatography.) *Sepu*, 11 (1993) 257.
- 26b. Boranes, silanes and related non-metallic compounds
- 377 Huang, B.: (Evidence for selenourea in human urine.) *Sepu*, 11 (1993) 221-222.
- 378 Li, S. and Li, X.: (Determination of trimethoxy(3-oxiranylmethoxypropyl)silane by gas chromatography.) *Fenxi Huaxue*, 21 (1993) 491; *C.A.*, 118 (1993) 246612w.
- 26c. Coordination compounds
- 379 Aggarwal, S.K., Kinter, M., Wills, M.R., Savory, J. and Herold, D.A.: Analysis of nickel in biological materials by isotope dilution gas chromatography/mass spectrometry. *Adv. Environ. Sci. Technol.*, 25 (1992) 77-87; *C.A.*, 119 (1993) 2541b.
- 380 Liu, Y., Lopez-Avila, V., Alcaraz, M., Beckert, W.F. and Heithmar, E.M.: Determination of metals in solid samples by complexation—supercritical fluid extraction and gas chromatography—atomic emission detection. *J. Chromatogr. Sci.*, 31 (1993) 310-316.
- See also 348.

## 27. VITAMINS AND VARIOUS ANIMAL GROWTH FACTORS (NON-PEPTIDIC)

- 381 Micali, G., Lanuzza, F. and Currò, P.: Analysis of tocopherols in margarine by on-line HPLC-HRGC coupling. *J. High Resolut. Chromatogr.*, 16 (1993) 536-538.
- 382 Snyder, J.M., Taylor, S.L. and King, J.W.: Analysis of tocopherols by capillary supercritical fluid chromatography and mass spectrometry. *J. Am. Oil Chem. Soc.*, 70 (1993) 349-354.

## 28. ANTIBIOTICS

- 383 Sokol, J. and Matisová, E.: (Chromatographic methods for residue determination of commonly used antibiotics.) *Chem. Listy*, 87 (1993) 786-794 - a review with 136 refs.
- 384 Zhang, Y., Agarwal, K.C., Beylot, M., Soloviev, M.V., David, F., Reider, M., Tserng, K.-Y. and Brunnengruber, H.: Assay of the acetyl-CoA probe acetyl-sulfomethoxazole and of sulfomethoxazole by gas chromatography-mass spectrometry. *Anal. Biochem.*, 212 (1993) 481-486.

## 29. INSECTICIDES, PESTICIDES AND OTHER AGROCHEMICALS

## 29a. General techniques

- 385 Atsuta, M. and Daigo, I.: (A study of automated screening for pesticides by differential polarity dual capillary column GC and multidetector retention index method. (DPDC-MDR)) *Kuromatogurafi*, 13 (1992) 303-304; *C.A.*, 118 (1993) 249723u.
- 386 Ceulemans, M., Lobinski, R., Dirx, W.M.R. and Adams, F.C.: Rapid sensitive speciation analysis of butyl- and phenyltin compounds in water by capillary gas chromatography-atomic emission spectrometry (GC-AES) after *in-situ* ethylation and in-line preconcentration. *Fresenius J. Anal. Chem.*, 347 (1993) 256-262.
- 387 Jin, R.: (Determination of pesticide quality and authenticity by gas chromatography.) *Faming Zhuanli Shenqing Gongkai Shuomingshu* CN 1,067,316 (Cl. G01N30/02), 23 Dec. 1992, Appl. 91,106,435, 28 May 1991; 10 pp.; *C.A.*, 119 (1993) 3053f.
- 388 Kondoh, H. and Murata, K.: (RP-C18 cartridge isolation of 17 controlled pesticides in wastewater.) *Hokkaido Kankyo Kagaku Kenkyu Senta Shoho*, 19 (1992) 53-59; *C.A.*, 118 (1993) 197659f.
- 389 Onji, Y., Uno, M. and Sasaki, M.: (Analysis of airborne pesticides by multidimensional GC/MS with large-volume sample injection.) *Kuromatogurafi*, 13 (1992) 301-302; *C.A.*, 118 (1993) 239768y.
- 390 Stan, H.-J. and Heberer, T.: Automated recognition of target compounds at low levels in environmental samples by means of capillary gas chromatography-mass spectrometry with dedicated mass spectral libraries and the macro program AUTARG. II. Application to pesticides in ground water samples. *J. Chromatogr. A*, 653 (1993) 55-62.
- 391 Stan, H.-J. and Linkerhägner, M.: Capillary gas chromatography-atomic emission detection: a useful instrumental method in pesticide residue analysis of plant foodstuffs. *J. High Resolut. Chromatogr.*, 16 (1993) 539-548.

See also 64, 157, 509, 513.

## 29b. Chlorinated insecticides

- 392 Barcarolo, R. and Tutta, C.: (Determination of organochlorine pesticides in dairy products using LC-GC.) *Riv. Soc. Ital. Sci. Aliment.*, 21 (1992) 141-150; *C.A.*, 118 (1993) 211499n.
- 393 Jablonska, A., Hansen, M., Ekeberg, D. and Lundanes, E.: Determination of chlorinated pesticides by capillary supercritical fluid chromatography-mass spectrometry with positive- and negative-ion detection. *J. Chromatogr.*, 647 (1993) 341-350.
- 394 Kuwabara, K., Matsumoto, H., Murakami, Y., Nishimune, T. and Sueki, K.: (Simultaneous determination of chlordanes, DDTs and other organochlorine pesticides in food in preparation for quantitative capillary gas chromatography.) *Osaka-furitsu Koshu Eisei Kenkyusho Kenkyu Hokoku, Shokuhin Eisei Hen*, 23 (1992) 41-45; *C.A.*, 118 (1993) 190224b.
- 395 Murakami, Y., Nishimune, T. and Sueki, K.: (Studies on environmental contaminants in food. V. Residues of oxadiazon and its metabolites in terrapin and corb shell processed foods.) *Shokuhin Eiseigaku Zasshi*, 33 (1992) 576-585; *C.A.*, 118 (1993) 232552q.
- 396 Murakami, Y., Nishimune, T., Sueki, K. and Tanaka, R.: (Studies on environmental contaminants in food. IV. Analysis of oxadiazon and its metabolites in blue mussel and pond snail.) *Shokuhin Eiseigaku Zasshi*, 33 (1992) 548-556; *C.A.*, 118 (1993) 232553r.
- 397 Neuzil, E.F. and Stone, D.J.: Detection of common chlorinated pesticides in water by gas chromatography. *J. Chem. Educ.*, 70 (1993) 167.

See also 83, 173, 205, 213, 400, 470, 474, 515, 516, 526.

## 29c. Phosphorus insecticides

- 398 Barcelo, D., Durand, G., de Bertrand, N. and Albaiges, J.: Determination of aquatic photodegradation products of selected pesticides by gas chromatography-mass spectrometry and liquid chromatography-mass spectrometry. *Sci. Total Environ.*, 132 (1993) 283-296; *C.A.*, 118 (1993) 240296z.
- 399 Bi, F., Wang, W., Huang, R. and Wei, Y.: (Gas chromatographic separation of the stereoisomers of 8 synthetic pyrethroids.) *Sepu*, 11 (1993) 158-160.
- 400 Erney, D.R. and Poole, C.F.: A study of single compound additives to minimize the matrix induced chromatographic response enhancement observed in the gas chromatography of pesticide residues. *J. High Resolut. Chromatogr.*, 16 (1993) 501-503.
- 401 Lacorte, S., Molina, C. and Barceló, D.: Screening of organophosphorus pesticides in environmental matrices by various gas chromatographic techniques. *Anal. Chim. Acta*, 281 (1993) 71-84.
- 402 Nishio, K. and Hayakawa, Y.: (Simultaneous analysis of organophosphorus pesticides in water by capillary gas chromatography.) *Mizu Kankyo Gakkaishi*, 16 (1993) 55-58; *C.A.*, 118 (1993) 240336n.
- 403 Nowak, J.: (Analysis of organophosphorus insecticides IPO-1391 and IPO-1401 soil residues by gas chromatography.) *Pestycydy (Warsaw)*, No. 2 (1992) 25-30; *C.A.*, 118 (1993) 207387g.

- 404 Yoshida, S., Moriguchi, Y., Konishi, Y., Taguchi, S. and Yakushi-ji, T.: (Analysis of organophosphorus termiticides in human milk.) *Jpn. J. Toxicol. Environ. Health*, 38 (1992) 52-56; *C.A.*, 118 (1993) 190189u.

See also 173, 406, 515, 516, 526.

#### 29d. Carbamates

- 405 Kokko, M.: Gas chromatographic screening for neostigmine and physostigmine using temperature-programmed retention indices. *J. Chromatogr.*, 648 (1993) 501-506.
- 406 Tonogai, Y., Tsumura, Y., Nakamura, Y. and Ito, Y.: (Sample preparation for simultaneous determination of 9 carbamate and 12 organophosphorus pesticides in agricultural products.) *Shokuhin Eiseigaku Zasshi*, 33 (1992) 449-457; *C.A.*, 118 (1993) 232437f.
- 407 Tsumura, Y., Tonogai, Y., Nakamura, Y., Miyata, M., Kamakura, K., Hashihata, N., Iwata, K., Ito, S., Minaba, K., et al.: (Analysis of pesticide residues in food by commercial kits and comparison with gas chromatography.) *Shokuhin Eiseigaku Zasshi*, 33 (1992) 458-466; *C.A.*, 118 (1993) 232436e.

#### 29e. Herbicides

- 408 Liu, R.: (Direct determination of naphthylacetic acid and phenoxyacetic acid pesticides by gas chromatography.) *Sepu*, 11 (1993) 111-112.
- 409 Nolte, J., Mayer, H., Khalifa, M.A. and Linscheid, M.: GC/MS of methylated phenoxyalkanoic acid herbicides. *Sci. Total Environ.*, 132 (1993) 141-146; *C.A.*, 118 (1993) 240356u.
- 410 Scott, C.: Determination of derivatized urea herbicides in water by solid-phase extraction, methylation and gas chromatography with nitrogen-phosphorus detector. *Analyst (Cambridge)*, 118 (1993) 1117-1122.
- 411 Tronczynski, J., Munsch, C., Durand, G. and Barcelo, D.: Monitoring of trace-levels of herbicides and their degradation products in the River Rhone, France, by gas chromatography-mass spectrometry. *Sci. Total Environ.*, 132 (1993) 327-337; *C.A.*, 118 (1993) 240297a.

See also 398, 512, 515, 519, 521.

#### 29f. Fungicides

- 412 Hajslova, J., Tahtah, W.H., Jehlickova, Z., Kocourek, V. and Cuhra, P.: Analysis of chlorophenoxy acids and other acidic contaminants in food crops. *Sci. Total Environ.*, 132 (1993) 259-274; *C.A.*, 118 (1993) 253530w.

See also 173, 203.

#### 29g. Other types of pesticides and various agrochemicals

- 413 Huang, Z., Xiong, F., Peng, S. and Nie, H.: (Rapid determination of procymidone residue in grape wine by gas chromatography with microchemical method.) *Sepu*, 11 (1993) 246-247.

See also 334, 516.

#### 30. SYNTHETIC AND NATURAL DYES

##### 30a. Synthetic dyes

See 533.

#### 31. PLASTICS AND THEIR INTERMEDIATES

- 414 Babenko, E.V., Kirpichnikov, P.A., Muratova, G.Y. and Manyurov, I.R.: (Use of mathematic modeling methods to calculate the strength characteristics of polymer composites.) *Zh. Prikl. Khim. (S.-Peterburg)*, 65 (1992) 2345-2349; *C.A.*, 119 (1993) 9693m.
- 415 Berg, B.E., Hegna, D.R., Orlien, N. and Greibrokk, T.: Determination of low levels of polymer additives migrating from polypropylene to food simulated liquids by capillary SFC and solvent venting injection. *Chromatographia*, 37 (1993) 271-276.
- 416 Cankurtaran, O. and Yilmaz, F.: The retention diagram of poly(*p*-chlorostyrene) and *tert*-butyl acetate by the inverse gas chromatography method. *Macromol. Rep.*, A30, Suppl. 1-2 (1993) 167-172; *C.A.*, 119 (1993) 9473q.
- 417 Evdokimova, S.P., Krylova, Yu.V., Beresnev, V.N. and Yatsenko, L.A.: (Determination of the composition of a five-component acrylate copolymer by pyrolytic gas chromatography.) *Kauch. Rezina*, No. 1 (1992) 16-18; *C.A.*, 118 (1993) 235661k.
- 418 Kojima, T., Takaku, H., Urata, Y. and Gotoh, K.: Pyrolysis gas chromatography-mass spectrometry studies on plasma-polymerized pyrrole. *J. Appl. Polym. Sci.*, 48 (1993) 1395-1398; *C.A.*, 118 (1993) 255668w.
- 419 Orav, A., Kuningas, K., Koplimets, E. and Rang, S.: Analysis of reaction by-products in ethylene oligomers by capillary gas chromatography. *Chromatographia*, 37 (1993) 259-263.
- 420 Romdhane, I.H. and Danner, R.P.: Polymer-solvent diffusion and equilibrium parameters by inverse gas-liquid chromatography. *AIChE J.*, 39 (1993) 625-635; *C.A.*, 118 (1993) 234887b.
- 421 Takatsu, M. and Yamamoto, T.: (Identification of soots from the combustion of various plastics using pyrolysis GC/MS.) *Bunseki Kagaku*, 42 (1993) 543-549.
- 422 Venema, A., van de Ven, H.J.F.M., David, F. and Sandra, P.: Supercritical fluid extraction of nylon 6: an investigation into the factors affecting the efficiency of extraction of caprolactam and its oligomers. *J. High Resolut. Chromatogr.*, 16 (1993) 522-524.
- 423 Webb, S.W.: Multicomponent inverse gas chromatography for analyses of sorption in polymers. *AIChE J.*, 39 (1993) 701-706; *C.A.*, 118 (1993) 234888c.
- 424 Wu, B., Yang, M., Liu, G., Pan, X. and Han, S.: (Study on chemical composition of combustion smoke from PS plastic.) *Sepu*, 11 (1993) 223-225.

See also 29, 207, 507.

## 32. DRUG ANALYSIS

## 32a. Drug analysis, general techniques

See 72, 227.

## 32b. Antirheumatics and antiinflammatory drugs

See 438.

## 32c. Autonomic and cardiovascular drugs

425 Foerster, H.J., Rominger, K.L., Ecker, E.M., Peil, H. and Wittrock, A.: Trace analysis of clenbuterol in biological fluids by automated capillary GC-MS. *Methodol. Surv. Biochem. Anal.*, 22 (1992) 231-232; C.A., 118 (1993) 182645z.

426 Sereda, K.M., Hardman, T.C., Dilloway, M.R. and Lant, A.F.: Development of a method for the detection of angiotensin converting enzyme inhibitors using electron capture-gas chromatography detection. *Anal. Proc.*, 30 (1993) 371-372.

427 Van Rhijn, J.A., Traag, W.A. and Heskamp, H.H.: Confirmatory analysis of clenbuterol using two different derivatives simultaneously. *J. Chromatogr.*, 619 (1993) 243-249.

428 Wang, Y., Zha, J., Liu, W., Liu, H., Gao, X. and Du, L.: (Determination of nitrendipine in rabbit plasma by gas chromatography.) *Hebei Yixueyuan Xuebao*, 13 (1992) 193-195; C.A., 118 (1993) 182663d.

See also 438.

## 32d. Central nervous system drugs

429 Cui, K., Zhou, Y., Zhang, C. and Chen, S.: (Determination of four ephedrine derivatives in urine by GC/MSD.) *Gaodeng Xuexiao Huaxue Xuebao*, 13 (1992) 1553-1554; C.A., 118 (1993) 246805m.

430 Fraser, A.D., Bryan, W. and Isner, A.F.: Urinary screening for  $\alpha$ -OH triazolam by FPIA and EIA with confirmation by GC/MS. *J. Anal. Toxicol.*, 16 (1992) 347-350; C.A., 118 (1993) 116100s.

431 Gaillard, Y., Gay-Montchamp, J.-P. and Ollagnier, M.: Gas chromatographic determination of zopiclone in plasma after solid-phase extraction. *J. Chromatogr.*, 619 (1993) 310-314.

432 Kauert, G., Herrle, I. and Wermeille, M.: Quantification of dimethindene in plasma by gas chromatography-mass fragmentography using ammonia chemical ionization. *J. Chromatogr.*, 617 (1993) 318-323.

433 Kojima, T., Taniguchi, T., Yashiki, M., Miyazaki, T., Iwasaki, Y. and Ohtani, M.: (A rapid method for detecting barbiturates in serum using EI-SIM.) *Nippon Iyo Masu Supokutoru Gakkai Koenshu*, 17 (1992) 203-206; C.A., 118 (1993) 246782b.

434 Martens, J. and Banditt, P.: Validation of the analysis of carbamazepine and its 10,11-epoxide metabolite by high-performance liquid chromatography from plasma: comparison with gas chromatography and the enzyme-multiplied immunoassay technique. *J. Chromatogr.*, 620 (1993) 169-173.

435 Park, K.H., Lee, M.H., Lee, M.G., Shim, C.K. and Park, J.S.: (Simultaneous determination of haloperidol and its metabolite, reduced haloperidol, in plasma by gas chromatography using nitrogen phosphorus selective detection.) *Yakche Hakhoechi*, 22 (1992) 197-204; C.A., 118 (1993) 204573d.

436 Rasanen, I., Ojanperä, I., Vuori, E. and Hase, T.A.: An homologous series of benzodiazepine retention index standards for gas chromatography. *J. High Resolut. Chromatogr.*, 16 (1993) 495-500.

437 Reimer, M.L.J., Mamer, O.A., Zavitsanos, A.P., Siddiqui, A.W. and Dadgar, D.: Determination of amphetamine, methamphetamine and desmethyldeprenyl in human plasma by gas chromatography/negative ion chemical ionization mass spectrometry. *Biol. Mass Spectrom.*, 22 (1993) 235-242; C.A., 118 (1993) 204624w.

438 Sane, R.T., Surve, S.R., Gangrade, M.G., Bapat, V.V. and Chonkar, N.L.: Simultaneous gas chromatographic determination of combined dosages, I) amitriptyline hydrochloride with chlor-diazepoxide, II) methocarbamol with ibuprofen, III) paracetamol with dichlofenac sodium and IV) paracetamol with promethazine hydrochloride. *Indian Drugs*, 30 (1993) 66-72; C.A., 118 (1993) 241066m.

439 Sano, T., Sato, K. and Katsumata, Y.: Simple deuterium labeling of methamphetamine and amphetamine, and its application to analysis of them in human blood by gas chromatography/chemical ionization-mass spectrometry. *Hochudoku*, 10 (1992) 247-255; C.A., 119 (1993) 2581q.

440 Sato, M. and Mitsui, T.: (Determination of the concentration of nitrazepam and diazepam with many fragment ions by multivariate analysis.) *Nippon Kagaku Kaishi*, (1993) 349-354; C.A., 118 (1993) 246609a.

441 Schmidt, N., Sittl, R., Brune, K. and Geisslinger, G.: Rapid determination of methadone in plasma, cerebrospinal fluid, and urine by gas chromatography and its application to routine drug monitoring. *Pharm. Res.*, 10 (1993) 441-444; C.A., 118 (1993) 182687q.

442 Schwartz, P.H. and Wasterlain, C.G.: Determination of serum and brain concentrations of neuroprotective and non-neuroprotective doses of MK-801. *J. Neurol. Sci.*, 115 (1993) 26-31; C.A., 118 (1993) 246806n.

443 Stetson, P.L., Domino, E.F. and Sneyd, J.R.: Determination of plasma propofol levels using gas chromatography-mass spectrometry with selected-ion monitoring. *J. Chromatogr.*, 620 (1993) 260-267.

444 West, R.E. and Ritz, D.P.: GC/MS analysis of five common benzodiazepine metabolites in urine as *tert*-butyldimethylsilyl derivatives. *J. Anal. Toxicol.*, 17 (1993) 114-116; C.A., 118 (1993) 204610p.

See also 454, 456, 463.

## 32e. Chemotherapeutics (exc. cytostatics and antibiotics)

445 Mooser, A.E. and Koch, H.: Confirmatory method for sulfonamide residues in animal tissues by gas chromatography and pulsed positive ion-negative ion-chemical ionization mass spectrometry. *J. Assoc. Off. Anal. Chem.*, 76 (1993) 976-982.

446 Tu, J., Xu, H., Sheng, L., Zhu, J., An, D. and Liu, G.: (Analysis of active ingredients in Wurenye.) *Zhongguo Zhongyao Zazhi*, 17 (1992) 538-540; C.A., 118 (1993) 198305z.

## 32f. Cytostatics

See 306.

32g. *Other drug categories*

- 447 Fang, X., Li, P., Tian, L. and An, D.: Classification of Chinese traditional drug Beimu (*Bulbus fritillariae*) by pyrolysis high resolution gas chromatography—pattern recognition. *J. Chin. Pharm. Sci.*, 1, No. 2 (1992) 65-72; C.A., 118 (1993) 219942x.
- 448 Ritter, W.: Determination of BAY u 3405, a novel thromboxane antagonist, in plasma and urine by HPLC and GC. *Methodol. Surv. Biochem. Anal.*, 22 (1992) 211-216; C.A., 118 (1993) 224773x.
- 449 Stanley, S.M.R., Wilhelmi, B.S. and Rodgers, J.P.: Comparison of immunoaffinity chromatography combined with gas chromatography—negative ion chemical ionisation mass spectrometry and radioimmunoassay for screening dexamethasone in equine urine. *J. Chromatogr.*, 620 (1993) 250-253.
- 450 Wu, Y., Chen, Q., Xu, Z., Liu, Z., Zhu, K. and Mao, Z.: (Exploration on identification of traditional Chinese medicinal materials by pyrolysis gas chromatography—numerical analysis.) *Sepu*, 11 (1993) 144-146.
- 451 Xu, Y.X., Shen, L. and Zhang, C.J.: (Analysis of dextromethorphan and its metabolites in human urine by using gas chromatography—mass spectrometry.) *Yaoxue Xuebao*, 28 (1993) 156-159; C.A., 118 (1993) 227687h.
- 452 Zhu, Y., Cheng, G. and Zhu, X.: (Quantitative determination of muscone in plasma by gas chromatography.) *Zhongguo Linchuang Yaoli Zazhi*, 8 (1992) 167-172; C.A., 118 (1993) 246802h.

See also 560.

32h. *Toxicological and forensic applications*

- 453 Brooks, K.E. and Smith, N.B.: Detection of acetonitrile and other polar solvents by capillary gas chromatography after direct injection of deproteinized serum. *J. Anal. Toxicol.*, 17 (1993) 127; C.A., 118 (1993) 249398s.
- 454 Coudore, F., Alazard, J.M., Paire, M., Andraud, G. and Lavarenne, J.: Rapid toxicological screening of barbiturates in plasma by wide-bore capillary gas chromatography and nitrogen-phosphorus detection. *J. Anal. Toxicol.*, 17 (1993) 109-113; C.A., 118 (1993) 227684e.
- 455 Gibb, R.P., Cockerham, H., Goldfogel, G.A., Lawson, G.M. and Raisys, V.: Substance abuse testing of urine by GC/MS in scanning mode evaluated by proficiency studies, TLC/GC, and EMIT. *J. Forensic Sci.*, 38 (1993) 124-133; C.A., 118 (1993) 141347k.
- 456 Gjerde, H., Hasvold, I., Pettersen, G. and Christophersen, A.S.: Determination of amphetamine and methamphetamine in blood by derivatization with perfluorooctanoyl chloride and gas chromatography/mass spectrometry. *J. Anal. Toxicol.*, 17 (1993) 65-68; C.A., 118 (1993) 227682c.
- 457 Goldberger, B.A., Darwin, W.D., Grant, T.M., Allen, A.C., Caplan, Y.H. and Cone, E.J.: Measurement of heroin and its metabolites by isotope-dilution electron-impact mass spectrometry. *Clin. Chem. (Washington)*, 39 (1993) 670-675.
- 458 Nakahara, Y., Kikura, R., Takahashi, K. and Konuma, K.: (GC/MS analysis of drugs and metabolites in hair for diagnosis of chronic drug abuse. On cocaine, heroin and amphetamines.) *Nippon Iyo Masu Supekutoru Gakkai Koenshu*, 17 (1992) 207-209; C.A., 118 (1993) 162590z.

- 459 Ohshita, T., Miyata, Y. and Ando, H.: (Application of capillary GC/MS for identification of drugs on forensic chemistry. II. Chemical confirmation of opium and opium poppy by simultaneous analyzing contained alkaloids and meconic acid.) *Shitsuryo Bunseki*, 40 (1992) 293-300; C.A., 118 (1993) 185242b.

See also 2, 3, 137, 149, 258, 276, 347, 430, 436, 444, 461.

32i. *Plant extracts*

- 460 Loughrin, J.H., Hamilton-Kemp, T.R., Burton, H.R. and Andersen, R.A.: Effect of diurnal sampling on the headspace composition of detached *Nicotiana suaveolens* flowers. *Phytochemistry*, 32 (1993) 1417-1419; C.A., 118 (1993) 251505t.
- 461 Zhu, J., Li, Y. and Bao, X.: GC determination of active cannabinoids in cannabis of Chinese origin and survey of their contents. *J. Chin. Pharm. Sci.*, 1, No. 2 (1992) 73-80; C.A., 118 (1993) 219943y.

See also 306, 310.

## 33. CLINICO-CHEMICAL APPLICATIONS

33b. *Complex mixtures and profiling (single compounds by cross-reference only)*

- 462 Atkinson, D.A., Hill, H.H. and Shultz, T.D.: Quantification of mammalian lignans in biological fluids using gas chromatography with ion mobility detection. *J. Chromatogr.*, 617 (1993) 173-179.
- 463 Kajita, M., Niwa, T., Wada, Y., Takeda, N., Yoshizumi, H., Tate-matsu, A. and Nagatsu, T.: (Detection of N-methyltyramine in parkinsonian brain by GC/MS.) *Nippon Iyo Masu Supekutoru Gakkai Koenshu*, 17 (1992) 187-190; C.A., 118 (1993) 205342q.
- 464 Lehmann, W.D., Stephan, M., Fuerstenberger, G. and Marks, F.: Profiling of monohydroxylated fatty acids in normal, hyperplastic and neoplastic mouse epidermis by gas chromatography—mass spectrometry. *Dev. Oncol.*, 71 (1993) 405-408; C.A., 118 (1993) 187296w.
- 465 Nishimi, T., Aramaki, S., Morita, J., Inokuchi, T., Yoshida, I. and Yamashita, F.: (Chronological changing of urinary organic acids profiling in neonates using the SIM method by GC/MS.) *Nippon Iyo Masu Supekutoru Gakkai Koenshu*, 17 (1992) 231-234; C.A., 118 (1993) 230900c.
- 466 Windsor, B.L., Radulovic, L.L., Bockbrader, H.N. and Chang, T.: Measurement of CI-979 (a candidate drug for the treatment of age-related disorders of cognition) in human plasma by capillary gas chromatography with nitrogen-selective detection. *J. Chromatogr.*, 619 (1993) 315-318.

See also 146, 183, 273, 282, 284, 342, 442.



## 34. FOOD ANALYSIS

## 34b. Complex mixtures (single compounds by cross-reference only)

- 467 Chung, T.Y., Eiserich, J.P. and Shibamoto, T.: Volatile compounds identified in headspace samples of peanut oil heated under temperatures ranging from 50 to 200 °C. *J. Agric. Food Chem.*, 41 (1993) 1467-1470.
- 468 Craske, J.D.: Separation of instrumental and chemical errors in the analysis of oils by gas chromatography – a collaborative evaluation. *J. Am. Oil Chem. Soc.*, 70 (1993) 325-334.
- 469 Fellous, R., George, G. and Schippa, C.: (Vanillin: anomalies in the carbon-13 identification protocol.) *Parfums, Cosmet., Aromes*, 106 (1992) 95-98; C.A., 118 (1993) 240427t.
- 470 Franich, P.A., Jakobsson, E., Jensen, S., Kroese, H.W. and Kylin, H.: Development of non-destructive methods for the determination of air-borne pollutants in pine noodles: Identification of trace constituents in radiate pine epicuticular wax. *Fresenius J. Anal. Chem.*, 347 (1993) 337-343.
- 471 Jambunathan, R., Sridhar, R., Raghunath, K., Dwivedi, S.L. and Nigam, S.N.: Oil quality characteristics and headspace volatiles of newly released groundnut (*Arachis hypogaea* L.) cultivars. *J. Sci. Food Agric.*, 61 (1993) 23-30; C.A., 118 (1993) 232618r.
- 472 Molnar-Perl, I. and Morvai, M.: Rapid method for the simultaneous GC quantitation of acids and sugars in fruits and vegetables. *Food Addit. Contam.*, 9 (1992) 505-514; C.A., 118 (1993) 253455a - a review with 9 refs.
- 473 Ortiz, M.C., Saez, J.A. and Palacios, J.L.: Typification of alcoholic distillates by multivariate techniques using data from chromatographic analysis. *Analyst (Cambridge)*, 118 (1993) 801-805.
- 474 Rahman, M.S., Montanarella, L., Johansson, B. and Larsen, B.R.: Trace levels of tris(4-chlorophenyl)-methanol and -methane in human milk. *Chemosphere*, 27 (1993) 1487-1497.
- 475 Righetti, B. and Facini, O.: Headspace gas composition in four *Prunus avium* cultivars with differing photosynthetic capabilities. *In Vitro Cell. Dev. Biol.: Plant*, 28P (1992) 179-182; C.A., 118 (1993) 143462t.
- 476 Rocchelli, V. and Baronciani, M.: (Volatile compounds in food packaging.) *Lab. 2000*, 6, No. 7 (1992) 68-72; C.A., 119 (1993) 10009z.
- 477 Sanchez-Muniz, F.J., Cuesta, C. and Garrido-Polonio, C.: Sunflower oil used for frying: combination of column, gas and high-performance size-exclusion chromatography for its evaluation. *J. Am. Oil Chem. Soc.*, 70 (1993) 235-240.
- 478 Terenina, M.B., Zhuravleva, I.L., Golovnya, R.V., Galustova, M.V., Solov'ev, G.V. and Enikeeva, N.G.: (Gas chromatographic profile of nitrogen-containing bases of white bread baked by a new technology using a lactic-acid bacterial preparation.) *Prikl. Biokhim. Mikrobiol.*, 28 (1992) 781-788; C.A., 118 (1993) 232687n.
- See also 203, 205, 208, 211, 217, 240, 246, 265, 271, 278, 297, 302, 304, 311, 353, 355, 356, 359, 360, 366, 367, 372, 381, 392, 394, 395, 406, 407, 412, 413, 485, 486, 493.
- 34c. Organoleptically important compounds (flavors, odors, volatiles)
- 479 Abbott, N., Etievant, P., Issanchou, S. and Langlois, D.: Critical evaluation of two commonly used techniques for the treatment of data from extract dilution sniffing analysis. *J. Agric. Food Chem.*, 41 (1993) 1698-1703.
- 480 Chung, T.Y., Eiserich, J.P. and Shibamoto, T.: Volatile compounds isolated from edible Korean chamchwi (*Aster scaber* Thumb). *J. Agric. Food Chem.*, 41 (1993) 1693-1697.
- 481 Collin, S., Osman, M., Delcambre, S., El-Zayat, A. and Dufour, J.-P.: Investigation of volatile flavor compounds in fresh and ripened Domiat cheeses. *J. Agric. Food Chem.*, 41 (1993) 1659-1663.
- 482 Egoif, L.M. and Jurs, P.C.: Quantitative structure-retention and structure-odor intensity relationships for a diverse group of odor-active compounds. *Anal. Chem.*, 65 (1993) 3119-3126.
- 483 Ferreira, V., Rapp, A., Cacho, J.F., Hastrich, H. and Yavas, I.: Fast and quantitative determination of wine flavor compounds using microextraction with Freon 113. *J. Agric. Food Chem.*, 41 (1993) 1413-1420.
- 484 Gomez, E., Ledbetter, C.A. and Hartsell, P.L.: Volatile compounds in apricot, plum, and their interspecific hybrids. *J. Agric. Food Chem.*, 41 (1993) 1669-1676.
- 485 Holscher, W. and Steinhart, H.: (Aroma changes during roasted coffee aging.) *Lebensmittelchemie*, 46 (1992) 106-111; C.A., 118 (1993) 232699t.
- 486 Hu, J. and Li, J.: (Gas chromatographic analysis of volatile organic acids in wine and fermented liquids.) *Sepu*, 11 (1993) 87-89.
- 487 Johnsen, P.B. and Lloyd, S.W.: Influence of fat content on uptake and depuration of the off-flavor 2-methylisoborneol by channel catfish (*Ictalurus punctatus*). *Can. J. Fish. Aquat. Sci.*, 49 (1992) 2406-2411; C.A., 118 (1993) 211614w.
- 488 Mills, O.E., Chung, S.-Y. and Johnsen, P.B.: Dehydration products of 2-methylisoborneol are not responsible for off-flavor in the channel catfish. *J. Agric. Food Chem.*, 41 (1993) 1690-1692.
- 489 Moio, L., Langlois, D., Etievant, P. and Addeo, F.: Powerful odorants in bovine, ovine, caprine and water buffalo milk determined by means of gas chromatography-olfactometry. *J. Dairy Res.*, 60 (1993) 215-222; C.A., 118 (1993) 253683y.
- 490 Ohsumi, C., Hayashi, T., Kubota, K. and Kobayashi, A.: Volatile flavor compounds formed in an interspecific hybrid between onion and garlic. *J. Agric. Food Chem.*, 41 (1993) 1808-1810.
- 491 Owuor, P.O.: Comparison of gas chromatographic volatile profiling methods for assessing the flavor quality of Kenyan black teas. *J. Sci. Food Agric.*, 59 (1992) 189-197; C.A., 118 (1993) 211669t.
- 492 Polesello, S., Lovati, F., Rizzolo, A. and Rovida, C.: Supercritical fluid extraction as a preparative tool for strawberry aroma analysis. *J. High Resolut. Chromatogr.*, 16 (1993) 555-559.
- 493 Sakamoto, K., Shimoda, M. and Osajima, Y.: (Concentration in Porapak Q column of volatile compounds in sake for analysis.) *Nippon Nogei Kagaku Kaishi*, 67 (1993) 685-692; C.A., 118 (1993) 232487x.
- 494 Shimoda, M.S., Shibamoto, T. and Noble, A.C.: Evaluation of headspace volatiles of Cabernet Sauvignon wines sampled by an on-column method. *J. Agric. Food Chem.*, 41 (1993) 1664-1668.

- 495 Stalcup, A.M., Ekborg, K.H., Gasper, M.P. and Armstrong, D.W.: Enantiomeric separation of chiral components reported to be in coffee, tea, or cocoa. *J. Agric. Food Chem.*, 41 (1993) 1684-1689.
- 496 Trevisan, A.: (Determination of volatile and semivolatile substances in wine by purge-and-trap and capillary gas chromatography.) *Vini Ital.*, 34, No. 5 (1992) 57-60; C.A., 118 (1993) 232465p.
- 497 Xiang, B.: (Use of gas chromatography in authentication of baijiu.) *Niangjiu*, No. 5 (1992) 26-32; C.A., 119 (1993) 7437g.

See also 460, 473, 475, 476.

### 35. ENVIRONMENTAL ANALYSIS

#### 35a. General papers and reviews

- 498 Douglas, G.S., McCarthy, K.J., Dahlen, D.T., Seavey, J.A., Steinhauer, W.G., Prince, R.C. and Elmendorf, D.L.: The use of hydrocarbon analyses for environmental assessment and remediation. In: Kostecki, P.T. and Calabrese, E.J. (Editors), *Contam. Soils*, Lewis, Boca Raton, 1992, pp. 1-21; C.A., 118 (1993) 204456t.
- 499 Merz, W.: (Analysis of environmental samples in chemical industry taking BASF Corp. as example.) *Zh. Anal. Khim.*, 48 (1993) 1296-1313.
- 500 Saiz-Jimenez, C.: Applications of pyrolysis-gas chromatography/mass spectrometry to the study of soils, plant materials and humic substances. A critical appraisal. *Dev. Agric. Managed-For. Ecol.*, 25 (1992) 27-38; C.A., 118 (1993) 204388x - a review with 58 refs.
- 501 Stan, H.-J. and Schwarzer, F.: Automated recognition of target compounds at low levels in environmental samples by means of capillary gas chromatography-mass spectrometry with dedicated mass spectral libraries and the macro program AUTARG. I. Description of the macro program AUTARG. *J. Chromatogr. A*, 653 (1993) 45-53.
- 502 Woolfenden, E.A., Broadway, G.M., Higham, P. and Seeley, I.: Alternative on-line and sorbent tube-based sampling procedures for 24-hr air pollution profiling. *Int. Lab.*, 23, No. 8 (1993) 29-36.

See also 147, 559.

#### 35b. Air pollution (complex mixtures; single compounds by cross-reference only)

- 503 Bertman, S.B., Buhr, M.P. and Roberts, J.M.: Automated cryogenic trapping techniques for capillary GC analysis of atmospheric trace compounds requiring no expandable cryogens: Application to the measurement of organic matter. *Anal. Chem.*, 65 (1993) 2944-2946.
- 504 Cao, X.-L. and Hewitt, C.N.: Passive sampling and gas chromatographic determination of low concentrations of reactive hydrocarbons in ambient air with reduction gas detector. *J. Chromatogr.*, 648 (1993) 191-197.
- 505 Fish, J.D., Huckins, J.N. and Zajicek, J.L.: Application of semipermeable membrane devices (SPMDs) as passive air sampler. *Chemosphere*, 27 (1993) 1609-1624.

- 506 Librando, V. and Fazzino, S.D.: Quantification of polycyclic aromatic hydrocarbons and their nitro derivatives in atmospheric particulate matter of Augusta city. *Chemosphere*, 27 (1993) 1649-1656.
- 507 Looock, F., Lampe, T. and Bahadir, M.: Examination of the emission of (semi)volatile organic compounds from polymer materials used as interior trim materials in automobiles. *Fresenius J. Anal. Chem.*, 347 (1993) 280-285.
- 508 Östermark, U. and Petersson, G.: Volatile hydrocarbons in exhaust from alkylate-based petrol. *Chemosphere*, 27 (1993) 1719-1728.
- 509 Roinestad, K.S., Louis, J.B. and Rosen, J.D.: Determination of pesticides in indoor air and dust. *J. Assoc. Off. Anal. Chem.*, 76 (1993) 1121-1126.
- 510 Soldatenkova, N.A. and Prokof'eva, V.G.: (Gas-chromatographic determination of methacrylonitrile in workplace air.) *Gig. Tr. Prof. Zabol.*, No. 1 (1992) 35-37; C.A., 118 (1993) 197000j.
- 511 Thulin, H.: Determination of nor-nitrogen mustard hydrochloride in air and on surfaces by using filter or wipe sampling and a derivatization-gas chromatographic procedure. *Am. Ind. Hyg. Assoc. J.*, 54 (1993) 76-81; C.A., 118 (1993) 196994z.

See also 187, 206, 213, 215, 251, 255, 329, 368, 389, 502, 530, 558.

#### 35c. Water pollution (complex mixtures; single compounds by cross-reference only)

- 512 Bulterman, A.-J., Vreuls, J.L., Ghijsen, R.T. and Brinkman, U.A.T.: Selective and sensitive detection of organic contaminants in water samples by on-line trace enrichment-gas chromatography-mass spectrometry. *J. High Resolut. Chromatogr.*, 16 (1993) 397-403.
- 513 Edgell, K.W., Erb, E.J., Wesselman, R.J. and Longbottom, J.E.: Gas chromatographic/electron capture detection method for determination of chlorinated acids in water: collaborative study. *J. Assoc. Off. Anal. Chem.*, 76 (1993) 1098-1112.
- 514 Eschke, H.D. and Traud, J.: (Analysis of sewage sludges and wastewaters by headspace gas chromatography with mass spectrometric detection.) *Vom Wasser*, 79 (1992) 269-280; C.A., 118 (1993) 197632s.
- 515 Foster, G.D., Gates, P.M., Foreman, W.T., McKenzie, S.W. and Rinella, F.A.: Determination of dissolved-phase pesticides in surface water from the Yakima River basin, Washington, using the goulden large-sample extractor and gas chromatography/mass spectrometry. *Environ. Sci. Technol.*, 27 (1993) 1911-1917.
- 516 Hernández, F., Morell, I., Beltrán, J. and López, F.J.: Multi-residue procedure for the analysis of pesticides in groundwater: application to samples from the Comunidad Valenciana, Spain. *Chromatographia*, 37 (1993) 303-312.
- 517 Janicki, W., Wolska, L., Wardencki, W. and Namiesnik, J.: Simple device for permeation removal of water vapour from purge gases in the determination of volatile organic compounds in aqueous samples. *J. Chromatogr. A*, 654 (1993) 279-285.
- 518 Linkiewicz, M.: Gas-chromatographic determination of the components in the synthesis of trimethylpropane. *Chem. Anal. (Warsaw)*, 37 (1992) 747-750.

- 519 Lintelmann, J., Günther, W.J., Rose, E. and Kettrup, A.: Behaviour of polycyclic aromatic hydrocarbons (PAHs) and triazine herbicides in water and aquifer material of a drinking water recharge plant. I. The area of investigation and the determination methods for PAHs and triazine herbicides in the aqueous matrix. *Fresenius J. Anal. Chem.*, 346 (1993) 988-994.
- 520 Longbottom, J.E., Edgell, K.W., Erb, E.J. and Lopez-Avila, V.: Gas chromatographic/nitrogen-phosphorus detection method for determination of ethylene thiourea in finished drinking waters: collaborative study. *J. Assoc. Off. Anal. Chem.*, 76 (1993) 1113-1120.
- 521 Readman, J.W., Kwong, L.L.W., Grondin, D., Bartocci, J., Ville-neuve, J.-P. and Mee, L.D.: Coastal water contamination from a triazine herbicide used in antifouling paints. *Environ. Sci. Technol.*, 27 (1993) 1940-1942.
- See also 49, 50, 57, 148, 151, 153, 155, 156, 192, 202, 205, 207, 212, 216, 222, 224, 232, 247, 283, 285, 323, 361, 365, 373, 375, 386, 388, 390, 397, 398, 401, 402, 411, 501.
- 35d. *Soil pollution (complex mixtures; single compounds by cross-reference only)*
- 522 Eiceman, G.A., Urquhart, N.S. and O'Connor, G.A.: Logistic and economic principles in gas chromatography-mass spectrometry use for plant uptake investigations. *J. Environ. Qual.*, 22 (1993) 167-173; *C.A.*, 118 (1993) 190659x.
- 523 Fukuda, M.: (Method and apparatus for microanalysis of volatile pollutants in soil.) *Jpn. Kokai Tokkyo Koho* JP 05 10,935 [93 10,935] (Cl. G01N30/04), 19 Jan. 1993, Appl. 91/184,026, 28 Jun. 1991; 5 pp.; *C.A.*, 118 (1993) 207379f.
- 524 Rastyanikov, E.G. and Drugov, Yu.S.: (Determination of volatile organic compounds in polluted soil by chromatography-mass spectrometry.) *Zh. Anal. Khim.*, 48 (1993) 1429-1434.
- 525 Rhodes, I.A.L., Olvera, R.Z. and Leon, J.A.: Determination of gasoline range total petroleum hydrocarbons (TPH) and approximate boiling point distribution in soil by gas chromatography. *Hydrocarbon Contam. Soils*, 1 (1991) 273-290; *C.A.*, 118 (1993) 190656u.
- 526 Wu, M., Liu, Z., Farnsworth, P.B. and Lee, M.L.: Comprehensive supercritical fluid extraction/gas chromatographic analysis of organic compounds in soil matrices with an element-selective radiofrequency plasma detector. *Anal. Chem.*, 65 (1993) 2185-2188.
- See also 194, 195, 197, 214, 221, 235, 248, 249, 501.
36. SOME TECHNICAL PRODUCTS AND COMPLEX MIXTURES
- 36a. *Surfactants*
- 527 Kane, M., Dean, J.R., Hitchen, S.M., Dowle, C.J. and Tranter, R.L.: Analysis of non-ionic surfactants using solid-phase extraction combined with supercritical fluid extraction and chromatography. *Anal. Proc.*, 30 (1993) 399-400.
- 528 Voelkel, A. and Janas, J.: Solubility parameters of broad and narrow distributed oxyethylates of fatty alcohols. *J. Chromatogr.*, 645 (1993) 141-151.
- 529 Voelkel, A., Janas, J. and Garcia-Dominguez, J.A.: Inverse gas chromatography in characterization of surfactants. Determination of binary parameter. *J. Chromatogr. A*, 654 (1993) 135-141.
- 36c. *Complex mixtures, technical products and unidentified compounds*
- 530 Arai, M.: (Determination of soluble organic fraction in particulates emitted from diesel engines by GC.) *Bunseki Kagaku*, 42 (1993) 593-597.
- 531 Aurela, B. and Räsänen, T.: Residual solvent content in heatset offset print. *J. High Resolut. Chromatogr.*, 16 (1993) 422-424.
- 532 Cai, Y.: (Pyrolysis-gas chromatography study on the type of kerogen and hydrocarbon generation from Member Shaihai 4 and 3 in Dongliang area.) *Diqiu Kexue*, 17 (1992) 411-419; *C.A.*, 118 (1993) 194741r.
- 533 Chiavari, G., Galletti, G.C., Lanterna, G. and Mazzeo, R.: The potential of pyrolysis-gas chromatography/mass spectrometry in the recognition of ancient painting media. *J. Anal. Appl. Pyrolysis*, 24 (1993) 227-242.
- 534 Fidalgo, M.L., Terron, M.C., Martinez, A.T., Gonzales, A.E., Gonzales-Vila, F.J. and Galletti, G.C.: Comparative study of fractions from alkaline extraction of wheat straw through chemical degradation, analytical pyrolysis, and spectroscopic techniques. *J. Agric. Food Chem.*, 41 (1993) 1621-1626.
- 535 Gao, X., Yang, J. and Hou, Y.: (Analysis of the composition of oxidation product of pseudocumene in vapor phase by GC/MS.) *Shiyou Huagong*, 21 (1992) 549-552; *C.A.*, 118 (1993) 182469v.
- 536 Giraldo, E.: Headspace monitoring and control of fixed-film anaerobic digesters. Avail. *Univ. Microfilms Int.*, Order No. DA9207403, 1991, 310 pp.; *C.A.*, 118 (1993) 153556r.
- 537 Greenwood, P.F., Zhang, E., Vastola, F.J. and Hatcher, P.G.: Laser micropyrolysis gas chromatography/mass spectrometry of coal. *Anal. Chem.*, 65 (1993) 1937-1946.
- 538 Havenga, W.J.: Rapid gas analyses for the investigation of spontaneous combustion using capillary gas chromatography. *Talanta*, 40 (1993) 213-219.
- 539 Huber, F.J.K., Kenndler, E., Reich, G., Hack, W. and Wolf, J.: Optimal selection of gas chromatographic columns for the analytical control of chemical warfare agents by application of information theory to retention data. *Anal. Chem.*, 65 (1993) 2903-2906.
- 540 Ishiwatari, M., Sakashita, H., Tatsumi, T. and Tominaga, H.: Thermal decomposition behavior of oil shale kerogens observed by stepwise pyrolysis-gas chromatography. *J. Anal. Appl. Pyrolysis*, 24 (1993) 273-290.
- 541 Kafka, Z. and Kura, M.: Characterization of organic compounds in industrial composts by gas chromatography-mass spectrometry. *Fresenius J. Anal. Chem.*, 346 (1993) 1000-1002.
- 542 Kobyakova, V.I. and Sokolov, O.A.: (Gas chromatographic analysis of volatile products of mycelial fungal biomass.) *Zh. Anal. Khim.*, 48 (1993) 1235-1241.
- 543 MacKay, G.A. and Smith, R.M.: Supercritical fluid extraction and chromatography-mass spectrometry of flame retardants from polyurethane foams. *Analyst (Cambridge)*, 118 (1993) 741-745.
- 544 Makavos, E.B. and Liu, C.C.: A method for creating predictable conditions in high temperature reactive gases. *High Temp. Mater. Processes*, 10 (1992) 127-134. *C.A.*, 118 (1993) 224484d.

- 545 Martin, L., Rubio, J. and Oteo, J.L.: (Characterization of porous glass by gas chromatography at zero coverage.) *Bol. Soc. Esp. Ceram. Vidrio*, 31 (1992) 233-237; *C.A.*, 119 (1993) 13747z.
- 546 Nikiforov, A.: Headspace constituents of shellac. *Z. Naturforsch., B: Chem. Sci.*, 48 (1993) 247-248; *C.A.*, 118 (1993) 193647c.
- 547 Olech, W. and Olejniczak, H.: (Evaluation of the technical state of transformers by gas chromatography. Evaluation criteria and research results.) *Energetyka*, 46 (1992) 385-388; *C.A.*, 119 (1993) 11722p.
- 548 Rastogi, S.C.: Gas chromatographic analysis of organic solvents mixtures on capillary columns of different polarity. *Chromatographia*, 37 (1993) 211-214.
- 549 Reeves, J.B., III. and Galletti, G.C.: Use of pyrolysis-gas chromatography/mass spectrometry in the study of lignin assays. *J. Anal. Appl. Pyrolysis*, 24 (1993) 243-255.
- 550 Tsai, C.H., Deo, M.D., Hanson, F.V. and Oblad, A.G.: Characterization and potential utilization of the Asphalt Ridge tar sand bitumen. I. Gas chromatography-mass spectrometry and pyrolysis-mass spectrometry analyses. *Fuel Sci. Technol. Int.*, 11 (1993) 475-506; *C.A.*, 118 (1993) 194817v.
- 551 Young, F.X.: Practical applications of gas chromatography in the paint and coatings industry. In: *Anal. Paints Relat. Mater.; ASTM Spec. Tech. Publ.*, Vol. STP 1119, ASTM, 1992, pp. 105-124; *C.A.*, 118 (1993) 214904b - a review with 22 refs.

See also 26, 90, 189, 193, 199, 200, 204, 210, 236, 250, 251, 252, 257, 259, 260, 286, 323, 424, 453, 507, 562.

### 37. CELLS, CELLULAR PARTICLES AND SUPRAMOLECULAR STRUCTURES

- 552 Smith, P.B. and Snyder, A.P.: Characterization of bacteria by oxidative and nonoxidative pyrolysis gas chromatography/ion-trap mass spectrometry. *J. Anal. Appl. Pyrolysis*, 24 (1993) 199-210.

See also 273.

### 38. INORGANIC COMPOUNDS

#### 38b. Anions

- 553 Haldimann, M. and Zimmerli, B.: Evaluation of ashing procedures for the gas chromatographic determination of fluoride in biological material. *Anal. Chim. Acta*, 282 (1993) 589-601.

#### 38c. Permanent and rare gases

- 554 Chulzhanova, M.G., Chulzhanov, Yu.A. and Popov, S.P.: (Gas chromatographic monitoring of the atmosphere of process boxes.) *Vysokochist. Veshchestva*, (1993) 158-161; *C.A.*, 118 (1993) 246478g.
- 555 Middleberg, R.A., Easterling, D.E., Zelonis, S.F., Rieders, F. and Rieders, M.F.: Estimation of perimortal percent carboxy-heme in nonstandard postmortem specimens using analysis of carbon monoxide by GC/MS and iron by flame atomic absorption spectrophotometry. *J. Anal. Toxicol.*, 17 (1993) 11-13; *C.A.*, 118 (1993) 95754r.
- 556 Takenouchi, S.: (Gas chromatographic analysis of fluid inclusions.) *Shigen Chishitsu*, 42 (1992) 289-299; *C.A.*, 118 (1993) 265609s.
- 557 Wang, S.: (Determination of trace argon gas in solid by vacuum melting gas chromatography.) *Gaodeng Xuexiao Huaxue Xuebao*, 13 (1992) 1521-1523; *C.A.*, 118 (1993) 265616s.

See also 132, 538.

#### 38d. Volatile inorganic compounds

- 558 Arakelyan, V.G.: (Gas chromatographic determination of sulphur hexafluoride in free air at a level of  $n \times 10^{-10}\%$ .) *Zh. Anal. Khim.*, 48 (1993) 1676-1682.
- 559 Bandy, A.R., Thornton, D.C., Ridgeway, R.G.J. and Blomquist, B.W.: Key sulfur-containing compounds in the atmosphere and ocean. Determination by gas chromatography-mass spectrometry and isotopically labeled internal standards. In: *Isot. Eff. Gas-Phase Chem.; ACS Symp. Ser.*, Vol. 502, ACS, Washington, 1992, pp. 409-422; *C.A.*, 118 (1993) 195439k.
- 560 Fan, X. and Ding, T.: (Determination of germanium in Chinese medicine herbs by gas chromatography (GC) through its hydride.) *Sepu*, 11 (1993) 227-228.
- 561 Ishigaki, H.: (Determination of sulfur dioxide in hexafluorosulfide decomposition gas.) *Jpn. Kokai Tokkyo Koho JP 05 34,331 [93 34,331] (Cl. G01N30/88)*, 9 Feb. 1993, Appl. 91/188,444, 29 Jul. 1991; 5 pp.; *C.A.*, 119 (1993) 19522u.
- 562 Sauers, I. and Cacheiro, R.A.: A cryogenic enrichment technique for gas chromatographic detection of disulfur decafluoride in sulfur hexafluoride discharges. In: *Conf. Rec. IEEE Int. Symp. Electr. Insul.*, 1992, pp. 340-344; *C.A.*, 118 (1993) 204212k.

See also 132, 538, 556.

### 39. RADIOACTIVE AND OTHER ISOTOPE COMPOUNDS

See 261, 262, 270, 277, 336, 340, 439.

## Planar Chromatography

### 1. REVIEWS AND BOOKS

- 1 Junker-Buchheit, A. and Jork, H.: Coupling methods in TLC. Part 1. Chromatographic methods. *CLB Chem. Labor Biotech.*, 44 (1993) 266-270; *C.A.*, 119 (1993) 151352t - a review with 28 refs.
- 2 Schuetz, H. and Erdmann, F.: (Quantitative high-pressure TLC in toxicology). *GIT Fachz. Lab.*, 37 (1993) 18-22; *C.A.*, 119 (1993) 132929v - a review with 13 refs.
- 3 Touchstone, J.C.: New developments in planar chromatography. *LC-GC*; 11 (1993) 404-411; *C.A.*, 119 (1993) 173250k - a review with 25 refs.

See also 172, 177.

### 2. FUNDAMENTALS, THEORY AND GENERAL

#### 2a. General

- 4 Ge, Z. and Lin, H.: (Measurement of the overall separation degree of thin-layer chromatography). *Gaodeng Xuexiao Huaxue Xuebao*, 13 (1992) 1383-1385; *C.A.*, 119 (1993) 103435a.

See also 8, 9.

#### 2b. Thermodynamics and theoretical relationships

- 5 Janicka, M. and Rózylo, J.K.: The use of TLC in sandwich chambers for theoretical prediction of mixture separation in HPLC. *J. Planar Chromatogr.*, 6 (1993) 362-367.

See also 17, 31.

#### 2c. Relationship between structure and chromatographic behaviour

See 21, 28.

### 3. GENERAL TECHNIQUES

#### 3a. Apparatus and accessories

- 6 Witkiewicz, Z., Mazurek, M. and Bladec, J.: A new instrument for overpressured layer chromatography. *J. Planar Chromatogr.*, 6 (1993) 407-412.

See also 139.

#### 3b. Detectors and detection reagents

- 7 Brose, C., Rochholz, G., Eermann, F. and Schuetz, H.: (A TLC detection program for 178 pesticides). *Beitr. Gerichtl. Med.*, 50 (1992) 221-228; *C.A.*, 119 (1993) 153487q.
- 8 Freeman, M., Stead, A. and Stroud, R.: The documentation of thin layer chromatography using a video image archival system. *J. Planar Chromatogr.*, 6 (1993) 419-421.
- 9 Végh, Z.: True color photodocumentation of UV-irradiated thin layer chromatograms. *J. Planar Chromatogr.*, 6 (1993) 341-345.
- 10 Wards, W. and Pyka, A.: New visualizing agents for selected fatty derivatives in thin layer chromatography. *J. Planar Chromatogr.*, 6 (1993) 320-322.
- 11 Zou, Z., Wang, H., Chen, D., Tang, Y. and Qiao, X.: (The applicability of a new stable fluorescent reagent CGE (N) for TLC measurements and spectrofluorometry). *Huaxi Yike Daxue Xuebao*, 24 (1993) 194-197; *C.A.*, 119 (1993) 176930t.

See also 57, 108, 110, 142, 180, 231.

#### 3c. Sorbents and columns, packing procedures

- 12 Arslanov, S.S. and Petropavlovskii, G.A.: (Chromatographic properties of esters of microcrystalline cellulose and pyromellitic acid). *Zh. Prikl. Khim. (S.-Peterburg)*, 65 (1992) 2625-2626; *C.A.*, 119 (1993) 134704s.
- 13 Creasy, K.E., Deng, Y.P., Park, J., Borgstedt, E.V.R., Davis, S.P., Suib, S.L. and Shaw, B.R.: Molecular design of aluminosilicate thin film devices. *Mater. Res. Soc. Symp. Proc.*, 233 (Synthesis/Characterization and Novel Applications of Molecular Sieve Material) (1991) 157-167; *C.A.*, 119 (1993) 108015y.
- 14 Flodberg, G. and Roeraade, J.: High pressure thin layer chromatography on 3 micron spherical particle beds. *J. Planar Chromatogr.*, 6 (1993) 252-255.
- 15 Hahn-Deinstrop, E.: TLC plates: initial treatment, pre-washing, activation, conditioning. *J. Planar Chromatogr.*, 6 (1993) 313-318.
- 16 Poole, C.F. and Fernando, W.P.N.: Comparison of the kinetic properties of commercially available precoated silica gel plates. *J. Planar Chromatogr.*, 6 (1993) 357-361.
- 17 Sarbu, C., Dumitrescu, D. and Pop, H.: (Optimum selection and combination of solvent systems for thin-layer chromatography using fuzzy functions). *Rev. Chim. (Bucharest)*, 44 (1993) 450-459; *C.A.*, 119 (1993) 130676m.

#### 3d. Quantitative analysis

- 18 Ferenczi-Fodor, K. and Végh, Z.: Validation of the quantitative planar chromatographic analysis of drug substances. 2: Comparison of some parameters in OPLC and TLC. *J. Planar Chromatogr.*, 6 (1993) 256-258.
- 19 Szepesi, G.: Some aspects of the validation of planar chromatographic methods used in pharmaceutical analysis. II. Ruggedness testing. *J. Planar Chromatogr.*, 6 (1993) 259-268.

- 20 Xiang, Y. and Wang, B.: (Quantitative principles of scanning thin-layer chromatography). *Daxue Huaxue*, 8 (1993) 34-40; C.A., 119 (1993) 173068g.

3e. *Preparative scale chromatography*

See 151, 195.

3f. *Programmed temperature, pressure, vapors, gradients*

- 21 Colthup, P.V., Bell, J.A. and Gadsdon, D.L.: Use of linear solvent gradients in manual and automated multiple development for the separation and characterization of drugs and metabolites by HPTLC. *J. Planar Chromatogr.*, 6 (1993) 386-393.

3g. *High performance procedures*

See 132.

4. SPECIAL TECHNIQUES

4a. *Automation*

- 22 Delvordre, P. and Postaire, E.: Discussion: automation and robotics in planar chromatography. *J. Planar Chromatogr.*, 6 (1993) 289-293.

See also 221.

4b. *Computerization and modelling*

- 23 Qin-Sun, W. and Bing-Wen, Y.: Integration of computer-assisted methods for optimization of HPTLC. *J. Planar Chromatogr.*, 6 (1993) 296-299.

See also 17.

4c. *Combination with other physico-chemical techniques (MS, IR etc.)*

- 24 Banks, C.T.: The development and application of coupled HPLC-TLC for pharmaceutical analysis. *J. Pharm. Biomed. Anal.*, 11 (1993) 705-710.
- 25 Plopkowiak, Z. and Popielarz, M.D.: (Application of chromatographic-spectrophotometric method for stability estimation of azlocillin sodium salt). *Farm. Pol.*, 49, No. 3-4 (1993) 35-39; C.A., 119 (1993) 146449a.
- 26 Tucker, S.A., Acree, W.E., Jr., Fetzer, J.C., Harcey, R.G., Tanga, M.J., Cheng, P.C. and Scott, L.T.: Spectroscopic investigation of fluorescence quenching agents. Part III: Effect of solvent polarity on the selectivity of nitromethane for discriminating between alternant versus nonalternant polycyclic aromatic hydrocarbons. *Appl. Spectrosc.*, 47 (1993) 715-722; C.A., 119 (1993) 82426f.
- 27 Wang, S., Jin, J., Sun, X. and Wang, X.: (Study on photothermal decay property of thin-layer sample concentration with photothermal deflection spectroscopy). *Guangpuxue Yu Guangpu Fenxi*, 12, No. 5 (1992) 7-9; C.A., 119 (1993) 84704u.

See also 3, 96, 107.

4g. *Enantiomers, separation*

- 28 Pyka, A.: A new optical topological index ( $I_{opt}$ ) for predicting the separation of D and L optical isomers by TLC part III. *J. Planar Chromatogr.*, 6 (1993) 282-288.

See also 110, 120.

4h. *Other special techniques*

See 159, 190.

5. HYDROCARBONS AND HALOGEN DERIVATIVES

5b. *Cyclic hydrocarbons, fullerenes*

See 107, 154.

5c. *Halogen derivatives*

- 29 Timofeeva, T.A.: (Method for determining hexachlorocyclohexane, chlorophos and 2,4-dichlorophenoxyacetic acid in water using thin-layer chromatography). *U.S.S.R. SU 1,778,683* (Cl. G01N30/90), 30 Nov. 1992, Appl. 4,862,855, 04 Jul. 1990; C.A., 119 (1993) 85085e.

7. PHENOLS

- 30 Barrero, A.F., Sánchez, J.F. and Rodríguez, I.: Use of alkenylresorcinols from *Ononis speciosa* as synthetic precursors of compounds with potential biological activity. *J. Natural Prod.*, 56 (1993) 1737-1746.
- 31 Dzido, T.H. and Polak, B.: A coadsorption effect in a liquid - solid TLC system of the type silica - heptane + dioxane in a horizontal DS chamber. *J. Planar Chromatogr.*, 6 91993) 378-381.
- 32 Futter, J.E. and Wall, P.: On-site identification of trace concentrations of phenols and cresols in contaminated land leachates using planar chromatography. *J. Planar Chromatogr.*, 6 (1993) 372-377.

See also 5, 139.

8. SUBSTANCES CONTAINING HETEROCYCLIC OXYGEN

8a. *Flavonoids*

- 33 Tschiersch, K. and Hoelzl, J.: Detection and analysis of acyl derivatives of apigenin 7- $\beta$ -D-glucoside. *PZ Wiss*, 5 (1992) 208-214; C.A., 119 (1993) 125292d.

8b. *Aflatoxins and other mycotoxins*

- 34 Leming, L., Jun, Z., Kai, S. and Wenbin, S.: Simultaneous thin layer chromatographic determination of zearalenone and patulin in maize. *J. Planar Chromatogr.*, 6 (1993) 274-277.

## 8c. Other compounds with heterocyclic oxygen (incl. tannins)

- 35 Speranza, G., Manitto, P., Cassara, P., Monti, D., de Castri, D. and Chialva, F.: Studies on aloe, 12. Furoaloesone, a new 5-methylchromone from Cape aloe. *J. Natural Prod.*, 56 (1993) 1089-1094.

## 9. OXO COMPOUNDS, ETHERS, EPOXIDES AND QUINONES

- 36 Kobow, M., Sprung, W.-D. and Schulz, E.: Zur Synthese einiger lägerkettiger phenylsubstituierter  $\alpha$ -Aminoketone. *Pharmazie*, 48 (1993) 697-699.
- 37 Maselennikova, T.A.: (Analysis of products of the reaction of anthraquinone, aniline hydrochloride, and aniline by thin-layer chromatography (TLC)). *Zavod. Lab.*, 59 (1993) 18-20; *C.A.*, 119 (1993) 151372z.
- 38 Novelo, M., Cruz, J.G., Hernández, L., Pereda-Miranda, R., Chai, H., Mar, W. and Pezzuto, J.M.: Cytotoxic constituents from *Hyp-tis verticillata*. *J. Natural Prod.*, 56 (1993) 1728-1736.

See also 221.

## 10. CARBOHYDRATES

## 10a. Mono and oligosaccharides. Structural studies

- 39 Dieckmann-Schuppert, A., Bause, E. and Schwarz, R.T.: Studies on O-glycans of *Plasmodium-falciparum*-infected human erythrocytes. Evidence for O-GlcNAc and O-GlcNAc-transferase in malaria parasites. *Eur. J. Biochem.*, 216 (1993) 779-788.
- 40 Kuczynski, T.W., Kendzierski, K.S. and Sewell, A.C.: Urinary oligosaccharides in pregnant or lactating women: pitfall in screening. *Clin. Chem. (Washington)*, 39 (1993) 2346-2347.
- 41 Lee, Y.-C., Kurosawa, N., Hamamoto, T., Nakaoka, T. and Tsuji, S.: Molecular cloning and expression of Gal $\beta$ 1,3GalNAc $\alpha$ 2,3-sialyltransferase from mouse brain. *Eur. J. Biochem.*, 216 (1993) 377-385.
- 42 Silvert, J.E., Sugumaran, G. and Cogburn, J.N.: Sulphation of proteochondroitin and 4-methylumbelliferyl  $\beta$ -D-xyloside-chondroitin formed by mouse mastocytoma cells cultured in sulphate-deficient medium. *Biochem. J.*, 296 (1993) 119-126.
- 43 Strang, A.-M., Allen, A.K., Holder, A.A. and van Halbeek, H.: The carbohydrate structures of *Trypanosoma brucei brucei* MITat 1.6 variant surface glycoprotein. A re-investigation of the C-terminal glycan. *Biochem. Biophys. Res. Commun.*, 196 (1993) 1430-1439.
- 44 Wilson, I.B.H., Taylor, J.P., Webberley, M.C., Turner, M.J. and Flitsch, S.L.: A novel mono-branched lipid phosphate acts as a substrate for dolichyl phosphate mannose synthetase. *Biochem. J.*, 295 (1993) 195-201.

See also 54, 75, 150.

## 11. ORGANIC ACIDS AND LIPIDS

## 11a. Organic acids and simple esters

- 45 Beyrich, T., Grünert, R. and Jira, T.: Vergleichende Untersuchungen zum katalytischen Einfluss von Cyclodextrinen auf Nitrophenylacetate und Nitrobenzoesäureester. *Pharmazie*, 48 (1993) 597-599.
- 46 Daret, D., Bliin, P., Dorian, B., Rigaud, M. and Larrue, J.: Synthesis of monohydroxylated fatty acids from linoleic acid by rat aortic smooth muscle cells and tissues: influence on prostacyclin production. *J. Lipid Res.*, 34 (1993) 1473-1482.
- 47 Pagou, M. and Koutselinis, A.: Lipophilicity and chromatographic behavior of benzoic acid derivatives. *Chem. Pharm. Bull.*, 41 (1993) 319-324; *C.A.*, 119 (1993) 167863w.
- 48 Smolarz, H.D. and Waksmundzka-Hajnos, M.: Two-dimensional TLC of phenolic acids on cellulose. *J. Planar Chromatogr.*, 6 (1993) 278-281.
- 49 Zeman, I., Schwarz, W., Brat, J. and Zajic, J.: (Chromatographic analyses of the products from fatty acid dimerization). *Potravin. Vedy*, 11 (1993) 1-12; *C.A.*, 119 (1993) 158489b.
- 50 Ziegler, K. and Hummelsiep, S.: Hepatoselective carrier-mediated sodium-independent uptake of pravastatin and pravastatin-lactone. *Biochim. Biophys. Acta*, 1153 (1993) 23-33.

See also 10, 28, 29, 64, 79, 83, 114.

## 11c. Lipids and their constituents

- 51 Aikens, J. and Dix, T.A.: Hydrodioxy (perhydroxy), peroxy and hydroxyl radical-initiated lipid peroxidation of large unilamellar vesicles (liposomes): comparative and mechanistic studies. *Arch. Biochem. Biophys.*, 305 (1993) 516-525.
- 52 Couto, A.S., de Lederkremer, R.M., Colli, W. and Alves, M.J.M.: The glycosylphosphatidylinositol anchor of the trypanosome-specific Tc-85 glycoprotein from *Trypanosoma cruzi*. Metabolic-labeling and structural studies. *Eur. J. Biochem.*, 217 (1993) 597-602.
- 53 Davila, A.M., Marchal, R., Monin, N. and Vandecasteele, J.P.: Identification and determination of individual sophorolipids in fermentation products by gradient elution high-performance liquid chromatography with evaporative light-scattering detection. *J. Chromatogr.*, 648 (1993) 139-149.
- 54 Endo, T., Oda, O., Yamanaka, N., Maeda, K., Yoshida, M. and Kobata, A.: Alterations in the carbohydrate structures of an abnormal protein from sera of patients with rheumatoid arthritis. *Arch. Biochem. Biophys.*, 307 (1993) 119-125.
- 55 Fanjul, L.F., Marrero, I., Gonzalez, J., Quintana, J., Santana, P., Estevez, F., Mato, J.M. and de Galarreta, C.M.R.: Does oligosaccharide-phosphatidylinositol (glycosyl-phosphatidylinositol) hydrolysis mediate prolactin signal transduction in granulosa cells? *Eur. J. Biochem.*, 216 (1993) 747-755.
- 56 Frenkel, R.A., Narahara, H., Eguchi, H., Toyoshima, K. and Johnston, J.M.: Metabolism of hexadecanol by rat type II pneumocytes. *Biochem. Biophys. Res. Commun.*, 196 (1993) 885-891.
- 57 Gérin, C. and Goutx, M.: Separation and quantification of phospholipids from marine bacteria with the Iatroscan Mark IV TLC-FID. *J. Planar Chromatogr.*, 6 (1993) 307-312.

- 58 Gopala Krishna, A.G.: Isolation and identification of the causative factors responsible for color fixation in rice bran oils. *J. Am. Oil Chem. Soc.*, 70 (1993) 785-788.
- 59 Hanisch, F.-G., Chai, W., Rosankiewicz, J.R., Lawson, A.M., Stoll, M.S. and Feizi, T.: Core-typing of O-linked glycans from human gastric mucins. Lack of evidence for the occurrence of the core sequence Gal1-6GalNAc. *Eur. J. Biochem.*, 217 (1993) 645-655.
- 60 He, P., Hu, J. and Macher, B.A.: Glycosphingolipids of rabbit, sheep, and pig thymus. *Arch. Biochem. Biophys.*, 305 (1993) 350-361.
- 61 Hidari, K.I.-P.J., Irie, F., Suzuki, M., Kon, K., Ando, S. and Hirabayashi, Y.: A novel ganglioside with a free amino group in bovine brain. *Biochem. J.*, 296 (1993) 259-263.
- 62 Horwitz, J. and Davis, L.L.: The substrate specificity of brain microsomal phospholipase D. *Biochem. J.*, 295 (1993) 793-798.
- 63 Hvattum, E., Grav, H.J. and Bremer, J.: Hormonal and substrate regulation of 3-thia fatty acid metabolism in Morris 7800 C1 hepatoma cells. *Biochem. J.*, 294 (1993) 917-921.
- 64 Jones, A.L., Lloyd, D. and Harwood, J.L.: Rapid induction of microsomal  $\Delta^{12}(\omega 6)$ -desaturase activity in chilled *Acanthamoeba castellanii*. *Biochem. J.*, 296 (1993) 183-188.
- 65 Li, Z.Y. and Ward, O.P.: Lipase-catalyzed esterification of glycerol and *n*-3 polyunsaturated fatty acid concentrate in organic solvent. *J. Am. Oil Chem. Soc.*, 70 (1993) 745-748.
- 66 Liepkalns, V.A., Myher, J.J., Kuksis, A., Leli, U., Freysz, N. and Hauser, G.: Molecular species of glycerophospholipids and diacylglycerols of cultured SH-S454 human neuroblastoma cells. *Biochem. Cell Biol.*, 71 (1993) 141-149.
- 67 Linard, A., Guesnet, P. and Durand, G.: Separation of the classes of phospholipid by overpressured layer chromatography (OPLC). *J. Planar Chromatogr.*, 6 (1993) 322-323.
- 68 Mukherjee, J.J., Jay, F.T. and Choy, P.C.: Purification, characterization and modulation of a microsomal carboxylesterase in rat liver for the hydrolysis of acyl-CoA. *Biochem. J.*, 295 (1993) 81-86.
- 69 Nakao, T., Kon, K., Ando, S., Miyatake, T., Yuki, N., Li, Y., Furuya, S. and Hirabayashi, Y.: Novel lacto-ganglio type gangliosides with  $G_{M2}$ -epitope in bovine brain which react with IgM from a patient of the amyotrophic lateral-sclerosis-like disorder. *J. Biol. Chem.*, 268 (1993) 21028-21034.
- 70 Norgauer, J., Dobos, G., Kownatzki, E., Dahinden, C., Burger, R., Kupper, R. and Gierschik, P.: Complement fragment C3a stimulates  $Ca^{2+}$  influx in neutrophils via a pertussis-toxin-sensitive G protein. *Eur. J. Biochem.*, 217 (1993) 289-294.
- 71 Ohmichi, M., Zhu, G. and Saltiel, A.R.: Nerve growth factor activates calcium-insensitive protein kinase C- $\epsilon$  in PC-12 rat pheochromocytoma cells. *Biochem. J.*, 295 (1993) 767-772.
- 72 Pörn, M.I., Ares, M.P.S. and Slotte, J.P.: Degradation of plasma membrane phosphatidylcholine appears not to affect the cellular cholesterol distribution. *J. Lipid Res.*, 34 (1993) 1385-1392.
- 73 Ren, S., Ariga, T., Scarsdale, J.N., Zhang, Y., Slominski, A., Livingston, P.O., Ritter, G., Kushi, Y. and Yu, R.K.: Characterization of a hamster melanoma-associated ganglioside antigen as 7-O-acetylated disialoganglioside GD3. *J. Lipid Res.*, 34 (1993) 1565-1572.
- 74 Sattler, W. and Stocker, R.: Greater selective uptake by Hep G2 cells of high-density lipoprotein cholesteryl ester hydroperoxides than of unoxidized cholesteryl esters. *Biochem. J.*, 294 (1993) 771-778.
- 75 Schneider, P., Rosat, J.-P., Ransijn, A., Ferguson, M.A.J. and McConville, M.J.: Characterization of glycoinositol phospholipids in the amastigote stage of the protozoan parasite *Leishmania major*. *Biochem. J.*, 295 (1993) 555-564.
- 76 Song, J. and Foster, D.A.: v-SRC Activates a unique phospholipase D activity that can be distinguished from the phospholipase D activity activated by phorbol esters. *Biochem. J.*, 294 (1993) 711-717.
- 77 Suetake, K., Gasa, S., Taki, T., Chiba, M., Yamaki, T., Ibayashi, Y. and Hashi, K.: Human blood group B-active ganglio-glycosphingolipid in rat glioma. *Biochim. Biophys. Acta*, 1170 (1993) 25-31.
- 78 Taberero, A., Bolanos, J.P. and Medina, J.M.: Lipogenesis from lactate in rat neurons and astrocytes in primary culture. *Biochem. J.*, 294 (1993) 635-638.
- 79 Takahashi, Y., Glasgow, W.C., Suzuki, H., Taketani, Y., Yamamoto, S., Anton, M., Kühn, H. and Brash, A.R.: Investigation of the oxygenation of phospholipids by the porcine leukocyte and human platelet arachidonate 12-lipoxygenases. *Eur. J. Biochem.*, 218 (1993) 165-171.
- 80 Takamatsu, K., Kamei, K., Kubushiro, K., Kiguchi, K., Nozawa, S. and Iwamori, M.: Luteal phase-characteristic induction of  $^{13}SO_3$ -GalCer in human cervical epithelia and uterine endometria, and follicular phase-characteristic formation of a ganglioside-derived negative charge gradient in different regions of fallopian tubes. *Biochim. Biophys. Acta*, 1170 (1993) 232-236.
- 81 Ueda, H., Kobayashi, T., Kishimoto, M., Tsutsumi, T., Watanabe, S. and Okuyama, H.: The presence of  $Ca^{2+}$ -independent phospholipase  $A_1$  highly specific for phosphatidylinositol in bovine brain. *Biochem. Biophys. Res. Commun.*, 195 (1993) 1272-1279.
- 82 Vered, A., Eugenio, M., Arie, D., Augusto, P., Sergio, M. and Shimon, G.: Fluorescence-based diagnosis of lipid storage diseases by analysis of the culture medium of skin fibroblasts. *Clin. Chim. Acta*, 218 (1993) 139-147.
- 83 Wang, X., Dyer, J.H. and Zheng, L.: Purification and immunological analysis of phospholipase D from castor bean endosperm. *Arch. Biochem. Biophys.*, 306 (1993) 486-494.
- 84 Weidner, E., Zhang, Z., Czech, B. and Peter, S.: Entölung von Rohlecithin mit Propan. *Fat Sci. Technol.*, 95 (1993) 347-351.
- 85 Wolucka, B.A., McNeil, M.R., Kalbe, L., Cocito, C. and Brennan, P.J.: Isolation and characterization of a novel glucuronosyl diacylglycerol from *Mycobacterium smegmatis*. *Biochim. Biophys. Acta*, 1170 (1993) 131-136.
- 86 Wright, L.C., Chen, S. and Roufogalis, B.D.: Regulation of the activity and phosphorylation of the plasma membrane  $Ca^{2+}$ -ATPase by protein kinase C in intact human erythrocytes. *Arch. Biochem. Biophys.*, 306 (1993) 277-284.
- 87 Yamada, K. and Sakane, F.: The different effects of sphingosine on diacylglycerol kinase isozymes in Jurkat cells, a human T-cell line. *Biochim. Biophys. Acta*, 1169 (1993) 211-216.

See also 10, 41.



## 12. ORGANIC PEROXIDES

See 51, 74, 92.

## 13. STEROIDS

## 13a. General techniques

- 88 Gazdag, M., Szepesi, G., Mihalyfi, K., Kemenes, B.P., Horvath, P., Vegh, Z., Renyei, M., Trischler, F. and Gorog, S.: (Analysis of steroids. Part 45. Analytical investigation of pipecuronium bromide (Arduan)). *Acta Pharm. Hung.*, 62, No. 3 (1992) 88-96; *C.A.*, 119 (1993) 80360f.
- 89 Lauko, A., Horvath, P., Trischler, F. and Gorog, S.: (Analysis of steroids. Part 44. Analytical investigation of the intermediates of the synthesis of pipecuronium bromide (Arduan)). *Acta Pharm. Hung.*, 62 (1992) 82-87; *C.A.*, 119 (1993) 103477r.

## 13b. Pregnane and androstane derivatives

- 90 Hoebus, J., Daneels, E., Roets, E. and Hoogmartens, J.: Identification of corticosteroid hormones by thin layer chromatography. *J. Planar Chromatogr.*, 6 (1993) 269-273.
- 91 Ichikawa-Haraguchi, M., Sumizawa, T., Yoshimura, A., Furukawa, T., Hiramoto, S., Sugita, M. and Akiyama, S.-i.: Progesterone and its metabolites: the potent inhibitors of the transporting activity of P-glycoprotein in the adrenal gland. *Biochim. Biophys. Acta*, 1158 (1993) 201-208.

## 13d. Sterols

- 92 Esworthy, R.S., Chu, F.-F., Geiger, P., Girotti, A.W. and Doroshov, J.H.: Reactivity of plasma glutathione peroxidase with hydroperoxide substrates and glutathione. *Arch. Biochem. Biophys.*, 307 (1993) 29-34.
- 93 Hwang, K.T. and Maerker, G.: Determination of 6-ketocholestanol in unirradiated and irradiated chicken meats. *J. Am. Oil Chem. Soc.*, 70 (1993) 789-792.
- 94 Swaminathan, S., Wilson, W.K., Pinkerton, F.D., Gerst, N., Ramser, M. and Schroepfer, G.J., Jr.: Inhibitors of sterol synthesis. Chemical synthesis and properties of 3 $\alpha$ -hydroxy-25,26,26,26,27,27,27-heptafluoro-5 $\alpha$ -cholest-8(14)-en-15-one and 25,26,26,26,27,27,27-heptafluorocholesterol and their effects on 3-hydroxy-3-methylglutaryl coenzyme A reductase activity in cultured mammalian cells. *J. Lipid Res.*, 34 (1993) 1805-1823.

See also 10, 74, 78, 98, 216.

## 13e. Bile acids and alcohols

- 95 Miki, S., Mosbach, E.H., Cohen, B.I., Mikami, T., Infante, R., Ayyad, N. and McSherry, C.K.: Metabolism of  $\beta$ -muricholic acid in the hamster and prairie dog. *J. Lipid Res.*, 34 (1993) 1709-1716.

## 13f. Ecdysones and other insect steroid hormones

- 96 Lafont, R., Porter, C.J., Williams, E., Read, H., Morgan, E.D. and Wilson, I.D.: The application of off-line HPTLC-MS-MS to the identification of ecdysteroids in plant and arthropod samples. *J. Planar Chromatogr.*, 6 (1993) 421-424.

## 13g. Other steroids

See 88.

## 14. STEROID GLYCOSIDES AND SAPONINS

See 33, 136.

## 15. TERPENES AND OTHER VOLATILE AROMATIC COMPOUNDS

## 15a. Terpenes

- 97 Achenbach, H., Waibel, R., Hefter-Bübl, U. and Constenla, M.A.: Constituents of *Fevillea cordifolia*: new norcucurbitacin and cucurbitacin glycosides. *J. Natural Prod.*, 56 (1993) 1506-1519.
- 98 Bondioli, P., Mariani, C., Lanzani, A., Fedeli, E. and Muller, A.: Squalene recovery from olive oil deodorizer distillates. *J. Am. Oil Chem. Soc.*, 70 (1993) 763-766.
- 99 Burnell, R.H., Dumont, N. and Théberge, N.: Synthesis of O-methyl nimbinone. *J. Natural Prod.*, 56 (1993) 1930-1936.
- 100 Chen, G.-F., Li, Z.-L., Pan, D.-J., Tang, C.-M., He, X., Xu, G.-Y., Chen, K. and Lee, K.-H.: The isolation and structural elucidation of four novel triterpene lactones, pseudolarolides A, B, C, and D, from *Pseudolarix kaempferi*. *J. Natural Prod.*, 56 (1993) 1114-1122.
- 101 Hufford, C.D., Jia, Y., Croom, E.M., Jr., Muhammed, I., Okunade, A.L., Clark, A.M. and Rogers, R.D.: Antimicrobial compounds from *Petalostemum purpureum*. *J. Natural Prod.*, 56 (1993) 1878-1889.
- 102 Inoue, H., Korenaga, T., Sagami, H., Koyama, T., Sugiyama, H. and Ogura, K.: Formation of farnesal and 3-hydroxy-2,3-dihydrofarnesal from farnesol by protoplasts of *Botryococcus braunii*. *Biochem. Biophys. Res. Commun.*, 196 (1993) 1401-1405.
- 103 Lou, H., Li, X., Onda, M., Konda, Y., Machida, T., Toda, Y. and Harigaya, Y.: Further isolation of glycosides from *Cynanchum hancockianum*. *J. Natural Prod.*, 56 (1993) 1437-1443.
- 104 Maatooq, G., El-Sharkawy, S., Afifi, M.S. and Rosazza, J.P.N.: Microbial transformation of cedrol. *J. Natural Prod.*, 56 (1993) 1039-1050.
- 105 Römer, S., Hugueney, P., Bouvier, F., Camara, B. and Kuntz, M.: Expression of the genes encoding the early carotenoid biosynthetic enzymes in *Capsicum annum*. *Biochem. Biophys. Res. Commun.*, 196 (1993) 1414-1421.
- 106 Samaranyake, G., Neidigh, K.A. and Kingston, D.G.I.: Modified taxols. 8. Deacylation and reacylation of baccatin III. *J. Natural Prod.*, 56 (1993) 884-898.

See also 151.

## 16. NITRO AND NITROSO COMPOUNDS

- 107 Hayes-Majstorovic, J., Guernet-Nivaud, E., Guernet, M. and Viel, C.: Mass spectrometry and liquid chromatography-mass spectrometry of  $\alpha$ -amino  $\beta$ -nitrostilbenes. *Anal. Lett.*, 26 (1993) 2239-2249.
- 108 Hiley, R.W.: Investigations of thin layer chromatographic techniques used for forensic explosives analysis in the early 1970s. *J. Forensic Sci.*, 38 (1993) 864-873; *C.A.*, 119 (1993) 133021e.
- 109 Tyrpien, K.: TLC identification of some nitrogen derivatives of PAH in airborne particulate matter. *J. Planar Chromatogr.*, 6 (1993) 413-415.

See also 45.

## 17. AMINES, AMIDES AND RELATED NITROGEN COMPOUNDS

## 17a. Amines and polyamines

- 110 Heuser, D. and Meads, P.: HPTLC separation of chiral amino alcohols and amines after derivatization with Marfey's reagent. *J. Planar Chromatogr.*, 6 (1993) 324-325.
- 111 Hodisan, T. and Hopartean, I.: Separation of some isomeric organic compounds by thin-layer chromatography. *Stud. Univ. Banes-Bolyai, Chem.*, 35 (1990) 25-29; *C.A.*, 119 (1993) 173274w.
- 112 Kotzabasis, K., Christakis-Hampas, M.D. and Roubelakis-Angelakis, K.A.: A narrow-bore HPLC method for the identification and quantitation of free, conjugated, and bound polyamines. *Anal. Biochem.*, 214 (1993) 484-489.
- 113 Whitaker, D.P. and Goosey, M.W.: Purification and properties of the enzyme arylamine N-acetyltransferase from the housefly *Musca domestica*. *Biochem. J.*, 295 (1993) 149-154.

See also 37.

## 17b. Catecholamines and their metabolites

- 114 Klaus, R., Fischer, W. and Hauck, H.E.: Extension of the application of the thermal *in-situ* reaction on  $\text{NH}_2$  layers to the detection of catecholamines in biological materials. *Chromatographia*, 37 (1993) 133-143.

## 17d. Other amine derivatives and amides (excl. peptides)

- 115 Amarowicz, R., Wanasundara, P.K.J.P.D. and Shahidi, F.: TLC separation of linamarin, linustatin and neolinustatin. *Nahrung*, 37 (1993) 88-90; *C.A.*, 119 (1993) 179468j.

## 18. AMINO ACIDS AND PEPTIDES; CHEMICAL STRUCTURE OF PROTEINS

## 18a. Amino acids and their derivatives

- 116 Chénais, B., Yapo, A., Lepoivre, M. and Tenu, J.-P.: *N $\omega$* -Hydroxy-L-arginine, a reactional intermediate in nitric oxide biosynthesis, induces cytostasis in human and murine tumor cells. *Biochem. Biophys. Res. Commun.*, 196 (1993) 1558-1565.
- 117 Das, B. and Sawant, S.: Quantitative HPTLC analysis of dansyl amino acids. *J. Planar Chromatogr.*, 6 (1993) 294-295.
- 118 Hill, R.E., White, R.L. and Smith, K.C.: The identification of 5-hydroxy-L-norvaline in cultures of pyridoxine auxotrophs of *Escherichia coli*. *J. Natural Prod.*, 56 (1993) 1246-1254.
- 119 Koopdonk-Kool, J.M., van Lopik-Peterse, M.C., Veenboer, G.J.M., Visser, T.J., Schoenmakers, C.H.H. and de Vijlder, J.J.M.: Quantification of type III iodothyrosine deiodinase activity using thin-layer chromatography and phosphor screen autoradiography. *Anal. Biochem.*, 214 (1993) 329-331.
- 120 LeFevre, J.W.: Reversed-phase thin-layer chromatographic separations of enantiomers of dansyl-amino acids using  $\beta$ -cyclodextrin as a mobile phase additive. *J. Chromatogr.*, 653 (1993) 293-302.
- 121 Preece, M.A., Sewell, I.J., Taylor, J.A. and Green, A.: Vigabatrin - interference with urinary amino acid analysis. *Clin. Chim. Acta*, 218 (1993) 113-116.
- 122 Xuan, H.T.K. and Lederer, M.: Adsorption chromatography on cellulose. X. Adsorption of tryptophan and derivatives from  $\text{CuSO}_4$ -containing eluents. *J. Chromatogr.*, 645 (1993) 185-188.

See also 6, 11, 12, 28, 123, 126.

## 18b. Peptides, peptidic and proteinous hormones, growth factors

- 123 Johnstone, S.A., Hubashy, I. and Waisman, D.M.: Regulation of annexin I-dependent aggregation of phospholipid vesicles by protein kinase C. *Biochem. J.*, 294 (1993) 801-807.
- 124 Lauritzen, E., Másson, M., Rubin, I., Bjerrum, O.J. and Holm, A.: Peptide dot immunoassay and immunoblotting: Electroblotting from aluminum thin-layer chromatography plates and isoelectric focusing gels to activated nitrocellulose. *Electrophoresis (Weinheim)*, 14 (1993) 852-859.
- 125 Moss, A.M. and Livingston, J.N.: Distinct  $\beta$ -subunits are present in hybrid insulin-like-growth-factor-1 receptors in the central nervous system. *Biochem. J.*, 294 (1993) 685-692.
- 126 Page, A.E., Fuller, K., Chambers, T.J. and Warburton, M.J.: Purification and characterization of a tripeptidyl peptidase I from human osteoclastomas: evidence for its role in bone resorption. *Arch. Biochem. Biophys.*, 306 (1993) 354-359.
- 127 Venema, R.C., Raynor, R.L., Noland, T.A., Jr. and Kuo, J.F.: Role of protein kinase C in the phosphorylation of cardiac myosin light chain 2. *Biochem. J.*, 294 (1993) 401-406.

## 21. PURINES, PYRIMIDINES, NUCLEIC ACIDS AND THEIR CONSTITUENTS

## 21a. Purines, pyrimidines, nucleosides, nucleotides

- 128 Johnson, D.F., Poksay, K.S. and Innerarity, T.L.: The mechanism for apo-B mRNA editing is deamination. *Biochem. Biophys. Res. Commun.*, 195 (1993) 1204-1210.
- 129 Leonard, S.A., Wong, S.C. and Nyce, J.W.: Quantitation of 5-methylcytosine by one-dimensional high-performance thin-layer chromatography. *J. Chromatogr.*, 645 (1993) 189-192.
- 130 Sakamoto, K., Kawai, G., Niimi, T., Satoh, T., Sekine, M., Yamaizumi, T., Nishimura, S., Miyazawa, T. and Yokoyama, S.: A modified uridine in the first position of the anticodon of a minor species of arginine tRNA, the argU gene product, from *Escherichia coli*. *Arch. Biochem. Biophys.*, 305 (1993) 369-375.
- 131 Um, H.-D. and Klein, C.: Evidence for allosteric regulation of succinyl-CoA synthetase. *Biochem. J.*, 295 (1993) 821-826.
- 132 Wang, S. and Ma, L.: (Quality evaluation of synthetic oligonucleotides by high performance thin-layer chromatography). *Seppu*, 11 (1993) 146-148; *C.A.*, 119 (1993) 85028p.

See also 114.

## 21c. Nucleic acids, DNA

- 133 Steinberg, J.J.: Measurement of DNA adducts by thin layer chromatography. *PCT Int. Appl. WO 93 16,198 (Cl. C12Q1/68)*, 19 Aug. 1993, US Appl. 831,278, 06 Feb. 1992, 41 pp.; *C.A.*, 119 (1993) 175384f.

## 22. ALKALOIDS

- 134 Boyes-Korkis, J.M., Gurney, K.A., Penn, J., Mantle, P.G., Bilton, J.N. and Sheppard, R.N.: Anacine, a new benzodiazepine metabolite of *Penicillium aurantiogriseum* produced with other alkaloids in submerged fermentation. *J. Natural Prod.*, 56 (1993) 1707-1717.
- 135 Cao, H.: (Determination of aconitine in Wutou analogs (roots of *Aconitum*) and their processed products in Hong Kong market). *Zhongguo Zhongyao Zazhi*, 18 (1993) 279-281; *C.A.*, 119 (1993) 167853t.
- 136 Ferreira, F., Moyna, P., Soule, S. and Vázquez, A.: Rapid determination of solanum glycoalkaloids by thin-layer chromatographic scanning. *J. Chromatogr.*, 653 (1993) 380-384.
- 137 Golkiewicz, W., Gadzikowska, M., Kuczynski, J. and Jusiak, L.: Micropreparative chromatography of some quaternary alkaloids from the roots of *Chelidonium majus* L. *J. Planar Chromatogr.*, 6 (1993) 382-385.
- 138 Hanuman, J.B. and Katz, A.: Isolation and identification of four norditerpenoid alkaloids from processed and unprocessed root tubers of *Aconitum ferox*. *J. Natural Prod.*, 56 (1993) 801-809.
- 139 Huang, T. and Jin, H.: (Paper roll chromatography). *Zhongguo Yaoxue Zazhi*, 27 (1992) 669-672; *C.A.*, 119 (1993) 103444c.
- 140 Kobayashi, J., Itagaki, F., Shigemori, H. and Sasaki, T.: Three new onnamide congeners from the Okinawan marine sponge *Theonella* sp. *J. Natural Prod.*, 56 (1993) 976-981.

- 141 Li, H., Zhang, W., Zhang, Q., Zhang, X., Wang, J., Chu, Y. and He, Q.: (Quantitative determination of berberine in Weichanghangtong capsule). *Zhongcaoyao*, 24 (1993) 244-245; *C.A.*, 119 (1993) 167849w.
- 142 Manjunath, M.N., Sattigeri, V.D. and Nagaraja, K.V.: Thin-layer chromatographic method for the detection of caffeine in non-alcoholic beverages. *Indian Food Packer*, 47 (1993) 51-52; *C.A.*, 119 (1993) 179469k.
- 143 Stennert, A. and Maier, H.G.: (Trigonelline in coffee. Part 1. Comparison of thin-layer with high-performance liquid chromatography. Simultaneous determination of caffeine.) *Z. Lebensm.-Unters. Forsch.*, 196 (1993) 430-434; *C.A.*, 119 (1993) 179522x.
- 144 Watson, A.B., Freer, I.K.A., Robins, D.J. and Walton, N.J.: Stereochemistry of the enzymic processes involved in the biosynthesis of scopolamine from hyoscyamine. *J. Natural Prod.*, 56 (1993) 1234-1238.

See also 208.

## 23. OTHER SUBSTANCES CONTAINING HETEROCYCLIC NITROGEN

## 23a. Porphyrins and other pyrroles

- 145 Jacob, K. and Doss, M.O.: Composition of urinary coproporphyrin isomers I-IV in human porphyrias. *Eur. J. Clin. Chem. Clin. Biochem.*, 31 (1993) 617-624.

## 23c. Indole derivatives and plant hormones (gibberelins)

See 222.

## 23e. Other N-heterocyclic compounds

- 146 Fujimoto, H., Ikeda, M., Yamamoto, K. and Yamazaki, M.: Structure of fischerin, a new toxic metabolite from an Ascomycete, *Neosartorya fischeri* var. *fischeri*. *J. Natural Prod.*, 56 (1993) 1268-1275.
- 147 Renz, P., Emdres, B., Kurz, B. and Marquart, J.: Biosynthesis of vitamin B12 in anaerobic bacteria. Transformation of 5-hydroxybenzimidazole and 5-hydroxy-6-methylbenzimidazole into 5,6-dimethylbenzimidazole in *Eubacterium limosum*. *Eur. J. Biochem.*, 217 (1993) 1117-1121.

See also 31, 109, 182.

## 24. ORGANIC SULPHUR COMPOUNDS (INCL. GLUCOSINOLATES)

- 148 Madelaine-Dupuich, C., Azema, J., Escoula, B., Rico, I. and Lattes, A.: Analysis of N-acyl aminonaphthalene sulphonic acid derivatives with potential anti-human immunodeficiency virus activity by thin-layer chromatography and flame ionization detection. *J. Chromatogr.*, 653 (1993) 178-180.
- 149 Muller, B. and Ruhnke, A.: Qualitätsprüfung von Knoblauchpulver. Vorschläge für eine Arzneibuchmonographie. *Dtsch. Apoth.-Ztg.*, 133 (1993) 37-47.

25. ORGANIC PHOSPHORUS COMPOUNDS (INCL. SUGAR PHOSPHATES)
- 150 Kellerer, M., Machicao, F., Berti, L., Sixt, B., Mushack, J., Seffer, E., Mosthaf, L., Ullrich, A. and Häring, J.U.: Inositol phospho-oligosaccharides from rat fibroblasts and adipocytes stimulate 3-O-methylglucose transport. *Biochem. J.*, 295 (1993) 699-704.
- 151 Kennedy Keller, R. and Thompson, R.: Rapid synthesis of isoprenoid diphosphates and their isolation in one step using either thin layer or flash chromatography. *J. Chromatogr.*, 645 (1993) 161-167.
- See also 6, 123, 125, 127.
27. VITAMINS AND VARIOUS ANIMAL GROWTH FACTORS (NON-PEPTIDIC)
- 152 Baldet, P., Gerbling, H., Axiotis, S. and Douce, R.: Biotin biosynthesis in higher plant cells. Identification of intermediates. *Eur. J. Biochem.*, 217 (1993) 479-485.
- 153 Madden, U.A. and Stahr, H.M.: Reverse phase thin layer chromatography assay of vitamin K in bovine liver. *J. Liq. Chromatogr.*, 16 (1993) 2825-2834.
- 154 Zhuravleva, E.D. and Kritskaya, S.L.: (Analytical method for monitoring the quality of a biological preparation of vitamins). *Pishch Prom-st. (Moscow)*, No. 10 (1992) 21; *C.A.*, 119 (1993) 137699e.
28. ANTIBIOTICS
- 155 Aburaki, S., Okuyama, S., Hoshi, H., Kamachi, H., Nishio, M., Hasegawa, T., Masuyoshi, S., Iimura, S., Konishi, M. and Oki, T.: Synthesis and antifungal activity of pradimicin derivatives. Modifications on the aglycone part. *J. Antibiot.*, 46 (1993) 1447-1457.
- 156 Blanchflower, S.E., Banks, R.M., Everett, J.R. and Reading, C.: Further novel metabolites of the paraherquamide family. *J. Antibiot.*, 46 (1993) 1355-1363.
- 157 Gould, S.J., He, W. and Cone, M.C.: New cyanocyclines from a cyanide-treated broth of *Streptomyces lusitanus*. *J. Natural Prod.*, 56 (1993) 1239-1245.
- 158 Isono, F., Sakaida, Y., Takahashi, S., Kinoshita, T., Nakamura, T. and Inukai, M.: Mureidomycins E and F, minor components of mureidomycins. *J. Antibiot.*, 46 (1993) 1203-1207.
- 159 Iwata-Reuyl, D., Basak, A., Silverman, L.S., Engle, C.A. and Townsend, C.A.: Synthesis and reaction of potential alternate substrates and mechanism-based inhibitors of clavaminic synthase. *J. Natural Prod.*, 56 (1993) 1373-1396.
- 160 Johdo, O., Tone, H., Okamoto, R., Yoshimoto, A., Naganawa, H., Sawa, T. and Takeuchi, T.: Microbial glycosidation of some anthracycline antibiotics by an antibiotic-negative mutant of acl-rubicin producer. *J. Antibiot.*, 46 (1993) 1219-1231.
- 161 Kageyama, S., Tsuchiya, T. and Umezawa, S.: Synthesis of 3-deoxy-3,4-didehydro derivatives of 5-O-mycaminosyltylonolide, 5-O-(4-deoxymycaminosyl)tylonolide, and desmycosin. *J. Antibiot.*, 46 (1993) 1265-1278.
- 162 Kamiyama, T., Itezono, Y., Umino, T., Satoh, T., Nakayama, N. and Yokose, K.: Tetrafibricin, a novel fibrinogen receptor antagonist II. Structural elucidation. *J. Antibiot.*, 46 (1993) 1047-1054.
- 163 Kamiyama, T., Umino, T., Fujisaki, N., Fujimori, K., Satoh, T., Yamashita, Y., Ohshima, S., Watanabe, J. and Yokose, K.: Tetrafibricin, a novel fibrinogen receptor antagonist I. Taxonomy, fermentation, isolation, characterization and biological activities. *J. Antibiot.*, 46 (1993) 1039-1046.
- 164 Kobrehel, G., Lazarevski, G., Kelneric, Z. and Dokic, S.: 9a,11-Cyclic carbamates of 15-membered azalides. *J. Antibiot.*, 46 (1993) 1239-1245.
- 165 Kumagai, K., Fukui, A., Tanaka, S., Ikemoto, M., Moriguchi, K. and Nabeshima, S.: PC-766B, A new macrolide antibiotic produced by *Nocardia brasiliensis*. II. Isolation, physico-chemical properties and structure elucidation. *J. Antibiot.*, 46 (1993) 1139-1144.
- 166 Kuroda, K., Yoshida, M., Uosaki, Y., Ando, K., Kawamoto, I., Oishi, E., Onuma, H., Yamada, K. and Matsuda, Y.: AS-183, a novel inhibitor of acyl-CoA:cholesterol acyltransferase, produced by *Scedosporium* sp. SPC-15549. *J. Antibiot.*, 46 (1993) 1196-1202.
- 167 Laatsch, H., Kellner, M., Wolf, G., Lee, Y.-S., Hansske, F., Koenetschny-Rapp, S., Pessara, U., Scheuer, W. and Stockinger, H.: Oligomycin F, a new immunosuppressive homologue of oligomycin A. *J. Antibiot.*, 46 (1993) 1334-1341.
- 168 Saito, A., Naito, S., Kobayashi, M., Tsuchiya, M., Toyama, T., Kaneko, S., Nanba, T. and Morisawa, Y.: Synthesis and anthelmintic activity of 13-alkoxymilbemycin derivatives. *J. Antibiot.*, 46 (1993) 1252-1264.
- 169 Shafiee, A., Chen, T.S., Arison, B.S., Dumont, F.J., Colwell, L. and Kaplan, L.: Enzymatic synthesis and immunosuppressive activity of novel desmethylated immunomycins (ascomycins). *J. Antibiot.*, 46 (1993) 1397-1405.
- 170 Ueno, M., Yoshinaga, I., Amemiya, M., Someno, T., Iinuma, H., Ishizuaka, M. and Takeuchi, T.: Delaminomycins, novel extracellular matrix receptor antagonists. V. Biosynthesis. *J. Antibiot.*, 46 (1993) 1390-1396.
- 171 Zhao, Y., Gao, H. and Dou, Z.: (Improved thin-layer chromatography in clindamycin synthesis). *Zhongguo Yiyao Gongye Zazhi*, 23 (1992) 506-507; *C.A.*, 119 (1993) 125320m.
- See also 25.
29. INSECTICIDES, PESTICIDES AND OTHER AGROCHEMICALS
- 172 Rathore, H.S. and Begum, T.: Thin-layer chromatographic methods for use in pesticide residue analysis. *J. Chromatogr.*, 643 (1993) 271-290 - a review with 290 refs.
- 29a. General techniques
- 173 Judge, D.N., Mullins, D.E. and Young, R.W.: High performance thin layer chromatography of several pesticides and their major environmental by-products. *J. Planar Chromatogr.*, 6 (1993) 300-306.
- See also 2, 6, 7, 231.

## 29b. Chlorinated insecticides

- 174 Sherma, J. and Rolfe, C.: Determination of diflubenuron residues in water by solid-phase extraction and quantitative high-performance thin-layer chromatography. *J. Chromatogr.*, 643 (1993) 337-339.

## 29c. Phosphorus insecticides

- 175 Xie, B., Zhou, Z., Song, R., Zhang, R. and Ji, J.: (Detection of a new organophosphorus pesticide). *Lihua Jianyan, Huaxue Fence*, 28 (1992) 296-297; C.A., 119 (1993) 153885t.

## 29d. Carbamates

- 176 Rathore, H.S. and Begum, T.: Thin-layer chromatographic behaviour of carbamate pesticides and related compounds. *J. Chromatogr.*, 643 (1993) 321-329.
- 177 Tekel, J. and Kovačičová, J.: Chromatographic methods in the determination of herbicide residues in crops food and environmental samples. *J. Chromatogr.*, 643 (1993) 291-303 - a review with 117 refs.

## 29e. Herbicides

- 178 Popova, T.V. and Zaplishny, V.N.: (Determination of dimethomorph residues in potatoes and tomatoes using the technique of thin-layer chromatography). *Izv. Timiryazevsk. S-kh. Akad.*, No. 6 (1992) 159-161; C.A., 119 (1993) 137708g.

## 29f. Fungicides

- 179 Dellacassa, E., Martinez, R. and Moyna, P.: Simultaneous determination of *o*-phenylphenol, imazalil, and thiabendazole residues in citrus fruits by TLC - densitometry. *J. Planar Chromatogr.*, 6 (1993) 326-327.

## 29g. Other types of pesticides and various agrochemicals

- 180 Pasha, A. and Vijayashankar, Y.N.: Thin-layer chromatographic detection of pyrethroid insecticides using *o*-tolidine. *Analyst (Cambridge)*, 118 (1993) 777-778; C.A., 119 (1993) 175475m.
- 181 Patil, V.B. and Shingare, M.S.: Thin-layer chromatographic detection of carbaryl using phenylhydrazine hydrochloride. *J. Chromatogr.*, 653 (1993) 181-183.

## 30. SYNTHETIC AND NATURAL DYES

## 30a. Synthetic dyes

- 182 Brivida, K., Fraser, G., Sies, H. and Ketterer, B.: Distribution of the monochlorobimane-glutathione conjugate between nucleus and cytosol in isolated hepatocytes. *Biochem. J.*, 294 (1993) 631-633.
- 183 Qi, D., Bian, Z. and Jin, W.: (Separation of impurities in crystal violet dye with TLC method). *Huaxue Tongbao*, No. 5 (1993) 40-42; C.A., 119 (1993) 97955b.
- 184 Singh, S., Das, M. and Khanna, S.K.: Bio-metabolism of Green S and Indigo Carmine through caecal microflora of rats. *Biochem. Biophys. Res. Commun.*, 195 (1993) 490-496.

- 185 Wall, P.E.: The value of planar chromatography for the analysis of triphenylmethane dyes. *J. Planar Chromatogr.*, 6 (1993) 394-403.

- 186 Yin, P., Li, H. and Yan, C.: (Separation of alkaline triphenylmethane dyer by ion-pair reversed phase thin-layer chromatography). *Fenxi Huaxue*, 21 (1993) 365; C.A., 119 (1993) 162364z.

## 30b. Chloroplast and other natural pigments

- 187 Drescher, J.N., Sherma, J. and Fried, B.: Thin layer chromatographic determination of alpha-carotene on magnesium oxide layers. *J. Liq. Chromatogr.*, 16 (1993) 3557-3561.
- 188 Shou, G.: (Quantitative determination of indirubin in Bingdun-ding oral liquid by TLC densitometry method). *Zhongcaoyao*, 24 (1993) 15-16; C.A., 119 (1993) 103447f.

## 32. DRUG ANALYSIS

## 32a. Drug analysis, general techniques

- 189 Hu, Y.Z., He, H., Lai, Y.H. and Ni, K.Y.: (New method for selectivity optimization in thin-layer chromatography and its application in pharmaceutical analysis). *Yaouxue Xuebao*, 28 (1993) 50-56; C.A., 119 (1993) 80318y.
- 190 Xia, S., Zhou, J., Zhou, J., Song, B. and Wu, Q.: (Determination of rivanol in human serum albumin by thin-layer fluorescent scanning technique). *Yaowu Fenxi Zazhi*, 13 (1993) 233-236; C.A., 119 (1993) 195033u.

See also 18, 19, 21, 24.

## 32b. Antirheumatics and antiinflammatory drugs

See 90.

## 32c. Autonomic and cardiovascular drugs

- 191 Ademola, J.I., Chow, C.A., Wester, R.C. and Maibach, H.I.: Metabolism of propranolol during percutaneous absorption in human skin. *J. Pharm. Sci.*, 82 (1993) 767-770.

See also 166.

## 32d. Central nervous system drugs

- 192 Auricchio, M.T., Batistic, M.A. and Markman, B.E.O.: (Detection of anorectics and benzodiazepines in pharmaceuticals used in the treatment of obesity). *Rev. Inst. Adolfo Lutz*, 51 (1991) 105-110; C.A., 119 (1993) 146711e.
- 193 El-Bardicy, M.G., El-Saharty, Y.S. and Tawakkol, M.S.: Bromometric determination of carbimazole. *Talanta*, 40 (1993) 577-583; C.A., 119 (1993) 125332s.
- 194 Matsumoto, S., Tagawa, M., Hatoyama, T., Fujii, T., Miyazaki, H. and Sekine, Y.: Absorption, distribution, metabolism and excretion of [carbonyl-<sup>14</sup>C]mosapride citrate after repeated oral administration in rats. *Arzneim.-Forsch.*, 43 (1993) 1103-1108.

- 195 Matsumoto, S., Yoshida, K., Itogawa, A., Tagawa, M., Fujii, T., Miyazaki, H. and Sekine, Y.: Metabolism of [carbonyl-<sup>14</sup>C]mosapride citrate after a single oral administration in rats, dogs and monkeys. *Arzneim.-Forsch.*, 43 (1993) 1095-1102.
- 196 Parodi, B., Caviglioli, G., Bachi, A., Cafaggi, S. and Romussi, G.: Herbal mixtures with claimed slimming activity: determination by TLC and HPLC of illegally added drugs. *Pharmazie*, 48 (1993) 678-681.

See also 121.

32e. *Chemotherapeutics (exc. cytostatics and antibiotics)*

- 197 Abjean, J.-P.: Qualitative screening for nitrofurans residues in food by planar chromatography. *J. Planar Chromatogr.*, 6 (1993) 319-320.
- 198 Abjean, J.-P. and Gaugain, M.: Planar chromatographic determination of ivermectin at low nanogram levels. *J. Planar Chromatogr.*, 6 (1993) 425-427.
- 199 Lawen, A. and Traber, R.: Substrate specificities of cyclosporin synthetase and peptidase SDZ 214-103 synthetase. Comparison of the substrate specificities of the related multifunctional polypeptides. *J. Biol. Chem.*, 268 (1993) 20452-20465.
- 200 Salomies, H.: Quantitative HPTLC of sulfonamides in pharmaceutical preparations. *J. Planar Chromatogr.*, 6 (1993) 337-340.
- 201 Shen, Z.F. and Xie, M.Z.: (Determination of berberine in biological specimen by high performance TLC and fluoro-densitometric method). *Yaoxue Xuebao*, 28 (1993) 532-536; C.A., 119 (1993) 195032t.

See also 156.

32f. *Cytostatics*

- 202 Boddy, A.V., Yule, S.M., Wyllie, R., Price, L., Pearson, A.D.J. and Idle, J.R.: Pharmacokinetics and metabolism of ifosfamide administered as a continuous infusion in children. *Cancer Res.*, 53 (1993) 3758-3764.
- 203 Borel, A.G. and Abbott, F.S.: Identification of carbamoylated thiol conjugates as metabolites of the antineoplastic 1-(2-chloroethyl)-3-cyclohexyl-1-nitrosourea, in rats and humans. *Drug Metab. Disp.*, 21 (1993) 889-901.

See also 18.

32g. *Other drug categories*

- 204 Konfino, M., Delcheva, S. and Mindjova, K.: The isolation of lovastatin and its determination by densitometric TLC and by HPLC. *J. Planar Chromatogr.*, 6 (1993) 404-406.
- 205 Song, J., Liu, X., Shi, Y., Zhou, L. and Hanying, Y.: (The formula design and the screen model *in vitro* for praziquantel spray). *Zhongguo Yaoke Daxue Xuebao*, 24 (1993) 109-111; C.A., 119 (1993) 125094r.
- 206 Zhou, C., Peng, S. and Liao, W.: (Method of quality control of Leignteng (*Tripterygium wilfordii*) topical liquids). *Zhongcaoyao*, 24 (1993) 191-192; C.A., 119 (1993) 103082h.

See also 167, 169, 196, 219.

32h. *Toxicological and forensic applications*

- 207 Taylor, M.R. and Westwood, S.A.: An investigation of drug detection in equine urine samples by modern TLC. *J. Planar Chromatogr.*, 6 (1993) 415-418.

See also 2, 6.

32i. *Plant extracts*

- 208 Feng, S., Bai, M. and Xie, Q.: (Quantitative analysis of Huangbai). *Shenyang Yaoxueyuan Xuebao*, 9 (1992) 212-214; C.A., 119 (1993) 103452d.
- 209 Liu, W., Zhou, W., Wu, D., Ying, L. and An, D.: (Separation techniques and quantitative determination of wuwei alcohol A contained in Sheng Mai Yin by TLC). *Zhongguo Yaoke Daxue Xuebao*, 24 (1993) 78-81; C.A., 119 (1993) 146715j.
- 210 Liu, X. and Bao, D.: (Preparation of Mianyishuangbao capsule). *Zhongcaoyao*, 24 (1993) 189-190; C.A., 119 (1993) 103081g.
- 211 Markowski, W. and Matysik, G.: Analysis of plant extracts by multiple development thin-layer chromatography. *J. Chromatogr.*, 646 (1993) 434-438.
- 212 Marston, A., Maillard, M. and Hostetmann, K.: Search for antifungal, molluscicidal and larvicidal compounds from African medicinal plants. *J. Ethnopharmacol.*, 38 (1993) 215-223; C.A., 119 (1993) 79882w.
- 213 Sane, R.T. and Kuber, V.V.: Standardization of folk medicine - an interdisciplinary approach. Part 1. *Indian Drugs*, 30 (1993) 220-224; C.A., 119 (1993) 79941q.
- 214 Shen, Y.-C. and Chen, C.-H.: New secoiridoid dilactones from *Fraxinus uhdei*. *J. Natural Prod.*, 56 (1993) 1905-1911.
- 215 Su, S., Xu, C., Sui, C., Sun, R., Wang, Y., Lui, H. and Shen, X.: (Analysis of the active components of Chinese drug Houpo and similar drugs). *Shenyang Yaoxueyuan Xuebao*, 9 (1992) 185-189, 223; C.A., 119 (1993) 125293e.
- 216 Tu, P., Xu, G., Xu, L. and Jin, R.: (Analysis of the fat soluble constituents in Chinese drugs shashen by TLC). *Zhongcaoyao*, 24 (1993) 128-130; C.A., 119 (1993) 146695c.
- 217 Wang, Z., Shi, D., Chen, Z., Zheng, L. and Chen, C.: (Quantitative determination of curculigoside in rhizomes of *Curculigo orchoides* by TLC-densitometry). *Shanghai Yike Daxue Xuebao*, 20, No. 1 (1993) 55-58; C.A., 119 (1993) 125306m.
- 218 Yamamoto, K., Yamashita, K., Hitomi, N., Suzuki, A. and Yoneda, K.: (Studies on the constituents of *Atractylodes* rhizome, constituents in the rhizome of *Atractylodes japonica* and TLC analysis of Jutsu). *Shoyakugaku Zasshi*, 47 (1993) 12-16; C.A., 119 (1993) 167573b.
- 219 Zhao, S. and Ma, C.: (Quality standard of Qingrejiadu oral liquid). *Zhongcaoyao*, 24 (1993) 242-243; C.A., 119 (1993) 125311j.
- 220 Zhou, Y., Xiao, M. and Zhang, H.: (Quality of *Fructus evodiae*). *Zhongguo Zhongyao Zazhi*, 18 (1993) 201-203; C.A., 119 (1993) 167848v.

See also 48, 135, 137, 141, 149, 206.

## 33. CLINICO-CHEMICAL APPLICATIONS

33b. *Complex mixtures and profiling (single compounds by cross-reference only)*

See 40, 54, 82, 121, 145, 201.

## 34. FOOD ANALYSIS

34b. *Complex mixtures (single compounds by cross-reference only)*

See 34, 49, 115, 142, 143, 154, 179, 197.

34c. *Organoleptically important compounds (flavors, odors, volatiles)*

221 Belay, M.T. and Poole, C.F.: Determination of vanillin and related flavor compounds in natural vanilla extracts and vanilla-flavored foods by thin layer chromatography and automated multiple development. *Chromatographia*, 37 (1993) 365-373.

## 35. ENVIRONMENTAL ANALYSIS

35b. *Air pollution (complex mixtures; single compounds by cross-reference only)*

See 109.

35c. *Water pollution (complex mixtures; single compounds by cross-reference only)*

222 Yevgenyev, M.I., Yevgenyeva, I.L., Moskva, N.A., Degterev, Y.V., Dublinsky, R.A. and Tsvunin, V.S.: (Extraction chromatographic determination of N-methyl-substituted indoles in wastewater). *Khim.-Farm. Zh.*, 26, No. 7-8 (1992) 89-91; C.A., 119 (1993) 79619r.

See also 29, 32, 173, 174.

35d. *Soil pollution (complex mixtures; single compounds by cross-reference only)*

See 229.

## 36. SOME TECHNICAL PRODUCTS AND COMPLEX MIXTURES

36b. *Antioxidants and preservatives*

223 Buergi, C. and Otz, T.: Determination of preservatives in body shampoos and bubble baths. *Mitt. Geb. Lebensmittelunters. Hyg.*, 83 (1992) 492-508; C.A., 119 (1993) 102961g.

## 38. INORGANIC COMPOUNDS

38a. *Cations*

224 Kumar, A., Asolkar, A. and Pandey, P.: Extractive spectrophotometric determination of lead(II), cadmium(II), zinc(II) and vanadium(IV) with hexamethyleneiminedithiocarbamate. *Indian J. Chem., Sect. A: Inorg., Bio-Inorg., Phys., Theor. Anal. Chem.*, 32A, No. 7 (1993) 644-666; C.A., 119 (1993) 151297d.

225 Leont'eva, L.B., Tselinskii, I.V., Biochinova, E.S. and Krauklish, I.V.: (Determination of barium and strontium by paper chromatography). *Zavod. Lab.*, 59 (1993) 16-17; C.A., 119 (1993) 151234f.

226 Leont'eva, L.B., Tselinskii, I.V., Biochinova, E.S., Krauklish, I.V. and Manevakaya, R.S.: (Determination of aluminum(III) and manganese(II) in the presence of iron(III) by peak solvent paper chromatography). *Zh. Prikl. Khim. (S.-Peterburg)*, 66 (1993) 914-916; C.A., 119 (1993) 130595j.

227 Morosanova, E.I., Maksimova, I.M. and Zolotov, Yu.A.: (Immobilized nitrogen-, oxygen- and sulfur-containing macrocyclic compounds for metal ion separation by reversed-phase thin-layer chromatography). *Zh. Anal. Khim.*, 47 (1992) 1845-1863; C.A., 119 (1993) 108044g.

228 Przeszlakowski, S. and Maliszewska, M.: Retention of some metal ions on silica gel modified with Alizarin Red S. *Chem. Anal. (Warsaw)*, 37 (1992) 545-550; C.A., 119 (1993) 130505e.

229 Sanchez-Camazano, M. and Sánchez-Martín, M.J.: Mobility of cadmium as influenced by soil properties, studied by soil thin-layer chromatography. *J. Chromatogr.*, 643 (1993) 357-362.

## 39. RADIOACTIVE AND OTHER ISOTOPE COMPOUNDS

230 Chen, F., Decristoforo, C., Rohrbacher, B. and Riccabona, G.: A simple two-strip method to determine the radiochemical purity of technetium-99m mercaptoacetyltriglycine. *Eur. J. Nucl. Med.*, 20 (1993) 334-338; C.A., 119 (1993) 80181y.

231 Klein, O. and Clark, T.: The advantages of a new bio-imaging analyzer for investigation of the metabolism of <sup>14</sup>C-radiolabeled pesticides. *J. Planar Chromatogr.*, 6 (1993) 368-371.

## Electrophoresis

### 1. REVIEWS AND BOOKS

- 1 Clifton, M.J., Roux de Balman, H. and Sanchez, V.: Continuous-flow electrophoresis in microgravity. *Analisis*, 21 (1993) M32-M34; *C.A.*, 119 (1993) 134597j - a review with 10 refs.
- 2 Engelhardt, H., Beck, W., Kohr, J. and Schmitt, T.: (Capillary electrophoresis: methods and possibilities). *Angew. Chem.*, 105 (1993) 659-680; *C.A.*, 119 (1993) 151353u - a review with 161 refs.
- 3 Nokihara, K., Kuriki, T., Morita, N. and Nishine, T.: (Sample preparation by using automated apparatus for two-dimensional electrophoresis and N-terminal and internal sequence analysis). *Shimadzu Hyoron*, 50 (1993) 89-94; *C.A.*, 119 (1993) 176826p - a review with 13 refs.

See also 22, 26, 31, 72, 74, 88, 101, 109, 110, 207, 226, 243, 371, 447, 448, 456, 468, 471, 484, 495, 498, 503, 507, 528, 533, 535, 538, 541, 543, 554, 558, 562, 581, 588, 609, 620, 624.

### 2. FUNDAMENTALS, THEORY AND GENERAL

#### 2a. General

- 4 Bier, M., Ostrem, J. and Marquez, R.B.: A new buffering system and its use in electrophoresis and isoelectric focusing. *Electrophoresis (Weinheim)*, 14 (1993) 1011-1018.
- 5 Burlatsky, S. and Deutch, J.: Influence of solid friction on polymer relaxation in gel electrophoresis. *Science (Washington)*, 260 (1993) 1782-1788; *C.A.*, 119 (1993) 118219g.
- 6 Costas, M.: Microcomputers in the comparative analysis of one-dimensional electrophoretic patterns. In: Bryce, C.F.A. (Editor), *Microcomput. Biochem.*, IRL, Oxford, 1992, pp. 189-213; *C.A.*, 119 (1993) 155291v.
- 7 Garcia, L.L. and Shihabi, Z.K.: Sample matrix effects in capillary electrophoresis. I. Basic considerations. *J. Chromatogr.*, 652 (1993) 465-469.
- 8 Gaš, B.: Axial temperature effects in electromigration. *J. Chromatogr.*, 644 (1993) 161-174.
- 9 Manz, A., Verpoorte, E., Effenhauser, C.S., Burggraf, N., Raymond, D.E., Harrison, D.J. and Widmer, H.M.: Miniaturization of separation techniques using planar chip technology. *J. High Resolut. Chromatogr.*, 16 (1993) 433-436.
- 10 Putnam, F.W.: Alpha-, beta-, gamma-globulin-Arne Tiselius and the advent of electrophoresis. *Perspect. Biol. Med.*, 36 (1993) 323-337; *C.A.*, 119 (1993) 94612b.
- 11 Riibe, H.: On the use of dimensionless parameters in acid-base theory. IV. The pH of water solutions of acids, bases, and simple ampholytes. *Electrophoresis (Weinheim)*, 14 (1993) 986-992.
- 12 Shihabi, Z.K.: Sample matrix effects in capillary electrophoresis. II. Acetonitrile deproteinization. *J. Chromatogr.*, 652 (1993) 471-475.

See also 50.

#### 2b. Thermodynamics and theoretical relationships

- 13 Hayashi, Y., Matsuda, R. and Terabe, S.: Optimization of precision and throughput in micellar electrokinetic chromatography. *Chromatographia*, 37 (1993) 149-155.

See also 8, 48, 50, 53, 61, 80, 85.

#### 2d. Measurement of physico-chemical and related values

- 14 Cleveland, J.A., Jr., Benko, M.H., Gluck, S.J. and Walbroehl, Y.M.: Automated pK<sub>a</sub> determination at low solute concentrations by capillary electrophoresis. *J. Chromatogr.*, 652 (1993) 301-308.

See also 67.

### 3. GENERAL TECHNIQUES

#### 3a. Apparatus and accessories

- 15 Ehr, T.G.J., Gorski, S.H. and Vitek, R.K.: Horizontal gel electrophoresis apparatus. *Brit. UK Pat. Appl. GB 2,263,339 (Cl. GO1N27/447)*, 21 Jul. 1993, US Appl. 820,558, 14 Jan. 1992; 80 pp.; *C.A.*, 119 (1993) 155496r.
- 16 Gombocz, E., Dwight, M. and Rammler, D.H.: Gel electrophoresis sample applicator/retriever. *U.S. US 5,217,591 (Cl. 204-299R; GO1N27/26)*, 08 Jun. 1993, US Appl. 522,325, 14 May 1990; 10 pp.; *C.A.*, 119 (1993) 135009f.
- 17 Harrison, D.J., Fluri, K., Seiler, K., Fan, Z., Effenhauser, C.S. and Manz, A.: Micromachining a miniaturized capillary electrophoresis-based chemical analysis system on a chip. *Science (Washington)*, 261 (1993) 895-897; *C.A.*, 119 (1993) 194744q.
- 18 Middendorf, L.R., Bruce, J.C., Bruce, R.C., Eckles, R.D., Roemer, S.C. and Sloniker, G.D.: A versatile infrared laser scanner/electrophoresis apparatus. *Proc. SPIE-Int. Soc. Opt. Eng.*, 1885(Proceedings of Advances in Fluorescence Sensing Technology, 1993) (1993) 423-434; *C.A.*, 119 (1993) 136646z.
- 19 Verhoef, M.: High-performance capillary electrophoresis. *Am. Lab. (Shelton)*, 25 (1993) 36E; *C.A.*, 119 (1993) 173092k.
- 20 Weber, G.: Method and apparatus for continuous carrier-free deflection electrophoresis. *Ger. DE 4,139,472 (Cl. B01D57/02)*, 11 Mar. 1993, Appl. 29 Nov. 1991; 8 p.; *C.A.*, 119 (1993) 194823q.

See also 34, 38, 54, 70, 86, 87, 188, 214, 253.

#### 3b. Detectors and detection procedures

- 21 Beck, W., van Hoek, R. and Engelhardt, H.: Application of a diode-array detector in capillary electrophoresis. *Electrophoresis (Weinheim)*, 14 (1993) 540-546.



- 22 Brazhnikov, V.V.: (Detectors for chromatography. Review). *Zavod Lab.*, 59 (1993) 16-22; C.A., 119 (1993) 194730g - a review with 17 refs.
- 23 Chen, F.-T.A., Tusak, A., Pentoney, S., Jr., Konrad, K., Lew, C., Koh, E. and Sternberg, J.: Semiconductor laser-induced fluorescence detection in capillary electrophoresis using a cyanine dye. *J. Chromatogr.*, 652 (1993) 355-360.
- 24 Cifuentes, A., de Frutos, M. and Diez-Masa, J.C.: (Modification of a HPLC UV-VIS detector for use in high-voltage capillary electrophoresis). *An. Quim.*, 89 (1993) 269-274; C.A., 119 (1993) 155453z.
- 25 Deng, Y., Yu, P., Ye, M. and Liu, H.: (Laser opto-thermal online detection for gel capillary electrophoresis). *Gaodeng Xuexiao Huaxue Xuebao*, 13 (1992) 1386-1388; C.A., 119 (1993) 141162s.
- 26 Earle, C.W., Baker, M.E., Denton, M.B. and Pomeroy, R.S.: Imaging applications for chemical analysis utilizing charge coupled device array detectors. *TrAC*, 12 (1993) 395-403 - a review with 49 refs.
- 27 Hara, T., Yokogi, J., Okamura, S., Kato, S. and Nakajima, R.: On-line chemiluminescence detection of proteins separated by capillary zone electrophoresis. *J. Chromatogr.*, 652 (1993) 361-367.
- 28 Mabuchi, T.: Rapid autoradiography of sequencing gels using imaging plates. *Anal. Biochem.*, 215 (1993) 158-159.
- 29 Papavassiliou, A.G.: *In vitro* (OP)<sub>2</sub>-Cu<sup>+</sup> mapping of electrophoretically resolved RNA-protein complexes. *Anal. Biochem.*, 214 (1993) 331-334.
- 30 Westerberg, G., Lundqvist, H., Kilár, F. and Langström, B.:  $\beta^+$ -Selective radiodetector for capillary electrophoresis. *J. Chromatogr.*, 645 (1993) 319-325.
- See also 52, 63, 70, 71, 84, 85, 111, 155, 159, 201, 206, 217, 404, 431, 520, 542, 646.
- 3c. *Stabilization media for electrophoresis*
- 31 Bae, Y.C. and Soane, D.: Polymeric separation media for electrophoresis: cross-linked systems or entangled solutions. *J. Chromatogr.*, 652 (1993) 17-22 - a review with 23 refs.
- 32 Caglio, S. and Righetti, P.G.: On the efficiency of methylene blue versus persulfate catalysis of polyacrylamide gels, as investigated by capillary zone electrophoresis. *Electrophoresis (Weinheim)*, 14 (1993) 997-1003.
- 33 Chuba, J.V.: Gel plates, equipment and kits for combined electrophoretic-immunoelectrophoretic analysis. U.S. US 5,238,651 (Cl. 422-461; BO1D61/42), 24 Aug. 1993, Appl. 555,874, 23 Jul. 1990; 25 pp.; C.A., 119 (1993) 177121s.
- 34 Fukushige, T., Toda, H., Akano, H., Ito, E., Fukuta, T., Fujimori, M. and Kawamura, Y.: Electrophoretic medium, gel holder, slab type electrophoretic apparatus and electrophoretic gel cutter. *Eur. Pat. Appl. EP 555,145* (Cl. BO1D57/02), 11 Aug. 1993, JP Appl. 92/62,781, 03 Feb. 1992; 23 pp.; C.A., 119 (1993) 177117v.
- 35 Guttman, A.: Capillary column containing a dynamically cross-linked composition and method of use. U.S. US 5,213,669 (Cl. 204-180.1; GO1N27/26), 25 May 1993, Appl. 829,638, 31 Jan. 1992; 10 pp.; C.A., 119 (1993) 90797t.
- 36 Kirkpatrick, F.H., Willis, T.C., Watt, W., Daum, H.A., III, Kaura, S. and Johnson, P.A.: Electrophoresis-gel container assemblies. *PCT Int. Appl. WO 93 13,410* (Cl. GO1N27/26), 08 Jul. 1993, US Appl. 816,448, 31 Dec. 1991; 33 pp.; C.A., 119 (1993) 155518z.
- 37 Notsu, K., Ebata, N., Udagawa, A. and Shiratori, M.: Process for preparing polyacrylamide gel plate for electrophoresis. *Eur. Pat. Appl. EP 555,119* (Cl. CO8F20/56), 11 Aug. 1993, JP Appl. 92/13,174, 28 Jan. 1992; 14 pp.; C.A., 119 (1993) 177145c.
- 38 Rosenvold, E.: Integral electrophoresis gel form. U.S. US 5,232,573 (Cl. 204-299R; GO1N27/26), 03 Aug. 1993, Appl. 961,480, 15 Oct. 1992; 7 pp.; C.A., 119 (1993) 155492m.
- See also 59, 69, 535, 584.
- 3d. *Quantitative analysis*
- See 219.
- 3e. *Preparative scale electrophoresis*
- See 58, 73.
4. SPECIAL TECHNIQUES
- 4a. *Automation*
- See 19.
- 4b. *Computerization and modelling*
- 39 Keil, L.B., Barbi, G., Comizio, R. and DeBari, V.A.: Mathematical modelling of the electrophoretic distribution of IgG and the detection of clonotype restriction. *Clin. Chem. (Washington)*, 39 (1993) 1985-1987.
- 40 Solomon, J.E. and Harrington, M.G.: A robust, high-sensitivity algorithm for automated detection of proteins in two-dimensional electrophoresis gels. *Comput. Appl. Biosci.*, 9 (1993) 133-139; C.A., 119 (1993) 112836j.
- 41 Wheeler, D. and Chrambach, A.: The fractional volume available to prolate spheroids in a network of randomly oriented fibers obtained by computer modeling: Correlation with the Ogston equation. *Electrophoresis (Weinheim)*, 14 (1993) 993-996.
- 42 Yonemoto, T., Zheng, S.N., Hanai, T. and Tadaki, T.: Separation and concentration of ampholytes by recycling free-flow electrophoresis. In: Furusaki, S., Endo, I. and Matsuno, R. (Editors), *Biochem. Eng. 2001, Proc. Asia-Pac. Biochem. Eng. Conf.*, Springer, Tokyo, 1992, pp. 572-574; C.A., 119 (1993) 115363p.
- See also 6, 49, 92, 199, 495, 542.
- 4c. *Combination with other physicochemical techniques, (MS, IR etc.)*
- 43 Hara, H. and Suzuki, T.: Reactor containing capillary electrophoresis fluorescence detectors. *Jpn. Kokai Tokkyo Koho JP 05,133,898* (93,133,898) (Cl. GO1N21/73), 28 May 1993, Appl. 91/295,302, 12 Nov. 1991; 7 p.; C.A., 119 (1993) 173327r.

- 44 Varghese, J. and Cole, R.B.: Cetyltrimethylammonium chloride as a surfactant buffer additive for reversed-polarity capillary electrophoresis-electrospray mass spectrometry. *J. Chromatogr.*, 652 (1993) 369-376.
- See also 84, 179, 191, 224, 324, 622, 642.
- 4d. *Affinity electrophoresis*
- 45 Smith, N.W.: Separation of positional isomers and enantiomers using capillary zone electrophoresis with neutral and charged cyclodextrins. *J. Chromatogr.*, 652 (1993) 259-262.
- 46 Sun, P., Barker, G.E., Hartwick, R.A., Grinberg, N. and Kaliszan, R.: Chiral separations using an immobilized protein-dextran polymer network in affinity capillary electrophoresis. *J. Chromatogr.*, 652 (1993) 247-252.
- 47 Takeo, K.: (Affinity electrophoresis. Its application to analysis of antigen-antibody interactions). *Seibutsu Butsuri Kagaku*, 37 (1993) 141-146; *C.A.*, 119 (1993) 157667w.
- See also 129, 162, 448.
- 4e. *Capillary zone electrophoresis and electrokinetic chromatography*
- 48 Andreev, V.P. and Lisin, E.E.: On the mathematical model of capillary electrophoresis. *Chromatographia*, 37 (1993) 202-210.
- 49 Bello, M.S., Levin, E.I. and Righetti, P.G.: Computer-assisted determination of the inner temperature and peak correction for capillary electrophoresis. *J. Chromatogr.*, 652 (1993) 329-336.
- 50 Bello, M.S., de Besi, P. and Righetti, P.G.: Thermally induced fluctuations of the electric current and baseline in capillary electrophoresis. *J. Chromatogr.*, 652 (1993) 317-327.
- 51 Caglio, S. and Righetti, P.G.: On the pH dependence of polymerization efficiency, as investigated by capillary zone electrophoresis. *Electrophoresis (Weinheim)*, 14 (1993) 554-558.
- 52 Chen, I.-C. and Whang, C.-W.: Capillary electrophoresis with amperometric detection using a porous cellulose acetate joint. *J. Chromatogr.*, 644 (1993) 208-212.
- 53 Dose, E.V. and Guiochon, G.: Timescales of transient processes in capillary electrophoresis. *J. Chromatogr.*, 652 (1993) 263-275.
- 54 Effenhauser, C.S., Manz, A. and Widmer, H.M.: Glass chips for high-speed capillary electrophoresis separations with sub-micrometer plate heights. *Anal. Chem.*, 65 (1993) 2637-2642.
- 55 Fu, X., Lu, J. and Xia, H.: (Determination of compound drugs by micellar electrokinetic capillary chromatography). *Fenxi Ceshi Tongbao*, 11 (1992) 38-40; *C.A.*, 119 (1993) 103434z.
- 56 Fuchigami, T., Higashijima, T. and Imasaka, T.: (Trace analysis of biological substances based on capillary electrophoresis/semiconductor laser fluorometry). *Kyushu Daigaku Chuo Bunseki Senta Hokoku*, 10 (1993) 34-40; *C.A.*, 119 (1993) 155083d.
- 57 Fuchigami, T., Imasaka, T. and Shiga, M.: Ultratrace analysis of biological substances by capillary electrophoresis/semiconductor laser fluorometry. *Proc. SPIE-Int. Soc. Opt. Eng.*, 1885 (Proceedings of Advances in Fluorescence Sensing Technology, 1993) (1993) 435-438; *C.A.*, 119 (1993) 155284v.
- 58 Hathaway, G.M.: Preparative capillary electrophoresis for off-line sequence, composition, and mass analysis of peptides. *Methods (San Diego)*, 4 (1992) 244-251; *C.A.*, 119 (1993) 155279x.
- 59 Hjertén, S.: Hydrophilic polymer coating for capillary electrophoresis of high pH stability for silica surfaces for suppression of electroendosmosis and solute adsorption. *U.S. US 5,221,447* (Cl. 204-180.1; GO1N27/26), 22 Jun. 1993, Appl. 804,313, 06 Dec. 1991; 5 pp.; *C.A.*, 119 (1993) 90798u.
- 60 Hjertén, S. and Kubo, K.: A new type of pH- and detergent-stable coating for elimination of electroendosmosis and adsorption in (capillary) electrophoresis. *Electrophoresis (Weinheim)*, 14 (1993) 390-395.
- 61 Huang, T.-L., Tsai, P., Wu, C.-T. and Lee, C.S.: Mechanistic studies of electroosmotic control at the capillary-solution interface. *Anal. Chem.*, 65 (1993) 2887-2893.
- 62 Jacquier, J.C., Rony, C. and Desbene, P.L.: Computer-assisted pH optimization for the separation of geometric isomers in capillary zone electrophoresis. *J. Chromatogr.*, 652 (1993) 337-345.
- 63 Jansson, M., Roeraade, J. and Laurell, F.: Laser-induced fluorescence detection in capillary electrophoresis with blue light from a frequency-doubled diode laser. *Anal. Chem.*, 65 (1993) 2766-2769.
- 64 Jorgenson, J.W.: Capillary electrophoresis: an introduction. *Methods (San Diego)*, 4 (1992) 179-190; *C.A.*, 119 (1993) 90354w.
- 65 Kaneta, T., Tanaka, S. and Taga, M.: Effect of cetyltrimethylammonium chloride on electroosmotic and electrophoretic mobilities in capillary zone electrophoresis. *J. Chromatogr.*, 653 (1993) 313-319.
- 66 Keely, C.A., Holloway, R.R., van de Goor, T.A.A.M. and McManigill, D.: Dispersion in capillary electrophoresis with external flow control methods. *J. Chromatogr.*, 652 (1993) 283-289.
- 67 Kita, H., Kitagishi, K. and Oka, K.: (Evaluation of partition coefficients by micellar electrokinetic capillary chromatography). *Kurumatogurafi*, 13 (1992) 329-330; *C.A.*, 119 (1993) 85318h.
- 68 Klepárník, K., Šlais, K. and Boček, P.: Determination of the isoelectric points of low and high molecular mass ampholytes by capillary electrophoresis. *Electrophoresis (Weinheim)*, 14 (1993) 475-479.
- 69 Kohr, J. and Engelhardt, H.: Characterization of quartz capillaries for capillary electrophoresis. *J. Chromatogr.*, 652 (1993) 309-316.
- 70 Kok, W.T. and Sahin, Y.: Solid-state field decoupler for off-column detection in capillary electrophoresis. *Anal. Chem.*, 65 (1993) 2497-2501.
- 71 Miller, K.J. and Lytle, F.E.: Capillary zone electrophoresis with time-resolved fluorescence detection using a diode-pumped solid-state laser. *J. Chromatogr.*, 648 (1993) 245-250.
- 72 Moretto, C. and Grandi, C.: (Capillary electrophoresis: Applications and potential usefulness in clinical chemistry). *G. Ital. Chem. Clin.*, 17 (1992) 77-94; *C.A.*, 119 (1993) 134586e - a review with 13 refs.
- 73 Nashabeh, W., Smith, J.T. and El Rassi, Z.: Studies in capillary zone electrophoresis with a post-column multiple capillary device for fraction collection and stepwise increase in electroosmotic flow during analysis. *Electrophoresis (Weinheim)*, 14 (1993) 407-416.

- 74 Paulus, A., Gassmann, E. and Widner, H.M.: Capillary electrophoresis and what it can do for you. *Analysis*, 21 (1993) M36, M38, M40, M41; C.A., 119 (1993) 173249s - a review with 6 refs.
- 75 Reinhoud, N.J., Tjaden, U.R. and van der Greef, J.: Strategy for setting up single-capillary isotachopheresis-zone electrophoresis. *J. Chromatogr.*, 653 (1993) 303-312.
- 76 Smith, J.T. and El Rassi, Z.: Capillary zone electrophoresis of biological substances with fused silica capillaries having zero or constant electroosmotic flow. *Electrophoresis (Weinheim)*, 14 (1993) 396-406.
- 77 Sun, P., Wu, N., Barker, G. and Hartwick, R.A.: Chiral separations using dextran and bovine serum albumin as run buffer additives in affinity capillary electrophoresis. *J. Chromatogr.*, 648 (1993) 475-480.
- 78 Tanaka, M., Ishida, T., Araki, T., Masuyama, A., Nakatsuji, Y., Okahara, M. and Terabe, S.: Double-chain surfactant as a new and useful micelle-forming reagent for micellar electrokinetic chromatography. *J. Chromatogr.*, 648 (1993) 469-473.
- 79 Tatsumi, N.: Capillary electrophoresis system and method. *U.S. Pat. 5,240,578* (Cl. 204-180.1; GO1N27/26), 31 Aug. 1993, JP Appl. 91/242,209, 24 Sep. 1991; 8 pp.; C.A., 119 (1993) 177123u.
- 80 Taylor, J.A. and Yeung, E.S.: Imaging of hydrodynamic and electrokinetic flow profiles in capillaries. *Anal. Chem.*, 65 (1993) 2928-2932.
- 81 Van Veelen, P.A., Tjaden, U.R., van der Greef, J., Ingendoh, A. and Hillenkamp, F.: Off-line coupling of capillary electrophoresis with matrix-assisted laser desorption mass spectrometry. *J. Chromatogr.*, 647 (1993) 367-374.
- 82 Wanders, B.J., van de Goor, T.A.A.M. and Everaerts, F.M.: On-line measurement of electroosmosis in capillary electrophoresis using a conductivity cell. *J. Chromatogr.*, 652 (1993) 291-294.
- 83 Wu, C.-T., Huang, T.-L. and Lee, C.S.: Leakage current consideration of capillary electrophoresis under electroosmotic control. *J. Chromatogr.*, 652 (1993) 277-281.
- 84 Wu, J., Ma, W., Shen, X., Huang, S., Xia, L. and Rao, M.: (Study on conductivity detection and ultraviolet photometric detection for high-performance capillary electrophoresis). *Fenxi Huaxue*, 21 (1993) 727-731; C.A., 119 (1993) 151240e.
- 85 Xue, Y. and Yeung, E.S.: Double-beam laser indirect absorption detection in capillary electrophoresis. *Anal. Chem.*, 65 (1993) 2923-2927.
- 86 Zhao, Z., Malik, A. and Lee, M.L.: Solute adsorption on polymer-coated fused-silica capillary electrophoresis columns using selected protein and peptide standards. *Anal. Chem.*, 65 (1993) 2747-2752.
- 87 Zhu, T., Fang, X. and Sun, Y.: (A modular system for capillary electrophoresis). *Sepu*, 11 (1993) 242-243; C.A., 119 (1993) 194745r.
- See also 2, 7, 8, 9, 12, 13, 14, 17, 19, 21, 22, 23, 24, 25, 27, 30, 31, 32, 35, 43, 44, 45, 46, 88, 90, 96, 98, 100, 101, 102, 104, 105, 106, 107, 108, 111, 112, 114, 116, 125, 127, 128, 130, 131, 132, 134, 149, 151, 152, 153, 154, 155, 156, 157, 158, 159, 162, 164, 165, 166, 167, 168, 169, 171, 175, 176, 177, 183, 186, 188, 189, 191, 197, 202, 207, 212, 213, 215, 220, 221, 224, 225, 226, 236, 251, 279, 286, 306, 313, 317, 318, 319, 320, 324, 325, 372, 411, 436, 446, 447, 449, 450, 451, 452, 484, 486, 487, 492, 497, 501, 508, 512, 514, 516, 527, 548, 549, 550, 551, 556, 563, 564, 565, 570, 571, 572, 575, 576, 579, 584, 587, 590, 601, 602, 603, 604, 605, 606, 607, 608, 609, 610, 611, 612, 614, 615, 616, 617, 618, 619, 620, 621, 622, 623, 624, 625, 626, 627, 628, 629, 630, 631, 632, 634, 635, 636, 637, 641, 642, 643, 644, 646, 647, 648, 649.
- 4f. *Isotachopheresis*
- 88 Nielen, M.W.F.: Trace enrichment of environmental samples in capillary zone electrophoresis. *TrAC*, 12 (1993) 345-356 - a review with 42 refs.
- 89 Zusková, I., Gaš, B. and Vacík, J.: Electromigration in systems with additives in background electrolytes. II. Ionic admixture. *J. Chromatogr.*, 648 (1993) 233-244.
- 4g. *Enantiomers, separation*
- 90 Rawjee, Y.Y., Williams, R.L. and Vigh, G.: Capillary electrophoretic chiral separations using  $\beta$ -cyclodextrin as resolving agent. II. Bases: chiral selectivity as a function of pH and the concentration of  $\beta$ -cyclodextrin. *J. Chromatogr.*, 652 (1993) 233-245.
- 91 Schmitt, T. and Engelhardt, H.: Derivatized cyclodextrins for the separation of enantiomers in capillary electrophoresis. *J. High Resolut. Chromatogr.*, 16 (1993) 525-529.
- See also 77, 175, 451, 621.
- 4h. *Two dimensional electrophoresis*
- 92 Appel, R.D., Sanchez, J.-C., Bairoch, A., Golaz, O., Miu, M., Vargas, J.R. and Hochstrasser, D.F.: SWISS-2DPAGE: A database of two-dimensional gel electrophoresis images. *Electrophoresis (Weinheim)*, 14 (1993) 1232-1238.
- 93 Burstin, J., Zivy, M., de Vienne, D. and Damerval, C.: Analysis of scaling methods to minimize experimental variations in two-dimensional electrophoresis quantitative data: Application to the comparison of maize inbred lines. *Electrophoresis (Weinheim)*, 14 (1993) 1067-1073.
- See also 40, 136, 137, 190, 213, 233, 235, 263, 264, 283, 284, 292, 297, 302, 308, 321, 340, 355, 366, 438, 442, 539, 567.
- 4i. *Other special techniques*
- 94 Chebotarev, V.Yu.: (Programmed-apparatus complex for the study of biopolymers by pulse gel electrophoresis). *Prib. Tekh. Eksp.*, (1992) 178-185; C.A., 119 (1993) 90392g.

- 95 Harmon, B.J., Patterson, D.H. and Regnier, F.E.: Mathematical treatment of electrophoretically mediated microanalysis. *Anal. Chem.*, 65 (1993) 2655-2662.
- 96 Hori, A., Matsumoto, T., Nimura, Y., Ikedo, M., Okada, H. and Tsuda, T.: Electroconcentration by using countercurrent due to pressurized flow and electrophoretic mobility. *Anal. Chem.*, 65 (1993) 2882-2886.
- See also 9, 28.
6. ALCOHOLS
- 97 Atamas, S.P. and Troitsky, G.V.: Isotachophoretic separation of polyols in boric acid solutions. *J. Chromatogr.*, 644 (1993) 407-411.
7. PHENOLS
- 98 Gonnord, M.F. and Collet, J.: Optimized capillary zone electrophoretic separation of chlorinated phenols. *J. Chromatogr.*, 645 (1993) 327-336.
- 99 Tanabe, T., Yoshioka, H. and Murakami, K.: Methods for removing phenolic substances from plant proteins by gel electrophoresis. *Jpn. Kokai Tokkyo Koho* JP 05,168,476 [93,168,476] (Cl. C12N9/10), 02 Jul. 1993, Appl. 91/356,322, 24 Dec. 1991; 4 pp.; C.A., 119 (1993) 155519a.
8. SUBSTANCES CONTAINING HETEROCYCLIC OXYGEN
- 8a. Flavonoids
- 100 Bjerregaard, C., Michaelsen, S., Mortensen, K. and Sørensen, H.: Determination of flavonoids by micellar electrokinetic capillary chromatography. *J. Chromatogr.*, 652 (1993) 477-485.
- See also 634.
- 8b. Aflatoxins and other mycotoxins
- 101 Scott, P.M.: Recent developments in methods of analysis for mycotoxins in foodstuffs. *TrAC*, 12 (1993) 373-381 - a review with 37 refs.
- See also 603.
9. OXO COMPOUNDS, ETHERS, EPOXIDES AND QUINONES
- 102 Aguin, M.L., Vindevogel, J. and Sandra, P.: Utilisation of the bisulfite addition reaction for the separation of neutral aldehydes by capillary electrophoresis. *Chromatographia*, 37 (1993) 451-454.
- 103 Cherr, G.N., Fan, T.W.-M., Pillai, M.C., Shields, T. and Higashi, R.M.: Electrophoretic separation, characterization, and quantitation of biologically active lignin-derived macromolecules. *Anal. Biochem.*, 214 (1993) 521-527.
10. CARBOHYDRATES
- 10a. Mono and oligosaccharides. Structural studies
- 104 Arentoft, A.M., Michaelsen, S. and Sørensen, H.: Determination of oligosaccharides by capillary zone electrophoresis. *J. Chromatogr.*, 652 (1993) 517-524.
- 105 Bergholdt, A., Overgaard, J., Colding, A. and Frederiksen, R.B.: Separation of D-galactonic and D-gluconic acids by capillary zone electrophoresis. *J. Chromatogr.*, 644 (1993) 412-415.
- 106 Chiesa, C. and Horváth, C.: Capillary zone electrophoresis of malto-oligosaccharides derivatized with 8-aminonaphthalene-1,3,6-trisulfonic acid. *J. Chromatogr.*, 645 (1993) 337-352.
- 107 Grill, E., Huber, C., Oefner, P., Vorndran, A. and Bonn, G.: Capillary zone electrophoresis of *p*-aminobenzoic acid derivatives of aldoses, ketoses and uronic acids. *Electrophoresis (Weinheim)*, 14 (1993) 1004-1010.
- 108 Honda, S., Suzuki, S., Nitta, A., Iwase, S. and Takechi, K.: Application of high-performance capillary electrophoresis to analysis of carbohydrates, especially those in glycoproteins. *Methods (San Diego)*, 4 (1992) 233-243; C.A., 119 (1993) 176958h.
- 109 Takechi, K. and Honda, S.: Analysis of carbohydrates in glycoproteins by high-performance liquid chromatography and high-performance capillary electrophoresis. *Methods Mol. Biol. (Totowa)*, 14(Glycoprotein Analysis in Biomedicine) (1993) 81-97; C.A., 119 (1993) 90318n - a review with 11 refs.
- 110 Linhardt, R.J., Liu, J. and Han, X.J.: Mapping and sequencing of oligosaccharides by electrophoresis. *Trends Glycosci. Glyco-technol.*, 5 (1993) 181-192; C.A., 119 (1993) 117618z - a review with 74 refs.
- 111 Lu, W. and Cassidy, R.M.: Pulsed amperometric detection of carbohydrates separated by capillary electrophoresis. *Anal. Chem.*, 65 (1993) 2878-2881.
- 112 Michaelsen, S., Schröder, M.-B. and Sørensen, H.: Separation and determination of glycosaminoglycan disaccharides by micellar electrokinetic capillary chromatography for studies of pelt glycosaminoglycans. *J. Chromatogr.*, 652 (1993) 503-515.
- 113 Schaumann, C., Oesch, F., Unger, K.K. and Wieser, R.J.: Analytical technique for studying the structure of glycoprotein N-glycans. *J. Chromatogr.*, 646 (1993) 227-234.
- 114 Shirota, O., Wiesler, D., Rice, D. and Novotny, M.: (Microcolumn separations of carbohydrates with laser-induced fluorescence detection). *Kuromatografii*, 13 (1992) 229-230; C.A., 119 (1993) 90555n.
- 115 Sugo, T., Mas, E., Abouakil, N., Endo, T., Escribano, M.-J., Kobata, A. and Lombardo, D.: The structure of N-linked oligosaccharides of human pancreatic bile-salt-dependent lipase. *Eur. J. Biochem.*, 216 (1993) 799-805.
- See also 116, 125.

## 10b. Polysaccharides, mucopolysaccharides, lipopolysaccharides

- 116 Taverna, M., Baillet, A. and Baylocq-Ferrier, D.: Analysis of neutral and sialylated N-linked oligosaccharides by micellar electrokinetic capillary chromatography with addition of a divalent cation. *Chromatographia*, 37 (1993) 415-422.

See also 112.

## 10c. Glycoproteins and their constituents

- 117 Besedin, A.G., Grekhov, R.A. and Zborovsky, A.B.: (Electrophoretic separation of blood serum seromucoid glycoproteins in polyacrylamide gel). *Klin. Lab. Diagn.*, (1992) 46-48; *C.A.*, 119 (1993) 112712r.
- 118 Dam, T.K., Sarkar, M., Ghosal, J. and Choudhury, A.: Purification and partial characterization of a N-glycolylneuraminic acid specific lectin from the clam *Anadara granosa* (L.) *Biochem. Biophys. Res. Commun.*, 196 (1993) 422-429.
- 119 Ichikawa-Haraguchi, M., Sumizawa, T., Yoshimura, A., Furukawa, T., Hiramoto, S., Sugita, M. and Akiyama, S.-i.: Progesterone and its metabolites: the potent inhibitors of the transporting activity of P-glycoprotein in the adrenal gland. *Biochim. Biophys. Acta*, 1158 (1993) 201-208.
- 120 Kamemura, K. Furuichi, Y., Umekawa, H. and Takahashi, T.: Purification and characterization of novel lectins from Great Northern bean, *Phaseolus vulgaris* L. *Biochim. Biophys. Acta*, 1158 (1993) 181-188.
- 121 Kuchheuser, W. and Krause, D.: A method for simultaneous determination of group-specific component (GC) and alpha<sub>2</sub>HSG-glycoprotein (AHSG) by ultrathin-layer isoelectric focusing and immunoblotting. *Electrophoresis (Weinheim)*, 14 (1993) 1082-1083.
- 122 Malette, B. and Bleau, G.: Biochemical characterization of hamster oviductin as a sulphated zona pellucida-binding glycoprotein. *Biochem. J.*, 295 (1993) 437-445.
- 123 Moser, T.L., Enghild, J.J., Pizzo, S.V. and Stack, M.S.: The extracellular matrix proteins laminin and fibronectin contain binding domains for human plasminogen and tissue plasminogen activator. *J. Biol. Chem.*, 268 (11993) 18917-18923.
- 124 Noguchi, S. and Nakano, M.: Structural characterization of the N-linked carbohydrate chains from mouse zona pellucida glycoproteins ZP2 and ZP3. *Biochim. Biophys. Acta*, 1158 (1993) 217-226.
- 125 Novotny, M.V. and Sudor, J.: High-performance capillary electrophoresis of glycoconjugates. *Electrophoresis (Weinheim)*, 14 (1993) 373-389.
- 126 Pilarsky, C., Haase, W. and Koch-Brandt, C.: Stable expression of gp80 (TRPM-2, clusterin), a secretory protein implicated in programmed cell death, in transfected BHK-21 cells. *Biochim. Biophys. Acta*, 1179 (1993) 306-310.
- 127 Schmerr, M.J. and Goodwin, K.R.: Characterization by capillary electrophoresis of the surface glycoproteins of ovine lentiviruses before and after treatment with glycosidic enzymes. *J. Chromatogr.*, 652 (1993) 199-205.

See also 109, 113.

## 11. ORGANIC ACIDS AND LIPIDS

## 11a. Organic acids and simple esters

- 128 Brumley, W.C. and Brownrigg, C.M.: Electrophoretic behavior of aromatic-containing organic acids and the determination of selected compounds in water and soil by capillary electrophoresis. *J. Chromatogr.*, 646 (1993) 377-389.
- 129 Futschik, K., Ammann, M., Bachmayer, S. and Kenndler, E.: Determination of ionic species formed during growth of *Escherichia coli* by capillary isotachopheresis. *J. Chromatogr.*, 644 (1993) 389-395.
- 130 Levi, V., Wehr, T., Talmadge, K. and Zhu, M.: Analysis of organic acids in wines by capillary electrophoresis and HPLC. *Int. Lab.*, 23, No. 8 (1993) P4-P7.
- 131 Levi, V., Wehr, T., Talmadge, K. and Zhu, M.: Analysis of organic acids in wines by capillary electrophoresis and HPLC. *Am. Lab. (Shelton)*, 25 (1993) 29-32; *C.A.*, 119 (1993) 115621w.
- 132 Kakeda, S., Wakida, S., Yamane, M., Kawahara, A. and Higashi, K.: Migration behavior of phthalate esters in micellar electrokinetic chromatography with or without added methanol. *Anal. Chem.*, 65 (1993) 2489-2492.
- 133 Worsham, L.M.S., Williams, S.G. and Ernst-Fonberg, M.L.: Early catalytic steps of *Euglena gracilis* chloroplast type II fatty acid synthase. *Biochim. Biophys. Acta*, 1170 (1993) 62-71.

See also 626, 635.

## 11c. Lipids and their constituents

- 134 Yoo, Y.S., Kim, Y.S., Jhon, G.-J. and Park, J.: Separation of gangliosides using cyclodextrin in capillary zone electrophoresis. *J. Chromatogr.*, 652 (1993) 431-439.

## 11d. Lipoproteins and their constituents

- 135 Abe, A., Noma, A. and Itakura, H.: Lipoprotein(a) phenotyping using a computerized micro scale and phenotype frequencies in a healthy Japanese population. *Clin. Chim. Acta*, 219 (1993) 149-157.
- 136 Asztalos, B.F., Sloop, C.H., Wong, L. and Roheim, P.S.: Comparison of apo A-I-containing subpopulations of dog plasma and prenodal peripheral lymph: evidence for alteration in subpopulations in the interstitial space. *Biochim. Biophys. Acta*, 1169 (1993) 301-304.
- 137 Asztalos, B.F., Sloop, C.H., Wong, L. and Roheim, P.S.: Two-dimensional electrophoresis of plasma lipoproteins: recognition of new apo A-I-containing subpopulations. *Biochim. Biophys. Acta*, 1169 (1993) 291-300.
- 138 Blanco-Vaca, F. and Pownall, H.J.: Disulfide linked dimers of apolipoprotein D in urine. *Electrophoresis (Weinheim)*, 14 (1993) 1086-1087.
- 139 Connelly, P.W., Maguire, G.F., Vezina, C., Hegele, R.A. and Little, J.A.: Identification of disulfide-linked apolipoprotein species in human lipoproteins. *J. Lipid Res.*, 34 (1993) 1717-1727.

- 140 Coresh, J., Kwiterovich, P.O., Jr., Smith, H.H. and Bachorik, P.S.: Association of plasma triglyceride concentration and LDL particle diameter, density, and chemical composition with premature coronary artery disease in men and women. *J. Lipid Res.*, 34 (1993) 1687-1697.
- 141 Craig, W.Y., Poulin, S.E., Ledue, T.B. and Kamboh, M.I.: Apolipoprotein (a): A comparison of isoforms identified by sodium dodecyl sulfate-polyacrylamide gel electrophoresis or by sodium dodecyl sulfate-agarose gel electrophoresis. *Electrophoresis (Weinheim)*, 14 (1993) 1038-1041.
- 142 Gjøen, T. and Berg, T.: Hepatic uptake and intracellular processing of LDL in rainbow trout. *Biochim. Biophys. Acta*, 1169 (1993) 225-230.
- 143 Greeve, J., Altkemper, I., Dieterich, J.-H., Greten, H. and Windler, E.: Apolipoprotein-B mRNA editing in 12 different mammalian species: hepatic expression is reflected in low concentrations of apoB-containing plasma lipoproteins. *J. Lipid Res.*, 34 (1993) 1367-1383.
- 144 Khouw, A.S., Parthasarathy, S. and Witztum, J.L.: Radiiodination of low density lipoprotein initiates lipid peroxidation: protection by use of antioxidants. *J. Lipid Res.*, 34 (1993) 1483-1496.
- 145 Lecomte, E., Artur, Y., Chancerelle, Y., Herbeth, B., Galteau, M.-M., Jeandel, C. and Siest, G.: Malondialdehyde adducts to, and fragmentation of, apolipoprotein B from human plasma. *Clin. Chim. Acta*, 218 (1993) 39-46.
- 146 Lins, L., Piron, S., Conrath, K., Vanloo, B., Brasseur, R., Rosseneu, M., Baert, J. and Ruyschaert, J.-M.: Enzymatic hydrolysis of reconstituted dimyristoylphosphatidylcholine-apo A-I complexes. *Biochim. Biophys. Acta*, 1151 (1993) 137-142.
- 147 Lynch, S.M. and Frei, B.: Mechanism of copper- and iron-dependent oxidative modification of human low density lipoprotein. *J. Lipid Res.*, 34 (1993) 1745-1753.
- 148 Sigalov, A.B.: Comparison of apolipoprotein A-I values assayed in lyophilized and frozen pooled human sera by a non-immunochemical electrophoretic method and by immunoassay. *Eur. J. Clin. Chem. Clin. Biochem.*, 31 (1993) 579-583.
- 149 Tadey, T. and Purdy, W.C.: Effect of detergents on the electrophoretic behaviour of plasma apolipoproteins in capillary electrophoresis. *J. Chromatogr.*, 652 (1993) 131-138.
- 150 Yang, C.-y., Gu, Z.-W., Valentinova, N., Pownall, H.J., Lee, B., Yang, M., Xie, Y.-h., Guyton, J.R., Vlasik, T.N., Fruchart, J.-C. and Gotto, A.M., Jr.: Human very low density lipoprotein structure: interaction of the C apolipoproteins with apolipoprotein B-100. *J. Lipid Res.*, 34 (1993) 1311-1321.
- See also 344, 345, 585.
13. STEROIDS
- 13c. Estrogens
- 151 Potter, K.J., Allington, R.J.B. and Algaier, J.: Separation of estrogens and rodenticides using capillary electrophoresis with aqueous-methanolic buffers. *J. Chromatogr.*, 652 (1993) 427-429.
14. STEROID GLYCOSIDES AND SAPONINS
- See 632.
16. NITRO AND NITROSO COMPOUNDS
- See 648.
17. AMINES, AMIDES AND RELATED NITROGEN COMPOUNDS
- 17a. Amines and polyamines
- 152 Hori, A., Matsumoto, T., Nimura, Y. and Tsuda, T.: Capillary electrophoresis on polyamine analysis. *Int. Congr. Ser. - Excerpta Med.*, 991 (Progress in Clinical Biochemistry) (1992) 149-150; C.A., 119 (1993) 155278w.
- 153 Takeda, S., Wakida, S.-i., Yamane, M., Kawahara, A. and Higashi, K.: Separation of aniline derivatives by micellar electrokinetic chromatography. *J. Chromatogr.*, 653 (1993) 109-114.
- See also 45, 646.
- 17d. Other amine derivatives and amides (excl. peptides)
- 154 Bjerregaard, C., Ingvarsen, L. and Sørensen, H.: Determination of aromatic choline esters by micellar electrokinetic capillary chromatography. *J. Chromatogr.*, 653 (1993) 99-108.
18. AMINO ACIDS AND PEPTIDES; CHEMICAL STRUCTURE OF PROTEINS
- 18a. Amino acids and their derivatives
- 155 Arriaga, E., Chen, D.Y., Cheng, X.L. and Dovichi, J.: High-efficiency filter fluorometer for capillary electrophoresis and its application to fluorescein thiocarbonyl amino acids. *J. Chromatogr.*, 652 (1993) 347-353.
- 156 Cappiello, M., Del Corso, A., Camici, M. and Mura, U.: Thiol and disulfide determination by free zone capillary electrophoresis. *J. Biochem. Biophys. Methods*, 26 (1993) 335-341; C.A., 119 (1993) 134839q.
- 157 Chan, K.C., Manini, G.M., Muschik, G.M. and Issaq, H.J.: Laser-induced fluorescence detection of 9-fluorenylmethyl chloroformate derivatized amino acids in capillary electrophoresis. *J. Chromatogr.*, 653 (1993) 93-97.
- 158 Hernandez, L., Joshi, N., Murzi, E., Verdeguer, P., Mifsud, J.C. and Guzman, N.: Collinear laser-induced fluorescence detector for capillary electrophoresis. Analysis of glutamic acid in brain dialysates. *J. Chromatogr.*, 652 (1993) 399-405.
- 159 Hernandez, L., Tucci, S., Guzman, N. and Paez, X.: *In vivo* monitoring of glutamate in the brain by microdialysis and capillary electrophoresis with laser-induced fluorescence detection. *J. Chromatogr.*, 652 (1993) 393-398.

- 160 Kitaoko, Y., Kobayashi, M. and Ujeno, Y.: Paper electrophoretic behavior of *p*-dihydroxyborylphenylalanine in aqueous solutions of di- and tricarboxylic acids as supporting solutions. *Annu. Rep. Res. React. Inst., Kyoto Univ.*, 25 (1992) 95-99; *C.A.*, 119 (1993) 188408s.
- 161 Meyer, H.E., Eisermann, B., Heber, M., Hoffmann-Posorske, E., Korte, H., Weight, C., Wegner, A., Hutton, T., Donella-Deana, A. and Perich, J.W.: Strategies for nonradioactive methods in the localization of phosphorylated amino acids in proteins. *FASEB J.*, 7 (1993) 776-782; *C.A.*, 119 (1993) 176961d.
- 162 Otsuka, K., Kashihara, M., Kawaguchi, Y., Koike, R., Hisamitsu, T. and Terabe, S.: Optical resolution by high-performance capillary electrophoresis. Micellar electrokinetic chromatography with sodium N-dodecanoyl-L-glutamate and digitonin. *J. Chromatogr.*, 652 (1993) 253-257.
- 163 Preece, M.A., Sewell, I.J., Taylor, J.A. and Green, A.: Vigabatrin - interference with urinary amino acid analysis. *Clin. Chim. Acta*, 218 (1993) 113-116.
- See also 9, 91, 172, 400, 401, 607.
- 18b. *Peptides, peptidic and proteinous hormones, growth factors*
- 164 Adamson, N., Riley, P.F. and Reynolds, E.C.: The analysis of multiple phosphoserine-containing casein peptides using capillary zone electrophoresis. *J. Chromatogr.*, 646 (1993) 391-396.
- 165 Castagnola, M., Di Pierro, D., Scatena, R., Tavazzi, B., Nocca, G., Rossetti, D.V. and Giardina, B.: Separation of reduced and oxidized glutathione by micellar electrokinetic capillary chromatography. *Biomed. Chromatogr.*, 7 (1993) 220-226.
- 166 Chen, N., Wang, L. and Zhang, Y.: Influence of column temperature and physico-chemical properties on the electrophoretic behaviour of polyglycine peptides in free-solution capillary electrophoresis. *J. Chromatogr.*, 644 (1993) 175-182.
- 167 Chen, N., Wang, L. and Zhang, Y.K.: Correlation free-solution capillary electrophoresis migration times of small peptides with physico-chemical properties. *Chromatographia*, 37 (1993) 429-432.
- 168 Chen, N., Wang, L., Zhang, Y. and Lu, P.: (Study on the electrophoretic behavior of dipeptides in high performance capillary zone electrophoresis). *Huaxue Xuebao*, 51 (1993) 697-701; *C.A.*, 119 (1993) 176976n.
- 169 Deacon, M., O'Shea, T.J., Lunte, S.M. and Smyth, M.R.: Determination of peptides by capillary electrophoresis-electrochemical detection using on-column Cu(II) complexation. *J. Chromatogr.*, 652 (1993) 377-383.
- 170 Greenberg, Z., Chorev, M., Muhlrad, A., Shteyer, A., Namdar, M., Mansur, N. and Bab, I.: Mitogenic action of osteogenic growth peptide (OGP): role of amino and carboxy-terminal regions and charge. *Biochim. Biophys. Acta*, 1178 (1993) 273-280.
- 171 Idei, M., Mezö, I., Vadász, Zs., Horváth, A., Teplán, I. and Kéri, G.: Comparison of high-performance liquid chromatography and capillary electrophoresis in the analysis of somatostatin analogue peptides. *J. Chromatogr.*, 648 (1993) 251-256.
- 172 Janknecht, R., Ernst, W.H., Houthaeve, T. and Nordheim, A.: C-Terminal phosphorylation of the serum-response factor. *Eur. J. Biochem.*, 216 (1993) 469-475.
- 173 Johnstone, S.A., Hubashy, I. and Waisman, D.M.: Regulation of annexin I-dependent aggregation of phospholipid vesicles by protein kinase C. *Biochem. J.*, 294 (1993) 801-807.
- 174 Kim, B.G. and Brooks, C.L.: Isolation and characterization of phosphorylated bovine prolactin. *Biochem. J.*, 296 (1993) 41-47.
- 175 Kirby, D.A., Miller, C.L. and Rivier, J.E.: Separation of neuropeptide Y diastereomers by high-performance liquid chromatography and capillary zone electrophoresis. *J. Chromatogr.*, 648 (1993) 257-265.
- 176 Kristensen, H.K. and Hansen, S.H.: Micellar electrokinetic chromatographic separation of basic polypeptides with equal mass to charge ratio using dynamically modified silica capillaries. *J. Liq. Chromatogr.*, 16 (1993) 2961-2975.
- 177 Landers, J.P., Oda, R.P., Liebenow, J.A. and Spelsberg, T.C.: Utility of high resolution capillary electrophoresis for monitoring peptide homo- and hetero-dimer formation. *J. Chromatogr.*, 652 (1993) 109-117.
- 178 Lauritzen, E., Måsson, M., Rubin, I., Bjerrum, O.J. and Holm, A.: Peptide dot immunoassay and immunoblotting: Electroblotting from aluminum thin-layer chromatography plates and isoelectric focusing gels to activated nitrocellulose. *Electrophoresis (Weinheim)*, 14 (1993) 852-859.
- 179 Le Maire, M., Deschamps, S., Moller, J.V., le Caer, J.P. and Rossier, J.: Electrospray ionization mass-spectrometry on hydrophobic peptides electroeluted from sodium dodecyl sulfate-polyacrylamide gel electrophoresis application to the topology of the sarcoplasmic reticulum Ca<sup>2+</sup>-ATPase. *Anal. Biochem.*, 214 (1993) 50-57.
- 180 Måsson, M., Lauritzen, E. and Holm, A.: Chemical activation of nitrocellulose membranes for peptide antigen-antibody binding studies: Direct substitution of the nitrate group with diaminoalkane. *Electrophoresis (Weinheim)*, 14 (1993) 860-865.
- 181 Miyagishima, T., Gasa, S., Honke, K., Sakai, M., Nishi, S., Yamamoto, M., Nishikawa, K., Miyazaki, T. and Makita, A.: Expression, purification and binding to the receptor of human insulin-like growth factor II. *Biochim. Biophys. Acta*, 1203 (1993) 155-161.
- 182 Moss, A.M. and Livingston, J.N.: Distinct  $\beta$ -subunits are present in hybrid insulin-like-growth-factor-1 receptors in the central nervous system. *Biochem. J.*, 294 (1993) 685-692.
- 183 Nitta, S., Sasagawa, H., Yomoto, C. and Okada, S.: (Analysis of insulin by capillary electrophoresis). *Kuromatogurafi*, 13 (1992) 339-340; *C.A.*, 119 (1993) 103453e.
- 184 Pedroni, S., Lecomte, M.C., Gautero, H. and Dhermy, D.: Heterogeneous phosphorylation of erythrocyte spectrin  $\beta$ -chain in intact cells. *Biochem. J.*, 294 (1993) 841-846.
- 185 Quandt, N., Stindl, A. and Keller, U.: Sodium dodecyl sulfate-polyacrylamide gel electrophoresis for *M<sub>r</sub>* estimations of high-molecular-weight polypeptides. *Anal. Biochem.*, 214 (1993) 490-494.
- 186 Tsuji, K.: Evaluation of sodium dodecyl sulfate non-acrylamide, polymer gel-filled capillary electrophoresis for molecular size separation of recombinant bovine somatotropin. *J. Chromatogr.*, 652 (1993) 139-147.
- 187 Venema, R.C., Raynor, R.L., Noland, T.A., Jr. and Kuo, J.F.: Role of protein kinase C in the phosphorylation of cardiac myosin light chain 2. *Biochem. J.*, 294 (1993) 401-406.
- See also 58, 86, 122, 207, 226, 329, 394, 402, 455.

## 18c. Elucidation of structure of proteins and enzymes

- 188 Amankwa, L.N. and Kuhr, W.G.: On-line peptide mapping by capillary zone electrophoresis. *Anal. Chem.*, 65 (1993) 2693-2697.
- 189 Chang, H.T. and Yeung, E.S.: On-column digestion of protein for peptide mapping by capillary zone electrophoresis with laser-induced native fluorescence detection. *Anal. Chem.*, 65 (1993) 2947-2951.
- 190 Hirano, H., Komatsu, S., Kajiwara, H., Takagi, Y. and Tsunasawa, S.: Microsequence analysis of the N-terminally blocked proteins immobilized on polyvinylidene difluoride membrane by Western blotting. *Electrophoresis (Weinheim)*, 14 (1993) 839-846.
- 191 Izumi, T., Isobe, T., Seta, K., Mylchreest, I. and Okuyama, T.: Direct online combination of capillary electrophoresis and ESI/TSQMS for trace analysis of enzymic digests of proteins. *Kuromatogurafi*, 13 (1992) 341-342; *C.A.*, 119 (1993) 90688h.
- 192 Tachibana, H., Seki, K. and Murakami, H.: Identification of hybrid-type carbohydrate chains on the light chain of human monoclonal antibody specific to lung adenocarcinoma. *Biochim. Biophys. Acta*, 1182 (1993) 257-263.
- 193 Tozawa, K., Miyauchi, M. and Yoshida, M.: Structure of the  $\alpha$  subunit of F<sub>1</sub>-ATPase probed by limited proteolysis. *J. Biol. Chem.*, 268 (1993) 19044-19054.
- 194 Wang, F., Su, C., Hollfelder, K., Waddington, D. and Pan, Y.-C.E.: Electroblothing proteolytic products from native gel for direct N-terminal sequence analysis: An approach for studying protein-protein interaction. *Electrophoresis (Weinheim)*, 14 (1993) 847-851.
- See also 146, 200, 213, 227, 274, 282, 308, 310, 311, 325, 365, 394, 422, 435.
19. PROTEINS
- 19a. General techniques
- 195 Acevedo, F.: Isotachopheresis of proteins in sucrose density gradients. *Electrophoresis (Weinheim)*, 14 (1993) 1019-1022.
- 196 Bjellqvist, B., Hughes, G.J., Pasquali, C., Paquet, N., Ravier, F., Sanchez, J.-C., Frutiger, S. and Hochstrasser, D.: The focusing positions of polypeptides in immobilized pH gradients can be predicted from their amino acid sequences. *Electrophoresis (Weinheim)*, 14 (1993) 1023-1031.
- 197 Cifuentes, A., Santos, J.M., de Frutos, M. and Diez-Masa, J.C.: High-efficiency capillary electrophoretic separation of basic proteins using coated capillaries and cationic buffer additives. Evaluation of protein-capillary wall interactions. *J. Chromatogr.*, 652 (1993) 161-170.
- 198 Davril, M., Ducourouble, M.-P. and Van-Seuningen, I.: Western blotting of basic proteins after non-denaturing electrophoresis in acid conditions using the Phast-System. *Electrophoresis (Weinheim)*, 14 (1993) 876-880.
- 199 Dunn, M.J.: The analysis of two-dimensional polyacrylamide gels for the construction of protein databases. In: Bryce, C.F.A. (Editor), *Microcomput. Biochem.*, IRL, Oxford, 1992, pp. 215-242; *C.A.*, 119 (1993) 155292w.
- 200 Eckerskorn, C. and Lottspeich, F.: Structural characterization of blotting membranes and the influence of membrane parameters for electroblotting and subsequent amino acid sequence analysis of proteins. *Electrophoresis (Weinheim)*, 14 (1993) 831-838.
- 201 Ferreras, M., Gavilanes, J.G. and Garcia-Segura, J.M.: A permanent Zn<sup>2+</sup> reverse staining method for the detection and quantitation of proteins in polyacrylamide gels. *Anal. Biochem.*, 213 (1993) 206-212.
- 202 Foret, F., Szökő, E. and Karger, B.L.: Trace analysis of proteins by capillary zone electrophoresis with on-column transient isotachopheretic preconcentration. *Electrophoresis (Weinheim)*, 14 (1993) 417-428.
- 203 Garcia-Carreno, F.L., Dimes, L.E. and Haard, N.F.: Substrate-gel electrophoresis for composition and molecular weight of proteinases or proteinaceous proteinase inhibitors. *Anal. Biochem.*, 214 (1993) 65-69.
- 204 Hodny, Z. and Syrový, I.: The interference of cysteine thiols in the detection of glycosylated and non-glycosylated proteins by modified silver staining after sodium dodecyl sulfate-polyacrylamide gel electrophoresis. *J. Biochem. Biophys. Methods*, 26 (1993) 273-281; *C.A.*, 119 (1993) 155459f.
- 205 Hoffman, W.L., Jump, A.A., Ruggles, A.O. and Kelly, P.J.: Antibodies bound to nitrocellulose in acidic buffers retain biological activity. *Electrophoresis (Weinheim)*, 14 (1993) 886-891.
- 206 Hong, H.-Y., Choi, J.-K. and Yoo, G.-S.: Detection of proteins on polyacrylamide gels using calconcarboxylic acid. *Anal. Biochem.*, 214 (1993) 96-99.
- 207 Jumppanen, J. and Riekkola, M.L.: (Capillary electrophoresis in protein and peptide analysis). *Kem.-Kemi*, 20 (1993) 202-208; *C.A.*, 119 (1993) 134581z - a review with 89 refs.
- 208 Kao, M.C.C. and Chung, M.C.M.: Pyridylethylation of cysteine residues in proteins. *Anal. Biochem.*, 215 (1993) 82-85.
- 209 Kapfhammer, J.P.: Protein blotting analysis of large sample numbers. *Electrophoresis (Weinheim)*, 14 (1993) 881-885.
- 210 Kubo, K.: Elimination of protein band "edge tailing" in Weber-Osborn-type slab gel electrophoresis by glycerol. *Anal. Biochem.*, 213 (1993) 200-205.
- 211 Kumar, T.K.S., Gopalakrishna, K., Prasad, V.V.H. and Pandit, M.W.: Multiple bands on the sodium dodecylsulphate-polyacrylamide gel electrophoresis gels of proteins due to intermolecular disulfide cross-linking. *Anal. Biochem.*, 213 (1993) 226-228.
- 212 Kurioka, S., Tomaru, K., Sato, Y. and Kurosu, Y.: (Protein analysis by untreated fused silica capillary electrophoresis). *Seibutsu Butsuri Kagaku*, 37 (1993) 175-181; *C.A.*, 119 (1993) 112710p.
- 213 Larmann, J.P., Jr., Lemmo, A.V., Moore, A.W., Jr. and Jorgenson, J.W.: Two-dimensional separations of peptides and proteins by comprehensive liquid chromatography-capillary electrophoresis. *Electrophoresis (Weinheim)*, 14 (1993) 439-447.
- 214 Lim, Y.-P., Callanan, H., Lin, S.-H., Thompson, N.L. and Hixon, D.C.: Preparative mini-slab gel continuous elution electrophoresis: application for the separation of two isoforms of rat hepatocyte cell adhesion molecule, Cell-CAM 105, and its associated proteins. *Anal. Biochem.*, 214 (1993) 156-164.



- 215 Lin, B., Chu, X., Zhang, Y. and Xu, D.: (Modification of capillary column of electrophoresis and its application in the separation of enzymes and antibodies). *Fenxi Huaxue*, 21 (1993) 247-250; C.A., 119 (1993) 112702n.
- 216 Liu, R.Y. and Li, X.: Preparation of metabolically labeled protein samples for quantitative analysis in SDS-polyacrylamide gel electrophoresis. *Anal. Biochem.*, 214 (1993) 340-343.
- 217 Nelson, T.J.: Destaining of nitrocellulose blots after staining with silver or colloidal gold. *Anal. Biochem.*, 214 (1993) 325-328.
- 218 O'Donoghue, G.V., Okarma, T.B. and Lee, Y.M.: An immune complex selective affinity column utilizing site-specific attachment of bovine conglutinin. *Anal. Biochem.*, 213 (1993) 310-317.
- 219 Ramasamy, R., Subanesan, N. and Ramasamy, M.: Measurement of protein mobility in sodium dodecyl sulfate polyacrylamide gel electrophoresis. *J. Natl. Sci. Counc. Sri Lanka*, 20 (1992) 123-126; C.A., 119 (1993) 155282t.
- 220 Rush, R.S., McGinley, M.D., Stoney, K.S. and Rohde, M.F.: High-performance capillary electrophoresis: Protein charge estimations and peptide mapping by complexation electrophoresis. *Methods (San Diego)*, 4 (1992) 191-204; C.A., 119 (1993) 90548n.
- 221 Schmalzing, D., Piggee, C.A., Foret, F., Carrilho, E. and Karger, B.L.: Characterization and performance of a neutral hydrophilic coating for the capillary electrophoretic separation of biopolymers. *J. Chromatogr.*, 652 91993) 149-159.
- 222 Shimoni, M., Reuveni, R. and Cais, M.: Non-dialysis method of rapid and facile sample preparation for the desalting and purification of enzymes and other proteins from plant extracts. *J. Chromatogr.*, 646 (1993) 90-105.
- 223 Strege, M.A. and Lagu, A.L.: Capillary electrophoresis as a tool for the analysis of protein folding. *J. Chromatogr.*, 652 (1993) 179-188.
- 224 Wahl, J.H., Goodlett, D.R., Udseth, H.R. and Smith, R.D.: Use of small-diameter capillaries for increasing peptide and protein detection sensitivity in capillary electrophoresis-mass spectrometry. *Electrophoresis (Weinheim)*, 14 (1993) 448-457.
- 225 Wu, J. and Pawliszyn, J.: Fast analysis of proteins by isoelectric focusing performed in capillary array detected with concentration gradient imaging system. *Electrophoresis (Weinheim)*, 14 (1993) 469-474.
- 226 Zhang, H.: (Recent advance in separation and analysis of proteins and peptides by HPCE). *Yaowu Fenxi Zazhi*, 13 (1993) 62-64; C.A., 119 (1993) 90290x - a review with 17 refs.
- See also 4, 27, 40, 46, 60, 73, 86, 92, 194, 284, 352, 366.
- 19b. *Proteins of cells, viruses and subcellular particles*
- 227 Addison, R.: The initial association of a truncated form of the *Neurospora* plasma membrane H<sup>+</sup>-ATPase and of the precursor of yeast invertase with microsomes are distinct processes. *Biochim. Biophys. Acta*, 1152 (1993) 119-127.
- 228 Azuma, E.K., Kitagawa, S., Yuo, A., Mizoguchi, H., Umezawa, K., Takaku, F. and Saito, M.: Activation of the respiratory burst and tyrosine phosphorylation of proteins in human neutrophils: no direct relationship and involvement of protein kinase C-dependent and -independent signaling pathways. *Biochim. Biophys. Acta*, 1179 (1993) 2113-223.
- 229 Bagai, S. and Sarkar, D.P.: Reconstituted Sendai virus envelopes as biological carriers: dual role of F protein in binding and fusion with liver cells. *Biochim. Biophys. Acta*, 1152 91993) 15-25.
- 230 Begault, B., Anagnostopoulos, T. and Edelman, A.: ATP-regulated chloride conductance in endoplasmic reticulum (ER)-enriched pig pancreas microsomes. *Biochim. Biophys. Acta*, 1152 (1993) 319-327.
- 231 Benkirane, A., Amghar, S. and Kirchhoff, H.: Analysis of membrane proteins of *Mycoplasma capricolum* strains by SDS-PAGE and immunoblotting. *J. Vet. Med., Ser. B*, 40 (1993) 119-124; C.A., 119 (1993) 134949a.
- 232 Billings, P.B., Martinez, A., Haselby, J.A. and Hoch, S.O.: Protein blot assays specific for the discrimination of the centromere autoantigen, CENP-A, from human cells. *Electrophoresis (Weinheim)*, 14 (1993) 909-916.
- 233 Celis, J.E., Rasmussen, H.H., Olsen, E., Madsen, P., Leffers, H., Honoré, B., Dejgaard, K., Gromov, P., Hoffmann, H.J., Nielsen, M., Vassilev, A. et al.: The human keratinocyte two-dimensional gel protein database: Update 1993. *Electrophoresis (Weinheim)*, 14 (1993) 1091-1198.
- 234 Desjardins, M. and Morse, D.: The polypeptide components of scintillons, the bioluminescence organelles of the dinoflagellate *Gonyaulax polyedra*. *Biochem. Cell Biol.*, 71 (1993) 176-182.
- 235 Franzén, B., Linder, S., Okuzawa, K., Kato, H. and Auer, G.: Nonenzymatic extraction of cells from clinical tumor material for analysis of gene expression by two-dimensional polyacrylamide gel electrophoresis. *Electrophoresis (Weinheim)*, 14 (1993) 1045-1053.
- 236 Grieve, P.A., Jones, A. and Alewood, P.F.: Analytical methods for differentiating minor sequence variations in related peptides. *J. Chromatogr.*, 646 (1993) 175-184.
- 237 Hafeez, A. and Sultana, K.: Polyacrylamide gel electrophoresis of whole cell and outer membrane proteins of enteropathogenic *E. coli*. *Pak. J. Zool.*, 24 (1992) 347-351; C.A., 119 (1993) 113031e.
- 238 Hatayama, T., Asai, Y., Wakatsuki, T., Kitamura, T. and Imahara, H.: Regulation of hsp70 induction in thermotolerant HeLa cells. *Biochim. Biophys. Acta*, 1179 (1993) 109-116.
- 239 Kapaklis-Deliyannis, G.P., Drummer, H.E., Brown, L.E., Tannock, G.A. and Jackson, D.C.: A study of the advantages and limitations of immunoblotting procedures for the detection of antibodies against influenza virus. *Electrophoresis (Weinheim)*, 14 (1993) 926-936.
- 240 Kaufmann, M., Fischer, J.A. and Muff, R.: Comparison of parathyroid hormone receptors in rat osteosarcoma cells and kidney. *Biochim. Biophys. Acta*, 1179 (1993) 197-202.
- 241 Minguell, J.J., Hardy, C.L. and Tavassoli, M.: Adhesive interaction of hemopoietic progenitor cell membrane with the RGD domain of fibronectin. *Biochim. Biophys. Acta*, 1151 (1993) 120-126.
- 242 Philipson, L.H., Malayev, A., Kuznetsov, A., Chang, C. and Nelson, D.J.: Functional and biochemical characterization of the human potassium channel Kv1.5 with a transplanted carboxyl-terminal epitope in stable mammalian cell lines. *Biochim. Biophys. Acta*, 1153 (1993) 111-121.
- 243 Stewart, G.W., Argent, A.C. and Dash, B.C.J.: Stomatins: a putative cation transport regulator in the red cell membrane. *Biochim. Biophys. Acta*, 1225 (1993) 15-25 - a review with 64 refs.

- 244 Taanman, J.-W., Hall, R.E., Tang, C., Marusich, M.F., Kennaway, N.G. and Capaldi, R.A.: Tissue distribution of cytochrome c oxidase isoforms in mammals. Characterization with monoclonal and polyclonal antibodies. *Biochim. Biophys. Acta*, 1225 (1993) 95-100.
- 245 Welling-Wester, S., Feijlbrief, M., Koedijk, D.G.A.M., Braaksm, M.A., Douma, B.R.K. and Welling, G.W.: Effect of different amounts of the non-ionic detergents C<sub>10</sub>E<sub>5</sub> and C<sub>12</sub>E<sub>5</sub> present in eluents for ion-exchange high-performance liquid chromatography of integral membrane proteins of Sendai virus. *J. Chromatogr.*, 646 (1993) 37-44.
- See also 213, 308, 333, 342, 350, 424, 470, 478.
- 19c. *Proteins synthesized by genetic manipulation, monoclonal antibodies*
- 246 Kluge-Beckerman, B., Song, M., Benson, M.D. and Liepnieks, J.J.: Recombinant murine serum amyloid A from baculovirus-infected insect cells: purification and characterization. *Biochim. Biophys. Acta*, 1182 (1993) 303-310.
- 247 Lala, A., Sharma, A., Sojar, H.T., Radel, S.J., Genco, R.J. and de Nardin, E.: Recombinant expression and partial characterization of the human formyl peptide receptor. *Biochim. Biophys. Acta*, 1178 (1993) 302-306.
- 248 Lill, H., Burkovski, A., Altendorf, K., Junge, W. and Engelbrecht, S.: Complementation of *Escherichia coli unc* mutant strains by chloroplast and cyanobacterial F<sub>1</sub>-ATPase subunits. *Biochim. Biophys. Acta*, 1144 (1993) 278-284.
- 249 Pollard, P., Owen, G. and Worwood, M.: PCR-based immunogenotyping at the Ig heavy chain CDR3 locus: improvements in resolution. *Br. J. Haematol.*, 84 (1993) 169-171; C.A., 119 (1993) 136980c.
- 250 Post, A.F., Ohad, I., Warner, K.M. and Bullerjahn, G.S.: Energy distribution between Photosystems I and II in the photosynthetic prokaryote *Prochlorothrix hollandica* involves a chlorophyll a/b antenna which associates with Photosystem I. *Biochim. Biophys. Acta*, 1144 (1993) 374-384.
- 251 Yowell, G.G., Fazio, S.D. and Vivilecchia, R.V.: Analysis of a recombinant granulocyte macrophage colony stimulating factor dosage form by capillary electrophoresis, capillary isoelectric focusing and high-performance liquid chromatography. *J. Chromatogr.*, 652 (1993) 215-224.
- See also 186, 272.
- 19d. *Microbial and plant proteins*
- 252 Bae, S., Myeong, H. and Park, C.: (Analysis of CheY acetylation by urea gel electrophoresis). *Han'guk Saenghwa Hakhoechi*, 26 (1993) 279-285; C.A., 119 (1993) 155308f.
- 253 Baumann, M. and Lauraeus, M.: Purification of membrane proteins using a micropreparative gel electrophoresis apparatus: purification of subunits of the integral membrane protein *Bacillus subtilis* aag-type quinol oxidase for low level amino acid sequence analysis. *Anal. Biochem.*, 214 (1993) 142-148.
- 254 Bowlby, N.R. and Yocum, C.F.: Effects of cholate on Photosystem II: selective extraction of a 22 kDa polypeptide and modification of Q<sub>B</sub>-site activity. *Biochim. Biophys. Acta*, 1144 (1993) 271-277.
- 255 Carpenter, S.D., Ohad, I. and Vermaas, W.F.J.: Analysis of chimeric spinach/cyanobacterial CP43 mutants of *Synechocystis* sp. PCC 6803: the chlorophyll-protein CP43 affects the water-splitting system of Photosystem II. *Biochim. Biophys. Acta*, 1144 (1993) 204-212.
- 256 Cassone, A., de Bernardis, F., Pontieri, E. and Carruba, G.: Molecular biotyping of *Candida* in experimental and clinical studies. *NATO ASI Ser., Ser. H*, 69(Molecular Biology and Its Application to Medical Mycology) (1993) 159-168; C.A., 119 (1993) 155951k.
- 257 Copin, M.P., Averbeck, D. and Bourgeois, C.M.: Development of a method based on the study of the size reduction of the DNA molecules. *Comm. Eur. Communities, [Rep.] EUR EUR* 14315, 1993, pp. 357-363; C.A., 119 (1993) 137705d.
- 258 Cremer, J., Putzker, M., Faulde, M. and Zöller, L.: Immunoblotting of *Yersinia* plasmid-encoded released proteins: A tool for serodiagnosis. *Electrophoresis (Weinheim)*, 14 (1993) 952-959.
- 259 Donovan, G.R. and Baldo, B.A.: Immunoaffinity analysis of cross-reacting allergens by protein blotting. *Electrophoresis (Weinheim)*, 14 (1993) 917-922.
- 260 Faulde, M., Cremer, J. and Zöller, L.: Humoral immune response against *Helicobacter pylori* as determined by immunoblot. *Electrophoresis (Weinheim)*, 14 (1993) 945-951.
- 261 Fawley, M.W.: Structure of a prasinoxanthin-chlorophyll a/b light-harvesting complex of the green flagellate *Pseudoscofieldia marina* (Micromonadophyceae). *Biochim. Biophys. Acta*, 1183 (1993) 85-90.
- 262 Ferreras, J.M., Barbieri, L., Girbés, T., Battelli, M.G., Rojo, M.A., Arias, F.J., Rocher, M.A., Soriano, F., Mendéz, E. and Stirpe, F.: Distribution and properties of major ribosome-inactivating proteins (28 S rRNA N-glycosidases) of the plant *Saponaria officinalis* L. (Caryophyllaceae). *Biochim. Biophys. Acta*, 1216 (1993) 31-42.
- 263 Flengsrud, R.: Separation of acidic barley endosperm proteins by two-dimensional electrophoresis. *Electrophoresis (Weinheim)*, 14 (1993) 1060-1066.
- 264 Friso, G., Giacometti, G.M., Barber, J. and Barbato, R.: Evidence for concurrent donor and acceptor side photoinduced degradation of the D1-protein in isolated reaction centres of Photosystem II. *Biochim. Biophys. Acta*, 1144 (1993) 265-270.
- 265 Funke, R.P., Kolchinsky, A. and Gresshoff, P.M.: Physical mapping of a region in the soybean (*Glycine max.*) genome containing duplicated sequences. *Plant Mol. Biol.*, 22 (1993) 437-446; C.A., 119 (1993) 174925w.
- 266 Goodwin, M.G. and Jackson, J.B.: Electrochromic responses of carotenoid absorbance bands in purified light-harvesting complexes from *Rhodobacter capsulatus* reconstituted into liposomes. *Biochim. Biophys. Acta*, 1144 (1993) 191-198.
- 267 Khelifi, D., Branlard, G. and Bourgoin-Greneche, M.: Diversity of some D zone omega gliadins of bread wheat as revealed by 2 step A-PAGE/SDS-PAGE technique. *J. Genet. Breed.*, 46 (1992) 351-357; C.A., 119 (1993) 137986w.
- 268 Koxsel, H., Celik, S. and Aktan, B.: Effects of gamma irradiation on gliadin electrophoregrams of some Turkish wheat cultivars. *Turk. J. Nucl. Sci.*, 19 (1992) 39-47; C.A., 119 (1993) 179695f.
- 269 Liang, O.D., Maccarana, M., Flock, J.-I., Paulsson, M., Preissner, K.T. and Wadström, T.: Multiple interactions between human vitronectin and *Staphylococcus aureus*. *Biochim. Biophys. Acta*, 1225 (1993) 57-63.

- 270 Mendes, E., Teixeira, N.A. and Ferreira, M.A.: (Application of PAGE electrophoresis and ELISA to the detection of gluten in foods for celiac disease patients). *Rev. Port. Farm.*, 43 (1993) 32-36; *C.A.*, 119 (1993) 115634c.
- 271 Miller, K.W., Konen, P.L., Olson, J. and Ratanavanich, K.M.: Membrane protein topology determination by proteolysis of maltose binding protein fusions. *Anal. Biochem.*, 215 (1993) 118-128.
- 272 Morris, S.A., Reville, W.P., Staunton, J. and Leadlay, P.F.: Purification and separation of *holo*- and *apo*-forms of *Saccharopolyspora erythraea* acyl-carrier protein released from recombinant *Escherichia coli* by freezing and thawing. *Biochem. J.*, 294 (1993) 521-527.
- 273 Parente, A., de Luca, P., Bolognesi, A., Barbieri, L., Battelli, M.G., Abbondanza, A., Sande, M.J.W., Gigliano, G.S., Tazzari, P.L. and Stirpe, F.: Purification and partial characterization of single-chain ribosome-inactivating proteins from the seeds of *Phytolacca dioica* L. *Biochim. Biophys. Acta*, 1216 (1993) 43-49.
- 274 Qu, Z.-H. and Finkelstein, R.A.: Checkerboard immunoblotting recognizes twenty epitopes among the B subunit proteins of the cholera enterotoxin family. *Electrophoresis (Weinheim)*, 14 (1993) 899-901.
- 275 Reniero, R., Piva, A., Morelli, L., Bottazzi, V. and Cocconcelli, P.S.: Purification of *Lactobacillus* secreted proteins. *Biotechnol. Tech.*, 7 (1993) 401-406; *C.A.*, 119 (1993) 176970f.
- 276 Silverstein, T., Cheng, L. and Allen, J.F.: Redox titration of multiple protein phosphorylations in pea chloroplast thylakoids. *Biochim. Biophys. Acta*, 1183 (1993) 215-220.
- 277 Xu, J., Lu, J. and Fang, Z.: (Polyacrylamide gel disc electrophoresis of soluble proteins and esterases from mycelia of six species of *Phytophthora*). *Zhenjun Xuebao*, 12 (1993) 54-64; *C.A.*, 119 (1993) 155295z.
- 278 Zöller, L., Cremer, J. and Faulde, M.: Western blot as a tool in the diagnosis of Lyme borreliosis. *Electrophoresis (Weinheim)*, 14 (1993) 937-944.
- For additional information see *C.A.*:  
119 (1993) 110662a.
- See also 190, 231, 237, 248, 340, 640.
- 19e. *Proteins of blood, serum and blood cells*
- 279 Arentoft, A.M., Frøkiær, H., Michaelsen, S., Sørensen, H. and Sørensen, S.: High-performance capillary electrophoresis for the determination of trypsin and chymotrypsin inhibitors and their association with trypsin, chymotrypsin and monoclonal antibodies. *J. Chromatogr.*, 652 (1993) 189-198.
- 280 Cullina, M.J. and Gready, J.F.: A novel lectin-based enzyme-linked immunosorbent assay for the measurement of IgA2 in serum and secretory IgA1 in secretions. *Clin. Chim. Acta*, 216 (1993) 23-38.
- 281 Fanger, B.O., Wade, A.C., Cashman, E.A. and Cardin, A.D.: Identification of a 63-kDa serum protein that binds somatostatin and gastrin-releasing peptide but not bombesin. *Biochim. Biophys. Acta*, 1179 (1993) 300-305.
- 282 Galliano, M., Minchiotti, L., Iadarola, P., Stoppini, M., Giagnoni, P., Watkins, S., Madison, J. and Putnam, F.W.: Protein and DNA sequence analysis of a 'private' genetic variant: albumin Ortonovo (Glu-505→Lys). *Biochim. Biophys. Acta*, 1225 (1993) 27-32.
- 283 Golaz, O., Hughes, G.J., Frutiger, S., Paquet, N., Bairoch, A., Pasquali, C., Sanchez, J.-C., Tissot, J.-D., Appel, R.D., Walzer, C. et al.: Plasma and red blood cell protein maps: Update 1993. *Electrophoresis (Weinheim)*, 14 (1993) 1223-1231.
- 284 Gulle, H., Kaufmann, S.H.E. and Moriarty, K.M.: Rapid electroelution of two-dimensionally separated protein mixtures: Its use in *in vitro* assays of T cell activities. *Electrophoresis (Weinheim)*, 14 (1993) 902-908.
- 285 Jeppsson, J.-O., Kristensson, H. and Fimiani, C.: Carbohydrate-deficient transferrin quantified by HPLC to determine heavy consumption of alcohol. *Clin. Chem. (Washington)*, 39 (1993) 2115-2120.
- 286 Liu, C.M., Wang, H.P., Chen, F.T.A., Sternberg, J.C. and Klein, G.L.: Detection of monoclonal gammopathy by capillary electrophoretic immunosubtraction (CEI). *U.S. US 5,228,960* (Cl. 204-182.8; C25B7/00), 20 Jul. 1993, Appl. 916,313, 17 Jul. 1992; 15 pp.; *C.A.*, 119 (1993) 155517y.
- 287 Lof, K., Koivula, T., Seppä, K., Fukunaga, T. and Sillanaukee, P.: Semi-automatic method for determination of different isoforms of carbohydrate-deficient transferrin. *Clin. Chim. Acta*, 217 (1993) 175-186.
- 288 Lubec, B., Weninger, M., Popow, C., Vierhapper, H., Lunec, J. and Lubec, G.: *In vivo* monitoring of serum protein cross linking in patients with diabetes mellitus. Evidence for pharmacological modification of immunoglobulin G cross links. *Amino Acids*, 4 (1993) 111-119; *C.A.*, 119 (1993) 90743x.
- 289 Lütkemeyer, D., Bretschneider, M., Büntemeyer, H. and Lehmann, J.: Membrane chromatography for rapid purification of recombinant antithrombin III and monoclonal antibodies from cell culture supernatant. *J. Chromatogr.*, 639 (1993) 57-66.
- 290 Nagata, K.-i. and Nozawa, Y.: Subcellular localization of a low  $M_r$  GTP-binding protein, c25KG, in resting and stimulated human platelets. *Biochem. Biophys. Res. Commun.*, 195 (1993) 1081-1088.
- 291 Shainoff, J.R., Urbanic, D.A., DiBello, P.M. and Valenzuela, V.: GPR-phoresis, a novel approach to determining fibrin monomer and other macromolecular derivatives of fibrinogen and fibrin in blood. *Blood Coagulation Fibrinolysis*, 4 (1993) 87-92; *C.A.*, 119 (1993) 90553k.
- 292 Stulík, J., Tichý, M. and Kovářová, H.: Two-dimensional gel electrophoresis of four serum samples from patients with IgD myeloma. *Clin. Chim. Acta*, 218 (1993) 149-158.
- 293 Szilágyi, A.K., Ha, N.T., Szelényi, J. and Puskás, E.: Differences in the isoelectric focusing patterns of serum and cerebrospinal fluid transthyretin. *Electrophoresis (Weinheim)*, 14 (1993) 1079-1081.
- 294 Uda, S.: Electrophoresis for identifying class of monoclonal immunoglobulins (M proteins). *Jpn. Kokai Tokkyo Koho JP 05 87,813* [93 87,813] (Cl. G01N33/561), 06 Apr. 1993, Appl. 91/173,510, 15 Apr. 1991; 5 pp.; *C.A.*, 119 (1993) 177144b.
- 295 Vostal, J.G. and Shulman, N.R.: Vinculin is a major platelet protein that undergoes  $Ca^{2+}$ -dependent tyrosine phosphorylation. *Biochem. J.*, 294 (1993) 675-680.
- 296 Zimmermann, C.W.: Repertoires of natural autoantibodies against muscle tissue are independent of age or gender in normal human adults. A Western blot study. *Clin. Chim. Acta*, 218 (1993) 29-38.
- See also 10, 39, 135, 205, 228, 243, 246, 278, 352, 412, 494.

19f. *Structural and muscle proteins*

- 297 Bárány, M. and Bárány, K.: Calponin phosphorylation does not accompany contraction of various smooth muscles. *Biochim. Biophys. Acta*, 1179 (1993) 229-233.
- 298 Fabbrizio, E., Bonet-Kerrache, A., Leger, J.J. and Mornet, D.: Actin-dystrophin interface. *Biochemistry*, 32 (1993) 10457-10463.
- 299 Lee, S. and Cote, G.P.: Isolation and characterization of three *Dictyostelium* myosin-I isozymes. *J. Biol. Chem.*, 268 (1993) 20923-20929.
- 300 Lehman, W., Denault, D. and Marston, S.: The caldesmon content of vertebrate smooth muscle. *Biochim. Biophys. Acta*, 1203 (1993) 53-59.
- 301 Lemos, Ana Lucia da Silva Corres, Moraes, M.A.C.: Electrophoretic identification of commercial river fish (family Pimelodidae). *Aliment. Nutr.*, 4 (1992) 57-63; *C.A.*, 119 (1993) 137722g.
- 302 Nagai, A., Yamada, S., Bunai, Y. and Ohya, I.: (Two-dimensional gel electrophoretic patterns of human and animal hair proteins). *Hanzaigaku Zasshi*, 59 (1993) 49-54; *C.A.*, 119 (1993) 153453a.
- 303 Valenta, C. and Gabor, F.: (Quality control of collagens: SDS-PAGE). *Sci. Pharm.*, 60 (1992) 267-272; *C.A.*, 119 (1993) 188241g.
- 304 Zavlin, P.M., Izmailova, V.N., Musaelyan, G.S., Sakvarelidze, M.A. and Yampol'skaya, G.P.: (Use of SDS gel electrophoresis for analysis of modification of gelatine). *Zh. Prikl. Khim. (S.-Peterburg)*, 66 (1993) 622-626; *C.A.*, 119 (1993) 183227x.

See also 400, 465, 482.

19g. *Protamines, histones and other nuclear proteins*

- 305 Chirat, F., Arkhis, A., Martinage, A., Jaquinod, M., Chevaillier, P. and Sautière, P.: Phosphorylation of human sperm protamines HP1 and HP2: identification of phosphorylation sites. *Biochim. Biophys. Acta*, 1203 (1993) 109-114.
- 306 Lindner, H., Wurm, M., Dirschlmayer, A., Sarg, B. and Helliger, W.: Application of high-performance capillary electrophoresis to the analysis of H1 histones. *Electrophoresis (Weinheim)*, 14 (1993) 480-485.
- 307 Reeves, R. and Nissen, M.S.: Interaction of high mobility group-I(Y) nonhistone proteins with nucleosome core particles. *J. Biol. Chem.*, 268 (1993) 21137-21146.
- 308 Wirth, P.J., Luo, L.-d., Benjamin, T., Hoang, T.N., Olson, A.D. and Parmelee, D.C.: The rat liver epithelial (RLE) cell nuclear protein database. *Electrophoresis (Weinheim)*, 14 (1993) 1199-1215.
- 309 Zagariya, A., Khrapunov, S. and Zacharias, W.: Rapid method for the fractionation of nuclear proteins and their complexes by batch elution from hydroxyapatite. *J. Chromatogr.*, 648 (1993) 275-278.

See also 232.

19h. *Chromoproteins and metalloproteins*

- 310 Apostolova, M., Bontchev, P.R., Nachev, C. and Sirakova, I.: Apometallothionein in rat liver. *J. Chromatogr.*, 620 (1993) 191-197.

- 311 Demuyneck, S., Li, K.W., van der Schors, R. and Dhainaut-Courtois, N.: Amino acid sequence of the small cadmium-binding protein (MP1) from *Nereis diversicolor* (annelida, polychaeta). Evidence for a myohemerythrin structure. *Eur. J. Biochem.*, 217 (1993) 151-156.
- 312 Pantaleso, Silmara de M., de Medeiros, F., Joao G., Numesmaia, Henrique G. da S. and Vieira, J.: (Screening for structural hemoglobinopathies in newborns from Joao Pessoa - PB). *Rev. Bras. Patol. Clin.*, 29 (1993) 8-13; *C.A.*, 119 (1993) 112706s.
- 313 Richards, M.P. and Beattie, J.H.: Characterization of metallothionein isoforms. Comparison of capillary zone electrophoresis with reversed-phase high-performance liquid chromatography. *J. Chromatogr.*, 648 (1993) 459-468.
- 314 Shen, J. and Inoue, Y.: Cellular localization of cytochrome c550. Its specific association with cyanobacterial Photosystem II. *J. Biol. Chem.*, 268 (1993) 20408-20413.
- 315 Sonoda, Y., Endo, M., Ishida, K., Sato, Y., Fukusen, N. and Fukuhara, M.: Purification of a human cytochrome P-450 isozyme catalyzing lanosterol 14 $\alpha$ -demethylation. *Biochim. Biophys. Acta*, 1170 (1993) 92-97.
- 316 Tamegai, H., Yamanaka, T. and Fukumori, Y.: Purification and properties of a 'cytochrome a<sub>1</sub>'-like hemoprotein from a magnetotactic bacterium, *Aquaspirillum magnetotacticum*. *Biochim. Biophys. Acta*, 1158 (1993) 237-243.
- 317 Wu, J. and Pawliszyn, J.: *In vitro* observation of interactions of iron and transferrin by capillary isoelectric focusing with a concentration gradient imaging detection system. *J. Chromatogr.*, 652 (1993) 295-299.
- 318 Zhu, M., Wehr, T., Levi, V., Rodriguez, R., Shiffer, K. and Cao, Z.A.: Capillary electrophoresis of abnormal hemoglobins associated with  $\alpha$ -thalassemias. *J. Chromatogr.*, 652 (1993) 119-129.
- 19i. *Proteins of glands, gland products, various zymogens (incl. milk proteins)*
- 319 Cifuentes, A., de Frutos, M. and Diez-Masa, J.C.: Analysis of whey proteins by capillary electrophoresis using buffer-containing polymeric additives. *J. Dairy Sci.*, 76 (1993) 1870-1875; *C.A.*, 119 (1993) 158533m.
- 320 De Jong, N., Visser, S. and Olieman, C.: Determination of milk proteins by capillary electrophoresis. *J. Chromatogr.*, 652 (1993) 207-213.
- 321 Hughes, G.J., Frutiger, S., Paquet, N., Pasquali, C., Sanchez, J.-C., Tissot, J.-D., Bairoch, A., Appel, R.D. and Hochstrasser, D.F.: Human liver protein map: Update 1993. *Electrophoresis (Weinheim)*, 14 (1993) 1216-1222.
- 322 Mori, E., Baba, T., Iwamatsu, A. and Mori, T.: Purification and characterization of a 38-kDa protein, sp38, with zona pellucida-binding property from porcine epididymal sperm. *Biochem. Biophys. Res. Commun.*, 196 (1993) 196-202.
- 323 Moufquie, J., Rothhut, B., Rainteau, D., Comera, C., Alfsen, A., Russo-Marie, F. and Lavialle, F.: Annexins from bovine adrenal cortex exhibit specific cytosol/membrane solvation. *Biochem. Biophys. Res. Commun.*, 195 (1993) 132-138.
- 324 Perkins, J.R., Parker, C.E. and Tomer, K.B.: The characterization of snake venoms using capillary electrophoresis in conjunction with electrospray mass spectrometry: Black Mambas. *Electrophoresis (Weinheim)*, 14 (1993) 458-468.

- 325 Vita, C., Bontems, F., Bouet, F., Tauc, M., Poujeol, P., Vatanpour, H., Harvey, A.L., Menez, A. and Toma, F.: Synthesis of charybdotoxin and of two N-terminal truncated analogues. Structural and functional characteristics. *Eur. J. Biochem.*, 217 (1993) 157-169.
- 326 Yoo, S.H.: pH-Dependent binding of chromogranin B and secretory vesicle matrix proteins to the vesicle membrane. *Biochim. Biophys. Acta*, 1179 (1993) 239-246.
- 327 Zhang, X., Duggan, L.J.L. and Gottlieb, P.A.: Loading dyes used in gel electrophoresis alter the apparent thermodynamic equilibrium of the lac repressor-operator complex. *Anal. Biochem.*, 214 (1993) 580-582.
- See also 29, 164, 308, 330, 337, 342, 475.
- 19j. *Proteins of brain, cerebrospinal fluid and eye*
- 328 Beretta, L., Dobránský, T. and Sobel, A.: Multiple phosphorylation of stathmin. Identification of four sites phosphorylated in intact cells and *in vitro* by cyclic AMP-dependent protein kinase and p34cdc2. *J. Biol. Chem.*, 268 (1993) 20076-20084.
- 329 Groenen, P.J.T.A., Seccia, M., Smulders, R.H.P.H., Gravela, E., Cheeseman, K.H., Bloemendal, H. and de Jong, W.W.: Exposure of  $\beta$ H-crystallin to hydroxyl radicals enhances the transglutaminase-susceptibility of its existing amine-donor and amino-acceptor sites. *Biochem. J.*, 295 (1993) 399-404.
- 330 Metz-Boutigue, M.-H., Garcia-Sablone, P., Hogue-Angeletti, R. and Aunis, D.: Intracellular and extracellular processing of chromogranin A. Determination of cleavage sites. *Eur. J. Biochem.*, 217 (1993) 247-257.
- 331 Ohba, S., Kamata, K. and Miki-Noumura, T.: Stabilization of microtubules by dynein-binding *in vitro*. Stability of microtubule-dynein complex. *Biochim. Biophys. Acta*, 1158 (1993) 323-332.
- 332 Riley, M.L. and Harding, J.J.: The reaction of malondialdehyde with lens proteins and the protective effect of aspirin. *Biochim. Biophys. Acta*, 1158 (1993) 107-112.
- 333 Tomarev, S.I., Zinovieva, R.D. and Piatigorsky, J.: Primary structure and lens-specific expression of genes for an intermediate filament protein and a  $\beta$ -tubulin in cephalopods. *Biochim. Biophys. Acta*, 1216 (1993) 245-254.
- See also 293, 385, 400, 438, 469.
- 19k. *Proteins of neoplastic tissue and transformed cells*
- See 235.
- 19l. *Specific binding and receptor proteins*
- 334 Alla, S.A., Buschko, J., Quitterer, U., Maidhof, A., Haasemann, M., Breipohl, G., Knolle, J. and Müller-Ester, W.: Structural features of the human bradykinin B2 receptor probed by agonists, antagonists, and anti-idiotypic antibodies. *J. Biol. Chem.*, 268 (1993) 17277-17285.
- 335 Bielinski, D.F., Pyun, H.Y., Linko-Stentz, K., Macara, I.G. and Fine, R.E.: Ral and Rab3a are major GTP-binding proteins of axonal rapid transport and synaptic vesicles and do not redistribute following depolarization stimulated synaptosomal exocytosis. *Biochim. Biophys. Acta*, 1151 (1993) 246-256.
- 336 Brady, M.J. and Palfrey, H.C.: Rapid and sustained phosphorylation of a calmodulin-binding protein (CaM-BP<sub>100</sub>) in NGF-treated PC12 cells. *J. Biol. Chem.*, 268 (1993) 17951-17958.
- 337 Ehrlich, K.C.: Partial purification of a pea seed DNA-binding protein that specifically recognizes 5-methylcytosine. *Prepar. Biochem.*, 23 (1993) 423-438.
- 338 Friedman, E., Butkerait, P. and Wang, H.-Y.: Analysis of receptor-stimulated and basal guanine nucleotide binding to membrane G proteins by sodium dodecyl sulfate-polyacrylamide gel electrophoresis. *Anal. Biochem.*, 214 (1993) 171-178.
- 339 Grosbois, M., Guerbette, F., Jolliot, A., Quintin, F. and Kader, J.-C.: Control of maize lipid transfer protein activity by oxidoreducing conditions. *Biochim. Biophys. Acta*, 1170 (1993) 197-203.
- 340 Hanlon, D.W., Ying, C. and Ordal, G.W.: Purification and reconstitution of the methyl-accepting chemotaxis proteins from *Bacillus subtilis*. *Biochim. Biophys. Acta*, 1158 (1993) 345-351.
- 341 Heinrichs, A.A.J., Bortell, R., Rahman, S., Stein, J.L., Slnemri, E.S., Litwack, G., Lian, J.B. and Stein, G.S.: Identification of multiple glucocorticoid receptor binding sites in the rat osteocalcin gene promoter. *Biochemistry*, 32 (1993) 11436-11444.
- 342 Iwaki, M., Kumagai, S., Akamatsu, Y. and Kitagawa, T.: Aflatoxin B<sub>1</sub>-binding proteins in primary cultured hepatocytes of chicken embryo: studies *in vivo* and *in vitro*. *Biochim. Biophys. Acta*, 1225 (1993) 83-88.
- 343 Jochen, A. and Hays, J.: Purification of the major substrate for palmitoylation in rat adipocytes: N-terminal homology with CD 36 and evidence for cell surface acylation. *J. Lipid Res.*, 34 (1993) 1783-1792.
- 344 Kushwaha, R.S., Hasan, S.Q., McGill, H.C., Jr., Getz, G.S., Dunham, R.G. and Kanda, P.: Characterization of cholesterol transfer protein inhibitor from plasma of baboons (*Papio* sp.). *J. Lipid Res.*, 34 (1993) 1285-1297.
- 345 Lelli, N., Garuti, R., Zambelli, F., Cassanelli, S., Tiozzo, R., Corsini, A., Bertolini, S., Riva, E., Ortisi, M.T., Bellu, R. and Calandra, S.: Alternative splicing of mutant LDL-receptor mRNA in an Italian patient with familial hypercholesterolemia due to a partial deletion of LDL-receptor gene (FH<sup>Potenza</sup>). *J. Lipid Res.*, 34 (1993) 1347-1354.
- 346 Liu, J., Han, J.R., Liu, C.-C., Suiko, M. and Liu, M.-C.: Identification of a putative tyrosine-O-sulphate (TyrS) receptor possibly functioning in the biosynthetic transport of tyrosine-sulphated proteins in Madin-Darby canine kidney cells. *Biochem. J.*, 294 (1993) 407-417.
- 347 Mathews, L.S. and Vale, W.W.: Characterization of type II activin receptors. Binding, processing, and phosphorylation. *J. Biol. Chem.*, 268 (1993) 19013-19018.
- 348 Melki, S.A. and Abumrad, N.A.: Expression of the adipocyte fatty acid-binding protein in streptozotocin-diabetes: effects of insulin deficiency and supplementation. *J. Lipid Res.*, 34 (1993) 1527-1534.
- 349 Memon, A.R., Herrin, D.L. and Thompson, G.A., Jr.: Intracellular translocation of a 28 kDa GTP-binding protein during osmotic shock-induced cell volume regulation in *Dunaliella salina*. *Biochim. Biophys. Acta*, 1179 (1993) 11-22.
- 350 Milligan, G., Mullaney, I. and McCallum, J.F.: Distribution and relative levels of expression of the phosphoinositidase-C-linked G-proteins G<sub>q</sub> and G<sub>11</sub> in human platelets and haemopoietically derived cell lines. *Biochim. Biophys. Acta*, 1179 (1993) 208-212.

- 351 Nielsen, S.U. and Spener, F.: Fatty acid-binding protein from rat heart is phosphorylated on Tyr<sup>19</sup> in response to insulin stimulation. *J. Lipid Res.*, 34 (1993) 1355-1366.
- 352 Pan, J., Pfeiffer, D.C., Reimer, C.L., Crawford, B.J., Vogl, A.W. and Auersperg, N.: Lack of inter-species reactivity between antigens and antibodies is overcome by protease treatment of Western blots. *Electrophoresis (Weinheim)*, 14 (1993) 892-898.
- 353 Pick, E., Gorzalczy, Y. and Engel, S.: Role of the rac1 p21-GDP-dissociation inhibitor for rho heterodimer in the activation of the superoxide-forming NADPH oxidase of macrophages. *Eur. J. Biochem.*, 217 (1993) 441-455.
- 354 Ramkumar, V., Kwatra, M., Benovic, J.L. and Stiles, G.L.: Functional consequences of A<sub>1</sub> adenosine-receptor phosphorylation by the  $\beta$ -adrenergic receptor kinase. *Biochim. Biophys. Acta*, 1179 (1993) 89-97.
- 355 Sabbatini, G.P., Smith, P.J. and von Holt, C.: A 46 kDa NTPase common to rat liver nuclear envelope, mitochondria, plasma membrane, and endoplasmic reticulum. *Biochim. Biophys. Acta*, 1153 (1993) 132-134.
- 356 Shinohara, Y., Kamida, M., Yamazaki, N. and Terada, H.: Isolation and characterization of cDNA clones and a genomic clone encoding rat mitochondrial adenine nucleotide translocator. *Biochim. Biophys. Acta*, 1152 (1993) 192-196.
- 357 Smith, D.F., Baggenstoss, B.A., Marion, T.N. and Rimerman, R.A.: Two FKBP-related proteins are associated with progesterone receptor complexes. *J. Biol. Chem.*, 268 (1993) 18365-18371.
- 358 Srivastava, R.A.K., Baumann, D. and Schonfeld, G.: *In vivo* regulation of low-density lipoprotein receptors by estrogen differs at the post-transcriptional level in rat and mouse. *Eur. J. Biochem.*, 216 (1993) 527-538.
- 359 Vieira, A.V. and Schneider, W.J.: Transport and uptake of retinol during chicken oocyte growth. *Biochim. Biophys. Acta*, 1169 (1993) 250-256.

See also 29, 180, 240, 247, 281, 368, 455, 505.

#### 19m. Urinary proteins

- 360 Aguzzi, F., Gasparro, C., Bergami, M.R. and Merlini, M.: High-sensitivity electrophoretic method for the detection of Bence Jones protein and for the study of proteinuria in unconcentrated urines. *Ann. Clin. Biochem.*, 30 (1993) 287-292; *C.A.*, 119 (1993) 155304b.
- 361 Kolaja, G.J., VanderMeer, D.A., Packwood, W.H. and Satoh, P.S.: The use of sodium dodecyl sulfate-polyacrylamide gel electrophoresis to detect renal damage in Sprague-Dawley rats treated with gentamicin sulfate. *Toxicol. Pathol.*, 20 (1992) 603-607; *C.A.*, 119 (1993) 110676h.
- 362 Marshall, T. and Williams, K.M.: Sodium dodecyl sulfate-polyacrylamide gel electrophoresis of urine: concentration of urinary proteins by precipitation with Coomassie Blue. *Clin. Chem. (Washington)*, 39 (1993) 2314-2318.
- 363 Petrini, C., Rizza, V., Bezzi, C., Sabadini, E. and D'Amico, G.: Comparison of SDS-PAGE and immunoblotting with polyclonal and monoclonal antihuman albumin sera for the detection of urinary polymers of albumin. *Contrib. Nephrol.*, 101 (Kidney, Proteins and Drugs) (1993) 158-163; *C.A.*, 119 (1993) 134947y.

- 364 Trewick, A.L., Luxton, F., Patel, S. and Heys, A.D.: Urine protein electrophoresis without sample concentration: an evaluation of two methods. *Ann. Clin. Biochem.*, 30 (1993) 321-323; *C.A.*, 119 (1993) 155447a.

See also 138.

#### 19n. Other proteins (incl. proteinous inhibitors of enzymic activity)

- 365 Alagiri, S. and Singh, T.P.: Stability and kinetics of a bifunctional amylase/trypsin inhibitor. *Biochim. Biophys. Acta*, 1203 (1993) 77-84.
- 366 Collet, J. and Baguñá, J.: Optimizing a method of protein extraction for two-dimensional electrophoretic separation of proteins from planarians (*Platyhelminthes, Turbellaria*). *Electrophoresis (Weinheim)*, 14 (1993) 1054-1059.
- 367 O'Neill, G.M. and Baldo, B.A.: Intra-species cross-reactivity of house dust mite allergens separated by protein blotting and detected by selective elution of mite components and IgE antibodies. *Electrophoresis (Weinheim)*, 14 (1993) 923-925.
- 368 Silverman, M., Speight, P. and Ho, L.: Identification of two unique polypeptides from dog kidney outer cortex and outer medulla that exhibit different Na<sup>+</sup>/D-glucose cotransport functional properties. *Biochim. Biophys. Acta*, 1153 (1993) 43-52.
- 369 Sitizyo, K., Takeuchi, T., Suzuki, M. and Saito, T.: (Electrophoretic analysis of urinary proteins in domestic animals with acid violet 17 staining). *Seibutsu Butsuri Kagaku*, 37 (1993) 183-188; *C.A.*, 119 (1993) 112711q.
- 370 Zhao, J. and Benbow, R.M.: An inhibitor of DNA topoisomerase I from *Xenopus laevis* ovaries. *Biochemistry*, 32 (1993) 10622-10628.

See also 203, 344.

#### 20. ENZYMES AND ENZYME ACTIVITY ESTIMATION

- 371 Gabriel, O. and Gersten, D.M.: Detection of enzymic activity after polyacrylamide gel electrophoresis and agarose gel isoelectric focusing. In: Eisenthal, R. and Danson, M.J. (Editors), *Enzyme Assays*, IRL, Oxford, 1992, pp. 217-253; *C.A.*, 119 (1993) 111779f - a review with 83 refs.
- 372 Konse, T., Takahashi, T., Nagashima, H. and Iwaoka, T.: Blotting membrane micropreparation in capillary electrophoresis to evaluate enzyme purity and activity. *Anal. Biochem.*, 214 (1993) 179-181.

See also 198, 215, 222, 431.

#### 20a. Oxidoreductases

- 373 Bjorkhem, I., Andersson, U., Sudjama-Sugiaman, E., Eggertsen, G. and Hylemon, P.: Studies on the link between HMG-CoA reductase and cholesterol 7 $\alpha$ -hydroxylase in lymph-fistula rats: evidence for both transcriptional and post-transcriptional of the two enzymes by bile acids. *J. Lipid Res.*, 34 (1993) 1497-1503.

- 374 Bross, P., Andresen, B.S., Winter, V., Kräutle, F., Jensen, T.G., Nandy, A., Klvraa, S., Ghisla, S., Bolund, L. and Gregersen, N.: Co-overexpression of bacterial GroESL chaperonins partly overcomes non-productive folding and tetramer assembly of *E. coli*-expressed human medium-chain acyl-CoA dehydrogenase (MCAD) carrying the prevalent disease-causing K304E mutation. *Biochim. Biophys. Acta*, 1182 (1993) 264-274.
- 375 Consalvi, V., Chiaraluca, R., Politi, L., Pasquo, A., de Rosa, M. and Scandurra, R.: Glutamate dehydrogenase from the thermoacidophilic archaeobacterium *Sulfolobus solfataricus*: studies on thermal and guanidine-dependent inactivation. *Biochim. Biophys. Acta*, 1202 (1993) 207-215.
- 376 Hayes, J.D., Judah, D.J. and Neal, G.E.: Resistance to aflatoxin B<sub>1</sub> is associated with the expression of a novel aldo-keto reductase which has catalytic activity towards a cytotoxic aldehyde-containing metabolite of the toxin. *Cancer Res.*, 53 (1993) 3887-3894.
- 377 Jimenez-Cervantes, C., Garcia-Borron, J., Valverde, P., Solano, F. and Lozano, J.A.: Tyrosinase isoenzymes in mammalian melanocytes. 1. Biochemical characterization of two melanosomal tyrosinases from B16 mouse melanoma. *Eur. J. Biochem.*, 217 (1993) 549-556.
- 378 Jones, A.M. and Hollocher, T.C.: Nitric oxide reductase of *Achromobacter cycloclastes*. *Biochim. Biophys. Acta*, 1144 (1993) 359-366.
- 379 Keefe, R.G. and Maier, R.J.: Purification and characterization of an O<sub>2</sub>-utilizing cytochrome-c oxidase complex from *Bradyrhizobium japonicum* bacteroid membranes. *Biochim. Biophys. Acta*, 1183 (1993) 91-104.
- 380 Li, Z., Szittner, R. and Meighen, E.A.: Subunit interactions and the role of the luxA polypeptide in controlling thermal stability and catalytic properties in recombinant luciferase hybrids. *Biochim. Biophys. Acta*, 1158 (1993) 137-145.
- 381 Magradze, D.N., Kerkadze, I.G., Kvaliashvili, V.R. and Vashakidze, L.K.: Electrophoretic identification of dehydrogenase activities in peach leaves. *U.S.S.R. SU 1,788,969 (Cl. C12Q/32)*, 15 Jan. 1993, Appl. 4,923,273, 01 Apr. 1991; *C.A.*, 119 (1993) 134382k.
- 382 Maly, I.P. and Toranelli, M.: Ultrathin-layer zone electrophoresis of lactate dehydrogenase isoenzymes in microdissected liver samples. *Anal. Biochem.*, 214 (1993) 379-388.
- 383 Ohishi, N., Imaoka, S., Suzuki, T. and Funae, Y.: Characterization of two P-450 isozymes placed in the rat CYP2D subfamily. *Biochim. Biophys. Acta*, 1158 (1993) 227-236.
- 384 Poorman, J.A., Buck, R.A., Smith, S.A., Overturf, M.L. and Loose-Mitchell, D.S.: Bile acid excretion and cholesterol 7 $\alpha$ -hydroxylase expression in hypercholesterolemia-resistant rabbits. *J. Lipid Res.*, 34 (1993) 1675-1685.
- 385 Rajas, F. and Rousset, B.: A membrane-bound form of glutamate dehydrogenase possesses an ATP-dependent high-affinity microtubule-binding activity. *Biochem. J.*, 295 (1993) 447-455.
- 386 Rescigno, A., Sanjust, E., Curreli, N., Padiglia, A. and Floris, G.: Improved chromatographic purification of peroxidase and  $\beta$ -glucosidase from *Hordeum vulgare* seedlings. *Prepar. Biochem.*, 23 (1993) 485-492.
- 387 Vaidyanathan, V.V., Sastry, P.S. and Ramasarma, T.: Inverse relationship of the dehydrogenase and ADP-ribosylation activities in sodium-nitroprusside-treated glyceraldehyde-3-phosphate dehydrogenase is coincidental. *Biochim. Biophys. Acta*, 1203 (1993) 36-44.
- See also 95, 534.
- 20b. *Transferases (excl. E.C. 2.7.-.-)*
- 388 Bogin, E., Avidar, Y., Shenkler, S., Israeli, B.-A., Spiegel, N. and Cohen, R.: A rapid field test for the determination of colostral ingestion by calves, based on  $\gamma$ -glutamyltransferase. *Eur. J. Clin. Chem. Clin. Biochem.*, 31 (1993) 695-699.
- 389 Dannenberg, A.J., Yang, E.K. and Aharon, D.: Dietary lipids induce Phase 2 enzymes in rat small intestine. *Biochim. Biophys. Acta*, 1210 (1993) 8-12.
- 390 Derbel, M., Igarashi, T. and Satoh, T.: Differential induction of glutathione S-transferase subunits by phenobarbital, 3-methylcholanthrene and ethoxyquin in rat liver and kidney. *Biochim. Biophys. Acta*, 1158 (1993) 175-180.
- 391 Fast, D.G., Jamieson, J.C. and McCaffrey, G.: The role of the carbohydrate chains of Gal $\beta$ -1,4-GlcNAc $\alpha$ 2,6-sialyltransferase for enzyme activity. *Biochim. Biophys. Acta*, 1202 (1993) 325-330.
- 392 Hussey A.J. and Hayes, J.D.: Human Mu-class glutathione S-transferases present in liver, skeletal muscle and testicular tissue. *Biochim. Biophys. Acta*, 1203 (1993) 131-141.
- 393 Lessire, R., Bessoule, J.-J., Cook, L., Cinti, D.L. and Cassagne, C.: Occurrence and characterization of a dehydratase enzyme in the leek icosanoyl-CoA synthase complex. *Biochim. Biophys. Acta*, 1169 (1993) 243-249.
- 394 Vuorio, O.E., Kalkkinen, N. and Londesborough, J.: Cloning of the related genes encoding the 56-kDa and 123-kDa subunits of trehalose synthase from the yeast *Saccharomyces cerevisiae*. *Eur. J. Biochem.*, 216 (1993) 849-861.
- 20c. *Transferases transferring phosphorus containing groups (E.C. 2.7.-.-)*
- 395 Bikhhabhai, R., Carlsson, T., Unge, T., Lövgren, S. and Strandberg, B.: Increased yield of homogeneous HIV-1 reverse transcriptase (p66/p51) using a slow purification approach. *J. Chromatogr.*, 639 (1993) 67-74.
- 396 Burrows, J.A. and Goward, C.R.: Preparation of DNA polymerase from *Bacillus caldotenax*. *J. Chromatogr.*, 639 (1993) 75-80.
- 397 Cutler, R.E., Jr., Maizels, E.T., Brooks, E.J., Mizuno, K., Ohno, S. and Hunzicker-Dunn, M.: Regulation of  $\delta$  protein kinase C during rat ovarian differentiation. *Biochim. Biophys. Acta*, 1179 (1993) 260-270.
- 398 Ding, J. and Badwey, J.A.: Stimulation of neutrophils with a chemoattractant activates several novel protein kinases that can catalyze the phosphorylation of peptides derived from the 47-kDa protein component of the phagocyte oxidase and myristoylated alanine-rich C kinase substrate. *J. Biol. Chem.*, 268 (1993) 17326-17333.
- 399 Friedmann, D.L., Kesterson, R., Puleo, P., Wu, A.H.B. and Perryman, M.B.: Recombinant creatine kinase proteins and proposed standards for creatine kinase isoenzyme and subform assays. *Clin. Chem. (Washington)*, 39 (1993) 1598-1601.
- 400 Goldberg, G.S. and Lau, A.F.: Dynamics of connexin 43 phosphorylation in pp60<sup>v-src</sup> transformed cells. *Biochem. J.*, 295 (1993) 735-742.

- 401 Her, J.-H., Lakhani, S., Zu, K., Vila, J., Dent, P., Sturgill, T.W. and Weber, M.J.: Dual phosphorylation and autophosphorylation in mitogen-activated protein (MAP) kinase activation. *Biochem. J.*, 296 (1993) 25-31.
- 402 Liebenhoff, U., Brockmeier, D. and Presek, P.: Substrate affinity of the protein tyrosine kinase pp60<sup>c-src</sup> is increased on thrombin stimulation of human platelets. *Biochem. J.*, 295 (1993) 41-48.
- 403 Wang, L.-M., Kucera, G.L. and Capizzi, R.L.: Purification and characterization of deoxycytidine kinase from acute myeloid leukemia cell mitochondria. *Biochim. Biophys. Acta*, 1202 (1993) 309-316.
- 20d. *Hydrolases, acting on ester bonds (E.C. 3.1.-.-)*
- 404 Frank, P., Cazenave, C., Albert, S. and Toulme, J.-J.: Sensitive detection of low levels of ribonuclease H activity by an improved renaturation gel assay. *Biochem. Biophys. Res. Commun.*, 196 (1993) 1552-1557.
- 405 Gbenle, G.O.: Characterization of a filtrable 3'-nucleotidase of *Leishmania donovani*. *Biochim. Biophys. Acta*, 1203 (1993) 162-167.
- 406 Hohmann, J., Kowalewski, H., Vogel, M. and Zimmermann, H.: Isolation of a Ca<sup>2+</sup> or Mg<sup>2+</sup>-activated ATPase (ecto-ATPase) from bovine brain synaptic membranes. *Biochim. Biophys. Acta*, 1152 (1993) 146-154.
- 407 Kozaki, K., Gotoda, T., Kawamura, M., Shimano, H., Yazaki, Y., Ouchi, Y., Orimo, H. and Yamada, N.: Mutational analysis of human lipoprotein lipase by carboxy-terminal truncation. *J. Lipid Res.*, 34 (1993) 1765-1772.
- 408 Kroetz, D.L., McBride, O.W. and Gonzalez, F.J.: Glycosylation-dependent activity of baculovirus-expressed human liver carboxylesterases: cDNA cloning and characterization of two highly similar enzyme forms. *Biochemistry*, 32 (1993) 11606-11617.
- 409 Manco, L. and Amorim, A.: Human erythrocyte pyrimidine 5'-nucleotidase isozymes: Effect of sulfhydryl reagents and electrophoretic discrimination. *Electrophoresis (Weinheim)*, 14 (1993) 1084-1085.
- 410 Minami, T., Tojo, H., Shinomura, Y., Matsuzawa, Y. and Okamoto, M.: Purification and characterization of a phospholipase A<sub>2</sub> from human ileal mucosa. *Biochim. Biophys. Acta*, 1170 (1993) 125-130.
- 411 Pedersen, J., Pedersen, M., Soeberg, H. and Biedermann, K.: Separation of isoforms of *Serratia marcescens* nuclease by capillary electrophoresis. *J. Chromatogr.*, 645 (1993) 353-361.
- 412 Tate, B.F. and Rittenhouse, S.E.: Thrombin activation of human platelets causes tyrosine phosphorylation of PLC- $\gamma_2$ . *Biochim. Biophys. Acta*, 1178 (1993) 281-285.
- 413 Tenjo, E., Sawazaki, K., Yasuda, T., Nadano, D., Takeshita, H., Iida, R. and Kishi, K.: Salivary deoxyribonuclease I polymorphism separated by polyacrylamide gel-isoelectric focusing and detected by the dried agarose film overlay method. *Electrophoresis (Weinheim)*, 14 (1993) 1042-1044.
- 414 Tsujita, T. and Okuda, H.: Palmitoyl-coenzyme A hydrolyzing activity in rat kidney and its relationship to carboxylesterase. *J. Lipid Res.*, 34 (1993) 1773-1781.
- 415 Van Hoof, V.O., van Mullem, M., de Broe, M.E. and Lepoutre, L.G.: Comparison of two commercially available systems for the electrophoretic separation of alkaline phosphatase isoenzymes. *J. Chromatogr.*, 646 (1993) 235-243.
- 416 Vergnes, H.A., Denier, C.J., Brisson-Lougarre, A., Grozdea, J.G., Bourrouillou, G., Colombies, P., Klaebe, A.G. and Perie, J.J.: Biochemical and immunological characteristics of neutrophil alkaline phosphatase in Down's syndrome. *Clin. Chim. Acta*, 218 (1993) 105-112.
- 417 Vilella, E., Joven, J., Fernandez, M., Vilaro, S., Brunzell, J.D., Olivecrona, T. and Bengtsson-Olivecrona, G.: Lipoprotein lipase in human plasma is mainly inactive and associated with cholesterol-rich lipoproteins. *J. Lipid Res.*, 34 (1993) 1555-1564.
- 418 Yasuda, T., Sawazaki, K., Nadano, D., Takeshita, H., Nakanaga, M. and Kishi, K.: Human seminal deoxyribonuclease I (DNase I): purification, enzymological and immunological characterization and origin. *Clin. Chim. Acta*, 218 (1993) 5-16.
- 20e. *Hydrolases, acting on glycosyl compounds (E.C. 3.2.-.-)*
- 419 Chen, Y.Q. and Wenger, D.A.: Galactocerebrosidase from human urine: purification and partial characterization. *Biochim. Biophys. Acta*, 1170 (1993) 53-61.
- 420 Evans, B.R., Margalit, R. and Woodward, J.: Attachment of pentaamineruthenium(III) to *Trichoderma reesei* cellobiohydrolase I increases its catalytic activity. *Biochem. Biophys. Res. Commun.*, 195 (1993) 497-503.
- 421 Franzini, E., Sellak, H., Hakim, J. and Pasquier, C.: Oxidative damage to lysozyme by the hydroxyl radical: comparative effects of scavengers. *Biochim. Biophys. Acta*, 1203 (1993) 11-17.
- 422 Kim, C.-H. and Kho, Y.-H.: Domain structure and multiplicity of raw-starch-digesting amylase from *Bacillus circulans*: extensive proteolysis with proteinase K, endopeptidase Glu-C and thermolysin. *Biochim. Biophys. Acta*, 1202 (1993) 200-206.
- 423 Ring, M. and Huber, R.E.: The properties of  $\beta$ -galactosidases (*Escherichia coli*) with halogenated tyrosines. *Biochem. Cell Biol.*, 71 (1993) 127-132.
- See also 386, 433.
- 20f. *Other hydrolases*
- 424 Dormer, R.L., Capurro, D.E., Morris, R. and Webb, R.: Demonstration of two isoforms of the SERCA-2b type Ca<sup>2+</sup>,Mg<sup>2+</sup>-ATPase in pancreatic endoplasmic reticulum. *Biochim. Biophys. Acta*, 1152 (1993) 225-230.
- 425 Harcum, S.W. and Bentley, W.E.: Detection, quantification, and characterization of proteases in recombinant *Escherichia coli*. *Biotechnol. Tech.*, 7 (1993) 441-447; *C.A.*, 119 (1993) 89587t.
- 426 Kolkenbrock, H., Hecker-Kia, A., Orgel, D., Buchlow, G., Sörensen, H., Hauer, W. and Ulbrich, N.: A trypsin sensitive stromelysin isolated from rheumatoid synovial fluid as an activator for matrix metalloproteinases. *Eur. J. Clin. Chem. Clin. Biochem.*, 31 (1993) 625-631.
- 427 Kumar, S. and Pritchard, D.I.: New observations on proteases of the human hookworm *Necator americanus*. *Biochem. Mol. Biol. Int.*, 30 (1993) 13-19; *C.A.*, 119 (1993) 89684x.
- 428 Muno, D., Ishidoh, K., Ueno, T. and Kominami, E.: Processing and transport of the precursor of cathepsin C during its transfer into lysosomes. *Arch. Biochem. Biophys.*, 306 (1993) 103-110.



- 429 Ng, Y.-C.: Hypokalemia downregulates cardiac  $\alpha 1$  and skeletal muscle  $\alpha 2$  isoforms of  $\text{Na}^+, \text{K}^+$ -ATPase in ferrets. *Biochem. Biophys. Res. Commun.*, 196 (1993) 39-46.
- 430 Nielsen, F.S. and Foltmann, B.: Activation of porcine pepsinogen A. The stability of two non-covalent activation intermediates at pH 8.5. *Eur. J. Biochem.*, 217 (1993) 137-142.
- 431 Nishikata, M.: Synthesis of  $\text{N}\alpha$ -[ $^3\text{H}$ ]-acetyl-L-lysine chloromethyl ketone and its use in the fluorographic detection of proteases. *Anal. Biochem.*, 214 (1993) 222-226.
- 432 Peters, K., Fittkau, S., Steinert, A. and Ströhl, D.: Peptidyl methyl ketones as ligands in affinity chromatography of serine and cysteine proteinases. *J. Chromatogr.*, 648 (1993) 91-99.
- 433 Pshezhetsky, A.V. and Potier, M.: Stoichiometry of the human lysosomal carboxypeptidase- $\beta$ -galactosidase complex. *Biochem. Biophys. Res. Commun.*, 195 (1993) 354-362.
- 434 Satoyoshi, M., Teranaka, T., Iwamoto, T., Imai, K., Kawase, T. and Saito, S.: (Electrophoretic behavior of protease on enzymogram: estimation of the molecular weight). *Shika Kiso Igakkai Zasshi*, 35 (1993) 186-196; *C.A.*, 119 (1993) 154536s.
- 435 Tal, D.M., Goldshleger, R. and Karlish, S.J.D.: Separation of membrane-embedded tryptic peptides of  $\text{Na}, \text{K}$ -ATPase by size-exclusion chromatography. *J. Chromatogr.*, 646 (1993) 153-157.
- 436 Vinther, A., Adelhorst, K. and Kirk, O.: Using capillary electrophoresis in the optimization of a carboxypeptidase Y catalyzed transpeptidation reaction. *Electrophoresis (Weinheim)*, 14 (1993) 486-491.
- 437 Zhang, J., Fujimoto, N., Iwata, K., Sakai, T., Okada, Y. and Hayakawa, T.: A one-step sandwich enzyme immunoassay for human matrix metalloproteinase 1 (interstitial collagenase) using monoclonal antibodies. *Clin. Chim. Acta*, 219 (1993) 1-14.

See also 193, 431, 476.

#### 20g. Lyases

- 438 Fujii, H., Nakamura, K., Takeo, K. and Kawai, S.: Heterogeneity of carbonic anhydrase and 68 kDa neurofilament in nerve roots analyzed by two-dimensional electrophoresis. *Electrophoresis (Weinheim)*, 14 (1993) 1074-1078.
- 439 Martinou, A., Kafetzopoulos, D. and Bouriotis, V.: Isolation of chitin deacetylase from *Mucor rouxii* by immunoaffinity chromatography. *J. Chromatogr.*, 644 (1993) 35-41.
- 440 Ranganathan, S. and Tew, K.D.: Analysis of glyoxalase-I from normal and tumor tissue from human colon. *Biochim. Biophys. Acta*, 1182 (1993) 311-316.

#### 20h. Isomerases

- 441 Juenke, J.E.M. and Holden, J.A.: The distribution of DNA topoisomerase II isoforms in differentiated adult mouse tissues. *Biochim. Biophys. Acta*, 1216 (1993) 191-196.
- 442 Yokoi, T., Nagayama, S., Kajiwara, R., Kawaguchi, Y., Horiuchi, R. and Kamataki, T.: Identification of protein disulfide isomerase and calreticulin as autoimmune antigens in LEC strain of rats. *Biochim. Biophys. Acta*, 1158 (1993) 339-344.

See also 491.

#### 20i. Ligases

- 443 Menguito, C.A., Papaconstantinou, J. and Weigel, P.H.: The histidyl-tRNA synthetase from *Streptococcus equisimilis*: over expression in *Escherichia coli*, purification, and characterization. *Prepar. Biochem.*, 23 (1993) 449-472.

#### 20j. Complex mixtures and incompletely identified enzymes

- 444 Keim-Miller, C.A., Ream, W. and Mosbaugh, D.W.: DNA helicase activity detected *in situ* following polyacrylamide gel electrophoresis. *Methods Mol. Cell. Biol.*, 3 (1992) 259-269; *C.A.*, 119 (1993) 133961m.
- 445 Zomer, A.W.M., de Weerd, W.F.C., Langeveld, J. and van den Bosch, H.: Ether lipid synthesis: purification and identification of alkyl dihydroxyacetone phosphate synthase from guinea-pig liver. *Biochim. Biophys. Acta*, 1170 (1993) 189-196.

#### 21. PURINES, PYRIMIDINES, NUCLEIC ACIDS AND THEIR CONSTITUENTS

##### 21a. Purines, pyrimidines, nucleosides, nucleotides

- 446 Andrus, A.: Oligonucleotide analysis by gel capillary electrophoresis. *Methods (San Diego)*, 4 (1992) 213-226; *C.A.*, 119 (1993) 176957g.
- 447 Baba, Y.: Prediction of the behaviour of oligonucleotides in high-performance liquid chromatography and capillary electrophoresis. *J. Chromatogr.*, 618 (1993) 41-55 - a review with 65 refs.
- 448 Bruchelt, G., Niethammer, D. and Schmidt, K.H.: Isotachopheresis of nucleic acid constituents. *J. Chromatogr.*, 618 (1993) 57-77 - a review with 107 refs.
- 449 Norwood, C.B., Jackim, E. and Cheer, S.: DNA adduct research with capillary electrophoresis. *Anal. Biochem.*, 213 (1993) 194-199.
- 450 Paulus, A.: Oligonucleotides separation with capillary gel electrophoresis. *Boll. Chim. Farm.*, 131 (1992) 299-301; *C.A.*, 119 (1993) 134816e.
- 451 Van Schepdael, A., Vandeweyer, M., van Aerschot, A., Herdewijn, P., Roets, E. and Hoogmartens, J.: Separation of the anomers and isomers of 2'-deoxyuridine and thymidine by capillary zone electrophoresis. *J. Chromatogr.*, 648 (1993) 299-305.
- 452 Warren, W.J. and Vella, G.: Analysis of synthetic oligodeoxyribonucleotides by capillary gel electrophoresis and anion-exchange HPLC. *BioTechniques*, 14 (1993) 598-606; *C.A.*, 119 (1993) 112827g.
- 453 Yabuki, T., Ejiri, S.-i. and Tsutsumi, K.-i.: Ubiquitous factors that interact simultaneously with two distinct *cis*-elements on the rat aldolase B gene promoter. *Biochim. Biophys. Acta*, 1216 (1993) 15-19.

See also 29, 345, 601.

21b. *Nucleic acids, RNA*

- 454 Abe, H., Ohtake, A., Yamamoto, S., Satoh, Y., Takayanagi, M., Amaya, Y., Takiguchi, M., Sakuraba, H., Suzuki, Y., Mori, M. and Niimi, H.: Cloning and sequence analysis of a full length cDNA encoding human mitochondrial 3-oxoacyl-CoA thiolase. *Biochim. Biophys. Acta*, 1216 (1993) 304-306.
- 455 Abe, T., Nishiyama, F., Snajdar, R., He, X. and Misono, K.S.: Aortic smooth muscle contains guanylate-cyclase-coupled 130-kDa atrial natriuretic factor receptor as predominant receptor form. Spontaneous switching to 60-kDa  $\beta$ -receptor upon cell culturing. *Eur. J. Biochem.*, 217 (1993) 295-304.
- 456 Andrews, A.T.: Electrophoresis of nucleic acids. In: Brown, T.A. (Editor), *Essent. Mol. Biol.*, IRL, Oxford, 1991, pp. 89-126; C.A., 119 (1993) 134619h - a review with 71 refs.
- 457 Buiser, R.G., Bambara, R.A. and Fay, P.J.: Pausing by retroviral DNA polymerases promotes strand transfer from internal regions of RNA donor templates to homopolymeric acceptor templates. *Biochim. Biophys. Acta*, 1216 (1993) 20-30.
- 458 Cullingford, T.E., Clark, J.B. and Phillips, I.R.: Characterization of cDNAs encoding the rat testis-specific E1 $\alpha$  subunit of the pyruvate dehydrogenase complex: comparison of expression of the corresponding mRNA with that of the somatic E1 $\alpha$  subunit. *Biochim. Biophys. Acta*, 1216 (1993) 149-153.
- 459 DeWille, J.W. and Farmer, S.J.: Linoleic acid controls neonatal tissue-specific stearyl-CoA desaturase mRNA levels. *Biochim. Biophys. Acta*, 1170 (1993) 291-295.
- 460 Dixon, B., Sahely, B., Liu, L. and Pohajdak, B.: Cloning a cDNA from human NK/T cells which codes for an unusual leucine zipper containing protein. *Biochim. Biophys. Acta*, 1216 (1993) 321-324.
- 461 Drabent, B., Bode, C. and Doenecke, D.: Structure and expression of the mouse testicular H1 histone gene (H1t). *Biochim. Biophys. Acta*, 1216 (1993) 311-313.
- 462 During, K.: Non-radioactive detection methods for nucleic acids separated by electrophoresis. *J. Chromatogr.*, 618 (1993) 105-131 - a review with 72 refs.
- 463 Fogel-Petrovic, M., Kramer, D.L., Ganis, B., Casero, R.A., Jr. and Porter, C.W.: Cloning and sequence analysis of the gene and cDNA encoding mouse spermidine/spermine N<sup>1</sup>-acetyltransferase - a gene uniquely regulated by polyamines and their analogs. *Biochim. Biophys. Acta*, 1216 (1993) 255-264.
- 464 Fujii, G., Tashiro, K., Emori, Y., Saigo, K. and Shiokawa, K.: Molecular cloning of cDNAs for two *Xenopus* proteasome subunits and their expression in adult tissues. *Biochim. Biophys. Acta*, 1216 (1993) 65-72.
- 465 Gambacurta, A., Piro, M.C. and Ascoli, F.: Differential *in vitro* translation of the precursors of bovine pancreatic trypsin inhibitor and its iso-inhibitor II is controlled by the 5'-end region of their mRNAs. *Biochim. Biophys. Acta*, 1174 (1993) 267-273.
- 466 Knecht, D., Kern, H. and Scherzinger, C.: Bidirectional transcription from actin promoters in *Dictyostelium*. *Biochim. Biophys. Acta*, 1216 (1993) 105-109.
- 467 Koch, O.M., Danenberg, P.V., Volkenandt, M., Danenberg, K., Bertino, J. and Horikoshi, T.: Detection of mutations and monoclonal lineages by analysis of RNA conformation polymorphism(s). *PCT Int. Appl. WO 93 11,266 (Cl. C12Q1/68)*, 10 Jun. 1993, US Appl. 803,171, 05 Dec. 1991; 52 pp.; C.A., 119 (1993) 88351z.
- 468 Kroczeck, R.A.: Southern and Northern analysis. *J. Chromatogr.*, 618 (1993) 133-145 - a review with 57 refs.
- 469 Kuentzel, S.L., Ali, S.M., Altman, R.A., Greenberg, B.D. and Raub, T.J.: The Alzheimer  $\beta$ -amyloid protein precursor/protease nexin-II is cleaved by secretase in a *trans*-Golgi secretory compartment in human neuroglioma cells. *Biochem. J.*, 295 (1993) 367-378.
- 470 Kurie, J.M., Younes, A., Miller, W.H., Jr., Burchert, M., Chiu, C.-F., Kolesnick, R. and Dmitrovsky, E.: Retinoic acid stimulates the protein kinase C pathway before activation of its  $\beta$ -nuclear receptor during human teratocarcinoma differentiation. *Biochim. Biophys. Acta*, 1179 (1993) 203-207.
- 471 Lilley, D.M.J., Bhattacharyya, A. and McAteer, S.: Gel electrophoresis and the structure of RNA molecules. *Biotechnol. Genet. Eng. Rev.*, 10 (1992) 379-401; C.A., 119 (1993) 90298f - a review with 100 refs.
- 472 López, S., Vila, L., Breviaro, F. and de Castellarnau, C.: Interleukin-1 increases 15-hydroxyeicosatetraenoic acid formation in cultured human endothelial cells. *Biochim. Biophys. Acta*, 1170 (1993) 17-24.
- 473 Mantamadiotis, T., Sobieszczuk, P., Weinstock, J. and Baldwin, G.S.: Nucleotide sequence encoding a novel member of the hydratase/dehydrogenase family. *Biochim. Biophys. Acta*, 1170 (1993) 211-215.
- 474 Moran, T.V. and Walker, J.C.: Molecular cloning of two novel protein kinase genes from *Arabidopsis thaliana*. *Biochim. Biophys. Acta*, 1216 (1993) 9-14.
- 475 Ohgiya, S., Ishizaki, K. and Shinriki, N.: Molecular cloning of guinea pig CYP1A1: complete primary structure and fast mobility of expressed protein on electrophoresis. *Biochim. Biophys. Acta*, 1216 (1993) 237-244.
- 476 Pan, Y.-X., Gu, H.H., Xu, J. and Dean, G.E.: *Saccharomyces cerevisiae* expression of exogenous vacuolar ATPase subunits B. *Biochim. Biophys. Acta*, 1151 (1993) 175-185.
- 477 Pelle, R. and Murphy, N.B.: Northern hybridization: Rapid and simple electrophoretic conditions. *Nucleic Acids Res.*, 21 (1993) 2783-2784; C.A., 119 (1993) 87486k.
- 478 Rajendrakumar, G.V., Radha, V. and Swarup, G.: Stabilization of a protein-tyrosine phosphatase mRNA upon mitogenic stimulation of T-lymphocytes. *Biochim. Biophys. Acta*, 1216 (1993) 205-212.
- 479 Schrum, L.W. and Hassan, H.M.: Transcriptional activation of Mn-superoxide dismutase gene (*sodA*) of *Escherichia coli* by MnCl<sub>2</sub>. *Biochim. Biophys. Acta*, 1216 (1993) 186-190.
- 480 Shashidharan, P. and Plaitakis, A.: Cloning and characterization of a glutamate transporter cDNA from human cerebellum. *Biochim. Biophys. Acta*, 1216 (1993) 161-164.
- 481 Swanson, K.D., Dhar, M.S. and Joshi, J.G.: The human and bovine 14-3-3 $\gamma$  protein mRNAs are highly conserved in both their translated and untranslated regions. *Biochim. Biophys. Acta*, 1216 (1993) 145-148.
- 482 Thinakaran, G. and Bag, J.: Regulation of c-jun/AP-1 expression in rat L6 myoblasts. *Biochem. Cell Biol.*, 71 (1993) 197-204.
- 483 Wu, P., Skrede, S., Hvattum, E. and Bremer, J.: Substrate and hormone regulation of palmitoyl-CoA synthetase in 7800 C1 Morris hepatoma cells and cultured rat hepatocytes. *Biochim. Biophys. Acta*, 1170 (1993) 118-124.

See also 143, 238, 242, 373, 397, 407, 410, 485, 519, 531, 578.

## 21c. Nucleic acids, DNA

- 484 Baba, Y. and Tshako, M.: (Development of new technology for DNA analysis using capillary electrophoresis). *Bio Ind.*, 10 (1993) 299-305; *C.A.*, 119 (1993) 134601f - a review with 12 refs.
- 485 Bai, S., Spray, D.C. and Burk, R.D.: Identification of proximal and distal regulatory elements of the rat connexin32 gene. *Biochim. Biophys. Acta*, 1216 (1993) 197-204.
- 486 Barron, A.E., Soane, D.S. and Blanch, H.W.: Capillary electrophoresis of DNA in uncross-linked polymer solutions. *J. Chromatogr.*, 652 (1993) 3-16.
- 487 Bianchi, N., Mischiati, C., Feriotto, G. and Gambari, R.: Polymerase-chain reaction: analysis of DNA/DNA hybridization by capillary electrophoresis. *Nucleic Acids Res.*, 21 (1993) 3595-3596; *C.A.*, 119 (1993) 174902m.
- 488 Cardot, P., Chambaz, J., Kardassis, D., Cladaras, C. and Zannis, V.I.: Factors participating in the liver-specific expression of the human apolipoprotein A-II gene and their significance for transcription. *Biochemistry*, 32 (1993) 9080-9093.
- 489 Chen, M., Maerz, W., Manfras, B.J., Kuehl, P., Usadel, K.H. and Boehm, B.O.: Rapid and simple subtyping of the HLA-DRB3 gene in Graves' disease by using temperature-gradient gel electrophoresis. *Hum. Immunol.*, 36 (1993) 199-203; *C.A.*, 119 (1993) 136976f.
- 490 Chu, B., Sokolov, E. and Khokhlov, A.: Laser experiments on DNA electrophoretic mobilities. *SPIE-Int. Soc. Opt. Eng.*, 1922(Laser Study of Macroscopic Biosystems) (1993) 354-359; *C.A.*, 119 (1993) 134819h.
- 491 De la Tour, C.B., Portemer, C., Forterre, P., Huber, R. and Duguet, M.: ATP-independent DNA topoisomerase from *Feridobacterium islandicum*. *Biochim. Biophys. Acta*, 1216 (1993) 213-220.
- 492 DeDionisio, L.: Capillary gel electrophoresis and the analysis of DNA phosphorothioates. *J. Chromatogr.*, 652 (1993) 101-108.
- 493 Den Dunnen, J.T. and van Ommen, G.J.B.: Methods for pulsed-field gel electrophoresis. *Appl. Biochem. Biotechnol.*, 38 (1993) 161-177; *C.A.*, 119 (1993) 90567t.
- 494 Diamandis, E.P.: Time-resolved fluorometry in nucleic acid hybridization and Western blotting techniques. *Electrophoresis (Weinheim)*, 14 (1993) 866-875.
- 495 Duke, T.: Computer simulation of DNA electrophoresis. *Analisis*, 21 (1993) M29-M31; *C.A.*, 119 (1993) 109884t - a review with 6 refs.
- 496 Ferraboli, S., Negri, R., di Mauro, E. and Barlati, S.: One-lane chemical sequencing of 3'-fluorescent-labeled DNA. *Anal. Biochem.*, 214 (1993) 566-570.
- 497 Figeys, D. and Dovichi, N.J.: Mobility of single-stranded DNA as a function of cross-linker concentration in polyacrylamide capillary gel electrophoresis. *J. Chromatogr.*, 645 (1993) 311-317.
- 498 Frenkel, K. and Klein, C.B.: Methods used for analyses of "environmentally" damaged nucleic acids. *J. Chromatogr.*, 618 (1993) 289-314 - a review with 176 refs.
- 499 Fujiwara, Y., Oka, M., Hamamoto, T. and Sone, N.: Cytochrome c-551 of the thermophilic bacterium PS3, DNA sequence and analysis of the mature cytochrome. *Biochim. Biophys. Acta*, 1144 (1993) 213-219.
- 500 Gordillo, M.E., Singh, K.V., Baker, C.J. and Murray, B.E.: Typing of group B streptococci: comparison of pulsed-field gel electrophoresis and conventional electrophoresis. *J. Clin. Microbiol.*, 31 (1993) 1430-1434; *C.A.*, 119 (1993) 112707t.
- 501 Guszczynski, T., Pulyaeva, H., Tietz, D., Garner, M.M. and Chrambach, A.: Capillary zone electrophoresis of large DNA. *Electrophoresis (Weinheim)*, 14 (1993) 523-530.
- 502 Hampel, K.J. and Lee, J.S.: Two-dimensional pulsed-field gel electrophoresis of yeast chromosomes: evidence for triplex-mediated DNA condensation. *Biochem. Cell Biol.*, 71 (1993) 190-196.
- 503 Heller, C.: Pulsed field gel electrophoresis. *Analisis*, 21 (1993) M11-M12, M14; *C.A.*, 119 (1993) 134595g - a review with 11 refs.
- 504 Hensels, G.W., Janssen, E.A.M., Hoogendijk, J.E., Valentijn, L.J., Baas, F. and Bolhuis, P.A.: Quantitative measurement of duplicated DNA as a diagnostic test for Charcot-Marie-Tooth disease type 1a. *Clin. Chem. (Washington)*, 39 (1993) 1845-1849.
- 505 Imamura, Y., Nakagawa, T., Iguchi-Ariga, S.M.M. and Ariga, H.: Transcriptional regulation of the N-myc gene: identification of positive regulatory element and its double- and single-stranded DNA binding proteins. *Biochim. Biophys. Acta*, 1216 (1993) 273-285.
- 506 Jeltsch, A., Fritz, A., Alves, J., Wolfes, H. and Pingoud, A.: A fast and accurate enzyme-linked immunosorbent assay for the determination of the DNA cleavage activity of restriction endonucleases. *Anal. Biochem.*, 213 (1993) 234-240.
- 507 Kauc, L.: (DNA pulsed-field gel electrophoresis - PFGE). *Postepy Mikrobiol.*, 31 (1992) 35-56; *C.A.*, 119 (1993) 112442c - a review with 99 refs.
- 508 Konrad, K.D. and Pentoney, S.L., Jr.: Contribution of secondary structure to DNA mobility in capillary gels. *Electrophoresis (Weinheim)*, 14 (1993) 502-508.
- 509 Kostka, H., Gehrisch, S. and Jaros, W.: The use of temperature gradient gel electrophoresis as a screening method for point mutations of familial combined hyperlipidaemia. *Eur. J. Clin. Chem. Clin. Biochem.*, 31 (1993) A35.
- 510 Kozlowski, P., Tymowska, Z. and Toczko, K.: Nucleotide and predicted amino acid sequence of a new member of the ras gene family from the slime mold *Physarum polycephalum*. *Biochim. Biophys. Acta*, 1174 (1993) 299-302.
- 511 Liu, C.H.L., Liu, S. and Wang, E.: Expression of an EF-1 $\alpha$ -like rat cDNA, S1, in *E. coli* and production of a rabbit polyclonal antiserum to the recombinant protein. *Biochem. Biophys. Res. Commun.*, 195 (1993) 1371-1378.
- 512 Luckey, J.A. and Smith, L.M.: A model for the mobility of single-stranded DNA in capillary gel electrophoresis. *Electrophoresis (Weinheim)*, 14 (1993) 492-501.
- 513 MacConnell, W.P.: Nucleic acid purification system and method. *U.S. US 5,217,593 (Cl. 204-299R; GO1N27/26)*, 08 Jun. 1993, *US Appl. 668,856*, 13 Mar. 1991; 10 pp.; *C.A.*, 119 (1993) 132918r.
- 514 Martin, F., Vairalles, D. and Henrion, B.: Automated ribosomal DNA fingerprinting by capillary electrophoresis of PCR products. *Anal. Biochem.*, 214 (1993) 182-189.
- 515 Martin, W., Hoppe, H.H., Mucbe, M. and Henke, L.: Comparison of population genetics of the single locus probes pS194 and pL427-4. *Adv. Forensic Haemogenet.*, 4 (1992) 195-197; *C.A.*, 119 (1993) 87992x.

- 516 Maschke, H.E., Frenz, J., Belenkii, A., Karger, B.L. and Hancock, W.S.: Ultrasensitive plasmid mapping by high performance capillary electrophoresis. *Electrophoresis (Weinheim)*, 14 (1993) 509-514.
- 517 Masubuchi, Y., Oana, H., Ono, K., Matsumoto, M., Doi, M., Minagawa, K., Matsuzawa, Y. and Yoshikawa, K.: Periodic behavior of DNA molecules during steady field gel electrophoresis. *Macromolecules*, 26 (1993) 5269-5270; *C.A.*, 119 (1993) 134835k.
- 518 Mayer, P., Sturm, J. and Weill, G.: DNA deformation in gel electrophoresis. *Proc. SPIE-Int. Soc. Opt. Eng.*, 1922(Laser Study of Macroscopic Biosystems) (1993) 368-376; *C.A.*, 119 (1993) 134820b.
- 519 Michaelis, C. and Weeks, G.: The isolation from a unicellular organism, *Dictyostelium discoideum*, of a highly-related *cdc2* gene with characteristics of the PCTAIRE subfamily. *Biochim. Biophys. Acta*, 1179 (1993) 117-124.
- 520 Millar, D.B.: Fluorstamp. *BioTechniques*, 14 (1993) 764. *C.A.*, 119 (1993) 90388k.
- 521 Nielsen, P.S. and Gausing, K.: *In vitro* binding of nuclear proteins to the barley plastocyanin gene promoter region. *Eur. J. Biochem.*, 217 (1993) 97-104.
- 522 Peterson, C.A., Gray, D.M., Gray, H.B., Jr. and Legerski, R.J.: Evidence for a salt-induced conformational transition in UV-irradiated superhelical PM2 DNA. *Biochim. Biophys. Acta*, 1216 (1993) 265-272.
- 523 Reynolds, J.E. and Diehl, S.R.: A method for recovery of high-molecular-weight DNA suitable for field-inversion gel electrophoresis from frozen tumor cells. *Cancer Genet. Cytogenet.*, 65 (1993) 68-70; *C.A.*, 119 (1993) 90552j.
- 524 Ridanpaa, M. and Husgafvel-Pursiainen, K.: Denaturing gradient gel electrophoresis (DGGE) assay for K-ras and N-ras genes: Detection of K-ras point mutations in human lung tumor DNA. *Hum. Mol. Genet.*, 2 (1993) 639-644; *C.A.*, 119 (1993) 132474t.
- 525 Rye, H.S., Yue, S., Quesada, M.A., Haugland, R.P., Mathies, R.A. and Glazer, A.N.: Picogram detection of stable dye-DNA intercalation complexes with two-color laser excited confocal fluorescence gel scanner. *Methods Enzymol.*, 217(Recombinant DNA, Pt. H) (1993) 414-431; *C.A.*, 119 (1993) 134813b.
- 526 Sangwan, V., Lenvik, T.R. and Gantt, J.S.: The *Arabidopsis thaliana* ribosomal protein S15 (*rig*) gene. *Biochim. Biophys. Acta*, 1216 (1993) 221-226.
- 527 Satow, T., Akiyama, T., Machida, A., Utagawa, Y. and Kobayashi, H.: Simultaneous determination of the migration coefficient of each base in heterogeneous oligo-DNA by gel filled capillary electrophoresis. *J. Chromatogr.*, 652 (1993) 23-30.
- 528 Seki, S., Akiyama, K., Watanabe, S. and Tsutsui, K.: Activity gel and activity blotting methods for detecting DNA-modifying (repair) enzymes. *J. Chromatogr.*, 618 (1993) 147-166 - a review with 66 refs.
- 529 Shastry, B.S.: Site-specific interaction of a partially purified human lens factor(s) with conserved sequences of the human gamma crystallin gene. *Arch. Biochem. Biophys.*, 306 (1993) 476-485.
- 530 Smith, D.R.: Agarose gel electrophoresis. *Methods Mol. Biol. (Totowa)*, 18(Transgenesis Techniques) (1993) 433-438; *C.A.*, 119 (1993) 90559s.
- 531 Toguri, T., Kobayashi, O. and Umemoto, N.: The cloning of eggplant seedling cDNAs encoding proteins from a novel cytochrome P-450 family (CYP76). *Biochim. Biophys. Acta*, 1216 (1993) 165-169.
- 532 Towner, P.: Recovery of DNA from electrophoresis gels. In: Brown, T.A. (Editor), *Essent. Mol. Biol.*, IRL, Oxford, 1991, pp. 127-141; *C.A.*, 119 (1993) 134821c.
- 533 Townsend, K.M. and Dawkins, H.J.S.: Field alternation gel electrophoresis - status quo. *J. Chromatogr.*, 618 (1993) 223-249 - a review with 192 refs.
- 534 Tu, G.-C. and Israel, Y.: A new approach for the rapid detection of common and atypical aldehyde dehydrogenase alleles. *Eur. J. Clin. Chem. Clin. Biochem.*, 31 (1993) 591-594.
- 535 Upcroft, P. and Upcroft, J.A.: Comparison of properties of agarose for electrophoresis of DNA. *J. Chromatogr.*, 618 (1993) 79-93 - a review with 146 refs.
- 536 Van den Bogert, C., de Vries, H., Holtrop, M., Muus, P., Dekker, H.L., van Galen, M.J.M., Bolhuis, P.A. and Taanman, J.-W.: Regulation of the expression of mitochondrial proteins: relationship between mtDNA copy number and cytochrome-c oxidase activity in human cells and tissues. *Biochim. Biophys. Acta*, 1144 (1993) 177-183.
- 537 Wu, C., Friedlander, P., Lamoureux, C., Zannis-Hadjopoulos, M. and Price, G.B.: cDNA clones contain autonomous replication activity. *Biochim. Biophys. Acta*, 1174 (1993) 241-257.
- 538 Yagi, M., Matsumoto, M. and Doi, M.: Dynamics of DNA in gel electrophoresis. In: Doyama, M. (Editor), *Comput. Aided Innovation New Mater. 2, Proc. Int. Conf. Exhib. Comput. Appl. Mater. Mol. Sci. Eng.*, 2nd 1992, Pt. 1, North-Holland, Amsterdam, 1993, pp. 311-314; *C.A.*, 119 (1993) 176091v.
- 539 Yoshikawa, H., Nagai, H., Matsubara, K. and Fujiyama, A.: Two-dimensional gel electrophoretograms of human chromosome specific restriction DNA fragments. *Biochem. Biophys. Res. Commun.*, 196 (1993) 1566-1572.
- 540 Young, C.C., Burghoff, R.L., Keim, L.G., Minak-Bernero, V., Lute, J.R. and Hinton, S.M.: Polyvinylpyrrolidone-agarose gel electrophoresis purification of polymerase chain reaction-amplifiable DNA from soils. *Appl. Environ. Microbiol.*, 59 (1993) 1972-1974; *C.A.*, 119 (1993) 87491h.
- 541 Zastawny, T.H. and Olinski, R.: (Pulse-field electrophoresis of DNA). *Zagadnienia Biofiz. Wspolczesnej*, 15 (1991) 57-73; *C.A.*, 119 (1993) 112459p - a review with 59 refs.
- 542 Zeineh, M.M., Zeineh, J.A. and Zeineh, R.A.: Computing densitometer for electrophoresis and DNA. *Am. Lab. (Shelton)*, 25 (1993) 44-53; *C.A.*, 119 (1993) 155104m.
- 543 Zimm, B.H.: Mechanism of gel electrophoresis of DNA: Unexpected findings. *Curr. Opin. Struct. Biol.*, 3 (1993) 437-444; *C.A.*, 119 (1993) 176790x - a review with 26 refs.
- See also 221, 255, 257, 309, 449, 452, 456, 462, 463, 468, 555, 591.
- 21d. *Structural studies on RNA and RNA mapping*
- 544 Bai, G. and Kusiak, J.W.: Cloning and analysis of the 5' flanking sequence of the rat N-methyl-D-aspartate receptor 1 (NMDAR1) gene. *Biochim. Biophys. Acta*, 1152 (1993) 197-200.
- 545 Kille, P., Kay, J. and Sweeney, G.E.: Analysis of regulatory elements flanking metallothionein genes in Cd-tolerant fish (pike and stone loach). *Biochim. Biophys. Acta*, 1216 (1993) 55-64.

- 546 Marqués, S., Ramos, J.L. and Timmis, K.N.: Analysis of the mRNA structure of the *Pseudomonas putida* TOL meta fission pathway operon around the transcription initiation point, the xylTE and the xylFJ regions. *Biochim. Biophys. Acta*, 1216 (1993) 227-236.
- 547 Mordier, S., Béchet, D., Roux, M.-P., Obled, A. and Ferrara, M.: Nucleotide sequence of bovine preprocathepsin B. A study of polymorphism in the protein coding region. *Biochim. Biophys. Acta*, 1174 (1993) 305-311.
- See also 448, 463, 526, 558.
- 21e. *Structural studies on DNA and DNA mapping*
- 548 Baba, Y., Ishimaru, N., Samata, K. and Tshako, M.: High-resolution separation of DNA restriction fragments by capillary electrophoresis in cellulose derivative solutions. *J. Chromatogr.*, 653 (1993) 329-335.
- 549 Baba, Y., Tshako, M., Sawa, T. and Akashi, M.: Effect of urea concentration on the base-specific separation of oligodeoxynucleotides in capillary affinity gel electrophoresis. *J. Chromatogr.*, 652 (1993) 93-99.
- 550 Baba, Y., Tshako, M., Sawa, T., Akashi, M. and Yashima, E.: (Sequence-specific recognition of DNA by high-performance affinity chromatography and capillary affinity gel electrophoresis). *Kuromatogurafi*, 13 (1992) 333-334; *C.A.*, 119 (1993) 177061x.
- 551 Bay, S., Starke, H., Elliott, J. and Dovichi, N.: Accuracy of two-color peak height encoded DNA sequencing by capillary gel electrophoresis and laser-induced fluorescence. *Proc. SPIE-Int. Soc. Opt. Eng.*, 1891(Proceedings of Advances in DNA Sequencing Technology, 1993) (1993) 8-11; *C.A.*, 119 (1993) 174862y.
- 552 Beck, J.S., Kwitek, A.E., Cogen, P.H., Metzger, A.K., Duyk, G.M. and Sheffield, V.C.: A denaturing gradient gel electrophoresis assay for sensitive detection of p53 mutations. *Hum. Genet.*, 91 (1993) 25-30; *C.A.*, 119 (1993) 87397g.
- 553 Berg, E.S. and Olaisen, B.: PCR typing including high resolution gel electrophoresis reveals new alleles in the COL2A1 VNTR. *Adv. Forensic Haemogenet.*, 4 (1992) 44-46; *C.A.*, 119 (1993) 87409n.
- 554 Brdicka, R. and Nürnberg, P.: Checking of individuality by DNA profiling. *J. Chromatogr.*, 618 (1993) 167-179 - a review with 33 refs.
- 555 Chen, P.-H., Lin, S.-Y., Wang, C.-K., Chen, Y.-J., Chen, T.-C. and Chang, J.-G.: "Hot spots" mutation analysis of p53 gene in gastrointestinal cancers by amplification of naturally occurring and artificially created restriction sites. *Clin. Chem. (Washington)*, 39 (1993) 2186-2191.
- 556 Chiari, M., Nesi, M. and Righetti, P.G.: Movement of DNA fragments during capillary zone electrophoresis in liquid polyacrylamide. *J. Chromatogr.*, 652 (1993) 31-39.
- 557 DeLuca, C.I., Davies, P.L., Samis, J.A. and Elce, J.S.: Molecular cloning and bacterial expression of cDNA for rat calpain II 80 kDa subunit. *Biochim. Biophys. Acta*, 1216 (1993) 81-93.
- 558 Duro, G., Izzo, V. and Barbieri, R.: Methods for recovering nucleic acid fragments from agarose gels. *J. Chromatogr.*, 618 (1993) 95-104 - a review with 71 refs.
- 559 Halle, J.P., Wurst, H. and Schmidt, C.: Genomic sequencing by ligation mediated polymerase chain reaction using direct blotting and non-radioactive detection. *DNA Sequence*, 3 (1993) 283-289; *C.A.*, 119 (1993) 110048e.
- 560 Häne, B.G., Jäger, K. and Drexler, H.G.: The Pearson product-moment correlation coefficient is better suited for identification of DNA fingerprint profiles than band matching algorithms. *Electrophoresis (Weinheim)*, 14 (1993) 967-972.
- 561 Hanes, J., von der Kammer, H., Kristjansson, G.I. and Scheit, K.H.: The complete cDNA coding sequence for the mouse CDEI binding protein. *Biochim. Biophys. Acta*, 1216 (1993) 154-156.
- 562 Hartley, J.A., Souhami, R.L. and Berardini, M.D.: Electrophoretic and chromatographic separation methods used to reveal inter-strand crosslinking of nucleic acids. *J. Chromatogr.*, 618 (1993) 277-288 - a review with 37 refs.
- 563 Huang, X.C., Quesada, M.A. and Mathies, R.A.: Capillary array electrophoresis: an approach to high-speed, high-throughput DNA sequencing. *Proc. SPIE-Int. Soc. Opt. Eng.*, 189(Proceedings of Advances in DNA Sequencing Technology, 1993) (1993) 12-20; *C.A.*, 119 (1993) 132444h.
- 564 Izumi, T., Yamaguchi, M., Yoneda, K., Isobe, T., Okuyama, T. and Shinoda, T.: Use of glucomannan for the separation of DNA fragments by capillary electrophoresis. *J. Chromatogr.*, 652 (1993) 41-46.
- 565 Kleemiss, M.H., Gilges, M. and Schomburg, G.: Capillary electrophoresis of DNA restriction fragments with solutions of entangled polymers. *Electrophoresis (Weinheim)*, 14 (1993) 515-522.
- 566 Kretz, K.A. and O'Brien, J.S.: Direct sequencing of polymerase chain reaction products from low melting temperature agarose. *Methods Enzymol.*, 218(Recombinant DNA, Pt. I) (1993) 72-79; *C.A.*, 119 (1993) 110009t.
- 567 Kruse, L., Meyer, G. and Hildebrandt, A.: A highly conserved repetitive sequence from *Physarum polycephalum* contains nucleotide arrangements similar to replicator sequences. *Biochim. Biophys. Acta*, 1216 (1993) 129-133.
- 568 Lavoie, C.A., Harvey, M. and Lasko, P.F.: Dbp45A encodes a *Drosophila* DEAD box protein with similarity to a putative yeast helicase involved in ribosome assembly. *Biochim. Biophys. Acta*, 1216 (1993) 140-144.
- 569 Levi, L., Hanukoglu, I., Raikhin, M., Kohen, F. and Koch, Y.: Cloning of LL5, a novel protein encoding cDNA from a rat pituitary library. *Biochim. Biophys. Acta*, 1216 (1993) 342-344.
- 570 Luckey, J.A. and Smith, L.M.: Optimization of electric field strength for DNA sequencing in capillary gel electrophoresis. *Anal. Chem.*, 65 (1993) 2841-2850.
- 571 Luckey, J.A. and Smith, L.M.: Optimization of electric field strength for DNA sequencing in capillary gel electrophoresis. *Proc. SPIE-Int. Soc. Opt. Eng.*, 1891(Proceedings of Advances in DNA Sequencing Technology, 1993) (1993) 21-26; *C.A.*, 119 (1993) 132445j.
- 572 Luckey, J.A., Drossman, H., Kostichka, T. and Smith, L.M.: High-speed DNA sequencing by capillary gel electrophoresis. *Methods Enzymol.*, 218(Recombinant DNA, Pt. I) (1993) 154-172; *C.A.*, 119 (1993) 110013q.
- 573 Makowski, G.S., Aslanzadeh, J. and Hopfer, S.M.: Mini-gel PAGE for enhanced resolution of polymerase chain reaction detection of  $\Delta F_{508}$  deletion in cystic fibrosis. *Clin. Chem. (Washington)*, 39 (1993) 2204-2206.

- 574 Matsumoto, M., Kurata, S.-i., Fujimoto, H. and Hoshi, M.: Haploid specific activations of protamine 1 and hsc70t genes in mouse spermatogenesis. *Biochim. Biophys. Acta*, 1174 (1993) 274-278.
- 575 McCord, B.R., McClure, D.L. and Jung, J.M.: Capillary electrophoresis of polymerase chain reaction-amplified DNA using fluorescence detection with an intercalating dye. *J. Chromatogr.*, 652 (1993) 75-82.
- 576 McGregor, D.A. and Yeung, E.S.: Optimization of capillary electrophoretic separation of DNA fragments based on polymer filled capillaries. *J. Chromatogr.*, 652 (1993) 67-73.
- 577 Nobukuni, Y., Mitsubuchi, H., Hayashida, Y., Ohta, K., Indo, Y., Ichiba, Y., Endo, F. and Matsuda, I.: Heterogeneity of mutations in Maple Syrup Urine Disease (MSUD): screening and identification of affected E1 $\alpha$  and E1 $\beta$  subunits of the branched-chain  $\alpha$ -keto-acid dehydrogenase multienzyme complex. *Biochim. Biophys. Acta*, 1225 (1993) 64-70.
- 578 Ochoa, A., Bovard-Houppermans, S. and Zakin, M.M.: Human apolipoprotein A-IV gene expression is modulated by members of the nuclear hormone receptor superfamily. *Biochim. Biophys. Acta*, 1210 (1993) 41-47.
- 579 Pariat, Y.F., Berka, J., Heiger, D.N., Schmitt, T., Vilenchik, M., Cohen, A.S., Foret, F. and Karger, B.L.: Separation of DNA fragments by capillary electrophoresis using replaceable linear polyacrylamide matrices. *J. Chromatogr.*, 652 (1993) 57-66.
- 580 Pfaffinger, D., McLean, J. and Scanu, A.M.: Amplification of human APO(a) kringle 4-37 from blood lymphocyte DNA. *Biochim. Biophys. Acta*, 1225 (1993) 107-109.
- 581 Ribeiro, E.A.: Quantitative analysis of enzymic digests of DNA using gel electrophoresis. *J. Chromatogr.*, 618 (1993) 181-201 - a review with 43 refs.
- 582 Richterich, P. and Church, G.M.: DNA sequencing with direct transfer electrophoresis and nonradioactive detection. *Methods Enzymol.*, 218(Recombinant DNA, Pt. I) (1993) 187-222; *C.A.*, 119 (1993) 110015s.
- 583 Robertson, J., Schaefer, T., Kronick, M. and Budowle, B.: Automated analysis of fluorescent amplified fragment length polymorphisms for DNA typing. *Adv. Forensic Haemogenet.*, 4 (1992) 35-37; *C.A.*, 119 (1993) 87407k.
- 584 Ruiz-Martinez, M.C., Berka, J., Belenkii, A., Foret, F., Miller, A.W. and Karger, B.L.: DNA sequencing by capillary electrophoresis with replaceable linear polyacrylamide and laser-induced fluorescence detection. *Anal. Chem.*, 65 (1993) 2851-2858.
- 585 Ruzicka, V., März, W., Russ, A., Fisher, E., Mondorf, W. and Gross, W.: Characterization of the gene for apolipoprotein E5-Frankfurt (Gln<sup>81</sup> - Lys, Cys<sup>112</sup> - Arg) by polymerase chain reaction, restriction isotyping, and temperature gradient gel electrophoresis. *Electrophoresis (Weinheim)*, 14 (1993) 1032-1037.
- 586 Siedlaczek, I., Epplen, C., Riess, O. and Epplen, J.T.: Simple repetitive (GAA)<sub>n</sub> loci in the human genome. *Electrophoresis (Weinheim)*, 14 (1993) 973-977.
- 587 Singhal, R.P. and Xian, J.: Separation of DNA restriction fragments by polymer-solution capillary zone electrophoresis. Influence of polymer concentration and ion-pairing reagents. *J. Chromatogr.*, 652 (1993) 47-56.
- 588 Slater, G.W., Mayer, P. and Drouin, G.: Electrophoresis for DNA sequencing. *Analisis*, 21 (1993) M25-M28; *C.A.*, 119 (1993) 109883s - a review with 13 refs.
- 589 Slater, G.W., Mayer, P. and Drouin, G.: On the limits of near-equilibrium DNA gel electrophoretic sequencing. *Electrophoresis (Weinheim)*, 14 (1993) 961-966.
- 590 Srinivasan, K., Girard, J.E., Williams, P., Roby, R.K., Weedn, V.W., Morris, S.C., Kline, M.C. and Reeder, D.J.: Electrophoretic separations of polymerase chain reaction-amplified DNA fragments in DNA typing using a capillary electrophoresis-laser induced fluorescence system. *J. Chromatogr.*, 652 (1993) 83-91.
- 591 Takarada, Y., Kalanin, J., Yamashita, K., Ohtsuka, N., Kagawa, S. and Matsuoka, A.: Phenylketonuria mutant alleles in different populations: missense mutation in exon 7 of phenylalanine hydroxylase gene. *Clin. Chem. (Washington)*, 39 (1993) 2354-2355.
- 592 Tsai, M.Y., Suess, P., Schwichtenberg, K., Eckfeldt, J.H., Yuan, J., Tuchman, M. and Hunninghake, D.: Determination of apolipoprotein E genotypes by single-strand conformational polymorphism. *Clin. Chem. (Washington)*, 39 (1993) 2121-2124.
- 593 Wang, G.L. and Goldstein, E.S.: An AP-1 binding site in the upstream region of DEB-A is part of a developmentally regulated negative element. *Biochim. Biophys. Acta*, 1216 (1993) 94-104.
- 594 Wang, J.C. and van Dyke, M.W.: Initiator sequences direct downstream promoter binding by human transcription factor IID. *Biochim. Biophys. Acta*, 1216 (1993) 73-80.
- 595 Williams, P.E.: The analysis of DNA restriction fragments and polymerase chain reaction products by dynamic sieving electrophoresis. *Methods (San Diego)*, 4 (1992) 227-232; *C.A.*, 119 (1993) 109975y.
- 596 Wu, C., Zannis-Hadjopoulos, M. and Price, G.B.: *In vivo* activity for initiation of DNA replication resides in a transcribed region of the human genome. *Biochim. Biophys. Acta*, 1174 (1993) 258-266.
- 597 Yanagida, F., Oshida, A., Shinohara, T. and Goto, S.: (Chromosomal DNA pattern analyses of wine yeasts and wild yeasts by pulsed field gel electrophoresis). *J. Inst. Enol. Vitic., Yamanashi, Univ.*, 27 (1993) 13-19; *C.A.*, 119 (1993) 134824f.
- 598 Yang, M., Allen, H. and DiCioccio, R.A.: Pedigree analysis of  $\alpha$ -L-fucosidase gene mutations in a fucosidosis family. *Biochim. Biophys. Acta*, 1182 (1993) 245-249.
- 599 Yassouridis, A. and Epplen, J.T.: Calculating paternity probabilities from DNA multilocus fingerprints in some cases of deficiency. *Electrophoresis (Weinheim)*, 14 (1993) 978-985.

See also 282, 448, 505, 516.

#### 21f. Complex mixtures of nucleic acids and their fragments

- 600 Orchard, K. and May, G.E.: An EMSA-based method for determining the molecular weight of a protein-DNA complex. *Nucleic Acids Res.*, 21 (1993) 3335-3336; *C.A.*, 119 (1993) 112729b.

#### 22. ALKALOIDS

- 601 Guo, R. and Thormann, W.: Acetylator phenotyping via analysis of four caffeine metabolites in human urine by micellar electrokinetic capillary chromatography with multiwavelength detection. *Electrophoresis (Weinheim)*, 14 (1993) 547-553.

- 602 Johansson, I.M., Grön-Rydberg, M.-B. and Schmekel, B.: Determination of theophylline in plasma using different capillary electrophoretic systems. *J. Chromatogr.*, 652 (1993) 487-493.
- 603 Ma, Y., Meyer, K.G., Afzal, D. and Agena, E.A.: Isolation and quantification of ergovaline from *Festuca arundinacea* (tall fescue) infected with the fungus *Acremonium coenophialum* by high-performance capillary electrophoresis. *J. Chromatogr.*, 652 (1993) 535-538.

See also 7.

## 23. OTHER SUBSTANCES CONTAINING HETEROCYCLIC NITROGEN

### 23a. Porphyrins and other pyrroles

- 604 Kiyohara, C., Saitoh, K. and Suzuki, N.: Micellar electrokinetic capillary chromatography of haematoporphyrin, protoporphyrin and their copper and zinc complexes. *J. Chromatogr.*, 646 (1993) 397-403.

### 23b. Bile pigments

- 605 Harman, A.D., Kibbey, R.G., Sablik, M.A., Fintschenko, Y., Kurtin, W.E. and Bushey, M.M.: Micellar electrokinetic capillary chromatography analysis of the behavior of bilirubin in micellar solutions. *J. Chromatogr.*, 652 (1993) 525-533.

### 23d. Pyridine derivatives

- 606 Janini, G.M., Chan, K.C., Barnes, J.A., Muschik, G.M. and Issaq, H.J.: Separation of pyridinecarboxylic acid isomers and related compounds by capillary zone electrophoresis. Effect of cetyltrimethylammonium bromide on electroosmotic flow and resolution. *J. Chromatogr.*, 653 (1993) 321-327.

### 23e. Other N-heterocyclic compounds

- 607 Esaka, Y., Yamaguchi, Y., Kano, K. and Goto, M.: Separation of amino acid-oxazole derivatives of the redox coenzyme pyrroloquinoline quinone by capillary zone electrophoresis. *J. Chromatogr.*, 652 (1993) 225-232.

## 24. ORGANIC SULPHUR COMPOUNDS (INCL. GLUCOSINOLATES)

See 639.

## 25. ORGANIC PHOSPHORUS COMPOUNDS (INCL. SUGAR PHOSPHATES)

See 172, 173, 174, 182, 184, 187, 400, 401.

## 26. ORGANOMETALLIC AND RELATED COMPOUNDS

### 26a. Organometallic compounds

- 608 Ng, C.L., Lee, H.K. and Li, S.F.Y.: Determination of organolead and organoselenium compounds by micellar electrokinetic chromatography. *J. Chromatogr.*, 652 (1993) 547-553.

### 26c. Coordination compounds

- 609 Iki, N. and Yotsuyanagi, T.: (High-performance capillary electrophoresis of metal complexes). *Kagaku (Kyoto)*, 48 (1993) 434-435; *C.A.*, 119 (1993) 130425d - a review with 18 refs.
- 610 Iki, N., Hoshino, H. and Yotsuyanagi, T.: Ion-association capillary electrophoresis. New separation mode for equally and highly charged metal chelates. *J. Chromatogr.*, 652 (1993) 539-546.

## 27. VITAMINS AND VARIOUS ANIMAL GROWTH FACTORS (NON-PEPTIDIC)

- 611 Bempong, D.K., Honigberg, I.L. and Meltzer, N.M.: Separation of 13-*cis* and all-*trans* retinoic acid and their photodegradation products using capillary zone electrophoresis and micellar electrokinetic chromatography (MEC). *J. Pharm. Biomed. Anal.*, 11 (1993) 829-833.
- 612 Chan, K.C., Lewis, K.C., Phang, J.M. and Issaq, H.J.: Separation of retinoic acid isomers using micellar electrokinetic chromatography. *J. High Resolut. Chromatogr.*, 16 (1993) 560-561.
- 613 Chiari, M., Nesi, M., Carrea, G. and Righetti, P.G.: Determination of total vitamin C in fruits by capillary zone electrophoresis. *J. Chromatogr.*, 645 (1993) 197-200.
- 614 Jegle, U.: Separation of water-soluble vitamins via high-performance capillary electrophoresis. *J. Chromatogr.*, 652 (1993) 495-501.

See also 359.

## 28. ANTIBIOTICS

- 615 Altria, K.D. and Chanter, Y.L.: Validation of a capillary electrophoresis method for the determination of a quinolone antibiotic and its related impurities. *J. Chromatogr.*, 652 (1993) 459-463.

## 29. INSECTICIDES, PESTICIDES AND OTHER AGROCHEMICALS

### 29g. Other types of pesticides and various agrochemicals

See 151.

## 30. SYNTHETIC AND NATURAL DYES

## 30a. Synthetic dyes

- 616 Croft, S.N. and Hinks, D.: Analysis of dyes by capillary electrophoresis. *J. Soc. Dyers Colour.*, 108 (1992) 546-551; *C.A.*, 119 (1993) 119477b.

## 30b. Chloroplast and other natural pigments

- 617 Tomlinson, A.J., Landers, J.P., Lewis, I.A.S. and Naylor, S.: Buffer conditions affecting the separation of Maillard reaction products by capillary electrophoresis. *J. Chromatogr.*, 652 (1993) 171-177.

## 31. PLASTICS AND THEIR INTERMEDIATES

- 618 Bullock, J.: Application of capillary electrophoresis to the analysis of the oligomeric distribution of polydisperse polymers. *J. Chromatogr.*, 645 (1993) 169-177.

## 32. DRUG ANALYSIS

## 32a. Drug analysis, general techniques

- 619 Altria, K.D.: Optimization and improvement of sensitivity in capillary electrophoresis for quantitation of selected pharmaceuticals. *LC-GC*, 11 (1993) 438-442; *C.A.*, 119 (1993) 103512y.
- 620 Altria, K.D.: Quantitative aspects of the application of capillary electrophoresis to the analysis of pharmaceuticals and drug related impurities. *J. Chromatogr.*, 646 (1993) 245-257 - a review with 91 refs.
- 621 Heuermann, M. and Blaschke, G.: Chiral separation of basic drugs using cyclodextrins as chiral pseudo-stationary phases in capillary electrophoresis. *J. Chromatogr.*, 648 (1993) 267-274.
- 622 Kostianen, R., Franssen, E.J.F. and Bruins, A.P.: Capillary zone electrophoresis-ionspray mass spectrometry of a synthetic drug-protein conjugate mixture. *J. Chromatogr.*, 647 (1993) 361-365.
- 623 Lee, T.T., Lillard, S.J. and Yeung, E.S.: Screening and characterization of biopharmaceuticals by high-performance capillary electrophoresis with laser-induced native fluorescence detection. *Electrophoresis (Weinheim)*, 14 (1993) 429-438.
- 624 Steur, W.: A critical review of high-performance liquid chromatography, supercritical fluid chromatography and capillary zone electrophoresis in drug analysis. *Paperback APV*, 32 (1993) 95-105; *C.A.*, 119 (1993) 146685z - a review with 18 refs.
- 625 Tomita, M. and Okuyama, T.: (Analysis of drugs by capillary electrophoresis: effect of methanol in sample solution on the electropherogram). *Hochudoku*, 11 (1993) 140-141; *C.A.*, 119 (1993) 133008f.

See also 45, 55, 91, 633.

## 32b. Antirheumatics and antiinflammatory drugs

- 626 Kanda, S., Tsuchiya, M., Imai, K., Nishio, A. and Yoda, R.: (Rapid determination of aspirin in aspirin ointment by capillary electrophoresis). *Byoin Yakugaku*, 19 (1993) 244-247; *C.A.*, 119 (1993) 188726u.

## 32c. Autonomic and cardiovascular drugs

- 627 Lukkari, P., Vuorela, H. and Riekkola, M.-L.: Effect of buffer solution pH on the elution and separation of  $\beta$ -blockers by micellar electrokinetic capillary chromatography. *J. Chromatogr.*, 652 (1993) 451-457.

## 32d. Central nervous system drugs

- 628 Altria, K.D. and Conolly, P.C.: On-line solution stability determination of pharmaceuticals by capillary electrophoresis. *Chromatographia*, 37 (1993) 176-178.
- 629 Tomlinson, A.J., Benson, L.M., Landers, J.P., Scanlan, G.F., Fang, J., Gorrod, J.W. and Naylor, S.: Investigation of the metabolism of the neuroleptic drug haloperidol by capillary electrophoresis. *J. Chromatogr.*, 652 (1993) 417-426.
- 630 Wolfisberg, H., Schmutz, A., Stotzer, R. and Thormann, W.: Assessment of automated capillary electrophoresis for therapeutic and diagnostic drug monitoring: determination of bupivacaine in drain fluid and antipyrine in plasma. *J. Chromatogr.*, 652 (1993) 407-416.

See also 163.

## 32g. Other drug categories

- 631 Jumppanen, J., Sirén, H. and Riekkola, M.-L.: Screening for diuretics in urine and blood serum by capillary zone electrophoresis. *J. Chromatogr.*, 652 (1993) 441-450.

See also 251.

## 32h. Toxicological and forensic applications

See 599, 601.

## 32i. Plant extracts

- 632 Chen, H.-R. and Sheu, S.-J.: Determination of glycyrrhizin and glycyrrhetic acid in traditional Chinese medicinal preparations by capillary electrophoresis. *J. Chromatogr.*, 653 (1993) 184-188.

See also 603.

## 33. CLINICO-CHEMICAL APPLICATIONS

## 33a. General papers and reviews

See 72.



- 33b. *Complex mixtures and profiling (single compounds by cross-reference only)*
- 633 Kim, N.J., Moon, D.C., Heo, G.S. and Lee, K.J.: Application of micellar electrokinetic chromatography for therapeutic drug monitoring. *Int. Congr. Ser.-Excerpta Med.*, 991(Progress in Clinical Biochemistry) (1992) 791-793; C.A., 119 (1993) 85265p.
- See also 39, 117, 135, 139, 140, 145, 148, 163, 280, 285, 287, 288, 292, 312, 345, 362, 388, 399, 407, 416, 417, 426, 504, 509, 552, 555, 573, 591, 592.
34. FOOD ANALYSIS
- 34a. *General papers and reviews*
- See 101.
- 34b. *Complex mixtures (single compounds by cross-reference only)*
- 634 Cancelon, P.F. and Bryan, C.R.: Use of capillary electrophoresis for monitoring citrus juice composition. *J. Chromatogr.*, 652 (1993) 555-561.
- 635 Lalljie, S.P.D., Vindevogel, J. and Sandra, P.: Quantitation of organic acids in sugar refinery juices with capillary zone electrophoresis and indirect UV detection. *J. Chromatogr.*, 652 (1993) 563-569.
- See also 130, 131, 257, 301, 319, 613, 644, 647.
- 34c. *Organoleptically important compounds (flavors, odors, volatiles)*
- 636 Khaled, M.Y., Anderson, M.R. and McNair, H.M.: Micellar electrokinetic capillary chromatography of pungent compounds using simultaneous on-line ultraviolet and electrochemical detection. *J. Chromatogr. Sci.*, 31 (1993) 259-264.
35. ENVIRONMENTAL ANALYSIS
- 35a. *General papers and reviews*
- See 88.
- 35c. *Water pollution (complex mixtures; single compounds by cross-reference only)*
- 637 Bachmann, K., Haag, I., Prokop, T., Röder, A. and Wagner, P.: Chromatographic methods for the analysis of size-classified and individual raindrops. *J. Chromatogr.*, 643 (1993) 181-188.
- 638 Ratnaweera, H., Fettig, J. and Oedegaard, H.: Particle and phosphate removal mechanismus with prepolymerized coagulants. In: Klute, R. and Hahn, H.H. (Editors), *Chem. Water Waste-water Treat. II, Proc. Gothenburg Symp., 5th*, Springer, Berlin, 1992, pp. 3-17; C.A., 119 (1993) 167026a.
- See also 88, 98, 128.
- 35d. *Soil pollution (complex mixtures; single compounds by cross-reference only)*
- See 128.
36. SOME TECHNICAL PRODUCTS AND COMPLEX MIXTURES
- 36a. *Surfactants*
- 639 Chen, S. and Pietrzyk, D.J.: Separation of sulfonate and sulfate surfactants by capillary electrophoresis: effect of buffer cation. *Anal. Chem.*, 65 (1993) 2770-2775.
- See also 649.
- 36c. *Complex mixtures, technical products and unidentified compounds*
- 640 Kos, T., Hoffmann-Sommergruber, K., Ferreira, F., Hirschwehr, R., Ahorn, H., Horak, F., Jager, S., Sperr, W., Kraft, D. and Scheiner, O.: Purification, characterization and N-terminal amino acid sequence of a new major allergen from european chestnut pollen Cas s 1. *Biochem. Biophys. Res. Commun.*, 196 (1993) 1086-1092.
37. CELLS, CELLULAR PARTICLES AND SUPRAMOLECULAR STRUCTURES
- See 533.
38. INORGANIC COMPOUNDS
- 38a. *Cations*
- 641 Groh, T. and Bächmann, K.: Investigation of vanadate as a pH sensitive analyte anion using capillary zone electrophoresis. *J. Chromatogr.*, 646 (1993) 405-410.
- 642 Huggins, T.G. and Henion, J.D.: Capillary electrophoresis/mass spectrometry determination of inorganic ions using an ion spray-sheath flow interface. *Electrophoresis (Weinheim)*, 14 (1993) 531-539.
- 643 Jones, B. and Takahashi, Y.: (The separation of inorganic metal cations by capillary electrophoresis). *Kankyo Kagaku*, 3 (1993) 424-425; C.A., 119 (1993) 108045h.
- 644 Kajiwara, H., Sato, A. and Kaneko, S.: Analysis of calcium and magnesium ions in wheat flour by capillary zone electrophoresis. *Biosci., Biotechnol., Biochem.*, 57 (1993) 1010-1011; C.A., 119 (1993) 115672p.
- 645 Nasi, J.C., Pereira, E.C., Trindale, W.S. and Ignacio, I.G.: (Electrophoresis of ions: yttrium, lanthanum, cerium(III) and praseodymium). *Rev. Bras. Farm.*, 72 (1992) 89-91; C.A., 119 (1993) 108037g.
- 646 Riviello, J.M. and Harrold, M.P.: Capillary electrophoresis of inorganic cations and low-molecular-mass amines using a copper-based electrolyte with indirect UV detection. *J. Chromatogr.*, 652 (1993) 385-392.

## 38b. Anions

- 647 Cortes, M.A. and Martin, Y.: (Determination of nitrates and nitrites in cheeses by capillary ionic electrophoresis). *Tec. Lab.*, 15 (1993) 267-269; *C.A.*, 119 (1993) 137729q.
- 648 Kumooka, Y., Nakamura, J. and Kishi, T.: (Analysis of anions in explosive residues by capillary electrophoresis). *Kagaku Keisatsu Kenkyusho Hokoku, Hokagaku-hen*, 45 (1992) 159-161; *C.A.*, 119 (1993) 163572c.
- 649 Pretswell, E.L., Morrison, A.R. and Park, J.S.: Comparison of capillary zone electrophoresis with standard gravimetric analysis and ion chromatography for the determination of inorganic anions in detergent matrices. *Analyst (London)*, 118 (1993) 1265-1267.

See also 642.

## PUBLICATION SCHEDULE FOR THE 1994 SUBSCRIPTION

*Journal of Chromatography A and Journal of Chromatography B: Biomedical Applications*

MONTH	O 1993	N 1993	D 1993	J	F	M	A	
Journal of Chromatography A	652/1 652/2 653/1	653/2 654/1 654/2 655/1	655/2 656/1 + 2 657/1 657/2	658/1 658/2 659/1 659/2	660/1 + 2 661/1 + 2 662/1 662/2	663/1 663/2 664/1	664/2 665/1 665/2 666/1 + 2 667/1	The publication schedule for further issues will be published later.
Bibliography Section						681/1		
Journal of Chromatography B: Biomedical Applications				652/1	652/2 653/1	653/2 654/1	654/2 655/1	

### INFORMATION FOR AUTHORS

(Detailed *Instructions to Authors* were published in *J. Chromatogr. A*, Vol. 657, pp. 463–469. A free reprint can be obtained by application to the publisher, Elsevier Science B.V., P.O. Box 330, 1000 AH Amsterdam, Netherlands.)

**Types of Contributions.** The following types of papers are published: Regular research papers (full-length papers), Review articles, Short Communications and Discussions. Short Communications are usually descriptions of short investigations, or they can report minor technical improvements of previously published procedures; they reflect the same quality of research as full-length papers, but should preferably not exceed five printed pages. Discussions (one or two pages) should explain, amplify, correct or otherwise comment substantively upon an article recently published in the journal. For Review articles, see inside front cover under Submission of Papers.

**Submission.** Every paper must be accompanied by a letter from the senior author, stating that he/she is submitting the paper for publication in the *Journal of Chromatography A* or *B*.

**Manuscripts.** Manuscripts should be typed in **double spacing** on consecutively numbered pages of uniform size. The manuscript should be preceded by a sheet of manuscript paper carrying the title of the paper and the name and full postal address of the person to whom the proofs are to be sent. As a rule, papers should be divided into sections, headed by a caption (e.g., Abstract, Introduction, Experimental, Results, Discussion, etc.). All illustrations, photographs, tables, etc., should be on separate sheets.

**Abstract.** All articles should have an abstract of 50–100 words which clearly and briefly indicates what is new, different and significant. No references should be given.

**Introduction.** Every paper must have a concise introduction mentioning what has been done before on the topic described, and stating clearly what is new in the paper now submitted.

**Experimental conditions** should preferably be given on a *separate* sheet, headed "Conditions". These conditions will, if appropriate, be printed in a block, directly following the heading "Experimental".

**Illustrations.** The figures should be submitted in a form suitable for reproduction, drawn in Indian ink on drawing or tracing paper. Each illustration should have a caption, all the *captions* being typed (with double spacing) together on a *separate sheet*. If structures are given in the text, the original drawings should be provided. Coloured illustrations are reproduced at the author's expense, the cost being determined by the number of pages and by the number of colours needed. The written permission of the author and publisher must be obtained for the use of any figure already published. Its source must be indicated in the legend.

**References.** References should be numbered in the order in which they are cited in the text, and listed in numerical sequence on a separate sheet at the end of the article. Please check a recent issue for the layout of the reference list. Abbreviations for the titles of journals should follow the system used by *Chemical Abstracts*. Articles not yet published should be given as "in press" (journal should be specified), "submitted for publication" (journal should be specified), "in preparation" or "personal communication".

Vols. 1–651 of the *Journal of Chromatography*; *Journal of Chromatography, Biomedical Applications* and *Journal of Chromatography, Symposium Volumes* should be cited as *J. Chromatogr.* From Vol. 652 on, *Journal of Chromatography A* (incl. Symposium Volumes) should be cited as *J. Chromatogr. A* and *Journal of Chromatography B: Biomedical Applications* as *J. Chromatogr. B*.

**Dispatch.** Before sending the manuscript to the Editor please check that the envelope contains four copies of the paper complete with references, captions and figures. One of the sets of figures must be the originals suitable for direct reproduction. Please also ensure that permission to publish has been obtained from your institute.

**Proofs.** One set of proofs will be sent to the author to be carefully checked for printer's errors. Corrections must be restricted to instances in which the proof is at variance with the manuscript.

**Reprints.** Fifty reprints will be supplied free of charge. Additional reprints can be ordered by the authors. An order form containing price quotations will be sent to the authors together with the proofs of their article.

**Advertisements.** The Editors of the journal accept no responsibility for the contents of the advertisements. Advertisement rates are available on request. Advertising orders and enquiries can be sent to the Advertising Manager, Elsevier Science B.V., Advertising Department, P.O. Box 211, 1000 AE Amsterdam, Netherlands; courier shipments to: Van de Sande Bakhuyzenstraat 4, 1061 AG Amsterdam, Netherlands; Tel. (+31-20) 515 3220/515 3222, Telefax (+31-20) 6833 041, Telex 16479 els vi nl. UK: T.G. Scott & Son Ltd., Tim Blake, Portland House, 21 Narborough Road, Cosby, Leics. LE9 5TA, UK; Tel. (+44-533) 753 333, Telefax (+44-533) 750 522. USA and Canada: Weston Media Associates, Daniel S. Lipner, P.O. Box 1110, Greens Farms, CT 06436-1110, USA; Tel. (+1-203) 261 2500, Telefax (+1-203) 261 0101.

# Analytical Applications of Circular Dichroism

Edited by **N. Purdie** and **H.G. Brittain**

Techniques and Instrumentation in Analytical Chemistry Volume 14

Circular dichroism is a special technique which provides unique information on dissymmetric molecules. Such compounds are becoming increasingly important in a wide variety of fields, such as natural products chemistry, pharmaceuticals, molecular biology, etc. The content of this book has been selected in order to feature the unique aspects of circular dichroism, and how these strengths can be of assistance to workers in the field.

Substantial discussions have been provided regarding the particular phenomena associated with dissymmetric compounds which give rise to the circular dichroism effect. Reviews are also given of the type of instrumentation available for the measurement of these effects. A number of chapters cover the wide range of applications illustrating the power of the method.

Owing to its broad appeal, the book will be of interest to workers in all areas of chemistry and pharmaceutical science.

## Contents:

1. Introduction to chiroptical phenomena (H.G. Brittain).
2. Instrumentation for the measurement of circular dichroism; past, present and future developments (D.R. Bobbitt).
3. Instrumental methods of infrared and Raman vibrational optical activity (L.A. Nafie *et al.*).
4. Application of infrared CD to the analysis of the solution conformation of biological molecules (M. Diem).
5. Determination of absolute configuration by CD. Applications of the octant rule and the exciton chirality rule (D.A. Lightner).
6. Analysis of protein structure by circular dichroism spectroscopy (J.F. Towell III, M.C. Manning).
7. Chiroptical studies of molecules in electronically

- excited states (J.P. Riehl).
  8. Analytical applications of CD to forensic, pharmaceutical, clinical, and food sciences (N. Purdie).
  9. The use of circular dichroism as a liquid chromatographic detector (A. Gergely).
  10. Applications of circular dichroism spectropolarimetry to the determination of steroids (A. Gergely).
  11. Circular dichroism studies of the optical activity induced in achiral molecules through association with chiral substances (H.G. Brittain).
- Subject index.

© 1994 360 pages Hardbound  
Price: Dfl. 355.00 (US \$ 202.75)  
ISBN 0-444-89508-6

## ORDER INFORMATION

For USA and Canada  
**ELSEVIER SCIENCE INC.**

P.O. Box 945  
Madison Square Station  
New York, NY 10160-0757  
Fax: (212) 633 3880

In all other countries  
**ELSEVIER SCIENCE B.V.**

P.O. Box 330  
1000 AH Amsterdam  
The Netherlands  
Fax: (+31-20) 5862 845

US\$ prices are valid only for the USA & Canada and are subject to exchange rate fluctuations; in all other countries the Dutch guilder price (Dfl.) is definitive. Customers in the European Community should add the appropriate VAT rate applicable in their country to the price(s). Books are sent postfree if prepaid.



**ELSEVIER**  
**SCIENCE** B.V.



0021-9673(19940318)681:1;1-0