

Bibliography Section**Please note**

The pagination of Vol. 682 continues
from that of Vol. 681.

JOURNAL OF

CHROMATOGRAPHY A

INCLUDING ELECTROPHORESIS AND OTHER SEPARATION METHODS

EDITORS

U.A.Th. Brinkman (Amsterdam)
R.W. Giese (Boston, MA)
J.K. Haken (Kensington, N.S.W.)
L.R. Snyder (Orinda, CA)

EDITORS, SYMPOSIUM VOLUMES,

E. Heftmann (Orinda, CA), Z. Deyl (Prague)

EDITORIAL BOARD

D.W. Armstrong (Rolla, MO)
W.A. Aue (Halifax)
P. Boček (Brno)
A.A. Boulton (Saskatoon)
P.W. Carr (Minneapolis, MN)
N.H.C. Cooke (San Ramon, CA)
V.A. Davankov (Moscow)
G.J. de Jong (Weesp)
Z. Deyl (Prague)
S. Dilli (Kensington, N.S.W.)
Z. El Rassi (Stillwater, OK)
H. Engelhardt (Saarbrücken)
F. Erni (Basle)
M.B. Evans (Hatfield)
J.L. Glajch (N. Billerica, MA)
G.A. Guiochon (Knoxville, TN)
P.R. Haddad (Hobart, Tasmania)
I.M. Hais (Hradec Králové)
W.S. Hancock (Palo Alto, CA)
S. Hjertén (Uppsala)
S. Honda (Higashi-Osaka)
Cs. Horváth (New Haven, CT)
J.F.K. Huber (Vienna)
K.-P. Hupe (Waldbronn)
J. Janák (Brno)
P. Jandera (Pardubice)
B.L. Karger (Boston, MA)
J.J. Kirkland (Newport, DE)
E. sz. Kováts (Lausanne)
K. Macek (Prague)
A.J.P. Martin (Cambridge)
L.W. McLaughlin (Chestnut Hill, MA)
E.D. Morgan (Keele)
J.D. Pearson (Kalamazoo, MI)
H. Poppe (Amsterdam)
F.E. Regnier (West Lafayette, IN)
P.G. Righetti (Milan)
P. Schoenmakers (Amsterdam)
R. Schwarzenbach (Dübendorf)
R.E. Shoup (West Lafayette, IN)
R.P. Singhai (Wichita, KS)
A.M. Siouffi (Marseille)
D.J. Strydom (Boston, MA)
N. Tanaka (Kyoto)
S. Terabe (Hyogo)
K.K. Unger (Mainz)
R. Verpoorte (Leiden)
Gy. Vigh (College Station, TX)
J.T. Watson (East Lansing, MI)
B.D. Westerlund (Uppsala)

EDITORS, BIBLIOGRAPHY SECTION

Z. Deyl (Prague), J. Janák (Brno), V. Schwarz (Prague)

ELSEVIER

JOURNAL OF CHROMATOGRAPHY A

INCLUDING ELECTROPHORESIS AND OTHER SEPARATION METHODS

Scope. The *Journal of Chromatography A* publishes papers on all aspects of **chromatography, electrophoresis** and related methods. Contributions consist mainly of research papers dealing with chromatographic theory, instrumental developments and their applications. In the *Symposium volumes*, which are under separate editorship, proceedings of symposia on chromatography, electrophoresis and related methods are published. *Journal of Chromatography B: Biomedical Applications* —This journal, which is under separate editorship, deals with the following aspects: developments in and applications of chromatographic and electrophoretic techniques related to clinical diagnosis or alterations during medical treatment; screening and profiling of body fluids or tissues related to the analysis of active substances and to metabolic disorders; drug level monitoring and pharmacokinetic studies; clinical toxicology; forensic medicine; veterinary medicine; occupational medicine; results from basic medical research with direct consequences in clinical practice.

Submission of Papers. The preferred medium of submission is on disk with accompanying manuscript (see *Electronic manuscripts* in the Instructions to Authors, which can be obtained from the publisher, Elsevier Science B.V., P.O. Box 330, 1000 AH Amsterdam, Netherlands). Manuscripts (in English; four copies are required) should be submitted to: Editorial Office of *Journal of Chromatography A*, P.O. Box 681, 1000 AR Amsterdam, Netherlands, Telefax (+31-20) 5862 304, or to: The Editor of *Journal of Chromatography B: Biomedical Applications*, P.O. Box 681, 1000 AR Amsterdam, Netherlands. Review articles are invited or proposed in writing to the Editors who welcome suggestions for subjects. An outline of the proposed review should first be forwarded to the Editors for preliminary discussion prior to preparation. Submission of an article is understood to imply that the article is original and unpublished and is not being considered for publication elsewhere. For copyright regulations, see below.

Publication information. *Journal of Chromatography A* (ISSN 0021-9673): for 1994 Vols. 652–682 are scheduled for publication. *Journal of Chromatography B: Biomedical Applications* (ISSN 0378-4347): for 1994 Vols. 652–662 are scheduled for publication. Subscription prices for *Journal of Chromatography A*, *Journal of Chromatography B: Biomedical Applications* or a combined subscription are available upon request from the publisher. Subscriptions are accepted on a prepaid basis only and are entered on a calendar year basis. Issues are sent by surface mail except to the following countries where air delivery via SAL is ensured: Argentina, Australia, Brazil, Canada, China, Hong Kong, India, Israel, Japan, Malaysia, Mexico, New Zealand, Pakistan, Singapore, South Africa, South Korea, Taiwan, Thailand, USA. For all other countries airmail rates are available upon request. Claims for missing issues must be made within six months of our publication (mailing) date. Please address all your requests regarding orders and subscription queries to: Elsevier Science B.V., Journal Department, P.O. Box 211, 1000 AE Amsterdam, Netherlands. Tel.: (+31-20) 5803 642; Fax: (+31-20) 5803 598. Customers in the USA and Canada wishing information on this and other Elsevier journals, please contact Journal Information Center, Elsevier Science Inc., 655 Avenue of the Americas, New York, NY 10010, USA, Tel. (+1-212) 633 3750, Telefax (+1-212) 633 3764.

Abstracts/Contents Lists published in Analytical Abstracts, Biochemical Abstracts, Biological Abstracts, Chemical Abstracts, Chemical Titles, Chromatography Abstracts, Current Awareness in Biological Sciences (CABS), Current Contents/Life Sciences, Current Contents/Physical, Chemical & Earth Sciences, Deep-Sea Research/Part B: Oceanographic Literature Review, Excerpta Medica, Index Medicus, Mass Spectrometry Bulletin, PASCAL-CNRS, Referativnyi Zhurnal, Research Alert and Science Citation Index.

US Mailing Notice. *Journal of Chromatography A* (ISSN 0021-9673) is published weekly (total 52 issues) by Elsevier Science B.V., (Sara Burgerhartstraat 25, P.O. Box 211, 1000 AE Amsterdam, Netherlands). Annual subscription price in the USA US\$ 4994.00 (US\$ price valid in North, Central and South America only) including air speed delivery. Second class postage paid at Jamaica, NY 11431. **USA POSTMASTERS:** Send address changes to *Journal of Chromatography A*, Publications Expediting, Inc., 200 Meacham Avenue, Elmont, NY 11003. Airfreight and mailing in the USA by Publications Expediting.

See inside back cover for Publication Schedule, Information for Authors and information on Advertisements.

© 1994 ELSEVIER SCIENCE B.V. All rights reserved.

0021-9673/94/\$07.00

No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, without the prior written permission of the publisher. Elsevier Science B.V., Copyright and Permissions Department, P.O. Box 521, 1000 AM Amsterdam, Netherlands.

Upon acceptance of an article by the journal, the author(s) will be asked to transfer copyright of the article to the publisher. The transfer will ensure the widest possible dissemination of information.

Special regulations for readers in the USA – This journal has been registered with the Copyright Clearance Center, Inc. Consent is given for copying of articles for personal or internal use, or for the personal use of specific clients. This consent is given on the condition that the copier pays through the Center the per-copy fee stated in the code on the first page of each article for copying beyond that permitted by Sections 107 or 108 of the US Copyright Law. The appropriate fee should be forwarded with a copy of the first page of the article to the Copyright Clearance Center, Inc., 27 Congress Street, Salem, MA 01970, USA. If no code appears in an article, the author has not given broad consent to copy and permission to copy must be obtained directly from the author. The fee indicated on the first page of an article in this issue will apply retroactively to all articles published in the journal, regardless of the year of publication. This consent does not extend to other kinds of copying, such as for general distribution, resale, advertising and promotion purposes, or for creating new collective works. Special written permission must be obtained from the publisher for such copying.

No responsibility is assumed by the Publisher for any injury and/or damage to persons or property as a matter of products liability, negligence or otherwise, or from any use or operation of any methods, products, instructions or ideas contained in the materials herein. Because of rapid advances in the medical sciences, the Publisher recommends that independent verification of diagnoses and drug dosages should be made.

Although all advertising material is expected to conform to ethical (medical) standards, inclusion in this publication does not constitute a guarantee or endorsement of the quality or value of such product or of the claims made of it by its manufacturer.

Ⓢ The paper used in this publication meets the requirements of ANSI/NISO Z39.48-1992 (Permanence of Paper).

Printed in the Netherlands

Environmental Analysis

Techniques, Applications and Quality Assurance

Edited by **D. Barceló**

Techniques and Instrumentation in Analytical Chemistry Volume 13

Three aspects of environmental analysis are treated in this book:

- the use of various analytical techniques
- their applications to trace analysis of pollutants, mainly organic compounds
- quality assurance aspects, including the use of certified reference materials for quality control of the entire analytical process.

The book will serve as a general reference for post-graduate students as well as a practical reference for environmental chemists who need to use the analytical techniques for environmental studies. Analytical chemists needing information on the complexity of environmental sample matrices and interferences will also find this an invaluable reference.

Contents: Part 1. Field Sampling Techniques and Sample Preparation.

1. Sampling techniques for air pollutants (R. Niessner). 2. Sample handling strategies for the analysis of organic contaminants from environmental samples (M.-C. Hennion, P. Scribe). 3. Extraction, clean-up and recoveries of persistent trace organic contaminants from sediment and biota samples (D.E. Wells).

Part 2. Application Areas.

4. Current developments in the analysis of polychlorinated biphenyls (PCBs) including planar

and other toxic metabolites in environmental matrices (D.E. Wells). 5. Official methods of analysis of priority pesticides in water using gas chromatographic techniques (D. Barceló). 6. Coupled-column reversed phase liquid chromatography as a versatile technique for the determination of polar pesticides (E.A. Hogendoorn, P. van Zoonen). 7. Liquid chromatographic determination of phenols and substituted derivatives in water samples (G. Marko-Varga). 8. HPLC methods for the determination of mycotoxins and phycotoxins (J.F. Lawrence, P.M. Scott). 9. Determination of radionuclides in environmental samples (V. Valkovic).

Part 3. Quality Assurance and Reference Materials.

10. Quality assurance in environmental analysis (W.P. Cofino). 11. Certified reference materials for the quality control of measurements in environmental monitoring (E.A. Maier). 12. Standard reference materials for the determination of trace organic constituents in environmental samples (S.A. Wise).

Part 4. Emerging Techniques.

13. Application of fluorescence spectroscopic techniques in the determination of PAHs and PAH metabolites (F. Ariese, C. Gooijer, N.H. Velthorst). 14. Characterization of surfactants in water by desorption ionization methods (F. Ventura). 15. Utilization of various LC-MS interfacing systems in environmental analysis; application to polar pesticides (M.H. Lamoree, R.T. Ghijsen, U.A.Th. Brinkman). 16. Hyphenated techniques applied to the speciation of organometallic compounds in the environment (O.F.X. Donard, R. Ritsema). 17. The potential of capillary electrophoresis in environmental analysis (M.W.F. Nielen). Subject index.

© 1993 660 pages Hardbound
Price: Dfl. 465.00 (US \$ 265.75)
ISBN 0-444-89648-1

ORDER INFORMATION

For USA and Canada
ELSEVIER SCIENCE INC.

P.O. Box 945
Madison Square Station
New York, NY 10160-0757
Fax: (212) 633 3880


In all other countries
ELSEVIER SCIENCE B.V.

P.O. Box 330
1000 AH Amsterdam
The Netherlands
Fax: (+31-20) 5862 845

US\$ prices are valid only for the USA & Canada and are subject to exchange rate fluctuations; in all other countries the Dutch guilder price (Dfl.) is definitive. Customers in the European Community should add the appropriate VAT rate applicable in their country to the price(s). Books are sent postfree if prepaid.



**ELSEVIER
SCIENCE** B.V.



AUDIENCE

All Mass Spectrometrists and those interested in the chemistry of charged particles.



Elsevier Science B.V.

Attn. Eugene P.M. Wijnhoven
P.O. Box 330, 1000 AH Amsterdam
The Netherlands
Fax: +31 (20) 586 2845
e-mail: e.wijnhoven@elsevier.nl

In the USA & Canada

Attn. Judy Weislogel
P.O. Box 945, Madison Square Station
New York, NY 10159-0945, USA
Fax: +1 (212) 633 5800
e-mail: j.weislogel@elsevier.com

INTERNATIONAL JOURNAL OF MASS SPECTROMETRY AND ION PROCESSES

Editors

M.T. Bowers, *University of California, Department of Chemistry, Santa Barbara, CA 93106, USA,*

H. Schwarz, *Technical University, Department of Chemistry, Strasse des 17 Juni 135, W 1000 Berlin 12, Germany*

J.F.J. Todd, *University of Kent at Canterbury, University Chemical Laboratory, Canterbury, Kent CT2 7NH, UK*

AIMS AND SCOPE

The journal contains papers dealing with fundamental aspects of mass spectrometry and ion processes, and the study of the application of mass spectrometric techniques to specific problems. The following topics, amongst others, are included:

- ❖ Theoretical and experimental studies of ion formation, ion separation and ion detection processes.
- ❖ The design and performance of instruments and accessories.
- ❖ Measurements of natural isotopic abundances, precise isotopic masses, ionization, appearance and excitation energies, ionization cross-sections.
- ❖ Development of techniques related to determining molecular structures, geological age determination, studies of thermodynamic properties, chemical kinetics, surface phenomena, radiation chemistry, and chemical analyses.
- ❖ Theory of mass spectra, application of computer techniques to mass spectral data.
- ❖ Chemistry and physics of cluster ions.
- ❖ Spectroscopy of gaseous ions including studies related to interstellar chemistry.
- ❖ Mechanistic studies of unimolecular processes and ion/molecule reactions in the gas phase including computational aspects.
- ❖ Physical organic chemistry of isolated ions.
- ❖ Biological aspects of mass spectrometry.

ABSTRACTED/INDEXED IN

Chemical Abstracts, Current Contents: Physical, Chemical & Earth Sciences, Excerpta Medica, INSPEC, Mass Spectrometry Bulletin, PASCAL/CNRS.

1995 SUBSCRIPTION DATA

Volumes 141-151 (in 33 issues)

Subscription price: Dfl. 4620.00 (US\$2432.00) ISSN 0168-1176

I would like a free sample copy of **International Journal of Mass Spectrometry and Ion Processes**
 Instructions to Authors.
 information on (personal) subscription rates
 to enter a subscription for 1995. Please send me a Proforma Invoice

Name _____

Address _____

Tel. _____

Fax _____

CONTENTS

Bibliography Section

Liquid Column Chromatography	B239
Gas Chromatography	B294
Planar Chromatography	B323
Electrophoresis	B332

ห้องสมุดรวมวิทยาศาสตร์บริการ

๑๑ ต.ค. 2537

JOURNAL OF CHROMATOGRAPHY A

VOL. 682 (1994)

JOURNAL OF CHROMATOGRAPHY A

INCLUDING ELECTROPHORESIS AND OTHER SEPARATION METHODS

EDITORS

U.A.Th. BRINKMAN (Amsterdam), R.W. GIESE (Boston, MA), J.K. HAKEN (Kensington, N.S.W.),
L.R. SNYDER (Orinda, CA)

EDITORS, SYMPOSIUM VOLUMES

E. HEFTMANN (Orinda, CA), Z. DEYL (Prague)

EDITORIAL BOARD

D.W. Armstrong (Rolla, MO), W.A. Aue (Halifax), P. Boček (Brno), A.A. Boulton (Saskatoon), P.W. Carr (Minneapolis, MN), N.H.C. Cooke (San Ramon, CA), V.A. Davankov (Moscow), G.J. de Jong (Weesp), Z. Deyl (Prague), S. Dilli (Kensington, N.S.W.), Z. El Rassi (Stillwater, OK), H. Engelhardt (Saarbrücken), F. Erni (Basle), M.B. Evans (Hatfield), J.L. Glajch (N. Billerica, MA), G.A. Guiochon (Knoxville, TN), P.R. Haddad (Hobart, Tasmania), I.M. Hais (Hradec Králové), W.S. Hancock (Palo Alto, CA), S. Hjertén (Uppsala), S. Honda (Higashi-Osaka), Cs. Horváth (New Haven, CT), J.F.K. Huber (Vienna), K.-P. Hupe (Waldbronn), J. Janák (Brno), P. Jandera (Pardubice), B.L. Karger (Boston, MA), J.J. Kirkland (Newport, DE), E. sz. Kováts (Lausanne), K. Macek (Prague), A.J.P. Martin (Cambridge), L.W. McLaughlin (Chestnut Hill, MA), E.D. Morgan (Keele), J.D. Pearson (Kalamazoo, MI), H. Poppe (Amsterdam), F.E. Regnier (West Lafayette, IN), P.G. Righetti (Milan), P. Schoenmakers (Amsterdam), R. Schwarzenbach (Dübendorf), R.E. Shoup (West Lafayette, IN), R.P. Singhal (Wichita, KS), A.M. Siouffi (Marseille), D.J. Strydom (Boston, MA), N. Tanaka (Kyoto), S. Terabe (Hyogo), K.K. Unger (Mainz), R. Verpoorte (Leiden), Gy. Vigh (College Station, TX), J.T. Watson (East Lansing, MI), B.D. Westerlund (Uppsala)

EDITORS, BIBLIOGRAPHY SECTION

Z. Deyl (Prague), J. Janák (Brno), V. Schwarz (Prague)



ELSEVIER

Amsterdam – Lausanne – New York – Oxford – Shannon – Tokyo

J. Chromatogr. A, Vol. 682 (1994)

The *Bibliography Section 1994* will be published in two volumes (681 and 682) of two issues each. Combined indexes to both volumes will appear in the last issue of the year, Vol. 682, No. 2.

© 1994 ELSEVIER SCIENCE B.V. All rights reserved.

0021-9673/94/\$07.00

No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, without the prior written permission of the publisher, Elsevier Science B.V., Copyright and Permissions Department, P.O. Box 521, 1000 AM Amsterdam, Netherlands.

Upon acceptance of an article by the journal, the author(s) will be asked to transfer copyright of the article to the publisher. The transfer will ensure the widest possible dissemination of information.

Special regulations for readers in the USA – This journal has been registered with the Copyright Clearance Center, Inc. Consent is given for copying of articles for personal or internal use, or for the personal use of specific clients. This consent is given on the condition that the copier pays through the Center the per-copy fee stated in the code on the first page of each article for copying beyond that permitted by Sections 107 or 108 of the US Copyright Law. The appropriate fee should be forwarded with a copy of the first page of the article to the Copyright Clearance Center, Inc., 27 Congress Street, Salem, MA 01970, USA. If no code appears in an article, the author has not given broad consent to copy and permission to copy must be obtained directly from the author. The fee indicated on the first page of an article in this issue will apply retroactively to all articles published in the journal, regardless of the year of publication. This consent does not extend to other kinds of copying, such as for general distribution, resale, advertising and promotion purposes, or for creating new collective works. Special written permission must be obtained from the publisher for such copying.

No responsibility is assumed by the Publisher for any injury and/or damage to persons or property as a matter of products liability, negligence or otherwise, or from any use or operation of any methods, products, instructions or ideas contained in the materials herein. Because of rapid advances in the medical sciences, the Publisher recommends that independent verification of diagnoses and drug dosages should be made.

Although all advertising material is expected to conform to ethical (medical) standards, inclusion in this publication does not constitute a guarantee or endorsement of the quality or value of such product or of the claims made of it by its manufacturer.

♾ The paper used in this publication meets the requirements of ANSI/NISO 239.48-1992 (Permanence of Paper).

Printed in the Netherlands

BIBLIOGRAPHY SECTION

SUPPLEMENT TO THE
JOURNAL OF CHROMATOGRAPHY A

1994

EDITORS:

Z. DEYL (Prague)

J. JANÁK (Brno)

V. SCHWARZ (Prague)

Bibliography Section

Liquid Column Chromatography

1. REVIEWS AND BOOKS

- 2780 Blau, K. and Halket, J.M. (Editors): *Handbook of Derivatives for Chromatography* (Second Edition), Wiley, Chichester, 1993, 369 p.; C.A., 120 (1994) 145386m.
- 2781 Brown, P.R. and Grushka, E. (Editors): *Advances in Chromatography*. Vol. 33, Dekker, New York, 1993, 283 p.; C.A., 120 (1994) 87625a.
- 2782 Buyanovskaya, A.G. and Terent'eva, E.A.: (Application of ion chromatography in organic elemental analysis). *Zh. Anal. Khim.*, 48 (1993) 1414-1428; C.A., 120 (1994) 68392f - a review with 124 refs.
- 2783 Ding, L., Zhang, Z. and An, D.: (HPLC-GC multidimensional chromatography and its applications). *Zhongguo Yaoxue Zazhi*, 28 (1993) 460-465; C.A., 120 (1994) 62363a - a review with 26 refs.
- 2784 Garcia, F.A.P. and Pires, E.M.V.: Chromatography. In: Kennedy, J.F. and Cabral, J.M.S. (Editors), *Recovery Processes Biol. Mater.*, Wiley, Chichester, 1993, pp. 415-451; C.A., 120 (1994) 63082v - a review with 8 refs.
- 2785 Haidacher, D. and Horvath, C.: Physicochemical basis of hydrophobic interaction chromatography. *NATO ASI Ser., Ser. C*, 383(Theoretical Advancement in Chromatography and Related Separation Techniques) (1992) 443-480; C.A., 120 (1994) 49095v - a review with 127 refs.
- 2786 Holman, R.B.: Theoretical and practical aspects of high performance liquid chromatography. *IBRO Handb. Ser.*, 15(High Performance Liquid Chromatography in Neuroscience Research) (1993) 3-11; C.A., 120 (1994) 128677q - a review with 15 refs.
- 2787 Holman, R. B., Cross, A.J. and Joseph, M.H. (Editors): *High-Performance Liquid Chromatography in Neuroscience Research* In: *IBRO Handb. Ser.*, Vol. 15, Wiley, Chichester, 1993, 369 pp.; C.A., 120 (1994) 211986k.
- 2788 Joseph, M.H. and Cross, A.J.: Methods development: preparation, separation and detection. *IBRO Handb. Ser.*, 15(High Performance Liquid Chromatography in Neuroscience Research) (1993) 13-23; C.A., 120 (1994) 128678r - a review with 2 refs.
- 2789 Lawlis, V.B. and Heinsohn, H.: Industrial perspective on the role of chromatography in the purification of biological molecules. *LC-GC*, 11 (1993) 720-729; C.A., 120 (1994) 72630z.
- 2790 Uden, P.C.: Element specific chromatographic detection for trace inorganic analysis. *Anal. Proc.*, 30 (1993) 405-408; C.A., 120 (1994) 68007j - a review with 33 refs.

See also 2805, 2807, 2817, 2832, 2836, 2837, 2840, 2842, 2847, 2852, 2858, 2862, 2908, 2909, 2935, 2936, 2939, 2940, 2943, 2945, 2946, 2950, 2955, 2956, 2957, 2958, 2961, 2962, 2973, 2975, 2979, 2990, 2991, 2995, 2998, 2999, 3004, 3012, 3021, 3153, 3181, 3230, 3239, 3242, 3243, 3289, 3303, 3306, 3379, 3384, 3385, 3387, 3393, 3423, 3509, 3521, 3666, 3710, 3740, 3802, 3803, 3842, 3847, 3848, 3849, 3869, 3894, 4043, 4045, 4047, 4054, 4055, 4068, 4069, 4077, 4086, 4097, 4112, 4130.

2. FUNDAMENTALS, THEORY AND GENERAL

2a. General

- 2791 Andersson, R. and Haemaelaenen, M.D.: Simplex focusing of retention times and latent variable projections of chromatographic profiles. *Chemom. Intell. Lab. Syst.*, 22 (1994) 49-61; C.A., 120 (1994) 100917u.
- 2792 Bahowick, T.J., Dunphy, D.R. and Synovec, R.E.: Analysis of unresolved chromatograms by the absorbance ratio and sequential chromatogram ratio techniques coupled with peak suppression. *J. Chromatogr. A*, 663 (1994) 135-150.
- 2793 Barnes, N.: Developments in ion exchange chromatography. *Am. Lab. (Shelton)*, 25 (1993) 52BB, 52DD-FF; C.A., 120 (1994) 264941y.
- 2794 Bussemas, H.H., Harsch, G. and Ettore, L.S.: Friedlieb Ferdinand Lunge (1794-1867): "Self grown pictures" as precursors of paper chromatography. *Chromatographia*, 38 (1994) 243-254.
- 2795 Dolan, J.W.: Extra peaks. *LC-GC Int.*, 7 (1994) 74-76.
- 2796 Dolan, J.W.: Fixing problems on the cheap. *LC-GC Int.*, 7 (1994) 250-252.
- 2797 Dolan, J.W.: The dilemma. *LC-GC Int.*, 7 (1994) 204-208.
- 2798 Dolan, J.W.: Worth repeating. *LC-GC Int.*, 7 (1994) 14-15.
- 2799 Eiteman, M.A.: Temperature-dependent phase inversion and its effect on partitioning in the poly(ethylene glycol)-ammonium sulfate aqueous two-phase system. *J. Chromatogr. A*, 668 (1994) 13-19.
- 2800 Ettore, L.S.: New, unified nomenclature for chromatography. *Chromatographia*, 38 (1994) 521-526.
- 2801 Eudes, V., Arnaud, B. and Genty, C.: (Self-training software to learn general chromatography). *Ann. Falsif. Expert. Chim. Toxicol.*, 86 (1993) 119-132; C.A., 120 (1994) 76578m.
- 2802 Hatrik, Š., Hrouzek, J., Lehotay, J. and Krupčík, J.: Use of a threshold criterion in the computer-assisted optimization of chromatographic separations. *J. Chromatogr. A*, 665 (1994) 9-15.

- 2803 Hrouzek, J., Hatřík, Š., Krupčík, J. and Benická, E.: Evaluation of separation quality with a novel threshold criterion. *J. Chromatogr. A*, 665 (1994) 3-8.
- 2804 Kracke, C.: (MCLC - membrane convective liquid chromatography). *Bioforum*, 16 (1993) 480-481; *C.A.*, 120 (1994) 157797j.
- 2805 Marina, M.L. and García, M.A.: Micellar liquid chromatography with hybrid eluents. *J. Liq. Chromatogr.*, 17 (1994) 957-980 - a review with 99 refs.
- 2806 Meyer, V.R.: Chromatographic integration errors: a closer look at a small peak. *LC-GC Int.*, 7 (1994) 94-100.
- 2807 Morita, I.: (Development of direct injection technique for the analysis of biological fluids). *Kuromatogurafi*, 14 (1993) 238A-241A; *C.A.*, 120 (1994) 186342n - a review with 8 refs.
- 2808 Nosko, S. and Burger, D.: (Recycling of solvent mixtures. Acetonitrile wastes). *LaborPraxis*, 17 (1993) 51-55; *C.A.*, 120 (1994) 152645y.
- 2809 Seaton, P. and Zimmer, J.: An investigative undergraduate laboratory experiment using extraction, chromatography, and a biological activity test. *Biochem. Educ.*, 21 (1993) 216-217; *C.A.*, 120 (1994) 243544f.
- 2810 Van der Wielen, L.A.M., Straathof, A.J.J. and Luyben, K.C.A.M.: Adsorptive and chromatographic bioreactors. In: Weijnen, M.P.C. and Drinkenburg, A.A.H. (Editors), *Precis. Process Technol.*, 1st 1992, Kluwer, Dordrecht, 1993, pp. 353-379; *C.A.*, 120 (1994) 167797y.
- For additional information see *C.A.*:
120 (1994) 94154z, 101280t, 105917d, 145325r, 207631y, 212022e, 263140z.
- See also 2784, 2789, 2826, 2827, 2834, 2896, 2929, 3371, 3373, 3378, 3423, 3842.
- 2b. Thermodynamics and theoretical relationships**
- 2811 Bhalekar, A.A. and Engberts, J.B.F.N.: Theory of gel filtration chromatography. Elution patterns of solutes in the presence of micellar and premicellar aggregates of micelle-forming surfactants. *Colloid Polym. Sci.*, 271 (1993) 1068-1075; *C.A.*, 120 (1994) 201113j.
- 2812 Bourguignon, B., de Aguiar, P.F., Khots, M.S. and Massart, D.L.: Optimization in irregularly shaped regions: pH and solvent strength in reversed-phase high-performance liquid chromatography separations. *Anal. Chem.*, 66 (1994) 893-904.
- 2813 Breuer, M.R.P., Claessens, G.A. and Cramers, C.A.: Statistical evaluation of the validity of a method for characterizing stationary phases for reversed-phase liquid chromatography based on retention of homologous series. *Chromatographia*, 38 (1994) 137-146.
- 2814 Geng, X.D. and Regnier, F.E.: Convergence of retention for small solutes in reversed-phase liquid chromatography. *Chromatographia*, 38 (1994) 158-162.
- 2815 Giona, M., Adrover, A., Barba, D. and Spera, D.: Simplified analysis of chromatographic-column dynamics. *Chem. Eng. Sci.*, 49 (1994) 541-547; *C.A.*, 120 (1994) 100924u.
- 2816 Lebedev, Yu.Ya.: (Criteria for the chromatographic band mobility for a case of inner-diffusing kinetics of interphase mass transfer). *Zh. Fiz. Khim.*, 67 (1993) 1915-1917; *C.A.*, 120 (1994) 201330c.
- 2817 Lee, W.C., Tsai, G.J. and Tsao, G.T.: Analysis of chromatography by plate theory. *Sep. Technol.*, 3 (1993) 178-197; *C.A.*, 120 (1994) 174067p - a review with 116 refs.
- 2818 Martire, D.E.: Unified approach to the theory of chromatography. I. "Incompressible" binary mobile phase (liquid chromatography). *NATO ASI Ser., Ser. C*, 383 (1992) 261-273; *C.A.*, 120 (1994) 63271f.
- 2819 Samo, A.R., Khahawer, M.Y., Chippa, M.A., Arbani, S.A. and Aliqureshi, G.A.: Improvement in the electrostatic theory of reversed-phase ion-pair chromatography to include the role of organic modifier. *J. Chem. Soc. Pak.*, 15 (1993) 196-199; *C.A.*, 120 (1994) 201352e.
- 2820 Ståhlberg, J.: Electrostatic retention model for ion-exchange chromatography. *Anal. Chem.*, 66 (1994) 440-449.
- 2821 Wang, Q.-S., Gao, R.-Y., Yan, B.-W. and Fan, D.P.: Statistical scanning method for the optimization of gradient elution high performance liquid chromatography. *Chromatographia*, 38 (1994) 187-190.
- 2822 Yang, G. and Tao, Z.: Unified theoretical moment expressions for elution chromatography and frontal chromatography. *Chin. J. React. Polym.*, 1 (1992) 109-114; *C.A.*, 120 (1994) 174341y.
- For additional information see *C.A.*:
120 (1994) 68446b, 157788g, 201382w.
- See also 3000, 3709, 4081, 4090, 4099.
- 2c. Relationship between structure and chromatographic behaviour**
- 2823 Sun, X., Xin, M. and Zhao, J.: Method for determination of partition coefficients by high-performance liquid chromatography: application to *o*-hydroxybenzenesulfonanilides. *J. Liq. Chromatogr.*, 17 (1994) 1183-1194.
- 2824 Sun, Z.L., Liu, M.C. and Hu, Z.D.: Relationship between retention behavior and molecular structure in HPLC. Correlation between solubility parameter and molecular properties. *Chromatographia*, 38 (1994) 599-608.
- See also 2805, 2992, 3198, 3415, 3441, 3873.
- 2d. Measurement of physico-chemical and related values**
- 2825 Dayal, U. and Mehta, S.K.: Studies on molecular weight determination using refractive index and multi-angle laser light scattering detectors in size exclusion chromatography. *J. Liq. Chromatogr.*, 17 (1994) 303-316.
- 2826 Eiteman, M.A.: Predicting partition coefficients of multi-charged solutes in aqueous two-phase systems. *J. Chromatogr. A*, 668 (1994) 21-30.
- 2827 Guan, Y., Treffry, T.E. and Lilley, T.H.: Application of a statistical geometrical theory to aqueous two-phase systems. *J. Chromatogr. A*, 668 (1994) 31-45.
- 2828 McGuffin, V.L., Evans, C.E. and Chen, S.H.: Direct examination of separation processes in liquid chromatography: effect of temperature and pressure on solute retention. *J. Microcolumn Sep.*, 5 (1993) 3-10; *C.A.*, 120 (1994) 182097p.
- 2829 Tan, L.C., Carr, P.W., Fréchet, J.M.J. and Smigol, V.: Liquid chromatographic study of solute hydrogen bond basicity. *Anal. Chem.*, 66 (1994) 450-457.



2830 Wingren, C., Persson, B. and Hansson, U.-B.: Supports for liquid-liquid partition chromatography in aqueous two-phase systems: a comparison of LiChrospher and LiParGel. *J. Chromatogr. A*, 668 (1994) 65-73.

For additional information see C.A.:

120 (1994) 108223k, 108413x.

See also 2799, 2845, 2846, 3018, 3094, 3144, 3371, 3372, 3378, 3423, 3459, 3481, 3830, 3837, 3838, 3922, 4104.

3. GENERAL TECHNIQUES

3a. Apparatus and accessories

2831 Alebic-Kolbah, T. and Wainer, I.W.: Bioactive LC reactors for stereospecific synthesis with online enantioselective separations. *Am. Lab. (Shelton)*, 25 (1993) 52R, 52V-X, 52Z-AA; *C.A.*, 120 (1994) 186386e.

2832 Arkhipov, D.B. and Belenkii, B.G.: (Development of liquid chromatography instrumentation. Trends in 1958-1991). *Zh. Anal. Khim.*, 48 (1993) 1894-1905; *C.A.*, 120 (1994) 181695v - a review with 13 refs.

2833 Dolan, J.W.: Choosing the right instrument. *LC-GC Int.*, 7 (1994) 136-138.

2834 Jordan, J.R.: ISO 9000 and analytical instruments. *LC-GC*, 11 (1993) 546-549; *C.A.*, 120 (1994) 152356e.

2835 McDowall, R.D.: The right tool for the job? *LC-GC Int.*, 7 (1994) 146-148.

2836 Volkov, S.A., Slin'ko, M.G. and Sakodynskii, K.I.: (Prospects for the development of chromatographic reactors). *Khim. Prom-st. (Moscow)*, (1993) 308-311; *C.A.*, 120 (1994) 80008t - a review with 28 refs.

For additional information see C.A.:

120 (1994) 94319g, 123799w, 239651w.

See also 2883, 2907, 2924, 3022, 3797, 4107.

3b. Detectors and detection reagents

2837 Barisci, J.N. and Wallace, G.G.: Advances in electrochemical detection for HPLC. *Chem. Aust.*, 60 (1993) 538-546; *C.A.*, 120 (1994) 123802s - a review with 28 refs.

2838 Cabalin, L.M., Ruperez, A. and Laserna, J.J.: Surface-enhanced Raman spectrometry for detection in liquid chromatography using a windowless flow cell. *Talanta*, 40 (1993) 1741-1747; *C.A.*, 120 (1994) 71012n.

2839 Conte, E.D. and Barry, E.F.: The evaluation of aerosol alkali flame ionization detector for high performance liquid chromatography. *Microchem. J.*, 48 (1993) 365-371; *C.A.*, 120 (1994) 94360p.

2840 Faegerstam, L.G. and O'Shannessy, D.J.: Surface plasmon resonance detection in affinity technologies. *Chromatogr. Sci. Ser.*, 63(Hanbook of Affinity Chromatography) (1993) 229-252; *C.A.*, 120 (1994) 186327m. - a review with 18 refs.

2841 Gale, R.J. and Li, H.: An electrodeless, high frequency liquid chromatography detector. *Proc. - Electrochem. Soc.*, 93-97 (1993) 227-235; *C.A.*, 120 (1994) 68422r.

2842 Giuffre, R.W.: The use of diode array detectors in the pharmaceutical industry. *Chromatogr. Sci. Ser.*, 62 (1993) 163-197; *C.A.*, 120 (1994) 62365c - a review with 36 refs.

2843 Jimena Garcia, J.A., Gimenez Plaza, J. and Cano Pavon, J.M.: Use of derivative spectrophotometry in the resolution of overlapped peaks in liquid chromatography and its application in the analysis of active components in insecticide formulations. *J. Liq. Chromatogr.*, 17 (1994) 277-285.

2844 Kita, J.I.: Optical detectors for capillary chromatography. *Patentschrift (Switz.)* CH 682,846 (Cl. G01N30/74), 30 Nov. 1993, JP Appl. 90/416,041, 28 Dec. 1990; 10 p.; *C.A.*, 120 (1994) 207716e.

2845 Lesec, J.: Flow fluctuations in GPC-viscometry. *J. Liq. Chromatogr.*, 17 (1994) 1011-1028.

2846 Lesec, J., Millequant, M. and Havard, T.: High temperature GPC with a single capillary viscometer. *J. Liq. Chromatogr.*, 17 (1994) 1029-1055.

2847 Mahuzier, G., Prognon, P., Sargi, L., Kouwatli, H., Tod, M. and Farinotti, R.: (Luminescence and detection in liquid chromatography: part III. Chemiluminescence). *Ann. Pharm. Fr.*, 52 (1994) 25-42; *C.A.*, 120 (1994) 211673f - a review with 54 refs.

2848 Miyazawa, T., Lertsiri, S., Fujimoto, K. and Oka, M.: Luminol chemiluminescent determination of hydrogen peroxide at picomole levels using high-performance liquid chromatography with a cation-exchange resin gel column. *J. Chromatogr. A*, 667 (1994) 99-104.

2849 Niederländer, H.A.G., Gooijer, C. and Velthorst, N.H.: Chemiluminescence detection in liquid chromatography based on photo-oxygenation involving reactive oxygen intermediates. *Anal. Chim. Acta*, 285 (1994) 143-159.

2850 Rosenzweig, Z. and Yeung, E.S.: Laser-based double-beam circular dichroism detector of liquid chromatography. *Appl. Spectrosc.*, 47 (1993) 2017-2021; *C.A.*, 120 (1994) 68425u.

2851 Sato, H. and Dasgupta, P.K.: Analyte identification in ion chromatography. Electromigration governed chronoamperometric profiles. *Anal. Chim. Acta*, 284 (1993) 27-36.

2852 Saz, J.M. and Díez-Masa, J.C.: Thermo-optical spectroscopy: new and sensitive schemes for detection in capillary separation techniques. *J. Liq. Chromatogr.*, 17 (1994) 499-520 - a review with 42 refs.

2853 Sun, X.-Y. and Aue, W.A.: Constancy of spectral response ratios in the flame photometric detector. *J. Chromatogr. A*, 667 (1994) 191-203.

2854 Takeuchi, T. and Miwa, T.: Signal enhancement by on-column fluorimetric detection in microcolumn liquid chromatography with cyclodextrin-bonded stationary phase. *Chromatographia*, 38 (1994) 555-558.

2855 Wyatt, P.J. and Papazian, L.A.: The interdetector volume in modern light scattering and high performance size-exclusion chromatography. *LC-GC Int.*, 7 (1994) 266-274.

For additional information see C.A.:

120 (1994) 68299f, 186529d, 186809y, 207634b.

See also 2825, 2832, 2938, 2994, 2997, 3009, 3011, 3025, 3042, 3060, 3154, 3155, 3156, 3168, 3173, 3182, 3238, 3266, 3297, 3317, 3511, 3651, 3655, 3657, 3766, 3809, 3838, 3887, 4100, 4125.

3c. Sorbents and columns, packing procedures

- 2856 Albert, K., Brindle, R., Martin, P. and Wilson, I.D.: Characterisation of C₁₈-bonded silicas for solid-phase extraction by solid-state NMR spectroscopy. *J. Chromatogr. A*, 665 (1994) 253-258.
- 2857 Albert, K., Brindle, R., Schmid, J., Buszewski, B. and Bayer, E.: cP/MAS NMR investigations of silica gel surfaces modified with aminopropylsilane. *Chromatographia*, 38 (1994) 283-290.
- 2858 Allenmark, S.G. and Andersson, S.: Proteins and peptides as chiral selectors in liquid chromatography. *J. Chromatogr. A*, 666 (1994) 167-179 - a review with 48 refs.
- 2859 Anazawa, T.A. and Jardim, I.C.S.F.: Polymethyloctylsiloxane adsorbed on porous silica as a packing material for reversed phase high performance liquid chromatography. *J. Liq. Chromatogr.*, 17 (1994) 1265-1279.
- 2860 Bargmann-Leyder, N., Truffert, J.-C., Tambuté, A. and Caude, M.: Evaluation of Pirkle-type chiral stationary phases by liquid and supercritical fluid chromatography. Influence of the spacer length and the steric hindrance in the vicinity of the stereogenic centre. *J. Chromatogr. A*, 666 (1994) 27-40.
- 2861 Bilz, J., Schmidt, S., Froelich, P. and Renger, P.: The use of stationary phases based on poly(vinyl alcohol) in anion chromatography. *Ger. (East)* DD 301,629 (Cl. B01J41/06), 06 May 1993, Appl. 328,447, 10 May 1989; 4 p.; C.A., 120 (1994) 201428r.
- 2862 Boenke, A.: (Reference materials in HPLC). *GIT Fachz. Lab.*, 38 (1994) 8-10; C.A., 120 (1994) 182063z - a review with 10 refs.
- 2863 Boschetti, E., Girot, P., Guerrier, L. and Andre, B.: (Chromatographic sorbents for preparative protein separations). *Ann. Pharm. Fr.*, 51 (1993) 299-306; C.A., 120 (1994) 49244t.
- 2864 Chappell, I.: The effect of silane type and endcapping on the selectivity of cyanopropyl bonded silicas used in normal-phase mode. *LC-GC Int.*, 7 (1994) 282-284.
- 2865 El'tikova, N.A.: (Chromatographic properties of carbon sorbents). *Zh. Fiz. Khim.*, 67 (1993) 1862-1865; C.A., 120 (1994) 201329j.
- 2866 Firouztale, E., Scott, A.P., Dalvie, S.K. and von Blohn, G.M.: Experimental and theoretical study of key parameters of adsorption on reverse phase macroporous resins. *AIChE Symp. Ser.*, 290 (New Developments in Bioseparation) (1992) 25-33; C.A., 120 (1994) 29243s.
- 2867 Fonseca, L.P., Caldeira, J.C.G. and Cabral, J.M.S.: Improvement in the polyethylene glycol-Cibacron Blue purification method. *J. Chromatogr. A*, 668 (1994) 61-64.
- 2868 Ge, H., Gilmore, K.G. and Wallace, G.G.: Studies on poly (3-octadecyl pyrrole) modified silica as a reversed phase HPLC packing material. *J. Liq. Chromatogr.*, 17 (1994) 1301-1316.
- 2869 Ghijs, M. and Sandra, P.: Polymeric coating in packed fused silica micro-LC columns. *J. Microcolumn Sep.*, 3 (1991) 443-448; C.A., 120 (1994) 94378a.
- 2870 Gübitz, G., Mihellyes, S., Kobinger, G. and Wutte, A.: New chemically bonded chiral ligand-exchange chromatographic stationary phases. *J. Chromatogr. A*, 666 (1994) 91-97.
- 2871 Helburn, R.S., Rutan, S.C., Pompano, J., Mitchem, D. and Patterson, W.T.: Solvatochromic studies of solvated chromatographic stationary phases. *Anal. Chem.*, 66 (1994) 610-618.
- 2872 Hosoya, K., Kishii, Y., Kimata, K. Araki, T. and Tanaka, N.: Efficiency of porous hydrophilic polymer-based packing materials in chromatographic analysis of drugs existing with polypeptide. *Chromatographia*, 38 (1994) 177-182.
- 2873 Hosoya, K., Yoshizako, K., Tanaka, N., Kimata, K., Araki, T. and Fréchet, J.M.J.: One-pot preparation method for a uniform-sized polymer-based chiral stationary phase for high-performance liquid chromatography with polymethacrylamide as a chiral selector. *J. Chromatogr. A*, 666 (1994) 449-455.
- 2874 Houbenová, A., Claessens, H.A., de Haan, J.W., Cramers, C.A. and Stulik, K.: An investigation concerning the stability of a sterically protected cyanopropyl modified silica substrate. *J. Liq. Chromatogr.*, 17 (1994) 49-68.
- 2875 Inagaki, M., Kitagawa, S. and Tsuda, T.: New packing method for liquid chromatographic column by the aid of electrostatic field. *Kuromatogurafi*, 14 (1993) 55R-60R; C.A., 120 (1994) 63273h.
- 2876 Ivakhnyuk, G.K., Babkin, O.E. and Fedorov, N.F.: (Technological bases of direct synthesis of standard adsorbents for chromatography). *Zh. Fiz. Khim.*, 67 (1993) 2052-2055; C.A., 120 (1994) 137898q.
- 2877 Jinno, M. and Uemura, T.: Copper-phthalocyanine stationary phase in microcolumn liquid chromatography. *J. Microcolumn Sep.*, 4 (1993) 325-329; C.A., 120 (1994) 54121e.
- 2878 Kitagawa, S., Inagai, M. and Tsuda, T.: (Tiny porous ceramic bed supports for fused-silica packed capillary columns used in electrochromatography and liquid chromatography). *Kuromatogurafi*, 14 (1993) 39R-43R; C.A., 120 (1994) 114503t.
- 2879 Kříž, J., Adamcová, E., Knox, J.H. and Hora, J.: Characterization of adsorbents by high-performance liquid chromatography using aromatic hydrocarbons. Porous graphite and its comparison with silica gel, alumina, octadecylsilica and phenylsilica. *J. Chromatogr. A*, 663 (1994) 151-161.
- 2880 Kurganov, A., Puchkova, Yu., Davankov, V. and Eisenbeiss, F.: Polyvinylpyrrolidone-coated silica packings for chromatography of proteins and peptides. *J. Chromatogr. A*, 663 (1994) 163-174.
- 2881 Li, Q., Zhu, J., Daugulis, A.J., Hsu, C.C. and Goosen, M.F.A.: Polysaccharide microcapsules and macroporous beads for enhanced chromatographic separation. *Biomater., Artif. Cells, Immobilization Biotechnol.*, 21 (1993) 391-398; C.A., 120 (1994) 128758s.
- 2882 Majors, R.E.: Measuring particle-size distribution of HPLC packings. *LC-GC Int.*, 7 (1994) 8-13.
- 2883 Majors, R.E.: New chromatography columns and accessories at the 1994 Pittsburgh Conference, Part I. *LC-GC Int.*, 7 (1994) 190-203.
- 2884 Miller, K.G. and Poole, C.F.: Methodological approach for evaluating operational parameters and the characterization of a popular sorbent for solid-phase extraction by high pressure liquid chromatography. *J. High Resolut. Chromatogr.*, 17 (1994) 125-134.
- 2885 Montgomery, M.E., Jr. and Wirth, M.J.: Spectroscopic study of the molecular basis of wetting of a C₁₈ surface by long-chain *n*-alcohols. *Anal. Chem.*, 66 (1994) 680-684.
- 2886 Novák, I. and Berek, D.: Structural in homogeneities in wide-pore silica gels. *J. Chromatogr. A*, 665 (1994) 33-36.

- 2887 Ooga, K.: Preparation of chitosan derivatives as resolving agents for racemic compounds by chromatography. *Jpn. Kokai Tokkyo Koho* JP 05,163,164 [93,163,164] (Cl. CO7B57/00), 29 Jun. 1993, Appl. 91/333,280, 17 Dec. 1991; 8 pp.; C.A., 120 (1994) 107629k.
- 2888 Park, J.H., Chae, J.J., Nah, T.H. and Jang, M.D.: Characterization of some silica-based reversed-phase liquid chromatographic columns based on linear solvation energy relationships. *J. Chromatogr. A*, 664 (1994) 149-158.
- 2889 Petro, M., Gemeiner, P. and Berek, D.: Dextran-grafted silica gel for high-performance size-exclusion chromatography of proteins. *J. Chromatogr. A*, 665 (1994) 37-45.
- 2890 Podszun, W., Boemer, B., Grosser, R., Lange, W. and Grosse-Bley, M.: Filler-containing optically active polymeric beds, and their manufacture and use. *Eur. Pat. Appl. EP* 545,168 (Cl. C08F20/54), 09 Jun. 1993, DE Appl. 4,139,747, 03 Dec. 1991; 9 p.; C.A., 120 (1994) 136485r.
- 2891 Reif, O.-W., Nier, V., Bahr, U. and Freitag, R.: Immobilized metal affinity membrane adsorbents as stationary phases for metal interaction protein separation. *J. Chromatogr. A*, 664 (1994) 13-25.
- 2892 Saito, Y., Jinno, K., Pesek, J.J., Chen, Y.-L., Luehr, G., Archer, J., Fetzer, J.C. and Biggs, W.R.: Molecular shape recognition capability of liquid-crystal bonded phases in reversed-phase high performance liquid chromatography. *Chromatographia*, 38 (1994) 295-303.
- 2893 Samsonov, G.Y., Pisarev, O.A. and Melenevsky, A.T.: Chromatographic purification and super-purification of biologically active compounds using heterorecticular and composite ion exchange resins at low pressure. *Pure Appl. Chem.*, 65 (1993) 2287-2290; C.A., 120 (1994) 49214h.
- 2894 Šerys, A., Liesiene, J., Urbonavičiene, J., Maruška, A., Radzevičius, K., Valatkevičiute, D. and Bumelis, V.: Preparation and characterization of cellulose based adsorbents for large scale hydrophobic interaction chromatography. *J. Liq. Chromatogr.*, 17 (1994) 749-760.
- 2895 Škrabáková, S., Matisová, E., Benická, E., Novák, I. and Berek, D.: Use of a novel carbon sorbent for the adsorption of organic compounds from water. *J. Chromatogr. A*, 665 (1994) 27-32.
- 2896 Slais, K. and Horka, M.: Effect of surface modification and mobile phase velocity on the performance of parallel current open tubular liquid chromatography. *J. Microcolumn Sep.*, 5 (1993) 63-69; C.A., 120 (1994) 94404f.
- 2897 Smigol, V., Svec, F. and Fréchet, J.M.J.: Novel uniformly sized polymeric stationary phase with hydrophilized large pores for direct injection HPLC determination of drugs in biological fluids. *J. Liq. Chromatogr.*, 17 (1994) 891-911.
- 2898 Smigol, V., Svec, F. and Fréchet, J.M.J.: Uniformly sized poly(vinylphenol) porous beads: a versatile HPLC separation medium allowing fast switching between size-exclusion, normal phase, and reversed phase chromatography. *J. Liq. Chromatogr.*, 17 (1994) 259-276.
- 2899 Tani, K. and Suzuki, Y.: Syntheses of spherical silica and titania from alkoxides on a laboratory scale. *Chromatographia*, 38 (1994) 291-294.
- 2900 Thienpont, A. and Félix, G.: Preparation of chemically bonded stationary phases for HPLC by *ann in situ* process. *Chromatographia*, 37 (1993) 505-510.
- 2901 Tsvetkov, F., Heller-Kallai, L. and Mingelgrin, U.: Potassium halide-treated montmorillonite (KTM) as a solid phase in liquid chromatography. *Clays Clay Miner.*, 41 (1993) 527-536; C.A., 120 (1994) 145314m.
- 2902 Uray, G., Maier, N.M. and Lindner, W.: Diphenylethanedi-amine derivatives as chiral selectors. III. Comparison of four new diastereomeric chiral stationary phases prepared by addition of mono-3,5-dinitrobenzoyldiphenylethanedi-amine derivatives to optically pure epoxy silica. *J. Chromatogr. A*, 666 (1994) 41-53.
- 2903 Wu, M. and Deng, W.: Adsorbents for purification of active ingredients from Chinese traditional herb. *Faming Zguanli Shenqing Gongkai Shuomingshu* CN 1,072,089 (Cl. A61K35/78), 19 May 1993, Appl. 92,111,985, 14 Nov. 1992; 6 p.; C.A., 120 (1994) 62260q.
- 2904 Yamashita, H. and Maekawa, T.: Synthesis of microporous silica-gel particles in a W/O emulsion an application to the direct optical resolution of amino acid enantiomers. *Anal. Sci.*, 9 (1993) 545-548; C.A., 120 (1994) 94376y.
- 2905 Yang, S., Kruk, L.F.R. and Khaledi, M.G.: Fluorinated bonded stationary phases in micellar liquid chromatography. *J. Chromatogr. A*, 664 (1994) 1-11.
- For additional information see C.A.:
120 (1994) 49540m, 54888k, 57042r, 63272g, 68317k, 68535e, 68541d, 86531e, 87529x, 94395d, 100832n, 109085d, 135234w, 145390h, 152567z, 157528x, 157785d, 166330d, 166331e, 186764b, 186765c, 186766d, 186767e, 212021d, 215314n, 239649b, 239653y, 239685k, 265315j.
- See also 2793, 2813, 2941, 2951, 2953, 2956, 2960, 2980, 2984, 2987, 3004, 3120, 3326, 3376, 3386, 3597, 3646, 3758, 3845, 4093.
- 3d. *Quantitative analysis*
- 2906 Meyer, V.R.: (Accuracy in determination of peak areas). *GIT Fachz. Lab.*, 38 (1994) 4-5; C.A., 120 (1994) 182072b.
- For additional information see C.A.:
120 (1994) 68239m, 152623q, 165428e.
- See also 4106.
- 3e. *Preparative scale chromatography*
- 2907 Eleveld, J.T., Claessens, H.A., German, A.L. and Cramers, C.A.: Evaluation of the performance of the annular expansion system for preparative liquid chromatography. *LC-GC Int.*, 7 (1994) 219-226.
- 2908 Samsonov, G.V.: (Preparative chromatography as a system of bioseparation methods). *Khim. Prom-st. (Moscow)*, No. 7 (1993) 292-295; C.A., 120 (1994) 100838u - a review with 22 refs.
- 2909 Velayudhan, A., Ladisch, M.R. and Porter, J.E.: Modeling of nonlinear elution chromatography for preparative-scale separations. *AIChE Symp. Ser.*, 290 (New Developments in Bioseparation) (1992) 1-11; C.A., 120 (1994) 49122b - a review with 127 refs.

- 2910 Weisz, A., Scher, A.L., Shinomiya, K., Fales, H.M. and Ito, Y.: A new preparative-scale purification technique: pH-zone-refining countercurrent chromatography. *J. Am. Chem. Soc.*, 116 (1994) 704-708; *C.A.*, 120 (1994) 94394c.
- See also 2799, 2826, 2827, 2830, 2867, 2894, 2933, 2976, 2979, 2983, 3002, 3077, 3082, 3083, 3198, 3212, 3267, 3322, 3371, 3372, 3375, 3378, 3395, 3396, 3399, 3404, 3405, 3441, 3459, 3481, 3530, 3534, 3625, 4027, 4071, 4072, 4073, 4075, 4078, 4107.
- 3f. *Programmed temperature, pressure, vapors, gradients*
- 2911 Deinhammer, R.S., Ting, E.-Y. and Porter, M.D.: Electrochemically modulated liquid chromatography (EMLC): a new approach to gradient elution separation. *J. Electroanal. Chem.*, 362 (1993) 295-299; *C.A.*, 120 (1994) 152657d.
- 2912 Kaminski, N., Kandybowicz, B. and Szukalski, J.: Determination of the optimum volume of a mixer or the maximum proportioning time in the gradient liquid chromatography with proportioning valves. *Chem. Anal. (Warsaw)*, 38 (1993) 237-254; *C.A.*, 120 (1994) 63259h.
- For additional information see *C.A.*:
120 (1994) 207636d.
- See also 2828, 3717, 4106.
4. SPECIAL TECHNIQUES
- 4a. *Automation*
- 2913 Jönsson, J.Å., Mathiasson, L., Lindegård, B., Trocewicz, J. and Olsson, A.-M.: Automated system for the trace analysis of organic compounds with supported liquid membranes for sample enrichment. *J. Chromatogr. A*, 665 (1994) 259-268.
- For additional information see *C.A.*:
120 (1994) 182096n.
- See also 2938.
- 4b. *Computerization and modelling*
- 2914 Andrews, M.: Adapting to a chemist's environment: the future trend in chromatography data management. *Int. Lab.*, 24, No. 2 (1994) 11-16.
- 2915 Funasaki, N., Hada, S. and Neya, S.: Frontal chromatography for nonassociable and micellar systems. *Bull. Chem. Soc. Jpn.*, 67 (1994) 65-73; *C.A.*, 120 (1994) 201356r.
- 2916 Gillespie, H.: The AIA standard for data interchange. *LC-GC Int.*, 7 (1994) 20-28.
- 2917 Hanai, T., Hatano, H., Nimura, N. and Kinoshita, T.: Computational chemical analysis of the retention of saccharides on amino phase. *J. Liq. Chromatogr.*, 17 (1994) 241-248.
- 2918 Head, M. and Yellin, B.: Review of the benefits of a chromatography data system on Windows® NT™. *Int. Lab.*, 24, No. 4 (1994) P10-P19.
- 2919 Kallrath, J., Schreick, A., Brockmueller, B., Agar, D.W., Watzemberger, O., Falle, S.A.E.G. and Giddings, J.R.: Simulation of chromatographic reactors. *Comput. Chem. Eng.*, 18 (Suppl.) (1993) S331-S335; *C.A.*, 120 (1994) 57581r.
- 2920 Kilz, P.: (Gel permeation chromatographic (GPC) analysis under windows). *LaborPraxis*, 17 (1993) 64-68; *C.A.*, 120 (1994) 94353p.
- 2921 Klätte, S.J., Zhang, Z. and Beck, T.L.: Computer simulation of chain molecule-inorganic interphases: chromatographic stationary phases and rigid rod self-assembled monolayers. *Mater. Res. Soc. Symp. Proc.*, 304 (1993) 141-146; *C.A.*, 120 (1994) 174260w.
- 2922 Konstantinov, V.A., Rodin, V.N., Volzhinskii, A.I. and Smirnov, N.N.: (Motion of concentration peaks in channels of a membrane-chromatographic element). *Zh. Prikl. Khim.*, 66 (1993) 1553-1557; *C.A.*, 120 (1994) 167502y.
- 2923 Koppenhoefer, B., Nothdurft, A., Pierrot-Sanders, J., Piras, P., Popescu, C., Roussel, C., Stiebler, M. and Trettin, U.: CHIRBASE, a graphical molecular database on the separation of enantiomers by liquid-, supercritical fluid-, and gas chromatography. *Chirality*, 5 (1993) 213-219; *C.A.*, 120 (1994) 162897w.
- 2924 Lago, P., Rovati, L., Cuffaro, D. and Corti, M.: A multiple-input digital correlator for the weak light scattering signals of chromatographic detectors. *Rev. Sci. Instrum.*, 65 (1994) 253-264; *C.A.*, 120 (1994) 123619n.
- 2925 McDowall, R.D.: Evaluating a chromatography data system. *LC-GC Int.*, 7 (1994) 258-265.
- 2926 Molnar, I.: (Computer-aided optimization in chromatography). *LaborPraxis*, 17 (1993) 40-45; *C.A.*, 120 (1994) 152624r.
- 2927 Ouchi, G.I.: How to choose a chromatography data system. *LC-GC Int.*, 7 (1994) 210-214.
- 2928 Ouchi, G.I.: Managing laboratory information. *LC-GC Int.*, 7 (1994) 254-256.
- 2929 Rowe, R.C., Mulley, V.J., Hughes, J.C., Nabney, I.T. and Debenham, R.M.: Neural networks for chromatographic peak classification - a preliminary study. *LC-GC Int.*, 7 (1994) 36-42.
- 2930 Smith, K.: Developments in the field of chromatography data handling. *Am. Lab. (Shelton)*, 25 (1993) 34-36; *C.A.*, 120 (1994) 181785z.
- 2931 Solomon, B.P.: ChromGraph: The chromatography data system with "Windows" into the 21st century. *Curr. Sep.*, 12 (1993) 149-151; *C.A.*, 120 (1994) 186718q.
- 2932 Strauss, R.: (Chromatography-information-management-systems (CIMS)-alternative or consequence). *LaborPraxis*, 17 (1993) 46-48; *C.A.*, 120 (1994) 152625s.
- 2933 Wang, Q., Gao, R. and Yan, B.: Computer-assisted optimization for the selection of mobile phase in semi-preparative HPLC. *Chin. J. Chem.*, 11 (1993) 71-75; *C.A.*, 120 (1994) 109874k.
- 2934 Yanagi, K.: Distortion free digital filter for chromatography. In: Doyama, M. (Editor), *Comput. Aided Innovation New Mater. 2, Proc. Int. Conf. Exhib. Comput. Appl. Mater. Mol. Sci. Eng.*, 2nd 1992, North-Holland, Amsterdam, 1993, pp. 939-940; *C.A.*, 120 (1994) 123592y.
- For additional information see *C.A.*:
120 (1994) 207673p.
- See also 2791, 2801, 2802, 2806, 2821, 2855, 2912, 2942, 2988, 2990, 4055.

- 4c. *Combination with other physico-chemical techniques (MS, IR etc.)*
- 2935 Bruins, A.P.: Atmospheric-pressure-ionization mass spectrometry. II. Applications in pharmacy, biochemistry and general chemistry. *TrAC*, 13 (1994) 81-90 - a review with 39 refs.
- 2936 Dong, Q.: (The hyphenated techniques of chromatography and FTIR spectroscopy). *Guangpuxue Yu Guangpu Fenxi*, 13 (1993) 25-32; *C.A.*, 120 (1994) 152616q - a review with 33 refs.
- 2937 Games, D.E., Rontree, J.A. and Fowils, I.A.: The direct combination of HPLC solute focusing with supercritical fluid chromatography and mass spectrometry to enhance sensitivity and improve identification of trace components in polar solvent extracts. *J. High Resolut. Chromatogr.*, 17 (1994) 68-71.
- 2938 Gilliard, J.A., Cumps, J.L. and Tilquin, B.L.: Efficiency of an automatic peak purity control procedure in high-performance liquid chromatography-photodiode array coupling based on evolving factor analysis. *Chemom. Intell. Lab. Syst.*, 21 (1993) 235-242; *C.A.*, 120 (1994) 116902w.
- 2939 Hiraoka, K.: (Introduction of the LC/MS interfaces: their fundamental aspects). *J. Mass Spectrom. Soc. Jpn.*, 41 (1993) 101-120; *C.A.*, 120 (1994) 207615w - a review with 21 refs.
- 2940 Hofmann, M., Sproul, M., Streck, R., Wilson, I.D. and Rapp, A.: (Liquid chromatography-NMR coupling and its applications). *LaborPraxis*, 17 (1993) 36-41; *C.A.*, 120 (1994) 123806w - a review with refs.
- 2941 Karlsson, K.E.: Fused silica microcolumn LC/MS and MS/MS using an aeronebulized interface. *J. Microcolumn Sep.*, 3 (1991) 449-458; *C.A.*, 120 (1994) 94355r.
- 2942 Koelbl, G., Kalcher, K. and Irgolic, K.J.: Computerized data treatment for an HPLC-GFAAS system for the identification and quantitation of trace element compounds. *J. Autom. Chem.*, 15 (1993) 37-45; *C.A.*, 120 (1994) 94302w.
- 2943 Long, G.L., Ducatte, G.R. and Lancaster, E.D.: Helium microwave-induced plasmas for element specific detection in chromatography. *Spectrochim. Acta, Part B*, 49B (1994) 75-87; *C.A.*, 120 (1994) 207627b - a review with 48 refs.
- 2944 Nabiev, I. and Manfait, M.: Industrial applications of the surface-enhanced Raman Spectroscopy. *Rev. Inst. Fr. Pet.*, 48 (1993) 261-285; *C.A.*, 120 (1994) 182059c.
- 2945 Sakairi, M.: (Atmospheric pressure ionization method as an interface for liquid chromatograph/mass spectrometer). *Bunseki*, (1993) 952-956; *C.A.*, 120 (1994) 207610r - a review with 10 refs.
- 2946 Schwedt, G. and Meyers, A.: (Coupling techniques in chromatography). *LaborPraxis*, 17 (1993) 12-16; *C.A.*, 120 (1994) 94341h - a review with 16 refs.
- 2947 Spraul, M., Hofmann, M., Lindon, J.C., Nicholson, J.K. and Wilson, I.D.: Liquid chromatography coupled with high-field proton nuclear magnetic resonance spectroscopy: Current status and future prospects. *Anal. Proc.*, 30 (1993) 390-392; *C.A.*, 120 (1994) 123951q.
- For additional information see *C.A.*:
120 (1994) 49489b.
- See also 2839, 2841, 2850, 3070, 3072, 3073, 3097, 3150, 3222, 3262, 3280, 3302, 3308, 3325, 3346, 3380, 3383, 3401, 3656, 3659, 3675, 3780, 3799, 3811, 3832, 3879, 3895, 3915, 3933, 3936, 3939, 4011, 4022, 4060, 4062, 4067, 4085, 4095, 4105.
- 4d. *Affinity chromatography (advances)*
- 2948 Beitle, R.R. and Ataai, M.M.: Immobilized metal affinity chromatography and related techniques. *AIChE Symp. Ser.*, 290(New Developments in Bioseparation) (1992) 34-44; *C.A.*, 120 (1994) 49123c.
- 2949 Carlsson, F., Axelsson, A. and Zacchi, G.: Mathematical modeling and parametric studies of affinity chromatography. *Comput. Chem. Eng.*, 18(Suppl.) (1993) S657-S661; *C.A.*, 120 (1994) 52698z.
- 2950 Chaiken, I.: Affinity chromatography in biology and biotechnology. Probing macromolecular interactions using immobilized ligands. *Chromatogr. Sci. Ser.*, 63(Handbook of Affinity Chromatography) (1993) 219-227; *C.A.*, 120 (1994) 186326k - a review with 18 refs.
- 2951 Chang, R.M. and Lee, W.C.: An affinity adsorbent derived from aminopropyl silica for serine protease chromatography. *J. Chem. Technol. Biotechnol.*, 59 (1994) 133-139; *C.A.*, 120 (1994) 238827w.
- 2952 Flurer, C.L. and Novotny, M.: Immobilized metal affinity chromatography with fused silica microcolumns. *J. Microcolumn Sep.*, 4 (1993) 497-501; *C.A.*, 120 (1994) 157786e.
- 2953 Johnson, P.R., Stern, N.J., Eitzman, P.D., Rasmussen, J.K., Milbrath, D.S., Gleason, R.M. and Hogancamp, R.E.: Reproducibility of physical characteristics, protein immobilization and chromatographic performance of 3M Emphaze Biosupport Medium AB 1. *J. Chromatogr. A*, 667 (1994) 1-9.
- 2954 Kang, K.A., Ryu, D.D.Y. and Shi, Y.: Simulation of heterogeneous interaction between ligand and product biomolecules in affinity chromatography. *AIChE Symp. Ser.*, 290(New Developments in Bioseparation) (1992) 12-24; *C.A.*, 120 (1994) 29242r.
- 2955 Kauvar, L.M.: Investigating specificity via affinity chromatography. *Chromatogr. Sci. Ser.*, 63(Handbook of Affinity Chromatography) (1993) 315-327; *C.A.*, 120 (1994) 186329p - a review with 26 refs.
- 2956 Larsson, P.O.: Support materials for affinity chromatography. *Chromatogr. Sci. Ser.*, 63(Handbook of Affinity Chromatography) (1993) 61-75; *C.A.*, 120 (1994) 186323g - a review with 16 refs.
- 2957 Ohlson, S. and Zopf, D.: Weak affinity chromatography. *Chromatogr. Sci. Ser.*, 63(Handbook of Affinity Chromatography) (1993) 299-314; *C.A.*, 120 (1994) 186328n - a review with 35 refs.
- 2958 Villems, R. and Toomik, P.: Overview [of affinity chromatography]. *Chromatogr. Sci. Ser.*, 63(Handbook of Affinity Chromatography) (1993) 3-60; *C.A.*, 120 (1994) 128673k - a review with 499 refs.
- 2959 Winzor, D.J. and Jackson, C.M.: Determination of binding constants by quantitative affinity chromatography. Current and future applications. *Chromatogr. Sci. Ser.*, 63(Handbook of Affinity Chromatography) (1993) 253-298; *C.A.*, 120 (1994) 128676p.

2960 Ziegler, T., Eckhardt, E., Strayle, J. and Herzog, H.: Synthesis of 5-aminopentyl 4,6-O-[(R)-1-carboxyethylidene]- β -D-galactopyranoside and its use as a ligand for the affinity chromatography of human serum amyloid P protein. *Carbohydr. Res.*, 253 (1994) 167-183; *C.A.*, 120 (1994) 264956g.

For additional information see *C.A.*:

120 (1994) 48625f, 49572y, 129023k, 129024m, 129084f, 186760x, 186762z, 212037p, 244366m, 245340k.

See also 2840, 3049, 3109, 3119, 3123, 3376, 3384, 3385, 3387, 3393, 3431, 3434, 3462, 3467, 3506, 3509, 3520, 3592, 3606, 3625, 3658.

4e. Functional analysis

2961 Ricci, M.C. and Cross, R.F.: Capillary electrophoresis: how good is this separation technique and how does it compare with HPLC. *Chem. Aust.*, 60 (1993) 540-546; *C.A.*, 120 (1994) 123834d - a review without refs.

4f. Trace analysis and pre-separation techniques

2962 Brinkman, U.A.T.: On-line sample treatment for or *via* column liquid chromatography. *J. Chromatogr. A*, 665 (1994) 217-231.

2963 Campíns-Falcó, P., Herráez-Hernández, R. and Sevillano-Cabeza, A.: On-line trace enrichment for the determination of ethacrynic acid in urine by liquid chromatography and column-switching. *Anal. Chim. Acta*, 284 (1993) 67-71.

2964 Land, C.C.J.: Solid-phase extraction of triazines from environmental water samples using aromatic sulfonic acid. *LC-GC Int.*, 7 (1994) 215-218.

2965 Majors, R.E.: Comparison of extraction disc and packed-bed cartridge technology in SPE (sample preparation perspectives). *LC-GC Int.*, 7 (1994) 70-72.

2966 Moreno Cordero, B., Perez Pavon, J.L., Garcia Pinto, C. and Fernandez Laespada, M.: Cloud point methodology: a new approach for preconcentration and separation in hydrodynamic systems of analysis. *Talanta*, 40 (1993) 1703-1710; *C.A.*, 120 (1994) 123837g.

2967 Van der Vlis, E., Irth, H., Tjaden, U.R. and van der Greef, J.: Potential of receptor-ligand interactions for sample handling in liquid and gas chromatography. *J. Chromatogr. A*, 665 (1994) 233-241.

For additional information see *C.A.*:

120 (1994) 158154j.

See also 2790, 2856, 2884, 2913, 2942, 3061, 3229, 3245, 3678, 3799, 3804, 3811, 3862, 4057, 4091, 4097, 4099.

4g. Enantiomers, separation

2968 Abu-Lafi, S., Sterin, M., Levin, S. and Mechoulam, R.: Structural features affecting chiral discrimination of terpene derivatives on a carbamated amylose stationary phase. *J. Chromatogr. A*, 664 (1994) 159-167.

2969 Allenmark, S.: Chiral discrimination by albumin: A mechanistic study of liquid chromatographic optical resolution of nonaromatic carboxylic acids. *Chirality*, 5 (1993) 295-299; *C.A.*, 120 (1994) 239337y.

2970 Anton, K., Eppinger, J., Frederiksen, L., Francotte, E., Berger, T.A. and Wilson, W.H.: Chiral separations by packed-column super- and subcritical fluid chromatography. *J. Chromatogr. A*, 666 (1994) 395-401.

2971 Belov, Y.P.: Chiral mobile phase in the enantiomeric analysis and in the evaluation of stability constants of π - π coomplexes. *Pure Appl. Chem.*, 65 (1993) 2273-2280; *C.A.*, 120 (1994) 152649c.

2972 Bertucci, C., Salvadori, P. and Lopes Guimaraes, L.F.: Determination of optical purity by high-performance liquid chromatography upon non-chiral stationary phases with dual circular dichroism/absorption detection. *J. Chromatogr. A*, 666 (1994) 535-539.

2973 Burns, D.T.: Cyclodextrins as versatile chiral recognition reagents for use in a variety of optical and separative analyses. *J. Pharm. Biomed. Anal.*, 12 (1994) 1-3 - a review with 39 refs.

2974 Cserhádi, T. and Forgács, E.: Charge-transfer chromatographic study of the interaction of non-ionic surfactants with hydroxypropyl- β -cyclodextrin. *J. Chromatogr. A*, 665 (1994) 17-25.

2975 Davankov, V.A.: Chiral selectors with chelating properties in liquid chromatography: fundamental reflections and selective review of recent developments. *J. Chromatogr. A*, 666 (1994) 55-76 - a review with 92 refs.

2976 Dingenen, J. and Kinkel, J.N.: Preparative chromatographic resolution of racemates on chiral stationary phases on laboratory and production scales by closed-loop recycling chromatography. *J. Chromatogr. A*, 666 (1994) 627-650.

2977 Dogan, I., Pustet, N. and Mannschreck, A.: The enantiomers of N-aryl-2-thioxo-4-oxazolidinones and N-arylrhodanines. Investigation by liquid chromatography, circular dichroism and thermal racemization. *J. Chem. Soc., Perkin Trans. 2*, (1993) 1557-1560; *C.A.*, 120 (1994) 54100x.

2978 Duff, K.J., Gray, H.L., Gray, R.J. and Bahler, C.C.: Chiral stationary phases in concert with homologous chiral mobile phase additives: push/pull model. *Chirality*, 5 (1993) 201-206; *C.A.*, 120 (1994) 190982p.

2979 Francotte, E.: Contribution of preparative chromatographic resolution to the investigation of chiral phenomena. *J. Chromatogr. A*, 666 (1994) 565-601 - a review with 212 refs.

2980 Gasparrini, F., Maggio, F., Misiti, D., Villani, C., Andreollini, F. and Mapelli, G.P.: High performance liquid chromatography on the chiral stationary phase (R,R)-DACH-DNB using carbon dioxide-based eluents. *J. High Resolut. Chromatogr.*, 17 (1994) 43-45.

2981 Gil-Av, E. and Schurig, V.: Resolution of non-racemic mixtures in achiral chromatographic systems: a model for the enantioselective effects observed. *J. Chromatogr. A*, 666 (1994) 519-525.

2982 Haginaka, J., Murashima, T. and Seyama, C.: Separation of enantiomers on a lysozyme-bonded silica column. *J. Chromatogr. A*, 666 (1994) 203-210.

2983 Hardt, I. and König, W.A.: Preparative enantiomer separation with modified cyclodextrins as chiral stationary phases. *J. Chromatogr. A*, 666 (1994) 611-615.

2984 Hermansson, J. and Hermansson, I.: Dynamic modification of the chiral bonding properties of a CHIRAL-AGP column by organic and inorganic additives. Separation of enantiomers of anti-inflammatory drugs. *J. Chromatogr. A*, 666 (1994) 181-191.

- 2985 Hyun, M.H., Cho, S.M., Ryoo, J.-J. and Kim, M.S.: A mechanistic evaluation for the resolution of enantiomers of α -arylpropionic acid derivatives on π -basic chiral stationary phases. *J. Liq. Chromatogr.*, 17 (1994) 317-328.
- 2986 Kaida, Y., and Okamoto, Y.: Optical resolution on regioselectively carbamoylated cellulose and amylose with 3,5-dimethylphenyl and 3,5-dichlorophenyl isocyanates. *Bull. Chem. Soc. Jpn.*, 66 (1993) 2225-2232; *C.A.*, 120 (1994) 77546m.
- 2987 Kirkland, K.M. and McCombs, D.A.: Changes in chiral selectivity with temperature for an ovomucoid protein-based column. *J. Chromatogr. A*, 666 (1994) 211-219.
- 2988 Koppenhoefer, B., Graf, R., Holzschuh, H., Nothdurft, A., Trettin, U., Piras, P. and Roussel, C.: CHIRBASE, a molecular database for the separation of enantiomers by chromatography. *J. Chromatogr. A*, 666 (1994) 557-563.
- 2989 Kurganov, A., Davankov, V., Unger, K., Eisenbeiss, F. and Kinkel, J.: Unusual peak distortion in ligand-exchange chromatography of enantiomers under overloaded conditions. *J. Chromatogr. A*, 666 (1994) 99-110.
- 2990 Lipkowitz, K.B.: Atomistic molecular modeling in chiral chromatography. *Am. Lab. (Shelton)*, 25 (1993) 20D, 20F, 20H; *C.A.*, 120 (1994) 53937p - a review with 10 refs.
- 2991 Lipkowitz, K.B.: Theoretical studies of brush-type chiral stationary phases. *J. Chromatogr. A*, 666 (1994) 493-503 - a review with 40 refs.
- 2992 Luthman, K., Jensen, A.V., Hacksell, U., Karlsson, A. and Pettersson, C.: Chiral discrimination of complexes between benzyloxy-carbonylglycyl-L-proline and 4-hydroxy-2-dipropylaminoindan in ion-pair chromatography. A molecular mechanics study. *J. Chromatogr. A*, 666 (1994) 527-534.
- 2993 McCalley, D.V.: Influence of analyte stereochemistry and basicity on peak shape of basic compounds in high-performance liquid chromatography with reversed-phase columns, using pyridine and alkyl-substituted derivatives as probe compounds. *J. Chromatogr. A*, 664 (1994) 139-147.
- 2994 Nesterenko, P.N., Krotov, V.V. and Staroverov, S.M.: Effect of mobile phase composition on the enantioselectivity of chromatographic separation on a quinine-bonded silica stationary phase. *J. Chromatogr. A*, 667 (1994) 19-28.
- 2995 Oi, N.: (Enantiomer separation by HPLC with chiral stationary phases derived from low molecular selectors). *Kuromatogurafi*, 14 (1993) 61R-70R; *C.A.*, 120 (1994) 207613u - a review with 113 refs.
- 2996 Ôi, N., Kitahara, H. and Aoki, F.: Enantiomer separation by high-performance liquid chromatography with (*R,R*)-tartaric acid mono-amide derivatives as bifunctional chiral selectors. *J. Chromatogr. A*, 666 (1994) 457-462.
- 2997 Tichý, M., Holanová, J. and Závada, J.: 6,6'-Dinitrobiphenyl-2,2'-dicarboxylic acid ionically bonded to aminopropyl silica: new axially chiral phase of C₂ symmetry for high-performance liquid chromatographic separation of enantiomeric amino alcohol derivatives. *J. Chromatogr. A*, 667 (1994) 11-17.
- 2998 Wainer, I.W.: Enantioselective high-performance liquid affinity chromatography as a probe of ligand-biopolymer interactions: an overview of a different use for high-performance liquid chromatographic chiral stationary phases. *J. Chromatogr. A*, 666 (1994) 221-224 - a review with 43 refs.
- 2999 Welch, C.J.: Evolution of chiral stationary phase design in the Pirkle laboratories. *J. Chromatogr. A*, 666 (1994) 3-26 - a review with 96 refs.
- See also 2831, 2858, 2860, 2870, 2873, 2887, 2890, 2902, 2904, 3054, 3099, 3137, 3163, 3167, 3168, 3214, 3226, 3233, 3234, 3236, 3256, 3257, 3263, 3268, 3272, 3283, 3297, 3334, 3415, 3685, 3699, 3700, 3703, 3706, 3723, 3738, 3758, 3812, 3845, 3850, 3851, 3853, 3860, 3861, 3867, 3874, 3875, 3879, 3883, 3889, 3893, 3897, 3901, 3902, 3908, 3917, 3919, 3923, 3929, 3961, 3963, 3967, 3969, 3978, 3990, 4010, 4066.
- 4h. Other special techniques
- 3000 Foucault, A.P., Camacho Frias, E., Bordier, C.G. and le Goffic, F.: Centrifugal partition chromatography: stability of various bi-phasic systems and pertinence of the "Stoke's model" to describe the influence of the centrifugal field upon the efficiency. *J. Liq. Chromatogr.*, 17 (1994) 1-17.
- 3001 Mang, V.: Two-phase liquid chromatography. *Ger. DE* 4,232,781 (Cl. B01D57/02), 28 Oct. 1993, Appl. 30 Sep. 1992; 4 p.; *C.A.*, 120 (1994) 80538r.
- 3002 Meester, R., van Buel, M.J., van der Wielen, L.A.M., Verpoorte, R. and Luyben, K.C.A.M.: Hydrodynamics of centrifugal partition chromatography. In: Weijnen, M.P.C. and Drinkenburg, A.A.H. (Editors), *Precis. Process Technol.*, 1st 1992, Kluwer, Dordrecht, 1993, pp. 295-303; *C.A.*, 120 (1994) 167477u.
- 3003 Shiragami, N.: Particles size analysis for micro-particles by capillary hydrodynamic chromatography. *Kona (Hirakata)*, 9 (1991) 13-18; *C.A.*, 120 (1994) 87536x.
- For additional information see *C.A.*:
120 (1994) 157817r.
- See also 2783, 2804, 2911, 2949, 3258, 3679, 3810, 3985, 4070, 4093.
5. HYDROCARBONS AND HALOGEN DERIVATIVES
- 5a. Aliphatic hydrocarbons
- For additional information see *C.A.*:
120 (1994) 192338g.
- 5b. Cyclic hydrocarbons, fullerenes
- 3004 Braun, T.: (Analytical chemistry of fullerenes and chemical analysis with fullerenes). *Magy. Kem. Foly.*, 99 (1993) 486-493; *C.A.*, 120 (1994) 94071v - a review with 22 refs.
- 3005 Darwish, A.D., Krotov, H.W., Taylor, R. and Walton, D.R.M.: Improved chromatographic separation of C₆₀ and C₇₀. *J. Chem. Soc., Chem. Commun.*, (1994) 15-16; *C.A.*, 120 (1994) 181795c.
- 3006 Dong, M.W., Duggan, J.X. and Stefanou, S.: A quick-turnaround HPLC method for the analysis of polynuclear aromatic hydrocarbons in soil, water and waste oil. *LC-GC*, 11 (1993) 802-810; *C.A.*, 120 (1994) 68453b.

- 3007 Edwards, E.A., Edwards, A.M. and Grbic-Galic, D.: A method for detection of aromatic metabolites at very low concentrations: application to detection of metabolites of anaerobic toluene degradation. *Appl. Environ. Microbiol.*, 60 (1994) 323-327; *C.A.*, 120 (1994) 100920q.
- 3008 García Pinto, C., Pérez Pavón, J.L. and Moreno Cordero, B.: Cloud point preconcentration and high-performance liquid chromatographic determination of polycyclic aromatic hydrocarbons with fluorescence detection. *Anal. Chem.*, 66 (1994) 874-881.
- 3009 Li, H. and Westerholm, R.: Determination of mono- and di-nitro polycyclic aromatic hydrocarbons by on-line reduction and high-performance liquid chromatography with chemiluminescence detection. *J. Chromatogr. A*, 664 (1994) 177-182.
- 3010 Liu, G., Djordjevic, N. and Erni, F.: The application of 'Z' shaped UV cells in open-tubular capillary column liquid chromatography. *Chromatographia*, 38 (1994) 313-316.
- 3011 Liu, M., Thienpont, A., Delville, M.H., Félix, G. and Netter, C.: Rapid separations of fullerenes by donor - acceptor liquid and supercritical fluid chromatography. *J. High Resolut. Chromatogr.*, 17 (1994) 1040-106.
- 3012 Nabivach, V.M. and Dmitriyev, V.P.: (Chromatographic detection of polycyclic aromatic hydrocarbons in coking products. Review). *Koks Khim.*, (1993) 29-34; *C.A.*, 120 (1994) 80925q - a review with 64 refs.
- 3013 Reupert, R., Brausen, G. and Witkowski, S.: Separation of polycyclic aromatic hydrocarbons by HPLC with fluorescence detection. *Chem. Anal. (Warsaw)*, 38 (1993) 463-467; *C.A.*, 120 (1994) 182090f.
- 3014 Rodríguez Delgado, M.A., Sánchez, M.J., Gonzáles, V. and García Montelongo, F.: Solute-micelle association constants of some polynuclear aromatic hydrocarbons by unicellular liquid chromatography with alcohol additives to mobile phase. *Chromatographia*, 38 (1994) 342-348.
- 3015 Stalling, D.L., Saim, S., Guo, C. and Kuo, K.: Separation and recovery of C₆₀- and C₇₀-fullerenes from carbon soot by supercritical fluid extraction and adsorption chromatography. *U.S. Pat.* 5,281,406 (Cl. 423-445B; C01B31/00), 25 Jan. 1994, *Appl.* 874,473, 22 Apr. 1992; 29 p.; *C.A.*, 120 (1994) 168100c.
- 3016 Von Schnering, H.G., Hoenle, W., Mittelbach, A., Carlsen, J., Janiak, R. and Quast, H.: Manufacture and chromatographic separation of C₆₀, C₇₀- and higher fullerenes. *Ger. Offen. DE* 4,224,556 (Cl. C01B31/00), 27 Jan. 1994, *Appl.* 24 Jul. 1992; 13 p.; *C.A.*, 120 (1994) 138664x.
- 3017 Yabiku, H.Y., Martins, M.S. and Takahashi, M.Y.: Levels of benzo[a]pyrene and other polycyclic aromatic hydrocarbons in liquid smoke flavor and some smoked foods. *Food Addit. Contam.*, 10 (1993) 399-405; *C.A.*, 120 (1994) 161972e.
- For additional information see *C.A.*:
120 (1994) 152057h, 191002f.
- See also 2879, 2880, 2892, 2898, 2980, 3668.
- 5c. *Halogen derivatives*
- 3018 Gianotti, G., Levi, M. and Turri, S.: Molecular weight characterization of fluoropolyoxyalkylenes by gel permeation chromatography. *J. Appl. Polym. Sci.*, 51 (1994) 973-978; *C.A.*, 120 (1994) 192796e.
- 3019 Kočan, A., Petřík, J., Chovancová, J. and Drobná, B.: Method for the group separation of non-ortho-, mono-ortho- and multi-ortho-substituted polychlorinated biphenyls and polychlorinated dibenzo-p-dioxins/polychlorinated dibenzofurans using activated carbon chromatography. *J. Chromatogr. A*, 665 (1994) 139-153.
- 3020 Wells, D.E. and Echarrí, I.: Determination of chlorobiphenyls with the separation of non-ortho, mono-ortho and di-ortho chloro congeners in fish and sea mammals. *Anal. Chim. Acta*, 286 (1994) 431-449.
- See also 2901, 3010, 3027.
- 5d. *Complex hydrocarbon mixtures (incl. analysis of tars, bitumens and mineral oils)*
- 3021 Leont'eva, S.A., Timofeeva, A.N., Kvasova, V.A. and Shevchenko, M.G.: (Use of chromatography in the study of petroleum products). *Khim. Prom.-st. (Moscow)*, (1993) 295-300; *C.A.*, 120 (1994) 111003u - a review with 3 refs.
- 3022 Overfield, R.E.: Method and equipment for chromatographic analysis of hydrocarbons, and hydrocarbon refining operations using them. *Can. Pat.* 1,324,897 (Cl. G01N33/28), 07 Dec. 1993, *Appl.* 598,633, 03 May 1989; 66 p.; *C.A.*, 120 (1994) 168490e.
- 3023 Tishenko, V.A., Shabalina, T.N., Lobzin, E.V., Polyakova, A.A. and Kalinina, L.D.: (Evaluation of aging of hydraulic oils). *Khim. Tekhnol. Topl. Masel*, (1993) 35-36; *C.A.*, 120 (1994) 111224s.
- 3024 Zakupra, V.A., Tkachev, V.T., Krygina, P.M. and Dogadin, O.B.: (Evaluation of the degree of aging of motor oil in a diesel engine). *Khim. Tekhnol. Topl. Masel*, (1993) 26-28; *C.A.*, 120 (1994) 111231s.
6. ALCOHOLS
- 3025 Gamoh, K. and Brooks, C.J.W.: Stability and reversed-phase liquid chromatographic studies of cyclic boronates. *Anal. Sci.*, 9 (1993) 549-555; *C.A.*, 120 (1994) 68435x.
- 3026 Goss, J.: Reverse phase high performance liquid chromatographic determination of primary and secondary aliphatic alcohols as phthalate monoesters by UV detection. *Chromatographia*, 38 (1994) 417-420.
- 3027 Hoh, H., Ikeda, S. and Ichinose, N.: Determination of 2,2,2-trichloroethanol in plasma and urine by ion-exclusion chromatography. *Analyst (London)*, 119 (1994) 409-413.
- 3028 Jansen, E.H.J., van den Berg, R.H. and Kroese, E.D.: Liquid chromatographic analysis and stability of benzo[a]pyrene-tetrols in blood protein adducts in rats after exposure to benzo[a]pyrene. *Anal. Chim. Acta*, 290 (1994) 86-93.
- 3029 Johansson, K., Joensson-Pettersson, G., Gorton, L., Markovarga, G. and Csoregi, E.: A reagentless amperometric biosensor for alcohol detection in column liquid chromatography based on co-immobilized peroxidase and alcohol oxidase in carbon paste. *J. Biotechnol.*, 31 (1993) 301-316; *C.A.*, 120 (1994) 100871z.
- 3030 Trathnigg, B., Thamer, D., Yan, X., Maier, B., Holzbauer, H.-R. and Much, H.: Characterization of ethoxylated fatty alcohols using liquid chromatography with density and refractive index detection. II. Quantification in liquid chromatography under critical conditions. *J. Chromatogr. A*, 665 (1994) 47-53.

- 3031 Yamaguchi, S., Inazu, A., Deyashiki, Y., Nakayama, T., Sato, K., Miyabe, Y. and Hara, A.: Stereospecificity of *trans*-dihydrodiol oxidation by dimeric and monomeric dihydrodiol dehydrogenases from mammalian tissues. *J. Biochem. (Tokyo)*, 115 (1994) 493-496.

For additional information see C.A.:

120 (1994) 49219p, 68497u, 218245b.

See also 2864, 3074, 3176, 3183, 3717, 4042.

7. PHENOLS

- 3032 Lamprecht, G. and Huber, J.F.K.: Ultra-trace analysis of phenols in water using high-performance liquid chromatography with on-line reaction detection. *J. Chromatogr. A*, 667 (1994) 47-57.
- 3033 Makuch, B., Gazda, K. and Kaminski, M.: Determination of phenol and monochlorophenols in water by reversed-phase liquid chromatography. *Anal. Chim. Acta*, 284 (1993) 53-58.
- 3034 Parker, M.H., McCann, D.J. and Mangold, J.B.: Sulfation of di- and tricyclic phenols by rat liver aryl sulfotransferase isoenzymes. *Arch. Biochem. Biophys.*, 310 (1994) 325-331.
- 3035 Pezet, R., Pont, V. and Cuenat, P.: Method to determine resveratrol and pterostilbene in grape berries and wines using high-performance liquid chromatography and highly sensitive fluorimetric detection. *J. Chromatogr. A*, 663 (1994) 191-197.
- 3036 Pocerull, E., Calull, M., Marcé, R.M. and Borrull, F.: Comparative study of solid-phase extraction of phenolic compounds. Influence of the ion-pair reagent. *Chromatographia*, 38 (1994) 579-584.

See also 3063, 3828, 4053.

8. SUBSTANCES CONTAINING HETEROCYCLIC OXYGEN

8a. Flavonoids

- 3037 Delgado, C., Tomás-Barberán, F.A., Talou, T. and Gaset, A.: Capillary electrophoresis as an alternative to HPLC for determination of honey flavonoids. *Chromatographia*, 38 (1994) 71-78.
- 3038 Engelhardt, U.H., Finger, A. and Kuhr, S.: Determination of flavone C glycosides in tea. *Z. Lebensm.-Unters. Forsch.*, 197 (1993) 239-244; *C.A.*, 120 (1994) 132504y.
- 3039 Garcia-Viguera, C., Tomas-Barberan, F.A., Ferreres, F., Artes, F. and Tomas-Lorente, F.: Determination of citrus jams genuineness by flavonoid analysis. *Z. Lebensm.-Unters. Forsch.*, 197 (1993) 255-259; *C.A.*, 120 (1994) 132506a.
- 3040 Hadj-Mahammed, M., Badjah-Hadj-Ahmed, Y. and Meklati, B.Y.: Behavior of polymethoxylated and polyhydroxylated flavones by carbon dioxide supercritical fluid chromatography with flame ionization and Fourier transform infrared detectors. *Phytochem. Anal.*, 4 (1993) 275-278; *C.A.*, 120 (1994) 186404j.
- 3041 Mouly, P.P., Arzouyan, C.R., Gaydou, E.M. and Estienne, J.M.: Differentiation of citrus juices by factorial discriminant analysis using liquid chromatography of flavanone glycosides. *J. Agric. Food Chem.*, 42 (1994) 70-79.

- 3042 Nogata, Y., Ohta, H., Yoza, K.-I., Berhow, M. and Hasegawa, S.: High-performance liquid chromatographic determination of naturally occurring flavonoids in *Citrus* with a photodiode-array detector. *J. Chromatogr. A*, 667 (1994) 59-66.

3043 Schmidt, P.C. and Ortega, G.G.: Passionsblumenkraut. Bestimmung des Gesamtflavonoidgehaltes von *Passiflorae herba*. *Dtsch. Apoth.-Ztg.*, 133 (1993) 17-26.

3044 Treutter, D., Santos-Buelga, C., Gutmann, M. and Kolodziej, H.: Identification of flavan-3-ols and procyanidins by high-performance liquid chromatography and chemical reaction detection. *J. Chromatogr. A*, 667 (1994) 290-297.

3045 Van Sumere, C., Fache, P., Vande Castelee, K., de Cooman, L., Everaert, E., de Loose, R. and Hutsebaut, W.: Improved extraction and reversed phase-high performance liquid chromatographic separation of flavonoids and the identification of *Rosa* cultivars. *Phytochem. Anal.*, 4 (1993) 279-292; *C.A.*, 120 (1994) 211711s.

3046 Zhu, B.T., Ezell, E.L. and Liehr, J.G.: Catechol-O-methyltransferase-catalyzed rapid O-methylation of mutagenic flavonoids. Metabolic inactivation as a possible reason for their lack of carcinogenicity *in vivo*. *J. Biol. Chem.*, 269 (1994) 292-299.

8b. Aflatoxins and other mycotoxins

3047 Bartolomé, B., Bengoechea, M.L., Pérez-Illarbe, F.J., Hernández, T., Estrella, I. and Gómez-Cordovés, C.: Determination of patulin in apple juice by high-performance liquid chromatography with diode-array detection. *J. Chromatogr. A*, 664 (1994) 39-43.

3048 Vudathala, D.K., Prelusky, D.B. and Trenholm, H.L.: Analysis of trace levels of deoxynivalenol in cow's milk by high pressure liquid chromatography. *J. Liq. Chromatogr.*, 17 (1994) 673-683.

3049 Ware, G.M., Umrigar, P.P., Carman, A.S., Jr. and Kuan, S.S.: Evaluation of fumonist immunoaffinity columns. *Anal. Lett.*, 27 (1994) 693-715.

8c. Other compounds with heterocyclic oxygen (incl. tannins)

3050 Gao, L. and Mazza, G.: Rapid method for complete chemical characterization of simple and acylated anthocyanins by high-performance liquid chromatography and capillary gas-liquid chromatography. *J. Agric. Food Chem.*, 42 (1994) 118-125.

3051 Laakso, J.A., Narske, E.D., Gloer, J.B., Wicklow, D.T. and Dowd, C.F.: Isokotanins A-C: new bicoumarins from the sclerotia of *Aspergillus alliaceus*. *J. Natural Prod.*, 57 (1994) 128-133.

3052 Pfeffer, M. and Walenciak-Reddel, E.: Determination of 6-amino-2,2-dimethyl-1,3-dioxepan-5-ol by post column derivatization with *o*-phthaldehyde/2-mercaptoethanol after ion-pair chromatography. *Chromatographia*, 38 (1994) 479-484.

For additional information see C.A.:

120 (1994) 62389p.

9. OXO COMPOUNDS, ETHERS, EPOXIDES AND QUINONES

- 3053 Bagchi, D., Moser, J. and Stohs, S.J.: Quantitative determination of urinary lipid metabolites by high pressure liquid chromatography as indicators of menadione-induced *in vivo* lipid peroxidation. *Arch. Environ. Contam. Toxicol.*, 26 (1994) 387-391; C.A., 120 (1994) 207805h.
- 3054 Chen, J. and Shum, W.: A direct HPLC method for the determination of enantiomeric excess of some highly enantiomerically enriched derivatives of chiral glycidols. *Tetrahedron Lett.*, 34 (1993) 7663-7666; C.A., 120 (1994) 68464f.
- 3055 Guichardant, M., Valette-Talbi, L., Cavadini, C., Crozier, G. and Berger, M.: Malondialdehyde measurement in urine. *J. Chromatogr. B*, 655 (1994) 112-116.
- 3056 Hoke, S.H., II, Cooks, R.G., Chang, C.-J., Kelly, R.C., Qyalls, S.J., Alvarado, B., McGuire, M.T. and Snader, K.M.: Determination of taxanes in *Taxus brevifolia* extracts by tandem mass spectrometry and high-performance liquid chromatography. *J. Natural Prod.*, 57 (1994) 277-286.
- 3057 Lehotay, J. and Halmo, P.: Determination of aliphatic aldehydes C₁-C₄ in waste gas by HPLC. *J. Liq. Chromatogr.*, 17 (1994) 847-854.
- 3058 Lehotay, J. and Hromulaková, K.: HPLC determination of trace levels of aliphatic aldehydes C₁-C₄ in river and tap water using on-line preconcentration. *J. Liq. Chromatogr.*, 17 (1994) 579-588.
- 3059 Lo Coco, F., Valentini, C., Novelli, V. and Ceccon, L.: High performance liquid chromatographic determination of 2-furaldehyde and 5-hydroxymethyl-2-furaldehyde in processed citrus juices. *J. Liq. Chromatogr.*, 17 (1994) 603-617.
- 3060 Nohta, H., Sakai, F., Kai, M., Ohkura, Y. and Saito, M.: 2-Amino-4,5-ethylenedioxyphenol as fluorescence derivatization reagent for aromatic aldehydes in liquid chromatography. *Anal. Chim. Acta*, 287 (1994) 223-227.
- 3061 Sanchez, D., Dosque, J.P., Janvier, G., Betbeder, A.M., Gauret, M.F. and Creppy, E.E.: (Residual formaldehyde in hospital respirators). *Bull. Soc. Pharm. Bordeaux*, 131 (1992) 69-75; C.A., 120 (1994) 62242k.
- 3062 Yamaguchi, M., Ishida, J., Xuan-Xuan, Z., Nakamura, M. and Yoshitake, T.: Determination of glyoxal, methylglyoxal, diacetyl, and 2,3-pentanedione in fermented foods by high-performance liquid chromatography with fluorescence detection. *J. Liq. Chromatogr.*, 17 (1994) 203-211.

For additional information see C.A.:
120 (1994) 47502b.

See also 2980, 4042.

10. CARBOHYDRATES

10a. Mono and oligosaccharides. Structural studies

- 3063 Amarowicz, R. and Shahidi, F.: Application of Sephadex LH-20 chromatography for the separation of cyanogenic glycosides and hydrophilic phenolic fraction from flaxseed. *J. Liq. Chromatogr.*, 17 (1994) 1291-1299.

- 3064 Coquet, A., Haerdi, W., Agosti, R.D. and Veuthey, J.-L.: Determination of sugars by liquid chromatography with post-column catalytic derivatization and fluorescence detection. *Chromatographia*, 38 (1994) 12-16.
- 3065 Fukamizo, T., Ohkawa, T., Ikeda, Y. and Goto, S.: Specificity of chitosanase from *Bacillus pumilus*. *Biochim. Biophys. Acta*, 1205 (1994) 183-188.
- 3066 Hagen, S.R.: Determination of neutral sugars in mycobacterial cell walls by high-performance liquid chromatography. *J. Liq. Chromatogr.*, 17 (1994) 97-104.
- 3067 Huang, A.S., Lam, S.Y., Nakayama, T.M. and Lin, H.: Microbiological and chemical changes in poi stored at 20° C. *J. Agric. Food Chem.*, 42 (1994) 45-48.
- 3068 Kakehi, K. and Honda, S.: Analysis of carbohydrates in glycoproteins by high-performance liquid chromatography and high-performance capillary electrophoresis. *Appl. Biochem. Biotechnol.*, 43 (1993) 55-71; C.A., 120 (1994) 128942x.
- 3069 Kubelka, V., Altmann, F., Kornfeld, G. and März, L.: Structures of the N-linked oligosaccharides of the membrane glycoproteins from three lepidopteran cell lines (Sf-21, IZD-Mb-0503, Bm-N). *Arch. Biochem. Biophys.*, 308 (1994) 148-157.
- 3070 Marko-Varga, G., Buttler, T., Gorton, L., Olsson, L., Durand, G. and Barceló, D.: Qualitative and quantitative carbohydrate analysis of fermentation substrates and broths by liquid chromatographic techniques. *J. Chromatogr. A*, 665 (1994) 317-332.
- 3071 Matsumoto, K., Maeda, Y., Kato, S. and Yuki, H.: Alteration of asparagine-linked glycosylation in serum transferrin of patients with hepatocellular carcinoma. *Clin. Chim. Acta*, 224 (1994) 1-8.
- 3072 Niwa, T., Dewald, L., Sone, J., Miyazaki, T. and Kajita, M.: Quantification of serum 1,5-anhydroglucitol in uremic and diabetic patients by liquid chromatography/mass spectrometry. *Clin. Chem. (Washington)*, 40 (1994) 260-264.
- 3073 Niwa, T., Wada, Y., Niwa, Y., Koyama, M., Deguchi, R., Maeda, K. and Kajita, M.: (Quantification of 1,5-anhydroglucitol in uremic serum by LC/MS). *Nippon Iyo Masu Supokutoru Gakkai Koenshu*, 18 (1993) 185-188; C.A., 120 (1994) 101132w.
- 3074 Oehlike, J., Brudel, M. and Blasig, I.E.: Benzoylation of sugars, polyols and amino acids in biological fluids for high-performance liquid chromatographic analysis. *J. Chromatogr. B*, 655 (1994) 105-111.
- 3075 Omichi, K. and Hase, S.: Linkage position analysis of pyridyl-amino-disaccharides by HPLC of fluorogenic Smith degradation products. *J. Biochem. (Tokyo)*, 115 (1994) 429-434.
- 3076 Rothenberg, B.E., Hayes, B.K., Toomre, D., Manzi, A.E. and Varki, A.: Biotinylated diaminopyridine: an approach to tagging oligosaccharides and exploring their biology. *Proc. Natl. Acad. Sci. U.S.A.*, 90 (1993) 11939-11943; C.A., 120 (1994) 128807g.
- 3077 Saska, M., Wu, M.D., Clarke, S.J., Iqbal, K. and Mrini, M.: Applications of continuous chromatographic separation of salt-sugar solutions. *Int. Sugar J.*, 95 (1993) 247-254; C.A., 120 (1994) 79790k.
- 3078 Simms, P.J., Hicks, K.B., Haines, R.M., Hotchkiss, A.T., Jr. and Osman, S.F.: Separation of lactose, lactobionic acid and lactobionolactone by high-performance liquid chromatography. *J. Chromatogr. A*, 667 (1994) 67-73.
- 3079 Slaughter, L.H. and Livingston, D.P., III: Separation of fructan isomers by high performance anion exchange chromatography. *Carbohydr. Res.*, 253 (1994) 287-291; C.A., 120 (1994) 186429w.

- 3080 Smouter, H. and Simpson, R.J.: Extraction and purification of preparative amounts of 1-kestose, 6-kestose, neokestose, nystose and inulin-pentasaccharide. *Stud. Plant Sci.*, 3(Inulin and Inulin-Containing Crops) (1993) 107-113; *C.A.*, 120 (1994) 264942z.
- 3081 Stefansson, M. and Westerlund, D.: Ligand exchange chromatography of carbohydrates at alkaline pH: effects of mobile phase additives and temperature. *J. Chromatogr. Sci.*, 32 (1994) 46-49.
- 3082 Stred'anský, M., Tomáška, M., Tomašková, A. and Šturdík, E.: Lactose hydrolysis in an aqueous two-phase system by whole-cell β -galactosidase of *Kluyveromyces marxianus*: partition and separation characteristics. *J. Chromatogr. A*, 668 (1994) 179-183.
- 3083 Takahashi, Y. and Goto, S.: Continuous separation of fructooligosaccharides using an annular chromatograph. *Sep. Sci. Technol.*, 29 (1994) 1311-1318; *C.A.*, 120 (1994) 215635t.
- For additional information see *C.A.*:
120 (1994) 49332v, 79699n, 109820q, 161859y.
- See also 2910, 2917, 3088, 3091, 3095, 3108, 3111, 3114, 3131, 3154, 3613, 3717, 3844.
- 10b. *Polysaccharides, mucopolysaccharides, lipopolysaccharides*
- 3084 Arvaniti, A., Karamanos, N.K., Dimitracopoulos, G. and Anastasiou, E.D.: Isolation and characterization of a novel 20 kDa sulfated polysaccharide from the extracellular slime layer of *Staphylococcus epidermidis*. *Arch. Biochem. Biophys.*, 308 (1994) 432-438.
- 3085 Bradbury, A.G.W. and Bello, A.B.: Determination of molecular size distribution of starch and debranched starch by a single procedure using high-performance size-exclusion chromatography. *Cereal Chem.*, 70 (1993) 543-547; *C.A.*, 120 (1994) 132518f.
- 3086 Carrino, D.A., Dennis, J.E., Drushel, R.F., Haynesworth, S.E. and Caplan, A.I.: Identity of the core proteins of the large chondroitin sulphate proteoglycans synthesized of skeletal muscle and prechondrogenic mesenchyme. *Biochem. J.*, 298 (1994) 51-64.
- 3087 Chatterton, N.J., Harrison, P.A., Thornley, W.R. and Bennett, J.H.: Separation and quantification of fructan (inulin) oligomers by anion-exchange chromatography. *Stud. Plant Sci.*, 3(Inulin and Inulin-Containing Crops) (1993) 93-99; *C.A.*, 120 (1994) 53041s.
- 3088 Fedders, G., Kock, R., van de Leur, E. and Greiling, H.: The metabolism of 2-fluoro-2-deoxy-D-glucose in human chondrocytes and its incorporation into keratan sulfate proteoglycans. *Eur. J. Biochem.*, 219 (1994) 1063-1071.
- 3089 Fischer, D.-C., Kolbe-Busch, S., Stöcker, G., Hoffmann, A. and Haubeck, H.-D.: Development of enzyme immunoassays specific for keratan sulphate- and core-protein-epitopes of the large aggregating proteoglycan from human articular cartilage. *Eur. J. Clin. Chem. Clin. Biochem.*, 32 (1994) 285-291.
- 3090 Gardiner, E.E., Mok, S.S., Sriratana, A., Robinson, H.C., Veitch, B.J.A., Lowther, D.A. and Handley, C.J.: Polymorphonuclear neutrophils release ^{35}S -labelled proteoglycans into cartilage during frustrated phagocytosis. *Eur. J. Biochem.*, 221 (1994) 871-879.
- 3091 Ip, C.C.Y., Miller, W.J., Silverklang, M., Mark, G.E., Ellis, R.W., Huang, L., Glushka, J., van Halbeek, H., Zhu, J. and Alhadeff, J.A.: Structural characterization of the N-glycans of a humanized anti-CD18 murine immunoglobulin G. *Arch. Biochem. Biophys.*, 308 (1994) 387-399.
- 3092 Kasinath, B.S., Block, J.A., Singh, A.K., Terhune, W.C., Maldonado, R., Dalavath, S., Kallgren, M.J. and Wanna, L.: Regulation of rat glomerular epithelial cell proteoglycans by high-glucose medium. *Arch. Biochem. Biophys.*, 309 (1994) 149-159.
- 3093 Kocharova, N.A., Thomas-Oates, J.E., Knirel, Y.A., Shashkov, A.S., Dabrowski, U., Kochetkov, N.K., Stanislavsky, E.S. and Kholodkova, E.V.: The structure of the O-specific polysaccharide of *Citrobacter* O16 containing glycerol phosphate. *Eur. J. Biochem.*, 219 (1994) 653-661.
- 3094 Komatsu, H., Takahata, T., Tanaka, M., Ishimitsu, S. and Okada, S.: Determination of the molecular-weight distribution of low-molecular-weight heparins using high-performance gel permeation chromatography. *Biol. Pharm. Bull.*, 16 (1993) 1189-1193; *C.A.*, 120 (1994) 239357e.
- 3095 Li, Y.T., Yuzyuk, J.A., Li, S.-C., Nematalla, A., Hasegawa, A., Kimura, M. and Nakagawa, H.: A novel sialidase capable of cleaving 3-deoxy-D-glycero-D-galacto-2-nonulosonic acid (KDN). *Arch. Biochem. Biophys.*, 310 (1994) 243-246.
- 3096 Liu, G., Goodall, D.M. and Loran, J.S.: Quantification of trace levels of β -cyclodextrin and substitution patterns in hydroxypropyl- β -cyclodextrin using high-performance liquid chromatography with polarimetric detection. *Chirality*, 5 (1993) 220-223; *C.A.*, 120 (1994) 144329h.
- 3097 Malovikova, A., Rinaudo, M. and Milas, M.: On the characterization of polygalacturonate salts in dilute solution. *Carbohydr. Polym.*, 22 (1993) 87-92; *C.A.*, 120 (1994) 57043s.
- 3098 Mumtaz, S. and Bachhawat, B.K.: Enhanced intracellular stability and efficacy of PEG modified dextranase in the treatment of a model storage disorder. *Biochim. Biophys. Acta*, 1199 (1994) 175-182.
- 3099 Okamoto, Y. and Kaida, Y.: Resolution by high-performance liquid chromatography using polysaccharide carbamates and benzoates as chiral stationary phases. *J. Chromatogr. A*, 666 (1994) 403-419.
- 3100 Taeuchi, T. and Miwa, T.: Liquid chromatography of cyclodextrins with indirect photometric detection of phenolphthalein inclusion complexes. *Chromatographia*, 38 (1994) 453-455.
- 3101 Takaya, Y., Uchisawa, H., Hanamatsu, K., Narumi, F., Okuzaki, B.-i. and Matsue, H.: Novel fucose-rich glycosaminoglycans from squid ink bearing repeating unit of trisaccharide structure $(-6\text{GalNAc}\alpha 1-3\text{Glc}-\beta 1-3\text{Fuc}\alpha 1-)_n$. *Biochem. Biophys. Res. Commun.*, 198 (1994) 560-567.
- 3102 Terbojevich, M., Cosani, A., Foher, B. and Marsano, E.: High-performance gel-permeation chromatography of chitosan samples. *Carbohydr. Res.*, 250 (1993) 301-314; *C.A.*, 120 (1994) 167186y.
- 3103 Volpi, N.: Fractionation of heparin, dermatan sulfate, and chondroitin sulfate by sequential precipitation: a method to purify a single glycosaminoglycan species from a mixture. *Anal. Biochem.*, 218 (1994) 382-391.
- For additional information see *C.A.*:
120 (1994) 49225n, 186406m, 211732z, 239342w.
- See also 2986, 3077, 3111, 3604.

- 10c. *Glycoproteins and their constituents*
- 3104 Arden, S.D., Rutherford, N.G., Guest, P.C., Curry, W.J., Bailyes, E.M., Johnston, C.F. and Hutton, J.C.: The post-translational processing of chromogranin A in the pancreatic islet: involvement of the eukaryote subtilisin PC2. *Biochem. J.*, 298 (1994) 521-528.
- 3105 Babál, P., Pindak, F.F., Wells, D.J. and Gardner, W.A., Jr.: Purification and characterization of a sialic acid-specific lectin from *Trichomonas mobilensis*. *Biochem. J.*, 299 (1994) 341-346.
- 3106 Beninati, S., Senger, D.R., Cordella-Miele, E., Mukherjee, A.B., Chackalaparampil, I., Shanmugam, V., Singh, K. and Mukherjee, B.B.: Osteopontin: its transglutaminase-catalyzed posttranslational modifications and cross-linking to fibronectin. *J. Biochem. (Tokyo)*, 115 (1994) 675-682.
- 3107 Bergman, A.-S. and Carlsson, S.R.: Saponin-induced release of cell-surface-anchored Thy-1 by serum glycosylphosphatidylinositol-specific phospholipase D. *Biochem. J.*, 298 (1994) 661-668.
- 3108 Bhat, U.R. and Helgeson, E.A.: An automated glycan release system for oligosaccharide mapping of a monoclonal antibody. *Am. Biotechnol. Lab.*, 12 (1994) 16; *C.A.*, 120 (1994) 189038d.
- 3109 Chatterjee, B.P., Sikdar, S. and Guha, A.K.: Affinity chromatography is a useful tool for isolation of lectins. In: Mukherjee, R.N. (Editor), *Downstream Process. Biotechnol., Proc. Int. Semin.*, 1991, McGraw-Hill, New Delhi, 1993, pp. 54-60; *C.A.*, 120 (1994) 264943a.
- 3110 Chou, C.-F., Riopel, C.L. and Omary, M.B.: Identification of a keratin-associated protein that localizes to a membrane compartment. *Biochem. J.*, 298 (1994) 457-463.
- 3111 Edge, A.S.B., Karasik, A., Yamada, K. and Kahn, C.R.: Effect of dexamethasone on the carbohydrate chains of the insulin receptor. *Biochem. Biophys. Res. Commun.*, 200 (1994) 852-859.
- 3112 Gegg, C.V. and Etzler, M.E.: Photoaffinity labeling of the adenine binding sites of two *Dolichos biflorus* lectins. *J. Biol. Chem.*, 269 (1994) 5687-5692.
- 3113 Hemmerich, S., Bertozzi, C.R., Leffler, H. and Rosen, S.D.: Identification of the sulfated monosaccharides of GlyCAM-1, an endothelial-derived ligand for L-selectin. *Biochemistry*, 33 (1994) 4820-4829.
- 3114 Hokke, C.H., Damm, J.B.L., Penninkhof, B., Aitken, R.J., Kamerling, J.P. and Vliegenthart, J.F.G.: Structure of the O-linked carbohydrate chains of porcine zona pellucida glycoproteins. *Eur. J. Biochem.*, 221 (1994) 491-512.
- 3115 Hu, R.-H., Mellors, A. and Bhavanandan, V.P.: Cleavage of epithectin, a mucin-type sialoglycoprotein, from the surface of human laryngeal carcinoma cells by a glycoprotease from *Pasteurella haemolytica*. *Arch. Biochem. Biophys.*, 310 (1994) 300-309.
- 3116 Itano, N., Oguri, K., Nakanishi, H. and Okayama, M.: Membrane-intercalated proteoglycan of a stroma-inducing clone from Lewis lung carcinoma binds to fibronectin via its heparan sulfate chains. *J. Biochem. (Tokyo)*, 114 (1993) 862-873.
- 3117 Ito, T., Hiraiwa, J., Nishioka, J., Iino, M. and Suzuki, K.: Characterization of functionally important regions of tissue factor by using monoclonal antibodies. *J. Biochem. (Tokyo)*, 114 (1993) 691-696.
- 3118 Jackson, D.E., Tetaz, T.J., Salem, H.H. and Mitchell, C.A.: Purification and characterization of two forms of soluble thrombomodulin from human urine. *Eur. J. Biochem.*, 221 (1994) 1079-1087.
- 3119 Kobayashi, R., Mizutani, A. and Hidaka, H.: Isolation and characterization of a 36-kDa microfibril-associated glycoprotein by the newly synthesized isoquinoline-sulfonamide affinity chromatography. *Biochem. Biophys. Res. Commun.*, 198 (1994) 1262-1266.
- 3120 Lepiku, M. and Järvi, J.: Optimization of synthesis of agarose-based lectin affinity sorbents. *Prepar. Biochem.*, 24 (1994) 61-67.
- 3121 Lindblom, A., Marsh, T., Fauser, C., Engel, J. and Paulsson, M.: Characterization of native laminin from bovine kidney and comparison with other laminin variants. *Eur. J. Biochem.*, 219 (1994) 383-392.
- 3122 Mandal, D.K., Nieves, E., Bhattacharya, L., Orr, G.A., Roboz, J., Yu, Q.-t. and Brewer, C.F.: Purification and characterization of three isolectins of soybean agglutinin. Evidence for C-terminal truncation by electrospray ionization mass spectrometry. *Eur. J. Biochem.*, 221 (1994) 547-553.
- 3123 Masumi, A., Akamatsu, Y. and Kitagawa, T.: Alteration by transforming growth factor- β 1 of asparagine-linked sugar chains in glucose transporter protein in Swiss 3T3 cells. *Biochim. Biophys. Acta*, 1221 (1994) 330-338.
- 3124 Mo, H., van Damme, E.J.M., Peumans, W.J. and Goldstein, I.J.: Isolation and characterization of an N-acetyl-D-galactosamine-binding lectin from Dutch Iris bulbs which recognizes the blood group A disaccharide (GalNAc α 1-3Gal). *J. Biol. Chem.*, 269 (1994) 7666-7673.
- 3125 Nagai, K., Shibata, K. and Yamaguchi, H.: Role of intramolecular high-mannose chains in the folding and assembly of soybean (*Glycine max*) lectin polypeptides: studies by the combined use of spectroscopy and gel-filtration size analysis. *J. Biochem. (Tokyo)*, 114 (1993) 830-834.
- 3126 Nitta, K., Ozaki, K., Ishikawa, M., Furusawa, S., Hosono, M., Kawauchi, H., Sasaki, K., Takayanagi, Y., Tsuiki, S. and Hakomori, S.: Inhibition of cell proliferation by *Rana catesbeiana* and *Rana japonica* lectins belonging to the ribonuclease superfamily. *Cancer Res.*, 54 (1994) 920-927.
- 3127 Okazaki, J., Gonda, Y., Kamada, A., Fujita, A. and Sakaki, T.: High-performance liquid chromatography analysis of chondroitin sulfate isomers in human periodontium. *J. Osaka Dent. Univ.*, 27 (1993) 101-106; *C.A.*, 120 (1994) 264952c.
- 3128 Rohrer, J.S.: Improved fractionation of sialylated glycopeptides by pellicular anion-exchange chromatography. *J. Chromatogr. A*, 667 (1994) 75-83.
- 3129 Rusiniak, M.E., Wagh, P.V., Bedi, G.S. and Back, N.: Purification and characterization of acute phase rat plasma thiostatin. *Prepar. Biochem.*, 24 (1994) 41-59.
- 3130 Schmid, H.P., Vallon, R., Tomek, W., Kreutzer-Schmid, C., Pouch, M.N., Badaoui, S., Boissonnet, G., Briand, M., Briand, Y. and Buri, J.: Glycosylation and deglycosylation of proteasomes (prosome) from calf-liver cells: high abundance of neuraminic acid. *Biochimie*, 75 (1993) 905-110; *C.A.*, 120 (1994) 72359t.
- 3131 Spillmann, D. and Finne, J.: Identification of a major poly-N-acetylglucosamine-containing cell-surface glycoprotein of mouse teratocarcinoma cells. Appearance on cells induced to primitive endoderm but not parietal endoderm differentiation. *Eur. J. Biochem.*, 220 (1994) 385-394.

- 3132 Van Damme, E.J.M., Smeets, K., Torrekens, S., van Leuven, F. and Peumans, W.J.: Characterization and molecular cloning of mannose-binding lectins from the Orchidaceae species *Listera ovata*, *Epipactis helleborine* and *Cymbidium hybrid*. *Eur. J. Biochem.*, 221 (1994) 769-777.
- For additional information see C.A.:
120 (1994) 186768f, 239572w.
- See also 3068, 3322, 3402, 3449, 3497, 3505, 3527, 3554.
11. ORGANIC ACIDS AND LIPIDS
- 11a. Organic acids and simple esters
- 3133 Aghabeigi, B. and Henderson, B.: Observations on the measurement of 2,3- and 2,5-dihydroxybenzoic acid using high performance liquid chromatography. *Inflammopharmacology*, 1 (1992) 363-368; C.A., 120 (1994) 123974z.
- 3134 Ahrens, M., Schulte, E. and Weber, K.: Fettsäuremethylester, Umesterung mit Trimethylsulfoniumhydroxid (Schnellverfahren)-Gemeinschaftsarbeiten der DGF, 138. Mitteilung: Deutsche Einheitsmethoden zur Untersuchung von Fetten, Fettprodukten, Tensiden und verwandten Stoffen, 105. Mitt.: Analyse von Fette XXXI. *Fat Sci. Technol.*, 96 (1994) 67-68.
- 3135 Akasaka, K., Akama, T., Ohru, H. and Meguro, H.: Measurement of hydroxy and hydroperoxy fatty acids by a high pressure liquid chromatography with a column-switching system. *Biosci., Biotechnol., Biochem.*, 57 (1993) 2016-2019; C.A., 120 (1994) 157798k.
- 3136 Akoh, C.C.: Enzymatic synthesis of acetylated glucose fatty acid esters in organic solvent. *J. Am. Oil Chem. Soc.*, 71 (1994) 319-323.
- 3137 Borges, M.F.M. and Pinto, M.M.M.: Separation of the diastereoisomers of ethyl esters of caffeic, ferulic, and isoferulic acids by thin-layer and high performance liquid chromatography. *J. Liq. Chromatogr.*, 17 (1994) 1125-1139.
- 3138 Burini, G.: Determination of the alkyl esters of *p*-hydroxybenzoic acid in mayonnaise by high-performance liquid chromatography and fluorescence labelling. *J. Chromatogr. A*, 664 (1994) 213-219.
- 3139 Cecchini, F. and Morassut, M.: (HPLC determination of sorbic acid in wine). *Boll. Chim. Ig., Parte Sci.*, 44(S5) (1993) 315-321; C.A., 120 (1994) 189938x.
- 3140 Fischer, K., Corsten, C., Leidmann, P., Bieniek, D. and Ketrup, A.: Determination of short chain aliphatic acids in silage effluents by ion exclusion chromatography. *Chromatographia*, 38 (1994) 43-51.
- 3141 Gachanja, A. and Worsfold, P.: Determination of aliphatic carboxylic acids in non-aqueous matrices by liquid chromatography with peroxyoxalate chemiluminescence detection. *Anal. Chim. Acta*, 290 (1994) 226-232.
- 3142 Gordon, J.A., Heller, S.K., Kaduce, T.L. and Spector, A.A.: Formation and release of a peroxisome-dependent arachidonic acid metabolite by human skin fibroblasts. *J. Biol. Chem.*, 269 (1994) 4103-4109.
- 3143 Hada, T., Hagiya, H., Suzuki, H., Arakawa, T., Nakamura, M., Matsuda, S., Yoshimoto, T., Yamamoto, S., Azekawa, T., Morita, Y., Ishimura, K. and Kim, H.-Y.: Arachidonate 12-lipoxygenase of rat pineal glands: catalytic properties and primary structure deduced from its cDNA. *Biochim. Biophys. Acta*, 1211 (1994) 221-228.
- 3144 Haldna, U. and Pentchuk, J.: Selectivity coefficient of aliphatic organic acid anions on the methacrylic matrix resin HIKS-1. *Eesti Tead. Akad. Toim., Keem.*, 41 (1992) 101-104; C.A., 120 (1994) 94382x.
- 3145 Harris, R.K., Greaves, J., Alexander, D., Wilson, T. and Haggerty, W.J.: Development of stability-indicating analytical methods for flaxseed lignans and their precursors. *ACS Symp. Ser.*, 547 (Food Phytochemicals for Cancer Prevention II) (1994) 295-305; C.A., 120 (1994) 161852r.
- 3146 Hatanaka, H., Yasui, Y. and Matsushita, S.: (Direct determination of phthalic acid esters in vodka by HPLC). *Shokuhin Eiseigaku Zasshi*, 35 (1994) 51-55; C.A., 120 (1994) 268446p.
- 3147 Helser, M.A. and Hotchkiss, J.H.: Comparison of tomato phenolic and ascorbate fractions on the inhibition of N-nitroso compound formation. *J. Agric. Food Chem.*, 42 (1994) 129-132.
- 3148 Holmes, C., Eisenhofer, G. and Goldstein, D.S.: Improved assay for plasma dihydroxyphenylacetic acid and other catechols using high-performance liquid chromatography with electrochemical detection. *J. Chromatogr. B*, 653 (1994) 131-138.
- 3149 Isbell, T.A., Kleiman, R. and Plattner, B.A.: Acid-catalyzed condensation of oleic acid into estolides and polyestolides. *J. Am. Oil Chem. Soc.*, 71 (1994) 169-174.
- 3150 Ito, T., Kidouchi, K., Kajita, M. and Niwa, T.: (LC/APCI-MS analysis of urinary isovalerylglycine in isovaleric acidemia). *Nippon Iyo Masu Supekutoru Gakkai Koenshu*, 18 (1993) 147-150; C.A., 120 (1994) 49490v.
- 3151 Ivanov, A.A. and Medved, A.L.: (Ion chromatography of carboxylic acids). *Zh. Anal. Khim.*, 48 (1993) 1539-1545; C.A., 120 (1994) 123847k.
- 3152 Kondo, Y., Kawai, Y., Miyazawa, T. and Mizutani, J.: Chemiluminescence-HPLC analysis of fatty acid hydroperoxide isomers. *Biosci., Biotechnol., Biochem.*, 57 (1993) 1575-1576; C.A., 120 (1994) 49218n.
- 3153 Lee, C. and Hajra, A.K.: Quantitative analysis of molecular species of diacylglycerol in biological samples. *Methods Neurosci.*, 18 (Lipid Metabolism in Signaling Systems) (1993) 190-199; C.A., 120 (1994) 264869f - a review with 16 refs.
- 3154 Lehtonen, P. and Vuorela, H.: Application of normal and second-order derivative spectroscopy in identifying organic acids and sugar acids in liquid chromatography with on-line photodiode array detection. *J. Liq. Chromatogr.*, 17 (1994) 1245-1255.
- 3155 Lewis, S.W., Worsfold, P.J. and McKerrell, E.H.: Monitoring carboxylic acid formation in engine oils by liquid chromatography with fluorescence detection. *J. Chromatogr. A*, 667 (1994) 91-98.
- 3156 Mason, S.R., Ward, L.C. and Reilly, P.E.B.: Fluorimetric detection of microsomal lauric acid hydroxylations using high-performance liquid chromatography after selective solvent partitioning and esterification with 1-pyrenyldiazomethane. *J. Liq. Chromatogr.*, 17 (1994) 619-632.
- 3157 Morris, J. and Fritz, J.S.: Separation of hydrophilic organic acids and small polar compounds using macroporous resin columns. *LC-GC Int.*, 7 (1994) 43-47.

- 3158 Nieroz, A., Breant, V., Guillaumont, M. and Aulagner, G.: Determination of β -hydroxybutyrate sodium and stability study in parenteral nutrition solutions. *J. Pharm. Clin.*, 12 (1993) 190-193; C.A., 120 (1994) 62410p.
- 3159 Powell, W.S., Zhang, Y. and Gravel, S.: Effects of phorbol myristate acetate on the synthesis of 5-oxo-6,8,11,14-eicosatetraenoic acid by human polymorphonuclear leukocytes. *Biochemistry*, 33 (1994) 3927-3933.
- 3160 Read, G., Richardson, N.R. and Wickens, D.G.: Determination of octadecadienoic acids in human serum, a critical appraisal. *Analyst (London)*, 119 (1994) 393-396.
- 3161 Rettenmeier, A.W., Hennigs, R. and Wodarz, R.: Determination of butoxyacetic acid and N-butoxyacetylglutamine in urine of lacquerers exposed to 2-butoxyethanol. *Int. Arch. Occup. Environ. Health*, 65(1, Suppl., Proceedings of International Symposium on Biological Monitoring, 1992) (1993) S151-S153; C.A., 120 (1994) 127039w.
- 3162 Sandmann, B.W. and Grayeski, M.L.: Quinoxalinone derivatization of biological carboxylic acids for detection by peroxyoxalate chemiluminescence with high-performance liquid chromatography. *J. Chromatogr. A*, 653 (1994) 123-130.
- 3163 Sinibaldi, M., Flieger, M., Cvak, L., Messina, A. and Pichini, A.: Direct resolution of optically active isomers on chiral packings containing ergoline skeletons. II. Enantioseparation of carboxylic acids. *J. Chromatogr. A*, 666 (1994) 471-478.
- 3164 Thomas, C.E., Jackson, R.L., Ohlweiler, D.F. and Ku, G.: Multiple lipid oxidation products of low density lipoproteins induce interleukin-1 beta release from human blood mononuclear cells. *J. Lipid Res.*, 35 (1994) 417-427.
- 3165 Von Schnakenburg, C., Byrd, D.J., Latta, K., Reusz, G.S., Graf, D. and Brodehl, J.: Determination of oxalate excretion in spot urines of healthy children by ion chromatography. *Eur. J. Clin. Chem. Clin. Biochem.*, 32 (1994) 27-29.
- 3166 Wilde, J. and Eggerer, H.: Substrate and inhibitor specificity of mevalonate kinase determined with substrate analogues. *Eur. J. Biochem.*, 221 (1994) 463-473.
- 3167 Winkler, F.J., Medina, R., Winkler, J. and Krause, H.: Mass spectral and semi-empirical studies on chiral discrimination in gaseous aggregation products of protonated dialkyl tartrates. *J. Chromatogr. A*, 666 (1994) 549-556.
- 3168 Yamamoto, A., Matsunaga, A., Mizukami, E., Hayakawa, K. and Miyazaki, M.: Enantiomeric purity determination by high-performance liquid chromatography with coupled polarized photometric/UV detection. Analysis of adulterative addition of synthetic malic and tartaric acids. *J. Chromatogr. A*, 667 (1994) 85-89.
- 3169 Ye, M., Walkup, R. and Hill, K.: Separation of T-MAZ ethoxylated sorbitan fatty acid esters by reverse phase chromatography. *Chromatographia*, 38 (1994) 337-341.
- For additional information see C.A.:
120 (1994) 49213g, 52993s, 182094k.
- See also 2969, 3050, 3067, 3182, 3241, 3782, 4053, 4126, 4127.
- 11b. *Prostaglandins*
- 3170 Bell, J.G., Tocher, D.R. and Sargent, J.R.: Effect of supplementation with 20:3(n-6), 20:4(n-6) and 20:5(n-3) on the production of prostaglandins E and F of the 1-, 2- and 3-series in turbot (*Scophthalmus maximus*) brain astroglial cells in primary culture. *Biochim. Biophys. Acta*, 1211 (1994) 335-342.
- 3171 De Waart, D.R., Koomen, G.C.M. and Wanders, R.J.A.: Studies on the urinary excretion of thromboxane B₂ in Zellweger patients and control subjects: evidence for a major role for peroxisomes in the β -oxidative chain-shortening of thromboxane B₂. *Biochim. Biophys. Acta*, 1226 (1994) 44-48.
- 3172 Reinke, M.: Quantitative analysis of 11-dehydro-thromboxane B₂ in human fluids by a combination of immunoaffinity chromatography and ELISA. *Pharm. Pharmacol. Lett.*, 3 (1994) 186-189; C.A., 120 (1994) 236315d.
- 11c. *Lipids and their constituents*
- 3173 Abidi, S.L. and Mounts, T.L.: Reversed-phase retention behavior of fluorescence labeled phospholipids in ammonium acetate buffers. *J. Liq. Chromatogr.*, 17 (1994) 105-122.
- 3174 Antonopoulou, S., Andrikopoulos, N.K. and Demopoulos, C.A.: Separation of the main neutral lipids into classes and species by RP-HPLC and UV detection. *J. Liq. Chromatogr.*, 17 (1994) 633-648.
- 3175 Arnoldsson, K.C. and Kaufmann, P.: Lipid class analysis by normal phase high performance liquid chromatography, development and optimization using multivariate methods. *Chromatographia*, 38 (1994) 317-324.
- 3176 Bianchi, G. and Vlahov, G.: Composition of lipid classes in the morphologically different parts of the olive fruit, cv. *Coratina* (*Olea europaea* Linn.). *Fat Sci. Technol.*, 96 (1994) 72-77.
- 3177 Button, D., Rothman, A., Bongiorno, C., Kupperman, E., Wolner, B. and Taylor, P.: Agonist-selective regulation of polyphosphoinositide metabolism in pulmonary artery smooth muscle cells. *J. Biol. Chem.*, 269 (1994) 6390-6398.
- 3178 Dasgupta, S., Hogan, E.L., Glushka, J. and Halbeek, H.: Branched monosialo gangliosides of the lacto-series isolated from bovine erythrocytes: characterization of a novel ganglioside, NeuGc-isooctacosylceramide. *Arch. Biochem. Biophys.*, 310 (1994) 373-384.
- 3179 Filip, V. and Kleinova, M.: Determination of mono-, di-, and triglycerides by HPLC. *Z. Lebensm.-Unters. Forsch.*, 196 (1993) 532-535; C.A., 120 (1994) 105196t.
- 3180 Gelsema, W.J., Choma, I., den Ouden, T.A.F., Zander, R. and van den Bosch, Z.: Quantitation of the diacyl, alkylacyl, and alk-1-enylacyl subclasses of choline glycerophospholipids by chemical dephosphorylation and benzylation. *Anal. Biochem.*, 217 (1994) 265-276.
- 3181 Hara, S. and Totani, Y.: (HPLC analysis of unoxidized and peroxide lipid mixtures). *Jasco Rep.*, 35 (1993) 87-91; C.A., 120 (1994) 72581j - a review with 26 refs.
- 3182 Husain, S. and Devi, K.S.: Separation and identification of isomeric conjugated fatty acids by high-performance liquid chromatography with photodiode array detection. *Lipids*, 28 (1993) 1037-1040; C.A., 120 (1994) 68459h.
- 3183 Indyk, H.E. and Woolard, D.C.: Determination of free myo-inositol in milk and infant formula by high-performance liquid chromatography. *Analyst (London)*, 119 (1994) 397-401.

- 3184 Klein, B.H. and Dudenhausen, J.W.: Simultaneous determination of phospholipid classes and the major molecular species of lecithin in human amniotic fluid by HPLC. *J. Liq. Chromatogr.*, 17 (1994) 981-998.
- 3185 Nakagawa, Y.: Application of paired-ion high-performance liquid chromatography to the separation of molecular species of phosphatidylinositol. *Lipids*, 28 (1993) 1033-1035; *C.A.*, 120 (1994) 49235r.
- 3186 Sarney, D.B., Fregapane, G. and Vulfson, E.N.: Lipase-catalyzed synthesis of lysophospholipids in a continuous bioreactor. *J. Am. Oil Chem. Soc.*, 71 (1994) 93-96.
- 3187 Taylor, D.C., MacKenzie, S.L., McCurdy, A.R., McVetty, P.B.E., Giblin, E.M., Pass, E.W., Stone, S.J., Scarth, R., Rimmer, S.R. and Pickard, M.D.: Stereospecific analyses of seed triacylglycerols from high-erucic acid Brassicaceae: detection of erucic acid at the sn-2 position in *Brassica oleracea* L. genotypes. *J. Am. Oil Chem. Soc.*, 71 (1994) 163-167.
- 3188 Vesterqvist, O., Sargent, C.A., Grover, G.J., Warrack, B.M., DiDonato, G.C. and Ogletree, M.L.: Characterization of rabbit myocardial phospholipase A₂ activity using endogenous phospholipid substrates. *Anal. Biochem.*, 217 (1994) 210-219.
- 3189 Vlahos, C.J., Matter, W.F., Hui, K.Y. and Brown, R.F.: A specific inhibitor of phosphatidylinositol 3-kinase, 2-(4-morpholinyl)-8-phenyl-4H-1-benzopyran-4-one (LY294002). *J. Biol. Chem.*, 269 (1994) 5241-5248.
- 3190 Yang, Q. and Lundahl, P.: Steric immobilization of liposomes in chromatographic gel beads and incorporation of integral membrane proteins into their lipid bilayers. *Anal. Biochem.*, 218 (1994) 210-221.
- For additional information see C.A.:
120 (1994) 49196d, 161845r.
- See also 3152, 3208, 3495, 3711, 3712.
- 11d. *Lipoproteins and their constituents*
- 3191 Facchinetti, R., Ruzzenente, O., Poli, G. and Guidi, G.C.: (Lipoprotein(a) determination: Comparison of methods and standardization problems). *G. Ital. Chim. Clin.*, 17 (1992) 375-385; *C.A.*, 120 (1994) 72648m.
- 3192 Fazio, S., Horie, Y., Simonet, W.S., Weisgraber, K.H., Taylor, J.M. and Rall, S.C., Jr.: Altered lipoprotein metabolism in transgenic mice expressing low levels of a human receptor-binding-defective apolipoprotein E variant. *J. Lipid Res.*, 35 (1994) 408-416.
- 3193 Himber, J., Missano, B. and Kuhl, H.: Lack of effect on the low density lipoprotein receptor in hamsters treated with 17 α -ethinyl estradiol. *Biochim. Biophys. Acta*, 1211 (1994) 359-363.
- 3194 Marques-Vidal, P., Azéma, C., Collet, X., Vieu, C., Chap, H. and Perret, B.: Hepatic lipase promotes the uptake of HDL esterified cholesterol by the perfused rat liver: a study using reconstituted HDL particles of defined phospholipid composition. *J. Lipid Res.*, 35 (1994) 373-384.
- 3195 Myara, I., Pico, I., Védie, B. and Moatti, N.: A method to screen for the antioxidant effect of compounds on low-density lipoprotein (LDL): Illustration with flavonoids. *J. Pharmacol. Toxicol. Methods*, 30 (1993) 69-73; *C.A.*, 120 (1994) 207778b.
- 3196 Shibusawa, Y., Miwa, N., Hirashima, T. and Matsumoto, U.: Separation of low density and very low density lipoproteins from human serum by hydroxyapatite chromatography. *J. Liq. Chromatogr.*, 17 (1994) 1203-1217.
- 3197 Talmud, P.J., Krul, E.S., Pessah, M., Gay, G., Schonfeld, G., Humphries, S.E. and Infante, R.: Donor splice mutation generates a lipid-associated apolipoprotein B 27.6 in a patient with homozygous hypobetalipoproteinemia. *J. Lipid Res.*, 35 (1994) 468-477.
- 3198 Weigel, D., Richter, O. and Arnold, K.: Partitioning of chemically modified low-density lipoprotein in aqueous polymer two-phase systems. *J. Chromatogr. A*, 668 (1994) 107-116.
- 3199 Young, S.G., Koduri, R.K., Austin, R.K., Bonnet, D.J., Smith, R.S. and Curtiss, L.K.: Definition of a nonlinear conformational epitope for the apolipoprotein B-100-specific monoclonal antibody, MB 47. *J. Lipid Res.*, 35 (1994) 399-407.
- For additional information see C.A.:
120 (1994) 186797q.
- See also 3164.
12. ORGANIC PEROXIDES
- See 2848, 3143, 4035.
13. STEROIDS
- 13a. *General techniques*
- 3200 Iwata, T., Hirose, T., Nakamura, M. and Yamaguchi, M.: Determination of urinary glucuronide conjugates by high-performance liquid chromatography with pre-column fluorescence derivatization. *J. Chromatogr. B*, 654 (1994) 171-176.
- 13b. *Pregnane and androstane derivatives*
- 3201 Ishii, Y., Mukoyama, H. and Ohtawa, M.: *In vitro* biotransformation of finasteride in rat hepatic microsomes. Isolation and characterization of metabolites. *Drug Metab. Disp.*, 22 (1994) 79-84.
- 3202 Joellenbeck, L., Qian, Z., Zarba, A. and Groopman, J.D.: Urinary 6 β -hydroxycortisol/cortisol ratios measured by high-performance liquid chromatography for use as a biomarker for the human cytochrome P-450 3A4. *Cancer Epidemiol., Biomarkers Prev.*, 1 (1992) 567-572; *C.A.*, 120 (1994) 238744s.
- 3203 Leyssens, L., Royackers, E., Gielen, B., Missotten, M., Schoofs, J., Czech, J., Noben, J.P., Hendriks, L. and Raus, J.: Metabolites of 4-chlorotestosterone acetate in cattle urine as diagnostic markers for its illegal use. *J. Chromatogr. B*, 654 (1994) 43-54.
- 3204 Surmann, J.P., Strahl, B. and Hock, C.: Bestimmung von Dexamethason und Retinolpalmitat in einer Emulsionssalbe. *Pharmazie*, 49 (1994) 139-142.
- 3205 Valenta, C. and Janout, H.: Corticosteroid analysis by HPLC with increased sensitivity by use of precolumn concentration. *J. Liq. Chromatogr.*, 17 (1994) 1141-1146.

For additional information see C.A.:
120 (1994) 62457j, 183118h.

See also 2937, 3200.

13c. Estrogens

3206 Ikegawa, S., Itoh, M. and Goto, J.: Separatory determination of biliary metabolites of equilin in rat by high-performance liquid chromatography. *J. Liq. Chromatogr.*, 17 (1994) 223-239.

3207 Lamparczyk, H., Zarzycki, P.K., Nowakowska, J. and Ochocka, R.J.: Application of β -cyclodextrins for the analysis of estrogenic steroids in human urine by high-performance liquid chromatography. *Chromatographia*, 38 (1994) 168-172.

See also 3200.

13d. Sterols

3208 Akasaka, K., Ohru, H. and Meguro, H.: Measurement of cholesterol ester hydroperoxides of high and combined low and very low density lipoprotein in human plasma. *Biosci., Biotechnol., Biochem.*, 58 (1994) 396-399; C.A., 120 (1994) 186431r.

3209 Kamal-Eldin, A. and Appelqvist, L.Å.: Variations in the composition of sterols, tocopherols and lignans in seed oils from four *Sesamum* species. *J. Am. Oil Chem. Soc.*, 71 (1994) 149-156.

For additional information see C.A.:
120 (1994) 49205f, 49473s.

See also 3176, 4029.

13e. Bile acids and alcohols

3210 Coca, E., Ribas, B., Trigueros, G., Mtnez-Sarmiento, J., Borgue, M., Ortega, D., Sobrino, A., Mallen, A., de Pablos, I. and Fdez-Represa, J.A.: A method for quick determination of bile acids in bile of patients with biliary lithiasis. *J. Liq. Chromatogr.*, 17 (1994) 1349-1363.

3211 Scalia, S. and Pazzi, P.: HPLC assay of conjugated bile acids in human fluids using on-line sample pretreatment on a standard isocratic chromatograph. *Clin. Chim. Acta*, 224 (1994) 181-190.

For additional information see C.A.:
120 (1994) 49206q, 62401m, 116905z.

13f. Ecdysones and other insect steroid hormones

3212 Modlin, R.F., Alred, P.A. and Tjerneld, F.: Utilication of temperature-induced phase separation for the purification of ecdysone and 20-hydroxyecdysone from spinach. *J. Chromatogr. A*, 668 (1994) 229-236.

14. STEROID GLYCOSIDES AND SAPONINS

3213 Crespín, F., Calmes, M., Elias, R., Maillard, C. and Balansard, G.: High-performance liquid chromatographic determination of saponins from *Hedera helix* L. using a light-scattering detector. *Chromatographia*, 38 (1994) 183-186.

For additional information see C.A.:
120 (1994) 86553p, 116909d.

See also 4027, 4028, 4031, 4032, 4033, 4034.

15. TERPENES AND OTHER VOLATILE AROMATIC COMPOUNDS

15a. Terpenes

3214 Armstrong, D.W. and Zukowski, J.: Direct enantiomeric resolution of monoterpene hydrocarbons via reversed-phase high-performance liquid chromatography with an α -cyclodextrin bonded stationary phase. *J. Chromatogr. A*, 666 (1994) 445-448.

3215 Khoshkhoo, N., Hedin, P.A. and McCarty, J.C., Jr.: Terpenoid aldehydes in root-knot nematode susceptible and resistant cotton plants. *J. Agric. Food Chem.*, 42 (1994) 204-208.

See also 2968, 3176, 3213, 4027, 4034, 4035.

15b. Essential oils

3216 Piry, J. and Příbela, A.: Application of high-performance liquid chromatography to the analysis of the complex volatile mixture of blackcurrant buds (*Ribes nigrum* L.) *J. Chromatogr. A*, 665 (1994) 105-109.

15c. Bitter substances

3217 Hermans-Lokkerbol, A.C.J. and Verpoorte, R.: Preparative separation and isolation of three α bitter acids from hop, *Humulus lupulus* L., by centrifugal partition chromatography. *J. Chromatogr. A*, 664 (1994) 45-53.

3218 Semenova, O.P., Timerbaev, A.R. and Bonn, G.K.: Application of high-performance liquid chromatography to the determination of bitter principles of pharmaceutical relevance. *J. Chromatogr. A*, 667 (1994) 327-333.

16. NITRO AND NITROSO COMPOUNDS

3219 Bellomonte, G., Filesi, C., Macri, A., Mosca, M. and Sanzini, E.: High-performance liquid chromatographic determination of nitrofurans and free chloramphenicol in poultry muscle, liver and eggs. *Ital. J. Food Sci.*, 5 (1993) 247-253; C.A., 120 (1994) 268441h.

3220 Jenkins, T.F., Miyares, P.H., Myers, K.F., McCormick, E.F. and Strong, A.B.: Comparison of solid phase extraction with salting-out solvent extraction for preconcentration of nitroaromatic and nitramine explosives from water. *Anal. Chim. Acta*, 289 (1994) 69-78.

3221 Veigl, E., Posch, W., Lindner, W. and Tritthart, P.: Selective and sensitive analysis of 1-nitropyrene in Diesel exhaust particulate extracts by multidimensional HPLC. *Chromatographia*, 38 (1994) 199-206.

- 3222 Verweij, A.M.A., de Bruyn, P.C.A.M., Choufoer, C. and Lipman, P.J.L.: Liquid chromatographic, thermospray/negative ion, tandem mass spectrometric (LC/TSP/MS/MS) analysis of some explosives. *Forensic Sci. Int.*, 60 (1993) 7-13; *C.A.*, 120 (1994) 127068e.
- For additional information see *C.A.*:
120 (1994) 47471r.
- See also 2864.
17. AMINES, AMIDES AND RELATED NITROGEN COMPOUNDS
- 17a. *Amines and polyamines*
- 3223 Busto, O., Valero, Y., Guasch, J. and Borrull, F.: Solid phase extraction applied to the determination of biogenic amines in wines by HPLC. *Chromatographia*, 38 (1994) 571-578.
- 3224 Cann-Moisan, C., Caroff, J., Hourmant, A., Videau, C. and Rapt, F.: Quantitative analysis of polyamines at trace levels by high performance liquid chromatography in high salt solutions. Application to seawater. *J. Liq. Chromatogr.*, 17 (1994) 1413-1417.
- 3225 Danner, R.M., Reddy, T.V. and Guion, C.W.: A sensitive reversed-phase ion-pairing HPLC method for the determination of polyamines in rat liver. *LC-GC*, 12 (1994) 244-248; *C.A.*, 120 (1994) 211719a.
- 3226 Duchateau, A.L.L., Guns, J.J., Kubben, R.G.R. and van Tilburg, A.F.P.: High-performance liquid chromatography of diamine enantiomers as Schiff bases on a chiral stationary phase. *J. Chromatogr. A*, 664 (1994) 169-176.
- 3227 Hallak, A., Rosenberg, R., Gilat, T. and Semjen, G.J.: Determination of free polyamines in human bile by high-performance liquid chromatography. *Clin. Sci.*, 85 (1993) 451-454; *C.A.*, 120 (1994) 186401f.
- 3228 Hiramatsu, K., Kamei, S., Sugimoto, M., Kinoshita, K., Iwasaki, K. and Kawakita, M.: An improved method of determining free and acetylated polyamines by HPLC involving an enzyme reactor and an electrochemical detector. *J. Biochem. (Tokyo)*, 115 (1994) 584-589.
- 3229 Jandera, P., Ventura, K., Hladoniková, R. and Churáček, J.: Comparison of various sorbents for the enrichment of samples of aliphatic amines using solid-phase extraction prior to the determination by HPLC with fluorimetric detection. *J. Liq. Chromatogr.*, 17 (1994) 69-95.
- 3230 Juorio, A.V. and Sloley, B.D.: Determination of trace amines and related compounds by high performance liquid chromatography with electrochemical or fluorometric detection. *IBRO Handb. Ser.*, 15(High Performance Liquid Chromatography in Neuroscience Research) (1993) 217-242; *C.A.*, 120 (1994) 157692w - a review with 75 refs.
- 3231 Kawase, H., Hasegawa, T., Hashimoto, H., Noto, T. and Nakajima, T.: A study on the metabolism of spermidine in mammals: purification and identification of a newly identified metabolite, 2-oxo-1-pyrrolidinepropionic acid, in rat urine. *J. Biochem. (Tokyo)*, 115 (1994) 356-361.
- 3232 Martelli, A., Arlorio, M. and Tourn, M.L.: Determination of amines and precursor amino acids in Gorgonzola cheese by ion-pair HPLC. *Riv. Aliment.*, 22 (1993) 261-270; *C.A.*, 120 (1994) 75760c.
- 3233 Pawlowska, M., Zukowski, J. and Armstrong, D.W.: Sensitive enantiomeric separation of aliphatic and aromatic amines using aromatic anhydrides as non-chiral derivatizing agents. *J. Chromatogr. A*, 666 (1994) 485-491.
- 3234 Pettersson, C. and Gioeli, C.: Chiral separation of amines using reversed-phased ion-pair chromatography. *Chirality*, 5 (1993) 241-245; *C.A.*, 120 (1994) 207658n.
- 3235 Rissler, K.: Separation of acetylated polypropylene glycol di- and triamines by gradient reversed-phase high-performance liquid chromatography and evaporative light scattering detection. *J. Chromatogr. A*, 667 (1994) 167-174.
- 3236 Sato, K., Nakano, H. and Hobo, T.: Effect of sample derivatization in liquid chromatographic separations of amine and amino acid enantiomers using diamide-type stationary phases. *J. Chromatogr. A*, 666 (1994) 463-470.
- 3237 Yuan, Z., Egorin, M.J., Rosen, D.M., Simon, M.A. and Callery, P.S.: Cellular pharmacology of N¹- and N⁸-aziridinyl analogues of spermidine. *Cancer Res.*, 54 (1994) 742-748.
- For additional information see *C.A.*:
120 (1994) 100888k, 157791c, 211708w.
- See also 2898, 3245.
- 17b. *Catecholamines and their metabolites*
- 3238 Acworth, I.N., Yu, J., Ryan, E., Garipey, K.C., Gamache, P., Hull, K. and Maher, T.: Simultaneous measurement of monoamine, amino acid, and drug levels, using high performance liquid chromatography and coulometric array technology: application to *in vivo* microdialysis perfusate analysis. *J. Liq. Chromatogr.*, 17 (1994) 685-705.
- 3239 Anderson, G.M.: High performance liquid chromatographic analysis of monoamines and metabolites in body fluids. *IBRO Handb. Ser.*, 15(High Performance Liquid Chromatography in Neuroscience Research) (1993) 57-88; *C.A.*, 120 (1994) 186332j - a review with many refs.
- 3240 Betto, P., Ricciarello, G., Ruggeri, S., Stocchi, F., Baronti, F., Nappi, G., Martignoni, E., Melzi D'Eril, G.V., Blandini, F. and Lucarelli, C.: HPLC monitoring of plasma catechol levels in the characterization of Parkinson's disease. *G. Ital. Chim. Clin.*, 17 (1992) 441-453; *C.A.*, 120 (1994) 46079g.
- 3241 Cheng, F.-C., Kuo, J.-S., Chia, L.-G., Tsai, T.-H. and Chen, C.-F.: Rapid measurement of the monoamine content in small volumes of rat plasma. *J. Chromatogr. B*, 654 (1994) 177-183.
- 3242 Duda, C.T. and Kissinger, P.T.: Determination of biogenic amines, their metabolites, and other neurochemicals by liquid chromatography/electrochemistry. *Tech. Behav. Neural Sci.*, 11(Methods in Neurotransmitter and Neuropeptide Research, Pt. 1) (1993) 41-82; *C.A.*, 120 (1994) 236224y - a review with 157 refs.
- 3243 Freeman, K., Lin, P., Lin, L. and Blank, C.L.: Monoamines and metabolites in the brain. *IBRO Handb. Ser.*, 15(High Performance Liquid Chromatography in Neuroscience Research) (1993) 27-55; *C.A.*, 120 (1994) 128679s - a review with 39 refs.

- 3244 Gerlo, E.A.M. and Sevens, C.: Urinary and plasma catecholamines and urinary catecholamine metabolites in pheochromocytoma: diagnostic value in 19 cases. *Clin. Chem. (Washington)*, 40 (1994) 250-256.
- 3245 Ikarashi, Y., LeRoy Blank, C., Kawakubo, T., Suda, Y. and Maruyama, Y.: Competitive binding of catecholamines, indoleamines, acetylcholine, and related metabolites to various glassy carbon materials. *J. Liq. Chromatogr.*, 17 (1994) 287-301.
- 3246 Kafil, J.B. and Dhingra, B.S.: Stability-indicating method for the determination of levodopa, levodopa-carbidopa and related impurities. *J. Chromatogr. A*, 667 (1994) 175-181.
- 3247 Kubo, H., Umiguchi, Y. and Kinoshita, T.: In-line derivatization method for fluorimetric determination of catecholamines by high performance liquid chromatography. *Chromatographia*, 38 (1994) 591-594.
- 3248 Kuruma, K., Hirai, E., Uchida, K., Kikuchi, J. and Terui, Y.: Quantification of histamine by postcolumn fluorescence detection high-performance liquid chromatography using orthophthalaldehyde in tetrahydrofuran and reaction mechanism. *Anal. Sci.*, 10 (1994) 259-265; *C.A.*, 120 (1994) 264971h.
- 3249 Nappi, A.J. and Vass, E.: Chromatographic analyses of the effects of glutathione, cysteine and ascorbic acid on the monophenol and diphenol oxidase activities of tyrosinase. *J. Liq. Chromatogr.*, 17 (1994) 793-815.
- 3250 Saito, K., Horie, M. and Nakazawa, H.: Determination of urinary excretion of histamine and 1-methylhistamine by liquid chromatography. *J. Chromatogr. B*, 654 (1994) 270-275.
- 3251 Saito, K., Yamada, F., Horie, M. and Nakazawa, H.: Determination of histamine and 1-methylhistamine by liquid chromatography using on-column derivatization and a column-switching technique. *Anal. Sci.*, 9 (1993) 803-806; *C.A.*, 120 (1994) 46099p.
- 3252 Sanchis Malloles, J.M., Villanueva Camanas, R.M. and Ramis-Ramos, G.: Determination of catecholamines as aminochromes by micellar liquid chromatography with thermal lens spectrophotometric detection. *Chromatographia*, 38 (1994) 365-372.
- 3253 Straub, B., Schollenberger, M., Kicherer, M., Luckas, B. and Hammes, W.P.: Extraction and determination of biogenic amines in fermented sausages and other meat products with reversed-phase HPLC. *Z. Lebensm.-Unters. Forsch.*, 197 (1993) 230-232; *C.A.*, 120 (1994) 132503x.
- 3254 Volin, P.: Determination of free urinary catecholamines by high-performance liquid chromatography with electrochemical detection. *J. Chromatogr. B*, 655 (1994) 121-126.
- 3255 Wakamatsu, K. and Ito, S.: Improved HPLC determination of 5-S-cysteinyl-dopa in serum. *Clin. Chem. (Washington)*, 40 (1994) 495-496.

For additional information see *C.A.*:
120 (1994) 153904u.

See also 3148, 3230.

17c. Urea and guanidine derivatives

For additional information see *C.A.*:
120 (1994) 49200a.

- 17d. *Other amine derivatives and amides (excl. peptides)*
- 3256 Bazylak, G.: Chiral separation of some amino alcohols by addition of helical nickel(II) chelate to the mobile phase used in reversed-phase high-performance liquid chromatography. *J. Chromatogr. A*, 665 (1994) 75-86.
- 3257 Hansson, G. and Ahnoff, M.: Chromatographic separation of amide diastereomers: correlation with molecular descriptors. *J. Chromatogr. A*, 666 (1994) 505-517.
- 3258 Kabasawa, Y. and Takeba, K.: Purification of 2,2'-dihydroxy-3,3',5,5',6-pentachlorobenzanilide by counter alternative current chromatography. *Anal. Sci.*, 9 (1993) 875-876; *C.A.*, 120 (1994) 191260p.
- 3259 Prokai, A.M. and Ravichandran, R.K.: Simultaneous analysis of hydroxylamine, N-methylhydroxylamine and N,N-dimethylhydroxylamine by ion chromatography. *J. Chromatogr. A*, 667 (1994) 298-303.

See also 2997, 3226, 3235.

18. AMINO ACIDS AND PEPTIDES; CHEMICAL STRUCTURE OF PROTEINS

18a. Amino acids and their derivatives

- 3260 Abe, Y., Ueda, T. and Imoto, T.: Reduction of disulfide bonds in proteins by 2-aminothiophenol under weakly acidic conditions. *J. Biochem. (Tokyo)*, 115 (1994) 52-57.
- 3261 Acil, Y. and Müller, P.K.: Rapid method for the isolation of the mature collagen cross-links, hydroxylysylpyridinoline and lysylpyridinoline. *J. Chromatogr. A*, 664 (1994) 183-188.
- 3262 Boni, R.L., Simpson, J.T., Naritsin, D.B., Saito, K. and Markey, S.P.: Quantification of L-tryptophan and L-kynurenine by liquid chromatography/electron capture negative ion chemical ionization mass spectrometry. *Biol. Mass Spectrom.*, 23 (1994) 27-32; *C.A.*, 120 (1994) 72872e.
- 3263 Bruckner, H., Haasmann, S., Langer, M., Westhauser, T., Wittner, R. and Godel, H.: Liquid chromatographic determination of D- and L-amino acids by derivatization with o-phthalaldehyde and chiral thiols. Applications with reference to biosciences. *J. Chromatogr. A*, 666 (1994) 259-273.
- 3264 Burgoyne, R.F.: Determining amino acid composition. *BioTechnology*, 11 (1993) 1302-1304; *C.A.*, 120 (1994) 100902k.
- 3265 Calabresi, E., Lasagni, L., Franceschelli, F., Bartolini, L., Serio, M. and Brandi, M.L.: Use of an internal standard to measure pyridinoline and deoxypyridinoline in urine. *Clin. Chem. (Washington)*, 40 (1994) 336-337.
- 3266 Di Pietra, A.M., Gotti, R., Bonazzi, D., Andrisano, V. and Cavrini, V.: HPLC determination of glutathione and L-cysteine in pharmaceuticals after derivatization with ethacrynic acid. *J. Pharm. Biomed. Anal.*, 12 (1994) 91-98.
- 3267 Egorova, T.A., Eremin, S.V., Mitaner, B.I., Zvonkova, E.N. and Shvets, V.I.: (Method for preparative separation of benzyloxy-carbonyl amino acids derivatives from protein hydrolyzates). *Biotechnologiya*, (1993) 32-35; *C.A.*, 120 (1994) 49247w.

- 3268 Galli, B., Gasparri, F., Misiti, D., Villani, C., Corradini, R., Dosena, A. and Marchelli, R.: Enantiomeric separation of dansyl- and dansylamino acids by ligand-exchange chromatography with (S)- and (R)-phenylalaninamide-modified silica gel. *J. Chromatogr. A*, 666 (1994) 77-89.
- 3269 Jenssen, T., Nurjhan, N., Perriello, G., Bucci, A., Toft, I. and Gerich, J.: Determination of [¹⁴C] glutamine specific activity in plasma. *J. Liq. Chromatogr.*, 17 (1994) 1337-1348.
- 3270 Kamata, K., Takahashi, M., Terasima, K. and Nishijima, M.: Liquid chromatographic determination of carnitine by precolumn derivatization with pyrene-1-carbonyl cyanide. *J. Chromatogr. A*, 667 (1994) 113-118.
- 3271 Kamimori, H., Hamashima, Y. and Konishi, M.: Determination of carnitine and saturated-acyl group carnitines in human urine by high-performance liquid chromatography with fluorescence detection. *Anal. Biochem.*, 218 (1994) 417-424.
- 3272 Kersten, B.S.: HPLC chiral optimization of a unique β -amino acid and its ester. *J. Liq. Chromatogr.*, 17 (1994) 33-48.
- 3273 Kilpatrick, I.C., Waller, S.J., Harrison, P.S. and Evans, R.H.: Rapid, automated high performance liquid chromatographic analyses of excitatory sulfur-containing and some other neurotoxic amino acids. *IBRO Handb. Ser.*, 15 (High Performance Liquid Chromatography in Neuroscience Research) (1993) 139-162; *C.A.*, 120 (1994) 211701p.
- 3274 Landry, J. and Delhaye, S.: HPLC determination of tryptophan in foodstuffs using barium hydroxide hydrolysis at 140° C. *Amino Acids*, 5 (1993) 317-321; *C.A.*, 120 (1994) 161873y.
- 3275 Miyajima, K., Hirata, M., Yoshida, T., Kosaka, H. and Okayama, A.: Study on measurement of δ -aminolevulinic acid in plasma by high-performance liquid chromatography. *J. Chromatogr. B*, 654 (1994) 165-169.
- 3276 Morleo, M.A. and Musi, E.: Evaluation of a new product for hydroxyproline determination in HPLC. *Boll. Chim. Farm.*, 132 (1993) 281-284; *C.A.*, 120 (1994) 264951b.
- 3277 Pech, L.L. and Nelson, D.L.: Purification and characterization of calmodulin (lysine 115) N-methyltransferase from *Paramecium tetraurelia*. *Biochim. Biophys. Acta*, 1199 (1994) 183-194.
- 3278 Peel, N., Al-Dehaimi, A., Colwell, A., Russell, G. and Eastell, R.: Sulfasalazine may interfere with HPLC assay of urinary pyridinium crosslinks. *Clin. Chem. (Washington)*, 40 (1994) 167-168.
- 3279 Pendergrass, J.A., Jr., Clark, E.P. and Palazzolo, D.L.: Diethylenetriaminepentaacetic acid added to the mobile phase improves GSH peak symmetry. *Curr. Sep.*, 12 (1993) 137-140; *C.A.*, 120 (1994) 186405k.
- 3280 Potin-Gautier, M., Boucharat, C., Astruc, A. and Astruc, M.: Speciation of selenoamino acids by online HPLC ETAA spectrometry. *Appl. Organomet. Chem.*, 7 (1993) 593-598; *C.A.*, 120 (1994) 207686v.
- 3281 Ranfft, K. and Faure, U.: A comparison of methods for the determination of tryptophan in feeds. *Agribiol. Res.*, 46 (1993) 65-72; *C.A.*, 120 (1994) 52946d.
- 3282 Sebesta, I., Fairbanks, L.D., Davies, P.M., Simmonds, H.A. and Leonard, J.V.: The allopurinol loading test for identification of carriers for ornithine carbamoyl transferase deficiency: studies in a healthy control population and females at risk. *Clin. Chim. Acta*, 224 (1994) 45-54.
- 3283 Spondlin, C. and Küsters, E.: Direct determination of enantiomeric purity of FMOC amino acids with high performance liquid chromatography. *Chromatographia*, 38 (1994) 325-329.
- 3284 Stocchi, V., Palma, F., Piccoli, G., Biagiarelli, B., Cucchiari, L. and Magnani, M.: HPLC analysis of taurine in human plasma sample using the DABS-Cl reagent with sensitivity at picomole levels. *J. Liq. Chromatogr.*, 17 (1994) 347-357.
- 3285 Strydom, D.J. and Cohen, S.A.: Sensitive analysis of cystine/cysteine using 6-aminoquinolyl-N-hydroxysuccinimidyl carbamate (AQC) derivatives. In: Angeletti, R.H. (Editor), *Tech. Protein Chem. IV, [Pap. Protein Soc. Symp.]*, 6th, 1992, Academic, San Diego, 1993, pp. 299-306; *C.A.*, 120 (1994) 49194b.
- 3286 Teerlink, T., van Leeuwen, P.A.M. and Houdijk, A.: Plasma amino acids determined by liquid chromatography within 17 minutes. *Clin. Chem. (Washington)*, 40 (1994) 245-249.
- 3287 Van Gennip, A.H., Driedijk, P.C., Elzinga, A. and Abeling, N.G.G.M.: Screening for defects of dihydropyrimidine degradation by analysis of amino acids in urine before and after acid hydrolysis. *J. Inherited Metab. Dis.*, 15 (1992) 413-415; *C.A.*, 120 (1994) 211700n.
- 3288 Woo, K.-L. and Lee, S.-H.: Determination of protein amino acids as butylthiocarbonyl derivatives by reversed-phase high-performance liquid chromatography with precolumn derivatization. *J. Chromatogr. A*, 667 (1994) 105-111.
- 3289 Zhu, S. and Zhao, S.: (High performance liquid chromatographic analysis of amino acids). *Sepu*, 12 (1994) 202-204; *C.A.*, 120 (1994) 182064a - a review with 53 refs.

For additional information see C.A.:

120 (1994) 49245u, 135129r, 157892w, 158133b, 239339a.

See also 2854, 2904, 3074, 3150, 3161, 3232, 3236, 3238, 3246, 3249, 3290, 3432, 3454, 3540, 3782.

18b. *Peptides, peptidic and proteinous hormones, growth factors*

- 3290 Asensi, M., Sastre, J., Pallardó, F.V., de la Asunción, J.G., Estrella, J.M. and Vina, J.: A high-performance liquid chromatography method for measurement of oxidized glutathione in biological samples. *Anal. Biochem.*, 217 (1994) 323-328.
- 3291 Bauer, K. and Schulz, M.: Biosynthesis of carnosine and related peptides by skeletal muscle cells in primary structure. *Eur. J. Biochem.*, 219 (1994) 43-47.
- 3292 Boyes, B.E. and Kirkland, J.J.: Rapid, high-resolution HPLC separation of peptides using small particles at elevated temperatures. *Pept. Res.*, 6 (1993) 249-258; *C.A.*, 120 (1994) 186400e.
- 3293 Brattsand, G., Marklund, U., Nylander, K., Ross, G. and Gullberg, M.: Cell-cycle-regulated phosphorylation of oncoprotein 18 on Ser 16, Ser 25 and Ser 38. *Eur. J. Biochem.*, 220 (1994) 359-368.
- 3294 Brooks, C.L., Isaacs, L.A. and Wicks, J.R.: Preparative purification of phosphorylated and nonphosphorylated bovine prolactins. *Mol. Cell. Endocrinol.*, 99 (1994) 301-305; *C.A.*, 120 (1994) 236314c.
- 3295 Cardamone, M., Ashman, K., Brandon, M.R. and Puri, N.K.: Identification of cysteine-containing peptides during the peptide mapping of recombinant proteins. *Pept. Res.*, 6 (1993) 242-248; *C.A.*, 120 (1994) 186399m.
- 3296 Chang, C.H., Chey, W.Y., Sun, Q., Leiter, A. and Chang, T.-m.: Characterization of the release of cholecystokinin from a murine neuroendocrine tumor cell line, STC-1. *Biochim. Biophys. Acta*, 1221 (1994) 339-347.

- 3297 Chen, S., Pawlowska, M. and Armstrong, D.W.: HPLC enantio-separation of di- and tripeptides on cyclodextrin bonded stationary phases after derivatization with 6-aminoquinolyl-N-hydroxysuccinimidyl carbamate (AQC). *J. Liq. Chromatogr.*, 17 (1994) 483-497.
- 3298 Congote, L.F. and Li, Q.: Accurate processing and secretion in the baculovirus expression system of an erythroid-cell-stimulating factor consisting of a chimaera of insulin-like growth factor II and an insect insulin-like peptide. *Biochem. J.*, 299 (1994) 101-107.
- 3299 Contreras Martinez, S. and Vera-Avila, L.E.: Effect of mobile phase composition on the separation of thyrotropin-releasing hormone and some metabolites by reversed-phase ion-pair chromatography. *J. Chromatogr. A*, 667 (1994) 119-124.
- 3300 Cummings, J., MacLellan, A., Langdon, S.P., Rozengurt, E. and Smyth, J.F.: Determination of two neuropeptide growth factor antagonists, [D-Arg¹,D-Phe⁵,D-Trp^{7,9},Leu¹¹]-substance P and [Arg⁶,D-Trp^{7,9},N-MePhe⁸]-substance P(6-11), by high-performance liquid chromatography with electrochemical detection. *J. Chromatogr. B*, 653 (1994) 195-203.
- 3301 Dave, K.J.: Liquid chromatographic methods for the analysis of opioid peptides. Avail. *Univ. Microfilms Int.*, Order No. DA9210039, 1991, 270 pp.; C.A., 120 (1994) 94605x.
- 3302 Ding, J., Burkhart, W. and Kassel, D.R.: Identification of phosphorylated peptides from complex mixtures using negative-ion orifice-potential stepping and capillary liquid chromatography/electrospray ionization mass spectrometry. *Rapid Commun. Mass Spectrom.*, 8 (1994) 94-98; C.A., 120 (1994) 158102r.
- 3303 Evans, C.J., Maidment, N.T. and Newcomb, R.: Reverse phase high performance liquid chromatography of biological peptides: technical aspects and applications. *IBRO Handb. Ser.*, 15(High Performance Liquid Chromatography in Neuroscience Research) (1993) 265-296; C.A., 120 (1994) 128680k - a review with 42 refs.
- 3304 Funakoshi, S., Fukuda, H. and Fujii, N.: The affinity purification procedure for peptide synthesized by the solid-phase technique. In: Yanaihara, N. (Editor), *Pept. Chem. 1992, Proc. Jpn. Symp.*, 2nd, 1992, ESCOM, Leiden, 1993, pp. 99-102; C.A., 120 (1994) 54963f.
- 3305 Gray, W.R.: Disulfide structures of highly bridged peptides: A new strategy for analysis. *Protein Sci.*, 2 (1993) 1732-1748; C.A., 120 (1994) 239584b.
- 3306 Griffiths, E.C.: Application of high performance liquid chromatography in studies of the metabolism of neuropeptides. *IBRO Handb. Ser.*, 15(High Performance Liquid Chromatography in Neuroscience Research) (1993) 297-318; C.A., 120 (1994) 46050r - a review with 17 refs.
- 3307 Haniu, M., Hsieh, P., Rohde, M.F. and Kenney, W.C.: Characterization of disulfide linkages in platelet-derived growth factor AA. *Arch. Biochem. Biophys.*, 310 (1994) 433-439.
- 3308 Harada, K., Nagai, H., Kimura, Y., Suzuki, M., Park, H.D., Watanabe, M.F., Luukkainen, R., Sivonen, K. and Carmichael, W.W.: Liquid chromatography/mass spectrometric detection of anatoxin-a, a neurotoxin from cyanobacteria. *Tetrahedron*, 49 (1993) 9251-9260; C.A., 120 (1994) 70983t.
- 3309 Hayakawa, Y.: Inclusion of a β -aspartyl peptide bond causes the loss of inherent physiological activities. *J. Biochem. (Tokyo)*, 115 (1994) 15-17.
- 3310 Hendrickson, T.L. and Wilson, G.S.: Improved clean-up method for the enkephalins in plasma using immunoaffinity chromatography. *J. Chromatogr. B*, 653 (1994) 147-154.
- 3311 Hosoi, K., Tsunasawa, S., Kurihara, K., Aoyama, H., Ueha, T., Murai, T. and Sakiyama, F.: Identification of mK1, a true tissue (glandular) kallikrein of mouse submandibular gland: tissue distribution and a comparison of kinin-releasing activity with other submandibular kallikreins. *J. Biochem. (Tokyo)*, 115 (1994) 137-143.
- 3312 Irminger, J.C., Vollenweider, F.M., Neerman-Arbez, M. and Halban, P.A.: Human proinsulin conversion in the regulated and the constitutive pathways of transfected AtT20 cells. *J. Biol. Chem.*, 269 (1994) 1756-1762.
- 3313 James, S., Gibbs, B.F., Toney, K. and Bennett, H.P.J.: Purification of antimicrobial peptides from an extract of the skin of *Xenopus laevis* using heparin-affinity HPLC: characterization by ion-spray mass spectrometry. *Anal. Biochem.*, 217 (1994) 84-90.
- 3314 Janini, G.M., Issaq, H.J. and Lukszo, J.: Determination of the purity of synthetic peptides by capillary electrophoresis, high performance liquid chromatography, and laser desorption mass spectrometry. *J. High Resolut. Chromatogr.*, 17 (1994) 102-103.
- 3315 Jespersen, A.M., Christensen, T., Klausen, N.K., Nielsen, P.F. and Sorensen, H.H.: Characterisation of a trisulphide derivative of biosynthetic human growth hormone produced in *Escherichia coli*. *Eur. J. Biochem.*, 219 (1994) 365-373.
- 3316 Kojima, K., Suzuki, Y., Parvez, H., Parvez, S. and Nagatsu, T.: Analysis of neuropeptides by high-performance liquid chromatography. *Tech. Behav. Neural Sci.*, 11(Methods in Neurotransmitter and Neuropeptide Research, Pt. 2) (1993) 305-359; C.A., 120 (1994) 183055k.
- 3317 Kubo, H.: Coulometric detection of peptides by reversed-phase liquid chromatography with a solid-phase reactor containing copper metal. *Anal. Chim. Acta*, 287 (1994) 211-214.
- 3318 Kukucka, M.A. and Misra, H.P.: Determination of oxytocin in biological samples by isocratic high-performance liquid chromatography with coulometric detection using C₁₈ solid-phase extraction and polyclonal antibody-based immunoaffinity column purification. *J. Chromatogr. B*, 653 (1994) 139-145.
- 3319 Lee, W.-J. and Brey, P.T.: Isolation and identification of cecropin antibacterial peptides from the extracellular matrix of the insect integument. *Anal. Biochem.*, 217 (1994) 231-235.
- 3320 Li, H. and Wang, E.: Determination of reduced glutathione in human plasma using liquid chromatography with electrochemical detection with a glassy carbon electrode chemically modified with indium ferricyanide. *Microchem. J.*, 49 (1994) 91-98; C.A., 120 (1994) 157814n.
- 3321 Maule, A.G., Shaw, C., Bowman, J.W., Thompson, D.P., Geary, T.G. and Thim, L.: KSAYMRFamide: a novel FMRFamide-related heptapeptide from the free-living nematode, *Panagrellus redivivus*, which is myoactive in the parasitic nematode, *Ascaris suum*. *Biochem. Biophys. Res. Commun.*, 200 (1994) 973-980.
- 3322 Mehvar, R.: Preparative and analytical separation of insulin-dextran conjugates from native insulin: application to preparation and characterization of insulin-dextran conjugates. *Drug. Dev. Ind. Pharm.*, 20 (1994) 395-404; C.A., 120 (1994) 200281a.

- 3323 Ng, A.S., Lewis, S.D. and Shafer, J.A.: Quantifying thrombin-catalyzed release of fibrinopeptides from fibrinogen using high-performance liquid chromatography. *Methods Enzymol.*, 222(Proteolytic Enzymes in Coagulation, Fibrinolysis, and Complement Activation, Pt. A) (1993) 341-358; *C.A.*, 120 (1994) 100023z.
- 3324 Ng, T.B. and Lo, L.L.H.: Study on the partial purification of gonadotropins and thyrotropin from deer (*Cervus elaphus*) pituitaries. *Biochem. Mol. Biol. Int.*, 31 (1993) 1023-1030; *C.A.*, 120 (1994) 208958x.
- 3325 Nuwaysir, L.M. and Stults, J.T.: Electrospray ionization mass spectrometry of phosphopeptides isolated by online immobilized metal-ion affinity chromatography. *J. Am. Soc. Mass Spectrom.*, 4 (1993) 662-669; *C.A.*, 120 (1994) 128827p.
- 3326 Ohta, T., Horimai, T., Kaburaki, Y. and Takitani, S.: Improvement in selectivity of anhydrotypsin-immobilized diol silica column by the use of eluent containing calcium ion. *J. Liq. Chromatogr.*, 17 (1994) 1281-1290.
- 3327 Paranandi, M.V., Guzzetta, A.W., Hancock, W.S. and Aswad, D.W.: Deamidation and isoaspartate formation during *in vitro* aging of recombinant tissue plasminogen activator. *J. Biol. Chem.*, 269 (1994) 243-253.
- 3328 Ramage, R., Green J., Muir, T.W., Ogunjobi, O.M., Love, S. and Shaw, K.: Synthetic, structural and biological studies of the ubiquitin system: the total chemical synthesis of ubiquitin. *Biochem. J.*, 299 (1994) 1'51-158.
- 3329 Randhawa, Z.I., Witkowska, H.E., Cone, J., Wilkins, J.A., Hughes, P., Yamanishi, K., Yasuda, S., Masui, Y., Arthur, P., Kletke, C., Bitsch, F. et al.: Incorporation of norleucine at methionine positions in recombinant human macrophage colony stimulating factor (M-CSF, 4-153) expressed in *Escherichia coli*: structural analysis. *Biochemistry*, 33 (1994) 4352-4362.
- 3330 Reynolds, E.C., Riley, P.F. and Adamson, N.J.: A selective precipitation purification procedure for multiple phosphoserine-containing peptides and methods for their identification. *Anal. Biochem.*, 217 (1994) 277-284.
- 3331 Salzet, M., Bulet, P., Watzet, C. and Malecha, J.: FMRFamide-related peptides in the sex segmental ganglia of the Pharyngobdellid leech *Erpobdella octoculata*. Identification and involvement in the control of hydric balance. *Eur. J. Biochem.*, 221 (1994) 269-275.
- 3332 Sandberg, M., Li, X., Folestad, S., Weber, S.G. and Orwar, O.: Liquid chromatographic determination of acidic β -aspartyl and γ -glutamyl peptides in extracts of rat brain. *Anal. Biochem.*, 217 (1994) 48-61.
- 3333 Söllner, C., Mentele, R., Eckerskorn, C., Fritz, H. and Sommerhoff, C.P.: Isolation and characterization of hirustasin, an anti-stasin-type serine proteinase inhibitor from the medical leech *Hirudo medicinalis*. *Eur. J. Biochem.*, 219 (1994) 937-943.
- 3334 Sýkora, D., Vinš, I., Hermann, P. and Kesner, F.: Reversed-phase high-performance liquid chromatography of diastereomers of some phosphonodipeptides. *J. Chromatogr. A*, 665 (1994) 59-65.
- 3335 Thompson, S.A., Higashiyama, S., Wood, K., Pollitt, N.S., Damm, D., McEnroe, G., Garrick, B., Ashton, N., Lau, K., Hancock, N., Klagsbrun, M. and Abraham, J.A.: Characteriation of sequences within heparin-binding EGF-like growth factor that mediate interaction with heparin. *J. Biol. Chem.*, 269 (1994) 2541-2549.
- 3336 Till, J.H., Annan, R.S., Carr, S.A. and Miller, W.T.: Use of synthetic peptide libraries and phosphopeptide-selective mass spectrometry to probe protein kinase substrate specificity. *J. Biol. Chem.*, 269 (1994) 7423-7428.
- 3337 Van der Meer, J.R., Rollema, H.S., Siezen, R.J., Beerthuyzen, M.M., Kuipers, O.P. and de Vos, W.M.: Influence of amino acid substitutions in the nisin leader peptide on biosynthesis and secretion of nisin by *Lactococcus lactis*. *J. Biol. Chem.*, 269 (1994) 3555-3562.
- 3338 Vincent, B., Vincent, J.-P. and Checler, F.: Neurotensin and neuromedin N undergo distinct catabolic processes in murine astrocytes and primary cultured neurons. *Eur. J. Biochem.*, 221 (1994) 297-306.
- 3339 Yadav, S.P., Brew, K., Majercik, M.H. and Puett, D.: Label selection approach to assess the role of individual amino groups in human chorionadotropin receptor binding. *J. Biol. Chem.*, 269 (1994) 3991-3998.

For additional information see C.A.:

120 (1994) 72815p, 192316y, 265314h.

See also 3388, 3400, 3472, 3497, 3554, 3714.

18c. Elucidation of structure of proteins and enzymes

- 3340 Degerman, E., Moos, M., Jr., Rascón, A., Vasta, V., Meacci, E., Smith, C.J., Lindgren, S., Andersson, K.-E., Belfrage, P. and Manganiello, V.: Single-step affinity purification, partial structure and properties of human platelet cGMP inhibited cAMP phosphodiesterase. *Biochim. Biophys. Acta*, 1205 (1994) 189-198.
- 3341 Drummond, J.T. and Matthews, R.G.: Nitrous oxide inactivation of cobalamin-dependent methionine synthase from *Escherichia coli*: characterization of the damage to the enzyme and prosthetic group. *Biochemistry*, 33 (1994) 3742-3750.
- 3342 Du, G., Ng, C. and Prestwich, G.D.: Odorant binding by a pheromone binding protein: active site mapping by photoaffinity labeling. *Biochemistry*, 33 (1994) 4812-4819.
- 3343 Fischer, W.H., Behan, D.P., Park, M., Potter, E., Lowry, P.J. and Vale, W.: Assignment of disulfide bonds in corticotropin-releasing factor-binding protein. *J. Biol. Chem.*, 269 (1994) 4313-4316.
- 3344 Gao, G. and Fonda, M.L.: Identification of an essential cysteine residue in pyridoxal phosphatase from human erythrocytes. *J. Biol. Chem.*, 269 (1994) 8234-8239.
- 3345 Herold, M., Grimm, R., Heiger, D. and Winzer, L.: (Peptide mapping by reversed-phase-HPLC. Part 2. Applications in protein characterization). *BioTec (Wuerzburg)*, 4 (1993) 48-54; *C.A.*, 120 (1994) 72635e.
- 3346 Hess, D., Covey, T.C., Winz, R., Brownsey, R.W. and Aebersold, R.: Analytical and micropreparative peptide mapping by high performance liquid chromatography/electrospray mass spectrometry of proteins purified by gel electrophoresis. *Protein Sci.*, 2 (1993) 1342-1351; *C.A.*, 120 (1994) 128984n.
- 3347 Hilliard IV, G.M., Cook, R.G., Weigel, N.L. and Pike, J.W.: 1,25-Dihydroxyvitamin D₃ modulates phosphorylation of serine 205 in the human vitamin D receptor: site-directed mutagenesis of this residue promotes alternative phosphorylation. *Biochemistry*, 33 (1994) 4300-4311.

- 3348 Jayaram, B. and Haley, B.E.: Identification of peptides within the base binding domains of the GTP- and ATP-specific binding sites of tubulin. *J. Biol. Chem.*, 269 (1994) 3233-3242.
- 3349 Kajihara, J.-i., Shibata, K., Nakano, Y., Nishimuro, S. and Kato, K.: Physicochemical characterization of PEG-PPG conjugated human urokinase. *Biochim. Biophys. Acta*, 1199 (1994) 202-208.
- 3350 Kamei-Hayashi, K., Oshino, R. and Hara, S.: Amino acid sequence of glutathione S-transferase α from guinea pig liver. *J. Biochem. (Tokyo)*, 114 (1993) 835-841.
- 3351 Knaus, H.-G., Garcia-Calvo, M., Kaczorowski, G.J. and Garcia, M.L.: Subunit composition of the high conductance calcium-activated potassium channel from smooth muscle, a representative of the mSlo and slowpoke family of potassium channels. *J. Biol. Chem.*, 269 (1994) 3921-3924.
- 3352 Konami, Y., Ishida, C., Yamamoto, K., Osawa, T. and Irimura, T.: A unique amino acid sequence involved in the putative carbohydrate-binding domain of a legume lectin specific for sialylated carbohydrate chains: primary sequence determination of *Maackia amurensis* hemagglutinin (MAH). *J. Biochem. (Tokyo)*, 115 (1994) 767-777.
- 3353 Malyan, A.N., Girault, G. and Berger, G.: Examination of catalytic activity of the β subunit isolated from chloroplast coupling factor 1. *Biochim. Biophys. Acta*, 1184 (1994) 202-206.
- 3354 McClelland, P., Adam, L.P. and Hathaway, D.R.: Identification of a latent Ca^{2+} /calmodulin dependent protein kinase II phosphorylation site in vascular calpain II. *J. Biochem. (Tokyo)*, 115 (1994) 41-46.
- 3355 Miyagawa, M., Fukuhara, K.-i., Katayanagi, K., Ishimaru, K., Morikawa, K., Matsuzaki, T., Sato, Y., Hara, S., Ikenaka, T., Arai, M. and Murao, S.: Crystallization and preliminary crystallographic data of the α -amylase inhibitors, Haim I and Paim I. *J. Biochem. (Tokyo)*, 115 (1994) 168-170.
- 3356 Moritz, R.L. and Simpson, R.J.: Purification of proteins and peptides for sequence analysis using microcolumn liquid chromatography. *J. Microcolumn Sep.*, 4 (1993) 485-489; *C.A.*, 120 (1994) 128765s.
- 3357 Nielsen, S.U., Rump, R., Højrup, P., Roepstorff, P. and Spener, F.: Differentiation regulation and phosphorylation of the fatty acid-binding protein from rat mammary epithelial cells. *Biochim. Biophys. Acta*, 1211 (1994) 189-197.
- 3358 Nishita, K., Tanaka, H. and Ojima, T.: Amino acid sequence of troponin C from scallop striated adductor muscle. *J. Biol. Chem.*, 269 (1994) 3464-3468.
- 3359 Odoul, M., Bagrel, D., Peyrieras, N., Bodaud, H. and Siest, G.: Glycosylation of gamma-glutamyltransferase is modified by ethanol in H5-6 hepatoma cell line. *Clin. Chim. Acta*, 225 (1994) 1-15.
- 3360 Oyama, F., Hashino, K., Oyama, R., Kato, I. and Titani, K.: Improved method for expression of Kunitz-type serine proteinase inhibitor domain of β -amyloid protein precursor in *Escherichia coli* and characterization of disulfide bonds of the product. *J. Biochem. (Tokyo)*, 114 (1993) 813-819.
- 3361 Smith, D.L., Zhang, Z. and Liu, Y.: Amide hydrogen exchange and mass spectrometry: a probe of high-order structure in proteins. *Pure Appl. Chem.*, 66 (1994) 89-94; *C.A.*, 120 (1994) 265074e.
- 3362 Strömqvist, M.: Peptide mapping using combinations of size-exclusion chromatography, reversed-phase chromatography and capillary electrophoresis. *J. Chromatogr. A*, 667 (1994) 304-310.
- 3363 Terada, S., Katayama, H., Noda, K., Fujimura, S. and Kimoto, E.: Amino acid sequences of Kunitz family subtilisin inhibitors from seeds of *Canavalia lineata*. *J. Biochem. (Tokyo)*, 115 (1994) 397-404.
- 3364 Tran, C.M., Schneiner-Bobis, G., Schoner, W. and Farley, R.A.: Identification of an amino acid in the ATP binding site of Na^+/K^+ -ATPase after photochemical labeling with 8-azido-ATP. *Biochemistry*, 33 (1994) 4140-4147.
- 3365 Vollmer, S.H., Walner, M.B., Tarbell, K.V. and Colman, R.F.: Guanosine 5'-O-[S-(bromo-2,3-dioxobutyl)]thiophosphate and adenosine 5'-O-[S-(4-bromo-2,3-dioxobutyl)]thiophosphate. New nucleotide affinity labels which react with rabbit muscle pyruvate kinase. *J. Biol. Chem.*, 269 (1994) 8082-8090.
- 3366 Watanabe, H., Narumi, H., Inaba, T., Ohgi, K. and Irie, M.: Purification, some properties, and primary structure of a base non-specific ribonuclease from oyster (*Crassostrea gigas*). *J. Biochem. (Tokyo)*, 114 (1993) 800-807.
- 3367 Winzer, L.: (Peptide mapping by reversed-phase HPLC. Part 1. Chromatographic conditions). *BioTec (Wuerzburg)*, 4 (1992) 47-51; *C.A.*, 120 (1994) 72636f.
- 3368 Yamamoto, N., Akino, A. and Takano, T.: Purification and specificity of a cell-wall-associated proteinase from *Lactobacillus helveticus* CP790. *J. Biochem. (Tokyo)*, 114 (1993) 740-745.
- 3369 Yokota, A. and Tokai, H.: Lysine residues involved in the hysteresis and in the regulatory sites of spinach ribulose 1,5-bisphosphate carboxylase/oxygenase. *J. Biochem. (Tokyo)*, 114 (1993) 746-753.

For additional information see C.A.:
120 (1994) 186520u.

See also 3277, 3401, 3402, 3409, 3427, 3470, 3475, 3479, 3526, 3594, 3611, 3616.

19. PROTEINS

19a. General techniques

- 3370 Andreasen, P.A.: Molecular mass of proteins and their partitioning in aqueous two-phase systems. *J. Chromatogr. A*, 664 (1994) 99-103.
- 3371 Andrews, B.A. and Haywood, K.: Effect of pH, ion type and ionic strength on partitioning of proteins in reverse micelle systems. *J. Chromatogr. A*, 668 (1994) 55-60.
- 3372 Asenjo, J.A., Schmidt, A.S., Hachem, F. and Andrews, B.A.: Model for predicting the partition behaviour of proteins in aqueous two-phase systems. *J. Chromatogr. A*, 668 (1994) 47-54.
- 3373 Bergloef, J.H.: Validation aspects relating to the use of chromatographic media. *Colloq. INSERM*, 227(Biotechnology of Blood Proteins) (1993) 31-36; *C.A.*, 120 (1994) 49203d.

- 3374 Fernandez, J., Andrews, L. and Mische, S.M.: An improved procedure for enzymatic digestion of polyvinylidene difluoride-bound proteins for internal sequence analysis. *Anal. Biochem.*, 218 (1994) 112-117.
- 3375 Forciniti, D.: Protein refolding using aqueous two-phase systems. *J. Chromatogr. A*, 668 (1994) 95-100.
- 3376 Grasselli, M., Diaz, L.E. and Cascone, O.: Beaded matrixes from cross-linked alginate for affinity and ion exchange chromatography of proteins. *Biotechnol. Tech.*, 7 (1993) 707-712; *C.A.*, 120 (1994) 100895k.
- 3377 Herbst, F., Nokihara, K., Beck, R., Watanabe, T. and Truedinger, U.: Development and applications of novel stationary phases for the HPLC separation of proteins and peptides. Part 3. Derivatized titanium dioxide). In: Yanaihara, N. (Editor), *Pept. Chem. 1992, Proc. Jpn. Symp., 2nd 1992*, ESCOM, Leiden, 1993, pp. 204-207; *C.A.*, 120 (1994) 72631a.
- 3378 Huddleston, J.G., Wang, R., Flanagan, J.A., O'Brien, S. and Lyddiatt, A.: Variation of protein partition coefficients with volume ratio in poly(ethylene glycol)-salt aqueous two-phase systems. *J. Chromatogr. A*, 668 (1994) 3-11.
- 3379 Josic, D.: (HPLC of proteins: short review with practical tips). *BioTec (Wuerzburg)*, 4 (1992) 41-46; *C.A.*, 120 (1994) 72563e - a review with 14 refs.
- 3380 Kassel, D.B., Luther, M.A., Willard, D.H., Fulton, S.P. and Salzmann, J.P.: Rapid purification, separation and identification of proteins and enzyme digests using packed capillary perfusion column LC and LC/MS. In: Angeletti, R.H. (Ed.), *Tech. Protein Chem. IV, [Pap. Protein Soc. Symp.]*, 6th, 1992, Academic, San Diego, 1993, pp. 55-64; *C.A.*, 120 (1994) 49192z.
- 3381 Luo, R.G. and Hsu, J.T.: Intraparticle protein diffusion effect on gradient elution chromatography. *Sep. Technol.*, 3 (1993) 221-229; *C.A.*, 120 (1994) 211709x.
- 3382 Mhatre, R., Qian, R., Krull, I.S. and Gadam, S. and Cramer, S.M.: Characterization of proteins separated by displacement chromatography using low angle laser light scattering photometry. *Chromatographia*, 38 (1994) 349-354.
- 3383 Moseley, M.A. and Unger S.E.: Packed capillary liquid chromatography/electrospray ionization/mass spectrometry in the pharmaceutical sciences: characterization of protein mixtures. *J. Microcolumn Sep.*, 4 (1992) 393-398; *C.A.*, 120 (1994) 235230k.
- 3384 Nendza, R.: (Chromatographic methods for protein purification. 5. Affinity chromatography). *Bioforum*, 16 (1993) 426-432; *C.A.*, 120 (1994) 72594r.
- 3385 Rhoads, A.R. and Friedberg, F.: Affinity chromatography of regulatory and signal-transducing proteins. *Chromatogr. Sci. Ser.*, 63(Handbook of Affinity Chromatography) (1993) 111-142; *C.A.*, 120 (1994) 128674m - a review with 169 refs.
- 3386 Rippel, G. and Szepesy, L.: Hydrophobic interaction chromatography of proteins on an Alkyl-Superose column. *J. Chromatogr. A*, 664 (1994) 27-32.
- 3387 Schmitt, J., Hess, H. and Stunnenberg, H.G.: Affinity purification of histidine-tagged proteins. *Mol. Biol. Rep.*, 18 (1993) 223-230; *C.A.*, 120 (1994) 72599w - a review with 25 refs.
- 3388 Thiele, C. and Fahrenholz, F.: Photocleavable biotinylated ligands for affinity chromatography. *Anal. Biochem.*, 218 (1994) 330-337.
- See also 2784, 2868, 2880, 2889, 2891, 2898, 3431, 4071.
- 19b. *Proteins of cells, viruses and subcellular particles*
- 3389 Chalkley, G.E., Knowles, P.P., Whitehead, P.C. and Cofer, A.I.: Biochemical characterisation of purified human wild-type p53 overexpressed in insect cells. *Eur. J. Biochem.*, 221 (1994) 167-175.
- 3390 Higashi, K. and Ogawara, H.: Daidzein inhibits insulin- or insulin-like growth factor-1-mediated signaling in cell cycle progression of Swiss 3T3 cells. *Biochim. Biophys. Acta*, 1221 (1994) 29-35.
- 3391 Liu, M.Y., Torgrimson, A. and Colombini, M.: Characterization and partial purification of the VDAC-channel-modulating protein from calf liver mitochondria. *Biochim. Biophys. Acta*, 1185 (1994) 203-212.
- 3392 Okumura, N., Miyatake, Y., Takao, T., Tamaru, T., Nagai, K., Okada, M. and Nakagawa, H.: Vasoactive intestinal peptide induces differentiation and MAP kinase activation in PC12h cells. *J. Biochem. (Tokyo)*, 115 (1994) 304-308.
- 3393 Page, M.G.P.: Purification of membrane transport proteins and receptors by immobilized-ligand affinity chromatography. *Chromatogr. Sci. Ser.*, 63(Handbook of Affinity Chromatography) (1993) 143-160; *C.A.*, 120 (1994) 186324h - a review with 84 refs.
- 3394 Zimmermann, A., Mertens, T., Schulz, A., Kruppenbacher, J.P., Nelsen-Salz, B. and Eggers, H.J.: HPLC is an effective and fast method for analysis of viral proteins: a study of encephalomyocarditis virus mutants differing in pathogenicity. *J. Gen. Virol.*, 74 (1993) 2759-2763; *C.A.*, 120 (1994) 100915s.
- See also 3277, 3355, 3407, 3630.
- 19c. *Proteins synthesized by genetic manipulation, monoclonal antibodies*
- 3395 Asenjo, J.A., Turner, R.E., Mistry, S.L. and Kaul, A.: Separation and purification of recombinant proteins from *Escherichia coli* with aqueous two-phase systems. *J. Chromatogr. A*, 668 (1994) 129-137.
- 3396 Barthel, T. and Kula, M.R.: Rapid purification of DesPro(2)-Val15-Leu17-aprotinin from the culture broth of a recombinant *Saccharomyces cerevisiae*. *Biotechnol. Bioeng.*, 42 (1993) 1331-1336; *C.A.*, 120 (1994) 72645h.
- 3397 Caplin, B.E., Hettich, L.A. and Marshall, M.S.: Substrate characterization of the *Saccharomyces cerevisiae* protein farnesyltransferase and type-I protein geranylgeranyltransferase. *Biochim. Biophys. Acta*, 1205 (1994) 39-48.
- 3398 Guo, L. and Liu, S.: (Separation of recombinant human interferon gamma on gel exclusion chromatography and the effect of denaturing agents on separation results). *Fenxi Huaxue*, 22 (1994) 169-171; *C.A.*, 120 (1994) 241950m.
- 3399 Hassinen, C., Köhler, K. and Veide, A.: Polyethylene glycol-potassium phosphate aqueous two-phase systems. Insertion of short peptide units into a protein and its effects on partitioning. *J. Chromatogr. A*, 668 (1994) 121-128.
- 3400 Iwakura, M. and Morikawa, M.: Availability of dihydrofolate reductase affinity handle in expressing human prolactin as a soluble fusion protein. *Biosci., Biotechnol., Biochem.*, 57 (1993) 1955-1957; *C.A.*, 120 (1994) 46936j.

For additional information see C.A.:

120 (1994) 49104x, 72891k, 100891f, 100892g, 239341v.

- 3401 Lewis, D.A., Guzzetta, A.W., Hancock, W.S. and Costello, M.: Characterization of humanized anti-TAC, an antibody directed against the interleukin 2 receptor, using electrospray ionization mass spectrometry by direct infusion, LC/MS, and MS/MS. *Anal. Chem.*, 66 (1994) 585-595.
- 3402 Orita, T., Oh-eda, M., Hasegawa, M., Kuboniwa, H., Esaki, K. and Ochi, N.: Polypeptide and carbohydrate structure of recombinant human interleukin-6 produced in Chinese hamster ovary cells. *J. Biochem. (Tokyo)*, 115 (1994) 345-350.
- 3403 Pedersen, J., Emborg, C., Sorensen, J. and Biedermann, K.: Purification of recombinant proteins by immunoaffinity chromatography with preselected antibodies. *Biotechnol. Tech.*, 7 (1993) 847-852; *C.A.*, 120 (1994) 264947e.
- 3404 Sarmiento, M.J., Pires, M.J., Cabral, J.M.S. and Aires-Barros, M.R.: Liquid-liquid extraction of a recombinant protein, cytochrome *b₅*, with aqueous two-phase systems of polyethylene glycol and potassium phosphate salts. *J. Chromatogr. A*, 668 (1994) 117-120.
- 3405 Sebastião, M.J., Cabral, J.M.S. and Aires-Barros, M.R.: Partitioning of recombinant *Fusarium solani pisi* cutinase in polyethylene glycol-aqueous salt solution two-phase systems. *J. Chromatogr. A*, 668 (1994) 139-144.
- See also 2894, 3489, 3517.
- 19d. *Microbial and plant proteins*
- 3406 Benaroudj, N., Fang, B., Triniolles, F., Ghelis, C. and Ladjimi, M.M.: Overexpression in *Escherichia coli*, purification and characterization of the molecular chaperone HSC70. *Eur. J. Biochem.*, 221 (1994) 121-128.
- 3407 Damm, I. and Green, B.R.: Separation of closely related intrinsic membrane polypeptides of the photosystem II light-harvesting complex (LHC II) by reversed-phase high-performance liquid chromatography on a poly(styrene-divinylbenzene) column. *J. Chromatogr. A*, 664 (1994) 33-38.
- 3408 Dong, T.X., Ng, T.B., Yeung, H.W. and Wong, R.N.S.: Isolation and characterization of a novel ribosome-inactivating protein, β -kiriowin, from the seeds of *Trichosanthes kirilowii*. *Biochem. Biophys. Res. Commun.*, 199 (1994) 387-393.
- 3409 Ducret, A., Sidler, W., Frank, G. and Zuber, H.: The complete amino acid sequence of R-phycoyanin-I α and β subunits from the red alga *Porphyridium cruentum*. Structural and phylogenetic relationships of the phycocyanins within the phycobiliprotein families. *Eur. J. Biochem.*, 221 (1994) 563-580.
- 3410 Duval, M., Job, C., Alban, C., Douce, R. and Job, D.: Developmental patterns of free and protein-bound biotin during maturation and germination of seeds of *Pisum sativum*: characterization of a novel seed-specific biotinylated protein. *Biochem. J.*, 299 (1994) 141-150.
- 3411 Fritz, G. and Aktories, K.: ADP-ribosylation of Rho proteins by *Clostridium botulinum* exoenzyme C3 is influenced by phosphorylation of Rho-associated factors. *Biochem. J.*, 300 (1994) 133-139.
- 3412 Galleschi, L., Grilli, I., Capocchi, A. and Moroni, P.: Gliadin extraction for reversed-phase high-performance liquid chromatography: an assessment of methodology. *Phyton (Horn)*, 33 (1993) 7-14; *C.A.*, 120 (1994) 189936v.
- 3413 Ho, K.K. and Tan, S.: Use of adsorption chromatography on Sephacryl S-500 for improved separation of isoforms of soluble photosynthetic catalysts from cyanobacteria. *J. Liq. Chromatogr.*, 17 (1994) 833-845.
- 3414 Huebner, F.R. and Bietz, J.A.: Improved chromatographic separation and characterization of ethanol-soluble wheat proteins. *Cereal Chem.*, 70 (1993) 506-511; *C.A.*, 120 (1994) 161855u.
- 3415 Jacobson, S.C., Andersson, S., Allenmark, S.G. and Guiochon, G.: Estimation of the number of enantioselective sites of bovine serum albumin using frontal chromatography. *Chirality*, 5 (1993) 513-515; *C.A.*, 120 (1994) 239336x.
- 3416 Kerfeld, C.A., Yeates, T.O. and Thornber, J.P.: Biochemical and spectroscopic characterization of the reaction center-LH1 complex and the carotenoid-containing B820 subunit of *Chromatium purpuratum*. *Biochim. Biophys. Acta*, 1185 (1994) 193-202.
- 3417 Kimura, M., Kouzuma, Y., Abe, K. and Yamasaki, N.: On a Bowman-Birk family proteinase inhibitor from *Erythrina variegata* seeds. *J. Biochem. (Tokyo)*, 115 (1994) 369-372.
- 3418 Larionova, N.I., Gladysheva, I.P., Topchieva, I.N. and Kazanskaya, N.F.: (Conjugation of the classical soybean inhibitor of the Bowman-Birk type with a copolymer of ethylene oxide and propylene oxide). *Biokhimiya (Moscow)*, 58 (1993) 1658-1664.
- 3419 Latchinian-Sadek, L. and Thomas, D.Y.: Secretion, purification and characterization of a soluble form of the yeast KEX1-encoded protein from insect-cell cultures. *Eur. J. Biochem.*, 219 (1994) 647-652.
- 3420 Lynch, A.S., Tyrrell, R., Smerdon, S.J., Briggs, G.S. and Wilkinson, A.J.: Characterization of the CysB protein of *Klebsiella aerogenes*: direct evidence that N-acetylserine rather than O-acetylserine serves as the inducer of the cysteine regulon. *Biochem. J.*, 299 (1994) 129-136.
- 3421 Marcone, M.F., Bondi, M.C. and Yada, R.Y.: Isolation of soybean 11S globulin by isoelectric precipitation and Sephacryl S-300 gel filtration chromatography: a new purification technique. *Biosci. Biotechnol., Biochem.*, 58 (1994) 413-415; *C.A.*, 120 (1994) 211964b.
- 3422 McDougall, J. and Wittman-Liebold, B.: Comparative analysis of the protein components from 5S rRNA-protein complexes of halophilic archaeobacteria. *Eur. J. Biochem.*, 221 (1994) 779-785.
- 3423 Sánchez-Ferrer, A., Pérez-Gilbert, M., Núñez, E., Bru, R. and García-Carmona, F.: Triton X-114 phase partitioning in plant protein purification. *J. Chromatogr. A*, 668 (1994) 75-83 - a review with 44 refs.
- 3424 Schoel, B., Welzel, M. and Kaufmann, S.H.E.: Hydrophobic interaction chromatography for the purification of cytolytic bacterial toxins. *J. Chromatogr. A*, 667 (1994) 131-139.
- 3425 Smith, C.M., Duff, S.M.G. and Chollet, R.: Partial purification and characterization of maize-leaf pyruvate, orthophosphate dikinase regulatory protein: a low-abundance, mesophyll-chloroplast stromal protein. *Arch. Biochem. Biophys.*, 308 (1994) 200-206.
- 3426 Tanabe, M., Asano, T., Moriya, N., Sugino, H. and Kakinuma, A.: Isolation and characterization of *Streptovorticillium* anticoagulant (SAC), a novel protein inhibitor of blood coagulation produced by *Streptovorticillium cinnamoneum* subsp. *cinnamoneum*. *J. Biochem. (Tokyo)*, 115 (1994) 743-751.

- 3427 Tanabe, M., Kawahara, K., Asano, T., Kato, K. and Kakinuma, A.: Primary structure and reactive site of *Streptovorticillium* anticoagulant (SAC), a novel protein inhibitor of blood coagulation produced by *Streptovorticillium cinnamomeum* subsp. *cinnamomeum*. *J. Biochem. (Tokyo)*, 115 (1994) 752-761.
- 3428 Terada, S., Fujimura, S., Katayama, H., Nagasawa, M. and Kimoto, E.: Purification and characterization of two Kunitz family subtilisin inhibitors from seeds of *Canavalia lineata*. *J. Biochem. (Tokyo)*, 115 (1994) 392-396.
- For additional information see C.A.:
120 (1994) 239524g.
- See also 3352, 3363, 4071.
- 19e. *Proteins of blood, serum and blood cells*
- 3429 Andersson, K., Wingren, C. and Hansson, U.B.: Liquid-liquid partition chromatography as a method to examine surface properties of antibodies and antigen-antibody complexes. *Scand. J. Immunol.*, 38 (1993) 95-101; C.A., 120 (1994) 214660k.
- 3430 Baso, D., Panozzo, M.P., del Giudice, G., Carbone, M.D., Meggiato, T., del Favero, G. and Plebani, M.: Concanavalin A affinity pattern of CA 19-9: an *in vitro* and *in vivo* study. *Clin. Chim. Acta*, 224 (1994) 219-221.
- 3431 Burton, S.J.: Considerations in the use of synthetic dye-ligand affinity adsorbents for the manufacture of blood proteins. *Colloq. INSERM*, 227(Biotechnology of Blood Proteins) (1993) 19-24; C.A., 120 (1994) 49202c.
- 3432 Chai, Y.-C., Ashraf, S.S., Rokutan, K., Johnston, R.B., Jr. and Thomas, J.A.: S-Thiolation of individual human neutrophil proteins including actin by stimulation of the respiratory burst: evidence against a role for glutathione disulfide. *Arch. Biochem. Biophys.*, 310 (1994) 273-281.
- 3433 Cochet, S., Hasnaoui, M'H., Debbia, M., Kroviarski, Y., Lambin, P., Cartron, J.P. and Bertrand, O.: Chromatography of human immunoglobulin G on immobilized Drimarene Rubine R/K-5BL. Study of mild, efficient elution procedures. *J. Chromatogr. A*, 663 (1994) 175-186.
- 3434 Dawidowicz, A.L., Rauckyte, T. and Rogalski, J.: High performance affinity chromatography for analysis of human antithrombin III. *J. Liq. Chromatogr.*, 17 (1994) 817-831.
- 3435 Drif, L. and Brehelin, M.: Purification and characterization of an agglutinin from the hemolymph of *Locusta migratoria* (Orthoptera). *Insect Biochem. Mol. Biol.*, 24 (1994) 283-289; C.A., 120 (1994) 239356d.
- 3436 Feldman, P.A., Harris, L., Evans, D.R. and Evans, H.E.: Preparation of a high purity factor IX concentrate using metal chelate affinity chromatography. *Colloq. INSERM*, 227(Biotechnology of Blood Proteins) (1993) 63-68; C.A., 120 (1994) 49204e.
- 3437 Field, M.C., Amatayakul-Chantler, S., Rademacher, T.W., Rudd, P.M. and Dwek, R.A.: Structural analysis of the N-glycans from human immunoglobulin A1: comparison of normal human serum immunoglobulin A1 with that isolated from patients with rheumatoid arthritis. *Biochem. J.*, 299 (1994) 261-275.
- 3438 Hartmann, H., Stelling, O. and Schell, D.: Rapid and highly sensitive quantification of mouse IgG combining chromatographic and immunoanalytic procedures. *Proc. MoBBEL 1992*, 7 (1993) 107-110; C.A., 120 (1994) 214649p.
- 3439 Herring, S.W.: Purification of factor IX. *PCT Int. Appl. WO 94 01,120* (Cl. A61K35/16), 20 Jan. 1994, US Appl. 913,666, 14 Jul. 1992; 21 p.; C.A., 120 (1994) 144127r.
- 3440 Huang, C. and Liang, N.: (Purification and identification of protilin in pig platelets). *Shengwu Huaxue Zazhi*, 10 (1994) 55-57; C.A., 120 (1994) 211733a.
- 3441 Jensen, P.E.H., Stigbrand, T. and Shanbhag, V.P.: Use of hydrophobic affinity partitioning as a method for studying various conformational states of the human α -macroglobulins. *J. Chromatogr. A*, 668 (1994) 101-106.
- 3442 Kuraya, N., Omichi, K., Nishimura, H., Iwanaga, S. and Hase, S.: Structural analysis of O-linked sugar chains in human blood clotting factor IX. *J. Biochem. (Tokyo)*, 114 (1993) 763-765.
- 3443 Ninfali, P., Baronciani, L., Rapa, S., Marziani, D. and Mannello, F.: Goat immunoglobulin purification on phosphocellulose and DEAE Affi-Gel Blue. *Prepar. Biochem.*, 24 (1994) 1-13.
- 3444 Okubo, K., Kang, D., Hamasaki, N. and Jeenings, M.L.: Red blood cell band 3. Lysine 539 and lysine 851 react with the same H₂DIDS (4,4'-diisothiocyanodihydrostilbene-2,2'-disulfonic acid) molecule. *J. Biol. Chem.*, 269 (1994) 1918-1926.
- 3445 Pazur, J.H., Liu, B. and Li, N.Q.: Isolation of anti-mannose antibodies by affinity chromatography on mannosyl-Sepharose. *Nat. Prod. Lett.*, 1 (1992) 51-57; C.A., 120 (1994) 264957h.
- 3446 Regoeczi, E., Chindemi, P.A. and Hu, W.-L.: Interaction of transferrin and its iron-binding fragments with heparin. *Biochem. J.*, 299 (1994) 819-823.
- 3447 Sinha, D., Yang, X., Emig, F. and Kirby, E.P.: Isolation and characterization of two protease inhibitors from bovine plasma. *J. Biochem. (Tokyo)*, 115 (1994) 387-391.
- 3448 Turrini, F., Mannu, F., Cappadoro, M., Ulliers, D., Giribaldi, G. and Arese, P.: Binding of naturally occurring antibodies to oxidatively and nonoxidatively modified erythrocyte band 3. *Biochim. Biophys. Acta*, 1190 (1994) 297-303.
- 3449 Vincent, C., Dennonroy, L. and Revillard, J.-P.: Molecular variants of β_2 -microglobulin in renal insufficiency. *Biochem. J.*, 298 (1994) 181-187.
- 3450 Voss, T., Falkner, E., Ahorn, H., Krystek, E., Maurer-Fogy, I., Bodo, G. and Haputmann, R.: Periplasmic expression of human interferon- $\alpha 2c$ in *Escherichia coli* results in a correctly folded molecule. *Biochem. J.*, 298 (1994) 719-725.
- 3451 Yamaguchi, K., Kato, N., Sakai, N., Matsumoto, M., Nagasawa, S., Hatsuse, H., Toyoguchi, T., Moriya, H. and Sakiyama, H.: Immune complex independent activation of complement, C1s secreted from hamster embryo malignant fibroblasts, Nil2C2 in serum free culture medium. *Biochim. Biophys. Acta*, 1205 (1994) 133-138.
- For additional information see C.A.:
120 (1994) 211731y, 212023f, 242510e, 242511f, 264964h, 267609g.
- See also 2960, 3340, 3374, 3398.
- 19f. *Structural and muscle proteins*
- 3452 Bishop, P.N., Crossman, M.V., McLeod, D. and Ayad, S.: Excretion and characterization of the tissue forms of collagen types II and IX from bovine vitreous. *Biochem. J.*, 299 (1994) 497-505.

- 3453 Fertala, A., Sieron, A.L., Ganguly, A., Li, S.-W., Ala-Kokko, L., Anumula, K.R. and Prockop, D.J.: Synthesis of recombinant human procollagen II in a stably transfected tumor cell line (HT1080). *Biochem. J.*, 298 (1994) 31-37.
- 3454 Graham, L. and Gallop, P.M.: Covalent protein crosslinks: general detection, quantitation, and characterization via modification with diphenylborinic acid. *Anal. Biochem.*, 217 (1994) 298-305.
- 3455 Heeley, D.H.: Investigation of the effects of phosphorylation of rabbit striated muscle α , α -tropomyosin and rabbit skeletal muscle troponin-T. *Eur. J. Biochem.*, 221 (1994) 129-137.
- 3456 Iio, T.: Melittin-binding of troponin C. *J. Biochem. (Tokyo)*, 114 (1993) 773-778.
- 3457 Iversen, P., Ertbjerg, P., Larsen, L.M., Monllao, S. and Moeller, A.J.: An FPLC method for determination of calpains and calpastatin in porcine m longissimus dorsi. *Biochimie*, 75 (1993) 869-872; *C.A.*, 120 (1994) 72034h.
- 3458 Ngai, S. Sönnichsen, F.D. and Hodges, R.S.: Photochemical cross-linking between native rabbit skeletal troponin C and benzoylbenzoyl-troponin I inhibitory peptide, residues 104-115. *J. Biol. Chem.*, 269 (1994) 2165-2172.
- 3459 Stecker, J.J., Kramer, D.L. and Brand, J.S.: Preparative chromatography and analysis of molecular weight fractions of gelatin. *J. Photogr. Sci.*, 40 (1992) 134-137; *C.A.*, 120 (1994) 167217j.
- 3460 Tillet, E., Wiedemann, H., Golbik, R., Pan, T.-C., Zhang, R.-Z., Mann, K., Chu, M.-L. and Timpl, R.: Recombinant expression and structural and binding properties of α 1(VI) and α 2(VI) chains of human collagen type VI. *Eur. J. Biochem.*, 221 (1994) 177-187.
- 3461 Zaviin, P.M., Nus, V.P., Ovchinnikov, A.N., Sakverelidze, M.A. and Yampol'skaya, G.P.: (Analysis of industrial photographic gelatins by highly efficient liquid chromatography). *Zh. Nauchn. Prikl. Fotogr.*, 38 (1993) 8-13; *C.A.*, 120 (1994) 120508q.
- For additional information see *C.A.*:
120 (1994) 211928t.
- See also 3351, 3358.
- 19g. *Protamines, histones and other nuclear proteins*
- 3462 Inomata, Y., Wada, T., Handa, H., Fujimoto, K. and Kawaguchi, H.: Affinity latex for the purification of specific proteins. In: Takagi, T. (Editor), *Proc. Int. Conf. Intell. Mater.*, 1st, Technomic, Lancaster, 1993, pp. 442-445; *C.A.*, 120 (1994) 262376u.
- 3463 Mahajan, P.B.: A simple method for removing transcriptional inhibitors from nuclear extracts. *Methods Mol. Cell. Biol.*, 4 (1993) 1-9; *C.A.*, 120 (1994) 126244x.
- 3464 Pironcheva, G. and Rusev, G.: Isolation and purification of U7 snRNP particles. *Int. J. Biochem.*, 26 (1994) 269-272; *C.A.*, 120 (1994) 211730x.
- 3465 Stanzel, M., Schön, A. and Sprinzl, M.: Discrimination against misacylated tRNA by chloroplast elongation factor Tu. *Eur. J. Biochem.*, 219 (1994) 435-439.
- 3466 Wakim, B.T. and Aswad, G.D.: Ca^{2+} -Calmodulin-dependent phosphorylation of arginine in histone 3 by a nuclear kinase from mouse leukemia cells. *J. Biol. Chem.*, 269 (1994) 2722-2727.
- 19h. *Chromoproteins and metalloproteins*
- 3467 Baba, Y. and Kawano, Y.: Selective adsorption of hemoglobin with copper-phthalocyanine trisulfonate derivative immobilized on gel beads. *Chem. Lett.*, (1994) 181-184; *C.A.*, 120 (1994) 157807n.
- 3468 Cengiz, S., Cengiz, M. and Ozpeker, F.: Spectrophotometric wavelength selection during detection of hemoglobin components by high performance liquid chromatography. *Turk. J. Med. Sci.*, 19 (1993) 111-117; *C.A.*, 120 (1994) 211937v.
- 3469 Commission of the European Communities (Luxembourg): The determination of HbA_{1c} (glycated hemoglobin) in a human hemolyzate reference material RM 405. *Comm. Eur. Communities*, [Rep.] EUR 15136, 1993, 63 pp.; *C.A.*, 120 (1994) 128939b.
- 3470 Joshi, A.A. and McDonald, M.J.: Role of α and β carboxyl-terminal residues in the kinetics of human oxyhemoglobin dimer assembly. *J. Biol. Chem.*, 269 (1994) 8549-8553.
- 3471 Khoo, C. and Cousins, R.J.: Purification and properties of rat cysteine-rich intestinal protein. *Biochem. J.*, 299 (1994) 445-450.
- 3472 Matsuoka, K., Seta, K., Yamakawa, Y., Okuyama, T., Shinoda, T. and Isobe, T.: Covalent structure of bovine brain calreticulin. *Biochem. J.*, 298 (1994) 435-442.
- 3473 Pauwels, M., van Weyenbergh, J., Soumillion, A., Proost, P. and de Ley, M.: Induction by zinc of specific metallothionein isoforms in human monocytes. *Eur. J. Biochem.*, 220 (1994) 105-110.
- 3474 Simo Siso, J.M., Cliville, G.X., Sans Mateu, T. and Joven Maried, J.: (Evaluation of a new automated chromatographic method for the determination of glycohemoglobin). *Rev. Soc. Esp. Quim. Clin.*, 12 (1993) 242-245; *C.A.*, 120 (1994) 128752k.
- 3475 Suzuki, T., Ohta, T., Yuasa, H.J. and Takagi, T.: The giant extracellular hemoglobin from the polychaete *Neanthes diversicolor*. The cDNA-derived amino acid sequence of linker chain L2 and the exon/intron boundary conserved in linker genes. *Biochim. Biophys. Acta*, 1217 (1994) 291-296.
- 3476 Townson, S.M., Hanson, G.R., Umproft, J.A. and Umproft, P.: A purified ferredoxin from *Giardia duodenalis*. *Eur. J. Biochem.*, 220 (1994) 439-446.
- 3477 Wallace, C.J.A. and Tanaka, Y.: Improving cytochrome c function by protein engineering?: studies of site-directed mutants of the human protein. *J. Biochem. (Tokyo)*, 115 (1994) 693-700.
- 3478 Yamashita, K., Ohkura, T., Ideo, H., Ohno, K. and Kanai, M.: Electrospray ionization-mass spectrometric analysis of serum transferrin isoforms in patients with carbohydrate-deficient glycoprotein syndrome. *J. Biochem. (Tokyo)*, 114 (1993) 766-769.
- For additional information see *C.A.*:
120 (1994) 49195c, 49477w, 129022j, 187769g, 264976p.
- See also 3446, 3501.
- 19i. *Proteins of glands, gland products, various zymogens (incl. milk proteins)*
- 3479 Akama, K., Oka, S., Tobita, T. and Hayashi, H.: The amino acid sequence of a boar transition protein 3. *J. Biochem. (Tokyo)*, 115 (1994) 58-65.

- 3480 Check, J.H., Zavos, P.M., Katsoff, D. and Kiefer, D.: Effects of Percoll discontinuous density gradients vs. SpermPrep™II vs. Sephadex G-50 gel filtration on semen parameters. *Tohoku J. Exp. Med.*, 169 (1993) 225-231; *C.A.*, 120 (1994) 211868y.
- 3481 Dos Reis Coimbra, J., Thömmes, J. and Kula M.-R.: Continuous separation of whey proteins with aqueous two-phase systems in a Graesser contactor. *J. Chromatogr. A*, 668 (1994) 85-94.
- 3482 Imbert, A. and Nicolas, M.: (Detection in milk of wheys on enzymic origin by identification of their caseinomacropeptides). *Sci. Aliments*, 13 (1993) 545-558; *C.A.*, 120 (1994) 215528k.
- 3483 Kameswari, D.B. and Prasad, K.S.N.: A simple and efficient method for the purification of seminal plasmin, an antimicrobial protein from bovine seminal plasma. *Prepar. Biochem.*, 24 (1994) 15-24.
- 3484 Law, A.J.R., Leaver, J., Banka, J.M. and Horne, D.S.: Quantitative fractionation of whey proteins by gel permeation FPLC. *Milchwissenschaft*, 48 (1993) 663-666; *C.A.*, 120 (1994) 268594k.
- 3485 Ng-Kwai-Hang, K.F. and Chin, D.: Semipreparative isolation of bovine casein components by high-performance liquid chromatography. *Int. Dairy J.*, 4 (1994) 99-110; *C.A.*, 120 (1994) 161885d.
- 3486 Rougeot, C., Rosinski-Chupin, I., Njamkero, E. and Rougeon, F.: Selective processing of submandibular rat 1 protein at dibasic cleavage sites. Salivary and bloodstream secretion products. *Eur. J. Biochem.*, 219 (1994) 765-773.
- 3487 Solovyov, D.A. and Ugarova, T.P.: (Purification and characterization of α -specific thrombin-like enzymes from the venoms of Middle Asian pit viper snakes *Agkistrodon halys halys* and *Agkistrodon halys Blomhoffii*). *Biokhimiya (Moscow)*, 58 (1993) 1221-1233. *C.A.*,

For additional information see *C.A.*:
120 (1994) 129028r.

See also 3357, 3368.

19j. *Proteins of brain, cerebrospinal fluid and eye*

- 3488 Van Noort, J.M., El Ouagmiri, M., Boon, J. and van Sechel, A.C.: Fractionation of central nervous system myelin proteins by reversed-phase high-performance liquid chromatography. *J. Chromatogr. B*, 653 (1994) 155-161.

For additional information see *C.A.*:
120 (1994) 96954j.

See also 3348.

19k. *Proteins of neoplastic tissue and transformed cells*

- 3489 Paquet, A., Lévesque, A. and Pagée, M.: Rapid purification method for human recombinant tumor necrosis factor alpha. *J. Chromatogr. A*, 667 (1994) 125-130.

See also 3451.

19l. *Specific binding and receptor proteins*

- 3490 Abo, A., Webb, M.R., Grogan, A. and Segal, A.W.: Activation of NADPH oxidase involves the dissociation of p21^{rac} from its inhibitory GDP/GTP exchange protein (rhoGDI) followed by its translocation to the plasma membrane. *Biochem. J.*, 298 (1994) 585-591.
- 3491 Andreeva, L. and Crompton, A.: An ADP-sensitive cyclosporin-A-binding protein in rat liver mitochondria. *Eur. J. Biochem.*, 221 (1994) 261-268.
- 3492 Anuradha, P., Khan, S.M., Karthikeyan, N. and Thampan, R.V.: The nonactivated estrogen receptor (naER) of the goat uterus is a tyrosine kinase. *Arch. Biochem. Biophys.*, 309 (1994) 195-204.
- 3493 Bignon, C., Sakal, E., Belair, L., Chapnik-Cohen, N., Djiane, J. and Gertler, A.: Preparation of the extracellular domain of the rabbit prolactin receptor expressed in *Escherichia coli* and its interaction with lactogenic hormones. *J. Biol. Chem.*, 269 (1994) 3318-3324.
- 3494 Borg, C., Lim, C.T., Yeomans, D.C., Dieter, J.P., Komiotis, D., Anderson, E.G. and le Breton, G.C.: Purification of rat brain, rabbit aorta, and human platelet thromboxane A₂/prostaglandin H₂ receptors by immunoaffinity chromatography employing anti-peptide and anti-receptor antibodies. *J. Biol. Chem.*, 269 (1994) 6109-6116.
- 3495 Boulanger, J., Huesca, M., Arab, S. and Lingeood, C.A.: Universal method for the facile production of glycolipid/lipid matrices for the affinity purification of binding ligands. *Anal. Biochem.*, 217 (1994) 1-6.
- 3496 Cadena, D.L., Chan, C. and Gill, G.N.: The intracellular tyrosine kinase domain of the epidermal growth factor receptor undergoes a conformational change upon autophosphorylation. *J. Biol. Chem.*, 269 (1994) 260-265.
- 3497 Calvete, J.J., Mann, K., Schäfer, W., Fernandez-Lafuente, R. and Guisán, J.M.: Proteolytic degradation of the RGD-binding and non-RGD-binding conformers of human platelet integrin glycoprotein IIb/IIIa: clues for identification of regions involved in the receptor's activation. *Biochem. J.*, 298 (1994) 1-7.
- 3498 Cattini, R., Robinson, D., Gill, O., Jolley, N. and Bacarese-Hamilton, T.: Measurement of prostate-specific antigen in serum using four different immunoassays. *Eur. J. Clin. Chem. Clin. Biochem.*, 32 (1994) 181-185.
- 3499 Christensson, A. and Lilja, H.: Complex formation between protein C inhibitor and prostate-specific antigen *in vitro* and in human semen. *Eur. J. Biochem.*, 220 (1994) 45-53.
- 3500 Hocher, B., Dürr, J.A. and Henssen, J.: Effect of bacitracin on the degradation of a vasopressin receptor ligand with high affinity for the V₁ and V₂ vasopressin isoreceptors. *Eur. J. Clin. Chem. Clin. Biochem.*, 32 (1994) 61-64.
- 3501 Jost, M., Weber, K. and Gerke, V.: Annexin II contains two types of Ca²⁺-binding sites. *Biochem. J.*, 298 (1994) 553-559.
- 3502 Karthikeyan, N. and Thampan, R.V.: A DNA-binding (R-I) and a non-DNA-binding (R-II) estrogen receptor in the goat uterine nucleus: purification and characterization. *Arch. Biochem. Biophys.*, 309 (1994) 205-213.
- 3503 Kearse, K.P. and Singer, A.: Isolation of immature and mature T cell receptor complexes by lectin affinity chromatography. *J. Immunol. Methods*, 167 (1994) 75-81; *C.A.*, 120 (1994) 160906z.

- 3504 Kortt, A.A., Malby, R.L., Caldwell, J.B., Gruen, L.C., Ivancic, N., Lawrence, M.C., Howlett, G.J., Webster, R.G., Hudson, P.J. and Colman, P.M.: Recombinant anti-sialidase single-chain variable fragment antibody. Characterization, formation of dimer and higher-molecular-mass multimers and the solution of the crystal structure of the single-chain variable fragment/sialidase complex. *Eur. J. Biochem.*, 221 (1994) 151-157.
- 3505 Kuroki, M., Kuwahara, M., Murakami, M., Matsumoto, Y. and Matsuoka, Y.: Fractionation of carcinoembryonic antigen and related antigens in normal adult feces using a gradient medium percoll. *Prepar. Biochem.*, 24 (1994) 25-40.
- 3506 Lançar-Benba, J., Foucher, B. and Saint-Macary, M.: Purification of the rat-liver mitochondrial dicarboxylate carrier by affinity chromatography on immobilized malate dehydrogenase. *Biochim. Biophys. Acta*, 1190 (1994) 213-216.
- 3507 Lesca, P., Witkamp, R., Maurel, P. and Galtier, P.: The pig as a model for studying AH receptor and other PAH-binding proteins in man. *Biochem. Biophys. Res. Commun.*, 200 (1994) 475-481.
- 3508 Matsui, M., Kobayashi, A., Kubo, T. and Natori, S.: Purification and characterization of ATPB, a novel protein that binds to A/T stretches in three segments of the *Sarcophaga* lectin gene. *Eur. J. Biochem.*, 219 (1994) 449-454.
- 3509 Moncollin, V. and Egly, J.M.: Purification of nucleic acid-binding proteins by affinity chromatography. *Chromatogr. Sci. Ser.*, 63(Handbook of Affinity Chromatography) (1993) 161-175; C.A., 120 (1994) 186325j - a review with 37 refs.
- 3510 Osborn, A.H., Kelleher, M., Handman, E. and Goding, J.W.: Anti-galactosyl antibodies that react with unmodified agarose: a potential source of artifacts in immunoaffinity chromatography. *Anal. Biochem.*, 217 (1994) 181-184.
- 3511 Schenerman, M.A. and Collins, T.J.: Determination of a monoclonal antibody binding activity using immunodetection. *Anal. Biochem.*, 217 (1994) 241-247.
- 3512 Schletter, J., Krüger, C., Lottspeich, F. and Schütt, C.: Improved method for preparation of lipopolysaccharide-binding protein from human serum by electrophoretic and chromatographic separation techniques. *J. Chromatogr. B*, 654 (1994) 25-34.
- 3513 Strecker, A., Franke, P., Weise, C. and Hucho, F.: All potential glycosylation sites of the nicotinic acetylcholine receptor δ subunit from *Torpedo californica* are utilized. *Eur. J. Biochem.*, 220 (1994) 1005-1011.
- 3514 Takeda, S., Sagara, Y., Kita, K., Natori, S. and Sekimizu, K.: Purification of a GTP-binding protein localized in mitochondria. *J. Biochem. (Tokyo)*, 114 (1993) 684-690.
- 3515 Trifillieff, A., Lach, E., Mousli, M., Haddad, E.-B., Landry, Y. and Gies, J.-P.: Co-solubilization of bradykinin B₂ receptors and angiotensin-converting enzyme from guinea pig lung membranes. *Biochim. Biophys. Acta*, 1191 (1994) 109-116.
- 3516 Zhigis, L.S., Ivanov, A.E., Rapoport, E.M., Kovalenko, E.A., Getman, E.I. and Zubov, V.P.: Purification of sialic acid binding protein from saprophytic bacteria by hydrophobic-interaction chromatography on butyl-toyopearl and polymer-coated porous glass. *Biotechnol. Tech.*, 7 (1993) 667-670; C.A., 120 (1994) 100889m.

For additional information see C.A.:
120 (1994) 186728t.

See also 3123, 3192, 3342, 3343, 3347, 3357, 3388, 3599, 3841.

19m. Urinary proteins

For additional information see C.A.:
120 (1994) 49217m, 72820m.

19n. Other proteins (incl. proteinous inhibitors of enzymic activity)

- 3517 Lee, S.Y., Moon, H.J., Kurata, S., Kurama, T., Natori, S. and Lee, B.L.: Purification and molecular cloning of cDNA for an inducible antibacterial protein of larvae of a coleopteran insect, *Holotrichia diomphalia*. *J. Biochem. (Tokyo)*, 115 (1994) 82-86.
- 3518 Morelle, W., Capon, C., Balduyck, M., Sautiere, P., Kovach, M., Michalski, C., Fournet, B. and Mizon, J.: Chondroitin sulphate covalently cross-link the three polyphosphate chains of inter- α -trypsin inhibitor. *Eur. J. Biochem.*, 221 (1994) 881-888.
- 3519 Novak, A. and Lowden, J.A.: GM₂ Ganglioside activator occurs in multiple forms. *Biochim. Biophys. Acta*, 1199 (1994) 209-214.
- 3520 Shiomi, K., Yamaguchi, S., Shimakura, K., Nagashima, Y., Yamamori, K. and Matsui, T.: Affinity chromatographic purification of tetrodotoxin by use of tetrodotoxin-binding high molecular weight substances in the body fluid of shore crab (*Hemigrapsus sanguineus*) as ligands. *Toxicon*, 31 (1993) 1615-1618; C.A., 120 (1994) 184780e.

For additional information see C.A.:
120 (1994) 52827r, 72647k.

See also 3406, 3447.

20. ENZYMES AND ENZYME ACTIVITY ESTIMATION

- 3521 Friedberg, F. and Rhoads, A.R.: Affinity chromatography of enzymes. *Chromatogr. Sci. Ser.*, 63(Handbook of Affinity Chromatography) (1993) 79-109; C.A., 120 (1994) 72003x - a review with 98 refs.

See also 3380.

20a. Oxidoreductases

- 3522 Al-Kassim, L.S. and Tsai, C.S.: Purification and kinetic characterization of pickrel liver alcohol dehydrogenase with dual co-enzyme specificity. *Biochem. Cell Biol.*, 71 (1993) 421-426.
- 3523 Arieli, B., Shahak, Y., Taglicht, D., Hauska, G. and Padan, E.: Purification and characterization of sulfide-quinone reductase, a novel enzyme driving anoxygenic photosynthesis in *Oscillatoria limnetica*. *J. Biol. Chem.*, 269 (1994) 5705-5711.
- 3524 Ball, K.L., Dale, S., Weekes, J. and Hardie, D.G.: Biochemical characterization of two forms of 3-hydroxy-3-methyl-glutaryl-CoA reductase kinase from cauliflower (*Brassica oleracea*). *Eur. J. Biochem.*, 219 (1994) 743-750.
- 3525 Battu, S., Cook-Moreau, J. and Beneytout, J.L.: Stabilization of potato tuber lipoxygenase on talc. *Biochim. Biophys. Acta*, 1211 (1994) 270-276.
- 3526 Cabisco, E., Aguilar, J. and Ros, J.: Metal-catalyzed oxidation of Fe²⁺ dehydrogenases. Consensus target sequence between propanediol oxidoreductase of *Escherichia coli* and alcohol dehydrogenase II of *Zymomonas mobilis*. *J. Biol. Chem.*, 269 (1994) 6592-6597.

- 3527 Ehrenwald, E. and Fox, P.L.: Isolation of nonlabile human ceruloplasmin by chromatographic removal of a plasma metalloproteinase. *Arch. Biochem. Biophys.*, 309 (1994) 392-395.
- 3528 Fraser, P.D. and Bramley, P.M.: The purification of phytoene dehydrogenase from *Phycomyces blakesleeanus*. *Biochim. Biophys. Acta*, 1212 (1994) 59-66.
- 3529 Ghose, S. and Mattiasson, B.: A two step displacement and affinity chromatography purification in lactate dehydrogenase recovery from beef heart extract. *Biotechnol. Tech.*, 7 (1993) 615-620; *C.A.*, 120 (1994) 211704s.
- 3530 Guoqiang, D., Kaul, R. and Mattiasson, B.: Integration of aqueous two-phase extraction and affinity precipitation for the purification of lactate dehydrogenase. *J. Chromatogr. A*, 668 (1994) 145-152.
- 3531 Heiden, S., Hedderich, R., Setzke, E. and Thauer, R.K.: Purification of a two-subunit cytochrome-*b*-containing heterodisulfide reductase from methanol-grown *Methanosarcina barkeri*. *Eur. J. Biochem.*, 221 (1994) 855-861.
- 3532 Inouye, K., Osaki, A. and Tonomura, B.: Dissociation of dimer of bovine erythrocyte Cu,Zn-superoxide dismutase and activity of the monomer subunit: effects of urea, temperature, and enzyme concentration. *J. Biochem. (Tokyo)*, 115 (1994) 507-515.
- 3533 Joshi, M.S. and Rajagopalan, K.V.: Specific incorporation of molybdopterin in xanthine dehydrogenase of *Pseudomonas aeruginosa*. *Arch. Biochem. Biophys.*, 308 (1994) 331-334.
- 3534 Kirchberger, J., Erdmann, H., Hecht, H.-J. and Kopperschläger, G.: Studies of the interaction of NADH oxidase from *Thermus thermophilus* HB8 with triazine dyes. *J. Chromatogr. A*, 668 (1994) 153-164.
- 3535 Mandai, M., Iwaki, M. and Honda, Y.: Catalytic nature of tyrosine hydroxylase in bovine ocular tissues. *J. Biochem. (Tokyo)*, 115 (1994) 797-802.
- 3536 Ozols, J.: Isolation and structure of a third form of liver microsomal flavin monooxygenase. *Biochemistry*, 33 (1994) 3751-3757.
- 3537 QKasai, S.: Preparation of *P*-flavin-bound and *P*-flavin-free luciferase and *P*-flavin-bound β -subunit of luciferase from *Photobacterium phosphoreum*. *J. Biochem. (Tokyo)*, 115 (1994) 670-674.
- 3538 Richard-Forget, F., Goupy, P. and Nicolas, J.: New approaches for separating and purifying apple polyphenol oxidase isoenzymes: hydrophobic, metal chelate and affinity chromatography. *J. Chromatogr. A*, 667 (1994) 141-153.
- 3539 Rossmann, R., Sauter, M., Lottspeich, F. and Böck, A.: Maturation of the large subunit (HYCE) of *Escherichia coli* hydrogenase 3 requires nickel incorporation followed by C-terminal processing at Arg537. *Eur. J. Biochem.*, 220 (1994) 377-384.
- 3540 Samyn, B., Berks, B.C., Page, M.D., Ferguson, S.J. and van Beeumen, J.J.: Characterisation and amino acid sequence of cytochrome *c*-550 from *Thiosphaera pantotropha*. *Eur. J. Biochem.*, 219 (1994) 585-594.
- 3541 Skorkowski, E.F.: Specific activation of crustacean malic enzyme by substrate and cofactor in affinity chromatography. *J. Mar. Biotechnol.*, 1 (1993) 109-112; *C.A.*, 120 (1994) 185776v.
- 3542 Tsai, C.S. and Al-Kassim, L.S.: Multifunctional activities of pickerel liver alcohol dehydrogenase. *Biochem. Cell Biol.*, 71 (1993) 427-431.

For additional information see C.A.:
120 (1994) 48567p.

See also 3031.

20b. *Transferases (excl. E.C. 2.7.-)*

- 3543 Andersson, M., Wettsten, M., Borén, J., Magnusson, A., Sjöberg, A., Rustaeus, S. and Olofsson, S.O.: Purification of diacylglycerol: acyltransferase from rat liver to near homogeneity. *J. Lipid Res.*, 35 (1994) 535-545.
- 3544 Andreotti, G., Cubellis, M.V., Nitti, G., Sannia, G., Mai, X., Marino, G. and Adams, M.W.W.: Characterization of aromatic aminotransferases from the hyperthermophilic archaeon *Thermococcus litoralis*. *Eur. J. Biochem.*, 220 (1994) 543-549.
- 3545 Board, P., Russell, R.J., Marano, R.J. and Oakeshott, J.G.: Purification, molecular cloning and heterologous expression of a glutathione S-transferase from the Australian sheep blowfly (*Lucilia cuprina*). *Biochem. J.*, 299 (1994) 425-430.
- 3546 Fujiwara, S., Sakuraba, H. and Noguchi, T.: Identification of chicken liver mitochondrial alanine:2-oxoglutarate aminotransferase and its response to starvation. *Arch. Biochem. Biophys.*, 310 (1994) 497-503.
- 3547 Gerber, U., Jucknischke, U., Putzien, S. and Fuchsbaue, H.-L.: A rapid and simple method for the purification of transglutaminase from *Streptovorticillium mobaraense*. *Biochem. J.*, 299 (1994) 825-829.
- 3548 Koger, J.B., Howell, R.G., Kelly, M. and Jones, E.E.: Purification and properties of porcine liver ornithine transcarbamylase. *Arch. Biochem. Biophys.*, 309 (1994) 293-299.
- 3549 Saidha, T. and Schiff, J.A.: Purification and properties of a phenol sulphotransferase from *Euglena* using L-tyrosine as substrate. *Biochem. J.*, 298 (1994) 45-50.
- 3550 Tiainen, P. and Karhi, K.A.: Ultrasensitive time-resolved immunofluorometric assay of glutathione transferase alpha in serum. *Clin. Chem. (Washington)*, 40 (1994) 184-189.
- 3551 Tong, H. and Davis, L.: 2-Amino-3-ketobutyrate-CoA ligase from beef liver mitochondria. Purification and partial sequence. *J. Biol. Chem.*, 269 (1994) 4057-4064.

See also 3277, 3341, 3350, 3359.

20c. *Transferases transferring phosphorus containing groups (E.C. 2.7.-)*

- 3552 Angelov, I.: Characterization of a proline-directed casein kinase from bovine brain. *Arch. Biochem. Biophys.*, 310 (1994) 97-107.
- 3553 Bodenbach, L., Fauss, J., Robitzki, A., Krehan, A., Lorenz, P., Lozeman, F.J. and Pyerin, W.: Recombinant human, casein kinase II. A study with the complete set of subunits (α , α' and β), site-directed autophosphorylation mutants and a bicistronically expressed holoenzyme. *Eur. J. Biochem.*, 220 (1994) 263-273.
- 3554 Chambers, T.C., Pohl, J., Glass, D.B. and Kuo, J.F.: Phosphorylation by protein kinase C and cyclic AMP-dependent protein kinase of synthetic peptides derived from the linker region of human P-glycoprotein. *Biochem. J.*, 299 (1994) 309-315.
- 3555 Chilks, T.J. and Mak, A.S.: MAP kinases from bovine brain: purification and characterization. *Biochem. Cell Biol.*, 71 (1993) 544-555.

- 3556 Communi, D., Vanweyenberg, V. and Erneux, C.: Purification and biochemical properties of a high-molecular-mass inositol 1,4,5-trisphosphate 3-kinase isoenzyme in human platelets. *Biochem. J.*, 298 (1994) 669-673.
- 3557 Dementiev, A.A., Golyshin, P.N., Ryabchenko, N.F., Pustobaev, V.N. and Shlyapnikov, S.V.: (Two forms of extracellular low molecular mass ribonuclease of *Bacillus* sp. BCF 247: isolation and characterization of the enzyme). *Biokhimiya (Moscow)*, 58 (1993) 1258-1265.
- 3558 Hayashi, D.K. and Stark, B.C.: A novel tRNA precursor cleaving endoribonuclease from *Zea mays*. *Arch. Biochem. Biophys.*, 309 (1994) 123-128.
- 3559 Hei, Y., Pelech, S.L., Chen, X., Diamond, J. and McNeill, J.H.: Purification and characterization of a novel ribosomal S6 kinase from skeletal muscle of insulin-treated rats. *J. Biol. Chem.*, 269 (1994) 7816-7823.
- 3560 Lanciotti, R.A. and Bender, P.K.: Baculovirus-directed expression of the γ -subunit of phosphorylase kinase: purification and calmodulin dependence. *Biochem. J.*, 299 (1994) 183-189.
- 3561 Lindquist, L., Kaiser, R., Reeves, P.R. and Lindberg, A.A.: Purification, characterization, and high performance liquid chromatography assay of *Salmonella* glucose-1-phosphate cytidylyltransferase from the cloned rfbF gene. *J. Biol. Chem.*, 269 (1994) 122-126.
- 3562 Mohamed, A.H., Huang, W., Huang, W., Venkatachalam, K.V. and Wakil, S.J.: Isolation and characterization of a novel acetyl-CoA carboxylase kinase from rat liver. *J. Biol. Chem.*, 269 (1994) 6859-6865.
- 3563 Taylor, L.K., Swanson, K.D., Kerigan, J., Mobley, W. and Landreth, G.E.: Isolation and characterization of a nerve growth factor-regulated Fos kinase from PC12 cells. *J. Biol. Chem.*, 269 (1994) 308-318.
- 3564 Torres, R.J., Mateos, F.A., Puig, J.G. and Becker, M.A.: A simplified method for the determination of phosphoribosylpyrophosphate synthetase activity in hemolysates. *Clin. Chim. Acta*, 224 (1994) 55-63.
- 3565 Weekes, J., Hawley, S.A., Corton, J., Shugar, D. and Hardie, D.G.: Activation of rat liver AMP-activated protein kinase by kinase kinase in a purified, reconstituted system. Effects of AMP and AMP analogues. *Eur. J. Biochem.*, 219 (1994) 751-757.
- 3566 Yang, C., Yanagi, S., Wang, X., Sakai, K., Taniguchi, T. and Yamamura, H.: Purification and characterization of a protein-tyrosine kinase p72^{9k} from bovine spleen. *Eur. J. Biochem.*, 221 (1994) 973-978.
- See also 3354, 3365, 3390.
- 20d. *Hydrolases, acting on ester bonds (E.C. 3.1.-)*
- 3567 Ameis, D., Merkel, M., Eckerskorn, C. and Greten, H.: Purification, characterization and molecular cloning of human hepatic lysosomal acid lipase. *Eur. J. Biochem.*, 219 (1994) 905-914.
- 3568 Dietrich, A., Meister, M., Brazil, D., Camps, M. and Gierschik, P.: Stimulation of phospholipase C-2 by recombinant guanine-nucleotide-binding protein β , dimers produced in c baculovirus/insect cell expression system. Requirement of γ -subunit isoprenylation of phospholipase C. *Eur. J. Biochem.*, 219 (1994) 171-178.
- 3569 Ford, K., Waltho, J. and Hornby, D.: Component analysis and characterization of a nuclear deoxyribonucleotidase. *Biochem. J.*, 298 (1994) 727-732.
- 3570 Halgašova, N., Kutejová, E. and Timko, J.: Purification and some characteristics of the acetylglucosylase from *Schizopyllum commune*. *Biochem. J.*, 298 (1994) 751-755.
- 3571 Hallmann, A. and Sumper, M.: An inducible arylsulfatase of *Volvox carteri* with properties suitable for a reporter-gene system. *Eur. J. Biochem.*, 221 (1994) 143-150.
- 3572 Harder, K.W., Owen, P., Wong, L.K.H., Aebersold, R., Clark-Lewis, I. and Jirik, F.R.: Characterization and kinetic analysis of the intracellular domain of human protein tyrosine phosphatase β (HPTP β) using synthetic phosphopeptides. *Biochem. J.*, 298 (1994) 295-401.
- 3573 Itoh, R.: Purification and some properties of an IMP-specific 5'-nucleotidase from yeast. *Biochem. J.*, 298 (1994) 593-598.
- 3574 James, S.R., Demel, R.A. and Downes, C.P.: Interfacial hydrolysis of phosphatidylinositol 4-phosphate and phosphatidylinositol 4,5-bisphosphate by turkey erythrocyte phospholipase C. *Biochem. J.*, 298 (1994) 499-506.
- 3575 Lee, M.W. and Severson, D.L.: Partial purification of a diacylglycerol lipase from bovine aorta. *Biochem. J.*, 298 (1994) 213-219.
- 3576 Maga, G., Spadari, S., Wright, G.E. and Focher, F.: Identification, partial purification and inhibition by guanine analogues of a novel enzymic activity which phosphorylates guanosine to GMP in the protozoan parasite *Eimeria tenella*. *Biochem. J.*, 298 (1994) 289-294.
- 3577 Manco, G., di Gennaro, S., de Rosa, M. and Rossi, M.: Purification and characterization of a thermostable carboxylesterase from the thermoacidophilic *Bacillus acidocaldarius*. *Eur. J. Biochem.*, 221 (1993) 965-972.
- 3578 Marshall, P., Davis, T.B. and Lemieux, C.: The I-Ceul endonuclease: purification and potential role in the evolution of *Chlamydomonas* group I introns. *Eur. J. Biochem.*, 220 (1994) 855-859.
- 3579 Masuno, H. and Okuda, H.: Role of processing of the oligosaccharide chains in the affinity of lipoprotein lipase for heparin. *Biochim. Biophys. Acta*, 1212 (1994) 125-128.
- 3580 Matvienko, N.N., Kramarov, V.M., Zheleznyaya, L.A. and Matvienko, N.I.: (New site-specific endonuclease and methylase from *Bacillus licheniformis* 736). *Biokhimiya (Moscow)*, 58 (1993) 1139-1153.
- 3581 Murakami, T., Kobayashi, T., Terasawa, T., Ohnishi, M., Kato, S., Sasahara, Y., Itoh, M., Nakano, T. and Tamura, S.: Characterization of multiple molecular forms of Mg²⁺-dependent protein phosphatase from *Saccharomyces cerevisiae*. *J. Biochem. (Tokyo)*, 115 (1994) 762-766.
- 3582 Okazaki, T., Bielawska, A., Domae, N., Bell, R.M. and Hannun, Y.A.: Characteristics and partial purification of a novel cytosolic, magnesium-independent, neutral sphingomyelinase activated in the early signal transduction of $1\alpha,25$ -dihydroxyvitamin D₃-induced HL-60 cell differentiation. *J. Biol. Chem.*, 269 (1994) 4070-4077.
- 3583 Palmer, F.B.S.C., Théolis, R., Jr., Cook, H.W. and Byers, D.M.: Purification of two immunologically related phosphatidylinositol 4(4,5)-bisphosphate phosphatases from bovine brain cytosol. *J. Biol. Chem.*, 269 (1994) 3403-3410.
- 3584 Sugimoto, H. and Yamashita, S.: Purification, characterization, and inhibition by phosphatidic acid of lysophospholipase transacylase from rat liver. *J. Biol. Chem.*, 269 (1994) 6252-6258.

- 3585 Yamada, J., Matsumoto, I., Furihata, T., Sakuma, M. and Suga, T.: Purification and properties of long-chain acyl-CoA hydrolases from the liver cytosol of rats treated with peroxisome proliferator. *Arch. Biochem. Biophys.*, 308 (1994) 118-125.
- 3586 Zhang, Z., Zhao, S., Ziratto, S.D., Bai, G. and Lee, E.Y.C.: Expression of recombinant inhibitor-2 in *E. coli* and its utilization for the affinity chromatography of protein phosphatase-1. *Arch. Biochem. Biophys.*, 308 (1994) 37-41.
- For additional information see C.A.:
120 (1994) 100127m.
- See also 3107, 3344.
- 20e. *Hydrolases, acting on glycosyl compounds (E.C. 3.2.-.-)*
- 3587 Brochu, G., Shah, G.M. and Poirier, G.G.: Purification of poly(ADP-ribose) glycohydrolase and detection of its isoforms by a zymogram following one- or two-dimensional electrophoresis. *Anal. Biochem.*, 218 (1994) 265-272.
- 3588 Chen, X.-G., Nagai, T. and Yamada, H.: Sialidase in rabbit blood. Characterization of sialidase purified from rabbit erythrocyte membrane. *Eur. J. Biochem.*, 221 (1994) 655-664.
- 3589 Dalmia, B.K. and Nikolov, Z.L.: Characterization of a β -galactosidase fusion protein containing the starch-binding domain of *Aspergillus glucoamylase*. *Enzyme Microb. Technol.*, 16 (1994) 18-23; C.A., 120 (1994) 100910m.
- 3590 Hiraizumi, S., Spohr, U. and Spiro, R.G.: Ligand affinity chromatographic purification of rat liver Golgi endomannosidase. *J. Biol. Chem.*, 269 (1994) 4697-4700.
- 3591 Ito, K., Matura, Y. and Minamiura, N.: Purification and characterization of fungal nuclease composed of heterogeneous subunits. *Arch. Biochem. Biophys.*, 309 (1994) 160-167.
- 3592 Krebs, S.L. and Grumet, R.: Affinity purification and characterization of a β -1,3-glucanase from celery. *Plant Sci. (Limerick)*, 93 (1993) 31-39; C.A., 120 (1994) 185866z.
- 3593 Lerouge, P., O'Neill, M.A., Darvill, A.G. and Albersheim, P.: The purification of commercially available endo- α -L-arabinanases and α -L-arabinosidase for use in the structural analysis of pectic polysaccharides. *Carbohydr. Res.*, 243 (1993) 373-378; C.A., 120 (1994) 72841u.
- 3594 Mosolova, T.P., Kalyuzhny, S.V. and Velikodvorskaya, G.A.: (Purification and some properties of *Clostridium thermocellum* endoglucanase formed by a recombinant strain of *E. coli*). *Biochimiya (Moscow)*, 58 (1993) 1213-1220.
- 3595 Myrnes, B. and Johansen, A.: Recovery of lysozyme from scallop waste. *Prepar. Biochem.*, 24 (1994) 69-80.
- 3596 Nidetzky, B., Steiner, W., Hayn, M. and Cleayssens, M.: Cellulose hydrolysis by the cellulases from *Trichoderma reesei*: a new model for synergistic interaction. *Biochem. J.*, 298 (1994) 705-710.
- 3597 Pazos, C., Franco-Fraguas, L. and Batista-Viera, F.: Salt promoted adsorption chromatography of malted barley amylases. *Chromatographia*, 38 (1994) 232-234.
- 3598 Takematsu, H., Kawano, T., Koyama, S., Kozutsumi, Y., Suzuki, A. and Kawasaki, T.: Reaction mechanism underlying CMP-N-acetylneuraminic acid hydroxylation in mouse liver: formation of a ternary complex of cytochrome *b*₅, CMP-N-acetylneuraminic acid, and a hydroxylation enzyme. *J. Biochem. (Tokyo)*, 115 (1994) 381-386.
- 3599 Xia, J. and Dubin, P.L.: Chromatographic evaluation of the binding of lysozyme to poly(dimethylallylammonium chloride). *J. Chromatogr. A*, 667 (1994) 311-315.
- 3600 Yamauchi, J. and Tanuma, S.-i.: Occurrence of an NAD⁺ glycohydrolase in bovine brain cytosol. *Arch. Biochem. Biophys.*, 308 (1994) 327-329.
- For additional information see C.A.:
120 (1994) 48932d.
- 20f. *Other hydrolases*
- 3601 Arbona, J.R. and Koohmaria, M.: Technical note: quantification of multicatalytic proteinase complex (proteasome) activity by ion-exchange chromatography. *J. Anim. Sci.*, 71 (1993) 3301-3306; C.A., 120 (1994) 48548h.
- 3602 Betageri, R., Hopkins, J.L., Thibeault, D., Emmanuel, M.J., Chow, G.C., Skoog, M.T., de Dreu, P. and Cohen, K.A.: Rapid, sensitive and efficient HPLC assays for HIV-1 proteinase. *J. Biochem. Biophys. Methods*, 27 (1993) 191-197; C.A., 120 (1994) 72020a.
- 3603 Büchler, M., Tislijar, U. and Wolf, D.H.: Proteinase yscD (oligopeptidase yscD). Structure, function and relationship of the yeast enzyme with mammalian thimet oligopeptidase (metalloendopeptidase, EP 24.15). *Eur. J. Biochem.*, 219 (1994) 627-639.
- 3604 Ejler, G., Söderström, K. and Karlström, A.: Inactivation of thrombin by a complex between rat mast-cell protease 1 and heparin proteoglycan. *Biochem. J.*, 299 (1994) 507-513.
- 3605 Figueiredo, E. and Duque-Magalhaes, M.C.: Identification, purification and partial characterization of a carboxypeptidase from the matrix of rat liver mitochondria: a novel metalloenzyme. *Biochem. J.*, 300 (1994) 15-19.
- 3606 Ibrahim-Granet, O., Bertrand, O., Debeauvais, J.P., Planche-nault, T., Diaquin, M. and Dupont, B.: *Aspergillus fumigatus* metalloproteinase that hydrolyzes native collagen: purification by dye-binding chromatography. *Protein Expression Purif.*, 5 (1994) 84-88; C.A., 120 (1994) 238797m.
- 3607 Inokuchi, T., Kobayashi, K.-i. and Horiuchi, S.: Purification and characterization of cathepsin E type acid proteinase from gastric mucosa of bullfrog, *Rana catesbeiana*. *J. Biochem. (Tokyo)*, 115 (1994) 76-81.
- 3608 Kikugawa, M., Kaneko, M., Fujimoto-Sakata, S., Maeda, M., Kawasaki, K., Takagi, T. and Tamaki, N.: Purification, characterization and inhibition of dihydropyrimidinase from rat liver. *Eur. J. Biochem.*, 219 (1994) 393-399.
- 3609 Kitadokoro, K., Tsuzuki, H., Nakamura, E., Sato, T. and Teraoka, H.: Purification, characterization, primary structure, crystallization and preliminary crystallographic study of a serine proteinase from *Streptomyces fradiae* ATCC 14544. *Eur. J. Biochem.*, 220 (1994) 55-61.
- 3610 Lamango, N.S. and Isaac, R.E.: Identification and properties of a peptidyl dipeptidase in the house-fly, *Musca domestica*, that resembles mammalian angiotensin-converting enzyme. *Biochem. J.*, 299 (1994) 651-657.
- 3611 O, I. and Stitt, B.L.: 8-Azido-ATP inactivation of *Escherichia coli* transcription termination factor Rho. Modification of one subunit inactivates the hexamer. *J. Biol. Chem.*, 269 (1994) 5009-5015.

- 3612 Ohtsuki, S., Homma, K.-i., Kurata, S., Komano, H. and Natori, S.: A prolyl endopeptidase of *Sarcophaga peregrina* (flesh fly): its purification and suggestion for its participation in the differentiation of the imaginal discs. *J. Biochem. (Tokyo)*, 115 (1994) 449-453.
- 3613 Pfeiffer, G., Strube, K.-H., Schnidt, M. and Geyer, R.: Glycosylation of two recombinant human uterine tissue plasminogen activator variants carrying an additional N-glycosylation site in the epidermal growth-factor-like domain. *Eur. J. Biochem.*, 219 (1994) 331-348.
- 3614 Seki, M., Yanagisawa, J., Kohda, T., Sonoyama, T., Ui, M. and Enomoto, T.: Purification of two DNA-dependent adenosinetriphosphatases having DNA helicase activity from HeLa cells and comparison of the properties of the two enzymes. *J. Biochem. (Tokyo)*, 115 (1994) 523-531.
- 3615 Sugahara, K., Jianying, Z., Yamamoto, Y., Yasuda, K., Kodama, H. and Kodama, H.: Measurement of iminodipeptides in the serum of patients with prolidase deficiency using liquid chromatography-mass spectrometry. *Eur. J. Clin. Chem. Clin. Biochem.*, 32 (1994) 113-117.
- 3616 Tran, C.M., Huston, E.E. and Farley, R.A.: Photochemical labeling and inhibition of Na,K-ATPase by 2-azido-ATP. Identification of an amino acid located within the ATP binding site. *J. Biol. Chem.*, 269 (1994) 6558-6565.
- 3617 Tsuji, A., Akamatsu, T., Nagamune, H. and Matsuda, Y.: Identification of targeting proteinase for rat α_1 -macroglobulin *in vivo*. Mast-cell tryptase is a major component of the α_1 -macroglobulin-proteinase complex endocytosed into rat liver lysosomes. *Biochem. J.*, 298 (1994) 79-85.
- 3618 Waxman, L., Doshi, K.P., Gaul, S.L., Wang, S., Bednar, R.A. and Stern, A.M.: Identification and characterization of endothelin converting activity from EAHY 926 cells: evidence for the physiologically relevant human enzyme. *Arch. Biochem. Biophys.*, 308 (1994) 240-253.
- For additional information see C.A.:
120 (1994) 264281v.
- See also 2951, 3311, 3349, 3353, 3364.
- 20g. *Lyases*
- 3619 Hansson, M. and Hederstedt, L.: Purification and characterization of a water-soluble ferrocyclase from *Bacillus subtilis*. *Eur. J. Biochem.*, 220 (1994) 201-208.
- 3620 Jones, R.M. and Jordan, P.M.: Purification and properties of porphobilinogen deaminase from *Arabidopsis thaliana*. *Biochem. J.*, 299 (1994) 895-902.
- 3621 Nathan, B., Hsu, C. and Bao, J., Wu, R. and Wu, J.: Purification and characterization of a novel form of brain L-glutamate decarboxylase. A Ca^{2+} -dependent peripheral membrane protein. *J. Biol. Chem.*, 269 (1994) 7249-7254.
- 3622 Palumbo, A., d'Ischia, M., Misuraca, G., de Martino, L. and Prota, G.: A new dopachrome-rearranging enzyme from the ejected ink of the cuttlefish *Sepia officinalis*. *Biochem. J.*, 299 (1994) 839-844.
- 3623 Pecina, A. and Paneque, A.: Detection of alginate lyase by activity staining after sodium dodecyl sulfate-polyacrylamide gel electrophoresis and subsequent renaturation. *Anal. Biochem.*, 217 (1994) 124-127.
- 3624 Woodcock, S.C. and Jordan, P.M.: Evidence for participation of aspartate-84 as a catalytic group at the active site of porphobilinogen deaminase obtained by site-directed mutagenesis of the hemC gene from *Escherichia coli*. *Biochemistry*, 33 (1994) 2688-2695.
- 20h. *Isomerases*
- 3625 Khac, S.B.-P. and Moreau, N.J.: Interactions between fluoroquinolones, Mg^{2+} , DNA and DNA gyrase, studied by phase partitioning in an aqueous two-phase system and by affinity chromatography. *J. Chromatogr. A*, 668 (1994) 241-247.
- 3626 Kohl, L., Callens, M., Wierenga, R.K., Opperdoes, F.R. and Michels, P.A.M.: Triose-phosphate isomerase of *Leishmania mexicana mexicana*. Cloning and characterization of the gene, overexpression in *Escherichia coli* and analysis of the protein. *Eur. J. Biochem.*, 220 (1994) 331-338.
- 3627 Luo, M.J., Smeland, T.E., Shoukry, K. and Schulz, H.: $\Delta^{3,5}, \Delta^{2,4}$ -Dienoyl-CoA isomerase from rat liver mitochondria. Purification and characterization of a new enzyme involved in the β -oxidation of unsaturated fatty acids. *J. Biol. Chem.*, 269 (1994) 2384-2388.
- 3628 Proudfoot, A.E.I., Turcatti, G., Wells, T.N.C., Payton, M.A. and Smith, D.J.: Purification, cDNA cloning and heterologous expression of human phosphomannose isomerase. *Eur. J. Biochem.*, 219 (1994) 415-423.
- 3629 Slesarev, A.I., Lake, J.A., Stetter, K.O., Gellert, M. and Kozyavkin, S.A.: Purification and characterization of DNA topoisomerase V. An enzyme from the hyperthermophilic prokaryote *Methanopyrus kandleri* that resembles eukaryotic topoisomerase I. *J. Biol. Chem.*, 269 (1994) 3295-3303.
- 3630 Störl, K., Störl, J., Roth, M. and Zimmer, C.: DNA topoisomerases from *Streptomyces noursei*: Influence of coumarins and quinolones on the enzymic activity. *Biochim. Biophys. Acta*, 1199 (1994) 143-148.
- 20i. *Ligases*
- 3631 Diez, T.A., Wurtele, E.S. and Nikolau, B.J.: Purification and characterization of 3-methylcrotonyl-coenzyme-A carboxylase from leaves of *Zea mays*. *Arch. Biochem. Biophys.*, 310 (1994) 64-75.
- 3632 Dowton, M. and Kennedy, I.R.: Purification of glutamine synthetase by adenosine-affinity chromatography. *J. Chromatogr. A*, 664 (1994) 280-283.
- 3633 Kerjan, P., Cerini, C., Sémériva, M. and Mirande, M.: The multi-enzyme complex containing nine aminoacyl-tRNA synthetases is ubiquitous from *Drosophila* to mammals. *Biochim. Biophys. Acta*, 1199 (1994) 293-297.
- 3634 Nakagawa, C.W., Mutoh, N. and Hayashi, Y.: Glutathione synthetase from the fission yeast. Purification and its unique heteromeric subunit structure. *Biochem. Cell Biol.*, 71 (1993) 447-453.
- 3635 Paul, G.: Activity and inhibition of acetyl-CoA carboxylase determined by HPLC. In: Boeger, P. and Sandmann, G. (Editors), *Target Assays Mod. Herbic. Relat. Phytotoxic Compd.*, Lewis, Boca Raton, 1993, pp. 161-166; C.A., 120 (1994) 185748n.

20j. Complex mixtures and incompletely identified enzymes

- 3636 Corradini, C., Panari, G., Rampilli, M. and Barzaghi, S.: (Evaluation of the official method of analysis for the identification of chymosins obtained from genetically modified microorganisms). *Sci. Tec. Latt.-Casearia*, 44 (1993) 195-204; *C.A.*, 120 (1994) 75786r.
- 3637 Leikin, A. and Shinitzky, M.: Shedding and isolation of the Δ^6 -desaturase system from rat liver microsomes by application of high hydrostatic pressure. *Biochim. Biophys. Acta*, 1211 (1994) 150-155.
- 3638 Stepnaya, O.A., Ledova, L.A. and Kulaev, I.S.: (The bacteriolytic enzymatic complex 'lysoamidase'. I. Estimation of the mode of the enzyme-polysaccharide interaction within the complex). *Bio-khimiya (Moscow)*, 58 (1993) 1523-1528.
- 3639 Zuurbier, K.W.M., Fung, S.Y., Scheffer, J.J.C. and Verpoorte, R.: Assay of chalcone synthase activity by high-performance liquid chromatography. *Phytochemistry*, 34 (1993) 1225-1229; *C.A.*, 120 (1994) 48547g.

See also 3405.

21. PURINES, PYRIMIDINES, NUCLEIC ACIDS AND THEIR CONSTITUENTS

21a. Purines, pyrimidines, nucleosides, nucleotides

- 3640 Cross, D.R., Miller, B.J. and James, S.J.: A simplified HPLC method for simultaneously quantifying ribonucleotides and deoxyribonucleotides in cell extracts of frozen tissues. *Cell Proliferation*, 26 (1993) 327-336; *C.A.*, 120 (1994) 157790b.
- 3641 Demidov, V.V. and Potaman, V.N.: HPLC photofingerprinting of conformational peculiarities and transitions in oligonucleotide duplexes. *Nucleic Acids Res.*, 21 (1993) 2691-2696; *C.A.*, 120 (1994) 49189d.
- 3642 Desnoyers, S., Shah, G.M., Brochu, G. and Poirier, G.G.: Erasable blot of poly(ADP-ribose) polymerase. *Anal. Biochem.*, 218 (1994) 470-473.
- 3643 Ersoz, M., Yildiz, S. and Pehlivan, E.: Ligand-exchange chromatography and kinetics of nucleosides and nucleic acid bases on immobilized copper-, nickel-, and cobalt-bis-diaminoethyl-gluoximated (bDAEG)-sporopollenin. *Spec. Publ.-R. Soc. Chem.*, 122(Ion Exchange Processes: Advances and Application) (1993) 305-313; *C.A.*, 120 (1994) 128743h.
- 3644 Gibson, J.B., Reynolds, R.A., Rogers, S., Palmieri, M.J. and Segal, S.: Uridine diphosphoglucose content of human erythrocytes: assessment by conversion to uridine diphosphoglucuronate. *J. Pediatr. (St. Louis)*, 123 (1993) 906-914; *C.A.*, 120 (1994) 264933x.
- 3645 Hirayama, T., Katayama, Y. and Watanabe, T.: (High-performance liquid chromatographic determination of O⁶-methyldeoxyguanosine as its phenylethano derivative in the reaction of methylating agents and 2'-deoxyguanosine). *Jpn. J. Toxicol. Environ. Health*, 39 (1993) 324-331; *C.A.*, 120 (1994) 70976t.
- 3646 Inaki, Y., Ohsaki, Y., Wang, H., Matsukawa, K., Mochizuki, E. and Wada, T.: (High-performance liquid chromatographic separation of oligonucleotides using deoxyadenosine immobilized silica gels for the stationary phase. Effect of the deoxyadenosine ligand concentration). *Bunseki Kagaku*, 42 (1993) 825-830; *C.A.*, 120 (1994) 128761n.
- 3647 Jault, J.-M., Kaibara, C., Yoshida, M., Garrod, S. and Allison, W.S.: Probing the specificity of nucleotide binding to the F₁-ATPase from thermophilic *Bacillus* PS3 and its isolated α and β subunits with 2-N₃-[α , β -³²P]ATP. *Arch. Biochem. Biophys.*, 310 (1994) 282-288.
- 3648 Jochheim, C., Janning, P., Marggraf, U., Löffler, T.M., Hasse, F. and Linscheid, N.: A procedure for the determination of 5-fluorouracil in tissue using microbore HPLC and fluorescence detection. *Anal. Biochem.*, 217 (1994) 285-291.
- 3649 MacLeod, M.C., Evans, F.E., Lay, J., Chiarelli, P., Geacintov, N.E., Powell, K.L., Daylong, A., Luna, E. and Harvey, R.G.: Identification of a novel, N7-deoxyguanosine adduct as the major DNA adduct formed by a non-bay-region diol epoxide of benzo[a]pyrene with low mutagenic potential. *Biochemistry*, 33 (1994) 2977-2987.
- 3650 Murray, J.B., Collier, A.K. and Arnold, J.R.P.: A general purification procedure for chemically synthesized oligoribonucleotides. *Anal. Biochem.*, 218 (1994) 177-184.
- 3651 Ohba, Y., Kai, M., Nohta, H. and Ohkura, Y.: Alkoxyphenylglyoxals as fluorogenic reagents selective for guanine and its nucleosides and nucleotides in liquid chromatography. *Anal. Chim. Acta*, 287 (1994) 215-221.
- 3652 Palmer, S. and Cox, S.: Comparison of extraction procedures for high-performance liquid chromatographic determination of cellular deoxynucleotides. *J. Chromatogr. A*, 667 (1994) 316-321.
- 3653 Pehlivan, E. and Yildiz, S.: Column chromatography and kinetics of nucleosides and nucleic acid bases on immobilized nickel- and cobalt-CDAE-sporopollenin. *Sep. Sci. Technol.*, 29 (1994) 887-895; *C.A.*, 120 (1994) 100922s.
- 3654 Ralston, S.L., Lau, H.H.S., Seidel, A., Luc, A., Platt, K.L. and Baird, W.M.: The potent carcinogen dibenzo[a,f]pyrene is metabolically activated to fjord-region 11,12-diol 13,14-epoxides in human mammary carcinoma MCF-7 cell cultures. *Cancer Res.*, 54 (1994) 887-890.
- 3655 Sonoki, S., Sanda, A. and Hisamatsu, S.: Simultaneous determination of mono-, di-, and trinucleotides by high-performance liquid chromatography using N-(dansyl)ethylenediamine as a fluorescent derivatizing reagent. *J. Liq. Chromatogr.*, 17 (1994) 1057-1064.
- 3656 Takeda, N., Nakamura, M., Yoshizumi, H. and Tatematsu, A.: Frit-fast atom bombardment liquid chromatography/mass spectrometry, a useful method for sequencing dinucleotides. *Anal. Biochem.*, 217 (1994) 155-157.
- 3657 Traoré, F., Fente, C., Prognon, P. and Mahuzier, G.: Luminarin 3 as a derivatization reagent for the liquid chromatographic determination of cytidine, adenosine and related nucleotides with fluorimetric detection. *Anal. Chim. Acta*, 290 (1994) 94-102.
- 3658 Westmark, P.R., Valencia, L.S. and Smith, B.D.: Influence of eluent anions in boronate affinity chromatography. *J. Chromatogr. A*, 664 (1994) 123-128.

- 3659 Ye, M.Y. and Shen, Y.: Electrospray ionization mass spectrometry and HPLC determination of the products in the radiolysis of 5-bromouracil, its nucleoside and nucleotide derivatives. *J. Liq. Chromatogr.*, 17 (1994) 773-791.
- 3660 Yonekura, S., Iwasaki, M., Kai, M. and Ohkura, Y.: Determination of guanine and its nucleosides and nucleotides in human erythrocytes by high-performance liquid chromatography with postcolumn fluorescence derivatization using phenylglyoxal reagent. *J. Chromatogr. B*, 654 (1994) 19-24.
- 3661 Yonekura, S., Iwasaki, M., Kai, M. and Ohkura, Y.: High-performance liquid chromatographic determination of guanine and its nucleosides and nucleotides in biospecimens with precolumn fluorescence derivation using phenylglyoxal. *Anal. Sci.*, 10 (1994) 247-251; *C.A.*, 120 (1994) 264970g.
- For additional information see *C.A.*:
120 (1994) 49199g, 49419d, 239334v.
- See also 3282, 3490, 3667, 3671, 3681, 3684, 3697, 3957, 3980.
- 21b. *Nucleic acids, RNA*
- 3662 Blechschidt, B., Shirokov, V. and Sprinzl, M.: Undecagold cluster modified tRNA^{Phe} from *Escherichia coli* and its activity in the protein elongation cycle. *Eur. J. Biochem.*, 219 (1994) 65-71.
- 3663 Moriya, J., Yokogawa, T., Wakita, K., Ueda, T., Nishikawa, K., Crain, P.F., Hashizume, T., Pomerantz, S.C., McCloskey, J.A., Kawai, G., Hayashi, N. et al.: A novel modified nucleoside found at the first position of the anticodon of methionine tRNA from bovine liver mitochondria. *Biochemistry*, 33 (1994) 2234-2239.
- 3664 Poot, R.A., Brink, M.F., Pleij, C.W.A., de Boer, H.A. and van Duin, J.: Separation of mutant and wild-type ribosomes based on differences in their anti Shine-Dalgarno sequence. *Nucleic Acids Res.*, 21 (1993) 5398-5402; *C.A.*, 120 (1994) 128745k.
- 3665 Rao, A. and Steward, O.: Evaluation of RNAs present in synaptodendrosomes: Dendritic, glial, and neuronal cell body contribution. *J. Neurochem.*, 61 (1993) 835-844; *C.A.*, 120 (1994) 101129a.
- 3666 Schott, H.: Nucleic acid and its derivatives. *Chromatogr. Sci. Ser.*, 63(Handbook of Affinity Chromatography) (1993) 177-216; *C.A.*, 120 (1994) 128675n - a review with 318 refs.
- See also 3422.
- 21c. *Nucleic acids, DNA*
- 3667 Azadnia, A., Campbell, R. and Sharma, M.: The scope of dansyl vs fluorescein label in fluorescence postlabeling assay for DNA damage. *Anal. Biochem.*, 218 (1994) 444-448.
- 3668 Baird, W.M., Lau, H.H.S., Schmerold, I., Coffing, S.L., Brozich, S.L., Lee, H. and Harvey, R.G.: Analysis of polycyclic aromatic hydrocarbon-DNA adducts by postlabeling with the weak β -emitters ³⁵S-phosphorothioate and ³²P-phosphate, immobilized boronate chromatography and high-performance liquid chromatography. *IARC Sci. Publ.*, 124(Postlabelling Methods for Detection of DNA Adducts) (1993) 217-226; *C.A.*, 120 (1994) 127022k.
- 3669 Cole, A.D. and Kmiec, E.B.: ATP-independent DNA renaturation catalyzed by a protein from *Ustilago maydis*. *Eur. J. Biochem.*, 220 (1994) 75-82.
- 3670 Kmiec, E.B. and Holloman, W.K.: ATP-dependent DNA renaturation and DNA-dependent ATPase reactions catalyzed by the *Ustilago maydis* homologous pairing protein. *Eur. J. Biochem.*, 219 (1994) 865-875.
- 3671 Takenaka, S., Uto, Y., Kondo, H., Ihara, T. and Takagi, M.: Electrochemically active DNA probes: detection of target DNA sequences at femtomole level by high-performance liquid chromatography with electrochemical detection. *Anal. Biochem.*, 218 (1994) 436-443.
- For additional information see *C.A.*:
120 (1994) 49472r.
- See also 3666.
- 21d. *Structural studies on RNA and RNA mapping*
- 3672 Tanigawa, M., Mukaiyama, N., Shimokubo, S., Wakabayashi, K., Fujita, Y., Fukudome, K. and Yamaoka, K.: Molecular weight distribution and intrinsic viscosity of sonicated and successively fractionated double-stranded deoxyribonucleic acid and polyribonucleotides. *Polym. J. (Tokyo)*, 26 (1994) 291-302; *C.A.*, 120 (1994) 164962f.
- See also 3663.
- 21e. *Structural studies on DNA and DNA mapping*
- 3673 Joshi, R.R., Likhite, S.M., Kumar, R.K. and Ganesh, K.N.: DNA cleavage by Cu(II)-desferal: identification of C1'-hydroxylation as the initial event for DNA damage. *Biochim. Biophys. Acta*, 1199 (1994) 285-292.
- 3674 Tawa, R., Tamura, G., Sakurai, H., Ono, T. and Kurishita, A.: High-performance liquid chromatographic analysis of methylation changes of CCGG sequence in brain and liver DNA of mice during pre- and postnatal development. *J. Chromatogr. B*, 653 (1994) 211-216.
- For additional information see *C.A.*:
120 (1994) 128751j, 263135b.
- 21f. *Complex mixtures of nucleic acids and their fragments*
- 3675 Wolf, S.M. and Vouros, P.: Application of capillary liquid chromatography coupled with tandem mass spectrometric methods to the rapid screening of adducts formed by the reaction of N-acetoxy-N-acetyl-2-aminofluorene with calf DNA. *Chem. Res. Toxicol.*, 7 (1994) 82-88; *C.A.*, 120 (1994) 98557f.
22. ALKALOIDS
- 3676 Freytag, W.E. and Stapf, W.: (Greater celandine-*Chelidonium majus* L. Quantitative HPLC determination of Chelidonium alkaloids). *P2 Wiss.*, 6 (1993) 126-133; *C.A.*, 120 (1994) 62415u.
- 3677 Huang, J.L. and Morgan, D.J.: Influence of perfusion flow rate on uptake and pharmacodynamics of quinidine in isolated perfused rat heart. *J. Pharm. Sci.*, 83 (1994) 119-123.

- 3678 Iwase, H., Gondo, K., Koike, T. and Ono, I.: Novel precolumn deproteinization method using a hydroxyapatite cartridge for the determination of theophylline and diazepam in human plasma by high-performance liquid chromatography with ultraviolet detection. *J. Chromatogr. B*, 655 (1994) 73-81.
- 3679 Lee, S.-S.: Application of centrifugal partition chromatography to the separation of Lauraceous alkaloids. *J. Chromatogr. A*, 667 (1994) 322-326.
- 3680 Leveque, D., Jehl, F., Quoix, E. and Monteil, H.: Pharmacokinetic profile of vinorelbine, a new semi-synthetic vinca alkaloid, determined by high-performance liquid chromatography. *Xenobiotica*, 23 (1993) 1325-1333; *C.A.*, 120 (1994) 124017b.
- 3681 Pabst, G., Weber, W., Müller, M. and Barkworth, M.F.: Study on the influence of food on the absorption of theophylline from a controlled-release preparation. *Arzneim.-Forsch.*, 44 (1994) 333-337.
- 3682 Park, M.K., Park, J.H., Kim, B.Y., Kim, J.M., Liem, K.J. and Han, B.H.: (Analysis of alkaloids in *Polygala tenuifolia* by HPLC). *Anal. Sci. Technol.*, 6 (1993) 255-259; *C.A.*, 120 (1994) 173583s.
- 3683 Samo, A.R., Arbani, S.A., Khahawer, M.Y., Chippa, M.A. and Qureshi, G.A.: The simultaneous analysis of theophylline, caffeine and their metabolites in human plasma by HPLC. *J. Chem. Soc. Pak.*, 15 (1993) 182-186; *C.A.*, 120 (1994) 94596v.
- 3684 Sun, P., Mariano, G.J., Barker, G. and Hartwick, R.A.: Comparison of micellar electrokinetic capillary chromatography and high-performance liquid chromatography on the separation and determination of caffeine and its analogues in pharmaceutical tablets. *Anal. Lett.*, 27 (1994) 927-937.
- 3685 Watt, A.P., Verrier, H.M. and O'Connor, D.: Resolution of synthetic (+)- and (-)-epibatidine by chiral high performance liquid chromatography and identification of the natural enantiomer. *J. Liq. Chromatogr.*, 17 (1994) 1257-1264.
- For additional information see *C.A.*:
120 (1994) 62388n, 62390g, 182132w, 237651x.
- See also 3795, 3982, 4011, 4022.
23. OTHER SUBSTANCES CONTAINING HETEROCYCLIC NITROGEN
- 23a. *Porphyryns and other pyrroles*
- 3686 Beukenveld, G.J.J., Meerman, L., Huizenga, J.R., Venekamp-Hoolsema, E.E.A., Gips, C.H. and Wolthers, B.G.: Determination of porphyrins in bile using high performance liquid chromatography and some chemical applications. *Eur. J. Clin. Chem. Clin. Biochem.*, 32 (1994) 153-159.
- 3687 Ho, J.W. and Candy, L.Y.F.: Effects of injection solvents on the separation of porphyrin and metalloporphyrin in reversed-phase liquid chromatography. *J. Liq. Chromatogr.*, 17 (1994) 549-558.
- 3688 Mank, A.J.G., Gooijer, C., Lingeman, H., Velthorst, N.H. and Brinkman, U.A.T.: Selective and sensitive *in vitro* detection method for aluminum phthalocyanine photosensitizers, using liquid chromatography and diode-laser induced fluorescence. *Anal. Chim. Acta*, 290 (1994) 103-120.
- 3689 Schoenfeld, N. and Mamet, R.: Interference of ofloxacin with determination of urinary porphyrins. *Clin. Chem. (Washington)*, 40 (1994) 417-419.
- 3690 Wu, W. and Stalcup, A.: Separation of porphyrins using a γ -cyclodextrin stationary phase. *J. Liq. Chromatogr.*, 17 (1994) 1111-1124.
- For additional information see *C.A.*:
120 (1994) 152374j, 186391c.
- See also 3477, 3824.
- 23b. *Bile pigments*
- For additional information see *C.A.*:
120 (1994) 62405r.
- 23c. *Indole derivatives and plant hormones (gibberelins)*
- 3691 Cheng, F.-C. and Kuo, J.-S.: Plasma serotonin. Reply to E.F. Marshall and M. Leitch. *J. Chromatogr. B*, 655 (1994) 169-170.
- 3692 Horiuchi, K., Yonekawa, O., Iwahara, K., Kanno, T., Kurihara, T. and Fujise, Y.: A hydrophilic tetrahydro- β -carboline in human urine. *J. Biochem. (Tokyo)*, 115 (1994) 362-366.
- 3693 Marshall, E.F. and Leitch, M.: Plasma serotonin. *J. Chromatogr. B*, 655 (1994) 167-168.
- 3694 Sherwood, R. and Keating, J.: Plasma serotonin. Reply to E.F. Marshall and M. Leitch. *J. Chromatogr. B*, 655 (1994) 171.
- For additional information see *C.A.*:
120 (1994) 49229s.
- See also 3241, 3245.
- 23d. *Pyridine derivatives*
- 3695 Angeli, E., Carnelli, L. and Pistis, A.: (Quantitative determination by HPLC of free and added nicotinic acid in fresh sausages). *Boll. Chim. Ig., Parte Sci.*, 44 (1993) 17-23; *C.A.*, 120 (1994) 132522c.
- 3696 Fontaine, J. and Hoerr, J.: Determination of supplemented nicotinic acid or nicotinamide in compound feed by HPLC after clean-up of extracts with cationic sorbent extraction. *Agrobiol. Res.*, 46 (1993) 10-19; *C.A.*, 120 (1994) 52945c.
- For additional information see *C.A.*:
120 (1994) 186392d.
- See also 2993, 4049.
- 23e. *Other N-heterocyclic compounds*
- 3697 Eiteman, M.A., Gordillo, R.M. and Cabrera, M.L.: Analysis of oxonic acid, uric acid, creatinine, allantoin, xanthine and hypoxanthine in poultry litter by reversed phase HPLC. *Fresenius J. Anal. Chem.*, 348 (1994) 680-683.
- 3698 Visconti, A. and Solfrizzo, M.: Isolation, characterization and biological activity of visoltricin, a novel metabolite of *Fusarium tricinctum*. *J. Agric. Food Chem.*, 42 (1994) 195-199.

For additional information see C.A.:
120 (1994) 86585a, 94594t.

See also 3248, 3250, 3251.

24. ORGANIC SULPHUR COMPOUNDS (INCL. GLUCOSINOLATES)

- 3699 Allenmark, S. and Claeson, S.: Chromatographic resolution of a C₂-symmetric spiro-sulfurane: 1,1'-spirobi[3H-2,1-benzoxathio]-3,3'-dione. *Tetrahedron: Asymmetry*, 4 (1993) 2329-2332; C.A., 120 (1994) 217366e.
- 3700 Altomare, C., Carotti, A., Cellamare, S., Fanelli, F., Gasparrini, F., Villani, C., Carrupt, P.A. and Testa, B.: Enantiomeric resolution of sulfoxides on a DACH-DNB chiral stationary phase: a quantitative structure-enantioselective retention relationship (QSERR) study. *Chirality*, 5 (1993) 527-537; C.A., 120 (1994) 190640a.
- 3701 Bastian, B., Knepper, T.P., Hoffmann, P. and Ortner, H.M.: Determination of aromatic sulfonic acids in industrial waste water by ion-pair chromatography. *Fresenius J. Anal. Chem.*, 348 (1994) 674-679.
- 3702 Chan, C., Cole, E.R. and Southwell-Keely, P.T.: Sulfenamides and sulfenamides. VIII. The chromatography of aryl sulfenamides. *Phosphorus, Sulfur Silicon Relat. Elem.*, 80 (1993) 63-73; C.A., 120 (1994) 244003r.
- 3703 Diter, P., Taudien, S., Samuel, O. and Kagan, H.B.: Enantiomeric enrichment of sulfoxides by preparative flash chromatography on an achiral phase. *J. Org. Chem.*, 59 (1994) 370-373; C.A., 120 (1994) 76970h.
- 3704 Elfakir, C., Mercier, J.P. and Dreux, M.: Detection of some glucosinolates in anion exchange chromatography. Part II: qualitative study on a styrene-divinylbenzene copolymeric anion exchanger. *Chromatographia*, 38 (1994) 585-590.
- 3705 Hill, J.C., Flannery, G.M. and Fraser, B.A.: HPLC purification and separation of 5,5'-substituted-2,4-imidazolidinedithiones. *J. Chromatogr. A*, 664 (1994) 63-70.
- 3706 Küsters, E., Loux, V., Schmid, E. and Floersheim, P.: Enantiomeric separation of chiral sulphoxides. Screening of cellulose-based sorbents with particular reference to cellulose tribenzoate. *J. Chromatogr. A*, 666 (1994) 421-432.
- 3707 Lai, Y.-H. and Ang, S.-G.: Separation by permeation effect in high performance liquid chromatographic studies of selected dithia[3.3]cyclophanes and thia[3.2]cyclophanenes. *J. Liq. Chromatogr.*, 17 (1994) 1219-1230.
- 3708 Zou, H., Jia, Z., Zhang, Y. and Lu, P.: Separation of aqueous polythionates by reversed-phase ion-pair liquid chromatography with suppressor-conductivity detection. *Anal. Chim. Acta*, 284 (1993) 59-65.
- 3709 Zou, H., Zhang, Y., Hong, M. and Lu, P.: Retention values of sulphonic acids as a function of the nature and concentration of inorganic salt in reversed-phase ion-pair liquid chromatography. *J. Liq. Chromatogr.*, 17 (1994) 707-719.

For additional information see C.A.:
120 (1994) 152583b.

See also 2977, 3034.

25. ORGANIC PHOSPHORUS COMPOUNDS (INCL. SUGAR PHOSPHATES)

- 3710 Barker, C.J.: High performance liquid chromatographic separation of inositol phosphates. *IBRO Handab. Ser.*, 15(High Performance Liquid Chromatography in Neuroscience Research) (1993) 195-216; C.A., 120 (1994) 157691v - a review with 75 refs.
- 3711 Bird, I.M.: Analysis of cellular phosphoinositides and phosphoinositols by extraction and simple analytical procedures. *Methods Mol. Biol. (Totowa)*, 27(Biomembrane Protocols, II) (1994) 227-248; C.A., 120 (1994) 264945c.
- 3712 Bird, I.M.: Analysis of cellular phosphoinositides and phosphoinositols by high-performance liquid chromatography. *Methods Mol. Biol. (Totowa)*, 27(Biomembrane Protocols, II) (1994) 249-277; C.A., 120 (1994) 264946d.
- 3713 Boetker, H.E., Kimose, H.H., Helligsoe, P. and Nielsen, T.T.: Analytical evaluation of high energy phosphate determination by high performance liquid chromatography in myocardial tissue. *J. Mol. Cell. Cardiol.*, 26 (1994) 41-48; C.A., 120 (1994) 186428v.
- 3714 Gorbics, L., Urge, L., Lang, E., Szendrei, G.I. and Otvos, L., Jr.: Successful and rapid verification of the presence of a phosphate group in synthetic phosphopeptides using the conditions of standard DABS-Cl amino acid analysis. *J. Liq. Chromatogr.*, 17 (1994) 175-189.
- 3715 Kientz, C.E., Verweij, A., de Jong, G.J. and Brinkman, U.A.: Verification of nonproduction of chemical warfare agents: II. Large volume injections in microcolumn liquid chromatography using flame photometric detection. *J. Microcolumn Sep.*, 4 (1993) 477-483; C.A., 120 (1994) 155993w.
- 3716 Kientz, C.E., Verweij, A., de Jong, G.J. and Brinkman, U.A.T.: Verification of nonproduction of chemical warfare agents: I. Determination of organophosphorus compounds by microcolumn liquid chromatography with flame photometric or thermionic detection. *J. Microcolumn Sep.*, 4 (1993) 465-475; C.A., 120 (1994) 155992v.
- 3717 Lazar, S., Lafosse, M., Herbretreau, B. and Guillaumet, G.: (Synthetic inositol phosphate mixture analysis. Separation using gradient elution liquid chromatography and evaporative light scattering detection). *Analisis*, 21 (1993) 415-418; C.A., 120 (1994) 182104p.
- 3718 Parthasarathy, R., Parthasarathy, L. and Vadnal, R.E.: Identification of phosphatidylinositol trisphosphate in rat brain. *Methods Leurosci.*, 18(Lipid Metabolism in Signaling Systems) (1993) 113-124; C.A., 120 (1994) 265206z.
- 3719 Pfau, W., Lecoq, S., Hughes, N.C., Seidel, A., Platt, K.L., Grover, P.L. and Phillips, D.H.: Separation of ³²P-labelled nucleoside 3',5'-bisphosphate adducts by HPLC. *IARC Sci. Publ.*, 124(Postlabelling Methods for Detection of DNA Adducts) (1993) 233-242; C.A., 120 (1994) 98586q.
- 3720 Sasakawa, N., Nakaki, T. and Kato, R.: Characterization of inositol phosphates by high-performance liquid chromatography. *Methods Neurosci.*, 18(Lipid Metabolism in Signaling Systems) (1993) 213-223; C.A., 120 (1994) 130312k.
- 3721 Wu, S.Y. and Casida, J.E.: Neuropathy target esterase inhibitors: Enantiomeric separation and stereospecificity of 2-substituted-4H-1,3,2-benzodioxaphosphorin 2-oxides. *Chem. Res. Toxicol.*, 7 (1994) 77-81; C.A., 120 (1994) 127291x.

For additional information see C.A.:

120 (1994) 68445a, 75635r.

See also 3093, 3166, 3183, 3293, 3302, 3325, 3334, 3554, 3574, 4064, 4066.

26. ORGANOMETALLIC AND RELATED COMPOUNDS

26a. Organometallic compounds

3722 Sarzanini, C., Sacchero, G., Aceto, M., Abollino, O. and Mentasti, E.: Ion chromatographic separation and on-line cold vapour atomic absorption spectrometric determination of methylmercury, ethylmercury and inorganic mercury. *Anal. Chim. Acta*, 284 (1994) 661-667.

3723 Takeuchi, T. and Miwa, T.: Separation of ferrocenylethanol enantiomers by microcolumn liquid chromatography with β -cyclodextrin as mobile phase additive. *J. Chromatogr. A*, 666 (1994) 439-443.

26b. Boranes, silanes and related non-metallic compounds

See 3025.

26c. Coordination compounds

3724 Basova, E.M., Kazakova, O.V., Bondareva, L.G., Polenova, T.V. and Ivanov, V.M.: (Behavior of platinum-group metal chelates with 7-phenylazo-8-hydroxyquinoline-5-sulphonic acid in ion-pair HPLC). *Zh. Anal. Khim.*, 48 (1993) 1531-1538; C.A., 120 (1994) 94158d.

3725 Jen, J.-F. and Yang, S.-M.: Simultaneous speciation determination of vanadium(IV) and vanadium(V) as EDTA complexes by liquid chromatography with UV detection. *Anal. Chim. Acta*, 289 (1994) 97-104.

3726 Mueller, R. and Venclik, I.: Determination of complexed metal ions in a solution. *Eur. Pat. Appl. EP 562,337* (Cl. G01N30/96), 29 Sep. 1993, CH Appl. 92/943, 25 Mar. 1992; 7 p.; C.A., 120 (1994) 152588g.

3727 Wang, Y., Liu, Q., Liu, J. and Cheng, J.: (Separation of metal coordination complexes by reversed-phase high-performance liquid chromatography-application of solvophobic theory). *Sepu*, 11 (1993) 327-330; C.A., 120 (1994) 94155a.

See also 4080, 4083, 4099, 4104, 4134.

27. VITAMINS AND VARIOUS ANIMAL GROWTH FACTORS (NON-PEPTIDIC)

3728 Adewusi, S.R.A. and Bradbury, J.H.: Carotenoids in cassava: comparison of open-column and HPLC methods of analysis. *J. Sci. Food Agric.*, 62 (1993) 375-383; C.A., 120 (1994) 53517b.

3729 Akoh, C.C.: Oxidative stability of fat substitutes and vegetable oils by the oxidative stability index method. *J. Am. Oil Chem. Soc.*, 71 (1994) 211-216.

3730 Alyabis, A.M. and Simpson, K.L.: Comparison of reverse-phase C-18 open column with the Bio-Rex 70 column in the determination of thiamin. *J. Food Compos. Anal.*, 6 (1993) 166-171; C.A., 120 (1994) 161869b.

3731 Blau, N., Kierat, L., Heizmann, C.W., Endres, W., Giudici, T. and Wang, M.: Screening for tetrahydrobiopterin deficiency in newborns using dried urine on filter paper. *J. Inherited Metab. Dis.*, 15 (1992) 402-404; C.A., 120 (1994) 157780y.

3732 Boschi, V., de Siero, E., Ferrara, L., Pecoraro, P. and Falconi, C.: (Separation and identification of micro quantities of tocopherols). *Riv. Sci. Aliment.*, 22 (1993) 313-317; C.A., 120 (1994) 105209z.

3733 Bui, M.H.: Simple determination of retinol, α -tocopherol and carotenoids (lutein, all-trans-lycopene, α - and β -carotenes) in human plasma by isocratic liquid chromatography. *J. Chromatogr. B*, 654 (1994) 129-133.

3734 Felton, S.P., Grace, R. and Halver, J.E.: A non-ion-pairing HPLC method for measuring new forms of ascorbate and ascorbic acid. *J. Liq. Chromatogr.*, 17 (1994) 123-131.

3735 Gallice, P., Sarrazin, F., Polverelli, M., Cadet, J., Berland, Y. and Crevat, A.: Ascorbic acid-2-O- β -glucuronide, a new metabolite of vitamin C identified in human urine and uremic plasma. *Biochim. Biophys. Acta*, 1199 (1994) 305-310.

3736 Heintze, A. and Schultz, G.: High-performance liquid chromatographic determination of 3-hydroxy-3-methylglutaryl coenzyme A as diphenacyl ester of 3-hydroxy-3-methylglutarate. *J. Chromatogr. B*, 655 (1994) 117-120.

3737 Herve, C., Beyne, P. and Delacoux, E.: Determination of thiamine and its phosphate esters in human erythrocytes by high-performance liquid chromatography with isocratic elution. *J. Chromatogr. B*, 653 (1994) 217-220.

3738 Irgens, L.H., Farkas, G. and Vigh, G.: Displacement chromatography on cyclodextrin silicas. V. Separation of the enantiomers of 5,10-dideazatetrahydrofolic acid. *J. Chromatogr. A*, 666 (1994) 603-609.

3739 Jensen, S.K.: Retinol determination in milk by HPLC and fluorescence detection. *J. Dairy Res.*, 61 (1994) 233-240; C.A., 120 (1994) 268412z.

3740 Karnaukhova, E.N. and Mishina, I.M.: (Chromatographic separation of carotenoids). *Khim. Prom-st. (Moscow)*, No. 7 (1993) 287-292; C.A., 120 (1994) 100837t - a review with 85 refs.

3741 Kim, S.H. and Kim, G.R.: (Separation of several oil soluble vitamins by MHPLC). *Choson Minjujuui Inmin Konghwaguk Kwahagwan Tongbo*, (1993) 39-41; C.A., 120 (1994) 49207h.

3742 Kobayashi, T., Tsugawa, N., Okano, T., Masuda, S., Takeuchi, A., Kubodera, N. and Nishii, Y.: The binding properties, with blood proteins, and tissue distribution of 22-oxa-1 α ,25-dihydroxyvitamin D₃, a noncalcemic analogue of 1 α ,25-dihydroxyvitamin D₃, in rats. *J. Biochem. (Tokyo)*, 115 (1994) 373-380.

3743 Kurmann, A. and Indyk, H.: The endogenous vitamin D content of bovine milk: influence of season. *Food Chem.*, 50 (1994) 75-81; C.A., 120 (1994) 242897m.

3744 Manoharan, M. and Schuille, P.O.: Measurement of ascorbic acid in human plasma and urine by high-performance liquid chromatography. Results in healthy subjects and patients with idiopathic calcium urolithiasis. *J. Chromatogr. B*, 654 (1994) 134-139.

- 3745 Masuda, S., Strugnelli, S., Calverley, M.J., Makin, H.L.J., Kremer, R. and Jones, G.: *In vitro* metabolism of the anti-psoriatic vitamin D analog, calcipotriol, in two cultured human keratinocyte models. *J. Biol. Chem.*, 269 (1994) 4794-4803.
- 3746 Mattila, P., Piironen, V., Uusi-Rauva, E. and Koivistoinen, P.: Determination of 25-hydroxycholecalciferol content in egg yolk by HPLC. *J. Food Compos. Anal.*, 6 (1993) 250-255; *C.A.*, 120 (1994) 268422c.
- 3747 Meyer, E., Lambert, W.E. and de Leenheer, A.P.: Simultaneous determination of endogenous retinoic acid isomers and retinol in human plasma by isocratic normal-phase HPLC with ultraviolet detection. *Clin. Chem. (Washington)*, 40 (1994) 48-50.
- 3748 Miyauchi, Y., Sano, N. and Nakamura, T.: Simultaneous determination of nicotinic acid and its two metabolites in human plasma using solid-phase extraction in combination with high performance liquid chromatography. *Int. J. Vitam. Nutr. Res.*, 63 (1993) 145-149; *C.A.*, 120 (1994) 264980k.
- 3749 Moussa, F., Depasse, F., Lompret, V., Hautem, J.-Y., Girardet, J.-P., Fontaine, J.-L. and Aymard, P.: Determination of phylloquinone in intravenous fat emulsions and soybean oil by high-performance liquid chromatography. *J. Chromatogr. A*, 664 (1994) 189-194.
- 3750 Ono, S., Hirano, H. and Oikawa, K.: (Determination of vitamin B₂ in rat lens). *Bitamin*, 67 (1993) 243-247; *C.A.*, 120 (1994) 158070d.
- 3751 Peng, Y.S. and Peng, Y.M.: Simultaneous liquid chromatographic determination of carotenoids, retinoids, and tocopherols in human buccal mucosal cells. *Cancer Epidemiol., Biomarkers Prev.*, 1 (1992) 375-382; *C.A.*, 120 (1994) 264958j.
- 3752 Shibata, K.: (Measurement of 6-aminonicotinamide, an antagonist of niacin, by high-performance liquid chromatography). *Bitamin*, 67 (1993) 493-497; *C.A.*, 120 (1994) 211710r.
- 3753 Sowell, A.L., Huff, D.L., Yeager, P.R., Caudill, S.P. and Gunter, E.W.: Retinol, α -tocopherol, lutein/zeaxanthin, β -cryptoxanthin, lycopene, α -carotene, trans- β -carotene, and four retinyl esters in serum determined simultaneously by reversed-phase HPLC with multiwavelength detection. *Clin. Chem. (Washington)*, 40 (1994) 411-416.
- 3754 Urbach, J. and Rando, R.R.: Isomerization of all-trans-retinoic acid to 9-cis-retinoic acid. *Biochem. J.*, 299 (1994) 459-465.
- 3755 Van Leeuwen, J.P.T.M., Pols, H.A.P., van den Bermd, G.C.M., Kempenaar, J., Thio, H.B., Birkenhäger, J.C. and Ponc, M.: Role of extracellular calcium in the regulation of 1,25-dihydroxy-vitamin D₃ formation in cultured human keratinocytes. *Biochim. Biophys. Acta*, 1221 (1994) 167-170.
- 3756 Wang, X.: (Determination of water-soluble vitamins in natural seaweed by high performance liquid chromatography). *Sepu*, 11 (1993) 364-366; *C.A.*, 120 (1994) 53027s.
- 3757 Watanabe, F., Tamura, Y., Stupperich, E. and Nakano, Y.: Uptake of cobalamin by *Euglena* mitochondria. *J. Biochem. (Tokyo)*, 114 (1993) 793-799.
- 3758 Wongyai, S.: Separation of L-(+)-ascorbic and D-(-)-isoascorbic acids on multimodel phenylpropanolamine-coated silica. *Chromatographia*, 38 (1994) 600-612.
- 3759 Yamauchi, R., Yagi, Y. and Kato, K.: Isolation and characterization of addition products of α -tocopherol with peroxy radicals of dilinoleoylphosphatidylcholine in liposomes. *Biochim. Biophys. Acta*, 1212 (1994) 43-49.
- For additional information see *C.A.*:
120 (1994) 52994t, 144305x.
- See also 3204, 3209, 3249, 3695, 3696, 3967, 4029.
28. ANTIBIOTICS
- 3760 Al-Meshal, M.A.: Simple and sensitive high-performance liquid chromatographic analysis of cefetamet in plasma. *Anal. Lett.*, 27 (1994) 919-926.
- 3761 Amade, P., Mallea, M. and Bouaicha, N.: Isolation, structural identification and biological activity of two metabolites produced by *Penicillium olsonii* Bainer and Sartory. *J. Antibiot.*, 47 (1994) 201-207.
- 3762 Andersen, A., Warren, D.J. and Sloerdal, L.: A sensitive and simple high-performance liquid chromatographic method for the determination of doxorubicin and its metabolites in plasma. *Ther. Drug Monit.*, 15 (1993) 455-461; *C.A.*, 120 (1994) 182211w.
- 3763 Asukabe, H., Murata, H., Harada, K., Suzuki, M., Oka, H. and Ikai, Y.: Improvement of chemical analysis of antibiotics. 21. Simultaneous determination of three polyether antibiotics in feeds using high-performance liquid chromatography with fluorescence detection. *J. Agric. Food Chem.*, 42 (1994) 112-117.
- 3764 Blain, P., Demaesschalck, E., Rompaey, I.V. and Gosselé, F.: Optimization of the high-performance liquid chromatography for virginiamycin determination in mineral premixes. *Analyst (London)*, 119 (1994) 361-364.
- 3765 Faouzi, M.A., Dine, T., Luyckx, M., Goudaliez, F., Mallevais, M.L., Gressier, B., Brunet, C., Cazin, M. and Cazin, J.C.: Stability and compatibility studies of cephamandole nafate with PVC infusion bags. *J. Pharm. Biomed. Anal.*, 12 (1994) 99-104.
- 3766 Fejglóvá, Z., Dolezal, J., Hrdlicka, A. and Frgalová, K.: Microbore HPLC determination of polyether antibiotics using postcolumn derivatization with benzaldehyde reagents. *J. Liq. Chromatogr.*, 17 (1994) 359-372.
- 3767 Horie, M., Saito, K., Nose, N., Oka, H. and Nakazawa, H.: Determination of mirosamicin in animal tissues by high-performance liquid chromatography. *J. Chromatogr. B*, 655 (1994) 47-52.
- 3768 Horii, S.: Liquid chromatographic determination of oxytetracycline and chlortetracycline residues in animal tissues. *J. Liq. Chromatogr.*, 17 (1994) 213-221.
- 3769 Hosoe, K., Konishi, E., Hidaka, T., Yamane, T., Yamashita, K. and Ohashi, T.: High-performance liquid chromatographic determination of 3'-hydroxy-5'-(4-isobutyl-1-piperazinyl)benzoxazinorifamycin (KRM-1648) and its deacetyl metabolite in plasma, whole blood, urine and tissue samples in rats. *J. Chromatogr. B*, 653 (1994) 177-186.
- 3770 Iwasaki, K., Shiraga, T., Nagase, K., Tozuka, Z., Noda, K., Sakuma, S., Fujitsu, T., Shimatani, K., Sato, A. and Fujioka, M.: Isolation, identification, and biological activities of oxidative metabolites of FK506, a potent immunosuppressive macrolide lactone. *Drug Metab. Disp.*, 21 (1993) 971-977.
- 3771 Kamiyama, T., Shimma, N., Ohtsuka, T., Nakayama, N., Itezo, Y., Nakada, N., Watanabe, J. and Yokose, K.: Cyclothialidine, a novel DNA gyrase inhibitor. II. Isolation, characterization and structure elucidation. *J. Antibiot.*, 47 (1994) 37-45.

- 3772 Kuroda, K., Morishita, Y., Saito, Y., Ikuina, Y., Ando, K., Kawamoto, I. and Matsuda, Y.: AS-186 Compounds, new inhibitors of acyl-CoA-cholesterol acetyltransferase from *Penicillium aspersophorum* KY1635. *J. Antibiot.*, 47 (1994) 16-22.
- 3773 Lal, J., Paliwal, J.K., Grover, P.K. and Gupta, R.C.: Determination of ampicillin in serum by high-performance liquid chromatography with precolumn derivatization. *J. Chromatogr. B*, 655 (1994) 142-146.
- 3774 Lhoëst, G., Maton, N., Laurent, A. and Verbeeck, R.K.: Isolation and identification of a FK-506 C₃₆₋₃₇ dihydrodiol from erythromycin-induced rabbit liver microsomes. *J. Pharm. Biomed. Anal.*, 12 (1994) 235-241.
- 3775 Loftsson, T. and Johannesson, H.R.: The influence of cyclodextrin on the stability of cephalotin and aztreonam in aqueous solutions. *Pharmazie*, 49 (1994) 292-293.
- 3776 Lovdahl, M.J., Reher, K.E., Russlie, H.Q. and Canafax, D.M.: Determination of cefpodoxime levels in chinchilla middle ear fluid and plasma by high-performance liquid chromatography. *J. Chromatogr. B*, 653 (1994) 227-232.
- 3777 Mathur, A.K.: Determination of salinomycin by high-performance liquid chromatography using a precolumn derivatization technique. *J. Chromatogr. A*, 664 (1994) 284-288.
- 3778 Morishita, Y., Chiba, S., Tsukuda, E., Tanaka, T., Ogawa, T., Yamasaki, M., Yoshida, M., Kawamoto, I. and Matsuda, Y.: RES-701-1, a novel and selective endothelin type B receptor antagonist produced by *Streptomyces* sp. RE-701. I. Characterization of producing strain, fermentation, isolation, physico-chemical and biological properties. *J. Antibiot.*, 47 (1994) 269-275.
- 3779 Napoli, K.L. and Kahan, B.D.: Sample clean-up and high-performance liquid chromatographic techniques for measurement of whole blood rapamycin concentrations. *J. Chromatogr. B*, 654 (1994) 111-120.
- 3780 Newman, R.A., Fuentes, A., Minor, T.Y., McManus, K.T. and Garitez, D.A.: Determination of echinomycin (NSC-526417) in human plasma: comparison of conventional HPLC to a capillary HPLC, electrospray ionization and mass spectrometry system. *J. Liq. Chromatogr.*, 17 (1994) 403-417.
- 3781 Nishida, F., Nishimura, M., Harada, K.-I., Suzuki, M., Meevoort, V., Flegel, T.W., Thebtaranonth, Y. and Intararuangorn, S.: Structure elucidation of glykenin glycosidic antibiotics from *Basidiomycetes* sp. V. High-performance liquid chromatographic separation of components of glykenin. *J. Chromatogr. A*, 664 (1994) 195-202.
- 3782 Nishikawa, T., Hasumi, H., Suzuki, S., Kubo, H. and Ohtani, H.: Interconversion of cyclosporin molecular form inducing peak broadening, tailing and splitting during reversed-phase liquid chromatography. *Chromatographia*, 38 (1994) 359-364.
- 3783 Ohtsuka, T., Itezono, Y., Nakayama, N., Sakai, A., Shimma, N. and Yokose, K.: Hispidospermidin, a novel phospholipase C inhibitor produced by *Chaetosphaeronema hispidulum* (Cda) Moesz NR 7127. II. Isolation, characterization and structural elucidation. *J. Antibiot.*, 47 (1994) 6-15.
- 3784 Pouliquen, H., Pinault, L. and le Bris, H.: Determination of oxolinic acid in seawater, marine sediment, and Japanese oyster (*Crassostrea gigas*) by high-performance liquid chromatography. *J. Liq. Chromatogr.*, 17 (1994) 929-945.
- 3785 Prokša, B., Adamcová, J. and Fuska, J.: Detection and assay of secondary metabolites of *Penicillium vermiculatum* DANG. *J. Chromatogr. A*, 665 (1994) 185-190.
- 3786 Ramos, M., Reuvers, T., Aranda, A. and Gómez, J.: Determination of chloramphenicol in chicken muscle by high performance liquid chromatography and UV-diode array detection. *J. Liq. Chromatogr.*, 17 (1994) 385-401.
- 3787 Roggo, B.E., Petersen, F., Delmendo, R., Jenny, H.-B., Peter, H.H. and Roesel, J.: 3-Alkanoyl-5-hydroxymethyl tetronic acid homologues and resistomycins: new inhibitors of HIV-1 protease. I. Fermentation, isolation and biological activity. *J. Antibiot.*, 47 (1994) 136-142.
- 3788 Samouris, G., Nathanael, B., Tsoukali-Papadopoulou, H. and Papadimitriou, N.: Determination of chloramphenicol residues in eggs by high performance liquid chromatography (HPLC). *Vet. Hum. Toxicol.*, 35 (1993) 406-409; C.A., 120 (1994) 189950v.
- 3789 Scaloni, A., Barra, D. and Bossa, F.: Sequence analysis of dehydroamino acid-containing peptides. *Anal. Biochem.*, 218 (1994) 226-228.
- 3790 Takeda, K., Yano, S., Sakuma, Y. and Yamaguchi, T.: (High-performance liquid chromatography method for temafloxacin in body fluids). *Chemotherapy (Tokyo)*, 41(Suppl. 5) (1993) 122-127; C.A., 120 (1994) 152873w.
- 3791 Takhirov, M.T., Tarasov, V.V., Bekzhanova, E.E. and Likho, V.G.: (Determination of tylosin in water by chromatography). *Gig. Sanit.*, (1993) 76-70; C.A., 120 (1994) 143512a.
- 3792 Taylor, R.B., Richards, R.M.E., Low, A.S. and Hardie, L.: Chemical stability of polymyxin B in aqueous solution. *Int. J. Pharm.*, 102 (1994) 201-206; C.A., 120 (1994) 143954w.
- 3793 Videira, M. and Aires-Barros, M.R.: Liquid-liquid extraction of clavulanic acid using an aqueous two-phase system of polyethylene glycol and potassium phosphate. *J. Chromatogr. A*, 668 (1994) 237-240.
- 3794 Wong, B.K., Bruhin, P.J. and Lin, J.H.: Column-switching high-pressure liquid chromatographic method for the determination of a new carbapenem antibiotic, L-739,428, in rat and monkey plasma. *J. Chromatogr. B*, 655 (1994) 158-162.
- 3795 Zhang, H. and Stewart, J.T.: Determination of a cefuroxime and aminophylline/theophylline mixture by high-performance liquid chromatography. *J. Liq. Chromatogr.*, 17 (1994) 1327-1335.
- 3796 Zuhowski, E.G., Gutheil, J.C. and Egorin, M.J.: Rapid and sensitive high-performance liquid chromatographic assay for novobiocin in human serum. *J. Chromatogr. B*, 655 (1994) 147-152.

For additional information see C.A.:

120 (1994) 45054h, 62432x, 62469q, 68714n, 94549g, 132495w, 144320y, 144322a, 239602f, 260382u.

See also 3219, 3973.

29. INSECTICIDES, PESTICIDES AND OTHER AGROCHEMICALS

29a. General techniques

- 3797 Brinkman, U.A.T., Lingeman, H. and Slobodnik, J.: SAMOS LC: a fully automated liquid chromatographic method for trace-level monitoring of polar pesticides. *LC-GC Int.*, 7 (1994) 157-163.

- 3798 Cardinali, D.: (Use of SPE (solid-phase extraction) in the determination of fungicides, insecticides, chlorinated acaricides and pyrethroids, in legumes and fruits, using capillary GLC). *Riv. Soc. Ital. Sci. Aliment.*, 21 (1992) 461-469; C.A., 120 (1994) 215545p.
- 3799 Chiron, S., Dupas, S., Scribe, P. and Barceló, D.: Application of on-line solid-phase extraction followed by liquid chromatography-thermospray mass spectrometry to the determination of pesticides in environmental waters. *J. Chromatogr. A*, 665 (1994) 295-305.
- 3800 Farhat, S., Baig, M.M.H. and Khan, M.A.: Retention times and response data of 40 insecticides by HPLC. *Karachi Univ. J. Sci.*, 20 (1992) 143-149; C.A., 120 (1994) 127591v.
- 3801 Huen, J.M., Gillard, R., Mayer, A.G., Baltensperger, B. and Kern, H.: Automatic measurement of pesticides in drinking water using on-line solid-phase extraction, HPLC and chemometry. *Fresenius J. Anal. Chem.*, 348 (1994) 606-614.
- 3802 Klisenko, M.A. and Makartchuk, T.L.: (Determination of pesticide residues by high-performance liquid chromatography). *Agrokhimiya*, No. 1 (1992) 139-158; C.A., 120 (1994) 238092c - a review with 192 refs.
- 3803 Miroshnitchenko, I.I. and Kretova, L.G.: (Determination of pesticide residues by liquid chromatography). *Agrokhimiya*, No. 1 (1992) 169-176; C.A., 120 (1994) 238093d - a review with 73 refs.
- 3804 Pichon, V. and Hennion, M.-C.: Determination of pesticides in environmental waters by automated on-line trace-enrichment and liquid chromatography. *J. Chromatogr. A*, 665 (1994) 269-281.
- For additional information see C.A.:
120 (1994) 268395w.
- 29b. *Chlorinated insecticides*
- See 4059.
- 29c. *Phosphorus insecticides*
- 3805 Saito, Y., Yamamoto, A. and Mizukami, E.: (Analysis of pesticide residues in vegetables and fruits. Separation and purification of organophosphorus pesticides from vegetable lipids by gel permeation chromatography). *Toyama-ken Eisei Kenkyusho Nenpo* 1992, 16 (1993) 210-214; C.A., 120 (1994) 53043u.
- See also 2843, 4059.
- 29d. *Carbamates*
- 3806 Hiemstra, M. and de Kok, A.: Determination of N-methylcarbamate pesticides in environmental water samples using automated on-line trace enrichment with exchangeable cartridges and high-performance liquid chromatography. *J. Chromatogr. A*, 667 (1994) 155-166.
- 3807 Honing, M., Barceló, D., Ghijsen, R.T. and Brinkman, U.A.T.: Optimisation of the liquid chromatographic separation of pirimicarb and its metabolites V-VII: application to a soil sample used as a candidate reference material. *Anal. Chim. Acta*, 286 (1994) 457-468.
- 3808 Matsuda, R., Hayashi, Y., Suzuki, T. and Saito, Y.: (Optimization of the reversed-phase liquid chromatographic analysis of N-methylcarbamate pesticides on the basis of information theory). *Bunseki Kagaku*, 42 (1993) 881-886; C.A., 120 (1994) 210622b.
- 3809 Yang, S.S. and Smetena, I.: Determination of aldicarb, aldicarb sulfoxide and aldicarb sulfone in tobacco using high-performance liquid chromatography with dual post-column reaction and fluorescence detection. *J. Chromatogr. A*, 664 (1994) 289-294.
- 29e. *Herbicides*
- 3810 Chang, C.-D., Armstrong, D.W. and Fleischmann, T.J.: Fractionation of [¹⁴C]metolachlor metabolites by centrifugal partition chromatography. *J. Liq. Chromatogr.*, 17 (1994) 19-32.
- 3811 Chiron, S., Martinez, E. and Barceló, D.: On-line and off-line sample preparation of acidic herbicides and bentazone transformation products in estuarine waters. *J. Chromatogr. A*, 665 (1994) 283-293.
- 3812 Hamper, B.C., Dukeshere, D.R. and Moedritzer, K.: Analytical and preparative separation of the enantiomers of pyrazole phenyl ether herbicides on three chiral stationary phases. *J. Chromatogr. A*, 666 (1994) 479-484.
- 3813 Huart, B., Paillet, F.M., Trique, B. and Ricordel, I.: (Protocol for studying herbicide diffusion in an ecosystem). *Ann. Falsif. Expert. Chim. Toxicol.*, 86 (1993) 75-86; C.A., 120 (1994) 47966f.
- 3814 Laganà, A., Fago, G., Marwo, A. and Pardo-Martinez, B.: Rapid method for determination of phenylurea herbicides. *Chromatographia*, 38 (1994) 88-92.
- 3815 Pratap, K. and Lemley, A.T.: Electrochemical peroxide treatment of aqueous herbicide solutions. *J. Agric. Food Chem.*, 42 (1994) 209-215.
- 3816 Schewes, R., Wüst, S., Lepschy-v. Gleissenthal, J., Maidl, F.X., Süß, A., Hock, B. and Fischbeck, G.: Determination of weathered atrazine residues in soil by enzyme immunoassay and HPLC: an evaluation study. *Anal. Lett.*, 27 (1994) 487-494.
- See also 2964.
- 29f. *Fungicides*
- 3817 Jinénez, J.J., Atienza, J., Bernal, J.L. and Toribio, L.: Determination of carbendazim in lettuce samples by SFE-PLCC. *Chromatographia*, 38 (1994) 391-399.
- 29g. *Other types of pesticides and various agrochemicals*
- 3818 Koeduka, K., Kenmotsu, K., Ogino, Y., Matsunaga, K. and Mori, T.: (Determination of methomyl and oxamyl in the environmental and biological samples by HPLC with post column fluorescence derivation). *Kankyo Kagaku*, 3 (1993) 771-782; C.A., 120 (1994) 210623c.
- 3819 Rasmussen, H.T., Friedman, S.K., Mustilli, A.J., McDonough, R. and McPherson, B.P.: Analysis of methyl neodecanamide in lake water by reversed-phase high performance liquid chromatography and gas chromatography-mass spectrometry. *J. Liq. Chromatogr.*, 17 (1994) 589-601.
- See also 2843, 4042, 4049.

30. SYNTHETIC AND NATURAL DYES

30a. Synthetic dyes

- 3820 Achwal, W.B.: Assessment of cellulosic dyeing parameters by a fiber column. *Colourage*, 40 (1993) 16-17; C.A., 120 (1994) 79295c.
- 3821 Izquierdo-Hornillos, R. and Shepard, T.A.: HPLC separation and determination of phenolphthalein and its glucuronid as markers of enterohepatic circulation using acetonitrile-methanol as organic modifier and application to rat plasma. *J. Liq. Chromatogr.*, 17 (1994) 157-173.
- 3822 Izquierdo-Hornillos, R. and Shepard, T.A.: HPLC-isocratic study of the separation and determination of phenolphthalein and its metabolite, phenolphthalein-glucuronic acid, as markers of enterohepatic circulation. *Microchem. J.*, 48 (1993) 356-364; C.A., 120 (1994) 49238u.
- 3823 Loefgren, A. and Andrasko, J.: HPLC analysis of printing inks. *J. Forensic Sci.*, 38 (1993) 1151-1140; C.A., 120 (1994) 98627d.

For additional information see C.A.:
120 (1994) 132496x.

30b. Chloroplast and other natural pigments

- 3824 Drexler, D. and Ballschmiter, K.: Separation of chlorophyll and metalloporphyrins by HPLC in normal-phase mode using a diol-phase. *Fresenius J. Anal. Chem.*, 348 (1994) 590-594.
- 3825 Kuiters, A.T. and Mulder, W.: Complexometric gel permeation chromatography of soluble humic substances using Sephadex G-25. *J. Soil Sci.*, 44 (1993) 501-512; C.A., 120 (1994) 162531x.
- 3826 Liw, K.Y., Nordin, M.R. and Goh, L.S.: Reaction of carotenes in palm oil with acid. *J. Am. Oil Chem. Soc.*, 71 (1994) 303-306.
- 3827 Luo, W., Al-Abdulay, A.B., Yoon, K. and Simpson, K.L.: Rapid determination of blood serum retinol by reversed-phase open-column chromatography. *Int. J. Vitam. Nutr. Res.*, 63 (1993) 82-86; C.A., 120 (1994) 264979s.
- 3828 Powell, C., Clifford, M.N., Pie, S.C., Ford, M.A., Robertson, A. and Gibson, C.L.: Tea cream formation: the contribution of black tea phenolic pigments determined by HPLC. *J. Sci. Food Agric.*, 63 (1993) 77-86; C.A., 120 (1994) 52997w.
- 3829 Voitke, P., Martin, C.-D., Nicklish, S. and Kohl, J.-G.: HPLC determination of lipophilic photosynthetic pigments in algal culture and lake water samples using a non-encapped C18-RP-column. *Fresenius J. Anal. Chem.*, 348 (1994) 762-768.

For additional information see C.A.:
120 (1994) 129031m, 161842n.

See also 3050, 3728, 3733, 3753.

31. PLASTICS AND THEIR INTERMEDIATES

- 3830 Balke, S.T., Mourey, T.H. and Harrison, C.A.: Number-average molecular weight by size exclusion chromatography. *J. Appl. Polym. Sci.*, 51 (1994) 2087-2102; C.A., 120 (1994) 192802d.

- 3831 Fréchet, J.M.J., Lochman, L., Šmigol, V. and Švec, F.: Reversed-phase high-performance liquid chromatography of functionalized dendritic macromolecules. *J. Chromatogr. A*, 667 (1994) 284-289.
- 3832 Grubisic-Gallot, Z., Gallot, Y. and Sedlacek, J.: Study of polystyrene-block-poly(methyl methacrylate) micelles by size exclusion chromatography/low-angle laser light scattering. 1. Influence of copolymer concentration and flow rate. *Macromol. Chem. Phys.*, 195 (1994) 781-791; C.A., 120 (1994) 165426c.
- 3833 Mukoyama, Y., Sugitani, H. and Mori, S.: Effects of the addition of lithium bromide and phosphoric acid in size-exclusion chromatography of poly(amide imide) and polyamic acid. *J. Appl. Polym. Sci.: Appl. Polym. Symp.*, 52 (1993) 183-191; C.A., 120 (1994) 55481j.
- 3834 Pleshkova, A.P. and Volkovich, S.V.: (Chromatographic-mass spectrometric analysis of impurities in circulating 2-ethylhexanol for the production of dioctyl phthalate plasticizer). *Khim. Prom-st. (Moscow)*, (1993) 417-421; C.A., 120 (1994) 109915z.
- 3835 Shalliker, R.A., Kavanagh, P.E. and Russell, I.M.: Separations of high-molecular-mass polystyrenes on different pore size and particle size reversed-phase columns in dichloromethane-acetonitrile. *J. Chromatogr. A*, 664 (1994) 221-228.
- 3836 Sokolov, M.I., Kovshov, Yu.S., Zornikov, I.P. and Moiseiv, V.V.: (Use of gel chromatography to select the conditions for hydrogenation of nitrite rubber in the latex stage). *Proizvod. Ispol'z. Elastomerov*, (1992) 4-8; C.A., 120 (1994) 136756e.
- 3837 Thompson, C.M., Taylor, S.G. and McGee, W.W.: Gel permeation chromatographic analysis of polyurethane prepolymer synthesis kinetics II. The effect of stoichiometry and type of diisocyanate. *J. Polym. Sci., Part A: Polym. Chem.*, 32 (1994) 113-120; C.A., 120 (1994) 77718u.
- 3838 Volet, G. and Lesec, J.: Non-exclusion effects in aqueous SEC: behavior of some polyelectrolytes using on-line mass detectors. *J. Liq. Chromatogr.*, 17 (1994) 559-577.

For additional information see C.A.:
120 (1994) 78551w, 152706u, 165590b, 193232e.

See also 2825, 2846, 2855, 2867, 2898, 3235.

32. DRUG ANALYSIS

- 3839 Klein, J., Fernandes, D., Gazarian, M., Kent, G. and Koren, G.: Simultaneous determination of lidocaine, prilocaine and the prilocaine metabolite o-toluidine in plasma by high-performance liquid chromatography. *J. Chromatogr. B*, 655 (1994) 83-88.

32a. Drug analysis, general techniques

- 3840 Bernal, J.L., del Nozal, M.J., Rosas, V. and Villarino, A.: Extraction of basic drugs from whole blood and determination by high performance liquid chromatography. *Chromatographia*, 38 (1994) 617-623.
- 3841 Hsiao, C.L.: The study of drug-albumin binding by the Hummel-Dreyer method using size-exclusion high-performance liquid chromatography. Avail. *Univ. Microfilms Int., Order No. DA9317527*, 1993, 81 pp.; C.A., 120 (1994) 152852p.

- 3842 Hu, C. and Jin, S.: (Error analysis in high-performance liquid chromatography). *Zhongguo Kangshengsu Zazhi*, 18 (1993) 312-319; *C.A.*, 120 (1994) 86549s - a review with 4 refs.
- 3843 Hubert, P., Chiap, P., Moors, M., Bourguignon, B., Massart, D.L. and Crommen, J.: Knowledge-based system for the automated solid-phase extraction of basic drugs from plasma coupled with their liquid chromatographic determination. Application to the biodetermination of β -receptor blocking agents. *J. Chromatogr. A*, 665 (1994) 87-99.
- 3844 Kohl, C., Oelschläger, H. and Rothley, D.: Identification of drug glucuronides in human urine by RP-HPLC after derivatization. *J. Pharm. Biomed. Anal.*, 12 (1994) 249-254.
- 3845 Konishi, M., Okuno, K. and Hashimoto, H.: Interaction of cyclodextrins with drugs. Chromatographic investigation using a cyclodextrin bonded-phase and its application to chiral separations. *Chromatographia*, 38 (1994) 381-385.
- 3846 Kugel, C., Heintzelmann, B. and Wagner, J.: Determination of distribution coefficients for some 5-HT₃ receptor antagonists by reversed-phase high-performance liquid chromatography. *J. Chromatogr. A*, 667 (1994) 29-35.
- 3847 Lafont, O., Menager, S. and Lacroix, C.: (Preparation of reference substances. An indispensable contribution to HPLC). *Actual. Chim. Ther.*, 19 (1992) 233-242; *C.A.*, 120 (1994) 207746q - a review with 13 refs.
- 3848 Usselman, B.: (Methods and apparatus for the determination of drugs in body fluids). *Prax. Klin. Pharm.*, 1 (Klinische Pharmakokinetik) (1992) 29-43; *C.A.*, 120 (1994) 235186a - a review with 16 refs.
- 3849 Yamaguchi, M.: (Sensitive fluorescence and chemiluminescence HPLC of biological substances and drugs). *Dojin News*, 68 (1993) 10-16; *C.A.*, 120 (1994) 211664d - a review with 51 refs.
- For additional information see *C.A.*:
120 (1994) 62374e, 94562f, 158152g.
- See also 2783, 2842, 2872, 2897, 3383.
- 32b. *Antirheumatics and antiinflammatory drugs*
- 3850 Ahn, H.-Y., Shiu, G.K., Trafton, W.F. and Doyle, T.D.: Resolution of the enantiomers of ibuprofen; comparison study of diastereomeric method and chiral stationary phase method. *J. Chromatogr. B*, 653 (1994) 163-169.
- 3851 Chakir, S., Maurice, M.-H., Magdalou, J., Leroy, P., Dubois, N., Lapique, F., Abdelhamid, Z. and Nicolas, A.: High-performance liquid chromatographic enantioselective assay for the measurement of ketoprofen glucuronidation by liver microsomes. *J. Chromatogr. B*, 654 (1994) 61-68.
- 3852 Delbeke, F.T., Vynckier, L. and Debackere, M.: The disposition of suxibuzone in the horse. *J. Vet. Pharmacol. Ther.*, 16 (1993) 283-290; *C.A.*, 120 (1994) 182216b.
- 3853 Kempe, M. and Mosbach, K.: Direct resolution of naproxen on a non-covalently molecularly imprinted chiral stationary phase. *J. Chromatogr. A*, 664 (1994) 276-279.
- 3854 Kumbhat, S.: Study of ocular penetration of flurbiprofen in human aqueous humour. *J. Pharm. Biomed. Anal.*, 12 (1994) 131-133.
- 3855 Lo, C.S., Wang, C.F. and Hsu, H.C.: A stability-indicating HPLC assay method of indomethacin and the stability of indomethacin gels. *Zhonghua Yaoxue Zazhi*, 45 (1993) 321-328; *C.A.*, 120 (1994) 144332d.
- 3856 Mishra, P., Arif, S. and Shakya, A.K.: Determination of naproxen in tablets by HPLC. *J. Inst. Chem. (India)*, 65 (1993) 27; *C.A.*, 120 (1994) 62470h.
- 3857 Mohamed, F.A., Jun, H.W., Elfaham, T.H., Sayed, H.A. and Hafez, E.: An improved HPLC procedure for the quantitation of diclofenac in plasma. *J. Liq. Chromatogr.*, 17 (1994) 1065-1088.
- 3858 Riegel, M. and Ellis, P.P.: High-performance liquid chromatographic assay for antiinflammatory agents diclofenac and flurbiprofen in ocular fluids. *J. Chromatogr. B*, 654 (1994) 140-145.
- 3859 Sochor, J., Klimesh, J., Zahradniczek, M. and Sedlacek, J.: High-performance liquid chromatographic assay for ibuprofen in whole blood using solid-phase extraction. *J. Chromatogr. B*, 654 (1994) 282-286.
- See also 2984, 2985, 3278.
- 32c. *Autonomic and cardiovascular drugs*
- 3860 Aboul-Enein, H.Y., Bakr, S.A. and Nicholls, P.J.: Direct enantiomeric separation of phenglutarimide by chiral high performance liquid chromatography. *J. Liq. Chromatogr.*, 17 (1994) 1105-1110.
- 3861 Ader, U., Muschalek, V. and Schneider, M.P.: Chromatographic resolution of chiral intermediates in β -adrenergic blocker synthesis on chiral stationary phases. *Chirality*, 5 (1993) 554-559; *C.A.*, 120 (1994) 173605a.
- 3862 Brandšteterová, E., Romanová, D., Králiková, D., Božeková, L. and Kříška, M.: Automatic solid-phase extraction and high-performance liquid chromatographic determination of quinidine in plasma. *J. Chromatogr. A*, 665 (1994) 101-104.
- 3863 Chen, C.L., Sangiah, S., Roder, J.D., Chen, H., Berlin, K.D., Garrison, G.L., Scherlag, B.J. and Lazzara, R.: Determination of GLC-V-13, a novel antiarrhythmic agent, in plasma and urine by high-performance liquid chromatography. *J. Liq. Chromatogr.*, 17 (1994) 913-927.
- 3864 Colletti, A.E. and Krieter, P.A.: Disposition of the angiotensin II receptor antagonist L-158,809 in rats and rhesus monkeys. *Drug Metab. Disp.*, 22 (1994) 183-188.
- 3865 De Vries, J.X. and Schmitz-Kummer, E.: Determination of the coumarin anticoagulant phenprocoumon and metabolites in human plasma, urine and breast milk by high-performance liquid chromatography after solid-phase extraction. *J. Chromatogr. B*, 655 (1994) 63-71.
- 3866 Diamond, S., Rakestraw, D., O'Neil, J., Lam, G.N. and Christ, D.D.: Induction of cytochromes P-450 and 3A in mice following the dietary administration of the novel cognitive enhancer lincirdine. *Drug Metab. Disp.*, 22 (1994) 65-73.
- 3867 Facklam, C. and Modler, A.: Separation of some enantiomers and diastereomers of propranolol derivatives by high-performance liquid chromatography. *J. Chromatogr. A*, 664 (1994) 203-211.
- 3868 Falco, P.C., Herráez-Hernández, R. and Sevillano-Cabeza, A.: Determination of triamterene in urine by HPLC using fluorescence detection and column-switching. *Chromatographia*, 38 (1994) 29-34.

- 3869 Flanagan, R.J.: High-performance liquid chromatography of psychotropic and related drugs. *IBRO Handb. Ser.*, 15(High Performance Liquid Chromatography in Neuroscience Research) (1993) 321-356; *C.A.*, 120 (1994) 98508r - a review with 91 refs.
- 3870 Goicolea, M.A., Gomez de Balugera, Z., Portela, M.J. and Barrio, R.: Evaluation of amperometric detection at a glassy carbon electrode for the liquid chromatographic determination of anti-hypertensive substances. *Analyst (London)*, 119 (1994) 269-272.
- 3871 Grundy, J.S., Kherani, R. and Foster, R.T.: Sensitive high-performance liquid chromatographic assay for nifedipine in human plasma utilizing ultraviolet detection. *J. Chromatogr. B*, 654 (1994) 146-151.
- 3872 Halpin, R.A., Ulm, E.H., Till, A.E., Kari, P.H., Vyas, K.P., Hunninghake, D.B. and Duggan, D.E.: Biotransformation of lovastatin. V. Differences in *in vivo* metabolite profiles of mouse, rat, dog, and human. *Drug Metab. Disp.*, 21 (1993) 1003-1011.
- 3873 Hamoir, T., Verlinden, Y. and Massart, D.L.: Reversed-phase liquid chromatography of β -adrenergic blocking drugs in the presence of a tailing suppressor. *J. Chromatogr. Sci.*, 32 (1994) 14-20.
- 3874 Haupt, D., Pettersson, C. and Westerlund, D.: Enantiomeric separations of remoxipride, propranolol, and trimipramine on CHIRAL-AGP using micellar chromatography and anionic additives. *Chirality*, 5 (1993) 224-228; *C.A.*, 120 (1994) 207657m.
- 3875 Hoshino, M., Matsui, E., Yajima, K. and Okahira, A.: Direct high-performance liquid chromatographic separation of the racemates and diastereomers of nadolol. *J. Chromatogr. A*, 664 (1994) 104-110.
- 3876 Labhasetwar, V., Underwood, T., Gallagher, M., Murphy, G., Langberg, J. and Levy, R.J.: Sotalol controlled-release systems for arrhythmias: *in vitro* characterization, *in vivo* drug disposition and electrophysiologic effects. *J. Pharm. Sci.*, 83 (1994) 156-164.
- 3877 Lo, C.-F. and Chen, C.-M.: Determination of higenamine in plasma and urine by high-performance liquid chromatography with electrochemical detection. *J. Chromatogr. B*, 655 (1994) 33-39.
- 3878 Migulla, H., Bauer, S., Kewitz, H. and Roots, I.: Mikromethode zur Bestimmung von Amrinon und Acetylamrinon in Plasma und Urin durch HPLC. *Pharmazie*, 49 (1994) 290-291.
- 3879 Muck, W. and Bode, H.: Bioanalytics of nimodipine - an overview of methods. *Pharmazie*, 49 (1994) 130-139.
- 3880 Mutschler, E.: Simultaneous determination of (R)- and (S)-brefenolol in human plasma: high-performance liquid chromatographic assay on a chiral stationary phase. *Arch. Pharm. (Weinheim)*, 327 (1994) 193-196; *C.A.*, 120 (1994) 260395a.
- 3881 Nourooz-Zadeh, J., Gopaul, N.K., Forster, L.A., Ferns, G.A. and Ånggård, E.E.: Measurement of plasma probucol levels by high-performance liquid chromatography. *J. Chromatogr. B*, 654 (1994) 55-60.
- 3882 Oertel, R., Richter, K., Trausch, B. and Gramatté, T.: Eine schnelle, einfache und empfindliche HPLC-Methode zur Bestimmung von Talinolol im Serum. *Pharmazie*, 49 (1994) 291-292.
- 3883 Quaglia, M.G., Desideri, N., Bossù, E., Farina, A. and Morrone, L.A.: Chromatographic resolution and pharmacological investigation of ICI 118551, a new β_2 -blocker. *J. Chromatogr. A*, 666 (1994) 289-293.
- 3884 Rutledge, D.R., Abadi, A.H. and Lopez, L.M.: Liquid chromatographic determination of celirolol, diltiazem, desmethyl diltiazem and deacetyldiltiazem in plasma using a short alkyl chain silanol deactivated column. *J. Pharm. Biomed. Anal.*, 12 (1994) 135-140.
- 3885 Tateishi, T., Harada, K. and Ebihara, A.: Fluorescence detection of mexiletine and its *p*-hydroxylated and hydroxymethylated metabolites in human plasma and urine by high-performance liquid chromatography using post-column derivatization with *o*-phthalaldehyde. *J. Liq. Chromatogr.*, 17 (1994) 659-671.
- 3886 Thongnopnua, P. and Viwatwongsa, K.: Quantitative analysis of nifedipine in plasma by high-performance liquid chromatography. *J. Pharm. Biomed. Anal.*, 12 (1994) 119-125.
- 3887 Trenque, T., Barre, J., Lamiable, D. and Tillement, J.P.: Linear pharmacokinetics of a new inotropic agent. *Arzneim.-Forsch.*, 44 (1994) 471-473.
- 3888 Turcan, R.G., Hillbeck, D., Hartlay, T.E., Gilbert, P.J., Coe, R.A.J., Troke, J.A. and Vose, C.W.: Disposition of [14 C]velnacrine maleate in rats, dogs, and humans. *Drug Metab. Disp.*, 21 (1993) 1037-1047.
- 3889 Zimmer, D. and Muschalek, V.: Enantioselective assay for the determination of nisoldipine in dog, rat and mouse plasma by chiral microbore high-performance liquid chromatography combined with gas chromatography-mass spectrometry. *J. Chromatogr. A*, 666 (1994) 241-248.

For additional information see C.A.:

120 (1994) 62429b, 94558j, 98621x, 144295u, 200507d.

See also 2937, 3238, 3843, 4003, 4025.

32d. Central nervous system drugs

- 3890 Agostini, O., Bonacchi, G., Coppini, G., Toja, E., Triolo, A., Manzani, S., Pieraccini, G. and Moneti, G.: Metabolism of isbufylline in humans. Isolation, identification, and synthesis of plasma and urine metabolites. *Drug Metab. Disp.*, 22 (1994) 259-268.
- 3891 Bamio-Nuez, A., Artelejo-Ortega, B., del Pozo Carrascosa, A.: (Analytical validation: application of a method for the high-performance liquid chromatography in the analysis of benzodiazepines). *Circ. Farm.*, 50 (1992) 295-308; *C.A.*, 120 (1994) 62461f.
- 3892 Bokovikova, T.N., Klyuev, N.A., Gorozhankin, S.K., Stronova, L.A. and Suranova, A.V.: (Determination of impurities in fentanyl). *Khim.-Farm. Zh.*, 27 (1993) 58-60; *C.A.*, 120 (1994) 11630d.
- 3893 Cabrera, K. and Lubda, D.: Influence of temperature on chiral high-performance liquid chromatographic separations of oxazepam and Prominal on chemically bonded β -cyclodextrin as stationary phase. *J. Chromatogr. A*, 666 (1994) 433-438.
- 3894 Campins-Falcó, P., Sevillano-Cabeza, A. and Molins-Legua, C.: Amphetamine and methamphetamine determinations in biological samples by high performance liquid chromatography. A review. *J. Liq. Chromatogr.*, 17 (1994) 731-747 - a review with 44 refs.
- 3895 Celma, C.: Mass spectrometry and liquid chromatography/mass spectrometry of some derivatives of 6-(2-chlorophenyl)-1-methyl-7,8,9,10-tetrahydro-4H-pyrido(4',3'-4,5)thieno(3,2-f)(1,2,4)triazolo(4,3-1)(1,4)diazepines. *Biol. Mass Spectrom.*, 23 (1994) 13-19; *C.A.*, 120 (1994) 152676j.

- 3896 Chopineau, J., Rivault, F., Sautou, V. and Sommier, M.F.: Determination of temazepam and its active metabolite, oxazepam in plasma, urine and dialysate using solid-phase extraction followed by high performance liquid chromatography. *J. Liq. Chromatogr.*, 17 (1994) 373-383.
- 3897 Demian, I.: High-performance liquid chromatography (HPLC) chiral separations of guaifenesin, methocarbamol, and racemorphan. *Chirality*, 5 (1993) 238-240; *C.A.*, 120 (1994) 116931e.
- 3898 Foda, N.H.: Quantitative analysis of metoclopramide in tablet formulations by HPLC. *Anal. Lett.*, 27 (1994) 549-559.
- 3899 Fossler, M.J., Leslie, J., Burke, M., Devane, J. and Young, D.: A reversed-phase ion-pair liquid chromatography method for the determination of 4-aminopyridine in Sprague-Dawley rats. *J. Pharm. Biomed. Anal.*, 12 (1994) 281-285.
- 3900 Gagliardi, L., de Orsi, D., Cavazzutti, G., Tonelli, D. and Zappoli, S.: HPLC determination of oxiracetam, its impurities, and piracetam in pharmaceutical formulations. *Anal. Lett.*, 27 (1994) 879-885.
- 3901 Gildersleeve, D.L., Jung, Y.-Y. and Wieland, D.M.: Direct optical resolution of vesamicol and a series of benzovesamicol analogues by high-performance liquid chromatography. *J. Chromatogr. A*, 667 (1994) 183-189.
- 3902 Groen, K., Zeijlmans, P.W.M., Burm, A.G.L. and van Kleef, J.W.: Improved clean-up procedure for the high-performance liquid chromatographic assay of bupivacaine enantiomers in human plasma and ultrafiltrate in the nanogram per milliliter range. *J. Chromatogr. B*, 655 (1994) 163-166.
- 3903 Gupta, R.N. and Dziurdzy, S.A.: Therapeutic monitoring of sertraline. *Clin. Chem. (Washington)*, 40 (1994) 498-499.
- 3904 Gupta, R.N., Fuller, H.D. and Dolovich, M.B.: Optimization of a column liquid chromatographic procedure for the determination of plasma salbutamol concentration. *J. Chromatogr. B*, 654 (1994) 205-211.
- 3905 Harrison, F.J.J., Heisler, F., Scherer, J., Wulsch, A., Galiano, A., Hornauer, M. and Kelly, J.: Study of pharmacokinetics and pharmacodynamics of two preparations of metoclopramide. *Arzneim.-Forsch.*, 44 (1994) 519-521.
- 3906 Hayakawa, K., Miyoshi, Y., Kurimoto, H., Matsushima, Y., Takayama, N., Tanaka, S. and Miyazaki, M.: Simultaneous determination of methamphetamine and its metabolites in the urine samples of abusers by high performance liquid chromatography with chemiluminescence detection. *Biol. Pharm. Bull.*, 16 (1993) 817-821; *C.A.*, 120 (1994) 127076f.
- 3907 Hewala, I.I.: High-performance liquid chromatographic and derivative difference spectrophotometric methods for the determination of acetaminophen and its degradation product in aged pharmaceutical formulations. *Anal. Lett.*, 27 (1994) 561-582.
- 3908 Ho, M.Y.K., Erhard, K.F. and Chen, T.K.: Pre-treatment of chiral α -AGP column with triethylamine significantly improves the detection sensitivity of an enantiomeric leukotriene antagonist. *J. Liq. Chromatogr.*, 17 (1994) 761-772.
- 3909 Jang, W.C. and Lee, H.S.: On-line trace enrichment of mifenidine in plasma using column-switching high performance liquid chromatography. *J. Liq. Chromatogr.*, 17 (1994) 1375-1384.
- 3910 Janssen, P.A.J. and Awouters, F.H.L.: Is it possible to predict the clinical effects of neuroleptics from animal data? Part V: From haloperidol and pimiperone to risperidone. *Arzneim.-Forsch.*, 44 (1994) 269-277.
- 3911 Jin, L. and Lau, C.E.: Determination of alprazolam and its major metabolites in serum microsamples by high-performance liquid chromatography and its application to pharmacokinetics in rats. *J. Chromatogr. B*, 654 (1994) 77-83.
- 3912 Joseph-Charles, J. and Bertucat, M.: Separation, identification and assay of fluradoline hydrochloride in the presence of other tricyclic antidepressants or neuroleptics by spectral and thin-layer and liquid chromatographic methods. *Anal. Chim. Acta*, 284 (1993) 45-52.
- 3913 Kapil, R.P., Lee, W., Jr., Wong, N.Y., Saxton, P.L. and Lam, G.N.: Determination of a novel sigma receptor antagonist, DuP 734, in rat plasma by reversed-phase high-performance liquid chromatography with ultraviolet detection. *J. Chromatogr. B*, 653 (1994) 221-226.
- 3914 Karamanos, N.K., Sivvas, E. and Papaioannou, D.: Rapid assay for the determination of two photoactivatable kainic acid analogues by high-performance liquid chromatography. *J. Liq. Chromatogr.*, 17 (1994) 521-532.
- 3915 Kassahun, K. and Abbott, F.: *In vivo* formation of the thiol conjugates of reactive metabolites of 4-ene VPA and its analog 4-pentenoic acid. *Drug Metab. Disp.*, 21 (1993) 1098-1106.
- 3916 Khetarpal, V.K., Cariola, C.M., Collina, S.J. and Ziemniak, J.A.: Assay of RP 69698, a novel LTB₄-antagonist, in plasma by HPLC and its application for characterizing pharmacokinetics of the drug in dogs. *J. Liq. Chromatogr.*, 17 (1994) 329-346.
- 3917 Knoche, B. and Blaschke, G.: Investigations on the *in vitro* racemization of thalidomide by high-performance liquid chromatography. *J. Chromatogr. A*, 666 (1994) 235-240.
- 3918 Komatsu, S., Murata, S., Aoyama, H., Zenki, T. and Tateishi, M.: High-performance liquid chromatographic determination of triazolam in biological fluids. *Jpn. J. Toxicol. Environ. Health*, 40 (1994) 91-95; *C.A.*, 120 (1994) 263174p.
- 3919 Kristensen, K., Angelo, H.R. and Blemmer, T.: Enantioselective high-performance liquid chromatographic method for the determination of methadone in serum using an AGP and a CN column as chiral and analytical column, respectively. *J. Chromatogr. A*, 666 (1994) 283-287.
- 3920 Lau-Cam, C.A., Rahman, M. and Roos, R.W.: Rapid reverse phase high performance liquid chromatographic assay method for ranitidine hydrochloride in dosage forms. *J. Liq. Chromatogr.*, 17 (1994) 1089-1104.
- 3921 Lauber, R., Mosimann, M., Bührer, M. and Zbinden, A.M.: Automated determination of midazolam in human plasma by high-performance liquid chromatography using column switching. *J. Chromatogr. B*, 654 (1994) 69-75.
- 3922 Lehotay, J., Cizmarik, J., Pham Thi Viet Nga and Bednarikova, A.: Study of local anaesthetics. Part 121: Correlation between $\log k'$, C , R_F , R_M and local anaesthesia for perhydroazepinyl ethyl esters of 2-, 3- and 4-alkoxyphenylcarbamic acids. *Pharmazie*, 49 (1994) 286.
- 3923 Lu, X.-L. and Yang, S.K.: Enantiomer resolution of camazepam and its derivatives and enantioselective metabolism of camazepam by human liver microsomes. *J. Chromatogr. A*, 666 (1994) 249-257.
- 3924 Malavasi, B., Flaminio, L. and Ascalone, V.: Simple, rapid and specific identification and quantification of a metabolite of alpidem, a new imidazopyridine anxiolytic, in human urine, by direct injection into HPLC column with fluorescence detection. *J. Liq. Chromatogr.*, 17 (1994) 419-432.

- 3925 Mannens, G., Huang, M.-L., Meuldermans, W., Hendrickx, J., Woestenborghs, R. and Heykants, J.: Absorption, metabolism, and excretion of risperidone in humans. *Drug Metab. Disp.*, 21 (1993) 1134-1141.
- 3926 Mayol, R.F., Cole, C.A., Colson, K.E. and Kerns, E.H.: Isolation and identification of the major urinary metabolite of *m*-chlorophenylpiperazine in the rat. *Drug Metab. Disp.*, 22 (1994) 171-174.
- 3927 Mayol, R.F., Cole, C.A., Luke, G.M., Colson, K.L. and Kerns, E.H.: Characterization of the metabolites of the antidepressant drug nefazodone in human urine and plasma. *Drug Metab. Disp.*, 22 (1994) 304-311.
- 3928 Meuldermans, W., Hendrickx, J., Mannens, G., Lavrijsen, K., Janssen, C., Bracke, J., le Jeune, L., Lauwers, W. and Heykants, J.: The metabolism and excretion of risperidone after oral administration in rats and dogs. *Drug Metab. Disp.*, 22 (1994) 129-138.
- 3929 Meurisse, R.L. and de Ranter, C.J.: Enantiomer separation of trifluadem and analogues by liquid chromatography with cellulose-based chiral stationary phase. *Chromatographia*, 38 (1994) 629-632.
- 3930 Moachon, G. and Matinier, D.: Simultaneous determination of modafinil and its acid metabolite by high-performance liquid chromatography in human plasma. *J. Chromatogr. B*, 654 (1994) 91-96.
- 3931 Ohno, M., Shiono, Y. and Konishi, M.: Simultaneous determination of dihydrocodeine and its metabolites in dog plasma by high-performance liquid chromatography with electrochemical and ultraviolet detection. *J. Chromatogr. B*, 654 (1994) 213-219.
- 3932 Ohshima, N., Kotaki, H., Sawada, Y. and Iga, T.: Tissue distribution and metabolism of amitriptyline after repeated administration in rats. *Drug Metab. Disp.*, 22 (1994) 21-25.
- 3933 Qin, X.-Z., Ip, D.P., Chang, K.H.-C., Dradransky, P.M., Brooks, M.A. and Sakuma, T.: Pharmaceutical applications of LC-MS. 1 - Characterization of a famotidine degradate in a package screening study by LC-APCI MS. *J. Pharm. Biomed. Anal.*, 12 (1994) 221-233.
- 3934 Rogowsky, D., Marr, M., Long, G. and Moore, C.: Determination of sertraline and desmethylsertraline in human serum using copolymeric bonded-phase extraction, liquid chromatography and gas chromatography-mass spectrometry. *J. Chromatogr. B*, 655 (1994) 138-141.
- 3935 Roy, S.D., Roos, E. and Sharma, K.: Transdermal delivery of buprenorphine through cadaver skin. *J. Pharm. Sci.*, 83 (1994) 126-130.
- 3936 Ryan, T.W.: Identification of barbiturates using high performance liquid chromatography-particle beam EI/CI mass spectroscopy. *J. Liq. Chromatogr.*, 17 (1994) 867-881.
- 3937 Selinger, K., Lanzo, C. and Sekut, A.: Determination of remifentanyl in human and dog blood by HPLC with UV detection. *J. Pharm. Biomed. Anal.*, 12 (1994) 243-248.
- 3938 Speed, W., Parton, A.H., Martin, I.J. and Howard, M.R.: The use of liquid chromatography/thermospray mass spectrometry with online ultraviolet diode array and radiochemical detection: characterization of the putative metabolites of U-78875 in female rat feces. *Biol. Mass Spectrom.*, 23 (1994) 1-5; *C.A.*, 120 (1994) 124029g.
- 3939 Swart, P.J., Bronner, G.M., Bruins, A.P., Ensing, K., Tepper, P.G. and de Zeeuw, R.A.: HPLC-UV atmospheric pressure ionization mass spectrometric determination of the dopamine D2 agonist N-0923 and its major metabolites after oxidative metabolism by rat, monkey-, and human liver microsomes. *Toxicol. Methods*, 3 (1993) 279-290; *C.A.*, 120 (1994) 152857u.
- 3940 Terakita, A., Shibukawa, A., Nakao, C. and Nakagawa, T.: Use of a small-bore internal-surface reversed-phase silica column for the micro high-performance frontal analysis of carbamazepine in an albumin solution with a small sample volume. *Anal. Sci.*, 10 (1994) 11-15; *C.A.*, 120 (1994) 182233e.
- 3941 Yang, S.K.: Base-catalysed rearrangement of temazepam. *J. Pharm. Biomed. Anal.*, 12 (1994) 209-219.
- 3942 Yu, Z., Abdel-Rehim, M. and Westerlund, D.: Determination of amide-type local anaesthetics by direct injection of plasma in a column-switching high-performance liquid chromatographic system using a pre-column with a semipermeable surface. *J. Chromatogr. B*, 654 (1994) 221-230.

For additional information see *C.A.*:

120 (1994) 62458k, 68713m, 86587c, 144319e, 260327e.

See also 3238, 3678, 3844, 3856, 3874, 4017, 4018.

32e. *Chemotherapeutics (exc. cytostatics and antibiotics)*

- 3943 Abdelrahman, A.N., Karim, E.I.A. and Ibrahim, K.E.E.: Determination of chloroquine and its decomposition products in various brands of different dosage forms by liquid chromatography. *J. Pharm. Biomed. Anal.*, 12 (1994) 205-208.
- 3944 Aref, S.O. and Kittaneh, R.B.: High-performance liquid chromatography and spectrophotometric studies on clopidol. *Orient. J. Chem.*, 9 (1993) 1-10; *C.A.*, 120 (1994) 144335g.
- 3945 Balani, S.K., Kaufman, L.R., Arison, B.H., Olah, T.V., Goldman, M.E., Varga, S.L., O'Brien, J.A., Ramjit, H.G., Rooney, C.S., Hoffman, J.M. et al.: Metabolism of 3-[2-(benzoxazol-2-yl)ethyl]-5-ethyl-6-methylpyridin-2(1H)-one (L-696,229), an HIV-1 reverse transcriptase inhibitor, by rat liver, slices and in humans. *Drug Metab. Disp.*, 22 (1994) 200-205.
- 3946 Bekersky, I., Puhl, R.J., Hanson, G. and Mong, S.: The pharmacokinetics of 1,3-di(4-imidazolino-2-methoxyphenoxy)propane.lactate (DMP.lactate), a new agent against opportunistic infections, in male beagle dogs. *Drug Metab. Disp.*, 22 (1994) 233-236.
- 3947 Chaulet, J.-F., Grelaud, G., Bellemin-Magninot, P., Mounier, C. and Brazier, J.-L.: Simultaneous determination of chloroquine, proguanil and their metabolites in human biological fluids by high-performance liquid chromatography. *J. Pharm. Biomed. Anal.*, 12 (1994) 111-117.
- 3948 De Miranda, P. and Burnette, T.C.: Metabolic fate and pharmacokinetics of the acyclovir prodrug valaciclovir in Cynomolgus monkeys. *Drug Metab. Disp.*, 22 (1994) 55-59.
- 3949 Dean, R.A., Ochieng, W., Black, J., Queener, S.F., Bartlett, M.S. and Dumaual, N.G.: Simultaneous determination of primaquine and carboxyprimaquine in plasma using high-performance liquid chromatography with electrochemical detection. *J. Chromatogr. B*, 655 (1994) 89-96.

- 3950 Degroodt, J.M., Wyhowski de Bukanski, B. and Srebrnik, S.: Determination of ivermectin residues in meat and liver by HPLC and fluorometric detection. *J. Liq. Chromatogr.*, 17 (1994) 1419-1426.
- 3951 Doerge, D.R., Bajic, S. and Lowes, S.: Multiresidue analysis of sulfonamides using liquid chromatography with atmospheric pressure chemical ionization mass spectrometry. *Rapid Commun. Mass Spectrom.*, 7 (1993) 1126-1130; *C.A.*, 120 (1994) 53063a.
- 3952 Furusawa, N., Mukai, T. and Itoh, H.: (Analytical method for residual sulfonamides in various tissues of chicken by high-performance liquid chromatography). *Nippon Kakin Gakkaishi*, 30 (1993) 359-364; *C.A.*, 120 (1994) 68706m.
- 3953 Galeano Díaz, T., Lopez Martínez, L., Martínez Galera, M. and Salinas, F.: Rapid determination of nitrofurantoin, furazolidone and furaltadone in formulations, feed and milk by high performance liquid chromatography. *J. Liq. Chromatogr.*, 17 (1994) 457-475.
- 3954 Kaniou, I., Zacariadis, G., Kalligas, G., Tsoukali, H. and Stratis, J.: Separation and determination of carbadox, nitrofurazone, nitrofurantoin, furazolidone, and furaltadone in their mixtures by thin layer and high performance liquid chromatography. *J. Liq. Chromatogr.*, 17 (1994) 1385-1398.
- 3955 Kim, H., Lapiguera, A. and Lin, C.: Gas chromatographic and high-performance liquid chromatographic methods for the determination of genaconazole in biological fluids. *J. Chromatogr. B*, 655 (1994) 21-26.
- 3956 Mizuno, A., Uematsu, T. and Nakashima, M.: Simultaneous determination of ofloxacin, norfloxacin and ciprofloxacin in human hair by high-performance liquid chromatography and fluorescence detection. *J. Chromatogr. B*, 653 (1994) 187-193.
- 3957 Morris, D.M. and Selinger, K.: Determination of 2'-deoxy-3'-thiacytidine (3TC) in human urine by liquid chromatography: direct injection with column switching. *J. Pharm. Biomed. Anal.*, 12 (1994) 255-264.
- 3958 Nebinger, P. and Koel, M.: Determination of serum zidovudine by ultrafiltration and high-performance liquid chromatography. *J. Pharm. Biomed. Anal.*, 12 (1994) 141-143.
- 3959 Sato, Y., Mishima, Y. and Seki, T.: (Rapid determination of synthetic antibacterials by HPLC. Application of Sep-Pak cartridge clean-up to hog kidney). *Sendai-shi Eisei Kenkyushoho 1992*, 22 (1993) 241-247; *C.A.*, 120 (1994) 207967n.
- 3960 Shao, J., Yuan, Z., Nie, H. and Zhang, J.: (Simultaneous determination of veterinary drug residues of 10 sulfonamides in meat by high performance liquid chromatography). *Sepu*, 11 (1993) 373-375; *C.A.*, 120 (1994) 53029u.
- 3961 Uccello-Barretta, G., Pini, D., Rosini, C. and Salvadori, P.: Stereochemical features of the stereoselective interaction between (R)- or (S)-7,7'-bis(1-propen-3-oxy)-2,2'-dihydroxy-1,1'-binaphthyl and quinine. *J. Chromatogr. A*, 666 (1994) 541-548.
- 3962 Valois, M.E.C., Takayanagui, O.M., Bonato, P.S., Lanchote, V.L. and Carvalho, D.: Determination of albendazole metabolites in plasma by HPLC. *J. Anal. Toxicol.*, 18 (1994) 86-90.
- 3963 Wallén, L., Ericsson, Ö., Wikström, I. and Hellgren, U.: High-performance liquid chromatographic method for the enantioselective analysis of mefloquine in plasma and urine. *J. Chromatogr. B*, 655 (1994) 153-157.
- 3964 Webb, L.E., Gechman, M. and Story, K.: Determination of L-2-oxothiazolidine-4-carboxylic acid (Procysteine®) in human plasma by high-performance liquid chromatography. *J. Chromatogr. B*, 654 (1994) 257-262.
- 3965 Zeng, Y.L., Chi, H.T. and Baker, J.K.: HPLC/EC assay of the distribution of the antimalarial arteether into fat and muscle tissue following intravenous administration. *J. Liq. Chromatogr.*, 17 (1994) 1365-1373.

See also 2823, 3689, 3844, 3990.

32f. Cytostatics

- 3966 Bolton, M.G., Hilton, J., Robertson, K.D., Streeper, R.T., Colvin, O.M. and Noe, D.A.: Kinetic analysis of the reaction of melphalan with water, phosphate, and glutathione. *Drug Metab. Disp.*, 21 (1993) 986-996.
- 3967 Brandsteterova, E., Marcincinova, K., Lehotay, J., Zbojova, A. and Halko, J.: HPLC analysis of optical isomers of leucovorin and methotrexate using achiral-chiral system. *Neoplasma*, 40 (1993) 241-245; *C.A.*, 120 (1994) 152850m.
- 3968 Cooper, T.H., Clark, J.G. and Guzinski, J.A.: Analysis of curcuminoids by high-performance liquid chromatography. *ACS Symp. Ser.*, 547 (Food Phytochemicals for Cancer Prevention II) (1994) 231-236; *C.A.*, 120 (1994) 157784c.
- 3969 Corlett, S.A. and Chrystyn, H.: Enantiomeric separation of R- and S-ifosfamide and their determination in serum from clinical subjects. *J. Chromatogr. B*, 654 (1994) 152-158.
- 3970 Cresteil, T., Monsarrat, B., Alvinerie, P., Tréluier, J.M., Vieira, I. and Wright, M.: Taxol metabolism by human liver microsomes: identification of cytochrome P450 isozymes involved in its biotransformation. *Cancer Res.*, 54 (1994) 386-392.
- 3971 Farrish, H.H., Hsyu, P.-H., Pritchard, J.F., Brouwer, K.R. and Jarrett, J.: Validation of a liquid chromatography post-column derivatization assay for the determination of cisplatin in plasma. *J. Pharm. Biomed. Anal.*, 12 (1994) 265-271.
- 3972 Ficarra, R., Ficarra, P., Altavilla, G., Calabro, M.L., Tommasini, S. and Constantino, D.: Analytical profile and release of lisdamine from pharmaceutical formulation. *Acta Technol. Legis Med.*, 4 (1993) 9-20; *C.A.*, 120 (1994) 173618g.
- 3973 Gabizon, A., Catane, R., Uziely, B., Kaufman, B., Safra, T., Cohen, R., Martin, F., Huang, A. and Barenholz, Y.: Prolonged circulation time and enhanced accumulation in malignant exudates of doxorubicin encapsulated in polyethyleneglycol coated liposomes. *Cancer Res.*, 54 (1994) 987-992.
- 3974 Haas, N.B., LaCreta, F.P., Walczak, J., Hudes, G.R., Brennan, J.M., Ozols, R.F. and O'Dwyer, P.J.: Phase I/pharmacokinetic study of topotecan by 24-hour continuous infusion weekly. *Cancer Res.*, 54 (1994) 1220-1226.
- 3975 Hamilton, C.L. and Kirkwood, J.A.: Column-switching high-performance liquid chromatographic method for the determination of a thymidylate synthase inhibitor, LY231514, an investigational agent for the treatment of solid tumors, in human plasma. *J. Chromatogr. B*, 654 (1994) 297-303.
- 3976 King, D.T., Venkateshwaran, T.G. and Stewart, J.T.: HPLC determination of a vincristine, doxorubicin, and ondansetron mixture in 0.9% sodium chloride injection. *J. Liq. Chromatogr.*, 17 (1994) 1399-1411.

- 3977 Kratz, F., Hartmann, M., Keppler, B. and Messori, L.: The binding properties of two antitumor ruthenium(III) complexes to apo-transferrin. *J. Biol. Chem.*, 269 (1994) 2581-2588.
- 3978 Pettit, G.R., Kantoci, D., Herald, D.L., Barkóczy, J. and Slack, J.A.: Procedures for the analyses of dolastatins 10 and 15 by high performance liquid chromatography. *J. Liq. Chromatogr.*, 17 (1994) 191-202.
- 3979 Prat, J., Pujol, M., Girona, V., Muñoz, M. and Solé, L.-A.: Stability of carboplatin in 5% glucose solution in glass, polyethylene and polypropylene containers. *J. Pharm. Biomed. Anal.*, 12 (1994) 81-84.
- 3980 Reichelová, V., Albertioni, F. and Liliemark, J.: Hydrophobicity parameters of 2-chloro-2'-deoxyadenosine and some related analogues and retention in reversed-phase liquid chromatography. *J. Chromatogr. A*, 667 (1994) 37-45.
- 3981 Takimoto, C.H., Klecker, R.W., Dahut, W.L., Yee, L.K., Strong, J.M., Allegra, C.J. and Grem, J.L.: Analysis of the active lactone form of 9-aminocamptothecin in plasma using solid-phase extraction and high-performance liquid chromatography. *J. Chromatogr. B*, 655 (1994) 97-104.
- 3982 Van Tellingen, O., Beijnen, J.H., Nooijen, W.J. and Bult, A.: Tissue disposition, excretion and metabolism of vinblastine in mice as determined by high-performance liquid chromatography. *Cancer Chemother. Pharmacol.*, 32 (1993) 286-292; *C.A.*, 120 (1994) 152895e.
- 3983 Zucchetti, M., de Fusco, M., Sessa, C., Fröhlich, A., Reichert, S. and D'Incalci, M.: High-performance liquid chromatographic assay for the determination of the novel podophyllotoxin derivative dimethylaminoetoposide (NK611) in human plasma. *J. Chromatogr. B*, 654 (1994) 97-102.
- For additional information see *C.A.*:
120 (1994) 144299y.
- See also 3762.
- 32g. *Other drug categories*
- 3984 Blaih, S.M. and Galal, S.M.: Determination of allopurinol and benzbromarone in a two-component mixture using high-performance liquid chromatography and derivative spectrophotometry. *S. T. P. Pharma Sci.*, 3 (1993) 408-412; *C.A.*, 120 (1994) 173622d.
- 3985 Bonet-Domingo, E., Torres-Lapasió, J.R., Medina-Hernández, M.J. and García-Alvarez-Coque, M.C.: Chromatographic monitoring of diuretics in urine samples using a sodium dodecyl sulphate-propranolol micellar eluent. *Anal. Chim. Acta*, 287 (1994) 201-210.
- 3986 Campíns-Falcó, P., Herráez-Hernández, R. and Sevillano-Cabeza, A.: Application of column-switching techniques to the determination of medium polarity drugs: determination of acetazolamide in urine. *J. Chromatogr. B*, 654 (1994) 85-90.
- 3987 Carlucci, G., Mazzeo, P., Vetuschi, C. and di Giuseppe, E.: Reversed-phase high-performance liquid chromatographic and derivative UV spectrophotometric determination of famotidine in pharmaceutical preparations. *Int. J. Pharm.*, 102 (1994) 271-274; *C.A.*, 120 (1994) 144350h.
- 3988 Dal Bo, L., Broccali, G.P., Silingardi, S. and Coppi, G.: A new HPLC method for pidotimod plasma levels determination. *Boll. Chim. Farm.*, 132 (1993) 126-128; *C.A.*, 120 (1994) 68736w.
- 3989 Dwivedi, A.K., Sirkar, K.P., Bhatt, G.R., Seth, R.K., Singh, S. and Sarin, J.P.S.: Determination of *cis*- and *trans*-centchroman in its dosage forms by high-performance liquid chromatography. *J. Chromatogr. A*, 663 (1994) 187-190.
- 3990 Egginger, G., Blaschke, E., Lindner, W. and Olsson, A.-M.: Stereoselective high-performance liquid chromatographic assay of (\pm)-delmopinol in plasma using solid-phase extraction, a chiral derivatizing agent and electrochemical detection. *J. Chromatogr. A*, 666 (1994) 275-282.
- 3991 Farthing, D., Sica, D.A., Fakhry, I. and Gehr, T.W.B.: Novel high-performance liquid chromatographic method using solid-phase on-line elution for determination of metolazone in plasma and whole blood. *J. Chromatogr. B*, 653 (1994) 171-176.
- 3992 Fischer, V., Vickers, A.E.M., Heitz, F., Mahadevan, S., Baldeck, J.-P., Minery, P. and Tynes, R.: The polymorphic cytochrome P-4502D6 is involved in the metabolism of both 5-hydroxytryptamine antagonists, tropisetron and ondasetron. *Drug Metab. Disp.*, 22 (1994) 269-274.
- 3993 Jones, A.: High-performance liquid chromatographic determination of iophenoxic acid in serum. *J. Chromatogr. B*, 654 (1994) 293-296.
- 3994 Kiriya, A., Mimoto, T., Kisanuki, S., Kiso, Y. and Takada, K.: Comparison of a new orally potent tripeptide HIV-1 protease inhibitor (anti-AIDS drug) based on pharmacokinetic characteristics in rats after intravenous and intraduodenal administrations. *Biopharm. Drug Dispos.*, 14 (1993) 697-707; *C.A.*, 120 (1994) 94631c.
- 3995 Kiss, T., Mrs. and Kovacs Hadady, K.: (Analytical and stability studies carried out during the formulation development of Heviz ointment). *Acta Pharm. Hung.*, 63 (1993) 215-221; *C.A.*, 120 (1994) 62447f.
- 3996 Mandrou, B. and Blanchin, M.D.: (Determination of cinnaine in an ointment by liquid chromatography). *Analisis*, 21 (1993) 409-413; *C.A.*, 120 (1994) 173584t.
- 3997 Moncrieff, J.: Determination of dapsone in serum and saliva using reversed-phase high-performance liquid chromatography with ultraviolet or electrochemical detection. *J. Chromatogr. B*, 654 (1994) 103-110.
- 3998 Mulligan, T.E., Blain, R.W., Oldfield, N.F. and Mico, B.A.: The analysis of Ro 24-4736 in human plasma by multidimensional reversed phase microbore HPLC/UV. *J. Liq. Chromatogr.*, 17 (1994) 133-150.
- 3999 Naidong, W., Lee, J.W. and Hulse, J.D.: Development and validation of a high-performance liquid chromatographic method for the determination of methocarbamol in human plasma. *J. Chromatogr. B*, 654 (1994) 287-292.
- 4000 Nishikawa, T., Hasumi, H., Suzuki, S., Kubo, H. and Ohtani, H.: Kinetic analysis of molecular interconversion of immunosuppressant FK506 by high-performance liquid chromatography. *Pharm. Res.*, 10 (1993) 1785-1789; *C.A.*, 120 (1994) 45470r.
- 4001 Ratiney, R., Astier, A. and Chariot, P.: Determination of fenoferine in tissue samples by high-performance liquid chromatography. *J. Chromatogr. B*, 654 (1994) 159-163.
- 4002 Rozman, E., Galcerán, M.T., Anglada, L. and Albet, C.: Metabolites of ebrotidine, a new H₂-receptor antagonist, in human urine. *J. Pharm. Sci.*, 83 (1994) 252-254.
- 4003 Sakakihara, Y., Nakamura, G., Tokoeda, Y., Abe, T. and Kamoshita, S.: A rapid microassay for dichloroacetate in serum by gel-permeation chromatography. *Eur. J. Clin. Chem. Clin. Biochem.*, 32 (1994) 79-83.

- 4004 Shum, Y.Y., Black, A. and Chang, T.: Determination of 2,2-dimethyl-N-(2,4,6-trimethoxyphenyl) dodecanamide, CI-976, in rat plasma by reversed-phase high-performance liquid chromatography. *J. Chromatogr. B*, 653 (1994) 205-209.
- 4005 Sugioka, N., Odani, H., Ohta, T., Kishimoto, H., Yasumura, T. and Takada, K.: Determination of a new immunosuppressant, mycophenolate mofetil, and its active metabolite, mycophenolic acid, in rat and human body fluids by high-performance liquid chromatography. *J. Chromatogr. B*, 654 (1994) 249-256.
- 4006 Trenerry, V.C., Wells, R.J. and Robertson, J.: The analysis of illicit heroin seizures by capillary zone electrophoresis. *J. Chromatogr. Sci.*, 32 (1994) 1-6.
- 4007 Vree, T.B., van den Biggelaar-Marteau, M. and Verwey-van Wissen, C.P.W.G.M.: Determination of furesemide with its acyl glucuronide in human plasma and urine by means of direct gradient high-performance liquid chromatographic analysis with fluorescence detection. Preliminary pharmacokinetics and effect of probenecid. *J. Chromatogr. B*, 655 (1994) 53-62.
- 4008 Wang, K., Blain, R.W. and Szuna, A.J.: Multidimensional narrow bore liquid chromatography analysis of Ro 24-0238 in human plasma. *J. Pharm. Biomed. Anal.*, 12 (1994) 105-110.
- 4009 Wilke, T.L., Tien Huy, N. and Tyle, P.: HPLC method development for AG-85 and benzyl alcohol in anhydrous topical formulations. *Int. J. Pharm.*, 103 (1994) 187-197; *C.A.*, 120 (1994) 173620b.
- 4010 Zeng, J.-n., Dou, L., Duda, M. and Stuting, H.H.: New chiral high-performance liquid chromatographic methodology used for the pharmacokinetic evaluation of dexfenfluramine. *J. Chromatogr. B*, 654 (1994) 231-248.
- For additional information see *C.A.*:
120 (1994) 62444c, 62452d, 94555f, 173623e.
- See also 2963, 3158, 3201, 3770, 3844, 3865, 3897, 3901, 4025, 4134.
- 32h. *Toxicological and forensic applications*
- 4011 Byrd, G.D., Uhrig, M.S., deBethizy, J.D., Caldwell, W.S., Crooks, P.A., Ravard, A. and Riggs, R.M.: Direct determination of cotinine-N-glucuronide in urine using thermospray liquid chromatography/mass spectrometry. *Biol. Mass Spectrom.*, 23 (1994) 103-107; *C.A.*, 120 (1994) 184762a.
- 4012 Dickey, R.W., Granade, H.R. and Bencsath, F.A.: Improved analytical methodology for the derivatization and HPLC-fluorometric determination of okadaic acid in phytoplankton and shellfish. *Dev. Mar. Biol.*, 3(Toxic Phytoplankton Blooms in the Sea) (1993) 495-499; *C.A.*, 120 (1994) 237647a.
- 4013 Fernández, P., Rodríguez, P., Bermejo, A.M., López-Rivadulla, M. and Cruz, A.: Simultaneous determination of cocaine and benzoylecgonine in vitreous humor by HPLC. *J. Liq. Chromatogr.*, 17 (1994) 883-890.
- 4014 Guevremont, R. and Quigley, M.N.: Determination of paralytic shellfish poisons using liquid chromatography. *J. Chem. Educ.*, 7 (1994) 80-82; *C.A.*, 120 (1994) 243565p.
- 4015 Guitton, J., David, O., Mialon, A. and Manchon, M.: (Evaluation of an automated liquid chromatography system for the toxicological screening of biological fluids). *Ann. Biol. Clin.*, 51 (1993) 611-617; *C.A.*, 120 (1994) 70996z.
- 4016 Huber, L.: Toxicological applications of diode array detection in HPLC. *Chromatogr. Sci. Ser.*, 62(Diode Array Detection in HPLC) (1993) 221-239; *C.A.*, 120 (1994) 47436h.
- 4017 Kintz, P.: Determination of buprenorphine and its dealkylated metabolite in human hair. *J. Anal. Toxicol.*, 17 (1993) 443-444; *C.A.*, 120 (1994) 235229s.
- 4018 Longo, M., Martines, C., Rolandi, L. and Cavallaro, A.: Simple and fast determination of some phenethylamines in illicit tablets by base-deactivated reversed-phase HPLC. *J. Liq. Chromatogr.*, 17 (1994) 649-658.
- 4019 Qin, Y., Xu, Y. and Zhang, C.: (Determination of some thermolabile dope agents by high-performance liquid chromatography). *Fenxi Huaxue*, 22 (1994) 84-86; *C.A.*, 120 (1994) 263171k.
- 4020 Sato, S., Ogata, T. and Kodama, M.: Wide distribution of toxins with sodium channel blocking activity similar to tetrodotoxin and paralytic shellfish toxins in marine animals. *Dev. Mar. Biol.*, 3(Toxic Phytoplankton Blooms in the Sea) (1993) 429-434; *C.A.*, 120 (1994) 237902e.
- 4021 Stabell, O.B., Pedersen, K. and Aune, T.: Detection and separation of toxins accumulated by mussels during the 1988 bloom of *Chrysochromulina polylepis* in Norwegian coastal waters. *Mar. Environ. Res.*, 36 (1993) 185-196; *C.A.*, 120 (1994) 127033q.
- 4022 Tai, S.S.-C., Christensen, R.G., Paule, R.C., Sander, L.C. and Welch, M.J.: The certification of morphine and codeine in a human urine standard reference material. *J. Anal. Toxicol.*, 18 (1994) 7-12.
- 4023 Teass, A.W., DeBord, D.G., Brown, K.K., Cheever, K.L., Stettler, L.E., Savage, R.E., Weigel, W.W., Dankovic, D. and Ward, E.: Biological monitoring for occupational exposures to o-toluidine and aniline. *Int. Arch. Occup. Environ. Health*, 65(1, Suppl., Proceedings of International Symposium on Biological Monitoring, 1992) (1993) S115-S118; *C.A.*, 120 (1994) 156000g.
- 4024 Veress, T.: Study of the extraction of LSD from illicit blotters for HPLC determination. *J. Forensic Sci.*, 38 (1993) 1105-1110; *C.A.*, 120 (1994) 98625b.
- 4025 White, V., Canfield, D.V. and Hordinsky, J.: The identification and quantitation of triamterene in blood and urine from a fatal aircraft accident victim. *J. Anal. Toxicol.*, 18 (1994) 52-53.
- See also 3161, 3645, 3721, 3823, 3847, 3894, 3906, 3918, 3956.
- 32i. *Plant extracts*
- 4026 Artaud, J., Falque, V., Archavlis and Derbesy, M.: (Identification and determination of sclareol epimers). *Parfums, Cosmet., Aromes*, 112 (1993) 79-82; *C.A.*, 120 (1994) 61907u.
- 4027 Bader, G., Tujja, D., Wray, V. and Hiller, K.: New triterpenoid saponins from *Heteropappus altaicus* (Willd.) Noropokr. and *Heteropappus biennis* (Ldb.) Tamamsch. *Pharmazie*, 49 (1994) 209-212.
- 4028 Gu, G., Zhou, L. and Zhao, G.: (Improved high performance liquid chromatographic method for the analysis of ginsenosides in Panax ginseng extracts and products the chromatographic modifier effect of ammonium acetate in separation of ginsenosides). *Sepu*, 12 (1994) 40-42; *C.A.*, 120 (1994) 186408p.
- 4029 Kamal-Eldin, A., Appelqvist, L.Å. and Yousif, G.: Lignan analysis in seed oils from four *Sesamum* species: comparison of different chromatographic methods. *J. Am. Oil Chem. Soc.*, 71 (1994) 141-147.

- 4030 Ko, H.-C., Tsai, T.-H., Chou, C.-J., Hsu, S.-Y., Li, S.-Y. and Chen, C.-F.: High-performance liquid chromatographic determination of rutaecarpine in rat plasma: application to a pharmacokinetic study. *J. Chromatogr. B*, 655 (1994) 27-31.
- 4031 Kraehenbuehl, S., Hasler, F. and Krapf, R.: Analysis and pharmacokinetics of glycyrrhizic acid and glycyrrhetic acid in humans and experimental animals. *Steroids*, 59 (1994) 121-126; *C.A.*, 120 (1994) 207765v.
- 4032 Oleszek, W.: Alfalfa (*Medicago sativa* L.) root saponins - chemical structure, biological activity, determination). *Pamięt. Pulawski 1990*, 97 (1992) 201-209; *C.A.*, 120 (1994) 49190x.
- 4033 Park, M.K., Park, J.H., Lee, M.Y., Kim, S.J. and Park, I.J.: Micro-analysis of ginseng saponins by ion chromatography with pulsed amperometric detection. *J. Liq. Chromatogr.*, 17 (1994) 1171-1182.
- 4034 Raggi, M.A., Maggei, F., Bugamelli, F. and Cantelli Forti, G.: Bioavailability of glycyrrhizin and licorice extract in rat and human plasma as detected by a HPLC method. *Pharmazie*, 49 (1994) 269-272.
- 4035 Rücker, G., Neugebauer, M. and Kiefer, A.: Quantitative Bestimmung von α -Peroxyachifolid in der Schafgarbe durch HPLC mit amperometrischer Detektion. *Pharmazie*, 49 (1994) 167-169.
- 4036 Schütt, H. and Hölzl, J.: Vergleichende Qualitätsuntersuchung von Johanniskraut-Fertigarzneimitteln unter Verwendung verschiedener quantitativer Bestimmungsmethoden. *Pharmazie*, 49 (1994) 206-209.
- 4037 Sharma, A., Singh, R.T. and Handa, S.S.: Determination of phyllanthin and hypophyllanthin by high-performance liquid chromatography in *Phyllanthus amarus*. *Phytochem. Anal.*, 4 (1993) 226-229; *C.A.*, 120 (1994) 173588x.
- 4038 Slanina, J., Taborska, E. and Musil, P.: (Determination of cynarine in the decoctions of the artichoke (*Cynara cardunculus* L.) by the HPLC method). *Cesko-Slov. Farm.*, 42 (1993) 275-278; *C.A.*, 120 (1994) 239347b.
- 4039 Tarantilis, P.A., Polissiou, M. and Manfait, M.: Separation of picrocrocin, *cis-trans*-crocins and safranal of saffron using high-performance liquid chromatography with photodiode-array detection. *J. Chromatogr. A*, 664 (1994) 55-61.
- 4040 Tsai, T.-H., Chou, C.-J., Cheng, F.-C. and Chen, C.-F.: Pharmacokinetics of honokiol after intravenous administration in rats assessed using high-performance liquid chromatography. *J. Chromatogr. B*, 655 (1994) 41-45.
- 4041 Wang, W., Saito, K. and Kano, Y.: On the evaluation of the preparation of Chinese medicinal prescriptions. VII. HPLC analysis of the components in Inchinkoto and Inchingoreisan. *Shoyakugaku Zasshi*, 47 (1993) 243-248; *C.A.*, 120 (1994) 116906a.
- 4042 Wei, F., Chen, M., Hou, Y. and Sun, Y.: (Determination of gossypol in cottonseed oil by high performance liquid chromatography). *Sepu*, 11 (1993) 363-364; *C.A.*, 120 (1994) 53026r.
- For additional information see *C.A.*:
120 (1994) 62079n, 62402n, 62411q, 86554q, 86559v, 86562r, 144294t, 144303v, 173188s.
- See also 2903, 3209, 3877, 3968.
33. CLINICO-CHEMICAL APPLICATIONS
- 33a. *General papers and reviews*
- 4043 De Antonis, K.M. and Brown, P.R.: Determination of biochemical markers by HPLC and CE for the diagnosis of diseases. *Am. Lab. (Shelton)*, 25 (1993) 62-71; *C.A.*, 120 (1994) 186335n - a review with 35 refs.
- 4044 Radeck, W.: (Analysis of physiological fluids - solid phase extraction and HPLC-determination). *Labor-Med.*, 16 (1993) 97-103; *C.A.*, 120 (1994) 211948z.
- 33b. *Complex mixtures and profiling (single compounds by cross-reference only)*
- See 3071, 3072, 3074, 3142, 3164, 3165, 3184, 3196, 3197, 3210, 3211, 3228, 3244, 3255, 3265, 3278, 3282, 3284, 3286, 3430, 3437, 3449, 3474, 3498, 3550, 3564, 3615, 3689, 3690, 3692, 3747, 3753.
34. FOOD ANALYSIS
- See 3743.
- 34a. *General papers and reviews*
- 4045 Almarcha, M.M. and Rovira, L.J.: (Determination of additives and organic residues in food by chromatographic techniques (HRGC and HPLC)). *Aliment. Equipos Tecnol.*, 12 (1993) 85-93; *C.A.*, 120 (1994) 52901k - a review with no refs.
- 4046 Doneche, B.: *Recent Achievements in Wine Chromatography: Applications to the Sensory Analysis of Wines*, 1993, 256 pp.; *C.A.*, 120 (1994) 268786z.
- 4047 Huber, L. and Schuster, R.: Food and beverages. *Chromatogr. Sci. Ser.*, 62(Diode Array Detection in HPLC) (1993) 293-324; *C.A.*, 120 (1994) 75702k - a review with 27 refs.
- See also 3828.
- 34b. *Complex mixtures (single compounds by cross-reference only)*
- 4048 Bianco, L. and Marucchi, M.: (New chlorine-releasing disinfectants: trichloroisocyanuric acid and its salts. Determination by HPLC). *Ind. Aliment. (Pinerolo)*, 32 (1993) 817-824; *C.A.*, 120 (1994) 75748e.
- 4049 Cioni, F. and Bartolucci, G.: (Methods of analysis for natural products with insecticidal activity). *Boll. Chim. Ig., Parte Sci.*, 44 (1993) 293-303; *C.A.*, 120 (1994) 161858x.
- 4050 De Koning, P.J., van den Bedem, J.W. and Olieman, C.: Characterization of dried milk according to heat treatment and usage - HPLC method. *Bull. Int. Dairy Fed.*, 285 (1993) 46-52; *C.A.*, 120 (1994) 52971h.
- 4051 Garbarino, M.: (Determination of thickening and gelling agents in food by anion-exchange chromatography with pulsed amperometric detection). *Boll. Chim. Ig., Parte Sci.*, 44 (1993) 99-108; *C.A.*, 120 (1994) 105233c.

4052 Mattivi, F.: Solid-phase extraction of trans-resveratrol from wines for HPLC analysis. *Z. Lebensm.-Unters. Forsch.*, 196 (1993) 522-525; C.A., 120 (1994) 52962f.

For additional information see C.A.:

120 (1994) 53028t, 53052w, 105480f, 268781w.

See also 2848, 3017, 3035, 3038, 3039, 3041, 3048, 3059, 3062, 3085, 3138, 3139, 3146, 3168, 3174, 3179, 3183, 3219, 3223, 3232, 3253, 3274, 3482, 3484, 3485, 3636, 3695, 3728, 3732, 3739, 3746, 3756, 3768, 3786, 3798, 3805, 3950, 3951, 3952, 3953, 3960, 4021, 4039, 4084, 4117, 4119, 4124.

34c. *Organoleptically important compounds (flavors, odors, volatiles)*

4053 Taylor, S.: Improved determination of vanillin and related phenolic components in vanilla (*Vanilla fragrans*) (Salisb. Ames) by high-performance liquid chromatography. *Fragrance J.*, 8 (1993) 281-287; C.A., 120 (1994) 242738k.

See also 4039.

35. ENVIRONMENTAL ANALYSIS

35a. *General papers and reviews*

4054 Chmil, V.D.: (The use chromatographic parameters for prediction of the toxicity of substances and their fate in biosphere). *Gig. Sanit.*, (1993) 57-60; C.A., 120 (1994) 171701e - a review with 21 refs.

4055 Sergeev, S.K. and Prokhorova, E.K.: (Present status of computer chromatography for ecological and sanitary analytical monitoring). *Zavod. Lab.*, 59 (1993) 1-6; C.A., 120 (1994) 171702f - a review with 3 refs.

See also 2937, 3006.

35b. *Air pollution (complex mixtures; single compounds by cross-reference only)*

See 3057.

35c. *Water pollution (complex mixtures; single compounds by cross-reference only)*

4056 Golovnya, R.N. and Terenina, M.B.: (Effect of concentration of organic compounds in analysis of water samples by high efficiency capillary chromatography). *Zh. Anal. Khim.*, 48 (1993) 1921-1933; C.A., 120 (1994) 172714y.

4057 Genu, S. and Hennion, M.-C.: On-line sample handling of water-soluble organic pollutants in aqueous samples using porous graphitic carbon. *J. Chromatogr. A*, 665 (1994) 243-251.

4058 Okutani, T., Yamakawa, K., Sakuragawa, A. and Gotoh, R.: Determination of a microamount of sulphide by ion chromatography with amperometric detection after coprecipitation with basic zinc carbonate. *Anal. Sci.*, 9 (1993) 731-734; C.A., 120 (1994) 172534q.

See also 2849, 2895, 2964, 3032, 3033, 3058, 3220, 3784, 3799, 3801, 3804, 3806, 3811, 3819, 4060, 4067, 4088, 4095.

35d. *Soil pollution (complex mixtures; single compounds by cross-reference only)*

4059 Snyder, J.L., Grob, R.L., McNally, M.E. and Oostdyk, T.S.: A different approach - using solid-phase disks and cartridges to extract organochlorine and organophosphate pesticides from soils. *LC-GC*, 12 (1994) 230-242; C.A., 120 (1994) 243261m.

See also 3784, 3807, 3816.

36. SOME TECHNICAL PRODUCTS AND COMPLEX MIXTURES

For additional information see C.A.:

120 (1994) 63257f.

36a. *Surfactants*

4060 Evans, K.A., Dubey, S.T., Kravetz, L., Dzidic, I., Gumulka, J., Mueller, R. and Stork, J.R.: Quantitative determination of linear primary alcohol ethoxylate surfactants in environmental samples by thermospray LC/MS. *Anal. Chem.*, 66 (1994) 699-705.

4061 Márquez, N., Antón, R.E., Usabillaga, A. and Selager, J.L.: Optimization of HPLC conditions to analyze widely distributed ethoxylated alkylphenol surfactants. *J. Liq. Chromatogr.*, 17 (1994) 1147-1169.

4062 Walton, B.: Analysis of anionic detergents using reversed-phase HPLC and a mass detector with ammonium as the counterion. *LC-GC Int.*, 7 (1994) 149-156.

4063 Wang, Z. and Fingas, M.: Analysis of sorbitan ester surfactants. Part I: High performance liquid chromatography. *J. High Resolut. Chromatogr.*, 17 (1994) 15-19.

For additional information see C.A.:

120 (1994) 61800d.

36b. *Antioxidants and preservatives*

4064 Wu, K., Zhang, W., Addis, P.B., Epley, R.J., Salih, A.M. and Lehrfeld, J.: Antioxidant properties of wild rice. *J. Agric. Food Chem.*, 42 (1994) 34-37.

4065 Yang, S.S., Lee, J.J., Lee, S.C., Su, S.J. and Chou, S.S.: (Simultaneous determination of five synthetic antioxidants in fats and oils). *Yaowu Shipin Fenxi*, 1 (1993) 287-296; C.A., 120 (1994) 75770f.

For additional information see C.A.:

120 (1994) 215539q.

See also 3733.

36c. *Complex mixtures, technical products and unidentified compounds*

- 4066 Kientz, C.E., Langenberg, J.P. and Brinkman, U.A.T.: Microcolumn liquid chromatography with thermionic detection of the enantiomers of O-ethyl S-2-diisopropylaminoethyl methylphosphonothioate (VX). *J. High Resolut. Chromatogr.*, 17 (1994) 95-97.
- 4067 Kwade, C., Voigtländer, R. and Cammann, K.: HPLC/MS-screening of waste water samples from use of laquers. *Fresenius J. Anal. Chem.*, 348 (1994) 520-522.
- 4068 Lu, W. and Li, T.: (Application of ion chromatography technique in rock and mineral analysis). *Yankuang Ceshi*, 11 (1992) 268-272; *C.A.*, 120 (1994) 152305n - a review with 41 refs.
- 4069 Sakodynskii, K.I.: Chromatography in the chemical industry. *Khim. Prom-st. (Moscow)*, (1993) 283-287; *C.A.*, 120 (1994) 94337m - a review with 2 refs.

For additional information see *C.A.*:

120 (1994) 152573y, 166664r.

See also 3021, 3023, 3024, 3155, 3461, 3834.

37. CELLS, CELLULAR PARTICLES AND SUPRAMOLECULAR STRUCTURES

- 4070 Cardot, P.J.P., Elgéa, C., Guernet, M., Godet, D. and Andreux, J.P.: Size- and density-dependent elution of normal and pathological red blood cells by gravitational field-flow fractionation. *J. Chromatogr. B*, 654 (1994) 193-203.
- 4071 Lu, M., Albertsson, P.-Å., Johansson, G. and Tjerneld, F.: Partitioning of proteins and thylakoid membrane vesicles in aqueous two-phase systems with hydrophobically modified dextran. *J. Chromatogr. A*, 668 (1994) 215-228.
- 4072 Ollero, M., Pascual, M.L., Muiño-Blanco, T., Cebrián-Pérez, J.A. and López-Pérez, M.J.: Revealing surface changes associated with maturation of ram spermatozoa by centrifugal counter-current distribution in an aqueous two-phase system. *J. Chromatogr. A*, 668 (1994) 173-178.
- 4073 Pinilla, M., de la Fuente, J., García-Pérez, A.I., Jimeno, P., Sancho, P. and Luque, J.: Biochemical characterization of human erythrocytes fractionated by counter-current distribution in aqueous polymer two-phase systems. *J. Chromatogr. A*, 668 (1994) 165-171.
- 4074 Seddas, A., Meignoz, R., Daire, X., Boudon-Padieu, E. and Caudwell, A.: Purification of grapevine flavescence dorée MLO (mycoplasma-like organism) by immunoaffinity [chromatography]. *Curr. Microbiol.*, 27 (1993) 229-236; *C.A.*, 120 (1994) 158002h.
- 4075 Stefánsson, H., Wollenberger, L. and Albertsson, P.-Å.: Fragmentation and separation of the thylakoid membrane. Effect of light-induced protein phosphorylation on domain composition. *J. Chromatogr. A*, 668 (1994) 191-200.
- 4076 Turánek, J.: Fast-protein liquid chromatography system as a tool for liposome preparation by the extrusion procedure. *Anal. Biochem.*, 218 (1994) 352-357.

- 4077 Walter, A., Lesieur, S., Blumenthal, R. and Ollivon, M.: Size characterization of liposomes by HPLC. In: Gregoriadis, G. (Editor), *Liposome Technol. (2nd ed.)*, Vol. 1, CRC, Boca Raton, 1993, pp. 271-290; *C.A.*, 120 (1994) 239266z - a review with 34 refs.
- 4078 Walter, H. and Widen, K.E.: Cell partitioning in two-polymer aqueous phase systems and cell electrophoresis in aqueous polymer solutions. Red blood cells from different species. *J. Chromatogr. A*, 668 (1994) 185-190.
- 4079 Warnock, D.E.: Quantitative isolation of endosome and plasma membranes by high-gradient magnetic affinity chromatography. Avail. *Univ. Microfilms Int.*, Order No. DA9317407, 1992, 163 pp.; *C.A.*, 120 (1994) 211706u.

See also 3464.

38. INORGANIC COMPOUNDS

See 3423.

38a. Cations

- 4080 Abolt, R.B. and Knepper, G.D.: Separation of metal ions on an anion-exchange resin by chromatographic elution. *U.S. US* 5,246,591 (Cl. 210-656; B01D15/08), 21 Sep. 1993, US Appl. 553,444, 13 Jul. 1990; 14 p.; *C.A.*, 120 (1994) 82311k.
- 4081 Alumaa, P. and Pentšuk, J.: Prediction of chromatographic retention of metal complexes from stability constant measured by ion chromatography. *Chromatographia*, 38 (1994) 566-570.
- 4082 Baechmann, K., Steeg, K.H., Groh, T., Haumann, I., Boden, J. and Holthues, H.: Analysis for cations and anions in small volumes using micro-LC and CE. *J. Microcolumn Sep.*, 4 (1992) 431-438; *C.A.*, 120 (1994) 123733v.
- 4083 Basova, E.M., Bondareva, L.G. and Ivanov, V.M.: (Some azoxines as possible analytical reagents for platinum group and non-ferrous metals). *Zh. Anal. Khim.*, 48 (1993) 1590-1598; *C.A.*, 120 (1994) 123625m.
- 4084 Branch, S., Ebdon, L. and O'Neill, P.: Determination of arsenic species in fish by directly coupled high-performance liquid chromatography-inductively coupled plasma mass spectrometry. *J. Anal. At. Spectrom.*, 9 (1994) 33-36; *C.A.*, 120 (1994) 156038a.
- 4085 Brown, A.A., Ebdon, L. and Hill, S.J.: Development of a coupled liquid chromatography-isotope dilution inductively coupled plasma mass spectrometry method for lead speciation. *Anal. Chim. Acta*, 286 (1994) 391-399.
- 4086 Dauchy, X., Point-Gautier, M., Astruc, A. and Astruc, M.: Analytical methods for the speciation of selenium compounds: a review. *Fresenius J. Anal. Chem.*, 348 (1994) 792-805.
- 4087 De Beer, H. and Coetzee, P.P.: Vanadium separation by ion chromatography. *Fresenius J. Anal. Chem.*, 348 (1994) 806-809.
- 4088 Di Cintio, R. and de Carolis, G.: (High-performance liquid chromatography (HPLC) for determination of cobalt, nickel, and copper in seawater). *Inquinamento*, 35 (1993) 58-61; *C.A.*, 120 (1994) 143504z.

- 4089 Dycbczynski, R., Danko, B. and Maleszewska, H.: Separation scheme for selective and quantitative isolation of cobalt from neutron-irradiated biological materials by ion exchange and extraction chromatography. *Zh. Anal. Khim.*, 49 (1994) 37-44; C.A., 120 (1994) 265265t.
- 4090 Fedotov, P.S., Maryutina, T.A., Pichugin, A.A. and Spivakov, B.Ya.: (Effect of extraction system kinetic properties on element separations by liquid chromatography with a free mobile phase). *Zh. Neorg. Khim.*, 38 (1993) 1878-1884; C.A., 120 (1994) 68107s.
- 4091 Gong, Z., Song, J. and Luo, J.: (Application of decompressed chromatographic microcolumn to analysis of ores-photometric determination of trace gold after preconcentration and separation with P-350 extraction resin). *Yankuang Ceshi*, 12 (1993) 189-193; C.A., 120 (1994) 68147e.
- 4092 Goyal, S.S., Hafez, A.A.R. and Rains, D.W.: Simultaneous determination of total sodium, potassium, magnesium, and calcium in plant tissues using acid digestion and ion chromatography. *Agron. J.*, 85 (1993) 1192-1197; C.A., 120 (1994) 265227g.
- 4093 Hu, W. and Haraguchi, H.: Direct determination of sodium, potassium, magnesium, and calcium ions in human saliva by ion chromatography using a taurine-conjugated bile salt micelle-coated stationary phase. *Anal. Chim. Acta*, 289 (1994) 231-236.
- 4094 Hu, W. and Haraguchi, H.: Simultaneous separation of inorganic cations and anions by ion chromatography using a single column coated with weak/strong-charged zwitterionic bile salt micelles. *Anal. Chem.*, 66 (1994) 765-767.
- 4095 Hwang, C.-j. and Jiang, S.-J.: Determination of arsenic compounds in water samples by liquid chromatography-inductively coupled plasma mass spectrometry with an *in situ* nebulizer-hydride generator. *Anal. Chim. Acta*, 289 (1994) 205-213.
- 4096 Ivask, J. and Pentchuk, J.: (Increasing the selectivity in the ion chromatographic determination of sodium, ammonium, and potassium ions. Effect of amino acids as eluent modifiers). *Eesti Tead Akad. Toim., Keem.*, 41 (1992) 33-35; C.A., 120 (1994) 94152x.
- 4097 Jones, P., Challenger, O.J. and Hill, S.J.: Recent advances in high performance chelation ion chromatography for trace metal determinations. *Spec. Publ.-R. Soc. Chem.*, 122 (1993) 279-287; C.A., 120 (1994) 94060r - a review with 15 refs.
- 4098 Karandashev, V.K., Grazhulene, S.S. and Sadykov, I.I.: Extraction-chromatographic system TBP-HBr in radiochemical neutron activation analysis of high-purity materials. *Microchim. Acta*, 112 (1993) 77-86; C.A., 120 (1994) 181921r.
- 4099 Lee, D.W., Kim, K.S., Park, Y.H., Czea, M.Z. and Chung, K.S.: (Development of separation and trace analysis methods for platinum group elements. Separations and retention behavior of platinum metal acetylacetonates in reversed-phase liquid chromatography). *Anal. Sci. Technol.*, 6 (1993) 107-119; C.A., 120 (1994) 68099r.
- 4100 Lucy, C.A. and Dinh, H.N.: Kinetics and equilibria of the Zn-EDTA-PAR postcolumn reaction detection system for the determination of alkaline earth metals. *Anal. Chem.*, 66 (1994) 793-797.
- 4101 Manjarekar, S.V. and Chavan, J.D.: Cation-exchange chromatographic separation of bismuth(III) in mixed and nonaqueous solvent systems. *Orient J. Chem.*, 9 (1993) 71-75; C.A., 120 (1994) 207462u.
- 4102 Milacic, R. and Stupar, J.: Simultaneous determination of chromium(III) complexes and chromium(VI) by fast protein anion exchange liquid chromatography-atomic absorption spectrometry. *Analyst (London)*, 119 (1994) 627-632.
- 4103 Nesterenko, P.N., Amirova, G.Zh. and Bol'shova, T.A.: Separation of metal ions by reversed-phase liquid chromatography using on-column complexation with 2-pyridinecarboxylic acid. *Anal. Chim. Acta*, 285 (1994) 161-168.
- 4104 Park, Y.S., Han, S.H., Joe, K.S., Eom, T.Y. and Lee, G.H.: Retention behavior of transition metal ions with some complexing agents on cation exchanger. *Bull. Korean Chem. Soc.*, 14 (1993) 692-696; C.A., 120 (1994) 152379q.
- 4105 Seubert, A. and Meinke, R.: On-line coupled ion chromatography-ICP-(AES,MS) and electro thermal vaporisation-ICP-MS in use for ultra trace analysis of high purity rhenium. *Fresenius J. Anal. Chem.*, 348 (1994) 510-519.
- 4106 Song, J.M., Lee, C. and Chung, K.S.: A quantitative determination of overlapped chromatographic peaks of dysprosium and yttrium using target transformation factor analysis. *Bull. Korean Chem. Soc.*, 15 (1994) 74-78; C.A., 120 (1994) 207472x.
- 4107 Sun, D., Brungs, M., Trimm, D. and Navratil, J.D.: Separation and recovery of nickel and molybdenum using continuous rotating annular chromatography. *Sep. Sci. Technol.*, 29 (1994) 831-843; C.A., 120 (1994) 82230h.
- 4108 Tanaka, K., Oota, K. and Matsushita, S.: Simultaneous determination of anions and cations by ion chromatography. *Jpn. Kokai Tokkyo Koho JP 05,223,798 (93,223,798) (Cl. G01N30/02)*, 31 Aug. 1993, Appl. 92/58,937, 12 Feb. 1992; 6 p.; C.A., 120 (1994) 94329k.
- 4109 Tsai, S.-J.J. and Hsu, S.-J.: Speciation of vanadium(V) and vanadium(IV) with 4-(2-pyridylazo)-resorcinol by using high-performance liquid chromatography with spectrophotometric detection. *Analyst (London)*, 119 (1994) 403-407.
- 4110 Vieira, N.E., Yergey, A.L. and Abrams, S.A.: Extraction of magnesium from biological fluids using 8-hydroxyquinoline and cation-exchange chromatography for isotopic enrichment analysis using thermal ionization mass spectrometry. *Anal. Biochem.*, 218 (1994) 92-97.
- 4111 Wagh, D.N., Kumar, S.D. and Iyer, C.S.P.: Determination of tungsten in low-grade tungsten ores by dual-column ion chromatography. *Anal. Chim. Acta*, 287 (1994) 229-234.
- 4112 Weiss, J., Jensen, D. and Saweet, A.: (Ion chromatography and capillary electrophoresis for ion analysis). *LaborPraxis*, 17 (1993) 36-43; C.A., 120 (1994) 94067y - a review with 11 refs.
- 4113 Wood, D.J., Elshani, S., Wai, C.M., Bartsch, R.A., Huntley, M. and Hartenstein, S.: Column chromatographic separation of Y^{3+} from Sr^{2+} by polymeric ionizable crown ether resins. *Anal. Chim. Acta*, 284 (1993) 37-43.

For additional information see C.A.:
120 (1994) 45066p, 152377n, 152575a.

See also 2942, 3722, 3725, 3726.

38b. Anions

- 4114 Alumaa, P. and Pentsuk, J.: Separation of oxoanions by ion chromatography using anion-exchange resin HIKS-1. *Eesti Tead Akad. Toim. Keem.*, 42 (1993) 149-152; C.A., 120 (1994) 181793a.

- 4115 Bhatnagar, R.P. and Mishra, D.D.: Distribution studies of halides and thiocyanate ions in aqueous acetone media. *Asian J. Chem.*, 5 (1993) 902-906; *C.A.*, 120 (1994) 201326f.
- 4116 Bondoux, G. and Jones, T.: Analysis of ppb levels of anions and cations in deionized water. *LC-GC Int.*, 7 (1994) 29-34.
- 4117 Draisci, R., Boria, P., Lucentini, L. and Cavalli, S.: (Ion chromatography of nitrites and nitrates in meat products). *Riv. Soc. Ital. Sci. Aliment.*, 21 (1992) 415-421; *C.A.*, 120 (1994) 215544n.
- 4118 Gennaro, M.C., Abrigo, C. and Marengo, E.: Retention of anions as a function of mobile phase pH in ion-interaction RP-HPLC. *J. Liq. Chromatogr.*, 17 (1994) 1231-1244.
- 4119 Li, D., Guan, X., Song, S. and Xie, T.: (Simultaneous determination of chloride and sulfur in rice by ion chromatography). *Sepu*, 11 (1993) 377-379; *C.A.*, 120 (1994) 53030n.
- 4120 Liu, X. and Yun, Z.: (Determination of thiocyanate anion by α -bromo-2,3,4,5,6-pentafluorotoluene derivatization and high-performance liquid chromatography). *Fenxi Huaxue*, 21 (1993) 1199-1201; *C.A.*, 120 (1994) 70981r.
- 4121 Michigami, Y. and Ueda, K.: Sulphite stabilizer in ion chromatography. *J. Chromatogr. A*, 663 (1994) 255-258.
- 4122 Michigami, Y., Fujii, K. and Ueda, K.: Effect of functional groups on the retention behaviour of anions in ion chromatography using a coated silica column. *J. Chromatogr. A*, 664 (1994) 117-122.
- 4123 Muenchow, V. and Steudel, R.: The decomposition of aqueous dithionite and its reactions with polythionates $S_nO_6^{2-}$ ($n = 3-5$) studied by ion-pair chromatography. *Z. Anorg. Allg. Chem.*, 620 (1994) 121-126; *C.A.*, 120 (1994) 207372q.
- 4124 Ruiz, E., Santillana, M.I., de Alba, M., Nieto, M.T. and Garcia-Castellano, S.: High performance ion chromatography determination of total sulfites in foodstuffs. *J. Liq. Chromatogr.*, 17 (1994) 447-456.
- 4125 Sakai, H., Fujiwara, T. and Kumamaru, T.: Determination of silicate by ion-exclusion chromatography with luminol chemiluminescence detection. *Bull. Chem. Soc. Jpn.*, 66 (1993) 3401-3406; *C.A.*, 120 (1994) 68257r.
- 4126 Scheller, U., Siedenberg, D. and Schügerl, K.: Ion chromatography for monitoring biotechnological processes. *Anal. Chim. Acta*, 285 (1994) 169-176.
- 4127 Scheller, U., Siedenberg, D., Hübner, U., Siebold, M., Kretzmer, G. and Schügerl, K.: Ion chromatography for monitoring biotechnological processes. Part II. Applications for industrial cultivation media. *Anal. Chim. Acta*, 286 (1994) 179-187.
- 4128 Wang, Y., Zhao, G., Gu, M., Shron, J. and Yan, Z.: Ion chromatographic determination of phosphates in castables. *Fresenius J. Anal. Chem.*, 348 (1994) 700-702.
- 4129 Yoza, N., Nakashima, S., Ueda, N., Miyajima, T., Nakamura, T. and Vast, P.: High-performance liquid chromatographic characterization of monofluorophosphate, difluorophosphate and hexafluorophosphate. *J. Chromatogr. A*, 664 (1994) 111-116.
- 4130 Zhu, Y., Xu, S. and Xu, H.: (Ion chromatographic determination of fluorine). *Lihua Jianyan, Huaxue Fence*, 29 (1993) 60-61; *C.A.*, 120 (1994) 152290d - a review with 30 refs.

For additional information see *C.A.*:

120 (1994) 68164h, 123665z, 152486x, 152599m, 239397f.

See also 4058, 4082, 4094, 4108, 4112.

38c. Permanent and rare gases

- 4131 Twombly, B., Brzychcy, A.M., Albert, P. and Flaherty, E.: Ion chromatographic analysis of semiconductor gases. *Am. Lab. (Shelton)*, 25 (1993) 36F-36K; *C.A.*, 120 (1994) 181811e.

38d. Volatile inorganic compounds

- 4132 Willis, J.D., Dowle, C.J., Malyan, A.P., Liversidge, R.D. and Cullen, P.: Determination of low levels nitrogen oxides in gas streams by high-performance liquid chromatographic determination of the Saltzman complex. *Analyst (London)*, 119 (1994) 389-392.

39. RADIOACTIVE AND OTHER ISOTOPE COMPOUNDS

- 4133 Das, N.R., Singh, K. and Lahiri, S.: Liquid-liquid extraction and reversed phase chromatographic separation of parent-daughter, ^{95}Zr - ^{95}Nb with TOA. *Radiochim. Acta*, 62 (1993) 213-216; *C.A.*, 120 (1994) 202525g.
- 4134 Riley, A.L.M. and Nowotnik, D.P.: The determination of charge of cationic ^{99m}Tc -radiopharmaceuticals. *J. Liq. Chromatogr.*, 17 (1994) 533-548.

See also 3269.

Gas Chromatography

1. REVIEWS AND BOOKS

- 1255 Wittkowski, R.: Fundamental aspects of capillary gas chromatography. In: Wittkowski, R. and Matisek, R. (Editors), *Capillary Gas Chromatogr. Food Control Res.*, Technomic, Lancaster, 1993, pp. 19-49; C.A., 119 (1993) 194727m - a review with 20 refs.

See also 1281, 1285, 1300, 1305, 1381, 1382, 1395, 1427, 1435, 1436, 1439, 1444, 1445, 1509, 1510, 1511, 1570, 1883, 1884, 1886, 1893, 1901, 1946, 1947.

2. FUNDAMENTALS, THEORY AND GENERAL

2a. General

- 1256 Ettre, L.S.: New, unified nomenclature for chromatography. *Chromatographia*, 38 (1994) 521-526.

- 1257 Hrouzek, J., Hatrik, Š., Krupčík, J. and Benická, E.: Evaluation of separation quality with a novel threshold criterion. *J. Chromatogr. A*, 665 (1994) 3-8.

See also 1387, 1431, 1432, 1456.

2b. Thermodynamics and theoretical relationships

- 1258 Poole, S.K., Kollie, T.O. and Poole, C.F.: Influence of temperature on the mechanism by which compounds are retained in gas-liquid chromatography. *J. Chromatogr. A*, 664 (1994) 229-251.

- 1259 Synakh, V.S.: (Solutions for steady state problems gas chromatography.) *Prikl. Mekh. Tekh. Fiz.*, No. 3 (1993) 172-176; C.A., 119 (1993) 206051b.

- 1260 Wang, F., Tian, Q., Guo, Y. and Cao, T.: The theoretical study on intersection point rule for homologues in gas chromatography. *Chin. Chem. Lett.*, 4 (1993) 795-796; C.A., 120 (1994) 133346k.

- 1261 Xu, G., Zhang, Y. and Lu, P.: (Restricting factors in gas chromatography.) *Yingyong Huaxue*, 10, No. 5 (1993) 66-71; C.A., 120 (1994) 123839j.

See also 1354, 1448.

2c. Relationship between structure and chromatographic behaviour

- 1262 Hilal, S.H., Carreira, L.A., Karickhoff, S.W. and Melton, C.M.: Estimation of gas-liquid chromatographic retention times from molecular structure. *J. Chromatogr. A*, 662 (1994) 269-280.

- 1263 Stankova, I.N., Beresnev, A.N., Agureev, V.G. and Lanin, S.N.: (Relation between the structure and the retention characteristics of phosphorus-containing organic compounds on SE-30 liquid stationary phase.) *Zh. Fiz. Khim.*, 66 (1992) 3321-3328.

See also 1359, 1448, 1477, 1519, 1604.

2d. Measurement of physico-chemical and related values

- 1264 Bandosz, T.J., Putyera, K., Jagiello, J. and Schwarz, J.A.: Application of inverse gas chromatography to the study of the surface properties of modified layered minerals. *Microporous Mater.*, 1 (1993) 73-79; C.A., 119 (1993) 274440c.

- 1265 Bonifaci, L., Carnelli, L. and Cori, L.: Determination of infinite dilution diffusion and activity coefficients of solvents in polystyrene by inverse gas chromatography on a capillary column. *J. Appl. Polym. Sci.*, 51 (1994) 1923-1930; C.A., 120 (1994) 165531h.

- 1266 Chehimi, M.M., Abel, M.-L., Pigois-Landureau, E. and Delamar, M.: Characterization of conducting polymers by inverse gas chromatography. Part II. Effect of dopant on the dispersive and specific properties of polypyrrole. *Synth. Met.*, 60 (1993) 183-194; C.A., 120 (1994) 9560m.

- 1267 Cortesi, A., Fermeglia, M., Kikic, I. and Spicka, B.: On the use of supercritical fluid chromatography for the determination of thermophysical properties. *Lat. Am. Appl. Res.*, 22 (1992) 125-133; C.A., 120 (1994) 137892h.

- 1268 Dernovaya, L.I., Chalych, A.E. and Eltekov, Yu.A.: Adsorption properties of surface iron oxide modified with stearic acid and octadecylamine. *Pure Appl. Chem.*, 65 (1993) 2201-2204; C.A., 120 (1994) 15927c.

- 1269 Deutsch, J.C., Santhosh-Kumar, C.R., Hassell, K.L. and Kolhouse, J.F.: Variation in ascorbic acid oxidation routes in H₂O₂ and cupric ion solution as determined by GC/MS. *Anal. Chem.*, 66 (1994) 345-350.

- 1270 Dutta, S., Chakraborty, S.S., Mandal, B.M. and Bhattacharyya, S.N.: Polymer-polymer miscibility prediction from polymer (1) analogue probe-polymer (2) exchange interaction parameter using inverse gas chromatography: the miscibility of poly(vinyl methyl ether) with poly(vinyl esters). *Polymer*, 34 (1993) 3499-3503; C.A., 119 (1993) 226907p.

- 1271 Etxeberria, A., Elorza, J.M., Iruin, J.J., Marco, C., Gomez, M.A. and Fatou, J.G.: Miscibility of poly(vinyl chloride)/poly(ethylene oxide) blends. II. An inverse gas chromatography study. *Eur. Polym. J.*, 29 (1993) 1483-1487; C.A., 119 (1993) 272268d.

- 1272 Glass, A.S. and Larsen, J.W.: Inverse gas chromatography of glassy polymer surfaces. *Macromolecules*, 26 (1993) 6354-6358; C.A., 119 (1993) 250936t.

- 1273 Glass, A.S. and Larsen, J.W.: Surface thermodynamics for non-polar adsorbates on Illinois No. 6 coal by inverse gas chromatography. *Energy Fuels*, 7 (1993) 994-1000; C.A., 119 (1993) 229711f.

- 1274 Goenenc, Z.S., Yildirim, R., Beler-Baykal, A.B. and Oensan, Z.I.: Adsorption parameters of some C₅-C₈ paraffinic hydrocarbons on platinum-alumina by gas chromatographic pulse techniques. *Appl. Catal. A*, 103 (1993) 35-42; C.A., 119 (1993) 234711k.

- 1275 Guitart, R.: (Solubility of six gases in blood and in several liquid media.) *Rev. Esp. Fisiol.*, 49 (1993) 195-202; C.A., 120 (1994) 128749q.

- 1276 Habboush, A.E., Farroha, S.M. and Kreishan, A.-L.Y.: Gas-liquid chromatographic study of thermodynamics of solution of some alkanes on liquid crystal stationary phases. *J. Chromatogr. A*, 664 (1994) 71-76.
- 1277 Hegedus, C.R. and Kamel, I.L.: Thermodynamic analysis of pigment and polymer surfaces using inverse gas chromatography. *J. Coat. Technol.*, 65, No. 820 (1993) 31-43; *C.A.*, 120 (1994) 79472h.
- 1278 Hu, Q., Huang, D. and Wang, S.: (Prediction of ternary liquid-liquid equilibria by measuring the activity coefficients with gas chromatographic headspace analysis.) *Huagong Xuebao (Chin. Ed.)*, 44 (1993) 192-198; *C.A.*, 120 (1994) 174669m.
- 1279 Hua, G., Lu, Z., Zhang, Q., Lu, F. and Zhang, S.: (Study on the method of absorbing nitrogen oxide (NO_x) with 2-octanol by GC/MS.) *Huanjing Huaxue*, 12 (1993) 111-115; *C.A.*, 119 (1993) 255740j.
- 1280 Kamdem, D.P., Bose, S.K. and Luner, P.: Inverse gas chromatography characterization of birch wood meal. *Langmuir*, 9 (1993) 3039-3044; *C.A.*, 119 (1993) 252364d.
- 1281 Katsanos, N.A.: Studies of diffusion and other rate processes by gas chromatography. *Pure Appl. Chem.*, 65 (1993) 2245-2252; *C.A.*, 119 (1993) 278978g - a review with 30 refs.
- 1282 Kontominas, M.G., Gavara, R., Giacini, J.R.: The adsorption of hydrocarbons on polystyrene by inverse gas chromatography: infinite dilution concentration region. *Eur. Polym. J.*, 30 (1994) 265-269; *C.A.*, 120 (1994) 135704z.
- 1283 Kunaver, M., Guthrie, J.T. and Kamin, F.: The application of inverse gas chromatography in paint formulation. *Surf. Coat. Int.*, 76, No. 2 (1993) 62-69; *C.A.*, 119 (1993) 273434s.
- 1284 Kuruc, J., Sahoo, M.K. and Kubinec, R.: Radiolysis and photolysis of nitroaromatic compounds - halogenoalkane mixtures. Part X. GC-FTIR-MS analysis of volatile radiolytic products in the radiolysis of nitroaniline - carbon tetrachloride solutions. *J. Radioanal. Nucl. Chem.*, 176 (1993) 185-197; *C.A.*, 120 (1994) 44811r.
- 1285 Larionov, O.G., Belyakova, L.D., Vinogradova, N.I., Vorobjeva, L.M., Petrenko, V.V. and Platonova, N.P.: Chromatographic methods in solid surface investigations. *Pure Appl. Chem.*, 65 (1993) 2237-2243; *C.A.*, 120 (1994) 15617b - a review with 23 refs.
- 1286 Li, Q., Li, J., Zhang, Z. and Huang, K.: (Online GC system for catalytic kinetic studies. 1. Characterization of the kinetic performances of silver catalysts in the epoxidation of ethylene.) *Jisuanji Yu Yingyong Huaxue*, 9 (1992) 234-236; *C.A.*, 119 (1993) 202838r.
- 1287 Liu, C.: (Determination of the sojourn time of adsorption on gas-solid surface by gas chromatography.) *Wuli Huaxue Xuebao*, 9 (1993) 489-494; *C.A.*, 119 (1993) 257039m.
- 1288 Liu, L. and Guo, C.: (Application of gas chromatography in the cytochrome P-450 monooxygenase model system. I. Quantitative analysis and application in the kinetic study of the monooxygenation of cyclohexane by metalloporphyrin.) *Fenxi Huaxue*, 21 (1993) 470-473; *C.A.*, 120 (1994) 100318z.
- 1289 Makarova, M.A., Paukshtis, E.A., Thomas, J.M., Williams, C. and Zamaraev, K.I.: *In situ* FTIR and GC kinetic studies: complementary methods in the mechanistic study of butanol dehydration on zeolite HZSM 5. *Stud. Surf. Sci. Catal.*, 75 (1993) 1711-1714; *C.A.*, 120 (1994) 7962v.
- 1290 Malbrunot, P., Vidal, D. and Vermese, J.: A new apparatus for the measurement of physical adsorption of gas mixtures at elevated pressures. *Rev. Sci. Instrum.*, 64 (1993) 1984-1988.
- 1291 Matsuyama, Y., Yasuoka, T. and Mitsuzawa, S.: Visualization of chemical reaction with aid of gas chromatograph. Hydrolysis of ethyl acetate. *Proc. Sch. Sci. Tokai Univ.*, 28 (1993) 49-52; *C.A.*, 119 (1993) 249329r.
- 1292 Mattie, D.R., Bates, G.D., Jr., Jepson, G.W., Fisher, J.W. and McDougal, J.N.: Determination of skin/air partition coefficients for volatile chemicals: experimental method and applications. *Fundam. Appl. Toxicol.*, 22 (1994) 51-57; *C.A.*, 120 (1994) 156026v.
- 1293 Mukhopadhyay, P. and Schreiber, H.P.: Inverse gas chromatography for polymer surface characterization above and below T_g. *Macromolecules*, 26 (1993) 6391-6396; *C.A.*, 119 (1993) 251897t.
- 1294 Nikolakaki, S., Vassilakos, C. and Katsanos, N.A.: Chromatographic determination of the rate and extent of absorption of air pollutants by sea water. *Chromatographia*, 38 (1994) 191-198.
- 1295 Papirer, E., Balard, H. and Sidqi, M.: The grafting of perfluorinated silanes onto the surface of silica: characterization by inverse gas chromatography. *J. Colloid Interface Sci.*, 159 (1993) 238-242; *C.A.*, 120 (1994) 15779f.
- 1296 Pyda, M., Sidqi, M., Keller, D.S. and Luner, P.: An inverse gas chromatographic study of calcium carbonate treated with alkylketene dimer. *Tappi J.*, 76, No. 4 (1993) 79-85; *C.A.*, 119 (1993) 183145u.
- 1297 Rhue, R.D., Pennell, K.D., Reve, W.H. and Hornsby, A.G.: Sorption of para-xylene vapors on salt-treated soils measured by flow-equilibration and gas chromatography methods. *J. Environ. Qual.*, 22 (1993) 521-527; *C.A.*, 120 (1994) 30050e.
- 1298 Scholten, A.B., Janssen, H.-G., de Haan, J.W. and Cramers, C.A.: A study of the adsorption activities of silanol surface structures on a fused silica model substrate by combining ²⁹Si CP MAS NMR and inverse gas chromatographic data. *J. High Resolut. Chromatogr.*, 17 (1994) 77-84.
- 1299 Simpson, E.J., Abukhadra, R.K., Koros, W.J. and Schechter, R.S.: Sorption equilibrium isotherms for volatile organics in aqueous solution. Comparison of head-space gas chromatography and on-line UV stirred cell results. *Ind. Eng. Chem. Res.*, 32 (1993) 2269-2276; *C.A.*, 119 (1993) 187789e.
- 1300 Tarasevich, Y.I., Bondarenko, S.V., Zhukova, A.I., Nazarenko, A.V. and Aksenenko, E.V.: Chromatographic and molecular statistical studies of structural and energy characteristics of surface micropores of silicate and silica adsorbents. *Pure Appl. Chem.*, 65 (1993) 2257-2264; *C.A.*, 120 (1994) 15618c - a review with 15 refs.
- 1301 Veillon, C., Patterson, K.Y., Nagey, D.A. and Tehan, A.M.: Measurement of blood volume with an enriched stable isotope of chromium (⁵³Cr) and isotope dilution by combined gas chromatography-mass spectrometry. *Clin. Chem. (Washington)*, 40 (1994) 71-73.
- 1302 Voelkel, A., Szymanowski, J., Meissner, E. and Myszkowski, J.: Polarity of fluorine-containing oxyethylates as determined by inverse gas chromatography. *J. Fluorine Chem.*, 64 (1993) 177-186; *C.A.*, 119 (1993) 228518m.
- 1303 Vrentas, J.S., Vrentas, C.M. and Romdhane, I.H.: Analysis of inverse gas chromatography experiments. *Macromolecules*, 26 (1993) 6670-6672; *C.A.*, 119 (1993) 250905g.

- 1304 Wakharkar, R.D., Biswas, S.S., Borate, H.B. and Ponde, D.E.: Monitoring the conversion of cycloheptanone to methyl 7-oxoheptanoate by gas chromatography. *J. Chromatogr. A*, 662 (1994) 420-423.
- 1305 Wang, M.J. and Wolff, S.: Surface energy of carbon black. In: Donnet, J.-B., Bansal, R.C. and Wang, M.-J. (Editors), *Carbon Black (2nd Ed.)*, M. Dekker, New York, 1993, pp. 229-243; *C.A.*, 120 (1994) 15764x.
- 1306 Weidenhamer, J.D., Macias, F.A., Fischer, N.H. and Williamson, G.B.: Just how insoluble are monoterpenes? *J. Chem. Ecol.*, 19 (1993) 1799-1807; *C.A.*, 120 (1994) 8750m.
- 1307 Xie, L.Q.: Measurement of solute diffusion in polymers by inverse gas chromatography using fused-silica open tubular columns. *Polymer*, 34 (1993) 4579-4584; *C.A.*, 120 (1994) 31903x.
- 1308 Zenkevich, I.G.: (Chromatographic characteristics of organic reactions based on the additivity of changes in gas chromatographic retention parameters for reagents and products.) *Zh. Org. Khim.*, 28 (1992) 1827-1840; *C.A.*, 119 (1993) 225327n.
- See also 1380, 1470, 1528, 1787, 1788, 1790, 1791, 1801, 1807.
- ### 3. GENERAL TECHNIQUES
- #### 3a. Apparatus and accessories
- 1309 Anon.: (Method and apparatus for IR-pyrolytic preparation of samples for GC/MS.) *Ger. Offen.* DE 4,206,109 (Cl. G01N1/28), 2 Sep. 1993, Appl. 27 Feb. 1992; 13 pp.; *C.A.*, 120 (1994) 123789t.
- 1310 Asleson, G.L., Doig, M.T. and Heldrich, F.J.: Applications of autosampling GC-MS in an introductory organic chemistry laboratory. *J. Chem. Educ.*, 70 (1993) A290-A294; *C.A.*, 120 (1994) 76556c.
- 1311 Blanch, G.P., Ibáñez, E., Herraiz, M. and Regiero, G.: Use of a programmed temperature vaporizer for off-line SFE/GC analysis in food composition studies. *Anal. Chem.*, 66 (1994) 888-892.
- 1312 Durand, J.P. and Boscher, Y.: Automatic capillary GC analyzer for commercial gasoline characterization. *Chim. Oggi*, 11, No. 7-8 (1993) 49-51; *C.A.*, 119 (1993) 274695q.
- 1313 Flores, P. and Bellar, T.A.: Determination of volatile organic compounds in soils using equilibrium headspace analysis and capillary column gas chromatography/mass spectrometry. Evaluation of the Tekmar 7000 HA analyzer. *Report*, EPA/600/R-93/032; Order No. PB93-155992, 1993, 68 pp.; *C.A.*, 120 (1994) 76329f.
- 1314 Gerstel, E. and Bremer, R.: (Gas-chromatographic method and device for analyzing gaseous raw material.) *Ger.* DE 4,132,358 (Cl. G01N30/02), 26 Nov. 1992, Appl. 27 Sep. 1991; 4 pp.; *C.A.*, 120 (1994) 94428s.
- 1315 Grob, K. and Munari, F.: Process and device for vaporization injections in equipment for gas chromatographic analysis. *Eur. Pat. Appl.* EP 551,847 (Cl. G01N30/12), 21 Jul. 1993, IT Appl. 92/MI49, 14 Jan. 1992; 14 pp.; *C.A.*, 119 (1993) 194812k.
- 1316 Keen, B. and Otteman, J.: Minimizing volume of vaporizing regulators speeds analysis time. *Adv. Instrum. Control*, 48 (1993) 711-717; *C.A.*, 120 (1994) 167390k.
- 1317 Kovalev, L.V. and Denisova, L.V.: (Chromatograph for determination of trace impurities of oxygen, carbon dioxide, methane, and carbon monoxide in inert gases.) *U.S.S.R.* SU 1,755,179 (Cl. G01N30/08), 15 Aug. 1992, Appl. 4,854,499, 24 Jul. 1990; *C.A.*, 119 (1993) 285426d.
- 1318 Lasser, R., Grieveson, B., Hemmerich, J.L., Stagg, R., Dowhlyuk, T., Torr, K., Massey, R. and Chambers, P.: The gas-chromatographic analysis system in the JET active gas handling plant. *Rev. Sci. Instrum.*, 64 (1993) 2449-2458.
- 1319 Manura, J.J.: Trapping of volatiles at the front of the GC capillary column using a GC cryotrapping device. *Am. Lab. (Shelton)*, 26, No. 5 (1994) 32PP-32RR.
- 1320 Matsui, M.: (Gas chromatograph for determination of hydrogen and water in organic compounds.) *Jpn. Kokai Tokkyo Koho* JP 04,120,463 [92,120,463] (Cl. G01N30/88), 21 Apr. 1992, Appl. 90/240,355, 11 Sep. 1990; 3 pp.; *C.A.*, 119 (1993) 285539t.
- 1321 Mori, H.: (Gas chromatograph for simultaneous determination of inorganic and organic compounds.) *Jpn. Kokai Tokkyo Koho* JP 05,164,750 [93,164,750] (Cl. G01N30/46), 29 Jun. 1993, Appl. 91/332,280, 16 Dec. 1991; 5 pp.; *C.A.*, 119 (1993) 194816q.
- 1322 Okhotnikov, B.P., Simongauz, S.E., Vashchun, S.S., Iliina, N.B., Malisova, A.D., Kepke, V.L., Ulanov, Yu.N., Pyatkovskij, V.N. and Ruchnov, V.I.: (Chromatograph thermostat for approximate analysis of gas-air mixtures.) *U.S.S.R.* SU 1,803,858 (Cl. G01N30/30), 23 Mar. 1993, Appl. 4,865,190, 17 Jul. 1990; *C.A.*, 120 (1994) 152568a.
- 1323 Poshemanskij, V.M. and Skorniyakov, E.P.: (Capillary gas chromatograph.) *U.S.S.R.* SU 1,741,060 (Cl. G01N30/02), 15 Jun. 1992, Appl. 4,762,210, 27 Nov. 1989; *C.A.*, 119 (1993) 285546t.
- 1324 Sacks, R.D. and Peters, A.J.: Mechanical gas-chromatograph injection valves and column multiplexing techniques. *U.S.* US 5,205,845 (Cl. B01D15/08), 27 Apr. 1993, Appl. 590,174, 28 Sep. 1990; 14 pp.; *C.A.*, 119 (1993) 216526z.
- 1325 Skorniyakov, E.P. and Poshemanskij, V.M.: (Capillary gas chromatograph.) *U.S.S.R.* SU 1,741,061 (Cl. G01N30/02), 15 Jun. 1992, Appl. 4,762,219, 27 Nov. 1989; *C.A.*, 119 (1993) 285545s.
- 1326 Skorniyakov, E.P., Fisejskij, Yu.K. and Poshemanskij, V.M.: (Gas-chromatograph method for analysis of substance mixtures using pure solvent vapors.) *U.S.S.R.* SU 1,734,003 (Cl. G01N30/06), 15 May 1992, Appl. 4,291,207, 29 Jul. 1987; *C.A.*, 119 (1993) 240782a.
- 1327 Snyder, A.P., Smith, P.B.W., Dworzanski, J.P. and Meuzelaar, H.L.C.: Pyrolysis-gas chromatography-mass spectrometry. Detection of biological warfare agents. In: *Mass Spectrometry for the Characterization of Microorganisms. ACS Symp. Ser.*, Vol. 541, ACS, Washington, 1994, pp. 62-84; *C.A.*, 120 (1994) 72654k.
- 1328 Ueda, H. and Hiromoto, M.: Expired gas analytical method and device. *Eur. Pat. Appl.* EP 574,027 (Cl. G01N30/46), 15 Dec. 1993, JP Appl. 92/178,994, 12 Jun. 1992; 11 pp.; *C.A.*, 120 (1994) 101243h.
- 1329 Valcárcel, M., Ballesteros, E. and Gallego, M.: Continuous liquid-liquid extraction and derivatization module coupled on-line with gas chromatographic detection. *TrAC*, 13 (1994) 68-73.
- 1330 Wang, M., Shen, R. and Wen, Z.: (Gas-chromatograph apparatus for continuous sampling and analysis of methane.) *Faming Zhuanli Shenqing Gongkai Shuomingshu* CN 1,063,362 (Cl. G01N30/24), 5 Aug. 1992, Appl. 92,100,938, 24 Feb. 1992; 12 pp.; *C.A.*, 119 (1993) 194797j.

- 1331 Zelikman, A.M., Davydenkov, A.K. and Lipavskij, V.N.: (Gas chromatograph with unchanging sensitivity.) *U.S.S.R. SU* 1,741,063 (Cl. G01N30/46), 15 Jun. 1992, Appl. 4,184,813, 22 Mar. 1990; *C.A.*, 119 (1993) 285543q.
- See also 1290, 1396, 1405, 1458, 1459, 1698, 1912, 1956, 1957, 1961, 1964.
- 3b. *Detectors and detection reagents*
- 1332 Arnold, M., Cammann, K. and Kleinböhmer, W.: The plasma emission detector (PED) as an element-specific detector for supercritical fluid chromatography. *Fresenius J. Anal. Chem.*, 348 (1994) 546-548.
- 1333 Felt, K. and Low, N.: A gas chromatograph detector system for high-confidence identification. *Int. Lab.*, 24, No. 2 (1994) 17-19.
- 1334 Fujinari, E.M.: Gas and liquid chromatography chemiluminescent nitrogen detection (CLND) techniques and optimization. *Dev. Food Sci.*, 32 (1993) 31-53; *C.A.*, 119 (1993) 199048a.
- 1335 Ilyasov, L.V.: (Thermoconductometric detection of gases in chromatography.) *U.S.S.R. SU* 1,803,861 (Cl. G01N30/62), 23 Mar. 1993, Appl. 4,798,646, 5 Mar. 1990; *C.A.*, 120 (1994) 152574z.
- 1336 Juvancz, Z., Cserháti, T., Markides, K.E., Bradshaw, J.S. and Lee, M.L.: Characterization of some new polysiloxane stationary phases by principal component analysis. *Chromatographia*, 38 (1994) 227-231.
- 1337 Lasa, J., Drozdowicz, B. and Sliwka, I.: A theoretical model of the electron capture detector. *Chromatographia*, 38 (1994) 304-312.
- 1338 Nishina, T., Kabasawa, A., Matsue, T. and Uchida, I.: Gas sensing using microring-ring electrodes. Application to a gas chromatographic detector. *Proc. Electrochem. Soc.*, 93-97 (1993) 8-14; *C.A.*, 120 (1994) 94409m.
- 1339 Pedersen-Bjergaard, S. and Greibrokk, T.: On-column atomic emission detection in capillary gas chromatography using a radio frequency plasma. *J. Microcolumn Sep.*, 6 (1994) 11-18.
- 1340 Popp, P., Oppermann, G., Leonhardt, J., Baer, H. and Merten, H.: (Noble gas ionization detector.) *Ger. (East) DD* 301,774 (Cl. G01N27/62), 16 Dec. 1993, Appl. 326,053, 27 Feb. 1989; 6 pp.; *C.A.*, 120 (1994) 182029t.
- 1341 Ratfisch, W.: (Flame ionization detector apparatus.) *Ger. Offen. DE* 4,136,413 (Cl. G01N27/62), 6 May 1993, Appl. 5 Nov. 1991; 5 pp.; *C.A.*, 119 (1993) 194801f.
- 1342 Ratfisch, W.: (Flame ionization detector.) *Ger. Offen. DE* 4,136,412 (Cl. G01N27/62), 6 May 1993, Appl. 5 Nov. 1991; 4 pp.; *C.A.*, 119 (1993) 194802g.
- 1343 Ratfisch, W.: (Flame-ionization detector and its operation.) *Ger. Offen. DE* 4,138,660 (Cl. G01N27/62), 9 Jun. 1993, Appl. 25 Nov. 1991; 4 pp.; *C.A.*, 119 (1993) 194810h.
- 1344 Ratfisch, W.: (Flame-ionization detector with second glow plug.) *Ger. Offen. DE* 4,138,060 (Cl. G01N27/62), 3 Jun. 1993, Appl. 19 Nov. 1991; 4 pp.; *C.A.*, 119 (1993) 194809q.
- 1345 Ratfisch, W.: (Flame-ionization detector.) *Ger. Offen. DE* 4,131,412 (Cl. G01N21/72), 27 May 1993, Appl. 20 Sep. 1991; 6 pp.; *C.A.*, 119 (1993) 194807n.
- 1346 Ratfisch, W.: (Flame-ionization detection method and apparatus.) *Ger. Offen. DE* 4,120,246 (Cl. G01N27/62), 11 Mar. 1993, Appl. 19 Jun. 1991; 6 pp.; *C.A.*, 119 (1993) 194821n.
- 1347 Shearer, R.L.: Gas chromatography and flameless sulfur chemiluminescence detection. *Int. Lab.*, 24, No. 4 (1994) P4-P8.
- 1348 Sun, X.-Y. and Aue, W.A.: Constancy of spectral response ratios in the flame photometric detector. *J. Chromatogr. A*, 667 (1994) 191-203.
- 1349 Talasek, R.T. and Schoenke, M.P.: Comparison of universal chromatographic detectors for trace analysis. *J. Chromatogr. A*, 667 (1994) 205-211.
- 1350 Tsujino, Y. and Kuwata, K.: Sensitive flame ionization detector for the determination of traces of atmospheric hydrocarbons by capillary column gas chromatography. *J. Chromatogr.*, 642 (1993) 383-388.
- 1351 Vassilaros, D.L.: Contamination of electron-capture detectors with volatile and semivolatile halocarbon impurities in GC gas distribution systems. *LC-GC Int.*, 7 (1994) 139-144.
- 1352 Wan, H.B., Wong, M.K. and Mok, C.Y.: Simple method for preventing unsuitable solvents from entering gas chromatographic detectors. *J. Chromatogr. A*, 663 (1994) 123-126.
- 1353 Yu, A., Wang, X., Yang, W. and Jin, Q.: (Microwave-induced argon plasma photoionization detector for gas chromatography.) *Huaxue Xuebao*, 51 (1993) 1179-1184; *C.A.*, 120 (1994) 182083f.
- See also 1450, 1453, 1527, 1552, 1568, 1707, 1713, 1716.
- 3c. *Sorbents and columns, packing procedures*
- 1354 Abraham, M.H., Du, C.M., Grate, J.W., McHill, R.A. and Shuely, W.J.: Fullerene as an adsorbent for gases and vapors. *J. Chem. Soc., Chem. Commun.*, (1993) 1863-1864; *C.A.*, 120 (1994) 174286j.
- 1355 Akahoshi, T.: (Open tubular wide bore column for GC (G-column).) *Kuromatogurafi*, 14, No. 3-4 (1993) 68-71; *C.A.*, 120 (1994) 94375x.
- 1356 Attaran Rezaii, M. and Lattanzi, L.: Narrow bore silica capillary columns for liquid-modified adsorption gas chromatography. *Chromatographia*, 38 (1994) 235-238.
- 1357 Berezkin, V.G., Levin, Ya.A. and Mukhina, V.P.: Polar immobilized polyhydroxy stationary phase for capillary chromatography. *J. High Resolut. Chromatogr.*, 17 (1994) 280-282.
- 1358 Betts, T.J.: Evaluation of a "Chirasil-Val" capillary for the gas chromatography of volatile oil constituents, including sesquiterpenes in patchouli oil. *J. Chromatogr. A*, 664 (1994) 295-300.
- 1359 Cloux, R., Defayes, G., Foti, K., Dutoit, J.C. and Kovats, E.S.: Pair-wise interactions by gas chromatography. Part III. Synthesis of isosteric stationary phases for gas chromatography. *Synthesis*, (1993) 909-919; *C.A.*, 120 (1994) 163394s.
- 1360 Ejsymont, J.: (Method of obtaining a graphite support for gas chromatography.) *Pol. PL* 161,135 (Cl. B01J20/20), 31 May 1993, Appl. 279,172, 26 Apr. 1989; 3 pp.; *C.A.*, 120 (1994) 94434r.
- 1361 Engewald, W., Reinhardt, R. and Steiborn, A.: The use of shape-selective stationary phases in GC. Part 1: Cyclodextrin phases. *Pure Appl. Chem.*, 65 (1993) 2253-2256; *C.A.*, 120 (1994) 182089n.
- 1362 Guthrie, E.J. and Harland, J.J.: Overview of phase development in capillary gas chromatography. *LC-GC Int.*, 7 (1994) 130-134.

- 1363 Ichikawa, H., Yokoyama, A., Hirata, K., Wada, H. and Kodera, K.: (Packing materials for adsorption gas chromatography and manufacture thereof.) *Jpn. Kokai Tokkyo Koho*, JP 04,166,763 [92,166,763] (Cl. G01N30/48), 12 Jun. 1992, Appl. 90/293,144, 30 Oct. 1990; 5 pp.; C.A., 120 (1994) 182031n.
- 1364 Kartsova, L.A. and Stolyarov, B.V.: (Crown ethers as components of selective mobile and stationary phases in chromatography.) *Zh. Anal. Khim.*, 48 (1993) 582-591.
- 1365 Kim, K.R., Kim, J.H., Park, H.K. and Oh, C.H.: Dual capillary column system for the qualitative gas chromatography. 2. Comparison between splitless and on-column injection modes. *Bull. Korean Chem. Soc.*, 14 (1993) 250-255; C.A., 119 (1993) 240710a.
- 1366 Krestov, A.G., Blokhina, S.V., Sibrina, G.V., Ol'khovich, M.V. and Basaeva, Ya.V.: (Stationary phase for gas chromatography based on microencapsulated mesomorphic *p*-*n*-heptyloxy-*p*'-cyanobiphenyl.) *Izv. Vyssh. Uchebn. Zaved., Khim. Khim. Tekhnol.*, 36, No. 6 (1993) 17-20; C.A., 120 (1994) 182085h.
- 1367 Krupcik, J., Valachovicová, M. and Kraus, G.: Evaluation of shape selectivity of liquid crystals in capillary gas chromatography. *J. Chromatogr. A*, 665 (1994) 111-116.
- 1368 Long, Y., Kang, X. and Huang, T.: (2,6-Di-*O*-butyl-3-*O*-acetyl- β -cyclodextrin as a chiral stationary phase for capillary gas chromatographic separation of enantiomers.) *Fenxi Huaxue*, 21 (1993) 924-926; C.A., 120 (1994) 22793j.
- 1369 Malakhov, V.V., Sidel'nikov, V.N. and Utkin, V.A.: (Use of capillary packet as a chromatography column.) *Dokl. Akad. Nauk*, 329 (1993) 749-51; C.A., 119 (1993) 194763v.
- 1370 Markham, D.A. and Langvardt, P.W.: Coupling capillary gas chromatography to traditional liquid chromatography detectors. *U.S. US* 5,242,471 (Cl. 95-87; B01D15/08), 7 Sep. 1993, Appl. 887,996, 22 May 1992; 12 pp.; C.A., 119 (1993) 240775a.
- 1371 Matsui, M.: (Packing material for gas chromatograph in which air be used as a carrier gas.) *Jpn. Kokai Tokkyo Koho* JP 93,180,819 [05,180,819] (Cl. G01N30/48), 23 Jul. 1993, Appl. 91/360,299, 26 Dec. 1991; 4 pp.; C.A., 120 (1994) 44830w.
- 1372 Myano, S., Ooi, S. and Ochiai, Y.: (Packing material for optical resolution of racemic modification and isomer separation for gas chromatography.) *Jpn. Kokai Tokkyo Koho* JP 05,223,805 [93,223,805] (Cl. G01N30/48), 3 Sep. 1993, Appl. 92/61,397, 14 Feb. 1992; 6 pp.; C.A., 120 (1994) 123876u.
- 1373 Naidanov, O.D. and Ryazantsev, A.A.: (Gas-chromatographic separation of organic compounds using a solid support based on chalcodony.) *Zavod. Lab.*, 59, No. 4 (1993) 18-19.
- 1374 Nishikawa, Y.: (Gas chromatograph and analytical method using it.) *Jpn. Kokai Tokkyo Koho* JP 05,203,635 [93,203,635] (Cl. G01N30/32), 10 Aug. 1993, Appl. 92/12,371, 27 Jan. 1992; 4 pp.; C.A., 119 (1993) 240776b.
- 1375 Saturnin, C., Tabacchi, R. and Saxer, A.: (Gas-chromatographic analysis of racemic mixtures on peralkylated cyclodextrins.) *Chimia*, 47 (1993) 221-226; C.A., 120 (1994) 123832b.
- 1376 Shepelenko, T.S. and Zibarev, P.V.: (High-energy modification of porous polymeric sorbents. 2. Studies of analytical possibilities of radiation-modified sorbents containing nitro groups in chromatography.) *Sib. Khim. Zh.*, No. 2 (1993) 34-37; C.A., 119 (1993) 194751q.
- 1377 Skorniyakov, E.P., Rapoport, L.M. and Fisejskij, Yu.K.: (Chromatographic method for determination of trace impurities in a gas.) *U.S.S.R. SU* 1,734,005 (Cl. G01N30/08), 15 May 1992, Appl. 4,291,206, 29 Jul. 1987; C.A., 119 (1993) 240781z.
- 1378 Vonk, N.: (Gas-chromatographic (GC) analysis of solvents with wide-bore columns.) *Verkroniek*, 66, No. 4 (1993) 16-18; C.A., 119 (1993) 194789h.
- 1379 Wang, H. and Ou, Q.: (Enantiomer separation on a chiral fused silica capillary column with a mixed stationary phase of 2,6-*O*-dipentyl-3-*O*-trifluoroacetyl- β -cyclodextrin and OV-7.) *Fenxi Huaxue*, 21 (1993) 834-837; C.A., 119 (1993) 194764w.
- 1380 Yamane, Y., Miyaji, K., Hanafusa, K., Hanai, T. and Hatano, H.: Structure-retention correlation on methyl phenyl polysiloxane phases in capillary gas chromatography. *Bull. Chem. Soc. Jpn.*, 66 (1993) 1881-1885; C.A., 119 (1993) 261760g.
- 1381 Zhou, X.-C., Wu, C.-Y., Lu, X.-R. and Chen, Y.-Y.: Application of crown ether compounds as gas chromatographic stationary phases. *J. Chromatogr. A*, 662 (1994) 203-218 - a review with 42 refs.
- 1382 Zuo, Y.: (Recent progress of porous polymeric beads—stationary phase for gas chromatography.) *Fenxi Huaxue*, 21 (1993) 971-978; C.A., 119 (1993) 240688z - a review with 63 refs.
- See also 1425, 1426, 1429, 1430, 1438, 1445, 1449, 1669, 1676, 1678.

3d. Quantitative analysis

- 1383 Liang, Y.-z., Hämäläinen, M.D., Kvalheim, O.M. and Andersson, R.: Assessment of peak origin and purity in one-dimensional chromatography by experimental design and heuristic evolving latent projections. *J. Chromatogr. A*, 662 (1994) 113-122.
- 1384 Meyer, V.R.: Chromatographic integration errors: A closer look at a small peak. *LC-GC Int.*, 7 (1994) 94-100.
- 1385 Yi, G., Bradshaw, J.S., Rossiter, B.E., Malik, A., Li, W., Petersson, P., Markides, K.E. and Lee, M.L.: New permethyl-substituted β -cyclodextrin polysiloxanes for use as chiral stationary phases in open tubular column chromatography. *J. Org. Chem.*, 58 (1993) 4844-4850; C.A., 119 (1993) 226279k.

See also 1401, 1515, 1550, 1760.

3e. Preparative scale chromatography

See 1428, 1437, 1446, 1454, 1840.

3f. Programmed temperature, pressure, vapors, gradients

- 1386 Messadi, D. and Ali-Mokhnache, S.: (Comparison of retention indexes computed by various interpolations in programmed temperature gas chromatography.) *C. R. Acad. Sci., Ser. II*, 316 (1993) 605-607; C.A., 119 (1993) 194748u.

4. SPECIAL TECHNIQUES

4a. Automation

- 1387 Scheuer, K.: Knowledge-based interpretation of gas chromatographic data. *Chemom. Intell. Lab. Syst.*, 19 (1993) 201-216.

See also 1312, 1386, 1410.

4b. Computerization and modelling

- 1388 Akiyama, K.: (Software for analysis of hydrocarbon retention index data of automobile exhaust gases.) *Kagaku to Sofutowea*, 15 (1993) 239-242; C.A., 119 (1993) 232697m.
- 1389 Andersson, R. and Haemaelaenen, M.D.: Simplex focusing of retention times and latent variable projections of chromatographic profiles. *Chemom. Intell. Lab. Syst.*, 22 (1994) 49-61.
- 1390 Heitzmann, U.: (Problems with the computerization of analytics.) *GIIT Spez. Chromatogr.*, 13, No. 2 (1993) 82-86; C.A., 120 (1994) 37488n.
- 1391 Park, M.K., Cho, J.H., Kim, N.Y. and Park, J.H.: Chromatographic pattern recognition for the analysis of complex mixtures. *Anal. Chim. Acta*, 284 (1993) 73-78.
- 1392 Snyder, W.D. and Blumberg, L.M.: Calculating operating parameters for a gas chromatograph. *Eur. Pat. Appl.* EP 570,707 (Cl. G01N30/88), 24 Nov. 1993, US Appl. 883,995, 18 May 1992; 15 pp.; C.A., 120 (1994) 182012g.

See also 1308.

4c. Combination with other physico-chemical techniques (MS, IR etc.)

- 1393 Brittain, R. and Feigel, C.: Chemical ionization and MS-MS technologies for benchtop ion trap GC-MS systems. *Am. Lab. (Shelton)*, 26, No. 3 (1994) 44-48.
- 1394 Clark, L.B.: Application of on-column gas chromatography-mass spectrometry and particle beam/liquid chromatography/mass spectrometry for water analysis. Avail. *Univ. Microfilms Int.*, Order No. DA 9219919, 1992, 137 pp.; C.A., 120 (1994) 143528k.
- 1395 Dong, Q.: (The hyphenated techniques of chromatography and FTIR spectroscopy.) *Guangpuxue Yu Guangpu Fenxi*, 13 (1993) 25-32; C.A., 120 (1994) 152616q - a review with 33 refs.
- 1396 Enqvist, J. and Enqvist, M.: Comparison of two-stage retention index monitoring capillary gas chromatography and selected ion monitoring capillary gas chromatography-mass spectrometry as analytical tools. *J. High Resolut. Chromatogr.*, 17 (1994) 141-144.
- 1397 Games, D.E., Rontree, J.A. and Fowles, I.A.: The direct combination of HPLC solute focusing with supercritical fluid chromatography and mass spectrometry to enhance sensitivity and improve identification of trace components in polar solvent extracts. *J. High Resolut. Chromatogr.*, 17 (1994) 68-71.
- 1398 Haaland, D.M., Thomas, E.V. and Blair, D.S.: Improvements in methods for spectral combination of gas chromatography/Fourier transform infrared spectroscopic data. *Appl. Spectrosc.*, 47 (1993) 1612-1619.
- 1399 Janák, K., Östman, C., Carlsson, H., Bengård, A. and Colmsjö, A.: Instrument-induced effects in the analysis of polycyclic aromatic compounds by capillary gas chromatography with atomic emission detection (GC-AED). *J. High Resolut. Chromatogr.*, 17 (1994) 135-140.
- 1400 Jenkins, T.J., Kaplan, M. and Davidson, G.: Capillary SFC-FTIR - effects of the flow-cell on chromatographic resolution. *J. High Resolut. Chromatogr.*, 17 (1994) 160-164.

- 1401 Johansson, M., Berglund, M. and Baxter, D.C.: Improving accuracy in the quantitation of overlapping, asymmetric, chromatographic peaks by deconvolution: theory and application to coupled gas chromatography atomic absorption spectrometry. *Spectrochim. Acta B*, 48B (1993) 1393-1409; C.A., 119 (1993) 240699d.
- 1402 Kirschner, C.H. and Taylor, L.T.: Carbon dioxide versus xenon as a mobile phase for flow cell packed column SFC/FT-IR. *J. High Resolut. Chromatogr.*, 17 (1994) 61-67.
- 1403 Krock, K.A., Ragunathan, N., Klawun, C., Sasaki, T. and Wilkins, C.L.: Multidimensional gas chromatography-infrared spectrometry-mass spectrometry. *Analyst (Cambridge)*, 119 (1994) 483-489.
- 1404 Kusmierz, J.J. and Abramson, F.P.: Improved measurement of stable isotope ratios in gas chromatography/mass spectrometry using the microwave-powered chemical reaction interface for mass spectrometry. *Biol. Mass Spectrom.*, 22 (1993) 537-543; C.A., 120 (1994) 26923z.
- 1405 Laramée, J.A. and Deinzer, M.L.: Capillary gas chromatographic introduction of environmental compounds into a toroidal electron monochromator/mass spectrometer. *Anal. Chem.*, 66 (1994) 719-724.
- 1406 Leal, W.S., Kuwahara, Y., Matsuyama, S., Suzuki, T. and Ozawa, T.: GC-FTIR potential for structure elucidation. *J. Braz. Chem. Soc.*, 3 (1992) 25-29; C.A., 119 (1993) 181029k.
- 1407 Listemann, M.L., Waller, F.J. and Herman, F.L.: Survey analyses using gas chromatography coupled with infrared and mass spectrometric detection. *Spectroscopy (Eugene)*, 8, No. 5 (1993) 40-45; C.A., 120 (1994) 94407j.
- 1408 Maswadeh, W., Arnold, N.S., McClennen, W.H., Tripathi, A., DuBow, J. and Meuzelaar, H.L.C.: Development of a laser devolatilization gas chromatography/mass spectrometry technique for single coal particles. *Energy Fuels*, 7 (1993) 1006-1012; C.A., 119 (1993) 229751u.
- 1409 Musil, S. and Leško, J.: Isotopic ratio of molecular patterns via gas chromatography-mass spectrometry with selected-ion monitoring as a chemometric tool. *J. Chromatogr. A*, 664 (1994) 253-261.
- 1410 Nakagawa, K.: (Data processor for GC/MS.) *Jpn. Kokai Tokkyo Koho JP 05,164,751 [93,164,751]* (Cl. G01N30/72), 29 Jun. 1993, Appl. 91/336,753, 19 Dec. 1991; 4 pp.; C.A., 119 (1993) 261823e.
- 1411 Schoene, K., Bruckert, H.-J., Steinhanses, J. and König, A.: Two stage derivatization with N-(tert.-butyldimethylsilyl)-N-methyl-trifluoroacetamide (MTBSTFA) and N-methyl-bis-(trifluoroacetamide) (MBTFA) for the gas-chromatographic analysis of OH-, SH- and NH-compounds. *Fresenius J. Anal. Chem.*, 348 (1994) 364-370.
- 1412 Sturaro, A., Parvoli, G. and Doretti, L.: Standards and GC-MS analysis: an answer to the requirement of compound confirmation. *Chromatographia*, 38 (1994) 239-241.
- 1413 Wilkins, C.L.: Multidimensional GC for qualitative IR and MS of mixtures. *Anal. Chem.*, 66 (1994) 295A-301A.

See also 1327, 1656, 1698, 1728, 1799.

4e. Functional analysis

- 1414 Amjeeb, M. and Wells, R.J.: Methylation reagents for the direct on-column derivatisation of veterinary residues. *J. Chromatogr. A*, 662 (1994) 123-137.

4f. Trace analysis and pre-separation techniques

- 1415 Bruno, T.J.: Chromatographic cryofocusing and cryotrapping with the vortex tube. *J. Chromatogr. Sci.*, 32 (1994) 112-115.
- 1416 Contarini, G. and Leardi, R.: Use of experimental design to optimize the analysis of volatile compounds by dynamic head-space extraction followed by cold trapping and capillary GC. *J. High Resolut. Chromatogr.*, 17 (1994) 91-94.
- 1417 Hartman, T.G., Lech, J., Karmas, K., Salinas, J., Rosen, R.T. and Ho, C.T.: Flavor characterization using adsorbent trapping—thermal desorption or direct thermal desorption—gas chromatography and gas chromatography—mass spectrometry. *IFT Basic Symp. Ser.*, 8 (1993) 37-60; *C.A.*, 119 (1993) 179503s.
- 1418 Hiatt, M.H., Youngman, D.R. and Donnelly, J.R.: Separation and isolation of volatile organic compounds using vacuum distillation with GC/MS determination. *Anal. Chem.*, 66 (1994) 905-908.
- 1419 Mottagh, S. and Pawliszyn, J.: Online monitoring of flowing samples using solid phase microextraction—gas chromatography. *Anal. Chim. Acta*, 284 (1993) 265-273.
- 1420 Nerin, C., Pons, B., Martínez, M. and Cacho, J.: Behaviour of several solid adsorbents for sampling tetraalkyllead compounds in air. *Mikrochim. Acta*, 112 (1994) 179-188.
- 1421 Picó, Y., Vreuls, J.J., Ghijsen, R.T. and Brinkman, U.A.T.: Drying agents for water-free introduction of desorption solvent into a GC after on-line SPE of aqueous samples. *Chromatographia*, 38 (1994) 461-469.
- 1422 Škrabáková, S., Matisová, E., Benická, E., Novák, I. and Berek, D.: Use of a novel carbon sorbent for the adsorption of organic compounds from water. *J. Chromatogr. A*, 665 (1994) 27-37.

See also 1319, 1377, 1451, 1492, 1729, 1734, 1744, 1874.

4g. Enantiomers, separation

- 1423 Anton, K., Eppinger, J., Frederiksen, L., Francotte, E., Berger, T.A. and Wilson, W.H.: Chiral separations by packed-column super- and subcritical fluid chromatography. *J. Chromatogr. A*, 666 (1994) 395-401.
- 1424 Bargmann-Leyder, N., Truffert, J.-C., Tambuté, A. and Caude, M.: Evaluation of Pirkle-type chiral stationary phases by liquid and supercritical fluid chromatography. Influence of the spacer length and the steric hindrance in the vicinity of the stereogenic centre. *J. Chromatogr. A*, 666 (1994) 27-40.
- 1425 Bicchi, C., D'Amato, A., Manzin, V., Galli, A. and Galli, M.: Cyclodextrin derivatives in the gas chromatographic separation of racemic mixtures of volatile compounds. VII. The use of 2,6-di-O-methyl-3-O-pentyl- β -cyclodextrin diluted in phases with different polarity in the separation of racemates in complex mixtures. *J. Chromatogr. A*, 666 (1994) 137-146.

- 1426 Dietrich, A., Maas, B. and Mosandl, A.: Diluted modified cyclodextrins as chiral capillary gas chromatographic stationary phases: Influence of the polysiloxane solvents on enantioselectivity and column efficiency. *J. Microcolumn Sep.*, 6 (1994) 33-42.
- 1427 Engel, K.H.: Stereodifferentiation of chiral flavor and aroma compounds. In: Wittkowski, R. and Matissek, R. (Editors), *Capillary Gas Chromatogr. Food Control Res.*, Technomic, Lancaster, 1993, pp. 149-175; *C.A.*, 119 (1993) 179409r - a review with 110 refs.
- 1428 Hardt, I. and König, W.A.: Preparative enantiomer separation with modified cyclodextrins as chiral stationary phases. *J. Chromatogr. A*, 666 (1994) 611-615.
- 1429 Jaques, K., Buda, W., Dumortier, L., van der Eycken, J., Venema, A. and Sandra, P.: Enantioselective separations by capillary GC on derivatized cyclodextrins. Part 2: Determination of enantiomeric excess of some racemic 2,3-isopropylidene-1,2,3-cyclohexanetriol derivatives on permethyl α -, β -, and γ -cyclodextrins. *J. High Resolut. Chromatogr.*, 17 (1994) 72-76.
- 1430 Jaques, K., Buda, W.M., Venema, A. and Sandra, P.: Enantioselective separations by capillary gas chromatography on derivatized cyclodextrins. III. Separation of some racemic 2,2-dialkyl-4-alkoxycarbonyl-1,3-dioxolane derivatives on 2,3-di-O-acetyl-6-O-*tert*-butyldimethylsilyl- β - and γ -cyclodextrins. *J. Chromatogr. A*, 666 (1994) 131-136.
- 1431 Koppenhoefer, B., Graf, R., Holzschuh, H., Nothdurft, A., Trettin, U., Piras, P. and Roussel, C.: CHIRBASE, a molecular database for the separation of enantiomers by chromatography. *J. Chromatogr. A*, 666 (1994) 557-563.
- 1432 Koppenhoefer, B., Nothdurft, A., Pierot-Sanders, J., Piras, P., Popescu, C., Roussel, C., Stiebler, M. and Trettin, U.: CHIRBASE, a graphical molecular database on the separation of enantiomers by liquid-, supercritical fluid- and gas chromatography. *Chirality*, 5 (1994) 213-219.
- 1433 Maas, B., Dietrich, A. and Mosandl, A.: Collection of enantiomer separation factors obtained by capillary gas chromatography on chiral stationary phases. *J. High Resolut. Chromatogr.*, 17 (1994) 169-173.
- 1434 Maas, B., Dietrich, A. and Mosandl, A.: Collection of enantiomers separation factors obtained by capillary gas chromatography on chiral stationary phases. *J. High Resolut. Chromatogr.*, 17 (1994) 109-115.
- 1435 Petersson, P. and Markides, K.E.: Chiral separations performed by supercritical fluid chromatography. *J. Chromatogr. A*, 666 (1994) 381-384 - a review with 79 refs.
- 1436 Schurig, V.: Enantiomer separation by gas chromatography on chiral stationary phases. *J. Chromatogr. A*, 666 (1994) 111-119 - a review with 115 refs.
- 1437 Schurig, V. and Grosenick, H.: Preparative enantiomer separation of enflurane and isoflurane by inclusion gas chromatography. *J. Chromatogr. A*, 666 (1994) 617-625.
- 1438 Tang, Y., Zhou, Y. and Armstrong, D.W.: Examination of the enantioselectivity of wall-immobilized cyclodextrin copolymers in capillary gas chromatography. *J. Chromatogr. A*, 666 (1994) 147-159.
- 1439 Werkhoff, P., Brennecke, S., Bretschneider, W., Guentert, M., Hopp, R. and Surburg, H.: Chiral specific analysis in essential oil, fragrance, and flavor research. *Z. Lebensm.-Unters. Forsch.*, 196 (1993) 307-328; *C.A.*, 119 (1993) 179461b - a review with 161 refs.

See also 1358, 1368, 1372, 1375, 1379, 1385, 1578, 1645, 1646, 1648, 1649, 1676, 1678, 1683, 1704, 1749, 1812, 1825, 1840, 1845, 1897.

4h. Other special techniques

- 1440 Itani, M. and Giese, R.W.: Extractive hydrogenation: a new and versatile technique broadening gas chromatography applications. *Anal. Chem.*, 66 (1994) 1189-1197.
- 1441 Kumar, N. and Gow, J.G.: Residual solvent analysis by headspace gas chromatography. *J. Chromatogr. A*, 667 (1994) 235-240.
- 1442 Morales, M.T. and Aparicio, R.: Optimization by mathematical procedures of two dynamic headspace techniques for quantifying virgin olive oil volatiles. *Anal. Chim. Acta*, 282 (1993) 423-431.
- 1443 Nicoud, R.M., Perrut, M. and Hotier, G.: Method and device for fractionating a mixture in a simulated fluidized bed in the presence of a compressed gas, a supercritical fluid or a subcritical fluid. *PCT Int. Appl.* WO 93 22,022 (Cl. B01D15/02), 11 Nov. 1993, FR Appl. 92/5,304, 29 Apr. 1992; 55 pp.; C.A., 120 (1994) 167925p.

See also 1474, 1511, 1585.

4i. Supercritical fluid chromatography

- 1444 Artz, W.E.: Supercritical fluid chromatography: principles and applications. In: Baianu, I.C., Pessen, H. and Kumosinski, T.F. (Editors), *Phys. Chem. Food Processes*, Van Nostrand Reinhold, New York, 1993, pp. 518-541; C.A., 119 (1993) 179404k - a review with 128 refs.
- 1445 Blomberg, L.G., Demirbükler, M., Häggglund, I. and Andersson, P.E.: Supercritical fluid chromatography: open tubular vs. packed columns. *TrAC*, 13 (1994) 126-137 - a review with 129 refs.
- 1446 Cretier, G., Majdalani, R., Neffati, J. and Rocca, J.L.: Improvement in the solvent evaporation injection technique for preparative supercritical fluid chromatography. *Chromatographia*, 38 (1994) 330-336.
- 1447 Grover, V., van Puyvelde, F. and Chimowitz, E.H.: A variational formulation and numerical solution of the optimal resolution problem in supercritical fluid chromatography. *Comput. Chem. Eng.*, 18 (1994) 153-162; C.A., 120 (1994) 33585u.
- 1448 Hagege, A., Rocca, J.L. and Djerki, R.: Thermodynamic study of alkanes, alkylbenzenes and chloroalkanes retention on alkyl- and cyanoalkyl-bonded stationary phases in supercritical fluid chromatography. *Chromatographia*, 38 (1994) 373-380.
- 1449 Ibáñez, E., Tabera, J., Herraiz, M. and Reglero, G.: Effect of temperature and density on the performance of micropacked columns in supercritical fluid chromatography. *J. Chromatogr. A*, 667 (1994) 249-255.
- 1450 Lembke, P., Engelhardt, H. and Ecker, R.: On-line coupling of packed column supercritical fluid chromatography with multiple wavelength detector and radioactivity flow-through detector. *Chromatographia*, 38 (1994) 491-501.
- 1451 Li, K., Ong, C.P. and Li, S.F.Y.: Systematic multivariate optimization of supercritical fluid extraction. *J. Chromatogr. Sci.*, 32 (1994) 53-56.

- 1452 Page, S.H., Sumpter, S.R., Goates, S.R., Lee, M.L., Dixon, D.J. and Johnston, K.P.: Tri-*n*-butylphosphate/carbon dioxide and acetone/CO₂ phase behaviors and utilities in capillary supercritical-fluid chromatography. *J. Supercrit. Fluids*, 6 (1993) 95-101.
- 1453 Sandmann, B.W. and Grayeski, M.L.: Peroxyoxalate chemiluminescence detection with packed column supercritical fluid chromatography. *Chromatographia*, 38 (1994) 163-167.
- 1454 Sargentí, S.R. and Lanças, F.M.: Design and construction of a simple supercritical fluid extraction system with semi-preparative and preparative capabilities for application to natural products. *J. Chromatogr. A*, 667 (1994) 213-218.
- 1455 Shen, Y., Huang, H., Liu, X., Wang, Q. and Zhou, L.: Application of capillary supercritical fluid chromatography to the characterization of edible oils and traditional Chinese medicines. *J. High Resolut. Chromatogr.*, 17 (1994) 74-76.
- 1456 Smith, R.M.: Nomenclature for supercritical fluid chromatography and extraction. *Pure Appl. Chem.*, 65 (1993) 2397-2403; C.A., 120 (1994) 145118a.
- 1457 Sun, C., Cong, S., Xu, X. et al.: (Capillary supercritical-fluid chromatographic method.) *Faming Zhuanli Shenqing Gongkai Shuomingshu*, CN 1,065,529 (Cl. G01N30/16), 21 Oct. 1992, Appl. 91,106,366, 1 Apr. 1991; 12 pp.; C.A., 119 (1993) 194703a.
- 1458 Vejrosta, J., Mikeová, M., Ansořgová, A., Zátorsky, J., Hajer, M. and Janda, V.: Multichannel restrictor for supercritical fluid chromatography. *J. Microcolumn Sep.*, 6 (1994) 23-25.
- 1459 Zhou, L., Shen, Y., Huang, H. and Zhu, D.: (Supercritical fluid chromatographic system and its application.) *Fenxi Huaxue*, 21 (1993) 983-987; C.A., 119 (1993) 285504c.

See also 1267, 1311, 1332, 1385, 1397, 1400, 1402, 1423, 1424, 1435, 1443, 1472, 1475, 1480, 1482, 1483, 1492, 1510, 1511, 1547, 1617, 1635, 1705, 1713, 1728, 1732, 1744, 1752, 1764, 1777, 1799, 1803, 1811, 1830, 1831, 1928, 1929.

5. HYDROCARBONS AND HALOGEN DERIVATIVES

5a. Aliphatic hydrocarbons

- 1460 Devyatykh, G.G., Krylov, V.A., Kovalev, I.D., Osipova, L.I. and Yan'kov, S.V.: (Impurity composition of specimens of high-purity substances in a collection exhibition. 8. Permanent gases.) *Vysokochist. Veshchestva*, No. 4 (1993) 7-11; C.A., 119 (1993) 240603t.
- 1461 Farmer, C.T., Milne, P.J., Riemer, D.D. and Zika, R.G.: Continuous hourly analysis of C₂-C₁₀ non-methane hydrocarbon compounds in urban air by GC-FID. *Environ. Sci. Technol.*, 28 (1994) 238-245.
- 1462 Kenig, F., Sinninghe Damsté, J.S., de Leeuw, J.W. and Hayes, J.M.: Molecular palaeontological evidence for food-web relationships. *Naturwissenschaften*, 81 (1994) 128-130.
- 1463 Lo, J.-G., Chen, T.-Y. and Tso, T.-L.: Improved resolution and determination of ambient C₅-C₁₂ hydrocarbons by GC-coupled refocussing system. *Chromatographia*, 38 (1994) 151-157.

- 1464 Mariani, C., Venturinin, S. and Fedeli, E.: (Determination of hydrocarbons and free and esterified minor components in olive oil of different classes.) *Riv. Ital. Sostanze Grasse*, 70 (1993) 321-327; *C.A.*, 120 (1994) 75743z.
- 1465 Martinez, J.E.: Impact of natural and anthropogenic hydrocarbons on tropospheric ozone production: results from automated gas chromatography. Avail. *Univ. Microfilms Int.*, Order No. DA9223785, 1992, 171 pp.; *C.A.*, 120 (1994) 142633d.
- 1466 Massias, L., Postaire, E., Regnault, C. and Hazebroucq, G.: Thermal desorption-gas chromatographic determination of ethane and pentane in breath as potential markers of lipid peroxidation. *Biomed. Chromatogr.*, 7 (1993) 200-203.
- 1467 Orav, A. and Rang, S.: Capillary gas chromatography of C₆-C₁₀ isoalkenes on polyethylene glycol 20M. *Eesti Tead. Akad. Toim., Keem.*, 42 (1993) 77-82; *C.A.*, 120 (1994) 22803n.
- 1468 Ramnäs, O., Östermark, U. and Petersson, G.: Characterization of sixty alkenes in a cat-cracked gasoline naphtha by gas chromatography. *Chromatographia*, 38 (1994) 222-226.
- 1469 Rodinkov, O.V. and Gumerov, M.F.: (Determination of gaseous unsaturated hydrocarbons in water.) *U.S.S.R. SU 1,770,894* (Cl. G01N30/06), 23 Oct. 1992, Appl. 4,720,536, 19 Jul. 1989; *C.A.*, 119 (1993) 233612s.
- See also 1274, 1276, 1287, 1448, 1940.
- 5b. *Cyclic hydrocarbons, fullerenes*
- 1470 Kawaki, H.: On the solute-stationary phase interaction in gas liquid chromatography. Relative retention values for mono-substituted benzene derivatives and their energy partition by means of regression analysis. *Chem. Pharm. Bull.*, 41 (1993) 797-803; *C.A.*, 119 (1993) 249530z.
- 1471 Kiss, G., Compaan, H. and Hoekstra, E.J.: (Use of deuterated PAHs [polycyclic aromatic hydrocarbons] for the determination of PAHs in environmental samples.) *Magy. Kem. Foly.*, 99 (1993) 200-204; *C.A.*, 119 (1993) 179918f.
- 1472 Kot, A., Sandra, P. and David, F.: Selectivity tuning in packed column SFC. Separation of the sixteen priority polycyclic aromatic hydrocarbons as an example. *J. High Resolut. Chromatogr.*, 17 (1994) 277-279.
- 1473 Kurán, P., Kubinec, R., Ostrovsky, I. and Soják, L.: Purge-and-trap injection capillary gas chromatographic determination of volatile aromatic hydrocarbons in river sediment. *J. Chromatogr. A*, 665 (1994) 133-138.
- 1474 Le Sech, J., Ducruet, V. and Feigenbaum, A.: Influence of dissolved gases in the dynamic headspace analysis of styrene and other volatile organic compounds and improvement of their determination. *J. Chromatogr. A*, 667 (1994) 340-347.
- 1475 Liu, M., Thienpont, A., Delville, M.H. and Félix, G.: Rapid separations of fullerenes by donor-acceptor liquid and supercritical fluid chromatography. *J. High Resolut. Chromatogr.*, 17 (1994) 104-106.
- 1476 Luks-Betlej, K.: Application of gas chromatography-mass spectrometry in selected ion monitoring mode for benzo[a]pyrene quantitation in particulate material on ppb level. *Chem. Anal. (Warsaw)*, 38 (1993) 585-588.
- 1477 Nabivach, V.M. and Gerasimenko, V.A.: (Correlation-additive procedure for identification of alkylaromatic compounds.) *Koks Khim.*, No. 5 (1993) 28-29; *C.A.*, 119 (1993) 229804p.
- 1478 Nyman, P.J., Perfetti, G.A., Joe, F.L., Jr. and Diachenko, G.W.: Comparison of two clean-up methodologies for the gas chromatographic/mass spectrometric determination of low nanogram/gram levels of polynuclear aromatic hydrocarbons in seafood. *Food Addit. Contam.*, 10 (1993) 489-501; *C.A.*, 120 (1994) 132525f.
- 1479 Potter, D.W. and Pawliszyn, J.: Rapid determination of polycyclic aromatic hydrocarbons and polychlorinated biphenyls in water using solid-phase microextraction and GC/MS. *Environ. Sci. Technol.*, 28 (1994) 298-305.
- 1480 Stalling, D.L., Saim, S., Guo, C. and Kuo, K.: Separation and recovery of C₆₀- and C₇₀-fullerenes from carbon soot by supercritical fluid extraction and adsorption chromatography. *U.S. US 5,281,406* (Cl. 423-445B; C01B31/00), 25 Jan. 1994, Appl. 874,473, 22 Apr. 1992; 29 pp.; *C.A.*, 120 (1994) 168100c.
- 1481 Sturaro, A., Parvoli, G., Doretti, L., Gori, G. and Bartolucci, G.B.: GC-MS investigation of polycyclic aromatic compounds in the manufacture of rubber tubes. *Ann. Occup. Hyg.*, 37 (1993) 327-334; *C.A.*, 120 (1994) 85299y.
- 1482 Tena, M.T., Lague de Castro, M.D. and Valcárcel, M.: Systematic study of the influence on the CO₂-supercritical extraction of PAHs in soil. *Chromatographia*, 38 (1994) 431-435.
- 1483 Thomas, D., Sim, P.G. and Benoit, F.M.: Capillary column supercritical fluid chromatography/mass spectrometry of polycyclic aromatic compounds using atmospheric pressure chemical ionization. *Rapid Commun. Mass Spectrom.*, 8 (1994) 105-110; *C.A.*, 120 (1994) 182126x.
- 1484 Van der Velde, E.G., Marsman, J.A., de Jong, A.P.J.M., Hoogerbrugge, R. and Liem, A.K.D.: Analysis and occurrence of toxic planar PCBs, PCDDs and PCDFs in milk by use of carbosphere activated carbon. *Chemosphere*, 28 (1994) 693-702.
- 1485 Venkataraman, C., Lyons, J.M. and Friedlander, K.: Size distribution of polycyclic aromatic hydrocarbons and elemental carbon. 1. Sampling, measurement methods, and source characterization. *Environ. Sci. Technol.*, 28 (1994) 555-562.
- 1486 Wang, P., Hou, D. and Lin, R.: (Aromatic hydrocarbon composition and thermal evolution characteristics of extracts of lignite from Huangxian.) *Diqiu Huaxue*, (1993) 313-325; *C.A.*, 120 (1994) 168215u.
- 1487 Zinov'eva, N.P., Kartashova, A.V. and Kosogaeva, T.M.: (Gas chromatographic determination of aromatic compounds in water by equilibration with steam.) *Gig. Sanit.*, No. 7 (1993) 76-77; *C.A.*, 120 (1994) 116306e.
- See also 1292, 1399, 1418, 1448, 1502, 1814.
- 5c. *Halogen derivatives*
- 1488 Alawi, M.A. and Buthe, A.: Identification of PCB-individual congeners in Jordanian mother milk by GC-MS. *Fresenius Environ. Bull.*, 2 (1993) 206-213; *C.A.*, 120 (1994) 71066h.
- 1489 Boehm, V., Schulte, E. and Thier, H.P.: Polychlorinated biphenyl residues in food and human milk. Determination of co-planar and mono-ortho substituted congeners. *Z. Lebensm.-Unters. Forsch.*, 196 (1993) 435-440; *C.A.*, 119 (1993) 179523y.
- 1490 Böhm, V., Schulte, E. and Thier, H.P.: Carbon column HPLC and carborane-phase capillary GLC for the separation of polychlorinated biphenyl congeners. *Fresenius J. Anal. Chem.*, 348 (1994) 297-300.

- 1491 Bøwadt, S. and Johansson, B.: Analysis of PCBs in sulfur-containing sediments by off-line supercritical fluid extraction and HRGC-ECD. *Anal. Chem.*, 66 (1994) 667-673.
- 1492 Bøwadt, S., Johansson, B., Pelusio, F., Larsen, B.R., Rovida, C.: Solid-phase trapping of polychlorinated biphenyls in supercritical fluid extraction. *J. Chromatogr. A*, 662 (1994) 424-433.
- 1493 Kanno, S.: Determination method for methyl iodide using diffusive sampler and capillary GC with thermal desorption injector and FID. *Ind. Health*, 31 (1993) 39-50; *C.A.*, 120 (1994) 142547d.
- 1494 Langenfeld, J.J., Hawthorne, S.B., Miller, D.J. and Pawliszyn, J.: Role of modifiers for analytical-scale supercritical fluid extraction of environmental samples. *Anal. Chem.*, 66 (1994) 909-916.
- 1495 Lee, H.-B. and Peart, T.E.: Optimization of supercritical carbon dioxide extraction for polychlorinated biphenyls and chlorinated benzenes from sediments. *J. Chromatogr. A*, 663 (1994) 87-95.
- 1496 Li, L., Wu, J., Chen, Y. and Jiang, K.: (Determination of polychlorinated biphenyls in transformer oils.) *Huanjing Kexue*, 14 (1993) 69-72; *C.A.*, 119 (1993) 274709x.
- 1497 Mes, J., Arnold, D.L. and Bryce, F.: Determination of polychlorinated biphenyls in postpartum blood, adipose tissue, and milk from female rhesus monkeys and their offspring after prolonged dosing with Aroclor 1254. *J. Anal. Toxicol.*, 18 (1994) 29-35; *C.A.*, 120 (1994) 127059c.
- 1498 Morosini, M. and Ballschmiter, K.: Retention indices of 28 polychlorinated biphenyls in capillary gas chromatography referred to 2,4,6-trichlorophenyl alkyl ethers as RI-standards. *Fresenius J. Anal. Chem.*, 348 (1994) 595-597.
- 1499 Ozawa, H., Murooka, M. and Matsubara, H.: (Determination of low-boiling-point halogenated hydrocarbons in water samples by purge trap and capillary gas chromatography/mass spectrometry.) *Ehime-ken Kogai Gijyutsu Senta Shoho*, 13 (1993) 20-23; *C.A.*, 119 (1993) 233515n.
- 1500 Schall, C. and Heumann, K.G.: GC determination of volatile organoiodine and organobromine compounds in Arctic seawater and air samples. *Fresenius J. Anal. Chem.*, 346 (1993) 717-722.
- 1501 Schreitmüller, J. and Ballschmiter, K.: Levels of polychlorinated biphenyls in the lower troposphere of the South-Atlantic ocean. Studies of global baseline pollution. *Fresenius J. Anal. Chem.*, 348 (1994) 226-239.
- 1502 Wilson, S.C., Burnett, V., Waterhouse, K.S. and Jones, K.C.: Volatile organic compounds in digested United Kingdom sewage sludges. *Environ. Sci. Technol.*, 28 (1994) 259-266.
- 1503 Zhang, E., Ying, Y. and Gong, Y.: (Determination of total trace organic chlorine content in naphtha by gas chromatography.) *Shiyou Lianzhi*, No. 12 (1992) 47-50; *C.A.*, 120 (1994) 11409n.
- 1504 Zhang, S.: (Determination of TCB residues in soil and water by gas chromatography.) *Turang (Nanjing)*, 25 (1993) 112-113; *C.A.*, 120 (1994) 76319c.
- See also 1292, 1356, 1415, 1418, 1448, 1479, 1751, 1814, 1874.
- 5d. *Complex hydrocarbon mixtures (incl. analysis of tars, bitumens and mineral oils)*
- 1505 Baum, M.G.: (Applied geochemical surface investigations as supplemental exploration tool – case histories.) *Erdoel, Erdgas, Kohle*, 110 (1994) 9-15; *C.A.*, 120 (1994) 168218x.
- 1506 Ciupe, R., Spangenberg, J., Woelke, J., Wild, G. and Meyer, T.: (Simultaneous determination of aromatic and halogenated hydrocarbons by gas chromatography with FID and electron-capture detector.) *GIT Spez. Chromatogr.*, 13, No. 2 (1993) 60-62; *C.A.*, 120 (1994) 37644k.
- 1507 De Aquino Neto, F.R., Cardoso, J.N., dos Santos Pereira, A., Fernandes, M.C.Z., Caetano, C.A. and de Castro Machado, A.L.: Application of high temperature high resolution gas chromatography to paraffinic deposits in petroleum production pipelines. *J. High Resolut. Chromatogr.*, 17 (1994) 259-263.
- 1508 Kaiser, E.W. and Siegl, W.O.: High resolution gas chromatographic determination of the atmospheric reactivity of engine-out hydrocarbon emissions from a spark-ignited engine. *J. High Resolut. Chromatogr.*, 17 (1994) 264-270.
- 1509 Lasak, K.: (Analytical monitoring of BTX hydrocarbon removal from coke-oven gas by gas chromatography.) *Hutn. Listy*, 48 (1993) 41-44; *C.A.*, 120 (1994) 111001s - a review with 22 refs.
- 1510 Levy, J.M.: Fossil fuel applications of SFC and SFE: a review. *J. High Resolut. Chromatogr.*, 17 (1994) 212-216 - a review with 52 refs.
- 1511 Peaden, P.A.: Simulated distillation of petroleum and its products by gas and supercritical fluid chromatography. *J. High Resolut. Chromatogr.*, 17 (1994) 203-211 - a review with 57 refs.
- 1512 Šepić, E. and Leskovšek, H.: Rapid gas-chromatographic determination of hydrocarbon types by selection absorption and flame ionization detection. *Fresenius J. Anal. Chem.*, 348 (1994) 650-653.
- 1513 Steliga, T.: (Use of gas chromatography for the control of effectiveness or removal of oil derivatives from mine wastewater.) *Nafta-Gas*, 49 (1993) 338-343; *C.A.*, 120 (1994) 143513b.
- See also 1262, 1388, 1422, 1924, 1932.
6. ALCOHOLS
- 1514 Andersson, R.R. and White, C.M.: Analysis of Fischer-Tropsch by-products waters by gas chromatography. *J. High Resolut. Chromatogr.*, 17 (1994) 245-250.
- 1515 Buckee, G.K. and Mundy, A.P.: Determination of ethanol in beer by gas chromatography (direct injection) - collaborative trial. *J. Inst. Brew.*, 99 (1993) 381-384; *C.A.*, 120 (1994) 7002g.
- 1516 Edinboro, L.E., Nanco, C.R., Soghioan, D.M. and Poklis, A.: Determination of ethylene glycol in serum utilizing direct injection on a wide-bore capillary column. *Ther. Drug Monit.*, 15 (1993) 220-223; *C.A.*, 119 (1993) 175479r.
- 1517 Gul'yayeva, T.N.: (Ethylene glycol detection in biologic objects in the presence of propylene glycol.) *Sud.-Med. Ekspert.*, 35, No. 4 (1992) 33-34; *C.A.*, 119 (1993) 197371b.
- 1518 Krupčík, J., Špáník, I. and Sandra, P.: Separation of C₁₁-C₁₄ branched-chain alcohols by high-resolution gas chromatography on a modified β -cyclodextrin stationary phase. *J. Chromatogr. A*, 665 (1994) 163-168.
- 1519 Krupčík, J., Benická, E., Májek, P., Skacáni, I. and Sandra, P.: Relationship between structure and chromatographic behaviour of secondary alcohols and their derivatives separated by high-resolution gas chromatography with a modified β -cyclodextrin stationary phase. *J. Chromatogr. A*, 665 (1994) 175-184.

- 1520 Ledahudcova, K., Kvasnicka, F. and Velisek, J.: (Determination of 3-amino-1,2-propanediol in protein hydrolyzates.) *Potravin. Vedy*, 11 (1993) 121-133; C.A., 120 (1994) 7018s.
- 1521 Linkiewicz, M. and Poskrobko, H.: Determination of impurities in 2-methoxyethanol and 2-butoxyethanol by gas chromatography. *Chem. Anal. (Warsaw)*, 38 (1993) 429-432.
- 1522 Linkiewicz, M., Jaworski, M. and Poskrobko, J.: Analysis of alcohol ethoxylation products with gas chromatography. *Chem. Anal. (Warsaw)*, 38 (1993) 149-155.
- 1523 Morgan, S.E., Frink, E.J. and Gandolfi, A.J.: A simplified gas chromatographic method for quantifying the sevoflurane metabolite hexafluoroisopropanol. *Anesthesiology*, 80 (1994) 201-205; C.A., 120 (1994) 123979e.
- 1524 Muszynski, J., Jankowski, W., Dowsin, A., Terlikiewicz, I., Pojasek, A. and Kuchar, W.: (Gas chromatography method for analysis of alkylphenols.) *Pol. PL 161,082* (Cl. G01N30/02), 31 May 1993, Appl. 282,795, 13 Dec. 1989; 6 pp.; C.A., 120 (1994) 94439w.
- 1525 Otsuka, M. and Ohmori, S.: Simple and sensitive determination of 2,3-butanediol in biological samples by gas chromatography with electron-capture detection. *J. Chromatogr. B*, 654 (1994) 1-7.
- 1526 Williamson, S.A. and Iverson, W.G.: Determination of short-chain diols and selected fermentation by-products in beer. *J. Am. Soc. Brew. Chem.*, 51 (1993) 114-118; C.A., 119 (1993) 179532a.
- 1527 Yu, A., Yang, W., Jin, Q. and Wang, F.: (Study on the microwave-induced plasma ionization detector – simultaneous determination of methyl alcohol and water by gas chromatography.) *Fenxi Huaxue*, 21 (1993) 1337-1339; C.A., 120 (1994) 123863n.
- 1528 Zhou, X., Wang, Y., Li, S. and Zheng, G.: (Studies on self-association of *n*-alcohols by gas-liquid chromatography.) *Gaodeng Xuexiao Huaxue Xuebao*, 14 (1993) 817-820; C.A., 120 (1994) 16679k.
- See also 1289, 1564, 1567, 1582, 1684, 1720, 1799, 1896.
7. PHENOLS
- 1529 Boyd, T.J.: Identification and quantification of mono-, di- and trihydroxybenzenes (phenols) at trace concentrations in seawater by aqueous acetylation and gas chromatographic-mass spectrometric analysis. *J. Chromatogr. A*, 662 (1994) 281-292.
- 1530 Buchholz, K.D. and Pawliszyn, J.: Determination of phenols by solid-phase microextraction and gas chromatographic analysis. *Environ. Sci. Technol.*, 27 (1993) 2844-2848.
- 1531 Jubasek, P., Slimak, M., Brezny, R., Vodny, S. and Kosik, M.: Classification of chlorinated phenols by pattern recognition method on the basis of mass spectra losses. *Chem. Papers (Bratislava)*, 47 (1993) 93-98; C.A., 119 (1993) 233514m.
- 1532 Kim, T., Hong, J., Rhee, J. and Kim, K.: Analysis of nitrophenols by gas chromatography/mass spectrometry using flash heater derivatization. *Bull. Korean Chem. Soc.*, 14 (1993) 529-531; C.A., 120 (1994) 94391z.
- 1533 Mussman, P., Levsen, K. and Radeck, W.: Gas chromatographic determination of phenol in aqueous samples after solid phase extraction. *Fresenius J. Anal. Chem.*, 348 (1994) 654-659.
- 1534 Wild, S.R., Harrad, S.J. and Jones, K.C.: Chlorophenols in digested U.K. sewage sludges. *Water Res.*, 27 (1993) 1527-1534; C.A., 119 (1993) 209782b.
- 1535 Yakovlev, V.S., Kryuchkova, A.P., Kugucheva, E.E., Medvedev, A.R., Chulkov, N.G. and Kalitina, N.N.: (Gas-chromatography method for determining phenol derivatives in complex mixtures.) U.S.S.R. SU 1,798,681 (Cl. G01N30/06), 28 Feb. 1993, Appl. 4,816,232, 19 Apr. 1990; C.A., 120 (1994) 44834a.
- See also 1544, 1555.
8. SUBSTANCES CONTAINING HETEROCYCLIC OXYGEN
- 8a. Flavonoids
- 1536 Pommier, F., Ackermann, R., Sioufi, A. and Godbillon, J.: Determination of acenocoumarol in human plasma by capillary gas chromatography with mass-selective detection. *J. Chromatogr. B*, 654 (1994) 35-41.
- See also 1591.
- 8b. Aflatoxins and other mycotoxins
- 1537 Kanhere, S.R. and Weber, D.: Analysis of Canadian and imported beers for *Fusarium* mycotoxins by gas chromatography-mass spectrometry. *Food Addit. Contam.*, 10 (1994) 381-389; C.A., 120 (1994) 161888g.
- 1538 Young, J.C. and Games, D.E.: Analysis of *Fusarium* mycotoxins by gas chromatography-Fourier transform infrared spectroscopy. *J. Chromatogr. A*, 663 (1994) 211-218.
- 8c. Other compounds with heterocyclic oxygen (incl. tannins)
- 1539 Arakelyan, V.G.: (Gas chromatographic determination of furfural in transformer oil for transformer diagnostic.) *Elektrotehnika*, No. 2 (1993) 17-19; C.A., 119 (1993) 285527n.
- 1540 Ciccio, P., Brancalone, E. and Cecinato, A.: Updating the UNI-CHIM method for the determination of PCDD and PCDF in waste incineration plants. *Fresenius J. Anal. Chem.*, 348 (1994) 131-135.
- 1541 Darskus, R., Hagenmaier, H., von Holst, C. and Beckmann, L.: Rapid determination of 2,3,7,8-tetrachlorodibenzo-*p*-dioxin in thermally treated soil. *Fresenius J. Anal. Chem.*, 348 (1994) 148-149.
- 1542 Fraisse, D., Paisse, O., Hong, L.N. and Gonnord, M.F.: Improvements in GC/MS strategies and methodologies for PCDD and PCDF analysis. I. Evaluation of the non-polar DB-5ms column. *Fresenius J. Anal. Chem.*, 348 (1994) 154-158.
- 1543 Galletti, G.C. and Antonelli, A.: The potential of pyrolysis-(methylation)/gas chromatography/mass spectrometry in the analysis of wine polyphenolics. *Rapid Commun. Mass Spectrom.*, 7 (1993) 656-658; C.A., 119 (1993) 179518a.
- 1544 Grochowalski, A., Wybraniec, S., Gorski, L. and Sokolowski, M.: Solid phase extraction and capillary GC-ECD analysis of polychlorinated dibenzo-*p*-dioxins in chlorinated phenols. *Chem. Anal. (Warsaw)*, 38 (1993) 279-286.

- 1545 Kjeller, L.O., Kulp, S.E., Jonsson, B. and Rappe, C.: Methodology for the determination of polychlorinated dibenzo-*p*-dioxins and dibenzofurans in sediment samples. *Toxicol. Environ. Chem.*, 39 (1993) 1-12; C.A., 120 (1994) 37679a.
- 1546 Kocan, A., Petrik, J., Chovanová, J. and Drobná, B.: Method for the group separation of non-*ortho*-, mono-*ortho*- and multi-*ortho*-substituted polychlorinated biphenyls and polychlorinated dibenzo-*p*-dioxins/polychlorinated dibenzofurans using activated carbon chromatography. *J. Chromatogr. A*, 665 (1994) 139-153.
- 1547 Larsen, B. and Facchetti, S.: Use of supercritical fluid extraction in the analysis of polychlorinated dibenzodioxins and dibenzofurans. *Fresenius J. Anal. Chem.*, 348 (1994) 159-162.
- 1548 Luijk, R., Akkerman, D.M., Slot, P., Olie, K. and Kapteijn, F.: Mechanism of formation of polychlorinated dibenzo-*p*-dioxins and dibenzofurans in the catalyzed combustion of carbon. *Environ. Sci. Technol.*, 28 (1994) 312-321.
- 1549 Rappe, C.: Dioxin, pattern and source identification. *Fresenius J. Anal. Chem.*, 348 (1994) 63-75.
- 1550 Schimmel, H., Grieping, B., Maier, E.A., Kramer, G.N., Roos, A.H. and Tuinstra, L.G.M.T.: Intercomparison study on milk powder fortified with PCDD and PCDF. *Fresenius J. Anal. Chem.*, 348 (1994) 37-46.
- 1551 Steinwandter, H.: A simple and fast micro method for the determination of 1,2,3,4,6,7,8-heptachlorodibenzo-*p*-dioxin and octachloro-dibenzo-*p*-dioxin in sewage sludges by GC-ECD. *Fresenius J. Anal. Chem.*, 348 (1994) 385-389.
- 1552 Stief-Tauch, H.P.: The response of the EEC to the dioxin problem. *Fresenius J. Anal. Chem.*, 348 (1994) 47-52.
- 1553 Talka, E.: Experiences on the analysis of PCDDs and PCDFs in pulp and paper industry. *Fresenius J. Anal. Chem.*, 348 (1994) 121-125.
- 1554 Verotta, L. and Peterlongo, F.: Selective extraction of phenolic components from *Ginkgo biloba* extracts using supercritical carbon dioxide and off-line capillary gas chromatography/mass spectrometry. *Phytochem. Anal.*, 4 (1993) 178-182; C.A., 120 (1994) 3760e.
- 1555 Yajima, S., Kondou, T. and Hoshi, K.: (Analysis of dioxins and chlorophenols in flue gas from the municipal solid waste incinerator.) *Ishikawajima-Harima Giho*, 33 (1993) 292-297; C.A., 120 (1994) 13881j.
- See also 1412, 1484, 1814.
9. OXO COMPOUNDS, ETHERS, EPOXIDES AND QUINONES
- 1556 Bagchi, D., Bagchi, M., Hassoun, E.A. and Stohs, S.J.: Detection of paraquat-induced *in vivo* lipid peroxidation by gas chromatography/mass spectrometry and high-pressure liquid chromatography. *J. Anal. Toxicol.*, 17 (1993) 411-414; C.A., 120 (1994) 25281b.
- 1557 Dennis, K.J., Ichinose, T., Miller, M. and Shibamoto, T.: Gas chromatographic determination of vapor-phase biomarkers formed from rats dosed with carbon tetrachloride. *J. Appl. Toxicol.*, 13 (1993) 301-303; C.A., 119 (1993) 175489u.
- 1558 Ebeler, S.E., Hinrichs, S.H., Clifford, A.J. and Shibamoto, T.: Volatile carbonyl levels in tissues of transgenic mice with nerve sheath tumors. *J. Chromatogr. B*, 654 (1994) 9-18.
- 1559 Garner, C.M.: The stereochemistry of commercial 2,6-dimethylcyclohexanone: an application of capillary gas chromatography. *J. Chem. Educ.*, 70 (1993) A310-A311; C.A., 120 (1994) 105971s.
- 1560 Hewala, I.I.: Gas chromatographic and difference spectrophotometric techniques for the determination of benzaldehyde in benzyl alcohol. *J. Pharm. Biomed. Anal.*, 12 (1994) 73-79; C.A., 120 (1994) 173626h.
- 1561 Kanai, H., Inouye, V., Goo, R., Chow, R., Yazawa, L. and Maka, J.: GC/MS analysis of MTBE, ETBE, and TAME in gasolines. *Anal. Chem.*, 66 (1994) 924-927.
- 1562 Li, D., Yue, W., Liu, L., Tian, G., Li, J., Liu, D. and Li, Z.: (Gas chromatographic determination of acetone in expiratory air.) *Zhonghua Laodong Weisheng Zhiyebing Zazhi*, 11 (1993) 305-307; C.A., 120 (1994) 98571f.
- 1563 Manjarrez, A., Capella, S., Villarruel, E. and Pablo Garcia, F.: Investigation of methods for determining MTBE in gasolines. *Riv. Combust.*, 46 (1992) 337-340; C.A., 119 (1993) 206701p.
- 1564 Swarin, S.J., Loo, J.F., Chladek, E., Drouillard, M.S. and Tejada, S.B.: Advanced emissions speciation methodologies for the auto/oil quality improvement research program - II. Aldehydes, ketones, and alcohols. *Soc. Automot. Eng., Spec. Publ.*, SP-920 (1993) 199-207; C.A., 119 (1993) 209345m.
- 1565 Wang, M., Teng, W., Liu, K., Feng, J., Su, L., Xu, L. and You, G.: (Study of radiolytic effect on residual ethylene oxide in perfusion devices for single use using gas chromatography.) *Fushe Yanjiu Yu Fushe Gongyi Xuebao*, 11, No. 1 (1993) 31-35; C.A., 120 (1994) 100796d.
- See also 1412, 1514, 1539, 1904.
10. CARBOHYDRATES
- 10a. Mono and oligosaccharides. Structural studies
- 1566 Cases, M.R., Stortz, C.A. and Cerezo, A.S.: Separation and identification of partially ethylated galactoses as their acetylated aldonitriles and alditols by capillary gas chromatography and mass spectrometry. *J. Chromatogr. A*, 662 (1994) 293-299.
- 1567 Coll Hellin, L., Gutierrez Ruiz, L. and Zapata Revilla, A.: Gas chromatographic determination of sugars and polyalcohols in apple liqueurs. *Z. Lebensm.-Unters. Forsch.*, 196 (1993) 49-52; C.A., 119 (1993) 179526b.
- 1568 D'Ambr, A.J. and Gray, G.R.: Molar-response factors for the quantitative analysis of fully methylated methyl 2-acetamido-2-deoxyhexopyranosides by gas-liquid chromatography with flame-ionization detection. *Carbohydr. Res.*, 247 (1993) 299-303; C.A., 120 (1994) 22820r.
- 1569 Dumas, F., Aussel, C., Pernet, P., Martin, C. and Giboudeau, J.: Gas chromatography applied to the lactulose-mannitol intestinal permeability test. *J. Chromatogr. B*, 654 (1994) 276-281.
- 1570 Fox, A. and Black, G.E.: Identification and detection of carbohydrate markers for bacteria. Derivatization and gas chromatography-mass spectrometry. In: *Mass Spectrometry for the Characterization of Microorganisms. ACS Symp. Ser.*, Vol. 541, ACS, Washington, 1994, pp. 107-131; C.A., 120 (1994) 101303c - a review with 60 refs.

- 1571 Fox, A., Rosario, R.M.T. and Larsson, L.: Monitoring of bacterial sugars and hydroxy fatty acids in dust from air conditioners by gas chromatography-mass spectrometry. *Appl. Environ. Microbiol.*, 59 (1993) 4354-4360; *C.A.*, 120 (1994) 98561c.
- 1572 Moers, M.E.C., Jones, D.M., Eakin, P.A., Fallick, A.E., Griffiths, H. and Larter, S.R.: Carbohydrate diagenesis in hypersaline environments: applications of GC-IRMS to the stable isotope analysis of derivatized saccharides from surficial and buried sediments. *Org. Geochem.*, 20 (1993) 927-933; *C.A.*, 120 (1994) 112109v.
- 1573 Patz, C.D., Zimmer, E. and Dietrich, H.: (Gas chromatographic analysis of monosaccharides as the alditol acetates compared to high-performance ion chromatographic (HPIC) determinations.) *Dtsch. Lebensm.-Rundsch.*, 89 (1993) 1-6; *C.A.*, 119 (1993) 179467h.
- 1574 Previs, S.F., Ciralo, S.T., Fernandez, C.A., Beylot, M., Agarwal, K.C., Soloviev, M.V. and Brunengruber, H.: Use of [6,6-²H₂] glucose and of low-enrichment [U-¹³C₆] glucose for sequential or simultaneous measurements of glucose turnover by gas chromatography-mass spectrometry. *Anal. Biochem.*, 218 (1994) 192-196.
- 1575 Reznicek, G., Susman, O. and Boehm, K.: (Determination of the absolute configuration of monosaccharides from plant glycosides by gas chromatography-mass spectroscopy.) *Sci. Pharm.*, 61 (1993) 35-45; *C.A.*, 120 (1994) 134945s.
- 1576 Rocchiccioli, F., Auclair, M. and Bougneres, P.-F.: Determination of ¹³C enrichment at specific positions of plasma glucose by gas chromatography-mass spectrometric analysis of the 1,2:5,6-diisopropylidene-3-O-acetyl- α -furanosyl derivative. *Anal. Biochem.*, 218 (1994) 358-363.
- 1577 Rubin, L.J., Hsu, F.F. and Sherman, W.R.: Measurement of inositol triphosphate by gas chromatography/mass spectrometry: femtomole sensitivity provided by negative-ion chemical ionization mass spectrometry in submilligram quantities of issue. *Methods Neurosci.*, 18 (1993) 201-212; *C.A.*, 120 (1994) 130311j.
- 1578 Takano, R., Kamei-Hayashi, K., Hara, S. and Hirase, S.: Assignment of the absolute configuration of partially methylated galactoses by combined gas-liquid chromatography-mass spectrometry. *Biosci., Biotechnol., Biochem.*, 57 (1993) 1195-1197; *C.A.*, 120 (1994) 77534f.
- 1579 Wiesner, D.A. and Sweeley, C.C.: Microscale analysis of glycosphingolipids by methanolysis, peracetylation, and gas chromatography. *Anal. Biochem.*, 217 (1994) 316-322.
- 1580 Wu, Y. and Gretz, M.R.: Stable isotope labeling method for studies of saccharide metabolism in *Agardhiella subulata*. *Hydrobiologia*, 260-261 (1993) 595-600; *C.A.*, 119 (1993) 199108v.
11. ORGANIC ACIDS AND LIPIDS
- 11a. Organic acids and simple esters
- 1581 Ballesteros, E., Cárdenas, S., Gallego, M. and Valcárcel, M.: Determination of free fatty acids in dairy products by direct coupling of a continuous preconcentration ion-exchange-derivatization module to a gas chromatograph. *Anal. Chem.*, 66 (1994) 628-634.
- 1582 Beihoffer, J. and Ferguson, C.: Determination of selected carboxylic acids and alcohols in groundwater by GC-MS. *J. Chromatogr. Sci.*, 32 (1994) 102-106.
- 1583 Cartwright, I.J.: Analysis of fatty acids by gas liquid chromatography. *Methods Mol. Biol. (Totowa)*, 19 (1993) 183-195; *C.A.*, 120 (1994) 3770h.
- 1584 Christie, W.W., Brechany, E.Y., Sebedio, J.L. and le Quere, J.L.: Silver ion chromatography and gas chromatography-mass spectrometry in the structural analysis of cyclic monoenoic acids formed in frying oils. *Chem. Phys. Lipids*, 66 (1993) 143-153; *C.A.*, 120 (1994) 75775m.
- 1585 De Leeuw, J.W. and Baas, M.: The behavior of esters in the presence of tetramethylammonium salts at elevated temperatures; flash pyrolysis or flash chemolysis? *J. Anal. Appl. Pyrolysis*, 26 (1993) 175-184.
- 1586 Diskina, D.E. and Zanozina, I.I.: (Use of gas chromatography for analysis of industrial effluents.) *Khim. Tekhnol. Topl. Masel*, No. 7 (1993) 38-39; *C.A.*, 120 (1994) 85034h.
- 1587 Fraser, A.D. and MacNeil, W.: Colorimetric and gas chromatographic procedures for glycolic acid in serum: the major toxic metabolite of ethylene glycol. *J. Toxicol., Clin. Toxicol.*, 31 (1993) 397-405; *C.A.*, 120 (1994) 71017t.
- 1588 Gibson, K.M., ten Brink, H.J., Schor, S.M., Kok, R.M., Bootsma, A.H., Hoffmann, G.F. and Jakobs, C.: Stable-isotope dilution analysis of D- and L-2-hydroxyglutaric acid: application to the detection and prenatal diagnosis of D- and L-2-hydroxyglutaric acidemias. *Pediatr. Res.*, 34 (1993) 277-280; *C.A.*, 119 (1993) 244937c.
- 1589 Gilarova, R., Voldrich, M., Demnerova, K., Cerovsky, M. and Dobias, J.: Gas chromatographic analysis of cellular fatty acids in the identification of lactic acid bacteria. *Potravin. Vedy*, 11 (1993) 197-208; *C.A.*, 120 (1994) 7015p.
- 1590 Hall, K.C., Else, M.A. and Jackson, M.B.: Determination of 1-aminocyclopropane-1-carboxylic acid (ACC) in leaf tissue and xylem sap using capillary column gas chromatography and a nitrogen/phosphorus detector. *Plant Growth Regul.*, 13 (1993) 225-230; *C.A.*, 120 (1994) 26698e.
- 1591 Heimler, D. and Pieroni, A.: Capillary gas chromatography of plant tissues and soil phenolic acids. *Chromatographia*, 38 (1994) 475-478.
- 1592 Kelley, R.I.: Quantification of 3-methylglutaconic acid in urine, plasma, and amniotic fluid by isotope-dilution gas chromatography-mass spectrometry. *Clin. Chim. Acta*, 220 (1993) 157-164.
- 1593 Khalfallah, Y., Normand, S., Tissot, S., Pachiardi, C., Beylot, M. and Riou, J.P.: Determination of δ^{13} C values of plasma lactate by gas chromatography/isotope ratio mass spectrometry. *Biol. Mass Spectrom.*, 22 (1993) 707-711; *C.A.*, 120 (1994) 101154e.

See also 1383.

- 1594 Lee, B.Y. and Thurmon, T.F.: Compact gas chromatography used with thin-layer chromatography for assessment of abnormalities of organic acids. *Clin. Chim. Acta*, 218 (1993) 215-222.
- 1595 Li, X.-d. and Lin, Z.-s.: Determination of sulphonic compounds as their thiotrifluoroacetate derivatives by gas chromatography with ion trap detection. *J. Chromatogr. A*, 667 (1994) 219-223.
- 1596 Lu, J., Wu, X. and Zhou, Q.: (Gas chromatographic determination of 18 α -glycyrrhizic acid and related compounds in rat plasma.) *Zhongguo Yaoxue Zazhi*, 28 (1993) 552-554; *C.A.*, 119 (1993) 261921k.
- 1597 Mueller, M.J. and Brodschelm, W.: Quantification of jasmonic acid by capillary gas chromatography—negative ion chemical ionization—mass spectrometry. *Anal. Biochem.*, 218 (1994) 425-435.
- 1598 Patterson, B.W. and Wolfe, R.R.: Concentration dependence of methyl palmitate isotope ratios by electron impact ionization gas chromatography/mass spectrometry. *Biol. Mass Spectrom.*, 22 (1993) 481-486; *C.A.*, 119 (1993) 221082f.
- 1599 Read, G., Richardson, N.R. and Wickers, D.G.: Determination of octadecanoic acids in human serum: a critical appraisal. *Analyt. (Cambridge)*, 119 (1994) 393-396.
- 1600 Rivis, I.F. and Danylyk, B.B.: (Determination of the fatty acid composition of phospholipids, triacylglycerols, and esters of cholesterol in biologic material.) *S- kh. Biol.*, No. 6 (1992) 163-165; *C.A.*, 119 (1993) 177069f.
- 1601 Rozes, N., Garbay, S., Denayrolles, M. and Lonvaud-Funel, A.: A rapid method for the determination of bacterial fatty acid composition. *Lett. Appl. Microbiol.*, 17 (1993) 126-131; *C.A.*, 120 (1994) 128762.
- 1602 Šajbidor, J., Certik, M. and Grego, J.: Lipid analysis of baker's yeast. *J. Chromatogr. A*, 665 (1994) 191-195.
- 1603 Sarycheva, Ye.N.: (Gas chromatographic determination of maleic ester in the air.) *Khim.-Farm. Zh.*, 27, No. 5 (1993) 53-54; *C.A.*, 120 (1994) 13873h.
- 1604 Supaporn, P., Shuichi, A., Kaoru, O. and Hideaki, H.: Analysis of fatty acids in methyl ester form by using relative retention times obtained from gas chromatography techniques. *Kanzei Chuo Bunsekishoho*, 32 (1993) 61-85; *C.A.*, 119 (1993) 248278m.
- 1605 Thomas, B. and Schoentube, E.: (Determination of γ -hydroxybutyric acid in human plasma by gas chromatography.) *Pharmazie*, 48 (1993) 623-624; *C.A.*, 119 (1993) 240828v.
- 1606 Tisza, S. and Molnár-Perl, I.: GC-MS quantitation of isocitric acid in the presence of a large excess of citric acid. *J. High Resolut. Chromatogr.*, 17 (1994) 165-168.
- 1607 Van Loon, W.M.G.M. and Boon, J.J.: Quantitative pyrolysis—gas chromatography—mass spectrometry and pyrolysis—mass spectrometry of chlorolignosulphonic acids. *TrAC*, 13 (1994) 169-176.
- 1608 Yuan, S., Zhang, B. and Huang, H.: (GC-MS analysis of influence of processing on volatile oil in *Daphne genkwa* Sieb. et Zucc.) *Zhongguo Zhongyao Zazhi*, 18 (1993) 595-597; *C.A.*, 120 (1994) 144286s.
- 1609 Zelles, L. and Bai, Q.Y.: Fractionation of fatty acids by solid phase extraction and their quantitative analysis by GC-MS. *Soil Biol. Biochem.*, 25 (1993) 495-507; *C.A.*, 119 (1993) 244669s.
- 1610 Zhao, Q., Gong, B., Liu, J. and Hu, L.: (The effect of exogenous SOD on unsaturated fatty acid in cardiac and hepatic cell membrane in mouse measured by gas chromatography.) *Zhongguo Yiyuan Yaoxue Zazhi*, 13 (1993) 401-402; *C.A.*, 120 (1994) 153652k.
- 1611 Zhou, Y. and Zhuang, W.: (Capillary column gas chromatography for the determination of trace α -naphthalene acetic acid residual in fruits and vegetables.) *Fenxi Huaxue*, 21 (1993) 992; *C.A.*, 119 (1993) 269286c.
- See also 1269, 1365, 1514, 1571, 1620, 1638, 1639, 1650, 1664, 1665, 1684, 1876.
- 11b. *Prostaglandins*
- 1612 Daniel, V.C., Minton, T.A., Brown, N.J., Nadeau, J.H. and Morrow, J.D.: Simplified assay for the quantification of 2,3-dinor-6-keto-prostaglandin F_{1 α} by gas chromatography—mass spectrometry. *J. Chromatogr. B*, 653 (1994) 117-122.
- 1613 Obata, T., Nagakura, T., Kanmuri, M., Masaki, T., Maekawa, K. and Yamashita, K.: (Quantitative measurement of 9 α , 11 β -PGF₂ in human urine and plasma by gas chromatography—mass spectrometry.) *Nippon Iyo Masu Supekutoru Gakkai Koenshu*, 18 (1993) 97-100; *C.A.*, 120 (1994) 23691m.
- 1614 Yamashita, K., Enomoto, K., Ishibashi, M., Hiramatsu, M., Hayashi, Y. and Yamamoto, S.: (Development of microdetermination method of PG₁₂ metabolites by gas chromatography/selected ion monitoring.) *Nippon Iyo Masu Supekutoru Gakkai Koenshu*, 18 (1993) 119-122; *C.A.*, 120 (1994) 23692n.
- See also 1304.
- 11c. *Lipids and their constituents*
- 1615 Alonso, L.: Fat modification in Afuega'l Pitu cheese during ripening by capillary gas chromatography. *J. Am. Oil Chem. Soc.*, 70 (1993) 1035-1037.
- 1616 Antoniosi Filho, N.R., Carrilho, E. and Lancas, F.M.: Fast quantitative analysis of soybean oil in olive oil by high-temperature capillary gas chromatography. *J. Am. Oil Chem. Soc.*, 70 (1993) 1051-1053.
- 1617 Blomberg, L.G., Demirbaker, M. and Andersson, P.E.: Argention supercritical fluid chromatography for quantitative analysis of triacylglycerols. *J. Am. Oil Chem. Soc.*, 70 (1993) 939-946.
- 1618 Conte, L.S., Gangemi, G. and Boschelle, O.: (Analysis of fat used in bread manufacture. Application of capillary GC and HPLC.) *Riv. Ital. Sostanze Grasse*, 70 (1993) 395-400; *C.A.*, 120 (1994) 29621k.
- 1619 Dizabo, P. and Pepe, C.: (Study of wax esters of natural origin by gas chromatography/mass spectrometry.) *J. Chim. Phys. Phys.-Chim. Biol.*, 90 (1993) 1787-1795; *C.A.*, 119 (1993) 273806q.
- 1620 Ferreira-Dias, S. and Fonseca, M.M.R.: Enzymic glycerolysis of olive oil: a reaction system with major analytical problems. *Bio-technol. Tech.*, 7 (1993) 447-452; *C.A.*, 119 (1993) 269279c.
- 1621 Garcia Regueiro, J.A., Diaz, I., David, F. and Sandra, P.: Possibilities of PTV injection for the analysis of triglycerides. *J. High Resolut. Chromatogr.*, 17 (1994) 180-183.

- 1622 García Regueiro, J.A., Gibert, J. and Díaz, I.: Determination of neutral lipids from subcutaneous fat of cured ham by capillary gas chromatography and liquid chromatography. *J. Chromatogr. A*, 667 (1994) 225-233.
- 1623 Lercker, G., Cocchi, M., Turchetto, E., Capella, P., Zullo, C. and Caboni, M.F.: Determination of total lipid components by a high resolution gas chromatographic method. Application to the short-term fasting of rats. *Riv. Sci. Aliment.*, 22 (1993) 147-152; *C.A.*, 120 (1994) 105708m.
- 1624 Schneller, R. and Wullschleger, R.: (Quantitative determination of butter fat in fat mixtures by capillary gas chromatography.) *Mitt. Geb. Lebensmittelunters. Hyg.*, 83 (1992) 737-744; *C.A.*, 119 (1993) 179466g.
- 1625 Tognetti, C., Giovannoni, G. and Frega, N.: (Lipid composition of some safflower cultivars.) *Riv. Ital. Sostanze Grasse*, 70 (1993) 295-298; *C.A.*, 119 (1993) 248482y.
- 1626 Walker, J.T., Sonesson, A., Keevil, C.W. and White, D.C.: Detection of *Legionella pneumophila* in biofilms containing a complex microbial consortium by gas chromatography-mass spectrometry analysis of genus-specific hydroxy fatty acids. *FEMS Microbiol. Lett.*, 113 (1993) 139-144; *C.A.*, 120 (1994) 26964p.
- See also 1455, 1579, 1600, 1884, 1928.
- 11d. *Lipoproteins and their constituents*
- 1627 Van den Berg, J.J.M., Winterbourn, C.C. and Kuypers, F.A.: Hypochlorous acid-mediated modification of cholesterol and phospholipid: analysis of reaction products by gas chromatography-mass spectrometry. *J. Lipid Res.*, 34 (1993) 2005-2012; *C.A.*, 120 (1994) 25168v.
- See also 1637.
13. STEROIDS
- 13b. *Pregnane and androstane derivatives*
- 1628 Dikkeschei, L.D., Wolthers, B.G., Willemsse, P.H.B., van der Pol, H., de Ruyter-Buitenhuis, A.W. and Nagel, G.T.: The determination of delta-5-androstenediol and its sulphate in serum and urine by gas chromatography-mass spectrometry. *Clin. Endocrinol. (Oxford)*, 39 (1993) 469-474; *C.A.*, 120 (1994) 153903t.
- 1629 Iki, K.: (Measurement of 18-hydroxycortisol by gas chromatography equipped with single ion monitoring: establishment of the method and its clinical application.) *Kanazawa Daigaku Juzen Igakkai Zasshi*, 102 (1993) 238-247; *C.A.*, 120 (1994) 153881j.
- 1630 Leyssens, L., Royackers, E., Gielen, B., Missotten, M., Schoofs, J., Czech, J., Noben, J.P., Hendriks, L. and Raus, J.: Metabolites of 4-chlorotestosterone acetate in cattle urine as diagnostic markers for its illegal use. *J. Chromatogr. B*, 654 (1994) 43-54.
- 1631 Vermeulen, G.J., Lambert, J.G.D., Lenczowski, M.J.P. and Goos, H.J.T.: Steroid hormone secretion by testicular tissue from African catfish, *Clarias gariepinus*, in primary culture: Identification and quantification by gas chromatography-mass spectrometry. *Fish Physiol. Biochem.*, 12 (1993) 21-30; *C.A.*, 120 (1994) 102562y.
- 13c. *Estrogens*
- 1632 Iijima, M., Hoshino, Y., Nose, N., Tokumaru, Y., Miyazaki, T. and Matsumoto, M.: (Determination of diethylstilbestrol glucuronide in beef by gas chromatography.) *Jpn. J. Toxicol. Environ. Health*, 39 (1993) 311-316; *C.A.*, 119 (1993) 224533q.
- 1633 Nakagawa, Y., Enomoto, K., Nakano, K. and Yamashita, K.: (Microdetermination of estrogens by gas chromatography/selected ion recording. An application to the measurement of aromatase activity.) *Nippon Iyo Masu Supekutoru Gakkai Koenshu*, 18 (1993) 123-126; *C.A.*, 120 (1994) 23693p.
- 1634 Van der Vliis, E., Irth, H., Tjaden, U.R. and van der Greef, J.: Potential of receptor-ligand interactions for sample handling in liquid and gas chromatography. *J. Chromatogr. A*, 665 (1994) 233-241.
- 13d. *Sterols*
- 1635 Kim, D.H., Lee, K.-J. and Heo, G.S.: Analysis of cholesterol and cholesteryl esters in human serum using capillary supercritical fluid chromatography. *J. Chromatogr. B*, 655 (1994) 1-8.
- 1636 Mariani, C., Gasparoli, A., Venturini, S. and Fedeli, E.: (Analytical differentiation of virgin and refined lard.) *Riv. Ital. Sostanze Grasse*, 70 (1993) 275-278; *C.A.*, 119 (1993) 269288e.
- 1637 Tanaka, M. and Kanamaru, S.: Capillary gas chromatography quantification of cholesterol in copper-oxidized low-density lipoprotein. *Biol. Pharm. Bull.*, 16 (1993) 538-543; *C.A.*, 119 (1993) 220817n.
- 1638 Tazuma, S., Hatsushika, S., Yamashita, G., Aihara, N., Sasaki, M., Horikawa, K., Yamashita, Y., Teramen, K., Ochi, H., Hirano, N., Miura, H., Ohya, T., Hino, H. and Kajiyama, G.: Simultaneous microanalysis of biliary cholesterol, bile acids and fatty acids in lecithin using capillary column gas chromatography: an advantage to assess bile lithogeneity. *J. Chromatogr. B*, 653 (1994) 1-7.
- 1639 Wang, H., Hachey, D.L., Liu, D., Zhang, S. and Dedley, M.A.: Microanalysis of cholesterol, phospholipid and medium- and long-chain fatty acids in biological materials. *Anal. Biochem.*, 218 (1994) 74-79.
- 1640 Watanabe, H., Mano, N., Ikegawa, S., Goto, J. and Nambara, T.: (Study on biosynthesis of bile acids by gas chromatography-mass spectrometry with negative ion chemical ionization detection.) *Nippon Iyo Masu Supekutoru Gakkai Koenshu*, 18 (1993) 105-106; *C.A.*, 120 (1994) 27976f.
- See also 1464, 1600, 1627, 1877.
- 13e. *Bile acids and alcohols*
- See 1638.
15. TERPENES AND OTHER VOLATILE AROMATIC COMPOUNDS
- 15a. *Terpenes*
- 1641 Casabianca, H. and Graff, J.-B.: Separation of linalyl acetate enantiomers: application to the authentication of bergamot food products. *J. High Resolut. Chromatogr.*, 17 (1994) 184-186.

- 1642 Morin, P.: Separation of geometrical isomers of sesquiterpene and diterpene alcohols on bare silica by carbon dioxide supercritical fluid chromatography. *Fresenius J. Anal. Chem.*, 348 (1994) 227-228.
- 1643 Riemer, D.D., Milne, P.J., Farmer, C.T. and Zika, R.G.: Determination of terpenes and related compounds in semi-urban air by GC-MSD. *Chemosphere*, 28 (1994) 837-850.
- 1644 Staccioli, G., Mellerio, G. and Bambagiotti, A.M.: Investigation on terpene-related hydrocarbons from a pliocenic fossil wood. *Holzforchung*, 47 (1993) 339-342; *C.A.*, 120 (1994) 134841e.
- 1645 Sybilska, D., Cieslak, M. and Goronowicz, J.: (Method for micro-preparative separation of chiral enantiomers of terpene hydrocarbons using gas chromatography.) *Pol. PL 158,846 (Cl. C07C13/00)*, 30 Oct. 1992, *Appl. 277,627*, 8 Feb. 1989; 10 pp.; *C.A.*, 119 (1993) 271433s.
- 1646 Sybilska, D., Kowalczyk, J., Asztemborska, M., Ochocka, R.J., Lamparczyk, H.: Chromatographic studies of the enantiomeric composition of some therapeutic composition applied in the treatment of liver and kidney diseases. *J. Chromatogr. A*, 665 (1994) 67-73.
- See also 1311, 1426, 1431, 1434, 1867.
- 15b. Essential oils**
- 1647 Adedeji, B.A.: Flavor characterization of different varieties of vanilla beans (*Vanilla planifolia* Andrews and *Vanilla tahitensis* Moore) by direct thermal desorption-gas chromatography and gas chromatography-mass spectrometry. *Avail. Univ. Microfilms Int.*, Order No. DA9320513, 1993, 242 pp.; *C.A.*, 120 (1994) 132631n.
- 1648 Faber, B., Dietrich, A. and Mosandl, A.: Chiral compounds of essential oils. XV. Stereodifferentiation of characteristic compounds of *Mentha* species by multi-dimensional gas chromatography. *J. Chromatogr. A*, 666 (1994) 161-165.
- 1649 Ghizzoni, C. and Bottini, A.: Application of gas chromatographic chiral columns in some essential oils. *Riv. Ital. EPPOS*, 3, No. 8 (1992) 9-19; *C.A.*, 119 (1993) 210209b.
- 1650 Guo, R. and Xu, C.: (Isolation and identification of essential oil in American ginseng (*Panax quinquefolius*) cultured in Heilongjiang.) *Zhongcaoyao*, 24 (1993) 570-571; *C.A.*, 120 (1994) 144287t.
- 1651 Hawer, W.S., Ha, J., Hwang, J. and Nam, Y.: Effective separation and quantitative analysis of major heat principles in red pepper by capillary gas chromatography. *Food Chem.*, 49 (1993) 99-103; *C.A.*, 120 (1994) 29622m.
- 1652 He, R., Wang, J., Zhang, W., Yu, X. and Ding, J.: (Essential oil from *Micromeria biflora*.) *Zhongguo Yaoxue Zazhi*, 28 (1993) 536-537; *C.A.*, 120 (1994) 116548k.
- 1653 Ji, L., Xu, Z. and Pan, J.: (GC-MS analysis of essential oil from fruits of *Gardenia jasminoides*.) *Zhongguo Yaoxue Zazhi*, 28 (1993) 398-400; *C.A.*, 120 (1994) 116903x.
- 1654 Ji, L., Xu, Z. and Pan, J.: (GC-MS analysis of the essential oil from the root *Ligusticum brachylobum* Franch.) *Zhongguo Zhongyao Zazhi*, 18 (1993) 294-295; *C.A.*, 119 (1993) 210856k.
- 1655 Ji, N., Chang, L., Li, Y. and Hu, Y.: (Analysis of Baihuayou (white flower oil) by capillary gas chromatography.) *Zhongcaoyao*, 24 (1993) 236-237; *C.A.*, 119 (1993) 234159m.
- 1656 Krock, K.A., Ragunathan, N. and Wilkins, C.L.: Multidimensional gas chromatography coupled with infrared and mass spectrometry for analysis of eucalyptus essential oils. *Anal. Chem.*, 66 (1994) 425-430.
- 1657 Lewinsohn, E., Savage, T.J., Gijzen, M. and Croteau, R.: Simultaneous analysis of monoterpenes and diterpenoids of conifer oleoresin. *Phytochem. Anal.*, 4 (1993) 220-225; *C.A.*, 120 (1994) 157823q.
- 1658 Li, M. and Zherong, J.: (The volatile constituents of the essential oil and their distribution in different parts of drug sweetflag (*Acorus calamus*.) *Zhongcaoyao*, 24 (1993) 459-461; *C.A.*, 120 (1994) 144284q.
- 1659 Li, Z. and Zhang, Q.: (Separation and identification of lactone components in *Ligusticum chuanxiong* Hort. and *Cnidium officinale* Mak. by GC-MS.) *Yaowu Fenxi Zazhi*, 13 (1993) 187-189; *C.A.*, 119 (1993) 210861h.
- 1660 Pino, J.A., Borges, P. and Roncal, E.: (Differentiation of the essential oils of four species of oregano by using gas-liquid chromatography.) *Alimentaria (Madrid)*, 244 (1993) 105-107; *C.A.*, 119 (1993) 224514j.
- 1661 Sattar, A., Mahmud, S. and Khan, S.A.: Citrus oils. Part-II. Gas chromatographic analysis of the essential oil of *Citrus limetta* var. Mitha. *Pak. J. Sci. Ind. Res.*, 35 (1992) 449-450; *C.A.*, 119 (1993) 179585v.
- 1662 Vilegas, J.H.Y., Lancas, F.M., Vilegas, W. and Pozetti, G.L.: Off-line supercritical fluid extraction-high resolution gas chromatography applied to the study of *Moraceae* species. *Phytochem. Anal.*, 4 (1993) 230-234; *C.A.*, 120 (1994) 173589y.
- 1663 Yang, Y., Xiang, B., An, D., Liang, X., Yuan, C., Cao, R., Peng, J. and Sheng, L.: (Analysis of the volatile oils from the drugs *Bupleurum*.) *Zhongcaoyao*, 24 (1993) 289-291; *C.A.*, 119 (1993) 278876x.
- See also 1403, 1425, 1439, 1454, 1870.
- 15c. Bitter substances**
- See 1464.
- 16. NITRO AND NITROSO COMPOUNDS**
- 1664 Crowley, T.O. and Larson, R.A.: HPLC and GC methods for analysis of water-soluble nitrosulfonic acids. *J. Chromatogr. Sci.*, 32 (1994) 57-60.
- 1665 Soják, L., Perjéssy, A., Kubinec, R., Giumanini, A.G. and Strazolini, P.: Identification of the isomers from mono- and dinitration of phenyl- and diphenylacetic acids by gas chromatography with Fourier transform infrared and mass spectrometric detection. *J. Chromatogr. A*, 665 (1994) 169-173.
- 1666 Sun, J., Ma, J. and Sui, J.: (GC determination of trace amounts of nitrobenzene in water.) *Lihua Jianyan, Huaxue Fence*, 29 (1993) 275-277; *C.A.*, 120 (1994) 14512b.
- 1667 Xia, B., Feng, Y. and Wu, L.: (Gas chromatographic determination of nitroglycerin in the workplace air.) *Gongye Weisheng Yu Zhiyebing*, 18 (1992) 232-234; *C.A.*, 119 (1993) 187380q.
- See also 1405, 1532.

17. AMINES, AMIDES AND RELATED NITROGEN COMPOUNDS

17a. Amines and polyamines

- 1668 Durden, D.A. and Davis, B.A.: Determination of regional distributions of phenylethylamine and *meta*- and *para*-tyramine in rat brain regions and presence in human and dog plasma by an ultra-sensitive negative chemical ion gas chromatography-mass spectrometric (NCI-GC-MS) method. *Neurochem. Res.*, 18 (1993) 995-1002; *C.A.*, 119 (1993) 177105q.
- 1669 Mohnke, M., Schmidt, B., Schmidt, R., Buijten, J.C. and Mussche, P.: Application of a fused silica column to the determination of very volatile amines by gas-solid chromatography. *J. Chromatogr. A*, 667 (1994) 334-339.
- 1670 Nakamura, A. and Kisaki, H.: (Method for determination of trimethylamine in fish muscle by head space gas chromatography.) *Nippon Kasei Gakkaishi*, 44 (1993) 781-785; *C.A.*, 119 (1993) 269295e.
- 1671 Skarping, G., Dalene, M. and Lind, P.: Determination of toluenediamine isomers by capillary gas chromatography and chemical ionization mass spectrometry with special reference to the biological monitoring of 2,4- and 2,6-toluene diisocyanate. *J. Chromatogr. A*, 663 (1994) 199-210.
- 1672 Vatsala, S., Bansal, V., Tuli, D.K., Rai, M.M., Jain, S.K., Srivastava, S.P. and Bhatnagar, A.K.: Gas chromatographic determination of residual hydrazine and morpholine in boiler feed water and steam condensates. *Chromatographia*, 38 (1994) 456-460.

See also 1306, 1520, 1709, 1815.

17b. Catecholamines and their metabolites

- 1673 Watson, D.G., Zhou, P., Midgley, J.M., Milligan, C.D. and Kaiser, K.: The determination of biogenic amines in four strains of the fruit fly *Drosophila melanogaster*. *J. Pharm. Biomed. Anal.*, 11 (1993) 1145-1149; *C.A.*, 120 (1994) 101193s.

17d. Other amine derivatives and amides (excl. peptides)

- 1674 Guan, F., Zhang, Y. and Luo, Y.: (A gas-chromatographic method for the determination of tetramine in rodenticide baits.) *Junshi Yixue Kexueyuan Yuankan*, 17 (1993) 292-293; *C.A.*, 120 (1994) 71398t.
- 1675 Rasmussen, H.T., Friedman, S.K., Mustilli, A.J., McDonough, R. and McPherson, B.P.: Analysis of methyl neodecanamide in lake water by reversed-phase high performance liquid chromatography and gas chromatography-mass spectrometry. *J. Liq. Chromatogr.*, 17 (1994) 589-601.
- 1676 Wan, H. and Ou, Q.: (Enantiomer separation on perpentylated and partially pentylated trifluoroacetylated α -cyclodextrin chiral capillary columns.) *Huaxue Xuebao*, 51 (1993) 796-803; *C.A.*, 119 (1993) 240723g.

See also 1672, 1795, 1858, 1871.

18. AMINO ACIDS AND PEPTIDES; CHEMICAL STRUCTURE OF PROTEINS

18a. Amino acids and their derivatives

- 1677 Allen, R.H., Stabler, S.P. and Lindenbaum, J.: Serum betaine, N,N-dimethylglycine and N-methylglycine levels in patients with cobalamin and folate deficiency and related inborn errors of metabolism. *Metab., Clin. Exp.*, 42 (1993) 1448-1460; *C.A.*, 120 (1994) 7380d.
- 1678 Corradini, R., Marchelli, R. and Palla, G.: Properties of L-phenylalanine tetraamides as chiral selectors for D,L-amino acids by capillary gas chromatography. *Chromatographia*, 38 (1994) 173-176.
- 1679 Corso, G., Esposito, M., Gallo, M., Dello Ruso, A. and Antonio, M.: Transformation of arginine into ornithine during the preparation of its *tert*-butyldimethylsilyl derivative for analysis by gas chromatography/mass spectrometry. *Biol. Mass Spectrom.*, 22 (1993) 698-702; *C.A.*, 120 (1994) 101153d.
- 1680 David, F., Beylot, M., Reider, M.W., Anderson, V.E. and Brunen-gruber, H.: Assay of the concentration and ¹³C enrichment of acetate and acetyl-CoA by gas chromatography-mass spectrometry. *Anal. Biochem.*, 218 (1994) 143-148.
- 1681 Kataoka, H., Nakai, K., Katagiri, Y. and Makita, M.: Analysis of free and bound O-phosphoamino acids in urine by gas chromatography with flame photometric detection. *Biomed. Chromatogr.*, 7 (1993) 184-188.
- 1682 Patterson, B.W., Carraro, F. and Wolfe, R.R.: Measurement of nitrogen-15 enrichment in multiple amino acids and urea in a single analysis by gas chromatography/mass spectrometry. *Biol. Mass Spectrom.*, 22 (1993) 518-523; *C.A.*, 120 (1994) 26922y.
- 1683 Sato, K., Watabe, K., Ihara, T. and Hobo, T.: Preparative separation of amino acid enantiomers as N-TFA, O-isopropyl derivatives by continuous countercurrent gas-liquid chromatography. *Chirality*, 5 (1993) 246-249; *C.A.*, 120 (1994) 107684z.
- 1684 Šimek, P. and Heydová, A.: High resolution capillary gas chromatography and gas chromatography-mass spectrometry of protein and non-protein amino acids, amino alcohols, and hydroxycarboxylic acids as their *tert*-butyldimethylsilyl derivatives. *J. High Resolut. Chromatogr.*, 17 (1994) 145-152.
- 1685 Sorge, C., Schnitzer, M. and Schulten, H.R.: In-source pyrolysis-field ionization mass spectrometry and Curie-point pyrolysis-gas chromatography/mass spectrometry of amino acids in humic substances and soils. *Biol. Fertil. Soils*, 16 (1993) 100-110; *C.A.*, 120 (1994) 105758c.
- 1686 Takahashi, S., Matsubara, K., Hasegawa, M., Akane, A. and Shiono, H.: Detection and measurement of S-benzyl-N-acetylcysteine in urine of toluene sniffers using capillary gas chromatography. *Arch. Toxicol.*, 67 (1993) 647-650; *C.A.*, 120 (1994) 98619c.
- 1687 Wang, J., Huang, Z.-H., Gage, D.A. and Watson, J.T.: Analysis of amino acids by gas chromatography-flame ionization detection and gas chromatography-mass spectrometry: Simultaneous derivatization of functional groups by an aqueous-phase chloroformate-mediated reaction. *J. Chromatogr. A*, 663 (1994) 71-78.

- 1688 Williams, J., Lang, D., Smith, J.A. and Lewis, M.J.: Plasma L-arginine levels in a rabbit model of hypercholesterolemia. *Biochem. Pharmacol.*, 46 (1993) 2097-2099; *C.A.*, 120 (1994) 104027b.
- See also 1436, 1783, 1882.
- 18b. *Peptides, peptidic and proteinous hormones, growth factors*
- 1689 Stolowitz, M.L.: Thiobenzoylation method of peptide sequencing with gas chromatography and mass spectrometric detection. *U.S. US 5,246,865 (Cl. 436-89; G01N30/72)*, 21 Sep. 1993, Appl. 919,524, 24 Jul. 1992; 19 pp.; *C.A.*, 120 (1994) 27032v.
21. PURINES, PYRIMIDINES, NUCLEIC ACIDS AND THEIR CONSTITUENTS
- 21a. *Purines, pyrimidines, nucleosides, nucleotides*
- 1690 Conte, E.D. and Barry, E.F.: Gas chromatographic determination of caffeine in beverages by alkali aerosol flame ionization detection. *Microchem. J.*, 48 (1993) 372-376.
- 1691 Maekawa, H., Yamazaki, Y. and Yagasaki, K.: (Extraction of theobromine and caffeine from cocoa powder by DMSO [for gas chromatography]. (Report I.)) *Kanzei Chuo Bunsekishoho*, 32 (1993) 93-96; *C.A.*, 119 (1993) 269283z.
- 1692 Mandel, H.G., Kuzmiers, J.J., Dickens, B.F. and Anderson, L.W.: Quantitation of urinary 7-methyl adenine by gas chromatography-mass spectrometry using isotopically labeled internal standards. *Anal. Biochem.*, 217 (1994) 292-297.
- See also 1404.
22. ALKALOIDS
- 1693 Cooper, D.A. and Moore, J.M.: Femtogram on-column detection of nicotine by isotope dilution gas chromatography/negative ion detection mass spectrometry. *Biol. Mass Spectrom.*, 22 (1993) 590-594; *C.A.*, 119 (1993) 264244c.
- 1694 Garba, M., Mustapha, A. and Choudhury, B.K.: Quantitative estimation of carbazole alkaloids using gas liquid chromatography. *Ghana J. Chem.*, 1 (1993) 349-353; *C.A.*, 119 (1993) 210891t.
- 1695 Hasegawa, M., Matsubara, K., Maseda, C., Kagawa, M., Takata-shi, S. and Akane, A.: Morphine and codeine urinalyses after ingestion of bean-jam bun. *Jpn. J. Toxicol. Environ. Health*, 39 (1993) 415-420; *C.A.*, 120 (1994) 71008r.
- 1696 Li, Y., Liu, G., Mu, W., Zhang, Y., Zhang, J. and He, J.: (Gas chromatographic analysis of nicotine in indoor air.) *Huaxi Yike Daxue Xuebao*, 24 (1993) 209-212; *C.A.*, 119 (1993) 187377u.
- 1697 Urakawa, N., Nagata, T., Kudo, K., Kimura, K. and Imamura, T.: (Simultaneous determination of nicotine and cotinine in tobacco smoker's tissues by GC/MS.) *Nippon Iyo Masu Supekutoru Gak-kai Koenshu*, 18 (1993) 77-80; *C.A.*, 119 (1993) 264240y.
- See also 1855, 1864.
23. OTHER SUBSTANCES CONTAINING HETEROCYCLIC NITROGEN
- 23a. *Porphyrins and other pyrroles*
- 1698 Pretorius, W.G., Ebdon, L. and Rowland, S.J.: Development of a high-temperature gas chromatography-inductively coupled plasma mass spectrometry interface for the determination of metalloporphyrins. *J. Chromatogr.*, 646 (1993) 369-375.
- 1699 Zeng, Y. and Uden, P.C.: High temperature gas chromatography-atomic emission detection of metalloporphyrins in crude oils. *J. High Resolut. Chromatogr.*, 17 (1994) 223-229.
- 1700 Zeng, Y. and Uden, P.C.: Size exclusion chromatography sample pretreatment for GC-AED analysis of metalloporphyrins in crude oils. *J. High Resolut. Chromatogr.*, 17 (1994) 217-222.
- 23d. *Pyridine derivatives*
- 1701 Kajita, M., Niwa, T., Maruyama, W., Nakahara, D., Takeda, N., Yoshizumi, H., Tatematsu, A., Watanabe, K., Naoi, M. and Nagatsu, T.: Endogenous synthesis of N-methylnorsoanolinol in rat brain during *in vivo* microdialysis with epinine. *J. Chromatogr. B*, 654 (1994) 263-269.
- 1702 Kirchohoefer, R.D.: Gas chromatographic/mass spectrometric identification of a pyridinedicarboxylic acid impurity in niacin as the trimethylsilyl ester. *J. Assoc. Off. Anal. Chem.*, 77 (1994) 117-120.
- See also 1778, 1907.
- 23e. *Other N-heterocyclic compounds*
- 1703 Granda, M., Menendez, R., Moineo, S.R., Bermejo, J. and Snape, C.E.: Mass spectrometric characterization of polynuclear aromatic nitrogen compounds in coal tar pitches separated by extrography. *Fuel*, 72 (1993) 19-23; *C.A.*, 119 (1993) 184461f.
- 1704 Lickefett, H., Krohn, K., Koenig, W.A., Gehrcke, B. and Sylatk, C.: Enantioseparation of 5-monosubstituted hydantoin by capillary gas chromatography - investigation of chemical and enzymic racemization. *Tetrahedron: Asymmetry*, 4 (1993) 1129-1135; *C.A.*, 119 (1993) 249883s.
24. ORGANIC SULPHUR COMPOUNDS (INCL. GLUCOSINOLATES)
- 1705 Calvey, E.M., Roach, A.G. and Block, E.: Supercritical fluid chromatography of garlic (*Allium sativum*) extracts with mass spectrometric identification of allicin. *J. Chromatogr. Sci.*, 32 (1994) 93-96.
- 1706 Cooper, J.F., Cabanis, J.C., Blachere, M., Tourte, J. and Gros, P.: (Methyl isothiocyanate in wines. Study and analysis.) *Ann. Falsif. Expert. Chem. Toxicol.*, 86 (1993) 49-54; *C.A.*, 120 (1994) 75772h.
- 1707 Di Sanzo, F.P., Bray, W. and Chawla, B.: Determination of the sulfur components of gasoline streams by capillary column gas chromatography with chemiluminescence detection. *J. High Resolut. Chromatogr.*, 17 (1994) 255-258.

- 1708 Guo, H., Cui, L., Zhou, X. and Wang, L.: (GC/MS analysis of organic sulfide in volatile oil of several *Allium* plants.) *Fenxi Ceshi Tongbao*, 11, No. 5 (1992) 68-70; C.A., 119 (1993) 278878z.
- 1709 Kataoka, H., Imamura, Y., Tanaka, H. and Makita, M.: Determination of cysteamine and cystamine by gas chromatography with flame photometric detection. *J. Pharm. Biomed. Anal.*, 11 (1993) 963-969; C.A., 120 (1994) 26696c.
- 1710 McCormack, A.J., Sudhakar, C., Levine, S.A., Patel, M.S. and McCann, J.M.: Simultaneous determination of hydrogen sulfide sulfur, mercaptan sulfur, and total sulfur in hydrotreated naphtha by GC with atomic emission detection. *LC-GC*, 12 (1994) 30-35.
- 1711 Persson, C. and Leck, C.: Determination of reduced sulfur compounds in the atmosphere using a cotton scrubber for oxidant removal and gas chromatography with flame photometric detection. *Anal. Chem.*, 66 (1994) 983-987.
- 1712 Ruiz, R., Hartman, T.G., Karmas, K., Lech, J. and Rosen, R.T.: Breath analysis of garlic-borne phytochemicals in human subjects: combined adsorbent trapping and short-path thermal desorption gas chromatography-mass spectrometry. In: *Food Phytochemicals for Cancer Prevention I. ACS Symp. Ser.*, Vol. 546, ACS, Washington, 1994, pp. 102-119; C.A., 120 (1994) 128965g.
- 1713 Shearer, R.L. and Skelton, R.J.: Supercritical fluid chromatography of petroleum products using flameless sulfur chemiluminescence detection. *J. High Resolut. Chromatogr.*, 17 (1994) 251-254.
- 1714 Stenroos, L., Bates, M., Brennan, M., Burmeister, M., Davis, N., Hastings, D., Hutte, R., Nyarady, S., Organ, G., et al.: Dimethyl sulfide in beer by chemiluminescence detector. *J. Am. Soc. Brew. Chem.*, 51 (1993) 172; C.A., 120 (1994) 29628t.
- 1715 Sye, W.F. and Jou, W.S.: The analysis of odorous sulfur compounds in air by solid adsorbent preconcentration and PTV thermal desorption. *J. Chin. Chem. Soc. (Taipei)*, 40 (1993) 455-461; C.A., 120 (1994) 37044q.
- 1716 Yongzhi, L., Zhangping, C. and Xuanjie, W.: Allitridin analysis by FID/FPD dual channel gas chromatography. *J. Chromatogr. Sci.*, 32 (1994) 61-64.
- See also 1347, 1607, 1717, 1889, 1926.
25. ORGANIC PHOSPHORUS COMPOUNDS (INCL. SUGAR PHOSPHATES)
- 1717 Black, R.M., Clarke, R.J., Read, R.W. and Reid, M.T.J.: Application of gas chromatography-mass spectrometry and gas chromatography-tandem mass spectrometry to the analysis of chemical warfare samples, found to contain residue of the nerve agent sarin, sulphur mustard and their degradation products. *J. Chromatogr. A*, 662 (1994) 301-321.
- 1718 Kalski, A. and Smok, W.: (Determination of ethylbicyclophosphate in biological material by gas chromatography.) *Bromatol. Chem. Toksykol.*, 25 (1992) 69-71; C.A., 119 (1993) 243042v.
- 1719 Pang, J., Huang, L., Wu, G., Dai, D. and Zhu, G.: (Study on the reaction mechanisms of triphenylphosphine and triethoxyphosphine with octasulfur by MS and GC/MS.) *Youji Huaxue*, 13 (1993) 377-382; C.A., 119 (1993) 250056f.
- 1720 Sokolowski, M. and Witkiewicz, Z.: Gas chromatography-mass spectrometry analysis of products of *o*-isopropyl methylphosphonofluoridate transformation in aliphatic alcohols. *Chem. Anal. (Warsaw)*, 38 (1993) 139-147.
- See also 1263, 1681.
26. ORGANOMETALLIC AND RELATED COMPOUNDS
- 26a. Organometallic compounds
- 1721 Cai, Y., Alzaga, R. and Bayona, J.M.: *In situ* derivatization and supercritical fluid extraction for the simultaneous determination of butyltin and phenyltin compounds in sediment. *Anal. Chem.*, 66 (1994) 1161-1167.
- 1722 Filippelli, M. and Baldi, F.: Alkylation of ionic mercury to methylmercury and dimethylmercury by methylcobalamin: simultaneous determination by purge-and-trap GC in line with FTIR. *Appl. Organomet. Chem.*, 7 (1993) 487-492; C.A., 120 (1994) 107209y.
- 1723 Imagawa, T., Tao, H. and Miyazaki, A.: (Determination of lead isotope ratios in seawater by purge and trap GC/MS after ethylation.) *Shigen to Kankyo*, 2 (1993) 331-338; C.A., 119 (1993) 210042s.
- 1724 Stepanov, V.M., Krylov, V.A., Malyshev, K.K. and Chernova, O.Yu.: (Statistical method for determination of the completeness and accuracy of a data set on the impurity composition of high-purity substances as illustrated by chromatographic analysis of samples of diethyltellurium.) *Vysokochist. Veshchestva*, No. 3 (1993) 170-176; C.A., 119 (1993) 216515v.
- See also 1295, 1348, 1420.
- 26b. Boranes, silanes and related non-metallic compounds
- 1725 D'Ulivo, L., Lampugnani, L., Sfetsios, J., Zamboni, R. and Forte, C.: Studies on the breakdown of organoselenium compounds in a hydrobromic acid-bromine digestion system. *Analyst (Cambridge)*, 119 (1994) 633-640.
- 1726 Elaseer, A. and Nickless G.: Determination of selenium by gas chromatography-electron-capture detection using a rapid derivatization procedure. *J. Chromatogr. A*, 664 (1994) 77-87.
- 1727 Just, U., Mellor, F. and Keidel, F.: Supercritical fluid chromatography (SFC) of linear and cyclic components in technical silicone oils. *Fresenius J. Anal. Chem.*, 348 (1994) 745-748.
- See also 1959.
- 26c. Coordination compounds
- 1728 Carey, J.M., Vela, N.P. and Caruso, J.A.: Chromium determination by supercritical fluid chromatography with inductively coupled plasma mass spectrometric and flame ionization detection. *J. Chromatogr. A*, 662 (1994) 329-340.
- 1729 Rohr, U., Meckel, L. and Ortner, H.M.: Ultratrace analysis of uranium and thorium in glass. Part. 1. ICP-MS, classical photometry and chelate-GC. *Fresenius J. Anal. Chem.*, 348 (1994) 356-363.

- 1730 Veillon, C., Patterson, K.Y., Rubin, M.A. and Moser-Veillon, P.B.: Determination of natural and isotopically enriched chromium in urine by isotope dilution gas chromatography/mass spectrometry. *Anal. Chem.*, 66 (1994) 856-860.
27. VITAMINS AND VARIOUS ANIMAL GROWTH FACTORS (NON-PEPTIDIC)
- 1731 Komuro, S., Nakatsuka, I., Yoshitake, A. and Iba, K.: Quantitative determination of F₆-1,25(OH)₂ vitamin D₃ in human serum by gas chromatography/mass spectrometry with high-resolution selected ion monitoring. *Biol. Mass Spectrom.*, 23 (1994) 33-38; C.A., 120 (1994) 101180k.
- See also 1269, 1451, 1634.
28. ANTIBIOTICS
- 1732 Arimoto, H. and Adachi, A.: (Supercritical fluid chromatography for macrolide antibiotics determination.) *Jpn. Kokai Tokkyo Koho*, JP 05,322,871 [93,322,871] (Cl. G01N30/88), 7 Dec. 1993, Appl. 92/125,288, 19 May 1992; 4 pp.; C.A., 120 (1994) 158153h.
- 1733 Kijak, P.J.: Confirmation of chloramphenicol residues in bovine milk by gas chromatography/mass spectrometry. *J. Assoc. Off. Anal. Chem.*, 77 (1994) 34-40.
29. INSECTICIDES, PESTICIDES AND OTHER AGROCHEMICALS
- 29a. General techniques
- 1734 Alzaga, R., Durand, G., Barceló, D. and Bayona, J.M.: Comparison of supercritical fluid extraction and liquid-liquid extraction for isolation of selected pesticides stored in freeze-dried water samples. *Chromatographia*, 38 (1994) 502-508.
- 1735 Beltran, J., Lopez, F.J. and Hernandez, F.: Solid-phase extraction of pesticide residues from ground water: comparison between extraction cartridges and extraction disks. *Anal. Chim. Acta*, 283 (1993) 297-303.
- 1736 Hayakawa, S., Sarai, E. and Araki, K.: (Method for solid-phase extraction of pesticides from water. Comparison of the methods for the solid-phase and liquid-liquid extraction.) *Mie-ken Kankyo Kagaku Senta Kenkyu Hokoku*, 13 (1993) 73-77; C.A., 119 (1993) 233524q.
- 1737 Holland, P.T., McNaughton, D.E. and Malcolm, C.P.: Multi-residue analysis of pesticides in wines by solid-phase extraction. *J. Assoc. Off. Anal. Chem.*, 77 (1994) 79-86.
- 1738 Neidert, E., Trotman, R.B. and Sachsenbrecker, P.W.: Levels and incidences of pesticide residues in selected agricultural food commodities available in Canada. *J. Assoc. Off. Anal. Chem.*, 77 (1994) 18-24.
- 1739 Pavoni, G. and Sandirocco, C.: (Multiresidue method for the determination in honey of some pesticides used against *Varroa jacobsoni* and of organochlorine pesticides using a SPE membrane Empore extraction disk.) *Boll. Chim. Ig., Parte Sci.*, 44 (1993) 41-45; C.A., 120 (1994) 161865x.
- 1740 Sarai, E., Hayakawa, S. and Araki, K.: (Studies on the solid extraction method for the determination of pesticides in water. II.) *Mie-ken Kankyo Kagaku Senta Kenkyu Hokoku*, 13 (1993) 79-82; C.A., 119 (1993) 210036t.
- 1741 Suzuki, T., Yaguchi, K., Ohnishi, K. and Yamagashi, T.: Determination of pesticides in water by capillary gas chromatography with splitless injection of large sample volumes. *J. Chromatogr. A*, 662 (1994) 139-146.
- 1742 Tong, B. and Li, X.: (Determination of methylmercury and ethylmercury in environmental water by megabore capillary gas chromatography after preconcentration by sulfhydryl cotton.) *Yankuang Ceshi*, 12 (1993) 131-134; C.A., 119 (1993) 278224h.
- 29b. Chlorinated insecticides
- 1743 Aplada-Sarlis, P., Liapis, K.S. and Miliadis, G.E.: Contamination of potato tubers and carrots in Greece with lindane residues. *Bull. Environ. Contam. Toxicol.*, 52 (1994) 135-140; C.A., 120 (1994) 29677h.
- 1744 Barnabas, I.J., Dean, J.R., Hitchen, S.M. and Owen, S.P.: Supercritical fluid extraction of organochlorine pesticides from an aqueous matrix. *J. Chromatogr. A*, 665 (1994) 307-315.
- 1745 Cetinkaya, M.: (Gas chromatographic determination of pesticide residues in hair dyes.) *Parfuem. Kosmet.*, 74, No. 1 (1993) 23-25; C.A., 119 (1993) 233681p.
- 1746 Cruz, I., Wells, D.E. and Marr, I.L.: Determination of organochlorines in sea water: an assessment. *Anal. Chim. Acta*, 283 (1993) 280-286.
- 1747 Kajimura, K., Sakagami, Y., Yokoyama, H., Doi, S. and Yoshida, S.: (Analytical method for organochlorine pesticides residues in crude drugs. (1st Report).) *Osaka-furitsu Koshu Eisei Kenkyusho Kenkyu Hokoku, Yakuji Shido-hen*, 27 (1993) 21-24; C.A., 120 (1994) 173580p.
- 1748 Krechniak, J. and Loboda-Peplinska, T.: (Determination of permethrin in the blood by gas chromatography.) *Bromatol. Chem. Toksykol.*, 24 (1991) 323-324; C.A., 120 (1994) 2309c.
- 1749 Mössner, S. and Ballschmiter, K.: Separation of α -hexachlorocyclohexane (α -HCH) and pentachlorocyclohexene (PCCH) enantiomers on a cyclo-dextrin phase (Cyclodex-B) by HRGC/ECD. *Fresenius J. Anal. Chem.*, 348 (1994) 583-589.
- 1750 Onuska, F.I., Terry, K.A., Seech, A. and Antonic, M.: Determination of toxaphene in soil by electron-capture negative-ion mass spectrometry and capillary column gas chromatography. *J. Chromatogr. A*, 665 (1994) 125-132.
- 1751 Schenck, F.J., Wagner, R., Hennessy, M.K. and Okrasinski, J.L., Jr.: Screening of organochlorine pesticide and polychlorinated biphenyl residues in nonfatty seafood products by tandem solid-phase extraction cleanup. *J. Assoc. Off. Anal. Chem.*, 77 (1994) 102-106.
- 1752 Tilio, R., Kapila, S., Nam, K.S., Bossi, R. and Facchetti, S.: Reduction/elimination of sulfur interference in organochlorine residue determination by supercritical fluid extraction. *J. Chromatogr. A*, 662 (1994) 191-197.
- 1753 Turrio-Baldassarri, L., di Domenico, A., Fulgenzi, A., Iacovella, N., Bocca, A. and Larsen, B.R.: GC-MS determination of HCB, DDT, DDE, and selected PCB congeners in cow's milk. *Fresenius Environ. Bull.*, 2 (1993) 370-374; C.A., 120 (1994) 132521b.

- 1754 Wan, H.B., Wong, M.K., Lim, P.Y. and Mok, C.Y.: Small-scale multi-residue method for the determination of organochlorine and pyrethroid pesticides in vegetables. *J. Chromatogr. A*, 662 (1994) 147-152.
- 1755 Zhou, Y. and Zhao, R.: (Determination of organochlorine pesticides in ginseng (*Panax ginseng*.) *Zhongcaoyao*, 24 (1993) 304-305; *C.A.*, 119 (1993) 210864m.

See also 1757.

29c. Phosphorus insecticides

- 1756 Kawata, K. and Yakusara, A.: Determination of fenitrothion and diazinon in air. *Bull. Environ. Contam. Toxicol.*, 52 (1994) 419-424; *C.A.*, 120 (1994) 115241t.
- 1757 Milliadis, G.E.: Determination of pesticide residues in natural waters of Greece by solid phase extraction and gas chromatography. *Bull. Environ. Contam. Toxicol.*, 52 (1994) 25-30; *C.A.*, 120 (1994) 14531g.
- 1758 Nakamura, Y., Sekiguchi, Y., Tsumura, Y., Tonogai, Y. and Ito, Y.: (Studies on metabolites of pesticides. II. Simultaneous determination of naled and its reduced product dichlorvos by gas chromatography.) *Shokuhin Eiseigaku Zasshi*, 34 (1993) 426-433; *C.A.*, 120 (1994) 132497y.
- 1759 Nutley, B.P. and Cocker, J.: Biological monitoring of workers occupationally exposed to organophosphorus pesticides. *Pestic. Sci.*, 38 (1993) 315-322; *C.A.*, 119 (1993) 277822q.
- 1760 Parfitt, C.H.: Wide-bore capillary gas chromatographic determination of organophosphorus pesticide residues in foods: interlaboratory trial. *J. Assoc. Off. Anal. Chem.*, 77 (1994) 92-101.
- 1761 Psathaki, M., Manoussaridou, E. and Stephanou, E.G.: Determination of organophosphorus and triazine pesticides in ground- and drinking water by solid-phase extraction and gas chromatography with nitrogen-phosphorus or mass spectrometric detection. *J. Chromatogr. A*, 667 (1994) 241-248.
- 1762 Tsumura, Y., Sekiguchi, Y., Nakamura, Y., Tonogai, Y. and Ito, Y.: (Studies on metabolites of pesticides. I. Simultaneous determination of etrimfos and its metabolite by gas chromatography with flame thermionic detection.) *Shokuhin Eiseigaku Zasshi*, 34 (1993) 415-419; *C.A.*, 120 (1994) 105201r.
- 1763 Wang, W.-K. and Huang, S.-D.: Determination of organophosphorus pesticides in water using C₁₈ or Florisil Sep-Pak Cartridge and gas chromatography with flame-photometric detection. *J. Chin. Chem. Soc.*, 41 (1994) 109-113.
- 1764 Zegers, B.N., Hogenboom, A.C., Dekkers, S.E.G., Lingeman, H. and Brinkman, U.A.T.: Packed capillary supercritical fluid chromatography of organophosphorus pesticides: Selective detection and applications. *J. Microcolumn Sep.*, 6 (1994) 55-62.

See also 1745.

29d. Carbamates

- 1765 Albelda, C.A., Picó, Y. and Mañes, J.: Determination of aldicarb, aldicarb sulfoxide, and aldicarb sulfone in oranges by simple gas-liquid chromatography with nitrogen-phosphorus detection. *J. Assoc. Off. Anal. Chem.*, 77 (1994) 74-78.

- 1766 Dyer, R.H.: Determination of ethyl carbamate (urethane) in alcoholic beverages using capillary gas chromatography with thermal energy analyzer detection: collaborative study. *J. Assoc. Off. Anal. Chem.*, 77 (1994) 64-67.
- 1767 Fauhl, C., Catsburg, R. and Wittkowski, R.: Determination of ethyl carbamate in soy sauces. *Food Chem.*, 48 (1993) 313-316; *C.A.*, 119 (1993) 224578h.
- 1768 Howald, W.N. and Porubek, D.J.: Quantification of 3-(2-hydroxypropyl)-5-methyl-2-oxazolidone in groundwater and sludge by isotope dilution gas chromatography/mass spectrometry. *Environ. Sci. Technol.*, 28 (1994) 682-685.
- 1769 Sun, S. and Ma, Y.: Detection and quantitation of trace ethyl carbamate in alcoholic beverages by GC/GC and GC/GC/MS. *Chin. Chem. Lett.*, 4 (1993) 611-614; *C.A.*, 119 (1993) 224526q.
- 1770 Ye, M.: (Determination of pirimicarb by gas chromatography.) *Nongyao Kexue Yu Guanli*, No. 1 (1993) 26-28; *C.A.*, 119 (1993) 264565h.

See also 1866.

29e. Herbicides

- 1771 Dieckmann, H., Kreuzig, R. and Bahadir, M.: Significance of keto-enol tautomerism in the analysis of 1,2,4-triazinone metabolites. *Fresenius J. Anal. Chem.*, 348 (1994) 749-753.
- 1772 Hong, S., Lee, A.S. and Smith, A.E.: Gas chromatographic analysis of dithiopyr in soil leachate. *J. Chromatogr. Sci.*, 32 (1994) 83-85.
- 1773 Klaffenbach, P. and Holland, P.T.: Analysis of sulfonylurea herbicides by gas-liquid chromatography-mass spectrometry and multiresidue determination. *Biol. Mass Spectrom.*, 22 (1993) 565-578; *C.A.*, 120 (1994) 99227k.
- 1774 Liu, C., Liu, W. and Sheng, L.: (Assay of matrine in lightyellow sophora (*Sophora flavescens*) and its pharmaceutical formulations by capillary gas chromatography.) *Zhongcaoyao*, 24 (1993) 405-406; *C.A.*, 120 (1994) 14979j.
- 1775 Mortimer, R.D. and Weber, D.F.: Some comments on a recently proposed method of determining chlormequat residues by derivatization with pentafluorothiophenol. *Pestic. Sci.*, 40 (1994) 31-35; *C.A.*, 120 (1994) 127601y.
- 1776 Papilloud, S. and Haerdi, W.: Supercritical fluid extraction of triazine herbicides from solid matrices. *Chromatographia*, 38 (1994) 514-519.
- 1777 Steinheimer, T.R., Pfeiffer, R.L. and Scoggin, K.D.: Extraction of atrazine, cyanazine, desethylatrazine, desisopropylatrazine, and metolachlor from fortified western cornbelt soils by SFE with CO₂. *Anal. Chem.*, 66 (1994) 645-650.

See also 1761.

29f. Fungicides

- 1778 Cioni, F. and Bartolucci, G.: (Methods of analysis for natural products with insecticidal activity.) *Boll. Chim. Ig., Parte Sci.*, 44 (1993) 293-303; *C.A.*, 120 (1994) 161858x.
- 1779 Dethle, M.D., Kale, V.D. and Rane, S.D.: Gas-chromatographic studies on the residues of fenvalerate in chickpea. *J. Maharashtra Agric. Univ.*, 18 (1993) 149-150; *C.A.*, 120 (1994) 105280r.

- 1780 Kim, H., Lapiguera, A. and Lin, C.: Gas chromatographic and high-performance liquid chromatographic methods for the determination of genaconazole in biological fluids. *J. Chromatogr. B*, 655 (1994) 21-26.
- 1781 Moulder, R., Bartle, K.D. and Clifford, A.A.: Supercritical fluid chromatography with electron-capture detection for the determination of agrochemicals. *Analyst (Cambridge)*, 118 (1993) 737-739.
- 1782 Yuldashev, Z.A., Zainutdinov, K.S., Popkov, V.A., Ikramov, L.T., Reshetnyak, V.Yu. and Dustmukhamedov, T.T.: (Toluin measurements in biologic fluids.) *Sud.-Med. Ekspert.*, 35, No. 4 (1992) 31-33; *C.A.*, 119 (1993) 197370a.
- 29g. *Other types of pesticides and various agrochemicals*
- 1783 Natangelo, M., Benfenati, E., de Gregorio, G., Fanelli, R. and Ciotti, G.: GC-MS analysis of N-phosphonomethylglycine (glyphosate) samples through derivatization with a perfluoropropionic anhydride and trifluoroethanol: Identification of byproducts. *Toxicol. Environ. Chem.*, 38 (1993) 225-232; *C.A.*, 120 (1994) 25500x.
- 1784 Nie, X., Nie, R., Li, Z. and Qiu, M.: (The quantitative analysis of pyrethroids.) *Yunnan Zhiwu Yanjiu*, 15 (1993) 317-319; *C.A.*, 120 (1994) 25502z.
- 1785 Pang, G.-F., Fan, C.-L., Chao, Y.-Z. and Zhao, T.-S.: Rapid method for the determination of multiple pyrethroid residues in fruits and vegetables by capillary column gas chromatography. *J. Chromatogr. A*, 667 (1994) 348-353.
- See also 1754.
31. PLASTICS AND THEIR INTERMEDIATES
- 1786 Babenko, E.V., Zenitova, L.A., Safiullina, F.F. and Mukharlyamov, S.F.: (Optimization of the formulation of casting polyurethanes by inverse gas chromatographic method.) *Izv. Vyssh. Uchebn. Zaved., Khim. Khim. Tekhnol.*, 36, No. 6 (1993) 105-108; *C.A.*, 120 (1994) 165509u.
- 1787 Baltus, R.E., Alger, M.M. and Stanley, T.J.: Solubility and diffusivity of cyclic oligomers in poly(dimethylsiloxane) using capillary column inverse gas chromatography. *Macromolecules*, 26 (1993) 5651-5656; *C.A.*, 119 (1993) 181795g.
- 1788 Boinon, B., Raihane, M. and Montheard, J.P.: Kinetics of thermal degradation of an alternating copolymer: poly(methylvinylidene cyanide-co-vinyl acetate). Thermogravimetric and gas chromatography-mass spectroscopy studies. *Polym. Degrad. Stab.*, 43 (1994) 27-31; *C.A.*, 120 (1994) 165172s.
- 1789 Bosse, F., Eisenberg, A., Xu, R. and Schreiber, H.P.: IGC study of surface properties of adsorbed styrene/methacrylic acid diblock polymers. *J. Appl. Polym. Sci.*, 51 (1994) 521-527; *C.A.*, 120 (1994) 108230k.
- 1790 De Juana, R., Etxeberria, A., Cortazar, M. and Irui, J.J.: A study of mixtures of poly(hydroxy ether of bisphenol A) and poly(ϵ -caprolactone) by inverse gas chromatography. *Macromolecules*, 27 (1994) 1395-1400; *C.A.*, 120 (1994) 165741b.
- 1791 Etxeberria, A., Uriarte, C., Fernandez-Berridi, M.J. and Irui, J.J.: Probing polymer-polymer interaction parameters in miscible blends by inverse gas chromatography: solvent effects. *Macromolecules*, 27 (1994) 1245-1248; *C.A.*, 120 (1994) 135738p.
- 1792 Garner, C.M., Ma, Y. and Bartus, J.: Capillary gas chromatography analysis of low-molecular-weight oligomers. *Polym. Prepr.*, 33 (1992) 1010-1011; *C.A.*, 120 (1994) 44789q.
- 1793 Grenier-Loustalot, M.-F., Renotte-Greco, D. and Grenier, P.: (Study of the surface properties by inverse gas chromatography of grafted silica characterized by carbon-13 NMR and DSC.) *Eur. Polym. J.*, 29 (1993) 1185-1196; *C.A.*, 119 (1993) 182021p.
- 1794 Haken, K. and Iddamalgoda, P.: Analysis of liquid-crystal polyesters by gas chromatography and high-performance liquid chromatography after hydrolytic fusion. *J. Appl. Polym. Sci.*, 52 (1993) 225-234; *C.A.*, 120 (1994) 9333q.
- 1795 Hida, M.: Analytical study of pyrolysis gas chromatography using matrix modifiers. *Bunseki Kagaku*, 43 (1994) 257-258.
- 1796 Hutchinson, E.J., Bowley, B., Pan, W.-P. and Nguyen, C.: Calibration for the composition of polymeric composites using TG/GC. *Thermochem. Acta*, 223 (1993) 259-269; *C.A.*, 119 (1993) 227429w.
- 1797 Ito, Y., Ohtani, H., Ueda, S., Nakashima, Y. and Tsuge, S.: Quantification of end groups in polystyrenes by pyrolysis-gas chromatography. *J. Polym. Sci., Part A: Polym. Chem.*, 32 (1994) 383-388; *C.A.*, 120 (1994) 78129q.
- 1798 Jiang, Z., Luo, Y., Jin, X. and Lu, F.: Studies on the thermal degradation of poly(phenyl-as-triazines) by high resolution pyrolysis-gas chromatography. *J. Anal. Appl. Pyrolysis*, 26 (1993) 145-157.
- 1799 Just, U., Holzbauer, H.-R. and Resch, M.: Molar mass determination of oligomeric ethylene oxide adducts using supercritical fluid chromatography and matrix-assisted laser desorption-ionization time-of-flight mass spectrometry. *J. Chromatogr. A*, 667 (1994) 354-360.
- 1800 Kontominas, M.G. and Giacini, J.R.: Interaction of food grade polystyrene with styrene monomer using inverse gas chromatography: migration aspects. *Dev. Food Sci.*, 32 (1993) 769-779; *C.A.*, 119 (1993) 202025y.
- 1801 Kontominas, M.G., Gavara, R. and Giacini, J.R.: Gas-solid chromatographic study of polymer surfaces: adsorption on polystyrene. *Eur. Polym. J.*, 30 (1994) 271-277; *C.A.*, 120 (1994) 135705a.
- 1802 McGuire, J.M. and Suchanec, R.R.: Analysis of formaldehyde-treated rosin by high-resolution gas chromatography-mass spectrometry. *J. Chromatogr. Sci.*, 32 (1994) 139-143.
- 1803 Pasch, H., Much, H. and Schulz, G.: Characterization of functional homopolymers and block copolymers by chromatographic methods. *J. Appl. Polym. Sci.*, 52 (1993) 79-90. *C.A.*, 120 (1994) 9326q.
- 1804 Pattueli, M.E., Montalti, A., Viola, G.T. and Zazzetta, A.: Sequence distribution of styrene-butadiene copolymers by ozonolysis, high-performance liquid chromatographic and gas chromatographic-mass spectrometric techniques. *J. Chromatogr. A*, 665 (1994) 117-123.
- 1805 Tan, Z., Jaeger, R. and Vancso, G.J.: Studies of crosslinking of poly(dimethylsiloxane) networks by inverse gas chromatography. *Polym. Prepr.*, 34 (1993) 306-307; *C.A.*, 120 (1994) 136020k.

- 1806 Tóth, T. and Garay, F.: Separation of novolac resin oligomers and related industrial materials by high temperature capillary gas chromatography. *J. High Resolut. Chromatogr.*, 17 (1994) 177-179.
- 1807 Voelkel, A., Andrzejewska, E., Maga, R. and Andrzejewski, M.: Characterization of dispersive and acid-base properties of cross-linked polymers by inverse gas chromatography. *Polymer*, 34 (1993) 3109-3111; *C.A.*, 119 (1993) 181768a.
- 1808 Xu, Z. and Yang, J.: (Pyrolysis-gas chromatographic study of cyclized random copolymer of *cis*-1,4-butadiene and isoprene.) *Yingyong Huaxue*, 10, No. 4 (1993) 101-103; *C.A.*, 119 (1993) 181470x.
- 1809 Yan, W., Shi, E., Zhang, B. and Xu, F.: (Pyrolysis gas chromatographic analysis of polyblend formed in copolymer and homopolymer of styrene, methyl methacrylate and acrylonitrile.) *Sepu*, 11 (1993) 324-325.
- 1810 Zaiкин, V.G., Mardanov, R.G., Kleiner, V.I., Krentsel, B.A. and Bobrov, B.N.: Differentiation of block, random and block-random ethylene/vinylcyclohexane copolymers by pyrolysis-gas chromatography/mass spectrometry. *J. Anal. Appl. Pyrolysis*, 26 (1993) 185-190.
- See also 1265, 1266, 1270, 1271, 1272, 1277, 1282, 1293, 1302, 1307, 1949.
32. DRUG ANALYSIS
- 32a. *Drug analysis, general techniques*
- 1811 Jagota, N.K.: Supercritical fluid chromatography of selected pharmaceuticals. Avail. *Univ. Microfilms Int.*, Order No. DA9301200, 1992, 189 pp.; *C.A.*, 120 (1994) 86605g.
- 1812 Koenig, W.A.: The direct resolution of enantiomeric drugs by chiral-phase gas chromatography. *Clin. Pharmacol.*, 18 (1993) 107-137; *C.A.*, 119 (1993) 188686f.
- 1813 Zhuravleva, I.L., Terenina, M.B., Golovnya, R.V. and Filimonova, M.A.: (Analysis of aqueous solutions of nitrogen-containing drugs by capillary gas chromatography.) *Khim.-Farm. Zh.*, 27, No. 5 (1993) 58-63; *C.A.*, 119 (1993) 278929s.
- See also 1747.
- 32b. *Antirheumatics and antiinflammatory drugs*
- 1814 Bushby, B., Fernandes, A., Wallace, D. and Kibblewhite, M.: Determination of trace organic micropollutants in atmospheric deposition. *Sci. Total Environ.*, 135 (1993) 81-94; *C.A.*, 120 (1994) 22851b.
- See also 1461.
- 32c. *Autonomic and cardiovascular drugs*
- 1815 Blandford, D.E. and Desjardins, P.R.E.: Detection and identification of amphetamine and methamphetamine in urine by GC/MS. *Clin. Chem. (Washington)*, 40 (1994) 145-147.
- 1816 Gao, H., Sun, W. and Sha, Z.: (Quantitative determination of nifedipine and related impurities by GC.) *Yaowu Fenxi Zazhi*, 13 (1993) 249-252; *C.A.*, 119 (1993) 256620g.
- 1817 Ji, S., Kong, Q., Li, X. and Li, P.: Gas chromatographic determination of mexiletine in human plasma with flame ionization detection after reaction with carbon disulfide. *Biomed. Chromatogr.*, 7 (1993) 196-199.
- 1818 Jones, J.B. and Mell, L.D., Jr.: A simple wash procedure for improving chromatography of HFAA derivatized amphetamine extracts for GC/MS analysis. *J. Anal. Toxicol.*, 17 (1993) 447; *C.A.*, 120 (1994) 123959y.
- 1819 Kumazawa, T., Sato, K., Seno, H. and Suzuki, O.: (Positive- and negative-ion mass spectrometry and rapid extraction with Sep-Pak C₁₈ cartridges for dihydropyridine calcium antagonists and their photodecomposition products.) *Hochudoku*, 11 (1993) 128-129; *C.A.*, 119 (1993) 173474m.
- 1820 Marciniak, B., Kujawa, E. and Ogródowczyk, M.: Evaluation of nifedipine preparations by chromatographic-spectrophotometric methods. *Pharmazie*, 47 (1992) 502-504; *C.A.*, 119 (1993) 278940p.
- 1821 Melgar, R. and Kelly, R.C.: A novel GC/MS derivatization method for amphetamines. *J. Anal. Toxicol.*, 17 (1993) 399-402; *C.A.*, 120 (1994) 98617a.
- 1822 Micheel, A.P., Ko, C.Y. and Evans, C.R.: Assay of residual organic solvents in topiramate drug substance by capillary gas chromatography. *J. Pharm. Biomed. Anal.*, 11 (1993) 1233-1238; *C.A.*, 120 (1994) 116925f.
- 1823 Ohue, O., Sawaguchi, T., Nakamura, S., Sawaguchi, A., Shinozuka, T., Kurihara, K. and Yanagida, J.: (The significance of detection of methamphetamine from formalin fixed by GC/MS.) *Nippon Iyo Masu Supekutoru Gakkai Koenshu*, 18 (1993) 87-90; *C.A.*, 120 (1994) 71007q.
- 1824 Poletini, A., Groppi, A., Ricossa, M.C. and Montagna, M.: Gas chromatographic/electron impact mass spectrometric selective confirmatory analysis of clenbuterol in human and bovine urine. *Biol. Mass Spectrom.*, 22 (1993) 457-461; *C.A.*, 119 (1993) 197377h.
- 1825 Wang, X., Jia, C. and Wan, H.: Two-dimensional gas chromatography for the enantiomeric separation of ephedrine and phenaminum and their metabolites. *J. Chromatogr. B*, 653 (1994) 98-101.
- 1826 Yamahata, T., Dote, S., Ozawa, Y., Nishikawa, H. and Maeda, S.: Determination of clonidine in human plasma by gas chromatography-electron-impact mass spectrometry. *J. Chromatogr. B*, 653 (1994) 92-97.
- 1827 Yang, L.L. and Tu, X.D.: (Determination of pseudoephedrine in human plasma by gas chromatography with a mass selective detector (GC-MSD).) *Yaowu Xuebao*, 28 (1993) 709-713; *C.A.*, 120 (1994) 68711j.
- 1828 Yoo, Y., Chung, H. and Choi, H.: Determination of methamphetamine and its metabolite amphetamine in biological fluids from 11 fatal cases. *Arch. Pharmacol. Res.*, 16 (1993) 175-179; *C.A.*, 120 (1994) 156051z.
- 1829 Zimmer, D. and Muschalek, V.: Enantioselective assay for the determination of nisoldipine in dog, rat and mouse plasma by chiral microbore high-performance liquid chromatography combined with gas chromatography-mass spectrometry. *J. Chromatogr. A*, 666 (1994) 241-248.

32d. Central nervous system drugs

- 1830 Berger, T.A. and Wilson, W.H.: Separation of drugs by packed column supercritical fluid chromatography. 1. Phenothiazine antipsychotics. *J. Pharm. Sci.*, 83 (1994) 281-286; C.A., 120 (1994) 116922c.
- 1831 Berger, T.A. and Wilson, W.H.: Separation of drugs by packed column supercritical fluid chromatography. 2. Antidepressants. *J. Pharm. Sci.*, 83 (1994) 287-290; C.A., 120 (1994) 116917e.
- 1832 Cui, J.F., Li, L., Cui, K.R., Zhou, Y., Li, N., Wang, M.Z. and Zhou, T.H.: (Separation and identification of stimulants and their metabolites.) *Yaoxue Xuebao*, 28 (1993) 455-463; C.A., 119 (1993) 175494s.
- 1833 Delbeke, F.T., Teale, P., Debackere, M. and Houghton, E.: ELISA screening with GC-MS confirmation of the tranquilizer chlorprothixene administered in subtherapeutic doses to horses. *J. Pharm. Biomed. Anal.*, 11 (1993) 569-575; C.A., 119 (1993) 175500r.
- 1834 Fitzgerald, R.L., Rexin, D.A. and Herold, D.A.: Benzodiazepine analysis by negative chemical ionization gas chromatography/mass spectrometry. *J. Anal. Toxicol.*, 17 (1993) 342-347; C.A., 119 (1993) 264271j.
- 1835 Gerlitz, J.: GC/MS quantitation of benzoylecgonine following liquid-liquid extraction. *J. Forensic Sci.*, 38 (1993) 1210-1214; C.A., 120 (1994) 98603t.
- 1836 Goromaru, T., Maeda, H., Matsuki, Y. and Kiwada, H.: (Isotopic fractionation of aminopyrine from its deuterated analogs and application for evaluation of hepatic function by capillary gas chromatography.) *Radioisotopes*, 42 (1993) 511-516; C.A., 120 (1994) 30f.
- 1837 Lombardero, N., Casanova, O., Behnke, M., Eyler, F.D. and Bertholf, R.L.: Measurement of cocaine and metabolites in urine, meconium, and diapers by gas chromatography/mass spectrometry. *Ann. Clin. Lab. Sci.*, 23 (1993) 385-394; C.A., 120 (1994) 94606y.
- 1838 Moore, C., Long, G. and Marr, M.: Confirmation of benzodiazepines in urine as trimethylsilyl derivatives using gas chromatography-mass spectrometry. *J. Chromatogr. B*, 655 (1994) 132-137.
- 1839 Rogowsky, D., Marr, M., Long, G. and Moore, C.: Determination of sertraline and desmethylsertraline in human serum using copolymeric bonded-phase extraction, liquid chromatography and gas chromatography-mass spectrometry. *J. Chromatogr. B*, 655 (1994) 138-141.
- 1840 Staerk, D.U., Shitangkoon, A. and Vigh, G.: Gas chromatographic separation of the enantiomers of volatile fluoroether anesthetics by derivatized cyclodextrins. II. Preparative-scale separations for isoflurane. *J. Chromatogr. A*, 663 (1994) 79-85.
- 1841 Su, P., Coutts, R.T., Baker, G.B. and Daneshmand, M.: Analysis of imipramine and three metabolites produced by isoenzyme CYP2D6 expressed in a human cell line. *Xenobiotica*, 23 (1993) 1289-1298; C.A., 120 (1994) 94628g.
- 1842 Tonn, G.R., Mutlib, A., Abbott, F.S., Rurak, D.W. and Axelson, J.E.: Simultaneous analysis of diphenhydramine and a stable isotope analog ($^2\text{H}_{10}$)diphenhydramine using capillary gas chromatography with mass selective detection in biological fluids from chronically instrumented pregnant ewes. *Biol. Mass Spectrom.*, 22 (1993) 633-642; C.A., 120 (1994) 22985y.
- 1843 Zavitsanos, P.A., Siddiqui, W. and Curtis, M.: A high resolution gas chromatographic/mass spectrometric method for the determination of N-0923 in human and monkey plasma. *Rapid Commun. Mass Spectrom.*, 7 (1993) 1145-1149; C.A., 120 (1994) 68759f.

See also 1437, 1523, 1861, 1863, 1864.

32e. Chemotherapeutics (exc. cytostatics and antibiotics)

- 1844 Patterson, T.A.: Part I. The quantitation of arecoline in tissue and prolongation of its duration of action. Part II. Prevention of physostygmine-, DFP-, and diazinon-induced acute toxicity of monoethylcholine and N-aminodecanol. Avail. *Univ. Microfilms Int.*, Order No. DA9239080, 1992, 173 pp.; C.A., 120 (1994) 124711y.
- 1845 Peytavin, G., Gimenez, F., Genissel, B., Gillotin, C., Baillet, A., Wainer, I.W. and Farinotti, R.: Chiral resolution of some antimalarial agents by sub- and supercritical fluid chromatography on an (S)-naphthylurea stationary phase. *Chirality*, 5 (1993) 173-180.
- 1846 Rakestraw, D.: Determination of amantadine in human plasma by capillary gas chromatography using electron-capture detection following derivatization with pentafluorobenzoyl chloride. *J. Pharm. Biomed. Anal.*, 11 (1993) 699-703; C.A., 119 (1993) 240841u.
- 1847 Wynn, N.R., Cooper, J.F., Zheng, S.Q. and Cabanis, J.C.: (Determination of procymidone residues in wine: immunoassays and mass detection-gas chromatography.) *Bull. O.I.V.*, 66 (1993) 551-557; C.A., 120 (1994) 75749f.
- 1848 Zhang, Y., Xu, G., Jin, R., Xu, L. and Li, Y.: (Determination of alantolactone and isoalantolactone in *Tumuxiang* by GC.) *Zhongguo Yaoke Daxue Xuebao*, 24 (1993) 248-250; C.A., 119 (1993) 210879v.

See also 1414.

32f. Cytostatics

- 1849 Embree, L., Burns, R.B., Heggie, J.R., Phillips, G.L., Reece, D.E., Spinelli, J.J., Hartley, D.O., Hudon, N.J. and Goldie, J.H.: Gas-chromatographic analysis of busulfan for therapeutic drug monitoring. *Cancer Chemother. Pharmacol.*, 32 (1993) 137-142; C.A., 119 (1993) 195025t.
- 1850 Guan, S., Wang, D., Zhang, J. and Gu, X.: (GC analysis of polyphase liposome-entrapped cyclophosphamide in blood.) *Shenyang Yaoxueyuan Xuebao*, 10 (1993) 123-125; C.A., 120 (1994) 68716q.
- 1851 Ragno, G., Veronico, M. and Vetusch, C.: Gas chromatographic and UV derivative determination of nitrendipine and its photodegradation product. *Int. J. Pharm.*, 99 (1993) 351-355; C.A., 120 (1994) 15017f.

32g. Other drug categories

- 1852 Abadi, M. and Solomon, H.M.: Detection of tolmetin and oxaprozin by a GC/MS method for barbiturates. *J. Anal. Toxicol.*, 18 (1994) 62; C.A., 120 (1994) 127087k.

- 1853 Dulery, B.D., le Henanf, N. and Haegele, K.D.: Determination of an acetylcholinesterase inhibitor, MDL 73745, in dog plasma and urine by combined gas chromatography/negative ion chemical ionization mass spectrometry. *Biol. Mass Spectrom.*, 22 (1993) 377-382; *C.A.*, 119 (1993) 195022q.
- 1854 Terada, N., Inoue, F., Kodo, N., Morioka, H., Kinugasa, A., Sawada, T. and Ogawa, K.: (Acylcarnithine analysis by gas chromatography/mass spectrometry (GCMS).) *Nippon Iyo Masu Supekutoru Gakkai Koenshu*, 18 (1993) 127-128; *C.A.*, 120 (1994) 72674s.
- See also 1455, 1536.
- 32h. *Toxicological and forensic applications*
- 1855 Abusada, G.M., Abukhalaf, I.K., Alford, D.D., Vinzon-Bautista, I., Pramanik, A.K., Ansari, N.A., Manno, J.E. and Manno, B.R.: Solid-phase extraction and GC/MS quantitation of cocaine, ecgonine methyl ester, benzoylecgonine, and cocaethylene from meconium, whole blood, and plasma. *J. Anal. Toxicol.*, 17 (1993) 353-358; *C.A.*, 119 (1993) 243071d.
- 1856 Anderson, R.E. and Nixon, G.L.: Isolation of benzoylecgonine from urine using solid-phase extraction. *J. Anal. Toxicol.*, 17 (1993) 432-433; *C.A.*, 120 (1994) 68738y.
- 1857 Buratti, M., Pellegrino, O. and Colombi, C.V.: (Urinary nitrous oxide, alothane and isofluorane gas chromatographic determination for the biological monitoring of occupational exposure to inhalation anesthetics.) *Med. Lav.*, 84, No. 1 (1993) 66-73; *C.A.*, 119 (1993) 175465h.
- 1858 Guan, F., Liu, Y., Luo, Y., Hu, X., Liu, F., Li, Q. and Kang, Z.: GC/MS identification of tetramine in samples from human alimentary intoxication and evaluation of artificial carbonic kidneys for the treatment of the victims. *J. Anal. Toxicol.*, 17 (1993) 199-201; *C.A.*, 119 (1993) 197372c.
- 1859 Hornbeck, C.L., Carrig, J.E. and Czarny, R.J.: Detection of a GC/MS artifact peak as methamphetamine. *J. Anal. Toxicol.*, 17 (1993) 257-263; *C.A.*, 119 (1993) 264264j.
- 1860 Huo, X., Liu, F., He, W., Luo, Y. and Kang, Z.: (GC determination of morphine in urine of opiate addicts.) *Yaowu Fenxi Zazhi*, 13 (1993) 363-365; *C.A.*, 120 (1994) 98596t.
- 1861 Kalasinsky, K.S., Levine, B., Smith, M.L., Maglulio, J., J. and Schaefer, T.: Detection of amphetamine and methamphetamine in urine by gas chromatography/Fourier transform infrared (GC/FTIR) spectroscopy. *J. Anal. Toxicol.*, 17 (1993) 359-364; *C.A.*, 119 (1993) 243072e.
- 1862 Kostrzewski, P.: (Micromethods for determining organic solvents in blood by headspace analysis.) *Med. Pr.*, 44 (1993) 131-139; *C.A.*, 120 (1994) 2335h.
- 1863 Ohshita, T.: (Detection of diacetylmorphine in human urine by capillary gas-chromatograph/mass spectrometer (GC/MS).) *Jpn. J. Toxicol. Environ. Health*, 39 (1993) 431-436; *C.A.*, 120 (1994) 25115a.
- 1864 Welch, M.J., Sniegoski, L.T., Allgood, C.C. and Habram, M.: Hair analysis for drugs of abuse: evaluation of analytical methods, environmental issues, and development of reference materials. *J. Anal. Toxicol.*, 17 (1993) 389-398; *C.A.*, 120 (1994) 25125d.
- 1865 Wu, A.H.B., Liu, N., Cho, Y.J., Johnson, K.G. and Wong, S.S.: Extraction and simultaneous elution and derivatization of 11-nor-9-carboxy- Δ^9 -tetrahydrocannabinol using Toxi-Lab SPEC prior to GC/MS analysis of urine. *J. Anal. Toxicol.*, 17 (1993) 215-217; *C.A.*, 119 (1993) 175498w.
- 1866 Zhang, Y., Li, H., Yan, X., Ye, J. and Tang, X.: (Rapid qualitative and quantitative analysis of carbamate pesticides poisoning. II. GC method.) *Yaowu Fenxi Zazhi*, 13 (1993) 377-380; *C.A.*, 120 (1994) 98597u.
- See also 1587, 1686, 1748, 1759, 1828, 1955.
- 32i. *Plant extracts*
- 1867 Ahmad, U.K., Sirat, H.M., Sanagi, M.M. and Smith, R.M.: Supercritical fluid extraction and capillary gas chromatography of the rhizomes of *Z. zerumbet*. *J. Microcolumn Sep.*, 6 (1994) 27-32.
- 1868 Garcia-Viguera, C., Ferreres, F. and Tomas-Barberan, F.A.: Study of Canadian propolis by GC-MS and HPLC. *Z. Naturforsch., C: Biosci.*, 48 (1993) 731-735; *C.A.*, 120 (1994) 4006a.
- 1869 Ma, X.Z., Yu, X.B., Chen, J.J., Liang, Z.M., Cao, T.C. and Wang, Y.: (Determination of major volatile components in Suhexiangwan (storax pill) using analytical supercritical fluid extraction (SFE).) *Yaowu Xuebao*, 28 (1993) 537-542; *C.A.*, 119 (1993) 278879a.
- 1870 Yoshikawa, M., Hatakeyama, S., Chatani, N., Nishino, Y. and Yamahara, J.: (Qualitative and quantitative analysis of bioactive principles in *Zingiberis rhizoma* by means of high performance liquid chromatography and gas liquid chromatography. On the evaluation of *Zingiberis rhizoma* and chemical change of constituents during *Zingiberis rhizoma* processing.) *Yakugaku Zasshi*, 113 (1993) 307-315; *C.A.*, 119 (1993) 188694g.
- See also 1554, 1663, 1774.
33. CLINICO-CHEMICAL APPLICATIONS
- 33b. *Complex mixtures and profiling (single compounds by cross-reference only)*
- 1871 Andrew, R., Best, S.A., Watson, D.G., Midgley, J.M., Reid, J.L. and Squire, I.B.: Analysis of biogenic amines in plasma of hypertensive patients and a control group. *Neurochem. Res.*, 18 (1993) 1179-1182; *C.A.*, 120 (1994) 26945h.
- 1872 Aoki, K., Aramaki, S., Yoshida, I., Inokuchi, T., Tanaka, M., Soda, H., Tokunaga, Y., Yoshino, M., Kato, H. and Ishimatsu, J.: (Problems of prenatal diagnosis for inherited organic acid disorders by using GC/MS.) *Nippon Iyo Masu Supekutoru Gakkai Koenshu*, 18 (1993) 143-146; *C.A.*, 120 (1994) 6198p.
- 1873 Caboni, M.F., Boschelle, O., Conte, L. and Lercker, G.: (Gas chromatographic analysis of salami at varying stages of maturation.) *Ind. Aliment. (Pinerolo)*, 32 (1993) 950-955; *C.A.*, 120 (1994) 105303a.
- 1874 Ghoos, Y., Claus, D., Geypens, B., Hiele, M., Maes, B. and Rutgeerts, P.: Screening method for the determination of volatiles in biomedical samples by means of an off-line closed-loop trapping system and high resolution gas chromatography-ion trap detection. *J. Chromatogr. A*, 665 (1994) 333-345.

- 1875 Gleispach, H., Leis, H.-J. and Windischhofer, W.: Application of gas chromatography-mass spectrometry in clinical chemistry. *J. Chromatogr. A*, 665 (1994) 155-162.
- 1876 Pitt, J.J. and Poulos, A.: Excretion of 3,6-epoxydicarboxylic acids in peroxisomal disorders. *Clin. Chim. Acta*, 223 (1993) 23-29.
- 1877 Tamasawa, N., Tamasawa, A. and Takebe, K.: Higher levels of plasma cholesterol sulfate in patients with liver cirrhosis and hypercholesterolemia. *Lipids*, 28 (1993) 833-836; *C.A.*, 119 (1993) 265678j.
- See also 1292, 1328, 1442, 1466, 1556, 1558, 1569, 1588, 1592, 1594, 1599, 1628, 1629, 1633, 1677, 1688, 1701.
- ### 34. FOOD ANALYSIS
- #### 34b. Complex mixtures (single compounds by cross-reference only)
- 1878 Baitz, H. and Cacho, C.L.: Gas chromatographic-mass spectrometric (GC-MS) identification of compounds in concentrates of Puerto Rican rums. *J. Agric. Univ. P. R.*, 77 (1993) 69-93; *C.A.*, 120 (1994) 29576z.
- 1879 Bossett, J.O. and Gauch, R.: Comparison of the volatile flavor compounds of six European AOC cheeses by using a new dynamic headspace GC-MS method. *Int. Dairy J.*, 3 (1993) 359-377; *C.A.*, 119 (1993) 179630f.
- 1880 Buecherl, T., Baner, A.L. and Piringer, O.G.: (Extraction and separation of potential migrants from food-contact materials with supercritical carbon dioxide.) *Dtsch. Lebensm.-Rundsch.*, 89, No. 3 (1993) 69-71; *C.A.*, 119 (1993) 269275y.
- 1881 Buiarelli, F., Cartoni, G. and Coccioli, F.: HPLC and GC-MS detection of compounds released to mineral waters stored in plastic bottles of PET and PVC. *Ann. Chim. (Rome)*, 83 (1993) 93-104; *C.A.*, 119 (1993) 188034s.
- 1882 Eiserich, J., Shibamoto, T., Dinnauer, M. and Jennings, W.: Maillard reaction products from microwave heating of model systems: exploiting gas chromatographic variables. In: *Thermally Generated Flavors. ACS Symp. Ser.*, Vol. 543, ACS, Washington, 1994, pp. 55-70; *C.A.*, 120 (1994) 161853s.
- 1883 Froehlich, O.: Components in foods with a high carbohydrate content. In: Wittkowski, R. and Matissek, R. (Editors), *Capillary Gas Chromatogr. Food Control Res.*, Technomic, Lancaster, 1993, pp. 75-90; *C.A.*, 119 (1993) 179406n - a review with 53 refs.
- 1884 Froehlich, O.: Fats and compounds in fat-containing foods. In: Wittkowski, R. and Matissek, R. (Editors), *Capillary Gas Chromatogr. Food Control Res.*, Technomic, Lancaster, 1993, pp. 91-108; *C.A.*, 119 (1993) 179407p - a review with 74 refs.
- 1885 Girard, B. and Nakai, S.: Species differentiation by multivariate analysis of headspace volatile patterns from canned Pacific salmon. *J. Aquat. Food Prod. Technol.*, 2 (1993) 51-68; *C.A.*, 119 (1993) 201952m.
- 1886 Hardt, R.: High-molecular-weight compounds. In: Wittkowski, R. and Matissek, R. (Editors), *Capillary Gas Chromatogr. Food Control Res.*, Technomic, Lancaster, 1993, pp. 177-203; *C.A.*, 119 (1993) 179410j - a review with 74 refs.
- 1887 Moriyasu, T., Saito, K., Nakazato, M., Ishikawa, F., Fujinuma, K. and Nishima, T.: (Amounts of acesulfame K, saccharin, and aspartame used in foods.) *Shokuhin Eiseigaku Zasshi*, 34 (1993) 277-282; *C.A.*, 119 (1993) 224654e.
- 1888 Rosen, R.T., Roshdy, T.H., Hartman, T.G. and Ho, C.T.: Determination of free and glycosidically bound organic compounds in an umbelliferous vegetable drink. In: *Food Phytochemicals for Cancer Prevention I. ACS Symp. Ser.*, Vol. 546, ACS, Washington, 1994, pp. 249-257; *C.A.*, 120 (1994) 161851q.
- 1889 Sakurai, H., Kashima, N., Kumagai, H., Seki, T., Okuma, K., Ariga, T. and Ishii, K.: (Determination of volatile components of upland wasabi by GC-MS.) *Nihon Daigaku Nojuigakubu Gijutsu Kenkyu Hokoku*, 50 (1993) 34-43; *C.A.*, 119 (1993) 202058m.
- 1890 Schreiber, G.A., Helle, N., Schulzki, G., Spiegelberg, A., Linke, B., Wagner, U. and Boegl, K.W.: Intercomparisons to evaluate the suitability of gas-chromatographic, electron-spin-resonance spectrometric and thermoluminescence methods to detect irradiated foods in routine control. *Radiat. Phys. Chem.*, 42 (1993) 391-396; *C.A.*, 119 (1993) 201965t.
- 1891 Sun, J.B., Severson, R.F. and Kays, S.J.: Quantitative technique for measuring volatile components of baked sweetpotatoes. *HortScience*, 28 (1993) 1110-1113; *C.A.*, 120 (1994) 132509d.
- 1892 Vallejo-Cordoba, B.: Shelf-life prediction and flavor quality assessment of pasteurized milk by multivariate analysis of dynamic headspace gas chromatographic data. Avail. *NLC*, Order No. DANN75441, 1992, 209 pp.; *C.A.*, 120 (1994) 161881z.
- 1893 Vieths, S.: Headspace gas chromatography of highly volatile compounds. In: Wittkowski, R. and Matissek, R. (Editors), *Capillary Gas Chromatogr. Food Control Res.*, Technomic, Lancaster, 1993, pp. 277-324; *C.A.*, 119 (1993) 179411k.
- See also 1311, 1334, 1464, 1478, 1484, 1489, 1515, 1526, 1537, 1567, 1573, 1581, 1584, 1611, 1615, 1616, 1618, 1620, 1622, 1636, 1641, 1651, 1670, 1690, 1691, 1706, 1714, 1737, 1738, 1743, 1751, 1753, 1754, 1760, 1762, 1765, 1766, 1767, 1769, 1778, 1785, 1847, 1962.
- #### 34c. Organoleptically important compounds (flavors, odors, volatiles)
- 1894 Cadwallader, K.R. and Xu, Y.: Analysis of volatile components in fresh grapefruit juice by purge and trap/gas chromatography. *J. Agric. Food Chem.*, 42 (1994) 782-784.
- 1895 Careri, M., Manini, P., Spagnoli, S., Barbieri, G. and Bolzoni, L.: Simultaneous distillation-extraction and dynamic headspace methods in the gas chromatographic analysis of Parmesan cheese volatiles. *Chromatographia*, 38 (1994) 386-394.
- 1896 Cooper, J.F., Tourte, J. and Gros, P.: Determination of pentachlorophenol residues in wine and corks by solvent extraction methodology and specific gas chromatography detection. *Chromatographia*, 38 (1994) 147-150.
- 1897 Karl, V., Dietrich, A. and Mosandl, A.: Stereoisomeric flavor compounds. LXIV: the chiro-specific analysis of 2-alkyl-branched flavor compounds from apple headspace extracts and their sensory evaluation. *Phytochem. Anal.*, 4 (1993) 158-164; *C.A.*, 120 (1994) 6991s.
- 1898 Murota, A.: Canonical discriminant analysis applied to the headspace GC profiles of coffee cultivars. *Biosci., Biotechnol., Biochem.*, 57 (1993) 1043-1048; *C.A.*, 119 (1993) 201979a.

- 1899 Plumas, B., Chaveron, H. and Adenier, H.: Determination of olfactive intensity in the food industry with differential olfactometry and gas chromatography. *Dev. Food Sci.*, 32 (1993) 1009-1019; C.A., 119 (1993) 179501q.
- 1900 Robertson, G.W., Griffiths, D.W., Smith, W.M. and Butcher, R.D.: The application of thermal desorption-gas chromatography-mass spectrometry to the analyses of flower volatiles from five varieties of oilseed rape (*Brassica napus* spp. *oleifera*). *Phytochem. Anal.*, 4 (1993) 152-157; C.A., 119 (1993) 221638s.
- 1901 Takeoka, G.: Aroma compounds. In: Wittkowski, R. and Matissek, R. (Editors), *Capillary Gas Chromatogr. Food Control Res.*, Technomic, Lancaster, 1993, pp. 109-147; C.A., 119 (1993) 179408q - a review with 192 refs.
- 1902 Tateo, F., Brunelli, N., Cucurachi, S. and Ferrillo, A.: New trends in the study of the merits and shortcomings of olive oil in organoleptic terms, in correlation with the GC/MS analysis of the aromas. *Dev. Food Sci.*, 32 (1993) 301-311; C.A., 119 (1993) 179641k.
- See also 1417, 1427, 1439, 1879, 1885.
35. ENVIRONMENTAL ANALYSIS
- 35a. *General papers and reviews*
- 1903 Dunemann, L. and Hajimiragha, H.: Development of a screening method for the determination of volatile organic compounds in body fluids and environmental samples using purge and trap gas chromatography-mass spectrometry. *Anal. Chim. Acta*, 283 (1993) 199-206.
- 1904 Le Lacheur, R.M., Sonnenberg, L.B., Singer, P.C., Christman, R.F. and Charles, M.J.: Identification of carbonyl compounds in environmental samples. *Environ. Sci. Technol.*, 27 (1993) 2745-2753.
- 1905 Prasad, S.S.: Trends in quality assurance in the characterization of chemical contaminants in the environment. *TrAC*, 13 (1994) 157-164.
- See also 1329.
- 35b. *Air pollution (complex mixtures; single compounds by cross-reference only)*
- 1906 Armanino, C., Forina, M., Bonfanti, L. and Maspero, M.: Extracting information from complex chromatographic fingerprints for evaluation of organic air pollution. *Anal. Chim. Acta*, 284 (1993) 79-89.
- 1907 Asilbekova, K.T. and Khasanova, V.M.: (Gas-chromatographic determination of pinacol in air and water.) *Uzb. Khim. Zh.*, No. 3-4 (1992) 16-20; C.A., 119 (1993) 216519z.
- 1908 Khasanova, V.M. and Kulabdullaeva, F.: (Gas-chromatographic determination of intermediates in azocene and traidimenol production in air and water.) *Uzb. Khim. Zh.*, No. 5-6 (1992) 83-85; C.A., 119 (1993) 216518y.
- 1909 Kruschel, B.D., Bell, R.W., Chapman, R.E., Spencer, M.J. and Smith, K.V.: Analysis of ambient polar and non-polar volatile organic compounds (VOCs) by thermal desorption, high resolution gas chromatography-mass spectrometry (TD/HRGC/MS). *J. High Resolut. Chromatogr.*, 17 (1994) 187-190.
- 1910 Mattinen, M.L., Sandell, E. and Saarela, K.: Possibilities of analyzing indoor air by gas chromatography using retention index monitoring. *Toxicol. Environ. Chem.*, 40 (1993) 133-142; C.A., 120 (1994) 142535y.
- 1911 Russow, R. and Foerstel, H.: Use of GC-QMS for stable isotope analysis of environmentally relevant main and trace gases in the air. *Isotopenpraxis*, 29 (1993) 327-334; C.A., 120 (1994) 17198s.
- 1912 Vyskrebentsev, V.P. and Spektorman, E.A.: (Sampling of air pollutants.) *Dokl. Akad. Nauk Resp. Uzb.*, No. 12 (1992) 33-35; C.A., 119 (1993) 209346n.
- 1913 Zhang, S., Liu, Y., Fukasawa, T. and Iwatsuki, M.: Study on organic compounds in airborne particles in Chenkdu area, China, by GC and GC/MS. *Taiki Osen Gakkaiishi*, 28 (1993) 359-366; C.A., 120 (1994) 171888w.
- See also 1350, 1376, 1420, 1463, 1465, 1476, 1500, 1501, 1508, 1549, 1562, 1586, 1603, 1643, 1667, 1671, 1696, 1715, 1756, 1952, 1967.
- 35c. *Water pollution (complex mixtures; single compounds by cross-reference only)*
- 1914 Angelov, G. and Panchev, N.: Direct determination of the organic solvent in organic-in-water emulsions. *Dokl. Bulg. Akad. Nauk.*, 45 (1992) 61-64; C.A., 119 (1993) 194784c.
- 1915 Lesage, S. and Brown, S.: Dynamic headspace analysis of volatile organic solvents in water. *Anal. Chem.*, 66 (1994) 572-575.
- 1916 Milyukin, M.V. and Piliipenko, A.T.: (Identification of microimpurities of semivolatile organic compounds of neutral and acidic character in natural and drinking waters by chromatography-mass spectrometry.) *Khim. Tekhnol. Vody*, 15 (1993) 424-435; C.A., 119 (1993) 210044u.
- 1917 Mingram, W.: (Application of gas chromatography and HPLC for the surveillance of drinking water.) *GIT Spez. Chromatogr.*, 13, No. 2 (1993) 74-81; C.A., 120 (1994) 37645m.
- 1918 Nguyen, D.-K., Bruchet, A. and Arpino, P.: High resolution capillary GC-MS analysis of low molecular weight organic compounds in municipal wastewater. *J. High Resolut. Chromatogr.*, 17 (1994) 153-159.
- 1919 Pomorska, K.: On-line extraction method for the isolation of pesticides from industrial wastewater. *Chem. Anal. (Warsaw)*, 38 (1993) 297-301.
- 1920 Richardson, S.D., Thruston, A.D., Jr., Collette, T.W., Patterson, K.S., Lykins, B.W., Jr., Majetich, G. and Zhang, Y.: Multispectral identification of chlorine dioxide disinfection byproducts in drinking water. *Environ. Sci. Technol.*, 28 (1994) 592-599.
- 1921 Sebok, D.: (Problems of gas chromatographic testing of surface waters.) *Magy. Kem. Lapja*, 48 (1993) 169-172; C.A., 119 (1993) 210006h.
- 1922 Wu, C., Cheng, J., Ma, Z., Chen, X. and Li, Z.: (Analysis of toxic organic compounds in the East Lake by gas chromatography.) *Huanjing Huaxue*, 12 (1993) 59-65; C.A., 119 (1993) 278130z.
- 1923 Wu, X. and Shao, Z.: (Capillary gas chromatographic method for determining organic pollutants in water.) *Shanghai Huanjing Kexue*, 11, No. 5 (1992) 25-27; C.A., 119 (1993) 210017n.
- 1924 Xu, X.: (Identifying the source of spilled oil at sea by matching gas chromatogram pattern with correlation coefficients.) *Haiyang Yu Huzhao*, 23 (1992) 505-510; C.A., 119 (1993) 188026r.

See also 1294, 1299, 1390, 1394, 1422, 1474, 1479, 1486, 1487, 1499, 1500, 1504, 1513, 1514, 1529, 1530, 1533, 1551, 1582, 1666, 1675, 1723, 1736, 1740, 1741, 1742, 1744, 1746, 1757, 1761, 1763, 1768, 1907, 1908, 1927, 1931.

35d. *Soil pollution (complex mixtures; single compounds by cross-reference only)*

See 1297, 1313, 1469, 1471, 1473, 1482, 1491, 1495, 1502, 1504, 1505, 1534, 1541, 1545, 1591, 1609, 1685, 1717, 1721, 1750, 1772, 1776, 1777, 1966.

36. SOME TECHNICAL PRODUCTS AND COMPLEX MIXTURES

36a. *Surfactants*

1925 Field, J.A., Field, T.M., Poiger, T. and Giger, W.: Determination of secondary alkane sulfonates in sewage wastewaters by solid-phase extraction and injection-port derivatization gas chromatography/mass spectrometry. *Environ. Sci. Technol.*, 28 (1994) 497-503.

1926 Hida, M., Okuyama, S., Mitsui, T., Minami, Y. and Fujimura, Y.: Determination of sodium dodecylbenzene sulfonate in ternary mixtures by Curie-point pyrolysis gas chromatography coupled to multivariate analysis. *Chromatographia*, 38 (1994) 436-440.

1927 Schroeder, H.F.: Pollutants in drinking water and waste water. *J. Chromatogr.*, 643 (1993) 145-161.

1928 Wang, Z., and Fingas, M.: Analysis of sorbitan ester surfactants. Part II: Capillary supercritical fluid chromatography. *J. High Resolut. Chromatogr.*, 17 (1994) 85-90.

1929 Ye, M.Y., Hill, K.D. and Walkup, R.G.: Separation of T-MAZ ethoxylated sorbitan fatty acid esters by supercritical fluid chromatography. *J. Chromatogr. A*, 662 (1994) 323-327.

1930 Zhou, M.: (Determination of mean relative molecular mass for cationic surfactants by GC.) *Riyong Huaxue Gongye*, No. 1 (1993) 38-41; C.A., 120 (1994) 137688w.

36b. *Antioxidants and preservatives*

1931 Furtmann, K.: Phtalates in surface water – a method for routine trace level analysis. *Fresenius J. Anal. Chem.*, 348 (1994) 291-296.

36c. *Complex mixtures, technical products and unidentified compounds*

1932 Awan, L.A. and Ikramullah: Characterization of Pakistani coal by pyrolysis gas chromatography. Part II. Shariagh coal. *Sci. Int. (Lahore)*, 4 (1992) 367-372; C.A., 119 (1993) 184466m.

1933 Cutler, E.T., Magidman, P. and Marmer, W.N.: Pyrolysis-gas chromatography of Hercosett 125 finish on wool. *Text. Chem. Color.*, 25, No. 6 (1993) 27-29; C.A., 120 (1994) 109369z.

1934 Felix, J.M. and Gatenholm, P.: Characterization of cellulose fibers using inverse gas chromatography. *Nord. Pulp Pap. Res. J.*, 8 (1993) 200-203; C.A., 120 (1994) 109720g.

1935 Galletti, G.C. and Mazzeo, R.: Pyrolysis/gas chromatography/mass spectrometry and Fourier-transform infrared spectroscopy of amber. *Rapid Commun. Mass Spectrom.*, 7 (1993) 646-650; C.A., 119 (1993) 202615r.

1936 Galletti, G.C., Chiavari, G. and Kahie, Y.D.: Pyrolysis/gas chromatography/ion-trap mass spectrometry of the 'tooth brush' tree (*Salvadora persica* L.). *Rapid Commun. Mass Spectrom.*, 7 (1993) 651-655; C.A., 119 (1993) 221663w.

1937 Galletti, G.C., Reeves, J.B., III, Bloomfield, J., Vogt, K.A. and Vogt, D.J.: Analysis of leaf and fine root litter from a subtropical montane rain forest by pyrolysis-gas chromatography/mass spectrometry. *J. Anal. Appl. Pyrolysis*, 27 (1993) 1-14; C.A., 120 (1994) 76402z.

1938 Hama, H., Matsubayashi, K., Murata, S. and Nomura, M.: (Studies on chemical structure of coal. I. Analysis of chemical constituents of Illinois No. 6 and Miike coals with a Curie-point pyrolyzer.) *Nippon Enerugi Gakkaishi*, 72 (1993) 467-475; C.A., 119 (1993) 206805a.

1939 Hardell, H.L.: Characterization of impurities in pulp and paper products using pyrolysis-gas chromatography/mass spectrometry, including direct methylation. *J. Anal. Appl. Pyrolysis*, 27 (1993) 73-85.

1940 Kahle, G., Staedter, H., Seiler, T. and Wollensak, J.: (Combined gas chromatography/mass-spectroscopy analysis of photoablative products after excimer- or erbium: YAG laser treatment of human corneas.) *Ophthalmologe*, 89 (1992) 505-509; C.A., 119 (1993) 198641b.

1941 Komaitis, M.E. and Aggelousis, G.: Qualitative analysis of volatile compounds from the crust and crumb of anise white bread. *Sci. Aliments*, 13 (1993) 585-591; C.A., 119 (1993) 224509m.

1942 Kondo, T.: Preparation of 6-O-alkylcelluloses. *Carbohydr. Res.*, 238 (1993) 231-240; C.A., 119 (1993) 226287m.

1943 Kuroda, K.: (Analytical pyrolysis of wood.) *Mokuzai Kogyo*, 48 (1993) 460-465; C.A., 119 (1993) 252355b - a review with 29 refs.

1944 Lanças, F.M., Vilegas, J.H.Y., Martins, S. and Gobato, E.A.F.: Chromatographic characterization of thermally upgraded products from alternative fuels. *J. High Resolut. Chromatogr.*, 17 (1994) 237-244.

1945 Meier, D., Berns, J., Gruenwald, C. and Faix, O.: Analytical pyrolysis and semicontinuous catalytic hydrolysis of Organo-cell lignin. *J. Anal. Appl. Pyrolysis*, 25 (1993) 335-347.

1946 Mosandl, A.: (Authenticity control of natural flavors and fragrances.) *Kontakte (Darmstadt)*, No. 3 (1992) 38-49; C.A., 119 (1993) 179379f - a review with 64 refs.

1947 Raynor, M.W. and Bartle, K.D.: Supercritical fluid chromatography and extraction in surface coatings analysis – review of representative applications. *J. Supercrit. Fluids*, 6 (1993) 39-49.

1948 Roberts, D.D. and Acree, T.E.: Gas chromatography-olfactometry of glucose-proline Maillard reaction products. In: *Thermally Generated Flavors. ACS Symp. Ser.*, Vol. 543, ACS, Washington, 1994, pp. 71-79; C.A., 120 (1994) 132547q.

1949 Shedrinsky, A.M., Grimaldi, D.A., Boon, J.J. and Baer, N.S.: Application of pyrolysis-gas chromatography and pyrolysis-gas chromatography/mass spectrometry to the unmasking of amber forgeries. *J. Anal. Appl. Pyrolysis*, 25 (1993) 77-95.

1950 Song, C., Hou, L., Saini, A.K., Hatcher, P.G. and Schobert, H.H.: CPDAS carbon-13 NMR and pyrolysis-GC-MS studies of structure and liquefaction reactions of Montana subbituminous coal. *Fuel Process. Technol.*, 34 (1993) 249-276; C.A., 119 (1993) 184475p.

1951 Takeshita, Y., Iwata, F., Hinata, K. and Yoshida, H.: Study on the identification of rice bran oil. *Kokushikan Daigaku Kogakubu Kiyo*, 26 (1993) 21-31; C.A., 119 (1993) 248317y.

1952 Tanaka, T. and Shoji, Z.: (Analysis of volatile compounds in the natto-fermenting room by gas chromatography-mass spectrometry.) *Nippon Shokuhin Gogyo Gakkaishi*, 40 (1993) 656-660; C.A., 120 (1994) 75943q.

See also 1277, 1280, 1283, 1312, 1391, 1441, 1468, 1485, 1496, 1503, 1509, 1514, 1522, 1531, 1539, 1540, 1553, 1555, 1561, 1563, 1564, 1609, 1664, 1699, 1700, 1703, 1707, 1710, 1713, 1727, 1745, 1889, 1908, 1965.

37. CELLS, CELLULAR PARTICLES AND SUPRAMOLECULAR STRUCTURES

1953 Yang, H., Ye, H., Tong, M. and Li, Z.: (GC/IR spectrometry for anaerobe identification.) *Huazhong Ligong Daxue Xuebao*, 20, No. 5 (1992) 35-38; C.A., 120 (1994) 72858e.

See also 1327, 1570, 1589, 1601.

38. INORGANIC COMPOUNDS

38a. Cations

See 1911.

38c. Permanent and rare gases

1954 Brooks, P.D. and Atkins, G.J.: Rapid isotopic analysis of trace gases at atmospheric levels. *Isotopenpraxis*, 28 (1992) 106-109; C.A., 120 (1994) 123654v.

1955 Caraeal, Z.L., Pradeau, D., Hamon, M., Abdoulaye, I., Paillet, F.M. and Lejeune, B.: New calibration method for gas chromatographic assay of carbon monoxide in blood. *J. Anal. Toxicol.*, 17 (1993) 193-195; C.A., 119 (1993) 197341s.

1956 Kovalev, L.V., Denisova, L.V., Menshikova, N.V. and Timofeev, I.A.: (Gas-chromatography method and device for determination of the trace contaminants of permanent and rare gases in oxygen.) U.S.S.R. SU 1,761,231 (Cl. B01D53/02), 15 Sep. 1992, Appl. 4,702,006, 7 Jun. 1989; C.A., 120 (1994) 152578d.

1957 Nagayanagi, H. and Yuguchi, H.: (Method and apparatus for chromatographic analysis of inorganic gas.) *Jpn. Kokai Tokkyo Koho* JP 05,273,184 [93,273,184] (Cl. G01N30/14), 22 Oct. 1993, Appl. 92/97,190, 24 Mar. 1992; 4 pp.; C.A., 120 (1994) 152600e.

1958 Yasui, S.: (Determination of argon in oxygen by gas chromatography.) *Jpn. Kokai Tokkyo Koho* JP 05,180,822 [93,180,822] (Cl. G01N30/88), 23 Jul. 1993, Appl. 91/355,763, 20 Dec. 1991; 5 pp.; C.A., 120 (1994) 22731n.

See also 1317, 1318, 1320, 1349, 1363, 1460.

38d. Volatile inorganic compounds

1959 Cullen, W.R., Li, H., Pergantis, S.A., Eigendorf, G.K. and Harrison, L.G.: The methylation of arsenate by a marine alga *Polyphysa peniculus* in the presence of L-methionine-methyl-D₃. *Chemosphere*, 28 (1994) 1009-1019.

1960 Liu, J.X., Gilbert, S.G., Mannheim, C.H. and Daun, H.: Determination of critical moisture content of flour by inverse gas chromatography. *Dev. Food Sci.*, 32 (1993) 753-767; C.A., 119 (1993) 202177z.

1961 Makino, Y.: (Gas chromatograph for determination of gases produced from thermal decomposition of hexafluorosulfide in electric appliances.) *Jpn. Kokai Tokkyo Koho* JP 04,120,461 [92,120,461] (Cl. G01N30/78), 21 Apr. 1992, Appl. 90/240,931, 10 Sep. 1990; 4 pp.; C.A., 120 (1994) 123757f.

1962 Mohr, S.A., Zottola, E.A. and Reineccius, G.A.: The use of gas chromatography to measure carbon dioxide production by dairy starter cultures. *J. Dairy Sci.*, 76 (1993) 3350-3353; C.A., 120 (1994) 7070c.

1963 Morimoto, F. and Hirai, T.: (Method and column for removing corrosive gases in chromatography in inorganic gas analysis.) *Jpn. Kokai Tokkyo Koho*, JP 04 76,455 [92 76,455] (Cl. G01N30/14), 11 Mar. 1992, Appl. 90/190,038, 17 Jul. 1990; 6 pp.; C.A., 120 (1994) 181994s.

1964 Ogino, H., Aomura, Y. and Mizuno, T.: (Method and apparatus for the determination of impurities in nitrogen gas.) *Jpn. Kokai Tokkyo Koho* JP 04,143,657 [92,143,657] (Cl. G01N30/14), 18 May 1992, Appl. 90/268,162, 5 Oct. 1990; 8 pp.; C.A., 120 (1994) 123787r.

1965 Uchida, S.: (Studies on the evaluation of dry matter content and feeding value in silage. VI. Estimation and application of toluene distillation method, freeze drying method, and gas chromatography for moisture determination.) *Nippon Sochi Gakkaishi*, 39 (1993) 155-161; C.A., 120 (1994) 7000e.

1966 Walsh, M.E. and Taylor, S.: Analytical method for white phosphorus residues in munitions-contaminated sediments. *Anal. Chim. Acta*, 282 (1993) 55-61.

1967 Yamamoto, N., Nishiura, H., Honjo, T., Ishikawa, Y. and Suzuki, K.: Continuous determination of atmospheric ammonia by an automated gas chromatographic system. *Anal. Chem.*, 66 (1994) 756-760.

See also 1318, 1320, 1363, 1460, 1710, 1715, 1954.

39. RADIOACTIVE AND OTHER ISOTOPE COMPOUNDS

See 1301, 1318, 1404, 1409, 1450, 1572, 1574, 1576, 1580, 1593, 1598, 1682, 1836, 1842, 1911, 1954.

Planar Chromatography

1. REVIEWS AND BOOKS

- 452 Jork, H.: Advances in thin-layer chromatography: Part 1. *Am. Lab. (Shelton)*, 25, No. 7 (1993) 36L, 36N-36O; *C.A.*, 120 (1994) 94044p - a review with 58 refs.
- 453 Lin, L. and Zhang, J.: (Current status and trend of quantitative thin-layer chromatography). *Sepu*, 11 (1993) 339-345; *C.A.*, 120 (1994) 94336k - a review with 25 refs.
- 454 Okuda, K.-I.: Liver mitochondrial P 450 involved in cholesterol catabolism and vitamin D activation. *J. Lipid Res.*, 35 (1994) 361-372.
- 455 Wang, Y., Pu, F. and He, X.: (Chinese medicine "Qianghuo"). *Tianran Chanwu Yanjiu Yu Kaifa*, 5, No. 4 (1993) 40-46; *C.A.*, 120 (1994) 143749h - a review with 18 refs.

See also 605.

2. FUNDAMENTALS, THEORY AND GENERAL

2a. General

- 456 Bussemas, H.H., Harsch, G. and Etre, L.S.: Friedlieb Ferdinand Lunge (1794-1867): "Self grown pictures" as precursors of paper chromatography. *Chromatographia*, 38 (1994) 243-254.
- 457 Etre, L.S.: New, unified nomenclature for chromatography. *Chromatographia*, 38 (1994) 521-526.

2b. Thermodynamics and theoretical relationships

See 606.

2c. Relationship between structure and chromatographic behaviour

- 458 Dimov, N.: Application of quantitative structure-retention relationships for chromatographic interaction elucidation. *Bulg. Chem. Commun.*, 25 (1992) 367-374; *C.A.*, 120 (1994) 244034b.
- 459 Wang, F., Jiang, Y., Guo, Y. and Cao, T.: (Relation between homologous retention values and homologous factors in thin layer chromatography). *Fenxi Shiyanshi*, 12, No. 5 (1993) 22-24; *C.A.*, 120 (1994) 123838h.

2d. Measurement of physico-chemical and related values

See 573.

3. GENERAL TECHNIQUES

3a. Apparatus and accessories

- 460 Dzido, T.: Method and chamber for developing thin-layer chromatograms. *Pol. PL* 161,388 (Cl. G01N30/90), 30 Jun. 1993, *Appl.* 280,087, 17 Jun. 1989, 8 pp.; *C.A.*, 120 (1994) 94435s.

See also 560.

3b. Detectors and detection reagents

- 461 Maslowska, J. and Owczarek, A.: (Use of acid phosphatase for detection and determination of typical anions on chromatograms). *Przem. Chem.*, 72 (1993) 495-496; *C.A.*, 120 (1994) 181774v.
- 462 Yin, P., Li, H. and Yan, C.: (Application of surfactants to thin-layer chromatography with a mixed plate of GDX-102 and silica gel G). *Sepu*, 12, No. 1 (1994) 35-36; *C.A.*, 120 (1994) 182109u.

See also 518, 532, 549, 550, 551, 560.

3c. Sorbents and columns, packing procedures

- 463 Hauck, H.E., Junker-Buchheit, A. and Wenig, R.: Reproducibility of silica gel 60 precoated plates in TLC and high-performance TLC. *GIT Fachz. Lab.*, 37 (1993) 973-977; *C.A.*, 120 (1994) 94388d.
- 464 Nazli, S. and Naser-Ud-Din.: Utilization of marble powder and the clay (Gacci Mutti) as thin layer chromatography adsorbent. *Pak. J. Sci. Ind. Res.*, 36 (1993) 166-168; *C.A.*, 120 (1994) 174369p.

See also 607.

3d. Quantitative analysis

See 453, 565.

3f. Programmed temperature, pressure, vapors, gradients

See 525.

4. SPECIAL TECHNIQUES

4b. Computerization and modelling

- 465 Markowski, W., Czapinska, K.L. and Blaszcak, M.: Determination of the constants of the Snyder-Soczewinski equation by means of gradient multiple development. *J. Liq. Chromatogr.*, 17 (1994) 999-1009.

- 466 Matyska, M. and Soczewinski, E.: Optimization of chromatographic systems in TLC by graphical method and with a computer program. *Chem. Anal. (Warsaw)*, 38 (1993) 555-563; C.A., 120 (1994) 207653g.
- See also 560.
- 4c. *Combination with other physico-chemical techniques (MS, IR etc.)*
- See 559, 601.
- 4f. *Trace analysis and preseparation techniques*
- See 491.
- 4g. *Enantiomers, separation*
- See 477.
- 4h. *Other special techniques*
- See 502, 557, 611.
5. HYDROCARBONS AND HALOGEN DERIVATIVES
- 5d. *Complex hydrocarbon mixtures (incl. analysis of tars, bitumens and mineral oils)*
- 467 Friedbacher, E.C. and Schindlbauer, H.: Analysis of bitumen by TLC-FID compared with column chromatography. *Bitumen*, 55 (1993) 149-151; C.A., 120 (1994) 195460b.
6. ALCOHOLS
- See 486.
7. PHENOLS
- See 470.
8. SUBSTANCES CONTAINING HETEROCYCLIC OXYGEN
- 8c. *Other compounds with heterocyclic oxygen (incl. tannins)*
- 468 Zhang, H., Chen, W. and Yuan, C.: (A comparative studies on coumarins constituents of Chinese crude drug "Chuanduhuo" and "Zheduhuo" (*Angelica biserrata* Shan et Yuan) Yuan et Shan). *Zhiwu Zhiyuan Yu Huanjing*, 2, No. 4 (1993) 60-62; C.A., 120 (1994) 227072v.
9. OXO COMPOUNDS, ETHERS, EPOXIDES AND QUINONES
- 469 Wang, L., Liu, H. and Shu, L.: (Separation and identification of anthraquinone derivatives by thin-layer chromatography). *Zhejiang Gongxueyuj Xuebao*, No. 1 (1993) 66-71; C.A., 120 (1994) 152677k.
10. CARBOHYDRATES
- 10a. *Mono and oligosaccharides. Structural studies*
- 470 Amarowicz, R. and Shahidi, F.: Application of Sephadex LH-20 chromatography for the separation of cyanogenic glycosides and hydrophylic phenolic fraction from flaxseed. *J. Liq. Chromatogr.*, 17 (1994) 1291-1299.
- 471 Li, Y.T., Yuziuk, J.A., Li, S.-C., Nematalla, A., Hasegawa, A., Kimura, M. and Nakagawa, H.: A novel sialidase capable of cleaving 3-deoxy-D-glycero-D-galacto-2-nonulosonic acid (KDN). *Arch. Biochem. Biophys.*, 310 (1994) 243-246.
- See also 585.
- 10b. *Polysaccharides, mucopolysaccharides, lipopolysaccharides*
- 472 Pavao, M.S.G., Rodrigues, M.A. and Mourão, P.A.S.: Acidic polysaccharides of the ascidian *Styela plicata*. Biosynthetic studies on the sulfated L-galactans of the tunic, and preliminary characterization of a dermatan sulfate-like polymer in body tissues. *Biochim. Biophys. Acta*, 1199 (1994) 229-237.
- 10c. *Glycoproteins and their constituents*
- 473 Teneberg, S., Ångström, J., Jovall, P. and Karlsson, K.-A.: Characterization of binding of Gal β 4GlcNAc-specific lectins from *Erythrina cristagalli* and *Erythrina corallodendron* to glycosphingolipids. Detection, isolation, and characterization of a novel glycosphingolipid of bovine buttermilk. *J. Biol. Chem.*, 269 (1994) 8554-8563.
11. ORGANIC ACIDS AND LIPIDS
- 11a. *Organic acids and simple esters*
- 474 Akoh, C.C.: Enzymatic synthesis of acetylated glucose fatty acid esters in organic solvent. *J. Am. Oil Chem. Soc.*, 71 (1994) 315-323.
- 475 Aviram, M. and Rosenblat, M.: Macrophage-mediated oxidation of extracellular low density lipoprotein requires an initial binding of the lipoprotein to its receptor. *J. Lipid Res.*, 35 (1994) 385-398.
- 476 Bean, T.A., Zhuang, W.C., Tong, P.Y., Eick, J.D. and Yourtee, D.M.: Effect of esterase on methacrylates and methacrylate polymers in an enzyme simulator for biodurability and biocompatibility testing. *J. Biomed. Mater. Res.*, 28, No. 1 (1994) 59-63; C.A., 120 (1994) 226874w.

- 477 Borges, M.F.M. and Pinto, M.M.M.: Separation of the diastereoisomers of ethyl esters of caffeic, ferulic, and isoferulic acids by thin-layer and high performance liquid chromatography. *J. Liq. Chromatogr.*, 17 (1994) 1125-1139.
- 478 Leont'eva, L.B., Tselinskii, I.V., Buichinova, E.S. and Krauklish, I.V.: (Determination of *m*-nitrobenzoic acid by peak paper chromatography). *Zh. Prikl. Khim. (St.-Peterburg)*, 66 (1993) 1650-1652; *C.A.*, 120 (1994) 182130u.
- 479 Park, E.K.C. and Obendorf, S.K.: Chemical changes in unsaturated oils upon aging and subsequent effects on fabric yellowing and soil removal. *J. Am. Oil Chem. Soc.*, 71 (1994) 17-30.
- 480 Shi, Y. and Huang, W.: (Suitable developing solvents in silica gel thin-layer chromatography for separating and identifying abscisic acid in Barnyard grass). *Sepu*, 11 (1993) 360-388; *C.A.*, 120 (1994) 49253v.
- 481 Thomas, C.E., Jackson, R.L., Ohlweiler, D.F. and Ku, G.: Multiple lipid oxidation products of low density lipoproteins induce interleukin-1 beta release from human blood mononuclear cells. *J. Lipid Res.*, 35 (1994) 417-427.
- 482 Wilde, J. and Eggerer, H.: Substrate and inhibitor specificity of mevalonate kinase determined with substrate analogues. *Eur. J. Biochem.*, 221 (1994) 463-473.
- 483 Wolff, R.L.: Contribution of *trans*-18:1 acids from dairy fat to European diets. *J. Am. Oil Chem. Soc.*, 71 (1994) 277-283.
- 484 Wolff, R.L. and Sebedio, J.-L.: Characterization of γ -linolenic acid geometrical isomers in borage oil subjected to heat treatments (deodorization). *J. Am. Oil Chem. Soc.*, 71 (1994) 117-126.
- See also 458, 497.
- 11c. *Lipids and their constituents*
- 485 Ariga, T., Tao, R.V., Lee, B., Yamawaki, M., Yoshino, H., Scarsdale, N.J., Kasama, T., Kushi, Y. and Yu, R.K.: Glycolipid composition of human cataractous lenses. Characterization of Lewis^x glycolipids. *J. Biol. Chem.*, 269 (1994) 2667-2675.
- 486 Bianchi, G. and Vlahov, G.: Composition of lipid classes in the morphologically different parts of the olive fruit, cv. *Coratina* (*Olea europaea* Linn.) *Fat Sci. Technol.*, 96 (1994) 72-77.
- 487 Boulanger, J., Huesca, M., Arab, S. and Lingwood, C.A.: Universal method for the facile production of glycolipid/lipid matrices for the affinity purification of binding ligands. *Anal. Biochem.*, 217 (1994) 1-6.
- 488 Cartwright, I.J.: Separation and analysis of phospholipids by thin layer chromatography. *Methods Mol. Biol. (Totowa)*, (1993) 153-167; *C.A.*, 120 (1994) 26661n.
- 489 Chigorno, V., Valsecchi, M. and Sonnino, S.: Biosynthesis of gangliosides containing C18:1 and C20:1 [3-¹⁴C]sphingosine after administrating [1-¹⁴C]pamitic acid and [1-¹⁴C]stearic acid to rat cerebellar granule cells in culture. *Eur. J. Biochem.*, 221 (1994) 1095-1101.
- 490 Dasgupta, S., Hogan, E.L., Glushka, J. and Halbeek, H.: Branched monosialo gangliosides of the lacto-series isolated from bovine erythrocytes: characterization of a novel ganglioside, NeuGc-iso-octaacylceramide. *Arch. Biochem. Biophys.*, 310 (1994) 373-384.
- 491 Deleva, D.D., Ilinov, P.P. and Zaprianova, E.T.: Modified procedure for determination of lipid-bound sialic acid in rat brain. *Bulg. Chem. Commun.*, 25 (1992) 514-519; *C.A.*, 120 (1994) 186406m.
- 492 Djavaheri-Mergny, M., Mora, L., Mazière, C., Auclair, M., Santus, R., Dubertret, L. and Mazière, J.-C.: Inhibitor of diphosphatidylglycerol synthesis by UVA radiations in N.C.T.C. 2544 human keratinocytes. *Biochem. J.*, 299 (1994) 85-90.
- 493 Gelsema, W.J., Choma, I., den Ouden, T.A.F., Zander, R. and van den Bosch, H.: Quantitation of the diacyl, alkylacyl, and alk-1-enylacyl subclasses of choline glycerophospholipids by chemical dephosphorylation and benzoylation. *Anal. Biochem.*, 217 (1994) 265-276.
- 494 Gerold, P., Dieckmann-Schuppert, A. and Schwarz, R.T.: Glycosylphosphatidylinositols synthesized by asexual erythrocytic stages of the malarial parasite, *Plasmodium falciparum*. Candidates for plasmodial glycosylphosphatidylinositol membrane anchor precursors and pathogenicity factors. *J. Biol. Chem.*, 269 (1994) 2597-2606.
- 495 Govers-Riemslog, J.W.P., Janssen, M.P., Zwaal, R.F.A. and Rosing, J.: Prothrombin activation on dioleoylphosphatidylcholine membranes. *Eur. J. Biochem.*, 220 (1994) 131-138.
- 496 Hidari, K.I.-P., Jwa, Kawashima, I., Tai, T., Inagaki, F., Nagai, Y. and Sanai, Y.: *In vitro* synthesis of disialoganglioside (GD_{1a}) from asialo-G_{M1} using sialyltransferases in rat liver Golgi vesicles. *Eur. J. Biochem.*, 221 (1994) 603-609.
- 497 Kamal-Eldin, A. and Appelqvist, L.Å.: Variation in fatty acid composition of the different acyl lipids in seed oils from four *Sesamum* species. *J. Am. Oil Chem. Soc.*, 71 (1994) 135-139.
- 498 Kawakami, Y., Nakamura, K., Kojima, H., Suzuki, M., Inagaki, F., Suzuki, A., Sonoki, S., Uchida, A., Murata, Y. and Tamai, Y.: A novel fucosylated glycosphingolipid with a Ga β 1-4Glc β 1-3Gal sequence in plerocercoids of the parasite, *Spirometra erinacei*. *J. Biochem. (Tokyo)*, 114 (1993) 677-683.
- 499 Kodaki, T., Woscholski, R., Emr, S., Waterfield, M.D., Nurse, P. and Parker, P.J.: Mammalian phosphatidylinositol 3-kinase induces a lethal phenotype on expression in *Schizosaccharomyces pombe*; comparison with the VPS34 gene product. *Eur. J. Biochem.*, 219 (1994) 775-780.
- 500 Kotani, M., Kawashima, I., Ozawa, H., Ogura, K., Ariga, T. and Tai, T.: Generation of one set of murine monoclonal antibodies specific for globo-series glycolipids: evidence for differential distribution of the glycolipids in rat small intestine. *Arch. Biochem. Biophys.*, 310 (1994) 89-96.
- 501 Lingwood, C.A. and Nutikka, A.: A novel chemical procedure for the selective removal of nonreducing terminal N-acetyl hexosamine residues from glycolipids. *Anal. Biochem.*, 217 (1994) 119-123.
- 502 Matsuyama, N., Mian, I.H., Akanda, A.M. and Furuya, N.: (Comparative studies on thin-layer chromatograms of lipids from various phytopathogenic bacteria). *Nippon Shokubutsu Byori Gakkaiho*, 59 (1993) 528-534; *C.A.*, 120 (1994) 26686z.
- 503 Mohney, R.P., Knez, J.J., Ravi, L., Sevlever, D., Rosenberry, T.L., Hirose, S. and Medof, M.E.: Glycoinositol phospholipid anchor-defective K562 mutants with biochemical lesions distinct from those in Thy-1⁻ murine lymphoma mutants. *J. Biol. Chem.*, 269 (1994) 6536-6542.

- 504 Munnik, T., Irvine, R.F. and Musgrave, A.: Rapid turnover of phosphatidylinositol 3-phosphate in the green alga *Chlamydomonas eugametos*: signs of a phosphatidylinositide 3-kinase signalling pathway in lower plants? *Biochem. J.*, 298 (1994) 269-273.
- 505 Mùthing, J., Maurer, U., Šoštarić, K., Neumann, U., Brandt, H., Duvar, S., Peter-Katalinic, J. and Weber-Schürholz, S.: Different distributions of glycosphingolipids in mouse and rabbit skeletal muscle demonstrated by biochemical and immunohistological analyses. *J. Biochem. (Tokyo)*, 115 (1994) 248-256.
- 506 Nohara, K., Nakauchi, H. and Spiegel, S.: Glycosphingolipids of rat T cells. Predominance of asialo-GM1 and GD1c. *Biochemistry*, 33 (1994) 4661-4666.
- 507 Paiement, J., Lavoie, C., Gavino, G.R. and Gavino, V.C.: Modulation of GTP-dependent fusion by linoleic and arachidonic acid in derivatives of rough endoplasmic reticulum from rat liver. *Biochim. Biophys. Acta*, 1190 (1994) 199-212.
- 508 Reddy, S.Y. and Prabhakar, J.V.: Cocoa butter extenders from kokum (*Garcinia indica*) and Phulwara (*Madhuca butyracea*) butter. *J. Am. Oil Chem. Soc.*, 71 (1994) 217-219.
- 509 Šajbidor, J., Čertík, M. and Grego, J.: Lipid analysis of baker's yeast. *J. Chromatogr. A*, 665 (1994) 191-195.
- 510 Sasagasako, N., Kobayashi, T., Yamaguchi, Y., Shinnoh, N. and Goto, I.: Glucosylceramide and glucosylsphingosine metabolism in cultured fibroblasts deficient in acid β -glucosidase activity. *J. Biochem. (Tokyo)*, 115 (1994) 113-119.
- 511 Taylor, D.C., MacKenzie, S.I., McCurdy, A.R., McVetty, P.B.E., Giblin, E.M., Pass, E.W., Stone, S.J., Scarth, R., Rimmer, S.R. and Pickard, M.D.: Stereospecific analyses of seed triacylglycerols from high-erucic acid Brassicaceae: detection of erucic acid at the sn-2 position in *Brassica oleracea* L. genotypes. *J. Am. Oil Chem. Soc.*, 71 (1994) 163-167.
- 512 Tsvetnitsky, V. and Gibbons, W.A.: An assay for phosphatidylserine decarboxylase using BondElut columns. *Anal. Biochem.*, 217 (1994) 157-158.
- 513 Ucciani, E., Mallet, J.F. and Zahra, J.P.: Cyanolipids and fatty acids of *Sapindus trifoliatus* L. (Sapindaceae) seed oil. *Fat Sci. Technol.*, 96 (1994) 69-71.
- 514 Van Veldhoven, P.P. and Mannaerts, G.P.: Sphingosine 1-phosphate metabolism in cultured skin fibroblasts: evidence for the existence of a sphingosine phosphatase. *Biochem. J.*, 299 (1994) 597-601.
- 515 Xu, G., Suzuki, T., Tahara, H., Kiso, M., Hasegawa, A. and Suzuki, Y.: Specificity of sialyl-sugar chain mediated recognition by the hemagglutinin of human influenza B virus isolates. *J. Biochem. (Tokyo)*, 115 (1994) 202-207.
- 516 Yamane, T., Kang, S.T., Kawahara, K. and Koizumi, Y.: High-yield diacylglycerol formation by solid-phase enzymatic glycerolysis of hydrogenated beef tallow. *J. Am. Oil Chem. Soc.*, 71 (1994) 339-342.
- 517 Yang, X. and Ryals, P.E.: Cytodifferentiation in *Tetrahymena vorax* in linked to glycosyl-phosphatidylinositol-anchored protein assembly. *Biochem. J.*, 298 (1994) 697-703.
- 518 Zellmer, S. and Lasch, J.: Quantitative determination of phosphatidylethanolamine and phosphatidylserine in liposomes and on HPTLS plates with o-phthalaldehyde. *Anal. Biochem.*, 218 (1994) 229-231.
12. ORGANIC PEROXIDES
- 519 Chladek, J.: The identification of organic peroxides. *Adv. Anal. Detect. Explos., Proc. Int. Symp., 4th (1992)*, 1993, pp. 73-76; C.A., 120 (1994) 237678m.
13. STEROIDS
- 13b. Pregnane and androstane derivatives
- 520 Leyssens, L., Royackers, E., Gielen, B., Missotten, M., Schoofs, J., Czech, J., Noben, J.P., Hendriks, L. and Raus, J.: Metabolites of 4-chlorotestosterone acetate in cattle urine as diagnostic markers for its illegal use. *J. Chromatogr. B*, 654 (1994) 43-54.
- 521 Luo, Z., He, Y.-a. and Halpert, J.R.: Role of residues 363 and 206 in conversion of cytochrome P450 2B1 from a steroid 16-hydroxylase to a 15 α -hydroxylase. *Arch. Biochem. Biophys.*, 309 (1994) 52-57.
- 522 Sutorkova-Milosevic, L. and Dimitrovska, A.: (Densitometric determination of norgestrel in pharmaceutical preparations). *Arch. Farm.*, 42 (1992) 137-140; C.A., 120 (1994) 200494x.
- 13d. Sterols
- 523 Kamal-Eldin, A. and Appelqvist, L.Å.: Variations in the composition of sterols, tocopherols and lignans in seed oils from four *Sesamum* species. *J. Am. Oil Chem. Soc.*, 71 (1994) 149-156.
- See also 454, 479, 486, 497, 586.
14. STEROID GLYCOSIDES AND SAPONINS
- 524 Lei, P., Gou, Y. and Cong, Y.: (Determination of ginsenoside in Wuzhenshenjiaye by HPLC). *Zhongcaoyao*, 24 (1993) 631-642; C.A., 120 (1994) 227076z.
- 525 Matysik, G.: Gradient thin-layer chromatography of extracts from *Digitalis lanata* and *Digitalis purpurea*. *Chromatographia*, 38 (1994) 109-113.
- See also 583.
15. TERPENES AND OTHER VOLATILE AROMATIC COMPOUNDS
- 15a. Terpenes
- See 486, 583.
- 15b. Essential oils
- 526 Liu, X., Sun, H. and Zeng, X.: (Application of cluster analysis in chemical taxonomy of medicinal material Maidong). *Zhongguo Zhongyao Zazhi*, 18 (1993) 585-587; C.A., 120 (1994) 86567w.

16. NITRO AND NITROSO COMPOUNDS

See 478.

17. AMINES, AMIDES AND RELATED NITROGEN COMPOUNDS

17a. Amines and polyamines

- 527 Kawase, H., Hasegawa, T., Hashimoto, H., Noto, T. and Nakajima, T.: A study on the metabolism of spermidine in mammals: purification and identification of a newly identified metabolite, 2-oxo-1-pyrrolidinepropionic acid, in rat urine. *J. Biochem. (Tokyo)*, 115 (1994) 356-361.
- 528 Pavoni, G.: (Rapid high-performance thin-layer chromatographic (HPTLC) method for determination of amitraz and its transformation product 2,4-xylidine, in honey). *Boll. Chim. Ig., Parte Sci.*, 44 (1993) 283-287; *C.A.*, 120 (1994) 132512z.

18. AMINO ACIDS AND PEPTIDES; CHEMICAL STRUCTURE OF PROTEINS

18a. Amino acids and their derivatives

- 529 Norfolk, E., Khan, S.H., Fried, B. and Sherma, J.: Comparison of amino acid separations on high performance silica gel, cellulose, and C-18 reversed phase layers and application of HPTLC to the determination of amino acids in *Biomphalaria glabrata* snails. *J. Liq. Chromatogr.*, 17 (1994) 1317-1326.

See also 531, 533.

18b. Peptides, peptidic and proteinous hormones, growth factors

- 530 Brattsand, G., Marklund, U., Nylander, K., Roos, G. and Gullberg, M.: Cell-cycle-regulated phosphorylation of oncoprotein 18 on Ser 16, Ser 25 and Ser 38. *Eur. J. Biochem.*, 220 (1994) 359-368.
- 531 Chambers, T.C., Pohl, J., Glass, D.B. and Kuo, J.F.: Phosphorylation by protein kinase C and cyclic AMP-dependent protein kinase of synthetic peptides derived from the linker region of human P-glycoprotein. *Biochem. J.*, 299 (1994) 309-315.
- 532 El-Thaher, T.S. and Bailey, G.S.: A new staining method for cyclic peptides after thin-layer chromatography. *Anal. Biochem.*, 217 (1994) 335-337.
- 533 Georgopapadakou, N.H., Hall, C.C., Lambros, T., Liu, W. and Watkins, J.D.: A radiometric assay for Ras-processing peptidase using an enzymatically radiolabeled peptide. *Anal. Biochem.*, 218 (1994) 273-277.
- 534 Liu, F. and Roth, R.A.: Identification of serines 967/968 in the juxtamembrane region of the insulin receptor as insulin-stimulated phosphorylation sites. *Biochem. J.*, 298 (1994) 471-477.

20. ENZYMES AND ENZYME ACTIVITY ESTIMATION

20a. Oxidoreductases

See 454.

21. PURINES, PYRIMIDINES, NUCLEIC ACIDS AND THEIR CONSTITUENTS

21a. Purines, pyrimidines, nucleosides, nucleotides

- 535 Hayashi, D.K. and Stark, B.C.: A novel tRNA precursor cleaving endoribonuclease from *Zea mays*. *Arch. Biochem. Biophys.*, 309 (1994) 123-128.
- 536 Ito, K., Matsuura, Y. and Minamiura, N.: Purification and characterization of fungal nuclease composed of heterogeneous subunits. *Arch. Biochem. Biophys.*, 309 (1994) 160-167.
- 537 Maga, G., Spadari, S., Wright, G.E. and Focher, F.: Identification, partial purification and inhibition by guanine analogues of a novel enzymic activity which phosphorylates guanosine to GMP in the protozoan parasite *Eimeria tenella*. *Biochem. J.*, 298 (1994) 289-294.

22. ALKALOIDS

- 538 Jain, R.: Interference of adulterants in thin layer chromatography method for drugs of abuse. *Indian J. Pharmacol.*, 25 (1993) 240-242; *C.A.*, 120 (1994) 210175h.
- 539 Rashed, A., Lali, S.A. and Chaudhry, H.R.: Identification and quantitative determination of morphine in the urine of heroin dependents. *Pak. J. Pharm. Sci.*, 5 (1992) 101-109; *C.A.*, 120 (1994) 210178m.

23. OTHER SUBSTANCES CONTAINING HETEROCYCLIC NITROGEN

23e. Other N-heterocyclic compounds

- 540 Feng, Y., Zhang, Z., Chen, Z., Lu, Q. and Xia, C.: (Application of thin-layer chromatography and densitometry in the analysis of the products of $S_{RN}1$ reaction, N-benzylcarbazole). *Sepu*, 11 (1993) 379-380; *C.A.*, 120 (1994) 94396e.
- 541 Leont'eva, L.B., Tselinskii, I.V., Boichinova, E.S. and Krauklish, I.V.: (Determination of 3-nitro-1,2,4-triazole by solvent peak paper chromatography). *Izv. Vyssh. Zaved. Khim. Khim. Tekhnol.*, 36, No. 7 (1993) 17-20; *C.A.*, 120 (1994) 152693n.

See also 578.

24. ORGANIC SULPHUR COMPOUNDS (INCL. GLUCOSINOLATES)

- 542 Ojha, K.G., Mathur, N. and Adwani, P.: Thin-layer chromatographic studies of some substituted 4H-[1,4]-benzothiazines. *Asian J. Chem.*, 4 (1992) 924-926; *C.A.*, 120 (1994) 94380v.

25. ORGANIC PHOSPHORUS COMPOUNDS (INCL. SUGAR PHOSPHATES)
- 543 McClelland, P., Adam, L.P. and Hathaway, D.R.: Identification of a latent Ca²⁺/calmodulin dependent protein kinase II phosphorylation site in vascular calpain II. *J. Biochem. (Tokyo)*, 115 (1994) 41-46.
- See also 530, 531, 534, 551.
26. ORGANOMETALLIC AND RELATED COMPOUNDS
- 26a. *Organometallic compounds*
- See 608.
- 26c. *Coordination compounds*
- 544 Yao, G., Cui, D., Xing, S., Pang, J., Fang, S. and Mei, W.: (Inclusion complex of β -cyclodextrin and nitroglycerin). *Zhongguo Yiyao Gongyo Zazhi*, 24 (1993) 445-447; *C.A.*, 120 (1994) 200339a.
27. VITAMINS AND VARIOUS ANIMAL GROWTH FACTORS (NON-PEPTIDIC)
- See 454, 462, 523, 558.
28. ANTIBIOTICS
- 545 El-Adi, S., Lashine, S. and Shanawani, A.: (Structural elucidation of griseofulvin-2,3-dichloro-5,6-dicyanobenzoquinone complex). *Spectrosc. Lett.*, 27 (1994) 281-291; *C.A.*, 120 (1994) 227085b.
- 546 Prokša, B., Adamcová, J. and Fuska, J.: Detection and assay of secondary metabolites of *Penicillium vermiculatum* DANG. *J. Chromatogr. A*, 665 (1994) 185-190.
- 547 Yanagisawa, M., Sakai, A., Adachi, K., Sano, T., Watanabe, K., Tanaka, Y. and Okuda, T.: Hispidospermidin, a novel phospholipase C inhibitor produced by *Chaetosphaeronema hispidulum* (Cda) Moesz NR 7127. I. Screening, taxonomy, and fermentation. *J. Antibiot.*, 47 (1994) 1-5.
- See also 579.
29. INSECTICIDES, PESTICIDES AND OTHER AGROCHEMICALS
- 29a. *General techniques*
- 548 Darwish, Y., Cserhádi, T. and Forgács, E.: Reversed-phase thin-layer chromatography of various pesticides in the presence of water soluble beta-cyclodextrin polymer. *Chromatographia*, 38 (1994) 509-513.
- 549 Kasatkina, A.A., Ivanova, L.P., Trifonova, V. and Galeeva, G.K.: (Polyethylene polyamines as reagents for thin-layer chromatography). *Zh. Prikl. Khim. (S. Peterburg)*, 66 (1993) 1177-1178; *C.A.*, 120 (1994) 156492u.
- 29b. *Chlorinated insecticides*
- 550 Satapathy, K.P., Ratha, N.K. and Bhatta, D.: Isolation and identification of the insecticide endosulfan from biological materials by thin layer chromatographic technique. *J. Indian Chem. Soc.*, 120 (1994) 183-184; *C.A.*, 120 (1994) 238110g.
- 29c. *Phosphorus insecticides*
- 551 Kasatkina, A.A., Ivanova, L.P., Trifonova, V.N., Galeeva, A.G. and Gurkova, R.G.: (Use of 2-(4-nitrobenzyl)pyridine as a reagent for detection of toxic chemical residues in environmental samples). *Zh. Prikl. Khim. (S.-Peterburg)*, 65 (1992) 2830-2832; *C.A.*, 120 (1994) 207647h.
- 29d. *Carbamates*
- 552 McGinnis, S.C. and Sherma, J.: Determination of carbamate insecticides in water by C-18 solid phase extraction and quantitative HPTLC. *J. Liq. Chromatogr.*, 17 (1994) 151-156.
- 553 Patil, V.B. and Shingare, M.S.: Thin-layer chromatographic spray reagent for the screening of biological materials for the presence of carbaryl. *Analyst (London)*, 119 (1994) 415-416.
30. SYNTHETIC AND NATURAL DYES
- 30a. *Synthetic dyes*
- 554 Aginsky, B.V.: Forensic examination of "slightly soluble" ink pigments using thin-layer chromatography. *J. Forensic Sci.*, 38 (1993) 1131-1133; *C.A.*, 120 (1994) 127065b.
- 30b. *Chloroplast and other natural pigments*
- 555 Hayashi, T., Fukaya, Y., Tanaka, H., Kunimatsu, M., Hotta, I., Ikai, Y., Oka, H., Hayakawa, J. and Suzuki, R.: (A novel method for clean-up of coal tar dyes using a quaternary amine cartridge). *Shokuhin Eiseigaku Zasshi*, 34 (1993) 398-403; *C.A.*, 120 (1994) 132496x.
- 556 Johjima, T.: (Determination of *cis*- and *trans*-carotenes of tangerine and yellowish tangerine tomatoes by micro-thin-layer chromatography). *Engel Gakkai Zasshi*, 62 (1993) 567-574; *C.A.*, 120 (1994) 211707v.
- 557 Ozeki, N., Oka, H., Okai, Y., Ohno, T., Hayakawa, J., Hayashi, T., Aoyama, T., Kushibiki, Y. and Sato, T.: (Application of reversed-phase TLC to the analysis of coal tar dyes in food). *Shokuhin Eiseigaku Zasshi*, 34 (1993) 542-545; *C.A.*, 120 (1994) 215540h.
- 558 Rosas Romero, A.J., Herrera, J.C., Martinez de Aparicio, E. and Molina Cuevas, E.A.: Thin-layer chromatographic determination of β -carotene, cantaxanthin, lutein, violaxanthin and neoxanthin on Chromarods. *J. Chromatogr. A*, 667 (1994) 361-366.

32. DRUG ANALYSIS

32a. Drug analysis, general techniques

- 559 Chambuso, M., Kovar, K.-A. and Zimmermann, W.: The systematic separation of unknown mixtures of drugs by multi-column solid-phase extraction analogous to the Stas-Otto extraction process. *Pharmazie*, 49 (1994) 142-148.
- 560 Ojanperä, I. and Jänchen, P.: The application of instrumental qualitative thin-layer chromatography to drug screening. *LC-GC Int.*, 7 (1994) 164-170.
- 561 Schutz, H., Rochholz, G., Seno, H. and Weiler, G.: Sind dünn-schichtchromatographische Screeningmethoden noch zeitgemäss? Kritischer Vergleich mit immunochemischen Tests. *Pharmazie*, 49 (1994) 213-216.

See also 538.

32b. Antirheumatics and antiinflammatory drugs

- 562 Abdel, Rahman, A.A.; Stamm, A.E. Aboutalev, A.; Abdel, Ragman, S.I. and Samy, E.M.: Accelerated stability testing of ibuprofen-Eudragit RSPM sustained-release tablets using proton-NMR, HPLC and TLC. *Bull. Pharm. Sci., Assiut Univ.*, 16, No. 1 (1993) 1-11; *C.A.*, 120 (1994) 86308n.
- 563 Chowdary, K.P.R. and Babu, K.V.V. Suresh: Dissolution, bioavailability and ulcerogenic studies on solid dispersions of indomethacin in water-soluble cellulose polymers. *Drug Dev. Ind. Pharm.*, 20 (1994) 799-813; *C.A.*, 120 (1994) 226779u.
- 564 Shirke, P.P., Patel, M.K., Tamhane, V.A., Tirodkar, V.B. and Sethi, P.D.: Simultaneous estimation of ibuprofen and paracetamol in combined dosage formulation by high performance thin layer chromatography. *Indian Drugs*, 30 (1993) 653-654; *C.A.*, 120 (1994) 86602d.
- 565 Sun, S.W. and Fabre, H.: Practical approach for validating the TLC assay of an active ingredient in a pharmaceutical formulation. *J. Liq. Chromatogr.*, 17 (1994) 433-445.

32c. Autonomic and cardiovascular drugs

- 566 Fujii, T., Matsumoto, S., Amejima, H., Hatoyama, T., Nakao, M., Kagemoto, A., Tanaka, K. and Miyazaki, H.: Absorption, distribution, metabolism and excretion of [¹⁴C]ebastine after a single administration in rats. *Arzneim.-Forsch.*, 44 (1994) 527-538.
- 567 Fujii, T., Tanaka, K., Matsumoto, S., Hatoyama, T., Nomura, N., Tagawa, M. and Miyazaki, H.: Absorption, distribution, metabolism and excretion of [¹⁴C]ebastine after repeated oral administration in rats. *Arzneim.-Forsch.*, 44 (1994) 538-543.
- 568 Turcan, R.G., Hillbeck, D., Hartlay, T.E., Gilbert, P.J., Coe, R.A.J., Troke, J.A. and Vose, C.W.: Disposition of [¹⁴C]velnacrine maleate in rats, dogs, and humans. *Drug Metab. Disp.*, 21 (1993) 1037-1047.
- 569 Yabuki, M., Mine, T., Iba, K., Nakatsuka, I. and Yoshitake, A.: Pharmacokinetics of SM-10888 and its metabolites depending on their physicochemical properties. *Drug Metab. Disp.*, 22 (1994) 294-297.

32d. Central nervous system drugs

- 570 Aboul-Enein, H.Y. and Serignese, V.: Thin layer chromatographic (TLC) determination of phenytoin in pharmaceutical formulations and identification of its hydroxylated urinary metabolites. *Anal. Lett.*, 27 (1994) 723-729.
- 571 Andersin, R., Ovakainen, J. and Kaitia, S.: Photochemical decomposition of midazolam. III - Isolation and identification of products in aqueous solutions. *J. Pharm. Biomed. Anal.*, 12 (1994) 165-172.
- 572 Joseph-Charles, J. and Bertucat, M.: Separation, identification and assay of fluradoline hydrochloride in the presence of other tricyclic antidepressants or neuroleptics by spectral and thin-layer and liquid chromatographic methods. *Anal. Chim. Acta*, 284 (1993) 45-52.
- 573 Lehotay, J., Cizmarik, J., Pham Thi Viet Nga and Bednarikova, A.: Study of local anaesthetics, Part 121: Correlation between log *k'*, *C*, *R_F*, *R_M* and local anaesthesia for perhydroazepinyl ethyl esters of 2-, 3- and 4-alkoxyphenylcarbamic acids. *Pharmazie*, 49 (1994) 286.
- 574 Rochholz, G., Ahrens, B. and Schütz, H.: Modified screening procedure with fluorescence detection for flunitrazepam and its metabolites via acridine derivatives. *Arzneim.-Forsch.*, 44 (1994) 469-471.

See also 538.

32e. Chemotherapeutics (exc. cytostatics and antibiotics)

- 575 Abdelrahman, A.N., Karim, E.I.A. and Ibrahim, K.E.E.: Determination of chloroquine and its decomposition products in various brands of different dosage forms by liquid chromatography. *J. Pharm. Biomed. Anal.*, 12 (1994) 205-208.
- 576 Feng, C., Liu, Y., Luo, Y. and Kang, Z.: (TLC analysis of benzodiazepines and barbiturates in biological samples). *Junshi Yixue Kexueyuan Yuankan*, 17 (1993) 215-217; *C.A.*, 120 (1994) 68703h.
- 577 Kaniou, I., Zacariadis, G., Kalligas, G., Tsoukali, H. and Stratis, J.: Separation and determination of carbadox, nitrofurazone, nitrofurantoin, furazolidone, and furaltadone in their mixtures by thin layer and high performance liquid chromatography. *J. Liq. Chromatogr.*, 17 (1994) 1385-1398.

32g. Other drug categories

- 578 Cheng, Z.J. and Yang, Z.P.: (The degradation mechanism of penequine hydrochloride in aqueous solutions). *Yaoxue Xuebao*, 28 (1993) 633-638; *C.A.*, 120 (1994) 37921y.
- 579 Dittmann, J., Wenger, R.M., Kleinkauf, H. and Lawen, A.: Mechanism of cyclosporin A biosynthesis. Evidence for synthesis via a single linear undecapeptide precursor. *J. Biol. Chem.*, 269 (1994) 2841-2846.
- 580 Hu, X., Zhou, B., Luo, S., Cai, H. and Yin, W.: (Stability of liensinine injection). *Zhongguo Zhongyao Zazhi*, 18 (1993) 345-346; *C.A.*, 120 (1994) 62093n.

See also 564, 594.

32h. *Toxicological and forensic applications*

- 581 De Zeeuw, R.A., Franke, J.P., van Halem, M., Schaapman, S., Logawa, E. and Siregar, C.J.P.: Thin-layer chromatography under tropical conditions: impact of high temperatures and high humidities on screening systems for analytical toxicology. *J. Chromatogr. A*, 664 (1994) 263-270.
- 582 Moriya, F., Chan, K.-M., Noguchi, T.T. and Wu, P.Y.K.: Testing for drugs of abuse in meconium of newborn infants. *J. Anal. Toxicol.*, 18 (1994) 41-45.
- 32i. *Plant extracts*
- 583 Bader, G., Tuja, D., Wray, V. and Hiller, K.: New triterpenoid saponins from *Heteropappus altaicus* (Willd.) Noropokr. and *Heteropappus biennis* (Lbd.) Tamamsch. *Pharmazie*, 49 (1994) 209-212.
- 584 Doi, K., Ohmori, K., Kishi, M. and Horiguchi, Y.: (Analytical studies on crude drug component in pharmaceutical products. II. Component of epimedium grandiflorum Morr). *Kanagawa-ken Eisei Kenkyusho Kenkyu Hokoku*, 23 (1993) 35-38; *C.A.*, 120 (1994) 144302u.
- 585 Jin, Y., Zhang, J., He, L., Ma, B., Fan, G., Gu, H., Chen, D., Li, J., Liang, M. and Li, C.: (Analysis of D-mannitol in *Cordyceps sinensis* and *C. militaris*). *Baijiu Yike Daxue Xuebao*, 18 (1992) 47-49; *C.A.*, 119 (1993) 125308p.
- 586 Kamal-Eldin, A., Appelqvist, L.Å. and Yousif, G.: Lignan analysis in seed oils from four *Sesamum* species: comparison of different chromatographic methods. *J. Am. Oil Chem. Soc.*, 71 (1994) 141-147.
- 587 Khatoon, S., Mehrotra, S., Shome, U. and Mehrotra, B.N.: Analysis of commercial "Ratanjot" by TLC fluorescence fingerprinting. *Int. J. Pharmacogn.*, 31 (1993) 269-277; *C.A.*, 120 (1994) 200496z.
- 588 Li, P., Xu, G. and Xu, L.: (Determination of verticine and verticinone in Hupeh Fritillary (*Fritillaria hupehensis*)). *Zhongcaoyao*, 24 (1993) 579-580; *C.A.*, 120 (1994) 86570s.
- 589 Ma, L., Song, W. and Wu, F.: (Quality evaluation of roots of *Scutellaria* species as Chinese medicines). *Tianran Chanwu Yanjiu Yu Kaifa*, 5, No. 4 (1993) 9-14; *C.A.*, 120 (1994) 144297w.
- 590 Ma, S., Chen, D. and Zhao, S.: (Chemical components from the pericarps of Chinese drug Bayuezha). *Zhongcaoyao*, 24 (1993) 563-566; *C.A.*, 120 (1994) 143805y.
- 591 Ohmori, K., Doi, K., Kishi, M. and Horiguchi, Y.: (Analytical studies on erude drug component in pharmaceutical products. I. Component of Ginseng). *Kanagawa-ken Eisei Kenkyusho Kenkyu Hokoku*, 23 (1993) 30-34; *C.A.*, 120 (1994) 144303v.
- 592 Pal, S.K., Chattaraj, S.C. and Saha, B.P.: Standardization of tinctures by their cochemical and thin layer chromatographic characteristics. *Res. Ind.*, 38 (1993) 154-156; *C.A.*, 120 (1994) 86214d.
- 593 Schütt, H. and Hölzl, J.: Vergleichende Qualitätsuntersuchung von Johanniskraut-Fertigarzneimitteln unter Verwendung verschiedener quantitativer Bestimmungsmethoden. *Pharmazie*, 49 (1994) 206-209.
- 594 Wang, Y., Wang, S., Yu, L. and Ma, R.: (Quantitative analysis of tubeimoside I). *Zhongguo Zhongyao Zazhi*, 18 (1993) 358-360; *C.A.*, 120 (1994) 62384h.

- 595 Yang, X. and Luo, X.: (Quantitative determination of icariin in shorthorned epimedium (*Epimedium brevicornum*) and its preparation Xiongfengjiaonang by dual-wavelength HPTLC-scanning). *Zhongcaoyao*, 24 (1993) 520-521; *C.A.*, 120 (1994) 86563s.

See also 455, 523, 524.

33. CLINICO-CHEMICAL APPLICATIONS

33b. *Complex mixtures and profiling (single compounds by cross-reference only)*

See 454, 481, 539, 576.

34. FOOD ANALYSIS

34b. *Complex mixtures (single compounds by cross-reference only)*

See 528, 555, 556, 557, 599.

35. ENVIRONMENTAL ANALYSIS

35c. *Water pollution (complex mixtures; single compounds by cross-reference only)*

See 552.

36. SOME TECHNICAL PRODUCTS AND COMPLEX MIXTURES

36a. *Surfactants*

- 596 Yang, J., Zhang, L. and Li, X.: Quantitative determination of ethylene oxide oligomer distribution in ethoxylated alkylphenol carboxymethyl ether salts by thin-layer chromatography. *J. Am. Oil Chem. Soc.*, 71 (1994) 109-111.

36b. *Antioxidants and preservatives*

- 597 Cui, Y.: (Separation and determination of antioxidant by TLC, IR and NMR). *Shiyou Lianzhi*, 24 (1993) 59-63; *C.A.*, 120 (1994) 111235w.
- 598 Kanner, J., Frankel, E., Granit, R., German, B. and Kinsella, J.E.: Natural antioxidants in grapes and wines. *J. Agric. Food Chem.*, 42 (1994) 64-69.
- 599 Khan, S.H., Murawski, M.P. and Sherma, P.: Quantitative high performance thin layer chromatographic determination of organic acid preservatives in beverages. *J. Liq. Chromatogr.*, 17 (1994) 855-865.
- 600 Kronman, A.G., Ponomarev, A.N., Sitnikova, E.F., Kondakova, I.A. and Vodop'yanova, T.V.: (Analytical monitoring of low concentrations of additives in suspension polymerization of vinyl chloride). *Zavod. Lab.*, 59, No. 4 (1993) 29-30; *C.A.*, 120 (1994) 108522g.

36c. *Complex mixtures, technical products and unidentified compounds*

- 601 Aginsky, V.N.: Comparative examination of links by using instrumental thin-layer chromatography and microspectrophotometry. *J. Forensic Sci.*, 38 (1993) 1111-1130; *C.A.*, 120 (1994) 156044z.
- 602 Klein, E., Tsang, J.S. and Baker, M.: Non-instrumental techniques for characterization and identification of artist's acrylic paint. *Polym. Prepr. (Am. Chem. Soc., Div. Polym. Chem.)*, 33 (1992) 650-651; *C.A.*, 120 (1994) 194049n.
- 603 Paesold, D., van Drunen, A., Hoeglinger, M., Niedermayr, H., Cornely, R. and Koehler, M.: Optimization of rolling oils using modern analysis methods. *Stahl Eisen*, 114 (1994) 75-81; *C.A.*, 120 (1994) 138851f.
- 604 Weber, J.V., Krzton, A., Cebolla, V. and Swistek, M.: Chromatographic separation of heavy organic residues. TLC-FID: a useful monitoring tool. *Erdoel Kohle, Erdgas, Petrochem.*, 46 (1993) 411-414; *C.A.*, 120 (1994) 81152x.

See also 519, 554.

37. CELLS, CELLULAR PARTICLES AND SUPRAMOLECULAR STRUCTURES

See 488, 502.

38. INORGANIC COMPOUNDS

38a. *Cations*

- 605 Dauchy, X., Point-Gautier, M., Astruc, A. and Astruc, M.: Analytical methods for the speciation of selenium compounds: a review. *Fresenius J. Anal. Chem.*, 348 (1994) 792-805.

- 606 Janjic, T.J., Zivkovic, V. and Celap, M.B.: Linear relation between R_M values obtained in salting-out planar chromatography and mol % ammonium sulphate in aqueous solvent system. *Chromatographia*, 38 (1994) 355-358.
- 607 Janjic, T.J., Zivkovic, V. and Celap, M.B.: Salting-out thin-layer chromatography of transition metal complexes. IV: Effect of sorbent on the behaviour of mixed aminocarboxylato cobalt(II) complexes. *Chromatographia*, 38 (1994) 447-452.
- 608 Janjic, T.J., Zivkovic, V., Vuckovic, G. and Celap, M.B.: Salting-out thin-layer chromatography of transition metal complexes. III: investigation of the behaviour of mixed aminocarboxylato cobalt(III) complexes on cellulose. *Chromatographia*, 37 (1993) 534-538.
- 609 Volynets, M.P. and Stefantsev, L.A.: (Information-retrieval system IPS-TSKh - an automated database on thin-layer chromatography of inorganic ions). *Zavod. Lab.*, 59, No. 6 (1993) 11-13; *C.A.*, 120 (1994) 123587a.

See also 610.

39. RADIOACTIVE AND OTHER ISOTOPE COMPOUNDS

- 610 Djakov, A.A., Rostovsev, V.J., Betenekov, N.D. and Ipatova, E.G.: Development of high sensitive techniques for determination of concentrations of fission materials in water media using thin-layer inorganic sorbents and plastic track detectors. *Nucl. Tracks Radiat. Meas.*, 21 (1993) 349-351; *C.A.*, 120 (1994) 226316r.
- 611 Shigematsu, A.: Paste-mold method for study of radioactive concentration in tissues and quantitative analysis method for metabolic studies. *Yakubutsu Datai*, 8 (1993) 782-783; *C.A.*, 120 (1994) 128529t.
- 612 Wilson, M.E. and Hung, J.C.: Microwave preparation of and one-strip paper chromatography for technetium 99m teboroxime. *Am. J. Hosp. Pharm.*, 50 (1993) 2376-2379; *C.A.*, 120 (1994) 227098h.

Electrophoresis

1. REVIEWS AND BOOKS

- 1299 Guzman, N.A. (Editor): *Capillary Electrophoresis Technology*. In: *Chromatogr. Sci. Ser.*, Vol. 64, 1993, Dekker, New York, 857 p.; C.A., 120 (1994) 174466t.
- 1300 Ricci, M.C. and Cross, R.F.: Capillary electrophoresis: how good is this separation technique and how does it compare with HPLC. *Chem. Aust.*, 60 (1993) 540-546; C.A., 120 (1994) 123834d - a review without refs.
- 1301 Wätzig, H. and Dette, C.: Capillary electrophoresis (CE) - a review. Strategies for method development and applications related to pharmaceutical and biological sciences. *Pharmazie*, 49 (1994) 83-96 - a review with 378 refs.
- 1302 Weiss, J., Jansen, D. and Saweet, A.: (Ion chromatography and capillary electrophoresis for ion analysis). *LaborPraxis*, 17 (1993) 36-43; C.A., 120 (1994) 94067y - a review with 11 refs.

See also 1331, 1356, 1363, 1367, 1379, 1380, 1383, 1384, 1388, 1389, 1390, 1397, 1400, 1412, 1418, 1420, 1423, 1428, 1434, 1468, 1513, 1608, 1619, 1823, 1871, 1935, 1938, 1941, 1956, 1959, 1977.

2. FUNDAMENTALS, THEORY AND GENERAL

2a. General

- 1303 Harrison, D.J., Fan, Z., Seiler, K. and Fluri, K.: Miniaturized chemical analysis systems and their fabrication: an alternative to chemical sensors. *Proc.-Electrochem. Soc.*, 93-97(Proceedings of the Symposium on Chemical Sensors II, 1993) (1993) 546-552; C.A., 120 (1994) 128708a.
- 1304 Orzechowski, A. and Blaszczyk, T.: (Measurement of size distribution and electrophoretic mobility of particles (macromolecules) in colloidal systems by means of the Zetasizer-4 system). *Polimery (Warsaw)*, 38 (1993) 247-252; C.A., 120 (1994) 165421x.
- 1305 Rhodes, P.H., Snyder, R.S., Roberts, G.O. and Baygents, J.C.: Electrohydrodynamic effects in continuous flow electrophoresis. *Appl. Theor. Electrophor.*, 2 (1991) 87-91; C.A., 120 (1994) 63278p.
- 1306 Stevenson, R.: Back to basics: I'll never doubt the power of pK_a again. *Am. Lab. (Shelton)*, 25 (1993) 52GG-HH; C.A., 120 (1994) 186651n.
- 1307 Stockham, T.G. and Ives, J.T.: Methods and apparatus for sharpening signal peaks in analysis of chromatographic migration patterns. *U.S. US 5,273,632* (Cl. 204-180.1; BO1D57/02), 28 Dec. 1993, Appl. 978,915, 19 Nov. 1992; 20 pp.; C.A., 120 (1994) 212022e.
- 1308 Viovy, J.L. and Duke, T.: Solid friction and polymer relaxation in gel electrophoresis. *Science (Washington)*, 264 (1994) 112-113; C.A., 120 (1994) 239488y.

- 1309 Zimm, B.H.: Diffusion and electrophoresis of a polyelectrolyte in an irregular matrix. *Polym. Prepr. (Am. Chem. Soc., Div. Polym. Chem.)*, 34 (1993) 1054; C.A., 120 (1994) 192834r.

See also 1351, 1359, 1714, 1931.

2b. Thermodynamics and theoretical relationships

- 1310 Chen, X.: A new model and theory of gel electrophoresis of DNA. Avail. *Univ. Microfilms Int.*, Order No. DA9311782, 1992, 135 pp.; C.A., 120 (1994) 157941b.
- 1311 Dukhin, S.S. and Razilov, I.A.: Nonlinearity of electrophoresis due to the mobility of adsorbed ions. The near-to-threshold magnitude of the external field). *Kolloidn. Zh.*, 55 (1993) 32-38; C.A., 120 (1994) 201343j.
- 1312 Dukhin, S.S., Shchuyurtz, Yu. Pfranger, Yu. and Razilov, I.A.: (Nonlinearity of electrophoresis due to the mobility of adsorbed ions. Threshold value of the external field). *Kolloidn. Zh.*, 55 (1993) 21-31; C.A., 120 (1994) 201342h.
- 1313 Friedl, W. and Kenndler, E.: Limitation of the optimisation of the resolution by the buffer pH in capillary zone electrophoresis. *Fresenius J. Anal. Chem.*, 348 (1994) 576-582.
- 1314 Li, H. and Shen, Z.: (Studies on electroosmotic flow in high-performance capillary electrophoresis). *Gaodeng Xuexiao Huaxue Xuebao*, 14 (1993) 1528-1530; C.A., 120 (1994) 207468a.
- 1315 Muthukumar, M.: Theory of electrophoretic mobility of polyelectrolyte chains. *Macromol. Theory Simul.*, 3 (1994) 61-71; C.A., 120 (1994) 108196d.

See also 1394, 1395, 1396, 1403, 1832.

2d. Measurement of physico-chemical and related values

- 1316 Nakatani, M., Shibukawa, A. and Nakagawa, T.: Sodium dodecyl sulfate-linear polyacrylamide-bonded capillary electrophoresis for the molecular weight determination of proteins. *Anal. Sci.*, 10 (1994) 1-4; C.A., 120 (1994) 186557m.
- 1317 Yang, H., Zhang, P. and Xu, K.: (Improvement of method for determining polypeptide molecular weight by SDS-PAGE). *Yaowu Fenxi Zazhi*, 13 (1993) 325-327; C.A., 120 (1994) 49379r.

See also 1350, 1539, 1895.

3. GENERAL TECHNIQUES

3a. Apparatus and accessories

- 1318 Majors, R.E.: New chromatography columns and accessories at the 1994 Pittsburgh Conference, Part I. *LC-GC Int.*, 7 (1994) 190-203.

- 1319 Manz, A.: Apparatus and method for electrophoretic separation. *Eur. Pat. Appl. EP 544,969* (Cl. G01N27/447), 09 Jun. 1993, Appl. 91/810,952, 06 Dec. 1991; 14 p.; C.A., 120 (1994) 123763e.
- 1320 Manz, A., Verpoorte, E., Effenhauser, C.S., Burggraf, N., Raymond, D.E. and Widmer, M.H.: Planar chip technology for capillary electrophoresis. *Fresenius J. Anal. Chem.*, 348 (1994) 567-571.
- 1321 Schmid, P.: Apparatus and method for an automated electrophoresis system and transfer system. *U.S. US 5,279,721* (Cl. 204-182.8; GO1N27/26), 18 Jan. 1994, Appl. 52,079, 22 Apr. 1993; 13 pp.; C.A., 120 (1994) 129037t.
- 1322 Togusari, T., Irisawa, K. and Kambara, H.: Apparatus for gel electrophoresis. *U.S. US 5,294,323* (Cl. 204-299R; GO1N27/26), 15 Mar. 1994, Appl. 69,841, 01 Jun. 1993; 8 pp.; C.A., 120 (1994) 265297e.
- See also 1362, 1378, 1382, 1392, 1393, 1425, 1426, 1437.
- 3b. Detectors and detection procedures**
- 1323 Chen, D.Y., Adelhelm, K., Cheng, X.L. and Dovichi, N.J.: A simple laser-induced fluorescence detector for sulforhodamine 101 in a capillary electrophoresis system: detection limits of 10 yostomoles or six molecules. *Analyst (London)*, 119 (1994) 349-352.
- 1324 Hara, T., Kayama, S., Nishida, H. and Nakajima, R.: Simple apparatus for online chemiluminescence detection of proteins separated by capillary zone electrophoresis. *Anal. Sci.*, 10 (1994) 223-225; C.A., 120 (1994) 186373y.
- 1325 Hernandez, L., Joshi, N., Verdeguer, P. and Guzman, N.A.: Laser-induced fluorescence detection for capillary electrophoresis: a powerful analytical tool for the separation and detection of trace amounts of analytes. *Chromatogr. Sci. Ser.*, 64(Capillary Electrophoresis Technology) (1993) 605-614; C.A., 120 (1994) 72690u.
- 1326 Herranen, M. and Vuento, M.: Staining of immunoblots by immunoassay. *Anal. Biochem.*, 218 (1994) 468-470.
- 1327 Kambara, H., Anazawa, T., Nagai, K., Machida, H. and Nasu, H.: Fluorescence detection electrophoresis apparatus. *U.S. US 5,290,419* (Cl. 204-299R; GO1N27/26), 01 Mar. 1994, JP Appl. 92/94,568, 14 Apr. 1992; 9 pp.; C.A., 120 (1994) 239634t.
- 1328 Koutney, L.B.: Novel detection schemes and automated image analysis algorithms for planar chromatography and gel electrophoresis. *Report, IS-T-1630; Order No. DE93001219*, 1992, 211 pp.; C.A., 120 (1994) 72873f.
- 1329 Moring, S.E., Pairaud, C., Albin, M., Locke, S., Tribault, P. and Tindall, G.W.: Enhancement of UV detection sensitivity for capillary electrophoresis with a high-sensitivity optical cell. *Am. Lab. (Shelton)*, 25 (1993) 32-39; C.A., 120 (1994) 123817a.
- 1330 Nuske, J. and Grus, F.H.: (Modern densitometry). *BioTec (Wuerzburg)*, 5 (1993) 33-35; C.A., 120 (1994) 72683u.
- 1331 Saz, J.M. and Díez-Masa, J.C.: Thermo-optical spectroscopy: new and sensitive schemes for detection in capillary separation techniques. *J. Liq. Chromatogr.*, 17 (1994) 499-520 - a review with 42 refs.
- 1332 Schwedt, G. and Harms, J.: (Capillary electrophoresis with scanning UV-visible detector). *LaborPraxis*, 17 (1993) 51-52; C.A., 120 (1994) 152353b.
- 1333 Wang, T.: Absorption detection and capillary modification in capillary electrophoresis. Avail. *Univ. Microfilms Int.*, Order No. DA9231446, 1992, 205 p.; C.A., 120 (1994) 207645f.
- 1334 Wang, T.: Capillary multireflective cell for use in capillary zone electrophoresis. *U.S. US 5,273,633* (Cl. 204-180.1; GO1N27/26), 28 Dec. 1993, Appl. 911,607, 10 Jul. 1992; 17 pp.; C.A., 120 (1994) 129017m.
- See also 1416, 1423, 1439, 1499, 1721, 1765, 1827.
- 3c. Stabilization media for electrophoresis**
- 1335 Burlatsky, S. and Deutch, J.: Solid friction and polymer relaxation in gel electrophoresis. Reply to comments. *Science (Washington)*, 264 (1994) 113; C.A., 120 (1994) 265117w.
- 1336 Chu, D.Y. and Mardis, E.R.: Sample well insert with wedge-shaped profile for ultra-thin slab gels in electrophoresis. *U.S. US 5,284,565* (Cl. 204-299R; GO1N27/26), 08 Feb. 1994, Appl. 995,495, 23 Dec. 1992; 9 pp.; C.A., 120 (1994) 158146h.
- 1337 Gombocz, E.A. and Rammler, D.H.: Use of multiple short gels for elongated separation pattern in gel electrophoresis. *PCT Int. Appl. WO 93 23,746* (Cl. GO1N27/26), 25 Nov. 1993, US Appl. 883,111, 14 May 1992, 17 pp.; C.A., 120 (1994) 72896r.
- 1338 Gombocz, E.A. and Roth, A.T.: Gel electrophoresis system including optical stage, sample applicator and sample retriever. *U.S. US 5,275,710* (Cl. 204-299R; GO1N27/26), 04 Jan. 1994, US Appl. 522,325, 14 May 1990; 23 pp.; C.A., 120 (1994) 129010d.
- 1339 Griess, G.A., Guiseley, K.B. and Serwer, P.: The relationship of agarose gel structure to the sieving of spheres during agarose gel electrophoresis. *Biophys. J.*, 65 (1993) 138-148; C.A., 120 (1994) 72752r.
- 1340 Hamase, K., Nakamura, H. and Nakamura, T.: Curdlan gel, evaluated as a polysaccharide support by electrophoresis of substituted benzenes. *Anal. Sci.*, 9 (1993) 799-802; C.A., 120 (1994) 123842e.
- 1341 Liu, P.Z., Malik, A., Kuchar, M.C.J., Vorkink, W.P. and Lee, M.L.: Polymeric hollow fibers for capillary electrophoresis. *J. Microcolumn Sep.*, 5 (1993) 243-253; C.A., 120 (1994) 211847r.
- 1342 Menchen, S.M. and Winnik, M.: Viscous electrophoresis hydrophilic-hydrophobic polymer medium and method. *U.S. US 5,290,418* (Cl. 204-299R; BO1D39/00), 01 Mar. 1994, Appl. 950,863, 24 Sep. 1992; 31 pp.; C.A., 120 (1994) 265330k.
- 1343 Pukl, M., Prosek, M. and Kaiser, R.E.: Planar electrochromatography. Part I. Planar electrochromatography on non-wetted thin layers. *Chromatographia*, 38 (1994) 83-87.
- 1344 Righetti, P.G.: New formulations for polyacrylamide matrices in electrokinetic and chromatographic methodologies. *PCT Int. Appl. WO 93 11,174* (Cl. C08F20/58), 10 Jun. 1993, IT Appl. 91/Mi3271, 06 Dec. 1991; 9 pp.; C.A., 120 (1994) 72893n.
- 1345 Stellwagen, J. and Stellwagen, N.C.: Transient electric birefringence of agarose gels. I. Unidirectional electric fields. *Biopolymers*, 34 (1994) 187-201; C.A., 120 (1994) 157955j.
- 1346 Viljoen, C.D., Wingfield, B.D. and Wingfield, M.J.: Agar, an alternative to agarose in analytical gel electrophoresis. *Biotechnol. Tech.*, 7 (1993) 723-726; C.A., 120 (1994) 49375m.
- See also 1318, 1366, 1386, 1401, 1408, 1424, 1879.

3d. Quantitative analysis

See 1384, 1746.

3e. Preparative scale electrophoresis

1347 Lim, Y.P. and Hixson, D.C.: Preparative gel electrophoresis device and method. *PCT Int. Appl.* WO 93 25,896 (Cl. GO1N27/26), 23 Dec. 1993, US Appl. 899,460, 16 Jun. 1992; 25 pp.; C.A., 120 (1994) 186748z.

See also 1374, 1420, 1634, 1964.

3f. Programmed voltage and buffer gradients

1348 Celentano, F.C. and Gianazza, E.: Strategies for pH gradients optimization. *Appl. Theor. Electrophor.*, 3 (1993) 191-201; C.A., 120 (1994) 174407z.

1349 Powell, A.C. and Sepaniak, M.J.: Analytical instrumentation and computational methods for generating and optimizing solvent-gradients in micellar electrokinetic capillary chromatography. *Anal. Instrum. (N.Y.)*, 21 (1993) 25-41; C.A., 120 (1994) 182077g.

4. SPECIAL TECHNIQUES

4a. Automation

See 1328.

4b. Computerization and modelling

1350 Bello, M.S. and Polezhaev, V.I.: Hydrodynamics gravitational sensitivity, and transport phenomena, in continuous flow electrophoresis. *AIAA J.*, 31 (1993) 997-998; C.A., 120 (1994) 176748h.

1351 Locke, B.R. and Arce, P.: Modeling electrophoretic transport of polyelectrolytes in beds of nonporous spheres. *Sep. Technol.*, 3 (1993) 111-120; C.A., 120 (1994) 186535c.

1352 Malenkevitz, J.M.: Density functional theory of microphase separation in block copolymers and Monte Carlo simulations of polyelectrolyte electrophoresis. Avail. *Univ. Microfilms Int.*, Order No. DA9229635, 1992, 225 p.; C.A., 120 (1994) 192781w.

1353 Pelton, R., Miller, P., McPhee, W. and Rajaram, S.: Strategies for improving electrophoresis data from the Coulter DELSA. *Colloids Surf. A*, 80 (1993) 181-189; C.A., 120 (1994) 117746k.

1354 Trubuil, A.: Analysis of one-dimensional electrophoregrams. *Comput. Appl. Biosci.*, 9 (1993) 451-458; C.A., 120 (1994) 72765x.

1355 Vindevogel, J. and Sandra, P.: Interactive removal of spikes from digitized electrophoregrams. *LC-GC Int.*, 7 (1994) 276-281.

See also 1328, 1372, 1838, 1902.

4c. Combination with other physicochemical techniques, (MS, IR etc.)

1356 Bruins, A.P.: Atmospheric-pressure-ionization mass spectrometry. II. Applications in pharmacy, biochemistry and general chemistry. *TrAC*, 13 (1994) 81-90 - a review with 39 refs.

1357 Fenselau, C. and Vestling, M.: Laser desorption MS interfaced with gel electrophoresis to characterize proteins. *Am. Lab. (Shelton)*, 25 (1993) 72-78; C.A., 120 (1994) 186652p.

1358 Tomlinson, A.J., Benson, L.M. and Naylor, S.: Nonaqueous solvents in the on-line capillary electrophoresis-mass spectrometry analysis of drug metabolites. *J. High Resolut. Chromatogr.*, 17 (1994) 175-177.

See also 1377, 1411, 1413, 1414, 1417, 1499, 1534, 1536, 1939.

4d. Affinity electrophoresis

See 1457, 1754.

4e. Capillary zone electrophoresis and electrokinetic chromatography

1359 Ahuja, E.S. and Foley, J.P.: A retention index for micellar electrokinetic chromatography. *Analyst (London)*, 119 (1994) 353-360.

1360 Aiken, J.H. and Huie, C.W.: Effect of 1-alkanols on separation performance in micellar electrokinetic capillary chromatography. *J. Microcolumn Sep.*, 5 (1993) 95-99; C.A., 120 (1994) 207659p.

1361 Arai, T., Ichinose, M., Kuroda, H., Nimura, N. and Kinoshita, T.: Chiral separation by capillary affinity zone electrophoresis using an albumin-containing support electrolyte. *Anal. Biochem.*, 217 (1994) 7-11.

1362 Bauer, H., Gauglitz, G., Hofmann, W., Nahm, W., Mettler, M., Nagel, J. and Wolf, B.: Capillary electrophoresis apparatus. *Ger. Offen.* DE 4,218,721 (Cl. GO1N27/447), 09 Dec. 1993, Appl. 06 Jun. 1992; 10 pp.; C.A., 120 (1994) 129013g.

1363 Bereuter, T.L.: Enantioseparation by capillary electrophoresis. *LC-GC Int.*, 7 (1994) 78-93 - a review with 111 refs.

1364 Birrell, E.C., Camilleri, P. and Okafo, G.N.: Phytic acid can greatly enhance resolution in capillary electrophoresis. *J. Chem. Soc., Chem. Commun.*, (1994) 43-44; C.A., 120 (1994) 182107s.

1365 Brehm, G.: (Capillary electrophoresis - the philosophers' stone or an alternative analytical method?). *Proc. MoBBEL* 1991, 6 (1992) 102-107; C.A., 120 (1994) 157938f.

1366 Bruin, G.J.M., Wang, T., Xu, X., Kraak, J.C. and Poppe, H.: Preparation of polyacrylamide gel-filled capillaries by photopolymerization for capillary electrophoresis. *J. Microcolumn Sep.*, 4 (1992) 439-448; C.A., 120 (1994) 239460h.

1367 Cannon, J. and Rickman, M.: Enhancing the sensitivity of capillary electrophoresis. *Am. Biotechnol. Lab.*, 11 (1993) 12; C.A., 120 (1994) 100815j - a review with 1 ref.

1368 Demorest, D.M., Werner, W.E., Wiktorowicz, J.E., Wenz, H.M. and Oaks, F.L.: Capillary electrophoresis molecular weight separation of biomolecules using a polymer-containing solution. *PCT Int. Appl.* WO 93 22,665 (Cl. GO1N27/26), 11 Nov. 1993, US Appl. 877,956, 01 May 1992; 75 pp.; C.A., 120 (1994) 101275v.

- 1369 Dill, R. and Siebert, C.J.: High-performance capillary electrophoresis capillary cartridge for uncoiled capillaries with liquid coolant. *U.S. US 5,269,901* (Cl. 204-299R; GO1N27/26), 14 Dec. 1993, Appl. 975,763, 13 Nov. 1992; 9 pp.; *C.A.*, 120 (1994) 72884k.
- 1370 Dougherty, A.M. and Schure, M.R.: Covalent surface modification for capillary electrophoresis: characterization and effect of nonionic bondings on separations in capillary electrophoresis. *Chromatogr. Sci. Ser.*, 64(Capillary Electrophoresis Technology) (1993) 435-472; *C.A.*, 120 (1994) 72743p.
- 1371 El Rassi, Z. and Nashabeh, W.: Capillary zone electrophoresis of biopolymers with hydrophilic fused-silica capillaries. *Chromatogr. Sci. Ser.*, 64(Capillary Electrophoresis Technology) (1993) 383-434; *C.A.*, 120 (1994) 72570e.
- 1372 Ermakov, S.V. and Righetti, P.G.: Computer simulation for capillary zone electrophoresis. *J. Chromatogr. A*, 667 (1994) 257-270.
- 1373 Fujimoto, C.: Electrophoretic electrode, method of and system for capillary electrophoresis using the electrophoretic electrode and fraction collector assembled into the capillary electrophoresis system. *Eur. Pat. Appl. EP 576,361* (Cl. GO1N27/447), 29 Dec. 1993, JP Appl. 92/208,382, 26 Jun. 1992; 16 pp.; *C.A.*, 120 (1994) 158137f.
- 1374 Fujimoto, C. and Jinno, K.: Micropreparative capillary electrophoresis. *Chromatogr. Sci. Ser.*, 64(Capillary Electrophoresis Technology) (1993) 509-523; *C.A.*, 120 (1994) 72745r.
- 1375 Fujimoto, C., Sawada, H. and Jinno, K.: Capillary electrophoresis in non-crosslinked polyacrylamide columns containing hydrophobic additives. *J. High Resolut. Chromatogr.*, 17 (1994) 107-108.
- 1376 Gareil, P., Roldan-Assad, R. and Lelievre, F.: (Capillary electrophoresis: a rapid maturity). *Analisis*, 21 (1993) M35-M38; *C.A.*, 120 (1994) 128881b.
- 1377 Goodlett, D.R., Wahl, J.H., Udseth, H.R. and Smith, R.D.: Reduced elution speed detection for capillary electrophoresis/mass spectrometry. *J. Microcolumn Sep.*, 5 (1993) 57-62; *C.A.*, 120 (1994) 101218d.
- 1378 Guttman, A.: Capillary electrophoresis apparatus using time-varying field strength. *PCT Int. Appl. WO 93 25,899* (Cl. GO1N27/447), 23 Dec. 1993, US Appl. 900,223, 17 Jun. 1992; 26 pp.; *C.A.*, 120 (1994) 186749a.
- 1379 Guttman, A.: Capillary polyacrylamide gel electrophoresis. *Chromatogr. Sci. Ser.*, 64(Capillary Electrophoresis Technology) (1993) 715-729; *C.A.*, 120 (1994) 72574j - a review with 39 refs.
- 1380 Guzman, N.A., Gonzalez, C.L., Trebilcock, M.A., Hernandez, L., Berck, C.M. and Advis, J.P.: The use of capillary electrophoresis in clinical diagnosis. *Chromatogr. Sci. Ser.*, 65(Capillary Electrophoresis Technology) (1993) 643-672; *C.A.*, 120 (1994) 72573h - a review with 53 refs.
- 1381 Hayes, M.A., Gilman, S.D. and Ewing, A.G.: Prospects for the use of capillary electrophoresis in neuro science. *Chromatogr. Sci. Ser.*, 64(Capillary Electrophoresis Technology) (1993) 753-793; *C.A.*, 120 (1994) 72747t.
- 1382 Hogan, B.L., Lunte, M.S., Stobaugh, J.F. and Lunte, C.E.: On-line coupling of *in vivo* microdialysis sampling with capillary electrophoresis. *Anal. Chem.*, 66 (1994) 596-602.
- 1383 Honda, S.: (Capillary electrophoresis. Principle and characteristic features of capillary electrophoresis). *Rinsho Kagaku (Nippon Rinsho Kagakkai)*, 22 (1993) 204-208; *C.A.*, 120 (1994) 211667g - a review with 4 refs.
- 1384 Hoyt, A.M., Jr.: Quantitative analysis with capillary zone electrophoresis. *Chromatogr. Sci. Ser.*, 64 (1993) 705-714; *C.A.*, 120 (1994) 68390d - a review with 41 refs.
- 1385 Hoyt, A.M., Jr., Beale, S.C., Larmann, J.P., Jr. and Jorgenson, J.W.: Preparation and evaluation of an online preconcentrator for capillary electrophoresis. *J. Microcolumn Sep.*, 5 (1993) 325-330; *C.A.*, 120 (1994) 186371w.
- 1386 Huang, M. and Lee, M.L.: Hydrogel polymer coatings for capillary electrophoretic separation of proteins. *J. Microcolumn Sep.*, 4 (1993) 491-496; *C.A.*, 120 (1994) 157937e.
- 1387 Huang, M., Vorking, W.P. and Lee, M.L.: Evaluation of surface-bonded polyethylene glycol and polyethylene imine in capillary electrophoresis. *J. Microcolumn Sep.*, 4 (1992) 135-143; *C.A.*, 120 (1994) 123833c.
- 1388 Jandik, P. and Bonn, G.: *Capillary Electrophoresis of Small Molecules and Ions*. VCH, New York, 1993, 298 pp.; *C.A.*, 120 (1994) 49549w.
- 1389 Karger, B.L.: Capillary electrophoresis. Current and future directions. *Am. Lab. (Shelton)*, 25 (1993) 23-26; *C.A.*, 120 (1994) 182056z - a review with 4 refs.
- 1390 Karger, B.L. and Foret, F.: Capillary electrophoresis: introduction and assessment. *Chromatogr. Sci. Ser.*, 64(Capillary Electrophoresis Technology) (1993) 3-64; *C.A.*, 120 (1994) 72568k - a review with 194 refs.
- 1391 Katsuta, S. and Saitoh, K.: Selectivity control in micellar electrokinetic chromatography by modification of micellar surface with cyclohexanol. *Anal. Lett.*, 27 (1994) 743-749.
- 1392 Klein, G.L. and Katzmann, S.P.: Method and apparatus for displaying capillary electrophoresis data. *Eur. Pat. Appl. EP 581,412* (Cl. GO1N27/447), 02 Feb. 1994, US Appl. 916,307, 17 Jul. 1992; 30 pp.; *C.A.*, 120 (1994) 186752w.
- 1393 Klein, G.L. and Miller, G.A.: Multi-channel capillary electrophoresis system. *Eur. Pat. Appl. EP 581,413* (Cl. GO1N27/447), 02 Feb. 1994, US Appl. 916,308, 17 Jul. 1992; 21 pp.; *C.A.*, 120 (1994) 186753x.
- 1394 Knox, J.H. and McCormack, K.A.: Temperature effects in capillary electrophoresis. 1: Internal capillary temperature and effect upon performance. *Chromatographia*, 38 (1994) 207-214.
- 1395 Knox, J.H. and McCormack, K.A.: Temperature effects in capillary electrophoresis. 2: Some theoretical calculations and predictions. *Chromatographia*, 38 (1994) 215-221.
- 1396 Knox, J.H. and McCormack, K.A.: Volume expansion and loss of sample due to initial self-heating in capillary electroseparation (CES) system. *Chromatographia*, 38 (1994) 279-282.
- 1397 Kohr, J. and Engelhardt, H.: Capillary electrophoresis with coated capillaries. *Chromatogr. Sci. Ser.*, 64(Capillary Electrophoresis Technology) (1993) 357-381; *C.A.*, 120 (1994) 72569m - a review with 24 refs.
- 1398 Lee, T.T.: Instrumental and theoretical development of capillary electrophoresis and its application to biological systems. *Report*, IS-T-1629; Order No. DE93001220, 1992, 254 pp.; *C.A.*, 120 (1994) 128858z.
- 1399 Lin, B., Shi, W., Hu, L., Yang, G., Huang, J. and Zhang, N.: (Effect of the buffer pH in capillary zone electrophoresis). *Gaodeng Xuexiao Huaxue Xuebao*, 14 (1993) 1229-1231; *C.A.*, 120 (1994) 72751q.

- 1400 Looser, R.: (Capillary electrophoresis). *Schweiz. Lab.-Z.*, 51 (1994) 19-24; *C.A.*, 120 (1994) 182061x - a review with refs.
- 1401 Malik, A., Zhao, A. and Lee, M.L.: Simple method for the preparation of highly efficient polymer-coated capillary electrophoresis columns. *J. Microcolumn Sep.*, 5 (1993) 119-125; *C.A.*, 120 (1994) 157943d.
- 1402 Mizuno, M. and Tochigi, K.: Capillary electrophoresis column for high resolution protein analysis. *Jpn. Kokai Tokkyo Koho JP 05,264,502* [93,264,502] (Cl. GO1N27/447), 12 Oct. 1993, Appl. 92/61,819, 18 Mar. 1993; 4 pp.; *C.A.*, 120 (1994) 101235g.
- 1403 Muijselaar, P.G.H.M., Claessens, H.A. and Cramers, C.A.: Application of the retention index concept in micellar electrokinetic capillary chromatography. *Anal. Chem.*, 66 (1994) 635-644.
- 1404 Otsuka, K. and Terabe, S.: Chiral separation by capillary electrophoresis and electrokinetic chromatography. *Chromatogr. Sci. Ser.*, 64(Capillary Electrophoresis Technology) (1993) 617-629; *C.A.*, 120 (1994) 72572g.
- 1405 Palmer, C.D. and McNair, H.M.: Novel pseudostationary phase for micellar electrokinetic capillary chromatography. *J. Microcolumn Sep.*, 4 (1993) 509-514; *C.A.*, 120 (1994) 207655j.
- 1406 Regnier, F.E. and Wu, D.: Chemical derivatization of fused silica capillaries. *Chromatogr. Sci. Ser.*, 64(Capillary Electrophoresis Technology) (1993) 287-309; *C.A.*, 120 (1994) 72741m.
- 1407 Schomburg, G.: Technology of separation capillaries for capillary zone electrophoresis and capillary gel electrophoresis: the chemistry of surface modification and formation of gels. *Chromatogr. Sci. Ser.*, 64(Capillary Electrophoresis Technology) (1993) 311-355; *C.A.*, 120 (1994) 72742n.
- 1408 Schomburg, G. and Gilges, M.: Deactivation of the inner surfaces of capillaries by thermally fixing polar polymers for capillary electrophoresis. *Eur. Pat. Appl. EP 587,156* (Cl. BO1D57/02), 16 Mar. 1994, DE Appl. 4,230,403, 11 Sep. 1992; 15 pp.; *C.A.*, 120 (1994) 265332n.
- 1409 Shohat, D. and Grushka, E.: Use of calixarenes to modify selectivities in capillary electrophoresis. *Anal. Chem.*, 66 (1994) 747-750.
- 1410 Sjoeholm, E., Nilvebrant, N.O. and Colmsjoe, A.: Characterization of dissolved kraft lignin by capillary zone electrophoresis. *J. Wood Chem. Technol.*, 13 (1993) 529-544; *C.A.*, 120 (1994) 56943y.
- 1411 Smith, R.D. and Udseth, H.R.: Mass-spectrometric detection for capillary electrophoresis. *Chromatogr. Sci. Ser.*, 64(Capillary Electrophoresis Technology) (1993) 525-567; *C.A.*, 120 (1994) 72571f.
- 1412 Stevenson, R.: HPCE to the rescue. *Am. Lab. (Shelton)*, 25 (1993) 52-56; *C.A.*, 120 (1994) 207620u - a review with no refs.
- 1413 Tetler, L.W. and Cooper, P.: Evaluation of capillary electrophoresis: Continuous flow fast atom bombardment interfaces. *Anal. Proc.*, 30 (1993) 397-398; *C.A.*, 120 (1994) 72833t.
- 1414 Tetler, L.W., Cooper, P.A. and Carr, C.M.: The application of capillary electrophoresis/mass spectrometry using negative-ion electrospray ionization to areas of importance in the textile industry. *Rapid Commun. Mass Spectrom.*, 8 (1994) 179-182; *C.A.*, 120 (1994) 166660m.
- 1415 Timerbaev, A., Semenova, O. and Bonn, G.: Metal ion capillary zone electrophoresis with direct UV detection: comparison of different migration modes for negatively charged chelates. *Chromatographia*, 38 (1994) 497-500.
- 1416 Timerbaev, A.R., Buchberger, W., Semenova, O.P. and Bonn, G.K.: Metal ion capillary zone electrophoresis using direct UV detection. *Spec. Publ.-R. Soc. Chem.*, 122 (1993) 111-121; *C.A.*, 120 (1994) 152398v.
- 1417 Tomer, K.B.: Capillary zone electrophoresis-mass spectrometry: continuous flow fast-atom bombardment and electrospray ionization. *Chromatogr. Sci. Ser.*, 64(Capillary Electrophoresis Technology) (1993) 569-586; *C.A.*, 120 (1994) 100818n.
- 1418 Trissl, H.W.: Capillary electrophoresis: Pioneering new approaches for biomolecular analysis. *Trends Biochem. Sci.*, 18 (1993) 409-414; *C.A.*, 120 (1994) 72565g - a review with 33 refs.
- 1419 Tsai, P. and Lee, C.S.: Direct control of electroosmotic flow in capillary electrophoresis. *Chromatogr. Sci. Ser.*, 64(Capillary Electrophoresis Technology) (1993) 475-488; *C.A.*, 120 (1994) 72744q.
- 1420 Tsuda, T.: Semipreparative capillary electrophoresis and its advantages. *Chromatogr. Sci. Ser.*, 64 (1993) 489-508; *C.A.*, 120 (1994) 68388j - a review with 21 refs.
- 1421 Vindevogel, J., Szucz, R. and Sandra, P.: Reproducibility of migration times and propagation of error in micellar electrokinetic chromatography. *J. Microcolumn Sep.*, 4 (1992) 399-404; *C.A.*, 120 (1994) 174348f.
- 1422 Yaun-Dongxing, T.Q. and Yang, P.: (High performance capillary electrophoresis). *Xiamen Daxue Xuebao, Ziran Kexueban*, 32 (1993) 336-340; *C.A.*, 120 (1994) 207466y.
- 1423 Yeung, E.S.: Optical detection schemes for capillary electrophoresis. *Chromatogr. Sci. Ser.*, 64 (1993) 587-603; *C.A.*, 120 (1994) 68389k - a review with 42 refs.
- 1424 Young, J.E., Bateman, M. and Lux, J.A.: Method of making an electrophoretic capillary tube with resistive coating of carbon black and polymer. *U.S. US 5,290,587* (Cl. 427-122; C23C4/00), 01 Mar. 1994, Appl. 914,153, 14 Jul. 1992; 11 pp.; *C.A.*, 120 (1994) 239635u.
- 1425 Zheng, P., Fu, X., Lu, J. and Zu, A.: (Capillary electrophoresis apparatus and experimental research). *Zhejiang Daxue Xuebao, Ziran Kexueban*, 27 (1993) 85-91; *C.A.*, 120 (1994) 152641u.
- 1426 Zimmermann, H.P.: Apparatus for performing capillary electrophoresis. *Eur. Pat. Appl. EP 581,983* (Cl. GO1N27/447), 09 Feb. 1994, Appl. 92/113,244, 04 Aug. 1992; 16 pp.; *C.A.*, 120 (1994) 186755z.

See also 1299, 1300, 1301, 1302, 1306, 1313, 1314, 1316, 1318, 1320, 1324, 1325, 1329, 1331, 1332, 1333, 1334, 1341, 1349, 1355, 1356, 1358, 1427, 1429, 1431, 1432, 1433, 1434, 1440, 1441, 1442, 1444, 1447, 1448, 1450, 1459, 1462, 1468, 1470, 1472, 1487, 1493, 1494, 1496, 1498, 1499, 1505, 1506, 1507, 1508, 1509, 1510, 1511, 1513, 1522, 1531, 1533, 1537, 1538, 1539, 1575, 1608, 1612, 1654, 1657, 1660, 1665, 1672, 1790, 1792, 1793, 1794, 1813, 1823, 1824, 1840, 1870, 1871, 1873, 1875, 1876, 1877, 1882, 1885, 1899, 1913, 1916, 1917, 1919, 1920, 1921, 1922, 1926, 1929, 1933, 1934, 1935, 1936, 1938, 1939, 1940, 1941, 1943, 1944, 1946, 1947, 1948, 1949, 1951, 1952, 1953, 1954, 1955, 1956, 1957, 1966, 1967, 1972, 1973, 1974, 1975, 1976.

4f. Isotachophoresis

See 1385, 1469, 1968.

4g. *Enantiomers, separation*

- 1427 Belder, D. and Schomburg, G.: Chiral separations of basic and acidic compounds in modified capillaries using cyclodextrin-modified capillary zone electrophoresis. *J. Chromatogr. A*, 666 (1994) 351-365.
- 1428 Burns, D.T.: Cyclodextrins as versatile chiral recognition reagents for use in a variety of optical and separative analyses. *J. Pharm. Biomed. Anal.*, 12 (1994) 1-3 - a review with 39 refs.
- 1429 Fanali, S.: Use of cyclodextrins in capillary electrophoresis. *Chromatogr. Sci. Ser.*, 64(Capillary Electrophoresis Technology) (1993) 731-752; *C.A.*, 120 (1994) 72746s.
- 1430 Huynh, T.K.X., Ossicini, L. and Polcaro, C.: Chiral separations on cellulose in electrophoresis. *J. Chromatogr. A*, 663 (1994) 264-266.
- 1431 Ishihama, Y., Oda, Y., Asakawa, N., Yoshida, Y. and Sato, T.: Optical resolution by electrokinetic chromatography using ovomucoid as a pseudo-stationary phase. *J. Chromatogr. A*, 666 (1994) 193-201.
- 1432 Kuhn, R., Steinmetz, C., Bereuter, T., Haas, P. and Erni, F.: Enantiomeric separations in capillary zone electrophoresis using a chiral crown ether. *J. Chromatogr. A*, 666 (1994) 367-373.
- 1433 Li, S. and Lloyd, D.K.: Packed-capillary electrochromatographic separation of the enantiomers of neutral and anionic compounds using β -cyclodextrin as a chiral selector. Effect of operating parameters and comparison with free-solution capillary electrophoresis. *J. Chromatogr. A*, 666 (1994) 321-335.
- 1434 Terabe, S., Otsuka, K. and Nishi, H.: Separation of enantiomers by capillary electrophoretic techniques. *J. Chromatogr. A*, 666 (1994) 295-319 - a review with 109 refs.

See also 1361, 1363, 1404, 1498, 1920, 1926, 1934, 1936, 1940, 1943, 1944, 1946, 1951.

4h. *Two dimensional electrophoresis*

- 1435 Lemkin, P.F., Myrick, J.E. and Upton, K.M.: Splitting merged spots in two-dimensional polyacrylamide gel electrophoresis gel images. *Appl. Theor. Electrophor.*, 3 (1993) 163-172; *C.A.*, 120 (1994) 101035s.
- 1436 Ono, K. and Kawashima, S.: Multiple labels for two-dimensional electrophoresis for protein or DNA identification. *Jpn. Kokai Tokkyo Koho JP 05,281,191 [93,281,191]* (Cl. GO1N27/447), 29 Oct. 1993, *Appl.* 91/125,209, 25 Apr. 1991; 5 pp.; *C.A.*, 120 (1994) 101276w.

See also 1528, 1530, 1532 1548, 1587, 1588, 1590, 1591, 1595, 1603, 1611, 1638, 1643, 1647, 1664, 1669, 1710, 1733, 1861, 1902, 1961.

4i. *Other special techniques*

- 1437 Paladini, A.A., Weber, G. and Erijman, L.: Analysis of dissociation and unfolding of oligomeric proteins using a flat bed gel electrophoresis at high pressure. *Anal. Biochem.*, 218 (1994) 364-369.
- 1438 Schmidt, J.L. and Cheh, H.Y.: Free-flow electrophoresis method. *U.S. US 5,277,774* (Cl. 204-180.1; GO1N27/26), 11 Jan. 1994, *US Appl.* 721,303, 26 Jun. 1991; 10 pp.; *C.A.*, 120 (1994) 129038u.

- 1439 Wu, J. and Pawliszyn, J.: Dual detection for capillary isoelectric focusing with refractive index gradient and absorption imaging detectors. *Anal. Chem.*, 66 (1994) 867-873.

See also 1415.

5. HYDROCARBONS AND HALOGEN DERIVATIVES

5b. *Cyclic hydrocarbons, fullerenes*

See 1340.

7. PHENOLS

- 1440 Chao, Y.-C. and Whang, C.-W.: Capillary zone electrophoresis of eleven priority phenols with indirect fluorescence detection. *J. Chromatogr. A*, 663 (1994) 229-237.

See also 1410.

8. SUBSTANCES CONTAINING HETEROCYCLIC OXYGEN

8a. *Flavonoids*

- 1441 Delgado, C., Tomas-Barberán, F.A., Talou, T. and Gaset, A.: Capillary electrophoresis as an alternative to HPLC for determination of honey flavonoids. *Chromatographia*, 38 (1994) 71-78.

9. OXO COMPOUNDS, ETHERS, EPOXIDES AND QUINONES

See 1410.

10. CARBOHYDRATES

10a. *Mono and oligosaccharides. Structural studies*

- 1442 Chang, L., Yao, R., Chen, Y. and Kong, J.: (Determination of monosaccharides by capillary zone electrophoresis). *Fenxi Huaxue*, 22 (1994) 1-5; *C.A.*, 120 (1994) 207672n.
- 1443 Hu, G.-f. and Vallee, B.L.: A gel retardation assay for the interaction of proteins and carbohydrates by fluorophore-assisted carbohydrate electrophoresis. *Anal. Biochem.*, 218 (1994) 185-191.
- 1444 Huber, C., Grill, E., Oefner, P. and Bobleter, O.: Capillary electrophoretic determination of the component monosaccharides in hemicelluloses. *Fresenius J. Anal. Chem.*, 348 (1994) 825-831.
- 1445 Kakehi, K. and Honda, S.: Analysis of carbohydrates in glycoproteins by high-performance liquid chromatography and high-performance capillary electrophoresis. *Appl. Biochem. Biotechnol.*, 43 (1993) 55-71; *C.A.*, 120 (1994) 128942x.

- 1446 Sato, T., Furukawa, K., Greenwalt, D.E. and Kobata, A.: Most bovine milk fat globule membrane glycoproteins contain asparagine-linked sugar chains with GalNAc β 1 \rightarrow 4GlcNAc groups. *J. Biochem. (Tokyo)*, 114 (1993) 890-900.
- 1447 Yao, R., Chen, Y. and Chaing, L.: Capillary zone electrophoresis of monosaccharides. *Chin. Chem. Lett.*, 4 (1993) 343-346; *C.A.*, 120 (1994) 77527f.
- See also 1459.
- 10b. *Polysaccharides, mucopolysaccharides, lipopolysaccharides*
- 1448 Ip, C.C.Y., Miller, W.J., Silberklang, M., Mark, G.E., Elis, R.W., Huang, L., Glushka, J., van Halbeek, H., Zhu, J. and Alhadeff, J.A.: Structural characterization of the N-glycans of a humanized anti-CD18 murine immunoglobulin G. *Arch. Biochem. Biophys.*, 308 (1994) 387-399.
- 1449 Moller, H.J., Larsen, F.S., Ingemann-Hansen, T. and Poulsen, J.H.: ELISA for the core protein of the cartilage large aggregating proteoglycan, aggrecan: comparison with the concentrations of immunogenic keratan sulphate in synovial fluid, serum and urine. *Clin. Chim. Acta*, 225 (1994) 43-55.
- 1450 Novotny, M.V. and Sudor, J.: Pulsed-field separation of polysaccharides in capillaries. *U.S. US 5,292,416 (Cl. 204-182.8; C25B7/00)*, 08 Mar. 1994, Appl. 975,314, 13 Nov. 1992; 11 pp.; *C.A.*, 120 (1994) 265331m.
- 1451 Pavao, M.S.G., Rodrigues, M.A. and Mourão, P.A.S.: Acidic polysaccharides of the ascidian *Styela plicata*. Biosynthetic studies on the sulfated L-galactans of the tunic, and preliminary characterization of a dermatan sulfate-like polymer in body tissues. *Biochim. Biophys. Acta*, 1199 (1994) 229-237.
- 1452 Renn, D.W., Santos, G.A., Dumont, L.E., Parent, C.A., Stanley, N.F., Stancioff, D.J. and Guiseley, K.B.: β -Carrageenan: isolation and characterization. *Carbohydr. Polym.*, 22 (1993) 247-252; *C.A.*, 120 (1994) 157982r.
- 1453 Volpi, N.: "Fast moving" and "slow moving" heparins, dermatan sulfate, and chondroitin sulfate: qualitative and quantitative analysis by agarose-gel electrophoresis. *Carbohydr. Res.*, 247 (1993) 263-278; *C.A.*, 120 (1994) 49369n.
- 1454 Volpi, N.: Fractionation of heparin, dermatan sulfate, and chondroitin sulfate by sequential precipitation: a method to purify a single glycosaminoglycan species from a mixture. *Anal. Biochem.*, 218 (1994) 382-391.
- 10c. *Glycoproteins and their constituents*
- 1455 Beninati, S., Senger, D.R., Cordella-Miele, E., Mukherjee, A.B., Chackalaparampil, I., Shanmugam, V., Singh, K. and Mukherjee, B.B.: Osteopontin: its transglutaminase-catalyzed posttranslational modifications and cross-linking to fibronectin. *J. Biochem. (Tokyo)*, 115 (1994) 675-682.
- 1456 Chou, C.-F., Riopel, C.L. and Omary, M.B.: Identification of a keratin-associated protein that localizes to a membrane compartment. *Biochem. J.*, 298 (1994) 457-463.
- 1457 Heegaard, P.M.H. and Boeg-Hansen, T.C.: Concanavalin A affinity immunoelectrophoresis of slowly migrating glycoproteins by chemical charge modification. *Appl. Theor. Electrophor.*, 3 (1993) 213-217; *C.A.*, 120 (1994) 157999b.
- 1458 Ito, T., Hiraiwa, J., Nishioka, J., Iino, M. and Suzuki, K.: Characterization of functionally important regions of tissue factor by using monoclonal antibodies. *J. Biochem. (Tokyo)*, 114 (1993) 691-696.
- 1459 Kuhn, R., Frei, R. and Christen, M.: Use of capillary affinity electrophoresis for the determination of lectin-sugar interactions. *Anal. Biochem.*, 218 (1994) 131-135.
- 1460 Lindblom, A., Marsh, T., Fauser, C., Engel, J. and Paulsson, M.: Characterization of native laminin from bovine kidney and comparison with other laminin variants. *Eur. J. Biochem.*, 219 (1994) 383-392.
- 1461 Liu, W. and Jiang, R.-q.: The fluorescent labeling of glycoproteins on sodium dodecyl sulfate-polyacrylamide gel. *Bioorg. Chem.*, 22 (1994) 29-35; *C.A.*, 120 (1994) 265006j.
- 1462 Maeda, Y., Funakushi, K., Sato, T., Suzuki, S., Nakazawa, H., Ito, K., Yonekawa, O. and Kanno, T.: (Capillary zone electrophoresis for analysis of Tamm-Horsfall glycoprotein and albumin in human urine). *Shizuoka-ken Eisei Kankyo Senta Hokoku*, 35 (1992) 15-19; *C.A.*, 120 (1994) 72732j.
- 1463 Masumi, A., Akamatsu, Y. and Kitagawa, T.: Alteration by transforming growth factor- β 1 of asparagine-linked sugar chains in glucose transporter protein in Swiss 3T3 cells. *Biochim. Biophys. Acta*, 1221 (1994) 330-338.
- 1464 Poruchynsky, M.S. and Ling, V.: Detection of oligomeric and monomeric forms of P-glycoprotein in multidrug resistant cells. *Biochemistry*, 33 (1994) 4163-4174.
- 1465 Spillmann, D. and Finne, J.: Identification of a major poly-N-acetylglucosamine-containing cell-surface glycoprotein of mouse teratocarcinoma cells. Appearance on cells induced to primitive endoderm but not parietal endoderm differentiation. *Eur. J. Biochem.*, 220 (1994) 385-394.
- 1466 Tseung, H.B., Lai, C.K., Lin, D.T., Lin, Y.L., Chen, C.W., Lian, C., and Chen, W.F.: Purification of envelope glycoproteins of the Newcastle disease virus. *Zhonghua Minguo Shouyi Xuehui Zazhi*, 19 (1993) 11-18; *C.A.*, 120 (1994) 239458p.
- 1467 Yoshida, M., Mizuno, Y., Nonaka, I. and Ozawa, E.: A dystrophin-associated glycoprotein A3a (one of 43DAG doublets), is retained in duchenne muscular dystrophy muscle. *J. Biochem. (Tokyo)*, 114 (1993) 634-639.
- See also 1445, 1446, 1573, 1625, 1694.
11. ORGANIC ACIDS AND LIPIDS
- 11a. *Organic acids and simple esters*
- 1468 Hiraoka, A. and Sasaki, H.: (Capillary electrophoresis. Capillary electrophoretic analyses of organic acids in cerebrospinal fluid). *Rinsho Kagaku (Nippon Rinsho Kagakukai)*, 22 (1993) 220-224; *C.A.*, 120 (1994) 239289j - a review with 26 refs.
- 1469 Karovičová, J., Drdák, M., Polonský, J. and Rajniaková, A.: Dynamics of production of organic acids during lactic fermentation of vegetable juice. *J. Chromatogr. A*, 665 (1994) 55-58.
- 1470 Liu, Y.-M. and Sheu, S.-J.: Separation of aromatic acids by reversed electroosmotic flow capillary electrophoresis. *J. Chromatogr. A*, 663 (1994) 239-242.

11c. Lipids and their constituents

- 1471 Ingvarsdén, L., Michaelsen, S. and Sørensen, H.: Analysis of individual phospholipids by high-performance capillary electrophoresis. *J. Am. Oil Chem. Soc.*, 71 (1994) 183-188.
- 1472 Yu, Z., Chang, L. and Su, T.: Capillary electrophoretic separation of gangliosides using cyclodextrin as buffer additive. *Chin. Chem. Lett.*, 4 (1993) 827-828; *C.A.*, 120 (1994) 211832g.

See also 1633.

11d. Lipoproteins and their constituents

- 1473 Aviram, M. and Rosenblat, M.: Macrophage-mediated oxidation of extracellular low density lipoprotein requires an initial binding of the lipoprotein to its receptor. *J. Lipid Res.*, 35 (1994) 385-398.
- 1474 Bozhko, G.K. and Kulabukhov, V.M.: (Redistribution of rabbit blood serum lipoproteins following a single injection of cholesterol). *Biokhimiya (Moscow)*, 58 (1993) 1594-1603.
- 1475 De Silva, H.V., Lauer, S.J., Wang, J., Simonet, W.S., Weisgraber, K.H., Mahley, R.W. and Taylor, J.M.: Overexpression of human apolipoprotein C-III in transgenic mice results in an accumulation of apolipoprotein B48 remnants that is corrected by excess apolipoprotein E. *J. Biol. Chem.*, 269 (1994) 2324-2335.
- 1476 Fazio, S., Horie, Y., Simonet, W.S., Weisgraber, K.H., Taylor, J.M. and Rall, S.C., Jr.: Altered lipoprotein metabolism in transgenic mice expressing low levels of a human receptor-binding-defective apolipoprotein E variant. *J. Lipid Res.*, 35 (1994) 408-416.
- 1477 Himber, J., Missano, B. and Kuhl, H.: Lack of effect on the low density lipoprotein receptor in hamsters treated with 17 α -ethinyl estradiol. *Biochim. Biophys. Acta*, 1211 (1994) 359-363.
- 1478 Huby, T., Doucet, C., Dieplinger, H., Chapman, J. and Thillet, J.: Structural domains of apolipoprotein(a) and its interaction with apolipoprotein B-100 in the lipoprotein(a) particle. *Biochemistry*, 33 (1994) 3335-3341.
- 1479 Ikeda, H.: (The subfractions of the cholesterol in high-density lipoprotein determined by 5% disk polyacrylamide-gel electrophoresis). *Seibutsu Butsuri Kagaku*, 37 (1993) 381-388; *C.A.*, 120 (1994) 128863x.
- 1480 James, R.W., Ruiz, J., Blanché, H., Pometta, D., Passa, P. and Froguel, P.: Phenotyping is an accurate means of analysing the principal apolipoprotein E isoforms. *Clin. Chim. Acta*, 225 (1994) 77-82.
- 1481 Kataoka, S., Paidi, M. and Howard, B.V.: Simplified isoelectric focusing/immunoblotting determination of apoprotein E phenotype. *Clin. Chem. (Washington)*, 40 (1994) 11-13.
- 1482 Leonhardt, W., Pietzsch, J. and Nitzsche, S.: Very-fast ultracentrifugation of human plasma lipoproteins: influence of the centrifugal field on lipoprotein composition. *Clin. Chim. Acta*, 224 (1994) 21-32.
- 1483 Marques-Vidal, P., Azéma, C., Collet, X., Vieu, C., Chap, H. and Perret, B.: Hepatic lipase promotes the uptake of HDL esterified cholesterol by the perfused rat liver: a study using reconstituted HDL particles of defined phospholipid composition. *J. Lipid Res.*, 35 (1994) 373-384.
- 1484 Seman, L.J., Jenner, J.L., McNamara, J.R. and Schaefer, E.J.: Quantification of lipoprotein(a) in plasma by assaying cholesterol in lectin-bound plasma fraction. *Clin. Chem. (Washington)*, 40 (1994) 400-403.
- 1485 Shibusawa, Y., Miwa, N., Hirashima, T. and Matsumoto, U.: Separation of low density and very low density lipoproteins from human serum by hydroxyapatite chromatography. *J. Liq. Chromatogr.*, 17 (1994) 1203-1217.
- 1486 Talmud, P.J., Krul, E.S., Pessah, M., Gay, G., Schonfeld, G., Humphries, S.E. and Infante, R.: Donor splice mutation generates a lipid-associated apolipoprotein B 27.6 in a patient with homozygous hypobetalipoproteinemia. *J. Lipid Res.*, 35 (1994) 468-477.
- 1487 Weinmann, W., Maier, C., Baumeister, K., Przybylski, M., Parker, C.E. and Tomer, K.B.: Isolation of hydrophobic lipoproteins in organic solvents by pressure-assisted capillary electrophoresis for subsequent mass spectrometric characterization. *J. Chromatogr. A*, 664 (1994) 271-275.
- 1488 Wiegel, D., Richter, O. and Arnold, K.: Partitioning of chemically modified low-density lipoprotein in aqueous polymer two-phase systems. *J. Chromatogr. A*, 668 (1994) 107-116.
- 1489 Young, S.G., Koduri, R.K., Austin, R.K., Bonnet, D.J., Smith, R.S. and Curtiss, L.K.: Definition of a nonlinear conformation epitope for the apolipoprotein B-100-specific monoclonal antibody, MB 47. *J. Lipid Res.*, 35 (1994) 399-407.
- 1490 Zanelli, T., Catapano, A.L., Averna, M.R., Barbagallo, C.M., Liotta, A., Giardina, F.C. and Notarbartolo, A.: A new case of apo C-II deficiency with a nonsense mutation in the apo C-II gene. *Clin. Chim. Acta*, 224 (1994) 111-118.

See also 1612, 1826.

13. STEROIDS

13d. Sterols

- 1491 Berg, A., Baumstark, M.W., Grathwohl, D., Spielberger, B. and Keul, J.: (Practical experience in the enzymic determination of lipoprotein cholesterol using the Helena REP (rapid electrophoresis) system). *Labor-Med.*, 16 (1993) 359-366; *C.A.*, 120 (1994) 72761t.

13e. Bile acids and alcohols

- 1492 Mitsuhashi, Y., Nishimura, M., Hirose, M., Inomata, M., Hisaki, H. and Ueta, N.: (Electrophoresis of bile acids in human bile, Cellulose acetate membrane and agarose gel membrane). *Seibutsu Butsuri Kagaku*, 37 (1993) 325-328; *C.A.*, 120 (1994) 49372h.

13f. Ecdysones and other insect steroid hormones

- 1493 Yasuda, A., Ikeda, M. and Naya, Y.: Analysis of hemolymph ecdysteroids in the crayfish *Procambarus clarkii* by high performance capillary electrophoresis. *Nippon Suisan Gakkaishi*, 59 (1993) 1793-1799; *C.A.*, 120 (1994) 268408c.

16. NITRO AND NITROSO COMPOUNDS

- 1494 McCord, B.R. and Hargadon, K.A.: Explosive analysis by capillary electrophoresis. In: Yinon, J. (Editor), *Adv. Anal. Detect. Explos., Proc. Int. Symp., 4th 1992*, Kluwer, Dordrecht, 1993, pp. 133-144; *C.A.*, 120 (1994) 210154a.

17. AMINES, AMIDES AND RELATED NITROGEN COMPOUNDS

17a. Amines and polyamines

- 1495 Hawel III, L., Thandrawinata, R.R., Fukumoto, G.H. and Byus, C.V.: Biosynthesis and selective export of 1,5-diaminopentane (cadaverine) in mycoplasma-free cultured mammalian cells. *J. Biol. Chem.*, 269 (1994) 7412-7418.

17d. Other amine derivatives and amides (excl. peptides)

See 1975.

18. AMINO ACIDS AND PEPTIDES; CHEMICAL STRUCTURE OF PROTEINS

18a. Amino acids and their derivatives

- 1496 Heber, M. and Meyer, H.E.: Non-radioactive determination of phosphoamino acids by capillary electrophoresis. *NATO ASI Ser., Ser. H*, 76 (Tyrosine Phosphorylation/Dephosphorylation and Downstream Signalling) (1993) 241-246; *C.A.*, 120 (1994) 72737q.
- 1497 Le Phuc Thuy and Nyhan, W.L.: High-voltage electrophoresis of amino acids in urine containing ampicillin. *Clin. Biochem.*, 26 (1993) 389-390; *C.A.*, 120 (1994) 186553g.
- 1498 Werner, A., Nassauer, T., Kiechle, P. and Erni, F.: Chiral separation by capillary zone electrophoresis of an optically active drug and amino acids by host-guest complexation with cyclodextrins. *J. Chromatogr. A*, 666 (1994) 375-379.

See also 1375, 1502, 1672, 1951.

18b. Peptides, peptidic and proteinous hormones, growth factors

- 1499 Advis, J.P., Iqbal, K., Malick, A.W. and Guzman, N.A.: Capillary electrophoresis coupled to fluorescence detection for the determination of multiple neuropeptides. In: De Pablo, F., Scanes, C.G. and Weintraub, B.D. (Editors), *Handb. Endocr. Res. Techn.*, Academic, San Diego, 1993, pp. 127-144; *C.A.*, 120 (1994) 183124g.
- 1500 Andersson, M., Östman, A., Westmermark, B. and Heldin, C.-H.: Characterization of the retention motif in the C-terminal part of the long splice form of platelet-derived growth factor A-chain. *J. Biol. Chem.*, 269 (1994) 926-930.
- 1501 Brattsand, G., Marklund, U., Nylander, K., Ross, G. and Gullberg, M.: Cell-cycle-regulated phosphorylation of oncoprotein 18 on Ser 16, Ser 25 and Ser 38. *Eur. J. Biochem.*, 220 (1994) 359-368.

- 1502 Chambers, T.C., Pohl, J., Glass, D.B. and Kuo, J.F.: Phosphorylation by protein kinase C and cyclic AMP-dependent protein kinase of synthetic peptides derived from the linker region of human P-glycoprotein. *Biochem. J.*, 299 (1994) 309-315.

- 1503 Deng, A.Y., Martin, L.L., Balwierczak, J.L. and Jeng, A.Y.: Purification and characterization of an endothelin degradation enzyme from rat kidney. *J. Biochem. (Tokyo)*, 115 (1994) 120-125.

- 1504 Fukuoka, E., Suzuki, K., Yoshinari, S., Okunishi, H. and Miyazaki, M.: (Simultaneous analysis of bradykinin and its metabolites by capillary electrophoresis). *Bunseki Kagaku*, 43 (1994) 131-137; *C.A.*, 120 (1994) 183150n.

- 1505 Gruebler, G., Yildiz, E., Zimmerman, H., Mihelic, M., Echner, H., Stoeva, S., Spyropoulos, S. and Voelter, W.: Capillary electrophoresis, an attractive analytical tool in the laboratories of peptide chemists. In: Yanaihara, N. (Editor), *Pept. Chem. 1992, Proc. Jpn. Symp., 2nd 1992*, ESCOM, Leiden, 1993, pp. 208-211; *C.A.*, 120 (1994) 72677v.

- 1506 Janini, G.M., Issaq, H.J. and Lukszo, J.: Determination of the purity of synthetic peptides by capillary electrophoresis, high performance liquid chromatography, and laser desorption mass spectrometry. *J. High Resolut. Chromatogr.*, 17 (1994) 102-103.

- 1507 Kist, T.B.L., Termignoni, C. and Grieneisen, H.P.H.: Capillary zone electrophoresis separation of kinins using a novel laser fluorescence detector. *Braz. J. Med. Biol. Res.*, 27 (1994) 11-19; *C.A.*, 120 (1994) 261462g.

- 1508 Krizek, M.: (Determination of some polypeptides using the method of capillary zone electrophoresis). *Chem. Listy*, 87 (1993) 863-868; *C.A.*, 120 (1994) 211834j.

- 1509 Lee, H.G. and Desiderio, D.M.: Capillary zone electrophoresis of synthetic opioid and tachykinin peptides. *J. Chromatogr. A*, 667 (1994) 271-283.

- 1510 Lee, H.G. and Desiderio, D.M.: Capillary zone electrophoresis of two synthetic neuropeptides: examination of detectability and resolution as a function of peptide concentration and buffer concentration. *J. Chromatogr. B*, 655 (1994) 9-19.

- 1511 Lim, B.C. and Sim, M.K.: Determination of angiotensins by capillary electrophoresis. *J. Chromatogr. B*, 655 (1994) 127-131.

- 1512 Mashima, K., Okajima, Y., Usui, J., Shimizu, T. and Kimura, K.: Alteration in protein-tyrosine phosphatase of rat epithelial cells by RSV-transformation: application of phospho-tyrosyl glutamine synthetase to the study of protein-tyrosine phosphatase. *J. Biochem. (Tokyo)*, 115 (1994) 333-337.

- 1513 Perrett, D., Birch, A. and Ross, G.: Capillary electrophoresis for peptides, including neuropeptides. *Biochem. Soc. Trans.*, 22 (1994) 127-131; *C.A.*, 120 (1994) 208670x - a review with 28 refs.

- 1514 Wang, L., Chen, N. and Zhang, Y.: (Separation of hydrophilic small peptides (Gly)_n by high-performance capillary zone electrophoresis). *Huaxue Xuebao*, 51 (1993) 1174-1178; *C.A.*, 120 (1994) 186554h.

See also 1535, 1686.

18c. Elucidation of structure of proteins and enzymes

- 1515 Bonafé, N., Mathieu, M., Kassab, R. and Chaussepied, P.: Troponomyosin inhibits the glutaraldehyde-induced cross-link between the central 48-kDa fragment of myosin head and segment 48-67 in actin subdomain 2. *Biochemistry*, 33 (1994) 2594-2603.
- 1516 Hess, D., Covey, T.C., Winz, R., Brownsey, R.W. and Aebersold, R.: Analytical and micropreparative peptide mapping by high performance liquid chromatography/electrospray mass spectrometry of proteins purified by gel electrophoresis. *Protein Sci.*, 2 (1993) 1342-1351; *C.A.*, 120 (1994) 128984n.
- 1517 Houen, G.: Microsequencing of polypeptides after electrophoresis in modified Laemmli gels and recovery of polypeptides from polyvinyl difluoride membranes. *Methods Mol. Cell. Biol.*, 4 (1993) 105-114; *C.A.*, 120 (1994) 158085n.
- 1518 Le Goff, P., Montano, M.M., Schodin, D.J. and Katzenellenbogen, B.S.: Phosphorylation of the human estrogen receptor. Identification of hormone-regulated sites and examination of their influence on transcriptional activity. *J. Biol. Chem.*, 269 (1994) 4458-4466.
- 1519 Malyan, A.N., Girault, G. and Berger, G.: Examination of catalytic activity of the β subunit isolated from chloroplast coupling factor 1. *Biochim. Biophys. Acta*, 1184 (1994) 202-206.
- 1520 Miyatake, N., Satake, K., Kamo, M. and Tsugita, A.: Specific chemical cleavage of asparaginyl and glycyl-glycine bonds in peptides and proteins by anhydrous hydrazine vapor. *J. Biochem. (Tokyo)*, 115 (1994) 208-212.
- 1521 Odoul, M., Bagrel, D., Peyrieras, N., Bodaud, H. and Siest, G.: Glycosylation of gamma-glutamyltransferase is modified by ethanol in H5-6 hepatoma cell line. *Clin. Chim. Acta*, 225 (1994) 1-15.
- 1522 Strömqvist, M.: Peptide mapping using combinations of size-exclusion chromatography, reversed-phase chromatography and capillary electrophoresis. *J. Chromatogr. A*, 667 (1994) 304-310.
- 1523 Watabe, S., Kohno, H., Kouyama, H., Hiroi, T., Yago, N. and Nakazawa, T.: Purification and characterization of a substrate protein for mitochondrial ATP-dependent protease in bovine adrenal cortex. *J. Biochem. (Tokyo)*, 115 (1994) 648-654.
- 1524 Wenegieme, E.F., Babitch, J.A. and Naren, A.P.: Cation binding to chicken gizzard α -actinin. *Biochim. Biophys. Acta*, 1205 (1994) 308-316.
- 1525 Yamamoto, N., Akino, A. and Takano, T.: Purification and specificity of a cell-wall-associated proteinase from *Lactobacillus helveticus* CP790. *J. Biochem. (Tokyo)*, 114 (1993) 740-745.
- 1526 Yano, S., Fukunaga, K., Ushio, Y. and Miyamoto, E.: Activation of Ca^{2+} /calmodulin-dependent protein kinase II and phosphorylation of intermediate filament proteins by stimulation of glutamate receptors in cultured rat cortical astrocytes. *J. Biol. Chem.*, 269 (1994) 5428-5439.

See also 1538, 1610, 1661, 1706, 1744, 1853.

19. PROTEINS

19a. General techniques

- 1527 Fernandez, J., Andrews, L. and Mische, S.M.: An improved procedure for enzymatic digestion of polyvinylidene difluoride-bound proteins for internal sequence analysis. *Anal. Biochem.*, 218 (1994) 112-117.
- 1528 Goldman, D. and O'Brien, S.J.: Two-dimensional protein electrophoresis in phylogenetic studies. *Methods Enzymol.*, 224 (Molecular Evolution: Producing the Biochemical Data) (1993) 113-121; *C.A.*, 120 (1994) 239456m.
- 1529 Klarskov, K. and Roepstorff, P.: Plasma desorption mass spectrometry of proteins transferred from gels after sodium dodecyl sulfate-polyacrylamide gel electrophoresis. *Biol. Mass Spectrom.*, 22 (1993) 433-440; *C.A.*, 120 (1994) 49275d.
- 1530 Latham, K.E., Garrels, J.I. and Solter, D.: Two-dimensional gel analysis of protein synthesis. *Methods Enzymol.*, 225 (Guide to Techniques in Mouse Development) (1993) 473-489; *C.A.*, 120 (1994) 265099s.
- 1531 Nakatani, M., Shibukawa, A. and Nakagawa, T.: Sodium dodecyl sulfate-polyacrylamide solution-filled capillary electrophoresis of proteins using stable linear polyacrylamide-coated capillary. *Biol. Pharm. Bull.*, 16 (1993) 1185-1188; *C.A.*, 120 (1994) 265118x.
- 1532 Schägger, H., Cramer, W.A. and von Jagow, G.: Analysis of molecular masses and oligomeric states of protein complexes by blue native electrophoresis and isolation of membrane protein complexes by two-dimensional native electrophoresis. *Anal. Biochem.*, 217 (1994) 220-230.
- 1533 Strege, M.A. and Lagu, A.L.: Capillary electrophoretic analysis of the urea unfolding of bovine serum albumin. *Int. Lab.*, 24, No. 4 (1994) P16-P18.
- 1534 Strupat, K., Karas, M., Hillenkamp, F., Eckerskorn, C. and Lottspeich, F.: Matrix-assisted laser desorption ionization mass spectrometry of proteins electroblotted after polyacrylamide gel electrophoresis. *Anal. Chem.*, 66 (1994) 464-470.
- 1535 Udagawa, A., Nozu, K., Shiratori, M. and Ehata, Y.: Gel electrophoresis method for protein, peptide, and nucleic acid determination. *Jpn. Kokai Tokkyo Koho JP 06 34,602 [94 34,602]* (Cl. GO1N27/447), 10 Feb. 1994, Appl. 92/102,186, 20 Jul. 1992; 5 pp.; *C.A.*, 120 (1994) 265329s.
- 1536 Vestling, M.M. and Fenselau, C.: Poly(vinylidene difluoride) membranes as the interface between laser desorption mass spectrometry, gel electrophoresis, and *in situ* proteolysis. *Anal. Chem.*, 66 (1994) 471-477.
- 1537 Warren, W.J., Cheng, Y.F. and Fuchs, M.: Protein immunodetection using capillary electrophoresis with membrane fraction collection. *LC-GC*, 12 (1994) 22-28; *C.A.*, 120 (1994) 264948f.
- 1538 Zhao, Z., Malik, A. and Lee, M.L.: Separations of proteins and proteolytic digests of proteins by capillary electrophoresis on Superox-coated open tubular columns. *J. Microcolumn Sep.*, 4 (1992) 411-417; *C.A.*, 120 (1994) 157935c.
- 1539 Zhu, M., Levi, V., Wehr, T. and Blum, W.: (Separation of proteins according to the molecular weight by capillary electrophoresis). *LaborPraxis*, 17 (1993) 24-25; *C.A.*, 120 (1994) 152667g.

See also 1316, 1317, 1371, 1402, 1436, 1439.

19b. *Proteins of cells, viruses and subcellular particles*

- 1540 Egger, C. and Winkler, H.: Bovine chromaffin cells: studies on the biosynthesis of phospholipids in chromaffin granules. *Biochim. Biophys. Acta*, 1211 (1994) 277-282.
- 1541 Galli, G., Rodriguez, F., Cosmina, P., Pratesi, C., Nogarotto, R., de Ferra, F. and Grandi, G.: Characterization of the surfactin synthetase multi-enzyme complex. *Biochim. Biophys. Acta*, 1205 (1994) 19-28.
- 1542 Geimonen, E.R., Rubtsov, A.M. and Boldyrev, A.A.: (The mechanism of temperature uncoupling of the sarcoplasmic reticulum Ca-pump: changes in the functional properties and structural state of the enzyme molecule). *Biokhimiya (Moscow)*, 58 (1993) 1296-1306.
- 1543 Hare, J.F. and Holoher, A.: Solubility in non-ionic detergents distinguishes between slowly and rapidly degraded plasma membrane proteins. *J. Biol. Chem.*, 269 (1994) 5981-5988.
- 1544 Heidenreich, O., Benseler, F., Fahrenholz, A. and Eckstein, F.: High activity and stability of hammerhead ribozymes containing 2'-modified pyrimidine nucleosides and phosphorothioates. *J. Biol. Chem.*, 269 (1994) 2131-2138.
- 1545 Higashi, K. and Ogawara, H.: Daidzein inhibits insulin- or insulin-like growth factor-1-mediated signaling in cell cycle progression of Swiss 3T3 cells. *Biochim. Biophys. Acta*, 1221 (1994) 29-35.
- 1546 Jäckle, S., Beisiegel, U., Rinninger, F., Buck, F., Grigoleit, A., Block, A., Gröger, I., Greten, H. and Windler, E.: Annexin VI, a marker protein of hepatocytic endosomes. *J. Biol. Chem.*, 269 (1994) 1026-1032.
- 1547 Kerfeld, C.A., Yeates, T.O. and Thornber, J.P.: Biochemical and spectroscopic characterization of the reaction center-LH1 complex and the carotenoid-containing B820 subunit of *Chromatium purpuratum*. *Biochim. Biophys. Acta*, 1185 (1994) 193-202.
- 1548 Lee, E.S., Steiner, M. and Lin, R.: Thioallyl compounds: potent inhibitors of cell proliferation. *Biochim. Biophys. Acta*, 1221 (1994) 73-77.
- 1549 Liu, M.Y., Torgrimson, A. and Colombini, M.: Characterization and partial purification of the VDAC-channel-modulating protein from calf liver mitochondria. *Biochim. Biophys. Acta*, 1185 (1994) 203-212.
- 1550 Miyamoto, K., Kanaya, S., Morikawa, K. and Inokuchi, H.: Overproduction, purification, and characterization of ferrochelatase from *Escherichia coli*. *J. Biochem. (Tokyo)*, 115 (1994) 545-551.
- 1551 Muller, G., Korndörfer, A., Kornak, U. and Malaisse, W.J.: Porin proteins in mitochondria from rat pancreatic islet cells and white adipocytes: identification and regulation of hexokinase binding by the sulfonyleurea glimepiride. *Arch. Biochem. Biophys.*, 308 (1994) 8-23.
- 1552 Neshar, M. and Boneh, A.: Effect of fatty acids and their acyl-CoA esters on protein kinase C activity in fibroblasts: possible implications in fatty acid oxidation defects. *Biochim. Biophys. Acta*, 1221 (1994) 66-72.
- 1553 Nigam, S.K., Goldberg, A.L., Ho, S., Rohde, M.F., Bush, K.T. and Sherman, M.Y.: A set of endoplasmic reticulum proteins possessing properties of molecular chaperones includes Ca²⁺-binding proteins and members of the thioredoxin superfamily. *J. Biol. Chem.*, 269 (1994) 1744-1749.
- 1554 Okumura, N., Miyatake, Y., Takao, T., Tamaru, T., Nagai, K., Okada, M. and Nakagawa, H.: Vasoactive intestinal peptide induces differentiation and MAP kinase activation in PC12h cells. *J. Biochem. (Tokyo)*, 115 (1994) 304-308.
- 1555 Okwu, A.K., Mais, D.E. and Halushka, P.V.: Agonist-induced phosphorylation of human platelet TXA₂/PGH₂ receptors. *Biochim. Biophys. Acta*, 1221 (1994) 83-88.
- 1556 Pushkareva, M.Yu. and Hannun, Y.A.: Modulation of cytosolic protein phosphorylation by sphingosylphosphorylcholine. *Biochim. Biophys. Acta*, 1221 (1994) 54-60.
- 1557 Radomska, A., Paul, P., Treat, S., Towbin, H., Pratt, C., Little, J., Magdalou, J., Lester, R. and Drake, R.: Photoaffinity labeling for evaluation of uridinyl analogs as specific inhibitors of rat liver microsomal UDP-glucuronosyltransferases. *Biochim. Biophys. Acta*, 1205 (1994) 336-345.
- 1558 Scott, J.E.: Influence of protein kinase C activation by 4β-phorbol ester or 1-oleoyl-2-acetyl-glycerol on disaturated phosphatidylcholine synthesis and secretion, and protein phosphorylation in differentiating fetal rabbit type II alveolar cells. *Biochim. Biophys. Acta*, 1221 (1994) 297-306.
- 1559 Teshima, R., Ikebuchi, H., Sawada, J.-i., Furuno, T., Nakanishi, M. and Terao, T.: Effects of herbimycin A and ST638 on Fcε receptor-mediated histamine release and Ca²⁺ signals in rat basophilic leukemia (RBL-2H3) cells. *Biochim. Biophys. Acta*, 1221 (1994) 37-46.
- 1560 Tonouchi, A., Fujita, A. and Kuhara, S.: Molecular cloning of the gene encoding a highly expressed protein in SFL1 gene-disrupted flocculating yeast. *J. Biochem. (Tokyo)*, 115 (1994) 683-688.
- 1561 Woo, M.K. and Murray, B.A.: Solid-phase binding analysis of N-CAM interactions with brain fodrin. *Biochim. Biophys. Acta*, 1191 (1994) 173-180.
- 1562 Yamaguchi, T., Yamada, S. and Kimoto, E.: Effects of cross-linking of membrane proteins on vesiculation induced by dimyristoylphosphatidylcholine in human erythrocytes. *J. Biochem. (Tokyo)*, 115 (1994) 659-663.

See also 1532, 1570, 1580, 1682, 1697, 1734, 1779, 1825, 1962.

19c. *Proteins synthesized by genetic manipulation, monoclonal antibodies*

- 1563 Aki, T., Fujikawa, A., Wada, T., Jyo, T., Shigeta, S., Murooka, Y., Oka, S. and Ono, K.: Cloning and expression of cDNA coding for a new allergen from the house dust mite, *Dermatophagoides farinae*: homology with human heat shock cognate proteins in the heat shock protein 70 family. *J. Biochem. (Tokyo)*, 115 (1994) 435-440.
- 1564 Asundi, V.K., Stahl, R.C., Showalter, L., Conner, K.J. and Carey, D.J.: Molecular cloning and characterization of an isoprenylated 67 kDa protein. *Biochim. Biophys. Acta*, 1217 (1994) 257-265.
- 1565 Dhandayuthapani, S., Banu, M.J. and Kashiwabara, Y.: Cloning and sequence determination of the gene coding for the elongation factor Tu of *Mycobacterium leprae*. *J. Biochem. (Tokyo)*, 115 (1994) 664-669.
- 1566 Endo, T., Ishibashi, Y., Shiokawa, S., Fukumaki, Y. and Okano, H.: Differential induction of adult and fetal globin gene expression in the human CML cell subline KU-812F/33. *J. Biochem. (Tokyo)*, 115 (1994) 540-544.

- 1567 Fukuchi, T., Shimada, N., Hanai, N., Ishikawa, N., Watanabe, K. and Kimura, N.: Recombinant rat nucleoside diphosphate kinase isoforms (α and β): purification, properties and application to immunological detection of native isoforms in rat tissues. *Biochim. Biophys. Acta*, 1205 (1994) 113-122.
- 1568 Groenen, P.J.T.A., Grootjans, J.J., Lubsen, N.H., Blemendal, H. and de Jong, W.W.: Lys-17 is the amine-donor substrate site for transglutaminase in β A3-crystallin. *J. Biol. Chem.*, 269 (1994) 831-833.
- 1569 Harada, Y., Senda, T., Sakamoto, T., Takamoto, K. and Ishibashi, T.: Expression of octopus rhodopsin in *Escherichia coli*. *J. Biochem. (Tokyo)*, 115 (1994) 66-75.
- 1570 Ishida, N., Tsujimoto, M., Kanaya, T., Shimamura, A., Tsuruoka, N., Kodama, S., Katayama, T., Oikawa, S., Matsui, M., Nakanishi, T., Kobayashi, J. and Nakazato, H.: Expression and characterization of human bone morphogenetic protein-2 in silkworm larvae infected with recombinant *Bombyx mori* nuclear polyhedrosis virus. *J. Biochem. (Tokyo)*, 115 (1994) 279-285.
- 1571 Kurome, T., Katayama, M., Murakami, K., Hashino, K., Kamihagi, K., Yasumoto, M. and Kato, I.: Expression of recombinant mouse/human chimeric antibody specific to human GMP-140/P-selectin. *J. Biochem. (Tokyo)*, 115 (1994) 608-614.
- 1572 Nawa, K., Ono, M., Fujiwara, H., Sugiyama, N., Uchiyama, T. and Marumoto, Y.: Monoclonal antibodies against human thrombomodulin whose epitope is located in epidermal growth factor-like domains. *Biochim. Biophys. Acta*, 1205 (1994) 162-170.
- 1573 Orita, T., Oh-eda, M., Hasegawa, M., Kuboniwa, H., Esaki, K. and Ochi, N.: Polypeptide and carbohydrate structure of recombinant human interleukin-6 produced in Chinese hamster ovary cells. *J. Biochem. (Tokyo)*, 115 (1994) 345-350.
- 1574 Peng, H.-L., Hsieh, M.-J., Zao, C.-L. and Chang, H.-Y.: Nucleotide sequence and expression in *Escherichia coli* of the *Klebsiella pneumoniae* *dead* gene. *J. Biochem. (Tokyo)*, 115 (1994) 409-414.
- 1575 Srinivasan, K., Morris, S.C., Girard, J.E., Kline, M.C. and Reeder, D.J.: Enhanced detection of PCR products through use of TOTO and YOYO intercalating dyes with laser induced fluorescence-capillary electrophoresis. *Appl. Theor. Electrophor.*, 3 (1993) 235-239; *C.A.*, 120 (1994) 126278m.
- See also 1725, 1735, 1770.
- 19d. *Microbial and plant proteins*
- 1576 Abe, A., Ogawa, S., Kohno, T. and Watabe, K.: Purification of *Bacillus subtilis* spore coat protein by electrophoretic elution procedure and determination of amino-terminal amino acid sequences. *Microbiol. Immunol.*, 37 (1993) 809-812; *C.A.*, 120 (1994) 211934s.
- 1577 Bornstein, N., Dumaine, M.H., Forey, F. and Fleurette, J.: Complementary identification of *Legionella* species by electrophoretic characterization of soluble protein extracts. In: Barbaree, J.M., Breiman, R.F. and Dufour, A.P. (Editors), *Legionella*, Am. Soc. Microbiol., Washington, 1993, pp. 194-198; *C.A.*, 120 (1994) 72735n.
- 1578 Bult, C.J. and Kiang, Y.T.: One-dimensional electrophoretic comparisons of plant proteins. *Methods Enzymol.*, 224 (Molecular Evolution: Producing the Biochemical Data) (1993) 81-97; *C.A.*, 120 (1994) 239454j.
- 1579 Combe, M.L., Pons, J.L., Sesboue, R. and Martin, J.P.: Electrophoretic transfer from polyacrylamide gel to nitrocellulose sheets, a new method to characterize multilocus enzyme genotypes of *Klebsiella* strains. *Appl. Environ. Microbiol.*, 60 (1994) 26-30; *C.A.*, 120 (1994) 101051u.
- 1580 Damm, I. and Green, B.R.: Separation of closely related intrinsic membrane polypeptides of the Photosystem II light-harvesting complex (LHC II) by reversed-phase high-performance liquid chromatography on a poly(styrene-divinylbenzene) column. *J. Chromatogr. A*, 664 (1994) 33-38.
- 1581 De Moissac, Y.R., Ronald, S.L. and Peppler, M.S.: Use of pulsed-field gel electrophoresis for epidemiological study of *Bordetella pertussis* in whooping cough outbreak. *J. Clin. Microbiol.*, 32 (1994) 398-402; *C.A.*, 120 (1994) 211835k.
- 1582 Ejiri, S.-i., Kawamura, R. and Katsumata, T.: Interactions among four subunits of elongation factor 1 from rice embryo. *Biochim. Biophys. Acta*, 1217 (1994) 266-272.
- 1583 Enami, I., Tomo, T., Kitamura, M. and Katoh, S.: Immobilization of the three extrinsic proteins in spinach oxygen-evolving Photosystem II membranes: roles of the proteins in stabilization of binding of Mn and Ca^{2+} . *Biochim. Biophys. Acta*, 1185 (1994) 75-80.
- 1584 Ito, A., Yasuda, K., Yamada, K., Komori, T. and Kitaguni, H.: Determination of antigenic protein of *Echinococcus* by immunoassay and SDS-polyacrylamide electrophoresis. *Jpn. Kokai Tokkyo Koho JP 06 25,298* [94 25,298] (Cl. CO7K15/08), 01 Feb. 1994, Appl. 92/207,054, 13 Jul. 1992; 10 pp.; *C.A.*, 120 (1994) 268200d.
- 1585 Lioi, L.: Electrophoretic variation and geographical distribution of the seed protein phytohemagglutinin in cultivated *Phaseolus vulgaris* L. *J. Genet. Breed.*, 45 (1991) 97-102; *C.A.*, 120 (1994) 102035d.
- 1586 Liu, H., Xu, G. and Cai, J.: (Electrophoretic analysis of soluble proteins in Shechuangzi). *Zhongguo Zhonggyao Zazhi*, 18 (1993) 465-466; *C.A.*, 120 (1994) 62394m.
- 1587 Liu, L. and Xue, G.: (A rapid method for two-dimensional electrophoresis of rice leaf proteins). *Shengwu Huaxue Yu Shengwu Wuli Jinzhan*, 19 (1992) 387-390; *C.A.*, 120 (1994) 49359j.
- 1588 Lopez, M.F., Patton, W.F., Sawlisch, W.B., Erdjument-Bromage, H., Barry, P., Gmyrek, K., Hines, T., Tempst, P. and Skela, W.M.: A glutathione S-transferase (GST) isozyme from broccoli with significant sequence homology to the mammalian theta-class of GSTs. *Biochim. Biophys. Acta*, 1205 (1994) 29-38.
- 1589 Metakovsky, E.V., Ng, P.K.W., Chernakov, V.M., Pogna, N.E. and Bushuk, W.: Gliadin alleles in Canadian western red spring wheat cultivars: use of two different procedures of acid polyacrylamide gel electrophoresis for gliadin separation. *Genome*, 36 (1993) 743-749; *C.A.*, 120 (1994) 157940a.
- 1590 Michea-Hamzehpour, M., Sanchez, J.C., Epp, S.F., Paquet, N., Hughes, G.J., Hochstrasser, D. and Pechere, J.C.: Two-dimensional polyacrylamide gel electrophoresis isolation and microsequencing of *Pseudomonas aeruginosa* proteins. *Enzyme Protein*, 47 (1993) 1-8; *C.A.*, 120 (1994) 49378q.
- 1591 Miller, B.S.: Analysis of the heat-shock response in *Bacillus subtilis* by two-dimensional gel electrophoresis. Avail. *Univ. Microfilms Int.*, Order No. DA9230095, 1992, 112 pp.; *C.A.*, 120 (1994) 128868c.

- 1592 Moore, C.B. and Brooks, W.M.: An evaluation of SDS-polyacrylamide gel electrophoretic analysis for the determination of intragenetic relationships of *Vairimorpha* isolates. *J. Invertebr. Pathol.*, 62 (1993) 285-288; C.A., 120 (1994) 211845p.
- 1593 Nitta, I., Ueda, T. and Watanabe, K.: Template-dependent peptide formation on ribosomes catalyzed by pyridine. *J. Biochem. (Tokyo)*, 115 (1994) 803-807.
- 1594 Oyama, F., Hashino, K., Oyama, R., Kato, I. and Titani, K.: Improved method for expression of Kunitz-type serine proteinase inhibitor domain of β -amyloid protein precursor in *Escherichia coli* and characterization of disulfide bonds of the product. *J. Biochem. (Tokyo)*, 114 (1993) 813-819.
- 1595 Santoni, V., Bellini, C. and Caboche, M.: Use of two-dimensional protein-pattern analysis for the characterization of *Arabidopsis thaliana* mutants. *Planta*, 192 (1994) 557-566; C.A., 120 (1994) 212875k.
- 1596 Schoel, B., Welzel, M. and Kaufmann, S.H.E.: Hydrophobic interaction chromatography for the purification of cytolytic bacterial toxins. *J. Chromatogr. A*, 667 (1994) 131-139.
- 1597 Sloss, L.L.: Evaluation of sodium dodecyl sulfate-polyacrylamide gel electrophoresis of proteins for the identification and typing of medically important bacteria. Avail. *Univ. Microfilms Int.*, Order No. BRDX96152, 1991, 229 pp.; C.A., 120 (1994) 101052v.
- 1598 Tanabe, M., Asano, T., Moriya, N., Sugino, H. and Kakinuma, A.: Isolation and characterization of *Streptovorticillium* anticoagulant (SAC), a novel protein inhibitor of blood coagulation produced by *Streptovorticillium cinnamomeum* subsp. *cinnamomeum* *J. Biochem. (Tokyo)*, 115 (1994) 743-751.
- 1599 Tanabe, M., Kawahara, K., Asano, T., Kato, K. and Kakinuma, A.: Primary structure and reactive site of *Streptovorticillium* anticoagulant (SAC), a novel protein inhibitor of blood coagulation produced by *Streptovorticillium cinnamomeum* subsp. *cinnamomeum* *J. Biochem. (Tokyo)*, 115 (1994) 752-761.
- 1600 Terada, S., Fujimura, S., Katayama, H., Nagasawa, M. and Kimoto, E.: Purification and characterization of two Kunitz family subtilisin inhibitors from seeds of *Canavalia lineata*. *J. Biochem. (Tokyo)*, 115 (1994) 392-396.
- 1601 Wrigley, C.W.: Increased efficiency in the electrophoretic identification of cereal grains according to variety and quality type. *Am. Lab. (Shelton)*, 25 (1993) 52D, 52F-52I; C.A., 120 (1994) 75776n.
- 1602 Yadav, S.P. and Dogra, J.V.V.: Identification of Indian wheat cultivars by gel electrophoresis of seed proteins. *J. Indian Bot. Soc.*, 71 (1992) 285-286; C.A., 120 (1994) 265823y.
- 1603 Yu, S.-G., Romanowska, E., Xue, Z.-t. and Albertsson, P.-Å.: Evidence for two different Rieske iron-sulfur proteins in the cytochrome *bf* complex of spinach chloroplast. *Biochim. Biophys. Acta*, 1185 (1994) 239-242.
- See also 1326, 1563, 1565, 1840.
- 19e. *Proteins of blood, serum and blood cells*
- 1604 Allen, G.J., Long, B.J. and Wisdom, G.B.: Identification of human plasma proteins using the Laemmli method of gel electrophoresis. *Biochem. Soc. Trans.*, 22 (1994) 26S; C.A., 120 (1994) 211858v.
- 1605 Attanasio, R., Stunz, G.W. and Kennedy, R.C.: Folding patterns of immunoglobulin molecules identified by urea gradient electrophoresis. *J. Biol. Chem.*, 269 (1994) 1834-1838.
- 1606 Bassé, F., Gaffet, P. and Bienvenüe, A.: Correlation between inhibition of cytoskeleton proteolysis and anti-vasculature effect of calpeptin during A23187-induced activation of human platelets: are vesicles shed by filopod fragmentation? *Biochim. Biophys. Acta*, 1190 (1994) 217-224.
- 1607 Bialkowska, K., Zembroń, A. and Sikorski, A.F.: Ankyrin inhibits binding of erythrocyte spectrin to phospholipid vesicles. *Biochim. Biophys. Acta*, 1191 (1994) 21-26.
- 1608 Chen, F.T.A., Sternberg, J.C. and Satow, T.: (Capillary electrophoresis. Serum protein analysis by capillary electrophoresis). *Rinsho Kagaku (Nippon Rinsho Kagakkai)*, 22 (1993) 214-219; C.A., 120 (1994) 239288h - a review with 12 refs.
- 1609 Cochet, S., Hasnaoui, M'H., Debbia, M., Kroviarski, Y., Lambin, P., Cartron, J.P. and Bertrand, O.: Chromatography of human immunoglobulin G on immobilized Dimarene Rubine R/K-5BL. Study of mild, efficient elution procedures. *J. Chromatogr. A*, 663 (1994) 175-186.
- 1610 Degerman, E., Moos, M., Jr., Rascón, A., Vasta, V., Meacci, E., Smith, C.J., Lindgren, S., Andersson, K.-E., Belfrage, P. and Manganiello, V.: Single-step affinity purification, partial structure and properties of human platelet cGMP inhibited cAMP phosphodiesterase. *Biochim. Biophys. Acta*, 1205 (1994) 189-198.
- 1611 Devine, D.V., Wong, K., Serrano, K., Chonn, A. and Cullis, P.R.: Liposome-complement interactions in rat serum: implications for liposome survival studies. *Biochim. Biophys. Acta*, 1191 (1994) 43-51.
- 1612 Goux, A., Athias, A., Persegol, L., Lagrost, L., Gambert, P. and Lallemand, C.: Capillary electrophoresis analysis of apolipoproteins A-I and A-II in human high-density lipoproteins. *Anal. Biochem.*, 218 (1994) 320-324.
- 1613 Hashimoto, S. and Kawano, K.: (Studies on the substance which causes a double β fraction on serum protein cellulose-acetate membrane electrophoresis). *Seibutsu Butsuri Kagaku*, 38 (1994) 7-12; C.A., 120 (1994) 265111q.
- 1614 Heiden, S.M.: Electrophoretic screening for variant human lymphocyte proteins labeled with tritiated sodium cyanoborohydride. Avail. *Univ. Microfilms Int.*, Order No. DA9225598, 1992, 146 pp.; C.A., 120 (1994) 128861v.
- 1615 Hlady, V., Andrade, J.D., Ho, C.H., Feng, L. and Tingey, K.: Plasma protein adsorption on model biomaterial surfaces. *Clin. Mater.*, 13 (1993) 85-93; C.A., 120 (1994) 144034h.
- 1616 Islam, M., Qamar, S. and Tayyab, S.: Involvement of lysine residues of goat serum albumin in high-affinity binding of bilirubin. *Biochim. Biophys. Acta*, 1205 (1994) 171-177.
- 1617 Kontturi, A.K., Kontturi, K., Sievers, G. and Vuoristo, M.: Purification of a polyelectrolyte sample from complex-forming charged trace impurities by continuous convective electrophoresis: depyrogenation of human albumin. *Sep. Sci. Technol.*, 29 (1994) 1029-1042; C.A., 120 (1994) 157959p.
- 1618 Kunitake, S.T., Carilli, C.T., Lau, K., Protter, A.A., Naya-Vigne, J. and Kane, J.P.: Identification of proteins associated with apolipoprotein A-I-containing lipoproteins purified by selected-affinity immunosorption. *Biochemistry*, 33 (1994) 1988-1993.
- 1619 Ono, M., Tsutsui, K., Wakui, H. and Yamanouchi, H.: (Application of electrophoresis to blond group typing in forensic science). *Seibutsu Butsuri Kagaku*, 37 (1993) 363-366; C.A., 120 (1994) 210062u - a review with 8 refs.

- 1620 Sakai, K., Fujii, T. and Hayashi, T.: Cell-free formation of disulfide-bonded multimer from isolated plasma fibronectin in the presence of a low concentration of SH reagent under a physiological condition. *J. Biochem. (Tokyo)*, 115 (1994) 415-421.
- 1621 Sinha, D., Yang, X., Emig, F. and Kirby, E.P.: Isolation and characterization of two protease inhibitors from bovine plasma. *J. Biochem. (Tokyo)*, 115 (1994) 387-391.
- 1622 Taya, K.: Microporous membrane of cellulose acetate containing fatty acid esters of polyoxyethylene sorbitol for electrophoresis. *Jpn. Kokai Tokkyo Koho JP 06 66,768 [94 66,768]* (Cl. GO1N27/447), 11 Mar. 1994, Appl. 92/245,686, 22 Aug. 1992; 9 pp.; C.A., 120 (1994) 265333p.
- 1623 Turrini, F., Mannu, F., Cappadoro, M., Ulliers, D., Giribaldi, G. and Arese, P.: Binding of naturally occurring antibodies to oxidatively and nonoxidatively modified erythrocyte band 3. *Biochim. Biophys. Acta*, 1190 (1994) 297-303.
- 1624 Urano, T., Ikeda, C., Shimokawa, M. and Kinoshita, T.: Different effect of 5,6-*trans*-prostaglandin E₂ on plasminogen activation by urokinase and streptokinase. *Biochim. Biophys. Acta*, 1205 (1994) 258-261.
- 1625 Vincent, C., Denroy, L. and Revillard, J.-P.: Molecular variants of β_2 -microglobulin in renal insufficiency. *Biochem. J.*, 298 (1994) 181-187.
- 1626 Vlasov, V.V., Pautova, L.V., Rykova, E.Yu. and Yakubov, L.A.: (Oligonucleotide interaction with serum proteins). *Biokhimiya (Moscow)*, 58 (1993) 1247-1251.
- 1627 Wada, M., Senoo, H. and Shinozaki, H.: Computerized electrophoresis device for fractionating serum globulins, detg. monoclonal proteins, and diagnosing diseases. *Jpn. Kokai Tokkyo Koho JP 05,312,812 [93,312,812]* (Cl. GO1N33/68), 26 Nov. 1993, Appl. 92/121,939, 14 May 1992; 16 pp.; C.A., 120 (1994) 158130y.
- 1628 Yamaguchi, K., Kato, N., Sakai, N., Matsumoto, M., Nagasawa, S., Hatsuse, H., Toyoguchi, T., Moriya, H. and Sakiyama, H.: Immune complex independent activation of complement, C1s secreted from hamster embryo malignant fibroblasts, Nil2C2 in serum free culture medium. *Biochim. Biophys. Acta*, 1205 (1994) 133-138.
- 1629 Yamauchi, M., Mizuhara, Y., Takahashi, A. and Toda, G.: Increases in serum β_3 integrin in patients with chronic liver disease. *Clin. Chim. Acta*, 225 (1994) 71-76.
- 1630 Yoshida, N., Hirata, H., Imaoka, S., Matsuda, M., Yamazumi, K. and Asakura, S.: Effect of calcium on the mobility of γ -chain from fibrinogen Osaka V on sodium dodecyl sulfate-polyacrylamide gel electrophoresis. *Thromb. Res.*, 73 (1994) 79-82; C.A., 120 (1994) 128880a.
- 1631 Zhang, J. and Redman, C.M.: Role of interchain disulfide bonds on the assembly and secretion of human fibrinogen. *J. Biol. Chem.*, 269 (1994) 652-658.
- See also 1527, 1533, 1539, 1571, 1572, 1657.
- 19f. *Structural and muscle proteins*
- 1632 Chandrasekhar, S., Harvey, A.K., Johnson, M.G. and Becker, G.W.: Osteonectin/SPARC is a product of articular chondrocytes/cartilage and is regulated by cytokines and growth factors. *Biochim. Biophys. Acta*, 1221 (1994) 7-14.
- 1633 Deng, Y. and Meng, J.: (Analysis of the composition changes in protein and lipid of modified keratin fibers). *Fenxi Ceshi Tongbao*, 11 (1992) 74-77; C.A., 120 (1994) 166632d.
- 1634 Feng, Y., Chen, B. and Liu, H.: (Polyacrylamide gel electrophoresis of *Bombyx mori* silk proteins. II. Preparative polyacrylamide gel electrophoresis of the middle gland fibroin). *Zhejiang Sichou Gongxueyuan Xuebao*, 9, No. 1 (1992) 1-4; C.A., 120 (1994) 136898c.
- 1635 Feng, Y., Chen, B. and Liu, H.: (Polyacrylamide gel electrophoresis of *Bombyx mori* silk proteins. III. Isoelectric focusing and two-dimensional polyacrylamide gel electrophoresis of silk proteins). *Zhejiang Sichou Gongxueyuan Xuebao*, 9, No. 2 (1992) 1-5; C.A., 120 (1994) 136899d.
- 1636 Feng, Y., Chen, B. and Xu, H.: (Polyacrylamide gel electrophoresis of *Bombyx mori* silk proteins. I. Polyacrylamide gel electrophoresis of dissociated and non-dissociated systems of silk proteins). *Zhejiang Sichou Gongxueyuan Xuebao*, 8 (1991) 6-10; C.A., 120 (1994) 166627f.
- 1637 Kaune, R., Munson, S. and Bikle, D.D.: Regulation of calmodulin binding to the ATP extractable 110 kDa protein (myosin I) from chicken duodenal brush border by 1,25-(OH)₂D₃. *Biochim. Biophys. Acta*, 1190 (1994) 329-336.
- 1638 Kurioka, S., Ishioka, N., Sato, J., Ushigome, S. and Nakamori, K.: Identification of noncollagenous bone proteins and purification of osteonectin, using two-dimensional gel electrophoresis and high performance liquid chromatography. *Jikeikai Med. J.*, 40 (1993) 249-257; C.A., 120 (1994) 211928t.
- 1639 Kuroda, M., Kohira, Y. and Sasaki, M.: Conformational change of skeletal muscle α -actinin induced by salt. *Biochim. Biophys. Acta*, 1205 (1994) 97-104.
- 1640 Nakagawa, H., Ishihara, M. and Ohashi, K.: 33-kDa Peptides prepared from chicken gizzard smooth muscle bundle both actin and desmin filaments *in vitro*. *J. Biochem. (Tokyo)*, 114 (1993) 623-626.
- 1641 Sakurada, K., Ikuhara, T., Seto, M. and Sasaki, Y.: An antibody for phosphorylated myosin light chain of smooth muscle: application to a biochemical study. *J. Biochem. (Tokyo)*, 115 (1994) 18-21.
- 1642 Shiraishi, F. and Yamamoto, K.: The effect of partial removal of troponin I and C on the Ca²⁺-sensitive ATPase activity of rabbit skeletal myofibrils. *J. Biochem. (Tokyo)*, 115 (1994) 171-173.
- 1643 Takai, D., Kirinoki, M., Miyazaki, J.-I. and Hirabayashi, T.: Detection and characterization of muscle-specific nuclear proteins. *J. Biochem. (Tokyo)*, 115 (1994) 219-223.
- 1644 Tanabe, R., Tatsumi, R. and Takahashi, K.: Purification and characterization of the 1,200-kDa subfragment of connectin filaments produced by 0.1 mM calcium ions. *J. Biochem. (Tokyo)*, 115 (1994) 351-355.
- 1645 Tokimitsu, I. and Tajima, S.: Inhibition of elastin synthesis by high potassium salt is mediated by Ca²⁺ influx in cultured smooth muscle cells *in vitro*: reciprocal effects of K⁺ on elastin and collagen synthesis. *J. Biochem. (Tokyo)*, 115 (1994) 536-539.
- 1646 Tomita, M., Kinoshita, T., Izumi, S., Tomino, S. and Yoshizato, K.: Characterizations of sea urchin fibrillar collagen and its cDNA clone. *Biochim. Biophys. Acta*, 1217 (1994) 131-140.
- 1647 Yamamoto, H., Mizuno, Y., Hayashi, K., Nonaka, I., Yoshida, M. and Ozawa, E.: Expression of dystrophin-associated protein 35DAG (A4) and 50DAG (A2) is confined to striated muscles. *J. Biochem. (Tokyo)*, 115 (1994) 162-167.

1648 Yamamoto, K.: ATPase activities of the two cross-linked species of actin and myosin. *J. Biochem. (Tokyo)*, 114 (1993) 770-772.

See also 1515, 1524, 1606, 1719.

19g. *Protamines, histones and other nuclear proteins*

1649 Fletcher, T.M., Krishnan, U., Serwer, P. and Hansen, J.C.: Quantitative agarose gel electrophoresis of chromatin: nucleosome-dependent changes in charge, shape, and deformability at low ionic strength. *Biochemistry*, 33 (1994) 2226-2233.

1650 Peters, J.-M., Franke, W.W. and Kleinschmidt, J.A.: Distinct 19 S and 20 S subcomplexes of the 26 S proteasome and their distribution in the nucleus and the cytoplasm. *J. Biol. Chem.*, 269 (1994) 7709-7718.

1651 Poznanovic, G., Grujic, V., Ivanovic-Matic, S. and Sevaljevic, L.: Regions of the haptoglobin 5' flanking gene sequence show different binding affinities to nuclear matrix proteins during the acute phase response. *J. Biochem. (Tokyo)*, 115 (1994) 422-428.

1652 Tyulenev, V.I., Konoplich, L.A., Krivonos, A.A. and Khrapunov, S.N.: (ATP-dependent structural changes in chromatin). *Biokhimiya (Moscow)*, 58 (1993) 1206-1212.

1653 Wiekowski, M. and DePamphilis, M.L.: One-dimensional gel analysis of histone synthesis. *Methods Enzymol.*, 225 (Guide to Techniques in Mouse Development) (1993) 489-501; C.A., 120 (1994) 265100k.

See also 1643.

19h. *Chromoproteins and metalloproteins*

1654 Beattie, J.H. and Richards, M.P.: Separation of metallothionein isoforms by micellar electrokinetic capillary chromatography. *J. Chromatogr. A*, 664 (1994) 129-134.

1655 Lobanov, N.A., Hensey, C.E., Usanov, S.A. and Azzi, A.: (Phosphorylation of cytochrome P-450_{sc} by protein kinase C, protective effect of adrenodoxin and cytochrome bs). *Biokhimiya (Moscow)*, 58 (1993) 1529-1537.

1656 Okazaki, T., Arakawa, H., Nagai, T., Muraoka, M., Oda, T. and Kanno, T.: (Analysis of haptoglobin-variant by polyacrylamide gradient gel electrophoresis). *Seibutsu Butsuri Kagaku*, 38 (1994) 47-49; C.A., 120 (1994) 267452a.

1657 Seidel, A., Bill, E., Häggström, L., Nordblad, P. and Kilár, F.: Complementary Mössbauer and EPR studies of iron(III) in diferic human serum transferrin with oxalate or bicarbonate as synergistic anions. *Arch. Biochem. Biophys.*, 308 (1994) 52-63.

1658 Ujiie, M., Sanwar, G., Peace, R.W. and Watson, D.: Chemical and mouse growth tests for nutritional assessment of commercial lactoferrins. *Nutr. Res.*, 13 (1993) 1087-1097; C.A., 120 (1994) 132947v.

1659 Yamazaki, S., Sato, K., Suhara, K., Sakaguchi, M., Mihara, K. and Omura, T.: Importance of the proline-rich region following signal-anchor sequence in the formation of correct conformation of microsomal cytochrome P-450s. *J. Biochem. (Tokyo)*, 114 (1993) 652-657.

1660 Zhao, Z., Malik, A., Lee, M.L. and Watt, G.D.: A capillary electrophoresis method for studying apo, holo, recombinant, and subunit dissociated ferritins. *Anal. Biochem.*, 218 (1994) 42-54.

See also 1566.

19i. *Proteins of glands, gland products, various zymogens (incl. milk proteins)*

1661 Akama, K., Oka, S., Tobita, T. and Hayashi, H.: The amino acid sequence of a boar transition protein 3. *J. Biochem. (Tokyo)*, 115 (1994) 58-65.

1662 Bordin, L., Cattapan, F., Clari, G., Toninello, A., Siliprandi, N. and Moret, V.: Spermine-mediated casein kinase II-uptake by rat liver mitochondria. *Biochim. Biophys. Acta*, 1199 (1994) 266-270.

1663 Fujino, Y., Fujii, T. and Daimon, T.: Presence of laminin B chain-like protein in bovine and rat adrenal chromaffin granules. *J. Biochem. (Tokyo)*, 115 (1994) 615-621.

1664 Gerard, N. and Jegou, B.: *In-vitro* influence of germ cells on Sertoli cell-secreted proteins: a two-dimensional gel electrophoresis analysis. *Int. J. Androl.*, 16 (1993) 285-291; C.A., 120 (1994) 211833h.

1665 Lerner, E.A. and Nelson, R.J.: Recovery of biological activity and determination of N-terminal sequence of maxadilan following capillary electrophoresis. *LC-GC*, 12 (1994) 36-38; C.A., 120 (1994) 264949g.

1666 Maeda, H. and Kasahara, K.: Isolation and localization of the 120 kDa protein in the liver of genetically obese Zucker rats. *J. Biochem. (Tokyo)*, 115 (1994) 37-40.

1667 Matsumoto, K., Tajima, H., Okazaki, H. and Nakamura, T.: Heparin as an inducer of hepatocyte growth factor. *J. Biochem. (Tokyo)*, 114 (1993) 820-826.

1668 Motojima, K., Ohmori, A., Takino, Y. and Goto, S.: Increase in the amount of elongation factor 2 in rat liver by peroxisome proliferators. *J. Biochem. (Tokyo)*, 114 (1993) 779-785.

1669 Nielsen, S.U., Rump, R., Højrup, P., Roepstorff, P. and Spener, F.: Differentiation regulation and phosphorylation of the fatty acid-binding protein from rat mammary epithelial cells. *Biochim. Biophys. Acta*, 1211 (1994) 189-197.

1670 Solovyov, D.A. and Ugarova, T.P.: (Purification and characterization of α -specific thrombin-like enzymes from the venoms of Middle Asian pit viper snakes *Agkistrodon halys halys* and *Agkistrodon halys Blomhoffii*). *Biokhimiya (Moscow)*, 58 (1993) 1221-1233.

See also 1525, 1697.

19j. *Proteins of brain, cerebrospinal fluid and eye*

1671 Fujii, T., Watanabe, M., Ogoma, Y., Kondo, Y. and Arai, T.: Microtubule-associated proteins, MAP 1A and MAP 1B, interact with F-actin *in vitro*. *J. Biochem. (Tokyo)*, 114 (1993) 827-829.

1672 Hiraoka, A., Miura, I., Hattori, M., Tominaga, I. and Machida, S.: Capillary-zone electrophoretic analyses of the proteins and amino acid components in cerebrospinal fluid of central nervous system diseases. *Biol. Pharm. Bull.*, 16 (1993) 949-952; C.A., 120 (1994) 157952f.

1673 Kato, K., Goto, S., Hasegawa, K. and Inaguma, Y.: Coinduction of two low-molecular-weight stress proteins, β B crystallin and HSP28, by heat or arsenite stress in human glioma cells. *J. Biochem. (Tokyo)*, 114 (1993) 640-647.

- 1674 Mizuno, Y., Yoshida, M., Yamamoto, H., Hirai, S. and Ozawa, E.: Distribution of dystrophin isoforms and dystrophin-associated proteins 43DAG (A3a) and 50DAG (A2) in various monkey tissues. *J. Biochem. (Tokyo)*, 114 (1993) 936-941.
- 1675 Sheller, R.A. and Bittner, G.D.: Whole intact tissue electrophoresis of nerve proteins. *J. Neurosci. Methods*, 49 (1993) 185-191; *C.A.*, 120 (1994) 186551e.
- 1676 Van Noort, J.M., El Ouagmiri, M., Boon, J. and van Sechel, A.C.: Fractionation of central nervous system myelin proteins by reversed-phase high-performance liquid chromatography. *J. Chromatogr. B*, 653 (1994) 155-161.
- See also 1569.
- 19k. *Proteins of neoplastic tissue and transformed cells*
- 1677 Keyomarsi, K., O'Leary, N., Molnar, G., Lees, E., Fingert, H.J. and Pardee, A.B.: Cyclin E, a potential prognostic marker for breast cancer. *Cancer Res.*, 54 (1994) 380-385.
- 1678 Yamaguchi, N., Hattori, K., Oh-eda, M., Kojima, T., Imai, N. and Ochi, N.: A novel cytokine exhibiting megakaryocyte potentiating activity from a human pancreatic tumor cell line HPC-Y5. *J. Biol. Chem.*, 269 (1994) 805-808.
- See also 1564, 1628.
- 19l. *Specific binding and receptor proteins*
- 1679 Ciruela, F., Blanco, J., Canela, E.I., Lluís, C., Franco, R. and Mallol, J.: Solubilization and molecular characterization of the nitrobenzylthioinosine binding sites from pig kidney brush-border membranes. *Biochim. Biophys. Acta*, 1191 (1994) 94-102.
- 1680 Crepaldi, T., Prat, M., Gioradano, S., Medico, E. and Comoglio, P.M.: Generation of a truncated hepatocyte growth factor receptor in the endoplasmic reticulum. *J. Biol. Chem.*, 269 (1994) 1750-1755.
- 1681 Dziarski, R. and Gupta, D.: Heparin, sulfated heparinoids, and lipoteichoic acids bind to the 70-kDa peptidoglycan/lipopolysaccharide receptor protein on lymphocytes. *J. Biol. Chem.*, 269 (1994) 2100-2110.
- 1682 Inanobe, A., Takahashi, K. and Katada, T.: Association of the β subunits of trimeric GTP-binding proteins with 90-kDa heat shock protein, hsp90. *J. Biochem. (Tokyo)*, 115 (1994) 486-492.
- 1683 Iwashina, M., Mizuno, T., Hirose, S., Ito, T. and Hagiwara, H.: His¹⁴⁵-Trp¹⁴⁶ residues and the disulfide-linked loops in atrial natriuretic peptide receptor are critical for the ligand-binding activity. *J. Biochem. (Tokyo)*, 115 (1994) 563-567.
- 1684 Joseph, S.K.: Biosynthesis of the inositol triphosphate receptor in WB rat liver epithelial cells. *J. Biol. Chem.*, 269 (1994) 5673-5679.
- 1685 Kohira, T., Matsumoto, K., Ichihara, A. and Nakamura, T.: Identification of a biologically functional novel IL-1 β -specific receptor on adult rat hepatocytes. *J. Biochem. (Tokyo)*, 114 (1993) 658-662.
- 1686 Liu, F. and Roth, R.A.: Identification of serines-967/968 in the juxtamembrane region of the insulin receptor as insulin-stimulated phosphorylation sites. *Biochem. J.*, 298 (1994) 471-477.
- 1687 Mason, G.G.F., Witte, A.-M., Whitelaw, M.L., Antonsson, C., McGuire, J., Wilhelmsson, A., Poellinger, L. and Gustafsson, J.-A.: Purification of the DNA binding form of dioxin receptor. Role of the Arnt cofactor in regulation of dioxin receptor function. *J. Biol. Chem.*, 269 (1994) 4438-4449.
- 1688 Moore, M.A. and McCarthy, M.P.: The effects of drugs on the incorporation of a conformationally-sensitive, hydrophobic probe into the ion channel of the nicotinic acetylcholine receptor. *Biochim. Biophys. Acta*, 1190 (1994) 457-464.
- 1689 Paiement, J., Lavoie, C., Gavino, G.R. and Gavino, V.C.: Modulation of GTP-dependent fusion by linoleic and arachidonic acid in derivatives of rough endoplasmic reticulum from rat liver. *Biochim. Biophys. Acta*, 1190 (1994) 199-212.
- 1690 Preiss, T., Chrzanosowska-Lightowlers, Z.M.A. and Lightowlers, R.N.: The tissue-specific RNA-binding protein COLBP is differentially regulated during myogenesis. *Biochim. Biophys. Acta*, 1221 (1994) 286-289.
- 1691 Rodríguez-Henche, N., Rodríguez-Pena, M., Guizarro, L.G. and Prieto, J.C.: Characterization of vasoactive intestinal peptide receptors in human liver. *Biochim. Biophys. Acta*, 1221 (1994) 193-198.
- 1692 Schletter, J., Krüger, C., Lottspeich, F. and Schütt, C.: Improved method for preparation of lipopolysaccharide-binding protein from human serum by electrophoretic and chromatographic separation techniques. *J. Chromatogr. B*, 654 (1994) 25-34.
- 1693 Schwabe, M., Zhao, J. and Kung, H.: Differential expression and ligand-induced modulation of the human interleukin-6 receptor on interleukin-6-responsive cells. *J. Biol. Chem.*, 269 (1994) 7201-7209.
- 1694 Singh, K., Deonaraine, D., Shanmugam, V., Senger, D.R., Mukherjee, A.B., Chang, P.-L., Prince, C.W. and Mukherjee, B.B.: Calcium-binding properties of osteopontin derived from non-osteogenic sources. *J. Biochem. (Tokyo)*, 114 (1993) 702-707.
- 1695 Takasuka, T., Sakurai, T., Goto, K., Furuichi, Y. and Watanabe, T.: Human endothelin receptor ET β . Amino acid sequence requirements for super stable complex formation with its ligand. *J. Biol. Chem.*, 269 (1994) 7509-7513.
- 1696 Takeda, S., Sagara, Y., Kita, K., Natori, S. and Sekimizu, K.: Purification of a GTP-binding protein localized in mitochondria. *J. Biochem. (Tokyo)*, 114 (1993) 684-690.
- 1697 Uematsu, J., Nishizawa, Y., Sonnenberg, A. and Owaribe, K.: Demonstration of type II hemidesmosomes in a mammary gland epithelial cell line, BMGE-H. *J. Biochem. (Tokyo)*, 115 (1994) 469-476.
- 1698 Wei, L.L. and Miner, R.: Evidence for the existence of a third progesterone receptor protein in human breast cancer cell line T47D. *Cancer Res.*, 54 (1994) 340-343.
- See also 1463, 1476, 1518, 1564, 1669.
- 19m. *Urinary proteins*
- 1699 Belova, L.A., Ogloblina, O.G. and Bekkert, R.: (A novel method for isolation of an acid-resistant trypsin inhibitor from human urine). *Biokhimiya (Moscow)*, 58 (1993) 1199-1205.
- 1700 Hossein-Nia, M., Mellotte, G.J., Foxall, P., Bending, M.R. and Holt, D.W.: Urinary protein excretion and the diagnosis of graft rejection or renal dysfunction in renal transplant patients. *Hum. Exp. Toxicol.*, 12 (1993) 407-409; *C.A.*, 120 (1994) 101064a.

- 1701 Perry, J.K., Scott, G.K. and Tse, C.A.: Modulation of proliferation of cultured human cells by urinary trypsin inhibitor. *Biochim. Biophys. Acta*, 1221 (1994) 145-152.
- 1702 Tang, Z., Xu, W., Yang, Z., Huang, Y., Mao, L. and Wu, H.: (Rapid determination of urinary proteins by SDS-PAGE and its clinical application). *Shengwu Huaxue Yu Shengwu Wuli Jinzhan*, 20 (1993) 324-327; *C.A.*, 120 (1994) 49356f.
- 1703 Toepfer, G., Pfisterer, J., Seifert, A. and Schaefer, P.: (Immunoblotting of urinary proteins after SDS-PAGE). *Klin. Labor.*, 39 (1993) 212-214; *C.A.*, 120 (1994) 49411v.
- 19n. *Other proteins (incl. proteinous inhibitors of enzymic activity)*
- 1704 Delisle, M.-C., Boyer, C., Vachon, V., Giroux, S. and Béliveau, R.: Immunodetection and characterization of proteins implicated in renal sodium/phosphate cotransport. *Biochim. Biophys. Acta*, 1190 (1994) 289-296.
- 1705 Jin, Z.X., Nakajima, T., Morisawa, M. and Hayashi, H.: Isolation and properties of a protein complex containing flagellar movement-initiating phosphoprotein from testes of a white salmon. *J. Biochem. (Tokyo)*, 115 (1994) 552-556.
- 1706 Kamei, S., Kamikubo, Y.-i., Hamuro, T., Fujimoto, H., Ishihara, M., Yonemura, H., Miyamoto, S., Funatsu, A., Enyoji, K.-i., Abumiya, T., Miyata, T. and Kato, H.: Amino acid sequence and inhibitory activity of Rhesus monkey tissue factor pathway inhibitor (TFPI): comparison with human TFPI. *J. Biochem. (Tokyo)*, 115 (1994) 708-714.
- 1707 Mendez, R. and de Haro, C.: Casein kinase II is implicated in the regulation of heme-controlled translational inhibitor of reticulocyte lysates. *J. Biol. Chem.*, 269 (1994) 6170-6176.
- 1708 Nara, K., Ito, S., Suzuki, Y., Ghoneim, M.A., Tachibana, S. and Hirose, S.: Elastase inhibitor elafin is a new type of proteinase inhibitor which has a transglutaminase-mediated anchoring sequence termed "cementoin". *J. Biochem. (Tokyo)*, 115 (1994) 441-448.
- 1709 Novak, A. and Lowden, J.A.: GM₂ Ganglioside activator occurs in multiple forms. *Biochim. Biophys. Acta*, 1199 (1994) 209-214.
- 1710 Tsuji, N., Kawazu, S., Nakamura, Y. and Fujisaki, K.: Protein analysis of *Strongyloides venezuelensis* by two-dimensional polyacrylamide gel electrophoresis. *J. Vet. Med. Sci.*, 55 (1993) 881-883; *C.A.*, 120 (1994) 186550d.
- 1711 Yusufi, A.N.K., Murayama, N., Gapstur, S.M., Szczepanska-Konkel, M. and Dousa, T.P.: Differential properties of brush-border membrane vesicles from early and late proximal tubules of rat kidney. *Biochim. Biophys. Acta*, 1191 (1994) 117-132.
- 1712 Zhang, N., Pan, N. and Chen, Z.: (Purification and partial characterization of an antagonistic protein Tzl). *Zhiwu Xuebao*, 35 (1993) 342-348; *C.A.*, 120 (1994) 72736p.
- 1713 Zhang, Z., Winyard, P.G., Chidwick, K., Murphy, G., Wardell, M., Carrell, R.W. and Blake, D.R.: Proteolysis of human native and oxidised α_1 -proteinase inhibitor by matrilysin and stromelysin. *Biochim. Biophys. Acta*, 1199 (1994) 224-228.
20. ENZYMES AND ENZYME ACTIVITY ESTIMATION
- 1714 Ohshima, M., Tanaka, S. and Ohshima, T.: (Application of polyacrylamide gel electrophoresis to the experiment for teaching of enzyme specificities). *Kagaku to Kyoiku*, 42 (1994) 137-141; *C.A.*, 120 (1994) 269181k.
- 20a. *Oxidoreductases*
- 1715 Buffoli, D.K., Scacco, S.C., Vergari, R., Solarino, G., Santacroce, G. and Papa, S.: Decline with age of the respiratory chain activity in human skeletal muscle. *Biochim. Biophys. Acta*, 1226 (1994) 73-82.
- 1716 Diamond, S., Rakestraw, D., O'Neil, J., Lam, G.N. and Christ, D.D.: Induction of cytochromes P-450 and 3A in mice following the dietary administration of the novel cognitive enhancer linoiridine. *Drug Metab. Disp.*, 22 (1994) 65-73.
- 1717 Fraser, P.D. and Bramley, P.M.: The purification of phytoene dehydrogenase from *Phycomyces blakesleeanae*. *Biochim. Biophys. Acta*, 1212 (1994) 59-66.
- 1718 Fujii, H., Shimomura, Y., Tokuyama, K. and Suzuki, M.: Modulation of branched-chain 2-oxo acid dehydrogenase complex activity in rat skeletal muscle by endurance training. *Biochim. Biophys. Acta*, 1199 (1994) 130-136.
- 1719 Guoqiang, D., Kaul, R. and Mattiasson, B.: Integration of aqueous two-phase extraction and affinity precipitation for the purification of lactate dehydrogenase. *J. Chromatogr. A*, 668 (1994) 145-152.
- 1720 Hada, T., Hagiya, H., Suzuki, H., Arakawa, T., Nakamura, M., Matsuda, S., Yoshimoto, T., Yamamoto, S., Azekawa, T., Morita, Y., Ishimura, K. and Kim, H.-Y.: Arachidonate 12-lipoxygenase of rat pineal glands: catalytic properties and primary structure deduced from its cDNA. *Biochim. Biophys. Acta*, 1211 (1994) 221-228.
- 1721 Jewett, S.L. and Rocklin, A.M.: N,N'-Diethylthiocarbamate as a stain for copper-zinc superoxide dismutase in polyacrylamide gels of red cell extracts. *Anal. Biochem.*, 217 (1994) 236-240.
- 1722 Kasai, S.: Preparation of P-flavin-bound and P-flavin-free luciferase and P-flavin-bound β -subunit of luciferase from *Photobacterium phosphoreum*. *J. Biochem. (Tokyo)*, 115 (1994) 670-674.
- 1723 Mandai, M., Iwaki, M. and Honda, Y.: Catalytic nature of tyrosine hydroxylase in bovine ocular tissues. *J. Biochem. (Tokyo)*, 115 (1994) 797-802.
- 1724 Nagai, A. and Ohta, D.: Histidinol dehydrogenase loses its catalytic function through the mutation of His²⁶¹→Asn due to its inability to ligate and essential Zn. *J. Biochem. (Tokyo)*, 115 (1994) 22-25.
- 1725 Nagai, A., Kheirolomoos, A. and Ohta, D.: Site-directed mutagenesis shows that the conserved cysteine residues of histidinol dehydrogenase are not essential for catalysis. *J. Biochem. (Tokyo)*, 114 (1993) 856-861.
- 1726 Nellaippan, K. and Vinayakam, A.: A method for demonstrating prophenoloxidase after electrophoresis. *Biotech. Histochem.*, 68 (1993) 193-195; *C.A.*, 120 (1994) 238747v.
- 1727 Packdibamrung, K., Misono, H., Harada, M., Nagata, S. and Nagasaki, S.: An inducible NADP⁺-dependent D-phenylserine dehydrogenase from *Pseudomonas syringae* NK-15: purification and biochemical characterization. *J. Biochem. (Tokyo)*, 114 (1993) 930-935.

See also 1701.

- 1728 Rashkovetsky, L.G., Maret, W. and Klyosov, A.A.: Human liver aldehyde dehydrogenases: new method of purification of the major mitochondrial and cytosolic enzymes and re-evaluation of their kinetic properties. *Biochim. Biophys. Acta*, 1205 (1994) 301-307.
- 1729 Reshef, A., Meiner, V., Berginer, V.M. and Leitersdorf, E.: Molecular genetics of cerebrotendinous xanthomatosis in Jews of North African origin. *J. Lipid Res.*, 35 (1994) 478-483.
- 1730 Usui, E., Okuda, K., Kato, Y. and Noshiro, M.: Rat hepatic 3α -hydroxysteroid dehydrogenase: expression of cDNA and physiological function in bile acid biosynthetic pathway. *J. Biochem. (Tokyo)*, 115 (1994) 230-237.
- 20b. *Transferases (excl. E.C. 2.7.-.-)*
- 1731 Andersson, M., Wettsten, M., Borén, J., Magnusson, A., Sjöberg, A., Rustaeus, S. and Olofsson, S.-O.: Purification of diacylglycerol: acyltransferase from rat liver to near homogeneity. *J. Lipid Res.*, 35 (1994) 535-545.
- 1732 Kerner, J., Zaluzec, E., Gage, D. and Bieber, L.L.: Characterization of the malonyl-CoA-sensitive carnitine palmitoyltransferase (CTP_o) of a rat heart mitochondrial particle. Evidence that the catalytic unit is CTP_o. *J. Biol. Chem.*, 269 (1994) 8209-8219.
- 1733 Nikol'skaya, I.I., Sharkova, E.V., Zenina, E.N., Lopatina, N.G., Atakhanova, B.A., Adylova, A., Koneeva, A.E., Kovalev, I.I. and Shishkin, S.S.: (DNA-methylases from human thyroid formations - nodular and diffuse goiters). *Biokhimiya (Moscow)*, 58 (1993) 1616-1624.
- 1734 QPech, L.L. and Nelson, D.L.: Purification and characterization of calmodulin (lysine 115) N-methyltransferase from *Paramecium tetraurelia*. *Biochim. Biophys. Acta*, 1199 (1994) 183-194.
- 1735 Radominska, A., Little, J.M., Lester, R. and Mackenzie, P.I.: Bile acid glucuronidation by rat liver microsomes and cDNA-expressed UDP-glucuronosyltransferases. *Biochim. Biophys. Acta*, 1205 (1994) 75-82.
- 1736 Tanaka, T., Yamamoto, S., Moriya, T., Taniguchi, M., Hayashi, H., Kagamiyama, H. and Oi, S.: Aspartate aminotransferase from a thermophilic formate-utilizing methanogen, *Methanobacterium thermoformicicum* strain SF-4: relation to serine and phosphoserine aminotransferases, but not to the aspartate aminotransferase family. *J. Biochem. (Tokyo)*, 115 (1994) 309-317.
- 1737 Tiainen, P. and Karhi, K.A.: Ultrasensitive time-resolved immunofluorometric assay of glutathione transferase alpha in serum. *Clin. Chem. (Washington)*, 40 (1994) 184-189.
- See also 1521, 1588.
- 20c. *Transferases transferring phosphorus containing groups (E.C. 2.7.-.-)*
- 1738 Bogoyevitch, M.A., Glennon, P.E., Andersson, M.B., Clerk, A., Lazou, A., Marshall, C.J., Parker, P.J. and Sugden, P.H.: Endothelin-1 and fibroblast growth factors stimulate the mitogen-activated protein kinase signaling cascade in cardiac myocytes. The potential role of the cascade in the integration of two signaling pathways leading to myocyte hypertrophy. *J. Biol. Chem.*, 269 (1994) 1110-1119.
- 1739 Chapman, A.B. and Agabian, N.: *Trypanosoma brucei* RNA polymerase II is phosphorylated in the absence of carboxyl-terminal domain heptapeptide repeats. *J. Biol. Chem.*, 269 (1994) 4754-4760.
- 1740 Cheley, S., Panchal, R.G., Carr, D.W., Scott, J.D. and Bayley, H.: Type II regulatory subunits of cAMP-dependent protein kinase and their binding proteins in the nervous system of *Aplysia californica*. *J. Biol. Chem.*, 269 (1994) 2911-2920.
- 1741 Childs, T.J. and Mak, A.S.: MAP kinases from bovine brain: purification and characterization. *Biochem. Cell Biol.*, 71 (1993) 544-555.
- 1742 Craig, L., Johnson, J.E. and Cornell, R.B.: Identification of the membrane-binding domain of rat liver CTP: phosphocholine cytidylyltransferase using chymotrypsin proteolysis. *J. Biol. Chem.*, 269 (1994) 3311-3317.
- 1743 Dementiev, A.A., Golyshin, P.N., Ryabchenko, N.F., Pustobaev, V.N. and Shlyapnikov, S.V.: (Two forms of extracellular low molecular mass ribonuclease of *Bacillus* sp. BCF 247: isolation and characterization of the enzyme). *Biokhimiya (Moscow)*, 58 (1993) 1258-1265.
- 1744 Edman, C.F. and Schulman, H.: Identification and characterization of $\delta\beta$ -CaM kinase and $\delta\gamma$ -CaM kinase from rat heart, two new multifunctional Ca^{2+} /calmodulin-dependent protein kinase isoforms. *Biochim. Biophys. Acta*, 1221 (1994) 89-101.
- 1745 Hanauske-Abel, H.M., Park, M.-H., Hanauske, A.-R., Popowicz, A.M., Lalonde, M. and Folk, J.E.: Inhibition of the G1-S transition of the cell cycle by inhibitors of deoxyhypusine hydroxylation. *Biochim. Biophys. Acta*, 1221 (1994) 115-124.
- 1746 Henderson, A.R., Stark, J.A., McQueen, M.J., Patten, R.L., Krishnan, S., Wood, D.E. and Webb, S.: Is determination of creatine kinase-2 after electrophoretic separation accurate? *Clin. Chem. (Washington)*, 40 (1994) 177-183.
- 1747 Kalab, P., Visconti, P., Leclerc, P. and Kopf, G.S.: p95, The major phosphotyrosine-containing protein in mouse spermatozoa, is a hexokinase with unique properties. *J. Biol. Chem.*, 269 (1994) 3810-3817.
- 1748 Kitani, T., Okuno, S. and Fujisawa, H.: cDNA cloning and expression of human calmodulin-dependent protein kinase IV. *J. Biochem. (Tokyo)*, 115 (1994) 637-640.
- 1749 Lichtinghagen, R., Diedrich-Glaubitz, R. and von Hörsten, B.: Identification of *Bordetella pertussis* in nasopharyngeal swabs using the polymerase chain reaction: evaluation of detection methods. *Eur. J. Clin. Chem. Clin. Biochem.*, 32 (1994) 161-167.
- 1750 Pisani, F.M. and Rossi, M.: Evidence that an archaeal α -like DNA polymerase has a modular organization of its associated catalytic activities. *J. Biol. Chem.*, 269 (1994) 7887-7892.
- 1751 Schrader, J., Gruwel, M., Decking, U. and Alves, C.: Wozu benötigen Endothelzellen Adenosintriphosphat? *Arzneim.-Forsch.*, 44 (1994) 436-438.
- 1752 Sedova, V.M. and Vorobyev, V.I.: (DNA-dependent RNA polymerase II from human placenta nuclei). *Biokhimiya (Moscow)*, 58 (1993) 1580-1586.
- 1753 Watanabe, H., Narumi, H., Inaba, T., Ohgi, K. and Irie, M.: Purification, some properties, and primary structure of a base non-specific ribonuclease from oyster (*Crassostrea grigus*). *J. Biochem. (Tokyo)*, 114 (1993) 800-807.

See also 1437, 1526, 1545, 1567.

20d. *Hydrolases, acting on ester bonds (E.C. 3.1.-.-)*

- 1754 Calderon de la Barca, A.M., Jensen, A.L. and Bog-Hansen, T.C.: Affinity methods with lectins: A tool to identify canine alkaline phosphatase isoenzymes. *J. Biochem. Biophys. Methods*, 27 (1993) 169-180; *C.A.*, 120 (1994) 72019g.
- 1755 Dormann, P., Kridl, J.C. and Ohlogge, J.B.: Cloning and expression in *Escherichia coli* of a cDNA coding for the oleoyl-acyl carrier protein thioesterase from coriander (*Coriandrum sativum* L.). *Biochim. Biophys. Acta*, 1212 (1994) 134-136.
- 1756 Kikuno, A., Kitada, M., Watanabe, S., Takeuchi, T., Komoda, T. and Sakagishi, Y.: (Comparative studies on serum ALP isoenzyme fractions isolated by conventional cellulose acetate membrane and polyacrylamide disc gel electrophoresis). *Seibutsu Butsuri Kagaku*, 38 (1994) 31-36; *C.A.*, 120 (1994) 264237y.
- 1757 Masuno, H. and Okuda, H.: Role of processing of the oligosaccharide chains in the affinity of lipoprotein lipase for heparin. *Biochim. Biophys. Acta*, 1212 (1994) 125-128.
- 1758 Nakanishi, Y., Okamoto, K. and Isohashi, F.: Subcellular distribution of ATP-stimulated and ADP-inhibited acetyl-CoA hydrolase in livers from control and clofibrate-treated rats: comparison of the cytosolic and peroxisomal enzyme. *J. Biochem. (Tokyo)*, 115 (1994) 328-332.
- 1759 Panteghini, M.: Electrophoretic fractionation of 5'-nucleotidase. *Clin. Chem. (Washington)*, 40 (1994) 190-196.

20e. *Hydrolases, acting on glycosyl compounds (E.C. 3.2.-.-)*

- 1760 Brochu, G., Shah, G.M. and Poirier, G.G.: Purification of poly(ADP-ribose) glycohydrolase and detection of its isoforms by a zymogram following one- or two-dimensional electrophoresis. *Anal. Biochem.*, 218 (1994) 265-272.
- 1761 Rizzotti, P. and Klein, G.: Evaluation of a specific immunoinhibition methods for the determination of pancreatic α -amylase. *Eur. J. Clin. Chem. Clin. Biochem.*, 32 (1994) 97-106.
- 1762 Tanaka, H., Ito, F. and Iwasaki, T.: Two sialidases which preferentially hydrolyze sialyl α 2-8 linkage from *Bacteroides fragilis* SBT3182. *J. Biochem. (Tokyo)*, 115 (1994) 318-321.

20f. *Other hydrolases*

- 1763 Agarwal, C., Hembree, J.R., Rorke, E.A. and Eckert, R.L.: Transforming growth factor β 1 regulation of metalloproteinase production in cultured human cervical epithelial cells. *Cancer Res.*, 54 (1994) 943-949.
- 1764 Kajihara, J.-i., Shibata, K., Nakano, Y., Nishimuro, S. and Kato, K.: Physicochemical characterization of PEG-PPG conjugated human urokinase. *Biochim. Biophys. Acta*, 1199 (1994) 202-208.
- 1765 Kleiner, D.E. and Stetler-Stevenson, W.G.: Quantitative zymography: detection of picogram quantities of gelatinases. *Anal. Biochem.*, 218 (1994) 325-329.
- 1766 McClelland, P., Adam, L.P. and Hathaway, D.R.: Identification of latent Ca^{2+} /calmodulin dependent protein kinase II phosphorylation site in vascular calpain II. *J. Biochem. (Tokyo)*, 115 (1994) 41-46.

- 1767 Ohtsuki, S., Homma, K.-i., Kurata, S., Komano, H. and Natori, S.: A prolyl endopeptidase of *Sarcophaga peregrina* (flesh fly): its purification and suggestion for its participation in the differentiation of the imaginal discs. *J. Biochem. (Tokyo)*, 115 (1994) 449-453.
- 1768 Seki, M., Yanagisawa, J., Kohda, T., Sonoyama, T., Ui, M. and Enomoto, T.: Purification of two DNA-dependent adenosinetriphosphatases having DNA helicase activity from HeLa cells and comparison of the properties of the two enzymes. *J. Biochem. (Tokyo)*, 115 (1994) 523-531.
- 1769 Shikata, S., Shimada, K., Ohnishi, Y., Horinouchi, S. and Beppu, T.: Characterization of secretory intermediates of *Serratia marcescens* serine protease produced during its extracellular secretion from *Escherichia coli* cells. *J. Biochem. (Tokyo)*, 114 (1993) 723-731.
- 1770 Shirasawa, Y., Osawa, T. and Hirashima, A.: Molecular cloning and characterization of prolyl endopeptidase from human T cells. *J. Biochem. (Tokyo)*, 115 (1994) 724-729.

See also 1519, 1523, 1853.

20g. *Lyases*

- 1771 Kuske, C.R., Ticknor, L.O., Guzmán, E., Gurley, L.R., Valdez, J.G., Thompson, M.E. and Jackson, P.J.: Purification and characterization of O-acetylserine sulfhydrylase isoenzymes from *Datura innoxia*. *J. Biol. Chem.*, 269 (1994) 6223-6232.
- 1772 Pecina, A. and Paneque, A.: Detection of alginate lyase by activity staining after sodium dodecyl sulfate-polyacrylamide gel electrophoresis and subsequent renaturation. *Anal. Biochem.*, 217 (1994) 124-127.
- 1773 Reddy, S., Cochran, B.J., Worth, L.L., Knutson, V.P. and Hadcox, M.K.: High-resolution slab gel isoelectric focusing: methods for quantitative electrophoretic transfer and immunodetection of proteins as applied to the study of the multiple isoelectric forms of ornithine decarboxylase. *Anal. Biochem.*, 218 (1994) 149-156.

20h. *Isomerases*

- 1774 Abe, I. and Prestwich, G.D.: Active site mapping affinity-labeled rat oxidosqualene cyclase. *J. Biol. Chem.*, 269 (1994) 802-804.
- 1775 Feldhoff, P.W., Mirski, S.E.L., Cole, S.P.C. and Sullivan, D.M.: Altered subcellular distribution of topoisomerase II α in a drug-resistant human small cell lung cancer cell line. *Cancer Res.*, 54 (1994) 756-762.
- 1776 Mukherjee, S.K., Reddy, M.K., Kumar, D. and Tewari, K.K.: Purification and characterization of a eukaryotic type 1 topoisomerase from pea chloroplast. *J. Biol. Chem.*, 269 (1994) 3793-3801.
- 1777 Nadal, M., Couderc, E., Duguet, M. and Jaxel, C.: Purification and characterization of reverse gyrase from *Sulfolobus shibatae*. Its proteolytic product appears as an ATP-independent topoisomerase. *J. Biol. Chem.*, 269 (1994) 5255-5263.
- 1778 Quéméneur, E., Guthapfel, R. and Gueguen, P.: A major phosphoprotein of the endoplasmic reticulum is protein disulfide isomerase. *J. Biol. Chem.*, 269 (1994) 5485-5488.

- 1779 Storl, K., Störl, J., Roth, M. and Zimmer, C.: DNA Topoisomerases from *Streptomyces noursei*: influence of coumarins and quinolones on the enzymic activity. *Biochim. Biophys. Acta*, 1199 (1994) 143-148.
- 1780 Yamazaki, H., Schneider, E., Myers, C.E. and Sinha, B.K.: Oncogene overexpression and de novo drug-resistance in human prostate cancer cells. *Biochim. Biophys. Acta*, 1226 (1994) 89-96.
- 20i. *Ligases*
- 1781 Barnes, T.S., Bleskan, J.H., Hart, I.M., Walton, K.A., Barton, J.W. and Patterson, D.: Purification of, generation of monoclonal antibodies to, and mapping of phosphoribosyl N-formylglycine amidotransferase. *Biochemistry*, 33 (1994) 1850-1860.
- 1782 Chan, K.W. and Koeppe, R.E.: Role of the TIGN sequence in *E. coli* tryptophanyl-tRNA synthetase. *Biochim. Biophys. Acta*, 1205 (1994) 223-229.
- 1783 Kerjan, P., Cerini, C., Sémériva, M. and Mirande, M.: The multi-enzyme complex containing nine aminoacyl-tRNA synthetases is ubiquitous from *Drosophila* to mammals. *Biochim. Biophys. Acta*, 1199 (1994) 293-297.
- 1784 Roberts, E., Nash, R.A., Robins, P. and Lindahl, T.: Different active sites of mammalian DNA ligases I and II. *J. Biol. Chem.*, 269 (1994) 3789-3792.
- 20j. *Complex mixtures and incompletely identified enzymes*
- 1785 Belogradov, G. and Hatfeji, Y.: Catalytic sector of complex I (NADH:ubiquinone oxidoreductase): subunit stoichiometry and substrate-induced conformation changes. *Biochemistry*, 33 (1994) 4571-4576.
- 1786 Leikin, A. and Shinitzky, M.: Shedding and isolation of the Δ -desaturase system from rat liver microsomes by application of high hydrostatic pressure. *Biochim. Biophys. Acta*, 1211 (1994) 150-155.
- 1787 Seo, H.G., Tatsumi, H., Fujii, J., Nishikawa, A., Suzuki, K., Kangawa, K. and Taniguchi, N.: Nitric oxide synthase from rat colon: purification, peptide sequencing, partial PCR cloning, and immunohistochemistry. *J. Biochem. (Tokyo)*, 115 (1994) 602-607.
- 1788 Shibata, H., Nagaoka, M., Sakai, M., Sawada, H., Watanabe, T. and Yokokura, T.: Kinetic studies on the plasminogen activation by the staphylokinase-plasmin complex. *J. Biochem. (Tokyo)*, 115 (1994) 738-742.
- 1789 Wilding, N., Mardell, S.K., Brookes, C.P. and Loxdale, H.D.: The use of polyacrylamide gel electrophoresis of enzymes to identify entomophthorean fungi in aphid hosts. *J. Invertebr. Pathol.*, 62 (1993) 268-272; *C.A.*, 120 (1994) 211844n.
21. PURINES, PYRIMIDINES, NUCLEIC ACIDS AND THEIR CONSTITUENTS
- 21a. *Purines, pyrimidines, nucleosides, nucleotides*
- 1790 Andrus, A.: Gel-capillary electrophoresis analysis of oligonucleotides. *Methods Mol. Biol. (Totowa)*, 26(Protocols for Oligonucleotide Conjugates) (1994) 277-300; *C.A.*, 120 (1994) 239457n.
- 1791 Desnoyers, S., Shah, G.M., Brochu, G. and Poirier, G.G.: Erasable blot of poly(ADP-ribose) polymerase. *Anal. Biochem.*, 218 (1994) 470-473.
- 1792 Lecoq, A.F., Montanarella, L. and di Biase, S.: Separation of nucleosides and nucleotide-3'-monophosphates by micellar electrokinetic capillary chromatography. *J. Microcolumn Sep.*, 5 (1993) 105-118; *C.A.*, 120 (1994) 157795g.
- 1793 Ramsey, R.S., Kerchner, G.A. and Cadet, J.: Micellar electrokinetic capillary chromatography of nucleic acid constituents and dinucleoside monophosphate photoproducts. *J. High Resolut. Chromatogr.*, 17 (1994) 4-8.
- 1794 Yin, H.F., Kleemiss, M.H., Lux, J.A. and Schomburg, G.: Diffusion coefficients of oligonucleotides in capillary gel electrophoresis. *J. Microcolumn Sep.*, 3 (1993) 331-335; *C.A.*, 120 (1994) 152647a.
- See also 1371, 1917.
- 21b. *Nucleic acids, RNA*
- 1795 Benhizia, F., Lagrange, D., Malewiak, M.-I. and Griglio, S.: *In vivo* regulation of hepatic lipase activity and mRNA levels by diets which modify cholesterol influx to the liver. *Biochim. Biophys. Acta*, 1211 (1994) 181-188.
- 1796 Chang, C.-D., Mukai, H., Kuno, T. and Tanaka, C.: cDNA Cloning of an alternatively spliced isoform of the regulatory subunit of Ca^{2+} /calmodulin-dependent protein phosphatase (calcineurin Ba2). *Biochim. Biophys. Acta*, 1217 (1994) 174-180.
- 1797 Chen, H. and Cui, H.: (Simple, fast, sensitive silver staining method for chemical synthetic oligonucleic acid). *Shengwu Huaxue Yu Shengwu Wuli Jinzhan*, 20 (1993) 482-483; *C.A.*, 120 (1994) 157839z.
- 1798 DesGroseillers, L., Auclair, D., Wickham, L. and Maalouf, M.: A novel actin cDNA is expressed in the neurons of *Aplysia californica*. *Biochim. Biophys. Acta*, 1217 (1994) 322-324.
- 1799 Fackelmayer, F.O. and Richter, A.: hnRNP-U/SAF-A is encoded by two differentially polyadenylated mRNAs in human cells. *Biochim. Biophys. Acta*, 1217 (1994) 232-234.
- 1800 Fujimoto, T., Fujisawa, J.-i. and Yoshida, M.: Novel isoforms of human cyclic AMP-responsive element modulator (hCREM) mRNA. *J. Biochem. (Tokyo)*, 115 (1994) 298-303.
- 1801 Gallagher, R.A., McClean, P.M. and Malik, A.N.: Cloning and nucleotide sequence of a full length cDNA encoding ribosomal protein L27 from human fetal kidney. *Biochim. Biophys. Acta*, 1217 (1994) 329-332.
- 1802 Hannan, R.D., Stennard, F.A. and West, A.K.: Localization of c-myc protooncogene expression in the rat heart *in vivo* and in the isolated perfused heart following treatment with norepinephrine. *Biochim. Biophys. Acta*, 1217 (1994) 281-290.
- 1803 Hitomi, K. and Tsukagoshi, N.: cDNA Sequence for rKST1, a novel member of the sodium ion-dependent glucose cotransporter family. *Biochim. Biophys. Acta*, 1190 (1994) 469-472.
- 1804 Lazarus, D.S., Aschoff, J., Fanburg, B.L. and Lanzillo, J.J.: Angiotensin converting enzyme (kininase II) mRNA production and enzymatic activity in human peripheral blood monocytes are induced by GM-CSF but not by other cytokines. *Biochim. Biophys. Acta*, 1226 (1994) 12-18.

- 1805 Maruyama, K., Endo, T., Kume, H., Kawamura, Y., Kanzawa, N., Kimura, S., Kawashima, S. and Maruyama, K.: A partial connectin cDNA encoding a novel type of RSP motifs isolated from chicken embryonic skeletal muscle. *J. Biochem. (Tokyo)*, 115 (1994) 147-149.
- 1806 Miyamoto, H., Homma, M. and Hotta, H.: Molecular cloning of the murine homologue of CD63/ME491 and detection of its strong expression in the kidney and activated macrophages. *Biochim. Biophys. Acta*, 1217 (1994) 312-316.
- 1807 Moriizumi, S., Watanabe, T., Unno, M., Nakagawara, K.-i., Suzuki, Y., Miyashita, H., Yonekura, H. and Okamoto, H.: Isolation, structural determination and expression of a novel reg gene, human reg β . *Biochim. Biophys. Acta*, 1217 (1994) 199-202.
- 1808 Negoro, N., Kanayama, Y., Iwai, J., Umetani, N., Nishimura, M., Konishi, Y., Okamura, M., Inoue, T. and Takeda, T.: Angiotensin-converting enzyme inhibitor increases angiotensin type 1A receptor gene expression in aortic smooth muscle cells of spontaneously hypertensive rats. *Biochim. Biophys. Acta*, 1226 (1994) 19-24.
- 1809 Raju, V.S. and Maines, M.D.: Coordinated expression and mechanism of induction of HSP32 (heme oxygenase-1) mRNA by hyperthermia in rat organs. *Biochim. Biophys. Acta*, 1217 (1994) 273-280.
- 1810 Rao, A. and Steward, O.: Evaluation of RNAs present in synaptodendrosomes: Dendritic, glial, and neuronal cell body contribution. *J. Neurochem.*, 61 (1993) 835-844; *C.A.*, 120 (1994) 101129a.
- 1811 Sakuma, T., Takai, M., Yokoi, T. and Kamataki, T.: Molecular cloning and sequence analysis of hamster CYP2E1. *Biochim. Biophys. Acta*, 1217 (1994) 229-231.
- 1812 Sato, S., Ozeki, Y. and Kawaguchi, A.: Transcription of the faoAB operon which encodes the HDT multienzyme complex involved in fatty acid β -oxidation in *Pseudomonas fragi* B-0771. *J. Biochem. (Tokyo)*, 115 (1994) 286-292.
- 1813 Stalbam, B.-M., Torvén, A. and Lundberg, L.G.: Application of capillary electrophoresis to the post-polymerase chain reaction analysis of rat mRNA for gastric H⁺,K⁺-ATPase. *Anal. Biochem.*, 217 (1994) 91-97.
- 1814 Sunitha, I. and Avigan, M.I.: A newly identified tyrosine kinase is preferentially expressed in the gastrointestinal tract. *Biochim. Biophys. Acta*, 1221 (1994) 348-352.
- 1815 Tarasov, I.A., Entelis, N.S. and Krashennikov, I.A.: (Mitochondrially-imported cytoplasmic lysine tRNA of the yeast: *in vitro* and *in vitro* test systems). *Biokhimiya (Moscow)*, 58 (1993) 1493-1502.
- 1816 Tomida, M., Yamamoto-Yamaguchi, Y. and Hozumi, M.: Three different cDNAs encoding mouse D-factor/LIF receptor. *J. Biochem. (Tokyo)*, 115 (1994) 557-562.
- 1817 Vambutas, V. and Wolgemuth, D.J.: Identification and characterization of the developmentally regulated pattern of expression in the testis of a mouse gene exhibiting similarity to the family of phosphodiesterases. *Biochim. Biophys. Acta*, 1217 (1994) 203-206.
- 1818 Wang, T.W.M., Byrne, C.D. and Hales, C.N.: Effect of ethanol on hepatic apolipoprotein B synthesis and secretion *in vitro*. *Biochim. Biophys. Acta*, 1211 (1994) 234-238.
- 1819 Williams, K.P., Fujimoto, D.N. and Inoue, T.: Two universally conserved adenosines of the group I intron that are important for self-splicing but not for core catalytic activity. *J. Biochem. (Tokyo)*, 115 (1994) 126-130.
- 1820 Yokoyama, K., Suzuki, H., Yasumoto, K.-i., Tomita, Y. and Shibahara, S.: Molecular cloning and functional analysis of a cDNA coding for human DOPACHrome tautomerase/tyrosinase-related protein-2. *Biochim. Biophys. Acta*, 1217 (1994) 317-321.
- 1821 Yu, L., Goldberg, I.H. and Dedon, P.C.: Eneidyne-mediated DNA damage in nuclei is modulated at the level of the nucleosome. *J. Biol. Chem.*, 269 (1994) 4144-4151.
- 1822 Zheng, Y., Wang, D., Fu, W., Chen, S., Liu, X., Yuan, P., Yan, P., Zhang, G. and Zhang, H.: (Examining rotavirus RNA with agarose electrophoresis). *Baiqiuwen Yike Daxue Xuebao*, 19 (1993) 507-509; *C.A.*, 120 (1994) 157953g.
- See also 1371, 1535, 1564, 1646, 1673, 1720, 1745, 1770, 1889, 1909.
- 21c. *Nucleic acids, DNA*
- 1823 Baba, Y.: (DNA analysis using capillary electrophoresis). *Bunseki*, (1993) 861-867; *C.A.*, 120 (1994) 100823k - a review with 13 refs.
- 1824 Baba, Y., Tomisaki, R., Sumita, C., Tanaka, A., Hide, K. and Tsuhako, M.: (High-resolution separation of single- and double-stranded DNA using capillary gel electrophoresis). *Bunseki Kagaku*, 42 (1993) 853-857; *C.A.*, 120 (1994) 186527b.
- 1825 Banas, B., Blaschke, D., Fittler, F. and Hörz, W.: Analysis of the promoter of the human prostatic acid phosphatase gene. *Biochim. Biophys. Acta*, 1217 (1994) 188-194.
- 1826 Bruin, T., Tuzgol, S., Mulder, W.J., van den Ende, A.E., Jansen, H., Hayden, M.R. and Kastelein, J.J.P.: A compound heterozygote for lipoprotein lipase deficiency, Val⁶⁹→Leu and Gly¹⁰⁰→Glu: correlation between *in vitro* LPL-activity and clinical expression. *J. Lipid Res.*, 35 (1994) 438-445.
- 1827 Buel, E. and Schwartz, M.: DAPI, a simple sensitive alternative to ethidium bromide staining of DNA in agarose gels. *Appl. Theor. Electrophor.*, 3 (1993) 253-255; *C.A.*, 120 (1994) 157838y.
- 1828 Chan, K.C.: Novel separation and detection methods of DNA fragments in electrophoresis. Avail. *Univ. Microfilms Int.*, Order No. DA9311482, 1992, 116 pp.; *C.A.*, 120 (1994) 265274v.
- 1829 Desbois, C., Magre, J., Blanquet, V., Capeau, J., Goossens, M. and Besmond, C.: Detection of sequence variations in the human insulin-receptor gene by parallel denaturing gradient gel electrophoresis. *Hum. Mutat.*, 2 (1993) 395-403; *C.A.*, 120 (1994) 262386x.
- 1830 Drolet, M., Bi, X. and Liu, L.F.: Hypernegative supercoiling of the DNA template during transcription elongation *in vitro*. *J. Biol. Chem.*, 269 (1994) 2068-2074.
- 1831 Duke, T.: Molecular mechanisms of DNA electrophoresis. *Int. J. Genome Res.*, 1 (1993) 227-247; *C.A.*, 120 (1994) 209551w.
- 1832 Duke, T. and Viovy, J.L.: Theory of DNA electrophoresis in physical gels and entangled polymer solutions. *Phys. Rev. E: Stat. Phys., Plasmas, Fluids, Relat. Interdiscip. Top.*, 49 (1994) 2408-2416; *C.A.*, 120 (1994) 239487x.
- 1833 Duke, T., Viovy, J.L. and Semenov, A.N.: Electrophoretic mobility of DNA in gels. I. New biased reptation theory including fluctuations. *Biopolymers*, 34 (1994) 239-247; *C.A.*, 120 (1994) 158106v.

- 1834 Ehrlich, K.C., Montalbano, B.G. and Thompson, C.: Use of disposable pipet tips to recover DNA from polyacrylamide gels by electroelution. *BioTechniques*, 15 (1993) 246-247; *C.A.*, 120 (1994) 49351a.
- 1835 Fukuda, K., Ichinose, M., Saiga, H., Shiokawa, K. and Yasugi, S.: Developmental changes of DNA methylation pattern of embryonic chick pepsinogen gene. *J. Biochem. (Tokyo)*, 115 (1994) 502-506.
- 1836 Gong, J., Tranganos, F. and Darzynkiewicz, Z.: A selective procedure for DNA extraction from apoptotic cells applicable for gel electrophoresis and flow cytometry. *Anal. Biochem.*, 218 (1994) 314-319.
- 1837 Gonzalez-Tizon, A.M., Martinez-Lage, A. and Mendez, J.: Chromosomal DNA extraction after restriction endonuclease treatments: a study by microdensitometry and electrophoresis. *Cytobios*, 74 (1993) 177-188; *C.A.*, 120 (1994) 158064e.
- 1838 Green, D.K.: Driving multiple pulsed field electrophoresis devices from one personal computer. *Comput. Appl. Biosci.*, 9 (1993) 593-596; *C.A.*, 120 (1994) 211838p.
- 1839 Harris, L.C., Remack, J.S. and Brent, T.P.: *In vitro* methylation of the human O⁶-methylguanine-DNA methyltransferase promoter reduces transcription. *Biochim. Biophys. Acta*, 1217 (1994) 141-146.
- 1840 Hebenbrock, K., Schuegerl, K. and Rinas, U.: Analysis of plasmid-DNA and cell protein of recombinant *Escherichia coli* using capillary and gel electrophoresis. A comparison of analytical methods. *DEHEMA Biotechnol. Conf.*, 5(Pt B) (1992) 849-852; *C.A.*, 120 (1994) 52685t.
- 1841 Heller, C., Duke, T. and Vivoy, J.L.: Electrophoretic mobility of DNA in gels. II. Systematic experimental study in agarose gels. *Biopolymers*, 34 (1994) 249-259; *C.A.*, 120 (1994) 157956k.
- 1842 Herrmann, J.L. and Saint Girons, I.: Value of pulsed-field gel electrophoresis for genetic and epidemiological studies of *Leptospira* and *Borrelia*. *Methods Mol. Cell. Biol.*, 4 (1993) 63-67; *C.A.*, 120 (1994) 126249c.
- 1843 Horie, K., Nishiguchi, S., Maeda, S. and Shimada, K.: Structures of replacement vectors for efficient gene targeting. *J. Biochem. (Tokyo)*, 115 (1994) 477-485.
- 1844 Hoshinaga, H., Amuro, N., Goto, Y. and Okazaki, T.: Molecular cloning and characterization of the rat cytochrome c oxidase subunit Vb gene. *J. Biochem. (Tokyo)*, 115 (1994) 194-201.
- 1845 Jin, S. and Zhang, W.: (An attempt of direct DNA hybridization on dried agarose gel). *Shengwu Huaxue Yu Shengwu Wuli Jinzhan*, 19 (1992) 240-241; *C.A.*, 120 (1994) 70332e.
- 1846 Kasai, K., Sakai, I., Yoshida, K., Mizuno, N., Sekiguchi, K., Sato, H. and Seta, S.: (Methodological investigation of separation of PCR amplification products of MCT118 (DIS80) locus using gel electrophoresis and determination of MCT118 type using DNA size marker. Comparison of typings using 123 bp ladder and Cetus allelic ladder). *Kagaku Keisatsu Kenkyusho Hokoku, Hokagaku-hen*, 46 (1993) 121-128; *C.A.*, 120 (1994) 263178t.
- 1847 Ke, S.H. and Wartell, R.M.: Influence of nearest neighbor sequence on the stability of base pair mismatches in long DNA: Determination by temperature-gradient gel electrophoresis. *Nucleic Acids Res.*, 21 (1993) 5137-5143; *C.A.*, 120 (1994) 101043t.
- 1848 Kodaira, M., Hiyama, K., Karakawa, T., Kameo, H. and Satoh, C.: Duplication detection in Japanese Duchenne muscular dystrophy patients and identification of carriers with partial gene deletions using pulsed-field gel electrophoresis. *Hum. Genet.*, 92 (1993) 237-243; *C.A.*, 120 (1994) 239477u.
- 1849 Kolble, K.: Electrophoresis of megabase DNA molecules. Avail. *Univ. Microfilms Int.*, Order No. BRD-96828, 1991, 163 pp.; *C.A.*, 120 (1994) 157934b.
- 1850 Kotaka, T., Adachi, S., Igarashi, K. and Shikata, T.: Dynamics of DNA molecules under gel electrophoresis. *ACS Symp. Ser.*, 548(Macro-iron Characterization) (1994) 466-478; *C.A.*, 120 (1994) 239459q.
- 1851 Lange, U., Steiner, S., Grolig, F., Wagner, G. and Philippsen, P.: Cloning and sequencing of a gene coding for an actin binding protein of *Saccharomyces exiguus*. *Biochim. Biophys. Acta*, 1217 (1994) 214-218.
- 1852 Li, Q. and Ownby, C.L.: A rapid method for extraction of DNA from agarose gels using a syringe. *BioTechniques*, 15 (1993) 976-978; *C.A.*, 120 (1994) 239498b.
- 1853 Lill, H., Steinemann, D. and Nelson, N.: Mutagenesis of the b'-subunit of *Synechocystis* sp. PCC 6803 ATP-synthase. *Biochim. Biophys. Acta*, 1184 (1994) 284-290.
- 1854 Lognonne, J.L.: Introduction to pulsed-field gel electrophoresis. *Methods Mol. Cell. Biol.*, 4 (1993) 49-55; *C.A.*, 120 (1994) 157717h.
- 1855 Lopez-Garcia, P., Abad, J.P. and Amils, R.: Genome analysis of different *Haloflexax mediterranei* strains using pulsed-field gel electrophoresis. *Syst. Appl. Microbiol.*, 16 (1993) 310-321; *C.A.*, 120 (1994) 158558u.
- 1856 Luque, I., Flores, E. and Herrero, A.: Nitrate and nitrite transport in the cyanobacterium *Synechococcus* sp. PCC 7942 are mediated by the same permease. *Biochim. Biophys. Acta*, 1184 (1994) 296-298.
- 1857 Magnusdottir, S., Aakerman, B. and Jonsson, M.: DNA electrophoresis in agarose gels: three regimes of DNA migration identified and characterized by the electrophoretic orientational behavior of DNA. *J. Phys. Chem.*, 98 (1994) 2624-2633; *C.A.*, 120 (1994) 157942c.
- 1858 Meyers, C.D., Fairbairn, D.W. and O'Neill, K.L.: Measuring the repair of hydrogen peroxide-induced DNA single strand breaks using the single cell gel assay. *Cytobios*, 74 (1993) 147-153; *C.A.*, 120 (1994) 155996z.
- 1859 Nagai, H., Ponglikitmongkol, M., Mita, E., Ohmachi, Y., Yoshikawa, H., Saeki, R., Yumoto, Y., Nakanishi, T. and Matsubara, K.: Aberration of genomic DNA in association with human hepatocellular carcinomas detected by 2-dimensional gel analysis. *Cancer Res.*, 54 (1994) 1545-1550.
- 1860 Popa, L.L., Plesa, A., Repanovici, R. and Popa, L.M.: (Electrophoresis of biotinylated nucleic acids). *Rev. Roum. Virol.*, 43 (1992) 41-46; *C.A.*, 120 (1994) 211841j.
- 1861 Schwartzman, J.B., Martinez-Robles, M.L. and Hernandez, P.: The migration behavior of DNA replicative intermediates containing an internal bubble analyzed by two-dimensional agarose gel electrophoresis. *Nucleic Acids Res.*, 21 (1993) 5474-5479; *C.A.*, 120 (1994) 48420k.
- 1862 Smith, C.L., Klco, S., Zhang, T.Y., Fang, H., Oliva, R., Wang, D., Bremer, M. and Lawrence S.: Analysis of megabase DNA using pulsed-field gel electrophoresis techniques. *Methods Mol. Genet.*, 2(Gene and Chromosome Analysis, Pt. B) (1993) 155-194; *C.A.*, 120 (1994) 70373u.

- 1863 Smith-Soerensen, B., Gebhardt, M.C., Kloen, P., McIntyre, J., Aguilar, F., Cerutti, P. and Boerresen, A.L.: Screening for TP53 mutations in osteosarcomas using constant denaturant gel electrophoresis (CDGE). *Hum. Mutat.*, 2 (1993) 274-285; *C.A.*, 120 (1994) 160229f.
- 1864 Sugaya, R., Ishimaru, S., Saigo, K. and Emori, Y.: Identification of a novel *Drosophila* gene encoding a Cdc2-related protein kinase. *J. Biochem. (Tokyo)*, 115 (1994) 150-155.
- 1865 Takematsu, H., Kawano, T., Koyama, S., Kozutsumi, Y., Suzuki, A. and Kawasaki, T.: Reaction mechanism underlying CMP-N-acetylneuraminic acid hydroxylation in mouse liver: formation of a ternary complex of cytochrome *b₅*, CMP-N-acetylneuraminic acid, and a hydroxylation enzyme. *J. Biochem. (Tokyo)*, 115 (1994) 381-386.
- 1866 Yang, S., Mao, Y., Wang, X., Xu, X. and Lin, X.: (Isolation and amplification of nonspecific PCR (polymerase chain reaction) products). *Fudan Xuebao, Ziran, Kexueban*, 32 (1993) 68-72; *C.A.*, 120 (1994) 237142g.
- 1867 Zhou, T., Li, Y., Liu, F. and Wen, J.: (A rapid and simple method for plasmid DNA purification). *Weishengwuxue Tongbao*, 20 (1993) 51-63; *C.A.*, 120 (1994) 49485x.
- See also 1308, 1310, 1436, 1535, 1749, 1779, 1780, 1797, 1798, 1801, 1812, 1889.
- 21d. *Structural studies on RNA and RNA mapping*
- 1868 Balakin, A.G., Lempicki, R.A., Huang, G.M. and Fournier, M.J.: *Saccharomyces cerevisiae* U14 small nuclear RNA has little secondary structure and appears to be produced by post-transcriptional processing. *J. Biol. Chem.*, 269 (1994) 739-746.
- 1869 Washio, K. and Ishikawa, K.: Cloning and sequencing of the gene for type I carboxypeptidase in rice. *Biochim. Biophys. Acta*, 1199 (1994) 311-314.
- 21e. *Structural studies on DNA and DNA mapping*
- 1870 Arakawa, H., Uetanaka, K., Maeda, M. and Tsuji, A.: Analysis of polymerase chain reaction product by capillary electrophoresis and its application to the detection of single base substitution in genes. *J. Chromatogr. A*, 664 (1994) 89-98.
- 1871 Baba, Y.: (Capillary electrophoresis. Capillary electrophoretic gene analysis). *Rinsho Kagaku (Nippon Rinsho Kagakkai)*, 22 (1993) 209-213; *C.A.*, 120 (1994) 209552x - a review with 11 refs.
- 1872 Baba, Y.: (Jurassic Park and chromatography, try to apply molecular recognizability of DNA to liquid chromatography (LC) and capillary electrophoresis (CE)). *Kuromatogurafi*, 14 (1993) 72-75; *C.A.*, 120 (1994) 70273m.
- 1873 Clark, B.K. and Sepaniak, M.J.: Evaluation of on-column labeling with intercalating dyes for fluorescence detection of DNA fragments separated by capillary electrophoresis. *J. Microcolumn Sep.*, 5 (1993) 275-282; *C.A.*, 120 (1994) 265106s.
- 1874 Deneer, H.G. and Knight, I.: Inhibition of the polymerase chain reaction by mucolytic agents. *Clin. Chem. (Washington)*, 40 (1994) 171-172.
- 1875 Dieguez-Lucena, J.L., Ruiz-Galdón, M., Morell-Ocaña, M., Garcia-Villanova, J., Flores-Polanco, F.J. and Reyes-Engel, A.: Capillary electrophoresis compared with silver staining of polyacrylamide gels for quantification of PCR products. *Clin. Chem. (Washington)*, 40 (1994) 493-494.
- 1876 Dolnik, V. and Novotny, M.: Capillary electrophoresis of DNA fragments in entangled polymer solutions: a study of separation variables. *J. Microcolumn Sep.*, 4 (1993) 515-519; *C.A.*, 120 (1994) 128869d.
- 1877 Drossman, H.: Construction and evaluation of a capillary electrophoresis DNA sequencer. Avail. *Univ. Microfilms Int.*, Order No. DA9221906, 1992, 461 pp.; *C.A.*, 120 (1994) 262430g.
- 1878 Fekete, C., Nagy, R., Debets, A.J.M. and Hornok, L.: Electrophoretic karyotypes and gene mapping in eight species of the *Fusarium* sections *Arthrosporiella* and *Sporotrichiella*. *Curr. Genet.*, 24 (1993) 500-504; *C.A.*, 120 (1994) 47174w.
- 1879 Fujii, H.: Fluorescent gel electrophoresis device for DNA sequence determination. *Jpn. Kokai Tokkyo Koho JP 06 18,416 [94 18,416]* (Cl. GO1N21/64), 25 Jan. 1994, Appl. 92/197,890, 30 Jun. 1992; 12 pp.; *C.A.*, 120 (1994) 237585d.
- 1880 Giffillan, A., Axton, R. and Brock, D.J.H.: Mass screening for cystic fibrosis heterozygotes: two assay systems compared. *Clin. Chem. (Washington)*, 40 (1994) 197-199.
- 1881 Grigsby, R.V., Fairbairn, D. and O'Neill, K.L.: Differential DNA damage detected in hybridomas. *Hybridoma*, 12 (1993) 755-761; *C.A.*, 120 (1994) 265110p.
- 1882 Grossman, P.D.: Electrophoretic separation of DNA sequencing extension products using low-viscosity entangled polymer networks. *J. Chromatogr. A*, 663 (1994) 219-227.
- 1883 Guendisch, C., Kirchhof, G., Baur, M., Bode, W. and Hartmann, A.: Identification of *Azospirillum* species by RFLP and pulsed-field gel electrophoresis. *Microb. Releases*, 2 (1993) 41-45; *C.A.*, 120 (1994) 211949a.
- 1884 Houshmand, M., Larsson, N.-G., Holme, E., Oldfors, A., Tulinius, M.H. and Andersen, O.: Automatic sequencing of mitochondrial tRNA genes in patients with mitochondrial encephalomyopathy. *Biochim. Biophys. Acta*, 1226 (1994) 49-55.
- 1885 Khrapko, K., Hanekamp, J.S., Thilly, W.G., Belenkii, A., Foret, F. and Karger, B.L.: Constant denaturant capillary electrophoresis (CDCE): a high resolution approach to mutational analysis. *Nucleic Acids Res.*, 22 (1994) 364-369; *C.A.*, 120 (1994) 237171r.
- 1886 Kievan, L.: DNA sizing control standards for electrophoretic analyses. *PCT Int. Appl. WO 94 02,647* (Cl. C12Q1/68), 03 Feb. 1994, US Appl. 918,765, 27 Jul. 1992; 28 pp.; *C.A.*, 120 (1994) 263049b.
- 1887 Lee, H.-C., Pang, C.-Y., Hsu, H.-S. and Wei, Y.-H.: Differential accumulations of 4,977 bp deletion in mitochondrial DNA of various tissues in human ageing. *Biochim. Biophys. Acta*, 1226 (1994) 37-43.
- 1888 Liang, C. and Gerbi, S.A.: Analysis of an origin of DNA amplification in *Sciara coprophila* by a novel three-dimensional gel method. *Mol. Cell. Biol.*, 14 (1994) 1520-1529; *C.A.*, 120 (1994) 184147d.
- 1889 Lightfoot, T. and Snyder, F.F.: Gene amplification and dual point mutations of mouse IMP dehydrogenase associated with cellular resistance to mycophenolic acid. *Biochim. Biophys. Acta*, 1217 (1994) 156-162.
- 1890 Lin, K.-C. and Shiuian, D.: DNase I footprinting studies of *Escherichia coli* biotin repressor-operator interactions. *J. Biochem. (Tokyo)*, 114 (1993) 670-676.

- 1891 Matvienko, N.N., Kramarov, V.M., Zheleznyaya, L.A. and Matvienko, N.I.: (New site-specific endonuclease and methylase from *Bacillus licheniformis* 736). *Biokhimiya (Moscow)*, 58 (1993) 1139-1153.
- 1892 Mills, J.B., Cooper, J.P. and Hagerman, P.J.: Electrophoretic evidence that single-stranded regions of one or more nucleotides dramatically increase the flexibility of DNA. *Biochemistry*, 33 (1994) 1797-1803.
- 1893 Mochizuki, T.: Electrophoretic apparatus for DNA sequencing. *Eur. Pat. Appl. EP 572,023* (Cl. G01N27/447), 01 Dec. 1993, JP Appl. 92/138,568, 29 May 1992; 7 pp.; C.A., 120 (1994) 70893p.
- 1894 Noda, C., Fukushima, C., Fujiwara, T., Matsuda, K., Kobune, Y. and Ichihara, A.: Developmental regulation of rat serine dehydratase gene expression: evidence for the presence of a repressor in fetal hepatocytes. *Biochim. Biophys. Acta*, 1217 (1994) 163-173.
- 1895 Pannetier, C.: Method for determining the nucleotide size of DNA fragments. *PCT Int. Appl. WO 94 01,581* (Cl. C12Q1/68), 20 Jan. 1994, FR Appl. 92/8,253, 03 Jul. 1992; 34 pp.; C.A., 120 (1994) 155893p.
- 1896 Papavassiliou, A.G.: In gel DNase I protection analysis of electrophoretically fractionated DNA-binding reactions. *Methods Mol. Cell. Biol.*, 4 (1993) 95-104; C.A., 120 (1994) 184140w.
- 1897 Pugh, C.W., Ebert, B.L., Ebrahim, O. and Ratcliffe, P.J.: Characterisation of functional domains within the mouse erythropoietin 3' enhancer conveying oxygen-regulated responses in different cell lines. *Biochim. Biophys. Acta*, 1217 (1994) 297-306.
- 1898 Richard, P., Thomas, G., de Zulueta, M.P., de Gennes, J.-L., Thomas, M., Cassaigne, A., Béréziat, G. and Iron, A.: Common and rare genotypes of human apolipoprotein E determined by specific restriction profiles of polymerase chain reaction-amplified DNA. *Clin. Chem. (Washington)*, 40 (1994) 24-29.
- 1899 Rocheleau, M.J. and Dovichi, N.J.: Separation of DNA sequencing fragments at 53 bases/minute by capillary gel electrophoresis. *J. Microcolumn Sep.*, 4 (1992) 449-453; C.A., 120 (1994) 97343w.
- 1900 Scholz, R.B., Milde-Langosch, K., Jung, R., Schlechte, H., Kabisch, H., Wagener, C. and Loening, T.: Rapid screening for Tp53 mutations by temperature gradient gel electrophoresis: a comparison with SSCP analysis. *Hum. Mol. Genet.*, 2 (1993) 2155-2158; C.A., 120 (1994) 70383x.
- 1901 Schoonmaker, D. and Kondracki, S.F.: Investigation of nosocomial legionellosis using restriction enzyme analysis by pulsed-field gel electrophoresis. In: Barbaree, J.M., Breiman, R.F. and Dufour, A.P. (Editors), *Legionella*, Am. Soc. Microbiol., Washington, 1993, 180-194; C.A., 120 (1994) 49171s.
- 1902 Shifman, M.A.: PFGE MAPPER: a tool to aid in the analysis of pulse field gel electrophoresis maps. *Comput. Appl. Biosci.*, 9 (1993) 511-515; C.A., 120 (1994) 262388z.
- 1903 Thiele, D., Willems, H., Koepf, G. and Krauss, H.: Polymorphism in DNA restriction patterns of *Coxiella burnetii* isolates investigated by pulsed field gel electrophoresis and image analysis. *Eur. J. Epidemiol.*, 9 (1993) 419-425; C.A., 120 (1994) 184155e.
- 1904 Tsongalis, G.J., Kaufmann, W.K., Wilson, S.J., Friedman, K.J. and Silverman, L.M.: Rapid screening for p53 mutations with a sensitive heteroduplex detection technique. *Clin. Chem. (Washington)*, 40 (1994) 485-486.
- 1905 Tsongalis, G.J., McPhail, A.H., Lodge-Rigal, R.D., Chapman, J.F. and Silverman, L.M.: Localized *in situ* amplification (LISA): a novel approach to *in situ* PCR. *Clin. Chem. (Washington)*, 40 (1994) 381-384.
- 1906 Vom Stein, J.: (New standards for automated central DNA analysis). *BioTec (Wuerzburg)*, 5 (1993) 38-43; C.A., 120 (1994) 70342h.
- 1907 Walker, P.R., Kokileva, L., LeBlanc, J. and Sikorska, M.: Detection of the initial stages of DNA fragmentation in apoptosis. *Bio-Techniques*, 15 (1993) 1032-1040; C.A., 120 (1994) 239453h.
- 1908 Wang, G., Whittam, T.S., Berg, C.M. and Berg, D.E.: RAPD (arbitrary primer) PCR is more sensitive than multilocus enzyme electrophoresis for distinguishing related bacterial strains. *Nucleic Acids Res.*, 21 (1993) 5930-5933; C.A., 120 (1994) 155115m.
- 1909 Watanabe, Y., Kawakami, K., Hirayama, Y. and Nagano, K.: Transcription factors positively and negatively regulating the Na,K-ATPase $\alpha 1$ subunit gene. *J. Biochem. (Tokyo)*, 114 (1993) 849-855.
- 1910 Weizenegger, M.: (Fluorescence sequencing in the IR range). *BioTec (Wuerzburg)*, 5 (1993) 18-20; C.A., 120 (1994) 70339n.
- 1911 Weizenegger, M. and Pohl, T.M.: (Nonradioactive sequencing with direct-blotting-electrophoresis). *BioTec (Wuerzburg)*, 4 (1992) 27-30; C.A., 120 (1994) 70341g.
- 1912 Xing, G., He, F., Qu, C. and Wu, Z.: (Separation of DNA fragments from agarose gel by centrifugation). *Junshi Yixue Kexueyuan Yuankan*, 17 (1993) 304-306; C.A., 120 (1994) 128751j.
- 1913 Zhang, J., Chen, D.Y., Harke, H.R. and Dovichi, N.: The sheath-flow cuvette in DNA sequencing by capillary gel electrophoresis and two-spectral-channel, laser-induced fluorescence detection. *Chromatogr. Sci. Ser.*, 64(Capillary Electrophoresis Technology) (1993) 631-641; C.A., 120 (1994) 70350j.

See also 1490, 1574, 1825, 1826, 1837, 1844, 1858.

21f. Complex mixtures of nucleic acids and their fragments

- 1914 Schaefer, P., Braun, R.W., Moehring, K., Henco, K., Kang, J., Wendland, T. and Kuhn, J.E.: Quantitative determination of human cytomegalovirus target sequences in peripheral blood leukocytes by nested polymerase chain reaction and temperature gradient gel electrophoresis. *J. Gen. Virol.*, 74 (1993) 2699-2707; C.A., 120 (1994) 154999r.

22. ALKALOIDS

- 1915 Hsu, P.C. and Sheu, S.J.: (A comparative study on the processed Ephedrae herba). *Zhonghua Yaoxue Zazhi*, 45 (1993) 483-489; C.A., 120 (1994) 62412r.
- 1916 Hurst, W.J. and Martin, R.A., Jr.: The quantitative determination of caffeine in beverages using capillary electrophoresis. *Anal. Chem.*, 21 (1993) 389-391; C.A., 120 (1994) 161892d.
- 1917 Sun, P., Mariano, G.J., Barker, G. and Hartwick, R.A.: Comparison of micellar electrokinetic capillary chromatography and high-performance liquid chromatography on the separation and determination of caffeine and its analogues in pharmaceutical tablets. *Anal. Lett.*, 27 (1994) 927-937.

23. OTHER SUBSTANCES CONTAINING HETEROCYCLIC NITROGEN

23a. Porphyrins and other pyrroles

- 1918 Buchler, J.W., Künzel, F.M., Mayer, U. and Nawra, M.: Metal complexes with tetrapyrrole ligands. Part 64. Electrophoresis of water-soluble porphyrins and their metal complexes. *Fresenius J. Anal. Chem.*, 348 (1994) 371-376.

23b. Bile pigments

- 1919 Wu, N., Sweedler, J.V. and Lin, M.: Enhanced separation and detection of serum bilirubin species by capillary electrophoresis using a mixed anionic surfactant-protein buffer system with laser-induced fluorescence detection. *J. Chromatogr. B*, 654 (1994) 185-191.

23e. Other N-heterocyclic compounds

- 1920 Thuncke, F., Hartenstein, H., Sicker, D. and Vogt, C.: Separation of enantiomers and diastereomers of 4-hydroxy-2H-1,4-benzoxazin-3(4H)-one derivatives by capillary electrophoresis. *Chromatographia*, 38 (1994) 470-474.

24. ORGANIC SULPHUR COMPOUNDS (INCL. GLUCOSINOLATES)

- 1921 Feldl, C., Moller, P., Otte, J. and Sorensen, H.: Micellar electrokinetic capillary chromatography for determination of indolyl glucosinolates and transformation products thereof. *Anal. Biochem.*, 217 (1994) 62-69.

See also 1323.

25. ORGANIC PHOSPHORUS COMPOUNDS (INCL. SUGAR PHOSPHATES)

See 1496, 1501, 1502, 1548, 1552, 1555, 1556, 1686.

26. ORGANOMETALLIC AND RELATED COMPOUNDS

26a. Organometallic compounds

- 1922 Medina, I., Rubi, E., Mejuto, M.C. and Cela, R.: Speciation of organomercurials in marine samples using capillary electrophoresis. *Talanta*, 40 (1993) 1631-1636; C.A., 120 (1994) 70989z.

See also 1918.

26b. Boranes, silanes and related non-metallic compounds

- 1923 Kobayashi, M. and Kitaoka, Y.: (Electrophoretic behavior and infrared absorption spectra of dihydroxyboryl compounds in aqueous di- and tricarboxylic acids). *Kyoto Daigaku Genshiro Jikkensho, [Tech. Rep.]*, KURRI-TR-382 (1993) 47-54; C.A., 120 (1994) 264181a.

26c. Coordination compounds

- 1924 Mishra, U. and Singh, R.K.P.: Ionophoretic technique in the study of Cu(II), Ni(II), Co(II) and Mn(II)-hydrazine-nitritotriacetate mixed-ligand complexes. *J. Chromatogr. A*, 667 (1994) 371-376.

28. ANTIBIOTICS

- 1925 Flurer, C.L. and Wolnik, K.A.: Quantitation of gentamicin sulfate in injectable solutions by capillary electrophoresis. *J. Chromatogr. A*, 663 (1994) 259-263.
- 1926 Gu, J. and Fu, R.: Separation of chiral isomers of *p*-nitrophenyl-2-amino-3-hydroxypropanone by capillary zone electrophoresis using cyclodextrins as chiral selector. *J. Chromatogr. A*, 667 (1994) 367-370.
- 1927 Rybin, V.K., Kozlova, N.V. and Makarov, N.V.: (Determination of the reaction product of rubomycin with stelin B in a paper electrophoresis method). *Zh. Anal. Khim.*, 48 (1993) 144-146; C.A., 120 (1994) 200508e.

30. SYNTHETIC AND NATURAL DYES

30a. Synthetic dyes

- 1928 Gasparič, J. and Sedmíková, A.: Effect of the number of sulpho groups on the electrophoretic mobility of sulphonated azo dyes. *J. Chromatogr. A*, 665 (1994) 197-202.

See also 1958.

30b. Chloroplast and other natural pigments

- 1929 Rigol, A., López-Sánchez, J.F. and Rauret, G.: Capillary zone electrophoresis of humic acids. *J. Chromatogr. A*, 664 (1994) 301-305.
- 1930 Trubetskoi, O.A., Kudryavtseva, L.Yu. and Shirshova, L.T.: (Soil humic substance fractionation by electrophoresis in polyacrylamide gel in the presence of denaturing agents). *Pochvo-vedenie*, No. 8 (1993) 122-126; C.A., 120 (1994) 190378w.

See also 1343.

31. PLASTICS AND THEIR INTERMEDIATES

- 1931 Anderson, J.L. and Solomentsev, Y.: Electrophoresis of nonuniformly charged chains. *Polym. Prepr. (Am. Chem. Soc., Div. Polym. Chem.)*, 34 (1993) 1061-1062; C.A., 120 (1994) 192835s.
- 1932 Aoyanagi, O., Muramatsu, N., Ohshima, H. and Kondo, T.: Electrophoretic behavior of poly A-graft-poly B-type microcapsules. *J. Colloid Interface Sci.*, 162 (1994) 222-226; C.A., 120 (1994) 1193467k.

32. DRUG ANALYSIS

32a. *Drug analysis, general techniques*

- 1933 Altria, K.D., Goodal, D.M. and Rogan, M.M.: Quantitative determination of drug counter-ion stoichiometry by capillary electrophoresis. *Chromatographia*, 38 (1994) 637-642.
- 1934 Francotte, E., Cherkaoui, S. and Faupel, M.: Separation of the enantiomers of some racemic nonsteroidal aromatase inhibitors and barbiturates by capillary electrophoresis. *Chirality*, 5 (1993) 516-526; *C.A.*, 120 (1994) 173604z.
- 1935 Imazawa, M.: (Capillary electrophoresis. Analysis of drugs in body fluids). *Rinsho Kagaku (Nippon Rinsho Kagakkai)*, 22 (1993) 225-229; *C.A.*, 120 (1994) 182155f - a review with 19 refs.
- 1936 Pálmarsdóttir, S. and Edholm, L.-E.: Capillary zone electrophoresis for separation of drug enantiomers using cyclodextrins as chiral selectors. Influence of experimental parameters on separation. *J. Chromatogr. A*, 666 (1994) 337-350.
- 1937 Smith, N.W. and Evans, M.B.: The analysis of pharmaceutical compounds using electrochromatography. *Chromatographia*, 38 (1994) 649-657.
- 1938 Thormann, W.: Capillary electrophoresis and electrokinetic capillary chromatography of drugs in body fluids. *Chromatogr. Sci. Ser.*, 64(Capillary Electrophoresis Technology) (1993) 693-704; *C.A.*, 120 (1994) 44923d - a review with 39 refs.
- 1939 Tomlinson, A.J., Benson, L.M. and Naylor, S.: Advantages of nonaqueous solvents in the analysis of drug metabolites using CE and online CE-MS. *LC-GC*, 12 (1994) 122-130; *C.A.*, 120 (1994) 182212x.
- 1940 Williams, R.C., Edwards, J.F. and Ainsworth, C.R.: Analysis of diastereoisomer impurities in chiral pharmaceutical compounds by capillary electrophoresis. *Chromatographia*, 38 (1994) 441-446.
- 1941 Wong, S.H.Y.: Advances in chromatography for clinical drug analysis: supercritical fluid chromatography, capillary electrophoresis, and selected high-performance liquid chromatography techniques. *Ther. Drug. Monit.*, 15 (1993) 576-580; *C.A.*, 120 (1994) 260339k - a review with 32 refs.

See also 1498.

32b. *Antirheumatics and antiinflammatory drugs*

- 1942 Donato, M.G., Baeyens, W., van den Bossche, W. and Sandra, P.: The determination of non-steroidal antiinflammatory drugs in pharmaceuticals by capillary zone electrophoresis and micellar electrokinetic capillary chromatography. *J. Pharm. Biomed. Anal.*, 12 (1994) 21-26.
- 1943 Karger, A.E., Stoll, E. and Hänsel, W.: Direkte chirale Trennung von Tiaprofen, Carprofen und Flurbiprofen durch Cyclodextrin-modifizierte Kapillarelektrophorese. *Pharmazie*, 49 (1994) 155-159.

32c. *Autonomic and cardiovascular drugs*

- 1944 Dethy, J.-M., de Broux, S., Lesne, M., Longstreth, J. and Gilbert, P.: Stereoselective determination of verapamil and norverapamil by capillary electrophoresis. *J. Chromatogr. B*, 654 (1994) 121-127.

- 1945 Nasi, J.C. and Francasso, F.A.R.: (Electrophoresis of β -adrenergic antagonists). *Rev. Bras. Farm.*, 74 (1993) 7-8; *C.A.*, 120 (1994) 62413s.
- 1946 Quang, C. and Khaledi, M.G.: Direct separation of the enantiomers of β -blockers by cyclodextrin-mediated capillary zone electrophoresis. *J. High Resolut. Chromatogr.*, 17 (1994) 99-101.
- 1947 Ricci, M.C. and Cross, R.F.: Capillary electrophoresis separation of sulfonamides and dihydrofolate reductase inhibitors. *J. Microcolumn Sep.*, 5 (1993) 207-215; *C.A.*, 120 (1994) 200515e.
- 1948 Shafaati, A. and Clark, B.J.: Development of capillary zone electrophoresis method for atenolol and its related impurities in a tablet preparation. *Anal. Proc.*, 30 (1993) 481-483. *C.A.*, 120 (1994) 173617f.

See also 1361.

32d. *Central nervous system drugs*

See 1358, 1940.

32e. *Chemotherapeutics (exc. cytostatics and antibiotics)*

See 1361.

32f. *Cytostatics*

- 1949 Chan, K.C., Muschik, G.M., Issaq, H.J. and Snader, K.M.: Separation of taxol and related compounds by micellar electrokinetic chromatography. *J. High Resolut. Chromatogr.*, 17 (1994) 51-52.

32g. *Other drug categories*

- 1950 Bodiroga, M., Agbaba, D., Zivanov-Stakic, D. and Popovic, R.: Determination of pralidoxime chloride in pharmaceutical dosage forms by isotachophoresis. *J. Pharm. Biomed. Anal.*, 12 (1994) 127-129.
- 1951 Dette, C., Wätzig, H. and Aigner, A.: Analysis of acetylcysteine by capillary electrophoresis (CE). Part 1: Control of optical purity. *Pharmazie*, 49 (1994) 245-248.
- 1952 Trenerry, V.C., Wells, R.J. and Robertson, J.: The analysis of illicit heroin seizures by capillary zone electrophoresis. *J. Chromatogr. Sci.*, 32 (1994) 1-6.
- 1953 Wätzig, H., Dette, C., Aigner, A. and Willschowitz, L.: Analysis of acetylcysteine by capillary electrophoresis (CE). Part 2: Determination of side components. *Pharmazie*, 49 (1994) 249-252.

32h. *Toxicological and forensic applications*

See 1619, 1952.

32i. *Plant extracts*

- 1954 Hines, H.B. and Brueggemann, E.E.: Behavior of ricin on untreated and treated capillary electrophoresis columns. *Chromatogr. Sci. Ser.*, 64(Capillary Electrophoresis Technology) (1993) 819-841; *C.A.*, 120 (1994) 70987x.

- 1955 Lu, C.F. and Sheu, S.J.: (Determination of amugdalin in traditional Chinese medicinal preparation by capillary electrophoresis). *Zhonghua Yaoxue Zazhi*, 45 (1993) 495-501; C.A., 120 (1994) 173577t.

See also 1915.

33. CLINICO-CHEMICAL APPLICATIONS

33a. General papers and reviews

- 1956 De Antonis, K.M. and Brown, P.R.: Determination of biochemical markers by HPLC and CE for the diagnosis of diseases. *Am. Lab. (Shelton)*, 25 (1993) 62-71; C.A., 120 (1994) 186335n - a review with 35 refs.

See also 1380.

33b. Complex mixtures and profiling (single compounds by cross-reference only)

- See 1480, 1481, 1484, 1485, 1486, 1490, 1625, 1629, 1729, 1737, 1746, 1759, 1761, 1826, 1880, 1898, 1904, 1905.

34. FOOD ANALYSIS

See 1916.

34b. Complex mixtures (single compounds by cross-reference only)

- 1957 Cancalon P.F.: Rapid monitoring of fruit juice adulteration by capillary electrophoresis. *LC-GC*, 11 (1993) 748-751; C.A., 120 (1994) 75750z.
- 1958 Tomlinson, A.J., Mlotkiewicz, J.A. and Lewis, I.A.S.: Application of capillary electrophoresis to the separation of colored products of Maillard reactions. *Food Chem.*, 49 (1994) 219-223; C.A., 120 (1994) 189989q.

See also 1469.

35. ENVIRONMENTAL ANALYSIS

35c. Water pollution (complex mixtures; single compounds by cross-reference only)

See 1977.

36. SOME TECHNICAL PRODUCTS AND COMPLEX MIXTURES

36c. Complex mixtures, technical products and unidentified compounds

- 1959 Ishida, H.: (Electrophoretic forming and (its) control factors). *Kobutsugaku Zasshi*, 22 (1993) 79-86; C.A., 120 (1994) 114157h - a review with 23 refs.
- 1960 Kelly, R.G., Scully, H.S. and Stoner, G.E.: Using capillary electrophoresis for failure analysis. *Mater Perform.*, 32 (1993) 60-63; C.A., 120 (1994) 94174f.

See also 1410, 1414, 1633, 1636.

37. CELLS, CELLULAR PARTICLES AND SUPRAMOLECULAR STRUCTURES

- 1961 Aki, M., Shimbara, N., Takashina, M., Akiyama, K., Kagawa, S., Tamura, T., Tanahashi, N., Yoshimura, T., Tanaka, K. and Ichihara, A.: Interferon- γ induces different subunit organizations and functional diversity of proteasomes. *J. Biochem. (Tokyo)*, 115 (1994) 257-269.
- 1962 Fortsas, E., Petric, M. and Brown, M.: Electrophoretic migration of adenovirus hexon under non-denaturing conditions. *Virus Res.*, 31 (1994) 57-65; C.A., 120 (1994) 212161z.
- 1963 Kraevskii, V.A., Panin, V.M. and Razin, S.V.: (Analysis of chromatin decondensation by means of gel electrophoresis in low-density agarose). *Mol. Biol. (Moscow)*, 28 (1994) 76-81; C.A., 120 (1994) 211853q.
- 1964 Morré, D.J., Lawrence, J., Safranski, K., Hammond, T. and Morré, D.M.: Experimental basis for separation of membrane vesicles by preparative free-flow electrophoresis. *J. Chromatogr. A*, 668 (1994) 201-213.
- 1965 Walter, H. and Widen, K.E.: Cell partitioning in two-polymer aqueous phase systems and cell electrophoresis in aqueous polymer solutions. Red blood cells from different species. *J. Chromatogr. A*, 668 (1994) 185-190.

38. INORGANIC COMPOUNDS

38a. Cations

- 1966 Baechmann, K., Steeg, K.H., Groh, T., Haumann, I., Boden, J. and Holthues, H.: Analysis for cations and anions in small volumes using micro-LC and CE. *J. Microcolumn Sep.*, 4 (1992) 431-438; C.A., 120 (1994) 123733v.
- 1967 Boden, J., Ehmann, T., Groh, T., Haumann, I. and Bächmann, A.: Influencing of the migration time in CZE. *Fresenius J. Anal. Chem.*, 348 (1994) 572-575.
- 1968 Hirokawa, T., Xia, W., Nakamura, K.-i., Tanaka, I., Nishiyama, F., Kiso, Y., Gaš, B. and Vacík, J.: Study of isotachophoretic separation behaviour of metal cations by means of particle-induced X-ray emission. VI. Selective separation of twenty metal cations using tartaric acid as a complexing agent. *J. Chromatogr. A*, 663 (1994) 245-254.

- 1969 Majmudar, A.A. and Manohar, C.: Electrophoretic separation of dilute TiO₂ suspension. *Sep. Sci. Technol.*, 29 (1994) 845-854; C.A., 120 (1994) 80624r.
- 1970 Mang, M., Gehmecker, H., Trautmann, N. and Herrmann, G.: Separation of oxidation states of neptunium and plutonium by continuous electrophoretic ion focusing. *Radiochim. Acta*, 62 (1993) 49-54; C.A., 120 (1994) 146813k.
- 1971 Nasi, J.C. and Luiz, A.J.: (Electrophoresis of ions: iron(2+), cobalt(2+), copper(2+), tin(2+), lead(2+) and antimony(2+)). *Rev. Bras. Farm.*, 74 (1993) 47-49; C.A., 120 (1994) 152373h.
- 1972 Šimuničová, E., Kaniansky, D. and Lokšíková, K.: Separation of alkali and alkaline earth metal and ammonium cations by capillary zone electrophoresis with indirect UV absorbance detection. *J. Chromatogr. A*, 655 (1994) 203-209.
- 1974 Jimidar, M., Hamoir, T., Degezelle, W., Massart, D.L., Soykenç, S. and van de Winkel, P.: Method development and optimization for the determination of rare earth metal ions by capillary zone electrophoresis. *Anal. Chim. Acta*, 284 (1993) 217-225.
- 1975 Leone, A.M., Francis, P.L., Rhodes, P. and Moncada, S.: A rapid and simple method for the measurement of nitrite and nitrate in plasma by high performance capillary electrophoresis. *Biochem. Biophys. Res. Commun.*, 200 (1994) 951-957.
- 1976 López-Sánchez, J.F., Amram, M.B., Lakkis, M.D., Lagarde, F., Rauret, G. and Leroy, M.J.F.: Quantitative aspects of separation of arsenical species by free solution capillary electrophoresis. *Fresenius J. Anal. Chem.*, 348 (1994) 810-814.
- 1977 Romano, J.P.: Capillary ion analysis. A method for determining ions in water and solid waste leachates. *Am. Lab. (Shelton)*, 25 (1993) 48-51; C.A., 120 (1994) 207393x - a review with 8 refs.

See also 1415, 1416, 1924, 1973.

38b. Anions

- 1973 Bondoux, G. and Jones, T.: Analysis of ppb levels of anions and cations in deionized water. *LC-GC Int.*, 7 (1994) 29-34.

See also 1966, 1967.

39. RADIOACTIVE AND OTHER ISOTOPE COMPOUNDS

See 1970.

PUBLICATION SCHEDULE FOR THE 1994 SUBSCRIPTION

Journal of Chromatography A and *Journal of Chromatography B: Biomedical Applications*

MONTH	1993	J-J	J	A	S	O	
Journal of Chromatography A	652-657	Vols. 658-672	673/1 673/2 674/1 + 2 675/1 + 2 676/1	676/2 677/1 677/2 678/1	678/2 679/1 679/2 680/1	680/2	The publication schedule for further issues will be published later.
Bibliography Section		Vol. 681			682/1		
Journal of Chromatography B: Biomedical Applications		Vols. 652-656	657/1 657/2	658/1 658/2	659	660/1 660/2	

INFORMATION FOR AUTHORS

(Detailed *Instructions to Authors* were published in *J. Chromatogr. A*, Vol. 657, pp. 463-469. A free reprint can be obtained by application to the publisher, Elsevier Science B.V., P.O. Box 330, 1000 AH Amsterdam, Netherlands.)

Types of Contributions. The following types of papers are published: Regular research papers (full-length papers), Review articles, Short Communications and Discussions. Short Communications are usually descriptions of short investigations, or they can report minor technical improvements of previously published procedures; they reflect the same quality of research as full-length papers, but should preferably not exceed five printed pages. Discussions (one or two pages) should explain, amplify, correct or otherwise comment substantively upon an article recently published in the journal. For Review articles, see inside front cover under Submission of Papers.

Submission. Every paper must be accompanied by a letter from the senior author, stating that he/she is submitting the paper for publication in the *Journal of Chromatography A* or *B*.

Manuscripts. Manuscripts should be typed in **double spacing** on consecutively numbered pages of uniform size. The manuscript should be preceded by a sheet of manuscript paper carrying the title of the paper and the name and full postal address of the person to whom the proofs are to be sent. As a rule, papers should be divided into sections, headed by a caption (e.g., Abstract, Introduction, Experimental, Results, Discussion, etc.). All illustrations, photographs, tables, etc., should be on separate sheets.

Abstract. All articles should have an abstract of 50-100 words which clearly and briefly indicates what is new, different and significant. No references should be given.

Introduction. Every paper must have a concise introduction mentioning what has been done before on the topic described, and stating clearly what is new in the paper now submitted.

Experimental conditions should preferably be given on a *separate* sheet, headed "Conditions". These conditions will, if appropriate, be printed in a block, directly following the heading "Experimental".

Illustrations. The figures should be submitted in a form suitable for reproduction, drawn in Indian ink on drawing or tracing paper. Each illustration should have a caption, all the *captions* being typed (with double spacing) together on a *separate sheet*. If structures are given in the text, the original drawings should be provided. Coloured illustrations are reproduced at the author's expense, the cost being determined by the number of pages and by the number of colours needed. The written permission of the author and publisher must be obtained for the use of any figure already published. Its source must be indicated in the legend.

References. References should be numbered in the order in which they are cited in the text, and listed in numerical sequence on a separate sheet at the end of the article. Please check a recent issue for the layout of the reference list. Abbreviations for the titles of journals should follow the system used by *Chemical Abstracts*. Articles not yet published should be given as "in press" (journal should be specified), "submitted for publication" (journal should be specified), "in preparation" or "personal communication".

Vols. 1-651 of the *Journal of Chromatography*; *Journal of Chromatography, Biomedical Applications* and *Journal of Chromatography, Symposium Volumes* should be cited as *J. Chromatogr.* From Vol. 652 on, *Journal of Chromatography A* (incl. *Symposium Volumes*) should be cited as *J. Chromatogr. A* and *Journal of Chromatography B: Biomedical Applications* as *J. Chromatogr. B*.

Dispatch. Before sending the manuscript to the Editor please check that the envelope contains four copies of the paper complete with references, captions and figures. One of the sets of figures must be the originals suitable for direct reproduction. Please also ensure that permission to publish has been obtained from your institute.

Proofs. One set of proofs will be sent to the author to be carefully checked for printer's errors. Corrections must be restricted to instances in which the proof is at variance with the manuscript.

Reprints. Fifty reprints will be supplied free of charge. Additional reprints can be ordered by the authors. An order form containing price quotations will be sent to the authors together with the proofs of their article.

Advertisements. The Editors of the journal accept no responsibility for the contents of the advertisements. Advertisement rates are available on request. Advertising orders and enquiries can be sent to the Advertising Manager, Elsevier Science B.V., Advertising Department, P.O. Box 211, 1000 AE Amsterdam, Netherlands; courier shipments to: Van de Sande Bakhuyzenstraat 4, 1061 AG Amsterdam, Netherlands; Tel. (+31-20) 515 3220/515 3222, Telefax (+31-20) 6833 041, Telex 16479 els vi nl. UK: T.G. Scott & Son Ltd., Tim Blake, Portland House, 21 Narborough Road, Cosby, Leics. LE9 5TA, UK; Tel. (+44-533) 753 333, Telefax (+44-533) 750 522. USA and Canada: Weston Media Associates, Daniel S. Lipner, P.O. Box 1110, Greens Farms, CT 06436-1110, USA; Tel. (+1-203) 261 2500, Telefax (+1-203) 261 0101.

Flow-Through (Bio)Chemical Sensors

By **M. Valcárcel** and **M.D. Luque de Castro**, Department of Analytical Chemistry,
University of Córdoba, 14004 Córdoba, Spain

Techniques and Instrumentation in Analytical Chemistry Volume 16

Flow-through sensors are more suitable than classical probe-type sensors for addressing real (non-academic) problems. The external shape and operation of flow-through (bio)chemical sensors are of great practical significance as they facilitate sample transport and conditioning, as well as calibration and sensor preparation, maintenance and regeneration, all of which result in enhanced analytical features and a wider scope of application.

This is a systematic presentation of flow-through chemical and biochemical sensors based on the permanent or transient immobilization of any of the ingredients of a (bio)chemical reaction (i.e. the analyte, reagent, catalyst or product) where detection is integrated with the analytical reaction, a separation process (dialysis, gas diffusion, sorption, etc.) or both.

The book deals critically with most types of flow-through sensors, discussing their possibilities and shortcomings to provide a realistic view of the state-of-the-art in the field. The large numbers of figures, the wealth of literature references and the extensive subject index complement the text.

Contents: 1. Sensors in Analytical Chemistry. Analytical chemistry at the turn of the XXI

century. Analytical information. What is a sensor? Sensors and the analytical process. Types of sensors. General features of (bio)chemical sensors. (Bio)chemical sensors and analytical properties. Commercial availability. Trends in sensor development.

2. Fundamentals of Continuous-Flow (Bio)Chemical Sensors. Definition. Classification. The active microzone. Flow-through cells. Continuous configurations. Regeneration modes. Transient signals. Measurement modes. The role of kinetics. Requirements for proper sensor performance.

3. Flow-Through Sensors Based on Integrated Reaction and Detection. Introduction. Flow-through sensors based on an immobilized catalyst. Flow-through immunosensors. Flow-through sensors based on an immobilized reagent. Flow-through sensors based on an *in situ* produced reagent.

4. Flow-Through Sensors Based on Integrated Separation and Detection. Introduction. Integrated gas diffusion and detection. Integrated liquid-liquid separation and detection. Integrated retention and detection. Flow-through sensors for multi-determinations based on integrated retention and detection. Ion-selective electrodes (ISEs) and ion-sensitive field-effect transistors (ISFETs).

5. Flow-Through Sensors Based on Integrated Reaction, Separation and Detection. Introduction. Integration of gas-diffusion, reaction and detection. Integration of dialysis, reaction and detection. Integration of sorption, reaction and detection.

Index.

© 1994 332 pages **Hardbound**
Price: Dfl. 355.00 (US\$ 202.75)
ISBN 0-444-89866-2

ORDER INFORMATION
ELSEVIER SCIENCE B.V.

P.O. Box 330
1000 AH Amsterdam
The Netherlands
Fax: (+31-20) 5862 845

For USA and Canada

P.O. Box 945
Madison Square Station
New York, NY 10159-0945
Fax: (212) 633 3680

US\$ prices are valid only for the USA & Canada and are subject to exchange rate fluctuations; in all other countries the Dutch guilder price (Dfl.) is definitive. Customers in the European Union should add the appropriate VAT rate applicable in their country to the price(s). Books are sent postfree if prepaid.



**ELSEVIER
SCIENCE**



0021-9673(19940916)682:1;1-J

J. M. A. 2537
10 m a 20