

VOL. 713 NO. 2 16 JUNE 1995

THIS ISSUE COMPLETES VOL. 713

## Bibliography Section

JOURNAL OF

# CHROMATOGRAPHY A

INCLUDING ELECTROPHORESIS AND OTHER SEPARATION METHODS

### EDITORS

U.A.Th. Brinkman (Amsterdam)  
R.W. Giese (Boston, MA)  
J.K. Haken (Kensington, N.S.W.)  
C.F. Poole (London)  
L.R. Snyder (Orinda, CA)  
S. Terabe (Hyogo)

EDITORS, SYMPOSIUM VOLUMES,  
E. Heftmann (Orinda, CA), Z. Deyl (Prague)

### EDITORIAL BOARD

D.W. Armstrong (Rolla, MO)  
W.A. Aue (Halifax)  
P. Boček (Brno)  
P.W. Carr (Minneapolis, MN)  
J. Crommen (Liège)  
V.A. Davankov (Moscow)  
G.J. de Jong (Weesp)  
Z. Deyl (Prague)  
S. Dilli (Kensington, N.S.W.)  
Z. El Rassi (Stillwater, OK)  
H. Engelhardt (Saarbrücken)  
M.B. Evans (Hatfield)  
S. Fanali (Rome)  
G.A. Guiochon (Knoxville, TN)  
P.R. Haddad (Hobart, Tasmania)  
I.M. Hais (Hradec Králové)  
W.S. Hancock (Palo Alto, CA)  
S. Hjertén (Uppsala)  
S. Honda (Higashi-Osaka)  
Cs. Horváth (New Haven, CT)  
J.F.K. Huber (Vienna)  
J. Janák (Brno)  
P. Jandera (Pardubice)  
B.L. Karger (Boston, MA)  
J.J. Kirkland (Newport, DE)  
E. sz. Kováts (Lausanne)  
C.S. Lee (Ames, IA)  
K. Macek (Prague)  
A.J.P. Martin (Cambridge)  
E.D. Morgan (Keele)  
H. Poppe (Amsterdam)  
P.G. Righetti (Milan)  
P. Schoenmakers (Amsterdam)  
R. Schwarzenbach (Dübendorf)  
R.E. Shoup (West Lafayette, IN)  
R.P. Singhal (Wichita, KS)  
A.M. Siouffi (Marseille)  
D.J. Strydom (Boston, MA)  
T. Takagi (Osaka)  
N. Tanaka (Kyoto)  
K.K. Unger (Mainz)  
P. van Zoonen (Bilthoven)  
R. Verpoorte (Leiden)  
Gy. Vigh (College Station, TX)  
J.T. Watson (East Lansing, MI)  
B.D. Westerlund (Uppsala)

### EDITORS, BIBLIOGRAPHY SECTION

Z. Deyl (Prague), J. Janák (Brno), V. Schwarz (Prague)

ELSEVIER

# JOURNAL OF CHROMATOGRAPHY A

INCLUDING ELECTROPHORESIS AND OTHER SEPARATION METHODS

**Scope.** The *Journal of Chromatography A* publishes papers on all aspects of **chromatography, electrophoresis** and related methods. Contributions consist mainly of research papers dealing with chromatographic theory, instrumental developments and their applications. In the *Symposium volumes*, which are under separate editorship, proceedings of symposia on chromatography, electrophoresis and related methods are published. *Journal of Chromatography B: Biomedical Applications*—This journal, which is under separate editorship, deals with the following aspects: developments in and applications of chromatographic and electrophoretic techniques related to clinical diagnosis or alterations during medical treatment; screening and profiling of body fluids or tissues related to the analysis of active substances and to metabolic disorders; drug level monitoring and pharmacokinetic studies; clinical toxicology; forensic medicine; veterinary medicine; occupational medicine; results from basic medical research with direct consequences in clinical practice.

**Submission of Papers.** The preferred medium of submission is on disk with accompanying manuscript (see *Electronic manuscripts* in the Instructions to Authors, which can be obtained from the publisher, Elsevier Science B.V., P.O. Box 330, 1000 AH Amsterdam, Netherlands). Manuscripts (in English; four copies are required) should be submitted to: Editorial Office of *Journal of Chromatography A*, P.O. Box 681, 1000 AR Amsterdam, Netherlands, Telefax (+31-20) 485 2304, or to: The Editor of *Journal of Chromatography B: Biomedical Applications*, P.O. Box 681, 1000 AR Amsterdam, Netherlands. Review articles are invited or proposed in writing to the Editors who welcome suggestions for subjects. An outline of the proposed review should first be forwarded to the Editors for preliminary discussion prior to preparation. Submission of an article is understood to imply that the article is original and unpublished and is not being considered for publication elsewhere. For copyright regulations, see below.

**Publication information.** *Journal of Chromatography A* (ISSN 0021-9673): for 1995 Vols. 683–714 are scheduled for publication. *Journal of Chromatography B: Biomedical Applications* (ISSN 0378-4347): for 1995 Vols. 663–674 are scheduled for publication. Subscription prices for *Journal of Chromatography A*, *Journal of Chromatography B: Biomedical Applications* or a combined subscription are available upon request from the publisher. Subscriptions are accepted on a prepaid basis only and are entered on a calendar year basis. Issues are sent by surface mail except to the following countries where air delivery via SAL is ensured: Argentina, Australia, Brazil, Canada, China, Hong Kong, India, Israel, Japan, Malaysia, Mexico, New Zealand, Pakistan, Singapore, South Africa, South Korea, Taiwan, Thailand, USA. For all other countries airmail rates are available upon request. Claims for missing issues must be made within six months of our publication (mailing) date. Please address all your requests regarding orders and subscription queries to: Elsevier Science B.V., Journal Department, P.O. Box 211, 1000 AE Amsterdam, Netherlands. Tel.: (+31-20) 485 3642; Fax: (+31-20) 485 3598. Customers in the USA and Canada wishing information on this and other Elsevier journals, please contact Journal Information Center, Elsevier Science Inc., 655 Avenue of the Americas, New York, NY 10010, USA. Tel. (+1-212) 633 3750, Telefax (+1-212) 633 3764.

**Abstracts/Contents Lists** published in Analytical Abstracts, Biochemical Abstracts, Biological Abstracts, Chemical Abstracts, Chemical Titles, Chromatography Abstracts, Current Awareness in Biological Sciences (CABS), Current Contents/Life Sciences, Current Contents/Physical, Chemical & Earth Sciences, Deep-Sea Research/Part B: Oceanographic Literature Review, Excerpta Medica, Index Medicus, Mass Spectrometry Bulletin, PASCAL-CNRS, Referativnyi Zhurnal, Research Alert and Science Citation Index.

**US Mailing Notice.** *Journal of Chromatography A* (ISSN 0021-9673) is published weekly (total 52 issues) by Elsevier Science B.V., (Sara Burgerhartstraat 25, P.O. Box 211, 1000 AE Amsterdam, Netherlands). Annual subscription price in the USA US\$ 5389.00 (US\$ price valid in North, Central and South America only) including air speed delivery. Second class postage paid at Jamaica, NY 11431. **USA POSTMASTERS:** Send address changes to *Journal of Chromatography A*, Publications Expediting, Inc., 200 Meacham Avenue, Elmont, NY 11003. Airfreight and mailing in the USA by Publications Expediting.

**See inside back cover** for Publication Schedule, Information for Authors and information on Advertisements.

© 1995 ELSEVIER SCIENCE B.V. All rights reserved.

0021-9673/95 \$09.50

No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, without the prior written permission of the publisher, Elsevier Science B.V. Copyright and Permissions Department, P.O. Box 521, 1000 AM Amsterdam, Netherlands.

Upon acceptance of an article by the journal, the author(s) will be asked to transfer copyright of the article to the publisher. The transfer will ensure the widest possible dissemination of information.

**Special regulations for readers in the USA**—This journal has been registered with the Copyright Clearance Center, Inc. Consent is given for copying of articles for personal or internal use, or for the personal use of specific clients. This consent is given on the condition that the copier pays through the Center the per-copy fee stated in the code on the first page of each article for copying beyond that permitted by Sections 107 or 108 of the US Copyright Law. The appropriate fee should be forwarded with a copy of the first page of the article to the Copyright Clearance Center, Inc., 222 Rosewood Drive, Danvers, MA 01923, USA. If no code appears in an article, the author has not given broad consent to copy and permission to copy must be obtained directly from the author. The fee indicated on the first page of an article in this issue will apply retroactively to all articles published in the journal, regardless of the year of publication. This consent does not extend to other kinds of copying, such as for general distribution, resale, advertising and promotion purposes, or for creating new collective works. Special written permission must be obtained from the publisher for such copying.

No responsibility is assumed by the Publisher for any injury and/or damage to persons or property as a matter of products liability, negligence or otherwise, or from any use or operation of any methods, products, instructions or ideas contained in the materials herein. Because of rapid advances in the medical sciences, the Publisher recommends that independent verification of diagnoses and drug dosages should be made.

Although all advertising material is expected to conform to ethical (medical) standards, inclusion in this publication does not constitute a guarantee or endorsement of the quality or value of such product or of the claims made of it by its manufacturer.

Ⓢ The paper used in this publication meets the requirements of ANSI/NISO Z39.48-1992 (Permanence of Paper).

Printed in the Netherlands

## CONTENTS

*Bibliography Section*

|                                    |      |
|------------------------------------|------|
| Liquid Column Chromatography ..... | B135 |
| Gas Chromatography .....           | B188 |
| Planar Chromatography .....        | B209 |
| Electrophoresis .....              | B220 |



## Bibliography Section

### Liquid Column Chromatography

#### 1. REVIEWS AND BOOKS

- 1292 Bertrand, A.: (Ion chromatography). In: *Acquis. Recent. Chromatogr. Vin Cours Eur. Form. Contin.* 1992, 1993, pp. 33-43; C.A., 121 (1994) 299381p - a review with 1 ref.
- 1293 Bryant, C.H., Adam, A., Taylor, D.R. and Rowe, R.C.: A review of expert systems for chromatography. *Anal. Chim. Acta*, 297 (1994) 317-347 - a review with 88 refs.
- 1294 Dughan, L.: Perfusion confusion: how does high-speed chromatography work? *Bio/Technology*, 12 (1994) 1090 - a review with no refs.
- 1295 Feistner, G.J.: Profiling of bacterial metabolites by liquid chromatography-electrospray mass spectrometry: a perspective. *Am. Lab. (Shelton)*, 26 (1994) 32L, 32M, 32N, 32O, 32P, 32Q; C.A., 121 (1994) 296211x - a review with 11 refs.
- 1296 Kumar, M.: Recent trends in chromatographic procedures for separation and determination of rare earth elements. A review. *Analyst (Cambridge)*, 119 (1994) 2013-2024.
- 1297 Lederer, M.: *Chromatography for Inorganic Chemistry* Wiley, Chichester, 1994, 221 p.; C.A., 121 (1994) 220570t.
- 1298 Miyazaki, M.: (Contribution of ion chromatography directed to the field of public health). *Hokuriku Kosho Eisei Gakkaishi*, 19 (1992) 1-7; C.A., 121 (1994) 296217d - a review with 38 refs.
- 1299 Szepesy, L.: (Development and importance of separation techniques in instrumental analysis). *Magy. Kem. Foly.*, 100 (1994) 340-353; C.A., 121 (1994) 270751y.
- 1300 Watanabe, Y. and Yamada, N.: (Column chromatography). *Lipid*, 5 (1994) 285-287; C.A., 122 (1995) 4559z - a review with 7 refs.
- 1301 Welinder, B.S., Kornfelt, T. and Sorensen, H.H.: Stationary phases: the Achilles heel in HPLC of protein pharmaceuticals? *Anal. Chem.*, 67 (1995) 39A-43A.
- 1302 Zhou, X., Wu, C. and Chen, Y.: (Application of crown ether compounds to chromatographic analysis). *Sepu*, 12 (1994) 175-179; C.A., 121 (1994) 220614k - a review with 68 refs.

See also 1304, 1316, 1321, 1322, 1325, 1346, 1366, 1368, 1369, 1375, 1380, 1382, 1392, 1407, 1425, 1434, 1436, 1440, 1446, 1447, 1449, 1452, 1456, 1458, 1460, 1468, 1470, 1471, 1473, 1474, 1479, 1481, 1482, 1484, 1491, 1492, 1507, 1518, 1523, 1551, 1579, 1639, 1640, 1647, 1651, 1660, 1817, 1856, 1876, 1895, 1914, 2102, 2118, 2179, 2252, 2270, 2272, 2460, 2461, 2469, 2496, 2525.

#### 2. FUNDAMENTALS, THEORY AND GENERAL

##### 2a. General

- 1303 Arutyunov, Yu.A. and Vigdergaus, M.S.: (Instrumental and methodological realization of some specific features of chromatography). *Zh. Anal. Khim.*, 49 (1994) 796-803; C.A., 121 (1994) 314871u.
- 1304 Beinert, W.D.: (Principles and advantages of the narrow-bore HPLC). *GIT Spez. Chromatogr.*, 14 (1994) 17-20; C.A., 122 (1995) 22799g - a review without refs.
- 1305 Brooks, C.A. and Cramer, S.M.: Investigation of displacer equilibrium properties and mobile phase operating conditions in ion-exchange displacement chromatography. *J. Chromatogr. A*, 693 (1995) 187-196.
- 1306 Cuesta Sánchez, F. and Massart, D.L.: Application of SIMPLISMA for the assessment of peak purity in liquid chromatography with diode array detection. *Anal. Chim. Acta*, 298 (1994) 331-339.
- 1307 De Ligny, C.L.: Some uncommon applications of chromatography. *Acta Chromatogr.*, 1 (1992) 7-22; C.A., 122 (1995) 65617p.
- 1308 Deinhammer, R.S., Ting, E.-Y. and Porter, M.D.: Dynamic modification of separations using electrochemically modulated liquid chromatography. *Anal. Chem.*, 67 (1995) 237-246.
- 1309 Dolan, J.W.: Case study - irreproducible retention times. *LC-GC Int.*, 8, No. 1 (1995) 10-13.
- 1310 Dolan, J.W.: Obtaining separations, part II: adjusting selectivity. *LC-GC*, 12 (1994) 446-450; C.A., 121 (1994) 314879c.
- 1311 Foti, G., Hajos, P. and Kovats, E.Sz.: Retention in analytical ion exchange chromatography-I. Strong electrolytes on strong ion exchangers. *Talanta*, 42 (1994) 1073-1081; C.A., 121 (1994) 270553k.
- 1312 Freeman, D.C. and Byrd, D.W.: Statistical treatment of large digital chromatographic data sets. *J. Chromatogr. A*, 686 (1994) 225-233.
- 1313 Giona, M. and Spera, D.: An integral approximation for the optimization of size-exclusion chromatographic column performance. *Spec. Publ. - R. Soc. Chem.*, 158 (Separations for Biotechnology 3) (1994) 193-199; C.A., 121 (1994) 296300a.
- 1314 Grize, Y.-L., Schmidli, H. and Born, J.: Effect of integration parameters on high-performance liquid chromatographic method development and validation. *J. Chromatogr. A*, 686 (1994) 1-10.

- 1315 Hamoir, T. and Massart, D.L.: Strategic approach for method selection in high-performance liquid chromatography. *Anal. Chim. Acta*, 298 (1994) 319-329.
- 1316 Hubert, P. and Dellacherie, E.: Molecular interactions in hydrophobic chromatography. *Mol. Interact. Biosep.*, (1993) 333-359; *C.A.*, 121 (1994) 249863b - a review with no refs.
- 1317 Lee, S.T. and Olesik, S.V.: Comparison of enhanced-fluidity and elevated-temperature mobile phases in reversed-phase high-performance liquid chromatography. *Anal. Chem.*, 66 (1994) 4498-4506.
- 1318 Mizrotsky, N., Kristol, L. and Grushka, E.: Use of chromatographic system peaks for continuous quantitative analysis. *J. Chromatogr. A*, 691 (1995) 21-27.
- 1319 Nelson, D.C.: A status report on the Analytical Instruments Association data communications standards program. *Chemom. Intell. Lab. Syst.*, 26 (1994) 43-46; *C.A.*, 121 (1994) 124081p.
- 1320 Olsen, B.A. and Sullivan, G.R.: Chemometric categorization of octadecylsilyl bonded-phase silica columns using test mixtures and confirmation of results with pharmaceutical compound separations. *J. Chromatogr. A*, 692 (1995) 147-159.
- 1321 Scott, R.P.W.: LC applications. *Chromatogr. Sci. Ser.*, 67 (1994) 281-320; *C.A.*, 122 (1995) 45011h - a review with no refs.
- 1322 Sofer, G.: Validation of chromatographic processes. *Prog. Biotechnol.*, 9(CEB6: Proceedings of the 6th European Congress on Biotechnology, 1993, Pt. 2) (1994) 977-980; *C.A.*, 121 (1994) 249789g - a review with no refs.
- 1323 Walczak, B.: Study on the mobile and stationary phase interactions in the reversed-phase liquid chromatography. *Pol. J. Chem.*, 66 (1992) 2015-2023; *C.A.*, 121 (1994) 264520f.
- 1324 Wang, R., Ottomar, K., Huddleston, J. and Lyddiatt, A.: Salting out phenomena in aqueous two-phase partition, hydrophobic interaction chromatography, and fractional precipitation. *Spec. Publ.-R.Soc. Chem.*, 158(Separations for Biotechnology 3) (1994) 532-538; *C.A.*, 122 (1995) 4846r.
- For additional information see *C.A.*:  
121 (1994) 287381a, 308504k.
- See also 1299, 1335, 1376, 1431, 1588.

## 2b. Thermodynamics and theoretical relationships

- 1325 Davis, J.M.: Statistical theories of peak overlap in chromatography. *Adv. Chromatogr. (N.Y.)*, 34 (1994) 109-176; *C.A.*, 121 (1994) 238951c - a review with 52 refs.
- 1326 Hansen, R.L. and Harris, J.M.: Lateral diffusion of molecules partitioned into C-18 ligands on silica surfaces. *Anal. Chem.*, 67 (1995) 492-498.
- 1327 Jenke, D.R.: Prediction of retention characteristics of multiprotic anions in ion chromatography. *Anal. Chem.*, 66 (1994) 4466-4470.
- 1328 Jiang, D.G., Wang, F., Zhou, C.R., Cao, T.Z. and Zhao, Y.: The relationship between retention values of homologs and the column temperature in reverse-phase liquid chromatography. *Chin. Chem. Lett.*, 5 (1994) 321-324; *C.A.*, 121 (1994) 220655z.

- 1329 Kalinichev, A.I.: (Describing the stationary concentration waves for Langmuir isotherms in the sorption of a mixture based on the theory for multicomponent chromatography). *Zh. Fiz. Khim.*, 68 (1994) 1658-1662; *C.A.*, 122 (1995) 65255u.
- 1330 Lebedev, Yu.Ya.: (Conditions of motion of the chromatographic zone for the case of intradiffusion kinetics of interfacial mass transfer). *Zh. Fiz. Khim.*, 68 (1994) 1733-1739; *C.A.*, 122 (1995) 12761z.
- 1331 Li, J. and Pardue, H.L.: Predictive steady-state chromatography. 1. Algorithms for leading and trailing edges of resolved and unresolved peaks in liquid chromatography. *Anal. Chem.*, 66 (1994) 3765-3772.
- 1332 Lochmüller, C.H., Reese, C. and Hsu, S.-H.: Cross-column retention prediction in reversed-phase liquid chromatography using factor analytical modeling. *Anal. Chem.*, 66 (1994) 3806-3813.
- 1333 Lübke, M., le Quéré, J.-L. and Barron, D.: Normal-phase high-performance liquid chromatography of volatile compounds. Selectivity and mobile phase effects on polar bonded silica. *J. Chromatogr. A*, 690 (1995) 41-54.
- 1334 Schure, M.R.: Molecular dynamics of liquid chromatography: Chain and solvent structure visualization. *Spec. Publ.-R. Soc. Chem.*, 139 (1994) 181-189; *C.A.*, 121 (1994) 239081f.
- 1335 Tchaplá, A. and Héron, S.: Property-structure relationship of solute-stationary phase complexes occurring in a molecular mechanism by penetration of elute in bonded alkyl chains in reversed-phase liquid chromatography. *J. Chromatogr. A*, 684 (1994) 175-188.
- 1336 Toft, J. and Kvalheim, O.M.: Multi-array resolution parameter for multidetermination chromatography. Performance of alternating regression, iterative target transformation factor analysis and heuristic evolving latent projections. *Chemom. Intell. Lab. Syst.*, 25 (1994) 61-75; *C.A.*, 121 (1994) 314695q.
- 1337 Wasiak, W.: Thermodynamic characteristics of the specific interactions between alkenes and chromatographic packings with chemically bonded copper(II) complexes. *Chem. Anal. (Warsaw)*, 39 (1994) 325-333; *C.A.*, 121 (1994) 287144a.
- 1338 Yang, S. and Khaledi, M.G.: Linear solvation energy relationships in micellar liquid chromatography and micellar electrokinetic capillary chromatography. *J. Chromatogr. A*, 692 (1995) 301-310.

For additional information see *C.A.*:

121 (1994) 220417y, 239104r, 239105s, 287069e, 296322j.

See also 1306, 1387, 1408, 1421, 1499, 1525, 1879, 2503, 2532.

## 2c. Relationship between structure and chromatographic behaviour

- 1339 Varga-Defterdarovic, L., Horvat, S., Skuric, M. and Horvat, J.: Correlation of structure and retention behaviour in reversed-phase high-performance liquid chromatography. I. Leucine-enkephalin-related glycoconjugates. *J. Chromatogr. A*, 687 (1994) 101-106.
- 1340 Varga-Defterdarovic, L., Horvat, S., Skuric, M. and Horvat, J.: Correlation of structure and retention behaviour in reversed-phase high-performance liquid chromatography. II. Methionine-enkephalin-related glycoconjugates. *J. Chromatogr. A*, 687 (1994) 107-112.

For additional information see C.A.:  
121 (1994) 270777m.

See also 1335, 1559, 1586, 1794.

## 2d. Measurement of physico-chemical and related values

- 1341 Abraham, M.H., Chadha, H.S. and Leo, A.J.: Hydrogen bonding. XXXV. Relationship between high-performance liquid chromatography capacity factors and water-octanol partition coefficients. *J. Chromatogr. A*, 685 (1994) 203-211.
- 1342 Barnikol, W.K.R. and Pötzschke, H.: Calibration of gel-permeation columns in the high-molecular-mass range: Fixed human thrombocytes for the estimation of interstitial volume and the haemocyanin of the Vineyard snail *Helix pomatia* as a molecular mass calibration substance. *J. Chromatogr. A*, 685 (1994) 221-227.
- 1343 Bignotti, F., Sozzani, P., Ranucci, E. and Ferruti, P.: NMR Studies, molecular characterization and degradation behavior of poly(amino amine)s. 1. Poly(amido amine) deriving from the polyaddition of 2. Methylpiperazine to 1,4-bis(acryloyl)piperazine. *Macromolecules*, 27 (1994) 7171-7178; C.A., 121 (1994) 281336t.
- 1344 Fischer, C.-H., Berek, D. and Macko, T.: Determination of complexation equilibria of macromolecules with small molecules by means of size exclusion chromatography. *Polym. Bull. (Berlin)*, 33 (1994) 339-346; C.A., 121 (1994) 256692y.
- 1345 Hsieh, M.-M. and Dorsey, J.G.: Bioavailability estimation by reversed-phase liquid chromatography: high bonding density C-18 phases for modeling biopartitioning processes. *Anal. Chem.*, 67 (1995) 48-57.
- 1346 Kiso, Y.: (Application of HPLC to the research of semipermeable membranes. HPLC using cellulose acetate as a stationary phase). *Shinsozai*, 5 (1994) 22-25; C.A., 121 (1994) 258114k - a review with 17 refs.
- 1347 Ong, S., Liu, H., Qiu, X., Bhat, G. and Pidgeon, C.: Membrane partition coefficients chromatographically measured using immobilized artificial membrane surfaces. *Anal. Chem.*, 67 (1995) 755-762.
- 1348 Pasch, H.: (Liquid chromatography at critical point of adsorption. A new method for block copolymer analysis). *GIT Fachz. Lab.*, 37 (1993) 1068-1070; C.A., 121 (1994) 206437y.
- 1349 Rosset, R., Hui, F., Xie, J., Machtalere, G., Kolodziejczyk, H. and Sassiati, P.: (New calibration methods in size exclusion chromatography). *Analisis*, 22 (1994) 293-304; C.A., 121 (1994) 206447b.
- 1350 Shortt, D.W.: Measurement of narrow-distribution polydispersity using multiangle light scattering. *J. Chromatogr. A*, 686 (1994) 11-20.
- 1351 Wallis, K.H. and Mueller, R.H.: Determination of the surface hydrophobicity of colloidal dispersions by miniaturized hydrophobic-interaction chromatography. *Pharm. Ind.*, 55 (1993) 1124-1128; C.A., 121 (1994) 238221q.

For additional information see C.A.:  
121 (1994) 206698j, 282576h, 291602r.

See also 1375, 1523, 1541, 1543, 1584, 1585, 1735, 1863, 1899, 1945, 2245, 2248, 2250, 2251, 2252, 2253, 2255, 2256, 2259, 2260, 2481.

## 3. GENERAL TECHNIQUES

### 3a. Apparatus and accessories

- 1352 De Santis, F.: Comment on wet effluent denuder coupled liquid/ion chromatography systems: annular and parallel plate denuders. *Anal. Chem.*, 66 (1994) 3503-3504.
- 1353 Kalousek, J., Gunter, J., Vanek, T. and Rosenberg, I.: Liquid chromatography column head. *Czech. Rep. CZ* 278,574 (Cl. B01D15/08), 16 Mar. 1994, Appl. 1,075, 16 Apr. 1991; 5 p.; C.A., 122 (1995) 45397p.
- 1354 Njamfa, S. and Joron, L.: High performance liquid chromatography instrument with controlling device and process for controlling its functioning. *Eur. Pat. Appl.* EP 628,813 (Cl. G0130/32), 14 Dec. 1994, FR Appl. 93/6,825, 08 Jun. 1993; 10 p.; C.A., 122 (1995) 71025k.
- 1355 Riegger, H.: Sample dosing system. *Ger. Offen.* DE 4,318,919 (Cl. G01N1/14), 08 Dec. 1994, Appl. 07 Jun. 1993; 13 p.; C.A., 122 (1995) 71024j.
- 1356 Schick, H.G.: Inlet filter for solvents in liquid chromatography. *U.S. US* 5,366,620 (Cl. 210-198.2; B01D15/08), 22 Nov. 1994, Appl. 41,481, 01 Apr. 1993; 10 p.; C.A., 122 (1995) 71008g.
- 1357 Simpson, R.C.: Modification of a conventional high-performance liquid chromatography autoinjector for use with capillary liquid chromatography. *J. Chromatogr. A*, 691 (1995) 163-170.
- 1358 Takahashi, T., Shizukuishi, K., Katho, Y., Senda, N. and Bando, T.: Liquid chromatograph mass spectrometer. *U.S. US* 5,368,727 (Cl. 210-198.2; B01D15/08), 29 Nov. 1994, JP Appl. 91/256,409, 03 Oct. 1991; 8 p.; C.A., 122 (1995) 71010b.

For additional information see C.A.:  
121 (1994) 220589f, 220591a, 220595e, 233885z, 270702h, 270713n, 282781w, 291557e, 296600y, 296603b, 314846q, 314928t;  
122 (1995) 45326q.

See also 1343, 1392, 1413, 1452, 1457, 2118.

### 3b. Detectors and detection reagents

- 1359 Abbas, A.A. and Shelly, D.C.: Optical properties of axial-illumination flow cells for simultaneous absorbance-fluorescence detection in micro liquid chromatography. *J. Chromatogr. A*, 691 (1995) 37-53.
- 1360 Allen, L.B., Koropchak, J.A. and Szostek, B.: Condensation nucleation light scattering detection for conventional reversed-phase liquid chromatography. *Anal. Chem.*, 67 (1995) 659-666.
- 1361 Aue, W.A., Singh, H. and Sun, X.-Y.: Fundamental noise in three chromatographic detectors. *J. Chromatogr. A*, 687 (1994) 283-290.
- 1362 Frolov, F.Ya. and Vrazhnikov, V.V.: Spectrophotometric detector for liquid chromatograph. *U.S.S.R. SU* 1,821,692 (Cl. G01N21/03), 15 Jun. 1993, Appl. 4,890,352, 14 Nov. 1990; C.A., 122 (1995) 45398q.
- 1363 Kurata, S. and Hirano, H.: (Fluorescent substances for development of multipurpose markers). *Hanzaigaku Zasshi*, 60 (1994) 95-100; C.A., 121 (1994) 275699x.

- 1364 Lee, K.S., Shin, J.H., Cha, M.J., Cha, G.S., Trojanowicz, M., Liu, D., Goldberg, H.D., Hower, R.W. and Brown, R.B.: Multiionophore-based solid-state potentiometric ion sensor as a cation detector for ion chromatography. *Sens. Actuators, B*, 20 (1994) 239-246; C.A., 122 (1995) 22659m.
- 1365 Lima, L.R., III, Synovec, R.E.: Laser-based dynamic surface tension detection for liquid chromatography by probing a repeating drop radius. *J. Chromatogr. A*, 691 (1995) 195-204.
- 1366 Lingeman, H. and Brinkman, U.A.T.: (Post-column) reaction-detection: an alternative to improve sensitivity and selectivity in column liquid chromatographic analysis. In: Stevenson, D. and Wilson, I.D. (Editors), *Sample Prep. Biomed. Environ. Anal., [Proc. Chromatogr. Soc. Int. Symp.]*, Plenum, New York, 1994, pp. 21-36; C.A., 121 (1994) 275562x - a review with 35 refs.
- 1367 Novak, T.J. and Grayeski, M.L.: Acridinium-based chemiluminescence for high-performance liquid chromatographic detection of chlorophenols. *Microchem. J.*, 50 (1994) 151-160; C.A., 121 (1994) 291660h.
- 1368 Rozhitskii, N.N., Belash, E.M. and Bykh, A.I.: (Electrochemiluminescent analysis: advances, problems, and prospects). *Zh. Anal. Khim.*, 49 (1994) 920-925; C.A., 122 (1995) 22803d - a review with 58 refs.
- 1369 Scott, R.P.W.: Liquid chromatography detectors. *Chromatogr. Sci. Ser.*, 67 (1994) 157-193; C.A., 122 (1995) 45009p - a review with 6 refs.
- 1370 Siddiqui, A. and Shelly, D.C.: Amperostatic-potentiometric detection for micro high-performance liquid chromatography. *J. Chromatogr. A*, 691 (1995) 55-65.
- 1371 Singh, H., Millier, B. and Aue, W.A.: Dual-channel response ratios from an integrative algorithm. *J. Chromatogr. A*, 687 (1994) 291-301.
- 1372 Sun, X.-Y., Singh, H., Millier, B., Warren, C.H. and Aue, W.A.: Noise, filters and detection limits. *J. Chromatogr. A*, 687 (1994) 259-281.
- 1373 Traoré, F., Prognon, P. and Mahuzier, G.: Solvent study on the 9-substituted quinolinizocoumarins used as precolumn fluorescent and chemiluminescent reagents in liquid chromatography. *Anal. Chim. Acta*, 294 (1994) 75-84.
- For additional information see C.A.:  
121 (1994) 220358e, 270873q, 291695y.
- See also 1350, 1532, 1590, 1593, 1613, 1638, 1671, 1717, 1744, 1757, 1814, 1883, 2088, 2100, 2108, 2141, 2161, 2194, 2297, 2302, 2353, 2359, 2439, 2511.
- 3c. *Sorbents and columns, packing procedures*
- 1374 Alvarez, C., Bertorello, H., Strumia, M. and Sanchez, E.I.: Synthesis and characterization of biospecific adsorbents containing glucose, usable to retain concanavalin A. *J. Chromatogr. A*, 686 (1994) 333-338.
- 1375 Baba, N.: (Gel permeation chromatography (GPC). Determination of molecular weight of macromolecular material). *Shin-sozai*, 5 (1994) 35-39; C.A., 121 (1994) 231696q - a review with 11 refs.
- 1376 Barry, A.R. and Chojnacki, R.: Biotechnology product validation. Part 8: Chromatography media and column quantification. *BioPharm (Eugene)*, 7 (1994) 43-47; C.A., 122 (1995) 38800m.
- 1377 Ciucanu, I. and König, W.A.: Immobilization of peralkylated  $\beta$ -cyclodextrin on silica gel for high-performance liquid chromatography. *J. Chromatogr. A*, 685 (1994) 166-171.
- 1378 Cummings, L.J., Ogawa, T. and Tunon, P.: Macro-Prep ceramic hydroxyapatite-new life for an old chromatographic technique. *Spec. Publ.-R.Soc. Chem.*, 158 (Separations for Biotechnology 3) (1994) 134-140; C.A., 122 (1995) 4617s.
- 1379 Daub, J. and Schlieper, T.: Manufacture of solid stationary phases for liquid chromatography in environmental sampling. *Ger. Offen. DE 4,413,319 (Cl. B01D15/08)*, 08 Sep. 1994, Appl. 18 Apr. 1994; 3 p.; C.A., 121 (1994) 262473u.
- 1380 Dimov, N.: High-performance liquid chromatography. The column. *Acta Chromatogr.*, 2 (1993) 21-32; C.A., 121 (1994) 314597j - a brief survey with 9 refs.
- 1381 Dumont, P.J. and Fritz, J.S.: Effect of resin sulfonation on the retention of polar organic compounds in solid-phase extraction. *J. Chromatogr. A*, 691 (1995) 123-131.
- 1382 Forgács, E. and Cserhádi, T.: Retention strength and selectivity of porous graphitized carbon columns. Theoretical aspects and practical applications. *TrAC*, 14 (1995) 23-29 - a review with 30 refs.
- 1383 Freitag, R., Frey, D. and Horváth, C.: Effect of bed compression on high-performance liquid chromatography columns with gigaporous polymeric packings. *J. Chromatogr. A*, 686 (1994) 165-177.
- 1384 Grieb, S.J., Matlin, S.A., Phillips, J.G., Belenguer, A.M. and Ritchie, H.J.: Chiral HPLC with carbohydrate carbamates: influence of support structure on enantioselectivity. *Chirality*, 6 (1994) 129-134; C.A., 122 (1995) 45352v.
- 1385 Guan, H. and Guiochon, G.: Properties of some  $C_{18}$  stationary phases for preparative liquid chromatography. I. Equilibrium isotherms. *J. Chromatogr. A*, 687 (1994) 179-200.
- 1386 Guan, H. and Guiochon, G.: Properties of some  $C_{18}$  stationary phases for preparative liquid chromatography. II. Column efficiency. *J. Chromatogr. A*, 687 (1994) 201-212.
- 1387 Hashimoto, K., Kawase, M., Adachi, S. and Shirai, Y.: Comparison of efficiency in the separation of binary components between a conventional batch chromatographic system and a simulated moving bed adsorber. In: *Off. Proc. Comb. Conf., 6th Conf. Asia Pac. Confed. Chem. Eng., 21st Australas Chem. Eng. Conf., Inst. Eng., Aust., Barton, 1993*, pp. 307/2-312/2; C.A., 122 (1995) 59134n.
- 1388 Hosoya, K., Kishii, Y., Kimata, K., Araki, T., Tanaka, N., Svec, F. and Fréchet, J.M.J.: Uniform-size hydrophobic polymer-based separation media selectively modified with a hydrophilic external polymeric layer. *J. Chromatogr. A*, 690 (1995) 21-28.
- 1389 Ichikawa, H., Yokoyama, A., Kawai, T., Wakizaka, H., Moriyama, H. and Komiya, K.: Packing for liquid chromatography and process for producing them. *Eur. Pat. Appl. EP 610,490 (Cl. G01N30/52)*, 12 Oct. 1994, JP Appl. 93/105,904, 09 Apr. 1993; 13 p.; C.A., 122 (1995) 22864z.
- 1390 Jinno, K., Nakagawa, K., Saito, Y., Ohta, H., Nagashima, H., Itoh, K., Archer, J. and Chen, Y.-L.: Nano-scale design of novel stationary phases to enhance selectivity for molecular shape and size in liquid chromatography. *J. Chromatogr. A*, 691 (1995) 91-99.
- 1391 Kimata, K., Tsuboi, R., Hosoya, K. and Tanaka, N.: Chemically bonded chiral stationary phase prepared by the polymerization of cellulose  $\pi$ -vinylbenzoate. *Anal. Methods Instrum.*, 1 (1993) 23-29; C.A., 121 (1994) 220637v.

- 1392 Kirkland, J.J.: Trends in HPLC column design for improved method development. *Am. Lab. (Shelton)*, 26 (1994) 28K, 28L, 28N-28R; C.A., 122 (1995) 45332p - a review with 20 refs.
- 1393 Kirkland, J.J. and Henderson, J.W.: Reversed-phase HPLC selectivity and retention characteristics of conformationally different bonded alkyl stationary phases. *J. Chromatogr. Sci.*, 32 (1994) 473-480.
- 1394 Kirkland, J.J., van Straten, M.A. and Claessens, H.A.: High pH mobile phase effects on silica-based reversed-phase high-performance liquid chromatographic columns. *J. Chromatogr. A*, 691 (1995) 3-19.
- 1395 Kobayashi, T., Yamaguchi, A., Imai, Y. and Kakimoto, M.: Optically active polyurethanes, their preparation and use as stationary phase for chromatographic resolution of enantiomers. *Jpn. Kokai Tokkyo Koho JP* 06,100,648 (94,100,648) (Cl. C08G18/32), 12 Apr. 1994, Appl. 92/247, 640, 17 Sep. 1992; 8 p.; C.A., 122 (1995) 56848n.
- 1396 Loth, F. and Fanter, C.: (Pearl-shaped cellulose products and their use as separating and supporting material). *Papier (Darmstadt)*, 47 (1993) 703-710; C.A., 121 (1994) 258149a.
- 1397 Moriyama, H., Anegayama, M., Komiya, K. and Kato, Y.: Characterization of a new reversed-phase chromatographic column on a 2- $\mu$ m porous microspherical silica gel. *J. Chromatogr. A*, 691 (1995) 81-89.
- 1398 Neue, U.D., Phillips, J., Walter, T.H., Capparella, M., Alden, B. and Fisk, R.P.: Reversed-phase column quality and its effect on the quality of a pharmaceutical analysis. *LC-GC Int.*, 8, No. 1 (1995) 26-33.
- 1399 Nimura, N., Itoh, H. and Kinoshita, T.: Diol-bonded silica gel as a restricted access packing forming a binary-layered phase for direct injection of serum for the determination of drugs. *J. Chromatogr. A*, 689 (1995) 203-210.
- 1400 Oscarsson, S., Angulo-Tatis, D., Chaga, G. and Porath, J.: Amphiphilic agarose-based adsorbents for chromatography. Comparative study of adsorption capacities and desorption efficiencies. *J. Chromatogr. A*, 689 (1995) 3-12.
- 1401 Pesek, J.J. and Matyska, M.T.: Synthesis and spectrometric characterization of a true diol bonded phase. *J. Chromatogr. A*, 687 (1994) 33-44.
- 1402 Petro, M., Berek, D. and Novak, I.: Composite sorbents for liquid chromatography. A size exclusion study of dextran gel incorporated into porous solid particles. *React. Polym.*, 23 (1994) 173-182; C.A., 122 (1995) 11701t.
- 1403 Rhee, D., Markovich, R., Chae, W.G., Qiu, X. and Pidgeon, C.: Chromatographic surfaces prepared from lyso phosphatidylcholine ligands. *Anal. Chim. Acta*, 297 (1994) 377-386.
- 1404 Rodrigues, A.E., Loureiro, J.M., Chenou, C., de la Vega, M.R.: Bioseparations with permeable particles. *J. Chromatogr. B*, 664 (1995) 233-240.
- 1405 Sarker, M. and Guiochon, G.: The packing stability of dynamically compressed preparative chromatography columns. *LC-GC*, 12 (1994) 300-306; C.A., 122 (1995) 70796a.
- 1406 Scholten, A.B., de Haan, J.W., Claessens, H.A., van de Ven, L.J.M. and Cramers, C.A.:  $^{29}\text{Si}$  NMR model dissolution study of the degradation of reversed phases for high-performance liquid chromatography. *Anal. Chem.*, 66 (1994) 4085-4092.
- 1407 Seely, R.J., Wight, H.D., Fry, H.H., Rudge, S.R. and Slaff, G.F.: Biotechnology product validation. Part 7: Validation of chromatography resin useful life. *BioPharm (Eugene)*, 7 (1994) 41-48; C.A., 121 (1994) 278870g - a review with 7 refs.
- 1408 Sentell, K.B., Bliesner, D.M. and Shearer, S.T.:  $^2\text{H}$  and  $^{13}\text{C}$  NMR studies of reversed phase liquid chromatographic stationary phases: Solvation and temperature effects. *Spec. Publ. - R. Soc. Chem.*, 139 (1994) 190-202; C.A., 121 (1994) 239082g.
- 1409 Shurygina, V.V., Stepnova, Z.N., Sturchak, S.V., Ryltsev, V.V. and Fulatov, V.N.: Method of sorbent preparation for affinity chromatography. *U.S.S.R. SU* 1,816,143 (Cl. B01J20/22), 30 May 1993, Appl. 4,488,422, 30 Sep. 1988; C.A., 122 (1995) 45399r.
- 1410 Smigol, V., Svec, F. and Frechet, J.M.J.: Two-dimensional high-performance liquid chromatography using monodisperse polymer beads containing segregated chemistries prepared by pore size specific functionalization. Single-column combinations of size exclusion or ion exchange with reversed-phase chromatography. *Anal. Chem.*, 66 (1994) 43-8-4315.
- 1411 Steenackers, D. and Sandra, P.: Preparation of small diameter thick film capillary columns for microcolumn separations. *J. Microcolumn Sep.*, 6 (1994) 361-367; C.A., 122 (1995) 22819p.
- 1412 Swart, R., Kraak, J.C. and Poppe, H.: Performance of an ethoxyethylacrylate stationary phase for open-tubular liquid chromatography. *J. Chromatogr. A*, 689 (1995) 177-187.
- 1413 Upchurch, P.E. and Schick, H.G.: Column for liquid chromatography. *Eur. Pat. Appl. EP* 624,795 (Cl. G01N30/60), 17 Nov. 1994, US Appl. 62,402, 14 May 1993; 14 p.; C.A., 122 (1995) 45289e.
- 1414 Van Zanten, J.H.: Terminally anchored chain interphases: their chromatographic properties. *Macromolecules*, 27 (1994) 6797-6807; C.A., 121 (1994) 264590d.
- 1415 Verhulst, H.A.M., van de Ven, L.J.M., de Haan, J.W., Claessens, H.A., Eisenbeiss, F. and Schick, H.G.: Patching in reversed-phase high-performance liquid chromatographic materials studied by solid-state NMR spectrometry. *J. Chromatogr. A*, 687 (1994) 213-221.
- 1416 Vermeulen, D.M. and Cantwell, F.F.: Slow change in the electrical potential at glass and silica surfaces due to  $\text{Na}^+$  sorption in the hydrated layer. *J. Chromatogr. A*, 693 (1995) 205-216.
- 1417 Welch, C.J.: Imprintable brush-type chiral stationary phase. *J. Chromatogr. A*, 689 (1995) 189-193.
- 1418 Whitman, D.A., Weber, T.P. and Blackwell, J.A.: Chemometric characterization of Lewis base-modified zirconia for normal phase chromatography. *J. Chromatogr. A*, 691 (1995) 205-212.
- 1419 Yu, J. and El Rassi, Z.: Chromatographic properties of zirconia-based stationary phases for ion exchange chromatography having surface bound cationic functions. *J. High Resolut. Chromatogr.*, 17 (1994) 705-712.
- 1420 Yu, J. and El Rassi, Z.: Preparation of amino-zirconia bonded phases and their evaluation in hydrophilic interaction chromatography of carbohydrates with pulsed amperometric detection. *J. High Resolut. Chromatogr.*, 17 (1994) 773-778.

For additional information see C.A.:

121 (1994) 233318s, 234008c, 244520c, 244549u, 270714p, 270858p, 270859q, 270864n, 270866q, 270867r, 270868s, 270869t, 270871n, 281074f, 283012h, 291616y, 291695y, 296323k, 301965c;  
122 (1995) 22863y, 32663x, 39752j, 45400j, 70989r.

See also 1301, 1326, 1346, 1353, 1435, 1464, 1466, 1469, 1482, 1492, 1493, 1496, 1499, 1504, 1511, 1512, 1549, 1567, 1588, 1595, 1623, 1746, 1759, 1767, 1792, 1796, 1853, 1872, 1873, 1884, 1917, 2220, 2263, 2269, 2497, 2520, 2534.

### 3d. Quantitative analysis

- 1421 Bahowick, T.J. and Synovec, R.E.: Correlation of quantitative analysis precision to retention time precision and chromatographic resolution for rapid, short column analysis. *Anal. Chem.*, 67 (1995) 631-640.
- 1422 Van den Bogaert, B., Boelens, H.F.M. and Smit, H.C.: Quantification of overlapping chromatographic peaks using a matched filter. *Chemom. Intell. Lab. Syst.*, 25 (1994) 297-311; *C.A.*, 122 (1995) 22813g.

For additional information see *C.A.*:  
121 (1994) 258979w.

See also 1704, 2364.

### 3e. Preparative scale chromatography

- 1423 Charton, F., Bailly, M. and Guiochon, G.: Recycling in preparative liquid chromatography. *J. Chromatogr. A*, 687 (1994) 13-31.
- 1424 Crétier, G., Neffati, J. and Rocca, J.L.: Preparative LC and preparative SFC: two complementary techniques in the fractionation of an impurity from a major component. *J. Chromatogr. Sci.*, 32 (1994) 449-454.
- 1425 Unger, K.K. (Editor): *Handbook of HPLC, Part 2: Preparative Liquid Column Chromatography*. GIT-Verlag, Darmstadt, 1994, 280 pp.; *C.A.*, 121 (1994) 229992q.

See also 1386, 1405, 1433, 1494, 1536, 1538, 1553, 1601, 1610, 1761, 1851, 1852, 1858, 1864, 1874, 1914, 1920, 1929, 1933, 1948, 1982, 2017, 2116, 2237, 2345, 2451, 2495.

### 3f. Programmed temperature, pressure, vapors, gradients

- 1426 Dolan, J.W.: Obtaining separations, part III: adjusting column conditions. *LC-GC*, 12 (1994) 520-524; *C.A.*, 122 (1995) 70797b.
- 1427 Lukulay, P.H. and McGuffin, V.L.: Solvent modulation in liquid chromatography: extension to serially coupled columns. *J. Chromatogr. A*, 691 (1995) 171-185.

See also 1354, 1530, 1547, 1797.

## 4. SPECIAL TECHNIQUES

### 4a. Automation

- 1428 Cooper, J.D.H., Buick, A.R., Sheung, C.T.F.: The development of a new automated configuration to extend the analytical capability of the ASTED system using the high-pressure liquid chromatographic analysis of a nucleoside in plasma as a model. *Lab. Rob. Autom.*, 6 (1994) 15-20; *C.A.*, 121 (1994) 250049x.
- 1429 Smead, F.: A guide to the next step in data management. *Int. Lab.*, 24, No. 11 (1994) 14-16.
- 1430 Wieling, J., Jonkman, J.H.G., Hempenius, J. and Mensink, C.K.: Integration of robotics and chemometrics for the automated optimization of drug extractions. *Chemom. Intell. Lab. Syst.*, 25 (1994) 355-366; *C.A.*, 122 (1995) 17325v.

See also 1668, 1980, 2210, 2227, 2277, 2298, 2336, 2343, 2412, 2421, 2435.

### 4b. Computerization and modelling

- 1431 Andrews, M.: Adapting to a chemist's environment: the future trend in chromatography data management. *Am. Lab. (Shelton)*, 26 (1994) 49-55; *C.A.*, 121 (1994) 314863t.
- 1432 Cepria, G. and Castillo, J.R.: Influence of the detection system on chromatographic profiles. Comparison of a refractive index and surface plasmon resonance based detector for HPLC. *Quim. Anal. (Barcelona)*, 13 (1994) 63-66; *C.A.*, 122 (1995) 71066z.
- 1433 Felinger, A. and Guiochon, G.: Computer simulations in non-linear chromatography. *TrAC*, 14 (1995) 6-10.
- 1434 Guillemin, C.L.: The deferred standard. *Process Control Qual.*, 6 (1994) 9-25; *C.A.*, 122 (1995) 45331n - a review with 19 refs.
- 1435 Hamoir, T., Sanchez, F.C., Bourguignon, B. and Massart, D.L.: Spectral mapping analysis: a method for the characterization of stationary phases. *J. Chromatogr. Sci.*, 32 (1994) 488-498.
- 1436 Head, M. and Yellin, B.: Review of the benefits of a chromatography data system on Windows NT. *Am. Lab. (Shelton)*, 26 (1994) 48T, 48U, 48V, 48X, 48Y; *C.A.*, 121 (1994) 291609y.
- 1437 Malmquist, G. and Danielsson, R.: Alignment of chromatographic profiles for principal component analysis: a prerequisite for fingerprinting methods. *J. Chromatogr. A*, 687 (1994) 71-88.
- 1438 Mowry, D. and Doll, G.: How software beta testing enhances product quality. *Int. Lab.*, 24, No. 11 (1994) 17-20.
- 1439 Reh, E.: Peak-shape analysis for unresolved peaks in chromatography: comparison of algorithms. *TrAC*, 14 (1995) 1-5.
- 1440 Tobis, J.: (Methods for evaluation of channeling effect in fixed-bed columns with granular packing). *Przem. Chem.*, 73 (1994) 257-259; *C.A.*, 121 (1994) 233595e - a review with 12 refs.
- 1441 Wiblin, D.J.: Computer aided desk-top scale-up and optimization of chromatographic processes. *Spec. Publ.-R. Soc. Chem.*, 158(Separations for Biotechnology 3) (1994) 572-578; *C.A.*, 122 (1995) 8106x.
- 1442 Zhao, G.-l., Zhao, Y.-m. and Liu, L.-n.: Microcomputer simulation of elution behavior for dicarboxylic acids in ion-exclusion chromatography. *Chem. Res. Chin. Univ.*, 10 (1994) 79-84; *C.A.*, 122 (1995) 45358b.

For additional information see C.A.:

- 121 (1994) 291557e;  
122 (1995) 11185c.

See also 1293, 1325, 1330, 1331, 1332, 1345, 1421, 1429,  
1430, 1498, 1586, 1830, 2249, 2253, 2270, 2290,  
2442, 2515, 2533.

4c. *Combination with other physico-chemical techniques (MS, IR etc.)*

- 1443 Bendicho, C.: Evaluation of an automated thermospray interface for coupling electrothermal atomization atomic absorption spectrometry and liquid chromatography. *Anal. Chem.*, 66 (1994) 4375-4381.
- 1444 Capiello, A. and Famiglini, G.: Evaluation of the performance of a microflow rate LC/MS particle beam interface. *Anal. Chem.*, 66 (1994) 3970-3976.
- 1445 Capiello, A. and Famiglini, G.: Analysis of thermally unstable compounds by a liquid chromatography/mass spectrometry particle beam interface with a modified ion source. *Anal. Chem.*, 67 (1995) 412-419.
- 1446 Caprioli, R.M.: Online LC/MS and CE/MS. In: Matsuo, T. (Editor), *Biol. Mass Spectrom.: Present Future, [Proc. Kyoto '92 Int. Conf.]*, Wiley, Chichester, 1994, pp. 75-100; C.A., 121 (1994) 249811h - a review with 65 refs.
- 1447 Engelhardt, U.H. and Wagner-Redeker, W.: (HPLC-MS (mass spectrometry). Techniques and applications. Part 1. HPLC/MS coupling techniques). *GIT Spez. Chromatogr.*, 14 (1994) 5-9; C.A., 122 (1995) 22801b - a review with 8 refs.
- 1448 Gumpersbach, J.: (Chromatography-mass spectrometry combination widened. The MD 800 family from Fison: routine mass spectrometric systems for chromatography). *GIT Fachz. Lab.*, 38 (1994) 846-847; C.A., 121 (1994) 270768j.
- 1449 Harada, K., Suzuki, M. and Oka, H.: Chromatography and mass spectrometry for analysis of natural products. In: Matsuo, T. (Editor), *Biol. Mass Spectrom.: Present Future, [Proc. Kyoto '92 Int. Conf.]* 1992, Wiley, Chichester, 1994, pp. 605-630; C.A., 121 (1994) 225275k - a review with many refs.
- 1450 Hutta, M., Kaniansky, D., Kovalcikova, E., Marak, J., Chalanyova, M., Madajova, V. and Simunicova, E.: Preparative capillary isotachopheresis as a sample pretreatment technique for complex ionic matrices in high-performance liquid chromatography. *J. Chromatogr. A*, 689 (1995) 123-133.
- 1451 Krull, I.S., Mhatre, R. and Cuniff, J.: Biopolymer detection, Part I - Mass spectrometry. *LC-GC Int.*, 8, No. 1 (1995) 14-20.
- 1452 Linscheid, M. and Westmoreland, D.G.: Analytical techniques for trace organic compounds. VI. Application of liquid chromatography-mass spectrometry. *Pure Appl. Chem.*, 66 (1994) 1913-1930; C.A., 122 (1995) 71049w - a review with 203 refs.
- 1453 Pálmarasdóttir, S. and Edholm, L.-E.: Enhancement of selectivity and concentration sensitivity in capillary zone electrophoresis by on-line coupling with column liquid chromatography and utilizing a double stacking procedure allowing for microliter injections. *J. Chromatogr. A*, 693 (1995) 131-143.
- 1454 Rottmann, L. and Heumann, K.G.: Development of an on-line isotope dilution technique with HPLC/ICP-MS for accurate determination of elemental species. *Fresenius J. Anal. Chem.*, 350 (1994) 221-227.

- 1455 Seubert, A.: On-line coupling between chromatography and inductively coupled plasma mass spectrometry - a current assessment. *Fresenius J. Anal. Chem.*, 350 (1994) 210-220.
- 1456 Stroh, J.G. and Rinehart, K.L.: Liquid chromatography/fast atom bombardment mass spectrometry. *Top Mass Spectrom.*, 1 (Experimental Mass Spectrometry) (1994) 287-311; C.A., 122 (1995) 4564x - a review with 54 refs.
- 1457 Wang, A.P.L., Guo, X. and Li, L.: Liquid chromatography/time-of-flight mass spectrometry with a pulsed sample introduction interface. *Anal. Chem.*, 66 (1994) 3664-3675.

For additional information see C.A.:

- 121 (1994) 220589f, 220591a, 244386p, 291568j;  
122 (1995) 19999s.

See also 1358, 1432, 1517, 1535, 1634, 1641, 1649, 1695,  
1728, 1736, 1741, 1742, 1760, 1772, 1775, 1777,  
1780, 1819, 1824, 1849, 1850, 1861, 1877, 1985,  
1996, 2053, 2060, 2073, 2077, 2117, 2128, 2143,  
2183, 2185, 2191, 2211, 2221, 2233, 2245, 2258,  
2260, 2265, 2271, 2303, 2324, 2353, 2357, 2362,  
2368, 2377, 2385, 2402, 2411, 2416, 2423, 2446,  
2447, 2461, 2462, 2487, 2517, 2521, 2527.

4d. *Affinity chromatography (advances)*

- 1458 Benes, M.J., Stambergova, A. and Scouten, W.H.: Affinity chromatography with immobilized benzenboronates. *Mol. Interact. Biosep.*, (1993) 313-322; C.A., 121 (1994) 249861z - a review with many refs.
- 1459 Boschetti, E.: New diffusive matrix to improve throughput in affinity chromatography applications. *Prog. Biotechnol.*, 9(CEB6: Proceedings of the 6th European Congress on Biotechnology, 1993, Pt. 1) (1994) 547-550; C.A., 121 (1994) 225372q.
- 1460 Chaiken, I.: Affinity chromatography using immobilized anti-sense-family peptides. *Mol. Interact. Biosep.*, (1993) 169-177; C.A., 121 (1994) 249854z - a review with many refs.
- 1461 Clonis, Y.D.: Affinity methods in bioseparation. *Prog. Biotechnol.*, 9(CEB6: Proceedings of the 6th European Congress on Biotechnology, 1993, Pt. 1) (1994) 527-533; C.A., 121 (1994) 225265g.
- 1462 Galaev, I.Yu. and Mattiasson, B.: Polymer-shielded dye-affinity chromatography. *Spec. Publ. - R. Soc. Chem.*, 158(Separations for Biotechnology 3) (1994) 179-185; C.A., 121 (1994) 296299g.
- 1463 Godfrey, M.A.J., Kwasowski, P., Clift, R. and Marks, V.: Rivanol-silica: a dye-ligand solid phase for affinity purifying proteins. *Spec. Publ. - R. Soc. Chem.*, 158(Separations for Biotechnology 3) (1994) 200-206; C.A., 121 (1994) 296301b.
- 1464 Gu, J., Stephenson, G. and Iadarola, M.J.: Recombinant proteins attached to a nickel-NTA column: use in affinity purification of antibodies. *BioTechniques*, 17 (1994) 257-262; C.A., 121 (1994) 228115u.
- 1465 Hage, D.S., Noctor, T.A.G. and Wainer, I.W.: Characterization of the protein binding of chiral drugs by high-performance affinity chromatography. Interactions of R- and S-ibuprofen with human serum albumin. *J. Chromatogr. A*, 693 (1995) 23-32.

- 1466 Hermanson, G.T., Mattson, G.R. and Krohn, R.I.: Preparation and use of immunoglobulin-binding affinity supports on Emphaze beads. *J. Chromatogr. A*, 691 (1995) 113-122.
  - 1467 Katoh, S., Terashima, M., Sada, E., Utsumi, H., Kamiya, Y., Yamada, K. and Majima, T.: Characteristics of a perfusion-type support in affinity chromatography. *J. Ferment. Bioeng.*, 78 (1994) 246-249; C.A., 121 (1994) 225415f.
  - 1468 Lindgren, G.E.S.: Immobilized metal affinity chromatography (IMAC). *Am. Biotechnol. Lab.*, 12 (1994) 36; C.A., 121 (1994) 225267j - a review with 7 refs.
  - 1469 Liu, X.-C. and Scouten, W.H.: New ligands for boronate affinity chromatography. *J. Chromatogr. A*, 687 (1994) 61-69.
  - 1470 Mattiasson, B., Linne, E. and Kaul, R.: Use of heterobifunctional ligands in affinity chromatographic processes. *Mol. Interact. Biosep.*, (1993) 395-401; C.A., 121 (1994) 249866e - a review with many refs.
  - 1471 Nachman-Clewner, M., Spence, C. and Bailon, P.: Receptor-affinity chromatography (RAC). *Mol. Interact. Biosep.*, (1993) 139-149; C.A., 121 (1994) 249853y - a review with many refs.
  - 1472 Narinesingh, D. and Ngo, T.T.: 2-Fluoro-1-methylpyridinium (FMP) salt-activated gels: properties and uses in affinity chromatography and enzyme immobilization for analytical applications. *Mol. Interact. Biosep.*, (1993) 49-68; C.A., 121 (1994) 275501b.
  - 1473 Ohlson, S. and Zopf, D.: Weak-affinity chromatography. *Mol. Interact. Biosep.*, (1993) 15-25; C.A., 121 (1994) 249849b - a review with many refs.
  - 1474 Parikh, I. and Cuatrecasas, P.: Affinity chromatography: an overview. *Mol. Interact. Biosep.*, (1993) 3-13; C.A., 121 (1994) 249848a - a review with many refs.
  - 1475 Sada, E. and Katoh, S.: Suitable antibodies as ligands in affinity chromatography of biomolecules. *Mol. Interact. Biosep.*, (1993) 205-211; C.A., 121 (1994) 250405k.
  - 1476 Safi, A., Saunier, M., Gastin, I., Alibada, Y., Dugue, B. and Gueant, J.-L.: Intrinsic factor covalently bound to Sepharose as affinity medium for the purification of a soluble intrinsic factor receptor from human urine. *J. Chromatogr. B*, 644 (1995) 253-259.
  - 1477 Santambien, P., Sdiqui, S., Hubert, E., Girot, P., Roche, A.C., Monsigny, M. and Boschetti, E.: *In vitro* toxicity assays for dye ligands used in affinity chromatography. *J. Chromatogr. B*, 664 (1995) 241-246.
  - 1478 Stellwagen, E.: Affinity chromatography with immobilized dyes. *Mol. Interact. Biosep.*, (1993) 247-255; C.A., 121 (1994) 250060u.
  - 1479 Tosa, T., Sato, T., Watanabe, T. and Minobe, S.: Affinity chromatographic removal of pyrogens. *Mol. Interact. Biosep.*, (1993) 323-332; C.A., 121 (1994) 249862a - a review with many refs.
  - 1480 Velander, W.H., Kaster, J.A. and Glasser, W.G.: Lignocellulosic and cellulosic beads for use in affinity and immunoaffinity chromatography of high molecular weight proteins. *U.S. US 5,328,603* (Cl. 210-198.2; B01D15/08), 12 Jul. 1994, US Appl. 496,314, 20 Mar. 1990; 24 pp.; C.A., 121 (1994) 276172g.
  - 1481 Vijayalakshmi, M.A.: Pseudo-biospecific affinity ligand chromatography: the case of immobilized histidine as a universal ligand. *Mol. Interact. Biosep.*, (1993) 257-275; C.A., 121 (1994) 249859e - a review with many refs.
  - 1482 Weetall, H.H.: Affinity chromatography on inorganic support materials. *Mol. Interact. Biosep.*, (1993) 27-48; C.A., 121 (1994) 249850v - a review with many refs.
  - 1483 West, I. and Goldring, O.: Lectin affinity chromatography. *Mol. Biotechnol.*, 2 (1994) 147-155; C.A., 122 (1995) 4625t.
  - 1484 Wulff, G.: Biorecognition in molecularly imprinted polymers: concept, chemistry, and application. *Mol. Interact. Biosep.*, (1993) 363-381; C.A., 121 (1994) 249864c - a review with many refs.
  - 1485 Zhao, C., Wang, L., Xie, S.-s. and Long, Z.-z.: (Preparation of Cibacron Blue Sephadex and its application on affinity chromatography). *Shengwu Huaxue Zazhi*, 10 (1994) 621-625; C.A., 121 (1994) 298620d.
- For additional information see C.A.:  
121 (1994) 250632g, 250635k.
- See also 1374, 1409, 1789, 1796, 1848, 1867, 1868, 1881, 1882, 1893, 1904, 1914, 1915, 1919, 1928, 1935, 1947, 1948, 1978, 1980, 1987, 1989, 2031, 2043, 2044, 2066, 2226, 2227.
- 4e. *Functional analysis*
- 1486 Morley, J.A., Elrod, L., Jr. and Bauer, J.F.: Determination of the active hydrogen content in pyridine-borane complex by Schiff base reduction and high-performance liquid chromatography. *Anal. Chem.*, 66 (1994) 4283-4287.
- See also 1385.
- 4f. *Trace analysis and pre separation techniques*
- 1487 Capparella, M., Foster, W., III, Larrousse, M., Phillips, D.J., Pomfret, A. and Tuvim, Y.: Characteristics and applications of a new high-performance liquid chromatography guard column. *J. Chromatogr. A*, 691 (1995) 141-150.
  - 1488 Fritz, J.S., Dumont, P.J. and Schmidt, L.W.: Methods and materials for solid-phase extraction. *J. Chromatogr. A*, 691 (1995) 133-140.
  - 1489 Gelencsér, A., Kiss, G., Krivácsy, Z., Varga-Puchony, Z. and Hlavay, J.: A simple method for the determination of capacity factor on solid-phase extraction cartridges. I. *J. Chromatogr. A*, 693 (1995) 217-225.
  - 1490 Schlabach, M., Biseth, A., Gundersen, H. and Oehme, M.: On-line GPC/carbon clean-up method for determination of PCDD/F in sediment and sewage sludge samples. *Organohalogen Compd.*, 11 (1993) 71-74; C.A., 121 (1994) 212154v.
- For additional information see C.A.:  
121 (1994) 220508d.
- See also 1757, 2218, 2225, 2507, 2515, 2525, 2526, 2529.
- 4g. *Enantiomers, separation*
- 1491 Allenmark, S. and Andersson, S.: Chromatographic resolution of chiral compounds by means of immobilized proteins. *Mol. Interact. Biosep.*, (1993) 179-187; C.A., 121 (1994) 249855a - a review with many refs.

- 1492 Andersson, L.I., Ekberg, B. and Mosbach, K.: Bioseparation and catalysis in molecularly imprinted polymers. *Mol. Interact. Biosep.*, (1993) 383-394; *C.A.*, 121 (1994) 249865d - a review with no refs.
  - 1493 Aubry, A.-F., Markoglou, N., Descorps, V., Wainer, I.W. and Félix, G.: Evaluation of a chiral stationary phase based on mixed immobilized proteins. *J. Chromatogr. A*, 685 (1994) 1-6.
  - 1494 Camacho-Torralba, P.L. and Vigh, G.: High-performance chiral displacement chromatographic separations in the normal-phase mode. III. Separation of the enantiomers of 5-vinylpyrrolidin-2-one using the Chiralcel-OD stationary phase. *J. Chromatogr. A*, 691 (1995) 213-216.
  - 1495 Hu, R., Takeuchi, T., Jin, J.-Y. and Miwa, T.: Separation of enantiomers by microcolumn liquid chromatography with methylated  $\beta$ -cyclodextrin as mobile phase additive. *Anal. Chim. Acta*, 295 (1994) 173-179.
  - 1496 Hyun, M.H. and Min, C.S.: A new chiral stationary phase bearing both  $\pi$ -acidic and  $\pi$ -basic sites derived from (S)-tyrosine for the liquid chromatographic resolution of racemates. *Chem. Lett.*, (1994) 1463-1466; *C.A.*, 121 (1994) 229986r.
  - 1497 Hyun, M.H., Ryoo, J.-J., Cho, Y.J. and Jin, J.S.: Unusual examples of the liquid chromatographic resolution of racemates. Resolution of  $\pi$ -donor analytes on a  $\pi$ -donor chiral stationary phase. *J. Chromatogr. A*, 692 (1995) 91-96.
  - 1498 Lipkowitz, K.B. and Anderson, A.G.: Computational aspects in supramolecular chemistry: Chiral discrimination in chromatography. *NATO ASI Ser., Ser. C*, 426 (1994) 183-198; *C.A.*, 122 (1995) 71070w.
  - 1499 Loun, B. and Hage, D.S.: Chiral separation mechanisms in protein-based HPLC columns. 1. Thermodynamic studies of (R)- and (S)-warfarin binding to immobilized human serum albumin. *Anal. Chem.*, 66 (1994) 3814-3822.
  - 1500 Mano, N., Oda, Y., Asakawa, N., Yoshida, Y., Sato, T. and Miwa, T.: Studies of ovomucoid-, avidin-, conalbumin- and flavoprotein-conjugated chiral stationary phases for separation of enantiomers by high-performance liquid chromatography. *J. Chromatogr. A*, 687 (1994) 223-232.
  - 1501 Pullen, R.H., Brennan, J.J. and Patonay, G.: Chiral separation retention mechanisms in high-performance liquid chromatography using bare silica stationary phase and  $\beta$ -cyclodextrin as a mobile phase additive. *J. Chromatogr. A*, 691 (1995) 187-193.
  - 1502 Quintero, G., Vo, M., Farkas, G. and Vigh, G.: Series of homologous displacers for preparative chiral displacement chromatographic separations on Cyclobond-II columns. *J. Chromatogr. A*, 693 (1995) 1-5.
  - 1503 Sellergren, B. and Shea, K.J.: Origin of peak asymmetry and the effect of temperature on solute retention in enantiomer separations on imprinted chiral stationary phases. *J. Chromatogr. A*, 690 (1995) 29-39.
  - 1504 Simek, Z. and Vespalec, R.: Interpretation of enantioselective activity of albumin used as the chiral selector in liquid chromatography and electrophoresis. *J. Chromatogr. A*, 685 (1994) 7-14.
  - 1505 Tittelbach, V. and Gilpin, R.K.: Species dependency of the liquid chromatographic properties of silica-immobilized serum albumins. *Anal. Chem.*, 67 (1995) 44-47.
  - 1506 Welch, C.J. and Perrin, S.R.: Improved chiral stationary phase for  $\beta$ -blocker enantioseparations. *J. Chromatogr. A*, 690 (1995) 218-225.
  - 1507 Williams, K.L. and Stalcup, A.M.: Chromatography with cyclodextrin-based stationary phases. *Mol. Interact. Biosep.*, (1993) 189-202; *C.A.*, 121 (1994) 249856b - a review with many refs.
- For additional information see *C.A.*:  
121 (1994) 244520c, 270805u, 270869t, 291695y, 301311m.
- See also 1377, 1384, 1391, 1395, 1417, 1453, 1545, 1610, 1611, 1623, 1630, 1726, 1737, 1738, 1739, 1746, 1753, 1766, 1773, 1776, 1785, 1788, 1800, 1811, 2086, 2149, 2173, 2267, 2269, 2276, 2281, 2282, 2291, 2297, 2299, 2302, 2303, 2305, 2306, 2315, 2337, 2344, 2348, 2349, 2354, 2403, 2433.
- #### 4h. Other special techniques
- 1508 Chen, A. and Lunte, C.E.: Microdialysis sampling coupled on-line to fast microbore liquid chromatography. *J. Chromatogr. A*, 691 (1995) 29-35.
  - 1509 Cui, Y. and Olesik, S.V.: Reversed-phase high-performance liquid chromatography using enhanced-fluidity mobile phases. *J. Chromatogr. A*, 691 (1995) 151-162.
  - 1510 Goto, M., Imamura, T. and Hirose, T.: Axial dispersion in liquid magnetically stabilized fluidized beds. *J. Chromatogr. A*, 690 (1995) 1-8.
  - 1511 Guerrier, L., Flayeux, I., Boschetti, E. and Radosevich, M.B.: Specific sorbent to remove solvent-detergent mixtures from virus-inactivated biological fluids. *J. Chromatogr. B*, 664 (1995) 119-125.
  - 1512 Jungbauer, A., Lettner, H.P., Guerrier, L. and Boschetti, E.: Chemical sanitization in process chromatography. Part 2: *In situ* treatment of packed columns and long-term stability of resins. *BioPharm (Eugene)*, 7 (1994) 37-42; *C.A.*, 121 (1994) 225374s.
  - 1513 Kitagawa, S. and Tsuda, T.: Effects of pH and organic solvent on chromatographic behavior in capillary electrochromatography. *J. Microcolumn Sep.*, 6 (1994) 91-96; *C.A.*, 121 (1994) 148022w.
  - 1514 Mayer, M.L. and Poole, C.F.: Identification of the procedural steps that affect recovery of semi-volatile compounds by solid-phase extraction using cartridge and particle-loaded membrane (disk) devices. *Anal. Chim. Acta*, 294 (1994) 113-126.
  - 1515 Oosterkamp, A.J., Irth, H., Tjaden, U.R. and van der Greef, J.: On-line coupling of liquid chromatography to biochemical assays based on fluorescent-labeled ligands. *Anal. Chem.*, 66 (1994) 4295-4301.
  - 1516 Ricker, R.D., Sandoval, L.A., Justice, J.D. and Geiser, F.O.: Multivariate visualization in the size-exclusion chromatography and pattern recognition of biological samples. *J. Chromatogr. A*, 691 (1995) 67-79.
  - 1517 Roessner, D. and Kulicke, W.-M.: On-line coupling of flow field-flow fractionation and multi-angle laser light scattering. *J. Chromatogr. A*, 687 (1994) 249-258.
  - 1518 Strobel, G.J.: (Perfusion chromatography). *GIT Spez. Chromatogr.*, 14 (1994) 25-28; *C.A.*, 122 (1995) 22802c.

- 1519 Van Buel, M.J., van der Wielen, L.A.M., Meester, R., Verpoorte, R. and Luyben, K.C.A.M.: Modeling of centrifugal partition chromatography. *Prog. Biotechnol.*, 9(ECB6: Proceedings of the 6th European Congress on Biotechnology, 1993, Pt. 1) (1994) 515-519; C.A., 121 (1994) 225370n.
- 1520 Wenzel, U. and Deron, S.: Design parameters of a decontamination process by solvent extraction chromatography. *Solvent Extr. Ion Exch.*, 12 (1994) 789-801; C.A., 121 (1994) 266078s.
- 1521 Zhang, J. and Fang, Q.: Application of high speed counter-current chromatography to the separation of stilbene derivatives from the roots of *Lindera reflexa*. *Planta Med.*, 60 (1994) 190-191; C.A., 121 (1994) 65385r.

For additional information see C.A.:

121 (1994) 244386p, 270705m, 270805u.

See also 1308, 1338, 1351, 1495, 1544, 1549, 1610, 1668, 1873, 1880, 1996, 2116, 2134, 2141, 2452, 2489.

## 5. HYDROCARBONS AND HALOGEN DERIVATIVES

### 5a. Aliphatic hydrocarbons

- 1522 Thelin, A., Peterson, E., Hutson, J.L., McCarthy, A.D., Ericsson, J. and Dallner, G.: Effect of squalstatin 1 on the biosynthesis of the mevalonate pathway lipids. *Biochim. Biophys. Acta*, 1215 (1994) 245-249.

See also 1337.

### 5b. Cyclic hydrocarbons, fullerenes

- 1523 Diack, M. and Quiochon, G.: Analysis and purification of the fullerenes by liquid chromatography. *Quim. Anal. (Barcelona)*, 12 (1993) 75-82; C.A., 121 (1994) 291379y - a review with 39 refs.
- 1524 Fladung, N.C.: Optimization of automated solid-phase extraction for quantitation of polycyclic aromatic hydrocarbons in aqueous media by high-performance liquid chromatography-UV detection. *J. Chromatogr. A*, 692 (1995) 21-26.
- 1525 Garcia, M.A. and Marina, M.L.: Study of the  $k'$  or  $\log k'$ -log  $P_{ow}$  correlation for a group of benzene derivatives and polycyclic aromatic hydrocarbons in micellar liquid chromatography with a C<sub>8</sub> column. *J. Chromatogr. A*, 687 (1994) 233-239.
- 1526 Heymann, D., Chibante, L.P.F. and Smalley, R.E.: Determination of C<sub>60</sub> and C<sub>70</sub> fullerenes in geologic materials by high-performance liquid chromatography. *J. Chromatogr. A*, 689 (1995) 157-163.
- 1527 Inukai, Y., Attalla, M.I., Pang, L.S.K. and Wilson, M.A.: Separation of fullerenes by chromatography on coal. *Fuel*, 74 (1995) 83-87; C.A., 122 (1995) 59888m.
- 1528 Kayali, M.N., Rubio-Barroso, S. and Polo-Diez, L.M.: Determination of PAHs in particulate air by micellar liquid chromatography. *J. Liq. Chromatogr.*, 17 (1994) 3623-3640.
- 1529 Lai, F. and White, L.: Automated precolumn concentration and high-performance liquid chromatographic analysis of polynuclear aromatic hydrocarbons in water using a single pump and a single valve. *J. Chromatogr. A*, 692 (1995) 11-20.

- 1530 Lubda, D., Mueller, M., Battermann, G. and Mayer, B.: (A stationary phase HPLC-separation of 18 polycyclic aromatic hydrocarbons (PAH's)). *GIT Fachz. Lab.*, 38 (1994) 12-15; C.A., 121 (1994) 307772j.
- 1531 McKinney, D.E., Clifford, D.J., Hou, L., Bogdan, M.R. and Hatcher, P.G.: High performance liquid chromatography of coal liquefaction process streams using normal-phase separation with diode array detection. *Prepr. Pap.-Am. Chem. Soc., Div. Fuel Chem.*, 39 (1994) 835-839; C.A., 121 (1994) 259415w.
- 1532 Niederländer, H.A.G., Nuijens, M.J., Dozy, E.M., Gooijer, C. and Velthorst, N.H.: Dioxetane chemiluminescence detection in liquid chromatography based on photosensitized on-line generation of singlet molecular oxygen; a thorough examination of experimental parameters and application to polychlorinated biphenyls. *Anal. Chim. Acta*, 297 (1994) 349-368.
- 1533 Rodríguez Delgado, M.A., Sánchez, M.J., González, V. and García Montelongo, F.: Influence of alcoholic modifiers on the selectivity of the separation of a group of polycyclic aromatic hydrocarbons by micellar liquid chromatography. *Anal. Chim. Acta*, 298 (1994) 423-430.
- 1534 Selegue, J.P., Shaw, J.P., Guarr, T.F. and Meier, M.S.: Purification and characterization of the larger fullerenes: New aspects of C<sub>76</sub>, C<sub>78</sub> and C<sub>84</sub>. *Proc. Electrochem. Soc.*, 94-24 (1994) 1274-1291; C.A., 122 (1995) 22362j.
- 1535 Sim, P.G., Quilliam, M.A., Leblanc, M.D. and Karunanithy, S.: Development and application of isotope dilution internal standard solutions for the chromatographic/mass spectrometric determination of polycyclic aromatic compounds in complex mixtures. *Polycyclic Aromat. Compd.*, 3 (1993) 55-63; C.A., 121 (1994) 270821w.
- 1536 Story, J.N.: Preparative separation and characterization of coal liquid aromatics. *Prepr. Pap.-Am. Chem. Soc., Div. Fuel Chem.*, 39 (1994) 772-776; C.A., 121 (1994) 283358a.
- 1537 Visser, T., Vredenburg, M.J. and de Jong, A.P.J.M.: Confirmational analysis of polycyclic aromatic hydrocarbons in soil extracts by cryotrapping gas chromatography-Fourier transform infrared spectrometry. *J. Chromatogr. A*, 687 (1994) 303-313.
- 1538 Wu, Y., Sun, Y., Gu, Z., Zhou, X., Xiong, Y., Sun, B. and Jin, Z.: A combined recrystallization and preparative liquid chromatographic method for the isolation of pure C<sub>70</sub> fullerene. *Carbon*, 32 (1994) 1180-1182; C.A., 121 (1994) 291198p.

For additional information see C.A.:

121 (1994) 220664b, 220693k, 233989m, 258979w.

See also 1418, 1427, 1435.

### 5c. Halogen derivatives

- 1539 Johansen, H.R., Becher, G. and Greibrokk, T.: Determination of planar PCBs by combining on-line SFE-HPLC and GC-ECD or GC/MS. *Anal. Chem.*, 66 (1994) 4068-4073.
- 1540 Picó, Y., Redondo, M.J., Font, G. and Mañes, J.: Solid-phase extraction on C<sub>18</sub> in the trace determination of selected polychlorinated biphenyls in milk. *J. Chromatogr. A*, 693 (1995) 339-346.

For additional information see C.A.:

121 (1994) 259019v.

See also 1381, 2220.

5d. *Complex hydrocarbon mixtures (incl. analysis of tars, bitumens and mineral oils)*

- 1541 Acevedo, S., Escobar, G., Ranaudo, M.A. and Gutierrez, L.B.: Discotic shape of asphaltenes obtained from GPC (gel permeation chromatography) data. *Fuel*, 73 (1994) 1807-1809; C.A., 121 (1994) 259312k.
- 1542 Andersen, S.I.: Concentration effects in HPLC-SEC analysis of petroleum asphaltenes. *J. Liq. Chromatogr.*, 17 (1994) 4065-4079.
- 1543 Garrick, N.W.: Use of gel-permeation chromatography in predicting properties of asphalt. *J. Mater. Civ. Eng.*, 6 (1994) 376-389; C.A., 121 (1994) 262029d.
- 1544 Saini, A.K. and Song, C.: Two-dimensional HPLC and GC-MS of oils from catalytic coal liquefaction. *Prepr. Pap.-Am. Chem. Soc., Div. Fuel Chem.*, 39 (1994) 796-800; C.A., 121 (1994) 259410r.

6. ALCOHOLS

- 1545 Kim, J.-H., Nishida, Y., Ohnui, H. and Meguro, H.: Simple and highly sensitive high-performance liquid chromatographic method for separating enantiomeric diacylglycerols by direct derivatization with a fluorescent chiral agent, (S)-(+)-2-tert.-butyl-2-methyl-1,3-benzodioxole-4-carboxylic acid. *J. Chromatogr. A*, 693 (1995) 241-249.
- 1546 Rollag, J., Liu, T. and Hage, D.S.: Determination of 3-methoxy-4-hydroxyphenylethylene glycol in urine using reversed-phase liquid chromatography with column switching and electrochemical detection. *J. Chromatogr. B*, 663 (1995) 193-200.

See also 1381, 1514, 1565, 1569, 1570, 1606, 2254.

7. PHENOLS

- 1547 Leal, E.R., Rodriguez-Vazquez, R. and Galindo, T.: Separation of phenolic compounds from sugarcane bagasse pith and their determination by HPLC. *J. Wood Chem. Technol.*, 14 (1994) 369-382; C.A., 121 (1994) 207794z.
- 1548 Liu, Y., Lopez-Avila, V., Alcaraz, M. and Jones, T.L.: Centrifugal partition chromatographic extraction of phenols and organochlorine pesticides from water sample. *Anal. Chem.*, 66 (1994) 4483-4489.
- 1549 Yang, S. and Khaleli, M.G.: Stationary phase effects on retention behavior of phenols in micellar liquid chromatography: Perfluorooctane vs. C<sub>18</sub>. *Anal. Chim. Acta*, 294 (1994) 135-143.

See also 1367, 1381, 1418, 1525.

8. SUBSTANCES CONTAINING HETEROCYCLIC OXYGEN

8a. *Flavonoids*

- 1550 Costa, S.S., Jossang, A., N'Bodo, B., Souza, M.L.M. and Moraes, V.L.G.: Patuletin acetylramnosides from *Kalanchoe brasiliensis* as inhibitors of human lymphocyte proliferative activity. *J. Natural Prod.*, 57 (1994) 1503-1510.
- 1551 Das, D.K.: Naturally occurring flavonoids: structure, chemistry, and high-performance liquid chromatography methods for separation and characterization. *Methods Enzymol.*, 234 (Oxygen Radicals in Biological Systems, Pt. D) (1994) 410-420 - a review with 45 refs.
- 1552 Franke, A.A. and Custer, L.J.: High-performance liquid chromatographic assay of isoflavonoids and coumestrol from human urine. *J. Chromatogr. B*, 662 (1994) 47-60.
- 1553 Häberlein, H. and Tschiersch, K.-P.: 2,5-Dihydroxy-7-methoxy-6,8-dimethylflavan-3-one, a new flavonoid from *Leptospermum scoparium*: *in vitro* affinity to the benzodiazepine binding site of the GABA<sub>A</sub> receptor-chloride channel complex. *Pharmazie*, 49 (1994) 860.
- 1554 Schulz, H.-U., Schürer, M., Krumbiegel, G., Wächter, W., Weyhenmeyer, R. and Seidel, G.: Untersuchungen zum Freisetungsverhalten und zur Bioäquivalenz von Silymarin-Präparaten. *Arzneim.-Forsch.*, 45 (1995) 61-64.

For additional information see C.A.:

121 (1994) 213096q, 213107u, 308460t.

See also 2233.

8b. *Aflatoxins and other mycotoxins*

- 1555 Gelosa, M. and Buzzetti, I.: (Chromatographic techniques for aflatoxin determination in milk). *Ind. Aliment. (Pinerolo)*, 33 (1994) 315-317; C.A., 121 (1994) 229127m.
- 1556 Holcomb, M., Thompson, H.C., Jr., Lipe, G. and Hankins, L.J.: HPLC with electrochemical and fluorescence detection of the OPA/2-methyl-2-propanethiol derivative of fumonisin B<sub>1</sub>. *J. Liq. Chromatogr.*, 17 (1994) 4121-4129.
- 1557 Shephard, G.S., Thiel, P.G. and Sydenham, E.W.: Liquid chromatographic determination of the mycotoxin fumonisin B<sub>2</sub> in physiological samples. *J. Chromatogr. A*, 692 (1995) 39-43.

See also 2458.

8c. *Other compounds with heterocyclic oxygen (incl. tannins)*

- 1558 Steinert, J., Khalaf, H. and Rimpler, M.: HPLC separation and determination of naphthol[2,3-b]furan-4,9-diones and related compounds in extracts of *Tabebuia avellanedae* (Bignoniaceae). *J. Chromatogr. A*, 693 (1995) 281-287.

See also 1373, 1514, 2415, 2451.

## 9. OXO COMPOUNDS, ETHERS, EPOXIDES AND QUINONES

- 1559 Baj, S. and Dawid, M.: Correlation between the chemical structures of dialkyl peroxides and their retention in reversed-phase high-performance liquid chromatography. *J. Liq. Chromatogr.*, 17 (1994) 3933-3949.
- 1560 Nakashima, K., Hidaka, Y., Yoshida, T., Kuroda, N. and Akiyama, S.: High-performance liquid chromatographic determination of short-chain aliphatic aldehydes using 4-(N,N-dimethylaminosulphonyl)-7-hydrazino-2,1,3-benzoxadiazole as a fluorescence reagent. *J. Chromatogr. B*, 661 (1994) 205-210.
- 1561 Wei, D., Sun, A., Jing, X. and Geng, A.: (Analysis of the reaction solution of bromination and methoxylation of *p*-hydroxybenzaldehyde by high performance liquid chromatography). *Sepu*, 12 (1994) 371-372; C.A., 122 (1995) 22858a.
- 1562 Wen, Y.-H., Lin, S.-J., Wu, S.-S. and Wu, H.-L.: Trace analysis of acetaldehyde as fluorogenic derivative by high performance liquid chromatography. *Gaoxiong Yixue Kexue Zazhi*, 10 (1994) 295-300; C.A., 121 (1994) 275687s.
- 1563 Wu, R. and White, L.B.: Automated procedure for determination of trace amounts of aldehydes in drinking water. *J. Chromatogr. A*, 692 (1995) 1-9.

For additional information see C.A.:  
121 (1994) 308489j.

See also 1373, 1381, 1418, 1514, 1522, 1658, 1885, 2002, 2361, 2370, 2455, 2486.

## 10. CARBOHYDRATES

## 10a. Mono and oligosaccharides. Structural studies

- 1564 Abelian, V.H., Yamamoto, T. and Afrikan, E.G.: (On the action mechanism of cyclomaltoextrin glucanotransferases from alkalophilic, thermophilic and mesophilic microorganisms). *Biochimiya (Moscow)*, 59 (1994) 1122-1129.
- 1565 Adachi, S., Panintrarux, C., Araki, Y., Kimura, Y. and Matsuno, R.: Separation of alkyl  $\beta$ -D-glucosides and *n*-alcohols by using a porous trimethylolpropane trimethacrylate homopolymer gel. *Biosci., Biotechnol., Biochem.*, 58 (1994) 1558-1563; C.A., 121 (1994) 250082c.
- 1566 Andersen, D.C., Goochee, C.F., Cooper, G., Weitzhandler, M.: Monosaccharide and oligosaccharide analysis of isoelectric focusing-separated and blotted granulocyte colony-stimulating factor glycoforms using high-pH anion-exchange chromatography with pulsed amperometric detection. *Glycobiology*, 4 (1994) 459-467; C.A., 121 (1994) 225404b.
- 1567 Corradini, C., Corradini, D., Huber, C.G. and Bonn, G.K.: Synthesis of a polymeric-based stationary phase for carbohydrate separation by high-pH anion-exchange chromatography with pulsed amperometric detection. *J. Chromatogr. A*, 685 (1994) 213-220.
- 1568 Corradini, C., Cristalli, A. and Corradini, D.: Determination of carbohydrates in fruit-based beverages by high-performance anion-exchange chromatography with pulsed amperometric detection (HPAEC-PAD). *Ital. J. Food Sci.*, 6 (1994) 103-111; C.A., 121 (1994) 254042n.

- 1569 Dennis, M.J., Massey, R.C. and Bigwood, T.: Investigation of sorbitol content in wines and an assessment of its authenticity using stable isotope ratio mass spectrometry. *Analyst (Cambridge)*, 119 (1994) 2057-2060.
- 1570 Dona, A.-M. and Verchere, J.-F.: High-performance liquid chromatography of alditols with indirect photometric detection. *J. Chromatogr. A*, 689 (1995) 13-21.
- 1571 Herbreteau, B., Villete, V., Lafosse, M. and Dreux, M.: Analysis of oligosaccharides using aminobonded silica gel and ternary eluent with evaporative scattering detection. *Fresenius J. Anal. Chem.*, 351 (1995) 246-250.
- 1572 Lehtonen, P. and Hurme, R.: Liquid chromatographic determination of sugars in beer by evaporative light scattering detection. *J. Inst. Brew.*, 100 (1994) 343-346; C.A., 122 (1995) 8303j.
- 1573 Matsui, T., Mizuochi, T., Titani, K., Okinaga, T., Hoshi, M., Bousfield, G.R., Sugino, H. and Ward, D.N.: Structural analysis of N-linked oligosaccharides of equine chorionic gonadotropin and lutropin  $\beta$ -subunits. *Biochemistry*, 33 (1994) 14039-14048.
- 1574 Mattsson, P., Battchikova, N., Sippola, K. and Korpela, T.: The role of histidine residues in the catalytic act of cyclomaltoextrin glucanotransferase from *Bacillus circulans* var. *alkalophilus*. *Biochim. Biophys. Acta*, 1247 (1995) 97-103.
- 1575 Nishida, Y., Bai, C., Ohnui, H. and Meguro, H.: A highly sensitive method to identify the DL-configurations of monosaccharides based on (-)-TBMB carboxylic acid and HPLC. *J. Carbohydr. Chem.*, 13 (1994) 1003-1008; C.A., 121 (1994) 250078f.
- 1576 Pitkanen, E. and Kanninen, T.: Determination of mannose and fructose in human plasma using deuterium labelling and gas chromatography/mass spectrometry. *Biol. Mass Spectrom.*, 23 (1994) 590-595; C.A., 121 (1994) 225794d.
- 1577 Ruan, S. and Lloyd, K.O.: Analysis of glycosphingolipid-derived oligosaccharides by high pH anion exchange chromatography. *Glycoconjugate J.*, 11 (1994) 249-256; C.A., 122 (1995) 4614p.
- 1578 Senn, H., Lendenmann, U., Snozzi, M., Hamer, G. and Egli, T.: The growth of *Escherichia coli* in glucose-limited chemostat cultures: a re-examination of the kinetics. *Biochim. Biophys. Acta*, 1201 (1994) 424-436.
- 1579 Tsuji, T., Yamamoto, K. and Osawa, T.: Affinity chromatography of oligosaccharides and glycopeptides with immobilized lectins. *Mol. Interact. Biosep.*, (1993) 113-126; C.A., 121 (1994) 249852x - a review with many refs.
- 1580 Zuercher, U., Muehleemann, C. and Amado, R.: (Determination of the neutral sugar constituents of dietary fiber as N-p-methoxyphenyl-glycosylamines by HPLC with automated precolumn derivatization). *Mitt. Geb. Lebensmittelunters. Hyg.*, 85 (1994) 31-45; C.A., 121 (1994) 253991j.

For additional information see C.A.:  
121 (1994) 299329c.

See also 1420, 1432, 1590, 1592, 1606, 1981, 2484.

## 10b. Polysaccharides, mucopolysaccharides, lipopolysaccharides

- 1581 Altmann, F.: (Analysis of glycoprotein N-glycans in the form of fluorescent pyridylamino-oligosaccharides by 2D-HPLC. An example of glycoproteins of insect cells). *Lebensm.-Biotechnol.*, 11 (1994) 63-66; C.A., 121 (1994) 250047v.
- 1582 Duflot, P.: Preparation of hypocaloric soluble glucose polymers. *Eur. Pat. Appl.* EP 593,368 (Cl. C08B37/00), 20 Apr. 1994, FR Appl. 92/12,447, 16 Oct. 1992; 17 p.; C.A., 121 (1994) 303414w.
- 1583 Hermann, K., Frank, G. and Ring, J.: Contamination of heparin by histamine: measurement and characterization by high-performance liquid chromatography and radioimmunoassay. *Allergy (Copenhagen)*, 49 (1994) 569-572; C.A., 121 (1994) 286698x.
- 1584 Huber, A. and Praznik, W.: Characterization of branching characteristics of starch-glucans by means of combined application of complexation, enzymatically catalyzed modification, and liquid-chromatography. *J. Liq. Chromatogr.*, 17 (1994) 4031-4056.
- 1585 Kennedy, J.F., Melo, E.H.M., Lloyd, L.L., Warner, F.P. and Jumel, K.: Gel permeation and gel filtration chromatography in the analysis of cellulose derivatives. In: Kennedy, J.F., Phillips, G.O. and Williams, P.A. (Editors), *Ligno-Cellul.* 1990, Horwood, Chichester, 1992, pp. 529-539; C.A., 121 (1994) 258158c.
- 1586 Nowakowski, R., Cardot, P.J.P., Coleman, A.W., Billard, E. and Guiochon, G.: Elution mechanisms of cyclodextrins in reversed phase chromatography. *Anal. Chem.*, 67 (1995) 259-266.
- 1587 Ronkko, R., Pennanen, T., Smolander, A., Kitunen, V., Kortemaa, H. and Haahtela, K.: Quantification of *Frankia* strains and other root-associated bacteria in pure cultures and in the rhizosphere of axenic seedlings by high-performance liquid chromatography-based muramic acid assay. *Appl. Environ. Microbiol.*, 60 (1994) 3672-3678; C.A., 121 (1994) 250081b.
- 1588 Shah, G. and Dubin, P.L.: Adsorptive interaction of Ficoll standards with porous glass size-exclusion chromatography columns. *J. Chromatogr. A*, 693 (1995) 197-203.
- 1589 Shimada, E. and Nakamura, K.T.: Determination of Stokes radii and molecular masses of sodium hyaluronates by Sephacryl gel chromatography. *J. Chromatogr. A*, 685 (1994) 172-177.
- 1590 Thomas, J. and Mort, A.J.: Continuous postcolumn detection of underivatized polysaccharides in high-performance liquid chromatography by reaction with permanganate. *Anal. Biochem.*, 223 (1994) 99-104.
- 1591 Vilim, V. and Fosang, A.J.: Proteoglycans isolated from dissociative extracts of differently aged human articular cartilage: characterization of naturally occurring hyaluronan-binding fragments of aggrecan. *Biochem. J.*, 304 (1994) 887-894.

For additional information see C.A.:  
121 (1994) 212862z, 213081f.

## 10c. Glycoproteins and their constituents

- 1592 Almeida, I.C., Ferguson, M.A.J., Schenkman, S. and Travassos, L.R.: Lytic anti- $\alpha$ -galactosyl antibodies from patients with chronic Chagas disease recognize novel O-linked oligosaccharides on mucin-like glycosylphosphatidylinositol-anchored glycoproteins of *Trypanosoma cruzi*. *Biochem. J.*, 304 (1994) 793-802.

- 1593 Goso, Y. and Hotta, K.: Dot blot analysis of rat gastric mucin using histochemical staining methods. *Anal. Biochem.*, 223 (1994) 274-279.
- 1594 Gulino, D., Martinez, P., Delachanal, E., Concord, E., Duperray, A., Alemany, M. and Marguerie, G.: Expression and purification of a soluble functional form of the platelet  $\alpha$ IIb $\beta$ 3 integrin. *Eur. J. Biochem.*, 227 (1995) 108-115.
- 1595 Ivanov, A.E., Zhigis, L.S., Rapoport, E.M., Lisyutina, O.E. and Zubov, V.P.: Characterization of weak hydrophobic composite sorbents and their application to the isolation of bacterial lectin. *J. Chromatogr. B*, 664 (1995) 219-223.
- 1596 Matsui, T., Ozeki, Y., Suzuki, M., Hino, A. and Titani, K.: Purification and characterization of two  $\text{Ca}^{2+}$ -dependent lectins from coelomic plasma of sea cucumber, *Stichopus japonicus*. *J. Biochem. (Tokyo)*, 116 (1994) 1127-1133.
- 1597 Nikitina, V.E., Alenkina, S.A., Italyanskaya, Yu.V. and Ponomareva, E.G.: (Purification and comparison of lectins from the cell surface of active and inactive in hemagglutination azo-spirilla). *Biokhimiya (Moscow)*, 59 (1994) 656-662.
- 1598 Saito, T., Nakamura, T., Kabuki, T., Kitazawa, H. and Itoh, T.: Fractionation of glycopeptides from bovine  $\kappa$ -casein and hen ovomucin via lysozyme affinity chromatography. *Anim. Sci. Technol.*, 65 (1994) 624-630; C.A., 121 (1994) 250075c.
- 1599 Smith, K.D., Elliott, M.A., Elliott, H.G., McLaughlin, C.M., Wightman, P. and Wood, G.C.: Heterogeneity of  $\alpha_1$ -acid glycoprotein in rheumatoid arthritis. *J. Chromatogr. B*, 661 (1994) 7-14.
- 1600 Van de Lest, C.H.A., Versteeg, E.M.M., Veerkamp, J.H. and van Kuppevelt, T.H.: A spectrophotometric method for the determination of heparan sulfate. *Biochim. Biophys. Acta*, 1201 (1994) 305-311.
- 1601 Zenteno, E., Ortega, M., Qin, Z., Montrueil, J., Debray, H.: Fast purification of *Phaseolus vulgaris* isolectins. *Prep. Biochem.*, 24 (1994) 175-183; C.A., 121 (1994) 275704v.

For additional information see C.A.:  
121 (1994) 222989k, 276046u, 296287b.

See also 1339, 1340, 1579, 1581, 1875, 2048.

## 11. ORGANIC ACIDS AND LIPIDS

## 11a. Organic acids and simple esters

- 1602 Adlof, R.O. and Emken, E.A.: Silver ion high-performance liquid chromatographic separation of fatty acid methyl esters labelled with deuterium atoms on the double bonds. *J. Chromatogr. A*, 685 (1994) 178-181.
- 1603 Bernwieser, I., Paetzold, R., Galensa, R. and Sontag, G.: (HPLC with coulometric electrode array detection. Determination of 1-O-trans-p-coumaroylglycerol in sorghum, maize, and beer). *Z. Lebensm.-Unters. Forsch.*, 198 (1994) 40-43; C.A., 121 (1994) 299391s.
- 1604 Cano, M.P., Torija, E., Marin, M.A. and Camara, M.: A simple ion-exchange chromatographic determination of non-volatile organic acids in some Spanish exotic fruits. *Z. Lebensm.-Unters. Forsch.*, 199 (1994) 214-218; C.A., 122 (1995) 8296j.

- 1605 Chase, G.W., Jr., Akoh, C.C. and Eitenmiller, R.R.: Evaporative light-scattering mass detection for high-performance liquid chromatographic analysis of sucrose polyester blends in cooking oils. *J. Am. Oil Chem. Soc.*, 71 (1994) 1273-1276.
- 1606 Chong, R.W. and Moore, B.J.: Determination of citrate, inositol and gentisic acid in a pharmaceutical diagnostic formulation by ion-moderated partition chromatography. *J. Chromatogr. A*, 692 (1995) 203-205.
- 1607 Czajkowska, T., Hrabovsky, I., Buszewski, B., Gilpin, R.K. and Jaroniec, M.: Comparison of the retention of organic acids on alkyl and alkylamide chemically bonded phases. *J. Chromatogr. A*, 691 (1995) 217-224.
- 1608 Daoud, H.G., Biacs, P.A., Dakar, M.A. and Hajdu, F.: Ion-pair chromatography and photodiode-array detection of vitamin C and organic acids. *J. Chromatogr. Sci.*, 32 (1994) 481-487.
- 1609 De Swaef, S.I., de Beer, J.O. and Vlietinck, A.J.: Quantitative determination of *p*-coumaric acid in *Echinacea purpurea* press juice and urgenin. A validated method. *J. Liq. Chromatogr.*, 17 (1994) 4169-4183.
- 1610 Denekamp, C., Mandelbaum, A., Weisz, A. and Ito, Y.: Preparative separation of stereoisomeric 1-methyl-4-methoxymethylcyclohexanecarboxylic acids by pH-zone-refining counter-current chromatography. *J. Chromatogr. A*, 685 (1994) 253-257.
- 1611 Gübitz, G. and Mihellyes, S.: Resolution of 2-hydroxydicarboxylic acid enantiomers by ligand exchange chromatography on chemically bonded chiral phases. *J. High Resolut. Chromatogr.*, 17 (1994) 733-734.
- 1612 Hagen, S.R. and Thompson, J.D.: Analysis of mycolic acids by high-performance liquid chromatography and fluorimetric detection. Implications for the identification of mycobacteria in clinical samples. *J. Chromatogr. A*, 692 (1995) 167-172.
- 1613 Iwata, T., Hirose, T., Nakamura, M. and Yamaguchi, M.: 4-(1-Methylphenanthro[9,10-d]imidazol-2-yl)-benzohydrazide as derivatization reagent for carboxylic acids in high-performance liquid chromatography with conventional and laser-induced fluorescence detection. *Analyst (Cambridge)*, 119 (1994) 1747-1751.
- 1614 Jackson, P.E.: Analysis of oxalate in Bayer liquors: a comparison of ion chromatography and capillary electrophoresis. *J. Chromatogr. A*, 693 (1995) 155-161.
- 1615 Lagoutte, D., Lombard, G., Nisseron, S., Papet, M.P. and Saint-Jalm, Y.: Determination of organic acids in cigarette smoke by high-performance liquid chromatography and capillary electrophoresis. *J. Chromatogr. A*, 684 (1994) 251-257.
- 1616 Lange, M. and Mályusz, M.: Fast method for the simultaneous determination of 2-oxo acids in biological fluids by high-performance liquid chromatography. *J. Chromatogr. B*, 662 (1994) 97-102.
- 1617 Lee, B.L., Ong, H.Y., Koh, D. and Ong, C.N.: High-performance liquid chromatographic method for determination of dehydroabietic and abietic acids, the skin sensitizers in bindi adhesive. *J. Chromatogr. A*, 685 (1994) 263-269.
- 1618 Lian, H.-Z., Mao, L. and Miao, J.: Determination of hexadecanoic and octadecanoic acids in stearin for industrial use by reversed-phase ion suppression high-performance liquid chromatography. *J. Liq. Chromatogr.*, 17 (1994) 4081-4086.
- 1619 MacMillan, D.K., Hill, E., Sala, A., Sigal, E., Shuman, T., Henson, P.M. and Murphy, R.C.: Eosinophil 15-lipoxygenase is a leukotriene A<sub>4</sub> synthase. *J. Biol. Chem.*, 269 (1994) 26663-26668.
- 1620 Nikolova-Damyanova, B., Christie, W.W. and Herslöf, B.: Silver ion high-performance liquid chromatography of esters of isomeric octadecenoic fatty acids with short-chain monounsaturated alcohols. *J. Chromatogr. A*, 693 (1995) 235-239.
- 1621 Notsumoto, S., Kobayashi, K., Hirauchi, K., Igimi, H. and Uchida, K.: (Determination of vanilmandelic acid in urine by high-performance liquid chromatography with ultraviolet detection). *Rinsho Kagaku (Nippon Rinsho Kagakkai)*, 23 (1994) 243-247; *C.A.*, 121 (1994) 272379g.
- 1622 O'Flaherty, J.T., Cordes, J.F., Lee, S.L., Samuel, M. and Thomas, M.J.: Chemical and biological characterization of oxo-eicosatetraenoic acids. *Biochim. Biophys. Acta*, 1201 (1994) 505-515.
- 1623 Oi, N., Kitahara, H., Aoki, F. and Kisu, N.: Direct separation of carboxylic acid enantiomers by high-performance liquid chromatography with amide and urea derivatives bonded to silica gel as chiral stationary phases. *J. Chromatogr. A*, 689 (1995) 195-201.
- 1624 Patthy, M., Király, I. and Sziráki, I.: Separation of dihydroxybenzoates, indicators of *in vivo* hydroxyl free radical formation, in the presence of transmitter amines and some metabolites in rodent brain, using high-performance liquid chromatography with electrochemical detection. *J. Chromatogr. B*, 664 (1995) 247-252.
- 1625 Pourfarzam, M., Schaefer, J., Turnbull, D.M. and Bartlett, K.: Analysis of fatty acid oxidation intermediates in cultured fibroblasts to detect mitochondrial oxidation disorders. *Clin. Chem. (Washington)*, 40 (1994) 2267-2275.
- 1626 Powell, W.S., Gravelle, F. and Gravel, S.: Phorbol myristate acetate stimulates the formation of 5-oxo-6,8,11,14-eicosatetraenoic acid by human neutrophils by activating NADPH oxidase. *J. Biol. Chem.*, 269 (1994) 25373-25380.
- 1627 Reid, R.H.P.: Studies on the retention behaviour of a group of organic anions of biochemical interest on quaternary bonded silica columns equilibrated with a functionally coherent series of counterions. Use of 2-(N-morpholino)ethanesulphonate as a counterion and N-tris(hydroxymethyl)methyl-3-aminopropane-sulphonate as an eluent. *J. Chromatogr. A*, 684 (1994) 221-234.
- 1628 Reynaud, D., Demin, P. and Pace-Asciak, C.R.: Hepoxilin A<sub>3</sub> formation in the rat pineal gland selectively utilizes (12S)-hydroperoxyeicosatetraenoic acid (HPETE), but not (12R)-HPETE. *J. Biol. Chem.*, 269 (1994) 23976-23980.
- 1629 Schnurr, K., Kühn, H., Rapoport, S.M. and Schewe, T.: 3,5-Di-*t*-butyl-4-hydroxytoluene (BHT) and probucol stimulate selectively the reaction of mammalian 15-lipoxygenase with biomembranes. *Biochim. Biophys. Acta*, 1254 (1995) 66-72.
- 1630 Takagi, T.: Resolution of vic-dihydroxy acid diastereomers to four enantiomers by high-performance liquid chromatography. *Am. Oil Chem. Soc.*, 71 (1994) 547-548; *C.A.*, 121 (1994) 169294b.
- 1631 Toyoshima, S., Saido, H., Watanabe, F., Miyatake, K. and Nakano, Y.: Assay for urinary methylmalonic acid by high-pressure liquid chromatography. *Biosci., Biotechnol., Biochem.*, 58 (1994) 1882-1883; *C.A.*, 121 (1994) 296320g.
- 1632 Ueda, N., Yamamoto, K., Yamamoto, S., Tokunaga, T., Shirakawa, E., Shinkai, H., Ogawa, M., Sato, T., Kudo, I., Inoue, K., Takizawa, H., Nagano, T. et al.: Lipoxygenase-catalyzed oxygenation of arachidonylethanolamide, a cannabinoid receptor agonist. *Biochim. Biophys. Acta*, 1254 (1995) 127-134.

- 1633 Weimar, A. and Stuurman, H.W.: (HPLC as analysis method for quality control of food. Determination of organic acids and the preservatives benzoic and sorbic acid in beverages and other foods). *GIT Fachz. Lab.*, 37 (1993) 652-654; *C.A.*, 121 (1994) 253981f.
- 1634 Yamane, M., Abe, A. and Nakajima, M.: High-performance liquid chromatography-thermospray mass spectrometry of  $\omega$ -hydroxy polyunsaturated fatty acids from rat brain homogenate. *J. Chromatogr. B*, 662 (1994) 91-96.
- 1635 Yurawecz, M.P., Hood, J.K., Roach, J.A.G., Mossoba, M.M., Daniels, D.H., Ku, Y., Pariza, M.W. and Chin, S.F.: Conversion of allylic hydroxy oleate to conjugated linoleic acid and methoxy oleate by acid-catalyzed methylation procedures. *J. Am. Oil Chem. Soc.*, 71 (1994) 1149-1155.
- For additional information see *C.A.*:  
121 (1994) 230483f, 253892c;  
122 (1995) 4610j.
- See also 1373, 1381, 1442, 1504, 1647, 1724, 2012, 2232, 2280.
- 11b. Prostaglandins**
- 1636 Hardcastle, J.E., He, M., Begum, B. and Vermillion-Salsbury, R.: Use of secondary chemical equilibria in liquid chromatography to determine dissociation constants of leukotriene B<sub>4</sub> and prostaglandin B<sub>2</sub>. *J. Chromatogr. A*, 691 (1995) 225-229.
- 1637 Lloyd-Evans, P., Barrow, S.E., Hill, D.J., Bowden, L.A., Rainer, G.E., Knight, J. and Rowley, A.F.: Eicosanoid generation and effects on the aggregation of thrombocytes from the rainbow trout, *Oncorhynchus mykiss*. *Biochim. Biophys. Acta*, 1215 (1994) 291-299.
- 1638 Rosenfeld, J.M. and Fang, X.: Simultaneous sorption and analytical derivatization on a polystyrene-divinylbenzene polymer. Preparation of chromophoric and fluorophoric derivatives of the prostaglandins. *J. Chromatogr. A*, 691 (1995) 231-237.
- See also 1993.
- 11c. Lipids and their constituents**
- 1639 Bartle, K.D. and Clifford, T.A.: Supercritical fluid extraction and chromatography of lipids and related compounds. *Adv. Appl. Lipid Res.*, 1 (1992) 217-264; *C.A.*, 121 (1994) 249949j - a review with 141 refs.
- 1640 Blank, M.L. and Snyder, F.L.: Chromatographic analysis of ether-linked glycerolipids, including platelet-activating factor and related cell mediators. *Chromatogr. Sci. Ser.*, 65(Lipid Chromatographic Analysis) (1994) 291-316; *C.A.*, 121 (1994) 249909w - a review with 118 refs.
- 1641 Borgeat, P., Picard, S., Braquet, P., Allen, M. and Shushan, B.: LC-MS-MS with ion spray: a promising approach for analysis of underivatized platelet activating factor (PAF). *J. Lipid Mediators Cell Signalling*, 10 (1994) 11-12; *C.A.*, 121 (1994) 250140v.
- 1642 Bruch, J., Gono, E., Malkusch, W. and Rehn, B.: Improved method for quantitative analysis of lung surfactant phospholipids in bronchoalveolar lavage fluids by high-performance liquid chromatography. *Clin. Chim. Acta*, 231 (1994) 193-204.
- 1643 Brütting, R. and Spiteller, G.: Produkte der Dimerisierung ungesättigter Fettsäuren XII: Die Dimerisierung von Konjunsäure. *Fat. Sci. Technol.*, 96 (1994) 445-451.
- 1644 Chang, M.-K., Konkerton, E.J., Chapital, D. and Wan, P.J.: Behavior of diglycerides and conjugated fatty acid triglycerides in reverse-phase chromatography. *J. Am. Oil Chem. Soc.*, 71 (1994) 1173-1175.
- 1645 Damiani, P., Santinelli, F., Simonetti, M.S., Castellini, M. and Rosi, M.: Comparison between two procedures for stereospecific analysis of triacylglycerols from vegetable oils - I: olive oil. *J. Am. Oil Chem. Soc.*, 71 (1994) 1157-1162.
- 1646 Hansen, S.L., Myers, M.R. and Artz, W.E.: Nonvolatile components produced in triolein during deep-fat frying. *J. Am. Oil Chem. Soc.*, 71 (1994) 1239-1243.
- 1647 Hayakawa, M., Sugiyama, S. and Ozawa, T.: HPLC analysis of lipids: analysis of fatty acids and their derivatives by a microcolumn HPLC system. *Chromatogr. Sci. Ser.*, 65(Lipid Chromatographic Analysis) (1994) 273-290; *C.A.*, 121 (1994) 249908v - a review with 23 refs.
- 1648 Hopia, A.: Application of high-performance size exclusion chromatography (HPSEC) when analyzing the quality of edible oils. In: Maikki, Y. and Lambertsen, G. (Editors), *Nord. Lipid Symp., Proc.*, 17th, Scand. Forum Lipid Res. Technol., Bergen, 1993, pp. 222-225; *C.A.*, 121 (1994) 279157s.
- 1649 Hori, M., Sahashi, Y., Koike, S., Yamaoka, R. and Sato, M.: Molecular species analysis of polyunsaturated fish triacylglycerol by high-performance liquid chromatography/fast atom bombardment mass spectrometry. *Anal. Sci.*, 10 (1994) 719-724; *C.A.*, 122 (1995) 8306n.
- 1650 Hussain, A.A., Jona, J.A., Yamada, A. and Dittert, L.W.: Chloramine-T in radiolabeling techniques. II. A nondestructive method for radiolabeling biomolecules by halogenation. *Anal. Biochem.*, 224 (1995) 221-226.
- 1651 Kukis, A.: Modern trends in the analysis of fats and oils. In: Maikki, Y. and Lambertsen, G. (Editors), *Nord. Lipid Symp., Proc.*, 17th, Scand. Forum Lipid Res. Technol., Bergen, 1993, pp. 188-195; *C.A.*, 121 (1994) 279107a - a review with 57 refs.
- 1652 Mahungu, S.M., Hansen, S.L. and Artz, W.E.: Quantitation of volatile compounds in heated trilinolein by static headspace-capillary gas chromatography/infrared spectroscopy - mass spectrometry. *J. Am. Oil Chem. Soc.*, 71 (1994) 1169-1171.
- 1653 Miyazawa, T., Fujimoto, K., Suzuki, T. and Yasuda, K.: Determination of phospholipid hydroperoxides using luminol chemiluminescence-high-performance liquid chromatography. *Methods Enzymol.*, 233(Oxygen Radicals in Biological Systems, Pt. C) (1994) 324-332; *C.A.*, 121 (1994) 225360j.
- 1654 Neff, W.E., Adlof, R.O., List, G.R. and El-Agaimy, M.: Analyses of vegetable oil triacylglycerols by silver ion high performance liquid chromatography with flame ionization detection. *J. Liq. Chromatogr.*, 17 (1994) 3951-3968.
- 1655 Neff, W.E., El-Agaimy, M.A. and Mounts, T.L.: Oxidative stability of blends and interesterified blends of soybean oil and palm olein. *J. Am. Oil Chem. Soc.*, 71 (1994) 1111-1116.
- 1656 Neff, W.E., Mounts, T.L., Rinsch, W.M., Konishi, H. and El-Agaimy, M.A.: Oxidative stability of purified canola oil triacylglycerols with altered fatty acid compositions as affected by triacylglycerol composition and structure. *J. Am. Oil Chem. Soc.*, 71 (1994) 1101-1109.

- 1657 Nilsson, Å., Chen, Q. and Dahlman, E.: Metabolism of chylomicron phosphatidylinositol in the rat: fate *in vivo* and hydrolysis with lipoprotein lipase and hepatic lipase *in vitro*. *J. Lipid Res.*, 35 (1994) 2151-2160.
- 1658 Piironen, V. and Kankare, M.: Analysis of lipid hydroperoxides using high-performance liquid chromatography with chemiluminescence detection. In: Maikki, Y. and Lambertsen, G. (Editors), *Nord. Lipid Symp., Proc., 17th*, Scand. Forum Lipid Res. Technol., Bergen, 1993, pp. 226-229; C.A., 121 (1994) 279158t.
- 1659 Schenk, T., Schuphan, I. and Schmidt, B.: High-performance liquid chromatographic determination of the rhamnolipids produced by *Pseudomonas aeruginosa*. *J. Chromatogr. A*, 693 (1995) 7-13.
- 1660 Shibamoto, T.: *Lipid Chromatographic Analysis*. Dekker, New York, 1994, 412 pp.; C.A., 121 (1994) 225828t.
- 1661 Smith, K.W., Perkins, J.M., Jeffrey, B.S.J. and Phillips, D.L.: Separation of molecular species of *cis* and *trans*-triacylglycerols in *trans*-hardened confectionery fats by silver ion high-performance liquid chromatography. *J. Am. Oil Chem. Soc.*, 71 (1994) 1219-1222.
- 1662 Swe, P.Z., Che Man, Y.B., Ghazali, H.M. and Wei, L.S.: Identification of major triglycerides causing the clouding of palm olein. *J. Am. Oil Chem. Soc.*, 71 (1994) 1141-1144.
- 1663 Weete, J.D.: Preferential degradation of noncholine phosphatides in soybean lecithin by thermalization. *J. Am. Oil Chem. Soc.*, 71 (1994) 1195-1199.
- 1664 Zhang, F., Koseoglu, S.S. and Rhee, K.C.: Effects of expander process on the phospholipids in soybean oil. *J. Am. Oil Chem. Soc.*, 71 (1994) 1145-1148.
- See also 1577, 1672, 1693, 1700, 1702, 2142, 2143, 2399, 2482, 2493.
- 11d. *Lipoproteins and their constituents*
- 1665 Maeba, R., Shimasaki, H. and Ueta, N.: Conformational changes in oxidized LDL recognized by mouse peritoneal macrophages. *Biochim. Biophys. Acta*, 1215 (1994) 79-86.
- 1666 Pursiainen, M., Jauhiainen, M. and Ehnholm, C.: Low-density lipoprotein activates the protease region of recombinant apo(a). *Biochim. Biophys. Acta*, 1215 (1994) 170-175.
- 1667 Sattler, W., Mohr, D. and Stocker, R.: Rapid isolation of lipoproteins and assessment of their peroxidation by high-performance liquid chromatography postcolumn chemiluminescence. *Methods Enzymol.*, 233(Oxygen Radicals in Biological Systems, Pt. C) (1994) 469-489; C.A., 121 (1994) 225365q.
- 1668 Shibusawa, Y., Mugiyama, M., Matsumoto, U. and Ito, Y.: Complementary use of counter-current chromatography and hydroxyapatite chromatography for the separation of three main classes of lipoproteins from human serum. *J. Chromatogr. B*, 664 (1995) 295-301.
- 1669 Verges, B., Rader, D., Schaefer, J., Zech, L., Kindt, M., Fariwell, T., Gambert, P. and Breser, H.B., Jr.: *In vivo* metabolism of apolipoprotein A-IV in severe hypertriglyceridemia: a combined radiotracer and stable isotope kinetic study. *J. Lipid Res.*, 35 (1994) 2280-2291.
- 1670 Wiegel, D., Meyer, C. and Arnold, K.: Partitioning of high-density lipoproteins in charge-sensitive two-phase systems. *J. Chromatogr. B*, 661 (1994) 159-164.
- 1671 Yamamoto, Y.: Chemiluminescence-based high-performance liquid chromatography assay of lipid hydroperoxides. *Methods Enzymol.*, 233(Oxygen Radicals in Biological Systems, Pt. C) (1994) 319-324; C.A., 121 (1994) 225359r.
- For additional information see C.A.:  
121 (1994) 296610b.
- See also 1694.
12. ORGANIC PEROXIDES
- 1672 Thomas, J.P., Kalyanaraman, B. and Girotti, A.W.: Involvement of preexisting lipid hydroperoxides in Cu<sup>2+</sup>-stimulated oxidation of low-density lipoprotein. *Arch. Biochem. Biophys.*, 315 (1994) 244-254.
- See also 1635, 1653, 1656, 1658, 1671.
13. STEROIDS
- 13a. *General techniques*
- 1673 Boschi, S., de lasio, R., Mesini, P., Bolelli, G.F., Sciajno, R., Pasquali, R. and Capelli, M.: Measurement of steroid hormones in plasma by isocratic high performance liquid chromatography coupled to radioimmunoassay. *Clin. Chim. Acta*, 231 (1994) 107-113.
- 1674 Hooijerink, D., Schilt, R., van Bennekom, E. and Brouwer, B.: Determination of anabolic esters in oily formulation and plasma in husbandry using high-performance liquid chromatography and gas chromatography-mass selective detection. *Analyst (Cambridge)*, 119 (1994) 2617-2622.
- 1675 Lensmeyer, G.L., Onsager, C., Carlson, I.H. and Wiebe, D.A.: Use of particle-loaded membranes to extract steroids for high-performance liquid chromatographic analyses. Improved analyte stability and detection. *J. Chromatogr. A*, 691 (1995) 239-246.
- 13b. *Pregnane and androstane derivatives*
- 1676 Dolezalova, M.: Routine high-performance liquid chromatographic determination of urinary unconjugated cortisol using solid-phase extraction and ultraviolet detection. *Clin. Chim. Acta*, 231 (1994) 129-137.
- 1677 El-Dawy, M.A., Habeeb, A.A., Mabrouk, M.M. and Bastawissy, I.A.E.: HPLC assay of lidocaine hydrochloride and prednisolone acetate in anti-hemorrhoidal preparations. *Egypt. J. Pharm. Sci.* 1993, 34 (1994) 683-690; C.A., 122 (1995) 64547d.
- 1678 Hansen-Møller, J.: Rapid high-performance liquid chromatographic method for simultaneous determination of androstenedione, skatole and indole in back fat from pigs. *J. Chromatogr. B*, 661 (1994) 219-230.

- 1679 Inoue, S., Inokuma, M., Harada, T., Shibutani, Y., Yoshitake, T., Charles, B., Ishida, J. and Yamaguchi, M.: Simultaneous high-performance liquid chromatographic determination of 6 $\beta$ -hydroxycortisol and cortisol in urine with fluorescence detection and its application for estimating hepatic drug-metabolizing enzyme induction. *J. Chromatogr. B*, 661 (1994) 15-23.
- 1680 Los, L.E., Pitzemberger, S.M., Ramjit, H.G., Coddington, A.B. and Colby, H.D.: Hepatic metabolism of spironolactone. Production of 3-hydroxy-thiomethyl metabolites. *Drug Metab. Disp.*, 22 (1994) 903-908.
- 1681 Prein, A., Mehnert, W. and Frömming, K.-H.: Penetration of hydrocortisone into excised human skin under the influence of cyclodextrins. *Pharmazie*, 50 (1995) 121-126.
- 1682 Valenta, C. and Schmatzberger-Wagerer, M.: Stabilitätsuntersuchungen von Progesteron-Hydrogelen. *Pharmazie*, 50 (1995) 69-70.
- 1683 Wong, Y.N., Chien, B.M. and D'mello, A.P.: Analysis of corticosterone in rat plasma by high-performance liquid chromatography. *J. Chromatogr. B*, 661 (1994) 211-218.

### 13c. Estrogens

- 1684 Guo, L. and Ma, S.: (Determination of urinary estriol by high performance liquid chromatography with fluorescence detection). *Zhonghua Yixue Jianyan Zazhi*, 17 (1994) 158-160; C.A., 121 (1994) 292940m.
- 1685 Paris, A., Dolo, L., Debrauwer, L., Rao, D. and Terqui, M.: Analysis of [ $^3$ H]estradiol-17 $\beta$  metabolites in calf perirenal fat. *Analyst (Cambridge)*, 119 (1994) 2623-2626.
- 1686 Smales, C.M., Cooke, D. and Blackwell, L.F.: Use of ion-exchange and hydrophobic-interaction chromatography for the rapid purification of lysozyme-estrone glucuronide conjugates. *J. Chromatogr. B*, 662 (1994) 3-14.

For additional information see C.A.:  
121 (1994) 213109w.

### 13d. Sterols

- 1687 Ansari, G.A.S. and Smith, L.L.: Assay of cholesterol autooxidation. *Methods Enzymol.*, 233(Oxygen Radicals in Biological Systems, Pt. C) (1994) 332-338; C.A., 121 (1994) 225361k.
- 1688 Bragagnolo, N., Rodríguez-Amaya, D.B.: (Evaluation of the cholesterol concentration as a quality control parameter for egg noodles). *Rev. Inst. Adolfo Lutz*, 53 (1993) 21-26; C.A., 121 (1994) 299406a.
- 1689 Caboni, M.F., Gallina, T.T., Lercker, G. and Capella, P.: Analytical problems of oxisterols determination in lipid-containing foods. *Riv. Ital. Sostanze Grasse*, 71 (1994) 243-246; C.A., 121 (1994) 299599r.
- 1690 Gu, Y.-F., Chen, Y. and Hammond, E.G.: Use of cyclic anhydrides to remove cholesterol and other hydroxy compounds from fats and oils. *J. Am. Oil Chem. Soc.*, 71 (1994) 1205-1209.
- 1691 Iorizzi, M., de Riccardis, F., Minale, L., Palgiano, E., Riccio, R., Debitus, C. and Duhet, D.: Polyoxygenated marine steroids from the deep water starfish *Styracaster caroli*. *J. Natural Prod.*, 57 (1994) 1361-1373.

- 1692 Kermasha, S., Kubow, S. and Goetghebeur, M.: Comparative high-performance liquid chromatographic analyses of cholesterol and its oxidation products using diode-array ultraviolet and laser light-scattering detection. *J. Chromatogr. A*, 685 (1994) 229-235.
- 1693 Nota, G., Naviglio, D., Romano, R., Spagna, M.S. and Improta, C.: (HPLC determination of trienanthoin,  $\beta$ -sitosterol, stigmasterol and capsaicin as anhydrous butter denaturing agents). *Riv. Ital. Sostanze Grasse*, 71 (1994) 197-202; C.A., 122 (1995) 8273z.
- 1694 Okazaki, M., Sasamoto, K., Muramatsu, T. and Hosaki, S.: (Binding capacities of immunoassay reagents for remnant-like particles to LDL and HDL by the improved HPLC method). *Rinsho Kagaku (Nippon Rinsho Kagakkai)*, 23 (1994) 236-242; C.A., 121 (1994) 275697v.
- 1695 Sevanian, A., Seraglia, R., Traldi, P., Rossato, P., Ursini, F. and Hodis, H.: Analysis of plasma cholesterol oxidation products using gas- and high-performance liquid chromatography/mass spectrometry. *Free Radical Biol. Med.*, 17 (1994) 397-409; C.A., 121 (1994) 250594w.
- 1696 Teng, J.I. and Smith, L.L.: High-performance liquid chromatographic analysis of human erythrocyte oxysterols as  $\Delta^4$ -3-ketone derivatives. *J. Chromatogr. A*, 691 (1995) 247-254.

See also 1522, 1990.

### 13e. Bile acids and alcohols

- 1697 Chaplin, M.F.: Analysis of bile acids and their conjugates using high-pH anion-exchange chromatography with pulsed amperometric detection. *J. Chromatogr. B*, 664 (1995) 431-434.
- 1698 Roda, A., Pellicciari, R., Cerre, C., Polimeni, C., Sadeghpour, B., Marinozzi, M., Forti, G.C. and Sapigni, E.: New 6-substituted bile acids: physico-chemical and biological properties of 6 $\alpha$ -methyl ursodeoxycholic acid and 6 $\alpha$ -methyl-7-epicholic acid. *J. Lipid Res.*, 35 (1994) 2268-2279.

## 14. STEROID GLYCOSIDES AND SAPONINS

For additional information see C.A.:  
122 (1995) 38963s.

See also 2448, 2450, 2455.

## 15. TERPENES AND OTHER VOLATILE AROMATIC COMPOUNDS

### 15a. Terpenes

- 1699 Ando, M., Arai, K., Kikuchi, K. and Isogai, K.: Synthetic studies of sesquiterpenes with a cis-fused decalin system. 4. Synthesis of (+)-5 $\beta$ H-eudesma-3,11-diene, (-)-5 $\beta$ H-eudesmane-4 $\beta$ ,11-diol, and (+)-5 $\beta$ H-eudesmane-4 $\alpha$ ,11-diol, and structure revision of a natural eudesmane-4,11-diol isolated from *Pluchea arguta*. *J. Natural Prod.*, 57 (1994) 1189-1199.

- 1700 Moreau, R.A., Powell, M.J., Osman, S.F., Whitaker, B.D., Fett, W.F., Roth, L. and O'Brien, D.J.: Analysis of intact hopanoids and other lipids from the bacterium *Zymomonas mobilis* by high-performance liquid chromatography. *Anal. Biochem.*, 224 (1995) 293-301.
- 1701 O'Mathona, D.P. and Dосkotch, R.W.: Five new labdane diterpenes from *Amphiachyris amoena*. *J. Natural Prod.*, 57 (1994) 1382-1390.
- 1702 Roth, L., Moreau, R.A., Powell, M.J. and O'Brien, D.J.: Semi-preparative separation of intact hopanoids from *Zymomonas mobilis*. *Anal. Biochem.*, 224 (1995) 302-308.
- 1703 Stadler, M., Anke, H. and Sterner, O.: New nematocidal and antimicrobial compounds from the basidiomycete *Cheimomophyllum candidissimum* (Berch & Curt.) Sing. *J. Antibiot.*, 47 (1994) 1284-1289.
- 1704 Yoshikawa, M., Yamaguchi, S., Chatani, N., Nishino, Y., Matsuo, T., Yamahara, J., Murakami, N., Matsuda, H. and Kubo, M.: (Crude drugs from aquatic plants. III. Quantitative analysis of triterpene constituents in *Alismatis rhizoma* by means of high performance liquid chromatography on the chemical change of the constituents during *Alismatis rhizoma* processing) *Yakugaku Zasshi*, 114 (1994) 241-247; *C.A.*, 121 (1994) 65665g.

See also 2154, 2393.

#### 15b. Essential oils

For additional information see *C.A.*:  
121 (1994) 213090h.

#### 15c. Bitter substances

- 1705 Buckee, G.K.: Determination of iso-alpha-, alpha- and beta-acids in hop and isomerized hop extracts. A collaborative trial. *Cerevisia Biotechnol.*, 19 (1994) 46-49; *C.A.*, 121 (1994) 229122f.
- 1706 Szucs, R., Vindevogel, J., Everaert, E., de Cooman, L., Sandra, P. and de Keukeleire, D.: Separation and quantification of all main hop acids in different hop cultivars by microemulsion electrokinetic chromatography. *J. Inst. Brew.*, 100 (1994) 293-296; *C.A.*, 122 (1995) 8305m.

#### 16. NITRO AND NITROSO COMPOUNDS

- 1707 Ahmad, F. and Roberts, D.J.: Use of narrow-bore high-performance liquid chromatography-diode array detection for the analysis of intermediates of the biological degradation of 2,4,6-trinitrotoluene. *J. Chromatogr. A*, 693 (1995) 167-175.
- 1708 Barek, J., Hai, P.T., Pacakova, V., Stulik, K., Svagrova, I. and Zima, J.: A study of HPLC separation and spectrophotometric, polarographic and voltametric detection of 4-substituted derivatives of N-nitroso-N-methylaniline. *Fresenius J. Anal. Chem.*, 350 (1994) 678-683.
- 1709 Chen, C.-Z. and Xiao, A.: Development of GPC method for characterizing aging behavior of propellants. *Int. Annu. Conf. ICT*, 25th (1994) 80/1-80/12; *C.A.*, 121 (1994) 304146x.

- 1710 Posyniak, A., Niedzielska, J., Semeniuk, S. and Zmudzki, J.: Solid-phase extraction and liquid chromatography analysis of nitrofurans compounds in meat. *Chem. Anal. (Warsaw)*, 39 (1994) 289-294; *C.A.*, 121 (1994) 254039s.

See also 1418, 2301.

#### 17. AMINES, AMIDES AND RELATED NITROGEN COMPOUNDS

##### 17a. Amines and polyamines

- 1711 Hornero-Mendez, D. and Garrido-Fernandez, A.: Biogenic amines in table oil. Analysis by high-performance liquid chromatography. *Analyst (Cambridge)*, 119 (1994) 2037-2041.
- 1712 Lee, H.K. and Hoffman, N.E.: Retention of some simple organic cations on an anion exchange column. *J. Liq. Chromatogr.*, 17 (1994) 4273-4284.
- 1713 Skarping, G., Dalene, M.D. and Tinnerberg, H.: Biological monitoring hexamethylene- and isophorone-diisocyanate by the determination of hexamethylene- and isophorone-diamine in hydrolysed urine using liquid chromatography and mass spectrometry. *Analyst (Cambridge)*, 119 (1994) 2051-2055.

For additional information see *C.A.*:  
122 (1995) 71121p.

See also 1343, 1373, 1758, 2353, 2417, 2485.

##### 17b. Catecholamines and their metabolites

- 1714 Abate, V.M., Overdorf, G.W. and Huang, T.: The determination of free urinary catecholamines using a new kit for liquid chromatography/electrochemistry. *Curr. Sep.*, 13 (1994) 53-55; *C.A.*, 121 (1994) 246527j.
- 1715 Egger, D., Reischbach, G. and Hüttner, L.: Simultaneous determination of histamine and serotonin in mast cells by high-performance liquid chromatography. *J. Chromatogr. B*, 662 (1994) 103-107.
- 1716 Fornstedt-Wallin, B. and Bergh, I.: Sensitive high-performance liquid chromatographic method for the determination of 5-S-cysteinyldopamine, 5-S-cysteinyl-3,4-dihydroxyphenylacetic acid and 5-S-cysteinyl-3,4-dihydroxyphenylalanine. *J. Chromatogr. B*, 663 (1995) 9-14.
- 1717 Kubalec, P., Brandsteterova, E., Lehotay, J. and Cizmarik, J.: Electrochemical detection in HPLC determination of catecholamines and their main metabolites. *Pharmazie*, 49 (1994) 897-899.
- 1718 Lowe, D.R., March, C., James, J.E. and Karnes, H.T.: A high performance liquid chromatographic method for histamine in plasma using solid phase extraction and fluorecamine derivatization. *J. Liq. Chromatogr.*, 17 (1994) 3563-3570.
- 1719 Murai, S., Saito, H., Masuda, Y., Ohkubo, N. and Itoh, T.: Rapid HPLC assay with coulometric detection for norepinephrine and 3-methoxy-4-hydroxyphenylglycol in the mouse brain. *J. Pharmacol. Toxicol. Methods*, 32 (1994) 99-103; *C.A.*, 122 (1995) 1212v.
- 1720 Nappi, A.J. and Vass, E.: The effects of glutathione and ascorbic acid on the oxidations of 6-hydroxydopa and 6-hydroxydopamine. *Biochim. Biophys. Acta*, 1201 (1994) 498-504.

- 1721 Pastoris, A., Cerutti, L., Sacco, R., de Vecchi, L. and Sbaffi, A.: Automated analysis of urinary catecholamines by high-performance liquid chromatography and on-line sample pretreatment. *J. Chromatogr. B*, 664 (1995) 287-293.
  - 1722 Ragab, G.H., Nohta, H., Kai, M. and Ohkura, Y.: Chemiluminescence determination of catecholamines in human blood plasma and urine using 1,2-diphenylethylenediamine as precolumn derivatization reagent in liquid chromatography. *Anal. Chim. Acta*, 298 (1994) 431-438.
  - 1723 Yamaguchi, M., Monji, H., Aoki, I. and Yashiki, T.: High-performance liquid chromatographic determination of phenylephrine in human serum using column switching with fluorescence detection. *J. Chromatogr. B*, 661 (1994) 93-99.
  - 1724 Zumarraga, M., Andia, I., Davila, R. and Zamolloa, M.I.: Homovanillic acid in plasma determined by HPLC with direct injection of plasma filtrates. *Clin. Chem. (Washington)*, 40 (1994) 2119-2120.
- See also 1583, 1627, 1769, 2074.
- 17c. *Urea and guanidine derivatives*
- 1725 Koebel, M. and Elsener, M.: Determination of urea and its thermal decomposition products by high-performance liquid chromatography. *J. Chromatogr. A*, 689 (1995) 164-169.
  - 1726 Nicholson, L.W., Pfeiffer, C.D., Goralski, C.T., Singaram, B. and Fisher, G.B.: High-performance liquid chromatographic separation of  $\beta$ -amino alcohols. I. Separation of (R,S)-1-(dialkylamino)-2-alkanols on an amylose-based chiral stationary phase. *J. Chromatogr. A*, 687 (1994) 241-248.
  - 1727 Shintani, H. and Inoue, T.: (Determination of blood urea using cation exchange column solid phase extraction combined with HPLC). *Bunseki Kagaku*, 43 (1994) 805-807; *C.A.*, 121 (1994) 275696u.
- See also 1751, 2098.
- 17d. *Other amine derivatives and amides (excl. peptides)*
- 1728 Jemal, M. and Hawthorne, D.: High performance liquid chromatography/ion-spray mass spectrometry of N-ethylmaleimide and acrylic acid ester derivatives for bioanalysis of thiol compounds. *Rapid Commun. Mass Spectrom.*, 8 (1994) 854-857; *C.A.*, 122 (1995) 4878c.
  - 1729 Ståhlberg, O. and Arvidsson, T.: Liquid chromatographic determination of ethylenediaminetetraacetic acid as metal complexes on a porous graphitic carbon column. *J. Chromatogr. A*, 684 (1994) 213-219.
  - 1730 Valladao, M. and Sandine, W.E.: Quaternary ammonium compounds in milk: detection by reverse-phase high performance liquid chromatography and their effect on starter growth. *J. Dairy Sci.*, 77 (1994) 1509-1514; *C.A.*, 121 (1994) 229123g.
- For additional information see *C.A.*:  
121 (1994) 220927q.
- See also 1418, 1514, 1632.
18. AMINO ACIDS AND PEPTIDES; CHEMICAL STRUCTURE OF PROTEINS
  - 18a. *Amino acids and their derivatives*
  - 1731 Alfano, M., Casari, E., Stenico, A., Murone, M. and Grazioli, V.: Improved HPLC determination of urinary hydroxylysine glycosides to study turnover rate of bone in humans and rats. *Clin. Chem. (Washington)*, 40 (1994) 2113-2114.
  - 1732 Alonso, M.L., Alvarez, A.I. and Zapico, J.: Rapid analysis of free amino acids in infant foods. *J. Liq. Chromatogr.*, 17 (1994) 4019-4030.
  - 1733 Amin, M.R., Tomita, Y. and Onodera, R.: Rapid determination of phenylalanine and its related compounds in rumen fluid by high-performance liquid chromatography. *J. Chromatogr. B*, 663 (1995) 201-207.
  - 1734 Baffi, F., Ianni, M.C., Ravera, M. and Magi, E.: Study of the influence of free dissolved amino acids on copper(II) adsorption/remobilization from inorganic fractions of marine sediments using a reversed-phase liquid chromatographic procedure. *Anal. Chim. Acta*, 294 (1994) 127-134.
  - 1735 Basink, V.A., Gromovoy, T.Yu. and Khil'chevskaya, E.G.: Free energies of amino acid adsorption on silica in neutral aqueous medium: estimation from high-performance liquid-chromatographic retention data. *Pol. J. Chem.*, 68 (1994) 777-781; *C.A.*, 121 (1994) 287143z.
  - 1736 Bonnot, G. and Febvay, G.: Fourier processing of liquid chromatograms using flow-radioactive detection. *Anal. Biochem.*, 224 (1995) 254-263.
  - 1737 Brueckner, H., Keller-Hoehl, C. and Wittner, R.: New reagents for chiral amino acid analysis by HPLC. *Chem. Pept. Proteins*, 5/6(Pt. A) (1993) 145-156; *C.A.*, 122 (1995) 4619u.
  - 1738 Brueckner, H., Strecker, B. and Langer, M.: Design of new reagents for the HPLC separation of DL-amino acids. *Chem. Pept. Proteins*, 5/6(Pt. A) (1993) 157-169; *C.A.*, 121 (1994) 296304e.
  - 1739 Brueckner, H., Strecker, B. and Wachsmann, M.: Use of chiral monohalo-s-triazines as novel derivatizing reagents for DL-amino acids. In: Schneider, C.H. and Eberle, A.N. (Editors), *Pept. 1992, Proc. Eur. Pept. Symp.*, 22nd 1992, ESCOM, Leiden, 1993, pp. 449-450; *C.A.*, 122 (1995) 4608q.
  - 1740 Carratu, B., Boniglia, C., Fiesi, C. and Bellomonte, G.: (Effects of technological treatments on food: ion-exchange chromatographic determination of pyridosine and furosine in food and dietetic products). *Riv. Sci. Aliment.*, 22 (1993) 455-457; *C.A.*, 121 (1994) 229115f.
  - 1741 Chen, S.: Partial characterization of the molecular species of phosphatidylserine from human plasma by high-performance liquid chromatography and fast atom bombardment mass spectrometry. *J. Chromatogr. B*, 661 (1994) 1-5.
  - 1742 Chowdhury, S.K., Eshraghi, J., Wolfe, H., Forde, D., Hlavac, A.G. and Johnston, D.: Mass spectrometric identification of amino acid transformations during oxidation of peptides and proteins: modification of methionine and tyrosine. *Anal. Chem.*, 67 (1995) 390-398.
  - 1743 De Vos, F. and Slegers, G.: High-performance liquid chromatographic determination of (4-[ $^{14}$ C]methoxyphenyl)-(5-fluoro-2-hydroxyphenyl)-methyleneaminobutyric acid and its benzophenone metabolite. *J. Chromatogr. A*, 692 (1995) 97-102.

- 1744 DeSilva, K., Strojek, J. and Kuwana, T.: Dual wavelength excitation method for the off-line liquid chromatographic analysis of derivatized amino acids. *Anal. Sci.*, 10 (1994) 573-578; *C.A.*, 122 (1995) 22830k.
- 1745 Gorenweg, M.G.M., van de Merbel, N.C., Slobodnik, J., Lingeman, H. and Brinkman, U.A.T.: Automated determination of weakly acid and basic pollutants in surface water by on-line electroanalysis sample treatment and column liquid chromatography. *Analyst (Cambridge)*, 119 (1994) 1752-1758.
- 1746 Hyun, M.H., Yang, D.H., Kim, H.J. and Ryoo, J.-J.: Mechanistic evaluation of the resolution of  $\alpha$ -amino acids on dynamic chiral stationary phases derived from amino alcohols by ligand-exchange chromatography. *J. Chromatogr. A*, 684 (1994) 189-200.
- 1747 Irvine, G.B.: Amino acid analysis. *Methods Mol. Biol. (Totowa)*, 32(Basic Protein and Peptide Protocols) (1994) 257-265; *C.A.*, 121 (1994) 250057y.
- 1748 Iwase, H., Ozawa, S., Ikuta, M. and Ono, I.: Determination of amino acids in human plasma by liquid chromatography with postcolumn ninhydrin derivatization using a hydroxyapatite cartridge for precolumn deproteinization. *J. Chromatogr. B*, 663 (1995) 15-24.
- 1749 Kamel, S., Brazier, M., Picard, C., Boitte, F., Samson, L., Desmet, G. and Sebert, J.L.: Urinary excretion of pyridinolines crosslinks measured by immunoassay and HPLC techniques in normal subjects and in elderly patients with vitamin D deficiency. *Bone Miner.*, 26 (1994) 197-208; *C.A.*, 121 (1994) 296565r.
- 1750 Kempe, M. and Mosbach, K.: Separation of amino acids, peptides and proteins on molecularly imprinted stationary phases. *J. Chromatogr. A*, 691 (1995) 317-323.
- 1751 Khan, J.K., Kuo, Y.-H., Kebede, N. and Lambein, F.: Determination of non-protein amino acids and toxins in *Lathyrus* by high-performance liquid chromatography with precolumn phenyl isothiocyanate derivatization. *J. Chromatogr. A*, 687 (1994) 113-119.
- 1752 Kim, Y.T., Glerum, C., Noland, T.L. and Hickie, D.: Use of Sep-Pak C<sub>18</sub> cartridges to clean up free amino acids from coniferous needles. *J. Chromatogr. A*, 690 (1995) 226-229.
- 1753 Krishnamurthy, T.: Chirality in microcystins. *J. Am. Soc. Mass Spectrom.*, 5 (1994) 724-730; *C.A.*, 122 (1995) 10561k.
- 1754 Lebet, V., Schneider, H., Arrigoni, E. and Amado, R.: (Critical evaluation of protein determination in food by the Kjeldahl method, based on amino acid analysis). *Mitt. Geb. Lebensmittelforsch. Hyg.*, 85 (1994) 46-58; *C.A.*, 121 (1994) 253993m.
- 1755 Martinez, M., Casillas, J.L., Addo-Yobo, F., Kenney, C.N. and Aracil, J.: An HPLC technique for the study of adsorption of amino acids on functionalized resins. *Spec. Publ. - R. Soc. Chem.*, 158(Separations for Biotechnology 3) (1994) 294-300; *C.A.*, 121 (1994) 296302c.
- 1756 Moret, S., Cherubin, S., Rodriguez-Estrada, M.T. and Lercker, G.: Determination of lysinoalanine by high performance liquid chromatography. *J. High Resolut. Chromatogr.*, 17 (1994) 827-830.
- 1757 Orwar, O., Sandberg, M., Jacobson, I., Sundahl, M. and Folestad, S.: Photochemical characterization and optimization of argon ion laser-induced fluorescence detection of  $\alpha$ -phthalaldehyde/ $\beta$ -mercaptoethanol-labeled amino acids and  $\gamma$ -glutamyl peptides in liquid chromatography: ultratrace analysis with neurobiological samples. *Anal. Chem.*, 66 (1994) 4471-4482.
- 1758 Puig-Deu, M. and Buxaderas, S.: Analytical conditions for the determination of 23 phenylthiocarbonyl amino acids and ethanolamine in musts and wines by high-performance liquid chromatography. *J. Chromatogr. A*, 685 (1994) 21-30.
- 1759 Rothmund, S., Krause, E., Beyermann, M., Dathe, M., Engelhardt, H. and Bienert, M.: Recognition of  $\alpha$ -helical peptide structures using high-performance liquid chromatographic retention data for D-amino acid analogues: influence of peptide amphipathicity and of stationary phase hydrophobicity. *J. Chromatogr. A*, 689 (1995) 219-226.
- 1760 Schmeer, K., Khalifa, M., Császár, J., Farkas, G., Bayer, E. and Molnár-Perl, I.: Compositional analysis of the phenylthiocarbonyl amino acids by liquid chromatography-atmospheric pressure ionization mass spectrometry with particular attention to the cyst(e)ine derivatives. *J. Chromatogr. A*, 691 (1995) 285-299.
- 1761 Simon, G., Hanak, L., Grevillot, G., Szanya, T. and Marton, G.: Preparative-scale amino acid separation by thermal parametric pumping on an ion-exchange resin. *J. Chromatogr. B*, 664 (1995) 17-31.
- 1762 Sormiachi, K., Ikeda, M., Akimoto, K. and Niwa, A.: Rapid determination of dabsylated hydroxyproline from cultured cells by reversed-phase high-performance liquid chromatography. *J. Chromatogr. B*, 664 (1995) 435-439.
- 1763 Takechi, M., Kanda, M., Hori, K., Kurotsu, T. and Saito, Y.: Purification and properties of L-ornithine  $\delta$ -aminotransferase from gramicidin S-producing *Bacillus brevis*. *J. Biochem. (Tokyo)*, 116 (1994) 955-959.
- 1764 Tani, S. and Tani, T.: Determination of methionine and its oxides in gelatin. *J. Photogr. Sci.*, 41 (1993) 172-175; *C.A.*, 121 (1994) 303479w.
- 1765 Tordjman, C., Lhumeau, A., Pastoureaux, P., Meunier, F., Serkiz, B., Volland, J.P. and Bonnet, J.: Evaluation and comparison of urinary pyridinium crosslinks in two rat models of bone loss - ovariectomy and adjuvant polyarthritis - using a new automated HPLC method. *Bone Miner.*, 26 (1994) 155-167; *C.A.*, 121 (1994) 225378w.
- 1766 Toyo'oka, T. and Liu, Y.-M.: High-performance liquid chromatographic resolution of amino acid enantiomers derivatized with fluorescent chiral Edman reagents. *J. Chromatogr. A*, 689 (1995) 23-30.
- 1767 Wallbroehl, Y. and Wagner, J.: Chiral separations of amino acids by capillary electrophoresis and high-performance liquid chromatography employing chiral crown ethers. *J. Chromatogr. A*, 685 (1994) 321-329.
- 1768 Warlow, R.S., Gooley, A., Rajasekariah, P., Oszarac, N. and Walls, R.S.: A preparative method for sequencing proteins and peptides: *In situ* gel staining with subsequent passive elution onto polyvinylidene difluoride membranes. *Electrophoresis (Weinheim)*, 16 (1995) 84-91.
- 1769 Yi, Y.-H., Liao, W.-P. and Lu, X.: Simultaneous determination of tryptophan, 5-hydroxytryptophan, 5-hydroxytryptamine, 5-hydroxyindoleacetic acid, 4-hydroxy-3-methoxyphenylacetic acid and 3-methoxy-4-hydroxyphenylglycol in human cerebrospinal fluid. *J. Chromatogr. B*, 661 (1994) 143-148.
- 1770 Yoshida, H., Arai, N., Sugoh, M., Iwabuchi, J., Shiomi, K., Shinose, M., Tanaka, Y. and Omura, S.: 4-Chlorothreonine, a herbicidal antimetabolite produced by *Streptomyces* sp. OH-5093. *J. Antibiot.*, 47 (1994) 1165-1166.

- 1771 Young, P.B., Molloy, A.M., Scott, J.M. and Kennedy, D.G.: A rapid high performance liquid chromatographic method for determination of homocysteine in porcine tissue. *J. Liq. Chromatogr.*, 17 (1994) 3553-3561.
- 1772 Zhang, J., Tsuboyama, N. and Kodama, H.: Simultaneous determination of cystathionine and NAC-dystathionine using liquid chromatography-atmospheric pressure chemical ionization mass spectrometry. *Anal. Biochem.*, 224 (1995) 17-20.
- 1773 Zhao, M. and Bada, J.L.: Determination of  $\alpha$ -dialkylamino acids and their enantiomers in geological samples by high-performance liquid chromatography after derivatization with a chiral adduct of o-phthalaldehyde. *J. Chromatogr. A*, 690 (1995) 55-63.
- For additional information see C.A.:  
121 (1994) 286690p.
- See also 1786, 1807, 1837, 1843, 1860, 1941, 1943, 2268, 2417.
- 18b. *Peptides, peptidic and proteinous hormones, growth factors*
- 1774 Acedo, M.I., Pueyo, E. and Polo, M.C.: Preliminary studies on peptides in wine by HPLC. *Am. J. Enol. Vitic.*, 45 (1994) 167-172; C.A., 121 (1994) 279168w.
- 1775 Affolter, M., Watts, J.D., Krebs, D.L. and Aebersold, R.: Evaluation of two-dimensional phosphopeptide maps by electrospray ionization mass spectrometry of recovered peptides. *Anal. Biochem.*, 223 (1994) 74-81.
- 1776 Aldrich-Wright, J.R., Newman, P.D., Rao, K.R.N., Vagg, R.S. and Williams, P.A.: Chiral metal complexes. XLII. Reversed-phase high-performance liquid chromatographic separation of racemic dipeptides as their ternary Co(III) complexes with a chiral triamine. *J. Chromatogr. A*, 693 (1995) 366-370.
- 1777 Andren, P.E. and Caprioli, R.M.: Microdialysis/mass spectrometry of neuropeptides. In: Matsuo, T. (Editor), *Biol. Mass Spectrom.: Present Future, [Proc. Kyoto '92 Int. Conf.]* 1992, Wiley, Chichester, 1994, pp. 355-367; C.A., 121 (1994) 225740h.
- 1778 Apparailly, F. and Combarnous, Y.: Role of sialic acid residues in the *in vitro* superactivity of human choriongonadotropin (hCG) in rat Leydig cells. *Biochim. Biophys. Acta*, 1224 (1994) 559-565.
- 1779 Ball, H.L., Bertolini, G., Levi, S. and Mascagni, P.: Purification of synthetic peptides with the aid of reversible chromatographic probes. *J. Chromatogr. A*, 686 (1994) 73-83.
- 1780 Battersby, J.E., Guzzetta, A.W. and Hancock, W.S.: Application of capillary high-performance liquid chromatography to biotechnology, with reference to the analysis of recombinant DNA-derived human growth hormone. *J. Chromatogr. B*, 662 (1994) 335-342.
- 1781 Battersby, J.E., Mukku, V.R., Clark, R.G. and Hancock, W.S.: Affinity purification and microcharacterization of recombinant DNA-derived human growth hormone isolated from an *in vivo* model. *Anal. Chem.*, 67 (1995) 447-455.
- 1782 Brakch, N., Cohen, P. and Boileau, G.: Processing of human prosomatostatin in AbT cells: S-28 and S-14 are generated in different secretory pathways. *Biochem. Biophys. Res. Commun.*, 205 (1994) 221-229.
- 1783 Cartier, P.G., Maikner, J.J., Deissler, K.C., Fisher, J.R., Barstow, L.E. and Fuentes, G.E.: Optimization of preparative RP purification of peptides by correctly matching media pore size and capacity to the needs of the purification. In: Hodges, R.S. and Smith, J.A. (Editors), *Pept.: Chem., Struct. Biol., Proc. Am. Pept. Symp., 13th 1993*, ESCOM, Leiden, 1994, pp. 232-234; C.A., 121 (1994) 250062w.
- 1784 Chen, J.-G., Woltman, S.J. and Weber, S.G.: Sensitivity and selectivity of the electrochemical detection of the copper(II) complexes of bioactive peptides, and comparison to model studies by rotating ring-disc electrode. *J. Chromatogr. A*, 691 (1995) 301-315.
- 1785 Davies, J.S., Enjalbal, C. and Llewellyn, G.: Progress towards a routine method for chiral analysis of synthetic peptides. In: Schneider, C.H. and Eberle, A.N. (Editors), *Pept. 1992, Proc. Eur. Pept. Symp., 22nd, 1992*, ESCOM, Leiden, 1993, pp. 453-454; C.A., 121 (1994) 296273u.
- 1786 De Antonis, K.M., Brown, P.R. and Cohen, S.A.: High-performance liquid chromatographic analysis of synthetic peptides using derivatization with 6-aminoquinolyl-N-hydroxysuccinimidyl carbamate. *Anal. Biochem.*, 223 (1994) 191-197.
- 1787 Furka, A., Sebestyen, F. and Dibo, G.: Computer program facilitating separation of peptides from mixtures prepared by the 'portioning-mixing' method. In: Schneider, C.H. and Eberle, A.N. (Editors), *Pept. 1992, Proc. Eur. Pept. Symp., 22nd, 1992*, ESCOM, Leiden, 1993, pp. 463-464; C.A., 121 (1994) 276033n.
- 1788 Griehl, C., Weigt, J. and Jeschkeit, H.: Use of tryptophan peptides for an HPLC racemization test with fluorescence detection. *J. High Resolut. Chromatogr.*, 17 (1994) 700-704.
- 1789 Hansen, P. and Lindeberg, G.: Importance of the  $\alpha$ -amino group in the selective purification of synthetic histidine peptides by immobilised metal ion affinity chromatography. *J. Chromatogr. A*, 690 (1995) 155-159.
- 1790 Hoermann, R., Berger, P., Spoettl, G., Gillesberger, F., Kardana, A., Cole, L.A. and Mann, K.: Immunological recognition and clinical significance of nicked human chorionic gonadotropin in testicular cancer. *Clin. Chem. (Washington)*, 40 (1994) 2306-2312.
- 1791 Klyushnichenko, V.E., Yakimov, S.A., Arutyunyan, A.M., Ivanov, A.E., Maltsev, K.V. and Wulfson, A.N.: Recombinant human insulin V. Optimization of the reversed-phase high-performance liquid chromatographic separation. *J. Chromatogr. B*, 662 (1994) 363-369.
- 1792 Korakas, D. and Valkó, K.: Comparative study of the retention behavior of lipidic peptides on RP-18 and Supelcosil<sup>TM</sup> LC-ABZ stationary phases. *J. Liq. Chromatogr.*, 17 (1994) 3571-3584.
- 1793 Krause, E. and Bienert, M.: Separation of peptides by reversed-phase liquid chromatography on polyalkylene columns. *Chem. Pept. Proteins*, 5/6(Pt. A) (1993) 135-143; C.A., 121 (1994) 296303d.
- 1794 Krause, E., Beyermann, M., Dathe, M., Rothmund, S. and Bienert, M.: Location of an amphipathic  $\alpha$ -helix in peptides using reversed-phase HPLC retention behavior of D-amino acids analogs. *Anal. Chem.*, 67 (1995) 252-258.

- 1795 Krause, E., Beyermann, M., Dathe, M., Wenschuh, H., Rothe-mund, S. and Bienert, M.: Correlation of  $\alpha$ -helical secondary peptide structure and retention behavior in reversed-phase HPLC. In: Hodges, R.S. and Smith, J.A. (Editors), *Pept.: Chem., Struct. Biol., Proc. Am. Pept. Symp., 13th 1993*, ESCOM, Leiden, 1994, pp. 147-148; C.A., 121 (1994) 250063x.
  - 1796 Lakhari, H., Legendre, E., Muller, D. and Jozefonvicz, J.: High-performance affinity chromatography of insulin on coated silica grafted with sialic acid. *J. Chromatogr. B*, 664 (1995) 163-173.
  - 1797 Lee, K.C., Yoon, J.Y., Woo, B.H., Kim, C.K. and DeLuca, P.P.: Post-column fluorescence HPLC for salmon calcitonin formula-tions. *Int. J. Pharm.*, 114 (1995) 215-220; C.A., 122 (1995) 64521r.
  - 1798 Melzig, M.F. and Heder, G.: Dexamethasone induced enhanced enkephalin degradation by angiotensin-converting enzyme (ACE) of endothelial cells. *Pharmazie*, 50 (1995) 1139-141.
  - 1799 Murata, H., Shoji, H., Oshikata, M., Harada, K.-I., Suzuki, M., Kondo, F. and Goto, H.: High-performance liquid chroma-tography with chemiluminescence detection of derivatized mi-crocystins. *J. Chromatogr. A*, 693 (1995) 263-270.
  - 1800 Nicholls, I.A., Ramström, O. and Mosbach, K.: Insights into the role of the hydrogen bond and hydrophobic effect on recogni-tion in molecularly imprinted polymer synthetic peptide recep-tor mimics. *J. Chromatogr. A*, 691 (1995) 349-353.
  - 1801 Ono, M., Raab, G., Lau, K., Abraham, J.A. and Klagsbrun, M.: Purification and characterization of transmembrane forms of heparin-binding EGF-like growth factor. *J. Biol. Chem.*, 269 (1994) 31315-31321.
  - 1802 Perrin, E., Miclo, L., Driou, A. and Linden, G.: Rapid determina-tion of the ratios of three aromatic residues in peptides by reversed-phase high-performance liquid chromatography with a high-resolution photodiode-array detector. *J. Chromatogr. B*, 664 (1995) 267-276.
  - 1803 Piemonte, F., Bernardini, S., Massoud, R. and Corasaniti, M.T.: Determination of serum concentrations of adamantyl dipeptide by high performance liquid chromatography after intravenous injection in rats. *Clin. Chem. Enzymol. Commun.*, 6 (1994) 157-162; C.A., 121 (1994) 244901c.
  - 1804 Reimeier, C., Schneider, I., Schneider, W., Schäfer, H.-L. and Elstner, E.F.: Effects of ethanolic extracts from *Eschscholtzia californica* and *Corydalis cava* on dimerization and oxidation of enkephalins. *Arzneim.-Forsch.*, 45 (1995) 132-136.
  - 1805 Roboti, A., Livaniou, E., Evangelatos, G.P., Tsoupras, G., Tsolas, O. and Ithakissios, D.S.: Large-scale chromatofocusing-based method for isolating thymosin  $\beta_4$  and thymosin  $\beta_9$  from bovine tissues. *J. Chromatogr. B*, 662 (1994) 27-34.
  - 1806 Roher, A.E., Kasunic, T.C., Woods, A.S., Cotter, R.J., Ball, M.J. and Fridman, R.: Proteolysis of A $\alpha$  peptide from Alzheimer disease brain by gelatinase A. *Biochem. Biophys. Res. Commun.*, 205 (1994) 1755-1761.
  - 1807 Schepky, A.G., Schulz-Knappe, P. and Forssmann, W.-G.: High-performance liquid chromatographic determination of sulfated peptides in human hemofiltrate using a radioactivity monitor. *J. Chromatogr. A*, 691 (1995) 255-261.
  - 1808 Shadle, P.J., Silverness, K.B.: Process for purification of basic fibroblast growth factor. U.S. US 5,331,095 (Cl. 530-399; C07K3/22), 19 Jul. 1994, Appl. 45,929, 12 Apr. 1993; 9 p.; C.A., 122 (1995) 17184y.
  - 1809 She, Q.-B., Nagao, I., Hayakawa, T. and Tsuge, H.: A simple HPLC method for the determination of S-adenosylmethionine and S-adenosylhomocysteine in rat tissues: the effect of vitamin B6 deficiency on these concentrations in rat liver. *Biochem. Biophys. Res. Commun.*, 205 (1994) 1748-1754.
  - 1810 Shon, K., Hasson, A., Spira, M.E., Cruz, L.J., Gray, W.R. and Olivera, B.M.:  $\delta$ -Conotoxin GmVIA, a novel peptide from the venom of *Conus gloriamaris*. *Biochemistry*, 33 (1994) 11420-11425.
  - 1811 Szokan, G., Majer, Z., Kollat, E., Hollosi, M., Kajtar, M. and Peredy-Kajtar, M.: Chromatographic and racemization studies on thio peptides. In: Schneider, C.H. and Eberle, A.N. (Editors), *Pept. 1992, Proc. Eur. Pept. Symp., 22nd 1992*, ESCOM, Leiden, 1993, pp. 455-456; C.A., 121 (1994) 275683n.
  - 1812 Tarelli, E., Corran, P.H., Bingham, B.R., Mollison, H. and Wait, R.: Lysine vasopressin undergoes rapid glycation in the pres-ence of reducing sugars. *J. Pharm. Biomed. Anal.*, 12 (1994) 1355-1361.
  - 1813 Wicar, S., Mulkerrin, M.G., Bathory, G., Khundkar, L.H. and Karger, B.L.: Conformational changes in the reversed phase liquid chromatography of recombinant human growth hor-mone as a function of organic solvent: the molten globule state. *Anal. Chem.*, 66 (1994) 3908-3915.
  - 1814 Yang, C.-S., Chou, S.-T., Lin, N.-N., Liu, L., Tsai, P.-J., Kuo, J.-S. and Lai, J.-S.: Determination of extracellular glutathione in rat brain by microdialysis and high-performance liquid chroma-tography with fluorescence detection. *J. Chromatogr. B*, 661 (1994) 231-235.
- For additional information see C.A.:  
121 (1994) 246510y.
- See also 1750, 1753, 1757, 1759, 1830, 1853, 1855, 1870, 2052, 2053, 2169, 2345, 2420.
- 18c. *Elucidation of structure of proteins and enzymes*
- 1815 Baldwin, J.S., Lee, L., Leung, T.K., Muruganandam, A. and Mu-tus, B.: Identification of the site of non-enzymatic glycation of glutathione peroxidase: rationalization of the glycation-related catalytic alterations on the basis of three-dimensional protein structure. *Biochim. Biophys. Acta*, 1247 (1995) 60-64.
  - 1816 Ball, K. and Preiss, J.: Allosteric sites of the large subunit of the spinach leaf ADP-glucose pyrophosphorylase. *J. Biol. Chem.*, 269 (1994) 24706-24711.
  - 1817 Bischoff, R. and Kolbe, H.V.J.: Deamidation of asparagine and glutamine residues in proteins and peptides: structural determi-nants and analytical methodology. *J. Chromatogr. B*, 662 (1994) 261-278 - a review with 58 refs.
  - 1818 Bouchon, B., Jaquinod, M., Klarskov, K., Trottein, F., Klein, M., van Dorsselaer, A., Bischoff, R. and Roitsch, C.: Analysis of the primary structure and post-translational modifications of the *Schistosoma mansoni* antigen Smp28 by electrospray mass spectrometry. *J. Chromatogr. B*, 662 (1994) 279-290.
  - 1819 Clerc, F.F., Monégier, B., Faucher, D., Cuiné, F., Pourcet, C., Holt, J.C., Tang, S.-Y., van Dorsselaer, A., Becquart, J. and Vuilhorgne, M.: Primary structure control of recombinant proteins using high-performance liquid chromatography, mass spectrometry and microsequencing. *J. Chromatogr. B*, 662 (1994) 245-259.

- 1820 Dell'Angelica, E.C., Schleicher, C.H. and Santome, J.A.: Primary structure and binding properties of calgranulin C, a novel S100-like calcium-binding protein from pig granulocytes. *J. Biol. Chem.*, 269 (1994) 28929-28936.
- 1821 Fields, C.G. and Fields, G.B.: Edman degradation sequence analysis of resin-bound peptides: characterization of unusual and side chain protected PTH-amino acids. In: Schneider, C.H. and Eberle, A.N. (Editors), *Pept. 1992, Proc. Eur. Pept. Symp., 22nd 1992*, ESCM, Leiden, 1993, pp. 447-448; C.A., 121 (1994) 275682m.
- 1822 Fisichella, S., Foti, S., Maccarrone, G. and Saletti, R.: Tryptic peptide mapping of sequence 299-585 of human serum albumin by high-performance liquid chromatography and fast atom bombardment mass spectrometry. *J. Chromatogr. A*, 693 (1995) 33-44.
- 1823 Garmroudi, F. and MacDonald, R.G.: Localization of the insulin-like growth factor II (IGF-II) binding/cross-linking site of the IGF-II/mannose 6-phosphate receptor to extracellular repeats 10-11. *J. Biol. Chem.*, 269 (1994) 26944-26952.
- 1824 Gaucheron, F., Mollé, D., Léonil, J. and Maubois, J.-L.: Selective determination of phosphopeptide  $\beta$ -CN(1-25) in a  $\beta$ -casein digest by adding iron: characterization by liquid chromatography with on-line electrospray-ionization mass spectrometric detection. *J. Chromatogr. B*, 664 (1995) 193-200.
- 1825 Hancock, W.S., Chloupek, R.C., Kirkland, J.J. and Snyder, L.R.: Temperature as a variable in reversed-phase high-performance liquid chromatographic separations of peptide and protein samples. I. Optimizing the separation of a growth hormone tryptic digest. *J. Chromatogr. A*, 686 (1994) 31-43.
- 1826 Honma, Y., Niimi, M., Uchiyama, T., Takahashi, Y. and Odani, S.: Evidence for conformational change of fatty acid-binding protein accompanying binding of hydrophobic ligands. *J. Biochem. (Tokyo)*, 116 (1994) 1025-1029.
- 1827 Kataoka, K., Tanizawa, K., Fukui, T., Ueno, H., Yoshimura, T., Esaki, N. and Soda, K.: Identification of active site lysyl residues of phenylalanine dehydrogenase by chemical modification with methyl acetyl phosphate combined with site-directed mutagenesis. *J. Biochem. (Tokyo)*, 116 (1994) 1370-1376.
- 1828 Kumazaki, T., Ishii, S.-i. and Yokosawa, H.: Identification of the reactive site of ascidian trypsin inhibitor. *J. Biochem. (Tokyo)*, 116 (1994) 787-793.
- 1829 Lemaire, M., Schmitter, J.-M., Issakidis, E., Miginiac-Maslow, M., Gadal, P. and Decottignies, P.: Essential histidine at the active site of sorghum leaf NADP-dependent malate dehydrogenase. *J. Biol. Chem.*, 269 (1994) 27291-27296.
- 1830 Malmquist, G.: Multivariate evaluation of peptide mapping using the entire chromatographic profile. *J. Chromatogr. A*, 687 (1994) 89-100.
- 1831 McKay, G.A., Robinson, R.A., Lane, W.S. and Wright, G.D.: Active-site labeling of an aminoglycoside antibiotic phosphotransferase (APH(3')-IIIa). *Biochemistry*, 33 (1994) 14115-14120.
- 1832 Michel, L., Garin, J., Vincon, M., Gagnon, J. and Vignais, P.: Mapping of the pyrophosphate binding sites of beef heart mitochondrial F<sub>1</sub>-ATPase by photolabelling with azidonitrophenyl [ $\alpha$ -<sup>32</sup>P]pyrophosphate. *Biochim. Biophys. Acta*, 1228 (1995) 67-72.
- 1833 Miyata, T., Inagi, R., Wada, Y., Yeda, Y., Iida, Y., Takahashi, M., Taniguchi, N. and Maeda, K.: Glycation of human  $\beta_2$ -microglobulin in patients with hemodialysis-associated amyloidosis: identification of the glycosylated sites. *Biochemistry*, 33 (1994) 12215-12221.
- 1834 Morishima-Kawashima, M., Hasegawa, M., Takio, K., Suzuki, M., Yoshida, H., Titani, K. and Ihara, Y.: Proline-directed and non-proline-directed phosphorylation of PHF-tau. *J. Biol. Chem.*, 270 (1995) 823-829.
- 1835 Mott, J.D., Pramanik, B.C., Moomaw, C.R., Afendis, S.J., DeMartino, G.N. and Slaughter, C.A.: PA28, An activator of the 20 S proteasome, is composed of two nonidentical but homologous subunits. *J. Biol. Chem.*, 269 (1994) 31466-31471.
- 1836 Robertson, J.G., Adams, G.W., Medzihradszky, K.F., Burlingame, A.L. and Villafranca, J.J.: Complete assignment of disulfide bonds in bovine dopamine  $\beta$ -hydroxylase. *Biochemistry*, 33 (1994) 11563-11575.
- 1837 Ronca, F., Ranieri-Raggi, M., Brown, P.E., Moir, A.J.G. and Raggi, A.: Evidence of a species-differentiated regulatory domain within the N-terminal region of skeletal muscle AMP deaminase. *Biochim. Biophys. Acta*, 1209 (1994) 123-129.
- 1838 Shahrokhi, Z., Eberlein, G., Buckley, D., Parandhi, M.V., Aswad, D.W., Stratton, P., Mischak, R. and Wang, Y.J.: Major degradation products of basic fibroblast growth factor: detection of succinimide and iso-aspartate in place of aspartate. *Pharm. Res.*, 11 (1994) 936-944; C.A., 121 (1994) 263389h.
- 1839 Shapiro, M.J., Panomitos, D. and Koshland, D.E., Jr.: Interactions between the methylation sites of the *Escherichia coli* aspartate receptor mediated by the methyltransferase. *J. Biol. Chem.*, 270 (1995) 751-755.
- 1840 Warburton, R.J. and Seybert, D.W.: Structural and functional characterization of bovine adrenodoxin reductase by limited proteolysis. *Biochim. Biophys. Acta*, 1246 (1995) 39-46.
- 1841 Watts, J.D., Affolter, M., Krebs, D.L., Wange, R.L., Samelson, L.E. and Aebersold, R.: Identification by electrospray ionization mass spectrometry of the sites of tyrosine phosphorylation induced in activated Jurkat T cells on the protein tyrosine kinase ZAP-70. *J. Biol. Chem.*, 269 (1994) 29520-29529.
- 1842 Xue, J., Kalafatis, M., Silveira, J.R., Kung, C. and Mann, K.G.: Determination of the disulfide bridges in factor Va heavy chain. *Biochemistry*, 33 (1994) 13109-13116.
- 1843 Yamada, H., Seno, M., Kobayashi, A., Moriyama, T., Kosaka, M., Ito, Y. and Imoto, T.: An S-alkylating reagent with positive charges as an efficient solubilizer of denatured disulfide-containing proteins. *J. Biochem. (Tokyo)*, 116 (1994) 852-857.
- 1844 Yamakawa, Y., Omori-Satoh, T. and Mebs, D.: Hemorrhagic principles in the venom of *Bitis arietans*, a viperous snake. II. Enzymatic properties with special reference to substrate specificity. *Biochim. Biophys. Acta*, 1247 (1995) 17-23.
- 1845 Zhang, Y., Beck, C.A., Poletti, A., Edwards, D.P. and Weigel, N.L.: Identification of phosphorylation sites unique to the B form of human progesterone receptor. *In vitro* phosphorylation by casein kinase II. *J. Biol. Chem.*, 269 (1994) 31034-31040.
- 1846 Zheng, Y., Bergold, A. and Duffel, M.W.: Affinity labeling of aryl sulfotransferase IV. Identification of a peptide sequence at the binding site for 3'-phosphoadenosine-5'-phosphosulfate. *J. Biol. Chem.*, 269 (1994) 30313-30319.

See also 1598, 1766, 1768, 1852, 1855, 1911, 1932, 1956, 1983, 2042.

## 19. PROTEINS

## 19a. General techniques

- 1847 Adrover, A. and Barba, D.: Analysis of size exclusion chromatographic properties of proteins. *Spec. Publ. - R. Soc. Chem.*, 158(Separations for Biotechnology 3) (1994) 15-21; C.A., 121 (1994) 296295c.
- 1848 Assouline, Z., Coutinho, J.B., Gilkes, N.R., Greenwood, J.M., Kilburn, D.G., Le, K.D., Miller, R.C., Jr., Ong, E., Ramirez, C. and Warren, R.A.J.: Cellulose-binding domains as affinity tags for protein purification. *Spec. Publ. - R. Soc. Chem.*, 158(Separations for Biotechnology 3) (1994) 43-47; C.A., 121 (1994) 296296d.
- 1849 Banks, J.F., Jr.: Separation and analysis of proteins by perfusion liquid chromatography and electrospray ionization mass spectrometry. *J. Chromatogr. A*, 691 (1995) 325-330.
- 1850 Banks, J.F., Jr., Quinn, J.P. and Whitehouse, C.M.: LC/ESI-MS determination of proteins using conventional liquid chromatography and ultrasonically assisted electrospray. *Anal. Chem.*, 66 (1994) 3688-3695.
- 1851 Bloomingburg, G.F. and Carta, G.: Separation of protein mixtures by continuous annular chromatography with step elution. *Chem. Eng. J. (Lausanne)*, 55 (1994) 19B-27B; C.A., 121 (1994) 225379x.
- 1852 Boschetti, E., Guerrier, L., Girot, P. and Horvath, J.: Preparative high-performance liquid chromatographic separation of proteins with HyperD ion-exchange supports. *J. Chromatogr. B*, 664 (1995) 225-231.
- 1853 Boyes, B.E. and Walker, D.G.: Selectivity optimization of reversed-phase high-performance liquid chromatographic peptide and protein separations by varying bonded-phase functionality. *J. Chromatogr. A*, 691 (1995) 337-347.
- 1854 Cai, C.-h., Romano, V.A. and Dubin, P.L.: Ionic strength dependence of protein retention on Superose 12 in SEC-IEC mixed mode chromatography. *J. Chromatogr. A*, 693 (1995) 251-261.
- 1855 Chloupek, R.C., Hancock, W.S., Marchylo, B.A., Kirkland, J.J., Boyes, B.E. and Snyder, L.R.: Temperature as a variable in reversed-phase high-performance liquid chromatographic separations of peptide and protein samples. II. Selectivity effects observed in the separation of several peptide and protein mixtures. *J. Chromatogr. A*, 686 (1994) 45-59.
- 1856 Compton, B.J. and Kreilgaard, L.: Chromatographic analysis of therapeutic proteins. *Anal. Chem.*, 66 (1994) 1175A-1180A - a review with 41 refs.
- 1857 Freitag, R. and Breier, J.: Development of an automated displacement chromatography purification process for biotechnologically produced proteins. *Spec. Publ. - R. Soc. Chem.*, 158(Separations for Biotechnology 3) (1994) 166-172; C.A., 121 (1994) 296297e.
- 1858 Freitag, R. and Breier, J.: Displacement chromatography in biotechnological downstream processing. *J. Chromatogr. A*, 691 (1995) 101-112.
- 1859 Guo, L.: (Study on the role of formic acid in formic acid-isopropanol system in protein separation). *Sepu*, 12 (1994) 278-280; C.A., 121 (1994) 225377v.
- 1860 Hellman, U., Wernstedt, C., Gonez, J. and Heldin, C.-H.: Improvement of an "in-gel" digestion procedure for the micro-preparation of internal protein fragments for amino acid sequencing. *Anal. Biochem.*, 224 (1995) 451-455.
- 1861 Hess, D., Nika, H., Chow, D.T., Bures, E.J., Morrison, H.D. and Aebersold, R.: Liquid chromatography-electrospray ionization mass spectrometry of 4-(3-pyridinylmethylaminocarboxypropyl) phenylhydantoins. *Anal. Biochem.*, 224 (1995) 373-381.
- 1862 Hotz, N.J. and Bauer, W.F.: Determination of strongly protein-bound borocaptate species by high-performance liquid chromatography with online inductively coupled plasma atomic emission spectroscopy detection of boron. In: Soloway, A.H., Barth, R.F. and Carpenter, D.E. (Editors), *Adv. Neutron Capture Ther.*, [Proc. Int. Symp.], 5th 1992, Plenum, New York, 1993, pp. 439-443; C.A., 121 (1994) 250042q.
- 1863 Irvine, G.B.: Molecular-weight estimation for native proteins using size-exclusion high-performance liquid chromatography. *Methods Mol. Biol. (Totowa)*, 32(Basic Protein and Peptide Protocols) (1994) 267-274; C.A., 121 (1994) 250058z.
- 1864 Jacob, L.R., Rathgeber, G. and Mack, M.: Scale-up of purification of proteins from egg white by tentacle ion-exchange chromatography. *Chim. Oggi*, 12 (1994) 25-28; C.A., 121 (1994) 275710u.
- 1865 Jing, G., Zhou, B., Liu, L., Zhou, J. and Liu, Z.: Resolution of proteins on a phenyl-Superose HR5/5 column and its application to examining the conformation homogeneity of refolded recombinant staphylococcal nuclease. *J. Chromatogr. A*, 685 (1994) 31-37.
- 1866 Kalef, E. and Gitler, C.: Purification of vicinal dithiol-containing proteins by arsenical-based affinity chromatography. *Methods Enzymol.*, 233(Oxygen Radicals in Biological Systems, Pt. C) (1994) 395-403; C.A., 121 (1994) 225364p.
- 1867 Khamilich, S., Loirat, M.J., Blanchard, D., le Maire, M., Bailly, P., Cartron, J.P. and Bertrand, O.: Influence of the size of the polar head of non-ionic detergents on membrane proteins immunoaffinity purification. *J. Biochem. Biophys. Methods*, 29 (1994) 123-134; C.A., 122 (1995) 4792v.
- 1868 Kim, Y.J. and Cramer, S.M.: Experimental studies in metal affinity displacement chromatography of proteins. *J. Chromatogr. A*, 686 (1994) 193-203.
- 1869 Kleinmann, I., Plicka, J., Smidl, P. and Vins, I.: Hydrophobic interaction chromatography of proteins on HEMA-based sorbents. *Am. Lab. (Shelton)*, 26 (1994) 34J, 34L; C.A., 121 (1994) 225368t.
- 1870 Kopaciewicz, W., Kellard, E. and Cox, G.B.: High velocity reversed-phase chromatography of proteins and peptides: use of conventional C18, 300 Å, 15 µm particles. *J. Chromatogr. A*, 690 (1995) 9-19.
- 1871 Laubscher, J.C.: (Technique of fast protein liquid chromatography (FPLC)). *Schweiz. Lab.-Z.*, 51 (1994) 80-85; C.A., 121 (1994) 225231t.
- 1872 Leonard, M., Fournier, C. and Dellacherie, E.: Polyvinyl alcohol-coated macroporous polystyrene particles as stationary phases for the chromatography of proteins. *J. Chromatogr. B*, 664 (1995) 39-46.
- 1873 Li, Y.-M., Liao, J.-L., Nakazato, K.-i., Mohammad, J., Terenius, L. and Hjerten, S.: Continuous beds for microchromatography: cation-exchange chromatography. *Anal. Biochem.*, 223 (1994) 153-158.

- 1874 Mao, Q.-M., Prince, I.G. and Hearn, M.T.W.: High-performance liquid chromatography of amino acids, peptides and proteins. CXXXIX. Impact of operating parameters in large-scale chromatography of proteins. *J. Chromatogr. A*, 691 (1995) 273-283.
- 1875 Millot, M.-C., Hervé, F. and Sébille, B.: Retention behaviour of proteins on poly(vinylimidazole)-copper(II) complexes supported on silica: application to the fractionation of desialylated human  $\alpha_1$ -acid glycoprotein variants. *J. Chromatogr. B*, 664 (1995) 55-67.
- 1876 Mueller, W.: Separation of proteins and nucleic acids. *Methods Enzymol.*, 228(Aqueous Two-Phase Systems) (1994) 193-206; C.A., 121 (1994) 249898s - a review with 20 refs.
- 1877 Nakayama, H., Kanai, M., Seta, K., Isobe, T. and Okuyama, T.: (Analysis of micro-quantities of proteins by combined two-dimensional electrophoresis-capillary column HPLC-ESI/TSQMS). *Kuromatogurafi*, 15 (1994) 72-73; C.A., 121 (1994) 275701s.
- 1878 Ngo, T.T.: Rapid purification of immunoglobulin G using azareophilic chromatography: novel mode of protein-solid phase interactions. *J. Chromatogr. B*, 662 (1994) 351-356.
- 1879 Noinville, V., Craescu, C.T., Vidal-Madjar, C. and Sebille, B.: Molecular approach to protein-polymer interactions in ion-exchange chromatography. *J. Chromatogr. B*, 664 (1995) 33-38.
- 1880 Reif, O.-W., Nier, V. and Freitag, R.: High-performance membrane chromatography (HPMC), superior to conventional protein-HPLC? *Prog. Biotechnol.*, 9(ECB6: Proceedings of the 6th European Congress on Biotechnology, 1993, Pt. 1) (1994) 521-524; C.A., 121 (1994) 250044s.
- 1881 Roe, S.D.: Protein A leakage from affinity adsorbents. *Spec. Publ.-R. Soc. Chem.*, 158(Separations for Biotechnology 3) (1994) 427-432; C.A., 121 (1994) 296294b.
- 1882 Shi, J.Y. and Goffe, R.A.: Comprehensive study on binding capacity of human immunoglobulin G to Avid A1 affinity gel. *J. Chromatogr. A*, 686 (1994) 61-71.
- 1883 Tongta, A., Liapis, A.I. and Siehr, D.J.: Equilibrium and kinetic parameters of the adsorption of  $\alpha$ -chymotrypsinogen A onto hydrophobic porous adsorbent particles. *J. Chromatogr. A*, 686 (1994) 21-29.
- 1884 Tsuneda, S., Saito, K., Furusaki, S. and Sugo, T.: High-throughput processing of proteins using a porous and tentacle anion-exchange membrane. *J. Chromatogr. A*, 689 (1995) 211-218.
- 1885 Uchida, K. and Stadtman, E.R.: Quantitation of 4-hydroxynonenal protein adducts. *Methods Enzymol.*, 233(Oxygen Radicals in Biological Systems, Pt. C) (1994) 371-380; C.A., 121 (1994) 225363n.
- 1886 Watanabe, E., Tsoka, S. and Asenjo, J.A.: Selection of chromatographic protein purification operations based on physicochemical properties. *Ann. N.Y. Acad. Sci.*, 721(Recombinant DNA Technology II) (1994) 348-364; C.A., 121 (1994) 275680j.
- For additional information see C.A.:  
121 (1994) 253752g, 275707y.
- See also 1301, 1419, 1451, 1463, 1466, 1471, 1475, 1479, 1480, 1516, 1518, 1750, 1796, 1899, 2049, 2052, 2492.
- 19b. *Proteins of cells, viruses and subcellular particles*
- 1887 Avliyakov, N.K., Sedel'nikova, S.E. and Shikaeva, O.S.: (Isolation of proteins from ribosomal 50S-subparticles of *Thermus thermophilus*. *Biokhimiya (Moscow)*, 59 (1994) 1003-1010.
- 1888 Claus, P., Schulze, E. and Wisniewski, J.R.: Insect proteins homologous to mammalian high mobility group proteins IY (HMG I/Y). Characterization and binding to linear and four-way junction DNA. *J. Biol. Chem.*, 269 (1994) 33042-33048.
- 1889 Fransen, M., Brees, C.A.E., de Béthune, B., van Veldhoven, P.P. and Mannaerts, G.P.: The 80 kDa cytosolic protein that binds the C-terminal part of rat acyl-CoA oxidase is not a peroxisomal import receptor but a prolyl-endopeptidase. *Biochim. Biophys. Acta*, 1201 (1994) 157-164.
- 1890 Ichimura, T., Ikuta, N., Uda, Y., Horigome, T. and Omata, S.: Separation of membrane proteins solubilized with a non-denaturing detergent and a high salt concentration by hydroxyapatite high-performance liquid chromatography. *Anal. Biochem.*, 224 (1995) 250-255.
- 1891 Ogawa, W. and Roth, R.A.: Characterization of a protein which binds phosphatidylinositol 3,4,5-trisphosphate and 4,5-bisphosphate. *Biochim. Biophys. Acta*, 1224 (1994) 533-540.
- For additional information see C.A.:  
121 (1994) 247365s, 253911h.
- See also 1597, 1808.
- 19c. *Proteins synthesized by genetic manipulation, monoclonal antibodies*
- 1892 De Collongue-Poyet, B., Vidal-Madjar, C., Sebille, B. and Unger, K.K.: Study of conformational effects of recombinant interferon  $\gamma$  adsorbed on a non-porous reversed-phase silica support. *J. Chromatogr. B*, 664 (1995) 155-161.
- 1893 Freitag, R. and Reif, O.-W.: Multi-dimensional affinity membrane chromatography in down-stream processing of antithrombin III. *Spec. Publ. - R. Soc. Chem.*, 158(Separations for Biotechnology 3) (1994) 173-178; C.A., 121 (1994) 296298f.
- 1894 Joyce, J.G., Cook, J.C., Przysiecki, C.T. and Lehman, E.D.: Chromatographic separation of low-molecular-mass recombinant proteins and peptides on Superdex 30 prep grade. *J. Chromatogr. B*, 662 (1994) 325-334.
- 1895 Jungbauer, A. and Boschetti, E.: Manufacture of recombinant proteins with safe and validated chromatographic sorbents. *J. Chromatogr. B*, 662 (1994) 143-179 - a review with 111 refs.
- 1896 Nishihira, J., Kuriyama, T., Sakai, M., Nishi, S., Ohki, S.-y. and Hikichi, K.: The structure and physicochemical properties of rat liver macrophage migration inhibitory factor. *Biochim. Biophys. Acta*, 1247 (1995) 159-162.
- 1897 Phillips, T.M.: Immunoaffinity measurement of recombinant granulocyte colony stimulating factor in patients with chemotherapy-induced neutropenia. *J. Chromatogr. B*, 662 (1994) 307-313.
- 1898 Rapoport, E.M., Zhigis, L.S., Vlasova, E.V., Piskarev, V.E., Khraltsova, L.S., Vorozhaikina, M.M. and Zubov, V.P.: (Use of thiophilic adsorption chromatography for recovery of monoclonal antibodies to group-specific antigen H (type I)). *Zh. Fiz. Khim.*, 68 (1994) 1809-1815; C.A., 122 (1995) 7343k.

- 1899 Zerovnik, E., Jerala, R., Poklar, N., Kroon-Zitko, L. and Turk, V.: Compactness of the molten globule in comparison to unfolded states as observed by size-exclusion chromatography. *Biochim. Biophys. Acta*, 1209 (1994) 140-143.

For additional information see C.A.:  
121 (1994) 253734c.

See also 1780, 1791, 1817, 1857, 1913, 1968, 1970, 1974, 2017, 2052, 2061.

#### 19d. Microbial and plant proteins

- 1900 De Noni, I., de Bernardi, G. and Pellegrino, L.: Detection of common-wheat (*Triticum aestivum*) flour in durum-wheat (*Triticum durum*) semolina by reverse-phase high-performance liquid chromatography (RP-HPLC) of specific albumins. *Food Chem.*, 51 (1994) 325-329; C.A., 121 (1994) 229157w.
- 1901 Dunaevsky, Ya.E., Pavliukova, E.B., Beliakova, G.A. and Belozersky, M.A.: (Anionic trypsin inhibitors from dry buckwheat seeds: isolation, specificity of action and effect on micromycetes growth). *Biokhimiya (Moscow)*, 59 (1994) 990-996.
- 1902 Friis, S., Anthonsen, D., Noren, O. and Sjöström, H.: Gamma-type gliadins cause secretion of prostaglandin E<sub>2</sub> in patients with coeliac disease. *Clin. Chim. Acta*, 231 (1994) 173-183.
- 1903 Gühnemann-Schäfer, K., Engeland, K., Linder, D. and Kindl, H.: Evidence for domain structures of the trifunctional protein and the tetrafunctional protein acting in glyoxysomal fatty acid  $\beta$ -oxidation. *Eur. J. Biochem.*, 226 (1994) 909-915.
- 1904 Kiser, K.B., Arnaud, P. and Schmidt, M.G.: Rapid purification of native SecA from *Escherichia coli*: development of a new affinity chromatography procedure. *Curr. Microbiol.*, 29 (1994) 323-329; C.A., 122 (1995) 4634v.
- 1905 Moutete, H.F., Olszewski, A., Gastin, I., Namour, F., Moneret-Vautrin, D.A. and Guéant, J.L.: Purification of allergenic proteins from peanut for preparation of the reactive solid phase of a specific IgE radioimmunoassay. *J. Chromatogr. B*, 664 (1995) 211-217.
- 1906 Pouillet, P., Crechet, J.-B., Bernardi, A. and Parmeggiani, A.: Properties of the catalytic domain of Sdc25p, a yeast GDP/GTP exchange factor of Ras proteins. Complexation with wild-type Ras2p, [S24N]Ras2p and [R80D, N81D]Ras2p. *Eur. J. Biochem.*, 227 (1995) 537-544.
- 1907 Rocher, A., Soriano, F., Molina, E., González-Limas, G. and Méndez, E.: Characterization of distinct  $\alpha$ - and  $\gamma$ -type gliadins and low molecular weight components from wheat endosperm as coeliac immunoreactive proteins. *Biochim. Biophys. Acta*, 1247 (1995) 143-148.
- 1908 Santucci, D.M., Haas, B. and Smarrelli, J., Jr.: Regulation of the inducible soybean nitrate reductase isoform in mutants lacking constitutive isoform(s). *Biochim. Biophys. Acta*, 1247 (1995) 46-50.
- 1909 Senyuk, V.I., Yattara, H.B. and Vaintraub, I.A.: Purification of some soybean proteins by hydrophobic chromatography. *Nahrung*, 38 (1994) 369-372; C.A., 121 (1994) 275711v.
- 1910 Sharman, G.J., Williams, D.H., Ewing, D.F. and Ratledge, C.: Isolation, purification and structure of exochelin M5, the extracellular siderophore from *Mycobacterium smegmatis*. *Biochem. J.*, 305 (1995) 187-196.

- 1911 Terabe, M., Kojima, S., Taguchi, S., Momose, H. and Miura, K.-i.: Three novel subtilisin-trypsin inhibitors from *Streptomyces*: primary structures and inhibitory properties. *J. Biochem. (Tokyo)*, 116 (1994) 1156-1163.

See also 1951, 1972, 2055.

#### 19e. Proteins of blood, serum and blood cells

- 1912 Borkow, G., Gutierrez, J.M. and Ovadia, M.: A potent antihemorrhagin in the serum of the non-poisonous water snake *Matrix tessellata*: isolation, characterization and mechanism of neutralization. *Biochim. Biophys. Acta*, 1201 (1994) 482-490.
- 1913 Büntemeyer, H., Tebbe, H., Lüttemeyer, D. and Lehmann, J.: Rapid high-performance liquid chromatographic quantification of recombinant human antithrombin III during production and purification. *J. Chromatogr. B*, 662 (1994) 209-216.
- 1914 Burnouf, T.: Chromatography in plasma fractionation: benefits and future trends. *J. Chromatogr. B*, 664 (1995) 3-15 - a review with 52 refs.
- 1915 Chang, J.Y.C. and Chang, H.M.: (Purification of plasminogen from swine plasma by affinity chromatography). *Shin Kexue (Taipei)*, 21 (1994) 14-20; C.A., 121 (1994) 295717e.
- 1916 Cullen, P.J., Dawson, A.P. and Irvine, R.F.: Purification and characterization of an Ins(1,3,4,5)P<sub>4</sub> binding protein from pig platelets: possible identification of a novel non-neuronal Ins(1,3,4,5)P<sub>4</sub> receptor. *Biochem. J.*, 305 (1995) 139-143.
- 1917 Dahri, L., Boisson-Vidal, C., Regnault, V., Muller, D., Sultan, Y. and Stoltz, J.F.: Synthetic sorbents for removal of factor VIII inhibitors from haemophilic A plasma. *J. Chromatogr. B*, 664 (1995) 47-54.
- 1918 Era, S., Kuwata, K., Imai, H., Nakamura, K., Hayashi, T. and Sogami, M.: Age-related change in redox state of human serum albumin. *Biochim. Biophys. Acta*, 1247 (1995) 12-16.
- 1919 Finger, U.B., Thömmes, J., Kinzelt, D. and Kula, M.-R.: Application of thiophilic membranes for the purification of monoclonal antibodies from cell culture media. *J. Chromatogr. B*, 664 (1995) 69-78.
- 1920 Haung, Y.C. and Lee, T.Y.: Purification of immunoglobulin G in the presence of albumin and transferrin by zeolites. In: *Off. Proc. Comb. Conf., 6th Conf. Asia Pac. Confed. Chem. Eng., 21st Australas. Chem. Eng. Conf.*, Inst. Eng. Aust., Barton, 1, 1993, pp. 147/1-152/1; C.A., 122 (1995) 16938k.
- 1921 Josic, D., Schwinn, H., Stadler, M. and Strancar, A.: Purification of factor VIII and von Willebrand factor from human plasma by anion-exchange chromatography. *J. Chromatogr. B*, 662 (1994) 181-190.
- 1922 Katoh, S. and Terashima, M.: Purification of secreted proteins by immunoaffinity chromatography with cross-reactive and anti-peptide antibodies. *Prog. Biotechnol.*, 9(CEB6: Proceedings of the 5th European Congress on Biotechnology, 1993, Pt. 1) (1994) 539-542; C.A., 121 (1994) 225371p.
- 1923 Lutomski, D., Joubert-Caron, R., Bourin, P., Bladier, D. and Caron, M.: Use of thiophilic adsorption in the purification of biotinylated Fab fragments. *J. Chromatogr. B*, 664 (1995) 79-82.
- 1924 Matveevskaya, N.S., Alyoshkin, V.A. and Rozina, M.N.: Isolation of the C3 complement component and its C3d subunit from IV-1 fraction of Cohn's fractionation of human plasma. *J. Chromatogr. B*, 664 (1995) 261-266.

- 1925 Müller, A.M.F., Hallier, E., Westphal, G., Schröder, K.R. and Bolt, H.M.: Determination of methylated globin and albumin for bio-monitoring of exposure to methylating agents using HPLC with precolumn fluorescent derivatization. *Fresenius J. Anal. Chem.*, 350 (1994) 712-715.
  - 1926 Nikolic, J.A. and Jankovic, M.: Purification of human alpha-fetoprotein. *J. Serb. Chem. Soc.*, 59 (1994) 525-530; *C.A.*, 121 (1994) 250076d.
  - 1927 Oxvig, C., Sand, O., Kristensen, T., Kristensen, L. and Sottrup-Jensen, L.: Isolation and characterization of circulating complex between human pregnancy-associated plasma protein-A and proform of eosinophil major basic protein. *Biochim. Biophys. Acta*, 1201 (1994) 415-423.
  - 1928 Palmieri, G., Cassani, G. and Fassina, G.: Peptide immobilization on calcium alginate beads: applications to antibody purification and assay. *J. Chromatogr. B*, 664 (1995) 127-135.
  - 1929 Schwarz, A., Kohen, F. and Wilchek, M.: Novel heterocyclic ligands for the thiophilic purification of antibodies. *J. Chromatogr. B*, 664 (1995) 83-88.
  - 1930 Soedjanaatmadja, U.M.S., Hofsteenge, J., Jeronimus-Stratingh, C.M., Bruins, A.P. and Beintema, J.J.: Demonstration by mass spectrometry that pseudo-hevein and hevein have ragged C-terminal sequences. *Biochim. Biophys. Acta*, 1209 (1994) 144-148.
  - 1931 Teh, L.-C. and Froger, M.: Direct capture of plasma factor VIII: C by ion chromatography. *Vox Sang*, 67 (1994) 8-13; *C.A.*, 122 (1995) 64081x.
  - 1932 Tokunaga, F., Goto, T., Wakabayashi, S. and Koide, T.: Amino acid sequence of porcine antithrombin III. *J. Biochem. (Tokyo)*, 116 (1994) 1164-1170.
  - 1933 Vallar, L., Costa, P.M.P., Teixeira, A., Pfister, M., Barrois, R., Costa, P.P. and Rivat, C.: Immunoadsorption procedure as a potential method for the specific  $\beta$ 2-microglobulin removal from plasma of patients with chronic renal failure. *J. Chromatogr. B*, 664 (1995) 97-106.
  - 1934 Van Noort, W.L., de Jong, G. and van Eijk, H.G.: Purification of isotransferrins by concanavalin A Sepharose chromatography and preparative isoelectric focusing. *Eur. J. Clin. Chem. Clin. Biochem.*, 32 (1994) 885-892.
  - 1935 Verdoliva, A., Cassani, G. and Fassina, G.: Affinity purification of polyclonal antibodies using immobilized multimeric peptides. *J. Chromatogr. B*, 664 (1995) 175-183.
- For additional information see *C.A.*:  
121 (1994) 228808x, 286573c;  
122 (1995) 8167t.
- See also 1404, 1822, 1842, 1858, 1878, 1882, 1898, 1948, 1950, 2058.
- 19f. *Structural and muscle proteins*
- 1936 Guo, X.-F., Nakaya, M. and Watabe, S.: Myosin subfragment-1 isoforms having different heavy chain structures from fast skeletal muscle of thermally acclimated carp. *J. Biochem. (Tokyo)*, 116 (1994) 728-735.
  - 1937 Hayashibara, T. and Miyanishi, T.: Binding of the amino-terminal region of myosin alkali 1 light chain to actin and its effect on actin-myosin interaction. *Biochemistry*, 33 (1994) 12821-12827.
  - 1938 Hill, P.A., Reynolds, J.J. and Meikle, M.C.: The purification and partial characterization of bone resorptive polypeptides from bovine bone matrix. *Biochim. Biophys. Acta*, 1201 (1994) 193-202.
  - 1939 Katoh, T., Tanahashi, K., Hasegawa, Y. and Morita, F.: Porcine aorta smooth-muscle myosin contains three species made of different combinations of two 17-kDa essential light-chain isoforms. *Eur. J. Biochem.*, 227 (1995) 459-465.
  - 1940 Mizuno, K. and Hayashi, T.: Peculiar effect of urea on the interaction of type I collagen with heparin on chromatography. *J. Biochem. (Tokyo)*, 116 (1994) 1257-1263.
  - 1941 Nakamura, F. and Suyama, K.: Analysis of aldosterone, an amino acid derived from aldol crosslink of elastin and collagen by high-performance liquid chromatography. *Anal. Biochem.*, 223 (1994) 21-25.
  - 1942 Ohashi, K., Nishimura, M., Terasaki, A.G. and Nakagawa, H.: A 36-kDa protein of the dense bodies of smooth muscle cells. *J. Biochem. (Tokyo)*, 116 (1994) 1354-1359.
  - 1943 Sato, K., Taira, T., Takayama, R., Ohtsuki, K. and Kawabata, M.: Improved chromatographic purification of human and bovine type V collagen sub-molecular species and their subunit chains from conventional crude preparations. Application to cell-substratum adhesion assay for human umbilical vein endothelial cell. *J. Chromatogr. B*, 663 (1995) 25-33.
  - 1944 Van Hoof, C., Cayla, X., Bosch, M., Merlevede, W. and Goris, J.: The phosphotyrosyl phosphatase activation of protein phosphatase 2A. A novel purification method, immunological and enzymic characterization. *Eur. J. Biochem.*, 226 (1994) 899-907.
- See also 1313.
- 19g. *Protamines, histones and other nuclear proteins*
- 1945 Berger, R.G., Hoffmann, R., Waidelich, D., Bayer, E., Ingendoh, A., Hillenkamp, F. and Zeppezauer, M.: Separation and determination of molecular masses of histone H1 subtypes from calf thymus. In: Hodges, R.S. and Smith, J.A. (Editors), *Pept.: Chem., Struct. Biol., Proc. Am. Pept. Symp.*, 13th 1993, ESCOM, Leiden, 1994, pp. 228-229; *C.A.*, 121 (1994) 275684p.
  - 1946 Wang, B. and Williamson, G.: Detection of a nuclear protein which binds specifically to the antioxidant responsive element (ARE) of the human NAD(P)H:quinone oxidoreductase gene. *Biochim. Biophys. Acta*, 1219 (1994) 645-652.
- 19h. *Chromoproteins and metalloproteins*
- 1947 Calvo, A. and Batistaviera, F.: Isolation of lactoferrin by immobilized metal ion affinity chromatography (IMAC). *Biochem. Edic.*, 22 (1994) 50-52; *C.A.*, 121 (1994) 254778g.
  - 1948 Chiancone, E., Gattoni, M., Fronticelli, C., Termine, A., Silvestri, L. and Sarti, P.: Preparative and analytical applications of immobilized haemoglobin. *J. Chromatogr. B*, 664 (1995) 89-95.
  - 1949 Eckerbom, S., Bergqvist, Y. and Jeppsson, J.-O.: Improved method for analysis of glycated hemoglobin by ion exchange chromatography. *Ann. Clin. Biochem.*, 31 (1994) 355-360; *C.A.*, 121 (1994) 250070x.

- 1950 Itälä, L., Seppä, K., Turpeinen, U. and Sillanaukee, P.: Separation of hemoglobin acetaldehyde adducts by high-performance liquid chromatography - cation-exchange chromatography. *Anal. Biochem.*, 224 (1995) 323-329.
- 1951 Moore, G.R., Kadir, F.H.A., Al-Massad, F.K., le Brun, N.E., Thomson, A.J., Greenwood, C., Keen, J.N. and Findlay, J.B.C.: Structural heterogeneity of *Pseudomonas aeruginosa* bacterioferritin. *Biochem. J.*, 304 (1994) 493-497.
- 1952 O'Donnell, J.K., Birch, P., Parsons, C.T., White, S.P., Okabe, J., Martin, M.J., Adams, C., Sundarapandian, K., Manjula, B.N., Acharya, A.S. et al.: Influence of the chemical nature of side chain at  $\beta 108$  of hemoglobin A on the modulation of the oxygen affinity by chloride ions. Low oxygen affinity variants of human hemoglobin expressed in transgenic pigs: hemoglobins Presbyterian and Yoshizuka. *J. Biol. Chem.*, 269 (1994) 27692-27699.
- 1953 Williams, J.P., Jo, H., Sacks, D.B., Crimmins, D.L., Thoma, R.S., Hunnicutt, R.E., Radding, W., Sharma, R.K. and McDonald, J.M.: Tyrosine-phosphorylated calmodulin has reduced biological activity. *Arch. Biochem. Biophys.*, 315 (1994) 119-126.
- 1954 Xie, J., Aguilar, M.-I. and Hearn, M.T.W.: High-performance liquid chromatography of amino acids, peptides and proteins. CXXV. Adsorption of horse heart cytochrome c onto a tentacle-type cation exchanger. *J. Chromatogr. A*, 691 (1995) 263-271.
- See also 1934, 2049.
- 19i. *Proteins of glands, gland products, various zymogens (incl. milk proteins)*
- 1955 Bergmann, J.E. and Guelzow, H.J.: (Separation of salivary proteins by high-pressure liquid chromatography). *Dtsch. Zahnärztl. Z.*, 48 (1993) 636-638; C.A., 121 (1994) 296269x.
- 1956 Kuo, K.-W., Chang, L.-S., Lin, P.-M., Leaber, R.J. and Chang, C.-C.: Immunological neutralization of cobrotoxin by its homologous precipitin and non-precipitin antibodies. *J. Biochem. (Tokyo)*, 116 (1994) 1227-1232.
- 1957 Madjisav, M., Armstrong, D.T. and Seamark, R.F.: Purification of a cell-cell adhesion regulator from porcine seminal vesicle fluid. *Biochem. Biophys. Res. Commun.*, 205 (1994) 1206-1216.
- 1958 Omori-Satoh, T., Yamakawa, Y., Nagaoka, Y. and Mebs, D.: Hemorrhagic principles in the venom of *Bitis arietans*, a viperous snake. I. Purification and characterization. *Biochim. Biophys. Acta*, 1246 (1995) 61-66.
- 1959 Rhee, S.K.: Purification and characterization of a major gap junction channel-forming protein in rat liver. *Korean Biochem. J.*, 27 (1994) 335-341; C.A., 121 (1994) 225391v.
- 1960 Sun, Y., Zhu, J., Gu, Y., Zhong, E., Liu, Y., Zhang, X., Gao, Y. and Cheng, L.: (Purification and biological activity assay of hepatic chalone extracted from rat liver). *Jieyou Xuebao*, 25 (1994) 308-312; C.A., 121 (1994) 275715z.
- 1961 Syvaöja, E.-L. and Korhonen, H.: Determination of colostral immunoglobulins by gel filtration chromatography. *Int. Dairy Fed. Spec. Issue*, 9404 (Indigenous Antimicrobial Agents of Milk: Recent Developments) (1994) 216-219; C.A., 122 (1995) 4626u.
- 1962 Valdivia, H.H., Martin, B.M., Ramirez, A.N., Fletcher, P.L. and Possani, L.D.: Isolation and pharmacological characterization of four novel  $\text{Na}^+$  channel-blocking toxins from the scorpion *Centruroides noxius* Hoffmann. *J. Biochem. (Tokyo)*, 116 (1994) 1383-1391.
- 1963 Wilde, C.J., Addey, C.V.P., Boddy, L.M. and Peaker, M.: Autocrine regulation of milk secretion by a protein in milk. *Biochem. J.*, 305 (1995) 51-58.
- See also 1824, 1896, 1984.
- 19j. *Proteins of brain, cerebrospinal fluid and eye*
- 1964 Kieselbach, T., Irrgang, K.-D. and Rüppel, H.: A segment corresponding to amino acids Val170-Arg182 of bovine arrestin is capable of binding to phosphorylated rhodopsin. *Eur. J. Biochem.*, 226 (1994) 87-97.
- See also 1834.
- 19k. *Proteins of neoplastic tissue and transformed cells*
- 1965 Ponder, R.D., Stafford, C.T., Kiefer, C.R., Ford, J.L., Thompson, W.O. and Hoffman, D.R.: Development of an enzyme-linked immunosorbent assay for measurement of fire ant venom-specific IgE. *Ann. Allergy*, 72 (1994) 329-332; C.A., 121 (1994) 278341k.
- 19l. *Specific binding and receptor proteins*
- 1966 Antonijczuk, K., Kroftova, O.S., Varghese, A.H., Antonijczuk, A., Henjum, D.C., Korza, G., Ozols, J. and Sunderman, F.W., Jr.: The 40 kDa  $^{63}\text{Ni}^{2+}$ -binding protein (pNiXc) on Western blots of *Xenopus laevis* oocytes and embryos is the monomer of fructose-1,6-bisphosphate aldolase A. *Biochim. Biophys. Acta*, 1247 (1995) 81-89.
- 1967 Higazi, A.A.-R. and Barghouti, I.I.: Regulation of neutrophil activation by oleic acid. *Biochim. Biophys. Acta*, 1201 (1994) 442-446.
- 1968 Josic, D., Lim, Y.-P., Strancar, A. and Reutter, W.: Application of high-performance membrane chromatography for separation of annexins from the plasma membranes of liver and isolation of monospecific polyclonal antibodies. *J. Chromatogr. B*, 662 (1994) 217-226.
- 1969 Kodentsova, V.M., Vrzhesinskaya, O.A., Risnik, V.V., Sokol'nikov, A.A. and Spirichev, V.B.: (Isolation of a riboflavin-binding apoprotein from egg white and its use for riboflavin detection in biological objects). *Prikl. Biokhim. Mikrobiol.*, 30 (1994) 603-609; C.A., 122 (1995) 4635w.
- 1970 Koppel, R., Litvak, M. and Solomon, B.: Affinity purification of a mannose-binding protein, a sensitive tool in the diagnostics of IgM, via site-directed phosphorylated mannan bound to alumina. *J. Chromatogr. B*, 662 (1994) 191-196.
- 1971 Kozma, M.M., Chowrimootoo, G., Debnam, E.S., Epstein, O. and Srai, S.K.S.: Developmental changes in mucosal iron binding proteins in the guinea pig. Expression of transferrin, H and L ferritin and binding of iron to a low molecular weight protein. *Biochim. Biophys. Acta*, 1201 (1994) 229-234.

- 1972 Neumann, G.M., Condron, R., Thomas, I. and Polya, G.M.: Purification and sequencing of a family of wheat lipid transfer protein homologues phosphorylated by plant calcium-dependent protein kinase. *Biochim. Biophys. Acta*, 1209 (1994) 183-190.
- 1973 Peracchia, F., Tamburro, A., Zanni, M. and Rotilio, D.: Effects of thrombin and thrombin peptide activating receptor (SFLLRN) on proteoglycan synthesis and distribution in human endothelial cells. *Biochem. Biophys. Res. Commun.*, 205 (1994) 1625-1631.
- 1974 Santes, C., Galinha, A., Bouchard, C., Mazieres, N., Spagnoli, R. and Fridman, W.H.: Recombinant soluble Fcγ receptors: production, purification and biological activities. *J. Chromatogr. B*, 662 (1994) 197-207.
- 1975 White, T.K. and Schnaar, R.L.: Solubilization of a membrane-associated protein from rat nervous system tissues which binds anionic glycolipids and phospholipids. *Biochim. Biophys. Acta*, 1196 (1994) 218-226.
- 1976 Yohn, J.J., Smith, C., Stevens, T., Morelli, J.G., Shurnas, L.R., Walchak, S.J., Hoffman, T.A., Kelley, K.K., Escobedo-Morse, A., Yanagisawa, M. et. al.: Autoregulation of endothelin-1 secretion by cultured human keratinocytes via the endothelin B receptor. *Biochim. Biophys. Acta*, 1224 (1994) 454-458.
- For additional information see C.A.:  
121 (1994) 250087h.
- See also 1476, 1592, 1820, 1823, 1826, 1839, 1845, 1891, 1916, 1964.
- 19m. Urinary proteins**
- 1977 Baer, J.C. and Hjelm, M.: Analysis of urinary proteins by liquid chromatography. *Ann. Clin. Biochem.*, 31 (1994) 315-326; C.A., 121 (1994) 249945e.
- 1978 Büeler, M.R., Wiederkehr, F. and Vonderschmitt, D.J.: Electrophoretic, chromatographic and immunological studies of human urinary proteins. *Electrophoresis (Weinheim)*, 16 (1995) 124-134.
- 1979 Keely, E.J. and Faiman, C.: Measurement of human urinary prolactin as a noninvasive study tool. *Clin. Chem. (Washington)*, 40 (1994) 2017-2021.
- 1980 Ruhn, P.F., Taylor, J.D. and Hage, D.S.: Determination of urinary albumin using high-performance immunoaffinity chromatography and flow injection analysis. *Anal. Chem.*, 66 (1994) 4265-4271.
- See also 1833.
- 19n. Other proteins (incl. proteinous inhibitors of enzymic activity)**
- 1981 Corradi da Silva, M.L., Tamura, T. and Rice, K.G.: Derivatization and purification of bisecting tyrosinamide oligosaccharides from ovotransferrin. *Arch. Biochem. Biophys.*, 315 (1994) 460-466.
- 1982 Libera, L.D., Massimino, M.L., Arslan, P., Beltrame, M. and Cantini, M.: Analysis of muscle cell culture medium by size-exclusion chromatography. *J. Chromatogr. B*, 664 (1995) 185-191.
- 1983 Morioka, M., Muraoka, H., Yamamoto, K. and Ishikawa, H.: An endosymbiont chaperonin is a novel type of histidine protein kinase. *J. Biochem. (Tokyo)*, 116 (1994) 1075-1081.
- 1984 Perales, J., Vilella, C., Domont, G.B., Choumet, V., Saliou, B., Moussatche, H., Bon, C. and Faule, G.: Molecular structure and mechanism of action of the crotoxin inhibitor from *Crotalus durissus terrificus* serum. *Eur. J. Biochem.*, 227 (1995) 19-26.
- See also 1596, 1828, 1835, 2049, 2135.
- 20. ENZYMES AND ENZYME ACTIVITY ESTIMATION**
- 20a. Oxidoreductases**
- 1985 Cendrin, F., Jouve, H.M., Gaillard, J., Thibault, P. and Zaccari, G.: Purification and properties of a halophilic catalase-peroxidase from *Haloarcula marismortui*. *Biochim. Biophys. Acta*, 1209 (1994) 1-9.
- 1986 Chae, H.Z., Chung, S.J. and Rhee, S.G.: Thioredoxin-dependent peroxide reductase from yeast. *J. Biol. Chem.*, 269 (1994) 27670-27678.
- 1987 Créminon, C., Habib, A., Maclouf, J., Pradelles, P., Grassi, J. and Frobert, Y.: Differential measurement of constitutive (COX-1) and inducible (COX-2) cyclooxygenase expression in human umbilical vein endothelial cells using specific immunometric enzyme immunoassays. *Biochim. Biophys. Acta*, 1254 (1995) 341-348.
- 1988 Del Bello, B., Maellaro, E., Sugherini, L., Santucci, A., Comporti, M. and Casini, A.F.: Purification of NADPH-dependent dehydroascorbate reductase from rat liver and its identification with 3α-hydroxysteroid dehydrogenase. *Biochem. J.*, 304 (1994) 385-390.
- 1989 Dowd, V. and Yon, R.J.: Further examination of a "concerted cluster" model of multivalent affinity. Heterogeneous adsorption of lactate dehydrogenase to Cibacron Blue immobilised on cellulose. *J. Chromatogr. A*, 693 (1995) 15-21.
- 1990 Johnson, K.A., Morrow, C.J., Knight, G.D. and Scallen, T.J.: *In vivo* formation of 25-hydroxycholesterol from endogenous cholesterol after a single meal, dietary cholesterol challenge. *J. Lipid Res.*, 35 (1994) 2241-2253.
- 1991 Maltseva, O.V., Solyanikova, I.P. and Golovleva, L.A.: Chlorocatechol 1,2-dioxygenase from *Rhodococcus erythropolis* 1CP. Kinetic and immunochemical comparison with analogous enzymes from Gram-negative strains. *Eur. J. Biochem.*, 226 (1994) 1053-1061.
- 1992 Niemetz, R., Altenschmidt, U., Brucker, S. and Fuchs, G.: Benzoyl-coenzyme-A 3-monooxygenase, a flavin-dependent hydroxylase. Purification, some properties and its role in aerobic benzoate oxidation via gentisate in a denitrifying bacterium. *Eur. J. Biochem.*, 227 (1995) 161-168.
- 1993 Noguchi, M., Miyano, M., Matsumoto, T. and Noma, M.: Human 5-lipoxygenase associates with phosphatidylcholine liposomes and modulates LTA<sub>4</sub> synthetase activity. *Biochim. Biophys. Acta*, 1215 (1994) 300-306.

- 1994 Novikov, D.K., Vanhove, G.F., Carchon, H., Asselberghs, S., Eyssen, H.J., van Veldhoven, P.P. and Mannaerts, G.P.: Peroxisomal  $\beta$ -oxidation. Purification of four novel 3-hydroxyacyl-CoA dehydrogenases from rat liver peroxisomes. *J. Biol. Chem.*, 269 (1994) 27125-27135.
- 1995 Padilla, C.A., Martinez-Galisteo, E., Barcena, J.A., Spyrou, G. and Holmgren, A.: Purification from placenta, amino acid sequence, structure comparisons and cDNA cloning of human glutaredoxin. *Eur. J. Biochem.*, 227 (1995) 27-34.
- 1996 Percival, M.D., Ouellet, M., Vincent, C.J., Yergey, J.A., Kennedy, B.P. and O'Neill, G.P.: Purification and characterization of recombinant human cyclooxygenase-2. *Arch. Biochem. Biophys.*, 315 (1994) 111-118.
- 1997 Regdel, D., Schewe, T. and Kuehn, H.: (Tomato fruit lipoxygenase: isolation, partial purification, characterization and interaction with biological membranes). *Biokhimiya (Moscow)*, 59 (1994) 788-796.
- 1998 Sailesh, S., Kumar, Y.V.K., Prasad, M. and Reddanna, P.: Sheep uterus dual lipoxygenase in the synthesis of 14,15-leukotrienes. *Arch. Biochem. Biophys.*, 315 (1994) 362-368.
- 1999 Sarkar, M.A. and Jackson, B.J.: Theophylline-N-demethylations as probes for P4501A1 and P4501A2. *Drug Metab. Disp.*, 22 (1994) 827-834.
- 2000 Soulimane, T. and Buse, G.: Integral cytochrome-c oxidase. Preparation and progress towards a three-dimensional crystallization. *Eur. J. Biochem.*, 227 (1995) 588-595.
- 2001 Tao, Y. and Chen, K.Y.: Purification of deoxyhypusine synthase from *Neurospora crassa* to homogeneity by substrate elution affinity chromatography. *J. Biol. Chem.*, 270 (1995) 383-386.
- 2002 Van Dijk, J., Boomsma, F., Alberts, G., Man in't Veld, A.J. and Schalekamp, M.A.D.H.: Determination of semicarbazide-sensitive amine oxidase activity in human plasma by high-performance liquid chromatography with fluorimetric detection. *J. Chromatogr. B*, 663 (1995) 43-50.
- 2003 Van Veldhoven, P.P., van Rompuy, P., Vanhooren, J.C.T. and Mannaerts, G.P.: Purification and further characterization of peroxisomal trihydroxycoprostanoyl-CoA oxidase from rat liver. *Biochem. J.*, 304 (1994) 195-200.
- 2004 Wang, X.T., Dumoulin, M.J., Befani, O., Mondovi, B. and Mateescu, M.A.: Joint chromatographic purification of bovine serum ceruloplasmin and amine oxidase. *Prep. Biochem.*, 24 (1994) 237-250; *C.A.*, 122 (1995) 4120z.
- See also 1815, 1827, 1829, 1836, 1840, 2219.
- 20b. *Transferases (excl. E.C. 2.7.-.)*
- 2005 Aigner, A., Jäger, M., Weber, P. and Wolf, S.: A nonradioactive assay for microsomal cysteine-S-conjugate N-acetyltransferase activity by high-pressure liquid chromatography. *Anal. Biochem.*, 223 (1994) 227-231.
- 2006 Ishiguro, H., Kawata, S., Yamasaki, E., Matsuda, Y., Fujii, S. and Matsuzawa, Y.: High-performance liquid chromatographic assay for farnesyl-protein transferase activity with dabsylated peptide. *J. Chromatogr. B*, 663 (1995) 35-42.
- 2007 Langkamp-Henken, B., Geller, A.M., LeGros, H.L., Jr., Price, J.O., de la Rosa, J. and Kotb, M.: Characterization of distinct forms of methionine adenosyltransferase in nucleated, and mature human erythrocytes and erythroleukemic cells. *Biochim. Biophys. Acta*, 1201 (1994) 397-404.
- 2008 Liebergesell, M., Sonomoto, K., Madkour, M., Mayer, F. and Steinbüchel, A.: Purification and characterization of the poly(hydroxyalkanoic acid) synthase from *Chromatium vinosum* and localization of the enzyme at the surface of poly(hydroxyalkanoic acid) granules. *Eur. J. Biochem.*, 226 (1994) 71-80.
- 2009 Mack, M., Bendrat, K., Zelder, O., Eckel, E., Linder, D. and Buckel, W.: Location of the two genes encoding glutaconate coenzyme A-transferase at the beginning of the hydroxyglutamate operon in *Acidaminococcus fermentans*. *Eur. J. Biochem.*, 226 (1994) 41-51.
- 2010 Mohsen, A.-W.A., Aull, J.L., Payne, D.M. and Daron, H.H.: Ligand-induced conformational changes of thymidylate synthase detected by limited proteolysis. *Biochemistry*, 34 (1995) 1669-1677.
- 2011 Pace, M., Agnellini, D., Gardana, C., Mauri, P.L. and Pietta, P.G.: High-performance liquid chromatographic assay of glycosyltransferases using flavonoids as substrate. *J. Chromatogr. A*, 691 (1995) 331-336.
- 2012 Reenilä, I., Tuomainen, P. and Männistö, P.T.: Improved assay of reaction products to quantitate catechol-O-methyltransferase activity by high-performance liquid chromatography with electrochemical detection. *J. Chromatogr. B*, 663 (1995) 137-142.
- 2013 Vancura, A. and Haldar, D.: Purification and characterization of glycerophosphate acyltransferase from rat liver mitochondria. *J. Biol. Chem.*, 269 (1994) 272009-27215.
- See also 1846.
- 20c. *Transferases transferring phosphorus containing groups (E.C. 2.7.-.)*
- 2014 Allen, B.G., Andrea, J.E. and Walsh, M.P.: Identification and characterization of protein kinase C $\xi$ -immunoreactive proteins. *J. Biol. Chem.*, 269 (1994) 29288-29298.
- 2015 Clasper, S., Chelvarajan, R.E.L., Easterby, J.S. and Powls, R.: Isolation of multiple dimeric forms of phosphoribulokinase from an alga and a higher plant. *Biochim. Biophys. Acta*, 1209 (1994) 101-106.
- 2016 Huot, J., Lambert, H., Lavoie, J.N., Guimond, A., Houle, F. and Landry, J.: Characterization of 45-kDa/54-kDa HSP27 kinase, a stress-sensitive kinase which may activate the phosphorylation-dependent protective function of mammalian 27-kDa heat-shock protein HSP27. *Eur. J. Biochem.*, 227 (1995) 416-427.
- 2017 Koller, G., Graumann, K., Kramer, W., Sara, M. and Jungbauer, A.: Laboratory-scale production and purification of recombinant HIV-1 reverse transcriptase. *J. Chromatogr. B*, 664 (1995) 107-118.
- 2018 Martinez-Costa, O.H., Estevez, A.M., Sanchez, V. and Aragon, J.J.: Purification and properties of phosphofructokinase from *Dictyostelium discoideum*. *Eur. J. Biochem.*, 226 (1994) 1007-1017.
- 2019 Rempel, A., Bannasch, P. and Mayer, D.: Differences in expression and intracellular distribution of hexokinase isoenzymes in rat liver cells of different transformation stages. *Biochim. Biophys. Acta*, 1219 (1994) 660-668.
- 2020 Sharova, N.P., Eliseeva, E.D. and Mikhailov, V.S.: (The presence of DNA polymerase  $\zeta$ -type in the eggs of the teleost fish *Misgurnus fossilis* L.) *Biokhimiya (Moscow)*, 59 (1994) 1067-1075.

- 2021 Shi, G.-Y., Chang, B.-I., Chen, S.-M., Wu, D.-H. and Wu, H.-L.: Function of streptokinase fragments in plasminogen activation. *Biochem. J.*, 304 (1994) 235-241.
  - 2022 Shiokawa, D., Ohyama, H., Yamada, T., Takahashi, K. and Tanuma, S.-i.: Identification of an endonuclease responsible for apoptosis in rat thymocytes. *Eur. J. Biochem.*, 226 (1994) 23-30.
  - 2023 Simanov, A.L.: High-performance liquid chromatographic analysis of low-molecular-mass products synthesized by polynucleotide phosphorylase in polymerization reaction. *J. Chromatogr. A*, 685 (1994) 39-44.
  - 2024 Song, W. and Jackowski, S.: Kinetics and regulation of pantothenate kinase from *Escherichia coli*. *J. Biol. Chem.*, 269 (1994) 27051-27058.
  - 2025 Uchida, T.: Immunologically and enzymatically distinct rat choline kinase isozymes. *J. Biochem. (Tokyo)*, 116 (1994) 1241-1250.
  - 2026 Vermeij, P., Detmers, F.J.M., Broers, F.J.M., Keltjens, J.T. and van der Drift, C.: Purification and characterization of coenzyme F<sub>390</sub> synthetase from *Methanobacterium thermoautotrophicum* (strain ΔH). *Eur. J. Biochem.*, 226 (1994) 185-191.
  - 2027 Zang, R., Müller, H.-J., Kielbassa, K., Marks, F. and Gschwendt, M.: Partial purification of a type η protein kinase C from murine brain: separation from other protein kinase C isoenzymes and characterization. *Biochem. J.*, 304 (1994) 641-647.
  - 2028 Zhang, J., Sanchez, R.J., Wang, S., Guarnaccia, C., Tossi, A., Zahariev, S. and Poonor, S.: Substrate specificity of CDC2 kinase from human HeLa cells as determined with synthetic peptides and molecular modeling. *Arch. Biochem. Biophys.*, 315 (1994) 415-424.
- See also 1816, 1831, 1841, 1983.
- 20d. *Hydrolases, acting on ester bonds (E.C. 3.1.-.-)*
- 2029 Amici, A., Emanuelli, M., Ferretti, E., Raffaelli, N., Ruggieri, S. and Magni, G.: Homogenous pyrimidine nucleotidase from human erythrocytes: enzymic and molecular properties. *Biochem. J.*, 304 (1994) 987-992.
  - 2030 Caruccio, N. and Ross, J.: Purification of a human polyribosome-associated 3' to 5' exoribonuclease. *J. Biol. Chem.*, 269 (1994) 31814-31821.
  - 2031 Guo, W., Shang, Z., Yu, Y. and Zhou, L.: Membrane affinity chromatography of alkaline phosphatase. *J. Chromatogr. A*, 685 (1994) 344-348.
  - 2032 Marcinkeviciene, L.Yu., Bachmatova, I.V., Brazenas, G.R., Baratova, L.A. and Revina, L.P.: (Purification and properties of cholesterol esterase from *Pseudomonas mendocina* 3121). *Biokhimiya (Moscow)*, 59 (1994) 648-655.
  - 2033 Masuno, H., Nakabayashi, H., Kobayashi, J., Saito, Y. and Okuda, H.: Reduced dimerization of lipoprotein lipase in post-heparin plasma of a patient with hyperchylomicronemia. *Biochim. Biophys. Acta*, 1254 (1995) 30-36.
  - 2034 Matts, P.J., White, G.F. and Payne, W.J.: Purification and characterization of the short-chain alkylsulphatase of coryneform B1a. *Biochem. J.*, 304 (1994) 937-943.
  - 2035 Morgan, E.W., Yan, B., Greenway, D., Petersen, D.R. and Parkinson, A.: Purification and characterization of two rat liver microsomal carboxylesterases (hydrolase A and B). *Arch. Biochem. Biophys.*, 315 (1994) 495-512.
  - 2036 Okamura, S. and Yamashita, S.: Purification and characterization of phosphatidylcholine phospholipase D from pig lung. *J. Biol. Chem.*, 269 (1994) 31207-31213.
  - 2037 Sahsah, Y., Thi, A.T.P., Roy-Macauley, H., d'Arcy-Lameta, A., Repellin, A. and Zuily-Fodil, Y.: Purification and characterization of a soluble lipolytic acylhydrolase from Cowpea (*Vigna unguiculata* L.) leaves. *Biochim. Biophys. Acta*, 1215 (1994) 66-73.
  - 2038 Shirazi, A., Iizuka, K., Fadden, P., Mosse, C., Somlyo, A.P., Somlyo, A.V. and Haystead, T.A.J.: Purification and characterization of the mammalian myosin light chain phosphatase holoenzyme. The differential effects of the holoenzyme and its subunits on smooth muscle. *J. Biol. Chem.*, 269 (1994) 31598-31606.
  - 2039 Takasaki, Y.: Two forms of restriction enzyme HindIII. *J. Biochem. (Tokyo)*, 116 (1994) 1281-1286.
  - 2040 Wang, R., Dodia, C.R., Jain, M.K. and Fisher, A.B.: Purification and characterization of a calcium-independent acidic phospholipase A<sub>2</sub> from rat lung. *Biochem. J.*, 304 (1994) 131-137.
- For additional information see C.A.:  
121 (1994) 250631f.
- See also 1944, 2022.
- 20e. *Hydrolases, acting on glycosyl compounds (E.C. 3.2.-.-)*
- 2041 Heng, M.H. and Glatz, C.E.: Charged fusions for β-galactosidase retention in anion-exchange chromatography. *J. Chromatogr. A*, 689 (1995) 227-234.
  - 2042 Itoh, Y., Hirashima, M., Yamada, H. and Imoto, T.: Colonic lysozymes of rabbit (Japanese White): recent divergence and functional conversion. *J. Biochem. (Tokyo)*, 116 (1994) 1346-1353.
  - 2043 Katoh, S. and Terashima, M.: Effective purification of α-amylases from fermentation broth by immunoaffinity chromatography. *Spec. Publ. - R. Soc. Chem.*, 158 (Separations for Biotechnology C) (1994) 228-234; C.A., 121 (1994) 295722c.
  - 2044 Mislovicova, D., Chudinova, M., Gemeiner, P. and Docolomansky, P.: Affinity chromatography of invertase on Concanavalin A-bead cellulose matrix: the case of an extraordinary strong binding glycoenzyme. *J. Chromatogr. B*, 664 (1995) 145-153.
  - 2045 Munoz, M.D., Hernández, L.M., Basco, R., Andaluz, E. and Larriba, G.: Glycosylation of yeast exoglucanase sequons in alg mutants deficient in the glucosylation steps of the lipid-linked oligosaccharide. Presence of glucotriose unit in Dol-PP-GlcNAc<sub>2</sub>Man<sub>9</sub>Glc<sub>3</sub> influences both glycosylation efficiency and selection of N-linked sites. *Biochim. Biophys. Acta*, 1201 (1994) 361-366.
  - 2046 Pritchard, D.G., Lin, B., Willingham, T.R. and Baker, J.R.: Characterization of the group B streptococcal hyaluronate lyase. *Arch. Biochem. Biophys.*, 315 (1984) 431-437.
  - 2047 Sakamoto, M., Shirane, Y., Narabayashi, I., Kimura, K., Morishita, N., Sakamoto, T. and Sakai, T.: Purification and characterization of a rhamnogalacturonase with protopectinase activity from *Trametes sanguinea*. *Eur. J. Biochem.*, 226 (1994) 285-291.

- 2048 Tulsiani, D.R.P., Skudlarek, M.D., Araki, Y. and Orgebin-Crist, M.-C.: Purification and characterization of two forms of  $\beta$ -D-galactosidase from rat epididymal luminal fluid: evidence for their role in the modification of sperm plasma membrane glycoprotein(s). *Biochem. J.*, 305 (1995) 41-50.
- 2049 Vachier, M.C., Piot, M. and Awadé, A.C.: Isolation of hen egg white lysozyme, ovotransferrin and ovalbumin, using a quaternary ammonium bound to a highly crosslinked agarose matrix. *J. Chromatogr. B*, 664 (1995) 201-210.
- For additional information see C.A.:  
121 (1994) 275502e.
- See also 1843.
- 20f. *Other hydrolases*
- 2050 Bogomolov, A.A., Danilenko, A.N. and Antonov, Yu.A.: (Precipitation of alfalfa (*Medicago sativa* L.) leaf juice enzyme and the mechanism of its effect on RuBISCO hydrolysis). *Biokhimiya (Moscow)*, 59 (1994) 689-693.
- 2051 Canales, J., Pinto, R.M., Costas, J., Hernández, M.T., Mirí, A., Bernet, D., Fernández, A. and Cameselle, J.C.: Rat liver nucleoside diphosphosugar or diphosphoalcohol pyrophosphatases different from nucleotide pyrophosphatase or phosphodiesterase I: substrate specificities of  $Mg^{2+}$ - and/or  $Mn^{2+}$ -dependent hydrolases acting on ADP-ribose. *Biochim. Biophys. Acta*, 1246 (1995) 167-177.
- 2052 Christianson, T. and Paech, C.: Peptide mapping of subtilisins as a practical tool for locating protein sequence errors during extensive protein engineering projects. *Anal. Biochem.*, 223 (1994) 119-129.
- 2053 Davis, M.T., Lee, T.D., Ronk, M. and Hefta, S.A.: Microscale immobilized protease reactor columns for peptide mapping by liquid chromatography/mass spectral analyses. *Anal. Biochem.*, 224 (1995) 235-244.
- 2054 Graham-Siegenthaler, K., Gauthier, S., Davies, P.L. and Elce, J.S.: Active recombinant rat calpain II. Bacterially produced large and small subunits associate both *in vivo* and *in vitro*. *J. Biol. Chem.*, 269 (1994) 30457-30460.
- 2055 Huang, C.-C., Huang, K.-F. and Chiou, S.-H.: Characterization of one novel venom protease with  $\beta$ -fibrinogenase activity from the Taiwan habu (*Trimeresurus mucrosquamatus*): purification and cDNA sequence analysis. *Biochem. Biophys. Res. Commun.*, 205 (1994) 1707-1715.
- 2056 Ivashchuk-Kienbaum, Yu.A., and Apell, H.-J.: Separation and characterization of  $Na^+, K^+$ -ATPase containing vesicles. *Biochim. Biophys. Acta*, 1196 (1994) 29-37.
- 2057 Jane, D.T. and Dufresne, M.J.: Expression and regulation of three lysosomal cysteine protease activities during growth of a differentiating LG rat myoblast cell line and its nonfusing variant. *Biochem. Cell Biol.*, 72 (1994) 267-274.
- 2058 Khan, M.T., Wang, K., Auland, M.E., Kable, E.P.W. and Roufogalis, B.D.: Membrane-bound high molecular mass proteinases from human erythrocytes. *Biochim. Biophys. Acta*, 1209 (1994) 215-221.
- 2059 Klabunde, T., Stahl, B., Suerbaum, H., Hahner, S., Karas, M., Hillenkamp, F., Krebs, B. and Witzel, H.: The amino acid sequence of the red kidney bean Fe(III)-Zn(II) purple acid phosphatase. Determination of the amino acid sequence by a combination of matrix-assisted laser desorption/ionization mass spectrometry and automated Edman sequencing. *Eur. J. Biochem.*, 226 (1994) 369-375.
- 2060 Li, Y.-M., Brostedt, P., Hjertén, S., Nyberg, F. and Silberring, J.: Capillary liquid chromatography-fast atom bombardment mass spectrometry using a high-resolving cation exchanger, based on a continuous chromatographic matrix. Application to studies on neuropeptide peptidases. *J. Chromatogr. B*, 664 (1995) 426-430.
- 2061 Loayza, S.L., Trikha, M., Markland, F.S., Riquelme, P. and Kuo, J.: Resolution of isoforms of natural and recombinant fibrolase, the fibrinolytic enzyme from *Agkistrodon contortrix contortrix* snake venom, and comparison of their EDTA sensitivities. *J. Chromatogr. B*, 662 (1994) 227-243.
- 2062 Muta, T., Seki, N., Takaki, Y., Hashimoto, R., Oda, T., Iwanaga, A., Tokunaga, F. and Iwanaga, S.: Purified horseshoe crab factor G. Reconstitution and characterization of the (1 $\rightarrow$ 3)- $\beta$ -D-glucan-sensitive serine protease cascade. *J. Biol. Chem.*, 270 (1995) 892-897.
- 2063 Ohkubo, I., Huang, K., Ochiai, Y., Takagaki, M. and Kani, K.: Dipeptidyl peptidase IV from porcine seminal plasma: purification, characterization, and N-terminal amino acid sequence. *J. Biochem. (Tokyo)*, 116 (1994) 1182-1186.
- 2064 Rossmann, R., Maier, T., Lottspeich, F. and Böck, A.: Characterisation of a protease from *Escherichia coli* involved in hydro-genase maturation. *Eur. J. Biochem.*, 227 (1995) 545-550.
- 2065 Shi, X.-j. and Knowles, A.F.: Prevalence of the marcurial-sensitive ectoATPase in human small cell lung carcinoma: characterization and partial purification. *Arch. Biochem. Biophys.*, 315 (1994) 177-184.
- 2066 Suen, S.-Y. and Etzel, M.R.: Sorption kinetics and breakthrough curves for pepsin and chymosin using pepstatin A affinity membranes. *J. Chromatogr. A*, 686 (1994) 179-192.
- 2067 Tsuchiya, Y., Takahashi, T., Sakurai, Y., Iwamatsu, A. and Takahashi, K.: Purification and characterization of a novel membrane-bound arginine-specific serine proteinase from porcine intestinal mucosa. *J. Biol. Chem.*, 269 (1994) 32985-32991.
- 2068 Ward, R.V., Atkinson, S.J., Reynolds, J.J. and Murphy, G.: Cell-surface-mediated activation of progelatinase A: demonstration of the involvement of the C-terminal domain of progelatinase A in cell surface binding and activation of progelatinase A by primary fibroblasts. *Biochem. J.*, 304 (1994) 263-269.
- For additional information see C.A.:  
121 (1994) 274928j.
- See also 1832, 1837, 1889.
- 20g. *Lyases*
- 2069 Choi, Y.S. and Cho, Y.D.: A new S-adenosylmethionine decarboxylase from soybean axes. *Biochim. Biophys. Acta*, 1201 (1994) 466-472.

- 2070 Harteneck, C., Klatt, P., Schmidt, K. and Mayer, B.: Expression of rat brain nitric oxide synthase in baculovirus-infected insect cells and characterization of the purified enzyme. *Biochem. J.*, 304 (1994) 683-686.
- 2071 Mankowitz, L., Staffas, L., Bakke, M. and Lund, J.: Adrenocorticotrophic-hormone-dependent regulation of a  $\mu$ -class glutathione transferase in mouse adrenocortical cells. *Biochem. J.*, 305 (1995) 111-118.
- 20h. *Isomerases*
- 2072 Ding, L., Seto, B.L., Ahmed, S.A. and Coleman, W.G., Jr.: Purification and properties of the *Escherichia coli* K-12 NAD-dependent nucleotide diphosphosugar epimerase, ADP-L-glycero-D-mannoheptose 6-epimerase. *J. Biol. Chem.*, 269 (1994) 24384-24390.
- 20i. *Ligases*
- For additional information see C.A.: 122 (1995) 4164r.
21. PURINES, PYRIMIDINES, NUCLEIC ACIDS AND THEIR CONSTITUENTS
- See 1876.
- 21a. *Purines, pyrimidines, nucleosides, nucleotides*
- 2073 Almudaris, A., Ashton, D.S., Ray, A. and Valko, K.: Trace analysis of impurities in 3'-azido-3'-deoxythymidine by reversed-phase high-performance liquid chromatography and thermospray mass spectrometry. *J. Chromatogr. A*, 689 (1995) 31-38.
- 2074 Betto, P., Popoli, P., Ricciarello, G., Caporali, M.G. and Antonini, R.: Simultaneous high-performance liquid chromatographic determination of adenosine and dopamine in rat striatal tissue with combined ultraviolet absorbance and electrochemical detection. *J. Chromatogr. B*, 662 (1994) 21-25.
- 2075 Bourque, A.J. and Cohen, A.S.: Quantitative analysis of phosphorothioate oligonucleotides in biological fluids using direct injection fast anion-exchange chromatography and capillary gel electrophoresis. *J. Chromatogr. B*, 662 (1994) 343-349.
- 2076 Brüggermann, B., Schlüter, H., Verspohl, E.J. and Zidek, W.: Rapid separation of dinucleotides with ion-pair high-performance liquid chromatography. *Fresenius J. Anal. Chem.*, 350 (1994) 719-721.
- 2077 Cong, H.N., Bertaux, O., Valencia, R., Becue, T., Fournier, T., Biou, D. and Porquet, D.: Separation and characterization of the main methylated nucleobases from nuclear, cytoplasmic and poly (A)\* RNA by high-performance liquid chromatography and mass spectrometry. *J. Chromatogr. B*, 661 (1994) 193-204.
- 2078 Dolan, M.E., Chae, M., Pegg, A.E., Mullen, J.H., Friedman, H.S. and Moschel, R.C.: Metabolism of O<sup>6</sup>-benzylguanine, an inactivator of O<sup>6</sup>-alkylguanine-DNA alkyltransferase. *Cancer Res.*, 54 (1994) 5123-5130.
- 2079 Douki, T. and Cadet, J.: Formation of cyclobutane dimers and (6-4) photoproducts upon far-UV photolysis of 5-methylcytosine-containing dinucleoside monophosphates. *Biochemistry*, 33 (1994) 11942-11950.
- 2080 Faraone-Mennella, M.R., de Lucia, F., de Maio, A., Gambacorta, A., Quesada, P., de Rosa, M., Nicolaus, B. and Farina, B.: ADP-ribosylation reactions in *Sulfolobus solfataricus*, a thermoacidophilic archaeon. *Biochim. Biophys. Acta*, 1246 (1995) 151-159.
- 2081 Fennell, T.R.: Development of methods for measuring biological markers of formaldehyde exposure. *Res. Rep.-Health Eff. Inst.*, 67 (1994) 1-26; C.A., 121 (1994) 223733j.
- 2082 Johnson, S.W., Swiggard, P.A., Handel, L.M., Brennan, J.M., Godwin, A.K., Ozols, R.F. and Hamilton, T.C.: Relationship between platinum-DNA adduct formation and removal and cisplatin cytotoxicity in cisplatin-sensitive and -resistant human ovarian cancer cells. *Cancer Res.*, 54 (1994) 5911-5916.
- 2083 Kock, R., Delvoux, B., Sigmund, M. and Greiling, H.: A comparative study of the concentrations of hypoxanthine, xanthine, uric acid and allantoin in the peripheral blood of normals and patients with acute myocardial infarction and other ischaemic diseases. *Eur. J. Clin. Chem. Clin. Biochem.*, 32 (1994) 837-842.
- 2084 Lennard, L. and Singleton, H.J.: High-performance liquid chromatographic assay of human red blood cell thiopurine methyltransferase activity. *J. Chromatogr. B*, 661 (1994) 25-33.
- 2085 Lu, T. and Gray, H.B., Jr.: Combined pH gradient and anion-exchange high-performance liquid chromatographic separation of oligodeoxyribonucleotides. *J. Chromatogr. A*, 686 (1994) 339-343.
- 2086 Murakami, A., Tamura, Y., Wada, H. and Makino, K.: Separation and characterization of diastereoisomers of antisense oligodeoxyribonucleoside phosphorothioates. *Anal. Biochem.*, 223 (1994) 285-290.
- 2087 Nithipatikorn, K., Mizumura, T. and Gross, G.J.: Determination of plasma adenosine by high-performance liquid chromatography with column switching and fluorometric detection. *Anal. Biochem.*, 223 (1994) 280-284.
- 2088 Oefner, P.J., Huber, C.G., Umlauf, F., Berti, G.-H., Stimpfl, E. and Bonn, G.K.: High-resolution liquid chromatography of fluorescent dye-labeled nucleic acids. *Anal. Biochem.*, 223 (1994) 39-46.
- 2089 Pecina-Slaus, N., Pavelic, J. and Pavelic, K.: Comparison of reverse-phase HPLC and gel electrophoretic purification of synthetic oligonucleotides. *Period. Biol.*, 96 (1994) 161-164; C.A., 121 (1994) 296539k.
- 2090 Potaman, V.N. and Soyfer, V.N.: Oligonucleotide model with non-identical complementary strands for chromatographic studies of structure-dependent photosusceptibility. *J. Chromatogr. A*, 690 (1995) 65-70.
- 2091 Ruther, J. and Baltes, W.: Analysis of purine compounds and creatinine by ion-pair high-performance liquid chromatography (HPLC) as a method for the detection of yeast extracts in commercial meat flavorings. *Z. Lebensm.-Unters. Forsch.*, 199 (1994) 307-310; C.A., 121 (1994) 299425f.
- 2092 Schuette, J.M., Cole, D.L. and Srivatsa, G.S.: Development and validation of a method for routine base composition analysis of phosphorothioate oligonucleotides. *J. Pharm. Biomed. Anal.*, 12 (1994) 1345-1353.

- 2093 Simanov, A.L., Gusakov, A.V., Simanova, N.S. and Sinitsyn, A.P.: (Analysis of the product composition in the polymerization reaction catalyzed by polynucleotide phosphorylase from *Thermus thermophilus*). *Biokhimiya (Moscow)*, 59 (1994) 726-731.
- 2094 Stout, M.L. and Ravindranath, Y.: Alkylsulphonic acid ion pairing with radial compression columns for determining plasma or cerebrospinal fluid 1- $\beta$ -D-arabinofuranosylcytosine in pediatric pharmacokinetic analysis. *J. Chromatogr. A*, 692 (1995) 59-66.
- 2095 Stratford, M.R.L. and Dennis, M.F.: Determination of adenine nucleotides by fluorescence detection using high-performance liquid chromatography and post-column derivatisation with chloroacetaldehyde. *J. Chromatogr. B*, 662 (1994) 15-20.
- 2096 Strobel, S.A., Cech, T.R., Usman, N. and Beigelman, L.: The 2,6-diaminopurine riboside•5-methylisocytidine wobble base pair: an isoenergetic substitution for the study of G•U pairs in RNA. *Biochemistry*, 33 (1994) 13824-13835.
- 2097 Terzuoli, L., Pandolfi, M., Arezzini, L., Pizzichini, M., Marinello, E. and Pagani, R.: Separation and determination of liver uric acid and allantoin. *J. Chromatogr. B*, 663 (1995) 143-147.
- 2098 Terzuoli, L., Pizzichini, M., Arezzini, L., Pandolfi, M.L., Marinello, E. and Pagani, R.: Separation and determination of allantoin, uric acid, hydantoin and urea. *J. Chromatogr. A*, 684 (1994) 350-353.
- 2099 Thoithi, G., van Schepdael, A., Herdewijn, P., Roets, E. and Hoogmartens, J.: Liquid chromatographic separation of diamino analogues of 2'- or 3'-deoxyadenosine from adenine on a poly(styrene-divinylbenzene) polymer column. *J. Chromatogr. A*, 689 (1995) 247-254.
- 2100 Vaca, C.E., Conradi, M., Sievertzon, M. and Bergman, J.: Synthesis of fluorescent derivatives of 7-methylguanine through reaction with 2-aryl-substituted malondialdehydes: analysis by HPLC with fluorescence detection. *Chem.-Biol. Interact.*, 93 (1994) 235-249; *C.A.*, 122 (1995) 4819j.
- For additional information see *C.A.*:  
121 (1994) 225390u, 250077e, 308507p.
- See also 1420, 2023, 2106, 2391, 2435.
- 21b. *Nucleic acids, RNA*
- See 1908, 2088.
- 21c. *Nucleic acids, DNA*
- 2101 Fadlallah, S., Cooper, S.K. and Fournier, M.: Assessment of N-nitrosodimethylamine DNA adducts in rat hepatocytes by high performance liquid chromatography and immunosorbent assay. *Int. J. Environ. Anal. Chem.*, 56 (1994) 165-174; *C.A.*, 121 (1994) 223747s.
- 2102 Hirabayashi, J. and Kasai, K.: Slalom chromatography: a size-dependent separation method for DNA molecules based on a hydrodynamic principle. *Mol. Interact. Biosep.*, (1993) 69-87; *C.A.*, 121 (1994) 249851w - a review with many refs.
- 2103 Hofman, Y.L., Petrillo, K.L., Greenblatt, H.C., Lehrer, R. and Payne, M.S.: Rapid scaleable chromatographic purification of nucleic acids from proteinaceous mixtures. *J. Chromatogr. A*, 692 (1995) 233-238.
- 2104 Huber, C.G., Oefner, P.J. and Bonn, G.K.: Rapid and accurate sizing of DNA fragments by ion-pair chromatography on alkylated nonporous poly(styrene-divinylbenzene) particles. *Anal. Chem.*, 67 (1995) 578-585.
- 2105 Mischiati, C., Feriotto, G., Fiorentino, D. and Gambari, R.: Chromatography in DNA radiolabeling: hands-off automation using a robotic workstation. *J. Chromatogr. B*, 664 (1995) 303-310.
- 2106 Shigenaga, M.K., Aboujaoude, E.N., Chen, Q. and Ames, B.N.: Assays of oxidative DNA damage biomarkers 8-oxo-2'-deoxyguanosine and 8-oxoguanine in nuclear DNA and biological fluids by high-performance liquid chromatography with electrochemical detection. *Methods Enzymol.*, 234(Oxygen Radicals in Biological Systems, Pt. D) (1994) 16-33; *C.A.*, 121 (1994) 294506y.
- 2107 Wainwright, N. and Boyd, S.H.: Aligned fiber diagnostic chromatography tip having a capture element in a micropipet tip assembly. *PT Int. Appl. WO 94 20,831 (Cl. G01N1/18)*, 15 Sep. 1994, US Appl. 27, 813, 08 Mar. 1993; 24 pp.; *C.A.*, 121 (1994) 296601z.
- See also 2088, 2108.
- 21e. *Structural studies on DNA and DNA mapping*
- 2108 Shealy, D.B., Lipowska, M., Lipowski, J., Narayanan, N., Sutter, S., Strekowski, L. and Patonay, G.: Synthesis, chromatographic separation, and characterization of near-infrared-labeled DNA oligomers for use in DNA sequencing. *Anal. Chem.*, 67 (1995) 247-251.
- See also 2106.
22. ALKALOIDS
- 2109 Alvarez-Fuentes, J., Fernandez-Arevalo, M., Holgado, M.A., Caraballo, I., Llera, J.M. and Rabasco, A.M.: Characterization of morphine polymeric coprecipitates. A biopharmaceutical study. *Pharmazie*, 49 (1994) 834-839.
- 2110 Carey, R.J. and DePalma, G.: A simplified method for the measurement of caffeine in plasma and brain: evidence for a cortical-subcortical caffeine concentration differential in brain. *J. Neurosci. Methods*, 53 (1994) 19-22; *C.A.*, 121 (1994) 244925p.
- 2111 Golkiewicz, W., Kuczynski, J., Markowski, W. and Jusiak, L.: High-performance liquid chromatography of some alkaloids on unmodified silica gel with aqueous-organic solvent mixtures. *J. Chromatogr. A*, 686 (1994) 85-91.
- 2112 Hirobe, C., Kunugi, A. and Tabei, K.: (Chemistry exercises for daily life. 2. Determination of caffeine in commercial health drinks by HPLC). *Kagaku to Kyoiku*, 42 (1994) 632-635; *C.A.*, 122 (1995) 9034j.
- 2113 Keller, A. and Schulz, H.-U.: Pharmakokinetik von zwei Theophyllin-Präparaten in unterschiedlichen galenischen Retardzubereitungen bei gesunden Probanden. *Arzneim.-Forsch.*, 44 (1994) 1233-1236.
- 2114 Krause, W., Reissmann, F. and Schöbel, C.: Circadian plasma level profile of lisuride in rats and mice after continuous administration via the diet. *Arzneim.-Forsch.*, 45 (1995) 6-10.

- 2115 Liu, L., Riese, J., Resch, K. and Kaever, V.: Impairment of macrophage eicosanoid and nitric oxide production by an alkaloid from *Sinomenium acutum*. *Arzneim.-Forsch.*, 44 (1994) 1223-1226.
- 2116 Ma, Y., Ito, Y., Sokolosky, E. and Fales, H.M.: Separation of alkaloids by pH-zone-refining counter-current chromatography. *J. Chromatogr. A*, 685 (1994) 259-262.
- 2117 Pacifici, R., Pichini, S., Altieri, I., Caronna, A., Passa, A.R. and Zuccaro, P.: High-performance liquid chromatographic-electrospray mass spectrometric determination of morphine and its 3- and 6-glucuronides: application to pharmacokinetic studies. *J. Chromatogr. B*, 664 (1995) 329-334.
- 2118 Papadoyannis, I.N.: Instrumental analytical techniques used for datura alkaloid analyses. *Instrum. Sci. Technol.*, 22 (1994) 241-258; C.A., 122 (1995) 17282d - a review with 18 refs.
- 2119 Pohjola, J. and Harpf, M.: Determination of atropine and obidoxime in automatic injection devices used as antidotes against nerve agent intoxication. *J. Chromatogr. A*, 686 (1994) 350-354.
- 2120 Rivory, L.P. and Robert, J.: Reversed-phase high-performance liquid chromatographic method for the simultaneous quantitation of the carboxylate and lactone forms of the camptothecin derivative irinotecan, CPT-11, and its metabolite SN-38 in plasma. *J. Chromatogr. B*, 661 (1994) 133-141.
- 2121 Rop, P.P., Grimaldi, F., Burle, J., de Saint Leger, M.N. and Viala, A.: Determination of 6-monoacetylmorphine and morphine in plasma, whole blood and urine using high-performance liquid chromatography with electrochemical detection. *J. Chromatogr. B*, 661 (1994) 245-253.
- 2122 Seregina, E.V., Laipanov, A.Kh., Slanskii, V.E. and Voronin, I.A.: (Chromatographic separation of tropane-group alkaloids). *Zh. Fiz. Khim.*, 68 (1994) 1823-1825; C.A., 122 (1995) 17309t.
- 2123 Sochor, J., Klimes, J. and Srurnova, M.: High-performance liquid chromatographic determination of terguride in solid dosage forms and plasma. *J. Chromatogr. B*, 663 (1995) 309-313.
- 2124 Suvegh, G., Zsador, B. and Szejtli, J.: Separation of diastereomer vincamine derivatives by chromatography on  $\beta$ -cyclodextrin polymer stationary phase. In: Hedges, A.R. (Editor), *Minutes Int. Symp. Cyclodextrins*, 6th, Ed. Sante, Paris, 1992, pp. 584-587; C.A., 122 (1995) 71061u.
- 2125 Taborska, E., Bochorakova, H., Paulova, H. and Dostal, J.: Separation of alkaloids in *Chelidonium majus* by reversed phase HPLC. *Planta Med.*, 60 (1994) 380-381; C.A., 121 (1994) 238500e.
- 2126 Tsuchiya, H., Ohtani, S., Yamada, K., Takagi, N. and Hayashi, T.: Determination of tetrahydro- $\beta$ -carboline in urine by high-performance liquid chromatography with suppression of artefact formation. *J. Pharm. Biomed. Anal.*, 12 (1994) 1547-1553.
- 2127 Van der Merwe, P.J., Brown, L.W. and Hendrikz, S.E.: Simultaneous quantification of ephedrine in urine by high-performance liquid chromatography. *J. Chromatogr. B*, 661 (1994) 357-361.
- For additional information see C.A.:  
121 (1994) 213105s, 220898f, 244910e;  
122 (1995) 39001p.
- See also 2292, 2359, 2459.
23. OTHER SUBSTANCES CONTAINING HETEROCYCLIC NITROGEN
- 23a. Porphyrins and other pyrroles
- 2128 Ebdon, L., Evans, E.H., Pretorius, W.G. and Rowland, S.J.: Analysis of geoporphyrins by high-temperature gas chromatography inductively coupled plasma mass spectrometry and high-performance liquid chromatography inductively coupled plasma mass spectrometry. *J. Anal. At. Spectrom.*, 9 (1994) 939-943; C.A., 121 (1994) 234254c.
- 2129 Habdas, J.: Chromatographic characterization of isomers of (4-methoxyphenyl)tolylporphyrin. *Acta Chromatogr.*, 2 (1993) 78-83; C.A., 122 (1995) 71068b.
- 2130 Ho, J.W. and Candy, L.Y.F.: Analysis of metalloporphyrins using cyclodextrin stationary phases with photodiode array UV detection. *J. Liq. Chromatogr.*, 17 (1994) 4111-4119.
- See also 1494, 2240.
- 23c. Indole derivatives and plant hormones (gibberelins)
- 2131 Jinchang, L., Shi, J., Xiaoliang, Z., Wang, G., Huifen, Y. and Yanjun, R.: (Separation and determination of three kinds of plant hormones by high-performance liquid chromatography). *Fenxi Huaxue*, 22 (1994) 801-804; C.A., 121 (1995) 250091e.
- See also 1678, 1715.
- 23d. Pyridine derivatives
- 2132 Defilippi, A., Piancone, G., Tibaldi, G.P., Tartaglino, L. and Viretto, A.: (Determination of isonicotinic acid hydrazide in dairy products). *Ind. Aliment. (Pinerolo)*, 33 (1994) 737-740; C.A., 122 (1995) 8299n.
- 2133 Iwaki, M., Ogiso, T., Hayashi, H., Lin, E.T. and Benet, L.Z.: Simultaneous measurement of nicotinic acid and its major metabolite, nicotinuric acid in urine using high-performance liquid chromatography: application of solid-liquid extraction. *J. Chromatogr. B*, 661 (1994) 154-158.
- 2134 Saz, J.M. and Marina, M.L.: Retention mechanism and implications for selectivity for a group of dihydropyridines in ionic micellar liquid chromatography. *J. Chromatogr. A*, 687 (1994) 1-12.
- 23e. Other N-heterocyclic compounds
- 2135 Denissen, J.F., Grabowski, B.A., Johnson, M.K., Boyd, S.A., Uchic, J.T., Stein, H., Cepa, S. and Hill, P.: The orally active renin inhibitor A-74273. *In vivo* and *in vitro* morpholine ring metabolism in rats, dogs, and humans. *Drug Metab. Disp.*, 22 (1994) 880-888.
- 2136 Gennaro, M.C., Abrigo, C., Marengo, E., Baldin, C. and Martelletti, M.T.: Determination of creatinin in human serum. Statistical intercalibration of methods. *Analyst (Cambridge)*, 120 (1995) 47-51.
- 2137 Imai, J., Murayama, K., Kawai, M., Yamaguchi, S. and Matsuura, K.: Determination of 8-methyl ether of xanthurenic acid in human urine by high-performance liquid chromatography. *J. Chromatogr. B*, 661 (1994) 149-153.

- 2138 Körner, A. and Peter, A.: Alternative methods for the determination of trace amounts of 4-aminomorpholine in molsidomine and linsidomine. *J. Chromatogr. A*, 689 (1995) 235-245.

For additional information see C.A.:  
121 (1994) 244909m.

See also 1373, 1627, 1718, 2431.

## 24. ORGANIC SULPHUR COMPOUNDS (INCL. GLUCOSINOLATES)

- 2139 Andersson, J.T. and Schmid, B.: Polycyclic aromatic sulfur heterocycles. IV. Determination of polycyclic aromatic compounds in a shale oil with the atomic emission detector. *J. Chromatogr. A*, 693 (1995) 325-338.
- 2140 Dangi, R.S., Jeevaratnam, K., Sugendran, K., Malhotra, R.C. and Raghuvveeran, C.D.: Solid-phase extraction and reversed-phase high-performance liquid chromatographic determination of sulphur mustard in blood. *J. Chromatogr. B*, 661 (1994) 341-345.
- 2141 Zhou, J. and Wang, W.: Electrocatalytic oxidation and amperometric determination of sulphhydryl compounds at a copper hexacyanoferrate film glassy carbon electrode in liquid chromatography. *Electroanalysis (N.Y.)*, 6 (1994) 29-35; C.A., 121 (1994) 244531g.

For additional information see C.A.:  
121 (1994) 238504j.

See also 1764, 2084, 2092, 2467.

## 25. ORGANIC PHOSPHORUS COMPOUNDS (INCL. SUGAR PHOSPHATES)

- 2142 Franz, H. and Maier, H.G.: (Inositol phosphates in coffee and coffee products. Part I. Identification and method of determination). *Dtsch. Lebensm.-Rundsch.*, 89 (1993) 276-282; C.A., 121 (1994) 279141g.
- 2143 Kim, H.-Y., Wang, T.-C.L. and Ma, Y.-C.: Liquid chromatography/mass spectrometry of phospholipids using electrospray ionization. *Anal. Chem.*, 66 (1994) 3977-3982.

See also 1642, 1741, 1775, 2075, 2092.

## 26. ORGANOMETALLIC AND RELATED COMPOUNDS

### 26a. Organometallic compounds

- 2144 Feistner, G.J., Gabrik, A.H. and Beer, S.V.: Application of capillary liquid chromatography-electrospray mass spectrometry to identify major siderophores of *Erwinia amylovora* as proferrioxamines and their potential role in virulence. *Dev. Plant Pathol.*, 3(Molecular Mechanisms of Bacterial Virulence) (1994) 429-444; C.A., 121 (1994) 225788e.
- 2145 Nigge, W., Marggraf, U. and Linscheid, M.: The quantitative determination of tin alkylates in sediments. *Fresenius J. Anal. Chem.*, 350 (1994) 533-537.

See also 2518.

### 26b. Boranes, silanes and related non-metallic compounds

- 2146 Born, S.B. and Skelly Frame, E.M.: Development of high-performance liquid-chromatographic-inductively coupled plasma method for speciation and quantification of silicones: from silanols to polysiloxanes. *Analyst (Cambridge)*, 119 (1994) 1687-1694.

### 26c. Coordination compounds

- 2147 Liao, J.P. and Saleh, F.Y.: Retention and spectral characteristics of copper-natural ligand complexes in reverse phase C18-HPLC with UV-PDA and fluorescence detection. *Int. J. Environ. Anal. Chem.*, 56 (1994) 239-259; C.A., 121 (1994) 209696t.
- 2148 Ohtsuka, C., Matsuzawa, K., Wada, H. and Nakagawa, G.: Retention behaviour of metal chelates in ion-pair reversed-phase liquid chromatography as a function of mobile phase composition with methanol-water and acetonitrile-water mobile phases. *Anal. Chim. Acta*, 294 (1994) 69-74.
- 2149 Villani, C. and Pirkle, W.H.: Direct high-performance liquid chromatographic resolution of planar chiral tricarbonyl ( $\eta^6$ -arene)-chromium(0) complexes. *J. Chromatogr. A*, 693 (1995) 63-68.
- 2150 Wang, S.F. and Wai, C.M.: Separation of metal dithiocarbamate complexes by high-performance liquid chromatography. *J. Chromatogr. Sci.*, 32 (1994) 506-510.

See also 1729, 2520.

## 27. VITAMINS AND VARIOUS ANIMAL GROWTH FACTORS (NON-PEPTIDIC)

- 2151 Anderson, P., Davidson, C.M., Littlejohn, D. and Ure, A.M.: Extraction of ergosterol from peaty soils and determination by high performance liquid chromatography. *Talanta*, 41 (1994) 711-720; C.A., 121 (1994) 220681e.
- 2152 Brinkmann, E., Dehne, L., Oei, H.B., Tiebach, R. and Baltes, W.: Separation of geometrical retinol isomers in food samples by using narrow-bore high-performance liquid chromatography. *J. Chromatogr. A*, 693 (1995) 271-279.
- 2153 Chuang, C.-Z., Trosclair, G. and Lopez-S, A.: Adaptation of a carotenoid procedure to analyze carotenoids, retinol, and alpha-tocopherol simultaneously. *J. Liq. Chromatogr.*, 17 (1994) 3613-3622.
- 2154 Diack, M. and Saska, M.: Separation of vitamin E and  $\gamma$ -oryzanols from rice bran by normalphase chromatography. *J. Am. Oil Chem. Soc.*, 71 (1994) 1211-1217.
- 2155 Dueker, S.R., Jones, A.D., Smith, G.M. and Clifford, A.J.: Stable isotope methods for the study of  $\beta$ -carotene- $d_8$  metabolism in humans utilizing tandem mass spectrometry and high-performance liquid chromatography. *Anal. Chem.*, 66 (1994) 4177-4185.
- 2156 El-Agaimy, M.A., Neff, W.E., El-Sayed, M. and Awatif, I.I.: Effect of saline irrigation water on olive composition. *J. Am. Oil Chem. Soc.*, 71 (1994) 1287-1289.

- 2157 Filimonov, V.N., Kolosova, I.F. and Balyatinskaya, L.N.: (Separation of synthetic fat-soluble vitamins by multicolumn normal-phase high-performance liquid chromatography). *Zh. Fiz. Khim.*, 68 (1994) 1835-1838; C.A., 122 (1995) 64524u.
- 2158 Finkel'shtein, E.I., Morev, S.N., Efremova, M.N. and Krasnokutskaya, I.S.: (Determination of retinyl acetate isomers with a multiwave-length detection regime). *Zh. Fiz. Khim.*, 68 (1994) 1829-1831; C.A., 122 (1995) 64522s.
- 2159 Ghareghbagh, R.K. and Ebel, S.: Stabilitätsanalytik von Dexpanthenol, Pantolacton und Pantosäure. *Pharmazie*, 50 (1995) 39-40.
- 2160 Giuliano, A.R., Neilson, E.M., Yap, H.-H., Baier, M. and Canfield, L.M.: Quantitation of and inter/intra-individual variability in major carotenoids of mature human milk. *J. Nutr. Biochem.*, 5 (1994) 551-556; C.A., 121 (1994) 299440g.
- 2161 Haroon, Y.: Reaction detection methods for K vitamins and their 2',3'-epoxy metabolite in liquid chromatography. In: Shearer, M.J. and Seghatchian, M.J. (Editors), *Vitam. K Vitam. K-Depend. Proteins*, CRC, Boca Raton, 1993, 55-90; C.A., 121 (1994) 254413j.
- 2162 Horst, R.L., Reinhardt, T.A., Goff, J.P., Nonnecke, B.J., Gambhir, V.K., Fiorella, P.D. and Napoli, J.L.: Identification of 9-*cis*,13-*cis*-retinoic acid as a major circulating retinoid in plasma. *Biochemistry*, 34 (1995) 1203-1209.
- 2163 Irby, R.B. and Adair, W.L., Jr.: Intermediates in the folic acid biosynthetic pathway are incorporated into molybdopterin the yeast, *Pichia canadensis*. *J. Biol. Chem.*, 269 (1994) 23981-23987.
- 2164 Jezequel-Cuer, M., Moel, G.L., Covi, G., Lepage, S., Peynet, J., Gousson-Evstigneeff, T., Laureaux, C. and Troupel, S.: (Effects of preanalytical conditions on  $\alpha$ -tocopherol stability in blood samples). *Ann. Biol. Clin.*, 52 (1994) 271-276; C.A., 121 (1994) 225405c.
- 2165 Kojima, R., Fujimori, T., Kiyota, N., Toriya, Y., Fukuda, T., Ohashi, T., Sato, T., Yoshizawa, Y., Takeyama, K., Mano, H., Masushige, S. and Kato, S.: *In vivo* isomerization of retinoic acids. Rapid isomer exchange and gene expression. *J. Biol. Chem.*, 269 (1994) 32700-32707.
- 2166 Kolosova, I.F., Filimonov, V.N. and Balyatinskaya, L.N.: (Separation of synthetic B group vitamins by ion-pair high-performance chromatography). *Zh. Fiz. Khim.*, 68 (1994) 1832-1834; C.A., 122 (1995) 64523t.
- 2167 Konings, E.J.M.: Estimation of vitamin D in baby foods with liquid chromatography. *Neth. Milk Dairy J.*, 48 (1994) 31-38; C.A., 121 (1994) 229137q.
- 2168 Kurlandsky, S.B., Xiao, J., Duell, E.A., Voorhees, J.J. and Fisher, G.J.: Biological activity of *all-trans* retinol requires metabolic conversion to *all-trans* retinoic acid and is mediated through activation of nuclear retinoid receptors in human keratinocytes. *J. Biol. Chem.*, 269 (1994) 32821-32827.
- 2169 Mitton, K.P. and Trevithick, J.R.: High-performance liquid chromatograph-electrochemical detection of antioxidants in vertebrate lens: glutathione, tocopherol, and ascorbate. *Methods Enzymol.*, 233(Oxygen Radicals in Biological Systems, Pt. C) (1994) 523-539; C.A., 121 (1994) 225366r.
- 2170 Motoe, K.: (Improved determination method of thiamin and 2-(1-hydroxyethyl)thiamin in foods by high performance liquid chromatography). *Bitamin*, 68 (1994) 379-384; C.A., 121 (1994) 279156r.
- 2171 Perlman, K.L., Pahl, J.M., Smith, C., Kutner, A. and DeLuca, H.F.: 26,27-Dihomo-1 $\alpha$ -hydroxy- and 26,27-dihomo-24-epi-1 $\alpha$ ,25-dihydroxyvitamin D<sub>2</sub> analogs that differ markedly in biological activity *in vivo*. *J. Biol. Chem.*, 269 (1994) 24014-24010.
- 2172 Saikatsu, S., Akeno, N. and Horiuchi, N.: Comparison of single cartridge (C18/OH) and high performance liquid chromatography sample purification for the thymus radioreceptor assay of 1,25(OH)<sub>2</sub>D<sub>3</sub>. *Klin. Labor.*, 39 (1993) 737-744; C.A., 121 (1994) 296505w.
- 2173 Seki, T., Isono, K., Ito, M. and Katsuta, Y.: Flies in the group Cyclorrhapha use (3S)-3-hydroxyretinal as a unique visual pigment chromophore. *Eur. J. Biochem.*, 262 (1994) 691-696.
- 2174 Shimoda, M., Shin, H.-C. and Kokue, E.-i.: Simultaneous determination of tetrahydrofolate, 10-formyltetrahydrofolate and 5-methyltetrahydrofolate in rat bile by high-performance liquid chromatography with electrochemical detection. *J. Vet. Med. Sci.*, 56 (1994) 701-705; C.A., 121 (1994) 250080a.
- 2175 Shin, H.-C., Shimoda, M. and Kokue, E.: Identification of 5,10-methylenetetrahydrofolate in rat bile. *J. Chromatogr. B*, 661 (1994) 237-244.
- 2176 Tzimas, G., Sass, J.O., Wittfoht, W., Elmazar, M.M.A., Ehlers, K. and Nau, H.: Identification of 9,13-*cis*-retinoic acid as a major plasma metabolite of 9-*cis*-retinoic acid and limited transfer of 9-*cis*-retinoic acid and 9,13-*cis*-retinoic acid to the mouse and rat embryos. *Drug Metab. Disp.*, 22 (1994) 928-936.
- 2177 Ueno, H., Yokota, Y. and Tono, T.: (Analysis of cyanocobalamin in pharmaceuticals by high performance liquid chromatography). *Toyama-ken Jakui Kenkyusho Nenpo* 1993, 21 (1994) 85-89; C.A., 122 (1995) 17308s.
- 2178 Werner, E.R., Schmid, M., Werner-Felmayer, G., Mayer, B. and Wachter, H.: Synthesis and characterization of <sup>3</sup>H-labelled tetrahydrobiopterin. *Biochem. J.*, 304 (1994) 189-193.
- 2179 Yakushina, L.M., Beketova, N.A., Bender, Ye.D. and Kharitonchik, L.A.: (Utilization of high-performance liquid chromatography for vitamin determination in biological fluids and food products). *Vopr. Pitan.*, (1993) 43-48; C.A., 121 (1994) 249876h - a review with 57 refs.

For additional information see C.A.:  
121 (1994) 308459z;  
122 (1995) 4612m, 8276c.

See also 1608, 2241.

## 28. ANTIBIOTICS

- 2180 Andersen, A., Warren, D.J. and Slordal, L.: Quantitation of cell-associated doxorubicin by high-performance liquid chromatography after enzymic desequstration. *Cancer Chemother. Pharmacol.*, 34 (1994) 197-202; C.A., 121 (1994) 220900a.
- 2181 Barrow, C.J., Sedlock, D.M., Sun, H.H., Cooper, R. and Gillum, A.M.: Win 66306, a new neurokinin antagonist produced by an *Aspergillus* species: fermentation, isolation and physico-chemical properties. *J. Antibiot.*, 47 (1994) 1182-1187.

- 2182 Bencsath, F.A., Plakas, S.M. and Long, A.R.: Optimization of the analytical performance of the magnetic sector mass spectrometer for the identification of residual chloramphenicol in shrimp. *Biol. Mass Spectrom.*, 23 (1994) 665-674; *C.A.*, 121 (1994) 299439p.
- 2183 Blanchflower, W.J., Hewitt, S.A. and Kennedy, D.G.: Confirmatory assay for the simultaneous detection of five penicillins in muscle, kidney and milk using liquid chromatography-electrospray mass spectrometry. *Analyst (Cambridge)*, 119 (1994) 2595-2601.
- 2184 Bland, J.M., Lax, A.R. and Klich, M.A.: HPLC separation and antifungal activity of the naturally occurring iturin-A homologs. In: Schneider, C.H. and Eberle, A.N. (Editors), *Pept. 1992, Proc. Eur. Pept. Symp.*, 22nd 1992, ESCOM, Leiden, 1993, pp. 332-333; *C.A.*, 121 (1994) 276517y.
- 2185 Delepine, B., Hurtaud, D. and Sanders, P.: Identification of tylosin in bovine muscle at the maximum residue limits level by liquid chromatography-mass spectrometry using a particle beam interface. *Analyst (Cambridge)*, 119 (1994) 2717-2721.
- 2186 Fabre, D., Bressolle, F., Kinowski, J.M., Bouvet, O., Paganin, F. and Galtier, M.: A reproducible, simple and sensitive HPLC assay for determination of ofloxacin in plasma and lung tissue. Application in pharmacokinetic studies. *J. Pharm. Biomed. Anal.*, 12 (1994) 1463-1469.
- 2187 Funabashi, Y., Horiguchi, T., Iinuma, S., Tanida, S. and Harada, S.: TAN-1496 A, C and E, diketopiperazine antibiotics with inhibitory activity against mammalian DNA topoisomerase I. *J. Antibiot.*, 47 (1994) 1202-1218.
- 2188 Hayashi, K.-i., Nakagawa, M. and Nakayama, M.: Nisamycin, a new manumycin group antibiotic from *Streptomyces* sp. K106. Taxonomy, fermentation, physico-chemical and biological properties. *J. Antibiot.*, 47 (1994) 1104-1109.
- 2189 Hedenmo, M. and Eriksson, B.-M.: Liquid chromatographic determination of the macrolide antibiotics roxithromycin and clarithromycin in plasma by automated solid-phase extraction and electrochemical detection. *J. Chromatogr. A*, 692 (1995) 161-166.
- 2190 Hsu, M.-C., Lin, Y.-S. and Chung, H.-C.: High-performance liquid chromatographic method for potency determination of cephalixin in commercial preparations and for stability studies. *J. Chromatogr. A*, 692 (1995) 67-72.
- 2191 Hurtaud, D., Delepine, B. and Sanders, P.: Particle beam liquid chromatography-mass spectrometry method with negative ion chemical ionization for the confirmation of oxacillin, cloxacillin and dicloxacillin residues in bovine muscle. *Analyst (Cambridge)*, 119 (1994) 2731-2736.
- 2192 Hussain, M.S., Chukwumaeze-Obiajunwa, V. and Micetich, R.G.: Sensitive high-performance liquid chromatographic assay for norfloxacin utilizing fluorescence detection. *J. Chromatogr. B*, 663 (1995) 379-384.
- 2193 Iwamoto, T., Fujie, A., Sakamoto, K., Tsurumi, Y., Shigematsu, N., Yamashita, S., Hashimoto, S., Okuhara, M. and Kohsaka, M.: WF 11899A, B and C, novel antifungal lipopeptides I. Taxonomy, fermentation, isolation and physico-chemical properties. *J. Antibiot.*, 47 (1994) 1084-1091.
- 2194 Khan, K., Paesen, J., Roets, E. and Hoogmartens, J.: Analysis of erythromycin A and its metabolites in biological samples by liquid chromatography with post-column ion-pair extraction. *J. Liq. Chromatogr.*, 17 (1994) 4195-4213.
- 2195 Lee, H., Jang, W.C. and Lee, H.S.: Hydrolysis kinetics of metampicillin by high performance liquid chromatography. *Arch. Pharmacol. Res.*, 17 (1994) 378-380; *C.A.*, 121 (1994) 308461u.
- 2196 Lee, H., Lee, J.S. and Lee, H.S.: Simultaneous determination of ampicillin and metampicillin in biological fluids using high-performance liquid chromatography with column switching. *J. Chromatogr. B*, 664 (1995) 335-340.
- 2197 Leroy, P., Decolin, D., Nicolas, A. and Archimbault, P.: Determination of josamycin residues in porcine tissues using high-performance liquid chromatography with pre-column derivatization and spectrofluorometric detection. *Analyst (Cambridge)*, 119 (1994) 2743-2747.
- 2198 Lyon, D.J., Cheung, S.W., Chan, C.Y. and Cheng, A.F.B.: Rapid HPLC assay of clinafloxacin, fleroxacin, levofloxacin, sparfloxacin and tosofloxacin. *J. Antimicrob. Chemother.*, 34 (1994) 446-448; *C.A.*, 121 (1994) 244956z.
- 2199 Mangold, J.B., Schran, H.F. and Tse, F.L.S.: Pharmacokinetics and metabolism of cyclosporin G in humans. *Drug Metab. Disp.*, 22 (1994) 973-979.
- 2200 Moretti, V.M., Maggi, G.L., Albertini, A., Bellagamba, F., Luzzana, U., Serrini, G. and Valfre, F.: High-performance liquid chromatographic determination of oxytetracycline in channel catfish (*Ictalurus punctatus*) muscle tissue. *Analyst (Cambridge)*, 119 (1994) 2749-2751.
- 2201 Morino, T., Masuda, A., Yamada, M., Nishimoto, M., Nishikiori, T., Saito, S. and Shimada, N.: Stevastelins, novel immunosuppressants produced by *Penicillium*. *J. Antibiot.*, 47 (1994) 1341-1343.
- 2202 Muritu, J.W., Kibwage, I.O., Maitai, C.K. and Hoogmartens, J.: Evaluation of tetracycline raw materials and finished products found on the Kenyan market. *J. Pharm. Biomed. Anal.*, 12 (1994) 1483-1488.
- 2203 Paesen, J., Khan, K., Roets, E. and Hoogmartens, J.: Study of the stability of erythromycin in neutral and alkaline solutions by liquid chromatography on poly(styrene-divinylbenzene). *Int. J. Pharm.*, 113 (1995) 215-222; *C.A.*, 122 (1995) 64115m.
- 2204 Parker, R.M. and Patel, R.K.P.: Determination of tilimicosin in ovine milk using high-performance liquid chromatography. *Analyst (Cambridge)*, 119 (1994) 2577-2579.
- 2205 Pinto, C.G., Pavon, J.L.P. and Cordero, B.M.: Micellar liquid chromatography of zwitterions: retention mechanism of cephalosporins. *Analyst (Cambridge)*, 120 (1995) 53-62.
- 2206 Potthast, H., Schug, B., Elze, M., Schwerdtle, R. and Blume, H.: Vergleich der Bioverfügbarkeiten von Erythromycinestolat bzw. Erythromycinethylsuccinat enthaltenden Trockensaft-Zubereitungen im steady state. *Pharmazie*, 50 (1995) 56-60.
- 2207 Saraya, S., Kraisintu, K. and Boonpooh, Y.: High-performance liquid chromatographic determination of tetracycline antibiotics in animal feeds. *Warasan Phesatchasat*, 20 (1993) 25-32; *C.A.*, 121 (1994) 229140k.
- 2208 Seki-Asano, M., Okazaki, T., Yamagishi, M., Sakai, N., Hanada, K. and Mizoue, K.: Isolation and characterization of new 18-membered macrolides FD-891 and FD-892. *J. Antibiot.*, 47 (1994) 1226-1233.
- 2209 Seneviratne, A.K., Jayewardene, A.L. and Gambertoglio, J.G.: Determination of ceftiozime in human abscess fluid by paired ion reversed-phase HPLC. *J. Liq. Chromatogr.*, 17 (1994) 4157-4167.

- 2210 Snippe, N., van de Merbel, N.C., Ruiter, F.P.M., Steijger, O.M., Lingeman, H. and Brinkman, U.A.T.: Automated column liquid chromatographic determination of amoxicillin and cefadroxil in bovine serum and muscle tissue using on-line dialysis for sample preparation. *J. Chromatogr. B*, 662 (1994) 61-70.
- 2211 Straub, R., Linder, M. and Voyksner, R.D.: Determination of  $\beta$ -lactam residues in milk using perfusive-particle liquid chromatography combined with ultrasonic nebulization electrospray mass spectrometry. *Anal. Chem.*, 66 (1994) 3651-3658.
- 2212 Tamaki, M., Akabori, S. and Muramatsu, I.: Studies of gramicidin S analogues having various ring sizes by reversed-phase high-performance liquid chromatography. *J. Chromatogr. A*, 685 (1994) 237-241.
- 2213 Touraki, M., Rigas, P., Pergandas, P. and Kastritsis, C.: Determination of oxytetracycline in the live fish feed *Artemia* using high-performance liquid chromatography with ultraviolet detection. *J. Chromatogr. B*, 663 (1995) 167-171.
- 2214 Wetterich, U. and Mutschler, E.: Quality of cefotaxime sodium preparations. *Arzneim.-Forsch.*, 45 (1995) 74-80.

For additional information see C.A.:

121 (1994) 213091j, 220904e, 244908k.

See also 2400, 2422, 2432, 2437.

## 29. INSECTICIDES, PESTICIDES AND OTHER AGROCHEMICALS

- 2215 Parrilla, P., Vidal, J.L.M., Galera, M.M. and Frenich, A.G.: Simple and rapid screening procedure for pesticides in water using SPE and HPLC/DAD detection. *Fresenius J. Anal. Chem.*, 350 (1994) 633-637.

### 29a. General techniques

- 2216 Parrilla, P., Galera, M.M., Vidal, J.L.M. and Frenich, A.G.: Determination of fenamiphos and folpet in water by time-domain differentiation of high-performance liquid chromatographic peaks. *Analyst (Cambridge)*, 119 (1994) 2231-2236.
- 2217 Sundaram, K.M.S., Nott, R. and Lewin, E.E.: Comparative study of the determination of tebufenozide in formulated products by gas chromatographic and liquid chromatographic methods. *J. Chromatogr. A*, 687 (1994) 323-332.

See also 2470.

### 29b. Chlorinated insecticides

- 2218 Barnabas, I.J., Dean, J.R., Hitchen, S.M. and Owen, S.P.: Selective supercritical fluid extraction of organochlorine pesticides and herbicides from aqueous samples. *J. Chromatogr. Sci.*, 32 (1994) 547-551.
- 2219 Dehal, S.S. and Kupfer, D.: Metabolism of the proestrogenic pesticide methoxychlor by hepatic P450 monooxygenases in rats and humans. Dual pathway involving novel *ortho* ring-hydroxylation by CYP2B. *Drug Metab. Disp.*, 22 (1994) 937-946.

- 2220 Grimvall, E. and Östman, C.: Fractionation of non-*ortho*-substituted toxic polychlorinated biphenyls on two nitro-containing liquid chromatographic stationary phases. *J. Chromatogr. A*, 685 (1994) 338-343.

See also 1532, 1540, 1548.

### 29c. Phosphorus insecticides

- 2221 Molina, C., Honig, M. and Barcelo, D.: Determination of organophosphorus pesticides in water by solid-phase extraction followed by liquid chromatography/high-flow pneumatically assisted electrospray mass spectrometry. *Anal. Chem.*, 66 (1994) 4444-4449.

For additional information see C.A.:

121 (1994) 254020d.

### 29e. Herbicides

- 2222 Dankwardt, A., Pullen, S., Rauchalles, S., Kramer, K., Just, F., Hock, B., Hofmann, R., Schewes, R. and Maidl, F.X.: Atrazine residues in soil two years after the atrazine ban - a comparison of enzyme immunoassay with HPLC. *Anal. Lett.*, 28 (1994) 621-634.
- 2223 Frebortova, J. and Tatarkovicova, V.: Determination of phenoxyacid herbicides in water. *Acta Univ. Palacki, Olomuc., Fac. Rerum Nat.*, 112 (1993) 63-68; C.A., 121 (1994) 294950p.
- 2224 Galletti, G.C., Bonetti, A. and Dinelli, G.: High-performance liquid chromatographic determination of sulfonylureas in soil and water. *J. Chromatogr. A*, 692 (1995) 27-37.
- 2225 Hatik, S., Lehotay, J. and Tekel, J.: Simultaneous HPLC determination of phenylurea herbicides and their corresponding anilines in water after on-line preconcentration. *J. High Resolut. Chromatogr.*, 17 (1994) 756-758.
- 2226 Marx, A., Giersch, T. and Hock, B.: Immunoaffinity chromatography of s-triazines. *Anal. Lett.*, 28 (1995) 267-278.
- 2227 Thomas, D.H., Beck-Westermeyer, M. and Hage, D.S.: Determination of atrazine in water using tandem high-performance immunoaffinity chromatography and reversed-phase liquid chromatography. *Anal. Chem.*, 66 (1994) 3823-3829.

For additional information see C.A.:

121 (1994) 274366z.

See also 1770, 2218.

### 29f. Fungicides

- 2228 Alawi, M.A.: Development of a new approach for the determination of folpet in grapes. *Anal. Lett.*, 28 (1995) 349-356.
- 2229 Jiménez, J.J., Atienza, J. and Bernal, J.L.: HPLC with fluorescence detection for the study of benomyl dissipation on treated lettuce. *J. Liq. Chromatogr.*, 17 (1994) 3999-4017.

29g. *Other types of pesticides and various agrochemicals*

- 2230 Brouwer, E.R., Struys, E.A., Vreuls, J.J. and Brinkman, U.A.T.: Automated determination of pyrethroid insecticides in surface water by column liquid chromatography with diode array UV detection, using on-line micelle-mediated sample preparation. *Fresenius J. Anal. Chem.*, 350 (1994) 487-495.
- 2231 Darwish, A.: Determination of deltamethrin levels in wool by reversed-phase high performance liquid chromatography. *J. Liq. Chromatogr.*, 17 (1994) 4215-4228.
- 2232 Guenther, K., Drauz, K., Klenk, H., Kriechbaum, D., Merget, S. and Schaefer, M.: (Semipreparative chromatographic enantiomeric resolution of a cyanazine homolog, using cellulose triacetate and benzoate). *GIT Spez. Chromatogr.*, 13 (1993) 96-101; *C.A.*, 121 (1994) 224188x.
- 2233 Ho, J.S. and Budde, W.L.: Investigation of the natural pesticide rotenone in water using liquid-solid disk extraction, supercritical fluid elution, and liquid chromatography/particle beam mass spectrometry. *Anal. Chem.*, 66 (1994) 3716-3722.

See also 1490, 2217.

30. SYNTHETIC AND NATURAL DYES

30a. *Synthetic dyes*

- 2234 Akins, D.L. and Kumar, V.T.: High-performance liquid chromatography of cyanine dyes. Multiphase separation, purification, and substitution of the counter ion. *J. Chromatogr. A*, 689 (1995) 269-273.
- 2235 Fink, W. and Auch, J.: (Malachite green, crystal violet, and brilliant green residues determination in edible fish by HPLC). *Dtsch. Lebensm.-Rundsch.*, 89 (1993) 246-251; *C.A.*, 121 (1994) 253976h.

See also 2482.

30b. *Chloroplast and other natural pigments*

- 2236 Bekasova, O.D., Sineshchekov, V.A., Brykina, G.D., Sineshchekova, E.V. and Grishina, L.E.: (Minor components in the pigment system of *Nostoc muscorum* and *Mastigodadus laminosus*). *Biokhimiya (Moscow)*, 59 (1994) 1042-1053.
- 2237 Fiorini, M.: Preparative high-performance liquid chromatography for the purification of natural anthocyanins. *J. Chromatogr. A*, 692 (1995) 213-219.
- 2238 Minguez-Mosquera, M.I. and Gandul-Rojas, B.: High-performance liquid chromatographic study of alkaline treatment of chlorophyll. *J. Chromatogr. A*, 690 (1995) 161-176.
- 2239 Misawa, N., Truesdale, M.R., Sandmann, G., Fraser, P.D., Bird, C., Schuch, W. and Bramley, P.M.: Expression of a tomato cDNA coding for phytoene synthase in *Escherichia coli*, phytoene formation *in vivo* and *in vitro*, and functional analysis of the various truncated gene products. *J. Biochem. (Tokyo)*, 116 (1994) 980-985.
- 2240 Saitoh, K., Awaka, I. and Suzuki, N.: Determination of chlorophylls by reversed-phase high-performance liquid chromatography with isocratic elution and the column-switching technique. *J. Chromatogr. A*, 693 (1995) 176-180.

- 2241 Sass, J.O. and Nau, H.: Single-run analysis of isomers of retinoyl- $\beta$ -D-glucuronide and retinoic acid by reversed-phase high-performance liquid chromatography. *J. Chromatogr. A*, 685 (1994) 182-188.
- 2242 Schoefs, B., Bertrand, M. and Lemoine, Y.: Separation of photosynthetic pigments and their precursors by reversed-phase high-performance liquid chromatography using a photodiode-array detector. *J. Chromatogr. A*, 692 (1995) 239-245.
- 2243 Steinberg, G. and Sheves, M.: Factors affecting the formation of an M-like intermediate in the photocycle of 13-*cis*-bacteriorhodopsin. *Biochemistry*, 33 (1994) 12439-12450.

For additional information see C.A.:  
122 (1995) 5047t.

See also 2152, 2468.

31. PLASTICS AND THEIR INTERMEDIATES

- 2244 Aignasse, M.F., Prognon, P., Stachowicz, M., Gheyouché, R. and Pradeau, D.: A new simple and rapid HPLC method for determination of DEHP in PVC packing and releasing studies. *Int. J. Pharm.*, 113 (1995) 241-246; *C.A.*, 122 (1995) 17326w.
- 2245 Dayal, U. and Mehta, S.K.: Determination of long chain branching in polyethylene by high temperature SEC coupled with a MALLS photometric detector. In: Bhardwaj, I.S. (Editor), *Polym. Sci.*, 2, Allied Publ., New Delhi, 1994, pp. 737-742; *C.A.*, 122 (1995) 32517c.
- 2246 Estrin, Ya.I. and Kasumova, L.T.: (Chromatography of functional oligobutadienes under critical conditions on silica gel). *Zh. Fiz. Khim.*, 68 (1994) 1784-1788; *C.A.*, 122 (1995) 11088y.
- 2247 Fung, V.S. and Tang, A.S.K.: Liquid chromatographic determination of plasticizer di(2-ethylhexyl)phthalate (DEHP) in PVC plastics. *Fresenius J. Anal. Chem.*, 350 (1994) 721-723.
- 2248 Gur'yanova, V.V. and Prudskova, T.N.: (Features of the chromatographic behavior of macromolecules in a binary eluent). *Vysokomol. Soedin., Ser. A, Ser. B*, 36 (1994) 1731-1736; *C.A.*, 122 (1995) 32520y.
- 2249 Huguenard, C., Elaissari, A. and Pefferkorn, E.: Surface area exclusion chromatography of diblock copolymers. *Polym. Prepr. (Am. Chem. Soc., Div. Polym. Chem.)*, 35 (1994) 643-644; *C.A.*, 121 (1994) 301814c.
- 2250 Lederer, K., Beytollahi-Amtmann, I. and Billiani, J.: Determination and correction of peak broadening in size exclusion chromatography of controlled rheology polypropylene. *J. Appl. Polym. Sci.*, 54 (1994) 47-55; *C.A.*, 121 (1994) 231821b.
- 2251 Menon, S.K., Dhoble, D.A. and Kulkarni, R.A.: Characterization of polymers using GPC coupled with online multi-angle laser light scattering photometry. In: Bhardwaj, I.S. (Editor), *Polym. Sci.*, 2, Allied Publ., New Delhi, 1994, pp. 749-754; *C.A.*, 122 (1995) 57013y.
- 2252 Mukesh, D.: Molecular graphics of polymers. In: Bhardwaj, I.S. (Editor), *Polym. Sci.*, 2, Allied Publ., New Delhi, 1994, pp. 719-724; *C.A.*, 121 (1994) 301734b.

- 2253 Netopilik, M.: Model analysis of the determination of the Mark-Houwink-Kuhn-Sakurada parameters and molecular weight distribution by means of size exclusion chromatography with dual detection. *Polymer*, 35 (1994) 4799-4803; C.A., 121 (1994) 301797z.
- 2254 Parlesak, A., Bode, J.C. and Bode, C.: Parallel determination of gut permeability in man with  $M_r$  400,  $M_r$  1500,  $M_r$  4000 and  $M_r$  10000 polyethylene glycol. *Eur. J. Clin. Chem. Clin. Biochem.*, 32 (1994) 813-820.
- 2255 Podzimek, S.: The use of GPC coupled with a multiangle laser light scattering photometer for the characterization of polymers. On the determination of molecular weight, size, and branching. *J. Appl. Polym. Sci.*, 54 (1994) 91-103; C.A., 121 (1994) 231822c.
- 2256 Puskas, J.E. and Hutchinson, R.: GPC calibration for the molecular weight measurement of butyl rubbers. *Rubber Chem. Technol.*, 66 (1993) 742-748; C.A., 122 (1995) 11888j.
- 2257 Schunk, T.C. and Long, T.E.: Compositional distribution characterization of poly(methyl methacrylate)-graft-polydimethylsiloxane copolymers. *J. Chromatogr. A*, 692 (1995) 221-232.
- 2258 Spahl, W. and Budzikiewicz, H.: Qualitative analysis of dental resin composites by gas and liquid chromatography/mass spectrometry. *Fresenius J. Anal. Chem.*, 350 (1994) 684-691.
- 2259 Wang, Q.C., Svec, F. and Frechet, J.M.J.: Fine control of the porous structure and chromatographic properties of monodisperse macroporous poly(styrene-co-divinylbenzene) beads prepared using polymer porogens. *J. Polym. Sci., Part A: Polym. Chem.*, 32 (1994) 2577-2588; C.A., 121 (1994) 232182f.
- 2260 Zigon, M., Leben, S., Sedlacek, J. and Grubisic-Gallot, Z.: Characterization of poly(phenylacetylenes) by SEC coupled with continuous viscometry and LALLS. *Acta Chim. Slov.*, 40 (1993) 255-266; C.A., 121 (1994) 256762w.
- For additional information see C.A.:  
121 (1994) 257618r, 302036n.
- See also 1344, 1348, 1349, 1375.
32. DRUG ANALYSIS
- 32a. Drug analysis, general techniques
- 2261 Bonet-Domingo, E., Medina-Hernandez, M.J. and Garcia-Alvarez-Coque, M.C.: On the direct injection of urine samples in micellar liquid chromatography. *Quim. Anal. (Barcelona)*, 12 (1993) 167-172; C.A., 121 (1994) 220901b.
- 2262 Bourquin, T., Bordi, D., Bounine, J.P., Compagnon, P.A., Drouin, J.E., Fatras, A., Lecoindre, L., Lepabic, C., Molt, P. et al.: (Conformity tests for liquid-phase chromatography: report of an SFSTP commission). *A.T.P. Pharma Prat.*, 4 (1994) 170-176; C.A., 121 (1994) 238491d.
- 2263 Castensson, S.: Standardized chromatographic media products by operative change control. *Prog. Biotechnol.*, 9 (1994) 973-975; C.A., 121 (1994) 238491c.
- 2264 Croes, K., McCarthy, P.T. and Flanagan, R.J.: HPLC of basic drugs and quaternary ammonium compounds on microparticulate strong cation-exchange materials using methanolic or aqueous methanol eluents containing an ionic modifier. *J. Chromatogr. A*, 693 (1995) 289-306.
- 2265 Franssen, E.J.F., Luurtsema, G., Medema, J., Visser, G.M., Jeronismus-Shaling, C.M., Bruins, A.P. and Vaalburg, W.: Application of liquid chromatography combined with mass-spectrometry (LC-MS) to establish identity and purity of PET-radiopharmaceuticals. *Appl. Radiat. Isot.*, 45 (1994) 937-940; C.A., 121 (1994) 286707z.
- 2266 Hagan, R.L.: High-performance liquid chromatography for small-scale studies of drugs stability. *Am. J. Hosp. Pharm.*, 51 (1994) 2162-2175; C.A., 121 (1994) 286684q.
- 2267 Haginaka, J., Seyama, C. and Murashima, T.: (Direct injection analysis of drug enantiomers in serum on protein-bonded silica materials by liquid chromatography). *Kuromatogurafi*, 15 (1994) 54-55; C.A., 121 (1994) 291876h.
- 2268 Hashimoto, A., Yamasaki, K., Kokusanya, Y., Miyamoto, T. and Sato, T.: Investigation of "Signal" constituents for the evaluation of animal crude drugs. I. Free aminoacids and total amino acids. *Chem. Pharm. Bull.*, 42 (1994) 1636-1641; C.A., 122 (1995) 64560c.
- 2269 Hermansson, J. and Grahn, A.: Resolution of racemic drugs on a new chiral column based on silica-immobilized cellobiohydrolase. Characterization of the basic properties of the column. *J. Chromatogr. A*, 687 (1994) 45-59.
- 2270 Hsu, H.-C. and Chien, C.-S.: (Validation of analytical methods: A simple model for HPLC assay methods). *Yaowu Shipin Fenxi*, 2 (1994) 161-176; C.A., 121 (1994) 308435p - a review with 39 refs.
- 2271 Iwabuchi, H., Kitazawa, E., Kobayashi, N., Watanabe, H., Kanai, M. and Nakamura, K.-i.: Studies on drug metabolism using liquid chromatography/mass spectrometry: comparison of three liquid chromatographic/mass spectrometric interfaces. *Biol. Mass Spectrom.*, 23 (1994) 540-546; C.A., 122 (1995) 167x.
- 2272 Li, Z. and Guo, P.: (Application of high performance liquid chromatography column switching technique in the study of pharmacokinetics and bioavailability). *Sepu*, 12 (1994) 378-379; C.A., 122 (1995) 132g - a review with 9 refs.
- 2273 Messer, D.C., Taylor, L.T., Weiser, W.E., McRae, J.W. and Cook, C.C.: A novel strictly anaerobic recovery and enrichment system incorporating lithium for detection of heat-injured *Listeria monocytogenes* in pasteurized milk containing background microflora. *Pharm. Res.*, 11 (1994) 1545-1548; C.A., 121 (1994) 299436k.
- 2274 Nasal, A., Sznitowska, M., Bucinski, A. and Kaliszan, R.: Hydrophobicity parameter from high-performance liquid chromatography on an immobilized artificial membrane column and its relationship to bioactivity. *J. Chromatogr. A*, 692 (1995) 83-89.
- 2275 Ong, C.P., Chow, K.K., Ng, C.L., Ong, F.M., Lee, H.K. and Li, S.F.Y.: Use of overlapping resolution mapping scheme for optimization of the high-performance liquid chromatographic separation of pharmaceuticals. *J. Chromatogr. A*, 692 (1995) 207-212.

- 2276 Ponder, G.W., Butram, S.L., Adams, A.G., Ramanathan, C.S. and Stewart, J.T.: Resolution of promethazine, ethopropazine, trimeprazine and trimipramine enantiomers on selected chiral stationary phases using high-performance liquid chromatography. *J. Chromatogr. A*, 692 (1995) 173-182.
- 2277 Shah, K.P., Chang, M. and Riley, C.M.: Automated analytical systems for drug development studies. II - A system for dissolution testing. *J. Pharm. Biomed. Anal.*, 12 (1994) 1519-1527.
- 2278 Van Molle, S., Vanbel, P. and Tilquin, B.: Mixture desing statistical technique optimization procedure applied to HPLC and its limitations. *J. Pharm. Belg.*, 49 (1994) 293-300; *C.A.*, 121 (1994) 263789p.

For additional information see *C.A.*:  
121 (1994) 213104r, 213180n.

See also 1398, 1399, 1501, 1606, 1726, 2315, 2341, 2469.

### 32b. Antirheumatics and antiinflammatory drugs

- 2279 Faouzi, M.A., Dine, T., Luyckx, M., Brunet, C., Gressier, B., Cazin, M., Wallaert, B. and Cazin, J.C.: High-performance liquid chromatographic method for the determination of budesonide in bronchoalveolar lavage of asthmatic patients. *J. Chromatogr. B*, 664 (1995) 463-467.
- 2280 Fischer, C. and Klotz, U.: Radical-derived oxidation products of 5-aminosalicylic acid and N-acetyl-5-aminosalicylic acid. *J. Chromatogr. B*, 661 (1994) 57-68.
- 2281 Hayball, P.J., Holman, J.W. and Nation, R.L.: Influence of octanoic acid on the reversible protein binding of ketorolac enantiomers to human serum albumin (HSA): comparative liquid chromatographic studies using a HSA chiral stationary phase. *J. Chromatogr. B*, 662 (1994) 128-133.
- 2282 Jones, D.J. and Bjorksten, A.R.: Detection of ketorolac enantiomers in human plasma using enantioselective liquid chromatography. *J. Chromatogr. B*, 661 (1994) 165-167.
- 2283 King, J.N., Mauron, C., Voirol, M.J., le Goff, C. and Hauffe, S.A.: High performance liquid chromatography and pharmacokinetics of the non-steroidal anti-inflammatory drug oxindanac in calves. *J. Vet. Pharmacol. Ther.*, 17 (1994) 186-192; *C.A.*, 121 (1994) 220956y.
- 2284 Peng, W., Li, T., Li, H. and Wang, E.: Direct injection of urine and determination of acetaminophen by micellar liquid chromatography with a wall-jet cell/carbon fibre microelectrode. *Anal. Chim. Acta*, 298 (1994) 415-421.
- 2285 Vajda, M., Otta-Horvath, K. and Toth-Forgacs, E.: Development of analytical methods for the determination of indosol as a pure substance or in eye drops. *Acta Pharm. Hung.*, 64 (1994) 131-134; *C.A.*, 121 (1994) 308483c.

For additional information see *C.A.*:  
121 (1994) 244911f, 308487g, 308506n.

See also 1465, 2429.

### 32c. Autonomic and cardiovascular drugs

- 2286 Agbaba, D., Janjic, V., Zivanov-Stakic, D. and Vladimirov, S.: High performance liquid chromatographic assay for isosorbide 5-mononitrate and impurities of inorganic nitrates in pharmaceuticals. *J. Liq. Chromatogr.*, 17 (1994) 3983-3988.
- 2287 Alton, K.B., Chan, T.-M. and Pramanik, B.N.: Urinary metabolites of (R),(R)-labetalol. *Drug Metab. Disp.*, 22 (1994) 866-872.
- 2288 Bansal, R., Louridas, A.T., Gottesman, R.D. and Aranda, J.V.: Determination of amrinone in human plasma by high-performance liquid chromatography with ultraviolet detection. *J. Liq. Chromatogr.*, 17 (1994) 3531-3539.
- 2289 Calucci, G., di Giuseppe, E. and Mazzeo, P.: Determination of alfuzosin in human plasma by high-performance liquid chromatography with column-switching. *J. Liq. Chromatogr.*, 17 (1994) 3989-3997.
- 2290 De Beer, J.O., Vandembroucke, C.V. and Massart, D.L.: Experimental design for the rapid selection of separation conditions for methyl and propyl parahydroxybenzoate, phenylephrine hydrochloride and chlorphenamine maleate by ion-pair liquid chromatography. *J. Pharm. Biomed. Anal.*, 12 (1994) 1379-1396.
- 2291 Egginger, G., Lindner, W., Brunner, G. and Stoschitzky, K.: Direct enantioselective determination of (R)- and (S)-propranolol in human plasma. Application to pharmacokinetic studies. *J. Pharm. Biomed. Anal.*, 12 (1994) 1537-1545.
- 2292 El-Dawa, M.A., Habeeb, A.A., Mabrouk, M.M. and El-Bastawissy, I.A.: HPLC assay of reproterol hydrochloride. *Egypt. J. Pharm. Sci.* 1993, 34 (1994) 691-698; *C.A.*, 122 (1995) 64548e.
- 2293 Grundy, J.S., Kherani, R. and Foster, R.T.: Photostability determination of commercially available nifedipine oral dosage formulations. *J. Pharm. Biomed. Anal.*, 12 (1994) 1529-1535.
- 2294 Hasegawa, T., Hara, K., Kenmochi, T. and Hata, S.: *In vitro* metabolism of dorzolamide, a novel potent carbonic anhydrase inhibitor, in rat liver microsomes. *Drug Metab. Disp.*, 22 (1994) 916-921.
- 2295 He, H., Edeki, T.I. and Wood, A.J.J.: Determination of low plasma timolol concentrations following topical application of timolol eye drops in humans by high-performance liquid chromatography with electrochemical detection. *J. Chromatogr. B*, 661 (1994) 351-356.
- 2296 Höld, K.M., de Boer, D., Zuidema, J. and Maes, R.A.A.: Evaluation of the Salivette as sampling device for monitoring  $\beta$ -adrenoceptor blocking drugs in saliva. *J. Chromatogr. B*, 663 (1995) 103-110.
- 2297 Hoshino, M., Yajima, K., Suzuki, Y. and Okahira, A.: Determination of nadolol diastereomers in dog plasma using chiral derivatization and reversed-phase high-performance liquid chromatography with fluorescence detection. *J. Chromatogr. B*, 661 (1994) 281-289.
- 2298 Hsieh, J.Y.-K., Lin, C., Matuszewski, B.K. and Dobrinska, M.R.: Fully automated methods for the determination of hydrochlorothiazide in human plasma and urine. *J. Pharm. Biomed. Anal.*, 12 (1994) 1555-1562.
- 2299 Ishii, K., Minato, K., Nishimura, N., Miyamoto, T. and Sato, T.: Direct chromatographic resolution of four optical isomers of diltiazem hydrochloride on a Chiralcel OF column. *J. Chromatogr. A*, 686 (1994) 93-100.

- 2300 Iyer, E.K. and Tipnis, H.P.: Quantitative determination of oxprenolol in plasma using high performance liquid chromatography. *Indian Drugs*, 31 (1994) 440-441; C.A., 122 (1995) 173w.
  - 2301 Kaniwa, N., Shameem, M., Katori, N., Aoyagi, N. and Kojima, S.: The suitable temperature specification for the evaluation of controlled release products *in vitro*. *Pharmazie*, 50 (1995) 53-55.
  - 2302 Kwok, D.K.W., Igwemezie, L., Kerr, C.R. and McErlane, K.M.: High-performance liquid chromatographic analysis using a highly sensitive fluorogenic reagent, 2-anthroyl chloride, and stereoselective determination of the enantiomers of mexiletine in human serum. *J. Chromatogr. B*, 661 (1994) 271-280.
  - 2303 Lankford, S.M. and Bai, S.A.: Determination of the stereochemical composition of the major metabolites of verapamil in dog urine with enantioselective liquid chromatographic techniques. *J. Chromatogr. B*, 663 (1995) 91-101.
  - 2304 Laugel, C., Chaminade, P., Baillet, A. and Ferrier, D.: Ion-pair reversed-phase liquid chromatographic determination of dihydralazine. *J. Chromatogr. A*, 686 (1994) 344-349.
  - 2305 McCarthy, J.P.: Direct enantiomeric separation of the four stereoisomers of nadolol using normal-phase and reversed-phase high-performance liquid chromatography with Chiralpak AD. *J. Chromatogr. A*, 685 (1994) 349-355.
  - 2306 McNulty, M.J., Page, T.L. and Deal, D.L.: Development and validation of a chiral LC method for analysis of the four stereoisomers of 1045U85 in plasma. *J. Pharm. Biomed. Anal.*, 12 (1994) 1453-1461.
  - 2307 Miller-Stein, C., Boppana, V.K. and Rhodes, G.R.: High-performance liquid chromatographic determination of the N-ethyl tricarbamate ester pro-drug of fenoldopam utilizing simultaneous post-column hydrolysis and fluorescence derivatization. *J. Chromatogr. B*, 661 (1994) 291-297.
  - 2308 Modamio, P., Montejó, O., Lastra, C.F. and Marino, E.L.: A valid high-performance liquid chromatography method for oxprenolol stability studies. *Int. J. Pharm.*, 112 (1994) 93-96; C.A., 121 (1994) 263462b.
  - 2309 Neau, B., Roux, J., Lambrey, B., Courtiol, C., Marty, J.-P. and Grislain, L.: Metabolism of <sup>14</sup>C-moxisylylate in hairless rat. Comparison between intravenous and oral route. *Arzneim.-Forsch.*, 45 (1995) 10-14.
  - 2310 Saarinen, M.T., Sirén, H., Riekkola, M.-L.: Screening and determination of  $\beta$ -blockers, narcotic analgesics and stimulants in urine by high-performance liquid chromatography with column switching. *J. Chromatogr. B*, 664 (1995) 341-346.
  - 2311 Santoro, M.I.R.M., Cho, H.S. and Kedor-Hackmann, E.R.M.: Determination of  $\beta$ -blockers in ophthalmic solutions by high performance liquid chromatography. *Anal. Lett.*, 28 (1995) 71-81.
  - 2312 Trausch, B., Oertel, R., Richter, K. and Gramatte, T.: Die Proteinbindung von Talinolol. *Pharmazie*, 50 (1995) 72.
  - 2313 Ulvi, V. and Keski-Hyynilä, H.: Analysis of chlorothiazide in the presence of its photodecomposition products. *Am. Lab. (Shelton)*, 26 (1994) 56-59; C.A., 122 (1995) 17322s.
  - 2314 Vlcek, J., Macek, K., Hulek, P., Bratova, M. and Fendrich, Z.: Pharmacokinetic parameters of verapamil and its active metabolite norverapamil in patients with hepatopathy. *Arzneim.-Forsch.*, 45 (1995) 146-149.
  - 2315 Wallerstein, S. and Cserhati, T.: Interaction of some anti-hypoxia drugs with hydroxypropyl- $\beta$ -cyclodextrin studied by means of charge transfer chromatography. *J. Biochem. Biophys. Methods*, 29 (1994) 49-60; C.A., 121 (1994) 238256e.
  - 2316 Wu, C., Akiyama, A. and Straub, J.A.: High-performance liquid chromatographic reversed-phase and normal-phase separation of diastereomeric  $\alpha$ -ketoamide calpain inhibitors. *J. Chromatogr. A*, 684 (1994) 243-249.
  - 2317 Yang, J.M. and Chan, K.: Simultaneous determination of moricizine and its sulphoxidation metabolites in biological fluids by high-performance liquid chromatography. *J. Chromatogr. B*, 663 (1995) 172-176.
  - 2318 Yazan, Y. and Bozan, B.: Rapid analysis of verapamil in plasma by reversed phase HPLC. *Pharmazie*, 50 (1995) 117-119.
- For additional information see C.A.:  
 121 (1994) 213149j, 244923m;  
 122 (1995) 38999h.
- See also 1506, 1680, 1723, 2113, 2114, 2124, 2358, 2447.
- 32d. *Central nervous system drugs*
- 2319 Abdel-Hay, M.H.: High-performance liquid chromatographic determination of floctafenine in presence of floctafenine acid. *Alexandria J. Pharm. Sci.*, 7 (1993) 25-28; C.A., 121 (1994) 308482b.
  - 2320 Agostini, O. and Toja, E.: Determination of a new antibrachospastic agent, MX2/120, in guinea pig plasma by high-performance liquid chromatography in a pharmacokinetic study. *J. Chromatogr. B*, 661 (1994) 313-318.
  - 2321 Amjad, F., Ansari, A.F. and Bhangar, M.I.: A simple method for the determination of chlorpromazine in pharmaceutical preparations and blood serum. *Pak. J. Pharm. Sci.*, 6 (1993) 79-82; C.A., 121 (1994) 308468b.
  - 2322 Annesley, T.M. and Clayton, L.T.: Determination of felbamate in human serum by high-performance liquid chromatography. *Ther. Drug Monit.*, 16 (1994) 419-424; C.A., 121 (1994) 244914j.
  - 2323 Atta-Politou, J., Tsalpalis, K. and Koutselinis, A.: A modified simple and rapid reversed phase high performance liquid chromatographic method for quantification of amitriptyline and nortriptyline in plasma. *J. Liq. Chromatogr.*, 17 (1994) 3969-3982.
  - 2324 Balizs, G., Börner, S. and Wennemar, A.: Determination of amperozide residues in swine liver using liquid chromatography-mass spectrometry. *Analyst (Cambridge)*, 119 (1994) 2687-2690.
  - 2325 Blume, H., Ali, S.L., Elze, M., Krämer, J., Wendt, G. and Scholz, M.E.: Relative Bioverfügbarkeit von Paracetamol in Suppositorien-Zubereitungen im Vergleich zu Tabletten. *Arzneim.-Forsch.*, 44 (1994) 1333-1338.
  - 2326 Boppana, V.K.: Simultaneous determination of granisetron and its 7-hydroxy metabolite in human plasma by reversed-phase high-performance liquid chromatography utilizing fluorescence and electrochemical detection. *J. Chromatogr. A*, 692 (1995) 195-202.
  - 2327 Dash, A.K.: Determination of pseudoephedrine hydrochloride in dosage forms by liquid chromatography. *J. Pharm. Biomed. Anal.*, 12 (1994) 1601-1606.

- 2328 Deeks, N.J., Abbott, R.W., Allen, G.D., Hollis, F.J. and Rhodes, G.: Determination of BRL 46470 in human plasma by high-performance liquid chromatography with ultraviolet absorbance detection followed by post-column photochemical reaction and fluorescence detection. *Analyst (Cambridge)*, 119 (1994) 2043-2050.
- 2329 Ebling, W.F., Russell Wada, D. and Stanski, D.R.: From piecewise to full physiologic pharmacokinetic modeling: applied to thipental disposition in the rat. *J. Pharmacokin. Biopharm.*, 22 (1994) 259-292.
- 2330 El-Yazigi, A., Wahab, F.A. and Afrane, B.: Stability study and content uniformity of prochlorperazine in pharmaceutical preparations by liquid chromatography. *J. Chromatogr. A*, 690 (1995) 71-76.
- 2331 Falcó, P.C., Legua, C.M., Hernandez, R.H. and Cabeza, A.S.: Improved amphetamine and methamphetamine determination in urine by normal-phase high-performance liquid chromatography with sodium 1,2-naphthoquinone 4-sulphonate as derivatizing agent and solid-phase extraction for sample clean-up. *J. Chromatogr. B*, 663 (1995) 235-245.
- 2332 Ficarra, R., Ficarra, P., Tommasini, S., Carulli, M., Costatino, D. and Calbro, M.L.: Chromatographic investigations on brotizolam. *Farmaco*, 49 (1994) 437-440; C.A., 121 (1994) 308503j.
- 2333 Franssen, R.M.E., Stolk, L.M.L., Brand, W.V.D. and Smit, B.J.: Determination of morphine and amphetamine. *Ziekenhuisfarmacie*, 10 (1994) 111-112; C.A., 122 (1995) 17321p.
- 2334 Gupta, R.N.: Column liquid chromatographic determination of paroxetine in human serum using solid-phase extraction. *J. Chromatogr. B*, 661 (1994) 362-365.
- 2335 Gurley, B.J., Zermatten, S. and Skelton, D.: Determination of antipyrine in human serum by direct injection restricted access media liquid chromatography. *J. Pharm. Biomed. Anal.*, 12 (1994) 1591-1595.
- 2336 Hempenius, J., Hendriks, G., Hingstman, J., Mensink, C.K., Jonkman, J.H.G. and Lin, C.C.: An automated analytical method for the determination of felbamate in human plasma by robotic sample preparation and reversed-phase high performance liquid chromatography. *J. Pharm. Biomed. Anal.*, 12 (1994) 1443-1451.
- 2337 Husain, S., Sekar, R. and Rao, R.N.: Enantiomeric separation and determination of antiparkinsonian drugs by reversed-phase ligand-exchange high-performance liquid chromatography. *J. Chromatogr. A*, 687 (1994) 351-355.
- 2338 Jurima-Romet, M., Crawford, K., Cyr, T. and Inaba, T.: Terfenadine metabolism in human liver. *In vitro* inhibition by macrolide antibiotics and azole antifungals. *Drug Metab. Disp.*, 22 (1994) 849-857.
- 2339 Kountourellis, J.E. and Ebete, K.O.: Reversed-phase high performance liquid chromatographic determination of cyproheptadine from urine by solid-phase extraction. *J. Chromatogr. B*, 664 (1995) 468-471.
- 2340 Lacroix, M., Tu, T.M., Donati, F. and Varin, F.: High-performance liquid chromatographic assays with fluorometric detection for mivacurium isomers and their metabolites in human plasma. *J. Chromatogr. B*, 663 (1995) 297-307.
- 2341 Larew, L.A., Olsen, B.A., Stafford, J.D. and Wilhelm, M.V.: Comparison of theory-based and empirical modeling for the prediction of chromatographic behavior in the ion-pairing separation of benzodiazepine-derived pharmaceutical compounds. *J. Chromatogr. A*, 692 (1995) 183-193.
- 2342 Lau, G.S.N. and Critchley, J.A.J.H.: The estimation of paracetamol and its major metabolites in both plasma and urine by a single high-performance liquid chromatography assay. *J. Pharm. Biomed. Anal.*, 12 (1994) 1563-1572.
- 2343 Lindegård, B., Björk, H., Jönsson, J.Å., Matthiasson, L. and Olsson, A.-M.: Column liquid chromatographic determination of a basic drug in blood plasma using the supported liquid membrane technique for sample pretreatment. *Anal. Chem.*, 66 (1994) 4490-4497.
- 2344 Mangoni, P., Sioufi, P. and Godbillon, J.: Stereospecific high-performance liquid chromatographic determination of an S(-)-benzopyran methyl ester derivative (CGP 50 068), its (-)-carboxylic acid metabolite (CGP 55 461) and the related (+)-enantiomer (CGP 54 228) in human and dog plasma. *J. Chromatogr. B*, 664 (1995) 393-400.
- 2345 Marastoni, M., Salvadori, S., Balboni, G., Scaranari, V., Borea, P.A. and Tomatis, A.: Synthesis and opioid activity of tyrosine sulfate containing dermorphin and deltorphin peptides. *Arzneim.-Forsch.*, 45 (1995) 116-119.
- 2346 Morris, R.G., Jones, T.E. and Saccoia, N.C.: Liquid chromatographic assay for dextromoramide in human plasma. *J. Liq. Chromatogr.*, 17 (1994) 4185-4193.
- 2347 Nicolle, E., Michaut, S., Serre-Debeauvais, F. and Bessard, G.: Rapid and sensitive high-performance liquid chromatographic assay for nalbuphine in plasma. *J. Chromatogr. B*, 663 (1995) 111-117.
- 2348 Nieto, C., Ramis, J., Conte, L., Fernández, J.M. and Forn, J.: On-line fully automated solid-phase extraction-liquid chromatography analysis of 1,2-dihydro-4-(1,2-dihydro-2-oxo-1-pyridyl)-2,2-dimethyl-1-oxonaphthalene-6-carbonitrile (UR-8225), a new potassium channel opener, in plasma samples. *J. Chromatogr. B*, 661 (1994) 319-325.
- 2349 Norris, R.L.G., Ravenscroft, P.J. and Pond, S.M.: Sensitive high-performance liquid chromatographic assay with ultraviolet detection of methadone enantiomers in plasma. *J. Chromatogr. B*, 661 (1994) 346-350.
- 2350 Rudolphi, C. and Blaschke, G.: Determination of the stereoselective aspects in *in vitro* and *in vivo* metabolism of the analgesic meptazinol by high-performance liquid chromatography. *J. Chromatogr. B*, 663 (1995) 315-326.
- 2351 Russel, F.G.M., Creemers, M.C.W., Tan, Y., van Riel, P.L.C.M. and Gribnau, F.W.J.: Ion-pair solid-phase extraction of cimetidine from plasma and subsequent analysis by high-performance liquid chromatography. *J. Chromatogr. B*, 661 (1994) 173-177.
- 2352 Sadjak, A. and Wintersteiger, R.: Compatibility of morphine, baclofen, floxuridine and fluorouracil in an implantable medication pump. *Arzneim.-Forsch.*, 45 (1995) 93-98.
- 2353 Shimada, K., Mino, T., Nakajima, M., Wakabayashi, H. and Yamato, S.: Application of the desulfurization of phenothiazines for a sensitive detection method by high-performance liquid chromatography. *J. Chromatogr. B*, 661 (1994) 85-91.
- 2354 Thuaud, N. and Sebillé, B.: Structural factors affecting the enantiomeric separation of barbiturates and thiobarbiturates with a chiral side-chain by various  $\beta$ -cyclodextrin supports. Effects of the presence of hydroxypropyl substituents on the chiral selector. *J. Chromatogr. A*, 685 (1994) 15-20.
- 2355 Tzeng, T.-B., Stamm, G. and Chu, S.-y.: Sensitive method for the assay of sertindole in plasma by high-performance liquid chromatography and fluorimetric detection. *J. Chromatogr. B*, 661 (1994) 299-306.

- 2356 Vermeij, T.A.C. and Edelbroek, P.M.: High-performance liquid chromatographic and megabore gas-liquid chromatographic determination of levetiracetam (ucb L059) in human serum after solid-phase extraction. *J. Chromatogr. B*, 662 (1994) 134-139.
- 2357 Verweij, A.M.A., Hordijk, M.L. and Lipman, P.J.L.: Quantitative liquid chromatography, thermospray/tandem mass spectrometric (LC/TSP/MS/MS) analysis of some tranquilizers of the thioxanthene group in whole-blood. *J. Liq. Chromatogr.*, 17 (1994) 4099-4110.
- 2358 Wahbi, A.-A.M., El-Walily, A., Bedair, M. and El-Gendy, A.: Determination of cinnarizine in the presence of its degradation products. *Egypt. J. Pharm. Sci.*, 34 (1993) 35-46; *C.A.*, 121 (1994) 308477d.
- 2359 Wilson, J.M., Cohen, R.I., Kezer, E.A. and Smith, E.R.: Analysis of dezocine in serum and urine by high performance liquid chromatography and pre-column derivatization. *J. Liq. Chromatogr.*, 17 (1994) 4245-4257.
- 2360 Yamaguchi, M., Monji, H., Yamashita, K., Aoki, I. and Yashiki, T.: Sensitive high-performance liquid chromatographic determination of chlorpheniramine in human serum using column switching. *J. Chromatogr. B*, 661 (1994) 168-172.
- For additional information see *C.A.*:  
121 (1994) 220905f, 263817w, 272147c, 308488h;  
122 (1995) 64555e.
- See also 1393, 1430, 1677, 2123, 2277, 2285, 2290, 2310, 2316, 2398.
- 32e. *Chemotherapeutics (exc. cytostatics and antibiotics)*
- 2361 Al-Angary, A.A., Bayomi, M.A., Khidr, S.H., Al-Meshal, M.A. and Lutfi, K.M.: Determination of arteether in plasma using a simple and rapid high-performance liquid chromatographic assay. *Anal. Lett.*, 27 (1994) 2689-2702.
- 2362 Balizs, G., Benesch-Girke, L., Börner, S. and Hewitt, S.A.: Comparison of the determination of four sulphonamides and their N<sup>4</sup>-acetyl metabolites in swine muscle tissue using liquid chromatography with ultraviolet and mass spectral detection. *J. Chromatogr. B*, 661 (1994) 75-84.
- 2363 Barbato, F., Morrica, P., Seccia, S. and Ventriglia, M.: High performance liquid chromatographic analysis of quinolone antibacterial agents. *Farmaco*, 49 (1994) 407-410; *C.A.*, 121 (1994) 308500f.
- 2364 Berger, B.J. and Fairlamb, A.H.: High-performance liquid chromatographic method for the separation and quantitative estimation of anti-parasitic melaminophenyl arsenical compounds. *Trans. R. Soc. Trop. Med. Hyg.*, 88 (1994) 357-359; *C.A.*, 121 (1994) 213160f.
- 2365 Botsoglou, N.A., Fletouris, D.J., Psomas, I.E. and Vassilopoulos, V.N.: Chromatographic behavior of the anthelmintic fenbendazole and its major metabolite oxfendazole in various ion-pair liquid chromatographic systems. *J. Liq. Chromatogr.*, 17 (1994) 4229-4243.
- 2366 Csiba, A. and Lombai, G.: (Liquid chromatographic determination of quinine in soft drinks). *Elelmiszervizsgalati Kozl.*, 40 (1994) 112-119; *C.A.*, 121 (1994) 299402w.
- 2367 Denouel, J., Keller, H.P., Schaub, P., Delaborde, C. and Humbert, H.: Determination of terbinafine and its desmethyl metabolite in human plasma by high-performance liquid chromatography. *J. Chromatogr. B*, 663 (1995) 353-359.
- 2368 Eckers, C., Hutton, K.A., de Biasi, V., East, P.B., Haskins, N.J. and Jacewicz, V.W.: Determination of clavam-2-carboxylate in clavulanate potassium and tablet material by liquid chromatography-tandem mass spectrometry. *J. Chromatogr. A*, 686 (1994) 213-218.
- 2369 El-Gizawy, S.M.: HPLC analysis of metronidazole and diloxanide furoate in its dosage forms. *Anal. Lett.*, 28 (1995) 83-92.
- 2370 El-Sayed, Y.M.: A simple high-performance liquid chromatographic assay for sparfloxacin in human plasma. *Anal. Lett.*, 28 (1995) 279-293.
- 2371 Ferioli, V., Gamberini, G., Rustichelli, C. and Vezzadini, F.: Direct determination of non-UV-absorbing compounds by high-performance liquid chromatography. *Farmaco*, 49 (1994) 411-413; *C.A.*, 122 (1995) 17331u.
- 2372 Fletouris, D., Botsoglou, N.A., Psomas, I.E. and Mantis, A.I.: Rapid ion-pair liquid chromatographic method for the determination of bendazole in cows' milk. *Analyst (Cambridge)*, 119 (1994) 2801-2804.
- 2373 Koks, C.H.W., Rosing, H., Meenhorst, P.L., Bult, A. and Beijnen, J.H.: High-performance liquid chromatographic determination of the antifungal drug fluconazole in plasma and saliva of human immunodeficiency virus-infected patients. *J. Chromatogr. B*, 663 (1995) 345-351.
- 2374 Nagaraja, N.V., Singh, S.K. and Gupta, R.C.: Sensitive high-performance liquid chromatographic method for the determination of methyl N-[5-[[4-(2-pyridinyl)-1-piperazinyl]carbonyl]-1H-benzimidazol-2-yl] carbamate in rat blood. *J. Chromatogr. B*, 664 (1995) 472-477.
- 2375 Nakamura, K. and Nobuso, M.: (Rapid determination of hydrophilic synthetic antibacterial drugs in meat and fish by high-performance liquid chromatography). *Hiroshima-ken Hoken Kankyo Senta Kenkyu Hokoku*, 1 (1993) 67-70; *C.A.*, 121 (1994) 279149r.
- 2376 Nakos, D.S., Botsoglou, N.A. and Psomas, I.E.: Ion-pair isolation and liquid chromatographic determination of albendazole, oxfendazole, oxiabendazole, and thiabendazole residues in milk. *J. Liq. Chromatogr.*, 17 (1994) 4145-4155.
- 2377 Porter, S.: Confirmation of sulfonamide residues in kidney tissue by liquid chromatography-mass spectrometry. *Analyst (Cambridge)*, 119 (1994) 2753-2756.
- 2378 Tanaka, M., Oshima, Y., Aoki, H. and Hakusui, H.: Determination of a new fluoroquinolone antimicrobial agent, (S)-10-[(S)-(8-amino-6-azaspiro[3,4]octan-6-yl)]-9-fluoro-2,3-dihydro-3-methyl-7-oxo-7H-pyrido[1,2,3-de][1,4]benzoxazine-6-carboxylic acid hemihydrate, DV-7751a, in human serum and urine using solid-phase extraction and high-performance liquid chromatography with fluorescence detection. *J. Chromatogr. B*, 664 (1995) 401-407.
- 2379 Thomas, K.M., Dabholkar, D.A. and Jain, C.L.: Simultaneous determination of pyrimethamine and sulphalene in pharmaceutical dosage forms by HPLC. *Indian Drugs*, 31 (1994) 343-345; *C.A.*, 122 (1995) 38990y.
- 2380 Uno, K. and Maeda, I.: Simultaneous determination of sulphamonomethoxine and its N<sup>4</sup>-acetyl metabolite in blood serum by high-performance liquid chromatography with direct injection. *J. Chromatogr. B*, 663 (1995) 177-180.

- 2381 Whitehouse, L.W., Menzies, A., Dawson, B., Cyr, T.D., By, A.W., Black, D.B. and Zamecnik, J.: Mouse hepatic metabolites of ketoconazole: isolation and structure elucidation. *J. Pharm. Biomed. Anal.*, 12 (1994) 1425-1441.

For additional information see C.A.:

121 (1994) 308491d.

See also 1703, 2277, 2426, 2438.

### 32f. Cytostatics

- 2382 Albertioni, F., Pettersson, B., Reichelova, V., Juliusson, G. and Lillemark, J.: Analysis of 2-chloro-2'-deoxyadenosine in human blood plasma and urine by high-performance liquid chromatography using solid-phase extraction. *Ther. Drug. Monit.*, 16 (1994) 413-418; C.A., 121 (1994) 244913h.
- 2383 Belfayol, L., Guillemin, L., Louchahi, K., Lortholary, O., Bosio, A.M. and Fauvel, F.: Measurement of 4-hydroxycyclophosphamide in serum by reversed-phase high-performance liquid chromatography. *J. Chromatogr. B*, 663 (1995) 395-399.
- 2384 Belz, S., Frickel, C., Wolfrom, C., Nau, H. and Henze, G.: High-performance liquid chromatographic determination of methotrexate, 7-hydroxymethotrexate, 5-methyltetrahydrofolic acid and folic acid in serum and cerebrospinal fluid. *J. Chromatogr. B*, 661 (1994) 109-118.
- 2385 Carins, W.R.L., Ebdon, L. and Hill, S.J.: Development of an HPLC-ICP-MS method for the determination of platinum species from new anti-tumor drugs. *Anal. Proc.*, 31 (1994) 295-297; C.A., 121 (1994) 308510j.
- 2386 De Fusco, M., D'Incalci, M., Gentili, D., Reichert, S. and Zucchetti, M.: High-performance liquid chromatographic assay for the determination of the novel etoposide derivative dimethylaminoetoposide (NK611) and its metabolites in urine of cancer patients. *J. Chromatogr. B*, 664 (1995) 409-414.
- 2387 Forgács, E. and Cserhádi, T.: Determination of hydrophobicity of non-homologous series of anticancer drugs by reversed-phase high-performance liquid chromatography. *J. Chromatogr. B*, 664 (1995) 277-285.
- 2388 Hanada, K., Nagai, N. and Ogata, H.: Quantitative determination of unchanged cisplatin in rat kidney and liver by high-performance liquid chromatography. *J. Chromatogr. B*, 663 (1995) 181-186.
- 2389 Huizing, M.T., Rosing, H., Koopman, F., Keung, A.C.F., Pinedo, H.M. and Beijnen, J.H.: High-performance liquid chromatographic procedures for the quantitative determination of paclitaxel (Taxol) in human urine. *J. Chromatogr. B*, 664 (1995) 373-382.
- 2390 Mouchard-Delmas, C., Gourdier, B. and Vistelle, R.: Determination of vinorelbine in rabbit plasma by high-performance liquid chromatography with coulometric detection. *J. Chromatogr. B*, 663 (1995) 390-394.
- 2391 Nattouf, H., Davrinche, C., Sinet, M. and Farinotti, R.: High-performance liquid chromatographic assay for 3'-amino-3'-deoxythymidine in plasma, with 9-fluorenyl methyl chloroformate as the derivatization agent. *J. Chromatogr. B*, 664 (1995) 365-371.
- 2392 Okuda, T., Motohashi, M., Aoki, I. and Yashiki, T.: Sensitive determination of methotrexate in monkey plasma by high-performance liquid chromatography using on-line solid-phase extraction. *J. Chromatogr. B*, 662 (1994) 79-84.
- 2393 Song, D. and Au, J.L.-S.: Isocratic high-performance liquid chromatographic assay of taxol in biological fluids and tissues using automated column switching. *J. Chromatogr. B*, 663 (1995) 337-344.
- 2394 Sparreboom, A., van Tellingen, O., Nooijen, W.J. and Beijnen, J.H.: Determination of paclitaxel and metabolites in mouse plasma, tissues, urine and faeces by semi-automated reversed-phase high-performance liquid chromatography. *J. Chromatogr. B*, 664 (1995) 383-391.
- 2395 Stafford, C.G. and St. Claire, R.L., III.: High-performance liquid chromatographic analysis of the lactone and carboxylate forms of a topoisomerase I inhibitor (the antitumor drug GI147211) in plasma. *J. Chromatogr. B*, 663 (1995) 119-126.
- 2396 Wedge, S.R., Laohavinij, S., Taylor, G.A. and Newell, D.R.: Measurement of 5,10-dideaza-5,6,7,8-tetrahydrofolate (lometrexol) in human plasma and urine by high-performance liquid chromatography. *J. Chromatogr. B*, 663 (1995) 327-335.
- 2397 Wright, J.E., Tretyakov, O., Ayashi, L.J., Elias, A., Rosowsky, A. and Frei, E. III.: Analysis of 4-hydroxycyclophosphamide in human blood. *Anal. Biochem.*, 224 (1995) 151-158.

For additional information see C.A.:

121 (1994) 308492e;

122 (1995) 38978a.

See also 1772, 2084, 2120, 2180, 2352, 2409, 2428, 2459.

### 32g. Other drug categories

- 2398 Akkers, M.R.J., Holtmaat, I. and Niel, J.C.C.V.: (Analysis stability and a (ring) survey of physostigmine salicylate injection 1 ml=1mg FNA). *Ziekenhuisfarmacie*, 10 (1994) 44-49; C.A., 121 (1994) 213088p.
- 2399 Anonymous: (Activity of the Oils and Fats Experimental Station in the cosmetics field). *Riv. Ital. Sostanze Grasse*, 71 (1994) 437-450; C.A., 121 (1994) 307960u.
- 2400 Bandak, S., Czejka, M., Schillter, J. and Schernhammer, E.: Pharmacokinetic drug interaction between epirubicin and interferon- $\alpha$ -2b in serum and red blood cells. *Arzneim.-Forsch.*, 45 (1995) 212-215.
- 2401 Barra, J., Taburet, A.M., Jardel, A., Fessi, H. and Puisieux, F.: High-performance liquid chromatographic assay for the simultaneous determination of ethyl clofibrate and clofibric acid in plasma. Evaluation of plasma stability of ethyl clofibrate polylactic nanocapsules in human and rat plasmas. *J. Chromatogr. B*, 661 (1994) 178-182.
- 2402 Benson, L.M., Veverka, K.A., Mays, D.C., Nelson, A.N., Shriver, Z.H., Lipsky, J.J. and Naylor, S.: Simultaneous structure-activity determination of disulfiram photolysis products by on-line continuous-flow liquid secondary ion mass spectrometry and enzyme inhibition assay. *J. Chromatogr. A*, 693 (1995) 162-166.

- 2403 Cai, W.M., Hatton, J., Pettigrew, L.C., Dempsey, R.J. and Chandler, M.H.H.: A simplified high-performance liquid chromatographic method for direct determination of warfarin enantiomers and their protein binding in stroke patients. *Ther. Drug Monit.*, 16 (1994) 509-512; C.A., 121 (1994) 291887n.
- 2404 Coppi, G. and Silingardi, S.: Pharmacokinetics of pidotimod in rats and dogs. *Arzneim.-Forsch.*, 44 (1994) 1460-1464.
- 2405 Crimella, T., Orlandi, R., Bocchiola, G., Anders, U. and Stradi, R.: Analytical and chemical profile of pidotimod. *Arzneim.-Forsch.*, 44 (1994) 1405-1410.
- 2406 Cutié, S.S. and Smith, C.G.: Determination of Methocel A15-LV cellulose ether in blends with microcrystalline cellulose. *J. Chromatogr. A*, 693 (1995) 371-375.
- 2407 D'Angelo, L., de Ponti, F., Crema, F., Caravaggi, M. and Crema, A.: Effect of food on the bioavailability of pidotimod in healthy volunteers. *Arzneim.-Forsch.*, 44 (1994) 1473-1475.
- 2408 Depin, J.C., Vigie, A., Chavernac, G., Rousselot, C., Lardy, C. and Guerrier, D.: Pharmacodynamics and antithrombotic effects after intravenous administration of the new thromboxane A<sub>2</sub> receptor antagonist sodium 4-[[1-[[[(4-chlorophenyl)sulfonyl]amino]ethyl]cyclopentyl]methyl]benzeneacetate. *Arzneim.-Forsch.*, 44 (1994) 1203-1207.
- 2409 Ezan, E., Carde, P., le Kerneau, J., Ardouin, T., Thomas, F., Isnard, F., Deschamps de Paillette, E. and Grognet, J.-M.: Pharmacokinetics in healthy volunteers and patients of NAc-SDKP (seraspénide), a negative regulator of hematopoiesis. *Drug Metab. Disp.*, 22 (1994) 943-948.
- 2410 Ferioli, V., Rustichelli, C., Vezzadini, F. and Gamberini, G.: Determination of azalaic acid in pharmaceuticals and cosmetics by RP-HPLC after pre-column derivatization. *Farmaco*, 49 (1994) 421-425; C.A., 121 (1994) 308502h.
- 2411 Hayes, F.J., Baker, T.R., Dobson, R.L.M. and Tsueda, M.S.: Rapid liquid chromatographic-mass spectrometric assay for oxymetazoline in whole rat blood. *J. Chromatogr. A*, 692 (1995) 73-81.
- 2412 Hsieh, J.Y.-K., Lin, C. and Matuszewski, B.K.: Robotic sample preparation and high-performance liquid chromatographic analysis of verlukast in human plasma. *J. Chromatogr. B*, 661 (1994) 307-312.
- 2413 Hung, B.-F., Hsu, Y.-N. and Yang, J.-M.: Determination of betanecol chloride by indirect photometric chromatography. *Chin. Pharm. J. (Taipei)*, 46 (1994) 123-133; C.A., 121 (1994) 308484d.
- 2414 Jones, G.D., Hoff, T.A., Gunraj, P.E., Facchini, V. and Chauhan, P.B.: Assay and single dose pharmacokinetics of a novel systemic acyl coenzyme A cholesterol O-acyltransferase inhibitor, RP 73163, in rat plasma using automated solid-phase extraction with high-performance liquid chromatography. *J. Chromatogr. B*, 661 (1994) 119-131.
- 2415 Kakchenko, O.I. and Ignatyev, N.V.: (High-performance liquid chromatography in the analysis of tromexan). *Khim.-Farm. Zh.*, 28 (1994) 63-64; C.A., 121 (1994) 213158m.
- 2416 Kim, Y., Park, S., Park, J. and Lee, W.: Detection of benzthiazide by high-performance liquid chromatography-thermospray mass spectrometry. *J. Chromatogr. A*, 689 (1995) 170-174.
- 2417 Laryea, M.D., Zass, R., Ritgen, J. and Wendel, U.: Simultaneous determination of betaine and N,N-dimethylglycine in urine. *Clin. Chim. Acta*, 230 (1994) 169-175.
- 2418 Lau, O.-W. and Mok, C.-S.: High-performance liquid chromatographic determination of active ingredients in cough-cold syrups with indirect conductometric detection. *J. Chromatogr. A*, 693 (1995) 45-54.
- 2419 Mailland, F., Coppi, G. and Silingardi, S.: Pharmacokinetics and oral bioavailability of pidotimod in humans. *Arzneim.-Forsch.*, 44 (1994) 1465-1469.
- 2420 Naylor, P.H., Naylor, C.W., Sasaki, D. and Mutchnick, M.G.: Utilization of HPLC-ELISA to assess serum levels of thymosin  $\alpha_1$  following subcutaneous administration to human subjects. *J. Liq. Chromatogr.*, 17 (1994) 3541-3551.
- 2421 Pfeffer, M. and Wykhoff, B.: An automated analytical high-performance liquid chromatographic procedure for iopamidol solutions using a benchmate workstation. *J. Liq. Chromatogr.*, 17 (1994) 4259-4271.
- 2422 Poirier, J.-M., Lebot, M. and Cheymol, G.: Cyclosporine in whole blood: Drug monitoring difficulties and presentation of a reliable normal-phase liquid chromatographic assay. *Ther. Drug Monit.*, 16 (1994) 388-394; C.A., 121 (1994) 271112c.
- 2423 Qin, X.-Z., Tsai, E.W., Sakuma, T. and Ip, D.P.: Pharmaceutical application of liquid chromatography-mass spectrometry. II. Ion chromatography-ion spray mass spectrometric characterization of alendronate. *J. Chromatogr. A*, 686 (1994) 205-212.
- 2424 Radulovic, D., Kocic-Pesic, V., Pecanac, D. and Zivanovic, L.: HPLC determination of sodium chromoglycate in pharmaceutical dosage forms. *Farmaco*, 49 (1994) 375-376; C.A., 121 (1994) 213131x.
- 2425 Rastogi, S.C. and Johansen, S.S.: Comparison of high-performance liquid chromatographic methods for the determination of 1,2-dibromo-2,4-dicyanobutane in cosmetic products. *J. Chromatogr. A*, 692 (1995) 53-57.
- 2426 Santoni, G., Medica, A., Gratterer, P., Furlanetto, S. and Pinzauti, S.: High-performance liquid chromatographic determination of benzalkonium and naphazoline or tetrahydrozoline in nasal and ophthalmic solutions. *Farmaco*, 49 (1994) 751-754; C.A., 122 (1995) 17336z.
- 2427 Sanwald, P., Huebert, N.D. and Haegele, K.D.: Simultaneous measurement of the major metabolites of dolasetron mesilate in human urine using solid-phase extraction and high-performance liquid chromatography. *J. Chromatogr. B*, 661 (1994) 101-107.
- 2428 Shaw, L.M., Bonner, H.S. and Brown, D.Q.: Metabolic pathways of WR-2721 (ethyol, amifostine) in the Balb/c mouse. *Drug Metab. Disp.*, 22 (1994) 895-902.
- 2429 Spöhrer, U. and Eyer, P.: Separation of geometric *syn* and *anti* isomers of obidoxime by ion-pair high-performance liquid chromatography. *J. Chromatogr. A*, 693 (1995) 55-61.
- 2430 Stevenson, A.J., Weber, M.P., Trudel, R., Leavitt, R., Woodard, D., Todi, F., Mendonca, M., Robillo, V., Young, L. et al.: Monitoring furosemide in race horses participating in an EIPH program. *J. Vet. Pharmacol. Ther.*, 17 (1994) 163-168; C.A., 121 (1994) 223773x.
- 2431 Tan, H.S.-I., Xu, J. and Zheng, Y.: Cation-exchange high-performance liquid chromatographic assay of piperazine in some pharmaceutical formulations. *J. Chromatogr. A*, 693 (1995) 307-314.

- 2432 Taylor, P.J., Salm, P., Norris, R.L.G., Ravenscroft, P.J. and Pond, S.M.: Comparison of high-performance liquid chromatography and monoclonal fluorescence polarization immunoassay for the determination of whole-blood cyclosporin A in liver and heart transplant patients. *Ther. Drug Monit.*, 16 (1994) 526-530; C.A., 121 (1994) 291888p.
- 2433 Ushio, T. and Yamamoto, K.: High-performance liquid chromatography of enantiomers of {2-[4-(3-ethoxy-2-hydroxypropoxy)phenylcarbamoyl]ethyl}-dimethylsulfonium *p*-toluenesulfonate (suplatast tosilate) on a cellulose tris-3,5-dimethylphenylcarbamate column. *J. Chromatogr. A*, 684 (1994) 235-242.
- 2434 Van Waarde, A., Anthonio, R.L., Visser, T.J., Elsinga, P.H., Posthumus, H., Weemaes, A.-M.A., Blanksma, P.K., Visser, G.M., Paans, A.M.J. and Vaalburg, W.: Quantification of an  $^{11}\text{C}$ -labelled  $\beta$ -adrenoceptor ligand, S-(-)CGP 12177, in plasma of humans and rats. *J. Chromatogr. B*, 663 (1995) 361-369.
- 2435 Whigan, D.B. and Schuster, A.E.: Manual and automated determination of 1- $\beta$ -D-arabinofuranosyl-E-5-(2-bromovinyl)uracil and its metabolite (E)-5-(2-bromovinyl)uracil in urine. *J. Chromatogr. B*, 664 (1995) 357-363.
- 2436 Woolf, E., Au, T., Haddix, H. and Matuszewski, B.: Determination of L-735 524, an human immunodeficiency virus protease inhibitor, in human plasma and urine via high-performance liquid chromatography with column switching. *J. Chromatogr. A*, 692 (1995) 45-52.
- 2437 Zeder-Lutz, G., van Regenmortel, M.H.V., Wenger, R. and Altschuh, D.: Interaction of cyclosporin A and two cyclosporin analogs with cyclophilin: relationship between structure and binding. *J. Chromatogr. B*, 662 (1994) 301-306.
- 2438 Zehender, H., Denouël, J., Roy, M., le Saux, L. and Schaub, P.: Simultaneous determination of terbinafine (Lamisil) and five metabolites in human plasma and urine by high-performance liquid chromatography using on-line solid-phase extraction. *J. Chromatogr. B*, 664 (1995) 347-355.
- For additional information see C.A.:  
121 (1994) 213081f, 213176r, 263263n, 308512m;  
122 (1995) 63956z.
- See also 1499, 1554, 1677, 2073, 2119, 2135, 2199, 2201, 2326, 2339.
- 32h. *Toxicological and forensic applications*
- 2439 Below, E. and Burrmann, M.: Application of HPLC equipment with rapid scan detection to the identification of drugs in toxicological analysis. *J. Liq. Chromatogr.*, 17 (1994) 4131-4144.
- 2440 Bonino, N., Diodati, J. and Cena, M.R.: Monitoring of urine by extraction chromatography with tri-N-octylphosphine oxide supported on a polypropylene column. *J. Radioanal. Nucl. Chem.*, 182 (1994) 103-110; C.A., 121 (1994) 275325x.
- 2441 Draisci, R., Lucentini, L., Giannetti, L. and Stacchini, A.: (Algal biotoxins in mussels: optimization of the HPLC method for okadaic acid determination). *Riv. Sci. Aliment.*, 22 (1993) 443-454; C.A., 121 (1994) 229114e.
- 2442 Grogg-Sulser, K., Helmlin, J.-J. and Clerc, J.-T.: Qualitative and quantitative determination of illicit heroin street samples by reversed-phase high-performance liquid chromatography: method development by CARTAGO-S. *J. Chromatogr. A*, 692 (1995) 121-129.
- 2443 Guittou, J., Durand, D., Bouillod, N. and Manchon, M.: Development of a solid/liquid extraction method for analysis of toxic drugs in serum using an automated liquid chromatograph: the Remedi. *Ann. Biol. Clin.*, 52 (1994) 111-115; C.A., 121 (1994) 220903d.
- 2444 Koves, E.M.: Use of high-performance liquid chromatography-diode array detection in forensic toxicology. *J. Chromatogr. A*, 692 (1995) 103-119.
- 2445 Low, A.S. and Taylor, R.B.: Analysis of common opiates and heroin metabolites in urine by high-performance liquid chromatography. *J. Chromatogr. B*, 663 (1995) 225-233.
- 2446 Nishikawa, M., Nakajima, K., Tatsuno, M., Kasuya, F., Igarashi, K., Fukui, M. and Tsuchihashi, H.: The analysis of cocaine and its metabolites by liquid chromatography/atmospheric-pressure chemical-ionization mass spectrometry (LC/APCI-MS). *Forensic Sci. Int.*, 66 (1994) 149-158; C.A., 121 (1994) 294522a.
- 2447 Tenczer, J., Lappenbergh-Pelzer, M., Schneider, V., Demme, U. and Köppel, C.: Fatal poisoning with detajmium: identification of detajmium and its metabolites and artifacts by gas chromatography-mass spectrometry and quantification by high-performance liquid chromatography. *J. Chromatogr. B*, 661 (1994) 47-55.
- For additional information see C.A.:  
121 (1994) 273951z.
- See also 1719, 1803, 2106, 2117, 2121, 2521.
- 32i. *Plant extracts*
- 2448 Barthomeuf, C., Chmielowiec, C. and Pourrat, H.: TLC - densitometry for quality control of *Hedera helix* L. (Araliaceae). *J. Planar Chromatogr.*, 7 (1994) 474-476.
- 2449 Cai, Y., Phillipson, J.D., Harper, J.I. and Corne, S.J.: High-performance liquid chromatographic and proton magnetic resonance spectroscopic method for quality evaluation of *Paeonia* roots. *Phytochem. Anal.*, 5 (1994) 183-189; C.A., 121 (1994) 263791h.
- 2450 Chuang, W.-C. and Sheu, S.-J.: Determination of ginsenosides in ginseng crude extracts by high-performance liquid chromatography. *J. Chromatogr. A*, 685 (1994) 243-251.
- 2451 Du, Q., Li, M., Cheng, Q., Zhang, T.Y. and Ito, Y.: Purification of (-)-epigallocatechin from enzymatic hydrolysate of its gallate using high-speed counter-current chromatography. *J. Chromatogr. A*, 687 (1994) 174-177.
- 2452 Durand, A., Gerbaud, A. and Margraff, R.: Centrifugal partition chromatography method for purifying taxoids. *PCT Int. Appl. WO 94 21, 622 (Cl. C07D305/14)*, 29 Sep. 1994, FR Appl. 93/3,251, 22 Mar. 1993; 14 p.; C.A., 122 (1995) 38810q.
- 2453 Graves, S.W. and Runyon, S.: Determination of methyleugenol in rodent plasma by high-performance liquid chromatography. *J. Chromatogr. B*, 663 (1995) 255-262.

- 2454 Kruger, P.B., de V. Albrecht, C.F., Liebenberg, R.W. and van Jaarsveld, P.P.: Studies on hypoxoside and rooperol analogues from *Hypoxis rooperi* and *Hypoxis latifolia* and their biotransformation in man by using high-performance liquid chromatography with in-line sorption enrichment and diode-array detection. *J. Chromatogr. B*, 662 (1994) 71-78.
- 2455 Lee, Y.-C., Huang, C.-Y., Wen, K.-C. and Suen, T.-T.: Determination of liquiritin, glycyrrhizin, hesperidin, cinnamic acid, cinnamaldehyde, honokiol and magnolol in the traditional Chinese medicinal preparation Wu-Ji-San by high-performance liquid chromatography. *J. Chromatogr. A*, 692 (1995) 137-145.
- 2456 Pande, A., Uniyal, G.C. and Shukla, Y.N.: Determination of chemical constituents of *Valeriana wallichii* by reverse phase HPLC. *Indian J. Pharm. Sci.*, 56 (1994) 56-58; C.A., 121 (1994) 225355m.
- 2457 Pietta, P.G., Mauri, P.L., Bruno, A. and Merfort, I.: MEKC as an improved method to detect calcifications in the flowers of *Arnica montana* and *A. chamissonis*. *Planta Med.*, 60 (1994) 369-372; C.A., 121 (1994) 238499m.
- 2458 Reif, K. and Metzger, W.: Determination of aflatoxins in medicinal herbs and plant extracts. *J. Chromatogr. A*, 692 (1995) 131-136.
- 2459 Sener, B. and Kucukboaci, N.: High-pressure liquid chromatographic determination of taxol in *Taxus baccata*. *J. Fac. Pharm. Gazi Univ.*, 11 (1994) 11-15; C.A., 121 (1994) 308452s.

For additional information see C.A.:

121 (1994) 213110q, 286692r, 286693s, 308454u;  
122 (1995) 64528y, 64553c.

See also 1521, 1553, 1554, 1558, 1609, 1704.

### 33. CLINICO-CHEMICAL APPLICATIONS

#### 33a. General papers and reviews

See 2179.

#### 33b. Complex mixtures and profiling (single compounds by cross-reference only)

For additional information see C.A.:

122 (1995) 4944w.

See also 1625, 1631, 1642, 1669, 1673, 1676, 1718, 1724, 1727, 1731, 1790, 1949, 1979, 2083, 2087, 2126, 2153, 2164, 2254.

### 34. FOOD ANALYSIS

#### 34a. General papers and reviews

2460 Bertrand, A.: (Primary chemical substances [in wine] currently determined by chromatography). *Acquiss. Recent. Chromatogr. Vin Cours Eur. Form. Contin.*, (1993) 1-16; C.A., 122 (1995) 8259z - a review with 8 refs.

2461 Nakazawa, H., Horie, M. and Shida, Y.: (Overview of liquid chromatography-mass spectrometry in food analysis). *Foods Food Ingredients J. Jpn.*, 159 (1994) 4-15; C.A., 121 (1994) 253954z - a review with 56 refs.

#### 34b. Complex mixtures (single compounds by cross-reference only)

2462 Bailey, R.G., Nursten, H.E. and McDowell, I.: A liquid chromatography-mass spectrometry study of a black tea liquor using the plasmaspay interface. *J. Sci. Food Agric.*, 66 (1994) 203-208; C.A., 121 (1994) 279188c.

2463 Calero Barcoj, J., Martinez Nieto, L. and Garcia-Granados Lopez de Hierro, A.: Countercurrent attraction process for recovering acids, phenols alcohol and their derivatives, from olive-pressing waste. *Span. ES 2,051,238* (Cl. B01D15/08), 01 Jun. 1994, Appl. 9,202,366, 24 Nov. 1992; 6 p.; C.A., 121 (1994) 208409q.

2464 Robards, K. and Antolovich, M.: Application of chromatography and pattern recognition to quality assessment of orange juice. *Chem. Aust.*, 61 (1994) 392-395; C.A., 121 (1994) 299413a.

2465 Sontag, G. and Bernwieser, I.: HPLC coupled with a colometric electrode-array detector. Determination of phloretin glucoside in juices for adulteration detection. *Dtsch. Lebensm.-Rundsch.*, 90 (1994) 72-73; C.A., 121 (1994) 253989q.

For additional information see C.A.:

121 (1994) 254006d, 254022f, 254024h, 279150j.

See also 1540, 1547, 1555, 1568, 1572, 1580, 1603, 1604, 1608, 1633, 1648, 1649, 1654, 1678, 1688, 1689, 1693, 1705, 1706, 1710, 1711, 1730, 1732, 1740, 1754, 1758, 1774, 1900, 1909, 2091, 2132, 2142, 2152, 2160, 2167, 2170, 2179, 2182, 2207, 2211, 2229, 2235, 2273, 2362, 2366, 2375, 2376, 2441, 2541.

#### 34c. Organoleptically important compounds (flavors, odors, volatiles)

See 1678.

### 35. ENVIRONMENTAL ANALYSIS

#### 35a. General papers and reviews

See 1379, 1436.

#### 35b. Air pollution (complex mixtures; single compounds by cross-reference only)

2466 Possanzini, M. and Di Palo, V.: Determination of volatile carbonyl compounds in air by HPLC analysis of fluorescent derivatives. *Comm. Eur. Communities, (Rep.)*, 2 (1994) 731-736; C.A., 121 (1994) 285552h.

For additional information see C.A.:

121 (1994) 307322n.

See also 1528, 2542.

35c. *Water pollution (complex mixtures; single compounds by cross-reference only)*

- 2467 Fiehn, O., Reemtsma, T. and Jekel, M.: Extraction and analysis of various benzothiazoles from industrial wastewater. *Anal. Chim. Acta*, 295 (1994) 297-305.
- 2468 Fuchs, F. and Heidt, A.: Application of hydrophobic interaction chromatography (HIC) in water analysis. *Acta Hydrochim. Hydrobiol.*, 22 (1994) 121-129; C.A., 121 (1994) 237937r.
- 2469 Nivet, J.M. and Dessin, H.: (Application of ion-exchange liquid chromatography to the analysis of anions in pharmaceutical-quality water). *S.T.P. Pharma Prat.*, 4 (1994) 177-192; C.A., 121 (1994) 238489h - a review with 2 refs.
- 2470 Volmer, D.: (Analysis of pesticides in aqueous environmental samples by thermospray HPLC-MS). *GI F Fachz. Lab.*, 38 (1994) 16-24; C.A., 121 (1994) 307773k.

See also 1490, 1529, 1548, 1563, 1745, 2215, 2216, 2218, 2221, 2223, 2224, 2225, 2227, 2230, 2233, 2483, 2501, 2511, 2517, 2531, 2544.

35d. *Soil pollution (complex mixtures; single compounds by cross-reference only)*

See 1537, 1734, 2222, 2224.

36. SOME TECHNICAL PRODUCTS AND COMPLEX MIXTURES

36a. *Surfactants*

- 2471 Jandera, P. and Urbanek, J.: Comparison of chromatographic behaviour of oligoethylene glycol nonylphenyl ether non-ionic and anionic surfactants in reversed-phase high-performance liquid chromatography. *J. Chromatogr. A*, 689 (1995) 255-267.
- 2472 Lemr, K., Zanette, M. and Marcomini, A.: Reversed-phase high-performance liquid chromatographic separation of 1-naphthyl isocyanate derivatives of linear alcohol polyethoxylates. *J. Chromatogr. A*, 686 (1994) 219-224.
- 2473 Trathnigg, B., Maier, B. and Thamer, D.: Analysis of polyethers by isocratic HPLC with universal detectors. III. A study on reproducibility. *J. Liq. Chromatogr.*, 17 (1994) 4285-4302.
- 2474 Ye, M.Y., Walkup, R.G. and Hill, K.D.: Determination of surfactant sodium lauryl ether sulfate by ion pairing chromatography with suppressed conductivity detection. *J. Liq. Chromatogr.*, 17 (1994) 4087-4097.

See also 1659.

36b. *Antioxidants and preservatives*

- 2475 Ashida, S., Noguchi, S.F. and Suzuki, T.: Antioxidative components, xanthone derivatives, in *Swertia japonica*, Makino. *J. Am. Oil Chem. Soc.*, 71 (1994) 1095-1099.
- 2476 Gopala Krishna, A.G. and Prabhakar, J.V.: Antioxidant constituents of peanut oil. *J. Am. Oil Chem. Soc.*, 71 (1994) 1245-1249.

- 2477 Motchnik, P.A., Frei, B. and Ames, B.N.: Measurement of antioxidants in human blood plasma. *Methods Enzymol.*, 234(Oxygen Radicals in Biological Systems, Pt. D) (1994) 269-279; C.A., 121 (1994) 296288c.
- 2478 Nerin, C., Salafranca, J., Cacho, J. and Rubio, C.: Separation of polymer and on-line determination of several antioxidants and UV stabilizers by coupling size-exclusion and normal-phase high-performance liquid chromatography columns. *J. Chromatogr. A*, 690 (1995) 230-236.
- 2479 Tian, L.L. and White, P.J.: Antipolymerization activity of oat extract in soybean and cottonseed oils under frying conditions. *J. Am. Oil Chem. Soc.*, 71 (1994) 1087-1094.

For additional information see C.A.:

122 (1995) 63961x.

See also 2290, 2425.

36c. *Complex mixtures, technical products and unidentified compounds*

- 2480 Alen, R.: Analysis of low-molecular-mass compounds as an aid for monitoring oxygen delignification. *Cellul. Chem. Technol.*, 27 (1993) 281-286; C.A., 121 (1994) 233288g.
- 2481 Balogh, D.T., Curvelo, A.A.S. and de Groote, R.A.M.C.: Molecular weight of organosolv lignins. In: Kennedy, J.F. and Phillips, G.O. (Editors), *Cellul.: Chem. Biochem. Mater Aspects*, Horwood, London, 1993, pp. 279-284; C.A., 122 (1995) 12333m.
- 2482 Erhan, S.Z. and Bagby, M.O.: Polymerization of vegetable oils and their uses in printing inks. *J. Am. Oil Chem. Soc.*, 71 (1994) 1223-1226.
- 2483 Karpuyk, A.D., Smirnova, G.B., Poletaeva, I.L., Betekhtin, S.F. and Ignatova, Z.N.: (Ion chromatography in monitoring of aqueous coolants). *Energetik*, (1994) 17-18; C.A., 121 (1994) 265972s.
- 2484 Lameloise, M.-L. and Lewandowski, R.: Purification of beet molasses by ion-exclusion chromatography: fixed-bed modelling. *J. Chromatogr. A*, 685 (1994) 45-52.
- 2485 Lord, B.S. and Stringham, R.W.: Monitoring formation of a diamine process intermediate by reversed-phase liquid chromatography. *Anal. Chem.*, 66 (1994) 4370-4374.
- 2486 Suckling, I.D., Pasco, M.F., Hortling, B. and Sundquist, J.: Assessment of lignin condensation by GPC analysis of lignin thioacidolysis products. *Holzforschung*, 48 (1994) 501-503; C.A., 122 (1995) 58502u.
- 2487 Xu, B., Madrali, E.S., Wu, F., Li, C.-Z., Herod, A.A. and Kandiyoti, R.: Characterization of successive extract fractions released from a sample of coal during liquefaction in a flowing-solvent reactor. *Energy Fuels*, 8 (1994) 1360-1369; C.A., 121 (1994) 234332d.
- 2488 Zeisler, P., Hamm, U. and Goettsching, L.: (Application of ion chromatography in the pulp and paper industry). *Papier (Darmstadt)*, 48 (1994) 18-26; C.A., 122 (1995) 58533e.

For additional information see C.A.:

121 (1994) 258552b, 270866q, 283079k, 291602r;  
122 (1995) 33436n, 38677b.

See also 1511, 1536, 1617, 2512.

## 37. CELLS, CELLULAR PARTICLES AND SUPRAMOLECULAR STRUCTURES

- 2489 Bernard, A., Bories, C., Loiseau, P.M. and Cardot, P.J.P.: Selective elution and purification of living *Trichomonas vaginalis* using gravitational field-flow fractionation. *J. Chromatogr. B*, 664 (1995) 444-448.
- 2490 Davis, L.S., McIlraith, M.J., Paecheo, T., Becker, B., Adix, L.M., Thomas, R., Wacholtz, M.C. and Lipsky, P.E.: Assessment of a positive selection technique using an avidin column to isolate human peripheral blood T cell subsets. *J. Immunol. Methods*, 175 (1994) 247-257; C.A., 121 (1994) 275965f.
- 2491 Lucas, L., García-Pérez, A.-I., Jimeno, P., Pérez, M.T., Pinilla, M., Sancho, P. and Luque, J.: Surface properties of crosslinked erythrocytes as studied by counter-current distribution in aqueous polymer two-phase systems. *J. Chromatogr. B*, 664 (1995) 137-144.
- 2492 Petric, R., Nicholson, D.W. and Ford-Hutchinson, A.W.: Renal leukotriene C<sub>4</sub> synthase: characterization, partial purification and alterations in experimental glomerulonephritis. *Biochim. Biophys. Acta*, 1254 (1995) 207-215.
- 2493 Tazuma, S., Ochi, H., Teramen, K., Yamashita, Y., Horikawa, K., Miura, H., Hirano, N., Sasaki, M., Aihara, N., Hatsushika, S., Tao, S., Ohya, T. and Kajiyama, G.: Degree of fatty acyl chain unsaturation in biliary lecithin dictates cholesterol nucleation and crystal growth. *Biochim. Biophys. Acta*, 1215 (1994) 74-78.
- 2494 Vemuri, S. and Rhodes, C.T.: Separation of liposomes by a gel filtration chromatographic technique: a preliminary evaluation. *Pharm. Acta Helv.*, 69 (1994) 107-113; C.A., 121 (1994) 308270f.

See also 1481, 2056.

## 38. INORGANIC COMPOUNDS

- 2495 Fischer, C.-H.: Analysis of colloids. VI. Semiconductor colloids of high monodispersity by preparative size exclusion chromatography. *J. Liq. Chromatogr.*, 17 (1994) 3593-3611.

## 38a. Cations

- 2496 Cheng, J., Liu, Q. and Liu, J.: (Platinum family metal analysis by high-performance liquid chromatography). *Fenxi Huaxue*, 22 (1994) 630-635; C.A., 121 (1994) 244306n - a review with 51 refs.
- 2497 Chuang, Y.S., Lee, K.W., Hwang, J.Y. and Lee, Y.M.: Separation of chromium(III) and chromium(VI) by carboxymethylated polyamine-polyurea resin column. *Anal. Sci. Technol.*, 7 (1994) 205-211; C.A., 121 (1994) 314693n.
- 2498 Chuveleva, E.A., Kharitonov, O.V. and Firsova, L.A.: (Modeling the behavior of curium and americium with separation by the method of elution-complexing agent chromatography). *Radiokhimiya*, 36 (1994) 167-170; C.A., 121 (1994) 266016v.
- 2499 Dadachova, E., Mirzadeh, S., Lambrecht, R.M., Hetherington, E.L. and Knapp, F.F., Jr.: Separation of carrier-free holmium-166 from neutron-irradiated dysprosium targets. *Anal. Chem.*, 66 (1994) 4272-4277.

- 2500 Donald, S., Spires, K. and Vincent, J.B.: Potential for decontamination of plutonium-containing solutions using transferrin metalloprotein affinity metal chromatography. *J. Inorg. Biochem.*, 56 (1994) 167-171; C.A., 121 (1994) 310102w.
- 2501 Groschner, M. and Appriou, P.: Three-column system for pre-concentration and speciation determination of trace metals in natural waters. *Anal. Chim. Acta*, 297 (1994) 369-376.
- 2502 Haldna, U. and Yakovleva, I.: Glycine-sodium carbonate eluents for ion chromatography. *Eesti Tead Akad. Toim., Keem.*, 43 (1994) 55-60; C.A., 122 (1995) 22654f.
- 2503 Inaba, K., Freiser, H. and Muralidharan, S.: Effect of kinetic factors on the efficiencies of centrifugal partition chromatographic separations of trivalent lanthanides with bis(2,4,4-trimethylpentyl)phosphinic acid extractant. *Solvent Extr. Res. Dev., Jpn.*, 1 (1994) 13-29; C.A., 122 (1995) 70802z.
- 2504 Janos, P.: Ion-interaction chromatographic separation of metal cations in the presence of complexing agents. *Fresenius J. Anal. Chem.*, 350 (1994) 646-648.
- 2505 Khuhawar, M., Lanjwani, S.N. and Khaskhely, G.Q.: High-performance liquid chromatographic determination of vanadium in crude petroleum oils using bis(salicylaldehyde)tetramethylethylenediamine. *J. Chromatogr. A*, 689 (1995) 39-43.
- 2506 Kieber, R.J. and Jones, S.B.: An undergraduate laboratory for the determination of sodium, potassium, and chloride: using an ion-exchange resin, ion chromatography, and atomic absorption spectrophotometry. *J. Chem. Educ.*, 71 (1994) A218, A220, A222; C.A., 121 (1994) 229900h.
- 2507 Kittel, N.E.V.D., Seuvart, A. and Wuench, G.: (Analysis for trace metals in high-purity refractory metals with matrix separation). *An. Quim.*, 90 (1994) 116-124; C.A., 121 (1994) 314732z.
- 2508 Korovin, V.Yu., Yagodin, G.A. and Saveleva, V.I.: (Separation of zirconium and hafnium with a solid extractant based on tributylphosphate). *Zh. Prikl. Khim. (S.-Peterburg)*, 67 (1994) 758-761; C.A., 122 (1995) 60663r.
- 2509 Mehra, M.C., Cormier, J.-G. and Ayikoe, K.: Ion chromatographic determination of cations with iodic acid. *Chim. Acta Turc.*, 22 (1994) 161-167; C.A., 122 (1995) 70924r.
- 2510 Mohite, B.S., Zambare, D.N. and Mahadik, B.E.: Separation of barium from associated elements using poly(dibenzo-18-crown-6) and column chromatography. *Analyst (Cambridge)*, 119 (1994) 2033-2036.
- 2511 Nagaosa, Y. and Segawa, S.: Reversed-phase HPLC determination of titanium(IV) and iron(II) with sodium 1,2-dihydroxybenzene-3,5-disulfonic acid. *J. High Resolut. Chromatogr.*, 17 (1994) 770-772.
- 2512 Padaruskas, A.V.: (Analysis of superconducting La-Sr-Cu based materials by ion-pair chromatography). *Izv. Vyssh. Uchebn. Zaved., Khim. Khim. Tekhnol.*, 36 (1993) 27-30; C.A., 121 (1994) 291455v.
- 2513 Panesar, K.S., Singh, O.V. and Tandon, S.N.: Liquid-liquid extraction and reversed phase chromatographic behavior of some 3d metal ions using bis(2,4,4-trimethylpentyl)dithiophosphinic acid (Cynex 301). *Talanta*, 41 (1994) 1341-1344; C.A., 122 (1995) 70799d.
- 2514 Pesavento, M., Biesuz, R. and Cortina, J.L.: Sorption of metal ions on a weak acid cation-exchange resin containing carboxylic groups. *Anal. Chim. Acta*, 298 (1994) 225-232.

- 2515 Pesavento, M., Profumo, A. and Sastre, A.: Chromatographic behavior of trace metal ions on a strong base anion exchange resin functionalized by azo ligands. *Talanta*, 41 (1994) 1689-1697; C.A., 122 (1995) 70804b.
- 2516 Pin, C., Briot, D., Bassin, C. and Poitrasson, F.: Concomitant separation of strontium and samarium-neodymium for isotopic analysis in silicate samples, based on specific extraction chromatography. *Anal. Chim. Acta*, 298 (1994) 209-217.
- 2517 Rottmann, L. and Heumann, K.G.: Determination of heavy metal interactions with dissolved organic materials in natural aquatic systems by coupling a high-performance liquid chromatography system with an inductively coupled plasma mass spectrometer. *Anal. Chem.*, 66 (1994) 3709-3715.
- 2518 Rubio, R., Padro, A. and Rauret, G.: LC-HG-QCAAS versus LC-HG-ICP/OES in arsenic speciation. *Fresenius J. Anal. Chem.*, 351 (1995) 331-333.
- 2519 San Andrés, M.P., Vera, S. and Marina, M.L.: Determination of Ni(II), Co(II) and Cu(II) as diethyldithiocarbamate complexes by high-performance liquid chromatography using hexadecyltrimethylammonium bromide in the mobile phase. *J. Chromatogr. A*, 685 (1994) 271-278.
- 2520 Słeboda, M., Wodecki, Z., Kolodziejczyk, A.M. and Nowicki, W.: Chromatographic separation of some transition metal ions on chelating column bearing acetylacetone moiety. *Chem. Anal. (Warsaw)*, 39 (1994) 149-152; C.A., 122 (1995) 45124x.
- 2521 Tomlinson, M.J., Wang, J. and Caruso, J.A.: Speciation of toxicologically important transition metals using ion chromatography with inductively coupled plasma mass spectrometric detection. *J. Anal. At. Spectrom.*, 9 (1994) 957-964; C.A., 121 (1994) 223766x.
- 2522 Umile, C. and Huber, J.F.K.: Determination of hexavalent chromium and accompanying anions in one run using isocratic ion chromatography with column switching. *Talanta*, 41 (1994) 1101-1106; C.A., 121 (1994) 270646t.
- 2523 Wu, J., Zhou, C.Y., Chi, H., Wong, M.K., Lee, H.K., Ong, H.Y. and Ong, C.N.: Determination of serum aluminium using an ion-pair reversed-phase high-performance liquid chromatographic-fluorimetric system with lumogallion. *J. Chromatogr. B*, 663 (1995) 247-253.
- 2524 Wuping, L. and Qiping, L.: Study on retention behaviour of platinum metal complexes in reversed-phase high-performance liquid chromatography. *Fresenius J. Anal. Chem.*, 350 (1994) 671-673.
- 2525 Yamane, T.: (Ion chromatography). *Materia*, 33 (1994) 341-345; C.A., 121 (1994) 244297k - a review with 19 refs.
- 2526 Yamane, T., Aruga, Y. and Ogawa, T.: Flow-injection system for sensitive and rapid determination of trace molybdenum(VI) and tungsten(VI) based on catalytic reaction and its application to ion chromatography. *Monatsh. Chem.*, 125 (1994) 20-24; C.A., 121 (1994) 220560q.
- 2527 Zhang, X. and Jiang, X.: Study on the combination of electrography and ion chromatography. *Int. Lab.*, 24, No. 11 (1994) 10-12.
- 2528 Zhang, X., Lin, C., Liu, L. and Qiao, G.: Determination of platinum by ion-pair reversed-phase high-performance liquid chromatography with 4,4'-bis(dimethylamino)thiobenzophenone. *J. Chromatogr. A*, 684 (1994) 354-359.
- 2529 Zhuravlev, A.A., Ribnov, V.V. and Ganiev, A.G.: (Separation of platinum and silver from hydrochloric acid solutions on aluminum oxide). *Dokl. Akad. Nauk Resp. Uzb.*, (1993) 37-38; C.A., 121 (1994) 314694p.
- For additional information see C.A.:  
121 (1994) 220424y, 220449k, 220508d, 225395z, 270551h, 314845p;  
122 (1995) 22648g, 22656h, 22756r, 11640x.
- See also 1296, 1364, 1454, 2150, 2385, 2544, 2545.
- 38b. Anions
- 2530 Albrich, H., Mueller, M. and Pinhack, H.: (Determination of fluorine, chlorine and bromine after ashing). *GIT Fachz. Lab.*, 38 (1994) 837-838; C.A., 121 (1994) 270850e.
- 2531 Grennaro, M.C., Abrigo, C., Saini, G., Muntau, H., Serrini, G. and Geiss, H.: Sulfate determination in lagoon water (Venice) by reversed-phase ion-interaction chromatography - a method validation. *Fresenius J. Anal. Chem.*, 350 (1994) 702-705.
- 2532 Hajos, P., Horvath, O. and Denke, V.: Prediction of retention for halide anions and oxoanions in suppressed ion chromatography using multiple species eluent. *Anal. Chem.*, 67 (1995) 434-441.
- 2533 Marengo, E., Gennaro, M.C., Abrigo, C. and Dinardo, A.: Chemometric optimization of azide sensitivity in reversed-phase ion-interaction HPLC. *Anal. Chem.*, 66 (1994) 4229-4235.
- 2534 McNeff, C., Zhao, Q. and Carr, P.W.: High-performance anion exchange of small anions with polyethyleneimine-coated porous zirconia. *J. Chromatogr. A*, 684 (1994) 201-211.
- 2535 Miura, Y., Tsubamoto, M. and Koh, T.: Ion chromatographic determination of sulphide, sulphite and thiosulphate in mixtures by means of their post column reactions with iodine. *Anal. Sci.*, 10 (1994) 595-600; C.A., 121 (1994) 220554r.
- 2536 Mura, P., Papet, Y., Sanchez, A. and Piriou, A.: Rapid and specific high-performance liquid chromatographic method for the determination of iodide in urine. *J. Chromatogr. B*, 664 (1995) 440-443.
- 2537 Popa, M.I.: High performance liquid chromatography of chloride in polymers. *Bul. Inst. Politeh. Iasi, Sect. 2: Chim. Ing. Chim.*, 39 (1993) 89-93; C.A., 122 (1995) 64565h.
- 2538 Stahl, R.: Routine determination of anions by capillary electrophoresis and ion chromatography. *J. Chromatogr. A*, 686 (1994) 143-148.
- 2539 Sumiyoshi, K., Yagi, T. and Nakamura, H.: Determination of cyanide by high-performance liquid chromatography using postcolumn derivatization with o-phthalaldehyde. *J. Chromatogr. A*, 690 (1995) 77-82.
- 2540 Vasconcelos, M.T.S.D., Gomes, C.A.R. and Machado, A.A.S.C.: Ion chromatographic determination of fluoride in welding fumes with elimination of high contents of iron by solid-phase extraction. *J. Chromatogr. A*, 685 (1994) 53-60.
- 2541 Yuan, Z., Dai, H. and Nie, H.: (HPLC determination of residue of inorganic bromide in oranges). *Fenxi Ceshi Xuebao*, 13 (1994) 53-57; C.A., 121 (1994) 299399a.
- For additional information see C.A.:  
121 (1994) 291441n, 291602r;  
122 (1995) 45320h, 45323m.

See also 1725, 1862, 2286, 2469, 2483, 2502, 2522.

38c. *Permanent and rare gases*

- 2542 Hekmat, M., Smith, R. and Fung, P.: An evaluation of the Occupational Health and Safety Administration method for the "determination of chlorine dioxide in workplace atmosphere". *Am. Ind. Hyg. Assoc. J.*, 55 (1994) 1087-1089; *C.A.*, 122 (1995) 37517n.

38d. *Volatile inorganic compounds*

- 2543 Gamoh, K. and Sakamoto, M.: (Indirect liquid chromatographic determination of hydroxyl radical based on the production of methanesulfinic acid). *Bunseki Kagaku*, 43 (1994) 691-696; *C.A.*, 121 (1994) 250073a.

See also 1352.

39. RADIOACTIVE AND OTHER ISOTOPE COMPOUNDS

- 2544 Cobb, J., Warwick, P., Carpenter, R.C. and Morrison, R.T.: Determination of strontium-90 in water and urine samples using ion chromatography. *Analyst (Cambridge)*, 119 (1994) 1759-1764.
- 2545 Vajda, N., Kis-Benedek, G., Bodizs, D. and Vodicska, M.: (Radiochemical determination of  $^{210}\text{Pb}$  by using a crown ether). *Izotoptech., Diagn.*, 37 (1994) 25-30; *C.A.*, 121 (1994) 270584w.

See also 1650, 1736, 2265, 2440.

## Gas Chromatography

### 1. REVIEWS AND BOOKS

- 717 Chakraborty, J.: Gas chromatography in clinical biochemistry. In: Baugh, P.J. (Editor), *Gas Chromatography*, IRL, Oxford, 1993, pp. 213-236; C.A., 121 (1994) 249886m - a review with 23 refs.
- 718 Grosch, W.: (New development in flavor research.) *Lebensmittelchemie*, 47, No. 6 (1993) 129-134; C.A., 121 (1994) 229044g.
- 719 Hinshaw, J.V. and Ettre, L.S.: *Introduction to Open-Tubular Column Gas Chromatography*. Advanced Communications, Cleveland, 1994, 190 p.
- 720 Szepes, L.: (Development and importance of separation techniques in instrumental analysis.) *Magy. Kem. Foly.*, 100 (1994) 340-353; C.A., 121 (1994) 270751y - a review with 121 refs.
- 721 Weston, P.C.: Gas chromatography - evolution and revolution. *TrAC*, 14, No. 2 (1995) VII-XII.

See also 766, 819, 951, 952, 953, 954, 957, 959, 1013, 1017, 1054, 1087, 1123, 1151, 1164, 1170.

### 2. FUNDAMENTALS, THEORY AND GENERAL

#### 2a. General

- 722 Felinger, A. and Guichon, G.: Computer simulations in non-linear chromatography. *TrAC*, 14 (1995) 6-10.
- 723 McNally, M.E.P. and Barefoot, A.C. III.: Chromatography: Future directions in the industrial laboratory. *Am. Lab. (Shelton)*, 26, No. 18 (1994) 28N-28W.
- 724 Meyer, V.R.: Errors in the area determination of incompletely resolved chromatographic peaks. *J. Chromatogr. Sci.*, 33 (1995) 26-33.
- 725 Reh, E.: Peak-shape analysis for unresolved peaks in chromatography: comparison of algorithms. *TrAC*, 14 (1995) 1-6.

#### 2b. Thermodynamics and theoretical relationships

- 726 Abraham, M.H., Andonian-Haftven, J., Du, C.M., Osei-Owusu, J.P., Sakellariou, P., Shuley, W.J., Poole, C.F. and Poole, S.K.: Comparison of uncorrected retention data on a capillary and a packed hexadecane column with corrected retention data on a packed squalane column. *J. Chromatogr. A*, 688 (1994) 125-134.

See also 827.

#### 2c. Relationship between structure and chromatographic behaviour

See 825, 846, 861, 1142.

#### 2d. Measurement of physico-chemical and related values

- 727 Cortesi, A., Kikic, I. and Spicka, B.: Supercritical fluid chromatography determination of partial molar volumes of alcohols. *Chem. Biochem. Eng. Q.*, 8 (1994) 17-20; C.A., 121 (1994) 187945u.
- 728 Dai, M.-G. and Yang, S.: (Studies on binary gas solid adsorption. (I). A new system for the rapid determination of adsorption isotherm with frontal chromatography.) *Gaodeng Xuexiao Huaxue Xuebao*, 15 (1994) 495-498; C.A., 121 (1994) 287066b.
- 729 Donnet, J.B., Wang, W., Vidal, A. and Wang, M.J.: Study of surface activity of carbon black by inverse gas chromatography. Part II: Effect of carbon black thermal treatment on its surface characteristics and rubber reinforcement. *Kautsch. Gummi, Kunstst.*, 46 (1993) 866-871; C.A., 121 (1994) 181608q.
- 730 Dulamita, N., Stanca, M. and Buciuman, F.: Characterization of Pd/Al<sub>2</sub>O<sub>3</sub> catalyst using the gas-chromatography method. *Prog. Catal.*, 2, No. 2 (1993) 15-19; C.A., 121 (1994) 264679q.
- 731 Grájek, H.: Gas chromatographic estimation of adsorption properties of activated carbons. *Process Technol. Proc.*, 11 (1994) 799-812; C.A., 121 (1994) 264490w.
- 732 Grajek, H. and Witkiewicz, Z.: Comparison of the chromatographic and static methods of the investigation of activated carbons. *Pol. J. Chem.*, 67 (1993) 911-921; C.A., 121 (1994) 238997x.
- 733 Khalfaoui, B. and Newsham, D.M.T.: Determination of second cross virial coefficients from gas-liquid chromatographic data. II. Dilute mixtures of water and brominated hydrocarbons. *J. Chromatogr. A*, 688 (1994) 117-123.
- 734 Martens, J.A., Ghys, D., van de Voorde, M., Verelst, H., Baron, G. and Jacobs, P.A.: Chromatographic study of the adsorption of O<sub>2</sub>, N<sub>2</sub>, Ar and CH<sub>4</sub> in cation exchanged A, X and Y zeolites. *Process Technol. Proc.*, 11 (1994) 819-824; C.A., 121 (1994) 264491x.
- 735 Qin, R.-Y. and Schreiber, H.P.: Application of inverse gas chromatography to molecular diffusion in polymers. *Langmuir*, 10 (1994) 4153-4156; C.A., 121 (1994) 256779g.
- 736 Shcherbina, A.E., Dolinskaya, R.M. and Shcherbina, E.I.: (Use of gas-chromatographic technique for comparative evaluation of extractive properties of new binary solvents.) *Neftekhimia*, 34 (1994) 280-284.
- 737 Spicka, B., Cortesi, A., Fermeglia, M. and Kikic, I.: Determination of partial molar volumes at infinite dilution using SFC technique. *J. Supercrit. Fluids*, 7 (1994) 171-176.
- 738 Wang, W., Vidal, A., Donnet, J.-B. and Wang, M.-J.: Study of surface activity of carbon black by inverse gas chromatography, Part III. Supercritical plasma treatment of carbon black and its surface activity. *Kautsch. Gummi Kunstst.*, 46 (1993) 933-934; C.A., 121 (1994) 233919p.
- 739 Webster, H.F. and Dillard, J.G.: Analysis of phosphonic acid treated aluminium oxide surfaces by inverse gas chromatography. *Polym. Mater. Sci. Eng.*, 70 (1993) 501-502; C.A., 121 (1994) 260977n.

- 740 Xu, Y., Xu, L., Sun, M. and Lu, J.: (Study of reaction degree for isoprene-maleic anhydride addition reaction by capillary gas chromatography.) *Shiyou Huagong* 23 (1994) 324-328; C.A., 121 (1994) 183079s.
- 741 Yoshino, A., Murate, K., Yoshida, T., Okabayashi, H., Krishna, P.R., Kamaya, H. and Ueda, I.: Surface-oriented saturable binding of halothane with micelles: paramagnetic relaxation of  $^{19}\text{F}$  NMR spin-lattice relaxation rate and gas chromatography studies. *J. Colloid Interface Sci.*, 166 (1994) 375-382; C.A., 121 (1994) 221725J.
- 742 Zhong, B. and Yang, N.: (Study on hydration rate of cement and polymeric structure of  $[\text{SiO}_4]^{4-}$  tetrahedra using  $^{29}\text{Si}$ -NMR and TMS-GC.) *Nanjing Huagong Xueyuan Xuebao*, 16, No. 3 (1994) 26-32; C.A., 121 (1994) 236983r.

See also 817, 1080, 1086.

### 3. GENERAL TECHNIQUES

#### 3a. Apparatus and accessories

- 743 Annino, R., Lewis, E.L., Lueck, D.E., Phillips, M.L. and Villalobos, R.: Modular gas chromatography device. U.S. US 5,340,543 (Cl. 422-89; G01N30/02), 23 Aug. 1994, Appl. 570,947, 22 Aug. 1990; 26 pp.; C.A., 121 (1994) 220692j.
- 744 Arnold, M. and Kleiböhmer, W.: A simple large volume injection system with a solid-phase sample loop for packed column supercritical fluid chromatography. *Chromatographia*, 39 (1994) 612-618.
- 745 Bot, D.S.M. and Schrijvers, H.L.: Dual-column gas chromatography retrieval system. *Dev Food Sci.*, 35 (1994) 75-88; C.A., 121 (1994) 279177y.
- 746 Boyd-Boland, A.A., Chai, M., Luo, Y.Z., Zhang, Z., Yang, M.J., Pawliszyn, J.B. and Görecki, T.: New solvent-free sample preparation techniques based on fiber and polymer technologies. *Environ. Sci. Technol.*, 28, No. 13 (1994) 569A-574A.
- 747 Byrd, D.W. and Freeman, D.C.: Automated static headspace sampler for gas chromatography. *J. Chromatogr. A*, 686 (1994) 235-243.
- 748 Cheh, C.H.S., Chang, S.D., Weng, C., Tang, Y., Holzhuetter, I.B. and Hawthorne, S.H.: Gas chromatographic method and apparatus. *Eur. Pat. Appl.* EP 602,924 (Cl. G01N30/30), 22 Jun. 1994, US Appl. 990,054, 14 Dec. 1992; 13 pp.; C.A., 121 (1994) 244480g.
- 749 Grob, K. and Fröhlich D.: Splitless injection of large volumes of aqueous samples: Further experience. Experiments on analysing triazines by direct injection of drinking water. *J. High Resolut. Chromatogr.*, 17 (1994) 792-794.
- 750 Harrington, P.D.B., Whittenberg, H.P.: Gas chromatography sample injector and apparatus using it. *PCT Int. Appl.* WO 94 20,848 (Cl. G01N30/18), 15 Sep. 1994, US Appl. 27,282, 05 Mar. 1993; 42 pp.; C.A., 121 (1994) 291689z.
- 751 Hong, L. and Wang, M.: (A convenient cold on-column injection method for capillary gas chromatography.) *Fenxi Ceshi Xuebao*, 13 (1994) 62-65; C.A., 121 (1994) 244527k.
- 752 Janak, K., Colmsjö, A. and Östman, C.: Increased on-column injection temperature for gas chromatography. *J. High Resolut. Chromatogr.*, 18 (1995) 117-120.

- 753 Klick, S.: Evaluation of different injection techniques in the gas chromatographic determination of thermolabile trace impurities in a drug substance. *J. Chromatogr. A*, 689 (1995) 69-76.
- 754 Mol, H.G.J., Hendriks, P.J.M., Janssen, H.-G., Cramers, C.A. and Brinkman, U.A.T.: Large volume injection in capillary GC using PTV injectors: comparison of inertness of packing materials. *J. High Resolut. Chromatogr.*, 18 (1995) 124-128.
- 755 Mol, H.G.J., Janssen H.-G., Cramers, C.A. and Brinkman, U.A.T.: Large volume sample introduction using temperature programmable injectors: Implications of linear diameter. *J. High Resolut. Chromatogr.*, 18 (1995) 19-27.
- 756 Morabito, P.L., McCabe, T. and Hiller, J.F.: Vented capillary gas chromatography apparatus. U.S. US 5,338,514 (Cl. 422-89; G01N30/00), 16 Aug. 1994, Appl. 111,808, 25 Aug. 1993; 9 pp; C.A., 121 (1994) 244472p.
- 757 Nakagawa, K., Miyoshi, S. and Fukushima, T.: (Gas chromatograph and method of using it.) U.S. US 5,339,673 (Cl. 73-23.36; G01N30/10), 23 Aug. 1994, JP Appl. 92/114,159, 06 Apr. 1992; 5 pp.; C.A., 121 (1994) 244476t.
- 758 Schubert, R., Pfaff, H., Mills, S. and Wagner, B.: Validation of a novel HRGC/HRMS instrument and the TACO user interface in respect to USEPA methods 1613 and 8290. *Organohalogen Compd.*, 11 (1993) 109-114; C.A., 121 (1994) 186604v.
- 759 Shude, C.: (Industrial chromatograph in a comprehensive utilization unit of liquefied petroleum gas.) *Huagong Zidonghua Ji Yibiao*, 20, No. 4 (1993) 41-48; C.A., 121 (1994) 259292d.
- 760 Wessels, P. and Schwinger, G.: Pressure-balanced headspace sampling using an electronic pressure-controlled, on column injector. *LC-GC Int.*, 8 (1995) 39-42.
- 761 Zheng, Y. and Shi, J.: (The application of bunched retention gap interface to ordinary reversed-phase high performance liquid chromatography (RP HPLC)/capillary gas chromatography (CGC).) *Chin. J. Chromatogr.*, 12 (1994) 395-397.

See also 823, 831, 835, 900, 1127.

#### 3b. Detectors and detection reagents

- 762 Acree, T.E. and Barnard, J.: Gas chromatography-olfactometry and CharmAnalysis. *Dev. Food Sci.*, 35 (1994) 211-220; C.A., 121 (1994) 225282k - a review with 21 refs.
- 763 Aue, W.A., Eisener, C.G., Gebhardt, J.A. and Lowery, N.B.: Holophotal flame photometric detection. *J. Chromatogr. A*, 688 (1994) 153-159.
- 764 Aue, W.A., Singh, H. and Sun, X.-Y.: Fundamental noise in three chromatographic detectors. *J. Chromatogr. A*, 687 (1994) 283-290.
- 765 Cai, H.: The development and characterization of pulsed discharge helium ionization/electron capture detectors for use in a gas chromatography system. Avail. *Univ. Microfilms Int.*, Order No. DA9415277, 1993, 146 pp.; C.A., 121 (1994) 220377k.
- 766 Colmsjö, A.L. and Berngaard, A.K.: Dual detection (PID-FID) in gas chromatography for the identification of PAH. *Polycyclic. Aromat. Compd.*, 3 (1993) 313-320; C.A., 121 (1994) 220604g - a review with 12 refs.
- 767 Da Silva, M.A.A.P., Lundahl, D.S. and McDaniel, M.R.: The capability and psychophysics of Osme: a new GC-olfactometry technique. *Dev Food Sci.*, 35 (1994) 191-209; C.A., 121 (1994) 279215j.

- 768 Farnsworth, P.B., Wu, M., Tacquard, M. and Lee, M.L.: Background correction device for enhanced element-selective gas chromatographic detection by atomic emission spectroscopy. *Appl. Spectrosc.*, 48 (1994) 742-746.
- 769 Ishii, M., Mutoh, M. and Nakamura, T.: A chemiluminescence reaction analyzer for a supercritical fluid. *Bunseki Kagaku*, 44 (1995) 91-96.
- 770 Jiang, G. and Xu, F.: (New flame photometric detector for gas chromatography.) *Huaxue Tongbao*, 3 (1994) 28-29; C.A., 121 (1994) 220631p.
- 771 Liu, E., Hu, Y. and Sun, J.: (Study on the relative response of flame ionization detector (FID) by regression analysis.) *Chin. J. Chromatogr.*, 12 (1994) 414-416.
- 772 Osipova, O.A., Kirichenko, V.E. and Pashkevich, K.I.: (Sensitivity of flame-ionization and electron-capture detectors to perfluoroacyl derivatives of amino, hydroxy, and mercapto compounds.) *Zh. Anal. Khim.*, 49 (1994) 715-718.
- 773 Pedersen-Bjergaard, S. and Greibrokk, T.: Evaluation of a low-resolution near-IR monochromator for atomic emission detection in capillary gas chromatography. *J. High Resolut. Chromatogr.*, 18 (1995) 9-14.
- 774 Pedersen-Bjergaard, S. and Greibrokk, T.: N-, O- and P-selective on column atomic emission detection in capillary gas chromatography. *J. Chromatogr. A*, 686 (1994) 109-119.
- 775 Singh, H., Millier, B. and Aue, W.A.: Constant-current a.c. electron-capture detector. *J. Chromatogr. A*, 690 (1995) 103-108.
- 776 Singh, H., Millier, B. and Aue, W.A.: Dual-channel response ratios from an integrative algorithm. *J. Chromatogr. A*, 687 (1994) 291-301.
- 777 Singh, H., Millier, B. and Aue, W.A.: Polarization, relaxation and unrestrictedly linear response in a bipolar, constant-frequency electron-capture detector. *J. Chromatogr. A*, 689 (1995) 45-61.
- 778 Sun, X.-Y., Singh, H., Millier, B., Warren, C.H. and Aue, W.A.: Noise, filters and detection limits. *J. Chromatogr. A*, 687 (1994) 259-281.
- 779 Valente, A.L.P. and Uden, P.C.: Comparison of a combined helium-argon plasma with pure helium plasmas for gas chromatography with atomic emission detection. *Analyst (Cambridge)*, 120 (1995) 419-421.
- 780 Wentworth, W.E., Cai, H. and Stearns, S.: Pulsed discharge helium ionization detector. Universal detector for inorganic and organic compounds at the low picogram level. *J. Chromatogr. A*, 688 (1994) 135-152.
- See also 834, 1016, 1030, 1174.
- 3c. *Sorbents and columns, packing procedures*
- 781 Forgács, E. and Cserhádi, T.: Retention strength and selectivity of porous graphitized carbon columns. Theoretical aspects and practical applications. *TrAC*, 14 (1995) 23-29.
- 782 Gonzales-Lara, R. and Gonzales, L.M.: (Methods for monitoring the quality in fabricating capillary columns for gas chromatography.) *Tec. Lab.*, 16 (1994) 178-185; C.A., 121 (1994) 244528m.
- 783 Heijmans, H., de Zeeuw, J., Buyten, J. Peene, J. and Mohnke, M.: The use of PLOT columns in gas chromatography. *Curr. Top. Membr.*, 41 (1994) 19-42; C.A., 121 (1994) 270506x.
- 784 Johnson, D.F.: Synthesis of novel polar and chiral polysiloxane stationary phases for gas and supercritical fluid chromatography. Avail. *Univ. Microfilms Int.*, Order No. DA9210202, 1991, 287 pp.; C.A., 121 (1994) 180820r.
- 785 Liu, X., Lou, X. and Zhou, L.: (Cyclodextrin derivatives as chiral stationary phases in high resolution capillary gas chromatography.) *Huaxue Shiji*, 16 (1994) 90-95; C.A., 121 (1994) 194557x.
- 786 Moriguchi, S., Naito, K. and Takei, S.: Adsorption effects on retention behaviour of hydrocarbons in gas-liquid and gas solid chromatography with the use of modified alumina coated with diphenyl phthalate as column packings. *J. Chromatogr. A*, 686 (1994) 101-108.
- 787 Ruan, Z. and Liu, H.: (An equation for calculation of the layer thickness in preparing porous polymer PLOT column by *in situ* copolymerization.) *Chin. J. Chromatogr.*, 13, No. 1 (1995) 1-3.
- 788 Steenackers, D. and Sandra, P.: Capillary GC on 50 micrometer i.d. columns coated with thick films. Theory and selected practical results. *J. High Resolut. Chromatogr.*, 18 (1995) 77-82.
- 789 Takayama, Y. and Takeichi, T.: Preparation of deactivated metal capillary for gas chromatography. *J. Chromatogr. A*, 685 (1994) 61-78.
- 790 Theory, L., Xu, G.-W., Zhang, Y.-K. and Lu, P.-Z.: (Intelligent optimization of column temperatures in gas chromatography.) *Huaxue Xuebao*, 52 (1994) 902-909; C.A., 121 (1994) 291656m.
- 791 Wasiak, W.: Chemically bonded chelates as selective complexing sorbents for gas chromatography. III. Silica chemically modified by N-benzoylthiourea groups. *J. Chromatogr. A*, 690 (1995) 93-102.
- 792 Wasiak, W., Voelkel, A. and Rykowska, I.: Physico-chemical characterization of chemically bonded stationary phases including metal complexes by inverse gas chromatography. *J. Chromatogr. A*, 690 (1995) 83-91.
- 793 Xu, G.-W., Zhang, X.-M., Yang, L., Zhang, Y.-K. and Lu, P.-Z.: (Intelligent optimization of column temperatures in gas chromatography. II. Program setup theory and application.) *Huaxue Xuebao*, 52 (1994) 910-916; C.A., 121 (1994) 291657n.
- 794 Xu, H. and Hong, H.: (Study on the tangential gradient column of gas chromatography.) *Xiangtan Daxue Ziran Kexue Xuebao*, 16, No. 2 (1994) 29-36; C.A., 121 (1994) 270801q.
- 795 Zhang, G., Xie, H., Ma, Y. and Liu, H.: (Coating of polar and highly polar stationary phase on flexible glass capillary column stationary phase.) *Fenxi Huaxue*, 22 (1994) 485-487; C.A., 121 (1994) 291632a.
- 796 Zhou, X., Yan, H., Wu, C. and Chen, Y.: (Two new chiral polysiloxanes containing chiral crown ether for capillary gas chromatography (CGC) separation of enantiomers.) *Chin. J. Chromatogr.*, 12 (1994) 404-405.
- See also 719, 816, 818, 830, 861, 862, 867, 877, 1047, 1097.
- 3d. *Quantitative analysis*
- 797 Freeman, D.C. and Byrd, D.W.: Statistical treatment of large digital chromatographic data sets. *J. Chromatogr. A*, 686 (1994) 225-233.
- See also 724, 875, 966, 1156.

3e. *Preparative scale chromatography*

See 826.

3f. *Programmed temperature, pressure, vapors, gradients*

- 798 Ferchichi, L., Nouar, L., Gheid, A. and Messadi, D.: (Advantages of retention indexes calculated by cubic spline interpolation in temperature-programmed gas chromatography.) *Bull. Soc. Chim. Belg.*, 103 (1994) 151-156; C.A., 121 (1994) 291659q.
- 799 Jain, V. and Phillips, J.B.: Fast temperature programming on fused-silica open-tubular capillary columns by direct resistive heating. *J. Chromatogr. Sci.*, 33 (1995) 55-59.
- 800 Santiuste, J.M.: Utilization of Rohrschneider's concept and Tekler equation in linear temperature programmed GC. Precalculation of retention data. *Chromatographia*, 40 (1995) 28-32.

See also 794, 836, 846, 863, 874.

## 4. SPECIAL TECHNIQUES

4b. *Computerization and modelling*

- 801 Du, H., Lahiri, S., Huang, G. and Stillman, M.J.: Developing an expert system for diagnosis of problem gas chromatographic data. *Anal. Chim. Acta*, 296 (1994) 21-31.

See also 722, 725, 1092, 1129.

4c. *Combination with other physico-chemical techniques (MS, IR etc.)*

- 802 Baykut, G.: Modular sampling and inlet systems for mobile environmental mass spectrometry. *TrAC*, 14 (1995) 10-23.
- 803 Bücherl, T., Eschler, M., Gruner, A., Palibroda, N. and Wolff, E.: Direct coupling of capillary supercritical fluid chromatography with double-focusing high resolution mass spectrometry. *J. High Resolut. Chromatogr.*, 17 (1994) 765-769.
- 804 Dirinck, P. and de Winne, A.: Advantages of instrumental procedures for measurement of flavor characters. *Dev Food Sci.*, 35 (1994) 259-265; C.A., 121 (1994) 279183x.
- 805 Goodman, K.J. and Brenna, J.T.: High-precision gas chromatography-combustion isotope ratio mass spectrometry at low signal levels. *J. Chromatogr. A*, 689 (1995) 63-68.
- 806 McClure, G.L.: General methods of sample preparation for infrared hyphenated techniques. In: Coleman, P.B. (Editor), *Pract. Sampling Tech. Infrared Anal.*, CRC, Boca Raton, 1993, pp. 165-215; C.A., 121 (1994) 220605h - a review with 80 refs.
- 807 Merritt, D.A., Brand, W.A. and Hayes, J.M.: Isotope-ratio-monitoring gas chromatography-mass spectrometry: methods for isotopic calibration. *Ogr. Geochem.*, 21 (1994) 573-583; C.A., 121 (1994) 291665p.
- 808 Mondello, L., Dugo, P., Bartle, K.D., Frere, B. and Dugo, G.: On-line high performance liquid chromatography coupled with high resolution gas chromatography and mass spectrometry (HPLC-HRGC-MS) for the analysis of complex mixtures containing highly volatile compounds. *Chromatographia*, 39 (1994) 529-538.

- 809 Rozenblum, A. and Brunerie, P.: Automatic processing of GC-MS analysis using a two-dimensional search system. *Dev Food Sci.*, 35 (1994) 133-137; C.A., 121 (1994) 279180u.

See also 756, 758, 847, 930, 950, 971, 997, 1031, 1035, 1052, 1183.

4e. *Functional analysis*

- 810 Dasgupta, A., Thompson, W.C. and Malik, S.: Use of microwave irradiation for rapid synthesis of perfluorooctanoyl derivatives of fatty alcohols, a new derivative for gas chromatography-mass spectrometric and fast atom bombardment mass spectrometric study. *J. Chromatogr. A*, 685 (1994) 279-285.

4f. *Trace analysis and pre separation techniques*

- 811 Barcarolo, R. and Casson, P.: Volatile enrichment on automatic static headspace using a precolumn and stopping the carrier gas flow during injection. *J. High Resolut. Chromatogr.*, 18 (1995) 5-8.
- 812 Fatoki, O.S. and Ogunfowokan, A.O.: Procedural cleanup technique for determination of phthalate esters in an aquatic environment. *Int. J. Environ. Stud.*, 44 (1993) 237-243; C.A., 121 (1994) 263070x.
- 813 Ho, J.S., Tang, P.H., Eichelberger, J.W. and Budde, W.L.: Liquid-solid disk extraction followed by SFE and GC-ion-trap MS for the determination of trace organic pollutants in water. *J. Chromatogr. Sci.*, 33 (1995) 1-8.
- 814 Meyer, V.R.: Quantitation of chromatographic peaks in the 0.1 to 1.0 % range. *Chromatographia*, 40 (1995) 15-22.

See also 780, 927, 946, 1052, 1071, 1152, 1174.

4g. *Enantiomers, separation*

- 815 Chang, A.C.-D.: Centrifugal partition, liquid, and gas chromatography separation of biologically important chiral molecules. Avail. *Univ. Microfilms Int.*, Order No. DA9417885, 1993, 139 pp.; C.A., 121 (1994) 270805u.
- 816 Dönnecke, J., König, W.A., Gyllenhaal, O., Vessman, J. and Schulze, C.: Enantiomer separation by capillary SFC and GC on immobilized octakis(2,6-di-O-methyl-3-O-pentyl)- $\gamma$ -cyclodextrin. *J. High Resolut. Chromatogr.*, 17 (1994) 779-783.
- 817 Jung, M., Fluck, M. and Schurig, V.: Enantiomerization of 2,2'-diisopropylbiphenyl during chiral inclusion gas chromatography: determination of the rotational energy barrier by computer simulation of dynamic chromatographic elution profiles. *Chirality*, 6 (1994) 510-512; C.A., 121 (1994) 230083a.
- 818 Liu, X.-L., Guo, F.-Q., Lou, X.-W. and Zhou, L.-M.: (A new chiral stationary phase-heptakis(2,6-di-O-n-butyl-3-O-trifluoroacetyl)- $\beta$ -cyclodextrin) heptakis(2,6-di-O-n-butyl-3-O-trifluoroacetyl)- $\beta$ -cyclodextrin.) *Huaxue Xuebao*, 52 (1994) 588-594; C.A., 121 (1994) 270788r.
- 819 Srinivas, N.R., Shyu, W.C. and Barbhuiya, R.H.: Gas chromatography determination of enantiomers as diastereomers following pre-column derivatization and applications to pharmacokinetic studies: A review. *Biomed. Chromatogr.*, 9 (1995) 1-9 - a review with 79 refs.

See also 785, 796, 918, 931, 982, 983, 1043, 1115.

#### 4h. Other special techniques

- 820 Monnet, C., Schmitt, P. and Divies, C.: Method for assaying volatile compounds by headspace gas chromatography and application to growing starter cultures. *J. Dairy Sci.*, 77 (1994) 1809-1815; C.A., 121 (1994) 229135n.

See also 832.

#### 4i. Supercritical fluid chromatography

- 821 Ashraf-Khorassani, M. and Levy, J.M.: Addition of modifier in supercritical fluid chromatography using a microbore reciprocating pump. *Chromatographia*, 40 (1995) 78-84.
- 822 Daimon, H. and Hirata, Y.: Direct coupling of capillary supercritical fluid chromatography with supercritical fluid extraction using modified carbon dioxide. *J. High Resolut. Chromatogr.*, 17 (1994) 809-813.
- 823 Hancock, S.A.: A novel device for cooling HPLC pump heads pumping supercritical fluids. *J. High Resolut. Chromatogr.*, 17 (1994) 795-796.
- 824 Hansen, S.L. and Artz, W.E.: Supercritical fluid fractionation of thermally oxidized canola oil. *J. Am. Oil Chem. Soc.*, 71 (1994) 615-618; C.A., 121 (1994) 229172x.
- 825 Heaton, D.M., Bartle, K.D., Clifford, A.A., Klee, M.S. and Berger, T.A.: Retention prediction based on molecular interaction in packed-column supercritical fluid chromatography. *Anal. Chem.*, 66 (1994) 4253-4257.
- 826 Hirata, Y., Kawaguchi, Y. and Kitano, K.: Large volume injection for preparative supercritical fluid chromatography. *Chromatographia*, 40 (1995) 42-46.
- 827 Köhler, U. and Klesper, E.: Influence of linear velocity, column length, and pressure drop in SFC. II. Plate numbers, effective plate numbers, and resolutions. *J. Chromatogr. Sci.*, 32 (1994) 525-534.
- 828 Mulcahey, L.J., Rankin, C.L. and McNally, M.E.P.: Environmental applications of supercritical fluid chromatography. *Adv. Chromatogr. (N.Y.)*, 34 (1994) 251-308; C.A., 121 (1994) 237653v.
- 829 Neu, P. and Hilscher, W.: (High-purity carbon dioxide for supercritical extractions.) *Gas Aktuell*, 47 (1994) 21-25; C.A., 121 (1994) 220404s.
- 830 Shariff, S.M., Tong, D. and Bartle, K.D.: Simulated distillation by supercritical fluid chromatography on packed capillary columns. *J. Chromatogr. Sci.*, 32 (1994) 541-546.
- 831 Thompson, P.G. and Taylor, L.T.: Liquid trapping after supercritical fluid extraction with modified carbon dioxide. *J. High Resolut. Chromatogr.*, 17 (1994) 759-764.
- 832 Vejrosta, J., Planeta, J., Mikešová, M., Ansorgová, A., Karásek, P., Fanta, J. and Janda, V.: Solute collection after off-line supercritical fluid extraction into a moving liquid layer. *J. Chromatogr. A*, 685 (1994) 113-119.
- 833 Vérillon, F., Hamstra, A. and Netter, C.: Enhanced chromatography with fluids containing carbon dioxide under automated pressure control. *LC-GC Int.*, 7 (1994) 710-716.
- 834 Wang, J. and Marshall, W.D.: Metal speciation by supercritical fluid extraction with on-line detection by atomic absorption spectrometry. *Anal. Chem.*, 66 (1994) 3900-3907.

- 835 Yang, Y., Hawthorne, S.B. and Miller, D.J.: Estimating flow-rates for sub- and supercritical fluid extractions with linear restrictors. *J. Chromatogr. A*, 690 (1995) 131-139.

- 836 Zhou, L.-M., Li, Z.-S., Wang, Q.-H. and Zhu, D.-Q.: (Approach to the theory of linear density programming in supercritical fluid chromatography.) *Huaxue Xuebao*, 52 (1994) 797-806; C.A., 121 (1994) 244519j.

See also 727, 737, 744, 769, 784, 803, 816, 849, 858, 859, 862, 881, 888, 902, 924, 933, 954, 962, 981, 984, 987, 994, 1031, 1039, 1065, 1091, 1097, 1156.

## 5. HYDROCARBONS AND HALOGEN DERIVATIVES

### 5a. Aliphatic hydrocarbons

- 837 Bergaentzle, M., Sanquer, F., Hasselmann, C. and Marchioni, E.: Detection of  $\gamma$ -irradiated raw-milk Camembert cheeses by capillary gas chromatographic analysis of volatile hydrocarbons. *Food. Chem.*, 51 (1994) 177-182; C.A., 121 (1994) 178043w.
- 838 Dégoussée, N., Triantaphylidés, C., Starek, S., Iacazio, G., Martini, D., Bladier, C., Voisine, R. and Montillet, J.-C.: Measurement of thermally produced volatile alkanes: An assay for plant hydroperoxy fatty acid evaluation. *Anal. Biochem.*, 224 (1995) 524-531.
- 839 Denha, A.M., Bartle, K.D. and Pilling, M.J.: Simplified gas chromatographic procedures for the determination of  $C_1$ - $C_7$  hydrocarbons in urban atmospheres. *Anal. Proc.*, 31 (1994) 329-331.
- 840 Jemma, C.A., Shore, P.R. and Widdicombe, K.A.: Analysis of  $C_1$ - $C_{16}$  hydrocarbons using dual-column capillary GC: Application to exhaust emissions from passenger car and motorcycle engines. *J. Chromatogr. Sci.*, 33 (1995) 34-48.
- 841 Mennie, D., Moffat, C.F. and McGill, A.S.: Identification of sterene compounds produced during the processing of edible oils. *J. High Resolut. Chromatogr.*, 17 (1994) 831-838.
- 842 Sojak, L. and Ostrovsky, I.: (Method of the separation and analysis of isomers containing from 7 to 20 carbon atoms in molecule and having a functional group in the middle of the carbon chain of the molecule.) *Czech. CS 277,050 (Cl. G01N30/00)*, 18 Nov. 1992, Appl. 89/5,276, 13 Sep. 1989; 5 pp.; C.A., 121 (1994) 270870m.
- 843 Szafranek, J., Malinski, E., Dubis, E., Hebanowska, E., Nawrot, J., Oksman, P. and Pihlaja, K.: Identification of branched alkanes in lipids of *Leptinotarsa decemlineata* Say and *Tribolium destructor* by GC-MS: a comparison of main-beam and link-scanned spectra. *J. Chem. Ecol.*, 20 (1994) 2197-2212; C.A., 121 (1994) 276066a.
- 844 Xu, X., Qu, H., Chang, Y. and Lü, J.: (Study on the method for determination of content of conjugated dienes in petroleum products by combination of diene synthesis reaction with gas chromatography. I. Packed column gas chromatography.) *Chin. J. Chromatogr.*, 13, No. 1 (1995) 56-58.
- 845 Zhu, L., Zhu, R. and Chen, L.: (Low-temperature enrichment method for the determination of trace lower hydrocarbons in air.) *Shanghai Huanjing Kexue*, 12, No. 5 (1993) 21-22; C.A., 121 (1994) 185837m.

See also 759, 786, 862.

#### 5b. Cyclic hydrocarbons, fullerenes

- 846 Bemgard, A.M., Colmsjö, A. and Wrangskog, K.: Prediction of temperature-programmed retention indexes for polynuclear aromatic hydrocarbons in gas chromatography. *Anal. Chem.*, 66 (1994) 4288-4294.
- 847 Budzinski, H., Pierard, C., Garrigues, P. and Bellocq, J.: Analysis for aromatic compounds by capillary gas chromatography/Fourier transform infrared spectroscopy. *Polycyclic Aromat. Compd.*, 3 (1993) 329-335; *C.A.*, 121 (1994) 220668f.
- 848 Elke, A. and Mueller, A.: A simple method of sample preparation for analysis of biphenyl residues in citrus fruit peels by gas chromatography. *Z. Lebensm.-Unters. Forsch.*, 198 (1994) 329-330; *C.A.*, 121 (1994) 177962h.
- 849 Heaton, D.M., Bartle, K.D., Clifford, A.A., Myers, P. and King, B.W.: Rapid separation of polycyclic aromatic hydrocarbons by packed column supercritical fluid chromatography. *Chromatographia*, 39 (1994) 607-611.
- 850 Huang, W. and Lin, Y.: (Analysis of styrene in atmosphere by gas chromatographic method.) *Shanghai Huanjing Kexue*, 12, No. 11 (1993) 18-20; *C.A.*, 121 (1994) 262176z.
- 851 Karabanov, N.T., Vetrova, Z.P., Shuvalova, T.N., Ivanova, L.A., Gur'ev, I.A. and Kutsovskaya, V.V.: (Extraction-gas-chromatographic analysis of environmental samples.) *Zh. Anal. Khim.*, 49 (1994) 760-763.
- 852 Kawamura, K., Suzuki, I., Fujii, Y. and Watanabe, O.: Ice core record of polycyclic aromatic hydrocarbons over the past 400 years. *Naturwissenschaften*, 81 (1994) 502-505.
- 853 Li, X.: (Simultaneous determination of boiling range and purity of petroleum toluene by gas chromatography (GC).) *Chin. J. Chromatogr.*, 13, No. 1 (1995) 66-67.
- 854 Sim, P.G., Quilliam, M.A., Pleasance, S., Anacleto, J., Perreault, H., Ramaley, L. and Benoit, F.M.: Application of multidimensional chromatographic/mass spectrometric techniques to the analysis of complex mixtures of polycyclic aromatic compounds (PAC). *Polycyclic Aromat. Compd.*, 3 (1993) 321-328; *C.A.*, 121 (1994) 182985x.
- 855 Sturaro, A., Parvoli, G., Rella, R. and Doretto, L.: Gas chromatographic-mass spectrometric, high-performance liquid chromatographic-UV and gas chromatographic-Fourier transform IR responses to an industrial mixture of diisopropylnapthalenes. *J. Chromatogr. A*, 688 (1994) 211-219.
- 856 Visser, T., Vredenburg, M.J. and de Jong, A.P.J.M.: Confirmational analysis of polycyclic aromatic hydrocarbons in soil extracts by cryotrapping gas chromatography-Fourier transform infrared spectrometry. *J. Chromatogr. A*, 687 (1994) 303-313.
- 857 Xiu, Z., Xu, F., Zhang, X. and Zhang, Y.: (Direct determination of benzene in air by gas chromatography.) *Chin. J. Chromatogr.*, 12 (1994) 440 and 448.
- 858 Yanagisawa, K. and Kikuzaki, H.: (Analysis for benzene in gasolines by supercritical fluid chromatography.) *Sekiyu Gakkaishi*, 37, No. 6 (1994) 628-631; *C.A.*, 121 (1994) 283237k.
- 859 Yang, Y. and Baumann, W.: Seasonal and areal variations of polycyclic aromatic hydrocarbon concentration in street dust determined by supercritical fluid extraction and gas chromatography-mass spectrometry. *Analyst (Cambridge)*, 120 (1995) 243-248.

See also 766, 1153.

#### 5c. Halogen derivatives

- 860 Brown, J.F., Jr.: Determination of PCBs metabolic, excretion, and accumulation rates for use as indicators of biological response and relative risk. *Environ. Sci. Technol.*, 28 (1994) 2295-2305.
- 861 Bruno, T.J. and Caciari, M.: Retention of halocarbons on a hexafluoropropylene epoxidemodified graphitized carbon black. III. Ethene-based compounds. *J. Chromatogr. A*, 686 (1994) 245-251.
- 862 Burford, M.D., Hawthorne, S.B. and Miller, D.J.: Analysis of volatile organics by supercritical fluid extraction coupled to gas chromatography. I. Optimization of chromatographic parameters. *J. Chromatogr. A*, 685 (1994) 79-94.
- 863 Chen, C.-Y. and Ling, Y.-C.: Optimizing the gas chromatographic separation and detection of polychlorinated biphenyls by use of electronic pressure programming and experimental design. *J. High Resolut. Chromatogr.*, 17 (1994) 784-791.
- 864 De Boer, J., van der Meer, J., Reutergårdh, L. and Calder, J.A.: Determination of chlorobiphenyls in cleaned-up seal blubber and marine sediment extracts: Interlaboratory study. *J. Assoc. Off. Anal. Chem. Int.*, 77 (1994) 1411-1422.
- 865 Harrison, I., Leader, R.U., Higgo, J.J.W. and Tjell, J.C.: Determination of organic pollutants in small samples of groundwaters by liquid-liquid extraction and capillary gas chromatography. *J. Chromatogr. A*, 688 (1994) 181-188.
- 866 Helmig, D. and Greenberg, J.: Artifact formation from the use of potassium-iodide-based ozone traps during atmospheric sampling of trace organic gases. *J. High Resolut. Chromatogr.*, 18 (1995) 15-18.
- 867 Hillery, B.R., Girard, J.E., Schantz, M.M. and Wise, S.A.: Quantitative analysis of selected PCB congeners in marine matrix reference materials using a novel cyanobiphenyl stationary phase. *J. High Resolut. Chromatogr.*, 18 (1995) 89-96.
- 868 Hino, T., Nakanishi, S. and Hobo, T.: (Determination of volatile organic compounds in water by purge and trap capillary GC/MS without using cryofocus; utilizing advantage of characteristics of absorption and desorption on Tenax GC adsorbent.) *Bunseki Kagaku*, 1 (1995) 55-62.
- 869 Hirose, Y., Matsumoto, R., Yamada, S., Nozaka, T., Ishino, M. and Tanaka, A.: (The determination of volatile halo organic compounds in pharmaceuticals.) *Jpn. J. Toxicol. Environ. Health*, 40 (1994) 298-301; *C.A.*, 121 (1994) 187407v.
- 870 Jiang, C. and Wang, C.: (Gas chromatographic analysis of 1,3-dichloro-2-butene chlorinate.) *Hecheng Xiangjiao Gongye*, 17 (1994) 44-46; *C.A.*, 121 (1994) 257604h.
- 871 Johansen, H.R., Becher, G. and Greibrokk, T.: Determination of planar PCBs by combining on-line SFE-HPLC and GC-ECD or GC/MS. *Anal. Chem.*, 66 (1994) 4068-4073.
- 872 Krokos, F., Creaser, C.S., Wright, C. and Startin, J.R.: The development and validation of a congener specific method for the concurrent determination of *ortho*, non-*ortho* chlorobiphenyls, PCDDs and PCDFs. *Organohalogen Compd.*, 11 (1993) 61-65; *C.A.*, 121 (1994) 229150p.

- 873 Meng, Y., Zhao, M., Wang, Z., Shen, L. and Zhang, J.: (Qualitative and quantitative analyses of major components of PCBs and PCB<sub>5</sub> standart samples.) *Jilin Daxue Ziran Kexue Xuebao*, 3 (1994) 105-109; C.A., 121 (1994) 273948d.
- 874 Pietrogrande, M.C., Pasti, L., Dondi, F., Bollain Rodriguez, M.H. and Carro Diaz, M.A.: PCB separation by HRGC-MS. Fourier analysis for characterizing aroclor chromatograms. *J. High Resolut. Chromatogr.*, 17 (1994) 839-850.
- 875 Sauvain, J.-J., de Alencastro, L.F., Tarradellas, J., Camenzino, R., Karlaganis, G. and Vuilleumier, C.: Comparison of four quantification methods for the determination of PCB in transformer oil. *Fresenius J. Anal. Chem.*, 350 (1994) 555-562.
- 876 Takasuga, T., Inoue, T., Ohi, E.: Identification of chlordanes related compounds as potential interferences to PCDFs in the analysis of ambient air sample in Japan by HRGC/HRMS. *Organohalogen Compd.*, 11 (1993) 97-100; C.A., 121 (1994) 185847q.
- 877 Vetter, W., Luckas, B., Biermans, F., Mohnke, M. and Rotzsche, H.: Gas chromatographic separation of polychlorinated biphenyls on a new apolar capillary column. *J. High Resolut. Chromatogr.*, 17 (1994) 851-858.
- 878 Vollrath, A., Hohl, C. and Seiler, H.: Trace analysis of chlorofluorocarbons/partially halogenated chlorofluorohydrocarbons (CFC/HCFC) in polymeric foams by headspace capillary gas chromatography with electron-capture detection or ion trap detection combined with preconcentration on a cold trap. *Fresenius J. Anal. Chem.*, 351 (1995) 251-259.

See also 733, 741, 894, 898, 903, 1043, 1050.

5d. *Complex hydrocarbon mixtures (incl. analysis of tars, bitumens and mineral oils)*

- 879 Akard, M. and Sacks, R.D.: High-speed GC air monitor using cryointegration for sample collection. *J. Chromatogr. Sci.*, 32 (1994) 499-505.
- 880 Armstrong, D.W., Le, K., Reid, G.L. III., Lee, S.C., Beutelmann, K.K., Horak, M. and Tran, P.: Gas-solid chromatographic analysis of automobile tailpipe emissions as a function of different engine and exhaust system modifications. *J. Chromatogr. A*, 688 (1994) 201-209.
- 881 Burford, M.D., Hawthorne, S.B. and Miller, D.J.: Analysis of volatile organics by supercritical fluid extraction coupled to gas chromatography. II. Quantitation of petroleum hydrocarbons from environmental sample. *J. Chromatogr. A*, 685 (1994) 95-111.
- 882 Lynch, T.P. and Heyward, M.P.: Coupled packed SFC and capillary GC for the quantitative analysis of complex petroleum fractions. *J. Chromatogr. Sci.*, 32 (1994) 534-540.
- 883 Schubert, J.: Post-mortem test for low-boiling arson residues of gasoline by gas chromatography-ion-trap mass spectrometry. *J. Chromatogr. B*, 662 (1994) 113-117.
- 884 Sheff, L.M. and Siegel, J.A.: Fluorescence of petroleum products V: three-dimensional fluorescence spectroscopy and capillary gas chromatography of neat and evaporated gasoline samples. *J. Forensic Sci.*, 39 (1994) 1201-1214; C.A., 121 (1994) 273981j.

- 885 Van Delft, R.J., Doveren, A.S.M.J. and Snijders, A.G.: The determination of petroleum hydrocarbons in soil using a miniaturized extraction method and gas chromatography. *Fresenius J. Anal. Chem.*, 350 (1994) 638-641.

See also 736, 791, 1150, 1158.

6. ALCOHOLS

- 886 Lee, D.-S., Park, H.-S., Kim, K., Lee, T.-S. and Noh, B.-S.: (Gas chromatographic and mass spectrometric determination of alcohol homologs in Korean folk soju (distilled liquor).) *J. Korean Chem. Soc.*, 38 (1994) 640-652; C.A., 121 (1994) 279202c.
- 887 Sadeg, N. and Fourcault-Eddi, M.: (Blood alcohol determination by the EMIT ST method. Comparison with gas chromatography.) *Ann. Biol. Clin.*, 52 (1994) 463-465; C.A., 121 (1994) 273949e.
- 888 Smith, R.M. and Briggs, D.A.: Separation of homologous aromatic alcohols and carboxylic acids by packed column supercritical fluid chromatography. *J. Chromatogr. A*, 688 (1994) 261-271.

See also 727, 810, 914, 926, 930, 1003, 1162, 1179.

7. PHENOLS

- 889 Chalaux, N., Bayona, J.M. and Albaigés, J.: Determination of nonylphenols as pentafluorobenzyl derivatives by capillary gas chromatography with electron-capture and mass spectrometric detection in environmental matrices. *J. Chromatogr. A*, 686 (1994) 275-281.
- 890 Ge, X. and Chu, Q.: (Rapid determination of pentachlorophenol residues in leather and leather products by acetylation gas chromatography.) *Fenxi Huaxue*, 22 (1994) 81-83; C.A., 121 (1994) 233386n.
- 891 Krämer, A. and Angerer, J.: Comparison of two different derivatization procedures for the determination of urinary chlorophenol excretion. *Fresenius J. Anal. Chem.*, 351 (1995) 327-330.

See also 865, 888, 1073, 1161.

8. SUBSTANCES CONTAINING HETEROCYCLIC OXYGEN

8a. Flavonoids

- 892 Adlercreutz, H., Fotsis, T., Watanabe, S., Lampe, J., Wahala, K., Makela, T. and Hase, T.: Determination of lignans and isoflavonoids in plasma by isotope dilution gas chromatography-mass spectrometry. *Cancer Detect. Prev.*, 18 (1994) 259-271; C.A., 121 (1994) 276084e.

## 8b. Aflatoxins and other mycotoxins

- 893 Takino, M. and Yamaguchi, K.: (Analysis of trichothecene mycotoxins by gas chromatography/mass spectrometry with electric pressure control splitless injection.) *Shokuhin Eiseigaku Zasshi*, 35 (1994) 244-252; C.A., 121 (1994) 178000e.

## 8c. Other compounds with heterocyclic oxygen (incl. tannins)

- 894 Berlincioni, M., Groce, G., di Domenico, A., la Rocca, C., Lolini, M., Megli, A., Pupp, M. and Rizzi, L.: Reliability assessment of a method for detecting priority organic microcontaminants in urban air. *Organohalogen Compd.*, 11 (1993) 171-174; C.A., 121 (1994) 185849s.
- 895 Chittim, B.G., Bunce, N.J., Hu, K., Tashiro, C.H.M. and Yeo, B.R.: Comparison of GC-MS with an *in vitro* bioassay for PCDDs and related compounds in environmental samples. *Chemosphere*, 29 (1994) 1783-1788.
- 896 Chittim, B.G., Bunce, N.J., Hu, K., Tashiro, C.H.M. and Yeo, B.R.: Comparison of GC-MS with an *in vitro* bioassay for PCDDs and related compounds in environmental samples. *Organohalogen Compd.*, 11 (1993) 5-8; C.A., 121 (1994) 185842j.
- 897 Dettmer, K. and Stieglitz, L.: PARC an automated apparatus for clean-up procedures used in PCDD/F- and chlorobenzene-routine-analysis. *Organohalogen Compd.*, 11 (1993) 203-206; C.A., 121 (1994) 186306z.
- 898 Kimber, M.L., Wight, H., Pratt, J.A.E., Ireland, P. and Stern, G.: HRGC and HRMS analysis of PSDD/PCDF, PCBs and toxaphenes in environmental samples. *Organohalogen Compd.*, 11 (1993) 105-108; C.A., 121 (1994) 186603u.
- 899 Ritterbusch, J., Lorenz, W. and Bahadir, M.: Determination of polyhalogenated dibenzo-*p*-dioxins and dibenzofurans in analytical laboratory waste and their decomposition by UV-photolysis. *Chemosphere*, 29 (1994) 1829-1838.
- 900 Sivils, L.D., Kapila, S., Yan, Q. and Elseewi, A.A.: Application of two-dimensional chromatography system for gas-phase photodegradation studies of polychlorinated dibenzo-*p*-dioxins. *J. Chromatogr. A*, 688 (1994) 221-230.
- 901 Takasuga, T., Inoue, T. and Ohi, E.: Identification of chlordanes related compounds as potential interferences to PCDFs in the HRGC/HRMS analysis of ambient air samples in Japan. *Chemosphere*, 29 (1994) 1839-1847.
- 902 Tilio, R., Krishnan, K., Kapila, S., Nam, K.S. and Facchetti, S.: A simple analytical methodology for multiresidue pollutant determinations. *Chemosphere*, 29 (1994) 1849-1858.
- 903 Tuinstra, L.G.M.T., van Rhijn, J.A., Traag, W.A., v.d. Spreng, P., Zuidema, T. and Horstman, H.J.: Method for the determination of dioxins, planar and other PCB's in human milk. *Organohalogen Compd.*, 11 (1993) 181-183; C.A., 121 (1994) 186304x.
- 905 Cueto, R., Squadrito, G.L. and Pryor, W.A.: Quantifying aldehydes and distinguishing aldehydic product profiles from autooxidation and ozonation of unsaturated fatty acids. *Methods Enzymol.*, 233 (1994) 174-182; C.A., 121 (1994) 225357p.
- 906 Franke, S., Hildebrandt, S., Francke, W. and Reincke, H.: The occurrence of chlorinated bis-(propyl)ethers in the Elbe river and tributaries. *Naturwissenschaften*, 82 (1995) 80-83.
- 907 Gao, Y.: (GC analytical method for isopropyl ether in workplace air.) *Gongye Weisheng Yu Zhiyebing*, 19 (1993) 360-362; C.A., 121 (1994) 211671z.
- 908 Husain, S., Sarma, N.P., Rao, K.S.R. and Lakshmi, V.V.S.: Separation and determination of reaction mixtures of anthraquinone by gas chromatography. *J. Chromatogr. A*, 686 (1994) 356-359.
- 909 Kurz, J. and Ballschmiter, K.: Isomer-specific determination of 79 polychlorinated diphenylethers (PCDE) in cod liver oil, chlorophenols and in fly ash. *Fresenius J. Anal. Chem.*, 351 (1995) 98-109.
- 910 Ojala, M., Kotiaho, T., Siirila, J. and Sihvonen, M.-L.: Analysis of aldehydes and ketones from beer as O-(2,3,4,5,6-pentafluorobenzyl)hydroxylamine derivatives. *Talanta*, 41 (1994) 1297-1309.
- 911 Poskrobko, J., Linkiewicz, M. and Jaworski, M.: Analysis of high-boiling ethoxylates of methyl, ethyl, and butyl alcohols with the use of gas chromatography. *Chem. Anal. (Warsaw)*, 39 (1994) 153-159.
- 912 Slowick, J.J. and Uden, P.C.: Gas chromatography with atomic emission spectrometric detection for the determination of fluoroethers. *J. Anal. At. Spectrom.*, 9 (1994) 951-956; C.A., 121 (1994) 291686w.
- 913 Van Kuijk, F.J.G.M., Siakotos, A.N., Fong, J.G., Stephens, R.J. and Thomas, D.W.: Quantitative measurement of 4-hydroxy-alkenals in oxidized low-density lipoprotein by gas chromatography-mass spectrometry. *Anal. Biochem.*, 224 (1995) 420-424.
- 914 Viskrebentsev, V.P. and Spektorman, E.A.: (Gas-chromatographic determination of cyclohexanone and cyclohexanol at the level of maximum allowable concentrations in air.) *Dokl. Akad. Nauk Resp. Uzb.*, 3 (1993) 46-48; C.A., 121 (1994) 262174x.
- 915 Zeng, C. and Qiao, X.: (Analysis of product in methylal synthesis by GC.) *Tianranqi Huagong*, 19, No. 3 (1994) 50-52; C.A., 121 (1994) 270843e.
- 916 Zou, Q. and Su, J.: (Quantitative analysis of muscone in Shexiang Baixin pills by gas chromatography.) *Zhongguo Zhongyao Zazhi*, 19 (1994) 418-419; C.A., 121 (1994) 213148h.

See also 872, 876, 892.

See also 930, 1147, 1162.

## 9. OXO COMPOUNDS, ETHERS, EPOXIDES AND QUINONES

- 904 Bakola-Christianopoulou, M.N., Papageorgiou, V.P. and Apazidou, K.K.: Synthesis and gas chromatographic/mass spectrometric study of silylated hydroxyquinones on all hydroxyl groups. *Phosphorus, Sulfur Silicon Relat. Elem.*, 88, No. 1-4 (1994) 53-65; C.A., 121 (1994) 230837z.

## 10. CARBOHYDRATES

## 10a. Mono and oligosaccharides. Structural studies

- 917 Baker, J.R., Zyzak, D.V., Thorpe, S.R. and Baynes, J.W.: Chemistry of the fructosamine assay: D-glucosamine is the product of oxidation of amadori compounds. *Clin. Chem. (Washington)*, 40 (1994) 1950-1955.
- 918 Larsson, L., Pehrson, C., Wiebe, T. and Christensson, B.: Gas chromatographic determination of D-arabinitol/L-arabinitol ratios in urine: A potential method for diagnosis of disseminated candidiasis. *J. Clin. Microbiol.*, 32 (1994) 1855-1859; C.A., 121 (1994) 199934c.
- 919 Liu, S.Q. and Davis, C.R.: Analysis of wine carbohydrates using capillary gas liquid chromatography. *Am. J. Enol. Vitic.*, 45 (1994) 229-234; C.A., 121 (1994) 225794d.
- 920 Pitkanen, E. and Kanninen, T.: Determination of mannose and fructose in human plasma using deuterium labelling and gas chromatography/mass spectrometry. *Biol. Mass Spectrom.*, 23 (1994) 590-595; C.A., 121 (1994) 225794d.
- 921 Shetty, H.U., Holloway, H.W. and Rapoport, S.J.: Capillary gas chromatography combined with ion trap detection for quantitative profiling in cerebrospinal fluid and plasma. *Anal. Biochem.*, 224 (1995) 279-285.
- 922 Wang, Y., Mu, H. and Qiu, J.: (Gas chromatographic (GC) determination of sugars in soils.) *Chin. J. Chromatogr.*, 12 (1994) 438-439.
- 923 Yang, Y. and Guo, S.: (Analysis of ethyl sorbate and sorbic acid in disinfectants for livestock and fowl by gas chromatography.) *Huaxue Shijie*, 34 (1993) 549-550; C.A., 121 (1994) 263813s.

## 10b. Polysaccharides, mucopolysaccharides, lipopolysaccharides

- 924 Boutte, T.T. and Swanson, B.G.: Supercritical fluid extraction and chromatography of sucrose and methyl glucose polyester. *Food Sci. Technol. (N.Y.)*, 62 (1994) 65-93; C.A., 121 (1994) 250068c.
- 925 Pastorova, I., Botto, R.E., Arisz, P.W. and Boon, J.J.: Cellulose char structure: a combined analytical pyrolysis gas chromatography-mass spectrometry, FTIR, and NMR study. *Carbohydr. Res.*, 262 (1994) 27-47; C.A., 121 (1994) 182152y.

## 11. ORGANIC ACIDS AND LIPIDS

## 11a. Organic acids and simple esters

- 926 Alvarez, P. and Malcorps, P.: Analysis of free fatty acids, fusel alcohols, and esters in beer: an alternative to CS<sub>2</sub> extraction. *J. Am. Soc. Brew. Chem.*, 52 (1994) 127-134; C.A., 121 (1994) 178026t.
- 927 Frank, H., Renschen, D., Klein, A. and Scholl, H.: Trace analysis of airborne haloacetates. *J. High Resolut. Chromatogr.*, 18 (1995) 83-88.
- 928 Gradowska, W. and Larsson, L.: Determination of absolute configurations of 2- and 3-hydroxy fatty organic acids in organic dust by gas chromatography/mass spectrometry. *J. Microbiol. Methods*, 20 (1994) 55-67; C.A., 121 (1994) 197742q.

- 929 Guneral, F. and Bachmann, C.: Age-related reference values for urinary organic acids in a healthy Turkish pediatric population. *Clin. Chem. (Washington)*, 40 (1994) 862-868.
- 930 Jackson, P., Carter, D., Dent, G., Cook, B.W., Chalmers, J.M. and Dunkin, I.R.: Selective thermolysis of the enol forms of acetates during gas chromatography, revealed by combined matrix-isolation Fourier transform infrared and mass spectrometry. *J. Chromatogr. A*, 686 (1994) 287-293.
- 931 Kaunzinger, A., Podebrad, F., Liske, R., Maas, B., Dietrich, A. and Mosandl, A.: Stereochemical differentiation and simultaneous analysis of 3-, 4-, and 5-hydroxyalkanoic acids from biopolyesters by multidimensional gas chromatography. *J. High Resolut. Chromatogr.*, 18 (1995) 49-53.
- 932 Label, P., Imbault, N. and Villar, M.: ELIZA quantitation and GC-MS identification of abscisic acid in stigma, ovary and pedicel of pollinated poplar flowers (*Populus nigra* L.). *Tree Physiol.*, 14, No. 5 (1994) 521-530; C.A., 121 (1994) 174576z.
- 933 Lancas, F.M., Queiroz, M.E.C. and da Silva, I.C.E.: Seed oil extraction with supercritical carbon dioxide modified with pentane. *Chromatographia*, 39 (1994) 687-692.
- 934 Laplante, A., Comte, B. and Des Rosiers, C.: Assay of blood and tissue oxaloacetate and  $\alpha$ -ketoglutarate by isotopic dilution gas chromatography-mass spectrometry. *Anal. Biochem.*, 224 (1995) 580-587.
- 935 Lehmann, W.D., Metzger, K., Stephan, M., Wittig, U., Zalán, I., Habenicht, A.J.R. and Fűstenberger, G.: Quantitative lipooxygenase product profiling by gas chromatography negative-ion chemical ionization mass spectrometry. *Anal. Biochem.*, 224 (1995) 227-234.
- 936 Maya, M., Carvalhas, P. and Morais, E.J.: (Determination of valproic acid in blood plasma using gas-liquid chromatography.) *Rev. Pot. Farm.*, 44 (1994) 35-39; C.A., 121 (1994) 194784u.
- 937 Nishikawa, Y. and Okumura, T.: Determination of nitrilotriacetic acid and ethylenediaminetetraacetic acid in environmental samples as their methyl esters derivatives by gas chromatography-mass spectrometry. *J. Chromatogr. A*, 690 (1995) 109-118.
- 938 Park, M.K., Park, J.H., Kim, K.H., Han, S.B. and Han, B.H.: (Analysis of aromatic acids in Panax ginseng by gas chromatography.) *Yakhak Hoechi*, 38 (1994) 389-393; C.A., 121 (1994) 225407e.
- 939 Patel, A.S.: Fractionation of salad dressings for fatty acid profile and soluble sugar determination using the ultracentrifuge. *J. Assoc. Off. Anal. Chem. Int.*, 77 (1994) 1539-1542.
- 940 Sakaguchi, F. and Kawamura, K.: Identification of 4-oxoheptanedioic acid in the marine atmosphere by capillary gas chromatography-mass spectrometry. *J. Chromatogr. A*, 687 (1994) 315-321.
- 941 Schafer, K.: Analysis of short chain fatty acids from different intestinal samples. I: By packed column gas chromatography. *Chromatographia*, 39 (1994) 706-712.
- 942 Scotter, M.J., Thorpe, S.A., Reynolds, S.L., Wilson, L.A. and Strutt, P.R.: Rapid capillary gas chromatographic method for the determination of propionates in bread and bread products. *Food Addit. Contam.*, 11 (1994) 295-300; C.A., 121 (1994) 254001y.
- 943 Smid, I. and Salfinger, M.: Mycobacterial identification by computer-aided gas-liquid chromatography. *Diagn. Microbiol. Infect. Dis.*, 19, No. 2 (1994) 81-88; C.A., 121 (1994) 275713x.

- 944 Stoakes, L., John, M.A., Lannigan, R., Schieven, B.C., Ramos, M., Harley, D. and Hussain, Z.: Gas-liquid chromatography of cellular fatty acids for identification of staphylococci. *J. Clin. Microbiol.*, 32 (1994) 1908-1910; C.A., 121 (1994) 199935d.
- 945 Stoll, U.: (Techniques for phospholipid analyses applied to sera and egg yolk.) *Fat. Sci. Technol.*, 96 (1994) 188-194; C.A., 121 (1994) 174257w.
- 946 Tomkins, B.A.: Screening procedure for sodium fluoroacetate (compound 1080) at sub-microgram/gram concentration in soils. *Anal. Lett.*, 27 (1994) 2703-2718; C.A., 121 (1994) 279666g.
- 947 Vetter, W. and Walther, W.: Pyrrolidides as derivatives for the determination of the fatty acids of triacylglycerols by gas chromatography. *J. Chromatogr. A*, 686 (1994) 149-154.
- 948 Wolff, R.L.: Analysis of alpha-linoleic acid geometrical isomers in deodorized oils by capillary gas-liquid chromatography on cycloalkyl polysiloxane stationary phases: a note of caution. *J. Am. Oil Chem. Soc.*, 71 (1994) 907-909; C.A., 121 (1994) 178017r.
- 949 Zhou, X., Zhang, D. and Wang, T.: Study on the structure of fulvic acid from weathered coal using modified sodium dichromate oxidation and GC-MS. *Fuel. Sci. Technol. Int.*, 12 (1994) 1495-1508; C.A., 121 (1994) 283316k.

See also 888, 1028, 1067.

#### 11b. Prostaglandins

- 950 Ishibashi, M., Nakagawa, Y., Harima, N., Hishinuma, T. and Mizugaki, M.: Analysis of 11-dehydrothromboxane B<sub>2</sub> in human urine improved by use of gas chromatography/high-resolution selected ion monitoring with <sup>18</sup>O<sub>2</sub>-labeled analog as an internal standard. *Biol. Mass Spectrom.*, 23 (1994) 612-620; C.A., 121 (1994) 246526h.

#### 11c. Lipids and their constituents

- 951 Ebeler, S.E., Shibamoto, T. and Osawa, T.: Gas and high-performance liquid chromatographic analysis of lipid peroxidation products. *Chromatogr. Sci. Ser.*, 65 (1994) 223-249; C.A., 121 (1994) 249906t.
- 952 Jones, A.D.: Quantitative capillary gas chromatography-mass spectrometry of lipids using stable isotope dilution methods. *Chromatogr. Sci. Ser.*, 65 (1994) 347-364; C.A., 121 (1994) 249911r - a review with 38 refs.
- 953 Kuksis, A.: GLC and HPLC of neutral glycerolipids. *Chromatogr. Sci. Ser.*, 65 (1994) 177-222; C.A., 121 (1994) 249905s - a review with 123 refs.
- 954 Matsumoto, K. and Taguchi, M.: Supercritical fluid chromatographic analysis of lipids. *Chromatogr. Sci. Ser.*, 65 (1994) 365-396; C.A., 121 (1994) 249912s - a review with 49 refs.
- 955 Murphy, R.C.: GC/MS of PAF and related lipids. *J. Lipid Mediators Cell Signalling*, 10 (1994) 183-184; C.A., 121 (1994) 250135x.
- 956 Staby, A., Borch-Jensen, C., Balchen, S. and Møllerup, J.: Quantitative analysis of marine oils by capillary supercritical fluid chromatography. *Chromatographia*, 39 (1994) 697-705.
- 957 Trzicka, E. and Mares, P.: Gas-liquid chromatography of neutral lipids. *Chromatogr. Sci. Ser.*, 65 (1994) 103-176; C.A., 121 (1994) 249904r - a review with 202 refs.

See also 824, 947.

#### 11d. Lipoproteins and their constituents

- 958 Cummings, M.W. and Watts, G.V.: Increased hepatic secretion of very-low-density lipoprotein apolipoprotein B-100 in cholesteryl ester storage disease. *Clin. Chem. (Washington)*, 41 (1995) 111-114.
- 959 Das, D.K., Maulik, N., Jones, R.M. and Bagchi, D.: Gas chromatography-mass spectrometric detection of plasmogen phospholipids in mammalian heart. *Chromatogr. Sci. Ser.*, 65 (1994) 317-145; C.A., 121 (1994) 249190q - a review with 42 refs.

See also 913, 945.

#### 12. ORGANIC PEROXIDES

- 960 Bortolomeazzi, R., Pizzale, L., Vichi, S. and Lercker, G.: Analysis of isomers of cholesteryl acetate hydroperoxides HPLC and GC-ITDMS. *Chromatographia*, 39 (1994) 577-580.
- 961 Schmidt, R.W.H., Schurath, U., Slemr, F. and Seiler, W.: A fast capillary GC/ECD suitable for airborne peroxyacetyl nitrate (PAN) and peroxypropionyl nitrate (PPN) measurements. *Comm. Eur. Communities, [Rep. 15609] EUR*, 2 (1994) 724-730; C.A., 121 (1994) 262169z.

See also 1029.

#### 13. STEROIDS

##### 13a. General techniques

- 962 Hanson, M.: Aspects of retention behaviour of steroids in packed column supercritical fluid chromatography. *Chromatographia*, 40 (1995) 58-68.

##### 13b. Pregnane and androstane derivatives

- 963 Batjoens, P., de Brabander, H.F., Smets, F. and Pottle, G.: Gas chromatography-mass spectrometric confirmation of anabolic compounds in injection sites. *Analyst (Cambridge)*, 119 (1994) 2607-2610.
- 964 Homma, K., Masumoto, M., Takeshita, E., Iri, H., Anzo, M., Sato, S. and Matsuo, N.: (Assessment of the urinary steroid profiles in newborn infants by GC/MS selected ion monitoring system.) *Horumon to Rinsho*, 42 (1994) 623-629; C.A., 121 (1994) 221156m.
- 965 Hooijerink, D., Schilt, R., van Bennekom, E. and Brouwer, B.: Determination of anabolic esters in oily formulations and plasma in husbandry using high-performance liquid chromatography and gas chromatography-mass selective detection. *Analyst (Cambridge)*, 119 (1994) 2617-2622.

- 966 Petersen, H.W., Christensen, F. and Vad, M.: Method validation in relation to accuracy and precision for the gas chromatography/mass spectrometric analysis of 19-nortestosterone in urine with heptafluorobutyric anhydride and N-methyl-N-(trimethylsilyl)trifluoroacetamide. *Analyst (Cambridge)*, 119 (1994) 2627-2630.
- 967 Schoene, C., Nedderman, A.N.R. and Houghton, E.: Preliminary study of the metabolism of 17 $\alpha$ -methyltestosterone in horses utilizing gas chromatography-mass spectrometric techniques. *Analyst (Cambridge)*, 119 (1994) 2537-2542.
- 968 Thienpont, L.M., de Brabandere, V.I., Stöckl, D. and de Leenheer, A.P.: Use of cyclodextrins for prepurification of progesterone and testosterone from human serum prior to determination with isotope dilution gas chromatography/mass spectrometry. *Anal. Chem.*, 66 (1994) 4116-4119.
- See also 1124.
- 13c. *Estrogens*
- 969 Ye, L., Zhang, C.J., Zhang, Y.Z. and Liu, X.: (Trenbolone and its metabolites in human urine by GC/MS analysis.) *Yaoxue Xuebao*, 29 (1994) 61-67; C.A., 121 (1994) 221153h.
- 13d. *Sterols*
- 970 Das, D.K., Gangopadhyay, H. and Cordis, G.A.: Capillary gas chromatography of myocardial cholesterol oxides. *Chromatogr. Sci. Ser.*, 65 (1994) 75-101; C.A., 121 (1994) 249903q - a review with 81 refs.
- 971 Dionne, S., Tuchweber, B., Plaa, G.L. and Yousef, I.M.: Phase I and phase II metabolism of lithocholic acid in hepatic acinar zone 3 necrosis. Evaluation in rats by combined radiochromatography and gas-liquid chromatography/mass spectrometry. *Biochem. Pharmacol.*, 48 (1994) 1187-1197; C.A., 121 (1994) 227821j.
- 972 Dyer, R.G., Hetherington, C.S., Alberti, K.G.M.M., Laker, M.F.: Simultaneous measurement of phytosterols (campesterol and  $\beta$ -sitosterol) and 7-ketocholesterol in human lipoproteins by capillary column gas chromatography. *J. Chromatogr. B*, 663 (1995) 1-7.
- 973 Guardiola, F., Codony, R., Rafecas, M. and Boatella, J.: Selective gas chromatographic determination of cholesterol in eggs. *J. Am. Oil Chem. Soc.*, 71 (1994) 867-871.
- 974 Linnet, K.: An HPLC-GC/MS reference method for serum total cholesterol with control for ester hydrolysis. *Clin. Biochem.*, 27 (1994) 177-182; C.A., 121 (1994) 200315k.
- 975 Ollivier, D., Tisse, C., Dordonnat, M., Tisse, C. and Guerere, M.: (Differences in sterol composition and microstructure between fruits from Cucurbitaceae and roots from Cruciferae in the candy industry and products studied using gas chromatography and microscopy.) *Sci. Aliments*, 14 (1994) 189-202; C.A., 121 (1994) 178237n.
- 976 Paterson, E., Zuercher, U. and Amado, R.: (Determination of cholesterol in single foods and complete meals containing animal and plant components, by gas chromatography.) *Mitt. Geb. Lebensmittelunters. Hyg.*, 85 (1994) 132-145; C.A., 121 (1994) 253994n.
- 977 Richardson, R.K., Evers, J.M. and Gray, I.K.: Determination of cholesterol in dairy products by whole sample saponification. *Int. Dairy J.*, 4 (1994) 521-532; C.A., 121 (1994) 203640y.
- 978 Sevanian, A., Seraglia, R., Traldi, P., Rossato, P., Ursini, F. and Hodis, H.: Analysis of plasma cholesterol oxidation products using gas- and high-performance liquid chromatography/mass spectrometry. *Free Radical Biol. Med.*, 17 (1994) 397-409; C.A., 121 (1994) 250594w.
- See also 933, 956, 957, 960.
- 13e. *Bile acids and alcohols*
- 979 Iida, T., Tazawa, S., Tamaru, T., Goto, J. and Nambara, T.: Gas chromatographic separation of bile acid 3-glucosides and 3-glucuronides without prior deconjugation on a stainless-steel capillary column. *J. Chromatogr. A*, 689 (1995) 77-84.
- 13f. *Ecdysones and other insect steroid hormones*
- See 1041.
15. TERPENES AND OTHER VOLATILE AROMATIC COMPOUNDS
- 15a. *Terpenes*
- 980 Diehl, B.H., Combs, G.L., Bade, R.K. and Trimble, S.H.: New applications in process gas chromatography. *Proc Annu. ISA Anal. Div. Symp.*, 27 (1994) 2413-2419; C.A., 121 (1994) 262194d.
- 981 Kallio, H., Kerrola, K. and Alhonenmaki, P.: Carvone and limonene in caraway fruits (*Carum carvi* L.) analyzed by supercritical carbon dioxide extraction-gas chromatography. *J. Agric. Food Chem.*, 42 (1994) 2478-2485.
- 982 Ravid, U., Putievski, E. and Katzir, E.: Chiral GC analysis of (1R)-(+)-pulegone with high enantiomeric purity in essential oils of some Lamiaceae aromatic plants. *Flavour Fragrance J.*, 9 (1994) 205-207; C.A., 121 (1994) 286323w.
- 983 Wang, X., Jia, C. and Wan, H.: The direct chiral separation of some optically active compounds in essential oils by multidimensional gas chromatography. *J. Chromatogr. Sci.*, 33 (1995) 22-25.
- See also 797, 1152.
- 15b. *Essential oils*
- 984 Barth, D., Chouchi, D., Della Porta, G. and Reverchon, E.: Desorption of lemon peel oil by supercritical carbon dioxide: Deterpenation and psoralenes elimination. *J. Supercrit. Fluids*, 7 (1994) 177-183.
- 985 Buchbauer, G., Jirovetz, L., Wasicky, M. and Kikiforov, A.: Comparative investigation of Douglas fir headspace samples, essential oils, and extracts (needles and twigs) using GC-FID and GC-FTIR-MS. *J. Agric. Food Chem.*, 42 (1994) 2852-2854.
- 986 Hartmann, G. and Koestner, S.: (Quality of ginger and ginger extracts. Effects of processing on quality with respect to the contents of gingerols.) *Fleischwirtschaft*, 74 (1994) 313-315; C.A., 121 (1994) 254189r.

- 987 Reverchon, E., Osséo, L.S. and Gorgoglione, D.: Supercritical CO<sub>2</sub> extraction of basil oil: Characterization of products and process modeling. *J. Supercrit. Fluids*, 7 (1994) 185-190.
- 988 Stashenko, E., Wiame, H., Dassy, S., Martinez, J.R. and Shibamoto, T.: Catalytic transformation of Copaiba (*Copaifera officinalis*) oil over zeolite ZSM-5. *J. High Resolut. Chromatogr.*, 18 (1995) 54-58.
- 989 Stashenko, E.E., Torres, W., Morales, J.R.M.: A study of the compositional variation of the essential oil of Ylang-ylang (*Cananga odorata* Hook Fil. et Thomson, *forma genuina*) during flower development. *J. High Resolut. Chromatogr.*, 18 (1995) 101-104.
- 990 Vilegas, J.H.Y. and Lanças, F.M.: HRGC-MS identification of low-polarity compounds from Gomortega keule. *J. High Resolut. Chromatogr.*, 18 (1995) 129-130.
- 991 Wang, X., Li, J. and Wei, L.: (Chemical constituents of *Sophora flavescens* volatile oils.) *Zhongguo Zhongyao Zazhi*, 19 (1994) 552-553; C.A., 121 (1994) 263802n.
- 992 Zekovic, Z., Pekic, B., Lepojevic, Z. and Petrovic, L.: Chromatography in our investigation of camomile (*Matricaria chamomilla* L.). *Chromatographia*, 39 (1994) 587-590.
- See also 808, 982, 1143.
16. NITRO AND NITROSO COMPOUNDS
- 993 Feltes, J. and Koll, J.: GC and HPLC with selective detection for the determination of explosives in water and soil. *LC-GC Int.*, 7 (1994) 698-701.
- 994 Francis, E.S.: Supercritical fluid chromatography and extraction: Instrumental techniques for the analysis of nitro and nitroso compounds. Avail. *Univ. Microfilms Int.*, Order No. DA9418451, 1994, 142 pp.; C.A., 121 (1994) 270806v.
- 995 Lauro-Marty, C., Lartigue-Mattei, C., Chabard, J.L., Beyssac, E., Aiache, J.M. and Madesclaire, M.: Gas chromatographic-mass spectrometric determination of isosorbide 5-mononitrate in human plasma. *J. Chromatogr. B*, 663 (1995) 153-159.
- 996 Linkerhägner, M., Stan, H.-J. and Rimkus, G.: Detection of nitro musks in human fat by capillary gas chromatography with atomic emission detection (AED) using programmed temperature vaporization (PTV). *J. High Resolut. Chromatogr.*, 17 (1994) 821-826.
- 997 Plomey, J.B., Koester, C.J. and March, R.E.: Determination of N-nitrosodimethylamine in complex environmental matrices by quadrupole ion storage tandem mass spectrometry enhanced by unidirectional ion ejection. *Anal. Chem.*, 66 (1994) 4437-4443.
- 998 Wang, D., He, Y., Cui, Q. and Niu, M.: (Gas chromatographic analysis of the nitration products of o-dichlorobenzene.) *Chin. J. Chromatogr.*, 13, No. 1 (1995) 68-69.
- 999 Wisneski, H.H., Yates, R.L. and Havery, D.C.: Determination of musk ambrett in fragrance products by capillary gas chromatography with electron capture detection: Interlaboratory study. *J. Assoc. Off. Anal. Chem. Int.*, 77 (1994) 1467-1471.
- 1000 Xia, G.Z., Fu, Y.K., Dong, S.N. and Li, R.: (Gas chromatographic method with EC detection to determine isosorbide-5-mononitrate (IS-5-MN) in human serum and its pharmacokinetic parameters.) *Yaoxue Xuebao*, 29 (1994) 634-638; C.A., 121 (1994) 271124h.
- See also 865, 961, 1001, 1153, 1169.
17. AMINES, AMIDES AND RELATED NITROGEN COMPOUNDS
- 17a. Amines and polyamines
- 1001 Hou, D. and Tang, J.: (Determination of acetonitrile, acrylonitrile, aniline and nitrobenzene in water and discharged industrial water by direct aqueous injection gas chromatography.) *Huanjing Kexue*, 15, No. 2 (1994) 69-70,74; C.A., 121 (1994) 237948v.
- 1002 Miyamoto, Y., Kataoka, H., Ohnui, S. and Makita, M.: (Determination of primary amines as their N-benzenesulfonyl-N-trifluoroacetyl derivatives by GC with electron capture detection.) *Bunseki Kagaku*, 12 (1994) 1113-1117.
- 1003 Zhukov, A.V. and Vereshchagin, A.G.: Nonenzymic isolation of ethanolamine from the phospholipids of soybean seeds and its identification. *Fiziol. Rast. (Moscow)*, 41, No. 3 (1994) 420-424; C.A., 121 (1994) 174359f.
- See also 1153.
- 17d. Other amine derivatives and amides (excl. peptides)
- 1004 Brunmark, P., Dalene, M. and Skarping, G.: Gas chromatography-negative-ion chemical ionization mass spectrometry of hydrolysed human urine and blood plasma for the biomonitoring of occupational exposure to 4,4'-methylenedianiline. *Analyst (Cambridge)*, 120 (1995) 41-45.
- 1005 Kameyama, H., Shimada, R., Nakagome, S., Watanabe, M. and Takeyama, T.: (Gas chromatographic determination of aliphatic amines, and evaluation of offensive odor in fish meal plant by odor organoleptic test and semiconductor sensor.) *Bunseki Kagaku*, 12 (1994) 1071-1076.
- 1006 Kolb, M. and Badahir, M.: Formation of Schiff bases with acetone as a solvent in the determination of anilines. *J. Chromatogr. A*, 685 (1994) 189-194.
- 1007 Muro, H., Tatsuhara, T., Sugimoto, T., Woo, M., Nishida, N., Murakami, K. and Yamaguchi, Y.: Determination of urinary valproylcarnitine by gas chromatography-mass spectrometry with selected-ion monitoring. *J. Chromatogr. B*, 663 (1995) 83-89.
- 1008 Promyshlennikova, E.P., Kirichenko, V.E., Paskievich, K.I., Golgor'eva, D.N. and Golovnya, R.V.: (Gas-chromatographic retention of polyfluorinated carboxylic acid amines.) *Zh. Anal. Khim.*, 49 (1994) 729-734.
- 1009 Taguchi, V.Y., Jenkins, S.W.D., Wang, D.T., Palmentier, J.-P.F.P. and Reiner, E.J.: Determination of N-nitrosodimethylamine by isotope dilution high-resolution mass spectrometry. *Can. J. Appl. Spectrosc.*, 39 (1994) 87-93; C.A., 121 (1994) 244538q.
- See also 1028, 1041.

# 18. AMINO ACIDS AND PEPTIDES; CHEMICAL STRUCTURE OF PROTEINS

## 18a. Amino acids and their derivatives

- 1010 Howell, S.E., Miller, S.R., McCallister, J.D., Cherkofsky, S.C. and Patrick, K.S.: Gas chromatographic-mass spectrometric determination of urinary 1-aminocyclopropanecarboxylic acid in mice using a deuterated internal standard. *J. Chromatogr. B*, 663 (1995) 148-152.
- 1011 Palaty, J., Burton, R. and Abbott, F.S.: Rapid assay for  $\gamma$ -aminobutyric acid in mouse brain synaptosomes using gas chromatography-mass spectrometry. *J. Chromatogr. B*, 662 (1994) 85-90.

See also 1131.

# 21. PURINES, PYRIMIDINES, NUCLEIC ACIDS AND THEIR CONSTITUENTS

## 21a. Purines, pyrimidines, nucleosides, nucleotides

- 1012 Blount, B.C. and Ames, B.N.: Development of a sensitive assay for detection of uracil in DNA. *Adv. Exp. Med. Biol.*, 338 (1993) 741-744; C.A., 121 (1994) 225380r.

See also 1013.

## 21c. Nucleic acids, DNA

- 1013 Shuker, D.E.G. and Bartsch, H.: Detection of human exposure to carcinogens by measurement of alkyl-DNA adducts using immunoaffinity clean-up in combination with gas chromatography-mass spectrometry and other methods of quantification. *Mutat. Res.*, 313 (1994) 263-268; C.A., 121 (1994) 248135d - a review with 21 refs.

# 22. ALKALOIDS

- 1014 Bekkouche, K., Lazrek, H.B., and Jana, M.: (Quantitative determination of atropine and scopolamine in Moroccan Mandragora.) *Plant. Med. Phytoter.*, 26 (1993) 309-318; C.A., 121 (1994) 238503h.
- 1015 Budzinski, H., Garrigues, P., Connan, J. and Bellocq, J.: Identification of alkylated dibenzothiophenes: Application to organic geochemistry. *Polycyclic Aromat. Compd.*, 3 (1993) 611-620; C.A., 121 (1994) 182949p.
- 1016 Hattori, H., Kurono, S., Kimura, M., Ishii, A., Suzuki, O. and Yamada, T.: Determination of cocaine in body fluids by gas chromatography with surface ionization detection: comparison with nitrogen-phosphorus detection. *Hochudoku*, 12 (1994) 26-32; C.A., 121 (1994) 223732h.
- 1017 Meissner, C., Krauss, D. and Wink, M.: (Identification and quantification of alkaloids of lupines by capillary gas chromatography and coupled mass spectrometry (GC and GC/MS).) *CLB Chem. Labor Biotech.*, 44 (1993) 331-336; C.A., 121 (1994) 225229y - a review with 13 refs.

- 1018 Wu, Y., Cui, K., Wang, S., Xu, Y. and Shen, L.: (Determination of codeine and its metabolite in urine by GC-MS.) *Fenxi Ceshi Xuebao*, 13, No. 3 (1994) 71-74; C.A., 121 (1994) 271132j.

See also 1127.

# 23. OTHER SUBSTANCES CONTAINING HETEROCYCLIC NITROGEN

## 23a. Porphyrins and other pyrroles

- 1019 Ebdon, L., Evans, E.H., Pretorius, W.G. and Rowland, S.J.: Analysis of geoporphyryns by high-temperature gas chromatography inductively coupled plasma mass spectrometry and high-performance liquid chromatography inductively coupled plasma mass spectrometry. *J. Anal. At. Spectrom.*, 9 (1994) 939-943; C.A., 121 (1994) 234245c.

## 23d. Pyridine derivatives

- 1020 Zahlsen, K. and Nilsen, O.G.: Nicotine in hair of smokers and non-smokers: sampling procedure and gas chromatographic/mass spectrometric analysis. *Pharmacol. Toxicol. (Copenhagen)*, 75 (1994) 143-149; C.A., 121 (1994) 198365f.
- 1021 Zhang, M. and Phillips, J.B.: Applications of multiplex gas chromatography to the determination of organics in solid samples. *J. Chromatogr. A*, 689 (1995) 275-284.
- 1022 Zhou, J. and Ogiso, T.: (Determination of betahistidine in plasma with GC-alkaline flame ionization detector.) *Zhongguo Yaoke Daxue Xuebao*, 25 (1994) 224-226; C.A., 121 (1994) 271126k.

See also 1096.

## 23e. Other N-heterocyclic compounds

- 1023 Galceran, M.T., Curto, M.J., Puignou, L. and Moyano, E.: Determination of acridine derived compounds in charcoal-grilled meat and creosote oils by liquid chromatographic and gas chromatographic analysis. *Anal. Chim. Acta*, 295 (1994) 307-313.
- 1024 Haber, H., Haber, H.M. and Melzig, M.F.: A new rapid method for the analysis of catecholic tetraisoquinolines from biological samples by gas chromatography/mass spectrometry. *Anal. Biochem.*, 224 (1995) 256-262.
- 1025 Jouret, C., Pace, E. and Parfait, A.: (Alkylpyrazines in the analytical differentiation of rums obtained from sugarcane juice and molasses.) *Ann. Falsif. Expert. Chim. Toxicol.*, 87, No. 926 (1994) 85-90; C.A., 121 (1994) 203646e.
- 1026 Vander Meer, R.K. and Morel, L.: Ant queens deposit pheromones and antimicrobial agent on eggs. *Naturwissenschaften*, 82 (1995) 93-95.

See also 1079.

## 24. ORGANIC SULPHUR COMPOUNDS (INCL. GLUCOSINOLATES)

- 1027 Chen, Y.-C., Her, J.-S. and Lo, J.-G.: (Determination of reduced sulfur compounds in the industrial environment.) *Huaxue*, 52 (1994) 115-122; C.A., 121 (1994) 237219h.
- 1028 Kopka, J., Ohlrogge, J.B. and Jaworski, J.G.: Analysis *in vivo* levels of acyl-thioesters with gas chromatography-mass spectrometry of the butylamide derivative. *Anal. Biochem.*, 224 (1995) 51-66.

See also 752, 1180.

## 25. ORGANIC PHOSPHORUS COMPOUNDS (INCL. SUGAR PHOSPHATES)

- 1029 Perkel, A.L., Krutskaya, L.V. and Freidin, B.G.: (Application of triphenylphosphine to the gas chromatographic determination of peroxides in the oxidation products of organic compounds.) *Zh. Anal. Khim.*, 49 (1994) 768-772.

## 26. ORGANOMETALLIC AND RELATED COMPOUNDS

## 26a. Organometallic compounds

- 1030 Alli, A., Jaffé, R. and Jones, R.: Analysis of organomercury compounds in sediments by capillary GC with atomic fluorescence detection. *J. High Resolut. Chromatogr.*, 17 (1994) 745-748.
- 1031 Blake, E., Raynor, M.W. and Cornell, D.: On-line capillary supercritical fluid chromatography-inductively coupled plasma mass spectrometry for the analysis of organometallic compounds. *J. High Resolut. Chromatogr.*, 18 (1995) 33-37.
- 1032 Cai, X.-J., Uden, P.C., Sullivan, J.J., Quimby, B.D. and Block, E.: Head-space gas chromatography with atomic emission and mass selective detection for the determination of organoselenium compounds in elephant garlic. *Anal. Proc.*, 31 (1994) 325-327.
- 1033 Ceulemans, M., Witte, C., Lobinski, R. and Adams, F.C.: Simplified sample preparation for GC speciation analysis of organotin in marine biomaterials. *Appl. Organomet. Chem.*, 8 (1994) 451-461; C.A., 121 (1994) 248183t.
- 1034 Dafflon, O., Gobet, H. and Koch, H.: (Quantification of methyl mercury in fish by gas chromatography.) *Mitt. Geb. Lebensmitelunters. Hyg.*, 84 (1993) 557-565; C.A., 121 (1994) 253985k.
- 1035 Sarradin, D.M., Leguille, F., Astrue, A., Pinel, R. and Astrue, M.: Optimization of atomization parameters in the separation of organotin compounds by hydride generation-gas chromatography-electrothermal atomic absorption spectrometry. *Analyst (Cambridge)*, 120 (1995) 79-83.

## 26b. Boranes, silanes and related non-metallic compounds

- 1036 Cai, X.-J., Uden, P.C., Block, E., Zhang, X., Quimby, B.D. and Sullivan, J.J.: Allium chemistry: Identification of natural abundance organoselenium volatiles from garlic, elephant garlic, onion, and Chinese chive using headspace gas chromatography with atomic emission detection. *J. Am. Oil Chem. Soc.*, 71 (1994) 2081-2084.
- 1037 Delgado-Morales, W., Zingaro, R.A. and Mohan, M.S.: Arsenic in natural gas: Analysis and characterization of pipeline solids by <sup>1</sup>H NMR and other methods. *Intern. J. Environ. Anal. Chem.*, 57 (1994) 313-328.
- 1038 Ishizuka, C., Omae, K., Tanaka, S. and Nakashima, H.: Determination of tetraethoxysilane (TEOS) in air. *Sangyo Igaku*, 36 (1994) 201-202; C.A., 121 (1994) 185856s.
- 1039 Wenclawiak, B.W. and Krah, M.: Reactive supercritical fluid extraction and chromatography of arsenic species. *Fresenius J. Anal. Chem.*, 351 (1995) 134-138.

## 26c. Coordination compounds

See 1170.

## 27. VITAMINS AND VARIOUS ANIMAL GROWTH FACTORS (NON-PEPTIDIC)

- 1040 Hui, B., Young, A.J., Booth, L.A., Briton, G., Evershed, R.P. and Bilton, R.F.: Detection of carotenoids on supercritical fluid chromatography (SFC). A preliminary investigation on the spectral shifts of carotenoids in supercritical carbon dioxide. *Chromatographia*, 39 (1994) 549-556.
- 1041 Sundaram, K.M.S., Nott, R. and Lewin, E.E.: Comparative study of the determination of tebufenozide in formulated products by gas chromatographic and liquid chromatographic methods. *J. Chromatogr. A*, 687 (1994) 323-332.
- 1042 Tomiyama, S., Nitta, T. and Yamada, S.: Gas chromatography/mass spectrometric analysis of 24R, 25-dihydroxyvitamin D<sub>3</sub> using 24R, 25-dihydroxy[6,19,19-<sup>2</sup>H]vitamin D<sub>3</sub> as internal standard. *Steroids*, 59 (1994) 559-563.

## 29. INSECTICIDES, PESTICIDES AND OTHER AGROCHEMICALS

## 29a. General techniques

- 1043 Hardt, I.H., Wolf, C., Gehrcke, B., Hochmuth, D.H., Pfaffenberger, B., Hühnerfuss, H. and König, W.A.: Gas chromatographic enantiomer separation of agrochemicals and polychlorinated biphenyls (PCBs) using modified cyclodextrins. *J. High Resolut. Chromatogr.*, 17 (1994) 859-864.
- 1044 Lin, R. and Li, N.: (Analysis of multi-pesticides formulations with flame ionization detector and hydrogen carrier gas by gas chromatography (GC).) *Chin. J. Chromatogr.*, 12 (1994) 419-420.

- 1045 Miyata, M., Kamakura, K., Narita, M., Hirahara, Y., Okamoto, K., Hasegawa, M., Koiguchi, S., Miyoshi, T., Yamana, T. and Tonogai, Y.: (Studies on systematic determination of organophosphorus, carbamate, organochlorine, and pyrethroid pesticides in polished rice by gas chromatography.) *Shokuhin Eiseigaku Zasshi*, 35 (1994) 276-286; C.A., 121 (1994) 178001f.
- 1046 Nakamura, Y., Tonogai, Y., Sekiguchi, Y., Tsumura, Y., Nishida, N., Takakura, K., Isechi, M., Yuasa, K. and Nakamura, M.: Multiresidue analysis of 48 pesticides in agricultural products by capillary gas chromatography. *J. Agric. Food Chem.*, 42 (1994) 2508-2518.
- 1047 Sippola, E., David, F. and Sandra, P.: Comparison of polydimethyl siloxane stationary phases with 5% and 50% phenyl substitution for the separation of organophosphorus and organonitrogen pesticides under optimized gas chromatographic conditions. *J. High Resolut. Chromatogr.*, 18 (1995) 111-114.
- 1048 Suzuki, S.: (Determination of airborne acephate and trichlorfon by GC/chemical ionization MS.) *Bunseki Kagaku*, 1 (1995) 41-48.
- See also 822, 1156.
- 29b. Chlorinated insecticides**
- 1049 Barnabas, I.J., Dean, J.R. and Hitchen, S.M.: Selective supercritical fluid extraction of organochlorine pesticides and herbicides from aqueous samples. *J. Chromatogr. Sci.*, 32 (1994) 547-551.
- 1050 Cao, X. and Yao, W.: (Separation of polychlorinated biphenyls and chlorinated pesticides by normal phase high performance liquid chromatography (HPLC)-gas chromatography (GC).) *Chin. J. Chromatogr.*, 12 (1994) 401-403.
- 1051 Fernandez-Alba, A.R., Valverde, A., Agüera, A. and Contreras, M.: Gas chromatographic determination of organochlorine and pyrethroid pesticides of horticultural concern. *J. Chromatogr. A*, 686 (1994) 263-274.
- 1052 Goosens, E.C., de Jong, D., de Jong, G.J., Rinkema, F.D. and Brinkman, U.A.T.: Continuous liquid-liquid extraction combined on-line with capillary gas chromatography-atomic emission detection for environmental analysis. *J. High Resolut. Chromatogr.*, 18 (1995) 38-44.
- 1053 Hainzel, D., Burhenne, J., Barlas, H. and Parlar, H.: Spectroscopic characterization of environmentally relevant C<sub>10</sub>-chloroterpenes from a photochemically modified toxaphene standard. *Fresenius J. Anal. Chem.*, 351 (1995) 271-285.
- 1054 Muir, D.G.C. and de Boer, J.: Recent developments in the analysis and environmental chemistry of toxaphene with emphasis on the marine environment. *TrAC*, 14 (1995) 56-66 - a review with 79 refs.
- 1055 Popp, P., Kalbitz, K. and Oppermann, G.: Application of solid-phase microextraction and gas chromatography with electron-capture and mass spectrometric detection for the determination of hexachlorocyclohexanes in soil solutions. *J. Chromatogr. A*, 687 (1994) 133-140.
- 1056 Sato, Y., Sugawara, M., Watabe, R., Kikuchi, M., Mishima, Y., Kato, T. and Oba, M.: (Simultaneous determination of organochlorine and pyrethroid pesticides by capillary gas chromatography with an ECD detector. Clean-up by solid phase extraction method for agricultural products.) *Shokuhin Eiseigaku Zasshi*, 35 (1994) 390-396; C.A., 121 (1994) 203655g.
- 1057 Zhu, Y., Lu, Y. and Xu, J.: (Analysis of oxyfluorfen residue by gas chromatographic method.) *Shanghai Huanjing Kexue*, 12, No. 11 (1993) 21-22; C.A., 121 (1994) 198103u.
- See also 877, 898, 901, 902.
- 29c. Phosphorus insecticides**
- 1058 Cooper, J.-F., Zheng, S.-Q., Wynn, N.R. and Palcy, L.: Determination of organophosphorus nematocidal residues in banana-field soils by gas chromatography/mass spectrometry. *J. Assoc. Off. Anal. Chem. Int.*, 77 (1994) 1580-1586.
- 1059 Erney, D.R.: Determination of organophosphorus pesticides in whole/chocolate/skim-milk and infant formula using solid-phase extraction with capillary gas chromatography/flame photometric detection. *J. High Resolut. Chromatogr.*, 18 (1995) 59-62.
- 1060 Grob, K., Biedermann, M. and Giuffrè, A.M.: Determination of organophosphorus insecticides in edible oils and fats by splitless injection of the oil into a gas chromatograph (injector-internal headspace analysis). *Z. Lebensm.-Unters. Forsch.*, 198, No. 4 (1994) 325-328; C.A., 121 (1994) 178054a.
- 1061 Suzuki, T., Yaguchi, K., Ohnishi, K. and Yamagishi, T.: Gas chromatographic detection of tris(2-chloroethyl) and tris(2-butoxyethyl)phosphate in groundwater by large-sample-volume injection. *J. Assoc. Anal. Chem. Int.*, 77 (1994) 1647-1651.
- 1062 Yang, D., Zhang, Y. and Fang, C.: (Capillary gas chromatography method for determination of organophosphorus and carbamate pesticide residues in rice.) *Weisheng Yanjiu*, 23 (1994) 97-100; C.A., 121 (1994) 254008f.
- See also 1047.
- 29d. Carbamates**
- 1063 Bakowski, R.S., Ali, M.S., White, J.D., Phillippo, T. and Ellis, R.L.: Gas chromatographic/mass spectrometric confirmation of ten N-methylcarbamate insecticides in liver. *J. Assoc. Off. Anal. Chem. Int.*, 77 (1994) 1568-1574.
- 1064 Canas, B.J., Joe, F.L., Diachenko, G.W. and Burns, G.: Determination of ethyl carbamate in alcoholic beverages and soy sauce by gas chromatography with mass selective detection: Collaborative study. *J. Assoc. Off. Anal. Chem. Int.*, 77 (1994) 1530-1536.
- 1065 Evans, M.B. and Smith, M.S.: A comparative study of the chromatography of hydroxylated dialkylthiocarbamates as models for drug metabolites. *Chromatographia*, 39 (1994) 569-576.
- See also 1062.

29e. *Herbicides*

- 1066 Del Valle, P.L. and Nelson, J.O.: Evaluation of atrazine soil extraction methods for the determination by enzyme immunoassay and gas chromatography. *Arch. Environ. Contam. Toxicol.*, 27 (1994) 375-383; C.A., 121 (1994) 172938v.
- 1067 Heberer, T., Butz, S. and Stan, H.-J.: Detection of 30 acidic herbicides and related compounds as their pentafluorobenzylic derivatives using gas chromatography/mass spectrometry. *J. Assoc. Off. Anal. Chem. Int.*, 77 (1994) 1587-1604.
- 1068 Lou, J. and Zhao, D.: (Determination of chlorothalonil residue on vegetables by gas chromatography.) *Huanjing Wuran Yu Fangzhi*, 15, No. 5 (1993) 38-39; C.A., 121 (1994) 229148u.
- 1069 Sanchez-Brunette, C., Martinez, L. and Tadeo, J.L.: Determination of corn herbicides by GC-MS and GC-NPD in environmental samples. *J. Agric. Food Chem.*, 42 (1994) 2210-2214.
- 1070 Ting, K.-C. and Lee, C.-S.: Gas chromatographic determination of triclopyr in fruits and vegetables. *J. Chromatogr. A*, 690 (1995) 119-129.
- 1071 Zhang, H. and Luo, Y.: (Use of X-5 resin miniature cartridge in concentration of six trace pyrethrins insecticides in biological samples.) *Chin. J. Chromatogr.*, 13, No. 1 (1995) 40-42.

See also 749, 832.

29f. *Fungicides*

- 1072 Chichila, T.M.P. and Erney, D.R.: Fullscan confirmation of tetrahydrophtalimide in whole milk using gas chromatography/ion trap mass spectrometry. *J. Assoc. Off. Anal. Chem. Int.*, 77 (1994) 1574-1580.
- 1073 Goldberg, D.M., Yan, J., Ng, E., Diamandis, E.P., Karumanchiri, A., Soleas, G. and Waterhouse, A.L.: Direct injection gas chromatographic mass spectrometric assay for trans-resveratrol. *Anal. Chem.*, 66 (1994) 3959-3963.

See also 890.

29g. *Other types of pesticides and various agrochemicals*

- 1074 Li, C.: Determination of methomyl 20 EC by gas chromatography. *Nongyao Kexue Yu Guanli*, 400, No. 3 (1994) 24-26; C.A., 121 (1994) 248551m.
- 1075 Nishizawa, H., Shigemura, M., Suzuki, T. and Matazaemon, U.: Simple cleanup procedure for analysis of buprofezin residues and its metabolite in crops by gas chromatography. *J. Assoc. Off. Anal. Chem. Int.*, 77 (1994) 1631-1633.
- 1076 Pang, G.-F., Zhao, T.-S., Chao, Y.-Z. and Fan, C.-L.: Cleanup with two florisil columns for gas chromatographic determination of multiple pyrethroid insecticides in products of animal origin. *J. Assoc. Anal. Chem. Int.*, 77 (1994) 1634-1638.
- 1077 Runguo, L.: (Determination of pyrethroids with same column gas chromatography.) *Huaxue Shijie*, 34 (1993) 491-494; C.A., 121 (1994) 172930m.
- 1078 Terreni, M., Benfenati, E., Natangelo, M., Facchini, G. and Pagani, G.: Synthesis and use of pentadeuteroethyl ethofumesate as an internal standard for the determination of ethofumesate and its metabolites in water by gas chromatography-mass spectrometry. *J. Chromatogr. A*, 688 (1994) 243-250.

- 1079 Wu, Y., Ping, X. and Liang, T.: (The residual determination of fenpyroximate and its *cis*-isomer by gas chromatography.) *Chin. J. Chromatogr.*, 12 (1994) 421-422.

See also 1051.

31. *PLASTICS AND THEIR INTERMEDIATES*

- 1080 Coskun, M., Ozdemir, E., Celik, S. and Cansiz, A.: Inverse gas chromatography study of *n*-alcohols-poly(acrylonitrile-co-methyl acrylate) systems. *Macromol. Rep.*, A31 (1994) 63-72; C.A., 121 (1994) 256756x.
- 1081 Jackson, P.L., Huglin, M.B. and Cervenka, A.: Use of the inverse gas chromatography to quantify interactions in anhydride cured epoxy resins. *Polym. Int.*, 35 (1994) 135-143; C.A., 121 (1994) 281573t.
- 1082 Judge, M.D.: Investigation of the thermal oxidation of a polybutadiene by headspace sampling-gas chromatography. *Talanta*, 41 (1994) 755-759; C.A., 121 (1994) 180840x.
- 1083 Karlsson, S., Hakkarainen, M. and Albertsson, A.-C.: Identification by headspace gas chromatography-mass spectrometry of *in vitro* degradation products of homo- and copolymers of L- and D,L-lactide and 1,5-dioxepan-2-one. *J. Chromatogr. A*, 688 (1994) 251-259.
- 1084 Menescal, R., Eveland, J., West, R. and Blaszo, M.: Poly(di-*n*-alkylsilylene) copolymers: investigation of the microstructure by <sup>1</sup>H and <sup>29</sup>Si solution NMR and pyrolysis-gas chromatography/mass spectrometry. *Macromolecules*, 27 (1994) 5893-5899; C.A., 121 (1994) 180356u.
- 1085 Spahl, W. and Budzikiewicz, H.: Quantitative analysis of dental resin composites by gas and liquid chromatography/mass spectrometry. *Fresenius J. Anal. Chem.*, 350 (1994) 684-691.
- 1086 Tazaki, M., Hashida, T., Onodera, A. and Homma, T.: Inverse gas chromatographic study of rubbers. Ethylene propylene rubber (EPR) and acrylonitrile butadiene rubber (NBR). *Nippon Gomu Kyokaishi*, 67 (1994) 55-63; C.A., 121 (1994) 257634t.
- 1087 Tsuge, S. and Ohtani, H.: (Pyrolysis gas chromatography. Composition analysis of macromolecular material.) *Shinsozai*, 5, No. 1 (1994) 30-34; C.A., 121 (1994) 231695p.

See also 735, 803.

32. *DRUG ANALYSIS*32a. *Drug analysis, general techniques*

- 1088 Mulligan, K.J. and McCauley, H.: Factors that influence the determination of residual solvents in pharmaceuticals by automated static headspace sampling coupled to capillary GC-MS. *J. Chromatogr. Sci.*, 33 (1995) 49-54.
- 1089 Rasanen, I., Ojanperä, I. and Vuori, E.: Comparison of internal and external retention index standards in drug screening by gas chromatography. *J. High Resolut. Chromatogr.*, 18 (1995) 66-67.
- 1090 Russo, M.V.: Static headspace gas chromatography of residual solvents in pharmaceutical products. *Chromatographia*, 39 (1994) 645-648.

- 1091 Wilson, I.D., Davis, P. and Ruane, R.J.: Supercritical fluid chromatography and extraction of pharmaceuticals. In: Dean, J.R. (Editor), *Appl. Supercrit. Fluids Ind. Anal.*, Blackie, Glasgow, 1993, pp. 74-103; C.A., 121 (1994) 213076h.
- 1092 Yang, Y.J., Xiang, B.R. and An, D.K.: (Expert system for gas chromatography used in pharmaceutical analysis.) *Yaoxue Xuebao*, 29 (1994) 603-609; C.A., 121 (1994) 213174p.

See also 753.

### 32b. Antirheumatics and antiinflammatory drugs

- 1093 Kim, K.-R., Shin, Y.-J., Shim, W.-H. and Myung, S.-W.: Rapid gas chromatographic profiling and screening of acidic non-steroidal antiinflammatory drugs in biological samples. *Arch. Pharmacol. Res.*, 17 (1994) 175-181; C.A., 121 (1994) 244900b.
- 1094 Laakkonen, U.-M., Leinonen, A. and Savonen, L.: Screening of non-steroidal anti-inflammatory drugs, barbiturates and methyl xanthines in equine urine by gas chromatography-mass spectrometry. *Analyst (Cambridge)*, 119 (1994) 2695-2696.

### 32c. Autonomic and cardiovascular drugs

- 1095 Adam, A., Gervais, N., Panoyan, A., Ong, H., Beliveau, L., Ayotte, C. and Delahaut, P.: Detection of clenbuterol in hair. *Analyst (Cambridge)*, 119 (1994) 2663-2666.
- 1096 Barbato, F., Grumetto, L. and Morrica, P.: Analysis of calcium channel blocking dihydropyridines and their degradation products by gas chromatography. *Farmaco*, 49 (1994) 461-466; C.A., 121 (1994) 213152e.
- 1097 Bargmann-Leyder, N., Thiébaud, D., Vergne, F., Bégos, A., Tambuté, A. and Caude, M.: High-speed chiral separation of beta-blockers by supercritical fluid chromatography on ChyRoSine-A. *Chromatographia*, 39 (1994) 673-681.
- 1098 Liu, F., Hu, X. and Li, Q.: Quantitative analysis of bencyonate in human plasma using a deuterated internal standard by gas chromatography-mass spectrometry with selected-ion monitoring. *J. Chromatogr. B*, 662 (1994) 123-127.
- 1099 Liu, X., Zhang, Y.Z., Zhang, C.J. and Ye, L.: (Determination of salbutamol in human urine.) *Yaoxue Xuebao*, 29 (1994) 454-458; C.A., 121 (1994) 194817g.
- 1100 Minnigh, M.B., Alvin, J.D. and Zemaitis, M.A.: Determination of plasma mexiletine levels with gas chromatography-mass spectrometry and selected-ion monitoring. *J. Chromatogr. B*, 662 (1994) 118-122.
- 1101 Ticehurst, M.D., Rowe, R.C. and York, P.: Determination of the surface properties of two batches of salbutamol sulfate by inverse gas chromatography. *Int. J. Pharm.*, 111, No. 3 (1994) 241-249; C.A., 121 (1994) 212823n.
- 1102 Van der Merwe, P.J. and Hendrikz, S.E.: Identification of ephedrine as their carbon disulfide derivatives. *J. Chromatogr. B*, 663 (1995) 160-166.
- 1103 Visser, T., Vredenburg, M.J., de Jong, A.P.J.M., van Rossum, H.J., Stephany, R.W. and van Ginkel, L.A.: Confirmation analysis of beta-agonists by cryotrapping gas chromatography-Fourier transform infrared spectrometry. *Analyst (Cambridge)*, 119 (1994) 2681-2685.

See also 995, 1000.

### 32d. Central nervous system drugs

- 1104 Cui, J., Li, N., Cui, K., Zhou, Y. and Zhou T.: (Identification of amphetamines in urine by GC and GC-MS.) *Yaowu Fenxi Zazhi*, 14, No. 4 (1994) 3-7; C.A., 121 (1994) 244907j.
- 1105 DeRuiter, J., Clark, C.R. and Noggle, F.T.: Gas chromatographic and mass spectral analysis of amphetamine products synthesized from 1-phenyl-2-nitropropene. *J. Chromatogr. Sci.*, 32 (1994) 511-519.
- 1106 DeRuiter, J., Hayes, L., Valaer, A., Clark, C.R. and Noggle, F.T.: Methcathinone and designer analogues: Synthesis, stereochemical analysis, and analytical properties. *J. Chromatogr. Sci.*, 32 (1994) 552-564.
- 1107 Edinboro, L.E. and Poklis, A.: Detection of benzodiazepines and tribenzazams by TRIAGE: confirmation by solid-phase extraction utilizing SPEC.3ML.MP3 microcolumns and GC-MS. *J. Anal. Toxicol.*, 18 (1994) 312-316; C.A., 121 (1994) 197806p.
- 1108 Hayes, M.J., Khemani, L. and Powell, M.L.: Quantitative determination of the NMDA antagonist cis-4-phosphonomethyl-2-piperidine carboxylic acid (CGS 19755) in human plasma using capillary gas chromatography/mass spectrometry. *Biol. Mass Spectrom.*, 23, No. 9 (1994) 555-561; C.A., 121 (1994) 194814d.
- 1109 Humbert, L., Jacquemont, M.C., Leroy, E., Leclerc, F., Houdret, N. and Lhermitte, M.: Determination of chloral hydrate and its metabolites (trichloroethanol and trichloroacetic acid) in human plasma and urine using electron capture gas chromatography. *Biomed. Chromatogr.*, 8 (1994) 273-277.
- 1110 Lartigue-Mattei, C., Galmier, M.J., Chabard, J.L., Beyssac, E. and Aiache, J.M.: Determination of oxeladin in human plasma by gas chromatography-mass spectrometry. *Biomed. Chromatogr.*, 9 (1995) 32-35.
- 1111 Leis, H.J., Windischhofer, W. and Wintersteiger, R.: Quantitative measurement of amphetamine in human plasma by gas chromatography/negative ion chemical ionization mass spectrometry using (<sup>2</sup>H<sub>5</sub>)amphetamine as internal standard. *Biol. Mass Spectrom.*, 23 (1994) 637-641; C.A., 121 (1994) 271147t.
- 1112 Li, B.: (Quantitative determination of isoflurane by gas chromatography (GC).) *Chin. J. Chromatogr.*, 12 (1994) 453-454.
- 1113 Rao, M.L., Staberock, U., Baumann, P., Hiemke, C., Deister, A., Cuendet, C., Arney, M., Härtler, S. and Kraemer, M.: Monitoring tricyclic antidepressant concentrations in serum by fluorescence polarization immunoassay compared with gas chromatography and HPLC. *Clin. Chem. (Washington)*, 40 (1994) 929-933.
- 1114 Shin, H.S., Park, J.-S., Park, P.-B. and Yun, S.J.: Detection and identification of famprofazone and its metabolite in human urine. *J. Chromatogr. B*, 661 (1994) 255-261.
- 1115 Shinohara, Y., Nagao, K., Akutsu, N. and Baba, S.: Assessment of the metabolic chiral inversion of suprofen in rat by gas chromatography-mass spectrometry combined with a stable isotope technique. *J. Pharm. Sci.*, 83 (1994) 1521-1523; C.A., 121 (1994) 194898j.
- 1116 Thompson, W.C. and Dasgupta, A.: Microwave-induced rapid preparation of fluoroderivatives of amphetamine, methamphetamine and 3,4-methylenedioxymethamphetamine for GC-MS confirmation assays. *Clin. Chem. (Washington)*, 40 (1994) 1703-1706; C.A., 121 (1994) 223776a.

- 1117 Tonn, G.R., Abbott, F.S., Rurak, D.W. and Axelson, J.E.: Simultaneous analysis of diphenylmethoxyacetic acid, a metabolite of diphenhydramine, and its deuterium-labeled stable isotope analog in ovine plasma and urine. *J. Chromatogr. B*, 663 (1995) 67-81.
- 1118 Zhang, Q., Wu, L. and Zhao, X.: (Determination of compound amidopyrine injection by capillary gas chromatography.) *Xiangtan Daxue Ziran Kexue Xuebao*, 16 (1994) 37-39; C.A., 121 (1994) 213137d.
- 1119 Zhang, Y. and Ouyang, X.: (Quantitative determination of phenobarbital and its preparations by gas chromatography.) *Chin. J. Chromatogr.*, 13, No. 1 (1995) 38-39.

See also 1018, 1024, 1126.

### 32e. Chemotherapeutics (exc. cytostatics and antibiotics)

- 1120 Awata, N., Toba, F., Ando, M., Shimada, H., Miyairi, S., Kato, T., Goto, J. and Nambara, T.: Immunoaffinity extraction of 4-hydroxy-2-(4-methylphenyl)benzothiazole and its metabolites for determination by gas chromatography-mass spectrometry. *Biol. Pharm. Bull.*, 17 (1994) 843-845; C.A., 121 (1994) 244904f.
- 1121 Neal, J.M., Howald, W.N., Kunze, K.L., Lawrence, R.F. and Trager, W.F.: Application of negative-ion chemical ionization isotope dilution gas chromatography-mass spectrometry to single-dose bioavailability studies of mefloquine. *J. Chromatogr. B*, 661 (1994) 263-269.
- 1122 Yan, J.-s. and Sun L.-l.: (Gas chromatography for determination of febuprol and its preparation.) *Zhongguo Yiyao Gongye Zazhi*, 25 (1994) 215-216; C.A., 121 (1994) 213167p.

### 32g. Other drug categories

See 1127.

### 32h. Toxicological and forensic applications

- 1123 Flanagan, R.J.: Gas chromatography in analytical toxicology: principles and practice. In: Baugh, P.J. (Editor), *Gas chromatography*, IRL, Oxford, 1993, pp. 171-212; C.A., 121 (1994) 172318t - a review with 100 refs.
- 1124 Kicman, A.T., Cowan, D.A., Myhre, L., Nilsson, S., Tomten, S. and Oftebro, H.: Effect on sports drug tests of ingesting meat from steroids (methenolane)-treated livestock. *Clin. Chem. (Washington)*, 40 (1994) 2084-2087.
- 1125 Pedroso, R.C., Salvadori, M.C., Andraus, M.H. and Lopez, N.M.R.: Determination of succinonitrile in horse urine by gas chromatography-nitrogen-phosphorus detector and gas chromatography-mass spectrometry. *Analyst (Cambridge)*, 119 (1994) 2697-2699.
- 1126 Peytavin, G., Dautrey, S., Chaminade, P., Chastre, F. and Fariotti, R.: (Carbamazepine overdose and toxicological screening.) *J. Pharm. Clin.*, 13 (1994) 131-139; C.A., 121 (1994) 271987s.
- 1127 Pohjola, J. and Harpf, M.: Determination of atropine and obidoxime in automatic injection devices used as antidotes against nerve agent intoxication. *J. Chromatogr. A*, 686 (1994) 350-354.

- 1128 Rowland, B.J., Irvin, J. and Keith, E.S.: Increased detection of marijuana use with a 50 µg/L urine screening cutoff. *Clin. Chem. (Washington)*, 40 (1994) 2114-2115.
- 1129 Yin, L.-B.: Classification results of mass spectra of toxic compounds by class modeling. *J. Chem. Inf. Comput. Sci.*, 34 (1994) 1232-1234; C.A., 121 (1994) 248141c.

See also 887, 1004.

### 32i. Plant extracts

- 1130 Zhang, L. and Fang, H.: (GC analysis of volatile oils in medicinal plants.) *Yaowu Fenxi Zazhi*, 14, No. 3 (1994) 52-60; C.A., 121 (1994) 213075g - a review with 10 refs.

See also 992.

## 33. CLINICO-CHEMICAL APPLICATIONS

### 33b. Complex mixtures and profiling (single compounds by cross-reference only)

- 1131 Bennett, M.J., Powell, S., Swartling, D.J. and Gibson, K.M.: Tiglylglycine excreted in urine in disorders of isoleucine metabolism and the respiratory chain measured by stable isotope dilution GC-MS. *Clin. Chem. (Washington)*, 40 (1994) 1879-1893.
- 1132 Lennart, L.: Determination of microbial chemical markers by gas chromatography-mass spectrometry - potential for diagnosis and studies on metabolism *in situ*. *APMIS*, 102 (1994) 161-169; C.A., 121 (1994) 174156n.

See also 905, 917, 918, 920, 929, 958, 964, 971, 1007, 1022.

## 34. FOOD ANALYSIS

### 34b. Complex mixtures (single compounds by cross-reference only)

- 1133 Chisholm, M.G., Guiher, L.A., Vonah, T.M. and Beaumont, J.L.: Comparison of some French-American hybrid wines with White Riesling using gas chromatography-olfactometry. *Am. J. Enol. Vitic.*, 45 (1994) 201-212; C.A., 121 (1994) 279411v.
- 1134 Chung, M.S., Lee, M. and Kang, T.S.: (DHS/GC/MSD analysis of volatile compounds of commercial raw ground beef). *Han'guk Ch'uksan Hanhoechi*, 36 (1994) 192-198; C.A., 121 (1994) 279143j.
- 1135 Grigoryeva, D.N., Golovnya, R.V., Syomina, L.A., Vasilyev, A.V. and Samusenko, A.L.: Correlation between sensory and gas chromatographic characteristic of Dutch cheese. *Dev Food Sci.*, 35 (1994) 283-286; C.A., 121 (1994) 279321r.
- 1136 Iverson, W.G.: Ethyl acetate extraction of beer: quantitative determination of additional fermentation byproducts. *J. Am. Soc. Brew. Chem.*, 52 (1994) 91-95; C.A., 121 (1994) 178024r.

- 1137 Riganakos, K.A. and Kontominas, M.G.: GC study of the effect of specific heat treatment on the sorption of water by wheat and soy flour. *Z. Lebensm.-Unters. Forsch.*, 198 (1994) 47-51; C.A., 121 (1994) 178253q.
- 1138 Robards, K. and Antolovich, M.: Methods for assessing the authenticity of orange juice. A review. *Analyst (Cambridge)*, 120 (1995) 1-28 - a review with 338 refs.
- 1139 Tateo, F., Ferrillo, A. and Orlandi, A.: Correlation between aromatic qualities and GC-MS composition of ready-made tomato sauces. *Dev Food Sci.*, 35 (1994) 107-112; C.A., 121 (1994) 279360c.

See also 820, 837, 841, 848, 864, 886, 903, 919, 926, 939, 942, 948, 956, 975, 976, 977, 981, 1023, 1025, 1036, 1045, 1051, 1059, 1060, 1063, 1064, 1070, 1141, 1160.

34c. *Organoleptically important compounds (flavors, odors, volatiles)*

- 1140 Ang, C.Y.W., Liu, F. and Sun, T.: Development of a dynamic headspace GC method for assessing the influence of heating end-point temperature on volatiles of chicken breast meat. *J. Agric. Food Chem.*, 42 (1994) 2493-2498; C.A., 121 (1994) 254065x.
- 1141 Aylott, R.I., Clyne, A.H., Fox, A.P. and Walker, D.A.: Analytical strategies to confirm Scotch whisky authenticity. *Analyst (Cambridge)*, 119 (1994) 1741-1746.
- 1142 Farkas, P., le Quere, J.L., Maarse, H. and Kovac, M.: The standard GC retention index library of flavor compounds. *Dev Food Sci.*, 35 (1994) 145-149; C.A., 121 (1994) 279181v.
- 1143 Forster, A. and Schmidt, R.: (Characterization and classification of hop varieties.) *Brauwelt*, 133 (1993) 2036-2038, 2049-2057; C.A., 121 (1994) 254308d.
- 1144 Imhof, R. and Bosset, J.O.: Quantitative GC-MS analysis of volatile flavor compounds in pasteurized milk products applying a standard addition method. *Food Sci. Technol. (London)*, 27 (1994) 265-269; C.A., 121 (1994) 253999t.
- 1145 King, B.M., de Heij, H.T. and Jagers, P.P.J.M.: Flavor concentration adjustment: correlation between GC-headspace measurement and sensory evaluation. *Dev Food Sci.*, 35 (1994) 277-281; C.A., 121 (1994) 279274c.
- 1146 Legger, A. and Roozen, J.P.: Gas chromatography and sensory analysis of chocolate flavor: intensity changes in time. *Dev Food Sci.*, 35 (1994) 287-291; C.A., 121 (1994) 279275d.
- 1147 Mahungu, S.M., Hansen, S.L. and Artz, W.E.: Quantification of volatile compounds in heated trilinolein by static headspace-capillary gas chromatography/infrared spectroscopy-mass spectrometry. *J. Am. Oil Chem. Soc.*, 71 (1994) 1169-1171.
- 1148 Schreier, P.: (Aromaforschung aktuell.) *Naturwissenschaften*, 82 (1995) 21-27.
- 1149 Van Ruth, S.M. and Roozen, J.P.: Gas chromatographic/sniffing port analysis and sensory evaluation of commercially dried bell pepper (*Capsicum annuum*) after rehydration. *Food. Chem.*, 51 (1994) 165-170; C.A., 121 (1994) 203820g.

See also 745, 767, 804, 809, 811, 986, 1133, 1138.

35. ENVIRONMENTAL ANALYSIS

35a. *General papers and reviews*

- 1150 Ilias, A.M. and Jaeger, C.: Evaluation of sampling techniques for the analyses of volatile and total recoverable petroleum hydrocarbons (TRPH) by IR, GC, and GC/MS methods. *Hydrocarbon Contam. Soils*, 3 (1994) 147-165; C.A., 121 (1994) 204122t.
- 1151 Reeve, R.N.: *Environmental Analysis*. Wiley, Winchester, 1994, 263 p.

See also 828, 852, 895.

35b. *Air pollution (complex mixtures; single compounds by cross-reference only)*

- 1152 Hoffmann, T.: Adsorptive preconcentration technique including oxidant scavenging for the measurement of reactive natural hydrocarbons in ambient air. *Fresenius J. Anal. Chem.*, 351 (1995).
- 1153 Patil, S.F. and Lonkar, S.T.: Determination of benzene, aniline and nitrobenzene in workplace air: a comparison of active and passive sampling. *J. Chromatogr. A*, 688 (1994) 189-199.

See also 839, 840, 845, 850, 857, 859, 866, 876, 879, 880, 894, 901, 907, 914, 927, 940, 961, 980, 1005, 1027, 1038, 1048, 1129, 1180.

35c. *Water pollution (complex mixtures; single compounds by cross-reference only)*

- 1154 House, W.A.: Sampling techniques for organic substances in surface waters. *Intern. J. Environ. Anal. Chem.*, 57 (1994) 207-214.
- 1155 Xu, Y.H. and Mitra, S.: Continuous monitoring of volatile organic compounds in water using on-line membrane extraction and microtrap gas chromatography system. *J. Chromatogr. A*, 688 (1994) 171-180.

See also 749, 812, 813, 851, 865, 868, 906, 993, 1001, 1049, 1054, 1061, 1069, 1078, 1159, 1181.

35d. *Soil pollution (complex mixtures; single compounds by cross-reference only)*

- 1156 Hawthorne, S.B. and Miller, D.J.: Direct comparison of Soxhlet and low- and high-temperature supercritical CO<sub>2</sub> extraction efficiencies of organic from environmental solids. *Anal. Chem.*, 66 (1994) 4005-4012.
- 1157 Kolb, M., Böhm, H.B. and Bahadir, M.: Analytical multimethod for the determination of low volatile organic pollutants in sediments and sewage sludges. *Fresenius J. Anal. Chem.*, 351 (1995) 286-296.
- 1158 Somitsch, W., Jerney, J. and Robra, K.-H.: Quantitative analysis and characterization of mineral oil contaminated soil by GC-data processing. *Fresenius Environ. Bull.*, 3, No. 6 (1994) 358-364; C.A., 121 (1994) 204127y.

See also 851, 856, 864, 881, 885, 899, 922, 946, 993, 1021, 1030, 1055, 1058, 1066, 1069, 1150, 1165.

### 36. SOME TECHNICAL PRODUCTS AND COMPLEX MIXTURES

#### 36a. Surfactants

1159 Kane, M., Dean, J.R., Hitchen, S.M., Dowle, C.J. and Tranter, R.L.: Extraction of surfactants from aqueous media by supercritical fluid extraction. *Analyst (Cambridge)*, 120 (1995) 355-357.

See also 889.

#### 36b. Antioxidants and preservatives

1160 Ishiwata, H., Sugita, T. and Yamada, T.: (Determination of tetramethylsuccinonitrile in foods by NPD-GC.) *Shokuhin Eiseigaku Zasshi*, 35 (1994) 382-389; C.A., 121 (1994) 203654f.

See also 798, 812.

#### 36c. Complex mixtures, technical products and unidentified compounds

1161 Banba, A., Nishita, A., Tajima, R., Mine, T., Amachi, T., Masuda, M. and Izumi, T.: (Prevention of musty odor (TCA) migration into food.(1). Development of a method for the determination of the substances related to musty odor (TCA,TCP) and a survey of corrugated cardboard boxes.) *Nippon Hosho Gakkaishi*, 3 (1994) 35-44; C.A., 121 (1994) 279167v.

1162 Diehl, J.W., Finkbeiner, J.W. and di Sanzo, F.P.: Determination of ethers and alcohols in reformulated gasolines by gas chromatography/atomic emission detection. *J. High Resolut. Chromatogr.*, 18 (1995) 108-110.

1163 Ehrmann, U. and Izquierdo, A.: Characterization of vacuum resids, hydrocracked products and asphaltenes by pyrolysis-gas chromatography (FID-FPD). *Fuel Sci. Technol. Int.*, 12 (1994) 1193-1220; C.A., 121 (1994) 234214s.

1164 Harriman, G.E.: The role of gas chromatography in petroleum exploration. In: Baugh, P.J. (Editor), *Gas Chromatography*, IRL, Oxford, 1993, pp. 331-357; C.A., 121 (1994) 208584t - a review with 13 refs.

1165 Kuckuk, R., Hill, W., Burba, P. and Davies, A.N.: Pyrolysis-GC-FTIR for structural elucidation of aquatic humic substances. *Fresenius J. Anal. Chem.*, 350 (1994) 528-532.

1166 Marchand, M., Termonia, M., Caprais, J.C. and Wybauw, M.: Purge and trap GC-MS analysis of volatile organic compounds from the Guaymas Basin hydrothermal site (Gulf of California). *Analisis*, 22 (1994) 326-331.

1167 Oono, A., Kosakai, N. and Inokawa, T.: (Qualitative and quantitative analyses of organic substances in cement-organic additive mixtures and organic additive-containing cement products.) *Jpn. Kokai Tokkyo Koho* JP 06,194,358 [94,194,358] (Cl. G01N30/86), 15 Jul. 1994, Appl. 92/342,111, 22 Dec. 1992; 13 pp.; C.A., 121 (1994) 262048j.

1168 Peters, R.J.B., de Leer, E.W.B. and Versteegh, J.F.M.: Identification of halogenated compounds produced by chlorination of humic acid in the presence of bromide. *J. Chromatogr. A*, 686 (1994) 253-261.

1169 Yinon, J., Yost, R.A. and Bulusu, S.: Thermal decomposition characterization of explosives by pyrolysis-gas chromatography-mass spectrometry. *J. Chromatogr. A*, 688 (1994) 231-242.

See also 729, 853, 855, 869, 875, 882, 890, 999, 1174.

### 37. CELLS, CELLULAR PARTICLES AND SUPRAMOLECULAR STRUCTURES

See 741, 943, 944.

### 38. INORGANIC COMPOUNDS

#### 38a. Cations

1170 Oguma, K., Shimizu, T. and Kuroda, R.: (Advances in chromatography of the rare earth elements (Review).) *Bunseki Kagaku*, 1 (1995) 1-17 - a review with 161 refs.

See also 834.

#### 38b. Anions

1171 Tsikas, D., Böger, R.H., Bode-Böger, S.M., Gutzki, F.-M. and Frölich, J.C.: Quantification of nitrite and nitrate in human urine and plasma as pentafluorobenzyl derivatives by gas chromatography-mass spectrometry using their <sup>15</sup>N-labelled analogs. *J. Chromatogr. B*, 661 (1994) 185-191.

See also 742.

#### 38c. Permanent and rare gases

1172 Atanasiu, F., Mihailescu, G.: Hydrogen purification by PSA method for gas chromatography uses. *Process Technol. Proc.*, 11 (1994) 259-263; C.A., 121 (1994) 182803m.

1173 De Vanssay, E., Zubrzycki, S., Sternberg, R., Raulin, F., Sergeant, M. and Phan-Tan-Luu, R.: Gas chromatography of Titan's atmosphere. V. Determination of permanent gases in the presence of hydrocarbons and nitriles with molecular sieve micropacked column and optimization of the GC parameters using a Doehlert experimental design. *J. Chromatogr. A*, 688 (1994) 161-170.

1174 Frisina, G., Testoni, F., Furlan, V. and Ostrander, C.: Determination of carbon monoxide in ethylene and propylene at concentration levels as low as 1 part per billion. *LC-GC Int.*, 8 (1995) 96-101.

1175 Yin, G. and Yao, T.: (Determination of dissolved argon in water by gas chromatography.) *Hedongli Gongcheng*, 15 (1994) 361-364; C.A., 121 (1994) 286191b.

See also 748.

## 38d. Volatile inorganic compounds

- 1176 Behar, O., Deri, F., Gueriane, S. and Oukrid, N.: The study by gas chromatography of the products of pyrolysis of fire retarded cellulose with chlorides and bromides. *J. Soc. Alger. Chim.*, 3 (1993) 1-8; C.A., 121 (1994) 257791s.
- 1177 Ducros, V., Ruffieux, D., Belin, N. and Favier, A.: Comparison of two digestion methods for the determination of selenium in biological samples. *Analyst (Cambridge)*, 119 (1994) 1715-1717.
- 1178 Shahi, P. and Chakma, A.: A gas chromatographic technique for the accurate determination of CO<sub>2</sub> in gas treating solvents. *Process Technol. Proc.*, 11 (1994) 937-943; C.A., 121 (1994) 208727s.
- 1179 Shahi, P., Hu, Y.-F. and Chakma, A.: Gas chromatographic analysis of acid gases and single/mixed alkanolamines. *J. Chromatogr. A*, 687 (1994) 121-132.
- 1180 Swan, H.B. and Ivey, J.P.: Analysis of atmospheric sulfur gases by capillary gas chromatography with atomic emission detection. *J. High Resolut. Chromatogr.*, 17 (1994) 814-820.
- 1181 Wang, Y., Li, X., Zhang, J., Wang, X. and Shi, T.: (Determination of carbon disulfide by gas chromatography.) *Huanjing Huaxue*, 13 (1994) 359-362; C.A., 121 (1994) 263060u.
- 1182 Wasiak, W. and Szczepaniak, W.: (Gas chromatography as a method for moisture determination in water-unstable pharmaceutical preparations.) *Acta Pol. Pharm.*, 50 (1993) 423-426; C.A., 121 (1994) 213177s.

See also 1027.

## 39. RADIOACTIVE AND OTHER ISOTOPE COMPOUNDS

- 1183 Meier-Augenstein, W.: On-line recording of <sup>13</sup>C/<sup>12</sup>C ratios and mass spectra in one gas chromatographic analysis. *J. High Resolut. Chromatogr.*, 18 (1995) 28-32.

See also 748, 805, 807, 1171.

## Planar Chromatography

### 1. REVIEWS AND BOOKS

- 213 Barcza, L.: (Molecular recognition and its relation to pharmaceutical analysis). *Gyógyszerészet*, 38 (1994) 477-484; C.A., 122 (1995) 17280b - a review with 18 refs.
- 214 Davis, J.M.: Statistical theories of peak overlap in chromatography. *Adv. Chromatogr. (N.Y.)*, 34 (1994) 109-176; C.A., 121 (1994) 238951c - a review with 52 refs.
- 215 Fried, B. and Sherma, J. (Editors): *Thin-Layer Chromatography: Techniques and Applications*, third-edition revised and expanded. In: *Chromatogr. Sci. Ser.*, 1994, Vol. 66, Dekker, New York, 1994, 451 pp.; C.A., 121 (1994) 270855k.
- 216 Kumar, M.: Recent trends in chromatographic procedures for separation and determination of rare earth elements. *Analyst (Cambridge)*, 119 (1994) 2013-2024.
- 217 Szepeszy, L.: (Development and importance of separation techniques in instrumental analysis). *Magy. Kem. Foly.*, 1100 (1994) 340-353; C.A., 121 (1994) 270751y - a review with 121 refs.

### 2. FUNDAMENTALS, THEORY AND GENERAL

#### 2a. General

- 218 Nyiredy, S., Fatér, Z. and Szabady, B.: Identification in planar chromatography by use of retention data measured using characterized mobile phases. *J. Planar Chromatogr.*, 7 (1994) 406-409.
- 219 Rozylo, J.K., Janicka, M. and Siembida, R.: Advantages of TLC as a pilot technique for HPLC. *J. Liq. Chromatogr.*, 17 (1994) 3641-3653.

#### 2b. Thermodynamics and theoretical relationships

- 220 Kunugi, A. and Tabei, K.: (Basic experiments on TLC. 1. Effect of temperature on  $R_f$  values). *Kagaku to Kyoiku*, 42 (1994) 206-210; C.A., 121 (1994) 270773g.

#### 2c. Relationship between structure and chromatographic behaviour

- 221 Pyka, A.: A new stereoisomeric topological index ( $I_{STI}$ ) for predicting the separation of stereoisomers in TLC. Part VII. *J. Planar Chromatogr.*, 7 (1994) 389-393.

See also 250, 302.

### 3. GENERAL TECHNIQUES

#### 3a. Apparatus and accessories

- 222 Partney, D.W., Jr.: Apparatus for thin layer chromatography. U.S. US 5,350,510 (Cl. 210-198.3; B01D15/08), 27 Sep. 1994, Appl. 16,789, 11 Feb. 1993; 7 pp.; C.A., 121 (1994) 270730r.

See also 231.

#### 3b. Detectors and detection reagents

- 223 Fijalek, Z. and Snyckerski, A.: Electrochemical and TLC-densitometric analysis of chlorambucil and its decomposition product. *Acta Pol. Pharm.*, 50 (1993) 301-306; C.A., 121 (1994) 291623y.

See also 255, 261, 262, 296, 348, 351, 358.

#### 3c. Sorbents and columns, packing procedures

- 224 Chowdary, K.P.R. and Rao, P.V.: Biopharmaceutical studies on solid dispersions of nalidixic acid in modified starches. *Drug Dev. Ind. Pharm.*, 20 (1994) 3015-3022; C.A., 121 (1994) 263486n.
- 225 Coman, V. and Marutoiu, C.: The preparation of *n*-octyl-bonded volcanic tuff as stationary phase for TLC. *J. Planar Chromatogr.*, 7 (1994) 450-452.
- 226 Hauck, H.E., Junker-Buchheit, A. and Wenig, R.: Reproducibility of Silica gel 60 precoated plates in TLC and HPTLC. *LC-GC Int.*, 8, No. 1 (1995) 34-38.
- 227 Kaku, S.: (Development of an optical immunoassay method). *SSM Tech. Rep.*, 5 (1993) 72-76; C.A., 121 (1994) 296463f.
- 228 Marutoiu, C. and Cenan, A.: (Production of cellulose triacetate). *Rom. RO 103,691 (Cl. C08B3/06)*, 15 Dec. 1993, Appl. 141,516, 08 Sep. 1989; 3 pp.; C.A., 121 (1994) 303354b.
- 229 Mudasingana, M.L., Noeva, V.A., Nechaeva, E.B., Prokof'eva, V.I. and Sokolova, T.M.: (Thin-layer chromatographic identification of phenothiazine derivatives on various sorbents). *Farmatsiya (Moscow)*, 42, No. 1 (1993) 58-59; C.A., 121 (1994) 263810p.

See also 213, 258, 303, 306, 337, 349, 397.

#### 3d. Quantitative analysis

- 230 Kang, J.S.: (Normalization and search of the UV/VIS spectra measured from TLC/HPTLC). *Yakhak Hoechi*, 38 (1994) 366-371; C.A., 121 (1994) 250180h.

See also 350.

3e. *Preparative scale chromatography*

- 231 Läufer, K., Lehmann, J., Petry, S., Scheuring, M. and Schmidt-Schuchardt, M.: Simple, inexpensive system for using thin-layer chromatography for micro-preparative purposes. *J. Chromatogr. A*, 684 (1994) 370-373.

See also 375.

3f. *Programmed temperature, pressure, vapors, gradients*

See 220, 233.

3g. *High performance procedures*

- 232 Belay, M.: *Method Development and Solvent Optimization for the Separation of Pharmaceutically and Biologically Important Compounds by High-Performance Planar Chromatography and Automated Multiple Development*. Wayne State Univ., 1993, 221 pp.; C.A., 121 (1994) 270804t.

## 4. SPECIAL TECHNIQUES

4a. *Automation*

See 347.

4b. *Computerization and modelling*

- 233 Qin-Sun, W., Bing-Wen, Y. and Zhi-Chao, Z.: Optimization of stepwise gradient HPTLC by statistical scanning. *J. Planar Chromatogr.*, 7 (1994) 477-480.
- 234 Wang, Q.-S. and Zhan, Z.-P.: A method for optimizing the HPTLC separation of a mixture of unknown composition. *J. Planar Chromatogr.*, 7 (1994) 394-398.

See also 302, 357.

4c. *Combination with other physico-chemical techniques (MS, IR etc.)*

- 235 Van de Nesse, R.J., Vinkenburg, I.H., Jonker, R.H.J., Hoornweg, G.Ph., Gooijer, C., Brinkman, U.A.T. and Velthorst, N.H.: Fluorescence line-narrowing spectroscopy as an off-line identification method for narrow-bore column liquid chromatography. *Appl. Spectrosc.*, 48 (1994) 788-795; C.A., 121 (1994) 270838g.
- 236 Wollf, S.C. and Kovar, K.-A.: Direct HPTLC-FTIR measurement in combination with AMD. *J. Planar Chromatogr.*, 7 (1994) 344-348.

See also 243, 281, 298.

4g. *Enantiomers, separation*

- 237 Aboul-Enein, J.Y. and Serignese, V.: Optical purity determination of thyroxine enantiomers in bulk materials by chiral thin layer chromatography. *Biomed. Chromatogr.*, 8 (1994) 317-318; C.A., 122 (1995) 17310m.

- 238 Lepri, L., Coas, V., Desideri, P.G. and Zocchi, A.: Reversed phase planar chromatography of enantiomeric compounds on triacetylcellulose. *J. Planar Chromatogr.*, 7 (1994) 376-381.

See also 221, 305, 306.

4h. *Other special techniques*

See 227, 359.

## 5. HYDROCARBONS AND HALOGEN DERIVATIVES

5b. *Cyclic hydrocarbons, fullerenes*

See 390.

5d. *Complex hydrocarbon mixtures (incl. analysis of tars, bitumens and mineral oils)*

- 239 Dhole, V.R. and Ghosal, G.K.: Rapid detection and characterization of petroleum products and residues by TLC: Part I. *J. Planar Chromatogr.*, 7 (1994) 469-471.
- 240 Whittaker, M. and Pollard, S.J.T.: Characterization of refractory wastes at hydrocarbon-contaminated sites: 1. Rapid column fractionation and thin layer chromatography of reference oils. *J. Planar Chromatogr.*, 7 (1994) 354-361.

## 6. ALCOHOLS

See 238.

## 7. PHENOLS

See 238, 260, 394.

## 8. SUBSTANCES CONTAINING HETEROCYCLIC OXYGEN

- 241 Gizella, P., Krawczyk, U. and Agnes, Es Kery: (Spectrophotometric and chromatographic investigation of bilberry anthocyanins for qualification purposes). *Acta Pharm. Hung.*, 64 (1994) 117-122; C.A., 121 (1994) 308449w.

8a. *Flavonoids*

- 242 Costa, S.S., Jossang, A., Bodo, B., Souza, M.L.M. and Moraes, V.L.G.: Patuletin acetylramnosides from *Kalanchoe brasiliensis* as inhibitors of human lymphocyte proliferative activity. *J. Natural Prod.*, 57 (1994) 1503-1510.

See also 394.

8b. *Aflatoxins and other mycotoxins*

- 243 Gelosa, M. and Buzzetti, I.: (Chromatographic techniques for aflatoxin determination in milk). *Ind. Aliment. (Pinerolo)*, 33 (1994) 315-317; C.A., 121 (1994) 229127m.

8c. *Other compounds with heterocyclic oxygen (incl. tannins)*

See 331.

## 9. OXO COMPOUNDS, ETHERS, EPOXIDES AND QUINONES

- 244 Li, G. and Zhen, H.: (Determination of hesperidin and total anthraquinones in Chenxianglubailu tablets by dual-wavelength TLC scanning). *Zhongcaoyao*, 25 (1994) 359-361; C.A., 121 (1994) 263825x.
- 245 Wu, P.: (Thin layer chromatography and quantitative determination of total anthraquinone in Jiiejin oral liquid). *Huaxue Yaoxue Zazhi*, 9 (1994) 165-168; C.A., 121 (1994) 213108v.

See also 378, 388.

## 10. CARBOHYDRATES

10a. *Mono and oligosaccharides. Structural studies*

- 246 Guo, X., Sun, H. and Yu, K.: (Determination of physical and chemical constants and separation as well as quantification of isosorbide nitrates). *Yaowu Fenxi Zazhi*, 14, No. 4 (1994) 37-41; C.A., 121 (1994) 213146f.
- 247 Taylor, W.G., Danielson, T.J. and Orcutt, R.L.: Thin-layer chromatographic detection of ivermectin in cattle serum. *J. Chromatogr. B*, 661 (1994) 327-333.

10c. *Glycoproteins and their constituents*

- 248 Cook, D.G., Fantini, J., Spitalnik, S.L. and Gonzales-Scarano, F.: Application of high-performance thin-layer chromatography to investigate interactions between human immunodeficiency virus type 1 glycoprotein gp 120 and galactosylceramide. *Methods Mol. Genet.*, (1994) 86-92; C.A., 121 (1994) 276116s.

## 11. ORGANIC ACIDS AND LIPIDS

11a. *Organic acids and simple esters*

- 249 Chen, G. and Wang, Q.: (TLC-UV method for quantitative determination of salicylic acid in Suxiaozhiyang tincture). *Zhongguo Yiyuan Yaoxue Zazhi*, 14 (1994) 271-272, 288; C.A., 121 (1994) 308514p.
- 250 Gocan, S., Cimpan, G. and Panea, T.: Normal phase TLC and reversed phase HPTLC of some plant growth stimulators. *J. Planar Chromatogr.*, 7 (1994) 435-439.
- 251 Herzog, R. and Leuschner, J.: Pharmacokinetic studies of benzalazine. *Arzneim.-Forsch.*, 44 (1994) 1381-1383.

- 252 Jung, S. and Hollingsworth, R.L.: Structures and stereochemistry of the very long  $\alpha,\omega$ -bifunctional alkyl species in the membrane of *Sarcina ventriculi* indicate that they are formed by tail-to tail coupling of normal fatty acids. *J. Lipid Res.*, 35 (1994) 1932-1945.

- 253 Li, Z., Wei, H., Liu, Y., Hao, C. and Fan, G.: (Quantitative determination of oleanolic acid in *Swertia pseudochinensis* Hara by TLC-densitometry method). *Hebei Yixueyuan Xuebao*, 15 (1994) 68-71; C.A., 121 (1994) 213084j.

- 254 Mueller, V., Peterhaensel, M. and von Lengeken, J.: (A rapid method for the semiquantitative determination of unwanted fermentation acids in silages). *Agribiol. Res.*, 46 (1993) 340-345; C.A., 121 (1994) 253988p.

- 255 Nikolova-Damyanova, B., Christie, W.W. and Herslof, B.: Improved separation of some positional isomers of monounsaturated fatty acids, as their phenacyl derivatives, by silver-ion thin layer chromatography. *J. Planar Chromatogr.*, 7 (1994) 382-385.

- 256 Suzuki, M. and Ishihara, K. and Takitani, S.: (Study on purity test of isocitric acid in citric acid by thin-layer chromatography). *Iyakuhi Kenkyo*, 25 (1994) 202-204; C.A., 121 (1994) 213114u.

- 257 Ueda, N., Yamamoto, K., Yamamoto, S., Tokunaga, T., Shirakawa, E., Shinkai, H., Ogawa, M., Sato, T., Kudo, I., Inoue, K., Takizawa, H., Nagano, T. et al.: Lipoxygenase-catalyzed oxygenation of arachidonylethanolamide, a cannabinoid receptor agonist. *Biochim. Biophys. Acta*, 1254 (1995) 127-134.

- 258 Vulimiri, S.V., Smith, C.V., Randerath, E. and Randerath, K.:  $^{32}\text{P}$ -Postlabeling of bile components: bulky adduct-like behavior in polyethyleneimine-cellulose thin layer chromatography. *Carcinogenesis*, 15 (1994) 2061-2064; C.A., 121 (1994) 248156m.

- 259 Wolff, R.L.: *Cis-trans* isomerization of octadecatrienoic acids during heating. Study of pinolenic (*cis*-5,*cis*-9,*cis*-12 18:3) acid geometrical isomers in heated pine seed oil. *J. Am. Oil Chem. Soc.*, 71 (1994) 1129-1134.

- 260 Zhou, W.: (Determination of water-soluble phenolic acid in Danshen oral liquid by TLC-densitometry). *Zhongcaoyao*, 25 (1994) 460-461; C.A., 122 (1995) 17291f.

- 261 Zou, Z., Wang, H., Zhang, A. and Qiao, X.: (Stable new fluorescent reagent CGE(N) used to determine *p*-aminosalicylic acid by TLC with fluorimetric detection). *Huaxi Yaoxue Zazhi*, 8 (1993) 70-73; C.A., 121 (1994) 263812r.

11b. *Prostaglandins*

- 262 Bruno, P., Caselli, M., Mangone, A., Traini, A. and Trisolini, C.: OPLC separation and quantitative determination of fluorescent derivatives of prostaglandins at picogram level. *J. Planar Chromatogr.*, 7 (1994) 362-367.

11c. *Lipids and their constituents*

- 263 Ariga, T., Kasai, N., Miyoshi, I., Yamawaki, M., Scarsdale, J.N., Yu, R.K., Kasama, T. and Taki, T.: Accumulation of isogloboside and ganglio-N-tetraosyl ceramide having blood group B determinant in the hepatomas of female LEC rats. *Biochim. Biophys. Acta*, 1254 (1995) 257-266.

- 264 Bonte, F., Pinguet, P., Chevalier, J.M. and Meybeck, A.: Analysis of all stratum corneum lipids by automated multiple development high-performance thin-layer chromatography. *J. Chromatogr. B*, 664 (1995) 311-316.
  - 265 Boumendjel, A. and Miller, P.F.: Synthesis of sphingosine-1-phosphate and dihydrosphingosine-1-phosphate. *J. Lipid Res.*, 35 (1994) 2305-2311.
  - 266 Brailoiu, E., Saila, L., Huhurez, G., Costuleanu, M., Filipeanu, C.M., Slatineanu, S., Cotrutzu, C. and Branisteanu, D.D.: TLC-a rapid method for liposome characterization. *Biomed. Chromatogr.*, 8 (1994) 193-195; C.A., 121 (1994) 199949m.
  - 267 Damiani, P., Santinelli, F., Simonetti, M.S., Castellini, M. and Rosi, M.: Comparison between two procedures for stereospecific analysis of thiacylglycerols from vegetable oils - I: olive oil. *J. Am. Oil Chem. Soc.*, 71 (1994) 1157-1162.
  - 268 Engelmann, B., Bräutigam, C. and Thieri, J.: Plasmalogen phospholipids as potential protectors against lipid peroxidation of low density lipoproteins. *Biochem. Biophys. Res. Commun.*, 204 (1994) 1235-1242.
  - 269 Field, F.J., Born, E., Chen, H., Murthy, S. and Mathur, S.N.: Lysophosphatidylcholine increases the secretion of cholesteryl ester-poor triacylglycerol-rich lipoproteins by CaCo-2 cells. *Biochem. J.*, 304 (1994) 35-42.
  - 270 Freeman, E.J. and Tallant, E.A.: Vascular smooth-muscle cells contain AT<sub>1</sub> angiotensin receptors coupled to phospholipase D activation. *Biochem. J.*, 304 (1994) 543-548.
  - 271 Furuya, S., Irie, F., Hashikawa, T., Nakazawa, K., Kozakai, A., Hasegawa, A., Sudo, K. and Hirabayashi, Y.: Ganglioside G<sub>D1a</sub> in cerebellar Purkinje cells. Its specific absence in mouse mutants with Purkinje cell abnormality and altered immunoreactivity in response to conjunctive stimuli causing long term desensitization. *J. Biol. Chem.*, 269 (1994) 32418-32425.
  - 272 Hartmann, S., Mallet, A.L., Minnikin, D.E. and Ridell, M.: Thin-layer chromatography and mass spectrometry of pentafluorobenzylidene derivatives of members of the phthiocerol family from *Mycobacterium tuberculosis* and *Mycobacterium marinum*. *Biochim. Biophys. Acta*, 1201 (1994) 129-134.
  - 273 Huang, K. and Akoh, C.C.: Lipase-catalyzed incorporation of n-3 polyunsaturated fatty acids into vegetable oils. *J. Am. Oil Chem. Soc.*, 71 (1994) 1277-1280.
  - 274 Kagehara, M., Tachi, M., Harii, K. and Iwamori, M.: Programmed expression of cholesterol sulfotransferase and transglutaminase during epidermal differentiation of murine skin development. *Biochim. Biophys. Acta*, 1215 (1994) 183-189.
  - 275 Lanne, B., Schierbeck, B. and Karlsson, K.-A.: On the role of the carboxyl group of sialic acid in binding of cholera toxin to the receptor glycosphingolipid, GM1. *J. Biochem. (Tokyo)*, 116 (1994) 1269-1274.
  - 276 Matsuda, K., Kasama, T., Ishizuka, I., Handa, S., Yamamoto, N. and Taki, T.: Structure of a novel phosphocholine-containing glycosphingolipid from *Mycoplasma fermentans*. *J. Biol. Chem.*, 269 (1994) 33123-33128.
  - 277 Mordarska, H. and Pasciak, M.: A simple method for differentiation of *Propionibacterium acnes* and *Propionibacterium propionicum*. *FEMS Microbiol. Lett.*, 123 (1994) 325-330; C.A., 121 (1994) 296310d.
  - 278 Nores, G.A., Mizutamari, R.K. and Kremer, D.M.: Chromatographic tank designed to obtain highly reproducible high-performance thin-layer chromatograms of gangliosides and neutral glycosphingolipids. *J. Chromatogr. A*, 686 (1994) 155-157.
  - 279 Perez, M.K., Fried, B. and Sherma, J.: Comparison of mobile phases and HPTLC qualitative and quantitative analysis, on preadsorbent silica gel plates, of phospholipids in *Biomphalaria glabrata* (gastropoda) infected with *Echinostoma caproni* (trematoda). *J. Planar Chromatogr.*, 7 (1994) 340-343.
  - 280 Rabkin, S.W.: Effect of hypoxia on choline metabolism and phosphatidylcholine biosynthesis in isolated adult rat ventricular myocytes: effects of trifluoperazine. *Biochem. Cell Biol.*, 72 (1994) 289-296.
  - 281 Sallin, C., Sieber, R., Bosset, J.O. and Tabacchi, R.: (Oxidized cholesterol determination in milk and milk products. Part 2. Possibilities and limits of TLC). *Mitt. Geb. Lebensmittelunters. Hyg.*, 84 (1993) 566-580; C.A., 121 (1994) 253984j.
  - 282 Steele, W. and Banks, W.: Triglyceride distribution in hydrogenated milk fat and its effects on separation by thin-layer chromatography. *Milchwissenschaft*, 49 (1994) 372-376; C.A., 121 (1994) 299572b.
  - 283 Sugita, M., Hayata, C., Yoshida, T., Suzuki, M., Suzuki, A., Takeda, T., Hori, T. and Nakatani, F.: A novel fucosylated glycosphingolipid from the millipede, *Parafontaria laminata armigera*. *Biochim. Biophys. Acta*, 1215 (1994) 163-169.
  - 284 Taki, T., Kasama, T., Handa, S. and Ishikawa, D.: A simple and quantitative purification of glycosphingolipids and phospholipids by thin-layer chromatography blotting. *Anal. Biochem.*, 223 (1994) 232-238.
  - 285 Tercé, F., Brun, H. and Vance, D.E.: Requirement of phosphatidylcholine for normal progression through the cell cycle in C3H/10T 1/2 fibroblasts. *J. Lipid Res.*, 35 (1994) 2130-2142.
  - 286 Watarai, S., Kiura, K., Shigeto, R., Shibayama, T., Kimura, I. and Yasuda, T.: Establishment of monoclonal antibodies specific for ganglioside GM1: detection of ganglioside GM1 in small cell lung carcinoma cell lines and tissues. *J. Biochem. (Tokyo)*, 116 (1994) 948-954.
  - 287 Weete, J.D.: Preferential degradation of noncholine phosphatides in soybean lecithin by thermalization. *J. Am. Oil Chem. Soc.*, 71 (1994) 1195-1199.
  - 288 Zlatanov, M.: Untersuchungen über die Phospholipidzusammensetzung der Glyceridöle einiger Vertreter der Familie Apiaceae. *Fat. Sci. Technol.*, 96 (1994) 456-457.
- See also 289, 330.
- 11d. *Lipoproteins and their constituents*
- 289 Nilsson, Å., Chen, Q. and Dahlman, E.: Metabolism of chylomicron phosphatidylinositol in the rat: fate *in vivo* and hydrolysis with lipoprotein lipase and hepatic lipase *in vitro*. *J. Lipid Res.*, 35 (1994) 2151-2160.
13. STEROIDS
- 13a. *General techniques*
- 290 Vanoosthuyze, K., Daeseleire, E., van Overbeke, A., van Peteghem, C. and Ermens, A.: Survey of hormones used in cattle fattening based on the analysis of belgian injection sites. *Analyst (Cambridge)*, 119 (1994) 2655-2658.
- See also 226.

## 13b. Pregnane and androstane derivatives

- 291 Agarwal, A.K., Pampori, N.A. and Shapiro, B.H.: Thin-layer chromatographic separation of regioselective and stereospecific androgen metabolites. *Anal. Biochem.*, 224 (1995) 455-457.
- 292 Auricchio, M.T., Markman, B.E.O. and Batistic, M.A.: (Detection of the presence of corticosteroids in alternative pharmaceutical preparations used for bronchial asthma and bronchitis treatment). *Rev. Inst. Adolfo Lutz*, 53, No. 1-2 (1993) 37-40; *C.A.*, 121 (1994) 18799u.
- 293 Richardson, T.H. and Johnson, E.F.: Alterations of the regioselectivity of progesterone metabolism by the mutagenesis of two key amino acid residues in rabbit cytochrome P450 2C3v. *J. Biol. Chem.*, 269 (1994) 23937-23943.
- 294 Smets, F., Pottier, G., de Brabander, H.F., Batjoens, P., Hendriks, L. and Courtheyn, D.: By-products of steroid synthesis: a cause of interferences in thin-layer chromatography. *Analyst (Cambridge)*, 119 (1994) 2571-2575.

## 13d. Sterols

- 295 Johnson, K.A., Morrow, C.J., Knight, G.D. and Scallen, T.J.: *In vivo* formation of 25-hydroxycholesterol from endogenous cholesterol after a single meal, dietary cholesterol challenge. *J. Lipid Res.*, 35 (1994) 2241-2253.
- 296 Wardas, W. and Pyka, A.: Visualizing agents for cholesterol in TLC. *J. Planar Chromatogr.*, 7 (1994) 440-443.

## 13e. Bile acids and alcohols

- 297 Roda, A., Pellicciari, R., Cerre, C., Polimeni, C., Sadeghpour, B., Marinozzi, M., Forti, G.C. and Sapigni, E.: New 6-substituted bile acids: physico-chemical and biological properties of 6 $\alpha$ -methyl ursodeoxycholic acid and 6 $\alpha$ -methyl-7-epicholic acid. *J. Lipid Res.*, 35 (1994) 2268-2279.

## 14. STEROID GLYCOSIDES AND SAPONINS

- 298 Zhongping, Y., Guoqiao, L., Hanhui, W., Weide, H., Shankai, Z. and Wenmei, H.: Determination of the glycosyl sequence of glycosides by enzymatic hydrolysis followed by TLC-FABMS. *J. Planar Chromatogr.*, 7 (1994) 410-412.

See also 376, 385.

## 15. TERPENES AND OTHER VOLATILE AROMATIC COMPOUNDS

## 15a. Terpenes

- 299 Ando, M., Arai, K., Kikuchi, K. and Isogai, K.: Synthetic studies of sesquiterpenes with a *cis*-fused decalin system. 4. Synthesis of (+)-5 $\beta$ H-eudesma-3,11-diene, (-)-5 $\beta$ H-eudesmane-4 $\beta$ ,11-diol, and (+)-5 $\beta$ H-eudesmane-4 $\alpha$ ,11-diol, and structure revision of a natural eudesmane-4,11-diol isolated from *Pluchea arguta*. *J. Natural Prod.*, 57 (1994) 1189-1199.

## 15b. Essential oils

See 388.

## 16. NITRO AND NITROSO COMPOUNDS

- 300 Miszcak, M. and Bladek, J.: Investigation of double-base propellants' compositions. *Int. Annu. Conf. ICT*, 25th (Energetic Materials-Analysis Characterization and Test Techniques), (1994) 57 (1-57) 10; *C.A.*, 121 (1994) 304145w.
- 301 Zou, H., Zhou, S., Hu, X., Zhang, Y. and Lu, P.: Identification of 2,4,6-trinitrotoluene and its biodegradation products by thin layer chromatography. *J. Planar Chromatogr.*, 7 (1994) 461-463.

See also 246, 363.

## 17. AMINES, AMIDES AND RELATED NITROGEN COMPOUNDS

## 17a. Amines and polyamines

- 302 Bazyłak, G.: Chemometrically aided description of imino-diethanol retention on hydrocarbonaceous silica and cellulose layers. *J. Planar Chromatogr.*, 7 (1994) 428-434.

See also 301.

## 17d. Other amine derivatives and amides (excl. peptides)

See 257, 393.

## 18. AMINO ACIDS AND PEPTIDES; CHEMICAL STRUCTURE OF PROTEINS

## 18a. Amino acids and their derivatives

- 303 Bhushan, R. and Parshad, V.: TLC of amino acids on thin silica gel layers impregnated with transition metal ions and their anions. *J. Planar Chromatogr.*, 7 (1994) 480-484.
- 304 DeSilva, K., Strojek, J. and Kuwana, T.: Dual wavelength excitation method for the off-line liquid chromatographic analysis of derivatized amino acids. *Anal. Sci.*, 10 (1994) 573-578; *C.A.*, 122 (1995) 22830k.
- 305 Golkiewicz, W. and Polak, B.: Relationship between capacity factors and mobile phase composition in thin layer ligand exchange chromatography of the enantiomers of some amino acids. *J. Planar Chromatogr.*, 7 (1994) 453-457.
- 306 Mathur, V., Kanoongo, N., Mathur, R., Narang, C.K. and Mathur, N.K.: Resolution of amino acid racemates on borate-gelled guaran-impregnated silica gel thin-layer chromatographic plates. *J. Chromatogr. A*, 685 (1994) 360-364.
- 307 Morioka, M., Muraoka, H., Yamamoto, K. and Ishikawa, H.: An endosymbiont chaperonin is a novel type of histidine protein kinase. *J. Biochem. (Tokyo)*, 116 (1994) 1075-1081.

- 308 Nakamura, F. and Suyama, K.: Analysis of aldose, an amino acid derived from aldol crosslink of elastin and collagen by high-performance liquid chromatography. *Anal. Biochem.*, 223 (1994) 21-25.
- 309 Roten, C.-A.H., Pagni, M., Margot, P., Touri, F. and Karamata, D.: Specific labeling of diaminopimelate: a radioassay for the determination of the peptidoglycan cross-linking index. *Anal. Biochem.*, 223 (1994) 208-211.
- 310 Sakakibara, Y., Suiko, M. and Liu, M.-C.: *De novo* sulfation of L-tyrosine in HepG2 human hepatoma cells and its possible functional implication. *Eur. J. Biochem.*, 226 (1994) 293-301.
- 311 Vladimirova, N.M., Potapenko, N.A., Sachs, G. and Modyanov, N.N.: Determination of the sidedness of the carboxy-terminus of the Na<sup>+</sup>/K<sup>+</sup>-ATPase  $\alpha$ -subunit using lactoperoxidase iodination. *Biochim. Biophys. Acta*, 1233 (1995) 175-184.
- See also 237, 238, 330.
- 18b. *Peptides, peptidic and proteinous hormones, growth factors*
- 312 Guesdon, F., Waller, R.J. and Saklatvala, J.: Specific activation of  $\beta$ -casein kinase by the inflammatory cytokines interleukin 1 and tumour necrosis factor. *Biochem. J.*, 304 (1994) 761-768.
- 313 Jähnen-Dechent, W., Trindl, A., Godovac-Zimmermann, J. and Müller-Esterl, W.: Posttranslational processing of human  $\alpha_2$ -HS glycoprotein (human fetuin). Evidence for the production of a phosphorylated single-chain form by hepatoma cells. *Eur. J. Biochem.*, 226 (1994) 59-69.
- 314 Suzuki, M., Oda, K., Takatudi, T. and Takitani, S.: (Study on purity test for related compounds in liothyronine sodium and levothyroxine sodium by thin-layer chromatography). *Iyakuhin Kenkyu*, 25 (1994) 205-209; C.A., 121 (1994) 213115v.
- 315 Walaas, S.I., Czernik, A.J., Olstad, O.K., Sletten, K. and Walaas, O.: Protein kinase C and cyclic AMP-dependent protein kinase phosphorylate phospholemman, and insulin and adrenaline-regulated membrane phosphoprotein, at specific sites in the carboxy terminal domain. *Biochem. J.*, 304 (1994) 635-640.
19. PROTEINS
- 19e. *Proteins of blood, serum and blood cells*
- 316 Li, H. and Zhang, R.X.: Measurement of platelet-activating factor in human blood by high performance thin layer chromatography and its clinical application. *J. Tongji Med. Univ.*, 13 (1993) 226-230; C.A., 121 (1994) 199938g.
21. PURINES, PYRIMIDINES, NUCLEIC ACIDS AND THEIR CONSTITUENTS
- 21a. *Purines, pyrimidines, nucleosides, nucleotides*
- 317 Faraone-Mennella, M.R., de Lucia, F., de Maio, A., Gambacorta, A., Quesada, P., de Rosa, M., Nicolaus, B. and Farina, B.: ADP-ribosylation reactions in *Sulfolobus solfataricus*, a thermoacidophilic archaeon. *Biochim. Biophys. Acta*, 1246 (1995) 151-159.
- 318 Salvadori, M.C., Rieser, E.M., Neto, L.M.R. and Nascimento, E.S.: Determination of xanthines by high-performance liquid chromatography and thin-layer chromatography in horse urine after ingestion of guarana powder. *Analyst (Cambridge)*, 119 (1994) 2701-2703.
- See also 335, 350.
- 21c. *Nucleic acids, DNA*
- 319 Park, W.J. and Adler, P.N.: Thin layer chromatography assay for the incorporation of digoxigenin into nucleic acid. *BioTechniques*, 17 (1994) 656-657; C.A., 121 (1994) 275692q.
- 21d. *Structural studies on RNA and RNA mapping*
- 320 Selvamurugan, N., Nag, M.K. and Elieiri, G.L.: Intron-encoded small nucleolar RNAs: new RNA sequence variants and genomic loci. *Biochim. Biophys. Acta*, 1260 (1995) 230-234.
22. ALKALOIDS
- 321 Bah, M., Bye, R. and Pereda-Miranda, R.: Biologically active natural products from Mexican medicinal plants. Part III. Hepatotoxic pyrrolizidine alkaloids in the Mexican medicinal plant *Packera candidissima* (Asteraceae: Senecioneae). *J. Ethnopharmacol.*, 43 (1994) 19-30; C.A., 121 (1994) 212761r.
- 322 Liu, Y., Ye, G. and Qian, W.: (The effect on alkaloids in tincture *Leonuri sibirici* by irradiative sterilization and the study of sterilization effect). *Zhongguo Yaoxue Zazhi*, 29, 8, Suppl. (1994) 12-14; C.A., 121 (1994) 308103d.
- 323 Papadoyannis, I.N.: Instrumental analytical techniques used for datura alkaloid analyses. *Instrum. Sci. Technol.*, 22 (1994) 241-258; C.A., 122 (1994) 17282d.
- 324 Shao, S., Xu, L., Li, Y. and Ge, W.: (Determination of atropine in *gucilasungao* by dual wavelength TLC scanning). *Zhongcaoyao*, 25 (1994) 357-358; C.A., 121 (1994) 263824w.
- 325 Shinde, V.M., Tendolkar, N.M. and Desai, B.S.: Simultaneous determination of theophylline and etophylline in pharmaceutical dosages by HPTLC. *Anal. Lett.*, 28 (1995) 45-58.
- 326 Yavich, P.A., Kakhtelidze, M.B. and Sarabunovich, A.G.: (Quantitative determination of berberine in *Phellodendron lavallei* bark). *Farmatsiya (Moscow)*, 42, No. 3 (1993) 49-50; C.A., 121 (1994) 141848m.
- 327 Zhang, T., Zhang, S. and Wei, L.: (Determination of alkaloids in *Evodia rutaecarpa* (Juss.) Benth. and its processed products by TLC-densitometry). *Zhongguo Zhongyao Zazhi*, 19 (1994) 409-411; C.A., 121 (1994) 213094n.
- See also 236, 402.

## 23. OTHER SUBSTANCES CONTAINING HETEROCYCLIC NITROGEN

## 23a. Porphyrins and other pyrroles

- 328 Lai, C.-K., Lam, C.-W. and Chan, Y.-W.: High-performance thin-layer chromatography of free porphyrins for diagnosis of porphyria. *Clin. Chem. (Washington)*, 40 (1994) 2026-2029.

## 23b. Bile pigments

See 243.

## 23d. Pyridine derivatives

- 329 Baranowska, I. and Swierczek, S.: A study of the retention of azines and diazines in RPTLC with bonded alkyl stationary phase. *J. Planar Chromatogr.*, 7 (1994) 399-405.

## 23e. Other N-heterocyclic compounds

- 330 Babizhayev, M.A., Seguin, M.-C., Gueyne, J., Evstigneeva, R.P., Ageyeva, E.A. and Zheltukhina, G.A.: L-Carnosine ( $\beta$ -alanyl-L-histidine) and carcinine ( $\beta$ -alanylhistamine) act as natural antioxidants with hydroxyl-radical-scavenging and lipid-peroxidase activities. *Biochem. J.*, 304 (1994) 509-516.
- 331 Shirke, P., Patel, M.K., Tirodkar, V.B., Tamhane, V. and Sethi, P.D.: Simultaneous determination of furazolidone and metronidazole in their combined dosage form by HPTLC. *Indian J. Pharm. Sci.*, 56 (1994) 108-109; C.A., 121 (1994) 308494g.
- 332 Wang, L. and Zhu, Q.: (Qualitative analysis of cuscohygrine and 1-methyl-2-pyrrolidone by TLC). *Huaxi Yaixue Zazhi*, 9 (1994) 119; C.A., 121 (1994) 187414v.

See also 229, 329, 366, 380.

## 24. ORGANIC SULPHUR COMPOUNDS (INCL. GLUCOSINOLATES)

- 333 Feng, Y., Zhang, S., Zhang, Y. and Shen, X.: (Simultaneous determination of three components of trimethoprim-sulfadiazine-sulfamethoxazole tablets by thin-layer chromatographic scanning). *Sepu*, 12 (1994) 364-366; C.A., 121 (1994) 208505m.
- 334 Maslankiewicz, M.J. and Pluta, K.: TLC separation of  $\beta$ - and  $\gamma$ -quinoliny sulfides. *J. Planar Chromatogr.*, 7 (1994) 484-486.

See also 329.

## 25. ORGANIC PHOSPHORUS COMPOUNDS (INCL. SUGAR PHOSPHATES)

- 335 Guranowski, A., Starzynska, E., Sillero, G. and Sillero, A.: Conversion of adenosine(5') oligophospho(5') adenosines into inosine(5') oligophospho(5') inosines by non-specific adenylate deaminase from the snail *Helix pomatia*. *Biochim. Biophys. Acta*, 1243 (1995) 78-84.

- 336 Tusarova, I., Halamek, E. and Kobliha, Z.: Thin-layer chromatography of a new group of toxic dialkylaminoethyl dialkyl-amido fluorophosphates. *J. Planar Chromatogr.*, 7 (1994) 372-375.

See also 279, 312, 315.

## 27. VITAMINS AND VARIOUS ANIMAL GROWTH FACTORS (NON-PEPTIDIC)

- 337 Brushan, R. and Parshad, V.: Separation of vitamin B complex and folic acid using TLC plates impregnated with some transition metal ions. *Biomed. Chromatogr.*, 8 (1994) 194-198; C.A., 121 (1994) 199950e.

See also 236.

## 28. ANTIBIOTICS

- 338 Funabashi, Y., Horiguchi, T., Iinuma, S., Tanida, S. and Harada, S.: TAN-1496 A, C and E, diketopiperazine antibiotics with inhibitory activity against mammalian DNA topoisomerase I. *J. Antibiot.*, 47 (1994) 1202-1218.
- 339 Ichihara, J., Shinohara, Y., Kogure, K. and Terada, H.: Nucleotide sequence of the 5'-flanking region of the rat type II hexokinase gene. *Biochim. Biophys. Acta*, 1260 (1995) 365-368.
- 340 Iwamoto, T., Fujie, A., Sakamoto, K., Tsurumi, Y., Shigematsu, N., Yamashita, S., Hashimoto, S., Okuhara, M. and Kohsaka, M.: WF 11899A, B and C, novel antifungal lipopeptides I. Taxonomy, fermentation, isolation and physico-chemical properties. *J. Antibiot.*, 47 (1994) 1084-1091.
- 341 Jindal, K.C., Chaudhary, R.S., Gangwal, S.S., Singla, A.K. and Khanna, S.: High-performance thin-layer chromatographic method for monitoring degradation products of rifampicin in drug excipient interaction studies. *J. Chromatogr. A*, 685 (1994) 195-199.
- 342 Khan, K., Paesen, J., Roets, E. and Hoogmartens, J.: Quantitative TLC of erythromycin - application to commercial bulk samples and biological samples. *J. Planar Chromatogr.*, 7 (1994) 349-353.
- 343 Medina, M.B. and Unruh, J.J.: Solid-phase clean-up and thin-layer chromatographic detection of veterinary aminoglycosides. *J. Chromatogr. B*, 663 (1995) 127-135.
- 344 Morino, T., Masuda, A., Yamada, M., Nishimoto, M., Nishikiori, T., Saito, S. and Shimada, N.: Stevastelins, novel immunosuppressants produced by *Penicillium*. *J. Antibiot.*, 47 (1994) 1341-1343.
- 345 Tammilehto, S., Salomies, H. and Torniainen, K.: Qualitative and quantitative TLC for monitoring ciprofloxacin photodegradation in aqueous solutions. *J. Planar Chromatogr.*, 7 (1994) 368-371.
- 346 Wang, W., Lu, M. and Shen, L.: (Bioinactive impurities in ribostamycin). *Zhongguo Kangshengsu Zazhi*, 19 (1994) 219-220; C.A., 121 (1994) 238259h.

## 29. INSECTICIDES, PESTICIDES AND OTHER AGROCHEMICALS

## 29a. General techniques

- 347 Butz, S. and Stan, H.-J.: Screening of 265 pesticides in water by thin-layer chromatography with automated multiple development. *Anal. Chem.*, 67 (1995) 620-630.

See also 233.

## 29d. Carbamates

- 348 Bose, D., Shivhare, P. and Gupta, V.K.: Thin layer chromatographic separation and determination of carbaryl and propxur. *J. Planar Chromatogr.*, 7 (1994) 415-418.

## 29e. Herbicides

- 349 Cserháti, T. and Forgács, E.: Retention of 4-cyanophenyl herbicides on water-insoluble  $\beta$ -cyclodextrin support. *J. Chromatogr. A*, 685 (1994) 295-302.
- 350 Tekel', J., Vaverkova, S., Kovacicova, J., Holla, M. and Havranek, E.: A screening method for the determination of the residues of racil herbicides in *Melissa officinalis* L. *Pharmazie*, 49 (1994) 899-901.

## 29g. Other types of pesticides and various agrochemicals

- 351 Sabato, R., Pierdominici, E., Achene, L. and Brambilla, G.: Presence of eleiophyllin in "premixes" and feeds containing salinomycin by TLC and HPLC analysis. *Microbiol. Aliments. Nutr.*, 11 (1993) 429-432; *C.A.*, 121 (1994) 106782p.

See also 247, 348.

## 30. SYNTHETIC AND NATURAL DYES

## 30a. Synthetic dyes

- 352 Lenze, U. and Kallmayer, H.-J.: 4-[(4-Hydroxy-1-naphthyl)phenylmethylene]-1(4H)-naphthalinon - die Indikatorsubstanz des " $\alpha$ -Naphtholbenziens". *Pharmazie*, 50 (1995) 36-38.
- 353 Mariani, E., Bargagna, A., Longobardi, M. and Dorato, S.: Application of TLC and HPTLC in the analysis of semipermanent hair dyes. *Int. J. Cosmet. Sci.*, 16, No. 1 (1994) 17-27; *C.A.*, 121 (1994) 186750q.
- 354 Upadhyay, R.K., Sharma, R.K., Babu, G. and Mishra, G.: Chromatographic separation of thioindigoid thiazolidinones: Part I. *J. Planar Chromatogr.*, 7 (1994) 464-468.

See also 220.

## 30b. Chloroplast and other natural pigments

- 355 Mínguez-Mosquera, M.I. and Gandul-Rojas, B.: High-performance liquid chromatographic study of alkaline treatment of chlorophyll. *J. Chromatogr. A*, 690 (1995) 161-176.

## 31. PLASTICS AND THEIR INTERMEDIATES

- 356 Shi, Y., Zhou, H. and Wang, D.: (Study on the main component in the RTP [rosin diglycol plasticizer] plasticizer). *Linchan Huaxue Yu Gongye*, 12 (1992) 293-297; *C.A.*, 121 (1994) 281836f.

## 32. DRUG ANALYSIS

## 32a. Drug analysis, general techniques

- 357 Potuzak, M., Turan, J. and Matejek, S.: (Use of thin-layer chromatography for pharmaceutical analysis. III. Computing technology, radiochromatography). *Ceska Slov. Farm.*, 43 (1994) 245-251; *C.A.*, 122 (1995) 17286h.
- 358 Potuzak, M., Turan, J. and Matejek, S.: (Utilization of thin-layer chromatography in pharmaceutical control. II. The mobile phase, development of the chromatogram, detection). *Ceska Slov. Farm.*, 43 (1994) 171-178; *C.A.*, 121 (1994) 286686s.
- 359 Uneda, K., Ninomiya, S., Esumi, Y., Shimada, N. and Nagatsuka, S.: (Application of bioimage analyzer on drug metabolism studies using thin layer chromatography (II). Direct development of plasma samples containing [ $^{14}$ C]indomethacin and its metabolites). *Yakubutsu Dotai*, 8 (1993) 1273-1282; *C.A.*, 121 (1994) 98931u.

See also 213.

## 32b. Antirheumatics and antiinflammatory drugs

- 360 Sava, S., Curea, E. and Bojita, M.: (Piroxicam identity and purity using a thin layer chromatographic method). *Farmacia (Bucharrest)*, 41, No. 3-4 (1993) 23-26; *C.A.*, 121 (1994) 308469c.

See also 238.

## 32c. Autonomic and cardiovascular drugs

- 361 Alton, K.B., Chan, T.-M. and Pramanik, B.N.: Urinary metabolites of (R),(R)-labetalol. *Drug Metab. Disp.*, 22 (1994) 866-872.
- 362 Chandrashekhar, T.G., Rao, P.S.N., Smrita, K., Vyas, S.K. and Dutt, C.: Analysis of amlodipine besylate by HPTLC with fluorimetric detection: a sensitive method for assay of tablets. *J. Planar Chromatogr.*, 7 (1994) 458-460.
- 363 Marciniak, B. and Rychcik, W.: Kinetic analysis of nifedipine photodegradation in the solid state. *Pharmazie*, 49 (1994) 894-897.

See also 365.

## 32d. Central nervous system drugs

- 364 Albert, K.: Identification test for methadone hydrochloride. *Pharm. Ztg.*, 139, No. 14 (1994) 42; *C.A.*, 121 (1994) 238171y.

- 365 Han, L., Li, X. and Liu, Y.: (Detection method for the related substances of encainide HCl). *Zhongguo Yaoxue Zazhi*, 29 (1994) 486-488; C.A., 121 (1994) 308473z.

See also 229, 236.

### 32e. Chemotherapeutics (exc. cytostatics and antibiotics)

- 366 Chervenkov, S.K., Pavlov, A.I. and Slavova, S.T.: IR-, <sup>1</sup>H-NMR and TLC examination of new styrylquinolines with antibacterial activity. *Anal. Lett.*, 28 (1995) 59-70.
- 367 Shearan, P. and O'Keeffe, M.: Novel approach the "on-site" testing for sulfamethazine in pork carcasses. *Analyst (Cambridge)*, 119 (1994) 2761-2763.
- 368 Spell, J.C. and Stewart, J.T.: Quantitative analysis of para-aminosalicylic acid and its major impurity meta-aminophenol in para-aminosalicylic acid drug substance by HPTLC and scanning densitometry. *J. Planar Chromatogr.*, 7 (1994) 472-473.

See also 247, 333, 393.

### 32f. Cytostatics

See 223.

### 32g. Other drug categories

- 369 Farina, A., Doldo, A., Cotichini, V., Gallo, F.R. and Calandra, S.: Assay and purity control of ticlopidine by thin layer chromatography. *J. Planar Chromatogr.*, 7 (1994) 386-388.
- 370 Ficarra, R., Ficarra, P., Tommasini, S., Carulli, M., Costantino, D. and Calabro, M.L.: Chromatographic investigations on brotizolam. *Farmaco*, 49 (1994) 437-440; C.A., 121 (1994) 308503j.
- 371 Liu, J., Tu, X., Chen, Y. and Wu, D.: (Films for the treatment of cough and asthma). *Zhongguo Yaoxue Zazhi*, 25 (1994) 218-220; C.A., 121 (1994) 238307x.
- 372 Mara, K.M. and Omar, F.A.: Spectrophotometric determination of piroxicam using 2,2-diphenyl-1-picrylhydrazyl. *Alexandria J. Pharm. Sci.*, 8, No. 1 (1994) 24-28; C.A., 121 (1994) 213164k.
- 373 Wahbi, Abdel-Aziz, M., El-Walily, A., Bedair, M., El-Gendy, A.: Determination of cinnarizine in the presence of its degradation products. *Egypt. J. Pharm. Sci.*, 34, No. 1-3 (1993) 35-46; C.A., 121 (1994) 308477d.

See also 332, 344, 364, 377.

### 32h. Toxicological and forensic applications

- 374 De Zeeuw, R.A., Franke, J.P., van Halem, M., Schaapman, S., Logawa, E. and Siregar, C.J.P.: Impact of hot and humid tropical conditions of Toxi-Lab® thin-layer chromatographic systems for the screening of basic and neutral drugs. *J. Anal. Toxicol.*, 18 (1994) 402-406.

See also 239, 350.

### 32i. Plant extracts

- 375 Abdel Sattar, A., Bankova, V., Kujumgiev, A., Galabov, A., Agnato, A., Todorova, C. and Popov, S.: Chemical composition and biological activity of leaf exudates from some Lamiaceae plants. *Pharmazie*, 50 (1995) 62-65.
- 376 Barthomeuf, C., Chmielowiec, C. and Pourrat, H.: TLC - densitometry for quality control of *Hedera helix* L. (Araliaceae). *J. Planar Chromatogr.*, 7 (1994) 474-476.
- 377 Cai, M.: (Quantitative determination of arteannuin in liposomes and dextrans by TLC-scanning method). *Zhongcaoyao*, 25 (1994) 353-376; C.A., 121 (1994) 263804q.
- 378 Demirezer, L.Ö.: Quantitative determination of some Rumex species with regards of anthraquinone derivatives. *Pharmazie*, 49 (1994) 936-937.
- 379 Gao, B. and Wei, S.: (Determination method of jujuboside A in Suanzoren tea). *Hebeisheng Kexueyan Xuebao*, No. 4 (1992) 61-64; C.A., 122 (1995) 17289m.
- 380 Gorinsky, V.: Bisbenzylisoquinoline derivatives from *Ocotea rodiei* for pharmaceutical use. *Eur. Pat. Appl. EP 610,060* (Cl. C07D491/18), 10 Aug. 1994, GB App. 93/1,921, 01 Feb. 1993; 9 pp.; C.A., 121 (1994) 212962g.
- 381 Pal, S.K., Mukherjee, P.K. and Saha, B.P.: Co-chemical and thin-layer chromatographic characterization of the prepared tincture of *Moringa oleifera* Lam. (fac. *Moringaceae*). *Res. Ind.*, 39 (1994) 191-193; C.A., 122 (1995) 17296m.
- 382 Pietta, P.G., Mauri, P.L., Bruna, A. and Merfort, I.: MEKC as an improved method to detect falsification in the flowers of *Arnica montana* and *A. chamissonis*. *Planta Med.*, 60 (1994) 369-372; C.A., 121 (1994) 238499m.
- 383 Su, J., Wang, B. and Wang, G.: (Analysis of Guiqi mixture and its main ingredient *Astragalus* root). *Zhongguo Zhongyao Zazhi*, 19 (1994) 158-160; C.A., 121 (1994) 42864g.
- 384 Wu, S., Jiang, Y. and Wang, L.: (Preparation and criterion of Qiangliyoujizhe oral liquid). *Huaxi Yaoxue Zazhi*, 9 (1994) 91-93; C.A., 121 (1994) 187006p.
- 385 Xu, R., Han, G., Feng, L., Dai, J. and Wang, Y.: (Determination of lipophilic compositions in Taidangshen and cultivated Ludangshen of different ages). *Zhongguo Yaoxue Zazhi*, 29 (1994) 358-360; C.A., 121 (1994) 238495g.
- 386 Yang, G.: (Quality standard of Shenrongshanshen medicinal wine). *Zhongcaoyao*, 25 (1994) 458-458, 467; C.A., 121 (1994) 286700s.
- 387 Yang, H. and Yang, H.: (Comparative study with thin-layer chromatography on water and ethanol as extraction solvents for components of *Puerariae radix*). *Huaxi Yaoxue Zazhi*, 9, No. 1 (1994) 35-38; C.A., 121 (1994) 187392m.
- 388 Yoshizaki, F., Sakamoto, R., Ando, T. and Hisamichi, S.: (Simplified method to identify the crude drugs blended in Orengedokuto and Keishininjinto). *Annu. Rep. Tohoku Coll. Pharm.*, 40 (1993) 201-204; C.A., 121 (1994) 308442p.
- 389 Zhen, H., Chen, Y. and Zhong, X.: (Quality standards of Huanglianjiadu pills). *Zhongguo Zhongyao Zazhi*, 19 (1994) 414-417; C.A., 121 (1994) 213095p.

See also 241, 253, 260, 321, 322, 326, 327.

## 33. CLINICO-CHEMICAL APPLICATIONS

33b. *Complex mixtures and profiling (single compounds by cross-reference only)*

See 316, 328.

## 34. FOOD ANALYSIS

34b. *Complex mixtures (single compounds by cross-reference only)*

See 254, 281, 282, 348, 393.

## 35. ENVIRONMENTAL ANALYSIS

35c. *Water pollution (complex mixtures; single compounds by cross-reference only)*

390 Vorontsov, A.M.: (Continuous-flow thin-layer chromatography (FTLC) in analysis of ecotoxicants: method characteristics and result evaluation criteria). *Zh. Ekol. Khim.*, 1 (1993) 33-34; C.A., 121 (1994) 237940m.

See also 301, 347, 348.

35d. *Soil pollution (complex mixtures; single compounds by cross-reference only)*

See 240.

## 36. SOME TECHNICAL PRODUCTS AND COMPLEX MIXTURES

36a. *Surfactants*

391 Bratt, C.E. and Akerlund, H.E.: Quantification of *n*-octyl  $\beta$ -D-glucopyranoside and *n*-dodecyl  $\beta$ -D-maltoside in biological samples by thin layer chromatography. *Plant Physiol. Biochem. (Paris)*, 32 (1994) 313-315; C.A., 121 (1994) 199910s.

36b. *Antioxidants and preservatives*

392 Ashida, S., Noguchi, S.F. and Suzuki, T.: Antioxidative components, xanthone derivatives, in *Swertia japonica*, Makino. *J. Am. Oil Chem. Soc.*, 71 (1994) 1095-1099.

393 Martin, E., Duret, M. and Vogel, J.: (Determination of sulfonamide residues in eggs). *Mitt. Geb. Lebensmittelunters. Hyg.*, 84 (1993) 274-280; C.A., 121 (1994) 203609v.

394 Tian, L.L. and White, P.J.: Antioxidant activity of oat extract in soybean and cottonseed oils. *J. Am. Oil Chem. Soc.*, 71 (1994) 1079-1086.

See also 330.

36c. *Complex mixtures, technical products and unidentified compounds*

395 Cheng, X. and Ma, H.: (Application of thin-layer chromatography to the synthesis of organometallic compounds). *Sepu*, 12 (1994) 222-223; C.A., 121 (1994) 230961k.

See also 234, 336.

## 37. CELLS, CELLULAR PARTICLES AND SUPRAMOLECULAR STRUCTURES

See 266, 277.

## 38. INORGANIC COMPOUNDS

38a. *Cations*

396 Leont'eva, L.B., Tselinskii, I.V., Boichinova, E.S., Manevskaya, R.S. and Gritsai, G.I.: (Determination of trace amount of Cd, Cu(II), and Pb by paper chromatography). *Zavod. Lab.*, 60 (1994) 7-8, 62-63; C.A., 121 (1994) 270682b.

397 Mohammad, A., Ajal, M., Fatima, N. and Yousuf, R.: Chromatography of cations and anions on mixed sorbent layers developed with aqueous methanol containing tributyl phosphate and formic acid. *J. Planar Chromatogr.*, 7 (1994) 444-449.

398 Murinov, Yu.I., Starikova, L.I. and Ilyushina, N.N.: (Paper chromatographic study of aminoamide as an extraction reagent [for rare earth metals]). *Organ. Reagenty v Anal. Khimii, Perm. Gos. Un-t, Perm*, (1991) 88-92; C.A., 121 (1994) 194441e.

399 Shimizu, T., Jindo, S., Satoh, M. and Mura, Y.: TLC of inorganic ions on trimethylhydroxypropylamine cellulose eluted with sulfuric acid - organic solvent mobile phases. *J. Planar Chromatogr.*, 7 (1994) 412-415.

400 Takeda, Y., Nagai, T. and Ishida, K.: Effects of amino-pretreatment of silica gel on the TLC behaviour of rare earths. *Fresenius J. Anal. Chem.*, 351 (1995) 186-189.

See also 216, 390.

38b. *Anions*

See 397.

## 39. RADIOACTIVE AND OTHER ISOTOPE COMPOUNDS

401 Gattavecchia, E., Tonelli, D., Breccia, A., Fini, A. and Ferri, E.: Radio TLC in quality control and development of radiopharmaceuticals. *J. Radioanal. Nucl. Chem.*, 181 (1994) 77-84; C.A., 121 (1994) 238517r.

402 Huang, X.Y., Wang, Y. and De Lu, Z.: Isolation and preparation of  $^3\text{H}$ -tetrandrine. *J. Labelled Compd. Radiopharm.*, 34 (1994) 899-903; C.A., 121 (1994) 308101b.

- 403 Yoshino, T., Miyoshi, H., Kishi, T., Satoou, K. and Matumoto, T.: (Adsorption of  $^{99m}\text{Tc}$ -labeled HMDP and MDP on hydroxyapatite). *Raidoisotopes*, 43 (1994) 609-612; *C.A.*, 121 (1994) 308267k.

See also 258, 309, 319, 357, 359.

## Electrophoresis

### 1. REVIEWS AND BOOKS

- 1023 Dolnik, V.: Capillary gel electrophoresis. *J. Microcolumn Sep.*, 6 (1994) 315-330; C.A., 121 (1994) 225286q - a review with 178 refs.
- 1024 Landers, J.P. (Editor): *Handbook of Capillary Electrophoresis*. CRC, Boca Raton, 1994, 649 pp.; C.A., 121 (1994) 225829u.
- 1025 Lebo, V.M.M. and Ribeiro, A.C.F.: The electrophoretic effect of electrolyte solutions. *Port. Electrochim. Acta*, 11 (1993) 297-304; C.A., 121 (1994) 286973h - a lecture with 13 refs.
- 1026 Marina, M.L. and Torre, M.: Capillary electrophoresis. *Talanta*, 41 (1994) 1411-1433; C.A., 121 (1994) 225317a - a review with 221 refs.
- 1027 Michaelsen, S. and Soerensen, H.: High-performance capillary electrophoresis: Principles, present possibilities and future potentialities in studies of low and high molecular weight charged and uncharged biomolecules. *Pol. J. Food Nutr. Sci.*, 3 (1994) 5-44; C.A., 121 (1994) 275570y - a review with 114 refs.
- 1028 Silverman, C. and Shaw, C.: Capillary electrophoresis in the evaluation of pharmaceuticals. In: Landers, J.P. (Editor), *Handb. Capillary Electrophor.*, CRC, Boca Raton, 1994, pp. 233-254; C.A., 121 (1994) 238488g - a review with 86 refs.
- 1029 Sutherland, J.C.: Electronic imaging of electrophoretic gels and blots. *Adv. Electrophor.*, 6 (1993) 1-42; C.A., 121 (1994) 220607k - a review with 134 refs.
- 1030 Szepes, L.: (Development and importance of separation techniques in instrumental analysis). *Magy. Kem. Foly.*, 100 (1994) 340-353; C.A., 121 (1994) 270751y - a review with 121 refs.
- 1031 Tsuda, T.: Control of electroosmotic flow in capillary electrophoresis. In: Landers, J.P. (Editor), *Handb. Capillary Electrophor.*, CRC, Boca Raton, 1994, pp. 563-590; C.A., 121 (1994) 213714q - a review with 59 refs.
- 1032 Watanabe, Y. and Yamada, N.: (Electrophoresis. 7. Agarose gel electrophoresis and PAG disc electrophoresis). *Lipid*, 4 (1993) 457-460; C.A., 121 (1994) 225309z - a review with 6 refs.

See also 1061, 1086, 1090, 1095, 1097, 1110, 1115, 1119, 1121, 1124, 1126, 1130, 1132, 1151, 1167, 1232, 1244, 1270, 1281, 1369, 1370, 1490, 1643, 1659, 1680, 1714, 1773, 1776, 1777, 1785, 1788, 1799.

### 2. FUNDAMENTALS, THEORY AND GENERAL

#### 2a. General

- 1033 Bello, M.S., Zhukov, M.Yu. and Righetti, P.G.: Combined effects of non-linear electrophoresis and non-linear chromatography on concentration profiles in capillary electrophoresis. *J. Chromatogr. A*, 693 (1995) 113-130.
- 1034 Hirokawa, T.: Bidirectional isotachopheresis. II. Fifteen electrolyte systems covering the pH range 3.5-10. *J. Chromatogr. A*, 686 (1994) 158-163.

- 1035 Keh, H.J. and Huang, T.Y.: Diffusiophoresis and electrophoresis of elliptical cylindrical particles. *Colloid Polym. Sci.*, 272 (1994) 855-871; C.A., 121 (1994) 264568c.

- 1036 Zimm, B.H.: Diffusion and electrophoresis of a polyelectrolyte in an irregular matrix: II. *Polym. Prepr. (Am. Chem. Soc., Div. Polym. Chem.)*, 35 (1994) 173-174; C.A., 122 (1995) 11093w.

See also 1030, 1085, 1682.

#### 2b. Thermodynamics and theoretical relationships

- 1037 Ermakov, S.V., Zhukov, M.Y., Capelli, L. and Righetti, P.G.: Experimental and theoretical study of artifactual peak splitting in capillary electrophoresis. *Anal. Chem.*, 66 (1994) 4034-4042.
- 1038 Han, R.J., Moss, O.R. and Wong, B.A.: Airborne fiber separation by electrophoresis and dielectrophoresis: theory desing considerations. *Aerosol Sci. Technol.*, 21 (1994) 241-258; C.A., 121 (1994) 307113v.
- 1039 Harmon, B.J., Leesong, I. and Regnier, F.E.: Selectivity in electrophoretically mediated microanalysis by control of product detection time. *Anal. Chem.*, 66 (1994) 3797-3805.
- 1040 Slater, G.W. and Guo, H.L.: Ogston gel electrophoretic sieving: How is the fractional volume available to a particle related to its mobility and diffusion coefficient(s)? *Electrophoresis (Weinheim)*, 16 (1995) 11-15.
- 1041 Yang, S. and Khaledi, M.G.: Linear solvation energy relationships in micellar liquid chromatography and micellar electrokinetic capillary chromatography. *J. Chromatogr. A*, 692 (1995) 301-310.

See also 1086, 1146, 1767.

#### 2d. Measurement of physico-chemical and related values

- 1042 Alarcon-Waess, O. and Velasco, R.M.: Electrokinetic properties of a charged tracer in a time-dependent external field. *J. Chem. Phys.*, 101 (1994) 5110-5115; C.A., 122 (1995) 39708z.
- 1043 Fang, X.H., Zhu, T., Qiang, D.M. and Sun, Y.L.: Monitoring electrolytic reaction of nuclear fast red by high performance capillary electrophoresis. *Chin. Chem. Lett.*, 5 (1994) 597-600; C.A., 121 (1994) 215825a.
- 1044 Herbert, B.J. and Dorsey, J.G.: *n*-Octanol-water partition coefficient estimation by micellar electrokinetic capillary chromatography. *Anal. Chem.*, 67 (1995) 744-749.
- 1045 Krabi, A. and Donath, E.: Polymer depletion layers as measured by electrophoresis. *Colloids Sur., A*, 92 (1994) 175-182; C.A., 122 (1995) 38644p.
- 1046 Radeva, T., Petkanchin, I. and Varoqui, R.: Electric surface properties of oxide particles in presence of neutral polymer. In: Jennings, B.R. and Stoylov, S.P. (Editors), *Colloid Mol. Electro-Opt. Proc. Int. Electro-Opt. Symp.*, 6th 1991, Inst. Phys., Bristol, 1992, pp. 111-114; C.A., 121 (1994) 268632s.

- 1047 Yadav, A. and Yadav, N.R.: Evaluation of stability constants-Cu/Co complexes (simple and mixed) with AMP/NTA using paper electrophoresis. *J. Electrochem. Soc. India*, 42 (1993) 183-186; C.A., 122 (1995) 39983k.

See also 1107, 1174, 1252, 1507, 1766.

### 3. GENERAL TECHNIQUES

#### 3a. Apparatus and accessories

- 1048 Dadoo, R., Zare, R.N. and Colon, L.A.: System for electrokinetic separation and detection where detection is performed at other than separation electric field. U.S. US 5,342,492 (Cl. 204-108.1; G01N27/26), 30 Aug. 1994, Appl. 115,466, 01 Sep. 1993; 8 p.; C.A., 121 (1994) 244473q.
- 1049 Goodale, D.L. and Mack, S.D.: Sample handling system. U.S. US 5,356,525 (Cl. 204-299R; G01N27/26), 18 Oct. 1994, US Appl. 48,708, 16 Apr. 1993; 24 p.; C.A., 122 (1995) 71116r.
- 1050 Hanai, T. and Tsuruta, H.: 2nL Injector for capillary electrophoresis. *Instrum. Sci. Technol.*, 22 (1994) 151-155; C.A., 121 (1994) 220635t.
- 1051 Jacobson, S.C., Hergenröder, R., Moore, A.W., Jr. and Ramsey, J.M.: Precolumn reactions with electrophoretic analysis integrated on a microchip. *Anal. Chem.*, 66 (1994) 4127-4132.
- 1052 Kylo, J.H., Kirby-Keyser, L., Donohoue, P.A. and Porter, C.C.: A new way to seal glass plates for denaturing gradient gel electrophoresis. *BioTechniques*, 16 (1994) 792; C.A., 121 (1994) 275863w.
- 1053 Madden, D. and Schollar, J.: Disposable gel electrophoresis apparatus. Ger. Offen. DE 4,408,129 (Cl. B01D57/02), 15 Sep. 1994, GB Appl. 93/5,052, 12 Mar. 1993; 6 pp.; C.A., 121 (1994) 225846x.
- 1054 Nakamura, N.: Electrophoretic display device using fullerene. *Jpn. Kokai Tokkyo Koho* JP 06,148,693 (94,148,693) (Cl. G02F1/167), 27 May 1994, Appl. 92/297,379, 06 Nov. 1992; 6 p.; C.A., 122 (1995) 42871c.
- 1055 Nakayama, K., Yoshida, M., Taki, M. and Watanabe, Y.: Capillary gel electrophoretic apparatus. *Jpn. Kokai Tokkyo Koho* JP 06,118,058 [94,118,058] (Cl. G01N27/447), 28 Apr. 1994, Appl. 92/264,586, 02 Oct. 1992; 4 p.; C.A., 121 (1994) 239171k.
- 1056 Nilsson, K. and Sellergren, B.: Analysis method. *PCT Int. Appl.* WO 94 16,319 (Cl. G01N27/447), 21 Jul. 1994, SE Appl. 93/132, 14 Jan. 1993; 10 p.; C.A., 121 (1994) 270740u.
- 1057 Suzuki, H.: Automated electrophoresis apparatus and method for positioning substrate film therein. Ger. Offen. DE 4,411,160 (Cl. G01N27/447), 06 Oct. 1994, JP Appl. 93/73,561, 31 Mar. 1993; 14 pp.; C.A., 121 (1994) 276162d.
- 1058 Takada, Y., Nakayama, K., Sakairi, M. and Hirabayashi, T.: Capillary electrophoresis-mass spectrometer for biological analysis. *Jpn. Kokai Tokkyo Koho* JP 06,249,833 (94,249,833) (Cl. G01N27/62), 09 Sep. 1994, Appl. 93/39,639, 01 Mar. 1993; 5 p.; C.A., 122 (1995) 45301c.
- 1059 Zhou, J. and Lunte, S.: Membrane-based on-column mixer for capillary electrophoresis/electrochemistry. *Anal. Chem.*, 67 (1995) 13-18.
- 3b. Detectors and detection procedures
- 1060 Bron, C.R., Godinaud, S., Tissot, M. and Launay, D.: Method for automatically correcting and matching digitized electrophoresis and western blots tests. *PCT Int. Appl.* WO 94 20,844 (Cl. G01N27/447), 15 Sep. 1994, CH Appl. 93/698, 09 Mar. 1993; 17 pp.; C.A., 121 (1994) 276180h.
- 1061 Busch, K.L.: Mass spectrometric detectors for samples separated by planar electrophoresis. *J. Chromatogr. A*, 692 (1995) 275-290 - a review with 65 refs.
- 1062 Cannon, J. and Schallern, G.: Deep-UV laser-induced fluorescence detection in capillary electrophoresis. *Spectroscopy (Eugene)*, 9 (1994) 42-45; C.A., 121 (1994) 220671b.
- 1063 Desbene, P.L., Morin, C.J., Desbene Monvernay, A.M. and Groult, R.S.: Utilization of fluorescein sodium salt in laser-induced indirect fluorimetric detection of ions separated by capillary zone electrophoresis. *J. Chromatogr. A*, 689 (1995) 135-148.
- 1064 Ferris, S.S., Lou, G. and Ewing, A.G.: Scanning electrochemical detection in capillary electrophoresis. *J. Microcolumn Sep.*, 6 (1994) 263-268; C.A., 122 (1995) 45370z.
- 1065 Gilman, S.D., Silverman, C.E. and Ewing, A.G.: Electrogenerated chemiluminescence detection for capillary electrophoresis. *J. Microcolumn Sep.*, 6 (1994) 97-106; C.A., 121 (1994) 270810s.
- 1066 Hanning, A.: Method of liquid bulk refractive index detection. *PCT Int. Appl.* WO 94 17,393 (Cl. G01N21/41), 04 Aug. 1994, SE Appl. 93/231, 27 Jan. 1993; 24 p.; C.A., 121 (1994) 270873q.
- 1067 Mala, Z., Vespalec, R. and Bocek, P.: Capillary zone electrophoresis with indirect photometric detection in the visible range. *Electrophoresis (Weinheim)*, 15 (1994) 1526-1530.
- 1068 Malone, M.A., Weber, P.L., Smyth, M.R. and Lunte, S.M.: Reductive electrochemical detection for capillary electrophoresis. *Anal. Chem.*, 66 (1994) 3782-3787.
- 1069 Morrison, T.B. and Parkinson, J.S.: Quantifying radiolabeled macromolecules and small molecules on a single gel. *BioTechniques*, 17 (1994) 922-926; C.A., 121 (1994) 296578x.
- 1070 Ren, J.C., Li, G.C., Deng, Y.Z. and Cheng, J.K.: Indirect thermooptical detection for capillary electrophoresis separation of amino acids. *Chin. Chem. Lett.*, 5 (1994) 767-768; C.A., 121 (1994) 250334m.
- 1071 Shamsi, S.A. and Danielson, N.D.: Naphthalenesulfonates as electrolytes for capillary electrophoresis of inorganic anions, organic acids, and surfactants with indirect photometric detection. *Anal. Chem.*, 66 (1994) 3757-3764.
- 1072 Slater, J.M. and Watt, E.J.: On-chip microband array electrochemical detector for use in capillary electrophoresis. *Analyst (Cambridge)*, 119 (1994) 2303-2307.
- 1073 Soper, S.A., Legendre, B.L., Jr., Flanagan, J.H., Jr., Williams, D.C. and Hammer, R.P.: Ultrasensitive near-IR fluorescence detection in capillary zone electrophoresis. *Proc. SPIE-Int. Soc. Opt. Eng.*, 213(Biochemical Diagnostic Instrumentation) (1994) 244-254; C.A., 121 (1994) 275896j.
- 1074 Timperman, A.T., Khatib, K. and Sweedler, J.V.: Wavelength-resolved fluorescence detection in capillary electrophoresis. *Anal. Chem.*, 67 (1995) 139-144.

See also 1099, 1112, 1229, 1280, 1668, 1690, 1735.

See also 1029, 1059, 1089, 1093, 1109, 1120, 1123, 1207, 1256, 1257, 1258, 1261, 1277, 1390, 1585, 1589, 1649.

### 3c. Stabilization media for electrophoresis

- 1075 Ciarfardini, G.: A method for the long-term preservation of polyacrylamide gel electrophoresis. *Lett. Appl. Microbiol.*, 19 (1994) 221-224; C.A., 121 (1994) 296412p.
- 1076 Dunn, M.J. and Crisp, S.J.: Detection of proteins in polyacrylamide gels using an ultrasensitive silver staining technique. *Methods Mol. Biol. (Totowa)*, 32(Basic Protein and Peptide Protocols) (1994) 113-118; C.A., 121 (1994) 250265q.
- 1077 Guldberg, P., Henriksen, K.F. and Guettler, F.: Constant denaturant gel electrophoresis without formamide. *BioTechniques*, 16 (1994) 786-788; C.A., 121 (1994) 293868f.
- 1078 Laywood, A., Whittall, R., Gudnason, V., Humphries, S.E. and Day, I.N.M.: Handling of large (300 x 400 mm), thin (0.4 mm) polyacrylamide gels and recovery as dried gels. *BioTechniques*, 17 (1994) 850-854; C.A., 121 (1994) 296416t.
- 1079 Sandoval, J.E. and Pesek, J.J.: Stable covalently-bonded supports for chemical separation apparatus made through a hydride intermediate. U.S. US 5,326,738 (Cl. 502-401; B01J20/06), 05 Jul. 1994, US Appl. 407,816, 15 Sep. 1989; 15 p.; C.A., 121 (1994) 283012h.
- 1080 Smith, B.J.: Drying gels. *Methods Mol. Biol. (Totowa)*, 32(Basic Protein and Peptide Protocols) (1994) 157-161; C.A., 121 (1994) 250266r.
- 1081 Zahnd, A., Funk, M., Vaudaux, P., Lew, D., Scherrer, J.-R. and Hochstrasser, D.F.: A classification of one-dimensional electrophoresis gels using wave packet decomposition. *Appl. Theor. Electrophor.*, 4 (1994) 19-24; C.A., 121 (1994) 250285w.
- 1082 Zewert, T. and Harrington, M.G.: Poly(N-acylalkylenimine) electrophoresis support media. U.S. US 5,338,428 (Cl. 204-299R; C25B9/00), 16 Aug. 1994, Appl. 69,075, 28 May 1993; 10 pp.; C.A., 121 (1994) 225862z.

See also 1056, 1121, 1134, 1136, 1587, 1702, 1707.

### 3d. Quantitative analysis

- 1083 Damerval, C.: Quantification of silver-stained proteins resolved by two-dimensional electrophoresis: genetic variability as related to abundance and solubility in two maize lines. *Electrophoresis (Weinheim)*, 15 (1994) 1573-1579.

### 3e. Preparative scale electrophoresis

- 1084 Margolis, J., Corthals, G. and Horvath, Z.S.: Preparative reflux electrophoresis. *Electrophoresis (Weinheim)*, 16 (1995) 98-100.

See also 1163, 1229.

## 4. SPECIAL TECHNIQUES

### 4a. Automation

See 1690.

### 4b. Computerization and modelling

- 1085 Beckers, J.L.: Calculation of the composition of sample zones in capillary zone electrophoresis. I. Mathematical model. *J. Chromatogr. A*, 693 (1995) 347-357.
- 1086 Davis, J.M.: Statistical theories of peak overlap in chromatography. *Adv. Chromatogr. (N.Y.)*, 34 (1994) 109-176; C.A., 121 (1994) 238951c - a review with 52 refs.
- 1087 Thormann, W., Molteni, S., Stoffel, E., Mosher, R.A. and Chmelik, J.: Computer modeling and experimental validation of the dynamics of capillary isoelectric focusing with electroosmotic zone displacement. *Anal. Methods Instrum.*, 1 (1993) 177-184; C.A., 122 (1995) 22805f.
- 1088 Yokogawa, H.: Processing of electrophoretic data from successive samples from a single source. Ger. Offen. DE 4,405,251 (Cl. G01N27/447), 25 Aug. 1994, JP Appl. 93/30,688, 19 Feb. 1993; 26 pp.; C.A., 121 (1994) 225861y.

See also 1029, 1060, 1114, 1135, 1165, 1220, 1336, 1698.

### 4c. Combination with other physicochemical techniques, (MS, IR etc.)

- 1089 Bachteler, G., Drexhage, K.-H., Arden-Jacob, J., Han, K.-T., Koellner, M., Mueller, R., Sauer, M., Seeger, S. and Wolfrum, J.: Sensitive fluorescence detection in capillary electrophoresis using laser diodes and multiplex dyes. *J. Lumin.*, 62 (1994) 101-108; C.A., 122 (1995) 22845u.
- 1090 Caprioli, R.M.: Online LC/MS and CE/MS. In: Matsuo, T. (Editor), *Biol. Mass Spectrom.: Present Future, [Proc. Kyoto '92 Int. Conf.]* 1992, Wiley, Chichester, 1994, pp. 75-100; C.A., 121 (1994) 249811h - a review with 65 refs.
- 1091 Foret, F., Thompson, T.J., Vouros, P., Karger, B.L., Gebauer, P. and Bocek, P.: Liquid sheath effects on the separation of proteins in capillary electrophoresis/electrospray mass spectrometry. *Anal. Chem.*, 66 (1994) 4450-4458.
- 1092 Hofstadler, S.A., Wahl, J.H., Bakhtiar, R., Anderson, G.A., Bruce, J.E. and Smith, R.D.: Capillary electrophoresis/Fourier-transform ion-cyclotron-resonance mass spectrometry with sustained off-resonance irradiation for the characterization of protein and peptide mixtures. *J. Am. Soc. Mass Spectrom.*, 5 (1994) 894-899; C.A., 121 (1994) 250599b.
- 1093 Olesik, J.W., Kinzer, J.A. and Olesik, S.V.: Capillary electrophoresis inductively coupled plasma spectrometry for rapid elemental speciation. *Anal. Chem.*, 67 (1995) 1-12.
- 1094 Pálmarisdóttir, S. and Edholm, L.-E.: Enhancement of selectivity and concentration sensitivity in capillary zone electrophoresis by on-line coupling with column liquid chromatography and utilizing a double stacking procedure allowing for microliter injections. *J. Chromatogr. A*, 693 (1995) 131-143.

See also 1112, 1132, 1138, 1142, 1176, 1215, 1219, 1235, 1246, 1247, 1585, 1778.

### 4d. Affinity electrophoresis

- 1095 Igloi, G.L.: Affinity electrophoresis of macromolecules: general principles and their application to nucleic acids. *Mol. Interact. Biosep.*, (1993) 511-531; C.A., 121 (1994) 249873e - a review with many refs.

See also 1469.

4e. *Capillary zone electrophoresis and electrokinetic chromatography*

- 1096 Ahuja, E.S., Little, E.L., Nielsen, K.R. and Foley, J.P.: Infinite elution range in micellar electrokinetic capillary chromatography using a nonionic/anionic mixed micellar system. *Anal. Chem.*, 67 (1995) 26-33.
- 1097 Baba, Y.: (Capillary electrophoresis: new technology for high-performance analysis of drugs and biopolymers). *Pharma. Tech. Jpn.*, 10 (1994) 987-996; C.A., 121 (1994) 225312v - a review with 14 refs.
- 1098 Bächmann, K., Göttlicher, B., Haage, I. and Hensel, W.: Improvement of reproducibility in capillary zone electrophoresis by sequential application of high voltage and pressure. *Fresenius J. Anal. Chem.*, 359 (1994) 716-717.
- 1099 Baechmann, K., Haag, I., Haumann, I. and Reineke, F.: Control electroosmotic flow (EOF) during capillary electrophoresis. *Ger. DE 4,315,928* (Cl. G01N27/447), 08 Dec. 1994, Appl. 12 May 1993; 21 p.; C.A., 122 (1995) 71117s.
- 1100 Bello, M.S., Capelli, L. and Righetti, P.G.: Dependence of the electroosmotic mobility on the applied electric field and its reproducibility in capillary electrophoresis. *J. Chromatogr. A*, 684 (1994) 311-322.
- 1101 Burggraf, N., Manz, A., Verpoorte, E., Effenhauser, C.S., Widmer, H.M. and de Rooij, N.F.: A novel approach to ion separations in solution: synchronized cyclic capillary electrophoresis (SCCE). *Sens. Actuators, B*, 20 (1994) 103-110; C.A., 122 (1995) 45362y.
- 1102 Chen, Y., Hölte, J.-V. and Schwarz, U.: Preparation of highly condensed polyacrylamide gel-filled capillaries with low detection background. *J. Chromatogr. A*, 685 (1994) 121-129.
- 1103 Cohen, N. and Grushka, E.: Influence of the capillary edge on the separation efficiency in capillary electrophoresis. *J. Chromatogr. A*, 684 (1994) 323-328.
- 1104 Emmer, A. and Roeraade, J.: Performance of zwitterionic and cationic fluorosurfactants as buffer additives for capillary electrophoresis of proteins. *J. Liq. Chromatogr.*, 17 (1994) 3831-3846.
- 1105 Erim, F.B., Boelens, H.F.M. and Kraak, J.C.: Applicability of capillary zone electrophoresis to study metal complexation in solution. *Anal. Chim. Acta*, 294 (1994) 155-163.
- 1106 Ewing, A.G., Hayes, M.A. and Kheterpal, I.: Capillary electrophoresis apparatus with improved electroosmotic flow control. *U.S. US 5,358,618* (Cl. 204-299R; C25B9/00), 25 Oct. 1994, Appl. 7,214, 22 Jan. 1993; 11 pp.; C.A., 122 (1995) 4935u.
- 1107 Gelfi, C., Orsi, A., Leoncini, F. and Righetti, P.G.: Fluidified polyacrylamides as molecular sieves in capillary zone electrophoresis of DNA fragments. *J. Chromatogr. A*, 689 (1995) 97-105.
- 1108 Gilman, S.D. and Ewing, A.G.: Analysis of single cells by capillary electrophoresis with on-column derivatization and laser-induced fluorescence detection. *Anal. Chem.*, 67 (1995) 58-64.
- 1109 Hauser, P.C., Renner, N.D. and Hong, A.P.C.: Anion detection in capillary electrophoresis with ion-selective microelectrodes. *Anal. Chim. Acta*, 295 (1994) 181-186.
- 1110 Heegaard, N.H.H. and Robey, F.A.: The emerging role of capillary electrophoresis as a tool for the study of biomolecular non-covalent interactions. *Am. Lab. (Shelton)*, 26 (1994) 28T-28X; C.A., 121 (1994) 249913t - a review with 29 refs.
- 1111 Issaq, H.J.: Comparison of chiral separations in capillary zone electrophoresis with other methods. *Instrum. Sci. Technol.*, 22 (1994) 119-149; C.A., 121 (1994) 220618q.
- 1112 Jones, W.R. and Jandik, P.: Method for separating ionic species using capillary electrophoresis. *U.S. US 5,366,601* (Cl. 204-180.1; B01D57/02), 22 Nov. 1994, US Appl. 471,535, 29 Jan. 1990; 24 p.; C.A., 122 (1995) 71124s.
- 1113 Karim, M.R., Janson, J.-C. and Takagi, T.: Size-dependent separation of proteins in the presence of sodium dodecyl sulfate and dextran in capillary electrophoresis: effect of molecular weight of dextran. *Electrophoresis (Weinheim)*, 15 (1994) 1531-1534.
- 1114 Keely, C.A., van de Goor, T.A.A.M. and McManigill, D.: Modeling flow profiles and dispersion in capillary electrophoresis with nonuniform zeta potential. *Anal. Chem.*, 66 (1994) 4236-4242.
- 1115 Landers, J.P. and Spelsberg, T.C.: Future prospects for capillary electrophoresis. In: Landers, J.P. (Editor), *Handb. Capillary Electrophor.*, CRC, Boca Raton, 1994, pp. 593-601; C.A., 121 (1994) 249929c - a review with 20 refs.
- 1116 Larman, J.P., Jr.: Two dimensional liquid chromatography-capillary electrophoresis. Avail. *Microfilms Int.*, Order No. DA9415346, 1993, 312 p.; C.A., 121 (1994) 244386p.
- 1117 Li, H. and Shen, Z.: (Study on separation mechanism of high performance capillary electrophoresis). *Huagong Xuebao*, 45 (1994) 475-481; C.A., 121 (1994) 225411b.
- 1118 Lloyd, D.K. and Wätzig, H.: Sodium dodecyl sulfate solution is an effective between-run rinse for capillary electrophoresis of samples in biological matrices. *J. Chromatogr. B*, 663 (1995) 400-405.
- 1119 Manabe, T.: (Capillary electrophoresis: application to the analysis of biopolymers). *Bunseki*, (1994) 467-472; C.A., 121 (1994) 249844m - a review with 58 refs.
- 1120 Mank, A.J.G., Lingeman, H. and Gooijer, C.: Semiconductor-laser-induced fluorescence detection in capillary electrophoresis using precolumn thiol derivatization. *J. High Resolut. Chromatogr.*, 17 (1994) 797-798.
- 1121 Mazzeo, J.R. and Krull, I.S.: Modification of capillaries and buffers for enhanced separations in capillary zone electrophoresis and capillary isoelectric focusing of biopolymers. In: Landers, J.P. (Editor), *Handb. Capillary Electrophor.*, CRC, Boca Raton, 1994, pp. 495-512; C.A., 121 (1994) 249925y - a review with 45 refs.
- 1122 Mesaros, J.M. and Ewing, A.G.: Characterization of continuous electrophoretic separations in narrow channels coupled to small bore capillaries. *J. Microcolumn Sep.*, 6 (1994) 483-494; C.A., 121 (1994) 275911k.
- 1123 Nasu, H., Nakamura, Y., Fujimiya, H. and Yamamoto, K.: Chemiluminescence detection for capillary gel electrophoresis. *Eur. Pat. Appl. EP 616,212* (Cl. G01N27/447), 21 Sep. 1994, JP Appl. 93/78,569, 15 Mar. 1993; 15 pp.; C.A., 121 (1994) 250624f.
- 1124 Nelson, R.J. and Burgi, D.S.: Temperature control in capillary electrophoresis. In: Landers, J.P. (Editor), *Handb. Capillary Electrophor.*, CRC, Boca Raton, 1994, pp. 549-562; C.A., 121 (1994) 249928b - a review with 42 refs.
- 1125 Nielsen, K.R. and Foley, J.P.: Zone sharpening of neutral solutes in micellar electrokinetic chromatography with electrokinetic injection. *J. Chromatogr. A*, 686 (1994) 283-291.

- 1126 Oda, R.P. and Landers, J.P.: Introduction to capillary electrophoresis. In: Landers, J.P. (Editor), *Handb. Capillary Electrophor.*, CRC, Boca Raton, 1994, pp. 9-42; C.A., 121 (1994) 249915v - a review with 54 refs.
- 1127 Pietta, P.G., Mauri, P.L., Gardana, C. and Zini, L.: Micellar electrokinetic chromatographic/ultraviolet diode array analysis of *Eleutherococcus senticosus*. *Phytochem. Anal.*, 5 (1994) 305-308; C.A., 122 (1995) 17302k.
- 1128 Seiler, K., Fluri, K. and Harrison, D.J.: Micromachining a miniaturized capillary electrophoresis-based chemical analysis system on a chip. *Mitt. Geb. Lebensmittelunters. Hyg.*, 85 (1994) 59-68; C.A., 121 (1994) 225583j.
- 1129 Shi, Y. and Fritz, J.S.: Capillary zone electrophoresis of neutral organic molecules in organic - aqueous solution. *J. High Resolut. Chromatogr.*, 17 (1994) 713-718.
- 1130 Shihabi, Z.K. and Garcia, L.L.: Effects of sample matrix on separation by capillary electrophoresis. In: Landers, J.P. (Editor), *Handb. Capillary Electrophor.*, CRC, Boca Raton, 1994, p. 537-548; C.A., 121 (1994) 249927a - a review with 39 refs.
- 1131 Smith, J.T. and El Rassi, Z.: Micellar electrokinetic capillary chromatography with *in situ* charged micelles. IV. Influence of the nature of the alkylglycoside surfactant. *J. Chromatogr. A*, 685 (1994) 131-143.
- 1132 Smith, R.D., Goodlett, D.R. and Wahl, J.H.: Capillary electrophoresis-mass spectrometry. In: Landers, J.P. (Editor), *Handb. Capillary Electrophor.*, CRC, Boca Raton, 1994, pp. 185-206; C.A., 121 (1994) 249919z - a review with 52 refs.
- 1133 Sun, P., Landman, A. and Hartwick, R.A.: Chitosan coated capillary with reversed electroosmotic flow in capillary electrophoresis for the separation of basic drugs and proteins. *J. Microcolumn Sep.*, 6 (1994) 403-407; C.A., 122 (1995) 22832n.
- 1134 Sun, P., Landman, A., Barker, G.E. and Hartwick, R.A.: Synthesis and evaluation of anionic polymer-coated capillaries with pH-independent electroosmotic flows for capillary electrophoresis. *J. Chromatogr. A*, 685 (1994) 303-312.
- 1135 Tavares, M.F.M.: Computer-assisted optimization of separations in capillary zone electrophoresis. Avail. *Univ. Microfilms Int.*, Order No. DA9418074, 1993, 316 p.; C.A., 121 (1994) 244521d.
- 1136 Terebe, S., Ozaki, H. and Tanaka, Y.: New pseudostationary phases for electrokinetic chromatography: a high-molecular surfactant and proteins. *J. Chin. Chem. Soc. (Taipei)*, 41 (1994) 251-257; C.A., 121 (1994) 244503z.
- 1137 Tickle, D.C., Okafo, G.N., Camilleri, P., Jones, R.F.D. and Kirby, A.J.: Glucopyranoside-based surfactants as pseudostationary phases for chiral separations in capillary electrophoresis. *Anal. Chem.*, 66 (1994) 4121-4126.
- 1138 Tomlinson, A.J., Benson, L.M., Braddock, W.D. and Oda, R.P.: On-line preconcentration-capillary electrophoresis-mass spectrometry (PC-CE-MS). *J. High Resolut. Chromatogr.*, 17 (1994) 729-731.
- 1139 Van der Vlis, E., Mazereeuw, M., Tjaden, U.R., Irth, H. and van der Greef, J.: Combined liquid-liquid electroextraction and isotachopheresis as a fast on-line focusing step in capillary electrophoresis. *J. Chromatogr. A*, 687 (1994) 333-341.
- 1140 Vespalec, R., Fanali, S. and Bocek, P.: Consequences of a maximum existing in the dependence of separation selectivity on concentration of cyclodextrin added as chiral selector in capillary zone electrophoresis. *Electrophoresis (Weinheim)*, 15 (1994) 1523-1525.
- 1141 Vogt, C. and Conradi, S.: Complex equilibria in capillary zone electrophoresis and their use for the separation of rare earth metal ions. *Anal. Chim. Acta*, 294 (1994) 145-153.
- 1142 Wahl, J.H., Hofstadler, S.A. and Smith, R.D.: Direct electrospray ion current monitoring detection and its use with on-line capillary electrophoresis mass spectrometry. *Anal. Chem.*, 67 (1995) 462-465.
- 1143 Wojtusik, M.J.: Capillary electrophoresis, a new technique for the semiconductor and plating industries. *Proc. AESP Annu. Tech. Conf.*, 80th (1993) 61-67; C.A., 122 (1995) 70801y.
- 1144 Xue, Y. and Yeung, E.S.: Characterization of band broadening in capillary electrophoresis due to nonuniform capillary geometries. *Anal. Chem.*, 66 (1994) 3575-3580.
- 1145 Yamaguchi, M., Izumi, T. and Isobe, T.: (Separation of biomacromolecules by HPCE using natural polysaccharides). *Kuromatogurafi*, 15 (1994) 110-111; C.A., 121 (1994) 275895h.
- 1146 Yang, S. and Khaleli, M.G.: Chemical selectivity in micellar electrokinetic chromatography: characterization of solute - micelle interactions for classification of surfactants. *Anal. Chem.*, 67 (1995) 499-510.
- 1147 Yuan, D., Wang, X., Tong, Q. and Yang, P.: (Application of surfactants as electrolyte additives in micellar electrokinetic capillary chromatography). *Fenxi Ceshi Xuebao*, 13 (1994) 87-97; C.A., 121 (1994) 270468m.
- See also 1023, 1024, 1026, 1027, 1028, 1031, 1033, 1037, 1041, 1043, 1044, 1048, 1049, 1050, 1055, 1058, 1059, 1062, 1063, 1064, 1065, 1067, 1068, 1070, 1071, 1073, 1074, 1085, 1089, 1090, 1091, 1092, 1094, 1151, 1152, 1153, 1154, 1155, 1156, 1157, 1164, 1165, 1166, 1168, 1169, 1170, 1171, 1172, 1173, 1176, 1183, 1184, 1185, 1186, 1187, 1188, 1189, 1203, 1204, 1205, 1206, 1207, 1208, 1209, 1212, 1213, 1214, 1217, 1226, 1227, 1228, 1229, 1231, 1232, 1235, 1238, 1239, 1246, 1247, 1253, 1255, 1258, 1259, 1268, 1270, 1278, 1281, 1282, 1286, 1287, 1326, 1380, 1381, 1387, 1390, 1409, 1414, 1416, 1469, 1482, 1513, 1517, 1524, 1542, 1582, 1584, 1585, 1589, 1590, 1643, 1645, 1649, 1659, 1662, 1680, 1682, 1702, 1713, 1715, 1745, 1753, 1755, 1756, 1757, 1758, 1759, 1760, 1761, 1762, 1763, 1769, 1770, 1771, 1772, 1773, 1774, 1775, 1776, 1777, 1778, 1779, 1780, 1781, 1782, 1783, 1784, 1787, 1788, 1791, 1792, 1798, 1800, 1802, 1804, 1805, 1806, 1807, 1809, 1810, 1811, 1812, 1813, 1814.
- 4f. *Isotachopheresis*
- 1148 Fukushi, K. and Hiirio, K.: Behaviour of periodate ion in isotachopheresis using cyclodextrins. *J. Chromatogr. A*, 684 (1994) 343-349.

- 1149 Hutta, M., Kaniansky, D., Kovalcikova, E., Marak, J., Chalanyova, M., Madajova, V. and Simunicova, E.: Preparative capillary isotachopheresis as a sample pretreatment technique for complex ionic matrices in high-performance liquid chromatography. *J. Chromatogr. A*, 689 (1995) 123-133.
- 1150 Liu, W., Zheng, J., Hu, R., Mu, J. and Zhuo, X.: (Simultaneous analysis of inorganic and organic anions by isotachopheresis). *Sepu*, 12 (1994) 241-244; C.A., 121 (1994) 220552p.
- 1151 Wanders, B.J. and Everaerts, F.M.: Isotachopheresis in capillary electrophoresis. In: Landers, J.P. (Editor), *Handb. Capillary Electrophor.*, CRC, Boca Raton, 1994, pp. 111-127; C.A., 121 (1994) 249917x - a review with 29 refs.

See also 1087, 1130, 1790, 1801, 1811.

#### 4g. Enantiomers, separation

- 1152 Aubry, A.-F., Markoglou, N., Descorps, V., Wainer, I.W. and Félix, G.: Evaluation of a chiral stationary phase based on mixed immobilized proteins. *J. Chromatogr. A*, 685 (1994) 1-6.
- 1153 Gahm, K.-H. and Stalcup, A.M.: Capillary zone electrophoresis study of naphthylethylcarbamoylated  $\beta$ -cyclodextrins. *Anal. Chem.*, 67 (1995) 19-25.
- 1154 Quang, C. and Khaleidi, M.G.: Extending the scope of chiral separation of basic compounds by cyclodextrin-mediated capillary zone electrophoresis. *J. Chromatogr. A*, 692 (1995) 253-265.
- 1155 Rawjee, Y.Y., Williams, R.L. and Vigh, G.: Efficiency optimization in capillary electrophoretic chiral separations using dynamic mobility matching. *Anal. Chem.*, 66 (1994) 3777-3781.
- 1156 Tait, R.J., Thompson, D.O., Stella, V.J. and Stobaugh, J.F.: Sulfoethyl ether  $\beta$ -cyclodextrin as a chiral discriminator for use with capillary electrophoresis. *Anal. Chem.*, 66 (1994) 40113-4018.
- 1157 Wang, J. and Warner, I.M.: Chiral separations using micellar electrokinetic capillary chromatography and a polymerized chiral micelle. *Anal. Chem.*, 66 (1994) 3773-3776.

See also 1094, 1111, 1137, 1140, 1183, 1212, 1760, 1771, 1772, 1779, 1806.

#### 4h. Two dimensional electrophoresis

- 1158 Gross, M., Kosmowsky, I.J., Lorenz, R., Molitoris, H.P. and Jaenicke, R.: Response of bacteria and fungi to high-pressure stress as investigated by two-dimensional polyacrylamide gel electrophoresis. *Electrophoresis (Weinheim)*, 15 (1994) 1559-1565.
- 1159 Rabilloud, T., Valette, C. and Lawrence, J.J.: Sample application by in-gel rehydration improves the resolution of two-dimensional electrophoresis with immobilized pH gradients in the first dimension. *Electrophoresis (Weinheim)*, 15 (1994) 1552-1558.

See also 1083, 1254, 1268, 1273, 1276, 1314, 1329, 1337, 1356, 1362, 1371, 1375, 1394, 1395, 1439, 1572, 1661, 1698, 1701, 1731, 1734, 1736, 1737, 1741, 1747, 1749, 1785.

#### 4i. Other special techniques

- 1160 Danziger, R.S.: Gel electrophoresis measurement of dynamic reaction product by continuous sampling. U.S. US 5,344,534 (Cl. 204-182.8; C2587/00), 06 Sep. 1994, Appl. 27,284, 05 Mar. 1993; 8 pp.; C.A., 121 (1994) 225864b.
- 1161 Dasgupta, P.K. and Li-Yuan, B.: Electrophoresis with chemically suppressed detection. U.S. US 5,358,612 (Cl. 204-180.1; GO1N27/26), 25 Oct. 1994, US Appl. 764,645, 24 Sep. 1991; 12 pp.; C.A., 122 (1995) 4934t.
- 1162 Jorgenson, J.W. and Lemmo, A.V.: Two dimensional separation system containing a flow gating system, a liquid chromatography column, and an electrophoresis capillary. *PCT Int. Appl. WO 94 20,840*, (Cl. GO1N27/26), 15 Sep. 1994, US Appl. 28,678, 09 Mar. 1993; 39 pp.; C.A., 121 (1994) 296600y.
- 1163 Watson, J.: Continuous free-flow electrophoresis: a modified approach. *Sep. Technol.*, 4 (1994) 239-243; C.A., 122 (1995) 34155g.

See also 1051.

### 5. HYDROCARBONS AND HALOGEN DERIVATIVES

#### 5b. Cyclic hydrocarbons, fullerenes

See 1129, 1754.

### 7. PHENOLS

- 1164 Bachmann, K., Göttlicher, B., Haag, I., Hannina, M. and Hensel, W.: Sample stacking for charched phenol derivatives in capillary zone electrophoresis. *Fresenius J. Anal. Chem.*, 350 (1994) 368-371.
- 1165 Masselter, S.M. and Zemmann, A.J.: Influence of buffer electrolyte pH on the migration behavior of phenolic compounds in co-electroosmotic capillary electrophoresis. *J. Chromatogr. A*, 693 (1995) 359-365.

### 8. SUBSTANCES CONTAINING HETEROCYCLIC OXYGEN

#### 8c. Other compounds with heterocyclic oxygen (incl. tannins)

- 1166 Bogan, D.P., Deasy, B., O'Kennedy, R., Smyth, M.R. and Fuhr, U.: Determination of free and total 7-hydroxycoumarin in urine and serum by capillary electrophoresis. *J. Chromatogr. B*, 663 (1995) 371-378.

### 9. OXO COMPOUNDS, ETHERS, EPOXIDES AND QUINONES

See 1129.

## 10. CARBOHYDRATES

## 10a. Mono and oligosaccharides. Structural studies

- 1167 El Rassi, Z.: Capillary electrophoresis of carbohydrates. *Adv. Chromatogr. (N.Y.)*, 34 (1994) 177-250; C.A., 121 (1994) 301156q - a review with 147 refs.
- 1168 Guttman, A., Cooke, N. and Starr, C.M.: Capillary electrophoresis separation of oligosaccharides: I. Effect of operational variables. *Electrophoresis (Weinheim)*, 15 (1994) 1518-1522.
- 1169 Klockow, A., Widmer, H.M., Amado, R. and Paulus, A.: Capillary electrophoresis of ANTS labelled oligosaccharide ladders and complex carbohydrates. *Fresenius J. Anal. Chem.*, 350 (1994) 415-425.
- 1170 Olechno, J.D. and Ulfelder, K.J.: Carbohydrate analysis by capillary electrophoresis. In: Landers, J.P. (Editor), *Handb. Capillary Electrophor.*, CRC, Boca Raton, 1994, pp. 255-286; C.A., 121 (1994) 249920t.
- 1171 Ren, J.-C., He, J.-L., Deng, Y.-Z. and Cheng, J.-K.: (Rapid separation of disaccharides by capillary zone electrophoresis with interference refractive index detection). *Gaodeng Xuexiao Huaxue Xuebao*, 14 (1993) 1661-1664; C.A., 121 (1994) 244517g.
- 1172 Ye, J. and Baldwin, R.P.: Determination of carbohydrates, sugar acids and alditols by capillary electrophoresis and electrochemical detection at a copper electrode. *J. Chromatogr. A*, 687 (1994) 141-148.

## 10b. Polysaccharides, mucopolysaccharides, lipopolysaccharides

- 1173 Brewster, J.D. and Fishman, M.L.: Capillary electrophoresis of plant starches as the iodine complexes. *J. Chromatogr. A*, 693 (1995) 382-387.
- 1174 Burns, N.L., Harris, J.M. and van Alstine, J.M.: Electrokinetic characterization of polysaccharide and related coatings. *Front. Biomed. Biotechnol.*, 1 (1993) 298-309; C.A., 122 (1995) 33562a.
- 1175 Gohel, M.D.I., Shum, D.K.Y. and Li, M.K.: Crystallization of urinary calcium oxalate at standardized osmolality and pH in the frozen state. *Clin. Chim. Acta*, 231 (1994) 11-22.
- 1176 Kouznetsov, D.A., Ivanov, A.A. and Veletsky, P.R.: Detection of alkylated cellulose derivatives in several archaeological linen textile samples by capillary electrophoresis/mass spectrometry. *Anal. Chem.*, 66 (1994) 4359-4365.
- 1177 Li, X.A., Hatanaka, K., Guo, L., Kitamura, Y. and Yamamoto, A.: Binding of serum amyloid P component to heparin in human serum. *Biochim. Biophys. Acta*, 1201 (1994) 143-148.

See also 1352.

## 10c. Glycoproteins and their constituents

- 1178 Ivanov, A.E., Zhigis, L.S., Rapoport, E.M., Lisyutina, O.E. and Zubov, V.P.: Characterization of weak hydrophobic composite sorbents and their application to the isolation of bacterial lectin. *J. Chromatogr. B*, 664 (1995) 219-223.
- 1179 Matsui, T., Ozeki, Y., Suzuki, M., Hino, A. and Titani, K.: Purification and characterization of two  $\text{Ca}^{2+}$ -dependent lectins from coelomic plasma of sea cucumber, *Stichopus japonicus*. *J. Biochem. (Tokyo)*, 116 (1994) 1127-1133.

- 1180 Puri, K.D. and Surolia, A.: Amino acid sequence of the winged bean (*Psophocarpus tetragonolobus*) basic lectin. Adenine binding and identification of the active-site tryptophan residue. *J. Biol. Chem.*, 269 (1994) 30917-30926.
- 1181 Smith, K.D., Elliott, M.A., Elliott, H.G., McLaughlin, C.M., Wightman, P. and Wood, G.C.: Heterogeneity of  $\alpha_1$ -acid glycoprotein in rheumatoid arthritis. *J. Chromatogr. B*, 661 (1994) 7-14.
- 1182 Van der Linden, E.C.M., de Graaf, T.W., Anbergen, M.G., Dekker, R.M., van Ommen, E.C.R., van den Eijnden, D.H. and van Dijk, W.: Preparative affinity electrophoresis of different glycoforms of serum glycoproteins: application for the study of inflammation-induced expression of sialyl-Lewis groups on  $\alpha_1$ -acid glycoprotein (orosomucoid). *Glycosylation Dis.*, 1 (1994) 45-52; C.A., 121 (1994) 250283u.

See also 1223, 1267, 1358.

## 11. ORGANIC ACIDS AND LIPIDS

## 11a. Organic acids and simple esters

- 1183 Biggin, M.E., Williams, R.L. and Vigh, G.: Experimental evidence for the existence of duoselective (Type III) enantiomer separations in the capillary electrophoretic analysis of chiral weak acids. *J. Chromatogr. A*, 692 (1995) 319-325.
- 1184 Chen, H.-R. and Sheu, S.-J.: Determination of the aromatic acids in traditional Chinese medicinal preparations by reversed electroosmotic flow capillary electrophoresis. *Chin. Pharm. J. (Taipei)*, 46 (1994) 145-153; C.A., 121 (1994) 308485e.
- 1185 Dabek-Zlotorzynska, E. and Dlouhy, J.F.: Capillary zone electrophoresis with indirect UV detection of organic anions using 2,6-naphthalenedicarboxylic acid. *J. Chromatogr. A*, 685 (1994) 145-153.
- 1186 Jackson, P.E.: Analysis of oxalate in Bayer liquors: a comparison of ion chromatography and capillary electrophoresis. *J. Chromatogr. A*, 693 (1995) 155-161.
- 1187 Lagoutte, D., Lombard, G., Nisseron, S., Papet, M.P. and Saint-Jalm, Y.: Determination of organic acids in cigarette smoke by high-performance liquid chromatography and capillary electrophoresis. *J. Chromatogr. A*, 684 (1994) 251-257.
- 1188 Reid, R.H.P.: Studies on the retention behaviour of a group of organic anions of biochemical interest on quaternary bonded silica columns equilibrated with a functionally coherent series of counterions. Use of 2-(N-morpholino)ethanesulphonate as a counterion and N-tris(hydroxymethyl)methyl-3-aminopropane-sulphonate as an eluent. *J. Chromatogr. A*, 684 (1994) 221-234.
- 1189 Wiley, J.P.: Determination of polycarboxylic acids by capillary electrophoresis with copper complexation. *J. Chromatogr. A*, 692 (1995) 267-274.

See also 1071, 1152, 1813.

## 11c. Lipids and their constituents

- 1190 Madkell, J.P.: Electrophoretic analysis of the lipopolysaccharides of *Bacteroides* spp. *Antonie van Leeuwenhoek*, 65 (1994) 155-161; C.A., 121 (1994) 275887g.

## 11d. Lipoproteins and their constituents

- 1191 Calabresi, L., Vecchio, G., Longhi, R., Gianazza, E., Palm, G., Wadensten, H., Hammarström, A., Olsson, A., Karlström, A., Sejlitz, T. et al.: Molecular characterization of native and recombinant apolipoprotein A-I<sub>Milano</sub> dimer. The introduction of an interchain disulfide bridge remarkably alters the physicochemical properties of apolipoprotein A-I. *J. Biol. Chem.*, 269 (1994) 32168-32174.
- 1192 Heinemann, T., Metzger, S., Fisher, E.A., Breslow, J.L. and Huang, L.-S.: Alternative polyadenylation of apolipoprotein B RNA is a major cause of B-48 protein formation in rat hepatoma cell lines transfected with human apoB-100 minigenes. *J. Lipid Res.*, 35 (1994) 2200-2211.
- 1193 Ikeda, Y., Ohta, T. and Matsuda, I.: Interaction between apo A-I-containing lipoproteins and lecithin:cholesterol acyltransferase. *Biochim. Biophys. Acta*, 1215 (1994) 307-313.
- 1194 Levy, E., Roy, C.C., Thibault, L., Bonin, A., Brochu, P. and Seidman, E.G.: Variable expression of familial heterozygous hypobetalipoproteinemia: transient malabsorption during infancy. *J. Lipid Res.*, 35 (1994) 2170-2177.
- 1195 Nauck, M. and Wieland, H.: (Measurement of lipoprotein cholesterol with the cholesterol dehydrogenase method after electrophoretic lipoprotein separation. A comparison with quantitative lipoprotein electrophoresis and the ultracentrifugation method). *Klin. Labor.*, 39 (1993) 757-765; *C.A.*, 122 (1995) 1193q.
- 1196 O'Meara, N.M., Cabana, V.G., Lukens, J.R., Loharikar, B., Forte, T.M., Polonsky, K.S. and Getz, G.S.: Heparin-induced lipolysis in hypertriglyceridemic subjects results in the formation of atypical HDL particles. *J. Lipid Res.*, 35 (1994) 2178-2190.
- 1197 Roheim, P.S. and Asztalos, B.F.: Clinical significance of lipoprotein size and risk for coronary atherosclerosis. *Clin. Chem. (Washington)*, 41 (1995) 147-152.
- 1198 Schaefer, J.R., Hackler, R., Brand, S., Schwarz, S., Kleine, T.O. and Steinmetz, A.: Apolipoprotein AI, AII, and AIV isoforms in plasma determined by automated isoelectric focusing with PhastSystem™ and immunofixation. *Clin. Chem. (Washington)*, 41 (1995) 76-81.
- 1199 Shibusawa, Y., Mugiya, M., Matsumoto, U. and Ito, Y.: Complementary use of counter-current chromatography and hydroxyapatite chromatography for the separation of three main classes of lipoproteins from human serum. *J. Chromatogr. B*, 664 (1995) 295-301.
- 1200 Verges, B., Rader, D., Schaefer, J., Zech, L., Kindt, M., Fairwell, T., Gambert, P. and Brewer, H.B., Jr.: *In vivo* metabolism of apolipoprotein A-IV in severe hypertriglyceridemia: a combined radiotracer and stable isotope kinetic study. *J. Lipid Res.*, 35 (1994) 2280-2291.
- 1201 White, A.L. and Lanford, R.E.: Cell surface assembly of lipoprotein(a) in primary cultures of baboon hepatocytes. *J. Biol. Chem.*, 269 (1994) 28716-28723.
- 1202 Zakharova, T.S., Ivanov, A.S., Echkalov, A.P., Berezov, A.T., Khalilov, E.M. and Archakov, A.I.: (Cholesterol content in liposomes regulates their interaction with blood serum lipoproteins). *Biokhimiya (Moscow)*, 59 (1994) 712-719.

## 13. STEROIDS

## 13a. General techniques

- 1203 Milofsky, R.E., Malberg, M.G. and Smith, J.M.: Determination of steroids by micellar electrokinetic chromatography with sensitized room temperature lanthanide ion luminescence detection. *J. High Resolut. Chromatogr.*, 17 (1994) 731-732.
- 1204 Touchstone, J.C. and Levin, S.S.: Capillary electrophoresis of steroid conjugates. *J. Liq. Chromatogr.*, 17 (1994) 3925-3931.

## 13c. Estrogens

- 1205 Chan, K.C., Muschik, G.M., Issaq, H.J. and Siiteri, P.K.: Separation of estrogens by micellar electrokinetic chromatography. *J. Chromatogr. A*, 690 (1995) 149-154.

## 13d. Sterols

See 1204.

## 16. NITRO AND NITROSO COMPOUNDS

See 1129.

## 17. AMINES, AMIDES AND RELATED NITROGEN COMPOUNDS

## 17a. Amines and polyamines

- 1206 Qi, S., Huang, A. and Sun, Y.: (Purity control of *p*-aminophenol in electrochemical production and purification using capillary electrophoresis). *Sepu*, 12 (1994) 348-350; *C.A.*, 122 (1995) 22847w.

## 17b. Catecholamines and their metabolites

See 1108, 1188.

## 18. AMINO ACIDS AND PEPTIDES; CHEMICAL STRUCTURE OF PROTEINS

## 18a. Amino acids and their derivatives

- 1207 Flanagan, J.H., Jr., Legendre, B.L., Jr., Hammer, R.P. and Soper, S.A.: Binary solvent effects in capillary zone electrophoresis with ultrasensitive near-IR fluorescence detection of related tri-carbocyanine dyes and dye-labeled amino acids. *Anal. Chem.*, 67 (1995) 341-347.
- 1208 Gilman, S.D., Pietron, J.J. and Ewing, A.G.: Post-column derivatization in narrow-bore capillaries for the analysis of amino acids and proteins by capillary electrophoresis with fluorescence detection. *J. Microcolumn Sep.*, 6 (1994) 373-384; *C.A.*, 122 (1995) 22843s.

- 1209 Janini, G.M., Muschik, G.M. and Isaaq, H.J.: Capillary electrophoresis separation of heterocyclic nitrosoamino acid-conformers at sub-ambient temperatures. *J. High Resolut. Chromatogr.*, 17 (1994) 753-755.
- 1210 Oguri, S., Uchida, C., Miyake, Y., Miki, Y. and Kakehi, K.: 1-Methoxycarbonylindolizine-3,5-dicarbaldehyde as a derivatization reagent for aminocompounds in high-performance capillary electrophoresis. *Analyst (Cambridge)*, 120 (1995) 63-68.
- 1211 Sakakibara, Y., Suiko, M. and Liu, M.-C.: *De novo* sulfation of L-tyrosine in HepG2 human hepatoma cells and its possible functional implication. *Eur. J. Biochem.*, 226 (1994) 293-301.
- 1212 Walbroehl, Y. and Wagner, J.: Chiral separations of amino acids by capillary electrophoresis and high-performance liquid chromatography employing chiral crown ethers. *J. Chromatogr. A*, 685 (1994) 321-329.
- 1213 Zhang, L., Yang, R.H. and Jin, S.: Capillary zone electrophoresis and micellar electrokinetic capillary chromatography of some protected amino acids and peptides. *Chin. Chem. Lett.*, 5 (1994) 765-766; *C.A.*, 121 (1994) 250595x.
- 1214 Zhou, S.Y., Zuo, H., Stobnaugh, J.F., Lunte, C.E. and Lunte, S.M.: Continuous *in vivo* monitoring of amino acid neurotransmitters by microdialysis sampling with on-line derivatization and capillary electrophoresis separation. *Anal. Chem.*, 67 (1995) 594-599.
- See also 1070, 1108, 1120, 1140, 1152, 1264.
- 18b. *Peptides, peptidic and proteinous hormones, growth factors*
- 1215 Affolter, M., Watts, J.D., Krebs, D.L. and Aebersold, R.: Evaluation of two-dimensional phosphopeptide maps by electrospray ionization mass spectrometry of recovered peptides. *Anal. Biochem.*, 223 (1994) 74-81.
- 1216 Apparailly, F. and Combarnous, Y.: Role of sialic acid residues in the *in vitro* superactivity of human choriongonadotropin (hCG) in rat Leydig cells. *Biochim. Biophys. Acta*, 1224 (1994) 559-565.
- 1217 Begley, D.J.: Free zone capillary electrophoresis. *Methods Mol. Biol. (Totowa)*, 32(Basic Protein and Peptide Protocols) (1994) 467-482; *C.A.*, 121 (1994) 250271p.
- 1218 Elgort, A. and Shields, D.: Prosomatostatin processing in pituitary GH<sub>3</sub> cells. Identification and secretion of the intact propeptide. *J. Biol. Chem.*, 269 (1994) 30668-30675.
- 1219 Fang, L., Zhang, R., Williams, E.R. and Zare, R.N.: On-line time-of-flight mass spectrometric analysis of peptides separated by capillary electrophoresis. *Anal. Chem.*, 66 (1994) 3696-3701.
- 1220 Furka, A., Sebestyen, F. and Dibo, G.: Computer program facilitating separation of peptides from mixtures prepared by the 'portioning-mixing' method. In: Schneider, C.H. and Eberle, A.N. (Editors), *Pept. 1992, Proc. Eur. Pept. Symp.*, 22nd 1992, ESCOM, Leiden, 1993, pp. 463-464; *C.A.*, 121 (1994) 276033n.
- 1221 Guesdon, F., Waller, R.J. and Saklatvala, J.: Specific activation of  $\beta$ -casein kinase by the inflammatory cytokines interleukin 1 and tumour necrosis factor. *Biochem. J.*, 304 (1994) 761-768.
- 1222 Hoermann, R., Berger, P., Spoettl, G., Gillesberger, F., Kardana, A., Cole, L.A. and Mann, K.: Immunological recognition and clinical significance of nicked human chorionic gonadotropin in testicular cancer. *Clin. Chem. (Washington)*, 40 (1994) 2306-2312.
- 1223 Jahnen-Dechent, W., Trindl, A., Godovac-Zimmanmann, J. and Müller-Esterl, W.: Posttranslational processing of human  $\alpha_2$ -HS glycoprotein (human fetuin). Evidence for the production of a phosphorylated single-chain form by hepatoma cells. *Eur. J. Biochem.*, 226 (1994) 59-69.
- 1224 Judd, R.C.: Electrophoresis of peptides. *Methods Mol. Biol. (Totowa)*, 32(Basic Protein and Peptide Protocols) (1994) 49-57; *C.A.*, 121 (1994) 250258q.
- 1225 Kimball, S.R., Karinch, A.M., Feldhoff, R.C., Mellor, H. and Jefferson, L.S.: Purification and characterization of eukaryotic translational initiation factor eIF-2B from liver. *Biochim. Biophys. Acta*, 1201 (1994) 473-481.
- 1226 Koulich, D.M., Nazimov, I.V., Maltsev, K.V. and Wulfson, A.N.: Recombinant human insulin VI. Determination of recombinant human proinsulin by capillary zone electrophoresis: optimization of efficiency and selectivity. *J. Chromatogr. B*, 662 (1994) 357-362.
- 1227 Lamthanh, H., Bdolah, A., Wollberg, Z. and Gy, J.L.: Separation of two isoforms (Ser<sup>7</sup>/Thr<sup>7</sup>) of natural sarafotoxin-a by capillary electrophoresis: mass spectrometry and synthesis. *J. Liq. Chromatogr.*, 17 (1994) 3859-3867.
- 1228 Lee, H.G. and Desiderio, D.M.: Optimization of the loading limit for capillary zone electrophoresis of synthetic opioid and tachykinin peptides: a study of the interactions among the amount of peptide, resolution, saturation, injection volume and capillary diameter. *J. Chromatogr. B*, 662 (1994) 34-45.
- 1229 Lee, H.G. and Desiderio, D.M.: Preparative capillary zone electrophoresis of synthetic peptides. Conversion of an autosampler into a fraction collector. *J. Chromatogr. A*, 686 (1994) 309-317.
- 1230 Licklider, L. and Kuhr, W.G.: Optimization of on-line peptide mapping by capillary zone electrophoresis. *Anal. Chem.*, 66 (1994) 4400-4407.
- 1231 Lin, B., Chu, X., Gui, H. and Jin, J.: (Quantitative determination of gastrin with HPE-100 capillary electrophoresis). *Fenxi Ceshi Xuebao*, 12 (1993) 56-59; *C.A.*, 122 (1995) 1195s.
- 1232 McCormick, R.M.: Capillary zone electrophoresis of peptides. In: Landers, J.P. (Editor), *Handb. Capillary Electrophor.*, CRC, Boca Raton, 1994, pp. 287-323; *C.A.*, 121 (1994) 249921u - a review with 106 refs.
- 1233 Roboti, A., Livaniou, E., Evangelatos, G.P., Tsoupras, G., Tsolas, O. and Ithakissios, D.S.: Large-scale chromatofocusing-based method for isolating thymosin  $\beta_4$  and thymosin  $\beta_9$  from bovine tissues. *J. Chromatogr. B*, 662 (1994) 27-34.
- 1234 Svoboda, I., Kasicka, V., Machova, M., Jiracek, J., Prusik, Z. and Barth, T.: Separation of semisynthetic insulin analogs by capillary electrophoresis. In: Schneider, C.H. and Eberle, A.N. (Editors), *Pept. 1992, Proc. Eur. Pept. Symp.*, 2nd 1992, ESCOM, Leiden, 1993, pp. 471-472; *C.A.*, 121 (1994) 272373a.
- 1235 Takada, Y., Nakayama, K., Yoshida, M. and Sakairi, M.: Comparison of peptide analysis results for horse, bovine and dog cytochrome c using capillary electrophoresis/mass spectrometry with untreated capillary column and basic buffer condition. *Rapid Commun. Mass Spectrom.*, 8 (1994) 695-697; *C.A.*, 121 (1994) 250583s.

- 1236 Walaas, S.I., Czernik, A.J., Olsad, O.K., Sletten, K. and Walaas, O.: Protein kinase C and cyclic AMP-dependent protein kinase phosphorylate phospholemman, an insulin and adrenaline-regulated membrane phosphoprotein, at specific sites in the carboxy terminal domain. *Biochem. J.*, 304 (1994) 635-640.
  - 1237 Watts, N.R.M. and Singh, R.P.: Peptides as standards for denaturing isoelectric focusing. *Electrophoresis (Weinheim)*, 16 (1995) 22-27.
  - 1238 Witte, D.T., N  g  rd, S. and Larsson, M.: Improved sensitivity by on-line isotachopheretic preconcentration in the capillary zone electrophoretic determination of peptide-like solutes. *J. Chromatogr. A*, 687 (1994) 155-166.
  - 1239 Ye, B., Hadjmohammadi, M. and Khaledi, M.G.: Selectivity control in micellar electrokinetic chromatography of small peptides using mixed fluorocarbon-hydrocarbon anionic surfactants. *J. Chromatogr. A*, 692 (1995) 291-300.
- See also 1108, 1138, 1213, 1249, 1264.
- 18c. *Elucidation of structure of proteins and enzymes*
- 1240 Blaxter, M.L., Vanfleteren, J.R., Xia, J. and Moens, L.: Structural characterization of an *Ascaris* myoglobin. *J. Biol. Chem.*, 269 (1994) 30181-30186.
  - 1241 Braun, H.-P., Kruff, V. and Schmitz, U.K.: Primary structure, cell-free synthesis and mitochondrial targeting of the 8.2 kDa protein of cytochrome c reductase from potato. *Biochim. Biophys. Acta*, 1188 (1994) 367-372.
  - 1242 Clottes, E., Couthon, F., Denoroy, L. and Vial, C.: Creatine kinase compactness and thiol accessibility during sodium dodecyl sulfate denaturation estimated by resonance energy transfer and 2-nitro-5-thiocyanobenzoic acid cleavage. *Biochim. Biophys. Acta*, 1209 (1994) 171-176.
  - 1243 Michel, L., Garin, J., Vincon, M., Gagnon, J. and Vignais, P.: Mapping of the pyrophosphate binding sites of beef heart mitochondrial F<sub>1</sub>-ATPase by photolabelling with azidonitrophenyl [ $\alpha$ -<sup>32</sup>P]pyrophosphate. *Biochim. Biophys. Acta*, 1228 (1995) 67-72.
  - 1244 Nakaya, M.: (Microsequence analysis of proteins after electrophoresis). *Nippon Suisan Gakkaishi*, 60 (1994) 547-548; *C.A.*, 121 (1994) 249964k - a review with 18 refs.
  - 1245 Stepkowski, D., Babiychuk, E.B., Danilova, V.M. and Kakol, I.: Skeletal muscle myosin regulatory light chains conformation affects the papain cleavage of A1 light chains. *Biochim. Biophys. Acta*, 1209 (1994) 253-259.
  - 1246 Takada, Y., Nakayama, K., Yoshida, M. and Sakairi, M.: Capillary electrophoresis/mass spectrometry study of analytical conditions for tryptic digests of horse-heart cytochrome c. *Anal. Sci.*, 10 (1994) 713-717; *C.A.*, 121 (1994) 276070x.
  - 1247 Takada, Y., Yoshida, M. and Sakairi, M.: (Development of capillary electrophoresis/mass spectrometry (CE/MS) interface). *Kuromatogurafi*, 14 (1993) 146-147; *C.A.*, 121 (1994) 275626w.
  - 1248 Taylor, J.M., Jacob-Mosier, G.G., Lawton, R.G., Remmers, A.E. and Neubig, R.R.: Binding of an  $\alpha_2$  adrenergic receptor third intracellular loop peptide to G $\beta$  and the amino terminus of G $\alpha$ . *J. Biol. Chem.*, 269 (1994) 27618-27624.
  - 1249 Warlow, R.S., Gooley, A., Rajasekariah, P., Oszarac, N. and Walls, R.S.: A preparative method for sequencing proteins and peptides: *In situ* gel staining with subsequent passive elution onto polyvinylidene difluoride membranes. *Electrophoresis (Weinheim)*, 16 (1995) 84-91.
  - 1250 Wells, N.J., Addison, C.M., Fry, A.M., Ganapathi, R. and Hickson, I.D.: Serine 1524 is a major site of phosphorylation on human topoisomerase II $\alpha$  protein *in vivo* and is a substrate for casein kinase II *in vitro*. *J. Biol. Chem.*, 269 (1994) 29746-29751.
- See also 1230, 1327, 1342, 1375, 1388, 1391, 1404, 1408, 1504, 1567.
19. PROTEINS
- 19a. *General techniques*
- 1251 Boyes, B.E. and Walker, D.G.: Selectivity optimization of reversed-phase high-performance liquid chromatographic peptide and protein separations by varying bonded-phase functionality. *J. Chromatogr. A*, 691 (1995) 337-347.
  - 1252 Brownleader, M.D., Trevan, M. and Dey, P.M.: SDS-agarose gel electrophoresis as a simple procedure for determining high molecular weight protein oligomerization. *Biochem. Educ.*, 22 (1994) 155-1158; *C.A.*, 122 (1995) 9045p.
  - 1253 Chen, F.-T.A. and Tusak, A.: Characterization of food proteins by capillary electrophoresis. *J. Chromatogr. A*, 685 (1994) 331-337.
  - 1254 Dunn, M.J. and Corbett, J.M.: Two-dimensional polyacrylamide gel electrophoresis using immobilized pH gradients in the first dimension. *Methods Mol. Biol. (Totowa)*, 32(Basic Protein and Peptide Protocols) (1994) 87-96; *C.A.*, 121 (1994) 250262m.
  - 1255 Fang, X., Zhu, T. and Sun, V.: Use of zwitterionic buffer additives to improve the separation of proteins in capillary zone electrophoresis. *J. High Resolut. Chromatogr.*, 17 (1994) 749-752.
  - 1256 Fernandez-Patron, C., Calero, M., Collazo, P.R., Garcia, J.R., Madrazo, J., Musacchio, A., Soriano, F., Estrada, R., Frank, R. et al.: Protein reverse staining: high-efficiency microanalysis of unmodified proteins detected on electrophoresis gels. *Anal. Biochem.*, 224 (1995) 203-211.
  - 1257 Fernandez-Patron, C., Hardy, E., Sosa, A., Seoane, J. and Castellanos, L.: Double staining of Coomassie Blue-stained polyacrylamide gels by imidazole-sodium dodecyl sulfate-zinc reverse staining: sensitive detection of Coomassie Blue-undetected proteins. *Anal. Biochem.*, 224 (1995) 263-269.
  - 1258 Hara, T., Nishida, H. and Nakajima, R.: Highly sensitive detection of proteins separated by capillary zone electrophoresis using online chemiluminescence detection. *Anal. Sci.*, 10 (1994) 823-825; *C.A.*, 121 (1994) 250331h.
  - 1259 Huang, T.-L., Shieh, C.H., Koh, E.V. and Cooke, N.: Evaluation of a neutral hydrophilic coated capillary for capillary zone electrophoretic separation of proteins. *J. Chromatogr. A*, 685 (1994) 313-320.
  - 1260 Jen  , P., Mini, T., Moes, S., Hintermann, E. and Horst, M.: Internal sequences from proteins digested in polyacrylamide gels. *Anal. Biochem.*, 224 (1995) 75-82.

- 1261 Jie, K.-S.G., Gijsbers, B.L.M.G. and Vermeer, C.: A specific colorimetric staining method for  $\gamma$ -carboxyglutamic acid-containing proteins in polyacrylamide gels. *Anal. Biochem.*, 224 (1995) 163-165.
  - 1262 Karssen, G., van Hoenselaar, T., Verkerk-Bakker, B. and Janssen, R.: Species identification of cyst and root-knot nematodes from potato by electrophoresis of individual females. *Electrophoresis (Weinheim)*, 16 (1995) 105-109.
  - 1263 Ko, Y.-G. and Thompson, G.A., Jr.: Purification of glycosylphosphatidylinositol-anchored proteins by modified Triton X-114 partitioning and preparative gel electrophoresis. *Anal. Biochem.*, 224 (1995) 166-172.
  - 1264 Küllertz, G. and Fischer, G.: Influence of additives on resolution and focusing efficiency in free-flow isoelectric focusing. *J. Chromatogr. A*, 684 (1994) 329-341.
  - 1265 Margolis, J.: Improvements in protein separation by electrophoresis. *PCT Int. Appl. WO 94 22,904* (Cl. C07K3/14), 13 Oct. 1994, AU Appl. 93/8,221, 07 Apr. 1993; 22 pp.; C.A., 121 (1994) 296620e.
  - 1266 Marshall, T., Abbott, N.J., Fox, P. and Williams, K.M.: Protein concentration by precipitation with pyrogallol red prior to electrophoresis. *Electrophoresis (Weinheim)*, 16 (1995) 28-31.
  - 1267 Millot, M.-C., Hervé, F. and Sébille, B.: Retention behaviour of proteins on poly(vinylimidazole)-copper(II) complexes supported on silica: application to the fractionation of desialylated human  $\alpha_1$ -acid glycoprotein variants. *J. Chromatogr. B*, 664 (1995) 55-67.
  - 1268 Nakayama, H., Kanai, M., Seta, K., Isobe, T. and Okuyama, T.: (Analysis of micro-quantities of proteins by combined two-dimensional electrophoresis-capillary column HPLC/ESI/TSQMS). *Kuromatogurafi*, 15 (1994) 72-73; C.A., 121 (1994) 275701s.
  - 1269 Ngo, T.T.: Rapid purification of immunoglobulin G using azareophilic chromatography: novel mode of protein-solid phase interactions. *J. Chromatogr. B*, 662 (1994) 351-356.
  - 1270 Palmieri, R. and Nolan, J.A.: Protein capillary electrophoresis: theoretical and experimental considerations for methods development. In: Landers, J.P. (Editor), *Handb. Capillary Electrophor.*, CRC, Boca Raton, 1994, pp. 325-368; C.A., 121 (1994) 249922v - a review with 198 refs.
  - 1271 Paulson, J.R. and Moore, R.E.: Characterization of proteins in coated papers by sodium dodecyl sulphate (SDS)-polyacrylamide gel electrophoresis. *Tappi J.*, 76 (1993) 102-104; C.A., 121 (1994) 233306m.
  - 1272 Pollard, J.W.: The *in vivo* isotopic labelling of proteins for polyacrylamide gel electrophoresis. *Methods Mol. Biol. (Totowa)*, 32(Basic Protein and Peptide Protocols) (1994) 67-72; C.A., 121 (1994) 250260j.
  - 1273 Pollard, J.W.: Two-dimensional polyacrylamide gel electrophoresis of proteins. *Methods Mol. Biol. (Totowa)*, 32(Basic Protein and Peptide Protocols) (1994) 73-85; C.A., 121 (1994) 250261k.
  - 1274 Pot, B., Vandamme, P. and Kersters, K.: Analysis of electrophoretic whole-organism protein fingerprints. In: Goodfellow, M. and O'Donnell, A.G. (Editors), *Chem. Methods Prokaryotic Syst.*, Wiley, Chichester, 1994, pp. 493-521; C.A., 121 (1994) 275882b.
  - 1275 Pyun, J.-c., Choi, H.-s. and Park, J.-s.: Stopped lower buffer flow electrofractionation: simple electrofractionation for complex protein mixture. *Anal. Biochem.*, 223 (1994) 59-61.
  - 1276 Samuel, M.: "Phast 2D", a two-dimensional electrophoretic technique on a single gel under native and denaturing conditions using Pharmacia Phast System. *Anal. Biochem.*, 224 (1995) 457-459.
  - 1277 Sanchez, R.I., Darsley, M.J. and Martin, M.T.: Spectrophotometric quantitation of silver in X-ray film images and silver-stained electrophoresis gel bands. *Anal. Biochem.*, 224 (1995) 190-194.
  - 1278 Simo-Alfonso, E., Conti, M., Gelfi, C. and Righetti, P.G.: Sodium dodecyl sulfate capillary electrophoresis of proteins in entangled solutions of poly(vinyl alcohol). *J. Chromatogr. A*, 689 (1995) 85-96.
  - 1279 Smith, B.J.: SDS polyacrylamide gel electrophoresis of proteins. *Methods Mol. Biol. (Totowa)*, 32(Basic Protein and Peptide Protocols) (1994) 23-34; C.A., 121 (1994) 250255m.
  - 1280 Su, C., Wang, F., Ciolek, D. and Pan, Y.-C.E.: Electrophoresis of proteins and protein-protein complexes in native polyacrylamide gels using a horizontal gel apparatus. *Anal. Biochem.*, 223 (1994) 93-98.
  - 1281 Swedberg, S.A.: The impact of column technology on protein separations by open tubular capillary electrophoresis: past lessons, future promises. In: Landers, J.P. (Editor), *Handb. Capillary Electrophor.*, CRC, Boca Raton, 1994, pp. 513-536; C.A., 121 (1994) 249926z - a review with 32 refs.
  - 1282 Tsuji, K.: Sodium dodecyl sulfate polyacrylamide gel- and replaceable polymer-filled capillary electrophoresis for molecular mass determination of proteins of pharmaceutical interest. *J. Chromatogr. B*, 662 (1994) 291-299.
  - 1283 Walker, J.M.: Gradient SDS polyacrylamide gel electrophoresis of proteins. *Methods Mol. Biol. (Totowa)*, 32(Basic Protein and Peptide Protocols) (1994) 35-38; C.A., 121 (1994) 250256n.
  - 1284 Walker, J.M.: Nondenaturing polyacrylamide gel electrophoresis of proteins. *Methods Mol. Biol. (Totowa)*, 32(Basic Protein and Peptide Protocols) (1994) 17-22; C.A., 121 (1994) 250254k.
  - 1285 Waterborg, J.H. and Matthews, H.R.: The electrophoretic elution of proteins from polyacrylamide gels. *Methods Mol. Biol. (Totowa)*, 32(Basic Protein and Peptide Protocols) (1994) 169-175; C.A., 121 (1994) 250267s.
  - 1286 Wu, D.: Novel approaches to the separation and quantification of proteins by capillary electrophoresis. Avail. *Univ. Microfilms Int.*, Order No. DA9420922, 1993, 169 pp.; C.A., 121 (1994) 275899n.
  - 1287 Yukui, Z., Chen, N., Wang, L., Liren, M. and Xiaohong, Q.: (Separation of some standard proteins by capillary zone electrophoresis and the study on their migration behavior). *Fenxi Huaxue*, 22 (1994) 798-800; C.A., 121 (1994) 250338r.
- See also 1076, 1077, 1083, 1084, 1091, 1104, 1113, 1133, 1136, 1159, 1208.
- 19b. *Proteins of cells, viruses and subcellular particles*
- 1288 Brennan, P. and O'Neill, L.A.J.: Effects of oxidants and antioxidants on nuclear factor  $\kappa$ B activation in three different cell lines: evidence against a universal hypothesis involving oxygen radicals. *Biochim. Biophys. Acta*, 1260 (1995) 167-175.

- 1289 Clark, B.J., Wells, J., King, S.R. and Stocco, D.M.: The purification, cloning, and expression of a novel luteinizing hormone-induced mitochondrial protein in MA-10 mouse Leydig tumor cells. Characterization of the steroidogenic acute regulatory protein (StAR). *J. Biol. Chem.*, 269 (1994) 28314-28322.
- 1290 De Vienne, D., Maurice, A., Josse, J.-M., Leonardi, A. and Darnerval, C.: Mapping factors controlling genetic expression. *Cell. Mol. Biol. (Paris)*, 40 (1994) 29-39; *C.A.*, 121 (1994) 273122e.
- 1291 Desjardins, M., Celis, J.E., van Meer, G., Dieplinger, H., Jahraus, A., Griffiths, G. and Huber, L.A.: Molecular characterization of phagosomes. *J. Biol. Chem.*, 269 (1994) 32194-32200.
- 1292 Faraone-Mennella, M.R., de Lucia, F., de Maio, A., Gambacorta, A., Quesada, P., de Rosa, M., Nicolaus, B. and Farina, B.: ADP-ribosylation reactions in *Sulfolobus solfataricus*, a thermoacidophilic archaeon. *Biochim. Biophys. Acta*, 1246 (1995) 151-159.
- 1293 Fransen, M., Brees, C.A.E., de Béthune, B., van Veldhoven, P.P. and Mannaerts, G.P.: The 80 kDa cytosolic protein that binds the C-terminal part of rat acyl-CoA oxidase is not a peroxisomal import receptor but a prolyl-endoropeptidase. *Biochim. Biophys. Acta*, 1201 (1994) 157-164.
- 1294 Huckriede, A. and Agsteribbe, E.: Decreased synthesis and inefficient mitochondrial import of hsp60 in a patient with a mitochondrial encephalomyopathy. *Biochim. Biophys. Acta*, 1227 (1994) 200-206.
- 1295 Iljst, L., Wanders, R.J.A., Ushikubo, S., Kamijo, T. and Hashimoto, T.: Molecular basis of long-chain 3-hydroxyacyl-CoA dehydrogenase deficiency: identification of the major disease-causing mutation in the  $\alpha$ -subunit of the mitochondrial trifunctional protein. *Biochim. Biophys. Acta*, 1215 (1994) 347-350.
- 1296 Janssen, G.M.C., van Damme, H.T.F., Kriek, J., Amons, R. and Möller, W.: The subunit structure of elongation factor 1 from *Artemia*. Why two  $\alpha$ -chains in this complex? *J. Biol. Chem.*, 269 (1994) 31410-31417.
- 1297 Jordan, P., Heid, H., Kinzel, V. and Kübler, D.: Major cell surface-located protein substrates of an ecto protein kinase are homologs of known nuclear proteins. *Biochemistry*, 33 (1994) 14696-14706.
- 1298 Kaul, S.C., Wadhwa, R., Sugihara, T., Obuchi, K., Komatsu, Y. and Mitsui, Y.: Identification of genetic events involved in early steps of immortalization of mouse fibroblasts. *Biochim. Biophys. Acta*, 1201 (1994) 389-396.
- 1299 Kikuchi, H., Fujinawa, T., Kuribayashi, F., Nakanishi, A., Imajoh-Ohmi, S., Goto, M. and Kanegasaki, S.: Induction of essential components of the superoxide generating system in human monoclastic leukemia U937 cells. *J. Biochem. (Tokyo)*, 116 (1994) 742-746.
- 1300 Kobayashi, T., Urabe, K., Orlow, S.J., Higashi, K., Imokawa, G., Kwon, B.S., Potterf, B. and Hearing, V.J.: The Pmel 17/silver locus protein. Characterization and investigation of its melanogenic function. *J. Biol. Chem.*, 269 (1994) 29198-29205.
- 1301 Liu, L. and Chander, A.: Stilbene disulfonic acids inhibit synexin-mediated membrane aggregation and fusion. *Biochim. Biophys. Acta*, 1254 (1995) 274-282.
- 1302 McKee, E.E.: Mitochondrial gene expression in *Saccharomyces cerevisiae*. IV Effects of yeast cytosol on mitochondrial protein synthesis, degradation, and respiration. *Biochim. Biophys. Acta*, 1201 (1994) 235-244.
- 1303 Mentel, R., Meinsen, D., Pilgrim, H., Herrmann, B. and Lindequist, U.: *In vitro* antiviral effect of extracts of *Kuehneromyces mutabilis* on influenza virus. *Pharmazie*, 49 (1994) 859-860.
- 1304 Nakamura, K., Koga, Y., Yoshida, H., Tanaka, K., Sasaki, M., Kimura, G. and Nomoto, K.: Inhibition of the T-cell receptor-mediated signal transduction by microinjection of anti-Lek monoclonal antibody into T-cell. *Biochim. Biophys. Acta*, 1224 (1994) 495-505.
- 1305 Nilsson, A. and Nygard, O.: Effect of oxidizing agents and haemin on the phosphorylation of eukaryotic elongation factor 2 in rabbit reticulocyte lysates. *Biochim. Biophys. Acta*, 1260 (1995) 200-206.
- 1306 Ogawa, W. and Roth, R.A.: Characterization of a protein which binds phosphatidylinositol 3,4,5-trisphosphate and 4,5-bisphosphate. *Biochim. Biophys. Acta*, 1224 (1994) 533-540.
- 1307 Sakai, R., Iwamatsu, A., Hirano, N., Ogawa, S., Tanaka, T., Nishida, J., Yazaki, Y. and Hirai, H.: Characterization, partial purification, and peptide sequencing of p130, the main phosphoprotein associated with v-Crk oncoprotein. *J. Biol. Chem.*, 269 (1994) 32740-32746.
- 1308 Santos, M.J., Kawada, M.E., Espeel, M., Figueroa, C., Alvarez, A., Hidalgo, U. and Metz, C.: Characterization of human peroxisomal membrane proteins. *J. Biol. Chem.*, 269 (1994) 24890-24896.
- 1309 Schaeffer, H.: Electrophoretic isolation of membrane proteins from acrylamide gels. *Appl. Biochem. Biotechnol.*, 48 (1994) 185-203; *C.A.*, 121 (1994) 250328n.
- 1310 Schwaner, I., Offermanns, S., Spicher, K., Seifert, R. and Schultz, G.: Differential activation of G<sub>i</sub> and G<sub>s</sub> proteins by E- and I-type prostaglandins in membranes from the human erythroleukemia cell line, HEL. *Biochim. Biophys. Acta*, 1265 (1995) 8-14.
- 1311 Sugihara, T., Kaul, S., Mitsui, Y. and Wadhwa, R.: Enhanced expression of multiple forms of VEGF is associated with spontaneous immortalization of murine fibroblasts. *Biochim. Biophys. Acta*, 1224 (1994) 365-370.
- 1312 Temesvari, L., Rodriguez-Paris, J., Bush, J., Steck, T.L. and Cardelli, J.: Characterization of lysosomal membrane proteins of *Dictyostelium discoideum*. A complex population of acidic integral membrane glycoproteins, Rab GTP-binding proteins and vacuolar ATPase subunits. *J. Biol. Chem.*, 269 (1994) 25719-25727.
- 1313 Thelin, A., Peterson, E., Hutson, J.L., McCarthy, A.D., Ericsson, J. and Dallner, G.: Effect of squalstatin 1 on the biosynthesis of the mevalonate pathway lipids. *Biochim. Biophys. Acta*, 1215 (1994) 245-249.
- 1314 Wang, S.-L., Tam, M.-F., Ho, Y.-S., Pai, S.-H. and Kao, M.-C.: Isolation and molecular cloning of human sorcin a calcium-binding protein in vincristine-resistant HOB1 lymphoma cells. *Biochim. Biophys. Acta*, 1260 (1995) 285-293.
- 1315 Webb, R. and Dormer, R.L.: Photoaffinity labelling of the ATP-binding sites of two Ca<sup>2+</sup>, Mg-ATPase isoforms in pancreatic endoplasmic reticulum. *Biochim. Biophys. Acta*, 1233 (1995) 1-6.

See also 1324, 1334, 1342, 1355, 1371, 1548, 1627, 1749.

- 19c. *Proteins synthesized by genetic manipulation, monoclonal antibodies*
- 1316 Björquist, P., Brohlin, M., Ehnebom, J., Ericsson, M., Kristiansen, C., Pohl, G. and Deinum, J.: Plasminogen activator inhibitor type-1 interacts exclusively with the proteinase domain of tissue plasminogen activator. *Biochim. Biophys. Acta*, 1209 (1994) 191-202.
- 1317 Chibazakura, T., Kitajima, S., Yonaha, M. and Yasukochi, Y.: Enhancement of bacterial transcription initiation *in vitro* by the 74 kDa subunit of human general transcription factor IIF (RAP74). *Biochim. Biophys. Acta*, 1219 (1994) 592-600.
- 1318 Claffey, K.P., Senger, D.R. and Spiegelman, B.M.: Structural requirements for dimerization, glycosylation, secretion, and biological function of VPF/VEGF. *Biochim. Biophys. Acta*, 1246 (1995) 1-9.
- 1319 Gorin, M.B., To, A.C. and Narfstrom, K.: Sequence analysis and exclusion of phosducin as the gene for the recessive retinal degeneration of the abyssinian cat. *Biochim. Biophys. Acta*, 1260 (1995) 323-327.
- 1320 Hagendorff, G., Hanes, J., von der Kammer, H. and Scheit, K.H.: A monoclonal antibody generated against a recombinant peptide fragment of the B3 domain of carcinoembryonic antigen reacts with intact carcinoembryonic antigen. *Biochim. Biophys. Acta*, 1260 (1995) 259-268.
- 1321 Hensley, P., McDevitt, P.J., Brooks, I., Trill, J.J., Feild, J.A., McNulty, D.E., Connor, J.R., Griswold, D.E., Kumar, N.V., Kopple, K.D., Carr, S.A. et al.: The soluble form of E-selectin is an asymmetric monomer. Expression, purification, and characterization of the recombinant protein. *J. Biol. Chem.*, 269 (1994) 23949-23958.
- 1322 Illarionov, B., Illarionova, V., Lee, J., van Dongen, W. and Vervoort, J.: Expression and properties of the recombinant lumazine (riboflavin) protein from *Photobacterium leiognathi*. *Biochim. Biophys. Acta*, 1201 (1994) 251-258.
- 1323 Kataoka, K., Takada, H., Tanizawa, K., Yoshimura, T., Esaki, N., Ohshima, T. and Soda, K.: Construction and characterization of chimeric enzyme consisting of an amino-terminal domain of phenylalanine dehydrogenase and a carboxy-terminal domain of leucine dehydrogenase. *J. Biochem. (Tokyo)*, 116 (1994) 931-936.
- 1324 Nikitina, V.E., Alenkina, S.A., Italyanskaya, Yu.V. and Ponomareva, E.G.: (Purification and comparison of lectins from the cell surface of active and inactive in hemagglutination azospirilla). *Biokhimiya (Moscow)*, 59 (1994) 656-662.
- 1325 Nishihira, J., Kuriyama, T., Sakai, M., Nishi, S., Ohki, S.-y. and Hikichi, K.: The structure and physicochemical properties of rat liver macrophage migration inhibitory factor. *Biochim. Biophys. Acta*, 1247 (1995) 159-162.
- 1326 Schmalzing, D., Nashabeh, W., Yao, X.-W., Mhatre, R., Regnier, F.E., Afeyan, N.B. and Fuchs, M.: Capillary electrophoresis-based immunoassay for cortisol in serum. *Anal. Chem.*, 67 (1995) 606-612.
- 1327 Van Nostrand, W.E., Schmaier, A.H., Neiditch, B.R., Siegel, R.S., Raschke, W.C., Sisodia, S.S. and Wagner, S.L.: Expression, purification, and characterization of the Kunitz-type proteinase inhibitor domain of the amyloid  $\beta$ -protein precursor-like protein-2. *Biochim. Biophys. Acta*, 1209 (1994) 165-170.
- 1328 Whelan, A., Russell, W.P., Nawaz, F., Newell, D.G. and Hewinson, R.G.: Purification and characterization of recombinant MPB70. *Spec. Publ. - R. Soc. Chem.*, 158 (Separations for Biotechnology 3) (1994) 559-564; C.A., 121 (1994) 296411n.
- See also 1226, 1282, 1302, 1349, 1353, 1361, 1369, 1417, 1456, 1471, 1504, 1522, 1536, 1563, 1597, 1723.
- 19d. *Microbial and plant proteins*
- 1329 Cash, P., Argo, E. and Bruce, K.D.: Characterisation of *Haemophilus influenzae* proteins by two-dimensional gel electrophoresis. *Electrophoresis (Weinheim)*, 16 (1995) 135-148.
- 1330 Dupuy, N., Willems, A., Pot, B., Dewettinck, D., Vandenbruaene, I., Maestrojuan, G., Dreyfus, B., Kersters, K., Collins, M.D. et al.: Phenotypic and genotypic characterization of *Bradyrhizobia* nodulating the leguminous tree *Acacia albida*. *Int. J. Syst. Bacteriol.*, 44 (1994) 461-473; C.A., 122 (1995) 2350a.
- 1331 Fairchild, C.D. and Glazer, A.N.: Nonenzymatic bilin addition to the  $\alpha$  subunit of an apophycocerythrin. *J. Biol. Chem.*, 269 (1994) 28988-28996.
- 1332 Friis, S., Anthonsen, D., Noren, O. and Sjöström, H.: Gamma-type gliadis cause secretion of prostaglandin  $E_2$  in patients with coeliac disease. *Clin. Chim. Acta*, 231 (1994) 173-183.
- 1333 Gans, P. and Wollman, F.-A.: The effect of cyanide on state transitions in *Chlamydomonas reinhardtii*. *Biochim. Biophys. Acta*, 1228 (1995) 51-57.
- 1334 Georgakopoulos, J.H. and Argyroudi-Akoyunoglou, J.H.: On the question of lateral migration of LHC II upon thylakoid protein phosphorylation in isolated pea chloroplasts: the stroma lamellar fraction separated from phosphorylated chloroplasts is not homogeneous. *Biochim. Biophys. Acta*, 1188 (1994) 380-390.
- 1335 Gong, H.: Light-dependent degradation of the Photosystem II D1 protein is retarded by inhibitors of chloroplast transcription and translation: possible involvement of a chloroplast-encoded proteinase. *Biochim. Biophys. Acta*, 1188 (1994) 422-426.
- 1336 Hack, B., Hanstein, S., Schuetz, B., Zimmermann, K. and Rusch, V.: (Computer-controlled analysis of electrophoretic protein patterns in the differentiation of lactic acid bacteria according to species and subspecies). *Lebensmittelind. Milchwirtsch.*, 114 (1993) 880-883; C.A., 121 (1994) 299387v.
- 1337 Hansen, C.E., Flengsrud, R. and Kopperud, C.: Two-dimensional gel electrophoresis of proteins in healthy roots and clubroots of *Brassica oleracea*. *Acta Agric. Scand. Sect. B*, 44 (1994) 123-128; C.A., 121 (1994) 226598e.
- 1338 Henkel, A.W. and Bieger, S.C.: Quantification of proteins dissolved in an electrophoresis sample buffer. *Anal. Biochem.*, 223 (1994) 329-331.
- 1339 Kaji, D.A., Rosato, Y.B., Canhos, V.P. and Priest, F.G.: Characterization by polyacrylamide gel electrophoresis of whole-cell proteins of some *Bacillus thuringiensis* subsp. *Israelensis* strains isolated in Brazil. *Syst. Appl. Microbiol.*, 17 (1994) 104-107; C.A., 121 (1994) 225607v.
- 1340 Kalchgruber, R., Eder, R. and Barna, J.: (Cultivar identification of grape by electrophoretic separation of isoenzymes and proteins as demonstrated with the cultivars Chardonnay and White Burgundy). *Mitt. Klosterneuburg*, 44 (1994) 14-23; C.A., 121 (1994) 229407c.

- 1341 Kersters, K., Pot, B., Dewettinck, D., Torck, U. and Vauterin, L.: Protein electrophoretic patterns for high-resolution identification of micro-organisms. *Prog. Biotechnol.*, 9(ECB6: Proceedings of the 6th European Congress on Biotechnology, 1993, Pt. 1) (1994) 367-370; C.A., 121 (1994) 250043r.
  - 1342 Kim, S., Pichersky, E. and Yocum, C.F.: Topological studies of spinach 22 kDa protein of Photosystem II. *Biochim. Biophys. Acta*, 1188 (1994) 339-348.
  - 1343 Matsumoto, M. and Ishikawa, F.: Purification of *Treponema pallidum*, Nichols strain, by two-step column chromatography. *J. Chromatogr. B*, 663 (1995) 217-224.
  - 1344 Matsuo, T., Graham, D., Patterson, B.D. and Hockley, D.B.: An electrophoretic method to detect cold-induced dissociation of proteins in crude extracts of higher plants. *Anal. Biochem.*, 223 (1994) 181-184.
  - 1345 Moutete, H.F., Olszewski, A., Gastin, I., Namour, F., Moneret-Vautrin, D.A. and Guéant, J.L.: Purification of allergenic proteins from peanut for preparation of the reactive solid phase of a specific IgE radioimmunoassay. *J. Chromatogr. B*, 664 (1995) 211-217.
  - 1346 Nakamura, A.: Electrophoresis method for selecting hybrid rice seed. *Jpn. Kokai Tokkyo Koho JP 06,201,685* [94,201,685] (Cl. GO1N33/48), 22 Jul. 1994, Appl. 92/235,151, 12 Aut. 1992; 5 pp.; C.A., 121 (1994) 276179q.
  - 1347 Noguchi, S., Yamazaki, T., Yaginuma, A., Sakamoto, J. and Sone, N.: Over-expression of membrane-bound cytochrome c-551 from thermophilic *Bacillus* PS3 in *Bacillus stearothermophilus* K1041. *Biochim. Biophys. Acta*, 1188 (1994) 302-310.
  - 1348 Norbeck, J. and Blomberg, A.: Gene linkage of two-dimensional polyacrylamide gel electrophoresis resolved proteins from isogene families in *Saccharomyces cerevisiae* by microsequencing of in-gel trypsin generated peptides. *Electrophoresis (Weinheim)*, 16 (1995) 149-156.
  - 1349 Pearson, D.M., O'Reilly, C., Colby, J. and Black, G.W.: DNA sequence of the cut A, B and C genes, encoding the molybdenum containing hydroxylase carbon monoxide dehydrogenase, from *Pseudomonas thermocarboxydovorans* strain C2. *Biochim. Biophys. Acta*, 1188 (1994) 432-438.
  - 1350 Rocher, A., Soriano, F., Molina, E., González-Limas, G. and Méndez, E.: Characterization of distinct  $\alpha$ - and  $\gamma$ -type gliadins and low molecular weight components from wheat endosperm as coeliac immunoreactive proteins. *Biochim. Biophys. Acta*, 1247 (1995) 143-148.
  - 1351 Shevchenko, N.A., Shvedova, O.E., Pilipchuk, B.Z. and Semikhodsky, A.G.: (Use of electrophoresis of prolamines for the analysis of ryegrass-fescue hybrids). *Fiziol. Biokhim. Kul't. Rast.*, 25 (1993) 479-484; C.A., 121 (1994) 226521z.
  - 1352 Shih, F.F.: Interaction of soy isolate with polysaccharide and its effect on film properties. *J. Am. Oil Chem. Soc.*, 71 (1994) 1281-1285.
  - 1353 Takamatsu, H., Nakane, A., Oguro, A., Sadaie, Y., Nakamura, K. and Yamane, K.: A truncated *Bacillus subtilis* SecA protein consisting of the N-terminal 234 amino acid residues forms a complex with *Escherichia coli* SecA51(ts) protein and complements the protein translocation defect of the secA51 mutant. *J. Biochem. (Tokyo)*, 116 (1994) 1287-1294.
  - 1354 Van den Berg, B.M. and Gabillard, D.: Isoelectric focusing in immobilized pH gradient of melon (*Cucumis melo* L.) seed protein: methodical and genetic aspects, and application in breeding. *Electrophoresis (Weinheim)*, 15 (1994) 1541-1551.
  - 1355 Wolfe, G.R., Cunningham, F.X., Jr., Grabowski, B. and Gantt, E.: Isolation and characterization of Photosystems I and II from the red alga *Porphyridium cruentum*. *Biochim. Biophys. Acta*, 1188 (1994) 357-366.
  - 1356 Zakimi, S., Tsuji, N. and Fujisaki, K.: Protein analysis of *Anaplasma marginale* and *Anaplasma centrale* by two-dimensional polyacrylamide gel electrophoresis. *J. Vet. Med. Sci.*, 56 (1994) 1025-1027; C.A., 122 (1995) 4744f.
- See also 1158, 1465, 1466.
- 19e. *Proteins of blood, serum and blood cells*
- 1357 Altmüller, A. and Presek, P.: Rapid protein tyrosine phosphorylation in the cytoskeleton of stimulated human platelets. *Biochim. Biophys. Acta*, 1265 (1995) 61-66.
  - 1358 Bean, P. and Peter, J.B.: Allelic D variants of transferrin in evaluation of alcohol abuse: differential diagnosis by isoelectric focusing-immunoblotting-laser densitometry. *Clin. Chem. (Washington)*, 40 (1994) 2078-2083.
  - 1359 Beppu, M., Inoue, M., Ishikawa, T. and Kikugawa, K.: Presence of membrane-bound proteinases that preferentially degrade oxidatively damaged erythrocyte membrane proteins as secondary antioxidant defense. *Biochim. Biophys. Acta*, 1196 (1994) 81-87.
  - 1360 Borkow, G., Gutierrez, J.M. and Ovadia, M.: A potent antihemorrhagin in the serum of the non-poisonous water snake *Matrix tessellata*: isolation, characterization and mechanism of neutralization. *Biochim. Biophys. Acta*, 1201 (1994) 482-490.
  - 1361 Büntemeyer, H., Tebbe, H., Lüttemeyer, D. and Lehmann, J.: Rapid high-performance liquid chromatographic quantification of recombinant human antithrombin III during production and purification. *J. Chromatogr. B*, 662 (1994) 209-216.
  - 1362 Cao, M., He, Y., Lian, S. and Li, C.: (Miniature two-dimensional polyacrylamide gel electrophoresis of fish serum proteins). *Huan Shifan Daxue Ziran Kexue Xuebao*, 17 (1994) 80-83; C.A., 121 (1994) 250324h.
  - 1363 Chang, Chi., Hüslér, T., Zhao, J., Wiedmer, T. and Sims, P.J.: Identity of a peptide domain of human C9 that is bound by the cell-surface complement inhibitor, CD59. *J. Biol. Chem.*, 269 (1994) 26424-26430.
  - 1364 Dahri, L., Voisson-Vidal, C., Regnault, V., Muller, D., Sultan, Y. and Stoltz, J.F.: Synthetic sorbents for removal of factor VIII inhibitors from haemophilic A plasma. *J. Chromatogr. B*, 664 (1995) 47-54.
  - 1365 Finger, U.B., Thömmes, J., Kinzelt, D. and Kula, M.-R.: Application of thiophilic membranes for the purification of monoclonal antibodies from cell culture media. *J. Chromatogr. B*, 664 (1995) 69-78.
  - 1366 Harrison, T.M., Hudson, K., Munson, S.E., Uff, S. and Glassford, S.: Derivation and partial analysis of two highly active myeloma cell transfectants. *Biochim. Biophys. Acta*, 1260 (1995) 147-156.
  - 1367 Hatanaka, M., Seya, T., Inai, S. and Shimizu, A.: The functions of the ninth component of human complement are sustained by disulfide bonds with different susceptibilities to reduction. *Biochim. Biophys. Acta*, 1209 (1994) 117-122.

- 1368 Josic, D., Schwinn, H., Stadler, M. and Strancar, A.: Purification of factor VIII and von Willebrand factor from human plasma by anion-exchange chromatography. *J. Chromatogr. B*, 662 (1994) 181-190.
  - 1369 Kaplan, D.R.: Delivery of interleukin 2 for immunotherapy. *J. Chromatogr. B*, 662 (1994) 315-323 - a review with 73 refs.
  - 1370 Kyle, R.A.: The monoclonal gammopathies. *Clin. Chem. (Washington)*, 40 (1994) 2154-2161 - a review with 31 refs.
  - 1371 Lukashova, I.V., Kovalev, L.I., Efimochkin, A.S. and Shishkin, S.S.: (Study of human lymphocyte proteins by two-dimensional gel electrophoresis). *Biokhimiya (Moscow)*, 59 (1994) 663-674.
  - 1372 Lutomski, D., Joubert-Caron, R., Bourin, P., Bladier, D. and Caron, M.: Use of thiophilic adsorption in the purification of biotinylated Fab fragments. *J. Chromatogr. B*, 664 (1995) 79-82.
  - 1373 Matveevskaya, N.S., Alyoshkin, V.A. and Rozina, M.N.: Isolation of the C3 complement component and its C3d subunit from IY-1 fraction of Cohn's fractionation of human plasma. *J. Chromatogr. B*, 664 (1995) 261-266.
  - 1374 Millard, C.B., Meier, H.L. and Broomfield, C.A.: Exposure of human lymphocytes to bis-(2-chloroethyl)sulfide solubilizes truncated and intact core histones. *Biochim. Biophys. Acta*, 1224 (1994) 389-394.
  - 1375 Mimura, Y., Kabat, E.A., Tanaka, T., Fujimoto, M., Takeo, K. and Nakamura, K.: Microheterogeneity of mouse antidextran monoclonal antibodies. *Electrophoresis (Weinheim)*, 16 (1995) 116-123.
  - 1376 Myara, I., Asso, M., Tenenhaus, D. and Moatti, N.: Usefulness of the kappa/lambda ratio for detecting a new monoclonal component in a patient with a monoclonal gammopathy. *Clin. Chim. Acta*, 231 (1994) 115-116.
  - 1377 Nagai, K., Inazu, T. and Yamamura, H.: p72<sup>syk</sup> is activated by vanadate plus H<sub>2</sub>O<sub>2</sub> in porcine platelets and phosphorylates GTPase activating protein on tyrosine residue(s). *J. Biochem. (Tokyo)*, 116 (1994) 1176-1181.
  - 1378 Orlandi, M., Bartolini, G., Belletti, B., Spisni, E. and Tomasi, V.: Thromboxane A<sub>2</sub> synthase activity in platelet free human monocytes. *Biochim. Biophys. Acta*, 1215 (1994) 285-290.
  - 1379 Oxvig, C., Sand, O., Kristensen, T., Kristensen, L. and Sottrup-Jensen, L.: Isolation and characterization of circulating complex between human pregnancy-associated plasma protein-A and proform of eosinophil major basic protein. *Biochim. Biophys. Acta*, 1201 (1994) 415-423.
  - 1380 Reif, O.W., Lausch, R. and Freitag, R.: Application of CE to the quantitative and qualitative analysis of serum proteins. *Am. Lab. (Shelton)*, 26 (1994) 33-37; C.A., 121 (1994) 225375t.
  - 1381 Reif, O.W., Lausch, R. and Freitag, R.: High-performance capillary electrophoresis of human serum and plasma proteins. *Adv. Chromatogr. (N.Y.)*, 34 (1994) 1-56; C.A., 121 (1994) 296200t.
  - 1382 Schwarz, A., Kohen, F. and Wilchek, M.: Novel heterocyclic ligands for the thiophilic purification of antibodies. *J. Chromatogr. B*, 664 (1995) 83-88.
  - 1383 Shi, J.Y. and Goffe, R.A.: Comprehensive study on binding capacity of human immunoglobulin G to Avid AL affinity gel. *J. Chromatogr. A*, 686 (1994) 61-71.
  - 1384 Su, L., Keren, D.F. and Warren, J.S.: Failure of anti-lambda immunofixation reagent mimics alpha heavy-chain disease. *Clin. Chem. (Washington)*, 41 (1995) 121-123. C.A.,
  - 1385 Van Noort, W.L., de Jong, G. and van Eijk, H.G.: Purification of isotransferrins by concanavalin A Sepharose chromatography and preparative isoelectric focusing. *Eur. J. Clin. Chem. Clin. Biochem.*, 32 (1994) 885-892.
  - 1386 Wang, X., Sada, K., Yanagi, S., Yang, C., Rezaul, K. and Yamamura, H.: Intracellular calcium dependent activation of p72<sup>syk</sup> in platelets. *J. Biochem. (Tokyo)*, 116 (1994) 858-861.
  - 1387 Wätzig, H. and Lloyd, D.K.: Effect of pH and sodium dodecyl sulfate concentration on the analytical window in the direct-injection analysis of plasma samples by capillary electrophoresis. *Electrophoresis (Weinheim)*, 16 (1995) 57-63.
  - 1388 Yano, S., Suzuki, K.-i., Katoh, M., Sugita, Y., Kaku, S., Kawamura, K. and Masuho, Y.: Epitopes and biological activities of two monoclonal antibodies to platelet integrin  $\alpha_{11b}\beta_3$ . *J. Biochem. (Tokyo)*, 116 (1994) 778-786.
- See also 1269, 1413, 1416, 1561, 1793.
- 19f. *Structural and muscle proteins*
- 1389 Arai, K., Terasawa, R., Yokote, Y., Takahashi, R. and Akahane, K.: (Pattern analysis of avian feather keratins by high performance capillary electrophoresis). *Kuromatogurafi*, 15 (1994) 161-167; C.A., 122 (1995) 4746h.
  - 1390 Banks, P.R. and Paquette, D.M.: Monitoring of a conjugation reaction between fluorescein isothiocyanate and myoglobin by capillary zone electrophoresis. *J. Chromatogr. A*, 693 (1995) 145-154.
  - 1391 Guo, X.-F., Nakaya, M. and Watabe, S.: Myosin subfragment-1 isoforms having different heavy chain structures from fast skeletal muscle of thermally acclimated carp. *J. Biochem. (Tokyo)*, 116 (1994) 728-735.
  - 1392 Hill, P.A., Reynolds, J.J. and Meikle, M.C.: The purification and partial characterization of bone resorptive polypeptides from bovine bone matrix. *Biochim. Biophys. Acta*, 1201 (1994) 193-202.
  - 1393 Janmot, C. and d'Albis, A.: Electrophoretic separation of developmental and adult rabbit skeletal muscle myosin heavy chain isoforms: example of application to muscle denervation study. *FEBS Lett.*, 353 (1994) 13-15; C.A., 121 (1994) 275908q.
  - 1394 Khawar, S.L., Watson, K. and Jones, G.L.: High resolution one-dimensional electrophoretic separation and partial characterisation of human head hair proteins. *Electrophoresis (Weinheim)*, 16 (1995) 110-115.
  - 1395 Kovaleva, M.A., Kovalev, L.I., Khudaidatov, A.I., Efimochkin, A.S. and Shishkin, S.S.: (A comparative analysis of protein composition of human skeletal and heart muscle by two-dimensional electrophoresis). *Biokhimiya (Moscow)*, 59 (1994) 675-681.
  - 1396 Kvannes, J., Eikhom, T.S. and Flatmark, T.: The peroxisomal  $\beta$ -oxidation enzyme system of rat heart. Basal level and effect of the peroxisome proliferator clofibrate. *Biochim. Biophys. Acta*, 1201 (1994) 203-206.
  - 1397 Mizuno, K. and Hayashi, T.: Peculiar effect of urea on the interaction of type I collagen with heparin on chromatography. *J. Biochem. (Tokyo)*, 116 (1994) 1257-1263.
  - 1398 Murayama, T. and Ogawa, Y.: Relationships among ryanodine receptor isoforms expressed in vertebrate skeletal muscles based on immunologic cross-reactivities. *J. Biochem. (Tokyo)*, 116 (1994) 1117-1122.

- 1399 Ohashi, K., Nishimura, M., Terasaki, A.G. and Nakagawa, H.: A 36-kDa protein of the dense bodies of smooth muscle cells. *J. Biochem. (Tokyo)*, 116 (1994) 1354-1359.
- 1400 Pirner, M.A. and Linck, R.W.: Tektins are heterodimeric polymers in flagellar microtubules with axial periodicities matching the tubulin lattice. *J. Biol. Chem.*, 269 (1994) 31800-31806.
- 1401 Rossini, K., Rizzi, C., Sandri, M., Bruson, A. and Carraro, U.: High-resolution sodium dodecyl sulfate-polyacrylamide gel electrophoresis and immunochemical identification of the 2X and embryonic myosin heavy chains in complex mixtures of isomyosins. *Electrophoresis (Weinheim)*, 16 (1995) 101-104.
- 1402 Sato, K., Taira, T., Takayama, R., Ohtsuki, K. and Kawabata, M.: Improved chromatographic purification of human and bovine type V collagen sub-molecular species and their subunit chains from conventional crude preparations. Application to cell-substratum adhesion assay for human umbilical vein endothelial cell. *J. Chromatogr. B*, 663 (1995) 25-33.
- 1403 Tatsumi, R. and Hattori, A.: Detection of quant myofibrillar proteins connectin and nobulin by electrophoresis in 2% polyacrylamide slab gels strengthened with agarose. *Anal. Biochem.*, 224 (1995) 28-31.
- 1404 Tokimitsu, I., Takehana, M., Hori, H., Nagai, Y. and Tajima, S.: Identification of nondisulfided Pro $\alpha_1$ (IV) chain produced by cultured B16 melanoma cells. *J. Biochem. (Tokyo)*, 116 (1994) 1039-1043.
- 1405 Urbancikova, M. and Hitchcock-DeGregori, S.E.: Requirement of amino-terminal modification for striated muscle  $\alpha$ -tropomyosin function. *J. Biol. Chem.*, 269 (1994) 24310-24315.
- 1406 Wheeler, T.J., Fell, R.D. and Hauck, M.A.: Translocation of two glucose transporters in heart: effects of rotenone, uncouplers, workload, palmitate, insulin and anoxia. *Biochim. Biophys. Acta*, 1196 (1994) 191-200.
- 1407 Yamada, H., Tajima, S., Nishikawa, T., Murad, S. and Pinnell, S.R.: Tranilast, a selective inhibitor of collagen synthesis in human skin fibroblasts. *J. Biochem. (Tokyo)*, 116 (1994) 892-897.
- 1408 Yamakawa, Y., Omori-Sato, T. and Mebs, D.: Hemorrhagic principles in the venom of *Bitis arietans*, a viperous snake. II. Enzymatic properties with special reference to substrate specificity. *Biochim. Biophys. Acta*, 1247 (1995) 17-23.

See also 1240, 1245, 1452, 1468.

#### 19g. Protamines, histones and other nuclear proteins

- 1409 Bossi, A., Gelfi, C., Orsi, A. and Righetti, P.G.: Isoelectric focusing of histones in extremely alkaline immobilized pH gradients: comparison with capillary electrophoresis. *J. Chromatogr. A*, 686 (1994) 121-128.
- 1410 Macaulay, C., Meier, E. and Forbes, D.J.: Differential mitotic phosphorylation of proteins of the nuclear pore complex. *J. Biol. Chem.*, 270 (1995) 254-262.
- 1411 Nagaraja, S., Delcuve, G.P. and Davie, J.R.: Differential compaction of transcriptionally competent and repressed chromatin reconstituted with histone H1 subtypes. *Biochim. Biophys. Acta*, 1260 (1995) 207-214.
- 1412 Wang, B. and Williamson, G.: Detection of a nuclear protein which binds specifically to the antioxidant responsive element (ARE) of the human NAD(P)H:quinone oxidoreductase gene. *Biochim. Biophys. Acta*, 1219 (1994) 645-652.

See also 1738.

#### 19h. Chromoproteins and metalloproteins

- 1413 Chiancone, E., Gattoni, M., Fronticelli, C., Termine, A., Silvestri, L. and Sarti, P.: Preparative and analytical applications of immobilized haemoglobin. *J. Chromatogr. B*, 664 (1995) 89-95.
- 1414 Ferranti, P., Malorni, A. and Pucci, P.: Structural characterization of hemoglobin variants using capillary electrophoresis and fast atom bombardment mass spectrometry. *Methods Enzymol.*, 231(Hemoglobins, Pt. B) (1994) 45-65; C.A., 121 (1994) 275582d.
- 1415 Gusov, V.M., Zelko, I.N., Bylinskaya, S.A. and Usanov, S.A.: (Structure-functional analysis of cytochrome P450c21. Immunochemical approaches and limited proteolysis). *Biokhimiya (Moscow)*, 59 (1994) 1149-1163.
- 1416 Hempe, J.M. and Craver, R.D.: Quantification of hemoglobin variants by capillary isoelectric focusing. *Clin. Chem. (Washington)*, 40 (1994) 2288-2295.
- 1417 Hibino, T., Lee, R.H. and Takabe, T.: Role of transit peptide sequence of plastocyanin for its expression, processing, and copper-binding activity in *Escherichia coli*. *J. Biochem. (Tokyo)*, 116 (1994) 826-832.
- 1418 Kister, J., Kiger, L., Francina, A., Hannay, P., Szymanowicz, A., Blouquit, Y., Promé, D., Galastéros, F., Delaunay, J. and Wajzman, H.: Hemoglobin Roanne [ $\alpha$ 94(G1) Asp  $\rightarrow$  Glu]: a variant of the  $\alpha$ 1 $\beta$ 2 interface with an unexpected high oxygen affinity. *Biochim. Biophys. Acta*, 1246 (1995) 34-38.
- 1419 Miyatake, A., Tsubaki, M., Hori, H. and Ichikawa, Y.: Purification and comparative characterization of cytochrome P-450<sub>sc</sub> (CYP XIA1) from sheep adrenocortical mitochondria. *Biochim. Biophys. Acta*, 1215 (1994) 176-182.
- 1420 Shaeffer, R.: Heterogeneity in the structure of the ubiquitin conjugates of human  $\alpha$  globin. *J. Biol. Chem.*, 269 (1994) 29530-29536.

See also 1385.

#### 19i. Proteins of glands, gland products, various zymogens (incl. milk proteins)

- 1421 Bjerrum, K.B. and Prause, J.U.: Collection and concentration of tear proteins studied by SDS gel electrophoresis: presentation of a new method with special reference to dry eye patients. *Graefe's Arch. Clin. Exp. Ophthalmol.*, 232 (1994) 402-405; C.A., 121 (1994) 225606u.
- 1422 Cifuentes, A., de Frutos, M. and Diez-Masa, J.C.: Polymeric networks vs. cross-linked polyacrylamide bonded gels for CE separations of whey proteins. *Curr. Top. Membr.*, 41(Cell Biology and Membrane Transport Processes) (1994) 109-122; C.A., 121 (1994) 275866z.
- 1423 Ishihama, N.: (Electrophoretic analysis of the proteins in human saliva. 3. SDS-gel electrophoresis with the inclusion of soluble polymers and its application to parotid Ps typing). *Mie Igaku*, 38 (1994) 123-126; C.A., 122 (1995) 4741c.
- 1424 Kramer, W., Wess, G., Enhsen, A., Bock, K., Falk, E., Hoffmann, A., Neckermann, G., Gantz, D., Schulz, S., Nickau, L., Petzinger, E., Turley, S. and Dietschy J.M.: Bile acid derived HMG-CoA reductase inhibitors. *Biochim. Biophys. Acta*, 1227 (1994) 137-154.

- 1425 Malin, E.L., Basch, J.J., Shieh, J.J., Sullivan, B.C. and Holsinger, V.H.: Detection of adulteration of buttermilk powder by gel electrophoresis. *J. Dairy Sci.*, 77 (1994) 2199-2206; C.A., 121 (1994) 299397y.
  - 1426 Valdivia, H.H., Martin, B.M., Ramirez, A.N., Fletcher, P.L. and Possani, L.D.: Isolation and pharmacological characterization of four novel Na<sup>+</sup> channel-blocking toxins from the scorpion *Centruroides noxius* Hoffmann. *J. Biochem. (Tokyo)*, 116 (1994) 1383-1391.
- See also 1253, 1325, 1558.
- 19j. *Proteins of brain, cerebrospinal fluid and eye*
- 1427 Beaulieu, E., Demeule, M., Pouliot, J.-F., Averill-Bates, D.A., Murphy, G.F. and Béliveau, R.: P-glycoprotein of blood brain barrier: cross-reactivity of MAb C219 with a 190 kDa protein in bovine and rat isolated brain capillaries. *Biochim. Biophys. Acta*, 1233 (1995) 27-32.
  - 1428 Catling, A.D., Reuter, C.W.M., Cox, M.E., Parsons, S.J. and Weber, M.J.: Partial purification of a mitogen-activated protein kinase kinase activator from bovine brain. Identification as B-Raf or a B-Raf-associated activity. *J. Biol. Chem.*, 269 (1994) 30014-30021.
  - 1429 Giardini, G., Lorusso, L., Barletta, L., Malaspina, A., Bolzani, W., Savoldi, F. and Ceroni, M.: Myelin basic protein purification in non-denaturing conditions. *Boll.-Soc. Ital. Biol. Sper.*, 69 (1993) 579-585; C.A., 121 (1994) 225585m.
  - 1430 Hsu, Y. and Molday, R.S.: Interaction of calmodulin with the cyclic GMP-gated channel of rod photoreceptor cells. Modulation of activity, affinity purification, and localization. *J. Biol. Chem.*, 269 (1994) 29765-29770.
  - 1431 Ichida, S., Wada, T., Akimoto, T., Kasamatsu, Y., Tahara, M. and Hasimoto, K.: Characteristics of specific <sup>125</sup>I- $\omega$ -conotoxin GVIA binding and <sup>125</sup>I- $\omega$ -conotoxin GVIA labeling using bifunctional crosslinkers in crude membranes from chick whole brain. *Biochim. Biophys. Acta*, 1233 (1995) 57-67.
  - 1432 Liljelund, P., Ghosh, P. and van den Pol, A.N.: Expression of the neural axon adhesion molecule L1 in the developing and adult rat brain. *J. Biol. Chem.*, 269 (1994) 32886-32895.
  - 1433 Mawal-Dewan, M., Henley, J., van de Voorde, A., Trojanowski, J.Q. and Lee, V.M.-Y.: The phosphorylation state of tau in the developing rat brain is regulated by phosphoprotein phosphatases. *J. Biol. Chem.*, 269 (1994) 30981-30987.
  - 1434 McPherson, P.S., Takei, K., Schmid, S.L. and de Camilli, P.: p145, A major Grb2-binding protein in brain, is co-localized with dynamin in nerve terminals where it undergoes activity-dependent dephosphorylation. *J. Biol. Chem.*, 269 (1994) 30132-30139.
  - 1435 Singer, W.D., Parker, S.K., Himes, R.H. and Detrich, H.W., III.: Polymerization of antarctic fish tubulins at low temperatures: role of carboxy-terminal domains. *Biochemistry*, 33 (1994) 15389-15396.
  - 1436 Solca, F.F., Lurie, D.I., Diltz, C.D., Johnson, R.S., Kumar, S., Rubel, E.W. and Fischer, E.H.: Identification and purification of a chicken brain neuroglia-associated protein. *J. Biol. Chem.*, 269 (1994) 27559-27565.
  - 19k. *Proteins of neoplastic tissue and transformed cells*
  - 1437 Hipfner, D.R., Gaudie, S.D., Deeley, R.G. and Cole, S.P.C.: Detection of the Mr 190,000 multidrug resistance protein, MRP, with monoclonal antibodies. *Cancer Res.*, 54 (1994) 5788-5792.
  - 1438 Miyagi, Y., Koshikawa, N., Yasumitsu, H., Miyagi, E., Hirahara, F., Aoki, I., Misugi, K., Umeda, M. and Miyazaki, K.: cDNA Cloning and mRNA expression of a serine proteinase inhibitor secreted by cancer cells: identification as placental protein 5 and tissue factor pathway inhibitor-2. *J. Biochem. (Tokyo)*, 116 (1994) 939-942.
  - 1439 Mohammad, R.M., Maki, A., Vistisen, K. and Al-Katib, A.: Protein studies of human non-Hodgkin's B-lymphoma: appraisal by two-dimensional gel electrophoresis. *Electrophoresis (Weinheim)*, 15 (1994) 1566-1572.
  - 1440 Nakata, N. and Tozawa, T.: (Evaluation of AFP-bands separated by lectin-affinity electrophoresis and PIVKA-II in early diagnosis of hepatocellular carcinoma). *Seibutsu Butsuri Kagaku*, 38 (1994) 327-331; C.A., 121 (1994) 275867a.
  - 1441 Oyama, T., Kanai, Y., Ochiai, A., Akimoto, S., Oda, T., Yanagihara, K., Nagafuchi, A., Tsukita, S., Shibamoto, S., Ito, F., Takeichi, M. et al.: A truncated  $\beta$ -catenin disrupts the interaction between E-cadherin and  $\alpha$ -catenin: a cause of loss of intercellular adhesiveness in human cancer cell lines. *Cancer Res.*, 54 (1994) 6282-6287.
  - 1442 Pazur, J.H.: Coupled gel electrophoresis-agar diffusion method for the detection of tumor antigens. *J. Chromatogr. B*, 663 (1995) 51-57.
  - 1443 Wojda, U. and Kuznicki, J.: Calcyclin from mouse Ehrlich ascites tumor cells and rabbit lung form non-covalent dimers. *Biochim. Biophys. Acta*, 1209 (1994) 248-252.
- See also 1314, 1619.
- 19l. *Specific binding and receptor proteins*
- 1444 Antonijczuk, K., Kroftova, O.S., Varghese, A.H., Antonijczuk, A., Henjum, D.C., Korza, G., Ozols, J. and Sunderman, F.W., Jr.: The 40 kDa <sup>63</sup>Ni<sup>2+</sup>-binding protein (pNiXc) on Western blots of *Xenopus laevis* oocytes and embryos is the monomer of fructose-1,6-bisphosphate aldolase A. *Biochim. Biophys. Acta*, 1247 (1995) 81-89.
  - 1445 Arnold, M., Ringler, P. and Brisson, A.: A quantitative electrophoretic migration shift assay for analyzing the specific binding of proteins to lipid ligands in vesicles or micelles. *Biochim. Biophys. Acta*, 1233 (1995) 198-204.
  - 1446 Barrett, P., MacLean, A. and Morgan, P.J.: Evidence for multiple forms of melatonin receptor-G-protein complexes by solubilization and gel electrophoresis. *J. Neuroendocrinol.*, 6 (1994) 509-515; C.A., 121 (1994) 272671w.
  - 1447 Blaikie, P., Immanuel, D., Wu, J., Li, N., Yajnik, V. and Margolis, B.: A region in Shc distinct from the SH2 domain can bind tyrosine-phosphorylated growth factor receptors. *J. Biol. Chem.*, 269 (1994) 32031-32034.
  - 1448 Bommakanti, R.K., Dratz, E.A., Siemsen, D.W. and Jesaitis, A.J.: Characterization of complex formation between G $\alpha$ 2 and octyl glucoside solubilized neutrophil N-formyl peptide chemoattractant receptor by sedimentation velocity. *Biochim. Biophys. Acta*, 1209 (1994) 69-76.

- 1449 Burg, D.L., Furlong, M.T., Harrison, M.L. and Geahlen, R.L.: Interaction of Lyn with the antigen receptor during B cell activation. *J. Biol. Chem.*, 269 (1994) 28136-28142.
- 1450 Chauchereau, A., Cohen-Solal, K., Jolivet, A., Bailly, A. and Milgrom, E.: Phosphorylation sites in ligand-induced and ligand-independent activation of the progesterone receptor. *Biochemistry*, 33 (1994) 13295-13303.
- 1451 Guo, B., Yu, Y. and Leibold, E.A.: Iron regulates cytoplasmic levels of a novel iron-responsive element-binding protein without aconitase activity. *J. Biol. Chem.*, 269 (1994) 24252-24260.
- 1452 Hansen, C.A., Joseph, S.K. and Robishaw, J.D.: Ins 1,4,5-P<sub>3</sub> and Ca<sup>2+</sup> signaling in quiescent neonatal cardiac myocytes. *Biochim. Biophys. Acta*, 1224 (1994) 517-526.
- 1453 Imamura, T., Takata, Y., Sasaoka, T., Takada, Y., Morioka, H., Haruta, T., Sawa, T., Iwanishi, M., Hu, Y.G., Suzuki, Y., Hamada, J. and Kobayashi, M.: Two naturally occurring mutations in the kinase domain of insulin receptor accelerate degradation of the insulin receptor and impair the kinase activity. *J. Biol. Chem.*, 269 (1994) 31019-31027.
- 1454 Inano, K., Curtis, S.W., Korach, K.S., Omata, S. and Horigome, T.: Heat shock protein 90 strongly stimulates the binding of purified estrogen receptor to its responsive element. *J. Biochem. (Tokyo)*, 116 (1994) 759-766.
- 1455 Johnson, J.L. and Toft, D.O.: A novel chaperone complex for steroid receptors involving heat shock proteins, immunophilins, and p23. *J. Biol. Chem.*, 269 (1994) 24989-24993.
- 1456 Josic, D., Lim, Y.-P., Strancar, A. and Reutter, W.: Application of high-performance membrane chromatography for separation of annexins from the plasma membranes of liver and isolation of monospecific polyclonal antibodies. *J. Chromatogr. B*, 662 (1994) 217-226.
- 1457 Kimura, Y., Sonehara, K., Kuramoto, E., Makino, T., Yamamoto, S., Yamamoto, T., Kataoka, T. and Tokunaga, T.: Binding of oligoguanylate to scavenger receptors is required for oligonucleotides to augment NK cell activity and induce IFN. *J. Biochem. (Tokyo)*, 116 (1994) 991-994.
- 1458 Kotomura, N., Okada, M., Ninomiya, Y., Tsukiyama, T., Umesono, K., Evans, R.M. and Niwa, O.: Repression of retinoic acid-induced transactivation by embryonal LTR binding protein. *J. Biochem. (Tokyo)*, 116 (1994) 1309-1316.
- 1459 Kumar, K.N., Babcock, K.K., Johnson, P.S., Chen, X., Eggeman, K.T. and Michaelis, E.K.: Purification and pharmacological and immunochemical characterization of synaptic membrane proteins with ligand-binding properties of N-methyl-D-aspartate receptors. *J. Biol. Chem.*, 269 (1994) 27384-27393.
- 1460 Kurose, T., Pashmforoush, M., Yoshimasa, Y., Carroll, R., Schwartz, G.P., Burke, G.T., Katsoyannis, P.G. and Steiner, D.F.: Cross-linking of a B25 azidophenylalanine insulin derivative to the carboxyl-terminal region of the  $\alpha$ -subunit of the insulin receptor. Identification of a new insulin-binding domain in the insulin receptor. *J. Biol. Chem.*, 269 (1994) 29190-29197.
- 1461 Kusui, T., Benya, R.V., Battey, J.F. and Jensen, R.T.: Glycosylation of bombesin receptors: characterization, effect on binding, and G-protein coupling. *Biochemistry*, 33 (1994) 12968-12980.
- 1462 Lee, C. and Yamada, K.M.: Alternatively spliced juxtamembrane domain of a tyrosine kinase receptor is a multifunctional regulatory site. Deletion alters cellular tyrosine phosphorylation pattern and facilitates binding of phosphatidylinositol-3-OH kinase to the hepatocyte growth factor receptor. *J. Biol. Chem.*, 270 (1995) 507-510.
- 1463 Mao, S., Yamashita, T. and Metzger, H.: Chemical cross-linking of IgE-receptor complexes in RBL-2H3 cells. *Biochemistry*, 34 (1995) 1968-1977.
- 1464 Moren, A., Olofsson, A., Stenman, G., Sahlin, P., Kanzaki, T., Claesson-Welsh, L., ten Dijke, P., Miyazono, K. and Heldin, C.-H.: Identification and characterization of LTBP-2, a novel latent transforming growth factor- $\beta$ -binding protein. *J. Biol. Chem.*, 269 (1994) 32469-32478.
- 1465 Mori, H., Yamashita, Y., Akasaka, T. and Yamamoto, Y.: Further characterization of the loss of antenna chlorophyll-binding protein CP43 from Photosystem II during donor-side photoinhibition. *Biochim. Biophys. Acta*, 1228 (1995) 37-42.
- 1466 Neumann, G.M., Condron, R., Thomas, I. and Polya, G.M.: Purification and sequencing of a family of wheat lipid transfer protein homologues phosphorylated by plant calcium-dependent protein kinase. *Biochim. Biophys. Acta*, 1209 (1994) 183-190.
- 1467 Pashmforoush, M., Yoshimasa, Y. and Steiner, D.F.: Exon 11 enhances insulin binding affinity and tyrosine kinase activity of the human insulin proreceptor. *J. Biol. Chem.*, 269 (1994) 32639-32648.
- 1468 Prinsen, C.F.M., Werten, P.J.L., Maassen, J.A. and Veerkamp, J.H.: No significant tyrosine phosphorylation of muscle fatty acid-binding protein. *Biochim. Biophys. Acta*, 1215 (1994) 103-108.
- 1469 Reif, O.-W., Lausch, R., Scheper, T. and Freitag, R.: Fluorescein isothiocyanate-labeled protein G as an affinity ligand in affinity/immunocapillary electrophoresis with fluorescence detection. *Anal. Chem.*, 66 (1994) 4027-4033.
- 1470 Safi, A., Saunier, M., Gastin, I., Alibada, Y., Dugue, B. and Gueant, J.-L.: Intrinsic factor covalently bound to Sepharose as affinity medium for the purification of a soluble intrinsic factor receptor from human urine. *J. Chromatogr. B*, 664 (1995) 253-259.
- 1471 Sautes, C., Galinha, A., Bouchard, C., Mazieres, N., Spagnoli, R. and Fridman, W.H.: Recombinant soluble Fc $\gamma$  receptors: production, purification and biological activities. *J. Chromatogr. B*, 662 (1994) 197-207.
- 1472 Singh, L., Wadhwa, R., Naidu, S., Nagaraj, R. and Ganesan, M.: Sex- and tissue-specific Bkm(GATA)-binding protein in the germ cells of heterogametic sex. *J. Biol. Chem.*, 269 (1994) 25321-25327.
- 1473 Staubs, P.A., Reichart, D.R., Saltiel, A.R., Milarski, K.L., Maegawa, H., Berhanu, P., Olersky, J.M. and Seely, B.L.: Localization of the insulin receptor binding sites for the SH2 domain proteins p85, Syp, and GAP. *J. Biol. Chem.*, 269 (1994) 27186-27192.
- 1474 Stroop, S.D., Kuestner, R.E., Serwold, T.F., Chen, L. and Moore, E.E.: Chimeric human calcitonin and glucagon receptors reveal two dissociable calcitonin interaction sites. *Biochemistry*, 34 (1995) 1050-1057.

- 1475 Sugimoto, K., Yata, H., Muro, Y. and Himeno, M.: Human centromere protein C (CENP-C) is a DNA-binding protein which possesses a novel DNA-binding motif. *J. Biochem. (Tokyo)*, 116 (1994) 877-881.
- 1476 Swope, S.L. and Haganir, R.L.: Binding of the nicotinic acetylcholine receptor to SH2 domains of Fyn and Fyk protein tyrosine kinases. *J. Biol. Chem.*, 269 (1994) 29817-29824.
- 1477 Take, M., Tsutsui, J.-i., Obama, H., Ozawa, M., Nakayama, T., Maruyama, I., Arima, T. and Muramatsu, T.: Identification of nucleolin as a binding protein for midkine (MK) and heparin-binding growth associated molecule (HB-GAM). *J. Biochem. (Tokyo)*, 116 (1994) 1063-1068.
- 1478 Takeda, E., Miyamoto, K.-i., Kubota, M., Minami, H., Yokota, I., Saijo, T., Naito, E., Ito, M. and Kuroda, Y.: Vitamin D-dependent rickets type II: regulation of human osteocalcin gene expression in cells with defective vitamin D receptors by 1,25-dihydroxyvitamin D-3, retinoic acid, and triiodothyronine. *Biochim. Biophys. Acta*, 1227 (1994) 195-199.
- 1479 Visai, L., Bozzini, S., Raucci, G., Toniolo, A. and Speziale, P.: Isolation and characterization of a novel collagen-binding protein from *Streptococcus pyogenes* strain 6414. *J. Biol. Chem.*, 270 (1995) 347-353.
- See also 1248, 1301, 1306, 1314, 1434.
- 19m. *Urinary proteins*
- 1480 Büeler, M.R., Wiederkehr, F. and Vonderschmitt, D.J.: Electrophoretic, chromatographic and immunological studies of human urinary proteins. *Electrophoresis (Weinheim)*, 16 (1995) 124-134.
- 1481 Gultepe, M., Avsar, K., Ozcan, A., Tokgoz, S. and Karaca, L.: (Sodium dodecyl sulfate-polyacrylamide-agarose composite gels in the electrophoretic examination of urinary proteins). *Bi-yokim. Derg.*, 18 (1993) 71-82; C.A., 121 (1994) 225608w.
- 1482 Jenkins, M.A., O'Leary, T.D. and Guerin, M.D.: Identification and quantitation of human urine proteins by capillary electrophoresis. *J. Chromatogr. B*, 662 (1994) 108-112.
- 1483 Keely, E.J. and Faiman, C.: Measurement of human urinary prolactin as a noninvasive study tool. *Clin. Chem. (Washington)*, 40 (1994) 2017-2021.
- 19n. *Other proteins (incl. proteinous inhibitors of enzymic activity)*
- 1484 Creuwels, L.A.J.M., Demel, R.A., van Golde, L.M.G. and Haagsman, H.P.: Characterization of a dimeric canine form of surfactant protein C (SP-C). *Biochim. Biophys. Acta*, 1254 (1995) 326-332.
- 1485 Delaria, K., Fiorentino, L., Wallace, L., Tamburini, P., Brownmell, E. and Muller, D.: Inhibition of cathepsin L-like cysteine proteases by cytotoxic T-lymphocyte antigen-2 $\beta$ . *J. Biol. Chem.*, 269 (1994) 25172-25177.
- 1486 Libera, L.D., Massimino, M.L., Arslan, P., Beltrame, M. and Cantini, M.: Analysis of muscle cell culture medium by size-exclusion chromatography. *J. Chromatogr. B*, 664 (1995) 185-191.
- 1487 Liu, B., Nakashima, S., Kanoh, H., Takano, T., Shimizu, T. and Nozawa, Y.: Activation of phospholipase D in Chinese hamster ovary cells expressing platelet-activating factor receptor. *J. Biochem. (Tokyo)*, 116 (1994) 882-891.
- 1488 Vachier, M.C., Piot, M. and Awadé, A.C.: Isolation of hen egg white lysozyme, ovotransferrin and ovalbumin, using a quaternary ammonium bound to a highly crosslinked agarose matrix. *J. Chromatogr. B*, 664 (1995) 201-210.
- 1489 Vaintraub, I.A., Bulmaga, V.P. and Musalem, M.: (The mechanism of ovalbumin proteolysis). *Biokhimiya (Moscow)*, 59 (1994) 1140-1148.
- See also 1179, 1327.
20. **ENZYMES AND ENZYME ACTIVITY ESTIMATION**
- 1490 Murphy, P.G.: Multilocus enzyme electrophoresis. *Soc. Appl. Bacteriol. Tech. Ser.*, 31 (New Techniques in Food and Beverage Microbiology) (1993) 211-221; C.A., 121 (1994) 249822n - a review with many refs.
- See also 1039.
- 20a. *Oxidoreductases*
- 1491 Barnett, J., Chow, J., Ives, D., Chiou, M., Mackenzie, R., Osen, E., Nguyen, B., Tsing, S., Bach, C., Freire, J., Chan, H., Sigal, E. and Ramesha, C.: Purification, characterization and selective inhibition of human prostaglandin G/H synthase 1 and 2 expressed in the baculovirus system. *Biochim. Biophys. Acta*, 1209 (1994) 130-139.
- 1492 Camadro, J.-M., Thome, F., Brouillet, N. and Labbe, P.: Purification and properties of protoporphyrinogen oxidase from the yeast *Saccharomyces cerevisiae*. Mitochondrial location and evidence for a precursor form of the protein. *J. Biol. Chem.*, 269 (1994) 32085-32091.
- 1493 Carper, D.A., Hohman, T.C. and Old, S.E.: Residues affecting the catalysis and inhibition of rat lens aldose reductase. *Biochim. Biophys. Acta*, 1246 (1995) 67-73.
- 1494 Cendrin, F., Jouve, H.M., Gaillard, J., Thibault, P. and Zaccari, G.: Purification and properties of a halophilic catalase-peroxidase from *Haloarcula marismortui*. *Biochim. Biophys. Acta*, 1209 (1994) 1-9.
- 1495 Ciriolo, M.R., Civitareale, P., Carri, M.T., de Martino, A., Galianzo, F. and Rotilio, G.: Purification and characterization of Ag,Zn-superoxide dismutase from *Saccharomyces cerevisiae* exposed to silver. *J. Biol. Chem.*, 269 (1994) 25783-25787.
- 1496 Créminon, C., Frobert, Y., Habib, A., Maclouf, J., Pradelles, P. and Grassi, J.: Immunological studies of human constitutive cyclooxygenase (COX-1) using enzyme immunometric assay. *Biochim. Biophys. Acta*, 1254 (1995) 333-340.
- 1497 Créminon, C., Habib, A., Maclouf, J., Pradelles, P., Grassi, J. and Frobert, Y.: Differential measurement of constitutive (COX-1) and inducible (COX-2) cyclooxygenase expression in human umbilical vein endothelial cells using specific immunometric enzyme immunoassays. *Biochim. Biophys. Acta*, 1254 (1995) 341-348.
- 1498 Freire-Moar, J., Alavi-Nassab, A., Ng, M., Mulkins, M. and Sigal, E.: Cloning and characterization of a murine macrophage lipoygenase. *Biochim. Biophys. Acta*, 1254 (1995) 112-116.

- 1499 Imaishi, H., Yamada, T. and Ohkawa, H.: Purification and immunochemical characteristics of NADPH-cytochrome P-450 oxidoreductase from tobacco cultured cells. *Biochim. Biophys. Acta*, 1246 (1995) 53-60.
  - 1500 Nakai, T., Mera, Y., Yasuhara, T. and Ohashi, A.: Divalent metal ion-dependent mitochondrial degradation of unassembled subunits 2 and 3 of cytochrome c oxidase. *J. Biochem. (Tokyo)*, 116 (1994) 752-758.
  - 1501 Noguchi, M., Miyano, M., Matsumoto, T. and Noma, M.: Human 5-lipoxygenase associates with phosphatidylcholine liposomes and modulates LTA<sub>4</sub> synthetase activity. *Biochim. Biophys. Acta*, 1215 (1994) 300-306.
  - 1502 Novikov, D.K., Vanhove, G.F., Carchon, H., Asselberghs, S., Eyssen, H.J., van Veldhoven, P.P. and Mannaerts, G.P.: Peroxisomal  $\beta$ -oxidation. Purification of four novel 3-hydroxyacyl-CoA dehydrogenases from rat liver peroxisomes. *J. Biol. Chem.*, 269 (1994) 27125-27135.
  - 1503 Ohara, H., Nakayama, T., Deyashiki, Y., Hara, A., Miyabe, Y. and Tsukada, F.: Reduction of prostaglandin D<sub>2</sub> to 9 $\alpha$ ,11 $\beta$ -prostaglandin F<sub>2</sub> by a human liver 3 $\alpha$ -hydroxysteroid/dihydrodiol dehydrogenase isozyme. *Biochim. Biophys. Acta*, 1215 (1994) 59-65.
  - 1504 Simon, M., Séraphin, B. and Faye, G.: The nuclear-encoded MSS2 gene is involved in the expression of the mitochondrial cytochrome-c oxidase subunit 2 (Cox2). *Biochim. Biophys. Acta*, 1228 (1995) 95-98.
  - 1505 Veijola, J., Koivunen, P., Annunen, P., Pihlajaniemi, T. and Kivirikko, K.I.: Cloning, baculovirus expression, and characterization of the  $\alpha$  subunit of propyl 4-hydroxylase from the nematode *Caenorhabditis elegans*. This  $\alpha$  subunit forms and active  $\alpha\beta$  dimer with the human protein disulfide isomerase/ $\beta$  subunit. *J. Biol. Chem.*, 269 (1994) 26746-26753.
  - 1506 Warburton, R.J. and Seybert, D.W.: Structural and functional characterization of bovine adrenodoxin reductase by limited proteolysis. *Biochim. Biophys. Acta*, 1246 (1995) 39-46.
- See also 1241, 1323, 1340.
- 20b. *Transferases (excl. E.C. 2.7.-.)*
  - 1507 Andreotti, G., Cubellis, M.V., Nitti, G., Sannia, G., Mai, X., Adams, M.W.W. and Marino, G.: An extremely thermostable aromatic aminotransferase from the hyperthermophilic archaeon *Pyrococcus furiosus*. *Biochim. Biophys. Acta*, 1247 (1995) 90-96.
  - 1508 Awald, P.D., Frost, D., Drake, R.R. and Selitrennikoff, C.P.: (1,3)  $\beta$ -Glucan synthase activity of *Neurospora crassa*: identification of a substrate-binding protein. *Biochim. Biophys. Acta*, 1201 (1994) 312-320.
  - 1509 Bouchon, B., Jaquinod, M., Klarskov, K., Trottein, F., Klein, M., van Dorsselaer, A., Bischoff, R. and Roitsch, C.: Analysis of the primary structure and post-translational modifications of the *Schistosoma mansoni* antigen Smp28 by electrospray mass spectrometry. *J. Chromatogr. B*, 662 (1994) 279-290.
  - 1510 Langkamp-Henken, B., Geller, A.M., LeGros, H.L., Jr., Price, J.O., de la Rosa, J. and Kotb, M.: Characterization of distinct forms of methionine adenosyltransferase in nucleated, and mature human erythrocytes and erythroleukemic cells. *Biochim. Biophys. Acta*, 1201 (1994) 397-404.
  - 1511 Mattsson, P., Battchikova, N., Sippola, K. and Korpela, T.: The role of histidine residues in the catalytic act of cyclomaltodextrin glucanotransferase from *Bacillus circulans* var. *alkalophilus*. *Biochim. Biophys. Acta*, 1247 (1995) 97-103.
  - 1512 Nguyen, T.T.M., Hinton, D.A. and Shur, B.D.: Expressing murine  $\beta$ 1,4-galactosyltransferase in HeLa cells produces a cell surface galactosyltransferase-dependent phenotype. *J. Biol. Chem.*, 269 (1994) 28000-28009.
  - 1513 Strege, M.A., Schmidt, D.F., Kreuzman, A., Dotzla, J., Yeh, W.K., Kaiser, R.E. and Lagu, A.L.: Capillary electrophoretic analysis of serine hydroxymethyltransferase in *Escherichia coli* fermentation broth. *Anal. Biochem.*, 223 (1994) 198-204.
- See also 1578.
- 20c. *Transferases transferring phosphorus containing groups (E.C. 2.7.-.)*
  - 1514 Allen, B.G., Andrea, J.E. and Walsh, M.P.: Identification and characterization of protein kinase C $\zeta$ -immunoreactive proteins. *J. Biol. Chem.*, 269 (1994) 29288-29298.
  - 1515 Boscoboinik, D.O., Chatelain, E., Bartoli, G.-M., Stäubli, B. and Azzi, A.: Inhibition of protein kinase C activity and vascular smooth muscle cell growth by d- $\alpha$ -tocopherol. *Biochim. Biophys. Acta*, 1224 (1994) 418-426.
  - 1516 Clerk, A., Bogoyevitch, M.A., Andersson, M.B. and Sugden, P.H.: Differential activation of protein kinase C isoforms by endothelin-1 and phenylephrine and subsequent stimulation of p42 and p44 mitogen-activated protein kinases in ventricular myocytes cultured from neonatal rat hearts. *J. Biol. Chem.*, 269 (1994) 32848-32857.
  - 1517 Fasco, M.J., Treanor, C.P., Spivack, S., Fiege, H.L. and Kaminsky, L.S.: Quantitative RNA-polymerase chain reaction-DNA analysis by capillary electrophoresis. *Anal. Biochem.*, 224 (1995) 140-147.
  - 1518 Fialkow, L., Chan, C.K., Rotin, D., Grinstein, S. and Downey, G.P.: Activation of the mitogen-activated protein kinase signaling pathway in neutrophils. Role of oxidants. *J. Biol. Chem.*, 269 (1994) 31234-31242.
  - 1519 Ishino, Y., Ueno, T., Miyagi, M., Uemori, T., Imamura, M., Tsunasawa, S. and Kato, I.: Overproduction of *Thermus aquaticus* DNA polymerase and its structural analysis by ion-spray mass spectrometry. *J. Biochem. (Tokyo)*, 116 (1994) 1019-1024.
  - 1520 Kakuda, H., Hosono, K., Shiroishi, K. and Ichihara, S.: Identification and characterization of the ackA (acetate kinase A)-pta (phosphotransacetylase) operon and complementation analysis of acetate utilization by an ackA-pta deletion mutant of *Escherichia coli*. *J. Biochem. (Tokyo)*, 116 (1994) 916-922.
  - 1521 Kanemitsu, F. and Okigaki, T.: Characterization of human creatine kinase BB and MB isoforms by means of isoelectric focusing. *Clin. Chim. Acta*, 231 (1994) 1-9.
  - 1522 Koller, G., Graumann, K., Kramer, W., Sara, M. and Jungbauer, A.: Laboratory-scale production and purification of recombinant HIV-1 reverse transcriptase. *J. Chromatogr. B*, 664 (1995) 107-118.

- 1523 McMahan, S.A. and Burgess, R.R.: Use of aryl azide cross-linkers to investigate protein-protein interactions: an optimization of important conditions as applied to *Escherichia coli* RNA polymerase and localization of a  $\sigma^{70}$ - $\alpha$  cross-link to the C-terminal region of  $\alpha$ . *Biochemistry*, 33 (1994) 12092-12099.
- 1524 Mitchell, C.E., Belinsky, S.A. and Lechner, J.F.: Detection and quantitation of mutant  $\kappa$ -ras codon 12 restriction fragment by capillary electrophoresis. *Anal. Biochem.*, 224 (1995) 148-153.
- 1525 Nadeau, O.W. and Carlson, G.M.: Zero length conformation-dependent cross-linking of phosphorylase kinase subunits by transglutaminase. *J. Biol. Chem.*, 269 (1994) 29670-29676.
- 1526 Ogg, S., Gabrielli, B. and Piwnicka-Worms, H.: Purification of a serine kinase that associates with and phosphorylates human Cdc25C on serine 216. *J. Biol. Chem.*, 269 (1994) 30461-30469.
- 1527 Okuno, S., Kitani, T. and Fujisawa, H.: Purification and characterization of  $\text{Ca}^{2+}$ /calmodulin-dependent protein kinase IV kinase from rat brain. *J. Biochem. (Tokyo)*, 116 (1994) 923-930.
- 1528 Raffaniello, R.D. and Raufman, J.-P.: Protein kinase C expression and translocation in dispersed chief cells from guinea-pig stomach. *Biochim. Biophys. Acta*, 1224 (1994) 551-558.
- 1529 Rempel, A., Bannasch, P. and Mayer, D.: Differences in expression and intracellular distribution of hexokinase isoenzymes in rat liver cells of different transformation stages. *Biochim. Biophys. Acta*, 1219 (1994) 660-668.
- 1530 Sharova, N.P., Eliseeva, E.D. and Mikhailov, V.S.: (The presence of DNA polymerase  $\zeta$ -type in the eggs of the teleost fish *Misgurnus fossilis* L.) *Biokhimiya (Moscow)*, 59 (1994) 1067-1075.
- 1531 Stack, J.H. and Emr, S.D.: VPS34p required for yeast vacuolar protein sorting is a multiple specificity kinase that exhibits both protein kinase and phosphatidylinositol-specific PI 3-kinase activities. *J. Biol. Chem.*, 269 (1994) 31552-31562.
- 1532 Stover, D.R., Liebetanz, J. and Lydon, N.B.: Cdc2-mediated modulation of pp60<sup>csrc</sup> activity. *J. Biol. Chem.*, 269 (1994) 26885-26889.
- 1533 Tokumitsu, H., Brickey, D.A., Glod, J., Hidaka, H., Sikela, J. and Soderling, T.R.: Activation mechanisms for  $\text{Ca}^{2+}$ /calmodulin-dependent protein kinase IV. Identification of a brain CaM-kinase IV kinase. *J. Biol. Chem.*, 269 (1994) 28640-28647.
- 1534 Uchida, T.: Immunologically and enzymatically distinct rat choline kinase isozymes. *J. Biochem. (Tokyo)*, 116 (1994) 1241-1250.
- 1535 Lee, M.W., Kraemer, F.B. and Severson, D.L.: Characterization of a partially purified diacylglycerol lipase from bovine aorta. *Biochim. Biophys. Acta*, 1254 (1995) 311-318.
- 1539 Lin, J.-L., Wang, W.-Y. and Liao, T.-H.: Thermal inactivation of shrimp deoxyribonuclease with and without sodium dodecyl sulfate. *Biochim. Biophys. Acta*, 1209 (1994) 209-214.
- 1540 Masuno, H., Nakabayashi, H., Kobayashi, J., Saito, Y. and Okuda, H.: Reduced dimerization of lipoprotein lipase in post-heparin plasma of a patient with hyperchylomicronemia. *Biochim. Biophys. Acta*, 1254 (1995) 30-36.
- 1541 Mitchell, J., Gutierrez, J. and Northup, J.K.: Purification, characterization, and partial amino acid sequence of a G protein-activated phospholipase C from squid photoreceptors. *J. Biol. Chem.*, 270 (1995) 854-859.
- 1542 Nellore, R. and Bhagat, H.R.: Development and validation of capillary electrophoresis assay for ribonuclease A. *J. Pharm. Biomed. Anal.*, 12 (1994) 1363-1367.
- 1543 Nishihara, Y., Hayashi, Y., Fujii, T., Adachi, T., Stigbrand, T. and Hirano, K.: The alkaline phosphatase in human plexus chorioideus. *Biochim. Biophys. Acta*, 1209 (1994) 274-278.
- 1544 Park, J.-W., Oh, M.-S., Yang, J.-Y., Park, B.-H., Rho, H.-W., Lim, S.-N., Jhee, E.-C. and Kim, H.-R.: Glycosylation, dimerization, and heparin affinity of lipoprotein lipase in 3T3-L1 adipocytes. *Biochim. Biophys. Acta*, 1254 (1995) 45-50.
- 1545 Pervin, A., Gu, K. and Linhardt, R.J.: Capillary electrophoresis to measure sulfoesterase activity on chondroitin sulfate and heparin derived disaccharides. *Appl. Theor. Electrophor.*, 3 (1993) 297-303; *C.A.*, 121 (1994) 274906a.
- 1546 Rao, G.N., Runge, M.S. and Alexander, R.W.: Hydrogen peroxide activation of cytosolic phospholipase A<sub>2</sub> in vascular smooth muscle cells. *Biochim. Biophys. Acta*, 1265 (1995) 67-72.
- 1547 Sahas, Y., Thi, A.T.P., Roy-Macauley, H., d'Arcy-Lameta, A., Repellin, A. and Zuily-Fodil, Y.: Purification and characterization of a soluble lipolytic acylhydrolase from Cowpea (*Vigna unguiculata* L.) leaves. *Biochim. Biophys. Acta*, 1215 (1994) 66-73.
- 1548 Vervoordeldonk, M.J.B.M., Schalkwijk, C.G., Vishwanath, B.S., Aarsman, A.J. and van den Bosch, H.: Levels and localization of group II phospholipase A<sub>2</sub> and annexin I in interleukin- and dexamethasone-treated rat mesangial cells: evidence against annexin mediation of the dexamethasone-induced inhibition of group II phospholipases A<sub>2</sub>. *Biochim. Biophys. Acta*, 1224 (1994) 541-550.

See also 1236, 1242.

#### 20d. Hydrolases, acting on ester bonds (E.C. 3.1.-.-)

- 1535 Alonso, J., Rodriguez, M.T. and Canet, W.: Detection of pectinesterase in polyacrylamide gels. *Electrophoresis (Weinheim)*, 16 (1995) 39-42.
- 1536 Bucht, G., Häggström, B., Radic, Z., Osterman, A. and Hjalmarsson, K.: Residues important for folding and dimerisation of recombinant *Torpedo californica* acetylcholinesterase. *Biochim. Biophys. Acta*, 1209 (1994) 265-273.
- 1537 Chambers, R.S. and Dahmus, M.E.: Purification and characterization of a phosphatase from HeLa cells which dephosphorylates the C-terminal domain of RNA polymerase II. *J. Biol. Chem.*, 269 (1994) 26243-26248.

See also 1340, 1596, 1685.

#### 20e. Hydrolases, acting on glycosyl compounds (E.C. 3.2.-.-)

- 1549 Ashmarina, L.I., Muronets, V.I. and Nagradova, N.K.: (Glycolytic enzymes in human erythrocytes: glyceraldehyde-3-phosphate dehydrogenase association with 3-phosphoglycerate kinase). *Biokhimiya (Moscow)*, 59 (1994) 873-880.
- 1550 Duewel, H.S., Daub, E. and Honek, J.F.: Investigations of the interactions of saccharides with the lysozyme from bacteriophage lambda. *Biochim. Biophys. Acta*, 1247 (1995) 149-158.
- 1551 Maehama, T., Nishina, H. and Katada, T.: ADP-Ribosylarginine glycohydrolase catalyzing the release of ADP-ribose from the cholera toxin-modified  $\alpha$ -subunits of GTP-binding proteins. *J. Biochem. (Tokyo)*, 116 (1994) 1134-1138.

- 1552 Munoz, M.D., Hernández, L.M., Basco, R., Andaluz, E. and Larriba, G.: Glycosylation of yeast exoglucanase sequons in alg mutants deficient in the glucosylation steps of the lipid-linked oligosaccharide. Presence of glucotriose unit in Dol-PP-GlcNAc<sub>2</sub>Man<sup>9</sup>Glc<sub>3</sub> influences both glycosylation efficiency and selection of N-linked sites. *Biochim. Biophys. Acta*, 1201 (1994) 361-366.
- 1553 Roy, R., Brooks, C. and Mitra, S.: Purification and biochemical characterization of recombinant N-methylpurine-DNA glycosylase of the mouse. *Biochemistry*, 33 (1994) 15131-15140.
- 1554 Seog, D.-H., Kito, M., Yoda, K. and Yamasaki, M.: Uso1 protein contains a coiled-coil rod region essential for protein transport from the ER to the Golgi apparatus in *Saccharomyces cerevisiae*. *J. Biochem. (Tokyo)*, 116 (1994) 1341-1345.
- See also 1340.
- 20f. Other hydrolases**
- 1555 Bogomolov, A.A., Danilenko, A.N. and Antonov, Yu.A.: (Precipitation of alfalfa (*Medicago sativa* L.) leaf juice enzyme and the mechanism of its effect on RuBISCO hydrolysis). *Biokhimiya (Moscow)*, 59 (1994) 689-693.
- 1556 Bossi, A., Righetti, P.G., Vecchio, G. and Severinsen, S.: Focusing of alkaline proteases (subtilisins) in pH 10-12 immobilized gradients. *Electrophoresis (Weinheim)*, 15 (1994) 1535-1540.
- 1557 Crabbe, T., O'Connell, J.P., Smith, B.J. and Docherty, A.J.P.: Reciprocated matrix metalloproteinase activation: a process performed by interstitial collagenase and progelatinase A. *Biochemistry*, 33 (1994) 14419-14425.
- 1558 Fernandez-Resa, P., Mira, E. and Quesada, A.R.: Enhanced detection of casein zymography of matrix metalloproteinases. *Anal. Biochem.*, 224 (1995) 434-435.
- 1559 Fujimoto, N., Hosokawa, N., Iwawata, K., Shinya, T., Okada, Y. and Hayakawa, T.: A one-step sandwich enzyme immunoassay for inactive precursor and complexed forms of human matrix metalloproteinase 9 (92 kDa gelatinase/type IV collagenase, gelatinase B) using monoclonal antibodies. *Clin. Chim. Acta*, 231 (1994) 79-88.
- 1560 Kakinuma, Y. and Igarashi, K.: Purification and characterization of the catalytic moiety of vacuolar-type Na<sup>+</sup>-ATPase from *Enterococcus hirae*. *J. Biochem. (Tokyo)*, 116 (1994) 1302-1308.
- 1561 Khan, M.T., Wang, K., Auland, M.E., Kable, E.P.W. and Roufogalis, B.D.: Membrane-bound high molecular mass proteinases from human erythrocytes. *Biochim. Biophys. Acta*, 1209 (1994) 215-221.
- 1562 Kitazono, A., Kitano, A., Tsuru, D. and Yoshimoto, T.: Isolation and characterization of the prolyl aminopeptidase gene (pap) from *Aeromonas sobria*: comparison with the *Bacillus coagulans* enzyme. *J. Biochem. (Tokyo)*, 116 (1994) 818-825.
- 1563 Loayza, S.L., Trikha, M., Markland, F.S., Riquelme, P. and Kuo, J.: Resolution of isoforms of natural and recombinant fibronase, the fibrinolytic enzyme from *Agkistrodon contortrix contortrix* snake venom, and comparison of their EDTA sensitivities. *J. Chromatogr. B*, 662 (1994) 227-243.
- 1564 Lundy, F.T., Magee, A.C., Blair, I.S. and McDowell, D.A.: A new method for the detection of proteolytic activity in *Pseudomonas lundensis* after sodium dodecyl sulfate-polyacrylamide gel electrophoresis. *Electrophoresis (Weinheim)*, 16 (1995) 43-45.
- 1565 Morel, F., Dewald, B., Berthier, S., Zaoui, P., Dianoux, A.C., Vignais, P.V. and Baggolini, M.: Further characterization of the gelatinase-containing particles of human neutrophils. *Biochim. Biophys. Acta*, 1201 (1994) 373-380.
- 1566 Ohkubo, I., Huang, K., Ochiai, Y., Takagaki, M. and Kani, K.: Dipeptidyl peptidase IV from porcine seminal plasma: purification, characterization, and N-terminal amino acid sequence. *J. Biochem. (Tokyo)*, 116 (1994) 1182-1186.
- 1567 Park, M.-Y., Omote, H., Maeda, M. and Futai, M.: Conserved Glu-181 and Arg-182 residues of *Escherichia coli* H<sup>+</sup>-ATPase (ATP synthase)  $\beta$  subunit are essential for catalysis: properties of 33 mutants between  $\beta$ Glu-161 and  $\beta$ Lys-201 residues. *J. Biochem. (Tokyo)*, 116 (1994) 1139-1145.
- 1568 Ronca, F., Ranieri-Raggi, M., Brown, P.E., Moir, A.J.G. and Raggi, A.: Evidence of a species-differentiated regulatory domain within the N-terminal region of skeletal muscle AMP deaminase. *Biochim. Biophys. Acta*, 1209 (1994) 123-129.
- 1569 Roth, R. and Bachofen, R.: Characterization and partial purification of an ATPase and inorganic pyrophosphatase of the archaebacterium *Methanobacterium thermoautotrophicum*. *Biochim. Biophys. Acta*, 1201 (1994) 271-283.
- 1570 Rozhin, J., Sameni, M., Ziegler, G. and Sloane, B.F.: Pericellular pH affects distribution and secretion of cathepsin B in malignant cells. *Cancer Res.*, 54 (1994) 6517-6525.
- 1571 Siebert, A., Lottspeich, F., Nelson, N. and Betz, H.: Purification of the synaptic vesicle-binding protein physophilin. Identification as 39-kDa subunit of the vacuolar H<sup>+</sup>-ATPase. *J. Biol. Chem.*, 269 (1994) 28329-28334.
- 1572 Vladimirova, N.M., Potapenko, N.A., Sachs, G. and Modyanov, N.N.: Determination of the sidedness of the carboxy-terminus of the Na<sup>+</sup>/K<sup>+</sup>-ATPase  $\alpha$ -subunit using lactoperoxidase iodination. *Biochim. Biophys. Acta*, 1233 (1995) 175-184.
- 1573 Zavalova, L.L., Kuzina, E.V. and Baskova, I.P.: (Sodium dodecyl sulfate stimulated the appearance of a nonspecific proteolytic activity of destabilase-endo- $\epsilon$ -( $\gamma$ -glutamyl)-lysyl-isopeptidase). *Biokhimiya (Moscow)*, 59 (1994) 905-910.
- See also 1243, 1293.
- 20g. Lyases**
- 1574 Jaffe, E.K., Ali, S., Mitchell, L.W., Taylor, K.M., Volin, M. and Markham, G.D.: Characterization of the role of the stimulatory magnesium of *Escherichia coli* porphobilinogen synthase. *Biochemistry*, 34 (1995) 244-251.
- 1575 Rault-Leonardon, M., Atkinson, M.A.L., Slaughter, C.A., Moomaw, C.R. and Srere, P.A.: *Azotobacter vinelandii* citrate synthase. *Biochemistry*, 34 (1995) 257-263.
- 1576 Sindern, S., van Eldik, R. and Bartholmes, P.: Influence of  $\alpha$ -subunits on the high-pressure stability of apo and holo  $\beta$ -subunits in the bienzyme complex tryptophan synthase from *Escherichia coli*. *Biochemistry*, 34 (1995) 1959-1967.
- 20h. Isomerases**
- See 1250.

## 20i. Ligases

- 1577 Keon, B.H., Ankrapp, D.P. and Keenan, T.W.: Cytosolic lipoprotein particles from milk-secreting cells contain fatty acid synthase and interact with endoplasmic reticulum. *Biochim. Biophys. Acta*, 1215 (1994) 327-336.
- 1578 Petric, R., Nicholson, D.W. and Ford-Hutchinson, A.W.: Renal leukotriene C<sub>4</sub> synthase: characterization, partial purification and alterations in experimental glomerulonephritis. *Biochim. Biophys. Acta*, 1254 (1995) 207-215.

## 20j. Complex mixtures and incompletely identified enzymes

- 1579 Inamoto, S., Fukuda, H., Abo, T. and Ohtsubo, E.: Site- and strand-specific nicking at oriT of plasmid R100 in a purified system: enhancement of the nicking activity of TraI (helicase I) with TraY and IHF. *J. Biochem. (Tokyo)*, 116 (1994) 838-844.
- 1580 King, B.J., Lee, L.S., Rackemann, R.G. and Scott, P.T.: Electrophoretic conditions for high resolution citrus isozymes in polyacrylamide gel electrophoresis. *Electrophoresis (Weinheim)*, 16 (1995) 32-38.
- 1581 Saud, H.B.M. and Jones, D.G.: Multilocus enzyme electrophoresis as a tool for studying genetic diversity in natural populations of *Rhizobium*. *Proc. Malays. Biochem. Soc. Conf.*, 16th (1991) 67-66; *C.A.*, 122 (1995) 4058k.

## 21. PURINES, PYRIMIDINES, NUCLEIC ACIDS AND THEIR CONSTITUENTS

## 21a. Purines, pyrimidines, nucleosides, nucleotides

- 1582 Bourque, A.J. and Cohen, A.S.: Quantitative analysis of phosphorothioate oligonucleotides in biological fluids using direct injection fast anion-exchange chromatography and capillary gel electrophoresis. *J. Chromatogr. B*, 662 (1994) 343-349.
- 1583 Colombatto, S., Giribaldi, G., Vargiu, C. and Grillo, M.A.: Modulation of ornithine aminotransferase activity by oxygen in rat hepatocyte cultures. *Biochim. Biophys. Acta*, 1224 (1994) 329-332.
- 1584 Heller, C. and Viovy, J.L.: Electrophoretic separation of oligonucleotides in replenishable polyacrylamide-filled capillaries. *Appl. Theor. Electrophor.*, 4 (1994) 39-41; *C.A.*, 121 (1994) 250286x.
- 1585 Krattiger, B., Bruno, A.E., Widmer, H.M. and Dändliker, R.: Hologram-based thermooptical absorbance detection in capillary electrophoresis: separation of nucleosides and nucleotides. *Anal. Chem.*, 67 (1995) 124-130.
- 1586 Lappalainen, K., Urtti, A., Söderling, E., Jääskeläinen, I., Syrjänen, K. and Syrjänen, S.: Cationic liposomes improve stability and intracellular delivery of antisense oligonucleotides into CaSki cells. *Biochim. Biophys. Acta*, 1196 (1994) 201-208.
- 1587 Manabe, T., Chen, T., Terabe, S., Yohda, M. and Endo, I.: Effect of linear polyacrylamide concentrations and applied voltages on the separation of oligonucleotides and DNA sequencing fragments by capillary electrophoresis. *Anal. Chem.*, 66 (1994) 4243-4252.
- 1588 Pecina-Slaus, N., Pavelic, J. and Pavelic, K.: Comparison of reverse-phase HPLC and gel electrophoretic purification of synthetic oligonucleotides. *Period. Biol.*, 96 (1994) 161-164; *C.A.*, 121 (1994) 296539k.

- 1589 Rabel, S.R., Stobaugh, J.F. and Trueworthy, R.: Determination of intracellular levels of 6-mercaptopurine metabolites in erythrocytes utilizing capillary electrophoresis with laser-induced fluorescence detection. *Anal. Biochem.*, 224 (1995) 315-322.
- 1590 Van Schepdael, A., Smets, K., Vandendriessche, F., van Aerschot, A., Herdewijn, P., Roets, E. and Hoogmartens, J.: Comparative stability study of thymidine and (dideoxy-D-erythrohexopyranosyl)thymine analogues monitored by capillary electrophoresis. *J. Chromatogr. A*, 687 (1994) 167-173.

See also 1606.

## 21b. Nucleic acids, RNA

- 1591 Ahn, K., Chen, D. and Kemper, B.: Inverse relationship of cotranslational translocation with the hydrophobic moment of the bovine preproparathyroid hormone signal sequence. *Biochim. Biophys. Acta*, 1224 (1994) 459-462.
- 1592 Allen, R.C. and Reeder, D.J.: Matrix modification for electrophoretic separation of nucleic acids or polypeptides. *U.S. US 5,354,442* (Cl. 204-182.8; GO1N27/26), 11 Oct. 1994, Appl. 886,28, 21 May 1992; 16 pp.; *C.A.*, 122 (1995) 4955a.
- 1593 Altieri, D.C.: Splicing of effector cell protease receptor-1 mRNA is modulated by an unusual retained intron. *Biochemistry*, 33 (1994) 13848-13855.
- 1594 Cameron, D.N., Khambaty, F.M., Wachsmuth, I.K., Tauxe, R.V. and Barrett, T.J.: Molecular characterization of *Vibrio cholerae* O1 strains by pulsed-field gel electrophoresis. *J. Clin. Microbiol.*, 32 (1994) 1685-1690; *C.A.*, 121 (1994) 225586n.
- 1595 Cenedella, R.J. and Shi, H.: Spatial distribution of 3-hydroxy-3-methylglutaryl coenzyme A reductase messenger RNA in the ocular lens: relationship to cholesterologenesis. *J. Lipid Res.*, 35 (1994) 2232-2240.
- 1596 Chen, J., Engle, S.J., Seilhamer, J.J. and Tischfield, J.A.: Cloning, expression and partial characterization of a novel rat phospholipase A<sub>2</sub>. *Biochim. Biophys. Acta*, 1215 (1994) 115-120.
- 1597 Deng, C.-I. and Cornett, L.E.: Regulation of  $\alpha_1$ -adrenergic receptor gene expression in rat liver cell lines. *Biochim. Biophys. Acta*, 1219 (1994) 669-676.
- 1598 Einzenberger, E., Eller, N., Heberle-Bors, E. and Vicente, O.: Isolation and expression during pollen development of a tobacco cDNA clone encoding a protein kinase homologous to shaggy/glycogen synthase kinase-3. *Biochim. Biophys. Acta*, 1260 (1995) 315-319.
- 1599 Fukumoto, S., Allan, E.H. and Martin, T.J.: Regulation of plasminogen activator inhibitor-1 (PAI-1) expression by 1,25-dihydroxyvitamin D-3 in normal and malignant rat osteoblasts. *Biochim. Biophys. Acta*, 1201 (1994) 223-228.
- 1600 Gast, F.-U. and Sängler, H.L.: Gel dependence of electrophoretic mobilities of double-stranded and viroid RNA and estimation of the contour length of a viroid by gel electrophoresis. *Electrophoresis (Weinheim)*, 15 (1994) 1493-1498.
- 1601 Gonzales, L.W., Ballard, P.L. and Gonzales, J.: Glucocorticoid and cAMP increase fatty acid synthetase mRNA in human fetal lung explants. *Biochim. Biophys. Acta*, 1215 (1994) 49-58.
- 1602 Grossman, L.I., Rosenthal, N.H., Akamatsu, M. and Erickson, R.P.: Cloning, sequence analysis, and expression of a mouse cDNA encoding cytochrome c oxidase subunit VIa liver isoform. *Biochim. Biophys. Acta*, 1260 (1995) 361-364.

- 1603 Gutierrez, J.A. and Guerriero, V., Jr.: Relative abundance of bovine Hsp70 mRNA and protein. *Biochim. Biophys. Acta*, 1260 (1995) 239-242.
- 1604 Hla, T., Jackson, A.Q., Appleby, S.B. and Maciag, T.: Characterization of edg-2, a human homologue of the *Xenopus* maternal transcript G10 from endothelial cells. *Biochim. Biophys. Acta*, 1260 (1995) 227-229.
- 1605 Holmberg, L. and Nygard, O.: Interaction sites of ribosome-bound eukaryotic elongation factor 2 in 18S and 28S rRNA. *Biochemistry*, 33 (1994) 15159-15167.
- 1606 Hölzl, B., Huber, R., Paulweber, B., Patsch, J.R. and Sandhofer, F.: Lipoprotein lipase deficiency due to a 3'splice site mutation in intron 6 of the lipoprotein lipase gene. *J. Lipid Res.*, 35 (1994) 2161-2169.
- 1607 Ide, H., Itoh, H. and Nawa, Y.: Sequencing of cDNAs encoding  $\alpha$ 1-microglobulin/bikunin of Mongolian gerbil and Syrian golden hamster in comparison with man and other species. *Biochim. Biophys. Acta*, 1209 (1994) 1275-1280.
- 1608 Inokuchi, Y., Kajitani, M. and Hirashima, A.: A study on the function of the glycine residue in the YGDD motif of the RNA-dependent RNA polymerase  $\beta$ -subunit from RNA coliphage Q $\beta$ . *J. Biochem. (Tokyo)*, 116 (1994) 1275-1280.
- 1609 Iwase, K., Yamauchi, K. and Ishikawa, K.: Cloning of cDNAs encoding argininosuccinate lyase and arginase from *Rana catesbeiana* liver and regulation of their mRNAs during spontaneous and thyroid hormone-induced metamorphosis. *Biochim. Biophys. Acta*, 1260 (1995) 139-146.
- 1610 Kaczmarek, E., Liu, Y., Berse, B., Chen, C.S. and McDonagh, J.: Biosynthesis of plasma factor XIII: evidence for transcription and translation in hepatoma cells. *Biochim. Biophys. Acta*, 1247 (1995) 127-134.
- 1611 Kosuga, K., Hata, S., Osumi, T., Sakakibara, J. and Ono, T.: Nucleotide sequence of a cDNA for mouse squalene epoxidase. *Biochim. Biophys. Acta*, 1260 (1995) 345-348.
- 1612 Kotani, E., Yamakawa, M., Iwamoto, S.-i., Tashiro, M., Mori, H., Sumida, M., Matsubara, F., Tani, K., Kadono-Okuda, K., Kato, Y. and Mori, H.: Cloning and expression of the gene of hemocytin, an insect humoral lectin which is homologous with the mammalian von Willebrand factor. *Biochim. Biophys. Acta*, 1260 (1995) 245-258.
- 1613 Kozma, M.M., Chowrimootoo, G., Debnam, E.S., Epstein, O. and Srai, S.K.S.: Developmental changes in mucosal iron binding proteins in the guinea pig. Expression of transferrin, H and L ferritin and binding of iron to a low molecular weight protein. *Biochim. Biophys. Acta*, 1201 (1994) 229-234.
- 1614 Kukuruzinska, M.A. and Lennon, K.: Diminished activity of the first N-glycosylation enzyme, dolichol-P-dependent N-acetylglucosamine-1-P transferase (GPT), gives rise to mutant phenotypes in yeast. *Biochim. Biophys. Acta*, 1247 (1995) 51-59.
- 1615 Li, X. and Chang, Y.-H.: Molecular cloning of a human complementary DNA encoding an initiation factor 2-associated protein (p<sup>67</sup>). *Biochim. Biophys. Acta*, 1260 (1995) 333-336.
- 1616 Määttä, A., Ekholm, E. and Penttinen, R.P.K.: Effect of the 3'-untranslated region on the expression levels and mRNA stability of  $\alpha$ 1(I) collagen gene. *Biochim. Biophys. Acta*, 1260 (1995) 294-300.
- 1617 Mellor, H., Flowers, K.M., Kimball, S.R. and Jefferson, L.S.: Cloning and characterization of a cDNA encoding rat PKR, the double-stranded RNA-dependent eukaryotic initiation factor-2 kinase. *Biochim. Biophys. Acta*, 1219 (1994) 693-696.
- 1618 Mulder, W., Scholten, I.H.J.M., van Roon, H. and Grivell, L.A.: Isolation and characterisation of the linked genes APA2 and QCR7, coding for Ap $\mu$ A phosphorylase II and the 14 kDa subunit VII of the mitochondrial bc<sub>1</sub>-complex in the yeast *Kluyveromyces lactis*. *Biochim. Biophys. Acta*, 1219 (1994) 719-723.
- 1619 Munkonge, F.M., Osborne, L.R., Geddes, D.M. and Alton, E.W.F.W.: Lack of inhibition by dideoxy-forskolin and verapamil of DIDS-sensitive volume-activated Cl<sup>-</sup> secretion in human squamous lung carcinoma epithelial cells. *Biochim. Biophys. Acta*, 1224 (1994) 342-348.
- 1620 Netzker, R., Hermfisse, U., Wein, K.-H. and Brand, K.: Expression of glycolytic isozymes in rat thymocytes during cell cycle progression. *Biochim. Biophys. Acta*, 1224 (1994) 371-376.
- 1621 Nishijima, M., Hara-Kuge, S., Takasuka, N., Akagawa, K., Setouchi, M., Matsuura, K., Yamamoto, S. and Akamatsu, Y.: Identification of a biochemical lesion, and characteristic response to lipopolysaccharide (LPS) of a cultured macrophage-like cell mutant with defective LPS-binding. *J. Biochem. (Tokyo)*, 116 (1994) 1082-1087.
- 1622 Paillart, J.-C., Marquet, R., Skripkin, E., Ehresmann, B. and Ehresmann, C.: Mutational analysis of the bipartite dimer linkage structure of human immunodeficiency virus type 1 genomic RNA. *J. Biol. Chem.*, 269 (1994) 27486-27494.
- 1623 Rozario, C., Smith, M.W. and Müller, M.: Primary sequence of a putative pyrophosphate-linked phosphofructokinase gene of *Giardia lamblia*. *Biochim. Biophys. Acta*, 1260 (1995) 218-222.
- 1624 Saiki, T., Kohno, K., Goshima, N., Kano, Y. and Imamoto, F.: Construction of a gyrA-gyrB-carrying plasmid overproducing functional DNA gyrase. *J. Biochem. (Tokyo)*, 116 (1994) 1208-1211.
- 1625 Sasaki, Y., Usui, T., Tanaka, I., Nakagawa, O., Sando, T., Takahashi, T., Namba, T., Narumiya, S. and Nakao, K.: Cloning and expression of a cDNA for rat prostacyclin receptor. *Biochim. Biophys. Acta*, 1224 (1994) 601-605.
- 1626 Sato, M., Mizobuchi, M., Murao, K., Tamaki, M. and Takahara, J.: Simple quantitative reverse transcribed-polymerase chain reaction (RT-PCR) method involving recombinant RNA generated by a false-priming PCR product. *J. Biochem. (Tokyo)*, 116 (1994) 1198-1201.
- 1627 Schillace, R., Preiss, T., Lightowlers, R.N. and Capaldi, R.A.: Developmental regulation of tissue-specific isoforms of subunit VIa of beef cytochrome c oxidase. *Biochim. Biophys. Acta*, 1188 (1994) 391-397.
- 1628 Schumann, G., Zündorf, I., Schmidt, A., Marschalek, R. and Dingermann, T.: Characterization of transcripts from the *Dicotyostelium discoideum* retrotransposable genetic element DRE. *Pharmazie*, 49 (1994) 923-925.
- 1629 Selvamurugan, N., Nag, M.K. and Eliceiri, G.L.: Intron-encoded small nucleolar RNAs: new RNA sequence variants and genomic loci. *Biochim. Biophys. Acta*, 1260 (1995) 230-234.
- 1630 Sharma, H.S., Tang, Z.H., Gho, B.C.G. and Verdouw, P.D.: Nucleotide sequence and expression of the porcine vascular endothelial growth factor. *Biochim. Biophys. Acta*, 1260 (1995) 235-238.
- 1631 Strotkamp, D., Roos, P.H. and Hanstein, W.G.: A novel CYP3 gene from female rats. *Biochim. Biophys. Acta*, 1260 (1995) 341-344.

- 1632 Sun, H.B., Neff, A.W., Mescher, A.L. and Malacinski, G.M.: Expression of the axolotl homologue of mouse chaperonin *t*-complex protein-1 during early development. *Biochim. Biophys. Acta*, 1260 (1995) 157-166.
- 1633 Tada, K., Fukunaga, T., Wakabayashi, Y., Masumi, S., Sato, Y., Izumi, H., Kohno, K. and Kuwano, M.: Inhibition of tubular morphogenesis in human microvascular endothelial cells by co-culture with chondrocytes and involvement of transforming growth factor  $\beta$ : a model for avascularity in human cartilage. *Biochim. Biophys. Acta*, 1201 (1994) 135-142.
- 1634 Takimoto, K., Gealy, R. and Levitan, E.S.: Multiple protein kinases are required for basal Kv1.5 K<sup>+</sup> channel gene expression in GH<sub>3</sub> clonal pituitary cells. *Biochim. Biophys. Acta*, 1265 (1995) 22-28.
- 1635 Takimoto, Y. and Kuramoto, A.: Mechanism of regulation of PDGF-A chain gene expression by serum and TPA. *Biochim. Biophys. Acta*, 1260 (1995) 176-182.
- 1636 Tasken, K.A., Jacobsen, F.W., Eikvar, L., Hansson, V. and Haugen, T.B.: The  $\alpha$ -subunit mRNAs for G<sub>s</sub> and G<sub>o2</sub> are differentially regulated by protein kinase A and protein kinase C in rat Sertoli cells. *Biochim. Biophys. Acta*, 1260 (1995) 269-275.
- 1637 Tsuji, S., Kaji, K. and Nagasawa, S.: Activation of the alternative pathway of human complement by apoptotic human umbilical vein endothelial cells. *J. Biochem. (Tokyo)*, 116 (1994) 794-800.
- 1638 Wetterauer, B.W., Hamker, U., von Haeseler, A., MacWilliams, H.K., Simon, M.-N. and Veron, M.: A protein kinase from *Dicystoteliu discoideu* with an unusual acidic repeat domain. *Biochim. Biophys. Acta*, 1265 (1995) 97-101.
- 1639 Yamamoto, Y., Takimoto, K., Izumi, S., Toriyama-Sakurai, M., Kageyama, T. and Takahashi, S.Y.: Molecular cloning and sequencing of cDNA that encodes cysteine proteinase in the eggs of the silkworm, *Bombyx mori*. *J. Biochem. (Tokyo)*, 116 (1994) 1330-1335.
- 1640 Yanagimoto, T., Itoh, S., Sawada, M., Hashimoto, H. and Kamataki, T.: Molecular cloning and functional expression of a mouse cytochrome P-450 (Cyp3a-13): examination of Cyp3a-13 enzyme to activate aflatoxin B<sub>1</sub> (AFB<sub>1</sub>). *Biochim. Biophys. Acta*, 1201 (1994) 405-410.
- 1641 Yohn, J.J., Smith, C., Stevens, T., Morelli, J.G., Shurnas, L.R., Walchak, S.J., Hoffman, T.A., Kelley, K.K., Escobedo-Morse, A., Yanagisawa, M. et al.: Autoregulation of endothelin-1 secretion by cultured human keratinocytes via the endothelin B receptor. *Biochim. Biophys. Acta*, 1224 (1994) 454-458.
- 1642 Zhou, Y., Lin, G. and Murtaugh, M.P.: Interleukin-4 suppresses the expression of macrophage NADPH oxidase heavy chain subunit (gp91-phox). *Biochim. Biophys. Acta*, 1265 (1995) 40-48.
- See also 1095, 1192, 1314, 1438, 1529, 1656, 1662, 1749.
- 21c. *Nucleic acids, DNA*
- 1643 Arakawa, H., Maeda, M. and Tsuji, A.: (Gene diagnosis by capillary electrophoresis). *Farumashia*, 30 (1994) 998-1003; *C.A.*, 121 (1994) 249951d - a review with 23 refs.
- 1644 Baba, Y.: (Design of HPCE capillary columns for high-resolution analysis of DNA). *Kuromatogurafi*, 15 (1994) 106-107; *C.A.*, 121 (1994) 275642y.
- 1645 Barron, A.E., Sunada, W.M. and Blanch, H.W.: The use of coated and uncoated capillaries for the electrophoretic separation of DNA in dilute polymer solutions. *Electrophoresis (Weinheim)*, 16 (1995) 64-74.
- 1646 Bolstad, A.I.: Sizing the *Fusobacterium nucleatum* genome by pulsed-field gel electrophoresis. *FEMS Microbiol. Lett.*, 123 (1994) 145-152; *C.A.*, 121 (1994) 294234h.
- 1647 Cassidy, S.A., Strekowski, L., Wilson, W.D. and Fox, K.R.: Effect of triplex-binding ligand on parallel and antiparallel DNA triple helices using short unmodified and acridine-linked oligonucleotides. *Biochemistry*, 33 (1994) 15338-15347.
- 1648 Chachaty, E., Saulnier, P., Martin, A., Mario, N. and Andremon, A.: Comparison of ribotyping, pulsed-field gel electrophoresis and random amplified polymorphic DNA for typing *Clostridium difficile* strains. *FEMS Microbiol. Lett.*, 122 (1994) 61-68; *C.A.*, 121 (1994) 273185c.
- 1649 Clark, B.K., Vo-Dinh, T. and Sepaniak, M.J.: Capillary electrophoresis-laser fluorometry instrumentation for the facile optimization of DNA separations using *in situ* size-selective gradients and adjustable detection zone. *Anal. Chem.*, 67 (1995) 680-683.
- 1650 Cooney, C.A.: Techniques and high resolution DNA size markers for pulsed field gel electrophoresis. *Mol. Biotechnol.*, 2 (1994) 119-127; *C.A.*, 122 (1995) 4745g.
- 1651 Correia, A., Martin, J.F. and Castro, J.M.: Pulsed-field gel electrophoresis analysis of the genome of amino acid producing corynebacteria: chromosome sizes and diversity of restriction patterns. *Microbiology (Reading)*, 140 (1994) 2841-2847; *C.A.*, 122 (1995) 2089x.
- 1652 Couette, B., le Ricousse, S., Fortin, D., Rafestin-Oblin, M.E. and Richard-Foy, H.: The establishment of the long terminal repeat of the mouse mammary tumor virus into CV-1 cells allows a functional analysis of steroid receptors. *Biochim. Biophys. Acta*, 1219 (1994) 607-612.
- 1653 Esposito, F., Cuccovillo, F., Morra, F., Russo, T. and Cimino, F.: DNA binding activity of the glucocorticoid receptor is sensitive to redox changes in intact cells. *Biochim. Biophys. Acta*, 1260 (1995) 308-314.
- 1654 Fernandez-Busquets, X. and Daban, J.-R.: Histones associated with single-stranded DNA do not preclude the formation of double-helical DNA. *Biochim. Biophys. Acta*, 1260 (1995) 132-138.
- 1655 Figey, D., Renborg, A. and Dovichi, N.J.: Labeling of double-stranded DNA by ROX-dideoxycytosine triphosphate using terminal deoxynucleotidyl transferase and separation by capillary electrophoresis. *Anal. Chem.*, 66 (1994) 4382-4383.
- 1656 Furuya, K.N., Gebhardt, R., Schuetz, E.G. and Schuetz, J.D.: Isolation of rat pgp3 cDNA: evidence for gender and zonal regulation of expression in the liver. *Biochim. Biophys. Acta*, 1219 (1994) 636-644.
- 1657 Gong, Z. and Hew, C.L.: Several splicing variants of isl-1 like genes in the chinook salmon (*Oncorhynchus tshawytscha*) encode truncated transcription factors containing a complete LIM domain. *Biochim. Biophys. Acta*, 1260 (1995) 349-354.
- 1658 Griesbach, R.J.: An improved method for transforming plants through electrophoresis. *Plant. Sci. (Shannon)*, 102 (1994) 81-89; *C.A.*, 121 (1994) 247420f.

- 1659 Guttman, A.: Separation of DNA by capillary electrophoresis. In: Landers, J.P. (Editor), *Handb. Capillary Electrophor.*, CRC, Boca Raton, 1994, pp. 185-206; C.A., 121 (1994) 249918y - a review with 55 refs.
- 1660 Hofman, Y.L., Petrillo, K.L., Greenblatt, H.C., Lehrer, R. and Payne, M.S.: Rapid scaleable chromatographic purification of nucleic acids from proteinaceous mixtures. *J. Chromatogr. A*, 692 (1995) 233-238.
- 1661 Huberman, J.A.: Analysis of DNA replication origins and directions by two-dimensional gel electrophoresis. In: Fantes, P. and Brooks, R. (Editors), *Cell Cycle*, IRL, Oxford, 1994, pp. 213-234; C.A., 121 (1994) 250275t.
- 1662 Kamahori, M. and Yamada, T.: Capillary electrophoresis of nucleic acids. *Eur. Pat. Appl. EP 607,495* (Cl. G01N27/447), 27 Jul. 1994, JP Appl. 92/226,894, 26 Aug. 1992; 24 pp.; C.A., 121 (1994) 225859d.
- 1663 Kimura, M., Takatsuki, A. and Yamaguchi, I.: Blastidicin S deaminase gene from *Aspergillus terreus* (BSD): a new drug resistance gene for transfection of mammalian cells. *Biochim. Biophys. Acta*, 1219 (1994) 653-659.
- 1664 Kodaira, K.-I., Godson, N.G. and Taketo, A.: Comparative studies on the minus origin mutants of *Escherichia coli* spherical single-stranded DNA phages. *Biochim. Biophys. Acta*, 1260 (1995) 191-199.
- 1665 Kodaira, K.-I., Oki, M., Taketo, A., Yasukawa, H. and Masamune, Y.: Determination of the single strand origin of *Shigella sonnei* plasmid pKYM. *Biochim. Biophys. Acta*, 1260 (1995) 183-190.
- 1666 Komiyama, M. and Inokawa, T.: Selective hydrolysis of tRNA by ethylenediamine bound to a DNA oligomer. *J. Biochem. (Tokyo)*, 116 (1994) 719-720.
- 1667 Krajewski, W.A., Panin, V.M. and Razin, S.V.: (Temperature dependence of chromatin condensation at different ionic strength of the medium: a study by low-density agarose gel electrophoresis). *Biofizika*, 39 (1994) 619-627; C.A., 121 (1994) 276081b.
- 1668 Lee, H.H.: An inexpensive cooling fan for SSCP gel electrophoresis. *BioTechniques*, 16 (1994) 790; C.A., 121 (1994) 293869g.
- 1669 Lim, L. and Howlett, B.J.: Linear plasmids, pLm9 and pLm10, can be isolated from the phytopathogenic ascomycete *Leptosphaeria maculans* by pulsed-field gel electrophoresis. *Curr. Genet.*, 26 (1994) 276-280; C.A., 121 (1994) 250330g.
- 1670 Locke, J. and McDermid, H.E.: Analysis of *Drosophila* chromosome 4 using pulsed field gel electrophoresis. *Chromosoma*, 102 (1993) 718-723; C.A., 121 (1994) 247520p.
- 1671 Mäueler, W., Kyas, A., Bröcker, F. and Epplen, J.T.: Altered electrophoretic behavior of DNA due to short-time UV-B irradiation. *Electrophoresis (Weinheim)*, 15 (1994) 1499-1505.
- 1672 Maule, J.: Electrophoretic karyotype analysis: pulsed field gel electrophoresis. *Methods Mol. Biol. (Totowa)*, 29(Chromosome Analysis Protocols) (1994) 221-252; C.A., 122 (1995) 2036c.
- 1673 Mickey, G., Sim, J., Durand, J. and Wills, C.: Use of denaturing gradient electrophoresis to determine the distributions of polymorphisms in entire genes in natural populations. *Biochim. Biophys. Acta*, 1260 (1995) 123-131.
- 1674 Mulder, W., Scholten, I.H.J.M., Nagelkerken, B. and Grivell, L.A.: Isolation and characterisation of the linked genes, FPS1 and QCR8, coding for farnesyl-diphosphate synthase and the 11 kDa subunit VIII of the mitochondrial bc<sub>1</sub>-complex in the yeast *Kluyveromyces lactis*. *Biochim. Biophys. Acta*, 1219 (1994) 713-718.
- 1675 Pignon, J.M., Vinatier, I., Fanen, P., Jonveaux, P., Tournilhac, O., Imbert, M., Rochant, H. and Goossens, M.: Exhaustive analysis of the p53 gene coding sequence by denaturing gradient gel electrophoresis: Application to the detection of point mutations in acute leukemias. *Hum. Mutat.*, 3 (1994) 126-132; C.A., 121 (1994) 293858c.
- 1676 Reese, H.R.: Effects of DNA charge and length on the electrophoretic mobility of intercalated DNA. *Biopolymers*, 34 (1994) 1349-1358; C.A., 122 (1995) 4737f.
- 1677 Revoltella, R.P., Dal Canto, B., Caracciolo, L. and D'Urso, C.M.: L-Carnitine and some of its analogs delay the onset of apoptotic cell death initiated in murine C2.8 hepatocytic cells after hepatocyte growth factor deprivation. *Biochim. Biophys. Acta*, 1224 (1994) 333-341.
- 1678 Riveron Rojas, A.M., Lopez-Canovas, L., Herrera-Isidron, J., Ruiz-Esquivel, L., Higginson-Clarke, D., Blanco, M.N. and Valdes-Acosta, F.: Molecular weight and kinetic parameters of DNA undergoing pulsed field gel electrophoresis. *Arch. Med. Res.*, 25 (1994) 193-198; C.A., 121 (1994) 250282t.
- 1679 Routman, E. and Cheverud, J.: A rapid method of scoring simple sequence repeat polymorphisms with agarose gel electrophoresis. *Mamm. Genome*, 5 (1994) 187-188; C.A., 121 (1994) 247299y.
- 1680 Satow, T.: (Analysis of DNA by capillary electrophoresis). *Bunseki Kagaku*, 43 (1994) 731-743; C.A., 121 (1994) 296207a - a review with 58 refs.
- 1681 Shuman, S.: Novel approach to molecular cloning and polynucleotide synthesis using vaccinia DNA topoisomerase. *J. Biol. Chem.*, 269 (1994) 32678-32684.
- 1682 Slater, G.W., Mayer, P. and Grossman, P.D.: Diffusion, Joule heating, and band broadening in capillary gel electrophoresis of DNA. *Electrophoresis (Weinheim)*, 16 (1995) 75-83.
- 1683 Sugano, K., Fukayama, N., Ohkura, H., Shimosato, Y., Yamada, Y., Inoue, T., Sekiya, T. and Hayashi, K.: Single-strand conformation polymorphism analysis by perpendicular temperature-gradient gel electrophoresis. *Electrophoresis (Weinheim)*, 16 (1995) 8-10.
- 1684 Suni, Y. and Rigas, B.: Detection of biotinylated nucleic acids directly on agarose gels. *BioTechniques*, 16 (1994) 782-784; C.A., 121 (1994) 296407r.
- 1685 Takasaki, Y.: Two forms of restriction enzyme HindIII. *J. Biochem. (Tokyo)*, 116 (1994) 1281-1286.
- 1686 Tan, X., Sun, X., Gonzalez-Crussi, F.X., Gonzalez-Crussi, F. and Hsueh, W.: PAF and TNF increase the precursor of NF-kappa B p50 mRNA in mouse intestine: quantitative analysis by competitive PCR. *Biochim. Biophys. Acta*, 1215 (1994) 157-162.
- 1687 Terajima, M., Matsui, Y., Copeland, N.G., Gilbert, D.J., Jenkins, N.A. and Obinata, M.: Structural organization of the mouse glycophorin A gene. *J. Biochem. (Tokyo)*, 116 (1994) 1105-1110.
- 1688 Venditti, P., Costanzo, G., Negri, R. and Camilloni, G.: ABFI contributes to the chromatin organization of *Saccharomyces cerevisiae* ARS1 B-domain. *Biochim. Biophys. Acta*, 1219 (1994) 677-689.

- 1689 Wang, M. and Lai, E.: Pulsed field separation of large supercoiled and open-circular DNAs and its application to bacterial artificial chromosome cloning. *Electrophoresis (Weinheim)*, 16 (1995) 1-7.
- 1690 Zakharov, S.F., Garner, M.M. and Chrambach, A.: Fluorescent labeling of DNA with ethidium homodimer without measurable decrease in DNA mobility: application to automated gel electrophoresis apparatus. *Anal. Biochem.*, 224 (1995) 195-198.
- See also 1095, 1314, 1366, 1498, 1606, 1610, 1616, 1630, 1637, 1642, 1730.
- 21d. Structural studies on RNA and RNA mapping**
- 1691 Annoni, G., Arosio, B., Santambrogio, D., Cullura, D., Gagliano, N. and Uslenghi, C.: Gene expression for interleukin-2 and tumor necrosis factor- $\alpha$  in the spleen of old rats under physiological condition and during septic shock. Possible pharmacological modulation. *Arzneim.-Forsch.*, 44 (1994) 1433-1436.
- 1692 Ichiara, J., Shinohara, Y., Kogure, K. and Terada, H.: Nucleotide sequence of the 5'-flanking region of the rat type II hexokinase gene. *Biochim. Biophys. Acta*, 1260 (1995) 365-368.
- 1693 Lin, A. and Huang, R.-G.: Hydrolysis of ribosomes by electrophoretically blotted ribotoxin. *BioTechniques*, 17 (1994) 636-638; *C.A.*, 121 (1994) 276061v.
- 1694 Matsubara, T., Takiguchi, M., Matsuda, I. and Mori, M.: Rat argininosuccinate lyase promoter: the dyad-symmetric CCAAT box sequence CCAATTGG in the promoter is recognized by NF- $\gamma$ . *J. Biochem. (Tokyo)*, 116 (1994) 1044-1055.
- 1695 Trackman, P.C., Feres-Filho, E.J. and Choi, Y.-J.: The 3'-untranslated region of rat lysyl oxidase cDNA. *Biochim. Biophys. Acta*, 1260 (1995) 355-360.
- 1696 Uchiyama, Y., Iwai, S., Ueno, Y., Ikehara, M. and Ohtsuka, E.: Role of the Mg<sup>2+</sup> ion in the *Escherichia coli* ribonuclease HI reaction. *J. Biochem. (Tokyo)*, 116 (1994) 1322-1329.
- 1697 Yamada, K., Kato, H., Kanda, N., Fujii-Kuriyama, Y., Utakoji, T. and Itoh, R.: Sequence homology of Chinese hamster metallothionein genes I and II to those of the mouse and rat, and their amplification in Cd-resistant cells. *Biochim. Biophys. Acta*, 1219 (1994) 581-591.
- 21e. Structural studies on DNA and DNA mapping**
- 1698 Appel, R.D., Sanchez, J.-C., Bairoch, A., Golaz, O., Ravier, F., Pasquali, C., Hughes, G.J. and Hochstrasser, D.F.: The SWISS-2DPAGE database of two-dimensional polyacrylamide gel electrophoresis. *Nucleic Acids Res.*, 22 (1994) 3581-3582; *C.A.*, 121 (1994) 273824k.
- 1699 Arai, T., Tsukada, T. and Nakayama, T.: Simple and rapid detection of cholesterol ester transfer protein deficiency by using nonradioisotopic single-strand conformation polymorphism analysis. *Clin. Chem. (Washington)*, 40 (1994) 2227-2229.
- 1700 Asakawa, J.-i., Duick, R., Neel, J.V., Kodaira, M., Satoh, C. and Hanash, S.M.: Genetic variation detected by quantitative analysis of end-labeled genomic DNA fragments. *Proc. Natl. Acad. Sci. U.S.A.*, 91 (1994) 9052-9056; *C.A.*, 121 (1994) 247343h.
- 1701 Asakawa, J.-i., Kuick, R., Neel, J.V., Kodaira, M., Satoh, C. and Hanash, S.M.: Quantitative and qualitative genetic variation in two-dimensional DNA gels of human lymphocytoid cell lines. *Electrophoresis (Weinheim)*, 16 (1995) 241-252.
- 1702 Best, N., Arriaga, E., Chen, D.Y. and Dovichi, N.J.: Separation of fragments up to 570 bases in length by use of 6% T non-cross-linked polyacrylamide for DNA sequencing in capillary electrophoresis. *Anal. Chem.*, 66 (1994) 4063-4067.
- 1703 Bowie, L.J., Reddy, P.L., Nagabhushan, M. and Sevigny, P.: Detection of  $\alpha$ -thalassemias by multiplex polymerase chain reaction. *Clin. Chem. (Washington)*, 40 (1994) 2260-2266.
- 1704 Brilliant, M.H., Gondo, Y. and Magliocco, A.: One-dimensional genome scanning: Identification of the basis of a mouse mutation and identification of genomic changes in ovarian carcinoma. *Electrophoresis (Weinheim)*, 16 (1995) 163-167.
- 1705 Brossette, S. and Wartell, R.M.: A program for selecting DNA fragments to detect mutations by denaturing gel electrophoresis methods. *Nucleic Acids Res.*, 22 (1994) 4321-4325; *C.A.*, 122 (1995) 2116d.
- 1706 Chen, J. and Thilly, W.G.: Use of denaturing-gradient gel electrophoresis to study chromium-induced point mutations in human cells. *Environ. Health Perspect. Suppl.*, 102(Suppl. 3) (1994) 227-229; *C.A.*, 121 (1994) 294821x.
- 1707 Cheng, J. and Mitchelson, K.R.: Glycerol-enhanced separation of DNA fragments in entangled solution capillary electrophoresis. *Anal. Chem.*, 66 (1994) 4210-4214.
- 1708 Chiquero, M.J., Olmo, A., Navarro, P., Ruiz-Perez, L., Castanys, S., Gonzalez-Pacanowska, D. and Gamarro, F.: Amplification of the H locus in *Leishmania infantum*. *Biochim. Biophys. Acta*, 1227 (1994) 188-194.
- 1709 Clark, B.K., Nickles, C.L., Morton, K.C., Kovac, J. and Sepaniak, M.J.: Rapid separation of DNA restriction digests using size selective capillary electrophoresis with application to DNA fingerprinting. *J. Microcolumn Sep.*, 6 (1994) 503-513; *C.A.*, 121 (1994) 275912m.
- 1710 Cole, D.E.C., Clarke, L.A., Riddell, D.C., Samson, K.A., Seltzer, W.K. and Salisbury, S.: Congenital adrenal hypoplasia, Duchenne muscular dystrophy, and glycerol kinase deficiency: importance of laboratory investigations in delineating a contiguous gene deletion syndrome. *Clin. Chem. (Washington)*, 40 (1994) 2099-2103.
- 1711 Dallinga-Thie, G.M., van Linde-Sibenius Trip, M., Kock, L.A.W. and de Bruin, T.W.A.: Apolipoprotein E2/E3/E4 genotyping with agarose gels. *Clin. Chem. (Washington)*, 41 (1995) 73-75.
- 1712 Dorval, I., Fergelot, P., Dubourg, C., Jezequel, P., Chauvel, B., le Gall, J.Y. and le Treut, A.: Rapid detection of 1078 delT mutation by PCR-mediated site-directed mutagenesis: detection of cystic fibrosis carriers in a Celtic population. *Clin. Chem. (Washington)*, 40 (1994) 2318-2319.
- 1713 Figeys, D., Renborg, A. and Dovichi, N.J.: Spatial and temporal depletion of ions from noncrosslinked denaturing polyacrylamide in capillary electrophoresis. *Electrophoresis (Weinheim)*, 15 (1994) 1512-1517.
- 1714 Fodde, R. and Losekoot, M.: Mutation detection by denaturing gradient gel electrophoresis (DGGE). *Hum. Mutat.*, 3 (1994) 83-94; *C.A.*, 121 (1994) 273043e - a review with 93 refs.
- 1715 Gelfi, C., Righetti, P.G., Cremonesi, L. and Ferrari, M.: Detection of point mutations by capillary electrophoresis in liquid polymers in temporal thermal gradients. *Electrophoresis (Weinheim)*, 15 (1994) 1506-1511.

- 1716 Gomada, M., Imaishi, H., Miura, K., Inouye, S., Nakazawa, T. and Nakazawa, A.: Analysis of DNA bend structure of promoter regulatory regions of xylene-metabolizing genes on the *Pseudomonas* TOL plasmid. *J. Biochem. (Tokyo)*, 116 (1994) 1096-1104.
- 1717 Gondo, Y.: Size-selected genomic libraries: The distribution and size-fractionation of restricted genomic DNA fragments by gel electrophoresis. *Electrophoresis (Weinheim)*, 16 (1995) 168-173.
- 1718 Gondo, Y. and Brilliant, M.H.: Theoretical basis of one-dimensional genome scanning: A direct method to identify the site of a mutation. *Electrophoresis (Weinheim)*, 16 (1995) 174-178.
- 1719 Guida, T., Marger, R.S., Papp, A.C., Snyder, P.J., Sedra, M.S., Kissel, J.T., Mendell, J.R. and Prior, T.W.: A molecular protocol for diagnosing myotonic dystrophy. *Clin. Chem. (Washington)*, 41 (1995) 69-72.
- 1720 Hara, A., Zhang, W., Kobayashi, H., Niikawa, S., Sakai, N. and Yamada, H.: A single cell gel electrophoresis technique for the detection of DNA damage induced by ACNU, an alkylating agent or irradiation in murine glioma cell lines. *Neurol. Res.*, 16 (1994) 234-240; C.A., 121 (1994) 294494t.
- 1721 Hitotsumatsu, T., Kitamoto, T., Iwaki, T., Fukui, M. and Tateishi, J.: An exon 8-spliced out transcript of neurofibromatosis 2 gene is constitutively expressed in various human tissues. *J. Biochem. (Tokyo)*, 116 (1994) 1205-1207.
- 1722 Hori, H., Kanamori, T., Mizuta, T., Yamaguchi, N., Liu, Y. and Nagai, Y.: Human laminin M chain: epitope analysis of its monoclonal antibodies by immunoscreening of cDNA clones and tissue expression. *J. Biochem. (Tokyo)*, 116 (1994) 1212-1219.
- 1723 Jeffers, M. and Pellicer, A.: Identification of multiple promoters within the N-ras proto-oncogene. *Biochim. Biophys. Acta*, 1219 (1994) 623-635.
- 1724 Joshi, R.R. and Ganesh, K.N.: Duplex and triplex directed DNA cleavage by oligonucleotide-Cu(II)/Co(III) metallodesferal conjugates. *Biochim. Biophys. Acta*, 1201 (1994) 454-460.
- 1725 Kapelner, S.N., Turner, R.T., Sarkar, G. and Bolander, M.E.: Deletion mutation can be an unsuspected gel artifact. *BioTechniques*, 17 (1994) 64-67; C.A., 121 (1994) 293895n.
- 1726 Kok, R.G., Christoffels, V.M., Vosman, B. and Hellingwerf, K.J.: A gene of *Acinetobacter calcoaceticus* BD413 encodes a periplasmic peptidyl-prolyl *cis-trans* isomerase of the cyclophilin sub-class that is not essential for growth. *Biochim. Biophys. Acta*, 1219 (1994) 601-606.
- 1727 Krawczak, M. and Lubjuhn, T.: An informativity index for multilocus DNA fingerprints. *Electrophoresis (Weinheim)*, 16 (1995) 16-21.
- 1728 Lueders, K.K.: Multilocus genomic mapping with intracisternal A-particle proviral oligonucleotide probes hybridized to mouse DNA in dried agarose gels. *Electrophoresis (Weinheim)*, 16 (1995) 179-185.
- 1729 Matsuda, Y. and Chapman, V.M.: Application of fluorescence *in situ* hybridization in genome analysis of the mouse. *Electrophoresis (Weinheim)*, 16 (1995) 261-272.
- 1730 Mischiati, C., Feriotto, G., Fiorentino, D. and Gambari, R.: Chromatography in DNA radiolabeling: hands-off automation using a robotic workstation. *J. Chromatogr. B*, 664 (1995) 303-310.
- 1731 Miwa, W., Yashima, K., Sekine, T. and Sekiya, T.: Demethylation of a repetitive DNA sequence in human cancers. *Electrophoresis (Weinheim)*, 16 (1995) 227-232.
- 1732 Musarrat, J., Arezina, J., Uddin, S. and Wani, A.A.: Induction and processing of promutagenic O<sup>6</sup>-ethylthymine lesion in specific gene segments of plasmid DNA. *Biochim. Biophys. Acta*, 1260 (1995) 276-284.
- 1733 Nelson, S.F.: Genomic mismatch scanning: Current progress and potential applications. *Electrophoresis (Weinheim)*, 16 (1995) 279-285.
- 1734 Ohsumi, T., Okazaki, Y., Hirotsune, S., Shibata, H., Muramatsu, M., Suzuki, H., Taga, C., Watanabe, S. and Hayashizaki, Y.: A spot cloning method for restriction landmark genomic scanning. *Electrophoresis (Weinheim)*, 16 (1995) 203-209.
- 1735 Okazaki, Y., Okuizumi, H., Sasaki, N., Ohsumi, T., Kuromitsu, J., Hirota, N., Muramatsu, M. and Hayashizaki, Y.: An expanded system of restriction landmark genomic scanning (RLGS Ver. 1.8). *Electrophoresis (Weinheim)*, 16 (1995) 197-202.
- 1736 Okuizumi, H., Okazaki, Y., Ohsumi, T., Hanami, T., Mizuno, Y., Muramatsu, M., Hayashizaki, Y., Plass, C. and Chapman, V.M.: A single gel analysis of 575 dominant and codominant restriction landmark genomic scanning loci in mice interspecific back-cross progeny. *Electrophoresis (Weinheim)*, 16 (1995) 253-260.
- 1737 Okuizumi, H., Okazaki, Y., Ohsumi, T., Hayashizaki, Y., Plass, C. and Chapman, V.M.: Genetic mapping of restriction landmark genomic scanning loci in the mouse. *Electrophoresis (Weinheim)*, 16 (1995) 233-240.
- 1738 Rozé, A., Zakim, M. and Duchange, N.: Regulation of melanotransferrin gene in melanoma cells. Analysis of the enhancer region. *Biochim. Biophys. Acta*, 1260 (1995) 301-307.
- 1739 Sagami, I., Kikuchi, H., Ikawa, S. and Watanabe, M.: Characterization of hamster CYP1A1 gene: inducible expression and negative regulation. *J. Biochem. (Tokyo)*, 116 (1994) 801-810.
- 1740 Sakuma, Y. and Nishigaki, K.: Computer prediction of general PCR products based on dynamical solution structures of DNA. *J. Biochem. (Tokyo)*, 116 (1994) 736-741.
- 1741 Shibata, H., Yoshino, K., Muramatsu, M., Plass, C., Chapman, V.M. and Hayashizaki, Y.: The use of restriction landmark genomic scanning to scan the mouse genome for endogenous loci with imprinted patterns of methylation. *Electrophoresis (Weinheim)*, 16 (1995) 210-217.
- 1742 Steffens, L., Franke, S., Nickel, S., Schwarzkopf, A., Flemmig, T.F. and Karch, H.: DNA fingerprinting of *Eikenella corrodens* by pulsed-field gel electrophoresis. *Oral Microbiol. Immunol.*, 9 (1994) 95-98; C.A., 122 (1995) 2024x.
- 1743 Stone, E.M., Vandenburgh, K., Nichols, B.E. and Sheffield, V.C.: Identification of rhodopsin gene mutations using GC-clamped denaturing gradient gel electrophoresis. *Methods Neurosci.*, 15(Photoreceptor Cells) (1993) 377-392; C.A., 122 (1995) 1979a.
- 1744 Swerdlow, H., Dew-Jager, K. and Gesteland, R.F.: Reloading and stability of polyacrylamide slab gels for automated DNA sequencing. *BioTechniques*, 16 (1994) 688-693; C.A., 121 (1994) 273112b.
- 1745 Tomisaki, R., Baba, Y., Tshako, M., Takahashi, S., Murakami, K., Anazawa, T. and Kambara, H.: High-speed DNA-sequencer using capillary gel electrophoresis with a laser-induced four-color fluorescent DNA detector. *Anal. Sci.*, 10 (1994) 817-820; C.A., 121 (1994) 293919y.

- 1746 Tsai, M.Y., Hanson, N.Q., Copeland, K.R., Beheshti, I. and Garg, U.: Determination of a T/G polymorphism at nucleotide 3206 of the apolipoprotein C III gene by amplification refractory mutation system. *Clin. Chem. (Washington)*, 40 (1994) 2235-2239.
- 1747 Uitterlinden, A.G.: Gene and genome scanning by two-dimensional DNA typing. *Electrophoresis (Weinheim)*, 16 (1995) 186-196.
- 1748 Van Mechelen, J.R., Smits, M., Douma, A.C., Rouster, J., Cameron-Mills, V., Heidekamp, F. and Valk, B.E.: Primary structure of a lipoxygenase from barley grain as deduced from its cDNA sequence. *Biochim. Biophys. Acta*, 1254 (1995) 221-225.
- 1749 Watanabe, S., Kawai, J., Hirotsune, S., Suzuki, H., Hirose, K., Taga, C., Ozawa, N., Fushiki, S. and Hayashizaki, Y.: Accessibility to tissue-specific genes from methylation profiles of mouse brain genomic DNA. *Electrophoresis (Weinheim)*, 16 (1995) 218-226.
- 1750 Windle, B., Silvas, E. and Parra, I.: High resolution microscopic mapping of DNA using multicolor fluorescent hybridization. *Electrophoresis (Weinheim)*, 16 (1995) 273-278.
- 1751 Yokota, H., Amano, S., Yamane, T., Ataka, K., Kikuya, E. and Oishi, M.: Differential cloning using in-gel competitive reassociation. *Electrophoresis (Weinheim)*, 16 (1995) 286-290.
- 1752 Yoshioka, H., Greenwel, P., Inoguchi, K., Truter, S., Inagaki, Y., Ninomiya, Y. and Ramirez, F.: Structural and functional analysis of the promoter of the human  $\alpha 1(XI)$  collagen gene. *J. Biol. Chem.*, 270 (1995) 418-424.
- See also 1107, 1145, 1295, 1498, 1587, 1629, 1632, 1663, 1673, 1675, 1688, 1695, 1697.
22. ALKALOIDS
- 1753 Rodopoulos, N. and Norman, A.: Determination of caffeine and its metabolites in urine by high-performance liquid chromatography and capillary electrophoresis. *Scand. J. Clin. Lab. Invest.*, 54 (1994) 305-315; C.A., 121 (1994) 220894b.
23. OTHER SUBSTANCES CONTAINING HETEROCYCLIC NITROGEN
- 23a. *Porphyryns and other pyrroles*
- See 1763.
- 23e. *Other N-heterocyclic compounds*
- 1754 Galceran, M.T., Curto, M.J., Puignou, L. and Moyano, E.: Determination of acridine derived compounds in charcoal-grilled meat and creosote oils by liquid chromatographic and gas chromatographic analysis. *Anal. Chim. Acta*, 295 (1994) 307-313.
- 1755 Ong, C.P., Ng, C.L., Lee, H.K. and Li, S.F.Y.: Separation of imidazole and its derivatives by capillary electrophoresis. *J. Chromatogr. A*, 686 (1994) 319-324.
- See also 1188, 1807.
24. ORGANIC SULPHUR COMPOUNDS (INCL. GLUCOSINOLATES)
- 1756 Stamler, J. and Loscalzo, J.: Capillary zone electrophoresis of thiols and their derivatives. *PCT Int. Appl. WO 94 17,398* (Cl. GO1N27/26), 04 Aug. 1994, US Appl. 9,469, 26 Jan. 1993; 39 pp.; C.A., 121 (1994) 225863a.
- See also 1120.
25. ORGANIC PHOSPHORUS COMPOUNDS (INCL. SUGAR PHOSPHATES)
- 1757 Oehrle, S.A. and Bossie, P.C.: Analysis of nerve agent degradation products using capillary ion electrophoresis. *J. Chromatogr. A*, 692 (1995) 247-252.
- See also 1215, 1221, 1236, 1305, 1357, 1582.
26. ORGANOMETALLIC AND RELATED COMPOUNDS
- 26a. *Organometallic compounds*
- See 1105.
- 26c. *Coordination compounds*
- 1758 Buchberger, W. and Haddad, P.R.: Separation of metallo-cyanide complexes by capillary zone electrophoresis. *J. Chromatogr. A*, 687 (1994) 343-349.
- 1759 Timerbaev, A.R. and Semenova, O.P.: Theoretical estimation of capillary zone electrophoresis behaviour of metal complexes using multivariate regression analysis. *J. Chromatogr. A*, 690 (1995) 141-148.
- See also 1047.
28. ANTIBIOTICS
- 1760 Armstrong, D.W., Gasper, M.P. and Rundlett, K.L.: Highly enantioselective capillary electrophoretic separations with dilute solutions of the macrocyclic antibiotic ristocetin A. *J. Chromatogr. A*, 689 (1995) 285-304.
- 1761 Tavares, M.F.M. and McGuffin, V.L.: Separation and characterization of tetracycline antibiotics by capillary electrophoresis. *J. Chromatogr. A*, 686 (1994) 129-142.
29. INSECTICIDES, PESTICIDES AND OTHER AGROCHEMICALS
- 29e. *Herbicides*
- See 1131.

## 30. SYNTHETIC AND NATURAL DYES

## 30a. Synthetic dyes

- 1762 Croft, S.N. and Hinks, D.: Analysis of dyes by capillary electrophoresis. *Book Pap. - Int. Conf. Exhib., AATCC*, (1992) 17-22; C.A., 121 (1994) 207647d.

## 30b. Chloroplast and other natural pigments

- 1763 Saitoh, K., Kato, H. and Teramae, N.: Separation of chlorophyll- $c_1$  and - $c_2$  by micellar electrokinetic capillary chromatography. *J. Chromatogr. A*, 687 (1994) 149-153.
- 1764 Singhal, R.M. and Kumar, P.: Electrophoretic fractionation of the soil humic acids of Chakrata Himalayas (U.P.) *Indian For.*, 120 (1994) 262-267; C.A., 121 (1994) 299950y.
- 1765 Stone, E.M., Vandenburgh, K., Nichols, B.E. and Sheffield, V.C.: Identification of rhodopsin gene mutations using GC-clamped denaturing gradient gel electrophoresis. *Methods Neurosci.*, 15(Photoreceptor Cells) (1993) 377-392; C.A., 121 (1994) 223085z.

## 31. PLASTICS AND THEIR INTERMEDIATES

- 1766 Bastos, D., Santos, R., Forcada, J., Hidalgo-Alvarez, R. and Nieves, F.J.: Surface and electrokinetic characterization of functional aldehyde polymer colloids. *Colloids Surf., A*, 92 (1994) 137-146; C.A., 122 (1995) 57305v.
- 1767 Baugartner, A. and Skolnick, J.: Polymer electrophoresis across a model membrane. *J. Phys. Chem.*, 98 (1994) 10655-10658; C.A., 121 (1994) 207178b.
- 1768 Paulke, B.-R., Moeglich, P.M., Knippel, E., Budde, A., Nitzsche, R. and Mueller, R.H.: Electrophoretic 3D-mobility profiles of latex particles with different surface groups. *Langmuir*, 11 (1995) 70-74; C.A., 122 (1995) 57061n.
- 1769 Radko, S.P., Garner, M.M., Caiafa, G. and Chrambach, A.: Molecular sieving of polystyrene carboxylate of a diameter up to 10  $\mu$ m in solutions of uncrosslinked polyacrylamide of  $M_r$  5  $\times$  10<sup>6</sup> using capillary zone electrophoresis. *Anal. Biochem.*, 223 (1994) 82-87.

See also 1045.

## 32. DRUG ANALYSIS

## 32a. Drug analysis, general techniques

- 1770 Aumatell, A., Wells, R.J. and Wong, D.K.Y.: Enantiomeric differentiation of a wide range of pharmacologically active substances by capillary electrophoresis using modified  $\beta$ -cyclodextrins. *J. Chromatogr. A*, 686 (1994) 293-307.
- 1771 Guttman, A. and Cooke, N.: Practical aspects in chiral separation of pharmaceuticals by capillary electrophoresis. II. Quantitative separation of naproxen enantiomers. *J. Chromatogr. A*, 685 (1994) 155-159.

- 1772 Lurie, I.S., Klein, R.F.X., Dal Cason, T.A., LeBelle, M.J., Brenneisen, R. and Weinberger, R.E.: Chiral resolution of cationic drugs of forensic interest by capillary electrophoresis with mixtures of neutral and anionic cyclodextrins. *Anal. Chem.*, 66 (1994) 4019-4026.
- 1773 Naylor, S., Benson, L.M. and Tomlinson, A.J.: Monitoring drug metabolism by capillary electrophoresis. In: Landers, J.P. (Editor), *Handb. Capillary Electrophor.*, CRC, Boca Raton, 1994, pp. 459-491; C.A., 121 (1994) 270981y - a review with 89 refs.
- 1774 Quaglia, M.G., Bossu, E., Desiderio, C. and Fanali, S.: (Use of capillary electrophoresis for testing the stability of a drug mixture in perfusion solution). *Farmaco*, 49 (1994) 403-406; C.A., 121 (1994) 308458y.
- 1775 Sun, P., Hoops, A. and Hartwick, R.A.: Enhanced albumin protein separations and protein-drug binding constant measurements using anti-inflammatory drugs as run buffer additives in affinity capillary electrophoresis. *J. Chromatogr. B*, 661 (1994) 335-340.
- 1776 Szoko, E., Kalasz, H. and Magyar, K.: (Application of capillary electrophoresis in drug analysis. Part I. Description of capillary electrophoresis technique). *Gyogyszereszet*, 38 (1994) 707-710; C.A., 122 (1995) 64494j - a review with 24 refs.
- 1777 Szoko, E., Kalasz, H. and Magyar, K.: (Application of capillary electrophoresis in drug analysis. Part II). *Gyogyszereszet*, 38 (1994) 711-713; C.A., 122 (1995) 64495k - a review with 61 refs.
- 1778 Tomlinson, A.J., Benson, L.M. and Naylor, S.: Effects of organic solvent in the CE and online CE-MS analysis of drug metabolite mixtures. *Am. Lab. (Shelton)*, 26 (1994) 29-36; C.A., 121 (1994) 291866e.

See also 1028, 1097, 1133, 1788.

## 32b. Antirheumatics and antiinflammatory drugs

See 1771, 1775.

## 32c. Autonomic and cardiovascular drugs

- 1779 Branch, S.K., Holzgrave, U., Jefferies, T.M., Mallwitz, H. and Matchett, M.W.: Chiral discrimination of phenethylamines with  $\beta$ -cyclodextrin and heptakis(2,3-di-O-acetyl)- $\beta$ -cyclodextrin by capillary electrophoresis and NMR spectroscopy. *J. Pharm. Biomed. Anal.*, 12 (1994) 1507-1517.
- 1780 Jumppanen, J.H. and Riekkola, M.-L.: Optimization of resolution of 13 diuretics in CZE by controlling capillary length and electroosmotic flow velocity. *J. Microcolumn Sep.*, 6 (1994) 595-604; C.A., 122 (1995) 64505p.

See also 1120.

## 32d. Central nervous system drugs

See 1138.

## 32e. Chemotherapeutics (exc. cytostatics and antibiotics)

- 1781 Yang, S. and Khaledi, M.G.: Micellar liquid chromatographic separation of sulfonamides in physiological samples using direct on-column injection. *J. Chromatogr. A*, 692 (1995) 311-318.

## 32f. Cytostatics

- 1782 Ng, C.L., Lee, H.K. and Li, S.F.Y.: Separation of tamoxifen and metabolites by capillary electrophoresis with non-aqueous buffer system. *J. Liq. Chromatogr.*, 17 (1994) 3847-3857.
- 1783 Quaglia, M.G., Farina, A., Kilar, F., Fanali, S., Bossu, E. and Dell'Aquila, C.: Analysis of a new doxorubicin derivative (FCE 23762) and related compounds by high performance capillary electrophoresis. *J. Liq. Chromatogr.*, 17 (1994) 3911-3923.

## 32g. Other drug categories

- 1784 Schmutz, A and Thormann, W.: Rapid determination of the antimycotic drug flucytosine in human serum by micellar electrokinetic capillary chromatography with direct sample injection. *Ther. Drug Monit.*, 16 (1994) 483-490; *C.A.*, 121 (1994) 291882g.

## 32h. Toxicological and forensic applications

- 1785 Anderson, N.L., Esquer-Blasco, R. and Anderson, N.G.: Two-dimensional electrophoresis as applied to problems of *in vitro* and *in vivo* toxicology. *Methods Toxicol.*, 1B(*In Vitro* Toxicity Indicators) (1994) 463-473; *C.A.*, 121 (1994) 249958m - a review with 3 refs.
- 1786 Oltra, J., Ortiz, M., de la Barra, V. and Stange, E.: (Blood typing of biochemical polymorphisms and application to paternity test in horses). *Arch. Med. Vet.*, 25 (1993) 147-153; *C.A.*, 121 (1994) 273986q.

## 32i. Plant extracts

- 1787 Pietta, P.G., Mauri, P.L., Bruno, A. and Merfort, I.: MEKC as an improved method to detect falsifications in the flowers of *Arnica montana* and *Arnica chamissonis*. *Planta Med.*, 60 (1994) 369-372; *C.A.*, 121 (1994) 238499m.

See also 1127.

## 33. CLINICO-CHEMICAL APPLICATIONS

## 33a. General papers and reviews

- 1788 Klein, G.L. and Jolliff, C.R.: Capillary electrophoresis for the routine clinical laboratory. In: Landers, J.P. (Editor), *Handb. Capillary Electrophor.*, CRC, Boca Raton, 1994, pp. 419-457; *C.A.*, 121 (1994) 249924x - a review with 33 refs.

See also 1714.

## 33b. Complex mixtures and profiling (single compounds by cross-reference only)

- 1789 Ridgeway, T.M., Hayes, D.B., Anderson, A.L., Levasseur, J.H., Demaine, P.D., Kenty, B.E. and Laue, T.M.: Possible clinical applications for direct molecular charge determination by equilibrium electrophoresis. *Proc. SPIE-Int. Soc. Opt. Eng.*, 213(Biochemical Diagnostic Instrumentation) (1994) 263-274; *C.A.*, 121 (1994) 275897k.

See also 1166, 1175, 1194, 1195, 1196, 1197, 1198, 1200, 1204, 1222, 1231, 1358, 1370, 1376, 1384, 1416, 1483, 1521, 1606, 1699, 1703, 1710, 1711, 1712, 1719, 1746, 1800.

## 34. FOOD ANALYSIS

## 34b. Complex mixtures (single compounds by cross-reference only)

See 1253, 1336, 1340, 1422, 1425.

## 35. ENVIRONMENTAL ANALYSIS

## 35b. Air pollution (complex mixtures; single compounds by cross-reference only)

- 1790 Pavlas, P.: (Determination of ionogenic pollutants in the air by capillary isotachopheresis). *Prac. Lek.*, 44 (1992) 63-65; *C.A.*, 121 (1994) 262183z.

See also 1038, 1804.

## 35c. Water pollution (complex mixtures; single compounds by cross-reference only)

See 1812.

## 36. SOME TECHNICAL PRODUCTS AND COMPLEX MIXTURES

## 36a. Surfactants

- 1791 Desbene, P.L. and Rony, C.M.: Determination by high-performance capillary electrophoresis of alkylaromatics used as bases of sulfonation in the preparation of industrial surfactants. *J. Chromatogr. A*, 689 (1995) 107-121.

See also 1071.

## 36b. Antioxidants and preservatives

- 1792 Maier, F.: Capillary electrophoresis analysis of preservation agents in cosmetics. *Seifen, Oele, Fette, Wachse*, 119 (1993) 991-997; *C.A.*, 122 (1995) 16821s.

## 36c. Complex mixtures, technical products and unidentified compounds

- 1793 Xu, Y., Ge, R. and Xu, S.: (Preparation of fetuin-gold probe). *Lanzhou Daxue Xuebao, Ziran Kexueban*, 29 (1993) 204-207; *C.A.*, 121 (1994) 225624y.

See also 1143, 1150, 1271.

## 37. CELLS, CELLULAR PARTICLES AND SUPRAMOLECULAR STRUCTURES

- 1794 Bondy, B., Bauer, J., Seuffert, I. and Weber, G.: Sodium chloride in separation medium enhances cell compatibility of free flow electrophoresis. *Electrophoresis (Weinheim)*, 16 (1995) 92-97.
- 1795 Dunstan, D.E.: Temperature dependence of the electrokinetic properties of two disparate surfaces. *J. Colloid. Interface Sci.*, 166 (1994) 472-475; *C.A.*, 121 (1994) 309069j.
- 1796 Kondo, T. and Ohshima, H.: Analysis of electrophoretic mobility data for biological cells with a new membrane model. *Prog. Colloid Polym. Sci.*, 93(Trends in Colloid and Interface Science VII) (1993) 280-283; *C.A.*, 121 (1994) 250278w.
- 1797 Ohshima, H.: Electrophoresis of soft particles. *Maku*, 19 (1994) 284-291; *C.A.*, 121 (1994) 308900m.
- 1798 Sloss, S. and Ewing, A.G.: Capillary electrophoresis for the analysis of single cells. In: Landers, J.P. (Editor), *Handb. Capillary Electrophor.*, CRC, Boca Raton, 1994, pp. 391-417; *C.A.*, 121 (1994) 249923w.
- 1799 Thomson-Carter, F.M., Carter, P.E. and Pennington, T.H.: Pulsed-field gel electrophoresis for the analysis of bacterial populations. *Soc. Appl. Bacteriol. Tech. Ser.*, 31(New Techniques in Food and Beverage Microbiology) (1993) 251-264; *C.A.*, 121 (1994) 249823p - a review with many refs.

See also 1045.

## 38. INORGANIC COMPOUNDS

## 38a. Cations

- 1800 Hiraoka, A., Miura, I., Akai, J.-i., Tominaga, I. and Hattori, M.: Capillary electrophoretic separation and determination of calcium and magnesium in cerebrospinal fluid and serum. *Manguneshumu (Kyoto)*, 13 (1994) 105-110; *C.A.*, 121 (1994) 225609x.
- 1801 Hirokawa, T., Xia, W. and Kiso, Y.: Isotachophoretic separation of rare earth ions. I. Separation behaviour of yttrium and fourteen lanthanide ions forming complexes with tartaric acid and  $\alpha$ -hydroxyisobutyric acid. *J. Chromatogr. A*, 689 (1995) 149-156.
- 1802 Morin, P., Francois, C. and Dreux, M.: Capillary electrophoresis of alkali and alkaline-earth cations with imidazole or benzylamine buffers. *J. Liq. Chromatogr.*, 17 (1994) 3869-3888.
- 1803 Motomezu, S., Oshima, M., Kavaahara, M. and Obata, Y.: Separation and sensitive determination of metal ions by capillary electrophoresis with 2-(5-bromo-2-pyridylazo)-5-(N-propyl-N-sulfopropylamino)phenol. *Analyst (Cambridge)*, 119 (1994) 1787-1792.

- 1804 Röder, A. and Bächmann, K.: Simultaneous determination of organic and inorganic anions in the sub- $\mu\text{mol/l}$  range in rain water by capillary zone electrophoresis. *J. Chromatogr. A*, 689 (1995) 305-311.
- 1805 Timerbaev, A.R., Semenova, O.P. and Bonn, G.K.: Capillary zone electrophoresis of lanthanoid elements after complexation with aminopolycarboxylic acids. *Analyst (Cambridge)*, 119 (1994) 2795-2799.
- 1806 Vogt, C. and Werner, G.: Speciation of heavy metals by capillary electrophoresis. *J. Chromatogr. A*, 686 (1994) 325-332.
- 1807 Xu, X., Kok, W.T., Kraak, J.C. and Poppe, H.: Simultaneous determination of urinary creatinine, calcium and other inorganic cations by capillary zone electrophoresis with indirect ultraviolet detection. *J. Chromatogr. B*, 661 (1994) 35-45.
- 1808 Yang, Y., Li, L. and Huang, H.: (Separation and determination of Y, Gd and Lu by high-voltage paper electrophoresis-fluorescence analysis method). *Yejin Fenxi*, 14 (1994) 48-50; *C.A.*, 122 (1995) 70882a.

See also 1063, 1067, 1093, 1141.

## 38b. Anions

- 1809 Harakuwe, A.H., Haddad, P.R. and Buchberger, W.: Optimisation of separation selectivity in capillary zone electrophoresis of inorganic anions using binary cationic surfactant mixtures. *J. Chromatogr. A*, 685 (1994) 161-165.
- 1810 Jimidar, M. and Massart, D.L.: Prediction of the migration behaviour of anions in capillary ion analysis. *Anal. Chim. Acta*, 294 (1994) 165-176.
- 1811 Kaniansky, D., Zelensky, I., Hybenova, A. and Onuska, F.I.: Determination of chloride, nitrate, sulfate, nitrite, fluoride, and phosphate by on-line coupled capillary isotachopheresis-capillary zone electrophoresis with conductivity detection. *Anal. Chem.*, 66 (1994) 4258-4264.
- 1812 Li, K. and Li, S.F.Y.: Determination of anions in water samples by capillary zone electrophoresis with indirect UV detection. *J. Liq. Chromatogr.*, 17 (1994) 3889-3910.
- 1813 Oefner, P.J.: Surface-charge reversed capillary zone electrophoresis of inorganic and organic anions. *Electrophoresis (Weinheim)*, 16 (1995) 46-56.
- 1814 Stahl, R.: Routine determination of anions by capillary electrophoresis and ion chromatography. *J. Chromatogr. A*, 686 (1994) 143-148.

See also 1071, 1109, 1148, 1150, 1804.

## 39. RADIOACTIVE AND OTHER ISOTOPE COMPOUNDS

See 1069.



## PUBLICATION SCHEDULE FOR THE 1995 SUBSCRIPTION

*Journal of Chromatography A* and *Journal of Chromatography B: Biomedical Applications*

| MONTH                                                | 1994          | J                                | F                                        | M                                | A                                        | M <sup>a</sup>                                   | J                                |                                                                      |
|------------------------------------------------------|---------------|----------------------------------|------------------------------------------|----------------------------------|------------------------------------------|--------------------------------------------------|----------------------------------|----------------------------------------------------------------------|
| Journal of Chromatography A                          | Vols. 683–688 | 689/1<br>689/2<br>690/1<br>690/2 | 691/1 + 2<br>692/1 + 2<br>693/1<br>693/2 | 694/1<br>694/2<br>695/1<br>695/2 | 696/1<br>696/2<br>697/1 + 2<br>698/1 + 2 | 699/1 + 2<br>700/1 + 2<br>702/1 + 2<br>703/1 + 2 | 704/1<br>704/2<br>705/1<br>705/2 | The publication schedule for further issues will be published later. |
| Bibliography Section                                 |               |                                  |                                          | 713/1                            |                                          |                                                  | 713/2                            |                                                                      |
| Journal of Chromatography B: Biomedical Applications |               | 663/1<br>663/2                   | 664/1<br>664/2                           | 665/1<br>665/2                   | 666/1<br>666/2                           | 667/1<br>667/2                                   | 668/1<br>668/2                   |                                                                      |

<sup>a</sup> Vol. 701 (Cumulative Indexes Vols. 652–700) expected in October.

### INFORMATION FOR AUTHORS

(Detailed *Instructions to Authors* were published in *J. Chromatogr. A*, Vol. 657, pp. 463–469. A free reprint can be obtained by application to the publisher, Elsevier Science B.V., P.O. Box 330, 1000 AH Amsterdam, Netherlands.)

**Types of Contributions.** The following types of papers are published: Regular research papers (full-length papers), Review articles, Short Communications and Discussions. Short Communications are usually descriptions of short investigations, or they can report minor technical improvements of previously published procedures; they reflect the same quality of research as full-length papers, but should preferably not exceed five printed pages. Discussions (one or two pages) should explain, amplify, correct or otherwise comment substantively upon an article recently published in the journal. For Review articles, see inside front cover under Submission of Papers.

**Submission.** Every paper must be accompanied by a letter from the senior author, stating that he/she is submitting the paper for publication in the *Journal of Chromatography A* or *B*.

**Manuscripts.** Manuscripts should be typed in **double spacing** on consecutively numbered pages of uniform size. The manuscript should be preceded by a sheet of manuscript paper carrying the title of the paper and the name and full postal address of the person to whom the proofs are to be sent. As a rule, papers should be divided into sections, headed by a caption (e.g., Abstract, Introduction, Experimental, Results, Discussion, etc.). All illustrations, photographs, tables, etc., should be on separate sheets.

**Abstract.** All articles should have an abstract of 50–100 words which clearly and briefly indicates what is new, different and significant. No references should be given.

**Introduction.** Every paper must have a concise introduction mentioning what has been done before on the topic described, and stating clearly what is new in the paper now submitted.

**Experimental conditions** should preferably be given on a *separate* sheet, headed "Conditions". These conditions will, if appropriate, be printed in a block, directly following the heading "Experimental".

**Illustrations.** The figures should be submitted in a form suitable for reproduction, drawn in Indian ink on drawing or tracing paper. Each illustration should have a caption, all the *captions* being typed (with double spacing) together on a *separate sheet*. If structures are given in the text, the original drawings should be provided. Coloured illustrations are reproduced at the author's expense, the cost being determined by the number of pages and by the number of colours needed. The written permission of the author and publisher must be obtained for the use of any figure already published. Its source must be indicated in the legend.

**References.** References should be numbered in the order in which they are cited in the text, and listed in numerical sequence on a separate sheet at the end of the article. Please check a recent issue for the layout of the reference list. Abbreviations for the titles of journals should follow the system used by *Chemical Abstracts*. Articles not yet published should be given as "in press" (journal should be specified), "submitted for publication" (journal should be specified), "in preparation" or "personal communication".

Vols. 1–651 of the *Journal of Chromatography*; *Journal of Chromatography, Biomedical Applications* and *Journal of Chromatography, Symposium Volumes* should be cited as *J. Chromatogr.* From Vol. 652 on, *Journal of Chromatography A* (incl. Symposium Volumes) should be cited as *J. Chromatogr. A* and *Journal of Chromatography B: Biomedical Applications* as *J. Chromatogr. B*.

**Dispatch.** Before sending the manuscript to the Editor please check that the envelope contains four copies of the paper complete with references, captions and figures. One of the sets of figures must be the originals suitable for direct reproduction. Please also ensure that permission to publish has been obtained from your institute.

**Proofs.** One set of proofs will be sent to the author to be carefully checked for printer's errors. Corrections must be restricted to instances in which the proof is at variance with the manuscript.

**Reprints.** Fifty reprints will be supplied free of charge. Additional reprints can be ordered by the authors. An order form containing price quotations will be sent to the authors together with the proofs of their article.

**Advertisements.** The Editors of the journal accept no responsibility for the contents of the advertisements. Advertisement rates are available on request. Advertising orders and enquiries can be sent to the Advertising Manager, Elsevier Science B.V., Advertising Department, P.O. Box 211, 1000 AE Amsterdam, Netherlands; Tel: 31 (20) 485 3796; Fax: 31 (20) 485 3810. Courier shipments to street address: Molenwerf 1, 1014 AG Amsterdam, Netherlands. UK: T.G. Scott & Son Ltd., Tim Blake, Portland House, 21 Narborough Road, Cosby, Leics. LE9 5TA, UK; Tel: (0116) 2750 521/2753 333; Fax: (0116) 2750 522. USA and Canada: Weston Media Associates, Daniel S. Lipner, P.O. Box 1110, Greens Farms, CT 06436-1110, USA; Tel: (203) 261 2500; Fax: (203) 261 0101.

# Carbohydrate Analysis

## High Performance Liquid Chromatography and Capillary Electrophoresis

Edited by Z. El Rassi

Journal of Chromatography Library, Volume 58

The objective of the present book is to provide a comprehensive review of carbohydrate analysis by HPLC and HPCE by covering analytical and preparative separation techniques for all classes of carbohydrates including mono- and disaccharides; linear and cyclic oligosaccharides; branched heterooligosaccharides (e.g., glycans, plant-derived oligosaccharides); glycoconjugates (e.g., glycolipids, glycoproteins); carbohydrates in food and beverage; compositional carbohydrates of polysaccharides; carbohydrates in biomass degradation; etc.

The book will be of interest to a wide audience, including analytical chemists and biochemists, carbohydrate, glycoprotein and glycolipid chemists, molecular biologists, biotechnologists, etc. It will also be a useful reference work for both the experienced analyst and the newcomer as well as for users of HPLC and HPCE, graduates and postdoctoral students.

### Contents: Part I. The Solute.

1. Preparation of carbohydrates for analysis by HPLC and HPCE (A.J. Mort, M.L. Pierce).

### Part II. Analytical and Preparative Separations.

2. Reversed-phase and hydrophobic interaction chromatography of carbohydrates and glycoconjugates (Z. El Rassi).  
3. High performance hydrophilic interaction chromatography of carbohydrates with

polar solvents (S.C. Churms).  
4. HPLC of carbohydrates with cation- and anion-exchange silica and resin-based stationary phases (C.G. Huber, G.K. Bonn).  
5. Analysis of glycoconjugates using high-pH anion-exchange chromatography (R.R. Townsend).  
6. Basic studies on carbohydrate - protein interaction by high performance affinity chromatography and high performance capillary affinity electrophoresis using lectins as protein models (S. Honda).  
7. Modern size exclusion chromatography of carbohydrates and glycoconjugates (S.C. Churms).  
8. High performance capillary electrophoresis of carbohydrates and glycoconjugates (Z. El Rassi, W. Nashabeh).  
9. Preparative HPLC of carbohydrates (K.B. Hicks).

### Part III. The Detection.

10. Pulsed electrochemical detection of carbohydrates at gold electrodes following liquid chromatographic separation (D.C. Johnson,

W.R. LaCourse).  
11. On-column refractive index detection of carbohydrates separated by HPLC and CE (A.E. Bruno, B. Krattiger).  
12. Mass spectrometry of carbohydrates and glycoconjugates (C.A. Settineri, A.L. Burlingame).  
13. Evaporative light scattering detection of carbohydrates in HPLC (M. Dreux, M. Lafosse).  
14. Chiroptical detectors for HPLC of carbohydrates (N. Purdie).  
15. Pre- and post-column detection-oriented derivatization techniques in HPLC of carbohydrates (S. Hase).  
16. Post-column enzyme reactors for the HPLC determination of carbohydrates (L.J. Nagels, P.C. Maes).  
17. Other direct and indirect detection methods of carbohydrates in HPLC and HPCE (Z. El Rassi, J.T. Smith).  
Subject index.

©1995 692 pages Hardbound  
Price: Dfl. 425.00 (US\$250.00)  
ISBN 0-444-89981-2

### ORDER INFORMATION

ELSEVIER SCIENCE B.V.  
P.O. Box 330  
1000 AH Amsterdam  
The Netherlands  
Fax: +31 (20) 485 2845

For USA and Canada:  
P.O. Box 945, New York  
NY 10159-0945  
Fax: +1 (212) 633 3680

*US\$ prices are valid only for the USA & Canada and are subject to exchange rate fluctuations; in all other countries the Dutch guilder price (Dfl.) is definitive. Customers in the European Union should add the appropriate VAT rate applicable in their country to the price(s). Books are sent postfree if prepaid.*



**ELSEVIER**

An imprint of Elsevier Science



0021-9673(19950616)713:2;1-2

- 5 N.A. 2538  
570.58