

Bibliography Section**Please note**

The pagination of Vol. 714 continues
from that of Vol. 713.

JOURNAL OF

CHROMATOGRAPHY A

INCLUDING ELECTROPHORESIS AND OTHER SEPARATION METHODS

EDITORS

U.A.Th. Brinkman (Amsterdam)
R.W. Giese (Boston, MA)
J.K. Haken (Kensington, N.S.W.)
C.F. Poole (London)
L.R. Snyder (Orinda, CA)
S. Terabe (Hyogo)

EDITORS, SYMPOSIUM VOLUMES,

E. Heftmann (Orinda, CA), Z. Deyl (Prague)

EDITORIAL BOARD

D.W. Armstrong (Rolla, MO)
W.A. Aue (Halifax)
P. Boček (Brno)
P.W. Carr (Minneapolis, MN)
J. Crommen (Liège)
V.A. Davankov (Moscow)
G.J. de Jong (Weesp)
Z. Deyl (Prague)
S. Dilli (Kensington, N.S.W.)
Z. El Rassi (Stillwater, OK)
H. Engelhardt (Saarbrücken)
M.B. Evans (Hatfield)
S. Fanali (Rome)
G.A. Guiochon (Knoxville, TN)
P.R. Haddad (Hobart, Tasmania)
I.M. Hais (Hradec Králové)
W.S. Hancock (Palo Alto, CA)
S. Hjertén (Uppsala)
S. Honda (Higashi-Osaka)
Cs. Horváth (New Haven, CT)
J.F.K. Huber (Vienna)
J. Janák (Brno)
P. Jandera (Pardubice)
B.L. Karger (Boston, MA)
J.J. Kirkland (Newport, DE)
E. sz. Kováts (Lausanne)
C.S. Lee (Ames, IA)
K. Macek (Prague)
A.J.P. Martin (Cambridge)
E.D. Morgan (Keele)
H. Poppe (Amsterdam)
P.G. Righetti (Milan)
P. Schoenmakers (Amsterdam)
R. Schwarzenbach (Dübendorf)
R.E. Shoup (West Lafayette, IN)
R.P. Singhal (Wichita, KS)
A.M. Siouffi (Marseille)
D.J. Strydom (Boston, MA)
T. Takagi (Osaka)
N. Tanaka (Kyoto)
K.K. Unger (Mainz)
P. van Zoonen (Bilthoven)
R. Verpoorte (Leiden)
Gy. Vigh (College Station, TX)
J.T. Watson (East Lansing, MI)
B.D. Westerlund (Uppsala)

EDITORS, BIBLIOGRAPHY SECTION

Z. Deyl (Prague), J. Janák (Brno), V. Schwarz (Prague)

ELSEVIER

JOURNAL OF CHROMATOGRAPHY A

INCLUDING ELECTROPHORESIS AND OTHER SEPARATION METHODS

Scope. The *Journal of Chromatography A* publishes papers on all aspects of **chromatography, electrophoresis** and related methods. Contributions consist mainly of research papers dealing with chromatographic theory, instrumental developments and their applications. In the *Symposium volumes*, which are under separate editorship, proceedings of symposia on chromatography, electrophoresis and related methods are published. *Journal of Chromatography B: Biomedical Applications*—This journal, which is under separate editorship, deals with the following aspects: developments in and applications of chromatographic and electrophoretic techniques related to clinical diagnosis or alterations during medical treatment; screening and profiling of body fluids or tissues related to the analysis of active substances and to metabolic disorders; drug level monitoring and pharmacokinetic studies; clinical toxicology; forensic medicine; veterinary medicine; occupational medicine; results from basic medical research with direct consequences in clinical practice.

Submission of Papers. The preferred medium of submission is on disk with accompanying manuscript (see *Electronic manuscripts* in the Instructions to Authors, which can be obtained from the publisher, Elsevier Science B.V., P.O. Box 330, 1000 AH Amsterdam, Netherlands). Manuscripts (in English; four copies are required) should be submitted to: Editorial Office of *Journal of Chromatography A*, P.O. Box 681, 1000 AR Amsterdam, Netherlands, Telefax (+31-20) 485 2304, or to: The Editor of *Journal of Chromatography B: Biomedical Applications*, P.O. Box 681, 1000 AR Amsterdam, Netherlands. Review articles are invited or proposed in writing to the Editors who welcome suggestions for subjects. An outline of the proposed review should first be forwarded to the Editors for preliminary discussion prior to preparation. Submission of an article is understood to imply that the article is original and unpublished and is not being considered for publication elsewhere. For copyright regulations, see below.

Publication information. *Journal of Chromatography A* (ISSN 0021-9673): for 1995 Vols. 683–714 are scheduled for publication. *Journal of Chromatography B: Biomedical Applications* (ISSN 0378-4347): for 1995 Vols. 663–674 are scheduled for publication. Subscription prices for *Journal of Chromatography A*, *Journal of Chromatography B: Biomedical Applications* or a combined subscription are available upon request from the publisher. Subscriptions are accepted on a prepaid basis only and are entered on a calendar year basis. Issues are sent by surface mail except to the following countries where air delivery via SAL is ensured: Argentina, Australia, Brazil, Canada, China, Hong Kong, India, Israel, Japan, Malaysia, Mexico, New Zealand, Pakistan, Singapore, South Africa, South Korea, Taiwan, Thailand, USA. For all other countries airmail rates are available upon request. Claims for missing issues must be made within six months of our publication (mailing) date. Please address all your requests regarding orders and subscription queries to: Elsevier Science B.V., Journal Department, P.O. Box 211, 1000 AE Amsterdam, Netherlands. Tel.: (+31-20) 485 3642; Fax: (+31-20) 485 3598. Customers in the USA and Canada wishing information on this and other Elsevier journals, please contact Journal Information Center, Elsevier Science Inc., 655 Avenue of the Americas, New York, NY 10010, USA, Tel. (+1-212) 633 3750, Telefax (+1-212) 633 3764.

Abstracts/Contents Lists published in Analytical Abstracts, Biochemical Abstracts, Biological Abstracts, Chemical Abstracts, Chemical Titles, Chromatography Abstracts, Current Awareness in Biological Sciences (CABS), Current Contents/Life Sciences, Current Contents/Physical, Chemical & Earth Sciences, Deep-Sea Research/Part B: Oceanographic Literature Review, Excerpta Medica, Index Medicus, Mass Spectrometry Bulletin, PASCAL-CNRS, Referativnyi Zhurnal, Research Alert and Science Citation Index.

US Mailing Notice. *Journal of Chromatography A* (ISSN 0021-9673) is published weekly (total 52 issues) by Elsevier Science B.V., (Sara Burgerhartstraat 25, P.O. Box 211, 1000 AE Amsterdam, Netherlands). Annual subscription price in the USA US\$ 5389.00 (US\$ price valid in North, Central and South America only) including air speed delivery. Second class postage paid at Jamaica, NY 11431. **USA POSTMASTERS:** Send address changes to *Journal of Chromatography A*, Publications Expediting, Inc., 200 Meacham Avenue, Elmont, NY 11003. Airfreight and mailing in the USA by Publications Expediting.

See inside back cover for Publication Schedule, Information for Authors and information on Advertisements.

© 1995 ELSEVIER SCIENCE B.V. All rights reserved.

0021-9673/95/\$09.50

No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, without the prior written permission of the publisher, Elsevier Science B.V., Copyright and Permissions Department, P.O. Box 521, 1000 AM Amsterdam, Netherlands.

Upon acceptance of an article by the journal, the author(s) will be asked to transfer copyright of the article to the publisher. The transfer will ensure the widest possible dissemination of information.

Special regulations for readers in the USA—This journal has been registered with the Copyright Clearance Center, Inc. Consent is given for copying of articles for personal or internal use, or for the personal use of specific clients. This consent is given on the condition that the copier pays through the Center the per-copy fee stated in the code on the first page of each article for copying beyond that permitted by Sections 107 or 108 of the US Copyright Law. The appropriate fee should be forwarded with a copy of the first page of the article to the Copyright Clearance Center, Inc., 222 Rosewood Drive, Danvers, MA 01923, USA. If no code appears in an article, the author has not given broad consent to copy and permission to copy must be obtained directly from the author. The fee indicated on the first page of an article in this issue will apply retroactively to all articles published in the journal, regardless of the year of publication. This consent does not extend to other kinds of copying, such as for general distribution, resale, advertising and promotion purposes, or for creating new collective works. Special written permission must be obtained from the publisher for such copying.

No responsibility is assumed by the Publisher for any injury and/or damage to persons or property as a matter of products liability, negligence or otherwise, or from any use or operation of any methods, products, instructions or ideas contained in the materials herein. Because of rapid advances in the medical sciences, the Publisher recommends that independent verification of diagnoses and drug dosages should be made.

Although all advertising material is expected to conform to ethical (medical) standards, inclusion in this publication does not constitute a guarantee or endorsement of the quality or value of such product or of the claims made of it by its manufacturer.

Ⓢ The paper used in this publication meets the requirements of ANSI/NISO Z39.48-1992 (Permanence of Paper).

Printed in the Netherlands

CONTENTS

Bibliography Section

Liquid Column Chromatography	B253
Gas Chromatography	B308
Planar Chromatography	B334
Electrophoresis	B347

ห้องสมุดวิทยาศาสตร์และเทคโนโลยี

19 ต.ค. 2538

JOURNAL OF CHROMATOGRAPHY A

VOL. 714 (1995)

JOURNAL OF CHROMATOGRAPHY A

INCLUDING ELECTROPHORESIS AND OTHER SEPARATION METHODS

EDITORS

U.A.Th. BRINKMAN (Amsterdam), R.W. GIESE (Boston, MA), J.K. HAKEN (Kensington, N.S.W.),
C.F. POOLE (London), L.R. SNYDER (Orinda, CA), S. TERABE (Hyogo)

EDITORS, SYMPOSIUM VOLUMES

E. HEFTMANN (Orinda, CA), Z. DEYL (Prague)

EDITORIAL BOARD

D.W. Armstrong (Rolla, MO), W.A. Aue (Halifax), P. Boček (Brno), P.W. Carr (Minneapolis, MN), J. Crommen (Liège), V.A. Davankov (Moscow), G.J. de Jong (Weesp), Z. Deyl (Prague), S. Dilli (Kensington, N.S.W.), Z. El Rassi (Stillwater, OK), H. Engelhardt (Saarbrücken), M.B. Evans (Hatfield), S. Fanali (Rome), G.A. Guiochon (Knoxville, TN), P.R. Haddad (Hobart, Tasmania), I.M. Hais (Hradec Králové), W.S. Hancock (Palo Alto, CA), S. Hjertén (Uppsala), S. Honda (Higashi-Osaka), Cs. Horváth (New Haven, CT), J.F.K. Huber (Vienna), J. Janák (Brno), P. Jandera (Pardubice), B.L. Karger (Boston, MA), J.J. Kirkland (Newport, DE), E. sz. Kováts (Lausanne), C.S. Lee (Ames, IA), K. Macek (Prague), A.J.P. Martin (Cambridge), E.D. Morgan (Keele), H. Poppe (Amsterdam), P.G. Righetti (Milan), P. Schoenmakers (Amsterdam), R. Schwarzenbach (Dübendorf), R.E. Shoup (West Lafayette, IN), R.P. Singhal (Wichita, KS), A.M. Siouffi (Marseille), D.J. Strydom (Boston, MA), T. Takagi (Osaka), N. Tanaka (Kyoto), K.K. Unger (Mainz), P. van Zoonen (Bilthoven), R. Verpoorte (Leiden), Gy. Vigh (College Station, TX), J.T. Watson (East Lansing, MI), B.D. Westerlund (Uppsala)

EDITORS, BIBLIOGRAPHY SECTION

Z. Deyl (Prague), J. Janák (Brno), V. Schwarz (Prague)



ELSEVIER

Amsterdam – Lausanne – New York – Oxford – Shannon – Tokyo

J. Chromatogr. A, Vol. 714 (1995)

The *Bibliography Section 1995* will be published in two volumes (713 and 714) of two issues each. Combined indexes to both volumes will appear in the last issue of the year, Vol. 714, No. 2.

© 1995 ELSEVIER SCIENCE B.V. All rights reserved.

0021-9673/95/\$09.50

No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, without the prior written permission of the publisher, Elsevier Science B.V., Copyright and Permissions Department, P.O. Box 521, 1000 AM Amsterdam, Netherlands.

Upon acceptance of an article by the journal, the author(s) will be asked to transfer copyright of the article to the publisher. The transfer will ensure the widest possible dissemination of information.

Special regulations for readers in the USA – This journal has been registered with the Copyright Clearance Center, Inc. Consent is given for copying of articles for personal or internal use, or for the personal use of specific clients. This consent is given on the condition that the copier pays through the Center the per-copy fee stated in the code on the first page of each article for copying beyond that permitted by Sections 107 or 108 of the US Copyright Law. The appropriate fee should be forwarded with a copy of the first page of the article to the Copyright Clearance Center, Inc., 222 Rosewood Drive, Danvers, MA 01923, USA. If no code appears in an article, the author has not given broad consent to copy and permission to copy must be obtained directly from the author. The fee indicated on the first page of an article in this issue will apply retroactively to all articles published in the journal, regardless of the year of publication. This consent does not extend to other kinds of copying, such as for general distribution, resale, advertising and promotion purposes, or for creating new collective works. Special written permission must be obtained from the publisher for such copying.

No responsibility is assumed by the Publisher for any injury and/or damage to persons or property as a matter of products liability, negligence or otherwise, or from any use or operation of any methods, products, instructions or ideas contained in the materials herein. Because of rapid advances in the medical sciences, the Publisher recommends that independent verification of diagnoses and drug dosages should be made.

Although all advertising material is expected to conform to ethical (medical) standards, inclusion in this publication does not constitute a guarantee or endorsement of the quality or value of such product or of the claims made of it by its manufacturer.

Ⓜ The paper used in this publication meets the requirements of ANSI/NISO Z39.48-1992 (Permanence of Paper).

Printed in the Netherlands

BIBLIOGRAPHY SECTION

SUPPLEMENT TO THE
JOURNAL OF CHROMATOGRAPHY A

1995

EDITORS:

Z. DEYL (Prague)

J. JANÁK (Brno)

V. SCHWARZ (Prague)

Bibliography Section

Liquid Column Chromatography

1. REVIEWS AND BOOKS

- 2546 Beelaram, A. and Sadana, A.: High-resolution fractionation processes: chromatographic techniques. *Isol. Purif.*, 2 (1994) 45-62; C.A., 122 (1995) 54016r - a review with 33 refs.
- 2547 Diez-Masa, J.C., Crego, A., de Frutos, M. and Polo, M.C.: High performance liquid chromatography with microcolumns. *Acquis. Recent. Chromatogr. Vin Cours Eur. Form. Contin.*, 1992, (1993) 27-32; C.A., 122 (1995) 75621z - a review with 18 refs.
- 2548 Hayashi, Y. and Matsuda, R.: Uncertainty structure information theory, and optimization of quantitative analysis in separation science. *Adv. Chromatogr. (N.Y.)*, 34 (1994) 347-427; C.A., 122 (1995) 150209b - a review with 198 refs.
- 2549 Janos, P.: Complex-forming equilibria in ion chromatography. *J. Chromatogr. A*, 699 (1995) 1-10 - a review with 84 refs.
- 2550 Jellum, E.: Chromatography and capillary electrophoresis in biomedicine. *Spec. Publ.-R. Soc. Chem.*, 154 (Reviews on Analytical Chemistry-Euroanalysis VIII) (1994) 308-316; C.A., 122 (1995) 50306n - a review with 14 refs.
- 2551 Kitazume, E.: (Countercurrent chromatography). *Bunseki*, (1994) 711-717; C.A., 122 (1995) 170873s - a review with 99 refs.
- 2552 Scott, R.P.W.: Liquid chromatography phase systems. *Chromatogr. Sci. Ser.*, 67 (1994) 51-92; C.A., 122 (1995) 143233b - a review with 15 refs.
- 2553 Scott, R.P.W.: Resolution retention and selectivity. *Chromatogr. Sci. Ser.*, 67 (1994) 15-49; C.A., 122 (1995) 143232a - a review with 13 refs.
- 2554 Sewell, P.A.: High-performance liquid chromatography. In: Hamilton, R.J. and Hamilton, S. (Editors), *Lipid Anal.*, IRL, Oxford, 1992, pp. 153-203; C.A., 122 (1995) 127737p - a review with 53 refs.
- 2555 Tachibana, K.: (Overview and basic theory of chromatography). *Bunseki*, (1995) 2-11; C.A., 122 (1995) 143283t - an elementary review with 5 refs.

See also 2556, 2578, 2602, 2630, 2638, 2655, 2656, 2662, 2676, 2695, 2707, 2709, 2717, 2734, 2737, 2742, 2744, 2754, 2755, 2783, 2819, 2914, 2934, 2936, 2953, 2992, 3009, 3022, 3080, 3081, 3123, 3129, 3423, 3491, 3518, 3519, 3520, 3533, 3549, 3623, 3631, 3634, 3675, 3764, 3806, 3809, 3820, 3822, 3832, 3849, 3882.

2. FUNDAMENTALS, THEORY AND GENERAL

2a. General

- 2556 Anonymous: (Ninety years of history of chromatography (1903-1993)). *Zh. Anal. Khim.*, 49 (1994) 1047-1058; C.A., 122 (1995) 80297c - a review with 106 refs.
- 2557 Bakken, G.A. and Kalivas, J.H.: Assessing chromatographic peak purity using condition index and singular value evolving profiles. *Anal. Chim. Acta*, 300 (1995) 173-181.
- 2558 Buerigisser, C.S., Scheidegger, A.M., Borkovec, M. and Sticher, H.: Chromatographic charge density determination of materials with low surface area. *Langmuir*, 10 (1994) 855-860; C.A., 120 (1994) 174398x.
- 2559 Dolan, J.W.: Case study - irreproducible retention times. *LC-GC*, 12 (1994) 900-904; C.A., 122 (1995) 100747e.
- 2560 Dolan, J.W.: Obtaining separations. Part IV: Putting it all together. *LC-GC*, 12 (1994) 666-668; C.A., 122 (1995) 100740x.
- 2561 Dolan, J.W.: Problems with water. *LC-GC Int.*, 8, No. 3 (1995) 134-136.
- 2562 Dolan, J.W.: Reader mailbag. *LC-GC Int.*, 8, No. 5 (1995) 250-252.
- 2563 Dolan, J.W.: Reader questions. *LC-GC Int.*, 8, No. 2 (1995) 76-82.
- 2564 Edlen, B.B.: Consistent chromatographic performance: fundamentals in media selection. *Am. Biotechnol. Lab.*, 12 (1994) 40, 42; C.A., 122 (1995) 50389s.
- 2565 Gagnon, P., Grund, E. and Lindbaeck, T.: Large-scale process development for hydrophobic interaction chromatography, part I - gel selection and development of binding conditions. *LC-GC*, 13 (1995) 318-328; C.A., 122 (1995) 234545w.
- 2566 Gennaro, M.C., Abrigo, C., Pobozy, E. and Marengo, E.: Retention dependence on organic modifier and interaction reagent concentration in reversed-phase ion-interaction HPLC. *J. Liq. Chromatogr.*, 18 (1995) 311-330.
- 2567 Glód, B.K., Alexander, P.W., Chen, Z.L. and Haddad, P.R.: Thermodynamic investigation of sample retention mechanism in ion-exclusion chromatography with inclusion compound in the mobile phase. *Anal. Chim. Acta*, 306 (1995) 267-272.
- 2568 Grung, B. and Kvalheim, O.M.: Retention time shift adjustments of two-way chromatograms using Bessel's inequality. *Anal. Chim. Acta*, 304 (1995) 57-66.

- 2569 Hanson, M., Unger, K.K., Denoyel, R. and Rouquerol, J.: Interactions of lysozyme with hydrophilic and hydrophobic polymethacrylate stationary phases in reversed phase chromatography (RPC). *J. Biochem. Biophys. Methods*, 29 (1994) 283-293; *C.A.*, 122 (1995) 208266m.
- 2570 Hawkes, S.J.: Distorted chromatographic peaks. *J. Chem. Educ.*, 71 (1994) 1032-1033; *C.A.*, 122 (1995) 55249f.
- 2571 Hill, H.M., Smith, I., Fairbrother, I., Tennant, K. and Robson, J.: Protein and other interference problems, and some solutions. *Methodol. Surv. Bioanal. Drugs*, 23(Biofluid and Tissue Analysis for Drugs, Including Hypolipidaemics) (1994) 327-334; *C.A.*, 122 (1995) 229994n.
- 2572 Kaiser, E. and Wojtusik, M.J.: Improvements in on-line ion chromatography. *Off. Proc.-Int. Water Conf.*, 54th (1993) 266-274; *C.A.*, 122 (1995) 150268v.
- 2573 Kohn, A.: Automating method validation and system suitability testing in HPLC and CE. *Am. Biotechnol. Lab.*, 12 (1994) 44-47; *C.A.*, 122 (1995) 26972f.
- 2574 Kozulic, B.: On the "donor-corridor" model of gel electrophoresis. II. Developments related to new gels capillary gel electrophoresis. *Appl. Theor. Electrophor.*, 4 (1994) 137-148; *C.A.*, 122 (1995) 234664j.
- 2575 Löre, A., Szász, G. and Budvári-Bárány, Z.: Further investigations in HPLC systems containing silica and polar bonded silica packing dynamically modified by camphorsulfonic acid. *J. Liq. Chromatogr.*, 18 (1995) 287-295.
- 2576 McDowall, R.D.: A suitable system for chromatography? *LC-GC Int.*, 8, No. 4 (1995) 196-200.
- 2577 McDowall, R.D.: Change? I've always worked this way. *LC-GC Int.*, 8, No. 2 (1995) 89-91.
- 2578 Mori, S.: (Size exclusion chromatography). *Bunseki*, (1995) 43-50; *C.A.*, 122 (1995) 143284u - a review with 81 refs.
- 2579 Ouchi, G.I.: Statistical techniques for chromatography data handling. *LC-GC*, 12 (1994) 826-830; *C.A.*, 122 (1995) 101065t.
- 2580 Poppe, H.: The role of miniaturization in the development of analytical fractionation methods. *Analisis*, 22 (1994) M22-M24; *C.A.*, 122 (1995) 177132h.
- 2581 Rowe, R.C., Mulley, V.J., Hughes, J.C., Nabney, I.T. and Debenham, R.M.: Neural networks for chromatographic peak classification - a preliminary study. *LC-GC*, 12 (1994) 690-698; *C.A.*, 122 (1995) 100741y.
- 2582 Rózylo, J.K., Janicka, M. and Siembida, R.: Advantages of TLC as a pilot technique for HPLC. *J. Liq. Chromatogr.*, 17 (1994) 3641-3653.
- 2583 Ryan, T.W.: HPLC method transfer to narrow bore columns: an evaluation. *J. Liq. Chromatogr.*, 18 (1995) 51-62.
- 2584 Seaver, C. and Przybytek, J.: Solvent selection, Part II - physical properties. *LC-GC Int.*, 8, No. 4 (1995) 190-195.
- 2585 Seaver, C. and Sadek, P.: Solvent selection. Part I. UV absorption characteristics. *LC-GC*, 12 (1994) 742; *C.A.*, 122 (1995) 177383r.
- 2586 Sharina, I.G., Cherkasov, A.V. and Nedospasov, A.A.: (Quantitative evaluations of the degree of similarity of chromatograms and electrophoretograms). *Zh. Fiz. Khim.*, 68 (1994) 1752-1757; *C.A.*, 122 (1995) 25434v.
- 2587 Shibukawa, M.: Theoretical interpretation of the retention of system peaks in partition chromatography with a mobile phase containing electrolytes. *J. Chromatogr. A*, 696 (1995) 165-172.
- 2588 Tanimura, M. and Tamura, M.: Chromatographic separation of three or more compositions with different adsorbability by cyclic pseudo-moving bed-like systems. *Jpn. Kokai Tokkyo Koho JP 06,170,112 [94,170,112]* (Cl. B01D15/00), 21 Jun. 1994, Appl. 92/329,516, 09 Dec. 1992; 7 p.; *C.A.*, 122 (1995) 137316q.
- 2589 Uhl, S., Rempfer, K., Egelhaaf, H.-J., Lehr, B. and Oelkrug, D.: Fluorescence characterization of acid-base interaction and mobility at chromatographic interfaces. *Anal. Chim. Acta*, 303 (1995) 17-23.
- 2590 Woodbury, C.P., Jr., Fitzloff, J.F. and Vincent, S.S.: Sample multiplexing for greater throughput in HPLC and related methods. *Anal. Chem.*, 67 (1995) 885-890.

For additional information see *C.A.*:

122 (1995) 27168k, 30628r, 177436k.

See also 2548, 2549, 2555, 2676, 2753, 3075, 3408.

2b. Thermodynamics and theoretical relationships

- 2591 Baumeister, E., Klose, U., Albert, K., Bayer, E. and Guiochon, G.: Determination of the apparent transverse and axial dispersion coefficients in a chromatographic column by pulsed field gradient nuclear magnetic resonance. *J. Chromatogr. A*, 694 (1995) 321-331.
- 2592 Blumberg, L.M.: Limits of resolution and speed of analysis in linear chromatography with and without focusing. *Chromatographia*, 39 (1994) 719-728.
- 2593 Borowko, M.: Isoelutotropic diagrams for ternary mobile phases systems in liquid-solid chromatography. *Chem. Anal. (Warsaw)*, 39 (1994) 319-324; *C.A.*, 122 (1995) 143536c.
- 2594 Bosch, E., Bou, P. and Rosés, M.: Linear description of solute retention in reversed-phase liquid chromatography by a new mobile phase polarity parameter. *Anal. Chim. Acta*, 299 (1994) 219-229.
- 2595 Cheong, W.J. and Choi, J.D.: Limitations of the linear solvation energy relationships in reversed phase liquid chromatography. *Bull. Korean Chem. Soc.*, 15 (1994) 868-873; *C.A.*, 122 (1995) 90393v.
- 2596 Foti, G. and sz. Kovats, E.: To the relationship of excess adsorption and retention in liquid/solid chromatography. In: Czempesz, F., Horvolygyi, Z. and Paszli, I. (Editors), *Proc.-Conf. Colloid Chem.*, 6th 1992, Hung. Chem. Soc., Budapest, 1993, pp. 96-99; *C.A.*, 122 (1995) 171021z.
- 2597 Furman, W.B., Layloff, T.P. and Tetzlaff, R.F.: Validation of computerized liquid chromatographic systems. *J. Assoc. Off. Anal. Chem.*, 77 (1994) 1314-1318.
- 2598 Glod, B.K.: Retention mechanism of ion exclusion chromatography. *Chem. Anal. (Warsaw)*, 39 (1994) 399-422; *C.A.*, 122 (1995) 171043h.
- 2599 Harlan, J.E., Picot, D., Loll, P.J. and Garavito, R.M.: Calibration of size-exclusion chromatography: use of a double Gaussian distribution function to describe pore sizes. *Anal. Biochem.*, 224 (1995) 557-563.
- 2600 Koo, Y.-M. and Hong, J.: Analysis of a nonlinear, coupled chromatographic system. *Korean J. Chem. Eng.*, 11 (1994) 227-231; *C.A.*, 122 (1995) 177075s.
- 2601 Meyer, V.R.: Quantitation of chromatographic peaks in the 0.1 to 1.0% range. *Chromatographia*, 40 (1995) 15-22.

- 2602 Miyabe, K.: Adsorption equilibrium and rate in reverse-phase liquid chromatography separation system. *Adsorpt. News*, 8 (1994) 16-20; *C.A.*, 122 (1995) 143205u - a review with 8 refs.
- 2603 Nicolas-Simonnot, M.-O., Sardin, M. and Grévillet, G.: Consequences of ion-exchange and sorption effects on pH waves propagating in a strong-base anion-exchange column. *J. Liq. Chromatogr.*, 18 (1995) 463-487.
- 2604 Parriott, P.J. and Lai, Y.-H.: Dynamic secondary equilibria in chromatography columns. *Chem. Aust.*, 61 (1994) 386-389; *C.A.*, 122 (1995) 177345e.
- 2605 Patrickios, C.S. and Patrickios, E.S.: Stoichiometric mass-action ion-exchange model. Explicit isotherms for mono-, di-, tri- and tetrameric ions. *J. Chromatogr. A*, 694 (1995) 480-485.
- 2606 Shatirishvili, Sh.I., Zakalashvili, G.N. and Sirbiladze, A.L.: (Evaluation of the minimum separation time of complex systems on high-performance chromatographic columns). *Khim. Prom-st (Moscow)*, (1994) 189-191; *C.A.*, 122 (1995) 203934t.
- 2607 Tan, L.C.: Study of retention mechanism in reversed-phase liquid chromatography. Avail. *Univ. Microfilms Int.*, Order No. DA9428946, 1994, 305 p.; *C.A.*, 122 (1995) 177131g.
- 2608 Thompson, R.A., Ge, Z., Grinberg, N., Ellison, D. and Tway, P.: Mechanistic aspects of the stereospecific interaction for aminoindanol with a crown ether column. *Anal. Chem.*, 67 (1995) 1580-1587.
- 2609 Zhu, J. and Guiochon, G.: Production rate of an isotachic train in displacement chromatography. *AIChE J.*, 41 (1995) 45-57; *C.A.*, 122 (1995) 84525y.
- See also 2568, 2581, 2648, 2657, 2688, 2695, 2739, 2761, 3495, 3845.
- 2c. *Relationship between structure and chromatographic behaviour*
- 2610 Gerasimenko, V.A. and Nabivach, V.M.: (Sorption-structural correlations in chromatography of aromatic compounds). *Zh. Fiz. Khim.*, 68 (1994) 1740-1748; *C.A.*, 122 (1995) 55540u.
- 2611 Hahai, T., Hatano, H., Nimura, N. and Kinoshita, T.: Molecular recognition in chromatography aided by computational chemistry. *Supramol. Chem.*, 3 (1994) 243-247; *C.A.*, 122 (1995) 171039m.
- 2612 Kumskov, M.I., Ponomareva, L.A., Smolenskii, E.A., Mitushev, D.F. and Zefirov, N.S.: (Automated method for formation of structure descriptors of organic compounds for quantitative structure-properties relations (QSPR)). *Izv. Akad. Nauk, Ser. Khim.*, (1994) 1391-1393; *C.A.*, 122 (1995) 104998k.
- 2613 Takács-Novák, K., Szász, G., Budvári-Bárány, Z., Józán, M. and Löre, A.: Relationship study between reversed phase HPLC retention and octanol/water partition among amphoteric compounds. *J. Liq. Chromatogr.*, 18 (1995) 807-825.
- 2614 Wilce, M.C.J., Aguilar, M.-I. and Hearn, M.T.W.: Physicochemical basis of amino acid hydrophobicity scales: evaluation of four new scales of amino acid hydrophobicity coefficients derived from RP-HPLC of peptides. *Anal. Chem.*, 67 (1995) 1210-1219.
- See also 3493.
- 2d. *Measurement of physico-chemical and related values*
- 2615 Bodzek, M. and Korus, I.: Characterization of ultrafiltration membranes using gel permeation chromatography. *Acta Chromatogr.*, 2 (1993) 40-55; *C.A.*, 122 (1995) 107741g.
- 2616 Miyabe, K. and Suzuki, M.: Chromatographic study on liquid-phase adsorption on octadecylsilyl-silica gel. *AIChE J.*, 41 (1995) 548-558; *C.A.*, 122 (1995) 191286t.
- 2617 Miyakoda, G., Nakayama, N., Matsui, K. and Yabuuchi, Y.: Mechanisms of the potent and long-lasting antihypertensive action of the new calcium channel blocker pranidipine. Evidence for strong affinity to the calcium channels or membranes. *Arzneim.-Forsch.*, 45 (1995) 471-475.
- See also 2614, 2650, 2681, 2735, 2762, 2874, 3057, 3156, 3446, 3447, 3607, 3610, 3611, 3612, 3620, 3842.
3. GENERAL TECHNIQUES
- 3a. *Apparatus and accessories*
- 2618 Goosens, E.C., de Jong, D., de Jong, G.J. and Brinkman, U.A.T.: Reversed-phase liquid chromatography coupled online with capillary gas chromatography II. Use of a solvent vapor exit to increase introduction volumes and introduction rates into the gas chromatograph. *J. Microcolumn Sep.*, 6 (1994) 207-215. *C.A.*, 122 (1995) 95569s.
- 2619 Karlsson, K. and Hellstroem, K.-G.: Method and apparatus for liquid chromatography. *PCT Int. Appl. WO 94 25,134 (Cl. B01D15/08)*, 10 Nov. 1994, SE Appl. 93/1,395, 26 Apr. 1993; 17 p.; *C.A.*, 122 (1995) 109480v.
- 2620 Rothchild, R.D.: Method and apparatus for moving-bed or simulated moving-bed continuous chromatographic separation. *U.S. US 5,387,347 (Cl. 210-659; B01D15/08)*, 07 Feb. 1995, Appl. 120,777, 15 Sep. 1993; 7 p.; *C.A.*, 122 (1995) 191538b.
- 2621 Stanin, V.V., Petrov, D.B., Isaev, V.B. and Ilyushin, A.A.: Thermostat for a chromatograph. *U.S.S.R. SU 1,807,385 (Cl. G01N30/30)*, 07 Apr. 1993, Appl. 4,933,992, 05 May 1991; *C.A.*, 122 (1995) 95640h.
- 2622 Wenk, M. and Haefeli, W.E.: A user-built automatic solid phase extraction device for parallel sample preparation prior to HPLC or RIA. *Eur. J. Clin. Chem. Clin. Biochem.*, 33 (1995) A18.
- For additional information see *C.A.*:
122 (1995) 155722y, 164202f, 164232r, 177425f.
- See also 3090.
- 3b. *Detectors and detection reagents*
- 2623 Grego, A.L., Diez-Masa, J.C. and Dabrio, M.V.: (Criteria for the modification of a UV-VIS HPLC detector to permit detection in open capillary columns). *Quim. Anal. (Barcelona)*, 12 (1993) 83-87; *C.A.*, 122 (1995) 150524a.
- 2624 Fallick, G. and Romano, J.: Enhanced photodiode array detection: HPLC applications in environmental research, development, and quality assurance using a spectral contrasting technique. *Int. Lab.*, 25, No. 3 (1995) 12C-12H.

- 2625 Glod, B.K., Alexander, P.W., Haddad, P.R. and Chen, Z.L.: Potentiometric detection using a metallic copper electrode in reversed-phase and ion-exclusion chromatography with eluents containing ion-interaction reagents. *J. Chromatogr. A*, 699 (1995) 31-37.
- 2626 Gorenstein, M.V., Li, J.B., Antwerp, J.V. and Chapman, D.: Detecting coeluted impurities by spectral comparison. *LC-GC*, 12 (1994) 768-772; *C.A.*, 122 (1995) 100744b.
- 2627 Jane, Y.S. and Shih, J.S.: Application of crown ether coated piezoelectric crystal as a detector for ion chromatography. *Analyst (Cambridge)*, 120 (1995) 517-522.
- 2628 Kim, J.-S. and Raines, R.T.: Dibromobimane as a fluorescent crosslinking reagent. *Anal. Biochem.*, 225 (1995) 174-176.
- 2629 Krull, I.S., Mhatre, R. and Cunniff, J.: Biopolymer detection, Part II - low-angle laser light scattering. *LC-GC Int.*, 8, No. 2 (1995) 83-88.
- 2630 Laubscher, J.C.: UV-VIS detection in HPLC. *Schweiz. Lab.-Z.*, 51 (1994) 239-243; *C.A.*, 122 (1995) 122195a - a review with 13 refs.
- 2631 Mank, A.J.G., Velthorst, N.H., Brinkman, U.A.T. and Gooijer, C.: Near-infrared laser-induced fluorescence detection in column liquid chromatography. A comparison of various lasers and detection systems. I. Continuous wave lasers. *J. Chromatogr. A*, 695 (1995) 165-174.
- 2632 Mank, A.J.G., Velthorst, N.H., Brinkman, U.A.T. and Gooijer, C.: Near-infrared laser-induced fluorescence detection in column liquid chromatography. A comparison of various lasers and detection systems. II. Pulsed lasers. *J. Chromatogr. A*, 695 (1995) 175-183.
- 2633 Roepke, W.: (Measurement in the femtomol range. The chemiluminescence detector in HPLC). *LaborPraxis*, 18 (1994) 52-55; *C.A.*, 122 (1995) 203943v.
- 2634 Smalley, M.B. and McGown, L.B.: Limits of detection and resolution for on-the-fly fluorescence lifetime detection in HPLC. *Anal. Chem.*, 67 (1995) 1371-1376.
- 2635 Takeuchi, T. and Miwa, T.: Effect of sodium dodecyl sulfate as stationary phase on signal intensities of dansylamino acids in microcolumn liquid chromatography with on-column fluorimetric detection. *J. Chromatogr. A*, 696 (1995) 185-192.
- 2636 Toyooka, T. and Liu, Y.-M.: Development of optically active fluorescent Edman-type reagents. *Analyst (Cambridge)*, 120 (1995) 385-390.
- 2637 Tran, C.D., Huang, G. and Grishko, V.I.: Direct and indirect detection of liquid chromatography by infrared thermal lens spectrometry. *Anal. Chim. Acta*, 299 (1995) 361-369.
- 2638 Wirth, M.J.: Spectroscopic probing of reversed-phase chromatographic retention. *LC-GC*, 12 (1994) 656-664; *C.A.*, 122 (1995) 100689n - a review with 39 refs.
- 2639 Wodecki, Z., Slebioda, M. and Kolodziejczyk, A.M.: Application of polarimetric detector for the high-performance liquid chromatographic determination of the optical purity of 5(4H)-oxazolones. *J. Chromatogr. A*, 696 (1995) 149-152.
- 2640 Yamamoto, A., Matsunaga, A., Mizukami, E., Hayakawa, K., Miyazaki, M., Nishimura, M., Kitaoka, M. and Fujita, T.: Evaluation of split-type flow cell for a polarized spectrophotometric detector. *Analyst (Cambridge)*, 120 (1995) 1137-1139.
- 2641 Yang, L., Janle, E., Huang, T., Gitzen, J., Kissinger, P.T., Vreeke, M. and Heller, A.: Applications of "wired" peroxidase electrodes for peroxide determination in liquid chromatography coupled to oxidase immobilized enzyme reactors. *Anal. Chem.*, 67 (1995) 1326-1331.
- 2642 Zegers, B.N., de Brouwer, J.F.C., Poppema, A., Lingeman, H. and Brinkman, U.A.T.: Electron-capture detection in reversed-phase liquid chromatography using packed-capillary columns. *Anal. Chim. Acta*, 304 (1995) 47-56.
- 2643 Zegers, B.N., Hessels, R., Jagesar, J., Rozenbrand, J., Lingeman, H. and Brinkman, U.A.T.: Photoionization detection in packed-capillary liquid and supercritical-fluid chromatography. *J. Liq. Chromatogr.*, 18 (1995) 413-440.

For additional information see C.A.:

122 (1995) 122158r, 177389x, 204235c.

See also 2768, 2820, 2921, 2933, 2966, 2984, 2993, 3007, 3017, 3027, 3034, 3035, 3050, 3052, 3473, 3509, 3541, 3613, 3717, 3737, 3745, 3773, 3872, 3895, 3896, 3899, 3902.

3c. Sorbents and columns, packing procedures

- 2644 Afeyan, N.B., Hodjat, H.K. and Varady, L.: Sulphonamide-bonded hydrophilic coatings for preparing chromatography matrix. *U.S. US 5,389,449 (Cl. 428-523; B01D15/08)*, 14 Feb. 1995, US Appl. 637,482, 04 Jan. 1991; 11 p.; *C.A.*, 122 (1995) 191540w.
- 2645 Alvarez, C., Bertorello, H. and Strumia, M.: Preparation of new synthetic hydrogels containing galactose, applicable in affinity chromatography. *J. Appl. Polym. Sci.*, 54 (1994) 1887-1891; *C.A.*, 122 (1995) 50602f.
- 2646 Anonymous: Chromatographic hydrations of sulphonated cation exchange resins. *Res. Discl.*, 361 (1994)36129; *C.A.*, 122 (1995) 135179y.
- 2647 Anspach, F.B. and Altmann-Haase, G.: Immobilized-metal-chelate regenerable carriers: (II) stability of copper chelate:conavalin A sandwich affinity-chromatographic sorbents. *Biotechnol. Appl. Biochem.*, 20 (1994) 323-333; *C.A.*, 122 (1995) 50411t.
- 2648 Arenas, R.V. and Foley, J.P.: Polymer coat4d alumina columns in reversed-phase liquid chromatography. *Am. Lab. (Shelton)*, 26 (1994) 32CC, 32DD; *C.A.*, 122 (1995) 150540c.
- 2649 Benmoussa, A., Mikou, M., Lacout, J.L. and Siouffi, A.M.: Lead phosphate hydroxyapatite high-performance liquid chromatography. *J. Chromatogr. A*, 694 (1995) 486-491.
- 2650 Biffis, A., Corain, B., Zecca, M., Corvaja, C. and Jerabek, K.: On the macromolecular structure and molecular accessibility of swollen microporous resins: a combined ESR-ISEC approach. *J. Am. Chem. Soc.*, 117 (1995) 1603-1606; *C.A.*, 122 (1995) 82759d.
- 2651 Bluemel, J.: Reaction of ethoxysilanes with silica: a solid state NMR study. *J. Am. Chem. Soc.*, 117 (1995) 2112-2113; *C.A.*, 122 (1995) 133271y.
- 2652 Buszewski, B., Gilpin, R.K. and Jaroniec, M.: Comparative studies of thermal properties of alkylamide and alkyl bonded phases. *Chem. Anal. (Warsaw)*, 39 (1994) 673-679; *C.A.*, 122 (1995) 80670u.

- 2653 Da, S.-L., Yue, W.-G., Wen, Y.-F., Da, H.-L. and Wang, Z.-H.: Preparation and characterization of bonded stationary phases of nitrogen-containing crown ether for high-performance liquid chromatography. *Anal. Chim. Acta*, 299 (1994) 239-247.
- 2654 Favrou, A. and Roussel, C.: Use of γ -cyclodextrin in chiral chromatography. In: Hedges, A.R. (Editor), *Minutes Int. Symp. Cyclodextrins, 6th*, Ed. Sante, Paris, 1992, pp. 603-606; C.A., 122 (1995) 143530w.
- 2655 Francisco Lorenzano Porras, C., Carr, P.W. and McCormick, A.V.: Pore characterization of SiO₂ chromatographic gels using nuclear magnetic resonance. In: Adair, J.H., Casey, J.A. and Venigalla, S. (Editors), *Handb. Charact. Tech. Solid-Solution Interface*, Am. Ceram. Soc., Westerville, 1993, pp. 273-281; C.A., 122 (1995) 197686v - a review with 22 refs.
- 2656 Giddings, J., Ratanathanawongs, S.K., Barman, B.N., Moon, M.H., Liu, G., Tjelta, B.L. and Hansen, M.E.: Characterization of colloidal and particulate silica by field-flow fractionation. *Adv. Chem. Ser.*, 234 (1994) 309-340; C.A., 122 (1995) 197692u - a review with 23 refs.
- 2657 Göhlin, K., Buschke, A. and Larsson, M.: Kinetic performance of open-tubular and packed columns in LC using the same stationary phase: immobilized polymethyloctadecylsiloxane. *Chromatographia*, 39 (1994) 729-739.
- 2658 He, B., Yan, H., Yu, K., Cheng, X. and Ni, A.: Synthesis of chiral ligand exchange resin. *Chin. Sci. Bull.*, 38 (1993) 1451-1453; C.A., 122 (1995) 134561e.
- 2659 Honda, F., Honda, H. and Koishi, M.: Utilization of the dry impact blending method to prepare irregularly shaped particles for high-performance liquid chromatographic column packings. *J. Chromatogr. A*, 696 (1995) 19-30.
- 2660 Kaminski, M.: Reasons for non-uniformity of bed structure in wet packed preparative liquid chromatographic columns. *Isol. Purif.*, 2 (1994) 1-17; C.A., 122 (1995) 30712p.
- 2661 Kitagawa, S. and Tsuda, T.: New packing method for liquid chromatographic columns by the aid of electrostatic field (II): application for packing of silica gels in short period. *Kuromatogurafi*, 15 (1994) 137-140; C.A., 122 (1995) 171044j.
- 2662 Levison, P.R.: Cellulosics as ion-exchange materials. In: Kennedy, J.F., Phillips, G.O. and Williams, P.A. (Editors), *Cellulosics*, Horwood, Chichester, 1993, pp. 25-36; C.A., 122 (1995) 131038d - a review with 25 refs.
- 2663 Lodkowski, R., Kruszynska, E., Suprynowicz, Z. and Buszewski, B.: Narrow bore HPLC column with chemically bonded phase on porous glass. *Chem. Anal. (Warsaw)*, 39 (1994) 543-554; C.A., 122 (1995) 171061n.
- 2664 Majors, R.E.: New chromatography columns and accessories at the 1995 Pittsburgh Conference, Part I. *LC-GC Int.*, 8, No. 5 (1995) 266-283.
- 2665 Maslowska, J. and Gasinska, E.: (Chelexes as a polymeric column packing in the chromatographic separation of amino acids). *Zesz. Nauk.-Politech. Lodz., Technol. Chem. Spozyw.*, 589 (1993) 41-55; C.A., 122 (1995) 150502s.
- 2666 Miyabe, K. and Suzuki, M.: Adsorption characteristics of reversed-phase liquid chromatography with various alkyl bonded phases. *J. Chem. Eng. Jpn.*, 27 (1994) 785-789; C.A., 122 (1995) 115973z.
- 2667 Myers, T.A., Healy, M.G., Murphy, W.R. and Al-Duri, B.: The assessment of chitosan beads as support materials for use in dye-ligand affinity chromatography. In: *ICHEM Res. Event, Two-Day Symp.*, 1, Inst. Chem. Eng., Rugby, 1994, pp. 210-213; C.A., 122 (1995) 26974h.
- 2668 Nagae, N.: (Packed column for reversed-phase liquid chromatography: "Develosil ODS-UG"). *Kuromatogurafi*, 15 (1994) 175-179; C.A., 122 (1995) 171045k.
- 2669 Noel, R., Sanderson, A. and Spark, L.: A monolithic ion-exchange material suitable for downstream processing of bio-products. In: Kennedy, J.F., Phillips, G.O. and Williams, P.A. (Editors), *Cellulosics*, Horwood, Chichester, 1993, pp. 17-24; C.A., 122 (1995) 131047f.
- 2670 Ogino, K., Sato, H., Aihara, Y., Suzuki, H. and Moriguchi, S.: Preparation and characterization of monodisperse oligo(ethylene glycol) dimethacrylate polymer beads for aqueous high-performance liquid chromatography. *J. Chromatogr. A*, 699 (1995) 67-72.
- 2671 Ogino, K., Sato, H., Tsuchiya, K., Suzuki, H. and Moriguchi, S.: Synthesis of monodisperse macroreticular styrene-divinylbenzene gel particles by a single-step swelling and polymerization method. *J. Chromatogr. A*, 699 (1995) 59-66.
- 2672 Pesek, J.J. and Tang, V.H.: The modification of alumina, zirconia, thoria and titania for potential use as HPLC stationary phases through a silanization/hydrosilation reaction scheme. *Chromatographia*, 39 (1994) 649-654.
- 2673 Pesek, J.J., Matyska, M.T., Soczewinski, E. and Christensen, P.: Spectroscopic studies of butylphenyl, mono-ol and perfluorinated bonded phases. *Chromatographia*, 39 (1994) 520-528.
- 2674 Schmid, J., Albert, K. and Bayer, E.: Synthesis and characterization of unsaturated bonded phases for high-performance liquid chromatography. *J. Chromatogr. A*, 694 (1995) 333-345.
- 2675 Schurig, F.V., Schleimer, M. and Mayer, S.: Preparation and of chiral, cyclodextrin-containing polymers immobilized on supports. *Ger. Offen DE 4,324,636 (Cl. C08B37/16)*, 11 May 1994, Appl. 22 Jul. 1993; 5 p.; C.A., 122 (1995) 84129x.
- 2676 Scott, R.P.W.: The liquid chromatography column. *Chromatogr. Sci. Ser.*, 67 (1994) 93-122; C.A., 122 (1995) 39513g - a review with 14 refs.
- 2677 Shalliker, R.A., Kavanagh, P.E. and Russell, I.M.: Reversed phase behaviour of high molecular mass polystyrenes in ethyl acetate solvent mixture. *Chromatographia*, 39 (1994) 663-667.
- 2678 Terabe, S., Ozaki, H., Tanaka, Y. and Okada, Y.: (Pseudo-stationary phases for electrokinetic chromatography). *Kuromatogurafi*, 15 (1994) 114-115; C.A., 122 (1995) 155375n.
- 2679 Tong, D., Bartle, K.D. and Clifford, A.A.: Preparation and evaluation of supercritical carbon dioxide-packed capillary columns for HPLC and SFC. *J. Microcolumn Sep.*, 6 (1994) 249-255; C.A., 122 (1995) 95583s.
- 2680 Wang, Q.: Desing preparation and characterization of novel polymeric separation media for high-performance liquid chromatography. Avail. *Univ. Microfilms Int.*, Order No. DA9501304, 1994, 172 p.; C.A., 122 (1995) 162792z.
- 2681 Yamamoto, F. and Rokushika, S.: (Characteristics of stationary phase on fluorocarbon bonded silica column). *Kuromatogurafi*, 15 (1994) 100-101; C.A., 122 (1995) 171103c.
- 2682 Zusman, R. and Zusman, I.: Gel fiberglass as a new support for affinity chromatography. *Biotechnol. Appl. Biochem.*, 21 (1995) 161-172; C.A., 122 (1995) 234537v.

For additional information see C.A.:

122 (1995) 50735b, 76031u, 82648s, 83166v, 90358n, 101130k, 122270w, 134188p, 161725t, 177435j, 177437m.

See also 2591, 2615, 2635, 2699, 2708, 2711, 2712, 2726, 2728, 2730, 2731, 2733, 2738, 2739, 2740, 2742, 2743, 2749, 2750, 2751, 2771, 2830, 2981, 3033, 3112, 3116, 3120, 3124, 3126, 3144, 3158, 3193, 3215, 3234, 3322, 3614, 3715, 3776, 3873.

3d. Quantitative analysis

2683 Meyer, V.R.: Errors in the area determination of incompletely resolved chromatographic peaks. *J. Chromatogr. Sci.*, 33 (1995) 26-33.

For additional information see C.A.:

122 (1995) 177349j.

See also 2548, 2779, 2781, 2805, 2809, 2980, 3624, 3667, 3752.

3e. Preparative scale chromatography

2684 Tharakan, J.P. and Belizaire, M.: Protein band dispersion in axial and radial flow chromatography. *J. Liq. Chromatogr.*, 18 (1995) 39-49.

See also 2620, 2746, 2842, 2928, 2981, 2983, 3026, 3173, 3193, 3481, 3499, 3563.

3f. Programmed temperature, pressure, vapors, gradients

2685 Treiber, L.R.: Normal-phase high-performance liquid chromatography with relay gradient elution. I. Description of the method. *J. Chromatogr. A*, 696 (1995) 193-199.

See also 2698, 3606.

4. SPECIAL TECHNIQUES

4a. Automation

2686 Olivè, J., Månuel, A., Rodriguez, P. and Grimalt, J.O.: A low cost, open and flexible on-line data acquisition system for instrumental chromatography. *Anal. Chim. Acta*, 306 (1995) 285-295.

2687 Pullen, F.S. and Richards, D.S.: Automated liquid chromatography/mass spectrometry for chromatographers. *Rapid Commun. Mass Spectrom.*, 9 (1995) 188-190; C.A., 122 (1995) 142700q.

For additional information see C.A.:

122 (1995) 204188q, 234488e.

See also 2573, 3017, 3078, 3463, 3579, 3587, 3658, 3821.

4b. Computerization and modelling

2688 Carta, G.: Linear driving force approximation for intraparticle diffusion and convection in permeable supports. *Chem. Eng. Sci.*, 50 (1995) 887-889; C.A., 122 (1995) 191295v.

2689 De Aguiar, P.F., Burquignon, B., Khots, M.S., Penninckx, W. and Massart, D.L.: The use of logistic transformation in HPLC optimization. *Quim. Anal. (Barcelona)*, 12 (1993) 177-183; C.A., 122 (1995) 150546j.

2690 Galushko, S.V., Kamenchuk, A.A. and Pit, G.L.: Software for method development in reversed-phase liquid chromatography. *Int. Lab.*, 25, No. 2 (1995) 14-17.

2691 Hanai, T., Hatano, H., Nimura, N. and Kinoshita, T.: Computational chemical analysis of the chiral recognition of binuclear copper(II) of N-salicylidene (R)-2-amino-1,2-bis(2-butoxy-5-tert-butylphenyl)-3-phenyl-1-propanol in liquid chromatography. *J. Liq. Chromatogr.*, 17 (1994) 4327-4334.

2692 Kevra, S.A., Bergman, D.L. and Maloy, J.T.: A computational introduction to chromatographic bandshape analysis. *J. Chem. Educ.*, 71 (1994) 1023-1028; C.A., 122 (1995) 55248e.

2693 Levy, D. and Miller, C.: Implementing client/server computing in the chromatography laboratory. *Int. Lab.*, 25, No. 2 (1995) 29-31.

2694 Mew, C.D.: A three dimensional surface viewer for Microsoft Windows. *Lab. Microcomput.*, 13 (1994) 17-24; C.A., 122 (1995) 204182h.

2695 Nabivach, V.M.: (Topological models and chromatographic retention of aromatic and heterocyclic compounds. Review. 1. Characteristics of design calculations and comparative characteristics of topological indexes). *Koks Khim.*, (1994) 16-22; C.A., 122 (1995) 150509z - a review with 32 refs.

2696 Naef, O.: (Analysis of a chromatogram). *Chimia*, 48 (1994) 409-412; C.A., 122 (1995) 204169j.

2697 Popov, A.A., Sergeev, S.K. and Prokhorova, E.K.: (Computerized chromatographic system for air quality monitoring). *Ekol. i Bezopas. Truda v Prom.-sti: Mater. Semin. Tsent. Ros. Dom Znani, M.*, (1991) 43-47; C.A., 120 (1994) 306103j.

2698 Soczewinski, E. and Markowski, W.: Application of stepwise gradient elution chromatography to preliminary separation in short columns. *Chem. Anal. (Warsaw)*, 39 (1994) 467-474; C.A., 122 (1995) 177127k.

2699 Van Straten, M.A., Lynch, R.J. and Claessens, H.A.: The role of stationary phases in reversed-phase liquid chromatography in the application of solvent optimization procedures. *LC-GC Int.*, 8, No. 3 (1995) 143-150.

2700 Villioth, O.: (Processing of chromatographic data under Windows NT). *LaborPraxis*, 18 (1994) 22-27; C.A., 122 (1995) 204171d.

See also 2597, 2598, 2611, 2686, 2718, 2752, 2754, 2761, 3062, 3122.

4c. Combination with other physico-chemical techniques (MS, IR etc.)

2701 Debets, A.J.J., Mekes, T.J.L., Ritburg, A. and Jacobs, P.L.: Column liquid chromatography coupled on-line with mass spectrometry using post-column phosphate suppression. *J. High Resolut. Chromatogr.*, 18 (1995) 45-48.

- 2702 Eaton, A.N. and Rao, R.: Mass detectors: analyte-specific detection for HPLC. *Int. Lab.*, 25, No. 2 (1995) 18-21.
- 2703 Gagne, J.P.: Systematic approach to the development and optimization of mass spectrometry-liquid chromatography systems for the analysis of organic substances. Avail. *NLC Order No. DANN88921*, 1993, 268 p.; *C.A.*, 122 (1995) 177404y.
- 2704 Joos, P.E.: Electrospray and APCI-mass spectrometry. *LC-GC Int.*, 8, No. 2 (1995) 92-95.
- 2705 Kaltashov, I.A., Yu, X. and Fenselau, C.: Simple interface for microbore LC and electrospray ionization mass spectrometry and analysis of melphalan-alkylation sites in metallothionein. *J. Pharm. Biomed. Anal.*, 13 (1995) 279-284.
- 2706 Manne, R.: On the resolution problem in hyphenated chromatography. *Chemom. Intell. Lab. Syst.*, 27 (1995) 89-94; *C.A.*, 122 (1995) 150308h.
- 2707 Murakoshi, R.: (Recent development of liquid chromatography (LC)-mass spectrometry (MS)). *Idemitsu Giho*, 37 (1994) 732-736; *C.A.*, 122 (1995) 150511u - a review with 9 refs.
- See also 2687, 2776, 2876, 2925, 2936, 2946, 2952, 2960, 2962, 2968, 2976, 2977, 2985, 3002, 3003, 3092, 3108, 3162, 3243, 3341, 3372, 3461, 3467, 3487, 3488, 3489, 3490, 3492, 3495, 3500, 3538, 3570, 3576, 3581, 3582, 3583, 3587, 3603, 3615, 3641, 3646, 3647, 3701, 3704, 3714, 3723, 3746, 3758, 3779, 3786, 3792, 3803, 3824, 3827, 3837, 3840, 3850, 3852, 3855, 3858, 3859, 3864, 3871, 3883, 3884.
- 4d. Affinity chromatography (advances)**
- 2708 Brena, B.M., Ryden, L.G. and Porath, J.: Immobilization of β -galactosidase on metal-chelate-substituted gels. *Biotechnol. Appl. Biochem.*, 19 (1994) 217-231; *C.A.*, 122 (1995) 208657q.
- 2709 De Frutos, M.: Chromatography-immunology coupling, a powerful tool for environmental analysis. *TrAC*, 14 (1995) 133-140 - a review with 49 refs.
- 2710 Galaev, I.Yu. and Mattiasson, B.: Shielding affinity chromatography. *BioTechnology*, 12 (1994) 1086; *C.A.*, 122 (1995) 26983k.
- 2711 Glatz, Z., Psotova, J., Janiczek, O. and Novotny, M.V.: Immobilized lectins as a means of reversible immobilization of ligands for affinity chromatography. *Int. J. Bio-Chromatogr.*, 1 (1994) 143-149; *C.A.*, 122 (1995) 75706f.
- 2712 Guo, W., Shang, Z.H., Yu, Y.N. and Zhou, L.M.: Cellulose membrane used as stationary phase of membrane affinity chromatography. *Chin. Chem. Lett.*, 5 (1994) 869-872; *C.A.*, 122 (1995) 75691x.
- 2713 Heywood, S.P. and Hooper, N.M.: Purification of GPI anchors by immunoaffinity chromatography. *Biochem. Soc. Trans.*, 23 (1995) 101S; *C.A.*, 122 (1995) 155591e.
- 2714 Jennissen, H.P. and Botzet, G.: The binding of phosphorylase kinase to immobilized calmodulin. *J. Mol. Recognit.*, 6 (1993) 117-130; *C.A.*, 122 (1995) 26386t.
- 2715 Kasai, K.-L., Sakurai, R. and Ishii, S.-I.: Frontal affinity chromatography using microcolumns as investigative tools for molecular recognition. *Int. J. Bio-Chromatogr.*, 1 (1994) 83-91; *C.A.*, 122 (1995) 50420v.
- 2716 Pang, I.-H., Smrcka, A.V. and Sternweis, P.C.: Synthesis and applications of affinity matrix containing immobilized β subunits of G proteins. *Methods Enzymol.*, 237(Heterotrimeric G Proteins) (1994) 164-174; *C.A.*, 122 (1995) 26970d.
- 2717 Pepper, D.S.: Some alternative coupling chemistries for affinity chromatography. *Mol. Biotechnol.*, 2 (1994) 157-178; *C.A.*, 122 (1995) 26909r - a review with 11 refs.
- 2718 Sridhar, P. and Sastri, N.V.S.: Modeling of bioseparations in an affinity percolation column packed with porous beads. *Indian Chem. Eng., Sect. A*, 36 (1994) 72-78; *C.A.*, 122 (1995) 50630p.
- 2719 Yoshihara, Y. and Kuroda, M.: Use of an SDS-gel-separated protein band as a ligand for affinity chromatography: procedure and application to the purification of domain-specific antibodies against α -actinin. *J. Biochem. (Tokyo)*, 117 (1995) 443-446.
- For additional information see *C.A.*:
122 (1995) 27255m, 155421z, 212170t.
- See also 2645, 2647, 2667, 2682, 2848, 2959, 3029, 3061, 3153, 3154, 3158, 3164, 3169, 3175, 3234, 3257, 3334, 3349, 3362, 3397, 3846, 3847, 3848, 3851.
- 4e. Functional analysis**
- See 3475, 3609.
- 4f. Trace analysis and preseparation techniques**
- 2720 Buttler, T., Jarskog, H., Gorton, L., Marko-Varga, G. and Ramnemark, L.: The use of microdialysis for sampling in column liquid chromatography. *Curr. Top. Membr.*, 41(Cell Biology and Membrane Transport Processes) (1994) 45-69; *C.A.*, 122 (1995) 127783a.
- 2721 Gy, P.M.: Sampling - diving into the unknown. *LC-GC Int.*, 8, No. 2 (1995) 70-74.
- 2722 Majors, R.E.: New approaches to sample preparation. *LC-GC Int.*, 8, No. 3 (1995) 128-133.
- 2723 Nguyen, V.S. and Nguyen, M.S.: Neutron activation analysis for determination of trace impurities of light rare earth elements in high purity Y_2O_3 . *Tap. Chi Hoa Hoc*, 32 (1994) 72-73; *C.A.*, 122 (1995) 122009t.
- See also 2792, 2797, 3451, 3576, 3632, 3823, 3825, 3826, 3873, 3882, 3884.
- 4g. Enantiomers, separation**
- 2724 Bielejewska, A., Kozbial, M., Nowakowski, R., Duszczyk, K. and Sybilska, D.: Studies on the behaviour of α -, β - and γ -cyclodextrins and some derivatives under reversed-phase liquid chromatographic. *Anal. Chim. Acta*, 300 (1995) 201-206.
- 2725 Bunton, B.M. and Walker, T.A.: Chiral separation of MDL 73,005EF enantiomers using an α_1 -acid glycoprotein column. *J. Chromatogr. A*, 699 (1995) 389-394.
- 2726 Chankvetadze, B., Yashima, E. and Okamoto, Y.: Dimethyl-, dichloro- and chloromethylphenylcarbamates of amylose as chiral stationary phases for high-performance liquid chromatography. *J. Chromatogr. A*, 694 (1995) 101-109.

- 2727 Cleveland, T.: Pirkle-concept chiral stationary phases for the HPLC separation of pharmaceutical racemates. *J. Liq. Chromatogr.*, 18 (1995) 649-671.
- 2728 Cortes, H.J. and Nicholson, L.W.: Chiral separations using packed capillary liquid chromatography. *J. Microcolumn Sep.*, 6 (1994) 257-263; C.A., 122 (1995) 122211c.
- 2729 Gasparrini, F., Lunazzi, L., Misiti, D. and Villani, C.: Organic stereochemistry and conformational analysis from enantioselective chromatography and dynamic nuclear magnetic resonance measurements. *Acc. Chem. Res.*, 28 (1995) 163-170; C.A., 122 (1995) 213225b.
- 2730 Haginaka, J. and Kanasugi, N.: Enantioselectivity of bovine serum albumin-bonded columns produced with isolated protein fragments. *J. Chromatogr. A*, 694 (1995) 71-80.
- 2731 Hyun, M.H., Cho, Y.J., Ryoo, J.-J., Jung, K.K. and Heo, G.S.: Preparation and application of an (S)-naproxen chiral stationary phase. *J. Chromatogr. A*, 696 (1995) 173-183.
- 2732 Hyun, M.H., Min, C.S. and Cho, Y.J.: Examples of liquid chromatographic resolution of π -acidic racemates on a π -acidic chiral stationary phase. *J. High Resolut. Chromatogr.*, 18 (1995) 63-65.
- 2733 Ihara, T., Sugimoto, Y., Asada, M., Nakagama, T. and Hobo, T.: Influence of the method of preparation of chiral stationary phases on enantiomer separations in high-performance liquid chromatography. *J. Chromatogr. A*, 694 (1995) 49-56.
- 2734 Kempe, M. and Mosbach, K.: Molecular imprinting used for chiral separations. *J. Chromatogr. A*, 694 (1995) 3-13 - a review with 88 refs.
- 2735 Levin, S. and Abu-Lafi, S.: Experimental studies of competition between enantiomers in chiral liquid chromatography through their system peak. *Chirality*, 6 (1994) 148-155; C.A., 122 (1995) 150539j.
- 2736 Li, S.: Liquid chromatographic separation of enantiomers and structurally-related compounds on β -cyclodextrin stationary phases. Avail. NCL Order No. DANN74572, 1992, 205 pp.; C.A., 120 (1994) 280434r.
- 2737 Lipkowitz, K.B.: Theoretical studies of type II-V chiral stationary phases. *J. Chromatogr. A*, 694 (1995) 14-37 - a review with 56 refs.
- 2738 Maier, N.M., Uray, G., Kleidernigg, O.P. and Lindner, W.: Diphenylethanediamine (DPEDA) derivatives as chiral selectors: IV. A comparison of 3,5-dinitrobenzoylated with Pirkle's standard(R)-phenylglycine-derived phase in normal phase HPLC. *Chirality*, 6 (1994) 116-128; C.A., 122 (1995) 95579v.
- 2739 Matlin, S.A., Tiritan, E.M., Crawford, A.J., Cass, Q.B. and Boyd, D.: HPLC with carbohydrate carbamate chiral phases: influence of chiral phase structure on enantioselectivity. *Chirality*, 6 (1994) 135-140; C.A., 122 (1995) 150538h.
- 2740 Nakamura, K., Fujima, H., Kitagawa, H., Wada, H. and Makino, K.: Preparation and chromatographic characteristics of a chiral-recognizing perphenylated cyclodextrin column. *J. Chromatogr. A*, 694 (1995) 111-118.
- 2741 Oda, Y., Mano, N., Asakawa, N., Yoshida, Y. and Sato, T.: Investigation of retention behavior for racemate drugs on avidin- and modified avidin-column. *J. Liq. Chromatogr.*, 17 (1994) 3393-3409.
- 2742 Oguni, K., Oda, H. and Ichida, A.: Development of chiral stationary phases consisting of polysaccharide derivatives. *J. Chromatogr. A*, 694 (1995) 91-100 - a review with 18 refs.
- 2743 Ôi, N., Kitahara, H. and Aoki, F.: Direct enantiomer separations by high-performance liquid chromatography with chiral urea derivatives as stationary phases. *J. Chromatogr. A*, 694 (1995) 129-134.
- 2744 Perrin, S.R.: Fast liquid chromatography for the analysis of enantiomers. *Prog. Pharm. Biomed. Anal.*, 1 (1994) 91-124; C.A., 122 (1995) 234469z - a review with refs.
- 2745 Rustum, A.M.: Determination of chiral purity of ethyl nipeccate using a Chiralcel-OG column. *J. Chromatogr. A*, 696 (1995) 75-81.
- 2746 Sato, Y., Shinomiya, K. and Kabasawa, Y.: (Aqueous two-phase partitioning method by using rotation locular countercurrent chromatograph. An application to enantiomeric separation). *Nippon Kagaku Kaishi*, (1994) 1067-1071; C.A., 122 (1995) 75692y.
- 2747 Schurig, V., Jung, M., Mayer, S., Fluck, M., Negura, S. and Jakubetz, H.: Unified enantioselective capillary chromatography on a Chirasil-DEX stationary phase. Advantages of column miniaturization. *J. Chromatogr. A*, 694 (1995) 119-128.
- 2748 Shibukawa, A., Kadohara, M., He, J.-y., Nishimura, M., Naito, S. and Nakagawa, T.: Study of the enantioselective binding between BOF-4272 and serum albumins by means of high-performance frontal analysis. *J. Chromatogr. A*, 694 (1995) 81-89.
- 2749 Stalcup, A.M. and Williams, K.L.: Comparison 1-(1-naphthyl)ethylcarbamate derivatives of a carbohydrate bonded chiral stationary phase. *J. Chromatogr. A*, 695 (1995) 185-193.
- 2750 Veigl, E., Böhs, B., Mandl, A., Krametter, D. and Lindner, W.: Comparison of (S)-N-(3,5-dinitrobenzoyl)tyrosine derivatives as chiral selectors for high-performance liquid chromatographic enantioseparations. *J. Chromatogr. A*, 694 (1995) 135-150.
- 2751 Veigl, E., Böhs, B., Mandl, A., Krametter, D. and Lindner, W.: Evaluation of silica gel-based brush type chiral cation exchangers with (S)-N-(3,5-dinitrobenzoyl)tyrosine as chiral selector: attempt to interpret the discouraging results. *J. Chromatogr. A*, 694 (1995) 151-161.
- 2752 Yashima, E., Yamada, M., Kaida, Y. and Okamoto, Y.: Computational studies on chiral discrimination mechanism of cellulose trisphenylcarbamate. *J. Chromatogr. A*, 694 (1995) 347-354.
- See also 2567, 2608, 2654, 2658, 2664, 2675, 2691, 2765, 2893, 3007, 3032, 3033, 3054, 3087, 3361, 3471, 3475, 3494, 3549, 3558, 3623, 3626, 3633, 3634, 3635, 3636, 3638, 3642, 3661, 3670, 3672, 3673, 3675, 3676, 3686, 3697, 3701, 3703, 3705, 3706, 3715, 3739, 3756, 3770, 3771, 3773, 3837, 3904.
- 4h. Other special techniques
- 2753 Abbasi, W.A. and Streat, M.: Comments on extraction chromatography. *J. Radioanal. Nucl. Chem.*, 188 (1994) 163-167; C.A., 122 (1995) 95404j.
- 2754 Deckert, P. and Artl, W.: (Simulated countercurrent chromatography). *Chem. - Ing. - Tech.*, 66 (1994) 1334-1340; C.A., 122 (1995) 84473e - a review with 36 refs.
- 2755 Foucault, A.P.: Solvent systems in centrifugal partition chromatography. *Chromatogr. Sci. Ser.*, 68(Centrifugal Partition Chromatography) (1995) 71-97; C.A., 122 (1995) 155313r - a review with 7 refs.

- 2756 Foucault, A.P. (Editor): *Centrifugal Partition Chromatography. Chromatogr. Sci. Ser.*, Vol. 68, Dekker, New York, 1995, 415 pp.; C.A., 122 (1995) 177422c.
- 2757 Wijnhoven, J.E.G.J., Koorn, J.P., Poppe, H. and Kok, W.T.: Hol-low-fibre flow field-flow fractionation of polystyrene sulphonates. *J. Chromatogr. A*, 699 (1995) 119-129.
- 2758 Yuan, N., Liu, Z. and Zhu, D.: Chromatograph-type electrophoresis separation method. *Faming Zhuanli Shenqing Gongkai Shuomingshu* CN 1,083,072 (Cl. C07K3/14), 02 Mar. 1994; Appl. 93,108,629, 16 Jul. 1993; 8 pp.; C.A., 122 (1995) 50769r.
- 2759 Zibas, S.A. and Love, L.J.C.: Solute-micelle interactions in zwitterionic micellar chromatography. *Anal. Chim. Acta*, 299 (1994) 17-27.
- See also 2551, 2565, 2580, 2620, 2712, 2746, 2763, 2859, 2944, 3036, 3110, 3225, 3278, 3505, 3805, 3856, 3869.
5. HYDROCARBONS AND HALOGEN DERIVATIVES
- 5b. *Cyclic hydrocarbons, fullerenes*
- 2760 Belous, E.F., Lanin, S.N. and Nikitin, Yu.S.: Combined application of normal and reversed-phase high-performance liquid chromatography to determining the group composition of aromatic hydrocarbons in petroleum products. *J. Anal. Chem.*, 50 (1995) 93-100; C.A., 122 (1995) 137743b.
- 2761 Cooper, C.L., Whitaker, K.W. and Sepaniak, M.J.: Separation behavior of common fullerenes in cyclodextrin-HPLC based on computationally-derived interaction energies. *J. Liq. Chromatogr.*, 17 (1994) 4311-4325.
- 2762 Davydov, V.Ya., Filatova, G.N., Khrustaleva, N.M. and Roshchina, T.M.: Separation of fullerenes and study of their intermolecular interactions with organic compounds by chromatography. *Proc.-Electrochem. Soc.*, 94-24 (1994) 1588-1594; C.A., 122 (1995) 204204s.
- 2763 Gasper, M.P., Berthod, A., Talabardon, K. and Armstrong, D.W.: An evaluation of the differential partitioning and separation of C₆₀ and C₇₀ fullerenes in a biphasic system using centrifugal partition chromatography (CPC). *J. Liq. Chromatogr.*, 18 (1995) 1019-1034.
- 2764 Heaton, D.M., Bartle, K.D., Clifford, A.A., Myers, P. and King, B.W.: Rapid separation of polycyclic aromatic hydrocarbons by packed column supercritical fluid chromatography. *Chromatographia*, 39 (1994) 607-611.
- 2765 Husain, N., Christian, A.Y. and Warner, I.M.: Effect of co-modifiers in cyclodextrin-modified mobile phases on the reversed-phase high-performance liquid chromatographic separation of polyaromatic hydrocarbons. *J. Chromatogr. A*, 699 (1995) 73-83.
- 2766 Lan, W.G., Chee, K.K., Wong, M.K. and Lee, H.K.: Orthogonal array design as a chemometric method for the optimization of analytical procedures. Part 4. Mixed-level design and its application to the high-performance liquid chromatographic determination of polycyclic aromatic hydrocarbon. *Analyst (Cambridge)*, 120 (1995) 281-287.
- 2767 Lintelmann, J., Boos, K.-S. and Kettrup, A.: (Biomonitoring of exposure to PAH using a coupled-column HPLC-method for the determination of 1-hydroxypyrene - part 1). *Bioforum*, 17 (1994) 442-445; C.A., 122 (1995) 232766v.
- 2768 Mäkelä, M. and Pyy, L.: Effect of temperature on retention time reproducibility and on the use of programmable fluorescence detection of fifteen polycyclic aromatic hydrocarbons. *J. Chromatogr. A*, 699 (1995) 49-57.
- 2769 Masterov, V.F., Chudnovski, F.A., Kozyrev, S.V., Zaidenberg, A.S., Rozhkova, N.N., Podbsenova, N.G. and Stefanovich, G.B.: (Microwave absorption in fullerene-containing shungites). *Mol. Cryst. Liq. Cryst. Sci. Technol., Sect. C*, 4 (1994) 213-216; C.A., 122 (1995) 121996a.
- 2770 Mueller, W., Broell, R., Weber, E., Daimer, J. and Mueller, R.: Chromatographic separation of fullerenes with a nonpolar arom. solvent as the elutriating agent. *PCT Int. Appl. WO 94 06,715* (Cl. C01B31/02), 31 Mar. 1994, DE Appl. 4,230,915, 16 Sep. 1992; 14 pp.; C.A., 120 (1994) 248667h.
- 2771 Ohmacht, R., Kele, M. and Matus, Z.: Reversed phase C₁₈ packings for the separation of polycyclic aromatic hydrocarbons by high performance liquid chromatography. *Chromatographia*, 39 (1994) 668-672.
- 2772 Podosenova, N.G.: (Separation of fullerenes by liquid chromatography). *Zh. Fiz. Chim.*, 68 (1994) 1298-1303; C.A., 122 (1995) 121984v.
- 2773 Reupert, R. and Brausen, G.: (Determination of polycyclic aromatic hydrocarbons in water, sludges, sediments, and soils by high-performance liquid chromatography). *Acta Hydrochim. Hydrobiol.*, 22 (1994) 202-215; C.A., 122 (1995) 122246t.
- 2774 Rozbeh, M. and Hurtubise, R.J.: Reversed-phase chromatographic separation of benzo[a]pyrene metabolites with β -cyclodextrin as a mobile phase modifier. *J. Liq. Chromatogr.*, 18 (1995) 17-37.
- 2775 Takeuchi, T., Hu, R. and Miwa, T.: Retention behavior of aromatic hydrocarbons in reversed-phase liquid chromatography with ion-exchange-induced stationary phases. *Chromatographia*, 39 (1994) 597-601.
- 2776 Thomas, D., Crain, S.M., Sim, P.G. and Benoit, F.M.: Use of immobilized copper ion chromatography and on-line mass spectrometry with atmospheric pressure chemical ionization for the profiling of complex mixtures of polycyclic aromatic compounds. *J. Chromatogr. A*, 695 (1995) 1-9.
- 2777 Whiton, R.S., Witherspoon, C.L. and Buckley, T.J.: Improved high-performance liquid chromatographic method for the determination of polycyclic aromatic hydrocarbon metabolites in human urine. *J. Chromatogr. B*, 665 (1995) 390-394.
- 2778 Wise, S.A., Deissler, A. and Sander, L.C.: Liquid chromatographic detection of polycyclic aromatic hydrocarbon isomers of molecular weight 278 and 302 in environmental standard reference materials. *Polycyclic Aromat. Compd.*, 3 (1993) 169-184; C.A., 122 (1995) 150587y.
- 2779 Wise, S.A., Schantz, M.M., Benner, B.A., Jr., Hays, M.J. and Schiller, S.B.: Certification of polycyclic aromatic hydrocarbons in a marine sediment standard reference material. *Anal. Chem.*, 67 (1995) 1171-1178.
- 2780 Yamamoto, K., Funasaka, H., Takahashi, T. and Akasaka, T.: Isolation of an ESR-active metallofullerene of La@C₈₂. *J. Phys. Chem.*, 98 (1994) 2008-2011; C.A., 120 (1994) 207225m.

2781 Zobel, H., Nerlich, B. and Panning, H.: Liquid chromatographic procedure and apparatus for separation and qualitative and quantitative determination of polycyclic aromatic hydrocarbon-containing mixtures. *Ger. Offen.* DE 4,312, 925 (Cl. G01N33/22), 10 Oct. 1994, Appl. 15 Apr. 1993; 7 p.; C.A., 122 (1995) 85295k.

See also 2634.

5c. Halogen derivatives

2782 Finizio, A., di Guardo, A. and Vighi, M.: Improved RP-HPLC determination of Kow for some chloroaromatic chemicals using molecular connectivity indices. *SAR QSAR Environ. Res.*, 2 (1994) 249-260; C.A., 122 (1995) 98877b.

2783 Golba, W., Przybylski, Z., Ceckiewicz, S. and Krasodomski, M.: (Chromatographic methods for determination of polychlorinated biphenyls). *Nafta-Gaz*, 50 (1994) 405-410; C.A., 122 (1995) 204161a a review with 33 refs.

2784 Takasuga, T., Ohi, E. and Inoue, T.: Complete isolation and determination of *mono-ortho* and *non-ortho* substituted PCBs and PCDDs/PCDFs by HPLC, on porous graphitic carbon with HRGC/HRMS. *Organohalogen Compd.*, 11 (1993) 101-104; C.A., 122 (1995) 204237e.

See also 3267.

5d. Complex hydrocarbon mixtures (incl. analysis of tars, bitumens and mineral oils)

2785 Fabinska, M.: (Identification of the group components from extrographic separation of the low-temperature coal tar). *Karbo-Energochem.-Ekol.*, 38 (1993) 55-57; C.A., 122 (1995) 165202t.

2786 Whittaker, M. and Pollard, S.J.T.: Characterization of refractory wastes at hydrocarbon-contaminated sites: 1. Rapid column fractionation and thin layer chromatography of reference oils. *J. Planar Chromatogr.*, 7 (1994) 354-361.

See also 3467, 3838.

6. ALCOHOLS

2787 Bourdeaux, R., Capolaghi, B., Lacroix, C., Younos, C. and Lehr, P.R.: Assay of urinary free and conjugated 3-methoxy-4-hydroxyphenylethyleneglycol by high-performance liquid chromatography with amperometric detection. *J. Chromatogr. B*, 665 (1995) 45-51.

2788 Kok, S.J., Posthumus, R., Bakker, I., Gooijer, C., Brinkman, U.A.T. and Velthorst, N.H.: Identification of stereoisomeric benzo[a]pyrene tetrols by reversed-phase liquid chromatography coupled semi-on-line to fluorescence line-narrowing spectroscopy. *Anal. Chim. Acta*, 303 (1995) 3-10.

2789 Tjioe, S.W. and Hurtubise, R.J.: Separation and characterization of tetrol metabolites of benzo[a]pyrene-DNA adducts using HPLC and solid-matrix room temperature luminescence. *Talanta*, 42 (1995) 59-64; C.A., 122 (1995) 125436f.

2790 Van Damme, F. and Oomens, A.C.: Determination of residual free epoxide in polyether polyols by derivatization with diethylammonium N,N-diethylthiocarbamate and liquid chromatography. *J. Chromatogr. A*, 696 (1995) 41-47.

2791 Wanaka, K. and Murui, T.: (Determination of fluorescence labeled alcohols from waxes in vegetable oils by high performance liquid chromatography). *Yukagaku*, 44 (1995) 43-48; C.A., 122 (1995) 104198z.

For additional information see C.A.:

122 (1995) 105252z.

See also 3784, 3830.

7. PHENOLS

2792 Alymova, A.T., Korenman, Ya.I., Bobrinskaya, E.V., Khizver, D.R. and Li, A.: (Separation and preconcentration of traces of aromatic hydroxy compounds by extraction and chromatography with mixed stationary phases). *Zh. Anal. Khim.*, 49 (1994) 1197-1201; C.A., 122 (1995) 204215w.

2793 Amarowicz, R., Wanasundara, P.K.J.P.D. and Shahidi, F.: Chromatographic separation of flaxseed phenolics. *Nahrung*, 38 (1994) 520-526; C.A., 122 (1995) 104164k.

2794 Galceran, M.T. and Jáuregui, O.: Determination of phenols in sea water by liquid chromatography with electrochemical detection after enrichment by using solid-phase extraction cartridges and disks. *Anal. Chim. Acta*, 304 (1995) 75-84.

2795 Kawahara, A., Yamada, M., Kato, Y., Nakamura, M., Nishijima, M. and Ito, Y.: (Systematic determination of diphenyl, *o*-phenylphenol, thiabendazole and imazalil in citrus fruits). *Foods Food Ingredients J. Jpn.*, 163 (1995) 104-110; C.A., 122 (1995) 185656x.

2796 Krämer, P.M., Li, Q.X. and Hammock, B.D.: Integration of liquid chromatography with immunoassay: an approach combining the strengths of both methods. *J. Assoc. Off. Anal. Chem.*, 77 (1994) 1275-1287.

2797 Pocerull, E., Sánchez, G., Borrull, F. and Marcé, R.M.: Automated on-line trace enrichment and determination of phenolic compounds in environmental waters by high-performance liquid chromatography. *J. Chromatogr. A*, 696 (1995) 31-39.

2798 Rozbeh, M. and Hurtubise, R.J.: Optimum separation and compound class separation of the metabolites of benzo[a]pyrene-DNA adducts with reversed-phase liquid chromatography. *J. Liq. Chromatogr.*, 17 (1994) 3351-3367.

2799 Sakuma, S., Kikuchi, C. and Kowaka, M.: Automated analysis of the total polyphenols content of beer and wort using an intelligent autosampler. *J. Am. Soc. Brew. Chem.*, 53 (1995) 29-32; C.A., 122 (1995) 158915z.

2800 Variyar, P.S. and Bandyopadhyay, C.: Estimation of phenolic compounds in green pepper berries (*Piper nigrum* L.) by high-performance liquid chromatography. *Chromatographia*, 39 (1994) 743-746.

2801 Yang, Y. and He, J.: (Direct determination of microamount of phenol in urine by reversed-phase high performance liquid chromatography). *Sepu*, 13 (1995) 70-71; C.A., 122 (1995) 153472t.

See also 2653, 2774, 2825, 2833, 3000, 3799, 3804, 3823, 3825.

8. SUBSTANCES CONTAINING HETEROCYCLIC OXYGEN

8a. Flavonoids

- 2802 Bongartz, D. and Hesse, A.: Boronic acid affinity chromatography as a tool for determination of quercetrin in urine. *Eur. J. Clin. Chem. Clin. Biochem.*, 33 (1995) A17.
- 2803 Cui, J.M., Chung, H.S. and Woo, W.S.: Determination of isoflavonoid glucosides in rhizomes of *Belamcanda chinensis* by high-performance liquid chromatography. *Saengyak Hakhoechi*, 24 (1993) 309-312; *C.A.*, 120 (1994) 280388d.
- 2804 Del Rio, J.A., Castillo, J. and Benavente-García, O.: Elucidation by reverse phase HPLC of some citrus flavanones and their respective dihydrochalcones: structural study. *J. Liq. Chromatogr.*, 17 (1994) 3461-3477.
- 2805 Hammouda, F.M., El-Nasr, M.M.S., Ismail, S.I. and Shahat, A.A.: Quantitative determination of the active constituent in Egyptian cultivated *Cynara scolymus*. *Int. J. Pharmacogn.*, 31 (1993) 299-304; *C.A.*, 120 (1994) 200497a.
- 2806 Kurkin, V.A., Braslavskii, V.B. and Zapesochynaya, G.G.: (Study of extracts of propolis and *Populus balsamifera* buds by high-performance liquid chromatography). *Zh. Fiz. Khim.*, 68 (1994) 1816-1818; *C.A.*, 122 (1995) 75690w.
- 2807 Liu, B., Anderson, D., Ferry, D.R., Seymour, L.W., de Takats, P.G. and Kerr, D.J.: Determination of quercetin in human plasma using reversed-phase high-performance liquid chromatography. *J. Chromatogr. B*, 666 (1995) 149-155.
- 2808 Pietrogrande, M.C. and Kahie, Y.D.: Effect of the mobile and stationary phases on RP-HPLC retention and selectivity of flavonoid compounds. *J. Liq. Chromatogr.*, 17 (1994) 3655-3670.
- 2809 Rehwald, A., Meier, B. and Sticher, O.: Qualitative and quantitative reversed-phase high-performance liquid chromatography of flavonoids in *Passiflora incarnata* L. *Pharm. Acta Helv.*, 69 (1994) 153-158; *C.A.*, 122 (1995) 142702s.
- 2810 Supko, J.G. and Phillips, L.R.: High-performance liquid chromatographic assay for genistein in biological fluids. *J. Chromatogr. B*, 666 (1995) 157-167.

See also 3769, 3803, 3804.

8b. Aflatoxins and other mycotoxins

- 2811 Alachaher, L., Abdeddaim, K. and Guermouche, M.H.: (Behavior of aflatoxins in liquid chromatography). *J. Soc. Alger. Chim.*, 4 (1994) 57-96; *C.A.*, 122 (1995) 212352d.
- 2812 Barbieri, G., Bergamini, C., Ori, E. and Resca, P.: Aflatoxin M₁ in Parmesan cheese: HPLC determination. *J. Food Sci.*, 59 (1994) 1313, 1331; *C.A.*, 122 (1995) 131397v.
- 2813 Dragacci, S., Gleizes, E., Fremy, J.M. and Candish, A.A.G.: Use of immunoaffinity chromatography as a purification step for the determination of aflatoxin M₁ in cheeses. *Food Addit. Contam.*, 12 (1995) 59-65; *C.A.*, 122 (1995) 238045f.

- 2814 Guo, Y. and Fu, C.: (HPLC determination of trace trichothecene mycotoxin in corn). *Fenxi Ceshi Xuebao*, 13 (1994) 59-62; *C.A.*, 122 (1995) 79317c.
- 2815 Maksimovic, M.Z. and Vuksanovic, M.: Application of some extraction techniques in analysis of aflatoxins (B₁, B₂, G₁ and G₂) in animal food by HPLC using UV and fluorescence detectors connected in series. *Eur. J. Clin. Chem. Clin. Biochem.*, 33 (1995) A23.
- 2816 Maragos, C.M. and Richard, J.L.: Quantitation and stability of fumonisins B₁ and B₂ in milk. *J. Assoc. Off. Anal. Chem.*, 77 (1994) 1162-1167.
- 2817 Milat, M.-L. and Blein, J.-P.: *Cercospora beticola* toxins. III. Purification, thin-layer and high-performance liquid chromatographic analyses. *J. Chromatogr. A*, 699 (1995) 277-283.
- 2818 Roch, O.G., Blunden, G., Coker, R.D. and Nawaz, S.: The validation of a solid phase clean-up procedure for the analysis of aflatoxins in groundnut cake using HPLC. *Food Chem.*, 52 (1995, Pub. 1994) 93-98; *C.A.*, 122 (1995) 79369w.
- 2819 Trucksess, M.W.: Mycotoxins. *J. Assoc. Off. Anal. Chem.*, 78 (1995) 135-141 - a review with 120 refs.
- 2820 Wilkes, J.G., Sutherland, J.B., Churchwell, M.I. and Williams, A.J.: Determination of fumonisins B₁, B₂, B₃ and B₄ by high-performance liquid chromatography with evaporative light-scattering detection. *J. Chromatogr. A*, 695 (1995) 319-323.
- 2821 Zimmerli, B. and Dick, R.: Determination of ochratoxin A at the ppt level in human blood, serum, milk and some foodstuffs by high-performance liquid chromatography with enhanced fluorescence detection and immunoaffinity column cleanup: methodology and Swiss data. *J. Chromatogr. B*, 666 (1995) 85-99.

8c. Other compounds with heterocyclic oxygen (incl. tannins)

- 2822 Clement, B. and Mladek, C.: Biotransformationsuntersuchungen im Hühnerrei. Teil 1: Metabolische Umwandlung von 7-Ethoxycoumarin. *Pharmazie*, 50 (1995) 207-210.
- 2823 Hiller, D.L. and Cole, R.O.: Short-columns gradient elution high-performance liquid chromatography for the rapid detection of 7-ethoxycoumarin metabolites in liver-slice incubates. *Anal. Biochem.*, 227 (1995) 251-254.
- 2824 Maroni, T. and Lazzari, P.: (Isocratic determination of furosine using HPLC). *Ind. Aliment. (Pinerolo)*, 33 (1994) 964-966; *C.A.*, 122 (1995) 238011s.
- 2825 Wolbis, M. and Nowak, S.: Coumarins and phenolic acids from the underground part of *Scopolia lurida*. *Acta Pol. Pharm.*, 51 (1994) 171-174; *C.A.*, 122 (1995) 114731g.
- 2826 Xu, Y., Mirocha, C.J. and Xie, W.: Analysis by liquid chromatography of fusarochromanone (TDP-1) added to corn and wheat. *J. Assoc. Off. Anal. Chem.*, 77 (1994) 1179-1183.

For additional information see *C.A.*:

122 (1995) 100764h, 127804h, 142705v, 155422a.

See also 3776, 3805.

9. OXO COMPOUNDS, ETHERS, EPOXIDES AND QUINONES

- 2827 Aboull-Enein, H.Y. and Abou-Basha, L.I.: Simple HPLC method for the determination of thymoquinone in black seed oil (*Nigella sativa* Linn). *J. Liq. Chromatogr.*, 18 (1995) 895-902.

- 2828 Berger, G., Kléo, J., Breton, J., Gilles, N. and Lirsac, P.N.: ^{18}O and ^{13}C labeling of photosynthetic and related quinones and their purification by high performance liquid chromatography. *J. Liq. Chromatogr.*, 17 (1994) 4531-4539.
- 2829 Coco, F.L., Valentini, C., Novelli, V. and Ceccon, L.: Liquid chromatographic determination of 2-furaldehyde and 5-hydroxymethyl-2-furaldehyde in beer. *Anal. Chim. Acta*, 306 (1995) 57-64.
- 2830 Forgacs, E.: Use of porous graphitized carbon column for determining taxol in *Taxus baccata*. *Chromatographia*, 39 (1994) 740-742.
- 2831 Granados, J.W., Mir, M.V., Lopez Garcia Serrana, H. and Lopez Martinez, M.C.: Comparison of spectrophotometric and chromatographic methods of determination of furanic aldehydes in wine distillates. *Food Chem.*, 52 (1995, Pub. 1994) 203-208; *C.A.*, 122 (1995) 79364r.
- 2832 Liu, G.: Oxidation of acetophenone in a high temperature open-tubular capillary column liquid chromatographic system. *Chromatographia*, 40 (1995) 99-101.
- 2833 Masselter, S., Zemmann, A. and Bobleter, O.: Analysis of lignin degradation products by capillary electrophoresis. *Chromatographia*, 40 (1995) 51-57.
- 2834 Sitholé, B.B., Nyarku, S. and Allen, L.H.: Chromatographic methods for the determination of alkylketene dimer in pulp and paper matrices. *Analyst (Cambridge)*, 120 (1995) 1163-1169.
- 2835 Toyooka, T. and Liu, Y.-M.: Determination of aldehydes by high-performance liquid chromatography with fluorescence detection after labelling with 4-(2-carbazoylpyrrolidin-1-yl)-7-(N,N-dimethylaminosulfonyl)-2,1,3-benzoxadiazole. *J. Chromatogr. A*, 695 (1995) 11-18.
- 2836 Winkler, R. and Maier, K.: (Testing an HPLC method for determination of malondialdehyde (MDA) in urine, plasma and serum with regard to possible interferences by glucose and streptozotocin (STZ)). *Klin. Labor.*, 40 (1994) 351-355; *C.A.*, 122 (1995) 182302t.
- 2837 Wu, H.-Y. and Jin, J.-K.: Determination of aldehydic lipid peroxidation products with dabsylhydrazine by high-performance liquid chromatography. *Anal. Chem.*, 67 (1995) 1603-1612.
- See also 2643, 2798, 2817, 2896, 3616, 3799, 3830.
10. CARBOHYDRATES
- 10a. Mono and oligosaccharides. Structural studies
- 2838 Buongiorno, A., Morviducci, L., Zorretta, D., Leonetti, F., Tamburrano, G. and Giaccari, A.: Novel method for the determination of hepatic glycogenolysis and gluconeogenesis *in vivo* in rats: preliminary results in hypoglycemia. *Front. Endocrinol.*, 9(Highlights in Molecular and Clinical Endocrinology) (1994) 383-386; *C.A.*, 122 (1995) 234517p.
- 2839 Chusney, G.D., Philippa, M. and Pickup, J.C.: Comparison of micro-enzymatic and high-performance liquid chromatographic methods for the assay of serum 1,5-anhydroglucitol. *Clin. Chim. Acta*, 235 (1995) 91-99.
- 2840 Cooper, G.A. and Rohrer, J.S.: Separation of neutral asparagine-linked oligosaccharides by high-pH anion-exchange chromatography with pulsed amperometric detection. *Anal. Biochem.*, 226 (1995) 182-184.
- 2841 Corradini, C.: Recent advances in carbohydrate analysis by high-performance anion-exchange chromatography coupled with pulsed amperometric detection (HPAEC-PAD). *Ann. Chim. (Rome)*, 84 (1994) 385-396; *C.A.*, 122 (1995) 187876t.
- 2842 Evangelisto, M.F., Adams, R.E., Murray, W.V. and Caldwell, G.W.: Preparative high-performance liquid chromatographic separation of fluorodeoxy sugars. *J. Chromatogr. A*, 695 (1995) 128-131.
- 2843 Fleming, S.C., Gill, M. and Laker, M.F.: A rapid HPLC method for the determination of free fucose in urine. A marker of malignancy in children. *J. Liq. Chromatogr.*, 18 (1995) 173-180.
- 2844 Hino, K., Yamamoto, S., Sano, O., Taniguchi, Y., Kohno, K., Usui, M., Fukuda, S., Hanzawa, H., Haruyama, H. and Kurimoto, M.: Carbohydrate structures of the glycoprotein allergen *Cry j I* from Japanese cedar (*Cryptomeria japonica*) pollen. *J. Biochem. (Tokyo)*, 117 (1995) 289-295.
- 2845 Hooper, L.V., Beranek, M.C., Manzella, S.M. and Baenziger, J.U.: Differential expression of GalNAc-4-sulfotransferase and GalNAc-transferase results in distinct glycoforms of carbonic anhydrase VI in parotid and submaxillary glands. *J. Biol. Chem.*, 270 (1995) 5985-5993.
- 2846 Karamanos, N.K., Vanky, P., Syrokou, A. and Hjerpe, A.: Identity of dermatan and chondroitin sequences in dermatan sulfate chains determined by using fragmentation with chondroitinases and ion-pair high-performance liquid chromatography. *Anal. Biochem.*, 225 (1995) 220-230.
- 2847 Karlsson, N.G. and Hansson, G.C.: Analysis of monosaccharide composition of mucin oligosaccharide alditols by high-performance anion-exchange chromatography. *Anal. Biochem.*, 224 (1995) 538-541.
- 2848 Kobata, A., Kochibe, N. and Endo, T.: Affinity chromatography of oligosaccharides on *Psathyrella velutina* lectin column. *Methods Enzymol.*, 247(Neoglycoconjugates, Pt. B) (1994) 228-237; *C.A.*, 122 (1995) 182313x.
- 2849 Lopez-Hernandez, J., Gonzalez-Castro, M.J., Vazquez-Blanco, M.E., Vazquez-Oderiz, M.L. and Simal-Lozano, J.: HPLC determination of sugars and starch in green beans. *J. Food Sci.*, 59 (1994) 1048-1049; *C.A.*, 122 (1995) 104173n.
- 2850 Molla, E., Camara, M.M., Diez, C. and Torija, M.E.: (Studies on the determination of sugars in fruits and their products). *Alimentaria (Madrid)*, 254 (1994) 87-93; *C.A.*, 122 (1995) 54322n.
- 2851 Nakagawa, H., Kawamura, Y., Kato, K., Shimada, I., Arata, H. and Takahashi, N.: Identification of neutral and sialyl N-linked oligosaccharide structures from human serum glycoproteins using three kinds of high-performance liquid chromatography. *Anal. Biochem.*, 226 (1995) 130-138.
- 2852 Natunen, J., Niemela, R., Penttila, L., Seppo, A., Ruohtula, T. and Renkonen, O.: Enzymic synthesis of two lacto-N-neo-hexaose-related Lewis x heptasaccharides and their separation by chromatography on immobilized wheat germ agglutinin. *Glycobiology*, 4 (1994) 577-583; *C.A.*, 122 (1995) 133607n.
- 2853 Neville, D.C.A., Field, R.A. and Ferguson, M.A.J.: Hydrophobic glycosides of N-acetylglucosamine can act as primers for polylactosamine synthesis and can affect glycolipid synthesis *in vivo*. *Biochem. J.*, 307 (1995) 791-797.

- 2854 Nojima, K., Isogami, C. and Kobashi, M.: (Application of 4-phenyl-1,2,4-triazoline-3,5-dione for the analysis of sorbic acid in foods). *Jpn. J. Toxicol. Environ. Health*, 40 (1994) 467-471; *C.A.*, 122 (1995) 29975v.
- 2855 Price, N.P.J. and Carlson, R.W.: Rhizobial lipo-oligosaccharide nodulation factors: multidimensional chromatographic analysis of symbiotic signals involved in the development of legume root nodules. *Glycobiology*, 5 (1995) 233-242; *C.A.*, 122 (1995) 238564z.
- 2856 Rodríguez-Aparicio, L.B., Ferrero, M.A. and Reglero, A.: N-Acetyl-D-neuraminic acid synthesis in *Escherichia coli* K1 occurs through condensation of N-acetyl-D-mannosamine and pyruvate. *Biochem. J.*, 308 (1995) 501-505.
- 2857 Romppanen, J. and Mononen, I.: Age-related reference values for urinary excretion of sialic acid and deoxysialic acid: application to diagnosis of storate disorders of free sialic acid. *Clin. Chem. (Washington)*, 41 (1995) 544-547.
- 2858 Saitoh, H., Takagaki, K., Majima, M., Nakamura, T. and Matsuki, A., Kasai, M., Narita, H. and Endo, M.: Enzymic reconstruction of glycosaminoglycan oligosaccharide chains using the transglycosylation reaction of bovine testicular hyaluronidase. *J. Biol. Chem.*, 270 (1995) 3741-3747.
- 2859 Solbrig-Lebuhn, H.: (Two dimensional HPLC of fluorescent labeled oligosaccharides). *BioTec (Wuerzburg)*, 6 (1994) 22; *C.A.*, 122 (1995) 75704d.
- 2860 Takahashi, N., Nakagawa, H., Fujikawa, K., Kawamura, Y. and Tomia, N.: Three-dimensional elution mapping of pyridylaminated N-linked neutral and sialyl oligosaccharides. *Anal. Biochem.*, 226 (1995) 139-146.
- 2861 Takeuchi, T., Kusakabe, I., Yoshida, S.Y.S. and Murata, K.: Separation of oligomannuronic acid based on degree of polymerization. *Jpn. Kokai Tokkyo Koho JP 06,172,375 [94,172,375] (Cl. CO7H1/06)*, 21 Jun. 1994, Appl. 92/325,289, 04 Dec. 1992; 4 pp.; *C.A.*, 122 (1995) 106386h.
- 2862 Yutani, S., Toh, G. and Honda, S.: (Analysis of fructooligosaccharides by HPLC and HPCE). *Kuromatogurafi*, 15 (1994) 210-211; *C.A.*, 122 (1995) 127992t.

For additional information see C.A.:

122 (1995) 79347n, 101143s, 158794j, 187881r, 187890t.

See also 2588, 2869, 2872, 2876, 2939, 3206, 3226, 3232, 3405, 3417, 3596, 3812.

10b. Polysaccharides, mucopolysaccharides, lipopolysaccharides

- 2863 Dubin, P.L.: Nonionic polysaccharides as calibration standards for aqueous size exclusion chromatography. *Carbohydr. Polym.*, 25 (1994) 295-303; *C.A.*, 122 (1995) 208918a.
- 2864 Hashimoto-Uoshima, M., Hascall, V.C., MacCallum, D.K. and Yanagishita, M.: Biosynthesis of proteoglycans and hyaluronic acid by rat oral epithelial cells (keratinocytes) *in vitro*. *Arch. Biochem. Biophys.*, 316 (1995) 724-732.
- 2865 Kasinath, B.S.: Effect of insulin on high-glucose medium-induced changes in rat glomerular epithelial cell metabolism of glycoconjugates. *Arch. Biochem. Biophys.*, 318 (1995) 286-294.

- 2866 Lagana, A., Marino, A., Fago, G. and Pardo Martinez, B.: Determination of free N-acetyl neuraminic acid by ultrafiltration and liquid chromatography in the serum of normal and cancer patients. *Anal. Chim. Acta*, 306 (1995) 65-71.
- 2867 Midura, R.J., Salustri, A., Calabro, A., Yanagishita, M. and Hascall, V.C.: High-resolution separation of disaccharide and oligosaccharide alditols from chondroitin sulfate, dermatan sulfate and hyaluronan using CarboPac PA1 chromatography. *Glycobiology*, 4 (1994) 333-342; *C.A.*, 122 (1995) 75676w.
- 2868 Müller, A., Pretus, H.A., McNamee, R.B., Jones, E.L., Browder, I.W. and Williams, D.L.: Comparison of the carbohydrate biological response modifiers Krestin, schizophyllan and glucan phosphate by aqueous size exclusion chromatography with inline argon-ion multi-angle laser light scattering photometry and differential viscometry detectors. *J. Chromatogr. B*, 666 (1995) 283-290.
- 2869 Ng, C.K., Handley, C.J., Preston, B.N., Robinson, H.C., Bolis, S. and Parker, G.: Effect of exogenous hyaluronan and hyaluronan oligosaccharides on hyaluronan and aggrecan synthesis and catabolism in adult articular cartilage explants. *Arch. Biochem. Biophys.*, 316 (1995) 596-606.
- 2870 Shao, M.-C. and Wold, F.: The effect of the protein matrix proximity on glycan reactivity in a glycoprotein model. *Eur. J. Biochem.*, 228 (1995) 79-85.
- 2871 Van Det, N.F., van den Born, J., Tamsma, J.T., Verhagen, N.A.M., van den Heuvel, L.P.W.J., Berden, J.H.M., Bruijn, J.A., Daha, M.R. and van der Woupe, F.J.: Proteoglycan production by human glomerular visceral epithelial cells and mesangial cells *in vitro*. *Biochem. J.*, 307 (1995) 759-768.
- 2872 Volpi, N., Cusmano, M. and Venturelli, T.: Qualitative and quantitative studies of heparin and chondroitin sulfates in normal human plasma. *Biochim. Biophys. Acta*, 1243 (1995) 49-58.
- 2873 Wong, K.S. and Jane, J.: Effects of pushing agents on the separation and detection of debranched amylopectin by high-performance anion-exchange chromatography with pulsed amperometric detection. *J. Liq. Chromatogr.*, 18 (1995) 63-80.
- 2874 Yang, J. and Z.: (Determination of molecular weights and their distributions of polysaccharose in astragalus by using 150-C ALC/GPC gel permeation chromatography). *Taiyuan Gongye Daxue Xuebao*, 25 (1994) 77-84; *C.A.*, 122 (1995) 100763g.

For additional information see C.A.:

122 (1995) 185680a.

See also 2629, 2846, 2949, 3139, 3206, 3232, 3475, 3818.

10c. Glycoproteins and their constituents

- 2875 Altman, E., Schäffer, C., Brisson, J.-R. and Messner, P.: Characterization of the glycan structure of a major glycopeptide from the surface layer glycoprotein of *Clostridium thermosaccharolyticum*. *Eur. J. Biochem.*, 229 (1995) 308-315.
- 2876 Ashton, D.S., Beddell, C.R., Cooper, D.J. and Lines, A.C.: Determination of carbohydrate heterogeneity in the humanised antibody CAMPATH 1H by liquid chromatography and matrix-assisted laser desorption ionisation mass spectrometry. *Anal. Chim. Acta*, 306 (1995) 43-48.

- 2877 Cho, M. and Cummings, R.D.: Galectin-1, a β -galactoside-binding lectin in chinese hamster ovary cells. I. Physical and chemical characterization. *J. Biol. Chem.*, 270 (1995) 5198-5206.
- 2878 Coppolino, M., Migliorini, M., Argraves, W.S. and Dedhar, S.: Identification of a novel form of the α_3 integrin subunit: covalent association with transferrin receptor. *Biochem. J.*, 306 (1995) 129-134.
- 2879 Hokke, C.H., Bergwerff, A.A., van Dedem, G.W.K., Kamerling, J.P. and Vliegthart, J.F.G.: Structural analysis of the sialylated N- and O-linked carbohydrate chains of recombinant human erythropoietin expressed in Chinese hamster ovary cells. Sialylation patterns and branch location of dimeric N-acetylglucosamine units. *Eur. J. Biochem.*, 228 (1995) 981-1008.
- 2880 Nakazawa, K., Isomura, T. and Shimoeda, S.: Proteoglycan synthesis by chicken corneal explants. *J. Biochem. (Tokyo)*, 117 (1995) 697-706.
- 2881 Nakazawa, K., Suzuki, S., Wada, K. and Nakazawa, K.: Proteoglycan synthesis by corneal explants from developing embryonic chicken. *J. Biochem. (Tokyo)*, 117 (1995) 707-718.
- 2882 Osman, M.E., Menzies, A.R., Martin, B.A., Williams, P.A., Phillips, G.O. and Baldwin, T.C.: Characterization of gum arabic fractions obtained by anion-exchange chromatography. *Phytochemistry*, 38 (1995) 409-417; *C.A.*, 122 (1995) 209779t.
- 2883 Parekh, R. and Ventom, A.: Glycotechnology: an unambiguous technique for glycan analysis. *Am. Biotechnol. Lab.*, 13 (1995) 9-10; *C.A.*, 122 (1995) 155376p.
- 2884 Paszkiewicz-Gadek, A., Gindzienski, A. and Porowska, H.: The use of preparative polyacrylamide gel electrophoresis and electroelution for purification of mucus glycoproteins. *Anal. Biochem.*, 226 (1995) 263-267.
- 2885 Sareneva, T., Pirhonen, J., Cantell, K. and Julkunen, I.: N-Glycosylation of human interferon- γ glycans at Asn-25 are critical for protease resistance. *Biochem. J.*, 308 (1995) 9-14.
- 2886 Sato, T., Takio, K., Kobata, A., Greenwalt, D.E. and Furukawa, K.: Site-specific glycosylation of bovine butyrophilin. *J. Biochem. (Tokyo)*, 117 (1995) 147-157.
- 2887 Sharom, F.J., Yu, X., Chu, J.W.K. and Doige, C.A.: Characterization of the ATPase activity of P-glycoprotein from multidrug-resistant Chinese hamster ovary cells. *Biochem. J.*, 308 (1995) 381-390.
- 2888 Weber, P.L., Kornfelt, T., Klausen, N.K. and Lunte, S.M.: Characterization of glycopeptides from recombinant coagulation factor VIIa by high-performance liquid chromatography and capillary zone electrophoresis with ultraviolet and pulsed electrochemical detection. *Anal. Biochem.*, 225 (1995) 135-142.
- 2889 Young, N.M., Watson, D.C., Yaguchi, M., Adar, R., Arango, R., Rodriguez-Arango, E., Sharon, N., Blay, P.K.S. and Thibault, P.: C-terminal post-translational proteolysis of plant lectins and their recombinant forms expressed in *Escherichia coli*. Characterization of "ragged ends" by mass spectrometry. *J. Biol. Chem.*, 270 (1995) 2563-2570.
11. ORGANIC ACIDS AND LIPIDS
- 11a. Organic acids and simple esters
- 2890 Bertolini, L., Castagnetti, G.B. and Gambini, G.: (Organic acids in vinegars: HPLC identification). *Ind. Bevande*, 23 (1994) 324-326; *C.A.*, 122 (1995) 54327t.
- 2891 Bieganska, M.L., Petruczynik, A. and Glowinski, K.: Reversed phase retention behavior of β -cyclodextrin complexes of phenolic and cinnamic acids. *J. Planar Chromatogr.*, 8 (1995) 63-68.
- 2892 Buiarelli, F., Cartoni, G., Coccioli, F. and Levetsovitu, Z.: Determination of phenolic acids in wine by high-performance liquid chromatography with a microbore column. *J. Chromatogr. A*, 695 (1995) 229-235.
- 2893 Castellani, L., Flieger, M. and Sinibaldi, M.: Enantiomer separation of 2-arylpropionic acids on an ergot alkaloid-based stationary phase microbore column application. *J. Liq. Chromatogr.*, 17 (1994) 3695-3703.
- 2894 Chavis, C., Chanez, P., Vachier, I., Boosquet, J., Michel, F.B. and Godard, P.: 5-15-Dihete and lipoxins generated by neutrophils from endogenous arachidonic acid as asthma biomarkers. *Biochem. Biophys. Res. Commun.*, 207 (1995) 273-279.
- 2895 Coudray, C., Talla, M., Martin, S., Fatome, M. and Favier, A.: High-performance liquid chromatography-electrochemical determination of salicylate hydroxylation products as an *in vivo* marker of oxidative stress. *Anal. Biochem.*, 227 (1995) 101-111.
- 2896 Demini, P., Reynaud, D. and Pace-Asciac, C.R.: Extractive derivatization of the 12-lipoxygenase products, hepxilins, and related compounds into fluorescent anthryl esters for their complete high-performance liquid chromatography profiling in biological systems. *Anal. Biochem.*, 226 (1995) 252-255.
- 2897 Ding, M.-Y., Koizumi, H. and Suzuki, Y.: Comparison of three chromatographic systems for determination of organic acids in wine. *Anal. Sci.*, 11 (1995) 239-243; *C.A.*, 122 (1995) 238049k.
- 2898 Fernandez-Garcia, E. and McGregor, J.U.: Determination of organic acids during the fermentation and cold storage of yogurt. *J. Dairy Sci.*, 77 (1994) 2934-2939; *C.A.*, 122 (1995) 79319e.
- 2899 Gao, Z. and Fu, C.: (Determination of organic acids and inorganic anions by single-column ion chromatography). *Fenxi Ceshi Xuebao*, 13 (1994) 32-36; *C.A.*, 122 (1995) 122251r.
- 2900 Gensler, M. and Schmidt, H.-L.: Isolation of the main organic acids from fruit juices and nectars for carbon isotope ratio measurements. *Anal. Chim. Acta*, 299 (1994) 231-237.
- 2901 Gronert, K., Virk, S.M. and Herman, C.A.: Endogenous sulfidopeptide leukotriene synthesis and 12-lipoxygenase activity in bullfrog (*Rana catesbeiana*) erythrocytes. *Biochim. Biophys. Acta*, 1255 (1995) 311-319.
- 2902 Hesse, A., Heynck, H. and Berg, W.: Measurement of urinary oxalic acid: a comparison of four methods. *Eur. J. Clin. Chem. Clin. Biochem.*, 33 (1995) A19.
- 2903 Huang, S.-Y. and Jin, J.-D.: Purification of methyl esters of polyunsaturated fatty acids by displacement chromatography. *Bioseparation*, 4 (1994) 343-351; *C.A.*, 122 (1995) 50412u.
- 2904 Jansen, E.H.J.M., Laan, C.A. and de Fluiter, P.: Determination of phthalate-induced rat liver cytochrome P-450 IVA1 by a fluorimetric enzymatic assay and by chemiluminescence detection on Western blots. *Anal. Chim. Acta*, 303 (1995) 99-102.
- See also 2851.

- 2905 Kadkhodae, M., Endre, Z.H., Towner, R.A. and Cross, M.: Hydroxyl radical generation following ischaemia-reperfusion in cell-free perfused rat kidney. *Biochim. Biophys. Acta*, 1243 (1995) 169-174.
- 2906 Kondo, Y., Hashidoko, Y. and Mizutani, J.: An enzymatic formation of 13-oxo-trideca-9,11-dienoic acid from 13-hydroperoxylinolenic acid by a homolytic hydroperoxide lyase in elicitor-treated soybean cotyledons. *Biochim. Biophys. Acta*, 1255 (1995) 9-15.
- 2907 Kroumova, A.B. and Wagner, G.J.: Methods for separation of free, short, medium, and long chain fatty acids and for their decarboxylation. *Anal. Biochem.*, 225 (1995) 270-276.
- 2908 Lee, H.S.: Liquid chromatographic determination of benzoic acid in orange juice: interlaboratory study. *J. Assoc. Off. Anal. Chem.*, 78 (1995) 80-82.
- 2909 Lin, J.-T., McKeon, T.A. and Stafford, A.E.: Gradient reversed-phase high-performance liquid chromatography of saturated, unsaturated and oxygenated free fatty acids and their methyl esters. *J. Chromatogr. A*, 699 (1995) 85-91.
- 2910 Olawoye, T.L., Minnikin, D.E. and Christie, W.W.: HPLC separation of *tert*.-butyl dimethylsilyl (TBDMS) ether homologs of mycolic acids, using a silver impregnated normal phase column. *Biochim. Biophys. Acta*, 122 (1995) 26980g.
- 2911 Park, P.W. and Goins, R.E.: *In situ* preparation of fatty acid methyl esters for analysis of fatty acid composition in foods. *J. Food Sci.*, 59 (1994) 1262-1266; *C.A.*, 122 (1995) 131279h.
- 2912 Perrut, M., Nicoud, R.-M. and Breivik, H.: Process for chromatographic fractionation of polyunsaturated fatty acids and their derivatives. *PCT Int. Appl. WO 94 25,552 (Cl. C11C1/08)*, 10 Nov. 1994, GB Appl. 93/8,912, 29 Apr. 1993; 70 p.; *C.A.*, 122 (1995) 109324x.
- 2913 Pouliot, J.-F. and Béliveau, R.: Palmitoylation of the glucose transporter in blood-brain barrier capillaries. *Biochim. Biophys. Acta*, 1234 (1995) 191-196.
- 2914 Rao, M.S., Hidajat, K. and Ching, C.B.: Reversed-phase HPLC: the separation method for the characterization and purification of long chain polyunsaturated fatty acids - a review. *J. Chromatogr. Sci.*, 33 (1995) 9-21 - a review with 108 refs.
- 2915 Saccani, G., Gherardi, S., Trifiro, A., Bordini, C.S., Calza, M., Zoni, C. and Freddi, C.: (Determination of organic acids in fruit juices by ion chromatography). *Ind. Bevande*, 23 (1994) 423-429; *C.A.*, 122 (1995) 212369q.
- 2916 Saito, M., Ushijima, T., Sasamoto, K., Ohkura, Y. and Ueno, K.: 2-(5-Hydrazinocarbonyl-2-furyl)-5,6-dimethoxybenzothiazole as a precolumn fluorescence derivatization reagent for carboxylic acids in high-performance liquid chromatography and its application to the assay of fatty acids in human serum. *Anal. Sci.*, 11 (1995) 103-107; *C.A.*, 122 (1995) 208892n.
- 2917 See, M.M., Thompson, R., Grinberg, N., Perpall, H.J., Bicker, G. and Tway, P.: Chromatographic analysis of residual acetate in bulk drugs. *J. Liq. Chromatogr.*, 18 (1995) 137-154.
- 2918 Skotty, D.R. and Nieman, T.A.: Determination of oxalate in urine and plasma using reversed-phase ion-pair high-performance liquid chromatography with tris(2,2'-bipyridyl)ruthenium(II)-electrogenerated chemiluminescence detection. *J. Chromatogr. B*, 665 (1995) 27-36.
- 2919 Sushchik, N.N., Gladyshev, M.I., Kalachova, G.S. and Guseynova, V.E.: Rapid assay of fatty acid composition using a portable high-performance liquid chromatograph for monitoring aquatic ecosystems. *J. Chromatogr. A*, 695 (1995) 223-228.
- 2920 Vichot, R. and Furton, K.G.: Variables influencing the direct determination of haloacetic acids in water by reversed-phase ion-pair chromatography with indirect UV detection. *J. Liq. Chromatogr.*, 17 (1994) 4405-4429.
- 2921 Walker, T.A.: A liquid chromatographic assay for citric acid in over-the-counter carbamide peroxide products using indirect UV detection. *J. Pharm. Biomed. Anal.*, 13 (1995) 171-176.
- 2922 Yu, W. and Powell, W.S.: Analysis of leukotrienes, lipoxins, and monooxygenated metabolites of arachidonic acid by reversed-phase high-pressure liquid chromatography. *Anal. Biochem.*, 226 (1995) 241-251.

For additional information see C.A.:
122 (1995) 50452g, 208914w.

See also 2625, 3033, 3268, 3361, 3410, 3565, 3616, 3802, 3810, 3816, 3888.

11b. Prostaglandins

- 2923 Capdevila, J.H., Morrow, J.D., Belosludtsev, Y.Y., Beauchamp, D.R., DuBois, R.N. and Falck, J.R.: The catalytic outcomes of the constitutive and the mitogen inducible isoforms of prostaglandin H₂ synthase are markedly affected by glutathione and glutathione peroxidase(s). *Biochemistry*, 34 (1995) 3325-3337.
- 2924 Gatti, R., Gotti, R., Cavrini, V., Soli, M., Bertaccini, A. and Carparelli, F.: Stability study of prostaglandin E₁ (PGE₁) in physiological solutions by liquid chromatography (HPLC). *Int. J. Pharm.*, 115 (1995) 113-117; *C.A.*, 122 (1995) 142298q.
- 2925 Yamane, M., Shimizu, S., Abe, A., Sugiura, H., Miyaoka, M. and Saitoh, T.: High-performance liquid chromatography-thermospray mass spectrometry of ω -carboxyleukotriene B₄ and ω -hydroxyleukotriene B₄ from an incubation mixture of human colonic well-differentiated adenocarcinoma homogenate. *J. Chromatogr. B*, 666 (1995) 197-202.

See also 2922.

11c. Lipids and their constituents

- 2926 Abidi, S.L., Mounts, T.L. and Rennick, K.A.: Separations of major soybean phospholipids on β -cyclodextrin-bonded silica. *J. Liq. Chromatogr.*, 17 (1994) 3705-3725.
- 2927 Adlof, R.O.: Analysis of triacylglycerol positional isomers by silver ion high performance liquid chromatography. *J. High Resolut. Chromatogr.*, 18 (1995) 105-107.
- 2928 Adlof, R.O.: The preparative separation of lipids by silver resin chromatography. *Process Technol. Proc.*, 11 (1994) 77-781; *C.A.*, 122 (1995) 189497u.
- 2929 Akasaka, K., Ohata, A., Ohnui, H. and Meguro, H.: Automatic determination of hydroperoxides of phosphatidylcholine and phosphatidylethanolamine in human plasma. *J. Chromatogr. B*, 665 (1995) 37-43.
- 2930 Alter, C.A. and Wolf, B.A.: Identification of phosphatidylinositol 3,4,5-trisphosphate in pancreatic islets and insulin secreting β -cells. *Biochem. Biophys. Res. Commun.*, 208 (1995) 190-197.

- 2931 Bernhard, W., Postle, A.D., Linck, M. and Sewing, K.-F.: Composition of phospholipid classes and phosphatidylcholine molecular species of gastric mucosa and mucus. *Biochim. Biophys. Acta*, 1255 (1995) 99-104.
- 2932 Christie, W.W. and Urwin, R.A.: Separation of lipid classes from plant tissues by high performance liquid chromatography on chemically bonded stationary phases. *J. High Resolut. Chromatogr.*, 18 (1995) 97-100.
- 2933 Eaton, S., Shmueli, E., Al-Mardini, H., Bartlett, K. and Record, C.O.: An HPLC assay for sn-1,2-diacylglycerol. *Clin. Chim. Acta*, 234 (1995) 71-78.
- 2934 Firestone, D.: Fats and oils. *J. Assoc. Off. Anal. Chem.*, 78 (1995) 150-155 - a review with 26 refs.
- 2935 Fukami, M.H. and Holmsen, H.: Diacylglycerol elevations in control platelets are unaccompanied by pleckstrin phosphorylation. Implications for the role of diacylglycerol in platelet activation. *Eur. J. Biochem.*, 228 (1995) 579-586.
- 2936 Grob, K. and Mariani, C.: LC-GC method for the determination of adulterated edible oils and fats. *Spec. Publ. - R. Soc. Chem.*, 160(Developments in the Analysis of Lipids) (1994) 73-81; *C.A.*, 122 (1995) 131220g - a review with 13 refs.
- 2937 Hayashi, K., Ohnishi, T., Tabata, M., Yoshita, K., Nakagawa, H., Kawano, S., Nagase, H., Okazawa, T., Yoshida, M. et al.: Studies on the analysis of serum alkenal and serum lipid peroxides in migrant workers exposed to dust. *Hokuriku Kosho Eisei Gakkaishi*, 20 (1993) 29-36; *C.A.*, 122 (1995) 48562z.
- 2938 Heron, S., Lesellier, E. and Tchaplal, A.: Analysis of triacylglycerols of borage oil by RPLC identification by coinjection. *J. Liq. Chromatogr.*, 18 (1995) 599-611.
- 2939 Holst, O., Bock, K., Brade, L. and Brade, H.: The structure of oligosaccharide bisphosphates isolated from the lipopolysaccharide of a recombinant *Escherichia coli* strain expressing the gene *gseA* [3-deoxy-D-manno-octulopyranosonic acid (Kdo) transferase I] of *Chlamydia psittaci* 6B. *Eur. J. Biochem.*, 229 (1995) 194-200.
- 2940 Itonori, S., Shiray, T., Kiso, Y., Oha, Y., Shiota, K. and Ogawa, T.: Glycosphingolipid composition of rat placenta: changes associated with stage of pregnancy. *Biochem. J.*, 307 (1995) 399-405.
- 2941 Jalink, K., Hengeveld, T., Mulder, S., Postma, F.R., Simon, M.-F., Chap, H., van der Marel, G.A., van Boorn, J.H., van Blitterswijk, W.J. and Moolenaar, W.H.: Lysophosphatidic acid-induced Ca^{2+} mobilization in human A431 cells: structure-activity analysis. *Biochem. J.*, 307 (1995) 609-616.
- 2942 Justin, A.-M., Hmyene, A., Kader, J.-C. and Mazliak, P.: Compared selectivities of the phosphatidylinositol-synthase from maize coleoptiles either in microsomal membranes or after solubilization. *Biochim. Biophys. Acta*, 1255 (1995) 161-166.
- 2943 Krainev, A.G., Ferrington, D.A., Williams, T.D., Squier, T.C. and Bigelow, D.J.: Adaptive changes in lipid composition of skeletal sarcoplasmic reticulum membranes associated with aging. *Biochim. Biophys. Acta*, 1235 (1995) 406-418.
- 2944 Leopold, K.: (Hydrophobic interaction chromatography (HIC) of lipid macroamphiphiles). *GIT Fachz. Lab.*, 38 (1994) 455-460; *C.A.*, 122 (1995) 182303u.
- 2945 Luciani, S., Antolini, M., Bova, S., Cargnelli, G., Cusinato, F., Debetto, P., Trevisi, L. and Varotto, R.: Inhibition of cardiac sarcolemmal sodium-calcium exchanger by glycerophosphoinositol 4-phosphate and glycerophosphoinositol 4,5-bisphosphate. *Biochem. Biophys. Res. Commun.*, 206 (1995) 674-680.
- 2946 Ma, Y.-C. and Kim, H.-Y.: Development of the on-line high-performance liquid chromatography/thermospray mass spectrometry method for the analysis of phospholipid molecular species in rat brain. *Anal. Biochem.*, 226 (1995) 293-301.
- 2947 Meyuhas, D. and Lichtenberg, D.: The effect of albumin on the state of aggregation and phase transformations in phosphatidylcholine-sodium cholate mixtures. *Biochim. Biophys. Acta*, 1234 (1995) 203-213.
- 2948 Nikolova-Damyanova, B., Christie, W.W. and Herslöf, B.G.: Retention properties of triacylglycerols on silver ion high-performance liquid chromatography. *J. Chromatogr. A*, 694 (1995) 375-380.
- 2949 Proudfoot, L., Schneider, P., Ferguson, M.A.J. and McConville, M.J.: Biosynthesis of the glycolipid anchor of lipophosphoglycan and the structurally related glycoinositolphospholipids from *Leishmania major*. *Biochem. J.*, 308 (1995) 45-55.
- 2950 Quarfordt, S.H., Landis, B.A. and Oswald, B.S.: The influence of cholesteryl ester content on hepatocyte triolein emulsion uptake. *Biochim. Biophys. Acta*, 1255 (1995) 82-86.
- 2951 Redden, P.R., Huang, Y.-S., Lin, X. and Horrobin, D.F.: Separation and quantification of the triacylglycerols in evening primrose and borage oils by reversed-phase high-performance liquid chromatography. *J. Chromatogr. A*, 694 (1995) 381-389.
- 2952 Spanos, G.A., Schwartz, S.J., van Breemen, R.B. and Huang, C.-H.: High-performance liquid chromatography with light-scattering detection and desorption chemical-ionization tandem mass spectrometry of milk fat triacylglycerols. *Lipids*, 30 (1995) 85-90; *C.A.*, 122 (1995) 131316t.
- 2953 Xu, G. and Chang, L.: (Progress in high performance liquid chromatography-precolum derivatization for gangliosides). *Sepu*, 13 (1995) 26-29; *C.A.*, 122 (1995) 234472v - a review with 5 refs.

For additional information see C.A.:

122 (1995) 27201r, 155416b, 212490x.

See also 2791, 2853, 2855, 2909, 2971, 3144, 3477.

11d. Lipoproteins and their constituents

- 2954 Anchisi, C., Batetta, B., Sanna, F., Fadda, A.M., Maccioni, A.M. and Dessi, S.: HDL subfractions as altered in cancer patients. *J. Pharm. Biomed. Anal.*, 13 (1995) 65-71.
- 2955 Atwal, A.S., Blais, L. and Jenkins, K.J.: Direct determination of apolipoproteins in plasma by high performance liquid chromatography. *J. Liq. Chromatogr.*, 18 (1995) 579-587.
- 2956 Nagayoshi, A., Matsuki, N., Saito, H., Tsukamoto, K., Kaneko, K., Wakashima, M., Kinoshita, M., Yamanaka, M. and Teramoto, T.: Defect in assembly process of very-low-density lipoprotein in suncus liver: an animal model of fatty liver. *J. Biochem. (Tokyo)*, 117 (1995) 787-793.
- 2957 Panteghini, M., Bonora, R. and Pagani, F.: Determination of glycated apolipoprotein B in serum by a combination of affinity chromatography and immunonephelometry. *Ann. Clin. Biochem.*, 31 (1994) 544-549; *C.A.*, 122 (1995) 75915y.
- 2958 Rinninger, F., Brundert, M., Jäckle, S., Kaiser, T. and Greten, H.: Selective uptake of low-density lipoprotein-associated cholesteryl esters by human fibroblasts, human HepG2 hepatoma cells and J774 macrophages in culture. *Biochim. Biophys. Acta*, 1255 (1995) 141-153.

2959 Tricerri, A., Garda, H.A. and Brenner, R.R.: A rapid and efficient procedure for the purification of human apolipoprotein A-I using gel filtration HPLC. *Int. J. Bio-Chromatogr.*, 1 (1994) 159-166; *C.A.*, 122 (1995) 75707g.

12. ORGANIC PEROXIDES

2960 Zhang, J.-R., Cazars, A.R., Lutzke, B.S. and Hall, E.D.: HPLC-chemiluminescence and thermospray LC/MS study of hydroperoxides generated from phosphatidylcholine. *Free Radical Biol. Med.*, 18 (1995) 1-10; *C.A.*, 122 (1995) 75987y.

See also 2641, 2929, 2971.

13. STEROIDS

13a. General techniques

See 2685.

13b. Pregnane and androstane derivatives

2961 Bidart, M. and Lesgards, G.: Direct injection analysis of 6 β -hydroxycortisol and cortisol in urine by HPLC-UV with on-line ISRP precolumn. *J. Liq. Chromatogr.*, 18 (1995) 725-738.

2962 Dwyer, J.L., Chapman, A.E. and Liu, X.: Analysis of steroids by combined chromatography-infrared spectroscopy. *LC-GC*, 13 (1995) 240-250; *C.A.*, 122 (1995) 197098e.

2963 Heilmann, P., Vielemeyer, O., Lessing, U. and Schöneshöfer, M.: Fully automated determination of testosterone levels in male urine using the ALCA™ system. *Eur. J. Clin. Chem. Clin. Biochem.*, 33 (1995) A20.

2964 Loftsson, T., Gudmundsdottir, T.K., Fridriksdottir, H., Sigurdardottir, A.M., Thorkelsson, J., Gudmundsson, G. and Hjaltason, B.: Fatty acids from cod-liver oil as skin penetration enhancers. *Pharmazie*, 50 (1995) 188-190.

2965 Santos-Montes, A., Gasco-Lopez, A.I. and Izquierdo-Hornillos, R.: Simultaneous determination of dexamethasone and betamethasone in pharmaceuticals by reversed-phase HPLC. *Chromatographia*, 39 (1994) 539-542.

2966 Shimada, K., Satoh, Y. and Nishimura, S.: Studies on neurosteroids. I. Retention behavior of derivatized 17-oxosteroids using high-performance liquid chromatography. *J. Liq. Chromatogr.*, 18 (1995) 713-723.

2967 Torres Cartas, S., García Alvarez-Coque, M.C. and Villanueva Camañas, R.M.: Determination of anabolic steroids in pharmaceuticals by liquid chromatography with a microemulsion of sodium dodecyl sulfate and pentanol as mobile phase. *Anal. Chim. Acta*, 302 (1995) 163-172.

2968 Wienkers, L.C., Stenwyk, R.C., Myszak, S.A. and Pearson, P.G.: *In vitro* metabolism of tirilazad mesylate in male and female rats. Contribution of cytochrome P4502C11 and 5 α -reductase. *Drug Metab. Disp.*, 23 (1995) 383-392.

For additional information see *C.A.*:
122 (1995) 115142c.

See also 3648, 3754, 3819.

13c. Estrogens

2969 Miller, R.B., Delker, G., Chen, C. and Sherwood, C.H.: Development of a HPLC method for the determination of 17 β -estradiol-3-phosphate in pharmaceutical preparations. *J. Liq. Chromatogr.*, 18 (1995) 127-136.

2970 Sheikh, S.U. and Touchstone, J.C.: Determination of free estriol in amniotic fluid by high performance liquid chromatography. *J. Liq. Chromatogr.*, 17 (1994) 3813-3820.

See also 2964.

13d. Sterols

2971 Bartolomeazzi, R., Pizzale, L., Vichi, S. and Lercker, G.: Analysis of isomers of cholesteryl acetate hydroperoxides by HPLC and GC-ITDMS. *Chromatographia*, 39 (1994) 577-580.

2972 Dinchuk, J., Hart, J., Gonzalez, G., Karmann, G., Schmidt, D. and Wirak, D.O.: Remodelling of lipoproteins in transgenic mice expressing human cholesteryl ester transfer protein. *Biochim. Biophys. Acta*, 1255 (1995) 301-310.

2973 Hamill, T.W. and Soliman, A.M.: Determination of cholesterol by *p*-nitrobenzoate derivatization and liquid chromatography. *J. Assoc. Off. Anal. Chem.*, 77 (1994) 1190-1196.

2974 Nielsen, J.H., Olsen, C.E., Duedahl, C. and Skibsted, L.H.: Isolation and quantification of cholesterol oxides in dairy products by selected ion monitoring mass spectrometry. *J. Dairy Res.*, 62 (1995) 101-113; *C.A.*, 122 (1995) 131321r.

13e. Bile acids and alcohols

2975 Iida, T., Yamaguchi, Y., Maruyama, J., Nishio, M., Goto, J. and Nambara, T.: High performance liquid chromatographic separation of nonamidated and glycine- and taurine-amidated bile acid 3-glucosides. *J. Liq. Chromatogr.*, 18 (1995) 701-711.

2976 Roda, A., Gioacchini, A.M., Cerrè, C. and Baraldini, M.: High-performance liquid chromatographic-electrospray mass spectrometric analysis of bile acids in biological fluids. *J. Chromatogr. B*, 665 (1995) 281-294.

2977 Scalia, S., Tirendi, S., Pazzi, P. and Bousquet, E.: Assay of free bile acids in pharmaceutical preparations by HPLC with electrochemical detection. *Int. J. Pharm.*, 115 (1995) 249-253; *C.A.*, 122 (1995) 170319r.

13g. Other steroids

2978 Gamoh, K., Yamaguchi, I. and Takatsuto, S.: Rapid and selective sample preparation for the chromatographic determination of brassinosteroids from plant material using solid-phase extraction method. *Anal. Sci.*, 10 (1994) 913-917; *C.A.*, 122 (1995) 50408x.

14. STEROID GLYCOSIDES AND SAPONINS

- 2979 Fujii, Y., Ikeda, Y. and Yamazaki, M.: High-performance liquid chromatographic determination of lanatosides in *Digitalis lutea* and *Digitalis ambigua* leaves. *J. Liq. Chromatogr.*, 17 (1994) 4451-4461.
- 2980 Kockar, O.M., Kara, M., Kara, S., Bozan, B. and Baser, K.H.: Quantitative determination of escin. A comparative study of HPLC and TLC densitometry. *Fitoterapia*, 65 (1994) 439-443; C.A., 122 (1995) 170314k.
- 2981 Muhr, P., Michelitsch, A., Likussar, W. and Schubert-Zsilavec, M.: Ein einfacher Zugang zu reinem Digitonin und Digalolin. *Pharmazie*, 50 (1995) 295-296.

For additional information see C.A.:
122 (1995) 150713m.

15. TERPENES AND OTHER VOLATILE AROMATIC COMPOUNDS

15a. Terpenes

- 2982 Cai, J.J., Tao, X. and Lipsky, P.E.: High performance liquid chromatographic determination of triptolide and triptolide in an ethyl acetate extract of *Tripterygium wilfordii* Hook F. *J. Liq. Chromatogr.*, 17 (1994) 4479-4487.
- 2983 Cheung, H.T.A., Fu, S.-L. and Smal, M.A.: Inhibition of platelet aggregation by diterpene acids from *Pinus massoniana* resin. *Arzneim.-Forsch.*, 44 (1994) 17-25.
- 2984 Green, M.D., Mount, D.L., Todd, G.D. and Capomacchia, A.C.: Chemiluminescent detection of artemisinin. Novel endoperoxide analysis using luminol without hydrogen peroxide. *J. Chromatogr. A*, 695 (1995) 237-242.
- 2985 Mondello, L., Dugo, P., Bartle, K.D., Frere, B. and Dugo, G.: On-line high performance liquid chromatography coupled with high resolution gas chromatography and mass spectrometry (HPLC-HRGC-MS) for the analysis of complex mixtures containing high volatile compounds. *Chromatographia*, 39 (1994) 529-538.
- 2986 Nowak, G.: Isolation and chromatography of 15-deoxyrepen and 25 other sesquiterpene lactones from *Centaurea bella*. *Acta Soc. Bot. Pol.*, 62 (1993) 33-36; C.A., 122 (1995) 89205d.
- 2987 O'Mathuna, D.P. and Doskotch, R.W.: New labdane diterpene glycosides from *Amphiachyris amoena*. *J. Natural Prod.*, 58 (1995) 82-92.
- 2988 Rodriguez, A.D., Cobar, O.M. and Martinez, N.: Isolation and structures of sixteen new asbestinin diterpenes from the caribbean gorgonian *Briareum asbestinum*. *J. Natural Prod.*, 57 (1994) 1638-1655.

15b. Essential oils

- 2989 Zekovic, Z., Pekic, B., Ledojevic, Z. and Petrovic, L.: Chromatography in our investigations of camomile (*Metricaria chamomilla* L.). *Chromatographia*, 39 (1994) 587-590.

See also 2985.

15c. Bitter substances

- 2990 Buckee, G.K.: Determination of iso-alpha-, alpha-, and beta-acids in hop and isomerized hop extracts. A collaborative trial. *Monatsschr. Brauwiss.*, 47 (1994) 284-286; C.A., 122 (1995) 185640n.

16. NITRO AND NITROSO COMPOUNDS

- 2991 Fu, C.-G. and Xu, H.-D.: High-performance liquid chromatography with post-column chemiluminescence detection for simultaneous determination of trace N-nitrosamines and corresponding secondary amines in groundwater. *Analyst (Cambridge)*, 120 (1995) 1147-1151.

See also 2653, 2796.

17. AMINES, AMIDES AND RELATED NITROGEN COMPOUNDS

17a. Amines and polyamines

- 2992 Cheng, F.-C. and Kuo, J.-S.: High-performance liquid chromatographic analysis with electrochemical detection of biogenic amines using microbore columns. *J. Chromatogr. B*, 665 (1995) 1-13 - a review with 141 refs.
- 2993 Dobberpuhl, D.A. and Johnson, D.C.: Pulsed electrochemical detection of alkanolamines separated by multimodal high-performance liquid chromatography. *J. Chromatogr. A*, 694 (1995) 391-398.
- 2994 Gennaro, M.C., Giacosa, D. and Abrigo, C.: The role of pH of the mobile-phase in ion-interaction RP-HPLC. *J. Liq. Chromatogr.*, 17 (1994) 4365-4380.
- 2995 Hayashi, T., Amino, M., Uchida, G. and Sato, M.: High-performance liquid chromatographic determination of primary aromatic amines in urine after derivatization to an azo dye with 2-aminoanthracene. *J. Chromatogr. B*, 665 (1995) 209-212.
- 2996 Hermann, K., Frank, G. and Ring, J.: High performance liquid chromatography for the separation of histamine, its precursor, and metabolites: application to biological samples. *J. Liq. Chromatogr.*, 18 (1995) 189-204.
- 2997 Hiramatsu, K., Sugimoto, M., Kamei, S., Hoshino, M., Kinoshita, K., Iwasaki, K. and Kawakita, M.: Determination of amounts of polyamines excreted in urine: demonstration of N¹,N⁸-diacetylspermidine and N¹,N¹²-diacetylspermine as components commonly occurring in normal human urine. *J. Biochem. (Tokyo)*, 117 (1995) 107-112.
- 2998 Huhn, G., Mattusch, J. and Schulz, H.: Determination of polyamines in biological materials by HPLC with 9-fluorenylmethyl chloroformate precolumn derivatization. *Fresenius J. Anal. Chem.*, 351 (1995) 563-566.
- 2999 Iniguez Crespo, M and Vazquez Lasa, B.: Determination of biogenic amines and other amines in wine by an optimized HPLC method with polarity gradient elution. *Am. J. Enol. Vitic.*, 45 (1994) 460-463; C.A., 122 (1995) 158852b.

- 3000 Lu, C.-S. and Huang, S.-D.: Trace determination of aromatic amines or phenolic compounds in dyestuffs by high-performance liquid chromatography with on-line preconcentration. *J. Chromatogr. A*, 696 (1995) 201-208.
- 3001 Marcé, M., Brown, D.S., Capell, T., Figueras, X. and Tiburcio, A.F.: Rapid high-performance liquid chromatographic method for the quantitation of polyamines as their dansyl derivatives: application to plant and animal tissues. *J. Chromatogr. B*, 666 (1995) 329-335.
- 3002 Qi, X. and Baldwin, R.P.: Liquid chromatography/electrochemical detection of hydroxylamines by oxidation at a cobalt phthalocyanine chemically modified electrode. *Electroanalysis (N.Y.)*, 6 (1994) 353-360; *C.A.*, 122 (1995) 150578w.
- 3003 Skarping, G., Dalene, M. and Brunmark, P.: Liquid chromatography and mass spectrometry determination of aromatic amines in hydrolysed urine from workers exposed to thermal degradation products of polyurethane. *Chromatographia*, 39 (1994) 619-623.
- 3004 Tang, Q., Zhuang, L., Xu, X., Shi, Y. and Li, S.: (Measurement of urinary polyamines by high performance liquid chromatography with electrochemical detection). *Sepu*, 12 (1994) 431-432; *C.A.*, 122 (1995) 50423y.
- 3005 Tomaselli, M., Cozzolino, A., Calemme, I. and Stasio, A.: (Determination of benzidine and other aromatic amines in dyes and dyed leathers). *Cuoio, Pelli, Mater. Concianti*, 69 (1993) 253-263; *C.A.*, 122 (1995) 190912g.
- 3006 Wintersteiger, R., Barary, M.H., El-Yazbi, F.A., Sabry, S.M. and Wahbi, A.-A.M.: Reversed phase-liquid chromatography of primary and secondary aliphatic amines after derivatization combined with electrochemical detection. *Anal. Chim. Acta*, 306 (1995) 273-283.
- 3007 Yamazaki, S., Ozaki, K., Saito, K. and Tanimura, T.: Enantiomeric separation of β -amino alcohols with a primary or tertiary amine moiety by reversed phase liquid chromatography with a chiral mobile phase containing copper(II) and (R)-mandelic acid. *J. High Resolut. Chromatogr.*, 18 (1995) 68-70.
- See also 2566, 2608, 2636, 2653, 3037, 3054, 3616.
- 17b. *Catecholamines and their metabolites*
- 3008 Coward, S., Boa, F.G. and Sherwood, R.A.: Sulfasalazine interference with HPLC assay of 5-hydroxyindole-3-acetic acid. *Clin. Chem. (Washington)*, 41 (1995) 765-766.
- 3009 Dryhurst, G.: Experimental techniques to measure low molecular weight neurotransmitters and their precursors and metabolites. *NATO ASI Ser., Ser. A*, 267(Bioelectrochemistry IV) (1994) 171-193; *C.A.*, 122 (1995) 151453p - a review with 41 refs.
- 3010 Elkihel, L., Grosclaude, J.M., Nembo, B., Jacquot, C., Gayral, P., Goudey-Perriere, F., Letourneux, Y. and Barreteau, H.: Synthesis and determination of N-acetyloctopamine by HPLC with electrochemical detection. Bioassay in *Nippostrongylus brasiliensis* (nematoda). *J. Liq. Chromatogr.*, 17 (1994) 3421-3435.
- 3011 Feier, U. and Goetsch, P.H.: Interlaboratory studies on precision characteristics of analytical methods. Determination of biogenic amines in fish and fish products - HPLC method. *Arch. Lebensmittelhyg.*, 44 (1993) 134-135; *C.A.*, 122 (1995) 131262x.
- 3012 Kirschbaum, J., Luckas, B. and Beinert, W.D.: (Application of automatic precolumn derivatization with 9-fluorenylmethyl chloroformate (FMOC-C1) for the determination of biogenic amines and amino acids in food by HPLC). *Dtsch. Lebensm.-Rundsch.*, 90 (1994) 224-228; *C.A.*, 122 (1995) 29955p.
- 3013 Kirschbaum, J., Luckas, B. and Beinert, W.D.: HPLC analysis of biogenic amines and amino acids in food after automatic precolumn derivatization with 9-fluorenylmethyl chloroformate. *Am. Lab. (Shelton)*, 26 (1994) 28C-28D, 28F, 28H; *C.A.*, 122 (1995) 104168q.
- 3014 Murai, S., Saito, H., Masuda, Y. and Itoh, T.: An improved method for the rapid assay of brain monoaminergic makers using HPLC coupled with coulometric detection. *Biog. Amines*, 10 (1994) 499-508; *C.A.*, 122 (1995) 26991m.
- 3015 Radjaipour, M. and Raster, H.: Separation of urine-catecholamines and their abundant metabolites by uniform HPLC-conditions and fluorimetric detection. *Eur. J. Clin. Chem. Clin. Biochem.*, 33 (1995) A28.
- 3016 Sanchis Mallols, J.M., Torres Lapasio, J.R., Villanueva Camanas, R.M. and Ramis-Ramos, G.: Determination of catecholamines in urine by micellar liquid chromatography with coulometric detection. *Chromatographia*, 39 (1994) 591-596.
- For additional information see *C.A.*:
122 (1995) 204407k.
- See also 2992, 2999.
- 17c. *Urea and guanidine derivatives*
- See 2566.
- 17d. *Other amine derivatives and amides (excl. peptides)*
- See 2643, 2685, 2994.
18. AMINO ACIDS AND PEPTIDES; CHEMICAL STRUCTURE OF PROTEINS
- 18a. *Amino acids and their derivatives*
- 3017 Bartók, T., Szalai, G., Lőrincz, Z., Börcsök, G. and Sági, F.: High-speed RP-HPLC/FL analysis of amino acids after automated two-step derivatization with *o*-phthaldialdehyde/3-mercaptopropionic acid and 9-fluorenylmethyl chloroformate. *J. Liq. Chromatogr.*, 17 (1994) 4391-4403.
- 3018 Bianchi, V. and Mazza, L.: Rapid reversed-phase high-performance liquid chromatographic method with double derivatization for the assay of urinary hydroxyproline. *J. Chromatogr. B*, 665 (1995) 295-302.
- 3019 Campos, K.L., Giovanelli, J. and Kaufman, S.: Characteristics of the nitric oxide synthase-catalyzed conversion of arginine to N-hydroxyarginine, the first oxygenation step in the enzymic synthesis of nitric oxide. *J. Biol. Chem.*, 270 (1995) 1721-1728.
- 3020 Carta, G. and Dinerman, A.A.: Displacement chromatography of amino acids: effects of selectivity reversal. *AIChE J.*, 40 (1994) 1618-1628; *C.A.*, 122 (1995) 31898r.

- 3021 Castelain, S., Kamel, S., Picard, C., Desmet, G., Seberty, J.L. and Brazier, M.: A simple and automated HPLC method for determination of total hydroxyproline in urine. Comparison with excretion of pyridinolines. *Clin. Chim. Acta*, 235 (1995) 81-90.
- 3022 Chaves das Neves, H.J.: Amino acids. The new methods in chromatography: GLC and SFC. *Acquis. Recent. Chromatogr. Vin Cours Eur. Form. Contin.*, 1992, (1993) 67-79; *C.A.*, 122 (1995) 79290p - a review with 6 refs.
- 3023 Collar, C. and Martínez-Anaya, M.A.: Influence of the microbial starter and the breadmaking step on the free amino acid profiles of wheat sours, doughs, and breads by reversed-phase high-performance liquid chromatography. *J. Liq. Chromatogr.*, 17 (1994) 3437-3460.
- 3024 Daniels, D.H., Joe, F.L., Jr. and Diachenko, G.W.: Determination of free glutamic acid in a variety of foods by high-performance liquid chromatography. *Food Addit. Contam.*, 12 (1995) 21-29; *C.A.*, 122 (1995) 238044e.
- 3025 De Witte, P.A., Cuveele, J.F., Merlevede, W.J. and Vandenheede, J.R.: Analysis of phosphorylhydroxyamino acids present in hydrolyzed cell extracts using dansyl derivatization. *Anal. Biochem.*, 226 (1995) 1-9.
- 3026 Egorova, T.A., Eremin, S.V., Mitsner, B.I., Zvonkova, E.N. and Shvets, V.I.: Isolation of individual amino acids from various microbiological sources using reversed-phase high-performance liquid chromatography. *J. Chromatogr. B*, 665 (1995) 53-62.
- 3027 Evrovski, J., Callaghan, M. and Cole, D.E.C.: Determination of homocysteine by HPLC with pulsed integrated amperometry. *Clin. Chem. (Washington)*, 41 (1995) 757-759.
- 3028 Fujimura, S., Yamamoto, A., Kobayashi, K. and Ishibashi, T.: Comparison of performance of amino acid analysis with HPLC equipped with sodium- and lithium-type columns. *Niigata Daigaku Nogakubu Kenkyu Hokoku*, 46 (1994) 71-77; *C.A.*, 122 (1995) 234532q.
- 3029 Gasinska, E.: (Affinity chromatography on special polymer beds loaded with ions of transition metals and its usability in the process of separating amino acids in protein hydrolyzates). *Zesz. Nauk. - Politech. Lodz., Technol. Chem. Spozyw.*, 657 (1994) 83-91; *C.A.*, 122 (1995) 158884p.
- 3030 Gerrits, M.I., Thijssen, J.H.H. and van Rijn, H.J.M.: Determination of pyridinoline and deoxypyridinoline in urine, with special attention to retaining their stability. *Clin. Chem. (Washington)*, 41 (1995) 571-574.
- 3031 Inou, H., Moritani, K., Date, Y., Kohashi, K. and Tsuruta, Y.: Determination of free hydroxyproline and proline in human serum by high-performance liquid chromatography using 4-(5,6-dimethoxy-2-phthalimidinyl)phenyl-sulfonyl chloride as pre-column fluorescent labelling reagent. *Analyst (Cambridge)*, 120 (1995) 1141-1145.
- 3032 Kera, Y., Aoyama, H., Matsumura, H., Hasegawa, A., Nagasaki, H. and Yamada, R.-h.: Presence of free D-glutamate and D-aspartate in rat tissues. *Biochim. Biophys. Acta*, 1243 (1995) 282-286.
- 3033 Knox, J.H. and Wan, W.-H.: Chiral chromatography of amino- and hydroxy-acids on surface modified porous graphite. *Chromatographia*, 40 (1995) 9-14.
- 3034 Lau, O.-W. and Mok, C.-S.: Indirect conductometric detection of amino acids after liquid chromatographic separation. *Anal. Chim. Acta*, 300 (1995) 183-191.
- 3035 Lau, O.-W. and Mok, C.-S.: Indirect conductometric detection of amino acids after liquid chromatographic separation. Part II. Determination of monosodium glutamate in foods. *Anal. Chim. Acta*, 302 (1995) 45-52.
- 3036 Ma, Y., Ito, Y., Torok, D.S. and Ziffer, H.: Separation of the 2- and 6-nitro-3-acetamido-4-chlorobenzoic acid precursors of a potent hydroxy-anthranilic acid oxygenase inhibitor by pH-zone-refining-counter-current chromatography. *J. Liq. Chromatogr.*, 17 (1994) 3507-3517.
- 3037 McAllister, T.A., Samuels, S.E., Sedgwick, G.W., Fenton, T.W. and Thompson, J.R.: High-performance liquid chromatographic analysis of β -phenylethylamine for the estimation of *in vivo* protein synthesis. *J. Chromatogr. B*, 666 (1995) 336-341.
- 3038 Michalsky, M., Stepan, J.J., Wilczek, H., Formankova, J. and Moro, L.: Galactosyl hydroxylysine in assessment of Paget's bone disease. *Clin. Chim. Acta*, 234 (1995) 101-108.
- 3039 Miles, D.R.: Amino acid and protein chromatography using immobilized platinum(II) ions. Avail. *Univ. Microfilms Int.*, Order No. DA9424133, 1994, 225 p.; *C.A.*, 122 (1995) 150568t.
- 3040 Misra, I., Charlier, H.A., Jr. and Mizioro, H.M.: Avian cytosolic 3-hydroxy-3-methylglutaryl-CoA synthase: evaluation of the role of cysteines in reaction chemistry. *Biochim. Biophys. Acta*, 1247 (1995) 253-259.
- 3041 Mollé, D., Léonil, J. and Bouhallab, S.: Separation of two-proline-containing peptides by capillary electrophoresis. *Anal. Biochem.*, 225 (1995) 161-162.
- 3042 Mornstad, H., Pfeiffer, H. and Teivens, A.: Estimation of dental age using HPLC-technique to determine the degree of aspartic acid racemization. *J. Forensic Sci.*, 39 (1994) 1425-1431; *C.A.*, 122 (1995) 48621t.
- 3043 Niwa, T., Shiobara, K., Hamada, T., Miyazaki, T., Tsukushi, S., Uema, K. and Tsuzuki, T.: Serum pyridinolines as specific markers of bone resorption in hemodialyzed patients. *Clin. Chim. Acta*, 235 (1995) 33-40.
- 3044 Peaper, D.R. and Jones, H.L.: Determination of S-methylmethionine in malt by alkaline hydrolysis and headspace gas chromatography - a collaborative trial. *J. Inst. Brew.*, 100 (1994) 397-398; *C.A.*, 122 (1995) 158849f.
- 3045 Rawal, N., Rajpurohit, R., Lischwe, M.A., Williams, K.R., Paik, W.K. and Kim, S.: Structural specificity of substrate for S-adenosylmethionine:protein arginine N-methyltransferases. *Biochim. Biophys. Acta*, 1248 (1995) 11-18.
- 3046 Sarvar, G. and Botting, H.G.: Reversed-phase liquid chromatographic determination of hypoglycin A (HG-1) in canned ackee fruit samples. *J. Assoc. Off. Anal. Chem.*, 77 (1994) 1175-1179.
- 3047 Sillanpaa, M., Kokkonen, R. and Sihvonen, M.-L.: Determination of EDTA and DTPA as their Fe(III) complexes in pulp and paper mill process and waste waters by liquid chromatography. *Anal. Chim. Acta*, 303 (1995) 187-192.
- 3048 Sottrup-Jensen, L.: A low-pH reverse-phase high-performance liquid chromatography system for analysis of the phenylthiohydantoins of S-carboxymethylcysteine and S-carboxyamidomethylcysteine. *Anal. Biochem.*, 225 (1995) 187-188.
- 3049 Sprongl, L.: Determination of carnitine by high performance liquid chromatography after derivatization with anthryldiazomethan. *Eur. J. Clin. Chem. Clin. Biochem.*, 33 (1995) A17.

- 3050 Takeuchi, T. and Miwa, T.: (On-column fluorimetric detection of dansylamine acids with bovine serum albumin in liquid chromatography). *Kuromatogurafi*, 15 (1994) 236-237; *C.A.*, 122 (1995) 127788f.
- 3051 Te Poele-Pothoff, M.T.W.B., van den Berg, M., Franken, D.G., Boers, G.H.J., Jakobs, C., de Kroon, I.F.I., Eskes, T.K.A.B., Trijbels, J.M.F. and Blom, H.J.: Three different methods for the determination of total homocysteine in plasma. *Ann. Clin. Biochem.*, 32 (1995) 218-220; *C.A.*, 122 (1995) 234526r.
- 3052 Uchikura, K. and Kirisawa, M.: (Ru(bpy)₃²⁺ chemiluminescence detection HPLC for D,L-amino acid). *Kuromatogurafi*, 15 (1994) 232-233; *C.A.*, 122 (1995) 155448p.
- 3053 Van de Merbel, N.C., Zuur, P., Frijlink, M., Holthuis, J.J.M., Lingeman, H. and Brinkman, U.A.T.: Automated monitoring of amino acids during fermentation processes using on-line ultrafiltration and column liquid chromatography: application to fermentation medium improvement. *Anal. Chim. Acta*, 303 (1995) 175-185.
- 3054 Yasuhara, F., Yamaguchi, S., Takeda, M., Ochiai, Y. and Miyano, S.: Gas chromatographic separation of enantiomeric amino acids and amines with α -methoxy- α -trifluoromethylpropionic acid as a chiral derivatizing agent. *J. Chromatogr. A*, 694 (1995) 227-236.
- For additional information see *C.A.*:
122 (1995) 104191s, 182748t.
- See also 2590, 2635, 2653, 2658, 2665, 2732, 2745, 2824, 2994, 3007, 3012, 3176, 3409, 3462, 3465, 3476, 3481, 3489, 3586, 3802, 3811.
- 18b. Peptides, peptidic and proteinous hormones, growth factors*
- 3055 Abdel-Kader, H., Kobykevich, M.M., Wigman, L.S. and Menon, G.K.: Gradient multipeak liquid chromatographic method for determination of ardacin in bulk chemical and premix formulations. *J. Assoc. Off. Anal. Chem.*, 78 (1995) 289-293.
- 3056 Aguilar, M.-I., Richards, K.L., Round, A.J. and Hearn, M.T.W.: Molecular definition of the retention parameters of peptides separated by RP-HPLC. *Pept. Res.*, 7 (1994) 207-217; *C.A.*, 122 (1995) 161287b.
- 3057 Basiuk, V.A. and Gromovoy, T.Yu.: Free energies of adsorption of dipeptides and 2,5-piperazinediones on silica from neutral aqueous solutions as estimated from high-performance liquid-chromatographic retention data. *Collect. Czech. Chem. Commun.*, 59 (1994) 1721-1728; *C.A.*, 122 (1995) 90309x.
- 3058 Cherkasov, A.V., Nedospasov, V.A., Yakhimovich, A.D. and Nedospasov, A.A.: (ANSA-analysis. VI. Activation, inhibition and interaction of proteases in enzyme mixtures). *Biokhimiya (Moscow)*, 59 (1994) 1574-1588.
- 3059 Congote, L.F.: Increased heparin binding by site directed mutagenesis of a recombinant chimera of bombyxin and insulin-like growth factor II. *Biochim. Biophys. Acta*, 1243 (1995) 538-542.
- 3060 Crimmins, D.L., Saylor, M., Rush, J. and Thoma, R.S.: Facile, *in situ* matrix-assisted laser desorption ionization-mass spectrometry analysis and assignment of disulfide pairings in heteropeptide molecules. *Anal. Biochem.*, 226 (1995) 355-361.
- 3061 De Ceuninck, F., Willeput, J. and Corvol, M.: Purification and characterization of insulin-like growth factor II (IGF II) and an IGF II variant from human placenta. *J. Chromatogr. B*, 666 (1995) 203-214.
- 3062 De Collongue, B., Gosselet, N.M., Sébille, N. and Schoot, B.: Comparative study of different predictive methods of peptides retention time on chromatographic reversed-phase columns. *J. Liq. Chromatogr.*, 17 (1994) 4349-4364.
- 3063 Fainzilber, M., van der Schors, R., Lodder, J.C., Li, K.W., Geraerts, W.P.M. and Kits, K.S.: New sodium channel-blocking conotoxins also affect calcium currents in *Lymnaea* neurons. *Biochemistry*, 34 (1995) 5364-5371.
- 3064 Fialaire, A. and Postaire, E.: Hydrolysis of peptide binding by phosphotungstic acid. *J. Assoc. Off. Anal. Chem.*, 77 (1994) 1338-1340.
- 3065 Goetze, J.P. and Rehfeld, J.F.: Displaced chromatographic elution of gastrin-17 in normal human serum. *Scand. J. Clin. Lab. Invest.*, 54 (1994) 673-676; *C.A.*, 122 (1995) 151799f.
- 3066 Hänslér, M., Keilhack, H. and Jakubke, H.-D.: Die Verwendung von Ficin zur Knüpfung von Peptidbindungen im gefrorenen wässrigen System. *Pharmazie*, 50 (1995) 184-187.
- 3067 Hoyer, G.L., Nolan, P.E., Jr., LeDoux, J.H. and Moore, L.A.: Selective stability-indicating high-performance liquid chromatographic assay for recombinant human regular insulin. *J. Chromatogr. A*, 699 (1995) 383-388.
- 3068 Johnston, R.N., Shaw, C., Halton, D.W., Verhaert, P. and Baguna, J.: GYIRFamide: a novel FMRFamide-related peptide (FaRP) from the triclad turbellarian, *Dugesia tigrina*. *Biochem. Biophys. Res. Commun.*, 209 (1995) 689-697.
- 3069 Klyushnichenko, V.E., Yakimov, S.A., Arutyunyan, A.M., Ivanov, A.E., Maltsev, K.V. and Wulfson, A.N.: (Recombinant human insulin. IV. Design and optimization of the system of control by reverse-phase high performance liquid chromatography). *Bioorg. Khim.*, 20 (1994) 1080-1088; *C.A.*, 122 (1995) 79157a.
- 3070 Kos, P., Gorzo, G., Suranyi, G. and Borbely, G.: Simple and efficient method for isolation and measurement of cyanobacterial hepatotoxins by plant tests (*Sinapis alba* L.). *Anal. Biochem.*, 225 (1995) 49-53.
- 3071 Li, K.W., El Filali, Z., van Golen, F.A. and Geraerts, W.P.M.: Identification of a novel amide peptide, GLTPNMNSLFF-NH₂, involved in the control of vas deferens motility in *Lymnaea stagnalis*. *Eur. J. Biochem.*, 229 (1995) 70-72.
- 3072 Manita, H.: Simple immunochromatographic method and apparatus for semiquantitative assay. *PCT Int. Appl. WO 94 28,415 (Cl. GO1N33/543)*, 08 Dec. 1994, JP Appl. 93/131,590, 02 Jun. 1993; 54 pp.; *C.A.*, 122 (1995) 101155x.
- 3073 Mitev, V., le Panse, R., Coulomb, B., Miteva, L. and Houdebine, L.-M.: Epidermal growth factor stimulates mitogen-activated protein kinase by PKC-dependent pathway in human keratinocytes. *Biochem. Biophys. Res. Commun.*, 208 (1995) 245-252.
- 3074 Nedospasov, A.A., Rodina, E.V. and Cherkasov, A.V.: (ANSA-analysis. V. Secondary specificity of peptidases for the structure of detectable groups of aminonaphthalenesulfonamide chromogenic substrates). *Biokhimiya (Moscow)*, 59 (1994) 1560-1573.
- 3075 Nguyen, T.S., Hossain, M. and Do, D.D.: Interference phenomena and non-ideal effects in displacement chromatography: experimental study of dipeptides. *Isol. Purif.*, 2 (1994) 27-44; *C.A.*, 122 (1995) 82037s.

- 3076 Nirenberg, U.: Reversed-phase HPLC. Analytical procedure. *Methods Mol. Biol. (Totowa)*, 36(Peptide Analysis Protocols) (1994) 23-35; *C.A.*, 122 (1995) 75701a.
- 3077 Okuno, S., Kitani, T. and Fujisawa, H.: Full activation of brain calmodulin-dependent protein kinase IV requires phosphorylation of the amino-terminal serine-rich region by calmodulin-dependent protein kinase IV kinase. *J. Biochem. (Tokyo)*, 117 (1995) 686-690.
- 3078 Paroni, R., de Vecchi, E., Cighetti, G., Arcelloni, C., Fermo, I., Grossi, A. and Bonini, P.: HPLC with o-phthalaldehyde precolumn derivatization to measure total, oxidized, and protein-bound glutathione in blood, plasma, and tissue. *Clin. Chem. (Washington)*, 41 (1995) 448-454.
- 3079 Ploemen, J.H.T.M., Wormhoudt, L.W., van Ommen, B., Commandeur, J.N.M., Vermeulen, N.P.E. and van Bladeren, P.J.: Polymorphism in the glutathione conjugation activity of human erythrocytes towards ethylene dibromide and 1,2-epoxy-3-(p-nitrophenoxy)-propane. *Biochim. Biophys. Acta*, 1243 (1995) 469-476.
- 3080 Polo, M.C., Gonzalez de Llano, D. and Ramos, M.: Peptides analysis by HPLC. *Acquis. Recent. Chromatogr. Vin Cours Eur. Form. Contin.*, 1992, (1993) 81-85; *C.A.*, 122 (1995) 75622a - a review with 18 refs.
- 3081 Rissler, K.: Sample preparation, high-performance liquid chromatographic separation and determination of substance P-related peptides. *J. Chromatogr. B*, 665 (1995) 233-270 - a review with 186 refs.
- 3082 Rodina, E.V., Cherkasov, A.V. and Nedospasov, A.A.: (ANSA-analysis. IV. Specificity spectra in characterization of proteases and their mixtures). *Biokhimiya (Moscow)*, 59 (1994) 1544-1559.
- 3083 Roturier, J.M., le Bars, D. and Gripon, J.C.: Separation and identification of hydrophilic peptides in dairy products using FMOc derivatization. *J. Chromatogr. A*, 696 (1995) 209-217.
- 3084 Sereda, T.J., Mant, C.T. and Hodges, R.S.: Selectivity due to conformational differences between helical and non-helical peptides in reversed-phase chromatography. *J. Chromatogr. A*, 695 (1995) 205-221.
- 3085 Shon, K., Grilley, M.M., Marsh, M., Yoshikami, D., Hall, A.R., Kurz, B., Gray, W.R., Imperial, J.S., Hillyard, D.R. and Olivera, B.M.: Purification, characterization, synthesis, and cloning of the lockjaw peptide from *Conus purpurascens* venom. *Biochemistry*, 34 (1995) 4913-4918.
- 3086 Szókán, G., Horváth, J., Almás, M., Saftics, G. and Palócz, A.: Liquid chromatographic analysis and separation of polypeptide components from honey bee venoms. *J. Liq. Chromatogr.*, 17 (1994) 3333-3349.
- 3087 Szókán, G., Majer, Z., Kollát, E., Kajtár, M., Hollósi, M. and Peredy-Kajtár, M.: Chromatographic studies on the racemization of thiopeptides. *J. Liq. Chromatogr.*, 18 (1995) 941-964.
- 3088 Thim, L., Wöldike, H.F., Nielsen, P.F., Christensen, M., Lynch-Devaney, K. and Podolsky, D.K.: Characterization of human and rat intestinal trefoil factor produced in yeast. *Biochemistry*, 34 (1995) 4757-4764.
- 3089 Waldmann, T., Nimmegern, E., Mitsch, M., Peters, J., Pfeifer, G., Müller, S., Kellermann, J., Engel, A., Hartl, F.-U. and Baumeister, W.: The thermosome of *Thermoplasma acidophilum* and its relationship to the eukaryotic chaperonin TRiC. *Eur. J. Biochem.*, 227 (1995) 848-856.
- 3090 Woodward, C. and Geiser, F.O.: Minimization of baseline disturbances in peptide mapping by means of principal-component multivariate visualization. *J. Chromatogr. A*, 699 (1995) 11-20.
- 3091 Zhao, Q., Sannier, F., Ricart, G. and Piot, J.M.: A rapid detection and identification of hemorphins released from bovine hemoglobin enzymatic hydrolysis by use of HPLC coupled with photodiode array detector. *J. Liq. Chromatogr.*, 18 (1995) 93-103.

For additional information see *C.A.*:
122 (1995) 29801k, 178566q.

See also 2614, 2629, 2639, 2705, 3109, 3120, 3125, 3199, 3243, 3244, 3387, 3795.

18c. Elucidation of structure of proteins and enzymes

- 3092 Boots, J.-W.P., van Dongen, W.D., Verheij, H.M., de Haas, G.H., Haverkamp, J. and Slotboom, A.J.: Identification of the active site histidine in *Staphylococcus hyicus* lipase using chemical modification and mass spectrometry. *Biochim. Biophys. Acta*, 1248 (1995) 27-34.
- 3093 Carrea, G., D'Arrigo, P., Piergianni, V., Roncaglio, S., Secundo, F. and Servi, S.: Purification and properties of two phospholipases D from *Streptomyces* sp. *Biochim. Biophys. Acta*, 1255 (1995) 273-279.
- 3094 Chang, L.-S., Kuo, K.-W., Lin, J., Lin, S.-R. and Chang, C.-C.: Analysis of a conformation-independent epitope and a conformational epitope in a protein: a study on cobrotoxin from Taiwan cobra venom. *J. Biochem. (Tokyo)*, 117 (1995) 863-868.
- 3095 Chobert, J.-M., Briand, L., Grinberg, V. and Haertlé, T.: Impact of esterification on the folding and the susceptibility to peptic proteolysis of β -lactoglobulin. *Biochim. Biophys. Acta*, 1248 (1995) 170-176.
- 3096 Giorgianni, F., Beranova, S., Wesdemiotis, C. and Viola, R.E.: Elimination of the sensitivity of L-aspartase to active-site-directed inactivation without alteration of catalytic activity. *Biochemistry*, 34 (1995) 3529-3535.
- 3097 Huang, D.-Y. and Ichikawa, Y.: Identification of essential lysyl and cysteinyl residues, and the amino acid sequence at the substrate-binding site of retinal oxidase. *Biochim. Biophys. Acta*, 1243 (1995) 431-436.
- 3098 Jaquet, K., Fukunaga, K., Miyamoto, E. and Meyer, H.E.: A site phosphorylated in bovine cardiac troponin T by cardiac CaM kinase II. *Biochim. Biophys. Acta*, 1248 (1995) 193-195.
- 3099 Mellet, P., Ermolieff, J. and Bieth, J.G.: Mapping the heparin-binding site of mucus proteinase inhibitor. *Biochemistry*, 34 (1995) 2645-2652.
- 3100 Miller, C.G.: Adsorptive biphasic column technology for protein sequence analysis and protein chemical modification. *Methods (San Diego)*, 6 (1994) 315-333; *C.A.*, 122 (1995) 26994q.
- 3101 Murphy, C.M. and Fenselau, C.: Recognition of the carboxy-terminal peptide in cyanogen bromide digests of proteins. *Anal. Chem.*, 67 (1995) 1644-1645.
- 3102 Ojima, T., Tanaka, H. and Nishita, K.: Amino acid sequence of C-terminal 17 kDa CNBr-fragment of Akazara scallop troponin-I. *J. Biochem. (Tokyo)*, 117 (1995) 158-162.

- 3103 Pederson, J., Filimonova, M., Roepstorff, P. and Biedermann, K.: (Isoforms of *Serratia marcescens* nuclease produced by natural and recombinant strains. Comparative characterization by plasma desorption mass spectrometry). *Biokhimiya (Moscow)*, 160 (1995) 450-461.
- 3104 Suzuki, K., Mizuguchi, M., Gomi, T. and Itagaki, E.: Identification of a lysine residue in the NADH-binding site of salicylate hydroxylase from *Pseudomonas putida* S-1. *J. Biochem. (Tokyo)*, 117 (1995) 579-585.
- 3105 Swoboda, I., Jilek, A., Ferreira, F., Engel, E., Hoffmann-Sommergruber, K., Scheiner, O., Kraft, D., Breiteneder, H., Pittenauer, E., Schmid, E. et al.: Isoforms of Bet v 1, the major birch pollen allergen, analyzed by liquid chromatography, mass spectrometry, and cDNA cloning. *J. Biol. Chem.*, 270 (1995) 2607-2613.
- 3106 Vaintraub, I.A. and Lapteva, N.A.: (Kinetics of one-by-one proteolysis. Ovalbumin proteolysis). *Biokhimiya (Moscow)*, 59 (1994) 1406-1413.
- 3107 Yamashita, T., Yomogida, S., Nagaoka, I. and Saito, K.: Effect of a chemical or proteolytic modification on the biological activity of guinea-pig cationic peptide. *Biochim. Biophys. Acta*, 1243 (1995) 295-299.
- 3108 Young, P.M. and Gorenstein, M.V.: Tryptic mapping by reversed-phase HPLC with photodiode-array detection incorporating the spectral-contrast technique. *LC-GC*, 12 (1994) 832-838; *C.A.*, 122 (1995) 208891m.
- See also 3090, 3176, 3194, 3200, 3206, 3208, 3222, 3318.
19. PROTEINS
- 19a. General techniques
- 3109 Bailey, J.M., Tu, O., Issai, G., Ha, A. and Shively, J.E.: Automated carboxy-terminal sequence analysis of polypeptides containing C-terminal proline. *Anal. Biochem.*, 224 (1995) 588-596.
- 3110 Belenkii, B.G. and Malt'sev, V.G.: High-performance membrane chromatography. *BioTechniques*, 18 (1995) 288-291; *C.A.*, 122 (1995) 127808n.
- 3111 Bollag, D.M.: Ion-exchange chromatography. *Methods Mol. Biol. (Totowa)*, 36 (Peptide Analysis Protocols) (1994) 11-22; *C.A.*, 122 (1995) 75700z.
- 3112 Chang, J.-H. and Han, Y.-P.: (Studies on new packings for high-performance ion-exchange chromatography for the separation of proteins. (II). Synthesis and characteristics of new packings for weak cation-exchange chromatography). *Gaodeng, Xuexiao Huaxue Xuebao*, 15 (1994) 1317-1321; *C.A.*, 122 (1995) 100738c.
- 3113 Corradini, D., Capitani, D. and Cellai, L.: Salt concentration effects in high-performance hydrophobic-interaction chromatography in comparison with NMR of proteins in solution. *J. Liq. Chromatogr.*, 17 (1994) 4335-4347.
- 3114 Feng, W., Jiang, X. and Geng, X.: The influence of urea on the retention of protein in hydrophobic interaction chromatography. *J. Liq. Chromatogr.*, 18 (1995) 217-226.
- 3115 Gerstner, J.A., Morris, J., Hunt, T., Hamilton, R. and Afeyan, N.B.: Rapid ion-exchange displacement chromatography of proteins on perfusive chromatographic supports. *J. Chromatogr. A*, 695 (1995) 195-204.
- 3116 Jacob, L.R., Mack, M. and Rathgeber, G.: Large-scale purification of proteins from egg white by tentacle ion-exchangers. *Am. Biotechnol. Lab.*, 12 (1994) 37; *C.A.*, 122 (1995) 75698e.
- 3117 Johnson, E.M., Berk, D.A., Jain, R.K. and Deen, W.M.: Diffusion and partitioning of proteins in charged agarose gels. *Biophys. J.*, 68 (1995) 1561-1568; *C.A.*, 122 (1995) 209067j.
- 3118 Kabzinski, A.K.M., Tagaki, T. and Tanaka, K.: The application of HPLC-LALLS method for determination of changes in thiolproteins and metallothioprotenes during separations by covalent affinity chromatography in example of metallothioneine. *Eur. J. Clin. Chem. Clin. Biochem.*, 33 (1995) A22-A23.
- 3119 Kasai, K.-i.: (Intelligent chromatography and electrophoresis as tools for understanding of biomolecules). *Kuromatogurafi*, 15 (1994) 206-209; *C.A.*, 122 (1995) 127787e.
- 3120 Kawakatsu, M., Kotaniguchi, H., Freiser, H.H. and Gooding, K.M.: The effects of pore diameter and ligand chain length on fast liquid chromatography of proteins and peptides. *J. Liq. Chromatogr.*, 18 (1995) 633-647.
- 3121 Kulish, D.M., Klyushnichenko, V.E. and Vulfson, A.N.: (Separation of proteins by liquid chromatography and capillary electrophoresis). *Zh. Fiz. Khim.*, 68 (1994) 1841-1847; *C.A.*, 122 (1995) 50407w.
- 3122 Kurnik, R.T., Yu, A.W., Blank, G.S., Burton, A.R., Smith, D., Athalye, A.M. and van Reis, R.: Buffer exchange using size exclusion chromatography, countercurrent dialysis, and tangential flow filtration: models development, and industrial application. *Biotechnol. Bioeng.*, 45 (1995) 149-157; *C.A.*, 122 (1995) 170139g.
- 3123 Laubscher, J.C.: (Optimization of HPLC [for protein separations]). *Schweiz. Lab.-Z.*, 51 (1994) 162-166; *C.A.*, 122 (1995) 208797k - a review with no refs.
- 3124 Nakazato, K., Mohammad, J. and Hjerten, S.: Studies of some experimental conditions for the polymerization of the continuous beads (cross-linked polyacrylamide gels) in order to improve their chromatographic properties. *Chromatographia*, 39 (1994) 655-662.
- 3125 Narayanan, S.R.: Application of liquid chromatography to the purification of proteins and peptides. *Prog. Pharm. Biomed. Anal.*, 1 (1994) 279-312; *C.A.*, 122 (1995) 234471u.
- 3126 Oscarsson, S.: Influence of salts on protein interactions at interfaces of amphiphilic polymers and adsorbents. *J. Chromatogr. B*, 666 (1995) 21-31.
- 3127 Patrickios, C.S., Gadam, S.D., Cramer, S.M., Hertler, W.R. and Hatton, T.A.: Block methacrylic polyampholytes as protein displacers in ion-exchange chromatography. *Biotechnol. Prog.*, 11 (1995) 33-38; *C.A.*, 122 (1995) 100732w.
- 3128 Tauer, C., Buchacher, A. and Jungbauer, A.: DNA clearance in chromatography of proteins, exemplified by affinity chromatography. *J. Biochem. Biophys. Methods*, 30 (1995) 75-78; *C.A.*, 122 (1995) 182347m.
- 3129 Uversky, V.N.: Gel-permeation chromatography as a unique instrument for quantitative and qualitative analysis of protein denaturation and unfolding. *Int. J. Bio-Chromatogr.*, 1 (1994) 103-114; *C.A.*, 122 (1995) 50340u - a review with 45 refs.

3130 Yu, Q. and Wang, F.-Y.: Optimization of ionic strength gradient elution chromatography of proteins. In: *Off. Proc. Comb. Conf., 6th Conf. Asia Pac. Confed. Chem. Eng., 21st Australas. Chem. Eng. Conf., Inst. Eng. Aust., Barton, 1993*, pp. 113/3-118/3; *C.A.*, 122 (1995) 26979p.

For additional information see C.A.:

122 (1995) 26973g, 50428d, 75710c, 182241x, 212099b.

See also 2571, 2629, 2649, 2684, 2758, 3039, 3084, 3218, 3225, 3814.

19b. *Proteins of cells, viruses and subcellular particles*

3131 Cai, S.-J., McAndrew, R.S., Leonard, B.P., Chapman, K.D. and Pidgeon, C.: Rapid purification of cotton seed membrane-bound N-acylphosphatidylethanolamine synthase by immobilized artificial membrane chromatography. *J. Chromatogr. A*, 696 (1995) 49-62.

3132 Kandrór, K.V., Coderre, L., Pushkin, A.V. and Pilch, P.F.: Comparison of glucose-transporter-containing vesicles from rat fat and muscle tissue: evidence for a unique endosomal compartment. *Biochem. J.*, 307 (1995) 383-390.

3133 Krantz, S., Salazar, R., Brandt, R., Kellermann, J. and Lottspeich, F.: Purification and partial amino acid sequencing of a fructosyllysine-specific binding protein from cell membranes of the monocyte-like cell line U937. *Biochim. Biophys. Acta*, 1266 (1995) 109-112.

3134 Naylor, D.J., Ryan, M.T., Condrón, R., Hoogenraad, N.J. and Høj, P.B.: Affinity-purification and identification of GrpE homologues from mammalian mitochondria. *Biochim. Biophys. Acta*, 1248 (1995) 75-79.

3135 Suzuki, T. and Kawakita, M.: Sites of labeling with N-(3-pyrene)maleimide on Ca²⁺-transporting ATPase of the sarcoplasmic reticulum. *J. Biochem. (Tokyo)*, 117 (1995) 881-887.

For additional information see C.A.:

122 (1995) 155383p.

19c. *Proteins synthesized by genetic manipulation, monoclonal antibodies*

3136 Crowe, J., Doebeli, H., Gentz, R., Hochuli, E., Stueber, D. and Henco, K.: 6xHis-Ni-NTA chromatography as a superior technique in recombinant protein expression/purification. *Methods Mol. Biol. (Totowa)*, 31 (Protocols for Gene Analysis) (1994) 371-387; *C.A.*, 122 (1995) 97826x.

3137 Klyushnichenko, V.E., Wulfson, A.N., Titel, C., Ehwald, R. and Miroshnikov, A.I.: (Separation of recombinant proteins by vesicle chromatography). *Bioorg.Khim.*, 20 (1994) 894-898; *C.A.*, 122 (1995) 100769p.

3138 Kwon, K.-S., Lee, S. and Yu, M.-H.: Refolding of α -antitrypsin expressed as inclusion bodies in *Escherichia coli*: characterization of aggregation. *Biochim. Biophys. Acta*, 1247 (1995) 179-184.

See also 3059, 3067, 3069, 3271.

19d. *Microbial and plant proteins*

3139 Adam, O., Vercellone, A., Paul, F., Monsan, P.F. and Puzo, G.: A nondegradative route for the removal of endotoxin from exopolysaccharides. *Anal. Biochem.*, 225 (1995) 321-327.

3140 Ahmed, Md.S., Khan, S., Arakawa, O. and Onoue, Y.: Properties of hemagglutinins newly separated from toxic phytoplankton. *Biochim. Biophys. Acta*, 1243 (1995) 509-512.

3141 Feldwisch, J., Zettl, R., Campos, N. and Palme, K.: Identification of a 23 kDa protein from maize photoaffinity-labelled with 5-azido-[7-³H]indol-3-ylacetic acid. *Biochem. J.*, 305 (1995) 853-857.

3142 Gassenschmidt, U., Jany, K.D., Tauscher, B. and Niebergall, H.: Isolation and characterization of a flocculating protein from *Moringa oleifera* Lam. *Biochim. Biophys. Acta*, 1243 (1995) 477-481.

3143 Hamato, N., Koshiba, T., Pham, T.-N., Tatsumi, Y., Nakamura, D., Takano, R., Hayashi, K., Hong, Y.-M. and Hara, S.: Trypsin and elastase inhibitors from bitter gourd (*Momordica charantia* Linn.) seeds: purification, amino acid sequences, and inhibitory activities of four new inhibitors. *J. Biochem. (Tokyo)*, 117 (1995) 432-437.

3144 Hirayama, C., Sakata, M., Morimoto, S., Tachibe, H., Ihara, H., Ohkuma, K. and Miyachi, Y.: The removal of endotoxins from protein solutions using column packings with aminated poly(γ -methyl L-glutamate) spheres. *Chromatographia*, 40 (1995) 33-38.

3145 Jekow, P., Fromme, P., Witt, H.T. and Saenger, W.: Photosystem I from *Synechococcus elongatus*: preparation and crystallization of monomers with varying subunit compositions. *Biochim. Biophys. Acta*, 1229 (1995) 115-120.

3146 Ko, J.-L., Hsu, C.-I., Lin, R.-H., Kao, C.-L. and Lin, J.Y.: A new fungal immunomodulatory protein, FIP-five isolated from the edible mushroom, *Flammulina velutipes* and its complete amino acid sequence. *Eur. J. Biochem.*, 228 (1995) 244-249.

3147 Nitti, G., Orru, S., Bloch, C., Jr., Morhy, L., Marino, G. and Pucci, P.: Amino acid sequence and disulphide-bridge pattern of three γ -thionins from *Sorghum bicolor*. *Eur. J. Biochem.*, 228 (1995) 250-256.

3148 Osman, A.A., Börner, H. and Mothes, T.: Isolation of two gamma-gliadins and determination of their N-terminal amino acid sequence. *Eur. J. Clin. Chem. Clin. Biochem.*, 33 (1995) A20.

3149 Terabe, M., Kojima, S., Taguchi, T., Momose, H. and Miura, K.-i.: A subtilisin inhibitor produced by *Streptomyces bikiniensis* possesses a glutamine residue at reactive site P1. *J. Biochem. (Tokyo)*, 117 (1995) 609-613.

3150 Weegels, P.L., Marseille, J.P., Bosveld, P. and Harner, R.J.: Large-scale separation of gliadins and their bread-making quality. *J. Cereal Sci.*, 20 (1994) 253-264; *C.A.*, 122 (1995) 159163q.

3151 Xu, Y.-H., Lu, Z.-Y. and Ihler, G.M.: Purification of deformin, an extracellular protein synthesized by *Bartonella bacilliformis* which causes deformation of erythrocyte membranes. *Biochim. Biophys. Acta*, 1234 (1995) 173-183.

3152 Zeidler, W., Egle, C., Ribeiro, S., Wagner, A., Katunin, V., Kreutzer, R., Rodnina, M., Wintermeyer, W. and Sprinzl, M.: Site-directed mutagenesis of *Thermus thermophilus* elongation factor Tu. Replacement of His85, Asp 81 and Arg300. *Eur. J. Biochem.*, 229 (1995) 596-604.

For additional information see C.A.:
122 (1995) 98897h, 207166s.

See also 3070, 3105, 3187, 3239, 3240.

19e. *Proteins of blood, serum and blood cells*

- 3153 Altman, E. and Bundle, D.R.: Polysaccharide affinity columns for purification of lipopolysaccharide-specific murine monoclonal antibodies. *Methods Enzymol.*, 247(Neoglycoconjugates, Pt. B) (1994) 243-253; C.A., 122 (1995) 184908u.
- 3154 Asano, S.: Antibody purification by affinity chromatography. *Jpn. Kokai Tokkyo Koho* JP 06,172,399 [94,172,399] (Cl. CO7K15/14), 21 Jun. 1994, Appl. 92/325,416, 04 Dec. 1992; 4 pp.; C.A., 122 (1995) 130998e.
- 3155 Brumeanu, T.D., Zaghouani, H. and Bona, C.: Purification of antigenized immunoglobulins derivatized with monomethoxy-polyethylene glycol. *J. Chromatogr. A*, 696 (1995) 219-225.
- 3156 Chenou, C. and Fick, M.: (Kinetics and modeling of albumin adsorption in ion-exchange chromatography). *Recent Prog. Genie Procedes*, 7(27, Phenomenes Fondamentaux: Developpements Methodologiques) (1993) 37-42; C.A., 122 (1995) 155399y.
- 3157 Chroneos, Z.C., Baynes, J.W. and Thorpe, S.R.: Identification of liver endothelial cells as a primary site of IgM catabolism in the rat. *Arch. Biochem. Biophys.*, 319 (1995) 63-73.
- 3158 Denizli, A. and Piskin, E.: DNA-immobilized polyhydroxyethylmethacrylate microbeads for affinity sorption of human immunoglobulin G and anti-DNA antibodies. *J. Chromatogr. B*, 666 (1995) 215-222.
- 3159 Ernst, A., Helmhold, M., Brunner, C., Pethö-Schramm, A., Armstrong, V.W. and Müller, H.-J.: Identification of two functionally distinct lysine-binding sites in kringle 37 and in kringles 32-36 of human apolipoprotein(a). *J. Biol. Chem.*, 270 (1995) 6227-6234.
- 3160 Garcia de Frutos, P. and Dahlbäck, B.: cDNA structure of rabbit C4b-binding protein α -chain. Preserved sequence motive in complement regulatory protein modules which bind C4b. *Biochim. Biophys. Acta*, 1261 (1995) 285-289.
- 3161 Higgins, D.A., Cromie, R.L., Liu, S.S., Magor, K.E. and Warr, G.W.: Purification of duck immunoglobulins: an evaluation of protein A and protein G affinity chromatography. *Vet. Immunol. Immunopathol.*, 44 (1995) 169-180; C.A., 122 (1995) 211516y.
- 3162 Ingendoh, A., Nakayama, H., Isobe, T., Ishioka, N., Kanai, M., Seta, K. and Okuyama, T.: (Analysis of human plasma proteins utilizing HPLC-ESI/MS and MALDI/TOFMS). *Kuromatogurafi*, 15 (1994) 260-261; C.A., 122 (1995) 127793d.
- 3163 Jacob, L., Schmitt, E. and Bruemmer, W.: Purification of antibodies by new chromatography techniques. *Am. Biotechnol. Lab.*, 12 (1994) 44-45; C.A., 122 (1995) 50390k.
- 3164 Kent, U.M.: Purification of antibodies using affinity chromatography. *Methods Mol. Biol. (Totowa)*, 34(Immunocytochemical Methods and Protocols) (1994) 29-35; C.A., 122 (1995) 103414e.
- 3165 Kent, U.M.: Purification of antibodies using ion-exchange chromatography. *Methods Mol. Biol. (Totowa)*, 34(Immunocytochemical Methods and Protocols) (1994) 23-27; C.A., 122 (1995) 103413d.
- 3166 Kuroma, T., Kurata, S. and Natori, S.: Molecular characterization of an insect transferrin and its selective incorporation into eggs during oogenesis. *Eur. J. Biochem.*, 228 (1995) 229-235.
- 3167 Litvinovich, S.V., Henschen, A.H., Krieglstein, K.G., Ingham, K.C. and Medved, L.V.: Structural and functional characterization of proteolytic fragments derived from the C-terminal regions of bovine fibrinogen. *Eur. J. Biochem.*, 229 (1995) 605-614.
- 3168 Loun, B. and Hage, D.S.: Characterization of thyroxine-albumin binding using high-performance affinity chromatography. II. Comparison of the binding of thyroxine, triiodothyronines and related compounds at the warfarin and indole sites of human serum albumin. *J. Chromatogr. B*, 665 (1995) 303-314.
- 3169 Martin, S.E.V., Thomas, O.R.T., Owens, R., Hill, C.R. and Dunnill, P.: Engineering and expression of a single-chain antibody for purification by immobilized metal affinity chromatography. In: *ICHEME Res. Event, Two-Day Symp.*, 1, Inst. Chem. Eng., Rugby, 1994, pp. 85-87; C.A., 122 (1995) 29504r.
- 3170 Miyata, T., Funatsu, A. and Kato, H.: Chemical cross-linking of activated coagulation factor VII with soluble tissue factor: calcium ions are not essential for full amidolytic activity of the factor VIIA-tissue factor complex after complex formation. *J. Biochem. (Tokyo)*, 117 (1995) 836-844.
- 3171 Motojima, K. and Goto, S.: Characterization of serum proteins down-regulated by peroxisome proliferators: transient repression of apoE gene expression in the rat liver. *J. Biochem. (Tokyo)*, 117 (1995) 597-602.
- 3172 Nassander, U.K., Steerenberg, P.A., de Jong, W.H., van Overveld, W.O.W.M., te Boekhorst, C.M.E., Poels, L.G., Jap, P.H.K. and Storm, G.: Design of immunoliposomes directed against human ovarian carcinoma. *Biochim. Biophys. Acta*, 1235 (1995) 126-139.
- 3173 Oono, Y.: (Preparation of active blood coagulation factor XII). *Jpn. Kokai Tokkyo Koho* JP 06,174,727 [94,174,727] (Cl. G01N33/86), 24 Jun. 1994, Appl. 92/329,180, 09 Dec. 1992; 5 p.; C.A., 122 (1995) 142481u.
- 3174 Pirie-Shepherd, S.R., Jett, E.A., Andon, N.L. and Pizzo, S.V.: Sialic acid content of plasminogen 2 glycoforms as a regulator of fibrinolytic activity. Isolation, carbohydrate analysis, and kinetic characterization of six glycoforms of plasminogen 2. *J. Biol. Chem.*, 270 (1995) 5877-5881.
- 3175 Tharakan, J.P. and Ayers, V.: Immobilized ligand penetration and capacity in resin-based immunoaffinity purification of factor IX. *Int. J. Bio-Chromatogr.*, 1 (1994) 93-102; C.A., 122 (1995) 50421w.
- 3176 Wang, X., DeVries, A.L. and Cheng, C.-H.: Antifreeze peptide heterogeneity in an antarctic eel pout includes an unusually large major variant comprised of two 7 kDa type III AFPs linked in tandem. *Biochim. Biophys. Acta*, 1247 (1995) 163-172.
- 3177 Yasuda, F., Hayashi, T., Tanitame, K., Nishioka, J. and Suzuki, K.: Molecular cloning and functional characterization of rat plasma protein S. *J. Biochem. (Tokyo)*, 117 (1995) 374-383.

For additional information see C.A.:
122 (1995) 29509w, 50736c, 184909v.

See also 2713, 3138, 3186, 3189, 3192, 3197, 3198, 3229, 3351.

19f. *Structural and muscle proteins*

- 3178 Ahmad, F. and Goldstein, B.J.: Purification, identification and subcellular distribution of three predominant protein-tyrosine phosphatase enzymes in skeletal muscle tissue. *Biochim. Biophys. Acta*, 1248 (1995) 57-69.
- 3179 Bailey, A.J., Sims, T.J., Avery, N.C. and Halligan, E.P.: Non-enzymic glycation of fibrous collagen: reaction products of glucose and ribose. *Biochem. J.*, 305 (1995) 385-390.
- 3180 Di Lisa, F., De Tullio, R., Salamino, F., Barbato, R., Melloni, E., Siliprandi, N., Schiaffino, S. and Pontremoli, S.: Specific degradation of troponin T and I by μ -calpain and its modulation by substrate phosphorylation. *Biochem. J.*, 308 (1995) 57-61.
- 3181 Postnikova, G.B.: (Chemical modification of myoglobin by isothiocyanate reagents. Effect of the N-terminal amino group modification on protein conformation). *Biokhimiya (Moscow)*, 59 (1994) 1445-1457.
- 3182 Potter, J.D., Sheng, Z., Pan, B. and Zhao, J.: A direct regulatory role for troponin T and a dual role for troponin C in the Ca^{2+} regulation of muscle contraction. *J. Biol. Chem.*, 270 (1995) 2557-2562.
- 3183 Skolysheva, L.K., Smirnova, E.A., Medvedeva, M.V. and Gusev, N.B.: (Isolation and some properties of bovine aorta caldesmon). *Biokhimiya (Moscow)*, 60 (1995) 339-348.
- 3184 Tillet, E., Mann, K., Nischt, R., Pan, T.-c., Chu, M.-L. and Timpl, R.: Recombinant analysis of human $\alpha 1(XVI)$ collagen. Evidence for processing of the N-terminal globular domain. *Eur. J. Biochem.*, 228 (1995) 160-168.

For additional information see C.A.:

122 (1995) 159742c.

See also 2719, 3098, 3132, 3185, 3387.

19h. *Chromoproteins and metalloproteins*

- 3185 Baksh, S., Spamer, C., Oikawa, K., McCubbin, W.D., Heilmann, C., Kay, C.M. and Michalak, M.: Zn^{2+} binding to cardiac calsequestrin. *Biochem. Biophys. Res. Commun.*, 209 (1995) 310-315.
- 3186 Bissé, E., Huaman-Guillen, P. and Wieland, H.: Chromatographic evaluation of minor hemoglobins: clinical significance of hemoglobin A_{1d}, comparison with hemoglobin A_{1c}, and possible interferences. *Clin. Chem. (Washington)*, 41 (1995) 658-663.
- 3187 Chan, C., Willis, A.C., Robinson, C.V., Aplin, R.T., Radford, S.E. and Ferguson, S.J.: The complete amino acid sequence confirms the presence of pseudoazurin in *Thiosphaera pantotropa*. *Biochem. J.*, 308 (1995) 585-590.
- 3188 De Pina, K., Navarro, C., McWalter, L., Boxer, D.H., Price, N.C., Kelly, S.M., Mandrand-Berthelot, M.-A. and Wu, L.-F.: Purification and characterization of the periplasmic nickel-binding protein NikA of *Escherichia coli* K12. *Eur. J. Biochem.*, 227 (1995) 857-865.
- 3189 Huaman-Guillen, P., Bissé, E. and Wieland, H.: Determination of carbamylated hemoglobin (CarHb) by HPLC. *Eur. J. Clin. Chem. Clin. Biochem.*, 33 (1995) A19-A20.
- 3190 Hunziker, P.E., Kaur, P., Wan, M. and Känzig, A.: Primary structures of seven metallothioneins from rabbit tissue. *Biochem. J.*, 306 (1995) 265-270.

- 3191 Lee, W.-Y., Brune, D.C., Lobrutto, R. and Blankenship, R.E.: Isolation, characterization and primary structure of rubredoxin from the photosynthetic bacterium, *Heliobacillus mobilis*. *Arch. Biochem. Biophys.*, 318 (1995) 80-88.
- 3192 Li, M., Lin, Z. and Johnson, M.E.: Photoaffinity labelling of cyanomethaemoglobin with derivatives of tryptophan and 5-bromotryptophan. *Biochem. J.*, 308 (1995) 251-260.
- 3193 Mateescu, M.A., Chakine, R., Roger, S., Atanasiu, R., Yamaguchi, N., Lamumiere, G. and Nadeau, R.: Protection of myocardial tissue against deleterious effects of oxygen free radicals by ceruloplasmin. *Arzneim. Forsch.*, 45 (1995) 476-480.
- 3194 Qian, Z.-Y., Jollès, P., Migliore-Samour, D. and Fiat, A.-M.: Isolation and characterization of sheep lactoferrin, an inhibitor of platelet aggregation and comparison with human lactoferrin. *Biochim. Biophys. Acta*, 1243 (1995) 25-32.
- 3195 Robinson, K., Jones, D., Howell, S., Soneji, Y., Martin, S. and Aitken, A.: Expression and characterization of maize ZBP14, a member of a new family of zinc-binding proteins. *Biochem. J.*, 307 (1995) 267-272.
- 3196 Tashiro, H. and Sone, N.: Preparation and characterization of the hydrophilic Cu₂-cytochrome c domain of subunit II of cytochrome c oxidase from thermophilic *Bacillus* P53. *J. Biochem. (Tokyo)*, 117 (1995) 521-526.
- 3197 Turpeinen, U., Karjalainen, U. and Stenman, U.-H.: Three assays for glycohemoglobin compared. *Clin. Chem. (Washington)*, 41 (1995) 191-195.
- 3198 Turpeinen, U., Sipilä, I., Anttila, P., Karjalainen, U., Kuronen, B., Kalkkinen, N., Ahola, T. and Stenman, U.-H.: Two α -chain hemoglobin variants, Hb Broussais and Hb Cemenelum, characterized by cation-exchange HPLC, isoelectric focusing, and peptide sequencing. *Clin. Chem. (Washington)*, 41 (1995) 532-536.

For additional information see C.A.:

122 (1995) 26987q, 50419b.

See also 3118, 3166, 3313.

19i. *Proteins of glands, gland products, various zymogens (incl. milk proteins)*

- 3199 Cervenansky, C., Engström, A. and Karlsson, E.: Role of arginine residues for the activity of fasciculins. *Eur. J. Biochem.*, 229 (1995) 270-275.
- 3200 Chu, C.-C., Li, S.-H. and Chen, Y.-H.: Resolution of isotoxins in the β -bungarotoxin family. *J. Chromatogr. A*, 694 (1995) 492-497.
- 3201 Datta, G., Dong, A., Witt, J. and Tu, A.T.: Biochemical characterization of basilase, a fibrinolytic enzyme from *Crotalus basiliscus basiliscus*. *Arch. Biochem. Biophys.*, 317 (1995) 365-373.
- 3202 Escriva, H., Pierce, A., Coddeville, B., Gonzalez, F., Benaissa, A., Leger, D., Wieruszkeski, J.-M., Spik, G. and Pamblanco, M.: Rat mammary-gland transferrin: nucleotide sequence, phylogenetic analysis and glycan structure. *Biochem. J.*, 307 (1995) 47-55.
- 3203 Fainzilber, M., Lodder, J.C., Kits, K.S., Kofman, O., Vinnitsky, I., van Rietschoten, J., Zlotkin, E. and Gordon, D.: A new conotoxin affecting sodium current inactivation interacts with the δ -conotoxin receptor site. *J. Biol. Chem.*, 270 (1995) 1123-1129.

- 3204 Fiorucci, L., Erba, F., Falasca, L., Dini, L. and Ascoli, F.: Localization and interaction of bovine pancreatic trypsin inhibitor and trypsin in the granules of bovine mast cells. *Biochim. Biophys. Acta*, 1243 (1995) 407-413.
- 3205 Hay, D.I., Ahern, J.M., Schluckebier, S.K. and Schlesinger, D.H.: Human salivary acidic proline-rich protein polymorphisms and biosynthesis studied by high-performance liquid chromatography. *J. Dent. Res.*, 73 (1994) 1717-1726; *C.A.*, 122 (1995) 182305w.
- 3206 Lochnit, G. and Geyer, R.: Carbohydrate structure analysis of batroxobin, a thrombin-like serine protease from *Bothrops moojeni* venom. *Eur. J. Biochem.*, 228 (1995) 805-816.
- 3207 Machold, J., Weise, C., Utkin, Y.N., Franke, P., Tsetlin, V.I. and Hucho, F.: A new class of photoactivable and cleavable derivatives of neurotoxin II from *Naja naja oxiana*. Synthesis, characterizations, and application for affinity labelling of the nicotinic acetylcholine receptor from *Torpedo californica*. *Eur. J. Biochem.*, 228 (1995) 947-954.
- 3208 Marangoni, S., Toyama, M.H., Arantes, E.C., Giglio, J.R., da Silva, C.A., Carneiro, E.M., Goncalves, A.A. and Oliveira, B.: Amino acid sequence of TsTX-V, an α -toxin from *Tityus serrulatus* scorpion venom, and its effect on K⁺ permeability of β -cells from isolated rat islets of Langerhans. *Biochim. Biophys. Acta*, 1243 (1995) 309-314.
- 3209 Nikai, T., Ohara, A., Komori, Y., Fox, J.W. and Sugihara, H.: Primary structure of a coagulant enzyme, bilineobin, from *Agkistrodon bilineatus* venom. *Arch. Biochem. Biophys.*, 318 (1995) 89-96.
- 3210 Ota, Y., Ido, K., Kimura, K., Fukui, E. and Ikemoto, S.: Electrophoretic analysis of bile proteins from patients with and without gallstones. *Electrophoresis (Weinheim)*, 16 (1995) 402-406.
- 3211 Palumbo, A. and Jackson, I.J.: Peroxidase activity in the ink gland of *Sepia officinalis* and partial nucleotide sequence of a candidate cDNA encoding the enzyme. *Biochim. Biophys. Acta*, 1247 (1995) 173-178.
- 3212 Schulze-Forster, K., Eckert, K. and Maurer, H.R.: The thymus extract Thymex-L potentiates the retinoic acid-induced differentiation of the human myeloid leukemia cell line HL-60. *Biochim. Biophys. Acta*, 1265 (1995) 110-116.
- 3213 Urbanke, W. and Luf, W.: (Identification of α -lactalbumin and β -lactoglobulin in sheep, goat, and mare milk by reversed-phase HPLC and derivative spectroscopy). *Lebensmittelind. Milchwirtsch.*, 114 (1993) 403-407; *C.A.*, 122 (1995) 131253v.
- 3214 Yu, H., Diamandis, E.P., Monne, M. and Croce, C.M.: Oral contraceptive-induced expression of prostate-specific antigen in the female breast. *J. Biol. Chem.*, 270 (1995) 6615-6618.
- 3215 Zietlow, M.F. and Etzel, M.R.: Evaluation of sulfopropyl ion-exchange membrane cartridges for isolation of proteins from bovine whey. *J. Liq. Chromatogr.*, 18 (1995) 1001-1018.
- See also 3094, 3095.
- 19j. *Proteins of brain, cerebrospinal fluid and eye*
- 3216 Borghini, I., Barja, F., Pometta, D. and James, R.W.: Characterization of subpopulations of lipoprotein particles isolated from human cerebrospinal fluid. *Biochim. Biophys. Acta*, 1255 (1995) 192-200.
- 3217 Chao, D., Severson, D.L., Zwiers, H. and Hollenberg, M.D.: Radiolabelling of bovine myristoylated alanine-rich protein kinase C substrate (MARCKS) in an ADP-ribosylation reaction. *Biochem. Cell Biol.*, 72 (1994) 391-396.
- 3218 Isobe, T., Nakayama, H., Ishioka, N., Kanai, M., Ingendoh, A., Seta, K. and Okuyama, T.: (Analysis of brain proteins utilizing HPLC-ESI/MS and MALDI/TOFMS). *Kuromatogurafi*, 15 (1994) 262-263; *C.A.*, 122 (1995) 127794e.
- 3219 Lambrechts, A., van Damme, J., Goethals, M., Vandekerckhove, J. and Ampe, C.: Purification and characterization of bovine profilin II. Actin, poly(L-proline) and inositolphospholipid binding. *Eur. J. Biochem.*, 230 (1995) 281-286.
- 3220 Melloni, E., Sparatore, B., Patrone, M., Pessino, A., Passalacqua, M. and Pontremoli, S.: Identity in molecular structure between "differentiation enhancing factor" of murine erythroleukemia cells and the 30 kD heparin-binding protein of developing rat brain. *Biochem. Biophys. Res. Commun.*, 210 (1995) 82-89.
- 3221 Smith, J.B., Yang, Z., Lin, P., Zaidi, Z., Abbasi, A. and Russell, P.: The complete sequence of human lens γ -S crystallin. *Biochem. J.*, 307 (1995) 407-410.
- 3222 Strub, J.-M., Garcia-Sablone, P., Lonning, K., Taupenot, L., Hubert, P., van Dorsselaer, A., Aunis, D. and Metz-Boutigue M.-H.: Processing of chromogranin B in bovine adrenal medulla. Identification of secretolytin, the endogenous C-terminal fragment of residues 614-626 with antibacterial activity. *Eur. J. Biochem.*, 229 (1995) 356-368.
- See also 3157, 3315.
- 19k. *Proteins of neoplastic tissue and transformed cells*
- 3223 Agrez, M.V., Meldrum, C.J., Sim, A.T.R., Aebersold, R.H., Clark, I.M., Cawston, T.E. and Burns, G.F.: A fibroblast elongation factor purified from colon carcinoma cells shares sequence identity with TIMP-1. *Biochem. Biophys. Res. Commun.*, 206 (1995) 590-600.
- 19l. *Specific binding and receptor proteins*
- 3224 Berggren Söderlund, M., Johannesson, G. and Fex, G.: Expression of humal *all-trans*-retinoic acid receptor and its ligand-binding domain in *Escherichia coli*. *Biochem. J.*, 308 (1995) 353-359.
- 3225 Buus, S., Stryhn, A., Winther, K., Kirkby, N. and Pedersen, L.O.: Receptor-ligand interactions measured by an improved spun column chromatography technique. A high efficiency and high throughput size separation method. *Biochim. Biophys. Acta*, 1243 (1995) 453-460.
- 3226 Dostalova, Z., Calvete, J.J., Sanz, L. and Töpfer-Petersen, E.: Boar spermadhesin AWN-1. Oligosaccharide and zona pellucida binding characteristics. *Eur. J. Biochem.*, 230 (1995) 329-336.
- 3227 Dotzlaw, J.E., Ghag, S.K. and Yeh, W.-K.: Process for purification of water-soluble penicillin-binding protein 2a. *Eur. Pat. Appl. EP 627,445 (Cl. C07K15/00)*, 07 Dec. 1994, US Appl. 61,894, 14 May 1993; 14 pp.; *C.A.*, 122 (1995) 128084k.
- 3228 Feldwisch, O., Lamertz, M., Hartmann, E., Feldwisch, J., Palme, K., Jastorff, B. and Jaenicke, L.: Purification and characterization of a cAMP-binding of *Volvox carteri* f. *nagariensis* lyengar. *Eur. J. Biochem.*, 228 (1995) 480-489.

- 3229 Greengard, J.S., Fernandez, J.A., Radtke, K.-P. and Griffin, J.H.: Identification of candidate residues for interaction of protein S with C4b binding protein and activated protein C. *Biochem. J.*, 305 (1995) 397-403.
- 3230 Gregoriou, M., Jones, P.F., Timms, J.F., Yang, J.J., Radford, S.E. and Rees, A.R.: Physicochemical characterization of the cytoplasmic domain of the epidermal growth factor receptor and evidence for conformational changes associated with its activation by ammonium sulphate. *Biochem. J.*, 306 (1995) 667-678.
- 3231 Joseph, S.K., Pierson, S. and Samanta, S.: Trypsin digestion of the inositol trisphosphate receptor: implications for the conformation and domain organization of the proteins. *Biochem. J.*, 307 (1995) 859-865.
- 3232 Karamanos, N.K., Hjerpe, A., Aletas, A., Tsegenidis, T., Anastasiou, E.D. and Antonopoulos, C.A.: Antibodies to three chondroitin sulfate-containing proteoglycans in squid skin recognize hexa- or longer chondroitin oligosaccharides as major antigenic determinants. *Arch. Biochem. Biophys.*, 316 (1995) 100-109.
- 3233 Kuo, K.-W., Chang, L.-S. and Chang, C.-C.: The structural loop II of cobrotoxin is the main binding region for nAChR and epitope in the region is conformation-dependent. *J. Biochem. (Tokyo)*, 117 (1995) 438-442.
- 3234 Langland, J.O., Pettiford, S.M. Jacobs, B.L.: Nucleic acid affinity chromatography: preparation and characterization of double-stranded RNA agarose. *Protein Expression Purif.*, 6 (1995) 25-32; *C.A.*, 122 (1995) 127807m.
- 3235 Lesca, P., Peryt, B., Larrieu, G., Alvinerie, M., Galtier, P., Daujat, M., Maurel, P. and Hoogenboom, L.: Evidence for the ligand-independent activation of the AH receptor. *Biochem. Biophys. Res. Commun.*, 209 (1995) 474-482.
- 3236 Ren, L., Chen, H. and Sternberg, E.A.: Tethered bandshift assay and affinity purification of a new DNA-binding protein. *BioTechniques*, 16 (1994) 852-855; *C.A.*, 122 (1995) 75675v.
- 3237 Sasaki, T., Shimazawa, R., Sawada, T., Iijima, T., Fukasawa, H., Shudo, K., Hashimoto, Y. and Iwasaki, S.: Determination of the photoaffinity-labeled site on the ligand-binding domain of retinoic acid receptor α . *Biochem. Biophys. Res. Commun.*, 207 (1995) 444-451.
- 3238 Tani, A., Lee, S., Oishi, O., Aoyagi, H. and Ohno, M.: Interaction of the fragments characteristic of bactenecin 7 with phospholipid bilayers and their antimicrobial activity. *J. Biochem. (Tokyo)*, 117 (1995) 560-565.
- 3239 Tomioka, H. and Sasabe, H.: Isolation of photochemically active archaeobacterial photoreceptor, pharaonis phoborhodopsin from *Natronobacterium pharaonis*. *Biochim. Biophys. Acta*, 1234 (1995) 261-267.
- 3240 Yang, Z. and Yang, Y.: (Cytokinin binding proteins from *Phaseolus vulgaris* hypocotyls). *Zhishi Xuebao*, 36 (1994) 878-883; *C.A.*, 122 (1995) 235275v.
- 3241 Yu, H. and Diamandis, E.P.: Prostate-specific antigen immunoreactivity in amniotic fluid. *Clin. Chem. (Washington)*, 41 (1995) 204-210.
- 19m. *Urinary proteins*
- 3242 Jäckle-Meyer, I., Szukics, B., Neubauer, K., Metzke, V., Petzoldt, R. and Stolte, H.: Extracellular matrix protein as early markers in diabetic nephropathy. *Eur. J. Clin. Chem. Clin. Biochem.*, 33 (1995) 211-219.
- 19n. *Other proteins (incl. proteinous inhibitors of enzymic activity)*
- 3243 Heath, R.L., Barton, P.A., Simpson, R.J., Reid, B.E., Lim, G. and Anderson, M.A.: Characterization of the protease processing sites in a multidomain proteinase inhibitor precursor from *Nicotiana glauca*. *Eur. J. Biochem.*, 230 (1995) 250-257.
- 3244 Jean, F., Basak, A., DiMaio, J., Seidah, N.G. and Lazure, C.: An internally quenched fluorogenic substrate of prohormone convertase 1 and furin leads to a potent prohormone convertase inhibitor. *Biochem. J.*, 307 (1995) 689-695.
- 3245 Levison, P.R., Badger, S.E., Toome, D.W. and Jones, R.M.H.: Strategies in the scale-up of the anion-exchange chromatography of hen egg-white proteins. *Spec. Publ. - R. Soc. Chem.*, 158 (Separations for Biotechnology 3) (1994) 267-272; *C.A.*, 122 (1995) 131366j.
- 3246 Pinto, R.A., Hawgood, S., Clements, J.A., Benson, B.J., Naidu, A., Hamilton, R.L. and Wright, J.R.: Association of surfactant protein C with isolated alveolar type II cells. *Biochim. Biophys. Acta*, 1255 (1995) 16-22.
- 3247 Ribeiro, J.M.C., Schneider, M. and Guimaraes, J.A.: Purification and characterization of prolixin S (nitrophorin 2), the salivary anticoagulant of the blood-sucking bug *Rhodnius prolixus*. *Biochem. J.*, 308 (1995) 243-249.
- 3248 Rodgers, K.J., Melrose, J. and Ghosh, P.: Biotin-labeled potato chymotrypsin inhibitor-1: a useful probe to the detection and quantitation of chymotrypsin-like serine proteinases on western blots and its application in the detection of a serine proteinase synthesized by articular chondrocytes. *Anal. Biochem.*, 227 (1995) 129-134.
- 3249 Subramanian, N. and Adiga, P.R.: Simultaneous purification of biotin-binding proteins-I and -II from chicken egg yolk and their characterization. *Biochem. J.*, 308 (1995) 573-577.

See also 3099, 3106, 3143, 3238.

20. ENZYMES AND ENZYME ACTIVITY ESTIMATION

- 3250 Rudometova, N.V., Glazova, N.V. and Dmitrenko, L.V.: (Use of sorption-chromatographic methods for purification and analysis of enzyme preparations). *Zh. Fiz. Khim.*, 68 (1994) 1848-1850; *C.A.*, 122 (1995) 54047b.

20a. Oxidoreductases

- 3251 Almira Correia, M., Yao, K., AllenToff, A.J., Wrighton, S.A. and Thompson, J.A.: Interactions of peroxyquinols with cytochromes P450 2B1, 3A1, and 3A5: influence of the apoprotein on heterolytic versus homolytic O-O bond cleavage. *Arch. Biochem. Biophys.*, 317 (1995) 471-478.

For additional information see *C.A.*:
122 (1995) 101133p.

See also 2878, 3168, 3249, 3302, 3315, 3828.

- 3252 Avezoux, A., Goodwin, M.G. and Anthony, C.: The role of the novel disulphide ring in the active site of the quinoprotein methanol dehydrogenase from *Methylobacterium extorquens*. *Biochem. J.*, 307 (1995) 735-741.
- 3253 Bagley, F.J., Hirschberger, L.L. and Stipanuk, M.H.: Evaluation and modification of an assay procedure for cysteine dioxygenase activity: high-performance liquid chromatography method for measurement of cysteine sulfinate and demonstration of physiological relevance of cysteine dioxygenase activity in cysteine catabolism. *Anal. Biochem.*, 227 (1995) 40-48.
- 3254 Bartholomew, B.A., Smith, M.J., Long, M.T., Darcy, P.J., Trudgill, P.W. and Hopper, D.J.: Tropine dehydrogenase: purification, some properties and an evaluation of its role in the bacterial metabolism of tropine. *Biochem. J.*, 307 (1995) 603-608.
- 3255 Camadro, J.-M., Matringe, M., Thome, F., Brouillet, N., Mornet, R. and Labbe, P.: Photoaffinity labeling of protoporphyrinogen oxidase, the molecular target of diphenylether-type herbicides. *Eur. J. Biochem.*, 229 (1995) 669-674.
- 3256 Camardella, L., Damonte, G., Carratore, V., Benatti, U., Tonetti, M. and Moneti, G.: Glucose 6-phosphate dehydrogenase from human erythrocytes: identification of N-acetyl-alanine at the N-terminus of the nature protein. *Biochem. Biophys. Res. Commun.*, 207 (1995) 331-338.
- 3257 Chaga, G.: A general method for immobilization of glycoproteins on regenerable immobilized metal-ion carriers: application to glucose oxidase from *Penicillium chrysogenum* and horseradish peroxidase. *Biotechnol. Appl. Biochem.*, 20 (1994) 43-53; C.A., 122 (1995) 154367z.
- 3258 De Stefano, G., Pizzuto, P. and Sciancalepore, V.: (Recovery of lactoperoxidase from acid whey by immobilized copper affinity chromatography). *Ind. Aliment. (Pinerolo)*, 33 (1994) 387-389; C.A., 122 (1995) 49731r.
- 3259 Emanuele, J.J., Heasley, C.J. and Fitzpatrick, P.F.: Purification and characterization of the flavoprotein tryptophan 2-monooxygenase expressed at high levels in *Escherichia coli*. *Arch. Biochem. Biophys.*, 31'6 (1995) 241-248.
- 3260 Gillam, E.M.J., Guo, Z., Ueng, Y.-F., Yamazaki, H., Cock, I., Reilly, P.E.B., Hooper, W.D. and Guengerich, F.P.: Expression of cytochrome P450 3A5 in *Escherichia coli*: effects of 5'-modification, purification, spectral characterization, reconstitution conditions, and catalytic activities. *Arch. Biochem. Biophys.*, 317 (1995) 374-384.
- 3261 Hallahan, D.L., West, J.M., Wallsgrave, R.M., Smiley, D.W.M., Dawson, G.W., Pickett, J.A. and Hamilton, J.G.C.: Purification and characterization of an acyclic monoterpene primary alcohol:NADP oxidoreductase from catmint (*Nepeta racemosa*). *Arch. Biochem. Biophys.*, 318 (1995) 105-112.
- 3262 Hiromasa, Y., Aso, Y., Yamashita, S. and Aikawa, Y.: Homogeneity of the pyruvate dehydrogenase multienzyme complex from *Bacillus stearothermophilus*. *J. Biochem. (Tokyo)*, 117 (1995) 467-470.
- 3263 Huang, Y.-C., Haselbeck, R.J., McAlister-Henn, L. and Colman, R.F.: N-Ethylmaleimide profiling of yeast NADP-dependent isocitrate dehydrogenase. *Arch. Biochem. Biophys.*, 316 (1995) 485-492.
- 3264 Il'chenko, A.P., Morgunov, I.G., Honeck, H., Mauersberger, S., Vasilkova, N.N. and Müller, H.-G.: (Purification and some properties of alcohol oxidase from the yeast *Yarrowia lipolytica* H-222). *Biokhimiya (Moscow)*, 59 (1994) 1312-1319.
- 3265 Jansson, I., Mole, J.E. and Schenkman, J.B.: The isolation and comparison of multiple forms of CYP-2B from untreated and phenobarbital-treated rabbit liver microsomes. *Arch. Biochem. Biophys.*, 316 (1995) 275-284.
- 3266 Kabzinski, A.K.N. and Paryjzak, T.: The preparation of enzymes electrodes with application of the fast immobilization of enzymes of affinity chromatography supports. *Eur. J. Clin. Chem. Clin. Biochem.*, 33 (1995) A22.
- 3267 Koga, N., Kikuichi-Nishimura, N., Hara, T., Harada, N., Ishii, Y., Yamada, H., Oguri, K. and Yoshimura, H.: Purification and characterization of a newly identified isoform of cytochrome P450 responsible for 3-hydroxylation of 2,5,2',5'-tetrachlorobiphenyl in hamster liver. *Arch. Biochem. Biophys.*, 317 (1995) 464-470.
- 3268 Lomnitski, L., Sklan, D. and Grossman, S.: Lipoxygenase activity in rat dermis and epidermis: partial purification and characterization. *Biochim. Biophys. Acta*, 1255 (1995) 351-359.
- 3269 Maier, T., Lottspeich, F. and Böck, A.: GTP hydrolysis by HypB is essential for nickel insertion into hydrogenases of *Escherichia coli*. *Eur. J. Biochem.*, 230 (1995) 133-138.
- 3270 Martinez, A., Knappskog, P.M., Olafsdottir, S., Doskeland, A.P., Eiken, H.G., Svebak, R.M., Bozzini, M.L., Apold, J. and Flatmark, T.: Expression of recombinant human phenylalanine hydroxylase as fusion protein in *Escherichia coli* circumvents proteolytic degradation by host cell proteases. Isolation and characterization of the wild-type enzyme. *Biochem. J.*, 306 (1995) 589-597.
- 3271 Notheis, C., Drewke, C. and Leistner, E.: Purification and characterization of the pyridoxol-5'-phosphate:oxygen oxidoreductase (deaminating) from *Escherichia coli*. *Biochim. Biophys. Acta*, 1247 (1995) 265-271.
- 3272 Ōba, K., Ishikawa, S., Nishikawa, M., Mizuno, H. and Yamamoto, T.: Purification and properties of L-galactono-γ-lactone dehydrogenase, a key enzyme for ascorbic acid biosynthesis, from sweet potato roots. *J. Biochem. (Tokyo)*, 117 (1995) 120-124.
- 3273 Obata, T., Shibusawa, Y. and Ito, Y.: Purification of soluble monoamine oxidase in rat liver cytosol by countercurrent chromatography. *Chromatographia*, 39 (1994) 624-626.
- 3274 Padiglia, A., Medda, R., Pazzaglia, G., Rescigno, A., Cruciani, E. and Floris, G.: Fractionation and characterization of two forms of peroxidase from *Oryza sativa*. *Prepar. Biochem.*, 25 (1995) 11-19.
- 3275 Peters, D., Frank, R. and Hengstenberg, W.: Lactose-specific enzyme II of the phosphoenolpyruvate-dependent phosphotransferase system of *Staphylococcus aureus*. Purification of the histidine-tagged transmembrane component IICB^{lac} and its hydrophilic IIB domain by metal-affinity chromatography, and functional characterization. *Eur. J. Biochem.*, 228 (1995) 798-804.
- 3276 Rescigno, A., Sollai, F., Masala, S., Porcu, M.C., Sanjust, E., Rinaldi, A.C., Curreli, N., Grifi, D. and Rinaldi, A.: Purification and characterization of an NAD(P)H:quinone oxidoreductase from *Glycine max* seedlings. *Prepar. Biochem.*, 25 (1995) 57-67.
- 3277 Riveros-Moreno, V., Heffernan, B., Torres, B., Chubb, A., Charles, I. and Moncada, S.: Purification to homogeneity and characterisation of rat brain recombinant nitric oxide synthase. *Eur. J. Biochem.*, 230 (1995) 52-57.

- 3278 Russell, R. and Scopes, R.K.: Use of hydrophobic chromatography for purification of the membrane-located choline dehydrogenase from a *Pseudomonas* strain. *Bioseparation*, 4 (1994) 279-284; *C.A.*, 122 (1995) 127113g.
- 3279 Schilling, B. and Lerch, K.: Amine oxidases from *Aspergillus niger*: identification of a novel flavin-dependent enzyme. *Biochim. Biophys. Acta*, 1243 (1995) 529-537.
- 3280 Schoch, S., Helfrich, M., Wiktorsson, B., Sundqvist, C., Rüdiger, W. and Ryberg, M.: Photoreduction of zinc protoporphorbide with NADPH-protochlorophyllide oxidoreductase from etiolated wheat (*Triticum aestivum* L.) *Eur. J. Biochem.*, 229 (1995) 291-298.
- 3281 Sharma, M.C. and Shapiro, B.H.: Purification and characterization of constituent testosterone 2 α -hydroxylase (cytochrome P450_{2a}) from mouse liver. *Arch. Biochem. Biophys.*, 316 (1995) 478-484.
- 3282 Simons, B.H., Barnett, P., Vollenbroek, E.G.M., Dekker, H.L., Muijsers, A.O., Messerschmidt, A. and Wever, R.: Primary structure and characterization of the vanadium chloroperoxidase from the fungus *Curvularia inaequalis*. *Biochem. J.*, 229 (1995) 566-574.
- 3283 Sinhababu, A.K., Boehlert, C.C., Gan, L.-S., Yanni, S.B. and Thakker, D.R.: High-performance liquid chromatographic purification, optimization of the assay, and properties of ribonucleoside diphosphate reductase from rabbit bone marrow. *Arch. Biochem. Biophys.*, 317 (1995) 285-291.
- 3284 Soukri, A., Valverde, F., Hafid, N., Elkebbaj, M.S. and Serrano, A.: Characterization of muscle glyceraldehyde-3-phosphate dehydrogenase isoforms from euthermic and induced hibernating *Jaculus orientalis*. *Biochim. Biophys. Acta*, 1243 (1995) 161-168.
- 3285 Swarts, J.C., Aquino, M.A.S., Han, J.-Y., Lam, K.-Y. and Sykes, A.G.: Kinetic studies on the reduction of the tyrosyl radical of the R2 subunit of *E. coli* ribonucleotide reductase. *Biochim. Biophys. Acta*, 1247 (1995) 215-224.
- 3286 Tachibana, M., Onoe, H., Okubo, A. and Yamazaki, S.: Application of micellar electrokinetic chromatography as a novel assay method for dimethyl sulfoxide reductase. *Biosci., Biotechnol., Biochem.*, 59 (1995) 282-284; *C.A.*, 122 (1995) 208205r.
- 3287 Takeuchi, A., Miyamoto, T., Yamaji, K., Masuho, Y., Hayashi, M., Hayashi, H. and Onozaki, K.: A human erythrocyte-derived growth-promoting factor with a wide target cell spectrum: identification as catalase. *Cancer Res.*, 55 (1995) 1586-1589.
- 3288 Wan, L., Gijzen, M. and van Huystee, R.S.: Heterogeneous glycosylation of cationic peanut peroxidase. *Biochem. Cell Biol.*, 72 (1994) 411-417.
- 3289 Wang, H.-t., Liu, W.-y. and Ulbrich, N.: Isolation and characterization of a tyrosinase from the skin of the white silky fowl (*Gallina lanigera*) employing copper saturated diethylaminoethyl-cellulose. *Biochim. Biophys. Acta*, 1243 (1995) 251-255.
- For additional information see *C.A.*:
122 (1995) 100181r, 127125n.
- See also 3104, 3193.
- 20b. *Transferases (excl. E.C. 2.7.-.)*
- 3290 Bico, P., Erhardt, J., Kaplan, W. and Dirr, H.: Porcine class π glutathione S-transferase: anionic ligand binding and conformational analysis. *Biochim. Biophys. Acta*, 1247 (1995) 225-230.
- 3291 Deigner, H.P., Mato, J.M. and Pajares, M.A.: Study of the rat liver S-adenosylmethionine synthetase active site with 8-azido ATP. *Biochem. J.*, 308 (1995) 565-571.
- 3292 Egaas, E., Falls, J.G., Svendsen, N.O., Ramstad, H., Skaare, J.U. and Dauterman, W.C.: Strain- and sex-specific differences in the glutathione S-transferase class pi in the mouse examined by gradient elution of the glutathione-affinity matrix and reverse-phase high performance liquid chromatography. *Biochim. Biophys. Acta*, 1243 (1995) 256-264.
- 3293 Habuchi, H., Habuchi, O. and Kimata, K.: Purification and characterization of heparan sulfate 6-sulfotransferase from the culture medium of chinese hamster ovary cells. *J. Biol. Chem.*, 270 (1995) 4172-4179.
- 3294 Herold, M., Grimm, R., Heiger, D.N., Bayerbach, S., Fischer, S. and Miller, C.E.: Purification and characterization of aspartate aminotransferase from *Escherichia coli* by chromatographic methods. *Int. J. Bio-Chromatogr.*, 1 (1994) 29-45; *C.A.*, 122 (1995) 154858k.
- 3295 Liu, Y., van Heeswijk, R., Hoj, P. and Hoogenraad, N.: Purification and characterization of ornithine acetyltransferase from *Saccharomyces cerevisiae*. *Eur. J. Biochem.*, 228 (1995) 291-296.
- 3296 Palmerini, C.A., Datti, A., Alunni, S., Vabder Elst, I.E. and Orlandino, A.: A fluorescent assay for the determination of UDPGlcNAc: Gal β 1,3GalNAc-R(GlcNAc to GalNAc) β 1,6-N-acetylglucosaminyltransferase activity. *Anal. Biochem.*, 225 (1995) 315-320.
- 3297 Slany, R.K. and Müller, S.O.: tRNA-guanine transglycosylase from bovine liver. Purification of the enzyme to homogeneity and biochemical characterization. *Eur. J. Biochem.*, 230 (1995) 221-228.
- 3298 Stadie, T.R.E., Chai, W., Lawson, A.M., Byfield, P.G.H. and Hanisch, F.-G.: Studies on the order and site specificity of GalNAc transfer to MUC1 tandem repeats by UDP-GalNAc: polypeptide N-acetylgalactosaminyltransferase from milk or mammary carcinoma cells. *Eur. J. Biochem.*, 229 (1995) 140-147.
- 3299 Taniguchi, M., Kobe, A., Kato, M. and Sugiyama, T.: Aspartate aminotransferase isozymes in *Panicum miliaceum* L., an NAD-malic enzyme-type C₄ plant: comparisons of enzymatic properties, primary structures, and expression patterns. *Arch. Biochem. Biophys.*, 318 (1995) 295-306.
- 3300 Tsvetnitsky, V., Auchli, L., Nicolaou, A. and Gibbons, W.A.: Characterization of phospholipid methylation in rat brain myelin. *Biochem. J.*, 307 (1995) 239-244.
- 3301 Varone, C.L., Canépa, E.T., Llambías, E.B.C. and Grinstein, M.: cAMP regulation of phenobarbital-mediated induction of δ -aminolevulinic synthase mRNA in hepatocytes from normal and experimental diabetic rats. *Biochem. Cell Biol.*, 72 (1994) 381-390.
- 3302 Weifold, P.A. and Feldman, D.A.: Fatty acids promote the formation of complexes between choline-phosphate cytidylyltransferase and cytidylyltransferase binding protein. *Arch. Biochem. Biophys.*, 318 (1995) 147-156.

- 3303 Yamamoto, T., Moriwaki, Y., Takahashi, S., Nasako, Y., Yamakita, J.-i., Hiroishi, K. and Higashino, K.: Determination of plasma purine nucleosides phosphorylase activity by high-performance liquid chromatography. *Anal. Biochem.*, 227 (1995) 135-139.
- 3304 Yeh, H.-I., Hsieh, C.-H., Wang, L.-Y., Tsai, S.-P., Hsu, H.-Y. and Tam, M.F.: Mass spectrometric analysis of rat liver cytosolic glutathione S-transferases: modifications are limited to N-terminal processing. *Biochem. J.*, 308 (1995) 69-75.
- See also 3420.
- 20c. *Transferases transferring phosphorus containing groups (E.C. 2.7.-.-)*
- 3305 Asano, Y., Mizumoto, K., Maruyama, T. and Ishihama, A.: Photoaffinity labeling of influenza virus RNA polymerase PB1 subunit with 8-azido GTP. *J. Biochem. (Tokyo)*, 117 (1995) 677-682.
- 3306 Bergwerff, A.A., van Oostrum, J., Kamerling, J.P. and Vliegthart, J.F.G.: The major N-linked carbohydrate chains from human urokinase. The occurrence of 4-O-sulfated, (α 2-6)-sialylated or (α 1-3)-fucosylated N-acetylgalactosamine(β 1-4)-N-acetylglucosamine elements. *Eur. J. Biochem.*, 228 (1995) 1009-1019.
- 3307 Brunati, A.M., Ruzzene, M., James, P., Guerra, B. and Pinna, L.A.: Hierarchical phosphorylation of a 50-kDa protein by protein tyrosine kinases TPK-IIb and C-Fgr, and its identification as hematopoietic-lineage cell-specific protein. *Eur. J. Biochem.*, 229 (1995) 164-170.
- 3308 Chernov, A.V., Matvienko, N.N., Zheleznyaya, L.A. and Matvienko, N.I.: (A new site-specific endonuclease-methylase from the thermophilic strain of *Bacillus* species LU11). *Biokhimiya (Moscow)*, 59 (1994) 1714-1729.
- 3309 Garg, N., Galev, I.Yu. and Mattiasson, B.: Effect of poly(vinyl alcohol) treatment of the dye-matrix on the chromatography of pyruvate kinase. *Biotechnol. Tech.*, 8 (1994) 645-650; *C.A.*, 122 (1995) 233947s.
- 3310 Hsuan, J.J., Totty, N.F., Truong, O. and Waterfield, M.D.: Purification of phosphatidylinositol 3-kinase by SDA-agarose gel electrophoresis and HPEC. *Protein Pept. Lett.*, 1 (1994) 25-32; *C.A.*, 122 (1995) 154733r.
- 3311 Ipata, P.L., Gualerzi, C., Scolozzi, C., Tozzi, M.G., Trinei, M. and Barsacchi, D.: Occurrence of inosine kinase as a distinct enzyme in *Spirulina platensis*. *Biochem. Biophys. Res. Commun.*, 209 (1995) 547-553.
- 3312 Joshi, B., Morley, S.J., Rgoads, R.E. and Pain, V.M.: Inhibition of protein synthesis by the heme-controlled eIF-2 α kinase leads to the appearance of mRNA-containing 48S complexes that contain eIF-4E but lack methionyl-tRNA_f. *Eur. J. Biochem.*, 228 (1995) 31-38.
- 3313 Lecocq, R., Lamy, F., Erneux, C. and Dumont, J.E.: Rapid purification and identification of calyphosine, a Ca²⁺-binding protein phosphorylated by protein kinase A. *Biochem. J.*, 306 (1995) 147-151.
- 3314 Price, D.J., Kawakami, Y., Kawakami, T. and Rivnay, B.: Purification of a major tyrosine kinase from RBL-2H3 cells phosphorylating Fc ϵ RI γ -cytoplasmic domain and identification as the Btk tyrosine kinase. *Biochim. Biophys. Acta*, 1265 (1995) 135-142.
- 3315 Seki, K., Chen, H.-C. and Huang, K.-P.: Dephosphorylation of protein kinase C substrates, neurogranin, neuromodulin, and MARCKS, by calcineurin and protein phosphatases 1 and 2A. *Arch. Biochem. Biophys.*, 316 (1995) 673-679.
- 3316 Shapovalova, N.I., Zheleznyaya, L.A. and Matvienko, N.I.: (A new site-specific endonuclease and methylase from the thermophilic strain of *Bacillus* species KT6). *Biokhimiya (Moscow)*, 59 (1994) 1730-1738.
- 3317 Stathopoulos, C., Kalpaxis, D.L. and Drains, D.: Partial purification and characterization of RNase P from *Dictyostelium discoideum*. *Eur. J. Biochem.*, 228 (1995) 976-980.
- 3318 Zhang, J., Fukui, T. and Ichikawa, A.: A third type of nucleoside diphosphate kinase from spinach leaves: purification, characterization and amino-acid sequence. *Biochim. Biophys. Acta*, 1248 (1995) 19-26.
- See also 3180.
- 20d. *Hydrolases, acting on ester bonds (E.C. 3.1.-.-)*
- 3319 Basu, P.S., Batabyal, S.K., Bhattacharya, A. and Datta, T.K.: Cholinesterase activities in cerebrospinal fluid of patients with idiopathic convulsive disorders. *Clin. Chim. Acta*, 235 (1995) 107-112.
- 3320 Beil, S., Kehrl, H., James, P., Staudenmann, W., Cook, A.M., Leisinger, T. and Kertesz, M.A.: Purification and characterization of the arylsulfatase synthesized by *Pseudomonas aeruginosa* PAO during growth in sulfate-free medium and cloning of the arylsulfatase gene (atsA). *Eur. J. Biochem.*, 229 (1995) 385-394.
- 3321 Cal, S., Aparicio, J.F., de Los Reyes-Gavilan, C.G., Nicieza, R.G. and Sanchez, J.: A novel exocytosolic endonuclease from *Streptomyces antibioticus*. *Biochem. J.*, 306 (1995) 93-100.
- 3322 Cernia, E., Ortaggi, G., Soro, S. and Castagnola, M.: Esters of crosslinked poly(vinyl alcohol) with fatty acids: new stationary phases for the separation of lipolytic enzymes. *Tetrahedron Lett.*, 35 (1994) 9051-9054; *C.A.*, 122 (1995) 181711p.
- 3323 Craxton, A., Ali, N. and Shears, S.B.: Comparison of the activities of a multiple inositol polyphosphate phosphatase obtained from several sources: a search for heterogeneity in this enzyme. *Biochem. J.*, 305 (1995) 491-498.
- 3324 Gijon, M.A., Perez, C., Mendez, E. and Sanchez Crespo, M.: Phospholipase A₂ from plasma of patients with septic shock is associated with high-density lipoproteins and C3 anaphylatoxin: some implications for its functional role. *Biochem. J.*, 306 (1995) 167-175.
- 3325 Johansson, F., Sommarin, M. and Larsson, C.: Rapid purification of the plasma membrane H⁺-ATPase in its non-activated form using FPLC. *Physiol. Plant.*, 92 (1994) 389-396; *C.A.*, 122 (1995) 49754a.
- 3326 Koblan, K.S., Schaber, M.D., Edwards, G., Gibbs, J.B. and Pompiano, D.L.: src-Homology 2 (SH2) domain ligation as an allosteric regulator: modulation of phosphoinositide-specific phospholipase C γ 1 structure and activity. *Biochem. J.*, 305 (1995) 745-751.

- 3327 Larnkjaer, A., Nykjaer, A., Olivecrona, G., Thogersen, H. and Ostergaard, P.B.: Structure of heparin fragments with high affinity for lipoprotein lipase and inhibition of lipoprotein lipase binding to α_2 -macroglobulin-receptor/low density lipoprotein-receptor-related protein by heparin fragments. *Biochem. J.*, 307 (1995) 205-214.
- 3328 Lindberg, A. and Olivecrona, G.: Lipase evolution: trout, *Xenopus* and chicken have lipoprotein lipase and apolipoprotein C-II-like activity but lack hepatic lipase-like activity. *Biochim. Biophys. Acta*, 1255 (1995) 205-211.
- 3329 McIntosh, J.M., Ghomashchi, F., Gelb, M.H., Dooley, D.J., Stoehr, S.J., Giordani, A.B., Naisbitt, S.R. and Olivera, B.M.: Conodipine-M, a novel phospholipase A₂ isolated from the venom of the marine snail *Conus magus*. *J. Biol. Chem.*, 270 (1995) 3518-3526.
- 3330 Mehrani, H. and Storey, K.B.: Effects of anoxia on protein phosphatase in turtle organs: purification and properties of protein phosphatase type-1 from turtle liver. *Arch. Biochem. Biophys.*, 316 (1995) 836-843.
- 3331 Mingarro, I., González-Navarro, H. and Braco, L.: Direct HPLC monitoring of lipase activity in reverse micellar media. *J. Liq. Chromatogr.*, 18 (1995) 235-244.
- 3332 Moorhead, G., MacKintosh, R.W., Morrice, N., Gallagher, T. and MacKintosh, C.: Purification of type 1 protein (serine/threonine) phosphatases by microcystin-Sepharose affinity chromatography. *FEBS Lett.*, 356 (1994) 46-50; *C.A.*, 122 (1995) 181618p.
- 3333 Pedersen, J., Andersen, J., Roepstorff, P., Filimonova, M. and Biedermann, K.: (Characterization of *Serrata marcescens* nuclease isoforms by electrospray mass spectrometry). *Biokhimiya (Moscow)*, 160 (1995) 462-469.
- 3334 Pozidis, C., Vlatakis, G. and Bouriotis, V.: Sequence-specific DNA affinity chromatography: application of a group specific adsorbent for the isolation of restriction endonucleases. *Prog. Biotechnol.*, 9(ECB6: Proceedings of the 6th European Congress on Biotechnology, 1993, Pt. 1) (1994) 543-546; *C.A.*, 122 (1995) 127098f.
- 3335 Recktenwald, A., Vernet, T., Storer, A.C. and Ziomek, E.: Reduction of strong lipase-polyclonal antibodies binding by limited proteolysis. *Anal. Biochem.*, 226 (1995) 31-34.
- 3336 Rugani, N., Carrière, F., Thim, L., Borgstrom, B. and Sarda, L.: Lipid binding and activating properties of porcine pancreatic colipase split at the Ile⁷⁹-Thr⁸⁰ bond. *Biochim. Biophys. Acta*, 1247 (1995) 185-194.
- 3337 Sunaga, H., Sugimoto, H., Nagamachi, Y. and Yamashita, S.: Purification and properties of lysophospholipase isoenzymes from pig gastric mucosa. *Biochem. J.*, 308 (1995) 551-557.
- 3338 Talesa, V., Grauso, M., Giovannini, E., Rosi, G. and Toutant, J.-P.: Solubilization, molecular forms, purification and substrate specificity of two acetylcholinesterases in the medicinal leech (*Hirudo medicinalis*). *Biochem. J.*, 306 (1995) 687-692.
- 3339 Teissère, M., Borel, M., Caillol, B., Nari, J., Gardies, A.M. and Noat, G.: Purification and characterization of a fatty acyl-ester hydrolase from post-germinated sunflower seeds. *Biochim. Biophys. Acta*, 1255 (1995) 105-112.
- 3340 Thomson, F.J. and Clark, M.A.: Purification of a phosphatidic-acid-hydrolysing phospholipase A₂ from rat brain. *Biochem. J.*, 306 (1995) 305-309.
- 3341 Witmer, M.R., Micanovic, R., Tredup, J., Lin, E., Hail, M. and Villafranca, J.J.: Active recombinant human cytosolic phospholipase A₂ is expressed in *Escherichia coli*. *Arch. Biochem. Biophys.*, 318 (1995) 430-438.
- 3342 Yamazaki, Y., Ogawa, Y., Afify, A.S., Kageyama, Y.-i., Okada, T., Okuno, H., Yoshii, Y. and Nose, T.: Difference between cancer cells and the corresponding normal tissue in view of stereoselective hydrolysis of synthetic esters. *Biochim. Biophys. Acta*, 1243 (1995) 300-308.
- See also 3092, 3093, 3103, 3275.
- 20e. *Hydrolases, acting on glycosyl compounds (E.C. 3.2.-)*
- 3343 Ara, K., Saeki, K., Igarashi, K., Takaiwa, M., Uemura, T., Hagi-hara, H., Kawai, S. and Ito, S.: Purification and characterization of an alkaline amylopullulanase with both α -1,4 and α -1,6 hydrolytic activity from alkalophilic *Bacillus* sp. KSM-1378. *Biochim. Biophys. Acta*, 1243 (1995) 315-324.
- 3344 Blaak, H. and Schrempf, H.: Binding and substrate specificities of a *Streptomyces olivaceoviridis* chitinase in comparison with its proteolytically processed form. *Eur. J. Biochem.*, 229 (1995) 132-139.
- 3345 Braithwaite, K.L., Black, G.W., Hazlewood, G.P., Ali, B.R.S. and Gilbert, H.J.: A non-modular endo- β -1,4-mannanase from *Pseudomonas fluorescens* subspecies *cellulosa*. *Biochem. J.*, 305 (1995) 1005-1010.
- 3346 Christakopoulos, P., Kekos, D., Macris, B.J., Claeysens, M. and Bhat, M.K.: Purification and characterization of a less randomly acting endo-1,4- β -D-glucanase from the culture filtrates of *Fusarium oxysporum*. *Arch. Biochem. Biophys.*, 316 (1995) 428-433.
- 3347 Gottvaldova, M., Hrabova, H. and Kucera, J.: (Amylase isolation by affinity chromatography on immobilized lectin of pea (*Pisum sativum* L.) saturated with beta-cyclodextrin). *Potravin. Vedy*, 12 (1994) 137-142; *C.A.*, 122 (1995) 100171n.
- 3348 Kato, S. and Terashima, M.: Purification of secreted α -amylases by immunoaffinity chromatography with cross-reactive antibody. *Appl. Microbiol. Biotechnol.*, 42 (1994) 36-39; *C.A.*, 122 (1995) 181674d.
- 3349 Kucera, J.: (Tandem column bioaffinity separation of amylases, agglutinin and chymotrypsin inhibitor of rye). *Potravin. Vedy*, 12 (1994) 339-350; *C.A.*, 122 (1995) 104189x.
- 3350 Martino, S., Emiliani, C., Orlacchio, A., Hosseini, R. and Stirling, J.L.: β -N-Acetylhexosaminidases A and S have similar sub-cellular distributions in HL-60 cells. *Biochim. Biophys. Acta*, 1243 (1995) 489-495.
- 3351 Moriyama, T., Tozawa, T., Nobuoka, M. and Ikeda, H.: Sialyl salivary-type amylosemia associated with immunoglobulin D-type multiple myeloma. *Clin. Chim. Acta*, 233 (1995) 127-134.
- 3352 Ribeiro, M.G., Pinto, R., Sa Miranda, M.C. and Suzuki, K.: Tay-Sachs disease: intron 7 splice junction mutation in two Portuguese patients. *Biochim. Biophys. Acta*, 1270 (1995) 44-51.
- 3353 Sova, V.V., Elyakova, L.A., Ivanova, E.P., Fedosov, Yu.V. and Mikhailov, V.V.: (Induced β -1,3-glucanase from the marine bacterium *Alteromonas* sp.) *Biokhimiya (Moscow)*, 59 (1994) 1369-1377.
- 3354 Tomizawa, H., Yamada, H., Tanigawa, K. and Imoto, T.: Effects of additives on irreversible inactivation of lysozyme at neutral pH and 100° C. *J. Biochem. (Tokyo)*, 117 (1995) 369-373.

- 3355 Vandersall, A.S., Cameron, R.G., Nairn C.J., III, Yelenosky, G. and Wodzinski, R.J.: Identification, characterization, and partial purification of glucoamylase from *Aspergillus niger* (syn. *A. ficum*) NRRL 3135. *Prepar. Biochem.*, 25 (1995) 29-55.
- 3356 Wood, P.J., Erfle, J.D., Teather, R.M., Weisz, J. and Miller, S.S.: Comparison of (1→3)(1→4)-β-D-glucan-4-glucanohydrolases (E.C. 3.2.1.73) from *Fibrobacter succinogenes* and from *Bacillus subtilis* and use of high-performance anion exchange chromatography in product characterization. *J. Cereal Sci.*, 19 (1994) 65-75; *C.A.*, 122 (1995) 208354p.
- See also 2569, 3371.
- 20f. *Other hydrolases*
- 3357 Akasaki, K., Yoshimoto, H., Nakamura, A., Shiomi, H. and Tsuji, H.: Purification and characterization of a major kyotorphin-hydrolyzing peptidase of rat brain. *J. Biochem. (Tokyo)*, 117 (1995) 897-902.
- 3358 Ardo, Y. and Jonsson, L.: Chromatographic profiling of aminopeptidolytic activities in lactobacilli as a tool for strain identification. *J. Dairy Res.*, 61 (1994) 573-579; *C.A.*, 122 (1995) 29988b.
- 3359 Becker, C., Shutov, A.D., Nong, V.H., Senyuk, V.I., Jung, R., Horstmann, C., Eisner, J., Nielsen, N.C. and Müntz, K.: Purification, cDNA cloning and characterization of proteinase B, an asparagine-specific enolpeptidase from germinating vetch (*Vicia sativa* L.) seeds. *Eur. J. Biochem.*, 228 (1995) 456-462.
- 3360 Belling, H.F., Steghens, J.P. and Collombel, C.: Firefly luciferase purification using polyethylene glycol and Dyematrix Orange A. *J. Chromatogr. A*, 695 (1995) 33-40.
- 3361 Blée, E. and Schuber, F.: Stereocontrolled hydrolysis of the linoleic acid monoepoxide regioisomers catalyzed by soybean epoxide hydrolase. *Eur. J. Biochem.*, 230 (1995) 229-234.
- 3362 Buttle, D.J.: Affinity chromatography of cysteine peptidases. *Methods Enzymol.*, 244 (Proteolytic Enzymes: Serine and Cysteine Peptidases) (1994) 639-648; *C.A.*, 122 (1995) 181495w.
- 3363 Chiba, Y., Midorikawa, T. and Ichishima, E.: Cloning and expression of the carboxypeptidase gene from *Aspergillus saitoi* and determination of the catalytic residues by site-directed mutagenesis. *Biochem. J.*, 308 (1995) 405-409.
- 3364 Clark, I.M., Mitchell, R.E., Posell, L.K., Bigg, H.F., Cawston, T.E. and O'Hare, M.C.: Recombinant porcine collagenase: purification and autolysis. *Arch. Biochem. Biophys.*, 316 (1995) 123-127.
- 3365 De Saizieu, A., Vankan, P. and van Loon, A.P.G.M.: Enzymic characterization of *Bacillus subtilis* GTP cyclohydrolase I. Evidence for a chemical dephosphorylation of dihydroneopterin triphosphate. *Biochem. J.*, 306 (1995) 371-377.
- 3366 Deperthes, D., Gauthier, E.R., Chapdelaine, P., Lazure, C., Tremblay, R.R. and Dubé, J.Y.: Identification of glandular kallikrein in dog pancreas and determination of its tissue distribution. *Biochim. Biophys. Acta*, 1243 (1995) 291-294.
- 3367 Diefenbach-Jagger, H., Brenner, C., Kemp, B.E., Baron, W., McLean, J., Martin, T.J. and Moseley, J.M.: Arg21 is the preferred kexin-cleavage site in parathyroid-hormone-related protein. *Eur. J. Biochem.*, 229 (1995) 91-98.
- 3368 Ding, Z. and Xu, Y.: A rapid sensitive purification of bovine thrombin. *Prepar. Biochem.*, 25 (1995) 21-28.
- 3369 Fujimura, S., Rikimaru, T., Baba, S., Hori, J., Hao, X.-Q., Terada, S. and Kimoto, E.: Purification and characterization of a non-hemorrhagic metalloprotease from *Agkistrodon halys brevicaudus* venom. *Biochim. Biophys. Acta*, 1243 (1995) 94-100.
- 3370 Gary, J.D. and Clarke, S.: Purification and characterization of an isoaspartyl dipeptidase from *Escherichia coli*. *J. Biol. Chem.*, 270 (1995) 4076-4087.
- 3371 Gey, M.H. and Unger, K.K.: Calculation of the molecular masses of two newly synthesized thermostable enzymes isolated from thermophilic microorganisms. *J. Chromatogr. B*, 666 (1995) 188-193.
- 3372 Grigoleit, V., Möder, M. and Reissmann, S.: Studies on pseudo-peptidic inhibitors of renin as a potential antihypertensives by HPLC and mass spectrometry. *Eur. J. Clin. Chem. Clin. Biochem.*, 33 (1995) A25.
- 3373 Hashimoto, T., Yamamoto, Y., Yoshida, Y. and Tagawa, K.: Cleavage of bovine mitochondrial ATPase inhibitor with endopeptidases, and binding of the resulting peptides to the interface between the α- and β-subunits of F₁ATPase. *J. Biochem. (Tokyo)*, 117 (1995) 641-647.
- 3374 Hirose, T., Yamasaki, K. and Yamamoto, T.: Irradiation with ultraviolet light in the presence of vanadate increases Ca²⁺ permeability of the sarcoplasmic reticulum membrane via Ca²⁺-ATPase. *J. Biochem. (Tokyo)*, 117 (1995) 324-330.
- 3375 Inaba, K. and Morisawa, M.: Identification of a 10S trypsin-like protease that cross-reacts with anti-proteasome antibody in sea urchin egg jelly. *Arch. Biochem. Biophys.*, 319 (1995) 177-184.
- 3376 Itoh, C. and Nagamatsu, A.: An aminopeptidase activity from porcine kidney that hydrolyzes oxytocin and vasopressin: purification and partial characterization. *Biochim. Biophys. Acta*, 1243 (1995) 203-208.
- 3377 Itoh, Y., Binner, S. and Nagase, H.: Steps involved in activation of the complex of pro-matrix metalloproteinase 2 (progelatinase A) and tissue inhibitor of metalloproteinases (TIMP)-2 and 4-aminophenylmercuric acetate. *Biochem. J.*, 308 (1995) 645-651.
- 3378 Katsumi, Y., Kihara, H., Ochiai, M. and Ashida, M.: A serine protease zymogen in insect plasma. Purification and activation by microbial cell wall components. *Eur. J. Biochem.*, 228 (1995) 870-877.
- 3379 Kim, Y.-T., Muramatsu, T. and Takahashi, K.: Leader peptidase from *Escherichia coli*: overexpression, characterization, and inactivation by modification of tryptophan residues 300 and 310 with N-bromosuccinimide. *J. Biochem. (Tokyo)*, 117 (1995) 535-544.
- 3380 Mäkinen, K.K., Mäkinen, P.-L., Loesche, W.J. and Syed, S.A.: Purification and general properties of an oligopeptidase from *Treponema denticola* ATCC 35405. A human oral spirochete. *Arch. Biochem. Biophys.*, 316 (1995) 689-698.
- 3381 Mars, I. and Monnet, V.: An aminopeptidase P from *Lactococcus lactis* with original specificity. *Biochim. Biophys. Acta*, 1243 (1995) 209-215.
- 3382 Meyer, W., Moll, R., Kath, T. and Schäfer, G.: Purification, cloning, and sequencing of archaeobacterial pyrophosphatase from the extreme thermoacidophile *Sulfolobus acidocaldarius*. *Arch. Biochem. Biophys.*, 319 (1995) 149-156.
- 3383 Moser, T.L., Enghild, J.J., Pizzo, S.V. and Stack, M.S.: Specific binding of urinary-type plasminogen activator (u-PA) to vitronectin and its role in mediating u-PA-dependent adhesion of U937 cells. *Biochem. J.*, 307 (1995) 867-873.

- 3384 Proudfoot, A.E.I., Magneat, E., Haley, T.M., Maione, T.E. and Wells, T.N.C.: The complete primary structure of glycosylated porcine platelet factor 4. *Eur. J. Biochem.*, 228 (1995) 658-664.
- 3385 Stewart, A.J., Piggott, N.H., May, F.E.B. and Westley, B.R.: Mitogenic activity of procathepsin D purified from conditioned medium of breast-cancer cells by affinity chromatography on pepstatinyl agarose. *Int. J. Cancer*, 57 (1994) 715-718; *C.A.*, 122 (1995) 77980q.
- 3386 Thomas, M.P., Verma, C., Boyd, S.M. and Brocklehurst, K.: The structural origins of the unusual specificities observed in the isolation of chymopapain M and actinidin by covalent chromatography and the lack of inhibition of chymopapain M by cystatin. *Biochem. J.*, 306 (1995) 39-46.
- 3387 Tyagi, S.C. and Simon, S.R.: Elastin-derived peptide binding to a hydrophobic domain on neutrophil elastase. *Biochem. Cell Biol.*, 72 (1994) 419-427.
- 3388 Vergas Romero, C., Neudorfer, I., Mann, K. and Schäfer, W.: Purification and amino acid sequence of aminopeptidase P from pig kidney. *Eur. J. Biochem.*, 229 (1995) 262-269.
- 3389 Yamaguchi, N., Koyama, K., Yamamura, Y., Okui, A., Konishi, E., Katayama, T., Katsuragi, N., Kurihara, T., Nakazato, H. and Imanishi, J.: A novel proteinase, glucagon-degrading enzyme, secreted by a human pancreatic cancer cell line, HPC-YO. *J. Biochem. (Tokyo)*, 117 (1995) 7-10.
- 3390 Zamolodchikova, T.S., Vorotyntseva, T.I. and Antonov, C.K.: Duodenase, a new serine protease of unusual specificity from bovine duodenal mucosa. Purification and properties. *Eur. J. Biochem.*, 227 (1995) 866-872.
- See also 3058, 3074, 3082, 3206.
- 20g. *Lyases*
- 3391 Broux, M., Ravel, S. and Douce, R.: Methionine biosynthesis in higher plants. II. Purification and characterization of cystathionine β -lyase from spinach chloroplasts. *Arch. Biochem. Biophys.*, 316 (1995) 585-595.
- 3392 De Waal, A., Meijer, A.H. and Verpoorte, R.: Strictosidine synthase from *Catharanthus roseus*: purification and characterization of multiple forms. *Biochem. J.*, 306 (1995) 571-580.
- 3393 Euverink, G.J.W., Wolters, D.J. and Dijkhuizen, L.: Prephenate dehydratase of the actinomycete *Amycolatopsis methanolica*: purification and characterization of wild-type and deregulated mutant proteins. *Biochem. J.*, 308 (1995) 313-320.
- 3394 He, Z. and Wiegel, J.: Purification and characterization of an oxygen-sensitive reversible 4-hydroxybenzoate decarboxylase from *Clostridium hydroxybenzoicum*. *Eur. J. Biochem.*, 229 (1995) 77-82.
- 3395 Law, R.D. and Plaxton, W.C.: Purification and characterization of a novel phosphoenolpyruvate carboxylase from banana fruit. *Biochem. J.*, 307 (1995) 807-816.
- 3396 Ravel, S., Droux, M. and Douce, R.: Methionine biosynthesis in higher plants. I. Purification and characterization of cystathionine γ -synthase from spinach chloroplasts. *Arch. Biochem. Biophys.*, 316 (1995) 572-584.
- 3397 Shelton, M.C. and Toone, E.J.: Differential dye-ligand chromatography as a general purification protocol for 2-keto-3-deoxy-6-phosphogluconate aldolases. *Tetrahedron: Asymmetry*, 6 (1995) 207-211; *C.A.*, 122 (1995) 154855g.
- 3398 Zhang, Z., Schofield, C.J., Baldwin, J.E., Thomas, P. and John, P.: Expression, purification and characterization of 1-aminocyclopropane-1-carboxylate oxidase from tomato in *Escherichia coli*. *Biochem. J.*, 307 (1995) 77-85.
- See also 2856, 3096.
- 20h. *Isomerases*
- 3399 Ashiuchi, M., Yoshimura, T., Kitamura, T., Kawata, Y., Nagai, J., Gorlatov, S., Esaki, N. and Soda, K.: *In vivo* effect of GroESL on the folding of glutamate racemase of *Escherichia coli*. *J. Biochem. (Tokyo)*, 117 (1995) 495-498.
- 3400 Huang, Y. and Dennis, D.T.: Histidine residues 139, 363 and 500 are essential for catalytic activity of cofactor-independent phosphoglyceromutase from developing endosperm of the castor plant. *Eur. J. Biochem.*, 229 (1995) 395-402.
- 3401 Saito, M., Hori, K., Kurotsu, T., Kanda, M. and Saito, Y.: Three conserved glycine residues in valine activation of gramicidin S synthetase 2 from *Bacillus brevis*. *J. Biochem. (Tokyo)*, 117 (1995) 276-282.
- 3402 Stein, T., Kluge, B., Vater, J., Franke, P., Otto, A. and Wittmann-Liebold, B.: Gramicidin S synthetase 1 (phenylalanine racemase), a prototype of amino acid racemases containing the cofactor 4'-phosphopantetheine. *Biochemistry*, 34 (1995) 4633-4642.
- 20i. *Ligases*
- See 3131.
- 20j. *Complex mixtures and incompletely identified enzymes*
- 3403 Alexiou, M. and Leese, H.J.: Enzymes of purine salvage and catabolism in the mouse preimplantation embryo measured by high performance liquid chromatography. *J. Reprod. Fert.*, 101 (1994) 151-158; *C.A.*, 122 (1995) 100153h.
- 3404 Gao, X.-D., Katsumoto, T. and Onodera, K.: Purification and characterization of chitin deacetylase from *Absidia coerulea*. *J. Biochem. (Tokyo)*, 117 (1995) 257-263.
21. PURINES, PYRIMIDINES, NUCLEIC ACIDS AND THEIR CONSTITUENTS
- 21a. *Purines, pyrimidines, nucleosides, nucleotides*
- 3405 Bedford, C.T., Hickman, A.D. and Logan, C.J.: Mixed ion-pair high-performance liquid chromatography of uridine 5'-diphospho- α -D-glucuronic acid and its hydrolysis products. *J. Chromatogr. A*, 695 (1995) 123-127.
- 3406 Bouliou, R. and Lenoir, A.: Determination of thiopurine nucleotides in human lung tissue by high-performance liquid chromatography. *J. Chromatogr. B*, 665 (1995) 213-216.
- 3407 Chimi, E.N. and Dousa, T.P.: Enzymatic synthesis and degradation of nicotinate adenine dinucleotide phosphate (NAADP), a Ca^{2+} -releasing agonist, in rat tissues. *Biochem. Biophys. Res. Commun.*, 209 (1995) 167-174.
- 3408 Dolan, J.W.: Questions about ion pairing. *LC-GC*, 12 (1994) 820-824; *C.A.*, 122 (1995) 100761e.

- 3409 Edwards, R.: Determination of S-adenosyl-L-methionine and S-adenosyl-L-homocysteine in plants. *Phytochem. Anal.*, 6 (1995) 25-30; C.A., 122 (1995) 127811h.
- 3410 Fernandez, B., Centeno, M.L., Sanchez-Tames, R. and Rodriguez, A.: Simultaneous analysis of cytokinins, auxins and abscisic acid by combined immunoaffinity chromatography, high-performance liquid chromatography and immunoassay. *Phytochem. Anal.*, 6 (1995) 49-54; C.A., 122 (1995) 127812j.
- 3411 Fischer, J.H.: Specific detection of nucleotides, creatine phosphate, and their derivatives from tissue samples in a simple, isocratic, recycling, low-volume system. *LC-GC Int.*, 8, No. 5 (1995) 254-264.
- 3412 Gajewska, M., Pawinski, T., Dziargowska-Szmidt, A., Grieb, P. and Ryba, M.: Determination of 2-bromo and 2-chloro derivatives of 2'-deoxyadenosine in human blood serum by high performance liquid chromatography. *Acta Pol. Pharm.*, 51 (1994) 109-114; C.A., 122 (1995) 100746d.
- 3413 Irisawa, M. and Komiyama, M.: Hydrolysis of DNA through cooperation of two metal ions: a novel mimic of phosphoesterases. *J. Biochem. (Tokyo)*, 117 (1995) 465-466.
- 3414 Iwai, S., Maeda, M., Shirai, M., Shimada, Y., Osafune, T., Murata, T. and Ohtsuka, E.: Reaction mechanism of T4 endonuclease V determined by analysis using modified oligonucleotide duplexes. *Biochemistry*, 34 (1995) 4601-4609.
- 3415 Krohn, M. and Wagner, R.: A procedure for the rapid preparation of guanosine tetraphosphate (ppGpp) from *Escherichia coli* ribosomes. *Anal. Biochem.*, 225 (1995) 188-190.
- 3416 Kyossef, Z.N., Drake, R.R., Kyosseva, S.V. and Elbein, A.D.: Synthesis of a new photoaffinity probe, 5-azido-[³²P]UDP xylose, by UDPglucuronate carboxylase from wheat germ. *Eur. J. Biochem.*, 228 (1995) 109-112.
- 3417 Liljebjelke, K., Adolphson, R., Baker, K., Doong, R.L. and Mohnen, D.: Enzymatic synthesis and purification of uridine diphosphate [¹⁴C]galacturonic acid: a substrate for pectin biosynthesis. *Anal. Biochem.*, 225 (1995) 296-304.
- 3418 Lou, S.N. and Montag, A.: (Determination of the status of nucleoside compounds in mushrooms). *Dtsch. Lebensm.-Rundsch.*, 90 (1994) 212-217; C.A., 122 (1995) 29956q.
- 3419 Meredith, M., Rabaglia, M. and Metz, S.: Cytosolic biosynthesis of GTP and ATP in normal rat pancreatic islets. *Biochim. Biophys. Acta*, 1266 (1995) 16-22.
- 3420 Meynial, Y., Paquet, V. and Combes, D.: Simultaneous separation of nucleotides and nucleotide sugars using an ion-pair reversed-phase HPLC: application for assaying glycosyltransferase activity. *Anal. Chem.*, 67 (1995) 1627-1631.
- 3421 Rush, J.S. and Waechter, C.J.: Method for the determination of cellular levels of guanosine-5'-diphosphate-mannose based on a weak interaction with concanavalin A at low pH. *Anal. Biochem.*, 224 (1995) 494-501.
- 3422 Schubert, B.: Enzymatic synthesis of ATP analogs and their purification by reverse-phase high-performance liquid chromatography. *Anal. Biochem.*, 226 (1995) 288-292.
- 3423 Shah, G.M., Poirier, D., Duchaine, C., Brochu, G., Desnoyers, S., Laguex, J., Verreault, A., Hoflack, J.-C., Kirkland, J.B. and Poirier, G.G.: Methods for biochemical study of poly(ADP-ribose) metabolism *in vitro* and *in vivo*. *Anal. Biochem.*, 227 (1995) 1-13, - review with some papers cited.
- 3424 Stolk, J.N., de Abreu, R.A., Boerbooms, A.M.T., de Koning, D.G.M., de Graaf, R., Kerstens, P.J.S.M. and van de Putte, L.B.A.: Purine enzyme activities in peripheral blood mononuclear cells: comparison of a new non-radiochemical high-performance liquid chromatography procedure and a radiochemical thin-layer chromatography procedure. *J. Chromatogr. B*, 666 (1995) 33-43.
- 3425 Streeper, R.T., Cotter, R.J., Colvin, M.E., Hilton, J. and Colvin, O.M.: Molecular pharmacology of hepsulfam, NSC 3296801: identification of alkylated nucleosides, alkylation site, and site of DNA cross-linking. *Cancer Res.*, 55 (1995) 1491-1498.
- 3426 Suzuki, J., Knoue, Y. and Suzuki, S.: Changes in the urinary excretion level of 8-hydroxyguanine by exposure to reactive oxygen-generating substances. *Free Radical Biol. Med.*, 18 (1995) 431-436; C.A., 122 (1995) 127810g.
- 3427 Warren, D.J., Andersen, A. and Slordal, L.: Quantitation of 6-thioguanine residues in peripheral blood leukocyte DNA obtained from patients receiving 6-mercaptopurine-based maintenance therapy. *Cancer Res.*, 55 (1995) 1670-1674.
- 3428 Yhang, Y., Lu, H. and Wang, J.: (Determination of nucleotides in food by HPLC). *Shipin Kexue (Beijing)*, (1994) 59-62; C.A., 122 (1995) 79324c.

For additional information see C.A.:

122 (1995) 50384m, 101064s, 150721h, 208913v.

See also 2994, 3466, 3483, 3721, 3727, 3742, 3748.

21b. Nucleic acids, RNA

- 3429 Dewanjee, M.K., Ghafouripour, A.K., Kapadvanjwala, M. and Samy, A.T.: Kinetics of hybridization of mRNA of c-myc oncogene with ¹¹¹In-labeled antisense oligodeoxynucleotide probes by high-pressure liquid chromatography. *BioTechniques*, 16 (1994) 844-850; C.A., 122 (1995) 73083b.
- 3430 Feuser, P., Hermann, R., Schorr, J., Colpan, M. and Bastian, H.: Chromatographic purification and separation of nucleic acid mixtures. *Ger. Offen. DE 4,321,904 (Cl. CO7H21/04)*, 12 Jan. 1995, Appl. 01 Jul. 1993; 9 pp.; C.A., 122 (1995) 101132n.
- 3431 Kanduc, D., Bracalello, A. and Mosavi, A.: HPLC fraction of rat liver tRNA. Isolation of tRNA^{Glu}, tRNA^{Gly}, tRNA^{Phe}, and tRNA^{Thr}. *Biochem. Arch.*, 10 (1994) 243-248; C.A., 122 (1995) 50388r.
- 3432 Kanduc, D., Bracalello, A., Troccoli, G., de Benedetto, C., Mosavi, A. and Azzarone, A.: Separation of rat liver cytoplasmic tRNAs by high performance liquid chromatography. *J. Liq. Chromatogr.*, 18 (1995) 505-513.
- 3433 Oefner, P.J. and Bonn, G.K.: High-resolution liquid chromatography of nucleic acids. *Am. Lab. (Shelton)*, 26 (1994) 28C, 28D, 28F-28J; C.A., 122 (1995) 155374m.
- 3434 Yamakawa, Y. and Yamada, Y.: (Separation of *E. coli* tRNAs by two dimensional chromatography on hydroxyapatite and hydrophobic interaction chromatography). *Kuromatogurafi*, 15 (1994) 246-249; C.A., 122 (1995) 127790a.

See also 3119, 3413, 3418.

21c. Nucleic acids, DNA

- 3435 Douki, T., Voituriez, L. and Cadet, J.: Measurement of pyrimidine (6-4) photoproducts in DNA by a mild acidic hydrolysis-HPLC fluorescence detection assay. *Chem. Res. Toxicol.*, 8 (1995) 244-253; C.A., 122 (1995) 127596s.
- 3436 Milliman, A., Huang, C.-R., Kim, Y. and LeBreton, P.R.: HPLC preparation of highly purified single-stranded M13 DNA. *BioTechniques*, 18 (1995) 308-315; C.A., 122 (1995) 127809p.
- 3437 Oefner, P.J., Huber, C.G., Puchhammer-Stoeckl, E., Umlauf, F., Gruenwald, K., Bonn, G.K. and Kunz, C.: High-performance liquid chromatography for routine analysis of hepatitis C virus cDNA/PCR products. *BioTechniques*, 16 (1994) 898-908; C.A., 122 (1995) 47659z.
- 3438 Sundaram, K.M.S. and Sloane, L.: Liquid chromatographic assay for the separation of single- and double-stranded DNA by using UV and UV diode-array detectors and hydroxylapatite column. *J. Liq. Chromatogr.*, 18 (1995) 925-939.

See also 3119, 3128, 3413, 3418, 3430, 3433.

21e. Structural studies on DNA and DNA mapping

- 3439 Ferrero, E.A., Fitch, M.M.M., Moore, P.H. and Wu, K.-K.: Resin-purified digoxigenin-labeled DNA probes: an efficient protocol for mapping and fingerprinting sugarcane. *BioTechniques*, 18 (1995) 266-273; C.A., 122 (1995) 206312z.

See also 2629.

22. ALKALOIDS

- 3440 Asano, N., Kato, A., Oseki, K., Kizo, H. and Matsui, K.: Calistegins of *Physalis alkekengi* var. *francheti* (Solanaceae). Structure determination and their glycosidase inhibitory activities. *Eur. J. Biochem.*, 229 (1995) 269-376.
- 3441 Chen, C.-M. and Chang, H.-C.: Determination of berberine in plasma, urine and bile by high-performance liquid chromatography. *J. Chromatogr. B*, 665 (1995) 117-123.
- 3442 Draeger, B.: Identification and quantification of calystegines, polyhydroxyl nortropane alkaloids. *Phytochem. Anal.*, 6 (1995) 31-37; C.A., 122 (1995) 155671f.
- 3443 Fernandez, P., Bermejo, A.M., Lopez-Rivadulla, M. and Landeira, A.C.: Determination of colchicine in biological fluids by reverse-phase HPLC: variation of colchicine levels in rats. In: Oliver, J.S. (Editor), *Forensic Toxicol., Proc. Int. Meet. Int. Assoc. Forensic Toxicol.*, 26th 1992, Scott. Acad. Press, Edinburgh, 1992, pp. 240-243; C.A., 122 (1995) 232820h.
- 3444 Gontier, E., Sangwan, B.S. and Barbotin, J.N.: Simultaneous determination of some tropane alkaloids by reversed-phase HPLC of *Datura innoxia* Mill. plant material. *Agro-Food-Ind. Hi-Tech*, 5 (1994) 26-28; C.A., 122 (1995) 182311v.
- 3445 Imamura, T., Nagata, T., Tokunaga, H., Kudo, K., Jitsufuchi, N. and Nakasono, I.: (Forensic toxicological analysis of colchicine in human samples using liquid chromatography-mass spectrometry). *Nippon Iyo Masu Supekutoru Gakkai Koenshu*, 19 (1994) 155-158; C.A., 122 (1995) 74091q.

- 3446 Kuczynski, J.: (Retention of some alkaloids in aqueous chromatographic systems with reversed phases. I. The acetate buffer-methanol system (RP-19). *Acta Pol. Pharm.*, 50 (1993) 131-136; C.A., 122 (1995) 150536f.
- 3447 Kuczynski, J.: (Retention of some alkaloids in aqueous chromatographic systems with reversed phases. II. The phosphate buffer-methanol system with a counterion (RP-18). *Acta Pol. Pharm.*, 50 (1993) 137-141; C.A., 122 (1995) 150537g.
- 3448 Lame, M.W., Jones, A.D., Morin, D., Segall, H.J. and Wilson, D.W.: Biliary excretion of pyrrolic metabolites of [¹⁴C]monocrotaline in the rat. *Drug Metab. Disp.*, 23 (1995) 422-429.
- 3449 Mwalupindi, A.G. and Warner, I.M.: Determination of theophylline by liquid chromatography with sensitized lanthanide luminescence detection. *Anal. Chim. Acta*, 306 (1995) 49-56.
- 3450 Perez-Martinez, I., Sagrado, S., Medina-Hernández, M.J.: A rapid procedure for the determination of caffeine, theophylline and theobromine in urine by micellar liquid chromatography and direct sample injection. *Anal. Chim. Acta*, 304 (1995) 195-201.
- 3451 Rehwald, A., Sticher, O. and Meier, B.: Trace analysis of Harman alkaloids in *Passiflora incarnata* by reversed-phase high-performance liquid chromatography. *Phytochem. Anal.*, 6 (1995) 96-100; C.A., 122 (1995) 197108h.
- 3452 Reynolds, J. and Albazi, S.J.: Simultaneous determination of nicotine and cotinine in untreated human urine by micellar liquid chromatography. *J. Liq. Chromatogr.*, 18 (1995) 537-552.
- 3453 Seregina, Ye.V. and Laipanov, A.Kh.: (Use of HPLC in analysis of tropane alkaloids). *Farmatsiya (Moscow)*, 43 (1994) 44-47; C.A., 122 (1995) 170327s.
- 3454 Weingarten, B., Wang, H.-Y. and Roberts, D.M.: Determination of codeine in human plasma by high-performance liquid chromatography with fluorescence detection. *J. Chromatogr. A*, 696 (1995) 83-92.
- 3455 Yevstratova, K.I. and Kudryashov, M.N.: (Analysis of scopolamine from *Datura innoxia* seeds). *Farmatsiya (Moscow)*, 43 (1994) 37-44; C.A., 122 (1995) 170326r.

For additional information see C.A.:

122 (1995) 75715h, 75717k, 98906k, 115093n, 115095q, 177547x, 197102b.

See also 3679, 3681, 3688, 3700, 3709, 3782, 3785, 3786, 3789, 3810.

23. OTHER SUBSTANCES CONTAINING HETEROCYCLIC NITROGEN

23a. Porphyrins and other pyrroles

- 3456 Beukeveld, G.J.J. and Wolthers, B.G.: Cholestyramine orally administered to patients with erythropoietic protoporphyria results in urinary excretion of protoporphyrin: to be explained by the Herbst-Volkheimer effect? *Clin. Chim. Acta*, 233 (1995) 119-126.
- 3457 Hori, H., Ishitani, O. and Ibusuki, T.: High performance liquid chromatographic isolation and characterization of (pheophytin)mercury(II). *J. Liq. Chromatogr.*, 18 (1995) 779-789.

3458 Hua, Z., Gibson, S.L., Foster, T.H. and Hilf, R.: Effectiveness of δ -aminolevulinic acid-induced protoporphyrin as a photosensitizer for photodynamic therapy *in vivo*. *Cancer Res.*, 55 (1995) 1723-1731.

23c. *Indole derivatives and plant hormones (gibberelins)*

3459 Hansen-Moeller, J. and Andersen, J.R.: Boar taint. Analytical alternatives. *Fleischwirtschaft*, 74 (1994) 963-966, 1005-1009; *C.A.*, 122 (1995) 212340y.

3460 Huang, T. and Kissinger, P.T.: Determination of serotonin in microdialyzates using UniJet fused silica microbore columns. *Curr. Sep.*, 13 (1995) 114-118; *C.A.*, 122 (1995) 230953e.

3461 Julliard, J., Maldiney, R., Sotta, B., Miginiac, E., Kerhoas, L. and Einhorn, J.: HPLC-ELISA and MS as complementary techniques to study plant hormone metabolites. *Analysis*, 22 (1994) 483-489; *C.A.*, 122 (1995) 182656m.

3462 Kagedal, B., Konradsson, P., Shibata, T. and Mishima, Y.: High-performance liquid-chromatographic analysis of dopachrome and dihydroxyphenylalanine. *Anal. Biochem.*, 225 (1995) 264-269.

3463 Opper, C., Wesemann, W., Barka, G., Kerkdijk, H. and Haak, G.: Analysis of serotonin in whole-blood samples - a novel fully automated method. *LC-GC Int.*, 8, No. 2 (1995) 104-106.

For additional information see C.A.:

122 (1995) 237916d.

See also 2992, 3008, 3410.

23e. *Other N-heterocyclic compounds*

3464 Bellstedt, K. and Seidel, J.: Determination of orotic acid in urine by means of HPLC. *Eur. J. Clin. Chem. Clin. Biochem.*, 33 (1995) A21-A22.

3465 Busse-Grawitz, A., Bissé, E. and Wieland, H.: HPLC method for a rapid and accurate determination of 3-methylhistidine (3-MH). *Eur. J. Clin. Chem. Clin. Biochem.*, 33 (1995) A19.

3466 Kock, R., Seitz, S., Delvoux, B. and Greiling, H.: A method for the simultaneous determination of creatinine and uric acid in serum by high-performance liquid chromatography evaluated versus reference method. *Eur. J. Clin. Chem. Clin. Biochem.*, 33 (1995) 23-29.

3467 Mao, J., Pacheco, C.R., Traficante, D.D. and Rosen, W.: Analysis of benzoquinolines and acridines in a Brazilian diesel oil by particle beam LC/MS and HPLC/UV. *J. Liq. Chromatogr.*, 18 (1995) 903-916.

3468 Mawatari, K.-i., Oshida, K., Iinuma, F. and Watanabe, M.: Determination of quinolinic acid in human urine by liquid chromatography with fluorimetric detection. *Anal. Chim. Acta*, 302 (1995) 179-183.

3469 Scholl, J.P.: An evaluation of column-to-column retention variability for compounds related to panadiplon (U-78,875) using three different reverse-phase HPLC separation modes. *J. Liq. Chromatogr.*, 17 (1994) 3369-3382.

3470 Thienpont, L.M., van Landuyt, K.G., Stöckl, D., Saeyens, W., de Keukeleire, D. and de Leenheer, A.P.: Evaluation of 2-iminoimidazolidin-4-one and thymine as respective internal standards for normal-phase and reversed-phase high-performance liquid chromatographic determination of creatinine in human serum. *J. Chromatogr. B*, 665 (1995) 63-69.

See also 2639, 2795, 3057, 3715, 3798.

24. ORGANIC SULPHUR COMPOUNDS (INCL. GLUCOSINOLATES)

3471 Gasparrini, F., Misiti, D. and Villani, C.: Behaviour of allyl aryl sufoxides in high-performance liquid chromatography on a chiral stationary phase. *J. Chromatogr. A*, 694 (1995) 163-167.

3472 Haj-Yehia, A.I. and Benet, L.Z.: 2-(4-N-Maleimidophenyl)-6-methoxybenzofuran: a superior derivatizing agent for fluorimetric determination of aliphatic thiols by high-performance liquid chromatography. *J. Chromatogr. B*, 666 (1995) 45-53.

3473 Howard, A.G. and Russell, D.W.: HPLC-FPD instrumentation for the measurement of the atmospheric dimethyl sulfide precursor β -(dimethylsulfonio)propionate. *Anal. Chem.*, 67 (1995) 1293-1295.

3474 Majima, E., Goto, S., Hori, H., Shinohara, Y., Hong, Y.-M. and Terada, H.: Stabilities of the fluorescent SH-reagent eosin-5-maleimide and its adducts with sulfhydryl compounds. *Biochim. Biophys. Acta*, 1243 (1995) 336-342.

3475 Okafo, G.N., Rana, K.K. and Camilleri, P.: Improved separation of diastereoisomers in capillary electrophoresis using a mixture of β -cyclodextrin and sodium taurodeoxycholate. *Chromatographia*, 39 (1994) 627-630.

3476 Winters, R.A., Zukowski, J., Ercal, N., Matthews, R.H. and Spitz, D.R.: Analysis of glutathione, glutathione disulfide, cysteine, homocysteine, and other biological thiols by high-performance liquid chromatography following derivatization by N-(1-pyrenyl)maleimide. *Anal. Biochem.*, 227 (1995) 14-21.

For additional information see C.A.:

122 (1995) 27151z.

See also 2795, 3715, 3836.

25. ORGANIC PHOSPHORUS COMPOUNDS (INCL. SUGAR PHOSPHATES)

3477 Abidi, S.L. and Mounts, T.L.: Separations of molecular species of phosphatidic acid by high-performance liquid chromatography. *J. Chromatogr. A*, 694 (1995) 365-373.

3478 Altrichter, J., Guse, A.H., Resch, K., Brock, J., Daeron, M. and Hüchel, C.: Phosphatidylinositol hydrolysis and an increase in Ca^{2+} concentration in the signal-transduction process triggered by murine Fc γ RIII are not required for protein kinase C translocation. *Eur. J. Biochem.*, 228 (1995) 587-595.

3479 Barker, C.J., French, P.J., Moore, A.J., Milsson, T., Berggren, P.-O., Bunce, C.M., Kirk, C.J. and Michell, R.H.: Inositol 1,2,3-trisphosphate and inositol 1,2- and/or 2,3-bisphosphate are normal constituents of mammalian cells. *Biochem. J.*, 306 (1995) 557-564.

- 3480 Burbano, C., Muzquiz, M., Osagie, A., Ayet, G. and Cuadrado, C.: Determination of phytate and lower inositol phosphates in Spanish legumes by HPLC methodology. *Food Chem.*, 52 (1995) 321-325; *C.A.*, 122 (1995) 185705n.
- 3481 Chen, S. and Claeys, M.: Approach to the large-scale preparation of highly pure phosphatidylserine from bovine brain. *J. Chromatogr. B*, 666 (1995) 178-182.
- 3482 Edwards, Y.S. and Murray, A.W.: Accumulation of phosphatidylalcohol in cultured cells: use of subcellular fractionation to investigate phospholipase D activity during signal transduction. *Biochem. J.*, 308 (1995) 473-480.
- 3483 Guranowski, A., Starzyńska, E., Sillero, M.A.G. and Sillero, A.: Conversion of adenosine(5') oligophospho(5') adenosines into inosine(5') oligophospho(5') inosines by non-specific adenylate deaminase from the snail *Helix pomatia*. *Biochim. Biophys. Acta*, 1243 (1995) 78-84.
- 3484 Tul'skaya, E.M., Shashkov, A.S., Evtushenko, L.I., Taran, V.V. and Naumova, I.B.: (Cell wall teichoic acids of *Nocardia albida* ssp. *albida* DSM 43120). *Biokhimiya (Moscow)*, 60 (1995) 179-186.
- 3485 Van Dijken, P., Lammers, A.A. and van Haastert, P.J.M.: In *Dicystostelium discoideum* inositol 1,3,4,5-tetrakisphosphate is dephosphorylated by a 3-phosphatase and a 1-phosphatase. *Biochem. J.*, 308 (1995) 127-130.
- See also 2926, 2929, 2930, 2931, 2939, 2942, 2945, 2949, 3025, 3077, 3405, 3417, 3419.
26. ORGANOMETALLIC AND RELATED COMPOUNDS
- 26a. *Organometallic compounds*
- 3486 Compañó, R., Granados, M., Leal, C. and Prat, M.D.: Determination of triphenyltin in sea water samples by liquid chromatography with fluorimetric detection. *Anal. Chim. Acta*, 302 (1995) 185-191.
- 3487 Gamoh, K., Ketuly, K.A., Cole, W.J., Brooks, C.J.W. and Anderson, R.A.: Chromatographic mass-spectrometric studies of cyclic 2-(N,N-dimethylaminomethyl)ferroceneboronates and related esters. *Anal. Sci.*, 10 (1994) 705-711; *C.A.*, 122 (1995) 204200n.
- 3488 Hempel, M., Chau, Y.K., Dutka, R.J., McInnis, R., Kwan, K.K. and Liu, D.: Toxicity of organomercury compounds: Bioassay results as a basis for risk assessment. *Analyst (Cambridge)*, 120 (1995) 721-724.
- 3489 Matni, G., Azani, R., van Calsteren, M.R., Bissonnette, M.C. and Blais, J.S.: Determination of free selenomethionine in nutritional supplements by high-performance liquid chromatography coupled with thermochemical hydride generation atomic absorption spectrometry. *Analyst (Cambridge)*, 120 (1995) 395-401.
- 3490 Padró, A., Rubio, R. and Rauret, G.: Germanium speciation by LC-HG-ICP/OES. *Fresenius J. Anal. Chem.*, 351 (1995) 449-453.
- 3491 Spurnar-Lobinska, J., Witte, C., Lobinski, R. and Adams, F.C.: Separation techniques in speciation analysis for organometallic species. *Fresenius J. Anal. Chem.*, 351 (1995) 351-377.
- 3492 Suzuki, K.T., Itoh, M. and Ohmichi, M.: Detection of selenium-containing biological constituents by high-performance liquid chromatography-plasma source mass spectrometry. *J. Chromatogr. B*, 666 (1995) 13-19.
- See also 3854.
- 26c. *Coordination compounds*
- 3493 Kumar, K., Sukumaran, K., Taylor, S., Chang, C.A., Nunn, A.D. and Tweedle, M.F.: Partition coefficients (log P) and HPLC capacity factors (k') of some Gd(III) complexes of linear and macrocyclic polyamino carboxylates. *J. Liq. Chromatogr.*, 17 (1994) 3735-3746.
- 3494 Nakata, R., Nagayama, M., Naitoh, T., Yasuma, S. and Oki, H.: Liquid chromatography of diastereoisomers of neutral tris- and bis(L- α -amino acidato)chromium(III) complexes. *Anal. Sci.*, 10 (1994) 663-665; *C.A.*, 122 (1995) 150547k.
- 3495 Tallarek, U., Baumeister, E., Albert, K., Bayer, E. and Guiochon, G.: NMR imaging of the chromatographic process. Migration and separation of bands of gadolinium chelates. *J. Chromatogr. A*, 696 (1995) 1-18.
- 3496 Tan, K.-C., Hor, T.S.A. and Lee, H.K.: High-performance liquid chromatography of group 6 dimetal decacarbonyl complexes singly-bridged by diphosphine ligands. *J. Liq. Chromatogr.*, 17 (1994) 3671-3680.
- See also 3862.
- 27 VITAMINS AND VARIOUS ANIMAL GROWTH FACTORS (NON-PEPTIDIC)
- 3497 Acilu, M. and Font, J.: (Identification of tocopherol isomers or vitamin E in vegetable oil samples by HPLC with photodiode detector). *Tec. Lab.*, 16 (1994) 866-869; *C.A.*, 122 (1995) 185660u.
- 3498 Adamson, P.C., Murphy, R.F., Godwin, K.A., Uim, E.H. and Balis, F.M.: Pharmacokinetics of 9-cis-retinoic acid in the rhesus monkey. *Cancer Res.*, 55 (1995) 482-485.
- 3499 Ambrosini, L. and Sala, B.: Method for the industrial preparation of (6S) folic acid derivatives by chromatographic separation. *Eur. Pat. Appl. EP 627,435 (Cl. C07D475/04)*, 07 Dec. 1994, Appl. 93/108,752, 01 Jun. 1993; 6 p.; *C.A.*, 122 (1995) 109342b.
- 3500 Bempong, D.K., Honigberg, I.L. and Meltzer, N.M.: Normal phase LC-MS determination of retinoic acid degradation products. *J. Pharm. Biomed. Anal.*, 13 (1995) 285-291.
- 3501 Bergaentzle, M., Arella, F., Bourguignon, J.B. and Hasselmann, C.: Determination of vitamin B6 in foods by HPLC - a collaborative study. *Food Chem.*, 52 (1995, Pub. 1994) 81-86; *C.A.*, 122 (1995) 79368v.
- 3502 Blanco, D., Pajares, M., Escotet, V.J. and Gutiérrez, M.D.: Determination of fat-soluble vitamins by liquid chromatography in pediatric parenteral nutrition. *J. Liq. Chromatogr.*, 17 (1994) 4513-4530.
- 3503 Cahnmann, H.J.: A fast photoisomerisation method for the preparation of tritium-labeled 9-cis-retinoic acid of high specific activity. *Anal. Biochem.*, 227 (1995) 49-53.

- 3504 Delgado-Zamarreño, M.M., Sanchez-Perez, A., Gomez-Perez, M.C. and Hernandez-Mendez, J.: Directly coupled sample treatment-high-performance liquid chromatography for on-line automatic determination of liposoluble vitamins in milk. *J. Chromatogr. A*, 694 (1995) 399-406.
- 3505 Du, Q., Zhang, T.-Y. and Ito, Y.: Purification of 25-hydroxy-cholecalciferol from irradiation of cholesta-5,7-diol by high-speed countercurrent chromatography. *J. Liq. Chromatogr.*, 18 (1995) 181-188.
- 3506 Esteve, M.J., Farre, R., Frigola, A., Lopez, J.C., Romera, J.M., Ramirez, M. and Gil, A.: Comparison of voltammetric and high performance liquid chromatographic methods for ascorbic acid determination in infant formulas. *Food Chem.*, 52 (1995, Pub. 1994) 99-102; *C.A.*, 122 (1995) 79370q.
- 3507 Fujiwara, K., Asao, T., Ikeda, M., Kisara, K. and Ono, S.: (Determination of vitamin K in plasma by HPLC with electrochemical detection). *Bitamin*, 68 (1994) 711-719; *C.A.*, 122 (1995) 100739d.
- 3508 Gehring, T.A., Cooper, W.M., Holder, C.L. and Thompson, H.C., Jr.: Liquid chromatographic determination of thiamine in rodent feed by postcolumn derivatization and fluorescence detection. *J. Assoc. Off. Anal. Chem.*, 78 (1995) 307-309.
- 3509 Hentz, N.G. and Bachas, L.G.: Class-selective detection system for liquid chromatography based on the streptavidin-biotin interaction. *Anal. Chem.*, 67 (1995) 1014-1018.
- 3510 Hu, O.Y.-P., Wu, C.-Y., Chan, W.-K. and Wu, F.Y.-H.: Determination of anticancer drug vitamin D₃ in plasma by high-performance liquid chromatography. *J. Chromatogr. B*, 666 (1995) 299-305.
- 3511 Huber, J.F.K. and Lamprecht, G.: Assay of neopterin in serum by means of two-dimensional high-performance liquid chromatography with automated column switching using three retention mechanisms. *J. Chromatogr. B*, 666 (1995) 223-232.
- 3512 Kraft, J.C., Kimelman, D. and Juchau, M.R.: *Xenopus laevis*: a model system for the study of embryonic retinoid metabolism. I. Embryonic metabolism of 9-*cis* and all-*trans*-retinals and retinols to their corresponding acid form. *Drug Metab. Disp.*, 23 (1995) 72-82.
- 3513 Kraft, J.C., Kimelman, D. and Juchau, M.R.: *Xenopus laevis*: a model system for the study of embryonic retinoid metabolism. II. Embryonic metabolism of all-*trans*-3,4-didehydroretinol to all-*trans*-3,4-didehydroretinoic acid. *Drug Metab. Disap.*, 23 (1995) 83-89.
- 3514 Lefebvre, P., Agadir, A., Cornic, M., Gourmel, B., Hue, B., Dreux, C., Degos, L. and Chomienne, C.: Simultaneous determination of all-*trans* and 13-*cis* retinoic acids and their 4-oxo metabolites by adsorption liquid chromatography after solid-phase extraction. *J. Chromatogr. B*, 666 (1995) 55-61.
- 3515 Li, K. and Zhao, F.-L.: Determination of α -tocopherol in human and rabbit plasma or serum by high-performance liquid chromatography. *J. Chin. Pharm. Sci.*, 3 (1994) 164-170; *C.A.*, 122 (1995) 155381m.
- 3516 Liaaen-Jensen, S. and Hertzberg, S.: Example 11: carotenoid sulfates. In: Britton, G., Liaaen-Jensen, S. and Pfander, H. (Editors), *Carotenoids*, Birkhaeuser, Basel, 1995, pp. 283-286; *C.A.*, 122 (1995) 234521k.
- 3517 Manan, F. and Ryley, J.: Development of an improved HPLC method for vitamin A assay in foods. *Discovery Innovation*, 6 (1994) 287-290; *C.A.*, 122 (1995) 104172m.
- 3518 Pfander, H.: Chromatography: Part I. General aspects. In: Britton, G., Liaaen-Jensen, S. and Pfander, H. (Editors), *Carotenoids*, Birkhaeuser, Basel, 1995, pp. 109-116; *C.A.*, 122 (1995) 234411z - a review with 5 refs.
- 3519 Pfander, H. and Niggli, U.: Chromatography: Part V. Supercritical-fluid chromatography. In: Britton, G., Liaaen-Jensen, S. and Pfander, H. (Editors), *Carotenoids*, Birkhaeuser, Basel, 1995, pp. 191-198; *C.A.*, 122 (1995) 234415d - a review with 14 refs.
- 3520 Pfander, H. and Riesen, R.: Chromatography: Part IV. High-performance liquid chromatography. In: Britton, G., Liaaen-Jensen, S. and Pfander, H. (Editors), *Carotenoids*, Birkhaeuser, Basel, 1995, pp. 145-190; *C.A.*, 122 (1995) 234414c - a review with 108 refs.
- 3521 Poon, R., Chu, L., Lecavalier, P., Bergman, A. and Villeneuve, D.C.: Urinary ascorbic acid - HPLC determination and application as a noninvasive biomarker of hepatic response. *J. Biochem. Toxicol.*, 9 (1994) 297-304; *C.A.*, 122 (1995) 74070g.
- 3522 Randolph, R.K. and Simon, M.: Metabolic regulation of active retinoid concentrations in cultured human epidermal keratinocytes by exogenous fatty acids. *Arch. Biochem. Biophys.*, 318 (1995) 6-14.
- 3523 Scalia, S., Renda, A., Ruberto, G., Bonina, F. and Menegatti, E.: Assay of vitamin A palmitate and vitamin E acetate in cosmetic creams and lotions by supercritical fluid extraction and HPLC. *J. Pharm. Biomed. Anal.*, 13 (1995) 273-277.
- 3524 Schuep, W., Hess, D. and Schierle, J.: Example 7: human plasma. Simultaneous determination of carotenoids, retinol and tocopherols. In: Britton, G., Liaaen-Jensen, S. and Pfander, H. (Editors), *Carotenoids*, Birkhaeuser, Basel, 1995, pp. 261-264; *C.A.*, 122 (1995) 234520j.
- 3525 Siddiqui, F.Q., Malik, F. and Fazli, F.R.Y.: Determination of serum retinol by reversed-phase high-performance liquid chromatography. *J. Chromatogr. B*, 666 (1995) 342-346.
- 3526 Skurikhin, V.N.: (Determination of vitamins D₂ and D₃ in feed preparations by microcolumn high-performance liquid chromatography with ultraviolet detection). *Byull. Vseross. Nauchno-Issled. Inst. Fiziol. Biokhim. Pitan. S-kh. Zhivotn.*, (1992) 70-76; *C.A.*, 122 (1995) 158878q.
- 3527 Skurikhin, V.N., Dvinskaya, L.M. and Riabikh, T.E.: (Carotenoids determination by high-performance liquid chromatography in plasma and tissues of farm animals). *Byull. Vseross. Nauchno-Issled. Inst. Fizio', Biokhim. Pitan. S-kh. Zhivotn.*, (1992) 76-82; *C.A.*, 122 (1995) 100770g.
- 3528 Stein, J., Hahn, A. and Rehner, G.: High-performance liquid chromatographic determination of nicotinic acid and nicotinamide in biological samples applying post-column derivatization resulting in bathochrome absorption shifts. *J. Chromatogr. B*, 665 (1995) 71-78.
- 3529 Sugimoto, T., Yoshida, A., Teradaïra, R., Fujita, K. and Nagatsu, T.: Isolation of deoxybiopterin from urine of cancer patients. *Biog. Amines*, 11 (1995) 1-6; *C.A.*, 122 (1995) 234528t.
- 3530 Thompson, C.O. and Trenery, V.C.: A rapid method for the determination of total L-ascorbic acid in fruits and vegetables by micellar electrokinetic capillary chromatography. *Food Chem.*, 53 (1995) 43-50; *C.A.*, 122 (1995) 238024y.

- 3531 Wakabayashi, H., Yamato, S., Nakajima, M. and Shimada, K.: Simultaneous determination of oxidized and reduced coenzyme Q and α -tocopherol in biological samples by high performance liquid chromatography with platinum catalyst reduction and electrochemical detection. *Biol. Pharm. Bull.*, 17 (1994) 997-1002; *C.A.*, 122 (1995) 154758c.
- 3532 Yakushina, L.M. and Taranova, A.G.: (Rapid method for determination of fat-soluble vitamins in human blood serum). *Zh. Fiz. Khim.*, 68 (1994) 1826-1828; *C.A.*, 122 (1995) 50406v.
- 3533 Zhang, Y. and Wang, J.: (Determination of vitamin D in food by high performance liquid chromatography). *Sepu*, 12 (1994) 411-413; *C.A.*, 122 (1995) 54275z - a review with 15 refs.
- 3534 Zhao, H., Li, K., Cao, Y. and Zheng, J.: (Simultaneous determination of vitamins A and D₂ in fortified milk by high performance liquid chromatography). *Sepu*, 13 (1995) 61-62; *C.A.*, 122 (1995) 158912w.
- For additional information see *C.A.*:
122 (1995) 115138f, 155417c, 158871g, 185697m, 197103c.
- See also 2566, 2828, 3464, 3598.
28. ANTIBIOTICS
- 3535 Anderson, K.L., Moats, W.A., Rushing, J.E., Wesen, D.P. and Papich, M.G.: Potential for oxytetracycline administration by three routes to cause milk residues in lactating cows, as detected by radioimmunoassay (Charm II) and high-performance liquid chromatography test methods. *Am. J. Vet. Res.*, 56 (1995) 70-77; *C.A.*, 122 (1995) 122373g.
- 3536 Baertschi, S.W., Dorman, D.E., Spangle, L.A., Collins, M.W. and Lorenz, L.J.: Formation of fluorescent pyrazine derivatives via a novel degradation pathway of the carbacephalosporin loracarbef. *J. Pharm. Biomed. Anal.*, 13 (1995) 323-328.
- 3537 Becker, M., Junker-Buchheit, A. and Wotschokowsky, M.: (Solid-phase extraction and HPLC analysis of cyclosporin A in blood plasma). *Labor-Med.*, 17 (1994) 169-170; *C.A.*, 122 (1995) 71214w.
- 3538 Blanchflower, W.J. and Kennedy, D.G.: Determination of lasalocid in eggs using liquid-chromatography - electrospray mass spectrometry. *Analyst (Cambridge)*, 120 (1995) 1129-1132.
- 3539 Cobin, J.A. and Johnson, N.A.: Liquid chromatographic method for rapid determination of total avermectin B₁ and 8,9-z-avermectin B₁ residues in apples. *J. Assoc. Off. Anal. Chem.*, 78 (1995) 419-423.
- 3540 Coleman, M.R., Macy, T.D., Moran, J.W. and Rodewald, J.M.: Ruggedness of the monensin and narasin liquid chromatographic assays. *J. Assoc. Off. Anal. Chem.*, 77 (1994) 1065-1072.
- 3541 Croubels, S., Baeyens, W. and van Peteghem, C.: Post-column zirconium chelation and fluorescence detection for the liquid chromatographic determination of tetracyclines. *Anal. Chim. Acta*, 303 (1995) 11-16.
- 3542 Emara, S., Morita, I., Tamura, K., Razee, S., Masujima, T., Mohamed, H.A., El-Gizawy, S.M. and El-Rabbat, N.A.: Simultaneous determination of free and bound adriamycin, adriamycinol, adriamycinone and daunorubicin in plasma using column-switching technique and protein-coated pre-columns. *Talanta*, 41 (1994) 1973-1980; *C.A.*, 122 (1995) 71228d.
- 3543 Fagan, S.V.B., Gombatz, C., Abdel-Kader, H. and Menon, G.: Determination of ardacin in various silage food diets by a rapid liquid chromatographic assay. *J. Assoc. Off. Anal. Chem.*, 78 (1995) 16-21.
- 3544 Gräfe, U., Ihn, W., Ritzau, M., Schade, W., Stengel, C., Fleck, W.F., Künkel, W., Härtl, A. and Gutsche, W.: Helioferins, a novel antifungal lipopeptides from *Mycogone rosea*: screening, isolation, structures and biological properties. *J. Antibiot.*, 48 (1995) 126-133.
- 3545 Gräfe, U., Schlegel, R., Ritzau, M., Ihn, W., Dornberger, K., Stengel, C., Fleck, W.F., Gutsche, W., Härtl, A. and Paulus, E.F.R.: Aurantiamicins, new depsipeptide antibiotics from *Streptomyces aurantiacus* IMET 43917. Production, isolation, structure elucidation, and biological activity. *J. Antibiot.*, 48 (1995) 119-125.
- 3546 Guyonnet, J., Richard, M. and Hellings, P.: Determination of fumagillin in muscle tissue of rainbow trout using automated ion-pairing liquid chromatography. *J. Chromatogr. B*, 666 (1995) 354-359.
- 3547 Hormazabal, V. and Yndestad, M.: Rapid assay for monitoring residues of enrofloxacin in milk and meat tissues by HPLC. *J. Liq. Chromatogr.*, 17 (1994) 3775-3782.
- 3548 Ikai, Y., Oka, H., Hayakawa, J., Matsumoto, M., Saito, M., Harada, K.-i., Mayumi, T. and Suzuki, M.: Total structures and antimicrobial activity of bacitracin minor components. *J. Antibiot.*, 48 (1995) 233-242.
- 3549 Itoh, T. and Yamada, H.: Diastereomeric β -lactam antibiotics. Analytical methods, isomerization and stereoselective pharmacokinetics. *J. Chromatogr. A*, 694 (1995) 195-208 - a review with 37 refs.
- 3550 Kobayashi, H., Sunaga, R., Furihota, K., Morisaki, N. and Iwasaki, S.: Isolation and structures of an antifungal antibiotic, fusarieline A, and related compounds produced by a *Fusarium* sp. *J. Antibiot.*, 48 (1995) 42-52.
- 3551 Kominato, K., Watanabe, Y., Hirano, S.-i., Kioka, T., Terasawa, T., Yoshioka, T., Okamura, K. and Tone, H.: Mer-A2026A and B, novel piericidins with vasodilating effect I. Producing organism, fermentation, isolation and biological properties. *J. Antibiot.*, 48 (1995) 106-112.
- 3552 La Follette, G., Jayewardene, A.L., Seneviratne, A.K., Lin, E.T. and Gambertoglio, J.G.: Determination of ticarcillin in human plasma and in urine by reversed-phase LC. *J. Pharm. Biomed. Anal.*, 13 (1995) 159-164.
- 3553 Lau, R.C.M. and Rinehart, K.L.: Berninamycins B, C and D, minor metabolites from *Streptomyces bernensis*. *J. Antibiot.*, 47 (1994) 1466-1472.
- 3554 Lee, C.-H., Kim, S., Hyun, B., Suh, J.-W., Yon, C., Kim, C., Lim, C. and Kim, C.: Cepacidina A, a novel antifungal antibiotic produced by *Pseudomonas cepacia*. I. Taxonomy, production, isolation and biological activity. *J. Antibiot.*, 47 (1994) 1402-1405.
- 3555 Mathew, M. and Das Gupta, V.: Stability of vancomycin hydrochloride solutions at various pH values as determined by high-performance liquid chromatography. *Drug. Dev. Ind. Pharm.*, 21 (1995) 257-264; *C.A.*, 122 (1995) 114783a.

- 3556 Matsumoto, K., Tanka, K., Matsutani, S., Sakazaki, R., Hino, H., Uotani, N., Tanimoto, T., Kawamura, Y., Nakamoto, S. and Yoshida, T.: Isolation and biological activity of thielocins: novel phospholipase A₂ inhibitors produced by *Thielavia terricola* RF-143. *J. Antibiot.*, 48 (1995) 106-112.
- 3557 McMullin, C.M., White, L.O., MacGowan, A.P., Holt, H.A., Lovering, A.M. and Reeves, D.S.: Assay of serum teicoplanin concentrations in clinical specimens: a comparison of isocratic high performance liquid chromatography with polarization fluoroimmunoassay and bioassay. *J. Antimicrob. Chemother.*, 34 (1994) 425-429; *C.A.*, 122 (1995) 45512x.
- 3558 Mikisch, E.: (Significance of the 4-epimers in the determination of oxytetracycline, tetracycline, and chlorotetracycline residues in animal-derived food, by HPLC). *Dtsch. Lebensm.-Rundsch.*, 90 (1994) 39-41; *C.A.*, 122 (1995) 54280x.
- 3559 Moats, W.A. and Harik-Khan, R.: Liquid chromatographic determination of β -lactam antibiotics in milk: a multiresidue approach. *J. Assoc. Off. Anal. Chem.*, 78 (1995) 49-54.
- 3560 Nishikawa, T., Suzuki, S. and Ohtani, H.: Reversible isomerization of rapamycin demonstrated by liquid chromatography. *J. Antibiot.*, 47 (1994) 1554-1556.
- 3561 Nozawa, O., Okazaki, T., Sakai, N., Komurasaki, T., Hanada, K., Morimoto, S., Chen, Z.-X., He, P.-M. and Mizoue, K.: A novel bioactive δ lactone FD-211. Taxonomy, isolation and characterization. *J. Antibiot.*, 48 (1995) 113-118.
- 3562 Paesen, J., Claeys, P., Cypers, W., Roets, E. and Hoogmartens, J.: Liquid chromatography of tylosin A and related substances on poly(styrene-divinylbenzene). *J. Chromatogr. A*, 699 (1995) 93-97.
- 3563 Paesen, J., Cypers, W., Busson, R., Roets, E. and Hoogmartens, J.: Isolation of decomposition products of tylosin using liquid chromatography. *J. Chromatogr. A*, 699 (1995) 99-106.
- 3564 Petersen, F., Fredenhagen, A., Mett, H., Lydon, N.B., Delmendo, R., Jenny, H.-B. and Peter, H.H.: Paeciloliquinones A, B, C, D, E and F: new potent inhibitors of protein tyrosine kinases produced by *Paecilomyces carneus*. I. Taxonomy, fermentation, isolation and biological activity. *J. Antibiot.*, 48 (1995) 191-198.
- 3565 Proksa, B. and Fuska, J.: Role of aliphatic acids in the biosynthesis of vermiculin in *Penicillium vermiculatum*. *Pharmazie*, 50 (1995) 215-216.
- 3566 Reeves, V.S.: Liquid chromatographic procedure for the determination of novobiocin residues in bovine milk: interlaboratory study. *J. Assoc. Off. Anal. Chem.*, 78 (1995) 55-58.
- 3567 Saitoh, K., Furumai, T., Oki, T., Nishida, F., Harada, K.-i. and Suzuki, M.: Pradimicin S, a new prodimicin analog. III. Application of the Frit-FAB LC/MS technique to the elucidation of the pradimicin S biosynthetic pathway. *J. Antibiot.*, 48 (1995) 162-168.
- 3568 Sancho-Chust, V., Fabra-Campos, S., Gómez-Meseguer, V., Bengochea, M. and Martín-Villodre, A.: Experimental studies on the influence of surfactants on intestinal absorption of drugs. Cefadroxil as model drug and sodium lauryl sulfate as model surfactant: studies in rat. *Arzneim.-Forsch.*, 45 (1995) 595-601.
- 3569 Seki-Asano, M., Okazaki, T., Yamagishi, M., Sakai, N., Takayama, Y., Hanada, K., Morimoto, S., Takatsuki, A. and Mizoue, K.: Isolation and characterization of a new 12-membered macrolide FD-895. *J. Antibiot.*, 47 (1994) 1395-1401.
- 3570 Tyczkowska, K.L., Voyksner, R.D., Straub, R.F. and Aronson, A.L.: Simultaneous multiresidue analysis of β -lactam antibiotics in bovine milk by liquid chromatography with ultraviolet detection and confirmation by electrospray mass spectrometry. *J. Assoc. Off. Anal. Chem.*, 77 (1994) 1122-1131.
- 3571 Vickers, A.E.M., Fisher, R.L., Brendel, K., Guertler, J., Dannecker, R., Keller, B. and Fisher, V.: Sites of biotransformation for the cyclosporin derivative SD2 IMM 125 using human liver and kidney slices and intestine. Comparison with rat liver slices and cyclosporin A metabolism. *Drug Metab. Disp.*, 23 (1995) 327-333.
- 3572 Vickers, A.E.M., Meyer, E., Dannecker, R., Keller, B., Tynes, R.E. and Maurer, G.: Human liver cytochrome P4503A biotransformation of the cyclosporin derivative SD2 IMM 125. *Drug Metab. Disp.*, 23 (1995) 321-326.
- 3573 Yang, C.F., Stassinopoulos, A. and Goldberg, I.H.: Specific binding of the biradical analog of neocarzinostatin chromophore to bulged DNA: implications for thiol-independent cleavage. *Biochemistry*, 34 (1995) 2267-2275.

For additional information see C.A.:

122 (1995) 115150d, 204532x, 209170n.

See also 3741, 3766.

29. INSECTICIDES, PESTICIDES AND OTHER AGROCHEMICALS

29a. General techniques

- 3574 Beneventi, G., Barbieri, C., del Carlo, G., Dondi, C. and Forti, S.: (Pesticides in wine: simultaneous determination of ethylenethiourea (ETU), propylenethiourea (PTU), fungicides and insecticides by SPE extraction with a C18 cartridge). *Ind. Bevande*, 23 (1994) 317-323; *C.A.*, 122 (1995) 54326s.
- 3575 Holstege, D.M., Scharberg, D.L., Tor, E.R., Hart, L.C. and Galey, F.D.: A rapid multiresidue screen for organophosphorus, organochlorine, and N-methyl carbamate insecticides in plant and animal tissues. *J. Assoc. Off. Anal. Chem.*, 77 (1994) 1263-1274.
- 3576 Marcé, R.M., Prosen, H., Crespo, C., Calull, M., Borrull, F. and Brinkman, U.A.T.: On-line trace enrichment of polar pesticides in environmental waters by reversed-phase liquid chromatography-diode array detection-particle beam mass spectrometry. *J. Chromatogr. A*, 696 (1995) 63-74.
- 3577 Newsome, W.H.: An evaluation of pesticide residue methodology. *J. Assoc. Off. Anal. Chem.*, 78 (1995) 4-8.
- 3578 Nouri, B., Fouillet, B., Toussaint, G., Chambon, P. and Chambon, R.: High-performance liquid chromatography with diode-array detection for the determination of pesticides in water using automated solid-phase extraction. *Analyst (Cambridge)*, 120 (1995) 1133-1136.
- 3579 Pffab, G. and York, H.: Application of AMD (automated multiple development) for the determination of pesticides in drinking water. Part 3. Solid phase extraction and influencing factors. *Acta Hydrochim. Hydrobiol.*, 22 (1994) 216-223; *C.A.*, 122 (1995) 88874j.

- 3580 Sannino, A.: Investigation into contamination of processed fruit products by carbendazim, methyl thiophanate and thiazobenzazole. *Food Chem.*, 52 (1995, Pub. 1994) 57-61; *C.A.*, 122 (1995) 79366t.
- 3581 Sennert, S., Volmer, D., Levsen, K. and Wünsch, G.: Multi-residue analysis of polar pesticides in surface and drinking water by on-line enrichment and thermospray LC-MS. *Fresenius J. Anal. Chem.*, 351 (1995) 642-649.
- 3582 Volmer, D., Levsen, K. and Engewald, W.: Analysis of polar pesticides in aqueous samples by combined online trace enrichment and thermospray liquid chromatography mass spectrometry. *Vom Wasser*, 82 (1994) 335-384; *C.A.*, 122 (1995) 114385x.
- 3583 Wilson, S.B. and Rontree, J.: (Analysis of pesticides by electro-spray LC-MS: a sensitive, specific, routine and robust analysis technique). *Foods Food Ingredients J. Jpn.*, 162 (1994) 81-86; *C.A.*, 122 (1995) 158873j.
- 3584 Yamamoto, A., Matsunaga, A., Ohto, M., Mizukami, E., Hayakawa, K. and Miyazaki, M.: Real-time analysis of multicomponent chromatograms: application to high-performance liquid chromatography. *Analyst (Cambridge)*, 120 (1995) 377-380.

For additional information see *C.A.*:

122 (1995) 48963f, 100762f, 212345d.

See also 3824, 3826.

29c. Phosphorus insecticides

For additional information see *C.A.*:

122 (1995) 99232z.

29d. Carbamates

- 3585 Sundaram, K.M.S. and Curry, J.: Fluorometric determination of aminocarb and mexacarbamate and some of their metabolites by liquid chromatography: influence of structural factors on fluorescence intensity. *J. Liq. Chromatogr.*, 17 (1994) 3759-3774.

For additional information see *C.A.*:

122 (1995) 29977x.

29e. Herbicides

- 3586 Brandsteterova, E., Kralikova, D., Lehotay, J. and Tekel, J.: HPLC analysis of some substituted anilines and phenylurea herbicides in milk. *Nahrung*, 38 (1994) 439-441; *C.A.*, 122 (1995) 54311h.
- 3587 Chiron, S., Papilloud, S., Haerdi, W. and Barceló, D.: Automated on-line liquid-solid extraction followed by liquid chromatography - high-flow pneumatically assisted electrospray mass spectrometry for the determination of acidic herbicides in environmental waters. *Anal. Chem.*, 67 (1995) 1637-1643.
- 3588 Reichert, J., Gernikeites, T. and Winkler, M.: (Application of HPLC with fluorescence detection to environmental analysis exemplified with the herbicide dalapon). *Vom Wasser*, 82 (1994) 37-48; *C.A.*, 122 (1995) 114387z.

29f. Fungicides

- 3589 Bushway, R.J., Young, B.E.S., Paradis, L.R. and Perkins, L.B.: Determination of thiabendazole in fruits and vegetables by competitive-inhibition enzyme immunoassay. *J. Assoc. Off. Anal. Chem.*, 77 (1994) 1243-1248.
- 3590 Bushway, R.J., Young, B.E.S., Paradis, L.R., Perkins, L.B., Martin, S.K. and Brown, M.P.: Determination of methyl 2-benzimidazolecarbamate in bulk fruit juice concentrates by competitive-inhibition enzyme immunoassay. *J. Assoc. Off. Anal. Chem.*, 77 (1994) 1237-1243.

29g. Other types of pesticides and various agrochemicals

- 3591 Blanco Gomis, D., Castano Fernandez, A., Megido Bernardo, V. and Gutierrez Alvarez, M.D.: High performance liquid chromatographic determination of cymiazole in honey with UV and electrochemical detection. *Chromatographia*, 39 (1994) 602-606.
- 3592 Sundaram, K.M.S., Sloane, L. and Curry, J.: Kinetics of azadirachtin hydrolysis in model aquatic systems by high-performance liquid chromatography. *J. Liq. Chromatogr.*, 18 (1995) 363-376.
- 3593 Tamiya, M., Kashiwabara, F., Watanabe, Y., Ando, T., Tsutsumi, T., Norizuki, H., Minamisawa, M. and Hisai, S.: (Development of an analytical method for methoprene in rice by HPLC). *Shokuhin Eiseigaku Zasshi*, 35 (1994) 593-598; *C.A.*, 122 (1995) 131286h.

See also 3808.

30. SYNTHETIC AND NATURAL DYES

30a. Synthetic dyes

- 3594 Castle, M. and Neuteboom, E.: High-performance liquid chromatography of the fluorescent dyes Fura-2 and Mag-Fura. Stability in organic solvents. *J. Chromatogr. A*, 696 (1995) 93-99.
- 3595 Roybal, J.E., Pfenning, A.P., Munns, R.K., Holland, D.C., Hurlbut, J.A. and Long, A.R.: Determination of malachite green and its metabolite, leucomalachite green, in catfish (*Ictalurus punctatus*) tissue by liquid chromatography with visible detection. *J. Assoc. Off. Anal. Chem.*, 78 (1995) 453-457.

See also 2628, 3000, 3005.

30b. Chloroplast and other natural pigments

- 3596 Bento, L.S.M.: Separation of sugar colorants using chromatographic columns containing sucrose crystals. *Publ. Tech. Pap. Proc. Annu. Meet. Sugar Ind. Technol.*, 53 (1994) 17-39; *C.A.*, 122 (1995) 109191b.
- 3597 Gao, L. and Mazza, G.: Characterization of acetylated anthocyanins in lowbush blueberries. *J. Liq. Chromatogr.*, 18 (1995) 245-259.
- 3598 Klein, R.: Identification, stereoconfiguration, chromatographic and fluorescence properties of natural pterins. *Adv. Exp. Med. Biol.*, 338(Chemistry and Biology of Pteridines and Folates) (1993) 43-46; *C.A.*, 122 (1995) 155614q.

- 3599 Mínguez-Mosquera, M.I., Hornero-Méndez, D. and Garrido-Fernández, J.: Detection of bixin, lycopene, cantaxanthin, and β -apo-8'-carotenal in products derived from red pepper. *J. Assoc. Off. Anal. Chem.*, 78 (1995) 491-498.
- 3600 Moskalenko, A.A., Toropygina, O.A. and Kuznetsova, N.Yu.: (Pigment-protein complexes of *Chromatium minutissimum* with extracted carotenoids). *Biokhimiya (Moscow)*, 59 (1994) 1414-1420.
- 3601 Szabo, G., Prosser, S.L. and Bulman, R.A.: Extraction and high-performance size-exclusion chromatography of humic substances from soils. In: Kalasz, H., Ettre, L.S. and Pick, J. (Editors), *New Approaches Chromatogr. '93, [Pap. Budapest Chromatogr. Conf.]*, 1990, Fekete Sas Diado, Budapest, 1993, pp. 171-186; *C.A.*, 122 (1995) 212895h.
- 3602 Van Heukelem, L., Lewitus, A.J., Kana, T.M. and Craft, N.E.: Improved separations of phytoplankton pigments using temperature-controlled high performance liquid chromatography. *Mar. Ecol.: Prog. Ser.*, 114 (1994) 303-313; *C.A.*, 122 (1995) 155379s.
- For additional information see *C.A.*:
122 (1995) 158879r.
- See also 3514, 3518, 3519, 3520, 3527, 3803.
31. PLASTICS AND THEIR INTERMEDIATES
- 3603 Allan, D.S., Birchmeier, M., Pribish, J., Priddy, D.B. and Smith, P.: Application of GPC-UV/VIS analysis to investigate the mechanism of styrene-co-acrylonitrile thermal discoloration. *Polym. Prepr. (Am. Chem. Soc., Div. Polym. Chem.)*, 34 (1993) 258-259; *C.A.*, 121 (1994) 10841f.
- 3604 Bullock, J.: High-temperature reversed-phase high-performance liquid chromatographic analysis of a synthetic copolymer on a non-porous support. *J. Chromatogr. A*, 694 (1995) 415-423.
- 3605 Cutie, S.S. and Martin, S.J.: Size-exclusion chromatography of crosslinked superabsorbent polymers. *J. Appl. Polym. Sci.*, 55 (1995) 605-609; *C.A.*, 122 (1995) 161960r.
- 3606 Glöckner, G., Wolf, D. and Engelhardt, H.: Control of adsorption and solubility in gradient high-performance liquid chromatography. Part V. Separation of styrene/methyl methacrylate copolymers by sudden-transition gradients in normal-phase and reversed-phase mode. *Chromatographia*, 39 (1994) 557-563.
- 3607 Henry, R.A., Noguchi, K. and Stevenson, R.: Solvent scouting in steric exclusion chromatography. Characterization of poly(vinyl acetate) and poly(vinyl alcohol) on the same organic gel column. *Am. Lab. (Shelton)*, 25 (1993) 36Z, 368B-368D; *C.A.*, 120 (1994) 299734k.
- 3608 Janco, M., Proudkova, T. and Berek, D.: Liquid chromatography of polymer mixtures applying combination of exclusion and full adsorption mechanisms. I. Analysis of polystyrene in its mixture with poly(methyl methacrylate) single column/single eluent approach. *J. Appl. Polym. Sci.*, 55 (1995) 393-397; *C.A.*, 122 (1995) 161958w.
- 3609 Kuzaev, A.I., Ol'khova, O.M., Solomko, S.I. and Baturin, S.M.: (Use of liquid chromatography in the critical region for separation of oligo(epichlorohydrins) with respect to functionality types). *Zh. Fiz. Khim.*, 68 (1994) 1794-1798; *C.A.*, 122 (1995) 106976a.
- 3610 Lehrle, R.S. and Sarson, D.S.: Polymer molecular weight distribution: results from matrix-assisted laser desorption ionization compared with those from gel-permeation chromatography. *Rapid Commun. Mass Spectrom.*, 9 (1995) 91-92; *C.A.*, 122 (1995) 162029f.
- 3611 Martin, L. and Balke, S.T.: Cyclic oligomer considerations in the size exclusion chromatography of poly(ethylene terephthalate). *J. Liq. Chromatogr.*, 18 (1995) 1-15.
- 3612 Mendichi, R., Audisio, G. and Giacometti Schieroni, A.: Molecular mass characterization of poly(4-vinylpyridine). *J. Liq. Chromatogr.*, 17 (1994) 3323-3331.
- 3613 Nagy, D.J.: Characterization of poly(vinyl alcohol) using SEC multiangle laser light scattering. *Int. Lab.*, 25, No. 2 (1995) P6-P11.
- 3614 Okada, T.: Multifunctional separation with polyamine-bonded resin. *Anal. Chim. Acta*, 303 (1995) 193-197.
- 3615 Pasch, H. and Rode, K.: Use of matrix-assisted laser desorption/ionization mass spectrometry for molar mass-sensitive detection in liquid chromatography of polymers. *J. Chromatogr. A*, 699 (1995) 21-29.
- 3616 Shintani, H.: HPLC analysis of toxic additives and residual monomer from dental plate. *J. Liq. Chromatogr.*, 18 (1995) 613-626.
- 3617 Staal, W.J., Cools, P., van Herk, A.M. and German, A.L.: Online extraction of polymers, oligomers, additives and monomers by multiple solvents on packed HPLC columns. *Process Technol. Proc.*, 11 (1994) 857-859; *C.A.*, 122 (1995) 162162u.
- 3618 Sturman, H.W., Geissler, M., Weimar, A., Paul, A. and Nickel, B.: (High performance liquid chromatography (HPLC) and gas chromatography (GC) as methods in plastic material analysis). *Kausch. Gummi Kunstst.*, 47 (1994) 812-815; *C.A.*, 122 (1995) 107514k.
- 3619 Terneuil, G., Hotier, G. and Lonchamp, D.: Chromatographic separation of mixtures of polyglycerols. *Ger. Offen. DE 4,404,291 (Cl. C07C43/10)*, 18 Aug. 1994, FR Appl. 93/1,755, 15 Feb. 1993; 9 p.; *C.A.*, 122 (1995) 82342n.
- 3620 Van Every, K.W. and Long, W.J.: High-temperature GPC/IR characterization of polyolefines. *Proc. SPIE-Int. Soc. Opt. Eng.*, 2089 (9th International Conference on Fourier Transform Spectroscopy, 1993) (1993) 186-187; *C.A.*, 122 (1995) 82482h.
- For additional information see *C.A.*:
122 (1995) 82466f, 107275h, 204273p.
- See also 2671, 2680, 2757, 3003, 3841.

32. DRUG ANALYSIS

32a. Drug analysis, general techniques

- 3621 Balogh, G., Csizer, E., Ferenczy, G.G., Halmos, Z., Herenyi, B., Horvath, P., Lauko, A. and Gorog, S.: Estimation of impurity profiles of drugs and related materials. 12. Isolation and identification of an isomeric impurity in danazol. *Pharm. Res.*, 12 (1995) 295-298; *C.A.*, 122 (1995) 197141p.
- 3622 Bieganowska, M.L. and Petruczynik, A.: Ion pair chromatography and its application in analysis of pharmaceuticals. *Chem. Anal. (Warsaw)*, 39 (1994) 525-541; *C.A.*, 122 (1995) 26925t.
- 3623 Caldwell, J.: Stereochemical determinants of the nature and consequences of drug metabolism. *J. Chromatogr. A*, 694 (1995) 39-48 - a review with 47 refs.
- 3624 Glazova, N.V., Efimova, T.B. and Dmitrenko, L.V.: (Use of gel chromatography for quantitative analysis of pyrogenic impurities in drugs). *Zh. Fiz. Khim.*, 68 (1994) 1803-1805; *C.A.*, 122 (1995) 89575z.
- 3625 Gluck, S.J., Martin, E. and Benko, M.H.: Centrifugal partition chromatography for the determination of octanol-water partition coefficients. *Chromatogr. Sci. Ser.*, 68(Centrifugal Partition Chromatography) (1995) 199-218; *C.A.*, 122 (1995) 155377q.
- 3626 Hermansson, J. and Grahn, A.: Optimization of the separation of enantiomers of basic drugs. Retention mechanisms and dynamic modification of the chiral bonding properties on an α_1 -acid glycoprotein column. *J. Chromatogr. A*, 694 (1995) 57-69.
- 3627 Herraiz-Hernández, R., van de Merbel, N.C. and Brinkman, U.A.T.: Determination of the total concentration of highly protein-bound drugs in plasma by on-line dialysis and column liquid chromatography: application to non-steroidal anti-inflammatory drugs. *J. Chromatogr. B*, 666 (1995) 127-137.
- 3628 Itoh, H., Kinoshita, T. and Nimura, N.: (Direct serum injection analysis on ODS silica gel column). *Kuromatogurafi*, 15 (1994) 256-257; *C.A.*, 122 (1995) 127791b.
- 3629 Leahy, D.E., Lynch, J., Finney, R.E. and Taylor, D.C.: Estimation of sieving coefficients of convective absorption of drugs in perfused rat jejunum. *J. Pharmacokin. Biopharm.*, 22 (1994) 411-429.
- 3630 Pidgeon, C., Orig, S., Liu, H., Qiu, X., Pidgeon, M., Dantzig, A.H., Munroe, J., Hornback, W.J., Kasher, J.S. et al.: IAM chromatography: an *in vitro* screen for predicting drug membrane permeability. *J. Med. Chem.*, 38 (1995) 590-594; *C.A.*, 122 (1995) 95739x.
- 3631 Qin, Z.: Advances in biopharmaceutical analysis in the People's Republic of China: 1991-1993. *J. Pharm. Biomed. Anal.*, 13 (1995) 1-7 - a review with 58 refs.
- 3632 Steele, K.M. and Lunte, C.E.: Microdialysis sampling coupled to on-line microbore liquid chromatography for pharmacokinetic studies. *J. Pharm. Biomed. Anal.*, 13 (1995) 149-154.
- 3633 Wainer, I.W., Ducharme, J., Granvil, C.P., Parenteau, H. and Abdullah, S.: Using chirality as a unique probe of pharmacological properties. *J. Chromatogr. A*, 694 (1995) 169-179.
- 3634 Zeng, S.: (Separations of drug enantiomers by high-performance liquid chromatography with chiral mobile phase additives). *Sepu*, 13 (1995) 21-25; *C.A.*, 122 (1995) 197088b - a review with 47 refs.
- 3635 Zeng, S., Shen, X. and Liu, Z.: (Pharmacology of chiral pharmaceuticals and HPLC determination of their enantiomers *in vivo*). *Zhejiang Yike Daxue Xuebao*, 23 (1994) 234-238; *C.A.*, 122 (1995) 95711g.

For additional information see *C.A.*:

122 (1995) 122271x, 229996q, 230008a.

See also 2575, 2613, 2722, 2727, 2741, 2748, 3780, 3781, 3796, 3831.

32b. Antirheumatics and antiinflammatory drugs

- 3636 Azzolina, O. and Collina, S.: Chiral recognition of alkyl 2-aryloxypropionates by HPLC. *J. Liq. Chromatogr.*, 18 (1995) 81-92.
- 3637 Blondino, F.E. and Byron, P.R.: The quantitative determination of aspirin and its degradation products in a model solution aerosol. *J. Pharm. Biomed. Anal.*, 13 (1995) 111-119.
- 3638 Caccamese, S., Principato, G., Ottanà, R., Previtara, T. and Zappalà, C.: Liquid chromatographic separation of the enantiomers of antihistaminic 3,3'-di(1,3-thiazolidin-4-one) derivatives with two and four stereogenic centres. *J. Chromatogr. A*, 694 (1995) 355-363.
- 3639 Daffonchio, L., Bestetti, A., Clavenna, G., Fedele, G., Ferrari, M.P. and Omini, C.: Effect of a new foam formulation of keto-profen lysine salt in experimental models of inflammation and hyperalgesia. *Arzneim.-Forsch.*, 45 (1995) 590-594.
- 3640 Deigner, H.P., Freyberg, C.E. and Laufer, S.: Distribution and excretion of [14 C]-labelled [2,2-dimethyl-6-(4-chlorophenyl)-7-phenyl-2,3-dihydro-1H-pyrrolizine-5-yl]-[2'- 14 C]-acetic acid in rats. *Arzneim.-Forsch.*, 45 (1995) 272-276.
- 3641 Ding, A., Zia-Amirhosseini, P., McDonagh, A.F., Burlingame, A.L. and Benet, L.Z.: Reactivity of tolmetin glucuronide with human serum albumin. Identification of binding sites and mechanisms of reaction by tandem mass spectrometry. *Drug Metab. Disp.*, 23 (1995) 369-376.
- 3642 Hamdoune, M., Mounie, J., Magdalou, J., Masmoudi, T., Goudonnet, H. and Escousse, A.: Characterization of the *in vitro* glucuronidation of flurbiprofen enantiomers. *Drug Metab. Disp.*, 23 (1995) 343-348.
- 3643 Hartmann, D., Stief, G., Lingenfelder, M., Güzelman, C. and Horsch, A.K.: Study on the possible interaction between tenoxicam and atenolol in hypertensive patients. *Arzneim.-Forsch.*, 45 (1995) 494-498.
- 3644 Mason, J.L. and Hobbs, G.J.: Simple method for the analysis of tenoxicam in human plasma using high-performance liquid chromatography. *J. Chromatogr. B*, 665 (1995) 410-415.
- 3645 Maya, M.T., Pais, J.P. and Morais, J.A.: A rapid method for the determination of piroxicam in plasma by high-performance liquid chromatography. *J. Pharm. Biomed. Anal.*, 13 (1995) 319-322.

- 3646 Teffera, Y. and Abramson, F.: Application of high-performance liquid chromatography/chemical reaction interface mass spectrometry for the analysis of conjugated metabolites: a demonstration using deuterated acetaminophen. *Biol. Mass Spectrom.*, 23 (1994) 776-783; C.A., 122 (1995) 45531c.
- 3647 Velic, I., Metzler, M., Hege, H.G. and Weymann, J.: Separation and identification of phase I and phase II [¹⁴C]antipyrine metabolites in rat and dog urine. *J. Chromatogr. B*, 666 (1995) 139-147.
- 3648 Volin, P.: Simple and specific reversed-phase liquid chromatographic method with diode-array detection for simultaneous determination of serum hydroxychloroquine, chloroquine and some corticosteroids. *J. Chromatogr. B*, 666 (1995) 347-353.
- For additional information see C.A.:
122 (1995) 89603g, 170366d, 170370a, 170372c, 170373d.
- See also 2741, 2893, 2925, 3008, 3627, 3711, 3755.
- 32c. *Autonomic and cardiovascular drugs*
- 3649 Berndt, A., Oertel, R., Richter, K., Terhaag, B. and Gramatté, T.: Relative Bioverfügbarkeit des Antiarrhythmikums Tircacizin und seiner drei Metabolite. *Arzneim.-Forsch.*, 45 (1995) 250-253.
- 3650 Cavazzutti, G., Gagliardi, L., de Orsi, D. and Tonelli, D.: Simultaneous determination of buzipide, phenylpropanolamine, and clocinizine in pharmaceutical preparations by ion-pair reversed-phase HPLC. *J. Liq. Chromatogr.*, 18 (1995) 227-234.
- 3651 Chatterjee, D.J., Li, W.Y., Hurst, A.K. and Koda, R.T.: High-performance liquid chromatographic method for determination of atenolol from human plasma and urine: simultaneous fluorescence and ultraviolet detection. *J. Liq. Chromatogr.*, 18 (1995) 791-806.
- 3652 Chen, C.L., Sangiah, S., Chen, H., Berlin, K.D., Garrison, G.L., Scherlag, B.J. and Lazzara, R.: Simultaneous determination of a novel antiarrhythmic agent, 7-benzyl-3-thia-7-azabicyclo[3.3.1]nonane and its sulfoxidation metabolite in plasma and urine by HPLC. *J. Liq. Chromatogr.*, 17 (1994) 3681-3694.
- 3653 Chilton, A.S., Godward, R.E. and Carey, P.F.: The determination in human plasma of 1-hydroxy-2-naphthoic acid following administration of salmeterol xinafoate. *J. Pharm. Biomed. Anal.*, 13 (1995) 165-169.
- 3654 Cieri, U.R.: Determination of reserpine, hydralazine HCl, and hydrochlorothiazide in tablets by liquid chromatography on a short, normal-phase column. *J. Assoc. Off. Anal. Chem.*, 77 (1994) 1104-1108.
- 3655 El Waliy, A.F.M., El-Yazbi, F.A., Belal, S.F. and Abdel Razak, O.: High-performance liquid chromatographic and derivative ultraviolet spectrophotometric determination of pindolol and clopamide mixture in tablets. *Anal. Lett.*, 28 (1995) 893-907.
- 3656 Elm, T. and Hansen, E.L.: Simultaneous determination of lofepramine and desipramine by a high-performance liquid chromatographic method used for therapeutic drug monitoring. *J. Chromatogr. B*, 665 (1995) 355-361.
- 3657 Frey, H.O., Kinkeldei, J. and Quast, M.: (Simple and rapid determination for determining the anticonvulsant lamotrigine). *Labor-Med.*, 17 (1994) 166-168; C.A., 122 (1995) 71213v.
- 3658 Herrmann, R.: Automated HPLC assay for sotalol in human plasma. *J. Pharm. Biomed. Anal.*, 13 (1995) 329-333.
- 3659 Hutt, V., Pabst, G., Salama, Z., Kappler, J. and Jaeger, H.: Untersuchungen zur Pharmakokinetik und Bioverfügbarkeit einer neuen Mexiletin-Zubereitung und gesunden Freiwilligen. *Arzneim.-Forsch.*, 45 (1995) 254-257.
- 3660 Jahn, P., Eckrich, C., Schneidrowski, B., Volz-Zang, C., Schulte, B., Mutschler, E. and Palm, D.: β_1 -Adrenoceptor subtype selective antagonism of esmolol and its major metabolite *in vitro* and in man. Investigations using tricresylphosphate as red blood cell carboxylesterase inhibitor. *Arzneim.-Forsch.*, 45 (1995) 536-541.
- 3661 Koytchev, R., Alken, R.-G., Mayer, O., Böhm, R., Ellrich, A. and Waldner-Kölblin, R.G.: Untersuchungen zur Bioäquivalenz von zwei oralen Propafenon-Zubereitungen. *Arzneim.-Forsch.*, 45 (1995) 542-545.
- 3662 Kugler, A.R., Olson, S.C. and Smith, D.E.: Determination of quinapril and quinaprilat by high-performance liquid chromatography with radiochemical detection, coupled to liquid scintillation counting spectrometry. *J. Chromatogr. B*, 666 (1995) 360-367.
- 3663 Leube, J. and Fischer, G.: Determination of the renin inhibitor Ro 42-5892 in human plasma by automated pre-column derivatization, reversed-phase high-performance liquid chromatographic separation and electrochemical detection after post-column irradiation. *J. Chromatogr. B*, 665 (1995) 373-381.
- 3664 Lozano, R., Warren, F.V., Jr., Perlman, S. and Joseph, J.M.: Quantitative analysis of fosinopril sodium by capillary zone electrophoresis and liquid chromatography. *J. Pharm. Biomed. Anal.*, 13 (1995) 139-148.
- 3665 Misztal, G. and Przyborowski, L.: Determination of aprindine in human plasma using reversed phase HPLC. *Pharmazie*, 50 (1995) 187-188.
- 3666 Miyazaki, M., Hosoki, K., Okunishi, H., Ishii, K., Ikeno, A. and Okazaki, Y.: Anti-atherosclerotic and plasma lipid lowering effects of the novel calcium blocker with α_1 -adrenoreceptor antagonistic activity, monatepil, in high cholesterol diet-fed Japanese *Macaca fuscata* monkeys. *Arzneim.-Forsch.*, 44 (1994) 288-297.
- 3667 Mustafa, A.A., Al-Rashood, K.A., Hagga, M.E.M., Gad-Kariem, E.A. and Al-Awady, M.E.: High-performance liquid chromatographic and polarographic methods for the quantitation of nicardipine hydrochloride in capsules. *Pak. J. Sci. Ind. Res.*, 37 (1994) 1-4; C.A., 122 (1995) 142704u.
- 3668 Mutoh, M., Nakada, N., Matsukuma, S., Ohshima, S., Yoshinari, K., Watanabe, J. and Arisawa, M.: Panclitins, novel pancreatic lipase inhibitors. I. Taxonomy, fermentation, isolation and biological activity. *J. Antibiot.*, 47 (1994) 1369-1375.
- 3669 Niazy, E.M., El-Sayed, Y.M. and Khidr, S.H.: Analysis of prazosin in plasma by high-performance liquid chromatography using fluorescence detection. *J. Liq. Chromatogr.*, 18 (1995) 977-989.
- 3670 Pham-Huy, C., Radenen, B., Sahui-Gnassi, A. and Claude, J.-R.: High-performance liquid chromatographic determination of (S)- and (R)-propranolol in human plasma and urine with a chiral β -cyclodextrin bonded phase. *J. Chromatogr. B*, 665 (1995) 125-132.
- 3671 Ring, J.A., Ghabrial, H., Ching, M.S., Shulkes, A., Smallwood, R.A. and Morgan, D.J.: Fetal hepatic propranolol metabolism. Studies in the isolated fetal sheep liver. *Drug Metab. Disp.*, 23 (1995) 190-196.

- 3672 Sharma, S.C., Evans, M.B. and Evans, S.J.: The enantiomeric separation of metipranolol and desacetylmepipranolol on a cellulose tris-3,5-dimethylphenyl-carbamate chiral stationary phase. *J. Pharm. Biomed. Anal.*, 13 (1995) 129-137.
- 3673 Siret, L., Macaudiere, P., Bargmann-Leyder, N., Tambute, A., Caude, M. and Goudeon, E.: Separation of the optical isomers of a new 1,4-dihydropyridine calcium channel blocker (LF 2.0254) by liquid and supercritical fluid chromatography. *Chirality*, 6 (1994) 440-445; *C.A.*, 122 (1995) 150543f.
- 3674 Stearns, R.A., Chakravarty, P.K., Chen, R. and Chiu, S.-H.L.: Biotransformation of lasartan to its active carboxylic acid metabolites in human liver microsomes. Role of cytochrome P4502C and 3A subfamily members. *Drug Metab. Disp.*, 23 (1995) 207-215.
- 3675 Tokuma, Y. and Noguchi, H.: Stereoselective pharmacokinetics of dihydropyridine calcium antagonists. *J. Chromatogr. A*, 694 (1995) 181-193 - a review with 96 refs.
- 3676 Volz, M., Mitrovic, V., Thiemer, J. and Schlepper, M.: Steady-state plasma kinetics of slow-release propafenone, its two isomers and its main metabolites. *Arzneim.-Forsch.*, 45 (1995) 246-249.
- For additional information see *C.A.*:
122 (1995) 45519e, 45530b, 71230y, 197104d.
- See also 2617, 2741, 3372, 3551, 3643, 3793.
- 32d. Central nervous system drugs**
- 3677 Al-Khamis, K.I., El-Sayed, Y.M., Al-Rashood, K.A. and Bawazir, S.A.: High performance liquid chromatographic determination of ranitidine in human plasma. *J. Liq. Chromatogr.*, 18 (1995) 277-286.
- 3678 Amin, R.D., Cheng, H. and Rogers, J.D.: Determination of MK-0476 in human plasma by liquid chromatography. *J. Pharm. Biomed. Anal.*, 13 (1995) 155-158.
- 3679 Bacracheva, N., Drenska, A., Gorantcheva, J., Tyutyulkova, N. and Vlahov, V.: Comparative bioavailability of two oral metamizole formulations. Influence of the acetylation phenotype. *Arzneim.-Forsch.*, 45 (1995) 282-285.
- 3680 Bansal, R. and Aranda, J.V.: Simultaneous microassay of alfentanil, fentanil, and sufentanil by high performance liquid chromatography. *J. Liq. Chromatogr.*, 18 (1995) 339-348.
- 3681 Barrett, D.A. and Shaw, P.N.: A stability indicating HPLC assay for diamorphine in aqueous solution. *J. Liq. Chromatogr.*, 17 (1994) 3727-3733.
- 3682 Berliz, G., Wintersteiger, R. and Pacha, W.: Determination of a novel xanthine derivative, MKS 213-492, in plasma by high-performance liquid chromatography with electrochemical detection. *J. Chromatogr. B*, 666 (1995) 368-373.
- 3683 Boughton, B.J., Sheehan, T.M.T., Wood, J., O'Brien, D., Butler, M., Simpson, A. and Hale, K.A.: High-performance liquid chromatographic assay of plasma thalidomide: stabilization of specimens and determination of a tentative therapeutic range for chronic graft-versus-host disease. *Ann. Clin. Biochem.*, 32 (1995) 79-83; *C.A.*, 122 (1995) 204389f.
- 3684 Bowyer, J.F., Clauging, P. and Newport, G.D.: Determination of d-amphetamine in biological samples using high-performance liquid chromatography after precolumn derivatization with o-phthalaldehyde and 3-mercaptopropionic acid. *J. Chromatogr. B*, 666 (1995) 241-250.
- 3685 Browne, T.R., Szabo, G.K., Davoudi, H. and Josephs, E.G.: Determination of pharmacokinetic drug interactions with carbamazepine and phenytoin using stable isotope labeling and simple high-performance liquid chromatography/ultraviolet detection technique. *Neurology*, 44 (1994) 2410-2411; *C.A.*, 122 (1995) 122371e.
- 3686 Cairns, A.M., Chiou, R.H.-Y., Rogers, C.J.D. and Demetriades, J.L.: Enantioselective high-performance liquid chromatographic determination of omeprazole in human plasma. *J. Chromatogr. B*, 666 (1995) 323-328.
- 3687 Chauert, N., Li, C., Ducharme, Y., Trimble, L.A., Ramachandran, C. and Nicoll-Griffith, D.A.: *In vitro* and *in vivo* biotransformations of the naphthalenic lignan lactone 5-lipoxygenase inhibitor, L-702,539. *Drug Metab. Disp.*, 23 (1995) 65-71.
- 3688 Chen, X., Lu, J. and Li, D.: (Simultaneous determination of theophylline, phenobarbital, phenytoin, and carbamazepine in human serum by high-performance liquid chromatography). *Zhongguo Yiyuan Yaoxue Zazhi*, 14 (1994) 389-391; *C.A.*, 122 (1995) 122396s.
- 3689 Delon, A., Favreliere, S., Couet, W., Courtois, P. and Bouquet, S.: Rapid and sensitive determination of thalidomide in human plasma by high performance liquid chromatography. *J. Liq. Chromatogr.*, 18 (1995) 297-309.
- 3690 Dobrev, D., Berndt, A., Oertel, R., Richter, K., Terhaag, B. and Gramatté, T.: Untersuchung zur Bioäquivalenz von Metoclopramid in zwei Tablettenformulierungen. *Arzneim.-Forsch.*, 45 (1995) 304-306.
- 3691 El-Sayed, Y.M., Niazy, E.M. and Khidr, S.H.: High-performance liquid chromatographic method for the quantitative determination of pheniramine in plasma. *J. Liq. Chromatogr.*, 18 (1995) 763-777.
- 3692 Fadiran, E.O., Leslie, J., Fossler, M. and Young, D.: Determination of clozapine and its major metabolites in human serum and rat plasma by liquid chromatography using solid-phase extraction and ultraviolet detection. *J. Pharm. Biomed. Anal.*, 13 (1995) 185-190.
- 3693 Feng, M.R., Siersma, P.R., Strenkoski, C.A. and Wright, D.S.: Sensitive high-performance liquid chromatographic method for a dopamine receptor agonist, CI-1007, and its metabolite PD 147693 in monkey plasma. *J. Chromatogr. B*, 665 (1995) 193-199.
- 3694 Garner, D.M., Pieniaszek, H.J., Jr., Connell, J.M. and Fiske, W.D.: Determination of linopirdine and its mono-N-oxide metabolite in human plasma and urine by high-performance liquid chromatography. *J. Liq. Chromatogr.*, 17 (1994) 4541-4555.
- 3695 Jonczyk, A. and Wilczynska-Wojtulewicz, I.: Determination of diphenhydramine and naphazoline in pharmaceutical mixtures by high-performance liquid chromatography. *Acta Pol. Pharm.*, 51 (1994) 115-118; *C.A.*, 122 (1995) 88606k.
- 3696 Kapil, R.P., Lee, W., Jr., Padovani, P.K., Wong, N.Y., Holm, K.A. and Lam, G.N.: Determination of a novel cognitive enhancer, X9121, and its mono N-oxide metabolite, XG696, in dog plasma by reversed-phase high-performance liquid chromatography with ultraviolet detection. *J. Chromatogr. B*, 665 (1995) 226-231.

- 3697 Kondo, T., Hamasaki, K., Yamaguchi, M., Aoki, I., Yoshida, K., Yoshimura, Y. and Tanayama, S.: Enantioselective determination of pazinaclone, a new isoindoline anxiolytic, and its active metabolite in rat plasma by high-performance liquid chromatography. *J. Chromatogr. B*, 666 (1995) 291-297.
- 3698 Lacroix, P.M., Yat, P.N. and Lovering, E.G.: Liquid chromatographic methods for fluoxetine hydrochloride, its meta isomer, and related compounds in raw materials. *J. Assoc. Off. Anal. Chem.*, 78 (1995) 334-339.
- 3699 Lee, D.-S., Kwon, K.-H. and Shin, H.-S.: Determination of cimetidine in human urine by column switching HPLC and UV-detection. *Bull. Korean Chem. Soc.*, 16 (1995) 77-79; *C.A.*, 122 (1995) 177582e.
- 3700 Lehr, G.J., Yuen, S.M. and Lawrence, G.D.: Liquid chromatographic determination of atropine in nerve gas antidotes and other dosage forms. *J. Assoc. Off. Anal. Chem.*, 78 (1995) 339-343.
- 3701 Matsui, K., Oda, Y., Ohe, H., Tanaka, S. and Asakawa, N.: Direct determination of E2020 enantiomers in plasma by liquid chromatography-mass spectrometry and column-switching techniques. *J. Chromatogr. A*, 694 (1995) 209-218.
- 3702 Miratsuka, A., Chu, T.Y.Y., Distefano, E.W., Lin, L.Y., Schmitz, D.A. and Cho, A.K.: Inactivation of constitutive hepatic cytochromes P450 by phenylcyclidine in the rat. *Drug Metab. Disp.*, 23 (1995) 201-206.
- 3703 Naidong, W., Lee, J.W. and Hulse, J.D.: Development and validation of a chiral HPLC method for the quantitation of methocarbamol enantiomers in human plasma. *J. Liq. Chromatogr.*, 17 (1994) 3747-3758.
- 3704 Nishikawa, M., Tatsuno, M., Katagi, M. and Tsuchihashi, H.: (Determination of triazolam and its metabolites in human urine by EMIT, gas chromatography-mass spectrometry, and thermospray liquid chromatography-mass spectrometry). *Jpn. Toxicol. Environ. Health*, 41 (1995) 59-66; *C.A.*, 122 (1995) 232809m.
- 3705 Sueyasu, M., Ikeda, T., Otsubo, K., Taniyama, T., Aoyama, T. and Oishi, R.: Enantioselective determination of thiamylal in human serum by high-performance liquid chromatography. *J. Chromatogr. B*, 665 (1995) 133-137.
- 3706 Terhechte, A. and Blaschke, G.: Investigation of the stereoselective metabolism of the chiral H₁-antihistaminic drug terfenadine by high-performance liquid chromatography. *J. Chromatogr. A*, 694 (1995) 219-225.
- 3707 Torchio, L., Lombardi, F., Visconti, M. and Doyle, E.: Determination of the polar drug dimiracetam in human plasma and serum by column-switching high-performance liquid chromatography. *J. Chromatogr. B*, 666 (1995) 169-177.
- 3708 Tosunoglu, S. and Ersoy, L.: Determination of baclofen in human plasma and urine by high-performance liquid chromatography with fluorescence detection. *Analyst (Cambridge)*, 120 (1995) 373-375.
- 3709 Venkateshwaran, T.G. and Stewart, J.T.: HPLC determination of morphine-hydromorphone-bupivacaine and morphine-hydromorphone-tetracaine mixtures in 0.9% sodium chloride injection. *J. Liq. Chromatogr.*, 18 (1995) 565-578.
- 3710 Venkateshwaran, T.G., King, D.T. and Stewart, J.T.: HPLC determination of a metoclopramide and ondansetron mixture in 0.9% sodium chloride injection. *J. Liq. Chromatogr.*, 18 (1995) 117-126.
- 3711 Wailly, A.E., Belal, S.F., Yakout, M.H. and Zamel, S.: Determination of mixture of acetaminofen and ibuprofen in tablets using derivative spectrophotometry and high-performance liquid chromatography. *Bull. Fac. Pharm. (Cairo Univ.)*, 31 (1993) 331-337; *C.A.*, 122 (1995) 197127p.
- 3712 Wang, M.Z., Klee, M.S. and Yang, S.K.: Achiral and chiral analysis of camazepam and metabolites by packed-column supercritical fluid chromatography. *J. Chromatogr. B*, 665 (1995) 139-146.
- 3713 Wu, W.-N., Masucci, J.A., McKown, L.A. and Rodgers, J.D.: Identification of four plasma metabolites of etoperidone in dogs. *J. Pharm. Biomed. Anal.*, 13 (1995) 177-184.

For additional information see *C.A.*:

122 (1995) 23084g, 115098t, 115128c, 177562y, 204420j.

See also 2583, 2701, 2741, 3621, 3632, 3650, 3659, 3788.

32e. Chemotherapeutics (exc. cytostatics and antibiotics)

- 3714 Cannavan, A., Blanchflower, W.J. and Kennedy, D.G.: Determination of levamisole in animal tissues using liquid chromatography - thermospray mass spectrometry. *Analyst (Cambridge)*, 120 (1995) 331-333.
- 3715 Ficarra, R., Calabro, M.L., Tommasini, S., Grasso, S., Carulli, M., Monforte, A.M., Constantino, D. and Ficarra, P.: Enantiomeric resolution of anti-HIV agents on a cellulose derivative chiral stationary phase. Note II. *Chromatographia*, 40 (1995) 39-41.
- 3716 Flores-Murrieta, F.J., Granados-Soto, V. and Hong, E.: A simple and rapid method for determination of fluconazole in human plasma samples by high-performance liquid chromatography. *J. Liq. Chromatogr.*, 17 (1994) 3803-3811.
- 3717 Garcia-Alvarez-Coque, M.C., Simo-Alfonso, E.F., Ramis-Ramos, G. and Esteve-Romero, J.S.: High-performance micellar liquid chromatography determination of sulphonamides in pharmaceuticals after azodye precolumn derivatization. *J. Pharm. Biomed. Anal.*, 13 (1995) 237-245.
- 3718 Holland, D.C., Munns, R.K., Roybal, J.E., Hurlbut, J.A. and Long, A.R.: Liquid chromatographic determination of the anticoccidial drug halofuginone hydrobromide in eggs. *J. Assoc. Off. Anal. Chem.*, 78 (1995) 37-40.
- 3719 Horie, M., Saito, K., Nose, N. and Nakazawa, H.: (Simultaneous determination of sulfonamides and their N^a-acetylated metabolites in meat by semi-micro high-performance liquid chromatography). *Kuromatogurafi*, 15 (1994) 254-255; *C.A.*, 122 (1995) 158856f.
- 3720 Hormazábal, V. and Yndestad, M.: High-performance liquid chromatographic determination of praziquantel in plasma and tissues of cultured fish for residue and pharmacokinetic studies. *J. Liq. Chromatogr.*, 18 (1995) 589-597.
- 3721 Huelssewede, J.W.: Comparison of high-performance liquid chromatography and bioassay for the determination of 5-fluorocytosine in serum. *Zentralbl. Bakteriol.*, 28 (1994) 513-518; *C.A.*, 122 (1995) 177548y.
- 3722 Humberstone, A.J., Currie, G.J., Porter, C.J.H., Scanlon, M.J. and Charman, W.N.: A simplified liquid chromatography assay for the quantitation of halofantrine and desbutylhalofantrine in plasma and identification of a degradation product of desbutylhalofantrine formed under alkaline conditions. *J. Pharm. Biomed. Anal.*, 13 (1995) 265-272.

- 3723 Idowu, O.R., Peggins, J.O., Brewer, T.G. and Kelley, C.: Metabolism of a candidate 8-aminoquinoline antimalarial agent, WR 238605, by rat liver microsomes. *Drug Metab. Disp.*, 23 (1995) 1-17.
- 3724 Kaka, J.S.: Rapid simplified analysis of pyrazinamide in rat plasma by HPLC. *J. Liq. Chromatogr.*, 17 (1994) 3793-3801.
- 3725 Kotecka, B.M. and Rieckmann, K.H.: *In vivo-in vitro* model to assess chloroquine activity in monkeys. *Chemotherapy (Basel)*, 41 (1995) 134-140; C.A., 122 (1995) 204419r.
- 3726 Munns, R.K., Turnipseed, S.B., Pfenning, A.P., Roybal, J.E., Holland, D.C. and Long, A.R.: Determination of residues of flumequine and nalidixic, oxolinic, and piromidic acids in catfish by liquid chromatography with fluorescence and UV detection. *J. Assoc. Off. Anal. Chem.*, 78 (1995) 343-352.
- 3727 Nicolas, F., de Sousa, G., Thomas, P., Placidi, M., Lorenzon, G. and Rahmani, R.: Comparative metabolism of 3'-azido-3'-deoxythymidine in cultured hepatocytes from rats, dogs, monkeys, and humans. *Drug Metab. Disp.*, 23 (1995) 308-313.
- 3728 Nishimura, K., Hirama, Y., Kawano, J., Hiratsuka, A. and Nakano, M.: (Simultaneous determination of synthetic antibacterials in meat and fish by HPLC. II. Studies of cleanup methods with liquid-liquid extraction). *Hokkaidoritsu Eisei Kenkyushoho*, 44 (1994) 66-68; C.A., 122 (1995) 79331c.
- 3729 Poirier, J.-M., Lebot, M., Descamps, P., Levy, M. and Cheymol, G.: Determination of itraconazole and its active metabolite in plasma by column liquid chromatography. *Ther. Drug Monit.*, 16 (1994) 596-601; C.A., 122 (1995) 71208x.
- 3730 Rifai, N., Sakamoto, M., Law, T., Platt, O., Mikati, M., Armsby, C.C. and Brugnara, C.: HPLC measurement, blood distribution, and pharmacokinetics of oral clotrimazole, potentially useful anticisplating agent. *Clin. Chem. (Washington)*, 41 (1995) 387-391.
- 3731 Roth, J.S. and Kelley, J.A.: Determination of the anti-HIV drug 2'- β -fluoro-2',3'-dideoxyadenosine in biological fluids by reversed-phase HPLC. *J. Liq. Chromatogr.*, 18 (1995) 441-462.
- 3732 Schenck, F.J.: Isolation and quantification of ivermectin in bovine milk by matrix solid phase dispersion (MSPD) extraction and liquid chromatographic determination. *J. Liq. Chromatogr.*, 18 (1995) 349-362.
- 3733 Smedley, M.D.: Liquid chromatographic determination of multiple sulfonamide residues in bovine milk. Collaborative study. *J. Assoc. Off. Anal. Chem.*, 77 (1994) 1112-1122.
- 3734 Tachibana, M., Aoyama, M., Taniguchi, R., Anabuki, K. and Kumagai, K.: (Simultaneous determination of residual synthetic antibacterials in chicken by HPLC). *Tokyo-to Suginami-ku Eisei Shikensho Nenpo*, 12 (1994) 86-89; C.A., 122 (1995) 103961f.
- 3735 Takahashi, Y., Sekiya, T., Nishikawa, M. and Endoh, Y.S.: Simultaneous high-performance liquid chromatographic determination of amprolium, ethopabate, sulfaquinoxaline and N4-acetylsulfaquinoxaline in chicken tissues. *J. Liq. Chromatogr.*, 17 (1994) 4489-4512.
- 3736 Tendolkar, N.M., Desai, B.S. and Shinde, V.M.: Simultaneous determination of tinidazole and clotrimazole from tablets by RP-HPLC. *Indian Drugs*, 31 (1994) 551-553; C.A., 122 (1995) 89592c.
- 3737 Tsai, C.-E. and Kondo, F.: A sensitive high-performance liquid chromatographic method for detecting sulfonamide residues in swine serum and tissues after fluorescamine derivatization. *J. Liq. Chromatogr.*, 18 (1995) 965-976.
- 3738 Vergin, H., Kikuta, C., Macher, H. and Metz, R.: Pharmacokinetics and bioavailability of different formulations of aciclovir. *Arzneim.-Forsch.*, 45 (1995) 508-515.
- 3739 Wei, Y., Nygard, G.A. and Khalil, S.K.W.: A HPLC method for the separation and quantification of the enantiomers of hydroxychloroquine and its three major metabolites. *J. Liq. Chromatogr.*, 17 (1994) 3479-3490.

For additional information see C.A.:

122 (1995) 115116x, 115143d, 197142q, 197143r, 197151s.

See also 3540, 3591, 3794.

32f. Cytostatics

- 3740 Albertioni, F., Pettersson, B., Beck, O., Rask, C., Seideman, P. and Peterson, C.: Simultaneous quantitation of methotrexate and its two main metabolites in biological fluids by a novel solid-phase extraction procedure using high-performance liquid chromatography. *J. Chromatogr. B*, 665 (1995) 163-170.
- 3741 Burk, M., Volmer, M., Fartash, K. and Schneider, W.: Ion-pair liquid chromatography of cytarabine and uracil-araboside in human plasma. *Arzneim.-Forsch.*, 45 (1995) 616-619.
- 3742 Freeman, K.B., Anliker, S., Hamilton, M., Osborne, D., Dahir, P.H., Nelson, R. and Allerheiligen, S.R.B.: Validated assays for the determination of gemcitabine in human plasma and urine using high-performance liquid chromatography with ultraviolet detection. *J. Chromatogr. B*, 665 (1995) 171-181.
- 3743 Himberg, J.-P., Koskinen, M., Riekkola, M.-L. and Elo, H.: High-performance liquid chromatographic separation and determination of salicylaldoxime and β -resorcyaldoxime in aqueous solutions and in human urine. *J. Liq. Chromatogr.*, 17 (1994) 3519-3529.
- 3744 Lu, G. and Jun, H.W.: Determination of trace methotrexate and 7-OH-methotrexate in plasma by high-performance liquid chromatography with fluorimetric detection. *J. Liq. Chromatogr.*, 18 (1995) 155-171.
- 3745 Mank, A.J.G., Molenaar, E.J., Gooijer, C., Lingeman, H., Velthorst, N.H. and Brinkman, U.A.T.: Determination of the anticancer drug metabolite WR1065 using pre-column derivatization and diode laser induced fluorescence detection. *J. Pharm. Biomed. Anal.*, 13 (1995) 255-263.
- 3746 Poon, G.K., Walter, B., Lonning, P.E., Horton, M.N. and McCague, R.: Identification of tamoxifen metabolites in human Hep G2 cell line, human liver homogenate, and patients on long-term therapy for breast cancer. *Drug Metab. Disp.*, 23 (1995) 377-382.
- 3747 Qian, M., O'Dwyer, P.J. and Gallo, J.M.: High-performance liquid chromatographic method for the determination of AG-331, a novel anti-cancer agent, in human serum and urine using solid-phase extraction and photodiode-array detection. *J. Chromatogr. B*, 666 (1995) 307-314.
- 3748 Ramsauer, B., Braess, J., Unterhalt, M., Kaufmann, C.C., Hidemann, W. and Schleyer, E.: Highly sensitive high-performance liquid chromatographic assay for 1- β -D-arabinofuranosylcytosine-5'-stearyl phosphate (cytarabine-ocfosfate). *J. Chromatogr. B*, 665 (1995) 183-192.
- 3749 Saini, P., Abu-Izza, K. and Lu, D.R.: High-performance liquid chromatographic assay for sodium mercaptoundecahydrododecaborate in rat tissues. *J. Chromatogr. B*, 665 (1995) 155-161.

- 3750 Walle, T., Walle, U.K., Kumar, G.N. and Bhalla, K.N.: Taxol metabolism and disposition in cancer patients. *Drug Metab. Disp.*, 23 (1995) 506-512.
- 3751 Webster, L.K., Ellis, A.G., Kestell, P. and Rewcastle, G.W.: Metabolism and elimination of 5,6-dimethylxanthenone-4-acetic acid in the isolated perfused rat liver. *Drug Metab. Disp.*, 23 (1995) 363-368.
- 3752 Yatsevich, O.V., Arzastmasev, A.P., Zimkova, N.I., Abrazakov, M.M. and Rodionova, G.M.: Quantitative determination of sarcolysin and its hydrolysis products by high-performance liquid chromatography). *Khim.-Farm. Zh.*, 28 (1994) 56-59; C.A., 122 (1995) 89589g.
- For additional information see C.A.:
122 (1995) 71221w, 204406j, 230000s, 230001t.
- See also 3510, 3542.
- 32g. *Other drug categories*
- 3753 Bo, L.D., Verga, F., Marzo, A., Ambrosoli, L. and Poli, A.: Determination of ticlopidine in human plasma by high-performance liquid chromatography and ultraviolet absorbance detection. *J. Chromatogr. B*, 665 (1995) 404-409.
- 3754 Brunn, H. and Georgii, S.: (Qualitative and quantitative detection of dexamethasone and related xenobiotic corticosteroids in bovine urine by ELISA and HPLC/ELISA). *Arch. Lebensmittelhyg.*, 45 (1994) 66-71; C.A., 122 (1995) 178543e.
- 3755 Caraballo, I., Fernandez-Arevalo, M., Holgado, M.A., Alvarez-Fuentes, J. and Rabasco, A.M.: Simultaneous HPLC determination of some drugs commonly used in cold medications: dextromethorphan, diphenhydramine, phenylephrine, phenylpropanolamine and pseudoephedrine. *Drug Dev. Ind. Pharm.*, 21 (1995) 605-613; C.A., 122 (1995) 142731a.
- 3756 Chiarotto, J.A. and Wainer, I.W.: Determination of metyrapone and the enantiomers of its chiral metabolite metyrapol in human plasma and urine using coupled achiral-chiral liquid chromatography. *J. Chromatogr. B*, 665 (1995) 147-154.
- 3757 Cohen, H., Armstrong, F. and Campbell, H.: Sensitive fluorescence detection of robenidine by derivatization with dansyl chloride and high-performance liquid chromatography. *J. Chromatogr. A*, 694 (1995) 407-413.
- 3758 Constanzer, M., Chavez, C. and Matuszewski, B.: Development and comparison of high-performance liquid chromatographic methods with tandem mass spectrometric and ultraviolet absorbance detection for the determination of cyclobenzaprine in human plasma and urine. *J. Chromatogr. B*, 666 (1995) 117-126.
- 3759 Crimella, T., Orlandi, R., Bocchiola, G., Anders, U. and Stradi, R.: Analytical and chemical profile of pidotimod. *Arzneim.-Forsch.*, 44 (1994) 1405-1410; C.A., 122 (1995) 197136r.
- 3760 Davis, M.R. and Quigley, M.N.: Liquid chromatographic determination of UV absorbers in sunscreens. *J. Chem. Educ.*, 72 (1995) 279-281; C.A., 122 (1995) 213167j.
- 3761 Dias, V.C., Lucien, J., LeGat, D.F. and Yatscoff, R.W.: Measurement of the active leflunomide metabolite (A77 1726) by reverse-phase high-performance liquid chromatography. *Ther. Drug Monit.*, 17 (1995) 84-88; C.A., 122 (1995) 177553w.
- 3762 El-Ealily, A.F.M., El-Anwar, F.M. and El-Swify, M.: Ion-pair HPLC analysis of multi-ingredient sedative-cough syrup. *Bull. Fac. Pharm. (Cairo Univ.)*, 31 (1993) 297-301; C.A., 122 (1995) 197123j.
- 3763 Farag, S.A.: Liquid chromatographic assay of diatrizoic acid and its diiodo degradation products in radio-opaque solutions. *J. Assoc. Off. Anal. Chem.*, 78 (1995) 328-333.
- 3764 Glinski, J.A. and Caviness, G.O.: Centrifugal partition chromatography in assay guided isolation of natural products. A case study of immunosuppressive components of *Tripterygium wilfordii*. *Chromatogr. Sci. Ser.*, 68(Centrifugal Partition Chromatography) (1995) 133-165; C.A., 122 (1995) 170309n - a review with 42 refs.
- 3765 Gordon, B., Marchant, N., Richards, R. and Walther, B.: Methods for the analysis of benfluorex metabolites. *Methodol. Surv. Bioanal. Drugs*, 23(Biofluid and Tissue Analysis for Drugs, Including Hypolipidaemics) (1994) 103-112; C.A., 122 (1995) 229986m.
- 3766 Guillaume, Y., Peyrin, E. and Guinchard, C.: Rapid determination of sulbactam and tazobactam in human serum by high-performance liquid chromatography. *J. Chromatogr. B*, 665 (1995) 363-371.
- 3767 Gumieniczek, A. and Przyborowski, L.: Determination of benzobromarone and benzarone in pharmaceuticals by high-performance liquid chromatography. *J. Liq. Chromatogr.*, 17 (1994) 3411-3419.
- 3768 Herzog, R. and Leuschner, J.: Experimental studies on the pharmacokinetics and toxicity of 5-aminosalicylic acid-O-sulfate following local and systemic application. *Arzneim.-Forsch.*, 45 (1995) 300-303.
- 3769 Ho, Y., Lee, Y.-L. and Hsu, K.-Y.: Determination of (+)-catechin in plasma by high-performance liquid chromatography using fluorescence detection. *J. Chromatogr. B*, 665 (1995) 383-389.
- 3770 Noroski, J.E. and Mayo, D.J.: Chiral LC of a cholesterol-lowering drug using serum albumin mobile phases. *J. Pharm. Biomed. Anal.*, 13 (1995) 83-87.
- 3771 Nowotnik, D.P., Nanjappan, P., Zeng, W. and Ramalingam, K.: Separation of the stereoisomers of hexamethyl-propyleneamine oxime (HM-PAO) by high-performance liquid chromatography. *J. Liq. Chromatogr.*, 18 (1995) 673-687.
- 3772 Pesonen, T., Holmalahti, J. and Pohjola, J.: Determination of chlorhexidine in saliva using high-performance liquid chromatography. *J. Chromatogr. B*, 665 (1995) 222-225.
- 3773 Rajewski, R.A., Kosednar, D.G., Matches, T.A., Wong, O.S., Burchett, K. and Thakker, K.: Stereo-specific analysis of a novel protein kinase C inhibitor. *J. Pharm. Biomed. Anal.*, 13 (1995) 247-253.
- 3774 Rastogi, S.C., Schouten, A., de Kruijff, N. and Weijland, J.W.: Contents of methyl-, ethyl-, propyl-, butyl- and benzylparaben in cosmetic products. *Contact Dermatitis*, 32 (1995) 28-30; C.A., 122 (1995) 142007u.
- 3775 Sallustio, B.C. and Fairchild, B.A.: Biosynthesis, characterisation and direct high-performance liquid chromatographic analysis of gemfibrozil 1-O- β -acylglucuronide. *J. Chromatogr. B*, 665 (1995) 345-353.

- 3776 Vielhauer, S., Rudolphi, A., Boos, K.-S. and Seidel, D.: Evaluation and routine application of the novel restricted-access precolumn packing material Alkyl-Diol Silica: coupled-column high-performance liquid chromatographic analysis of the photoreactive drug 8-methoxypsoralen in plasma. *J. Chromatogr. B*, 666 (1995) 315-322.
- 3777 Wallner, P.: HPLC analysis of UVA and -B filters in cosmetics. *Dtsch. Lebensm.-Rundsch.*, 89 (1993) 375-379; *C.A.*, 122 (1995) 142002p.
- 3778 Wang, C.P., Chan, K.W., Schiksnis, R.A., Scatina, J. and Sisenwine, S.F.: High performance liquid chromatographic isolation, spectroscopic characterization, and immunosuppressive activities of two rapamycin degradation products. *J. Liq. Chromatogr.*, 17 (1994) 3383-3392.
- For additional information see *C.A.*:
122 (1995) 45518d, 89602f, 229990h, 229991j, 230002u.
- See also 2725, 2822, 2868, 3406, 3464, 3537, 3560, 3571, 3572, 3650, 3655, 3699, 3729, 3746, 3819, 3835.
- 32h. *Toxicological and forensic applications*
- 3779 Ardrey, R.E., Kimber, K.A. and Kilpatrick, G.: LC-MS - a routine analytical tool in forensic toxicology. In: Oliver, J.S. (Editor), *Forensic Toxicol., Proc. Int. Meet. Int. Assoc. Forensic Toxicol.*, 26th 1992, Scot. Acad. Press, Edinburgh, 1992, pp. 103-119; *C.A.*, 122 (1995) 232815k.
- 3780 Bogusz, M. and Erkens, M.: Influence of biological matrix on chromatographic behavior and detection of selected acidic, neutral, and basic drugs examined by means of a standardized HPLC-DAD system. *J. Anal. Toxicol.*, 19 (1995) 49-55.
- 3781 Carretero, L., Maldonado, M. and Laserna, J.J.: (Laser-induced fluorescence for detection of drugs banned in sports by micellar liquid chromatography). *Quim. Anal. (Barcelona)*, 11 (1992) 107-119; *C.A.*, 122 (1995) 48615u.
- 3782 Casale, J.F. and Moore, J.M.: Detection and determination of pseudococaine in coca leaves and illicit cocaine samples. *J. Forensic Sci.*, 39 (1994) 1537-1543; *C.A.*, 122 (1995) 48624w.
- 3783 Chen, H., Ron, N., Ma, Q., Han, N. and Fan, D.: (Reverse phase liquid chromatography for analyzing Saikuzuo residue in field atmosphere, brown rice, hull and straw). *Huanjing Wuran Yu Fangzhi*, 16 (1994) 35-36; *C.A.*, 122 (1995) 74269d.
- 3784 Chen, H.-M. and Peterson, C.M.: Quantifying ethanol by high-performance liquid chromatography with precolumn enzymic conversion and derivatization with fluorometric detection. *Alcohol (N.Y.)*, 11 (1994) 577-582; *C.A.*, 122 (1995) 48567e.
- 3785 Debruyne, D., Albessard, F., Bigot, M.C. and Moulin, M.: Comparison of three advanced chromatographic techniques for cannabis identification. *Bull. Narc.*, 46 (1994) 109-121; *C.A.*, 122 (1995) 74099y.
- 3786 Furness, J.F., Weekes, A., Clench, M., Wolff, K., Barnes, I., Hay, A.W.M. and Cooke, M.: The determination of methadone and metabolites in human urine by HPLC with ultraviolet, and particle beam mass spectrometric detection. *J. Liq. Chromatogr.*, 17 (1994) 4431-4444.
- 3787 Hallbach, J. and Guder, W.G.: Experiences with HPLC and/or GC-MS as a first step for drugs of abuse testing and clinical toxicology. *Eur. J. Clin. Chem. Clin. Biochem.*, 33 (1995) A29.
- 3788 Lambert, W.E., Meyer, E., Xue-Ping, Y. and de Leenheer, A.P.: Screening, identification, and quantitation of benzodiazepines in postmortem samples by HPLC with photodiode array detection. *J. Anal. Toxicol.*, 19 (1995) 35-40.
- 3789 Lehman, T. and Brenneisen, R.: High performance liquid chromatographic profiling of cannabis products. *J. Liq. Chromatogr.*, 18 (1995) 689-700.
- 3790 Logan, B.K. and Stafford, D.T.: In-line solid phase sample preparation for analysis of drugs in postmortem fluids by high performance liquid chromatography. In: Oliver, J.S. (Editor), *Forensic Toxicol., Proc. Int. Meet. Int. Assoc. Forensic Toxicol.*, 26th 1992, Scot. Acad. Press, Edinburgh, 1992, pp. 288-294; *C.A.*, 122 (1995) 207218k.
- 3791 Marr, J.C., McDowell, L.M. and Quilliam, M.A.: Investigation of derivatization reagents for the analysis of diarrhetic shellfish poisoning toxins by liquid chromatography with fluorescence detection. *Nat. Toxins*, 2 (1994) 302-311; *C.A.*, 122 (1995) 25421p.
- 3792 Nishikawa, M., Tatsuno, M., Tsuchihashi, H., Igarashi, K., Kasuya, F. and Fukui, M.: (The analysis of quaternary ammonium compounds in human urine by thermospray liquid chromatography-mass spectrometry). *Jpn. J. Toxicol. Environ. Health*, 40 (1994) 534-541; *C.A.*, 122 (1995) 125463x.
- 3793 Polettini, A., Montagna, M., Hogendoorn, E.A., Dijkman, E., van Zoonen, P. and van Ginkel, L.A.: Applicability of coupled-column liquid chromatography to the analysis of β -agonists in urine by direct sample injection. I. Development of a single-residue reversed-phase liquid chromatography-UV method for clenbuterol and selection of chromatographic conditions suitable for multi-residue analysis. *J. Chromatogr. A*, 695 (1995) 191-31.
- 3794 Roenn, A.M., Lemnge, M.M., Angelo, H.R. and Bygbjerg, I.C.: High-performance liquid chromatography determination of dapsone, monoacetyldapsone, and pyrimethamine in filter paper blood spots. *Ther. Drug Monit.*, 17 (1995) 79-83; *C.A.*, 122 (1995) 177552v.
- 3795 Rositano, J. and Nicholson, B.C.: A method for the detection of cyanobacterial peptide toxins by HPLC. *Spec. Publ. - R. Soc. Chem.*, 149 (Detection Methods for Cyanobacterial Toxins) (1994) 155-157; *C.A.*, 122 (1995) 74064h.
- 3796 Sabater Montesinos, J., Garcia Alvarez-Coque, M.C., Ramis Ramos, G. and Laserna Vazquez, J.J.: On the use of micellar liquid chromatography for detection of drug misuse in sport. *Quim. Anal. (Barcelona)*, 11 (1992) 163-171; *C.A.*, 122 (1995) 48616v.
- 3797 Stockemer, J. and Guerne, M.: (Optimization of a HPLC method for the determination of DSP toxins). *Arch. Lebensmittelhyg.*, 44 (1993) 146-147; *C.A.*, 122 (1995) 131263y.
- 3798 Walker, J.A., Kkrueger, S.T., Lurie, I.S., Marche, H.L. and Newby, N.: Analysis of heroin drug seizures by micellar electrokinetic capillary chromatography (MECC). *J. Forensic Sci.*, 40 (1995) 6-9; *C.A.*, 122 (1995) 207200y.
- For additional information see *C.A.*:
122 (1995) 95762z.
- See also 2777, 2789, 2801, 2937, 3042, 3443, 3445, 3450, 3521, 3704.

32i. *Plant extracts*

- 3799 Basha, L.I.A., Rashed, M.S. and Aboul-Enein, H.Y.: TLC assay of thymoquinone in black seed oil (*Nigella sativa* L.) and identification of dithymoquinone and thymol. *J. Liq. Chromatogr.*, 18 (1995) 105-115.
- 3800 Collins, M.D., Wasmund, L.M. and Bosland, P.W.: Improved method for quantifying capsaicinoids in *Capsicum* using HPLC. *HortScience*, 30 (1995) 137-139; *C.A.*, 122 (1995) 182312w.
- 3801 Graenicher, F., Christen, P. and Vuagnat, P.: Rapid high performance liquid chromatographic quantification of valepatriates in hair root cultures of *Valeriana officinalis* L. var. *sambucifolia* Mikan. *Phytochem. Anal.*, 5 (1994) 297-301; *C.A.*, 122 (1995) 89574y.
- 3802 Singh, B.K., Teclé, B. and Shaner, D.L.: Determination of 2-keto acids and amino acids in plant extracts. *J. Liq. Chromatogr.*, 17 (1994) 4469-4477.
- 3803 Tarantilis, P.A., Tsoupras, G. and Polissiou, M.: Determination of saffron (*Crocus sativus* L.) components in crude plant extract using high-performance liquid chromatography-UV-visible photodiode-array detection-mass spectrometry. *J. Chromatogr. A*, 699 (1995) 107-118.
- 3804 Tzakou, O., Verekokidou, E. and Harvala, C.: Comparative HPLC analysis of polyphenolic compounds in four species of *Galium* L. *J. Liq. Chromatogr.*, 17 (1994) 4463-4467.
- 3805 Yang, F., Ou, Q. and Yu, W.: Semi-preparative separation of taraxeryl-acetate and coumarins from *Artemisia dalailamae* Kraschen by high-speed countercurrent chromatography. *J. Liq. Chromatogr.*, 18 (1995) 395-403.

For additional information see C.A.:

122 (1995) 89595f, 170330n, 170332q, 196659b, 197105e.

See also 2805, 2827, 2882, 2979, 2982, 3764, 3785, 3789.

33. CLINICO-CHEMICAL APPLICATIONS

33a. *General papers and reviews*

- 3806 Riley, C.M., Lough, W.J. and Wainer, I.W. (Editors): *Pharmaceutical and Biomedical Applications of Liquid Chromatography Prog. Pharm. Biomed. Anal.*, Vol. 1, Pergamon, Oxford, 1994, 379 pp.; *C.A.*, 122 (1995) 234817m.

33b. *Complex mixtures and profiling (single compounds by cross-reference only)*

- 3807 Idei, M., Pajor, A. and Menyhart, J.: Bioanalysis of human amniotic fluid: chromatographical and biological characterization of a component inhibiting contraction of the myometrium. In: Kalasz, H., Ettre, L.S. and Pick, J. (Editors), *New Approaches Chromatogr. '93, [Pap. Budapest Chromatogr. Conf.]*, 1990, Fekete Sas Kiado, Budapest, 1993, pp. 103-123; *C.A.*, 122 (1995) 210391s.

See also 2787, 2836, 2839, 2843, 2857, 2902, 2918, 2933, 2954, 2955, 2961, 2963, 2970, 2996, 2997, 3008, 3021, 3027, 3030, 3038, 3043, 3051, 3078, 3186, 3189, 3197, 3198, 3241, 3242, 3319, 3351, 3426, 3456, 3464, 3468, 3470, 3515, 3746, 3750, 3865, 3899.

34. FOOD ANALYSIS

- 3808 Yueming, Q. and Wuji, Z.: (Chemical reaction and chromatographic method for the determination of dimethyl residues in vegetables). *Fenzi Huaxue*, 22 (1994) 899-901; *C.A.*, 122 (1995) 79348p.

See also 3022.

34a. *General papers and reviews*

- 3809 Doneche, B. (Editor): *Modern Advances in the Chromatography of Wine: Applications to the Sensory Analysis of Wines, Porto, Port., Mar. 31-33 April, 1992. Tec & Doc - Lavoisier, Paris, 1993, 253 pp.*; *C.A.*, 122 (1995) 104523b.

34b. *Complex mixtures (single compounds by cross-reference only)*

- 3810 De Maria, C.A.B., Trugo, L.C., Moreira, R.F.A. and Petracco, M.: Simultaneous determination of total chlorogenic acid, trigonelline and caffeine in green coffee samples by high performance gel filtration chromatography. *Food Chem.*, 52 (1995) 447-449; *C.A.*, 122 (1995) 185709s.
- 3811 Golc-Wondra, A., Skocir, E. and Prosek, M.: Determination of monosodium glutamate in food products. *J. Planar Chromatogr.*, 8 (1995) 117-121.
- 3812 Juelich, E. and Lind, G.: (Determination of sugar, GdL, and sodium ascorbate in seasonings for sausages). *Fleischwirtschaft*, 74 (1994) 637-638; *C.A.*, 122 (1995) 185635q.
- 3813 Kermasha, S., Goetghebeur, M. and Dumont, J.: Separation and characterization of pepper-contaminated cinnamon using HPLC analyses. *Food Sci. Technol. (London)*, 27 (1994) 578-582; *C.A.*, 122 (1995) 54386m.
- 3814 Krueger, E., Radau, B. and Roenn, D.: (Investigation of beers and beer heads by methods employed in protein chemistry). *Monatsschr. Brauwiss.*, 46 (1993) 312-319; *C.A.*, 122 (1995) 131260v.
- 3815 Lancaster, F.E. and Lawrence, J.F.: Determination of annatto in high-fat dairy products, margarine and hard candy by solvent extraction followed by high-performance liquid chromatography. *Food Addit. Contam.*, 12 (1995) 9-19; *C.A.*, 122 (1995) 238043d.
- 3816 Madigan, D., McMurrough, I. and Smyth, M.R.: Rapid determination of 4-vinyl guaiacol and ferulic acid in beers and worts by high-performance liquid chromatography. *J. Am. Soc. Brew. Chem.*, 52 (1994) 152-155; *C.A.*, 122 (1995) 54374f.
- 3817 Rimedio, N.T. and Chen, L.F.: Chromatographic separation of organic nonsugars, colloidal material and inorganic-organic complexes from juice, liquors, syrups and/or molasses. *U.S. US 5,382,294 (Cl. 127-42; C13J1/06)*, 17 Jan. 1995, US Appl. 749,667, 26 Aug. 1991; 25 p.; *C.A.*, 122 (1995) 190875x.

- 3818 Van Loo, J., Booten, K. and Smits, G.: Chromatographic separation of polydispersed saccharide compositions to give products useful in foods. *PCT Int. Appl.* WO 94 12,541 (Cl. C08B37/18), 09 Jun. 1994, BE Appl. 92/1,021, 24 Nov. 1992; 47 p.; C.A., 122 (1995) 109228u.
- 3819 Zhu, Z.: (Rapid determination of dexamethasone acetate residue in pork by high-performance liquid chromatography). *Fenxi Huaxue*, 22 (1994) 1163-1166; C.A., 122 (1995) 185693g.
- For additional information see C.A.:
122 (1995) 104175q, 158880j, 163938p, 212360e.
- See also 2793, 2795, 2799, 2812, 2813, 2814, 2815, 2816, 2818, 2824, 2826, 2829, 2831, 2849, 2850, 2854, 2890, 2892, 2897, 2898, 2900, 2908, 2911, 2915, 2926, 2936, 2952, 2974, 2990, 2999, 3011, 3013, 3023, 3024, 3035, 3044, 3083, 3245, 3358, 3428, 3459, 3480, 3501, 3504, 3506, 3517, 3526, 3530, 3533, 3534, 3535, 3546, 3547, 3558, 3559, 3566, 3574, 3580, 3586, 3589, 3590, 3591, 3593, 3719, 3728, 3732, 3733, 3734, 3735, 3737, 3797, 3844, 3850, 3861, 3889, 3891, 3901.
- 34c. *Organoleptically important compounds (flavors, odors, volatiles)*
- See 3459.
35. ENVIRONMENTAL ANALYSIS
- 35a. *General papers and reviews*
- 3820 Conrad, E.E., Kelly, K.P. and Schwartz, N.L.: GPC and oily samples. *Environ. Test. Anal.*, 3 (1994) 34-39; C.A., 122 (1995) 204159f - a review with 3 refs.
- 3821 Fort, D.J., Delphon, J., Powers, C.R., Helems, R., Gonzalez, R. and Stover, E.L.: Development of automated methods of identifying toxicants in the environment. *Bull. Environ. Contam. Toxicol.*, 54 (1995) 104-111; C.A., 122 (1995) 232799h.
- 3822 Hayakawa, K.: (Development of new photometric detection in liquid chromatography for the determination of environmental compounds). *Yakugaku Zasshi*, 114 (1994) 669-680; C.A., 122 (1995) 182254d - a review with 39 refs.
- See also 2624, 2709, 2767, 2773, 2778, 2781, 3867.
- 35b. *Air pollution (complex mixtures; single compounds by cross-reference only)*
- See 2697, 3473, 3900.
- 35c. *Water pollution (complex mixtures; single compounds by cross-reference only)*
- 3823 Elvira-Cozar, C., Cano-Faura, P., Perez-Arribas, L.V., Leon-Gonzalez, M.E. and Polo-Diez, L.M.: Trace priority pollutant phenols enrichment from water by ion chromatography. *Chromatographia*, 40 (1995) 91-95.
- 3824 Minnaard, W.A., Slobodnik, J., Vreuls, J.J., Hupe, K.-P. and Brinkman, U.A.T.: Rapid liquid chromatographic screening of organic micropollutants in aqueous samples using a single short column for trace enrichment and separation. *J. Chromatogr. A*, 696 (1995) 333-340.
- 3825 Pocurull, E., Marce, R.M. and Borrull, F.: Liquid chromatography of phenolic compounds in natural water using on-line trace enrichment. *Chromatographia*, 40 (1995) 85-90.
- 3826 Zapf, A., Heyer, R. and Stan, H.-J.: Rapid micro liquid-liquid extraction method for trace analysis of organic contaminants in drinking water. *J. Chromatogr. A*, 694 (1995) 453-461.
- See also 2764, 2794, 2797, 2919, 2920, 3047, 3576, 3578, 3579, 3582, 3585, 3587, 3878, 3882, 3885, 3898.
- 35d. *Soil pollution (complex mixtures; single compounds by cross-reference only)*
- 3827 Osipov, G.A., Nazina, T.N. and Ivanova, A.E.: (Analysis of microbial community from water-flooded oil field by chromatography-mass spectrometry). *Mikrobiologiya*, 63 (1994) 876-882; C.A., 122 (1995) 75689c.
- See also 2764, 2786, 3857.
36. SOME TECHNICAL PRODUCTS AND COMPLEX MIXTURES
- 36a. *Surfactants*
- 3828 Brenner, C., Jan, G., Chevalier, Y. and Wroblewski, H.: Evaluation of the efficacy of zwitterionic dodecyl carboxybetaine surfactants for the extraction and the separation of mycoplasma membrane protein antigens. *Anal. Biochem.*, 224 (1995) 515-523.
- 3829 Forgács, E. and Cserhádi, T.: Retention behaviour of nonyl-phenyl ethyleneoxide oligomers on alumina columns. *Fresenius J. Anal. Chem.*, 351 (1995) 688-689.
- 3830 Marcomini, A. and Zanette, M.: Derivatization procedures and HPLC separations for environmental analysis of aliphatic alcohols polyethylenglycol ethers (AE). *Riv. Ital. Sostanze Grasse*, 71 (1994) 203-208; C.A., 122 (1995) 136796j.
- 3831 Matsuzaki, M., Ishii, K., Yoshimura, H. and Hashimoto, S.: HPLC determination of cationic and amphoteric surfactants in cosmetics. *J. SCCJ*, 27 (1993) 494-497; C.A., 120 (1994) 307031w.
- 3832 McPherson, B.P. and Rasmussen, H.T.: Chromatography of cationic surfactants: HPLC, TLC and GLC. *Surfactant Sci. Ser.*, 53 (1994) 289-326; C.A., 122 (1995) 136787g - a review with 89 refs.
- See also 2934, 3843.
- 36b. *Antioxidants and preservatives*
- 3833 Parhizkari, G., Miller, R.B. and Chen, C.: A stability-indicating HPLC method for the determination of benzalkonium chloride in phenylephrine HCl 10% ophthalmic solution. *J. Liq. Chromatogr.*, 18 (1995) 553-563.

- 3834 Rose, R.C. and Bode, A.M.: Analysis of water-soluble antioxidants by high-pressure liquid chromatography. *Biochem. J.*, 306 (1995) 101-105.
- 3835 Weyland, J.W., Stern, A. and Rooselaar, J.: Determination of bonopol, bronidox, and methyl dibromo glutaronitrile in cosmetics by liquid chromatography with electrochemical detection. *J. Assoc. Off. Anal. Chem.*, 77 (1994) 1132-1136.
- 36c. *Complex mixtures, technical products and unidentified compounds*
- 3836 Heikkilä, H., Hyeok, G. and Kuisma, J.: Fractionating sulphite cooking liquors by simulated moving bed chromatography. *PCT Int. Appl. WO 94 26,380 (Cl. B01D15/08) 24 Nov. 1994, FI Appl. 93/2,108, 10 May 1993; 38 p.; C.A.*, 122 (1995) 109957f.
- 3837 Lawrence, J.F., Ménard, C. and Cleroux, C.: Evaluation of pre-chromatographic oxidation for liquid chromatographic determination of paralytic shellfish poisons in shellfish. *J. Assoc. Off. Anal. Chem.*, 78 (1995) 514-520.
- 3838 Lehotay, J. and Oktavec, D.: Application of HPLC with diode array detection in tribology. *J. Liq. Chromatogr.*, 17 (1994) 3783-3792.
- 3839 Oshima, Y.: Postcolumn derivatization liquid chromatographic method for paralytic shellfish toxins. *J. Assoc. Off. Anal. Chem.*, 78 (1995) 528-532.
- 3840 Quilliam, M.A.: Analysis of diarrhetic shellfish poisoning toxins in shellfish tissue by liquid chromatography with fluorometric and mass spectrometric detection. *J. Assoc. Off. Anal. Chem.*, 78 (1995) 555-570.
- 3841 Sassi, P., Machtalere, G., Hui, F., Kolodziejczyk, H. and Rosset, R.: Liquid chromatographic determination of base oil composition and content in lubricating oils containing dispersants of the polybutenylsuccinimide type. *Anal. Chim. Acta*, 306 (1995) 73-79.
- 3842 Swerin, A. and Waagberg, L.: Size-exclusion chromatography for characterization of cationic polyelectrolytes used in papermaking. *Nord. Pulp. Pap. Res. J.*, 9 (1994) 18-25; *C.A.*, 122 (1995) 84033m.
- 3843 Taylor, C.D., Anderson, H.A. and Brydon, D.L.: Characterization of textile lubricants by HPLC. Part III: investigation of emulsions. *J. Text. Inst.*, 85 (1994) 49-56; *C.A.*, 122 (1995) 204239g.
- 3844 Yasumoto, T., Fukui, M., Sasaki, K. and Sugiyama, K.: Determination of marine toxins in foods. *J. Assoc. Off. Anal. Chem.*, 78 (1995) 574-582.
- For additional information see C.A.:
122 (1995) 75712e, 85174v, 165048x.
- See also 2588, 2615.
37. CELLS, CELLULAR PARTICLES AND SUPRAMOLECULAR STRUCTURES
- 3845 Chun, J.-H. and Yang, S.-M.: Electrokinetic effect on separation of colloidal particles in capillary hydrodynamic chromatography (CHDC). In: *Off. Proc. Comb. Conf., 6th Conf. Asia Pac. Confed. Chem. Eng., 21st Australas Chem. Eng. Conf., Inst. Eng., Aust., Barton, 1993, pp. 283/2-287/2; C.A.*, 122 (1995) 84499t.
- 3846 Mandrusov, E., Houg, A., Klein, E. and Leonard, E.F.: Membrane-based cell affinity chromatography to retrieve viable cells. *Biotechnol. Prog.*, 11 (1995) 208-213; *C.A.*, 122 (1995) 208889s.
- 3847 Nandan, D., Daubenberger, C., Mpimbaza, G. and Pearson, T.W.: A rapid single-step purification method for immunogenic members of the hsp 70 family: validation and application. *J. Immunol. Methods*, 176 (1994) 255-263; *C.A.*, 122 (1995) 26992n.
- 3848 Qin, A. and Cui, Z.: (Purification of Marek's disease virus pp38 expressed in insect cells infected with recombinant baculovirus through affinity column chromatography). *Shengwu Gongcheng Xuebao*, 10 (1994) 239-243; *C.A.*, 122 (1995) 75705e.
- 3849 Quilliam, M.A.: Seafood toxins. *J. Assoc. Off. Anal. Chem.*, 78 (1995) 144-148 - a review with 56 refs.
- 3850 Quilliam, M.A., Xie, M. and Hardstaff, W.R.: Rapid extraction and cleanup for liquid chromatographic determination of domoic acid in unsalted seafood. *J. Assoc. Off. Anal. Chem.*, 78 (1995) 543-554.
- 3851 Shibusawa, Y., Suzuki, K., Kinoshita, H. and Matsumoto, U.: Selective separation of human peripheral platelets, granulocytes and lymphocytes by surface affinity chromatography. *J. Chromatogr. B*, 666 (1995) 233-239.
- For additional information see C.A.:
122 (1995) 50734a.
38. INORGANIC COMPOUNDS
- 38a. *Cations*
- 3852 Alberti, J., Rubio, R. and Rauret, G.: Arsenic speciation in marine biological materials by LC-UV-HG-ICP/OES. *Fresenius J. Anal. Chem.*, 351 (1995) 415-419.
- 3853 Arakawa, K.: Analysis of cations and anions by ion-exchange chromatography. *Jpn. Kokai Tokkyo Koho JP 06,273,401 [94,273,401] (Cl. G01N30/02), 30 Sep. 1994, Appl. 93/57,372, 17 Mar. 1993; 6 p.; C.A.*, 122 (1995) 122162n.
- 3854 Cobo-Fernández, M.G., Palacios, M.A., Chakraborti, D., Quevauviller, P. and Cámara, C.: On line speciation of Se(VI), Se(IV) and trimethylselenium by HPLC - microwave oven - hydride generation - atomic absorption spectrometry. *Fresenius J. Anal. Chem.*, 351 (1995) 438-442.
- 3855 Ding, H., Wang, J., Dorsey, J.G. and Caruso, J.A.: Arsenic speciation by micellar liquid chromatography with inductively coupled plasma mass spectrometric detection. *J. Chromatogr. A*, 694 (1995) 425-431.
- 3856 Fedotov, P.S., Maryutina, T.A., Pukhovskaya, V.M. and Spivakov, B.Y.: Influence of kinetic properties of extraction systems on the separation and preconcentration of some elements by countercurrent chromatography. *J. Liq. Chromatogr.*, 17 (1994) 3491-3506.
- 3857 Fodor, P. and Fischer, L.: Problems of the speciation of chromium soil samples. *Fresenius J. Anal. Chem.*, 351 (1995) 454-455.

- 3858 High, K.A., Blais, J.-S., Methven, B.A.J. and McLaren, J.W.: Probing the characteristics of metal-binding proteins using high-performance liquid chromatography - atomic absorption spectroscopy and inductively coupled plasma mass spectrometry. *Analyst (Cambridge)*, 120 (1995) 629-634.
- 3859 High, K.A., Methven, B.A., McLaren, J.W., Siu, K.W.M., Wang, J., Klaverkamp, J.F. and Blais, J.S.: Physico-chemical of metal binding proteins using HPLC-ICP-MS, HPLC-MA-AAS and electrospray-MS. *Fresenius J. Anal. Chem.*, 351 (1995) 393-402.
- 3860 Hu, W., Tao, H., Tominaga, M., Miyazaki, A. and Haraguchi, H.: A new approach for the simultaneous determination of inorganic cations and anions using ion chromatography. *Anal. Chim. Acta*, 299 (1994) 249-256.
- 3861 Jeter, H.W. and Grob, B.: Determination of radiostrontium in milk using an extraction chromatography column. *Radioact. Radiochem.*, 5 (1994) 8-17; *C.A.*, 122 (1995) 79342g.
- 3862 Khuahawar, M.Y. and Lanjwani, S.N.: High-performance liquid chromatographic separation and determination of cobalt(II), cobalt(III) and iron(II) using bis(salicylaldehyde)tetramethylethylenediamine. *J. Chromatogr. A*, 695 (1995) 132-135.
- 3863 Lintschinger, J., Kalcher, K., Gössler, W., Kölbl, G. and Novic, M.: Simultaneous determination of chromium(III) and chromium(VI) by reversed-phase ion-pair HPLC with chromium-specific detection. *Fresenius J. Anal. Chem.*, 351 (1995) 604-609.
- 3864 Martín, I., López-González, M.A., Gómez, M., Cámara, C. and Palacios, M.A.: Evaluation of high-performance liquid chromatography for the separation and determination of arsenic species by on-line high-performance liquid chromatographic-hydride generation-atomic absorption spectrometry. *J. Chromatogr. B*, 666 (1995) 101-109.
- 3865 Mathieu, C. and Mathieu, H.: Aluminium in serum determined by HPLC. *Clin. Chem. (Washington)*, 41 (1995) 625.
- 3866 Mimura, H., Kobayashi, T. and Akiba, K.: Chromatographic separation of strontium and cesium with mixed zeolite column. *J. Nucl. Sci. Technol.*, 32 (1995) 60-67; *C.A.*, 122 (1995) 145124r.
- 3867 Mohite, B.S., Patil, J.M. and Zambare, D.N.: Column chromatographic separation of lead from alloy and environmental samples using poly(dibenzo-18-crown-6). *J. Indian Chem. Soc.*, 71 (1994) 289-291; *C.A.*, 122 (1995) 121985w.
- 3868 Myasoedov, B.F., Chmutova, M.K., Smirnov, I.V. and Shadrin, A.Yu.: Isolation of actinides and separation from rare earth elements and selected fission products by bidentate neutral organophosphorus compounds. *Proc. Int. Conf. Technol. Expo. Future Nucl. Syst.: Emerging Fuel Cycles Waste Disposal Options*, 1 (1993) 581-587; *C.A.*, 120 (1994) 333498y.
- 3869 Nakamura, S., Watanabe, M. and Akiba, K.: Separation of yttrium from lanthanoid elements by centrifugal partition chromatography. *J. Liq. Chromatogr.*, 18 (1995) 377-394.
- 3870 Neely, F.L.: Determination of Fe(III) in fermentation broth by ion-interaction chromatography. *J. Liq. Chromatogr.*, 17 (1994) 4381-4390.
- 3871 Owen, L.M.W., Crews, H.M., Massey, R.C. and Bishop, N.J.: Determination of copper, zinc and aluminium from dietary sources in femur, brain and kidney and guinea pigs and study of some elements in *in vivo* intestinal digests by size-exclusion chromatography inductively coupled plasma mass spectrometry. *Analyst (Cambridge)*, 120 (1995) 705-712.
- 3872 Peterson, J.A. and Risley, D.S.: Validation of an HPLC method for the determination of sodium in LY293111 sodium, a novel LTB_4 receptor antagonist, using evaporative light scattering detection. *J. Liq. Chromatogr.*, 18 (1995) 331-338.
- 3873 Prasad, B.B., Kumar, A. and Sundd, S.: Preparation, characterization and performance of silica gel-immobilized poly(N-chloranil-N,N,N',N'-tetramethylethylenediammonium disulphosalicylate) for chromatographic separation of heavy metals. *React. Polym.*, 23 (1994) 229-238; *C.A.*, 122 (1995) 177128m.
- 3874 Renko, M., Osojnik, A. and Hudnik, V.: ICP emission spectrometric analysis of rare earth elements in permanent magnet alloys. *Fresenius J. Anal. Chem.*, 351 (1995) 610-613.
- 3875 Schilling, T., Schramel, P., Michalke, B. and Knapp, G.: Separation of noble metal ions on silica with chemically bonded ligands. *Microchim. Acta*, 116 (1994) 83-90; *C.A.*, 122 (1995) 150303c.
- 3876 Subramanian, S. and Woittiez, J.R.W.: Determination of cobalt in biological samples by radiochemical neutron activation analysis employing reversed-phase chromatography. *Biol. Trace Elem. Res.*, 43-45 (1994) 117-124; *C.A.*, 122 (1995) 50697r.
- 3877 Szczepaniak, W. and Szymanski, A.: Chromatographic separation of platinum metals on silica modified with macrocyclic compounds. *Acta Chromatogr.*, 2 (1993) 7-20; *C.A.*, 122 (1995) 150301a.
- 3878 Vasconcelos, M.T. and Gomes, C.A.R.: Limitations of ion chromatography with post-column reaction for determination of heavy metals in waters containing strong chelating agents. *J. Chromatogr. A*, 696 (1995) 227-234.
- 3879 Voloshchik, I.N.: Method of determination of ammonium and alkali metal cations. *U.S.S.R. SU 1,810,822 (Cl. G01N30/06)*, 23 Apr. 1993, Appl. 4,757,658, 10 Nov. 1989; *C.A.*, 122 (1995) 122186y.
- 3880 Wang, H., Zhang, H.S. and Cheng, J.K.: Determination of V(V), Nb(V) and Ta(V) as their 2-(5-nitro-2-pyridylazo)-5-diethylaminophenol complexes by reversed-phase high-performance liquid chromatography. *Fresenius J. Anal. Chem.*, 351 (1995) 683-685.
- 3881 Wang, J. and Marshall, W.D.: Supercritical fluid extraction with carbon dioxide for the characterisation of cadmium, copper and zinc bound to metallothioneins. *Analyst (Cambridge)*, 120 (1995) 623-628.
- 3882 Watanabe, K.: (Determination of trace level chromium(VI) and cadmium ions by ion chromatography). *Kogyo Yosui*, 433 (1994) 55-62; *C.A.*, 122 (1995) 88657r - a review with 15 refs.
- 3883 Wróbel, K., González, E.B., Wróbel, K. and Sanz-Medel, A.: Aluminium and silicon speciation in human serum by ion-exchange high-performance liquid chromatography - electrothermal atomic absorption spectrometry and gel electrophoresis. *Analyst (Cambridge)*, 120 (1995) 809-815.
- 3884 Yang, X.-J.: Extractive chromatographic separation and inductively coupled plasma atomic emission spectrometric determination of trace impurities in high purity europium oxide. *Talanta*, 41 (1994) 1807-1813; *C.A.*, 122 (1995) 150412n.
- 3885 Zernichow, L. and Lund, W.: Size exclusion chromatography of aluminium species in natural waters. *Anal. Chim. Acta*, 300 (1995) 167-171.

- 3886 Zhang, X.S., Lin, X.S., Quiao, G.J. and Liu, L.Z.: Determination of palladium and gold as 4,4'-bis(dimethylamino)thiobenzophenone complex by ion-pair reversed phase liquid chromatography. *Chromatographia*, 39 (1994) 693-696.
- 3887 Zuo, X.J., Chen, L.R. and Wang, Q.: System peaks in single column ion chromatography (SCIC) with indirect spectrophotometric detection. *Chromatographia*, 39 (1994) 713-718.
- For additional information see C.A.:
122 (1995) 95400e, 122169v.
- See also 2627, 2685, 2723, 3906.
- 38b. Anions**
- 3888 Albazi, S.J., Gharib, T. and Cortes, J.C.: Chromate as a mobile phase for the determination of anions using cetyltrimethylammonium-coated columns and indirect photometric detection. *J. Liq. Chromatogr.*, 18 (1995) 489-503.
- 3889 Anonymous: Anions in beer by ion chromatography. *J. Am. Soc. Brew. Chem.*, 52 (1994) 178-180; C.A., 122 (1995) 54378k.
- 3890 DeVargas, M.C., DeMedina, H.L., Villalobos, E., Gutierrez, E. and Mazurek, R.: Determination of total nitrogen in sediments by using high-pressure bombs and ion chromatography. *Analyst (Cambridge)*, 120 (1995) 761-763.
- 3891 Dolenc, J. and Gorenc, D.: Ion chromatographic determination of inorganic anions in vinegar samples. *Nahrung*, 38 (1994) 434-438; C.A., 122 (1995) 54310g.
- 3892 Edgell, K.W., Longbottom, J.E. and Pfaff, J.D.: Determination of inorganic anions in water by ion-chromatography: collaborative study. *J. Assoc. Off. Anal. Chem.*, 77 (1994) 1253-1263.
- 3893 Fung, Y.S. and Dao, K.L.: Ion chromatographic determination of traces of some oxoanions with direct spectrophotometric detection. *Anal. Chim. Acta*, 300 (1995) 207-214.
- 3894 Karpyuk, A.D., Vakhrusheva, M.V. and Sazhina, N.K.: (Determination of chloride ions by ion chromatography in the coolant of the primary circuit of a nuclear power plant). *Zavod. Lab.*, 60 (1994) 8-9, 62; C.A., 122 (1995) 177260y.
- 3895 Li, W. and Rossi, T.M.: Determination of sulfamate and sulfate as degradation products in an antiepileptic drug using ion chromatography and indirect UV detection. *J. Liq. Chromatogr.*, 18 (1995) 917-923.
- 3896 Liu, A., Xu, L., Li, T., Dong, S. and Wang, E.: Electrocatalytic oxidation and ion chromatographic detection of Br⁻, I⁻, SO₃²⁻, S₂O₃²⁻ and SCN⁻ at a platinum particle-based glassy carbon modified electrode. *J. Chromatogr. A*, 699 (1995) 39-47.
- 3897 Mehra, M.C. and Landry, E.: Novel UV absorbing eluents for speciation and ion chromatographic determinations of anions. *Fresenius J. Anal. Chem.*, 351 (1995) 584-586.
- 3898 Murayama, T., Oohara, H. and Aoyama, Y.: Determination of free cyanide in water. *Jpn. Kokai Tokkyo Koho* JP 06,300,742 [94,300,742] (Cl. G01N30/14), 28 Oct. 1994, JP Appl. 93/26,403, 16 Feb. 1993; 7 p.; C.A., 122 (1995) 114538z.
- 3899 Nagashima, K., Fukushima, K. and Kamaya, M.: Determination of trace amounts of sulfide in human serum by high-performance liquid chromatography with fluorometric detection after derivatization with 2-amino-5-N,N-diethylaminotoluene and iron(III). *J. Liq. Chromatogr.*, 18 (1995) 515-526.
- 3900 Poovey, H.G. and Rando, R.J.: Determination of chlorine and chlorine dioxide by non-suppressed ion chromatography and application to exposure assessment in the paper industry. *J. Liq. Chromatogr.*, 18 (1995) 261-275.
- 3901 Ruiz, E., Santillana, M.I., Nieto, M.T. and Sastre, I.: Determination of water-soluble inorganic phosphates in fresh vegetables by ion chromatography. *J. Liq. Chromatogr.*, 18 (1995) 989-1000.
- 3902 Sakai, H., Fujiwara, T. and Kumamaru, T.: Sensitivity enhancement in the determination of silicate by ion-exclusion chromatography using luminol chemiluminescence detection. *Anal. Chim. Acta*, 302 (1995) 173-177.
- For additional information see C.A.:
122 (1995) 95458e, 95459f, 122163p, 122164q, 177336c.
- See also 2566, 2685, 2899, 3836, 3853, 3860, 3879, 3887.
- 38d. Volatile inorganic compounds**
- 3903 Charbonneau, S., Gilbert, R. and Lépine, L.: Determination of carbonic acid in steam-condensate cycle samples using nonsuppressed ion chromatography. *Anal. Chem.*, 67 (1995) 1204-1209.
- 3904 Huszthy, P., Bradshaw, J.S., Bordunov, A.V. and Izatt, R.M.: Enantiomeric separation of chiral α -(1-naphthyl)ethylammonium perchlorate by silica gel-bound chiral-pyridino-18-crown-6 ligands. *ACH - Models Chem.*, 131 (1994) 445-454; C.A., 122 (1995) 159871u.
- For additional information see C.A.:
122 (1995) 122165r.
- See also 2641.
- 39. RADIOACTIVE AND OTHER ISOTOPE COMPOUNDS**
- 3905 Dadachova, E., Mirzadeh, S., Lambrecht, R.M., Hetherington, E.L. and Knapp, F.F., Jr.: Separation of carrier-free ¹⁶⁶Ho from Dy₂O₃ targets by partition chromatography and electrophoresis. *J. Radioanal. Nucl. Chem.*, 199 (1995) 155-123; C.A., 122 (1995) 144930p.
- 3906 Mathus, J.N., Murali, M.S., Iyer, R.H., Ramanujam, A., Dhama, P.S., Gopalakrishnan, V., Badheka, L.P. and Banerji, A.: Extraction chromatographic separation of minor actinides from PUREX high-level wastes using CMPO. *Nucl. Technol.*, 109 (1995) 216-225; C.A., 122 (1995) 117198z.
- For additional information see C.A.:
122 (1995) 117178t.
- See also 2828, 3861, 3866.

Gas Chromatography

1. REVIEWS AND BOOKS

- 1184 Grob, K.: *Einspritztechniken in der Kapilar-Gaschromatographie*. Hühig, Heidelberg, 1995, 465 p.
- 1185 Miller, R.J.: Industrial gas chromatography - "A personal perspective". *J. High. Resolut. Chromatogr.*, 18 (1995) 213-216.
- 1186 Samukawa, K. and Oguri, N.: (Pyrolysis gas chromatography.) Gihodo Shuppan, Tokyo, 1994, 278 p.; *C.A.*, 122 (1995) 122115z.
- 1187 Verhoeven, N.M., Wanders, R.J.A., van den Heuvel, C.M.M. and Jakobs, C.: Stable isotope dilution gas chromatography-mass spectrometry for detection of peroxisomal metabolites: pre- and postnatal diagnosis. *Nippon Iyo Masu Supekutoru Gakkai Koenshu*, 19 (1994) 3-20; *C.A.*, 122 (1995) 50293f - a review with 33 refs.

See also 1198, 1298, 1301, 1308, 1336, 1343, 1498, 1603, 1609, 1741, 1784.

2. FUNDAMENTALS, THEORY AND GENERAL

2a. General

- 1188 Bartle, K.D.: Introduction to the theory of chromatographic separations with reference to gas chromatography. In: Baugh, P.J. (Editor), *Gas Chromatography*, IRL, Oxford, 1993, pp.1-14.
- 1189 Berthod, A., Zhou, E.Y., Le, K. and Armstrong, D.W.: Determination and use of Rohrschneider-McReynolds constants for chiral stationary phases used in capillary gas chromatography. *Anal. Chem.*, 67 (1995) 849-857.
- 1190 Jöhl, M., Bowlin, D. and Davis, J.M.: Verification of the extended statistical model of overlap: application to gas chromatograms containing variable peak densities. *Anal. Chim. Acta*, 302 (1995) 133-146; *C.A.*, 122 (1995) 150533c.
- 1191 Lebrón-Aguilar, R., Quintanilla-López, J.E., Tello, A.M., Fernández-Torres, A. and García-Domínguez, J.A.: New equation for specific retention volumes in capillary column gas chromatography. *J. Chromatogr. A*, 697 (1995) 441-451.
- 1192 Olivé, J. and Grimalt, J.O.: Relationship between different chromatographic peak description functions and numerical solutions of the mass balance equation. *J. Chromatogr. Sci.*, 33 (1995) 194-203.
- 1193 Zhang, X., Zhao, L. and Zhang, G.: (Prediction of gas chromatographic retention index by neural network.) *Yingyong Huaxue*, 11, No. 5 (1994) 54-59; *C.A.*, 122 (1995) 95603y.
- 1194 Zou, G. and Liu, Z.: (Demonstrating the tailing of normal chromatographic peak by theory of diffusion and mass transfer.) *Chin. J. Chromatogr.*, 13 (1995) 107.

See also 1268.

2b. Thermodynamics and theoretical relationships

- 1195 Afzal, M., Khan, M. and Ahmad, H.: Kinetics of the sorption of organic vapors in chromatographic silica gel. *Adsorpt. Sci. Technol.*, 11, No. 2 (1994) 113-122; *C.A.*, 122 (1995) 143511r.
- 1196 Kowalski, W.J.: Gas chromatographic enantioseparation of nucleophilic compounds on a stationary phase containing europium chelate. *Acta Chromatogr.*, 2 (1993) 84-88; *C.A.*, 122 (1995) 150550f.

See also 1342.

2c. Relationship between structure and chromatographic behaviour

- 1197 Défayes, G., Reddy, K.S., Dallos, A. and Kováts, E.sz.: Pair-wise interactions by gas chromatography V. Interaction free enthalpies of solutes with primary chloro- and bromoalkanes. *J. Chromatogr. A*, 699 (1995) 131-154.
- 1198 Jing, P. and Fu, R.: Supramolecular chemistry for gas chromatography. I. Relationship between supramolecular chemistry and gas chromatography. *Fenxi Huaxue*, 23 (1995) 104-110; *C.A.*, 122 (1995) 177343c - a review with 55 refs.

See also 1280, 1281, 1341, 1345, 1366, 1395, 1459, 1692.

2d. Measurement of physico-chemical and related values

- 1199 Balard, H., Wang, W.D. and Papirer, E.: Energetic surface heterogeneity of carbon black determined by IGC. *Polym. Mater. Sci. Eng.*, 70 (1993) 468-469; *C.A.*, 122 (1995) 39671g.
- 1200 Berezkina, L.G. and Sarycheva, E.A.: (Gas chromatographic study of dehydration of dicaliumphosphate dihydrate.) *Zh. Fiz. Khim.*, 68 (1994) 1304-1306.
- 1201 Berezkina, L.G., Sarycheva, E.A. and Sukhdolova, V.I.: (Investigation of de- and rehydration in system CaSO₄-water vapours by chromatographic methods.) *Zh. Fiz. Khim.*, 68 (1994) 1678-1681.
- 1202 Boneva, S. and Tashkova, K.: Gas chromatographic investigation of the interaction between vinyltriethoxysilane and aerosil and active charcoal surfaces. *Anal. Lab.*, 2, No. 1 (1993) 53-55; *C.A.*, 122 (1995) 204243d.
- 1203 Carraffo, H.M., Simon, L.D., Daly, J.W., Spande, T.F. and Jones, T.H.: Cis- and trans-configuration of α,α -disubstituted piperidines and pyrrolidines by GC-FTIR; application to decahydroquinoline stereochemistry. *Tetrahedron*, 50 (1994) 11329-11338; *C.A.*, 122 (1995) 10323j.
- 1204 Chickos, J.S., Hosseini, S. and Hesse, D.G.: Determination of vaporization enthalpies of simple organic molecules by correlations of changes in gas chromatographic net retention times. *Thermochim. Acta*, 249 (1995) 41-61; *C.A.*, 122 (1995) 171385w.

- 1205 Fafard, M., El-Kindi, M., Schreiber, H.P., Dipaola-Baranyi, G. and Hor, A.M.: Estimating surface energy variations of solids by inverse gas chromatography. *J. Adhes. Sci. Technol.*, 8 (1994) 1383-1394; C.A., 122 (1995) 82532z.
- 1206 Garnier, G. and Glasser, W.G.: Measurement of the surface free energy of amorphous cellulose by alkane adsorption: a critical evaluation of inverse gas chromatography (IGC). *J. Adhes.*, 46 (1994) 165-180; C.A., 122 (1995) 136443s.
- 1207 Genkin, A.N. and Petrova, N.A.: Some peculiarities of the reaction between Rh(I) carbonyl complexes and unsaturated hydrocarbons under the conditions of gas-liquid chromatography. *Rhodium Express*, 2 (1993) 12-18; C.A., 122 (1995) 45351u.
- 1208 Genkin, A.N. and Petrova, N.A.: The competition between unsaturated hydrocarbons in the course of their reactions with rhodium(I) carbonyl complexes under the conditions of GLC. *Rhodium Express*, 6 (1994) 19-26; C.A., 122 (1995) 133379q.
- 1209 Guan, Y.-A., Zhou, L.-M. and Li, L.-E.: Live retention database for identification of compounds in capillary gas chromatographic analysis. Principle and structure. *Chin. J. Chem.*, 12 (1994) 265-269; C.A., 121 (1994) 314902e.
- 1210 Hussam, A., Basu, S.G., Hixon, M. and Olumee, Z.: General method for the study of solute-surfactants association equilibria of volatile solutions by head space gas chromatography. *Anal. Chem.*, 67 (1995) 1459-1464.
- 1211 Jackson, P.L. and Huglin, M.B.: Use of inverse gas chromatography to measure diffusion coefficients in crosslinked polymers at different temperatures. *Eur. Polym. J.*, 31, No.1 (1995) 63-65; C.A., 122 (1995) 82786k.
- 1212 Jiang, X., Fan, X. and Yang, Z.: (Gas chromatography observation of enantioselective autoinduction effect in the synthesis of chiral alcohols.) *Chin. J. Chromatogr.*, 13 (1995) 110-112.
- 1213 Lai, C.-C. and Tan, C.-S.: Measurement of molecular diffusion coefficients in supercritical carbon dioxide using a coated capillary column. *Ind. Eng. Chem. Res.*, 34 (1995) 674-680; C.A., 122 (1995) 115575w.
- 1214 Langleron, M., Cavadias, S. and Amouroux, J.: Study by mass spectrometry and gas chromatography of toluene oxidation in low-pressure plasma reactor. *Rapid Commun. Mass Spectrom.*, 9 (1995) 18-22; C.A., 122 (1995) 164023y.
- 1215 Mahne, I. and Tiedje, J.M.: Criteria and methodology for identifying respiratory denitrifiers. *Appl. Environ. Microbiol.*, 61, No. 3 (1995) 1110-1115; C.A., 122 (1995) 186324z.
- 1216 Marinichev, A.N., Puchkova, L.V. and Stolyarov, B.V.: (Use of vapor-phase gas chromatographic analysis to study the solvolysis of tert.-butyl chloride.) *Zh. Prikl. Khim. (S.-Peterburg)*, 67, No. 7 (1994) 1132-1138.
- 1217 Nikolic, A.D., Ivancev, I.I. and Petrovic, S.D.: A thermodynamic study of N-n-pentylbenzamide solutions using gas-liquid chromatography. *J. Serb. Chem. Soc.*, 59 (1994) 633-638; C.A., 122 (1995) 40008c.
- 1218 Pershina, V., Fricke, B. and Ionova, G.V.: Theoretical study of the physicochemical properties of the light transactinides. *J. Alloys Compd.*, 213/214 (1994) 33-37; C.A., 121 (1994) 308532t.
- 1219 Raihane, M., Montheard, J.P. and Boinon, B.: Kinetics of thermal degradation of copolymers methylvinylidene cyanide with substituted styrenes. Thermogravimetric and gas chromatography-mass spectroscopy studies. *Polym. Degrad. Stab.*, 46 (1994) 171-179; C.A., 122 (1995) 82275t.
- 1220 Sandercock, P.M.L.: Comparison of passive charcoal adsorption with a dynamic charcoal adsorption technique. *J. - Can. Soc. Forensic Sci.*, 27 (1994) 179-201; C.A., 122 (1995) 2902g.
- 1221 Shaoxian, P., Wu, T., Qu, J. and Ji, Y.: (Study on adsorption of organic compounds on river sediment using gas chromatography.) *Fenxi Huaxue*, 22 (1994) 656-659; C.A., 122 (1995) 88770x.
- 1222 Sotiropoulou, V., Vassilev, G.P., Katsanos, N.A., Hetaxa, H. and Roubani-Kalantzopoulou, F.: Simple determination of experimental isotherms using diffusion denuder tubes. *J. Chem. Soc. Faraday Trans.*, 91 (1995) 485-492; C.A., 122 (1995) 109618w.
- 1223 Tsutsumi, K. and Ban, K.: Analysis of adsorption and adhesion behaviors of modified carbon fibers by means of inverse gas chromatography. *Stud. Surf. Sci. Catal.*, 80 (1993) 679-684; C.A., 122 (1995) 33446r.
- 1224 Voelkel, A. and Janas, J.: Elimination of adsorption effects of polarity parameters determined by inverse gas chromatography. *J. Chromatogr. A*, 693 (1995) 315-323.
- 1225 Wang, W.-D., Haidar, B., Vidal, A. and Donnet, J.-B.: Study of surface activity of carbon black by inverse gas chromatography. Part IV: Effect of mechanical action on surface activity of carbon black. *Kautsch. Gummi Kunstst.*, 47 (1994) 238-241; C.A., 122 (1995) 108308h.
- 1226 Wang, W.-D., Vidal, A., Nanse, G. and Donnet, J.-B.: Study of surface activity of carbon black by inverse gas chromatography. Part V: Modification of carbon black surface by silane. *Kautsch. Gummi Kunstst.*, 47 (1994) 493-501; C.A., 122 (1995) 135717x.
- 1227 Zechmeister-Boltenstern, S.: Measurement of actual and potential denitrification and soil respiration with an automated gas chromatographic system. *Bodenkultur*, 45 (1994) 219-226; C.A., 122 (1995) 55065t.

See also 1344, 1370, 1412, 1675, 1677, 1678, 1680, 1786.

3. GENERAL TECHNIQUES

3a. Apparatus and accessories

- 1228 Arutyunov, Yu.A., and Vigdergauz, M.S.: (Some specific features of chromatography. Apparatus and methodical aspects.) *Zh. Anal. Khim.*, 49 (1994) 796-803.
- 1229 Fulton, D.B.: Method of producing a gas chromatogram. *Can. Pat. Appl.* CA 2,091,279 (Cl. G01N30/00), 10 Sep. 1994, Appl. 09 Mar. 1993; 24 pp.; C.A., 122 (1995) 122273z.
- 1230 Ghaoui, L. and Green, L.S.: Dual vapor and liquid injector for gas chromatography. *J. Chromatogr. A*, 696 (1995) 235-244.
- 1231 Gonzáles, M.I., Carnicero, M., de la Torre, R., Ortuño, J. and Segura, J.: Influence of the injection technique on the thermal degradation of cocaine and its metabolites in gas chromatography. *J. Chromatogr. B*, 664 (1995) 317-327.
- 1232 Grob, K. and Munari, F.: Process and device for gas chromatographic analysis of large volume samples. *PCT Int. Appl.* WO 94 28,409 (Cl. G01/N30/12), 8 Dec. 1994, IT Appl. 93/Mi1058, 21 May 1993; 33 pp.; C.A., 122 (1995) 95644n.

- 1233 Hayashi, T.: (Gas chromatograph for separation of low boiling point components.) *Jpn. Kokai Tokkyo Koho JP 06,288,998* [94,288,998] (Cl. G01N30/46), 18 Oct. 1994, Appl. 93/78,112, 05 Apr. 1993; 5 pp.; C.A., 122 (1995) 122272y.
- 1234 Higdon, W.R.: Connector for capillary tubes. *Eur. Pat. Appl. EP 636,882* (Cl. G01N30/60), 01 Feb. 1995, US Appl. 100,022, 30 Jul. 1993; 17 pp.; C.A., 122 (1995) 177430d.
- 1235 Hinshaw, J.V.: Measuring flow. *LC-GC Int.*, 8 (1995) 137-142.
- 1236 Iida, K. and Yoshihara, T.: (Gas chromatograph.) *Jpn. Kokai Tokkyo Koho JP 06,324,028* [94,324,028] (Cl. G01N30/66), 25 Nov. 1994, Appl. 93/111,375, 13 May 1993; 4 pp.; C.A., 122 (1995) 177426g.
- 1237 Ivanov, T., Balkanski, A. and Barbov, D.: (Automated system for gas chromatographic diagnostic of oil-filled high-voltage equipment in electric power systems.) *Energetika (Sofia)*, 45, No. 3 (1994) 44-46; C.A., 122 (1995) 110323c.
- 1238 Jiao, K.: Development of field analysis for semi-volatile organic compounds by gas chromatography/mass spectrometry. Avail. *Univ. Microfilms Int.*, Order No. DA9425789, 1994, 194 pp.; C.A., 122 (1995) 150567s.
- 1239 Kolesar, E.S.Jr. and Reston, R.R.: Silicon micromachined gas chromatography system with a thin-film copper phthalocyanine stationary phase and its resulting performance. *Surf. Coat. Technol.*, 68/69 (1994) 679-685; C.A., 122 (1995) 95379d.
- 1240 Krylov, V.A., Krasotskii, S.G. and Yan'kov, S.V.: (Dosing system for aggressive volatile substances for highly sensitive gas-chromatographic determination of impurities.) *Vysokochist. Veshchestva*, 4 (1994) 157-160; C.A., 122 (1995) 203969h.
- 1241 Majors, R.E.: New approaches to sample preparation. *LC-GC Int.*, 8 (1995) 128-133.
- 1242 Mattsson, J. and Wirth, K.P.: (Automatic flow control - a promising technique in gas chromatography.) *Kem. Tidskr./Kemi-vaerlden*, 2 (1994) 20-22; C.A., 122 (1995) 121993x.
- 1243 Sivchenko, V.Ya., Maltsev, E.K., Ovcharenko, N.P., Bondarchuk, A.I., Kritskij, V.E. and Singovskij, S.V.: (Gas chromatograph.) *U.S.S.R. SU 1,807,384* (Cl. G01N30/10), 07 Apr. 1993, Appl. 4,913,720, 25 Feb. 1991; C.A., 122 (1995) 95639q.
- 1244 Tipler, A.: Gas chromatography instrumentation, operation, and experimental considerations. In: Baugh, P.J. (Editor), *Gas Chromatography*, IRL, Oxford, 1993, pp.15-70.
- 1245 Tirillini, B. and Stoppini, A.M.: Injection of a sample by means of microneedles followed by capillary gas chromatography. *J. Chromatogr. Sci.*, 33 (1995) 139-142.
- 1246 Torres, L. and Amiet, F.: Method for continuous analysis of volatile organic compounds in an atmosphere to be analyzed and corresponding concentration unit and chromatograph. *PCT Int. Appl. WO 94 28,408* (Cl. G01N30/08), 08 Dec. 1994, FR Appl. 93/6,628, 01 Jun. 1993; 32 pp.; C.A., 122 (1995) 122264x.
- 1247 Wiegand P.M., Fisher, J.F., Parrish, J.R. and Ballard, E.S.: (Fast gas chromatography method, apparatus and applications.) *Eur. Pat. Appl. EP 633,469* (Cl. G01N30/30), 11 Jan. 1995, US Appl. 88,677, 07 Jul. 1993; 27 pp.; C.A., 122 (1995) 95646q.
- See also 1227, 1294, 1326, 1357, 1385, 1392.
- 3b. *Detectors and detection reagents*
- 1248 Aue, W.A. and Thurbide, K.B.: Reactive-flow luminiscence with flame ionization detection. *J. Chromatogr. A*, 699 (1995) 183-193.
- 1249 Aue, W.A., Eisener, C.G., Gebhardt, J.A. and Lowery, N.B.: Second-channel design for a single-channel flame photometric detector. *J. Chromatogr. A*, 699 (1995) 195-201.
- 1250 Blumberg, L.M. and Dandeneau, R.D.: The effect of capillary column diameter on the performance of thermal conductivity detectors. *J. High. Resolut. Chromatogr.*, 18 (1995) 235-242.
- 1251 Goode, S.R. and Thomas, C.L.: Characterizing the factors that influence oxygen selectivity in gas chromatography-microwave-induced plasma atomic emission spectrometry. *J. Anal. At. Spectrom.*, 9 (1994) 965-970; C.A., 121 (1994) 314637x.
- 1252 Hackett, M., Wang, H., Miller, G.C. and Bornhop, D.J.: Ultraviolet-visible detection for capillary gas chromatography and combined ultraviolet-mass spectrometry using a remote flow cell. *J. Chromatogr. A*, 695 (1995) 243-257.
- 1253 Jones, M.E.: A simple-to-build thermal-conductivity GC detector. *J. Chem. Educ.*, 71 (1994) 995-996; C.A., 122 (1995) 9084a.
- 1254 Kalontarov, L., Jing, H., Amirav, A. and Cheskis, S.: Mechanism of sulfur emission quenching in flame photometric detectors. *J. Chromatogr. A*, 696 (1995) 245-256.
- 1255 Lacey, R.F.: Thermal conductivity detector. *U.S. US 5,379,630* (Cl. 73-25.03; G01N25/18), 10 Jan. 1995, Appl. 84,652, 28 Jun. 1993; 8 pp.; C.A., 122 (1995) 150608f.
- 1256 Mistry, B.S., Reineccius, G.A. and Jasper, B.L.: Comparison of gas chromatographic detectors for the analysis of volatile sulfur compounds in foods. *ACS Symp. Ser.*, 564 (1994) 8-21; C.A., 121 (1994) 299395w.
- 1257 Roth, E. and Kiefer, W.: Surface-enhanced Raman spectroscopy as a detection in gas chromatography. *Appl. Spectrosc.*, 48 (1994) 1193-1195.
- 1258 Sandmann B.W.: Peroxyoxalate chemiluminiscence detection with chromatography: 1. Development and evaluation of an interface for packed column supercritical fluid chromatography. 2. Quinoxalinone derivatization of carboxylic acids for high-performance liquid chromatography. Avail. *Univ. Microfilms Int.*, Order No. DA9420284, 1993, 95 pp.; C.A., 122 (1995) 150569u.
- 1259 Seo, Y., Umemura, K. and Ichise, M.: (Gas electrode for selective detection of sulfur compounds in gasoline.) *Bunseki Kagaku*, 44 (1995) 277-282.
- 1260 Steely, J.S.: Chemiluminiscence detection of sulfur compounds in cooked milk. *ACS Symp. Ser.*, 564 (1994) 22-35; C.A., 121 (1994) 299396x.
- 1261 Thurbide, K.B. and Aue, W.A.: Quenching-free reactive-flow photometry. *J. Chromatogr. A*, 694 (1995) 433-440.
- 1262 Tzanani, N. and Amirav, A.: Combined pulsed flame photometric ionization detector. *Anal. Chem.*, 67 (1995) 167-173.
- 1263 Witkiewicz, Z.: (Atomic emission detector for gas chromatography.) *Przem. Chem.*, 73 (1994) 460-462 C.A., 122 (1995) 121973r.
- 1264 Yu, A., Tian, X., Yang, W. and Jin, Q.: (Microwave-induced plasma ionization detector for gas chromatography.) *Jilin Daxue Ziran Kexue Xuebao*, 2 (1994) 105-108; C.A., 122 (1995) 150531a.

- 1265 Zhu, J.: (Diagnosis and treatment of contamination in gas chromatographical detectors.) *Huagong Zidonghua Ji Yibiao*, 21, No. 2 (1994) 58-59; C.A., 122 (1995) 150521x.
- See also 1236, 1329, 1332, 1339, 1346.
- 3c. *Sorbents and columns, packing procedures*
- 1266 Bao, L., Dai, R., Deng, Y. and Fu, R.: (The preparation and characterization of high temperature gas chromatographic column.) *Chin. J. Chromatogr.*, 13 (1995) 108-109.
- 1267 Baugh, P.J.: Applications of packed and capillary GC. In: Baugh, P.J. (Editor), *Gas Chromatography*, IRL, Oxford, 1993, pp.103-131.
- 1268 Berezkin, V.G., Malyukowa, I.V. and Alishoev, V.R.: Dependence of retention of organic compounds on carrier gas pressure in capillary adsorption chromatography. *J. Chromatogr. A*, 699 (1995) 155-159.
- 1269 Cui, T. and Fu, C.: (High performance liquid chromatographic detection of trace polyamines in plant tissue by pre-column derivatization with phthalimidylbenz磺onyl chloride.) *Chin. J. Chromatogr.*, 13 (1995) 117-119.
- 1270 Dawes, P.A.: Development, technology, and utilization of capillary columns for gas chromatography. In: Baugh, P.J. (Editor), *Gas Chromatography*, IRL, Oxford, 1993, pp. 71-102; C.A., 122 (1995) 150504u.
- 1271 Dimandja, J.-M., Valentin, J.R. and Phillips, J.B.: Gas chromatographic column for the storage of sample profiles. *Anal. Chim. Acta.*, 299 (1994) 29-36.
- 1272 GavriloVA, T.B., Nikitin, Yu, S. and Rudnitskaya, T.A.: (Adsorbents on the basis of sym-Heptazine.) *Zh. Fiz. Khim.*, 68 (1994) 545-550.
- 1273 Gawdzik, B. and Matynia, T.: Studies of the selectivity of di(methacryloxymethyl)-*m*-xylene-divinylbenzene porous copolymers for gas chromatography. *Chem. Anal. (Warsaw)*, 39 (1994) 161-166.
- 1274 Huang, Z., Cheng, Y., Cai, G., Fu, R. and Jing, P.: (Synthesis of crown ethers. Part XXIV. Synthesis of liquid crystal crown ether and its use as stationary phase for capillary gas chromatography.) *Wuhan Daxue Xuebao, Ziran Kexueban*, 6 (1993) 93-99; C.A., 122 (1995) 57221q.
- 1275 Lao, H. and Detellier, C.: Gas chromatographic separation of linear hydrocarbons on microporous organo-smectites. *Clays Clay Miner.*, 42 (1994) 477-481; C.A., 121 (1994) 304341g.
- 1276 Li, W., Malik, A. and Lee, M.L.: Fused silica packed capillary columns in supercritical fluid chromatography. *J. Microcolumn Separ.*, 6 (1994) 557-563.
- 1277 Liu, L., Tang, M., Han, J. and Shen, H.: (Epichlorhydrin crosslinked oligopolymer of β -cyclodextrin as a stationary phase in gas chromatography.) *Fenxi Huaxue*, 22 (1994) 905-908; C.A., 122 (1995) 95602x.
- 1278 Mñuk, P. and Feltl, L.: Gas chromatographic study of the inclusion properties of calixarenes I. *p*-tert-Butylcalix[4]arene in a micropacked column. *J. Chromatogr. A*, 696 (1995) 101-112.
- 1279 Naohiro, N.K.: (Inorganic porous chromatographic column.) *Jpn. Kokkai Tokkyo Koho* JP 06,265,534 [94,265,534] (Cl. G01N30/60), 22 Sep. 1994, JP Appl. 93/23,670, 18 Jan. 1993; 8 pp.; C.A., 122 (1995) 70989r.
- 1280 Poole, S.K. and Poole, C.F.: Application of principal component factor analysis to the cavity model of solvation to identify factors important in characterizing the solvent properties of gas chromatographic stationary phases. *J. Chromatogr. A*, 697 (1995) 429-440.
- 1281 Poole, S.K. and Poole, C.F.: Chemometric classification of the solvent properties (selectivity) of commonly used gas chromatographic stationary phases. *J. Chromatogr. A*, 697 (1995) 415-427.
- 1282 Ruan, Z. and Liu, H.: Preparation of 4-vinylpyridine and divinylbenzene porous-layer open tubular columns by *in situ* copolymerization. *J. Chromatogr. A*, 693 (1995) 79-88.
- 1283 Tambe, A.S., Biswas, S.S. and Ayyangar, N.R.: Synthesis of novel asters of 4-(2'-phenylisopropyl)phenol used as column packings for gas liquid chromatography. *Indian J. Chem. Technol.*, (1994) 371-372; C.A., 122 (1995) 136681t.
- 1284 Xu, H.-S. and Hong, H.: (Study of the gradient effect of a new-kind of gradient loaded GC column.) *Gaodeng Xuexiao Huaxue Xuebao*, 15 (1994) 1152-1154; C.A., 122 (1995) 22841q.
- 1285 Xu, Y.: (Sample capacity of capillary column.) *Hebei Shifan Daxue Xuebao, Ziran Kexueban*, 18, No. 4 (1994) 68-71; C.A., 122 (1995) 177384s.
- See also 1189, 1195, 1239, 1314, 1315, 1317, 1320, 1322, 1331, 1340, 1384, 1556.
- 3d. *Quantitative analysis*
- 1286 Below, L.: (Methods for certification of calibration gases for the operation of process gas chromatographs.) *Gas-Wasserfach: Gas/Erdgas*, 135 (1994) 1-7; C.A., 122 (1995) 191817s - a review with 5 refs.
- 1287 Gu, H., Zhang, L., Li, B. and Liu, L.: (Quality control and quality assurance of GC-MS in international interlaboratory comparison test.) *Huaxue Tongbao*, 10 (1994) 37-40; C.A., 122 (1995) 95565n.
- 1288 Hachenberg, H.: (Calibration and result evaluation in statistic head-space gas chromatography.) *LaborPraxis* 18, No 8 (1994) 32-34.
- 1289 Schwertner, H.A.: Method with addition of fatty acid analog internal standard to blood for concentration-based analysis of lipid fatty acids, and its use in determining the likelihood that a patient is at risk for coronary artery disease or diabetes. *U.S. US 5,362,649* (Cl. 436-71; G01N33/92), 08 Nov. 1994, Appl. 19,582, 16 Feb. 1993; 7 pp.; C.A., 122 (1995) 27254k.
- See also 1311, 1631, 1647.
- 3e. *Preparative scale chromatography*
- See 1333, 1710.
- 3f. *Programmed temperature, pressure, vapors, gradients*
- 1290 Ferchichi, L. and Messadi, D.: (Retention and column efficiency simulations in programmed temperature gas chromatography: theory and application.) *Analisis*, 22 (1994) 392-399; C.A., 122 (1995) 122224j.

- 1291 Gerbino, T.C. and Castello, G.: Prediction of programmed temperature retention indices on capillary columns of different polarities. *J. Chromatogr. A*, 699 (1995) 161-171.
- 1292 Krylov, A.I., Vasil'eva, I.A. and Khlebnikova, N.S.: (Effect of parameters of chromatographic separation on the accuracy and precision in the determination of retention indexes with temperature programming). *Zh. Anal. Khim.*, 49 (1994) 1097-1103.
- 1293 Rubey, W.A.: An instrumentation assembly for studying operational behavior of thermal gradient programmed gas chromatography. *Rev. Sci. Instrum.*, 65 (1994) 2802-2807.
- See also 1268, 1327.
4. SPECIAL TECHNIQUES
- 4a. Automation
- 1294 Amamya, M., Shibata, S. and Matsura, T.: (Process gas chromatography.) *Jpn. Kokai Tokkyo Koho JP 06,242,096 [94,242,096]* (Cl. G01N30/68), 02 Sep. 1994, Appl. 93/25,320, 15 Feb. 1993; 4 pp.; C.A., 122 (1995) 45312g.
- See also 1286, 1392, 1722.
- 4b. Computerization and modelling
- 1295 Dimov, N. and Lekova, K.: (Software products in chromatographic analytical activities.) *Anal. Lab.*, 3, No. 2 (1994) 110-115; C.A., 122 (1995) 203901e - a review with 28 refs.
- See also 1395, 1543.
- 4c. Combination with other physico-chemical techniques (MS, IR etc.)
- 1296 Ardrey, R.E., Bukowski, N. and Gummersbach, J.: (Negative-chemical ionization eliminates matrix effects.) *Laborpraxis*, 18, No. 11 (1994) 92-94.
- 1297 Braumann, U., Händel, H., Albert, K., Ecker, R. and Spraul, M.: On-line monitoring of the supercritical fluid extraction process with proton nuclear magnetic resonance spectroscopy. *Anal. Chem.*, 67 (1995) 930-935.
- 1298 Collman, W.M., III and Gordon, B.M.: Analysis of natural products by gas chromatography/matrix isolation/infrared spectrometry. *Adv. Chromatogr. (N.Y.)*, 34 (1994) 57-108; C.A., 122 (1995) 204147a - a review with 51 refs.
- 1299 Dagan, S. and Amirav, A.: Fast, high temperature and thermolabile GC-MS in supersonic molecular beams. *Int. J. Mass Spectrom. Ion Processes*, 133 (1994) 187-210; C.A., 122 (1995) 22827q.
- 1300 Feigel, C.: Ion trap GC-MS-MS with a benchtop mass spectrometer. *Spectroscopy (Eugene)*, 9, No. 5 (1994) 31-33; C.A., 122 (1995) 150515y.
- 1301 Niwa, M. and Imai, M.: (Gas chromatograph/mass spectrometer and its applications to the analysis of lubricating oils.) *Sumitomo Keikinokoku Gihō*, 35 (1994) 103-109; C.A., 122 (1995) 122192x.
- 1302 Seeterlin, M.A.: A tandem time-of-flight mass spectrometer: MS/MS on the chromatographic time scale. Avail. *Univ. Microfilms Int.*, Order No. DA9431311, 1994, 128 pp.; C.A., 122 (1995) 173242w.
- 1303 Ye, M., Ye, H., Zeng, X. and Chen, Z.: (Study on combined gas chromatography and infrared spectrometry.) *Huazhong Ligong Daxue Xuebao*, 22, No. 10 (1994) 24-28; C.A., 122 (1995) 150532b.
- See also 1252, 1361, 1366, 1465, 1526, 1547, 1554, 1585, 1593, 1596, 1673.
- 4e. Functional analysis
- 1304 Watson, D.G.: Chemical derivatization in gas chromatography. In: Baugh, P.J. (Editor), *Gas Chromatography*, IRL, Oxford, 1993, pp.133-170.
- See also 1473, 1782, 1788.
- 4f. Trace analysis and pre-separation techniques
- 1305 Benfenati, E., Garofani, S., Natangelo, M., Mangiapani, S. and Fanelli, R.: Impurities released from extractive phases used in the analysis of pesticides. *Intern. J. Environ. Anal. Chem.*, 58 (1995) 23-30.
- 1306 Crowley, X.W., Murugaiah, V., Naim, A. and Giese, R.W.: Masking as a mechanism for evaporative loss of trace analyte, especially after solid-phase extraction. *J. Chromatogr. A*, 699 (1995) 395-402.
- 1307 Ferreira, V., Fernández, P., Meléndez, J. and Cacho, J.: Analytical characteristics of sample evaporation with the micro-Kuderna-Danish concentrator. *J. Chromatogr. A*, 695 (1995) 41-55.
- 1308 Pawliszyn, J.: New directions in sample preparation for analysis of organic compounds. *TrAC*, 14 (1995) 113-122 - a review with 30 refs.
- 1309 Skrabakova, S. and Matisova, E.: (Application of carbon sorbents for enrichment of pollutants in environmental samples.) *Chem. Listy*, 89 (1995) 180-191 - a review with 104 refs.
- 1310 Skrabakova, S., Matisova, E., Onderova, M., Novak, I. and Berek, D.: The testing of carbon sorbent for preconcentration of volatile organic trace compounds. *Chem. Zvesti*, 48 (1994) 169-174.
- 1311 Sunesson, A.-L., Nilsson, C.-A. and Andersson, B.: Evaluation of adsorbents for sampling and quantitative analysis of microbial volatiles using thermal desorption-gas chromatography. *J. Chromatogr. A*, 699 (1995) 203-214.
- 1312 Zapf, A., Heyer, R. and Stan, H.-J.: Rapid micro liquid-liquid extraction method for trace analysis of organic contaminants in drinking water. *J. Chromatogr. A*, 694 (1995) 453-461.
- See also 1802.
- 4g. Enantiomers, separation
- 1313 Koppenhoefer, B. and Epperlein, U.: Overview of the gas chromatographic separation of enantiomers containing a three-membered ring. *J. Chromatogr. Sci.*, 33 (1995) 244-255.

- 1314 Koppenhoefer, B., Mühleck, U. and Lohmiller, K.: Backbone modification of Chirasil-Val I. Effect of loading on the separation of enantiomers by gas chromatography. *J. Chromatogr. A*, 699 (1995) 215-221.
- 1315 Koppenhoefer, B., Mühleck, U., Walsler, M. and Lohmiller, K.: Backbone modification of Chirasil-Val, part II: Introduction of a rigid cyclohexyl spacer. *J. Chromatogr. Sci.*, 33 (1995) 217-222.
- 1316 Leal, W.S., Shi, X., Liang, D., Schal, C. and Meinwald, J.: Application of chiral gas chromatography with electroantennographic detection to the determination of the stereochemistry of a cockroach sex pheromone. *Proc. Natl. Acad. Sci. U. S. A.*, 92 (1995) 1033-1037; *C.A.*, 122 (1995) 182318c.
- 1317 Maas, B., Dietrich, A., Bartschat, D. and Mosandl, A.: Tert-butylidimethylsilylated cyclodextrins: Versatile chiral stationary phases in capillary gas chromatography. *J. Chromatogr. Sci.*, 33 (1995) 223-228.
- 1318 Reinhardt, R., Engewald, W., Himmelreich, U., Christian, B. and Koppenhoefer, B.: Enantiomer separation of cycloalkanols by capillary gas chromatography on cyclodextrin phases. *J. Chromatogr. Sci.*, 33 (1995) 236-243.
- 1319 Smith, R.J., Taylor, D.R. and Wilkins, S.M.: Temperature dependence of chiral discrimination in supercritical fluid chromatography and high-performance liquid chromatography. *J. Chromatogr. A*, 697 (1995) 591-596.
- 1320 Takeichi, T., Toriyama, H., Shimura, S., Takayama, Y. and Morikawa, M.: Cyclodextrin carbamates as novel chiral stationary phase for capillary gas chromatography. *J. High Resolut. Chromatogr.*, 18 (1995) 179-190.
- 1321 Taylor, D.R.: Chiral separations by gas chromatography. In: Baugh, P.J. (Editor), *Gas Chromatography*, IRL, Oxford, 1993, pp.237-281.
- 1322 Wilkins, S.M., Taylor, D.R. and Smith, R.J.: Enantiomer separation by supercritical fluid chromatography on a chiral stationary phase physically anchored to porous graphitic carbon. *J. Chromatogr. A*, 697 (1995) 587-590.
- 1323 Zhou, X., Wan, H. and Ou, Q.: Gas chromatographic separation mechanism of enantiomers on 2,6-O-dibutyl-3-O-trifluoroacetylated- γ -cyclodextrin. *Fenxi Huaxue*, 22 (1994) 560-564; *C.A.*, 121 (1994) 314884a.
- See also 1189, 1196, 1212, 1419, 1431, 1447, 1450, 1468, 1470, 1494, 1523, 1540, 1541, 1570, 1579, 1644, 1687, 1693, 1710, 1711, 1741.
- 4h. *Other special techniques*
- 1324 Akrad, M.L.: Turnable selectivity strategies for high-speed gas chromatography. Avail. *Univ. Microfilms Int.*, Order No. DA9423132, 1994, 192 pp.; *C.A.*, 122 (1995) 150566r.
- 1325 Williams, D.: Inverse gas chromatography. In: Ishida, H. (Editor), *Charact. Compos. Mater.*, Butterworth-Heinemann, Boston, 1994, pp. 80-104; *C.A.*, 122 (1995) 143229e.
- See also 1787.
- 4i. *Supercritical fluid chromatography*
- 1326 Allington, R.W., Walters, H.L., Jameson, D.G. and Tehrani, Y.: Apparatus and method for supercritical fluid extraction or supercritical fluid chromatography. *PCT Int. Appl.* WO 95 03,106 (Cl. B01D15/08), 02 Feb. 1995, US Appl. 96,919, 23 Jul. 1993; 212 pp.; *C.A.*, 122 (1995) 191211q.
- 1327 Bartle, K.D., Bevan, C.D., Clifford, A.A., Jafar, S.A., Naila, M. and Verrall, M.S.: Preparative-scale supercritical fluid chromatography. *J. Chromatogr. A*, 697 (1995) 579-585.
- 1328 Blackwell, J.A. and Schallinger, L.E.: Hydrofluorocarbon and perfluorocarbon mobile phases for capillary supercritical fluid chromatography. *J. Microcolumn Separ.*, 6 (1994) 551-556.
- 1329 Brown, F.R., Roehl, R., Tabajonda, J. and Garcha, J.S.: Evaluation and application of supercritical fluid chromatography with electrolytic conductivity detection. *Anal. Lett.*, 28 (1995) 703-729.
- 1330 Champagne, A.T. and Bienkowski, P.R.: The supercritical fluid extraction of anthracene and pyrene from a model soil: Equilibrium study. *Sep. Sci. Technol.*, 30 (1995) 1289-1307.
- 1331 Dressman, S.F. and Michael, A.C.: On-line electrochemical detection for supercritical fluid chromatography. *Anal. Chem.*, 67 (1995) 1339-1345.
- 1332 Duccate, G.R.: Aspects of the microwave-induced plasma as an element selective detector for supercritical fluid chromatography and supercritical fluid extraction. Avail. *Univ. Microfilms Int.*, Order No. DA9419207, 1993, 115 pp.; *C.A.*, 122 (1995) 177100w.
- 1333 Hanson, M.: Small-scale preparative supercritical fluid chromatography of cyproterone acetate. *Chromatographia*, 40 (1995) 139-142.
- 1334 Lanças, F.M., Queiroz, M.E.C. and Silva, I.C.E.: Simplex optimization of extraction of soybean oil by supercritical pentane. *Chromatographia*, 40 (1995) 421-424.
- 1335 Maxwell, R.J., Lightfield, A.R. and Stolker, A.A.M.: An SPE column-teflon sleeve assembly for in-line retention during supercritical fluid extraction of analytes from biological matrices. *J. High. Resolut. Chromatogr.*, 18 (1995) 231-234.
- 1336 McNally, M.E.P.: Advances in environmental SFE. *Anal. Chem.*, 67, No. 9 (1995) 308A-315A - a review with 30 refs.
- 1337 Messer, D.C., Taylor, L.T., Weiser, W.E., McRae, J.W. and Cook, C.C.: Quantitative analysis of a drug in an animal feed employing sample preparation by supercritical fluid extraction. *Pharm. Res.*, 11 (1994) 1545-1548; *C.A.*, 121 (1994) 299436k.
- 1338 Mikesova, M. and Vejrosta, J.: (Publication activity in supercritical fluid extraction (SFE).) *Chem. Listy*, 89 (1995) 217-222.
- 1339 Miller, D.J. and Hawthorne, S.B.: Determination of solubilities of organic solutes in supercritical CO₂ by online flame ionization detection. *Anal. Chem.*, 67 (1995) 273-279.
- 1340 Morrison, J.F. and Lee, M.L.: Effect of phase behavior on supercritical fluid chromatographic and extraction performance. *NATO ASI Ser., Ser. E*, 273 (1994) 641-652; *C.A.*, 122 (1995) 204208w.
- 1341 Nishikawa, Y.: Studies on evaluation of retention and separation of pesticides by supercritical fluid chromatography. *Bunseki Kagaku*, 44 (1995) 247-248.
- 1342 Roth, M.: Current challenges in thermodynamics of supercritical fluid chromatography. *NATO ASI Ser., Ser. E*, 273 (1994) 631-639; *C.A.*, 122 (1995) 170851h.

- 1343 Saito, M., Yamauchi, Y. and Okuyama, T.: *Fractionation by Packed-Column SFC and SFE: Principles and Applications*. VCH, New York, 1994, 276 p.; C.A., 122 (1995) 204267q.
- 1344 Simoes, P.C., Matos, H.A., Carmelo, P.J., Gomes de Azevedo, E. and Nunes da Ponte, M.: Mass transfer on countercurrent packed columns: Application to supercritical CO₂ extraction of terpenes. *Ind. Eng. Chem. Res.*, 34, No. 2 (1995) 613-618; C.A., 122 (1995) 89055e.
- 1345 Wang, F.A., Jiang, Y.L., Jiang, D.G. and Wang, W.C.: The relationship between the retention value and carbon number of homologues in supercritical fluid chromatography. *J. Chromatogr. Sci.*, 33 (1995) 71-74.
- 1346 Zegers, B.N., Hessels, R., Jagesar, J., Rozenbrand, J., Lingeman, H. and Brinkman, U.A.T.: Photoionization detection in packed-capillary liquid and supercritical-fluid chromatography. *J. Liq. Chromatogr.*, 18 (1995) 413-440; C.A., 122 (1995) 177139r.
- See also 1213, 1258, 1276, 1297, 1319, 1322, 1367, 1373, 1414, 1498, 1503, 1579, 1603, 1611, 1621, 1629, 1638, 1664, 1673, 1693, 1711, 1717, 1758, 1761, 1768, 1780.
5. HYDROCARBONS AND HALOGEN DERIVATIVES
- 5a. *Aliphatic hydrocarbons*
- 1347 Bechtold, W.E., Strunk, M.R., Thornton-Manning, J.R. and Henderson, R.F.: Analysis of butadiene, butadiene monoxide, and butadiene dioxide in blood by gas chromatography/gas chromatography/mass spectrometry. *Chem. Res. Toxicol.*, 8 (1995) 182-187; C.A., 122 (1995) 125434p.
- 1348 Choi, Y.-W. and Choe, K.-H.: (Determination of natural gas components by gas chromatographic multicolumn system.) *Anal. Sci. Technol.*, 7 (1994) 339-347; C.A., 122 (1995) 85124d.
- 1349 Grob, K. and Bronz, M.: Analytical problems in determining 3,5-stigmastadiene and campestadiene in edible oils. *Riv. Ital. Sostanze Grasse*, 71 (1994) 291-295; C.A., 122 (1995) 54325r.
- 1350 Kawano, T. and Shimokawa, K.: A simple method for vacuum extraction and quantitative determination of internal ethylene of excised apple tissue. *Engei Gakkai Zasshi*, 63 (1994) 453-459; C.A., 122 (1995) 79321z.
- 1351 Kohlmüller, D. and Kochen, W.: (Analysis of the human expiration air: a challenge for medicine and environmental toxicology.) *CLB Chem. Labor Biotech.*, 45 (1994) 17-21; C.A., 122 (1995) 195380s.
- 1352 Reis Alves, L.A., Grob, R., Min, M. and de Desmolin, M.: (Improvement for the natural gas analysis.) *Analisis*, 22 (1994) 490-494; C.A., 122 (1995) 165020g.
- 1353 Rho, B.R., Park, B.E., Park, D.S. and Son, M.R.: Determination of trace methane impurities in ultra-high purity gases by gas chromatography. *Bull. Korean Chem. Soc.*, 15 (1994) 811-814; C.A., 122 (1995) 71103j.
- 1354 Wu, X.-m.: (A technique for the evaluation of shallow gaseous hydrocarbons and results of its application.) *Shiyou Kantan Yu Kaifa*, 21, No. 1 (1994) 40-45; C.A., 122 (1995) 110256h.
- See also 1275, 1522, 1744, 1796, 1802.
- 5b. *Cyclic hydrocarbons, fullerenes*
- 1355 Ciupe, R., Spangenberg, J., Meyer, T. and Wild, G.: (Solid-phase extraction system for gas chromatographic determination of PAHs (polycyclic aromatic hydrocarbons) in oil samples.) *GIT Fachz. Lab.*, 38 (1994) 825-829; C.A., 121 (1994) 304262g.
- 1356 Ehret-Henry, J., Cucruet, V., Luciani, A. and Feigenbaum, A.: Styrene and ethylbenzene migration from polystyrene into dairy products by dynamic purge-and-trap gas chromatography. *J. Food Sci.*, 59 (1994) 990-992 and 1001; C.A., 122 (1995) 7935b.
- 1357 Górecki, T. and Pawliszyn, J.: Solid phase microextraction/isothermal GC for rapid analysis of complex organic samples. *J. High Resolut. Chromatogr.*, 18 (1995) 161-166.
- 1358 Heo, G.S. and Kim, D.H.: (A rapid method for analyzing polycyclic aromatic hydrocarbons (PAHs) in urban dust using supercritical fluid extraction (SFE) and gas chromatography/mass spectrometry (GC/MS).) *J. Korean Chem. Soc.*, 38, No. 10 (1994) 726-733; C.A., 121 (1994) 307085n.
- 1359 Kok, P.W. and Ong, C.N.: Blood and urinary benzene determined by headspace gas chromatography with photoionization detection: application in biological monitoring of low-level non-occupational exposure. *Int. Arch. Occup. Environ. Health*, 66 (1994) 195-201; C.A., 122 (1995) 48566d.
- 1360 Komolprasert, V., Hargraves, W.A. and Armstrong, D.J.: Determination of benzene residues in recycled polyethylene terephthalate (PETE) by dynamic headspace-gas chromatography. *Food Addit. Contam.*, 11 (1994) 605-614; C.A., 122 (1995) 79344j.
- 1361 Kruge, M.A. and Bensley, D.F.: Flash pyrolysis-gas chromatography-mass spectrometry of lower Kittanning vitrinites. Changes in distributions of polyaromatic hydrocarbons as a function of coal rank. *ACS Symp. Ser.*, 570 (1994) 136-148; C.A., 122 (1995) 34854j.
- 1362 Landrock, A., Richter, H. and Merten, H.: Water Cl⁺, a new selective and highly sensitive method for the detection of environmental components using ion trap mass spectrometer. *Fresenius J. Anal. Chem.*, 351 (1995) 536-543.
- 1363 Li, R., Wang, Y. and Cao, L.: (Determination of benzene and C₇-C₁₀ alkyl substitute benzenes in No.2 mixed benzene by capillary gas chromatography.) *Shiyou Huagong*, 23 (1994) 83-815; C.A., 122 (1995) 110343j.
- 1364 Peschke, L. and Haegel, F.-H.: Improved analysis of polycyclic aromatic hydrocarbons in aqueous solutions by gas chromatography. *Fresenius J. Anal. Chem.*, 351 (1995) 622-624.
- 1365 Pilchowski, K. and Richartz, T.: Automatic gas chromatographic determination of BTX aromatics in ambient air. *J. Prakt. Chem./Chem.-Ztg.*, 336 (1994) 617-619; C.A., 121 (1994) 307074h.
- 1366 Radke, M., Budzinski, H., Pierard, C., Willsch, H. and Garrigues, P.: Chromatographic retention behavior of trimethylphenanthrenes and their identification by GC-FTIR in geological samples. *Polycyclic. Aromat. Compd.*, 3(Suppl.) (1993) 467-474; C.A., 121 (1994) 314878b.
- 1367 Reimer, G. and Suarez, A.: Comparison of supercritical fluid extraction and Soxhlet extraction for the analysis of native polycyclic aromatic hydrocarbons in soils. *J. Chromatogr. A*, 699 (1995) 253-263.

- 1368 Skrbic, B.D.: Unified retention indexes of some alkylbenzenes and bicyclic and related compounds on OV-101 and SE-30. *J. Serb. Chem. Soc.*, 59 (1994) 153-155; *C.A.*, 121 (1994) 314883z.
- 1369 Wilgoerth, U. and Schwedt, G.: (Relating to practice of PAH determination in waste oil. Method comparison GC-HPLC.) *J. Prakt. Chem./Chem.-Ztg.*, 336 (1994) 703-706; *C.A.*, 122 (1995) 113999p.
- See also 1345, 1394, 1686.
- 5c. *Halogen derivatives*
- 1370 Brunazzi, E., Pagliani, A. and Petarea, L.: Measurement of activity coefficients at infinite dilution of chlorinated solvents in commercial polyethylene glycol ethers. *J. Chem. Eng. Data*, 40 (1995) 443-447.
- 1371 Brush, R.C. and Rice, G.W.: Trihalomethanes produced in humic acid reactions: a GC-MS experiment for instrumental analysis. *J. Chem. Educ.*, 71, No. 12 (1994) A293-296; *C.A.*, 122 (1995) 55240w.
- 1372 Buethe, A. and Denker, E.: Qualitative and quantitative determination of PCB congeners by using a HT-5 GC column and an efficient quadrupole MS. *Chemosphere*, 30 (1995) 753-771.
- 1373 Castellano, T.M., Giglio, J.J., Evans, E.H. and Caruso, J.A.: Evaluation of low pressure inductively coupled plasma mass spectrometry for the analysis of gaseous samples. *J. Anal. At. Spectrom.*, 9 (1994) 1335-1340; *C.A.*, 122 (1995) 122208g.
- 1374 Chu, S., Yang, C. and Xu, X.: (Separation of polychlorobiphenyls by alumina-silica(Ag⁺) combined column for capillary GC analysis.) *Huanjing Huaxue*, 13 (1994) 555-560; *C.A.*, 122 (1995) 196364b.
- 1375 Cooper, C.M., Chang, S.F., Harrison, L.I.: Determination of the chlorofluorocarbon substitute 1,1,1,2-tetrafluoroethane (HFA-134a) in human and animal blood using gas chromatography with headspace analysis. *J. Chromatogr. B*, 667 (1995) 85-93.
- 1376 De Boer, J., Dao, Q.T., Wester, P.G., Bøwadt, S. and Brinkman, U.A.T.: Determination of mono-ortho substituted chlorobiphenyls by multidimensional gas chromatography and their contribution to TCDD equivalents. *Anal. Chim. Acta*, 300 (1995) 135-165.
- 1377 Djozan, D. and Assadi, Y.: Optimization of the gas stripping and cryogenic trapping method for capillary gas chromatographic analysis of traces of volatile halogenated compounds in drinking water. *J. Chromatogr. A*, 697 (1995) 525-532.
- 1378 Geissler, M. and Weimar, A.: (Determination of polychlorinated biphenyls (PCBs) by gas chromatography-mass spectrometry (GC-MS).) *GIT Spez. Chromatogr.*, 14, No. 1 (1994) 29-33; *C.A.*, 122 (1995) 22852u.
- 1379 Hajslová, J., Cuhra, P., Kempný, M., Poustka, J., Holadová, K. and Kocourek, V.: Determination of polychlorinated biphenyls in biotic matrices using gas chromatography-microwave-induced plasma atomic emission spectrometry. *J. Chromatogr. A*, 699 (1995) 231-239.
- 1380 Hale, R.C. and Taylor, M.O.: Determination of PCBs in fish tissues using supercritical fluid extraction. *Environ. Sci. Technol.*, 29 (1995) 1043-1047.
- 1381 Imagawa, T.: Synthesis of unidentified hexachloronaphthalene isomers and their retention characteristics. *Kankyo Kagaku*, 4, No. 3 (1994) 671-676; *C.A.*, 122 (1995) 105376t.
- 1382 Janák, K., Grimvall, E., Östman, C., Colmsjö, A., Athanasiadou, M. and Bergman, Å.: Gas chromatography-atomic emission detection (GC-AED) set up for bio-monitoring of PCBs and methylsulfonyl-PCBs. *J. Microcolumn Separ.*, 6 (1994) 605-616.
- 1383 Keymeulen, R., Voutetaki, A. and van Langenhove, H.: Determination of volatile chlorinated hydrocarbons in plant leaves by gas chromatography-mass spectrometry. *J. Chromatogr. A*, 699 (1995) 223-229.
- 1384 Larsen, B.R.: HRGC separation of PCB congeners. *J. High Resolut. Chromatogr.*, 18 (1995) 141-151.
- 1385 LeBlanc, A., Levesque, B. and Allaire, S.: Rapid, sensitive, and noninvasive sampling technique for determination of chloroform in alveolar breath. *J. Anal. Toxicol.*, 19 (1995) 56-57; *C.A.*, 122 (1995) 125435q.
- 1386 Ma, Y. and Guo, C.: (Gas-chromatographic determination of Freon 12.) *Huanjing Wuran Yu Fangzhi*, 16, No. 3 (1994) 43-44; *C.A.*, 122 (1995) 168511d.
- 1387 Natzeck, C., Luckas, B., Buyten, J. and Moskopp, G.: HRGC determination of toxic PCBs in marine organisms after SPE with Carboxpack B. *Organohalogen Compd.*, 11 (1993) 143-146; *C.A.*, 122 (1995) 4607p.
- 1388 Ogino, H. and Sakai, N.: Determination of trace impurities in high purity carbon tetrachloride using capillary gas chromatography-mass spectrometry. *Kuromatogurafi*, 15, No. 1 (1994) 11-15; *C.A.*, 122 (1995) 177398z.
- 1389 Picó, Y., Redondo, M.J., Font, G. and Manes, J.: Solid-phase extraction on C₁₈ in the trace determination of selected polychlorinated biphenyls in milk. *J. Chromatogr. A*, 693 (1995) 339-346.
- 1390 Pyle, S.M. and Gurka, D.F.: Volatile organic analysis by direct aqueous injection. *Talanta*, 41 (1994) 1845-1852.
- 1391 Raccanelli, S., Pavoni, B., Maroli, L. and Sfriso, A.: One step clean-up and separation of chlorinated, aliphatic and polycyclic aromatic hydrocarbons in environmental samples, prior to gas chromatographic quantification. *Toxicol. Environ. Chem.*, 45 (1994) 121-137; *C.A.*, 122 (1995) 87882y.
- 1392 Simmonds, P.G., O'Doherty, S., Nickless, G., Sturrock, G.A., Swaby, R., Knight, P., Ricketts, J., Woffending, G. and Smith, R.: Automated gas chromatograph/mass spectrometer for routine atmospheric field measurements of the CFC replacement compounds, the hydrofluorocarbons and hydrochlorofluorocarbons. *Anal. Chem.*, 67 (1995) 717-723.
- 1393 Takasuga, T., Ohi, E. and Inoue, T.: Complete isolation and determination of mono-ortho and non-ortho substituted PCBs and PCDDs/PCDFs by HPLC, on porous graphitic carbon, with HRGC/HRMS. *Organohalogen Compd.*, 11 (1993) 101-104; *C.A.*, 122 (1995) 204237e.
- 1394 Tanaka, N., Kimata, K., Hosoya, K., Araki, T., Patteraon, D.G., Barnhart, E.R. and Alexander, L.R.: (Sample preparation for GC/MS analysis of hydrophobic environmental contaminants by direct serum injection into restricted-access reversed-phase HPLC columns.) *Kankyo Kagaku*, 4 (1994) 484-485; *C.A.*, 122 (1995) 180385y.
- 1395 Vetter, W. and Luckas, B.: Calculation algorithm for the retention times of polychlorinated biphenyls and the control of its efficiency on a non-polar capillary column. *J. Chromatogr. A*, 699 (1995) 173-182.
- 1396 Wang, H.-w., Simmons, M.S. and Deiningor, R.A.: Application of dynamic headspace analysis for trihalomethanes in flowing water. *J. Chromatogr. Sci.*, 33 (1995) 109-115.

- 1397 Wester, P.G. and de Boer, J.: Determination of polychlorinated terphenyls in biota and sediments with gas chromatography/mass spectrometry. *Organohalogen Compd.*, 14 (1993) 121-124; *C.A.*, 122 (1995) 45380c.
- 1398 Yokouchi, Y., Akimoto, H., Barrie, L.A., Bottenheim, J.W., Anlauf, K. and Jobson, B.T.: Serial gas chromatographic/mass spectrometric measurements of some volatile organic compounds in the Arctic atmosphere during the 1992 Polar Sunrise Experiment. *J. Geophys. Res., [Atmos.]*, 99 (1994) 23,379-23,389; *C.A.*, 122 (1995) 166005t.
- 1399 Zhang, X.: (Determination of chlorobenzenes in water by headspace-GC.) *Kuangye Gongcheng*, 14, No.3 (1994) 52-55; *C.A.*, 122 (1995) 169493m.
- See also 1197, 1296, 1635, 1636, 1639, 1643, 1647.
- 5d. *Complex hydrocarbon mixtures (incl. analysis of tars, bitumens and mineral oils)*
- 1400 Alajbeg, A. and Kovac, T.: HR GC and GC-MS study of steroids in northern Croatian petroleum. *Croat. Chem. Acta*, 67 (1994) 393-405; *C.A.*, 122 (1995) 85067n.
- 1401 Barman, B.N.: Determination of hydrocarbon types in gasoline range samples by multidimensional gas chromatography, fluorescent indicator adsorption and bromine number methods. *Fuel*, 74 (1995) 401-406; *C.A.*, 122 (1995) 191909y.
- 1402 Chen, B., Qin, Y. and Li, M.: (Analysis of organic compounds in particulate phases of combustion products in domestic fuel gas.) *Weisheng Yanjiu*, 23 (1994) 286-289; *C.A.*, 122 (1995) 62948m.
- 1403 De Juan, G.D.: (Applications of gas chromatography in the refining industry. Part I. Determination of the types of hydrocarbons in middle petroleum fractions.) *Quim Ind.*, 41, No. 3 (1994) 25-32; *C.A.*, 122 (1995) 191947j.
- 1404 Dinkov, N. and Shopnikolov, E.: (Possibilities of gas chromatography for determination of mineral oils and greases for forensic practice purposes.) *Anal. Lab.*, 3 (1994) 199-202; *C.A.*, 122 (1995) 125469d.
- 1405 Hall, P.B., Stoddart, D., BJOROEY, M., Larter, L.S. and Brasher, J.E.: Detection of petroleum heterogeneity in Elfisk and satellite fields using thermal extraction, pyrolysis-GC, GC-MS and isotope techniques. *Org. Geochem.*, 22 (1994) 383-402; *C.A.*, 122 (1995) 164954c.
- 1406 Letoffe, J.M., Claudy, P., Garcin, M. and Volle, J.L.: Evaluation of crystallized fractions of crude oils by differential scanning calorimetry. Correlation with gas chromatography. *Fuel*, 74 (1995) 92-95; *C.A.*, 122 (1995) 59844u.
- 1407 Li, W., Malik, A., Lee, M.L., Jones, B.A., Porter, N.L. and Richter, B.E.: Group-type separation of diesel fuels using packed capillary column supercritical fluid chromatography. *Anal. Chem.*, 67 (1995) 647-654.
- 1408 Luo, Z., Hsia, Y.-p., Xie, K. and Zhang, Y.: Analysis of diesel components in soil and water contaminated by semivolatle synthetic organic compounds. *J. Chromatogr. Sci.*, 33 (1995) 263-267.
- 1409 Mitsui, T. and Fujimura, Y.: Determination of blend composition of heavy oil and kerosene by GC using principal component analysis. *Bunseki Kagaku*, 44 (1995) 383-389.
- 1410 Sepic, E., Leskovsek, H. and Trier, C.: Aerobic bacterial degradation of selected polyaromatic compounds and n-alkanes found in petroleum. *J. Chromatogr. A*, 697 (1995) 515-523.
- 1411 Tang, Y. and Stauffer, M.: Multiple cold trap pyrolysis gas chromatography: a new technique for modeling hydrocarbon generation. *Org. Geochem.*, 22 (1994) 863-872; *C.A.*, 122 (1995) 164966h.
- 1412 Tegelaar, E.W. and Noble, R.A.: Kinetics of hydrocarbon generation as a function of the molecular structure of kerogen as revealed by pyrolysis-gas chromatography. *Org. Geochem.*, 22 (1994) 543-574; *C.A.*, 122 (1995) 191833u.
- 1413 Xiang, Y., Morgan, S.L. and Watt, B.E.: TPH and BTEX quantitation in gasoline and diesel contaminated soils by capillary gas chromatography-mass spectrometry. *J. Chromatogr. Sci.*, 33 (1995) 98-108.
- 1414 Yang, Y., Hawthorne, S.B. and Miller, D.J.: Comparison of sorbent and solvent trapping after supercritical fluid extraction of volatile petroleum hydrocarbons from soil. *J. Chromatogr. A*, 699 (1995) 265-276.
- See also 1786, 1794.
6. ALCOHOLS
- 1415 Bureiko, A.S., Dorofeikov, V.V., Scherbak, I.G. and Pavlov, I.P.: (Gas chromatographic estimation of the esterase activity of proteolytic enzymes by means of continuous gas extraction of a volatile product.) *Vopr. Med. Khim.*, 40, No. 3 (1994) 57-58; *C.A.*, 122 (1995) 4100t.
- 1416 Dasgupta, A. and Macaulay, R.: Microwave-induced rapid synthesis of 4-carbethoxyhexafluorobutyl derivatives of fatty alcohols - a novel derivative for gas chromatography-chemical ionization mass spectrometric study. *J. Chromatogr. A*, 695 (1995) 136-141.
- 1417 Fomichev, A.E. and Klyuev, A.E.: (Method for ethylene glycol measurements in biological and technical fluids.) *Sud.-Med. Ekspert.*, 37, No. 3 (1994) 28-30; *C.A.*, 122 (1995) 98910g.
- 1418 Livase, J.F., Perkins, S.L., Tokessy, N.E. and Maddock, M.J.: Simultaneous determination of alcohols and ethylene glycol in serum by packed or capillary-column gas chromatography. *Clin. Chem. (Washington)*, 41 (1995) 300-305.
- 1419 Reinhardt, R., Engewald, W. and Görög, S.: Gas chromatographic separation of the enantiomers of flumecinol and some related compounds. *J. High. Resolut. Chromatogr.*, 18 (1995) 259-262.
- 1420 Tsukamoto, S., Kanegae, T., Matsumura, Y., Uchigasaki, S., Kitazawa, M., Muto, T., Chiba, S., Oshida, S., Nagoya, T. et al.: Simultaneous measurement of alcohols and hydrogen cyanide in biological specimens using headspace chromatography. *Nippon Hoigaku Zasshi*, 48 (1994) 336-342; *C.A.*, 122 (1995) 153476x.
- 1421 Yang, Y., Yao, G. and Li, H.: (The rapid determination of normal alcohols in fatty alcohols produced from hydroformylation unit by capillary gas chromatographic (GC) method.) *Chin. J. Chromatogr.*, 13 (1995) 126-127.
- 1422 Zhang, Q.: (Determination of methanol in perfume by headspace gas chromatography (GC).) *Chin. J. Chromatogr.*, 13 (1995) 144.

See also 1212, 1269, 1317, 1318, 1443, 1479.

7. PHENOLS

- 1423 Ahmed, N. and Hale, K.: A microassay for urinary phenol using capillary gas chromatography and optimized enzymic hydrolysis. *Clin. Chim. Acta*, 230 (1994) 201-208; *C.A.*, 121 (1994) 294507z.
- 1424 Bayonove, C. and Leroy, F.: (Detection of chlorophenols in wines and corks.) *Ind. Bevande*, 23 (1994) 231-242; *C.A.*, 122 (1995) 29965s.
- 1425 Besner, A., Gilbert, R., Tetreault, P., Lepine, L. and Archambault, J.-F.: Determination of pentachlorophenol and its hydrocarbon solvent in wood, soil, and water by gas chromatography and FT-IR spectroscopy in a single-sample treatment. *Anal. Chem.*, 67 (1995) 442-446.
- 1426 Janku, J., Kubes, V., Horsak, J. and Kuras, M.: Determination of phenol and phenol derivatives in polluted soils and sludges. *Fresenius Environ. Bull.*, 3 (1994) 475-480; *C.A.*, 122 (1995) 204244e.
- 1427 Pendergrass, S.M.: An alternative method for the analysis of phenol and *o*-,*m*- and *p*-cresol by capillary GC/FID. *Am. Ind. Hyg. Assoc. J.*, 55 (1994) 1051-1054; *C.A.*, 122 (1995) 62718m.
- 1428 Smith, T.J., Wearne, R.H. and Wallis, A.F.A.: Properties of chlorinated dihydroxybenzenes-components of pulp bleaching effluents. *Chemosphere*, 29 (1994) 1555-1560.
- 1429 Veningerova, M., Prachar, V., Uhnak, J. and Kovacicova, J.: Polychlorinated phenols in total diet. *Z. Lebensm.-Unters. Forsch.*, 199 (1994) 317-321; *C.A.*, 121 (1994) 299426g.

See also 1312, 1784.

8. SUBSTANCES CONTAINING HETEROCYCLIC OXYGEN

8a. Flavonoids

- 1430 Adlercreutz, H., Fotsis, T., Kurzer, M.S., Wähälä, K., Mäkelä, T. and Hase, T.: Isotope dilution gas chromatographic-mass spectrometric method for the determination of unconjugated lignans and isoflavonoids in human feces, with preliminary results in omnivorous and vegetarian women. *Anal. Biochem.*, 225 (1995) 101-105.

8b. Aflatoxins and other mycotoxins

- 1431 Haugen, J.-E., Oehme, M. and Mueller, M.D.: Enantiomer-specific analysis of homoanatoxin-a, a cyanophyte neurotoxin. *Spec. Publ. - R. Soc. Chem.*, 149 (1994) 40-44; *C.A.*, 122 (1995) 74058j.
- 1432 Jefferis, T.M., Brammer, G., Zotou, A., Brough, P.A. and Gallagher, T.: Determination of anotoxin-a, homoanotoxin and propylanotoxin in cyanobacterial extracts by HPLC, GC/mass spectrometry and capillary electrophoresis. *Spec. Publ. - R. Soc. Chem.*, 149 (1994) 34-39; *C.A.*, 122 (1995) 74057h.

8c. Other compounds with heterocyclic oxygen (incl. tannins)

- 1433 Bestmann, H.J., Haak, U., Kern, F. and Hölldobler, B.: 2,4-Dimethyl-5-hexanolide, a trail pheromone component of the carpenter ant *Camponotus herculeanus*. *Naturwissenschaften*, 82 (1995) 142-144.
- 1434 Galletti, G.C. and Bocchini, P.: Behavior of catechin and ellagic acid subjected to thermally-assisted hydrolysis-methylation/gas chromatography/mass spectrometry. *Rapid Commun. Mass Spectrom.*, 9 (1995) 250-254; *C.A.*, 122 (1995) 182710z.
- 1435 Kim, J.-Y., Chang, Y.-S. and Lee, D.-W.: (Sample pretreatment and HRGC/HRMS analysis for determination of dioxins in environmental samples.) *J. Korean Chem. Soc.*, 38 (1994) 819-826; *C.A.*, 122 (1995) 95634j.
- 1436 Klyuev, N.A., Golovkov, V.F., Chernov, S.A., Brodsky, E.S. and Shuisky, G.M.: Synthesis and properties of halo derivatives of dibenzo-*p*-dioxin containing a trifluoromethyl group. *Khim. Geterotsykl. Soedin.*, 7 (1994) 902-907; *C.A.*, 122 (1995) 187504v.
- 1437 Samsonov, D.P., Kiryukhin, V.P. and Zhiryukhina, N.P.: (Determination of polychlorinated dibenzo-*p*-dioxins and dibenzofurans in biological materials using gas chromatography-mass spectrometry in the negative ion-chemical ionization mode.) *Zh. Anal. Khim.*, 49 (1994) 868-873.
- 1438 Scilingo, A., Gigantiello, N., Guidugli, F. and Munari, F.: Analysis of polychlorinated dibenzo-*p*-dioxins (PCDDs) and polychlorinated dibenzofurans (PCDFs) by online LC-GC-MS. *Organohalogen Compd.*, 19 (1004) 55-59; *C.A.*, 122 (1995) 195394z.
- 1439 Soyfer, V.S., Murenets, N.V., Brodsky, E.S. and Klyuev, N.A.: A unified extraction and clean-up procedure for PCDD/PCDF determination in various matrixes by GC-MS. *Organohalogen Compd.*, 11 (1993) 79-81; *C.A.*, 122 (1995) 150585w.

See also 1203, 1393, 1394.

9. OXO COMPOUNDS, ETHERS, EPOXIDES AND QUINONES

- 1440 Buckee, G.K. and Mundy, A.P.: Determination of vicinal diketones in beer by gas chromatography (headspace technique)-collaborative trial. *J. Inst. Brew.*, 100 (1994) 247-253; *C.A.*, 122 (1995) 8304k.
- 1441 Dagaut, P., Reuillon, M., Cathonnet, M. and Presvots, D.: Gas chromatography and mass spectrometry identification of cyclic ethers formed from reference fuels combustion. *Chromatographia*, 40 (1995) 147-154.
- 1442 Ichinose, T., Miller, M.G. and Shibamoto, T.: Determination of free malonaldehyde formed in liver microsomes upon CCl₄ oxidation. *J. Appl. Toxicol.*, 14 (1994) 453-455; *C.A.*, 122 (1995) 25438z.
- 1443 Logan, B.K., Case, G.A. and Kiesel, E.L.: Differentiation of diethyl/acetone and ethano/acetone nitrile solvent pairs, and other common volatiles by dual column headspace gas chromatography. *J. Forensic Sci.*, 39 (1994) 1544-1551; *C.A.*, 122 (1995) 48625x.
- 1444 Miyake, T. and Shibamoto, T.: Quantitative analysis by gas chromatography of volatile carbonyl compounds in cigarette smoke. *J. Chromatogr. A*, 693 (1995) 376-381.

- 1445 Nishikawa, Y., Shimono, M., Kakagawa, K., Kato, T., Adachi, A. and Arimoto, H.: (Analysis for aldehydes in air using GC/MS; a comparison with the GC-FTD method.) *Kankyo Kagaku*, 4 (1994) 518-519; *C.A.*, 122 (1995) 195401z.
- 1446 Noguchi, K., Nagai, K., Fukita, K. and Maeda, H.: Determination of acetone and acetaldehyde in biological materials by gas chromatography combined with a pulse heating Curie-point hydrolyzer. *Hochudoku*, 12 (1994) 199-206; *C.A.*, 122 (1995) 74085r.
- 1447 Reinhardt, R., Steinborn, A., Engewald, W., Anhalt, K. and Schulze, K.: Enantiomer separation of α -campholene and fencholene derivatives by capillary gas chromatography on permethylated cyclodextrin phases. I. Compounds separable with single columns. *J. Chromatogr. A*, 697 (1995) 475-484.
- 1448 Rosell-Mel , A., Carter, J., Parry, A.T. and Eglinton, G.: Determination of the U_{K37} index in geological samples. *Anal. Chem.*, 67 (1995) 1283-1289.
- 1449 Sithole, B.B., Nyarku, S. and Allen, L.H.: Chromatographic methods for the determination of alkylketene dimer in pulp and paper matrices. *Analyst (Cambridge)*, 120 (1995) 1163-1169.
- 1450 Steinborn, A., Reinhardt, R., Engewald, W., Wyssuwa, K. and Schulze, K.: Enantiomer separation of α -campholene and fencholene derivatives by capillary gas chromatography on permethylated cyclodextrins. II. Compounds separable with coupled techniques. *J. Chromatogr. A*, 697 (1995) 485-494.
- 1451 Thomas, M.J., Robison, T.W., Samuel, M. and Forman, H.J.: Detection and identifying volatile aldehydes as dinitrophenylhydrazones using gas chromatography mass spectrometry. *Free Radical Biol. Med.*, 18 (1995) 553-557; *C.A.*, 122 (1995) 128036w.
- See also 1347, 1754.
10. CARBOHYDRATES
- 10a. *Mono and oligosaccharides. Structural studies*
- 1452 Deutsch, J.C., Kolli, V.R., Santosh-Kumar, C.R. and Kolhouse, J.F.: Serum xylose analysis by gas chromatography/mass spectrometry. *Am. J. Clin. Pathol.*, 102 (1994) 595-599; *C.A.*, 122 (1995) 50620k.
- 1453 Kim, Y.S., Park, K.S., Park, H.K. and Kim, S.W.: Compositional sugar analysis of antitumor polysaccharides by high-performance liquid chromatography and gas chromatography. *Arch. Pharmacol. Res.*, 17 (1994) 337-342; *C.A.*, 122 (1995) 23103n.
- 1454 Siccha, M.A., Lock De Ugaz, O. and Molina, S.M.: (Determination of galactomannan in gums of Tara, Charan and Cat's claw by gas chromatography.) *Bol. Soc. Quim. Peru*, 60 (1994) 39-43; *C.A.*, 122 (1995) 109208n.
- 1455 Zhu, X., Zeng, Z., Tang, X. and Zhuang, H.: (Analysis of monosaccharides in polyhedrons and granules of insect virus by gas chromatography.) *Bingdu Xuebao*, 10 (1994) 278-284; *C.A.*, 122 (1995) 100760d.
- 10b. *Polysaccharides, mucopolysaccharides, lipopolysaccharides*
- 1456 Chassagne, D., Crouzet, J., Baumes, R.L., Lepoutre, J.-P. and Bayonove, C.L.: Determination of trifluoroacetylated glycosides by gas chromatography coupled to methane negative chemical ionization mass spectrometry. *J. Chromatogr. A*, 694 (1995) 441-451.
- 1457 Chester, T.L. and Innis, D.P.: Determination of maltodextrin in psyllium-based bulk laxatives by *in situ* silylation and supercritical fluid chromatography. *Anal. Chem.*, 67 (1995) 1290-1292.
- 1458 Kleen, M. and Lindstroem, K.: Polysaccharides and lignins in newsprint process waters studied by pyrolysis-gas chromatography/mass spectrometry. *Nord. Pulp. Pap. Res. J.*, 9 (1994) 111-119; *C.A.*, 122 (1995) 190774p.
11. ORGANIC ACIDS AND LIPIDS
- 11a. *Organic acids and simple esters*
- 1459 Antoniosi Filho, N.R. and Lan as, F.M.: Identification of FAMES using ECL values and a three-dimensional Kov ts retention index system. *J. High Resolut. Chromatogr.*, 18 (1995) 167-170.
- 1460 Asakawa, F., Jitunari, F., Suna, S., Choi, J.O., Manabe, Y. and Takeda, N.: (Determination of 3-phenoxybenzoic acid in urine of workers exposed to permethrin using GC-MS). *Sangyo Igaku*, 36 (1994) 324-325; *C.A.*, 122 (1995) 125423j.
- 1461 De Jong, C., Palma, K. and Neeter, R.: Sample preparation before capillary gas-chromatographic estimation of free fatty acids in fermented dairy products. *Neth. Milk Dairy J.*, 48 (1994) 151-156; *C.A.*, 122 (1995) 158845b.
- 1462 Engan, T., Bjerve, K.S., Hoe, A.L. and Krane, J.: Characterization of plasma lipids in patients with malignant disease by ^{13}C nuclear magnetic resonance spectroscopy and gas liquid chromatography. *Blood*, 85 (1995) 1323-1330; *C.A.*, 122 (1995) 182659q.
- 1463 Fox, A. and Rosario, R.M.T.: Quantification of muramic acid, a marker for bacterial peptidoglycan, in dust collected from hospital and home air-conditioning filters using gas chromatography-mass spectrometry. *Indoor Air*, 4 (1994) 239-247; *C.A.*, 122 (1995) 195430h.
- 1464 Franotovic, M. and Sipus, D.: Rapid identification of *M. tuberculosis* by analysing tuberculostearic acid with gas chromatography: Determination of the method's sensitivity. *Period. Biol.*, 96 (1994) 27-30; *C.A.*, 121 (1994) 296284y.
- 1465 Goosens, E.C., Beerthuis, I.M., de Jong, D., de Jong, G.J. and Brinkman, U.A.T.: Reversed-phase liquid chromatography coupled on-line with capillary gas chromatography. Use of an anion-exchange membrane to remove an ion-pair reagent from the eluent. *Chromatographia*, 40 (1995) 267-271.
- 1466 Huifen, G., Guanliang, C., Liming, Z. and Yuzheng, Z.: (Determination of benzylic acid in water by gas chromatography-mass spectrometry.) *Fenxi Huaxue*, 22 (1994) 826-828; *C.A.*, 122 (1995) 88867j.
- 1467 Jan, S., Roblot, C., Goethals, G., Courtois, J., Courtois, B., Nava Suacedo, J.E., Sequin, J.-P. and Barbotin, J.-N.: Study of parameters affecting poly(3-hydroxybutyrate) quantification by gas chromatography. *Anal. Biochem.*, 225 (1995) 258-263.

- 1468 Karl, V., Gutser, J., Dietrich, A., Maas, B. and Mosandl, A.: Stereoisomeric flavor compounds. LXVIII. 2-,3- And 4-alkyl-branched acids, part 2: chiro-specific analysis and sensory evaluation. *Chirality*, 6 (1994) 427-434; *C.A.*, 122 (1995) 54281y.
- 1469 Katayama-Hirayama, K., Tobita, S. and Hirayama, K.: (Analysis of the methoxime-trimethylsilyl derivative of β -keto adipic acid by the GC/MS method.) *Mizu Shori Gijutsu*, 35 (1994) 401-404; *C.A.*, 122 (1995) 122244r.
- 1470 Kaunzinger, A., Thomsen, M., Dietrich, A. and Mosandl, A.: Stereodifferentiation and simultaneous analysis of 2- and 3-hydroxyalkanoic acids from biomembranes by multidimensional gas chromatography. *J. High Resolut. Chromatogr.*, 18 (1995) 191-194.
- 1471 Li, Y., Zhang, L., Liu, G., Hou, Y., Chen, L. and Zeng, P.: (Gas chromatographic analysis of C_1 - C_6 fatty acids in rat fecal samples.) *Huaxi Yike Daxue Xuebao*, 25 (1994) 456-459; *C.A.*, 122 (1995) 100768n.
- 1472 Li, Z. and Li, Q.: (Determination of fatty acids in red scalefish meat by gas chromatography/mass spectrometry method.) *Chin. J. Chromatogr.*, 13 (1995) 113-114.
- 1473 Liu, K.-S.: Preparation of fatty acid methyl esters for gas-chromatographic analysis of lipids in biological materials. *J. Am. Oil Chem. Soc.*, 71, No. 11 (1994) 1179-1187; *C.A.*, 122 (1995) 4527y.
- 1474 Manni, G. and Caron, F.: Calibration and determination of volatile fatty acids in waste leachates by gas chromatography. *J. Chromatogr. A*, 690 (1995) 237-242.
- 1475 Michaelis, W., Richnow, H.H. and Seifert, R.: Chemically bound chlorinated aromatics in humic substances. *Naturwissenschaften*, 82 (1995) 139-142.
- 1476 Mo, G.Q., Zhang, X.M., Cheng, W.B., Li, D. and Yang, X.Y.: (Determination of β -phenyllactic acid in rat plasma by derivatization capillary gas chromatography (DCGC).) *Yaouxue Xuebao*, 30 (1995) 70-73; *C.A.*, 122 (1995) 177592h.
- 1477 Murase, M., Kimura, Y. and Nagata, Y.: Determination of portal short-chain fatty acids in rats fed various dietary fibers by capillary gas chromatography. *J. Chromatogr. B*, 664 (1995) 415-420.
- 1478 Nikkari, T., Malo-Ranta, U., Hiltunen, T., Jaakkola, O. and Yla-Herttua, S.: Monitoring of lipoprotein oxidation by gas chromatographic analysis of hydroxy fatty acids. *J. Lipid Res.*, 36 (1995) 200-207; *C.A.*, 122 (1995) 100765j.
- 1479 Perez, J.J., Fauville-Duffaux, M., Dossogne, J.L., de Hoffmann, E. and Pouthier, F.: Faster identification of mycobacteria using gas liquid and thin layer chromatography. *Eur. J. Clin. Microbiol. Infect. Dis.*, 13 (1994) 717-725; *C.A.*, 122 (1995) 4624s.
- 1480 Park, P.W. and Goins, R.E.: *In situ* preparation of fatty acid methyl esters for analysis of fatty acid composition in foods. *J. Food Sci.*, 59 (1994) 1262-1266; *C.A.*, 122 (1995) 131279h.
- 1481 Pena, A. and Sandra, P.: Chemotaxometric characterization of yeast cell. *J. Chromatogr. Sci.*, 33 (1995) 116-122.
- 1482 Procida, G., Gabrielli-Favretto, L., Pertoldi-Marletta, G. and Ceccon, L.: Modified esterification procedure employing boron trifluoride/methanol complex for the determination of fatty acids and oils. *Riv. Ital. Sostanze Grasse*, 71 (1994) 547-553; *C.A.*, 122 (1995) 185711m.
- 1483 Robles, V.E., Gonzales, A.M., Sainz, M.M., Ramirez, G.P. and Martinez, R.B.: Determination of cellular fatty acids of *Klebsiella ozaenae* by gas chromatography. *Rev. Roum. Biochim.*, 31 (1994) 51-55; *C.A.*, 121 (1994) 296274v.
- 1484 Sakai, T., Araki, T., Morita, Y. and Masuyama, Y.: Gas chromatographic determination of butoxyacetic acid after hydrolysis of conjugated metabolites in urine from workers exposed to 2-butoxyethanol. *Int. Arch. Occup. Environ. Health*, 66 (1994) 249-254; *C.A.*, 122 (1995) 88110g.
- 1485 Santosh-Kumar, C.R., Deutsch, J.C., Hassel, K.L., Kolhouse, N.M. and Kolhouse, J.F.: Quantitation of red blood cell folates by stable isotope dilution gas chromatography-mass spectrometry utilizing a folate internal standard. *Anal. Biochem.*, 225 (1995) 1-7.
- 1486 Schuetz, B., Hack, B., Hanstein, S., Zimmermann, K. and Rusch, V.: (Quality control of starter cultures by gas chromatographic analysis of cell wall fatty acids.) *Lebensmittelind. Milchwirtsch.*, 114 (1993) 432-434; *C.A.*, 122 (1995) 131254w.
- 1487 Schwaiger, I. and Vojir, F.: (Application of multivariate statistical methods for the authenticity control of the edible oils.) *Dtsch. Lebensm.-Rundsch.*, 90, No. 5 (1994) 143-146; *C.A.*, 122 (1995) 131497c.
- 1488 Shao, X. and Liu, Y.: (Gas chromatographic analysis of sesame oil and sauce adulterated with peanut oil and sauce.) *Zhongguo Tiaoweipin*, 9 (1994) 29-30; *C.A.*, 122 (1995) 185690d.
- 1489 Sturaro, A., Parvoli, G., Doretti, L., Bancomina, C. and Neonato, C.: Gas chromatographic/mass spectrometric determination of vinyl acetate levels in air. *Org. Mass Spectrom.*, 29 (1994) 575-577; *C.A.*, 121 (1994) 307087q.
- 1490 Wahl, H.G., Chrzanowski, A., Müller, C., Liebich, H.M. and Hoffmann, A.: Identification of furan fatty acids in human blood cells and plasma by multi-dimensional gas chromatography-mass spectrometry. *J. Chromatogr. A*, 697 (1995) 453-459.
- 1491 Wheelan, P., Zirrolli, J.A. and Murphy, R.C.: Analysis of hydroxy fatty acids as pentafluorobenzyl ester, trimethylsilyl ether derivatives by electron ionization gas chromatography/mass spectrometry. *J. Am. Soc. Mass Spectrom.*, 6 (1995) 40-51; *C.A.*, 122 (1995) 187229j.
- 1492 Xiong, D., Zhu, X. and Sheng, Z.: (Analysis of metabolic products of anaerobes with gas-liquid chromatography.) *Zhonghua Yixue Jianshan Zazhi*, 17 (1994) 172-175; *C.A.*, 121 (1994) 296286a.
- 1493 Zhancai, Z.: (Gas chromatographic determination of trace propionic acid in glutinous broomcorn millet, non-glutinous broomcorn millet and grain sorghum.) *Fenxi Huaxue*, 22, No. 7 (1994) 753; *C.A.*, 122 (1995) 54300d.
- 1494 Zhang, L.-Y. and Hamberg, M.: A gas-liquid chromatographic method for steric analysis of 2-hydroxy, 3-hydroxy, and 2,3-dihydroxy acids. *Chem. Phys. Lipids*, 74 (1994) 151-161; *C.A.*, 122 (1995) 100734y.
- 1495 Zhou, X. and Zhang, D.: Structural investigation of fulvic acid from weathered coal by using capillary GC-MS. *Fuel Sci. Technol. Int.*, 12 (1994) 1169-1182; *C.A.*, 122 (1995) 13448q.
- 1496 Zielinska-Stasiek, M. and Cisowski, W.: Gas-liquid chromatographic analysis of fatty acids obtained from the seeds of common trees. *Acta Chromatogr.*, 3 (1994) 37-47; *C.A.*, 122 (1995) 155378r.

See also 1289, 1448, 1449, 1554, 1613.

11b. *Prostaglandins*

- 1497 Schweer, H., Cawello, W. and Seyberth, H.W.: Gas chromatography/negative ion chemical ionization triple quadrupole mass spectrometry determination and pharmacokinetics of 11 α -hydroxy-9,15-dioxo-2,3,4,5,20-pentano-19-carboxyprostanic acid in plasma. *Prostaglandins, Leukotrienes Essent. Fatty Acids* 51 (1994) 401-405; *C.A.*, 122 (1995) 152104n.

See also 1611.

11c. *Lipids and their constituents*

- 1498 Bartle, K.D. and Clifford, A.A.: Supercritical fluid extraction and chromatography of lipid materials. *Spec. Publ. - R. Soc. Chem.*, 160 (1994) 1-41; *C.A.*, 122 (1995) 50308q - a review with 112 refs.
- 1499 Bereuter, T.L. and Lorbeer, E.: Monitoring of lipase-catalyzed cleavage of acylglycerols by high-temperature gas chromatography. *J. Chromatogr. A*, 697 (1995) 469-474.
- 1500 Blomberg, L.G. and Demibuker, M.: Analysis of triacylglycerols by argentation supercritical fluid chromatography. *Spec. Publ. - R. Soc. Chem.*, 160 (1994) 42-58; *C.A.*, 122 (1995) 50418a.
- 1501 Ellerbe, P., Sniegowski, L.T. and Welch, M.J.: Isotope dilution mass spectrometry as a candidate definitive method for determining total glycerides and triglycerides in serum. *Clin Chem. (Washington)*, 41 (1995) 397-404.
- 1502 Gulumian, M., Ambaram, K. and Verdoon, G.H.: Identification and quantification of fiber-induced lipid peroxidation products by GC-MSD. *NATO ASI Ser., Ser. H*, 85 (1994) 417-424; *C.A.*, 121 (1994) 294504w.
- 1503 Nakajima, N. and Yamamoto, T.: (Isolation of triglycerides from plant oil by supercritical chromatography.) *Jpn. Kokai Tokkyo Koho JP 06,293,895 [94,293,895]* (Cl. C11B3/10), 21 Oct. 1994, Appl. 91/106,410, 12 Apr. 1991; 4 pp.; *C.A.*, 122 (1995) 104556q.
- 1504 Plank, C. and Lorbeer, E.: Simultaneous determination of glycerol, and mono-, di- and triglycerides in vegetable oil methyl esters by capillary gas chromatography. *J. Chromatogr. A*, 697 (1995) 461-468.
- 1505 Precht, D. and Frede, E.: Determination of the solid fat content in milk fat by gas chromatographic triglyceride analysis. *Fat Sci. Technol.*, 96 (1994) 324-330; *C.A.*, 121 (1994) 299423d.
- 1506 Tetens, V., Kristensen, N.B. and Calder, A.G.: Measurement of ¹³C enrichment of plasma lactate by gas chromatography/isotope ratio mass spectrometry. *Anal. Chem.*, 67 (1995) 858-862.
- 1507 Zoellner, P., Lorbeer, E. and Remberg, G.: Utility of nicotinoyl derivatives in structural studies on mono- and diacylglycerols by gas chromatography/mass spectrometry. Part. 1. *Org. Mass Spectrom.*, 29 (1994) 253-259; *C.A.*, 122 (1995) 106267v.

See also 1334, 1416.

13. STEROIDS

13b. *Pregnane and androstane derivatives*

- 1508 Holder, G.: External quality assessment of urinary-free cortisol measurements in the UK against a gas chromatography mass spectroscopy reference method. *Ann. Clin. Biochem.*, 32, No. 1 (1995) 84-90; *C.A.*, 122 (1995) 178552g.
- 1509 Legrand, C., Dousset, B., Tronel, H., Belleville, F. and Nabet, P.: Measurement of plasma testosterone by gas chromatography-negative-ion mass spectrometry using pentafluoropropionic derivatives. *J. Chromatogr. B*, 663 (1995) 187-192.
- 1510 Mgrd, M., Berg, H.E.B., Tagesson, V., Jaeremo, M.L.G., Karlsson, L.L.H., Mathiasson, L.J.E., Bonneau, M. and Hansen-Møller, J.: Determination of androstenone in pig fat using supercritical fluid extraction and gas chromatography-mass spectrometry. *J. Agric. Food Chem.*, 43 (1995) 114-120.
- 1511 Reisch, J. and Cramer, M.: (The detection of biogenic steroids in sheep thymus tissue by GC-MS.) *Sci. Pharm.*, 61 (1993) 185-191; *C.A.*, 121 (1994) 292927n.
- 1512 Steinhart, H. and Hartwig, M.: (Determination of naturally-occurring steroids (androgens and progestogens) in beef.) *GIT Fachz. Lab.*, 38 (1994) 893-898; *C.A.*, 122 (1995) 185639u.
- 1513 Vermeulen, G.J., Lambert, J.G.D., van der Looy, M.J.W. and Goos, H.J.T.: Gas chromatographic-mass spectrometric (GC-MS) analysis of gonadal steroids in plasma of the male African catfish, *Clarias gariepinus*: effects of castration or treatment with gonadotropin releasing hormone analog. *Gen. Comp. Endocrinol.*, 96 (1994) 288-297; *C.A.*, 122 (1995) 6005q.
- 1514 Vladimirova, V. and Tsoutsoulova-Draganova, A.: Gas chromatographic-mass spectrometric investigation of fluoxymestron in urine samples. *Anal. Lab.*, 2, No. 3 (1993) 184-189; *C.A.*, 122 (1995) 23963t.
- 1515 Ye, L., Zhang, C. and Zhang, Y.: (Gas chromatography-mass spectrometric analysis of metabolites of nandronolone and norethiosterone in human urine.) *Fenxi Huaxue* 23 (1995) 32-35; *C.A.*, 122 (1995) 96621q.

13c. *Estrogens*

- 1516 Moretti, G., Citti, G., Cammarata, P., Ferretti, E., Fiori, M. and di Giovannandrea, R.: (Residue analysis of urinary trenbolone in cattle by thin-layer chromatography and gas chromatography-mass spectrometry.) *Riv. Sci. Aliment.*, 23 (1994) 227-232; *C.A.*, 122 (1995) 29974u.

13d. *Sterols*

- 1517 Benfenati, E., Cools, E., Fattore, E. and Fanelli, R.: A GC-MS method for the analysis of fecal and plant sterols in sediment samples. *Chemosphere*, 29 (1994) 1393-1405.
- 1518 Grob, K. and Mariani, C.: LC-GC methods for the determination of adulterated edible oils and fats. *Spec. Publ. - R. Soc. Chem.*, 160 (1994) 73-81; *C.A.*, 122 (1995) 131220g - a review with 13 refs.
- 1519 Kisza, J., Staniewski, B. and Januskiewicz, M.: Determination of cholesterol in milk fat by gas chromatography (GC). *Pol. J. Food Nutr. Sci.*, 3 (1994) 75-81; *C.A.*, 122 (1995) 104162h.

- 1520 Klatt, L.V., Mitchell, B.A. and Smith, R.L.: Cholesterol analysis in foods by direct saponification-gas chromatographic method: Collaborative study. *J. Assoc. Off. Anal. Chem. Int.*, 78 (1995) 75-79.
- 1521 Toschi, T.G., Caboni, M.F., Capella, P. and Lercker, G.: High performance liquid chromatography and capillary gas chromatography in the analysis of oxysterols. *Riv. Ital. Sostanze Grasse*, 71 (1994) 329-333; *C.A.*, 122 (1995) 54328u.
- See also 1481.
15. TERPENES AND OTHER VOLATILE AROMATIC COMPOUNDS
- 15a. Terpenes
- 1522 Bosset, J.O., Buetikofer, U., Gauch, R and Sieber, R.: (Occurrence of terpenes and aliphatic hydrocarbons in Swiss Gruyere and Etivaz alpine cheese using dynamic headspace GC-MS analysis of their volatile flavor compounds.) *Schweiz. Milchwirtschaft. Forsch.*, 23, No. 2 (1994) 37-41; *C.A.*, 122 (1995) 8443e.
- 1523 Dietrich, A., Maas, B. and Mosandl, A.: Diluted modified cyclodextrins as chiral stationary phases - Influence of the polysiloxane solvent: heptakis(2,3-di-O-acetyl-6-O-tert.-butyldimethylsilyl)- β -cyclodextrin. *J. High Resolut. Chromatogr.*, 18 (1995) 152-156.
- 1524 Fan, H., Xu, K., Yang, F., Wang, S. and Wu, L.: Determination of menthol and borneol in Houyanqing pill by capillary gas chromatography. *Zhongguo Yaoxue Zazhi*, 29, No. 8 (1994) 51-52; *C.A.*, 121 (1994) 308490c.
- 1525 Grob, K., Biedermann, M., Bronz, M. and Schmid, J.P.: Recognition of mild deodorization of edible oils by the loss of volatile components. *Z. Lebensm.-Unters. Forsch.*, 199 (1994) 191-194; *C.A.*, 122 (1995) 8294g.
- 1526 Kawai, N., Tsubokura, H. and Nishikawa, Y.: Structural analysis of phytol group of protochlorophylls by gas chromatography. *Kinki Daigaku Rikogakubu Kenkyu Hokoku*, 30 (1994) 25-29; *C.A.*, 122 (1995) 80977z.
- 1527 Kimball, B.A., Craver, R.K., Johnston, J.J. and Nolte, D.L.: Quantitative analysis of the mono- and sesquiterpenoids of Douglas-fir sapwood by solvent extraction and gas chromatography with mass selective detection. *J. High. Resolut. Chromatogr.*, 18 (1995) 221-225.
- 1528 Li, X., Shen, X. and Yao, T.: (Rapid assay of borneol in Zhenzhumingmu eye drops.) *Zhejiang Yike Daxue Xuebao*, 23, No. 1 (1994) 43-44; *C.A.*, 122 (1995) 38979b.
- 1529 Patitucci, M.L., Pinto, A.C. and Cordoso, J.N.: Analysis of crude extracts and fractions of Brazilian Polypodiaceae by high-resolution gas chromatography-mass spectrometry. I. Triterpenes. *Phytochem. Anal.*, 6 (1995) 38-44; *C.A.*, 122 (1995) 155672g.
- 1530 Ravid, U., Putievsky, E. and Katzir, I.: Chiral GC analysis of enantiomerically pure (R)(-)-linalyl acetate in some Lamiaceae, Myrtle and Petitgrain essential oils. *Flavour Fragrance J.*, 9 (1994) 275-276; *C.A.*, 122 (1995) 63966c.
- 1531 Rossini, C., Pandolfi, E., Dellacassa, E. and Moyna, P.: Determination of 1,8-cineole in *Eucalyptus* essential oils by thin-layer chromatography and densitometry. *J. Assoc. Off. Anal. Chem. Int.*, 78 (1995) 115-117.
- 1532 Vilegas, J.H.Y., Lanças, F.M. and Antoniosi Filho, N.R.: High temperature capillary GC analysis of phytopreparations of "Espinheira Santa" (*Maytenus ilicifolia* M. and *Maytenus aquifolium* M.-Celastraceae), a Brazilian antiulcer plant. *Chromatographia*, 40 (1995) 341-344.
- See also 1344, 1447, 1450, 1719.
- 15b. Essential oils
- 1533 Alfonso, D. and Kapetanidis, I.: (Volatile constituents of *Isochroma gesnerioides*: qualitative analysis by GC-MS.) *Sci. Pharm.*, 62 (1994) 59-65; *C.A.*, 122 (1995) 156263m.
- 1534 Blanco Tirado, C., Stashenko, E.E., Combariza, M.Y. and Martinez, J.R.: Comparative study of Colombian citrus oils by high-resolution gas chromatography and gas chromatography-mass spectrometry. *J. Chromatogr. A*, 697 (1995) 501-513.
- 1535 Dung, N.X., Luu, H.V., Khoi, T.T. and Leclercq, P.A.: GC and GC/MS analysis of the leaf oil of *Cleistocalyx operculatus* Roxb. Merr. e Perry (Syn. *Eugenia operculata* Roxb.; *Syzygium mervosum* DC.) *J. Essent. Oil Res.*, 6, No. 6 (1994) 661-662; *C.A.*, 122 (1995) 101642d.
- 1536 Garcia, V.M.C., Nascimento, E.A. and Morais, S.A.L.: Volatile wood oils of the Brazilian *Pinus caribaea* var. *Hondurensis* and Spanish *Pinus pinaster* var. *mediterranea*. *J. Braz. Chem. Soc.*, 5 (1994) 107-112; *C.A.*, 122 (1995) 63967d.
- 1537 Hener, U., Faulhaber, S., Kreis, P. and Mosandl, A.: Evaluation of authenticity of balm oil (*Melissa officinalis* L.) *Pharmazie*, 50 (1995) 60-62; *C.A.*, 122 (1995) 196656y.
- 1538 Hethelyi, I., Ceseko, I., Grosz, M., Mark, G. and Palinkas, J.: (Capillary gas chromatographic investigation of *Artemisia annua* essential oils.) *Olaj, Szappan, Kozmet.*, 43 (1994) 103-106; *C.A.*, 122 (1995) 89057g.
- 1539 Jirovetz, L., Buchbauer, G., Remberg, G. and Nikiforov, A.: (Analysis of grapevine secretion (grapevine tears) using GC-FTIR-MS.) *Vitis*, 33, No. 2 (1994) 97; *C.A.*, 122 (1995) 76513c.
- 1540 Kreis, P. and Mosandl, A.: Chiral compounds of essential oils. Part XVI. Enantioselective multidimensional gas chromatography in authenticity control of balm oil (*Melissa officinalis* L.) *Flavour Fragrance J.*, 9 (1994) 249-256; *C.A.*, 122 (1995) 38982x.
- 1541 Kreis, P. and Mosandl, A.: Chiral compounds of essential oils. Part XVII. Simultaneous stereoanalysis of Cymbopogon oil constituents. *Flavour Fragrance J.*, 9 (1994) 257-260; *C.A.*, 122 (1995) 63957a.
- 1542 Ma, X.-y., Pan, H.-p. and Chen, Y.-z.: Off-line two-dimensional gas chromatography-mass spectrometry for analysis of essential oil of medicinal herb. *Chem. Res. Chin Univ.*, 10 (1994) 291-296; *C.A.*, 122 (1995) 182310u.
- 1543 Orav, A., Kuningas, K. and Kailas, T.: Computerized capillary gas chromatographic identification and determination of Siberian fir oil constituents. *J. Chromatogr. A*, 697 (1995) 495-499.
- 1544 Vernin, G., Ghiglione, C. and Parkanyi, C.: GC-MS-SPEMA bank analysis of *Thymus serpyllum praecox* (Opiz) Wollm (wild thyme) from Hautes Alpes (France). *Dev. Food Sci.*, 34 (1994) 50-515; *C.A.*, 122 (1995) 8390k.

- 1545 Wang, J., Yang, C., Da, W. and Zhang, C.: (GC-MS analysis of essential oil from pericaps of *Illicium difengpi*.) *Zhongguo Zhongyao Zazhi*, 19 (1994) 422-423; C.A., 122 (1995) 76504a.
- 1546 Wang, Y., Zhang, H. and Zhang, J.: (GC-MS analysis of the volatile oil in the stem of *Actinidia kolomikta* Planch.) *Baiqiuwen Yike Daxue Xuebao*, 20 (1994) 358-359; C.A., 122 (1995) 156308e.
- 1547 Yan, J.-C., Zhang, H., Xu, S.-M., Wei, Y.-D. and Yan, X.: (Studies on the volatile oil composition of *Rhodiola* using GC/MS/DS and GC/IS/DS.) *Gaodeng Xuexiao Huaxue Xuebao*, 15 (1994) 1175-1177; C.A., 122 (1995) 17300h.
- 1548 Yang, S. and Lin, D.: (GC-MS study on the essential oil of citrus peel. III. Analysis of the essential oil from *Meixian shaddock* peel.) *Fenxi Ceshi Xuebao*, 13, No. 6 (1994) 56-59; C.A., 122 (1995) 185664y.
- 1549 Ye, C., Wu, X., Rong, Y., Liu, W., Zhu, R. and Mao, X.: (The fingerprint spectrum analysis of capillary gas chromatographic retention indices for essential oil of the genus *Atractylodes*.) *Yaowu Fenxi Zazhi*, 14, No. 6 (1994) 13-16; C.A., 122 (1995) 115092m.

See also 1530.

16. NITRO AND NITROSO COMPOUNDS

- 1550 Akiyama, K. and Sakamoto, K.: (Analytical method of nitro-PAHs in automobile exhaust by GC-SID.) *Kankyo Kagaku*, 4 (1994) 647-654; C.A., 122 (1995) 87870t.
- 1551 Frassanito, R., Benfenati, E., Ciotti, G. and Fanelli, R.: A simple method for determination of N-nitrosamine traces in trifluralin samples by gas chromatography-mass spectrometry. *Toxicol. Environ. Chem.*, 45 (1994) 199-204; C.A., 122 (1995) 125831r.
- 1552 Ma, M., Xu, X., Kang, Z. and Bian, Y.: (Detection of trace level nitropolycyclic aromatic hydrocarbons in urban surface water for the first time.) *Huanjing Huaxue*, 13 (1994) 543-549; C.A., 122 (1995) 196362z.
- 1553 Mussmann, P., Preiss, A., Levsen, K. and Wuensch, G.: Development of methods for the analysis of nitrotoluenes, nitramines and other organic compounds in wastewater of former ammunition manufacturing sites. *Vom Wasser*, 82 (1994) 79-90; C.A., 122 (1995) 114386y.
- 1554 Soják, L., Perjéssy, A., Kubinec, R., Giumanini, A.G. and Strazolini, P.: Identification of the isomers from mono- and dinitration of α -hydroxydiphenylacetic acid by capillary gas chromatography with Fourier transform infrared and mass spectrometric detection. *J. Chromatogr. A*, 695 (1995) 57-64.
- 1555 You, J., Li, H. and Sun, X.: (The determination of nitrated products of toluene by gas chromatography (GC).) *Chin. J. Chromatogr.*, 13 (1995) 128-129.

17. AMINES, AMIDES AND RELATED NITROGEN COMPOUNDS

17a. Amines and polyamines

- 1556 Ghaoui, L., Pell, K., Chritz, K. and Loomis, R.N.: Analysis of amine compounds by capillary gas chromatography using ammonia-helium carrier gas. *J. High Resolut. Chromatogr.*, 18 (1995) 157-160.
- 1557 Lin, D., Lay, J.O.Jr., Bryant, M.S., Malaveille, C., Friesen, M., Bartsch, H., Lang, N.P. and Kadlubar, F.F.: Analysis of 4-amino-biphenyl-DNA adducts in human urinary bladder and lung by alkaline hydrolysis and negative ion gas chromatography-mass spectrometry. *Environ. Health Perspect. Suppl.*, 102 (1994) 11-16; C.A., 122 (1995) 98874y.
- 1558 Wilson, P.M., Que Hee, S.S. and Froines, J.R.: Determination of hemoglobin adduct levels of the carcinogen 2,4-diaminotoluene using gas chromatography-electron impact positive-ion mass spectrometry. *J. Chromatogr. B*, 667 (1995) 166-172.

See also 1312, 1570.

17c. Urea and guanidine derivatives

See 1562.

17d. Other amine derivatives and amides (excl. peptides)

- 1559 Avtushenko, Yo.Ye., Sulman, E.M., Ankudinova, T.V. and Sidorov, A.I.: (GLC analysis of reaction mixtures in the reductive condensation of 3,4-xylidine with D-ribose.) *Khim.-Farm. Zh.*, 28, No.5 (1994) 59-60; C.A., 122 (1995) 187958w.
- 1560 Dong, X.-D., Lu, J. and Cha, C.: Determination of 4,4'-methylenedianiline (MDA) and identification of isomers in technical-grade MDA in hydrolyzed plasma and urine from workers exposed to methylenediphenyldiisocyanate by gas chromatography-mass spectrometry. *J. Electroanal. Chem.*, 381 (1995) 195-201; C.A., 122 (1995) 155193b.
- 1561 Kataoka, H., Shindoh, S. and Makita, M.: Determination of secondary amines in various foods by gas chromatography with flame photometric detection. *J. Chromatogr. A*, 695 (1995) 142-148.
- 1562 Zell, C.H., Neidlein, R., Strein, K., Bartsch, W. and Michel, H.: Pharmacokinetics of the organic nitrates trans-N-(4-nitrocyclohexyl)-urea in dogs and trans-(4-nitrocyclohexyl)-acetamide in dogs and man. *Arzneim.-Forsch.*, 44 (1994) 1186-1190; C.A., 122 (1995) 95801m.

See also 1322, 1766.

18. AMINO ACIDS AND PEPTIDES; CHEMICAL STRUCTURE OF PROTEINS

18a. Amino acids and their derivatives

- 1563 Adachi, J., Mio, T., Ueno, Y., Nishimura, A., Sumino, K. and Tatsuno, Y.: (Gas chromatography-mass spectrometry analysis for 3-(phenylamino) alanine and its metabolites.) *Nippon Iyo Masu Supekutoru Gakkai Koenshu*, 19 (1994) 103-106; C.A., 122 (1995) 71258p.

- 1564 Bartels, M.J.: Quantitation of the tetrachloroethylene metabolite N-acetyl-S-(trichlorovinyl)cysteine in rat urine via negative ion chemical ionization gas chromatography/tandem mass spectrometry. *Biol. Mass Spectrom.*, 23 (1994) 689-694; *C.A.*, 121 (1994) 294513y.
- 1565 Kataoka, H., Takagi, K. and Makita, M.: Determination of total plasma homocysteine and related aminothiols by gas chromatography with flame photometric detection. *J. Chromatogr. B*, 664 (1995) 421-425.
- 1566 Li, D., Yin, Q. and Yao, Y.: (Determination of amino acids in radish.) *Yanbian Daxue Xuebao, Ziran Kexueban*, 20, No. 3 (1994) 47-49; *C.A.*, 122 (1995) 159138k.
- 1567 Oh, C.-H., Brownson, D.M. and Mabry, T.J.: Screening for non-protein amino acids in seeds of the Guam cycad, *Cycas circinalis*, by an improved GC-MS method. *Planta Med.*, 61 (1995) 66-70; *C.A.*, 122 (1995) 182671n.
- 1568 Peaper, D.R. and Jones, H.L.: Determination of S-methylmethionine in malt by alkaline hydrolysis and headspace gas chromatography - a collaborative trial. *J. Inst. Brew.*, 100 (1994) 397-398; *C.A.*, 122 (1995) 158849f.
- 1569 Williams, B.D. and Wolfe, R.R.: Determination of amino- and amide-¹⁵N glutamine enrichment with tertiary butyldimethylsilyl derivatives. *Biol. Mass Spectrom.*, 23, No. 11 (1994) 682-688; *C.A.*, 122 (1995) 27179q.
- 1570 Yasuhara, F., Yamaguchi, S., Takeda, M., Ochiai, Y. and Miyano, S.: Gas chromatographic separation of enantiomeric amine acids and amines with α -methoxy- α -trifluoromethylpropionic acid as a chiral derivatizing agent. *J. Chromatogr. A*, 694 (1995) 227-236.
- See also 1314, 1315, 1323.
- 18b. *Peptides, peptidic and proteinous hormones, growth factors*
- 1571 Kataoka, H., Takagi, K. and Makita, M.: Determination of glutamine and related aminothiols by gas chromatography with flame photometric detection. *Biomed. Chromatogr.*, 9 (1995) 85-89.
20. ENZYMES AND ENZYME ACTIVITY ESTIMATION
- 20a. *Oxidoreductases*
- See 1716.
- 20d. *Hydrolases, acting on ester bonds (E.C. 3.1.-.-)*
- See 1415.
21. PURINES, PYRIMIDINES, NUCLEIC ACIDS AND THEIR CONSTITUENTS
- 21a. *Purines, pyrimidines, nucleosides, nucleotides*
- 1572 Malins, D.C. and Gunselman, S.J.: Fourier-transform infrared spectroscopy and gas chromatography-mass spectrometry reveal a remarkable degree of structural damage in the DNA of wild fish exposed to toxic chemicals. *Proc. Natl. Acad. Sci. U. S. A.*, 91 (1994) 13038-13041; *C.A.*, 122 (1995) 63783r.
- 1573 Ohba, S., Kidouchi, K., Sumi, S., Takeda, N., Yoshizumi, H., Tatematsu, A., Kobayashi, M. and Wada, Y.: Chemical diagnosis of dihydropyrimidinuria (the first case in Japan) by using HPLC and GC-MS. *Nippon Iyo Masu Supekutoru Gakkai Koenshu*, 19 (1994) 147-150; *C.A.*, 122 (1995) 103244z.
- 1574 Teixeira, A.J.R., Ferreira, M.R., van Dijk, W.J. van de Werken, G. and de Jong, A.P.J.M.: Analysis of 8-hydroxy-2'-deoxyguanosine in rat urine and liver DNA by stable isotope dilution gas chromatography/mass spectrometry. *Anal. Biochem.*, 227 (1995) 307-319.
- 21c. *Nucleic acids, DNA*
- 1575 Dizdaroglu, M.: Chemical determination of oxidative DNA damage by gas chromatography-mass spectrometry. *Methods Enzymol.*, 234 (1994) 1-16; *C.A.*, 121 (1994) 294482n.
- See also 1572.
22. ALKALOIDS
- 1576 Draeger, B.: Identification and quantification of calystegines, polyhydroxyl nortropane alkaloids. *Phytochem. Anal.*, 6 (1995) 31-37; *C.A.*, 122 (1995) 155671f.
- 1577 Gallagher, C.A., Hough, L.B., Keefner, S.M., Seyed-Mozaffari, A., Archer, S. and Glick, S.D.: Identification and quantification of the indole alkaloid ibogaine in biological samples by gas chromatography-mass spectrometry. *Biochem. Pharmacol.*, 49 (1995) 73-79; *C.A.*, 122 (1995) 122391m.
- 1578 Mizugaki, M.: Analysis of aconitum alkaloids by means of gas chromatography/mass spectrometry. *Nippon Iyo Masu Supekutoru Gakkai Koenshu*, 19 (1994) 65-72; *C.A.*, 122 (1995) 74046d.
- See also 1568.
23. OTHER SUBSTANCES CONTAINING HETEROCYCLIC NITROGEN
- 23c. *Indole derivatives and plant hormones (gibberelins)*
- See 1577.

23d. Pyridine derivatives

- 1579 Siret, L., Macaudiere, P., Bargmann-Leyder, N., Tambute, A., Caude, M. and Gougeon, E.: Separation of the optical isomers of a new 1,4-dihydropyridine calcium channel blocker (LF 2.0254) by liquid and supercritical fluid chromatography. *Chirality*, 6 (1994) 440-445.
- 1580 Titon, M.C. and Nardillo, A.M.: Separation of low-boiling pyridine bases by gas chromatography. *J. Chromatogr. A*, 699 (1995) 403-407.

23e. Other N-heterocyclic compounds

- 1581 Benetti, G. and Renzetti, A.: (Alkylpyrazines: important aroma components of licorice extracts.) *Ind. Aliment. (Pinerolo)*, 33 (1994) 639-642; *C.A.*, 122 (1995) 29983w.
- 1582 Taylor, G.W., Machan, Z.A., Mehmet, S., Cole, P.J. and Wilson, R.: Rapid identification of 4-hydroxy-3-alkylquinolines produced by *Pseudomonas aeruginosa* using gas chromatography-electron-capture mass spectrometry. *J. Chromatogr. B*, 664 (1995) 458-462.

See also 1203.

24. ORGANIC SULPHUR COMPOUNDS (INCL. GLUCOSINOLATES)

- 1583 Andersson, J.T. and Schmid, B.: Polycyclic aromatic sulfur heterocycles IV. Determination of polycyclic aromatic compounds in a shale oil with the atomic emission detector. *J. Chromatogr. A*, 693 (1995) 325-338.
- 1584 Anonymous: Dimethyl sulfide precursor in malt by headspace gas chromatography. *J. Am. Soc. Brew. Chem.*, 52 (1994) 193; *C.A.*, 122 (1995) 54381f.
- 1585 Betti, A., Lodi, G., Bigli, C., Chiorboli, G. and Coppi, S.: Use of overpressured layer chromatography and coupled OPLC-GC-MS for the analysis of acetylenic thiophene derivatives in extracts of *Tagete patula*. *J. Planar Chromatogr.*, 7 (1994) 301-304; *C.A.*, 122 (1995) 27180h.
- 1586 Choi, Y.-W., Kim, J.-H., Choe, K.-H. and Shin, S.-S.: (Determination of sulfur-containing odorants in natural gas by gas chromatography/flame photometric detection.) *Anal. Sci. Technol.*, 7 (1994) 349-359; *C.A.*, 122 (1995) 85125e.
- 1587 Heberer, T. and Stan, H.-J.: N-(Phenylsulfonyl)-sarcosine a new contaminant in sewage from ground water. *Fresenius Environ. Bull.*, 3 (1994) 639-643; *C.A.*, 122 (1995) 88883m.

See also 1256, 1259, 1260, 1565, 1571, 1748.

25. ORGANIC PHOSPHORUS COMPOUNDS (INCL. SUGAR PHOSPHATES)

- 1588 Habboush, A.E., Farroha, S.M. and Khalaf, H.I.: Extraction-gas chromatographic method for the determination of organophosphorus compounds as lubricating oil additives. *J. Chromatogr. A*, 696 (1995) 257-263.

See also 1718.

26. ORGANOMETALLIC AND RELATED COMPOUNDS

26a. Organometallic compounds

- 1589 Cai, Y. and Bayona, J.M.: Determination of methylmercury in fish and river water samples using *in situ* sodium tetraethylborate derivatization following by solid-phase microextraction and gas chromatography-mass spectrometry. *J. Chromatogr. A*, 696 (1995) 113-122.
- 1590 Cai, Y. and Bayona, J.M.: Simultaneous speciation of butyl-, phenyl-, and cyclohexyltin compounds in aqueous matrices using ethylation followed by solid-phase trace enrichment, SFC and GC determination. *J. Chromatogr. Sci.*, 33 (1995) 89-97.
- 1591 Chiavarini, S., Cremisini, C., Ingraio, G. and Morabito, R.: Determination of methylmercury in human hair by capillary GC with electron-capture detection. *Appl. Organomet. Chem.*, 8 (1994) 563-570; *C.A.*, 122 (1995) 125451s.
- 1592 Dirx, W.M.R.: Speciation of organotin compounds in waters and sediments using gas chromatography quartz furnace atomic absorption spectrometry (GC-QFAAS) and/or atomic emission detection (GC-AED). Avail. *Univ. Microfilms Int.*, Order No. DA9419247, 1993, 224 pp.; *C.A.*, 122 (1995) 114389b.
- 1593 Filippelli, M.: Methylmercury determination by purge and trap-GC-FTIR-AAS after NaBH₄ derivatization of an environmental thiosulfate extract. *Appl. Organomet. Chem.*, 8 (1994) 687-691; *C.A.*, 122 (1995) 204264m.
- 1594 Forsyth, D.S. and Hayward, S.: Further development and optimization of a quartz tube atomizer for a gas chromatography-atomic absorption spectrometer system. *Fresenius J. Anal. Chem.*, 351 (1995) 403-409.
- 1595 Harms, U.: Improvement of sample pretreatment for gas chromatographic determination of methylmercury in marine biota. *Appl. Organomet. Chem.*, 8 (1994) 645-648; *C.A.*, 122 (1995) 125453u.
- 1596 Mena, M.L., McLeod, C.W., Jones, P., Withers, A., Minganti, V., Capelli, R. and Quevauviller, P.: Microcolumn preconcentration and gas chromatography-microwave induced plasma-atomic emission spectrometry (GC-MIP-AES) for mercury speciation in waters. *Fresenius J. Anal. Chem.*, 351 (1995) 456-460.
- 1597 Minganti, V., Capelli, R. and de Pellegrini, R.: Evaluation of different derivatization methods for the multi-element analysis of Hg, Pb and Sn compounds by gas chromatography-microwave induced plasma-atomic emission spectrometry in environmental samples. *Fresenius J. Anal. Chem.*, 351 (1995) 471-477.
- 1598 Nerin, C. and Pons, B.: Speciation of organolead compounds in air by GC-MS-SIM. *Appl. Organomet. Chem.*, 8 (1994) 607-614; *C.A.*, 122 (1995) 140980g.
- 1599 Plzák, Z., Polanská, M. and Suchánek, M.: Identification and determination of butyltin compounds in water by ion trap gas chromatography-mass spectrometry after conversion to methyl or hydride derivatives. *J. Chromatogr. A*, 699 (1995) 241-252.
- 1600 Rubi, E., Lorenzo, R.A., Casais, M.C. and Cela, R.: Quality control in the routine analysis of methylmercury in biological and environmental materials using gas chromatography with electron capture detection. *Appl. Organomet. Chem.*, 8 (1994) 677-686; *C.A.*, 122 (1995) 125454v.

- 1601 Sanderson, R.D., Roediger, A.H.A. and Summers, G.J.: Hydrocarbon-soluble difunctional organolithium anionic initiators. A gas-liquid chromatographic study of the reaction of sec.-butyllithium with *m*-divinylbenzene. *Polym. Int.*, 35 (1994) 263-272; *C.A.*, 122 (1995) 10785m.
- 1602 Stab, J.A., Brinkman, U.A.T. and Cofino, W.P.: Validation of the analysis of organotin compounds in biological tissues using alkylation and gas chromatography. *Appl. Organomet. Chem.*, 8 (1994) 577-585; *C.A.*, 122 (1995) 125452t.
- 1603 Szpunar-Lobińska, J., Witte, C., Lobiński, R. and Adams, F.C.: Separation techniques in speciation analysis for organometallic species. *Fresenius J. Anal. Chem.*, 351 (1995) 351-377.
- 1604 Wang, K., Zhong, L. and Liu, S.: (Determination of aquatic butyltin compounds by chromatography with flame photometric detection.) *Huanjing Huaxue*, 13 (1994) 550-554; *C.A.*, 122 (1995) 196363a.
- 1605 Witte, C., Szpunar-Lobinska, J., Lobinski, R. and Adams, F.C.: Determination of trimethyllead in rainwater and road dust by capillary GC MIP-AE spectrometry after *in situ* ethylation and extraction. *Appl. Organomet. Chem.*, 8 (1994) 621-627; *C.A.*, 122 (1995) 141853e.
- 1606 Xu, F., Jiang, G. and Han, H.: (Determination of butyltin species in sediments by gas chromatography.) *Huanjing Kexue*, 15 (1994) 85-87; *C.A.*, 122 (1995) 141810p.
- See also 1609.
- 26b. *Boranes, silanes and related non-metallic compounds*
- 1607 Badescu, V., Radu, M., Zaharescu, M. and Vasilescu, A.: Hydrolysis-condensation of tetraethoxysilane in nonparental solvents studied by GC-MS. *J. Sol-Gel Sci. Technol.*, 2 (1994) 43-49; *C.A.*, 122 (1995) 133257y.
- 1608 Wachholz, S., Keidel, F., Just, U., Geissler, H. and Käßler, K.: Analysis of a mixture of linear and cyclic siloxanes by cryo-gas chromatography-Fourier transform infrared spectroscopy and gas chromatography-mass spectrometry. *J. Chromatogr. A*, 693 (1995) 89-99.
- 26c. *Coordination compounds*
- 1609 Lin, Y., Smart, N.G. and Wai, C.M.: Supercritical fluid extraction and chromatography on metal chelates and organometallic compounds. *TrAC*, 14 (1995) 123-132 - a review with 56 refs.
- See also 1207.
27. VITAMINS AND VARIOUS ANIMAL GROWTH FACTORS (NON-PEPTIDIC)
- 1610 Ballester, E., Gallego, M. and Valcárcel, M.: A gas chromatographic flow method for preconcentration and determination of vitamins D₂ and D₃ in pharmaceutical preparations. *Chromatographia*, 40 (1995) 425-431.
- 1611 Brunner, G.H. and Upmooor, D.: Scale up of supercritical fluid chromatography (SFC). *NATO ASI Ser., Ser. E*, 273 (1994) 653-668; *C.A.*, 122 (1995) 50391m.
- 1612 Ibáñez, E., Tabera, J., Reglero, G. and Herraiz, M.: Optimization of fat-soluble vitamin separation by supercritical fluid chromatography. *Chromatographia*, 40 (1995) 448-452.
- 1613 Norman, E.J. and Morrison, J.A.: Screening elderly population of cobalamin (vitamin B₁₂) deficiency using the urinary methylmalonic acid assay by gas chromatography mass spectrometry. *Am. J. Med.*, 94, No. 6 (1993) 589-594; *C.A.*, 122 (1995) 50676h.
- 1614 Riss, G., Kormann, A.W., Glinz, E., Walther, W. and Ranald, U.B.: Separation of the eight stereoisomers of all-rac- α -tocopherol from tissues and plasma: chiral phase HPLC and capillary gas chromatography. *Methods Enzymol.*, 234 (1994) 302-310; *C.A.*, 122 (1995) 4615q.
- 1615 Scalia, S., Ruberto, G. and Bonina, F.: Determination of vitamin A, vitamin E, and their esters in tablet preparations using supercritical fluid extraction and HPLC. *J. Pharm. Sci.*, 84 (1995) 433-436; *C.A.*, 122 (1995) 197095b.
- 1616 Yamada, S., Nitsuta, T. and Tomyama, S.: (Gas chromatography for determination of vitamin D₃ derivatives.) *Jpn. Kokai Tokkyo Koho JP 06,341,980 [94,341,980]* (Cl. G01N30/88), 13 Dec. 1994, Appl. 93/58,033, 24 Feb. 1993; 4 pp.; *C.A.*, 122 (1995) 128085m.
28. ANTIBIOTICS
- 1617 Akhtar, M.H., Danis, C., Sauve, A. and Barry, C.: Gas chromatographic determination of incurred chloramphenicol residues in eggs following optical extraction. *J. Chromatogr. A*, 696 (1995) 123-130.
29. INSECTICIDES, PESTICIDES AND OTHER AGROCHEMICALS
- 29a. *General techniques*
- 1618 Brooks, M.W.: Application of supercritical fluid extraction and chromatography in residue analysis. Avail. *Univ. Microfilms Int.*, Order No. DA9420603, 1994, 185 pp.; *C.A.*, 122 (1995) 154004d.
- 1619 Cheng, C.-L., Shih, S.-J., Yen, G.-C., Chou, S.-S. and Liu, C.-H.: Multiresidue determination of pesticides in fruits and vegetables by gas chromatography/mass spectrometry/selected ion monitoring. *Yaowu Shipin Fenxi*, 2 (1994) 113-122; *C.A.*, 122 (1995) 54312j.
- 1620 Eisert, R., Levsen, K. and Wünsch, G.: Analysis of polar thermally labile pesticides using different solid-phase extraction (SPE) materials with GC and HPLC techniques. *Intern. J. Environ. Anal. Chem.*, 58 (1995) 103-120.
- 1621 Hillmann, R. and Bächmann, K.: Extraction of pesticides using supercritical trifluoromethane and carbon dioxide. *J. Chromatogr. A*, 695 (1995) 149-154.
- 1622 Hirahara, Y., Narita, M., Okamoto, K., Miyoshi, T., Miyata, M., Koiguchi, S., Hasegawa, M., Kamakura, K., Yamana, T. and Tonogai, Y.: (Simple and rapid simultaneous determination of various pesticides of polished rice by gas chromatography.) *Shokuhin Eiseigaku Zasshi*, 35 (1994) 517-524; *C.A.*, 122 (1995) 8309r.
- 1623 Itano, K. and Okamoto, A.: (Multiresidue gas chromatographic method for determining pesticides in agricultural products.) *Kankyo Kagaku*, 4 (1994) 458-459; *C.A.*, 122 (1995) 158890n.

- 1624 Kawasaki, M., Fukuhara, K. and Uchiyama, S.: (Gas chromatographic-mass spectrometric (GC/MS) screening of pesticides.) *Shokuhin Eiseigaku Zasshi*, 35 (1994) 479-496; C.A., 122 (1995) 29978y.
- 1625 Li, L., Guan, Y. and Zhou, L.: (Standart Kovats indexes of 20 pesticides and their identification in temperature-programmed capillary gas chromatography.) *Fenxi Huaxue*, 23 (1995) 14-19; C.A., 122 (1995) 99228c.
- 1626 Mojasevic, M. and Helling, C.S.: Use of multiresidue gas chromatographic analysis to determine pesticide mobility on soil TLC plates. *J. Environ. Sci. Health, Part B*, 30, No. 2 (1995) 163-173; C.A., 122 (1995) 180905t.
- 1627 Newsome, W.H.: An evolution of pesticide residue methodology. *J. Assoc. Off. Anal. Chem. Int.*, 78 (1995) 4-8.
- 1628 Scaroni, I., Previati, M.P. and Bovolenta, A.: (Evaluation of a multiresidue method for the determination of pesticide residues in wine - Note 1.) *Ind. Bevande*, 23 (1994) 222-226; C.A., 122 (1995) 29964r.
- 1629 Snyder, J.L., Grob, R.L., McNally, M.E. and Oostdyk, T.S.: Supercritical fluid extraction of selected pesticides from fortified soils and determination by gas chromatography with electron capture detection. *J. Environ. Sci. Health*, A29 (1994) 1801-1816; C.A., 121 (1994) 294955u.
- 1630 Tuinstra, L.G.M.T., van de Spreng, P. and Gaikhorst, P.: Ion trap detection for development of a multi residue/multi matrix method for pesticide residues in agricultural products. *Intern. J. Environ. Anal. Chem.*, 58 (1995) 81-91.
- 1631 Wan, H.B., Wong, M.K. and Mok, C.Y.: Use of statistically designed experiments to minimize the effect of water on gas chromatographic analysis. *J. Chromatogr. Sci.*, 33 (1995) 66-70.
- 1632 Williams, C.S., Eastoe, B.V., Slaiding, I.R. and Walker, M.D.: Analysis of pesticide residues in hops and their extraction by liquid CO₂ during the production of hop extracts. *Food Addit. Contam.*, 11 (1994) 615-619; C.A., 122 (1995) 79345k.
- See also 1312, 1551.
- 29b. *Chlorinated insecticides*
- 1633 Bentabol, A. and Jodral, M.: Determination of organochlorine pesticides in cheese. *J. Assoc. Off. Anal. Chem. Int.*, 78 (1995) 94-98.
- 1634 Hainzl, D., Burhenne, J. and Parlar, H.: (Isolation of single components for toxaphene analysis.) *GIT Fachz. Lab.*, 38 (1994) 285-294; C.A., 121 (1994) 294490p.
- 1635 Hirama, Y., Nakano, M. and Nishimura, K.: (Analytical method for PCBs and organochlorine pesticides by dual-column capillary gas chromatography.) *Hokkaidoritsu Eisei Kenkyushoho*, 44 (1994) 73-76; C.A., 122 (1995) 79333e.
- 1636 Ling, Y.-C. and Huang, I.-P.: Multiresidue-matrix solid-phase dispersion method for determining 16 organochlorine pesticides and polychlorinated biphenyls in fish. *Chromatographia*, 40 (1995) 259-266.
- 1637 Liu, C.: Field detection of organochlorine pesticides, explosives, and volatile organic compounds. Avail. *Univ. Microfilms Int.*, Order No. DA9425790, 1994, 133 pp.; C.A., 122 (1995) 150594y.
- 1638 Macnaughton, S.J., Kikic, I., Rovedo, G., Foster, N.R. and Alessi, P.: Solubility of chlorinated pesticides in supercritical carbon dioxide. *J. Chem. Eng. Data*, 40 (1995) 593-597.
- 1639 Mariani, E., Bargagna, A., Longobardi, M. and Bruschi, E.: Gas chromatographic determination of chlorinated pesticides and PCBs in complex cosmetic matrixes. *Farmaco*, 49 (1994) 441-442; C.A., 122 (1995) 16814s.
- 1640 Mariani, G., Benfenati, E. and Fanelli, R.: A Nici-GC-MS method to analyze endosulfan in biological samples. *Intern. J. Environ. Anal. Chem.*, 58 (1995) 67-72.
- 1641 Natangelo, M., Romagnano, S., Guzzella, L. and Benfenati, E.: GC analysis of some organochlorine pesticides using a brominated internal standard. *Intern. J. Environ. Anal. Chem.*, 58 (1995) 55-66.
- 1642 Nowak, J.: Determination of phosalone and methoxychlor residues in soil by gas chromatography. *Chem. Anal. (Warsaw)*, 39 (1994) 681-685.
- 1643 Ong, V.S. and Hites, R.A.: Determination of pesticides and polychlorinated biphenyls in water: A low-solvent method. *Environ. Sci. Technol.*, 29 (1995) 1259-1266.
- 1644 Pfaffenberger, B., Hühnerfuss, H., Gehrcke, B., Hardt, I., König, W.A. and Rimkus, G.: Gas chromatographic separation of the enantiomers of bromocyclen in fish samples. *Chemosphere*, 29 (1994) 1385-1391.
- 1645 Polese, L., Minelli, E.V. and Ribeiro, M.L.: An online method for the determination of organochlorine pesticide residues in soybean derivative. *J. Braz. Chem. Soc.*, 5 (1994) 19-22; C.A., 122 (1995) 29966t.
- 1646 Rao, V.S.: Detection of organochlorine pesticide residues using gas-chromatography in village drinking water ponds and in biota of Guntur (Dist.) of Andhra Pradesh. *Geobios (Jodhpur)*, 21 (1994) 68-73; C.A., 121 (1994) 307730u.
- 1647 Schantz, M.M., Koster, B.J., Oakley, L.M., Schiller, S.B. and Wise, S.A.: Certification of polychlorinated biphenyl congeners and chlorinated pesticides in a whale blubber standart reference material. *Anal. Chem.*, 67 (1995) 901-910.
- 1648 Yuan, L., Chen, Q., Wang, Y. and Jing, Z.: (Method for the determination of residual PCNB and benzene hexachloride isomers in ginseng (*Panax ginseng*) by capillary column gas chromatography.) *Zhongcaoyao*, 25 (1994) 519-520; C.A., 122 (1995) 64554d.
- 1649 Yuanhong, W. and Rong, H.: (Determination of organic chloride pesticide residues in ginseng with capillary gas chromatography.) *Fenxi Huaxue*, 22 (1994) 931-934; C.A., 122 (1995) 17328y.
- 29c. *Phosphorus insecticides*
- 1650 Barra, R. and Carmi, J.: A simple method for extraction and GC-ECD analysis of chlorpyrifos from water and environmental samples. *J. High Resolut. Chromatogr.*, 18 (1995) 194-195.
- 1651 Cai, C.P., Liang, M. and Wen, R.R.: Rapid multiresidue screening method for organophosphate pesticides in vegetables. *Chromatographia*, 40 (1995) 417-420.
- 1652 Camel, V., Caude, M. and Tambute, A.: SFE of an organophosphorus compound from soils with capillary GC analysis. *J. Chromatogr. Sci.*, 33 (1995) 123-132.

- 1653 Eisert, R. and Levsen, K.: Determination of organophosphorus, triazine and 2,6-dinitroaniline pesticides in aqueous samples via solid-phase microextraction (SPME) and gas chromatography with nitrogen-phosphorus detection. *Fresenius J. Anal. Chem.*, 351 (1995) 555-562.
- 1654 Hankemeier, T., Louter, A.J.H., Rinkema, F.D. and Brinkman, U.A.T.: On-line coupling of solid-phase extraction and gas chromatography with atomic emission detection for analysis of trace pollutants in aqueous samples. *Chromatographia*, 40 (1995) 119-124.
- 1655 Saenz Barrio, C., Sanz Asensio, J., Plaza Medina, M. and Perez Clavijo, M.: Evaluation of the decay of malathion, dichlorofluanid and fenitrothion pesticides in apple samples, using gas chromatography. *Food Chem.*, 52 (1995) 305-309; *C.A.*, 122 (1995) 185788s.
- 1656 Sakai, T., Imamura, T., Sato, M., Ogawa, M., Shimura, K., Ohkuma, K., Mori, Y., Kurata, H., Ohida, T. et al.: (Rapid and simultaneous determination of organophosphorus pesticides in agricultural products by FDP-GC.) *Shokuhin Eiseigaku Zasshi*, 35 (1994) 636-644; *C.A.*, 122 (1995) 131287j.
- 29d. *Carbamates*
- 1657 Ma, Y.-P., Deng, F.-Q., Chen, D.-Z. and Sun, S.-W.: Determination of ethyl carbamate in alcoholic beverages by capillary multi-dimensional gas chromatography with thermionic specific detection. *J. Chromatogr. A*, 695 (1995) 259-265.
- 29e. *Herbicides*
- 1658 Antonovic, D.G. and Boncic-Caricic, G.A.: (Gas chromatographic retention indexes for N-substituted amino s-triazines on DB-1 and DB-5 capillary columns.) *J. Serb. Chem. Soc.*, 59 (1994) 993-996; *C.A.*, 122 (1995) 203979m.
- 1659 Bonwick, G.A., Abdul-Latif, P., Sun, C., Baugh, P.J., Smith, C.J., Armitage, R. and Davies, D.H.: Comparison of chemical methods and immunoassay for the detection of pesticide residues in various matrices. *Food Agric. Immunol.*, 6 (1994) 267-276; *C.A.*, 122 (1995) 74472q.
- 1660 Brady, J.F., LeMasters, G.S., Williams, R.K., Pittman, J.H., Daubert, J.P., Cheung, M.W., Skinner, D.H., Turner, J. and Rowland, M.A.: Immunoassay analysis and gas chromatography confirmation of atrazine residues in water samples from a field study conducted in the state of Wisconsin. *J. Agric. Food Chem.*, 43, No. 1 (1995) 268-274.
- 1661 Gruessner, B., Shambaugh, N.C. and Watzin, M.C.: Comparison of an enzyme immunoassay and gas chromatography/mass spectrometry for the detection of atrazine in surface waters. *Environ. Sci. Technol.*, 29 (1995) 251-254.
- 1662 Heberer, T., Butz, S. and Stan, H.-J.: Analysis of phenoxycarboxylic acids and other acidic compounds in tap, ground, surface and sewage water at the low ng/l level. *Intern. J. Environ. Anal. Chem.*, 58 (1995) 43-53.
- 1663 Kumazawa, T., Sato, K., Seno, H., Ishii, A. and Suzuki, O.: Capillary gas chromatography with four different detectors for dinitroaniline herbicides in human body fluids. *J. Anal. Toxicol.*, 19 (1995) 95-98; *C.A.*, 122 (1995) 180436r.
- 1664 Lanças, F.M., Barbirato, M.A. and Galhiane, M.S.: Extraction of norflurazon residues in cotton seeds with supercritical CO₂. *Chromatographia*, 40 (1995) 432-434.
- 1665 Thio, A.P. and Kornet, M.J.: Confirmation and analysis of s-chlorotriazines by derivatization with pyrrolidine. *Anal. Lett.*, 28 (1995) 677-684.
- 1666 Ting, K.-C. and Lee, C.-S.: Gas chromatographic determination of triclopyr in fruits and vegetables. *J. Chromatogr. A*, 690 (1995) 119-129.
- See also 1429, 1653.
- 29f. *Fungicides*
- 1667 Van Doorn, C., Vink, M. and van der Poll, J.M.: Gas chromatographic determination of chlorothalonil and its metabolite 4-hydroxy-2,5,6-trichlorisophthalonitrile (HTI) in water. *Chromatographia*, 40 (1995) 458-462.
- See also 1425.
- 29g. *Other types of pesticides and various agrochemicals*
- 1668 Bingkun, X., Yimin, S. and Lianying, L.: (Analysis of the chemical components of TY germicide for crops by gas chromatography.) *Fenxi Huaxue*, 22 (1994) 829-832; *C.A.*, 122 (1995) 3432r.
- 1669 Ling, Y.-C. and Huang, L.-P.: Multi-residue matrix solid-phase dispersion method for the determination of six synthetic pyrethroids in vegetables followed by gas chromatography with electron capture detection. *J. Chromatogr. A*, 695 (1995) 75-82.
- 1670 Mallinson, E.T., Dreas, J.S., Wilson, R.T. and Henry, A.C.: Determination of dexamethasone in liver and muscle by liquid chromatography and gas chromatography/mass spectrometry. *J. Agric. Food Chem.*, 43 (1995) 140-145.
- 1671 Zhou, Y.: (Determination of chlordimeform residue in honey by using GC with wide-bore capillary column.) *Fenxi Ceshi Xuebao*, 13, No. 4 (1994) 60-62; *C.A.*, 122 (1995) 54295f.
31. PLASTICS AND THEIR INTERMEDIATES
- 1672 Albertsson, A.-C., Barenstedt, C. and Karlsson, S.: Solid-phase extraction and gas chromatographic-mass spectrometric identification of degradation products from enhanced environmentally degradable polyethylene. *J. Chromatogr. A*, 690 (1995) 207-217.
- 1673 Calvey, E., M., Begley, T.H. and Roach, J.A.G.: SFC-FTIR spectroscopy and SFC-MS of nonvolatile extractants of microwave susceptor packaging. *J. Chromatogr. Sci.*, 33 (1995) 61-65.
- 1674 Choi, Y., Lee, H., Fountain, S.T. and Lubman, D.M.: Direct chemical analysis of UV laser ablation products of organic polymers by using selective ion monitoring mode in gas chromatography/mass spectrometry. *J. Am. Soc. Mass Spectrom.*, 5 (1994) 106-112; *C.A.*, 122 (1995) 32730s.
- 1675 Hadj Romdhane, I.: Polymer-solvent diffusion and equilibrium parameters by inverse gas-liquid chromatography. Avail. *Univ. Microfilms Int.*, Order No. DA9428107, 1994, 235 pp.; *C.A.*, 122 (1995) 161994e.
- 1676 Hiramatsu, S., Akieda, T. and Inoma, S.: (Quantitative analysis of polybutene.) *Kanzei Chuo Bunsekishoho*, 33 (1994) 107-115; *C.A.*, 122 (1995) 134374w.

- 1677 Jackson, P.L., Huglin, M.B. and Cervenka, A.: Use of inverse gas chromatography to quantify interactions in amine cured epoxy resins. *Polym. Int.*, 35 (1994) 119-133; *C.A.*, 121 (1994) 302187n.
- 1678 Judge, M.D.: Investigation of the thermal oxidation of a polybutadiene by head-space sampling-gas chromatography. *Talanta*, 41 (1994) 755-759.
- 1679 Kwong, E., Baert, L. and Bechard, S.: Gas-liquid chromatographic quantitation of polyethylene glycol 400 in pharmaceutical preparations. *J. Pharm. Biomed. Anal.*, 13 (1995) 77-81; *C.A.*, 122 (1995) 142724a.
- 1680 Lezcano, E.G., Prolongo, M.G. and Coll, C.S.: Characterization of the interactions in the poly(4-hydroxystyrene)/poly(ϵ -caprolactone) system by inverse gas chromatography. *Polymer*, 36 (1995) 565-573; *C.A.*, 122 (1995) 188829s.
- 1681 MacMahon, T. and Chace, M.: Poly(amic acid) and polyimide characterization using gas chromatography/mass spectrometry and particle beam liquid chromatography/mass spectrometry. *J. Am. Soc. Mass Spectrom.*, 5 (1994) 299-304; *C.A.*, 122 (1995) 161983a.
- 1682 Pleshkova, A.P. and Volkovich, S.V.: (Analysis of polycarbonate compositions of unknown composition using a chromatography/mass spectrometry method.) *Khim. Prom-st. (Moscow)*, 2 (1994) 86-89.
- 1683 Pokorny, J., Zchoval, J. and Kubat, J.: (Study of migration of volatile components from plastics by the headspace method.) *Plasty Kauc.*, 31 (1994) 106-109.
- 1684 Sithole, B.B. and Allen, L.H.: Quantitative analysis of synthetic polymers in pulp and paper mill pitch deposits by pyrolysis gas chromatography. *J. Pulp Pap. Sci.*, 20 (1994) 168-172; *C.A.*, 122 (1995) 109031z.
- 1685 Sturman, H.W., Geissler, M., Weimar, A., Paul, A. and Nickel, B.: (High performance liquid chromatography (HPLC) and gas chromatography (GC) as methods in plastic material analysis.) *Kautsch. Gummi Kunstst.*, 47 (1994) 812-815; *C.A.*, 122 (1995) 107514k.
- 1686 Yang, X.: (Reinforced plastics based on unsaturated polyester resins - determination of residual styrene monomer content by gas chromatography. *Huaxue Shijie*, 34 (1993) 220-223; *C.A.*, 122 (1995) 82675y.

See also 1211, 1219, 1360, 1765.

32. DRUG ANALYSIS

32a. Drug analysis, general techniques

- 1687 Abe, I., Nishiyama, T., Nakahara, T. and Frank, H.: Gas chromatographic enantiomer separation of pharmaceuticals on capillary columns coated with novel chiral polysiloxanes. *J. Chromatogr. A*, 694 (1995) 237-243.
- 1688 Feng, C., Liu, Y. and Luo, Y.: (Systematic analysis for twenty-one hypnotic and sedative drugs by two different GC columns and detectors.) *Yaowu Fenxi Zazhi*, 14, No. 5 (1994) 3-6; *C.A.*, 122 (1995) 17334x.
- 1689 Ferry, M.A.: GC/MS for volatile organics. *Environ. Test. Anal.*, 3, No. 5 (1994) 46-50; *C.A.*, 122 (1995) 204167g.
- 1690 Noggle, F.T., Clark, C.R. and DeRuiter, J.: GC-MS and LC of addition products formed from the reaction of allylbenzene and related arylpropenes with acetonitrile and sulfuric acid. *J. Chromatogr. Sci.*, 33 (1995) 255-262.
- 1691 Ohayon-Courtes, C.: Significance of gas chromatography coupled with mass spectrometry for the pharmaceutical product control laboratory. *Ann. Falsif. Expert. Chim. Toxicol.*, 87 (1994) 135-144; *C.A.*, 121 (1994) 308439t.
- 1692 Rasanen, I., Ojanperä, I. and Vuori, E.: Comparison of four homologous retention index standard series for gas chromatography of basic drugs. *J. Chromatogr. A*, 693 (1995) 69-78.

See also 1610, 1679.

32b. Antirheumatics and antiinflammatory drugs

See 1392.

32c. Autonomic and cardiovascular drugs

- 1693 Bargmann-Leyder, N., Sella, C., Bauer, D., Tambuté, A. and Caude, M.: Supercritical fluid chromatographic separation of β -blockers on Chyrosine-A: Investigation of the chiral recognition mechanism using molecular modeling. *Anal. Chem.*, 67 (1995) 952-958.
- 1694 Nishikawa, M., Ishii, A., Watanabe, K., Hiroshi, S., Kumazawa, T. and Suzuki, O.: Determination of diltiazem in body fluids by gas chromatography with surface ionization detection. *Hochudoku*, 12 (1994) 247-251; *C.A.*, 122 (1995) 71235d.
- 1695 Xu, Y., Shen, L., Wu, Y., Wang, S. and Zhang, C.: (Analysis of beta-blockers and their metabolites in urine by gas chromatogr. II. Analysis of carteolol, pindolol, timolol and their metabolites.) *Fenxi Ceshi Xuebao*, 13, No. 6 (1994) 85-87; *C.A.*, 122 (1995) 150732s.

See also 1579.

32d. Central nervous system drugs

- 1696 Berger, T.A. and Wilson, W.H.: Separation of basic drugs by packed-column supercritical fluid chromatography 3. Stimulants. *J. Pharm. Sci.*, 84 (1995) 489-492; *C.A.*, 122 (1995) 197135q.
- 1697 Chiarotti, M. and Marsili, R.: Gas chromatographic analysis of methadone in urine samples after solid phase microextraction. *J. Microcolumn Separ.*, 6 (1994) 577-580.
- 1698 Feng, N., Vollenweider, F.X., Minder, E.I., Rentsch, K., Grampp, T. and Vonderschmitt, D.J.: Development of a gas chromatography-mass spectrometry method for determination of ketamine in plasma and its application to human samples. *Ther. Drug Monit.*, 17 (1995) 95-100; *C.A.*, 122 (1995) 177554x.
- 1699 Hu, X., Liu, F. and Luo, Y.: (Analysis for local anesthetics in plasma.) *Yaowu Fenxi Zazhi*, 14, No. 6 (1994) 7-10; *C.A.*, 122 (1995) 95761y.
- 1700 Ishii, A., Nishikawa, M., Seno, H., Kumazawa, T., Watanabe, K. and Suzuki, O.: Determination of trihexyphenidyl in body fluids by gas chromatography with surface ionization detection. *Hochudoku*, 12 (1994) 241-246; *C.A.*, 122 (1995) 71236e.

- 1701 Jacob, P., Ill, Tisdale, E.C., Panganiban, K., Cannon, D., Zabel, K., Mendelson, J.E. and Jones, R.T.: Gas chromatographic determination of methamphetamine and its metabolite in human plasma and urine following conversion to N-propyl derivatives. *J. Chromatogr. B*, 664 (1995) 449-457.
- 1702 Jeannot, C. and Rohrbasser, C.: (Gas chromatographic/mass spectrometric analysis of β -agonists using an immunoaffinity column.) *Chimia*, 48 (1994) 413-416; *C.A.*, 122 (1995) 71239h.
- 1703 Kingsbury, D.P., Makowski, G.S. and Stone, J.A.: Quantitative analysis of fentanyl in pharmaceutical preparations by gas chromatography-mass spectrometry. *J. Anal. Toxicol.*, 19 (1995) 27-30; *C.A.*, 122 (1995) 89600d.
- 1704 Li, H., Zhang, Y., Yan, X., Tang, X. and Jing, Y.: (Capillary gas chromatographic analysis of central nerve system sedatives in blood.) *Yaowu Fenxi Zazhi*, 14, No. 5 (1994) 7-9; *C.A.*, 122 (1995) 23105q.
- 1705 Lorec, A.-M., Bruguerolle, B., Attolini, L. and Roucoules, X.: Rapid simultaneous determination of lidocaine, bupivacaine, and their two main metabolites using capillary gas-liquid chromatography with nitrogen phosphorus detector. *Ther. Drug Monit.*, 16 (1994) 592-595; *C.A.*, 122 (1995) 71207w.
- 1706 Luo, Y., Feng, J.L., Liu, F. and Hu, X.Y.: (Determination of dihydroetorpine hydrochloride in biological fluids.) *Yaoxue Xuebao*, 29 (1994) 702-706; *C.A.*, 122 (1995) 23104p.
- 1707 Meatherall, R.: GC-MS conformation of urinary benzodiazepine metabolites. *J. Anal. Toxicol.*, 18 (1994) 369-381; *C.A.*, 122 (1995) 25448c.
- 1708 Noggle, F.T., Clark, C.R. and DeRuiter, J.: Gas chromatographic and mass spectral analysis of methamphetamine synthesized from allylbenzene. *J. Chromatogr. Sci.*, 33 (1995) 153-161.
- 1709 Sautou, V., Constant, H., Bonnefous, J.-L., Chopineau, J. and Aulagner, G.: (Analysis of opiates in urine of children by gas chromatography/mass spectrometry.) *J. Pharm. Belg.*, 49 (1994) 383-389; *C.A.*, 122 (1995) 2918s.
- 1710 Schurig, V., Green, B.S. and Grosenick, H.: (Preparative gas phase chromatographic resolution of inhalation anesthetics.) *Ger. Offen.* DE 4,317,139 (Cl. C01B57/00), 24 Nov. 1994, Appl. 21 May 1993; 4 pp.; *C.A.*, 122 (1995) 132557j.
- 1711 Schurig, V., Jung, M., Mayer, S., Negura, S., Fluck, M. and Jakubetz, H.: (Enantioselective separation of hexobarbital by gas chromatography (GC), supercritical fluid chromatography (SFC), liquid chromatography (LC), and cation-exchange chromatography (CEC) on capillary column packed with Chirasil-Dex.) *Angew. Chem.*, 106 (1994) 2265-2267; *C.A.*, 122 (1995) 142721x.
- 1712 Song, D., Khaykis, V. and Kohlhof, K.: Determination of flumazenil in plasma by gas chromatography-negative chemical ionization mass spectrometry. *J. Chromatogr. B*, 663 (1995) 263-273.
- 1713 Ulrich, S., Meyer, F.P., Neuhofer, S. and Knorr, W.: Megabore capillary gas-liquid chromatographic method with nitrogen-phosphorus selective detection for the assay of haloperidol and reduced haloperidol in serum: results of therapeutic drug monitoring during acute therapy of eight schizophrenics. *J. Chromatogr. B*, 663 (1995) 289-296.
- 1714 Xie, X., Yang, Z., Li, J., Cai, C. and Pang, J.: (Study on the relative bioavailability in human body of tramadol hydrochloride tablets by gas chromatography.) *Zhongguo Yaoxue Zazhi*, 29 (1994) 671-673; *C.A.*, 122 (1995) 71293w.
- 1715 Yamaguchi, Y., Yasuki, Y., Maeda, T., Suzuki, T., Yamamoto, N. and Morita, M.: (Automated analysis of trace organic compounds in the atmosphere by thermal desorption-GC dual detector (photo ionization detector with electrolytic conductivity detector.) *Kankyo Kagaku*, 4 (1994) 510-511; *C.A.*, 122 (1995) 195400y.

See also 1231, 1391, 1688, 1724, 1726, 1731.

32e. Chemotherapeutics (exc. cytostatics and antibiotics)

- 1716 Fraser, I.J., Clare, R.A. and Pleasance, S.: Determination of the acylcoenzyme A cholesterol acyltransferase inhibitor 447C88 in plasma using gas chromatography-mass spectrometry and liquid chromatography atmospheric pressure chemical ionization tandem mass spectrometry. *J. Chromatogr. B*, 663 (1995) 275-287.
- 1717 Tena, M.T., Lague de Castro, M.D. and Valcárcel, M.: Improved supercritical fluid extraction of sulphonamides. *Chromatographia*, 40 (1995) 197-203.

See also 1668.

32f. Cytostatics

- 1718 Steelant, W.F.A., Bruyneel, E.A., Mareel, M.M. and van den Eeckhout, E.G.: Capillary gas chromatography of hexadecylphosphocholine in Caco-2T cells and cell culture media. *Anal. Biochem.*, 227 (1995) 246-250.

See also 1453.

32g. Other drug categories

- 1719 Huang, Q., Liang, B., Lu, H. and Su, H.: (Gas chromatographic determination of menthol and menthone in Yinqiao-Jiedu granules.) *Guangdong Yiyao Xueyuan Xuebao*, 10, No. 2 (1994) 90-91; *C.A.*, 122 (1995) 197100z.
- 1720 Wu, G. and Liu, C.: (Determination of Norcantharidi in tablets by gas chromatography (GC).) *Chin. J. Chromatogr.*, 13 (1995) 142-143.
- 1721 Yang, L.L., Yuan, Y.S., Tong, Y.L. and Guo, C.Y.: Determination of tetramethylpyrazine and borneol in human plasma by gas chromatography-mass spectrometry. *Yaoxue Xuebao*, 29 (1994) 697-701; *C.A.*, 122 (1995) 200c.

See also 1524, 1528.

32h. Toxicological and forensic applications

- 1722 Karkkainen, M., Sippola, E., Pikkariainen, A.-L., Rautio, T. and Himberg, K.: Automated gas chromatographic amphetamine profiling. *Forensic Sci. Int.*, 69 (1994) 55-64; *C.A.*, 122 (1995) 180423j.
- 1723 Kudo, K., Nagata, T., Kimura, K., Imamura, T. and Jitsufuchi, N.: Sensitive determination of Δ^9 -tetrahydrocannabinol in human tissues by GC-MS. *J. Anal. Toxicol.*, 19 (1995) 87-90; *C.A.*, 122 (1995) 180435q.

- 1724 Liu, R.H., McKeehan, A.M., Edwards, C., Foster, G., Bensley, W.D., Langner, J.G. and Walia, A.S.: Improved gas chromatography/mass spectrometry analysis of barbiturates in urine using centrifuge-based solid-phase extraction, methylation, with *d*₅-pentobarbital as internal standard. *J. Forensic Sci.*, 39 (1994) 1504-1514; *C.A.*, 122 (1995) 48623v.
- 1725 Maurer, H.H.: Toxicological analysis of drugs and poisons by GC/MS. *Spectrosc. Eur.*, 6, No. 6 (1994) 21-23; *C.A.*, 122 (1995) 125464y.
- 1726 Mieczkowski, T.: A research note: The outcome of GC/MS/MS confirmation of hair assays on 93 cannabinoid (+) cases. *Forensic Sci. Int.*, 70 (1995) 83-91; *C.A.*, 122 (1995) 153484y.
- 1727 Moeller, M.R. and Mueller, C.: The detection of 6-monoacetylmorphine in urine, serum and hair by GC/MS and RIA. *Forensic Sci. Int.*, 70 (1995) 125-133; *C.A.*, 122 (1995) 153486a.
- 1728 Neumann, H.: Comparison of heroin by capillary gas chromatography in Germany. *Forensic Sci. Int.*, 69 (1994) 7-16; *C.A.*, 122 (1995) 180419n.
- 1729 Ripani, L., Schiavone, S. and Garofano, L.: GC quantitative determination of illicit LSD. *J. Forensic Sci.*, 39 (1994) 512-517; *C.A.*, 122 (1995) 48601m.
- 1730 Seto, Y.: (Headspace gas chromatography in forensic toxicology.) *Hochudoku*, 12 (1994) 175-191; *C.A.*, 122 (1995) 74106y.
- 1731 Smith, M.L., Hughes, R.O., Levine, B., Dickerson, S., Darwin, W.D. and Cone, E.J.: Forensic drug testing for opiates. VI. Urine testing for hydromorphone, hydrocodone, oxycodone, and oxycodone with commercial opiate immunoassays and gas chromatography-mass spectrometry. *J. Anal. Toxicol.*, 19 (1995) 18-26; *C.A.*, 122 (1995) 74109b.
- 1732 Tanaka, K., Ohmori, T., Inoue, T. and Seta, S.: Impurity profiling analysis of illicit metamphetamine by capillary gas chromatography. *J. Forensic Sci.*, 39 (1994) 500-511; *C.A.*, 122 (1995) 48600k.
- 1733 Watson, D.G., Su, Q., Midgley, J.M., Doyle, E. and Morton, N.S.: Analysis of unconjugated morphine, codeine, and normorphine and of morphine glucuronides in small volumes of plasma from children. *J. Pharm. Biomed. Anal.*, 13 (1995) 27-32; *C.A.*, 122 (1995) 150728m.
- See also 1351, 1359, 1404, 1418, 1460, 1484, 1560.
- 32i. *Plant extracts*
- 1734 O'Rear, C.E. and Llewellyn, G.C.: A rapid analytical procedure for Δ^9 -tetrahydrocannabinol in plant material. In: Llewellyn, G.C., Dashek, W.V. and O'Rear, C.E. (Editors), *Biodeterior. Res.* 4, Plenum, New York, 1994, pp. 493-497; *C.A.*, 122 (1995) 180417k.
- See also 1542.
34. FOOD ANALYSIS
- 34b. *Complex mixtures (single compounds by cross-reference only)*
- 1735 Laber, H., Schulz, J., Sponholz, W.R. and Bremser, W.: Use of a combined GC/FTIR/MS system for the analysis of spirits. *Freisenius J. Anal. Chem.*, 351 (1995) 530-535.
- See also 1349, 1356, 1380, 1389, 1424, 1429, 1440, 1461, 1480, 1487, 1488, 1493, 1504, 1505, 1512, 1518, 1519, 1520, 1561, 1584, 1589, 1617, 1619, 1623, 1628, 1655, 1657, 1666, 1669, 1790, 1791, 1793, 1807.
- 34c. *Organoleptically important compounds (flavors, odors, volatiles)*
- 1736 Akochi, K.-E., Alli, L., Kermasha, S. and Yaylayan, V.: Quantitation of alkylpyrazines in maple syrup, maple flavors and non-maple syrups. *Food Res. Int.*, 27, No. 5 (1994) 451-457; *C.A.*, 122 (1995) 8298m.
- 1737 Apps, P.J. and Veary, C.M.: (High precision profiling of volatiles from raw meat.) *Arch. Lebensmittelhyg.*, 44 (1993) 92-95; *C.A.*, 121 (1994) 299386u.
- 1738 Blank, I., Staempfli, A. and Eisenreich, W.: Analysis of food flavorings by gas chromatography-olfactometry. *Dev. Food Sci.*, 35 (1994) 271-276; *C.A.*, 121 (1994) 299526q.
- 1739 Garcia-Jares, C., Garcia-Martin, S. and Cela-Torrijos, R.: Analysis of some highly volatile compounds of wine by means of purge and cold trapping injector capillary gas chromatography. Application to the differentiation of Rias Baixas Spanish white wines. *J. Agric. Food Chem.*, 43 (1995) 764-768.
- 1740 Harayama, K., Hayase, F. and Kato, H.: New method for analysing the volatiles in beer. *Biosci., Biotechnol., Biochem.*, 58 (1994) 2246-2247; *C.A.*, 122 (1995) 79365s.
- 1741 Koppenhoefer, B., Behnisch, R., Epperlein, U., Holzschuh, H., Bernreuther, A., Piras, P. and Roussel, C.: Enantiomeric odor differences and gas chromatographic properties of flavors and fragrances. *Perfum. Flavor.*, 19, No. 5 (1994) 1-14; *C.A.*, 121 (1994) 307955w.
- 1742 Milo, C. and Grosch, W.: Detection of odor defects in boiled cod and trout by gas chromatography-olfactometry of headspace samples. *J. Agric. Food Chem.*, 43 (1995) 459-462.
- 1743 Mosandl, A.: (Gas chromatography-isotope ratio mass spectrometry (GC-IRMS) in analysis of odorous compounds.) *GIT Fachz. Lab.*, 38 (1994) 882-888; *C.A.*, 122 (1995) 196518e.
- 1744 Schulzki, G., Spiegelberg, A., Boegl, K.W. and Schreiber, G.A.: Detection of radiation-induced hydrocarbons in camembert irradiated before and after the maturing process-comparison of florisil column chromatography and online coupled liquid chromatography-gas chromatography. *J. Agric. Food Chem.*, 43 (1995) 372-376.
- 1745 Villen, J., Senorans, F.J., Reglero, G. and Herraiz, M.: Analysis of wine aroma by direct injection and gas chromatography without previous extraction. *J. Agric. Food Chem.*, 43 (1995) 717-722.
- 1746 Waldhams, L.J., Blight, M.M., Kerguelen, V., le Metayer, M., Marion-Poll, F., Masson, C., Pham-Delegue, M. and Woodcock, C.M.: Discrimination of oilseed rape volatiles by honey bee: novel combined gas chromatographic-electrophysiological behavioral assay. *J. Chem. Ecol.*, 20 (1994) 3221-3231; *C.A.*, 122 (1995) 50705s.
- 1747 Wood, A.F., Aston, J.W. and Douglas, G.K.: A cold-trap method for the collection and determination of headspace compounds from cheese. *Aust. J. Dairy Technol.*, 49 (1994) 42-47; *C.A.*, 122 (1995) 54286d.

1748 Wyllie, S.G., Leach, D.N., Wang, Y. and Shewfelt, R.L.: Sulfur volatiles in *Cucumis melo* cv. *makdimon* (Muskmelon) aroma. Sensory evaluation by gas chromatography-olfactometry. *ACS Symp. Ser.*, 564 (1994) 36-48; *C.A.*, 121 (1994) 299609u.

See also 1256, 1468, 1522, 1525, 1581.

35. ENVIRONMENTAL ANALYSIS

35a. General papers and reviews

1749 Madden, A. and Lewis, E.T.: From purge and trap to ambient air. *Environ. Test. Anal.*, 4, No. 1 (1995) 56-57 and 89; *C.A.*, 122 (1995) 195388a.

See also 1597.

35b. Air pollution (complex mixtures; single compounds by cross-reference only)

1750 Chai, M. and Pawliszyn, J.: Analysis of environmental air samples by solid-phase microextraction and gas chromatography/ion trap mass spectrometry. *Environ. Sci. Technol.*, 29 (1995) 693-701.

1751 Davoli, E., Cappellini, L., Moggi, M. and Fanelli, R.: Automated high-speed analysis of selected organic compounds in urban air by online isotopic dilution cryofocusing gas chromatography/mass spectrometry. *J. Am. Soc. Mass Spectrom.*, 5 (1994) 1001-1007; *C.A.*, 122 (1995) 37519q.

1752 Milina, R., Tsatsova, D. and Pavlova, A.: (Analysis of volatile organic pollutants in air.) *Anal. Lab.*, 2 (1993) 104-108; *C.A.*, 122 (1995) 62727p.

1753 Scharf, J. and Sarafin, R.: Analysis of airborne substances by means of microwave thermal desorption. *J. High. Resolut. Chromatogr.*, 18 (1995) 253-258.

See also 1246, 1365, 1398, 1445, 1463, 1489, 1598, 1802, 1803, 1806.

35c. Water pollution (complex mixtures; single compounds by cross-reference only)

1754 Amalric, L., Guillard, C. and Pichat, P.: The GC-MS identification of some aliphatic intermediates from the TiO₂ photocatalytic degradation of dimethoxybenzenes in water. *Res. Chem. Intermed.*, 21 (1995) 33-46; *C.A.*, 122 (1995) 196432x.

1755 Tandom, R.K., Finlayson, P. and Adeloju, S.B.: Testing the waters. *Educ. Chem.*, 31 (1994) 159-161; *C.A.*, 122 (1995) 132189r.

1756 Tera, M. and Kitsuwu, T.: (Analysis of volatile organic compounds in water using purge and trap GC/MS without cryofocus technique.) *Kogyo Yosui*, 433 (1994) 63-68; *C.A.*, 122 (1995) 196317p.

1757 Wan, H.B., Chi, H., Wong, M.K. and Mok, C.Y.: Solid phase microextraction using pencil lead as sorbent for analysis of organic pollutants in water. *Anal. Chim. Acta*, 298 (1994) 219-223.

See also 1312, 1362, 1377, 1390, 1396, 1399, 1408, 1425, 1466, 1552, 1553, 1587, 1589, 1592, 1605, 1643, 1646, 1650, 1654, 1660, 1661, 1770.

35d. Soil pollution (complex mixtures; single compounds by cross-reference only)

1758 Camel, V., Tambuté, A. and Caude, M.: Influence of ageing on the supercritical fluid extraction of pollutants in soils. *J. Chromatogr. A*, 693 (1995) 101-111.

1759 Nouredin, S., Vandenbroucke, M. and Courtot, et P.: (Study of sedimentary humic matter by application of the pyrolysis-liquid chromatography-gas chromatography technique.) *Oceanol. acta*, 17 (1994) 43-53; *C.A.*, 122 (1995) 114147w.

See also 1221, 1330, 1358, 1367, 1397, 1408, 1413, 1414, 1425, 1426, 1517, 1606, 1629, 1642, 1652, 1784.

36. SOME TECHNICAL PRODUCTS AND COMPLEX MIXTURES

36a. Surfactants

1760 Almen, P. and Ericsson, I.: Quantification of sodium dodecyl sulfate in a two-phase system using filament-pulse-pyrolysis-gas chromatography. *Langmuir*, 11 (1995) 108-110; *C.A.*, 122 (1995) 164086w.

1761 Arzt, W.E. and Meyers, M.R.: Supercritical fluid extraction and chromatography of emulsifiers. *J. Am. Oil Chem. Soc.*, 72 (1995) 219-224; *C.A.*, 122 (1995) 185672z.

1762 Fendinger, N., Begley, W.M., McAvoy, D.C. and Eckhoff, W.S.: Measurement of alkyl ethoxylate surfactants in natural waters. *Environ. Sci. Technol.*, 29 (1995) 856-863.

1763 McPherson, B.P. and Rasmussen, H.T.: Chromatography of cationic surfactants: HPLC, TLC, and GLC. *Surfactant Sci. Ser.*, 53 (1994) 289-326; *C.A.*, 122 (1995) 136787g - a review with 89 refs.

1764 Nace, V.M. and Knoel, J.C.: Nonionic surfactant polarity index determination by inverse gas chromatography. *J. Am. Oil Chem. Soc.*, 72 (1995) 89-95; *C.A.*, 122 (1995) 109358m.

See also 1210.

36b. Antioxidants and preservatives

1765 Ali, S., Bhatt, B.D. and Nigam, R.N.: Determination of additives in polyolefines using megabore capillary gas chromatography. *Polym. Sci.*, 2 (1994) 922-926; *C.A.*, 122 (1995) 11080q.

1766 Duffield, P., Bourne, D., Tan, K., Garruto, R.M. and Duncan, M.W.: Analysis of the neurotoxic plasticizer *n*-butylbenzene-sulfonamide by gas chromatography combined with accurate mass selected ion monitoring. *J. Anal. Toxicol.*, 18 (1994) 361-368; *C.A.*, 122 (1995) 71104k.

See also 1588.

- 36c. *Complex mixtures, technical products and unidentified compounds*
- 1767 Carbini, M., Volpin, S. and Traldi, P.: Curie-point pyrolysis gas chromatography/mass spectrometry in the identification of paint media. *Org. Mass Spectrom.*, 29 (1994) 561-565; C.A., 122 (1995) 9023e.
- 1768 Carey, J.M. and Sutton, G.P.: Polyol ester synthetic refrigeration lubricant analysis by supercritical fluid chromatography. *Anal. Chem.*, 67 (1995) 1632-1636.
- 1769 Cassista, A.R. and Sandercock, P.M.L.: Comparison and identification of automotive topcoats: microchemical spot tests, microspectrophotometry, pyrolysis-gas chromatography, and diamond anvil cell FTIR. *J. - Can. Soc. Forensic Sci.*, 27 (1994) 209-223; C.A., 122 (1995) 2903h.
- 1770 Eschke, H.-D., Traud, J. and Dibowski, H.-J.: (Studies on the occurrence of polycyclic musk odors in different environmental compartments. 1st Communication: detection and analysis by GC/MS in surface and sewage waters and fish.) *Umweltwiss. Schadst.-Forsch.*, 6 (1994) 183-189; C.A., 122 (1995) 114425k.
- 1771 Lognay, G., Marlier, M., Seck, D., Haubruge, E., Wathelet, J.P., Coulibaly, A.D., Gaspar, C. and Severin, M.: (Head space sampling and GC-MS detection of the biocide molecules from *Boscia senegalensis* leaves.) *Bull. Rech. Agron. Gembloux*, 29, No. 1 (1994) 117-124; C.A., 122 (1995) 51369x.
- 1772 Mellon, F.A., Waldron, K.W., Selvendran, R.R. and Galletti, G.C.: The potential of pyrolysis/gas chromatography/mass spectrometry for determining the progress of lignification: an investigation of maturing asparagus sclerenchyma tissue. *Org. Mass Spectrom.*, 29, No. 10 (1994) 556-560; C.A., 122 (1995) 8313n.
- 1773 Nakashima, H., Miyano, N. and Matsunaga, I.: (Analysis of tributyl phosphate in flame retardant textiles by capillary gas chromatography.) *Jpn. J. Toxicol. Environ. Health*, 40 (1994) 461-466; C.A., 122 (1995) 12008j.
- 1774 Nali, M., Calemma, V. and Montanari, L.: Pyrolysis/gas chromatography/mass spectrometry of asphaltene fractions. *Org. Mass Spectrom.*, 29 (1994) 607-614; C.A., 122 (1995) 137788v.
- 1775 Nali, M., Corana, F., Montanari, L. and Pellegrini, L.: A pyrolysis-gas chromatography/mass spectrometry study on coals. *J. Anal. Appl. Pyrolysis*, 29 (1994) 15-23; C.A., 122 (1995) 852226p.
- 1776 Neer, L.A. and Deo, M.D.: Simulated distillation of oils with a wide carbon number distribution. *J. Chromatogr. Sci.*, 33 (1995) 133-138.
- 1777 Ohtani, H., Kato, T., Tsuge, S. and Yano, T.: Determination of rosin sizing agents in paper by pyrolysis-gas chromatography combined with online methylation. *Tappi J.*, 77 (1994) 177-183; C.A., 122 (1995) 190767p.
- 1778 Oono, A. and Inokawa, T.: (Quantitative analysis of pulverization agent in inorganic powder materials.) *Jpn. Kokai Tokkyo Koho JP 06,308,110 [94,308,110]* (Cl. G01/N30/88), 04 Nov. 1994, Appl. 93/94,333, 21 Apr. 1993; 8 pp.; C.A., 122 (1995) 95556k.
- 1779 Qi, Y., Wang, F., Liu, S. and Fan, Y.: (Rapid evaluation of anti-aging properties of asphalt by inverse gas-liquid chromatography.) *Shiyou Huagong*, 24 (1995) 44-49 C.A., 122 (1995) 140762n.
- 1780 Schubert, A.J.: Determining fractions of petroleum fuels by supercritical fluid chromatography. *U.S. US 5,345,029* (Cl. 585-825; C07C7/12), 06 Sep. 1994, Appl. 949,198, 22 Sep. 1992; 5 pp.; C.A., 121 (1994) 304424m.
- 1781 Senthil, V. and Soundararajan, N.: Determination of molecular structures of photolysis products of substituted aromatic azides in diethylamine solvent by gas chromatography/mass spectrometry. *Rapid Commun. Mass Spectrom.*, 8 (1994) 755-758; C.A., 121 (1994) 300525x.
- 1782 Smith D.L. and Piepmeier, E.H.: Multivariate approach to fingerprint identification of organic compounds using an oscillating glow discharge detector for gas chromatography. *Anal. Chem.*, 67 (1995) 1084-1091.
- 1783 Smolenskaya, N.Yu. and Sapozhnikov, Yu.M.: (Gas chromatographic analysis of transformer oil for air, water, oxygen and nitrogen.) *Elektr. Stn.*, 8 (1994) 34-37; C.A., 122 (1995) 137768p.
- 1784 Tavendale, M.H., Wilkins, A.L., Langdon, A.G., Mackie, K.L., Stuthridge, T.R. and McFarlane, P.N.: Analytical methodology for the determination of freely available bleached kraft mill effluent-derived organic constituents in recipient sediments. *Environ. Sci. Technol.*, 29 (1995) 1407-1414.
- 1785 Uematsu, Y., Hatooka, Y., Hirata, K., Hirokado, M. and Kazama, M.: (Determination of residual organic solvents in flavor by standard addition head-space gas chromatography.) *Shokuhin Eiseigaku Zasshi*, 35 (1994) 645-651; C.A., 122 (1995) 131288k.
- 1786 Van Leeuwen, J.A., Jonker, R.J. and Gill, R.: Octane number prediction based on gas chromatographic analysis with non-linear regression techniques. *Chemom. Intell. Lab. Syst.*, 25 (1994) 325-339.
- 1787 Ye, L., Luo, A. and Fu, R.: (The latest developments of the application of pyrolysis gas chromatography to explosives.) *Chin. J. Chromatogr.*, 13 (1995) 104-106.
- 1788 Young, R.J. and Fujinari, E.M.: Nitrogen detection for simulated distillation of refinery streams. *Int. Lab.*, 25, No. 2 (1995) 22-28.
- See also 1206, 1214, 1223, 1237, 1259, 1361, 1369, 1370, 1400, 1401, 1402, 1403, 1405, 1406, 1407, 1409, 1421, 1422, 1444, 1449, 1474, 1475, 1637, 1684, 1799.
37. CELLS, CELLULAR PARTICLES AND SUPRAMOLECULAR STRUCTURES
- See 1479, 1481, 1485, 1486, 1718.
38. INORGANIC COMPOUNDS
- 38b. Anions
- 1789 Cardeal, Z.L., Gallet, J.P., Astier, A. and Pradeau, D.: Cyanide assay: statistical comparison of a new gas chromatographic calibration method versus the classical spectrophotometric method. *J. Anal. Toxicol.*, 19 (1995) 31-34; C.A., 122 (1995) 74110v.

- 1790 Dennis, M.J., Burrell, A., Mathieson, K., Willetts, P. and Massey, R.C.: The determination of the flour improver potassium bromate in bread by gas chromatographic and ICP-MS methods. *Food Addit. Contam.*, 11 (1994) 633-639; C.A., 122 (1995) 79359t.
- 1791 Himata, K., Kuwahara, T., Ando, S. and Maruoka, H.: Determination of bromate in bread by capillary gas chromatography with a mass detector (GC/MS). *Food Addit. Contam.*, 11 (1994) 559-569; C.A., 122 (1995) 79343h.
- 1792 Van Kreel, B.K.: An improved bromide assay for the estimation of extracellular water volume by capillary gas chromatography. *Clin. Chim. Acta*, 231 (1994) 117-128.
- 38c. *Permanent and rare gases*
- 1793 Abe, M., Miyazaki, H., Nagai, Y., Nakajima, M. and Miyabe, M.: Determination of carbon monoxide in dark-fleshed fish by gas chromatography. *Nagoya-shi Eisei Kenkyushoho*, 40 (1994) 10-15; C.A., 122 (1995) 158857g.
- 1794 Czaplicka, M. and Kubica, K.: (A single step determination of coal pyrolysis gases.) *Karbo-Energochem.-Ekol.*, 38 (1993) 254-256; C.A., 122 (1995) 137860n.
- 1795 Goyette, B., Vigneault, C. and Raghavan, G.S.V.: Effect of argon on gas chromatographic analysis for controlled atmosphere storage. *Trans. ASAE*, 37 (1994) 1221-1224; C.A., 122 (1995) 54324q.
- 1796 Hiyama, T., Sugaya, S., Yamamoto, J. and Kamimura, K.: Determination of occluded gases in mixed oxide fuel pellets using gas sampling unit and gas chromatography. *J. Nucl. Mater.*, 218 (1995) 1-7; C.A., 122 (1995) 122061d.
- 1797 Lasa, J., Drozdowicz, B. and Korus, A.: Determination of the nitrogen concentration in argon on ppm level by gas chromatography with thermal conductivity detector. *Chem. Anal. (Warsaw)*, 39 (1994) 735-739;.
- 1798 Lazutkin, A.E., Sokolov, B.K. and Sorokin, F.A.: (Method for determination of carbon oxides in inert gases and air.) *U.S.S.R. SU 1,814,061* (Cl. G01N30/08), 07 May 1993, Appl. 4,938,231, 12 Apr. 1991; C.A., 122 (1995) 95547h.
- 1799 Levin, M.D., Ellis, C., Springfield, J. and Engel, R.R.: Carbon monoxide generation from hydrocarbons at ambient and physiological temperature: a sensitive indicator of oxidant damage? *J. Chromatogr. A*, 695 (1995) 324-328.
- 1800 Van Dam, J. and Daenens, P.: Microanalysis of carbon monoxide in blood by head-space capillary gas chromatography. *J. Forensic Sci.*, 39 (1994) 473-478; C.A., 122 (1995) 48597q.
- 1801 Yoshino, N., Nii, Y. and Hirata, T.: (A separation system of nitrogen, oxygen, carbon dioxide, and argon by capillary gas chromatography and its application to in-package atmosphere. *Nippon Shokuhin Kogyo Gakkaishi*, 41 (1994) 865-870; C.A., 122 (1995) 158846c.
- 38d. *Volatile inorganic compounds*
- 1802 Arah, J.R.M., Crichton, I.J., Smith, K.A., Clayton, H. and Skiba, U.: Automated gas chromatographic analysis system for micrometeorological measurements of trace gas fluxes. *J. Geophys. Res., [Atmos]*, 99, No. D8 (1994) 16593-16598; C.A., 121 (1994) 314755j.
- 1803 Borossay, J., Torkos, K. and Bakos, E.: Gas chromatographic CO₂/O₂ determinations in manholes of a telephone network. *Microchem. J.*, 51 (1995) 122-126; C.A., 122 (1995) 195407f.
- 1804 Brunner, U., Chasteen, T.G., Ferloni, P. and Bachofen, R.: Chromatographic determination of phosphine (PH₃) and hydrogen sulfide (H₂S) in the headspace of anaerobic bacterial enrichments using flame photometric detection. *Chromatographia*, 40 (1995) 399-403.
- 1805 Cheng, J., Yin, B., Lu, D. and Cha, H.: (Method with bacterial culture and headspace gas chromatography for detecting environmental toxicants.) *Huanjing Kexue*, 15, No. 3 (1994) 71-79; C.A., 122 (1995) 180406f.
- 1806 DeAndrade, J.B., Araujo, F.R.J. and de Aragao, N.M.: (Determination of nitric acid vapor in air.) *Quim. Nova*, 18, No. 1 (1995) 5-10; C.A., 122 (1995) 168516j.
- 1807 Munar, M., Maurice, M.-J. and Kluessendorf, A.: Quantitation of sulfur dioxide residues in malt and beer by headspace gas chromatography. *J. Am. Soc. Brew. Chem.*, 52 (1994) 168-171; C.A., 122 (1995) 54376h.
- 1808 Semiletov, I.P., Pozdnyakov, N.V. and Kosolapkin, G.Yu.: (Method of determination of the composition of gas inclusions in ice.) *U.S.S.R. SU 1,809,380* (Cl. G01N30/06), 15 Apr. 1993, Appl. 4,654,383, 23 Dec. 1988; C.A., 122 (1995) 122180s.
- See also 1215, 1420, 1796.
39. RADIOACTIVE AND OTHER ISOTOPE COMPOUNDS
- 1809 Fujioka, J., Chiba, K. and Kayane, M.: (Product analysis in reaction involving carbon-13 isotope ratio variation.) *Jpn. Kokai Tokkyo Koho JP 06,265,524*, [94,265,524] (Cl. G01N27/62), 22 Sep. 1994, Appl. 92/75,855, 28 Feb. 1992; 7 pp.; C.A., 122 (1995) 122172r.
- 1810 Merritt, D.A. and Hayes, J.M.: Nitrogen isotopic analyses by isotope-ratio-monitoring gas chromatography/mass spectrometry. *J. Am. Soc. Mass Spectrom.*, 5, No. 5 (1994) 387-397; C.A., 121 (1994) 314910f.
- 1811 Ricapito, I., Malara, C., Pierini, G. and Spelta, B.: Near real-time diagnostic analysis of hydrogen isotopes by gas chromatography. *Process Technol. Proc.*, 11 (1994) 465-473; C.A., 122 (1995) 22680m.
- See also 1506, 1569, 1616.
- See also 1215, 1352, 1353, 1803, 1811.

Planar Chromatography

1. REVIEWS AND BOOKS

- 404 Nicholson, J.: Paperwork made easy. *Chem. Br.*, 30 (1994) 658-660; *C.A.*, 122 (1995) 204153z - a review with 13 refs.
 405 Volynets, M.P.: (*Quantitative Thin-Layer Chromatography in Inorganic Analysis*). Nauka, Moscow, 1993, 240 pp.; *C.A.*, 121 (1994) 220574x.

See also 443, 445, 468, 473, 558, 641.

2. FUNDAMENTALS, THEORY AND GENERAL

2a. General

- 406 Hauck, H.E., Junker-Buchheit, A., Weinig, R. and Schoemer, S.: (Good laboratory practice (GLP) in chromatography). *Labor-Praxis*, 70-72 (1995) 109-110; *C.A.*, 122 (1995) 204170c.
 407 Huynh, T.K.X., Lederer, M. and Leipzig-Pagani, E.: Adsorption chromatography on cellulose. XIII. General effects of aqueous solutions of α -cyclodextrin as eluent. *J. Chromatogr. A*, 695 (1995) 155-159.
 408 Huynh, T.K.X., Lederer, M. and Leipzig-Pagani, E.: Adsorption chromatography on cellulose. XIII. Chromatography with aqueous solutions of carbohydrates as eluents. *J. Chromatogr. A*, 695 (1995) 160-164.
 409 Malinowska, I., Rózyło, J.K. and Gumieniak, A.: A simple method for optimizing multi-component mobile phases in adsorption TLC. *J. Planar Chromatogr.*, 8 (1995) 23-32.

2b. Thermodynamics and theoretical relationships

- 410 Merkkü, P., Ylirussi, J. and Vuorela, H.: Evaluation of diffusion and evaporation of ternary mobile phases for TLC. *J. Planar Chromatogr.*, 8 (1995) 112-116.

2c. Relationship between structure and chromatographic behaviour

- 411 Pyka, A.: Topological indices and R_M values of isomeric phenol derivatives in structure - biological activity studies: Part VI. *J. Planar Chromatogr.*, 8 (1995) 52-62.
 412 Waksmundzka-Hajnos, M., Zareba, S. and Chabros, A.: Retention behavior of some heteroazophenols in normal phase systems of TLC. I. Comparison of adsorption properties of Florisil and silica. *Chem. Anal. (Warsaw)*, 39 (1994) 121-132; *C.A.*, 122 (1995) 143531x.

See also 413, 584, 595.

2d. Measurement of physico-chemical and related values

- 413 Woolston, C.R.J., Lee, J.B. and Swinbourne, F.J.: Partition coefficients and chromatographic parameters of some substituted 1,3-thiazolidin-4-ones. *Indian J. Chem., Sect. B.: Org. Chem. Incl. Med. Chem.*, 33b (1994) 590-592; *C.A.*, 121 (1994) 34737c.

See also 433.

3. GENERAL TECHNIQUES

3a. Apparatus and accessories

- 414 Bharati, S., Roistrum, G.A. and Loeberg, R.: Calibration and standardization of latroscan (TLC-FID) using standards derived from crude oils. *Org. Geochem.*, 22 (1994) 835-862; *C.A.*, 122 (1995) 164965g.
 415 Flodberg, G. and Roeraade, J.: A device for high pressure thin layer chromatography. *J. Planar Chromatogr.*, 8 (1995) 10-13.

See also 581.

3b. Detectors and detection reagents

- 416 Klaus, R., Fischer, W. and Hauck, H.E.: Reagent-free detection of substances on amino-modified silica gel layers - thin-layer chromatographic separation and detection of fruit acids. *LC-GC Int.*, 8, No. 3 (1995) 151-156.
 417 Navas, D.A. and Garcia, S.F.: Fiber optic fluorescence scanning in thin-layer chromatography. *Instrum. Sci. Technol.*, 22 (1994) 273-281; *C.A.*, 122 (1995) 204180f.
 418 Roth, E. and Kiefer, W.: Surface-enhanced Raman spectroscopy as a detection method in gas chromatography. *Appl. Spectrosc.*, 48 (1994) 1193-1195; *C.A.*, 122 (1995) 45395m.
 419 Wardas, W., Pyka, A. and Jedrzejczak, M.: Visualizing agents for aromatic carboxylic acids in TLC. *J. Planar Chromatogr.*, 8 (1995) 148-151.
 420 Wintersteiger, R., Barary, M.H., El-Yazbi, F.A., Sabry, S.M. and Wahbi, Abdel-Aziz, M.: Fluorimetric determination of morphine in codeine. *Spectosc. Lett.*, 27 (1994) 1447-1460; *C.A.*, 122 (1995) 38974w.

See also 466, 472, 563, 567, 569, 574, 576, 588.

3c. Sorbents and columns, packing procedures

- 421 Casale, S.F.J.: (Manufacture of abrasion- and cracking-resistant plates and films for thin-layer chromatography). *Span. ES 2,052,444 (Cl. B01D15/08)*, 01 Jul. 1994, Appl. 9,202,416, 30 Oct. 1992; 8 pp.; *C.A.*, 122 (1995) 136905u.

- 422 Hajouj, Z., Thomas, J. and Siouffi, A.M.: Two-dimensional TLC on mechanically blended silica-based bonded phases. Evaluation of the behaviors of C8-diol mixtures using polynuclear aromatic hydrocarbons and comparison with C-18 cyano mixtures. *J. Liq. Chromatogr.*, 18 (1995) 887-894.
- 423 Junker-Buchheit, A., Hauck, H.E. and Wenig, R.: Environmentally friendly aluminum-backed TLC sheets. *Int. Lab.*, 25, No. 3 (1995) 121-12U.
- 424 Marutoiu, C., Constantiescu, R. and Dogar, F.: (Characterization of silica gel chemically bonded to an R-NH₂ support used in thin-layer chromatography). *Rev. Chim. (Bucharest)*, 45 (1994) 600-603; *C.A.*, 122 (1995) 150555m.
- 425 Mayer, M.L., Poole, C.F. and Henry, M.P.: Sampling characteristics of octadecylsiloxane-bonded silica particle-embedded glass fiber discs for solid-phase extraction. *J. Chromatogr. A*, 695 (1995) 267-277.
- 426 Okamoto, Y.: Thin layer chromatography for separating optical isomers. *Jpn. Kokai Tokkyo Koho JP 05,232,099* [93,232,099] (Cl. G01N30/93), Appl. 92/37,711, 25 Feb. 1992, 5 pp.; *C.A.*, 120 (1994) 49568b.
- 427 Rozylo, J.K., Gwis-Chomicz, D. and Malinowska, I.: Possibility of using chitin support in chromatographic control of environment pollution. *Environ. Prot. Eng.*, 16 (1990, Publ. 1992) 121-130; *C.A.*, 120 (1994) 260038m.

See also 406, 439, 473, 516, 523, 566, 639.

3e. Preparative scale chromatography

See 502.

3f. Programmed temperature, pressure, vapors, gradients

- 428 Soczewinski, E. and Matysik, G.: Method of an apparatus for producing a step-wise gradient of eluted composition in thin-layer chromatography. *Pol. PL 161,007* (Cl. G01N30/90), 31 May 1993, Appl. 268, 134, 07 Oct. 1987; 4 pp.; *C.A.*, 122 (1995) 204270k.

3g. High performance procedures

- 429 Chauhan, B.L., Mitra, S.K., Mohan, A.R., Gopumadhavan, S. and Anturlikar, S.D.: Development of high performance thin layer chromatography (HPTLC) fingerprint technique and bioassay to establish the shelf-life of a proprietary herbal formulation (PHF), an Ayurvedic formulation containing multiple ingredients of herbal origin. *Indian Drugs*, 31 (1994) 333-338; *C.A.*, 122 (1995) 38956s.

See also 431, 542, 606.

4. SPECIAL TECHNIQUES

4b. Computerization and modelling

See 432.

- 4c. *Combination with other physico-chemical techniques (MS, IR etc.)*

- 430 Fanibanda, T., Milnes, J. and Gormally, J.: Thin layer chromatography-mass spectrometry using infrared laser desorption. *Int. J. Mass Spectrom. Ion Processes*, 140 (1994) 127-132; *C.A.*, 122 (1995) 64550z.
- 431 Koglin, E., Kramer, H., Sawatski, J., Lehner, C. and Hellman, J.: *In situ* identification of high-performance thin-layer chromatography spots by FT-surface-enhanced Raman scattering. *Proc. SPIE-Int. Soc. Opt. Eng.*, 2089(9th International Conference on Fourier Transform Spectroscopy) (1993) 304-305; *C.A.*, 121 (1994) 194619u.

See also 418, 471, 494, 510, 532, 536, 576, 598, 640.

4f. Trace analysis and preseparation techniques

- 432 Matyska, M.T., Siouffi, A.-M. and Volpe, N.: Computer-aided optimization of a sample clean-up procedure - application to nitrosamines and amines. *J. Planar Chromatogr.*, 8 (1995) 39-46.

4g. Enantiomers, separation

See 426.

4h. Other special techniques

- 433 Dross, K. and Sonntag, C.: The estimation of physicochemical properties by RP₁₈-TLC. In: Wermuth, C.-G. (Editor), *Trends QSAR Mol. Modell.* 92, *Proc. Eur. Symp. Struct.-Act. Relat.: QSAR Mol. Moled.*, 9th 1992, ESCOM, Leiden, 1993, pp. 370-372; *C.A.*, 121 (1994) 157039j.

See also 406, 494, 568.

5. HYDROCARBONS AND HALOGEN DERIVATIVES

5a. Aliphatic hydrocarbons

- 434 Friedbacher, E.C. and Schindlbauer, H.: Quantitative evaluation of bitumen analysis by TLC/FID. *Bitumen*, 56 (1994) 105-108; *C.A.*, 122 (1995) 85193a.

5b. Cyclic hydrocarbons, fullerenes

See 422, 425, 434.

5d. Complex hydrocarbon mixtures (incl. analysis of tars, bitumens and mineral oils)

- 435 Dhole, V.R. and Ghosal, G.K.: Adulteration of petrol with kerosene: detection and semiquantitative determination by TLC: Part II. *J. Planar Chromatogr.*, 8 (1995) 80-81.

- 436 Lian, H.J., Lee, C. and Yen, T.F.: Fractionation of asphalt by thin-layer chromatography interfaced with flame ionization detector (TLC-FID) and subsequent characterization by FTIR. In: Sharma, M.K. and Yen, T.F. (Editors), *Asphaltene Part. Fossil Fuel Explor., Recovery, Refin., Prod. Processes, [Proc. Int. Symp.]*, 1992, Plenum, New York, 1994, pp. 229-238; C.A., 122 (1995) 137783q.
6. ALCOHOLS
- 437 Yao, Y.: (Comparative study on yinqiao detoxicating decoction in new convenient and old traditional forms). *Zhongguo Zhongyao Zazhi*, 19 (1994) 60406; C.A., 122 (1995) 64098h.
- See also 446, 455, 517.
7. PHENOLS
- 438 Baranowska, I. and Skotniczna, A.: Behaviour of phenols in normal and reversed-phase thin-layer chromatography. *Chromatographia*, 39 (1994) 564-568.
- 439 Hinze, W.L., Feng, L., Moreno, B., Quina, F.H., Suzuki, Y. and Wang, H.: Utilization of micelle-mimetic intramolecular ionene aggregates as the mobile phase in pseudophase thin-layer liquid chromatography. *Anal. Sci.*, 11 (1995) 183-187; C.A., 122 (1995) 229738g.
- See also 411, 412, 425.
8. SUBSTANCES CONTAINING HETEROCYCLIC OXYGEN
- 8a. *Flavonoids*
- 440 Fater, Z., Samu, Z. and Kernoczi, Z.: (Identification of the flavonoid compounds of DIURO drops based on retention data and UV/VIS spectra). *Acta Pharm. Hung.*, 64 (1994) 193-198; C.A., 122 (1995) 170376g.
- See also 626.
- 8b. *Aflatoxins and other mycotoxins*
- 441 Grabarkiewicz-Szczesna, J.: (Analysis of alternariotoxins by thin-layer chromatography method). *Rocz. Akad. Roln. Poznaniu*, 256 (1993) 17-23; C.A., 122 (1995) 153464s.
- 442 Milat, M.-L. and Blein, J.-P.: *Cercospora beticola* toxins. III. Purification, thin-layer and high-performance liquid chromatographic analyses. *J. Chromatogr. A*, 699 (1995) 277-283.
- 443 Nawaz, S., Coker, R.D. and Haswell, S.J.: HPTLC - a valuable chromatographic tool for the analysis of aflatoxins. *J. Planar Chromatogr.*, 8 (1995) 4-9 - a review with 69 refs.
- 444 Stockenstrom, S., Sydenham, E.W. and Thiel, P.G.: Determination of fumonisins in corn: evaluation of two purification procedures. *Mycotoxin Res.*, 10 (1994) 9-14; C.A., 122 (1995) 79337j.
- 445 Trucksess, M.W.: Mycotoxins. *J. Assoc. Off. Anal. Chem.*, 78 (1995) 135-141 - a review with 120 refs.
- 8c. *Other compounds with heterocyclic oxygen (incl. tannins)*
- See 560, 561, 616.
9. OXO COMPOUNDS, ETHERS, EPOXIDES AND QUINONES
- 446 Baranowska, I., Maslinska-Solich, J. and Swierczek, S.: TLC investigation of copoly(imino-acetals) from terephthaldehyde and aliphatic amino-diols. *Acta Chromatogr.*, 2 (1993) 56-54; C.A., 122 (1995) 162290j.
- 447 Onodera, S., Uchida, A. and Nakajima, C.: Thin-layer chromatographic fractionation of Ames mutagenic chlorosemiquinone in chlorinated 4-methylphenol solution. *J. Chromatogr. A*, 699 (1995) 291-296.
- See also 425, 442, 455, 456, 479.
10. CARBOHYDRATES
- 10a. *Mono and oligosaccharides. Structural studies*
- 448 Neville, D.C.A., Field, R.A. and Ferguson, M.A.J.: Hydrophobic glycosides of N-acetylglucosamine can act as primers for polylactosamine synthesis and can affect glycolipid synthesis *in vivo*. *Biochem. J.*, 307 (1995) 791-797.
- 449 Rodríguez-Aparicio, L.B., Ferrero, M.A. and Reglero, A.: N-Acetyl-D-neuraminic acid synthesis in *Escherichia coli* K1 occurs through condensation of N-acetyl-D-mannosamine and pyruvate. *Biochem. J.*, 308 (1995) 501-505.
- 450 Sato, T., Takio, K., Kobata, A., Greenwalt, D.E. and Furukawa, K.: Site-specific glycosylation of bovine butyrophilin. *J. Biochem. (Tokyo)*, 117 (1995) 147-157.
- 451 Vandersall, A.S., Cameron, R.G., Nairn, C.J., III, Yelenosky, G. and Wodzinski, R.J.: Identification, characterization, and partial purification of glucoamylase from *Aspergillus niger* (syn. *A. ficum*) NRRL 3135. *Prepar. Biochem.*, 25 (1995) 29-55.
- See also 483, 485, 639.
- 10b. *Polysaccharides, mucopolysaccharides, lipopolysaccharides*
- 452 Mourao, P.A.S. and Assreuy, A.-M.: Trehalose as a possible precursor of the sulfated L-galactan in the ascidian tunic. *J. Biol. Chem.*, 270 (1995) 3132-3140.
- 10c. *Glycoproteins and their constituents*
- 453 Treumann, A., Lively, M.R., Schneider, P. and Ferguson, M.A.J.: Primary structure of CD52. *J. Biol. Chem.*, 270 (1995) 6088-6099.

11. ORGANIC ACIDS AND LIPIDS

11a. Organic acids and simple esters

- 454 Bieganowska, M.L., Petruczyński, A. and Glowński, K.: Reversed phase retention behavior of β -cyclodextrin complexes of phenolic and cinnamic acids. *J. Planar Chromatogr.*, 8 (1995) 63-68.
- 455 Blee, E. and Schuber, F.: Stereocontrolled hydrolysis of the linoleic acid monoepoxide regioisomers catalyzed by soybean epoxide hydrolase. *Eur. J. Biochem.*, 230 (1995) 229-234.
- 456 Demini, P., Reynaud, D. and Pace-Assiac, C.R.: Extractive derivatization of the 12-lipoxygenase products, hepoxilins and related compounds into fluorescent anthryl esters for their complete high-performance liquid chromatography profiling in biological systems. *Anal. Biochem.*, 226 (1995) 252-255.
- 457 Kondo, Y., Hashidoko, Y. and Mizutani, J.: An enzymatic formation of 13-oxo-trideca-9,11-dienoic acid from 13-hydroperoxylinolenic acid by a homolytic hydroperoxide lyase in elicitor-treated soybean cotyledons. *Biochim. Biophys. Acta*, 1255 (1995) 9-15.

See also 416, 419, 481, 483, 507, 597, 606, 637, 640.

11b. Prostaglandins

- 458 Capote, R., Francisco, G., González, J.A. and Nuñez, A.J.: Quantitative analysis of (15S)-15-methylprostaglandin F₂ α methyl ester by high performance thin layer chromatography. *J. Planar Chromatogr.*, 8 (1995) 155-156.

11c. Lipids and their constituents

- 459 Alter, C.A. and Wolf, B.A.: Identification of phosphatidylinositol 3,4,5-trisphosphate in pancreatic islets and insulin-secreting β -cells. *Biochem. Biophys. Res. Commun.*, 208 (1995) 190-197.
- 460 Alvarez, J.G., Slomovic, B. and Ludmir, J.: Analysis of dipalmitoyl phosphatidylcholine in amniotic fluid by enzymatic hydrolysis and high-performance thin-layer chromatography reflectance spectrodensitometry. *J. Chromatogr. B*, 665 (1995) 79-87.
- 461 Bouhours, D., Hansson, G.C. and Bouhours, J.-F.: Structure and genetic polymorphism of blood group A-active glycosphingolipids of the rat large intestine. *Biochim. Biophys. Acta*, 1255 (1995) 131-140.
- 462 Buche, S.D., Patil, M.N. and Jaiswal, P.K.: Detection of mahua oil in groundnut and sesame oils. *J. Oil Technol. Assoc. India (Bombay)*, 26 (1994) 83; *C.A.*, 122 (1995) 79339m.
- 463 Cao, S.G. and Hatch, G.M.: Cardiolipin is synthesized from exogenous phosphatidylglycerol in rat heart. *Biochim. Biophys. Acta*, 1256 (1995) 97-102.
- 464 Carsberg, C.J., Ohanian, J. and Friedmann, P.S.: Ultraviolet radiation stimulates a biphasic pattern of 1,2-diacylglycerol formation in cultured human melanocytes and keratinocytes by activation of phospholipases C and D. *Biochem. J.*, 305 (1995) 471-477.
- 465 Choi, B.O., Yamaki, T., Ibayashi, Y., Maeda, Y., Gasa, S. and Hashi, K.: Interleukin 4 enhances ganglioside GD3 expression on the human fibroblast cell line WI-38. *J. Biochem. (Tokyo)*, 117 (1995) 315-320.

- 466 Davani, B. and Olsson, N.U.: Detection of natural galactolipids by thin layer chromatography with 8-anilino-1-naphthalenesulfonate (ANS) as fluorogenic visualization reagent. *J. Planar Chromatogr.*, 8 (1995) 33-35.
- 467 El Bawab, S., Macovschi, O., Lagarde, M. and Prigent, A.-F.: Time-course changes in content and fatty acid composition of phosphatidic acid from rat thymocytes during concanavalin A stimulation. *Biochem. J.*, 308 (1995) 113-118.
- 468 Firestone, D.: Fats and oils. *J. Assoc. Off. Anal. Chem.*, 78 (1995) 150-155 - a review with 26 refs.
- 469 Fujiwaki, T., Hamanaka, S., Tate, S., Inagaki, F., Suzuki, M., Suzuki, A. and Mori, C.: Tissue accumulation of sulfatide and GM₃ ganglioside in a patient with variant Farber disease. *Clin. Chim. Acta*, 234 (1995) 23-36.
- 470 Fukami, M.H. and Holmsen, H.: Diacylglycerol elevations in control platelets are unaccompanied by pleckstrin phosphorylation. Implications for the role of diacylglycerol in platelet activation. *Eur. J. Biochem.*, 228 (1995) 579-586.
- 471 Goto, M., Kato, M., Asaumi, M., Shirai, K. and Venkateswaran, K.: TLC/FID method for evaluation of the crude oil-degrading capability of marine microorganisms. *J. Mar. Biotechnol.*, 2 (1994) 45-50; *C.A.*, 122 (1995) 127786d.
- 472 Gracheva, N.Y., Pchelkin, V.P. and Tsydendambaev, V.D.: Method for determining boundaries of chromatographic zones of individual classes of phospholipids when their mixture is separated by thin-layer chromatography. *U.S.S.R. SU 1,778,684*, (Cl. GO1N30/90), 30 Nov. 1992, Appl. 4,872,327, 10 Oct. 1990; *C.A.*, 120 (1994) 49567a.
- 473 Henderson, R.J. and Tocher, D.R.: Thin-layer chromatography. *Lipid Anal.*, (1992) 63-111; *C.A.*, 122 (1995) 127735m - a review with 63 refs.
- 474 Hung, C.-Y., Ko, Y.-G. and Thompson, G.A., Jr.: Temperature induced alteration of inositolphosphorylceramides in the putative glycosylated lipid precursors of *Tetrahymena mimbres* glycosylphosphatidylinositol-anchored proteins. *Biochem. J.*, 307 (1995) 107-113.
- 475 Itonori, S., Shirai, T., Kiso, Y., Oha, Y., Sjiota, K. and Ogawa, T.: Glycosphingolipid composition of rat placenta: changes associated with stage of pregnancy. *Biochem. J.*, 307 (1995) 399-405.
- 476 Jalink, K., Hengeveld, T., Mulder, S., Postma, F.R., Simon, M.-F., Chap, H., van der Marel, G.A., van Boom, J.H., van Blitterswijk, W.J. and Moolenaar, W.H.: Lysophosphatidic acid-induced Ca²⁺ mobilization in human A431 cells: structure-activity analysis. *Biochem. J.*, 307 (1995) 609-616.
- 477 Ko, Y.G., Hung, C.-Y. and Thompson, G.A., Jr.: Temperature regulation of the *Tetrahymena mimbres* glycosylphosphatidylinositol-anchored protein lipid composition. *Biochem. J.*, 307 (1995) 115-121.
- 478 Krasil'nikova, M.A., Shatskaya, V.A. and Shtutman, M.S.: (Regulation of phospholipid turnover in hamster fibroblasts transformed by oncogenes v-src and N-ras). *Biokhimiya (Moscow)*, 59 (1994) 1766-1774.
- 479 Kuksis, A., Myher, J.J., Marai, L. and Geher, K.: Analyses of hydroperoxides and core aldehydes of triacylglycerols. In: Maikki, Y. and Lambertsen, G. (Editors), *Nord. Lipid. Symp. Proc.*, 17th, Scand. Forum Lipid Res. Technol., Bergen, 1993, pp. 630-638; *C.A.*, 122 (1995) 8268b.

- 480 Leenders, R.G.G., de Jong, J.G.N. and Wevers, R.A.: Extraction and purification of gangliosides from plasma and fibroblasts before analysis by thin layer chromatography. *Ann. Clin. Biochem.*, 32 (1995) 68-73; *C.A.*, 122 (1995) 182637f.
- 481 Lomnitski, L., Sklan, D. and Grossman, S.: Lipoygenase activity in rat dermis and epidermis: partial purification and characterization. *Biochim. Biophys. Acta*, 1255 (1995) 351-359.
- 482 Michova, V.: (Analysis of individual phospholipids in blood serum by thin layer and gas chromatography). *Anal. Lab.*, 2 (1993) 60-63; *C.A.*, 122 (1995) 4620n.
- 483 Muething, J. and Unland, F.: A comparative assessment of TLC overlay technique and microwell adsorption assay in the examination of influenza A and Sendai virus specificities towards oligosaccharides and sialic acid linkages of gangliosides. *Glycoconjugate J.*, 11 (1994) 486-492; *C.A.*, 122 (1995) 127977s.
- 484 Persat, F., Bouhours, J.-F., Petavy, A.-F. and Mojon, M.: Free ceramides of *Echinococcus multilocularis* metacestodes. *Biochim. Biophys. Acta*, 1255 (1995) 280-284.
- 485 Proudfoot, L., Schneider, P., Ferguson, M.A.J. and McConville, M.J.: Biosynthesis of the glycolipid anchor of lipophosphoglycan and the structurally related glycoinositolphospholipids from *Leishmania major*. *Biochem. J.*, 308 (1995) 45-55.
- 486 Qanbar, R. and Possmayer, F.: On the surface activity of surfactant-associated protein C (SP-C): effects of palmitoylation and pH. *Biochim. Biophys. Acta*, 1255 (1995) 251-259.
- 487 Ridgway, N.D. and Merriam, D.L.: Metabolism of short-chain ceramide and dihydroceramide analogues in Chinese hamster ovary (CHO) cells. *Biochim. Biophys. Acta*, 1256 (1995) 57-70.
- 488 Schmid, P.C., Deli, E. and Schmid, H.H.O.: Generation and remodeling of phospholipid molecular species in rat hepatocytes. *Arch. Biochem. Biophys.*, 319 (1995) 168-176.
- 489 Seibel, W.: (Determination of soft wheat in Durum wheat semolina and pasta. Part 4. Detection by thin-layer chromatography in high-temperature-dried pasta). *Getreide, Mehl Brot*, 47 (1993) 40-44; *C.A.*, 122 (1995) 131261w.
- 490 Seyama, Y., Otsuka, H., Ohashi, K., Vivien-Roels, B. and Pevet, P.: Sexual dimorphism of lipids in Harderian glands of golden hamsters. *J. Biochem. (Tokyo)*, 117 (1995) 661-670.
- 491 Sharom, F.J., Yu, X., Chu, J.W.K. and Doige, C.A.: Characterization of the ATPase activity of P-glycoprotein from multidrug-resistant Chinese hamster ovary cells. *Biochem. J.*, 308 (1995) 381-390.
- 492 Smith, M.C., Webster, C.L., Sherma, J. and Fried, B.: Determination of neutral lipids in regular and lowfat eggs by high performance TLC with densitometry. *J. Liq. Chromatogr.*, 18 (1995) 527-535.
- 493 Sugiyama, T., Kudo, N., Ojima, T., Mabuchi-Itoh, K., Yamashita, A. and Waku, K.: Coenzyme A-dependent cleavage of membrane phospholipids in several rat tissues: ATP-independent acyl-CoA synthesis and the generation of lysophospholipids. *Biochim. Biophys. Acta*, 1255 (1995) 167-176.
- 494 Taki, T., Ishikawa, D., Habda, S. and Kasama, T.: Direct mass spectrometric analysis of glycosphingolipid transferred to a polyvinylidene difluoride membrane by thin layer chromatography blotting. *Anal. Biochem.*, 225 (1995) 24-27.
- 495 Tarandjiiska, R., Marekov, I., Nikolova-Damyanova, B. and Amidzhin, B.: Determination of molecular species of triacylglycerols from highly unsaturated plant oils by successive application of silver ion and reversed phase TLC. *J. Liq. Chromatogr.*, 18 (1995) 859-872.
- 496 Watanabe, K. and Nishiyama, M.: A new type of two-dimensional thin-layer chromatography (TLC-mapping) for analysis of acidic glycosphingolipid molecular species. *Anal. Biochem.*, 227 (1995) 195-200.
- 497 Yamada, S., Araki, S., Abe, S., Kon, K., Ando, S. and Satake, M.: Structural analysis of a novel triphosphoglycosphingolipid from the egg of the sea hare, *Aplysia kurodai*. *J. Biochem. (Tokyo)*, 117 (1995) 794-799.

See also 448.

12. ORGANIC PEROXIDES

See 456, 479.

13. STEROIDS

13d. Sterols

See 489.

13f. Ecdysones and other insect steroid hormones

498 Kálász, H., Báthori, M. and Máthé, I.: Displacement thin-layer chromatography of some ecdysteroids. *J. Liq. Chromatogr.*, 18 (1995) 837-848.

13g. Other steroids

499 Moretti, G., Citti, G., Cammarata, P., Ferretti, E., Fiori, M. and di Giovannandrea, R.: (Residue analysis of urinary trenbolone in cattle by thin-layer chromatography and gas chromatography-mass spectrometry). *Riv. Sci. Aliment.*, 23 (1994) 227-232; *C.A.*, 122 (1995) 29974u.

See also 627.

14. STEROID GLYCOSIDES AND SAPONINS

500 Fujii, Y., Ikeda, Y. and Yamazaki, M.: High-performance liquid chromatographic determination of lanatosides in *Digitalis lutea* and *Digitalis ambigua* leaves. *J. Liq. Chromatogr.*, 17 (1994) 4451-4461.

501 Liang, S., Zhang, G., Li, Y. and Zheng, X.: (Determination of magnolol, honokiol, chlorogenic acid and *Gardenia* glycosides in qingganlian injection by TLC-densitometry). *Yaowu Fenxi Zazhi*, 15 (1995) 24-26; *C.A.*, 122 (1995) 222988e.

15. TERPENES AND OTHER VOLATILE AROMATIC COMPOUNDS

15a. Terpenes

- 502 Abdallah, O.M. and Ibraheim, Z.Z.: Sesquiterpene lactones from *Pluchea dioscoridis*. *Pharmazie*, 50 (1995) 277-278.
- 503 Nowak, G.: Isolation and chromatography of 15-deoxyrepin and 25 other sesquiterpene lactones from *Centaurea bella*. *Acta Soc. Bot. Pol.*, 62, No. 1-2 (1993) 33-36; *C.A.*, 122 (1995) 89205d.
- 504 O'Mathuna, D.P. and Doskotch, R.W.: New labdane diterpene glycosides from *Amphiachyris amoena*. *J. Natural Prod.*, 58 (1995) 82-92.
- 505 Rossini, C., Pandolfi, É., Dellacassa, E. and Moyna, P.: Determination of 1,8-cineole in *Eucalyptus* essential oils by thin-layer chromatography and densitometry. *J. Assoc. Off. Anal. Chem.*, 78 (1995) 115-117.

15b. Essential oils

- 506 Faruq, M.O., Chowdhury, J., Yusuf, M., Farooque, S.M., Chowdhury, M.M. and Khuda, M.M.: TLC technique in the component characterization and quality determination of Bangladeshi lemongrass oil (*Cymbopogon citratus* Stapf.). *Bangladesh J. Sci. Ind. Res.*, 29, No. 2 (1994) 27-38; *C.A.*, 122 (1995) 196519f.
- 507 He, Z., Liu, Y., Li, J. and Cheng, Y.: (Quality specification of compound Xisojin liquid). *Zhongguo Yiyuan Yaoxue Zazhi*, 14 (1994) 306-308; *C.A.*, 122 (1995) 38679d.
- 508 Zekovic, Z., Pekic, B., Lehojevic, Z. and Petrovic, L.: Chromatography in our investigations of camomile (*Matricaria chamomilla* L.). *Chromatographia*, 39 (1994) 587-590.

See also 505.

16. NITRO AND NITROSO COMPOUNDS

- 509 Chen, Z.: (TLC determination of N-dimethylnitrosamine and N-diethylnitrosamine in soy sauce). *Zhongguo Tiaoweipin*, 11 (1994) 27-30; *C.A.*, 122 (1995) 185694h.
- 510 Juskiewicz, T., Posyniak, A., Semeniuk, S. and Niedzielska, J.: Determination of 5-nitrofurans in meat by high-performance thin layer and liquid chromatography. In: Haagsma, N., Ruiter, A. and Czedik-Eysenberg, P.B. (Editors), *Residues Vet. Drugs Food, Proc. EuroResidue Conf., 2nd, 2*, Utrecht Univ. Fac. Vet. Med., Utrecht, 1993, pp. 401-403; *C.A.*, 121 (1994) 132503t.

See also 407, 425, 432, 511.

17. AMINES, AMIDES AND RELATED NITROGEN COMPOUNDS

17a. Amines and polyamines

- 511 Bladek, J. and Mischak, M.: Rapid methods for quantitation of stabilizers and their reaction products in propellants. In: Yinon, J. (Editor), *Adv. Anal. Detect. Explos., Proc. Int. Symp., 4th*, 1992, Kluwer, Dordrecht, 1993, pp. 199-207; *C.A.*, 120 (1994) 221769f.
- 512 Gonmori, K., Yoshioka, N. and Chattopadhyay, P.K.: (Identification of human semen stains by thin-layer chromatography). *Hanzaigaku Zasshi*, 60 (1994) 149-154; *C.A.*, 122 (1995) 2906m.
- 513 Weelderden-Visser, E.M.J.V. and Horst, H.J.T.: (Formulation and shelf-life of dobutamine concentrate for infusion). *Ziekenhuisfarmacie*, 10 (1994) 128-133; *C.A.*, 122 (1995) 170016q.
- 514 Wojtczak, A.B., Brdiczka, D. and Wojtczak, L.: Is monoamine oxidase activity in the outer mitochondrial membrane influenced by the mitochondrial respiratory state? *Biochim. Biophys. Acta*, 1229 (1995) 249-255.

See also 432.

17d. Other amine derivatives and amides (excl. peptides)

- 515 Mao, L. and Que, P.: (Determination of adenosine in Zhonghuashushibao (*Sennematum sinensis*)). *Shongcaoyao*, 24 (1993) 467-468; *C.A.*, 120 (1994) 62391h.
- 516 Perisic-Janjic, N.V., Djakovic, T.L. and Petrovic, S.M.: Thin-layer chromatography of benzamides on cellulose and unconventional starch and aminoplast supports. *Chromatographia*, 40 (1995) 36-38.
- 517 Pettit, G.R., Pettit, G.R., III, Backhaus, R.A. and Boettner, F.E.: Antineoplastic agents, 294. Variations in the formation of pancratistatin and related isocarbostryls in *Hymenocallis littoralis*. *J. Natural Prod.*, 58 (1995) 37-43.

See also 512.

18. AMINO ACIDS AND PEPTIDES; CHEMICAL STRUCTURE OF PROTEINS

18a. Amino acids and their derivatives

- 518 Baumbach, E.: (Hydrolysis of glutathione and TLC of amino acids). *Math. Naturwiss. Unterr.*, 47 (1994) 270-273; *C.A.*, 122 (1995) 80376c.
- 519 De Witte, P.A., Cuveele, J.F., Merlevede, W.J. and Vandenhede, J.R.: Analysis of phosphorylhydroxyamino acids present in hydrolyzed cell extracts using dansyl derivatization. *Anal. Biochem.*, 226 (1995) 1-9.
- 520 Ejiri, S.: Simple method for analyzing phosphoamino acids by thin-layer chromatography: application to elongation factor 1 from wheat embryo. *Biosci. Biotechnol. Biochem.*, 58 (1994) 2066-2067; *C.A.*, 122 (1995) 26995r.
- 521 Kobow, M., Sprung, W.-D., Schulz, E., Facklam, C. and Michalik, M.: Synthese einiger Hydroxyphenylaminosäureesterhydrochloride. *Pharmazie*, 50 (1995) 297-298.

- 522 Shi, Y. and Wang, D.: (Suitable developing solvent system for paper chromatography in cystine identification). *Sepu*, 12 (1995) 455; *C.A.*, 122 (1995) 229722x.
- 523 Siddiqi, Z.M. and Rani, S.: Studies of tin(IV) selenoarsenate III: thin layer chromatography of α -amino acids. *J. Planar Chromatogr.*, 8 (1995) 141-143.

See also 546, 632.

18b. Peptides, peptidic and proteinous hormones, growth factors

- 524 Fuchs, B., Hecker, D. and Scheidtmann, K.H.: Phosphorylation studies on rat p53 using the baculovirus expression system. Manipulation of the phosphorylation state with okadaic acid and influence on DNA binding. *Eur. J. Biochem.*, 228 (1995) 625-639.
- 525 Moyers, J.S., Linder, M.E., Shannon, J.D. and Parsons, S.J.: Identification of the *in vitro* phosphorylation sites on G_s α mediated by pp60^{c-src}. *Biochem. J.*, 305 (1995) 411-417.
- 526 Ojanperä, I., Pelander, A., Vuori, E., Himberg, K., Waris, M. and Niinivaara, K.: Detection of cyanobacterial hepatotoxins by TLC. *J. Planar Chromatogr.*, 8 (1995) 69-72.
- 527 Sakakibara, Y., Suiko, M., Nakajima, H. and Liu, M.-C.: Sulphation of L-tyrosine in mammalian cells: a comparative study. *Biochem. J.*, 305 (1995) 993-998.
- 528 Tyagi, S.C. and Simon, S.R.: Elastin-derived peptide binding to a hydrophobic domain on neutrophil elastase. *Biochem. Cell Biol.*, 72 (1994) 419-427.

See also 518.

21. PURINES, PYRIMIDINES, NUCLEIC ACIDS AND THEIR CONSTITUENTS

21a. Purines, pyrimidines, nucleosides, nucleotides

- 529 Foresti, A., Staffas, J. and Hemminki, K.: Comparison of TLC- and HPLC-32P-postlabelling assay for cisplatin-DNA adducts. *Carcinogenesis*, 15 (1994) 2829-2834; *C.A.*, 122 (1995) 71218a.
- 530 Imai, T. and Okuda, S.: Isolation and identification of adenosine triphosphoribosyl nicotinamide adenine dinucleotide phosphate from *Azotobacter vinelandii*. *J. Biochem. (Tokyo)*, 117 (1995) 163-168.
- 531 Lee, C.K., Brown, B.G., Bombick, B.R. and Doolittle, D.J.: Effect of pH on the T4 polynucleotide kinase reaction of the ³²P-post-labeling assay of DNA adducts. *Anal. Biochem.*, 227 (1995) 156-161.
- 532 Morden, W. and Wilson, I.D.: Separation of nucleotides and bases by high performance thin layer chromatography with identification by tandem mass spectrometry. *J. Planar Chromatogr.*, 8 (1995) 98-102.
- 533 Stolk, J.N., de Abreu, R.A., Boerbooms, A.M.T., de Koning, D.G.M., de Graaf, R., Kerstens, P.J.S.M. and van de Putte, L.B.A.: Purine enzyme activities in peripheral blood mononuclear cells: comparison of a new non-radiochemical high-performance liquid chromatography procedure and a radiochemical thin-layer chromatography procedure. *J. Chromatogr. B*, 666 (1995) 33-43.

- 534 Wang, M.-Y. and Liehr, J.G.: Lipid hydroperoxide-induced endogenous DNA adducts in hamster: possible mechanism of lipid hydroperoxide-mediated carcinogenesis. *Arch. Biochem. Biophys.*, 316 (1995) 38-46.

See also 515.

22. ALKALOIDS

- 535 Bieganowska, A.I. and Petruczynik, A.: Thin-layer reversed-phase chromatography of some alkaloids in ion-association systems. II. *Chem. Anal. (Warsaw)*, 39 (1994) 445-454; *C.A.*, 122 (1995) 204207v.
- 536 Draeger, B.: Identification and quantification of calysteginet polyhydroxyl nortropane alkaloids. *Phytochem. Anal.*, 6 (1995) 31-37. *C.A.*, 122 (1995) 155671f.
- 537 Qiu, N., Ye, J., Yuan, S. and Liu, J.: (Determination of guanfu base A hydrochloride in Anxin injection by TLC). *Zhongguo Yaokexue Xuebao*, 25 (1994) 309-310; *C.A.*, 122 (1995) 89604h.
- 538 Sun, X., Zhang, Z., Liu, Z. and Zhang, W.: (Quality standards of effervescent granules for arresting cold pain). *Zhongguo Zhongyao Zazhi*, 20, No. 1 (1995) 26-30; *C.A.*, 122 (1995) 196637t.
- 539 Tam, M.N., Nikolova-Damyanova, B. and Pyuskyulev, B.: Quantitative thin layer chromatography of indole alkaloids. II. Catharanthine and vindoline. *J. Liq. Chromatogr.*, 18 (1995) 849-858.

See also 420, 507, 589, 599.

23. OTHER SUBSTANCES CONTAINING HETEROCYCLIC NITROGEN

23e. Other N-heterocyclic compounds

- 540 Hedrera, M.E., Perillo, I.A. and Fernández, B.M.: HPTLC: a complementary successful method for the elucidation of organic reaction mechanisms. Cyclic amidines. *J. Planar Chromatogr.*, 8 (1995) 129-133.
- 541 Papandopulo, P.Kh., Utezhonov, K.S. and Uzakova, K.G.: (Identification of metronidazole, tinidazole and nitrazole in mixtures by thin-layer chromatography). *Aktual. Probl. Farmatsii. Alma-At. Gos. Med. In-t, Alma-Ata*, (1991) 40-42; *C.A.*, 120 (1994) 260251a.
- 542 Ray, S. and de Prabir, K.: HPTLC and TLC method for rapid quantification and identification of omeprazole. *Indian Drugs*, 31 (1994) 543-547; *C.A.*, 122 (1995) 89591b.
- 543 Tyrpien, K., Bodzek, D. and Janoszka, B.: Application of TLC and GC-MS to the identification of azaarenes in sewage sludges. *J. Planar Chromatogr.*, 8 (1995) 75-77.
- 544 Zarakar, S.S., Dhanvate, A.A., Doshi, V.J., Salunkhe, U.B. and Sawant, S.V.: Simultaneous determination of furazolidone and metronidazole from its pharmaceutical preparations using high performance thin layer chromatography. *Indian Drugs*, 31 (1994) 468-471; *C.A.*, 122 (1995) 38983y.

24. ORGANIC SULPHUR COMPOUNDS (INCL. GLUCOSINOLATES)

- 545 Zhao, S., Li, L. and Wang, B.: (Quality of standard of multicomponent Dibiling (nasal drops)). *Zhongcaoyao*, 25 (1994) 517-518, 526; C.A., 122 (1995) 114754s.

See also 413.

25. ORGANIC PHOSPHORUS COMPOUNDS (INCL. SUGAR PHOSPHATES)

- 546 Chen, S. and Claeys, M.: Approach to the large-scale preparation of highly pure phosphatidylserine from bovine brain. *J. Chromatogr. B*, 666 (1995) 178-182.

See also 459, 519, 520, 525, 530.

26. ORGANOMETALLIC AND RELATED COMPOUNDS

26c. Coordination compounds

- 547 Vuckovic, G., Miljevic, D., Janjic, T.J., Solujic, L., Juranic, N. and Celap, M.B.: Salting-out thin-layer chromatography of dipeptidato cobalt(III) complexes. *J. Serb. Chem. Soc.*, 58 (1993) 921-924; C.A., 122 (1995) 204197s.

28. ANTIBIOTICS

- 548 Abjean, J.P.: Screening of chloramphenicol residues in pork muscle by planar chromatography. *J. Assoc. Off. Anal. Chem.*, 77 (1994) 1101-1104.
- 549 Cserhati, T. and Oros, G.: Interaction of some antibiotics with hydroxypropyl- β -cyclodextrin. *J. Inclusion Phenom. Mol. Recognit. Chem.*, 18 (1994) 265-274; C.A., 122 (1995) 89253t.
- 550 Gräfe, U., Ihn, W., Ritzau, M., Schade, W., Stengel, C., Fleck, W.F., Künkel, W., Härtl, A. and Gutsche, W.: Helioferins, novel antifungal lipopeptides from *Mycogone rosea*: screening, isolation, structures and biological properties. *J. Antibiot.*, 48 (1995) 126-133.
- 551 Gräfe, U., Schlegel, R., Ritzau, M., Ihn, W., Dornberger, K., Stengel, C., Fleck, W.F., Gutsche, W., Härtl, A. and Paulus, E.F.: Aurantiamycins, new depsipeptide antibiotics from *Streptomyces aurantiacus* IMET 43917. Production, isolation, structure elucidation, and biological activity. *J. Antibiot.*, 48 (1995) 119-125.
- 552 He, H., Yu, R., Yang, Q., Yu, Z. and Wang, X.: (Quantitative analysis of midecamycin by TLC). *Zhongguo Yaoke Daxue Xuebao*, 25 (1994) 283-285; C.A., 122 (1995) 89578c.
- 553 Kinashi, H., Mori, E., Hatani, A. and Nimi, O.: Isolation and characterization of linear plasmids from lankacidin-producing *Streptomyces* species. *J. Antibiot.*, 47 (1994) 1447-1455.
- 554 Ko, H.C., Lin, H.Y. and Chen, L.H.: (Determination of monensin residues in chicken and beef by TLC and bioautography). *Zhongguo Nanyue Huaxue Huizhi*, 32 (1994) 497-506; C.A., 122 (1995) 104279b.

- 555 Lee, C.-H., Kim, S., Hyun, B., Suh, J.-W., Yon, C., Kim, C., Lim, Y. and Kim, C.: Cepacidine A, a novel antifungal antibiotic produced by *Pseudomonas cepacia*. I. Taxonomy, production, isolation and biological activity. *J. Antibiot.*, 47 (1994) 1402-1405.
- 556 Matsumoto, K., Tanaka, K., Matsutani, S., Sakazaki, R., Hino, H., Uotani, N., Tanimoto, T., Kawamura, Y., Nakamoto, S. and Yoshida, T.: Isolation and biological activity of thielocins: novel phospholipase A₂ inhibitors produced by *Thielavia terricola* RF-143. *J. Antibiot.*, 48 (1995) 106-112.
- 557 Medina, M.B., Unruh, J.J., Jones, E. and Bueso, C.E.: Solid phase clean-up and TLC analysis of hygromycin B in bovine plasma and swine serum. In: Haagsma, N., Ruiter, A. and Czedik-Eysenberg, P.B. (Editors), *Residues Vet. Drugs Food, Proc. EuroResidue Conf., 2nd, 2*, Utrecht Univ. Fac. Vet. Med., Utrecht, 1993, pp. 490-494; C.A., 121 (1994) 156018q.
- 558 Plotkowiak, Z., Popielarz-Brzezinska, M., Borkowska, M. and Plocienniczak, T.: Methods of mezlocillin analysis in presence of breakdown products. *Pharmazie*, 50 (1995) 260-262 - a review with 13 refs.
- 559 Roets, E., Adams, E., Muriithi, I.G. and Hoogmartens, J.: Determination of the relative amounts of the B and C components of neomycin by thin-layer chromatography using fluorescence detection. *J. Chromatogr. A*, 696 (1995) 131-138.
- 560 Stadler, M., Anke, H. and Sterner, O.: Metabolites with nematocidal and antimicrobial activities from the ascomycete *Lachnum papyraceum* (Karst.) Karst. III. Production of novel isocoumarin derivatives, isolation, and biological activities. *J. Antibiot.*, 48 (1995) 261-266.
- 561 Stadler, M., Anke, H. and Sterner, O.: Metabolites with nematocidal and antimicrobial activities from the ascomycete *Lachnum papyraceum* (Karst.) Karst. III. Structural elucidation of novel isocoumarin derivatives. *J. Antibiot.*, 48 (1995) 267-273.
- 562 Stadler, M., Anke, H., Shan, R. and Sterner, O.: New metabolites with nematocidal and antimicrobial activities from the ascomycete *Lachnum papyraceum* (Karst.) Karst. VI. Structure determination of non-halogenated metabolites structurally related to mycorrhizin A. *J. Antibiot.*, 48 (1995) 154-157.

See also 581, 610.

29. INSECTICIDES, PESTICIDES AND OTHER AGROCHEMICALS

29a. General techniques

- 563 Braccini, G. and Carrozza, G.P.: (Separation and identification of pesticides by thin-layer chromatography. Preliminary studies). *Inquinamento*, 36 (1994) 68-70; C.A., 122 (1995) 131294j.
- 564 Cserh ti, T. and Forg cs, E.: Charge transfer chromatographic study of the binding of commercial pesticides to various albumins. *J. Chromatogr. A*, 699 (1995) 285-290.
- 565 Kofman, I.S.: (Separation and identification of multicomponent pesticide mixtures by spectrodensitometric thin-layer chromatography). *Zagryaznenie Atmosf. i Pochvy: Tr. 4 Vses. Soveshch., Obninsk, Mai, 1989, M*, (1991) 199-208; C.A., 120 (1994) 291970p.

- 566 Mojasevic, M. and Helling, C.S.: Use of multiresidue gas chromatographic analysis to determine pesticide mobility on soil TLC plates. *J. Environ. Sci. Health, Part B*, B30 (1995) 163-173; *C.A.*, 122 (1995) 180905t.
- See also 423.
- 29c. *Phosphorus insecticides*
- 567 Falah, I.I. and Hammers, W.E.: Separation, activation and quantitation of organophosphorus insecticides in the TLC-EI technique. *Toxicol. Environ. Chem.*, 42 (1994) 35-49; *C.A.*, 122 (1995) 25760y.
- 568 Patil, B.V. and Shingare, M.S.: A new spray reagent for selective detection of dichlorvos by thin-layer chromatography. *Talanta*, 41 (1994) 367-369; *C.A.*, 120 (1994) 238116p.
- 569 Patil, V.B. and Singare, M.S.: Thin-layer chromatographic detection of monocrotophos in biological materials. *Talanta*, 41 (1994) 2127-2130; *C.A.*, 122 (1995) 125470x.
- 29e. *Herbicides*
- 570 Dailey, O.D., Jr. and Johnson, R.M.: Quantitative analysis of alachlor and atrazine in polymeric microcapsules determined by reversed-phase high performance thin layer chromatography with densitometry. *J. Liq. Chromatogr.*, 18 (1995) 873-885.
- 571 Johnson, R.M. and Pepperman, A.B.: Analysis of metribuzin and associated metabolites in soil and water samples by solid phase extraction and reversed phase thin layer chromatography. *J. Liq. Chromatogr.*, 18 (1995) 739-753.
- 29f. *Fungicides*
- 572 Sherma, J. and McGinnis, S.C.: Determination of pentachlorophenol and cymiazole in water and honey by C-18 solid phase extraction and quantitative HPTLC. *J. Liq. Chromatogr.*, 18 (1995) 755-761.
- 29g. *Other types of pesticides and various agrochemicals*
- 573 Gerunova, L.K. and Rabinovich, M.I.: (Determination of ramrod in biological systems by thin-layer chromatography). *Zh. Anal. Khim.*, 49 (1994) 1163-1165; *C.A.*, 122 (1995) 25435w.
- 574 Patil, V.B. and Shingare, M.S.: Use of cobalt acetate and o-tolidine reagent for the detection of pyrethroid insecticides by thin layer chromatography. *J. Planar Chromatogr.*, 8 (1995) 160-161.
- See also 572.
30. SYNTHETIC AND NATURAL DYES
- 30a. *Synthetic dyes*
- 575 Dugar, S.M., Leibowitz, J.N. and Dyer, R.H.: Identification of synthetic colors in beverage alcohol products by solid-phase extraction and thin-layer chromatography. *J. Assoc. Off. Anal. Chem.*, 77 (1994) 1335-1337.
- 576 Munoz, P., Reuvers, T., Martin de Pozuelo, M. and Marcos, M.V.: A screening and confirmatory method to detect malachite green residues in fish by HPTLC and HPLC-diode array. In: Haagsma, N., Ruiter, A. and Czedik-Eysenberg, P.B. (Editors), *Residues Vet. Drugs Food, Proc. EuroResidue Cong. 2nd, 2*, Utrecht Univ. Fac. Vet. Med., Utrecht, 1993, pp. 504-508; *C.A.*, 121 (1994) 156021k.
- 577 Rendle, D.F., Crabtree, S.R., Wiggins, K.G. and Salter, M.T.: Cellulase digestion of cotton dyed with reactive dyes and analysis of the products. *J. Soc. Dyers Colour.*, 110 (1994) 338-341; *C.A.*, 122 (1995) 25499d.
- 578 Sweeny, E.G., Schmidt, C.H., Zimin, A., Caputo, P.A. and Anderson, P.M.: Determination of dyes in diesel fuels. *Soc. Automot. Eng., [Spec. Publ.] SP*, SP-1056(Diesel Fuel) (1994) 1-9; *C.A.*, 122 (1995) 218059k.
- See also 407, 408, 412, 415.
- 30b. *Chloroplast and other natural pigments*
- 579 Mínguez-Mosquera, M.I., Hornerp-Méndez, D. and Garrido-Fernández, J.: Detection of bixin, lycopene, canthaxanthin, and β -apo-8'-carotenal in products derived from red pepper. *J. Assoc. Off. Anal. Chem.*, 78 (1995) 491-498.
32. DRUG ANALYSIS
- 32a. *Drug analysis, general techniques*
- 580 Heinisch, G., Kopelent-Frank, H. and Nixdorf, C.: The identity control of formulated drugs. Part 8. *Sci. Pharm.*, 61 (1993) 251-264; *C.A.*, 122 (1995) 38975x.
- 581 Kenyon, A.S., Flinn, P.E. and Layloff, T.P.: Rapid screening of pharmaceuticals by thin-layer chromatography: analysis of essential drugs by visual methods. *J. Assoc. Off. Anal. Chem.*, 78 (1995) 41-49.
- See also 423, 654.
- 32c. *Autonomic and cardiovascular drugs*
- 582 Posyniak, A., Niedzielska, J., Semeniuk, S. and Zmudzki, J.: Screening for clenbuterol and salbutamol residues in urine by planar chromatography. *J. Planar Chromatogr.*, 8 (1995) 157-159.
- 32d. *Central nervous system drugs*
- 583 Briellmann, T. and Rippstein, S.: Detection of tetracyclic benzodiazepines with reactive thin-layer chromatography. In: Daldrup, T. (Editor), *Spurenanal. Hum.-Umweltbereich, Beitr. Symp. Ges. Toxikol. Forensische Chem.*, 1991, Verlag, Heppenheim, 1992, pp. 93-98; *C.A.*, 120 (1994) 173596y.
- 584 Cserháti, T.: Interaction of monoamine oxidase inhibitors with glutamic acid, and the effect of salt on the strength of the interaction. *J. Planar Chromatogr.*, 8 (1995) 107-111.

- 585 El-Bardicy, M.G., El-Gendy, A.E., Loutfy, H.M. and Ellaithy, M.M.: Determination of nortryptiline hydrochloride and fluphenazine hydrochloride in combination. *Bull. Fac. Pharm. (Cairo Univ.)*, 31 (1993) 291-296; *C.A.*, 122 (1995) 197122h.
- 586 El-Gendy, A.E., El-Bardicy, M.G., Loutfy, H.M. and El-Tarras, M.F.: Stability indicating methods for the determination of carbamazepine. *Bull. Fac. Pharm. (Cairo Univ.)*, 31 (1993) 315-320; *C.A.*, 122 (1995) 197125m.
- 587 Fatér, Z., Szabady, B. and Nyiredy, S.: Two-dimensional over-pressured layer chromatographic separation of barbiturate derivatives. *J. Planar Chromatogr.*, 8 (1995) 145-147.
- 588 Hazai, I., Ürmös, I. and Klebovich, I.: Application of TLC - digital autoradiography as a rapid method in a pilot study of deramciclane metabolism. *J. Planar Chromatogr.*, 8 (1995) 92-97.
- 589 Kalász, H., Kerecsen, L., Csermely, T., Götz, H., Friedmann, T. and Hosztafi, S.: TLC investigation of some morphine derivatives. *J. Planar Chromatogr.*, 8 (1995) 17-22.
- 590 Kamble, V.W., Garad, M.V. and Padalikar, S.V.: Identification and estimation of diazepam in Ayurvedic preparation. *J. Planar Chromatogr.*, 8 (1995) 143-144.
- 591 Kovács-Hadady, K.: Study of the retention behavior of barbiturates by overpressured layer chromatography on silica gel layers impregnated with dodecyltrimethylammonium bromide. *J. Planar Chromatogr.*, 8 (1995) 47-51.
- 592 Spell, J.C. and Stewart, J.T.: Quantitative analysis of chloroprocaine and its major impurity 4-amino-2-chlorobenzoic acid in drug substance and injection dosage form by HPTLC and scanning densitometry. *J. Planar Chromatogr.*, 8 (1995) 72-74.
- 593 Wu, W.-N., Masucci, J.A., McKown, L.A. and Rodgers, J.D.: Identification of four plasma metabolites of etoperidone in dogs. *J. Pharm. Biomed. Anal.*, 13 (1995) 177-184.
- 32e. *Chemotherapeutics (exc. cytostatics and antibiotics)*
- 594 Balogh, G., Csizer, E., Ferenczy, G.G., Halmos, Z., Herenyi, B., Horvath, P., Lauko, A. and Gorog, S.: Estimation of impurity profiles of drugs and related materials. 12. Isolation and identification of an isomeric impurity in danazol. *Pharm. Res.*, 12 (1995) 295-298; *C.A.*, 122 (1995) 197141p.
- 595 Bieganowska, M.L., Doraczynska-Szopa, A. and Petruczynik, A.: The retention behavior of some sulfonamides on different TLC plates. 2. Comparison of the selectivity of the systems and quantitative determination of hydrophobicity parameters. *J. Planar Chromatogr.*, 8 (1995) 122-128.
- 596 Gaugain, M. and Abjean, J.-P.: A specific and sensitive fluorescent detection method for the planar chromatographic analysis of nitroimidazoles. *J. Planar Chromatogr.*, 8 (1995) 78-80.
- 597 Sadykov, Z.K. and Ikramov, L.T.: (Detection of novocaine and its metabolite - *p*-aminobenzoic acid). *Anal. Sintez i Farmakol. Izuch. Nekotor. Fiziol. Aktiv. Veshchestv. I Tashk. Gos. Med. In-t, Tashkent*, (1991) 84-86; *C.A.*, 120 (1994) 260439t.
- 598 Yang, Y., Liu, Z., Kong, Q., Shen, M. and Wang, Y.: (Pyrolytic mechanism and determination of benzodiazepine drugs by pyrolytic-TLC). *Yaowu Fenxi Zazhi*, 14 (1994) 29-32; *C.A.*, 122 (1995) 150722p.
- 599 Zhang, Q., Wu, L. and Lin, F.: (Determination of three components in APC tablets by thin layer chromatography-ultraviolet spectrophotometry). *Xiangtan Daxue Ziran Kexue Xuebao*, 15 (1993) 94-97; *C.A.*, 122 (1995) 38981w.
- See also 544, 560, 561, 586.
- 32f. *Cytostatics*
- 600 Cserhati, T.: Charge-transfer chromatographic study of the complex formation of some anticancer drugs with γ -cyclodextrin. *Anal. Biochem.*, 225 (1995) 328-332.
- 601 Forgács, E. and Cserhádi, T.: Binding of anticancer drugs to human serum albumin studied by reversed-phase chromatography. *J. Chromatogr. A*, 696 (1995) 265-272.
- 602 Li, X. and Fu, X.: (Preparation of anticancer medicines containing sophoridine). *Faming Zhuanli Shenqing Gongkai Shuomingshu* CN 1,089,846 (Cl. A61K35/78), 27 Jul. 1994, Appl. 93,100,881, 18 Jan. 1993; 13 pp.; *C.A.*, 122 (1995) 170171m.
- 603 Luo, Y. and Yang, L.: (Quality of antitumor agent etoposide VP₁₆). *Yaowu Fenxi Zazhi*, 14, No. 5 (1994) 25-27; *C.A.*, 122 (1995) 39000n.
- 32g. *Other drug categories*
- 604 Gajewska, M., Radon, B. and Glass, G.: Quantitative and quantitative analysis of estazolam. *Acta Pol. Pharm.*, 51 (1994) 207-210; *C.A.*, 122 (1995) 115163k.
- 605 Kirsten, D., Kirsten, A. and Oremek, G.M.: (Improvement in the quantitative analysis of propranolol in human blood plasma. Fluorometric determination by thin-layer chromatography). *Laboratoriumsmedizin*, 18 (1994) 466-468; *C.A.*, 122 (1995) 177536t.
- 606 Sanghavi, N.M., Puranik, K.A. and Samarth, M.M.: Analysis of ketotifen fumarate by high-performance thin-layer chromatography. *Indian Drugs*, 32, No. 1 (1995) 53-54; *C.A.*, 122 (1995) 197130j.
- 607 Smith, M.C. and Sherma, J.: Quantitative HPTLC determination of benzoyl peroxide in acne medications. *J. Planar Chromatogr.*, 8 (1995) 14-16.
- 608 Tuszyńska, E., Kwiatkowska, B., Wiczorkowska, A. and Kaniewska, T.: Investigation on chemical changes in overdue pharmaceuticals. I. Investigations on decomposition of atenolol. *Chem. Anal. (Warsaw)*, 39 (1994) 563-569; *C.A.*, 122 (1995) 38721m.
- 609 Yang, D. and Fang, T.: (Determination of amiodarone in human serum by HPTLC). *Yaowu Fenxi Zazhi*, 15 (1995) 35-37; *C.A.*, 122 (1995) 177591g.
- See also 542, 545, 585, 594, 629.
- 32i. *Plant extracts*
- 610 Chellal, A. and Lukasova, E.: Evidence for antibiotics in the two Algerian truffles *Terfezia* and *Tirmania*. *Pharmazie*, 50 (1995) 228-229.
- 611 Hong, S.P., Kim, Y.C., Kim, K.H., Park, J.H. and Park, M.K.: Characteristic component of *Rehmanniae radix preparata* compared to *Rehmanniae radix* and *Rehmanniae radix crudus*. *Anal. Sci. Technol.*, 6 (1993) 401-404; *C.A.*, 122 (1995) 222591v.

- 612 Hou, Z., Song, X. and Zhao, Y.: (Method of identification and determination of the main components in traditional Chinese medicine Kening Koufuye). *Shenyang Yaoxueyuan Xuebao*, 11 (1994) 263-268; C.A., 122 (1995) 222985b.
- 613 Kockar, O.M., Kara, M., Kara, S., Bozan, B. and Baser, K.H.C.: Quantitative determination of escin. A comparative study of HPLC and TLC-densitometry. *Fitoterapia*, 65 (1994) 439-443; C.A., 122 (1995) 170314k.
- 614 Li, K., Yang, X., Yu, M., Xie, C. and Xu, L.: (Determination of loganin in *Cornus officinalis* Sieb et Zucc. by TLC scanner). *Zhongguo Zhongyao Zazhi*, 12 (1994) 738; C.A., 122 (1995) 170374e.
- 615 Li, Y., Li, J. and Zhang, J.: (Qualitative and quantitative determination of paeonoside in therapeutic *Paeonia suffruticosa*). *Zhongcaoyao*, 25 (1994) 580-581; C.A., 122 (1995) 170317p.
- 616 Liu, M., Bai, M., Bai, R. and Feng, H.: (Processing of adhesive *Rehmannia* (*Rehmannia glutinosa*). I. Extraction, separation, identification and assay of 5-hydroxymethyl-furfural). *Zhongcaoyao*, 26 (1995) 13-14; C.A., 122 (1995) 196660v.
- 617 Ma, G., Xu, G., Jin, R. and Xu, L.: (Determination of dryocrassin and filixic acid ABA in Guanzhong by TLC scanning method). *Zhongguo Yaokexue Xuebao*, 25 (1994) 376-377; C.A., 122 (1995) 197097d.
- 618 Mo, Z.: (Quality specification of Diessheng tincture). *Yaowu Fenxi Zazhi*, 14, No. 6 (1994) 25-28; C.A., 122 (1995) 115094p.
- 619 Shou, G. and Lu, G.: (Quantitative determination of paeonin in Shuganzhitong Pian by macroporous resin pre-separation-TLC densitometry method). *Zhongcaoyao*, 25 (1994) 573-574; C.A., 122 (1995) 115096r.
- 620 Soewandhi, S.N., Soetarno, S., Troskialina, L. and Lidya, B.: Etherification of inositol mannan to improve its usage as excipient in pharmaceutical technology. *Asahi Garasu Zaidan Josei Kenkyu Seika Hokoku*, (1994) 673-682; C.A., 122 (1995) 196655x.
- 621 Song, J. and Yang, Z.: (TLC-scanning determination of liensinine in *Lotus plumule*). *Zhongguo Yiyuan Yaokexue Zazhi*, 13 (1993) 351-353; C.A., 120 (1994) 62398r.
- 622 Sun, H., Xue, X. and Lou, X.: (Comparison of constituents of five species of crude drugs "Baqi" harvested in Tianmushan mountain, Zhejiang). *Zhejiang Yike Daxuebao*, 22 (1993) 255-256; C.A., 122 (1995) 38671v.
- 623 Wang, Z. and Luo, S.: (Identification and assay of Shenxiang Vangwei Chonji). *Zhongguo Yiyuan Yaokexue Zazhi*, 14 (1994) 308-310; C.A., 122 (1995) 38973v.
- 624 Xia, Z., Xu, R., Guo, S. and Dang, F.: (TLC identification of Leigongteng (*Tripterygium wilfordii*) and Kunmiminsanhaitang (*T. hypoglaucom*). *Zhongcaoyao*, 25 (1994) 464-465; C.A., 122 (1995) 38953x.
- 625 Xiong, T., Cheng, L., Deng, H., Yang, J. and Xu, J.: (Identification of *Polygonum multiflorum* extract by TLC). *Huaxi Yaokexue Zazhi*, 9 (1994) 246-247; C.A., 122 (1995) 142149s.
- 626 Yan, Y., Li, D. and Wu, G.: (TLC-scanning determination of baicalin content in Niu Huang Qingxin Wan). *Zhongguo Zhongyao Zazhi*, 20 (1995) 33-34; C.A., 122 (1995) 222986c.
- 627 Zhang, L. and He, W.: (TLC determination of ursolic acids in Hubei Shanzha (*Crataegus hupenhensis* Sarg.)). *Yaowu Fenxi Zazhi*, 15 (1995) 30-33; C.A., 122 (1995) 222989f.
- 628 Zhang, S., Piao, H., Shen, Z., Meng, F., Zhang, Y., Zheng, Y. and Li, J.: (Detection of lignanoids content in *Sorbus pohusshanensis* (Hance) Hedl by the thin-layer chromatographic colorimetry). *Yanbian Yixueyuan Xuebao*, 17, No. 1 (1994) 32-34; C.A., 122 (1995) 115088q.
- 629 Zhao, C. and Yang, X.: (Dynamic analysis of paeoniflorin from Guizhi liquid). *Zhongguo Yaokexue Zazhi*, 29 (1994) 650-652; C.A., 122 (1995) 64551a.
- 630 Zhao, H., Xu, J. and Liu, R.: (Determination of γ -schisandrin in traditional Chinese medicine Wuweizi (*Schisandra chinensis*) and its preparation Sanlingdan by TLC-densitometry). *Yaowu Fenxi Zazhi*, 15 (1995) 33-35; C.A., 122 (1995) 222990z.
- 631 Zhou, L. and Zhong, S.: (Determination of geniposide in cape jasmine (*Gardenia jasminoides*) and its oral liquid). *Zhongcaoyao*, 25 (1994) 627-628; C.A., 122 (1995) 170333r.

See also 429, 500, 503, 537.

33. CLINICO-CHEMICAL APPLICATIONS

33b. *Complex mixtures and profiling (single compounds by cross-reference only)*

See 469, 480, 482, 638.

34. FOOD ANALYSIS

34a. *General papers and reviews*

See 443.

34b. *Complex mixtures (single compounds by cross-reference only)*

See 441, 444, 462, 479, 489, 492, 499, 509, 510, 548, 554, 557, 572, 576, 636, 637.

34c. *Organoleptically important compounds (flavors, odors, volatiles)*

632 Golc-Wondra, A., Skocir, E. and Prosek, M.: Determination of monosodium glutamate in food products. *J. Planar Chromatogr.*, 8 (1995) 117-121.

35. ENVIRONMENTAL ANALYSIS

35a. *General papers and reviews*

See 427.

35c. *Water pollution (complex mixtures; single compounds by cross-reference only)*

See 571, 572.

35d. *Soil pollution (complex mixtures; single compounds by cross-reference only)*

See 543, 571.

36. SOME TECHNICAL PRODUCTS AND COMPLEX MIXTURES

36a. *Surfactants*

633 Buschmann, N. and Wodarczak, S.: Analysis methods for carbohydrate surfactants. *Comun. Jorn. Com. Esp. Deterg.*, 25 (1994) 203-207; C.A., 122 (1995) 191034j.

634 McPherson, B.P. and Rasmussen, H.P.: Chromatography of cationic surfactants: HPLC, TLC, and GLC. *Surfactant Sci. Ser.*, 53(Cationic Surfactants) (1994) 289-326; C.A., 122 (1995) 136787g.

635 Trapani, G., Altomare, C., Franco, M., Latrofa, A. and Liso, G.: Determination of hydrophile-lipophile balance of some polyethoxylated non-ionic surfactants by reversed-phase thin layer chromatography. *Int. J. Pharm.*, 116 (1995) 95-99; C.A., 122 (1995) 169902u.

See also 468, 511, 627.

36b. *Antioxidants and preservatives*

636 Gutorska, A. and Helbrecht, A.: TLC determination of Tinuvin 622 in extracts derived from polyolefinic foodstuffs-packaging materials. *J. Planar Chromatogr.*, 8 (1995) 36-38.

637 Smith, M.C. and Sherma, J.: Determination of benzoic and sorbic acid preservatives by solid phase extraction and quantitative TLC. *J. Planar Chromatogr.*, 8 (1995) 103-106.

36c. *Complex mixtures, technical products and unidentified compounds*

638 Costongs, G.M.P.J., Janson, P.C.W., Hermans, W.J.T.A., van Oers, R.J.M. and Leerkes, B.: Evaluation of performance characteristics of automated measurement systems for allergy testing. *Eur. J. Clin. Chem. Clin. Biochem.*, 33 (1995) 295-300.

See also 426, 434.

37. CELLS, CELLULAR PARTICLES AND SUPRAMOLECULAR STRUCTURES

639 Hara, N. and Saito, K.: (Analysis of yeast cell constitution by thin layer chromatography using diatomaceous earth as a support). *Nippon Kingakkai Kaiho*, 35 (1994) 181-188; C.A., 122 (1995) 50385n.

640 Perez, J.J., Fauville-Dufaux, M., Dosogne, J.L., de Hoffmann, E. and Pouthier, F.: Faster identification of mycobacteria using gas liquid and thin layer chromatography. *J. Clin. Microbiol. Infect. Dis.*, 13 (1994) 717-725; C.A., 122 (1995) 4624s.

641 Quilliam, M.A.: Seafood toxins. *J. Assoc. Off. Anal. Chem.*, 78 (1995) 144-148 - a review with 56 refs.

38. INORGANIC COMPOUNDS

642 McLaughlin, J.R. and Sherma, J.: Quantitative HPTLC determination of elemental sulfur in sulfur topical medications. *J. Liq. Chromatogr.*, 17 (1994) 4445-4450.

See also 405.

38a. *Cations*

643 Celap, M.B.: Application of the salting-out thin-layer chromatography to transition metal coordination chemistry. *Conf. Coord. Chem.*, 14th(Contributions to Development of Coordination Chemistry) (1993) 51-52; C.A., 120 (1994) 314570z.

644 Golubev, A.V., Chekhovskaya, L.M. and Matkovskij, K.I.: (Method of determination of heavy metals in aqueous solution). *U.S.S.R. SU* 1,820,939 (Cl. G01N30/90), 07 Jun. 1993, Appl. 4,823,795, 07 May 1990; C.A., 122 (1995) 229655c.

645 Mohammad, A. and Khan, M.A.M.: TLC separation and microgram detection of metal ions on lithium chloride-impregnated silica gel and alumina layers. *J. Planar Chromatogr.*, 8 (1995) 134-140.

646 Shimizu, T., Kaneko, M. and Toyoshima, Y.: Reversed phase extraction chromatography of inorganic ions on silica gel layers impregnated with a primary alkylamine and developed with sulfuric acid and sulfate mobile phases. *J. Planar Chromatogr.*, 8 (1995) 152-154.

647 Soljic, Z., Hrestak, Z. and Eskinja, I.: (Identification of Au³⁺, V⁵⁺, Tl⁺, Tl³⁺, Ce⁴⁺, Th⁴⁺, UO₂²⁺, Pt⁴⁺, Se(IV) and Te(IV) in a mixture by thin layer chromatography). *Kem. Ind.*, 43 (1994) 415-417; C.A., 122 (1995) 177144p.

648 Vershney, K.G., Khan, A.A. and Siddiqui, M.S.: Thin-layer and paper chromatography of metal ions on some inorganic ion exchangers using nicotine as a developer at different pH values. *Proc. Natl. Acad. Sci., India, Sect. A.*, 63 (1993) 495-499; C.A., 122 (1995) 150299f.

39. RADIOACTIVE AND OTHER ISOTOPE COMPOUNDS

649 Hung, J.C., Wilson, M.E. and Silberstein, E.B.: Pitfalls in the standard radiochemical purity testing for technetium-99m-exametazime. *J. Nucl. Med. Technol.*, 22 (1994) 229-231; C.A., 122 (1995) 38991z.

650 Hung, J.C., Wilson, M.E., Gebhard, M.W. and Gibbons, R.J.: Comparison of four alternative radiochemical purity testing methods for ^{99m}Tc-sestamibi. *Nucl. Med. Commun.*, 16 (1995) 99-104; C.A., 122 (1995) 223045g.

651 Lambert, F., Slegers, G., Hermanne, A. and Mertens, J.: Production and purification of bromine-77-suitable for labeling monoclonal antibodies used in tumor imaging. *Radiochim. Acta*, 65 (1994) 223-226; C.A., 122 (1995) 89364e.

652 Molochnikova, N.P. and Myasoedov, B.F.: (Identification of the oxidation state of americium by thin-layer chromatography using Russian plates). *Zh. Anal. Khim.*, 49 (1994) 1104-1106; C.A., 122 (1995) 150319n.

653 Szentgyorgiy, P., Rakias, F. and Bodor, E.: (Radiochromatographic analysis of ^{99m}Tc -labeling of Cardio-Spect, a heart-imaging radiopharmaceutical). *Izotoptech., Diagn.*, 37 (1994) 89-98; *C.A.*, 122 (1995) 142698v.

654 Vucina, J.L., Vuga, D.A. and Vukicevic-Nikolic, N.S.: Radiochemical purity of ^{99m}Tc radiopharmaceuticals. *J. Radioanal. Nuc. Chem.*, 199 (1995) 135-142; *C.A.*, 122 (1995) 115169s.

See also 588.

Electrophoresis

1. REVIEWS AND BOOKS

- 1815 Byers, C.H. and Amarnath, A.: Understand the potential of electro-separations. *Chem. Eng. Prog.*, 91 (1995) 63-69; *C.A.*, 122 (1995) 191266m - a review with 25 refs.
- 1816 Chrambach, A., Dunn, M.J. and Radola, B.J. (Editors): *Advances in Electrophoresis*, Vol. 7, VCH, Weinheim, 1994, 487 pp.; *C.A.*, 122 (1995) 128041u.
- 1817 Hayashi, Y. and Matsuda, R.: Uncertainty structure information theory, and optimization of quantitative analysis in separation science. *Adv. Chromatogr. (N.Y.)*, 34 (1994) 347-427; *C.A.*, 122 (1995) 150209b - a review with 198 refs.
- 1818 Hilsner, V.J. and Freire, E.: Quantitative analysis of conformational equilibrium using capillary electrophoresis: applications to protein folding. *Anal. Biochem.*, 224 (1995) 465-485 - a review with 50 refs.
- 1819 Jones, W.R.: Electrophoretic capillary ion analysis. In: Landers, J.P. (Editor), *Handb. Capillary Electrophor.*, CRC, Boca Raton, 1994, pp. 209-232; *C.A.*, 122 (1995) 150208a - a review with 55 refs.
- 1820 Khaledi, M.G.: Micellar electrokinetic capillary chromatography. In: Landers, J.P. (Editor), *Handb. Capillary Electrophor.*, CRC, Boca Raton, 1994, pp. 43-93; *C.A.*, 122 (1995) 150506w - a review with 108 refs.
- 1821 Pentoney, S.L., Jr. and Sweedler, J.V.: Optical detection techniques for capillary electrophoresis. In: Landers, J.P. (Editor), *Handb. Capillary Electrophor.*, CRC, Boca Raton, 1994, pp. 147-183; *C.A.*, 122 (1995) 150507x - a review with 159 refs.
- 1822 Yonemoto, T.: (Separation-purification of biochemical substances by electrophoretic method). *Petrotech (Tokyo)*, 17 (1994) 846-851; *C.A.*, 122 (1995) 75642g - a review with 21 refs.
- 1823 Zhu, T., Fang, X. and Sun, Y.: (High performance capillary electrophoresis. Part I. Frontier in analytical chemistry). *Huaxue Jinzhan*, 6 (1994) 229-243; *C.A.*, 122 (1995) 121922y - a review with 108 refs.

See also 1841, 1843, 1864, 1868, 1869, 1872, 1876, 1879, 1886, 1894, 1895, 1896, 1898, 1901, 1903, 1905, 1909, 1911, 1917, 1921, 1922, 1925, 1928, 1934, 1935, 1940, 1953, 1956, 1969, 1982, 1999, 2034, 2042, 2043, 2049, 2065, 2114, 2128, 2151, 2157, 2212, 2352, 2367, 2390, 2435, 2449, 2452, 2501, 2519, 2540, 2597, 2600, 2617, 2618, 2620, 2621, 2634, 2647, 2652.

2. FUNDAMENTALS, THEORY AND GENERAL

2a. General

- 1824 Blasband, A.J.: Notched spacer for slab-gel electrophoresis. *U.S. US 5,384,025* (Cl. 204-299R; C25B9/00), 24 Jan. 1995, Appl. 206,995, 07 Mar. 1994; 8 pp.; *C.A.*, 122 (1995) 155707x.
- 1825 Brooks, H.B. and Brooks, D.W.: Small scale electrophoresis. *J. Chem. Educ.*, 72 (1995) A28-A29; *C.A.*, 122 (1995) 159748j.
- 1826 Cooke, M. and Wilson, I.D.: Separation science in the 21st Century. *LC-GC Int.*, 8, No. 3 (1995) 158-162.
- 1827 Kohn, A.: Automating method validation and system suitability testing in HPLC and CE. *Am. Biotechnol. Lab.*, 12 (1994) 44-47; *C.A.*, 122 (1995) 26972f.
- 1828 Razilov, I.A. and Dukhin, S.S.: (Nonlinearity of electrophoresis determined by the mobility of adsorbed ions. Strong external field). *Kolloidn. Zh.*, 56 (1994) 697-702; *C.A.*, 122 (1995) 90371m.
- 1829 Saul, A.J. and Stowers, A.: Electrophoretic resolution of charged molecules. *PCT Int. Appl. WO 94 25,144* (Cl. B01D57/02), 10 Nov. 1994, AU Appl. 93/8,659, 05 May 1993; 69 pp.; *C.A.*, 122 (1995) 209207e.
- 1830 Sharina, I.G., Cherkasov, A.V. and Nedospasov, A.A.: (Quantitative evaluations of the degree of similarity of chromatograms and electrophoretograms). *Zh. Fiz. Khim.*, 68 (1994) 1752-1757; *C.A.*, 122 (1995) 25434v.
- 1831 Terabe, S., Ozaki, H., Tanaka, Y. and Okada, Y.: (Pseudo-stationary phases for electrokinetic chromatography). *Kuromatogurafi*, 15 (1994) 114-115; *C.A.*, 122 (1995) 155375n.

See also 1892, 1893.

2b. Thermodynamics and theoretical relationships

- 1832 Allison, S.A. and Nambi, P.: Electrophoresis of spheres by a discretized integral equation/finite difference approach. *Macromolecules*, 27 (1994) 1413-1422; *C.A.*, 120 (1994) 165497b.
- 1833 Boese, F.G.: Contributions to a mathematical theory of free flow electrophoresis. In: Bauer, J. (Editor), *Cell Electrophor.*, CRC, Boca Raton, 1994, pp. 3-32; *C.A.*, 122 (1995) 50313n.
- 1834 Grateful, T.M., Athalye, A.M. and Lightfoot, E.N.: Numerical description of zone electrophoresis in the continuous flow electrophoresis device. In: Bauer, J. (Editor), *Cell Electrophor.*, CRC, Boca Raton, 1994, pp. 33-53; *C.A.*, 122 (1995) 50314p.
- 1835 Ishihama, Y., Oda, Y., Uchikawa, K. and Asakawa, N.: Evaluation of solute hydrophobicity by microemulsion electrokinetic chromatography. *Anal. Chem.*, 67 (1995) 1588-1595.
- 1836 Jumppanen, J.H. and Riekkola, M.-L.: Marker techniques for high-accuracy identification in CZE. *Anal. Chem.*, 67 (1995) 1060-1066.
- 1837 Kozulic, B.: A model of gel electrophoresis. *Appl. Theor. Electrophor.*, 4 (1994) 117-123; *C.A.*, 122 (1995) 234662g.

- 1838 Kozulic, B.: On the "donor-corridor" model of gel electrophoresis. I. Equations describing the relationship between mobility and size of DNA fragments and protein-SDS complexes. *Appl. Theor. Electrophor.*, 4 (1994) 125-136; C.A., 122 (1995) 234663h.
- 1839 Kozulic, B.: On the "donor-corridor" model of gel electrophoresis. III. The gel constant and resistance, and the net charge, friction, diffusion and electrokinetic force of the migrating molecules. *Appl. Theor. Electrophor.*, 4 (1994) 149-159; C.A., 122 (1995) 234665k.
- 1840 Kozulic, B.: On the "door-corridor" model of gel electrophoresis. II. Developments related to new gels, capillary gel electrophoresis and gel chromatography. *Appl. Theor. Electrophor.*, 4 (1994) 137-148; C.A., 122 (1995) 234664j.
- 1841 Rhodes, P.H. and Snyder, R.S.: Theoretical and experimental studies on the stabilization of hydrodynamic flow in free fluid electrophoresis. In: Bauer, J. (Editor), *Cell Electrophor.*, CRC, Boca Raton, 1994, pp. 57-74; C.A., 122 (1995) 50315q - a review with 14 refs.
- 1842 Zydny, A.L.: Boundary effects on the electrophoretic motion of a charged particle in a spherical cavity. *J. Colloid Interface Sci.*, 169 (1995) 476-485; C.A., 122 (1995) 197899s.
- See also 1818, 1930, 1944, 1994, 2035, 2645.
- 2c. *Relationship between structure and electrophoretic behaviour*
- See 1835, 2408.
- 2d. *Measurement of physico-chemical and related values*
- 1843 Kirkland, J.J.: New separation method for characterizing the size of silica sols. *Adv. Chem. Ser.*, 234 (1994) 287-308; C.A., 122 (1995) 197691t - a review with 36 refs.
- 1844 Yarmola, E. and Chrambach, A.: Nonlinear "Ferguson curves" by two runs of the commercial automated HPGE-1000 gel electrophoresis apparatus with intermittent scanning of fluorescence. *Electrophoresis (Weinheim)*, 16 (1995) 350-353.
- See also 1902, 1919, 2592, 2619.
3. GENERAL TECHNIQUES
- 3a. *Apparatus and accessories*
- 1845 Alam, A.: Electroelution apparatus and method for isolation and recovery of biomolecules. *PCT Int. Appl. WO 94 28,406* (Cl. G01N27/447), 08 Dec. 1994, US Appl. 63,731, 20 May 1993; 30 pp.; C.A., 122 (1995) 101110d.
- 1846 Arai, A.: Device for capillary electrophoresis chiral separation. *Jpn. Kokai Tokyo Koho* JP 06,288,983 [94,288,983] (Cl. G01N27/447), 18 Oct. 1994, Appl. 93/98,664, 31 Mar. 1993; 5 pp.; C.A., 122 (1995) 204271m.
- 1847 Bevan, C.D. and Mutton, I.M.: Use of freeze-thaw flow management for controlling and switching fluid flow in capillary tubes. *Anal. Chem.*, 67 (1995) 1470-1473.
- 1848 Cognard, D. and Hache, J.: Method and apparatus for multiple electrophoresis for the migration and transfer of macromolecules. *Eur. Pat. Appl. EP 605,269* (Cl. B01D57/02), 06 Jul. 1994, FR Appl. 92/15,451, 22 Dec. 1992; 9 pp.; C.A., 122 (1995) 76001j.
- 1849 Hagiwara, H., Irisawa, K., Myazaki, J. and Ozaki, T.: Gel electrophoresis device. *Jpn. Kokai Tokkyo Koho* JP 06,288,982 [94,288,982] (Cl. G01N27/447), 18 Oct. 1994, Appl. 93/889, 30 Mar. 1993; 4 p.; C.A., 122 (1995) 122255v.
- 1850 Inomata, T. and Takahashi, Y.: Capillary electrophoresis apparatus. *Jpn. Kokai Tokkyo Koho* JP 06,324,019 [94,324,019] (Cl. G01N27/447), 25 Nov. 1994, Appl. 93/111,399, 13 May 1993; 7 p.; C.A., 122 (1995) 177429k.
- 1851 Kambara, H., Anazawa, T., Takahashi, S. and Murakami, K.: Electrophoresis analyzer with wavelength-selective detection. *Eur. Pat. Appl. EP 619,483* (Cl. G01N21/64), 12 Oct. 1994, JP Appl. 93/80,927, 07 Apr. 1993; 14 pp.; C.A., 122 (1995) 50717x.
- 1852 Kontush, S.M., Dukhin, S.S. and Vidov, O.I.: (Aperiodic electrophoresis). *Kolloidn. Zh.*, 56 (1994) 654-660; C.A., 122 (1995) 90369s.
- 1853 Liu, D.: (A new apparatus for electrophoresis). *Daxue Huaxue*, 9 (1994) 46; C.A., 122 (1995) 238781t.
- 1854 Moring, S.E., Albin, M.S., Kowallis, R.B., Lee, T.E., Mead, D.E., Nickel, J.H., Oldham, M.F., Reel, R.T., Orpin, T.S. and Woods, J.C.: Capillary electrophoresis instrument. *U.S. US 5,384,024* (Cl. 204-299R; B01D61/46), 24 Jan. 1995, Appl. 850,764, 13 Mar. 1992; 25 pp.; C.A., 122 (1995) 155706w.
- 1855 Nakazato, T. and Shinozaki, T.: Multiple sample-applicator for gel electrophoresis. *Jpn. Kokai Tokkyo Koho* JP 06,281,621 [94,281,621] (Cl. G01N27/447), 07 Oct. 1994, Appl. 92/232,206, 31 Aug. 1992; 11 pp.; C.A., 122 (1995) 128048b.
- 1856 Sabanayagam, C.R., Holzwarth, G.M. and Lai, E.H.: Method and apparatus for gel electrophoresis. *Eur. Pat. Appl. EP 644,420* (Cl. G01N27/447), 22 Mar. 1995, US Appl. 119,033, 09 Sep. 1993; 22 pp.; C.A., 122 (1995) 234831m.
- 1857 Sarrine, R.J.: Platform for conducting electrophoresis, and an electrophoresis plate for use with the platform. *Eur. Pat. Appl. EP 631,134* (Cl. G01N27/447), 28 Dec. 1994, US Appl. 79,229, 21 Jun. 1993; 10 pp.; C.A., 122 (1995) 101114h.
- 1858 Sarrine, R.J., Garsee, H.A., Kelley, C.D., Everitt, M.T., Golias, T.L., Boone, E.W., Guadagno, P.A. and Petersen, E.H.: Automatic electrophoresis method and apparatus. *Eur. Pat. Appl. EP 631,132* (Cl. G01N27/447), 28 Dec. 1994, US Appl. 79,378, 21 Jun. 1993; 56 pp.; C.A., 122 (1995) 101115j.
- See also 1824, 1907, 1912, 1925, 2000, 2060.
- 3b. *Detectors and detection procedures*
- 1859 Andersson, P.E., Pfeffer, W.D. and Blomberg, L.G.: Indirect detection in capillary electrophoresis. Comparison between indirect UV and indirect laser-induced fluorescence detection for the determination of isoprenyl pyrophosphates. *J. Chromatogr. A*, 699 (1995) 323-330.
- 1860 Gunsheski, M.A.: Indirect fluorometric detection of arsenic and selenium oxyanions separated by capillary zone electrophoresis and ion chromatography. Avail. *Univ. Microfilms Int.*, Order No. DA 9431955, 1993, 157 p.; C.A., 122 (1995) 203978k.

- 1861 Lim, H.B., Lee, J.J. and Lee, K.-J.: Simple and sensitive laser-induced fluorescence detection for capillary electrophoresis and its application to protein separation. *Electrophoresis (Weinheim)*, 16 (1995) 674-678.
- 1862 Lohmann, J., Schickle, H. and Bosch, T.C.G.: REN display, a rapid and efficient method for nonradioactive differential display and mRNA isolation. *BioTechniques*, 18 (1995) 200-202; *C.A.*, 122 (1995) 127919z.
- 1863 Park, S., Lunte, S.M. and Lunte, C.E.: A perfluorosulfonated ionomer joint for capillary electrophoresis with on-column electrochemical detection. *Anal. Chem.*, 67 (1995) 911-918.
- 1864 Patton, W.F.: Biologist's perspective on analytical imaging systems as applied to protein gel electrophoresis. *J. Chromatogr. A*, 698 (1995) 55-87 - a review with 98 refs.
- 1865 Potter, C.G. and Anderson, L.J.: Method of measuring chemiluminescence of multiple samples on a continuous matrix. *PCT Int. Appl. WO 94 25,855* (Cl. G01N21/76), 10 Nov. 1994, *FI Appl. 93/1,858*, 23 Apr. 1993; 37 pp.; *C.A.*, 122 (1995) 76013q.
- 1866 Saz, J.M., Krattiger, B., Bruno, A.E., Diez-Masa, J.C. and Widmer, H.M.: Thermo-optical absorbance detection of native proteins separated by capillary electrophoresis in 10 μ m I.D. tubes. *J. Chromatogr. A*, 699 (1995) 315-322.
- 1867 Shear, J.B., Fishman, H.A., Allbritton, N.L., Garigan, D., Zare, R.N. and Scheller, R.H.: Single cells as biosensors for chemical separations. *Science (Washington)*, 267 (1995) 74-77; *C.A.*, 122 (1995) 75657r.
- 1868 Takagi, T.: (New aspects of electrophoresis and concern about manufacturing). *Tanpakushitsu Kakusan Koso*, 40 (1995) 350-357; *C.A.*, 122 (1995) 127722e - a review with 25 refs.
- 1869 Toda, T. and Ohashi, M.: Current status and perspectives of computer-aided two-dimensional densitometry. *J. Chromatogr. A*, 698 (1995) 41-54 - a review with 35 refs.
- See also 1821, 1849, 1943, 2018, 2066, 2084, 2352, 2581, 2667.
- 3c. *Stabilization media for electrophoresis*
- 1870 Akashi, M., Inami, Y., Kishida, A. and Baba, Y.: (Novel hydrogels for electrophoresis). *Kuromatogurafi*, 15 (1994) 108-109; *C.A.*, 122 (1995) 182485e.
- 1871 Buettner, C. and Beck, W.: A method of producing a capillary for an electrophoresis device. *Eur. Pat. Appl. EP 634,651* (Cl. G01N27/447), 18 Jan. 1995, *Appl. 93/111,150*, 12 Jul. 1993; 13 pp.; *C.A.*, 122 (1995) 101113g.
- 1872 Destro-Bisol, G. and Santini, S.A.: Electrophoresis on cellulose acetate and Cellogel: current status and perspectives. *J. Chromatogr. A*, 698 (1995) 33-40 - a review with 43 refs.
- 1873 Guadagno, P.A., Rayachoti, R. and Petersen, E.H.: An electrophoresis plate and a method of making an electrophoresis plate package. *Eur. Pat. Appl. EP 631,133* (Cl. G01N27/447), 28 Dec. 1994, *US Appl. 79,228*, 21 Jun. 1993; 8 pp.; *C.A.*, 122 (1995) 128056c.
- 1874 Lin, B., Chu, X. and Yang, Y.: Sym-triazines dye affinity capillary electrophoresis column and its preparation. *Faming Zhuanli Shenqing Gongkai Shuomingshu CN 1,085,318* (Cl. G01N30/02), 13 Apr. 1994, *Appl. 92,109,938*, 29 Sep. 1992; 12 pp.; *C.A.*, 122 (1995) 50708v.
- 1875 Mechref, Y. and El Rassi, Z.: Fused-silica capillaries with surface-bound dextran layer crosslinked with diepoxypolyethylene glycol for capillary electrophoresis of biological substances at reduced electroosmotic flow. *Electrophoresis (Weinheim)*, 16 (1995) 617-624.
- 1876 Righetti, P.G.: Macroporous gels: facts and misfacts. *J. Chromatogr. A*, 698 (1995) 3-17 - a review with 53 refs.
- 1877 Shimizu, M.: Electrophoresis gel for separation and recovery of DNA and proteins. *Jpn. Kokai Tokkyo Koho JP 06,273,382* [94,273,382] (Cl. G01N27/447), 30 Sep. 1994, *Appl. 93/57,235*, 17 Mar. 1993; 6 p.; *C.A.*, 122 (1995) 177433g.
- 1878 Wheeler, T.T., Ford, H.C. and Wilkins, R.J.: Rehydration of dried polyacrylamide gels. *BioTechniques*, 18 (1995) 233-234; *C.A.*, 122 (1995) 128033t.
- 1879 Wirth, P.J. and Romano, A.: Staining methods in gel electrophoresis, including the use of multiple detection methods. *J. Chromatogr. A*, 698 (1995) 123-143 - a review with 166 refs.
- See also 1897, 1913, 1918, 1923, 1931, 2024, 2065.
- 3d. *Quantitative analysis*
- See 1817.
- 3e. *Preparative scale electrophoresis*
- See 2153, 2557.
- 3f. *Programmed voltage and buffer gradients*
- See 1901, 1934.
4. SPECIAL TECHNIQUES
- 4a. *Automation*
- See 1858, 1882.
- 4b. *Computerization and modelling*
- 1880 Barkema, G.T., Marko, J.F. and Widom, B.: Electrophoresis of charged polymers: simulation and scaling in a lattice model of reptation. *Phys. Rev. E: Stat. Phys., Plasmas, Fluids, Relat. Interdiscip. Top.*, 49 (1994) 5303-5309; *C.A.*, 121 (1994) 58450e.
- 1881 Horgan, G.W. and Glasbey, C.A.: Uses of digital image analysis in electrophoresis. *Electrophoresis (Weinheim)*, 16 (1995) 298-305.
- 1882 Riveron, A.M., Herrera, J.A., Ruiz-Esquivel, L., Baez-Camargo, M., Lopez-Canovas, L., Noa, M.D. and Orozco, E.: Hardware and a memory resident program for simultaneous control and switching of pulsed field gel electrophoresis. *Anal. Lett.*, 28 (1995) 945-860.
- 1883 Schubert, F.E.: Suspension for use electrophoretic image display systems. *U.S. US 5,380,362* (Cl. 106-493; C08K13/06), 10 Jan. 1995, *Appl. 92,749*, 16 Jul. 1993; 8 p.; *C.A.*, 122 (1995) 147657r.

- 1884 Sevick, E.M. and Williams, D.R.M.: Motion of a polyelectrolyte chain hooked around a post. *Phys. Rev. E: Stat. Phys., Plasmas, Fluids, Relat. Interdiscip. Top.*, 50 (1994) R3357-R3360; *C.A.*, 122 (1995) 115933m.
- See also 1837, 1838, 1839, 1840, 1887, 1893, 2072, 2124, 2383, 2439, 2473, 2482, 2487.
- 4c. *Combination with other physicochemical techniques, (MS, IR etc.)*
- 1885 Takada, Y., Sakairi, M. and Koizumi, H.: Atmospheric pressure chemical ionization interface for capillary electrophoresis/mass spectrometry. *Anal. Chem.*, 67 (1995) 1474-1476.
- See also 1929, 1995, 2002, 2029, 2044, 2141, 2369, 2490, 2599, 2610, 2661.
- 4d. *Affinity electrophoresis*
- 1886 Takeo, K.: Advances in affinity electrophoresis. *J. Chromatogr. A*, 698 (1995) 89-105 - a review with 54 refs.
- 4e. *Capillary zone electrophoresis and electrokinetic chromatography*
- 1887 Beckers, J.L.: Calculation of the composition of sample zones in capillary zone electrophoresis. II. Simulated electropherograms. *J. Chromatogr. A*, 696 (1995) 285-294.
- 1888 Busch, M.H.A., Kraak, J.C. and Poppe, H.: Cellulose acetate-coated fused-silica capillaries for the separation of proteins by capillary zone electrophoresis. *J. Chromatogr. A*, 695 (1995) 287-296.
- 1889 Castagnola, M., Cassiano, L., Lupi, A., Messina, I., Patamia, M., Rabino, R., Rossetti, D.V. and Giardina, B.: Ion-exchange electrokinetic capillary chromatography with Starburst (pamam) dendrimers: a route towards high-performance electrokinetic capillary chromatography. *J. Chromatogr. A*, 694 (1995) 463-469.
- 1890 Cole, L.J. and Kennedy, R.T.: Selective preconcentration for capillary zone electrophoresis using protein G immunoaffinity capillary chromatography. *Electrophoresis (Weinheim)*, 16 (1995) 549-556.
- 1891 Demorest, D.: Application-specific capillary electrophoresis. *PCT Int. Appl. WO 94 26,396 (Cl. B01D57/02)*, 24 Nov. 1994, US Appl. 59,946, 07 May 1993; 39 pp.; *C.A.*, 122 (1995) 155714x.
- 1892 Dunn, C.D., Hankins, M.G. and Ghowsi, K.: Reverse direction anion capillary electrophoresis: theory and application. *Sep. Sci. Technol.*, 29 (1994) 2419-2433; *C.A.*, 122 (1995) 121989a.
- 1893 Ermakov, S.V., Zhukov, M.Yu., Capelli, L. and Righetti, P.G.: Wall adsorption in capillary electrophoresis. Experimental study and computer simulation. *J. Chromatogr. A*, 699 (1995) 297-313.
- 1894 Guan, F.: (Enantiomeric separation by high-performance capillary electrophoresis). *Fenxi Huaxue*, 22 (1994) 731-737; *C.A.*, 122 (1995) 95559p - a review with 54 refs.
- 1895 Guan, F.: (Separation and determination of ions by high-performance capillary electrophoresis). *Fenxi Huaxue*, 23 (1995) 111-116; *C.A.*, 122 (1995) 177067r - a review with 50 refs.
- 1896 Guan, F.: (The role of surfactants in high performance capillary electrophoresis). *Sepu*, 13 (1995) 30-32; *C.A.*, 122 (1995) 204163c - a review with 38 refs.
- 1897 Guttman, A., Shieh, C.H., Karger, B.L., Pentoney, S.J., Jr., Konrad, K.D., Rampal, S. and Ganzler, K.: Capillary electrophoresis column containing removable separation gel composition, and method of use. *U.S. US 5,370,777 (Cl. 204-182.8; GO1N27/26)*, 06 Dec. 1994, US Appl. 829,638, 31 Jan. 1992; 23 pp.; *C.A.*, 122 (1995) 76042y.
- 1898 Heiger, D.N. and Majors, R.E.: Capillaries and chemistries for capillary electrophoresis. *LC-GC*, 13 (1995) 12-23; *C.A.*, 122 (1995) 75645k - a review with 15 refs.
- 1899 Hjertén, S., Valtcheva, L., Elenbring, K. and Liao, J.-L.: Fast, high-resolution (capillary) electrophoresis in buffers designed for high field strengths. *Electrophoresis (Weinheim)*, 16 (1995) 584-594.
- 1900 Huang, M., Dubrovackova-Schneiderman, E., Novotny, M.V., Fatunmbi, H.O. and Wirth, M.J.: Self-assembled alkyl silane monolayers for the preparation of stable and efficient coatings in capillary electrophoresis. *J. Microcolumn Sep.*, 6 (1994) 571-576; *C.A.*, 122 (1995) 100894a.
- 1901 Janini, G.M. and Issaq, H.J.: The buffer in capillary zone electrophoresis. *Chromatogr. Sci. Ser.*, 64 (1993) 119-160; *C.A.*, 120 (1994) 201066x - a review with 91 refs.
- 1902 Janini, G.M., Muschik, G.M. and Issaq, H.J.: Micellar electrokinetic chromatography in coated capillaries: determination of micelle-aqueous partition coefficients. *J. High Resolut. Chromatogr.*, 18 (1995) 171-174.
- 1903 Jankowski, J.A., Tracht, S. and Sweedler, J.V.: Assaying single cells with capillary electrophoresis. *TrAC*, 14 (1995) 170-176 - a review with 39 refs.
- 1904 Kano, K., Minami, K., Horiguchi, K., Ishimura, T. and Kodera, M.: Ability of non-cyclic oligosaccharides to form molecular complexes and its use for chiral separation by capillary zone electrophoresis. *J. Chromatogr. A*, 694 (1995) 307-313.
- 1905 Kennler, E.: Organic solvents in capillary electrophoresis. *Chromatogr. Sci. Ser.*, 64 (1993) 161-186; *C.A.*, 120 (1994) 201067x - a review with 46 refs.
- 1906 Khaledi, M.G., Quang, C., Sahota, R.S., Strasters, J.K. and Smith, S.C.: Controlling migration behavior in capillary electrophoresis: optimization strategies for method development. *Chromatogr. Sci. Ser.*, 64 (1993) 187-260; *C.A.*, 120 (1994) 205279r.
- 1907 Knoll, M. and Cammann, D.: Capillary electrophoretic separation method and apparatus for chemical and biochemical analysis. *Ger. Offen. DE 4,314,755 (Cl. GO1N27/447)*, 10 Nov. 1994, Appl. 05 May 1993; 15 pp.; *C.A.*, 122 (1995) 27247k.
- 1908 Krstanovic, S.: Connectorized capillaries for use with separation instrumentation components. *Eur. Pat. Appl. EP 617,278 (Cl. GO1N27/447)*, 28 Sep. 1994, US Appl. 31,007, 12 Mar. 1993; 25 pp.; *C.A.*, 122 (1995) 27245h.
- 1909 Lickl, E.: (Capillary electrophoresis in trace analysis). *Oester. Chem. Z.*, 95 (1994) 238-240; *C.A.*, 122 (1995) 122194z - a review with 19 refs.
- 1910 Manz, A., Effenhauser, C.S., Burggraf, N., Verpoorte, E., Raymond, D.E. and Widmer, H.M.: Capillary electrophoresis integrated onto a planar microstructure. *Analysis*, 22 (1994) M25-M28; *C.A.*, 122 (1995) 177104a.

- 1911 Manz, A., Effenhauser, C.S., Burggraf, N., Verpoorte, E., Raymond, D.E. and Widmer, H.M.: Capillary electrophoresis integrated onto a planar microstructure. A step towards miniaturized total analysis system (μ -TAS). *Kagaku to Kogyo (Tokyo)*, 47 (1994) 1536-1541; *C.A.*, 122 (1995) 95307e - a review with 38 refs.
- 1912 Manz, A., Harrison, D.J. and Effenhauser, C.S.: Method and sample applicator for controlling sample introduction in micro-column separation techniques. *Eur. Pat. Appl. EP 620,432*, (Cl. GO1N27/447), 19 Oct. 1994, Appl. 93/810,272, 15 Apr. 1993; 11 pp.; *C.A.*, 122 (1995) 50716w.
- 1913 Mayer, S., Schleimer, M. and Schurig, V.: Dual chiral recognition system involving cyclodextrin derivatives in capillary electrophoresis. *J. Microcolumn Sep.*, 6 (1994) 43-48; *C.A.*, 122 (1995) 122210b.
- 1914 Muijselaar, P.G.H.M., Claessens, H.A. and Cramers, C.A.: Parameters controlling the elution window and retention factors in micellar electrokinetic capillary chromatography. *J. Chromatogr. A*, 696 (1995) 273-284.
- 1915 Muijselaar, P.G.H.M., Claessens, H.A., Cramers, C.A., Jansen, J.F.G.A., Meijer, E.W., de Brabander-van den Berg, E.M.M. and van der Wal, S.: Dendrimers as pseudo-stationary phases in electrokinetic chromatography. *J. High Resolut. Chromatogr.*, 18 (1995) 121-123.
- 1916 Navin, M.J. and Morris, M.D.: Capillary electrophoresis separation techniques and mechanisms in dilute polymer matrices. *J. Chin. Chem. Soc. (Taipei)*, 42 (1995) 5-9; *C.A.*, 122 (1995) 234650b.
- 1917 Nishi, H. and Terabe, S.: Optical resolution of drugs by capillary electrophoretic techniques. *J. Chromatogr. A*, 694 (1995) 245-276 - a review with 146 refs.
- 1918 Okuyama, T., Izumi, T. and Yamaguchi, M.: Capillary column packed with glucomannan gel for capillary electrophoresis and methods for making and using same. *U.S. US 5,364,520* (Cl. 204-299R; C25B9/00), 15 Nov. 1994, JP Appl. 91/350,146, 10 Dec. 1991; 4 pp.; *C.A.*, 122 (1995) 50745e.
- 1919 Penn, S.G., Bergstrom, E.T., Knights, I., Liu, G., Ruddick, A. and Goodall, D.M.: Capillary electrophoresis as a method for determining binding constants: application to the binding of cyclodextrins and nitrophenolates. *J. Phys. Chem.*, 99 (1995) 3875-3880; *C.A.*, 122 (1995) 182502h.
- 1920 Quang, C.: Chemically selective interactions in capillary zone electrophoresis. Avail. *Univ. Microfilms Int.*, Order No. DA9431884, 1994, 129 p.; *C.A.*, 122 (1995) 170311g.
- 1921 Regnier, F.E., Patterson, D.H. and Harmon, B.J.: Electrophoretically-mediated microanalysis (EMMA). *TrAC*, 14 (1995) 177-181 - a review with 21 refs.
- 1922 Semba, T.: (Present status of capillary electrophoresis (CE). A view of studies on proteins by CE). *Meiji Yakka Daigaku Kenkyu Kiyu*, 24 (1994) 59-61; *C.A.*, 122 (1995) 234480w - a review with many refs.
- 1923 Shieh, C.H.: Concentration-variation methods for controlling electroosmotic flow in coated capillary electrophoresis columns. *U.S. US 5,391,274* (Cl. 204-180.1; B01D57/02), 21 Feb. 1995, Appl. 138,324, 18 Oct. 1993; 13 pp.; *C.A.*, 122 (1995) 109226k.
- 1924 Stathakis, C. and Cassidy, R.M.: Ionic polymers for selectivity control in the capillary electrophoretic separations of small ions. *Am. Lab. (Shelton)*, 26 (1994) 28J-28M; *C.A.*, 122 (1995) 100889c.
- 1925 Stevenson, R.: A critical review of the development of HPCE instrumentation. *Am. Lab. (Shelton)*, 26 (1994) 29-33; *C.A.*, 122 (1995) 127717g - a review with 24 refs.
- 1926 Sun, P. and Hartwick, R.A.: On-line kinetic monitoring for biochemical reactions using multi-point detection in high-performance capillary electrophoresis. *J. Chromatogr. A*, 695 (1995) 279-285.
- 1927 Tanaka, N., Fukutome, T., Tanigawa, T., Hosoya, K., Kimata, K., Araki, T. and Unger, K.K.: Structural selectivity provided by starburst dendrimers as pseudostationary phase in electrokinetic chromatography. *J. Chromatogr. A*, 699 (1995) 331-341.
- 1928 Terabe, S.: Micellar electrokinetic chromatography. *Chromatogr. Sci. Ser.*, 64 (1993) 65-87; *C.A.*, 120 (1994) 201064u - a review with 43 refs.
- 1929 Tomlinson, A.J., Benson, L.M. and Naylor, S.: Advantages of nonaqueous solvents in the analysis of drug metabolites using CE and on-line CE-MS. *LC-GC Int.*, 8, No. 4 (1995) 210-216.
- 1930 Winzor, D.J.: Measurement of binding constants by capillary electrophoresis. *J. Chromatogr. A*, 696 (1995) 160-163.
- 1931 Xu, H.: (Derivatized cyclodextrins for the separation of chiral drugs in capillary electrophoresis). *Yaowu Fenxi Zazhi*, 14 (1994) 48-50; *C.A.*, 122 (1995) 115144e.
- 1932 Zheng, J., Hu, R., Lin, H. and Liu, W.: (Application of column-coupling in capillary electrophoresis). *Fenxi Huaxue*, 22 (1994) 855-858; *C.A.*, 122 (1995) 121963n.
- See also 1817, 1818, 1819, 1820, 1821, 1823, 1831, 1835, 1836, 1840, 1843, 1846, 1847, 1850, 1854, 1859, 1860, 1861, 1863, 1866, 1871, 1874, 1875, 1876, 1885, 1886, 1933, 1935, 1939, 1940, 1943, 1946, 1947, 1948, 1950, 1954, 1955, 1956, 1957, 1961, 1962, 1976, 1977, 1978, 1990, 1991, 1993, 1994, 1995, 1996, 1997, 1998, 1999, 2000, 2001, 2002, 2005, 2008, 2009, 2012, 2015, 2016, 2017, 2018, 2019, 2020, 2021, 2022, 2023, 2024, 2025, 2026, 2032, 2037, 2038, 2039, 2041, 2043, 2047, 2048, 2053, 2056, 2057, 2063, 2068, 2069, 2074, 2106, 2109, 2110, 2126, 2134, 2141, 2143, 2145, 2162, 2167, 2178, 2181, 2187, 2190, 2194, 2195, 2196, 2201, 2258, 2285, 2297, 2307, 2309, 2313, 2335, 2348, 2363, 2364, 2369, 2407, 2429, 2430, 2453, 2454, 2459, 2460, 2483, 2490, 2506, 2507, 2511, 2515, 2530, 2552, 2564, 2569, 2571, 2572, 2574, 2577, 2578, 2579, 2581, 2582, 2583, 2585, 2587, 2588, 2590, 2591, 2593, 2594, 2595, 2596, 2597, 2598, 2599, 2600, 2601, 2602, 2603, 2604, 2605, 2606, 2607, 2608, 2609, 2610, 2611, 2612, 2613, 2614, 2617, 2620, 2621, 2624, 2625, 2626, 2629, 2631, 2632, 2653, 2654, 2655, 2657, 2658, 2659, 2660, 2663, 2664, 2665, 2666, 2667, 2668, 2669, 2670.
- 4f. *Isotachophoresis*
- 1933 Altria, K.D. and Hadgett, T.A.: An evaluation of the use of capillary electrophoresis to monitor trace drug residues following the manufacture of pharmaceuticals. *Chromatographia*, 40 (1995) 23-27.

1934 Righetti, P.G. and Chiari, M.: Conventional isoelectric focusing and immobilized pH gradients: an overview. *Chromatogr. Sci. Ser.*, 64 (1993) 89-116; *C.A.*, 120 (1994) 201065v - a review with 57 refs.

See also 1890, 1992, 2628.

4g. Enantiomers, separation

1935 Bereuter, R.L.: Enantioseparation by capillary electrophoresis. *LC-GC*, 12 (1994) 748-766; *C.A.*, 122 (1995) 100691g - a review with 119 refs.

See also 1846, 1894, 1904, 1913, 1917, 1997, 2578, 2585, 2598, 2602, 2603, 2605, 2611.

4h. Two dimensional electrophoresis

1936 Fritz, P., Mischlinski, A., Wicherek, C. and Saal, J.: Two-dimensional cryostat section electrophoresis: a novel method and its application to the diseased synovial membrane. *Histochem. J.*, 26 (1994) 804-816; *C.A.*, 122 (1995) 27068c.

1937 Poppe, H.: The role of miniaturization in the development of analytical fractionation methods. *Analisis*, 22 (1994) M22-M24; *C.A.*, 122 (1995) 177132h.

1938 Wang, T. and Tong, Z.: (The methods solving problems encountered in two-dimensional polyacrylamide gel electrophoresis). *Zhivü Shenglixue Tongxun*, 30 (1994) 283-285; *C.A.*, 122 (1995) 50518h.

See also 1864, 1869, 1879, 1881, 1886, 1982, 2007, 2027, 2029, 2034, 2036, 2044, 2046, 2060, 2072, 2082, 2085, 2089, 2096, 2099, 2114, 2121, 2124, 2135, 2151, 2157, 2166, 2173, 2179, 2199, 2200, 2202, 2212, 2217, 2225, 2245, 2255, 2256, 2277, 2302, 2336, 2388, 2487, 2508, 2533, 2560, 2618, 2647.

4i. Other special techniques

1939 Duro, G., Barbieri, R., Ribauda, M.R., Feo, S. and Izzo, V.: A downward capillary blotting procedure from hundred base pairs to hundred kilobases nucleic acids. *Anal. Biochem.*, 225 (1995) 360-362.

1940 Lunte, S.M., Malone, M.A. and Zuo, H.: Capillary electrophoresis/electrochemistry for the analysis of microdialysis samples. *Curr. Sep.*, 13 (1994) 75-79; *C.A.*, 122 (1995) 50304k - a review with 21 refs.

1941 Slais, K. and Friedl, Z.: Ampholytic dyes for spectroscopic determination of pH in electrofocusing. *J. Chromatogr. A*, 695 (1995) 113-122.

1942 Yuan, N., Liu, Z. and Zhu, D.: Chromatograph-type electrophoresis separation method. *Faming Zhuanli Shenqing Gongkai Shuomingshu* CN 1,083,072 (Cl. C07K3/14), 02 Mar. 1994, Appl. 93,108,629, 16 Jul. 1993; 8 pp.; *C.A.*, 122 (1995) 50769r.

See also 1852, 1882, 1910, 2453.

5. HYDROCARBONS AND HALOGEN DERIVATIVES

5b. Cyclic hydrocarbons, fullerenes

1943 Kaneta, T. and Imasaka, T.: Indirect detection of aromatic hydrocarbons by semiconductor laser fluorometry in micellar electrokinetic chromatography. *Anal. Chem.*, 67 (1995) 829-834.

See also 1927.

5d. Complex hydrocarbon mixtures (incl. analysis of tars, bitumens and mineral oils)

1944 Kokal, S., Tang, T., Schramm, L. and Sayegh, S.: Electrokinetic and adsorption properties of asphaltenes. *Colloids Surf., A*, 94 (1995) 253-265; *C.A.*, 122 (1995) 137690g.

1945 Samadijavan, A. and Loidl, D.: (Electrophoretic study of the refractive behavior of bitumen emulsion). *Oester. Chem. Z.*, 95 (1994) 272-277; *C.A.*, 122 (1995) 165062x.

6. ALCOHOLS

1946 Altria, K.D. and Howells, J.S.: Quantitative determination of organic solvents by capillary electrophoresis using indirect UV detection. *J. Chromatogr. A*, 696 (1995) 341-348.

7. PHENOLS

1947 Benz, N.J. and Fritz, J.S.: Optimization of separations of alkyl-substituted phenolate anions by capillary zone electrophoresis. *J. High Resolut. Chromatogr.*, 18 (1995) 175-178.

1948 Masselter, S.M. and Zemmann, A.J.: Influence of organic solvents in coelectroosmotic capillary electrophoresis of phenols. *Anal. Chem.*, 67 (1995) 1047-1053.

See also 1915, 1950, 2628.

8. SUBSTANCES CONTAINING HETEROCYCLIC OXYGEN

8a. Flavonoids

See 2616.

9. OXO COMPOUNDS, ETHERS, EPOXIDES AND QUINONES

1949 Iwagami, S. and Sawabe, Y.: (Simultaneous determination of sennoside A and B in rhei rhizoma of sennae folium by micellar electrokinetic capillary chromatography). *Osaka-furitsu Koshu Eisei Kenkyusho Kenkyu Hokoku, Yakuji Shido-hen*, 28 (1994) 17-21; *C.A.*, 122 (1995) 197093z.

1950 Masselter, S., Zemmann, A. and Bobleter, O.: Analysis of lignin degradation products by capillary electrophoresis. *Chromatographia*, 40 (1995) 51-57.

See also 1946, 2616.

10. CARBOHYDRATES

10a. Mono and oligosaccharides. Structural studies

- 1951 Charkoudian, J., Pluskal, M. and Horton, R.: Membrane surface for carbohydrate analysis. *Anal. Lett.*, 28 (1995) 1055-1070.
- 1952 Edens, R.E., Fromm, J.R., Fromm, S.J., Lonhardt, R.J. and Weiler, J.M.: Two-dimensional affinity resolution electrophoresis demonstrates that three distinct heparin populations interact with antithrombin III. *Biochemistry*, 34 (1995) 2400-2407.
- 1953 Jackson, P.: The analysis of fluorophore-labeled saccharides by high-resolution polyacrylamide gel electrophoresis. *Adv. Electrophor.*, 7 (1994) 255-262; *C.A.*, 122 (1995) 182215s - a review with 163 refs.
- 1954 Kerns, R.J., Vlahov, I.R. and Linhardt, R.J.: Capillary electrophoresis for monitoring chemical reactions: sulfation and synthetic manipulation of sulfated carbohydrates. *Carbohydr. Res.*, 267 (1995) 143-152; *C.A.*, 122 (1995) 240283a.
- 1955 Mechref, Y., Ostrander, G.K. and El Rassi, Z.: Capillary electrophoresis of carboxylated carbohydrates. I. Selective precolumn derivatization of gangliosides with UV absorbing and fluorescent tags. *J. Chromatogr. A*, 695 (1995) 83-95.
- 1956 Oefner, P. and Scherz, H.: Capillary electrophoresis and thin-layer electrophoresis of carbohydrates. *Adv. Electrophor.*, 7 (1994) 155-224; *C.A.*, 122 (1995) 182214r - a review with 130 refs.
- 1957 Shimura, K. and Kassai, K.-i.: Determination of the affinity constants of concanavalin A for monosaccharides by fluorescence affinity probe capillary electrophoresis. *Anal. Biochem.*, 227 (1995) 186-194.
- 1958 Starr, C.M., Klock, J.C., Skop, E., Masada, I. and Giudici, T.: Fluorophore-assisted electrophoresis of urinary carbohydrates for the identification of patients with oligosaccharidosis- and mucopolysaccharidosis-type lysosomal storage diseases. *Glycosylation Dis.*, 1 (1994) 165-176; *C.A.*, 122 (1995) 27067b.
- 1959 Tsen, G., Halfter, W., Kröger, S. and Cole, G.J.: Agrin is a heparan sulfate proteoglycan. *J. Biol. Chem.*, 270 (1995) 3392-3399.
- 1960 Yutani, S., Toh, G. and Honda, S.: (Analysis of fructooligosaccharides by HPLC and HPCE). *Kuromatogurafi*, 15 (1994) 210-211; *C.A.*, 122 (1995) 127992t.

See also 2578.

10b. Polysaccharides, mucopolysaccharides, lipopolysaccharides

- 1961 Hayase, S., Tsukamoto, T. and Ushio, T.: (Direct analysis of glycosaminoglycans by high-performance capillary electrophoresis and separation in the presence of amphoteric additive). *Kuromatogurafi*, 15 (1994) 216-217; *C.A.*, 122 (1995) 155524k.
- 1962 Karamanos, N.K., Axelsson, S., Vanky, P., Tzanakakis, G.N. and Hjerpe, A.: Determination of hyaluronan and galactosaminoglycan disaccharides by high-performance capillary electrophoresis at the attomole level. Applications to analyses of tissue and cell culture proteoglycans. *J. Chromatogr. A*, 696 (1995) 295-305.

- 1963 Volpi, N., Cusmano, M. and Venturilli, T.: Qualitative and quantitative studies of heparin and chondroitin sulfates in normal human plasma. *Biochim. Biophys. Acta*, 1243 (1995) 49-58.

See also 2047.

10c. Glycoproteins and their constituents

- 1964 Choi-Miura, N.-H., Sano, Y., Oda, E., Nakano, Y., Tobe, T., Yanagishita, T., Taniyama, M., Katagiri, T. and Tomita, M.: Purification and characterization of a novel glycoprotein which has significant homology to heavy chains of inter- α -trypsin inhibitor family from human plasma. *J. Biochem. (Tokyo)*, 117 (1995) 400-407.
- 1965 Clark, M.A. and Weiss, A.S.: Hutchinson-Gilford progeria types defined by differential binding of lectin DSA. *Biochim. Biophys. Acta*, 1270 (1995) 142-148.
- 1966 Fassbender, K., Fassbender, M., Schaberg, T., Sobieska, M. and Müller, W.: Glycosylation of α_1 -acid glycoprotein in bacterial lung infections: distinct pattern in tuberculosis. *Clin. Chem. (Washington)*, 41 (1995) 472-473.
- 1967 Hanski, C., Hanski, M.-L., Zimmer, T., Ogorek, D., Devine, P. and Riecken, E.-O.: Characterization of the major sialyl-Lex^x-positive mucins present in colon, colon carcinoma, and sera of patients with colorectal cancer. *Cancer Res.*, 55 (1995) 928-933.
- 1968 Irimura, T. and Kawashima, H.: Detection and characterization of glycoprotein carbohydrate chains after electrophoretic separation. In: Dunbar, B.S. (Editor), *Protein Blotting*, IRL, Oxford, 1994, pp. 155-161; *C.A.*, 122 (1995) 155523j.
- 1969 Lampreave, F., Alava, M.A. and Piñeiro, A.: Lectin affinity immunoelectrophoresis of serum glycoproteins. *J. Chromatogr. A*, 698 (1995) 107-122 - a review with 132 refs.
- 1970 Lenoir, D., Ruggiero-Lopez, D., Louisot, P. and Biol, M.-C.: Developmental changes in intestinal glycosylation: nutrition-dependent multi-factor regulation of the fucosylation pathway at weaning time. *Biochim. Biophys. Acta*, 1234 (1995) 29-36.
- 1971 Madsen, P., Rasmussen, H.H., Flint, T., Gromov, P., Kruse, T.A., Honoré, B., Vorum, H. and Celis, J.E.: Cloning, Expression, and chromosome mapping of human galectin-7. *J. Biol. Chem.*, 270 (1995) 5823-5829.
- 1972 Mohanraj, D. and Ramakrishnan, S.: Cytotoxic effects of ricin without an interchain disulfide bond: genetic modification and chemical crosslinking studies. *Biochim. Biophys. Acta*, 1243 (1995) 399-406.
- 1973 Sato, T., Takio, K., Kobata, A., Greenwalt, D.E. and Furukawa, K.: Site-specific glycosylation of bovine butyrophilin. *J. Biochem. (Tokyo)*, 117 (1995) 147-157.
- 1974 Satoh, T., Abe, H., Sendai, Y., Iwata, H. and Hoshi, H.: Biochemical characterization of a bovine oviduct-specific sialo-glycoprotein that sustains sperm viability *in vitro*. *Biochim. Biophys. Acta*, 1266 (1995) 117-123.
- 1975 Sudou, A., Ozawa, M. and Muramatsu, T.: Lewis X structure increases cell substratum adhesion in L cells. *J. Biochem. (Tokyo)*, 117 (1995) 271-275.
- 1976 Weber, P.L., Kornfelt, T., Klausen, N.K. and Lunte, S.M.: Characterization of glycopeptides from recombinant coagulation factor VIIIa by high-performance liquid chromatography and capillary zone electrophoresis using ultraviolet and pulsed electrochemical detection. *Anal. Biochem.*, 225 (1995) 135-142.

11. ORGANIC ACIDS AND LIPIDS

11a. Organic acids and simple esters

- 1977 Erim, F.B., Xu, X. and Kraak, J.C.: Application of micellar electrokinetic chromatography and indirect UV detection for the analysis of fatty acids. *J. Chromatogr. A*, 694 (1995) 471-479.
- 1978 Schneede, J. and Ueland, P.M.: Application of capillary electrophoresis with laser-induced fluorescence detection for routine determination of methylmalonic acid in human serum. *Anal. Chem.*, 67 (1995) 812-819.

See also 2593, 2626.

11d. Lipoproteins and their constituents

- 1979 Baadenhuijsen, H., Demacker, P.N.M., Hessels, M., Boerma, G.J.M., Penders, T.J., Weykamp, C. and Willems, H.L.: Testing the accuracy of total cholesterol assays in an external quality-control program. Effect of adding sucrose to lyophilized control sera compared with use of fresh or frozen sera. *Clin. Chem. (Washington)*, 41 (1995) 724-730.
- 1980 Decossin, C., Tailleux, A., Fruchart, J.-C. and Fiévet, C.: Prevention of *in vitro* low-density lipoprotein oxidation by an albumin-containing Lp A-I subfraction. *Biochim. Biophys. Acta*, 1255 (1995) 31-38.
- 1981 Dobrian, A. and Simionescu, M.: Irreversibly glycosylated albumin alters the physico-chemical characteristics of low density lipoproteins of normal and diabetic subjects. *Biochim. Biophys. Acta*, 1270 (1995) 26-35.
- 1982 Greenspan, P., Mao, F.-w., Ryu, B.-H. and Gutman, R.L.: Advances in agarose gel electrophoresis of serum lipoproteins. *J. Chromatogr. A*, 698 (1995) 333-339 - a review with 56 refs.
- 1983 Marcovina, S.M., Albers, J.J., Gabel, B., Koschinsky, M.L. and Gaur, V.P.: Effect of the number of apolipoprotein(a) kringle 4 domains on immunochemical measurements of lipoprotein(a). *Clin. Chem. (Washington)*, 41 (1995) 246-255.
- 1984 Nagayoshi, A., Matsuki, N., Saito, H., Tsukamoto, K., Kaneko, K., Wakashima, M., Kinoshita, M., Yamanaka, M. and Teramoto, T.: Defect in assembly process of very-low-density lipoprotein in suncus liver: an animal model of fatty liver. *J. Biochem. (Tokyo)*, 117 (1995) 787-793.
- 1985 Nauck, M., Winkler, K., Wittman, C., Mayer, H., Luley, C., März, W. and Wieland, H.: Direct determination of lipoprotein(a) cholesterol by ultracentrifugation and agarose gel electrophoresis with enzymatic staining for cholesterol. *Clin. Chem. (Washington)*, 41 (1995) 731-738.
- 1986 Patel, D.D., Soutar, A.K. and Knight, B.L.: Abnormal structure and co-operative binding of low-density lipoprotein receptors containing the Glu-80 → Lys mutation. *Biochim. Biophys. Acta*, 1255 (1995) 285-292.
- 1987 Plonné, D., Heller, H., Kahlert, U. and Dargel, R.: Quantitative and qualitative characterization of apolipoprotein B containing lipoproteins produced by the visceral rat yolk sac in two different *in vitro* systems: organ culture and isolated epithelial cells in suspension culture. *Biochim. Biophys. Acta*, 1256 (1995) 71-80.

- 1988 Sigalov, A.B.: Cryopreservation and long-term storage of human low density lipoproteins. *Eur. J. Clin. Chem. Clin. Biochem.*, 33 (1995) 73-81.
- 1989 Skarlatos, S.I., Diverger, N., Rader, D. and Kruth, H.S.: Cholesterol efflux from human monocyte-derived macrophages in the presence of LpA-I:A-II. *Biochim. Biophys. Acta*, 1270 (1995) 19-25.

See also 2260, 2374.

13. STEROIDS

13d. Sterols

See 1985.

16. NITRO AND NITROSO COMPOUNDS

- 1990 Cao, J.: (Separation of complex nitrophenol sodium salts by micellar electrokinetic capillary chromatography). *Fenxi Huaxue*, 22 (1994) 1009-1011; *C.A.*, 122 (1995) 154012e.
- 1991 Oehrle, S.A.: Analysis of cationic ingredients and degradation products in liquid propellants by capillary ion electrophoresis (CIE). *J. Energ. Mater.*, 12 (1994) 197-209; *C.A.*, 122 (1995) 137592b.
- 1992 Tsikas, D., Böger, R.H., Bode-Böger, S.M., Brunner, G. and Frölich, J.C.: Formation of S-nitroso compounds from sodium nitroprusside, nitric oxide or nitrite and reduced thiols: analysis by capillary isotachopheresis. *J. Chromatogr. A*, 699 (1995) 363-369.

See also 1994.

17. AMINES, AMIDES AND RELATED NITROGEN COMPOUNDS

17a. Amines and polyamines

See 2593.

17b. Catecholamines and their metabolites

- 1993 Chang, H.-T. and Yeung, E.S.: Determination of catecholamines in single adrenal medullary cells by capillary electrophoresis and laser-induced native fluorescence. *Anal. Chem.*, 67 (1995) 1079-1083.
- 1994 Pyell, V. and Bütthorn, U.: Optimization strategies in micellar electrokinetic capillary chromatography. Optimization of the temperature of the separation capillary. *Chromatographia*, 40 (1995) 69-77.
- 1995 Takada, Y., Yoshida, M., Sakairi, M. and Koizumi, H.: (Analysis of substances related to brain function using capillary electrophoresis/MS). *Bunseki Kagaku*, 44 (1995) 241-246; *C.A.*, 122 (1995) 234745m.

- 1996 Yu, X., Luo, Z., Song, C. and Lodder, R.A.: (Determination of norepinephrine and epinephrine by high performance capillary electrophoresis). *Shandong Yike Daxue Xuebao*, 32 (1994) 347-349; *C.A.*, 122 (1995) 205369m.
18. AMINO ACIDS AND PEPTIDES; CHEMICAL STRUCTURE OF PROTEINS
- 18a. *Amino acids and their derivatives*
- 1997 Chan, K.C., Muschik, G.M. and Issaq, H.J.: Enantiomeric separation of amino acids using micellar electrokinetic chromatography after pre-column derivatization with the chiral reagent 1-(9-fluorenyl)-ethyl chloroformate. *Electrophoresis (Weinheim)*, 16 (1995) 504-509.
- 1998 Guo, Y., Colón, L.A., Dadoo, R. and Zare, R.N.: Analysis of underivatized amino acids by capillary electrophoresis using constant potential amperometric detection. *Electrophoresis (Weinheim)*, 16 (1995) 493-497.
- 1999 Issaq, H.J. and Chan, K.C.: Separation and detection of amino acids and their enantiomers by capillary electrophoresis: a review. *Electrophoresis (Weinheim)*, 16 (1995) 467-480 - a review with 81 refs.
- 2000 Jacobson, S.C. and Ramsey, J.M.: Microchip electrophoresis with sample stacking. *Electrophoresis (Weinheim)*, 16 (1995) 481-486.
- 2001 Kawamata, M. and Ohba, M.: (Study on analysis of D- and L-amino acids by capillary electrophoresis a preliminary examination on composition of carrier buffer solution). *Taisei Kensetsu Gijutsu Kenkyushoho*, 27 (1994) 315-318; *C.A.*, 122 (1995) 182487g.
- 2002 Lu, W., Yang, G. and Cole, R.B.: Determination of amino acids by on-line capillary electrophoresis-electrospray ionization mass spectrometry. *Electrophoresis (Weinheim)*, 16 (1995) 487-492.
- 2003 Michalke, B.: Capillary electrophoresis methods for clear identification of seleno amino acids in complex matrices like human milk. *Fresenius J. Anal. Chem.*, 351 (1995) 670-677.
- 2004 Minamoto, S. and Kanazawa, K.: Electrochemical determination of enzymic production of ultimate carcinogen from tryptophan pyrolysate by rat hepatic microsomes. *Anal. Biochem.*, 225 (1995) 143-148.
- 2005 Molle, D., Leonil, J. and Bouhallab, S.: Separation of two-proline-containing peptides by capillary electrophoresis. *Anal. Biochem.*, 225 (1995) 161-162.
- 2006 Zhou, J. and Lunte, S.M.: Direct determination of amino acids by capillary electrophoresis/electrochemistry using a copper microelectrode and zwitterionic buffers. *Electrophoresis (Weinheim)*, 16 (1995) 498-503.
- See also 2026, 2625, 2627.
- 18b. *Peptides, peptidic and proteinous hormones, growth factors*
- 2007 Cazorla, P., Aldudo, J., Haas, C., Vázquez, J., Valdivieso, F. and Bullido, M.J.: Location of an epitope shared by Alzheimer's amyloid peptide and brain creatine kinase using a newly developed monoclonal antibody. *Biochim. Biophys. Acta*, 1270 (1995) 149-156.
- 2008 Chen, F.-T.A.: Characterization of protease-catalyzed hydrolysis of cyanine-labeled angiotensin using capillary electrophoresis with laser-induced fluorescence detection. *Anal. Biochem.*, 225 (1995) 341-345.
- 2009 Cifuentes, A. and Poppe, H.: Effect of pH and ionic strength of running buffer on peptide behavior in capillary electrophoresis: theoretical calculation and experimental evaluation. *Electrophoresis (Weinheim)*, 16 (1995) 516-524.
- 2010 De Ceuninck, F., Willeput, J. and Corvol, M.: Purification and characterization of insulin-like growth factor II (IGFII) and an IGF II variant from human placenta. *J. Chromatogr. B*, 666 (1995) 203-214.
- 2011 Fuchs, B., Hecker, D. and Scheidtman, K.H.: Phosphorylation studies on rat p53 using the baculovirus expression system. Manipulation of the phosphorylation state with okadaic acid and influence on DNA binding. *Eur. J. Biochem.*, 228 (1995) 625-639.
- 2012 Fukuoka, E.: (Bradykinin metabolism in plasma and species differences determined by capillary electrophoresis). *Osaka Ika Daigaku Zasshi*, 53 (1994) 46-54; *C.A.*, 122 (1995) 75816s.
- 2013 Hegedus, Z., Ando, I., Toth, G.K., Varadi, G. and Monostori, E.: Application of polyacrylamide gel electrophoresis for analysis of oligopeptide phosphorylation *in vitro*. *BioTechniques*, 18 (1995) 631-636; *C.A.*, 122 (1995) 234657j.
- 2014 Karaplis, A.C., Lim, S., Baba, H., Arnold, A. and Kronenberg, H.M.: Inefficient membrane targeting, translocation, and proteolytic processing by signal peptidase of a mutant preproparathyroid hormone protein. *J. Biol. Chem.*, 270 (1995) 1629-1635.
- 2015 Lin, B.-C., Xu, X. and Luo, G.-A.: Zone broadening and simulation of migration process of peptides in capillary zone electrophoresis. *Sci. China, Ser. B*, 37 (1994) 807-819; *C.A.*, 122 (1995) 204202q.
- 2016 Matsubara, N., Koezuka, K. and Terabe, S.: Separation of eleven angiotensin II analogs by capillary electrophoresis with a nonionic surfactant in acidic media. *Electrophoresis (Weinheim)*, 16 (1995) 580-583.
- 2017 Orwar, O., Weber, S.G., Sandberg, M., Folestad, S., Tivesten, A. and Sundahl, M.: Fluorescence, photodestruction, photoionization and thermal degradation of α -phthalaldehyde/ β -mercaptoethanol-labelled aliphatic α -oligopeptides. *J. Chromatogr. A*, 696 (1995) 139-148.
- 2018 Pinto, D.M., Arriaga, E.A., Sia, S., Li, Z. and Dovichi, N.J.: Solid-phase fluorescent labeling reaction of picomole amounts of insulin in very dilute solutions and their analysis by capillary electrophoresis. *Electrophoresis (Weinheim)*, 16 (1995) 534-540.
- 2019 Ruberto, M.A. and Grayeski, M.L.: Investigation of acridinium labeling for chemiluminescence detection of peptides separated by capillary electrophoresis. *J. Microcolumn Sep.*, 6 (1994) 545-550; *C.A.*, 122 (1995) 75776d.
- 2020 Schultz, N.M., Huang, L. and Kennedy, R.T.: Capillary electrophoresis-based immunoassay to determine insulin content and insulin secretion from single islets of Langerhans. *Anal. Chem.*, 67 (1995) 924-929.
- 2021 Strausbauch, M.A., Madden, B.J., Wettstein, P.J. and Landers, J.P.: Sensitivity enhancement and second-dimensional information from solid phase extraction-capillary electrophoresis of entire high-performance liquid chromatography fractions. *Electrophoresis (Weinheim)*, 16 (1995) 541-548.

- 2022 Tadey, T. and Purdy, W.C.: Capillary electrophoretic resolution of phosphorylated peptide isomers using micellar solutions and coated capillaries. *Electrophoresis (Weinheim)*, 16 (1995) 574-579.
- 2023 Thorsteinsdóttir, M., Beijersten, I. and Westerlund, D.: Capillary electroseparations of enkephalin-related peptides and protein kinase A peptide substrates. *Electrophoresis (Weinheim)*, 16 (1995) 564-573.
- 2024 Thorsteinsdóttir, M., Isaksson, R. and Westerlund, D.: Performance of amino-silylated fused-silica capillaries for the separation of enkephalin-related peptides by capillary zone electrophoresis and micellar electrokinetic chromatography. *Electrophoresis (Weinheim)*, 16 (1995) 557-563.

See also 1879, 1995, 2038, 2047, 2057, 2168, 2240, 2469.

18c. Elucidation of structure of proteins and enzymes

- 2025 Adamson, N.J. and Reynolds, E.C.: High performance capillary electrophoresis of casein phosphopeptides containing 2-5 phosphoserine residues: relationship between absolute electrophoretic mobility and peptide charge and size. *Electrophoresis (Weinheim)*, 16 (1995) 525-528.
- 2026 Kim, N.J., Kim, J.H. and Lee, K.-J.: Application of capillary electrophoresis to amino acid sequencing of peptide. *Electrophoresis (Weinheim)*, 16 (1995) 510-515.
- 2027 Lopez, M.F., Barry, P., Sawilovich, W.B., Hines, T. and Skea, W.M.: High resolution 2-D peptide mapping with subsequent analysis of peptides by microsequencing or lectin binding directly from PVDF membrane blots. *Appl. Theor. Electrophor.*, 4 (1994) 95-102; *C.A.*, 122 (1995) 50535m.
- 2028 Miyauchi, M., Tozawa, K. and Yoshida, M.: F₁-ATPase α -subunit made up from two fragments (1-395, 396-503) is stabilized by ATP and complexes containing it obey altered kinetics. *Biochim. Biophys. Acta*, 1229 (1995) 225-232.
- 2029 Nakayama, H., Kanai, M., Seta, K., Isobe, T. and Okuyama, T.: (Primary structure analysis of proteins by combined two-dimensional electrophoresis-capillary column HPLC-ESI/TSQMS). *Kuro-matogurafu*, 15 (1994) 264-265; *C.A.*, 122 (1995) 127993u.
- 2030 Pedersen, J., Filimonova, M., Roepstorff, P. and Biedermann, K.: (Isoforms of *Serratia marcescens* nuclease produced by natural and recombinant strains. Comparative characterization by plasma desorption mass spectrometry). *Biokhimiya (Moscow)*, 60 (1995) 450-461.
- 2031 Rawal, N., Rajpurohit, R., Lischwe, M.A., Williams, K.R., Paik, W.K. and Kim, S.: Structural specificity of substrate for S-adenosylmethionine:protein arginine N-methyltransferases. *Biochim. Biophys. Acta*, 1248 (1995) 11-18.
- 2032 Van Riel, J. and Olieman, C.: Determination of caseinomacropptide with capillary zone electrophoresis and its application to the detection and estimation of rennet whey solids in milk and buttermilk powder. *Electrophoresis (Weinheim)*, 16 (1995) 529-533.

See also 2047, 2057, 2169, 2186, 2190, 2213, 2306, 2351.

19. PROTEINS

19a. General techniques

- 2033 Ali, R., Sayeed, S.A. and Khan, A.A.: A sensitive novel staining agent for the resolved proteins on PAGE. *Int. J. Pept. Protein Res.*, 45 (1995) 97-99; *C.A.*, 122 (1995) 182371q.
- 2034 Cash, P.: Protein mutations revealed by two-dimensional electrophoresis. *J. Chromatogr. A*, 698 (1995) 203-224 - a review with 149 refs.
- 2035 Chae, K.S. and Lenhoff, A.M.: Computation of the electrophoretic mobility of proteins. *Biophys. J.*, 68 (1995) 1120-1127; *C.A.*, 122 (1995) 155538t.
- 2036 Cordwell, S.J., Wilkins, M.R., Cerpa-Poljak, A., Gooley, A.A., Duncan, M., Williams, K.L. and Humphery-Smith, I.: Cross-species identification of proteins separated by two-dimensional gel electrophoresis using matrix-assisted laser desorption ionisation/time-of-flight mass spectrometry and amino acid composition. *Electrophoresis (Weinheim)*, 16 (1995) 438-443.
- 2037 Corradini, D. and Cannarsa, G.: N,N,N',N'-Tetramethyl-1,3-butanediamine as effective running electrolyte additive for efficient electrophoretic separation of basic proteins in bare fused-silica capillaries. *Electrophoresis (Weinheim)*, 16 (1995) 630-635.
- 2038 Fadden, P. and Haystead, T.A.J.: Quantitative and selective fluorophore labeling of phosphoserine on peptides and proteins: characterization at the attomole level by capillary electrophoresis and laser-induced fluorescence. *Anal. Biochem.*, 225 (1995) 81-88.
- 2039 Gao, J., Gomez, F.A., Haerter, R. and Whitesides, G.M.: Determination of the effective charge of a protein in solution by capillary electrophoresis. *Proc. Natl. Acad. Sci. U.S.A.*, 91 (1994) 12027-12030; *C.A.*, 122 (1995) 27085f.
- 2040 Glatz, Z., Janiczek, O., Wimmerova, M. and Novotny, M.V.: Detection of quinoproteins after electrophoresis in the presence of urea or SDS. *Biochem. Mol. Biol. Int.*, 35 (1995) 1-10; *C.A.*, 122 (1995) 127923w.
- 2041 Guttman, A.: On the separation mechanism of capillary sodium dodecyl sulfate-gel electrophoresis of proteins. *Electrophoresis (Weinheim)*, 16 (1995) 611-616.
- 2042 Heegaard, N.H.H.: Characterization of biomolecules by electrophoretic analysis of reversible interactions. *Appl. Theor. Electrophor.*, 4 (1994) 43-63; *C.A.*, 122 (1995) 50346a - a review with many refs.
- 2043 Heller, C.: Capillary electrophoresis of proteins and nucleic acids in gels and entangled polymer solutions. *J. Chromatogr. A*, 698 (1995) 19-31 - a review with 105 refs.
- 2044 Henzel, W.J., Grimley, C., Bourell, J.H., Billeci, T.M., Wong, S.C. and Stults, J.T.: Analysis of two-dimensional gel proteins by mass spectrometry and microsequencing. *Methods (San Diego)*, 6 (1994) 239-247; *C.A.*, 122 (1995) 27209z.
- 2045 Hess, D. and Aebersold, R.: Internal sequence analysis of proteins separated by polyacrylamide gel electrophoresis. *Methods (San Diego)*, 6 (1994) 227-238; *C.A.*, 122 (1995) 272087.
- 2046 Hillebrandt, S. Streffer, C.: Quantitative computerized analysis of silver-stained and Coomassie Blue-stained two-dimensional protein maps. *Z. Naturforsch. C: Biosci.*, 49 (1994) 849-855; *C.A.*, 122 (1995) 74084q.

- 2047 Huang, M., Plocek, J. and Novotny, M.V.: Hydrolytically stable cellulose-derivative coatings for capillary electrophoresis of peptides, proteins and glycoconjugates. *Electrophoresis (Weinheim)*, 16 (1995) 396-401.
- 2048 Huang, T.-L., Shieh, P.C.H. and Cooke, N.: Isoelectric focusing of proteins in capillary electrophoresis with pressure-driven mobilization. *Chromatographia*, 39 (1994) 543-548.
- 2049 Jacobson, G.: Protein blotting using semi-dry electrophoretic transfer equipment. In: Dunbar, B.S. (Editor), *Protein Blotting*, IRL, Oxford, 1994, pp. 53-70; C.A., 122 (1995) 127725h - a review with 19 refs.
- 2050 Kasai, K.-i.: (Intelligent chromatography and electrophoresis as tools for understanding of biomolecules). *Kuromatogurafi*, 15 (1994) 206-209; C.A., 122 (1995) 127787e.
- 2051 Krueger, E., Radau, B. and Roenn, D.: (Investigation of beers and beer heads by methods employed in protein chemistry). *Monatsschr. Brauwiss.*, 46 (1993) 312-319; C.A., 122 (1995) 131260v.
- 2052 Kubo, K. and Takagi, T.: Modulation of the behavior of a protein in polyacrylamide gel electrophoresis in the presence of dodecyl sulfate by varying the cations. *Anal. Biochem.*, 224 (1995) 572-579.
- 2053 Kulish, D.M., Klyushnichenko, V.E. and Vul'fson, A.N.: (Separation of proteins by liquid chromatography and capillary electrophoresis). *Zh. Fiz. Khim.*, 68 (1994) 1841-1847; C.A., 122 (1995) 50407w.
- 2054 Kyd, J.M., Taylor, D. and Cripps, A.W.: Conservation of immune responses to proteins isolated by preparative polyacrylamide gel electrophoresis from the outer membrane of nontypeable *Haemophilus influenzae*. *Infect. Immun.*, 62 (1994) 5652-5658; C.A., 122 (1995) 27077e.
- 2055 Lee, V.H. and Dunbar, B.S.: Sample preparation for protein electrophoresis and transfer. In: Dunbar, B.S. (Editor), *Protein Blotting*, IRL, Oxford, 1994, pp. 87-103; C.A., 122 (1995) 127901n.
- 2056 Liu, Y., Fu, R. and Gu, J.: Capillary zone electrophoretic separation of proteins using a column coated with epoxy polymer. *J. Chromatogr. A*, 694 (1995) 498-506.
- 2057 Piccoli, G., Fiorani, M., Biagiarelli, B., Palma, F., Vallorani, L., de Bellis, R. and Stocchi, V.: High-performance capillary electrophoretic separation of proteins and peptides using a bonded hydrophilic phase capillary. *Electrophoresis (Weinheim)*, 16 (1995) 625-629.
- 2058 Rider, M.H., Puype, M., van Damme, J., Gevaert, K., de Boeck, S., D'Alayer, J., Rasmussen, H.H., Celis, J.E. and Vanderckhove, J.: An agarose-based gel-concentration system for microsequence and mass spectrometric characterization of proteins previously purified in polyacrylamide gels starting at low picomole levels. *Eur. J. Biochem.*, 230 (1995) 258-265.
- 2059 Salazar, R., Brandt, R. and Krantz, S.: Expression of fructosyllysine receptors on human monocytes and monocyte-like cell lines. *Biochim. Biophys. Acta*, 1266 (1995) 57-63.
- 2060 Sarmiento, M.: High resolution, standard format two-dimensional protein electrophoresis using disposable glass micropipettes and non-dedicated equipment. *Electrophoresis (Weinheim)*, 16 (1995) 431-437.
- 2061 Schaeffer, H.: Denaturing electrophoretic techniques. In: Von Jagow, G. and Schaeffer, H. (Editors), *Pract. Guide Membr. Protein Purif.*, Academic, San Diego, 1994, pp. 59-79; C.A., 122 (1995) 155328z.
- 2062 Schaeffer, H.: Native gel electrophoresis. In: Von Jagow, G. (Editor), *Pract. Guide Membr. Protein Purif.*, Academic, San Diego, 1994, pp. 81-104; C.A., 122 (1995) 155527p.
- 2063 Senda, M., Sasaki, T. and Hiyama, T.: Capillary electrophoresis of redox proteins and adenine nucleotides. *Kuromatogurafi*, 15 (1994) 169-174; C.A., 122 (1995) 27071y.
- 2064 Shaw, S.M. and Crabbe, M.J.C.: Monitoring the progress of nonenzymic glycation *in vitro*. *Int. J. Pept. Protein Res.*, 44 (1994) 594-602; C.A., 122 (1995) 50702p.
- 2065 Shoji, M., Kato, M. and Hashizume, S.: Electrophoretic recovery of proteins from polyacrylamide gel. *J. Chromatogr. A*, 698 (1995) 145-162 - a review with 99 refs.
- 2066 Sreeramulu, G. and Singh, N.K.: Destaining of Coomassie Brilliant Blue R-250-stained polyacrylamide gels with sodium chloride solutions. *Electrophoresis (Weinheim)*, 16 (1995) 362-363.
- 2067 Szewczyk, B. and Summers, D.F.: Efficient elution of purified proteins from polyvinylidene difluoride membranes (immobilon) after transfer from SDS-PAGE and their use as immunogens. *Mol. Biotechnol.*, 2 (1994) 129-134; C.A., 122 (1995) 27069d.
- 2068 Szoko, E.: (Separation of proteins by capillary zone electrophoresis-the effect of dextran polymers on the selectivity). *Gyogyaszerteszet*, 38 (1994) 715-718; C.A., 122 (1995) 75821q.
- 2069 Tang, J. and Huang, J.: (Effect of sample overloading on the separation of proteins by high performance capillary gel electrophoresis). *Gaojishu Tongxun*, 4 (1994) 29-32; C.A., 122 (1995) 75842x.
- 2070 Theos, C.W.: Protein electrophoresis in aqueous two-phase systems. Avail. *Univ. Microfilms Int.*, Order No. DA9434086, 1994, 236 pp.; C.A., 122 (1995) 182496j.
- 2071 Thornton, J.R., Daum, H.A., III and Case S.T.: Agarose gel electrophoresis of high molecular mass protein complexes. *BioTechniques*, 18 (1995) 324-327; C.A., 122 (1995) 127922v.
- 2072 Tsonis, P.A. and del Rio-Tsonis, K.: Protein separation techniques in the study of tissue regeneration. *J. Chromatogr. A*, 698 (1995) 361-367.
- 2073 Vuillard, L., Marret, N. and Rabilloud, T.: Enhancing protein solubilization with nondetergent sulfobetaines. *Electrophoresis (Weinheim)*, 16 (1995) 295-297.
- 2074 Wheat, T.E.: Principles and practice of peptide analysis with capillary zone electrophoresis. *Methods Mol. Biol. (Totowa)*, 36(Peptide Analysis Protocols) (1994) 65-83; C.A., 122 (1995) 75703c.
- See also 1818, 1838, 1861, 1866, 1868, 1875, 1877, 1879, 1888, 1890, 1899, 1922, 2084, 2181, 2384, 2482, 2618, 2648, 2659.
- 19b. *Proteins of cells, viruses and subcellular particles*
- 2075 Azem, A., Shaked, I., Rosenbusch, J.P. and Daniel, E.: Cross-linking of porin with glutaraldehyde: a test for the adequacy of premises of cross-linking theory. *Biochim. Biophys. Acta*, 1243 (1995) 151-156.

- 2076 Bentlage, H.A.C.M., Janssen, A.J.M., Chomyn, A., Attardi, G., Walker, J.E., Schägger, H., Sengers, R.C.A. and Trijbels, F.J.M.: Multiple deficiencies of mitochondrial DNA- and nuclear-encoded subunits of respiratory NADH dehydrogenase detected with peptide- and subunit-specific antibodies in mitochondrial myopathies. *Biochim. Biophys. Acta*, 1234 (1995) 63-73.
- 2077 Boensch, C., Kuo, M., Connolly, D.T., Huang, S.S. and Huang, J.S.: Identification, purification, and characterization of cell-surface retention sequence-binding proteins from human SK-Hep cells and bovine liver plasma membranes. *J. Biol. Chem.*, 270 (1995) 1807-1816.
- 2078 Cai, S.-J., McAndrew, R.S., Leonard, B.P., Chapman, K.D. and Pidgeon, C.: Rapid purification of cotton seed membrane-bound N-acylphosphatidylethanolamine synthase by immobilized artificial membrane chromatography. *J. Chromatogr. A*, 696 (1995) 49-62.
- 2079 Duncan, R.F., Cavener, D.R. and Qu, S.: Heat shock effects on phosphorylation of protein synthesis initiation factor proteins eIF-4E and eIF-2 α in *Drosophila*. *Biochemistry*, 34 (1995) 2985-2997.
- 2080 Eriksson, H., Ridderstråle, M., Degerman, E., Ekholm, D., Smith, C.J., Manganiello, V.C., Belfrage, P. and Tornqvist, H.: Evidence for the key role of the adipocyte cGMP-inhibited cAMP phosphodiesterase in the antilipolytic action of insulin. *Biochim. Biophys. Acta*, 1266 (1995) 101-107.
- 2081 Fine, A., Panchenko, M.P., Smith, B.D., Yu, Q. and Goldstein, R.H.: Discordant regulation of transforming growth factor- β receptors by prostaglandin E₂. *Biochim. Biophys. Acta*, 1261 (1995) 19-24.
- 2082 Freiburghaus, A.U.: Two-dimensional polyacrylamide gel electrophoresis of membrane proteins. *Mol. Biotechnol.*, 2 (1994) 281-293; *C.A.*, 122 (1995) 127900m.
- 2083 Gaigg, B., Simbeni, R., Hrastnik, C., Paltauf, F. and Daum, G.: Characterization of a microsomal subfraction associated with mitochondria of the yeast, *Saccharomyces cerevisiae*. Involvement in synthesis and import of phospholipids into mitochondria. *Biochim. Biophys. Acta*, 1234 (1995) 214-220.
- 2084 Gianazza, E., Coari, P., Lovati, M.R., Manzoni, C., Ghibaldi, E. and Salmons, M.: Basic proteins and basic membranes. Adjusting blotting and staining conditions to Immobilon CD. *J. Chromatogr. A*, 698 (1995) 351-359.
- 2085 Gimenez, B., Amarasekera, D., Argo, E. and Cash, P.: Analysis of protein synthesis by two-dimensional gel electrophoresis in T cells persistently infected with coxsackie B virus. *Electrophoresis (Weinheim)*, 16 (1995) 317-321.
- 2086 Hills, D. and Crane-Robinson, C.: Baculovirus expression of human basic fibroblast growth factor from a synthetic gene: role of the Kozak consensus and comparison with bacterial expression. *Biochim. Biophys. Acta*, 1260 (1995) 14-20.
- 2087 Kvannes, J., Eikhom, T.S. and Flatmark, T.: On the mechanism of stimulation of peroxisomal β -oxidation in rat heart by partially hydrogenated fish oil. *Biochim. Biophys. Acta*, 1255 (1995) 39-49.
- 2088 Lopez-Moratalla, N., del Mar Calonge, M., López-Zabalza, M.J., Pérez-Mediavilla, L.A., Subirá, M.L. and Santiago, E.: Activation of human lymphomononuclear cells by peptides derived from extracellular matrix proteins. *Biochim. Biophys. Acta*, 1265 (1995) 181-188.
- 2089 Manabe, T., Yamamoto, H. and Kawai, M.: Studies on the procedure for the construction of cellular protein databases employing micro two-dimensional electrophoresis: an HL-60 protein database. *Electrophoresis (Weinheim)*, 16 (1995) 407-422.
- 2090 Matsumoto, K., Okazaki, H. and Nakamura, T.: Novel function of prostaglandins as inducers of gene expression of HGF and putative mediators of tissue regeneration. *J. Biochem. (Tokyo)*, 117 (1995) 458-464.
- 2091 Mithieux, G., Ajzannay, A. and Minassian, C.: Identification of membrane-bound phosphoglucosyltransferase and glucose-6 phosphatase by ³²P-labeling of rat liver microsomal membrane proteins with ³²P-glucose-6 phosphate. *J. Biochem. (Tokyo)*, 117 (1995) 908-914.
- 2092 Naylor, D.J., Ryan, M.T., Condrón, R., Hoogenraad, N.J. and Høj, P.B.: Affinity-purification and identification of GrpE homologues from mammalian mitochondria. *Biochim. Biophys. Acta*, 1248 (1995) 75-79.
- 2093 Nonaka, T., Matsumoto, H., Shimada, W., Miyagi, I., Okada, K., Fukao, H., Ueshima, S., Kikuchi, H., Tanaka, S. and Matsuo, O.: Effect of cyclic AMP on urokinase-type plasminogen activator receptor and fibrinolytic factors in a human osteoblast-like cell line. *Biochim. Biophys. Acta*, 1266 (1995) 50-56.
- 2094 Ono, T.-a., Noguchi, T. and Nakajima, Y.: Characteristic changes of function and structure of Photosystem II during strong-light photoinhibition under aerobic conditions. *Biochim. Biophys. Acta*, 1229 (1995) 239-248.
- 2095 Palm, L., Andersen, J., Rahbek-Nielsen, H., Hansen, T.S., Kristiansen, K. and Hojrup, P.: The phosphorylated ribosomal protein S7 in *Tetrahymena* is homologous with mammalian S4 and the phosphorylated residues are located in the C-terminal region. Structural characterization of proteins separated by two-dimensional polyacrylamide gel electrophoresis. *J. Biol. Chem.*, 270 (1995) 6000-6005.
- 2096 Pemberton, P.W., Lobley, R.W., Holms, R., Soerensen, S.H., Simpson, K.W. and Batt, R.M.: Characterization of microvillar membrane proteins of dog small intestine by two-dimensional electrophoresis. *Comp. Biochem. Physiol., B: Biochem. Mol. Biol.*, 110B (1995) 483-492; *C.A.*, 122 (1995) 209055d.
- 2097 Pestonjamas, K.N. and Mehta, N.G.: Neutral polymers elicit, and antibodies to spectrin, band 4.1 protein and cytoplasmic domain of band 3 protein inhibit the concanavalin A-mediated agglutination of human erythrocytes. *Biochim. Biophys. Acta*, 1235 (1995) 10-20.
- 2098 Puchkova, L.V., Aleinikova, T.D., Zakharova, E.T., Konopistseva, L.A., Tsybalenko, N.V. and Gaitskhoki, V.S.: (Regulation of biosynthesis of molecular forms of ceruloplasmin in ontogenesis of the rats). *Biokhimiya (Moscow)*, 59 (1994) 1304-1311.
- 2099 Sato, E.F., Edashige, K., Inoue, M. and Utsumi, K.: Okadaic acid increased annexin I and induced differentiation of human promyelocytic leukemia cells. *Biochim. Biophys. Acta*, 1266 (1995) 23-30.
- 2100 Sugo, T., Nakamikawa, C., Tanabe, S. and Matsuda, M.: Activation of prothrombin by factor Xa bound to the membrane surface of human umbilical vein endothelial cells: its catalytic efficiency is similar to that of prothrombinase complex on platelets. *J. Biochem. (Tokyo)*, 117 (1995) 244-250.

- 2101 Takano, M., Yokoyama, K., Yayama, K. and Okamoto, H.: Murine fibroblasts synthesize and secrete kininogen in response to cyclic-AMP, prostaglandin E₂ and tumor necrosis factor. *Biochim. Biophys. Acta*, 1265 (1995) 189-195.
- 2102 Turner, G.P. and Kneale, G.G.: Site-directed mutagenesis of the M13 gene 5 protein: the role of Arg-21, Tyr-26 and Tyr-41. *Biochim. Biophys. Acta*, 1260 (1995) 79-84.
- 2103 Weil, R., Cloutier, J.-F., Fournel, M. and Veillette, A.: Regulation of Zap-70 by Src family tyrosine protein kinases in an antigen-specific T-cell line. *J. Biol. Chem.*, 270 (1995) 2791-2799.
- See also 1970, 2072, 2108, 2150, 2234, 2245, 2478.
- 19c. *Proteins synthesized by genetic manipulation, monoclonal antibodies*
- 2104 Bill, R.M., Winter, P.C., McHale, C.M., Hodges, V.M., Elder, G.E., Caley, J., Flitsch, S.L., Bicknell, R. and Lappin, T.R.J.: Expression and mutagenesis of recombinant human and murine erythropoietins in *Escherichia coli*. *Biochim. Biophys. Acta*, 1261 (1995) 35-43.
- 2105 Cvekl, A., McDermott, J.B. and Piatigorsky, J.: cDNA encoding a chicken protein (CRP1) with homology to hnRNP type A/B. *Biochim. Biophys. Acta*, 1261 (1995) 290-292.
- 2106 Kwon, K.-S., Lee, S. and Yu, M.-H.: Refolding of α -antitrypsin expressed as inclusion bodies in *Escherichia coli*: characterization of aggregation. *Biochim. Biophys. Acta*, 1247 (1995) 179-184.
- 2107 Mukai, H., Mori, K., Takanaga, H., Kitagawa, M., Shibata, H., Shimakawa, M., Miyahara, M. and Ono, Y.: *Xenopus* PKN: cloning and sequencing of the cDNA and identification of conserved domains. *Biochim. Biophys. Acta*, 1261 (1995) 296-300.
- 2108 Samal, B.B., Arakawa, T., Boone, T.C., Jones, T., Prestrelski, S.J., Narhi, L.O., Wen, J., Stearns, G.W., Crandall, C.A., Pope, J. and Suggs, S.: High level expression of human leukemia inhibitory factor (LIF) from a synthetic gene in *Escherichia coli* and the physical and biological characterization of the protein. *Biochim. Biophys. Acta*, 1260 (1995) 27-34.
- 2109 Strege, M.A. and Lagu, A.L.: Capillary electrophoretic separations of biotechnology-derived proteins in *E. coli* fermentation broth. *Electrophoresis (Weinheim)*, 16 (1995) 642-646.
- 2110 Yao, Y.J., Loh, K.C., Chung, M.C.M. and Li, S.F.Y.: Analysis of recombinant human tumor necrosis factor beta by capillary electrophoresis. *Electrophoresis (Weinheim)*, 16 (1995) 647-653.
- 2111 Zhu, A., Wang, Z.-K. and Goldstein, J.: Identification of tyrosine 108 in coffee bean α -galactosidase as an essential residue for the enzyme activity. *Biochim. Biophys. Acta*, 1247 (1995) 260-264.
- See also 2028, 2086, 2102, 2150, 2182, 2231, 2268, 2302, 2404, 2446.
- 19d. *Microbial and plant proteins*
- 2112 Black, J. and Durig, L.: Sequential method for the electrophoretic analysis of barley varieties, including computer storage and matching of banding patterns. *J. Am. Soc. Brew. Chem.*, 53 (1995) 19-23; C.A., 122 (1995) 158914y.
- 2113 Buchrieser, C., Gangar, V.V., Murphree, R.L., Tamplin, M.L. and Kaspar, C.W.: Multiple *Vibrio vulnificus* strains in oysters as demonstrated by clamped homogeneous electric field gel electrophoresis. *Appl. Environ. Microbiol.*, 61 (1995) 1163-1168; C.A., 122 (1995) 179593h.
- 2114 Cooke, R.J.: Gel electrophoresis for the identification of plant varieties. *J. Chromatogr. A*, 698 (1995) 281-299 - a review with 68 refs.
- 2115 Costas, M., Holmes, B., Ganner, M., On, S.L.W., Hoffman, P.N., Worsley, M.A. and Panigrahi, H.: Identification of outbreak-associated and other strains of *Clostridium difficile* by numerical analysis of SDS-PAGE protein patterns. *Epidemiol. Infect.*, 113 (1994) 1-12; C.A., 122 (1995) 27320d.
- 2116 Deb, A., Bhattacharyya, D. and Das, J.: A 25-kDa β -lactam-induced outer membrane protein of *Vibrio cholerae*. Purification and characterization. *J. Biol. Chem.*, 270 (1995) 2914-2920.
- 2117 Demirevska-Kepova, K., Metodiev, M. and Tsekov, T.: Comparison of gel electrophoretic and immunochemical measurements of Rubisco quantities in leaves. *Dokl. Bulg. Akad. Nauk.*, 46 (1993) 105-108; C.A., 122 (1995) 101720c.
- 2118 Descheemaeker, P., Pot, B., Ledebuer, A.M., Verrips, T. and Kersters, K.: Comparison of the *Lactococcus lactis* differential medium (DCL) and SDS-PAGE of whole-cell proteins for the identification of lactococci to subspecies level. *Syst. Appl. Microbiol.*, 17 (1994) 459-466; C.A., 122 (1995) 155621q.
- 2119 Durner, J. and Böger, P.: Ubiquitin in the prokaryote *Anabaena variabilis*. *J. Biol. Chem.*, 270 (1995) 3720-3725.
- 2120 Farber, S.P.: (Seed 12 S globulins in identification of cabbage cultivars). *Dokl. Ross. Akad. S-kh. Nauk.*, (1994) 8-10; C.A., 122 (1995) 235441w.
- 2121 Giometti, C.S., Tollaksen, S.L., Mukund, S., Zhou, Z.H., Ma, K., Mai, X. and Adams, M.W.W.: Two-dimensional gel electrophoresis mapping of proteins isolated from the hyperthermophile *Pyrococcus furiosus*. *J. Chromatogr. A*, 698 (1995) 341-349.
- 2122 Golitsyn, V.M., Teten'kin, V.L., Elanskaya, I.V. and Gulyaev, B.A.: (Spectral properties of the cyanobacterium *Cyanoecystis* sp. PCC 6803 mutants devoid of the PhotoSystem 2 activity). *Bio-khimiya (Moscow)*, 60 (1995) 485-490.
- 2123 Jekow, P., Fromme, P., Witt, H.T. and Saenger, W.: Photosystem I from *Synechococcus elongatus*: preparation and crystallization of monomers with varying subunit compositions. *Biochim. Biophys. Acta*, 1229 (1995) 115-120.
- 2124 Kamo, M., Kawakami, T., Miyatake, N. and Tsugita, A.: Separation and characterization of *Arabidopsis thaliana* proteins by two-dimensional gel electrophoresis. *Electrophoresis (Weinheim)*, 16 (1995) 423-430.
- 2125 Kondo, S., Matsumoto, T., Yokoyama, Y., Ohmori, I. and Suzuki, H.: The shortest isoform of human vascular endothelial growth factor/vascular permeability factor (VEGF/VPF₁₂₁) produced by *Saccharomyces cerevisiae* promotes both angiogenesis and vascular permeability. *Biochim. Biophys. Acta*, 1243 (1995) 195-202.
- 2126 Lookhart, G. and Bean, S.: A fast method for wheat cultivar differentiation using capillary zone electrophoresis. *Cereal Chem.*, 72 (1995) 42-47; C.A., 122 (1995) 155533n.
- 2127 Mendoza, J.A., Martinez, J.L. and Horowitz, P.M.: Tetradameric chaperonin 60 can be assembled *in vitro* from monomers in a process that is ATP independent. *Biochim. Biophys. Acta*, 1247 (1995) 209-214.

- 2128 Michaud, D. and Assetin, A.: Application to plant proteins of gel electrophoretic methods. *J. Chromatogr. A*, 698 (1995) 263-279 - a review with 145 refs.
- 2129 Phan-Thanh, L. and Gormon, T.: Analysis of heat and cold shock proteins in *Listeria* by two-dimensional electrophoresis. *Electrophoresis (Weinheim)*, 16 (1995) 444-450.
- 2130 Seki, K., Kobayashi, K. and Masuda, S.: A simple procedure for comparison of protein band patterns on SDS-PAGE between *Staphylococcus aureus* strains. *Zentralbl. Bakteriol., Suppl.*, 26 (1994) 61-63; *C.A.*, 122 (1995) 234647f.
- 2131 Takeda, T., Kurasawa, Y., Watanabe, Y. and Numata, O.: Polymerization of highly purified *Tetrahymena* 14-nm filament protein/citrate synthase into filaments and its possible role in regulation of enzymatic activity. *J. Biochem. (Tokyo)*, 117 (1995) 869-874.
- 2132 Tsiomenko, A.B., Tuymetova, G.P., Eldarov, M.A., Korolev, S.V., Skryabin, K.G. and Kulaev, I.S.: (The prosegment of yeast α -factor directs the human growth hormone to the culture medium of *Saccharomyces cerevisiae*). *Biokhimiya (Moscow)*, 59 (1994) 1675-1688.
- 2133 Wang, X., Yang, S., Xi, Y. and Cao, Y.: (Electrophoresis differentiation of oleosins between different family and genus of plants). *Xibei Shifan Daxue Xuebao, Ziran Kexueban*, 30 (1994) 53-56; *C.A.*, 122 (1995) 75841w.
- 2134 Werner, W.E., Wiktorowicz, J.E. and Kasarda, D.D.: Wheat varietal identification by capillary electrophoresis of gliadins and high molecular weight glutenin subunits. *Cereal Chem.*, 71 (1994) 397-402; *C.A.*, 122 (1995) 30176s.
- 2135 Yu, S.-G., Stefansson, H., Romanowska, E. and Albertsson, P.-A.: Two dimensional electrophoresis of thylakoid membrane proteins and its application to microsequencing. *Photosynth. Res.*, 41 (1994) 475-486; *C.A.*, 122 (1995) 50499c.
- 2136 Zimacheva, A.V. and Mosolov, V.V.: (Inhibitors of cysteine proteinases from soya seeds). *Biokhimiya (Moscow)*, 60 (1995) 118-123.
- See also 1957, 2066, 2104, 2109, 2395, 2623.
- 19e. *Proteins of blood, serum and blood cells*
- 2137 Brumeanu, T.-D., Zaghouni, H. and Bona, C.: Purification of antigenized immunoglobulins derivatized with monomethoxy-polyethylene glycol. *J. Chromatogr. A*, 696 (1995) 219-225.
- 2138 Cheema, A.A., Malik, M.A. and Javed, M.A.: Identification of plasma proteins of broiler chicks of *Gallus domesticus* by polyacrylamide gel electrophoresis using different dyes. *Pak. J. Zool.*, 26 (1994) 192-195; *C.A.*, 122 (1995) 127896q.
- 2139 Colls, J., Betterle, C., Volpato, M., Prentice, L., Smith, B.R. and Furmaniak, J.: Immunoprecipitation assay for autoantibodies to steroid 21-hydroxylase in autoimmune adrenal diseases. *Clin. Chem. (Washington)*, 41 (1995) 375-380.
- 2140 Fujimura, S., Rikimaru, T., Baba, S., Hori, J., Hao, X.-Q., Terada, S. and Kimoto, E.: Purification and characterization of a non-hemorrhagic metalloprotease from *Agkistrodon halys brevicaudus* venom. *Biochim. Biophys. Acta*, 1243 (1995) 94-100.
- 2141 Hofstadler, S.A., Swanek, F.D., Gale, D.C., Ewing, A.G. and Smith, R.D.: Capillary electrophoresis - electrospray ionization Fourier transform ion cyclotron resonance mass spectrometry for direct analysis of cellular proteins. *Anal. Chem.*, 67 (1995) 1477-1480.
- 2142 Janiak, A., Villar, R., Cassoly, R. and Rendu, F.: Tubulin is not phosphorylated in resting and thrombin-activated platelets. *J. Biochem. (Tokyo)*, 117 (1995) 296-302.
- 2143 Lausch, R., Reif, O.-W., Riechel, P. and Scheper, T.: Analysis of immunoglobulin G using a capillary electrophoretic affinity assay with protein A and laser-induced fluorescence detection. *Electrophoresis (Weinheim)*, 16 (1995) 636-641.
- 2144 Marchi, G., Dorizzi, R.M., Ferrari, V.D. and Jirillo, A.: Appearance of heavy chains and unfavorable prognosis in Hodgkin disease. *Clin. Chem. (Washington)*, 41 (1995) 761-762.
- 2145 Miura, T., Yabuki, S., Funato, T., Kawamura, T., Sasaki, T., Tsubo, S. and Machida, S.: (Capillary electrophoretic separation of human serum proteins by the addition of trimethylammonium propanesulfonate (Z1-methyl) to the buffer). *Seibutsu Kagaku*, 38 (1994) 425-431; *C.A.*, 122 (1995) 100891x.
- 2146 Miyata, T., Funatsu, A. and Kato, H.: Chemical cross-linking of activated coagulation factor VII with soluble tissue factor: calcium ions are not essential for full amidolytic activity of the factor VIIa-tissue factor complex after complex formation. *J. Biochem. (Tokyo)*, 117 (1995) 836-844.
- 2147 Ozaki, Y., Satoh, K., Yatomi, Y., Miura, S., Fujimura, Y. and Kume, S.: Protein tyrosine phosphorylation in human platelets induced by interaction between glycoprotein Ib and von Willebrand factor. *Biochim. Biophys. Acta*, 1243 (1995) 482-488.
- 2148 Prin, C., Bene, M.C., Gobert, B., Montagne, P. and Faure, G.C.: Isoelectric restriction of human immunoglobulin isotypes. *Biochim. Biophys. Acta*, 1243 (1995) 287-290.
- 2149 Stafeyeva, O.A., Khailova, L.S., Katrukha, A.G. and Bulargina, T.V.: (Preparation of anti-E1 monoclonal antibodies and their application in the study of the pyruvate dehydrogenase complex). *Biokhimiya (Moscow)*, 60 (1995) 124-130.
- 2150 Szamel, M., Leufgen, H., Kurrle, R. and Resch, K.: Differential signal transduction pathways regulating interleukin-2 synthesis and interleukin-2 receptor expression in stimulated human lymphocytes. *Biochim. Biophys. Acta*, 1235 (1995) 33-42.
- 2151 Tissot, J.-D. and Spertini, F.: Analysis of immunoglobulins by two-dimensional gel electrophoresis. *J. Chromatogr. A*, 698 (1995) 225-250 - a review with 237 refs.
- 2152 Ugarova, T.P., Zamarron, C., Veklich, Y., Bowditch, R.D., Ginsberg, M.H., Weisel, J.W. and Plow, E.F.: Conformational transitions in the cell binding domain of fibronectin. *Biochemistry*, 34 (1995) 4457-4466.
- 2153 Van den Berg, C.W., Harrison, R.A. and Morgan, B.P.: A rapid method for the isolation of analogs of human CD59 by preparative SDS-PAGE: application to pig CD59. *J. Immunol. Methods*, 179 (1995) 223-231; *C.A.*, 122 (1995) 184912r.
- 2154 Wang, X., DeVries, A. and Cheng, C.-H.: Antifreeze peptide heterogeneity in an antarctic eel pout includes an unusually large major variant comprised of two 7 kDa type III AFPs linked in tandem. *Biochim. Biophys. Acta*, 1247 (1995) 163-172.
- 2155 Yasuda, F., Hayashi, T., Tanitame, K., Nishioka, J. and Suzuki, K.: Molecular cloning and functional characterization of rat plasma protein S. *J. Biochem. (Tokyo)*, 117 (1995) 374-383.
- See also 1966, 2059, 2100, 2106, 2186, 2223, 2327.

19f. *Structural and muscle proteins*

- 2156 Asano, T., Shinohara, H., Morishita, R., Norota, I., Kato, K. and Endoh, M.: The G-protein G_o in mammalian cardiac muscle: localization and coupling to A₁ adenosine receptors. *J. Biochem. (Tokyo)*, 117 (1995) 183-189.
- 2157 Bárány, K., Bárány, M. and Giometti, C.S.: Polyacrylamide gel electrophoretic methods in the separation of structural muscle proteins. *J. Chromatogr. A*, 698 (1995) 301-332 - a review with 179 refs.
- 2158 Belkin, A.M. and Burridge, K.: Association of aciculin with dystrophin and utrophin. *J. Biol. Chem.*, 270 (1995) 6328-6337.
- 2159 Chan, D., Cole, W.R., Chow, C.W., Mundlos, S. and Bateman, J.F.: A COL2A1 mutation in a chondrogenesis type II results in the replacement of type II collagen by type I and III collagens in cartilage. *J. Biol. Chem.*, 270 (1995) 1747-1753.
- 2160 Claeys, E., Uytterhaegen, L., Buts, B. and Demeyer, D.: Quantification of beef myofibrillar proteins by SDS-PAGE. *Meat Sci.*, 39 (1995) 177-193; *C.A.*, 122 (1995) 104167p.
- 2161 Demignot, S., Borge, L. and Adolphe, M.: Transglutaminase activity in rabbit articular chondrocytes in culture. *Biochim. Biophys. Acta*, 1266 (1995) 163-170.
- 2162 Deyl, Z. and Miksik, I.: Separation of collagen type I chain polymers by electrophoresis in non-cross-linked polyacrylamide-filled capillaries. *J. Chromatogr. A*, 698 (1995) 369-373.
- 2163 Hayashi, A., Suzuki, T. and Tajima, S.: Modulations of elastin expression and cell proliferation by retinoids in cultured vascular smooth muscle cells. *J. Biochem. (Tokyo)*, 117 (1995) 132-136.
- 2164 Inaba, K.: ATP-Dependent conformational changes of dynein: evidence for changes in the interaction of dynein heavy chain with the intermediate chain 1. *J. Biochem. (Tokyo)*, 117 (1995) 903-907.
- 2165 Kawamura, Y., Kume, H., Itoh, Y., Ohtsuka, S., Kimura, S. and Maruyama, K.: Localization of three fragments of connectin in chicken breast muscle sarcomeres. *J. Biochem. (Tokyo)*, 117 (1995) 201-207.
- 2166 Knecht, M., Regitz-Zagrosek, V., Pleissner, K.-P., Jungblut, P., Steffen, C., Hildebrandt, A. and Fleck, E.: Characterization of myocardial protein composition in dilated cardiomyopathy by two-dimensional gel electrophoresis. *Eur. Heart J.*, 15(Suppl. D) (1994) 37-44; *C.A.*, 122 (1995) 182490c.
- 2167 LeBlanc, E.L., Singh, S. and LeBlanc, R.J.: Capillary zone electrophoresis of fish muscle sarcoplasmic proteins. *J. Food Sci.*, 59 (1994) 1267-1270; *C.A.*, 122 (1995) 212342a.
- 2168 Moyers, J.S., Linder, M.E., Shannon, J.D. and Parsons, S.J.: Identification of the *in vitro* phosphorylation sites on G_{sα} mediated by pp60^{c-src}. *Biochem. J.*, 305 (1995) 411-417.
- 2169 Schug, B.S. and Kalbhen, D.A.: Influence of chloroquine and other substances on the collagenolytic activity in human osteoarthritic cartilage *in vitro*. *Arzneim.-Forsch.*, 45 (1995) 285-289.
- 2170 Schumacher-Perdreau, F., Stefanik, D., Peters, G. and Pulverer, G.: Extracellular protein analysis: a reliable epidemiological tool for tracing methicillin-resistant Staphylococci. *Zentralbl. Bakteriol., Suppl.*, 26 (1994) 53-50; *C.A.*, 122 (1995) 234646e.
- 2171 Sires, U.I., Dublet, B., Aubert-Foucher, E., van der Rest, M. and Welgus, H.G.: Degradation of the COL1 domain of type XIV collagen by 92-kDa gelatinase. *J. Biol. Chem.*, 270 (1995) 1062-1067.
- 2172 Skolyseva, L.K., Smirnova, E.A., Medvedeva, M.V. and Gusev, N.B.: (Isolation and some properties of bovine aorta caldesmon). *Biokhimiya (Moscow)*, 60 (1995) 339-348.
- 2173 Sutton, C.W., Pemberton, K.S., Cottrell, J.S., Corbett, J.M., Wheeler, C.H., Dunn, M.J. and Pappin, D.J.: Identification of myocardial proteins from two-dimensional gels by peptide mass fingerprinting. *Electrophoresis (Weinheim)*, 16 (1995) 308-316.
- 2174 Yang, B., Jung, D., Rafael, J.A., Chamberlain, J.S. and Campbell, K.P.: Identification of α-syntrophin binding to syntrophin triplet, dystrophin, and utrophin. *J. Biol. Chem.*, 270 (1995) 4975-4978.
- 2175 Yoshihara, Y. and Kuroda, M.: Use of an SDS-gel-separated protein band as a ligand for affinity chromatography: procedure and application to the purification of domain-specific antibodies against α-actinin. *J. Biochem. (Tokyo)*, 117 (1995) 443-446.

19g. *Protamines, histones and other nuclear proteins*

- 2176 Annunziato, A.T., Eason, M.B. and Perry, C.A.: Relationship between methylation and acetylation of arginine-rich histones in cycling and arrested HeLa cells. *Biochemistry*, 34 (1995) 2916-2924.
- 2177 Buhrmester, H., von Kries, J.P. and Strätling, W.H.: Nuclear matrix protein ARBP recognizes a novel DNA sequence motif with high affinity. *Biochemistry*, 34 (1995) 4108-4117.
- 2178 Lindner, H., Helliger, W., Sarg, B. and Meraner, C.: Effect of buffer composition on the migration order and separation of histone H1 subtypes. *Electrophoresis (Weinheim)*, 16 (1995) 604-610.
- 2179 Santoro, R., D'Erme, M., Matroantonio, S., Reale, A., Marenzi, S., Saluz, H.-P., Strom, R. and Caiafa, P.: Binding of histone H1 e-c variants to CpG-rich DNA correlates with the inhibitory effect on enzymic DNA methylation. *Biochem. J.*, 305 (1995) 739-744.
- 2180 Thiriet, C. and Albert, P.: Rapid and effective Western blotting of histones from acid-urea-Triton and sodium dodecyl sulfate polyacrylamide gels: two different approaches depending on the subsequent qualitative or quantitative analysis. *Electrophoresis (Weinheim)*, 16 (1995) 357-361.

See also 2554.

19h. *Chromoproteins and metalloproteins*

- 2181 Beattie, J.H., Self, R. and Richards, M.P.: The use of solid phase concentrators for on-line preconcentration of metallothionein prior to isoform separation by capillary zone electrophoresis. *Electrophoresis (Weinheim)*, 16 (1995) 322-328.
- 2182 Hibino, T., Lee, B.H., Takabe, T. and Takabe, T.: Expression and characterization of Met92Gln mutant plastocyanin from *Silene pratensis*. *J. Biochem. (Tokyo)*, 117 (1995) 101-106.

- 2183 Jansen, E.H.J.M., Laan, C.A. and de Fluiter, P.: Determination of phthalate-induced rat liver cytochrome P-450 IVA1 by a fluorimetric enzymatic assay and by chemiluminescence detection on Western blots. *Anal. Chim. Acta*, 303 (1995) 99-102.
- 2184 Qian, Z.-Y., Jollès, P., Migliore-Samour, D. and Fiat, A.-M.: Isolation and characterization of sheep lactoferrin, an inhibitor of platelet aggregation and comparison with human lactoferrin. *Biochim. Biophys. Acta*, 1243 (1995) 25-32.
- 2185 Tashiro, H. and Sone, N.: Preparation and characterization of the hydrophilic Cu_A-cytochrome c domain of subunit II of cytochrome c oxidase from thermophilic *Bacillus* PS3. *J. Biochem. (Tokyo)*, 117 (1995) 521-526.
- 2186 Turpeinen, U., Sipilä, I., Anttila, P., Karjalainen, U., Kuronen, B., Kalkkinen, N., Ahola, T. and Stenman, U.-H.: Two α -chain hemoglobin variants, Hb Broussais and Hb Cemenelum, characterized by cation-exchange HPLC, isoelectric focusing, and peptide sequencing. *Clin. Chem. (Washington)*, 41 (1995) 532-536.
- 2187 Wu, J. and Pawliszyn, J.: Application of capillary isoelectric focusing with absorption imaging detection to the quantitative determination of human hemoglobin variants. *Electrophoresis (Weinheim)*, 16 (1995) 670-673.
- See also 1926.
- 19i. *Proteins of glands, gland products, various zymogens (incl. milk proteins)*
- 2188 Accatino, L., Pizarro, M., Solis, N. and Koenig, C.S.: Association of canalicular membrane enzymes with bile acid micelles and lipid aggregates in human and rat bile. *Biochim. Biophys. Acta*, 1243 (1995) 33-42.
- 2189 Chu, C.-C., Li, S.-H. and Chen, Y.-H.: Resolution of isotoxins in the β -bungarotoxin family. *J. Chromatogr. A*, 694 (1995) 492-497.
- 2190 Kristiansen, K.R., Otte, J., Zakora, M. and Qvist, K.B.: Capillary electrophoresis used to monitor the enzymic hydrolysis of caseins and the fractionation of hydrolysis products. *Milchwissenschaft*, 49 (1994) 683-688; *C.A.*, 122 (1995) 131350z.
- 2191 Leikin, A. and Shinitzky, M.: Characterization of the lipid surrounding the Δ^6 -desaturase of rat liver microsomes. *Biochim. Biophys. Acta*, 1256 (1995) 13-17.
- 2192 Marangoni, S., Toyama, M.H., Arantes, E.C., Giglio, J.R., da Silva, C.A., Carneiro, E.M., Goncalves, A.A. and Oliveira, B.: Amino acid sequence of TsTX-V, an α -toxin from *Tityus serrulatus* scorpion venom, and its effect on K⁺ permeability of β -cells from isolated rat islets of Langerhans. *Biochim. Biophys. Acta*, 1243 (1995) 309-314.
- 2193 Ota, Y., Ido, K., Kimura, K., Fukui, E. and Ikemoto, S.: Electrophoretic analysis of bile proteins from patients with and without gallstones. *Electrophoresis (Weinheim)*, 16 (1995) 402-406.
- 2194 Otte, J., Midtgaard, L. and Qvist, K.B.: Analysis of caseinomacropptide(s) by free solution capillary electrophoresis. *Milchwissenschaft*, 50 (1995) 75-79; *C.A.*, 122 (1995) 212373m.
- 2195 Otte, J.A.H.J., Kristiansen, K.R., Zakora, M. and Qvist, K.B.: Separation of individual whey proteins and measurement of α -lactalbumin and β -lactoglobulin by capillary zone electrophoresis. *Neth. Milk Dairy J.*, 48 (1994) 81-97; *C.A.*, 122 (1995) 29963q.
- 2196 Recio, I., Molina, E., Ramos, M. and de Frutos, M.: Quantitative analysis of major whey proteins by capillary electrophoresis using uncoated capillaries. *Electrophoresis (Weinheim)*, 16 (1995) 654-658.
- 2197 Saito, M., Kita, K., Sekiya, K., Omata, S. and Horigome, T.: Purification and molecular shape of a 144 kDa protein bearing N-acetylglucosamine residues from rat liver nuclear envelopes. *J. Biochem. (Tokyo)*, 117 (1995) 47-53.
- 2198 Sciacchitano, S., Danese, D., Andreoli, M., di Gregorio, A., Sacchi, A. and Pontecorvi, A.: Analysis of ras mutations in thyroid cytologic smears by denaturing gradient gel electrophoresis. *Front. Endocrinol.*, 9 (Highlights in Molecular and Clinical Endocrinology) (1994) 61-65; *C.A.*, 122 (1995) 234644c.
- 2199 Witzmann, F., Clack, J., Fultz, C. and Jarnot, B.: Two-dimensional electrophoretic mapping of hepatic and renal stress proteins. *Electrophoresis (Weinheim)*, 16 (1995) 451-559.
- 2200 Xu, C., Rigney, D.R. and Anderson, D.J.: Two-dimensional electrophoretic profile of human sperm membrane proteins. *J. Androl.*, 15 (1994) 595-602; *C.A.*, 122 (1995) 155525m.
- See also 2025, 2032, 2404.
- 19j. *Proteins of brain, cerebrospinal fluid and eye*
- 2201 Althaus, J.S., Fici, G.J. and VonVoigtlander, P.F.: Antibody transformation by peroxydinitrite as determined using capillary electrophoresis: a feasibility study. *Res. Commun. Mol. Pathol. Pharmacol.*, 87 (1995) 359-366; *C.A.*, 122 (1995) 237178q.
- 2202 Borghini, I., Barja, F., Pometta, D. and James, R.W.: Characterization of subpopulations of lipoprotein particles isolated from human cerebrospinal fluid. *Biochim. Biophys. Acta*, 1255 (1995) 192-200.
- 2203 Cameron, B.M., Sr., Merrill, C.R., Creed, G.J. and Vanderputten, D.: Methods for the diagnosis of peripheral nerve damage. *U.S. Pat. 5,364,793* (Cl. 436-86; GO1N33/49), 15 Nov. 1994, US Appl. 620,104, 30 Nov. 1990; 46 pp.; *C.A.*, 122 (1995) 27283u.
- 2204 Chao, D., Severson, D.L., Zwiers, H. and Hollenberg, M.D.: Radiolabelling of bovine myristoylated alanine-rich protein kinase C substrate (MARCKS) in an ADP-ribosylation reaction. *Biochem. Cell Biol.*, 72 (1994) 391-396.
- 2205 Goedert, M.: Molecular dissection of the neurofibrillary lesions of Alzheimer's disease. *Arzneim.-Forsch.*, 45 (1995) 403-409.
- 2206 Hosoi, T., Uchiyama, M., Okumura, E., Saito, T., Ishiguro, K., Uchida, T., Okuyama, A., Kishimoto, T. and Hisanaga, S.-i.: Evidence for cdk5 as a major activity phosphorylating tau protein in porcine brain extract. *J. Biochem. (Tokyo)*, 117 (1995) 741-749.
- 2207 Johnson, G., Ghanbari, H.A., Wolozin, B. and Merrill, C.R.: Methods for the diagnosis of Alzheimer's disease. *PCT Int. Appl. WO 95 05,604* (Cl. GO1N33/68), 23 Feb. 1995, US Appl. 105,922, 13 Aug. 1993; 28 pp.; *C.A.*, 122 (1995) 234861w.

- 2208 Kordeli, E., Lambert, S. and Bennett, V.: Ankyrin. A new ankyrin gene with neural-specific isoforms localized at the axonal initial segment and node of Ranvier. *J. Biol. Chem.*, 270 (1995) 2352-2359.
- 2209 Pley, U.M., Hill, B.L., Alibert, C., Brodsky, F.M. and Parham, P.: The interaction of calmodulin with clathrin-coated vesicles, triskelions, and light chains. Localization of a binding site. *J. Biol. Chem.*, 270 (1995) 2395-2402.
- 2210 Rahmatullah, M., Ginnan, R. and Robishaw, J.D.: Specificity of G protein α - γ subunit interactions. N-terminal 15 amino acids of γ subunit specifies interaction with α subunit. *J. Biol. Chem.*, 270 (1995) 2946-2951.
- 2211 Riley, M.L. and Harding, J.J.: The reaction of methylglyoxal with human and bovine lens proteins. *Biochim. Biophys. Acta*, 1270 (1995) 36-43.
- 2212 Shirao, T.: The roles of microfilament-associated proteins, drebrins, in brain morphogenesis: a review. *J. Biochem. (Tokyo)*, 117 (1995) 231-236 - a review with 42 refs.
- 2213 Strub, J.-M., Garcia-Sablone, P., Lonning, K., Taupenot, L., Hubert, P., van Dorsselaar, A., Aunis, D. and Metz-Boutigue, M.-H.: Processing of chromogranin B in bovine adrenal medulla. Identification of secretolytin, the endogenous C-terminal fragment of residues 614-626 with antibacterial activity. *Eur. J. Biochem.*, 229 (1995) 356-368.
- 2214 Urayama, O., Murakoshi, T. and Ikawa, Y.: Krev-1 protein is abundantly expressed in the rat spinal cord. *Biochim. Biophys. Acta*, 1243 (1995) 446-452.
- 19k. *Proteins of neoplastic tissue and transformed cells*
- 2215 Almquist, K.C., Loe, D.W., Hipfner, D.R., Mackie, J.E., Cole, S.P.C. and Deeley, R.G.: Characterization of the M_r 190,000 multidrug resistance protein (MRP) in drug-selected and transfected human tumor cells. *Cancer Res.*, 55 (1995) 102-110.
- 2216 Gorbachev, A.V., Egorova, S.G. and Myagkov, A.V.: (A biochemical study of antigen A2F4 associated with human lung adenocarcinoma). *Biokhimiya (Moscow)*, 59 (1994) 1401-1405.
- 2217 Kovarova, H., Stulik, J., Hochstrasser, D.F., Bures, J., Melichar, B. and Jandik, P.: Two-dimensional electrophoretic study of normal colon mucosa and colorectal cancer. *Appl. Theor. Electrophor.*, 4 (1994) 103-106; *C.A.*, 122 (1995) 234660e.
- 2218 Taylor, I.C.A., Roy, S., Yaswen, P., Stampfer, M.R. and Varmus, H.E.: Mouse mammary tumors express elevated levels of RNA encoding the murine homolog of SKY, a putative receptor tyrosine kinase. *J. Biol. Chem.*, 270 (1995) 6872-6880.
- 2219 Toyoshima, M., Nakajima, M., Yamori, T. and Tsuruo, T.: Purification and characterization of the platelet-aggregating sialoglycoprotein gp44 expressed by highly metastatic variant cells of mouse colon adenocarcinoma 26. *Cancer Res.*, 55 (1995) 767-773.
- See also 2110, 2198, 2229.
- 19l. *Specific binding and receptor proteins*
- 2220 Adam, R., Drummond, D.R., Solic, N., Holt, S.J., Sharma, R.P., Chamberlin, S.G. and Davies, D.E.: Modulation of the receptor binding affinity of amphiregulin by modification of its carboxyl terminal tail. *Biochim. Biophys. Acta*, 1266 (1995) 83-90.
- 2221 Akamura, F., Kato, M., Kameyama, K., Nukada, T., Haga, T., Kato, H., Takenawa, T. and Kikkawa, U.: Characterization of G α family G proteins G $_{L1\alpha}$ (G $_{14\alpha}$), G $_{L2\alpha}$ (G $_{11\alpha}$), and G $_{q\alpha}$ expressed in the baculovirus-insect cell system. *J. Biol. Chem.*, 270 (1995) 6246-6253.
- 2222 Brenner, C., Jan, G., Chevalier, Y. and Wroblewski, H.: Evaluation of the efficacy of zwitterionic dodecyl carboxybetaine surfactants for the extraction and the separation of mycoplasma membrane protein antigens. *Anal. Biochem.*, 224 (1995) 515-523.
- 2223 Cain, T.J., Liu, Y., Takizawa, T. and Robinson, J.M.: Solubilization of glycosyl-phosphatidylinositol-anchored proteins in quiescent and stimulated neutrophils. *Biochim. Biophys. Acta*, 1235 (1995) 69-78.
- 2224 Cardenas, A.M., Kuijpers, G.A.J. and Pollard, H.B.: Effect of protein synthesis inhibitors on synexin levels and secretory response in bovine adrenal medullary chromaffin cells. *Biochim. Biophys. Acta*, 1234 (1995) 255-260.
- 2225 Chan, K.C.: Characterization of rat serum insulin-like growth factor binding proteins by two-dimensional gel electrophoresis: identification of a potentially novel form. *Univ. Microfilm Int.*, Order No. DA9430421, 1993, 124 pp.; *C.A.*, 122 (1995) 179200c.
- 2226 Chantry, A.: The kinase domain and membrane localization determine intracellular interactions between epidermal growth factor receptors. *J. Biol. Chem.*, 270 (1995) 3068-3073.
- 2227 Chen, P.C., DuBois, G.C. and Chen, M.: Mapping the domain(s) critical for the binding of human tumor necrosis factor- α to its two receptors. *J. Biol. Chem.*, 270 (1995) 2874-2878.
- 2228 Denker, B.M., Boutin, P.M. and Neer, E.J.: Interactions between the amino- and carboxyl-terminal regions of G α subunits: analysis of mutated G α /G α_2 chimeras. *Biochemistry*, 34 (1995) 5544-5553.
- 2229 Di Renzo, M.F., Poulosom, R., Olivero, M., Comoglio, P.M. and Lemoine, N.R.: Expression of the Met/hepatocyte growth factor receptor in human pancreatic cancer. *Cancer Res.*, 55 (1995) 1129-1138.
- 2230 Jo, D.-W., Leren, T.-P., Yang, Z.-Y., Chung, Y.-H., Taylor, J.M. and Paik, Y.-K.: Characterization of an upstream regulatory element of the human apolipoprotein E gene, and purification of its binding protein from the human placenta. *J. Biochem. (Tokyo)*, 117 (1995) 915-922.
- 2231 Kishino, J., Kawamoto, K., Ishizaki, J., Verheij, H.M., Ohara, O. and Arita, H.: Pancreatic-type phospholipase A $_2$ activates prostaglandin E $_2$ production in rat mesangial cells by receptor binding reaction. *J. Biochem. (Tokyo)*, 117 (1995) 420-424.
- 2232 Kobayashi, A., Sogawa, K., Imataka, H. and Fujii-Kuriyama, Y.: Analysis of functional domains of a GC box-binding protein, BTEB. *J. Biochem. (Tokyo)*, 117 (1995) 91-95.
- 2233 Kovacina, K.S. and Roth, R.A.: Characterization of the endogenous insulin receptor-related receptor in neuroblastomas. *J. Biol. Chem.*, 270 (1995) 1881-1887.
- 2234 Krantz, S., Salazar, R., Brandt, R., Kellermann, J. and Lottspeich, F.: Purification and partial amino acid sequencing of a fructosyllysine-specific binding protein from cell membranes of the monocyte-like cell line U937. *Biochim. Biophys. Acta*, 1266 (1995) 109-112.

- 2235 Krupenko, S.A., Kolesnik, O.I., Krupenko, N.I. and Strel'chyonok, O.A.: Organization of the transcortin-binding domain on placental plasma membranes. *Biochim. Biophys. Acta*, 1235 (1995) 387-394.
- 2236 Lu, Z., Xia, L., Mesmer, O.T. and Lo, T.C.Y.: Use of hexose transport mutants to examine the expression and properties of the rat myoblast GLUT 1 transport process. *Biochim. Biophys. Acta*, 1234 (1995) 155-165.
- 2237 Moore, M.A. and McCarthy, M.P.: Snake venom toxins, unlike smaller antagonists, appear to stabilize a resting state conformation of the nicotinic acetylcholine receptor. *Biochim. Biophys. Acta*, 1235 (1995) 336-342.
- 2238 Ozcelebi, F. and Miller, L.J.: Phosphopeptide mapping of cholecystokinin receptors on agonist-stimulated native pancreatic acinar cells. *J. Biol. Chem.*, 270 (1995) 3435-3441.
- 2239 Pouliot, J.-F. and Béliveau, R.: Palmitoylation of the glucose transporter in blood-brain barrier capillaries. *Biochim. Biophys. Acta*, 1234 (1995) 191-196.
- 2240 Sakakibara, Y., Suiko, M., Nakajima, H. and Liu, M.-C.: Sulphation of L-tyrosine in mammalian cells: a comparative study. *Biochem. J.*, 305 (1995) 993-998.
- 2241 Shire, D., Carillon, C., Kaghad, M., Calandra, B., Rinaldi-Carmona, M., le Fur, G., Caput, D. and Ferrara, P.: An amino-terminal variant of the central cannabinoid receptor resulting from alternative splicing. *J. Biol. Chem.*, 270 (1995) 3726-37311.
- 2242 Stipani, I., Natuzzi, D., Daddabbo, L., Ritieni, A., Randazzo, G. and Palmieri, F.: Photoaffinity labeling of the mitochondrial oxoglutarate carrier by azido-phthalonate. *Biochim. Biophys. Acta*, 1234 (1995) 149-154.
- 2243 Szabo, T., Kadish, J.L. and Czop, J.K.: Biochemical properties of the ligand-binding 20-kDa subunit of the β -glucan receptors on human mononuclear phagocytes. *J. Biol. Chem.*, 270 (1995) 2145-2151.
- 2244 Tamei, H., Hoshino, T., Yoshida, S., Hayashi, T., Iwata, K. and Suzuki, K.: One-step sandwich enzyme immunoassays for human C4b-binding protein (C4BP) and protein S-C4BP complex using monoclonal antibodies. *Clin. Chim. Acta*, 234 (1995) 115-125.
- 2245 Thomson, M., Korn, M. and Hall, P.F.: GTP-binding proteins in adrenocortical mitochondria. *Biochim. Biophys. Acta*, 1248 (1995) 159-169.
- 2246 Wang, G.L. and Semenza, G.L.: Purification and characterization of hypoxia-inducible factor 1. *J. Biol. Chem.*, 270 (1995) 1230-1237.
- 2247 Waters, M.J., Daniel, N., Bignon, C. and Djiane, J.: The rabbit mammary gland prolactin receptor is tyrosine-phosphorylated in response to prolactin *in vivo* and *in vitro*. *J. Biol. Chem.*, 270 (1995) 5136-5143.
- 2248 Xu, J., McKeenan, K., Matsuzaki, K. and McKeenan, W.L.: Inhibin antagonizes inhibition of liver cell growth by activin by a dominant-negative mechanism. *J. Biol. Chem.*, 270 (1995) 6308-6313.
- 2249 Yamaguchi, T., Fermandez, R. and Roth, R.A.: Comparison of the signaling abilities of the *Drosophila* and human insulin receptors in mammalian cells. *Biochemistry*, 34 (1995) 4962-4968.
- 2250 Yokota, Y., Arai, T. and Kawasaki, T.: Oligomeric structures required for complement activation of serum mannan-binding proteins. *J. Biochem. (Tokyo)*, 117 (1995) 414-419.
- 2251 Yu, H. and Diamandis, E.P.: Prostate-specific antigen immunoreactivity in amniotic fluid. *Clin. Chem. (Washington)*, 41 (1995) 204-210.
- 2252 Zhang, Q.-X., Walker, F., Burgess, A.W. and Baldwin, G.S.: Reduction in platelet-derived growth factor receptor mRNA in v-src-transformed fibroblasts. *Biochim. Biophys. Acta*, 1266 (1995) 9-15.

See also 2081, 2105, 2259, 2399, 2648.

19m. Urinary proteins

- 2253 Burlando, L.: Factitious additional band simulating Bence Jones protein caused by contamination from face plate for ureteroscopy. *Clin. Chem. (Washington)*, 41 (1995) 477.
- 2254 Dussol, B., Daudon, M., Dupuy, P., Michel, R., Berland, Y., Dagon, J.C. and Verdier, J.M.: Analysis of the protein content of five different types of kidney stones. In: Ryall, R.L. (Editor), *Urolithiasis 2, [Proc. Int. Symp.], 7th 1994*, Plenum, New York, 1994, pp. 363-364; C.A., 122 (1995) 234744k.
- 2255 Grover, P.K. and Resnick, M.I.: Two-dimensional analysis of proteins in the urine of male and female stone formers. In: Ryall, R.L. (Editor), *Urolithiasis 2, [Proc. Int. Symp.], 7th 1992*, Plenum, New York, 1994, pp. 287-288; C.A., 122 (1995) 234645d.
- 2256 Grover, P.K. and Resnick, M.I.: Two-dimensional gel analysis of proteins in unprocessed human urine using double stain. In: Ryall, R.L. (Editor), *Urolithiasis 2, [Proc. Int. Symp.], 7th 1992*, Plenum, New York, 1994, pp. 285-286; C.A., 122 (1995) 109048d.
- 2257 Hiratsuka, N., Shiba, K., Shinomura, K., Hosaki, S., Cho, H., Nagasaki, A. and Kobayashi, S.: Urinary protein fractions in healthy subjects using cellulose acetate membrane electrophoresis followed by staining with Acid Violet 17. *Biol. Pharm. Bull.*, 17 (1994) 1355-1357; C.A., 122 (1995) 182524s.
- 2258 Liu, C.-M. and Wang, H.-P.: Method of sample preparation for urine protein analysis with capillary electrophoresis. *PCT Int. Appl. WO 95 02,182 (Cl. GO1N33/483)*, 19 Jan. 1995, US Appl. 91,844, 09 Jul. 1993; 40 pp.; C.A., 122 (1995) 155762m.
- 2259 Withold, W. and Reinauer, H.: An immunoblotting procedure following agarose gel electrophoresis for detection of Bence Jones proteinuria compared with immunofixation and quantitative light chain determination. *Eur. J. Clin. Chem. Clin. Biochem.*, 33 (1995) 135-138.

19n. Other proteins (incl. proteinous inhibitors of enzymic activity)

- 2260 Borisova, T.A., Los, G.V. and Lishko, V.K.: (Inclusion of secretory protein precursors into artificial phospholipid vesicles). *Biokhimiya (Moscow)*, 59 (1994) 1483-1489.
- 2261 Lomas, D.A., Elliott, P.R., Chang, W.W., Wardell, M.R. and Carrell, R.W.: Preparation and characterization of latent α_1 -antitrypsin. *J. Biol. Chem.*, 270 (1995) 5282-5288.
- 2262 Nadano, D., Yasuda, T., Takeshita, H. and Kishi, K.: Activity staining of mammalian ribonuclease inhibitors after electrophoresis in sealed vestrical slab polyacrylamide gels. *Anal. Biochem.*, 227 (1995) 210-215.

2263 Pinto, R.A., Hawgood, S., Clements, J.A., Benson, B.J., Naidu, A., Hamilton, R.L. and Wright, J.R.: Association of surfactant protein C with isolated alveolar type II cells. *Biochim. Biophys. Acta*, 1255 (1995) 16-22.

See also 2040, 2136, 2199, 2623, 2630.

20. ENZYMES AND ENZYME ACTIVITY ESTIMATION

2264 John, M.A. and Hussain, Z.: Multilocus enzyme electrophoresis using ultrathin polyacrylamide gels. *J. Microbiol. Methods*, 19 (1994) 307-313; *C.A.*, 122 (1995) 49687f.

See also 2052.

20a. Oxidoreductases

2265 Agarwal, A.K., Mune, T., Monder, C. and White, P.C.: Mutations in putative glycosylation sites of rat 11 β -hydroxysteroid dehydrogenase affect enzymatic activity. *Biochim. Biophys. Acta*, 1248 (1995) 70-74.

2266 Braun, H.-P. and Schmitz, U.K.: Molecular structure of the 8.0 kDa subunit of cytochrome-c reductase from potato and its $\Delta\Psi$ -dependent import into isolated mitochondria. *Biochim. Biophys. Acta*, 1229 (1995) 181-186.

2267 Esworthy, R.S., Baker, M.A. and Chu, F.: Expression of selenium-dependent glutathione peroxidase in human breast tumor cell lines. *Cancer Res.*, 55 (1995) 957-962.

2268 Fujita, Y., Uruga, Y. and Ichisima, E.: Molecular cloning and nucleotide sequence of the protyrosinase gene, melO, from *Aspergillus oryzae* and expression of the gene in yeast cells. *Biochim. Biophys. Acta*, 1261 (1995) 151-154.

2269 Hicks, D.B.: Purification of three catalase isozymes from facultatively alkaliphilic *Bacillus firmus* OF4. *Biochim. Biophys. Acta*, 1229 (1995) 347-355.

2270 Jiménez-Cervantes, C., García-Borrón, J.C., Lozano, J.A. and Solano, F.: Effect of detergents and endogenous lipids on the activity and properties of tyrosinase and its related proteins. *Biochim. Biophys. Acta*, 1243 (1995) 421-430.

2271 Kobayashi, T., Ishida, T., Horiike, K., Takahara, Y., Numao, N., Nakazawa, A., Nakazawa, T. and Nozaki, M.: Overexpression of *Pseudomonas putida* catechol 2,3-dioxygenase with high specific activity by genetically engineered *Escherichia coli*. *J. Biochem. (Tokyo)*, 117 (1995) 614-622.

2272 Krungkrai, J.: Purification, characterization and localization of mitochondrial dihydroorotate dehydrogenase in *Plasmodium falciparum*, human malaria parasite. *Biochim. Biophys. Acta*, 1243 (1995) 351-360.

2273 Lomnitski, L., Sklan, D. and Grossman, S.: Lipoygenase activity in rat dermis and epidermis: partial purification and characterization. *Biochim. Biophys. Acta*, 1255 (1995) 351-359.

2274 Marui, Y., Hayashi, C., Matsuda, Y., Matsuura, S., Eto, A., Ohba, Y. and Okuda, K.: Multi-enzyme reference material from established human cell lines and human sources. *Clin. Chim. Acta*, 233 (1995) 19-38.

2275 Mazhul', M.M. and Danilov, V.S.: (Study of properties of NAD(P)H:FMN oxidoreductase from the marine luminescent bacteria *Vibrio fischeri*). *Biokhimiya (Moscow)*, 59 (1994) 1608-1614.

2276 Narita, H. and Morishita, E.: Production and application of monoclonal antibodies specific to pyrroloquinoline quinone. *J. Biochem. (Tokyo)*, 117 (1995) 830-835.

2277 Nijtmans, L.G.J., Barth, P.G., Lincke, C.R., van Galen, M.J.M., Zwart, R., Klement, P., Bolhuis, P.A., Ruitenbeek, W., Wanders, R.J.A. and van den Bogert, C.: Altered kinetics of cytochrome c oxidase in a patient with severe mitochondrial encephalomyopathy. *Biochim. Biophys. Acta*, 1270 (1995) 193-201.

2278 Nijtmans, L.G.J., Spelbrink, J.N., van Galen, M.J.M., Zwaan, M., Klement, P. and van den Bogert, C.: Expression and fate of the nuclear encoded subunits of cytochrome-c oxidase in cultured human cells depleted of mitochondrial gene products. *Biochim. Biophys. Acta*, 1265 (1995) 117-126.

2279 Palumbo, A. and Jackson, I.J.: Peroxidase activity in the ink gland of *Sepia officinalis* and partial nucleotide sequence of a candidate cDNA encoding the enzyme. *Biochim. Biophys. Acta*, 1247 (1995) 173-178.

2280 Reddy, S.G., Sacchetti, J.C. and Blanchard, J.S.: Expression, purification, and characterization of *Escherichia coli* dihydrodipicolinate reductase. *Biochemistry*, 34 (1995) 3492-3501.

2281 Rubartelli, A., Bonifaci, N. and Sitia, R.: High rates of thioredoxin secretion correlate with growth arrest in hepatoma cells. *Cancer Res.*, 55 (1995) 675-680.

2282 Schilling, B. and Lerch, K.: Amine oxidases from *Aspergillus niger*: identification of a novel flavin-dependent enzyme. *Biochim. Biophys. Acta*, 1243 (1995) 529-537.

2283 Soukri, A., Valverde, F., Hafid, N., Elkebbaj, M.S. and Serrano, A.: Characterization of muscle glyceraldehyde-3-phosphate dehydrogenase isoforms from euthermic and induced hibernating *Jaculus orientalis*. *Biochim. Biophys. Acta*, 1243 (1995) 161-168.

2284 Suzuki, K., Mizuguchi, M., Gomi, T. and Itagaki, E.: Identification of a lysine residue in the NADH-binding site of salicylate hydroxylase from *Pseudomonas putida* S-1. *J. Biochem. (Tokyo)*, 117 (1995) 579-585.

2285 Tachibana, M., Onoe, H., Okubo, A. and Yamazaki, S.: Application of micellar electrokinetic chromatography as a novel assay method for dimethyl sulfoxide reductase. *Biosci., Biotechnol., Biochem.*, 59 (1995) 282-284; *C.A.*, 122 (1995) 208205r.

2286 Terada, K., Kawarada, Y., Miura, N., Yasui, O., Koyama, K. and Sugiyama, T.: Copper incorporation into ceruloplasmin in rat livers. *Biochim. Biophys. Acta*, 1270 (1995) 58-62.

2287 Uchida, K. and Kawakishi, S.: Formation of 2-oxo-histidine at the active site of Cu,Zn-superoxide dismutase by reaction with H₂O₂. *Int. Congr. Ser.-Excerpta Med.*, 1058 (Frontiers of Reactive Oxygen Species in Biology and Medicine) (1994) 175-176; *C.A.*, 122 (1995) 127419m.

2288 Wang, H.-t., Liu, W.-y. and Ulbrich, N.: Isolation and characterization of a tyrosinase from the skin of the white silky fowl (*Gallina lanigera*) employing copper saturated diethylaminoethyl-cellulose. *Biochim. Biophys. Acta*, 1243 (1995) 251-255.

See also 2076.

20b. Transferases (excl. E.C. 2.7.-.)

2289 Buki, K.G., Bauer, P.I., Hakam, A. and Kun, E.: Identification of domains of poly(ADP-ribose) polymerase for protein binding and self-association. *J. Biol. Chem.*, 270 (1995) 3370-3377.

- 2290 Egaas, E., Falls, J.G., Svendsen, N.O., Ramstad, H., Skaare, J.Y. and Dauterman, W.C.: Strain- and sex-specific differences in the glutathione S-transferase class pi in the mouse examined by gradient elution of the glutathione-affinity matrix and reverse-phase high performance liquid chromatography. *Biochim. Biophys. Acta*, 1243 (1995) 256-264.
- 2291 Hambartsumian, A.A. and Bezirjian, Kh.O.: (Valine-pyruvate transaminase of *Brevibacterium flavum*: isolation and primary characterization). *Biokhimiya (Moscow)*, 59 (1994) 1378-1384.
- 2292 Jeong, J., Murthy, S.N.P., Radek, J.T. and Lorand, L.: The fibronectin-binding domain of transglutaminase. *J. Biol. Chem.*, 270 (1995) 5654-5658.
- 2293 Saheki, T., Mori, K., Kobayashi, K., Horiuchi, M., Shige, T., Obara, T., Suzuki, S., Mori, M. and Yamamura, K.-i.: Importance of ornithine transcarbamylase (OTC) deficiency in small intestine for urinary orotic acid excretion: analysis of OTC-deficient spf-ash mice with OTC transgene. *Biochim. Biophys. Acta*, 1270 (1995) 87-93.
- 2294 Varone, C.L., Canépa, E.T., Llambías, E.B.C. and Grinstein, M.: cAMP Regulation of phenobarbital-mediated induction of δ -aminolevulinic synthase mRNA in hepatocytes from normal and experimental diabetic rats. *Biochem. Cell Biol.*, 72 (1994) 381-390.
- See also 2274, 2345, 2436, 2446.
- 20c. *Transferases transferring phosphorus containing groups (E.C. 2.7.-)*
- 2295 Ahmad, F. and Goldstein, B.J.: Purification, identification and subcellular distribution of three predominant protein-tyrosine phosphatase enzymes in skeletal muscle tissue. *Biochim. Biophys. Acta*, 1248 (1995) 57-69.
- 2296 Chernov, A.V., Matvienko, N.N., Zheleznaya, L.A. and Matvienko, N.I.: (A new site-specific endonuclease-methylase from the thermophilic strain of *Bacillus* species LU11). *Biokhimiya (Moscow)*, 59 (1994) 1714-1729.
- 2297 Craig, A.G., Hoeger, C.A., Miller, C.L., Goedken, T., Rivier, J.E. and Fischer, W.H.: Monitoring protein kinase and phosphatase reactions with matrix-assisted laser desorption/ionization mass spectrometry and capillary zone electrophoresis: comparison of the detection efficiency of peptide-phosphopeptide mixtures. *Biol. Mass Spectrom.*, 23 (1994) 519-528; *C.A.*, 122 (1995) 127051k.
- 2298 Heider, H. and Widmer, H.R.: Okadaic acid potentiates heat-induced activation of erk2. *Biochim. Biophys. Acta*, 1265 (1995) 196-200.
- 2299 Hsuan, J.J., Totty, N.F., Truong, O. and Waterfield, M.D.: Purification of phosphatidylinositol 3-kinase by SDA-agarose gel electrophoresis and HPEC. *Protein Pept. Lett.*, 1 (1994) 25-32; *C.A.*, 122 (1995) 154733r.
- 2300 Kameshita, I. and Fujisawa, H.: Preparation and characterization of calmodulin-dependent protein kinase IV (CaM-kinase IV) free of CaM-kinase IV kinase from rat cerebral cortex. *J. Biochem. (Tokyo)*, 117 (1995) 85-90.
- 2301 Kuroda, K. and Ueda, R.: A 130 kDa polypeptide immunologically related to the 180 kDa catalytic subunit of DNA polymerase α -primase complex is detected in early embryos of *Drosophila*. *J. Biochem. (Tokyo)*, 117 (1995) 809-818.
- 2302 Matsumoto, N., Kojima, S., Osawa, T. and Toyoshima, S.: Protein kinase C phosphorylates p50 LSP1 and induces translocation of p50 LSP1 in T lymphocytes. *J. Biochem. (Tokyo)*, 117 (1995) 222-229.
- 2303 Murthy, V.V.: Adenylate kinase mimics creatine kinase-MM isoenzyme in a CK isoenzyme electrophoresis assay. *J. Clin. Lab. Anal.*, 8 (1994) 140-143; *C.A.*, 122 (1995) 26347f.
- 2304 Ohgi, K., Iwama, M., Tada, K., Takizawa, R. and Irie, M.: Role of Lys108 in the enzymatic activity of RNase Rh from *Rhizopus niveus*. *J. Biochem. (Tokyo)*, 117 (1995) 27-33.
- 2305 Pertile, P. and Cantley, L.C.: Type 2 phosphatidylinositol 4-kinase is recruited to CD4 in response to CD4 cross-linking. *Biochim. Biophys. Acta*, 1248 (1995) 129-134.
- 2306 Price, D.J., Kawakami, Y., Kawakami, T. and Rivnay, B.: Purification of a major tyrosine kinase from RBL-2H3 cells phosphorylating Fc ϵ R1 γ -cytoplasmic domain and identification as the Btk tyrosine kinase. *Biochim. Biophys. Acta*, 1265 (1995) 133-142.
- 2307 Roberts, M.W., Preiss, J. and Okita, T.W.: A capillary zone electrophoresis assay for the nucleoside transfer enzyme adenosine diphosphate-glucose pyrophosphorylase. *Anal. Biochem.*, 225 (1995) 121-126.
- 2308 Shapovalova, N.I., Zheleznaya, L.A. and Matvienko, N.I.: (A new site-specific endonuclease and methylase from the thermophilic strain of *Bacillus* species KT6). *Biokhimiya (Moscow)*, 59 (1994) 1730-1738.
- 2309 Tsuchiya, M., Osago, H. and Shimoyama, M.: Assay of arginine-specific adenosine-5'-diphosphate-ribosyltransferase by capillary electrophoresis. *Anal. Biochem.*, 224 (1995) 486-489.
- 2310 Zhang, J., Fukui, T. and Ichikawa, A.: A third type of nucleoside diphosphate kinase from spinach leaves: purification, characterization and amino-acid sequence. *Biochim. Biophys. Acta*, 1248 (1995) 19-26.
- See also 2107, 2274.
- 20d. *Hydrolases, acting on ester bonds (E.C. 3.1.-)*
- 2311 Carrea, G., D'Arrigo, P., Piergianni, V., Roncaglio, S., Secundo, F. and Servi, S.: Purification and properties of two phospholipases D from *Streptomyces* sp. *Biochim. Biophys. Acta*, 1255 (1995) 273-279.
- 2312 Cheung, C.K., Panesar, N.S., Haines, C., Masarei, J. and Swaminathan, R.: Immunoassay of a tartrate-resistant acid phosphatase in serum. *Clin. Chem. (Washington)*, 41 (1995) 679-686.
- 2313 Kálmán, F., Ma, S., Hodel, A., Fox, R.O. and Horváth, C.: Charge and size effects in the capillary zone electrophoresis of nuclease A and its variants. *Electrophoresis (Weinheim)*, 16 (1995) 595-603.
- 2314 Pedersen, J., Andersen, J., Roepstorff, P., Filimonova, M. and Biedermann, K.: (Characterization of *Serratia marcescens* nuclease isoforms by electrospray mass spectrometry). *Biokhimiya (Moscow)*, 60 (1995) 462-469.
- 2315 Rugani, N., Carrière, F., Thim, L., Borgstrom, B. and Sarda, L.: Lipid binding and activating properties of porcine pancreatic colipase split at the Ile⁷⁹-Thr⁸⁰ bond. *Biochim. Biophys. Acta*, 1247 (1995) 185-194.
- 2316 Sembaj, A., Carriazo, C., Sanz, E. and Barral, J.M.: Determination of alkaline phosphatase isozymes in amniotic fluid. *Eur. J. Clin. Chem. Clin. Biochem.*, 33 (1995) 281-284.

- 2317 Teissère, M., Borel, M., Caillol, B., Nari, J., Gardies, A.M. and Noat, G.: Purification and characterization of a fatty acyl-ester hydrolase from post-germinated sunflower seeds. *Biochim. Biophys. Acta*, 1255 (1995) 105-112.
- 2318 Uritani, M., Takai, M. and Yoshinaga, K.: Protective effect of disaccharides on restriction endonucleases during drying under vacuum. *J. Biochem. (Tokyo)*, 117 (1995) 774-779.
- 2319 Withiam-Leitch, M., Rubin, R.P., Koshlukava, S.E. and Aletta, J.M.: Identification and characterization of carboxyl ester hydrolase as a phospholipid hydrolyzing enzyme of zymogen granule membranes from rat exocrine pancreas. *J. Biol. Chem.*, 270 (1995) 3780-3787.
- 2320 Yamazaki, Y., Ogawa, Y., Afify, A.S., Kageyama, Y.-i., Okada, T., Okuno, H., Yoshii, Y. and Nose, T.: Difference between cancer cells and the corresponding normal tissue in view of stereoselective hydrolysis of synthetic esters. *Biochim. Biophys. Acta*, 1243 (1995) 300-308.
- See also 2030, 2088, 2223, 2274, 2297.
- 20e. *Hydrolases, acting on glycosyl compounds (E.C. 3.2.-.-)*
- 2321 Ara, K., Saeki, K., Igarashi, K., Takaiwa, M., Uemura, T., Hagi-hara, H., Kawai, S. and Ito, S.: Purification and characterization of an alkaline amylopullulanase with both α -1,4 and α -1,6 hydrolytic activity from alkalophilic *Bacillus* sp. KSM-1378. *Biochim. Biophys. Acta*, 1243 (1995) 315-324.
- 2322 Beyer, E., Ivleva, T., Artykova, G. and Wiederschain, G.: Change of isoforms' spectra of α -L-fucosidase from human skin fibroblasts in intracellular storage of nonhydrolyzable substances. *Biochim. Biophys. Acta*, 1270 (1995) 7-11.
- 2323 Kwon, K.-S., Lee, J., Kang, H.G., Hah, Y.C.: Detection of β -glucosidase activity in polyacrylamide gels with esculin as substrate. *Appl. Environ. Microbiol.*, 60 (1994) 4584-4586; *C.A.*, 122 (1995) 26377r.
- 2324 Lee, E.-O. and Kim, J.-D.: Palmitoyl lysozyme-induced stabilization of PE (phosphatidylethanolamine) liposomes and their interaction with *Candida albicans*. *J. Biochem. (Tokyo)*, 117 (1995) 54-58.
- 2325 Martino, S., Emiliani, C., Orlicchio, A., Hosseini, R. and Stirling, J.L.: β -N-Acetylhexosaminidases A and S have similar sub-cellular distributions in HL-60 cells. *Biochim. Biophys. Acta*, 1243 (1995) 489-495.
- 2326 Miura, R.O., Yamagata, S., Miura, Y., Harada, T. and Yamagata, T.: Analysis of glycosaminoglycan-degrading enzymes by substrate gel electrophoresis (zymography). *Anal. Biochem.*, 225 (1995) 333-340.
- 2327 Moriyama, T., Tozawa, T., Nobuoka, M. and Ikeda, H.: Sialyl salivary-type amylasemia associated with immunoglobulin D-type multiple myeloma. *Clin. Chim. Acta*, 233 (1995) 127-134.
- 2328 Yoshigi, N., Sahara, H. and Koshino, S.: Role of the C-terminal region of α -amylase from barley. *J. Biochem. (Tokyo)*, 117 (1995) 63-67.
- See also 1974, 2111, 2408.
- 20f. *Other hydrolases*
- 2329 Akasaki, K., Yoshimoto, H., Nakamura, A., Shiomi, H. and Tsuji, H.: Purification and characterization of a major kyotorphin-hydrolyzing peptidase of rat brain. *J. Biochem. (Tokyo)*, 117 (1995) 897-902.
- 2330 Anner, B.M., Imesch, E. and Moosmayer, M.: Normal sensitivity of Na⁺/K⁺-ATPase isolated from brain and kidney of spontaneously hypertensive rats to sodium, ouabain or mercury. *Biochim. Biophys. Acta*, 1270 (1995) 95-99.
- 2331 Balaban, N.P., Sharipova, M.R., Itskovich, E.L., Leshchinskaya, I.B. and Rudenskaya, G.N.: (Secreted serine proteinase from sporogeneous bacteria *Bacillus intermedius* 3-19). *Biokhimiya (Moscow)*, 59 (1994) 1393-1400.
- 2332 Bañuelos, M., Quintero, F.J. and Rodríguez-Navarro, A.: Functional expression of the ENA1(PMR2)-ATPase of *Saccharomyces cerevisiae* in *Schizosaccharomyces pombe*. *Biochim. Biophys. Acta*, 1229 (1995) 233-238.
- 2333 Belling, H.F., Steghens, J.P. and Collombel, C.: Firefly luciferase purification using polyethylene glycol and Dyematex Orange A. *J. Chromatogr. A*, 695 (1995) 33-40.
- 2334 Bergmann, U., Tuuttila, A., Stetler-Stevenson, W.G. and Tryggvason, K.: Autolytic activation of recombinant human 72 kilodalton type IV collagenase. *Biochemistry*, 34 (1995) 2819-2825.
- 2335 Bockelmann, W., Hoppe-Seyler, T. and Heller, K.J.: Quantitative determination of endopeptidase activity from lactic acid bacteria with capillary electrophoresis. *Milchwissenschaft*, 50 (1995) 13-17; *C.A.*, 122 (1995) 208206s.
- 2336 Deperthes, D., Gauthier, E.R., Chapdelaine, P., Lazure, C., Tremblay, R.R. and Dubé, J.Y.: Identification of glandular kallikrein in dog pancreas and determination of its tissue distribution. *Biochim. Biophys. Acta*, 1243 (1995) 291-294.
- 2337 Gary, J.D. and Clarke, S.: Purification and characterization of an isoaspartyl dipeptidase from *Escherichia coli*. *J. Biol. Chem.*, 270 (1995) 4076-4087.
- 2338 Hashimoto, T., Yamamoto, Y., Yoshida, Y. and Tagawa, K.: Cleavage of bovine mitochondrial ATPase inhibitor with endopeptidases, and binding of the resulting peptides to the interface between the α - and β -subunits of F₁ATPase. *J. Biochem. (Tokyo)*, 117 (1995) 641-647.
- 2339 Hymes, J., Fleischhauer, K. and Wolf, B.: Biotinylation of biotinidase following incubation with biocytin. *Clin. Chim. Acta*, 233 (1995) 39-45.
- 2340 Imai, K., Yokohama, Y., Nakanishi, I., Ohuchi, E., Fujii, Y., Nakai, N. and Okada, Y.: Matrix metalloproteinase 7 (matrilysin) from human rectal carcinoma cells. Activation of the precursor, interaction with other matrix metalloproteinases and enzymic properties. *J. Biol. Chem.*, 270 (1995) 6691-6697.
- 2341 Inomata, M. and Kawashima, S.: The possible self-down-regulation of calpain triggered by cell membranes. *Biochim. Biophys. Acta*, 1235 (1995) 107-114.
- 2342 Itoh, C. and Nagamatsu, A.: An aminopeptidase activity from porcine kidney that hydrolyzes oxytocin and vasopressin: purification and partial characterization. *Biochim. Biophys. Acta*, 1243 (1995) 203-208.
- 2343 Jeng, A.Y., Wong, M., Duelfer, T., Shapiro, S.D., Kramer, R.A. and Hu, S.-i.: Mouse macrophage metalloelastase expressed in bacteria absolutely requires zinc for activity. *J. Biochem. (Tokyo)*, 117 (1995) 216-221.

- 2344 Kim, Y.-T., Muramatsu, T. and Takahashi, K.: Leader peptidase from *Escherichia coli*: overexpression, characterization, and inactivation by modification of tryptophan residues 300 and 310 with N-bromosuccinimide. *J. Biochem. (Tokyo)*, 117 (1995) 535-544.
- 2345 Kimoto, M., Whitley, G.S.J., Tsuji, H. and Ogawa, T.: Detection of N^G, N^G-dimethylarginine dimethylaminohydrolase in human tissues using a monoclonal antibody. *J. Biochem. (Tokyo)*, 117 (1995) 237-238.
- 2346 Lengyel, E., Gum, R., Juarez, J., Clayman, G., Seiki, M., Sato, H. and Boyd, D.: Induction of M_r 92,000 type IV collagenase expression in a squamous cell carcinoma cell line by fibroblasts. *Cancer Res.*, 55 (1995) 963-967.
- 2347 Mars, I. and Monnet, V.: An aminopeptidase P from *Lactococcus lactis* with original specificity. *Biochim. Biophys. Acta*, 1243 (1995) 209-215.
- 2348 Pedrosa, M.M. and Legaz, M.E.: Separation of arginase isoforms by capillary zone electrophoresis and isoelectric focusing in density gradient column. *Electrophoresis (Weinheim)*, 16 (1995) 659-669.
- 2349 Spruth, M., Reidlinger, J. and Müller, V.: Sodium ion dependence of inhibition of the Na⁺-translocating F₁F₀-ATPase from *Acetobacterium woodii*. Probing the site(s) involved in ion transport. *Biochim. Biophys. Acta*, 1229 (1995) 96-102.
- 2350 Srinivasan, M., Kalousek, F. and Curthoys, N.P.: *In vitro* characterization of the mitochondrial processing and the potential function of the 68-kDa subunit of renal glutaminase. *J. Biol. Chem.*, 270 (1995) 1185-1190.
- 2351 Ueno, S., Kusaba, M., Takeda, K., Maeda, M., Futai, M., Izumi, F. and Kawamura, M.: Functional consequences of substitution of the disulfide-bonded segment, Cys127-Cys150, located in the extracellular domain of the Na,K-ATPase β subunit: Arg148 is essential for the functional expression of Na,K-ATPase. *J. Biochem. (Tokyo)*, 117 (1995) 591-596.
- 2352 Weder, J.K.P. and Kaiser, K.-P.: Fluorogenic substrates for hydrolase detection following electrophoresis. *J. Chromatogr. A*, 698 (1995) 181-201 - a review with 92 refs.
- 2353 Yoshimoto, T., Tabira, J., Kabashima, T., Inoue, S. and Ito, K.: Protease II from *Moraxella lacunata*: cloning, sequencing, and expression of the enzyme gene, and crystallization of the expressed enzyme. *J. Biochem. (Tokyo)*, 117 (1995) 654-660.
- See also 2028, 2140, 2589.
- 20g. *Lyases*
- 2354 Zhang, R., Yatsuki, H., Kusakabe, T., Iwabe, N., Miyata, T., Imai, T., Yoshida, M. and Hori, K.: Structures of cDNAs encoding the muscle-type and non-muscle-type isozymes of lamprey fructose biphosphate aldolases and the evolution of aldolase genes. *J. Biochem. (Tokyo)*, 117 (1995) 545-553.
- 20h. *Isomerases*
- 2355 Ashiuchi, M., Yoshimura, T., Kitamura, T., Kawata, Y., Nagai, J., Gorlatov, S., Esaki, N. and Soda, K.: *In vivo* effect of GroESL on the folding of glutamate racemase of *Escherichia coli*. *J. Biochem. (Tokyo)*, 117 (1995) 495-498.
- 2356 Kang, M.-K., Kim, C.-K., Johng, T.-N. and Paik, Y.-K.: Cholesterol biosynthesis from lanosterol: regulation and purification of rat hepatic sterol 8-isomerase. *J. Biochem. (Tokyo)*, 117 (1995) 819-823.
- 2357 Kaufmann, S.H., Charron, M., Burke, P.J. and Karp, J.E.: Changes in topoisomerase I level and localization during myeloid mutation *in vitro* and *in vivo*. *Cancer Res.*, 55 (1995) 1255-1260.
- 2358 Saito, M., Hori, K., Kurotsu, T., Kanda, M. and Saito, Y.: Three conserved glycine residues in valine activation of gramicidin S synthetase 2 from *Bacillus brevis*. *J. Biochem. (Tokyo)*, 117 (1995) 276-282.
- 2359 Staron, K., Kowalska-Loth, B., Zabek, J., Czerwinski, R.M., Nieznanski, K. and Szumiel, I.: Topoisomerase I is differently phosphorylated in two sublines of L5178Y mouse lymphoma cells. *Biochim. Biophys. Acta*, 1260 (1995) 35-42.
- 2360 Urade, Y., Tanaka, T., Eguchi, N., Kikuchi, M., Kimura, H., Toh, H. and Hayaishi, O.: Structural and functional significance of cysteine residues of glutathione-independent prostaglandin D synthase. Identification of Cys⁶⁵ as an essential thiol. *J. Biol. Chem.*, 270 (1995) 1422-1428.
- 20i. *Ligases*
- See 2078.
- 20j. *Complex mixtures and incompletely identified enzymes*
- 2361 Gao, X.-D., Katsumoto, T. and Onodera, K.: Purification and characterization of chitin deacetylase from *Absidia coerulea*. *J. Biochem. (Tokyo)*, 117 (1995) 257-263.
- 2362 Misra, I., Charlier, H.A., Jr. and Miziorko, H.M.: Avian cytosolic 3-hydroxy-3-methylglutaryl-CoA synthase: evaluation of the role of cysteines in reaction chemistry. *Biochim. Biophys. Acta*, 1247 (1995) 253-259.
21. PURINES, PYRIMIDINES, NUCLEIC ACIDS AND THEIR CONSTITUENTS
- 21a. *Purines, pyrimidines, nucleosides, nucleotides*
- 2363 Baba, Y., Inoue, H., Tshako, M., Sawa, T., Kishida, A. and Akashi, M.: Evaluation of the selective binding ability of oligodeoxynucleotides to poly(9-vinyladenine) using capillary affinity gel electrophoresis. *Anal. Sci.*, 10 (1994) 967-969; *C.A.*, 122 (1995) 23137b.
- 2364 Chen, N., Manabe, T., Terabe S., Yohda, M. and Endo, I.: High-resolution separation of oligonucleotides and DNA sequencing reaction products by capillary electrophoresis with linear polyacrylamide and laser-induced fluorescence detection. *J. Microcolumn Sep.*, 6 (1994) 539-543; *C.A.*, 122 (1995) 73236d.
- 2365 Edel, C.M., Hartog, A.F. and Berden, J.A.: Analysis of the inhibitory non-catalytic ADP binding site on mitochondrial F₁, using NAP₃-2N₃ADP as probe. Effects of the modification of ATPase and ITase activity. *Biochim. Biophys. Acta*, 1229 (1995) 103-114.
- 2366 Mao, B.: Studies of site-specific and stereospecific BPDE-N²-dG oligodeoxynucleotide adducts by gel electrophoresis. Avail. *Univ. Microfilms Int.*, Order No. DA9422939, 1993, 249 pp.; *C.A.*, 122 (1995) 232767w.

- 2367 Shah, G.M., Poirier, D., Duchaine, C., Brochu, G., Desnoyers, S., Laguex, J., Verreault, A., Hoflack, J.-C., Kirkland, J.B. and Poirier, G.G.: Methods for biochemical study of poly(ADP-ribose) metabolism *in vitro* and *in vivo*. *Anal. Biochem.*, 227 (1995) 1-13 - review with some papers cited.
- 2368 Takenaka, A., Matsumoto, O., Chen, Y., Hasegawa, S.-i., Chatake, T., Tsunoda, M., Ohta, T., Komatsu, Y., Koizumi, M. and Ohtsuka, E.: Structural composition of hammerhead ribozymes. *J. Biochem. (Tokyo)*, 117 (1995) 850-855.
- 2369 Zhao, Z., Wahl, J.H., Udseth, H.R., Hofstadler, S.A., Fuciarelli, A.F. and Smith, R.D.: On-line capillary electrophoresis-electrospray ionization mass spectrometry of nucleotides. *Electrophoresis (Weinheim)*, 16 (1995) 389-395.
- See also 1926, 2063, 2488, 2528, 2642.
- 21b. *Nucleic acids, RNA*
- 2370 Aas-Eng, D.A., Åsheim, H.C., Deggerdal, A., Smeland, E. and Funderud, S.: Characterization of a promoter region supporting transcription of a novel human β -galactoside α -2,6-sialyltransferase transcript in HepG2 cells. *Biochim. Biophys. Acta*, 1261 (1995) 166-169.
- 2371 Boon, W.C., Roche, P.J., Hammond, V.E., Jeyaseelan, K., Crawford, R.J. and Coghlan, J.P.: Cloning and expression analysis of a cytochrome P-450_{11 β} cDNA in sheep. *Biochim. Biophys. Acta*, 1260 (1995) 109-112.
- 2372 Cary, J.W. and Bhatnagar, D.: Nucleotide sequence of a *Aspergillus parasiticus* gene strongly repressed by thiamine. *Biochim. Biophys. Acta*, 1261 (1995) 319-320.
- 2373 De Kleijn, D.P.V., de Leeuw, E.P.H., van den Berg, M.C., Martens, G.J.M. and van Herp, F.: Cloning and expression of two mRNAs encoding structurally different crustacean hyperglycemic hormone precursors in the lobster *Homarus americanus*. *Biochim. Biophys. Acta*, 1260 (1995) 62-66.
- 2374 Dinchuk, J., Hart, J., Gonzalez, G., Karmann, G., Schmidt, D. and Wirak, D.O.: Remodelling of lipoproteins in transgenic mice expressing human cholesteryl ester transfer protein. *Biochim. Biophys. Acta*, 1255 (1995) 301-310.
- 2375 Doll, R.F., Bruce, A. and Smith, F.I.: Regulation of the human acid β -glucosidase promoter in multiple cell types. *Biochim. Biophys. Acta*, 1261 (1995) 57-67.
- 2376 Ferreras, J.M., Iglesias, R., Barbieri, L., Alegre, C., Bolognesi, A., Rojo, M.A., Carbajales, M.L., Escarmis, C. and Girbes, T.: Effects and molecular action of ribosome-inactivating proteins on ribosomes from *Streptomyces lividans*. *Biochim. Biophys. Acta*, 1243 (1995) 85-93.
- 2377 Ganga, M.A., Gonzalez, M.P., Lopez-Lastra, M. and Sandino, A.M.: Polyacrylamide gel electrophoresis of viral genomic RNA as a diagnostic method for infectious pancreatic necrosis virus detection. *J. Virol. Methods*, 50 (1994) 227-236; *C.A.*, 122 (1995) 100882v.
- 2378 Gao, G., Widmer, J., Stapleton, D., Teh, T., Cox, T., Kemp, B.E. and Witters, L.A.: Catalytic subunits of the porcine nad rat 5'-AMP-activated protein kinase are members of the SHF1 protein kinase family. *Biochim. Biophys. Acta*, 1266 (1995) 73-82.
- 2379 Grombacher, T. and Kaina, B.: Constitutive expression and inducibility of O⁶-methylguanine-DNA methyltransferase and N-methylpurine-DNA glycosylase in rat liver cells exhibiting different status of differentiation. *Biochim. Biophys. Acta*, 1270 (1995) 63-72.
- 2380 Hata, S., Emi, Y., Iyanagi, T. and Osumi, T.: cDNA cloning of a putative G protein-coupled receptor from brain. *Biochim. Biophys. Acta*, 1261 (1995) 121-125.
- 2381 Hata, S., Inoue, T., Kosuga, K., Nakashima, T., Tsukamoto, T. and Osumi, T.: Identification of two splice isoforms of mRNA for mouse hepatocyte nuclear factor 4 (HNF-4). *Biochim. Biophys. Acta*, 1260 (1995) 55-61.
- 2382 Henco, K., Harders, J., Wiese, U. and Riesner, D.: Temperature gradient gel electrophoresis (TGGE) for the detection of polymorphic DNA and RNA. *Methods Mol. Biol. (Totowa)*, 31(Protocols for Gene Analysis) (1994) 211-228; *C.A.*, 122 (1995) 73179n.
- 2383 Hepburn, A.G.: SIZE: a program to determine the size of nucleic acid and protein molecules from gel mobilities. *BioTechniques*, 17 (1994) 1140-1143; *C.A.*, 122 (1995) 27216z.
- 2384 Hydén, H.: Isolation and biochemical mapping in the range of 10⁻⁷ to 10⁻¹² g of fresh, single mammalian neurons in brain. I. Techniques. *TrAC*, 14 (1995) 141-148.
- 2385 Inoguchi, K., Yoshioka, H., Khaleduzzaman, M. and Ninomiya, Y.: The mRNA for α 1(XIX) collagen chain, a new member of FACITs, contains a long unusual 3' untranslated region and displays many unique splicing variants. *J. Biochem. (Tokyo)*, 117 (1995) 137-146.
- 2386 Jaffe, J., Hochberg, M., Riss, J., Hasin, T., Reich, L. and Laskow, R.: Cloning, sequencing and expression of two isoforms of the murine oct-1 transcription factor. *Biochim. Biophys. Acta*, 1261 (1995) 201-209.
- 2387 Johnsen, L.B., Sørensen, E.S., Petersen, T.E. and Berglund, L.: Characterization of a bovine mammary gland PP3 cDNA reveals homology with mouse and rat adhesion molecule GlyCAM-1. *Biochim. Biophys. Acta*, 1260 (1995) 116-118.
- 2388 Kanbara, H., Okano, K., Kawamoto, K. and Furuyama, H.: Methods for determination and separation of mRNA using DNA probes and two-dimensional gel electrophoresis. *Jpn. Kokai Tokkyo Koho*, JP 06,294,796 [94,294,796] (Cl. GO1N33/50), 21 Oct. 1994, Appl. 93/84,433, 12 Apr. 1993; 8 pp.; *C.A.*, 122 (1995) 125336h.
- 2389 Kang, J., Harders, J., Riesner, D. and Henco, K.: TGGE in quantitative PCR of DNA and RNA. *Methods Mol. Biol. (Totowa)*, 31(Protocols for Gene Analysis) (1994) 229-235; *C.A.*, 122 (1995) 73180f.
- 2390 Kawakami, M.: (Gel electrophoresis of nucleic acids. Basic principle and applications). *Seibutsu Butsuri Kagaku*, 38 (1994) 349-353; *C.A.*, 122 (1995) 155308t - a review with 9 refs.
- 2391 Kikuchi, Y. and Suzuki-Fujita, K.: Synthesis and self-cleavage reaction of a chimeric molecule between RNase P-RNA and its model substrate. *J. Biochem. (Tokyo)*, 117 (1995) 197-200.
- 2392 Larochelle, S. and Suter, B.: Molecular cloning of the *Drosophila* homologue of the rat ribosomal protein L11 gene. *Biochim. Biophys. Acta*, 1261 (1995) 147-150.
- 2393 Li, H.-O., Tang, X., Kitabayashi, I., Gachelin, G., Chiu, R. and Yokoyama, K.: Induction by adenovirus-5 E1A of the differentiation phenotype of F9 teratocarcinoma cells involves a conserved region (CR1) of E1A. *Biochim. Biophys. Acta*, 1266 (1995) 148-156.

- 2394 Li, X., Wistow, G.J. and Piatigorsky, J.: Linkage and expression of the argininosuccinate lyase/ δ -crystallin genes of the duck: insertion of a CR1 element in the intergenic spacer. *Biochim. Biophys. Acta*, 1261 (1995) 25-34.
- 2395 Lina, B., Bornstein, N., Tuil, E., Meugnier, H., Bes, M. and Fleurette, J.: Use of pulsed-field gel electrophoresis, ribotyping and multilocus enzyme electrophoresis to characterize methicillin-resistant *Staphylococcus aureus* strains isolated from a nosocomial outbreak. *Zentralbl. Bakteriol., Suppl.*, 26 (1994) 104-106; C.A., 122 (1995) 234772t.
- 2396 Ludolph, D.C., Neff, A.W., Parker, M.A., Mescher, A.L., Smith, R.C. and Malacinski, G.M.: Cloning and expression of the axolotl proto-oncogene ski. *Biochim. Biophys. Acta*, 1260 (1995) 102-104.
- 2397 Lumelsky, N.L.: Decay of globin mRNA during megakaryocytic differentiation of erythroleukemic cell line is accomplished through shortening of the poly(A) tail and degradation from the 3' end of the transcript. *Biochim. Biophys. Acta*, 1261 (1995) 265-271.
- 2398 Martinotti, S., Toniato, E., Colagrande, A., Alesse, E., Alleva, C., Screpanti, I., Morrone, S., Scarpa, S., Frati, L., Hayday, A.C., Piovella, F. and Gulino, A.: Heavy-metal modulation of the human intercellular adhesion molecule (ICAM-1) gene expression. *Biochim. Biophys. Acta*, 1261 (1995) 107-114.
- 2399 Mercier, T., Gaillard-Sanchez, I., Martel, P., Seillan-Heberden, C.: Constitutive overexpression of c-fos protein in rat liver epithelial cells decreases TGF- β synthesis and increases TGF- β receptors. *Biochim. Biophys. Acta*, 1266 (1995) 64-72.
- 2400 Mizumoto, K., Muroya, K., Takagi, T., Omata-Yamada, T., Shibuta, H. and Iwasaki, K.: Protein factors required for *in vitro* transcription of Sendai virus genome. *J. Biochem. (Tokyo)*, 117 (1995) 527-534.
- 2401 Monteil, C., Fillastre, J.-P. and Morin, J.-P.: Expression and sub-cellular distribution of phosphoenolpyruvate carboxykinase in primary cultures of rabbit kidney proximal tubule cells: comparative study with renal and hepatic PEPCK *in vivo*. *Biochim. Biophys. Acta*, 1243 (1995) 437-445.
- 2402 Moor, N.A., Stepanov, V.G., Repkova, M.N., Venyaminova, A.G., Vratskikh, L.V., Yamkovo, V.I., Motorin, Yu.A. and Lavrik, O.I.: (The role of 3'-CCA end in the interaction of tRNA^{Phe} from *E. coli* and *Thermus thermophilus* with homologous phenylalanyl-tRNA synthetases). *Biochimiya (Moscow)*, 59 (1994) 1299-1303.
- 2403 Nacken, W., Kingsman, A.J., Kingsman, S.M., Sablitzky, F. and Sorg, C.: A homologue of the human MSS1 gene, a positive modulator of HIV-1 gene expression, is massively expressed in *Xenopus oocytes*. *Biochim. Biophys. Acta*, 1261 (1995) 293-295.
- 2404 Nagata, Y., Tashiro, F., Yi, S., Murakami, T., Maeda, S., Takahashi, K., Shimada, K., Okamura, H. and Yamamura, K.-i.: A 6-kb upstream region of the human transthyretin gene can direct developmental, tissue-specific, and quantitatively normal expression in transgenic mouse. *J. Biochem. (Tokyo)*, 117 (1995) 169-175.
- 2405 Ogata, K., Kurahashi, A., Ohno, R., Takahashi, K. and Terao, K.: Interaction of 5S rRNA-L5 protein complex, methionyl-tRNA, and methionyl-tRNA synthetase in the macromolecular ARS complex. *J. Biochem. (Tokyo)*, 117 (1995) 750-757.
- 2406 Oshiro, S. and Katsura, N.: An improved method for extraction of mRNA and protein from the fungus *Chaetomium gracile* with anhydrous hydrogen fluoride. *J. Biochem. (Tokyo)*, 117 (1995) 475-479.
- 2407 Reyes Engel, A. and Dieguez Lucena, J.L.: Determination of genetic expression by specific capture of RNA and its direct quantitation by free-zone capillary electrophoresis. *Span. ES 2,055,661 (Cl. C12Q1/68)*, 16 Aug. 1994, Appl. 9,300,103, 20 Jan. 1993; 5 pp.; C.A., 122 (1995) 101139v.
- 2408 Ribeiro, M.G., Pinto, R., Sa Miranda, M.C. and Suzuki, K.: Tay-Sachs disease: intron 7 splice junction mutation in two Portuguese patients. *Biochim. Biophys. Acta*, 1270 (1995) 44-51.
- 2409 Saito, S., Sakakura, S., Enomoto, M., Ichijo, M., Matsumoto, K. and Nakamura, T.: Hepatocyte growth factor promotes the growth of cytotrophoblasts by the paracrine mechanism. *J. Biochem. (Tokyo)*, 117 (1995) 671-676.
- 2410 Shimizu, M.: Specific aminoacylation of C4N hairpin RNAs with the cognate aminoacyl-adenylates in the presence of a dipeptide: origin of the genetic code. *J. Biochem. (Tokyo)*, 117 (1995) 23-26.
- 2411 Stromstedt, M. and Waterman, M.R.: A full-length cDNA encoding mouse adrenodoxin. *Biochim. Biophys. Acta*, 1261 (1995) 126-128.
- 2412 Tajima, S., Wachi, H. and Takehana, M.: Post-translational regulation of type I collagen synthesis by heparin in vascular smooth muscle cells. *J. Biochem. (Tokyo)*, 117 (1995) 353-358.
- 2413 Takeda, R., Mizobuchi, M., Muro, K., Sato, M. and Takahara, J.: Characterization of three cDNAs encoding two isozymes of an isoaspartyl protein carboxyl methyltransferase from human erythroid leukemia cells. *J. Biochem. (Tokyo)*, 117 (1995) 683-685.
- 2414 Tanaka, M., Maeda, K. and Nakashima, K.: Chicken α -enolase but not β -enolase has a Src-dependent tyrosine-phosphorylation site: cDNA cloning and nucleotide sequence analysis. *J. Biochem. (Tokyo)*, 117 (1995) 554-559.
- 2415 Terui, Y., Furukawa, Y., Sakoe, K., Ohta, M. and Saito, M.: Expression of differentiation-related phenotypes and apoptosis are independently regulated during myeloid cell differentiation. *J. Biochem. (Tokyo)*, 117 (1995) 77-84.
- 2416 Thomas, T., Macpherson, A. and Rogers, P.: Ceruloplasmin gene expression in the rat uterus. *Biochim. Biophys. Acta*, 1261 (1995) 77-82.
- 2417 Tomita, Y., Yuasa, C., Ni, R., Ishimura, K. and Ichihara, A.: Long-term maintenance of functional rat hepatocytes in primary culture by additions of pyruvate and various hormones. *Biochim. Biophys. Acta*, 1243 (1995) 329-335.
- 2418 Trayhurn, P., Duncan, J.S., Nestor, A., Thomas, M.E.A., Eastmond, N.C. and Rayner, D.V.: Rapid chemiluminescent detection of mRNAs on Northern blots with digoxigenin end-labelled oligonucleotides. *Electrophoresis (Weinheim)*, 16 (1995) 341-344.
- 2419 Van Neck, J.W., Medina, J.J., Onnekink, C., Schwartz, S.M. and Bloemers, H.P.J.: Expression of basic fibroblast growth factor and fibroblast growth factor receptor genes in cultured rat aortic smooth muscle cells. *Biochim. Biophys. Acta*, 1261 (1995) 210-214.

- 2420 Vasseur, S., Frigerio, J.-M., Dusetti, N.J., Keim, V., Dagorn, J.-C. and Iovanna, J.L.: Two transcripts are generated from the Pancreatitis Associated Protein II gene by alternative splicing in the 5' untranslated region. *Biochim. Biophys. Acta*, 1261 (1995) 272-274.
- 2421 Wachi, H., Seyama, Y., Yamashita, S. and Tajima, S.: Enhanced tropoelastin-degrading activity during cell passages in cultured smooth muscle cells. *Biochim. Biophys. Acta*, 1243 (1995) 393-398.
- 2422 Wilson, R., Allen, A.J., Oliver, J., Brookman, J.-L., High, S. and Bulleid, N.J.: The translocation, folding, assembly and redox-dependent degradation of secretory and membrane proteins in semi-permeabilized mammalian cells. *Biochem. J.*, 307 (1995) 679-687.
- 2423 Yasutake, H., Goda, T. and Takase, S.: Dietary regulation of sucrose-isomaltase gene expression in rat jejunum. *Biochim. Biophys. Acta*, 1243 (1995) 270-276.
- 2424 Yu, J., Nagarajan, S., Liu, J., Young, N. and Medof, M.E.: Cloning and characterization of the mouse PIG-A gene. *Biochim. Biophys. Acta*, 1255 (1995) 344-350.
- 2425 Zhang, H., Wang, J., Hwang, I. and Goodman, H.M.: Isolation and expression of an *Arabidopsis* 14-3-3-like protein gene. *Biochim. Biophys. Acta*, 1266 (1995) 113-116.
- See also 1865, 1868, 1939, 2050, 2090, 2093, 2279, 2436, 2441, 2455, 2457, 2484, 2503, 2516, 2523, 2527, 2558, 2567.
- 21c. *Nucleic acids, DNA*
- 2426 Alonso, A., Martin, P., Albarran, C. and Sancho, M.: Evaluation of the Tris-glycine/Tris-chloride discontinuous buffer system for the electrophoretic analysis of VNTR and STR loci. *Adv. Forensic Haemogenet.*, 5 (1994) 124-126; *C.A.*, 122 (1995) 179552u.
- 2427 Ashby, J., Tinwell, H., Lefevre, P.A. and Browne, M.A.: Comparative mutagenicity of 2-methylpropene (isobutene), its epoxide 2-methyl-1,2-epoxypropane and propylene oxide in the *in vitro* micronucleus test using human lymphocytes. *Mutagenesis*, 10 (1995) 85-89; *C.A.*, 122 (1995) 233163h.
- 2428 Aubert, D., Lapan, H., Lemaire, P., Foudrinier, F., Marx-Chemla, C., Bonhomme, A. and Pinon, J.-M.: Rapid detection of toxoplasmic nucleic acid by enzyme-linked immunofiltration-assay after membrane transfer. *Electrophoresis (Weinheim)*, 16 (1995) 354-356.
- 2429 Berka, J., Pariat, Y.F., Müller, O., Hebenbrock, K., Heiger, D.N., Foret, F. and Karger, B.L.: Sequence dependent migration behavior of double-stranded DNA in capillary electrophoresis. *Electrophoresis (Weinheim)*, 16 (1995) 377-388.
- 2430 Bianchi, N., Mischiati, C., Feriotto, G., Fiorentino, D., Di Biase, S., Apicella, N. and Gambari, R.: Capillary electrophoresis: detection of hybridization between synthetic oligonucleotides and HIV-1 genomic DNA amplified by polymerase-chain reaction. *J. Virol. Methods*, 47 (1994) 321-329; *C.A.*, 122 (1995) 73139z.
- 2431 Birren, B. and Lai, E.: Rapid pulsed field separation of DNA molecules up to 250 kb. *Nucleic Acids Res.*, 22 (1994) 5366-5370; *C.A.*, 122 (1995) 100900z.
- 2432 Callaghan, A. and Laias, N.: Pulsed-field gel electrophoresis of mosquito DNA. *Trends Genet.*, 11 (1995) 40-41; *C.A.*, 122 (1995) 206328j.
- 2433 Cane, P.A. and Pringle, C.R.: Molecular epidemiology of respiratory syncytial virus: a review of the use of reverse transcription-polymerase chain reaction in the analysis of genetic variability. *Electrophoresis (Weinheim)*, 16 (1995) 329-333.
- 2434 Carlsson, C., Larsson, A., Jonsson, M. and Norden, B.: Dancing DNA in capillary solution electrophoresis. *J. Am. Chem. Soc.*, 117 (1995) 3871-3872; *C.A.*, 122 (1995) 209053b.
- 2435 Carter, G.L., Wilson, L.J. and Dunn-Coleman, N.S.: Molecular characterization of chromosomes by pulsed field electrophoresis. *Prog. Ind. Microbiol.*, 29(Aspergillus: 50 Years on) (1994) 667-683; *C.A.*, 122 (1995) 97509w - a review with refs.
- 2436 Ciesla, J., Weiner, K.X.B., Weiner, R.S., Reston, J.T., Maley, G.F. and Maley, F.: Isolation and expression of rat thymidylate synthase cDNA: Phylogenetic comparison with human and mouse thymidylate synthases. *Biochim. Biophys. Acta*, 1261 (1995) 233-243.
- 2437 Den Dunnen, J.T., Grootsholten, P.M. and van Ommen, G.J.B.: Pulsed-field gel electrophoresis in the analysis of genomic DNA and YAC clones. In: Davies, K.E. (Editor), *Hum. Genet. Dis. Anal.*, (2nd ed.), IRL, Oxford, 1993, pp. 35-58; *C.A.*, 122 (1995) 97465d.
- 2438 Doria, A., Warram, J.H., Rich, S.S. and Krolewski, A.S.: Angiotensin I-converting enzyme (ACE): estimation of DNA haplotypes in unrelated individuals using denaturing gradient gel blots. *Hum. Genet.*, 94 (1994) 117-123; *C.A.*, 122 (1995) 231881s.
- 2439 Duewer, D.L., Currie, L.A., Reeder, D.J., Leigh, S.D., Liu, H.-K. and Mudd, J.L.: Interlaboratory comparison of autoradiographic DNA profiling measurements. 2. Measurement uncertainty and its propagation. *Anal. Chem.*, 67 (1995) 1220-1231.
- 2440 Fukunaga, M. and Takahashi, Y.: Pulsed field gel electrophoresis analysis of *Borrelia burgdorferi sensu lato* isolated in Japan and taxonomic implications with Lyme disease spirochetes. *Microbiol. Immunol.*, 38 (1994) 747-751; *C.A.*, 122 (1995) 27521v.
- 2441 Gow, J., Cash, P., Behan, W., McGarry, F., Simpson, K. and Behan, P.: Detection of picornavirus genomic and template RNA strands by a novel semi-nested polymerase chain reaction-technique and agarose gel electrophoresis. *Electrophoresis (Weinheim)*, 16 (1995) 338-340.
- 2442 Gyobu, Y., Hosorogi, S., Kodama, H. and Shimada, T.: (Pulsed-field gel electrophoresis of *Vibrio cholerae* O1 distribution in the environment). *Toyama-ken Eisei Kenkyusho Nenpo*, 17 (1994) 125-128; *C.A.*, 122 (1995) 128365c.
- 2443 Hellman, B., Vaghef, H. and Bostroem, B.: The concepts of tail moment and tail inertia in the single cell gel electrophoresis assay. *Mutat. Res.*, 336 (1995) 123-131; *C.A.*, 122 (1995) 180389c.
- 2444 Holland, M.M., Turni, L.A., DelRio, S., Marino, M., Lofts, R.S., Fisher, D.L., Ross, J., Schumm, J.W. and Williams, P.L.: Typing human DNA using capillary electrophoresis: comparison of slab gel and capillary formats. *Adv. Forensic Haemogenet.*, 5 (1994) 156-159; *C.A.*, 122 (1995) 155526n.
- 2445 Hutson, M.S., Holzwarth, G., Duke, T. and Viovy, J.-L.: Two-dimensional motion of DNA bands during 120° pulsed-field gel electrophoresis. I. Effect of molecular weight. *Biopolymers*, 35 (1995) 297-306; *C.A.*, 122 (1995) 155536r.

- 2446 Inoue, T., Osumi, T. and Hata, S.: Molecular cloning and functional expression of a cDNA for mouse squalene synthase. *Biochim. Biophys. Acta*, 1260 (1995) 49-54.
- 2447 Ishihara, H., Engel, J.D. and Yamamoto, M.: Structure and regulation of the chicken GATA-3 gene. *J. Biochem. (Tokyo)*, 117 (1995) 499-508.
- 2448 Johansson, E., Joensson, J.O., Henriksson, P., Brismar, K. and Heneen, W.K.: Comparison of electrophoretic methods for identification of the 1BL/1RS rye translocation in wheat, simultaneous screening for high-molecular-weight glutenins, and detection of chromosome deletions. *J. Genet. Breed.*, 48 (1994) 291-299; *C.A.*, 122 (1995) 182488h.
- 2449 Johnston, J.R.: Pulsed field gel electrophoresis. In: Johnston, J.R. (Editor), *Mol. Genet. Yeast*, IRL, Oxford, 1994, pp. 83-95; *C.A.*, 122 (1995) 208836x - a review with 54 refs.
- 2450 Jong, A.Y., Wang, B. and Zhang, S.Q.: Pulsed field gel electrophoresis labeling method to study the pattern of *Saccharomyces cerevisiae* chromosomal DNA synthesis during the G1/S phase of the cell cycle. *Anal. Biochem.*, 227 (1995) 32-39.
- 2451 Jonsson, M., Jacobsson, U., Norden, B. and Eriksson, T.: Electrophoretic orientation of large DNA in free solution. *Eur. Space Agency, [Spec. Publ.] ESA SP*, ESA SP-1132(Vol. 4 Final Reports of Sounding Rocket Experiments in Fluid Science and Materials Sciences) (1994) 87-97; *C.A.*, 122 (1995) 182499n.
- 2452 Kaufmann, M.E. and Pitt, T.L.: Pulsed-field gel electrophoresis of bacterial DNA. In: Chart, H. (Editor), *Methods Pract. Lab. Bacteriol.*, CRC, Boca Raton, 1994, pp. 83-92; *C.A.*, 122 (1995) 208817s - a review with 14 refs.
- 2453 Kim, Y. and Morris, M.D.: Rapid pulsed field capillary electrophoretic separation of megabase nucleic acids. *Anal. Chem.*, 67 (1995) 784-786.
- 2454 Kleparnik, K., Mala, Z., Doskar, J., Rosypal, S. and Bocek, P.: An improvement of restriction analysis of bacteriophage DNA using capillary electrophoresis in agarose solution. *Electrophoresis (Weinheim)*, 16 (1995) 366-376.
- 2455 Komura, J.-i., Okada, T. and Ono, T.: Repression of transient expression by DNA methylation in transcribed regions of reporter genes introduced into cultured human cells. *Biochim. Biophys. Acta*, 1260 (1995) 73-78.
- 2456 Laber, T.L., Giese, S.A., Iverson, J.T. and Liberty, J.A.: Validation studies on the forensic analysis of restriction fragment length polymorphism (RFLP) on LE agarose gels without ethidium bromide: effects of contaminants, sunlight, and the electrophoresis of varying quantities of deoxyribonucleic acid (DNA). *J. Forensic Sci.*, 39 (1994) 707-730; *C.A.*, 122 (1995) 74088u.
- 2457 Lee, Y.L., Helman, L., Hoffman, T. and Laborda, J.: dlk, pG2 and Pref-1 mRNAs encode similar proteins belonging to the EGF-like superfamily. Identification of polymorphic variants of this RNA. *Biochim. Biophys. Acta*, 1261 (1995) 223-232.
- 2458 Lim, R.W., Zhu, C.Y. and Stringer, B.: Differential regulation of primary response gene expression in skeletal muscle cells through multiple signal transduction pathways. *Biochim. Biophys. Acta*, 1266 (1995) 91-100.
- 2459 Lin, B., Diao, H., Zhang, Y. and Chu, X.: (Determination of polymerase chain reaction-amplified products of hepatitis B virus with non-gel sieving capillary electrophoresis). *Fenxi Huaxue*, 22 (1994) 449-452; *C.A.*, 122 (1995) 152414p.
- 2460 Liu, M.-S., Zang, J., Evangelista, R.A., Rampal, S. and Chen, F.-T.A.: Double-stranded DNA analysis by capillary electrophoresis with laser-induced fluorescence using ethidium bromide as an intercalator. *BioTechniques*, 18 (1995) 316-323; *C.A.*, 122 (1995) 128034u.
- 2461 Ma, C. and Bloomfield, V.A.: Gel electrophoresis measurement of counterion condensation on DNA. *Biopolymers*, 35 (1995) 211-216; *C.A.*, 122 (1995) 127907u.
- 2462 Manicardi, G.C., Bizzaro, D., Azzoni, P. and Bianchi, U.: Cytological and electrophoretic analysis of DNA methylation in the holocentric chromosomes of *Megoura viciae* (Homoptera, Aphididae). *Genome*, 37 (1994) 625-630; *C.A.*, 122 (1995) 179748n.
- 2463 Martin, T., Hughes, S., Hughes, K. and Dawson, M.: Direct sequencing of PCR amplified pig PrP genes. *Biochim. Biophys. Acta*, 1270 (1995) 211-214.
- 2464 Matsumoto, M. and Doi, M.: Brownian dynamics simulation of DNA gel electrophoresis. *Mol. Simul.*, 12 (1994) 219-226; *C.A.*, 122 (1995) 73160z.
- 2465 Maule, J.C.: Colored microparticles for clear visualization beads and plugs. *Trends Genet.*, 11 (1995) 127; *C.A.*, 122 (1995) 234658k.
- 2466 Moll, R., Schmidtke, S. and Schäfer, G.: Nucleotide sequence of a gene cluster encoding ribosomal proteins in the thermoacidophilic crenarchaeon *Sulfolobus acidocaldarius*. *Biochim. Biophys. Acta*, 1261 (1995) 315-318.
- 2467 Monaco, A.P., Larin, Z. and Lehrach, H.: Construction of yeast artificial chromosome libraries by pulsed-field gel electrophoresis. *Mol. Biotechnol.*, 1 (1994) 241-249; *C.A.*, 122 (1995) 97747x.
- 2468 Nagy, A., Vagvoelgyi, C. and Ferenczy, L.: Simple method for obtaining chromosomal DNA for orthogonal field alteration gel electrophoresis analysis from *Phaffia rhodozyma*. *BioTechniques*, 18 (1995) 63-65; *C.A.*, 122 (1995) 75838a.
- 2469 Nedved, M.L. and Moe, G.R.: The use of affinity coelectrophoresis to characterize cooperative, nonspecific maleic acid binding peptides. *Anal. Biochem.*, 227 (1995) 80-84.
- 2470 Nixon, G.I. and Slater, G.W.: DNA electrophoretic collisions with single obstacles. *Phys. Rev. E.: Stat. Phys., Plasmas, Fluids, Relat. Interdiscip. Top.*, 50 (1994) 5033-5038; *C.A.*, 122 (1995) 50522e.
- 2471 Nozawa, M.: Method with antibody for DNA determination. *Jpn. Kokai Tokkyo Koho JP 06,289,027 [94,289,027]* (Cl. GO1N33/561), 18 Oct. 1994, Appl. 93/77,926, 05 Apr. 1993; 3 pp.; *C.A.*, 122 (1995) 50752e.
- 2472 Ohizumi, H., Masuda, Y., Nakajo, S., Sakai, I., Ohsawa, S. and Nakaya, K.: Geranylgeraniol is a potent inducer of apoptosis in tumor cells. *J. Biochem. (Tokyo)*, 117 (1995) 11-13.
- 2473 Raghava, G.P.S.: DNAOPT: a computer program to aid optimization of DNA gel electrophoresis and SDS-PAGE. *BioTechniques*, 18 (1995) 274-280; *C.A.*, 122 (1995) 127921u.
- 2474 Robertson, J.M.: Evaluation of native and denaturing polyacrylamide gel electrophoresis for short tandem repeat analysis. *Adv. Forensic Haemogenet.*, 5 (1994) 320-322; *C.A.*, 122 (1995) 152564n.
- 2475 Rodriguez, P., Allardet-Servent, A., de Barbeyrac, B., Ramuz, M. and Bebear, C.: Genetic variability among *Chlamydia trachomatis* reference and clinical strains analyzed by pulsed-field gel electrophoresis. *J. Clin. Microbiol.*, 32 (1994) 2921-2928; *C.A.*, 122 (1995) 98493y.

- 2476 Schmidt, M., Svendsen, I. and Feierabend, J.: Analysis of the primary structure of the chloroplast isozyme of triosephosphate isomerase from rye leaves by protein and cDNA sequencing indicates a eukaryotic origin of its gene. *Biochim. Biophys. Acta*, 1261 (1995) 257-264.
- 2477 Semenov, A.N., Duke, T.A.J. and Viovy, J.-L.: Gel electrophoresis of DNA in moderate fields: the effect of fluctuations. *Phys. Rev. E: Stat. Phys., Plasmas, Fluids, Relat. Interdiscip. Top.*, 51 (1995) 1520-1537; *C.A.*, 122 (1995) 155532m.
- 2478 Sinha, B.K., Yamazaki, H., Eliot, H.M., Schneider, E., Borner, M.M. and O'Connor, P.M.: Relationships between proto-oncogene expression and apoptosis induced by anticancer drugs in human prostate tumor cells. *Biochim. Biophys. Acta*, 1270 (1995) 12-18.
- 2479 Slater, G.W., Mayer, P., Hubert, S.J. and Drouin, G.: The biased reptation model of DNA gel electrophoresis: a user guide for constant field mobilities. *Appl. Theor. Electrophor.*, 4 (1994) 71-79; *C.A.*, 122 (1995) 50534k.
- 2480 Spitzner, J.R., Chung, I.K. and Muller, M.T.: Determination of 5' and 3' DNA triplex interference boundaries reveals the core DNA binding sequence for topoisomerase II. *J. Biol. Chem.*, 270 (1995) 5932-5943.
- 2481 Stellwagen, N.C.: Toward understanding electrophoretic anomalies in polyacrylamide gels: kilobase-sized DNAs. In: Sarma, R.H. and Sarma, M.H. (Editors), *Struct. Biol. State of the Art, Proc. Conversation Discip. Biomol. Stereodyn.*, 8th 1993, Adenine, Schenectady, 1994, pp. 285-299; *C.A.*, 122 (1995) 126452e.
- 2482 Takasaki, Y., Iwanaga, T. and Tsuji, N.: Quantitative analysis of DNA and proteins on personal computer. *Sasego Kogyo Koto Senmon Gakko Kenkyu Hokoku*, 31 (1994) 23-27; *C.A.*, 122 (1995) 155684n.
- 2483 Tornar, A., Garofano, L., Vecchio, C., Pizzamiglio, M., Albonici, L. and Manzari, V.: The use of capillary electrophoresis in forensic DNA analysis. *Adv. Forensic Haemogenet.*, 5 (1994) 390-392; *C.A.*, 122 (1995) 153480u.
- 2484 Tulchinsky, E., Grigorian, M., Tkatch, T., Georgiev, G. and Lukanidin, E.: Transcriptional regulation of the mts1 gene in human lymphoma cells: the role of DNA-methylation. *Biochim. Biophys. Acta*, 1261 (1995) 243-248.
- 2485 Uesugi, M., Kusakabe, T. and Sugiura, Y.: Hydrophobic DNA binding of esperamicin requires conformational distortion of the host DNA. *Biochim. Biophys. Acta*, 1261 (1995) 99-106.
- 2486 Uhrberg, M., Enczmann, J. and Wernet, P.: Rapid DNA cross-match analysis of HLA class II genotypic polymorphisms by temperature gradient gel electrophoresis in unrelated bone marrow donor selection. *Eur. J. Immunogenet.*, 21 (1994) 313-324; *C.A.*, 122 (1995) 24765s.
- 2487 Vijg, J.: Detecting individual genetic variation. *BioTechnology*, 13 (1995) 137-139; *C.A.*, 122 (1995) 124426a.
- 2488 Wang, M.-Y. and Liehr, J.G.: Lipid hydroperoxide-induced endogenous DNA adducts in hamster: possible mechanism of lipid hydroperoxide-mediated carcinogenesis. *Arch. Biochem. Biophys.*, 316 (1995) 38-46.
- 2489 Wheeler, D.L. and Chrambach, A.: Computer simulation of the directional displacement of rod-shaped, arc-shaped, and circular objects in an array of obstacles, representing a simple model for the gel electrophoresis of small DNA. *Biopolymers*, 35 (1995) 179-185; *C.A.*, 122 (1995) 128022p.
- 2490 Wolf, S.M. and Vouros, P.: Incorporation of sample stacking techniques into the capillary electrophoresis CF-FAB mass spectrometric analysis of DNA adducts. *Anal. Chem.*, 67 (1995) 891-900.
- 2491 Woodward, T.M., Carlson, J., McClelland, C. and DeMartini, J.C.: Analysis of lentiviral genomic variation by denaturing gradient gel electrophoresis. *BioTechniques*, 17 (1994) 366-371; *C.A.*, 122 (1995) 97797p.
- 2492 Xu, W.-H., Sato, Y., Ikeda, M. and Yamashita, O.: Molecular characterization of the gene encoding the precursor protein of diapause hormone and pheromone biosynthesis activating neuropeptide (DH-PBAN) of the silkworm, *Bombyx mori* and its distribution in some insects. *Biochim. Biophys. Acta*, 1261 (1995) 83-89.
- 2493 Yakubu, D.E. and Pennington, T.H.: Epidemiological evaluation of *Neisseria meningitidis* serogroup B by pulsed-field gel electrophoresis. *FEMS Immunol. Med. Microbiol.*, 10 (1995) 185-190; *C.A.*, 122 (1995) 232010u.
- 2494 Yarmola, E. and Chrambach, A.: Band width measurement in automated gel electrophoresis apparatus: DNA dispersion in a discontinuous system and in a single buffer. *Electrophoresis (Weinheim)*, 16 (1995) 345-349.
- 2495 Yurovsky, V.V., Schulze, D.H. and White, B.: Analysis of diversity of T cell antigen receptor genes using polymerase chain reaction and sequencing gel electrophoresis. *J. Immunol. Methods*, 175 (1994) 227-236; *C.A.*, 122 (1995) 78532g.
- 2496 Zhizhina, G.P., Korobko, V.G. and Alessenko, A.V.: (Correlation of DNA degradation induced by the tumour necrosis factor α with nuclear sphingosin content and DNA peroxide accumulation). *Biokhimiya (Moscow)*, 59 (1994) 1756-1765.
- 2497 Zhou, Y.-H. and Ragan, M.A.: Characterization of the polyubiquitin gene in the marine red alga *Gracilaria verrucosa*. *Biochim. Biophys. Acta*, 1261 (1995) 215-222.
- See also 1838, 1844, 1862, 1865, 1868, 1877, 1881, 1882, 1939, 2050, 2265, 2279, 2370, 2382, 2383, 2386, 2389, 2390, 2395, 2425, 2524, 2554, 2647.
- 21d. *Structural studies on RNA and RNA mapping*
- 2498 Aoki, H., Tanaka, K. and Ida, S.: The genomic organization of the gene encoding a nitrate-inducible ferredoxin-NADP⁺ oxidoreductase from rice roots. *Biochim. Biophys. Acta*, 1229 (1995) 389-392.
- 2499 Bogdanova, S.L., Degtyarev, A.I., Baranov, P.V., Dokudovskaya, S.S., Lavrik, I.N., Dontsova, O.A., Oretskaya, T.S., Krynetskaya, N.F., Shabarova, Z.A. and Bogdanov, A.A.: (Site-directed cleavage of single internucleotide bonds of ribosomal RNA). *Biokhimiya (Moscow)*, 60 (1995) 297-307.
- 2500 Cramer, G.S., Wang, L.-F. and Eaton, B.T.: Differentiation of cognate dsRNA genome segments of bluetongue virus reassortants by temperature gradient gel electrophoresis. *J. Virol. Methods*, 51 (1995) 211-220; *C.A.*, 122 (1995) 182482b.
- 2501 Heller, C.: Field inversion gel electrophoresis. *Methods Mol. Biol. (Totowa)*, 31(Protocols for Gene Analysis) (1994) 135-146; *C.A.*, 122 (1995) 75617c - a review with 18 refs.

- 2502 Yamamoto, H., Naruse, A., Ohsaki, T. and Sekiguchi, J.: Nucleotide sequence and characterization of the large mitochondrial rRNA gene of *Penicillium urticae*, and its comparison with those of other filamentous fungi. *J. Biochem. (Tokyo)*, 117 (1995) 888-896.
- See also 2376, 2420, 2561.
- 21e. *Structural studies on DNA and DNA mapping*
- 2503 Allard, S., Labbé, M. and Falardeau, P.: Recombination of endogenous D₂ dopamine receptor gene with a metallothionein promoter in GH4C1 cells confers functional and inducible D₂ response. *Biochim. Biophys. Acta*, 1260 (1995) 43-48.
- 2504 Appel, E., Eisenberg, S. and Roitelman, J.: Improved PCR amplification/HhaI restriction for unambiguous determination of apolipoprotein E alleles. *Clin. Chem. (Washington)*, 41 (1995) 187-190.
- 2505 Athma, P., Liu, J. and Swift, M.: PCR detection of the TaqI restriction fragment length polymorphism linked to the ataxia telangiectasia locus. *Clin. Chem. (Washington)*, 41 (1995) 625-626.
- 2506 Baba, Y.: (PCR analysis and DNA sequencing by capillary electrophoresis in human genome project). *Dojin News*, 71 (1994) 8-14; *C.A.*, 122 (1995) 97534a.
- 2507 Baba, Y., Tshako, M., Hayashi, K., Okubo, K., Matsubara, K., Hattori, M., Sakaki, Y., Miki, T. and Ogihara, T.: (Human genome analysis and gene diagnosis using capillary electrophoresis with laser-induced fluorescence detector). *Kuromatogurafi*, 15 (1994) 218-219; *C.A.*, 122 (1995) 124376j.
- 2508 Bautsch, W.: Bacterial genome mapping by two-dimensional pulsed-field gel electrophoresis (2D-PFGE). *Mol. Biotechnol.*, 2 (1994) 29-44; *C.A.*, 122 (1995) 124298k.
- 2509 Boultonwood, J.: Physical mapping of the human genome by pulsed field gel electrophoresis. *Methods Mol. Biol. (Totowa)*, 31(Protocols for Gene Analysis) (1994) 121-133; *C.A.*, 122 (1995) 73178m.
- 2510 Brillante, R., Yang, V., Proos, A. and Burnett, L.: PCR analysis of hair root specimens to detect Tay-Sachs disease carriers in Ashkenazi Jews. *Clin. Chem. (Washington)*, 41 (1995) 321-322.
- 2511 Butler, J.M., McCord, B.R., Jung, J.M. and Allen, R.O.: Rapid analysis of the short tandem repeat HUMB01 by capillary electrophoresis. *BioTechniques*, 17 (1994) 1062-1070; *C.A.*, 122 (1995) 231925j.
- 2512 Chung, I.K., Soisson, S.M. and Muller, M.T.: Clustering of Sp1 sites near the promoter region of ICP34.5 in herpes simplex virus type 1. *J. Biochem. (Tokyo)*, 117 (1995) 19-22.
- 2513 Dahm-Daphi, J. and Dikomey, E.: Separation of DNA fragments induced by ionizing irradiation using a graded-field gel electrophoresis. *Int. J. Radiat. Biol.*, 67 (1995) 161-168; *C.A.*, 122 (1995) 234335c.
- 2514 Donabedian, S., Chow, J.W., Shlaes, D.M., Green, M. and Zervos, M.J.: DNA hybridization and contour-clamped homogeneous electric field electrophoresis for identification of enterococci to the species level. *J. Clin. Microbiol.*, 33 (1995) 141-145; *C.A.*, 122 (1995) 152550e.
- 2515 Dovichi, N.J.: Capillary gel electrophoresis for DNA sequencing: separation and detection. In: Landers, J.P. (Editor), *Handb. Capillary Electrophor.*, CRC, Boca Raton, 1994, pp. 369-387; *C.A.*, 122 (1995) 97473e.
- 2516 Duncan, M.K., Roth, H.J., Thompson, M., Kantorow, M. and Piatigorsky, J.: Chicken β B1 crystallin: gene sequence and evidence for functional conservation of promoter activity between chicken and mouse. *Biochim. Biophys. Acta*, 1261 (1995) 68-76.
- 2517 Eng, B., Ainsworth, P. and Wayne, J.S.: Anomalous migration of PCR products using nondenaturing polyacrylamide gel electrophoresis: the amelogenin sex-typing system. *J. Forensic Sci.*, 39 (1994) 1356-1359; *C.A.*, 122 (1995) 48617w.
- 2518 Fuller, C.W.: An improved polyacrylamide gel containing ethylene glycol for DNA sequencing. *PCT Int. Appl. WO 94 23,092* (Cl. C25B7/00), 13 Oct. 1994, US Appl. 37,608, 26 Mar. 1993; 8 pp.; *C.A.*, 122 (1995) 73953k.
- 2519 Gejman, P.V. and Weinstein, L.S.: Detection of mutations and polymorphisms of G_{3a} subunit gene by denaturing gradient gel electrophoresis. *Methods Enzymol.*, 237(Heterotrimeric G Proteins) (1994) 308-320; *C.A.*, 122 (1995) 124030s - a review with 46 refs.
- 2520 Greiner, T.C., Raffeld, M., Lutz, C., Dick, F. and Jaffe, E.S.: Technical advance. Analysis of T cell receptor- γ gene rearrangements by denaturing gradient gel electrophoresis of GC-clamped polymerase chain reaction products: correlation with tumor-specific sequences. *Am. J. Pathol.*, 146 (1995) 46-55; *C.A.*, 122 (1995) 211880n.
- 2521 Gyobu, Y., Isobe, J. and Tanaka, D.: (Use of pulsed field gel electrophoresis for epidemiology of *Salmonella*). *Toyama-ken Eisei Kenkyusho Nenpo*, 17 (1994) 129-132; *C.A.*, 122 (1995) 128469q.
- 2522 Hillier, A.J. and Davidson, B.E.: The use of pulsed-field gel electrophoresis for microbial strain identification. *Australas. Biotechnol.*, 4 (1994) 167-168; *C.A.*, 122 (1995) 100661x.
- 2523 Hoesche, C., Bartsch, P. and Kilimann, M.W.: The CRE consensus sequence in the synapsin I gene promoter region confers constitutive activation but no regulation by cAMP in neuroblastoma cells. *Biochim. Biophys. Acta*, 1261 (1995) 249-256.
- 2524 Huh, G.H., Matsuura, Y., Meshi, T. and Iwabuchi, M.: Differential expression of the two types of histone H2A genes in wheat. *Biochim. Biophys. Acta*, 1261 (1995) 155-160.
- 2525 Itoh, M., Nakamura, M., Suzuki, T., Kawai, K., Horitsu, H. and Takamizawa, K.: Mechanism of chromium(VI) toxicity in *Escherichia coli*: is hydrogen peroxide essential in Cr(VI) toxicity? *J. Biochem. (Tokyo)*, 117 (1995) 780-786.
- 2526 Jaakkola, S., Vornanen, T., Leinonen, J., Rannikko, S. and Stenman, U.-H.: Detection of prostatic cells in peripheral blood: correlation with serum concentrations of prostate-specific antigen. *Clin. Chem. (Washington)*, 41 (1995) 182-186.
- 2527 Kamatchi, G.L., Kofuji, P., Wang, J.B., Fernando, J.C.R., Liu, Z., Mathura, J.R., Jr. and Burt, D.R.: GABA_A receptor β ₁, β ₂, and β ₃ subunits: comparisons in DBA/2J and C57BL/6J mice. *Biochim. Biophys. Acta*, 1261 (1995) 134-142.
- 2528 Katsube, T. and Fukumaki, Y.: A role for the distal CCAAT box of the β -globin gene in Hb switching. *J. Biochem. (Tokyo)*, 117 (1995) 68-76.

- 2529 Kawabata, H., Anzai, N., Yoshida, Y. and Okuma, M.: A new method for quantitative estimation of the degree of DNA fragmentation utilizing agarose gel electrophoresis. *Int. J. Hematol.*, 59 (1994) 311-316; *C.A.*, 122 (1995) 100887a.
- 2530 Kleparnik, K., Garner, M. and Bocek, P.: Injection bias of DNA fragments in capillary electrophoresis with sieving. *J. Chromatogr. A*, 698 (1995) 375-383.
- 2531 Kneba, M., Bolz, I., Linke, B., Bertram, J., Rothaupt, D. and Hiddemann, W.: Characterization of clone-specific rearranged T-cell receptor γ -chain genes in lymphomas and leukemias by the polymerase chain reaction and DNA sequencing. *Blood*, 84 (1994) 574-581; *C.A.*, 122 (1995) 103446s.
- 2532 Knoblauch, H., Weiss, N., Eggersdorfer, I., Keller, C. and Schuster, H.: A nonisotopic single-strand conformation polymorphism protocol using a direct blotting electrophoresis, a chemiluminescent detection system, and a multiplex approach. *PCR Methods Appl.*, 4 (1994) 52-55; *C.A.*, 122 (1995) 97851b.
- 2533 Kuff, E.L. and Mietz, J.A.: Analysis of DNA restriction enzyme digests by two-dimensional electrophoresis in agarose gels. *Methods Mol. Biol. (Totowa)*, 31(Protocols for Gene Analysis) (1994) 177-186; *C.A.*, 122 (1995) 97815t.
- 2534 Kuick, R., Asakawa, J.-i., Neel, J.V., Satoh, C. and Hanash, S.M.: High yield of restriction fragment length polymorphisms in two-dimensional separations of human genomic DNA. *Genomics*, 25 (1995) 345-353; *C.A.*, 122 (1995) 179656f.
- 2535 Kurvinen, K., Hietanen, S., Syrjaenen, K. and Syrjaenen, S.: Rapid and effective detection of mutations in the p53 gene using nonradioactive single-strand conformation polymorphism (SSCP) technique applied on PhastSystem. *J. Virol. Methods*, 51 (1995) 43-54; *C.A.*, 122 (1995) 124389r.
- 2536 Lareu, M.V., Phillips, C.P., Pestoni, C., Barros, F., Munoz, J. and Carracedo, A.: Anomalous electrophoretic behavior of HUMACTBP2 (SE33). *Adv. Forensic Haemogenet.*, 5 (1994) 121-123; *C.A.*, 122 (1995) 152562k.
- 2537 Laubscher, K.H., Glew, R.H., Lee, R.E. and Okinaka, R.T.: Use of denaturing gradient gel electrophoresis to identify mutant sequences in the β -glucocerebrosidase gene. *Hum. Mutat.*, 3 (1994) 411-415; *C.A.*, 122 (1995) 97862f.
- 2538 Law, H.-Y., Ong, J., Yoon, C.-S., Cheng, H., Tan, C.-L. and Ng, I.: Rapid antenatal diagnosis in Chinese of β -thalassaemia caused by the common 4-bp deletion in codons 41/42 using high-resolution agarose gel electrophoresis and heteroduplex detection. *Biochem. Med. Metab. Biol.*, 53 (1994) 149-151; *C.A.*, 122 (1995) 124435c.
- 2539 Lefevre, J.C., Gasc, A.M., Lemozy, J., Sicard, A.M. and Faucon, G.: Pulsed field gel electrophoresis for molecular epidemiology of penicillin resistant *Streptococcus pneumoniae* strains. *Pathol. Biol.*, 42 (1994) 547-552; *C.A.*, 122 (1995) 27070x.
- 2540 Little, R.D. and Schildkraut, C.L.: Mapping sites of replication initiation and termination in circular viral genomes using two-dimensional agarose gel electrophoresis. *Methods Mol. Genet.*, 4(Molecular Virology T) (1994) 326-344; *C.A.*, 122 (1995) 97543c - a review with 26 refs.
- 2541 Loomans, D., Sokolov, I.M. and Blumen, A.: Tension and mobility of a DNA fragment in the lakes-straits model. *Macromol. Theory Simul.*, 4 (1995) 145-154; *C.A.*, 122 (1995) 75843y.
- 2542 Maekawa, M., Sudo, K., Kanno, T., Kotani, K., Key, D.C., Ishikawa, J., Izumi, M. and Etoh, K.: Genetic basis of the silent phenotype of serum butyrylcholinesterase in three compound heterozygotes. *Clin. Chim. Acta*, 235 (1995) 41-57.
- 2543 Makowski, G.S., Aslanzadeh, J. and Hopfer, S.M.: *In situ* PCR amplification of Guthrie card DNA to detect cystic fibrosis mutations. *Clin. Chem. (Washington)*, 41 (1995) 477-479.
- 2544 Murray, V., Moore, A.G., Matias, C. and Wickham, G.: The interaction of hedamycin and DC92-B in a sequence selective manner with DNA in intact human cells. *Biochim. Biophys. Acta*, 1261 (1995) 195-200.
- 2545 Na, W.-J. and Chung, Y.-B.: The variable number of tandem repeats allele distribution in the Korean population (II): analysis by polymerase chain reaction and high-resolution agarose gel electrophoresis. *Mol. Cells*, 4 (1994) 495-504; *C.A.*, 122 (1995) 152624g.
- 2546 Nair, S., Poh, C.L., Lim, Y.S., Tay, L. and Goh, K.T.: Genome fingerprinting of *Salmonella typhi* by pulsed-field gel electrophoresis for subtyping common phage types. *Epidemiol. Infect.*, 113 (1994) 391-402; *C.A.*, 122 (1995) 152560h.
- 2547 Nam, H.S., Kim, K.E., Lee, L.-J. and Kim, E.-H.: Polymerase chain reaction and denaturing gradient gel electrophoresis for screening new alleles in HLA-DQA. *Mol. Cells*, 5 (1995) 52-57; *C.A.*, 122 (1995) 184922u.
- 2548 Nissen, H., Hansen, P.S., Færgeman, O. and Hørder, M.: Mutation screening of the codon 3500 region of the apolipoprotein B gene by denaturing gradient-gel electrophoresis. *Clin. Chem. (Washington)*, 41 (1995) 419-423.
- 2549 Ochiai, Y., Inazawa, J., Ueyama, H. and Ohkubo, I.: Human gene for β -microseminoprotein: its promoter structure and chromosomal localization. *J. Biochem. (Tokyo)*, 117 (1995) 346-352.
- 2550 Okayama, N. and Noriyasu, H.: (PCR-SSCP of silver stain for analysis of β -thalassaemia mutation). *Igaku Kensa*, 44 (1995) 18-22; *C.A.*, 122 (1995) 152546h.
- 2551 Oldroyd, N.J., Urquhart, A.J., Kimpton, C.P., Millican, E.S., Watson, S.K., Downes, T. and Gill, P.D.: A highly discriminating octoplex short tandem repeat polymerase chain reaction system suitable for human individual identification. *Electrophoresis (Weinheim)*, 16 (1995) 334-337.
- 2552 Oto, M., Suehiro, T., Akiyama, Y. and Yuasa, Y.: Microsatellite instability in cancer identified by non-gel sieving capillary electrophoresis. *Clin. Chem. (Washington)*, 41 (1995) 482-483.
- 2553 Pan, J.-P., Chiang, A.-N., Tai, J.J., Wang, S.-P. and Chang, M.-S.: Restriction fragment length polymorphism of apolipoprotein B gene in Chinese population with coronary heart disease. *Clin. Chem. (Washington)*, 41 (1995) 424-429.
- 2554 Rutberg, S.E., Fuchs, S.Y. and Ronai, Z.: Ultraviolet irradiation and c-jun over-expression regulates replication of polyoma sequences in WOP cells through a PEBP2 binding site. *Biochim. Biophys. Acta*, 1261 (1995) 90-98.
- 2555 Saad, F.A., Halliger, B., Mueller, C.R., Roberts, R.G. and Danieli, G.A.: Single base substitutions are detected by double strand conformation analysis. *Nucleic Acids Res.*, 22 (1994) 4352-4353; *C.A.*, 122 (1995) 24770q.
- 2556 Samigullin, T.Kh., Miroshnichenko, G.P., Antonov, A.S., Yakovlev, S.P. and Yankovskaya, M.B.: (A comparative study of ribosomal DNA repeats of fruit plants of the *Rosaceae juss* family). *Biokhimiya (Moscow)*, 59 (1994) 1349-1359.

- 2557 Schlax, P.E., Jr., Capp, M.W. and Record, M.T., Jr.: Preparative-scale purification of DNA restriction fragments by continuous-flow gel electrophoresis. *BioTechniques*, 18 (1995) 94-100; C.A., 122 (1995) 75839b.
- 2558 Schuetz, J.D. and Guzelian, P.S.: Isolation of CYP3A5P cDNA from human liver: a reflection of a novel cytochrome P-450 pseudogene. *Biochim. Biophys. Acta*, 1261 (1995) 161-165.
- 2559 Shimadzu, M., Nunoi, H., Terasaki, H., Ninomiya, R., Iwata, M., Kanegasaka, S. and Matsuda, I.: Structural organization of the gene for CD40 ligand: molecular analysis for diagnosis of X-linked hyper-IgM syndrome. *Biochim. Biophys. Acta*, 1260 (1995) 67-72.
- 2560 Sidman, C.L. and Shaffer, D.J.: Large-scale genomic comparison using two-dimensional DNA gels. *Genomics*, 23 (1994) 15-22; C.A., 122 (1995) 179497e.
- 2561 Smit, J.J.M., Mol, C.A.A.M., van Deemter, L., Wagenaar, E., Schinkel, A.H. and Borst, P.: Characterization of the promoter region of the human MDR3 P-glycoprotein gene. *Biochim. Biophys. Acta*, 1261 (1995) 44-56.
- 2562 Snopkova, S., Goetz, F., Doskar, J. and Rosypal, S.: Pulsed-field gel electrophoresis of the genomic restriction fragments of coagulase-negative staphylococci. *FEMS Microbiol. Lett.*, 124 (1994) 131-140; C.A., 122 (1995) 24799f.
- 2563 Sokolova, I.A., Cowan, K.H. and Schneider, E.: Ca²⁺/Mg²⁺-dependent endonuclease activation is an early event in VP-16-induced apoptosis of human breast cancer MCF7 cells *in vitro*. *Biochim. Biophys. Acta*, 1266 (1995) 135-142.
- 2564 Srinivasan, K.: Development of capillary electrophoretic separation methods for PCR-amplified DNA products. Avail. *Univ. Microfilms Int.*, Order No. DA9426898, 1993, 244 pp.; C.A., 122 (1995) 155528q.
- 2565 Steffens-Nakken, H.M., Zwart, G. and van den Bergh, F.A.J.T.M.: Validation of allele-specific polymerase chain reaction for DNA typing of HLA-B27. *Clin. Chem. (Washington)*, 41 (1995) 687-692.
- 2566 Tanaka, M., Fukada, S., Matsuyama, M. and Nagahama, Y.: Structure and promoter analysis of the cytochrome P-450 aromatase gene of the teleost fish, medaka (*Oryzias latipes*). *J. Biochem (Tokyo)*, 117 (1995) 719-725.
- 2567 Tang, L., Pelech, S. and Berger, J.D.: Isolation of the cell cycle control gene *cdc2* from *Paramecium tetraurelia*. *Biochim. Biophys. Acta*, 1265 (1995) 161-167.
- 2568 Thonnard, J., Deldime, F., Heusterspreute, M., Delepaut, B., Hanon, F., de Bruyère, M. and Philippe, M.: HLA class II genotyping: two assay systems compared. *Clin. Chem. (Washington)*, 41 (1995) 553-556.
- 2569 Tomisaki, R., Baba, Y. and Tshako, M.: (Rapid separation of DNA sequencing reaction product using capillary electrophoresis). *Bunseki Kagaku*, 43 (1994) 1205-1207; C.A., 122 (1995) 75815r.
- 2570 Wan, Q. and Schuchman, E.H.: A novel polymorphism in the human acid sphingomyelinase gene due to size variation of the signal peptide region. *Biochim. Biophys. Acta*, 1270 (1995) 207-210.
- 2571 Wang, Y., Ju, J., Carpenter, B.A., Atherton, J.M., Sensabaugh, G.F. and Mathies, R.A.: Rapid sizing of short tandem repeat alleles using capillary array electrophoresis and energy-transfer fluorescent primers. *Anal. Chem.*, 67 (1995) 1197-1203.
- 2572 Wenz, H.M.: Capillary electrophoresis as a technique to analyze sequence-induced anomalously migrating DNA fragments. *Nucleic Acids Res.*, 22 (1994) 4002-4008; C.A., 122 (1995) 47727v.
- 2573 Wijnen, J., Fodde, R. and Khan, P.M.: DGGE polymorphism in intron 10 of MSH2, the HNPCC gene. *Hum. Mol. Genet.*, 3 (1994) 2268; C.A., 122 (1995) 98485x.
- 2574 Woolley, A.T. and Mathies, R.A.: Ultra-high-speed DNA fragment separations using microfabricated capillary array electrophoresis chips. *Proc. Natl. Acad. Sci. U.S.A.*, 91 (1994) 11348-11352; C.A., 122 (1995) 47745z.
- 2575 Yasuoka, A., Abe, K., Saigo, K., Arai, S. and Emori, Y.: Molecular cloning of a fish gene encoding a novel seven-transmembrane receptor related distantly to catecholamine, histamine, and serotonin receptors. *Biochim. Biophys. Acta*, 1235 (1995) 467-469.
- 2576 Zheleznyaya, L.A., Fedorov, O.V., Brik, A.F. and Matvienko, N.I.: (The nucleotide sequence of the flagellin gene of *Escherichia coli* B38). *Biokhimiya (Moscow)*, 59 (1994) 1621-1637.
- See also 1876, 1881, 2296, 2318, 2364, 2370, 2381, 2395, 2408, 2433, 2454, 2462, 2485, 2492.
22. ALKALOIDS
- 2577 Zhang, Z.-Y., Fasco, M.J. and Kaminsky, L.S.: Determination of theophylline and its metabolites in rat liver microsomes and human urine by capillary electrophoresis. *J. Chromatogr. B*, 665 (1995) 201-208.
23. OTHER SUBSTANCES CONTAINING HETEROCYCLIC NITROGEN
- 23c. *Indole derivatives and plant hormones (gibberelins)*
- See 2004.
24. ORGANIC SULPHUR COMPOUNDS (INCL. GLUCOSINOLATES)
- 2578 Okafo, G.N., Rana, K.K. and Camillery, P.: Improved separation of diastereoisomers in capillary electrophoresis using a mixture of β -cyclodextrin and sodium taurodeoxycholate. *Chromatographia*, 39 (1994) 627-630.
25. ORGANIC PHOSPHORUS COMPOUNDS (INCL. SUGAR PHOSPHATES)
- See 1859, 2013, 2022, 2080, 2099, 2168, 2365.
26. ORGANOMETALLIC AND RELATED COMPOUNDS
- 26a. *Organometallic compounds*
- See 2662.

26c. Coordination compounds

- 2579 Lee, H.J., Lee, S.H., Chung, K.S. and Lee, K.W.: Determination of Pd(II) and Pt(II) metacyanocomplexes using capillary electrophoresis. *Bull. Korean Chem. Soc.*, 15 (1994) 945-949; *C.A.*, 122 (1995) 150395j.
- 2580 Tewari, B.B., Nautiyal, A.P., Kamaluddin, and Srivastava, S.K.: (Studies on electrophoretic behavior of biologically important iron(III) and chromium(III) binary complexes with cysteine). *Zh. Neorg. Khim.*, 39 (1994) 1170-1172; *C.A.*, 122 (1995) 27074b.

See also 2658.

27. VITAMINS AND VARIOUS ANIMAL GROWTH FACTORS (NON-PEPTIDIC)

- 2581 Shi, H., Ma, Y., Humphrey, J.H. and Craft, N.E.: Determination of vitamin A in dried human blood spots by high-performance capillary electrophoresis with laser-excited fluorescence detection. *J. Chromatogr. B*, 665 (1995) 89-96.

28. ANTIBIOTICS

- 2582 Altria, K.D., Clayton, N.G., Harden, R.C., Makwana, J.V. and Portsmouth, M.J.: Inter-company cross validation exercise on capillary electrophoresis. Quantitative determination of drug counter-ion level. *Chromatographia*, 40 (1995) 47-50.
- 2583 Blais, B.W., Cunningham, A. and Yamazaki, H.: A novel immunofluorescence capillary electrophoresis assay system for the determination of chloramphenicol. *Food Agric. Immunol.*, 6 (1994) 409-417; *C.A.*, 122 (1995) 185653u.
- 2584 Dunn, D.S., Raghavan, S. and Volz, R.G.: Ciprofloxacin attachment to porous-coated titanium surfaces. *J. Appl. Biomater.*, 5 (1994) 325-331; *C.A.*, 122 (1995) 89315q.
- 2585 Gu, J.-L. and Fu, R.-N.: Separation of chiral isomers of chloramphenicol metabolite by capillary zone electrophoresis using cyclodextrin as chiral selector. *Chin. Sci. Bull.*, 39 (1994) 1801-1805; *C.A.*, 122 (1995) 150708p.
- 2586 Kinashi, H., Mori, E., Hatani, A. and Nimi, O.: Isolation and characterization of linear plasmids from lankacidin-producing *Streptomyces* species. *J. Antibiot.*, 47 (1994) 1447-1455.

29. INSECTICIDES, PESTICIDES AND OTHER AGROCHEMICALS

29e. Herbicides

- 2587 Galceran, M.T., Carneiro, M.C. and Puignou, L.: Capillary electrophoresis of quaternary ammonium ion herbicides: paraquat, diquat and difenzoquat. *Chromatographia*, 39 (1994) 581-586.
- 2588 Song, L., Ou, Q., Yu, W. and Li, G.: Separation of six phenylureas and chlorsulfuron standards by micellar, mixed micellar and microemulsion electrokinetic chromatography. *J. Chromatogr. A*, 699 (1995) 371-382.

29f. Fungicides

- 2589 Maillard, J.L., Favreau, C. and Reboud-Ravaux, M.: Role of monocyte/macrophage derived matrix-metalloproteinases (gelatinases) in prolonged skin inflammation. *Clin. Chim. Acta*, 233 (1995) 61-74.

30. SYNTHETIC AND NATURAL DYES

30a. Synthetic dyes

See 1941.

30b. Chloroplast and other natural pigments

- 2590 Coffey, J.S. and Castle, L.: Analysis for caramel color (Class III). *Food Chem.*, 51 (1994) 413-416; *C.A.*, 122 (1995) 30024r.
- 2591 Watanabe, T., Kotokawa, N., Yamamoto, A. and Terabe, S.: (Separation by capillary electrophoresis of natural yellow pigments used in food). *Kuromatogurafi*, 15 (1994) 212-213; *C.A.*, 122 (1995) 158855e.

31. PLASTICS AND THEIR INTERMEDIATES

- 2592 Gittings, M.R. and Saville, D.A.: Electrophoretic mobility and dielectric response measurements on electrokinetically ideal polystyrene latex particles. *Langmuir*, 11 (1995) 798-800; *C.A.*, 122 (1995) 162028u.
- 2593 McNair, H.M. and Sun, X.: Capillary zone electrophoresis for polyimide composition analysis. *J. High Resolut. Chromatogr.*, 18 (1995) 115-116.
- 2594 Okada, T.: Non-aqueous capillary electrophoretic separation of polyethers and evaluation of weak complex formation. *J. Chromatogr. A*, 695 (1995) 309-317.
- 2595 Ono, M., Tanaka, C. and Yoshinaga, K.: (The analysis of water-soluble components in latex by capillary electrophoresis). *Takeda Kenkyushoho*, 53 (1994) 104-114; *C.A.*, 122 (1995) 135701n.

See also 1880, 2652.

32. DRUG ANALYSIS

32a. Drug analysis, general techniques

- 2596 Altria, K.D. and Kersey, M.T.: Capillary electrophoresis and pharmaceutical analysis: a survey of industrial applications and their status in the United Kingdom and United States. *LC-GC Int.*, 8, No. 4 (1995) 201-208.
- 2597 Jellum, E.: Chromatography and capillary electrophoresis in biomedicine. *Spec. Publ. - R. Soc. Chem.*, 154 (Reviews on Analytical Chemistry-Euroanalysis VIII) (1994) 308-316; *C.A.*, 122 (1995) 50306n - a review with 14 refs.

- 2598 Lloyd, D.K., Li, S. and Ryan, P.: Protein chiral selectors in free-solution capillary electrophoresis and packed-capillary electrochromatography. *J. Chromatogr. A*, 694 (1995) 285-296.
- 2599 Naylor, S., Tomlinson, A.J., Benson, L.M. and Gorrod, J.W.: Capillary electrophoresis and capillary electrophoresis-mass spectrometry in drug and metabolite analysis. *Eur. J. Drug Metab. Pharmacokinet.*, 19 (1994) 235-240; *C.A.*, 122 (1995) 23106r.
- 2600 Rashid, A., Xiang, B., Zhang, Z. and Au, D.: (Recent advances in high performance capillary electrophoresis (HPCE) and its application in the field of pharmaceutical analysis). *Zhongguo Yaoke Daxue Xuebao*, 25 (1994) 380-384; *C.A.*, 122 (1995) 197089c - a review with 58 refs.
- 2601 Soini, H., Riekkola, M.-L. and Novotny, M.V.: A note on aspects of sample preparation for drug analysis in serum by capillary electrophoresis. In: Stevenson, D. and Wilson, I.D. (Editors), *Sample Prep. Biomed. Environ. Anal., [Proc. Chromatogr. Soc. Int. Symp.]*, 1991, Plenum, New York, 1994, pp. 167-169; *C.A.*, 122 (1995) 23086j.
- 2602 Tanaka, Y. and Terabe, S.: Partial separation zone technique for the separation of enantiomers by affinity electrokinetic chromatography with proteins as chiral pseudo-stationary phases. *J. Chromatogr. A*, 694 (1995) 277-284.
- 2603 Yoshinaga, M. and Tanaka, M.: Chromatographic optical resolution in pharmaceutical analysis. *Jpn. Kokai Tokyo Koho JP 06,256,226 [94/256,226]* (Cl. C07B57/00), 13 Sep. 1994, Appl. 93/62,667, 26 Feb. 1993; 8 p.; *C.A.*, 122 (1995) 122271x.
- See also 1917, 1929, 1931, 2580.
- 32b. Antirheumatics and antiinflammatory drugs**
- 2604 Boonkerd, S., Lauwers, M., Detaevernier, M.R. and Michotte, Y.: Separation and simultaneous determination of the components in an analgesic tablet formulation by micellar electrokinetic chromatography. *J. Chromatogr. A*, 695 (1995) 97-102.
- 2605 Fanali, S. and Aturki, Z.: Use of cyclodextrins in capillary electrophoresis for the chiral resolution of some 2-arylpropionic acid non-steroidal anti-inflammatory drugs. *J. Chromatogr. A*, 694 (1995) 297-305.
- 32c. Autonomic and cardiovascular drugs**
- 2606 Lozano, R., Warren, F.V., Jr., Perlman, S. and Joseph, J.M.: Quantitative analysis of fosinopril sodium by capillary zone electrophoresis and liquid chromatography. *J. Pharm. Biomed. Anal.*, 13 (1995) 139-148.
- 32d. Central nervous system drugs**
- 2607 Chicharro, M., Zapardiel, A., Bermejo, E., Lopez, J.A.P. and Hernandez, L.: Direct determination of sympathomimetic drugs in human urine by capillary zone electrophoresis. *Quim. Anal. (Barcelona)*, 13 (1994) 134-137; *C.A.*, 122 (1995) 74102u.
- See also 1933.
- 32e. Chemotherapeutics (exc. cytostatics and antibiotics)**
- 2608 Cao, J. and Cross, R.F.: The separation of dihydrofolate reductase inhibitors and the determination of $pK_{a,1}$ values by capillary zone electrophoresis. *J. Chromatogr. A*, 695 (1995) 297-308.
- 2609 Taylor, R.B. and Reid, R.G.: Analysis of basic antimalarial drugs by CZE; Part 2. Validation and application to bioanalysis. *J. Pharm. Biomed. Anal.*, 13 (1995) 21-26.
- 32g. Other drug categories**
- 2610 Hsieh, Y.-L., Cai, J., Li, Y.-T. and Henion, J.D.: Detection of noncovalent FKBP-FK506 and FKBP-rapamycin complexes by capillary electrophoresis-mass spectrometry and capillary electrophoresis-tandem mass spectrometry. *J. Am. Soc. Mass Spectrom.*, 6 (1995) 85-90; *C.A.*, 122 (1995) 177576f.
- 2611 Noroski, J.E., Mayo, D.J. and Moran, M.: Determination of the enantiomer of a cholesterol-lowering drug by cyclodextrin-modified micellar electrokinetic chromatography. *J. Pharm. Biomed. Anal.*, 13 (1995) 45-52.
- 2612 Pietta, P.G., Bruno, A., Mauri, P.L., Gardana, C., Maffei-Facino, R. and Carini, M.: Determination of sunscreen agents by micellar electrokinetic chromatography. *J. Pharm. Biomed. Anal.*, 13 (1995) 229-235.
- See also 1933, 2624.
- 32h. Toxicological and forensic applications**
- 2613 Jefferies, T.M., Brammer, G., Zotou, A., Brough, P.A. and Gallagher, T.: Determination of anatoxin-a, homoanatoxin and propylanatoxin in cyanobacterial extracts by HPLC, GC-mass spectrometry and capillary electrophoresis. *Spec. Publ. - R. Soc. Chem.*, 149 (Detection Methods for Cyanobacterial Toxins) (1994) 34-39; *C.A.*, 122 (1995) 74057h.
- 2614 Lurie, I.S.: Analysis of seized drugs by capillary electrophoresis. In: Adamovics, J.A. (Editor), *Anal. Addict. Misused Drugs*, Dekker, New York, 1995, pp. 151-219; *C.A.*, 122 (1995) 232724e.
- See also 2444, 2456, 2483, 2517, 2536, 2607.
- 32i. Plant extracts**
- 2615 King, B.J., Lee, L.S., Rackemann, R.G. and Scott, P.T.: Preparation of extracts for electrophoresis from citrus leaves. *J. Biochem. Biophys. Methods*, 29 (1994) 295-305; *C.A.*, 122 (1995) 100905e.
- 2616 Song, L.G., Zhang, S.M., Ou, Q.Y. and Yu, W.L.: Studies of crude chloroform extracts of roots of *Podophyllum emodii* var. *chinensis* Sprague by micellar electrokinetic chromatography. *Chromatographia*, 39 (1994) 682-686.
- 33. CLINICO-CHEMICAL APPLICATIONS**
- 33a. General papers and reviews**
- 2617 Landers, J.P.: Clinical capillary electrophoresis. *Clin. Chem. (Washington)*, 41 (1995) 495-509 - a review with 72 refs.

- 2618 Young, D.S. and Tracy, R.P.: Clinical applications of two-dimensional electrophoresis. *J. Chromatogr. A*, 698 (1995) 163-179 - a review with 117 refs.
- 33b. *Complex mixtures and profiling (single compounds by cross-reference only)*
- 2619 Kowalchuk, R.M., Pollack, S.R. and Corcoran, T.A.: Zeta potential of bone from particle electrophoresis: solution composition and kinetic effects. *J. Biomed. Mater. Res.*, 29 (1995) 47-57; *C.A.*, 122 (1995) 170113u.
- See also 1958, 1966, 1978, 1979, 1983, 1985, 1995, 2139, 2144, 2170, 2186, 2251, 2253, 2259, 2274, 2316, 2327, 2504, 2505, 2510, 2526, 2542, 2543, 2548, 2552, 2553, 2565, 2568, 2584, 2589.
34. FOOD ANALYSIS
- 34a. *General papers and reviews*
- 2620 Canclon, P.F.: Capillary electrophoresis: a new tool in food analysis. *J. Assoc. Off. Anal. Chem.*, 78 (1995) 12-15 - a review with 95 refs.
- 2621 Delgado, C., Talou, T. and Gaset, A.: (Capillary electrophoresis: application and development in the domain of the food science and agrosresource industry). *Spectra Anal.*, 23 (1994) 42-46; *C.A.*, 122 (1995) 237990y - a review with many refs.
- 34b. *Complex mixtures (single compounds by cross-reference only)*
- 2622 Craig, A., Ritchie, A.H. and Mackie, I.M.: Determining the authenticity of raw reformed breaded scampi (*Nephrops norvegicus*) by electrophoretic techniques. *Food Chem.*, 52 (1995) 451-454; *C.A.*, 122 (1995) 185710k.
- 2623 Scobbie, A.E. and Mackie, I.M.: The use of sodium dodecyl sulphate-polyacrylamide gel electrophoresis to confirm the presence of salmon (*Salmo salar*) eggs in an illegal fish bait. *Electrophoresis (Weinheim)*, 16 (1995) 306-307.
- 2624 Thompson, C.O., Trenery, V.C. and Kemmerly, B.: Micellar electrokinetic capillary chromatographic determination of artificial sweeteners in low-Joule soft drinks and other foods. *J. Chromatogr. A*, 694 (1995) 507-514.
- See also 2051, 2134, 2160, 2167, 2190, 2194, 2195, 2196, 2583, 2590, 2591.
- 34c. *Organoleptically important compounds (flavors, odors, volatiles)*
- 2625 Golc-Wondra, A., Skocir, E. and Prosek, M.: Determination of monosodium glutamate in food products. *J. Planar Chromatogr.*, 8 (1995) 117-121.
35. ENVIRONMENTAL ANALYSIS
- 35a. *General papers and reviews*
- 2626 DeWitt, C.M. and Moore, K.H.: Capillary electrophoretic assay of xenobiotic acyl-CoA formation. *Am. Lab. (Shelton)*, 27 (1995) 28C, 28D, 28F, 28G, 28H, 28I; *C.A.*, 122 (1995) 182495h.
- 2627 Magi, B., Marzocchi, B., Lazzeri, C., Bini, L., Santucci, A. and Pallini, V.: Detection of xenobiotic-protein adducts: electrophoretic and immunochemical approaches. In: Renzoni, A. (Editor), *Contam. Environ.*, Lewis, Boca Raton, 1994, pp. 129-133; *C.A.*, 122 (1995) 25427v.
- 35c. *Water pollution (complex mixtures; single compounds by cross-reference only)*
- 2628 Praus, P.: Possibilities of simple steam distillation in combination with capillary isotachopheresis for the determination of chlorophenols in river and industrial waste waters. *Anal. Chim. Acta*, 302 (1995) 39-44.
- 2629 Yonekubo, J., Sasaki, H. and Takahashi, Y.: (Capillary ion analysis applied for forest environment and products. (II). Analysis of rain and acid washing water). *Kankyo Kagaku*, 4 (1994) 394-395; *C.A.*, 122 (1995) 196346x.
- See also 2666.
- 35d. *Soil pollution (complex mixtures; single compounds by cross-reference only)*
- 2630 Verschaeve, L. and Gilles, J.: Single cell gel electrophoresis assay in the earthworm for the detection of genotoxic compounds in soils. *Bull. Environ. Contam. Toxicol.*, 54 (1995) 112-119; *C.A.*, 122 (1995) 238565a.
36. SOME TECHNICAL PRODUCTS AND COMPLEX MIXTURES
- 36c. *Complex mixtures, technical products and unidentified compounds*
- 2631 Chan, T., Carpio, R. and VanMeurs, R.: The use capillary electrophoresis for the evaluation of post metal etch cleaning processes. *Proc.-Electrochem. Soc.*, 94-29 (1994) 260-269; *C.A.*, 122 (1995) 177202f.
- 2632 Zemann, A.J. and Bobleter, O.: Separation of biomass degradation products by capillary electrophoresis. In: Bridgwater, A.V. (Editor), *Adv. Thermochem. Biomass Convers.*, Blackie, Glasgow, 1994, pp. 953-965; *C.A.*, 122 (1995) 190674f.
- See also 2595.

37. CELLS, CELLULAR PARTICLES AND SUPRAMOLECULAR STRUCTURES
- 2633 Akiba, T., Nishi, A., Takaoki, M., Nagaoka, S. and Tomita, F.: Electrophoretic free mobility and viability of microbial cells: a preliminary study in preparation for space experiments. *Appl. Theor. Electrophor.*, 4 (1994) 65-69; *C.A.*, 122 (1995) 75832u.
- 2634 Bauer, J. (Editor): *Cell Electrophoresis*. CRC, Boca Raton, 1994, 328 pp.; *C.A.*, 122 (1995) 75997b.
- 2635 Bruckheimer, E.M., Gillum, K.D. and Schroit, A.J.: Colocalization of Rh polypeptides and the aminophospholipid transporter in dilauroylphosphatidylcholine-induced erythrocyte vesicles. *Biochim. Biophys. Acta*, 1235 (1995) 147-154.
- 2636 Cohly, H.H.P. and Das, S.K.: Cell electrophoresis using antibodies and antigens as ligands. In: Bauer, J. (Editor), *Cell Electrophor.*, CRC, Boca Raton, 1994, pp. 113-142; *C.A.*, 122 (1995) 50318t.
- 2637 Friedrich, U., Ruyters, G. and Bauer, J.: Cell electrophoresis in microgravity. In: Bauer, J. (Editor), *Cell Electrophor.*, CRC, Boca Raton, 1994, pp. 315-319; *C.A.*, 122 (1995) 50328w.
- 2638 Goering, R.V.: Rapid epidemiological evaluation of staphylococci by field inversion gel electrophoresis. *Zentralbl. Bakteriolog., Suppl.*, 26 (1994) 97-103; *C.A.*, 122 (1995) 234648g.
- 2639 Golovanov, M.V.: Electrophoresis of cells at a physiological ionic strength. In: Bauer, J. (Editor), *Cell Electrophor.*, CRC, Boca Raton, 1994, pp. 181-196; *C.A.*, 122 (1995) 50321p.
- 2640 Heinrich, J. and Wagner, H.: Electrode compartment-separating membranes for cell electrophoresis. In: Bauer, J. (Editor), *Cell Electrophor.*, CRC, Boca Raton, 1994, pp. 103-112; *C.A.*, 122 (1995) 50317s.
- 2641 Knisley, K.A. and Rodkey, L.S.: New buffer systems for cell electrophoresis. In: Bauer, J. (Editor), *Cell Electrophor.*, CRC, Boca Raton, 1994, pp. 145-161; *C.A.*, 122 (1995) 50319u.
- 2642 Kuzmanova, M., Ivanov, S., Nankova, V. and Markov, M.: Effects of extremely high frequency electromagnetic fields on electrophoretic mobility and ATP content in rat erythrocytes. *Bioelectrochem. Bioenerg.*, 35 (1994) 53-56; *C.A.*, 122 (1995) 26810b.
- 2643 Matsumura, H., Mori, F., Kawahara, K., Obata, C. and Furusawa, K.: Effect of amino acids, polypeptides and proteins on electrophoretic mobilities of phospholipid liposomes. *Colloids Surf., A*, 92 (1994) 87-93; *C.A.*, 122 (1995) 49598c.
- 2644 Morrison, D.R.: Cell electrophoresis in microgravity: past and future. In: Bauer, J. (Editor), *Cell Electrophor.*, CRC, Boca Raton, 1994, pp. 283-313; *C.A.*, 122 (1995) 50327v.
- 2645 Ohshima, H.: A simple expression of Henry's function for the retardation effect in electrophoresis of spherical colloidal particles. *J. Colloid Interface Sci.*, 168 (1994) 269-271; *C.A.*, 122 (1995) 90352f.
- 2646 Schutt, W., Hashimoto, N. and Shimizu, M.: Application of cell electrophoresis for clinical diagnosis. In: Bauer, J. (Editor), *Cell Electrophor.*, CRC, Boca Raton, 1994, pp. 255-266; *C.A.*, 122 (1995) 50325t.
- 2647 Serwer, P., Khan, S.A. and Griess, G.A.: Non-denaturing gel electrophoresis of biological nanoparticles: viruses. *J. Chromatogr. A*, 698 (1995) 251-261 - a review with 84 refs.
- 2648 Seshi, B.: Cell adhesion to proteins separated by lithium dodecyl sulfate-polyacrylamide gel electrophoresis and blotted onto a polyvinylidene difluoride membrane: a new cell-blotting technique. *J. Immunol. Methods*, 176 (1994) 185-201; *C.A.*, 122 (1995) 50641t.
- 2649 Slivinsky, G.G.: Simultaneous two-parameter measurements of the electrophoretic features of cell subpopulations and their different sedimentation characteristics. In: Bauer, J. (Editor), *Cell Electrophor.*, CRC, Boca Raton, 1994, pp. 199-217; *C.A.*, 122 (1995) 50322q.
- 2650 Todd, P.: The loading and unloading of cells in electrophoretic separations. In: Bauer, J. (Editor), *Cell Electrophor.*, CRC, Boca Raton, 1994, pp. 75-101; *C.A.*, 122 (1995) 50316r.
- 2651 Walter, H. and Widen, K.E.: Differential electrophoretic behavior in aqueous polymer solutions of red blood cells from Alzheimer patients and from normal individuals. *Biochim. Biophys. Acta*, 1234 (1995) 184-190.
- 2652 Zeltner, W.A., Wang, J.-F., Omatete, O.O., Janney, M.A., Tejedor-Tejedor-M.I., Anderson, M.A., Riman, R.E., Shanefield, D.J. and Adair, J.H.: Model hydrous oxide colloids for particle electrophoresis. In: Adair, J.H., Casey, J.A. and Venigalla, S. (Editors), *Handb. Charact. Tech. Solid-Solution Interface*, Am. Ceram. Soc., Westerville, 1993, pp. 87-103; *C.A.*, 122 (1995) 170829g - a review with 10 refs.
38. INORGANIC COMPOUNDS
- 38a. Cations
- 2653 Altria, K.D., Wood, T., Kitscha, R. and Roberts-McIntosh, A.: Validation of a capillary electrophoresis method for the determination of potassium counter-ion levels in an acidic drug salt. *J. Pharm. Biomed. Anal.*, 13 (1995) 33-38.
- 2654 Baraj, B., Martínez, M., Sastre, A. and Aguilar, M.: Simultaneous determination of Cr(III), Fe(III), Cu(II) and Pb(II) as UV-absorbing EDTA complexes by capillary zone electrophoresis. *J. Chromatogr. A*, 695 (1995) 103-111.
- 2655 Chen, G.J., Lee, N.M., Hu, C.C. and Liu, C.Y.: Chemical modification of capillary column for electrophoretic separations of transition metal ions. *J. Chromatogr. A*, 699 (1995) 343-351.
- 2656 Dadachova, E., Mirzadeh, S., Lambrecht, R.M., Hetherington, E.L. and Knapp, F.F., Jr.: Separation of carrier-free ¹⁶⁶Ho from Dy₂O₃ targets by partition chromatography and electrophoresis. *J. Radioanal. Nucl. Chem.*, 199 (1995) 115-123; *C.A.*, 122 (1995) 144930p.
- 2657 Jia, L., Zhang, H. and Hu, Z.: Determination of platinum(II) in the presence of rhodium(III), ruthenium(III), osmium(III) and iridium(III) by micellar electrokinetic capillary chromatography. *Anal. Lett.*, 28 (1995) 917-930.
- 2658 Latva, M., Ala-Kleme, T., Bjennes, H., Kankare, J. and Haapakka, K.: Time-resolved luminescence detection of europium(III) chelates in capillary electrophoresis. *Analyst (Cambridge)*, 120 (1995) 367-372.
- 2659 O'Keeffe, M., Dunemann, L., Theobald, A. and Svehla, G.: Capillary electrophoresis in speciation analysis. Investigations of metal-polyaminopolycarboxylate complexes and effects of metals on proteins. *Anal. Chim. Acta*, 306 (1995) 91-97.

- 2660 Stathakis, C. and Cassidy, R.M.: Effect of electrolyte composition in the capillary electrophoretic separation of inorganic/organic anions in the presence of cationic polymers. *J. Chromatogr. A*, 699 (1995) 353-361.
- 2661 Wrobel, K., Gonzales, E.B., Wrobel, K. and Sanz-Medel, A.: Aluminium and silicon speciation in human serum by ion-exchange high-performance liquid chromatography - electrothermal absorption spectrometry and gel electrophoresis. *Analyst (Cambridge)*, 120 (1995) 809-815.
- See also 1887, 1991, 2579, 2582.
- 38b. Anions
- 2662 Albert, M., Demesmay, C. and Rocca, J.L.: Analysis of organic and non-organic arsenious or selenious compounds by capillary electrophoresis. *Fresenius J. Anal. Chem.*, 351 (1995) 426-432.
- 2663 Boden, J., Bächmann, K., Kotz, L., Fabry, L. and Pahlke, S.: Application of capillary zone electrophoresis with an isotachophoretic initial state to determine anionic impurities on as-polished silicon wafer surfaces. *J. Chromatogr. A*, 696 (1995) 321-332.
- 2664 Fu, X. and Lue, J.: (Capillary electrophoresis of inorganic anions). *Fenxi Huaxue*, 22 (1994) 1019-1021; *C.A.*, 122 (1995) 177140j.
- 2665 Kelly, L., Burgi, D.S. and Nelson, R.J.: Controlled temperature anion separation by capillary electrophoresis. *U.S. US* 5,385,654 (Cl. 204-180.1; G01N27/26), 31 Jan. 1995, *Appl.* 88,439, 07 Jul. 1993; 16 p.; *C.A.*, 122 (1995) 177331x.
- 2666 Li, K. and Li, S.F.Y.: Speciation of selenium and arsenic compounds in natural waters by capillary zone electrophoresis after on-column preconcentration with field-amplified injection. *Analyst (Cambridge)*, 120 (1995) 361-366.
- 2667 Salimi-Moosavi, H. and Cassidy, R.M.: Capillary electrophoresis of inorganic anions in nonaqueous media with electrochemical and indirect UV detection. *Anal. Chem.*, 67 (1995) 1067-1073.
- 2668 Song, L., Ou, Q., Yu, W. and Xu, G.: Effect of high concentrations of salts in samples on capillary electrophoresis of anions. *J. Chromatogr. A*, 696 (1995) 307-319.
- 2669 Tindall, G.W. and Perry, R.L.: Separation of fast anions by capillary electrophoresis without flow reversal. *J. Chromatogr. A*, 696 (1995) 349-352.
- See also 1860, 1892, 2660, 2670.
39. RADIOACTIVE AND OTHER ISOTOPE COMPOUNDS
- 2670 Lucy, C.A. and McDonald, T.L.: Separation of chloride isotopes by capillary electrophoresis based on the isotope effect on ion mobility. *Anal. Chem.*, 67 (1995) 1074-1078.
- See also 2656.

Quality Assurance for Environmental Analysis

Method Evaluation within the Measurements and Testing Programme (BCR)

Edited by Ph. Quevauviller, E.A. Maier and B. Griepink

Techniques and Instrumentation in Analytical Chemistry, Volume 17

Quality assurance (QA) for environmental analysis is a growing feature of the nineties as is illustrated by the number of QA guidelines and systems which are being implemented nowadays. This book focuses on the technical aspects of quality assurance. The techniques used in different analytical fields are critically reviewed and existing tools for evaluating their performance are described. Particular reference is made to the activities of the Measurements and Testing Programme (BCR) of the European Commission towards the improvement of quality control of environmental analysis.

Contents: 1. Quality assurance for environmental analysis (Ph. Quevauviller *et al.*). 2. Development of ICPMS and ID-ICPMS with the determination of Pb and Hg in environmental matrices as an example (M. Campbell). 3. Detection of sources of error in the determination of Cr in environmental matrices by FAAS and ETAAS (G. Rauret *et al.*). 4. Analysis of environmental and biological samples by atomic spectroscopic methods (M. Hoenig, M.F. Gunn). 5. Validation of neutron activation analysis techniques (K. Heydorn). 6. Flow-through (bio)chemical sensors in environmental analysis (M.D. Luque de Castro, M. Válcárcel). 7. Fiber optical sensors applied to field measurements (C. Cámara *et al.*). 8. Chromium speciation in environmental and biological

samples (K. Vercoutere, R. Cornelis). 9. Determination of aluminium species in natural waters (B. Fairman, A. Sanz-Medel). 10. Selenium speciation analyses in water and sediment matrices (C. Cámara *et al.*). 11. Antimony speciation in water (M.B. de la Calle-Guntiñas *et al.*). 12. Arsenic speciation in environmental matrices (A. Amran *et al.*). 13. Mercury speciation in biological matrices (I. Drabæk, Å Iverfeldt). 14. Speciation analysis of organolead compounds. Status and future prospects (R. Lobinski *et al.*). 15. Speciation analysis of organotin by GC-AAS and GC-AES after extraction and derivatization (W.M.R. Dirx *et al.*). 16. High performance liquid chromatography - isotope dilution - inductively coupled plasma - mass spectrometry for lead and tin speciation in environmental samples (S.J. Hill *et al.*). 17. Speciation of organotin compounds in environmental samples by GC-MS (R. Morabito *et al.*). 18. Development of supercritical fluid extraction

procedures for the determination of organotin compounds in sediment (J.M. Bayona). 19. Hydride generation for speciation analyses using GC/AAS (R. Ritsema *et al.*). 20. Single and sequential extraction schemes for trace metal speciation in soil and sediment (A.M. Ure *et al.*). 21. Methods for the determination of chlorinated biphenyls in air (M. Morosini, K. Balschmitter). 22. Sample handling and determination of carbamate pesticides and their transformation products in various matrices (M. Honing *et al.*). 23. Method development for the determination of polycyclic aromatic hydrocarbons (PAHs) in environmental matrices (J. Jacob). 24. Method validation for the determination of dioxins (T. Rymen). Subject index.

©1995 670 pages **Hardbound**
Price: Dfl. 475.00 (US\$279.50)
ISBN 0-444-89955-3

ORDER INFORMATION

ELSEVIER SCIENCE B.V.
P.O. Box 330
1000 AH Amsterdam
The Netherlands
Fax: +31 (20) 485 2845

For USA and Canada:
P.O. Box 945, New York
NY 10159-0945
Fax: +1 (212) 633 3680

US\$ prices are valid only for the USA & Canada and are subject to exchange rate fluctuations; in all other countries the Dutch guilder price (Dfl.) is definitive. Customers in the European Union should add the appropriate VAT rate applicable in their country to the price(s). Books are sent postfree if prepaid.



ELSEVIER

An imprint of Elsevier Science

Carbohydrate Analysis

High Performance Liquid Chromatography and Capillary Electrophoresis

Edited by **Z. El Rassi**

Journal of Chromatography Library, Volume 58

The objective of the present book is to provide a comprehensive review of carbohydrate analysis by HPLC and HPCE by covering analytical and preparative separation techniques for all classes of carbohydrates including mono- and disaccharides; linear and cyclic oligosaccharides; branched heterooligosaccharides (e.g., glycans, plant-derived oligosaccharides); glycoconjugates (e.g., glycolipids, glycoproteins); carbohydrates in food and beverage; compositional carbohydrates of polysaccharides; carbohydrates in biomass degradation; etc.

The book will be of interest to a wide audience, including analytical chemists and biochemists, carbohydrate, glycoprotein and glycolipid chemists, molecular biologists, biotechnologists, etc. It will also be a useful reference work for both the experienced analyst and the newcomer as well as for users of HPLC and HPCE, graduates and postdoctoral students.

Contents: Part I. The Solute.

1. Preparation of carbohydrates for analysis by HPLC and HPCE (A.J. Mort, M.L. Pierce).

Part II. Analytical and Preparative Separations.

2. Reversed-phase and hydrophobic interaction chromatography of carbohydrates and glycoconjugates (Z. El Rassi).
3. High performance hydrophilic interaction chromatography of carbohydrates with

polar solvents (S.C. Churms).
4. HPLC of carbohydrates with cation- and anion-exchange silica and resin-based stationary phases (C.G. Huber, G.K. Bonn).
5. Analysis of glycoconjugates using high-pH anion-exchange chromatography (R.R. Townsend).
6. Basic studies on carbohydrate - protein interaction by high performance affinity chromatography and high performance capillary affinity electrophoresis using lectins as protein models (S. Honda).
7. Modern size exclusion chromatography of carbohydrates and glycoconjugates (S.C. Churms).
8. High performance capillary electrophoresis of carbohydrates and glycoconjugates (Z. El Rassi, W. Nashabeh).
9. Preparative HPLC of carbohydrates (K.B. Hicks).

Part III. The Detection.

10. Pulsed electrochemical detection of carbohydrates at gold electrodes following liquid chromatographic separation (D.C. Johnson,

W.R. LaCourse).
11. On-column refractive index detection of carbohydrates separated by HPLC and CE (A.E. Bruno, B. Krattiger).
12. Mass spectrometry of carbohydrates and glycoconjugates (C.A. Settineri, A.L. Burlingame).
13. Evaporative light scattering detection of carbohydrates in HPLC (M. Dreux, M. Lafosse).
14. Chiroptical detectors for HPLC of carbohydrates (N. Purdie).
15. Pre- and post-column detection-oriented derivatization techniques in HPLC of carbohydrates (S. Hase).
16. Post-column enzyme reactors for the HPLC determination of carbohydrates (L.J. Nagels, P.C. Maes).
17. Other direct and indirect detection methods of carbohydrates in HPLC and HPCE (Z. El Rassi, J.T. Smith).
Subject index.

©1995 692 pages Hardbound
Price: Dfl. 425.00 (US\$250.00)
ISBN 0-444-89981-2

ORDER INFORMATION

ELSEVIER SCIENCE B.V.
P.O. Box 330
1000 AH Amsterdam
The Netherlands
Fax: +31 (20) 485 2845

For USA and Canada:
P.O. Box 945, New York
NY 10159-0945
Fax: +1 (212) 633 3680

US\$ prices are valid only for the USA & Canada and are subject to exchange rate fluctuations; in all other countries the Dutch guilder price (Dfl.) is definitive. Customers in the European Union should add the appropriate VAT rate applicable in their country to the price(s). Books are sent postfree if prepaid.



ELSEVIER

An imprint of Elsevier Science

PUBLICATION SCHEDULE FOR THE 1995 SUBSCRIPTION

Journal of Chromatography A and *Journal of Chromatography B: Biomedical Applications*

MONTH	1994	J-J ^a	A	S	O	N	D	
Journal of Chromatography A	Vols. 683-688	689-708/1	708/2 709/1 709/2 710/1	710/2 711/1 711/2	712/1 712/2			The publication schedule for further issues will be published later.
Bibliography Section		713		714/1			714/2	
Journal of Chromatography B: Biomedical Applications		663-669	670/1 670/2	671/1 + 2	672/1 672/2	673/1 673/2	674/1 674/2	

^a Vol. 701/1 + 2 (Cumulative Indexes Vols. 652-700) expected in December.

INFORMATION FOR AUTHORS

(Detailed *Instructions to Authors* were published in *J. Chromatogr. A*, Vol. 657, pp. 463-469. A free reprint can be obtained by application to the publisher, Elsevier Science B.V., P.O. Box 330, 1000 AH Amsterdam, Netherlands.)

Types of Contributions. The following types of papers are published: Regular research papers (full-length papers), Review articles, Short Communications and Discussions. Short Communications are usually descriptions of short investigations, or they can report minor technical improvements of previously published procedures; they reflect the same quality of research as full-length papers, but should preferably not exceed five printed pages. Discussions (one or two pages) should explain, amplify, correct or otherwise comment substantively upon an article recently published in the journal. For Review articles, see inside front cover under Submission of Papers.

Submission. Every paper must be accompanied by a letter from the senior author, stating that he/she is submitting the paper for publication in the *Journal of Chromatography A* or *B*.

Manuscripts. Manuscripts should be typed in **double spacing** on consecutively numbered pages of uniform size. The manuscript should be preceded by a sheet of manuscript paper carrying the title of the paper and the name and full postal address of the person to whom the proofs are to be sent. As a rule, papers should be divided into sections, headed by a caption (e.g., Abstract, Introduction, Experimental, Results, Discussion, etc.). All illustrations, photographs, tables, etc., should be on separate sheets.

Abstract. All articles should have an abstract of 50-100 words which clearly and briefly indicates what is new, different and significant. No references should be given.

Introduction. Every paper must have a concise introduction mentioning what has been done before on the topic described, and stating clearly what is new in the paper now submitted.

Experimental conditions should preferably be given on a *separate* sheet, headed "Conditions". These conditions will, if appropriate, be printed in a block, directly following the heading "Experimental".

Illustrations. The figures should be submitted in a form suitable for reproduction, drawn in Indian ink on drawing or tracing paper. Each illustration should have a caption, all the *captions* being typed (with double spacing) together on a *separate sheet*. If structures are given in the text, the original drawings should be provided. Coloured illustrations are reproduced at the author's expense, the cost being determined by the number of pages and by the number of colours needed. The written permission of the author and publisher must be obtained for the use of any figure already published. Its source must be indicated in the legend.

References. References should be numbered in the order in which they are cited in the text, and listed in numerical sequence on a separate sheet at the end of the article. Please check a recent issue for the layout of the reference list. Abbreviations for the titles of journals should follow the system used by *Chemical Abstracts*. Articles not yet published should be given as "in press" (journal should be specified), "submitted for publication" (journal should be specified), "in preparation" or "personal communication".

Vols. 1-651 of the *Journal of Chromatography*; *Journal of Chromatography, Biomedical Applications* and *Journal of Chromatography, Symposium Volumes* should be cited as *J. Chromatogr.* From Vol. 652 on, *Journal of Chromatography A* (incl. Symposium Volumes) should be cited as *J. Chromatogr. A* and *Journal of Chromatography B: Biomedical Applications* as *J. Chromatogr. B*.

Dispatch. Before sending the manuscript to the Editor please check that the envelope contains four copies of the paper complete with references, captions and figures. One of the sets of figures must be the originals suitable for direct reproduction. Please also ensure that permission to publish has been obtained from your institute.

Proofs. One set of proofs will be sent to the author to be carefully checked for printer's errors. Corrections must be restricted to instances in which the proof is at variance with the manuscript.

Reprints. Fifty reprints will be supplied free of charge. Additional reprints can be ordered by the authors. An order form containing price quotations will be sent to the authors together with the proofs of their article.

Advertisements. The Editors of the journal accept no responsibility for the contents of the advertisements. Advertisement rates are available on request. Advertising orders and enquiries can be sent to the Advertising Manager, Elsevier Science B.V., Advertising Department, P.O. Box 211, 1000 AE Amsterdam, Netherlands; Tel: 31 (20) 485 3796; Fax: 31 (20) 485 3810. Courier shipments to street address: Molenwerf 1, 1014 AG Amsterdam, Netherlands. UK: T.G. Scott & Son Ltd., Venessa Bird, Portland House, 21 Narborough Road, Cosby, Leics. LE9 5TA, UK; Tel: (0116) 2750 521/2753 333; Fax: (0116) 2750 522. USA and Canada: Weston Media Associates, Daniel S. Lipner, P.O. Box 1110, Greens Farms, CT 06436-1110, USA; Tel: (203) 261 2500; Fax: (203) 261 0101.

